SUPPORT OF AQUATIC LIFE USES
IN BIG SPRING CREEK, FERGUS COUNTY, MONTANA
BASED ON THE COMPOSITION AND STRUCTURE
OF THE BENTHIC ALGAE COMMUNITY

Prepared for:

State of Montana
Department of Environmental Quality
P.O. Box 200901
Helena, Montana 59620-0901

Project Officer: Rebecca Ridenour
DEQ Contract No. 200012-3

STATE DOCUMENTS COLLECTION

MAY 28 2002

MONTANA STATE LIBRARY
1515 E. 6th AVE.
HELENA, MONTANA 59620

Prepared by:

Loren L. Bahls, Ph.D.
Hannaea
1032 Twelfth Avenue
Helena, Montana 59601

December 14, 2001

SUMMARY

In May and August 2001, periphyton samples were collected at six stations on Big Spring Creek near Lewistown, Montana for the purpose of assessing whether the creek is water-quality limited and in need of TMDLs. The samples were collected following DEQ standard operating procedures, processed and analyzed using standard methods for periphyton, and evaluated following modified USEPA rapid bioassessment protocols for wadeable streams.

Results from the samples collected in May could not be compared to biological criteria Montana streams because these samples were collected outside of the summer sampling "window". The sample collected below Lewistown in May contained an unusually large number of motile, mesotrophic to eutrophic diatoms that may indicate siltation and nutrient problems.

In August, the site just below Big Springs supported a cold-water flora that is typical of a spring creek. The moderate impairment noted here is caused by natural thermal stress.

Diatom metrics for site 02 above Lewistown indicated minor impairment from disturbance and possible low levels of toxic chemicals. Aquatic life uses were fully supported. Site 03 below Lewistown also provided full support of aquatic life uses, with minor impairment indicated from siltation and toxics.

Diatom metrics for site 05 near the mouth of Big Spring Creek indicated **moderate impairment and partial support of aquatic life uses**. The probable cause of this impairment is nutrient enrichment. The bioassessment ratings for sites 03 and 05 would be the same if diatom metrics for these sites were compared to criteria for prairie streams instead of mountain streams.

INTRODUCTION

This report evaluates the biological integrity, support of aquatic life uses, and probable causes of impairment to those uses, in Big Spring Creek near Lewistown, Montana. The purpose of this report is to provide information that will help the State of Montana determine whether Big Spring Creek is water-quality limited and in need of TMDLs.

The federal Clean Water Act directs states to develop water pollution control plans (Total Maximum Daily Loads or TMDLs) that set limits on pollution loading to water-quality limited waters. Water-quality limited waters are lakes and stream segments that do not meet water-quality standards, that is, that do not fully support their beneficial uses. The Clean Water Act and USEPA regulations require each state to (1) identify waters that are water-quality limited, (2) prioritize and target waters for TMDLs, and (3) develop TMDL plans to attain and maintain water-quality standards for all water-quality limited waters.

Evaluation of use support in this report is based on the species composition and structure of the periphyton (benthic algae, phytobenthos) community at six stream sites that were sampled in May and August, 2001. The periphyton community is a basic biological component of all aquatic ecosystems. Periphyton accounts for much of the primary production and biological diversity in Montana streams (Bahls et al. 1992).

Plafkin et al. (1989) and Stevenson and Bahls (1999) list several advantages of using periphyton in biological assessments:

- Algae are universally present in large numbers in all streams and unimpaired periphyton assemblages typically support a large number (>30) of species;
- Algae have rapid reproduction rates and short life cycles, making them useful indicators of short-term impacts;

- As primary producers, algae are most directly affected by physical and chemical factors, such as temperature, nutrients, dissolved salts, and toxins;
- Sampling is quick, easy and inexpensive, and causes minimal damage to resident biota and their habitat;
- Standard methods and criteria exist for evaluating the composition, structure, and biomass of algal associations;
- Identification to species is straightforward for the diatoms, for which there is a large body of taxonomic and ecological literature;
- Excessive algae growth in streams is often correctly perceived as a problem by the public.
- Periphyton and other biological communities reflect the *biological integrity*¹ of waterbodies; restoring and maintaining the biological integrity of waterbodies is a goal of the federal Clean Water Act;
- Periphyton and other biological communities integrate the effects of different stressors and provide a measure of their aggregate impact; and
- Periphyton and other biological communities may be the only practical means of evaluating impacts from non-point sources of pollution where specific ambient criteria do not exist (e.g., impacts that degrade habitat or increase nutrients).

Periphyton is a diverse assortment of simple photosynthetic organisms, called algae, and other microorganisms that live attached to or in close proximity of the stream bottom. Most algae, such as the diatoms, are microscopic. Diatoms are distinguished by having a cell wall composed of opaline glass--hydrated amorphous silica. Diatoms often carpet a stream bottom with a slippery brown film.

¹ *Biological integrity* is defined as "the ability of an aquatic ecosystem to support and maintain a balanced, integrated, adaptive community of organisms having a species composition, diversity, and functional organization comparable to that of natural habitats within a region" (Karr and Dudley 1981).

Some algae, such as the filamentous greens, are conspicuous and their excessive growth may be aesthetically displeasing, deplete dissolved oxygen, interfere with fishing and fish spawning, clog water filters and irrigation intakes, create tastes and odors in drinking water, and cause other problems.

PROJECT AREA AND SAMPLING SITES

The project area is located in Fergus County near the city of Lewistown (pop. 6,368) in central Montana. The Big Springs, seven miles southeast of Lewistown, generate most of the streamflow in Big Spring Creek. From Big Springs, Big Spring Creek flows northwesterly for about 30 miles through Lewistown and on to its confluence with the Judith River.

Major tributaries of Big Spring Creek, including aquifers feeding the Big Springs, head in the Big Snowy Mountains (maximum elevation 8,730 feet), an outlier of the Northern Rockies Ecoregion (Woods et al. 1999). Although the segment of Big Spring Creek addressed here is located in the Montana Valley and Foothill Prairies Ecoregion, water quality is dominated by the Big Springs. In the Montana Surface Water Quality Standards, the stream is classified B-1 above Lewistown and B-2 below Lewistown.

The surface geology of the Big Spring Creek watershed consists of Big Snowy dolomite and limestone in the headwaters, metamorphic rocks of the Kootenai Formation in the middle reach, and shales of the Colorado Group in the lower reach (Renfro and Feray 1972). Vegetation is alpine tundra and spruce-fir forest in the headwaters, mixed forest and grassland in the middle reach, and mixed grassland at lower elevations (USDA 1976).

Periphyton samples were collected at 2 sites on May 25 and at another 4 sites on August 20 and 22, 2001. The 2 sites

sampled on May 25 bracket the City of Lewistown (Map 1, Table 1). The four sites sampled in August span the reach from just below Big Springs to near the mouth of the stream (Maps 2 and 3, Table 1). Elevations at the sampling sites range from 4250 feet below Big Springs to 3475 feet near the mouth of the creek.

Recreation, fish and aquatic life uses in upper Big Spring Creek are threatened by land development and discharges from a fish hatchery (MDEQ 1998). Below Lewistown, aquatic life, fish and recreational uses are partially impaired by agriculture, channelization, on-site domestic wastewater, the outfall from the Lewistown wastewater treatment plant, stormwater runoff, animal confinement facilities, and silviculture (MDEQ 1998).

Three previous reports on Big Spring Creek and tributaries have been prepared for MDEQ by this consultant (Bahls 1999a, 1999b, and 2001).

METHODS

Periphyton samples were collected in May by Tom Pick, NRCS/DNRC, and in August by Rebecca Ridenour, MDEQ Monitoring and Data Management Bureau, following standard operating procedures of the MDEQ Planning, Prevention, and Assistance Division.

Using appropriate tools, microalgae were scraped, brushed, or sucked from natural substrates in proportion to the rank of those substrates at the study site. Macroalgae were picked by hand in proportion to their abundance at the site. All collections of microalgae and macroalgae were pooled into a common container and preserved with Lugol's solution.

The samples were examined to estimate the relative abundance and rank by biovolume of diatoms and genera of soft (non-diatom)

algae according to the method described by Bahls (1993). Soft algae were identified using Dillard (1999), Prescott (1978), Smith (1950), and Whitford and Schumacher (1984). These books also served as references on the ecology of the soft algae, along with Palmer (1969, 1977).

After the identification of soft algae, the raw periphyton samples were cleaned of organic matter using sulfuric acid and potassium dichromate, and permanent diatom slides were prepared using Naphrax, a high refractive index mounting medium, following *Standard Methods for the Examination of Water and Wastewater* (APHA 1998).

Between 410 and 458 diatom cells (820 to 916 valves) were counted at random and identified to species. The four volume series by Krammer and Lange-Bertalot (1986, 1988, 1991a, 1991b) was used as the main taxonomic and autecological reference for the diatoms. Lowe (1974), Bahls et al. (1984), Van Dam et al. (1994), and Lange-Bertalot (1996) were also used as ecological references for the diatoms.

The diatom proportional counts were used to generate an array of diatom association metrics (Table 2). A metric is a characteristic of the biota that changes in some predictable way with increased human influence (Barbour et al. 1999).

Metric values generated from the August samples were compared to numeric biocriteria or threshold values developed for streams in the Rocky Mountain Ecoregion of Montana (Table 3). These criteria are based on metric values measured in least-impaired reference streams (Bahls et al. 1992) and on metric values measured in streams that are known to be impaired by various sources and causes of pollution (Bahls 1993). The biocriteria in Table 3 do not apply to the May samples, which were collected outside of the summer sampling "window".

The criteria in Table 3 distinguish among four levels of impairment and three levels of aquatic life use support: no impairment or only minor impairment (full support); moderate impairment (partial support); and severe impairment (nonsupport). These impairment levels correspond to excellent, good, fair, and poor *biological integrity*, respectively.

Quality Assurance. Several steps were taken to assure that the study results are accurate and reproducible.

Replicate periphyton samples were collected from site 03. Both samples from this site were analyzed by *Hannaea* and the results were compared side-by-side (Tables 4 and 5). Diatom metrics generated from the replicate samples resulted in identical bioassessment ratings for this site: minor impairment and full support of aquatic life uses.

Upon receipt of the samples, station and sample information were recorded in a laboratory notebook and the samples were assigned a unique number compatible with the Montana Diatom Database, e.g., 2250-01. The first part of this number (2250) designates the sampling site (Big Spring Creek at County Farm below Lewistown); the second part of this number (01) designates the number of periphyton samples that have been collected at this site to date for which data have been entered into the Montana Diatom Database.

Sample observations and analyses of soft (non-diatom) algae were recorded in a lab notebook along with station and sample information provided by MDEQ. A portion of the raw sample was used to make duplicate diatom slides. After completing the diatom proportional count, the slide used for the count will be deposited in the University of Montana Herbarium in Missoula.

be retained by Hannaea in Helena.

RESULTS AND DISCUSSION

SAMPLE NOTES

BS-2. Sample is silty; contains *Elodea* and other plant material.

M22BSPRC01. Some moss is present.

M22BSPRC02. Visible mats of *Phormidium* are present.

M22BSPRC05. Sample is silty.

NON-DIATOM ALGAE

Diatoms dominated periphyton samples collected from the

Burleigh and County Farm sites in May (Table 4). Green algae and cyanobacteria were also common at both sites. The Burleigh site above Lewistown supported more than twice the number of algal genera than did the County Farm site below Lewistown.

In August, diatoms (*Diatoma hyemalis*) dominated the site above Castle Creek just below the Big Springs. Two other chrysophycean algae--*Tribonema* and *Vaucheria*--ranked 2nd and 3rd in biomass here. All of these algae are typical of cold, spring-fed streams. Green algae were absent from this site.

The mat-forming cyanobacterium *Phormidium* dominated the algal flora below Pike Creek in August (Table 4). Diatoms were common here and ranked second. Green algae were also absent at this site, which supported only 2 genera of non-diatom algae.

The filamentous and potentially nuisance-forming green alga *Cladophora* dominated the algal flora of Big Spring Creek at the two sites below Lewistown that were sampled in August (Table 4). Diatoms ranked 2nd in biovolume at both sites and cyanobacteria were also present, especially as epiphytes on *Cladophora* at BSPRC05. Replicate samples from BSPRC03 generated very similar results (Table 4).

DIATOMS

May Samples. Since they were collected outside of the summer sampling "window", diatom metrics generated from the May samples cannot be compared with the biocriteria in Table 3. This is because samples collected during the cool seasons in Montana tend to be dominated by a single species.

Nitzschia fonticola dominated the diatom assemblages at both sites sampled in May (Table 5). This is a cool-season, but clean

water diatom that indicates meso-eutrophic conditions (Van Dam et al. 1994). *N. fonticola* is also motile and is adapted to living on unstable substrates.

The two periphyton samples that were collected in May from above and below Lewistown shared 64% of their diatom floras (Table 5). This indicates that the two sites were very similar, floristically, and that little or no environmental change occurred between them. However, the two sites both supported a handful of teratological diatom valves, which may indicate the presence of toxic chemicals in the water.

August Samples. The sample collected just below Big Springs in August was dominated by *Diatoma hyemalis* (Table 5). This oligotrophic diatom indicates cold waters that do not vary in temperature by more than a few degrees seasonally. Although the Shannon species diversity and percent dominant species metrics for this site both indicated moderate impairment and partial support of aquatic life uses, the thermal stresses that cause this "impairment" may be considered natural and result from the discharge of the Big Springs.

Diatom metrics at sites 02 and 03 indicated minor impairment but full use support in August (Table 5). At site 02, a slightly elevated disturbance index and percent dominant species were due to an abundance of *Achnanthes minutissimum*. Site 02 only shared about 12% of its flora with site 01, indicating that the flora of Big Spring Creek had changed from a spring flora to more of a stream flora at this point.

A somewhat elevated number of motile diatoms at site 03 indicated minor impairment from siltation here (Table 5). Sites 02 and 03 shared only about a quarter of their floras, indicating that a moderate amount of environmental change had occurred between them. These sites bracket the City of Lewistown and the

outfall from the Lewistown wastewater treatment plant.

The two replicate samples from site 03 generated a percent community similarity index of 84% (Table 5). In addition, diatom metrics from both sites yielded the same bioassessment rating: minor impairment but with full support of aquatic life uses. These results are acceptable from a quality assurance standpoint.

A large percent dominant species indicated moderate impairment and partial support of aquatic life uses at BSPRC05 (Table 5). The dominant species here was the eutrophic diatom *Cocconeis pediculus*. This diatom has a concave valve surface, by which it is adapted to living as an epiphyte on *Cladophora* and other filamentous algae. Its abundance at site 05 is directly related to dominance at this site by *Cladophora* (Table 4). *Cladophora* and *Cocconeis* indicate that the probable cause of impairment at this site is nutrient enrichment.

Sites 03 and 05 shared only about a third of their diatom floras, indicating that a moderate amount of environmental change (degradation) occurred between them. Both sites are downstream from the outfall of the Lewistown wastewater treatment plant.

All of the sites sampled in August, except the one just below Big Springs, had a few teratological diatom valves. This may indicate small concentrations of toxic chemicals, including heavy metals. None of the sites sampled in May and August supported a large number of diatoms in the family Epithemiaceae. This indicates that phosphorus, not nitrogen, was likely the limiting nutrient in Big Spring Creek.

The overall impairment ratings for Big Spring Creek would not change if metrics for sites 03 and 05 in the B-2 segment below Lewistown were compared to criteria for plains streams instead of mountain streams. Site 03 would still have good

biological integrity (minor impairment due to teratological cells) and site 05 would have fair biological integrity due to elevated percent dominant species and depressed diatom diversity.

REFERENCES

- APHA. 1998. Standard Methods for the Examination of Water and Wastewater. 20th Edition. American Public Health Association, Washington, D.C.
- Bahls, L.L. 1979. Benthic diatom diversity as a measure of water quality. Proc. Mont. Acad. Sci. 38:1-6.
- Bahls, L.L. 1993. Periphyton Bioassessment Methods for Montana Streams (Revised). Montana Department of Health and Environmental Sciences, Helena.
- Bahls, L.L. 1999a. Use Support in Big Spring Creek Based on Periphyton Composition and Community Structure. Prepared for the Montana Department of Environmental Quality, Helena.
- Bahls, L.L. 1999b. Effects Of An Oil Spill on The Composition and Structure of the Periphyton Community in Casino Creek and Big Spring Creek near Lewistown, Montana. Prepared for the Montana Department of Environmental Quality, Helena, Contract No. 200012.
- Bahls, L.L. 2001. Biological Integrity of Cottonwood Creek, Fergus County, Montana, Based on the Composition and Structure of the Benthic Algae Community. Prepared for the Montana Department of Environmental Quality, Helena, Contract No. 200012-3.
- Bahls, L.L., E.E. Weber, and J. O. Jarvie. 1984. Ecology and Distribution of Major Diatom Ecotypes in the Southern Fort Union Coal Region of Montana. U.S. Geological Survey Professional Paper 1289, U.S. Government Printing Office, Washington.
- Bahls, L.L., Bob Bukantis, and Steve Tralles. 1992. Benchmark Biology of Montana Reference Streams. Montana Department of Health and Environmental Sciences, Helena.
- Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. 1999. Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish. Second Edition. EPA/841-B-99-002. U.S. EPA, Office of Water, Washington, D.C.

- Dillard, G.E. 1999. Common Freshwater Algae of the United States. J. Cramer, Berlin.
- Johansen, J.R. 1999. Diatoms of Aerial Habitats. Chapter 12 in Stoermer, E.F., and J.P. Smol (eds.), The Diatoms, Cambridge University Press, New York.
- Karr, J.R., and D.R. Dudley. 1981. Ecological perspectives on water quality goals. Environmental Management 5:55-69.
- Krammer, K., and H. Lange-Bertalot. 1986. Bacillariophyceae, Part 2, Volume 1: Naviculaceae. In Ettl, H., J. Gerloff, H. Heynig, and D. Mollenhauer (eds.), Freshwater Flora of Middle Europe. Gustav Fischer Publisher, New York.
- Krammer, K., and H. Lange-Bertalot. 1988. Bacillariophyceae, Part 2, Volume 2: Bacillariaceae, Epithemiaceae, Surirellaceae. In Ettl, H., J. Gerloff, H. Heynig, and D. Mollenhauer (eds.), Freshwater Flora of Middle Europe. Gustav Fischer Publisher, New York.
- Krammer, K., and H. Lange-Bertalot. 1991a. Bacillariophyceae, Part 2, Volume 3: Centrales, Fragilariaceae, Eunotiaceae. In Ettl, H., J. Gerloff, H. Heynig, and D. Mollenhauer (eds.), Freshwater Flora of Middle Europe. Gustav Fischer Publisher, Stuttgart.
- Krammer, K., and H. Lange-Bertalot. 1991b. Bacillariophyceae, Part 2, Volume 4: Achnanthaceae, Critical Supplement to Navicula (Lineolatae) and Gomphonema, Complete List of Literature for Volumes 1-4. In Ettl, H., G. Gartner, J. Gerloff, H. Heynig, and D. Mollenhauer (eds.), Freshwater Flora of Middle Europe. Gustav Fischer Publisher, Stuttgart.
- Lange-Bertalot, Horst. 1979. Pollution tolerance of diatoms as a criterion for water quality estimation. Nova Hedwigia 64:285-304.
- Lange-Bertalot, Horst. 1996. Red list of limnic diatoms from Germany. Schr.-R. f. Vegetationskde., H. 28, pp. 633-677. BfN, Bonn-Bad Godesberg.
- Lowe, R.L. 1974. Environmental Requirements and Pollution Tolerance of Freshwater Diatoms. EPA-670/4-74-005.
- McFarland, B.H., B.H. Hill, and W.T. Willingham. 1997. Abnormal *Fragilaria* spp. (Bacillariophyceae) in streams impacted by mine drainage. Jour. of Freshwater Ecology 12(1):141-149.
- MDEQ. 1998. Waterbodies in Need of TMDL Development. Montana Department of Environmental Quality, Helena.

- Palmer, C.M. 1969. A composite rating of algae tolerating organic pollution. *Journal of Phycology* 5:78-82.
- Palmer, C.M. 1977. *Algae and Water Pollution: An Illustrated Manual on the Identification, Significance, and Control of Algae in Water Supplies and in Polluted Water.* EPA-600/9-77-036.
- Plafkin, J.L., M.T. Barbour, K.D. Porter, S.K. Gross, and R.M. Hughes. 1989. *Rapid Bioassessment Protocols for Use in Rivers and Streams: Benthic Macroinvertebrates and Fish.* EPA 440-4-89-001.
- Prescott, G.W. 1978. *How to Know the Freshwater Algae.* Third Edition. Wm. C. Brown Company Publishers, Dubuque, Iowa.
- Renfro, H.B., and D.E. Feray. 1972. *Geological Highway Map of the Northern Rocky Mountain Region.* American Association of Petroleum Geologists, Tulsa, Oklahoma.
- Smith, G.M. 1950. *the Fresh-Water Algae of The United States.* McGraw-Hill Book Company, New York.
- Stevenson, R.J., and L.L. Bahls. 1999. *Periphyton Protocols.* Chapter 6 in Barbour, M.T., J. Gerritsen, B.D. Snyder, and J.B. Stribling. *Rapid Bioassessment Protocols for Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish.* Second Edition. EPA/841-B-99-002. U.S. EPA, Office of Water, Washington, D.C.
- USDA. 1976. *Climax Vegetation of Montana (map).* U. S. Department of Agriculture, Soil Conservation Service, Cartographic Unit, Portland.
- Van Dam, H., A. Mertens, and J. Sinkeldam. 1994. A coded checklist and ecological indicator values of freshwater diatoms from the Netherlands. *Netherlands Journal of Aquatic Ecology*, 28(1):117-133.
- Whitford, L.A., and G.J. Schumacher. 1984. *A Manual of Fresh-Water Algae (Revised).* Sparks Press, Raleigh, North Carolina.
- Whittaker, R.H. 1952. A study of summer foliage insect communities in the Great Smoky Mountains. *Ecological Monographs* 22:1-44.
- Woods, A.J., Omernik, J.M., Nesser, J.A., Sheldon, J., and Azevedo, S.H. 1999. *Ecoregions of Montana (color poster with map),* U.S. Geological Survey, Reston, Virginia.

Map 3

Table 1. Location of periphyton sampling stations on Big Spring Creek in 2001.

Station	Station Code	Sample Number	Latitude/ Longitude	Sample Date
Big Spring Creek at Burleigh above Lewistown	BS-1	2251-01	47 01 40 109 22 53	5/25/01
Big Spring Creek at County Farm below Lewistown	BS-2	2250-01	47 05 10 109 27 13	5/25/01
Big Spring Creek headwaters above Castle Creek	M22BSPRC01	1038-03	47 00 12 109 20 40	8/20/01
Big Spring Creek below Pike Creek	M22BSPRC02	1756-02	47 01 57 109 22 57	8/20/01
Big Spring Creek below Lewistown	M22BSPRC03	1822-02	47 05 30 109 27 34	8/20/01
Big Spring Creek below Lewistown (replicate)	M22BSPRC03r	1822-03	47 05 30 109 27 34	8/20/01
Big Spring Creek near mouth in section 36	M22BSPRC05	1759-02	47 11 46 109 38 02	8/22/01

Table 2. Diatom association metrics used to evaluate biological integrity in Montana streams: reference, range of values in Montana streams, and expected direction of metric response to increasing anthropogenic perturbation or natural stress.

Metric	Reference	Range of Values	Expected Response
Shannon Species Diversity	Bahls 1979	0.00-5.00+	Decrease ¹
Pollution Index ²	Bahls 1993	1.00-3.00	Decrease
Siltation Index ³	Bahls 1993	0.00-90.0+	Increase
Disturbance Index ⁴	Barbour et al. 1999	0.00-100.0	Increase
No. Species Counted	Bahls 1979, 1993	0-100+	Decrease ¹
Percent Dominant Species	Barbour et al. 1999	5.0-100.0	Increase
Percent Abnormal Cells	McFarland et al. 1997	0.0-20.0+	Increase
Similarity Index	Whittaker 1952	0.0-80.0+	Decrease
Percent Epithemiaceae	Stevenson & Pan 1999	0.0-80.0+	Decrease
Percent Aerophiles	Johansen 1999	0.0-100	Increase

¹ Shannon diversity and species richness may increase somewhat in naturally nutrient-poor mountain streams in response to slight to moderate increases in nutrients or sediment.

² Composite numeric expression of the pollution tolerances assigned by Lange-Bertalot (1979) to the common diatom species.

³ Sum of the percent abundances of all species in the genera *Navicula*, *Nitzschia*, and *Surirella*.

⁴ Percent abundance of *Achnanthes minutissima*

Table 3. Criteria for rating levels of biological integrity, environmental impairment or natural stress, and aquatic life use support in Wadeable **mountain** streams of Montana using selected metrics for benthic diatom associations. The lowest rating for any one metric is the overall rating for the study site.

Biological Integrity/ Impairment or Natural Stress/Use Support	Diversity Index (Shannon)	Pollution Index	Siltation Index	Disturbance Index	Number of Species Counted	Percent Dominant Species	Percent Abnormal Cells	Percent Similarity Index ¹
Excellent None/Full Support	>2.99	>2.50	<20.0	<25.0	>29	<25.0	0.0	>59.9
Good/Minor Full Support	2.00- 2.99	2.01- 2.50	20.0- 39.9	25.0- 49.9	20- 29	25.0- 49.9	>0.0- <1.0	40.0- 59.9
Fair/Moderate Partial Support	1.00- 1.99	1.50- 2.00	40.0- 59.9	50.0- 74.9	10- 19	50.0- 74.9	1.0- 9.9	20.0- 39.9
Poor/Severe Nonsupport	<1.00	<1.50	>59.9	>74.9	<10	>74.9	>9.9	<20.0

¹ The Similarity Index or Percent Community Similarity (Whittaker 1952) may be used to compare a study site to an unimpacted upstream control site on the same stream. This metric measures the degree of floristic similarity between diatom associations at the two sites and is the sum of the smaller of the two percent abundance values for each species that is common to both sites. Adjacent riffles on the same stream, without intervening tributaries or environmental perturbations, will generally have at least 60% of their diatom floras in common (Bahls 1993). PCS may also be used to gauge the relative amount of impairment or recovery that occurs between adjacent study sites: >59.9% = very similar floras, no change; 40.0-59.9% = somewhat similar floras, minor change; 20.0-39.9% = somewhat dissimilar floras, moderate change; <20.0% = very dissimilar floras, major change.

Table 4. Relative abundance of cells and ordinal rank by biovolume of diatoms (Class Bacillariophyceae) and genera of non-diatom algae in periphyton samples collected from Big Spring Creek in 2001: d = dominant; a = abundant; f = frequent; c = common; o = occasional; r = rare.

Taxa	Station						
	BS-1	BS-2	BSPRC01	BSPRC02	BSPRC03	BSPRC03r	BSPRC05
Chlorophyta (green algae)							
Ankistrodesmus							
Cladophora	o/7	o/3			o/5	o/6	o/8
Hormidium	o/10				d/1	d/1	d/1
Mougeotia	o/8						
Oedogonium	a/2	o/5			a/3	a/3	o/7
Scenedesmus							
Spirogyra	f/3						
Ulothrix	o/9	c/4					
Zygogonium	c/6						
Chrysophyta (golden algae)							
Bacillariophyceae	a/1	d/1		c/2	d/2	d/2	f/2
Tribonema	c/5		d/1				
Vaucheria			c/3				
			o/2				
Cyanophyta (cyanobacteria)							
Calothrix							
Chamaesiphon			o/6				a/3
Chroococcus							a/4
Oscillatoria	f/4	f/2	o/4	c/3	c/4	o/4	o/6
Phormidium			o/5	a/1		o/5	c/5
No. of Non-Diatom Genera	9	4	5	2	4	5	7

Table 5. Percent abundance of major diatom species and values of selected diatom association metrics for periphyton samples collected from Big Spring Creek in 2001. Underlined values indicate full support of aquatic life uses with minor impairment; **bold values** indicate partial support and moderate impairment; **underlined and bold values** indicate nonsupport and severe impairment; all other values indicate full support of aquatic life uses and no impairment when compared to criteria for mountain streams in Table 3.

Species/Metric (Trophic State ¹)	Station						
	BS-1	BS-2	BSPRC01	BSPRC02	BSPRC03	BSPRC03r	BSPRC05
<i>Achnanthes biasolettianum</i> (m)	2.02	0.23	0.60	8.10	0.66	0.33	
<i>Achnanthidium minutissimum</i> (t)	17.02	5.25	2.28	46.31	4.29	6.54	9.63
<i>Amphora pediculus</i> (e)	2.26	1.82	1.20	2.38	5.05	6.65	4.63
<i>Cocconeis pediculus</i> (e)	0.00		0.00		6.04	4.58	50.85
<i>Cymbella affinis</i> (e)	2.74	1.37		10.60	0.99	0.87	0.24
<i>Diatoma hyemalis</i> (o)	0.00		70.98				
<i>Diatoma vulgaris</i> (m-e)	1.07	2.05	0.48	0.24	9.45	10.47	
<i>Encyonopsis microcephala</i> (m-e)	1.19	0.46	0.00	4.88	0.88	1.85	7.68
<i>Gomphonema minutum</i> (e)	0.24	2.51	0.36		12.31	12.76	0.12
<i>Navicula cryptotenella</i> (t)	7.14	2.05		1.55	2.86	1.64	2.20
<i>Navicula tripunctata</i> (e)	4.29	6.84	0.48	0.95	8.24	3.93	1.22
<i>Nitzschia fonticola</i> (m-e)	35.71	52.45	6.35	4.29	12.75	13.41	0.24
<i>Nitzschia palea</i> (e)	0.71	7.30		0.48		0.65	0.85
<i>Rhoicosphenia curvata</i> (e)		0.80	0.00		15.27	11.89	2.07
Number of Cells Counted	420	438	417	420	455	458	410
Shannon Species Diversity	3.65	3.06	1.94	3.13	4.01	4.17	<u>2.95</u>
Pollution Index	2.79	2.73	2.89	2.80	2.80	2.78	<u>2.78</u>
Siltation Index	59.40	78.90	6.84	8.69	<u>33.52</u>	<u>29.12</u>	17.68
Disturbance Index	17.02	5.25	2.28	<u>46.31</u>	4.29	6.54	9.63
Number of Species Counted	46	41	22	32	37	43	38
Percent Dominant Species	35.71	52.45	70.98	<u>46.31</u>	15.27	13.41	50.85
Percent Abnormal Cells	0.95	0.68	0.00	<u>0.24</u>	<u>0.33</u>	<u>0.11</u>	<u>0.98</u>
Percent Epithemiaceae	0.00	0.23	0.00	0.48	0.22	0.00	0.49
Similarity Index	63.70		12.07	26.62	83.61		33.01

¹ e = eutraphentic; m = mesotraphentic; m-e = meso-eutraphentic; o = oligotraphentic; t = tolerant of a wide range of trophic state from oligotraphentic to eutraphentic.

APPENDIX A: DIATOM PROPORTIONAL COUNTS

Sample	Genus/Species/Variety	Pollution Tolerance Class	Count	Percent
225101	Achnanthes lanceolata	2	3	0.36
225101	Achnanthidium biasolettianum	3	17	2.02
225101	Achnanthidium minutissimum	3	143	17.02
225101	Amphipleura pellucida	2	1	0.12
225101	Amphora libyca	3	1	0.12
225101	Amphora pediculus	3	19	2.26
225101	Aulacoseira distans	3	1	0.12
225101	Cocconeis pediculus	3	0	0.00
225101	Cocconeis placentula	3	16	1.90
225101	Cymbella affinis	3	23	2.74
225101	Cymbella cistula	3	1	0.12
225101	Diatoma hyemalis	3	0	0.00
225101	Diatoma moniliformis	2	7	0.83
225101	Diatoma vulgare	3	9	1.07
225101	Encyonema silesiacum	2	11	1.31
225101	Encyonopsis microcephala	2	10	1.19
225101	Frustulia vulgaris	2	2	0.24
225101	Gomphonema angustatum	2	1	0.12
225101	Gomphonema minutum	3	2	0.24
225101	Gomphonema olivaceum	3	9	1.07
225101	Gomphonema parvulum	1	2	0.24
225101	Gyrosigma acuminatum	3	1	0.12
225101	Melosira varians	2	8	0.95
225101	Navicula acceptata	2	0	0.00
225101	Navicula capitata	2	2	0.24
225101	Navicula capitatoradiata	2	11	1.31
225101	Navicula cryptotenella	2	60	7.14
225101	Navicula pupula	2	2	0.24
225101	Navicula reichardtiana	2	10	1.19
225101	Navicula tripunctata	3	36	4.29
225101	Nitzschia acicularis	2	5	0.60
225101	Nitzschia apiculata	2	2	0.24
225101	Nitzschia dissipata	3	34	4.05
225101	Nitzschia fonticola	3	300	35.71
225101	Nitzschia frustulum	2	2	0.24
225101	Nitzschia heufleriana	3	8	0.95
225101	Nitzschia linearis	2	1	0.12
225101	Nitzschia palea	1	6	0.71
225101	Nitzschia paleacea	2	2	0.24
225101	Nitzschia sociabilis	2	14	1.67
225101	Nitzschia vermicularis	2	1	0.12
225101	Pseudostaurosira brevistriata	3	8	0.95
225101	Reimeria sinuata	3	2	0.24
225101	Staurosira construens	3	2	0.24
225101	Staurosirella leptostauron	3	28	3.33
225101	Staurosirella pinnata	3	6	0.71
225101	Surirella minuta	2	3	0.36
225101	Synedra nana	3	4	0.48
225101	Synedra ulna	2	4	0.48

Sample	Genus/Species/Variety	Pollution Tolerance Class	Count	Percent
225001	Achnanthidium biasolettianum	3	2	0.23
225001	Achnanthidium minutissimum	3	46	5.25
225001	Amphipleura pellucida	2	1	0.11
225001	Amphora pediculus	3	16	1.82
225001	Cocconeis placentula	3	2	0.23
225001	Cymbella affinis	3	12	1.37
225001	Denticula tenuis	3	2	0.23
225001	Diatoma mesodon	3	1	0.11
225001	Diatoma moniliformis	2	8	0.91
225001	Diatoma vulgaris	3	18	2.05
225001	Encyonopsis microcephala	2	4	0.46
225001	Gomphonema minutum	3	22	2.51
225001	Gomphonema olivaceum	3	25	2.85
225001	Melosira varians	2	2	0.23
225001	Navicula acceptata	2	2	0.23
225001	Navicula capitatoradiata	2	9	1.03
225001	Navicula cryptotenella	2	18	2.05
225001	Navicula gregaria	2	2	0.23
225001	Navicula halophila	2	2	0.23
225001	Navicula menisculus	2	3	0.34
225001	Navicula pelliculosa	1	2	0.23
225001	Navicula reichardtiana	2	5	0.57
225001	Navicula tripunctata	3	60	6.84
225001	Navicula trivialis	2	2	0.23
225001	Nitzschia acicularis	2	9	1.03
225001	Nitzschia archibaldii	2	4	0.46
225001	Nitzschia dissipata	3	24	2.74
225001	Nitzschia fonticola	3	460	52.45
225001	Nitzschia heufleriana	3	3	0.34
225001	Nitzschia linearis	2	4	0.46
225001	Nitzschia palea	1	64	7.30
225001	Nitzschia pusilla	1	10	1.14
225001	Nitzschia recta	3	1	0.11
225001	Nitzschia sociabilis	2	6	0.68
225001	Nitzschia vermicularis	2	2	0.23
225001	Reimeria sinuata	3	3	0.34
225001	Rhoicosphenia curvata	3	7	0.80
225001	Staurosirella leptostauron	3	7	0.80
225001	Staurosirella pinnata	3	2	0.23
225001	Synedra nana	3	1	0.11
225001	Synedra ulna	2	4	0.46

Sample	Genus/Species/Variety	Pollution Tolerance Class	Count	Percent
103803	Achnanthes lanceolata	2	11	1.32
103803	Achnanthidium biasolettianum	3	5	0.60
103803	Achnanthidium minutissimum	3	19	2.28
103803	Amphora pediculus	3	10	1.20
103803	Cocconeis pediculus	3	0	0.00
103803	Cocconeis placentula	3	7	0.84
103803	Cyclotella distinguenda	2	3	0.36
103803	Cymbella cistula	3	2	0.24
103803	Cymbella falaisensis	3	36	4.32
103803	Denticula tenuis	3	0	0.00
103803	Diatoma hyemalis	3	592	70.98
103803	Diatoma mesodon	3	5	0.60
103803	Diatoma vulgare	3	4	0.48
103803	Encyonema silesiacum	2	2	0.24
103803	Encyonopsis microcephala	2	0	0.00
103803	Fragilaria capucina	2	8	0.96
103803	Fragilaria vaucheriae	2	0	0.00
103803	Gomphonema angustatum	2	37	4.44
103803	Gomphonema minutum	3	3	0.36
103803	Gomphonema parvulum	1	13	1.56
103803	Karayevia clevei	3	2	0.24
103803	Meridion circulare	3	6	0.72
103803	Navicula capitatoradiata	2	0	0.00
103803	Navicula decussis	3	0	0.00
103803	Navicula lundii	2	0	0.00
103803	Navicula minima	1	0	0.00
103803	Navicula tripunctata	3	4	0.48
103803	Nitzschia fonticola	3	53	6.35
103803	Rhoicosphenia curvata	3	0	0.00
103803	Staurosirella leptostauron	3	0	0.00
103803	Staurosirella pinnata	3	5	0.60
103803	Synedra ulna	2	7	0.84

Sample	Genus/Species/Variety	Pollution Tolerance Class	Count	Percent
175602	Achnanthyidum biasolettianum	3	68	8.10
175602	Achnanthyidum minutissimum	3	389	46.31
175602	Amphipleura pellucida	2	0	0.00
175602	Amphora inariensis	3	4	0.48
175602	Amphora pediculus	3	20	2.38
175602	Aulacoseira distans	3	3	0.36
175602	Cocconeis placentula	3	23	2.74
175602	Cymbella affinis	3	89	10.60
175602	Cymbella cistula	3	2	0.24
175602	Denticula kuetzingii	3	2	0.24
175602	Denticula tenuis	3	2	0.24
175602	Diatoma moniliformis	2	31	3.69
175602	Diatoma vulgaris	3	2	0.24
175602	Encyonema brehmii	2	6	0.71
175602	Encyonema silesiacum	2	22	2.62
175602	Encyonopsis microcephala	2	41	4.88
175602	Fragilaria vaucheriae	2	4	0.48
175602	Gomphonema angustatum	2	4	0.48
175602	Gomphonema olivaceum	3	2	0.24
175602	Gomphonema parvulum	1	3	0.36
175602	Gomphonema pumilum	3	9	1.07
175602	Navicula acceptata	2	0	0.00
175602	Navicula capitatoradiata	2	4	0.48
175602	Navicula cryptotenella	2	13	1.55
175602	Navicula pupula	2	0	0.00
175602	Navicula reichardtiana	2	2	0.24
175602	Navicula tripunctata	3	8	0.95
175602	Nitzschia acicularis	2	2	0.24
175602	Nitzschia dissipata	3	2	0.24
175602	Nitzschia fonticola	3	36	4.29
175602	Nitzschia palea	1	4	0.48
175602	Nitzschia sociabilis	2	2	0.24
175602	Reimeria sinuata	3	11	1.31
175602	Staurosirella leptostauron	3	7	0.83
175602	Synedra ulna	2	23	2.74

Sample	Genus/Species/Variety	Pollution Tolerance Class	Count	Percent
182202	Achnanthes lanceolata	2	2	0.22
182202	Achnanthidium biasolettianum	3	6	0.66
182202	Achnanthidium minutissimum	3	39	4.29
182202	Amphora inariensis	3	2	0.22
182202	Amphora pediculus	3	46	5.05
182202	Aulacoseira islandica	3	1	0.11
182202	Cocconeis pediculus	3	55	6.04
182202	Cocconeis placentula	3	31	3.41
182202	Cymbella affinis	3	9	0.99
182202	Denticula kuetzingii	3	2	0.22
182202	Diatoma moniliformis	2	18	1.98
182202	Diatoma vulgare	3	86	9.45
182202	Encyonema silesiacum	2	27	2.97
182202	Encyonopsis microcephala	2	8	0.88
182202	Gomphoneis herculeana	3	1	0.11
182202	Gomphonema minutum	3	112	12.31
182202	Gomphonema olivaceum	3	2	0.22
182202	Gomphonema parvulum	1	4	0.44
182202	Gyrosigma attenuatum	3	2	0.22
182202	Karayevia clevei	3	2	0.22
182202	Melosira varians	2	1	0.11
182202	Navicula acceptata	2	2	0.22
182202	Navicula capitatoradiata	2	30	3.30
182202	Navicula cryptotenella	2	26	2.86
182202	Navicula gregaria	2	3	0.33
182202	Navicula menisculus	2	4	0.44
182202	Navicula pupula	2	2	0.22
182202	Navicula reichardtiana	2	31	3.41
182202	Navicula tripunctata	3	75	8.24
182202	Nitzschia capitellata	2	0	0.00
182202	Nitzschia dissipata	3	8	0.88
182202	Nitzschia fonticola	3	116	12.75
182202	Nitzschia inconspicua	2	2	0.22
182202	Nitzschia sociabilis	2	2	0.22
182202	Nitzschia supralitoria	2	4	0.44
182202	Rhoicosphenia curvata	3	139	15.27
182202	Staurosirella leptostauron	3	2	0.22
182202	Staurosirella pinnata	3	0	0.00
182202	Synedra ulna	2	8	0.88

Sample	Genus/Species/Variety	Pollution Tolerance Class	Count	Percent
182203	Achnanthes lanceolata	2	2	0.22
182203	Achnanthidium biasolettianum	3	3	0.33
182203	Achnanthidium minutissimum	3	60	6.54
182203	Amphipleura pellucida	2	0	0.00
182203	Amphora inariensis	3	2	0.22
182203	Amphora pediculus	3	61	6.65
182203	Cocconeis pediculus	3	42	4.58
182203	Cocconeis placentula	3	45	4.91
182203	Cymbella affinis	3	8	0.87
182203	Diatoma moniliformis	2	14	1.53
182203	Diatoma vulgaris	3	96	10.47
182203	Encyonema auerswaldii	2	1	0.11
182203	Encyonema brehmii	2	5	0.55
182203	Encyonema silesiacum	2	16	1.74
182203	Encyonopsis microcephala	2	17	1.85
182203	Fragilaria capucina	2	2	0.22
182203	Fragilaria vaucheriae	2	2	0.22
182203	Gomphoneis herculeana	3	2	0.22
182203	Gomphonema minutum	3	117	12.76
182203	Gomphonema olivaceum	3	8	0.87
182203	Gomphonema parvulum	1	9	0.98
182203	Gomphonema pumilum	3	2	0.22
182203	Gomphonema truncatum	3	2	0.22
182203	Navicula acceptata	2	6	0.65
182203	Navicula capitatoradiata	2	29	3.16
182203	Navicula cryptotenella	2	15	1.64
182203	Navicula lundii	2	3	0.33
182203	Navicula menisculus	2	2	0.22
182203	Navicula recens	2	2	0.22
182203	Navicula reichardtiana	2	39	4.25
182203	Navicula stroemii	2	2	0.22
182203	Navicula tripunctata	3	36	3.93
182203	Nitzschia capitellata	2	2	0.22
182203	Nitzschia dissipata	3	0	0.00
182203	Nitzschia fonticola	3	123	13.41
182203	Nitzschia palea	1	6	0.65
182203	Nitzschia sociabilis	2	2	0.22
182203	Reimeria sinuata	3	2	0.22
182203	Rhoicosphenia curvata	3	109	11.89
182203	Stauroneis smithii	2	2	0.22
182203	Staurosira construens	3	2	0.22
182203	Staurosirella leptostauron	3	11	1.20
182203	Staurosirella pinnata	3	3	0.33
182203	Synedra acus	2	2	0.22
182203	Synedra ulna	2	3	0.33

Sample	Genus/Species/Variety	Pollution Tolerance Class	Count	Percent
175902	Achnanthyidium minutissimum	3	79	9.63
175902	Amphora inariensis	3	2	0.24
175902	Amphora libyca	3	0	0.00
175902	Amphora pediculus	3	38	4.63
175902	Aulacoseira canadensis	3	0	0.00
175902	Aulacoseira distans	3	1	0.12
175902	Aulacoseira islandica	3	2	0.24
175902	Cocconeis pediculus	3	417	50.85
175902	Cocconeis placentula	3	27	3.29
175902	Cyclotella meneghiniana	2	1	0.12
175902	Cymbella affinis	3	2	0.24
175902	Encyonema silesiacum	2	3	0.37
175902	Encyonopsis microcephala	2	63	7.68
175902	Epithemia sorex	3	2	0.24
175902	Gomphonema minutum	3	1	0.12
175902	Gomphonema parvulum	1	10	1.22
175902	Gyrosigma attenuatum	3	1	0.12
175902	Navicula capitata	2	0	0.00
175902	Navicula capitatoradiata	2	26	3.17
175902	Navicula cryptotenella	2	18	2.20
175902	Navicula menisculus	2	5	0.61
175902	Navicula peregrina	2	2	0.24
175902	Navicula perpusilla	2	2	0.24
175902	Navicula radiosa	3	1	0.12
175902	Navicula reichardtiana	2	14	1.71
175902	Navicula tripunctata	3	10	1.22
175902	Navicula veneta	1	0	0.00
175902	Nitzschia acicularis	2	2	0.24
175902	Nitzschia capitellata	2	0	0.00
175902	Nitzschia dissipata	3	39	4.76
175902	Nitzschia fonticola	3	2	0.24
175902	Nitzschia frustulum	2	6	0.73
175902	Nitzschia heufleriana	3	2	0.24
175902	Nitzschia palea	1	7	0.85
175902	Nitzschia paleacea	2	4	0.49
175902	Nitzschia perminuta	3	2	0.24
175902	Nitzschia sigmoidea	3	1	0.12
175902	Nitzschia sociabilis	2	2	0.24
175902	Reimeria sinuata	3	1	0.12
175902	Rhoicosphenia curvata	3	17	2.07
175902	Rhopalodia gibba	2	2	0.24
175902	Staurosirella leptostauron	3	2	0.24
175902	Staurosirella pinnata	3	4	0.49