UC-NRLF

C 2 871 501

EX-11111
FEB 25 1921

University of Texas Bulletin

No. 1856: October 5, 1918

The Weno and Pawpaw Formations of the Texas Comanchean

BY

W. S. Adkins

On A New Ammonite Fauna of the Lower Turonian of Mexico

BY

Emil Bose

Bureau of Economic Geology and Technology
Division of Economic Geology
J. A. Udden, Director of the Bureau and Head of the Division

PUBLISHED BY
THE UNIVERSITY OF TEXAS
AUSTIN

EXCHANGE

EX LIBRIS

PALEO
LIBRARY

EARTH
SCIENCES
LIBRARY

University of Texas Bulletin

No. 1856: October 5, 1918

The Weno and Pawpaw Formations of the Texas Comanchean

BY
W. S. Adkins

On A New Ammonite Fauna of the Lower Turonian of Mexico

BY
Emil Böse

Bureau of Economic Geology and Technology
Division of Economic Geology
J. A. Udden, Director of the Bureau and Head of the Division

PUBLISHED BY THE UNIVERSITY SIX TIMES A MONTH, AND ENTERED AS
SECOND-CLASS MATTER AT THE POSTOFFICE AT AUSTIN, TEXAS,
UNDER THE ACT OF AUGUST 24, 1912

to mind
abstract

The benefits of education and of useful knowledge, generally diffused through a community, are essential to the preservation of a free government.

Sam Houston

Cultivated mind is the guardian genius of democracy. . . . It is the only dictator that freemen acknowledge and the only security that freemen desire.

Mirabeau B. Lamar

EXCHANGE

**The Weno and Pawpaw Formations of the
Texas Comanchean**

BY

W. S. Adkins

CONTENTS

INTRODUCTION.	9
STRATIGRAPHY.	13
Rocks underlying the Weno and Pawpaw Formations.	13
Pre-Comanchean.	13
Comanchean.	14
Thickness changes.	14
Lithological changes.	16
Weno and Pawpaw Formations.	25
Thickness changes.	25
North-South changes.	25
East-West changes.	26
Lithological changes.	26
Pawpaw Formation.	27
Weno Formation.	35
Stratigraphic Correlation.	40
North Central Texas and Southern Oklahoma.	40
South Central Texas.	40
West Texas and Mexico.	41
Europe and North Africa.	42
PALEONTOLOGY.	44
The Weno Fauna.	45
The Pawpaw Fauna.	47
Comparison of Pyrite Faunae.	53
Texas.	53
Kiamitia Clay Fauna.	54
Duck Creek Marl Fauna.	54
Denton Clay Fauna.	55
Grayson Fauna.	56
Del Rio Clay Fauna.	58
Summary of Pyrite Faunae.	60
Europe and Africa.	64
DESCRIPTIONS OF SPECIES.	68
CEPHALOPODA.	68
Nautilus texanus Shumard.	63
Nautilus sp.	68
Hamites tenawa Adkins and Winton.	69
Hamites sp. aff. armatus Sowerby.	69
Ancyloceras bendirei n. sp.	70
Hamulina worthensis n. sp.	71
Baculites comanchensis n. sp.	74
Turrilites bosquensis n. sp.	76
Turrilites worthensis Adkins and Winton.	78
Turrilites sp.	78
Scaphites hilli Adkins and Winton.	79

Engonoceras serpentinum (Cragin)	84
Engonoceras sp.	85
Flickia boesei n. sp.	85
Flickia (?) bosquensis n. sp.	87
Schloenbachia wenoensis n. sp.	89
Schloenbachia wintoni n. sp.	90
Mortoniceras worthense n. sp.	91
Acanthoceras worthense n. sp.	93
ASTEROIDEA.	95
Pentagonaster texensis Adkins and Winton.	95
Metopaster hortensae Adkins and Winton.	97
Comptonia wintoni n. sp.	97
Pentaceros americanus n. sp.	99
ECHINOIDEA.	101
Leiocidaris sp.	101
Goniopygus sp.	102
Pedinopsis symmetrica (Cragin)	102
Peltastes sp.	102
Salenia sp.	103
Cyphosoma volanum Cragin.	103
Holectypus limitis Böse.	103
Holaster sp. aff. simplex Shumard.	104
Epiaster wenoensis n. sp.	105
Epiaster aguilerae Böse.	109
Epiaster subobesus n. sp.	110
Enallaster wenoensis n. sp.	112
Enallaster bravoensis Böse.	114
Enallaster sp. aff. texanus (Roemer).	114
Hemiaster calvini Clark.	114
Hemiaster rivistae n. sp.	115
PELECYPODA.	118
Nucula nokonis n. sp.	118
Nucula wenoensis n. sp.	120
Arca washitaensis n. sp.	121
Gervilliopsis invaginata (White)	122
Ostrea carinata ? Lamarek.	122
Ostrea sp. aff. diluviana Linnaeus.	122
Ostrea (Alectryonia) quadriplicata Shumard.	123
Exogyra sp. aff. arietina Roemer	123
Pecten inconspicuus Cragin.	123
Pecten georgetownensis Kniker.	125
Venericardia wenoensis n. sp.	125
Protocardia sp. aff. multistriata (Shumard).	126
Corbula wenoensis n. sp.	127
Corbula basiniformis n. sp.	130
Corbula littoralis n. sp.	133
Cyprimeria washitaensis n. sp.	134
Remondia ? acuminata (Cragin)	136

GASTROPODA.	137
<i>Amberleya graysonensis</i> n. sp.	137
<i>Trochus laticonicus</i> n. sp.	138
<i>Helicocryptus mexicanus</i> Böse.	139
<i>Nerita</i> sp.	139
<i>Neritina</i> sp.	139
<i>Anchura mudgeana</i> White.	139
<i>Natica</i> sp.	140
<i>Lunatia</i> sp.	140
<i>Globiconcha</i> sp.	140
<i>Turritella graysonensis</i> n. sp.	140
<i>Turritella worthensis</i> n. sp.	142
<i>Cinulia washitaensis</i> n. sp.	143
PROTOZOA.	145
<i>Nodosaria texana</i> Conrad.	145
BIBLIOGRAPHY	146
PLATES 1—11.	149-170

LIST OF ILLUSTRATIONS

FIGURES

- Figure 1. Map of Texas and part of Oklahoma showing outcrops of the Weno and Pawpaw formations, and the lithological facies of the Pawpaw formation 11
- Figure 2. *Ancycloceras bendirei* n. sp., suture of the type individual, camera lucida drawing, x 5 71
- Figure 3. *Hamulina worthense* n. sp., sutures of type individual, camera lucida drawing, x 15 73
- Figure 4. *Baculites comanchensis* n. sp., sutures of type individual, camera lucida drawing, x 8 75
- Figure 5. *Turrilites bosquensis* n. sp., sutures, type individual, camera lucida drawing, x 8 77
- Figure 6. *Scaphites hilli* Adkins and Winton, type individual, diagrammatic restoration showing orientation and sutures, camera lucida drawing, x 12 82
- Figure 7. *Scaphites hilli* Adkins and Winton, young individual, showing the last five sutures, camera lucida drawing, x 10 83
- Figure 8. *Scaphites hilli* Adkins and Winton, external and internal suture, camera lucida drawing, x 10 84
- Figure 9. *Flickia boesei* n. sp., type individual, sutures, camera lucida drawing, x 8 86
- Figure 10. *Flickia bosquensis* n. sp., diagrammatic projection of sutures of type individual, camera lucida drawing, x 5 88
- Figure 11. *Schloenbachia wintoni* n. sp., type individual, last suture, camera lucida drawing, x 5. Keel region slightly distorted in type individual. 91
- Figure 12. *Mortoniceras worthense* n. sp., suture of individual, Plate 1, figure 6, camera lucida drawing, x 10 93
- Figure 13. *Acanthoceras worthense* n. sp., suture, camera lucida drawing, x 8 94

PLATES

- Plates 1—11. Fossils of the Weno and Pawpaw formations.....149-170

THE WENO AND PAWPAP FORMATIONS OF THE TEXAS COMANCHEAN¹

By W. S. ADKINS

INTRODUCTION

The Weno and Pawpaw formations are two thin formations lying near the top of the Comanchean series of strata, and are most typically developed in the region between the Red and the Brazos rivers. The importance of these formations, especially the Pawpaw, is much greater than their thickness would seem to indicate. The Pawpaw is a formation of small thickness, sharply limited lithologically both above and below, and contains a remarkable series of small pyritic fossils which are so sharply characterized and so distinctive in appearance, even in minute fragments such as are encountered in drilling, that this formation should be recognized in wells lying to the east of its outcrop and should furnish at least one certain and dependable stratigraphic level in the Comanchean. In the hope that the formation may have this value, its fauna has been described in some detail and the lateral variations in its lithological character indicated.² The same statements hold to a lesser extent for the Weno formation. The stratigraphic position of these two formations in the Comanchean series and their equivalents in the Central Texas section are given in the following table.

The Weno and Pawpaw formations are visible over a small areal extent, their outcrops aggregating somewhat more than 100 square miles, mainly in a narrow strip in Johnson, Tarrant, Denton, Cooke and Grayson Counties, Texas, and Love, Marshall, Bryan and Choctaw Counties, Oklahoma. In addition they have stratigraphic equivalents in Central and Western Texas and in Mexico, that will be discussed later. Their north-south distribution in Texas is between Lat. 32°N. and 34°N.; and their east-west distribution is practically unknown. In Oklahoma their east-west distribution is between Long. 95°15' and 97°; and their north-south distribution is unknown. The total length of their outcrop north of the Brazos River

¹Manuscript accepted June 15, 1920, published November, 1920.

²It is hoped that operators and others having well samples suspected of being Upper Comanchean will submit them for examination to the Bureau of Economic Geology at Austin.

³I wish to express my indebtedness to Professor W. M. Winton, with whom the preliminary work on these and other Comanchean formations was done jointly. The results of these studies will appear in forthcoming papers by both authors.

CORRELATION TABLE OF THE COMANCHEAN FORMATIONS OF CENTRAL TEXAS¹

Series	Division	Formations North of the Brazos River	Formations South of the Brazos River	Stage
UPPER CRETACEOUS		Eagleford	Eagleford	TURONIAN
Unconformity				
COMANCHEAN	WASHITA	Woodbine Grayson Mainstreet Pawpaw Weno Denton Fort Worth Duck Creek marl Duck Creek Ls. Kiamitia	Buda	UPPER CENOMANIAN
			Del Rio	
			Georgetown	LOWER CENOMANIAN
	FREDERICKS- BURG	Goodland Walnut	Edwards Comanche Peak Walnut	VRACONIAN
	TRINITY	Paluxy Glenrose (Basement Sands) ²	Glenrose Travis Peak	ALBIAN
				APTIAN
Unconformity				
	PENNSYLVANIAN	PENNSYLVANIAN		
Unconformity				

is about 110 miles in Texas and 100 miles in Oklahoma. The outcrop is S-shaped, and north and west of its limits in North Central Texas the formations are eroded away; while east and south of the outcrop, the Weno-Pawpaw rock sheet continues underground for an unknown distance, being buried beneath the more recent Comanchean and Cretaceous strata. This concealed extension of these formations still awaits exploration by means of detailed well logs and the examination of samples from drillings. The formations show distinct north-south changes along their outcrops and equally distinct east-west changes. As a whole, they thicken moderately from the Brazos river (near Blum) north to the Red River (north of Gainesville), and thence eastward down the Red River valley through Den-

¹See also: Böse, University of Texas Bulletin, 1902, p. 16, fig. 1.

²The Comanchean sea was transgressive northwards over much of Central Texas, and the Basement sand is not a single formation, but at different places is the time equivalent of various off-shore formations.

Fig. 1. Map of Texas and part of Oklahoma, showing outcrops of the Weno and Pawpaw formations and the lithological facies of the Pawpaw formation. SI—Sandstone and Ironstone; CL—Clay; M—Marl; LS—Limestone.

ison and Bennington they thicken rapidly; so that by inference the main thickening is ENE. and the lines of equal thickness run about 60° E. of N.

Southward from the Red River in Cooke County the Weno and Pawpaw formations thin gradually to at least the Colorado River where they are probably represented by strata lying near the top of the Georgetown limestone. The two formations behave very differently over the area mentioned. The Weno formation, south of the change of direction of its outcrop in Cooke County decreases slightly in thickness, from about 75 feet near Gainesville to 65 feet near Fort Worth, 60 near Riovista, 40 near Waco, and 20(?) near Georgetown. The Pawpaw clay on the other hand, thins more rapidly from the Red River southward. Near Cedar Mills, Cooke County it is 60 feet thick, in Denton County 40 feet, at the north border of Tarrant County 27 feet, on Sycamore Creek near Fort Worth 24 feet, at the south border of Tarrant County 12 feet, at Riovista 5 feet, at Blum 3 feet, and near Waco about 2 feet thick. This thinning is a part of the general thinning southwards of the softer formations of the North Texas Washita, which south of the Brazos are represented in the Georgetown limestone, as is observable in the Kiamitia marl, Duck Creek marl, Denton marl, Pawpaw clay and Grayson marl. The alternate harder formations, the Duck Creek limestone, Fort Worth limestone, Weno limestone, Mainstreet limestone, undergo proportionately less thinning southwards and hence contribute relatively larger components to the Georgetown limestone.

There are equally distinct differences of lithology and facies, which are presented later.

The present paper is only an introduction to the large and interesting Weno and Pawpaw faunae, the majority of whose species are still undescribed. Much collecting and further study is required before any attempt can be made towards a monographic treatment of these fossils. It will be noted that the limits set in this paper coincide with the boundaries of horizons 25 to 33 inclusive, of Adkins and Winton's preliminary section of the North Texas Washita division.¹

¹Adkins and Winton, Paleontological Correlation of the Fredericksburg and Washita Formations in North Texas. Univ. Texas Bull. 1945, 1920.

STRATIGRAPHY

The stratigraphy of the Weno and Pawpaw formations involves the regional stratigraphy of the Comanchean area north of the Brazos River, and accordingly, certain lithological and structural features of this region will be described.

In almost the whole region in question, along the line of outcrop of these two formations between the Brazos and the Red Rivers, the Comanchean sediments are deposited over a large geosyncline whose axis is transverse to the direction of the Comanchean outcrops—i. e., roughly northwest-southeast—and whose bulk was largely filled in by pre-Comanchean deposits. However, there is a feeble surface reflection of this buried structure, which influences the thickness and lithologic characters of all Comanchean deposits, including the Weno and Pawpaw formations. The north edge of this deep trough is near the Red River and the south edge is south of the Brazos; the north slope is steep and the south slope more gentle. This depression, in which the Fort Worth region is located will be called the Fort Worth geosyncline.

ROCKS UNDERLYING THE WENO AND PAWPAW FORMATIONS

PRE-COMANCHEAN

Its extent may be indicated by the meager data available on the underground position of the Ellenburger limestone. At Muenster, Cooke County, it was reached at a depth of about 1800 feet, or 1050 feet below sea level. At Myra, Cooke County, the Ellenburger was reached at a depth of 1640 feet below the surface or about -900 feet (sea level).¹ In the Polytechnic (Byrens-Burchell No. 1) well near Fort Worth, the Ellenburger if present lies below -3950 feet, at which level the Bend has probably not been reached. The difference in elevations of the last two subsurface points is 3050 feet which represents the minimum change of elevation of the first Ellenburger reached between Myra and Fort Worth, in case the Ellenburger is present at the latter place. Coincidentally the thickness of Pennsylvanian deposits is not more than 600 feet at Myra, but has increased to at least 3600 feet at Fort Worth. Other wells in Tarrant and nearby counties have penetrated similar Pennsylvanian material.

Farther south there is doubt that the Ellenburger is present. In the Hillshoro well the pre-Cambrian rocks were said to be reached at a depth

¹Matteson, Econ. Geol., XIV, No. 2, p. 1919.

of 2100 feet, apparently without intervening Ellenburger, and at Waco the pre-Cambrian is present at about 2500 feet. However, at points still farther south and southwest, as at Gatesville and Leander, the Ellenburger is present but thin, and overlies pre-Cambrian shales or slate-graywacke which near Leander are stated to be identical in appearance with the Virginia shales and to be of upper Huronian age.¹ At Georgetown, Leon Springs and Camp Bullis,² the Trinity directly overlies the pre-Cambrian schists;³ the depths of the contact are respectively 1100 feet, 1015 feet, and 1790 feet.

The depression which is triangular shaped and narrower to the northwest, is flanked to the north, at least in Cooke County, by a structurally high area of Ellenburger at -900 feet, and a thin Pennsylvanian; and to the west by the structurally high Bend Arch in which the -3000 and -3500 Ellenburger contours have a trend east and south.⁴ Whether this Ellenburger depression opens to the southeast or to the northwest is not known to me. It was, however, invaded by the Pennsylvanian seas which deposited in its central part at least 3600 feet of sediments, and left a surface which was more largely levelled than the original floor.

Upon this surface the Comanchean sea transgressed from the southeast, depositing sediments which still show a feeble reflection of the underlying geosyncline.

COMANCHEAN

THICKNESS CHANGES

Any pre-Comanchean ridges and valleys over which the Comanchean seas spread would be indicated by deposition which is thicker in the valleys and thinner over the crests. If therefore we identified a regional thinning of the beds in two directions from a given location, this might be attributed to deposition on two sides of an existing valley in the ocean bottom, whether originally this valley was erosional or synclinal. This lensing is evident in the Comanchean formations, and particularly in the lower ones (Glenrose, Paluxy). For example the Glenrose fills in much of the depression by lensing; at Decatur it is about 35 feet thick but at Fort Worth about 475 feet,

¹By Professor A. W. Johnston, oral communication.

²Sellards: *The Geology and Mineral Resources of Bexar County*, Univ. Texas Bull. 1932, pp. 19-20.

³These facts were discovered by Dr. J. A. Udden, who has kindly permitted their use here.

⁴Sellards: *On the Underground Position of the Ellenburger Formation in North Central Texas*, Univ. Texas Bull. 1849, 1920.

and is increasing southeastward at the rate of about 8 feet per mile. (Hill). Likewise the Paluxy entirely disappears to the south. There are in addition two complicating circumstances, which affect the thickness and lithology, general regional thickening and depositional facies. The Fredericksburg division increases steadily in thickness from the Red River to beyond the Rio Grande. Its marl and marly limestone facies persist at most levels to

APPROXIMATE THICKNESS OF THE COMANCHEAN FORMATIONS OF
NORTH TEXAS, IN FEET

FORMATION	Bryan and Marshall Counties, Oklahoma	Grayson County, Texas	Cooke County	Denton County	Tarrant County	Parker County	Johnson County	Hill County
Woodbine	200†	500	400†	400—	325		100-300	100†
Buda	...	...	...	...	...			
Grayson	50	50†	60	82	80		100	
Mainstreet	5	10-15	40	50	50		50	35†
Pawpaw	56	50	45	27-40	12-27		9-18	5
Weno	114	125	110	70-100	62		40	25
Denton	25	25	25	25	25		20-25	5
Fort Worth	30	30	30	30	32	30	25-27	25
Duck Creek marl...	40	60	30†	30†	22	20	20	50
Duck Creek limestone	100	130	50†	50†	45	40	40	...
Kiamitia	130	61	40	30-40	27-31	30	18	19
Goodland	16	12-20	19		117			
Edwards						20	35	40
Comanche Peak						120	105	150
Walnut	8	(5)	5-15		125-150	150	100	150
Paluxy					150	150	90	150
Glenrose				?	470	100-300	400-500	
Basement Sand	250	250	200†	200?	150†	150†	150	
WASHITA (e x c l u - sive Woodbine ..	550†	541†	430	394	374	120	338	164†
FREDERICKS- BURG	25?	50	25?		242	290	330	350
TRINITY			?	200?	770†		650	
COMANCHEAN...	825	841	655		1386			

†Approximate thickness.

the region between the Brazos and the Colorado Rivers. South and west of this region the Rudistid facies invades the Fredericksburg division, beginning with the level representing the top of the Edwards limestone (see page 24). The Rudistid facies is characterized by considerable thickness of indurated massive limestone, so that southward as this facies vertically invades the stratigraphic column the invaded formations thicken. Rough estimates of the thickness of this division are: Red River (Denison, 50 feet; Goodland, Oklahoma, 25 feet); Trinity River (Fort Worth), 242 feet; Colorado River (Austin), 350 feet. And for the Washita division: Denison, 541 feet; Fort Worth, 374 feet; Austin, 220 feet.

In the following table of approximate thicknesses it should be noted that the basal Comanchean deposits are transgressive northwards in such a manner that we do not know with certainty what part if any of the Basement Sands of Cooke, Grayson and Denton Counties is of Trinity age. The lowest determined Comanchean fossils in this area indicate the level of *Exogyra texana* and its varieties of Walnut age. In the Duck Creek formation of the table the "marl" is taken to be the portion above the Mineralized Ledge¹ (Stratum 17) of the Fort Worth region, i. e., above the horizon of abundance of *Scaphites worthensis*.

LITHOLOGICAL CHANGES

The lithology of the Washita division is diverse and each formation must be considered separately. However, it is generally true in North-Central Texas that the formations decrease in thickness southwards to the Brazos-Colorado divide south of Waco, Texas; these changes in thickness and lithology may be considered as a unit, and are related to the depositional conditions in the large North-Central Texas trough already referred to. Crossing the Brazos-Colorado Divide in which the thinned Pennsylvanian and Ellenburger and pre-Cambrian floor are structurally high, there is in the Cretaceous another set of thickness changes which for certain formations, notably the Buda, amount to a regional increment in thickness as far south as the turning point of the Washita outcrops in Bexar County. Westward from this point there appears a little known set of changes in thickness and lithology of these formations. In West Texas again, the northern, near-shore facies (Cerro de Muleros, Finlay Mountains, Kent, Sierra Blanca) passes quickly into the southern, massive limestone, in part rudistid facies (Shafter, Terlingua, Fort Stockton, Sheffield) and the Texas and Pacific Railway is approximately the boundary line between these two facies. The transition zone from the Edwards

¹Winton and Adkins: The Geology of Tarrant County, Univ. Texas Bull. 1931, pp. 42-43.

limestone level passes between Fort Stockton and Sheffield and eastward to a point near Pecos, thence some distance north of Coke and Runnels counties, north of the Central Mineral Region, and approximately to the northern border of Hood County, near Comanche Peak. Here the Edwards outcrop disappears eastward under the later formations. An extended study of the rudistid facies will be necessary to define this boundary line precisely; but it is cited here as an illustration of the type of lithological variations seen within the extent of the Texas Comanchean outcrops. For the other Washita formations likewise, much field work will be necessary to map even approximately their different lithological facies; accordingly the following summary of these formations is brief and condensed.

GRAYSON FORMATION

Sandstone facies:

Unknown.

Clay facies:

Upper half of the Del Rio Clay (approximately the *Gryphea mucronata* zone, south of the Brazos River). The middle member of the Grayson is a clay as far north as Denton County, Texas.

Marl facies:

Near Bennington, Durant and Bokchito, Oklahoma, Denison, Fink, Gainesville, Denton, Fort Worth, and Burleson, Texas. This is the prevailing facies along the outcrop. The easternmost exposure in Central Texas is east of Roanoke, Texas, where the few, insignificant limy seams seen in the Fort Worth region are represented by eleven conspicuous limestone strata. There is little doubt that this formation grows rapidly more calcareous eastward and southeastward, and that its limestone facies lies in this direction from the outcrop in North-Central Texas and not far from the outcrop. Passing down the Red River valley, however, this does not hold, for the formation is marly in the Tishomingo quadrangle and near Bennington, Oklahoma, where it has essentially the same lithology and fossils as in Tarrant County, Texas.

WEST TEXAS

In the Cerro de Muleros section, Böse's subdivision 8¹ consisting of yellow marls containing *Schloenbachia* sp., *Pecten subalpinus* (Böse), *Exogyra whitneyi* Böse,² *Hemiaster calvini* Clark, *Protocardia texana* (Conrad), and *Enallaster bravoensis* Böse, is Grayson. It overlies the Mainstreet formation, here a sandstone, and underlies subdivision 9, a

¹Böse: Inst. Geol. Mex. Bol. 25, p. 27-28.

²Böse: Univ. Texas Bull. 1902, p. 10.

limestone which may be either Grayson or Buda, probably the latter; this underlies the Woodbine (Dakota) sandstone. The Kent section contains equivalents of Weno and Pawpaw in division 3 of Dumble and Cummins¹ section, which however does not permit of more precise definition.

A north-south line drawn two miles east of Roanoke, Denton County, therefore delimits the marl facies of the Grayson from the transitional beds to the limestone facies, since east of this line the Grayson becomes calcareous. This line however probably turns east since on going down the Red River valley it leaves the marl facies of the Grayson to the north as seen at Denison, Bennington and numerous other places in Southern Oklahoma. What relations exist east and south of these points depends on well records not yet investigated. South of the Brazos River near Waco there is at the Grayson outcrop a transitional zone to the southern clay facies (zone of *Gryphea mucronata*, upper half of the Del Rio Clay) which continues westward to the El Paso region. The sand facies is unknown but is to be expected in the removed areas of North Central Texas, in the isolated Upper Washita remnants of the Panhandle not yet investigated, and in the Tucumcari region.

Limestone facies:

The limestone facies, on the other hand, lying southwards in Sonora and Chihuahua is still unidentified in Texas, except at one point, the Mariscal Mountains, at the south tip of the Big Bend region.²

MAINSTREET

Sand facies:

At Cerro de Muleros, near El Paso, Texas; apparently near Santa Rosa, New Mexico (Dr. Böse).

Clay facies (Del Rio Clay, in part): The Del Rio Clay is poorly known, and therefore its correlation is tentative. It is equivalent to the Grayson and the upper part of the Mainstreet, and contains the following fossil levels:

(1) *Nodosaria texana* Conrad. A zone of abundance above that of *Exogyra cartledgei* Böse³ in the upper 30 feet of the formation, which at Shafter reaches a thickness of 190 feet. Below this zone of abundance *Nodosaria* is scattering.

(2) *Gryphea mucronata* Gabb. Upper half of the Del Rio Clay.

(3) *Exogyra arietina* Roemer. Rare near the middle of the clay in

¹Dumble and Cummins: Amer. Geol., xii, 1893, 309.

²Udden, Baker and Böse: Univ. Texas Bull. 44, 1919, p. 76.

³Böse, Univ. Texas Bull, 1902.

West Texas, abundant lower; in Central Texas (South Bosque, Round Rock, Austin) it is abundant in the basal half of the formation but rare or wanting above. It also occurs in the top of the Georgetown limestone at Austin.

(4) *Turrilites brazoensis* Roemer occurs in the lowest five feet of the Del Rio Clay at South Bosque and at Austin; its main zone of abundance is in the top of the underlying Georgetown limestone.

Zones (1) and (2) are Grayson; zones (3) and (4) are Mainstreet; or briefly the upper part containing *Gryphea mucronata* is Grayson, and the lower part containing *Exogyra arietina* is Mainstreet. The Mainstreet equivalents locally have sandy levels in Central Texas, as for instance at the top of the Georgetown limestone at South Bosque and in the *E. arietina* slabs at Austin. Westward from the turning point of its outcrop in Bexar County the Del Rio becomes sandier and more flaggy and contains sandy slabs with *Nodosaria* and other fossils. At the south end of the Quitman Mountains the Del Rio is transitional, showing a mixture of sand and clay facies, and at Cerro de Muleros the Mainstreet, represented by Böse's subdivision 7 is sandstone. North of the turning point the typical Central Texas section continues to the Brazos, where the Del Rio differentiates into two portions by the intercalation of limestone strata basally; this basal part becomes upper Mainstreet and the upper marlier part becomes Grayson. The lower Mainstreet however continues as a limestone into Central and West Texas.

Marl facies:

The upper Mainstreet formation in McLennan County is transitional from clay to marl, and at South Bosque is rather calcareous. North of the Brazos this calcareous marl contains limy seams which become more prominent and take on the interbedded appearance of the Mainstreet limestone of North Texas.

Limestone facies:

The typically developed Mainstreet limestone between the Brazos and the Red Rivers shows this lithological facies. There is considerable interbedded marl as far north as Gainesville. Eastward from Gainesville, down the Red River Valley, at Cedar Mills, Denison, Bennington, Bokchito and near Hugo, the marl is more reduced in amount, and the formation consists of compact, slightly marly, massive shell breccia or indurated limestone with typical Mainstreet fossils.

PAWPAW FORMATION

The Pawpaw formation shows along its north-south outcrop an almost idealized sequence of marine facies.

Sand facies:

Red River Valley, as far south as the southern border of Cooke County, Texas.

Clay facies:

Denton, Tarrant and Johnson counties, Texas, to near Riovista.

Marl facies:

Johnson and Hill counties, Texas.

Limestone facies:

The equivalents of the Pawpaw south of the Brazos are questionable. If it is represented in this region, it is the upper part of the Georgetown limestone. (See page 11.)

In West Texas its exact equivalent is unknown. At Cerro de Muleros however a distinct Weno and possibly Pawpaw fauna are present in Böse's subdivision 6, but not enough distinctive Pawpaw species are recorded to decide how much of this subdivision represents it. A hiatus at the base and at the top of the Pawpaw has been noted by Stephenson,¹ near the Red River, but it is unlikely that this accounts for any great thickness of Washita sediments.

WENO FORMATION

Sand facies:

There is sand in the upper half of the Weno, in the Red River Valley, Denison, Cedar Mills, Gainesville, Texas. In the lower half of the Weno: unknown. At Tucumcari, New Mexico, Dr. Böse found a sandstone, probably Weno-Pawpaw, containing *Ostrea quadriplicata* Shumard, *Turritella* sp., *Protocardia multistriata* (Shumard), *Protocardia* sp. aff. *texana* (Conrad), *Gryphea dilatata* Marcou² and other fossils.

Clay (shale) facies:

Red River Valley, Gainesville and Denison, Texas. Contains a few thin ironstone seams; southward and to a less extent eastward this facies becomes marly and is intercalated with thin limestone seams.

Marl facies:

Denton, Tarrant, Johnson and Hill counties, Texas. The upper half of the Weno is limy at places (as Fort Worth) where the lower half is marly. The formation is prevailingly of limestone throughout, south

¹Stephenson: U. S. G. S., Prof. Paper 120-H, p. 143.

²Marcou: Geology of North America, plate IV, figures 1, 1a, 3, 1842.

of the Brazos River. In West Texas at Cerro de Muleros, the Weno is represented in Böse's subdivision 6, which consists of marls, sandstone and limestone. This series is therefore composite and transitional, like the Weno at Gainesville and Denison, and contains many identical fossils.

Limestone facies:

Portion of the Georgetown limestone near top, south of the Brazos. An indurated, massive limestone facies is unknown but probably lies to the east and south of the outcrop in Central Texas.

DENTON FORMATION

Sand facies:

Unknown.

Clay facies:

From Blue Mound, near Haslet, Tarrant County, Texas, northward to Denison. Contains pyrite and crustacean fauna.

Marl facies:

In North-Central Texas along the outcrop the marl facies appears below Blue Mound, just north of Fort Worth and continues southward. In southern Oklahoma, the Red River Valley and Central Texas north of the Brazos, this formation is a clay, slightly arenaceous at the base and capped by shell conglomerate. This conglomerate is much more conspicuous at the Red River than at Fort Worth and southward, where it has largely disappeared, leaving a very shelly marl. South of the Brazos River, near Waco, there is a transitional zone between the marl and the limestone facies.

At Cerro de Muleros a portion of subdivision 5 is a gray marl containing great numbers of *Gryphea washitaensis* Hill, and represents the Denton.

Limestone facies:

The portion of the Georgetown limestone which represents the Denton formation is a consolidated shelly limestone with very little calcareous cementing material, and usually massive without marl interbedding. It extends from a point near Waco southwards to the turning point of the Comanchean outcrops in Bexar County and thence westward to beyond the Pecos River. In the El Paso region, however, the equivalent to the Denton belongs to the northern (littoral-bathyal) facies and is marly as in North Texas.

FORT WORTH FORMATION

This formation is calcareous over all of North and Central Texas but is somewhat marly in the Red River valley and is prevailingly marly near El

Paso. A line connecting these two areas will give very roughly the boundary between the marl and lime areas.

Sand facies:

Unknown.

Clay facies:

Unknown.

Marl facies:

Cerro de Muleros, near El Paso, gray to bluish marls; Denison and eastwards, marly limestone and marly interbedding.

Limestone facies:

South of the Red River and throughout North and Central Texas at the outcrop. The middle portion of the Georgetown limestone represents this facies. A deeper sea subphase found in southern Trans-Pecos Texas consists of hard, crystalline, sparsely fossiliferous, relatively pure, fine grained lithographic limestone, which composes most of the Georgetown and Buda. This is the purest and probably the deepest sea deposition known for this formation.

DUCK CREEK (Upper)

Sand facies:

Unknown.

Marl facies:

This group of strata is prevailingly a calcareous marl and contains the least limestone at Gainesville. At Fort Worth it has only slightly more lime than at Gainesville, while at Denison and at Caddo, Oklahoma, it is distinctly more limy. The change from marliness to liminess therefore appears to be in this region at least largely an east-west change. The Duck Creek marl thickens to the northwest, being thickest near Denison and thicker in the Tishomingo than in the Atoka area.

Limestone facies:

South of the Brazos-Colorado River divide, the Duck Creek marl is imbedded in the base of the Georgetown limestone, where it is a marly impure limestone.

The situation in West Texas is unknown.

DUCK CREEK (Lower)

Sand facies:

Unknown. There are some bituminous sandy layers in the Duck Creek in Cooke County, Texas, but these appear to be local.

Shale (clay) facies:

Seen at the type locality, three miles north of Denison, Texas. This shale is somewhat calcareous and is transitional to a marl.

Marl facies:

This facies invades the limestone series at Gainesville, Texas, where all of the lower Duck Creek except the basal ten feet is prevailingly marly. Probably the removed portion of the Duck Creek west and north of the outcrop was prevailingly marly.

Limestone facies:

Extends from Denton County southward to the turning point of the Comanchean outcrop in Bexar County, and thence westward to the Trans-Pecos region. The group of strata is composite and at a given point is more marly above than below. For instance, at Denison the Hamites and Desmoceras zones are prevailingly limy marl with some blue shale, and at Denison the zone of Desmoceras brazoense is marly at the top; at Fort Worth the section is limestone to the top of the Schloenbachia trinodosa horizon and limy even above this; while below the Brazos the whole lower Duck Creek section becomes limy.

South of the Brazos the Duck Creek formation lies at the base of the Georgetown limestone where it is a chalky white indurated rather unfossiliferous limestone, differing considerably in appearance at Austin and at Georgetown, Texas; while west of the southern turning point of the outcrop it becomes a hard, crystalline, consolidated, sparsely fossiliferous limestone. At Cerro de Muleros near El Paso, on the contrary, the northern littoral facies is present as at Gainesville and Denison and this level is represented by a series of clay, marl and limestone, which represents a transitional facies. Vraconian faunae from Chihuahua and Zaca-tecas (p. 60) represent in part this stratigraphic level.

KIAMITIA FORMATION**Sand facies:**

Unknown.

Clay (shale) facies:

From near Primrose, southwestern Tarrant County, south through Johnson County to beyond the Brazos. This formation at Blum, Hill County, is a yellowish clay 19 feet thick. This facies contains a conspicuous limonite fauna, Schloenbachia, Arca, small gastropods and pelecypods and other fossils.

Marl facies:

Southern Tarrant County to between Gainesville and Fink. The formation is increasing in thickness northwards and as the clay facies disappears, flag layers enter first as limestone in thin sheets and farther north as shell conglomerate sheets. The bulk of the formation is marl, and the limonite fauna has largely disappeared.

Lime facies (conglomerate):

Fink, Grayson County, Texas, to east of Hugo, Oklahoma. Coincidentally on passing eastwards and northwards from the marl facies the scattered shells in the formation become consolidated into conglomeratic sheets with shelly marl between, and these predominate at Gainesville and Denison. Turning east down the Red River Valley the thickness of the Kiamitia and the amount of this conglomerate greatly increase. Near Hugo, Oklahoma, the Kiamitia shell conglomerate is about 150 feet thick and is extensively used for crushed rock. This formation is a shallow water oyster bed deposit of mixed Fredericksburg and Washita fossils, mainly the former.

The formation disappears at the Brazos-Colorado uplift, and in West Texas its relations are unknown.

EDWARDS FORMATION

The Fredericksburg Division also must be considered sectionally, since its various parts behave differently as to their changes of thickness and lithology.

Sand facies:

Unknown.

Clay facies:

Unknown.

Marl facies:

Unknown. Böse's subdivision 3 at Cerro de Muleros contains some marl, but this subdivision is not positively known to be Edwards. Throughout the Red River region and North-Central Texas, the top of the Goodland, which is thought to correspond to the Edwards, is a non-marly limestone, usually massive and sometimes crystalline. North of Sheffield and Fort Stockton, the Fredericksburg is stated to be prevailingly marly.

Limestone facies:

This is known from Fort Worth south to the turning point of the outcrop in Bexar County, and thence west to El Paso. The Red River extension of this level is not identifiable with certainty because it is not clear whether the 12-20 feet of white limestone at the top of the Fredericksburg division as seen at Denison or north of Goodland, Oklahoma, represents only the Edwards or also in part still lower Fredericksburg. The Edwards is less than 10 feet thick at Fort Worth; it is 33 feet thick at Comanche Peak, and southwards thickens rapidly.

Rudistid facies:

The Rudistids invaded Texas most widely in upper Fredericksburg time, when they were scattered as far north as Fort Worth; however, they are very rare and inconspicuous north of the Brazos, in Central Texas. In

West Texas they are reported as far north as Bailey County, and they are common in Coke and Runnels counties. In Trans-Pecos Texas, the Texas and Pacific Railway roughly divides the southern rudistid facies of the Edwards from the northern littoral facies. The rudistids are rare or wanting at Cerro de Muleros, Kent, Sierra Blanca and the Finlay Mountains; and present at points between these and the Rio Grande. Gabb¹ describes from the Sierra de las Conchas near Arivechi, Sonora, fossils which if correctly identified include among diverse stratigraphic levels that of the Edwards limestone; the facies represented is unknown to me. Felix and Lenk² also record the great extent of the Fredericksburg reef facies in Mexico.

WENO AND PAWPAW FORMATIONS

The influence of the underlying syncline on the deposition of these two upper Washita formations is small, due to their small thickness. However, they show an appreciable syncline in the Fort Worth region, and a conspicuous thinning southward.

Since the maximum rate of change of thickness, like the maximum change in lithology, may not coincide in direction with the outcrop, the formation along the outcrop will in general show only a greater or less north-south or an east-west component of change in thickness or lithology and the maximum change will be in a direction lying at an angle to the outcrop.

THICKNESS CHANGES

North-South Changes

This is true of the north-south outcrop of the two formations, which does not exactly coincide with the direction of greatest thickness change. The Weno and Pawpaw strata outcrop in a line passing about a mile east of Gainesville, Cooke County, and thence nearly south to near Denton; thereafter the boundary between the two formations runs near Fort Worth, west of Cleburne, Rivovista and Blum.

The outcrop thus from the Red River to the Brazos has a general trend of east of north, while the direction of greatest thickness change is slightly east of north. (See figure 3).

¹Gabb: *Pal. Cal.*, vol. 2, p. 257 ff.

²Felix and Lenk: *Beitr. z. Geol. u. Pal. d. Rep. Mex.*, II, p. 28.

East-West Changes (North Texas and Oklahoma)

In the whole Red River region these two formations at their outcrops maintain an almost uniform thickness, but they are slightly thicker eastward as far as Choctaw County, Oklahoma. Their behavior east of this point is unknown to me.

TABLE OF APPROXIMATE THICKNESS OF WENO AND PAWPAW FORMATIONS

Locality	Weno		Total	Paw-paw
	Lower	Upper		
Denison	45	80	125	50
Gainesville	40	70	110	44.8
Blue Mound (Haslet)				27.3
Fort Worth	12.7	49.6	62.3	24.6
Riovista	10	25	35	5.0

LITHOLOGICAL CHANGES

The Weno and Pawpaw formations are marked by striking lithological changes which produce along their outcrops localized lithological regions each with a characteristic fauna. The Pawpaw formation, and to a less extent the Weno, passes from north to south along its outcrop through a "typical" series of lithological facies—sand-clay-marl-limestone, which aside from various complicating factors of deposition is usually taken to represent a progressive series of marine facies from near-shore to off-shore conditions. Likewise the problem of localized faunales is vividly impressed upon one by the situation in the Pawpaw clay (as also in the Weno), where within a few miles one fauna largely disappears and an equally rich and varied, but different one occupies its stratigraphic position. For example the Gainesville-Denison fauna (Nacreous, Arca), the Fort Worth fauna (Turrilites, Engonoceras, Hamites, Scaphites) and the Riovista fauna (Flickia, echinoids) of the Pawpaw formation represent three different marine facies. The narrow ribbon-like outcrop of these formations in North Central Texas and Southern Oklahoma gives only a limited opportunity for the study of these different marine phases, but the following regional differences in the sediments of these formations are apparent.

PAWPAW FORMATION

Nature of Material	Locality of Outcrop	Facies of Marine Deposition
(a) Semi-consolidated sandstone and ironstone	{ Choctaw county, Oklahoma Marshall county, Oklahoma Bryan county, Oklahoma Love county, Oklahoma Grayson county, Texas Cooke county, Texas	Neritic
(b) Clay with sandy layers.....	{ Denton county Tarrant county	Bathyal
(c) Marl with limy layers.....	{ Johnson county Hill county	
(d) Limestone.....	South of Brazos River	Deeper Sea

UPPER WENO

(a) Shale with sand layers and iron- stone	{ Grayson county Cooke county	Neritic
(b) Marl with lime layers.....	Denton county	Bathyal
(c) Limestone.....	Tarrant county and southward	Deeper Sea (in part Zoogenic)

LOWER WENO

(a) Shale with sand layers.....	Red River region	Neritic
(b) Marl and limestone layers.....	Cooke county to Brazos River	Bathyal
(c) Limestone.....	South of Brazos River	Deeper Sea

PAWPAW FORMATION

Eastward from the turning point of the Comanchean outcrops in the Red River uplifted area near Orlena, Cooke County, Texas, the Pawpaw is a ferruginous sandstone or an unconsolidated sand. This outcrop runs down the Red River valley in a direction a little south of east, lying in Cooke and Grayson Counties, Texas, and in Love, Marshall, Bryan, and Choctaw Counties, Oklahoma.

Sandstone Facies

(a) (Semi-consolidated Sandstone and Ironstone)

In general, the Pawpaw formation is prevailingly sandy and iron bearing at its outcrops north and east of the south border of Cooke County, Texas.

Near Gainesville, the formation is a series of thickbedded, hard, ferruginous sandstones with interbedded, laminated impregnated sandy layers and semi-consolidated iron-impregnated sand. The layers are locally a shell conglomerate containing great quantities of casts of *Arca*, *Turritella* and other bivalves and gastropods. The basal layers are characterized by an abundant pyrite fauna. *Turritiles*, *Scaphites*, *Hamites* and echinoids are rare or absent. This type of lithology continues around the turning point of the Comanchean formations south of Orlena to a point near Pottsboro; and at Denison the Pawpaw formation is almost entirely a coarse consolidated brownish-red sand with few fossils. At Durant the formation is similar to the Denison exposures. Eastward through Bennington and Bokchito the sandy phase continues, and is included in the Bokchito formation of Taff.

SECTION OF SUGAR LOAF MOUNTAIN, BRYAN COUNTY, OKLAHOMA
Section 22, R. 12 E., T. 5 S. (furnished by W. M. Winton)

MAINSTREET:

Limestone, iron stained, old cap of hill, practically removed by weathering.

Feet

PAWPAW:

Massive yellow red sandstone, soft in fresh exposures, indurated where long exposed; no fossils seen.....	5.0
Soft limonitic sand; no fossils seen.....	30.0
Ironstone ledge composed of fossils (<i>Arca</i> sp., <i>Ostrea quadriplicata</i> , small ammonites, gastropods, etc., a typical Pawpaw fauna).....	0.5
Red sand, cross bedded and containing lenticular hard masses of sandstone	10.0
Ironstone ledge composed of fossils (<i>Arca</i> sp., <i>Ostrea quadriplicata</i> , ammonites, <i>Nodosaria</i> , etc., a typical Pawpaw fauna).....	0.3
Red sand	10.0

WENO:

Yellowish hard limestone with typical Weno fossils.....	0.8
Soft marl; no fossils seen.....	0.3
Very hard massive limestone, pinkish in color. Large numbers of fossils having same hardness as matrix and giving uniform fracture. Ledge forms conspicuous topographic break traceable for miles. <i>Schloenbachia</i> sp. M., and other typical Weno fossils.....	1.9
Soft marl	15.0
Limestone ledge sandy; no fossils seen.....	1.0
Soft marl with a few thin limestone ledges, each less than 0.5 feet thick....	95.0

DENTON:

Conglomerate of <i>Gryphea washitaensis</i> , <i>Ostrea carinata</i> , and other Denton fossils. Typical Denton marl ("Caddo" limestone of Taff, top). Exposed, about	2.0
---	-----

In the creek, one-fourth mile west of the mountain, the contact between the Duck Creek marl and the Fort Worth limestone is exposed as at Caddo, Oklahoma.

In the second cut $1\frac{1}{2}$ miles north of Hugo, Oklahoma, there is 10 inches of shell conglomerate containing abundant juvenile and adult *Ostrea quadriplicata* Shumard, *Gryphea washitaensis* Hill, *Plicatula* sp., *Pecten subalpinus* (Böse), abundant *Leiocidaris* spines, *Corbula littoralis* Adkins, and other typical upper Weno fossils as found at Gainesville and Denison. This represents part of the Quarry limestone group. Above it is an iron-stone ledge, 4 inches thick, which is red-stained and conglomeratic and resembles the basal Pawpaw as seen at Gainesville.

Along the St. Louis and San Francisco Railway track between Bennington and Bokchito, Oklahoma, the upper Washita section, Woodbine to Weno, is seen to advantage. The Grayson marl with an estimated thickness of 50 feet is well developed with its characteristic fossils.¹

The Mainstreet limestone is seen in roadside cuts one mile northwest of Bennington, in the caprock of Sugarloaf Mountain, six miles north of Bennington, in the bed of Sulphur Creek, where it forms an extensive pavement limestone, and in the railroad cuts nearby, particularly in the long cut running west down to Bokchito Creek, which has it well exposed. Everywhere it is underlain by the brown consolidated Pawpaw sandstone. The nature of the Pawpaw and Weno outcrops in this region is seen from the following section.

SECTION OF CUT NEAR SULPHUR CREEK, ON THE ST. LOUIS AND SAN FRANCISCO RAILWAY, TWO MILES WEST OF BENNINGTON, OKLAHOMA

MAINSTREET:	Feet
Brown limestone with <i>Turrilites brazoensis</i> , <i>Pecten subalpinus</i> , <i>Exogyra</i> sp. (large), and ammonites. The top of the exposure is an <i>Exogyra arietina</i> conglomerate. Exposed	2
Brown-blue limestone with irregular sandy inclusions.....	1
Yellow limestone containing <i>Kingena wacoensis</i> , <i>Ostrea marcoui</i> , and <i>Pecten subalpinus</i>	1.5

¹Taff (U. S. G. S. Geologic Atlas of the United States, Atoka Folio, No. 79) has overlooked the Grayson marl, which in this quadrangle is typically developed as at Denison. It may be seen near the Woodbine ("Silo") sandstone knob mapped by Taff one mile northwest of Bennington, Oklahoma, where it overlies typical Mainstreet limestone containing *Turrilites brazoensis* Roemer, *Exogyra arietina* Roemer and other distinctive fossils, and contains *Gryphea mucronata* Gabb, *Gryphea*, sp. (truncate) *Pecten subalpinus* (Böse), *Plicatula* sp., *Ostrea* sp. aff. *subovata* Shumard, and *Engonoceras* sp. This locality is mentioned by Taff, *ibid.*, p. 6. It also forms the overburden in the Mainstreet quarry one mile north of Durant, Oklahoma, where it is very fossiliferous.

PAWPAW:

Consolidated brown sandstone, no fossils seen.....	5
Unconsolidated sand, iron stained	10
Blue jointed clay with scattered Gryphea.....	10
Red-brown sandstone—calcareous in part—with nacreous and ironstone fossils: Arca, Nucula sp., Corbula sp., Cerithium, Turritella, Anchura, like the Gainesville and Denison nacreous fauna.....	1
Marl, containing ironstone seams, no fossils seen.....	10

The cut east of the Bokchito Creek bridge shows the foregoing section much better exposed. Pawpaw exposures were also noted near Durant, Oklahoma. North of the railroad, 1½ miles east of Bennington, Oklahoma, about 50 feet of massive brown sandstone was noted underlying the Main-street limestone at the top of the hill. This sandstone contains limy layers bearing *Ostrea quadriplicata*, *O. subovata* (?), *marcoui*, *Trigonia* sp., *Gryphea* sp., and *Pecten* sp. (large). Taff gives the following section for his Bokchito formation in the Atoka and Tishomingo quadrangles¹:

SECTION OF THE BOKCHITO FORMATION IN THE ATOKA AND TISHOMINGO (OKLAHOMA) QUADRANGLES

J. A. Taff, 1902-1903

	Feet
4. Sandy and clay shales, and locally friable, cross bedded sandstone. In the clay and in some of the iron and lime concretions, shells are preserved with original nacre	50
3. Hard, semi-crystalline, bluish, oyster shell limestone, weathering yellowish and projecting as ledges separated by friable shales.....	10-20
2. Friable sandstone beds, locally cross bedded, alternating with and including deposits of sandy clay	20-30
1. Sandy clay shale, with ferruginous limestone segregations and ironstone nodules.	90
	170-190

This thickness is given as 140 feet in the texts and the Columnar Section sheets of both folios.

The upper member (4) of this section is probably Pawpaw. Sheets of nacreous shells in the Pawpaw have been noted in Grayson County. The other three members are Weno.

The Pawpaw formation is exposed in the hilltops east of Denison and

¹Taff: U. S. G. S., Geologic Atlas of the United States, Atoka Folio No. 79, 1902, p. 6; and, Tishomingo Folio, No. 98, 1903, p. 6.

may be seen in cuts on East Main Street, where it is a cross-bedded rather consolidated, brick red sandstone. No fossils were seen at this locality. The formation is stated by Stephenson¹ to be 50 or more feet thick near Denison. The Pawpaw is poorly exposed north of a west branch of Little Mineral Creek just south of Fink, where it is steeply tilted. On the Red River about 2 miles northwest of Cedar Mills, Grayson County, there is a long cliff dipping west (5°) which exposes near the top about 40 feet of loose sand only partially consolidated, which is probably Pawpaw.

SECTION OF BLUFF OF RED RIVER TWO MILES NORTHWEST OF CEDAR MILLS, GRAYSON COUNTY, TEXAS

MAINSTREET:	Feet
White limestone with <i>Ostrea quadriplicata</i> , <i>Turrillites brazoensis</i> , <i>Kingena</i> , <i>Exogyra arietina</i> , <i>Pecten subalpinus</i> , <i>Enallaster</i> sp. aff. <i>texanus</i> . Seen in ravine above east end of river bluff near road.....	10
PAWPAW:	
Sandstone with thin ferruginous layers.....	40
Blue marl with clay and sand seams and nacreous fossils: <i>Engonoceras</i> , <i>Corbula</i> , <i>Nucula</i> , <i>Ostrea quadriplicata</i> , <i>Pecten subalpinus</i>	30
QUARRY LIMESTONE (?):	
Massive grayish semi-crystalline limestone.....	2
WENO:	
Blue marl with nacreous fossils.....	40

The roadside cuts 2½ miles east of Gainesville, Cooke County, inconsecutive sections of the Pawpaw are seen. The interval from the top of the Quarry group to the base of the Mainstreet limestone is not less than 45 feet. The base of the Pawpaw overlying the Quarry limestone is seen in a cut of the Missouri, Kansas and Texas railway (Wichita Falls branch) at the north end of the brickyards pit 1¾ miles southeast of Gainesville. This exposes about 8 inches of an irregularly deposited reddish ferruginous shell conglomerate largely of *Ostrea quadriplicata*—cemented into the pitted surface of the Quarry limestone. The same level is exposed near the water tank on the southwest rim of the pit. In roadside cuts ¾ mile southeast of this locality about 45 feet of Pawpaw may be seen poorly exposed; it consists of thin layers of ferruginous dark red brown sandstone which, locally at least, consists of cemented masses of ironstone fossil casts: *Arca*, *Protocardia*, *Nucula*, *Corbula*, *Turritella* and rare *Ostrea quadriplicata* and *Gryphea*. These ironstone layers are interbedded with brown to

¹Stephenson, U. S. G. S., Prof. Paper 120-H, p. 142.

yellow, softer, ferruginous, very arenaceous clay. Their ironstone seams do not exceed 6 inches in thickness and locally are highly indurated.

SECTION OF THE PAWPAW FORMATION IN ROADSIDE CUTS TWO AND ONE-HALF MILES SOUTHEAST OF GAINESVILLE, TEXAS

Section furnished by W. M. Winton

MAINSTREET:

	Feet
White massive limestone with characteristic Mainstreet fossils. Exposed...	6

PAWPAW:

Alternating red ironstone and ferruginous sandy clay layers. About 16 compact ironstone layers each 3 to 4 inches thick, alternating with clay layers each about one foot thick. The ironstone layers are similar from bottom to top and the basal 10 feet is more fossiliferous than the upper portion. The ironstone layers contain: Remondia sp., Trigonina sp., Arca sp., Engonoceras sp., Nucula sp., Corbula sp., and other nacreous and ironstone fossils	21.5
Brown sandstone flag layer	0.5
Red ironstone layer with <i>Nodosaria texana</i>	0.3
Brown clay	5.0
Red ironstone with <i>Arca</i> sp., <i>Ostrea quadriplicata</i>	0.5
Brown clay, sparsely fossiliferous: <i>Arca</i> , gastropods.....	15.0
Red calcareous sandstone: <i>Ostrea quadriplicata</i>	2.0

This is underlain by the Quarry Limestone group (see page 36). The thick semi-consolidated sand seen in Grayson County is absent. The Pawpaw outcrop which has turned nearly south at the turning point near Orlena, makes a narrow north-south strip from Gainesville to the southern border of Tarrant County. The ironstone facies does not persist past the Trinity River, since interstratified clay lenses here largely make up the formation; as far south as Fort Worth there are large amounts of weathered ironstone fragments in the surface debris of the Pawpaw.¹

Clay Facies

This facies of the Pawpaw formation is best developed in the Fort Worth region. From Cooke County southward to the Brazos the Pawpaw is rapidly thinning, as previously described. At the same time, passing southward through Denton and Tarrant Counties the clay facies is encountered, and in the region of the Brazos the marl facies appears. The clay facies grades into each of the other two facies, so that the transitions are gradual. This gradation occurs through the invasion of one facies by seams of the other kind of material.

¹Winton and Adkins: Univ. Texas Bull. 1931, p. 67.

The transitional zone from the ironstone to the clay facies occurs in Denton and northern Tarrant Counties. As far south as the Trinity River considerable amounts of ironstone fragments occur in the Pawpaw exposures; these are the residue of numerous thin ironstone ledges similar to those of the Red River area but sparsely fossiliferous, which are scattered rather evenly throughout the formation. The interbedding material is an arenaceous clay. South of the Trinity the section is prevailingly clay but has thin ironstone, limestone and sand seams.

SECTION OF THE PAWPAW FORMATION AT BLUE MOUND ONE AND ONE-HALF MILES SOUTH OF HASLET, TARRANT COUNTY, TEXAS

MAINSTREET:	Feet
White chalky limestone, exposed	4
PAWPAW:	
Brown to yellow, sparsely fossiliferous clay, with thin ironstone seams and scattered sand lenses	23
Sandstone flag layer	0.3
Brownish yellow impure clay	4
Thickness of Pawpaw	27.3

On Sycamore Creek, three miles southeast of Fort Worth the Pawpaw has an average thickness from several measurements of 7.5 meters (24.6 feet).

At the southern border of Tarrant County the Pawpaw is 12 feet thick and is transitional to the marl facies. In northern Johnson County the transitional zone continues and the section consists of clay mixed with marl. Passing from the upper Noland's River southeast to Riovista the section changes considerably, for from Riovista southward the marl facies is found, with little clay and no ironstone; there is also a corresponding change in the Pawpaw fauna.

SECTION OF BRANCH OF NOLAND'S RIVER NEAR CLEBURNE-WEATHER-FORD ROAD, 10 MILES WEST OF CLEBURNE, TEXAS

Section furnished by W. M. Winton

MAINSTREET:	Feet
Massive limestone containing <i>Pachymya</i> sp., <i>Turrilites brazoensis</i> , <i>Holcotypus limitis</i> , <i>Pecten</i> sp.	32.4
PAWPAW:	
Reddish clay containing fragments of <i>Turrilites</i> sp., <i>Arca</i> sp., etc., pyritized	10.8
WENO:	
Limy marl capped by a hard lime ledge one foot thick, containing <i>Pecten georgetownensis</i> , <i>Nautilus texanus</i> , <i>Pinna</i> sp., <i>Kingena</i> sp.	7.8

Marl, containing <i>Nodosaria texana</i>	1.0
Massive limestone containing <i>Schloenbachia</i> sp. M. (notched tubercles), <i>Nautilus texanus</i> , <i>Epiaster</i> sp., <i>Protocardia</i> sp., <i>Gervilliopsis invaginata</i> (one)	7.4
CONCEALED BY BED OF STREAM, ESTIMATED.....	10.0
Soft whitish marl, no fossils seen.....	10.0

Marl Facies

The transition from clay facies to the marl facies of this formation takes place rather abruptly, within a distance of 2 miles along the outcrop and terminates in central Johnson County, south of which the formation is almost entirely marly. One mile southeast of Riovista, the Pawpaw is a distinctly marly uniform light yellow, fossiliferous deposit lying between the conspicuous Weno and Mainstreet limestones. It is about 5 feet thick. In the territory covered by this depositional phase the conspicuous pyrite fauna of the Fort Worth region is rare, except a few *Turrilites*, *Arca* and small gastropods, and other fossils, echinoids and ammonites are abundant.

SECTION ON WACO-RIOVISTA ROAD, ONE MILE SOUTH OF RIOVISTA, TEXAS

MAINSTREET:	Feet
Massive white limestone with characteristic Mainstreet fossils: <i>Turrilites brazoensis</i> , <i>Kingena</i> , <i>Exogyra arietina</i> . Exposed	5
PAWPAW:	
Homogeneous calcareous, straw-yellow marl with a few ironstone and calcareous fragments, containing <i>Enallaster bravoensis</i> , <i>E. wenoensis</i> , <i>E. riovistae</i> , <i>Holaster</i> sp., <i>Epiaster wenoensis</i> , <i>Plicatula</i> sp., * <i>Flickia boesei</i> , * <i>Schloenbachia</i> sp., * <i>Turrilites</i> sp., * <i>Arca washitaensis</i>	5.5
WENO:	
Limestone escarpment in two terraces, poorly exposed at the base, consisting of massive soft limestone with marl interbedding. Contains <i>Enallaster</i> , <i>Epiaster</i> , <i>Holaster</i> , <i>Hemiaster calvini</i> , <i>Pecten subalpinus</i> , <i>Plicatula</i> and many other fossils, <i>Trigonia clavigera</i> . About	40
DENTON:	
Soft light straw-colored marl containing <i>Gryphea washitaensis</i> , <i>Ostrea carinata</i> , <i>Pecten subalpinus</i> , <i>Plicatula</i> sp., and <i>Kingena</i> sp. (large).....	5

LIMESTONE FACIES: The equivalents if any of the Pawpaw formation in Central Texas are largely insignificant since they consist of only a minute thickness of very calcareous marl and limestone imbedded in the top of the Georgetown limestone, underneath the *Turrilites brazoensis* zone which represents the basal Mainstreet formation of North Texas. This situation presumably exists westward to the Trans-Pecos region wherever the

more southward section is found. However, in West Texas, on passing northward into the near-shore facies (as at Kent and Cerro de Muleros) the Weno and Pawpaw formations again thicken, as in passing from Austin to Fort Worth, and show characteristic and so far as can be judged from published accounts, similar fossils. This level at Muleros has echinoids and ammonites and other distinctive fossils (see page 41), but so far the large pyrite fauna of the North Texas Pawpaw has not been reported from West Texas.

WENO FORMATION

A summary of the marine facies of the Weno formation and several geological sections have already been given. The Weno formation is composite, and represents different phases of marine deposition both at different places and in its different levels at the same place. In the Red River region the formation is mostly a series of blue shales with clay-ironstone, limestone and irregular consolidated sand seams. The ironstone and shale layers are rich in nacreous fossils. The shale is here capped by the Quarry limestone group, which loses its massive character south of Denton County. Passing southward from the Red River, the basal Weno transforms from a clay into a calcareous marl with shelly, often slabby, limestone seams and the upper part after passing rapidly through a marl stage becomes pre-vaillingly a soft chalky fossiliferous limestone. This is the situation in Denton and Tarrant Counties. The basal division is also capped by a thin chalky limestone which forms a bench, and the Weno thus consists of two terraces, a persistent topographic feature which passes southwards across Denton, Tarrant, Johnson and Hill Counties and disappears only in north-western McLennan County where the Weno limestone consolidates with the Mainstreet and Fort Worth limestones (by the virtual disappearance of the intervening softer formations) to form the middle Georgetown limestone.

In the Red River region the shale (clay) facies predominates from southern Cooke County to east of Bennington, Oklahoma. It is conspicuous in the first deep cut of the St. Louis and San Francisco Railway track north of Denison, of which a photograph is given by Stephenson,¹ and in the pit of the brickyard $1\frac{3}{4}$ miles southeast of Gainesville, of which a section is here given.

¹Stephenson, U. S. G. S., Prof. Paper 120-H, pl. XXII B.

SECTION OF THE WENO FORMATION AT LOCALITY 601, PIT OF GAINESVILLE BRICK COMPANY, ONE AND THREE-FOURTHS MILES SOUTH-EAST OF GAINESVILLE, TEXAS

PAWPAW:	Feet
Shell limestone, consisting almost entirely of masses of comminuted shells...	0.2
Shelly calcareous marl, with ironstained arenaceous cement and masses of shells. O. QUADRIPLICATA LAYER	.8
This stratum contains: <i>O. quadriplicata</i> , <i>Pecten subalpinus</i> , <i>Plicatula</i> sp., <i>Leiocardis</i> sp. spines.	
Hard reddish limestone composed almost entirely of comminuted shells iron-stained	.3
Shell conglomerate of small packed shells, laminated.....	.5
QUARRY:	
Compact shelly blue to yellow limestone containing many comminuted shell fragments. QUARRY LIMESTONE	.9
This stratum contains: <i>Gryphea</i> sp., <i>Pecten subalpinus</i> , cidarid spines (rare), <i>Plicatula</i> sp.	
Shell conglomerate of <i>Gryphea washitaensis</i> chiefly, with considerable calcareous cement. GRYPHEA CONGLOMERATE	.7
This stratum, which is conspicuous as just underlying the cap rock of the rim of the pit, contains: <i>Ostrea carinata</i> , cidarid spines, <i>Ostrea marcoui</i> , <i>Plicatula</i> sp.	
WENO:	
Packed bluish to yellow clay, with two thin seams of <i>Gryphea</i> shell conglomerate, each one inch thick.....	1.0
Sand layer containing no fossils, irregular thickness, about.....	.2
Blue to yellow shale, with a considerable amount of sand in the form of irregular anastomosing sand lenses and streaks. Fossils in the shale..	4.7
Ironstone band	.1
Blue to yellowish iron stained shale, with irregular calcareous concretions, locally called "ginger shale"	2.5
The ginger concretions also occur in the sandy layer just above the ironstone seam.	
Fossiliferous Ironstone stratum	.5
The clay-ironstone of this and the overlying ironstone layers has been excavated and dumped in a large heap on the east rim of the pit. Material <i>in situ</i> in the pit and that in this heap bears the following nacreous fossils: ‡ <i>Protoecardia</i> sp. aff. <i>multistriata</i> (Shumard), very abundant as casts and molds, with the original shell; ‡ <i>Barbatia</i> sp., ‡ <i>Corbula basiniformis</i> Adkins, ‡ <i>Nucula nokonis</i> Adkins, ‡ <i>Nucula wenoensis</i> Adkins, ‡ <i>Globiconcha</i> sp., ‡ <i>Natica</i> sp., ‡ <i>Cinulia washitaensis</i> Adkins, <i>Cambarus</i> (?) sp., ‡ <i>Schloenbachia wintoni</i> Adkins, ‡ <i>Gervilliopsis invaginata</i> (White), ‡ <i>Pecten inconspicuus</i> Cragin, ‡ <i>Anchura mudgeana</i> (White), and many other fossils.	
Blue fossiliferous shale and calcareous bands.....	4.0
This material is an alternation of thin bands of yellow sandy lime-	

stone with packed laminated blue-gray shale, slate color when fresh, but weathering to blackish-blue.	
Fossiliferous ironstone band	.2
Blue shale, laminated, jointed on weathering, very fossiliferous.....	19.8
This layer contains an abundant nacreous shell fauna, including: <i>Gervilliopsis invaginata</i> (White), <i>Gryphea washitaensis</i> Hill (juvenile stages), <i>Enallaster</i> sp., <i>Tellina</i> sp., <i>Corbula basiniformis</i> Adkins, <i>Nucula wenoensis</i> Adkins, <i>Pecten inconspicuus</i> Cragin, <i>Anchura mudgeana</i> White, <i>Natica</i> sp., <i>Turritella</i> sp., <i>Cerithium</i> sp., etc.	
Gray-blue arenaceous laminated shale, calcareous in part, forming a terrace in the pit. <i>G. washitaensis</i> , <i>Gervilliopsis invaginata</i> , <i>Pecten</i> , <i>Plicatula</i>	.3
Laminated blue shale	.2
Ironstone seam	.1
Laminated gray-blue shale	1.9
This works down into kidney-shaped lumps, and is locally called "kidney shale."	
Arenaceous laminated marl. <i>Gryphea washitaensis</i> abundant at top and bottom. Main zone of <i>Gervilliopsis invaginata</i> (White), <i>Trigonia clavigera</i> Cragin, <i>Pecten subalpinus</i> Böse, <i>Leiocardis</i> spines. This layer forms a secondary terrace	.9
Blue fossiliferous marl. Locally called "buff marl," a desirable brick material, free from impurities, burns to buff color.....	11.2

51.0

This material has been thrown in heaps onto the terrace above and is rich in small calcitic fossils, including: *Venericardia wenoensis* Adkins, *Trochus laticonicus* Adkins, *Trigonia clavigera* Cragin, *Neritina*, *Nerita*, *Pecten* sp., worm tubes, minute corals.

In the bottom of the pit covered by water is about 30 feet more of Weno shale. It is stated that the pit has at one point been dug 70 feet deep. A water well nearby passed through the Denton marl into the Fort Worth limestone. The Denton marl is poorly exposed one-fourth mile west of the pit, and the Fort Worth limestone is finely exposed with its typical sequence of fossil zones, in the creeks between the brickyard and Gainesville.

The Quarry limestone is variable in thickness. Blocks two to three feet thick have been excavated from the pit and thrown in a heap above its south rim. An extensive outcrop, with quarried indurated blocks, three feet thick, occurs three miles east of Gainesville just north of the track. The quarry group forms a persistent outcrop eastward to the river near the the northeast corner of Grayson County, in which county its relations have been described by Stephenson. In Denton County on the Denton-Krum road, it outcrops as a bench, and is 15 inches thick. Here it is abundantly fossiliferous, and contains *Pecten subalpinus*, cidarid spines, *Ostrea marcoui*, *Gryphea washitaensis* and *Ostrea carinata*. Beneath it is a ferruginous and calcareous shell conglomerate layer 1 to 2 inches thick consisting almost entirely of the last two species named. In Tarrant

County it is consolidated with the top of the Weno formation, which is a chalky limestone.

SECTION OF PAWPAW AND WENO FORMATIONS AT LOCALITY 602 ON
SYCAMORE CREEK, FOUR MILES SOUTHEAST OF FORT WORTH, TEXAS

MAINSTREET:	Feet
White massive limestone, with <i>Kingena wacoensis</i> (?) (Roemer), <i>Turrilites brazoensis</i> Roemer, <i>Exogyra arietina</i> Roemer, <i>Pecten subalpinus</i> (Böse), <i>Homomya</i> , <i>Pachymya</i> , <i>Ostrea carinata</i> (?) Lamarck. Exposed	15.0

PAWPAW:

Marl and clay, with thin ironstone seams. The top marly portion contains <i>Enallaster</i> sp., <i>Hemiaster</i> sp., <i>Holaster</i> sp., <i>Epiaster</i> sp., <i>Pecten subalpinus</i> (Böse) and many other calcite fossils.....	18.0
Sandstone flag layer. STARFISH ZONE.....	.3
Ironstained reddish impure clay with many pyrite fossils, including: <i>Engonoceras</i> sp., <i>Turrilites</i> sp., <i>Hamites tenawa</i> Adkins and Winton, <i>Holaster</i> sp., <i>Enallaster</i> sp., <i>Hemiaster</i> sp. aff. <i>bexari</i> Clark, <i>Acanthoceras worthense</i> Adkins, <i>Arca</i> sp., <i>Nucula</i> sp., <i>Mortoniceras worthense</i> Adkins, <i>Schloenbachia</i> sp.	4.0

WENO:

Impure limestone, forms top of first terrace. Marlier beneath, <i>Nodosaria texana</i> Conrad, zone at the base.....	19.7
Limestone at top, forming second terrace, marl beneath. Contains a large fauna, rich in echinoids and pelecypods.....	31.0
Limestone ledge, indurated, fucoidal, <i>Gervilliopsis invaginata</i> (White) zone, <i>Gryphea washitaensis</i> Hill, <i>Pecten subalpinus</i> (Böse), <i>Pinna guadalupae</i> Böse	1.3
There is about 13 feet of fossiliferous Weno marl beneath the creek bed. This is well exposed at the locality 618 (q. v.).	

SECTION OF THE WENO AND PAWPAW FORMATIONS FOUR AND ONE-HALF
MILES SOUTHEAST OF FORT WORTH, TEXAS, AND JUST NORTH OF
THE INTERNATIONAL AND GREAT NORTHERN RAILWAY BRIDGE
ACROSS SYCAMORE CREEK

MAINSTREET:	Feet
White massive, in part marly, limestone, with <i>Kingena</i> (?) <i>wacoensis</i> (Roemer), <i>Exogyra arietina</i> Roemer, <i>Turrilites brazoensis</i> Roemer, <i>Ostrea carinata</i> Lamarck (?)	6.0

PAWPAW:

Brown clay, marlier ot the top, with an abundant pyrite fauna at the base, including: <i>Arca</i> sp., <i>Mortoniceras worthense</i> Adkins, <i>Acanthoceras worthense</i> Adkins, <i>Engonoceras</i> sp., <i>Schloenbachia</i> sp., <i>Hamites</i> sp., <i>Turrilites</i> sp., <i>Plicatula</i> sp., <i>Pecten subalpinus</i> (Böse), <i>Enallaster bravoensis</i> Böse, <i>E. wenoensis</i> Adkins, <i>Epiaster wenoensis</i> Adkins, and other species	20.7
---	------

WENO:

Chalky limestone; contains: <i>Ostrea</i> sp. aff. <i>diluviana</i> , <i>Ostrea curinata</i> (?) Lamarck, <i>Pachymya austinensis</i> Shumard, <i>Homomya</i> sp., <i>Schloenbachia</i> sp. (M.), <i>Pecten subalpinus</i> (Böse), <i>Pecten texanus</i> Roemer, <i>Plicatula</i> sp., <i>Nodosaria texana</i> Conrad, and many other species.....	13.5
Yellow marl with many limestone bands, poorly exposed.....	36.2
Hard white limestone, furoid, with <i>Gervilliopsis invaginata</i>	.5
Brownish yellow marl, about.....	15.0

SECTION OF PAWPAW AND WENO FORMATIONS EAST OF KATY LAKE
AND EAST OF SYCAMORE CREEK, ABOUT THREE AND ONE-HALF
MILES SOUTHEAST OF FORT WORTH, TEXAS

Section furnished by Gayle Scott

Section is from creek bed to upland; direction of section, 92 degrees from magnetic north.

MAINSTREET:

	Feet
Weathered limestone	15.6
Limestone ledge, with slight terrace above.....	2.7

PAWPAW:

Terrace, covered with soil and water-soaked, marl base; contains a sandstone ledge two feet below top	10.4
Red, ironstained, slightly sandy marl, with numerous small pyrite fossils; characteristic weathering of receding Pawpaw hillsides.....	12.0
Yellow marl, in part soil-covered overlying the flat narrow terrace formed by top member of Weno limestone.....	3.0

WENO:

Hard limestone ledge, partly concealed, about.....	3.2
Disintegrated thin limestone ledges alternating with marl layers of varying thickness, partly concealed by limestone debris and soil.....	10.5
Limestone ledge, impure, yellow, heavily ironstained, indurated ledge, containing <i>Pentagonaster texensis</i> Adkins and Winton (rare), and small starfish, indet. (fairly numerous). This ledge forms the base of the abrupt upper Weno terrace, and is easily recognizable in the Fort Worth region. PENTAGONASTER ZONE	1.0
Yellow marl, heavily ironstained, containing thin limestone ledges. NODOSARIA TEXANA HORIZON	6.0
Disintegrated limestone, capped by a thin limestone ledge.....	3.2
CONCEALED. Interval contains <i>Gervilliopsis</i> Zone	28.5
Heavy limestone ledge, impure, furoidal, yellow, and slightly less resistant than the one mentioned below.....	.9
Blue marl (pipe clay), weathering to gray, fractures on drying into small blocks.	3.5
Heavy limestone ledge, fairly pure, very resistant, containing <i>Pecten texanus</i> , <i>Pecten georgetownensis</i> . In many places forms the bed of Sycamore Creek	1.5
Blue marl (pipe clay), gray on weathering, and having a very fine texture	5.2

DENTON:

Top of *Gryphea washitaensis* conglomerate, in bed of creek..... 2.0

Further sections of the Weno formation in Denton, Johnson, and other counties have just been given in the discussion of the Pawpaw formation (q. v.).

STRATIGRAPHIC CORRELATION

NORTH CENTRAL TEXAS AND SOUTHERN OKLAHOMA

The Weno and Pawpaw formations can be traced continuously through these two regions and their correlation is obvious. As has been already noted, the sandstone facies of the Pawpaw formation has been included by Taff in his Bokchito formation. This facies is similar at Gainesville, Denison and Bokchito, and in itself affords insufficient differences to be considered as a different formation on the two sides of the Red River. The whole Weno formation is also included in the Bokchito formation of Taff. Taff's tabulation of equivalencies¹ makes clear that the Bokchito formation is to be considered equivalent to the undefined upper part of the Georgetown limestone in the Austin quadrangle, but to what formations in North Texas is not clear. However, his Caddo formation is described as having at its top a bed of oyster shells "similar to those occurring below and at the top of the Kiamichi formation" (ibid, p. 6). This is the Denton marl and has characteristic Denton fossils, and therefore the base of the Bokchito formation is to be considered as the base of the Weno.²

SOUTH CENTRAL TEXAS

The Weno and Pawpaw formations have been traced with certainty to the Brazos River south of Blum. The regional and facies differences in Central Texas make correlation difficult on account of the smallness of the known Del Rio fauna. It is presumed that the *Exogyra arietina* zone of the basal Del Rio clay is continuous with that of the Mainstreet limestone north of the Brazos near Blum, and that the *Nodosaria texana* zone present in West Texas at the top of the Del Rio just beneath the Buda limestone³ but reduced or absent in Central Texas does not correspond to the prominent *Nodosaria* zone found at the top of the Weno and the base of the Paw-

¹Taff: Atoka Folio, p. 6.

²Stephenson (U. S. G. S., Prof. Paper 120-H, pl. XVIII) places the Denton clay member and the *Ostrea carinata* bed capping it at the base of Taff's Bokchito formation; and also, very curiously, puts the "Gervilliopsis bed" near the base of the Denton clay.

³Böse: Univ. Texas Bull. 1902, p. 19.

paw formations in North Texas. *Turritiles brazoensis*, also a fossil of very limited vertical range, marks the top of the Georgetown limestone and the base of the Del Rio clay in Central Texas from the Brazos southwards. The basal Mainstreet then is represented by the uppermost Georgetown in Central Texas, and the Pawpaw if present by strata lying high in the Georgetown, while the bulk of the lower Del Rio (*Exogyra arietina* zone) is Mainstreet and the upper Del Rio is Grayson.

WEST TEXAS AND MEXICO

West of the southern turning point of the Comanchean outcrops in Bexar County, the Georgetown limestone and the Del Rio clay preserve their individuality into the Trans-Pecos region. At the type locality of the Del Rio clay, it is typically developed as a laminated, greenish-blue clay with thin intratified arenaceous clay seams, calcareous flags and shell breccia (*Exogyra arietina* and *Nodosaria texana*). In lithology this formation "varies from a clay to an arenaceous thin bedded limestone."¹ Here the limestone facies is beginning to appear. It increases in amount southwards and in the Mariscal Mountains the bulk of the formation is limestone. On the other hand, at Cerro de Muleros, Böse's subdivision 6, representing Weno and Pawpaw, is a marl with sand and lime, and his subdivision 7, representing the Mainstreet is sandstone. The lithological variations of the Weno and Pawpaw are summarized on page 27. The Georgetown, like the Buda, is over much of West Texas a fine grained crystalline semi-lithographic limestone, grading northwards into a marlier and sandier near shore phase. The same correlation as in Central Texas (page 40) holds for these strata.

The age of subdivision 6 of Cerro de Muleros is delimited by several of its fossils. The following fossils have not been found below this subdivision, nor below the Denton formation in North Texas: *Ostrea quadriplicata* Shumard, *Ostrea marcoui* Böse. *Pinna guadalupae* Böse is unknown below subdivision 6 and below the Weno of North Texas. *Enallaster bravoensis* Böse, known from subdivisions 5-6, is not known to occur lower than the Denton formation elsewhere. *Helicocryptus mexicanus* Böse of subdivision 6 is as yet known only from the Pawpaw of North Texas, and the same is true of *Placosmilia* spp., which are probably identical with *P. bravoensis* and *P. mexicana*. The zone of extreme abundance of *Gryphea washitaensis* Hill so conspicuous in this subdivision is paralleled in North Texas only by that at the top of the Denton formation.

¹Roberts and Nash: Univ. Texas Bull. 1803, pp. 14-15.

These facts indicate that subdivision 6 is equivalent to the Weno and Pawpaw and to part of the Denton formation. The overlying sand, lacking *Exogyra arietina*, contains near its top *Hemiaster calvini* Clark and *Exogyra whitneyi* Böse. The former fossil is upper Washita, Weno to Buda, and the latter is known from the Buda, although from the present record it is evident that it ranges as low as the Mainstreet formation.

It may be of interest to add a list of species described from Cerro de Muleros, which have been found in North Texas. The number cited is that of the corresponding division at Cerro de Muleros, and the following formation names refer to the location of the fossil in North Texas.

- Epiaster aguilaerae* Böse, 5; basal Fort Worth.
- Enallaster bravoensis* 5, 6, 8; Weno to Grayson.
- Holactypus limitis* 5; Weno to Grayson.
- Ostrea marcoui* 4, 5, 6; Weno to Grayson.
- Pinna guadalupae* 2, 6; basal Weno.
- Pecten subalpinus* 2, 3, 5, 6, 8, 9; Goodland to Grayson.
- Pecten irregularis* 1, 2; Walnut to Kiamitia.
- Plicatula subgurgitis* 2, 5, 6; Duck Creek to Pawpaw.
- Helicocryptus mexicanus* 6; Pawpaw.
- Turbo chihuahuensis*, La Encantada; Weno.
- Tylostoma chihuahuense* 1, 2; Glenrose to Goodland.
- Schloenbachia trinodosa* 5, 6; Duck Creek.

EUROPE AND NORTHERN AFRICA

It is unquestionable that the aggregation of Pawpaw pyrite ammonites here described parallels Pervinquièr's Vraconian fauna more closely than it does his Cenomanian fauna. This might mean that the Pawpaw fauna is Vraconian, or that some later (Cenomanian) species are included in his Vraconian fauna; or the Pawpaw fauna may be one which has persisted with little change from its Vraconian ancestors. *Scaphites aequalis*-like species in Texas are known to range from the Duck Creek marl to at least the Pawpaw clay; in Europe they are of Upper and Lower Cenomanian, Vraconian, or even Albian age, so that this species decides little; the Pawpaw or the Duck Creek *aequalis* may be contemporaneous with the Vraconian examples. *Flickia* decides nothing, since the range of the genus is unknown and is extended with each new species discovered. *Hamites simplex* ranges from Aptien to Mid-Cenomanian. Attempts have been made to correlate the Texas Washita division on the basis of *Schloenbachia inflata*. This brings us face to face with the question, What is *inflata*? Is it, as sometimes claimed, a *Schloenbachia* with low, sigmoidal ribs and two tubercles, one umbilical and one marginal; or is it a *Mortoniceras*, with square volutions and coarse short ribs having three tubercles, an umbilical

and a pair of twinned ventro-marginal ones, as indicated by Pervinquière? If the latter, there is no Duck Creek species known to me that approaches it as closely as do several Upper Washita species. For only above the base of the Weno do we begin to find fairly large (6-12 inches) coarse straight ribbed square coiled ammonites with an umbilical and a twinned marginal tubercle; this is the commonest and most characteristic Weno calcite species, and it abounds in slightly smaller size in the nacreous ironstone Weno fauna (Plate 3, figure 11). In the Pawpaw clay the same type is abundant as small pyritic casts (*M. worthense*, Plate 1, figures 6-10, 18-19, 26), having extremely short coarse ribs with very prominent twinned marginal tubercles and a narrow square volution. Again in the Mainstreet limestone the common calcite species has lower, somewhat sigmoid ribs with twinned marginal tubercles and a rectangular section. Are any of these *inflata*? If the Vraconian is to be placed in the Weno, which seems possible, the whole Cenomanian will have to lie between the Pawpaw and the base of the Eagleford shales, which is generally claimed to be the base of the Turonian. Again a Duck Creek species similar to *Schloenbachia elobensis* has been noted; but a comparison of Szajnocha's figures and description indicate that this species is different from his; its section is much lower, and the shell bears spiral circllets which form regular radial rows containing numerous imbricated lips, while in the Texas species (known only as internal casts) there are radial rows of only a few rounded tubercles; in addition, the stratigraphic data given by Szajnocha are inadequate for correlation with the Texas strata.

A comparison of the Duck Creek limonite fauna with the African Vraconian yields only slight results. On the theory that the Pawpaw is Vraconian, the Duck Creek could scarcely be placed earlier than the Albian, since the Glenrose is with some certainty assigned to the Aptian. But Albian species¹ are not apparent in the Duck Creek limonite fauna, whose ammonites are preponderantly small species of *Schloenbachia* and *Hamites*, with scattering *Scaphites* sp. aff. *aequalis*, *Desmoceras*, etc. *Hoplites* and other abundant Albian genera are so far unknown. The fauna on the other hand is predominantly Vraconian. This Vraconian aspect appears to end with the Duck Creek limestone or marl, since with the next limonite fauna, that of the Denton marl, there seem to be some species with Cenomanian affinities. Comparison of the Texas Washita faunae with those of Western Europe, examples of which are before me, shows much less resemblance than with the Northern African fauna. Common European species, as *Douvilleiceras mammilare*, *Schloenbachia varians*, *S. mantelli*, and *Holaster subglobosus* have no representatives in the Texas material so far known. On the other hand, the Mainstreet, Grayson and Buda faunae are

¹Listed among other places in: Tomitch: Contributions à la Connaissance de l'étage Albien dans le Sud-Est de la France. Le Mans, 1918.

of a higher Cenomanian type, containing for example *Codiopsis* sp. aff. *doma*, *Tissotia* spp., and *Acanthoceras* spp.; and Berry states¹ that the Dakota (Woodbine?) flora is of Turonian age.

On the whole then the Pawpaw fauna is placed provisionally as Cenomanian with much uncertainty as to its exact position. This question will be settled when the rich ammonite faunae of the Lower Washita are critically studied. The Texas Comanchean sea was transgressive or at least not regressive for a long period so that the relatively stationary shallow-water conditions may not have encouraged a rapid evolution of these ammonites; whatever the cause, they present notable similarities from the earliest to the latest pyrite faunae known in the Texas Comanchean. On the theory of persistent and nearly stationary species from Vraconian to Cenomanian time some nearly similar species should be much extended vertically in the Lower Washita sediments; further study of the proper marine facies of each stratigraphical level should reveal these and by filling in the great gaps in our paleontological knowledge give a fairly complete faunal succession. It is hoped that information on the extent of the various marine facies in the Texas Comanchean formations will contribute to this result.

PALEONTOLOGY

The rich Weno and Pawpaw faunae initiate a new group of species after the considerable paleontological break which occurs at the top of the Denton marl. The Weno formation, besides containing characteristic ammonites, is notable in its marl facies for an exuberant echinoid fauna, and in its shale facies for a great abundance and variety of distinctive cephalopods, pelecypods and gastropods preserved with the original nacreous shell. Certain elements of this nacreous fauna forcibly remind one of the Eocene faunae, with which they have many genera in common. The fossils are perfectly preserved, often with iridescent luster, and the original shell shows the minutest details: in pelecypods, prodissoconch and other embryonic stages, ligament, and usually the finest features of dentition and external ornamentation; in the cephalopods, the original pearly or iridescent shell, and beneath this, finely preserved sutures etched into the ironstone interior.

The Pawpaw formation on the other hand contains a diverse and characteristic assemblage of small pyritic ammonites, echinoids, pelecypods,

¹Berry: U. S. G. S., Prof. Paper 84, pp. 71, 128.

gastropods, corals and other fossils.¹ This fauna has close resemblances to that described by Pervinquire² from the Vraconian of Tunis and Algeria, a subject that will be considered later.

The later Washita seas were transgressing over extensive land areas and a large numerical expansion in the marine fauna occurred, probably in part in the invaded trough already described. In addition, the portion of the ocean bottom which received the mixed clay and sand deposits was probably at a moderate depth and not far offshore, and permitted a wealth and variety of fauna. Note the complexion of life in the Pawpaw seas: swarms of sharks and dogfish; a multitude of bottom-loving crustacea, small crabs and lobsters, echinoids, oysters and scallops; sessile corals and worms; pelagic protozoa; a few large, free-swimming nautili and iridescent-shelled ammonites, and a great diversity of small ammonites.

THE WENO FAUNA

Of the large and well preserved Weno fauna only a few species have been mentioned in the literature, as follows:

Hill³ listed the following fauna from the North Denison sands (=Weno):

<i>Azinea</i> sp.	<i>Tapes</i> sp.
<i>Nuculaea</i> sp.	<i>Turritella</i> sp.
<i>Corbula</i> sp.	

And the following from the Pawpaw shales and the North Denison sands (not separated):

<i>Turritella</i> sp. (predominant)	<i>Trigonia emoryi</i> Conrad
<i>Corbula</i> sp.	<i>Pholadomya postextensa</i> Cragin
<i>Azinea</i> sp.	<i>Cyprimeria</i> (large sp.)
<i>Volsella</i> sp.	<i>Anchura mudgeana</i> White
<i>Tapes</i> sp.	<i>Leda</i> 2 spp.
<i>Cytherea</i> sp.	<i>Mactra</i> sp.
<i>Tellina</i> sp.	<i>Dentalium</i> sp.
<i>Avicula</i> sp.	<i>Scclaria</i> sp.
<i>Gervilliopsis</i> or <i>Gervillea</i>	<i>Ostrea quadriplicata</i> Shumard
<i>Arca</i> sp.	<i>Pecten inconspicuus</i> Cragin
<i>Nucula</i> sp.	<i>Yoldia microdonta</i> Cragin
<i>Ammonites emarginatus</i> Cragin	<i>Turritella seriatim-granulosa</i> Roemer
<i>Sphenodiscus belviderensis</i> var.	<i>Astarte</i> sp.
<i>serpentinum</i> Cragin	

¹Throughout this paper an asterisk (*) preceding the name of a fossil indicates that the fossil has pyrite, limonite or hematite preservation; a double dagger (‡) indicates nacreous preservation.

²Pervinquire: tudes de Palontologie Tunisienne. Paris, 1907.

³Hill: Bull. Geol. Soc. Amer., 5, 1894., and 21st Ann. Rept., U. S. G. S. pt. 7, p. 277, 1901.

The following species of the fauna have been described by Cragin:¹

<i>Pecten inconspicuus</i>	<i>Tellina subaequalis</i>
<i>Pholadomya postextensa</i>	<i>Sphenodiscus emarginatus</i>
<i>Corbula crassicistata</i>	<i>Engonoceras belviderensis</i> var. <i>serpentinum</i>
<i>Tapes denisonensis</i>	

The following species were listed by Stephenson:²

<i>Nucula</i> sp.	<i>Cymbophora</i> sp.
<i>Ostrea quadriplicata</i> Shumard	<i>Turritella</i> sp.
<i>Protocardia texana</i> (Conrad)	<i>Anchura mudgeana</i> White
<i>Cyprimeria</i> sp.	<i>Engonoceras serpentinum</i> Cragin
<i>Corbula</i> 3 spp.	Crustacea

An incomplete list of the Weno fossils on hand is given in the accompanying table (Table 1) ; many species were omitted for further study.

RANGES OF WENO FOSSILS

The range as given in this table will demonstrate that one of the major paleontological breaks in the Washita division lies at the base of the Weno formation. The contact between the Denton and Weno formations is apparently conformable, and many species cross it; but a considerable number of others do not range higher than this contact while another considerable group does not range lower. Without speculating as to the origins of the latter group, it may be delimited more closely in view of a more detailed study.

ABBREVIATIONS FOR TABLE OF WENO FOSSILS

* Pyrite, limonite or hematite preservation.

† Nacreous preservation.

A Abundant

O Occasional

R Rare

S Sand phase

SH Shale phase

M Marl phase

L Limestone phase

u upper part of formation

m middle

l lower

¹Cragin: Colo. Coll. Stud., 5, 1894, p. 49.

²Stephenson: U. S. G. S., Prof. Paper 120-H, 1918, p. 141.

- 601 Pit of Gainesville Brick Company, brickyards one and three-fourths miles south-east of Gainesville, Texas, just east of Rock Creek (branch of Pecan Creek) and just south of the Missouri, Kansas and Texas railway (Wichita Falls branch). Includes the ironstone dump on east rim of pit. Shale facies basally; ironstone and sand facies above.
- 602 East bank of Sycamore Creek, two and one-half miles southeast of Fort Worth, Texas. Marl facies basally; limestone facies above.
- 604 Cut of St. Louis and San Francisco railway, three-fourths mile north of Union Station, Denison, Texas. Shale facies basally; ironstone and sand facies above.
- 605 A branch of Duck Creek, three-fourths mile north of Union Station, Denison, Texas, and just west of locality 604. Shale facies.
- 606 Exposure in valley just south of St. Louis and San Francisco track, two miles north of Denison, Texas. Shale facies.
- 611 Waterfall 200 yards south of crossing of International and Great Northern railway over Houston and Texas Central railway track near Sycamore Creek, three miles southeast of Fort Worth, Texas. Marl facies.
- 612 Cliff on northeast bank of Sycamore Creek, 100 yards north of the Houston and Texas Central railway bridge across Sycamore Creek, three miles southeast of Fort Worth, Texas. Marl facies basally; limestone facies above.
- 618 Area in valley east of Armstrong Iron Works, South Hemphill street, Fort Worth, Texas, and west of the International and Great Northern railway track. Marl facies basally; limestone facies above.

THE PAWPAW FAUNA

As already stated, the bulk of this fauna is limonitic, pyritic or hematitic, but there are in addition some fossils with calcareous preservation. The calcareous fossils are echinoids, pelecypods, a few cephalopods and other miscellaneous fossils. The echinoids of the basal portion of the formation are mainly limonitic, but those of the upper, marlier portion over all of the Pawpaw outcrops except in the Red River region, are calcareous mud-filled tests.

LIST OF PAWPAW LOCALITIES

- 714 One-fourth mile south of the International and Great Northern railway bridge across Sycamore Creek, four and one-half miles southeast of Fort Worth, Texas. Entire thickness of Pawpaw, somewhat overwashed at top; north-eastward facing slope of hillside, and beneath the slope a considerable terrace on the top of the Weno limestone. Clay facies basally; some marl at top.
- 715 Amphitheater one-half mile northeast of locality 714. Entire thickness of Pawpaw formation exposed.
- 716 Pit of Cobb Brickyards, one-fourth mile east of Sycamore Creek and three miles southeast of Fort Worth, Texas.
- 718 Cut of International and Great Northern railway track one-fourth mile south of bridge across Sycamore Creek, four and one-half miles southeast of Fort

- Worth, Texas, and one-eighth mile east of locality 714. Upper part of Pawpaw formation, partly marl, overlain by Mainstreet limestone.
- 719 Crowley Road, one mile south of the Baptist Seminary, and four and one-half miles south of Fort Worth, Texas. Westward facing cut in hillside, exposing poorly most of the Pawpaw formation.
- 720 Rim of Weno escarpment one mile southeast of Riovista, Texas, and midway between the Riovista-Waco pike and the Santa Fé track. The Pawpaw, similarly exposed between the Mainstreet and Weno limestones, continues westward to the next locality.
- 721 East-west escarpment just east of Riovista-Waco road, one mile south of Riovista, Johnson County, Texas. The Pawpaw is much thinned, and is transitional between clay and marl facies.
- 722 Cut of Missouri, Kansas and Texas railway (Wichita Falls branch) at north end of pit of the Gainesville Brick Company, one and three-fourths miles southeast of Gainesville, Texas.
- 723 Westward facing hillside lying under Mainstreet upland on west side of Mansfield Road, halfway between Glen Garden Country Club and Sycamore Creek, three miles southeast of Fort Worth, Texas.
- 724 Hillside on north side of east-west road, three miles southeast of Haslet, Tarrant County, Texas. Whole thickness of Pawpaw.

FAUNA OF THE PAWPAW FORMATION

S: Sand facies; C: Clay facies; M: Marl facies; L: Limestone facies; A: Abundant; O: Occasional; R: Rare; l: lower third of Pawpaw formation; m: middle portion of Pawpaw formation; u: upper portion of Pawpaw formation; *: Pyrite or limonite preservation; †: nacreous preservation.

SPECIES	LOCALITY: 716	714	716	718	719	723	724	722	720	721
	C	C	C	M	C	C	CS	S	M	M
CEPHALOPODS:										
<i>Nautilus texanus</i> Shumard	R	R	R	O	--	R	--	--	R	R
<i>Nautilus</i> sp.	--	R	R	O	--	--	--	--	R	--
* <i>Stolteckia</i> sp. aff. <i>dispar</i>	--	R	--	--	--	R	--	--	--	--
* <i>Flickia boeseli</i> Adkins	--	--	--	--	--	--	--	--	--	Rl
* <i>Acanthoceras</i> <i>worthense</i> Adkins	A	A	O	--	O	A	O	--	--	--
* <i>Acanthoceras</i> sp. aff. <i>suzannae</i>	--	A	--	--	--	A	--	--	--	--
* <i>Hamites</i> <i>tenawa</i> Adkins and Winton	--	R	--	--	--	R	--	--	--	--
* <i>Hamites</i> sp. aff. <i>armatus</i>	--	R	--	--	--	R	--	--	--	--
* <i>Scaphites</i> <i>hilli</i> Adkins and Winton	--	Ol	O	--	O	A	R	--	--	--
* <i>Scaphites</i> sp. aff. <i>worthensia</i>	--	Ol	--	--	--	O	--	--	--	--
* <i>Baculites</i> <i>comanchensis</i> Adkins	--	R	--	--	O	--	--	--	--	--
* <i>Turrillites</i> <i>worthensis</i> Adkins and Winton	A	A	A	--	O	A	R	--	--	--
* <i>Turrillites</i> sp. B. Adkins and Winton	O	A	O	--	--	A	--	--	--	--
* <i>Turrillites</i> sp.	A	A	A	--	A	A	O	--	--	--
† <i>Schloenbachia</i> <i>wintoni</i> Adkins	R	R	R	--	--	--	--	--	--	--
* <i>Mortoniceras</i> sp.	--	O	--	--	--	O	--	--	--	--
* <i>Mortoniceras</i> <i>worthense</i> Adkins	O	A	O	--	O	A	R	--	--	--
* <i>Engonoceras</i> sp.	A	A	A	--	O	A	O	--	--	--
† <i>Engonoceras</i> <i>serpentinum</i> Cragin	--	R	R	--	--	R	--	--	--	--
* <i>Puzosia</i> sp.	--	R	--	--	--	--	--	--	--	--
* <i>Lytoceras</i> sp. aff. <i>marut</i>	--	R	--	--	--	R	--	--	--	--
* <i>Hamulina</i> <i>worthensia</i> Adkins	--	R	--	--	--	--	--	--	--	--
ASTEROIDEA:										
* <i>Pentaceros</i> <i>americanus</i> Adkins	--	Rl	--	--	--	--	--	--	--	--
* <i>Metopaster</i> <i>hortensae</i> Adkins and Winton	--	Rl	R	--	--	--	--	--	--	--
* <i>Comptonia</i> <i>wintoni</i> Adkins	--	Rl	--	--	--	--	--	--	--	--
*sp. indet.	--	Rl	--	--	--	--	--	--	--	--
OPHIUROIDEA:										
<i>Ophioglypha</i> sp.	--	R	--	--	--	--	--	--	--	--

SPECIES	LOCALITY: 716	714	715	718	719	723	724	722	720	721
ECHINOIDEA:										
*Goniophorus sp. -----	--	R	--	--	--	--	--	--	--	--
*Peltastes sp. -----	--	R	--	--	--	--	--	--	--	--
Enallaster bravoensis Bose -----	O	O	O	O	--	--	--	O	O	O
Enallaster sp. aff. texanus -----	O	--	O	O	--	--	--	O	O	O
Enallaster riovistae Adkins -----	O	--	--	--	--	--	--	--	A	A
Hemiaster wenoensis Adkins -----	--	--	--	--	--	--	--	--	--	O
Hemiaster calvini Clark -----	O	O	O	O	--	--	--	O	O	O
Epiaster wenoensis Adkins -----	O	--	O	O	--	--	--	O	O	O
Epiaster aubobesius Adkins -----	O	--	O	O	--	--	--	--	O	O
Holaster sp. -----	O	--	O	O	--	--	--	--	O	O
*Holaster sp. -----	--	R	O	O	--	--	--	--	O	O
Holactypus limitis Bose -----	O	O	O	O	O	O	R	--	A	O
*Goniopygus sp. -----	--	R	--	--	--	--	--	--	--	--
CRUSTACEA:										
crabs, indet. -----	--	R	--	--	--	R	--	--	--	--
lobsters, indet. -----	--	R	--	--	--	R	--	--	--	--
CORALS:										
*Placosmilia spp. -----	--	R	--	--	--	R	--	--	--	--
*Trochosmilia sp. -----	--	R	--	--	--	R	--	--	--	--
corals, indet. -----	--	R	--	--	--	--	--	--	--	--
PROTOZOA:										
*Nodosaria texana Conrad -----	O	A	O	--	O	O	--	--	O	--
PELECYPODS:										
*Arca washitaensis Adkins -----	A	A	A	--	A	A	--	--	R	--
Ostrea carinata (?) Lamarck -----	--	O	--	--	--	--	--	--	--	--
Ostrea quadriplicata Shumard -----	--	O	--	--	--	O	--	--	--	--
Gryphea washitaensis Hill -----	O	O	O	O	O	O	O	--	O	--
Remondia (?) acuminata (Cragin) -----	R	R	R	--	--	R	--	--	--	--
Plicatula spp. -----	A	A	A	A	A	A	A	--	A	--
Pecten subalpinus (Bose) -----	R	R	R	--	--	--	O	--	O	--
Nucula sp. -----	O	O	O	--	--	--	O	--	O	--
Leda sp. -----	O	O	O	--	--	--	R	--	--	--
Lima sp. -----	O	A	A	--	--	--	O	--	O	--
GASTROPODS:										
Helicocryptus mexicanus Bose -----	--	R	--	--	--	--	--	--	--	--
*Turritella sp. -----	O	R	O	O	--	R	--	--	--	--
*Turbo sp. -----	O	O	--	--	--	--	R	--	--	--
*Cinulia sp. -----	R	R	R	--	--	R	--	--	--	--
*Nerinea sp. -----	O	R	O	--	--	R	--	--	--	--
VERTEBRATES:										
sp. indet. (shark?, head bones) -----	--	R	--	--	--	R	--	--	--	--
sp. (Odontus appendiculatus?) -----	A	A	A	O	O	A	O	--	--	--
sp. indet. (teeth) -----	A	A	A	A	O	A	--	--	O	--
sp. indet. (vertebrae) -----	A	A	A	A	--	A	--	--	O	--

PARALLELS BETWEEN UPPER WASHITA AND LOWER WASHITA ECHINOIDS

LOWER WASHITA (below Weno formation)	UPPER WASHITA (Weno and higher formations)
Holactypus planatus Roemer	H. limitis Böse
	E. bravoensis Böse
Enallaster texanus Roemer	E. riovistae Adkins
	E. wenoensis Adkins
	E. sp. aff. texanus Roemer
Hemiaster whitei Clark and E. elegans Shumard	H. calvini Clark

<i>Epiaster aguilerae</i> Böse	<i>E. wenoensis</i> Adkins
<i>Holaster simplex</i> Shumard	<i>H. sp.</i>
<i>Cyphosoma texana</i> Roemer	<i>C. volanum</i> Cragin
<i>Salenia mexicana</i> Schlueter and <i>S. texana</i> Credner	<i>S. volana</i> Whitney
<i>Goniopygus</i> sp.	<i>G. budaensis</i> Whitney
<i>Goniophorus</i> sp. 1	<i>G. sp. 2.</i>

COMPARISON OF PYRITE FAUNA

TEXAS

It has been pointed out¹ that the North Texas Comanchean system contains a series of alternate lime and marl (or clay) formations, some of which contain conspicuous pyrite and limonite faunae; these are

- (1) The Kiamitia formation;
- (2) The Duck Creek marl;
- (3) The Denton marl;
- (4) The Pawpaw clay;
- (5) The Grayson formation;
- (6) The Del Rio clay.

Many pyrite and limonite species are common to these six formations, and still others have more restricted vertical ranges and are confined to limited levels within a single formation. In general it may be said that these fossils are of distinct species from those otherwise preserved in the same formation; that they are rare in the intervening formations, and that the complexion of the limonite faunae is different from that of the other faunal components of the formations in question. In addition there seem to be certain limonite species in the various intervening limestone formations, especially in the Fort Worth and Weno limestones. To assist in determining the age and correlation of these formations the following provisional fossil lists are given. The list for the Pawpaw formation will be found on page 51.

The rather regular alternation of marl (or clay) and lime formations in the North Texas section suggests some form of cyclic deposition, in which the same marine conditions were many times repeated, and this inference is equally strong for the contained pyrite faunae, when it is found that they occur prevailing in the alternate, clay, formations. This

¹Winton and Adkins: The Geology of Tarrant County, Univ. Texas Bull. 1931. 1920.

does not imply that in certain formations no pyrite faunae will be found; they existed somewhere during the whole Washita at least, and likely will be found in the clay facies of each formation, just as representatives of the Mainstreet pyrite fauna have been found in the clay facies (Del Rio) of Central Texas, though apparently absent in the limestone of North Texas. It is certain that exuberant pyrite faunae are associated with clay facies; and accordingly the greatest abundance of pyrite fossils in the Kiamitia is in southern Tarrant and Johnson counties; Denton formation, Denison to northern Tarrant County; Pawpaw formation, Tarrant County; Mainstreet formation, McLennan County and southward; middle Grayson formation, Denton County. In each instance the lithology of the region mentioned is largely of the clay facies. These facts suggest recurrent similar conditions which favored the spread of pyrite faunae. At a given locality this recurrence might be many times repeated; and such pyrite zones are still being discovered at new stratigraphic levels. What these conditions and their causes were will not be discussed here. They resulted in each zone in a highly characteristic pyrite and limonite fossil association, which is as follows: *Arca washitaensis*, *Engonoceras* sp., *Scaphites* spp., *Turritites* spp., *Hamites* spp., *Cinulia* sp., *Nerinea* sp., *Lunatia* sp., *Turritella* sp., pyritic starfishes, and abundant small, non-pyritic crustacea, especially crabs.

KIAMITIA CLAY FAUNA

The marl facies of the Kiamitia formation extends from Primrose, southwest of Fort Worth, to a point between Gainesville and Fink, on the Red River. The clay facies extend from near Primrose to at least the Brazos River; it is found throughout Johnson County, and on Cedar Creek near Blum is a reddish clay 19 feet thick. With the advent of the clay facies the shell conglomerates of the northern section disappear and a pyrite fauna similar to that of the Duck Creek marl appears. This little investigated fauna contains *Arca* sp., and other small pyritic and limonitic pelecypods and gastropods.

DUCK CREEK MARL FAUNA

This fauna is notable for its large number of small pyritic and limonitic gastropods and pelecypods. Such genera as *Cerithium*, *Cinulia*, *Turritella*, *Turbo*, *Lunatia*, *Arca*, *Nucula* and *Corbula* are highly characteristic of the lower Washita pyrite zones. Equally characteristic is

the exceptional variety of gerontic ammonites, especially *Hamites*; *Scaphites*, *Schloenbachia*, *Desmoceras* and other genera are also prominent in the fauna.

DUCK CREEK LIMONITE AND PYRITE FOSSILS

**Scaphites worthensis* Adkins and Winton.

**Scaphites* sp. aff. *worthensis*.

**Engonoceras* sp.

Hamites tanima Adkins and Winton.

**Hamites* spp.

**Crioceras* (?) sp.

**Schloenbachia* sp.

**Neolobites* (?) sp.

**Arca* sp.

**Nucula* sp.

**Turbo* sp.

**Lunatia* sp.

**Cerithium* sp.

**Nerinea* sp. aff. *pellucida* Cragin

**Cinulia* sp.

**Anchura* sp.

**Turritella* sp.

**Nerinea* sp.

**Placosmilia* sp.

DENTON CLAY FAUNA

The clay facies of the Denton formation extends from northern Tarrant County (Blue Mound, near Haslet) to beyond Denison. It is characterized by a great diversity of small pyrite and limonite ammonites, some of them nacreous, and by many small crustacea, mainly small pyritic crabs. The ammonites seem more abundant southwards than on the Red River and the crustacea the reverse.

FAUNA OF THE DENTON FORMATION

* Limonite or pyrite species.

U Upper.

L Lower.

**Acanthoceras* sp. aff. *aumalense* Coquand

**Engonoceras* sp.

**Mortoniceras* sp.

- *Schloenbachia sp.
- *Acanthoceras (?) sp.
- *Turrilites sp. (?)
- *Baculites (?) sp.
- Ophioglypha texana Clark
- *Starfish sp. (ray)
- *Goniophorus (?) sp. 2 specimens, one with the shell.
- Leiocidaris sp. (spine and plate).
- Leiocidaris hemigranosus Shumard.
- Hemiaster sp. (plate).

- Hoploparia sp.
- Cambarus (?) sp. (like sp. in Gainesville brickyards Weno).
- Crabs 9 spp.

- Ostrea quadriplicata Roemer.
- Ostrea carinata Lamarck.
- Gryphea washitaensis Hill.
- Ostrea sp. (small zigzag).
- Stearnsia robbinsi White (?).
- Gervilliopsis invaginata White (?)
- Trigonia emoryi Conrad.
- Plicatula dentonensis (?) Cragin.
- *Arca sp.
- Anomia sp.
- Plicatula spp.
- *Nucula sp.
- *Cardium sp.
- *Leda sp.
- *Corbula sp.
- Pecten sp. aff. inconspicuus Cragin.
- Crania sp.
- Dentalium sp.
- ?Porocystis-like masses.

- *Natica sp.
- *Cinulia sp.

- Fish teeth, vertebrae, skin, plates.

GRAYSON FAUNA

The pyrite and limonite fauna so far discovered in the Grayson formation is confined to the middle clay member of the formation in North Texas, and includes diverse ammonites, gastropods and pelecypods, as listed below. However, the pyrite fauna of the middle Del Rio clay which occurs in McLennan County and southward is of Grayson age,

and contains some identical species. This latter fauna is well developed on the South Bosque River, five miles west of Waco and includes *Turrilites bosquensis* Adkins, *Flickia* (?) *bosquensis* Adkins, *Acanthoceras worthense* Adkins, *Schloenbachia* spp. and other common Grayson and Pawpaw pyrite species. It is notable that the preservation of this fauna is almost exclusively pyritic.

FAUNA OF THE GRAYSON MARL

* Pyrite or limonite species.

(C) Cidarid zone.

(Co) Coral zone.

U Upper member.

M Middle member.

L Basal member.

CEPHALOPODS:

Turrilites sp. (medium size) (M).

**Turrilites* sp. aff. *worthensis* A and W.

**Turrilites* spp. (M).

**Turrilites* sp. (whiplash) (M).

*ammonite aff. *Flickia* (small, keelless) (M).

**Hamites* 3 spp. (M).

Acanthoceras ? sp. (Denison, WSA).

Acanthoceras spp.

Engonoceras spp. (L).

**Schloenbachia* spp.

ECHINODERMS:

Hemiaster calvini Clark.

starfish sp.

Goniophorus (?) sp.

cidarid spines 3 spp. (C).

Cyphosoma volanum (?) Cragin.

Enallaster texanus (?) Roemer.

Enallaster sp. aff. *traski* (?) Whitney.

Enallaster sp. aff. *bravoensis* Böse (?).

CORALS:

**Placosmilia* (?) sp. (Co).

GASTROPODS:

Cerithium sp.

Turritella marnochi (?) White.

Turritella sp.

Cinulia pelletti Whitney.

**Gyrodes* (?).

**Turbo* sp.

**Turritella* sp. (sharp spired) M.

PELECYPODS:

- Gryphea mucronata Gabb.
- Gryphea sp. aff. corrugata Say.
- Exogyra sp. aff. columbae Sowerby (L).
- Exogyra n. sp. (Denison, M).
- Exogyra sp. aff. texana (U, under Buda limestone, Bosque River, W of Waco).
- Pecten texanus Roemer.
- Pecten subalpinus Böse.
- *Arca sp.
- Ostrea sp. aff. subovata Shumard.
- Lima sp. (elongate).
- Lima sp. (quadrate).
- Lima sp. aff. wacoensis Roemer.
- Ostrea sp. (saucer).
- Anomia sp.
- Plicatula spp.
- Protocardia texana (?) Conrad.
- Pholadomya shattucki Böse.
- Tapes sp.
- Inoceramus sp. M.
- Corbula (?) sp.
- Trigonia sp.
- Cyprimeria sp. (small).
- Cyprimeria sp. aff. crassa Meek.
- *Nucula sp.
- *Crassatella (?) sp. ("Remondia").
- *Barbatia sp.
- Trigonia sp. (large).
- *Isocardia sp.
- Shark teeth, vertebrae, bones.

DEL RIO CLAY FAUNA

The Del Rio limonite fauna is very widespread in Texas, having been found at Waco, South Bosque, Austin, Quihi (Medina County), Del Rio, Terlingua, and the Solitario.

In McLennan County at the base of the Del Rio clay, which corresponds to the middle of the Mainstreet limestone, an extensive pyrite and limonite fauna occurs, including the following:

- *Exogyra arietina Roemer.
- *Turritella sp.
- *Cerithium sp.
- *Turritiles spp.
- *Schloenbachia sp.

- **Nerinea* sp.
- Lunatia* sp.
- **Natica* sp.
- Goniophorus* sp.
- Pecten subalpinus* Böse.
- Turrilites brazoensis* Roemer.

The Del Rio pyrite fauna corresponding to the base of the Grayson formation of North Texas has already been noted; it includes:

- **Turrilites bosquensis* Adkins.
- **Flickia* (?) *bosquensis* Adkins.
- **Acanthoceras worthense* Adkins.
- **Schloenbachia* sp.
- Exogyra arietina* Roemer.
- Gryphea mucronata* Gabb.
- Gryphea* sp.
- Pecten subalpinus* (Böse).

The first named limonite fauna with an extensive calcite microfauna immediately overlies the Georgetown limestone and is of middle Main-street age. It is seen at South Bosque, and will probably be discovered at many places in Central Texas.

Liddle states that the Del Rio clay throughout Medina County, contains great amounts of dark brown limonitic fragments and fossils.¹ Good localities are a small Del Rio inlier four miles slightly west of north of Quihi on the Bandera road, and the main area of exposure just north of the inlier. These localities contain limonitic gastropods, **Turrilites*, **Scaphites*, small ammonites, and loose **Nodosaria*.

At Loma de la Cruz and other localities in the Del Rio clay lowland two miles south and southeast of Del Rio, the basal clay contains limonitic **Schloenbachia* sp., **Hamites* sp., **Scaphites* sp., **Turrilites* (two species), *sp. aff. *Flickia* (?) *bosquensis* Adkins; **Pyrina* or *Cassidulus* sp., cidarid sp., **Leda* (two species), **Nucula* sp., **Arca* sp. **Plicatula* sp.; **Nerinea* sp. (abundant), **Cerithium* spp. and other fossils.

Along the base of the Reed Plateau, Terlingua, the basal Del Rio clay contains abundant brown limonite fossils, the great majority of which are various species of *Turrilites*. The fauna includes: **Nodosaria texana* Conrad, **Turrilites* (three species), **Acanthoceras* sp., **Schloenbachia* sp.; **Nucula* sp.

¹Liddle, The Geology and Mineral Resources of Medina County, Univ. Texas Bull. (in press).

The Del Rio clay surrounding the Solitario Uplift contains a rich assortment of brown limonitic fossils, including **Engonoceras* sp., **Turrilites* (two species), **Turritella* sp., gastropods, **Nucula* sp., **Tapes* sp., **Nodosaria texana* Conrad, and other fossils.

It may be mentioned that these limonite faunae in Texas are not confined to the Comanchean, since a rich fauna was found in the Terlingua beds (Taylor marl equivalent) about five miles north of the crossing of the Alpine-Terlingua road through Terlingua Creek, Brewster County, Texas. This fauna includes: **Turrilites* sp., **Baculites* sp., **Ptychoceras* sp., **Desmoceras* sp., **Trochosmilina* sp., **Lingula* sp., **Lunatia* sp., **Natica* sp., **Nerinea* sp., numerous other gastropods, and fish teeth and vertebrae.

Böse¹ found a rich Vraconian fauna in limestone blocks covering small hills just west of Camacho, Zacatecas, between this station and the Trinidad mine, and west of Opal, Zacatecas, in the core of an anticlinal hill consisting of thin-bedded limestone with lenses and concretions of chert. These fossils are mainly silicified, but are cited here on account of their striking resemblances to the pyrite fossils described in this paper. The fauna includes typical Vraconian genera, as *Phylloceras*, *Lytoceras*, *Macroscaphites*, *Hamites*, *Hamulina*, *Ptychoceras*, *Diptychoceras*, *Anisoceras*, *Turrilites*, *Baculites*, *Desmoceras*, *Acanthoceras*, *Ancycloceras*, *Toxoceras*, *Crioceras*, *Scaphites*, *Schloenbachia*, *Brancoceras* and *Exogyra*.

SUMMARY OF PYRITE FAUNAE

The Washita marl and clay faunae contain distinctive assemblages of ammonites, crustacea, starfishes, echinoids, gastropods, pelecypods, corals, and other fossils. The crustacea are preserved partly with the original integument and partly with limonite replacement. The other fossils are in part calcitic, but mainly limonite and hematite pseudomorphs and casts of the interior of the original shell.

Ammonites: These are very distinctive for each fauna. Species of *Schloenbachia* abound in each marl and clay formation. *Acanthoceras* is found in the Denton, Pawpaw and Grayson faunae, but not in the Duck Creek. *Hamites* are abundant in the Duck Creek marl and present in the other formations. *Turrilites* abounds in the Pawpaw and is not known with certainty below it. *Flickia*, *Hamulina*, *Baculites* and *Puzosia* are known only from the Pawpaw clay.

¹Böse, On some new Cretaceous Faunas from Mexico, Univ. Texas Bull. (in press).

TABULATION OF KNOWN AMMONITES OF THE WASHITA PYRITE AND LIMONITE FAUNAE

Species	Number of Known Species in:				
	Duck Creek marl	Denton marl	Pawpaw clay	Grayson marl	Del Rio clay
Hamites	10	1	3	3	..
Turrilites	..	1?	10	4	1
Acanthoceras	..	2	2	1	1
Scaphites	2	..	2	..	..
Flickia	..	..	1	..	1?
Baculites	..	..	1	..	..
Schloenbachia	3	1	2	2	1
Mortoniceras	1	..	2	..	..
Puzosia	..	..	1	..	..
Hamulina	..	..	1	..	..
Engonoceras	1	1	3	1	..
Crioceras (?)	1	..	..	..	..
Neolobites (?)	1	..	..	..	..

ECHINODERMS:

TABULATION OF THE KNOWN ECHINODERMATA OF THE WASHITA LIMONITE FAUNAE

Species	Number of Known Species in:				
	Duck Creek marl	Denton marl	Pawpaw clay	Grayson marl	Del Rio clay
ASTEROIDEA:					
Metopaster	..	..	1	..	..
Comptonia	..	..	1	..	..
Pentaceros	..	..	1	..	..
genus, indeterminate..	..	1	2	1	..
OPHIUROIDEA:					
Ophioglypha	..	..	1	..	..
ECHINOIDEA ¹ :					
Holaster	1	..	2	1	..
Hemiaster	2	1?	2	2	..
Stenonia	..	..	..	1	..
Epiaster	..	..	1	..	..
Enallaster	1	..	3	3	..
Goniophorus	1	1	1	1?	1
Goniopygus	..	..	1	..	..
Leiocidaris	..	1	..	..	..
Salenia	..	..	1	..	..
cidarid spines	..	..	2	3	..

¹Most of these have calcitic preservation.

Of these echinoderms the starfishes and brittle stars have so far been found only in the upper formations, Denton and above. The echinoids also seem more abundant and varied in the Upper Washita; *Enallaster* particularly shows a development of several species in the Weno and Pawpaw formations. *Stenonia supernus* (Cragin) is known only from the Grayson marl. Among the small echinoids, Regularia, especially *Salenidae*, predominate. The *Salenidae* have a wide distribution and their various species seem reliable as horizon markers.

CRUSTACEA:

There is a great variety of crabs belonging to many different genera. Of the lobsters a small species of *Hoploparia* is abundant in the Denton marl. Segments of appendages abound in these four formations; claws are found in the Duck Creek and Pawpaw and rarely in the intervening formations. These crustacea are for the most part calcareous and excellently preserved. In addition there are known in the Texas Comanchean, from the Goodland indeterminate limb segments; from the Duck Creek limestone *Callianassa* sp. claws; from the Fort Worth limestone a lobster related to *Homarus* (Dr. Shuler); and from the Weno, two lobsters and an indeterminate claw. A crab and a lobster claw have been reported from the Buda.

TABULATION OF KNOWN CRUSTACEA OF THE WASHITA PYRITE AND LIMONITE FAUNAE

Species	Duck Creek marl	Denton marl	Pawpaw clay	Grayson marl
ASTACURA:				
Cambarus ?	..	1	..	..
Hoploparia	1	1	..	..
appendages, indet.	..	..	3	..
species, indet.	..	..	4	..
ANOMURA:				
Callianassa	..	1	..	..
BRACHYURA:				
species indet.	..	9	2	..

PELECYPODA:

The Duck Creek limonite fauna is marked by the relative poverty of pelecypods and the relative abundance of gastropods. A small species of *Arca*, abundant in the Pawpaw also occurs in the other formations.

The Pawpaw and Grayson each has a considerable assemblage of small pyritic and limonitic pelecypod casts.

TABULATION OF KNOWN PYRITE OR LIMONITE PELECYPODS IN THE
WASHITA PYRITE FAUNAE

Species	Duck Creek marl	Denton marl	Pawpaw clay	Grayson marl
*Arca	1	1	1	1
*Nucula	1	1	1	1
*Leda	..	1	1	..
*Corbula	..	1	1	..
*Plicatula	1	2	2	1
*Barbatia	..	..	..	1
*Isocardia (?)	..	..	..	1
*Remondia (?)	..	..	1	1
*Lima	1	1	2	1

GASTROPODA:

As may be seen in the following table, a fauna of small gastropods is a feature of the pyrite and limonite fossils in these four formations. In the Duck Creek marl these small limonite gastropods are the most conspicuous element of the fauna.

TABULATION OF WASHITA LIMONITE AND PYRITE GASTROPODS

Species	Duck Creek marl	Denton marl	Pawpaw clay	Grayson marl
*Natica	1	1	1	1
*Cinulia	1	1	1	1
*Cerithium	1	..	..	1
*Turritella	1	..	1	2
*Gyrodes	1	..	..	1
*Turbo	1	..	1	1
*Lunatia	2	..	1	..
*Nerinea	2	..	1	..
*Anchura	1	..	..	..

OTHER COMPONENTS:

Small corals (*Trochosmilia*, *Placosmilia*) occur in the Duck Creek and Pawpaw formations, and probably in the Denton and Grayson. There is a notable abundance of small sharks (*Lamna*, *Oxyrhina*, *Ptychodus*) in the Pawpaw, where teeth, vertebrae, plates, skin and bones are found; the same fossils are found less abundantly in the other formations in

question. These sharks also occur in the intervening limestone formations but their abundance in the marls and clays testifies to the favorable conditions for their existence.

In view of the importance which these pyrite faunae may have for correlation purposes, it is considered useful to refer to certain pyrite faunae of approximately the same age known from Europe and Africa. Of these the Vraconian fauna investigated by Pervinquière shows the closest similarities to the Pawpaw fauna here described, notably in the ammonites. The bearings of these fossils on the correlation of the Pawpaw formation have been discussed elsewhere (page 42). It is evident that unless conditions for the pyritization of fossils are worldwide at a given time, it is useless to attempt to make exact correlations of Texas pyrite zones with particular zones abroad. Instead, those pyrite fossils which have limited vertical ranges may, like any other such fossil, be used to mark within narrow limits the age of the zone in which the fossil occurs. A list of similar Texas and foreign pyrite and limonite species is added.

EUROPE AND AFRICA

*European Limonite and Pyrite Faunae*¹

VALANGIAN

Sayn² has described a series of pyritic ammonites from the Valangian of Provence and Dauphiné; this fauna consists prevailingly of species of *Lytoceras*, *Phylloceras*, *Garnieria*, *Neocomites*, *Thurmannia*, *Mortoniceras* and *Leopoldia*.

ALBIAN

In the Haute-Marne, near Montiérender are two gray sand layers with pyritic ammonites, separated by one meter of clayey sand.

Upper Level:

Hoplites deluci
denarius
splendens
auritus
Turrilites catenatus
Hamites rotundus

Inoceramus concentricus
Nucula pectinata

Lower Level:

Desmoceras beudanti
Hoplites deluci
denarius
quercifolius
Douvillerias mamillatum
lyelli
Hamites alternotuberculatus
Belemnopsis minimus

¹Haug: *Traité de Géologie*, pp. 1232-1298.

²G. Sayn, *Les ammonites pyriteuses du sud-est de France*. *Mém. Soc. Géol. France*. No. 23, 1901.

Inoceramus concentricus
salomonis
Plicatula radiola
Nucula pectinata
Arca fibrosa
Trigonia fittoni
Dentalium decussatum
Solarium moniliferum
Cerithium trimonile

ALBIAN.

In the Rhone basin (les Baronnies and le Diois) the Albian is bathyal and contains a level of small pyritic ammonites:

Phylloceras alpinum
Tetragonites timotheanus
Kossmatella chabaudi mühlenbecki
Desmoceras latidorsatum

VRACONIAN

In the Rhone basin, region of Dieulefit, the Vraconian is represented by black marls with pyritic fossils:

Gaudryceras dozei
Turrillites bergeri
tuberculatus
Belemnopsis ultimus

*Cerithium lallerianum*¹
Avellana muratelli

African Pyrite Faunae

ALBIAN

In the province of Constantine Blayac² collected the following species, preserved as ferruginous molds:

Upper Level:

Tetragonites timotheanus
Kossmatella agassiziana
Desmoceras parandieri, etc.

Lower Level:

Phylloceras velledae
Puzosia mayoriana
paronae
Desmoceras beudanti
latidorsatum
Douvilleiceras mamillatum
Turrillites gresslyi

¹J. E. Fallot: Étude géologique sur les étages moyens et supérieurs du terrain crétacé dans le Sud-Est de la France. Ann. des Sciences géol., 1, 268 p., 41 fig., 8 pl., 1885.

²J. Blayac: Le Gault et le Cénomanien du bassin de la Seybouse et des hautes plaines limitrophes (Algérie) C. R. Acad. Sci., CXLIII, 252-5, 1906.

ALBIAN

In the province of Algiers, near Aumale, Peron¹ found marls with ferruginous fossils:

Natica	Phylloceras velledae
Solarium	Pozosia mayoriana
Cerithium	Desmoceras latidorsatum
Nucula	Desmoceras beudanti
Leda	Uhligella dupiniana
Astarte	

CENOMANIAN

The Cenomanian is uniformly represented in Northern Tunis by an alternation, indefinitely repeated, of marls and limestones. According to Pervinquièr,² only the lower part of the stage is fossiliferous. Here a Vraconian fauna is found, with the following species preserved as ferruginous molds:

Phylloceras ellipticum? Kossmat	Turrilites Bergeri Brongniart
—Velledae Mich. var. Seresitensis	—Wiesti? Sharpe
Pervinquièr	—Morrissi Sharpe
—Tanit Pervinquièr	—Scheuchzerianus Bosc.
—decipiens Kossmat	—costatus Lamarck
Lytoceras Flicki Pervinquièr	—Puzosianus d'Orbigny
cf. Marut Stoliczka	—Kerimensis Pervinquièr
—Timotheaum Mayor	Forbesiceras obtectum Sharpe
—cf. Kingianum Kossmat	Saynoceras Gazellae Pervinquièr
Hamites simplex d'Orbigny	Scaphites aequalis Sowerby
—virgulatus Brongniart	—obliquus Sowerby
—armatus Sowerby	—Thomasi Pervinquièr
Baculites baculoides Mantell	Puzosia Paronae Kilian
Puzosia Chirchensis Pervinquièr	Mortoniceras Nicaisei Coquand
—Mayoriana d'Orbigny	—proratum Coquand
Placenticeras Uhligi Choffat	Tunesites Salambo Pervinquièr
—Saadense Th. et. P.	Acanthoceras Brottianum d'Orbigny
Flickia simplex Pervinquièr	—Martimpreyi Coquand
Brancoceras Zrissense Pervinquièr	—Aumalense Coquand
Mortoniceras inflatum Sowerby with its var.	—Suzannae Pervinquièr
subinflata, spinosa and orientalis	Stoliczkaia dispar d'Orbigny
	Belemites (Hibolites) minimus Lister

Pervinquièr³ has described many Cenomanian ammonites from Algeria, some of them ferruginous; the fauna contains *Phylloceras*, *Lytoceras*, *Hamites*, *Scaphites*, *Puzosia*, *Acanthoceras*, *Turrilites*, *Mortoniceras*, and several other common genera, and is similar to that of Tunis.

¹A. Peron: Essai d'une description géologique d'Algérie pour servir de guide aux géologues dans l'Afrique française. Ann. des Sci. Géol., xiv, art. 4, 202 pp., figs. 1883.

²Pervinquièr: Et. pal. tun., Céphalopodes, pp. 417-418.

³Pervinquièr, Sur quelques ammonites du crétacé algérien. Mém. Soc. Géol. France, No. 42, 1910.

Boule, Lemoine and Thevenin¹ cite from Madagascar a series of sandy clays of middle and upper Cenomanian age, containing pyritic fossils. These clays are characterized by *Acanthoceras subvicinale* B. L. & T., *Scaphites aequalis* Sowerby, *Belemnites fibula* Forbes, *Ostrea foisseyi* Lemoine, and contain *Acanthoceras prenodosoides* B. L. & T., *Phylloceras forbesianum* d'Orbigny, *P. diegoi* B. L. & T., etc.

PARALLELS BETWEEN WENO-PAWPAW FOSSILS AND FOSSILS OF THE EUROPEAN AND AFRICAN CENOMANIAN AND VRACONIAN

Texas	Africa (Tunis)	Europe
*Scaphites hilli Adkins and Winton	*S. aequalis Sowerby	S. aequalis Sowerby
*Flickia boesei Adkins	*F. simplex Pervinquièrè	
*Acanthoceras worthense Adkins	*A. martimpreyi Coquand	A. martimpreyi Coquand
*Acanthoceras sp.	*A. suzannae Pervinquièrè	
*Hamites tenawa Adkins and Winton	*H. simplex d'Orbigny	H. simplex d'Orbigny
*Hamites sp.	*H. armatus Sowerby	H. armatus Sowerby
*Turrilites worthensis Adkins and Winton	*T. wiesti Sharpe	
*Baculites comanchensis Adkins	*B. baculoides Mantell	B. baculoides Mantell
*Mortoniceras worthense Adkins	*M. inflatum var. spinosum Pervinquièrè	
*Puzosia sp.	*P. paronae Kilian	
*Lytoceras sp.	L. marut Stoliczka	
*Metopaster hortensae Adkins and Winton		M. parkinsoni Forbes
*Comptonia-wintoni Adkin.		C. comptoni Forbes
Schloenbachia wintoni Adkins		S. rostrata
Corbula wenoensis Adkins		C. bicarinata Noetling
Pinna guadalupae Böse		P. decussata Goldfuss
Pecten texanus Roemer		P. alpinus d'Orbigny
Pecten subalpinus Böse		P. alpinus d'Orbigny
Plicatula subgurgitis Böse		P. gurgitis Pictet
Ostrea carinata ? Lamarck		O. carinata Lamarck and O. frons
Ostrea marcoui Böse		O. syphax Coquand
Gryphea washitaensis Hill		G. vesiculosa Sowerby

¹Boule, Lemoine and Thevenin, Paléontologie de Madagascar, Céphalopodes crétacés des Environs de Diego-Suarez, Ann. de Pal., vol. 1, fasc. 4, pp. 2, 4, 1906.

Helicocryptus mexicanus
Böse

Holectypus limitis Böse

Enallaster bravoensis Böse

Epiaster aguilerae Böse

Hemiaster calvini Clark

H. radiatus Sowerby

H. cenomanensis Guéranger

E. lepidus de Loriol

E. triangularis d'Orbigny

H. latigrunda Peron and
Gauthier

DESCRIPTION OF SPECIES¹

CEPHALOPODA

NAUTILUS TEXANUS Shumard

1860: *Nautilus texanus* Shumard, Trans. Acad. Sci., St. Louis, 1, 1856-60.

1889: *Nautilus texanus* Hill, Geol. Surv. Texas, Bull. 4, p. 21.

1893: *Nautilus texanus* Cragin, Geol. Surv. Texas, 4th Ann. Rept., p. 236.

1895: *Nautilus washitanus* Cragin, Colo. Coll. Stud., 5, p. 67.

1902: *Nautilus texanus* Shattuck, U. S. G. S., Bull. 205, p. 34, pl. XXIII, figs. 1-2,
pl. XXIV, figs. 1-2.

1920: *Nautilus texanus* Adkins and Winton, Univ. Texas Bull. 1945, p. 32, pl. 20,
figs. 1-2.

1920: *Nautilus texanus* Winton and Adkins, Univ. Texas Bull. 1931, pp. 58, 61, 66.

This is the commonest Comanchean Nautilus, and has a known range from the basal Duck Creek limestone (at Denison, Texas) to the Buda limestone (at Austin, Texas). It has a zone of abundance in the top of the Weno limestone, and one in the top of the Fort Worth limestone.

NAUTILUS sp.

The upper Pawpaw clay contains considerable numbers of small Nautili of a size about one-third that of *Nautilus texanus* Shumard. These Nautili are more closely coiled, have a smaller umbilicus, a much thicker and lower volution, and apparently are smooth. The species is not described here because of the poor preservation of the material at hand.

¹Types and figured material are deposited in the Bureau of Economic Geology, Austin, unless otherwise specified in the text. The types at Austin are in a metal locker in the fireproof vault of the University of Texas Library. Most of the photographs were made by Mr. F. Christiansen, Austin, but for some I am indebted to Professor John Davis, Fort Worth.

AMMONITES¹

HAMITES TENAWA Adkins and Winton

1920: *Hamites tenawa* Adkins and Winton, Univ. Texas Bull. 1945, p. 43, pl. 6, fig. 4.

1920: *Hamites tenawa* Winton and Adkins, Univ. Texas Bull. 1931, p. 21.

1920: *Hamites* sp., Winton and Adkins, *ibid.*, p. 69.

HORIZON: Pawpaw formation, clay facies, base.

LOCALITY: About three miles southeast of Haslet, Tarrant County, Texas (type locality); 714, near Fort Worth, Texas; 723, Glen Garden Country Club, near Fort Worth, Texas.

Among several *Hamites* in the Pawpaw clay is a straight equal ribbed species resembling *Hamites simplex*. The species and its suture have been figured elsewhere; the suture has six lobes and six saddles, mainly bifid and slightly dissected.

HAMITES sp. aff. ARMATUS Sowerby

HORIZON: Base of the Pawpaw formation, clay facies.

LOCALITY: 714, near Fort Worth, Texas.

Fragments of a small *Hamites* preserved in hematite and showing on every second or third rib four tubercles, two lateral and two ventral, are found in the basal third of the Pawpaw clay. The suture is rather similar to that of *Hamites armatus* Sowerby and has the following characteristics: Suture consists on each side of three saddles and two lobes, in addition to the siphonal and antisiphonal lobes; suture much more dissected than that of *Hamites tenawa*; siphonal saddle narrow and low, with three rounded subdivisions; siphonal lobe narrow, bifid, each lobule narrow at base, trifid terminally; first lateral saddle longer and much broader than siphonal lobe, flared terminally, bifid, each division twice bifid and laterally

¹In coiled ammonites the external margin of the volution (often keeled) is ventral and the internal, concealed margin is dorsal; the sides of the volution are the flanks, the terminal opening is the aperture. Lobes point backwards, away from the aperture, and saddles forwards, toward the aperture. The siphonal (external, ventral) lobe lies on the ventral mid-line and its angulated lobules point backwards; the antisiphonal lobe lies on the dorsal mid-line and its lobules point backwards. Next to the siphonal lobe and on each side of it is the first lateral saddle, and dorsal to it the first lateral lobe; still more dorsally the second lateral saddle, then the second lateral lobe. Further saddles and lobes are numbered seriatim and are often called adventitive elements. In straight shells (*Hamites*, *Baculites*, *Ptychoceras*, straight portions of *Scaphites*) and in fragments of *Turritiles*, the siphuncle should first be located; the siphonal lobe lies upon it.

dissected; first lateral lobe narrow, laterally dissected, twice bifid terminally; second lateral saddle slightly broader, twice bifid; second lateral lobe lower, simpler, bifid, dissected laterally; third lateral saddle low, broad, rather simple, twice bifid; antisiphonal lobe very simple, rounded, slightly trifid at tip, nearly twice as tall as broad.

ANCYCLOCERAS BENDIREI n. sp.

Pl. 11, fig. 1

1920: *Hamites* sp. B. Winton and Adkins, Univ. Texas Bull. 1931, p. 22.

MEASUREMENTS: Diameter of shell without tubercles, dorsoventral, at small end of fragment 19 mm., at middle of curve, 28 mm., at large end of curve 25 mm.; same diameters, right-left, respectively, 16.5 mm., 22 mm., 24.5 mm. Number of ribs in 5 cm. along straight portion, 7; on curve, 4; average number of dorsal costellae per centimeter, 6.

HORIZON: Base of Weno formation, marl facies.

LOCALITY: 618 (type locality), middle exposure, about 10 feet above the top of the Denton marl, near Fort Worth, Texas. The type, consisting of three fragments, and a fragment of another individual were found here.

DESCRIPTION: Two limbs connected by a curve; of these a greater part of the thicker limb and a part of the thinner limb are missing. There is not enough of the curve present to detect a spiral winding of the coil. The smaller, ascending limb increases moderately in diameter to the curve, where the fragment has its greatest thickness. There are four symmetrically placed rows of prominent, rather slender spines, two ventro-lateral and two mid-lateral. The spines of the two ventro-lateral rows are coarser and are laterally flattened at the base; those of the mid-lateral rows are more slender and are circular at the base. These spines are widely spaced, and a set of the four spines lies on each rib. The ribs are coarse, remote, and in a plane nearly at right angles to the long axis, however, they slant ventrally towards the aperture, and dorsally from the last line of tubercles are continued as obscure elevations which cross the dorsum as narrow, fine ribs. The four rows of tubercles bound three flat longitudinal strips, one mid-ventral and two ventro-lateral; the remainder of the circumference is evenly curved. The cross section of the shell is therefore hexagonal in its ventral half and short oval in its dorsal half. Shell and suture are preserved.

SUTURE: Similar to that of *Ancycloceras lineatus* Gabb¹ and *Hamites quadrinodosus* Jimbo.² Suture consists of medium sized siphonal lobe small antisiphonal lobe, and large, nearly equal, first and second lateral lobes and first and second lateral saddles; the third lateral saddle is smaller. Details of siphonal lobe indistinct; the next four elements are large, slender, spreading, subequal, much dissected, primarily bifid, the subdivi-

Fig. 2. *Ancycloceras bendirei* n. sp., suture of the type individual, camera lucida drawing, x 5. SL—Siphonal lobe; ASL—Antisiphonal lobe.

sions bifid, and the resulting inflections generally minutely trifid, sometimes bifid, at the tips. These four elements are subquadrate and subequal. The third lateral saddle is smaller, laterally dissected and twice bifid terminally. The antisiphonal lobe is about twice as tall as wide, laterally dissected, and trifid terminally, the central prolongation being the longest.

This species is not closely similar to any known to me. It has somewhat the form of *Hamites obstrictus* Jimbo, but differs in having four rows of prominent slender spines, and in the ribs, which are coarser and more widely spaced. Superficially also it resembles in form *Ancycloceras matheronianum* d'Orbigny (Neocomian), but differs from this species in the suture, especially in the size of the siphonal lobe and the amount of dissection of the lateral elements, in the diminished lateral ribs and in the presence of spines throughout the length of the coil instead of tubercles.

HAMULINA WORTHENSIS n. sp.

Pl. 2, figs. 23-26

This fossil has two non-contiguous straight limbs united by a single curve, and furthermore the suture has six lobes in all, like *Hamulina hamus*

¹Gabb, Geol. Surv. Calif., Paleontology. vol. 2, p. 139, pl. xxiii, fig. 18c.

²Jimbo, Pal. Abh., vi, 3, 1894.

(Quenstedt) and *H. quenstedti* Uhlig.¹ The juvenile portion of the shell, absent in the material at hand, may have been a close coil as in *Macroscaphites*, as indicated by the grooved impression on the long limb, but this is not certain, and in addition the suture forbids identification with *Macroscaphites*. The suture and the absence of ribbing prevent its reference to *Ancycloceras*. Böse² describes a similar *Hamulina* sp. from the Vraconian limestone blocks of Camacho, Zacatecas.

MEASUREMENTS:

	I (type)	II	III	IV
Length of fragment-----	7.8 mm.	10.0	8.9	8.1
Width, lower end-----	1.6	1.0	1.5	1.7
Width on bend-----	3.0	2.1	2.2	--

HORIZON: Base of the Pawpaw formation, clay facies.

LOCALITY: 714, near Fort Worth, Texas. The type and two other individuals have been found here.

Three fragments of a straight limbed ammonite, all lacking the aperture and part of the larger limb, were found in the basal Pawpaw clay. In the absence of better material their systematic position can not be settled, and I assign them with hesitation to the genus *Hamulina*. The curve connecting the larger limb is preserved, but this limb made no impression on the inner face of the smaller limb, as in some *Ptychoceras*, and therefore was not in close contact with it; in addition, the visible portion of the curve is open, indicating a distant, possibly short, thick limb. This might suggest *Hamulina* or *Ancycloceras*, which mostly differ in their sutures and their prominent ribbing. Whether there were three limbs as in *Diphychoceras*, is unknown; however, the suture is very different from, and less dissected than that figured by Gabb for his *Diphychoceras laevis* (*laeve*)³ and that species also differs greatly in form from ours.

Two other individuals (Plate 2, figures 23-25) show distinctly the curve and the proximal portion of the shorter, thicker limb. This is free, and is separated from the thinner limb by the space of about half of its thickness. The aperture is not visible. One individual (Plate 2, figures 23, 25) shows also a part of the curve at the lower end of the slender limb. This end turns with about the same curvature as that of the larger curve, and on the venter of the lower end of the slender limb is a wedge-

¹Pervinquière, Ét. pal. tun., Céph., pp. 88-89.

²Böse, On some new Cretaceous Faunas from Mexico, Univ. Texas Bull. (in press).

³Gabb, Pal. Cal., II, pp. 142-5, pl. xxv, fig. 21a-b.

shaped groove about 1 mm. long with the shallowing, pointed end towards the aperture, which indicates that the lower end of this ammonite was a close coil, as in *Ancycloceras*, or a closely apposed limb. These three individuals agree essentially in their sutures.

DESCRIPTION: Only fragments preserved, showing a small limb, the curve connecting with the larger limb, and a portion of this limb. Suture present; shell, aperture and lower portion of slender limb absent. The curve, in the type individual does not quite turn to a position parallel to the small limb; and the two limbs are not in contact, at least at the curve. Dorsal side with a central and two lateral flattened smooth strips (the antisiphonal lobe lies on the former), making the cross-section near the turn a very thick crescentic or kidney bean shape, slightly concave dorsally and broadly convex ventrally. The smaller limb is slender in its younger stages and thicker to a point near the curve; here there is a slight dorsal inflation beyond which are no sutures; farther up, at the curve, there is a slight dorsal constriction. The ventral mid-line bears a small ridge which apparently marks the siphuncle.

SUTURE: The suture is but slightly dissected, and consists of two

Fig. 3. *Hamulina worthensis* n. sp., suture of type individual, camera lucida drawing, x 15.

lateral lobes and three saddles, in addition to the siphonal and anti-siphonal lobes. The siphonal lobe is bifid and has two slender elongate tips; it is about twice as tall as broad. The first and second lateral saddles are nearly equal and similar; each is about half as tall as broad, and is bifid by a shallow, simple lobule. The first lateral lobe is trifid and progressively diminishes in size from the younger to the older sutures. The second lateral lobe is rather smaller than the first, and is bifid. The third lateral saddle is of the same height and about half the width of the other two, and is broadly rounded and unequally bifurcated by an angular notch

lying nearer the antisiphonal lobe and making the more dorsal division smaller than the more lateral one. The antisiphonal lobe is peculiarly narrow and elongated, and has a slightly inflated, bifid tip.

The suture has the same number of elements, six lobes and six saddles, as *Ptychoceras*, and the species seems to have some resemblance to *P. laeve*, var. *hamaimensis* Pervinquire¹ (Lower Gault, Tunis); however, the first two saddles of our species are broader and less dissected; the third saddle is unequally and shallowly bifid; the first lateral lobe is trifid instead of bifid; and the other lobes are bifid. The African species also differs greatly in form from ours, in having two apposed limbs and annular constriction and inflations on the thicker limb. The Texas Upper Cretaceous (Navarro) species *Ptychoceras texanum* (Shumard)² has the two limbs strongly ribbed and closely apposed.

BACULITES COMANCHENSIS n. sp.

Pl. 2, figs. 20-22

1920: *Baculites* sp., Winton and Adkins, Univ. Texas Bull. 1931, pp. 21, 69.

MEASUREMENTS:

	I (type)	II	III
Length -----	12.5	17.3	13.4
Smaller end, major diameter-----	3.3	4.0	3.9
Smaller end, minor diameter-----	2.9	3.4	3.0
Number of ribs in 5 mm.-----	5	1-2	3
Larger end, major diameter-----	4.1	6.1	4.1
Larger end, minor diameter-----	3.4	3.7	3.1
Number of ribs in 5 mm.-----	4	1-2	3

A characteristic Pawpaw baculite lacking the aperture and the coiled portion, will be described here on account of its stratigraphic importance. The straight portion is preserved with the sutures, as a limonite cast.

HORIZON: Pawpaw formation, clay facies, base, rare.

LOCALITIES: Individual II from locality 714, near Fort Wrth, Texas; type individual (Pl. 2, fig. 21) and the others from locality 719, west-facing hill to east of Crowley road, one mile south of the Baptist Seminary, four and a half miles south of Fort Wrth, Texas.

DESCRIPTION: Straight fragments preserved, slightly and uniformly tapering, lateral outlines slightly undulating but on the whole straight; cross-section a short oval; living chamber not preserved, but

¹Et. pal. tun., p. 90, pl. IV, figs. 5, 6a-b.

²Proc. Bost. Soc. Nat. Hist., 8, 1861 (1862), p. 190.

sutures suddenly more crowded at larger end. Ribs consist of evenly rounded, low, annular swellings in a plane nearly perpendicular to the long axis of the shell, more crowded near the enlarged end, almost equally steep on both sides of the annulation, tallest on the venter and on the ventral half of the sides, thence decreasing in height toward the dorsum, where they turn sharply toward the more constricted end of the shell and become obsolete upon crossing the dorsum. Over the more constricted lower end of the shell a suture lies in the lower half of the valley between each two annulations, but in the thicker part of the shell the relation of sutures to annulations is variable; near the thicker end in the type, the sutures are crowded.

SUTURE: Besides the siphonal and antisiphonal lobes, the sutural elements consist of three lateral saddles and two lateral lobes. The

Fig. 4. *Baculites comanchensis* n. sp., sutures of type individual, camera lucida drawing, x 8.

siphonal lobe is bifid, having two slender, rather acute points and a relatively broad, low, crenulate, external saddle; this lobe is nearly twice as tall as broad. The antisiphonal lobe is extremely small and simple; it consists of a single long, narrow, rounded point. The first lateral saddle is about twice as wide as the siphonal lobe and is deeply bifid by a rather simple lobule with one central and two incipient lateral points; each of these portions is again bifid. The first lateral lobe is very tall and bifurcated terminally.¹ The second saddle is bifid and not quite as wide as the first; of its divisions the ventral one is trifold and the dorsal one bifid. The second lobe is broader at the base than the first, and about two-thirds the height; it is bifid terminally. The third saddle is broadly rounded and shallowly bifid. The suture (Fig. 5) is peculiarly simplified and is very characteristic. It is somewhat similar to that of *Baculites vertebralis* Lamarck (Santonian), but differs in having a very simple and reduced antisiphonal lobe and third lateral saddle, and in the greater breadth and simplicity of its first and second saddles. It has even less resemblance to *Baculites baculoides* Mantell (Vraconian), whose saddles

¹If the lobes are trifold the fossil belongs to *Bochianites* Lory; however, the suture agrees better with *Baculites*.

and lobes are narrow, much dissected and constricted basally. I have not had opportunity to compare this species with other Vraconian and Albian species.

TURRILITES BOSQUENSIS n. sp.

Pl. 3, figs. 3, 7

MEASUREMENTS:

Height of spire.....	21 mm.
Diameter of last volution.....	12 mm.
Aperture, height	7.2 mm.
Aperture, breadth	6.1 mm.

HORIZON: Middle of Del Rio clay at top of abundant horizon of *Exogyra arietina* Roemer, in association with *Flickia* ? *bosquensis* Adkins, *Acanthoceras worthense* Adkins, *Pecten subalpinus* (Böse), *Gryphea mucronata* Gabb, and small pyrite fossils. The turrilite apparently is abundant.

LOCALITY: West bank of the South Bosque River, 150 yards south of the bridge on the Speegleville road, tall Del Rio cliff, 5.5 miles west of the courthouse at Waco, Texas (type locality).

DESCRIPTION: Shell conical, sharp spired, spiral angle 37 degrees near tip but spreading slightly on the last volution, sinistral, turreted, five volutions in the type, decreasing rapidly in size towards the apex. In side view the volutions are more angulated and the tubercles relatively more prominent than in any other known Texas turrilite. There are four spiral rows of tubercles, of which three are large and nearly equal and lie on the prominent flank of the volution and the fourth is diminutive and lies beneath the upper margin of the volution. The third row of tubercles lies at the exact margin between the volutions and is therefore covered as is the fourth small row, by the succeeding volution, so that on the earlier volutions only two rows of tubercles are exposed. Of these the lower row, nearest the siphuncle, is separated from the lower exposed margin of the volution by a broad smooth evenly curved spiral strip.

Tubercles of the first and second rows¹ are obliquely placed and are not connected by visible ridges. The third row is closely spaced, the tubercles being more numerous than in the first two rows, and this row

¹Counting from below when the turrilite is placed point down (See Pervinquière, Et. pal. tun., Ceph., p. 428). The siphuncle is then inferior and may be concealed by overlap of the next volution.

is separated from the second row by a sharply excavated spiral groove of only half the width of the space separating the first two rows. Finally, the small tubercles of the fourth row are paired with the tubercles of the third row, the tubercles of a pair being connected by an obsolete radial ridge.

The angularity of the volutions is caused by the lateral projection of the first and second rows of tubercles. In the type individual, the section of the last volution (pl. 3, fig. 7) is elliptical in its basal (dorsal) half, and the inner margin is evenly rounded. The superolateral margin is incised between the three prominent tubercles, and the superior margin in the region of the fourth line of small tubercles is almost straight.

SUTURE: The type individual shows a series of suture stages beginning on the earliest preserved volution. This volution is rounded and

Fig. 5. *Turrilites bosquensis* n. sp., sutures of type individual, camera lucida drawing, x 8. The numbers indicate the volutions on which the corresponding sutures occur.

is totally smooth and tubercles begin to appear only at a diameter of spiral of 2.8 mm. At this point the volution is 1.1 mm. thick and the first and second rows of tubercles appear simultaneously. The suture is simple and consists of a low, undissected siphonal lobe, a simple first lateral saddle, a slightly smaller and still undissected first lateral lobe, a slightly bifid but otherwise simple second saddle, and an undivided

second lateral lobe, of which half is concealed by the next volution. The siphonal lobe of the second volution is dissected, and the two lateral saddles are bifid with simple rounded subdivisions, but the lateral lobes are rounded and undivided. On the third volution the saddles become still more complicated, in general being trifold, but the lateral lobes have little inflection. The siphonal lobe has elongated to its final form but still lacks secondary inflections. In the fourth volution the siphonal lobe is subdivided laterally, and has mature inflections and form. The lateral lobes are slender and trifold, and the lateral saddles are dissected, in general twice trifold. The final volution of the type individual has all elements of the suture dissected (the siphonal lobe was poorly preserved and likely is more dissected than is shown in the drawing). The first saddle is twice as broad as the first lobe and is broadly bifid. The second saddle is bifid and the second lobe in this individual indeterminate.

Number of individuals: about 10.

TURRILITES WORTHENSIS Adkins and Winton

Pl. 3, figs. 1, 6

This common Pawpaw species marks the clay phase in the region from Denton to Johnson County, Texas. It is more abundant in the basal five feet of the formation. The species is rare in the marl transition phase, one individual having been found near Riovista. The species is very rare or missing in the Red River region.

TURRILITES sp.

Pl. 3, figs. 2, 4

There are several dextral and sinistral turrilites, preserved in hematite or pyrite, which characterize the Pawpaw clay, and material is being collected for a further study of their structure. In numbers these turrilites are abundant and are one of the most striking features of the fauna. They likewise are rare except in the clay facies, and have not been found so far, north of Denton County or south of Johnson County, Texas. *Turrilites* similar to these species have been found in the Del Rio clay just above the *Exogyra arietina* horizon (equivalent of Grayson formation) near Waco and Austin, Texas.

SCAPHITES HILLI Adkins and Winton

Pl. 2, figs. 1-12

1920: *Scaphites hilli* Adkins and Winton, Univ Texas Bull. 1945, p. 37, pl. 7, figs. 3-6.1920: *Scaphites hilli* Winton and Adkins, Univ. Texas Bull. 1931, p. 21.1920: *Scaphites* sp. A. Winton and Adkins, *ibid.*, pp. 21, 69.

	I	II	III	IV	V	VI
	Pl. 2,	Pl. 2,	Pl. 2,	Pl. 2,	Pl. 2,	Pl. 2,
	Fig. 1	Fig. 4	Fig. 7	Fig. 8	Fig. 12	Fig. 17
MEASUREMENTS						
Greatest length	9.5	9.5	13.5	11.0	8.0	7.5
Greatest thickness	8.2	4.4	7.3	4.8	7.0	6.5
Diameter of coil	7.8	3.8	6.0	6.5	7.0	6.2
Length of uncoiled portion.....		5.0	8.0	7.5		
Thickness of last turn.....	8.2	4.4	4.5	2.6	2.6	3.6
Thickness of next to last turn....	3.6	3.1	1.5			
Greatest length of umbilicus.....	1.8	1.2	2.1	2.0	1.5	1.2
Number of (ventral) ribs per centimeter, on coil	20	28	20	26	32	26
On uncoiled portion		24	26	30		

HORIZON: Basal half of Pawpaw formation, clay phase. This species has so far not been found outside of Tarrant County, Texas, but it may be expected throughout the extent of the clay phase of the Pawpaw, or farther.

LOCALITY: Most frequent at the type locality one-fourth mile south of the International and Great Northern Railway bridge across Sycamore Creek, Fort Worth, Texas (locality 714), and one-half mile west of the Glen Garden Country Club, Fort Worth, Texas (locality 723).

NUMBER OF INDIVIDUALS: About 25 individuals have been found. These are in the Bureau of Economic Geology, Austin, Texas, and Texas Christian University, Fort Worth, Texas.

TYPE INDIVIDUAL: From locality 714 (described above): deposited in Walker Museum, Chicago.

MODE OF PRESERVATION: Type, dark reddish brown hematite; other individuals light reddish yellow limonite; a few scattered pyrite cubes noted.

DESCRIPTION: This scaphite consists of a coiled portion and an elongate unrolled portion terminating in the aperture, and at their junction there is a prominent dorso-lateral tubercle or geniculation which may be taken as an orienting point in describing the ribs and sutures of this fossil.

The coiled portion consists of an unknown number (3 or 4 ?) volutions, and the umbilicus is deep and, due to the position of the dorso-lateral tubercle, has a kidney-bean shape with the end nearer the aperture somewhat pointed. There is a distinct elongation, which in the absence of the uncoiled portion will orient the shell.

For the most part there are no sutures nearer the aperture than the dorso-lateral tubercle, and the uncoiled portion represents the living chamber, and therefore lacks sutures.

Coiled portion ventricose to subglobular, very obese, subovate in outline, evenly and almost circularly rounded ventrally; dorsally the circular margin is broken by the incurved margin of the dorso-lateral tubercle, resulting in a more or less open crescentic or tear-shaped umbilicus, lying slightly oblique to the axis of the uncoiled portion; greatest transverse diameter of coil ventral to this tubercle; coil slightly constricted just above the tubercle. The uncoiled portion of which the part below the tubercle is the living chamber, is elongate and ornamented with numerous fine ribs which course obliquely from the dorso-lateral margins to the venter, being nearer the aperture at the latter point. The living chamber is inflated at the edges at the dorso-lateral tubercle where it is in contact with the coil, constricted dorsally at a point a little nearer the aperture and thereafter on approaching the aperture is constricted and curved dorsally. Its dorsal side is excavated into a prominent groove, whose width is about half that of the living chamber, and into which the coil fits at the upper end; the groove thus is exposed from the coil to the aperture. The resulting cross section of the living chamber is concave-convex, the short dorsal concave margin passing over the sharply rounded edges of the dorsal groove into the longer lateral and ventral margin. This section varies at different points of the scaphite; near the aperture it is more concave dorsally.

FORM: There is a variation in form, amount of embracing and rate of increase in size of the successive volutions, as is seen also in the intergrades between *Scaphites obliquus* and *Scaphites aequalis*.¹

The type of *Scaphites hilli* is an obese individual with thick, rapidly expanding volutions, which closely approaches the typical *S. aequalis* Sowerby as figured by Pervinquière. This expansion over the last volution of the coil amounts to a doubling of the breadth of the coil, the breadths at the dorso-lateral tubercle and at the point opposite it on the next volution being in the ratio of 8:3.6 in the type. The amount of thickening interiorly is also great, since the umbilicus is deep. The re-

¹Pervinquière, Et Pal. Tun., pp. 118-120.

sulting cross-section of the last turn is thickly crescentic (pl. 2, fig. 1).

The remaining individuals have thick coils but show a less rapid expansion of the last volution. One individual (pl. 2, fig. 7) has preserved most of the uncoiled portion which is a thick crescent with rounded sides and a relatively narrow non-ornamented dorsal groove; the aperture is destroyed. Usually only the coiled portion is preserved, and these individuals are of variable size, even when the dorso-lateral tubercle is also present. They have the same relative thickness and expansion of the coil as the individuals of *S. aequalis* figured by Pervinquière.¹ It is notable that the ribbing becomes obsolete over the lower half of the dorso-lateral tubercle and the entire apertural region.

A straight groove of variable prominence along the ventral midline is a feature of all material found.

RIBS: The ribs are branched or unbranched. They are most numerous at the ventral mid-line and thence laterally some stop short of the dorso-lateral margin and some pass over this margin with a pronounced sigmoidal flexure and describe across the dorsum an arcuate curve, convexed towards the aperture. These principal ribs are thick and elevated laterally. In addition there are shorter ribs which are best developed at the venter and disappear at both ends, or else branch from a principal rib laterally, and crossing the venter disappear laterally, on the opposite side of the coil. There is some irregularity in the distribution of these types of ribs, as follows: In one individual (pl. 2, fig. 7) in the region at the end of the uncoiled portion there are long and short simple ribs and branched ribs but the long simple ribs predominate. In the branched ribs the point of branching is far ventral, lying one-third the distance from the ventral midline to the dorsal midline. Near the dorso-lateral tubercle there are long and short simple ribs with rare incipient branching; the short ribs decrease in length, as their ends lie farther from the umbilicus. The same statements hold for the other individuals figured. In the distal half of the outer volution of the coiled portion a short simple rib alternates with a long simple rib; however, there is incipient branching in the long ribs opposite the dorso-lateral tubercle (pl. 2, fig. 7) sporadically near the tubercle (pl. 2, fig. 4) or even over the whole distal half of the volution (pl. 2, fig. 12). In one individual (pl. 2, fig. 8) there are alternate long and short simple ribs next to the tubercle, but more proximally, there is an irregular alternation of simple and branched ribs. In general, this region shows an alternation of long simple ribs with short simple ones, and incipient or developed branching. This is the situation in the

¹Et. Pal. Tun., plate IV, figures 24-26.

type individual (pl. 2, fig. 1) in which the ribs are at some places evidently branching, and at other places apparently of alternate length.

The remaining visible portion of the coil shows in general an alternation of one branched with one unbranched rib. This alternation is often irregular; in pl. 2, fig. 8 the simple ribs preponderate. The branching may be incipient giving the impression of an alternation of one branched rib with two simple ribs or of one long simple rib alternating with two short simple ribs (pl. 2, fig. 12). In one individual (pl. 2, fig. 12) a branch from a principal rib at one umbilical margin crosses the venter and fuses with the next principal rib, giving a zig-zag ribbing.

Briefly the ribbing is more branched on the coil and simpler on the straight portion. It is seen that the ornamentation of the coil somewhat resembles that of *S. meslei* Grossouvre (Coniacian), which, however, has the principal ribs more prominent and the secondary ribs largely unbranched; this species also unrolls much less rapidly than the type of *S. hilli*. The obesity of the typical *S. hilli* also removes it from *S. obliquus* as does the coarseness of its ribbing; in fact, no individual known approaches *S. obliquus* in either respect. In both respects it is similar to the thick coiled examples of *S. aequalis* Sowerby, from which it differs in the rate of increase of its turns, in the dorso-lateral tubercle and shape of the umbilicus, and in the suture. *S. aequalis* shows the same prominence of the primary ribs near the umbilical margin and the same branching laterally on the coil, as *S. hilli*.

Fig. 6. *Scaphites hilli* Adkins and Winton, suture, type individual, camera lucida drawing, x 12. The fossil is figured on Pl. 2, fig. 1.

SUTURE: Figure 6 shows the suture of the type, so far as can be seen without dismembering the scaphite. This well developed suture has externally besides the siphonal lobe, two conspicuous lobes and near the umbilical wall a third irregular, low, wide lobe. Of these, the first lateral

lobe is narrow, very tall, and trifold, with the central point strongly developed. The second lateral lobe is broad, nearly the same height as the siphonal lobe, its breadth being about six tenths that of the first saddle, and is bifid with each lobule split into rounded tips. The third lobe is twice as broad as tall and is obscurely split into four rounded divisions. The first saddle is large, subquadrate in outline and deeply bifid; each division is split by a lobule into two unequal bifid components. The second saddle is bifid. The third saddle, which lies on the umbilical wall, is wide and low.

Fig. 7. *Scaphites hilli* Adkins and Winton, young individual, showing the last five sutures, camera lucida drawing, x 10. The fossil is figured on Pl. 2, figs. 3-4.

A comparison of the mature suture of the type with juvenile suture stages is represented in figure 7. This individual differs from the type in many ways; it is less obese and its coils increase less rapidly; its ribbing is possibly coarser; its first lateral saddle is shallow and very wide and is split by a shallow lobule, whereas it is tall, narrow and split by a deep lobule in the type of *S. hilli*; the two have the same horizon and range and are apparently connected by intergrades. The situation is somewhat similar to that of *Scaphites aequalis* and *S. obliquus*, and until further material is available, I would refer this form to *S. hilli*. This suture is seen to be simpler and to lack the finer divisions of saddles and lobes seen in figure 6. The lobes on both individuals are, however, bifid.

The internal suture of a small individual whose form and ribbing mainly agree with the last named individual is shown in figure 8. This suture has six lateral and internal saddles and five lobes. The anti-siphonal lobe is trifid, the remaining ones bifid.

Fig. 8. *Scaphites hilli* Adkins and Winton, external and internal suture, camera lucida drawing, x 10.

Very closely related to this species is *Scaphites worthensis* Adkins and Winton (Duck Creek marl) which, however, lacks the ventro-lateral tubercle; its suture also differs in several respects. *Scaphites semicostatus* Roemer and *S. texanus* Roemer, have been described from the Eagleford formation near New Braunfels; *S. vermiculus* Shumard from the Eagleford near Woodlake, Grayson County; and *S. vermicosus* Shumard from the Navarro near Dresden, Navarro County.

ENGONOCERAS SERPENTINUM (Cragin)

Pl. 4, figs. 3, 5-6, 12

1900: *Sphenodiscus belviderensis*, var. *serpentinus* Cragin, Colo. Coll. Stud., viii, p. 31, pl. 2, figs. 4-6.

1903: *Engonoceras serpentinum* Hyatt, U. S. G. S., Mon. XLIV, p. 162, pl. XIX, figs. 7-14; pl. XX, figs. 1-5.

1918: *Engonoceras serpentinum* Stephenson, U. S. G. S., Prof. Paper 120-H, p. 143.

This distinctive *Engonoceras* is frequent in the shales of the Middle Weno formation in Grayson and Cooke counties, Texas. Shell nacreous in the material studied, with fine, close, sigmoid, radial striae; suture simple, as figured by Hyatt (U. S. G. S., Mon. XLIV, pl. XIX, fig. 7); venter rather square, angulated and rounded at edges, zigzag due to alternating marginal tubercles on the two flanks.

HORIZON: Lower and Middle Weno shale and ironstone; Pawpaw shale.

LOCALITIES: Frisco cut, three-fourths mile north of Union Station, Denison, Texas, and the cut of Duck Creek just west of this locality; pit of brickyard, one and three-fourths miles southeast of Gainesville, Texas; Pawpaw bluff on the Red River, northwest of Cedar Mills, Texas.

ENGONOCERAS sp.

Pl. 4, figs. 8-10

The most numerous fossil in the basal Pawpaw clay is *Engonoceras*; this genus is rare in the Duck Creek, Denton, Grayson, and Del Rio pyrite faunae, and its abundance is diagnostic for the Pawpaw clay as so far examined. There is probably more than one species in this formation. The individuals are small (one inch or less), hematitic or pyritic, with sharply etched sutures; flanks smooth and nearly flat; venter straight-edged, angulated, concaved on the midline (so far as observed never acute as in *Metengonoceras*); umbilicus narrow; sutures of varying complexity, mainly very simple, with numerous auxiliary elements.

HORIZON: Pawpaw formation, clay facies, abundant: Tarrant County, numerous localities; Denton County, Johnson County.

Grayson formation: marl facies and middle clay member, rare, Tarrant and Denton counties.

Denton clay: rare, Denison to Blue Mound, north of Fort Worth.

Duck Creek marl: rare, Grayson to Tarrant counties.

FLICKIA BOESEI n. sp.

Pl. 1, figs. 1-3

1920: *Flickia* sp. Winton and Adkins, Univ. Texas Bull. 1931, p. 69.

HORIZON: Pawpaw formation, clay facies, base.

LOCALITY: Type individual, base of Pawpaw formation, just or top of Weno limestone escarpment, one-fourth mile east of Riovista-Waco road and one mile south of Riovista, Texas. One eroded individual, the type, was found here.

MEASUREMENTS: Height of last whorl, 3.5 mm., width, 4.6 mm.; greatest diameter, 10.5 mm.; width of umbilicus, 3.7 mm.

DESCRIPTION: Form discoidal, inflated, volutions thick, with thick, crescentic section, moderately embracing, umbilicus deep, open, showing at least five volutions. Cast smooth and unornamented.

From the section of the end of the coil it is evident that the volution at this age embraces only about half of the preceding one; the volutions are also much broader and lower than in the adult *Flickia simplex*, having at the end of the fifth (?) volution a ratio of breadth to height of 3:2. The flank is broadly convex and passes into the umbilical wall by a sharp

convexity of more than a right angle. The curvature of the venter is nearly elliptical; a keel is lacking, but at the ventral midline of the latest part of the coil there is a perceptible rounded angulation. The dorsal region of the last volution has a more nearly circular curvature than the venter, and the venter of the next younger volution is therefore taller and more nearly circular than the outer volution.

SUTURE: Goniatic, as in *Flickia simplex* Pervinquière, from which it differs only in minor details. All of the sutural elements are less tall and relatively less slender than in the African species. As in that species there are three lateral saddles and three lateral lobes; of these the first and second lateral saddles and most of the second lateral lobe lie upon the flank, while the remainder of the visible suture lies on the steep umbilical wall. The siphonal lobe is bifid with two low, rounded points and a central (external) saddle; the saddle has a greater breadth and is considerably less pointed than the lobules, in fact it has a very broad, even, arcuate curvature. The first lateral saddle is broad, non-angulate, evenly curved and almost as tall as broad; its breadth occupies nearly half that of the flank. The first lateral lobe and the second lateral saddle have almost the same size and shape: each is rounded terminally, having almost a circular curvature, and slightly more constricted basally; and the second lateral saddle has in one instance a very slight suggestion of angularity, like *Neolobites*. The second lateral lobe and the third lateral saddle are low, broad and non-angular. The foregoing description is from the type individual.

Fig. 9. *Flickia boesei* n. sp., type individual, sutures, camera lucida drawing, x 8.

Pervinquière¹ established the genus and species *Flickia simplex* to include a small ammonite with simplified suture remotely like *Neolobites*, from the Vraconnian of Tunis. Comparing the latest sutures of the two types,² we find that the first lateral saddle of the Texas species is somewhat lower and broader than that of *Flickia simplex*; the second one is higher and lies entirely upon the flank, while in the African species it appears to lie partly upon the umbilical wall; the external saddle of

¹Pervinquière: Et. pal. tun., p. 212, pl. 9, figs. 2-5, text figs. 80, 82.

Flickia boesei is lower and more flattened than that of *F. simplex*; and the siphonal lobe is narrower and taller and instead of being inflated,¹ tapers slightly to the end. The Texas type is probably juvenile, since the living chamber is lacking, and since it is more similar to the younger than to the older sutures of *Flickia simplex*.

Such simplified ammonites afford few visible characters for comparison, and therefore separations must be made on the basis of slight superficial differences. Very likely better preserved material will afford a more secure basis for distinguishing the species and for defining its relationships. Provisionally it is placed in *Flickia* on account of its form and suture. Since the species differs only slightly in form, cross-section and suture from *Flickia simplex*, essentially the same arguments as to its generic position hold. It is notable, however, that in the Texas species the resemblance to *Neolobites* suggested by Pervinquière is even more striking than in *Flickia simplex*, especially in the very slight angularity of the saddles.

FLICKIA (?) BOSQUENSIS n. sp.

Pl. 1, fig. 4; Pl. 4, fig. 11

There has recently come to light a peculiar *Flickia*-like ammonite, which although not belonging to this genus as described by Pervinquière seems to stand closer to it than to any other genus, and is here described on account of the possible light that it may throw on the relations of this obscure group. This ammonite has prominent umbilical tubercles and a few widely spaced low, rounded, coarse ribs, and the marginal tubercles are lacking. In these respects it agrees with *Neolobites*, but is thicker, more discoidal, has a more rounded keel, a more open umbilicus, and the suture consists of three saddles and two lobes, the first saddle being much taller and broader than the second. It will be seen that the suture has fewer elements than in *Neolobites*, and in this respect is like *Flickia*. The suture resembles that of *Flickia* in the simplicity, the roundness and the non-angularity of its elements; it differs in having the siphonal lobe broader and its sides more convergent towards the tip, the first lateral lobe shorter than the siphonal lobe, and the first lateral saddle much broader than the siphonal lobe. It differs from *Flickia simplex* and *F. boesei* in having ribs and umbilical tubercles instead of being smooth. The sutures on this ammonite are more numerous and crowded than in either genus mentioned. Due to lack of literature, a more exact generic determination can not be given.

¹Ibid., Pervinquière, p. 214, fig. 82.

HORIZON: Top of lower half of the Del Rio clay, equivalent of Grayson formation (top of zone of *Exogyra arietina*, in association with *Turritites bosquensis*, *Acanthoceras worthense*, *Pecten subalpinus*, *Gryphea mucronata*).

TYPE LOCALITY: West bank of the South Bosque River, 150 yards south of the bridge on the Speegleville road; tall Del Rio cliff; 5.5 miles west of courthouse at Waco, Texas.

DESCRIPTION: Form inflated, section of volution rather tall, flanks and venter broadly rounded, keel absent, dorsum concave with a sharper curvature than the venter; volution covering about one-third of preceding one. Volutions with few widely spaced, very large, smooth, low umbilical tubercles each one-third to one-half the height of the volution; from each of these a single low broad rib runs ventrally, reducing in height and disappearing at the ventro-lateral margin. Ventral marginal tubercles absent. Umbilicus wide. Only three volutions visible in the type (interior damaged). The form is approximately that of *Flickia boesei* except for the presence of ribs and the much more open umbilicus. In the type the crowded sutures end abruptly near the end of the last volution, beyond which the living chamber has several oblique low ribs each terminating in a low, indistinct umbilical tubercle.

SUTURE: In form the suture is rather like that of *Flickia* or of *Neolobites*, with a slight angularity in some places. Three lobes and three saddles visible on each side, including the siphonal lobe. Siphonal lobe

Fig. 10. *Flickia* (?) *bosquensis* n. sp., diagrammatic projection of sutures of type individual, camera lucida drawing, x 5.

subquadrangle in outline, with two equal, simple, rounded lobules and a shallow external saddle. First lateral saddle about one and two-thirds times as broad as the siphonal lobe, more angular on its ventral margin, more evenly curved dorsally. First lateral lobe about half as broad as

first saddle, lower than first saddle or siphonal lobe. Second saddle still lower, about one and a half times as tall as the first lobe. Second lobe nearly as broad as second saddle, and somewhat lower. Most of third saddle visible externally, lying entirely on steep umbilical wall. Further elements of suture concealed by overlap on next inner volution. All sutural elements are simple and entire, as in *Flickia*.

SCHLOENBACHIA WENOENSIS n. sp.

Pl. 1, fig. 14

MEASUREMENTS: (Type individual).

Height of volution, 8.0 mm.

Width of volution, 4.7 mm.

HORIZON: Base of Pawpaw formation, clay phase.

LOCALITY: 723 (type locality); 714, both on Sycamore Creek, near Fort Worth, Texas. The species is rather rare; only a few fragments, most of them crushed, have been found.

The genus *Schloenbachia* Neumayr, in the restricted sense adopted by de Grossouvre, Pervinquièrre and others, includes ammonites with sinuous simple or branched ribs, umbilical and marginal tubercles, and sutures with the first lateral lobe usually trifold. *Mortoniceras* on the contrary usually has a square cross-section, coarse, almost straight, tuberculate ribs and the suture less dissected than in *Schloenbachia*, with the first lateral lobe more or less square and bifid. The type species are *Mortoniceras texanum* (Roemer) and *Schloenbachia varians* (Mantell). A rather rare pyritic species of the Pawpaw clay is here described on account of its stratigraphic importance.

DESCRIPTION: Form discoidal, volutions rather flat and broad moderately embracing, outer volution covering about one-third the width of the preceding one, umbilicus therefore wide and open and, on account of the flatness of the volutions, rather shallow; three coils only exposed in the type (fragment), the two inner ones being practically smooth, the outer one decorated with numerous, closely spaced, low, evenly rounded, sigmoidal, branched and simple ribs, most of them having distinct, sharply elevated umbilical tubercles, which are nearly of even height and evenly spaced around the inner margin of the volution; and near the ventral margin a low overturned, sharp-topped, oblique ridge representing the twin marginal tubercles. On this ridge the more dorsally situated tubercle is represented by a gently curved elevation and the more ventral tubercle is sharper-topped and descends abruptly by a steep ridge and becomes

obsolete at a point near the keel, leaving practically no groove bordering the keel. The ribs, on this ridge, have sigmoidal flexures directed forwards, while at the umbilical margin the ribs are directed backwards. The keel is sharp-topped, triangular in section, and elevated above the level of the lateral tubercles. The section of the volution is rectangular and slightly broader dorsally. At the widest point on the volution its height and breadth are in the ratio of 3:2. The section is thus truncate-cuneiform; the flanks are practically straight, the venter lightly arcuate, and the dorsum has a V-shaped excavation for the reception of the next inner volution.

SCHLOENBACHIA WINTONI n. sp.

Pl. 3, figs. 8-11

1920: *Schloenbachia* sp. M, Adkins and Winton, Univ. Texas Bull. 1945, p. 34, pl. 5, figs. 1-4.

1920: *Schloenbachia* sp. M, Winton and Adkins, Univ. Texas Bull. 1931, p. 22.

MEASUREMENTS: (Type individual).

Greatest height of last volution	-----	34 mm.
Greatest breadth of last volution, excluding tubercles	----	32 mm.
Greatest diameter of shell	-----	104 mm.

HORIZON: Upper Weno formation, shale and limestone facies, and ironstone bands in the blue shale of the Red River region. Known also from south-central Texas.

LOCALITIES: 601, clay-ironstone layers in blue shale of upper part of Weno formation, pit of brickyards, one and three-fourths miles south-east of Gainesville, Texas (type locality); 604, cut of Frisco track, three-fourth mile north of Union Station, Denison, Texas; 612, 618, 714, 715, 716, near Fort Worth, Texas; 720, 721, near Riovista, Texas.

DESCRIPTION: Shell discoidal, volutions thickened, umbilical wall steep, passing onto the flank by a rounded curve. Venter truncate, keel low with a parallel groove on each side. Flanks slightly convex between the ribs. Cross-section of volution subquadrate, slightly wider dorsally than ventrally. Flanks ornamented with ribs, each connected with a ventro-marginal tubercle. These ribs are either simple or bifid. The simple ribs disappear dorsally before reaching the umbilical angle. The bifid ribs end dorsally in a single, very tall umbilical tubercle and ventrally in two less elevated marginal tubercles.

Visible portion of suture consists of siphonal lobe, two lateral saddles

and two lateral lobes. The siphonal lobe is about twice as tall as broad, and laterally is minutely dissected, and has also two or three prominent lateral inflections. The terminal points are slender and slightly incised laterally. The first lateral saddle is nearly twice as broad as the siphonal lobe and is trifold, having the more dorsal division slightly dissected, the broader central division trifold, and the slender ventral division next to the siphonal lobe entire. The deepest incision in the saddle lies between the first two divisions. The first lateral lobe is slender and about two and

Fig. 11. *Schloenbachia wintoni* n. sp., type individual, last suture, camera lucida drawing, x 5. Keel region slightly distorted in type individual.

one-half times as tall as broad. It is bifid terminally and the ventral lobule is larger. Each lobule is trifold. The second saddle is lower and broader than the first lobe, and is bifid terminally; laterally it has an irregular outline. The second lobe is broad and bifid. The internal suture has not been examined.

This species differs from *Schloenbachia nodosa* Böse and *S. trinodosa* Böse in its suture and in the cross-section of the volution; it differs from the common *Schloenbachia* of the Fort Worth limestone (*S. leonensis* of various authors) in many respects, notably in its marginal tubercles.

MORTONICERAS WORTHENSE n. sp.

Pl. 1, figs. 6-10, 18-19, 26

1920: *Mortoniceras* sp. Winton and Adkins, Univ. Texas Bull. 1931, p. 68.

MEASUREMENTS:	I (tpe)	II	III	IV
Fragment, outer whorl; length----	15.0	16.5	14.4	10.8
Small end, height -----	3.5	4.3	3.0	2.4

Small end, width -----	4.0	4.0	4.0	3.1
Large end, height -----	4.5	5.5	5.3	4.0
Large end, width -----	4.3	4.8	4.7	3.6

HORIZON: Basal two-thirds of the Pawpaw formation, clay phase.

LOCALITY: 723 (type locality) ; 714, and other localities near Fort Worth, Texas.

DESCRIPTION: Shell discoidal, volutions slightly embracing, each covering one-fourth or less of the next inner volution, umbilicus consequently broad, volutions strongly ribbed with widely spaced, coarse, short, simple or bifurcated, straight ribs, each with an umbilical tubercle and a pair of coarse marginal tubercles which are connected by a concave topped ridge. These tubercles are abruptly raised above the level of the venter, but the oblique rapidly diminishing rib passes from the tubercle towards the keel and becomes obsolete; the keel is therefore bounded on each side by a narrow valley. Keel prominent, steep sided, triangular in section, taller than the outer ventro-lateral tubercle in the earlier volutions, lower in the later ones. Cross-section of the volutions roughly square, but in a few individuals even broader than tall; to be more detailed, the flanks are inflated, and in the region of a rib bicarinate, corresponding to the umbilical and lateral tubercles; venter tricarinate, corresponding to the keel and the two ventro-lateral tubercles; dorsal midline with a slight excavation, into which the next inner coil fits. Number of volutions six or more. At the 6 mm. stage (about four volutions) the shell is practically without ribs. The ribs then come in as straight, broad, low, wedge-shaped elevations, thicker and higher at the marginal end, where the pair of marginal tubercles is an elongate prominence with a slight tuberculate swelling at each end. Within the next half volution the umbilical tubercles have become prominent, and the marginal tubercles are as tall as the keel, giving to the cross-section of the volution a nearly square aspect, instead of the triangular section which exists in the younger stages, in which the keel is still prominent. The sutures coincidentally increase in complexity.

SUTURE: The siphonal lobe is nearly quadrate. It is bifid and each lobule is relatively simple. The first saddle is broad and is twice bifid. The remainder of the suture is invisible on the flanks of the individuals at hand, which are preserved as pseudomorphs of coarse grained hematite which has replaced all structures and destroyed the details. From the ends of fragments broken along suture lines it is inferred that the first lateral lobe is lower than the first saddle and about two-thirds as broad; terminally it is divided and possibly is trifid. The second saddle is lower

and narrower than the first lobe, and apparently is bifid; it lies at the level of the dorso-marginal tubercles. The second lobe lies on the umbilical wall and is broad and flat, but from the material at hand its details can not be described. Between it and the tall narrow anti-siphonal lobe there is a depressed saddle-like space. This description and figure of sutures are taken from the individual figured on plate 1, figure 6.

Fig. 12. *Mortoniceras worthense* n. sp., suture of individual, Plate 1, Figure 6, camera lucida drawing, x 10.

This species has similarities to *Mortoniceras inflatum* var. *spinosum* Pervinquière¹ (Vraconian of Tunis). One individual of five volutions has two gaps in the ribbing of the last volution; they have no obvious relation to the branching or doubling of ribs. The same feature is noted in the earlier volutions in this species.

ACANTHOCERAS WORTHENSE n. sp.

Pl. 1, figs. 11-13, 15-17, 20-25

1920: *Acanthoceras* sp. Winton and Adkins, Univ. Texas Bull. 1931, p. 69.

MEASUREMENTS:

	Pl. 1, fig. 12 Type (I)	Pl. 1, fig. 13 II
Greatest diameter -----	14.9 mm.	13.8
Height of last whorl -----	6.0 mm.	6.0
Width of last whorl -----	6.0 mm.	5.0
Width of umbilicus -----	1.4 mm. (?)	1.0

HORIZON: Pawpaw formation, basal two-thirds, clay facies.

LOCALITY: 723 (type locality), near Fort Worth, Texas; 714, 715, 716, 719, 724, near Fort Worth, Texas. The species is generally abundant in favorably weathered exposures of the clay phase.

DESCRIPTION: Shell discoidal, compressed, four or more volutions considerably embracing, umbilicus consequently small; shell keelless, sec-

¹Et. pal. tun., p. 229, pl. xi, fig. 3a-b.

tion of volution roughly oval, venter smooth in young individuals and very slightly convex. In older individuals, however, there is a median broad, low, very rounded swelling, similar to a median tubercle. In young individuals the ribs become obsolete just ventral to the ventral marginal tubercle, but in older individuals the ribs are continuous across the venter. Ventro-marginal tubercles acute, one on each rib, becoming obsolete in the later volutions. Ribs long and short, mainly unbranched. Dorso-lateral (umbilical) tubercles few and scattered. Cross-section of earlier whorls subrectangular, flanks straight, venter lightly convex, dorsum very concave; in later whorls the flanks are more rounded and make an even curve with the venter. In certain young individuals the ribs are flexuous, having a prominent sigmoidal curve, and are much thickened at the marginal tubercle. This individual shows no ribs at a diameter less than 3.8 mm. After that diameter, the intercalated ribs, one or two between some branches, are similar to those of the later volutions.

The suture of the species resembles that of young individuals of *A. martimpreyi* Coquand, as figured by Pervinquire.¹ The considerable

Fig. 13. *Acanthoceras worthense* n. sp., suture, camera lucida drawing, x 8.

amount of material at hand does not contain any sutures which are much more complicated than the ones figured here. The suture shows externally besides the siphonal lobe, four saddles and four lobes, which progressively decrease in size toward the anti-siphonal region. The bifid first saddle is more irregular and less square than in the examples of *A. martimpreyi* mentioned. All the remaining saddles are simple, or bifid, and the lobes are generally bifid. The first lobe is as tall as the first

¹Pervinquire, Et. pal. tun., p. 294, figs. 109-110.

saddle and about three-fourths as broad. The second and third saddles are taller than broad and rather angular in form. The remaining lobes likewise are subquadrate and shallowly bifid. The internal suture has not been examined.

This species greatly resembles *Acanthoceras martimpreyi* Coquand, *A. aumalense* Coquand, and *A. suzannae* Pervinquire, all from the Vraconian of Tunis.

ASTEROIDEA

The genera of Comanchean starfishes are in such great need of definition that it has been found impossible to assign the Texas species with much certainty to the proper genera; for in the absence of authentic material of described species it is difficult to make such definitions from figures and text alone. When the Texas starfishes are closely defined it will probably be necessary to form new genera for certain of the species, but in the meantime they have been provisionally assigned to known genera, avoiding when possible those founded on recent species, on account of the great amount of subdivision which the recent *Asteroidea* are receiving at the hands of modern workers.

PENTAGONASTER TEXENSIS Adkins and Winton

Pl. 7, fig. 7

1920: *Pentagonaster texensis* Adkins and Winton, Univ. Texas Bull. 1945, p. 47, pl. 10, figs. 5-6.

1920: *Pentagonaster texensis* Winton and Adkins, Univ. Texas Bull. 1931, p. 22.

A recently discovered individual shows the superomarginal plates, and the interior of the disk is eroded down to the oral surface, of which several adambulacral, circumoral, and a few paramarginal plates are exposed. All of the superomarginal plates are shown, but due to erosion and the thinness of the disk, certain oral plates come into view, and there is over the disk a resulting mixture of oral and aboral plates. This individual was found in a limestone slab, face down, at the type locality described below.

HORIZON: Upper five feet of the Weno limestone.

LOCALITY: 602, east slope of the valley of Sycamore Creek, four miles southeast of Fort Worth, Texas, at a level 29 feet below the base of the Mainstreet limestone.

DESCRIPTION:

SUPERO-MARGINAL PLATES: On each aboral interarc there are fourteen supero-marginal plates excluding the terminal plate, which is common to each two adjacent arcs. These plates, though separated, seem well preserved, and in this individual as in the type, lack ornamentation. It is entirely possible that better material will show superficial structure on the plates since most related species show it. In shape the plates are trapezoidal or cuneiform, as described for the type. They are tallest and widest at the center of the arc and decrease in size to the tips of the rays. The marginal plates of this species are relatively large and are taller than broad; in *Pentagonaster browni* Weller, they are broader than tall, and are relatively smaller.

PARAMARGINAL PLATES: These are a row of minute, nearly equal and similar subquadrate plates lying apposed to the inner edge of the marginal plates. Only a few, near the tips of the rays, are preserved in this individual; they apparently belong to the aboral surface.

ADAMBULACRAL PLATES: These plates, as described in the type, are mostly quadrate and are of about the same size as the other interior oral and aboral plates (about .5 mm. diameter). This individual has along one radius two nearly apposed rows of plates that belong to this class; they diminish in size somewhat toward the tip of the ray.

CIRCUMORAL PLATES: There is a group of plates massed around the mouth region, which are in part oral plates and in part dislocated central aboral plates. They are rounded and larger than the other interior plates, and a few are elongate. In the absence of better material, they can not be described more accurately.

INTERMEDIATE PLATES: There are a few continuous radial rows of plates lying on the radii and extending from the center of the disk to the tips of the rays. These are apparently aboral plates. They have a rounded, roughly quadrate to hexagonal shape, and are small, equal and similar. The madreporite and ambulacral details are not visible.

RELATED SPECIES: *Pentagonaster browni* Weller (Fox Hills formation, Lander, Wyoming) is described as having 16 supero-marginal plates which are relatively smaller and less tall than in this species. The inter-radial arcs of the Wyoming species are also deeper than in our material. *P. lunulatus* Woodward and *P. megaloplax* Sladen (Upper Chalk, Senonian) differ from the Texas species in having punctate or papillose marginal plates, in the concavity of the interarcs, and in the number of marginal plates. *P. robustus* Spencer (Upper Chalk) differs in the same characters and in the meeting of the supero-marginal plates along the midline

of the ray; and *P. obtusus* Forbes (Upper Chalk) is stated to have 18 marginal plates, the superior ones punctate and the two sets alternating.

	I (type)	II
R (average of five) -----	23.5 mm.	15.9 mm.
r (average of five) -----	14.8 mm.	9.9 mm.
R/r -----	1.58	1.6

METOPASTER HORTENSAE Adkins and Winton

Pl. 7, fig. 6

1920: *Metopaster hortensae* Adkins and Winton, Univ. Texas Bull. 1845, p. 46, pl. 10, figs. 2-4.

1920: *Metopaster* sp. Winton and Adkins, Univ. Texas Bull. 1931, p. 69.

1920: *Metopaster hortensae* Winton and Adkins, *ibid.*, p. 21.

This rare species, which characterizes the basal Pawpaw clay, has very tuberculated and spiny plates. The marginals are covered with closely set, low, rounded tubercles, and at places have adherent clumps of small thick spines.

One row of ambulacral pore plates is shown on the type individual. This has two rows of oblique pores, indicating four rows of tube-feet per groove.

As already stated, there are six paired supero- and infero-marginals in addition to the terminal unpaired supero-marginal plate, totalling eight supero-marginals and six infero-marginals.

The terminal marginal plate is similar to that of *Mitraster*, which differs from the Texas form in its cycloid contour and in the shape and ornamentation of its marginal plates.

COMPTONIA WINTONI n. sp

Pl. 7, figs. 4-5

1920: *Comptonia* sp. Adkins and Winton, Univ. Texas Bull. 1945, p. 49, pl. 10, fig. 1.

MEASUREMENTS:

Major radius, about ----- 24 mm.

Minor radius (average) ----- 6 mm.

DESCRIPTION: Disk flat although somewhat collapsed in the type. Rays elongate, tapering, incomplete in the type; probably the ratio R:r is about 4:1. Minor radius about 6 mm. in the type. Supero-marginal and infero-marginal plates numerous, usually paired, disk relatively small and

the interr radial arcs next to the disk paraboloid, sharply instead of broadly curved, and in this respect more similar to *Comptonia comptoni* Forbes than to *C. elegans* Gray. The two rows of supero-marginal plates are apparently nowhere in contact throughout the length of the ray. Oral and aboral sides of disk covered with rather small, unequal, polygonal to rounded raised plates.

ABORAL SIDE: The supero-marginal plates at the center of the arc are slightly wedge-shaped with rounded interior margins; their sides converge slightly, passing out from the center of the disk. The plates decrease in size and are more elongate, on approaching the tip of the ray. Over the central part of the arc seven plates have the breadth of about 10 mm. There are probably about 20 supero-marginal plates in one entire arc (the tips of the rays are absent in the type).

Along the center of each ray is a row of radial plates which are larger than the other central plates but not elevated above them. One elevated rounded plate on each interray bears a small circular pore (gonopore). In one interray there are two such plates. Between these plates are more depressed rows of plates arranged so as to make a pentagon with its corners at the interradii. The central radial rows of plates along the rays meet the middles of the sides of this pentagon. The small, raised, oval madreporite is situated excentrically, and bears 9 or 10 radiating mostly divaricate, V-shaped straight ridges.

ORAL SIDE: The infero-marginal plates are equal in width and in number to the supero-marginal and lie opposite them or nearly so. The oral surface of this starfish is distinctly broader than the aboral surface, so that the line of junction of the two sets of marginals is situated well aboral to the edge of the disk (and this does not seem to be due to crushing). The infero-marginal plates are subquadrate-cuneiform in profile, but are very thick and rounded at the external edge. No para-marginal rows of plates are visible. The other plates of the oral face of the disk, which are poorly preserved, are also not in rows. The five ambulacral grooves are distinct but without perceptible structure in our material. The number of rows of tube feet is unknown. The grooves are prominent, and have near the peristome a breadth of about .6 mm. The marginal and other larger plates are covered by rather fine, low, round, irregularly scattered tubercles. Evidences of spine pits and pedicellariae are absent.

The species differs from *Comptonia elegans* Gray in several noticeable features. The disk is practically flat in our species; it however, shows some evidence of flattening in preservation, since the central plates are considerably and abruptly depressed below the level of the supero-marginal

plates. *Comptonia elegans* on the other hand has a central ridge of very rounded elevated plates running the length of each ray and connecting to form on the disk a central, narrow, elevated ring. In our species the center of the disk is flat and the plates are very little disarranged, since the depression mentioned above has not disturbed the relations of the central plates to each other.

Calliderma and *Nymphaster* differ from our species in many notable features. Both have the aboral interradial areas in most species confined to the disk, so that the supero-and infero-marginal plates meet over most of the length of the ray. This is not true in the Texas species. Both genera named were described from recent species, the genotypes being *Nymphaster protentus* Sladen¹ and *Calliderma emma* Gray.²

This species differs from *Comptonia elegans* Gray in several important respects. The disk is practically flat in our species; it, however, shows some evidence of flattening in preservation, since the central plates are considerably and abruptly depressed below the level of the supero-marginal plates. *Comptonia elegans*, on the other hand, has a central ridge of very rounded elevated plates running the length of each ray and connecting to form on the disk a central narrow elevated ring. In our species the center of the disk is flat and the plates are very little disarranged, as the foregoing description shows.

PENTACEROS AMERICANUS n. sp.

Pl. 7, figs. 1-3

MEASUREMENTS:	R (average of two)-----	45.0 mm.
	r (average of five)-----	10.4 mm.
	R:r -----	4.33:1

A species having the form of *Pentaceros jurassicus* Quenstedt and in most respects agreeing with the described generic characters, is here referred to this genus.

HORIZON: Base of the Pawpaw formation, clay facies.

TYPE LOCALITY: 714, one-fourth mile south of the International and Great Northern Railway bridge across Sycamore Creek, four and one-half miles southeast of Fort Worth, Texas. One individual, the type, was found here.

¹Sladen: Narrative of Challenger Exp., 1885, vol. 1, p. 612; Zool. Chal. Exp., part 11, Report on the Asteroidea, 1889, p. 294.

Fisher, W. K., Starfishes of the Philippine seas and adjacent waters. U. S. N. M. Bull. 100, vol. 3, p. 261, 1919.

²Gray: Proc. Zool. Soc. Lond., part XV, 1847, p. 76; Ann. Mag. Nat. Hist., 1847, vol. XX, p. 198; Synopsis of Species of Starfish, British Museum, 1886, p. 7.

TYPE INDIVIDUAL: A well preserved individual showing both sides of the disk in great detail and having two complete and three fragmentary rays. Museum of Texas Christian University, Fort Worth, Texas.

DESCRIPTION: Marginal plates distinct, subrectangular on edge view, rounded on the face, bearing fine tubercles. Inter-marginalia present in the interarcs and the proximal half of the rays, decreasing in size outwards from the disk. Disk relatively small, arcs sharply rounded, rays long and slender; oral surface flat with numerous small plates; aboral surface with a central raised pentagonal system of plates and a row of radiating plates on the midline of each ray.

ABORAL SIDE: The rays are slender and the interarcs sharply curved. Each interarc contains from tip to tip of the rays 48 supero-marginal plates ($17+7$ in each half arc). Of these all except the terminal seven of each ray are elongate in the direction of the ray, sharply rounded on top and separated from those of the opposite side of the ray by radial plates. The terminal seven plates of each margin of the ray are apposed along the mid-line; the plates are transversely elongate and join by alternate facetting, along a zigzag line; the terminal plate is unpaired and is common to the two rows. The seven terminal plates are swollen and smooth and present a very different appearance from the others. In the type they are smooth and whitish, lacking the iron stain of the rest of the animal. A single row of radials extends from the inner pair of these terminal plates to a polygonal elevation in the center of the disk, the plates increasing in size towards the disk. Between this row of plates and the supero-marginal plates there are two parallel rows of plates, an adradial row of large plates fitting into the spaces between the radialia, and a para-marginal row of small cuboidal plates lying against the marginal plates, there being about three of these plates to one marginal plate.

The interradiar areas on the disk are triangular and are covered with three or four rows of small plates lying close to the marginals; at the apex of each area is a perforated, elevated polygonal plate, the genital plate with its gonopore. The radial ridges running on the mid-line of the ray near the disk are composed of three irregular rows of large rounded ossicles and numerous smaller scattered plates, which were apparently imbedded in a membrane. The central polygonal ring-like elevation at their union bears scattered large and small ossicles. The anus is sub-central and large; the madreporite is excentric, raised, hexagonal in shape, depressed in the center, and has coarse imbricated V-shaped ridges, and straight, simple or branching ridges, radiating from the center.

ORAL SIDE: The oral side is strikingly flattened, being only slightly compressed along the ambulacral grooves. Its prevailing smooth ap-

pearance is due to the smallness and flatness of the numerous plates which make up its area. Infero-marginal plates as described for the supero-marginals, and paired with them; however, there are only six inflated terminal plates separated to the tip of the ray by the ambulacral groove. The common terminal closing plate is restricted to the aboral side. Infero-marginal plates finely granulose.

Interambulacral areas of oral side broadly triangular with numerous crescentic, coarsely granulated small plates in roughly concentric rows. An adambulacral row of plates with combs of spines lies next to the ambulacral grooves. Between these and the marginals is one row of plates over most of the length of the ray, but on approaching the inter-radial areas other rows enter. Peristome central, apparently protected by fine incurved spines.

The species resembles in form *Nymphaster radiatus* Spencer (Lower Chalk, Upper Cenomanian), from which it differs in the shape and ornamentation of its marginal plates. It also has resemblances to *Pentaceros boysii* Forbes and *P. squamatus* Forbes, both from the Upper Chalk of Kent, England (Senonian), but it differs in these in its form and in the ornamentation of the plates.

ECHINOIDEA

LEIOCIDARIS SP.

Cidarids are known from all the Texas Washita formations above the Fort Worth limestone. These are undescribed and their tests and spines unrelated with one exception, *Leiocardaris hemigranosus* Shumard from the Denton formation, of which spines and test have been found in such close proximity that they almost certainly belong to the same animal. Cidarid plates, apparently of a *Leiocardaris*, are abundant in the Quarry limestone group of the Red River region and with them occur great numbers of spines. In addition, in the Pawpaw clay and in the Grayson marls there are several different species of large cidarid spines. In the Main-street limestone well preserved cidarid tests are known.

Cidarid sp. 1, spine: The commonest echinoid spine of the Quarry group has the following characteristics:

The shaft bears about sixteen carinae which are made up of rows of rounded, bead-like tubercles, whose tips are slightly pointed down the shaft. The rows are even topped and depressed. On some spines certain carinae are elevated considerably above others and are longer, more prominent, and have a sharp serrate keel and a triangular cross-section. The spine is a short oval in cross-section. Below the tip of the shaft several, in part alternate, carinae end, and the remaining ones become more elevated

and sharp keeled, giving to the shaft a *Cereus*-like appearance. At the base of the spine there is an inflated sharp topped milled collar; the condyle is constricted and the articulating surface deeply excavated. Locality: 601, near Gainesville, Texas; widely distributed in North Texas.

Cidarid sp. 2, spine: *Cidarid* spines of medium length, the shaft not inflated. The shaft bears about 30 depressed, even topped, equal and equally spaced continuous lines. The base is gently inflated and straight sided down to the collar. The collar is not milled and is rounded, making a wide angle with the shaft. The straight margin of the base below the collar descends to the rim of the circular articulating socket. This spine differs from the preceding in having numerous nonserrate carinae. Locality: 601, pit of brickyards, near Gainesville, Texas.

Cidarid sp. 3, spine: Spines with an inflated, club-like shaft bearing about 10 sharply but unequally elevated, smooth keeled carinae and a prominently constricted neck. The margins of the base above and below the collar are smooth and straight and make at the collar a rounded angle of about 135 degrees. The rim of the circular articulating socket is a ring-like thickening. Cross section of inflated shaft angular and subcircular. Locality: 601, pit of brickyards near Gainesville, Texas, Quarry limestone.

GONIOPYGUS sp.

A species of *Gonipygus* from the basal five feet of the Pawpaw formation clay facies, locality 714, near Fort Worth, Texas, is in the hands of Professor F. L. Whitney for description.

PEDINOPSIS SYMMETRICA (Cragin)

1893: *Dumbea symmetrica* Cragin, Geol. Surv. Texas, 4th Ann., Rept., p. 150, pl. xxiv,¹ fig. 12; pl. xxv, figs. 4-7; pl. xxvii, fig. 1.

1915: *Pedinopsis symmetrica* Clark, U. S. G. S., Mon. LIV, p. 64, pl. xxiii, figs. 1 a-h.

Three individuals of this species found in the basal Weno marl, locality 618, near Fort Worth, Texas, have been placed in the hands of Professor F. L. Whitney for description.

PELTASTES sp.

A poorly preserved individual somewhat similar to a *Peltastes* found in the topmost stratum of the Mainstreet limestone at Denison, Texas, was found at locality 714, near Fort Worth, Texas, in the basal stratum of the Pawpaw clay, and it is referred to this genus. These echinoids will be described by Professor F. L. Whitney.

¹Erroneously cited as pl. xxxiv by Cragin and Clark.

SALENIA sp.

A large flattened *Salenia*, resembling in form *S. mexicana* Schlüter, which is not known to range so high, was found in the basal stratum of the Weno formation, just on top of the *Gryphea washitaensis* shell conglomerate of the Denton marl, by the St. Louis and San Francisco Railway track, two and one-half miles north of Denison, Texas.

The following are some of the features of this individual: There are about 20 pairs of ambulacral tubercles, obliquely situated; the ambulacral lips are considerably wider than the inter-ambulacral lips; the apical system ("cap") is large, the plates ornately scalloped; the sutures are widely and irregularly excavated.

Salenidae have so far been found at the following levels in the Texas Comanchean:

Buda: *S. volana*.

Grayson: *Peltastes* sp.: *Goniophorus* sp.

Mainstreet: _____.

Pawpaw; *Peltastes*? sp., *Salenia volana*.

Weno: *Salenia* sp. aff. *texana*.

Denton: *Goniophorus* sp.

Duck Creek marl: *Goniophorus* sp.

Duck Creek limestone: *Salenia* sp.

Goodland limestone: *Salenia mexicana*, *S. texana*, *Salenia* n. sp.

Walnut: *S. mexicana*; *S. n.* sp.

Glenrose: *Salenia* n. sp.

CYPHOSOMA VOLANUM Cragin

This widespread Upper Washita species is occasional in the Weno formation, marl facies, in association with a considerable echinoid fauna and other fossils, such as *Turritella worthensis*, *Venericardia worthensis*, *Remondia* ? *acuminata* and *Ancycloceras bendirei*. It is known otherwise from the Weno to the Buda limestone.

HOLECTYPUS LIMITIS Böse

This species is distinguished from *Holactypus planatus* Roemer by having the periproct in length about half the radius of the test and situated about halfway between the centrally located peristome and the ambitus, while the *Holactypus planatus* the periproct occupies almost the entire length between the peristome and the ambitus. This species, described from subdivision 5 of Cerro de Muleros (Duck Creek and Fort Worth formations) occurs in the Upper Washita beds of North Texas as

Holectypus planatus does in the Lower Washita and Fredericksburg beds. It is scattering in the Weno and Pawpaw formations but more abundant in the echinoid horizon at the base of the marl phase of the Weno formation, in association with *Enallaster bravoensis*, *E. wenoensis*, *Hemiaster riovistae*, *Epiaster wenoensis*, *Pedinopsis symmetrica* and *Holaster*. It is sparse in the Mainstreet formation and has a zone of abundance near the base of the Grayson marl.

HOLASTER sp. aff. **SIMPLEX** Shumard

HORIZON: Weno formation, marl facies, abundant near base; Pawpaw formation, marl facies, occasional near top.

LOCALITIES: 611, 612, 618, 715, 716, 718, near Fort Worth, Texas; 721, 722, near Riovista, Texas; cut of Missouri, Kansas and Texas Railway, one mile north of Union Station, Denison, Texas, between localities 605 and 606.

The common *Holasters* of the Weno and Pawpaw formations have close similarities with the low phase of *Holaster simplex* Shumard of the Duck Creek and Fort Worth formations, but are specifically distinct. However, it is considered best to defer their description awaiting better material. These echinoids occur as eroded mud-filled, calcite tests and the ambulacral areas and apical system are almost invariably damaged.

Tests ovoid in outline, non-angular, anterior notch shallow, longer than tall, greatest perimeter generally above the base.

Ambulacra flush with surface, postero-laterals in some individuals more divergent than in *H. simplex*; peristome transverse, oval, lying in a depressed space near the ambitus. Periprocot low, ovoid, vertically elongate lying in tall individuals beneath a prominent posterior median projection. Apical system subcentral elongate.

It should be borne in mind that the *Holasters* of the Weno marl are diverse, and that among them *Holaster simplex* may occur, since its known stratigraphic range extends into the upper Fort Worth limestone.

The species of *Holaster* closely related to *H. simplex* Shumard of the Duck Creek and Fort Worth formations, and found in the Weno and Pawpaw formations shows the same variation in form as the lower species; sloping individuals with their greater perimeter at the base, and top-heavy individuals with the greatest perimeter above the base, are common, the latter being most abundant. This species seems to differ from *H. simplex* in the shape of the madreporite which is elongate instead of subquadrate and in details of the apical system. Since such species are not easily definable by form alone, it has been thought advisable not to describe the

Weno species until more material is available. Cragin¹ described three species of *Holaster*, some of which are valid species. *Holaster supernus* Cragin (Grayson marl, Argyle, Texas), a top-heavy echinoid with a circular apical system, is a *Stenonia*. The small low species, *H. nanus*, from the "Vola bed" (Mainstreet?) on the Denison-Bonham road, Choctaw Creek, Grayson County, Texas, is apparently the same low species that is common in the Weno and Pawpaw formations throughout North Texas. The other species, *H. completus*, which was found in the "Denison beds" and the Mainstreet limestone in Grayson County, has a very distinctive apical system, but has the periproct infra-marginal, and is not a *Holaster*. In addition, a very small pyritic *Holaster* whose characters are too imperfectly known to warrant description, occurs in the basal Pawpaw clay.

EPIASTER WENOENSIS n. sp.

Pl. 6, fig. 6

MEASUREMENTS:

Type: Length	54 mm.
Breadth	50 mm.
Height	29 mm.
Apical system to posterior border	32.5 mm.

HORIZON: Weno and Pawpaw formations, most abundant in the basal third of the Weno, marl facies, in association with *Holaster* sp. aff. *simplex* Shumard, *Enallaster bravoensis* Böse, *E. wenoensis* n. sp., *Hemiaster calvini* Clark and *H. sp. aff. bezari* Clark.

LOCALITIES: 618, near International and Great Northern Railway track, three miles southeast of Fort Worth, Texas (type locality). 602, 611, 612, near Fort Worth, Texas; near Riovista and Blum, Texas; three miles north of Gainesville, Texas; Missouri, Kansas and Texas Railway cut one mile north of Union Station, Denison, Texas; and elsewhere generally distributed in the North Texas Weno. Localities 714, 715, 718, 723, and other Pawpaw localities near Fort Worth, Texas, especially in the upper marly portion of the Pawpaw.

DESCRIPTION: Superficially similar to *Hemiaster elegans* Shumard, but smaller, smoother, less inflated, less bulged between the ambulacral grooves, and with straighter and narrower ambulacra. The species resembles *E. aguilerae* Böse, but is smaller and differs in the form and ambulacral details. The Texas Comanchean *Epiasters* may be distinguished

¹Cragin, Geol. Surv. Texas, 4th Ann. Rept., pp. 155-158.

from *Hemiaster calvini* and others by the absence of a conspicuous tubercle between the pores of a pair in the unpaired ambulacrum and usually by the larger size, the slighter depression of the ambulacral grooves, and the shape of the test. It is stated that *Hemiaster* has peripetalous fascioles while *Epiaster* lacks them; but this character is of no value in the usual Texas material, which entirely lacks fascioles. *Hemiaster elegans* and *Epiaster aguilerae* are of large size, while *Hemiaster whitei* and *Epiaster wenoensis* are of medium and small size.

Test: Inflated, rotund, smooth, ambulacra nearly straight, only slightly depressed. Widest point of test just anterior of apical system, tallest point on mid-line posterior to apical system and one-third the distance from it to the ambitus. Sides inflated, rather straight, greatest perimeter at bottom. Aboral side nearly flat, peristome region excavated, with a projecting keel on the posterior lip and the grooves of the antero-laterals and the anterior sulcus slanting into the peristome. Posteriorly on the oral mid-line the test is inflated, the most prominent point being near the posterior end of the test. Ambitus rounded, with only an antero-lateral angulation just back of the point where the antero-lateral ambulacra cross the ambitus; posteriorly the test has a rounded outline. Test narrowed to a pointed, very narrowly truncate posterior end.

Unpaired ambulacrum lies in the shallow anterior sulcus, which is similar in form to the other ambulacral grooves, but slightly narrower. Each one has about thirty-one pairs of short slit-like pores set at a slight angle to each other.

Antero-lateral ambulacra diverge at an angle of about 150 degrees, then turn forwards slightly, being thereafter almost straight to the ambitus. Pore zones similar, with the pores almost equal. Each zone has about fifty pore pairs, the pores being short, slit-like, and not set at an angle to each other. Both pores of the anterior zone and the anterior pores of the posterior zone are equal; the posterior pores of the posterior zone are about one and one-third times as long.

Postero-lateral ambulacra long, straight, diverge at an angle of about 60 degrees. About fifty-four pore pairs in each zone, pores slit-like, similar and nearly equal, the interior ones being, however, shorter than the exterior ones.

Peristomal ambulacral pores. The ambulacra after crossing the ambitus continue as smooth finely granulated strips of large plates with reduced pores, and end at the peristome. However, on approaching the peristome the pores become conspicuous and are borne on elevated knobs with a crater-like opening to one side of the top. These are present in the following Texas species examined: *Epiaster wenoensis* n. sp. (most

prominent), *E. aguilerae* Böse, *Hemiaster elegans* Shumard, *H. whitei* Clark, *H. sp.* (Fredericksburg division), *H. calvini* Clark. The antero-lateral rows are conspicuous, consist of eight to nine cup-like elevations decreasing in size outwards from the peristome, the cavities opening outwards and forwards in the anterior row, outwards and backwards in the posterior row. Each cavity bears two nearly circular ambulacral pores, placed obliquely. The antero-lateral and postero-lateral craters make two V-shaped rows at their junctions back of the peristome. The postero-lateral rows each contain about five similar, smaller craters which are lower and more widely spaced farther from the peristome.

Peristome oval, the long axis transverse, the anterior curvature being even, the posterior margin with a central, bluntly rounded, elevated, forwardly projecting lip.

Periproct rather similar to that of *Epiaster aguilerae* Böse: shape trapezoid with the bottom and top rounded, the two superior angles being sharply rounded and the superior margin but slightly arched; the sides diverge, making the periproct broader at the bottom than at the top; the inferior angles are broadly rounded and the inferior margin likewise, it being nearly the arc of a circle of the same diameter as the periproct.

RELATED SPECIES: *Epiaster wenoensis* n. sp. is smaller than *E. aguilerae* Böse; is more narrowed and pointed posteriorly; is less elevated medially just posterior to the apical system, but farther posteriorly is more straight-topped, the slope to the periproct being relatively abrupt; the notch made in the ambitus by the anterior sulcus is shallower and less pronounced than in *E. aguilerae*; the peristomal ambulacral pores are larger and more pronounced, more numerous and more crowded than in any other Texas species examined; the species is more smooth in appearance and more sparsely tuberculated aborally than *E. aguilerae* Böse or *H. elegans* Shumard.

The relations existing between *Hemiaster*, *Epiaster* and *Macraster* in the Texas Comanchean have not been satisfactorily solved. Hill¹ and Clark² have considered *Macraster texanus* Roemer³ to be synonymous with *Hemiaster elegans* Shumard.⁴ Roemer and Böse⁵ state that this species comes from the Fredericksburg division, but Hill⁶ says that it "makes a well defined horizon near the very top of the immense thickness of lower

¹Hill, Annotated Check List, etc., Geol. Surv. Texas Bull. 4, p. 2, 1889.

²Clark and Twitchell, U. S. G. S. Mon. LIV, p. 88, 1915.

³Roemer, Neues Jahrb. f. Min., Geol. p. Pal., Bd. I, pp. 191-195. Tf. VI, 1888.

⁴Shumard, Pal. Exp. Red River of La. in 1852, p. 210, pl. II, fig. 4a-c, 1853.

⁵Böse, Inst. Geol. Mex. Bol. 25, p. 173, 1910.

⁶Hill, On the Occurrence of *Macraster texanus*, Amer. Nat., XXIII, p. 68, 1889.

marine Cretaceous in Texas" (Buda limestone?) and is "not as Dr. Roemer infers from the specimens which accompanied it to Germany, with the *Exogyra texana* fauna, a statement which has been verified by Mr. George Stolley, the collector."

The distinction between *Hemiaster* and *Epiaster* revolves around the presence or absence of fascioles, and Clark¹ refers *Epiaster elegans* Auctt. to the genus *Hemiaster* "as better material has shown that peripetalous fascioles are present although poorly developed and commonly destroyed on most specimens." There seems no *a priori* reason for doubting this statement until well preserved material demonstrates the contrary.

Epiaster aguilerae Böse,² a distinctive species of the lower Fort Worth limestone, described from subdivision 5 of Cerro de Muleros near El Paso, is here redescribed by comparison with *E. wenoensis* n. sp., an analagous and equally distinct species from the Upper Washita beds of North Texas. In the collections of the Department of Geology, University of Texas, is an unlabeled echinoid, probably an *Epiaster wenoensis*, and marked with a red T, as many of Cragin's types were marked, but whether this is a type, and of what, is unknown. Its dimensions, length 36.5 mm., breadth 34.0 mm., height 17.0 mm., do not agree with those of any *Epiaster* mentioned by Cragin, who described *Epiaster electus* (Travis Peak formation), *E. elegans* var. *praenuntius* (Comanche Peak limestone), *E. hemiasterinus* (Grayson formation), and *E. whitei* Clark (Travis Peak, Comanche Peak and Fort Worth formations). Since Cragin's species are unfigured and the types apparently lost, they are considered invalid species. The present species differs from *Hemiaster whitei* Clark in being much less tuberculate, in having narrow straight ambulacra in shallow grooves instead of wide ambulacra in deep grooves; the form is much wider, more broadly rounded posteriorly, and distinctly lower; the slope anterior to the apical system is steeper. In appearance this species can not be confused with the common *H. whitei* Clark of the North Texas Goodland limestone. *Epiaster hemiasterinus* Cragin is considered by Clark³ as identical with *Hemiaster whitei*, and Cragin admits⁴ the "probable identity" of the two. It is described as being tall, with deep ambulacral grooves.

¹Clark and Twitchell, U. S. G. S. Mon. LIV, p. 88, 1915.

²Böse, Inst. Geol. Mex., Bol. 25, p. 173, pl. 47, figs. 2-4, 67; pl. 48, figs. 1, 2, 4, 1910

³Clark, U. S. G. S., Mon. LIV, p. 89, 1915.

⁴Cragin, Geol. Surv. Texas, 4th Ann. Rept., p. 155, 1893.

EPIASTER AGUILERAE Böse

Pl. 5, fig. 5; Pl. 8, fig. 7

1910: *Epiaster aguilerae* Böse, Inst. Geol. Méx., Bol. 25, p. 173, pl. 47, figs. 2-4, 6-7; pl. 48, figs. 1, 2, 4.

1920: *Epiaster aguilerae* Winton and Adkins, Univ. Texas Bull. 1931, p. 58.

MEASUREMENTS:

	Böse's		
	type	I	II
	mm.	mm.	mm.
Length -----	77.2	66.2	48.0
Width -----	69.1	65.5	43.5
Height -----	39.0	32.0	24.0
Apical system to posterior margin----	43.9	37.0	29.0

Two individuals from the basal Fort Worth limestone near Fort Worth are referred to this species, which was described from Subdivision 5 (Fort Worth formation) of Cerro de Muleros on the basis of one imperfectly preserved individual. The individual figured in this paper is of the same proportions as Böse's individual but about one-seventh smaller. It differs also in having the antero-lateral ambulacra diverge at a greater angle, and in being slightly taller and longer from the apical system back: but the last two features mentioned are due in part to a small amount of end-wise crushing suffered by this fossil. In other respects it agrees with Böse's description. This echinoid is proportionately broader and shorter than the typical Fort Worth limestone, *Hemiaster elegans* Shumard, of the form figured by Adkins and Winton¹ and has a decidedly rounded, non-angular contour as compared with that species. It will be noted also that the ambulacra are narrower, more flexuous (near the ambitus), and sunk in much shallower grooves than in *H. elegans*. The madreporite is smaller than in *H. elegans*² or in *Macraster texanus* Roemer.³ No fascioles are visible. In longitudinal section *H. elegans* is much taller, straight-sided and less evenly rounded posteriorly than *E. aguilerae*, and its inter-ambulacral areas, including the median posterior keel more elevated. In the individual at hand the periproct is situated lower than in material of *H. elegans* collected in the Fort Worth limestone at Fort Worth, but higher than in examples figured by Clark,⁴ and the anterior notch is deeper than in Clark's figures but shallower than in some material collected by me.

¹Univ. Texas Bull. 1945, pl. 8, fig. 3.

²Compare Clark, in Clark and Twitchell: U. S. G. S. Mon. LIV, pl. XLIII.

³Böse: Inst. Geol. Méx. Bol. 25, pl. XLVIII, figs. 1-3, 5.

⁴Clark, *ibid.*, pl. XLII, fig. 1-4.

It is evident that the group of *Hemiaster elegans* of the Fort Worth limestone is in an unsatisfactory taxonomic condition and contains more than one species.

The other individual of *Epiaster aguilerae* is smaller but similar in proportions to the one figured here. Like it also, the widest point is just anterior to the apical system and the contour is rounded; the form of the test is longer and somewhat more pointed posteriorly. The longitudinal section is essentially the same. The periproct is situated higher and is perceptibly taller and less rounded, being pointed at both ends.

HORIZON: Basal Fort Worth limestone.

LOCALITY: Subdivision 5 of Cerro de Muleros; Individual 1, one-half mile east of Texas Christian University, Fort Worth, Texas, locality 406.

Individual 2, one-half mile southeast of Mt. Olivet Cemetery, Fort Worth, Texas, lower 10 feet of Fort Worth limestone.

EPIASTER SUBOBESUS n. sp.

Pl. 11, fig. 3

MEASUREMENTS: Length, 89 mm.; Width, 78 mm.; Height, 52 mm.

HORIZON: Top five feet of Weno formation, marl facies; Pawpaw formation, marl facies.

TYPE LOCALITY: 720, top of Weno limestone, one mile southeast of Riovista, Texas.

DESCRIPTION: Test rotund, elevated, tapering posteriorly to a very narrow tall truncation sloping forwards at the bottom in the top of which the very small periproct is located. Contour same as *E. wenoensis* except more prolonged posteriorly; form straighter sided, much more elevated than that species, apical system proportionately farther forward. Base nearly flat, slightly raised on the posterior mid-line, but much less than in *E. wenoensis*, excavated for the peristome. Ambulacra straight, arranged as in *E. wenoensis*, inter-ambulacral areas notably more elevated. Unpaired ambulacrum with about 60 similar slit-like pore pairs obliquely set. Antero-laterals very long, with 90 or more pore pairs in each zone; all pores equal and similar; postero-laterals with 80 or more pore pairs, equal similar slits. Peristome medium, transversely oval, 6.5x6.0 mm.; periproct extremely small, circular, 3.5 mm. diameter, much smaller and situated relatively lower than in *E. wenoensis*. The description applies to the type individual. This species might be taken as an old individual of *E. wenoensis*, but this is precluded by the differences cited above. The species having this characteristic size and form, is common in North Texas.

ENALLASTER

The genus *Enallaster* ranges from the Neocomian to the Turonian, but is especially abundant in the subtropical Cenomanian seas, where its great development and exuberance of closely related species constitute a special and complicated problem. The littoral deposits of the Mediterranean and the Texas regions contain a sequence of poorly defined *Enallasters*, in which the separate species are probably closely restricted in vertical range and on careful study will prove to have great stratigraphic value; the two areas mentioned have closely similar species.

Enallaster differs from *Holaster* and *Stenonia* in having the ambulacral pores sunk in ambulacral grooves instead of being flush with the surface; it differs from *Hemiaster* and *Epiaster* in the exceptional depth of the anterior unpaired ambulacral groove and in the arrangement and length of the pore zones of the unpaired ambulacrum whose pore pairs are of alternate sizes instead of equal. *Enallasters* are of very diverse form, one group being flattened and short-ovoid, one group depressed but very elongate, and a third group rather tall and of elliptical contour.

As defined by Cotteau, *Enallaster* contains only species in which the anterior unpaired ambulacrum contains alternate long and short pore pairs. *Hemiaster* differs from *Enallaster* in having the pore pairs of the unpaired ambulacrum slit-like and similar, and the pores of a pair separated by a rounded tubercle. The Texas species of *Hemiaster* are diverse and fall into two general groups.

A. Postero-laterals very short, biconvex, apical system placed far back on test; anterior row of antero-laterals with pairs of minute circular pores, form low, with long anterior slope.

H. longisulcus (Adkins and Winton)¹.

H. riovistae Adkins.

B. Postero-laterals long, more nearly straight, apical system nearly central; anterior zone of antero-laterals with larger slit-like pores; form more elevated and inflated.

H. calvini Clark.

H. elegans Shumard.

H. whitei Clark.

H. sp. (Goodland limestone).

H. comanchei Clark.

¹Erroneously referred to the genus *Enallaster*.

ENALLASTER WENOENSIS n. sp.

Pl. 5, figs. 3

MEASUREMENTS:

	I (Type)
Length -----	25.0 mm.
Breadth -----	23.4 mm.
Height -----	14.3 mm.
Length from apical system to posterior border -----	9.0 mm.

HORIZON: Upper five feet of Weno formation, marl facies. Base of Weno: base of Pawpaw clay.

LOCALITY: 720 (type locality), one mile southeast of Riovista, Texas, and one-half mile east of the Waco road; 602, 618, near Fort Worth, Texas.

DESCRIPTION—TEST: Test oval in general outline, broadest point anterior to the center, less elongate and more abruptly truncated posteriorly than *H. riovistae* n. sp.: anterior notch more pronounced than in that species and anterior groove shorter. Apical system located a little less than two-thirds the way back on the mid-line. The outline of the test seen from above is ovate, with angulations; the anterior corners of the test are broadly round to the notch of the anterior sulcus, the posterior corners on the ambitus are abruptly rounded to the excavated periproct area. Oval surface inflated medially, the most prominent point being on the mid-line about two-thirds the way from the anterior end of the test. Aborally the highest point of the test is on the carina midway between the posterior ends of the short postero-lateral ambulacra and the test slopes gently anteriorly from this point, as in *E. bravoensis* Böse, and *H. riovistae*.

Anterior unpaired ambulacrum sunk in the deep anterior sulcus which is relatively shorter and narrower than that of *H. riovistae* n. sp.

Each zone has about 31 pairs of ambulacral pores. About the first twelve pore pairs are short, similar and equal except that they increase in size from the apical system anteriorly. They have each two equal short slits separated by a tubercle. Next come five pore pairs, three long and two short, alternating. Next come two groups of two short pairs each, alternating with long pairs. Thereafter anteriorly the pores are short and similar. Throughout this series in all the short pairs, the pores of a pair are separated by a tubercle, which in all the long pairs is lacking. The type agrees with this description. The narrow ambulacral plates, reaching the mid-line, have each two transverse rows of variable sized tubercles, irregularly arranged.

Anterolateral ambulacra: Diverge from apical system at angle of 180 degrees, then turn forward, their middle parts making, if produced, an angle of about 45 degrees with each other; then turn laterally and cross the ambitus at a point posterior to the anterior corner of the test. The anterior pore zone has 18 to 20 small, circular, closely spaced pore pairs. The posterior zone has about 27 pairs of slit-like pores, the anterior pore being much shorter.

Posterolateral ambulacra: Short, inflated centrally diverge at angle of about 130 degrees from each other. Each zone has about 12 pore pairs, those of the posterior zone being more elongate than those of the anterior zone. In each zone the anterior pore is shorter than the posterior.

Apical system: Four genital plates, their conspicuous thick-lipped perforations making the four corners of a square. Oculars small, perforated.

Peristome: Ovoid, depressed, posterior lip straight, with an elevated median carina behind it.

Periproct: Ovoid, inferior margin extended downward; situated at top of vertically excavated area in posterior truncated area.

This species is most similar to *E. bravoensis* Böse, from which it differs in many particulars. *E. bravoensis* in form resembles *E. texanus* (Roemer) and is the inflated texanus-like species which is widespread in the upper Washita and which has been so frequently mistaken for *E. texanus*. We have examples of it from the Weno to the Grayson formations, and it has been found in the Buda limestone (Whitney). *E. bravoensis* is not so tall as *E. texanus* and is not therefore so inflated as that species; yet it is more rotund than *E. wenoensis* and lacks the angular contour of this species. *E. wenoensis* is more ovate in form and does not have the long posterior narrowing of the test seen in many individuals of *E. bravoensis*. It is abruptly narrowed posteriorly and has a truncate posterior end. There is variation in the figured material of *E. bravoensis* in this regard, and in the absence of a designated type individual it is impossible to say which condition is typical for that species. Böse's plate XLI, figure 5, shows an individual in which the posterior narrowing is slight and the truncation is broad, as in *E. texanus* or *E. mexicanus*; in plate XLII, figure 9, the narrowing is prolonged and the truncation narrow, as in *E. wenoensis*. However, a constant and distinctive feature of *E. wenoensis* is that the posterior end of the test is not merely truncate but is deeply excavated vertically, and in this excavation the periproct lies. In *E. bravoensis* as figured, the apical system is subcentral or at least farther forward than in *E. wenoensis*; the main limits of the antero-laterals therefore diverge at a wider angle (about 80 degrees) in *E. bravoensis*;

and this position of the apical system is to some extent correlated with the elongation of the test. The anterior ambital notch in *E. wenoensis* is always sharper and deeper than in *E. bravoensis*; in the latter species it has the same form as in *H. riovistae* here described, while in this respect *E. wenoensis* and *H. longisulcus* (Adkins and Winton) are closely similar to each other. In *E. wenoensis* there is a conspicuous high point on the median carina halfway between the apical system and the posterior end of the test; in *E. bravoensis* and *H. riovistae* the carina is low and the highest point is variable, often lying near the apical system. *E. bravoensis* in addition has a distinctly angular inferior inflation to the test which in our species is evenly rounded.

ENALLASTER BRAVOENSIS Böse

Pl. 8, fig. 4

1910: *Enallaster bravoensis* Böse, Inst. Geol. Méx., Bol. 25, p. 168, pl. 41, figs. 5-10; pl. 42, figs. 2-12; pl. 43, figs. 1-2, 6-7.

1916: *Enallaster bravoensis* Whitney, Bull. Amer. Pal. No. 26, p. 16, pl. 6, figs. 3-5.

1920: *Enallaster bravoensis* Adkins and Winton, Univ. Texas Bull. 1945, p. 58, pl. 9, fig. 11.

This species is occasional in the Upper Washita formations, Weno to Buda, and in North Texas is still unknown from the lower formations. At Cerro de Muleros it is reported from subdivisions 5, 6, and 8 (Fort Worth to Grayson). This species differs from *Enallaster wenoensis*, which also is truncated and tapering posteriorly, by its more elevated and inflated form, smaller depth of ambulacral grooves, and its taller and differently shaped longitudinal section in which the anterior slope is shorter, more abrupt and more rounded.

ENALLASTER sp. aff. TEXANUS (Roemer)

There are in the Upper Washita formations undefined species similar in form to *Enallaster texanus* (Roemer). These are abundant in the Weno and Grayson marls, and are in need of study and revision.

HEMIASTER CALVINI Clark

Pl. 5, figs. 1, 2, 4; Pl. 6, fig. 3; Pl. 8, fig. 6

This species may be recognized by its tall rounded form and its short, deep ambulacral grooves. It is somewhat variable in form and size, but

the most common variants are here figured.

Form elevated; outline angular, broadly truncated and shallowly excavated posteriorly, with a shallow anterior ambital notch, widest just posterior to the antero-lateral ambulacra. The apical system is a little posterior to the center of the test, the distances in front of and behind it having the ratio 10:8. Ambulacra deeply excavated, the unpaired one being longest; the antero-laterals are longer than the postero-laterals (ratio 9:7); the antero-laterals diverge at an angle of about 105 degrees, the postero-laterals at about 45 degrees. The unpaired ambulacrum is slightly wider than the others. The posterior median keel and the interambulacral areas are elevated, the former being sharp-topped. Peripetalous fasciole visible but irregular. Pores of unpaired ambulacrum short, slit like, oblique, separated by a prominent tubercle. Pores of lateral ambulacra elongate transverse slits which are nearly equal in the forward and the rear zones. Peristome elongated transversely, with a prominent thickened posterior lip; periproct high, vertically elongate, ovate. The longitudinal section is distinctly elevated, posterior slope straight and inclined, anterior slope rounded, the highest point of the test being on the posterior median keel just back of the apical system.

HORIZON: Weno to Buda formations.

The Weno marl contains another species which has deeply excavated short ambulacra, much constricted at each end and flared in the middle. These ambulacra are remarkably short and inflated and have a very characteristic appearance. The whole ambulacral system is placed forwards on the test, the posterior median keel and the interambulacral areas are elevated, the outline is almost circular and non-angular, the form is low, and the anterior median notch is shallow.

This species has some resemblance to the poorly figured and described species *Hemiaster bexari* Clark.

HEMIASTER RIOVISTAE n. sp.

Pl. 6, fig. 4; Pl. 8, figs. 2-3, 5

MEASUREMENTS:	(Type)
Length on mid-line -----	25.5
Width -----	23.5
Height -----	14.0

HORIZON: Top of Weno formation, marl facies.

LOCALITY: 720 (type locality), one mile southeast of Riovista, Texas, and one-half mile east of the Waco road.

DESCRIPTION—Test: This species has nearly the form of *Enallaster bravoensis* Böse, but differs in many respects, notably in having the pore pairs of the unpaired ambulacrum all similar. Test low, elliptical in general outline, notched at the anterior ambitus, rounded laterally, with three distinct angulations, broadly truncate posteriorly. Anterior sulcus long, apical system situated slightly more than two-thirds the length of the mid-line from the notch of the anterior sulcus. Highest point of test on the low posterior median carina about one-fifth the length of the test from the posterior end. From this point the mid-line curves sharply posteriorly to the top of the truncated posterior end. This point is only slightly taller than the edges of the anterior sulcus just in front of the apical system. Slope of test forward from apical system gentle, almost a straight line, similar to that of *E. bravoensis* Böse. The middle is the most inflated point on the test. Inferiorly the most prominent point is on the mid-line three-fourths the way back from the anterior end; the shell therefore is tallest at this point and anteriorly is wedge-shaped, as in *E. bravoensis* Böse. There is an elevated median tuberculated area and the peristome lies in a deep depression of the recurved anterior sulcus and is bounded posteriorly by an overhanging pointed lip. Margins of shell sharply rounded on approaching ambitus, and coarsely tuberculate. Around each tubercle is a circlet of small tubercles, and the intervening spaces recovered by fine granules irregularly scattered. Oval projections of postero-lateral ambulacra consisting of large smooth elongate plates with scattered tubercles and fine granules. Laterally the test bears several fascicle-like bands each consisting of three crowded rows of the smallest sized granules.

Anterior unpaired ambulacrum: Long, shallow, sides nearly straight, making a rather shallow notch at the ambitus. Each pore zone has about 32 pairs of transverse slit-like pores, the pores of a pair being separated by a conspicuously elevated, transversely elongate tubercle. The pores are similar and the ten nearest the apical system show a graduation in size; the others are approximately equal, except near the ambitus, where they are more nearly circular. The pore pairs lie each on an elongate ambulacral plate which reaches the middle. Each plate has a double row of small granules.

Antero-lateral Ambulacra: Diverge at an angle of 180 degrees and turn forward making over the central part of their course an angle of about 47 degrees with each other. They then diverge and cross the ambitus far anteriorly. The anterior pore zone consists of about 22 pores of minute, closely spaced, circular pores, situated for the most part about

the diameter of a pore apart. These are situated in the antero-median quadrant of a quadrate or trapezoidal ambulacral plate which bears two or three large tubercles. The posterior zone consists of about 32 pairs, each having an anterior oval pore and a posterior wedge-shaped slit narrowest at the forward end.

Postero-lateral ambulacra: Short, rather wide and bulging in the center, axes straight diverging at an angle of about 115 degrees from each other. The anterior zone has about 12 pairs of short slits; the posterior about 14 pairs of nearly equal longer slightly oblique slits. All of these pores decrease in size toward each end of the zones.

Apical system: Four genital plates, the right anterior one being the elongated madreporite, the posterior pair slightly farther apart than the anterior pair; four perforated oculars.

Peristome: Transversely oval with a slightly emarginate lip. The lip posteriorly is elevated into a blunt high tip or carina.

Periproct: In shape an oval, transversely situated but with the rounded inferior margin slightly bulged downward; each diameter (on the type) about 2.8 mm., situated high on posterior truncated surface two-thirds the way up.

Aside from generic characters the following differences will assist in separating *H. riovistae* n. sp. from *Enallaster wenoensis* n. sp. *E. wenoensis* has the apical system farther forward; the antero-laterals diverge at a slightly greater angle; the anterior ambital notch is deeper and more sharply incised; the form is broader, less elongate, more nearly circular, distinctly less flat and more constricted posteriorly. The area in which the periproct lies is not merely truncate, but is sharply excavated giving a narrow vertical groove, while in *H. riovistae* the posterior end of the test is broadly truncate and the excavation is broad and shallow; and the pores of the anterior unpaired ambulacrum are in part alternating long and short pairs, instead of being all similar and separated by a tubercle as in *H. riovistae*. The more posterior position of the apical system of *H. riovistae* results in its having a longer anterior sulcus, it being intermediate in this respect between *E. wenoensis* and *H. longisulcus*. The antero-laterals are in many individuals which are here referred to *E. wenoensis* much more sunken than in *H. riovistae*. The only other described species with which *H. riovistae* might be confused is *E. bravoensis* which has long and short pore pairs in the anterior unpaired ambulacrum instead of having all the pore pairs similar. *H. longisulcus* (Adkins and Winton) has these pore pairs similar, but it is at once distinguished by its lower, more elongate form, and by the unusually long anterior sulcus and the consequent posterior position of the apical system.

PELECYPODA

NUCULA NOKONIS n. sp.

Pl. 10, figs. 12-16, 19-20

MEASUREMENTS:

	I
Length -----	17.0 mm.
Height -----	13.4 mm.
Thickness -----	8.3 mm.

HORIZON: Blue shale and red clay-ironstones of upper half of Weno formation, Red River region.

LOCALITY: 601, pit of brickyards, one and three-fourths miles southeast of Gainesville, Texas (type locality); 604, cut of Frisco track, three-fourths mile north of Union Station, Denison, Texas.

DESCRIPTION: Shell inequilateral, sub-equivalved, sub-triangular in contour, with fine radial ribs.

Right valve, exterior: Beak sharply rounded, angle 100 degrees; a straight rounded ridge runs from the umbo to the anterior angle, making with the anterior margin a long low triangle whose other two sides are almost straight portions of the anterior margin. These portions are in length as two to one, the dorsalmost portion being the longer, and make with each other an angle of 15 degrees. The ventralward portion, if extended, would make with the extended ventral side an anterior angle of 60 degrees; the actual angle is sharply rounded. The ventral margin is gently convex downward, and extends backward to the sharply rounded posterior angle (90 degrees). The postero-dorsal margin is straight.

The ornamentation consists of about 47 very flat low radial ribs which are unbranched and continuous from beak to ventral margin. Between these lie fine straight depressions one-sixth to one-eighth the width of a rib. The ribs are crossed by fine scalloped growth lines with the convexities pointed ventrally, giving to the growth lines a fine crenulate appearance. On the anterior area (lunule) the ribs disappear and the growth lines thicken slightly. This area contains a radial anterior ridge and a concurrent posterior depression; the growth lines in crossing these are crenulate. Two deeper concentric constrictions representing stages of growth lie in the ventral half of the valve.

On the shorter posterior area (escutcheon) the growth lines turn dorsally and passing over a radial depression and a radial ridge, converge to the margin.

Right valve, interior: The rounded, posteriorly directed beak is slightly elevated over the posterior margin. This margin is gently convex over

its dorsal three-fifths and bears a taxodont series of nine teeth of which the two dorsalmost and the one ventralmost are shorter and narrower than the others. The antero-ventral two-fifths of the margin is concave and lies just posterior to the truncately elliptical scar of the post-adductor muscle. The posterior end of this margin is the posterior angle, from which the rounded ventral margin runs to the anterior angle. This margin is more sharply rounded near the posterior angle. The antero-dorsal margin is distinctly bulged, and its dorsal two-thirds bears a taxodont series of twenty teeth, which taper in size at each end.

Beneath the umbo is a narrow triangular pit directed downward and anteriorly; and posterior to it is a parallel tooth containing along its length, except at the top, a sulcate depression.

Left valve, exterior: The beak is rounded and forms an angle of about 105 degrees. The posterior area is contained between the straight posterior margin and a curved ridge which is the arc of a circle. In this area the growth lines turn sharply dorsally and converge at the margin. In its center is a radial elevation on whose outer surface are irregular crenulations which are oblique to the growth lines and continuous with those on the main portion of the valve. Lateral to the elevation is a concurrent depression and dorsal to it a small plane area. The posterior angle is 100 degrees. The ventral margin is curved; and more sharply so at the posterior end. The anterior area is a triangular space included between the convex anterior margin and a concurrent but divergent ridge which runs from the beak to the anterior angle. It bears numerous coarse crenulations which are continuous with those over the main body of the valve.

The ornamentation of the valve consists of numerous radial simple flat ribs similar to those of the right valve; and concentric depressions and growth lines. The two valves fit together evenly with a slight ventral gape.

Left valve, interior: This valve is essentially similar to the right valve. Anterior to the beak and ventro-lateral to the taxodont series of teeth is an elongate tooth pointed antero-ventrally and fitting into a socket in the right valve. Its end lies below the tenth tooth anterior to the beak.

Type individual: This description is compiled from three individuals of which the first, showing the exterior of the right and left valves, is to be considered the type individual. It and the other two, as many of the other individuals, were found in the blue marl of the upper half of the Weno formation, 20 feet below Quarry limestone in cut of Frisco track, one mile north of Union Station, Denison, Texas (604).

NUCULA WENOENSIS n. sp.

Pl. 10, figs. 10-11

MEASUREMENTS:	I (type)	II	III
Breadth -----	8.5	10.2	13.0
Height -----	14.9	17.5	18.5
Length -----	20.5	23.7	26.0

HORIZON: Weno formation, blue shale, Red River region, occasional; clay ironstone, occasional.

LOCALITY: 604, Frisco cut, three-fourths mile north of Union Station, Denison, Texas (type locality); 601, pit of brickyard, one and three-fourth miles southeast of Gainesville, Texas.

DESCRIPTION: Shell sub-equivalved, moderately inflated, inequilateral, beaks placed back of the middle of the valve. Ventral margin a long, sub-elliptical curve, sharply angulated posteriorly, more rounded anteriorly. Anterior margin long, slightly convexed, beaks opposite, approximate; posterior margin shorter than anterior, nearly straight, lacking a prominent incision just posterior to the beaks. Anterior areas crescentic, together forming an elongate biconvex strip, posterior areas forming a strip proportionately shorter but equally wide. These are crossed by plain recurved growth lines.

Valves nearly smooth in appearance; ornamentation consists of numerous fine, simple radial striaeiform ridges, crossed by growth lines and growth rings. These ridges on reaching the ventral border form a slightly incised, non-crenulate edge. Hinge taxodont, essentially as in *Nucula nokonis*.

This species differs from *N. nokonis* in its form and ornamentation. The anterior margin is straighter and longer; the posterior margin is straighter and lacks the sharp notch just back of the beaks, having instead a gentle curvature; the posterior "areas" are much more pronounced in side view, and instead of lying flat are elevated at the line of junction of the valves by a posterior projecting carina. The ornamentation at once distinguishes the two species, for in *N. nokonis* the radial ribs are very pronounced and the concentric growth lines on crossing them form a system of prominent squares; the concentric ribs on approaching the anterior border become zigzag or wavy making a conspicuous V-shaped angulation, and in the same region the concentric ribs become crenulate and lamellated. The radial ribs form a prominent scalloped edge at the ventral margin of the valve.

ARCA WASHITAENSIS n. sp.

Pl. 10, fig. 5

MEASUREMENTS:	I (type)	II	III	IV	V
Length -----	9.2	12.6	8.1	6.7	8.0
Height -----	7.0	9.5	5.0	4.9	6.2
Thickness -----	6.6	9.5	5.6	4.6	6.4
Length of hinge line	7.3	9.9	6.8	5.7	7.0

HORIZON: Pawpaw formation, clay facies, widely distributed, Johnson to Denton counties; ironstone and sandstone facies, widely distributed, Denton County to east of Bennington, Oklahoma. A similar, and probably identical species of *Arca* occurs in the Grayson marl and clay at Burleson and Roanoke, Texas; in the Denton clay at Denison, Texas; and in the Duck Creek marl, Grayson to Johnson counties, Texas.

LOCALITIES: 714, near Fort Worth, Texas (type locality), all Pawpaw localities in the Fort Worth region. This fossil is found wherever Pawpaw fossils occur, in Denton, Tarrant and Johnson counties, and practically so in the Red River region. Its zone of abundance is one of the most reliable markers of the Pawpaw formation.

DESCRIPTION: Shell small, valves very inflated, subequal inequilateral, taller posteriorly than anteriorly. Beaks prominent, remote, projecting, rounded, situated slightly anterior to the center of the shell. Antero-dorsal margin short, concaved; posterior margin longer and nearly straight. Anterior margin short, sharply rounded, ventral margin long and nearly elliptical, dorsal margin broadly rounded. Hinge line long and straight and marked by a crenulated thin ridge in casts of the interior. The shell gapes slightly at its posterior end. The valves are ornamented by numerous low, fine, subequal radiating striae, which equal in width the valleys between them; and a few (7 or more) unequal, coarse concentric ridges lying mainly on the ventral half of the valve.

This *Arca* may be distinguished from others described from the Texas Comanchean strata by its form and size. It is a very characteristic pyrite and hematite fossil and is widespread geographically in the Pawpaw formation, and in addition appears to have considerable vertical range, reappearing in the Upper Washita in association with pyritic *Engonoceras*, *Turrilites*, *Hamites*, *Scaphites* and starfishes with each invasion of the clay and ironstone phase of deposition.

GERVILLIOPSIS INVAGINATA (White)

1888: *Dalliconcha invaginata* White, Proc. Acad. Nat. Sci., Phila., p. 35, pl. 2, figs. 4-5.

1889: *Gervilliopsis invaginata* Hill: Geol. Surv. Texas Bull. 4, p. 9.

1896: *Gervilliopsis invaginata* Stanton and Vaughan, Am. Journ. Sci., IV series, vol. 1, vol. 1, pp. 21-26.

1910: *Gervilleia invaginata* Böse, Inst. Geol. Méx., Bol. 25, p. 87.

1920: *Gervilliopsis invaginata* Adkins and Winton, Univ. Texas Bull. 1945, p. 67, pl. 18, fig. 1.

This shell abounds in the lower Weno of Cooke County, where on fresh exposures it is found with the nacreous shell intact. It has a restricted vertical range and is a reliable marker for the basal Weno north of the Brazos River. It has not been reported from South Central Texas.

Stanton and Vaughan, however, record the species from subdivision 6 (Weno-Pawpaw) of Cerro de Muleros.

OSTREA CARINATA? Lamarck

The known distribution of *carinata*-like species in Texas has already been published.¹ Pervinquièr² has figured types of *Ostrea carinata* Lamarck; and Texas material is now being assembled for a critical study and comparison with related European material. Oysters of the group of *O. carinata* and widespread in North, Central and West Texas at different stratigraphic levels, and among them a zone of abundance of the smaller forms in the upper part of the Weno limestone (Quarry group of North Texas) and a scattering representation in the Pawpaw formation.

OSTREA sp. aff. DILUVIANA Linnaeus

There are in the Weno and Pawpaw formations three oysters with zigzag coarse margins and ribbed valves, resembling the group of *Ostrea diluviana*. One of these occurs in the Weno and Mainstreet limestones and is similar to that figured by Hill from the Austin Chalk (U. S. G. S., 21st Ann. Rept., pt. 7, pl. XLV, fig. 2). Another species from the Weno marl is smaller and thinner, and the margins are crenulate (Univ. Texas Bull. 1945, pl. 16, fig. 1). A third unfigured species from the Weno marl has few prominent zigzag ribs of varying height. These species will likely be found to have considerable vertical range.

¹Winton and Adkins, Univ. Texas Bull. 1931, p. 57; Adkins and Winton, Univ. Texas Bull. 1945, p. 59.

²Pervinquièr, Pal. Univ., 3 me sér., 1910, fiches 197-198.

OSTREA (ALECTRYONIA) QUADRIPLICATA Shumard

- 1860: *Ostrea quadriplicata* Shumard, Trans. St. Louis Acad. Sci., vol. 1, p. 608.
1879: *Ostrea quadriplicata* White, 11th Ann. Rept. U. S. G. and G. Surv., Terr., pp. 275-276, pl. 5, fig. 6a.
1920: *Ostrea quadriplicata* Adkins and Winton, Univ. Texas Bull. 1945, p. 60, pl. 16, figs. 6-10.

This species has not been seen above the Mainstreet limestone, where it is rare, or below the Denton marl. It is abundant at the top of the Denton marl, with *Ostrea carinata* and *Gryphea washitaensis*; and at the Weno-Pawpaw contact in the Quarry limestone, with *Ostrea carinata*, *Gryphea washitaensis* and cidarid spines.

This species is not known in South Central Texas, but is abundant near El Paso in subdivision 6 (equivalent of Weno-Pawpaw).

EXOGYRA sp. aff. ARIETINA Roemer

A small species of *Exogyra*, juvenile and young adult stages, was found in the basal Weno at locality 611, near Fort Worth, Texas. The evolution of the *arietina* group leaves large gaps to be filled in; the group is known from the Goodland limestone, upper third, and the Kiamitia marl (*E. spp.* aff. *plexa*, ribbed and non-ribbed), from the Duck Creek limestone (*E. plexa*); the Mainstreet limestone (*E. arietina*) and the Grayson marl (two species aff. *E. arietina*).

PECTEN INCONSPICUUS Cragin

Pl. 11, fig. 4

- 1895: *Pecten inconspicuus* Cragin, Colo. Coll. Stud. 5, 1894 (1895), p. 51.

Cragin's description follows:

"Shell small, thin, subcircular, a trifle higher than long, slightly truncated anteriorly and posteriorly, right valve gently convex, its outer surface smooth except for faint concentric striae and a few remote, subimbricate growth-lines; umbonal angle sharp at apex, nearly a right angle; anterior ear (imperfect in the type) re-entrant below, as indicated by the direction of the striae upon it, outline of posterior ear marking an obtuse angle, its posterior margin rather more than one and a half times as long as its dorsal. Left valve unknown.

"MEASUREMENTS: Height 9.5, length 8.75, convexity of left valve 1 mm.

"OCCURRENCE: On slope of Pawpaw Creek, east of Denison, Texas, in red ochraceous shell-conglomerate of the Pawpaw clays. The associate fossils are *Ostrea quadriplicata*, *Tapes dentonensis*, *Yoldia microdonta*, *Turritella seriatim-granulata*, *Sphenodiscus*, *Turritiles*, etc."

Although Cragin's description is very brief and general, it fits a common Weno species of *Pecten* in many respects; the locality also makes it probable that our Gainesville and Denison material is to be referred to Cragin's species.¹

HORIZON: Ironstone and shale facies, upper two-thirds of the Weno formation. Basal Pawpaw formation, sand-ironstone facies. Very rare south of the Red River region.

LOCALITY: Cut of St. Louis and San Francisco Railway, one mile north of Denison, Texas (Pl. 11, fig. 1) and pit of brickyards, one and three-fourths miles east of Gainesville, Texas. Numerous localities in Cooke and Grayson counties, Texas. In the pit of the Gainesville brickyard there are extensive sheets of nacreous shells consisting largely of *Pecten inconspicuus*, *Gervilliopsis invaginata* and prodissoconchs and later embryonic stages of *Gryphea washitaensis*.

DESCRIPTION: Shell slightly elevated, smooth, inequivalve, inequilateral; ventral margin almost the arc of a circle whose diameter is the antero-posterior dimension of the shell. Antero-dorsal margin almost straight, postero-dorsal margin slightly concave near middle and more elevated by a sharp fold above the ear than the antero-dorsal margin: the two make an angle of about 95 degrees at the umbo, the angle being pointed and very slightly rounded at the tip; the angle made by lines drawn from the extremities of the two margins to the umbo is about a right angle. The umbo projects slightly over the hinge line. Anterior and posterior ears triangular, the latter having a longer base along the margin than the former, and having nearly twice the area. Hinge line straight, making an angle of about 46 degrees with the antero-dorsal margin and an angle of about 42 degrees with the postero-dorsal margin. The terminal angle of the anterior ear is about 95 degrees; of the posterior ear, about 100 degrees. The latter angle is broadly rounded.

The ears have fine parallel sigmoid growth line which bend toward the umbo at the hinge line, but at the dorsal margin bend anteriorly and posteriorly and after ascending these margins, course concentrically across

¹This species has never been figured, so far as I can discover; the type is presumably in the Colorado College museum, but efforts to get information about it have been misspent.

the valve. The growth lines on the valve are fine but unequal; there are also irregular coarser, more widely spaced imbricated rings.

In addition, the right valve and its ears are ornamented with numerous fine striaeiform radial curved elevated costellae which spread fan-wise as they approach the margin; these are irregularly truncated near the ventral margin, and on the ears, by intersection with the coarser radial striae, form a network of small squares.

The left valve is apparently slightly concave and has a similar structure to the right valve.

PECTEN GEORGETOWNENSIS (Kniker)

- 1919: *Neithea georgetownensis* Kniker, Comanchean and Cretaceous Penetridae of Texas. Univ. Texas Bull. 1817, p. 31, pl. VI, figs. 1-3.
 1920: *Pecten georgetownensis* Winton and Adkins, Geology of Tarrant County, Univ. Texas Bull. 1931, pp. 22, 64, 66.
 1920: *Pecten georgetownensis* Adkins and Winton, Univ. Texas Bull. 1945, p. 70, pl. 12, figs. 5-6.

This species may be distinguished by its form and by the radial grooves located on the center of the primary and secondary ribs of the right or both valves. These grooves, however, occur sporadically on other species of *Pecten* of the *P. subalpinus* type. It marks the basal third of the Weno formation, marl facies, and is associated with *Turritella worthensis*, *Ancycloceras bendirei*, *Remondia acuminata* Cragin, *Pedinopsis symmetrica* Cragin, and numerous widely ranging species. It is widespread and abundant.

VENERICARDIA WENOENSIS n. sp.

Pl. 6, fig. 2

MEASUREMENTS (type individual):

Antero-posterior	-----	23 mm.
Dorso-ventral	-----	23 mm.
Right-left	-----	18 mm.

HORIZON AND LOCALITY: Basal Weno formation, marl facies, locality 618, near Fort Worth, Texas (type locality); middle Weno shales, "buff marl," below the main *Gervilliaopsis invaginata* zone, locality 601, near Gainesville, Texas.

DESCRIPTION: This *Venericardia* somewhat resembles *V. alticostata* Conrad var.: of the Eocene, but has the ribs more strongly imbricated

and the ventral margin less nearly circular.¹ I know of no other similar species from the Comanchean.

Shell oblique, elliptical in outline except for the projecting, elevated umbonal region. Beaks greatly recurved. Ventral margin almost an ellipse, sharply curved below the beak, gently curved behind; dorsal-posterior margin nearly straight. The valves have about 24 sharply raised ribs, which bear remote but unevenly spaced, elevated, nodose imbrications. The lateral dimension of the shell is thick, and the species varies somewhat in outline. The hinge and other interior details are unknown.

PROTocardia sp. aff. MULTISTRIATA (Shumard)

Pl. 10, figs. 21-26, 32

This beautiful *Protocardia* occurs as casts and molds in the clay-ironstone of the Gainesville brickyards and shows in nacreous preservation the minutest details of the exterior and the interior of the shell. Most frequently the shell has crumbled into a friable powder and falls to pieces as the rock is broken open. It then leaves in the ironstone excellent molds of the exterior of the shells showing the radial and concentric striations of the exterior and the details of the hinge, and casts of the interior showing details of the dentition, the two muscle scars, pallial line, etc. Frequently the shell has been removed by weathering and the ironstone carries a cavity of its exact shape bounded by the cast and the mold. Rarely in the ironstone but usually in the blue shale, the original shell is preserved intact.

The *Protocardia* group of the Texas Comanchean has not been satisfactorily studied and its species are still poorly defined. As in many other Texas genera this uncertainty can be resolved only by a critical description and photographic illustration of the types. Here as in other genera where new species might have been founded, this has not been done pending a better understanding of the relations of the species already described.

Shell inequivalve, inequilateral, biconvex, ornamented with sixteen or more serrate radial lines and about 75 round topped subequal concentric ribs separated by narrow valleys. The dorsal margin of the shell is arcuate interiorly, and anteriorly is more thickened. It bears several imbricated lamellae parallel to and below the hinge line. The region under the beak is deeply excavated.

¹Harris, The Midway Stage, Bull. Amer. Pal., 4, pl. 4, fig. 12, 1896; and, The Lignitic Stage, Bull. Amer. Pal., 9, pl. 11, fig. 1, 1897.

HORIZON: Middle and Upper Weno shale and clay-ironstone.

Locality of figured material: 604, cut of St. Louis and San Francisco track, three-fourths mile north of the Union Station, Denison, Texas; the species occurs throughout the Red River region, and is abundant at locality 601, pit of brickyards, near Gainesville, Texas.

CORBULA WENOENSIS n. sp.

Pl. 10, figs. 1-4

MEASUREMENTS:	I	II	III
Length -----	10.6	11.6	11.2
Breadth (one valve) -----	3.8	4.4	4.0
Height -----	7.7	8.1	8.0
Length of rostrum -----	2.0	2.5	2.5

HORIZON: Weno formation, upper half, clay and ironstone phase, abundant; Pawpaw clay, rare. Abundant in the Red River region.

LOCALITIES: 604, near Denison, Texas (type locality); 601, near Gainesville, Texas.

Shell small, inequivalve, concentrically ribbed with a truncated rostrum.

Right valve, exterior: Form a very irregular quadrilateral; size small. Viewed directly from above, the valve is roughly pear shaped and consists of a very inflated rotund body and a narrowed and depressed posterior siphon tube or rostrum. The umbonal margin is roughly the arc of a circle whose diameter equals the height of the shell. This border passes by a sharp-pointed curve of 120 degrees into the anterior border, which is almost a straight line making an angle of about 45 degrees with the dorso-ventral axis of the valve. The anterior border passes into the ventral border by a sharply rounded curve which is approximately the arc of a circle one-half the diameter of the dorso-ventral dimension (height) of the shell; the anterior and ventral borders if extended would meet at an angle of about 55 degrees. The ventral border is very slightly convex and this approximates a straight line through one-half the greatest length of the shell, where it is perpendicular to the dorso-ventral axis. A little anterior to its middle there is slight concavity which involves the lower two or three ribs of the shell. The ventral border now curves upward and posteriorly, reaching the narrowed siphon tube.

The snout-shaped siphon tube lies opposite the middle one-third of the height of the shell and is roughly a right angled isosceles triangle whose hypotenuse is the line of attachment to the body of the valve. This line of attachment is a narrow constricted or depressed zone which notches

both the ventral and the dorsal margins of the valve and gives to the rostrum an inflated appearance; the notch in the dorsal margin is the deeper and more pronounced one. Posterior to the depressed zone and running slightly divergent to it is a rounded ridge which assumes a conical inflation as it approaches the postero-ventral extremity of the rostrum; on the rostrum dorsal to this ridge is a parallel narrow depression and beyond it a rounded inflation which forms the postero-dorsal extremity of the rostrum.

The ornamentation consists of thickened, rounded, concentric ribbings, which are broadly flattened on their tops and descend by almost right-angled sides to small plane-bottomed valleys lying between them. On the dorso-ventral axis the widths of ribbings and valleys are about as 2 to 1. The ribs and valleys decrease in width at the anterior margin and at the constriction of the siphonal tube.

The ventral one-half of the shell contains 12 rounded ribbings; the dorsal one-half contains 17 rounded ribbings of similar form but of constantly decreasing size, and in addition in the umbonal region 25-30 fainter ribbings which resemble growth lines.

These 29 heavier ribbings course posteriorly making broad downward convexities to the region of the siphon tube constriction; their posterior ends lie in the depressed zone mentioned. They then rise onto the elevated ridge which lies near the anterior end of the rostrum, and course over the inflated rostrum to its dorsal margin. On this elevated ridge the component growth lines in the ribbings appear as slight thin imbricated lamellae. Not all of the ribbings run continuously from the anterior to the posterior margin. Towards the anterior margin some of the ribbings, especially the dorsal ones fuse by twos into broader and more depressed flat bands. The ribbings on a whole are more flattened and more consolidated on the anterior margin, and more resolved into growth lines on the posterior margin, in the neighborhood of the siphon-tube. In one ribbing, splitting is seen over the middle half of the shell. In one place a small auxiliary ribbing between two of usual size, is seen over the middle half of the shell.

The right valve is somewhat larger than the opening into which the left valve fits, so that the arched, rotund margins of the right valve enclose an elliptical basin shaped visceral space upon which the reduced left valve lies.

Right valve, interior: Viewed directly from above, the cavity of this valve has an elliptical contour except at the siphonal end where the margins of the ellipse are drawn out so that each margin has an inward facing convexity; the dorsal margin then slants ventrally and posteriorly to meet

the ventral margin. There is also a deep rounded concealed recess under the umbo, from this viewpoint. The anterior adductor muscle scar is a small subcircular basin-like depression with a raised rim whose dorsal border connects with the pallial line, a faint narrow groove which approaches the ventral margin as it courses towards the post adductor scar. This scar occupies a rounded-triangular elevated area lying between two more elevated ridges.

The more anterior ridge is a shelf-like lamina which separates the body cavity proper from the siphonal (rostral) extension of the shell; and as it passes umbonally it unites with the more posterior ridge forming a U-shaped junction and enclosing between the two ridges the scar of the adductor muscle. The united ridge continues anteriorly as the ventral edge of a broad excavated pit or socket. The external (prismatic) layer of the valve forms a distinct umbonal imbrication which makes up the dorsal edge of this socket. This inner edge of the external layer passes anteriorly, including the umbo but excluding the tooth against which it is apposed; and proceeds to encircle and parallel the anterior and ventral margins of the valve as far as the ventral border of the rostrum, making an imbricated impression about midway between the pallial line and the ventral margin. The thick external portion is cancellated and penetrated by irregular concentric airspaces formed by the loose apposition of long lamellae. The external prismatic portion makes up .8 of the thickness of the shell; the inner portion is porcellanous and in places pearly-iridescent.

Rostrum: The inner layer emerges from the body cavity proper onto the rostrum, where it is thrown into two small elevated folds which cross each other almost at right angles and consequently form four depressed spaces sloping away from them at either side. The shorter fold is in the antero-posterior axes, and the longer one is continuous with the umbonal imbrication mentioned above.

DISCUSSION: The description of the unfigured Cragin species *C. crassicosata*, agrees with this species in some particulars, but disagrees especially in the number of concentric ribs. In Cragin's Kansas material this number is "7 or 8 on the basal half of a shell the same number of millimeters high," while in our material the number is 11 to 12. Cragin states that the Kansas and the Denison specimens show "no differences of specific value," but further information of a deciding nature is not forthcoming.

Cragin's description of the species, *Corbula crassicosata*, is as follows:¹

¹Cragin, Colo. Coll. Stud., 5, 1895, p. 61.

"*Corbula crassicostata*, sp. nov.

"Shell triangular-ovate, gibbous; nearly as broad as high, shorts; gaping posteriorly by a short, conically inflated, gently truncated rostrum, which is placed high above the base of the shell; unbones placed in advance of the middle, that of the right valve only moderately high arched, its summit obtuse; surface ornamented with very coarse, flattish-topped, concentric ribs, separated by abrupt, deep narrow intervals. There are seven or eight of the ribs on the basal half of a right valve the same number of millimeters high.

"Measurements: Height 7.5, length 10, breadth about 7 mm.

"Occurrence: In arenaceous limestone bands of the Kiowa shales at Belvidere, Kansas; in Nos. 2-4 of the writer's 'Belvidere Section.'

"So far as the writer can judge from material now in hand, the similar *Corbula* that abounds in the condition of casts and molds in Texas, presents no differences of specific value from the Kansas shell above described. The casts show that the pallial line is very sharply impressed."

CORBULA BASINIFORMIS n. sp.

Pl. 9, figs. 7-24; Pl. 10, figs. 7-9

MEASUREMENTS: Type—

Length	17.4 mm.
Height	12.1 mm.
Breadth	7.7 mm.
Difference in length of valves	2.4 mm.

HORIZON: Blue shale and clay-ironstone layers of upper half of Weno formation, Red River region, abundant; Pawpaw clay, occasional.

LOCALITIES: 604, cut of Frisco track, three-fourths mile north of Union station, Denison, Texas (type locality); 601, pit of brickyards one and three-fourths miles southeast of Gainesville, Texas; numerous other localities in Cooke and Grayson counties, Texas. Rare, as limonite stained casts in the Fort Worth region.

DESCRIPTION: Shell small, inequivalve, almost smooth, with a short pointed rostrum.

Right valve, exterior: Valve rather inflated, evenly rounded, its outline almost a truncated ellipse. As seen from directly above, the umbonal angle is about 120 degrees; the dorso-anterior margin is thereafter almost straight for about one-third the greatest length of shell; the anterior margin then describes an evenly rounded curve, a portion of an ellipse, which continues to the middle of the ventral margin; here the margin is more nearly straight, but again curves more sharply dorsally and posteriorly, and reaches the posterior (rostral) angle. This angle is 90 degrees and is very slightly rounded-acuminate. The postero-dorsal margin is almost

straight, having a roughened, striate protruding area—the rostrum—reaching to the umbo; and anterior to it is the straight posterior margin of the valve.

The greatest height of the valve is near its center. Its convexity is greater on the umbonal margin than on the ventral margin; from the center the surface slopes gently to a line, marked in places by a prominent growth line, below which the surface is curved at a sharply rounded angle of 90 degrees to the margins of the valve, giving to the valve a characteristic geniculated and basin-like appearance. The rim of the valve thus slopes inward at all points except on the dorsal margin of the rostrum, so that the margin of the valve has a smaller circumference than the prominent growth line mentioned above.

The ornamentation consists of radial striae and concentric growth lines. The radial striae are fine, unbranched, wavy, or punctate lines which end near the geniculation. The concentric growth lines are more conspicuous (toward the ventral margin they resemble fine ribs), and pass posteriorly, turning sharply dorsally at the line of junction of the main body of the valve and the slightly developed rostrum. They then pass dorsally in straight lines which give to the elongate triangular rostrum a roughened striated appearance.

Right valve, interior: As viewed directly from above the interior of the right valve has three prominent contours: the outer one is the margin of the valve; the middle one is a narrow groove, the junction between the thicker outer (prismatic) shell layer and the thinner inner (porcellaneous) shell layer; the innermost contour is the limit of the main body cavity. The first two are the shape of a long ellipse, and are situated a short distance apart; they begin at the antero-dorsal one-third of the prominent tooth, and run concentrically around the anterior and ventral margins of the valve, making an evenly rounded elliptical curve to the postero-ventral extremity of the rostrum. The innermost contour encloses roughly a subquadrate area whose boundary begins at the ventral margin of the tooth and describes a slight curve to the dorsal end of the anterior adductor scar. Here it runs almost straight ventrally with a slight posteriorward convexity formed by the scar, in such a manner that a sharp-edged, crescent shaped shelf, elevated above the main body cavity, is formed between it and the middle contour mentioned. The scar is pear-shaped and its inflated ventral end lies at the widest point of the shelf. The shelf continues along the interior of the ventral side of the visceral space to a point one-third the way from the anterior end, and then fuses with the wall of the valve. Just anterior to it lies the pallial line, a narrow groove with a slight and irregular interior elevated crest which descends to the bottom

of the visceral space and proceeds backward to a point underneath the posterior adductor muscle scar. Here it turns backward making a 45 degree angle, then turns straight dorsally, making with its former course a 90 degree angle, and reaches the lower angle of the scar. This rounded triangular scar likewise lies on an elevation formed by the inner contour mentioned, which passes dorsally and anteriorly, then turns anteriorly reaching the ventral border of the tooth. It thus leaves dorsal to it and posterior to the tooth, a kidney-shaped pit, into whose posterior border a small tooth projects.

Left valve, exterior: This valve is smaller and distinctly shorter than the right valve, but has almost the same depth. Its smaller length is due to the shortness of the rostrum which fails to cover about one-half the rostral area of the right valve, and thus leave a narrow wedge-shaped gap for the siphon tubes. The left valve is in contour an obliquely truncated ellipse. Seen from above, it has an umbonal angle of about 100 degrees and an almost straight antero-dorsal margin. This continues as a sharply and evenly rounded curve into the ventral margin which is gently curved, but which posteriorly curves more sharply, reaching the tip of the rostrum. The postero-dorsal margin is roughly a straight line, except for a slight bulging near the posterior half. The valve has a marginal geniculation of about two-thirds the height of the corresponding one on the right valve, and consequently a basin-like form for the entire valve.

The ornamentations consist of growth lines and weak concentric ribs, of which there are about seven below the geniculation, and 26 or more above it, the ribs decreasing in size in the dorsal one-third of the shell.

The ventral one-half of the valve, above the geniculation, bears about 12 ribs.

Left valve, interior: This valve has a truncate elliptical margin. It may be divided into two regions: the visceral cavity and the rostrum. The anterior part lacks the prominent shelf seen in the other valve. The visceral cavity is a subquadrate depressed area from which a shallow curved recess extends under the beak. Its margin is separated from the posterior margin of the valve by the triangular shaped rostrum whose inner surface is practically plane; the dorsal and narrow two-thirds of the triangle is occupied by the elliptical scar of the posterior adductor muscle; the anterior edge of this scar is smooth and forms the rim of the "shelf"; from it a perpendicular wall descends to the floor of the visceral cavity. The ventral and wider one-third of the rostrum is the space for the siphon tubes, and is plane with the floor of the visceral space.

From the lower end of the posterior adductor scar the narrow ribbon-like pallial line descends ventrally for a short distance, turns sharply pos-

teriorly and then makes an acute re-entrant angle (25 degrees) passing now anteriorly and almost perpendicularly to its former course; it runs in almost a straight line to the anterior one-third of the valve, where it gently curves upward meeting the postero-ventral corner of the scar of the anterior adductor muscle.

This scar is roughly pear-shaped and its tip is continuous postero-dorsally with an elevated ridge which passes along the bottom of the pit for the resilium. Dorsal to it, the elevated margin of the valve runs posteriorly, stopping short at the middle of this pit. This pit is subcircular and lies anterior to the central tooth just dorsal to its mid-dorsal point is the umbo. Posterior to it is a circular elevation which is separated by a constricted neck from the post-adductor muscle scar.

LOCALITY: Cut in Frisco railroad, three-fourths mile north of Union Station, Denison, Texas, in Weno marl, 20 feet below Quarry limestone (locality 604).

CORBULA LITTORALIS n. sp.

Pl. 10, fig. 5

MEASUREMENTS: (Type) length of valve ----- 18.0 mm.
Length of rostrum --- 4.0 mm.
Height of valve ----- 11.5 mm (?)

HORIZON: "Buff marl," near middle of the Weno formation, and immediately below zone of abundance of *Gervilliopsis invaginata* (White) in association with *Venericardia wenoensis*, *Trochus laticonicus*, cidarid spines, *Trigonia clavigera* and *Placosmilia* sp.

LOCALITY: 601, pit of brickyard, one and three-fourths miles southeast of Gainesville, Texas (type locality).

A fragmentary right valve, lacking umbo, antero-dorsal margin and posterior half of ventral margin is apparently different from any other described species, and it is considered best to describe it here. The individual is rather similar to *C. wenoensis* Adkins.

Right valve, exterior: The (restored) contour is elongate-triangular, with probably a nearly straight antero-dorsal margin, a sharply rounded anterior angle (probably about 60 degrees), a long nearly straight ventral side, curving up only slightly to meet the ventral margin of the rostrum. The rostrum is short, truncate, and the form of an isosceles triangle; its rounded postero-ventral angle is 100 degrees; its sharp postero-dorsal angle is 110 degrees; an almost obsolete shallow oblique groove separates it from the body of the valve, forming the base of the isosceles triangle. The postero-dorsal side is nearly straight except for a slight

convexity in its posterior one-third. The valve is distinctly rotund and elevated.

The ornamentation consists of 12 coarse rounded concentric ribs, and a number of fine growth lines in the umbonal region; the ventral half of the height contains nine of the coarse ribs. These ribs turn dorsally along the anterior end of the rostrum and just posterior to the oblique shallow groove, making an angle of 95 degrees to 100 degrees. The ribs near this angle bear each about four or five fine growth lines. The edges of the ribs appear as thin slightly imbricated and overlapping lamellae. Dorsal to the angle, the rostrum is conically inflated, the width of the fold increasing postero-ventrally and at the posterior end occupying the whole height of the rostrum. Dorsal to the inflation is a narrow concurrent and slightly divergent depression. The rostrum is half the height of the shell and is very close to the ventral margin. The ribs are unbranched and continuous from the anterior to the posterior ends of the valve. They are elevated with rather sharply rounded tops and are separated by plane bottomed, vertical-sided valleys whose width is one-sixth that of a rib.

Remarks: The following species of *Corbula* are to be found in the literature on the Comanchean: *C. pikensis* Hill (Trinity division); and *C. crassicostata* Cragin, Kiowa, Kansas, and (by implication) in the "ochreous shell-conglomerate of the Denison beds, Denison, Texas." *C. basiniformis* is distinguishable from the other species mentioned by its relatively unornamented surface and its geniculate margin. *C. littoralis* is distinguished from the others by its few coarse elevated and broad ribs; *C. wenoensis* is distinguished from *C. littoralis* by its smaller size, its different and less elongate form, its greater number of ribs, the more dorsal position of its rostrum, the shape and relative height and position of the rostrum, and by the internal characters as described.

CYPRIMERIA WASHITAENSIS n. sp.

Pl. 9, figs. 1-6

MEASUREMENTS:	Type	Pl. 9, fig. 4	Pl. 9, fig. 2
	Pl. 9, fig. 1		
Height -----	52.0	52.0	76
Length -----	58.5	59.0	77+
Breadth -----	15.0	17.5	25

HORIZON: Weno formation, upper and middle thirds, shale facies.

LOCALITIES: 604, cut of St. Louis and San Francisco Railway, one mile north of the Union Station, Denison, Texas (type locality); 601, pit of brickyards, three-fourths mile southeast of Gainesville, Texas.

DESCRIPTION: Shell subovate, inequilateral, subequivalve, beaks pointed anteriorly (prosogyrate); surfaces of both valves covered with numerous, simple, fine, striaeiform, concentric growth lines, and having a few widely spaced, coarser growth rings. Shell preserved, interior filled with ironstone. The beaks are inconspicuous, recurved anteriorly forming anterior to them a small notch in the margin and are rounded, incurved and approximate; anterior to them the margins of the valves are in contact and the growth lines are imbricated and separated, making several lines parallel to the margins of the valves. Posterior to the beaks is a much wider triangular depression between the margins of the valve. It runs from the umbo backwards to the postero-dorsal angle; its bottom is the apposed inner dorsal margins of the valves, its top the separated outer dorsal margins, while its sides consist of growth lamellae terminating abruptly and together constituting the thickness of the valves at this point. The groove in one individual contains over the medial half of its length a well preserved external hinge ligament. This has a chitinous appearance but is replaced by calcareous material.

The dentition is approximately that of *Cyprimeria* Conrad. The hinge of the left valve consists of two posterior lamellae, two teeth and an anterior lamella. The most dorsal posterior lamella is very broad and oblique, and is nearly parallel to the inner dorsal shell margin, which it has overgrown. It is roughly lanceolate or blade-shaped, the point being directed toward the umbo, the curved back directed ventrally, the edge directed dorsally and a narrowed portion directed postero-dorsally. It is separated by a narrow groove from the shorter second posterior lamella, which lies parallel to it and is coalescent with its umbonal half. Ventral to this lamella there is a narrow, deep, elongated pit, which separates it from the more posterior tooth. This pit in one individual contains a mass of brownish material chitinous in appearance but in fact calcareous. The two teeth are sharp, ridge-like and elongated, making an angle of about 60 degrees with each other; they are remote and are separated by a triangular area, raised centrally. Running anteriorly from the base of the tooth is an elongate elevated margin which may be taken as an anterior lamella. Dorsal and parallel to this is an elongate groove which is bounded dorsally by the upturned inner dorsal edge of the valve. Parallel and anterior to the anterior tooth is a narrow deeper depression the length of the tooth, which anteriorly connects with the groove. Below the teeth is an elongated area of small pits and punctations on the dorsal portion of the inner surface of each valve. *Cyprimeria crassa* Meek has been recorded¹

¹Hill, Annotated Check List, p. 14.

from the Eagleford shales near Denison, Texas and Cragin considers it to be widespread in the Fredericksburg division in South Texas. This species is stated to differ from *C. texana* (Roemer) in its large thickness and size. Cragin¹ also records *Cyprimeria* sp. aff. *excavata* Morton from the Eagleford shales ("top of *Ostrea quadriplicata* beds"). Still a larger *Cyprimeria*, called *C. gigantea*² by Cragin is recorded from the Grayson marl near Roanoke, Texas; we have found similar individuals in the Grayson at Roanoke, Denison, Burleson and elsewhere. Our species appears to be close to *Cyprimeria texana* (Roemer) which is known as casts from the Fredericksburg division. One individual of this species at hand shows a simple entire pallial line, some details of dentition, and papillae representing the pits on the inner surface of the valves below the hinge, as mentioned above in the description of *C. washitaensis* and as figured by Roemer and Conrad for *C. texana*. The two species seem to differ but slightly, yet in view of the uncertain systematic position of the Fredericksburg species and its poor preservation it is thought best to describe the present material separately. The relations of the pallial line to the posterior adductor muscle scar are entirely different in the two species and the scar is of a different shape. In a fragment of the left valve of a large individual of *C. washitaensis* from near Denison the scar is subquadrate in shape and the pallial line attaches directly to its lower anterior corner, while in Roemer's figure³ of the impression of the right valve, the scar is an inclined oval and its postero-ventral end lies in a prominent loop in the pallial line; this loop is missing in our species. Roemer's species also appears to have a considerably sharper marginal curvature just anterior to the umbo than has *C. washitaensis*, of which however I have not yet found any interior casts for direct comparison, all of the known material consisting of nacreous shells.

REMONDIA ? ACUMINATA (Cragin)

Pl. 6, fig. 1

This species does not have close similarities in form to that figured by Gabb,⁴ yet the hinge is rather similar. Stanton⁵ has figured and discussed the species of this genus. The hinge is here figured (Pl. 6. fig. 1). The species has an acuminate tip, nearly straight antero-dorsal and postero-dorsal margins and a gently rounded ventral margin. Posteriorly the

¹Cragin, Geol. Surv. Texas, 4th Ann. Report., pp. 176-7, 1893.

²Cragin, Ibid., p. 176.

³Roemer, Kr. Texas, pl. VI, fig. 8a.

⁴Gabb, Geol. Surv. Calif., Pal. II, p. 270, pl. 36, figs. 17-17a, 1869 (*Remondia furcata*).

⁵Stanton, Proc. U. S. N. M., XIX, 299-301, 1920.

shell is incurved to form a rostrum-like constriction. There are 8 to 15 roughly concentric, sharply elevated ridges which contract towards the posterior end of the shell.

The species is occasional in the lower Weno marl of the Fort Worth region, especially at locality 618, in association with *Pecten georgetownensis*, *Ancycloceras bendirei*, *Schloenbachia wintoni*, *Turritella worthensis*, and *Venericardia wenoensis*.

GASTROPODA

AMBERLEYA GRAYSONENSIS n. sp.

Pl. 6, fig. 5

MEASUREMENTS: (Type individual).

Height -----	21.0 mm.
Breadth (estimated) -----	17.2 mm.
Last whorl, height -----	11.0 mmf.
Last whorl, breadth -----	10.0 mm.

HORIZON: Basal stratum of Weno shale, in association with *Ostrea quadriplicata*, *Gryphea washitaensis*, *Salenia* sp. and various echinoids.

LOCALITY: 606, just south of Frisco track, two and one-half miles north of Denison, Texas.

DESCRIPTION: Shell conical, turreted, spiral angle about 63 degrees (estimated, shell of type distorted); volutions three or more, the terminal one being over one and one-half times the diameter of the next smaller one. Volutions angular, projecting, cross-section somewhat pentagonal. The volutions bear externally three coarse revolving ridges, of which the outer ones form the upper and lower shoulders of the volution. These ridges are equal and equally spaced, the distance between them being about two times the width of a ridge. The volution is crossed transversely by lanellar growth lines with thin overlapping irregular edges. From the center of the spire these trend sharply forwards, and crossing the first heavy ridge then pass obliquely across the outer face of the volution and after traversing the marginal strip of the volution on the other side of the ridges, disappear at the suture. They form slightly elevated imbricated low nodes on crossing the three spirals. Umbilicus absent, aperture indeterminate.

TROCHUS LATICONICUS n. sp.

Pl. 10, figs. 30-31

MEASUREMENTS:	I	II
Height -----	9.0	7.8
Greatest width -----	8.9	8.8
Aperture, height -----	3.0	2.5
Aperture, width -----	4.4	4.0

HORIZON: Weno formation, shale facies. Has so far been found only in the zone of *Gervilliopsis invaginata* (White).

TYPE LOCALITY: 601, pit of brickyard, one and three-fourths miles southeast of Gainesville, Texas.

DESCRIPTION: Shell small, conical, angulated. Volutions five of which, two earliest in this type, are nearly smooth. Form trochoid, apical angle 55 degrees, sutures impressed, ornamentation consists of a coarse, revolving spiral on the keel, ten finer spirals above, the alternate ones being coarser and consisting of equal low rounded tubercles, the others being straight fine revolving elevated ridges; and below the keel ten equal similar non-tuberculate spiral ridges, each about half the width of the intervening valley. Growth lines numerous, fine, sigmoid, oblique to the ridges. Umbilicus concealed, small if any; aperture broken; probably obliquely oval. The next tuberculate ridge above the keel is thicker than the others and the keel and it lie just above the suture on the younger whorls and form a distinct angulation at the base of each volution. The shell on the last volution is thick (one half mm.) and the external markings well preserved.

Trochus sp. Shattuck¹ from the Buda limestone at Austin differs from our species in having a larger apical angle (66 degrees) and straighter, non-angulated sides. *Trochus texanus* Roemer² from the Barton Creek (Austin, Texas) fauna of the Edwards limestone differs locally from our species in the nature of the ornamentation, the straightness of the sides and apparently in the shape of the aperture. This species is also described as having a pointed tooth on the inner margin of the outer lip.

¹Shattuck, U. S. G. S. Bull. 205, p. 31, pl. XIX, figs. 2-3.

²Roemer, Pal. Abh., vol. 4, pt. 4, p. 15, pl. 1, fig. 13.

HELICOCRYPTUS MEXICANUS Böse

1910: *Helicocryptus mexicanus* Böse, Inst. Geol. Méx., Bol. 25, p. 140, pl. 46, figs. 1-5; pl. 47, fig. 1.

A single individual lacking the mouth but otherwise well preserved and agreeing with the original description, was determined by Dr. Böse as belonging to this species.

LOCALITY: 714, near Fort Worth, Texas, in the basal Pawpaw clay.

NERITA sp.

Pl. 10, fig. 28

This fossil is found in the "buff marl" of the Weno brickyards at Gainesville, Texas, in association with *Trochus laticonicus*, *Venericardia wenoensis*, and other fossils listed in the geological section of this locality.

NERITINA sp.

Pl. 10, fig. 27

This small calcitic fossil is one of a group of peculiar small fossils associated in the "buff marl" of the brickyard pit, near Gainesville, Texas. The others are *Trochus laticonicus* Adkins, *Nerita* sp., *Venericardia wenoensis* Adkins, *Placosmilia* spp., *Trigonia clavigera* Cragin, *Leiocardis* spines, *Corbula littoralis* Adkins, *Pecten* sp. and numerous widely distributed Weno fossils.

ANCHURA MUDGEANA White

Pl. 10, figs. 39-40

1879: *Anchura mudgeana* White, 11th Ann. Rept. Geol. and Geogr. Surv. Terr., p. 312, pl. 7, fig. 3a-b.

1889: *Anchura (Drepanocheilus) mudgeana* Hill, Geol. Surv. Texas Bull. 4, p. 19.

1894: *Anchura mudgeana* Hill, Bull. Geol. Soc. Amer., vol. 5, p. —.

1918: *Anchura mudgeana* Stephenson, U. S. G. S. Prof. Paper 120-H, p. 141.

This common species of the Weno shale at Denison and Gainesville has not been found outside the Red River region, to my knowledge. It is conspicuous in the ironstone and blue shale layers of the Weno in association with a large nacreous fauna including *Corbula* spp., *Nucula* spp., *Turritella* sp., *Cerithium* sp., *Natica* sp., and others. It differs from *Anchura prolabiata* White in having a single instead of a double lip; and from *A. ruida*

White and *A. haydeni* White¹ in having a rounded, lobulated, externally directed lip instead of an upturned sharply pointed one. It differs from *Anchura monilifera* Gabb² found at Arivechi, Sonora, in the less elongate form and greater apical angle, in the greater width of the aperture, and in the lobulated lip.

NATICA sp.

Pl. 10, fig. 29

A small nacreous *Natica* is known from the ironstone layers of the Upper Weno shale of the Red River region. It is preserved with the slightly altered calcareous shell. Locality 601, pit of brickyards near Gainesville, Texas.

LUNATIA sp.

Pl. 10, fig. 38

This Weno species has a nacreous preservation and occurs in both the Upper Weno shale and the interspersed clay-ironstone seams. It is occasional in the ironstone heap thrown out of the brickyards pit near Gainesville, Texas.

GLOBICONCHA sp.

Pl. 10, fig. 41

This large species is rare in the Red River region, and is known from the clay-ironstone layers of the upper part of the Weno shale, in association with *Protocardia* sp. aff. *multistriata* (Shumard), *Turritella graysonensis*, *Cinulia washitaensis*, *Gervilliopsis invaginata*, *Schloenbachia wintoni*, *Cambarus* ? sp. and other Upper Weno fossils. Locality, 601, near Gainesville, Texas.

TURRITELLA GRAYSONENSIS n. sp.

Pl. 10, fig. 43

MEASUREMENTS:

Type, greatest height -----	36.0 mm.
Width of last volution -----	11.5 mm.
Section of last volution, height -----	7.5 mm.
Section of last volution, width -----	7.0 mm.

¹White, 11th Ann. Rept., U. S. G. and G. S. Terr., 1879.

²Gabb, Geol. Surv. Cal., Pal. II, p. 262, pl. 35, fig. 7, 1869.

HORIZON: Weno shale and ironstone, abundant; Pawpaw clay and sand, rare. Rare south of the Red River region.

LOCALITY: This is the common nacreous *Turritella* of the Weno ironstone and clay at Denison, Gainesville, and elsewhere in the Red River region. Locality 604, Frisco cut three-fourths mile north of Union Station, Denison, Texas (type locality) ; 601, near Gainesville, Texas.

DESCRIPTION: Shell turreted, steep, spiral angle 17 degrees; ten or more volutions, straight-sided, sutures sharply impressed, angles of volutions sharply rounded, surface almost smooth, bearing about fourteen lightly tuberculate spiral lines of unequal height. Of these, ten lie on the side of the volution and four below the shoulder. On the terminal volution of the type these are arranged as follows:

Beginning at the suture, the first ridge is strongest and bears widely spaced low obscure tubercles with flat spaces between; the thickness of this rib equals the width of the space separating it from the second rib. The second, fifth and seventh ribs are slightly narrower, but are ribbon-like and low in proportion to their breadth and have slight nodular elevations widely spaced from each other. The other ribs are low, subequal and have rounded tops and slight nodular elevations at wide intervals. The last four ribs lying beyond the shoulder of the volution are relatively narrower and more elevated than those on the flanks, and the inner three are closely spaced, being separated by about the width of a rib; the last rib is more widely separated but is the same size. The rest of the volution next to the umbilicus is smooth except for one or two obsolete spiral ribs. In the intervals between the ribs mentioned there are three to five fine equal raised lines. The transverse growth lines are prominent and sigmoid, and resemble those in *T. bravoensis* Böse, but are finer. Aperture unknown; cross-section of volution sub-quadrate, outer side straight, inner side rounded.

This species is separated from *Turritella bravoensis* Böse, *T. budaensis* Shattuck, and *T. planilateras* Conrad by having the tubercles very low, and depressed, instead of coarse and prominent. It is separated from *T. leonensis* Conrad, *T. marnochi* White, and *T. denisonensis* Cragin in having a greater number of spiral ridges and more tubercles in each ridge.

Cragin¹ describes without figuring, a variety under the name of *T. kansensis*, which has less crowded revolving ridges than our species. Likewise, *T. seriatim-granulata* Roemer² differs from *T. wenoensis* in having only five elevated tuberculated ridges, the middle one of which is flanked

¹Cragin, Notes on some fossils of the Comanche series, Science, n. s., vol. 6, pp. 134-136, 1897.

²Roemer, Kr. Texas, p. 39, pl. 4, fig. 12.

on each side by a low, non-tuberculate ridge. The ridges are wider and more prominent than in our species and the intervening spaces much narrower; the tubercles are more widely spaced; the fine secondary revolving lines are apparently lacking; and the contour is more angular. Roemer's species is also figured as having a smaller apical angle (about 12 degrees).

TURRITELLA WORTHENSIS n. sp.

Pl. 10, fig. 42

MEASUREMENTS: Type length of fragment 26 mm.; width (estimated) 14.5 mm.

HORIZON: Basal Weno formation, marl facies. Abundant in the Fort Worth region in association with *Ancycloceras bendirei*, *Remondia acuminata* and *Pecten georgetownensis*.

LOCALITIES: 618, near Fort Worth, Texas (type locality); extensively distributed in Tarrant County, Texas.

DESCRIPTION: Shell conical, steeply turreted, apical angle 23 degrees, with six or more volutions. Shell nearly straight sided, sutures impressed, surface with usually six nearly elevated spiral ridges bearing conspicuous, equal, equally-spaced, rounded tubercles. Beneath this are two parallel thin spiral ridges with obsolete tubercles. These form the bottom of the volution, and the uppermost of the six ridges mentioned, which is longer than the others, forms the top. Between each two ridges are four to six fine non-tuberculate spiral lines, but exceptionally one or two of these may be thickened, giving the appearance of an alternating series of fine and coarse revolving ridges on the volution. In most of the material at hand the six revolving ridges are equally spaced, and are separated by a depression only slightly wider than one of the ridges. The tubercles are equally spaced on all the ridges, and number about 65 on the last volution. These tubercles decrease in size on the earlier volutions, and all are present in the youngest stage examined.

Transverse growth lines are not visible. The details of the aperture are not preserved. The cross-section is rounded and triangular, the external side of the volution being longest. Below the shoulder of the usually tuberculate volution are about four impressed lines, of which the two already mentioned are not covered by the succeeding volution. Below this point the volution is smooth.

Many species of *Turritella* have been described from the Texas Comanchean, and some of these are indeterminate, on account of faulty figures and description, or lack of necessary information concerning stratigraphy and locality.

The present species is nearly straight-sided and is slightly angulated, while in *T. bravoensis* Böse¹ the middle rows of tubercles form a conspicuous angular projection on each volution. The number of rows of tubercles is greater than in Böse's species and the tubercles are more crowded in the rows. The conspicuous transverse striated growth lines of his species are absent in our material. *T. planilateras* Conrad² is somewhat similar in form to our species, but has a prominent alternation of tuberculate and smooth spiral ridges, while in our species the intermediate lines are tuberculate and do not approach the main ridges in thickness. *T. budaensis* Shattuck³ has the rows of tubercles diverse, the second row from the suture being most prominent, and the form is more angulated than in our species. *T. worthensis* is at once separated by its strong tubercles from the faintly tuberculate or granulated species, *T. seriatim-granulata* Roemer⁴ (Fredericksburg, Lower Washita), *T. marnochi* White,⁵ *T. leonensis* Conrad (Fredericksburg)⁶ and *T. denisonensis* Cragin⁷ (Mainstreet, Grayson).

Ellisor⁸ has described several species of *Turritella* of which only *T. manchacensis* (Buda) and *T. washitensis* (Buda) seem to approach the present species in ornamentation or form.

CINULIA WASHITAENSIS n. sp.

Pl. 10, figs. 33-37

1920: *Cinulia* sp. Winton and Adkins, Univ. Texas Bull. 1931, p. 66.

MEASUREMENTS:

Total height	15 mm.
Total breadth	14 mm.
Body whorl, excluding lip, height.....	10 mm.
Body whorl, excluding lip, breadth.....	10 mm.
Breadth of lip	4 mm.

HORIZON: Weno blue shale and clay-ironstone layers, middle and upper portions of the formation, especially in the Red River region.

LOCALITY: 601, pit of brickyards, one and three-fourths miles south-

¹Böse, Inst. Geol. Méx., Bol. 25, p. 149, pl. 31, figs. 8-9; pl. 32, figs. 1-2, 1910.

²Conrad, Mex. Bdry. Surv., Vol. 1, Pt. II, p. 158, pl. 14, figs. 1a-b, 1857.

³Shattuck, U. S. G. S. Bull. 205, p. 31, pl. XIX, figs. 4-6, 1902.

⁴Roemer, Kr. Texas, p. 39, pl. 4, figs. 12a-b, 1852.

⁵White, U. S. G. and G. S. Terr., 11th Ann. Rept., pp. 314-315, pl. 7, figs. 5a-b, 1879.

⁶Conrad, Mex. Bdry. Surv., Vol. 1, Pt. II, p. 165, pl. 21, figs. 7a-b, 1857.

⁷Cragin, Colo. Coll. Stud., 5, p. 65, 1895.

⁸Ellisor, Univ. Texas Bull. 1840, 1920.

east of Gainesville, Texas (type locality); 604, cut of St. Louis and San Francisco Railway, three-fourths mile north of Union Station, Denison, Texas. Occasional in Cooke and Grayson counties, Texas. About 25 individuals found.

DESCRIPTION: Shell of medium size, sub-globose, total outline sub-elliptical; four volutions, of which the last is much the largest and occupies two-thirds the height of the shell. Surface ornamented with evenly rounded equal revolving lines, about 25 on the last volution. Between the lines are concave depressions each about one and one-half times the width of a line. These contain numerous equally spaced vertical cross striations, which run perpendicular to the depressions and to the adjacent revolving lines. Aperture elongate, rounded-cuneate, narrowed at the top and enlarged at the bottom. Outer lip much thickened and rounded, elongate, extending the length of the two last volutions in the type (slightly shorter in other individuals), and bearing on the inner margin about 14 nearly equal blunt tubercles in two equal groups. The inner lip is flat and expanded and bears on its apertural margin three elevated spiral folds or laminae which project into the aperture; these are thickened and rounded at their free margins; the basal two lie almost horizontal while the upper one is directed somewhat downwards and is concealed underneath the inner lip. Viewed from outside the outer lip is very broad and thickened and bears about seven irregular coarse overlapping laminae.

This species resembles *Cinulia tarrantensis* Cragin¹ of the Goodland limestone near Fort Worth, but differs in several respects. *C. tarrantensis* is poorly described and figured and some of its critical points are indeterminate, no further material having been discovered from the type locality, Goodland limestone near Benbrook, Tarrant County, Texas. The type apparently is lost, or at least it is not in the other Dumble Survey material at Austin. *Cinulia* is extremely rare in the Goodland limestone of the Fort Worth region. The species appears to differ from *C. washitaensis* in having a narrow and much less imbricate lip and in having the spire lower in proportion to the length of the lip. In Cragin's species the lip does not seem to reach the top of the body whorl, while in the type of our species it is more elongate, extending the whole length of the terminal and next adjacent whorls instead of only three-fourths of the terminal whorl. In some other individuals it is lower, but never so low as in *C. tarrantensis*. The figure of *C. tarrantensis* shows the angulation of the turns above the body whorl to be pronounced, while in our species also differs in numerous details. The outer lip is broader in proportion

¹Cragin, Geol. Surv. Texas, 4th Ann. Rept., p. 223, pl. XLII, fig. 1.

these turns are rounded. The shape of the aperture and of the teeth to the body and bears much coarser imbricated lamellae than in Cragin's species; the aperture is more slender throughout and is noticeably more compressed below; the sub-central crenulate region of the inner edge of the outer lip is distinctly angular; the inner lip is broader and the three teeth placed differently, the two terminal ones being more closely spaced. Our species is apparently larger than *C. tarrantensis*, and comes from a higher horizon.

The unfigured and indeterminate species *Cinula ? texana* (Shumard)² was described from the Fredericksburg division of Bosque County; its description will apply to most species of the genus. *C. rectilabrum* Gabb¹ is a lower species and bears on its inner lip two teeth instead of three; it is more slender, more pointed apically, and has narrower lips and a more elongate aperture. *Cinulia pelletti* Whitney and *C. conradi* Whitney,³ both from the Buda limestone differ in several respects from the Weno species, the former in the proportions of the shell, the height and shape of the aperture, and the thickness of the outer lip, the latter in the shape, size and number of costellae.

PROTOZOA

NODOSARIA TEXANA Conrad

Pl. 11, fig. 2

This characteristic foraminiferean shell has already been described⁴ in some detail and the features mentioned apply to the Weno and Pawpaw material at hand. The fossil ranges in North Texas through the upper third of the Weno limestone and the base of the Pawpaw clay. The individuals are generally scattered, but also rarely occur in slabs. At Fort Worth there is a zone of abundance at a point about 18 feet below the top of the Weno formation, which is found at this stratigraphic level everywhere between the Red River and the Brazos. This zone appears to be different from the common Del Rio *Nodosaria* zone of West Texas, for prevailing in West Texas the zone of abundance of *Nodosaria* is in the top of the Del Rio clay just underneath the Buda limestone, and the fossil is scattering in the middle Del Rio clay. Near Del Rio, Texas, where the formation is about 200 feet thick the *Nodosaria* slabs are prominent near the top, and near the Chisos Mining Company, Brewster County, where

¹Gabb, Geol. Surv. Calif., Pal. II., p. 264, pl. 35, figs. 10-10a.

²Shumard, Trans. Acad. Sci., St. Louis, I, 1860, 597.

³Whitney, Trans. Texas Acad. Sci., XII, p. 23, pl. 10, figs. 9-11.

⁴Böse, Inst. Geol. Méx., Bol. 25, p. 177, 1910; Adkins and Winton, Univ. Texas Bull. 1945, p. 76, 1920.

the Del Rio clay is 120 feet or more thick, the zone of abundance of *Nodosaria* is above the occurrence of *Exogyra cartledgei* Böse, which lies 10-30 feet below the Buda limestone.¹ This level probably corresponds to the Grayson formation of North Texas. *Nodosaria* is rare at Austin in the middle Del Rio clay. At Cerro de Muleros, which is in the northern facies, it is reported as occurring in subdivision 5 (Duck Creek and Fort Worth), and as abundant in subdivision 6 (Denton, Weno and Pawpaw); this is its lowest recorded occurrence.³

BIBLIOGRAPHY

- Adkins, W. S., and Winton, W. M.: Paleontological Correlation of the Fredericksburg and Washita Formations in North Texas. Univ. Texas Bull. 1945, 1920. (Bibliography.)
- Berry, E. W.: The lower Cretaceous floras of the world. Maryland Geol. Surv., Lower Cretaceous, pp. 99-151. Baltimore, 1911.
- The upper Cretaceous floras of the world. Maryland Geol. Surv., Upper Cretaceous, pp. 183-313. Baltimore, 1916.
- Böse, Emil: Monografía geológica y paleontológica del Cerro de Muleros, etc. Inst. Geol. Méx., Bol. 25, 1910.
- On a new *Exogyra* from the Del Rio clay and some Observations on the Evolution of *Exogyra* in the Texas Cretaceous. Univ. Texas Bull. 1902, 1919.
- Cotteau, G.: Echinides, in Paléontologie française.
- Cragin, F. W.: A contribution to the Invertebrate Paleontology of the Texas Cretaceous. Geol. Surv. Texas., 4th Ann. Rept., pp. 141-294.
- Descriptions of Invertebrate fossils from the Comanche Series in Texas, Kansas, and Indian Territory. Colo. Coll. Stud., 5, 1895, 49.
- Dumble, E. T.: The Geology of East Texas. Univ. Texas Bull. 1869, 1920.
- Grossouvre, A.: Recherches sur la Craie Supérieure. I. Stratigraphie generale. II. Paléontologie. Paris, 1893-1901.
- Haug, Emile: Traité de Géologie. Paris, 1911.
- Hill, R. T.: The Geology of the Black and Grand Prairies of Texas. U. S. G. S. 21st Ann. Rept., Pt. 7, 1901.
- Geology of the Territory adjacent to the Red River in Arkansas, Texas, and Indian Territory. Bull. Geol. Soc. Amer., 5, pp. 297-338, 1893.
- Noetling, F.: Die Fauna der baltischen Cenoman-geschiebe. Pal. Abh., Bd. II, heft 4, 1885.
- Pervinquière, L.: Etudes de paléontologie tunisienne. I. Céphalopodes des terrains secondaires. II. Lamellibranches et gastropodes des terrains secondaires. Paris, 1907.
- Pervinquière, L.: Sur quelques ammonites du crétacé algérien. Mém. Soc. Géol. France, Paléontologie, tome xvii, fasc. 2-3, Mém. No. 42, pp. 1-86, pls. 1-7 (x-xvi), 1910.

¹Böse, Univ. Texas Bull. 1902, p. 19, 1919.

²Böse, Inst. Geol. Méx., Bol. 25, pp. 24-26, 1910.

- Richardson, G. B.: Report of a reconnaissance in Trans-Pecos Texas north of the Texas and Pacific Railway, Univ. Texas Mineral Surv., Bull. No. 9, 1904.
- Sayn, G.: Les ammonites pyriteuses des marnes valangiennes du sud-est de France. Mém. Soc. Géol. France, Paléontologie, tome 9, Mém. No. 23, pls. 1-6 (vi-x, pl. vii repeated), pp. 1-66, 1901.
- Sellards, E. H.: The Geology and Mineral Resources of Bexar County. Univ. Texas Bull. 1932, 1920.
- Sladen, P., and Spencer, W. K.: A monograph of the British fossil Echinodermata, Asteroidea. Paleontogr. Soc.
- Szajnocha, Ladislaus: Zur Kenntniss der mittelcretacischen Cephalopodea-fauna der Inseln Elobi an der Westküste Afrika. Denkschr. d. Kais. Akad. Wiss., Wien., XLIX, pp. 1-8, pls. 1-4, 1884.
- Whitney, F. L.: The Echinoidea of the Buda limestone. Bull. Amer. Pal., No. 26, 1916.
- Winton, W. M., and Adkins, W. S.: The Geology of Tarrant County. Univ. Texas Bull. 1931, 1920.

PLATE 1

Flickia, Schloenbachia, Mortonieras, Acanthoceras

PLATE 1.

- Flickia, Schloenbachia, Mortonicerias, Acanthoceras** **Plate 1**
- Figures 1-3. *Flickia boesei* n. sp.Page 85**
 Rare, basal Pawpaw formation, marl-clay transitional area. Fig. 1, type individual, side view, x 4.7; fig. 2, same individual, ventral view, x 4.7; fig. 3, same individual, x 1.33. Locality: 714, near Fort Worth, Texas.
- Figure 4. *Flickia* (?) *bosquensis* n. sp.Page 87**
 Rare, top of *Exogyra arietina* horizon, middle Del Rio clay (equivalent of base of the Grayson formation). Locality: cliff on west bank of the South Bosque River, 100 yards south of the bridge of the Speegleville road, and 5.5 miles west of the courthouse at Waco, Texas. Type individual, side view, x 2.0. (See Pl. 4, fig. 11.)
- Figure 5. *Schloenbachia* sp.**
 Rare, basal Pawpaw clay. Locality: 714, near Fort Worth, Texas.
- Figures 6-10, 18-19, 26. *Mortonicerias worthense* n. sp.Page 91**
 Abundant, Pawpaw formation, base, clay facies.
 Fig. 19, type individual, x 4.0. Locality: 723, near Fort Worth, Texas; figs. 10, 26, same locality, x 4.0. Other figures, x 2.0, same locality.
- Figures 11-13, 15-17, 20-25. *Acanthoceras worthense* n. sp.Page 93**
 Abundant, Pawpaw formation, base, clay facies.
 Fig. 12, type individual, x 2.0. Locality: 723, near Fort Worth, Texas. Other individuals, x 2.0, same locality.
- Figure 14. *Schloenbachia wenoensis* n. sp.Page 89**
 Rare, Pawpaw formation, base, clay facies. Fig. 14, type individual, x 2.0. Locality: 723, near Fort Worth, Texas.

PLATE 2

Scaphites, Hamulina, Metopaster, Comptonia

PLATE 2.

Scaphites, Baculites, Ptychoceras, Metopaster, Comptonia**Plate 2****Figures 1-12. *Scaphites hilli* Adkins and Winton.....Page 79**

Occasional, basal 10 feet of the Pawpaw formation, clay facies, sparse outside of Tarrant County region. Fig. 1, type individual, x 4.0, ventral view, showing suture and mid-ventral groove. Locality: 714, near Fort Worth, Texas. Fig. 2, individual showing venter with sutures outlined in the hematite and limonite areas, x 4.0. Locality: 714. Fig. 3, individual showing suture, dorso-lateral tubercle, and aperture, x 4.0. Locality: 714. Fig. 4, same individual, reverse side, x 5.0. Figs. 5, 6, individuals x 4.0. Locality: 714. Fig. 7, individual showing uncoiled portion and dorso-lateral tubercle, x 2.0. Locality: 723, near Fort Worth, Texas. Fig. 8, individual showing uncoiled portion, tubercle and juvenile suture, x 4.0. Locality: 723. Figs. 9-12, showing rib variations, x 4.0. Locality: 723.

Figures 13, 15-18. *Scaphites* sp. aff. *hilli* Adkins and Winton.....

Occasional, Pawpaw clay, base. Fig. 14, x 3.0; figs. 13, 16-18, x 2.0, showing rib variations. Locality: 713, near Fort Worth, Texas.

Figure 14. *Metopaster hortensae* Adkins and Winton.....Page 97

Rare, Pawpaw formation, base, clay facies. Type individual, aboral side, x 2.0. Locality: 714, near Fort Worth, Texas.

Figure 19. *Comptonia wintoni* n. sp.....Page 97

Rare, Pawpaw formation, base, clay facies. Type individual, aboral side, x 3.0. Locality: 714, near Fort Worth, Texas.

Figures 20-22. *Baculites comanchensis* n. sp.....Page 74

Occasional, Pawpaw formation, base, clay facies. Type individual. Fig. 21, x 4.0. Locality: 719, near Fort Worth, Texas. Fig. 20, x 2.0, same locality. Fig. 21, x 2.0. Locality: 714, near Fort Worth, Texas.

Figures 23-26. *Hamulina worthensis* n. sp.Page 71

Rare, basal Pawpaw clay. Type individual, Fig. 26, x 5.0, showing suture. Locality: 714. Fig. 23, individual showing form of curve and short limb, x 4.0. Locality: 714. Fig. 25, same individual, ventral view, x 2.0. Fig. 24, individual showing form of long limb and of curve, x 4.0. Locality: 714. Individuals of Figs. 23, 24 in museum of Texas Christian University, Fort Worth, Texas.

PLATE 3

Turritiles, Acanthoceras, Schloenbachia

PLATE 3.

Turrilites, Acanthoceras, Schloenbachia**Plate 3****Figures 1, 6. *Turrilites worthensis* Adkins and Winton.....Page 78**

Abundant, Pawpaw formation, base, clay facies. Fig. 1, x 4.0; locality: 714, near Fort Worth, Texas. Fig. 6, showing portion of suture, x 4.0; locality: 714, near Fort Worth, Texas.

Figures 2, 4. *Turrilites* sp.....Page 78

Abundant, Pawpaw formation, base, clay facies. x 4.0; locality: 714, near Fort Worth, Texas.

Figures 3, 7. *Turrilites bosquensis* n. sp.....Page 76

Rare, top of *Exogyra arietina* horizon, middle Del Rio clay (equivalent of base of the Grayson formation). Locality: cliff on west bank of the South Bosque River, 100 yards south of the bridge of the Speegleville road, and 5.5 miles west of the courthouse at Waco, Texas. Fig. 3, type individual, x 4.0, showing tubercles and angularity of profile. Fig. 7, same individual, x 2.0, showing aperture.

Figure 5. *Acanthoceras worthense* n. sp.....Page 93

Abundant, basal and middle Pawpaw clay. Individual showing venter, x 4.0. Locality: 714, near Fort Worth, Texas.

Figures 8-11. *Schloenbachia wintoni* n. sp.....Page 90

Occasional, Pawpaw formation; abundant, Weno formation. Figs. 8, 11, type individual, x 1.0; locality 601, near Gainesville, Texas, nacreous individual from ironstone in upper third of Weno formation. Figs. 9-10, nacreous individual, x 1.0, same locality. Fig. 9 shows the slightly bifid marginal tubercles. Type in museum of Texas Christian University, Fort Worth, Texas.

2000
2000

PLATE 4

Schloenbachia, Engonoceras, Flickia, Corbula

PLATE 4.

Schloenbachia Engonocreas, Flickia, Corbula

Plate 4

- Figure 1. *Schloenbachia* sp.
Rare, Pawpaw clay, x 4.0. Locality: 714, near Fort Worth, Texas. (See Pl. 1, fig. 5.)
- Figure 2. *Engonoceras* sp.Page 85
Rare, Pawpaw clay, x 2.5. Locality: 714, near Fort Worth, Texas.
- Figure 4. *Acanthoceras worthense* n. sp.Page 93
Abundant, Pawpaw clay, x 4.0. Locality: 714, near Fort Worth, Texas.
- Figures 3, 5-6, 12. *Engonoceras serpentinum* (Cragin)Page 84
Occasional, Weno shale, Grayson and Cooke counties, Texas and southern Oklahoma; rare, Pawpaw clay and shale. Figs. 3, 5, x 2.5; locality: 714, near Fort Worth, Texas. Figs. 6, 12, x 1.0; locality: 604, near Denison, Texas.
- Figures 8-10. *Engonoceras* sp.Page 85
Very abundant, Pawpaw clay, especially near the base; rare, Grayson, Denton, and Duck Creek formations, clay and marl facies, x 3.0, locality: 714, near Fort Worth, Texas.
- Figure 11. *Flickia* (?) *bosquensis* n. sp.Page 87
Rare, top of *Exogyra arietina* horizon, middle Del Rio clay (equivalent of base of Grayson formation). Type individual, x 4.0. Locality: cliff on west bank of South Bosque River, 100 yards south of the bridge of the Speegleville road, and 5.5 miles west of the courthouse at Waco, Texas. (See pl. 1, fig. 4.)

PLATE 5

Hemiaster, Epiaster, Enallaster

PLATE 5.

Hemiaster, Epiaster, Enallaster**Plate 5****Figures 1-2, 4. *Hemiaster calvini* Clark.....Page 114**

Occasional, Weno and Pawpaw formations; rare, Denton and Mainstreet formations; abundant, middle Grayson formation. Fig. 1, large sized individual with strongly developed ambulacral grooves, x 2.0; locality: Argyle, Texas, Grayson marl, Baylor University Museum. Fig. 2, individual in some respects resembling *Hemiaster bexari* Clark, x 3.0; locality: 720, basal Pawpaw marl, near Riovista, Texas. Fig. 4, x 2.0; locality: west branch of Little Mineral Creek, one mile northeast of Fink, Texas.

Figure 3. *Enallaster wenoensis* n. sp.....Page 112

Abundant, Weno formation, marl facies; rare, Pawpaw formation, marl and clay facies. Type individual, x 4.0; locality: 720, basal Pawpaw marl, near Riovista, Texas.

Figure 5. *Epiaster aguilerae* Bose.....Page 109

Rare, basal Fort Worth limestone. Locality: 406, one-half mile east of Texas Christian University, Fort Worth, Texas. Aboral side, x 1.0.

PLATE 6

Remondia(?), Venericardia, Amberleya, Hemiaster, Epiaster

PLATE 6.

Remondia, Venericardia, Amberleya, Hemiaster, Epiaster**Plate 6****Figure 1. *Remondia* (?) *acuminata* (Cragin).....Page 136**

Occasional, basal Weno marl; rare, Pawpaw clay. Locality: 618, near Fort Worth, Texas, individual, x 1, showing hinge structure.

Figure 2. *Venericardia wenoensis* n. sp.....Page 125

Occasional, Weno formation, shale facies, Red River region, and marl facies, Fort Worth region. Fig. 2, individual, x 2.0, showing ribbing. Locality: 618, near Fort Worth, Texas.

Figure 3. *Hemiaster calvini* Clark.....Page 114

Occasional, Weno and Pawpaw formations. Fig. 3, x 3.0. Locality: 903, Grayson marl, one-half mile southeast of Denison, Texas.

Rare, basal Pawpaw formation, clay-marl transition area; rare, Weno marl. Type individual, x 4.0. Locality: 720, near Riovista, Texas.

Figure 4. *Hemiaster riovistae* n. sp.....Page 115

Rare, Weno formation, shale facies. Locality: 606, basal stratum of Weno formation, near Denison, Texas. Type individual, x 2.0, showing carinae and ribbing.

Figure 5. *Amberleya graysonensis* n. sp.....Page 137

Abundant, basal Weno formation, marl facies; rare Pawpaw clay and marl. Type individual, x 1.0. Locality: 618, near Fort Worth, Texas.

Figure 6. *Epiaster wenoensis* n. sp.....Page 105

Abundant, basal Weno formation, marl facies; rare, Pawpaw clay and marl. Type individual, x 1.0. Locality: 618, near Fort Worth, Texas.

PLATE 7

Pentaceros, Comptonia, Metopaster, Pentagonaster

PLATE 7

Pentaceros, Comptonia, Metopaster, Pentagonaster**Plate 7****Figures 1-3. *Pentaceros americanus* n. sp.Page 99**

Pawpaw formation, clay facies, basal 5 feet, rare.

Fig. 1. Type individual, aboral side, x 2. Locality: 714, one-fourth mile south of the International and Great Northern Railway bridge across Sycamore Creek, four and one-half miles southeast of Fort Worth, Texas. The large plate above and to the left of the center of the disk is the madreporite. Type in museum of Texas Christian University, Fort Worth, Texas.

Fig. 2. Type individual, oral side, x 2.

Fig. 3. Type individual, aboral side, x 1.5.

Figures 4-5. *Comptonia wintoni* n. sp.Page 97

Pawpaw formation, clay facies, basal portion, rare.

Fig. 4. Type individual, aboral side, x 2. Locality: 714, near Fort Worth, Texas.

Fig. 5. Type individual, oral side, x 2.

Figure 6. *Metopaster hortensae* Adkins and Winton.Page 97

Pawpaw formation, clay facies, base, rare.

Type individual, oral side, x 2. Locality: 714, near Fort Worth, Texas.

Figure 7. *Pentagonaster texensis* Adkins and Winton.Page 95

Weno formation, limestone facies, upper 10 feet, rare

Locality: 602, east bank of Sycamore Creek, about four miles southeast of Fort Worth, Texas. Aboral side, x 1.5. Figured individual in museum of Texas Christian University, Fort Worth, Texas.

PLATE 8.

Hemiaster, Epiaster, Enallaster.**Plate 8**

- Figure 1. **Hemiaster longisulcus** (Adkins and Winton).....
Occasional, top of Fort Worth limestone. Type individual, x 2.0. Locality: On Cedar Creek, two miles southeast of Blum, Texas. Note the similar pores of the anterior unpaired ambulacrum, and the posteriorly placed apical system.
- Figures 2-3, 5. **Hemiaster riovistae** n. sp.....Page 115
x 2.0. Locality: 720, near Riovista, Texas.
- Figure 4. **Enallaster bravoensis** Bose.....Page 114
Occasional, Weno formation, marl and clay facies; rare, Pawpaw, Mainstreet and Buda formations; abundant, Grayson formation, marl and clay facies. Fig. 4, individual x 2.0. Locality 618, near Fort Worth, Texas.
- Figure 6. **Hemiaster calvini** Clark.....Page 114
Locality unknown, Cummins Collection. Note position of apical system, and the deep ambulacral grooves.
- Figure 7. **Epiaster aguilerae** Bose.....Page 109
Rare, basal Fort Worth limestone. Locality: 406, one-half mile east of Texas Christian University, Fort Worth, Texas. (Same individual as Pl. 5, fig. 5.) Apical system, x 4.0.

TO THE
MEMBERS OF THE

PLATE 9

Cyprimeria, Corbula

PLATE 9.

Cyprimeria, Corbula

Plate 9

Figures 1-6. *Cyprimeria washitaensis* n. sp.Page 134

Occasional, Weno shale, Cooke and Grayson counties, Texas. Fig. 1, type individual, x 1.0; locality: 604, near Denison, Texas. Figs. 2-4, x 1.0, same locality. Figs. 3-5, individuals showing hinge structure, x 1.0, same locality. Fig. 6, individual showing posterior adductor muscle scar and part of pallial line, x 1.0, same locality.

Figures 7-24. *Corbula basiniformis* n. sp.Page 130

Abundant, Weno shale and ironstone. Cooke and Grayson counties, Texas and southern Oklahoma. Fig. 21, type individual, x 1.5; locality: 604, near Denison, Texas. Other individuals, x 1.5, same locality.

PLATE 10

**Corbula, Arca, Protocardia, Neritina, Nerita, Natica, Lunatia,
Trochus, Anchura, Cinulia, Globicóncha, Turritella**

PLATE 10.

Corbula, Nucula, Arca, Protocardia; Neritina, Nerita, Natica, Lunatia, Trochus, Anchura, Cinulia, Globiconcha, Turritella Plate 10

Figures 1-4. *Corbula wenoensis* n. sp.Page 127

Weno shale, rare. Fig. 4, type. All individuals x 1.5. Locality: 601, pit of brickyards, one and three-fourths miles southeast of Gainesville, Texas.

Figure 5. *Corbula littoralis* n. sp.Page 133

Weno shale, rare. Fig. 5, type individual, x 1.5. Locality: 601, pit of brickyards, one and three-fourths miles southeast of Gainesville, Texas.

Figure 6. *Arca washitaensis* n. sp.Page 121

Abundant, Pawpaw clay; rare, Grayson, Denton and Duck Creek formations, clay facies and more rare in the marl facies. Fig. 6, type individual, x 1.5. Locality: 714, one-fourth mile south of the International and Great Northern Railway bridge across Sycamore Creek, near Fort Worth, Texas, at the base of the Pawpaw clay.

Figures 7-9. *Corbula basiniformis* n. sp.Page 130

Abundant, Weno shale. Locality: 604, cut of St. Louis and San Francisco Railway, one mile north of Union Station, Denison, Texas. (See Pl. 9, figs. 7-24.) x 1.5.

Figures 10-11. *Nucula wenoensis* n. sp.Page 120

Rare, Weno shale. Fig. 10, type individual, x 1.5. Fig. 11, x 1.5. Locality: 601, pit of brickyards, one and three-fourths miles southeast of Gainesville, Texas.

Figures 12-16, 19-20. *Nucula nokonis* n. sp.Page 118

Occasional, Weno shale, x 1.5. Locality: 604, near Denison, Texas. Type, Fig. 19, x 1.5, same locality.

Figures 21-26, 32. *Protocardia* sp. aff. *multistriata* (Shumard)Page 126

Abundant, Weno ironstone and shale, x 1. Locality: 601, near Gainesville, Texas.

Figure 27. *Neritina* sp.Page 139

Rare, middle Weno shale (buff marl). Locality: 601, near Gainesville, Texas. x 1.5.

Figure 28. *Nerita* sp.Page 139

Rare, middle Weno shale. Locality: 601, near Gainesville, Texas. x 1.5.

Figure 29. *Natica* sp.Page 140

Rare, ironstone and upper Weno shale. Locality: 601, near Gainesville, Texas.

- Figures 30-31. *Trochus laticonicus* n. sp.Page 138
 Rare, middle Weno shale (buff marl). Locality: 601, near Gainesville, Texas.
 Fig. 30, type individual, x 1.5. Fig. 31, x 1.5.
- Figures 33-37. *Cinulia washitaensis* n. sp.Page 143
 Occasional, middle and upper Weno shale. Locality: 601, near Gainesville, Texas.
 Fig. 33, type individual, x 1.5. Figs. 34-37, x 1.5.
- Figure 38. *Lunatia* sp.Page 140
 Rare, Weno ironstone and upper Weno shale. Locality: 601, near Gainesville,
 Texas. x 1.5.
- Figures 39-40. *Anchura mudgeana* White.Page 139
 Abundant, middle and upper Weno shale, Cooke and Grayson counties, Texas.
 Locality: 604, near Denison, Texas. x 1.5.
- Figure 41. *Globiconcha* sp.Page 140
 Rare, Weno ironstone. Locality: 601, near Gainesville, Texas. x 1.5.
- Figure 42. *Turritella worthensis* n. sp.Page 142
 Abundant, lower Weno marl, Tarrant County, Texas. Type. Locality, 618, near
 Fort Worth, Texas. Fig. 42, type individual, x 1.5.
- Figure 43. *Turritella graysonensis* n. sp.Page 140
 Abundant, Weno shale, Grayson County, Texas; occasional, Cooke County, Texas.
 Locality: 604, near Denison, Texas. Type, Fig. 43, x 1.5.

PLATE 11

Pecten, Epiaster, Nodosaria, Ancyloceras

PLATE 11.

Pecten, Epiaster, Nodosaria, Ancyloceras.

Plate 11

Figure 1. Ancyloceras bendirei n. sp.Page 70

Rare, Weno formation, base, marl facies. Fig. 4, type individual, x 1.2. Locality: 618, near Fort Worth, Texas.

Figure 2. Nodosaria texana Conrad.....Page 145

Abundant as isolated individuals and rarely in slabs, upper third of Weno and base of Pawpaw formations in north central Texas; abundant in flag layers, upper and to a less extent the middle Del Rio clay in West Texas. Fig. 3, x 5.0. Locality: Terlingua, Texas. (Compare Univ. Texas Bull 1945, pl. 21.)

Figure 3. Epiaster subobesus n. sp.....Page 110

Occasional, Weno formation, marl facies, and Pawpaw formation, marl facies. Abundant near base of Weno, in the "first terrace" of the Fort Worth region. Fig. 2. Locality: 618, near Fort Worth, Texas, type individual, x 1.0.

Figure 4. Pecten inconspicuus Cragin.....Page 123

Abundant, Weno shales and ironstone; rare, Denton clay; rare, Pawpaw formation, sand and ironstone facies. Cooke and Grayson counties, Texas, and southern Oklahoma. Fig. 1, clay-ironstone shell conglomerate from the middle Weno shales, x 4.0. Locality: 604, near Denison, Texas.

On A New Ammonite Fauna of the Lower Turonian
of Mexico

BY
Emil Böse

<i>Exogyra</i> Say.	230
<i>Exogyra haarmanni</i> sp. nov.	230
<i>Exogyra</i> cfr. <i>olisiponensis</i> Sharpe	230
GASTROPODA.	232
<i>Tylostoma</i> Sharpe.	232
<i>Tylostoma</i> aff. <i>ovatum</i> Sharpe.	232
ECHINODERMATA.	232
<i>Hemiaster</i> Desor.	232
<i>Hemiaster</i> sp.	232

LIST OF ILLUSTRATIONS

FIGURES

Figure 1.	<i>Metoecoceras</i> aff. <i>whitei</i> Hyatt, suture.....	204
Figure 2.	Suture, <i>Vascoceras</i> <i>angermanni</i> n. sp. (above) and <i>Vascoceras</i> aff. <i>adonense</i> Choffat (below)	215
Figure 3.	Suture, <i>Vascoceras</i> aff. <i>gamai</i> Choffat.....	216
Figure 4.	Sutures, <i>Vascoceras</i> <i>mohovanense</i> n. sp. (above), and <i>Mammites</i> <i>mohovaensis</i> n. sp. (below)	220
Figure 5.	Suture, <i>Neptychites</i> aff. <i>cephalotus</i> Courtiller.....	222
Figure 6.	Suture, <i>Neptychites</i> aff. <i>cephalotus</i> Courtiller.....	222
Figure 7.	Sutures, <i>Neptychites</i> aff. <i>xetriformis</i> Pervinqui�re (lower left hand corner) and <i>Hoplitoides</i> sp. (other three)	226

PLATES

Plates 12-20.	Lower Turonian Fossils from Cerro del Macho.....	235-252
---------------	--	---------

ON A NEW AMMONITE FAUNA OF THE LOWER TURONIAN OF MEXICO

BY EMIL BÖSE

INTRODUCTION

A number of years ago the existence of Turonian beds was proven in Mexico and later on it was shown that these beds have a great distribution in the country. At first very few localities with faunas of this age were known, but later the finds of fossils of Turonian age have augmented in such a manner that we now know that the Turonian has a vast distribution in Mexico, numerous localities having been discovered between Lat. 20°30' N. and Lat. 32° N. All of these beds are petrographically very uniform and consist of argillaceous shales and laminated limestones with intercalations of thin beds of limestone, all of black to light gray color. The fauna of these beds is also rather uniform and consists mostly of numerous specimens of *Inoceramus labiatus* Schlotheim, in some places accompanied by *Inoceramus hercynicus* Petraschek; in some localities fishes are relatively numerous; in others bivalves other than *Inoceramus* are found. Cephalopods have been found very rarely, all of them crushed and nearly indeterminable. The determination of the age had to be founded exclusively on the *Inoceramus* and on the position of the beds in relation to the Senonian and the Cenomanian.

About nine years ago I received the first collections of Turonian ammonites, which allowed a much better determination of the age of the beds mentioned here because these cephalopods were found together with well preserved specimens of the same *Inoceramus* which already had been determined by us as *I. labiatus*.

In February, 1911, Dr. Ernst Angermann sent me a small collection of fossils collected by him on the Cerro del Macho, Hacienda del Mohóvano, Municipality of San Pedro, District of Parras in the State of Coahuila. The greater part of these fossils consisted of internal molds of bivalves and gastropods, but there were also present three cephalopods relatively badly preserved and several indeterminable fragments of ammonites. I recognized at once that these fossils belonged to a facies and a horizon altogether unknown in Mexico, one of the ammonites being a *Vascoceras* of the group of *V. Kossmati* and another a *Neoptychites* of the group of *N. xettriformis* Pervinquière, the bivalves belonged mostly to *Trigonia*,

Avicula of the group of *A. gravis* Coquand, the gastropods to *Tylostoma* and similar genera.

A short time afterwards I showed these fossils to Dr. E. Haarmann, geologist of the Cia. Perforadora Mexicana, who then told me that he had a much larger collection of fossils from the same locality which he had made before Dr. Angermann had visited the place; and he offered me his material for a detailed paleontological study. The collection of Dr. Haarmann proved to be much larger than that of Dr. Angermann and contained a large number of ammonites some of which were rather well preserved. Haarmann in making his collection had separated the fossils of the different beds. The lowest of these beds was represented by a single piece of rock with impressions of bivalves, but as this rock had a very distinct character from that of the other beds and as in a locality of apparently the same age a badly preserved sea urchin had been found which appeared to belong to *Hemiaster*, I asked Dr. Haarmann to collect better material in the lowest horizon, because this horizon seemed to belong to the Cenomanian. Haarmann visited the locality again in August, 1911, collected ammonites in each of the three beds distinguished by him, and turned this material over to me in April, 1912.

The material thus collected proved to be of great stratigraphical and paleontological interest, as will be shown by the following description. But before I enter into a detailed study I wish to express my gratitude to Dr. E. Haarmann to whose amiability I owe the opportunity of studying this interesting fauna. I also wish to express my obligations to Professor Dr. W. Branca in Berlin who kindly sent me a great number of photographs of Turonian ammonites from Egypt, studied by Dr. Eck but at that time still undescribed.

LIST OF LITERATURE CITED IN THIS PAPER

1. Boule, Lemoine et Thevenin, Diego Suarez.—M. Boule, P. Lemoine et A. Thevenin, Paléontologie de Madagascar. Céphalopodes crétacés des environs de Diego-Suarez.—Ann. de Paléontologie t. I et II, Paris, 1906-07.
2. Choffat, Espèces nouv. ou peu conn.—P. Choffat, Recueil d'études paléontologiques sur la faune crétacique du Portugal. Vol. L, Espèces nouvelles ou peu connues. Ire. série. Section des Travaux géologiques du Portugal. Lisbon, 1886.
3. Choffat, Syst. crét. Portugal II.—P. Choffat, Recueil de Monographies stratigraphiques sur le système crétacique supérieur au nord du Tage. Service géol. du Portugal. Lisbonne, 1900.
4. Chudeau, Ammonites du Damergou.—R. Chudeau, Ammonites du Damergou (Sahara méridional).—Bull. Soc. géol. France, 4me sér., t. 9, fasc. 1-2, 1909.
5. Coquand, Géol. et Pal. de Constantine.—H. Coquand. Géologie et Paléontologie de la région sud de la Province de Constantine. Mém. Soc. d'Emulation de la Provence, II. Marseille, 1862.

6. Cotteau, Peron et Gauthier, Ech. foss. de l'Algérie.—Cotteau, Peron et Gauthier, Echinides fossiles de l'Algérie. Description des espèces déjà recueillies dans ce pays et considerations sur leur position stratigraphique. IV, Etage Cénomanién. Paris, 1878.
7. Eck, Turon in Aegypten.—O. Eck, Vorläufige Mitteilungen über die Bearbeitung der Cephalopoden der Schweinfurthschen Sammlung und über die Entwicklung des Turons in Aegypten. Monatsber. d. Deutsch. Geol. Ges. 1910. Berlin, 1911.
8. Fourtau, Faune crét. d'Egypte.—R. Fourtau, Contribution à l'étude de la faune crétacique d'Egypte. Bull. Institut Egyptien, sér. 4, vol. 4, 1903.
9. Fritsch, Ceph. böhm. Kreideform.—A. Fritsch, Cephalopoden der böhmischen Kreideformation (unter Mitwirkung des Dr. Urb, Schlönbach). Prag, 1872.
10. Geinitz, Elbthalgebirge.—H. Br. Geinitz, Das Elbthalgebirge in Sachsen. Palaeontographia Bd. 20, I u. II. 1872-75.
12. Guéranger, Album Paléontol.—Ed. Guéranger, Album Paléontologique de la Sarthe représentant au moyen de la photographie les fossiles recueillis dans cette circonscription. Le Mans, 1867.
13. Guillemain und Harbort, Profil d. Kreidesch. a. Mungo.—C. Guillemain und E. Harbort, Profil der Kreideschichten am Mungo. Abh. d. k. Preuss. Geol. Landesanstalt. Neu Folge, Heft 62, 1909.
14. Haug, Traité de Géologie. Paris, 1908-11.
15. Hyatt, Pseudocer.—A. Hyatt, Pseudoceratites of the Cretaceous. Edited by T. W. Stanton. U. S. Geol. Surv. Monogr. vol. 44, Washington, 1903.
16. Jack and Etheridge, Queensland.—R. L. Jack and R. Etheridge, Jr. The Geology and Paleontology of Queensland and New Guinea. Brisbane and London. 1892.
17. v. Koenen, Nachter. Foss. Mungo in Kamerun.—A. von Koenen, Nachtrag zu Ueber Fossilien der Unteren Kreide am Ufer des Mungo in Kamerun. Abh. K. Ges. d. Wiss. zu Göttingen, Math. Phys. Kl. Neue Folge, Bd. 1, No. 1. 1898.
18. Kossmat, Südind. Kreideform.—Fr. Kossmat, Untersuchungen über die südindische Kreideformation. Beiträge zur Palaeontologie und Geologie Oesterreich-Ungarns und des Orients, I Bd. 9, 1895; II u. III Bd. 11, 1897.
19. Lartet, Géol. de la Palestine.—L. Lartet, Essai sur la géologie de la Palestine et des contrées avoisinantes telles que l'Égypte et l'Arabie. II me partie Paléontologie. Ann. d. Sciences Géologiques publ. s. l. dir. de M. Hébert et A. Milne-Edwards. T. III. Paris, 1872.
20. Laube und Bruder, Amm. d. böhm. Kreide.—G. C. Laube und G. Bruder, Ammonites der böhmischen Kreide. Palaeontographica Bd. 33, 1887.
21. Lisson, Amm. del Perú.—C. I. Lisson, Contribución al conocimiento sobre algunos Ammonites del Perú. 4o Congreso científico Latino-Americano, 1o Panamericano, celebrado en Santiago de Chile. Lima, 1908.
22. Lisson, Terr. recon. en el Perú.—C. I. Lisson, Terrenos reconocidos hasta hoy en el Perú y sinopsis de su fauna y flora fósiles. Bol. de Minas, Ind. y Constr. publ. por la Escuela de Ingenieros, ser. II, t. IV. Lima, 1912.
23. Peron, Amm. du Crét. sup. de l'Algérie.—A. Peron, Les ammonites du Crétacé supérieur de l'Algérie. Mém. Soc. géol. France, Paléontologie, t. 6 et 7 mém. no. 17. 1896-97.
24. Pervinquière, Ét. géol. Tunisie.—L. Pervinquière, Etude géologique de la Tunisie

- Centrale. Thèses présentées à la Faculté des Sciences de Paris p. obt. le grade de Docteur-es-Sciences Naturelles. Paris, 1903.
25. Pervinquière, Paléontologie Tunisienne.—L. Pervinquière, Études de Paléontologie Tunisienne. I Céphalopodes des Terrains Secondaires. Dir. Gén. des Travaux Publics, Carte géol. de la Tunisie. Paris, 1907.
 26. Petrascheck, Amm. d. sächs. Kreideform.—W. Petrascheck, Die Ammoniten der sächsischen Kreideformation. Beitr. z. Pal. u. Geol. Oesterreich-Ungarns u. d. Orients. Bd. 14, 1902.
 27. Petrascheck, Inoc. a. d. Kr. Böhmens.—W. Petrascheck, Ueber Inoceramen aus der Kreide Böhmens und Sachsens. Jahrb. d. K. K. Geol. Reichsanstalt. Wien Bd. 53, 1903 (1904).
 28. Roemer, Kreidebild. v. Texas.—F. Roemer, Die Kreidebildungen von Texas und ihre organischen Einschlüsse. Bonn, 1852.
 29. Schlagintweit, Vracon u. Cenoman in Peru.—O. Schlagintweit, Die Fauna des Vracon und Cenoman in Peru. N. Jahrb. f. Min., Geol. u. Pal., Beilagebd. 33, 1912.
 30. Sharpe, Secondary distr. of Portugal.—D. Sharpe, On the Secondary District of Portugal which lies on the North of the Tagus. Quart. Jour. Geol. Soc. London, Vol. VI, 1850.
 31. Sharpe, On Tylostoma.—D. Sharpe, On Tylostoma, a proposed genus of Gastropodous Mollusks. Quart. Jour. Geol. Soc. London, Vol. 4, 1849.
 32. Solger, Mungokreide.—Fr. Solger, Die Fossilien der Mungokreide von Kamerun und ihre geologische Bedeutung, mit besonderer Berücksichtigung der Ammoniten. Beiträge zur Geologie von Kamerun herausgeg. v. E. Esch. Stuttgart, 1904.
 33. Stanton, Colorado Form.—T. W. Stanton, The Colorado Formation and its invertebrate Fauna. U. S. Geol. Surv. Bull. No. 16, 1893.
 34. Stoliczka, Ceph. Cret. Rocks India.—F. Stoliczka, The fossil cephalopoda of the Cretaceous Rocks of Southern India (Ammonitidae). Mem. of the Geological Surv. of India. Palaeontologia Indica, 1865.
 35. Thomas et Peron, Hauts-Plateaux de la Tunisie.—Ph. Thomas et A. Peron Description des mollusques fossiles des Terrains Crétacés de la région sud des Hantes—Plateaux de la Tunisie recueillis en 1885 et 1886. Exploration Scientifique de la Tunisie. Paris, 1889-90.
 36. White, Inv. foss. Nevada, Utah, etc.—Ch. A. White, Report upon the invertebrate fossils collected in portions of Nevada, Utah, Colorado, New Mexico, and Arizona, by parties of the expeditions of 1871, 1872, 1873 and 1874. Report upon Geogr. and Geol. Expl. and Surv. west of the 100th Meridian by Geo. A. Wheeler. Eng. Dept. U. S. Army, Part IV Palaeontology. Washington, 1875.
 37. White, Brazil.—Ch. A. White, Contribuções á Paleontologia do Brazil. Archivos do Museu Nacional do Rio de Janeiro, Vol. VII, 1887.
 38. Woods, Cret. dep. Northern Nigeria.—H. Woods, The palaeontology of the upper Cretaceous deposits of Northern Nigeria. The Geology and Geography of Northern Nigeria by J. D. Falconer. London and Bungay, 1911.
 39. Yabe, Cret. Ceph. Hokkaido II.—H. Yabe, Cretaceous Cephalopoda from the Hokkaido. Part II. Jour. of the College of Science, Imp. Univ., Tokyo, 1904, Vol. 20, art. 2.

40. Yabe, Strat. u. Pal. Hokkaido.—H. Yabe, Zur Stratigraphie und Paläontologie der oberen Kreide von Hokkaido und Sachalin. Zeitschr. d. Deutsch. geol. Ges. Bd. 61, 1909.
41. Zittel, Lib. Wüste.—K. A. Zittel, Beiträge zur Geologie und Palaeontologie der Libyschen Wüste und der angrenzenden Gebiete von Aegypten. I. Geologischer Theil. Palaeontographica Bd. 30, I. 1883.

STRATIGRAPHY

The Cerro del Macho is a small hill in the eastern part of the Hacienda del Mohóvano belonging to Mr. Frederick Ritter. The hacienda lies to the southeast of the station Carillo of the railway which connects Sierra Mojada, Coahuila with Escalon, Chihuahua (Mexican Central Railway) and part of its land belongs to the State of Chihuahua while the rest lies in the State of Coahuila. The Cerro del Macho belongs to this last region and lies near the boundary line of both states.

Dr. E. Haarmann gave me the following geological cross-section of the fossiliferous locality, to which I add the stratigraphical explanation obtained as the result of the study of the fauna.

n

Fig. 1

Fossil bearing beds on the Cerro del Macho, Hacienda del Mohóvano, Coahuila.

III. Gray hard limestones	}	Lower Turonian.
II. Gray-bluish marls		
I. Yellow and red marls and argillaceous limestones		
- Upper Cenomanian(?).

Three different petrographical horizons can be distinguished: the lowest beds consist of yellow and red marls with argillaceous limestones containing numerous fossils, especially bivalves; this horizon which for the present we shall designate with the number I has a thickness of only two

meters in this locality, but its base is covered. Above these marls lie gray-bluish marls with large cephalopods; the thickness of this horizon which we designate as Number II, is 2.5 meters. On top of these marls we find gray limestones with numerous cephalopods, bivalves and gastropods; the thickness of this horizon, which we call Number III, is about 5 to 6 meters, its upper limit being unknown. According to Dr. Haarmann the beds show a strike of N50-60°E and a dip of approximately 10°NW.¹

The whole outcrop has thus only a thickness of about 10 meters, but nevertheless we find two different horizons, as will be shown in the following. Among the fossils collected in this locality I have been able to distinguish twenty-one different species which are distributed in the three horizons in the following manner:

Horizon I.

Metoecoceras aff. *Whitei* Hyatt
Metoecoceras n. sp.
Exogyra *Haarmanni* n. sp.
Exogyra cfr. *olisiponensis* Sharpe
Hemiaster sp.

Horizon II.

Mammites *mohovanensis* n. sp.
Pseudaspidoceras aff. *Footeanum* Petrascheck
Pseudaspidoceras aff. *pedroanum* White
Vascoceras aff. *Adonense* Choffat
Fagesia *Haarmanni* n. sp.
Fagesia *Pervinquieri* n. sp.

Horizon III.

Vascoceras *Angermanni* n. sp.
Vascoceras ex. aff. *Gamai* Choffat
Vascoceras (?) sp.
Vascoceras *mohovanense* n. sp.
Neoptychites aff. *xetiformis* Pervinquièrè
Hoplitoïdes aff. *mirabilis* Pervinquièrè
Inoceramus *labiatus* Schlotheim
Avicula *Aguilerae* n. sp.
Trigonia sp.
Crassatella sp.
Tylostoma aff. *ovatum* Sharpe

To show the stratigraphic importance of the different species we shall have to compare them with related forms.

¹According to a sketch of Dr. Angermann the strike of the beds is N-S and the dip 10 degrees W.

Horizon I.

Species from Cerro del Macho	Related species	Upper	
		Cenomanian	Turonian
<i>Metoecoceras</i> aff. <i>Whitei</i> Hyatt	<i>M. Whitei</i> Hyatt	?	?
<i>Metoecoceras</i> n. sp.	<i>M. Gestianum</i> Petrascheck	+	—
<i>Exogyra</i> <i>Haarmanni</i> n. sp.			
<i>Exogyra</i> cfr. <i>olisiponensis</i> Sharpe	<i>Ex. olisiponensis</i> Sharpe	+	+
<i>Hemiaster</i> sp.			

The age of this horizon cannot be determined quite exactly. The fossils are few, the cephalopods among them indicate only that the horizon belongs either to the lower Turonian or the upper Cenomanian. *Metoecoceras* aff. *Whitei* is similar to forms found in the Colorado Formation of the United States, a division which up to now has not been subdivided stratigraphically with the necessary exactness; it probably represents in its larger part the Turonian, but may possibly contain some upper Cenomanian and a portion of Emscherian. The portion in which *M. Whitei* has been found belongs to the lower part of the formation, which is either the lower Turonian or the upper Cenomanian. Our *Metoecoceras* n. sp. is very similar to *M. Gestianum* Petrascheck,¹ which occurs in the upper Cenomanian of Saxony, but we do not know exactly if our specimen has been found in this lowest horizon or in one a little higher.

Interesting is the occurrence of an *Exogyra* certainly belonging to the group of *Exogyra olisiponensis* Sharpe. This species is found on the border of the Mediterranean in the Cenomanian as well as in the Turonian, although the variety which is most similar to our specimen seems to be limited principally to the Cenomanian. Our *Exogyra Haarmanni* n. sp. is a new species and not very characteristic, therefore without any stratigraphic value. The *Hemiaster* found in the same bed can not be used for the determination of the age as it does not appear to be identical with any described species nor to be closely related to any known species which might indicate the age of these beds.

The small number of fossils of this horizon does not allow an exact determination of its age. Taking into consideration the radical difference between the character of this fauna and that of Horizon II, and the presence of a form of *Metoecoceras* very similar to the one of the upper Cenomanian of Saxony, I feel inclined to consider this horizon provisionally as upper Cenomanian until other and more characteristic fossils allow a more exact determination of the age.

Much more numerous is the fauna of Cephalopods in Horizon II and

¹Petrascheck, *Amm. d. sächs. Kreideform.*, 1.140 (10) pl. 7 (1); figs. 3-5.

their forms are so characteristic that they easily allow the determination of the age of these beds. In the following table we shall compare them with related species of other countries.

Horizon II.

Species from Cerro del Macho	Related species	Turonian	
		Lower	Middle
<i>Mammites mohovanensis</i> n. sp.	<i>M. nodosoides</i> Schlotheim		
<i>Pseudaspidoceras</i> aff. <i>Footeanum</i>	<i>P. Footeanum</i> Petrascheck	+	—
<i>Pseudaspidoceras</i> aff. <i>pedroanum</i>	<i>P. Footeanum</i> Stoliczka	+	—
<i>Vascoceras</i> aff. <i>adonense</i>	<i>V. adonense</i> Choffat	+	—
<i>Fagesia Haarmanni</i> n. sp.	<i>F. superstes</i> Kossmat	+	—
<i>Fagesia Pervinquieri</i> n. sp.	<i>F. tevesthensis</i> Peron	+	—

All the species of this horizon indicate that it belongs to the lower Turonian or Salmurian. It is true that Choffat cites his *Vascoceras adonense* from the middle Turonian, but I believe that this middle Turonian of Portugal belongs in reality still to the Salmurian, i.e., the lower Turonian. The occurrence of forms belonging to the groups of *Fagesia superstes*, *Fagesia tevesthensis*, *Mammites nodosoides*, *Inoceramus labiatus*, etc., appears to be quite decisive for the determination of the age of those beds. In our own case we have to consider that this horizon lies immediately above what we take to be upper Cenomanian; there seems to be no reason to doubt that our Horizon II represents the very lowest Turonian. This idea is confirmed also by the fauna of Horizon III, which will be discussed farther on.

Our *Mammites mohovanensis* is very nearly related to *M. nodosoides* Schlotheim. It differs principally in its broader cross section; still more similar to our species is *Mammites conciliatus* Stoliczka, which belongs to the same group, the main difference consisting in its more evolute form and some details of ornamentation. In these details our form is still more similar to a form determined by Fritsch as *Mammites conciliatus* which in reality occupies an intermediate position between our species and that of India.

The species which we have called *Pseudaspidoceras* aff. *Footeanum* Petrascheck is much less related to the type of this group (*P. Footeanum* Stoliczka) than to the specimens from the lower Turonian of Saxony described by Petrascheck, on account of the more rounded and less subquadrangular cross section, as well as the greater height of the whorl. This difference may be explained in our section by its large size; comparing the different figured forms of *P. Footeanum* we see that the adult specimens tend to lose their quadrangular cross section. This group has been found

in the lower Turonian in many cases where an exact determination of the age of the beds has been possible. Varieties of this group have been described from India, from different parts of northern Africa, as Egypt¹ and Tunis²; from Portugal³, from Nigeria⁴, from Saxony⁵, and from Brazil.⁶ It is therefore a universally distributed form which characterizes the lower Turonian.

To the same group of *Pseudaspidoceras Footeanum* belongs also the species called by me *Pseudaspidoceras* aff. *pedroanum* White. It is still nearer related to the Indian type than the former species, but differs by its less strong ribs and the smooth ventral part. In all its characters it is similar to the *Ammonites pedroanus* White and only differs in some details of the ornamentation. The similarity is so great that I have not been able to separate the two forms specifically and have not united our specimen with the Brazilian species only because this latter form is still imperfectly known.

Our *Vascoreras* aff. *adonense* is certainly very similar to the Portuguese type from the Turonian, but differs by its suture and the form which is still more evolute.

The form of greatest stratigraphic interest in our fauna is perhaps our *Fagesia Haarmanni* n. sp. It belongs to the group of *Fagesia superstes* Kossmat, but differs from the type as well as from all the other species of this genus by its more evolute form. The species belonging to this group are limited to the Turonian and especially to the lower part of it. The type was described from the Utatur group of India,⁷ and a very typical form has also been found in Tunis⁸; a nearly related species has been cited from the Turonian of Portugal.⁹

Fagesia Pervinquieri n. sp. is a very characteristic species and belongs to that group of *Fagesia* which loses its ribs in a rather juvenile state of growth; the type of this group is *Fagesia tevesthensis* Peron. Our species differs from the type by its higher and less broad whorls. According to Pervinquière *Fagesia Rudra* Stoliczka¹⁰ belongs also to this group; this

¹Zittel, Lib. Wüste, p. LXXIX (A. cfr. *Footeanus*) Eck, Turon in Aegypten, p. 350 (*Acanthoceras* cfr. *Footeanum*).

²Pervinquière, Paléontologie Tunisienne, p. 314 (*Mammites Salmuriensis*).

³Choffat, Espèces nouv. ou peu conn., p. 66 (*Acanthoceras* cfr. *Footeanum*).

⁴Woods, Cret. dep. Northern Nigeria, p. 283 (*Mammites* [*Pseudaspidoceras*] sp.).

⁵Petrascheck, Amm. d. sächs. Kreideform., p. 144 (*Mammites Footeanus*).

⁶White, Brazil, p. 212 (*Ammonites pedroanus*).

⁷Kossmat, Südindische Kreideform., p. 133, pl. 17, fig. 1.

⁸Pervinquière, Paléontologie Tunisienne, p. 322, pl. 20, fig. 1-3.

⁹Choffat, Espèces nouv. ou peu conn., p. 69, pl. 10, fig. 4.

¹⁰Stoliczka, Ceph. Cret. Rocks India, p. 122, pl. 60.

species which occurs in the Utatur group of India is similar to our species on account of the missing umbilical nodules in a stage of development which still shows the ribs on the ventral part; it differs, though, by its much broader and larger whorls. Another form which by Pervinquire is considered as belonging to the same group is *Ammonites Kotoi* Yabe.¹ Pervinquire believes that this species is perhaps identical with *Fagesia tevesthensis* Peron, but it appears to me that we have here a different species although it is certainly related to the African form. According to Yabe² it is not quite certain from which beds this form has been collected; it occurs either in the Mammites beds or in the Scaphites beds. The age of *Fagesia Rudra* is not well known, although Kossmat presumes that this species belongs to the middle Utatur group. *Fagesia tevesthensis* characterizes the lower Turonian; a very similar form exists in beds of the same age in France and another in the Turonian of Portugal.

Still richer than the fauna of Horizon II is that of the upper limestones which we have distinguished as Horizon III. For their comparison with related species we shall again unite them in a table.

Species from Cerro del Macho	Related species	Lower	
		Turonian	Emscherian
<i>Vascoceras Angermanni</i> n. sp.	<i>V. Kossmati</i> Choffat	+	—
<i>Vascoceras</i> ex aff. <i>Gamai</i>	<i>V. Gamai</i> , Choffat	+	—
<i>Vascoceras</i> (?) sp.	<i>V. (?) arnesense</i> Choffat	+	—
<i>Vascoceras mohovanense</i> n. sp.	<i>V. polymorphum</i> Pervinqui�re	+	—
<i>Neoptychites</i> aff. <i>cephalotus</i>	<i>N. cephalotus</i> Courtiller	+	—
<i>Neoptychites</i> aff. <i>xetiformis</i>	<i>N. xetiformis</i> Pervinqui�re	+	—
<i>Hoplitoides</i> aff. <i>mirabilis</i>	<i>H. mirabilis</i> Pervinqui�re	+	—
<i>Inoceramus labiatus</i> Schlotheim	<i>I. labiatus</i> Schlotheim	+	—
<i>Avicula Aguilerae</i> n. sp.	<i>A. gravis</i> Coquand	+	+
<i>Tylostoma</i> aff. <i>ovatum</i> Sharpe	<i>T. ovatum</i> Sharpe	+	(Cenomanian to upper Turonian)

Nearly all of the groups of ammonites cited in this list occur only in the lower Turonian, to which, according to my opinion, belong also the beds with *Vascoceras* of the so-called middle Turonian of Portugal.

Vascoceras Angermanni n. sp. is very similar to *V. Kossmati* Choffat, but its form is still a little broader, there is no doubt that it belongs to the same group and this is until now only known in the Turonian of Portugal and the lower Turonian of Egypt.³

The fossil which we have called *Vascoceras* ex. aff. *Gamai* Choffat is a juvenile form and not very well preserved which is very similar to the

¹Yabe, Cret. Ceph. Hokkaido II, 26, pl. 6, figs. 3-4.

²Yabe, Strat. u. Pal. Hokkaido, p. 441.

³Eck, Turon in Tegypten, p. 381, 382.

small individuals figured by Choffat¹ in figures 3 and 4 of plate 7 and figure 2 of plate 10. Considering the size and state of preservation this specimen is not of great stratigraphic value.

Vascoceras (?) sp. is a pretty large specimen which is very similar to *Ammonites arnesensis* Choffat. The generic determination is not quite certain as is also the case with *A. arnesensis*. Our specimen is a little twisted; it does not have much importance for the determination of the age of the beds.

Vascoceras mohovanense n. sp. is a specimen relatively small but very characteristic; it belongs certainly to the group of *Vascoceras polymorphum* Pervinquièrè of the lower Turonian of Tunis. It differs from the type by a missing intermediate row of nodules but in its general character and also in its suture is nearly related to the type. Pervinquièrè considers *Vascoceras subconciliatum* Choffat from the Turonian of Portugal as a form of the same group but this species is different from ours on account of its general form and its suture.

The species described by me as *Neoptychites* aff. *cephalotus* Courtiller is a juvenile specimen which resembles greatly the small individuals of the cited species, as has been demonstrated in the paleontological part of this paper. The group of *Neoptychites cephalotus* characterizes the lower Turonian; it occurs frequently in Algiers and Tunis, also in the neighborhood of Saumur, France. Similar and perhaps identical forms have been found in India (*Neoptychites Telinga* Stoliczka)² and in Kamerun (*Neoptychites Telingaeformis* Solger³); the age of the beds where these latter species have been found is not exactly known.

The form which I compare with *Neoptychites xetrisformis* Pervinquièrè, is not very well preserved but resembles sufficiently the species from Tunis. Pervinquièrè considers as a nearly related form *Neoptychites crassus* Solger,⁴ the age of this latter species is not determined with certainty.

In the limestones of Horizon III we find frequently cephalopods which resemble very much the bicarinated *Hoplitoides* of Pervinquièrè. The greater part of these specimens is entirely corroded but in one I have been able to prepare a portion of the interior whorl which clearly shows the existence of the two lateral keels on the ventral portion; one also notes the suture in part, certainly much destroyed but showing the same elements as in the bicarinated *Hoplitoides*. A large specimen has the external form of

¹Choffat, Espèces nouv. ou peu conn.

²Stoliczka, Ceph. Cret. Rocks India, p. 125, pl. 62; Kossmat, Südind. Kreideform., p. 71, pl. 7, fig. 1, pl. 17, fig. 13.

³Solger, Mungokreide, p. 108.

⁴Solger, Mungokreide, p. 119, pl. 3, fig. 5.

Hoplitoides mirabilis but is a little less involute. This group is of some interest in so far as having been found only in the lower and upper Turonian. The types of *Hoplitoides mirabilis*¹ and *H. Munieri*² occur in Tunis mainly in the lower Turonian, *H. Munieri* also in the upper Turonian. A similar species has been found also in the Turonian of Egypt.³

In the same Horizon III, Dr. E. Haarmann also collected four specimens of *Inoceramus* of which three are typical individuals of *Inoceramus labiatus* Schlotheim. According to an observation made by José G. Aguilera, this fossil occurs in great numbers in beds north of the station of Carillo which apparently overlie limestones identical with those described here.

In Horizon III occurs very frequently *Avicula Aguilerae* n. sp. This species is very similar to *Avicula gravida* Coquand which occurs in northern Africa in the Turonian as well as in the lower Senonian or Emscherian. The form and size of this *Avicula* are so similar to those of ours that this find in Mexico has a certain stratigraphic interest.

In the material of Horizon III we find a great number of other bivalves, but all in the state of internal molds; a specific determination is entirely impossible. The most frequent form probably belongs to *Tylostoma* or some subgenus of *Natica*; the greater part of the material is badly preserved, but some specimens are a little more complete, although preserved as internal molds, and resemble strongly *Tylostoma ovatum* Sharpe which in Portugal occurs in beds from the Cenomanian to the upper Turonian.

From the character of the fauna of Horizon III we have to conclude that these beds still belong to the lower Turonian. This appears rather probable from the circumstance that Horizon II is extremely thin and is well confirmed especially by the cephalopod fauna. Interesting is the occurrence of *Inoceramus labiatus* Scholtheim. This fossil is extremely common in Mexico and generally is found in great numbers in calcareous shales and shaly limestones. The question now is, what relation exists between those beds and those of Cerro del Macho? In the described locality the shales, etc., do not exist, but according to the identification of J. G. Aguilera these are found north of the station of Carillo where they apparently overlie limestones similar to those of the Cerro del Macho. If we accept this explanation, our beds at Cerro del Macho have to be considered as the lowest part of the Turonian, while the shales with *Inoceramus labiatus* would be a little younger and perhaps represent the lower as well as the upper Turonian. We have no data with which to solve this problem, as up to now the beds with *Inoceramus labiatus* have not been found in contact

¹Pervinquière, Paléontologie Tunisienne, p. 218, pl. 10, fig. 3.

²Idem, ibid., p. 217, pl. 10, figs. 1, 2.

³Eck, Turon in Aegypten, p. 380, 386.

with the cephalopod beds described here; neither have those beds with *Inoceramus labiatus* been found in immediate contact with fossiliferous beds of the Emscherian. Only in one place, Opal, Zacatecas, I have found beds with *Inoceramus labiatus* (rare) and *I. hercynicus* (very frequent) in contact with sandstones containing an *Inoceramus* nearly related to *Inoceramus cycloides*, but the cross section of this locality is too incomplete and the fauna too poor for a decision of this rather important problem.

The locality of Cerro del Macho is stratigraphically important in so far as it gives us some data for the limitation of the upper Cenomanian and the lower Turonian. In a former work we have been able to subdivide the Cenomanian of Cerro de Muleros near Ciudad Juarez, but we have not been able to find fossils in the sandstones between the beds with *Inoceramus labiatus* and the marls with *Hemiaster Calvini*; thus it was impossible to decide if those sandstones belonged to the Cenomanian or the Turonian. In the present case we have very fossiliferous beds very near the limit between the Cenomanian and the Turonian. The fossils of the Cenomanian are very different from those found up to the present in Cenomanian rocks of Mexico. Of especial importance is the occurrence of *Metococeras*. The Turonian ammonite fauna is also entirely new for Mexico, the few ammonites so far found in the beds with *Inoceramus labiatus* belonging to entirely different groups.

Haarmann has distinguished two horizons, II and III, but these do not appear to be more than local subdivisions notwithstanding the circumstance that the fauna of the two beds seem to be rather different. We must not give too much importance to this circumstance, as the collections so far made are still rather small and the aspect of the fauna may change altogether with larger collections made. As far as we can see, the lower bed contains principally cephalopods while in the upper one bivalves and gastropods predominate.

With respect to the conditions of life in this locality we may say with some certainty that at the end of the Cenomanian in the Cerro del Macho region as well as in others of northern Mexico, especially in the State of Chihuahua, there existed a littoral facies or at least a very shallow sea; this is indicated by the great quantity of Ostreidae which nearly form beds or which have been carried away from neighboring beds. At the beginning of the Turonian age the sea seems to have deepened a little, the predominance of the ammonites and the absence of littoral bivalves indicating a less shallow sea. It is probable that during the time of the deposition of the upper part of the Salmurian the sea again became a little shallower; this is indicated by the predominance of Lamellibranchia and Gastropoda with thick shells, but the facies is not quite as much a littoral one as in

Cerro del Muleros near Ciudad Juarez at the same time, where the corresponding beds consist entirely of sandstones.

According to the observations of Haarmann and our palaeontological studies the stratigraphical cross section of the Cerro del Macho presents the following features:

5-6 m. of gray limestones with: <i>Vascoceras Angermanni</i> n. sp., <i>V.</i> ex aff. <i>Gamar</i> Choffat, <i>Vasc.</i> sp., <i>V. mohovanense</i> n. sp., <i>Neoptychites</i> aff. <i>cephalotus</i> Courtiller, <i>N.</i> aff. <i>xetiformis</i> Pervinquier, <i>Hoplitoides</i> aff. <i>mirabilis</i> Pervinquier, <i>Inoceramus labiatus</i> Schlotheim, <i>Avicula Aguilerae</i> n. sp., <i>Trigonia</i> sp., <i>Cras-satella</i> sp., <i>Tylastoma</i> aff. <i>ovatum</i> Sharpe, (Lamellibranchia and Gastropoda extremely frequent).	Hor. III	Lower Turonian
2.5 m. of gray-bluish marls with: <i>Mammites mohovanensis</i> n. sp., <i>Pseudaspidoceras</i> aff. <i>Footeanum</i> Petrascheck, <i>Ps.</i> aff. <i>pedroanum</i> White, <i>Vascoceras</i> aff. <i>adonense</i> Choffat, <i>Fagesia Haarmanni</i> n. sp. <i>F. Pervinquieri</i> n. sp.	Hor. II	(Salmurian)
2 m. yellow and reddish marls and limestones with: <i>Metaecoceras</i> aff. <i>Whitei</i> Hyatt, <i>Metaecoceras</i> nov. sp., <i>Exogyra Haarmanni</i> n. sp., <i>Ex.</i> cfr. <i>olisiponensis</i> Sharpe <i>Hemiaster</i> sp. (Lamellibranchia predominate).	Hor. I	Upper Cenomanian

If really the beds with *Inoceramus labiatus* lie immediately on top of the cephalopod beds, as the observations of J. G. Aguilera seem to indicate, another deepening of the sea may have followed and the general rising of the bottom of the ocean did not begin before the age of the lower Senonian.

ON THE RELATIONS BETWEEN THE FAUNA OF CERRO DEL MACHO AND ISOCHRONOUS FAUNAS OF OTHER REGIONS OF THE EARTH

The fauna of the upper Cenomanian of Cerro del Macho is too small to be compared with others; the most important fossil, *Metaecoceras* n. sp. belongs to a group which has been found in the upper Cenomanian. Of certain interest is the occurrence of *Exogyra* cfr. *olisiponensis* Sharpe because it indicates a faunistic similarity with the development of the Cenomanian along the border of the Mediterranean, a circumstance which to a certain degree can be noted also in the fauna of other Cenomanian beds of Mexico.

The fossils of the lower Turonian or Salmurian of the Cerro del Macho present a much better material for faunistic comparisons. Already while comparing our species with related ones from other regions we have seen that similar forms are found mainly around the Mediterranean, inclusive of Portugal. Comparing the faunas of the Salmurian in the above mentioned region, we find that all of them contain certain elements which either do not exist in most other regions or which occur only very rarely there. We shall see later on that this special fauna is not restricted to the Mediterranean but extends also to other parts of Africa and that certain elements belonging to it occur also in Asia and America. This indicates a large extension of the same horizon, but at the present time it is impossible to say if this distribution coincides with that of the same paleontological facies. But before we enter into such a discussion we must compare our fauna with the isochronous faunas of the Mediterranean and other parts of the earth in a somewhat more detailed manner.

Portugal: We begin with the region of Portugal. According to Choffat¹ we find in the lower part of the Turonian (beds with *Ostrea columba major*) the following cephalopods:

Puzosia cfr. *planulata* Sowerby
Vascoceras Mundaе Choffat
Vascoceras Gamai Choffat
Pseudaspidoceras cfr. *Footeanum* Stoliczka
Neolobites sp. nov.

In his middle Turonian we find the following species:

Vascoceras Gamai Choffat
Vascoceras Gamai Choffat var. *triangularis* Choffat
Vascoceras Mundaе Choffat
Vascoceras amieirense Choffat
Vascoceras silvanense Choffat
Vascoceras adonense Choffat
Vascoceras Grossouvrei Choffat
Vascoceras Barcoicense Choffat
Vascoceras Douvillei Choffat
Vascoceras subconciliatum Choffat
Vascoceras harttiforme Choffat
Vascoceras Kossmati Choffat
Vascoceras (?) *arnesense* Choffat

¹Choffat, Syst. crét. Portugal II. p. 162, 170, 173.

Choffat, Espèces nouv. ou peu conn., p. 44, 47.

Mammites pseudonodosoides Choffat
Pseudaspidoceras cfr. *Footeanum* Stoliczka
Fagesia aff. *superstes* Kossmat
Fagesia aff. *tevesthensis* Peron
Puzosia cfr. *Gaudama* Forbes
Puzosia sp.
Pachydiscus peramplus var. *beyrense* Choffat
Pseudotissotia Barjonai Choffat
Ammonites cfr. *Pachydiscus Rollandi* Peron
Ammonites sp. ind.

From the upper Turonian the author cites:

Vascoceras Gamai Choffat
Schloenbachia ?

Algeria: The Salmurian in northern Africa, especially in Algeria, Tunis and Egypt is very well developed. Still, in the year of 1904, Solger¹ could not cite from Algeria more than the following species of cephalopods: According to Peron:

Acanthoceras deverioide Grossouvre
Sphenodiscus Requieri d'Orbigny (*Hoplitoides ingens* according to Pervinquire)
Pachydiscus peramplus Mantell and other species of the same genus
Neptychites Telinga Stoliczka (*N. cephalotus* according to Pervinquire)
Puzosia Austeni Sharpe

According to Coquand:

Amm. Fleuriauxi d'Orbigny
Amm. papalis d'Orbigny
Heterammonites ammoniticas Coquand (*Hemitissotia Morreni* Coquand according to Pervinquire)

Peron cites further the following species:²

Mammites ? *tevesthensis* Peron (= *Fagesia tevesthensis* according to Pervinquire)
Pachydiscus Durandi Thomas et Peron (= *Vascoceras Durandi* according to Pervinquire)
Pachydiscus Rollandi Peron (= *Thomasites Rollandi* according to Pervinquire)

Tunis: In the present time we owe to the painstaking investigations of L. Pervinquire a much more complete knowledge of the fauna of the

¹Solger, Mungokreide p. 203.

²Peron, Amm. du Crt. sup. de l'Algrie.

Salmurian in northern Africa, especially of Tunis. Pervinquire cites and describes the following cephalopods:

Puzosia Austeni ? Sharpe
Pachydiscus peramplus Mantell
Hoplitoides Munieri Pervinquire
Hoplitoides mirabilis Pervinquire
Prionotropis Neptuni Geinitz
Acanthoceras Douvillei Pervinquire
Mammites nodosoides Schlotheim
Pseudaspidoceras salmuriense Courtiller
Pseudaspidoceras armatum Pervinquire
Fagesia superstes Kossmat
Fagesia tevesthensis Peron
Fagesia Peroni Pervinquire
Fagesia Fleuryi Pervinquire
Vascoceras Durandi Thomas et Peron
Vascoceras cfr. *barcoicense* Choffat
Vascoceras polymorphum Pervinquire
Thomasites Rollandi Peron
Thomasites Meslei Pervinquire
Thomasites Jordani Pervinquire
Pseudotissotia segnis Solger
Pseudotissotia Pavillieri Pervinquire
Pseudotissotia Luciae Pervinquire
Pseudotissotia Massipiana Pervinquire
Neoptychites cephalotus Courtiller
Neoptychites xetiriformis Pervinquire
Neoptychites Courguechoni Pervinquire

Egypt: Until a very few years ago our knowledge of Turonian faunas of Egypt was rather fragmentary, but in 1911 Eck published a preliminary study¹ of the collections made by Schweinfurth which contain faunas of the Cenomanian, Turonian and Senonian. From the lower Turonian Eck cites the following cephalopods:

Neolobites Schweinfurthi Eck
Fagesia bomba Eck
Fagesia cfr. *tevesthenensis* Peron
Vascoceras cfr. *amieirensis* Choffat
Vascoceras Kossmati Choffat

¹Eck, Turon in Aegypten. (On account of the war I have not been able to obtain the later work of the author.)

Vascoceras Durandi Thomas et Peron
Vascoceras barcoicense Choffat
Pseudotissotia segnis Solger
Pseudaspidoceras cfr. *Footeanum* Stoliczka
Hoplitoides sp. (?)
Hoplitoides cfr. *mirabilis* Pervinquière (?)

Southern Sahara: In Africa there are other localities where fossils indicating the existence of the Salmurian in a facies similar to that of Portugal, Tunis, Egypt, etc., have been found. Chudeau¹ describes some cephalopods from Damergou, a region which lies nearly exactly where Longitude 9°E Greenwich crosses Latitude 15°N. This author cites from that part:

Vascoceras Cauvini Chudeau
Acanthoceras Gadeni Chudeau

According to the author *Vascoceras Caudini* is similar to *V. Durandi* Thomas et Peron.

Nigeria: Woods² describes several cephalopods from northern Nigeria which seem to belong to Turonian forms. The species described are the following:

Vascoceras Nigeriense Woods (similar to *V. Durandi* Thomas et Peron)
Vascoceras Gongilense Woods (similar to *V. subconciliatum* Choffat)
Pseudaspidoceras sp. (similar to *Ps. Footeanum* Stoliczka)
Hoplitoides Nigeriensis Woods (similar to *H. Munieri* Pervinquière)

Kamerun: One can not doubt that the above cited fauna represents the Turonian and probably the Salmurian; therefore it is rather astonishing that a similar fauna has not been discovered in Kamerun, where only the *Neoptychites* of the Mungo remind us of Salmurian forms. Solger supposes that the Turonian exists in that region, but Guillemain and Harbort³ maintain that the beds in all their extension belong to the Emscherian.

From what we have seen in the foregoing description we conceive the idea that in northern and central Africa and in Portugal exists a facies of special features representing the Salmurian. This facies is distinguished by the frequency of the genera: *Vascoceras*, *Fagesia*, *Neoptychites*, *Pseudaspidoceras*, and in second place *Pseudotissotia*, *Thomasites* and *Hoplitoides*.

¹Chudeau, Ammonites du Damergou, p. 66.

²Woods, Cret. dep. northern Nigeria, pp. 281-286.

³Guillemain und Harbort. Profil der Kreidesch. a. Mungo, p. 431.

Syria and Palestine: Of this characteristic fauna nothing seems to be known in other parts of the Mediterranean. In Syria and Palestine, Turonian beds are scarcely known. Nötling cites from there a cephalopod similar to *Prionotropis Woolgari*, and Diener cites a *Mammites nodosoides*. This does not prove that the Salmurian in its north African facies does not exist there, because only a few years ago we did not know any more about the Turonian of Egypt, until Eck published his first lists of cephalopods.

Southern France: It seems that on the European border of the Mediterranean the north African facies of the Salmurian with its characteristic fauna of cephalopods is yet unknown. In France the same horizon has quite a different fauna. In Provence there are, according to Grossouvre,¹ beds with: *Mammites nodosoides*, *M. conciliatus* and *Pseudotissotia Douvillei*, but at least the first two of these species are nearly universally distributed types. Pervinquière adds to this list *Fagesia Boucheroni* Coquand. In Aquitania the following cephalopods have been found: *Pseudaspidoceras salmuriense*, *Fagesia superstes?*, *Fagesia Boucheroni* Coquand, *Fagesia tevesthensis*, and *Neoptychites cephalotus*. Thus this fauna contains quite a good number of the Salmurian fauna of the Mediterranean, although in the case of *Fagesia* and *Neoptychites* it seems that only very rare specimens have been found.

Saxony and Bohemia: The fauna of the Salmurian in Saxony and Bohemia contains a still smaller number of elements which might belong to the Mediterranean facies. Fritsch² describes from there as *Mammites conciliatus* Stoliczka a form which certainly belongs to this group, although as Kossmat already has noted, the species is different. The same author cites *Acanthoceras Neptuni*. Laube and Bruder³ figure *Mammites nodosoides* and other species of this genus, as well as *Puzosia Austeni*. Petrascheck⁴ cites *Mammites Footeanus* Stoliczka, *Mammites* cfr. *crassitesta* Stoliczka, *Barroisiceras* (?) *Fleuriausianum* d'Orbigny, *Acanthoceras* cfr. *Choffati* Kossmat; all forms of groups which in part occur around the Mediterranean and in part in India.

Northern Europe: The faunas of the Salmurian of northern France and northern Germany show much less similarity with the Mediterranean facies. They contain only ammonites belonging to groups of a universal distribution, like *Mammites nodosoides*, *Barroisiceras* (?) *Fleuriausianum*, *Acanthoceras Neptuni*, *Pachydiscus peramplus*, *Puzosia Austeni*, etc. Of course, we have to take into consideration that neither in northern Ger-

¹Grossouvre, Craie Supérieure, Stratigr., p. 507.

²Fritsch, Ceph. böhm. Kreideform., p. 35.

³Laube und Bruder, Amm. d. böhm. Kreide, p. 229.

⁴Petrascheck, Amm. d. sächs. Kreideform., p. 155.

many nor in Bohemia is there a modern stratigraphic-paleontological work on the Turonian fauna.

Less still do we know about England, from which Solgar could cite only a very few species as probably of Turonian age.

From the foregoing we see that the more we advance toward the north, the more the elements belonging to the Mediterranean facies of the Salmurian diminish.

India: Certain faunal relations exist between the Salmurian of India and that of the Mediterranean. In India the group of *Mammites nodosoides* is represented by two species, *Mammites conciliatus* Stoliczka and *M. crassitesta* Stoliczka; the group *Pseudaspidoceras* by one, *Ps. Footeanum* Stoliczka. The genus *Fagesia* is represented by two species, *Fagesia superstes* Kossmat, and *F. Rudra* Stoliczka. In the same manner we find *Neoptychites* represented by two species: *Neoptychites Telinga* Stoliczka (= *cephalotus* Courtiller according to Pervinquière) and *N. xetra* Stoliczka.

Japan: In Japan there is probably also Turonian with elements of the Mediterranean facies. Yabe cites the occurrence of beds with *Mammites*. In these he found a *Mammites* of the group of *M. nodosoides* Schlotheim.¹ The same author describes a *Fagesia Kotoi* which occurs either in the same beds or in those with *Scaphites*, which are a little younger.

Australia: It seems that in Australia the Turonian has not yet been found; at least the fossils found in the Desert Sandstone, considered by Jack and Etheridge as the representative of the upper Cretaceous,² do not allow such a conclusion.

Brazil: In South America only very few localities have been discovered where fossils of the Salmurian have been found. Some species were described from Brazil by White but without any indication of their age. Several authors have already noted that the Salmurian must exist there, but it seems nearly impossible to separate the Salmurian fossils from those of the other Cretaceous horizons. Pervinquière notes the similarity between *Vascoceras Durandi* and *Ammonites Harttii* Hyatt,³ while Choffat who had before that time called attention to the similarity between that species and his *Vascoceras harttiforme* Kossmat believes that *A. Harttii* might be a *Fagesia*, but the suture published by Hyatt⁴ shows that the species belongs to *Vascoceras*.

Kossmat has also indicated the similarity between *Ammonites pedroanus*

¹Yabe, *Strat. u. Pal. Hokkaido*, p. 441.

²Jack and Etheridge, *Queensland*, p. 557 et seq.

³White, *Brazil*, p. 226, pl. 19, figs. 1, 2; pl. 20, fig. 3 (erroneously called *A. Pedroanus* in the explanation of the plate and in the text).

⁴Hyatt, *Pseudocer.*, pl. 14, fig. 16.

White¹ and *Pseudaspidoceras Footeanum* Stoliczka. This species is found at the same locality (Bom Jesus, Larangeiras, Provincia de Sergipe) where *Vascoceras Harttii* occurs. It is therefore probably safe to say that the Salmurian exists in that region in a facies similar to that of northern Africa.

Pervinquire finally has noted that his *Vascoceras polymorphum* is very similar to *Ammonites offarcinatus* White,² a species found at Trapiche das Pedras Velho, Porto dos Barcos and Bom Jardim in the Province of Sergipe. It seems to me probable that this type belongs to the multi-tuberculated *Vascoceras*.

Peru: Species from the Salmurian have also been found in Peru. Lisson³ described and figured in 1908 two specimens of *Vascoceras* which he identifies with *Vasc. amieirens* Choffat. There can be no doubt that this form is a real *Vascoceras* very similar to *Vasc. amieirens*, although the Peruvian form is a little more involute and the cross-section different. Later on Lisson⁴ cites from Cuesta de Huanambra, in the province of Cajamarca, a *Mammites nodosoides* var. *Afra* Pervinquire, but notes that the fossil seems to occur in the lower Senonian (Emscherian). From the same locality he cites *Hoplitoides ingens* v. Koenen; this makes the determination of the first species doubtful or indicates that both the Salmurian and the Emscherian exist in this locality. This latter opinion is confirmed by a geological cross-section published by Schlagintweit,⁵ whose locality "Cuesta de Huanyanba," is probably identical with the "Cuesta de Huanambra" of Lisson, as both indicate that the place lies to the west of Celendin. Schlagintweit gives the following cross-section:

6. Senonian beds.
5. Few meters of marly limestones of yellow color with *Mammites nodosoides* Schlotheim. *Turonian*.
4. About 15 m. sandy dolomitic limestone.
3. Few meters of yellow marls of the Cenomanian with *Exogyra columba* Lamarck.
2. 100-150 meters of light-colored limestone. *Vraconian*.
1. Aptian-Gault with fossils.

United States: In the rest of America no fauna similar to the Mediterranean facies of the Salmurian appears to have been discovered. In the western United States the Turonian is represented by the Eagleford shales,

¹White, Brazil, p. 213, pl. 22, figs. 1, 2 (non pl. 20, fig. 3).

²White, Brazil, p. 219, pl. 23, figs. 3, 4.

³Lisson, Amm. del Per, p. 9, 9a, 9b.

⁴Lisson, Terr. recon. en el Per, II, p. 1.

⁵Schlagintweit, Vracon u. Cenoman i. Peru, p. 57.

similar to the beds of Mexico, with *Inoceramus labiatus*; and in the central United States by the Benton and the Colorado formation. This latter probably contains also the Emscherian. The cephalopods collected in these beds are entirely different from those of the Mediterranean Salmurian and seem to occur in slightly younger beds. Possibly the Salmurian may yet be found in Texas, as there is only a short distance between this state and our locality.

From the foregoing we reach the conclusion that our fauna of Cerro del Macho has very near relations with the African and Portuguese facies of the Salmurian and that a similar fauna probably exists in South America on the Pacific coast (Peru) as well as on the Atlantic border (Brazil). In South America, especially in Peru, the similarity with the faunas of north Africa appears to begin even in the Cenomanian (Schlagintweit) and to persist still in the Senonian (Lisson), while in Mexico we can not say the same, although in the Cenomanian as well as in the Senonian there exist certain elements which remind us of forms from north Africa. We certainly have to take into consideration that the fauna of the Cenomanian of Mexico is still very little known, only the faunas of Cerro del Muleros and of the Vraconian of Camacho, Zacatecas, having been studied in detail; and this latter fauna resembles entirely that of Europe. The few fossils contained in the probably Cenomanian beds of Cerro del Macho seem to indicate in a certain degree that the upper Cenomanian of this region has a fauna similar to that of north Africa; but this problem can only be solved in the future when larger collections have been made.

The Salmurian faunas of the African-Portuguese region as well as those of Cerro del Macho have some, although not very near, relations with the Salmurian of India and Japan, the connecting elements consisting of a few groups of *Mammites*, *Pseudaspidoceras*, *Fagesia* and *Neoptychites*; and we might even say that our facies is as much related to that of India and Japan, as it is to that of France and Saxony-Bohemia.

I do not pretend that these results are to be considered as definite, because the Salmurian in general is yet very little studied, not only in far away countries like India and Japan but also in a great part of Europe.

PALEONTOLOGY—CEPHALOPODA

METOECOCERAS Hyatt

Hyatt¹ proposed the name of *Metoecoceras*² for the forms belonging to the group of *Ammonites Swallovi* Shumard and united in the genus the

¹Hyatt, *Pseudocer.*, p. 116.

²I accept the proposal of Pervinquière to change the original name *Metoicoceras* into *Metoecoceras*.

following species: *M. Swallowi* Shumard, *M. gibbosum* Hyatt, *M. Whitei* Hyatt, *M. acceleratum* Hyatt, and *M. Kanabense* Hyatt (figured but not described). These are forms similar to *Acanthoceras* in one direction and to *Pulchellia* in the other; they are always involute, with relatively flat flanks, a nearly rectangular or lanceolate cross section, much higher than broad. The ornamentation consists of straight or slightly flexuous ribs which begin at the umbilical border without forming real nodules in that region, although in some cases they show slight thickening in that place. The ribs continue on the flanks, showing near the ventral border a thickening or very little pronounced nodule, and on the ventral shoulder a very sharp nodule which is somewhat longitudinally prolonged and very near the above mentioned thickening. Between the main ribs other secondary ribs are intercalated which begin above the umbilical border in different heights; the main ribs are often bifurcated. Between the two rows of nodules on the ventral shoulders there is a flattened zone, smooth in some cases while in others the very low ribs cross this zone. The nodules of the two ventral shoulders frequently form a kind of undulated or interrupted keels.

The suture of small individuals resembles that of *Heinzia* and *Pulchellia*, while that of larger specimens is similar to the suture of certain groups of *Acanthoceras* (*Mantelliceras*) and *Acompsoceras*. Characteristic is the external saddle divided by a secondary rather deep lobe; the first lateral lobe is broad and not very deep, the first lateral saddle is narrow but in general higher than the external saddle; the second lateral lobe is narrow and short, showing scarcely half of the depth of the first one. The second lateral saddle is similar to the first one but much lower; there are always three or four auxiliary saddles or even more in very large specimens. All the saddles are very little ramified, the auxiliary ones often entirely whole. The first lateral lobe seems to be always bifid.

Very characteristic also is the manner of involution in this genus. While the internal whorls are completely involute, the external whorl covering the greater part of the inner one, we see that the larger whorls and the animal chamber show a much wider curve and cover a much larger part of the next smaller whorl.

The species belonging to this group are so characteristic that one has to consider them as an independent genus or subgenus. Hyatt considers this group as a special family, giving it a position between the *Heinzidae* and the *Mantelliceratidae*.

To the species united by Hyatt in the genus *Metococeras* we probably have to add some European forms. One of these is *Pulchellia Gestiana* Petrascheck¹ from the upper Cenomanian of Saxony. Petrascheck con-

¹Petrascheck, *Amm. d. sächs. Kreideform.*, p. 140, pl. 7, figs. 3-5.

siders this species as a *Pulchellia* and it certainly has the greatest similarity with this genus if we do not take into account *Metoecoceras*. Comparing the figures of Petrascheck with those of the American *Metoecoceras* we note a surprising similarity; for example, between figure 4 of Petrascheck and figure 2 of Stanton,¹ and between figure 3 of Petrascheck and figure 10 of plate 15 of Hyatt.²

The suture of the specimens from Saxony is imperfectly known, but the elements which can be recognized in figure 4 of plate 7 and figure 5 of the text (p. 141) of Petrascheck, coincide very well with those of *Metoecoceras*.

Petrascheck identifies his specimens with *Ammonites Geslianus* d'Orbigny,³ but Sayn points out that the figure of d'Orbigny differs very much from the figures of Petrascheck. It is possible that *Ammonites Geslianus* d'Orbigny belongs also to *Metoecoceras* but this can not be decided as long as the suture is entirely unknown. The specimen figured by Guéranger⁴ under the name of *A. Geslianus* is so badly preserved that no details can be recognized. What Geinitz⁵ figures as *A. Geslianus d'Orbigny* seems to differ considerably from the type and belongs perhaps to *Stoliczkaia*. Petrascheck accepts the determination of Geinitz but does not figure any specimen which shows the transition between the smooth ventral part and the form in which the ribs pass the venter without interruption.

Kossmat⁶ has noted the external similarity between *Acanthoceras vicinale* Stoliczka⁷ and "*Buchiceras*" *Swallovi* Shumard, indicating at the same time the difference in the suture which is much more complicated, especially in the auxiliary saddles of the Indian species.

Still more similar to *M. Swallovi* is a cephalopod from Madagascar described by Boule, Lemoine and Thevenin⁸ under the name of *Acanthoceras (Prionotropis) subvicinale*. The suture of these specimens coincides perfectly with that of *Metoecoceras*, as has already been observed by the authors, but it also has a great similarity with *Pulchellia*. The specimens are very small and show an ornamentation very similar to that of *Metoecoce-*

¹Stanton, Colorado Form., p. 168, pl. 38 (*Buchiceras Swallovi* White; according to Hyatt, this is a new species, *M. Whitei*, Hyatt).

²Hyatt, Pseudocer.

³D'Orbigny, Pal. fran c., terr. crét., ceph. p. 325, pl. 97, figs. 1-2 (under the name of *Am. catillus* Sow.; this determination was corrected by d'Orbigny in Prodrôme II, p. 146).

⁴Guéranger, Album Paléontol., pl. 5, fig. 2.

⁵Geinitz, Elbthalgebirge I, p. 280, pl. 62, fig. 3.

⁶Kossmat, Südind. Kreideform., p. 201 (105).

⁷Stoliczka, Ceph. Cret. Rocks India, p. 84, pl. 44. Kossmat, Südind. Kreideform., p. 200 (104), pl. 25 (11), fig. 2.

⁸Boule, Lemoine et Thevenin, Diego-Suarez, p. 31, pl. 8, fig. 5; fig. 16 of the text.

ras; they differ principally by the presence of a median row of scarcely perceptible, very small tubercles on the ventral part. Unfortunately the specimens are all very small, therefore we can not know if this row of tubercles becomes stronger in larger specimens, or if it disappears completely, or if it persists in larger individuals in the same manner as in juvenile specimens. Hyatt does not mention the existence of a median row of tubercles in juvenile specimens of *Metoecoceras*. *Acanthoceras subvicinale* occurs in the Cenomanian of Diego-Suarez.

Thus we see that in Europe as well as in India and Madagascar there seem to exist forms which resemble the *Metoecoceras* of America. Probably we have to include in this genus at least *Pulchellia Gesliana* Petrascheck and perhaps also *Ammonites Geslianus* d'Orbigny. Very similar is also *Acanthoceras subvicinale* Boule, Lemoine et Thevenin, especially on account of the suture, while *Acanthoceras vicinale* Stoliczka, although showing a very similar ornamentation, differs by its more complicated suture. But we should note that the suture changes considerably in the different *Metoeceras* of America, that of *M. Whitei* being much more complicated than that of *M. Swallovi* (in the limitation of Hyatt). At present we can not decide how many forms should be included in the genus *Metoecoceras*, but can only indicate that this genus is probably represented in the Cenomanian of Europe and perhaps of Madagascar, and that similar forms but with a much more complicated suture, are found in the Cenomanian and Turonian of India.

METOECOCERAS aff. WHITEI Hyatt

Pl. 12 figures 4, 7

1876: *Buchiceras svallovi* White, Inv. foss. Nevada, Utah, etc., p. 202, pl. 20.

1893: *Buchiceras svallovi* Stanton, Colorado Form., p. 168, pl. 37, fig. 1 (?pl. 38, figs. 1-3).

1903: *Metoecoceras Whitei* Hyatt, Pseudocer., p. 122, pl. 13, figs. 3-5, pl. 14, figs. 1-10, fig. 15.

In the lower bed of Cerro del Macho occur two fragments of cephalopods which greatly resemble the group of *Metoecoceras Whitei*; they are especially similar to specimens figured by Hyatt. Our individuals show the following features:

Shell discoidal, very involute, with whorls of nearly rectangular cross section, much higher than broad. The flanks are very little convex, nearly flat, the ventral portion is flattened. The umbilicus is extremely narrow. The ornamentation consists of strong ribs which begin on the umbilical

border; they are slightly thickened near the umbilical border but do not form real nodules. Between these main ribs other secondary ones are intercalated; they begin above the umbilical border and are a little less strong on the flanks than the main ribs. The ribs are slightly bent forward on the flanks but do not considerably increase in strength. Near the ventral shoulder the ribs show a tuberculiform thickening and from this point bend more strongly forwards; they end on the ventral shoulder in a strong nodule which is longitudinally prolonged. Between the two rows of nodules on the ventral shoulders we observe an entirely flat and smooth zone of the venter; the ribs do not continue over the ventral portion, but every pair of the ventral nodules is connected by a slight thickening, which produces on the ventral zone a slight undulation.

The suture is only partly visible. We recognize the external saddle, the first lateral lobe, the first lateral saddle and the second lateral lobe. The suture is not very well preserved because this part of the specimen is somewhat corroded, but still we are able to note the general character. The external saddle (compare figure) is broad and divided into two parts by a rather deep secondary lobe; the first lateral lobe is broad and shallow, ending in two points, and on the sides are two rather large branches; the first lateral saddle is higher than the external one, its ramifications are few and simple; the second lateral lobe is much shallower than the first one, and it is also narrower and trifid. In this specimen the umbilical portion is not preserved but one sees that the second lateral saddle remains rather far from the umbilicus, therefore several auxiliary lobes and saddles must have existed in the intermediate space.

Figure 1. *Metoecoceras* aff. *whitei* Hyatt, suture

The ornamentation of our specimens coincides perfectly with that of *Metoecoceras Whitei* Hyatt, only on the ventral part we do not observe as

deep furrows as in Hyatt's plate 13, figure 3, but simply an undulation. As far as the suture is visible it corresponds with that of *M. Whitei*, especially with that of plate 14, figure 8 of Hyatt. I do not identify my specimens with *M. Whitei* because they are not sufficiently well preserved, but I am sure that they belong to the group of *M. Whitei*.

We know very little about the stratigraphic position of *M. Whitei*; some of Hyatt's originals have been found in the Kanab Valley, Utah. According to Walcott and Stanton¹ the species occurs there up to a height of 335 feet above the base of the Cretaceous. Stanton refers that division to his Colorado Formation which probably contains several horizons, perhaps from the upper Cenomanian, or at least from the lower Turonian to the Emschian. For the determination of the age of our beds the species found in the United States do not help us much. On the other hand, we do not know any European form which might be identified with our species, although this has a great similarity with *Pulchellia Gesliana* Petrascheck from the upper Cenomanian of Saxony, as we have shown in the stratigraphical part.

Number of specimens: 2.

Age: Upper Cenomanian (?), lower beds of Cerro del Macho.

METOECOCERAS sp. nov.

Pl. 12, figs. 1-3

The collection of Dr. Haarmann contains a specimen of *Metoecoceras* which is certainly different from the species described above; unfortunately we do not know in which horizon this specimen has been found; according to its petrographical character it belongs to Horizon I, but it might have been found also in Horizon II. Its features are:

Shell discoidal, very involute, with whorls of a lanceolate cross section truncated in the ventral part, much higher than broad. The flanks are a little convex, the ventral portion is flattened. The ornamentation consists of rather strong ribs, which are slightly flexuous and which bend forward. These begin near the umbilical border, where they are slightly thickened, but do not form nodules. Part of these ribs bifurcate below the middle of the height of the flank. Between these main ribs we find other secondary ones intercalated which begin in different heights of the flank, and which at their beginning are less strong than the main ribs. All the ribs show near the ventral shoulder a slight thickening of a kind of very low and rounded nodules; on the ventral shoulder the ribs end in a nodule which is longitudinally elongated. These nodules form together a kind of interrupted

¹Stanton, Colorado Form., p. 35.

keel on both sides of the ventral portion. Between the rows of nodules we observe on the ventral portion a rather broad, flat and smooth zone.

With respect to the involution we observe that the last whorl seems to show a tendency to form a wider curve than the preceding whorl.

No part of our specimen shows a trace of the suture.

According to its form and ornamentation our specimen belongs certainly to *Metoecoceras*, and I unite it provisionally with this genus although the suture is unknown.

The species differs from *M. Whitei* Hyatt *typus* in its flexuous ribs, but it resembles the specimens figured by Stanton¹ in figures 1-2 of plates 38 and called *Buchiceras Swallovi*. Hyatt unites it with his *Metoecoceras Whitei*, notwithstanding that it differs much from the type. It seems that our specimen has a smaller number of ribs than that figured by Stanton and that the ribs are more flexuous.

Our species resembles much more *Pulchellia Gesliniana* Petrascheck² than the above described species. This similarity is really very great but our species is distinguished by a smaller number of ribs which are also less flexuous. I do not doubt that both species belong to the same group. *Metoecoceras Geslinianum* Petrascheck occurs in the upper Cenomanian in Saxony.

Number of specimens: 1.

Age: Upper Cenomanian (?) or lower Turonian.

MAMMITES Laube et Bruder, Emend. Petrascheck

MAMMITES MOHOVANENSIS sp. nov.

Pl. 12, figs. 6, 8

In Horizon II of Cerro del Macho there has been found a well preserved specimen of *Mammites*, and also several fragments belonging to the same species. They show the following features:

Shell globose, rather involute, with whorls of a trapezoidal or nearly subquadrangular cross section, much broader than high. The umbilicus is rounded. On the umbilical border we observe six thick nodules, which are rather pointed and a little rounded. In these nodules begin broad, rounded, radial ribs, partly simple, partly in pairs. Between these ribs are intercalated generally two secondary ribs also low and rounded, which begin above the umbilical border, but without showing nodules at their beginning. On these secondary, as well as on the main, ribs we find two strong and rounded

¹Stanton, Colorado Form.

²Petrascheck, *Amm. d. sächs. Kreideform.*, p. 140 (10), pl. 7 (1), figs. 3-5.

nodules near and on the ventral part. These nodules form two rows, each of which is composed of 16 to 17 nodules on the last whorl; those of the row on the ventral portion are a little elongated longitudinally. Thus we find on each side of the shell three rows of nodules: the umbilical, the intermediate and the ventral row. It seems that the nodules of the intermediate row increase in thickness in the front part of the whorl, while those of the ventral row decrease a little and extend themselves more longitudinally, but none of the specimens is large enough to show the development of the nodules in an advanced age. Between the two ventral rows we observe a narrow concave zone in the inner whorl and a nearly flat one on the external whorl.

The suture is very much destroyed but its main features can still be recognized. The external lobe is rather deep and narrow, ending in two branches; the external saddle is broad and high, divided into two branches by an adventive lobe of relatively great depth; the first lateral lobe is deep and ends in two points; the first lateral saddle is apparently broad but its exact form can not be recognized because the shell is much destroyed in this part.

Dimensions:

Diameter	87.0 mm.	(1)
Height of the last whorl.....	33.0 mm.	0.38
Width of the last whorl.....	42.0 mm.	0.48
Diameter of the umbilicus.....	23.0 mm.	0.26
Height of the last whorl in the same diameter but half a whorl backwards	30.5 mm.	0.35
Width of the last whorl in the same diameter and half a whorl back- wards	39.0 mm.	0.45

Our species certainly belongs to the group of *Mammites nodosoides* Schlotheim. Its suture coincides in general fairly well with the one figured by Laube and Bruder¹, but its form and ornamentation resemble more that of *Mammites conciliatus* Stoliczka,² which differs from the type of the group especially by the greater thickness of the whorl and the more numerous nodules on the ventral portion.

Our species differs from *Mammites conciliatus* by its more involute form, the external whorl covering nearly two-thirds of the preceding one, barely leaving uncovered the row of umbilical tubercles. We find another difference in the number of intermediate and ventral nodules, which is much less in our specimen than in the species of Stoliczka. In this direction our individuals resemble more the type of *Mammites nodosoides*, but

¹Laube und Bruder, Amm. d. böhm. Kreide, p. 230.

²Stoliczka, Ceph. Cret. Rocks India, p. 99, pl. 50, fig. 4; pl. 51, fig. 1.

they differ from this as well as from *M. conciliatus* through the small distance between the two ventral rows of nodules. In these comparisons I take as type of the species *M. nodosoides*, the specimen figured by Laube and Bruder because this comes from the same locality as the originals of Schlotheim, which were never figured.

Similar to our specimens is also the small individual figured by Pervinquier¹ as a typical *M. nodosoides*.

Fritsch² figures a *Mammites* under the name of *Ammonites conciliatus* Stoliczka, and Kossmat³ confirms the conclusion that this specimen is very nearly related to the species from southern India; he notes that the main difference consists in the involution. In this direction the *Mammites conciliatus* Fritsch resembles more our species than that of Stoliczka. Though in this latter species the external whorl barely covers the preceding one up to the row of intermediate nodules, we observe that in the specimen figured by Fritsch the external whorl covers the preceding one nearly to the row of umbilical nodules.

The group of *Mammites nodosoides* has a quite universal distribution; it is represented in the lower Turonian of all Europe, of Africa, Asia, and South America. In North America we probably find it represented by *Ammonites Loewianus* White,⁴ an observation previously made by Pervinquier.

Number of specimens: 3.

Age: Lower Turonian (Salmurian), lower horizon.

PSEUDASPIDOCERAS Hyatt emend. Pervinquier

PSEUDASPIDOCERAS aff. FOOTEANUM Petrascheck

1902: *Mammites Footeanus* Stoliczka, Petrascheck, *Amm. d. sächs. Kreideform.*, p. 144, pl. 9, fig. 1.

Among the material collected by Dr. Haarmann in the lower horizon of the Turonian of Cerro del Macho is a rather large, not very well preserved but quite characteristic cephalopod, showing the following features:

Shell moderately evolute with whorls of a subrectangular-oval cross-section, much higher than broad. The flanks are flattened, the ventral portion rounded. The umbilicus is moderately wide, the umbilical wall is

¹Pervinquier, *Paléontologie Tunisienne*, pl. 18, fig. 1.

²Fritsch, *Ceph. böhm. Kreideform.*, pl. 7, figs. 1, 2.

³Kossmat, *Südind. Kreideform.*, p. 22 (129).

⁴Stanton, *Colorado Form.*, p. 178, pl. 43, figs. 3-4. Some authors write *Am. laevianus* but it must be *Loewianus* because the species was dedicated to Dr. Oscar Loew.

abrupt. The ornamentation consists in widely separated, low, rounded, nearly straight ribs which begin in a nodule at the umbilical border and which end in another very high and pointed nodule at the ventral shoulder. Between the two rows of nodules on the ventral part we observe a broad, slightly convex and smooth zone. In the last half whorl we count five ribs. The suture is not visible.

Dimensions:

Diameter	280 mm.	(1)
Height of the last whorl.....	117 mm.	0.42
Width of the last whorl.....	94 mm.	0.34
Diameter of the umbilicus.....	104 mm.	0.37

Our specimen certainly belongs to the group of *Pseudaspidoceras Footeanum* Stoliczka,² but differs from it by its cross-section which is much higher than wide. This is not an absolutely distinctive character because we should take into consideration the large size of our specimen and we do not know the latest development of the species of Stoliczka. Our specimen resembles *Pseudaspidoceras Footeanum* Petrascheck more than it does the type. The cross-section of the species of Petrascheck is more similar to ours and also has widely separated ribs.

The group of *Pseudaspidoceras Footeanum* Stoliczka has a very wide distribution; it is represented in all the northern part of Africa (we include in this group *Ps. salmuriense*), in Nigeria, in Portugal, in Saxony, and in India.

Number of specimens: 1.

Age: Lower Turonian (Salmurian), lower horizon.

PSEUDASPIDOCERAS aff. PEDROANUM White

Pl. 13, fig. 1; Pl. 15, fig. 1

1887: *Ammonites Pedroanus* White, Brazil, p. 212, pl. 22, figs. 1, 2 (not pl. 20, fig. 3).

In the lower horizon of the Turonian of Cerro del Macho we find a large and relatively well preserved specimen which belongs to the group of *Pseudaspidoceras Footeanum* Stoliczka. Its features are:

Shell large with whorls of a subquadrangular cross-section, nearly as high as wide. The flanks are flattened, the ventral portion is nearly flat. The umbilicus is relatively wide and deep, the umbilical wall is vertical but without a sharp umbilical shoulder. The external whorl covers only the ventral portion of the preceding one and leaves the nodules of the ventral shoulder free. The ornamentation consists of extremely low and

²Stoliczka, Ceph. Cret. Rocks India, p. 101, pl. 52, figs. 1, 2.

rounded, nearly straight ribs which sometimes are scarcely perceptible; on the last whorl we count thirteen of these ribs. The ribs begin on the umbilical border in a strong and pointed nodule and end with another and still more prominent but somewhat longitudinally elongated nodule on the ventral shoulder. Between the two rows on the ventral shoulders we observe a smooth and very slightly convex zone which does not show any signs of nodules or ribs; only the two last nodules seem to be connected by a kind of low rounded rib which crosses the smooth zone of the venter. Between the nodules of the rows on the ventral shoulder we find some ribs or nodules intercalated which do not correspond to umbilical nodules.

Traces of the suture are visible in different parts but I have not been able to draw a complete line.

Dimensions:

Diameter: 214 mm.

The last part of our specimen being somewhat crushed and twisted, we have taken the following dimensions from a little smaller diameter:

Diameter	164 mm.	(1)
Height of the last whorl.....	63 mm.	0.38
Width of the last whorl.....	70 mm.	0.43
Diameter of the umbilicus.....	59 mm.	0.36
Height of the last whorl on the same diameter but half a whorl back- wards	42 mm.	0.26
Width of the last whorl on the same point.....	45 mm.	0.27

Our specimen belongs, as we have already mentioned, to the group of *Pseudaspidoceras Footeanum* Stoliczka¹ which it resembles in its involution, the cross-section of the whorls, the two series of prominent nodules, etc. It differs from the type through its weaker ribs, and the entire absence of ribs and nodules on the ventral portion and of a concave zone along the line of symmetry on the ventral portion.

Our species shows much nearer relations to *Pseudaspidoceras pedroanum* White from Brazil than to the type of the group. It has the same kind of involution, a very similar cross-section, a nearly identical ornamentation; it only seems to differ a little with respect to the nodules on the ventral shoulder, which according to the figure of White are connected on the last part of the whorl by a kind of low crest, which is also mentioned in the text. White also says that in very large individuals the nodules on both sides of the venter are connected by a crest which crosses the venter. We have indicated a similar phenomenon in the last two nodules of our specimen. We thus see that our species resembles in a surprising manner *Ammonites pedroanus* Kossmat and others have already recognized

that this species belongs to the group of *Pseudaspidoceras Footeanum*. White compares his species with *Ammonites Leonensis* Conrad, which is probably a *Schloenbachia*, and also with *Prionotropis Woolgari*.

We know nothing about the age of the beds where the type of the species has been found but the similarity with *Pseudaspidoceras Footeanum* allows us to conclude that they probably belong to the Salmurian. In this idea we are confirmed by the circumstance that in the same locality *Vascoceras Harttii* Hyatt has been discovered.

Number of specimens: 1.

Age: Lower Turonian (Salmurian), lower horizon.

FAGESIA Pervinquière

FAGESIA HAARMANNI sp. nov.

Pl. 14, figs. 1, 2; Pl. 15, fig. 2

In the lower bed of the Salmurian of Cerro del Macho Dr. Haarmann found a not quite complete specimen of so characteristic a *Fagesia* that I have no hesitation in giving it a new name. Another more complete but somewhat deformed specimen collected by Dr. E. Wittich, was given to me by Mr. Federico Ritter. The features of the larger specimen are:

Shell moderately involute with whorls of transversally elliptical cross-section, much broader than high. The umbilicus is rather narrow and deep; the umbilical wall is abrupt, in the internal whorls nearly vertical. The ventral portion is broad and slightly convex, the region of the flanks is very narrow and one might say scarcely developed and in reality only represented by the umbilical nodules; the ventral part nearly touches the umbilical wall. The ornamentation consists on the umbilical border of some nine very strong, pointed and round nodules. From these nodules start rounded, broad, rather high ribs which cross the ventral portion without interruption in a curve directed toward the front; in general a pair of ribs starts from each nodule and one or two more become intercalated between every two pairs. This disposition can not be observed with entire certainty because the border of the whorl is a little broken.

The suture could not be made visible.

Dimensions:

Diameter	165 mm.	(1)
Height of the last whorl.....	52 mm.	0.32
Width of the last whorl.....	74 mm.	0.45
Diameter of the umbilicus.....	66 mm.	0.40

At the first view our species resembles greatly a *Stephanoceras*, a similarity which has been noted also in other species of *Fagesia* especially *Fagesia superstes* Kossmat.¹ From this species, which represents the type of the genus, our species differs by being much more evolute and by its stronger ribs; the umbilical nodules are less numerous and thicker, the ribs are also less numerous and a little stronger.

Our species is certainly the most evolute one of all the *Fagesia* so far described, but its nearest relative is without doubt *Fagesia superstes* Kossmat. According to the figure published by Kossmat no nodules can be seen on the internal whorls, while in our species these nodules can be distinguished even on the smallest whorls.

The occurrence of this species is of certain stratigraphical importance in so far as it belongs to a very characteristic group which seems at present to be limited to the Turonian. According to Pervinquire only six species are known and all of these belong to the Turonian. *Fagesia superstes*, which resembles ours more than any other species, has been found in India (upper Utatur group) and in Tunis (lower Turonian); similar forms have been discovered in the Turonian of Portugal and France.²

Number of specimens: 2.

Age: Lower Turonian (Salmurian), lower horizon.

FAGESIA PERVINQUIERI sp. nov.

Pl. 14, fig. 3

In the lower horizon of our Salmurian we find a relatively well preserved specimen which belongs to *Fagesia* but differs essentially from the foregoing species by its entirely different form and the weaker ornamentation. Its general features are:

Shell subglobose, very involute, with whorls of transversally semilunar cross-section. The umbilicus is narrow and deep, the umbilical wall is nearly vertical. On the umbilical border of the internal whorls appear not very strong nodules; on the last part of the external whorl the umbilical border seems to be entirely smooth. On the internal whorl we observe numerous broad and low ribs which are strongest in the central part of the ventral portion, while they seem to flatten out toward the umbilical border. On the ventral portion the ribs curve slightly toward the front. The number of ribs could not be counted because the ventral

¹Kossmat, Sdind. Kreideform., p. 26 (133), pl. 6 (17), fig. 1.

²Pervinquire, Palontologie Tunisienne, p. 325, note 5.

portion of the last whorl is a little corroded, but there are certainly more than forty on the last whorl. The suture is not visible.

Dimensions:

Diameter	94 mm.	(1)
Height of the last whorl.....	36 mm.	0.40
Width of the last whorl.....	60 mm.	.0.64
Diameter of the umbilicus.....	31 mm.	0.33
Height of the last whorl on the same diameter but half a whorl backwards	27 mm.	0.29
Width of the last whorl at the same point.....	55 mm.	0.59

On account of its general form and its ornamentation I place this specimen in the genus *Fagesia* although the suture could not be made visible. Our species is quite similar to *Fagesia tevesthensis* Peron¹ especially the large specimen figured by Pervinquière² (pl. 20, fig. 6). This individual shows on the umbilical border nodules quite similar to those of our specimen. But the African species has much broader and lower whorls which in part may be explained by its stage of development, because all the species of *Fagesia* change their cross-section of the whorls with their age and grow more rapidly in width than in height.

Another species which resembles ours is *Fagesia rudra* Stoliczka.³ This species shows very similar ribs and the umbilical border is entirely smooth. *Fagesia rudra* differs from our species by its much broader and lower whorls.

Unfortunately only a single specimen of our species has been found. It was impossible to study the internal whorls in their details and therefore we can not make more detailed comparisons with other species.

Number of specimens: 1.

Age: Lower Turonian (Salmurian), lower horizon.

VASCOCERAS Choffat

When Choffat⁴ established his new genus *Vascoceras* he subdivided it into four groups: 1. mono-tuberculated forms with wide umbilicus; 2. subglobose forms with rounded umbilical border; 3. globose forms with angular or subangular umbilical border; 4. multi-tuberculated forms. These groups could be called: 1. Group of *Vascoceras Gamai*; 2. Group

¹Peron, Amm. du Crét. sup. de l'Algérie, p. 23, pl. 7, figs. 2, 3.

²Pervinquière, Paléontologie Tunisienne, p. 325, pl. 20, figs. 5, 6.

³Stoliczka, Ceph. Cret. Rocks India, p. 122, pl. 60, fig. 1.

⁴Choffat, Espèces nouv. ou peu conn., p. 51.

of *Vascoceras Douvillei*; 3. Group of *Vascoceras harttiiforme*; 4. Group of *Vascoceras subconciliatum*.

Pervinquièr¹ unites the first two groups in one and distinguishes only three subdivisions. This seems to me to be the best solution, as the division of the first two groups appears to be rather artificial.

Among the material of Cerro del Macho we find forms representing the three groups. To the first one (in the sense of Pervinquièr) belongs our *Vascoceras* aff. *Gamai* Choffat and *Vascoceras* aff. *adonense*; to the second one *Vascoceras Angermanni* sp. nov., and perhaps *Vascoceras* aff. *arnesense* Choffat; to the third one *Vascoceras mohovanense* n. sp.

The genus *Vascoceras* so far has been found in Portugal, Spain, Algeria, Tunis, Egypt, the Sahara, Nigeria, Brazil, Peru, and Mexico.

VASCOCERAS sp. nov. ex aff. **V. ADONENSE** Choffat

Pl. 14, fig. 4; Pl. 17, fig. 2

1886: *Vascoceras adonense* Choffat, *Espèces nouv. ou peu conn.*, p. 59, pl. 9, fig. 3; pl. 21, fig. 12.

Our specimen shows the following features:

Internal mold rather large, a little oblique on account of deformation, quite evolute, with whorls of a transversally semi-lunar cross-section, much broader than high. The umbilicus is wide, the umbilical wall is quite abrupt, the umbilical border is rounded and scarcely subangular. In the external whorl no nodules exist on the umbilical border; the internal whorls are not visible. The surface of the species is smooth, no ornamentation can be distinguished.

The suture is very simple; it is not very well preserved but the general features can be well distinguished. The external lobe is quite wide and not very deep and ends in two branches; the external saddle is wide at the base, becoming narrower upward, and is very little ramified; the first lateral lobe is funnel shaped, very wide at the mouth; the first lateral saddle is similar to the external one but smaller and inclined toward the umbilicus while the external one rather bends over toward the venter; the second lateral lobe is funnel shaped, very wide at the mouth and much shallower than the first one; the second lateral saddle (on the umbilical border) is similar to the first one but smaller.

The dimensions could not be measured exactly on account of the deformation of the specimen.

¹Pervinquièr, *Paléontologie Tunisienne*, p. 331.

With respect to its external form our species resembles *Vascoceras adonense* Choffat, but this latter species shows nodules on its umbilical border, while these are entirely missing in our specimen. This might be explained by the stage of development, our individual being much larger than that of Choffat, but our species is certainly much more evolute. A

Figure 2. Suture, *Vascoceras angermanni* n. sp. (above) and *Vascoceras* aff. *adonense* Choffat (below).

fundamental difference can be found in the suture. In the type species we find on the umbilical border the first lateral saddle, but we see that in our specimen the second lateral saddle is in this place. In its general character our suture is much more similar to that of *Vascoceras Gamai*.

Our species therefore has to be considered as a new form which in its external figure resembles *Vascoceras adonense* while the suture places it in the vicinity of *Vascoceras Gamai*. We can not give it a new name because the specimen is too badly preserved.

Number of specimens: 1.

Age: Lower Turonian (Salmurian), lower horizon.

VASCOCERAS aff. GAMAI Choffat

Pl. 15, figs. 3-5

1886: *Vascoceras Gamai* Choffat, *Espèces nouv. ou peu conn.*, p. 54, pl. 7, figs. 1-4; pl. 8, fig. 1; pl. 10, fig. 2; pl. 21, figs. 1-5.

In the collection made by Dr. Angermann I found a cephalopod which probably represents a juvenile stage of some species belonging to the group of *Vascoceras Gamai*. Judging from the petrographical aspect of the specimen I suppose that it was found in the upper horizon of the Salmurian of Cerro del Macho. Its features are:

Shell relatively small, quite involute with whorls of a subquadrangular rounded cross-section, broader than high. The umbilicus is narrow and probably deep, the umbilical wall is abrupt. The flanks are flattened, the ventral part rounded. On the umbilical border we count seven thick, round, and pointed nodules. From each of these starts either a pair of ribs or a simple rib, which curves slightly forward and crosses without interruption the ventral portion. Between these main ribs one or two secondary ones are intercalated. These begin above the umbilical border, but on the venter are as strong as the main ones. The ribs are thick, broad, rounded, wider than the interstices by which they are separated. On the ventral shoulders the ribs seem to have a tendency toward a slight thickening but real nodules do not form there. On the anterior part of the last whorl, and especially on the ventral region, the ribs seem to disappear. This region is partially corroded but in a place which is well preserved one notes clearly that the ribs are very low and scarcely perceptible on the line of symmetry.

Figure 3. Suture, *Vascoceras* aff. *gamai* Choffat

The suture is very little ramified. The external lobe has two branches and is quite deep and narrow; the external saddle is little ramified, broad and rounded; the first lateral lobe is less deep than the external one and rather narrow. The first lateral saddle is little ramified and resembles in its form the external one, but is narrower.

Our specimen resembles the juvenile form figured by Choffat in figure 4

of plate 7; unfortunately no figure of the form viewed from the side has been published. The juvenile specimen of figure 3 of plate 7, the lateral view of which is given in figure 2 of plate 10 of Choffat is also similar. The main difference exists probably in the smaller number of umbilical nodules in our specimen and in the narrower cross-section, although this latter feature changes with the age.

In the juvenile specimens of *Vascoceras Mundae* the ribs are less numerous and thicker.

Vascoceras Gamai Choffat occurs in Portugal mainly in the lower and middle Turonian of Choffat, both of which according to my opinion belong to the Salmurian.

Number of specimens: 1.

Age: Lower Turonian (Salmurian), probably from the upper horizon.

VASCOCERAS ANGERMANNI sp. nov.

Pl. 16, figs. 1-4; Pl. 17, fig. 1

In the collection of Dr. Angermann as well as in that of Dr. Haarmann we find a globose cephalopod which at the first view resembles completely a Nautilus. In the collection of Angermann there is a small, quite well preserved specimen, and in Mr. Haarmann's collection a much larger one, but the features of both are the same. There are also a number of fragments which probably should be united with this singular species. Its features are:

Shell large, globose, extremely involute, with whorls of semi-lunar cross-section, much broader than high. The whorls cover the preceding ones nearly entirely. The umbilicus is narrow and deep, the umbilical wall abrupt, the umbilical border smooth and angular. The surface of the shell is entirely smooth without any nodules or ribs.

The suture is not very well preserved but its general features can be well recognized. The external lobe is quite deep and moderately wide, ending in two branches. The external saddle is moderately broad, quite high, very little ramified; the first lateral lobe is little ramified, broad, ends in two points and is deeper than the external one; the first lateral saddle is broad and rounded, less high than the external saddle. The second lateral lobe is probably similar to the first but less deep; it is not entirely preserved.

Dimensions of the smaller specimen:

Dimensions:

Diameter	75 mm.	(1)
Height of the last whorl.....	32 mm.	0.45
Width of the last whorl.....	69 mm.	0.92
Diameter of the umbilicus.....	14 mm.	0.19
Height of the last whorl on the same diameter but half a whorl backwards	27 mm.	0.36
Width of the last whorl at the same point.....	62 mm.	0.83
Diameter of the largest specimen: 124 mm.		

Our species resembles greatly *Vascoceras Kossmati* Choffat¹; it is of course evident that all the species belonging to this group must be very similar, as none of them shows any ornamentation. The main difference between our species and *Vascoceras Kossmati* consists in the different curve of the cross section of the internal whorls, which is always of greater radius. The suture is also different, the saddles of our species being relatively higher and the lobes deeper. One might add that our species grows to a larger size than the Portuguese species.

Vascoceras Kossmati Choffat has been found in Choffat's middle Turonian of Portugal, which, according to my opinion, belongs to the Salmurian. In Egypt it occurs in the lower Turonian.² Our larger specimens have been found in the upper bed of the Salmurian of Cerro del Macho and the specimen collected by Angermann belongs to the same horizon, judging from its petrographical character.

Number of specimens: 3 and several fragments.

Age: Lower Turonian (Salmurian), upper horizon.

VASCOCERAS sp.

Pl. 18, fig. 12

Dr. Haarmann found in the upper horizon of Cerro del Macho a cephalopod, the generic determination of which remains somewhat in doubt because its suture can not be made visible. Mr. Frederico Ritter gave me another and a little better preserved specimen which shows traces of the suture. Their features are:

Shell subglobose of moderate size, very involute with whorls of semi-lunar cross section, broader than high. Flanks and ventral portion rounded. The umbilicus is small and deep, the umbilical wall is vertical and even a little overhanging; the umbilical border is angular. The shell

¹Choffat, *Espèces nouv. ou peu conn.*, p. 63, pl. 13, figs. 8, 9; pl. 14, figs. 1, 2; pl. 21, figs. 26, 27.

²Eck, *Turon in Aegypten*, p. 384.

seems to be entirely smooth, no nodules can be observed on the umbilical border. The suture is similar to that of the group of *Vascoceras Kossmati* but it is not complete enough for a drawing.

Dimensions:

Diameter	106 mm.	(1)
Height of the last whorl.....	43 mm.	0.41
Width of the last whorl.....	72 mm.	0.68
Diameter of the umbilicus.....	22 mm.	0.21

One of the specimens is a little obliquely deformed which makes the determination still more difficult. The species resembles externally much *Ammonites arnesensis* Choffat¹, the generic position of which is equally doubtful; it seems to me to belong to *Vascoceras* or to a new subgenous, the suture being quite different from that of the type of the genus.

In one of our specimens the suture is not visible; in the other one we see parts of it which are similar to the elements of the suture of *Vascoceras Angermanni*. Therefore I place our species provisionally in the genus *Vascoceras*. Externally this species resembles somewhat *Vascoceras Angermanni* but the umbilicus is wider, the general form less globose, and the whorls are less broad in relation to their height.

Number of specimens: 2.

Age: Lower Turonian (Salmurian), upper horizon.

VASCOCERAS MOHOVANENSE sp. nov.

Pl. 18, figs. 1-2

Dr. Haarmann collected in the upper bed of our Salmurian a juvenile individual of a cephalopod which certainly represents a new species. Its features are:

Shell small, subglobose, quite involute with whorls of trapezoidal cross section, much broader than high. The flanks are somewhat flattened, as is also the ventral part. This gives the cross-section its angular form. The umbilicus is narrow and deep, the umbilical wall is abrupt, the umbilical border is angular. The ornamentation consists in the last whorl of about eleven high and pointed nodules (nearly all broken in the present specimen) on the umbilical border; from each of these generally starts a pair of ribs. Between this pair of ribs now and then secondary ribs are intercalated which begin above the umbilical border. The ribs are thick

¹Choffat, *Espèces nouv. ou peu conn.*, p. 68, pl. 13, fig. 10; pl. 14, fig. 3, pl. 22, fig. 39.

and high. They become a little lower on the venter but cross it without interruption; no longitudinal furrow exists on the venter. On the ventral shoulders the ribs thicken a little, forming on each shoulder a row of radially elongated nodules; these are not very strong and rather low. These nodules make the ventral shoulders appear angular and the ventral zone more flattened than concave.

Figure 4. Sutures, *Vascoceras mohovanense* n. sp. (above) and *Mammites mohovanensis* n. sp. (below).

The suture is very simple. The external lobe is quite wide and deep, and bifid; the external saddle is broad, low, or a rounded form and a little ramified; the first lateral lobe is wide, approximately as deep as the external lobe but much wider, and it ends in two points; the first lateral saddle is very low, rounded and broad and lies in the umbilical nodule; the form of the second lateral lobe, which lies mostly on the umbilical wall, can not be well distinguished.

Dimensions:

Diameter	56 mm.	(1)
Height of the last whorl.....	21 mm.	0.38
Width of the last whorl.....	45 mm.	0.80 (with the nodule)
Diameter of the umbilicus.....	19 mm.	0.34
Height of the last whorl on the same diameter but half a whorl backwards.....	16 mm.	0.29
Width of the last whorl at the same point.....	34 mm.	0.61

This very characteristic species certainly belongs to the group of *Vascoceras subconciliatum* Choffat¹, but it resembles *Vascoceras polymorphum*

¹Choffat, *Espèces nouv. ou peu conn.*, p. 64, pl. 15, figs. 1-3; pl. 16, fig. 4; pl. 22, figs. 28-31.

Pervinquièr¹ more than it does the type. From both species it differs through the absence of an intermediate row of nodules and consequently through the different cross section of the whorl. From *Vascoceras subconciliatum* our species is also distinguished by its very reduced and simple suture. In this case it resembles much more *Vascoceras polymorphum* Pervinquièr, which shows a very similar suture.

Pervinquièr has extensively discussed the relations of *Vascoceras polymorphum* with the other *Vascoceras*, as well as with *Mammites* and *Aconthoceras*, indicating the similarity of certain groups of *Aconthoceras* and *Mammites*, and emphasizing the singular nature of the suture, which is similar to that of other *Vascoceras*. I believe that Pervinquièr's reasons are well founded and that the present group should be placed in the genus *Vascoceras*, especially as this genus is nearly related to *Acanthoceras* and *Mammites*; therefore the similarity between certain groups of the three genera is not very surprising.

Vascoceras polymorphum Pervinquièr occurs in the lower Turonian of Tunis; *Vascoceras subconciliatum* Choffat is found in beds of a similar age in Portugal. Pervinquièr also emphasizes the similarity between this group and a cephalopod from Brazil, *Ammonites offaccinatus* White,² but this latter species is too little known to allow a detailed comparison. The suture is not figured but according to White it is rather complicated. It is therefore possible that it belongs to a different group, or even to a different genus.

Number of specimens: 1.

Age: Lower Turonian (Salmurian), lower horizon.

NEOPTYCHITES Kossmat

NEOPTYCHITES aff. CEPHALOTUS Courtiller

Pl. 18, figs. 3, 10, 13

1907: *Neptychites cephalotus* Courtiller Pervinquièr, Paléontologie Tunisienne, p. 393, pl. 27, figs. 1-3. Cum syn.

Among our material exists a cephalopod of small size which in its external form and its suture resembles the juvenile specimens of *Neptychites*. Its features are:

Shell subglobose, very involute, last whorl broader than high with nearly

¹Pervinquièr, Paléontologie Tunisienne, p. 336, pl. 21, figs. 2-5.

²White, Brazil, p. 219, pl. 23, figs. 3, 4.

semilunar cross-section. The umbilicus is very narrow; the wall can not be seen. The flanks are slightly convex, the venter is rounded. The ornamentation consists of low, rounded, broad, numerous ribs (about 32 to the whorl) which apparently do not reach down to the umbilical region and which are strongest on the ventral portion. On the internal whorl we note a well defined and rather broad constriction which toward the front is limited by a slight thickening; the constriction is especially evident on the venter. The ribs are curved forwards, as is the case with the constriction, although in less degree.

Figure 5. Suture, *Neoptychites* aff. *cephalotus* Courtillet

The suture is very similar to that of *Neoptychites* and corresponds in general to that of *Neoptychites cephalotus*, although the figure published by Pervinquire is taken from a much larger specimen; a little more similar still is the suture of the small specimen of *N. xetiformis*, in the above cited work of Pervinquire (p. 398, fig. 153), although there the external saddle is more slender than in our specimen. The suture of our individual is not complete and in some places can not be very well followed, but in general narrow, ending in two branches; the external saddle is quite broad and one recognizes the main features. The external lobe is deep and relatively high and little ramified; the first lateral lobe is a little shallower than the external one, ends in two points and shows also a branch directed obliquely toward the ventral region; the first lateral saddle is similar to the external one but is lower and narrower.

Figure 6. Suture, *Neoptychites* aff. *cephalotus* Courtillet

Dimensions:

Diameter	50 mm.	(1)
Height of the last whorl.....	26 mm.	0.52
Width of the last whorl.....	33 mm.	0.66

Our specimen represents the juvenile stage of a *Neoptychites* similar to the one figured by Pervinquire in pl. 27 fig. 2, which it resembles much in its external features; it differs mainly in its great width. But according to Pervinquire the dimensions change much in the genus, and we should not give too much importance to this feature. A specific determination of our

specimen is not possible on account of its small size; we can only say that it is a *Neoptychites* similar to *Neoptychites cephalotus* but probably a different species.

Neoptychites cephalotus occurs in the lower Turonian of Tunis, Algeria and France. Grossouvre proposes to unite it with *Neoptychites Telinga* Stoliczka and Pervinquire accepts this opinion. There is no doubt that the adult individuals of the two species resemble each other very much, although the suture seems to be somewhat different and a little more ramified in *N. cephalotus*; but this last character is not decisive as long as we know only the suture of a very large specimen of *N. Telinga*. A definite result can only be reached through the study of the development of *N. Telinga* or at least of that of small specimens.

Number of specimens: 1.

Age: Lower Turonian (Salmurian), upper horizon.

NEOPTYCHITES aff. XETRIFORMIS Pervinquire

Pl. 18, figs. 9, 11

1907: *Neoptychite xetriformis* Pervinquire, Palontologie Tunisienne, p. 398, pl. 27, figs. 5-7.

Among the material collected by Dr. Angermann is a very typical *Neoptychites*. The specimen is sufficiently well preserved for a generic determination. Its features are:

Shell discoidal, very involute, with whorls of nearly sagittiform cross section, much higher than broad. The greatest width is near the umbilical border. The flanks are quite flattened and very little convex, and the venter is rounded. The umbilicus is extremely narrow and deep; the umbilical wall is vertical; apparently a slight thickening exists on the umbilical border, at least in the last part of the external whorl. On the flanks we observe radial, broad, low, and rounded ribs which are strongest in the region near the venter and which decrease in strength toward the umbilical as well as toward the ventral region; they disappear near the umbilicus. On the venter the ribs are very low and scarcely discernible; near the ventral portion, they bend forward. The number of ribs could not be counted because part of the surface is corroded.

The suture is not very well preserved but the main features can be recognized. The external lobe is relatively wide and shallow, ending in two short branches; the external saddle is not very high nor very broad but little ramified. The first lateral lobe is wide and ends in two points; it also has a branch on each side, and is deeper than the external lobe. The first

lateral saddle is narrow, a little lower than the external saddle, and little ramified; second lateral lobe is similar to the first in its form but smaller and much shallower; the second lateral saddle is very low and very broad and of oblique form.

Dimensions:

Diameter	73 mm.	(1)
Height of the last whorl.....	40 mm.	0.55
Width of the last whorl.....	33 mm.	0.45
Diameter of the umbilicus.....	6.5 mm.	0.09

With respect to its external form our specimen is very similar to *Neoptychites xetrisformis*; even the dimensions coincide sufficiently, if we take into consideration the variability of *Neoptychites*. The main difference consists in the ornamentation; the ribs are much more numerous in our species (probably 25 to 30 on the last whorl) and they do not disappear completely on the venter. The specimen thus occupies an intermediate position between *N. cephalotus* and *N. xetrisformis*, or between this and *N. xetra*, if the small individual figured by Stoliczka¹ really belongs to *N. Xetra*. The suture of our specimen resembles that of *N. Xetra* still more than that of *N. xetrisformis*.

N. xetrisformis occurs in the lower Turonian of Tunis. *N. Xetra* Stoliczka belongs to the lower and middle Utatur group according to Kossmat, but we should not forget that the determination of the horizons of the Utatur group is in general based on the petrographical character of the fossils and on the locality. The determination of the age is therefore not always certain for each species. How doubtful it sometimes is we can see in our own case. On page 168 (72) Kossmat cites *N. Xetra* from the lower and middle Utatur group; on page 196 (131) he cites it from the lower and upper Utatur group. This doubt with respect to the age of the different species can not surprise us, because Kossmat could not base his determinations on well divided and stratigraphically well studied cross sections, but rather on general observations and subdivisions based more on the petrographical character of the rocks than on the faunas. Kossmat himself (loc. cit. p. 132 (197)) is convinced that a paleontological and stratigraphical subdivision of the Utatur group is possible, but until this has been executed in a modern manner, we should not give too much importance to the age which has been assigned to a certain species from India. We should always take into account that in such old collections labels

¹Stoliczka, Ceph. Cret. Rocks India, pl. 61, fig. 2.

may have been changed and that the fossils frequently have not been collected with the necessary exactness.

Number of specimens: 1.

Age: Lower Turonian (Salmurian), lower horizon.

HOPLITOIDES V. Koenen Emend. Solger et Pervinquière

The genus *Hoplitoides* has been established by von Koenen¹ for certain ammonites from the limestones of the Mungo in Kamerun; later Solger² studied the same fauna and made the definition of the genus more precise. Then Pervinquière¹ amplified this definition still more and distinguished two groups within the genus *Hoplitoides*: the bicarinate and the monocarinate forms. The first group seems to be limited to the Turonian, the second one to the Emscherian.

Among our material we find a few ammonites which very probably belong to the first group, the bicarinate *Hoplitoides*. The specimens are not very well preserved and the generic determination is not entirely certain. The external form corresponds completely with that of the *Hoplitoides*, but the suture is much destroyed and scarcely permits recognition of the general character of the line; this, however, coincides entirely with the suture of the bicarinate *Hoplitoides*. The character of the suture on the other hand shows that the specimens do not belong to *Placenticeras*, a genus which externally is quite similar to *Hoplitoides*.

HOPLITOIDES aff. **MIRABILIS** Pervinquière

Pl. 19, figs. 1-3

1907: *Hoplitoides mirabilis* Pervinquière, Paléontologie Tunisienne, p. 218, pl. 10, fig. 3

Among the material placed at my disposition I found two relatively complete specimens and four fragments which, according to their suture and external form, belong to the so-called Pseudoceratites of the Cretaceous. The best preserved specimen shows the following features:

Shell discoidal, entirely evolute, with whorls of lanceolate cross section truncated at the point, much higher than broad. The umbilicus is very narrow, the umbilical wall is vertical and the umbilical border rounded. The flanks are smooth, very little convex, nearly flat. The venter is flat with a sharp shoulder on each side; on the internal whorl we observe on both shoulders of the venter a sharp keel and the zone there is concave.

¹v. Koenen, Nachtr. Foss. Mungo i. Kamerun, p. 53.

²Solger, Mungokreide, p. 127.

The suture is badly preserved, because all the specimens are corroded on the surface, and no suture could be made visible on the internal whorls. We note a little ramified external saddle, which is rather broad and low; the first lateral lobe is broad, shallow and ends in three points, those on the umbilical side being longer than the one on the ventral side. Some smaller ramifications exist on the sides of this lobe. The first lateral

Figure 7. Sutures, *Neoptychites* aff. *xetiformis* Pervinquière (lower left hand corner) and *Hoplitoides* sp. (other three).

saddle is broad and relatively high, and a secondary lobe divides it in two parts. The following saddles are much destroyed and no longer show their ramifications. The lobes are much shallower than the first lateral lobe.

Dimensions:

Diameter	149 mm. (1)	119 mm. (1)
Height of the last whorl.....	76 mm. 0.51	58 mm. 0.49
Width of the last whorl.....	45 mm. 0.30	34 mm. 0.29
Diameter of the umbilicus.....	19 mm. 0.13	18 mm.? 0.15

Our specimens belong very probably to the genus *Hoplitoides* and to the bicarinate group of it. Externally they resemble this group in an extraordinary manner and especially *Hoplitoides mirabilis* Pervinquière. Pervinquière has united these forms, which distinguish themselves by a flat venter even in a very advanced stage, with the *Hoplitoides* of the Emscherian, which show a flat venter only in their youth, the venter in the adult age becoming sharp or a little rounded. Only the study of a large and

rich material could decide if it would not be better to separate these two groups at least subgenerically. It was probably the want of sufficient material which decided Pervinquière not to separate the two Turonian species from the rest of the genus. If such study should prove that all the species from the Turonian belong to the bicarinate group, it would be better to unite them in a different species.

The main difference between our species and *H. mirabilis* consists probably in the larger diameter of the umbilicus, but our material is not sufficiently well preserved to allow a more detailed comparison.

The two species of bicarinate *Hoplitoides* described by Pervinquière occur in the lower Turonian; one of them has also been found in the upper Turonian (*H. Munieri*).

Eck¹ cites with some doubt *Hoplitoides mirabilis*, or a similar form, from the Turonian of Egypt. Woods² describes a *Hoplites Nigeriensis* from Nigeria which he takes to belong to the bicarinate *Hoplitoides*. These specimens occur in probably Turonian beds. In the same locality have been found *Vascoceras Gongilense* Woods and *Pseudaspidoceras* sp. Woods mentions that *Hoplitoides* has also been found in another locality of Nigeria.

Number of specimens: 2, and 4 fragments.

Age: Lower Turonian (Salmurian), upper horizon.

LAMELLIBRANCHIATA

AVICULA Bruguière

AVICULA AGUILERAE sp. nov.

Pl. 20, figs. 1-2, 11-12

The material from the upper horizon of the Salmurian of Cerro del Macho contains a number of specimens of a large and well characterized *Avicula*. Its features are:

Shell thin, of subquadrate form, nearly as high as broad, not very convex; the right valve is a little more convex than the left one. On the anterior side is a prolongation in form of an auricula which is not quite completely preserved but which certainly was not very long; on the posterior side we note a prolongation in form of a wing, with a precise limit between this wing and the rest of the shell. The auricula as well as the wing

¹Eck, Turon in Aegypten, p. 382 and 386.

²Woods, Cret. dep. Northern Nigeria, p. 284, pl. 23, fig. 3; pl. 24, figs. 1-5; fig. 1 of the text.

form with their superior border approximately the prolongation of the cardinal border and do not elevate themselves above the beaks. The beaks are small and pointed. The cardinal border is straight, long, and vertical in relation to the longitudinal axis of the shell. The surface of the shell is entirely smooth and shows only fine concentric striae of growth. All our specimens are internal molds but some have preserved remains of the shell.

Our very characteristic species resembles *Avicula gravida* Coquand.¹ The main difference seems to consist in the greater convexity of the valves in this latter species, also in the different position of the beaks in relation to the upper border of the auricula and the cardinal border; but it is possible that this last difference may be explained as an error of the draftsman, because Coquand mentions expressly that neither the auricula nor the posterior wing elevate themselves above the beaks. In every case our species is nearly related to the African one.

Coquand mentions that his species occurs in the Turonian (Mornasian); later on Peron² corrected this determination of the horizon. He says that he has always found this species in the Santonian, at the base of the Senonian, in the beds with *Buchiceras* and *Hemiaster Fourneli*. But notwithstanding this correction it is possible that Coquand was right in his determination of the horizon, because Pervinquière³ also mentions that he has found *Avicula gravida* in the lower Turonian. The species seems to occur also in the Emscherian, from which horizon it is cited by Pervinquière⁴ who has also found it at the base of the Senonian immediately above the Turonian. The species thus probably lived from the Salmurian to the Senonian; it is of course possible that the specimens found in the different horizons really were different species, although these certainly belong to the same group.

At first view, one might believe that our species are identical with *Avicula pedernalis* Roemer¹ but *Avicula Aguilera* does not have the singular scars of that species, which remind us of those of *Meleagrina*. Roemer has figured them and I have observed them also on a specimen from the Vraconian of Arivechi, Sonora. The general form of Roemer's species is also a little different from ours.

Number of specimens: 6.

Age: Lower Turonian (Salmurian), upper horizon.

¹Coquand, Géol. et Pal. de Constantine, p. 216, pl. 13, figs. 17, 18.

²Thomas et Peron, Hauts-Plateaux de la Tunisie, p. 241.

³Pervinquière, Et. géol. Tunisie, pp. 101, 108.

⁴Pervinquière, Et. géol. Tunisie, p. 115, 117, 151.

INOCERAMUS Sowerby

INOCERAMUS LABIATUS Schlotheim

Pl. 20, fig. 5

- 1813: *Ostracites labiatus* Schlotheim in Leonhard's min. Taschenb. VII p. 93, fide Geinitz.
1875: *Inoceramus labiatus* Geinitz, Elbthalgebirge II, p. 46, pl. 12 cum syn.
1893: *Inoceramus labiatus* Stanton, Colorado Form., p. 77, pl. 10, fig. 4; pl. 14, fig. 2 cum syn.
1903: *Inoceramus labiatus* Petrascheck, Inoc. a. Kr. Böhmens, p. 156.

In the upper horizon of the Salmurian of Cerro del Macho, Dr. Haarmann found four *Inoceramus* of which at least three are typical *Inoceramus labiatus* Schlotheim. They are relatively small individuals but are entirely identical with those from Parras, Mexico, and different European localities. I have discussed this species extensively in Boletín 30 of the Instituto Geológico de México; a detailed description of the specimens found on Cerro del Macho does not seem to be necessary.

The occurrence of this species at our locality is of some importance because the determination of the age of the Mexican Turonian beds up to the present is entirely founded on bivalves, especially *Inoceramus*, and this class of fossils is always stratigraphically of less value than the cephalopods. My former determinations are now confirmed by the occurrence of *Inoceramus labiatus* together with typical ammonites of the lower Turonian.

Up to the present, *Inoceramus labiatus* has not yet been found in the lower beds of our Salmurian; it only occurs in the upper one, and even there it is not very frequent. Of course we can not yet draw any conclusions from this distribution. The stratigraphical conditions of the region of Mohóvano should first be studied with care, larger collections from the lower horizons should be made, so as to show if *Inoceramus labiatus* really does not appear in those beds; and finally the relation between the cephalopod beds and the shales with *Inoceramus* of other regions of northern Mexico should be ascertained.

Number of specimens: 3 (4?).

Age: Lower Turonian (Salmurian), upper horizon.

EXOYRA Say

EXOYRA HAARMANNI sp. nov.

Pl. 18, figs. 4-8

In the lowest beds of Cerro del Macho numerous specimens of an entirely smooth *Exogyra* have been found. This species resembles to a certain degree *Ex. columba* Lamarck but is not completely identical with it. Its character is:

Larger valve thin, very convex, especially in the central portion, thus forming a kind of rounded crest which goes from the beak to the lower margin. This crest gives the shell an asymmetric aspect because it is not entirely in the center, but a little nearer the anterior margin; this asymmetry is particularly noted in large specimens and much less in juvenile individuals. The beak is quite slender, spirally coiled at its point. The valve is broad, without forming wings and without a furrow in any of the sides. From the above mentioned crest the shell slopes rapidly toward the anterior margin and slowly toward the posterior margin. The surface is entirely smooth, showing only fine striae of growth and in a large specimen there is an indication of two slight concentric ridges, certainly produced by the manner of growth.

The smaller valve is nearly flat. In none of the specimens is the surface entirely preserved and a detailed description of this valve can not be made.

Our species resembles *Ex. columba* Lamarck but is much smaller and never shows a radial ornamentation, and the rounded crest is nearer to the anterior margin while in the European species it is nearer the posterior one.

In its shape our species resembles still more *Ex. columbella* Meek,¹ especially on account of the position of its crest, but the radial ornamentation of that species is entirely missing in ours.

Number of specimens: 12 and numerous fragments.

Age: Upper Cenomanian?, lower beds of Cerro del Macho.

EXOYRA cfr. OLISIPONENSIS Sharpe

1850: *Exogyra olisiponensis* Sharpe, Secondary Distr. of Portugal, p. 185, pl. 19, figs. 1, 2.

1911: *Exogyra olisiponensis* Woods, Cret. dep. Northern Nigeria, p. 278, pl. 20, figs. 1-3, cum syn.

In the upper part of the marls which we believe belong to the Cenomanian, Dr. Haarmann found a fragment of a large *Exogyra*. This

¹Stanton, Colorado Form., p. 63, pl. 8, figs. 2-4.

species is distinguished by a moderately broad and coiled beak which slopes quickly toward the anterior margin; this steep slope continues on the valve itself, while toward the posterior margin the valve slopes in a very regular curve. The ornamentation consists of radial, thick, and not very numerous ribs. Where these are crossed by specially strong concentric lamellae of growth, a kind of prolongations is formed which nearly resemble spines. The shell itself is very thick.

The specimen does not seem to be very nearly related to the *Exogyra* with radial ribs, of the Mexican middle and upper Cretaceous (*Ex. Whitneyi*, *Ex. costata*, etc.) but much more to the varieties with few ribs belonging to the group of *Ex. Olisiponensis* from the border of the Mediterranean.

Our fragment is especially similar to an individual from the Cenomanian of Wadi Am Rimpf figured by Fourtau¹; this specimen shows a very similar ornamentation.

Very similar also are some specimens from Deba Habe in Nigeria figured by Woods; these show particularly the strongly sloping region of the shell toward the anterior margin.

Exogyra olisiponensis occurs in the Cenomanian as well as in the Turonian. Many authors certainly take the species in a very wide sense and we do not yet know if it is possible to separate specifically the form of the Cenomanian from that of the Turonian. In general it seems to me that our specimen resembles more the varieties figured from the Cenomanian (compare among others Lartet, Géol. de la Palésthine, pl. 11, fig. 1) than those from the Turonian. With this view the layer in which our specimen has been found coincides well; it occurs in the upper part of the marly limestones directly below the Salmurian of Cerro del Macho. We have considered these marls with some doubt as upper Cenomanian. Unfortunately we do not know the exact stratigraphic position of the individuals from Nigeria. Woods presumes that they come from the Turonian, because in Gongila, in Nigeria, not only typical cephalopods of the Turonian (*Vascoceras*, *Pseudaspidoceras*, bi-carinated *Hoplitoides*) have been found, but also specimens of *Exogyra olisiponensis*, while in Deba Habe only *Ex. olisiponensis* has been collected. A solution of this problem is impossible for the moment, but there is the possibility that at Deba Habe the Cenomanian might be represented by beds with *Ex. olisiponensis*.

Number of specimens: 1.

Age: Upper Cenomanian (?), upper part of the lowest horizon of Cerro del Macho.

¹Fourtau, Faune crét. d'Égypte, p. 287, fig. 5.

GASTROPODA

TYLOSTOMA Sharpe

TYLOSTOMA aff. OVATUM Sharpe

Pl. 20, fig. 3

1849: *Tylostoma ovatum* Sharpe, On *Tylostoma*, p. 379, pl. 9, figs. 7, 8.

In the upper horizon of our Salmurian numerous specimens of gastropods have been collected; among these *Tylostoma* or similar genera seem to predominate. The greater part consists of internal molds which are badly preserved. In some specimens we note that the labrum is thickened on the inner side, thus these individuals very probably belong to *Tylostoma*. It is very possible that there are different species in the collection but in consideration of the bad state of preservation of the material it is impossible to distinguish them with certainty. The best specimen, as well as some less well preserved, has much similarity with *T. ovatum* Sharpe. Its features are:

Shell subglobose, of broad oval form, with a low spire of approximately six coils, mouth suboval and relatively narrow, labrum having an inner thickening. The surface is entirely smooth.

According to Choffat¹ *T. ovatum* occurs in all the beds from the Bellasian to the upper Turonian; it is thus not very surprising that a similar form is found in our beds.

Number of specimens: 5 and probably numerous fragments.

Age: Lower Turonian (Salmurian), upper horizon.

ECHINODERMATA

HEMIASTER Desor

HEMIASTER sp.

Pl. 20, figs. 6-10

In the lower marls of Cerro del Macho, Dr. Haarmann found a *Hemiaster* which probably represents a new species. He found another specimen loose below the hill which seems to belong to the same species, but it is too badly preserved for determination. The features of the better preserved specimen are:

¹Choffat, Syst. crét. Portugal II, p. 190.

Shell of small size, little convex, slightly polygonal in the ambitus, thinning toward the back and toward the front, the greatest width lying far in front of the center of the shell; slightly notched in front and truncated behind. The upper face is convex, the ambitus rounded, the lower face is nearly flat, a little thickened on the plastron and slightly depressed in the region of the peristome.

The impair ambulacrum lies in a relatively broad and moderately deep furrow which begins in the apex and from there regularly widens toward the ambitus, narrowing from there to the peristome; the furrow notches the contour considerably. The zones of pores are composed of pairs of small, nearly circular pores; the pores of each pair seem to be separated by a slight thickening. The zone is wide and nearly smooth; large tubercles are missing there and only the fine granulation can be observed. The zones of pores occupy less than half the furrow between the apex and the ambitus. Farther on follow pores separated by larger intervals and of an oblique position, arranged in such a manner that the pairs of the two zones alternate. The exact number of these pairs could not be counted.

The anterior paired ambulacra are moderately large and of lanceolate form; they lie in relatively deep furrows. The poriferous zones are broad and consist of pairs of elongated, nearly equal pores; the pores of each pair are connected by a very distinct furrow. The interporiferous zone is relatively narrow and has not quite the width of one of the poriferous zones.

The posterior paired ambulacra are a little shorter than the anterior ones. The poriferous zones consist of pairs of elongated nearly equal pores; the pores of each pair are connected by a slight furrow. The interporiferous zone is rather narrow and less wide than each of the poriferous zones. Outside of the petals the pores are smaller and the pairs are separated by large intervals.

On the whole surface of the interambulacra we observe apparently crenulated and perforated tubercles, enclosed by narrow areas and very near together; they are stronger on the anterior portion than on the posterior one. In the interporiferous zones of the ambulacra of the upper face, tubercles do not seem to exist. In the ambulacra of the lower face tubercles are infrequent, while the interambulacra of the same face are densely covered by tubercles, especially the plastron; they decrease in size from the center toward the outside. The whole surface of the shell is covered by a fine granulation.

The impair posterior inter-ambulacrum of the upper face has a sharp crest which is higher than the apical apparatus. The anterior interambulacra of the same face show much less pronounced crests; the lateral interambulacra are flattened on the center, the flattened zone being lim-

ited by well marked borders. These margins and the crests of the other interambulacra give the polygonal contour to the figure of the shell.

The peristome is small and of subpentagonal form. Its position is ex-centric toward the front; the lip is rather prominent. The periproct is not well preserved, but is of oval form and lies in the upper part of the posterior face. The apical apparatus is relatively large, the two posterior pairs are widely separated from the anterior ones; the details can not be clearly recognized.

Dimensions:

Antero-posterior diameter	27.8 mm.
Transversal diameter	27.3 mm.
Height	19.0 mm.
Distance between the apex and the posterior margin.....	16.2 mm.

Our species is not very characteristic but I have not found one with which it might be identified entirely. It resembles to a certain degree *Hemiaster Meslei* Peron et Gauthier,¹ but in this species the posterior ambulacra diverge much more than in ours, the contour is more notched in front, the furrow of the impair ambulacra is much narrower and deeper.

As we have not more than one well preserved individual I only describe and figure it without giving a new name to this species, or identifying it with a known one.

Number of specimens: 2.

Age: Upper Cenomanian (?), lower horizon of Cerro del Macho.

¹Cotteau, Peron et Gauthier, Ech, foss. de l'Algérie IV, p. 102, pl. 2, lgs. 5-8.

PLATE 12

Metoecoceras, Fagesia, Mammites

Metoecoceras, Fagesia, Mammites.**Plate 12**

Figures 1, 2, 3. **Metoecoceras** sp. nov.....page 205

Upper Cenomanian (?) or lower Turonian. Cerro del Macho, Hacienda de Mohóv-
vano, Coahuila.

Fig. 1. View from the opposite side from Figure 3.

Fig. 2. Venter of the same specimen.

Fig. 3. Side view.

Figures 4, 7. **Metoecoceras** aff. **Whitei** Hyatt.....page 203

Upper Cenomanian (?), lower horizon of Cerro del Macho, Hacienda de Mohó-
vano, Coahuila.

Fig. 4. Venter.

Fig. 7. Side view.

Figure 5. **Fagesia Pervinquieri** sp. nov.....page 212

Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóv-
vanto, Coahuila.

Venter and cross-section.

Figures 6, 8. **Mammites mohovanensis** sp. nov.....page 206

Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóv-
vanto, Coahuila.

Fig. 6. Cross-section and venter.

Fig. 8. Side view.

TO THE
LIBRARY OF THE
CONGRESS

1223

PLATE 13

Pseudaspidoceras

PLATE 13

Pseudaspidoceras**Plate 13**

Figure 1. **Pseudaspidoceras** aff. **Pedroanum** White sp.....page 209
Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Side view.

PLATE 14

Fagesia, Vascoceras

PLATE 14

Fagesia, Vascoceras**Plate 14**

- Figure 1. **Fagesia Haarmanni** sp. nov.page 211
Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
View of the other side of the specimen figured as Fig. 2, Pl. 15.
- Figure 2. **Fagesia Haarmanni** sp. nov.page 211
Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Cross-section and venter of specimen figured as Pls. 15 and 14.
- Figure 3. **Fagesia Pervinquiri** sp. nov.page 212
Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Side view of specimen figured on Pl. 12, fig. 5.
- Figure 4. **Vascoceras** n. sp. ex. aff. **V. adonense** Choffat.page 214
Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Cross-section and venter.

PLATE 15

Pseudaspidoceras, Fagesia, Vascoceras

PLATE 15

Pseudaspidoceras, Fagesia, Vascoceras**Plate 15**

- Figure 1. **Pseudaspidoceras** aff. **Pedroanum** White sp.....page 209
Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Cross-section and venter.
- Figure 2. **Fagesia Haarmanni** sp. nov.....page 211
Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Side view.
- Figures 3, 4, 5. **Vascoceras** aff. **Gamai** Choffat.....page 216
Lower Turonian, probably from upper horizon. Cerro del Macho, Hacienda de
Mohóvano, Coahuila.
Figs. 3 and 5, view of both sides.
Fig. 4. Cross-section and venter of the same specimen.

PLATE 16

Vascoceras

PLATE 16

Vascoceras**Plate 16**

- Figures 1, 2, 3, 4. **Vascoceras Angermanni** sp. nov. page 217
Lower Turonian, upper horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Fig. 1. Side view.
Fig. 2. Cross-section and venter of specimen figured on Fig. 4.
Fig. 3. Cross-section and venter.
Fig. 4. Side view of a larger specimen.

PLATE 17

Vascoceras

PLATE 17

Vascoceras**Plate 17**

- Figure 1. **Vascoceras Angermanni** sp. nov. page 217
Lower Turonian, upper horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Side view of a large specimen.
- Figure 2. **Vascoceras** n. sp. ex. aff. **V. adonense** Choffat. page 214
Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Side view of the specimen figured on Pl. 14, fig. 4.

PLATE 18

Vascoceras, Neoptychites, Exogyra

PLATE 18

Vascoceras, Neoptychites, Exogyra**Plate 18**

Figures 1, 2. **Vascoceras Mohovanense** sp. nov.page 219

Lower Turonian, lower horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.

Fig. 1. Side view.

Fig. 2. Cross-section and venter.

Figures 3, 13, 10. **Neoptychites** aff. **Cephalotus** Courtiller.page 221

Lower Turonian, upper horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.

Fig. 3. Side view.

Fig. 13. Cross-section and venter of the external whorl of the same specimen.

Fig. 10. Cross-section and venter of the inner whorl.

Figures 4, 5, 6, 7, 8. **Exogyra Haarmanni** sp. nov.page 230

Upper Cenomanian (?). Lowest horizon of Cerro del Macho, Hacienda de Mohóvano, Coahuila.

Fig. 4. Side view.

Fig. 5. Side view.

Fig. 6. Small valve.

Fig. 7. Large valve.

Fig. 8. Large valve.

Figures 9, 11. **Neoptychites** aff. **Xetriformis** Pervinquier.page 223

Lower Turonian, upper horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.

Fig. 9. Cross-section and venter.

Fig. 11. Side view.

Figure 12. **Vascoceras** sp.page 218

Lower Turonian, upper horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.

Side view.

PLATE 19

Hoplitoides

PLATE 19

Hoplitoides**Plate 19**

- Figures 1, 3. **Hoplitoides** aff. **mirabilis** Pervinquière.....page 225
Lower Turonian, upper horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Side view of two specimens.
- Figure 2. **Hoplitoides** aff. **mirabilis** Pervinquière.....page 225
Lower Turonian, upper horizon. Cerro del Macho, Hacienda de Mohóvano, Coahuila.
Cross-section and venter of specimen figured on Fig. 3.

PLATE 20

Avicula, Tylostoma, Exogyra, Inoceramus, Hemiaster

PLATE 20

Avicula, Tylostoma, Exogyra, Inoceramus, Hemiaster.**Plate 20**

- Figures 1, 2, 11, 12. **Avicula Aguilerae** sp. nov.page 227
 Lower Turonian, upper horizon. Cerro del Macho, Hacienda del Mohóvano, Coahuila.
 Fig. 1. Left valve of specimen.
 Fig. 2. Left valve.
 Fig. 11. Cardinal margin of same specimen.
 Fig. 12. Anterior side.
- Figure 3. **Tylostoma** aff. **ovatum** Sharpe.page 232
 Lower Turonian, upper horizon. Cerro del Macho, Hacienda del Mohóvano, Coahuila.
- Figure 4. **Exogyra** cfr. **olisiponensis** Sharpe.page 230
 Upper Cenomanian (?) Lowest horizon of Cerro del Macho, Hacienda del Mohóvano, Coahuila.
- Figure 5. **Inoceramus labiatus** Schlotheim.page 229
 Lower Turonian, upper horizon. Cerro del Macho, Hacienda del Mohóvano, Coahuila.
- Figures 6, 7, 8, 9, 10. **Hemiaster** sp.page 232
 Upper Cenomanian (?). Lowest horizon of Cerro del Macho, Hacienda del Mohóvano, Coahuila.
 Fig. 6. Front.
 Fig. 7. Posterior side.
 Fig. 8. Left side.
 Fig. 9. Lower face.
 Fig. 10. Upper face.

INDEX

Figures in italics denote description of species.

Abbreviations, tables of fossils	46, 51, 55, 57	<i>Barroisiceras</i> (?) <i>Fleuriausianum</i> d'Orbigny	197
<i>Acanthoceras</i>	201, 221	<i>Belemnites fibula</i> Forbes	67
cfr. Choffati Kossmat	197	Bend arch	14
<i>neptuni</i>	197	Bennington, Oklahoma	17, 19, 28
<i>preodosooides</i> B. L. & F.	67	Benton formation	200
sp.	44	Berry, E. W.	44, 146
subvicinale Bou'e, Lemoine and Thevenin	67, 202, 203	Bibliography, Ammonite Fauna of Lower Turonian of Mexico	180-183
vicinale Stollczka	202	Weno and Pawpaw Formations of Texas	146-147
worthense n. sp.	154, 150	Blayac, J.	65
<i>Acompeoceras</i>	201	Blue Mound, Tarrant County	21, 33, 65
Adkins, W. S.	12, 146, 147	Blum, Texas	12, 23, 25, 40, 54
Africa, northern	42	<i>Bochianites</i> Lory	75
limonite and pyrite fauna of	64, 65	Bohemia, fauna of	197
Turonian of	208, 209	Bokehito, Oklahoma	17, 19, 28
Aguilera, Jose G.	190, 192	creek	29
Albian, correlation of	10	formation	28, 29, 40
fauna of	64, 65, 66	Bom Jardin, Brazil	199
Algeria	45, 214, 223	Bom Jesus, Brazil	199
fauna of	66, 194	Boese, E.	20, 72, 146
Salmurian of	194	Boule, M.	67, 180, 202
Algiers	66, 189	Branca, W.	180
Alpine-Terlingua road	60	Brazil, Salmurian fauna of	198, 200, 210, 214, 221
<i>Amberleya graysonensis</i>	137, 160	Brazos-Colorado divide	16, 22, 24
Ammonite fauna, Lower Turonian	179	Brazos River	21, 41, 64
Ammonites, description of species	69, et. seq.	Brewster County, Texas	60
pyritic or limonitic	44, 59, 60, 64, 65	Bruder, G.	197, 207
tabulation of	61	Bryan County, Oklahoma	16, 27
<i>Ammonites acceleratum</i> Hyatt	201	thickness of formations in	16
<i>arnesensis</i> Choffat	189, 219	<i>Buchiceras</i>	228
<i>conciliatus</i> Stollczka	208	" <i>Buchiceras</i> " <i>Swalovi</i> Shumard	202, 206
<i>Gastianus</i> d'Orbigny	202, 203	Buda fauna	43
<i>gibbosum</i> Hyatt	201	formation	42, 62
<i>Harttii</i> Hyatt	198	Buff marl	37
<i>Kanabense</i> Hyatt	201	Burleson, Texas	17
<i>Kotoi</i> Yabe	188	Caddo, Oklahoma	22, 29
<i>leonensis</i> Conrad	211	Caddo limestone of Taff	28, 40
<i>Loewianus</i> White	208	Cajamarca, province of	199
<i>offacinatus</i> White	199, 221	Camacho, Zacatecas	60
<i>pedroanus</i> Kossmat	210	Vraconian of	200
<i>pedroanus</i> White	187, 198, 199	Camp Bullis, Texas	14
<i>Swalovi</i> Shumard	200	Carillo, Mexico	183, 190
<i>Whitei</i> Hyatt	201	Cedar Creek	54
<i>Anchura mudgeana</i> White	139, 169	Cedar Hills	31
<i>Ancyloceras bendirei</i> n. sp.	70, 170	Cedar Mills, Texas	12, 19
<i>lineatus</i> Gabb	71	Celendin, Peru	199
<i>matheronianum</i> d'Orbigny	71	Cenomanian	179, 190, 192, 200, 230, 231, 234
Angermann, Dr. Ernst	179, 184, 217, 218, 223	correlation of	10
Aptian, correlation of	10	fauna	42, 66, 67
Aptian-Gault	199	of Diego-Suarez	203
Aptian age	42	of Saxony	201, 205, 206
Aquitania, cephalopods of	197	of United States	206
Area	23, 26, 28, 29, 32, 33	Cephalopoda	68
<i>washitensis</i> n. sp.	54, 121, 168	<i>Cerithium</i>	29
Arivechi, Sonora	25, 228	Cerro del Macho	179, 190, 191, 192, 214, 218
Armstrong Iron Works	47	Cenomanian of	206, 230, 234
Asia, Turonian of	208	fauna of	192, 200, 203
Asterisk, explanation of	46	fossil-bearing beds of	183
Asteroides	95	marls of	232
Atoka, Oklahoma	22	Salmurian of	193, 211, 216, 227, 229, 231
Austin, Texas	19, 58, 68	Turonian of	208, 209
Australia, fauna of	198	Cerro de Muleros, N. M.	16, 17, 18, 19,
<i>Avicula aguilerae</i> n. sp.	190, 227, 252	20, 21, 22, 23, 24, 25, 35, 41, 42, 191, 192, 200	
<i>gracilis</i> Coquand	180, 190, 228	Chihuahua	18, 23, 183, 191
<i>pedernalis</i> Roemer	228	Choctaw County, Oklahoma	26, 27
<i>Baculites comanchensis</i> n. sp.	74, 152	Choffat, P.	186, 189, 198, 198, 213
Bailey County, Texas	25	Christiansen, F.	68
Bandera road	59	Chudeau, R.	196
Baptist Seminary, Fort Worth	61		

Cia. Perforadora Mexicana	180	<i>Enallaster</i>	111
<i>Cinula washitaensis</i> n. sp.	143, 168	<i>bravoensis</i> Boese	17, 41, 114, 164
<i>Cinulia</i> sp.	64	sp. aff. <i>texanus</i> (Roemer)	31, 114
Cleburne, Texas	25	<i>wenoensis</i> n. sp.	112, 158
Coahuila, Mexico	179, 183	<i>Engonoceras</i>	26, 31, 32
Cobb Brickyards, Fort Worth	47	<i>serpentinum</i> (Cragin)	84, 156
<i>Codiopsis</i> sp. aff. <i>doma</i>	44	sp.	54, 85, 156
Coke County, Texas	25	<i>Epiaster</i>	34
Colorado formation	186, 200, 205	<i>Aguilerae</i> Boese	109, 158
Comanche sea in Texas	44	<i>subobesus</i> n. sp.	110, 170
Comanche Peak, Texas	17, 24	<i>wenoensis</i> n. sp.	105, 160
Comanche series, thickness	15	Escalon, Mexico	183
<i>Comptonia wintoni</i> n. sp.	97, 162	Etheridge, R. Jr.	198
Constantine, province of	65	Europe, limonite and prite fauna of	64
Cooke County, Texas	9, 12, 20, 22, 27	<i>Ezogyr arietina</i>	18, 19, 31, 41, 42
thickness of formations	16	<i>cartledgei</i>	18
Coquand, H.	194, 228	cfr. <i>olistiponensis</i> Sharpe	192, 230
<i>Corbula</i>	29, 31, 32	<i>columba</i> Lamarck	199, 236
<i>basiniformis</i> n. sp.	130, 166	<i>columbella</i> Meek	230
<i>littoralis</i> n. sp.	29, 133, 166	<i>costata</i>	231
Correlation of formations	10	<i>Haarmanni</i> n. sp.	185, 230, 248
Cotteau, G.	146	<i>olistiponensis</i>	231
Cragin, F. W.	46, 146	sp. aff. <i>arietina</i> Roemer	123
Cretaceous, Mexican	231	<i>texana</i>	16
Crustacea, tabulation of pyrite and limonite	62	<i>Whitneyi</i>	17, 42, 231
Cuesta de Huanambra, Peru	199	Exposures of Pawpaw	47
Cuesta de Huanyanba	199		
Cummins, W. F., and Dumble, E. T.	18	Facies	16
<i>Cyphosoma volanum</i> Cragin	108	<i>Pagesia</i>	196, 200, 211
<i>Cyprina washitaensis</i> n. sp.	134, 166	<i>Boucheroni</i> Coquand	197
		<i>Haarmanni</i> n. sp.	187, 211, 240
Dagger, explanation of	45	<i>Kotoi</i>	198
Dakota flora, age of	44	<i>rudra</i> Stoliczka	187, 198, 213
Damergou	196	<i>superstes</i> Kosamat	186, 187, 197, 198, 212
Davis, John	68	<i>tevesthensis</i> Peron	186, 187, 188, 197, 212
Deba Habe, India	231	Fallot, J. E.	65
Decatur, Texas	14	Felix and Lenk	25
Del Rio, Texas	58, 59	Fink, Texas	15, 23, 24, 64
Del Rio clay	17, 18, 19, 41, 53, 56, 59	Finlay Mountains, Texas	16, 25
fauna of	68	<i>Flickia</i>	26, 42
limonite and pyrite fauna	61	<i>boesei</i> n. sp.	85, 150
Denison, Texas	17, 19, 20, 22, 24, 30, 47, 54, 55, 68	<i>bosquensis</i> n. sp.	87, 150
Denton County, Texas	9, 12, 17, 30, 35, 37, 54	Fort Stockton, Texas	16, 24
thickness of formations	15	Fort Worth, Texas	12, 13, 14, 17, 20, 21, 24, 32, 47, 51
Denton clay, fauna of	55, 60, 62	Fort Worth formation	21, 22
limonite and pyrite fauna of	43, 61, 62, 63	facies of	22
marl	41, 46, 53, 54, 62	geosyncline	13
Descriptions of species	68, et. seq.	limestone	22, 62
Desert sandstone	198	Fossil-bearing beds of Cerro del Macho	183
<i>Desmoceras brazoense</i>	23	Fossils, pyrite	9, 33, 42, 44, 53, 60
Diego-Suarez	203	Fourteau, R.	231
Diener, C.	197	France, Salmurian fauna of	200
Dieulefit, France	65	Turonian of	212, 223
<i>Douvilleiceras mamillare</i>	43	Fredericksburg division	24
Duck Creek marl	42, 53, 54, 62	facies	24
fauna of	54, 60	thickness of	15
limonite and pyrite fauna of	43, 61, 62, 63	Fritsch, A.	196, 197, 208
upper and lower facies	22		
Dumble, E. T.	146	Gabb, W. F.	25
and Cummins, W. F.	18	Gainesville, Texas, 12, 17, 19, 20, 26, 31, 32, 47, 51, 54	
Durant, Oklahoma	17, 28, 29	Gainesville Brick Company	31, 36, 47, 51
		Gastropoda	187, 191, 232
Eagleford shales	43, 199	Gastropods, tabulation of pyritic or limonitic	63
Echinodermata	232	Gatesville, Texas	14
tabulation of limonite	61	Geinitz, H. Br.	202
Echinoidea	101	Georgetown, Texas	12, 14
Echinoid fauna of Washita formation	62	limestone	19, 20, 21, 41, 69
of Weno formation	44, 52	Geosyncline, Fort Worth formation	13
Eck, O.	180, 195, 197, 227	<i>Gervilliois invaginata</i>	34
Edwards formation	24	(White)	122
facies	24	Ginger shale	36
Egypt	214, 218, 227	Glenrose formation	14, 43
ammonites from	180	<i>Globiconcha</i> sp.	140, 168
fauna of	196	Gongila, Nigeria	231
Salmurian of	193, 196	<i>Goniopygus</i> sp.	102
Turonian of	188, 190	Goodland, top of	24
Ellenburger limestone	13, 14, 16	fauna of	62
El Paso, Texas	21, 24		
Emscherian, the	190, 196, 200, 205, 225, 226, 228		

Grayson County, Texas	9, 27	clay, fauna of	64
thickness of formation in	16	facies	23, 24
Grayson formation	17, 19, 29, 41, 63	Kidney shale	37
facies of	17	<i>Kingena</i>	31, 33
fauna of	43, 66, 60, 63	Koenen, A. von	225
limonite and pyrite faunas of	61, 62, 63	Kossmat, A.	188, 198, 202, 208, 224
Graywacke	14		
<i>Gryphea</i>	29, 36	Lamellibranchia	191
<i>mucronata</i>	18, 19	Lamellibranchiata	227
<i>dilatata</i>	20	Lamna	63
<i>washitaensis</i>	21, 28, 29, 41	Lartet, L.	231
Grossouvre, A.	146, 223	Laube, G. C.	197, 207
Guillemain, C.	196	Lemoine, P.	67, 180, 202
		Leander, Texas	14
Haarmann, E.	180, 183, 184,	Lensing, in Comanchean formations	14
190, 191, 192, 206, 208, 211, 217, 218, 229, 230		<i>Leiocidaris</i>	29, 101
Hacienda del Mohovano	179, 183	Leon Springs, Texas	14
<i>Hamites</i>	23, 26, 28	Liddle, R. A.	59
<i>quadrinodosus</i> Jimbo	71	Limonite and pyrite fauna of Africa	66
<i>simplex</i>	42, 69	of Europe	64
sp.	64	of Madagascar	67
sp. aff. <i>armatus</i> Sowerby	69	of Texas	53-67
<i>tenawa</i> Adkins and Winton	69	summary of	60-67
<i>Hamulina worthensis</i> n. sp.	71, 162	Lisson, C. I.	199, 200
Harbort, E.	196	Lithological changes	16
Haslet, Texas	21, 83, 61, 56	Loew, Oscar	208
Haug, E.	64, 146	Love County, Oklahoma	27
<i>Heinzia</i>	201	<i>Lunatia</i> sp.	54, 140, 168
<i>Helicocryptus mexicanus</i>	41	McLennan County, Texas	19, 85, 54, 66
Boese	139	Madagascar	202, 203
<i>Hemiasper</i>	180, 232	pyritic fossils of	67
<i>Calvini</i> Clark	17, 42, 114, 168, 191	breccia	43, 47, 54, 69
<i>Fourneli</i>	228	facies	18, 19
<i>Meslei</i> Peron et Gauthier	232, 252	fauna of	43
sp.	115, 160	sandstone in	17
<i>riovistae</i> n. sp.	20, 35	<i>Mammites</i>	200, 221
Hill County, Texas	15	cfr. <i>crassitesta</i> Stoliczka	197, 198
thickness of formation	15	<i>conciatus</i>	197, 198
Hill, R. T.	16, 46, 146	Fritsch	208
Hillsboro, Texas	13	Stoliczka	186, 207, 208
<i>Holaster</i> sp. aff. <i>simplex</i> Shumard	104	<i>Footeanus</i> Stoliczka	197
<i>subglobosus</i>	43	Laube et Bruder, emend. Petrascheck	206
<i>Holotypus limitis</i>	33, 103	<i>Mohovanensis</i> n. sp.	186, 206, 220, 236
<i>Homarus</i>	62	<i>nodosoides</i> Schlotheim	186, 197, 198, 199, 207, 208
Hood County, Texas	17	var. <i>Afra</i> Pervinquière	199
<i>Hoplitoides</i>	189, 196, 225, 231	<i>Mantelliceras</i>	201
aff. <i>mirabilis</i> Pervinquière	225, 250	Mariscal Mountains, Texas	18, 41
<i>ingens</i> v. Koenen	199	Marshall County, Oklahoma	27
<i>mirabilis</i> Pervinquière	190, 226	thickness of formations	16
<i>Munieri</i>	190, 227	Medina County, Texas	58, 59
<i>nigeriensis</i>	227	<i>Meleagrina</i>	228
sp.	226	<i>Metococeras</i>	191, 200, 201, 202
<i>Hoploparia</i>	62	aff. <i>Whitei</i> Hyatt	185, 203, 236
Horizons, fossil	12	<i>Geslium</i> Petrascheck	185
Hugo, Oklahoma	19, 24, 29	sp. nov.	186, 192, 205, 235
Huronian slates	14	<i>Swalovi</i>	202, 203
Hyatt, A.	201, 202	<i>Whitei</i>	203, 205
		<i>Metopaster hortensae</i> Adkins and Winton	97, 162
India, fauna of	198, 200, 209, 212	Mexico	214
<i>Inoceramus</i> Sowerby	229	and West Texas	41
<i>hercynicus</i> Petrascheck	179, 191	Mornasian	228
<i>labiatus</i> Schlotheim	179, 186, 190, 192, 200, 229, 252	<i>Mortoniceras</i>	42
<i>cycloides</i>	191	n. sp.	91, 160
Instituto Geologico de Mexico	229	<i>worthense</i>	43
		Mungo region, of Africa	196, 225
Jack R. L.	198	Myra, Texas	13
Japan, Turonian fauna of	198		
Johnson County, Texas	9, 20, 35, 61, 54	Nacreous fossils, Weno	26, 44, 54
thickness of formations in	15	Nash, J. P.	41
Johnston, A. W.	14	<i>Natica</i> sp.	140, 168
Juarez, Mexico	191, 192	<i>Nautilus</i> sp.	68
		<i>texanus</i>	83, 34
Kamerun, Africa	189, 225	<i>Neptychites</i>	196, 200, 221
fauna of	196	aff. <i>cephalotus</i> Courtiller	189, 221, 248
Kanab Valley, Utah	205	aff. <i>retiformis</i> Pervinquière	179, 225, 226, 248
Kent, Texas	16, 18, 25, 35	<i>crassus</i> Solger	189
Kiamitia formation	23, 40, 53, 64	<i>telingaeformis</i> Solger	189
conglomerate	24		

<i>Telinga Stoliczka</i>	189, 198, 223	<i>guadalupe Boese</i>	41
<i>zetra</i>	224	<i>Placenticeras</i>	225
<i>zetriiformis</i>	189, 222, 224	<i>Placosmilia</i>	63
<i>Nerinea</i> sp.	54	<i>bravoensis</i>	41
<i>Nerita</i> sp.	139, 168	<i>mexicana</i>	41
<i>Neritina</i> sp.	139, 168	sp.	41
Nigeria, fauna of	196, 209, 214, 227, 231	<i>Plicatula</i> sp.	29
<i>Nodosaria texana</i>	18, 19, 28, 32, 34, 41, 146, 170	Polytechnic well, Fort Worth	13
Noetling, F.	146, 197	Porto dos Barcos	199
Noland's River	33	Portugal, fauna of	196, 209, 214, 217
North Central Texas	40	Turonian fauna of, 186, 187, 188, 193, 212, 218, 221	
North Denison sands	45	Pre-Cambrian strata	13, 14, 16
Northern Europe, fauna of	197	Primrose, Tarrant County	64
France, fauna of	197	<i>Prionotropis</i>	202
Germany, fauna of	197	<i>Woolgari</i>	197, 211
Tunis, pyritic fauna of	66	<i>Protocardia</i>	34
<i>Nucula</i>	29, 31, 32	<i>multistriata</i>	20
<i>nokonis</i> n. sp.	118, 168	sp. aff. <i>mltistriatum</i> (Shumard)	126, 168
<i>wenoensis</i> n. sp.	120, 168	sp. aff. <i>texana</i>	20
Oklahoma	40	<i>texana</i>	17
Opal, Zacatecas	60, 191	Protozoa	145
<i>Ostrea (Alectryonia) quadriplicata</i> Shumard	123	Province of Sergipe, Brazil	199
<i>carinata</i>	28	<i>Pseudaspidoceras</i>	196, 198, 200, 231
<i>carinata?</i> Lamarck	122	aff. <i>Footeanum</i> Petrascheck	186, 203
<i>columba major</i>	193, 194	aff. <i>pedroanum</i> White	187, 209, 238
<i>foisseyi</i> Lemoine	67	<i>Footeanum</i> Stoliczka	186, 198, 199, 209, 210
<i>marcoui</i>	29, 41	<i>pedroanum</i> White	210
<i>quadriplicata</i>	20, 28, 29, 31, 32, 36, 41	<i>salmuriense</i>	197, 209
sp. aff. <i>diluviana</i> Linnseus	122	sp.	227
<i>subovata</i>	29	<i>Pseudotissotia Douvillei</i>	197
Ostreidae	191	<i>Ptychoceras</i>	69
<i>Oxyrhina</i>	63	<i>Ptychodus</i>	63
<i>Pachydiscus peramplus</i>	197	<i>Puzosia Austeni</i>	197
<i>Pachymya</i>	33	Pyrite ammonites	42, 56, 66, 68, 69, 60-61, 64-67
Paleontology	44	fauna, comparison of	53
Palestine, fauna of	197	of North Texas Comanchean	63-67
Paluxy sand	15	of Pawpaw formation	44, 47-63
Parker County, Texas, thickness of formation	15	summary of	60-67
Parras, Mexico	229	Pyritic starfishes	54
Pawpaw formation	9, 41, 42, 43, 51, 53, 60, 62	Quarry limestone	29, 31, 32, 35, 36, 37
clay facies of	32	Quihi, Texas	58, 59
correlation of	10, 40	Red River	54, 55
facies of	11, 20, 26, 27	Red River valley	19, 20, 21, 27, 35
fauna of	44, 45, 60-63	Reed Plateau, Texas	59
fossils	9, 10	<i>Regularia</i>	62
ironstone of	27	<i>Remondia</i>	32
limonitic and pyritic fauna of	51-63	<i>Remondia? acuminata</i> (Cragin)	136, 160
lithology	25, 27	Rhone basin	65
location of outcrop	9, 10	Richardson, G. B.	147
marl facies	34	Rio Vista, Texas	12, 20, 26, 51
thickness	25, 26, 33	Ritter, Frederico	183, 211, 218
changes in	10, 12	Roanoke, Texas	17, 18
sandstone facies	27	Roberts, Jno. R.	41
shales	45	Roemer, F.	228
stratigraphy	13	Round Rock, Texas	19
Pecos, Texas	17	<i>Rudistid</i> facies	17, 24
<i>Pecten</i>	29, 33	Runnels County, Texas	25
<i>georgetownensis</i> (Kniker)	33, 125	Sahara desert, the	214
<i>inconspicuus</i> Cragin	123, 170	<i>Salenia</i> sp.	103
<i>subalpinus</i>	17, 29, 31	<i>Saleniidae</i>	62
<i>Pedinopsis symmetrica</i> (Cragin)	102	Salmurian, faunas of, 186, 193, 194, 195, 196, 208, 211, 212, 213, 215, 217, 218, 219, 227, 228, 231	
Pelecypoda	118	Santonian	228
Pelecypods, pyritic or limonitic, tabulation of	63	Saumur, France	189
<i>Peltastes</i> sp.	102	Saxony, Cenomanian of	186, 209
Pennsylvanian strata	13, 14, 16	fauna of	197
<i>Pentaceras americanus</i> n. sp.	99, 162	Saxony-Bohemia, Salmurian of	200
<i>Pentagonaster texensis</i> Adkins and Winton	95, 162	Sayn, G.	64, 147, 202
Peron, A.	66, 194	<i>Scaphites</i>	26, 28
Pern, Salmurian fauna of	199, 200, 214	<i>aequalis</i>	42, 67
Pervinquier, L.	42	<i>Hilli</i> Adkins and Winton	79, 162
45, 64, 66, 146, 189, 194, 196, 208, 212, 221, 222		sp.	54, 69
Petrasccheck, W.	186, 186, 197, 201, 202	sp. <i>aequalis</i>	43
Petrographical horizons	183	<i>worthensis</i>	16
<i>Phylloceras forbesianum</i> d'Orbigny	67	Schlagintweit, O.	199, 200
<i>diegoi</i> Boule, Lemoine and Thevenin	67	<i>Schloenbachia</i>	211
<i>Pinna</i>	33		

<i>elobiensis</i>	43	Turning point of outcrop	19
<i>inflata</i>	42	northern, at Orlena, Cooke County	27
<i>mantelli</i>	43	southern, Bexar County	19, 21, 24
sp.	17, 23	<i>Turritites</i>	26, 28
<i>triodosa</i>	23	<i>bosquensis</i> n. sp.	76, 154
<i>varians</i>	43	<i>brazoensis</i>	19, 31, 33, 41
<i>wenoensis</i> n. sp.	39, 150	sp.	33, 54, 69, 78, 154
<i>wintoni</i> n. sp.	90, 154	<i>worthensis</i> Adkins and Winton	73, 154
Schweinfurth, collections of	195	<i>Turritella</i>	28, 29
Sellards, E. H.	14, 147	<i>graysonensis</i> n. sp.	140, 168
Senonian	179, 190, 199, 209, 228	sp.	54
Shafter, Texas	16, 18	<i>worthensis</i> n. sp.	142, 168
Sharks, remains of	53	<i>Tylostoma ovatum</i>	232
Sheffield, Texas	16, 24	Type collection	68
Shuler, E. W.	62	Udden, J. A.	14
Sierra Blanca, Texas	16, 25	Union Station, Denison	47
Sierra Mojada, Mexico	183	United States, Salmurian lacking in	199
Sladen, P.	147	Turonian fauna of	199
Slate, Pre-Cambrian	14	Utatur group, of India	187, 212, 224
Solger, Fr.	194, 195, 198, 225	Valangian	61
Solitario, the	58, 60	<i>Vasoceras</i>	196, 213, 221, 231
Sonora, Mexico	18	<i>adonense</i>	186, 187
South America, Salmurian fauna of	198, 208	aff. <i>adonense</i>	214, 215
South Bosque, Texas	19, 58	aff. <i>arnesense</i> Choffat	214
South Bosque River	57	aff. <i>Gamai</i> Choffat	214, 216, 242
South Central Texas	40	<i>amieirensense</i> Choffat	199
Southern France, fauna of	197	<i>Angermannii</i>	219
Sahara, fauna of	196	sp. nov.	214, 215, 217, 244
Spain	214	<i>Caudini</i>	196
Spencer, W. K.	147	<i>Douvillei</i>	214
Stanton	201, 205, 206	<i>Durandi</i>	198
Starfishes, pyritic	54	ex. aff. <i>Gamai</i> Choffat	188
<i>Stenonia supernus</i> (Cragin)	62	<i>Gamai</i>	213, 215
<i>Stephanoceras</i>	212	<i>Gongileense</i> Woods	227
Stephenson, L. W.	20, 35, 40, 46	<i>Harttii</i> Hyatt	211
<i>Stoliczkaia</i>	202	<i>harttii</i> forme	214
Stoliczka, F.	207, 223, 224	<i>Kosemati</i>	179
Stratigraphy	13, 183	<i>mohovanense</i> n. sp.	189, 214, 219, 248
Sugar Loaf Mountain, Bryan County, Okla-		<i>Mundae</i>	216
homa	28, 29	n. sp. ex. aff. <i>V. adonense</i> Choffat	214, 240
Sulphur Creek	29	<i>polymorphum</i> Pervinquier	189, 199, 220, 221
Sycamore Creek	33, 38, 39, 47, 51	sp.	189, 218, 248
Syria, fauna of	197	<i>subconciatum</i>	214, 220, 221
Szjanocha, L.	43, 147	<i>Venericardia wenoensis</i> n. sp.	125, 160
Table of fossils:		Von Koenen, A.	225
of Del Rio clay	58-59	Vraconian, correlation of	19
of Denton clay	56	fauna	23, 42, 65, 66, 67
of Duck Creek formation	55	of Arivechi, Sonora	228
of Grayson marl	57-58	Waco	57, 58
of Pawpaw formation	51-52	Wadi Am Rimpf	231
of Weno formation	48-50	Walcott, C. G.	205
pyritic and limonitic	51-52	Washita division	42, 46
Taff, J. A.	29, 30	fauna of	43, 44, 52, 54, 60-62
Tarrant County, Texas	9, 12, 20, 23, 33, 35, 51, 54, 55	thickness of	15
thickness of formation	15	Weno formation	9, 20, 33, 43, 46, 51, 62
Terlingua, Texas	16, 58, 59	correlation of	10, 40
Creek	60	facies	11, 20, 26, 27
Thevenin, A.	67, 202	fauna	45-50
Thicknesses, of Comanchean	15	fossils	12
<i>Thomasites</i>	196	range of	46
Tishomingo, Okla	22	table of	48-50
<i>Tissotia</i> app.	44	lithology	26, 27
Trapiche das Pedras Velho, Brazil	199	location of outcrop	9, 10
<i>Trigonia</i>	29, 32, 179	sandstone in	20
Trinidad mine	60	stratigraphy of	13
Trinity division, thickness of	15	thickness of	25, 26
<i>Trochomilites</i>	63	changes in	10, 12
<i>Trochus laticinctus</i> n. sp.	158, 168	West Texas	41
Tucumcari, N. M.	18, 20	and Mexico	41
Tunis	46, 66, 187, 190, 196, 212, 214, 221, 223, 224	Whitney, F. L.	102, 147
fauna of	194	Winton, W. M.	9, 28, 32, 33, 146, 147
Turonian	205, 206, 208, 209, 212, 213, 215, 217, 219, 221, 223, 225, 228	and Adkins	16
correlation of	10	Wittich, E.	211
of Mexico	179, 220	Woodbine? flora	44
Turonian, Lower, Ammonite fauna of	179	Woods, H.	196, 227, 231
		Yabe, H.	198
		Zacatecas, Mexico	60

RETURN TO DESK FROM WHICH BORROWED

Renewed books are subject to immediate recall.

SEP 26 1977

LD 21-50m-4,'63
(D6471s10)476

General Library
University of California
Berkeley

711

C034616668