UC-NRLF

\$D 14 452

WHO AMONG THE WILD-FLOWERS

W. I. BEECROFT

Z
Main Lib. Agric. Dept.

LIBRARY
OF THE
UNIVERSITY OF CALIFORNIA.

Class

BIOLOGY
LIBRARY
6

DEP

DIV. C

DEPT. OF AGRICULTURE
DIV. OF AGR'L EDVCATION

WHO'S WHO
AMONG THE
WILD FLOWERS

WHO'S WHO AMONG THE WILD FLOWERS

COMPILED AND ILLUSTRATED

BY

W. I. BEECROFT

• WITH AN INTRODUCTION

BY

FRANCES DUNCAN

NEW YORK
MOFFAT, YARD AND COMPANY
1910

QK118
B4

BIOLOGY
LIBRARY
9

Main Lib.
Agric. Dept.

Copyright 1910, by
MOFFAT, YARD AND COMPANY
NEW YORK

All Rights Reserved

Published January, 1910

INTRODUCTION

There may be perverse and misguided folk, who, like myself, have an unaccountable distaste for looking at Mother Nature through the spectacles of Another. One is humbly aware that the "spectacles" offered by the various writers are admirable; that they clarify and enlarge the vision, place the object in an excellent light and show our much-inspected Mother as wearing the loveliest of complexions, and yet—one likes to use his own eyesight—his own spectacles. To such, this little book will prove a peculiar blessing. It goes no farther than to tell "Who's Who" among the wild flowers, merely giving an introduction and allowing one the privilege of making friends with them after the devices of his own heart.

Mr. Beecroft believes that a portrait if an excellent likeness and true in all its details will enable the most unskill'd, unpractis'd and unbotanical person to recognize his new acquaintance more speedily than much writing of many paragraphs. The drawings are supplemented by the briefest of notes on the personal appearance—only enough to identify, and in the blank spaces "remarks" can be made by the reader who is allowed the chance of making a flower book after the devices of his own

INTRODUCTION

heart; thus an idler will make it the record of a summer's wanderings and write in bits of verse and prose that fit the subjects; the exact and botanical soul can add descriptions, Bertillon measurements, variations of nomenclature, drawings of parts to heart's content: children will enjoy mounting the flowers they find opposite their pictures. While to the botanist it will be a most convenient field-book, for at the end of the season his various jottings will be found already classified and indexed.

In arrangement, the excellent method inaugurated by Mrs. Parsons, now customary in most non-technical flower-books, has been followed here, that of grouping the flowers according to color. Within the color-divisions the plants are arranged according to their time of appearance.

The nomenclature is that of the latest edition of Gray's "Manual." The pronunciation is indicated, since it sacrificed no space and may be a convenience to children and those of us whose instincts about the pronunciation of a botanical name are not always infallible.

The common names of plants are legion, of these, the best known and most generally used are in large type, the less important ones in small type and they are given lest some reader look for a flower under the alias by which he knows it, and missing this should think his friend omitted in this "Who's Who."

INTRODUCTION

It may be well to remind flower-loving folk that some of our plants are in sore need of a little intelligent affection. Unless speedily befriended, the exquisite Trailing Arbutus will quite vanish from our hillsides, leaving behind only a memory of its loveliness. The cutting of the blossoms would be harmless enough, but it is the ripping up of the creeping, slow-growing root-fibres for convenient picking which makes the extinction of this darling New-Englander so certain. The bunches for sale in the city streets represent the death of millions of little unborn Mayflowers. The next plant to go will be the Mountain Laurel. Other flowers whose existence is in peril are the Fringed Gentian and Hepatica, Orchids, Maiden Hair Fern, Sabatia, Ground Pine or Club Moss and the Hollies.

The nature love which manifests itself in a kind of pot-hunter's enthusiasm or sees in each rare flower only a kind of botanical scalp to be added to his belt, is a poor sort of affection, and though perhaps one cannot sing about that Millennium of Flowers when folk will be content to love the wood rose and leave it on its stalk, we may live to see it cut instead of torn from its stem and the last Fringed Gentian respected and left to perpetuate its lovely kind.

FRANCES DUNCAN.

BROOKLYN, NEW YORK, January, 1910.

CONTENTS

INTRODUCTION	v

I

WHITE AND GREENISH

	PAGE
BLOODROOT	2
DUTCHMAN'S BREECHES	4
SQUIRREL CORN	4
TOOTHWORT	6
CUT-LEAVED TOOTHWORT	6
PARTRIDGE BERRY	6
FOAM-FLOWER	10
MITREWORT	10
NAKED MITREWORT	10
EARLY SPRING EVERLASTING	12
EARLY SAXIFRAGE	12
RED BANEERRY	14
WHITE BANEERRY	14
SOLOMON'S SEAL	16
GREAT SOLOMON'S SEAL	16
RUE ANEMONE	18
WOOD ANEMONE	18
WILD SARSAPARILLA	20
DWARF GINSENG	20
ARROW ARUM	24
WATER ARUM	26
MAIANTHEMUM	26
PAINTED TRILLIUM	28
NODDING TRILLIUM	28
LARGE-FLOWERED TRILLIUM	28

	PAGE
STAR FLOWER	30
GOLDTHREAD	30
FALSE SOLOMON'S SEAL	32
STAR-FLOWERED SOLOMON'S SEAL	32
POISON SUMACH	34
POISON IVY	34
BRISTLY SARSAPARILLA	36
SHIN LEAF	38
ROUND-LEAVED PYROLA	38
ONE-FLOWERED PYROLA	38
RED-BERRIED ELDER	40
COMMON ELDER	40
SWEET PEPPERBUSH	40
REIN ORCHIS	42
THIMBLEWEED	44
FIELD CHICKWEED	44
INDIAN PIPE	46
WILD LEEK	46
BLADDER CAMPION	50
WHITE CAMPION	50
MUSK MALLOW	52
COMMON MALLOW	52
CHAMOMILE	54
BUNCHBERRY	54
DAISY FLEABANE (2)	56
YARROW	58
VIRGIN'S BOWER	60
RAGGED FRINGED ORCHIS	62
RATTLESNAKE PLANTAIN	62
GREATER GREEN ORCHIS	64
HOOKE'S ORCHIS	64
ROUND-LEAVED SUNDEW	66
LONG-LEAVED SUNDEW	66
SLENDER LADIES' TRESSES	66
THOROUGHWORT	68

CONTENTS

vii

	PAGE
WHITE SNAKEROOT	68
TURTLEHEAD	70
ARROW-HEAD	72
ROUGH BEDSTRAW	72
WILD BALSAM-APPLE	76
TALL MEADOW RUE	78
GRASS OF PARNASSUS	80
WATER PLANTAIN.	82
COMMON DODDER	84
WOOD SORREL	84
PEARLY EVERLASTING	86
WHITE MELILOT	86
WILD CARROT	88
BUTTONBUSH	90
POKEWEED	92
FLOWERING MOSS	94
MAYFLOWER	94
WHITE ADDER'S TONGUE	96
TWINLEAF	96
MAY APPLE	97
SWEET WHITE VIOLET	97
LANCE-LEAVED VIOLET	98
CANADA VIOLET	98
WATER CRESS	99
CREEPING SNOWBERRY	99
NEW JERSEY TEA	100
BLACK SNAKEROOT	100
CLAMMY AZALEA	100
SHOWY LADY'S SLIPPER	101
LIZARD'S TAIL	101
POKE MILKWEED	102
WATER HEMLOCK	102
WATER PARSNIP	102
DALIBARDA	103
WATER LILY	103

	PAGE
WHITE FRINGED ORCHIS	104
PRAIRIE WHITE FRINGED ORCHIS	104
WINTERGREEN	104
CULVER'S ROOT	105
JIMSON WEED	105

II

YELLOW AND ORANGE

YELLOW ADDER'S TONGUE	110
MARSH MARIGOLD	112
STAR GRASS	114
CINQUEFOIL	114
GOLDEN ALEXANDERS	116
WILD PARSNIP	116
CLINTONIA	118
INDIAN CUCUMBER-ROOT	120
OAKESIA	122
LARGE-FLOWERED BELLWORT	122
BELLWORT	122
YELLOW LADY'S SLIPPER	126
SMALL YELLOW LADY'S SLIPPER	126
GOLDEN RAGWORT	128
CARRION-FLOWER	130
YELLOW POND LILY	132
WILD YELLOW LILY	134
TURK'S-CAP LILY	134
FRINGED LOOSESTRIFE	136
AGRIMONY	138
CELANDINE	138
LOOSESTRIFE	140
FOUR-LEAVED LOOSESTRIFE	140
CONE-FLOWER	142
TALL CONE-FLOWER	142
COMMON MULLEIN	144

CONTENTS

ix

	PAGE
MOTH MULLEIN	144
WILD LETTUCE	146
YELLOW, or HOP CLOVER	148
LADY'S SORREL	148
BLACK MUSTARD	150
YELLOW ROCKET	150
COMMON EVENING PRIMROSE	154
SUNDROPS	154
SPOTTED TOUCH-ME-NOT	156
PALE TOUCH-ME-NOT	156
COMMON ST. JOHN'S WORT	158
BUTTER-AND-EGGS	160
YELLOW FRINGED ORCHIS	162
DOWNY FALSE FOXGLOVE	162
SMOOTH FALSE FOXGLOVE	162
ELECAMPANE	164
PARTRIDGE PEA	166
WILD SENNA	166
FALL DANDELION	166
TANSY	170
ORANGE HAWKWEED	172
CANADA HAWKWEED	172
HAWKWEED	174
HAIRY HAWKWEED	174
RATTLESNAKE-WEED	174
THIN-LEAVED SUNFLOWER	176
LARGER BUR-MARIGOLD	176
GOLDEN CORYDALIS	179
ROUND-LEAVED VIOLET	179
DOWNY YELLOW VIOLET	180
GOLDEN CLUB	180
INDIAN POKE	181
YELLOW MELILOT	181
FROSTWEED	182
BUTTERFLY WEED	183

	PAGE
GINSENG	183
HORSE BALM	184
SNEEZEWEED	184
FERN-LEAVED FALSE FOXGLOVE	185
GOLDEN ASTER	185
GOLDEN ROD	185

III

RED, PINK, CRIMSON AND MAGENTA

SPRING BEAUTY	188
SHOOTING STAR	190
ARETHUSA	190
WAKE ROBIN	192
SESSILE-FLOWERED WAKE ROBIN	192
WILD COLUMBINE	194
WILD PINK	196
MAIDEN PINK	196
DEPTFORD PINK	196
WILD CRANESBILL	198
HERB ROBERT	198
PITCHER-PLANT	202
STEMLESS LADY'S SLIPPER	204
SHOWY ORCHIS	206
WILD AZALEA	208
FRINGED MILKWORT	210
FIELD MILKWORT	210
MOUNTAIN LAUREL	212
TWISTED STALK	214
CLASPING-LEAVED TWISTED-STALK	214
WOOD BETONY	216
BOUNCING BET	216
PALE CORYDALIS	218
SHEEP LAUREL	220
PALE LAUREL	220

CONTENTS

xi

	PAGE
GRASS PINK	222
ROSE POGONIA	222
WILD ORANGE-RED LILY	224
SPREADING DOGBANE	226
TWIN-FLOWER	226
MEADOW-SWEET	228
HARDHACK	228
PURPLE FLOWERING RASPBERRY	230
PURPLE FRINGED ORCHIS	234
SMALLER PURPLE FRINGED ORCHIS	234
PRINCE'S PINE	234
SPOTTED WINTERGREEN	234
COMMON MILKWEED	236
PURPLE MILKWEED	236
HEDGE BINDWEED	238
FIELD BINDWEED	238
LARGE MARSH PINK	240
BITTER BLOOM	240
SEA PINK	240
SLENDER MARSH PINK	240
SWAMP MILKWEED	242
FOUR-LEAVED MILKWEED	242
GREAT WILLOW-HERB	244
CARDINAL FLOWER	244
MEADOW BEAUTY	246
MARYLAND MEADOW BEAUTY	246
OSWEGO TEA	246
PINK KNOTWEED	248
LADY'S THUMB	248
WATER PERSICARIA	248
JOE-PYE WEED	250
RABBIT'S-FOOT CLOVER	252
LIVE-FOR-EVER	254
GROUND PINK	258
DOWNY PHLOX	258

	PAGE
VIOLET WOOD SORREL	259
HAIRY BEARD-TONGUE	259
PAINTED CUP	260
QUEEN OF THE PRAIRIE	261
COMMON PASTURE ROSE	262
SWAMP ROSE	262
EARLY ROSE	262
SWEET-BRIER; EGLANTINE	263
WILD BERGAMOT	263
PURPLE CONE-FLOWER	264
CROSS-LEAVED MILKWORT	264
CORN COCKLE	265
SWAMP ROSE MALLOW	265
PURPLE GERARDIA	266

IV

BLUE AND PURPLE

LIVERLEAF	270
SHARP-LOBED LIVERLEAF	270
BIRD-FOOT VIOLET	272
ROBIN'S PLANTAIN	274
WILD LUPINE	274
BLUETS	276
GILL-OVER-THE-GROUND	276
LARGER BLUE FLAG	278
SLENDER BLUE FLAG	278
COW VETCH	280
AMERICAN VETCH	280
BITTERSWEET; NIGHTSHADE	282
SELF-HEAL	284
BLUE-EYED GRASS	284
BELLFLOWER	286
HAREBELL	286
PALE SPIKED LOBELIA	290

CONTENTS

xiii

	PAGE
MOTHERWORT	290
PICKEREL-WEED	292
MONKEY FLOWER	292
MAD-DOG SKULLCAP	294
SKULLCAP	294
BLUE VERVAIN	296
WILD MINT	298
SPEARMINT	298
PEPPERMINT	298
IRONWEED	300
CHICORY	302
BLAZING STAR	302
NEW ENGLAND ASTER	304
FRINGED GENTIAN	306
CLOSED GENTIAN	306
VIRGINIAN COWSLIP	310
EARLY BLUE VIOLET	310
ARROW-LEAVED VIOLET	311
DOG VIOLET	311
COMMON VIOLET	312
WILD BLUE PHLOX	312
FORGET-ME-NOT	313
SMALLER FORGET-ME-NOT	313
SPIDERWORT	313
BEACH PEA	314
COMMON SPEEDWELL	314
AMERICAN BROOKLIME	315
VIRGINIA DAY-FLOWER	315
BLUE TOADFLAX	316
VIPER'S BUGLOSS	316
TALL WILD LARKSPUR	317
DWARF LARKSPUR	317
PURPLE ORCHIS	317
SEA LAVENDER	318

V

MISCELLANEOUS

	PAGE
SKUNK CABBAGE	322
WILD GINGER	324
JACK-IN-THE-PULPIT	326
BRANCHING BUR-REED	328
SWEET FLAG	328
PINESAP	330
BEECHDROPS	330
LARGE CORAL ROOT	332
RATTLESNAKE-ROOT	332
GROUNDNUT	334
EARLY MEADOW RUE	338
BLUE COHOSH	338
COMMON ALUM ROOT	339
LARGE TWAYBLADE	340
CUT-TAIL FLAG	341
NARROW-LEAVED CAT-TAIL FLAG	341

LIST OF ILLUSTRATIONS

	PAGE
BLOODROOT	3
DUTCHMAN'S BREECHES	5
SQUIRREL CORN	5
TOOTHWORT	7
PARTRIDGE BERRY	7
FOAM-FLOWER	11
MITREWORT	11
EARLY EVERLASTING	13
EARLY SAXIFRAGE	13
RED BANEBERRY	15
SOLOMON'S SEAL	17
RUE ANEMONE	19
WOOD ANEMONE	19
WILD SARSAPARILLA	21
DWARF GINSENG	21
ARROW ARUM	25
WATER ARUM	27
MAIANTHEMUM	27
PAINTED TRILLIUM	29
STAR FLOWER	31
GOLDTHREAD	31
FALSE SOLOMON'S SEAL	33
POISON SUMACH	35
POISON IVY	35
BRISTLY SARSAPARILLA	37
SHIN LEAF	39
RED-BERRIED ELDER	41
SWEET PEPPERBUSH	41

	PAGE
REIN ORCHIS (2)	43
THIMBLEWEED	45
FIELD CHICKWEED	45
INDIAN PIPE	47
WILD LEEK	47
BLADDER CAMPION	51
WHITE CAMPION	51
MUSK MALLOW	53
COMMON MALLOW	53
CHAMOMILE	55
BUNCH BERRY	55
DAISY FLEABANE	57
YARROW	59
VIRGIN'S BOWER	61
RAGGED FRINGED ORCHIS	63
RATTLESNAKE PLANTAIN	63
GREATER GREEN ORCHIS	65
ROUND-LEAVED SUNDEW	67
LADIES' TRESSES	67
THOROUGHWORT	69
TURTLEHEAD	71
ARROW-HEAD	73
ROUGH BEDSTRAW	73
WILD BALSAM-APPLE	77
TALL MEADOW RUE	79
GRASS OF PARNASSUS	81
WATER PLANTAIN	83
DODDER	85
WOOD SORREL	85
PEARLY EVERLASTING	87
WHITE MELILOT	87
WILD CARROT	89
BUTTONBUSH	91
POKEWEED	93
YELLOW ADDER'S TONGUE	111

ILLUSTRATIONS

xvii

	PAGE
MARSH MARIGOLD	113
STAR GRASS	115
CINQUEFOIL	115
GOLDEN ALEXANDERS	117
WILD PARSNIP	117
CLINTONIA	119
INDIAN CUCUMBER-ROOT	121
OAKESIA	123
LARGE-FLOWERED BELLWORT	123
BELLWORT	123
YELLOW LADY'S SLIPPER	127
GOLDEN RAGWORT	129
CARRION-FLOWER	131
YELLOW POND LILY	133
WILD YELLOW LILY	135
FRINGED LOOSESTRIFE	137
AGRIMONY	139
CELANDINE	139
LOOSESTRIFE	141
FOUR-LEAVED LOOSESTRIFE	141
CONE-FLOWER	143
COMMON MULLEIN	145
MOTH MULLEIN	145
WILD LETTUCE	147
YELLOW CLOVER	149
LADY'S SORREL	149
BLACK MUSTARD	151
COMMON EVENING PRIMROSE	153
SUNDROPS	153
SPOTTED TOUCH-ME-NOT	157
COMMON ST. JOHN'S WORT	159
BUTTER-AND-EGGS	161
YELLOW FRINGED ORCHIS	163
DOWNY FALSE FOXGLOVE	163
ELECAMPANE	165

	PAGE
PARTRIDGE PEA	167
FALL DANDELION	167
TANSY	171
ORANGE HAWKWEED	173
HAWKWEED	175
RATTLESNAKE-WEED	175
THIN-LEAVED SUNFLOWER	177
LARGER BUR-MARIGOLD	177
SPRING BEAUTY	189
SHOOTING STAR	191
ARETHUSA	191
WAKE ROBIN	193
WILD COLUMBINE	195
WILD PINK	197
MAIDEN PINK	197
WILD CRANESBILL	199
HERB ROBERT	199
PITCHER-PLANT	203
STEMLESS LADY'S SLIPPER	205
SHOWY ORCHIS	207
WILD AZALEA	209
FRINGED MILKWORT	211
FIELD MILKWORT	211
MOUNTAIN LAUREL	213
TWISTED-STALK	215
WOOD BETONY	217
BOUNCING BET	217
PALE CORYDALIS	219
SHEEP LAUREL	221
GRASS PINK	223
ROSE POGONIA	223
WILD ORANGE-RED LILY	225
SPREADING DOGBANE	227
TWIN-FLOWER	227
MEADOW-SWEET	229

ILLUSTRATIONS

xix

PAGE

HARDHACK	229
PURPLE FLOWERING RASPBERRY	231
PURPLE FRINGED ORCHIS	235
PRINCE'S PINE	235
COMMON MILKWEED	237
HEDGE BINDWEED	239
LARGE MARSH PINK	241
SEA PINK	241
SWAMP MILKWEED	243
GREAT WILLOW-HERB	245
CARDINAL FLOWER	245
MEADOW BEAUTY	247
OSWEGO TEA	247
PINK KNOTWEED	249
WATER PERSICARIA	249
JOE-PYE WEED	251
RABBIT'S-FOOT CLOVER	253
LIVE-FOR-EVER	255
LIVERLEAF	271
BIRD-FOOT VIOLET	273
ROBIN'S PLANTAIN	275
WILD LUPINE	275
BLUETS	277
GILL-OVER-THE-GROUND	277
LARGER BLUE FLAG	279
COW VETCH	281
BITTERSWEET	283
SELF-HEAL	285
BLUE-EYED GRASS	285
BELLFLOWER	287
HAREBELL	287
PALE SPIKED LOBELIA	291
MOTHERWORT	291
PICKEREL-WEED	293
MONKEY FLOWER	293

	PAGE
MAD-DOG SKULLCAP	295
SKULLCAP	295
BLUE VERVAIN	297
WILD MINT	299
SPEARMINT	299
PEPPERMINT	299
IRONWEED	301
CHICORY	303
BLAZING STAR	303
NEW ENGLAND ASTER	305
FRINGED GENTIAN	307
CLOSED GENTIAN, ...	307
SKUNK CABBAGE	323
WILD GINGER	325
JACK-IN-THE-PULPIT	327
BRANCHING BUR-REED	329
SWEET FLAG	329
PINESAP	331
BEECHDROPS	331
CORAL ROOT	333
RATTLESNAKE-ROOT	333
GROUNDNUT	335

I

WHITE AND GREENISH

BLOODROOT: Indian Paint: Red Puccoon.

Sanguinària canadénsis.

Poppy Family

Apr.—May.

Found in open woods and borders of woodlands. Nova Scotia to Florida, west to Nebraska.

FLOWERS—Pure white, 8–12 divisions, solitary on stalk 6–12 inches high.

LEAVES— Rounded, deeply lobed, springing from thick rootstock.

Juice of root orange colored.

(*See Frontispiece.*)

NOTES.

Bloodroot.

1. DUTCHMAN'S BREECHES: Ear-drops:
White Hearts: Soldiers' Cap.

Dicentra Cucullària.

Fumitory Family.

Apr.—May.

Found in rich, rocky woods. Nova Scotia to the Carolinas, common westward.

FLOWERS—White, tipped with pale yellow, strung on slender stalk 5–10 inches high.

LEAVES— Finely cut, from a bulbous root.

2. SQUIRREL CORN: *Dicentra canadensis.*

Fumitory Family.

Apr.—May.

Found in rich woods. Nova Scotia to Virginia, west to Mississippi.

FLOWERS—White, tinged with magenta, fragrant; 4–8 on a stalk 6–12 inches high.

LEAVES— Finely cut, from the base. Roots bear small, yellow tubers.

NOTES.

1. Dutchman's Breeches.

2. Squirrel Corn.

1. TOOTHWORT; CRINKLE-ROOT:

Dentària diphýlla.

Mustard Family.

Apr.—May.

Found in rich woods. Nova Scotia, Maine to Minnesota, south to the Carolinas, west to the Mississippi.

FLOWERS—White, 4 divisions, borne in a loose cluster terminating a stalk 8–13 inches high.

LEAVES— 2 opposite, compound leaves borne on flower stalk, leaves at base larger. Rootstalk long, crinkled, possessing pungent flavor.

2. CUT-LEAVED TOOTHWORT: Pepper-root.

D. laciniàta.

Plant similar to the above.

FLOWERS—White or pinkish.

LEAVES— Stem leaves in whorls of 3, much divided, toothed. Roots tuberous, edible.

3. PARTRIDGE BERRY: Partridge Vine:

Twin-berry: Mitchella-vine: Squawberry.

Mitchélla rèpens.

Madder Family.

May—June.

Found in woods. Nova Scotia to Gulf of Mexico, westward to Minnesota and Texas.

FLOWERS—White, fragrant, in pairs at the tips of the sprays. Berries red, edible.

LEAVES— Rounded, evergreen, in pairs on a trailing plant. Stems, 6–12 inches long.

1. Toothwort.

2. Partridge Berry.

NOTES.

NOTES.

---

1. FOAM-FLOWER; FALSE MITREWORT:
Coolwort: Nancy-over-the-ground.

Tiarélla cordifolia.

Saxifrage Family.

Apr.—June.

Found in rich woods. Nova Scotia to Georgia, westward to Minnesota and Indiana.

FLOWERS—White, small, in clusters on stalk 6–12 inches high.

LEAVES— Lobed and toothed. Hairy throughout, springing from the base.

2. MITREWORT; BISHOP'S CAP:

Mitella diphylla.

Saxifrage Family.

Apr.—May.

Found in rich woods. Quebec to North Carolina, west to Minnesota and Missouri.

FLOWERS—White, tiny, on stalk 8–16 inches high.

LEAVES— Lower leaves coarsely toothed; 2 opposite leaves borne halfway up the flower stalk; hairy.

- *NAKED MITREWORT:

M. nuda.

Resembles preceding. Flowers greenish and few. No leaves on flower stalk. Found in cool, mossy woods. Labrador to New York, Michigan, Minnesota and northward.

1.

2.

1. Foam-flower: False Mitrewort.

2. Mitrewort.

1. EARLY SPRING, PLANTAIN-LEAVED,
or MOUSE-EAR EVERLASTING; PUS-
SY'S TOES: White Plantain, and Ladies To-
bacco. *Antennària plantaginifolia*.

Composite Family.

Apr.—June.

Found in dry fields and on hillsides. Com-
mon in U. S., Labrador to Gulf of Mexico,
west to Nebraska.

FLOWER-HEADS—Whitish, in small, tight clusters
terminating a white-woolly stem, 6–18
inches high.

LEAVES— Lower leaves broad near the tip, 3-
ribbed, stalks nearly as long as the
leaves; upper leaves narrow, downy.

2. EARLY SAXIFRAGE: *Saxifraga virginien̄sis*.

Saxifrage Family.

Apr.—June.

Found on dry, rocky hillsides. New Bruns-
wick to Georgia, west to Minnesota and Ten-
nessee.

FLOWERS—White, 5 divisions, small, in clusters
terminating a downy stalk 4–10 inches
high.

LEAVES— In a rosette at the base, toothed.

1. Early Everlasting.

2. Early Saxifrage.

14   WHITE AND GREENISH

RED BANEBERRY; Cohosh: Herb-Christopher.

Actaëa rubra.

Crowfoot Family.

Apr.—June.

Found in rich woods, an erect, bushy plant
1–2 feet high. Nova Scotia to Georgia and
far west. Common northward.

FLOWERS—White, small, in short, thick, terminal
cluster. Berries red, poisonous.

LEAVES—Compound, irregularly toothed and
lobed.

WHITE BANEBERRY:

A. álba.

Closely resembles the preceding except that
the poisonous berries are white with a black
eye. Common. Nova Scotia to Georgia and
far west.

NOTES.

Red Baneberry.

SOLOMON'S SEAL: Hairy, or True, or Twin
Flowered Solomon's Seal.

Polygonatum biflorum.

Lily Family.

Apr.—June.

Found in rich woods. A plant 1–3 feet high.

New England to Florida, west to Michigan,
Minnesota, Kansas and Texas.

FLOWERS—Greenish white, cylindrical, usually
borne in pairs from the leaf angles.
Berries blue-black.

LEAVES— Finely hairy beneath, set alternately on
stem.

Rootstalk jointed, marked with scars of
previous years' growths.

GREAT SOLOMON'S SEAL: *P. commutatum.*

Found in meadows and along river-banks. A
plant 2–7 feet high, resembles preceding, but
leaves smooth throughout, partly clasping and
flowers are in clusters of 2–8.

NOTES.

Solomon's Seal.

1. RUE ANEMONE: *Anemonélla thalictroides*.
Crowfoot Family. Apr.—June.

Found in thin woods, a small plant 5–9 inches high. New England to Minnesota, south to Tennessee.

FLOWERS—White, or pink tinged, 5–10 divisions, in clusters of 2–3.

LEAVES— In groups of 3, 3-lobed.
Roots tuberous.

2. WOOD ANEMONE: Wind-Flower:

Anemone quinquefòlia.

Crowfoot Family. Apr.—June.

A plant 4–8 inches high. Found in open woods. Canada, south to Georgia, west to Rocky Mountains.

FLOWERS—White, or magenta tinged, solitary, 4–9 divisions.

LEAVES— In a whorl of 3–5, each leaf variously divided and toothed.

NOTES.

---

1. Rue Anemone.

2. Wood Anemone.

1. WILD or FALSE SARSAPARILLA:

Aràlia nudicaùlis.

Ginseng Family.

May—June.

A plant 8–12 inches high. Found in woods.

New England to Dakota, south to mountains of Carolina.

FLOWERS—Greenish white, tiny, 3–7 clusters.
Berries purple-black in clusters.

LEAVES— In 3 divisions, each consisting usually of 5 finely toothed leaflets.

Roots long, slender, aromatic.

2. DWARF GINSENG: Ground Nut.

Panax trifolium.

Ginseng Family.

May—June.

A plant 4–8 inches tall. Found in woods.

Canada to the mountains of Georgia.

FLOWERS—White, tiny, in small clusters. Berries yellow.

LEAVES— In 3 divisions, each consisting of 3–5 toothed leaflets.

Root spherical, aromatic.

NOTES.

---

1. Wild Sarsaparilla.

2. Dwarf Ginseng.

NOTES.

NOTES.

ARROW ARUM: Green Arrow Arum.

Peltandra virginica.

Arum Family.

May—June.

A plant 1-3 feet high. Found growing in clumps in shallow water. Maine to Florida, west to Michigan and Missouri.

FLOWERS—Small, borne on club within a fleshy sheath; outer surface of sheath green. Berries green in close cluster.

LEAVES— Rich green, large, shaped like an arrow-head.

NOTES.

Arrow Arum.

1. WATER ARUM: Marsh Calla.

Calla palústris.

Arum Family.

May—June.

A plant 5–10 inches high. Found in cool bogs and swamps. Nova Scotia and southward to Virginia, westward to Minnesota, Iowa and Michigan.

FLOWERS—Tiny on short, thick club; outer envelope white above, greenish beneath, opening nearly flat. Berries red, in clusters.

LEAVES— Rich green, somewhat heart-shaped, from creeping rootstock.

2. MAIANTHEMUM: Canada Mayflower: Two-leaved Solomon's Seal: False Lily-of-the-Valley.

Maiáanthemum canadéense.

Found in rich woods. Labrador to North Carolina, west to Dakota and Manitoba.

FLOWERS—White, tiny, in small clusters terminating a stem, 3–6 inches high. Berries brown-spotted, turning red in autumn.

LEAVES— 2–3 seated on stem.

NOTES.

---

1. Water Arum.

2. Maianthemum.

PAINTED TRILLIUM. *Trillium undulatum.*

Lily Family.

May—June.

Found in rich woods. Quebec and Ontario to Georgia, west to Wisconsin and Missouri.

FLOWERS—Solitary, 3 wavy-edged, white divisions marked with a crimson V at base.

LEAVES— Broad, in a whorl of 3 at the summit of stem, 8–15 inches high.

NODDING TRILLIUM: Nodding Wake-Robin.

T. cernuum.

FLOWERS—White, flower stem recurved so that the blossom hangs beneath the leaves.

LARGE-FLOWERED TRILLIUM: Large-flowered Wake-Robin or White Wood-Lily.

T. grandiflorum.

Found in same habitat, Western Vermont and Quebec to North Carolina, west to Minnesota and Missouri.

FLOWERS—White, turning pink with age, divisions often 2 inches long.

NOTES.

Painted Trillium.

1. STAR FLOWER: Star Anemone: Chickweed Wintergreen. *Trientalis americana.*

Primrose Family. May—June.

A plant 3–7 inches high. Found in moist woods. From mountains of Virginia and Illinois north to Labrador and Manitoba.

FLOWERS—White, on slender stalks, 6–7 pointed divisions.

LEAVES— 5–10 in a circle at summit of stem.

2. GOLDTHREAD: Canker-root.

Cóptis trifolia.

Crowfoot Family.

May—July.

Found both in dry and wet woods. Labrador to Alaska, south to Michigan and mountains of North Carolina and Tennessee.

FLOWERS—White, 5–9 divisions, solitary, on stalk 3–6 inches high.

LEAVES— From the base, dark green, evergreen, finely toothed, divided into 3 leaflets.

Rootstock wiry, yellow, bitter.

NOTES.

1. Star Flower.

2. Goldthread.

FALSE SOLOMON'S SEAL: False Spikenard:

Solomon's Zig-zag.

Smilacina racemosa.

Lily Family.

May—June.

A plant 1–3 feet high. Found in moist woods and borders of woods. Nova Scotia to Georgia, westward Minnesota, Arizona and British Columbia.

FLOWERS—White, tiny, in clusters terminating plant stem. Berries first brown-spotted, finally red.

LEAVES— Alternate, with very short stems if any.

STAR-FLOWERED SOLOMON'S SEAL*S. stellata.*

Is a much smaller species.

FLOWERS—White, larger and fewer than those of the preceding. Berries black, or green, with black stripes.

LEAVES— Partly clasping.

NOTES.

False Solomon's Seal.

1. POISON SUMACH:

Rhús Vérnix.

Cashew Family.

June.

Shrub 6-15 feet high. Found on wet, low ground. Very poisonous. Western Maine to Ontario and southward.

FLOWERS—Small, whitish green in loose clusters from the angles of the leaves. Fruit small, dry, grayish white berries.

LEAVES—7-13 leaflets without teeth. Leaf-stalks red.

2. POISON IVY: Three-Leaved Ivy: Mercury.

Rhús Toxicodéndron.

Cashew Family.

June—July.

A plant sometimes vine-like in habit, sometimes bushy. Found along roadsides, on fences and stone walls, tree-trunks, and running over the ground. Poisonous, and very common.

FLOWERS—Tiny, greenish white, in clusters. Fruit dry, grayish white berries.

LEAVES—In groups of 3 leaflets (Common Ivy, Virginia Creeper or Woodbine has 5), toothless or irregularly notched.

1. Poison Sumach.

2. Poison Ivy.

BRISTLY SARSAPARILLA: *Aràlia hispida*.

Ginseng Family.

June—July.

Found in rocky woods. A hairy plant 2-3 feet high. Newfoundland to North Carolina, west to Indiana and Minnesota.

FLOWERS—Tiny, dull white, in clusters. Berries in clusters, bluish in effect, inedible.

LEAVES— Compound, toothed.

NOTES.

Bristly Sarsaparilla.

SHIN LEAF:

Pýrola elliptica.

Heath Family.

June—July.

Found in woods. Eastern United States and
Quebec to Maryland, west to Rocky mountains.

FLOWERS—Greenish white, 5 divisions, fragrant,
on stalk 5–10 inches high.

LEAVES— From the base, thin, exceeding their
stalks in length.

ROUND-LEAVED PYROLA:

P. americana.

FLOWERS—Similar to the preceding.

LEAVES— Nearly round, thick, shining green.

ONE-FLOWERED PYROLA:

Monèses uniflora.

Bears a single white or pink flower.

NOTES.

Shin Leaf.

1. RED-BERRIED ELDER:

Sambucus racemosa.

Honeysuckle Family.

June—July.

Found on borders of woods. A shrub 2–12 feet high. New Foundland to British Columbia, south to Georgia, Iowa and Colorado.

FLOWERS—Dull white, in large cluster, followed by red berries.

LEAVES— 5–7 finely-toothed leaflets.

COMMON ELDER: Elderberry: American, Sweet, or Black-berried Elder.

S. canadensis.

FLOWERS—White, in flat clusters; fruit purple-black.

2. SWEET PEPPERBUSH; White Alder: Alder-leaved Clethra.

Clèthra alnifolia.

Heath Family.

July—Aug.

Shrub 3–10 feet high. Found in low woodlands, thickets, chiefly near the Atlantic coast.

FLOWERS—White, in narrow clusters, fragrant.

LEAVES— Alternate, finely toothed, especially toward the tips.

NOTES.

1. Red-berried Elder.

2. Sweet Pepperbush.

1. REIN ORCHIS: *Habenària bracteàta.*

Orchis Family. June—Aug.

Found in damp woods. Nova Scotia to Alaska, south to Washington, Minnesota and Pennsylvania.

FLOWERS—Greenish, small, in spike terminating stem 6–20 inches high.

LEAVES— Long and broad. Long, narrow bracts or leaflets spring from bases of flowers.

2. REIN ORCHIS: *Habenària hyperbòrea.*

Orchis Family. June—Aug.

Found in wet woods. Newfoundland and Alaska, south to Pennsylvania and Nebraska.

FLOWERS—Greenish, in dense, narrow spike terminating stem 8–30 inches high.

LEAVES— Oblong to lance-shaped.

NOTES.

---

1. Rein Orchis.

2. Rein Orchis.

1. THIMBLEWEED: TALL or SUMMER
ANEMONE.

Anemone virginiana.

Crowfoot Family.

June—Aug.

Found along roadsides and borders of wood-lands. Maine to Minnesota and southward.

FLOWERS—White or greenish, 5 divisions, terminating stalks 2–3 feet high. Seeds form thimble-shaped burrs.

LEAVES— 3 in a whorl, 3-parted, divisions variously lobed and toothed. Later flowers bear 2 divided leaves midway of flower stalks.

2. FIELD CHICKWEED: *Cerástium arvense.*

Pink Family.

May—July.

Found in dry, rocky land. Common.

FLOWERS—White, 5 divisions, divisions 2-lobed, terminating stems 4–10 inches high.

LEAVES— Small, narrow.

NOTES.

---

1. Thimbleweed.

2. Field Chickweed.

1. INDIAN PIPE; Corpse Plant: Ice-plant:
Ghost-flower. *Monótropha uniflora*.

Heath Family. June—Aug.

Found in moist, heavily-shaded woods.
Throughout North America, except far south.

FLOWERS—Waxy white, solitary, nodding, terminating stem 4–10 inches high.

LEAVES— None. Plant stem colorless, turning black with age, bearing numerous scales, one or more stems arising from a mass of brittle, fibrous roots.

2. WILD LEEK: *Allium tricóccum*.

Lily Family. June—July.

Found in rich woods. New Brunswick to Minnesota, south to North Carolina.

FLOWERS—Greenish white, clustered at the end of stalk 4–15 inches high.

LEAVES— Two or three leaves 8–10 inches long, which wither before the flowers bloom, arise from a pointed bulb; strongly onion scented.

1. Indian Pipe.

2. Wild Leek.

NOTES.

NOTES.

---

1. BLADDER CAMPION: *Silène latifolia.*

Pink Family. June—Aug.

Found in fields, roadsides, waste places.

Canada south to New Jersey, Illinois and Iowa. Naturalized from Europe.

FLOWERS—White, 5 divisions, divisions deeply 2-lobed, flower cup much inflated; terminating branches 8–18 inches high.

LEAVES— Opposite, smooth.

2. WHITE CAMPION: Evening Lychnis.

Lychnis álba.

Pink Family. July—Sept.

Plant 1–2 feet high. Found in waste places.

Adventive from Old World.

FLOWERS—White or pink, fragrant, opening in the evening; divisions 5, deeply cleft, sticky and hairy beneath the flower. Small clusters terminate branches.

LEAVES— Opposite, leaves and stems hairy.

1. Bladder Campion.

2. White Campion.

1. MUSK MALLOW:

Málva moschàta.

Mallow Family.

June—Sept.

Found by roadsides and in waste places.

Canada, Eastern, Middle and Southern States.

Naturalized from Europe. Escaped from cultivation.

FLOWERS—White or pink, 5 divisions, in small cluster terminating stem, 1–2 feet high.

LEAVES— Deeply cut and lobed, faintly musk-scented.

2. COMMON MALLOW; Cheeses.

Málva rotundifòlia.

Mallow Family.

June—Oct.

Found in waste places. Throughout United States. Naturalized from Europe.

FLOWERS—White or pinkish, small, on long stalks from leaf angles.

LEAVES— Slightly lobed, long stemmed. Plant stem creeping, 4–10 inches long.

1. Musk Mallow.

2. Common Mallow.

1. CHAMOMILE: Mayweed: Dog-fennel: Dill-weed: Pig-sty Daisy: Dog's or Fetid Chamomile. *Anthemis Cótula.*

Composite Family.

June—Oct.

A plant 8–20 inches high. Found by roadsides and in dry waste land. Widely distributed throughout North America.

FLOWER-HEADS—Resemble small daisies, the white rays 3-toothed on the tip.

LEAVES— Finely dissected, ill-scented.

2. BUNCHBERRY: Low or Dwarf Cornel.

Córnus canadénsis.

Dogwood Family.

June—July.

Found in woods. Plant 4–8 inches high.

Labrador to Alaska, south to West Virginia and Minnesota.

FLOWERS—Dull white, tiny, in clusters surrounded by 4 white leaflets simulating petals. Berries red in close cluster, edible.

LEAVES— Smooth, oval, pointed, 4–6 in a whorl.

1
Chamomile.

2
Bunchberry.

DAISY FLEABANE: *Erigeron ramòsus.*

Composite Family. June—Oct.

A plant 1-2 feet high. Found in fields, waste lands, roadsides. Nova Scotia to Virginia, and west to Missouri.

FLOWER-HEADS—About $\frac{1}{2}$ inch across, with large, green-yellow disk, rays white, narrow.

LEAVES— Narrow, nearly or quite toothless, lower leaves broader at the tip.

DAISY FLEABANE; SWEET SCABIOUS:

E. ánnuus.

Similar to the preceding, with broader, coarsely-toothed leaves. Rays white or pale lilac.

NOTES.

Daisy Fleabane.

YARROW; MILFOIL: Old Man's Pepper; Nose-
bleed. *Achillèa Millefòlium.*

Composite Family.

June—Oct.

Found on roadsides and in waste places.

Common throughout North America.

FLOWER-HEADS—Grayish white, sometimes crim-
son, small, in flat-topped cluster ter-
minating stem, 1–2 feet high.

LEAVES— Finely dissected, aromatic.

NOTES.

Yarrow.

VIRGIN'S BOWER: Traveller's Joy: Old Man's Beard: Devil's Yarn.

Clématis virginiana.

Crowfoot Family.

July—Aug.

A vine 6–12 feet long. Found in rich, damp soil, climbing over fences and shrubs. From Canada south to Georgia and Kansas.

FLOWERS—White or greenish, 4 divisions, in loose clusters from the leaf angles. Seeds in plummy masses.

LEAVES— Opposite, consisting each of 3 coarsely-toothed leaflets.

NOTES.

Virgin's Bower.

1. RAGGED FRINGED ORCHIS: Fringed
Green Orchis. *Habenaria lácera.*

Orchis Family. July—Aug.

Found in swamps and wet woods from
Newfoundland to Minnesota, south to Mis-
souri and Alabama.

FLOWERS—Greenish or dingy white, with lower
lip cut to thread-like fineness; borne in
spike terminating stem, 10–20 inches
high.

LEAVES— Long and narrow, or oblong.

2. RATTLESNAKE PLANTAIN: Downy Rat-
tlesnake Plantain. *Epipáctis pubéscens.*

Orchis Family. July—Aug.

Found in dry woods. New England to Flor-
ida, west to the Mississippi.

FLOWERS—Small, white, numerous on downy stalk
6–18 inches high.

LEAVES— Mottled with light and dark green.

1. Ragged Fringed Orchis.

2. Rattlesnake Plantain.

GREATER GREEN ORCHIS: Large Round-leaved Orchis.

Habenària orbiculàta.

Orchis Family.

July—Aug.

Found in rich woods. Labrador to Alaska, south to South Carolina; Minnesota, and Washington.

FLOWERS—Greenish white, on stalk 1–2 feet high.

LEAVES— 2 large, broad leaves, shining green above, silvery beneath, lying flat on the ground. One or more tiny leaflets borne on the flower stalk.

HOOKEER'S ORCHIS:

H. Hookèri.

Similar to the preceding. Flowers yellowish green. No leaflets on flower stalk.

NOTES.

Greater Green Orchis.

1. ROUND-LEAVED SUNDEW: Dew-plant.

Drósera rotundifòlia.

Sundew Family.

July—Aug.

Found in bogs and sandy marshes. Labrador to the Gulf of Mexico and westward. From Alaska to California.

FLOWERS—Small, white, opening one at a time, on stalk 4–10 inches high.

LEAVES— Grow in a rosette, round, borne on long stems, and covered with red hairs.

LONG-LEAVED SUNDEW: *D. longifòlia.*

Leaves widest at the tip, tapering toward the stems. Leaf stems without hairs.

2. LADIES' TRESSES:

Spiránthes cérnua.

Orchis Family.

Aug.—Oct.

Found in low meadows and swamps. Newfoundland to Georgia, Minnesota and Nebraska.

FLOWERS—Small, white or yellowish, arranged spirally on stem 6–12 inches high.

LEAVES— Long and narrow, grass-like.

SLENDER LADIES' TRESSES:

S. grácilis.

Found in pastures and on dry hillsides. Bears flowers on one side of the stem, or spirally. Basal leaves wither before flowers bloom. Roots tuberous.

1. Round-leaved Sundew.

2. Ladies' Tresses.

THOROUGHWORT; BONESET: Ague-weed:

Indian Sage.

Eupatòrium perfoliàtum.

Composite Family.

July—Sept.

Found in low, wet ground. From Nebraska, Manitoba, New Brunswick, south to the Gulf of Mexico.

FLOWER-HEADS—Small, dull white, in large, flat-topped cluster terminating stem 2-5 feet high.

LEAVES— Opposite, often united at the base around the stem, wrinkled, toothed. Plant stem and under surface of leaves downy.

WHITE SNAKEROOT: White or Indian Sanicle: Deerwort Boneset.

E. urticaefòlium.

Found in rich moist woods. A plant 1-4 feet high.

FLOWER-HEADS—White, resembling the garden Ageratum, in spreading cluster.

LEAVES— Nearly heart-shaped, coarsely toothed, long-stemmed.

NOTES.

Thoroughwort.

TURTLEHEAD; BALMONEY: Snake-head:

Shell-flower: Cod-head.

Chelone glabra.

Figwort Family.

July—Sept.

Found in wet ground. Newfoundland to Florida and Texas, west to Mississippi and Minnesota.

FLOWERS—White or tinged with pink, terminating stem 1–3 feet high.

LEAVES— Opposite, toothed.

NOTES.

Turtlehead.

1. ARROW-HEAD: *Sagittària latifòlia.*

Water Plantain Family. July—Sept.

Found in shallow water. From Mexico northward throughout North America.

FLOWERS—White, divisions 3, in whorls of 3, on stalk 6 inches to 4 feet high.

LEAVES— Arrow-head-shaped, exceedingly variable, from short and broad to long and narrow.

2. ROUGH BEDSTRAW: *Gàlium aspréllum.*

Madder Family. June—Aug.

A plant 2-4 feet tall. Found in rich, moist ground. Eastern half of United States and Canada.

FLOWERS—White, tiny, in airy clusters. Flower stems 2-3 times forked.

LEAVES— In whorls of 6 on main stems, prickly-rough on edge and rib. Plant stems rough backwards with hooked prickles.

1. Arrow-head.

2. Rough Bedstraw.

NOTES.

NOTES.

WILD BALSAM-APPLE: *Echinocýstis lobàta*.
Gourd Family. July—Sept.

A rapid climbing vine 10–20 feet long.

Found in rich, moist soil, beside streams and in waste places. New Brunswick to Kentucky, west to Manitoba and Texas.

FLOWERS—Greenish white, small, in loose clusters from the leaf angles.

LEAVES— Deeply and sharply 5-lobed.

NOTES.

Wild Balsam-apple.

TALL MEADOW RUE:

Thalictrum polygamum

Crowfoot Family.

July—Sept.

Found in swamps and low meadows.

Newfoundland to Ohio and southward.

FLOWERS—White or greenish, in plume-like cluster terminating stem 3–10 feet high.

LEAVES—Compound, the leaflets small, lustreless, lobed.

NOTES.

Tall Meadow Rue.

GRASS OF PARNASSUS: Carolina Grass of
Parnassus. *Parnássia caroliniàna.*

Saxifrage Family.

July—Sept.

Found in swamps and wet meadows. New
Brunswick to Florida, west to Iowa.

FLOWERS—Creamy white with greenish veins, di-
visions 5, on slender stalks 8–20 inches
high.

LEAVES— From the base, somewhat heart-shaped,
a single leaf clasping the flower stalks.

NOTES.

Grass of Parnassus.

WATER PLANTAIN:

Alisma Plantago-aquatica.

Water Plantain Family.

July—Sept.

Found in shallow water. North America,
Europe, Asia.

FLOWERS—Small, white, in loose, symmetrical
cluster, on stalk 1–3 feet high.

LEAVES— From the base, variable in shape, broad
or narrow.

NOTES.

Water Plantain.

1. COMMON DODDER: Strangle-weed: Love-vine: Angel's Hair. *Cuscuta Gronovii.*

Convolvulus Family. July—Sept.

Found in moist soil. Nova Scotia and Minnesota south to Gulf of Mexico.

FLOWERS—Tiny, dull white, 5-lobed, in clusters.

LEAVES— None. Stems long, wiry, yellow or orange, twining over vegetation.

2. WOOD SORREL: White or True Wood-sorrel: Alleluia: *Oxalis Acetosella.*

Wood Sorrel Family. May—July.

A plant 3-4 inches tall. Found in cold, damp woods. Nova Scotia and Manitoba, south to North Carolina.

FLOWERS—White or pinkish, veined with deep pink lines, 5 divisions.

LEAVES— From the base, composed of 3 heart-shaped leaflets.

NOTES.

---

1. Dodder.

2. Wood Sorrel.

1. PEARLY, or LARGE-FLOWERED, EVER-LASTING: Immortelle: Silver-leaf: Cotton-weed: Moonshine: None-so-pretty.

Anáphalis margaritácea.

Composite Family.

July—Sept.

Found in dry fields, hillsides. North Carolina to Kansas and California, also far north.

FLOWER-HEADS—Center yellow surrounded by white scales, the cluster terminating a white-woolly stem 1–3 feet high.

LEAVES— Long, narrow, white-woolly.

2. WHITE MELILOT; SWEET CLOVER: Bokhara or Tree Clover: Honey Lotus.

Melilòtus álba.

Pulse Family.

June—Aug.

Found by roadsides and in waste places.

Naturalized from Europe: widespread in United States.

FLOWERS—White, small, fragrant, in slender clusters terminating branches 3–10 feet high.

LEAVES— Compound, consisting of 3 finely-toothed leaflets.

1. Pearly Everlasting.

2. White Melilot.

WILD CARROT; QUEEN ANNE'S LACE:

Bird's Nest.

Daucus Carota.

Parsley Family.

July—Sept.

A plant 2–3 feet high. Found in fields, roadsides, waste places. Naturalized from Europe. Eastern half of United States and Canada.

FLOWERS—White, tiny, in broad, rather flat cluster, curling up later, resembling a bird's nest.

LEAVES— Finely dissected.

NOTES.

Wild Carrot.

BUTTONBUSH: Globe-flower: Honey-balls:
Button-ball Shrub: River-bush.

Cephalánthus occidentális.

Madder Family.

July—Sept.

A shrub 3-12 feet high. Found in wet ground and beside ponds and streams. New Brunswick to Florida and Cuba, westward to Arizona and California.

FLOWERS—White, small, clustered in round heads.

LEAVES— Opposite, or in whorls of 3.

NOTES.

Buttonbush.

POKEWEED; PIGEON BERRY: Scape: Ink-
berry: Garget. *Phytolacca decandra.*

Pokeweed Family. July—Sept.

A plant 4–10 feet tall. Found by roadsides,
in clearings, waste places. Maine and On-
tario to Florida and Texas.

FLOWERS—Small, with 5 white divisions, pink
tinted outside, and a conspicuous green
center; borne in loose spikes. Berries
dark purple in long drooping clusters.
Ripe August—October.

LEAVES—Oblong to lance-shaped, tapering at
both ends.

Plant stems branching, reddening to-
ward fall. Root large, poisonous.

NOTES.

Pokeweed.

FLOWERING MOSS; PYXIE: Pine-barren
Beauty. *Pyxidanthèra barbulàta.*

Diapensia Family. Mch.—May.

A low plant, growing in mat-like patches,
creeping prostrate branches 6–10 inches
long. Found in dry, sandy soil, pine barrens.

New Jersey to North Carolina.

FLOWERS—White, often pink, about $\frac{1}{4}$ inch
across, 5-lobed, numerous.

LEAVES— Very small and narrow, seated on
stem, crowded toward the ends of the
branches, moss-like in appearance.

MAYFLOWER; TRAILING A R B U T U S:
Ground Laurel. *Epigàea rèpens.*

Heath Family. Mch.—May.

Found in woodlands or mossy, rocky places.

Newfoundland to Florida, west to Kentucky,

Minnesota and the Northwest.

FLOWERS—White or pink tinted, 5-lobed, fragrant,
in clusters at the ends of the woody,
creeping branches 6–10 inches long.

LEAVES— Oval, rounded at the base, old leaves
more or less rusty spotted.

NOTES.

WHITE ADDER'S TONGUE:

Erythrònium álbidum.

Lily Family.

Mch.—May.

Found in rich soil. Common west as far as Texas, Arkansas and Minnesota. Somewhat rare in Eastern States, New York to New Jersey.

FLOWERS—Pinkish-white, yellow at the center, purplish-tinged outside, 6 strongly recurved divisions, terminating stalks 5–8 inches high.

LEAVES— 2, from the base, smooth, more or less mottled, from solid bulb.

TWINLEAF: Rheumatism Root.

Jeffersònia diphýlla.

Barberry Family.

Apr.—May.

Found in rich, shady woods. New York to Wisconsin, south to Tennessee.

FLOWERS—White, 8 divisions, 1 inch broad, solitary, on stalk about 7 inches high, twice as tall in seed time.

LEAVES— From the root, long-stemmed, parted into 2 divisions.

MAY APPLE; MANDRAKE: Hog Apple:

Wild Lemon.

Podophýllum peltàtum.

Barberry Family.

Apr.—May.

A plant 12–18 inches high. Found in rich, moist woods. Western Quebec and western New England to Minnesota and southward.

FLOWERS—White, 6–9 divisions, solitary, unpleasantly scented. Fruit, large, edible berry.

LEAVES— Those of the flowerless stems terminated by a large, rounded, 7–9-lobed leaf supported by a stem in the middle; those of the flowering stem 2, much lobed, with the nodding flower in the fork.

SWEET WHITE VIOLET:

Viola pállens.

Violet Family.

Apr.—May.

Plant 3–5 inches tall. Found in moist, wooded places. Common.

FLOWERS—White with purple veins, side divisions usually bearing a small tuft of hairs, slightly fragrant.

LEAVES— Round heart-shaped, slightly scalloped.

LANCE-LEAVED VIOLET:

Viola lanceolata.

Violet Family.

Apr.—June.

Found in moist ground. Nova Scotia to Minnesota and southward.

FLOWERS—White, purple veined, side divisions beardless.

LEAVES— Long, narrow, indistinctly scalloped, tapering into a long stem.

CANADA VIOLET:

Viola canadensis.

Violet Family.

May—July.

Found in woodlands, very widely-distributed, mountainous and hilly country chiefly. Easily distinguished from other white violets by its tall leafy stem 6-24 inches high.

FLOWERS—White or pale lavender, violet tinged on the outside, side divisions bearded, lower one striped with fine dark lines.

LEAVES— Heart-shaped, pointed, toothed.

NOTES.

WATER CRESS:

Radícula Nastúrtium-aquáticum.

Mustard Family.

Apr.—Aug.

Found in brooks, ditches. Naturalized from
Europe. Common.

FLOWERS—White, 4 divisions, small, in terminal
clusters.

LEAVES—Compound, with 3–11 rounded leaflets.
Stems spreading and rooting, 4–10
inches high.

CREEPING SNOWBERRY:

Chiógenes hispídula.

Heath Family.

May—June.

A low, creeping plant. Found in mossy bogs
and woods. Canada, south to Minnesota,
Michigan and North Carolina.

FLOWERS—White, very small, bell-shaped, 4-lobed,
solitary, from the leaf angles. Berries
white.

LEAVES—Evergreen, small, with an abrupt point.
Leaves and berries aromatic.

NOTES.

---

100 WHITE AND GREENISH

NEW JERSEY TEA; RED-ROOT:

Ceanòthus americanus.

Buckthorn Family.

May—July.

A shrub 1–4 feet high. Found in dry, open woods. Gulf of Mexico to Ontario.

FLOWERS—White, small, in dense, oblong, terminal clusters.

LEAVES— Long egg-shaped, finely toothed, conspicuously 3-ribbed. Root reddish.

BLACK SNAKEROOT; BLACK COHOSH:

Tall Bugbane.

Cimicífuga racemòsa.

Crowfoot Family.

June—July.

Found in rich woods, hillsides. Southern New England and Wisconsin south to Georgia.

FLOWERS—White, ill-scented, fuzzy, borne on a long, slender spike terminating stem 3–8 feet high.

LEAVES— Compound, sharply toothed.

CLAMMY AZALEA; WHITE SWAMP HONEYSUCKLE:

Rhododéndron viscòsum.

Heath Family.

June—July.

A shrub 3–7 feet high, resembling the earlier-blooming Pink Azalea. Found in swamps, generally near the coast. Maine, south and west to Ohio and Arkansas.

FLOWERS—White or pink-tinged, 5-lobed; tube longer than the lobes, sticky-hairy outside.

LEAVES— Blunt lance-shaped, margins and midrib bristly.

SHOWY LADY'S SLIPPER:

Cypripedium hirsutum.

Orchis Family.

June—July.

Found in wet, mossy woods, swamps.

Newfoundland to Georgia, west to Mississippi.

FLOWERS—Pouch white, more or less stained with crimson-magenta, similar in shape to the Stemless Lady's Slipper, but differs in the pouch being open but not divided; other divisions of the flower white, wide, not longer than the pouch.

LEAVES—Broad, pointed, many ribbed. Plant stem leafy 1–2 feet high.

LIZARD'S TAIL:

Saururus cernuus.

Pepper Family.

June—Aug.

A plant 2–5 feet high. Found in swamps, shallow water. Rhode Island to Gulf of Mexico, west to Minnesota and Texas.

FLOWERS—White, fragrant, very small, crowded on slender spikes, nodding at the tip.

LEAVES—Heart-shaped, converging ribs, dark green, on stout stems.
Plant-stem jointed, sparingly branched, leafy.

NOTES.

POKE or TALL MILKWEED:

Asclèpias phytolaccoïdes.

Milkweed Family.

June—Aug.

A plant 3–6 feet high. Found in woodland borders, thickets from Maine to Georgia and far west.

FLOWERS—Similar in structure to the Common Milkweed, white and greenish, or magenta tinged, drooping, in a loose cluster.

LEAVES— Rather broad, pointed at each end.

WATER HEMLOCK; SPOTTED COW-BANE:

Musquash Root: Beaver Poison.

Cicuta maculata.

Parsley Family.

June—Aug.

Found in wet meadows, swampy places. A deadly poison contained in its fleshy roots.

New Brunswick to Virginia and westward.

FLOWERS—White, tiny, in thin, flat clusters terminating branches, 3–6 feet high.

LEAVES— Compound, leaflets toothed. Plant stem marked with purplish lines.

WATER PARSNIP: Hemlock Water Parsnip.

Sium cicutaefolium.

Is a similar species growing in wet places and also poisonous. It may be found from the Atlantic to the Pacific.

FLOWERS—White, in broad clusters.

LEAVES— Compound 7–15 sharply toothed leaflets, but finely dissected when growing under water.

DALIBARDA: Creeping Dalibarda.

Dalibárda rèpens.

Rose Family.

June—Sept.

Found in woodlands. Nova Scotia to Pennsylvania, west to the Mississippi.

FLOWERS—White, solitary, 5 divisions, on stalks 2–5 inches high.

LEAVES— From the base, heart-shaped, scalloped-toothed; leaves and stems finely hairy.

WATER LILY: Pond Lily: Water Nymph: Sweet-scented Water Lily.

Castàlia odoràta.

Water Lily Family.

June—Sept.

Found in shallow water. Common. Nova Scotia to the Gulf of Mexico, west to Mississippi.

FLOWERS—Pure white, sometimes pink tinged, center yellow, divisions many. Flower 3–5 inches across, floating, very fragrant.

LEAVES— Floating, mostly oblong in shape, cleft at the base, on long stems.

NOTES.

WHITE FRINGED ORCHIS:

Habenària blephariglóttis.

Orchis Family.

July—Aug.

Found in bogs, swamps. Eastern Canada to Michigan, south to Mississippi.

FLOWERS—Similar in habit to the Yellow Fringed Orchis, white, fragrant, lip fringed, in spike terminating stem 1–2 feet high.

LEAVES— Lance-shaped, parallel veined, clasping stem.

PRAIRIE WHITE FRINGED ORCHIS:

H. leucophaëa.

Is similar to the preceding, differing chiefly in having larger white or greenish flowers, lip divided into wedge-shaped segments, deeply fringed. It is more common west, being found in moist, open ground from Western New York to Minnesota and Arkansas.

WINTERGREEN; CHECKERBERRY: Partridge-berry: Deer, Box, or Spice-berry: Mountain or Ground Tea. *Gaulthèria procúmbens.*

Heath Family.

July—Aug.

Found in woodlands. Newfoundland to Northern Georgia, west to Minnesota, Michigan and Manitoba and far north.

FLOWERS—White, small, somewhat bell-shaped, nodding, from the leaf angles. Berries red, edible.

LEAVES— Dark green, evergreen, slightly toothed, aromatic, clustered at the top of stem, 2–5 inches high.

CULVER'S ROOT: Culver's Physic:

Verónica virginica.

Figwort Family.

July—Aug.

An erect plant with straight, usually unbranched stem 2–7 feet high. Found in moist woods, meadows. Western Massachusetts and Connecticut to Manitoba and southward.

FLOWERS—White or pale lavender, in dense, terminal spikes 3–6 inches long.

LEAVES— 4–7, in whorls, lance-shaped, pointed, finely toothed.

JAMESTOWN or JIMSON WEED: Thorn-apple: Stramonium: Devil's Trumpet.

Datùra Stramòonium.

Nightshade Family.

July—Sept.

A plant 1–5 feet tall. Found in fields, waste land. Nova Scotia to Gulf, west beyond the Mississippi Naturalized from Asia.

FLOWERS—White, about 4 inches long, trumpet-shaped, with 5 sharply-pointed lobes. Fruit large, egg-shaped, covered with spines.

LEAVES— Coarsely toothed and angled, rank-smelling.

ASTERS—See page 304.

NOTES.

NOTES.

NOTES.

NOTES.

II

YELLOW AND ORANGE

YELLOW AND ORANGE

YELLOW ADDER'S-TONGUE; DOG'S-
TOOTH VIOLET: Trout Lily.

Erythronium americanum.

Lily Family.

Apr.—May.

Found in moist woods, brooksides. New
Brunswick to Florida, west to Mississippi,
Minnesota and Arkansas.

FLOWERS—Lily-like, 6 divisions, yellow, more or
less tinged with brown-purple on the
outside, borne on stalk 6–12 inches
high.

LEAVES— 2 from the base, green, or mottled
with brown-purple.

NOTES.

Yellow Adder's Tongue.

MARSH MARIGOLD: Meadow-gowan: American Cowslip. *Caltha palústris*.

Crowfoot Family. Apr.—May.

A plant 8–24 inches high. Found in low, wet meadows. Newfoundland to South Carolina, Tennessee, and Nebraska.

FLOWERS—Yellow, 5–9 divisions.

LEAVES— Mostly from the root, round, kidney or heart-shaped.

NOTES.

Marsh Marigold.

1. STAR GRASS: *Hypóxis hirsùta*.
Amaryllis Family. Apr.—July.

Found in dry, open woods, meadows. Maine to Gulf of Mexico and west to Minnesota, Texas, Kansas.

FLOWERS—Yellow inside, greenish and hairy outside, 6 divisions.

LEAVES— Grass-like, 2–6 inches high.

2. CINQUEFOIL: Common Cinquefoil: Five-Finger. *Potentílla canadénsis*.

Rose Family. Apr.—Aug.

Found in dry fields, roadsides, waste places.

Quebec to Georgia, west to Minnesota, Iowa.

FLOWERS—Yellow, 5 divisions, growing singly on slender stalks from leaf angles.

LEAVES— Composed of 5 toothed leaflets. Plant stems wiry, prostrate, 6–20 inches long.

NOTES.

1. Star Grass.

2. Cinquefoil.

1. GOLDEN ALEXANDERS: Early Meadow Parsnip. *Zizia aurea.*

Parsley Family. May—June.

Found in meadows and river-banks. From New Brunswick and Dakota to the Gulf States.

FLOWERS—Yellow, tiny, in thin, flat clusters terminating branches, 8–30 inches high.

LEAVES—Compound, generally in 3 divisions, each composed of 3–7 toothed leaflets.

2. WILD PARSNIP; FIELD PARSNIP; Madnep: Tank. *Pastinàca sativa.*

Parsley Family. June—Sept.

A plant 2–5 feet tall, with noticeably grooved stem, found in roadsides, waste places. Common throughout United States and Canada. Naturalized from Europe.

FLOWERS—Dull yellow, tiny, forming flat clusters terminating branches, 2–5 feet high.

LEAVES—Compound, composed of many toothed leaflets.

NOTES.

---

1. Golden Alexanders.

2. Wild Parsnip.

CLINTONIA; YELLOW CLINTONIA:

Clintonia boreàlis.

Lily Family.

May—June.

Found in moist woods. Labrador to North
Carolina, west to Manitoba and Minnesota.

FLOWERS—Greenish yellow, 6 divisions, lily-like,
3–6 terminating stalk 6–15 inches
high. Berries pure blue.

LEAVES— Usually 3, from the base, oval or ob-
long.

NOTES.

Clintonia.

INDIAN CUCUMBER-ROOT:

Medèola virginiana.

Lily Family.

May—June.

A plant 1–3 feet high, found in rich woods.

New Brunswick to Minnesota, south to Gulf States.

FLOWERS—Greenish yellow, few at summit of stem. Berries purple-black.

LEAVES— In 2 whorls; lower whorl of 5–9 leaves, upper whorl of 3–5 smaller leaves.

Rootstock white, with slight cucumber flavor.

NOTES.

Indian Cucumber-root.

1. OAKESIA:

Oakèsia sessilifolia.

Lily Family.

May—June.

A plant 6–12 inches tall, found in rich woods.

New England to Florida, west to Minnesota, Nebraska and Arkansas.

FLOWERS—Pale yellow, 6 divisions, long and slender. Seed capsule 3-sided, resembling a beechnut.

LEAVES— Partly clasping, whitish beneath.

2. LARGE-FLOWERED BELLWORT:

Uvulària grandiflora.

Lily Family.

Apr.—June.

Found in rich woods. Western New England to Minnesota and Kansas, south to Georgia.

FLOWERS—Larger and more irregular than those of the preceding, pale or greenish yellow, inner surface smooth.

LEAVES— Leaves appear as if perforated by the plant stem.

3. BELLWORT:

U. perfoliata.

Resembles the preceding but may be distinguished by the rough inner surface of the flower. Found from Massachusetts to Ontario, Dakota, and southwest.

NOTES.

1. Oakesia.   2. Large-flowered Bellwort.   3. Bellwort.

NOTES.

NOTES.

YELLOW LADY'S SLIPPER: Large Yellow Lady's Slipper: Yellow Moccasin Flower: Whippoorwill's Shoe.

Cypripedium parviflorum Var. *pubescens*.

Orchis Family.

May—July.

Found in moist or boggy woods. Nova Scotia to Alabama, west to Minnesota, E. Kansas and Nebraska.

FLOWERS—Solitary, borne at the top of leafy stem 1–2 feet high; pouch golden yellow; other divisions of flower lined and marked with madder-purple.

LEAVES— Parallel veined, clasping stem.

SMALL YELLOW LADY'S SLIPPER:

C. parviflorum.

Similar but smaller than the preceding, with similar range. Intergrading forms frequent.

NOTES.

Yellow Lady's Slipper.

GOLDEN RAGWORT; Groundsel: Squaw-weed.

Senècio aúreus.

Composite Family.

May—July.

Found in wet meadows. Newfoundland and
Ontario to Gulf of Mexico, west to Missouri.

FLOWER-HEADS—Golden yellow, in loose clusters
terminating stem, 12–30 inches high.

LEAVES— Those from the root round or heart-
shaped, long-stemmed, toothed; stem
leaves variable, deeply lobed, upper-
most clasping.

NOTES.

Golden Ragwort.

CARRION-FLOWER: *Smilax herbàcea.*

Lily Family. May—July.

A handsome vine growing 4-15 feet long, found in moist soil. Canada to Gulf States, west to Nebraska.

FLOWERS—Greenish yellow, in inconspicuous clusters, ill-scented. Berries at first green, finally blue-black.

LEAVES— Somewhat heart-shaped, parallel veined.

NOTES.

Carrion-flower.

YELLOW POND LILY; COW LILY: Large

Yellow Pond or Water Lily: Spatter-dock.

Nymphaea advena.

Water Lily Family.

May—Sept.

Found in shallow, stagnant water. Nova Scotia to Gulf States, west to Rocky Mountains.

FLOWERS—Yellow, more or less tinged with green and purple, 6 unequal divisions; flowers borne erect on thick stems.

LEAVES— Large, deeply cleft at the base, floating.

NOTES.

Yellow Pond Lily.

WILD YELLOW LILY: Meadow or Field Lily:

Canada Lily.

Lilium canadense.

Lily Family.

June—July.

Found in wet meadows and swamps. Nova Scotia to Georgia, west beyond Mississippi, Minnesota and Missouri.

FLOWERS—Yellow to orange, with purple-brown spots inside, terminating stems 2–5 feet high.

LEAVES— Usually in whorls of 4–10.

TURK'S-CAP LILY: Turban Lily.

L. superbum.

Flowers orange-yellow, half drooping, divisions more reflexed than those of the preceding, thickly spotted. Upper leaves arranged alternately, lower leaves in whorls.

NOTES.

Wild Yellow Lily.

FRINGED LOOSESTRIFE:

Steironèma ciliatum.

Primrose Family.

June—Aug.

Found in low ground, thickets. Maine to
British Columbia, south to Gulf States.

FLOWERS—Yellow, divisions 5, a terra-cotta colored ring in the center.

LEAVES— Opposite, a fringe of hairs borne on leaf stems. 18–24 inches high.

NOTES.

Fringed Loosestrife.

1. AGRIMONY: *Agrimonia gryposépala.*

Rose Family.

June—Aug.

Found in roadsides, borders of woods and fields. Canada to North Carolina, west to California.

FLOWERS—Yellow, small, divisions 5, borne on slender spikes terminating branches, 2–4 feet high.

LEAVES— Composed of usually 7 toothed leaflets, with 1 or more pairs of tiny leaflets between; fragrant when crushed.

2. CELANDINE: Greater Celandine: Swallowwort. *Chelidonium majus.*

Poppy Family.

May—Aug.

A plant 1–2 feet high, found in roadsides, waste places. Eastern United States. Naturalized from Europe.

FLOWERS—Yellow, on slender stalks, divisions 4. Seed pods erect, 1–2 inches long.

LEAVES— Ornamentally cut and lobed. Stem contains a strong, yellow juice.

NOTES.

---

1. Agrimony.

2. Celandine.

1. LOOSESTRIFE: *Lysimàchia terrèstris*.
Primrose Family. June—Aug

Found in low, wet ground. Newfoundland to Northern Georgia, west to Minnesota, Arkansas and Manitoba.

FLOWERS—Yellow, with reddish dots near the center, borne in elongated cluster terminating stem, 1-2 feet high.

LEAVES— Opposite, tapering at both ends.

2. FOUR-LEAVED LOOSESTRIFE; Whorled Loosestrife: Crosswort. *Lysimàchia quadrifolia*.
Primrose Family. June—Aug.

A slender plant 1-3 feet tall, found in open woods, moist, sandy soil. New Brunswick to Georgia, west to Illinois and Minnesota.

FLOWERS—Yellow, dotted around the center or faintly streaked with red, borne on slender stems springing from the leaf angles.

LEAVES— In whorls of 3-7, usually 4.

NOTES.

---

1. Loosestrife.

2. Four-leaved Loosestrife.

CONE-FLOWER; PURPLE CONE-FLOWER:

Black-eyed Susan: Yellow or Ox-eye Daisy:

Nigger-head: Golden Jerusalem.

Rudbékia hirta.

Composite Family.

June—Sept.

A plant 1–3 feet tall, found in fields and open, sunny places. New England to Gulf States, Ontario and Northwest to Colorado.

FLOWER-HEADS—Rays 10–20, yellow or orange-yellow, surrounding a conical, dark purplish-brown disk.

LEAVES— Nearly or quite toothless, lower leaves broader at the tip, 3-ribbed. Leaves and stems rough-hairy.

TALL CONE-FLOWER: Green-headed Cone-flower: Thimble-weed.

R. laciniàta.

Rays 6–10, yellow, slightly drooping, surrounding a greenish yellow disk. Leaves variously divided, lobed and toothed. A plant 3–10 feet tall. Found in moist thickets. Quebec to Montana, south to Gulf States.

NOTES.

Cone-flower.

1. COMMON MULLEIN: Great Mullein: Velvet or Flannel Plant: Mullein Dock: Aaron's Rod. *Verbáscum Thápsus*.

Figwort Family.

June—Sept.

A plant 2-7 feet tall, found in dry fields, waste places. Nova Scotia to Florida, west to Minnesota and Kansas. Naturalized from Europe.

FLOWERS—Yellow, 5-parted, in dense, elongated spike.

LEAVES— Thick, velvety, in a rosette on the ground, others clasping the plant stem.

2. MOTH MULLEIN: *Verbáscum Blattària*.

Figwort Family.

June—Sept.

A plant 2-5 feet tall, found in dry fields, waste places. Fairly common through United States and Canada. Naturalized from Europe.

FLOWERS—Yellow, sometimes white, tinged on the back with lavender, fringed in the center with ruddy hairs; borne in loose spike.

LEAVES— Smooth, more or less deeply toothed.

NOTES.

1

1. Common Mullein.

2

2. Moth Mullein.

WILD LETTUCE: Tall Lettuce: Wild Opium:
Horse-weed. *Lactuca canadensis.*

Composite Family. June—Sept.

Found in roadsides, borders of fields, thickets, moist soil. Canada to Georgia, west to Arkansas.

FLOWER-HEADS—Pale-yellow, numerous, small, in loose, spreading cluster terminating stem, 4–10 feet high.

LEAVES— Lower leaves deeply toothed, often 1 foot long, narrowed toward the base; upper leaves lance-shaped, toothless.

NOTES.

Wild Lettuce.

1. YELLOW or HOP CLOVER:

Trifolium agrarium.

Pulse Family.

June—Sept.

Found in sandy fields, roadsides. Naturalized from Europe. Virginia to Iowa, far northward.

FLOWERS—In heads composed of many tiny, yellow florets, resembling small, dried hops when withered.

LEAVES— In groups of 3 leaflets, almost imperceptibly toothed.

2. LADY'S SORREL: Yellow Wood-sorrel.

Oxalis corniculata.

Wood Sorrel Family.

May—Sept.

Common weed in all situations. Nova Scotia to Gulf, west to Dakota.

FLOWERS—Yellow, small, 5 divisions.

LEAVES— Composed of 3 heart-shaped leaflets.

NOTES.

---

1. Yellow or Hop Clover.

2. Lady's Sorrel.

BLACK MUSTARD:*Brássica nìgra.*

Mustard Family.

June—Sept.

A plant 2-5 feet high, found in fields and roadsides. Naturalized from Europe. Common throughout United States.

FLOWERS—Yellow, 4-parted, in loose clusters terminating branches. Seed pods about $\frac{1}{2}$ inch long, 4-sided, lying close to the stem.

LEAVES— Variously lobed and toothed, terminal lobe largest.

YELLOW ROCKET; WINTER CRESS: Herb of St. Barbara: Yellow Bitter, or Rocket Cress.

Barbarèa vulgàris.

Mustard Family.

Apr.—June.

Flowers similar to those of preceding; seed pods long, curved. Upper leaves stemless, lower leaves usually in 5 divisions, terminal one largest. Lake Superior northward and westward; apparently introduced eastward.

NOTES.

Black Mustard.

NOTES.

NOTES.

---

154 YELLOW AND ORANGE

1. COMMON EVENING PRIMROSE: Evening Primrose: Night Willow-herb.

Oenothèra biénnis.

Evening Primrose Family. June—Sept.

A plant 1-5 feet high, found in fields and roadsides. Labrador to Gulf, west to Rockies.

FLOWERS—Yellow, 4-parted, 1-2 inches across, in terminal spikes, opening in the evening.

LEAVES— Alternate, lance-shaped, mostly seated on stem, obscurely toothed. Plant stem leafy, hairy, more or less branched.

2. SUNDROPS: Common Sundrops:

Oenothèra fruticòsa.

Evening Primrose Family. May—July.

Plant 1-3 feet tall, found in fields, roadsides.

New England to South Carolina, west to Indiana and Michigan.

FLOWERS—Yellow, $\frac{1}{2}$ -1 inch across, 4 divisions, terminating stem and from the leaf angles. Seed pods ribbed and winged.

LEAVES— Oblong to lance-shaped, minutely toothed.

O. pùmila is a similar but smaller species. Leaves toothless, basal leaves broader at the tip. More western in its range.

1, Common Evening Primrose.

2. Sundrops.

SPOTTED TOUCH-ME-NOT; JEWEL-WEED: Silver Cap: Lady's Ear-drop: Snap Weed: Wild Balsam: Wild Lady's Slipper.

Impatiens biflora.

Touch-me-not Family.

June—Sept.

A smooth-stemmed, branching plant, 2–5 feet high, found on brooksides and in moist, shady places. Nova Scotia to Oregon, south to Missouri and Florida.

FLOWERS—Yellow, more or less spotted with reddish brown; sac longer than it is broad, tapering to an incurved spur. Seed pods burst when touched.

LEAVES— Long-stemmed, coarsely toothed.

PALE TOUCH-ME-NOT:

I. pallida.

Flowers pale yellow, sparingly dotted with reddish brown; sac broader than long with short spur. Most abundant northward.

NOTES.

Spotted Touch-me-not.

COMMON ST. JOHN'S WORT:

Hypericum perforatum.

St. John's-wort Family.

June—Sept.

A plant 1-2 feet high, found in fields, road-sides. Common throughout the United States.

Naturalized from Europe.

FLOWERS—Yellow, in clusters terminating the branches; divisions 5, black-dotted along the margins.

LEAVES— Small, opposite, oblong, more or less black-dotted.

NOTES.

Common St. John's-wort.

BUTTER-AND-EGGS: Yellow Toad Flax: Flax-weed: Brideweed: Eggs and Bacon.

Linària vulgàris.

Figwort Family.

June—Oct.

A plant 1-3 feet tall, found in fields, roadsides, waste places. Nova Scotia to Virginia, west to Manitoba and Nebraska. Naturalized from Europe.

FLOWERS—Yellow and orange, 2-lipped, with a long spur, borne on terminal spikes.

LEAVES— Long, narrow, numerous.

NOTES.

Butter-and-eggs.

1. YELLOW FRINGED ORCHIS:

Habenària ciliàris.

Orchis Family.

July—Aug.

Found in moist meadows, sandy bogs. Vermont and Massachusetts to Michigan, Missouri, and southward.

FLOWERS—Golden or orange-yellow, lower lip fringed, borne in spike terminating a stem, 1–2 feet high.

LEAVES— Lance-shaped, clasping.

2. DOWNY FALSE FOXGLOVE:

Geràrdia flàva.

Figwort Family.

July—Aug.

A plant 2–4 feet tall, found in open woods. New England to Gulf States, west to Ontario and Wisconsin, Iowa, Arkansas.

FLOWERS—Yellow, trumpet-shaped, 5-lobed.

LEAVES— Opposite, lower ones more or less irregularly lobed and toothed. Plant stem and leaves downy.

SMOOTH FALSE FOXGLOVE: *G. virginica.*

Flowers slightly larger than those of preceding. Lower leaves deeply cut and toothed. Whole plant smooth.

1. Yellow Fringed Orchis.

2. Downy False Foxglove.

ELECAMPANE: Horseheal: Yellow Starwort.

Inula Helènum.

Composite Family.

July—Sept.

Plant 2-6 feet tall. Found in roadsides, fields, damp pastures. Naturalized from Europe. Minnesota and Missouri east to Nova Scotia, south to South Carolina.

FLOWER-HEADS—Yellow, solitary or few, 2-4 inches across, rays long and narrow.

LEAVES— Large, white-veined, rough above, woolly beneath, toothed; lower leaves stemmed, upper ones partly clasping plant stem.

NOTES.

Elecampane.

1. PARTRIDGE PEA: Large-flowered Sensitive Plant: *Cássia Chamaecrista.*

Pulse Family. July—Sept.

A plant 1–2 feet tall, found in dry or sandy fields. Massachusetts to Minnesota and southward.

FLOWERS—Yellow, 5-parted, about 1 inch across, 2–4 from the bases of the leaves.

LEAVES— Composed of 10–15 pairs of leaflets.

- WILD SENNA: *C. marilándica.*

Flowers yellow, divisions 5, 3 forming an upper lip, 2 a lower one. Leaflets 5–9 pairs. 3–8 feet high. Moist soil, swamps. New England to Ohio, Tennessee, and North Carolina.

2. FALL DANDELION: Autumnal Hawkbit.

Leóntodon autumnàlis.

Composite Family. July—Nov.

Found in fields, roadsides. Throughout northern United States, south to Ohio and New Jersey and Pennsylvania. Naturalized from Europe.

FLOWER-HEADS—Yellow, smaller than the common dandelion, on slender, branching stalks, 6–18 inches high.

LEAVES— Long, narrow, more or less coarsely toothed.

1. Partridge Pea.

2. Fall Dandelion.

NOTES.

NOTES.

TANSY: Bitter-buttons. *Tanacètum vulgàre.*

Composite Family. July—Sept.

Found in roadsides, waste places and around old buildings. Introduced from Europe.

Nova Scotia to North Carolina and Missouri, west to Minnesota.

FLOWER-HEADS—Yellow, in rather flat clusters terminating stem, $1\frac{1}{2}$ –3 feet tall.

LEAVES— Deeply cut into narrow, toothed divisions, aromatic.

NOTES.

Tansy.

ORANGE or TAWNY HAWKWEED; DEVIL'S
PAINT-BRUSH: Grim the Collier.

Hieracium aurantiacum.

Composite Family.

July—Sept.

Found in fields, pastures. Naturalized from
Europe. Canada south to Pennsylvania and
Middle Atlantic States.

FLOWER-HEADS—Reddish orange, in a group ter-
minating stem, 6–20 inches high.

LEAVES— Mostly at the base, blunt, lance-shaped.
Whole plant hairy.

CANADA HAWKWEED:

H. canadense.

FLOWERS—Yellow, terminating stem 1–4 feet tall.

LEAVES— Seated on stem, coarsely toothed, espe-
cially the lower ones.

NOTES.

Orange Hawkweed.

1. HAWKWEED: Rough Hawkweed.

Hieracium scabrum.

Composite Family.

July—Sept.

A plant 1–3 feet high, found in dry woods, pastures. Nova Scotia to Georgia, west to Nebraska.

FLOWER-HEADS—Yellow, in branching, terminal clusters.

LEAVES— Blunt, almost toothless, seated on stem. Whole plant hairy.

HAIRY HAWKWEED:

H. Gronovii.

A similar plant with slender stem, very leafy and hairy below.

2. RATTLESNAKE-WEED: Early or Vein-leaf Hawkweed: Snake or Poor Robin's Plantain.

Hieracium venosum.

Composite Family.

June—Sept.

Found in dry woods, open, sandy places. Atlantic States to Minnesota and Iowa.

FLOWER-HEADS—Yellow, resembling small dandelions, few on slender, branching stalk, 10–20 inches high.

LEAVES— In a rosette, hairy, with purplish veins.

NOTES.

1. Hawkweed.

2. Rattlesnake-weed.

1. THIN-LEAVED or TEN-PETALED SUN-
FLOWER: *Helianthus decapétalus.*

Composite Family.

Aug.—Sept.

A plant 2–5 feet tall, found in moist woods, beside streams. Eastern United States to Minnesota, Missouri and southward.

FLOWER-HEADS—Yellow, 2–3 inches broad, rays about 10.

LEAVES— Egg-shaped, pointed, opposite, smooth or slightly rough. Plant stem rough above; smooth below, 2–5 feet tall.

2. LARGER BUR-MARIGOLD: Brook Sun-
flower. *Bidens laevis.*

Composite Family.

Aug.—Oct.

Plant 1–2 feet high, found in swamps, wet places. New England to Minnesota and southward.

FLOWER-HEADS—Yellow, 1–2½ inches across, rays 8–10. Seed vessels with 2–4 prongs.

LEAVES— Opposite, toothed, narrow lance-shaped, seated on stem.

NOTES.

1. Thin-leaved Sunflower.

2. Larger Bur-Marigold.

NOTES.

GOLDEN CORYDALIS: *Corýdalis áurea.*

Fumitory Family. Mch.—May.

A smooth-stemmed, branching plant 6–14 inches high, found in woodlands. Nova Scotia to Pennsylvania, west to the Rocky Mountains.

FLOWERS—Golden yellow, irregular, about $1\frac{1}{2}$ inch long, borne in short, terminal clusters. Seed pods drooping or spreading, slightly curved, lumpy.

LEAVES—Compound, finely dissected.

ROUND-LEAVED, or EARLY YELLOW VIOLET:*Viola rotundifolia.*

Violet Family. Apr.—May.

Found in woodlands. A plant 2–4 inches high. New England to Georgia.

FLOWERS—Yellow, with purplish veins, small, side divisions bearded.

LEAVES—Round or long heart-shaped, indistinctly toothed.

DOWNY YELLOW VIOLET:

Viola pubescens.

Violet Family.

Apr.—May.

An erect leafy plant 6–15 inches high, common in dry woodlands. Maine to Ontario, south to Maryland and Kansas.

FLOWERS—Golden yellow, with purplish veins, borne on stalks from the leaf angles.

LEAVES— Broadly heart-shaped, somewhat toothed. Plant stem and leaves slightly downy.

GOLDEN CLUB:

Oróntium aquáticum.

Arum Family.

Apr.—May.

Found in shallow water, swamps, chiefly near the Atlantic coast. New England to Gulf of Mexico.

FLOWERS—Yellow, minute, crowded on a club 1–2 inches long, terminating stalk 6–36 inches high; the stalk flattened just below the flower cluster.

LEAVES— Dark green, oblong, long-stemmed, floating or erect.

INDIAN POKE; AMERICAN WHITE HEL-
LEBORE: Itch-weed. *Veràtrum víride.*

Lily Family. Apr.—June.

A plant 2–7 feet high, poisonous throughout;
found in low, wet places. Canada to British
Columbia, south to Georgia, Tennessee, Min-
nesota.

FLOWERS—Very small, forming a large, terminal,
yellow-green cluster.

LEAVES— Large, broad, with an uneven surface,
soon turning brown and unsightly.

YELLOW MELILOT or SWEET CLOVER:

Melilòtus officinàlis.

Pulse Family. June—Aug.

Found in waste places. Naturalized from
Europe.

FLOWERS—Resemble the White Melilot, except in
color, yellow, small, in slender spikes
terminating branches, 2–4 feet high.

LEAVES— Small, in groups of 3, blunt tipped,
toothed.

FROSTWEED; ROCKROSE: Frost-flower,
Frost-wort: Long-branched Frost-weed: Ca-
nadian Rockrose. *Helianthemum canadense*.

Rockrose Family. June—Aug.

Found in sandy soil. New England to South
Carolina, west to Wisconsin.

FLOWERS—Of 2 kinds: the earlier ones solitary,
about 1 inch across, yellow, 5 divisions;
later ones small, clustered at the base of
the leaves.

LEAVES— Lance-oblong, pale beneath, set close
to the stem.

NOTES.

BUTTERFLY WEED; PLEURISY ROOT:

Orange-root: Orange Milkweed.

Asclèpias tuberosa.

Milkweed Family.

June—Sept.

Found in dry fields. Maine to Gulf States, west to Ontario and Arizona. Common southward.

FLOWERS—Similar in structure to the Common Milkweed, reddish orange or orange-yellow, in erect clusters terminating stem, 2–3 feet high.

LEAVES—Narrow-oblong, or lance-shaped, toothless, hairy beneath, nearly or quite stemless.

GINSENG:

Panax quinquefolium.

Ginseng Family.

July—Aug.

Found in rich woods. Quebec to Alabama, west to Nebraska.

FLOWERS—Inconspicuous, greenish yellow, in a cluster terminating stem, 8–15 inches high. Berries red in a cluster.

LEAVES—Consisting of 3 sets in a whorl, each set composed of 5 somewhat egg-shaped, sharp-pointed, toothed leaflets.

HORSE BALM; RICH-WEED: Horse-weed:
Stone-root: Citronella.

Collinsònia canadénsis.

Mint Family.

July—Sept.

A plant 2-4 feet tall, found in rich, moist woods. New England to Florida, Ontario and Wisconsin to Kansas and Missouri.

FLOWERS—Pale yellow, about $1\frac{1}{2}$ inch long, with 3 long, protruding filaments, lemon-scented; 5-lobed, one lobe fringed and much larger than the others; borne in loose, spreading cluster.

LEAVES—Oval, toothed. Plant stem branching.

SNEEZÉWEED; SWAMP SUNFLOWER:

Helènum autumnàle.

Composite Family.

Aug.—Sept.

A plant 2-6 feet high, found in wet meadows, river banks. Quebec to Florida, west to Arizona, Minnesota and northwest.

FLOWER-HEADS—Yellow, 1-2 inches across, rays drooping, broad, 3-5 cleft, in terminal clusters.

LEAVES—Lance-shaped or oblong, alternate, toothed.

FERN-LEAVED FALSE FOXGLOVE:

Gerárdia pediculària.

Figwort Family.

Aug.—Sept.

A plant 1–3 feet high, found in borders of dry woods, thickets. Maine to Ontario, Minnesota, and West Virginia.

FLOWERS—Yellow, bell-shaped, with 5 spreading, rounded lobes, 1 inch or more across, somewhat finely hairy.

LEAVES— Fern-like, deeply cut into many toothed lobes. Plant stem leafy, branching, somewhat sticky-hairy.

GOLDEN ASTER:

Chrysópsis mariàna.

Composite Family.

Aug.—Oct.

Found in dry, sandy soil. Gulf of Mexico, north to Pennsylvania and Long Island.

FLOWER-HEADS—Yellow, aster-like, nearly 1 inch across, in loose, spreading clusters.

LEAVES— Oblong, seated on stem, toothless. Leaves and plant stem silky-hairy when young, becoming smooth with age.

GOLDEN-ROD. The golden-rods are familiar to everyone. They comprise a very extensive group, with many intergrading forms, and none but students care to puzzle out the different species.

NOTES.

III

RED, PINK, CRIMSON AND
MAGENTA

SPRING BEAUTY: Claytonia.

Claytonia virginica.

Purslane Family.

Mch.—May.

A plant 6–12 inches high, found in moist, open woods, low meadows. Common west and south.

FLOWERS—White veined with pink, or pink with veins of deeper pink; divisions 5.

LEAVES— Long and narrow, width variable.

Roots tuberous.

C. caroliniana is similar to the preceding, except that the flowers are smaller and fewer, and leaves more oblong.

Spring Beauty.

1. SHOOTING STAR; AMERICAN COW-SLIP: Pride of Ohio. *Dodecàtheon Meàdia.*

Primrose Family. Apr.—June.

Found in open woods, prairies, moist cliffs.

Wisconsin to Pennsylvania, south to Texas and Georgia.

FLOWERS—Magenta-pink or white, the 5 divisions turned backward, pendulous, grouped on stalk 8–20 inches high.

LEAVES— Oblong, narrowed toward the base.

2. ARETHUSA: Indian Pink.

Arethusa bulbosa.

Orchis Family.

May—June.

Found in bogs and swamps. Newfoundland south to South Carolina mountains, west to Indiana and Minnesota.

FLOWERS—Magenta-crimson, lower lip spotted with magenta crimson and crested with 3 white or yellow hairy ridges; solitary, on stalk 5–10 inches high.

LEAVES— Long and narrow, solitary, appearing after the flower opens.

Root bulbous.

1. Shooting Star.

2. Arethusa.

WAKE ROBIN: Purple Trillium: Ill-scented

Wake Robin: Birthroot. *Trillium erectum.*

Lily Family.

Apr.—June.

Plant 8–16 inches high, found in rich woods.

Eastern Quebec south to North Carolina, west to Minnesota and Missouri.

FLOWERS—Brown-purple, dull pink, white or greenish, divisions 3, on short stalk from junction of leaves.

LEAVES— Broad, 3 in a whorl.

SESSILE-FLOWERED WAKE ROBIN: Stemless Trillium. *T. sessile.*

Flowers stemless, dark purplish or greenish.

T. recurvatum. Outer divisions of flower reflexed; colored divisions narrowed at the base; leaves narrowed into a stem.

NOTES.

Wake Robin.

WILD COLUMBINE: Honeysuckle.

Aquilegia canadensis.

Crowfoot Family.

Apr.—June.

A slender plant 1–2 feet high, found in open, rocky woodlands, pastures. Common throughout the United States.

FLOWERS—Red outside, yellow within; 5 tubes terminating in spurs; borne on slender stalks from the leaf angles.

LEAVES—Compound, more or less divided, teeth rounded.

NOTES.

Wild Columbine

1. WILD PINK: Catchfly. *Silène pennsylvànica*.
Pink Family. Apr.—June.

Found in gravelly, rocky places. Eastern New England to New York and Kentucky, south to Georgia.

FLOWERS—Crimson-pink, divisions 5, somewhat sticky immediately below the flower, in terminal clusters, 4–10 inches high.

LEAVES— Blunt lance-shaped, mostly at the base; a few opposite, lance-shaped leaves seated on the stems.

2. MAIDEN PINK: *Diánthus deltoïdes*.
Pink Family. June—Aug.

Found in fields, waste places. New England to Michigan. Naturalized from Europe.

FLOWERS—Crimson-pink or white, 5 toothed divisions, blooming singly, terminating slender stems 6–12 inches high.

LEAVES— Narrow, erect.

- DEPTFORD PINK: *D. Armèria*.

Flowers crimson-pink, white-dotted, divisions toothed. Plant stem and leaves finely hairy. Naturalized from Europe.

1

1. Wild Pink.

2

2. Maiden Pink.

1. WILD CRANESBILL; WILD or SPOTTED GERANIUM:

Geranium maculatum.

Geranium Family.

Apr.—July.

A plant 1–2 feet high, found in open woods, roadsides. Maine to Manitoba and southward.

FLOWERS—Pale or deep magenta-pink or lavender, divisions 5.

LEAVES— Usually 5-parted variously toothed, stem leaves opposite. Leaves and plant stem hairy.

2. HERB ROBERT: Red Robin: Red Shanks: Dragon's Blood.

Geranium Robertianum.

Geranium Family.

June—Oct.

Found in moist, rocky woods. Eastern Quebec to Minnesota, south to New Jersey and Missouri.

FLOWERS—Deep or pale magenta or pink, about $\frac{1}{2}$ inch across, divisions 5.

LEAVES— 3–5 parted, variously cleft and toothed, strongly scented.

Stems spreading, hairy, 6–18 inches high.

1. Wild Cranesbill.

2. Herb Robert.

NOTES.

NOTES.

PITCHER-PLANT: Indian Dipper: Side-saddle
Flower: Huntsman's Cup.

Sarracènia purpùrea.

Pitcher-plant Family.

May—June.

Found in mossy bogs. Labrador to Minnesota, Rocky Mountains, southeast of Alleghenies to Florida.

FLOWERS—Variable in color, the 5 outer divisions greenish to madder purple, inner divisions dull pink; solitary, nodding, borne on stalk 1–2 feet high.

LEAVES—Hollow, usually green with reddish purple veins.

NOTES.

Pitcher-plant.

STEMLESS, PINK, or VENUS' LADY'S SLIP-

PER: Moccasin Flower. *Cyripedium acaúle.*

Orchis Family.

May—June.

Found in dry woods. Newfoundland to North Carolina, west to Indiana, Michigan, Minnesota.

FLOWERS—Pouch variable crimson, veined with deeper crimson, sometimes white, solitary, on stalk 6–12 inches high.

LEAVES— 2, from the base, 6–8 inches long.

NOTES.

Stemless Lady's Slipper.

SHOWY ORCHIS: Gay, or Spring Orchis.

Orchis spectábilis.

Orchis Family.

May—June.

Found in rich woods. New Brunswick to Georgia, west to Minnesota and Missouri.

FLOWERS—Magenta, lower lip almost white, 3-6 borne on stalk 4-12 inches high.

LEAVES— 2, broad, shining, from the base.

NOTES.

Showy Orchis.

WILD AZALEA: Pinxter Flower: Wild Honey-suckle: Pink or Purple Azalea.

Rhododéndron nudiflòrum.

Heath Family.

May—June.

A shrub 2–6 feet high, found in open woods, swamps. Massachusetts to Florida, west to Illinois, Missouri, Texas.

FLOWERS—Crimson-pink to nearly white, the base of the long, narrow tube deeper in color, divisions 5, 6 long, protruding filaments; flowers borne in small, terminal clusters, opening with or before the leaves.

LEAVES— Usually clustered, tapering at each end.

NOTES.

Wild Azalea.

1. FRINGED MILKWORT; FLOWERING WINTERGREEN: Gay Wings: Fringed Polygala. *Polygala paucifolia.*

Milkwort Family. May—July.

Found in moist woods. Eastern Quebec to Minnesota and Illinois, south in Alleghenies.

FLOWERS—Magenta, magenta-crimson, or rarely white, very irregular in form, fringed on the tip.

LEAVES— Clustered at the summit of the stem, 3–6 inches high.

2. FIELD MILKWORT: Common or Purple Milkwort: Purple Polygala.

Polygala sanguinea.

Milkwort Family. June—Sept.

Found in moist or sandy fields, roadsides.

Southern Canada to North Carolina, west to the Mississippi.

FLOWERS—Deep or pale magenta, rarely white, tiny, clustered in globular or oblong heads, terminating stems 6–12 inches high.

LEAVES— Small, narrow

1. Fringed Milkwort.

2

2. Field Milkwort.

MOUNTAIN or AMERICAN LAUREL: Calico

Bush: Spoonwood: Calmoun: Broad-leaved
Kalmia. *Kálmia latifolia*.

Heath Family.

May—July.

Found in rocky woods. Canada and Maine mountains to Gulf of Mexico, west to Ohio and Tennessee. Usually a shrub 3–6 feet high, but in its southern range, often tree-like.

FLOWERS—White or pink, bowl-shaped, 5-pointed; 10 filaments spring from the center, their tips lodged in little pockets; flowers borne in clusters terminating the branches.

LEAVES— Long, evergreen, tapering at both ends.

NOTES.

Mountain Laurel.

TWISTED-STALK: Sessile-leaved Twisted Stalk.

Stréptopus ròseus.

Lily Family.

May—July.

A plant 1-2½ feet high, found in moist woods. Newfoundland to Minnesota, south to Ohio, Pennsylvania, mountains of Georgia.

FLOWERS—Bell-shaped, dull, purplish pink, solitary, from the axils of the leaves.

LEAVES— Alternate, seated on stem.

CLASPING-LEAVED TWISTED-STALK:

S. amplexifolius.

Differs from the preceding in its greenish flowers with widely spreading segments, and whitish bloom on the under side of the leaves. Leaves strongly clasping.

NOTES.

Twisted-stalk.

1. WOOD BETONY; LOUSEWORT: Beefsteak

Plant: High Heal-all. *Pedicularis canadensis*.

Figwort Family.

May—July.

Found in open woods. Nova Scotia to Florida, west to Manitoba, Kansas, Colorado.

FLOWERS—Greenish yellow and purplish red, 2-lipped, in short, dense clusters, terminating stems 6–18 inches high.

LEAVES— Long, narrow, deeply toothed.

2. BOUNCING BET; SOAPWORT: Hedge

Pink: Bruisewort: Old Maid's Pink: Fuller's Herb.

Saponaria officinalis.

Pink Family.

June—Sept.

A plant 1–2 feet high, found in roadsides, waste places. Adventive from Europe.

Common, escaped from gardens.

FLOWERS—Magenta-pink to white, divisions 5, but often double, borne in terminal clusters.

LEAVES— Opposite, toothless, 3–5-ribbed. Plant stem stout, thick jointed.

1

1. Wood Betony

2

2. Bouncing Bet.

PALE CORYDALIS: *Corýdalis sempérvirens.*

Fumitory Family.

May—Aug.

Found in rich, rocky places. Eastern Quebec to Alaska, south to Georgia, Kentucky, and Montana.

FLOWERS—Crimson-pink about $1\frac{1}{2}$ inch long, in loose clusters terminating branches, 1-2 feet high.

LEAVES— Compound, variously cut and divided.

NOTES.

Pale Corydalis.

SHEEP LAUREL; LAMBKILL: Wicky: Sheep-poison: Calf-kill: Narrow-leaved Laurel.

Kálmia angustifolia.

Heath Family.

June—July.

A shrub 1–3 feet high, found on hillsides and in swamps. Labrador and Ontario to Michigan, south to Georgia.

FLOWERS—Crimson-pink, in clusters encircling plant stem.

LEAVES—Narrow, opposite, or in groups of 3. Stem terminated by the upright newer leaves.

PALE LAUREL: Swamp Laurel. *K. polifolia.*

May be distinguished by its flowers terminating the stems, and by its leaves, which are usually set opposite, edges rolled back, whitish beneath. Found in cold bogs and hillside swamps. Labrador to Alaska, south to Pennsylvania and Michigan, west to Pacific coast.

Sheep Laurel.

1. GRASS PINK: *Calopogon*.*Calopogon pulchellus*.

Orchis Family.

June—July.

Found in open bogs, meadows. Newfoundland to Florida, west to Minnesota and Missouri.

FLOWERS—Magenta-pink, the lip on the upper side of flower densely bearded with yellow and magenta-crimson hairs; 3–12 flowers terminate stalk 10–16 inches high.

LEAF—Solitary, grass-like, from bulb.

2. ROSE, or SWEET POGONIA; SNAKE MOUTH:

Pogonia ophioglossoides.

Orchis Family.

June—July.

Found in wet meadows, swamps. Newfoundland to Florida, west to Indiana and Minnesota.

FLOWERS—Crimson-pink, crest of lower lip yellow to white, solitary, terminating stem 8–15 inches high.

LEAVES—A small leaflet borne just below the flower, and a larger leaf midway of the stem.

1. Grass Pink.

2. Rose Pogonia.

WILD ORANGE-RED LILY; WOOD LILY;
Flame or Philadelphia Lily.

Lilium philadelphicum.

Lily Family.

June—July.

Found in dry, sandy soil, woodland borders.

New England to North Carolina, west to
Minnesota and Missouri.

FLOWERS—Orange or reddish orange, spotted inside with purplish brown; 1-3 borne erect at the summit of stem 1-3 feet high.

LEAVES— In whorls of 3-8.

NOTES.

Wild Orange-red Lily.

1. SPREADING DOGBANE: Fly-trap Dogbane: Bitter-root: Honey-bloom: Indian Hemp.

Apócynum androsaemifólium.

Dogbane Family.

June—Aug.

Found in open woods, roadsides. Northern Canada to Georgia, west to Nebraska.

FLOWERS—Pink tinted, veined inside with deeper pink, bell-shaped with 5 spreading lobes, in loose clusters on the tips of the branches.

LEAVES— Opposite, toothless, short-stemmed. Plant stem branching, shrubby, 1-4 feet high.

2. TWIN-FLOWER: Ground Vine.

Linnaèa boreàlis.

Honeysuckle Family.

June—Aug.

A trailing vine, stems 6-20 inches long, found in moist, mossy woods. Labrador to New Jersey and mountains of Maryland, west to Minnesota; also far west.

FLOWERS—Whitish, tinged and striped with crimson-pink, fragrant, slender bell-shaped, 5-lobed, borne in pairs, on stalk 3-4 inches high.

LEAVES— Opposite, rounded, scallop-toothed, evergreen.

1. Spreading Dogbane.

2. Twin-flower.

1. MEADOW-SWEET: Queen-of-the-Meadow:
Quaker Lady. *Spiraea latifolia*.

Rose Family. June—Aug.

A shrubby plant, 2-4 feet high, found in rocky pastures, borders of fields. Newfoundland to Georgia, west to Minnesota and Missouri.

FLOWERS—Small, flesh pink or whitish, in dense, terminal clusters.

LEAVES— Light green, oval or oblong, rather coarsely toothed. Plant stems red or purplish brown.

S. salicifolia differs from the preceding in its white flowers, yellowish brown stems, and finely toothed leaves. Chiefly in low ground.

2. HARDHACK; STEEPLE BUSH:

Spiraea tomentosa.

Rose Family. July—Sept.

A shrubby plant, with erect, woolly stems 2-4 feet high, found in low ground or on hillsides. New Brunswick to Georgia, west to Minnesota and Kansas.

FLOWERS—Pink or magenta, rarely white, in dense, steeple-like clusters.

LEAVES— Dark green above, whitish-woolly beneath, toothed.

1. Meadow-sweet.

2. Hardhack.

PURPLE FLOWERING RASPBERRY: Virginia Raspberry. *Rubus odoratus*.

Rose Family. June—Aug.

A sprawling shrub, 3-5 feet high, found in rocky woods, shaded roadsides. Nova Scotia to Georgia, west to Michigan.

FLOWERS—Crimson-pink, fading as they get older, 1-2 inches broad. Fruit a thin, red, scarcely edible berry.

LEAVES— Large, lobed and toothed, hairy, similar to those of the maple in shape. Plant stems hairy.

NOTES.

Purple Flowering Raspberry.

NOTES.

NOTES.

---

1. LARGE, or EARLY, PURPLE FRINGED ORCHIS: *Habenària fimbriàta.*

Orchis Family. June—Aug.

Found in wet woods, swamps. Newfoundland to North Carolina, west to Michigan.

FLOWERS—Magenta-pink, varying to almost white, lower lip 3-parted, deeply fringed, in spike terminating stem 1–5 feet high.

LEAVES— Lance-shaped above, somewhat oval below.

SMALLER PURPLE FRINGED ORCHIS:

H. psycòdes.

Bears smaller blossoms, less deeply fringed.

A plant 1–3 feet high, found in similar situations west to Indiana and Minnesota.

2. PRINCE'S PINE; PIPSISSEWA:

Chimáphila umbellàta.

Heath Family. June—Aug.

Found in dry woods. Nova Scotia to Georgia, west to Pacific.

FLOWERS—Flesh color or pinkish, waxy, 5-parted, in small groups terminating stem 3–10 inches high.

LEAVES— Dark, shining green, evergreen, sharply toothed, opposite or in whorls.

SPOTTED WINTERGREEN: Pipsissewa.

C. maculàta.

Leaves white veined, tapering toward the tip.

Found in similar situations west to Minnesota and Mississippi.

1

1. Purple Fringed Orchis.

2

2. Prince's Pine.

COMMON MILKWEED: Silkweed.

Asclèpias syriaca.

Milkweed Family.

June—Aug.

A plant 3–5 feet high, found in fields, roadsides, waste places. Common. New Brunswick south to North Carolina and Kansas; also westward.

FLOWERS—Variable in color, dull crimson-pink, lilac or yellowish; one or more thick clusters springing from the leaf angles. Seed pods large, filled with silky down and flat, overlapping seeds.

LEAVES— Opposite, broad oblong, toothless.

PURPLE MILKWEED:

A. purpurascens.

Similar in habit to the preceding. Found in woods, dry fields. New Hampshire to Ontario, Minnesota, Kansas, and southward.

FLOWERS—Crimson or crimson-magenta, in terminal clusters.

LEAVES— Elliptical or egg-shaped, upper ones taper-pointed, minutely downy beneath.

NOTES.

Common Milkweed.

HEDGE or GREAT BINDWEED; WILD
MORNING-GLORY: Bell-bind: Rutland
Beauty: Lady's Nightcap.

Convólvulus sèpium.

Convolvulus Family.

June—Sept.

A vine 3–10 feet long, found in thickets,
fields, waysides. North Atlantic States west-
ward.

FLOWERS—Pink or white, large, bell-shaped, on
long stems from leaf angles.

LEAVES— Triangular or arrowhead-shaped, long-
stemmed.

FIELD BINDWEED:

C. arvénis.

Similar to the preceding, but much smaller.

Flowers not over 1 inch long, pinkish or
white, often in pairs. Naturalized from Eu-
rope. North Atlantic States west to Kansas.

NOTES.

Hedge Bindweed,

1. LARGE MARSH PINK:

Sabàtia dodecándra.

Gentian Family.

July—Aug.

A plant 1–2 feet high, found on borders of brackish ponds near the Atlantic coast.

Massachusetts to North Carolina.

FLOWERS—Crimson-pink, about 2 inches across, 8–12 divisions.

LEAVES— Lower leaves blunt-tipped, tapering toward the base, upper leaves narrow, lance-shaped.

BITTER BLOOM: Rose Pink: Square-stemmed

Sabatia: Rose Centaury.

S. angulàris.

Flowers crimson-pink or white, greenish star in the center, 5-parted. Leaves opposite, 5-veined, clasping the square plant stem.

Found in rich soil. New York to Florida, west to Indian Territory, Michigan, Ontario.

2. SEA or MARSH PINK: Rose of Plymouth.

Sabàtia stellàris.

Gentian Family.

July—Sept.

Found along the Atlantic coast.

FLOWERS—Crimson-pink, about 1 inch across.

LEAVES— Upper leaves narrow, widening as they approach the root.

SLENDER MARSH PINK:

S. gràcilis.

Resembles the Sea Pink, but stems are more slender, leaves narrower.

1

1. Large Marsh Pink.

2

2. Sea Pink.

SWAMP MILKWEED: *Asclèpias incarnàta*.

Milkweed Family.

July—Aug.

Found in swampy places. New Brunswick,
west and southwest.

FLOWERS—Dull crimson or crimson-pink, in clusters, terminating stem 2–4 feet high.

LEAVES— Narrow, lance-shaped, short-stemmed; 2 downy lines on stems below flower clusters.

FOUR-LEAVED MILKWEED:

A. quadrifòlia.

Bears 1–2 whorls of 4 leaves each near the middle of plant stem. Flowers pinkish and white. New Hampshire to North Carolina, Ontario to Arkansas.

NOTES.

Swamp Milkweed.

1. GREAT or SPIKED WILLOW-HERB;
FIREWEED: *Epilobium angustifolium*.

Evening Primrose Family. July—Aug.

Found in dry soil, clearings, newly burnt land.

New England to North Carolina, west to Minnesota and Kansas, and, with few exceptions, to the Pacific.

FLOWERS—Magenta, rarely white, divisions 4, in spike terminating stem 2–8 feet high.

LEAVES— Narrow, quite toothless, tapering.

2. CARDINAL FLOWER: *Lobelia cardinalis*.

Lobelia Family. Aug.—Sept.

Found in low, moist ground. New Brunswick to Ontario and southward.

FLOWERS—Rich, deep red, lower lip 3-lobed, in somewhat 1-sided spike terminating stem 2–4 feet high.

LEAVES— Dark green, oblong to lance-shaped, slightly toothed, upper leaves stemless.

1. Great Willow-herb.

2. Cardinal Flower.

1. MEADOW BEAUTY; DEER-GRASS:

Rhèxia virginica.

Meadow Beauty Family.

July—Sept.

A square-stemmed plant 10–18 inches high, found in sandy marshes or near water. Maine to Florida, Ontario to Iowa and southward.

FLOWERS—Magenta or purplish, division 4, 8 prominent, golden yellow, floral organs.

LEAVES— Opposite, mostly 5-veined, toothed, seated on stem.

MARYLAND MEADOW BEAUTY:

R. mariàna.

May be distinguished from the preceding by its round stems and short-stemmed leaves. Found south of New Jersey.

2. OSWEGO TEA; BEE BALM:

Monárda didyma.

Mint Family.

July—Sept.

A square-stemmed plant 2–3 feet high, found in moist ground, especially near streams, from New England west and south; also cultivated in gardens.

FLOWERS—Scarlet, 2-lipped, clustered in terminal heads, with a number of leaflets below.

LEAVES— Opposite, toothed, dark green, aromatic. Plant stem hairy.

1. Meadow Beauty.

2. Oswego Tea.

1. PINK KNOTWEED; SMARTWEED: Common *Persicaria*: Jointweed.

Polygonum pennsylvanicum.

Buckwheat Family. July—Sept.

A plant of sprawling, branching habit, 1–3 feet high, common in waste places, moist soil.

FLOWERS—Pink or greenish, small, crowded on short, narrow, terminal spikes.

LEAVES— Long, narrow, toothless, tapering toward the tip. Plant stems enlarged at the joints.

LADY'S THUMB: *P. Persicaria.*

Is a similar species, leaves usually marked with a dark triangle. Found throughout North America. Naturalized from Europe.

2. WATER PERSICARIA:

Polygonum amphibium.

Buckwheat Family. July—Aug.

Found in shallow water. Quebec to New Jersey, west to Pacific.

FLOWERS—Pink, small, in short, dense, terminal clusters.

LEAVES— Usually floating, smooth, shining above, elliptical or oblong.

1. Pink Knotweed.

2. Water Persicaria.

JOE-PYE WEED: Trumpet Weed: Gravel or
Kidney-root: Purple Thoroughwort: Tall or
Purple Boneset. *Eupatòrium purpùreum.*

Composite Family. Aug.—Sept.

A tall plant 3–10 feet high, common in low,
swampy places. New England to the Gulf of
Mexico and westward.

FLOWER-HEADS—Dull magenta or magenta-crim-
son, in broad, terminal cluster.

LEAVES— Lance-shaped, toothed, in whorls of 3–6.
Plant stem green or purplish.

NOTES.

Joe-Pye Weed.

RABBIT'S-FOOT, OLD-FIELD, PUSSY, or
STONE CLOVER: *Trifolium arvense*.

Pulse Family.

Aug.—Sept.

A plant 4–10 inches high, with branching
stem, found in fields, waste places, dry soil.

Naturalized from Europe.

FLOWERS—In soft, fuzzy heads, gray or gray-
pink in effect.

LEAVES— In groups of 3 leaflets, widest toward
the tip.

NOTES.

Rabbit's-foot Clover.

LIVE-FOR-EVER; GARDEN ORPINE: Mid-summer-men: Pudding-bag Plant: Live-long: Witches' Money: Garden-stonecrop.

Sedum purpureum.

Orpine Family.

Aug.—Sept.

Found in fields, waysides. Quebec to Maryland, Michigan and westward. Introduced from Europe.

FLOWERS—Small, dull garnet red or reddish purple, in thick, terminal clusters.

LEAVES—Thick, smooth, toothed, seated on stem 10–20 inches high.

NOTES

Live-for-ever.

NOTES.

NOTES.

GROUND or MOSS PINK: *Phlór subulàta*.
Polemonium Family. Apr.—June.

A low plant, forming moss-like mats, sometimes carpeting the ground. Found in rocky ground, hillsides. Southern New York to Michigan, south to Florida and Kentucky.

FLOWERS—Crimson-pink, magenta to white, 5-parted, divisions usually notched on the tip, borne in small clusters terminating the stems.

LEAVES— Small, narrow, about $\frac{1}{2}$ inch long, growing in tufts at the joints of the plant stems, which are 2–5 inches long.

DOWNY PHLOX: *Phlór pilòsa*.
Polemonium Family. May—June.

A downy-stemmed plant 1–2 feet high, found in dry, sandy soil. Ontario to Manitoba, New Jersey, Florida, Arkansas and Texas.

FLOWERS—Borne in terminal clusters, crimson-pink, purplish or white, 5-parted; somewhat sticky immediately beneath the flower.

LEAVES— Narrow or lance-shaped, toothless, stemless.

VIOLET WOOD SORREL: *Oxalis violàcea*.

Wood Sorrel Family.

May—June.

Found in rocky places, open woods. Massachusetts to Minnesota and southward.

FLOWERS—Pale magenta, sometimes white, 5 divisions, borne in clusters of 3–12 terminating a slender stalk 4–8 inches high.

LEAVES— Composed of 3 clover-like leaflets borne on slender stems springing from the root.

HAIRY BEARD-TONGUE: *Pentstèmon hirsùtus*.

Figwort Family.

May—July.

A plant 1–3 feet high, found in dry, rocky ground. Maine to Georgia, westward to Wisconsin and Missouri.

FLOWERS—Dull magenta or white, tubular, upper lip 2-lobed, lower lip 3-lobed, throat nearly closed by a hairy palate on the lower lip; flowers borne in loose spike.

LEAVES— Oblong to lance-shaped, slightly toothed, upper leaves stemless, lower leaves stemmed.

PAINTED CUP: Scarlet Painted Cup: Indian
Paint-brush. *Castilleja coccinea.*

Figwort Family. May—July.

Found in wet meadows, moist, sandy soil.

Massachusetts to Manitoba, south to Virginia,
Tennessee and Texas.

FLOWERS—Small, greenish yellow, quite eclipsed
by the 3-cleft floral leaves which are
generally tipped with brilliant scar-
let; flowers borne in clusters terminat-
ing stem 8–10 inches high.

LEAVES— Lower leaves oblong, clustered, undi-
vided; upper ones generally deeply 3–5
cleft, stemless, stained on the tip with
scarlet.

QUEEN OF THE PRAIRIE:

Filipéndula rubra.

Rose Family.

June—July.

A tall, smooth-stemmed, branching plant,
2-8 feet high, found in meadows, prairies.
Pennsylvania to Michigan, Iowa, Kentucky
and Georgia.

FLOWERS—Deep pink, fragrant, 5 divisions, borne
in large clusters.

LEAVES—Compound 7-9-parted, terminal leaflet
largest, lobed and toothed.

COMMON PASTURE, or DWARF, WILD
ROSE:

Ròsa virginiana.

Rose Family.

June—July.

A shrub 1–5 feet high, found on margins of swamps, rocky shores. Newfoundland to New York and Eastern Pennsylvania.

FLOWERS—Crimson-pink, 5 divisions, outer divisions often with 1–2 small lobes.

LEAVES—Compound, usually 7 toothed leaflets, leaf-surface smooth and often shining above. Stipules (leafy formations at the bases of the leaves) more or less dilated. Prickles hooked.

SWAMP ROSE:

R. carolina.

A straight-stemmed shrub 2–7 feet high, found along borders of swamps and streams. Nova Scotia to Florida, westward to Minnesota and Mississippi. Leaflets 5–9, finely toothed, acute at each end, usually downy beneath. Stipules long and very narrow.

SMOOTH, EARLY or MEADOW ROSE:

R. blanda.

A bush 2–4 feet high, found in rocky, moist ground. Newfoundland to New England and westward. Outer divisions of the flowers covered with bristly hairs. Leaflets 5–7, toothed. Stipules dilated, usually toothless. Prickles usually conspicuous by their absence.

PASTURE ROSE

R. hùmilis—Outer divisions of the flower always more or less lobed. Leaflets usually 7, smooth and often shining, toothed. Stipules usually narrow. Stems generally low, bearing straight, slender prickles. Nova Scotia to Florida, westward to Minnesota, Missouri and Louisiana.

SWEETBRIER; EGLANTINE:

R. rubiginòsa.

Readily known by the aromatic odor of its leaves when crushed. Introduced from Europe.

WILD BERGAMONT:

Monárda fistulòsa.

Mint Family.

July—Aug.

A plant 2–3 feet high, found in open woods, dry, rocky soil. Maine to Ontario, southward to Tennessee.

FLOWERS—Purplish, magenta, pink or whitish, tubular, 1–1½ inch long; upper lip erect, toothed, lower lip spreading, 3-lobed; clustered in a solitary, terminal head.

LEAVES— Opposite, lance-shaped, toothed, upper leaves whitish or the color of the flowers.

PURPLE CONE-FLOWER: *Brauneria purpurea*.

Composite Family.

July—Sept.

Found in rich soil. Western Pennsylvania and Virginia to Michigan, Iowa and southward.

FLOWER-HEADS—15–20 magenta colored, drooping rays, disk madder purple, terminating stem 2–3 feet high.

LEAVES— More or less toothed, lower ones 5-ribbed, long-stemmed, upper ones 3-veined, stemless.

B. pallida. Flowers similar to those of the preceding. Rays slender, drooping. Leaves lance-shaped, rough, narrowed at each end, 3-ribbed, toothless. Michigan and Illinois to Alabama and Texas.

CROSS-LEAVED MILKWORT: Marsh Milk-
wort.

Polýgala cruciàta.

Milkwort Family.

July—Sept.

A plant 3–12 inches high, found on margins of swamps, in low ground. Maine to South Carolina, and from Michigan to Minnesota and Nebraska.

FLOWERS—Magenta, white or greenish, tiny, borne in terminal, clover-like heads.

LEAVES— Generally in whorls of 4. Plant stems square, branching.

CORN COCKLE; CORN ROSE: Corn or Red
Campion: Crown of the Field.

Agrostemma Githago.

Pink Family.

July—Sept.

Found in grain fields, dry, waste places
throughout the United States. Introduced
from Europe.

FLOWERS—Magenta, 5 divisions, the narrow lobes
of the outer divisions exceeding the
inner ones in length; solitary, terminat-
ing stems 1-3 feet high.

LEAVES— Opposite, seated on stem, long, narrow,
erect. Plant stem covered with fine,
silky hairs.

SWAMP ROSE MALLOW; MALLOW ROSE.

Hibiscus Moscheutos.

Mallow Family.

July—Sept.

A shrub-like plant 4-6 feet high, found in
marshes, on lake-shores, river-banks. Massa-
chusetts and southward; usually near the coast;
locally in the interior.

FLOWERS—Pink or white, 4-7 inches across, 5 di-
visions, conspicuously veined; flowers
borne singly, or in scant clusters at the
summit of the stems.

LEAVES— Egg-shaped, pointed, toothed, densely
white-woolly beneath. Lower leaves
and sometimes the upper ones 3-lobed.

PURPLE GERARDIA: Large Purple Gerardia.

Gerardia purpurea.

Figwort Family.

Aug.—Sept.

Found in low fields, moist, sandy soil. Massachusetts to Florida and Texas near the coast, westward along the Great Lakes to Wisconsin.

FLOWERS—Purplish, magenta to whitish, cup-shaped, with 5 flaring lobes, downy.

LEAVES— Opposite, small, narrow. Plant stem somewhat angled, with spreading branches.

NOTES.

NOTES.

NOTES.

IV

BLUE AND PURPLE

LIVERLEAF; HEPATICA: LIVERWORT:

Round-lobed or Kidney Liverleaf: Noble Liverwort: Squirrel Cup. *Hepática triloba.*

Crowfoot Family. Mch.—May.

Found in open woods. Nova Scotia to Florida, Missouri and Minnesota.

FLOWERS—Variable in color, blue-violet to white, 6–12 divisions, borne on hairy stems 4–6 inches high.

LEAVES— From the root, with 3 rounded lobes, green or reddish purple, new leaves appearing after the flowers bloom.

SHARP-LOBED LIVERLEAF: *H. acutíloba.*

Differs from the preceding in that the 3 leaflets just beneath the flower, and lobes of the leaves are more acute at the tips. Leaves sometimes 5-lobed. More common westward.

Liverleaf.

BIRD-FOOT VIOLET:

Viola pedata.

Violet Family.

Apr.—June.

Found in open, sunny fields and slopes.

Massachusetts to Minnesota and southward.

FLOWERS—Largest of all the violets, blue-violet or lilac, orange center conspicuous, borne on stalk 4-10 inches high.

LEAVES— Variously divided and toothed.

NOTES.

Bird-foot Violet.

1. ROBIN'S PLANTAIN: Poor Robin's or Robert's Plantain: Blue Spring Daisy.

Erigeron pulchellus.

Composite Family.

Apr.—June.

An erect, hairy-stemmed plant 10–20 inches high, found in fields, copses, along roadsides. Maine to Ontario, Minnesota and southward.

FLOWER-HEADS—Rays lilac or pale violet, disk greenish yellow, borne in small terminal groups.

LEAVES— Lower leaves broad near the tip, slightly toothed, upper leaves lance-shaped.

2. WILD LUPINE: Old Maid's Bonnets: Wild Pea: Sun Dial.

Lupinus perennis.

Pulse Family.

May—June.

An erect, branching, leafy plant, found in dry, sandy places. New England to Minnesota, southward to the Gulf of Mexico.

FLOWERS—Violet-blue, blossoms pea-like in long clusters.

LEAVES— Compound, 7–11 long, narrow leaflets radiating from a common center.

1. Robin's Plantain.

2. Wild Lupine.

1. **BLUETS; INNOCENCE:** Quaker Ladies:
Quaker Bonnets: Venus Pride: Houstonia.

Houstonia caerùlea.

Madder Family.

Apr.—July.

A delicate little plant found in patches, almost carpeting moist, grassy places. Nova Scotia to Ontario and Wisconsin, southward to Georgia and Alabama.

FLOWERS—Pale violet, lilac or white, center yellow, about $\frac{1}{2}$ inch across, 4-lobed, on slender stems 3–6 inches high.

LEAVES— Opposite, seated on stem, lower ones broader near the tip.

H. purpurea is a southern species. Flowers deep or pale lilac, in small clusters. Leaves broad or lance-shaped, 3–5 ribbed. Plant stems smooth or hairy. Maryland to Iowa and southwest.

2. **GILL-OVER-THE-GROUND:** Ground Ivy or Joy: Creeping Charlie: Field Balm.

Népeta hederàcea.

Mint Family.

Apr.—July.

A plant with creeping stem, 6–18 inches long, found in damp, shady places. Eastern half of United States and Canada, south to Kansas and Georgia. Naturalized from Europe.

FLOWERS—Small, pale purple, spotted darker near the throat; 2-lipped, upper lip 2-lobed, lower lip 3-lobed; growing from leaf angles.

LEAVES— Rounded, heart-shaped, scallop toothed.

1. Bluets.

2. Gill-over-the-Ground.

LARGER BLUE FLAG; FLEUR-DE-LIS:

Blue Iris: Flower-de-luce. *Iris versicolor.*

Iris Family. May—July.

Found in swamps, on margins of ponds and streams. Newfoundland to Manitoba and southward.

FLOWERS—Violet-blue, the 3 outer divisions recurved, yellow at the base, veined with purple; 1 or more flowers borne on stem 2–3 feet high.

LEAVES—Erect, sword-shaped.

SLENDER BLUE FLAG: *I. prismática.*

Flowers smaller than those of the preceding, borne on slender stems. Leaves narrow, grass-like. Marshes near the coast. Nova Scotia to Georgia.

NOTES.

Larger Blue Flag.

COW VETCH; TARE: Blue or Tufted Vetch:

Cat Peas: Tinegrass.

Vicia Cracca.

Pulse Family.

June—Aug.

Found in fields, waste land. Newfoundland to New Jersey, westward to Kentucky, Iowa and Minnesota.

FLOWERS—Small, violet-blue, closely set on a 1-sided spike. Fruit a small pod.

LEAVES—Tendrill-bearing, divided into 18–24 narrow leaflets with bristle-like points.

AMERICAN VETCH: Tare or Pea Vine.

V. americana.

Flowers 3–9, slightly larger than those of the preceding. Leaflets 8–14, blunt; distinctly veined. Found in moist places. New York to Virginia and westward.

NOTES.

Cow Vetch.

BITTERSWEET; NIGHTSHADE: Blue Bindweed: Felonwort: Scarlet or Snakeberry: Poison Flower: Woody Nightshade.

Solànum Dulcamàra.

Nightshade Family.

June—Sept.

A climbing or straggling, shrubby plant, stems 2–8 feet long, found in moist thickets.

Eastern United States. Naturalized from Europe.

FLOWERS—Violet or purple, with yellow conic center, 5 reflexed divisions, in loose clusters. Berries inedible, successively green, yellow, orange, then red; appear at the same time with flowers.

LEAVES—Toothless, variable, some bearing 2 leaflets at the base.

NOTES.

Bittersweet.

1. SELF-HEAL; HEAL-ALL: Blue   Curls:
Heart-of-the-Earth: Brunella.

Prunella vulgaris.

Mint Family.

June—Sept.

Found in fields, waste places, along roadsides, throughout the United States. Naturalized from Europe.

FLOWERS—Tiny, purple, varying to white, lower lip slightly fringed, borne in a dense head, terminating stem 6–12 inches high.

LEAVES— Opposite, oblong, narrowed toward the tip, slightly toothed.

2. BLUE-EYED GRASS: Eye-bright: Blue
Star.

Sisyrinchium angustifolium.

Iris Family.

May—July.

Found in moist fields, meadows. Newfoundland to the far West, southward to Virginia.

FLOWERS—Deep violet-blue with a yellow center, the 6 divisions tipped with a point; flowers borne erect on flat stems 6–14 inches high.

LEAVES— Grass-like.

1

2

1. Self-heal.

2. Blue-eyed Grass,

1. BELLFLOWER: Rampion, European, or Creeping Bellflower. *Campánula rapunculoïdes*. Bluebell Family. July—Aug.

A sturdy plant 1-3 feet high, found along roadsides and in waste places. New Brunswick to Ontario, Pennsylvania and Ohio. Introduced from Europe and Asia. Escaped from gardens.

FLOWERS—Purple-blue, bell-shaped, 5-lobed, hanging down mostly on one side of the stem.

LEAVES— Lower leaves arrowhead-shaped, with a heart-shaped base; upper leaves broad lance-shaped, short-stemmed, toothed.

2. HAREBELL; BLUEBELL: Bluebells of Scotland: Lady's Thimble.

Campánula rotundifolia.

Bluebell Family. June—Sept.

Found on dry or moist rocky cliffs, in meadows, along margins of ponds. Arctic regions south to New Jersey and westward.

FLOWERS—Violet-blue, bell-shaped, 5-lobed, terminating exquisitely slender stalks 6-18 inches high.

LEAVES— Lower leaves nearly round, usually withered before flowering time; stem leaves narrow, seated on stem.

1

1. Bellflower.

2

2. Harebell.

NOTES.

NOTES.

---

1. PALE SPIKED LOBELIA:

Lobelia spicata.

Lobelia Family.

July—Aug.

Found in dry, sandy soil, meadows. Prince Edward Island to Ontario, westward and southwest.

FLOWERS—Pale bluish violet, upper lip of 2 narrow, erect lobes, lower lip spreading, 3-lobed; flowers borne in loose spikes, terminating stem 1-4 feet high.

LEAVES— Lower leaves oblong, obtuse, upper leaves narrow, slightly toothed.

2. MOTHERWORT:

Leonurus Cardiaea.

Mint Family.

June—Aug.

An erect, branched plant 2-4 feet high, found in waste places, generally near dwellings. Nova Scotia to North Carolina, west to Minnesota and Nebraska. Naturalized from Europe.

FLOWERS—Pale lilac or white, small, upper lip bearded, clustered in the axils of the upper leaves.

LEAVES— Opposite, long-stemmed, lower leaves variously cut and toothed, upper leaves 3-cleft.

1. Pale Spiked Lobelia.

2. Motherwort.

1. PICKEREL-WEED: *Pontederia cordata.*

Pickerel-weed Family. July—Sept.

An erect, stout, fleshy plant 1–3 feet high, found in shallow water. Nova Scotia to Ontario, Minnesota and Texas.

FLOWERS—Violet-blue, 2-lipped, upper lip 3-lobed, with 2 yellowish spots within; lower lip of 3 spreading lobes; flowers borne on thick spike.

LEAVES— One large leaf shaped like an arrow-head, is borne midway of the flower stalk.

2. MONKEY FLOWER: *Mimulus ringens.*

Figwort Family. June—Sept.

A square-stemmed plant 1–3 feet high, found beside brooks and streams, swamps. New Brunswick to Manitoba and southward.

FLOWERS—Purple, with 2 yellow spots near the throat; 2-lipped, upper lip 2-lobed, erect; lower lip 3-lobed, spreading; flowers borne on slender stalks from the leaf angles.

LEAVES— Opposite, oblong or lance-shaped, toothed, clasping stem by a heart-shaped base.

1. Pickerel-weed.

2. Monkey Flower.

1. MAD-DOG SKULLCAP: Helmet Flower:
Madweed: Hoodwort. *Scutellària lateriflòra*.

Mint Family.

July—Sept.

A plant 1–2 feet high, found in damp, shady places throughout the United States and northward.

FLOWERS—Blue or purple, varying to whitish, small, about $\frac{1}{4}$ inch long, growing in 1-sided spikes from the angles of the upper leaves.

LEAVES— Opposite, pointed, rounded at the base, coarsely toothed. 1–2 feet high.

2. SKULLCAP: Larger or Hyssop Skullcap.

Scutellària integrifòlia.

Mint Family.

May—Aug.

Found in borders of fields and woods.

Massachusetts to Florida and Texas, chiefly near the coast.

FLOWERS—Blue, about 1 inch long, opposite each other at the top of a stem 6–20 inches high.

LEAVES— Oblong or narrower, mostly toothless. Plant stem and leaves covered with a fine down.

1. Mad-dog Skullcap.

2. Skullcap.

BLUE VERVAIN: Wild Hyssop: Simpler's Joy.

Verbena hastata.

Vervain Family.

July—Sept.

A plant 3-7 feet high, found in fields, waste places, along roadsides. Common in the United States and Canada.

FLOWERS—Purple or violet, small, 5-lobed, blossoming in circles from the base of the flower spikes upward.

LEAVES— Opposite, stemmed, lance-shaped, toothed, rough; lower leaves more or less lobed.

NOTES.

Blue Vervain.

1. WILD MINT:

Méntha arvensis var. *canadensis*.

Mint Family.

July—Sept.

Found in wet places. New Brunswick to California.

FLOWERS—Purplish to white, tiny in whorls at the angles of the leaves.

LEAVES—Tapering from the center toward each end, toothed, hairy throughout, aromatic odor like pennyroyal. Plant stems branching, 10–24 inches high.

2. SPEARMINT: Garden or Mackerel Mint.

Méntha spicata.

Mint Family.

July—Sept.

Found in wet places. Eastern half of United States and Canada. Naturalized from Europe.

FLOWERS—Purplish, variable in depth of color, in narrow, acute, usually interrupted terminal spikes.

LEAVES—Aromatic, opposite, narrowly oblong, acute, sharply toothed, stemless or short-stemmed.

3. PEPPERMINT:

M. piperita.

Flowers purplish, in dense or interrupted, blunt, terminal spikes. Leaves egg-shaped, smooth, toothed, acute, very pungent tasting. Plant stem purplish, 18–36 inches high. Naturalized from Europe.

1. Wild Mint.

2. Spearmint.

3. Peppermint.

IRONWEED: Flat Top.

Vernònia noveboracénsis.

Composite Family.

July—Sept.

Found in moist soil, meadows. Massachusetts to Virginia and Mississippi; usually near the coast.

FLOWER-HEADS—Reddish purple, the brownish purple scales bristle tipped, in broad, flat clusters terminating stem 3–9 feet high.

LEAVES— Long lance-shaped, toothed.

NOTES.

Ironweed.

1. CHICORY; SUCCORY: Blue Sailors: Bunk:

Cichòrium Intybus.

Composite Family.

July—Oct.

Found in fields, waste places, along roadsides.

Newfoundland to Minnesota and southward.

Naturalized from Europe.

FLOWER-HEADS—Blue, violet-blue, lavender, rays
finely toothed on the tips; flowers
borne at intervals on branching stems
1-3 feet high.

LEAVES— Oblong or lance-shaped, coarsely
toothed, partly clasping.

2. BLAZING STAR: Button Snakeroot: Gay

Feather.

Liàtris scariòsa.

Composite Family.

Aug.—Sept.

Found in fields, along roadsides, dry soil.

Maine to Ontario, Nebraska and southward.

FLOWER-HEADS—Variable purple, borne in a
spike terminating stem 2-6 feet high.

LEAVES— Narrow lance-shaped.

1. Chicory.

2. Blazing Star.

NEW ENGLAND ASTER: Starwort.

Aster novae-angliae.

Composite Family.

Aug.—Oct.

Most striking of the asters; found in swamps, roadsides, moist ground. Maine, westward and southwest. Like the golden-rods, the number of different species of asters, with their intergrading forms, is so great as to render their identification difficult, even for the experienced botanist.

FLOWER-HEADS—Rays 40–50, light purple or violet, disk yellow.

LEAVES— Lance-shaped, toothless, clasping stem by a broad base. Plant stem stout, branched, rough.

New England Aster.

1. FRINGED GENTIAN: *Gentiàna crinìta.*

Gentian Family. Sept.—Oct.

Found in moist meadows, woods. Maine to Quebec, south to Georgia and westward.

FLOWERS—Pale or deep violet-blue, erect, tubular, with 4 spreading, fringed lobes.

LEAVES— Opposite, acute at the tip, widening to a rounded base, seated on stem. Plant stem branching, erect, each bearing a single terminal flower, 1–3 feet high.

2. CLOSED GENTIAN: Blind or Bottle Gentian. *Gentiàna Andréwsii.*

Gentian Family. Aug.—Oct.

A plant 1–2 feet high, found in woodland borders, moist soil. Maine to Manitoba and southward.

FLOWERS—Tightly closed, deep violet blue at the apex, white at the base, in terminal clusters from the leaf angles.

LEAVES— Opposite, smooth, pointed at the tip, narrowed at the base.

1. Fringed Gentian.

2. Closed Gentian.

NOTES.

NOTES.

VIRGINIAN COWSLIP: Bluebells: Tree or
Smooth Lungwort. *Merténsia virginica*.

Borage Family. Apr.—May.

A plant 1–2 feet high, found in low meadows,
along streams. New York and Ontario to
Nebraska, and southward.

FLOWERS—Pinkish in the bud, afterward violet,
trumpet-shaped, 5-lobed, about 1 inch
long, borne in loose clusters which are
rolled up at the tip and straighten as
the flowers expand.

LEAVES—Veiny, toothless, upper leaves seated on
stem, lower ones long-stemmed.

EARLY BLUE VIOLET: *Viola palmata*.
Violet Family. Apr.—May.

Found in dry ground, chiefly woodlands.

Massachusetts to Minnesota, and southward.

FLOWERS—Light or deep violet.

LEAVES—Heart-shaped, first leaves of spring
sometimes undivided, later ones 5–9-
lobed, the segments variously toothed
or cleft, middle segment usually the
largest.

ARROW-LEAVED VIOLET:

Viola sagittata.

Violet Family.

Apr.—May.

Plant 2–8 inches high; found in wet meadows, dry borders. Massachusetts to Minnesota and southward.

FLOWERS—Light or deep violet, upper and side divisions bearded, lower one veined.

LEAVES— Arrowhead-shaped, often toothless above the middle, slightly lobed below.

DOG VIOLET: Running Violet. *Viola conspersa.*

Violet Family.

Apr.—June.

Plant 2–6 inches high, found in low, shaded ground. Eastern Quebec to Minnesota and southward.

FLOWERS—Pale purple, sometimes white, raised above the leaves on stems from the leaf angles, pansy-fashion.

LEAVES— Round heart-shaped, with barely noticeable scallops. Plant stems ascending.

COMMON VIOLET: Purple, Meadow or Hooded

Blue Violet.

Viola papilionacea.

Violet Family.

Apr.—June.

Plant 3–7 inches high, found in low ground.

Massachusetts to Minnesota and southward.

FLOWERS—Light purple, variable, divisions yellowish or white at the base.

LEAVES— Heart-shaped, scallop-toothed.

WILD BLUE PHLOX:

Phlox divaricata.

Polemonium Family.

May—June.

Plant 1–2 feet high, easily recognized by its likeness to the cultivated Phloxes; found in moist, thin woods. Western Quebec to Minnesota and southward.

FLOWERS—Pale violet or lilac, in loose, spreading clusters, 5 lobes, notched at the tips.

LEAVES— Those of flowering stem opposite, oblong, tapering to a point; those of sterile shoots oblong or egg-shaped, not pointed. Plant stem finely coated with rather sticky hairs.

FORGET-ME-NOT: Mouse-ear: Snake Grass:

Scorpion Grass: Love Me: *Myosòtis scorpioides*.

Borage Family.

May—July.

Escaped from gardens; a plant with leafy, branching, half-reclining stems 6–15 inches high, found in wet ground. Newfoundland to Pennsylvania and beyond. Naturalized from Europe.

FLOWERS—Small, light blue, with a golden eye, 5-lobed, in small clusters.

LEAVES— Oblong or lance-shaped, seated on stem; leaves and stem hairy.

SMALLER FORGET-ME-NOT: *M. láxa*.

A native species, found in wet ground from Newfoundland to Ontario and southward.

FLOWERS—Light blue, on long stems, loosely clustered; lobes of the flower cup as long as the flower tube.

LEAVES— Blunt, oblong.

SPIDERWORT: Widow's or Job's Tears.

Tradescántia virginiana.

Spiderwort Family.

May—Aug.

Found in rich, moist woods, thickets. Connecticut to South Carolina; also introduced northwestward.

FLOWERS—Purplish blue, 1–2 inches broad, clustered and seated upon the end of stem between blade-like leaflets, 3 parted.

LEAVES— Long, narrow, sheathing at the base. Plant stem erect, leafy, mucilaginous, 8–36 inches high.

BEACH PEA: Seaside, Sea, or Everlasting Pea.

Láthyrus marítimus.

Pulse Family.

May—Aug.

Found chiefly near seashores from New Jersey and Oregon to the Arctic Sea; also shores of Great Lakes.

FLOWERS—Purple, about 1 inch long or less, 5–12 in short clusters on stalk springing from the leaf angles.

LEAVES— Compound, consisting of 6–10 leaflets. A pair of arrowhead-shaped leaflets are borne at the base of the compound leaves. Plant stem stout, reclining, 1–2 feet long.

COMMON SPEEDWELL: Fluellin: Paul's Betony: Ground-hele.

Verónica officinális.

Figwort Family.

May—Aug.

A low growing plant, with downy, prostrate stem 3–10 inches long, found in dry fields, open woods. Newfoundland to Ontario, Michigan, and southward.

FLOWERS—Small, light lavender in color, with darker lines, 4 unequal divisions, borne in slender spikes from the leaf angles, hld erect from the hairy, prostrate stem.

LEAVES— Oval, toothed, rounded at the tip, tapering at the base into a short stem.

AMERICAN BROOKLIME:

Verónica americana.

Figwort Family.

May—Sept.

A smooth-stemmed plant, 6–15 inches high, its stem lying partly on the ground, rooting from lower joints; found on borders of brooks, ponds, damp places. Newfoundland to Alaska and southward to Virginia.

FLOWERS—Small, quickly fading; in color lavender-blue, with slightly darker markings, 4-lobed, the lower lobe narrower than the others; borne in loose, terminal clusters.

LEAVES— Opposite, long oval or oblong lance-shaped, toothed, short-stemmed.

VIRGINIA, or COMMON, DAY-FLOWER:

Commelina virginica.

Spiderwort Family.

June—Sept.

A plant with smooth, branched stems 18–36 inches high, found in moist, shady places. New York to Florida, west to Michigan, Kansas and Texas.

FLOWERS—Violet-blue, about 1 inch broad, of 3 divisions, one of which is small and inconspicuous.

LEAVES— Lance-shaped, sheathing. A small leaf more or less encloses flower cluster.

C. hirtella is similar to the preceding, but may be distinguished by its leaves, which are brown-bearded where they sheath the flower stem.

BLUE TOADFLAX: Wild Toadflax: Blue Linaria. *Linaria canadensis.*

Figwort Family. June—Sept.

A slender plant, 5–30 inches high, found in dry, sandy places. New England westward and southwest.

FLOWERS—Pale violet or lavender, small, in slender spikes; upper lip with 2 acute divisions, lower lip larger, 3-lobed, white at the throat.

LEAVES— Narrow, toothless.

VIPER'S BUGLOSS; BLUE-WEED: Viper's Herb or Grass: Blue Thistle: Snake-flower: Blue Devil. *Echium vulgare.*

Borage Family. June—Sept.

A plant 12–30 inches high, with bristly, hairy, spotted stem, found in fields, waste places, along roadsides. Naturalized from Europe.

FLOWERS—Variable purple, pink in the bud, 5 unequal lobes, borne on a 1-sided spike, which is coiled at first, unrolling as the flowers expand.

LEAVES— Lance-shaped, hairy, mostly stemless.

TALL WILD LARKSPUR:

Delphinium exaltatum.

Crowfoot Family.

July—Aug.

A plant 2–6 feet high, found in rich woods.

Pennsylvania to Minnesota, Nebraska and southward.

FLOWERS—Light purple, irregular, borne in long, slender spikes.

LEAVES— Deeply 3–5-cleft, divisions narrowly wedge-shaped, 3-cleft at the apex.

DWARF LARKSPUR: Stagger Weed.

D. tricorne.

A plant 2–3 feet high, appearing earlier in the season; leaves deeply 5-parted, their divisions unequally 3–5-cleft; root a tuberous cluster.

PURPLE ORCHIS: Fringeless Purple Orchis.

Habenaria peramoëna.

Orchis Family.

July—Aug.

Found in moist meadows. Pennsylvania and New Jersey to Illinois, and southward.

FLOWERS—Purple, clustered in dense spike terminating stem 12–30 inches high; lower lip fan-shaped, toothed.

LEAVES— Oblong to lance-shaped.

SEA LAVENDER; MARSH ROSEMARY:

Canker-root: Ink-root. *Limonium carolinianum*.

Leadwort Family.

July—Sept.

A plant 1-2 feet high, with much-branched stems, found in salt marshes. Labrador to Texas.

FLOWERS—Lavender, very tiny, borne in sprays on upper side of branches.

LEAVES— From the root, oblong, or blunt lance-shaped, midrib prominent.

NOTES.

NOTES.

NOTES.

V

MISCELLANEOUS

SKUNK CABBAGE: Swamp Cabbage.

Symplocarpus foetidus.

Arum Family.

Mch.—Apr.

An unmistakably ill-scented plant, found in low, wet ground. Nova Scotia to North Carolina, westward to Ontario, Minnesota and Iowa.

FLOWERS—Minute, scattered over a thick, fleshy club within its flower envelope, which is variously streaked with greenish yellow and madder purple, and appears just above the ground.

LEAVES— From the base, large, often 1 foot across, strongly veined.

NOTES.

Skunk Cabbage.

WILD GINGER: Canada Snakeroot: Asarabacca.

Asarum canadense.

Birthwort Family.

Apr.—May.

Found in rich woods. North Carolina, Missouri and Kansas, northward to New Brunswick and Manitoba.

FLOWERS—Brownish purple, with 3 long-pointed divisions, borne on short stem close to the ground, often hidden under the leaves.

LEAVES— 2, broadly heart-shaped, leaf stems hairy, 6–12 inches high, from aromatic rootstock.

NOTES.

Wild Ginger.

JACK-IN-THE-PULPIT; INDIAN TURNIP:

Arisaema triphýllum.

Arum Family.

Apr.—June.

A plant 10–30 inches high, springing from an acrid bulb, found in moist, shady woods.

Nova Scotia, westward to Minnesota, and southward to the Gulf States.

FLOWERS—Minute, clustered at the base of the smooth club within the flower envelope; latter pale green or marked with purplish stripes, variable in depth of color. Berries red, clustered in a head.

LEAVES— 1–2, long-stemmed, smooth, divided into 3 large leaflets.

NOTES.

Jack-in-the-Pulpit.

1. BRANCHING BUR-REED:

Sparganium americanum.

Bur-reed Family.

June—Aug.

Plant 1–2 feet high, stem having 1–2 branches, found on borders of ponds and streams. Newfoundland to Minnesota, southward to Missouri and Florida.

FLOWERS—Of 2 kinds, the upper ones in downy heads, lower ones round, bur-like, borne on stalks from the leaf angles.

LEAVES— Long, narrow.

2. SWEET FLAG:

Acorus Calamus.

Arum Family.

June—Aug.

Found in wet, swampy places. Maine, south and west to Minnesota, Iowa, and Kansas.

FLOWERS—Tiny, compactly covering a long, slender, greenish club, which grows from the side of a two-edged stem.

LEAVES— Long, narrow, sword-like. Rootstock aromatic.

1. Branching Bur-reed

2. Sweet Flag.

1. PINESAP; FALSE BEECHDROPS: Yellow
Bird's-nest. *Monótropa Hypópitys.*

Heath Family. June—Sept.

Found in dry woods. Florida and Arizona,
northward into British Possessions.

FLOWERS—Crimson and yellow, somewhat variable,
about $\frac{1}{2}$ inch long, slightly fragrant,
borne in drooping clusters (which be-
come erect later), terminating stem
4–12 inches high.

LEAVES— None. Stems clustered, scaly, arising
from a mass of fibrous roots.

2. BEECHDROPS: Cancer-root.

Epifàgus virginiana.

Broom-rape Family. Aug.—Oct.

Found usually under beech trees. New
Brunswick to Ontario, Wisconsin, and south-
ward.

FLOWERS—Light yellow or brownish, purple-
striped, scattered along the branches.

LEAVES— None. Plant stem brownish or red-
dish tinged, arising from brittle, fibrous
roots, 6–20 inches high.

1. Pinesap.

2. Beechdrops.

1. LARGE CORAL ROOT:

Corallorrhiza maculàta.

Orchis Family.

July—Aug.

Found in dry woods. Nova Scotia, westward to British Columbia, south to Florida, Missouri, and California.

FLOWERS—Light yellow, more or less purplish-stained; lower lip white with purplish spots and lines, 3-lobed; borne on spike terminating stem 8–20 inches high.

LEAVES—None. Plant stem madder-purple or yellowish.

2. RATTLESNAKE-ROOT: Wild Lettuce: Canker-weed: Lion's Foot. *Prenánthes altíssima.*

Composite Family.

July—Sept.

Plant 3–6 feet high, found in woodlands, thickets. Newfoundland to Manitoba, southward to Georgia and Tennessee.

FLOWER-HEADS—Inner divisions inconspicuous, whitish, pendulous in loose terminal cluster, with small clusters from the leaf angles.

LEAVES—Long-stemmed, variously shaped, toothed, and cleft.

1

1. Coral Root.

2

2. Rattlesnake-root.

GROUNDNUT: Wild Bean. *Apios tuberosa*.
Pulse Family. July—Sept.

A climbing vine, 3-6 feet high, found in low, moist ground. New Brunswick to Minnesota, south to Florida, Louisiana, and Kansas.

FLOWERS—Maroon and pale brown-lilac, fragrant, borne in short, thick clusters from the leaf angles.

LEAVES—Compound, 3-7 oval leaflets.
Rootstock bears edible, tuberous enlargements.

NOTES.

Groundnut.

NOTES.

NOTES.

EARLY MEADOW RUE:

Thalictrum dioicum.

Crowfoot Family.

Apr.—May.

Plant 1–2 feet high, found in thin woods.

Maine, westward and southward.

FLOWERS—Small, tassel-like, in airy clusters, green, or purplish in effect.

LEAVES—Compound, leaflets small, lustreless, 3–7-lobed.

BLUE COHOSH: Pappoose Root.

Caulophyllum thalictroides.

Barberry Family.

Apr.—May.

Plant 1–3 feet high, found in rich woods.

New Brunswick to Manitoba and southward.

FLOWERS—Greenish yellow, more or less tinged with the bluish color of the plant; flowers about $\frac{1}{2}$ inch across, 6 divisions, borne in small clusters, which appear before the leaves are fully expanded. Seeds blue.

LEAVES—Compound, leaflets lobed. Whole plant covered with a whitish bloom when young.

COMMON ALUM ROOT:

Heuchèra americana.

Saxifrage Family.

May—Aug.

Found in dry and rocky woods. Connecticut to North Carolina, westward to Minnesota, Kansas, and Mississippi.

FLOWERS—Greenish or purplish, 5-cleft, with several orange-tipped filaments protruding; borne in a long, narrow cluster on a more or less hairy stem, 2–3 feet high.

LEAVES— Heart-shaped, with rounded lobes and teeth.

NOTES.

LARGE TWAYBLADE: *Liparis liliifolia*.
Orchis Family. June—July.

Found in woodlands. New Hampshire to
Minnesota, Missouri, and Alabama.

FLOWERS—Madder-purple or greenish, lip broad,
translucent, other divisions exceedingly
narrow; 5–15 flowers borne on stem
4–9 inches high.

LEAVES— 2, from the base, broad, shining.

NOTES.

CUT-TAIL FLAG:

Typha latifolia.

Bur-reed Family.

June—July.

Found in swamps throughout temperate North America.

FLOWERS—Borne in a dense, cylindrical head terminating stem 4–8 feet high.

LEAVES— Long, narrow, usually exceeding the flower stem in height.

NARROW-LEAVED CAT-TAIL:

T. angustifolia.

Leaves narrower than those of the preceding, and the 2 parts of the flower-head are usually separated by a short gap. Plant 4–9 feet high. Maine to North Carolina and westward, mainly near the coast.

NOTES.

NOTES.

NOTES.

NOTES.

INDEX

	PAGE
Achillea Millefolium	58
Acorus Calamus	328
Actaea alba	14
Actaea rubra	14
Adder's Tongue, White	96
Adder's Tongue, Yellow	110
Agrimonia gryposepala	138
Agrimony	138
Agrostemma Githago	265
Alder, White	40
Alisma Plantago-aquatica	82
Allium tricoccum	46
Alum Root, Common	339
Anaphalis margaritacea	86
Anemone quinquefolia	18
Anemone, Rue	18
Anemone, Fall or Summer	44
Anemone virginiana	44
Anemone, Wood	18
Anemonella thalictroides	18
Antennaria plantaginifolia	12
Anthemis Cotula	54
Apios tuberosa	334
Apocynum androsaemifolium	226
Aquilegia canadensis	194
Aralia hispida	36
Aralia nudicaulis	20
Arbutus, Trailing	94
Arethusa bulbosa	190
Arisaema triphyllum	326
Arrow-head	72
Arum, Green Arrow	24
Arum, Water	26
Asarum canadense	324

	PAGE
<i>Asclepias incarnata</i>	242
<i>Asclepias phytolaccoides</i>	102
<i>Asclepias purpurascens</i>	236
<i>Asclepias quadrifolia</i>	242
<i>Asclepias syriaca</i>	236
<i>Asclepias tuberosa</i>	183
Aster, Golden	185
Aster, New England	304
Aster novae-angliae	304
Azalea, Clammy	100
Azalea, Wild or Pink	208
Balmony	70
Balsam-apple, Wild	76
Baneberry, Red	14
Baneberry, White	14
<i>Barbarea vulgaris</i>	150
Beard-tongue, Hairy	259
Bee Balm	246
Bedstraw, Rough	72
Beechdrops	330
Beechdrops, False	330
Bellflower	286
Bellwort	122
Bellwort, Large-flowered	122
Bergamot, Wild	263
Betony, Wood	216
<i>Bidens laevis</i>	176
Bindweed, Field	238
Bindweed, Hedge	238
Bird's Nest	88
Birthroot	192
Bishop's Cap	10
Bitter Bloom	240
Bittersweet	282
Black-eyed Susan	142
Blazing Star	302
Bloodroot	2
Bluebell	286
Bluets	276
Blue-weed	316
Boneset	68
Bouncing Bet	216
<i>Brassica nigra</i>	150

INDEX

347

	PAGE
Brauneria pallida	264
Brauneria purpurea	264
Brooklime, American	315
Bugloss, Viper's	316
Bunchberry	54
Bur-marigold, Larger	176
Bur-reed, Branching	328
Butter-and-Eggs	160
Butterfly Weed	183
Buttonbush	90
Calico Bush	212
Calla, Marsh	26
Calla palustris	26
Calopogon pulchellus	222
Caltha palustris	112
Campanula rapunculoides	286
Campanula rotundifolia	286
Campion, Bladder	50
Campion, White	50
Cancer-root	330
Canker-root	30
Cardinal Flower	244
Carrion-flower	130
Carrot, Wild	88
Cassia Chamaecrista	166
Cassia marilandica	166
Castalia odorata	103
Castilleja coccinea	260
Caulophyllum thalictroides	338
Ceanothus americanus	100
Celandine	138
Cephalanthus occidentalis	90
Cerastium arvense	44
Chamomile	54
Checkerberry	104
Cheeses	52
Chelidonium majus	138
Chelone glabra	70
Chickweed, Field	44
Chicory	302
Chimaphila maculata	234
Chimaphila umbellata	234
Chiogenes hispidula	99

	PAGE
Chrysopsis mariana	185
Cichorium Intybus	302
Cicuta maculata	102
Cimicifuga racemosa	100
Cinquefoil	114
Claytonia caroliniana	188
Claytonia virginica	188
Clematis virginiana	60
Clethra alnifolia	40
Clintonia borealis	118
Clintonia, Yellow	118
Clover, Sweet	86-181
Clover, Rabbit's-foot	252
Clover, Yellow or Hop	148
Cohosh, Black	100
Cohosh, Blue	338
Collinsonia canadensis	184
Columbine, Wild	194
Commelina virginica	315
Cone-flower	142
Cone-flower, Purple	142-264
Cone-flower, Tall	142
Convolvulus arvensis	238
Convolvulus sepium	238
Coptis trifolia	30
Coral Root, Large	332
Corallorrhiza maculata	332
Corn Cockle	265
Cornel, Dwarf	54
Corn Rose	265
Cornus canadensis	54
Corpse plant	46
Corydalis aurea	179
Corydalis, Golden	179
Corydalis, Pale	218
Corydalis sempervirens	218
Cowbane, Spotted	102
Cowslip, American	190
Cowslip, Virginian	310
Cranesbill, Wild	198
Cress, Water	99
Cress, Winter	150
Crinkle-root	6
Cucumber-root, Indian	120

INDEX

349

	PAGE
Culver's Root	105
Cuscuta Gronovii	84
Cypripedium acaule	204
Cypripedium hirsutum	101
Cypripedium parviflorum	126
Cypripedium parviflorum var. pubescens	126
Dalibarda repens	103
Dandelion, Fall	166
Datura Stramonium	105
Daucus Carota	88
Day-flower, Virginia	315
Deer-grass	246
Delphinium exaltatum	317
Delphinium tricornes	317
Dentaria diphylla	6
Dentaria laciniata	6
Devil's Paint-brush	172
Dianthus Armeria	196
Dianthus deltoides	196
Dicentra canadensis	4
Dicentra Cucullaria	4
Dodder, Common	84
Dodecatheon Meadia	190
Dogbane, Spreading	226
Drosera longifolia	66
Drosera rotundifolia	66
Dutchman's Breeches	4
Echinocystis lobata	76
Echium vulgare	316
Eglantine	263
Elder, Black-berried	40
Elder, Common	40
Elder, Red-berried	40
Elecampane	164
Epifagus virginiana	330
Epigaea repens	94
Epilobium angustifolium	244
Epipactis pubescens	62
Erigeron annuus	56
Erigeron pulchellus	274
Erigeron ramosus	56
Erythronium albidum	96

	PAGE
<i>Erythronium americanum</i>	110
<i>Eupatorium perfoliatum</i>	68
<i>Eupatorium purpureum</i>	250
<i>Eupatorium urticaefolium</i>	68
Everlasting, Early Spring	12
Everlasting, Pearly	86
Everlasting, Plantain-leaved	12
<i>Filipendula rubra</i>	261
Fireweed	244
Flag, Cat-tail	341
Flag, Larger Blue	278
Flag, Narrow-leaved Cat-tail	341
Flag, Slender Blue	278
Flag, Sweet	328
Fleabane, Daisy	56
Fleur-de-lis	278
Flower-de-luce	278
Foam-flower	10
Forget-me-not	313
Forget-me-not, Smaller	313
Foxglove, Downy False	162
Foxglove, Fern-leaved False	185
Foxglove, Smooth False	162
Frostweed	182
<i>Galium asprellum</i>	72
<i>Gaultheria procumbens</i>	104
Gentian, Closed	306
Gentian, Fringed	306
<i>Gentiana Andrewsii</i>	306
<i>Gentiana crinita</i>	306
<i>Geranium maculatum</i>	198
<i>Geranium Robertianum</i>	198
<i>Geranium</i> , Wild	198
<i>Gerardia flava</i>	162
<i>Gerardia pedicularia</i>	185
<i>Gerardia purpurea</i>	266
<i>Gerardia</i> , Purple	266
<i>Gerardia virginica</i>	162
Gill-over-the-Ground	276
Ginger, Wild	324
Ginseng	183
Ginseng, Dwarf	20

INDEX

351

	PAGE
Globe-flower	90
Golden Alexanders	116
Golden Club	180
Golden-rod	185
Goldthread	30
Grass, Blue-eyed	284
Grass of Parnassus	80
Groundnut	334
 Habenaria blephariglottis	104
Habenaria bracteata	42
Habenaria ciliaris	162
Habenaria fimbriata	234
Habenaria Hookeri	64
Habenaria hyperborea	42
Habenaria lacera	62
Habenaria leucophaea	104
Habenaria orbiculata	64
Habenaria peramoena	317
Habenaria psycodes	234
Hardhack	228
Harebell	286
Hawkweed, Canada	172
Hawkweed, Hairy	174
Hawkweed, Orange or Tawny	172
Hawkweed, Rough	174
Heal-all	284
Helenium autumnale	184
Helianthemum canadense	182
Helianthus decapetalus	176
Hellebore, American White	181
Hemlock, Water	102
Hepatica acutiloba	270
Hepatica triloba	270
Herb Robert	198
Heuchera americana	339
Hibiscus Moscheutos	265
Hieracium aurantiacum	172
Hieracium canadense	172
Hieracium Gronovii	174
Hieracium scabrum	174
Hieracium venosum	174
Honeysuckle	194
Honeysuckle, White Swamp	100

	PAGE
Horse Balm	184
Houstonia caerulea	276
Houstonia purpurea	276
Hypericum perforatum	158
Hypoxis hirsuta	114
Impatiens biflora	156
Impatiens pallida	156
Indian Hemp	226
Indian Pipe	46
Indian Poke	181
Indian Turnip	326
Innocence	276
Inula Helenium	164
Iris prismatica	278
Iris versicolor	278
Ironweed	300
Ivy, Poison	34
Jack-in-the-Pulpit	326
Jamestown or Jimson Weed	105
Jeffersonia diphylla	96
Jewel-weed	156
Joe-Pye Weed	250
Kalmia angustifolia	220
Kalmia latifolia	212
Kalmia polifolia	220
Knotweed, Pink	248
Lactuca canadensis	146
Ladies' Tresses	66
Ladies' Tresses, Slender	66
Lady's Slipper, Large Yellow	126
Lady's Slipper, Showy	101
Lady's Slipper, Small Yellow	126
Lady's Slipper, Stemless or Pink	204
Lady's Thumb	248
Lambkill	220
Larkspur, Dwarf	317
Larkspur, Tall Wild	317
Lathyrus maritimus	314
Laurel, Mountain	212
Laurel, Pale	226

	PAGE
Laurel, Sheep	220
Lavender, Sea	318
Leek, Wild	46
Leontodon autumnalis	166
Leonurus Cardiaca	290
Lettuce, Wild	146
Liatris scariosa	302
Lilium canadense	134
Lilium philadelphicum	224
Lilium superbum	134
Lily, Cow	132
Lily, Meadow or Field	134
Lily, Turk's Cap	134
Lily, White Water	103
Lily, Wild Orange-red	224
Lily, Wild Yellow	134
Lily, Wood	224
Lily, Yellow Pond	132
Limonium carolinianum	318
Linaria canadensis	316
Linaria vulgaris	160
Linnaea borealis	226
Liparis liliifolia	340
Live-for-ever	254
Liverleaf	270
Liverleaf, Sharp-lobed	270
Liverwort	270
Lizard's Tail	101
Lobelia cardinalis	244
Lobelia, Pale Spiked	290
Lobelia spicata	290
Loosestrife	140
Loosestrife, Four-leaved	140
Loosestrife, Fringed	136
Lousewort	216
Lupine, Wild	274
Lupinus perennis	274
Lychnis alba	50
Lychnis, Evening	50
Lysimachia quadrifolia	140
Lysimachia terrestris	140
Maianthemum canadense	26
Mallow, Common	52

	PAGE
Mallow, Musk	52
Mallow Rose	265
Mallow, Swamp Rose	265
Malva moschata	52
Malva rotundifolia	52
Mandrake	97
Marigold, Marsh	112
May Apple	97
Mayflower	94
Meadow Beauty	246
Meadow Beauty, Maryland	246
Meadow Rue, Early	338
Meadow Rue, Tall	78
Meadow-sweet	228
Medeola virginiana	120
Melilot, White	86
Melilot, Yellow	181
Melilotus alba	86
Melilotus officinalis	181
Mentha arvensis var. canadensis	298
Mentha piperita	298
Mentha spicata	298
Mercury	34
Mertensia virginica	310
Milfoil	58
Milkweed, Common	236
Milkweed, Four-leaved	242
Milkweed, Poke or Tall	102
Milkweed, Purple	236
Milkweed, Swamp	242
Milkwort, Cross-leaved	264
Milkwort, Field	210
Milkwort, Fringed	210
Mimulus ringens	292
Mint, Wild	298
Mitchella repens	6
Mitella diphylla	10
Mitella nuda	10
Mitrewort	10
Mitrewort, False	10
Mitrewort, Naked	10
Moccasin Flower	204
Monarda didyma	246
Monarda fistulosa	263

INDEX

355

	PAGE
Moneses uniflora	38
Monkey Flower	292
Monotropa Hypopitys	330
Monotropa uniflora	46
Morning-glory, Wild	238
Moss, Flowering	94
Motherwort	290
Mullein, Common	144
Mullein, Moth	144
Mustard, Black	150
Myosotis laxa	313
Myosotis scorpioides	313
Nepeta hederacea	276
New Jersey Tea	100
Nightshade	282
Nymphaea advena	132
Oakesia sessilifolia	122
Oenothera biennis	154
Oenothera fruticosa	154
Oenothera pumila	154
Orchis, Greater Green	64
Orchis, Hooker's	64
Orchis, Large Round-leaved	64
Orchis, Large Purple Fringed	234
Orchis, Prairie White Fringed	104
Orchis, Purple	317
Orchis, Ragged Fringed	62
Orchis, Rein	42
Orchis, Showy	206
Orchis, Smaller Purple Fringed	234
Orchis spectabilis	206
Orchis, White Fringed	104
Orchis, Yellow Fringed	162
Orontium aquaticum	180
Orpine, Garden	254
Oswego Tea	246
Oxalis Acetosella	84
Oxalis corniculata	148
Oxalis violacea	259
Painted Cup	260
Panax quinquefolium	183

	PAGE
<i>Panax trifolium</i>	20
<i>Parnassia caroliniana</i>	80
Parsnip, Early Meadow	116
Parsnip, Water	102
Parsnip, Wild or Field	116
Partridge Berry	6
<i>Pastinaca sativa</i>	116
Pea, Beach	314
Pea, Partridge	166
<i>Pedicularis canadensis</i>	216
<i>Peltandra virginica</i>	24
<i>Pentstemon hirsutus</i>	259
Pepperbush, Sweet	40
Peppermint	298
<i>Persicaria, Water</i>	248
<i>Phlox divaricata</i>	312
Phlox, Downy	258
<i>Phlox pilosa</i>	258
<i>Phlox subulata</i>	258
Phlox, Wild Blue	312
<i>Phytolacca decandra</i>	92
Pickerel-weed	292
Pigeon Berry	92
Pinesap	330
Pink, Deptford	196
Pink, Grass	222
Pink, Ground or Moss	258
Pink, Indian	190
Pink, Large Marsh	240
Pink, Maiden	196
Pink, Sea	240
Pink, Slender Marsh	240
Pink, Wild	196
Pinxter Flower	208
Pipsissewa	234
Pitcher-plant	202
Plantain, Robin's	274
Plantain, Water	82
Pleurisy Root	183
<i>Podophyllum peltatum</i>	97
<i>Pogonia ophioglossoides</i>	222
Pogona, Rose or Sweet	222
Pokeweed	92
<i>Polygala cruciata</i>	264

INDEX

357

	PAGE
Polygala, Field	210
Polygala, Fringed	210
Polygala paucifolia	210
Polygala sanguinea	210
Polygonatum biflorum	16
Polygonatum commutatum	16
Polygonum amphibium	248
Polygonum pennsylvanicum	248
Polygonum Persicaria	248
Pontederia cordata	292
Potentilla canadensis	114
Prenanthes altissima	332
Primrose, Common Evening	154
Prince's Pine	234
Prunella vulgaris	284
Pussy's Toes	12
Pyrola americana	38
Pyrola elliptica	38
Pyrola, One-flowered	38
Pyrola, Round-leaved	38
Pyxidanthera barbulata	94
Pyxie	94
Queen Anne's Lace	88
Queen of the Prairie	261
Radicula Nasturtium-aquaticum	99
Ragwort, Golden	128
Raspberry, Purple Flowering	230
Rattlesnake Plantain, Downy	62
Rattlesnake-root	332
Rattlesnake-weed	174
Red-root	100
Rheumatism Root	96
Rhexia mariana	246
Rhexia virginica	246
Rhododendron nudiflorum	208
Rhododendron viscosum	100
Rhus Toxicodendron	34
Rhus Vernix	34
Rich-weed	184
Rocket, Yellow	150
Rockrose	182
Rosa blanda	262

	PAGE
<i>Rosa carolina</i>	262
<i>Rosa humilis</i>	263
<i>Rosa rubiginosa</i>	263
<i>Rosa virginiana</i>	262
Rose, Pasture or Dwarf	262
Rose, Smooth	262
Rose, Swamp	262
Rosemary, Marsh	318
<i>Rubus odoratus</i>	230
<i>Rudbeckia hirta</i>	142
<i>Rudbeckia laciniata</i>	142
<i>Sabatia angularis</i>	240
<i>Sabatia dodecandra</i>	240
<i>Sabatia gracilis</i>	240
<i>Sabatia stellaris</i>	240
<i>Sagittaria latifolia</i>	72
<i>Sambucus canadensis</i>	40
<i>Sambucus racemosa</i>	40
<i>Sanguinaria canadensis</i>	2
<i>Saponaria officinalis</i>	216
<i>Sarracenia purpurea</i>	202
Sarsaparilla, Bristly	36
Sarsaparilla, False or Wild	20
<i>Saururus cernuus</i>	101
<i>Saxifraga virginensis</i>	12
Saxifrage, Early	12
Scabious, Sweet	56
<i>Scutellaria integrifolia</i>	294
<i>Scutellaria lateriflora</i>	294
<i>Sedum purpureum</i>	254
Self-heal	284
<i>Senecio aureus</i>	128
Senna, Wild	166
Shin Leaf	38
Shooting Star	190
<i>Silene latifolia</i>	50
<i>Silene pennsylvanica</i>	196
<i>Sisyrinchium angustifolium</i>	284
<i>Sium cicutae-folium</i>	102
Skullcap	294
Skullcap, Mad-dog	294
Skunk Cabbage	322
Smartweed	248

INDEX

359

	PAGE
<i>Smilacina racemosa</i>	32
<i>Smilacina stellata</i>	32
<i>Smilax herbacea</i>	130
Snake Mouth	222
Snakeroot, Black	100
Snakeroot, White	68
Sneezeweed	184
Snowberry, Creeping	99
Soapwort	216
<i>Solanum Dulcamara</i>	282
Solomon's Seal	16
Solomon's Seal, False	32
Solomon's Seal, Great	16
Solomon's Seal, Star-flowered	32
Sorrel, Lady's	148
Sorrel, Violet Wood	259
Sorrel, Wood	84
<i>Sparganium americanum</i>	328
Spatter-dock	132
Spearmint	298
Speedwell, Common	314
Spiderwort	313
Spikenard, False	32
<i>Spiraea latifolia</i>	228
<i>Spiraea tomentosa</i>	228
<i>Spiranthes cernua</i>	66
<i>Spiranthes gracilis</i>	66
Spring Beauty	188
Squirrel Corn	4
Star Flower	30
Star Grass	114
Steeple Bush	228
<i>Steironema ciliatum</i>	136
St. John's Wort, Common	158
<i>Streptopus amplexifolius</i>	214
<i>Streptopus roseus</i>	214
Succory	302
Sumach, Poison	34
Sundew, Long-leaved	66
Sundew, Round-leaved	66
Sundrops	154
Sunflower, Swamp	184
Sunflower, Thin-leaved	176
Sweetbrier	263

	PAGE
<i>Symplocarpus foetidus</i>	322
<i>Tanacetum vulgare</i>	170
Tansy	170
<i>Thalictrum dioicum</i>	338
<i>Thalictrum polygamum</i>	78
Thimbleweed	44
Thoroughwort	68
<i>Tiarella cordifolia</i>	10
Toadflax, Blue	316
Toadflax, Yellow	160
Toothwort	6
Touch-me-not, Pale	156
Touch-me-not, Spotted	156
<i>Tradescantia virginiana</i>	313
<i>Trientalis americana</i>	30
<i>Trifolium agrarium</i>	148
<i>Trifolium arvense</i>	252
<i>Trillium cernuum</i>	28
<i>Trillium erectum</i>	192
<i>Trillium grandiflorum</i>	28
Trillium, Large-flowered	28
Trillium, Nodding	28
Trillium, Painted	28
Trillium, Purple	192
<i>Trillium recurvatum</i>	192
<i>Trillium sessile</i>	192
Trillium, Stemless	192
<i>Trillium undulatum</i>	28
Turtlehead	70
Twayblade, Large	340
Twin-flower	226
Twinleaf	96
Twisted-stalk	214
Twisted-stalk, Clasping-leaved	214
<i>Typha angustifolia</i>	341
<i>Typha latifolia</i>	341
<i>Uvularia grandiflora</i>	122
<i>Uvularia perfoliata</i>	122
<i>Veratrum viride</i>	181
<i>Verbascum Blattaria</i>	144
<i>Verbascum Thapsus</i>	144

INDEX

361

	PAGE
Verbena hastata	296
Vernonia noveboracensis	300
Veronica americana	315
Veronica officinalis	314
Veronica virginica	105
Vervain, Blue	296
Vetch, American	280
Vetch, Cow	280
Vicia americana	280
Vicia Cracca	280
Viola canadensis	98
Viola conspersa	311
Viola lanceolata	98
Viola pallens	97
Viola palmata	310
Viola papilionacea	312
Viola pedata	272
Viola pubescens	180
Viola rotundifolia	179
Viola sagittata	311
Violet, Arrow-leaved	311
Violet, Bird-foot	272
Violet, Canada	98
Violet, Common	312
Violet, Dog	311
Violet, Dog's-tooth	110
Violet, Downy Yellow	180
Violet, Early Blue	310
Violet, Early Yellow	179
Violet, Lance-leaved	98
Violet, Round-leaved	179
Violet, Sweet White	97
Virgin's Bower	60
Wake Robin	98-192
Widow's or Job's Tears	313
Willow-herb, Great	244
Wind-flower	18
Wintergreen	104
Wintergreen, Flowering	210
Wintergreen, Spotted	234
Yarrow	58
Zizia aurea	116

THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW

TEL. NO. 642-2532

Renewed books are subject to immediate recall.

[illegible]

Who's who among the
Wild Flowers.

BIOLOGY
LIBRARY
G

JUN 30 1917 L.S. 6 JUL 1 1917

JUL 1 1917 cat. 6 JUL 11 1917

JUL 19 1919 Ehlers

APR 25 1922 Miller APR 27 1928

AUG 1 5 1946 Miller AUG 5 1946

RARIES

88

Beecroft

201899

QK118

BIOLOGY
LIBRARY
G

UN

LIBRARY