Zool. dept.

EX LIBRIS

Zool. dept.

BIOLOGY
LIBRARY
G

Q L 53
C 35

BIOLOGY
LIBRARY
G

Cambridge Eng. University
Laboratory guide in zoology

254175

PHYLUM PROTOZOA. *Animals in which there is one nucleus, or, if more than one nucleus, in which the nuclei are disposed apparently irregularly and are not differentiated in relation to the functional tissues of the animal.*

Class 1. **GYMNOMYXA.** Naked forms without distinct cortical layer and capable of emitting pseudopodia.

Order 1. **LOBOSA.** Simple forms with blunt pseudopodia.

Gymnamoebida. Without a shell. Ex. *Amoeba*.

Thecamoebida. With a shell. Ex. *Arcella*, *Diffugia*.

Order 2. **FORAMINIFERA.** Gymnomyxa with thread-like pseudopodia forming networks, and a shell the outside of which is covered by protoplasm. Ex. *Polystomella*, *Globigerina*.

Order 3. **RADIOLARIA.** Marine Gymnomyxa with radiating pseudopodia, central capsule and an additional skeleton of flinty needles. Ex. *Collozoum*, *Thalassicola*.

Order 4. **HELIOZOA.** Gymnomyxa with stiff radiating pseudopodia. Skeleton when present only in the form of isolated needles. Ex. *Actinosphaerium*, *Actinophrys*.

Order 5. **MYCETOZOA.** Gymnomyxa of large size with branching pseudopodia. Reproduction by means of cellulose-coated spores. The contents of many spores coalesce to form one individual. Ex. *Badhamia*, *Fuligo*.

Class 2. **CORTICATA.** Protozoa with a distinct cuticle and almost always a distinct cortical layer.

Order 1. **CILIATA.** Forms provided with cilia.

Holotricha. Ex. *Paramecium*.

Heterotricha. Ex. *Stentor*.

Hypotricha. Ex. *Stylonychia*, *Kerona*.

Peritricha. Ex. *Vorticella*.

Order 2. **ACINETARIA.** Forms provided with sucking tentacles. Ex. *Acineta*.

Order 3. **FLAGELLATA.** Forms provided with flagella.

Lissoflagellata. Ex. *Polytoma*.

Choanoflagellata. Ex. *Monosiga*.

Phytoflagellata. Ex. *Volvox*.

Dinoflagellata. Ex. *Ceratium*.

Cystoflagellata. Ex. *Noctiluca*.

Order 4. **SPOROZOA.** Parasitic forms, devoid of mouth, cilia, flagella or tentacles. The younger stages, at least, are cell parasites.

Gregarinida. Ex. *Monocystis*.

Coccidiida. Ex. *Coccidium*.

Haemosporidia. Ex. Malarial Parasite.

PROTOZOA.**GYMNOMYXA.****LOBOSA.**

1. **Amœba.** Examine alive, then irrigate with $\frac{1}{4}$ p.c. solution of osmic acid. Wash with water and stain by irrigating with picrocarmine. When the nucleus becomes bright red wash again with water. Then run in dilute glycerine and finally strong glycerine, sealing with gold size.

2. **Arcella.** *sh.* shell; *a.* aperture of shell; *n, n.* nuclei; *c. v.* contractile vacuole; *ps.* pseudopodia; *f. b.* food particles; *g. v.* gas vacuoles.

FORAMINIFERA.

3. **Polystomella.** Examine (*a*) alive in watch-glass, to see the external protoplasm and the pseudopodia, (*b*) transferred to slide using the high power of the microscope. Then kill with osmic acid, decalcify in 5 p.c. acetic acid, stain with picrocarmine and mount permanently, to see nuclei and cell-chambers.

4. Shells of Foraminifera. Calcareous.

HELIOZOA.

5. **Actinosphærium.** (a) Floating in water in watch-glass. Low power. (b) Mount on slide, supporting coverslip by two narrow strips of gummed paper. Examine with high power. (c) Stain as in case of *Amoeba*, to show nuclei. *ex.* ectoplasm; *en.* endoplasm; *ps.* pseudopodia; *c.v.* contractile vacuole; *ax.* axes of pseudopodia (*ps.*); *gr.* granular cortex of pseudopodia; *v.ex.* vacuoles of ectoplasm; *f.v.* food balls in endoplasm.

PROTOZOA

6. **Actinophrys.** Stained to show single nucleus with osmic and picocarmine. Draw with high power. *ex.* ectoplasm; *en.* endoplasm; *n.* nucleus; *ps.* pseudopodia; *c.v.* contractile vacuole.

7. Shells of Radiolaria. Mostly siliceous.

MYCETOZOA.

8. Various Mycetozoa to show (*a*) spore-fruit, (*b*) burst spore-fruit showing capillitium, (*c*) creeping plasmodium. Note rhythmic character of flow of protoplasm. (*d*) plasmodium killed and stained to show nuclei.

CORTICATA.

CILIATA.

1. (a) View of a **Paramœcium** mounted in gum to restrict movement. The animal is easily visible to the naked eye, if black paper be used as a background. (b) Various phases in the disappearance and reforming of the contractile vacuole. (c) **Paramœcium** stained by irrigation with methylene blue. (d) Fed with indian ink. Watch the rotation of the food vacuoles, the entrance of the ink masses into the endoplasm, and their extrusion from the temporary anus.

Long adoral cilia, cilia, ectoplasm, endoplasm; food vacuoles; outer part of ectoplasm with trichocysts; contractile vacuoles (two) in diastole; same in systole; vestibule; œsophagus; position of anus; macronucleus; micronucleus. (Chlorophyll grains are present in some species.)

2. **Opalina ranarum.** A mouthless holotrichous form.
3. **Nyctotherus cordiformis.** A heterotrichous form. pharynx, contractile vacuole, nucleus, anus (permanent).
4. **Balantidium entozoon.** A heterotrichous form. Four contractile vacuoles, mouth, nucleus.
(a) Examine alive in a drop of the rectal fluid. (b) Stain as in Amœba. (c) Irrigate fresh drop of fluid with methylene blue.

5. **Stentor.** A heterotrichous free-living form.

FLAGELLATA.

1. **Polytoma uvella.** Smear a drop of the fluid over the slide and examine the drying edges, where the animals will be more or less still. *f.* flagella; *c.v.* contractile vacuoles; *n.* nucleus.

Look for forms undergoing fission.

Add iodine. The flagella are seen more distinctly.

2. **Euglena viridis.** Gullet, pigment spot, flagellum, contractile vacuole, nucleus.

Kill with iodine.

PROTOZOA

3. **Volvox.** (*a*) Alive and rotating in the water. (*b*) Removed to slide and slightly compressed by coverslip, supported on strips of paper. (*c*) Killed with iodine.

Individuals of colony; connecting threads; flagella; parthenogonidia; female reproductive cell (macrogonidium); male reproductive cell dividing into microgonidia.

SPOROZOA.

GREGARINIDA.

1. Free **Monocystis** from the sperm rosettes in the vesicula seminalis of the Earthworm. High power. *cor.* cortical substance; *med.* medullary granular substance; *nu.* nucleus; *n, m.* successive changes of form of the Gregarine.

N.B. Open the Earthworm from the dorsal side in salt solution. Remove alimentary canal carefully and open median vesicula seminales. Note white threads streaming from the sperm rosettes which are the Monocystis. Cut off small part of rosette with attached parasite and examine in .75 p.c. salt solution.

2. Cysts from amongst developing spermatozoa in the lateral vesiculæ seminales. (*a*) With two unbroken individuals; (*b*) the encysted individuals fused and more or less broken up; (*c*) with young lemon-shaped pseudonavicellæ; (*d*) with complete pseudonavicellæ; (*e*) enlarged drawing of a single pseudonavicella seen under the highest available power. *es.* spore-case.

PROTOZOA

3. **Clepsidrina blattarum** from the chylic ventricle of the cockroach; anterior chamber (protomerite); posterior chamber (deuteromerite); anterior knob for attachment (epimerite) generally absent; nucleus; nucleolus; cortical substance; medullary substance.

PHYLUM PORIFERA. *Organisms consisting of a dermal layer of flattened cells and a gastral layer of flagellate collar cells (choanocytes), separated by a gelatinous mesoglaea containing generative and skeleton-secreting cells. Numerous canals communicate between the gastral cavity and the exterior.*

Class 1. **CALCAREA.** With calcareous spicules. Ex. *Leucosolenia*, *Grantia*.

Class 2. **HEXACTINELLIDA.** With siliceous spicules of triaxonic (sex-radiate) type. Ex. *Euplectella*.

Class 3. **DEMOSPONGIÆ.** Skeleton composed either of siliceous spicules of various types (never triaxonic) or of spongin, pure or combined with spicules or foreign bodies. Ex. *Oscarella*, *Spongilla* (freshwater), *Euspongia* (bath sponge).

PORIFERA.

1. **Leucosolenia.** Simple calcareous sponge (Ascon type). Osculum. The surface is rough with projecting calcareous spicules.
2. **Grantia.** Calcareous sponge (Sycon type): colonial form. Narrow elongated oscula.

3. Transverse section through **Leucosolenia.** Gastral cavity with collar-cells; radial canals; mesogloea with ova; spicules. Draw outline from low power, fill in details from high power of decalcified specimen.

PORIFERA

4. Transverse section through **Grantia**. Gastral cavity; flagellated chambers (with collar cells); inhalent pores (prosopyles); apopyles; inhalent canals.

5. Examples of Hexactinellida and Demospongiæ.

Class 4. **CTENOPHORA.** Coelenterates with a secondarily acquired bilateral symmetry. Eight meridional rows of vibratile plates formed of fused cilia. With oesophagus. Ex. *Pleurobrachia*, *Beroë*, *Cestus*.

HYDROZOA.

HYDROMEDUSÆ.

1. Colony of **Obelia** (as type of Calyptoblastic Hydroid). Cœnosarc; perisarc; hydrotheca; gonotheca; blastostyle; medusa-bud. The material of **Obelia** and other hydroids is given to the student preserved in 70 p.c. alcohol. It should be stained in borax carmine (in 50 p.c. alcohol), dehydrated and mounted in Canada balsam.

2. Medusa of **Obelia** (type of Leptomedusæ). Tentacles of edge of umbrella; manubrium; the four gonads; the four radial canals; the eight otocysts.

1880

1880

HYDROZOA

3. Colonies of other Calyptoblastea. (a) **Sertularia**, sessile polyps on both sides of branch.
(b) **Antennularia**, sessile polyps on one side of branch; nematocalyces.

4. **Pennaria**. A colony of hydroids of the Gymnoblastea.
5. **Lizzia**. (Type of the Anthomedusæ). Manubrial gonad; groups of marginal tentacles; ocelli.

HYDROZOA

6. A single polyp of **Tubularia** (Gymnoblasic Hydroid) removed from the colony and examined with a simple lens.

Circumoral tentacles; gonophores, on branched processes of the body wall; large tentacles; small globular swelling at the base of the body passing into cœnosarc.

7. Longitudinal section through a polyp of **Tubularia**. (Low and high powers.) Mouth; ectoderm; endoderm of enteron; *en. tent.* solid axis of the tentacle, consisting of modified, much vacuolated endoderm cells; nematocysts; circumoral tentacles; gonophores and stalk of gonophores; oral tentacles; globular swelling; cœnosarc.

HYDROZOA

8. Transverse section of a **Tubularian polyp** near the mouth. (Low power.) Axes of oral tentacles prolonged for a short distance down the body; ectoderm; endoderm.

9. Bunch of female gonophores of **Tubularia** stained slightly. In ripe gonophores observe umbrella cavity; rudimentary tentacles; manubrium with gonad; embryo in sub-umbrellar cavity. Look out for stages showing development of gonophores.

HYDROZOA

10. **Geryonia**. One of the Trachymedusæ, no hydroid phase. Long manubrium; six radial canals; six tentacles; velum.

11. View of part of margin of disc of **Geryonia** showing tentaculocyst.

12. Transverse section through the edge of the disc of **Geryonia**. (High power.) Thickened ectodermal ring; velum; tentaculocyst; *n.* 1 upper, *n.* 2 lower nerve-rings; circular canal; sub-umbrellar ectoderm; gelatinous tissue.

SIPHONOPHORA.

- (a) **Physophora hydrostatica**. Pneumatophore; nectocalyces; dactylozooids; nutritive polyps (hardly distinguishable) in the midst of *dact.*; tentacles with lateral branches terminated by batteries of thread cells.
- (b) **Physalia** with large pneumatophore and long tentacles.
- (c) **Velella** with diagonal crest on the pneumatophore.

ACALEPHÆ.

1. **Aurelia aurita.** Small marginal tentacles; 8 sense tentacles; 4 sub-genital pits and 4 groups of gastral filaments in the radii of four of the sense tentacles; 4 oral arms in the radii of the other four sense tentacles; the pseudo-velum; 16 radiating canals, those in the radii (radial) of the 8 sense tentacles branch, the others (adradial) do not; circular canal.

2. Examine a sense tentacle first in situ, and then in watch glass with low power to see protective folds of umbrella; otoliths.

ACALEPHÆ

3. **Rhizostoma.** Branching mouth-bearing processes from manubrium; sense tentacles. Absence of velum and tentacles.
4. **Pelagia noctiluca.** Note the 16 gastral pouches. Gonads; marginal notches and sense tentacles; 8 long marginal tentacles; sub-genital pits.
5. A **Hydratuba** or **Scyphistoma.**

ACTINOZOA.

ALCYONARIA (OCTACTINIA).

1. **Alcyonium palmatum.** Naked eye and hand lens. Coenenchyma; polyps extended; pinnate tentacles of polyps.

2. Transverse section through a polyp of **Alcyonium**, at the œsophageal region. Body-wall; œsophageal wall; ciliated groove of latter; mesenteries; mesogloea of œsophagus with spicules; longitudinal muscles of mesenteries.

3. Transverse section through a polyp of **Alcyonium**, below the end of the œsophagus. Mesenteric filaments.

ALCYONARIA

4. Section of undecalcified coenenchyma of **Alcyonium** made by hand; spicules; polyp tubes with mesenteries [and ova].

5. **Corallium rubrum**. Axial skeletal rod of red coral, covered by soft coenenchyma; polyps extended.

6. **Pennatula** (sea feather). Axis of colony containing a hard skeletal rod, and without branches on its basal portion; feather-like branches with polyps along one edge.

ZOANTHARIA (HEXACTINIA).

1. View of a **Peachia** opened along one side. Tentacles partly or entirely retracted. Body wall; mesenteries (twelve); secondary mesenteries; mesenteric filaments; gonads; œsophagus laid open; œsophageal groove, projecting lip of same.
2. View of the face of a mesentery. Gonad, filament as in 1; longitudinal muscle of mesentery.

3. Transverse sections of **Peachia** (*a*) at the level of the œsophagus, (*b*) below the end of the œsophagus.

ZOANTHARIA

4. A sea-anemone with tentacles expanded (**Anemonia sulcata**). Aboral disc for attachment ; tentacles ; mouth.
5. A colonial coral-forming Zoantharian (**Asteroides calycularis**). Polyyps, some expanded and some retracted, with mouth and tentacles ; thecæ or cups of calcareous matter with radiating septa.

6. The dried theca of a solitary coral-forming Zoantharian (**Fungia**). Septa.
7. Dried skeletons of different forms of corals. (a) **Lophohelia**, (b) **Oculina**. Theca ; septa ; costæ.

CTENOPHORA.

Cydippe. Rows of ciliated plates; wide oesophagus (stomach); tentacles; gastro-vascular canals.
Examine **Beroë** and **Cestus**.

PHYLUM PLATYHELMINTHES. *Flat worms without an anus. Excretory system as fine canaliculi ending in flame-cells. Generative organs complicated and generally provided with vitellaria. Hermaphrodite. No body cavity or vascular system.*

Class 1. **TURBELLARIA.** Free-living Platyhelminthes covered with cilia: with muscular protrusible pharynx. Classified according to the structure of the alimentary canal.

Order 1. **ACCELA.** Gut solid. Ex. *Convoluta*.

Order 2. **RHABDOCELA.** Gut hollow, rod-shaped. Ex. *Mesostoma*.

Order 3. **TRICLADA.** Three main branches to gut. Generative opening single. Ex. *Dendrocoelum*.

Order 4. **POLYCLADA.** Numerous branches to gut. Generative opening double. Ex. *Thysanozoon*.

Class 2. **TREMATODA.** Parasitic Platyhelminthes with suckers and hooks and forked alimentary canal. Without cilia.

Order 1. **MONOGENEA.** Trematodes with direct development. Ex. *Polystomum*.

Order 2. **DIGENEA.** Trematodes with alternation of hosts. Ex. *Distomum*.

Class 3. **CESTODA.** Parasitic Platyhelminthes with suckers and hooks on the head, and a chain of sexual proglottides breaking off when ripe; no mouth or alimentary canal. Ex. *Tenia*, *Bothriocephalus*.

TURBELLARIA**TURBELLARIA.**

1. **Dendrocœlum.** A fresh-water Triclad; naked eye or simple lens. Eyes; pharynx; terminal portions of generative ducts; enteric cæca.
2. The same compressed with low power to show genital organs, pharynx, etc. more in detail. With high power observe cilia and rhabdites.

TURBELLARIA

3. Transverse section of a **Planarian**, through pharyngeal region (low power:—the details to be drawn with a high power). Pharynx; pharynx-sheath; epidermis; enteric cæca; longitudinal nerve-cords; testes.

4. **Mesostoma** seen as a transparent object with a low power. Cerebral ganglia with two eyes; enteron; mouth; generative opening; uterus possibly containing winter or summer eggs; yolk-glands; testes; penis; ovary; muscular sac. Under high power notice flame-cells.

TREMATODA.

1. **Distoma hepaticum**, the liver fluke. (Simple lens.) Anterior, ventral sucker; genital pore.
2. Another specimen (stained), in which the excretory system has been injected. (Low power.)
Testes; yolk-glands; uterus.

TRANSMISSION

RECEIVED
JAN 10 1900
U. S. DEPT. OF AGRICULTURE
WASHINGTON, D. C.

3. **Redia** of Distoma. (Low power.) Pharynx; contained Cercariæ.
4. **Cercaria** under high power, with suckers and tail.
5. **Polystoma integerrimum**—from frog's bladder. (Simple lens.) Mouth on anterior sucker; disc with circle of suckers at hind end of ventral surface; two lateral papillæ.

ACIDOPHILUS

... ..
... ..
... ..

... ..
... ..
... ..

CESTODA

CESTODA.

1. **Tænia solium.** (Naked eye.) Head (scolex); chain of proglottides; genital opening.
2. Head of **Tænia.** (Low power.) Suckers; rostellum; hooks; budding zone.
3. **Cysticercus pisiformis**, from rabbit, being the cystic stage of *T. serrata* of the dog. (Lens.)

CESTODA

4. (a) Sexually mature proglottis of **Tænia**, stained, cleared in oil of cloves and mounted in Canada balsam. Draw (low power): genital pore; vas deferens; cirrus; vagina; testes; ovary; yolk-glands; excretory canals. With high power see calcareous bodies.
- (b) Old proglottis ready to separate from chain. Branching uterus full of eggs in various stages.

5. Transverse section through an adult proglottis of **Tænia**. Cuticle; cortical layer consisting of fusiform cells, intermediate layer, longitudinal muscles, circular muscles; inner zone with testes, uterus; large and small excretory canal; lateral nerve-cord; calcareous bodies.

1777

in the year 1777, the first year of the war, the British evacuated the city of Philadelphia and moved to Lancaster and then to York. The Continental Congress followed them and met in Lancaster and then in York. The British evacuated the city of Philadelphia and moved to Lancaster and then to York. The Continental Congress followed them and met in Lancaster and then in York.

CESTODA

6. **Cœnurus cerebralis.** The cystic stage of *Tænia cœnurus* from the brain of a sheep. Numerous scolices projecting from the wall of the vesicle.

7. **T. echinococcus;** cyst in human liver.

8. **Schistocephalus dimorphus** from the Stickleback (*Gasterosteus*).

PHYLUM NEMATODA. *Round worms, generally much elongated, with a single layer of longitudinal muscles in the body wall. Nervous system as a ring in the anterior part of the body giving off nerves backwards and forwards. Body cavity but no vascular system. Male generative organs open into cloaca. Amœboid spermatozoa. Often parasitic.*
Ex. Ascaris, Filaria, Trichina, Rhabditis.

NEMATODA.

1. **Ascaris lumbricoides.** Female dissected. Œsophagus; intestine; vagina dividing to form the two uteri, which pass backwards to the hind end of the body where they bend round and are continued forwards as the slender oviducts. The oviducts soon become continuous with the convoluted ovaries.

Male dissected. Œsophagus, intestine as above (hind end of intestine pinned on one side); vas deferens passing into testis—a convoluted tube; copulatory spicules in sheaths.

NEMATODA

2. Transverse section of **Ascaris**. (Low power.) Cuticle; epidermis; layer of longitudinal muscles; large lateral line of granular matter continuous with the epidermis and containing the excretory canal; less conspicuous dorsal and ventral lines; intestine (or oesophagus); sections of ovary or testis; section of the two uteri; or section of vas deferens.

3. Transverse section of muscular fibre of **Ascaris**. (High power.) Contractile cortical layer; granular inner mass; (nucleus).

4. **Trichocephalus dispar** in human colon.
5. Encysted **Trichina**.
6. Various free-living nematodes (**Rhabditis** or **Diplogaster**). In male note especially copulatory spicules and bursa.

PHYLUM ANNELIDA. *Segmented worms usually with well-developed coelom and metamerically segmented nephridia. Canalicular vascular system. Nervous system with cerebral ganglia, œsophageal ring and ventral nerve-cord with a ganglionic swelling in each segment.*

Class 1. **CHÆTOPODA.** Annelida with bristles (chætæ) secreted in pits of the ectoderm and serving as organs of locomotion.

Order 1. **OLIGOCHÆTA.** Chætopoda without parapodia or tentacles: hermaphrodite with localised generative organs. Development direct.

Sub-order 1. **Terricolæ.** Ex. *Lumbricus*, *Allolobophora*.

Sub-order 2. **Limicolæ.** Ex. *Nais*, *Tubifex*, *Chætogaster*.

Order 2. **POLYCHÆTA.** Chætopoda with chætæ generally arranged in bundles and carried on outgrowths of the body wall (parapodia). Head with tentacles and palps: the sexes are separate and the generative products are developed from the coelomic wall throughout the body. Generally with a Trochosphere larva. Ex. *Nereis*, *Aphrodite*, *Eunice* free-swimming forms: *Terebella*, *Arenicola*, *Filograna* tubicolous forms.

Class 2. **HIRUDINEA.** Annelida without chætæ and parapodia, moving by means of a terminal ventral sucker, the mouth acting as an anterior sucker, and living on the juices of other animals; coelom much reduced hermaphrodites with direct development.

Order 1. **RHYNCHOBDELLIDÆ.** Ex. *Clepsine*, *Pontobdella*.

Order 2. **GNATHOBDELLIDÆ.** Ex. *Hirudo*, *Nephelis*.

CHÆTOPODA.

OLIGOCHÆTA.

1. **Nais proboscidea.** (Low power.) Avoid too much pressure by using small slips of paper to support the cover.

Mouth; proboscis; eyes; groups of hooked ventral setæ; long dorsal hair-setæ; pharynx, segments 1—4; œsophagus, segments 5—7; crop; intestine; rectum; sacs in which the setæ are imbedded; muscles of setigerous sacs. (The hair-setæ are first found in the 5th segment, in which however they are shorter than in the succeeding.) Dorsal vessel; dilated lateral vessels placed immediately in front of a septum in segments 3, 4, and 5; nephridia—best seen in dorsal view (1st nephridium in 6th segment); [testes in 5th segment, ovaries in 6th, atrium and opening of vas deferens in segment 6; receptaculum seminis in segment 5; vesicula seminalis and ovarian sac unpaired and extending some way back]; zones of fission visible in some specimens.

2. With a high power draw

Brain; body-cavity septa; nephridium; funnel opening of nephridium into cœlom; tactile hairs.

POLYCHÆTA.

1. Dorsal view of head of **Nereis**. Præstomium, with eyes, tentacles and palps; tentacular cirri (cirri of 1st and 2nd segments fused with head).
2. Parapodium of **Nereis**. Notopodium and neuropodium, each with two or three lobes; notopodial cirrus; neuropodial cirrus; acicula; setæ.
3. **Heteronereis** (sexual) form. Enlarged eyes: modified posterior segments.
4. Parapodium of **Heteronereis** form: flattened expansions of parapodia: oar-shaped setæ.

POLYCHÆTA

- [illegible]

8. **Arenicola.** Mouth; anus; parapodia (very rudimentary); gills.
9. **Spirographis.** Tube; parapodia; branchial filaments.
10. **Filograna** (Serpulid). In 70 p.c. spirit, decalcify with weak nitric acid, stain in borax carmine and mount in Canada balsam. Tentacles (modified palps); parapodia; thoracic membrane (fused cirri) used in moulding tube; individual budded off posteriorly.

HIRUDINEA.

THE MEDICINAL LEECH.

1. **Hirudo medicinalis.** *A* dorsal, and *B* ventral aspect, of a medicinal leech killed by chloroform vapour. Annuli (95 in number), superficial constrictions, of which, in most regions, five appear to make up one segment or metamere as indicated by the deep-lying organs; oral sucker and mouth; posterior sucker; anus dorsad of the sucker; penis between 24th and 25th annulus, counting from the edge of the mouth; eyes (5 pairs).

HIRUDINEA

2. The Leech opened by a median dorsal incision extending from the posterior sucker to the prostomium. The eleven pairs of alimentary cæca (belonging to the same segments, respectively, as the 4th—14th post-oral ganglia); vessels (dorsal and lateral trunks) containing blood coloured red with hæmoglobin. Median rectum corresponding to the last eight ganglia; rectal cæca; anus; seventeen pairs of nephridia existing in the same segments as the 2nd to the 18th post-oral nerve ganglia inclusive, though the first is in the same horizontal line with the third post-oral ganglion.

3. Remove the alimentary canal and expose the nephridia, generative organs and nerve-cord. The nine pairs of testes, the first lying in the same segment as the 8th post-oral ganglion, and between the 6th and 7th pairs of nephridia; the epididymis on each side in the 7th post-oral; the vas deferens on each side; the penis opening in the middle line behind the 6th post-oral ganglion; the pair of ovaries; the glandular enlargement of the oviducts (albumen gland); the vagina opening in the middle line behind the seventh ganglion; the seventeen pairs of nephridia numbered; the twenty-three post-oral ganglia of the nerve-cord; the supracæsophageal ganglion; cæsophageal passage between the cerebral ganglion and the 1st post-oral; the double nerve-cord enclosed in ventral sinus.

HIRUDINEA

4. Transverse section through a Leech under low power. Epidermis ; alimentary canal ; connective tissue and muscles of the body ; dorsal sinus ; ventral sinus enclosing the nerve-cord ; lateral vessel ; fine capillary branches of vascular system ; nephridia ; vas deferens ; unicellular glands dipping into body muscles from the epidermis ; longitudinal muscular fibres ; external circular muscular fibres ; dorso-ventral muscles.
5. Sections from the anterior extremity to show the eyes and special sense organs.

CRUSTACEA. *Aquatic arthropoda usually with two pairs of antennæ on the head and breathing by gills.*

Class 1. **ENTOMOSTRACA.** Small Crustacea with a variable number of segments and appendages. Larva, when present, has the Nauplius form.

Order 1. **PHYLLOPODA.** Entomostraca with leaf-like swimming feet and usually a carapace or bivalve shell formed by a reduplication of the skin. Ex. *Apus*, *Daphnia*.

Order 2. **OSTRACODA.**

Order 3. **COPEPODA.** Entomostraca without shell-forming reduplication of the skin. Swimming feet biramous. Abdomen without appendages. Ex. *Cyclops*.

Order 4. **CIRRIPEDIA.** Fixed and for the most part hermaphrodite Entomostraca with indistinctly segmented body enclosed by a reduplication of the skin and a calcareous valved shell. Usually six pairs of biramous thoracic appendages. Ex. *Lepas*, *Balanus*, *Sacculina*.

Order 5. **TRILOBITA.** Palaeozoic Entomostraca with one pair of antennæ.

Class 2. **MALACOSTRACA.** Crustacea with a definite number (20) of segments and appendages. Larva when present usually has the Zoea form.

Order 1. **ARTHROSTRACA (Edriophthalmata).** Malacostraca with sessile eyes, one pair of maxillipeds, seven free thoracic segments, without thoracic carapace.

Sub-order 1. **Amphipoda.** Ex. *Gammarus*.

Sub-order 2. **Isopoda.** Ex. *Oniscus*, *Asellus*.

Order 2. **THORACOSTRACA (Podophthalmata).** Malacostraca with eyes which are often stalked; with a dorsal shield which connects all, or at least the anterior thoracic segments with the head.

Sub-order 1. **Cumacea.**

Sub-order 2. **Stomatopoda.** Ex. *Squilla*.

Sub-order 3. **Schizopoda.** Ex. *Mysis*, *Euphausia*.

Sub-order 4. **Decapoda.** Ex. *Astacus*, *Cancer*.

CRUSTACEA.

1. **Apus**, dorsal view. Carapace; backwardly projecting free portion of same; transverse groove marking junction of these two parts; eyes; externally segmented posterior part of body; terminal setæ.

Ventral view. Notice that the anterior legs are pediform and stout, the posterior foliaceous and delicate. The posterior six segments are without appendages.

2. Appendage of **Apus** (typical Phyllopod appendage). Corm or axis of limb, bearing 6 endites, and 2 exites—the flabellum and bract. Endites 5 and 6 correspond to the exopodite and endopodite of the appendages of *Astacus*, and the flabellum to the epipodite. In some legs there is an apical lobe beyond the endite 6.

PHYLLOPODA

3. **Daphnia**, the water-flea. Mount on slide, and keep in place by supported coverslip. Low power. Head; fused compound eyes; median simple eye; first pair of antennæ (rudimentary but seen with high power); second pair of antennæ, large and biramous with long bristles bearing delicate hairs; mouth; hepatic diverticula of anterior part of alimentary canal; heart; brood pouch, probably containing ova or embryos; thoracic appendages; abdomen curved ventralwards within the valves of the shell, which is formed by the prolonged lateral halves of the carapace; anus terminal and slightly dorsal; terminal setæ.

Notice the circulation of the blood as shown by the movement of the corpuscles.

COPEPODA

1. **Cyclops**, dorsal view. Steady the animal with a coverslip on a slide. (Low power.) Carapace; eye; 1st and 2nd pairs of antennæ (uniramous); four appendage-bearing thoracic segments behind the cephalo-thorax; 1st abdominal segment, double in the male, single in the female. The generative openings are on this segment and the ovisacs are attached to it; three posterior abdominal segments, of which the last bears the caudal fork; genital gland in the thorax; alimentary canal.

Segments as follows:

Head	{	ant. 1.
		ant. 2.
		mandibles.
		maxilla 1.
		maxilla 2.

Thorax. maxillipeds (uniramous).

5 pairs of biramous appendages (last much reduced).

Abdomen without appendages. 5 segments in males, 4 in females.

CIRRIPEDIA

1. **Lepas**, side view. Scutum; tergum; carina; opening of shell; peduncle.
2. **Lepas**, side view, with shell and mantle of one side removed. Oral cone; gills attached to base of anterior thoracic legs; thorax with six pairs of biramous legs; abdomen rudimentary and continued as penis, just dorsal to root of which is anus; testicular follicles, on sides of prosoma and thorax.

Cut open peduncle and observe ovaries. Examine the inner side of the uninjured mantle for ovigerous frenum, to which may be attached the ovarian lamella.

CRUSTACEAN LARVÆ.

Nauplius larva with three pairs of appendages, the future 1st antennæ (uniramous), 2nd antennæ and mandibles (biramous); the single median eye. The unjointed carapace may be prolonged into long spines.

Zoæa larva, cephalothoracic shield with spines; free abdominal segments; forked telson; compound stalked eyes.

The appendages in a typical Zoæa are:

- Two pairs of antennæ.
- Mandibles without palp.
- Two pairs of maxillæ.
- Two pairs of maxillipeds.

Gammarus fluviatilis. Side view.

1st antenna; 2nd antenna; head (with which the 1st thoracic segment is fused); oral appendages; eyes; the seven free thoracic segments; the six abdominal segments.

Appendages of *G. fluviatilis*, mounted and seen with low power:

antenna 1.

antenna 2.

mandible.

maxilla 1.

maxilla 2.

Asellus aquaticus.

1. Dorsal view of the entire animal, examined in a watch-glass. Head; eyes; the seven free terga of the thorax (the tergum of the 1st thoracic segment is obsolete); single piece representing the fused abdominal segments (the 1st abdominal segment may perhaps be seen as a minute free ring); 1st antennæ; the much longer 2nd antennæ; the seven pairs of ambulatory limbs; the bifurcated 6th abdominal appendages.

(In small and transparent specimens: alimentary canal; liver.)

2. Ventral view. Abdominal appendages; brood-pouch in female, formed by the oostegites, processes of the thoracic appendages.

DECAPODA

Carcinus, common shore-crab. Note the short 2nd antennæ, the reduced abdomen, wide in the female with four pairs of biramous ovigerous appendages, narrow in the male with two pairs of appendages only, forming copulatory styles.

Class 2. **ONYCHOPHORA.** Tracheates with a soft skin, paired segmental nephridia and numerous similar leg-bearing segments. The head is not marked off from the body and bears one pair of antennæ, one pair of jaws and one of oral papillæ. Tracheal stigmata diffusely scattered. Muscular fibres unstriated. Ex. *Peripatus*.

ONYCHOPHORA.

PERIPATUS CAPENSIS.

1. Ventral view.

Antennæ; oral papillæ; mouth; lips; jaws; the seventeen pairs of legs; generative opening; anus.

2. Dorsal view of head to show the eyes. (Simple lens.)

3. **Peripatus** dissected so as to show the general anatomy.

Brain; ventral nerve trunks; pharynx; œsophagus; intestine; rectum; slime glands; reservoirs of slime glands; salivary glands; testis and vas deferens, or ovary, uteri and vagina.

Class 3. **MYRIAPODA**. Tracheata with numerous similar leg-bearing appendages and with one pair of antennæ. Tracheal stigmata regularly and segmentally arranged.

Order 1. **CHILOPODA**, with three pairs of jaws (mandibles and two pairs of maxillæ) and a pair of poison claws. One pair of legs to each body segment. Ex. *Lithobius*, *Scolopendra*.

Order 2. **DIPLOPODA**, with two pairs of jaws (mandibles and maxillæ). Body consisting of numerous double segments. Ex. *Julus*.

MYRIAPODA.

1. **Lithobius.** Ventral view. Antennae; head; 1st body segment, with poison claws; broad sterna of body segments; ambulatory appendages.

In side view note stigmata in alternate segments.

2. **JAWS** of same. Mandibles; 2nd maxillæ; 1st maxillæ united in the median line; poison claw (appendage of 1st post-cephalic segment).

3. **Julus.** Note short antennæ, cylindrical body, narrow sterna, numerous legs (two to each segment of the body).

4. **JAWS** of same. Mandibles; maxillæ as a four-lobed plate.

Class 4. **INSECTA.** Tracheata with the body divided into three regions: (1) the head with a single pair of antennæ and mouth parts, (2) the thorax with three pairs of legs and usually two pairs of wings, and (3) the abdomen with 10 segments usually without appendages. Young always more or less different from adult.

Order 1. COLLEMBOLA. Spring-tails. }	Wingless insects whose ancestors are believed to have been also wingless.
Order 2. THYSANURA. Bristle-tails. }	
Order 3. SIPHONAPTERA. Fleas. Wingless animals supposed to have developed from winged ancestors.	
Order 4. ORTHOPTERA. Earwigs, locusts, crickets, grasshoppers, leaf- and stick-insects.	
Order 5. ISOPTERA. (Termitidæ) white-ants.	
Order 6. PARANEUROPTERA. (Odonata) dragon-flies.	
Order 7. HEMIPTERA. Bugs, plant-lice.	
Order 8. NEUROPTERA. Ant-lions, lace-wings.	
Order 9. LEPIDOPTERA. Moths and butterflies.	
Order 10. COLEOPTERA. Beetles.	
Order 11. DIPTERA. Flies.	
Order 12. HYMENOPTERA. Ants, saw-flies, bees, wasps.	

INSECTA.

THE COCKROACH. (PERIPLANETA AMERICANA.)

In *P. orientalis*—the common species—the wings are rudimentary in the female. In *P. americana*, both sexes have wings.

1. Dorsal or tergal aspect of a full grown male specimen pinned by the base of the legs. Head; thorax; abdomen; antennæ; prothorax; mesothorax; metathorax; mesothoracic wing, pinned forward; metathoracic wing, pinned forward. 1st to 7th abdominal segments or somites. Terga of the 8th and 9th somites overlapped by the 7th. Tergum of the 10th somite. Right and left styles attached to sternum of the 9th somite. Right and left jointed cerci anales attached to the tergum of the 10th somite.

2. Dorsal aspect of female specimen.

Look for the **heart** through the thin dorsal skin of the abdomen beneath the wings.

PERIPLANETA

3. Ventral or sternal aspect of abdomen of a male specimen. Sterna of 1st to 9th abdominal somites; right and left styles attached to the 9th sternum; bifid sternum of the 10th somite, called "podical plates"; tergum of 10th abdominal somite; genital aperture between the sterna of the 9th and 10th somites; anus between the sternum (podical pl.) and tergum of the 10th somite.

PERIPLANETA

4. Enlarged view of the posterior part of the sternal aspect of the abdomen of an adult female. Sterna of the 1st to 6th abdominal somites; sternum of the 7th abd. somite (elongate and bifid posteriorly); proper position for the sternal sclerites of somites 8 and 9, represented by a soft expansible membrane, the genital funnel; genital aperture placed beneath the podical plates (sternum of 10th somite) and leading into genital funnel.

5. A leg detached, and magnified three times linear. Coxa; trochanter; femur; tibia; the six joints of the tarsus; the two unguis or claws.

6. The head and neck as seen laterally (enlarged). Antenna; compound eye; fenestra; maxillary palp; labial palp; labrum; cervical sclerites.

7. Head detached and seen from above. Antenna; eyes; fenestræ; epicranial suture; clypeus; labrum; maxillary palp; labial palp.

62-9

PERIPLANETA

8. The 3rd post-oral pair of appendages or 2nd maxillæ (labium and labial palps). Submentum; mentum; bifid ligula; paraglossæ; labial palp.

[N.B. For preparations of insect mouth parts, the detached appendage should be boiled with potash to remove attached muscles, dehydrated and mounted in Canada balsam.]

9. One of the 2nd post-oral pair of appendages or 1st maxillæ. Cardo; stipes; palp; lacinia; galea. (Compare with 8 and with mandible and maxillæ of Crayfish.)

10. One of the mandibles.

11. Enlarged view of the two thoracic stigmata of one side placed between prothorax and mesothorax, and between mesothorax and metathorax.

PERIPLANETA

12. **Dissection of Heart and Alimentary Canal.** Remove the abdominal terga carefully so that the heart lying within the pericardium may be seen in situ. Then remove also the thoracic terga and free the alimentary canal from the adherent masses of the fat body and unravel its coils. The larger tracheal branches should be cut with fine scissors. Right salivary gland; right salivary receptacle; oesophagus; crop; proventriculus; pyloric cæca; chylific ventricle; Malpighian tubes, opening into the short small intestine or ileum; large intestine or colon; rectum; cerebral ganglion.

13. Right salivary gland and receptacle removed with ducts, &c. entire. Under low power: acini; tracheal tubes; ducts; nerves. This should be fixed in corrosive sublimate, stained in borax carmine, and mounted in balsam.

14. Portion of fatty body showing tracheæ and fatty tissue. High power.

PERIPLANETA

15. The remaining organs of the same cockroach as seen after the removal of the alimentary canal and fatty body. Cerebral ganglion; sub-oesophageal ganglion, closely attached to the cerebral and representing three fused ganglia (those of the mandibular and two maxillary segments); 1st, 2nd, and 3rd thoracic ganglia; 1st abdominal ganglion; 2nd to 5th abdominal ganglia; 6th and last abdominal ganglion. In FEMALES. Colleterial glands; oviduct; spermatheca; ovarian tubes. In immature MALES. Testis as a somewhat transparent lobed body lying on each side in the fat immediately below the 4th and 5th terga; vas deferens as a slender duct passing back to the small mushroom-shaped gland. In adult MALES. Mushroom gland (vesicula seminalis); short duct of same; accessory gland.

16. Ovarian tube under low power of microscope.
17. Contents of the mushroom gland and spermatheca. Spermatozoa.
18. Malpighian tubule under high power.
19. Muscular fibre under high power.

INSECTA.

1. Head of a female **Culex** [Diptera] from above. (Low power.) Eyes; antennæ; labrum or upper lip; the proboscis constituted by the labium or fused 2nd maxillæ, in a dorsal groove of which are contained (when not extruded) the styliform mandibles and 1st maxillæ and the hypopharynx or lingua; the groove is closed dorsally by the labrum; maxillary palps.

[In the male, the maxillary palps are much longer, and the antennæ bear very long hairs.]

2. Head of a **Notonecta** or Water-boatman [Hemiptera] from above. (Low power.) Eyes; antennæ; labrum; jointed proboscis or labium; the stylet-like mandibles and 1st maxillæ, contained in the proboscis. [There are no maxillary or labial palps.]

INSECTA

3. Mouth-parts of a **Bee** removed from head and mounted: (1) the labium with the ligula, bearing on each side a shorter process the paraglossa and a long five-jointed labial palp; (2) the 1st maxilla with stipes, lacinia folded back on the stipes and a short palp at end of stipes; (3) the mandible strong and platelike.

4. Mouth-parts of a **Butterfly**. (These can be best seen if the two eyes are removed from the head and the rest boiled in potash and mounted.) Spirally coiled proboscis formed by the union of the two 1st maxillæ; rudimentary maxillary palps; three-jointed palps of the rudimentary labium.

5. Side view of a **Lepidopterous caterpillar**. Head; the three thoracic segments, each with a pair of legs; the ten (9th small) abdominal segments, the 3rd to 6th being each provided with a pair of 'pro-legs'; the eight abdominal stigmata of one side.

6. Side view of a **Lepidopterous pupa**. Head; eyes; antennæ; maxillæ; prothorax; mesothorax; prothoracic leg; mesothoracic leg; mesothoracic wing; metathoracic wing; the ten abdominal segments; the eight abdominal stigmata.

ARACHNIDA.

1. **Scorpio**, dorsal view. Chelicerae; pedipalpi; carapace; the pair of large median eyes; group of small lateral eyes; terga of first seven abdominal segments; five post-abdominal segments; telson.

2. **Scorpio**, ventral view. Chelicerae; pedipalpi with stout basal joint; four pairs of ambulatory legs; jaw-like process of basal joints of the legs of the first two pairs; genital operculum (on 1st abdominal); pectines; four pairs of pulmonary stigmata; 7th abdominal segment; postabdomen and telson as in 1.

3. **Spider**, dorsal view. Chelicerae; pedipalpi; ambulatory legs 1—4; eyes; cephalothorax; unsegmented abdomen.

4. **Mite**, ventral view. Thorax and abdomen broadly fused; chelicerae; pedipalpi; ambulatory legs; reproductive opening and anus.

LIMULUS

1. **Limulus**, dorsal view. Cephalothorax; abdomen; spines (six on each side) on margins of abdomen; telson; the two compound eyes at the sides of the median region of the cephalothoracic carapace; the two simple eyes at the front end of the same region.

2. **Limulus**, ventral view. Pediform legs on cephalothorax (six pairs), the last five with gnathobase; spatulate digging processes on last leg; genital operculum; lamellar abdominal appendages (five pairs) with gill books; anus; telson; position of mouth.

3. Casts of fossil forms **Eurypterus** and **Prestwichia**.

MOLLUSCA.

1. **Helix.** View of the entire animal killed by immersion for two days in warm water. Shell; head; the two superior or oculiferous tentacles; the two inferior tentacles; foot; thickened edge of the mantle; apex of the shell; mouth of the shell; pore of the mucous gland of foot.

2. The animal after removal of the shell viewed from the right side. Head; foot; surface of mantle covering the visceral hump; mantle-fold, at the junction of the visceral hump with the head and foot; circular respiratory orifice leading into the pulmonary sac; anus; genital pore; columella (or spindle) muscle; visceral hump containing liver and hermaphrodite gland.

3. View of the respiratory chamber, opened by a cut extending from the respiratory orifice across to the left side, and by another cut on the right side passing backwards along the extreme right-hand limit of the chamber, so as to leave the anus, rectum, and kidney in continuity with the roof. Floor of the respiratory chamber formed by the thin dorsal integument; rectum; kidney; external opening of kidney; duct of the kidney by the left side of the rectum; auricle and ventricle of the heart lying in the pericardium; blood-vessels of the respiratory surface.

N.B. Remember that rectum, kidney, and heart are turned back and are now seen inverted.

4. Notice that the partition between the pericardium and the kidney sac is indented by a shallow bay. At the end of this bay is the reno-pericardial opening. Open the pericardium and pass a fine guarded bristle through the opening into the kidney.

5. Open the dorsal integument by a median cut extending from the head to the visceral hump, unravel generative organs and pin out to right side of animal. Generative pore on neck; hermaphrodite gland; hermaphrodite duct dividing into oviduct with thick glandular walls and slender sperm-duct; albumen gland opening into hermaphrodite duct; spermatheca; spermathecal duct and its long appendix; digitate glands and dart-sac opening into vagina; penis; flagellum of male duct.

1871
The first of the year was a very dry one
and the crops were much injured
by the drought. The wheat was
very poor and the corn was
also much injured. The
cattle and sheep were
also much injured by the
drought. The people were
very poor and the
country was very dry.

6. A complete view of the digestive organs obtained by carefully separating the alimentary tract from its attachments. In dissecting, the generative organs are pushed to the right side, the digestive to the left. Buccal mass; nerve-collar; ducts of the salivary glands; the two salivary glands; oesophagus; proventriculus; stomach; first bend of the intestine; rectum; anus; the kidney, between the rectum and the heart.
7. Radula removed from buccal mass and examined in glycerine under low power.

8. Nerve-collar and chief nerves shown by drawing the œsophagus forward through the collar, and cutting it in front of the collar, then turning the buccal mass upwards and forwards so as to expose its ventral surface. Cerebral or supra-œsophageal ganglion, giving off a pair of nerves to the buccal ganglia on the dorsal surface of the buccal mass but ventral to the œsophagus; nerves from cerebral ganglion to the tentacles; infra-œsophageal ganglion (pedal, pleural and visceral fused); nerves from pedal ganglion to foot.

9. Calcareous spicule removed from the dart-sac and enlarged three times.

10. Section of hermaphrodite gland, examined under high power. Ova; sperm mother-cells; spermatozoa.

1. **Chiton.** Ventral view. Head, foot, mantle-groove containing the numerous ctenidia.
2. Dorsal view. The eight shell plates ; girdle.

3. **Haliotis.** Dorsal view of shell. Spiral; row of pores.
4. Shell removed and mantle-cavity opened. The two ctenidia with anus between them. Mucous folds in mantle-roof.
5. **Patella.** The common Limpet. Ventral view. Shell; pallial branchiæ forming a ring round the foot in the mantle-groove. Head.

6. **Buccinum.** The common Whelk. Proboscis everted; tentacles. Foot with operculum. Shell.

7. Removed from shell, mantle-cavity opened. Rectum, anus; ctenidium, one only, the left (primitive right), persists. Mucous folds; complicated osphradium. Renal opening at hind end of mantle-chamber. In female, opening of oviduct. In male, long penis with convoluted vas deferens passing along its left side and opening on a small papilla near its apex.

8. **Aplysia.** Foot with lateral flaps (epipodia) projecting dorsalwards; mantle with shell in nearly closed sac; single ctenidium; spermatic groove leading from genital aperture to penis on head.
9. **Eolis.** Dorsal cerata: without ctenidia and mantle folds.
10. **Doris.** Dorsal view. Anus median and posterior; anal branchiæ.

1. **Anodonta.** View of the animal lying in the right valve of the shell, the left having been removed by carefully separating from it the edge of the mantle and cutting through the attachment of the two adductor muscles to the shell (keep the edge of the knife close to the shell). Left lobe of mantle; thickened free edge of same; right lobe of mantle; margin of posterior border of mantle-flap twice indented so as to form two incomplete siphons—dorsal and ventral, the latter with tentaculiferous margin; anterior adductor; posterior adductor; anterior retractor; posterior retractor; protractor.

2. Side view of animal with left lobe of mantle turned back. Foot; labial palps; gills; line of attachment of outer gill with mantle. Right mantle-lobe; dorsal siphon; ventral siphon.

3. Side view with the labial palps and gill plates of left side pinned back. Inner lamella of inner gill plate attached to the foot anteriorly, free medianly and fused with the corresponding lamella of the other side posteriorly. Inner and outer gill plates of the right side. Probe passed between the foot and the middle portion of the inner gill plate into the epibranchial chamber. Apertures of the left kidney exposed by cutting forward through the attachment of the inner gill plate to the foot, and of the generative gland slightly in front of the aperture of the kidney.

4. View of heart with dorsal wall of pericardium removed. Ventricle surrounding the rectum; left auricle; right auricle; rectum traversing ventricle.

5. View of kidneys when the auricles have been cut through laterally, the ventricle and rectum posteriorly and the heart turned forwards. Reno-pericardial apertures lying at extreme anterior limit of the pericardium on each side of the intestine. Dark green glandular portion of the kidney. Non-glandular ureter exposed by cutting through the floor of the pericardium. Aperture of kidney to exterior indicated by bristle near and below reno-pericardial aperture.

6. View of nervous system (ganglia are yellow in fresh specimens). Cerebral ganglion; pedal ganglion; cerebro-pedal commissure; parieto-splanchnic ganglia; visceral commissure running for the greater portion of its length through the glandular part of the kidney.

7. View of alimentary canal. Position of mouth; stomach; liver; coiled intestine; rectum; anterior adductor; posterior adductor; foot. This drawing may be made from a prepared specimen.

8. Three transverse sections of a spirit specimen as seen with the naked eye.

(a) Through the middle part of the foot. Mantle-lobe; foot; outer gill; inner gill; (notice attachment of lamellæ); liver; rectum; intestine in foot in several places.

(b) Through the hinder part of the foot. Mantle-lobes; foot; outer gill; inner gill; (notice attachment of lamellæ); epibranchial chamber; pericardium; ventricle; rectum; auricles; vena-cava; non-glandular part of kidney; glandular ditto.

(c) Behind the foot. Notice floor of epibranchial chamber continuous across the middle line.

GILLS OF LAMELLIBRANCHIATA

1. **Nucula.** Ctenidium with two rows of short filaments.
2. **Mytilus.** (a) Adjacent filaments of the gill mounted in sea-water ciliary junctions. Only one of the limbs of the filaments should be mounted.
(b) Single filament seen endwise. Ascending limb ; descending limb ; interlamellar concrescences.

GILLS OF LAMELLIBRANCH

3. **Mytilus.** Transverse section through filaments; ciliated epithelium; large epithelial cells and ciliary junction; chitinous rods; vascular space of filaments.
4. **Anodon.** Transverse section through gill. Filaments in section; ciliated epithelium, chitinous rods, vascular spaces in sub-filamentar tissue; interlamellar junctions; water pores leading into epibranchial space between gills.

5. External features of other Lamellibranchs.

Mytilus. Reduced foot with byssus; gills; mantle containing gonads.

Pecten. Single adductor muscle; rudimentary foot; mantle-edge with numerous eyes.

Cardium. Elongated foot used for jumping.

Solen. Siphons; the two mantle-edges fused save for the slit through which the elongated foot projects.

1. **Sepia.** View of the anterior face of the entire animal;—head; the eight short arms; the partly hectocotyliised fourth left arm of the male; the two long arms, retractile into special pits; eyes; visceral sac; mantle-edge; lateral fins.
2. View of posterior face. Funnel.

3. View of the posterior surface after the mantle-cavity and funnel have been opened by a median incision. Valve of funnel; sockets which receive the two articular knobs on the mantle-flap; anus; renal papillæ; gills; generative papilla on the left side; ink-gland, and its duct; stellate ganglia. [In the female, nidamental glands, and accessory glands.]

4. View of the excretory and vascular systems, obtained by cutting through the posterior body wall, removing the ink-sac, and opening the two renal sacs. Renal papilla; coelomic opening of kidney; the large vena cava bringing back blood from the head, and bifurcating on arriving at the kidneys; paired veins from posterior part of body, dilating into large sacs just before they reach the branchial hearts; genital vein entering the right branch of the anterior cava; branchial hearts each with its appendage; afferent branchial veins; efferent branchial veins, passing into the auricles; ventricle; the anterior and posterior aortæ.

5. View of the alimentary canal and its glands, obtained by turning the animal over, and dissecting through the anterior wall of the visceral sac.

Oesophagus leaving skull, and accompanied by the anterior aorta and the salivary ducts; the two salivary glands; liver; liver ducts with pancreatic cæca; stomach; spiral cæcum of stomach; intestine.

6. Male generative organs. Testis, lying in a coelomic sac into which opens the vas deferens, at first narrow and coiled, then dilating into a wide tube—the vesicula seminalis; prostate; receptacle of the spermatophores; genital papilla with terminal aperture.

7. Female organs. Ovary, lying in a coelomic sac, into which opens the oviduct; nidamental glands; accessory glands.

8. View of brain as exposed by a median longitudinal section through the skull. Supra-oesophageal ganglion; superior buccal ganglion; inferior buccal ganglion; oesophagus; salivary duct; oesophageal commissure; pedal ganglion giving off nerves to the arms and continuous behind with the visceral ganglion; the latter gives off on each side a large nerve to the stellate ganglia; sections of auditory sacs ramifying in the cranial floor.

9. View of radula and its sheath from the side, obtained by removing the jaws and dissecting away the muscles of the buccal mass. Radula ; sheath of radula ; cesophagus.
10. Spermatophore under low power ; spermatocysts ; spiral spring.
11. Transverse section of an eye. Cornea ; iris ; lens ; its ciliary attachments ; retina ; optic ganglion ; optic nerve.

Demonstration specimens of other forms.

SHELLS OF CEPHALOPODA

Ammonites and **Nautilus** (Tetrabranchiata). Septa; siphuncle.

Belemnites (extinct Dibranchiata). Phragmocone (rudiment of original chambered shell); rostrum or guard; proostracum. (Notice hooks on arms.)

Sepia. Internal 'cuttle bone.' Rudimentary rostrum; laminated region: lateral expansion.

Loligo. Internal horny 'pen.'

ECHINODERMATA.

1. **Asterias.** Sketch of the entire animal, from the ambulacral side. Mouth; ambulacra, with tube-feet; ocelli; spines fringing the ambulacral groove: with a lens observe large pedicellariæ situated on these spines.

2. Dorsal view of same. Madreporic plate; arms of the trivium; arms of the bivium.

In live specimens notice the dermal branchiæ.

3. Alimentary canal. Divide the animal into a dorsal and ventral half by cutting through the body-wall along the sides of each arm, the radial muscles of the stomach and the cesophagus: cut round the madreporic plate so that the stone-canal shall be left in the ventral half and then remove the dorsal half with the alimentary canal and note: the stomach with its five diverticula; the pyloric sac with its thin walls, and its five radial prolongations, which bifurcate to form the ten hepatic cæca; the rectal cæca; the ten gonads.

4. View of the organs exposed after removal of the alimentary canal. Stone-canal (with heart); position of water-vascular ring; Tiedemann's bodies; ambulacral ossicles; ampullæ of tube-feet; cut oral end of stomach.

For Polian vesicles see Solaster; there are none in Asterias.

5. Sketch of a pedicellaria, drawn with a low power.

6. Transverse section of the decalcified arm of a small specimen. Body-wall; spines; dermal branchiæ; ambulacral ossicles; neural epithelium; radial water-vascular canal; ampullæ; tube-feet; body-cavity; hepatic cæca and their mesenteries; in sections near base of arms, generative organs.

Examine sections of an undecalcified arm to show ambulacral ossicles.

7. Region of ambulacral groove under high power. Columnar epithelial cells; internal prolongations of same; sections of nerve-fibres and nerve-cells between the latter; septum containing the so-called radial blood-vessel; radial water-vascular trunk; ambulacral ossicles; ampullæ; tube-feet.

8. Transverse section of stone-canal and heart.

ECHINODERMATA

1. **Echinus.** Ventral view of live animal: peristomial membrane with mouth; teeth; ten branchial plumes of peristomium (diverticula of body-cavity).

2. **Shell** of Echinus cleaned. Dorsal view: anal opening; periproct; genital plates with genital openings, one of them perforated by the water pores of the madreporite; ocular plates; five double rows of ambulacral plates with ambulacral pores for the tube-feet; five double rows of interambulacral plates; tubercles to which the spines are articulated.

3. Demonstration specimens of Echinoids and Ophiuroids.

Demonstration specimens of Holothurians and Crinoids.

Auricularia form of Holothuroidea.

Bipennaria form of Asteroidea.

Pluteus form of Echinoidea and Ophiuroidea.

TUNICATA

1. **Ciona.** Animal removed from its test and drawn in side view. Mouth; atrial aperture; ganglion; longitudinal muscles of body-wall; red pigment spot marking termination of vas deferens; point of origin of test vessels.
2. Atrial cavity and body cavity opened along the dorsal middle line; body wall separated from the pharynx by cutting the cords passing from one to the other. Pharynx; œsophagus leading into stomach; intestine with typhlosole; rectum; heart in pericardium; oviduct and vas deferens opening into atrial cavity.

3. **Ciona.** Pharynx and buccal cavity opened and viscera further exposed. Tentacular ring; peripharyngeal groove; dorsal tubercle; subneural gland; endostyle; languets and hyperpharyngeal groove; aperture from pharynx to œsophagus; ovary; testicular follicles on intestine.

4. Portion of pharyngeal wall stained and mounted. Transverse bars; longitudinal bars; internal longitudinal bars from inner side of which project the papillæ; stigmata.

5. An Ascidian tadpole of **Ciona** or some other Tunicate.
6. Transverse section of **Ciona**. Test; body wall with longitudinal muscle bundles; atrial cavity; pharynx with stigmata; endostyle and hyperpharyngeal groove; rectum and generative ducts.

7. Demonstration specimens of other Tunicata.

Demonstration specimens of **Balanoglossus**.

HEMICHORDATA

THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW

AN INITIAL FINE OF 25 CENTS

WILL BE ASSESSED FOR FAILURE TO RETURN
THIS BOOK ON THE DATE DUE. THE PENALTY
WILL INCREASE TO 50 CENTS ON THE FOURTH
DAY AND TO \$1.00 ON THE SEVENTH DAY
OVERDUE.

MAY 25 1963 BIOLOGY LIBRARY

11MY'63RB

OP 31 3041 1963

BIOLOGY
LIBRARY

254175

QL
53
C35

Cambridge,

THE UNIVERSITY OF CALIFORNIA LIBRARY