S B

369

G32

UC-NRLF

\$B 307 692

YB 47626

GIFT OF

EX LIBRIS

MAR 28 1915

CITRUS CULTURE

in the

WEST INDIES

LIBRARY
OF THE
UNIVERSITY
OF CALIFORNIA

1880

1880

CITRUS CULTURE

IN THE
WEST INDIES

PUBLISHED BY
AGRICULTURAL BUREAU
GERMAN KALI WORKS,
HAVANA CUBA.

LA PRUEBA, PRINTERS, OBRAPIA 99
HAVANA.

STB 369

G 32

TO THE
LIBRARY OF THE
CONGRESS

10070

A. Y.
THIS PAMPHLET AS WELL AS THE FOLLOWING CAN BE
OBTAINED FREE OF CHARGE BY WRITING TO

GERMAN KALI WORKS

P. O. BOX 1007. HAVANA, CUBA.

PUBLISHED IN

ENGLISH

Bananas in the West Indies

Cacao Culture in the West Indies

Coffee Culture in the West Indies

Cotton Culture

Fertilizer Guide for the West Indies

Miscellaneous Economical Plants in the West Indies.

Pineapple Culture in the West Indies

Stassfurt Industry

Sugar Cane Culture in the West Indies

Tobacco Culture in the West Indies

SPANISH

Cultivo de la Caña en las Antillas

Cultivo del Café en las Antillas

Cultivo de la Piña en las Antillas

Cultivo del Tabaco en las Antillas

Diversas Plantas Utiles en las Antillas.

El Cultivo del Cacao en las Antillas

Guía de Fertilizantes para uso en las Antillas

Naranjos en las Antillas.

FRENCH

Guide des Engrais A L'usage des Antilles

La Potasse sous les Tropiques

La Culture du Café aux Antilles

La Culture du Tabac aux Antilles

Les Bananiers aux Antilles

CONTENTS

	<u>Page.</u>
FOREWORD	7
GENERAL INFORMATION	7
PROPAGATION	7
INSECTS AND DISEASES	7
COMMERCIAL IMPORTANCE	8
CHOOSING A LOCATION	10
Exposure	11
Soils	11
Water	12
Transportation	12
Labor	13
Social Conditions	13
The Man	13
The Capital	14
METHODS OF STARTING A GROVE	14
The Commercial Method	14
Modification No. 1.—Windbreaks	16
Modification No. 2.—Planting from Own Nursery	17
Modification No. 3.—Planting Seedlings and Budding in the Field	19
VARIETIES OF STOCK AND BUD	19
The Sour Orange	21
The Rough Lemon	21

	<u>Page.</u>
The Grapefruit and Shaddock	22
Commercial Varieties of Oranges	22
Varieties of Grapefruit	23
HOW TO PLANT A TREE	24
The Distance Apart	25
HOW THE WILD TREES GROW	25
Nature's Way	26
Imitating Nature	26
CULTURAL METHODS	28
Local Wind Protection	28
Mulching	28
Cultivation	30
Irrigation	31
Fertilizing	32
Lime	34
Potash	34
Phosphorous	34
Nitrogen	35
Cover Crops	35
INFLUENCE OF FERTILIZERS ON THE TREES	36
INFLUENCE OF FERTILIZERS ON THE FRUIT	39
FERTILIZER FORMULAS	44
HOW TO TAKE CARE OF A GROVE	51
CITRUS ENEMIES AND REMEDIES	56
Physiological Troubles	56
Insects	58
Formulas for Solutions	59

ILLUSTRATIONS

	Page.
WELL CULTIVATED AND FERTILIZED GROVE 6 YEARS OLD —BAYAMON, PORTO RICO	9
BANANAS USED AS LOCAL WIND PROTECTION,—PORTO RICO EXPERIMENT STATION	15
VIEW IN CORNER OF NURSERY OF C. D. ABBEY — ISLE OF PINES	20
OLD NEGLECTED GROVE,—AFTER BEING FERTILIZED AND MULCHED —SANTIAGO DE LAS VEGAS	29
YOUNG GROVE, SHOWING EFFECT OF FERTILIZER ON GRASS AROUND THE TREES —ISLE OF PINES	33
ORANGE TREE 4 YEARS OLD, UNFERTILIZED.—BAYAMON, PORTO RICO	37
ORANGE TREE 4 YEARS OLD, FERTILIZED,—BAYAMON, PORTO RICO	38
ORANGE TREE FROM UNFERTILIZED PART OF FISKE GROVE —CEBALLOS, CUBA	40
ORANGE TREE FROM FERTILIZED PART OF FISKE GROVE,—CEBALLOS, CUBA	41
GRAPEFRUIT TREE 3 YEARS OLD WELL FERTILIZED, —ISLE OF PINES	43
GRAPEFRUIT TREE 4 YEARS OLD, UNFERTILIZED — LAS TUNAS, CUBA	45
GRAPEFRUIT TREE 4 YEARS OLD, FERTILIZED —LAS TUNAS, CUBA	46
ORANGE TREE 5 YEARS OLD, NOT FERTILIZED —LA GLORIA, CUBA	49
ORANGE TREE 5 YEARS OLD, FERTILIZED WITH WOOD ASHES AND COVER CROPS	50
VIEW OF GROVE FERTILIZED AND UNFERTILIZED,— LAS TUNAS, CUBA	53
VIEW OF 2 GROVES, FERTILIZED AND UNFERTILIZED. HERRADURA, CUBA	55
TREE NEARLY GIRDLED FROM GUMMOSIS	57

FOREWORD

In presenting this pamphlet to citrus growers we realize that it is impossible to treat the subject as fully as we should like to, but we hope that the condensed information given will be of value to the majority, and those who desire more specific information can obtain it from the following publications:

General Information on the Whole Subject.

“Citrus Fruits and Their Culture”, by Hume
Glen St. Mary Nurseries, Glen St. Mary, Fla.
Annual Reports, Fla. State Horticultural Society,
Published by the Secretary, Jacksonville, Fla.
Annual Reports, Cuban National Horticultural
Society, published by the Secretary, Havana, Cuba.

Propagation:

Circular No. 24, Cuba Experiment Station
Bulletin No. 4, Porto Rico Experiment Station

Insects and Diseases:

Bulletin No. 9, Cuba Experiment Station
Bulletin No. 10, Porto Rico Experiment Station
Bulletins Nos. 51, 56, 67, 88, 94, 97 and 106, Florida Experiment Station.

Practically all citrus growers in Cuba and Porto Rico know by experience that we are willing to furnish them whatever information we can relating to the subject of citrus growing. We should like to impress upon planters in other Islands in the West Indies that we stand ready at all times to do the same for them. Those who do not find the information they need in this pamphlet can feel free to write us. We charge nothing for the information we give.

COMMERCIAL IMPORTANCE

Citrus fruits are not natives of America, but they were undoubtedly brought over soon after Columbus' first voyage, and all the different kinds, including oranges, grapefruit, lemons and limes, are now found growing wild in all of the islands in the West Indies. It is known that oranges were shipped from some of the Islands to the United States even during the time of sailing ships and although the trade was unimportant from a commercial standpoint, the West Indies was then the chief source of supply for the United States.

According to Lelong, the first commercial grove in California was two acres planted near Los Angeles in 1841, but on account of undeveloped shipping facilities the Eastern States were not supplied from there until about 1880. From Florida some of the wild fruit was shipped commercially, as from the West Indies, but very little cultivated fruit was shipped before 1880. From this it is readily apparent that no great amount of oranges were consumed in North America previous to 1880, but according to statistics, in 1900, just 20 years later, over 10,000,000 boxes were shipped out of Cali-

Well cultivated and fertilized grove 6 years old, Rio Honda Fruit Co., Bayamon, Porto Rico.

ifornia and Florida. About that time, citrus culture properly speaking commenced in Porto Rico and Cuba, and some attempts were made a few years later in Jamaica. In the rest of the Islands the cultivation is yet limited, although a few limes are cultivated in some of the smaller British Islands. Most of the Islands continue shipping fruit from the wild trees, but that is becoming less remunerative as the markets are being supplied with cultivated fruit. The fruit from the wild trees is as a rule good, but the trees being seedlings, it is difficult to grade it uniformly. It is also difficult to market it in good condition, because it is frequently not firm enough to stand being packed and shipped long distances. Through this as well as through ignorant and careless handling, the West Indian oranges have become unfavorably known in the world's markets, but the experience with the cultivated orange in Cuba and Porto Rico has clearly shown that a fruit can be produced which will in all respects meet the market requirements. Enough acreage is now planted in the two Islands, and enough fruit is already being shipped to make the business of considerable commercial importance, and in all probability cultivation will soon be extended to other Islands also.

CHOOSING A LOCATION

In choosing a location in which to start a citrus grove in the West Indies the following are the main considerations. Exposure, soil, water, transportation, labor and social conditions. In considering the last factor it is presumed that the owner wishes to live on the property and be his own manager, but in that case his fitness as

a planter will of course have to be considered, and still another consideration is the capital necessary for the undertaking.

Exposure is a factor seldom considered, but experience has shown that it is a very important one. Cyclones and storms are difficult to guard against, because such disturbances may occur in places reported to be free from them, it is well however to look up the weather records, if such are available. What the prospective orange grower is more interested in is the ordinary wind, its velocity, general direction and the **lay of** the land in relation to forests or mountains which may serve as a wind protection.

Soils, roughly speaking, are composed of sand, clay and vegetable matter. If a soil does not stick together when wet it contains mostly sand and is called a sandy soil. If it is sticky when wet and becomes hard and lumpy in drying, it is called a clay soil. A peaty soil does the same, but peaty soils are practically unknown in the West Indies. The vegetable matter in a soil, usually called humus, can be recognized by its dark color when it is wet. When found in large deposits it is called peat, but we are interested only in that contained in the upper 6 to 12 inches of soil. It is that which makes the soil dark and distinguishes the top soil from the subsoil. A mixture of sand and clay, about six to one, is called a sandy loam. One part clay to four parts sand is termed a clay loam, and with diminishing sand the soil is said to be clay or heavy clay.

The subsoil is of no less importance than the top soil. The roots of a citrus tree extend under normal conditions ten to fifteen feet downwards and the subsoil should be

such as to allow the roots free passage. A layer of impervious rock or hardpan within six to ten feet of the surface is detrimental in all cases. The trees do not usually show the effect of such before six to ten years old, but sooner or later there will be troubles of various kinds which can not readily be remedied. The same may be said of water. It is not safe to plant citrus trees on land in which the water stands within 6 to 10 feet of the surface for any length of time. The most desirable for citrus fruits is a sandy loam, containing a fair amount of humus. A very sandy soil is less desirable because it does not retain moisture so well and fertilizers leach through it too readily. A heavy clay soil is undesirable because it retains too much water when wet, and in dry weather it becomes too dry, and cracks open unless it is cultivated constantly.

Water is a very important factor and the following points should be considered: The record of rainfall, not alone the annual, but the monthly and the weekly as well, should be looked up. In that way it will be possible to determine whether irrigation will be needed or not, and to what extent, also to what extent draining and ditching will be needed for carrying off surplus water. In case irrigation is needed it is well to find out if water can be obtained in large enough quantity without great expense.

Transportation or rather the absence of transportation, is the one great drawback in many places. After the production becomes large enough transportation always follows, but frequently the pioneers fail because transportation is too costly. Ocean freight is fairly cheap, but railroad freight is as a rule high in the West Indies.

Furthermore the wagon roads are often poor, and as citrus fruit is a bulky crop, distance from a seaport should be considered.

Labor in the West Indies is not very good for horticultural work, and a prospective grower should be prepared to teach rather than boss.

Social Conditions may not properly be classed as a horticultural problem, but it has proved to be fully as important as any of the above mentioned. The majority of those engaged in citrus culture, not alone in the West Indies, but in Florida and California as well, are in the business because they like it. They are engaged in it because it is pleasant outdoor work in a tropical climate and they are willing to sacrifice some of the monetary gain for the pleasure of living "close to nature," but churches, schools, good neighbours and good government are all necessary factors. The climate is often asked about by prospective settlers, but that is really of minor importance. The climate is favorable enough in practically all parts of the West Indies for Northern people to live comfortably.

The Man whether he be the owner or the manager, is of more importance to the success of the undertaking, than any of the foregoing factors. It is often the case he knows nothing about citrus growing at the beginning, and of course he must be prepared to pay for his experience. This however he can always obtain cheaper by working for another man than for himself, and it is a very good plan to begin working in well kept orange groves in different places. A year's experience of this kind will be worth more than several years in one small grove. There are few branches of agriculture as highly develop-

ed as citrus fruit growing, and even experienced planters find it advisable to visit other groves, from time to time in order to compare notes.

The Capital necessary for developing a citrus grove is usually under estimated. Some of the earlier contracts in Porto Rico and Cuba were taken for \$300.00 per acre, guaranteeing the trees planted and brought into bearing in 3 years. This the contracting Companies either did not do, or else they lost money. The price was later raised to \$500.00 per acre, and again in some cases to \$1,000.00 per acre, extending the time to 5 years. As a matter of fact, no general rules can be laid down, because the capital needed will depend upon all the factors discussed above. Practically all the failures coming under the writer's observation could be attributed to the one thing, lack of knowledge. A planter who knows how, can accomplish more with a small capital, even under unfavorable conditions, than the one who does not know can with a large capital under the best of conditions.

METHODS OF STARTING A GROVE

There is one general method employed by most planters in all places, and also a number of modifications employed by more progressive planters wherever local conditions demand such changes.

The Commercial Method consists in clearing the land if it contains trees and shrubs, after which it is plowed usually without removing the larger stumps. It is then staked off in absolutely straight lines for which purpose a surveyor's instrument is often used, especially for large tracts. The holes are then dug and the trees planted

Bananas used as local wind protection, Porto Rico Experiment Station.

immediately. The trees are usually secured from a commercial nursery, often from some neighbouring island or from Florida.

Modification No. 1: Windbreaks.—It was observed in Porto Rico shortly after commercial planting commenced, that trees protected from the wind grew much better than those exposed, and since then, systems of wind protection for young groves have been tried out in various places with excellent result. Where the land is covered with thick forest, a successful method is to clear off the native growth in strips 200 to 500 feet wide, leaving uncleared strips 20 to 25 feet between. Such strips of forest growth at right angles to the prevailing wind direction is of great protection to the young citrus trees. Many planters object to the system because the forest trees take nourishment from the citrus trees, and the wind belts serve as a harbour for insects, such as the parasol ant. These objections are frequently very serious and should of course be considered. A windbreak of large trees always impoverishes the soil for some distance away, but this can to some extent be prevented by deep plowing and subsoil breaking in which way the roots are cut below the feeding limit of young trees. If ants are troublesome it may be necessary to clear off the underbrush in the windbreaks, and keep the surface soil clean until the nests can be destroyed. Where there are no forest trees on the land or where there are but a few scattered trees of no value for windbreaks it has been found advisable to plant artificial windbreaks. This has been done and is being done with more or less success in different places, but usually not until the planter sees the necessity of it. That is, after the young orange trees

have been planted a year or more, and do not show a satisfactory growth. Unfortunately a great deal of damage has then been done, and a great deal of time has been lost. A better method is to plan windbreaks as part of the grove and plant belts of quick growing trees suitable to the locality a year or two years before planting the citrus trees.

Modification No. 2: Planting from Own Nursery.—This method is frequently followed by the more experienced planters, and there can be no question about the advisability of it, for those who know how. It is claimed that it takes too long to start a grove in this way, but while it does take longer to start, it does not necessarily take longer to produce bearing trees and usually it does not take as long.

In following this method the planter selects a piece of land on his property that is close to a water supply, and on which the soil is preferably a clayey one. This land is cleared, plowed and cultivated thoroughly in order to make the soil mellow and fine for seed bed. Rows are then laid off 4 ft. apart and furrows made and a fertilizer composed of about equal amounts of Nitrogen Potash and Phosphoric Acid is applied at the rate of at least $\frac{1}{2}$ ton per acre. This is mixed thoroughly with the soil after which the seeds are planted 2 to 3 inches apart. The seeds should be covered with about an inch of soil and a light mulch of grass should be spread over, covering the rows. The ground should then be irrigated or watered by hand, so as to keep it constantly moist. The seeds will germinate in a few days and in about two months the plants can be thinned out, leaving the most vigorous at a distance of about 3 ft. apart. By constant

cultivation or mulching, together with irrigation when needed, and a sprinkle of Nitrate of Soda, the plants will have attained the size of a lead pencil in about 5 months from planting, and should then be budded. The buds should be tied to stakes in order to grow straight and when 2 ft. high the top should be nipped off to make the trees branch. In about 18 months from the time the seeds were planted these trees will be as large as the average nursery tree sold commercially today. At the time the nursery is planted the land should be prepared for the grove. It should be cleared, plowed and planted to legumes. The windbreaks should be planted and ants nests destroyed. Quickly maturing legumes such as cow-peas may be plowed under in 6 months and another crop planted, but the last crop should be plowed under about 4 months before the trees are to be set out. Then the land should be staked off, placing a stake where each tree is to be planted, and 3 to 4 rows of pidgeon peas or other similar plants should be planted between each tree row. The holes for the trees should be dug at least 4 ft. broad and 2 ft. deep and should again be filled up with vegetable matter and top soil. When the nursery trees are large enough they may be transplanted without being materially checked in growth. This is accomplished by digging a trench on each side of a row not closer than 12 inches to the trunks of the trees. Then by starting at one end of the row the trees can be removed one by one by digging horizontally at a depth of 12 to 15 inches, leaving a ball of earth around each tree two feet square and 12 to 15 inches deep. If the soil is clayey it will adhere when moist, and the tree can be transported to the grove if the ball is wrapped in burlap. The tree

needs no pruning before planting if it has been properly pruned in the nursery, and if planted with reasonable care it will in three years reach the size of a 5 year old tree planted in the old way.

Modification No. 3: Planting Seedlings and Budding in the Field.—This method can not be practiced under all conditions, and it has no real merit except in a case where a planter wishes to start a grove cheaply. In a forest all the larger trees may be taken out, leaving all the brush and small growth. The lines can then be staked out and a place 4 ft. square prepared at each stake. A few seeds can be planted there and some mulch spread over after which it can be left to take care of itself, except the surplus plants should be pulled up after a few months, leaving only one in a place and of course vines and tall grass should be cut off occasionally so as not to smother the young trees. The trees will not grow as fast as in a nursery, but they may be budded when about two years old, and will have the advantage over transplanted trees that they are never planted too deeply, also they will be less liable to gum diseases. Of course after the trees are budded they must be mulched or cultivated, and a space several feet wide must be kept clean around each one.

VARIETIES OF STOCK AND BUD

Citrus fruits can not usually be reproduced true to seed, and the commercial growers plant nothing but budded trees. A budded tree consists of two parts, the lower part originating from the seed which is called the stock, and the upper part, originating from the bark of another tree which is called the bud or the scion.

View in corner of nursery of Mr. C. D. Abbey, Santa Ana, Isle of Pines,
showing fertilized and unfertilized rows.

In selecting seeds for nursery stock the following points should be considered. The soil on which the grove is to be located, and the variety of fruit to be grown. All of the different species and varieties of the citrus family can be budded one on to another, but some succeed better on one stock than on another. The same is true of soil, all kinds of citrus trees may grow on any particular soil, but on account of different root systems and different feeding habits, some are much better adapted to certain soils than others.

The Sour Orange seems to be more universally adapted than the rough lemon, the grapefruit, etc., but it is less vigorous than these, and in light soil the growth is often slow for commercial purposes. During the first few years the sour orange makes a slower growth than the rough lemon, grapefruit or shaddock, even on heavy or medium heavy soil, but after the root system becomes well developed, the difference is not very great. The root system is deep, for which reason it is often recommended for dry soil, which is quite right, provided the soil water is not too far down, but even if the tree can reach water when it becomes old enough, it should be well remembered that moisture close to the surface is necessary during the first few years. Where the sour orange can be especially recommended is on heavy moist land, because it is less subject to gumming than the other stocks.

The sour orange is a good stock for the grapefruit and the common orange, although it can not be recommended for the navel. Neither should it be used for satsuma nor kumquat.

The Rough Lemon is more vigorous than the sour orange and is especially desirable for light soil poor in

plant food. It should never be planted on moist land, because it is very much subject to gumming. The root system is shallow and in some soils the tap root is entirely absent.

The rough lemon is suitable as a budding stock for any of the commercial citrus fruits. It induces early bearing because of its vigour, but the first couple of crops of fruit are usually large, coarse and juiceless. It produces a larger tree of any variety budded on to it, than does the sour orange during the first 4 to 5 years, but later there is almost no difference in size, unless the soil is very poor. The rough lemon is especially well suited as a stock on which to bud King, Tangerine and Kumquat varieties. When grapefruit and orange varieties are budded on to it the tree seems to be especially subject to gumming on heavy soil.

The Grapefruit and the Shaddock are very good budding stocks, the shaddock especially being very vigorous. The root system is deeper than that of the rough lemon, and it is suitable for the same class of soil as that. It is not much more subject to gumming than the sour orange, and less so than the rough lemon, and as it is suitable as a stock for all kinds of citrus fruits it is worthy of general use.

Commercial Varieties of Oranges as advertised in nursery catalogues are almost legion and practically all of them have been planted in the West Indies, the names are still retained but there is considerable variation, which is however of minor practical importance. The question is not what an orange is called, but what the quality and appearance is. The most desirable size is $2\frac{1}{8}$ inches in diameter, packing 176 to the box, form

round, color bright, peel thin smooth and silky. The flesh should be a rich orange color with but little fiber and few seeds. The juice should be plentiful, subacid and distinct orange flavor. The time of ripening is of no less importance. In October and November a good orange sells well, later the competition becomes great and those who can not have extra early oranges should plant late varieties, and fertilize with a large percentage of Potash so as to have them mature after March. The home consumption is becoming to be of some importance in some of the Islands, and for this purpose the Navel is one of the best on account of its large size and its sweetness. The Navel can not however, be successfully produced except on medium heavy soil and with irrigation.

Varieties of Grapefruit may be summarized like the oranges. A good size is 4 to 5 inches in diameter, color light yellow, form round to flat, not pear shaped, peel $\frac{1}{16}$ inches thick, bright and smooth. The flesh should be a bright yellow in color with not too much fiber or rag and few seeds. The juice should be plentiful with a flavor of bitter and subacid. A number of varieties described in nursery catalogues conform to this standard, and it may be said that any of the well known good varieties are desirable for planting in the West Indies. The soil, cultivation and especially the fertilization is frequently of more importance than the variety. For instance in Cuba the fruit from young trees on some of the heavier soils is large, coarse and pithy unless it is properly fertilized. It is probable that certain varieties are specially adapted to certain soils, but that has not been thoroughly worked out yet. Seedling trees of which the origin is unknown have been observed to yield better

fruit than well known Florida varieties which would indicate that varieties better adapted to the West Indies may be found. That however should not induce planters to buy native varieties unless they know the origin, because there are a number of so called grapefruits in the West Indies that are nothing more than improved types of the shaddock.

HOW TO PLANT A TREE

There are seemingly more mistakes made in planting trees than in any other operation connected with the business. If it were not for that this chapter would not have to be included. The following directions if followed will obviate any mistakes in planting. Dig the holes large and deep. Fill the soil in again with the top soil and vegetable matter in the bottom and heap it up. Leave it for some weeks until perfectly settled. When planting the tree without a ball of earth use a planting board, that is, a board 6 to 8 feet long with a notch in the middle and a block under each end. Make an excavation in the previously prepared hole. Place the board over this with the blocks resting on the level ground. Insert the tree in the notch with the roots close to the board. Then fill the soil in around the roots and firm it down well, using plenty of water. In this way the tree will stand as high above the level of the soil as the thickness of the blocks under the board. In sandy soil these blocks may be but 3 to 4 inches thick according to whether the soil is very wet or not. In soft and very wet soil the trees settle more than in drier soil. In clay soil when planting in rainy weather the blocks should be 6 inches thick, because the trees will settle a great deal

and they should stand with the crown roots about 2 inches above the level when fully settled.

In heavy clay soil or in land with impervious subsoil in which citrus trees are frequently planted, although it is unsuitable, the writer has observed standing water in the holes all through a rainy season. Trees planted under such conditions start gumming almost immediately, and frequently the roots decay. This can be remedied by subsoil draining or even surface draining, connecting each hole with a drainage ditch.

The Distance Apart at which trees should be planted is frequently a subject for discussion. The tendency is to crowd too many trees on to an acre, because they look so small when planted and it is hard to realize that they will grow so large. In general 25 x 25 feet apart (70 trees per acre) is the most suitable distance. Some have planted this way with a tree between each four trees making the number 126 per acre, but those extra trees had to be taken out before they bore fruit. Others have planted 20 x 20 (108 trees per acre) which is excusable on light soil, but on the heavier lands like eastern Cuba, and the banana soils of Jamaica, 70 trees per acre is enough.

HOW THE WILD TREES GROW

Citrus trees grow naturally in the West Indies without any care, which is fully attested by the great number of seedling trees found in all of the Islands. Why is it then that a planted grove requires so much care? It is because in the average grove the natural conditions have been lost sight of, and with an artificial start the artificial conditions must be kept up.

Nature's Way is to drop seeds promiscuously wherever they happen to fall, but only those dropping in favorable spots in the forest or in well protected fence corners survive germination, and after germination only the most vigorous plants overcome the obstacles to which the budded trees are subject. Unfortunately it is not possible for the planter to exercise such rigorous selection, but it is practical in many instances to copy nature's methods of growing trees. The tree growing in the cool moist forest atmosphere is not subject to scale insects, because those conditions are favourable to the growth of fungi, which will keep the scale in check. The surrounding trees and shrubs take up a great deal of nourishment and rob the citrus tree of food, for which reason it grows slowly. On the other hand those surrounding trees and shrubs shade the soil, keeping it cool and moist. The mulch or leaf mold protects the soil from baking by the sun, beating by the rains and cracking from drought.

Imitating Nature is the cheapest method of cultivation, and considering that a tree planted is capital invested it pays well to take advantage of it. In the commercial method of starting a grove spraying is one of the first operations, and in many of the earlier groves more money was spent on that, than on all other work combined. The more advanced planters who have practical experience and theoretical knowledge do not "lay awake nights" studying up new cures for tree diseases. They plan their work in such a way as to keep the trees healthy and that is the keynote to success.

The grove is in the first place started with windbreaks around and smaller plants between the rows. This is imitating forest conditions and the trees will not have

to be sprayed for scale. The next question is how to keep the soil moist. In this it is also possible to imitate nature, but it is not always practical. The air and soil in the forest are humid because of the dense shade and the leaf mold covering the ground but in the grove the shade must not be dense, and enough mulch to effectually protect the soil can seldom be obtained. The remedy is continual cultivation during the dry season, but experience has shown that that method does not fill all the requirements. Irrigation is frequently necessary and in many localities it is one of the best paying investments. Nature also teaches a lesson in feeding plants and in the case of citrus trees the lesson is a most valuable one. A self-planted seedling tree growing in uncultivated soil does not suffer from die-back, gummosis, foot-rot, etc. It takes up no more food than it can readily digest. It does not grow fast, but nature is not in a hurry. The fruit may be too soft to ship or too sweet to suit the palate of some people, but it suits the mocking bird and the wild parrot and nature is satisfied.

The budded tree planted by hand and forced into quick growth by constant cultivation of the soil is under many disadvantages. If the soil is rich enough to produce a tree it is usually too rich in nitrogen to produce a healthy tree. This may cause no harm if the soil is not cultivated, but in that case a young tree would not grow fast enough to suit commercial demands, and it would not survive because of the artificial conditions under which it is placed. In cultivating virgin soil the nitrogen is liberated faster than the other plant foods and the tree becomes unhealthy from improper food. If it survives and bears fruit it will usually not be firm enough

to ship and frequently too insipid to be marketable. Therefore when interfering with nature in trying to produce quick growth and marketable fruit, the planter must be prepared to feed his trees according to requirement.

CULTURAL METHODS

The proper method of treating a grove is a local problem. General directions can not be followed explicitly, but by the use of common sense a planter may adapt any of the usual cultural methods to suit his special conditions. In the young grove the main problems are how to keep the air and the soil cool and moist during the dry season, and how to keep the soil supplied with the proper amount of plant food.

Local Wind Protection is, as previously, stated of great benefit to young trees because in that way it is possible to duplicate forest conditions. Many different plants may be used, such as pidgeon peas, corn, castor beans, bananas and even sugar cane. One or more rows of such plants between each row of citrus trees will serve as an effective windbreak and soil shade, but like forest trees they will rob the citrus trees of plant food. This must be guarded against by plowing deeply close to the plants to prevent the roots from spreading too far. Such plants may also serve as a harbor for insects, such as leaf eating beetles or ants and if so the method is of course impractical.

Mulching consists of covering the soil with a layer of dry vegetation, thick enough to prevent evaporation of soil moisture. It is the most natural method of preserving the moisture and preventing the weeds from growing, but it is not always practical. A mulch to be ef-

Old neglected grove, Santiago de las Vegas, Cuba, recovering after
being fertilized and mulched.

fective must be thick enough to keep the soil from drying out during the dry season and such large amount of grass or any other kind of plants can seldom be obtained without too great cost. Another drawback is the danger of fire in a grove mulched all over. These objections can be overcome in young groves by mulching around the trees only and cultivating the middles. In this way less mulch is needed and the danger of fire is lessened. In mulching trees the roots form in the surface soil and for that reason it is necessary to remove the mulch and cultivate the ground during the rainy season in order to keep the root system from becoming too shallow or else to continue mulching without interruption during the life of the tree.

Cultivation is a broad term denoting all work in connection with weeding and stirring the soil. In young groves it is strictly necessary to keep the weeds away from the tree and unless mulching is used it is necessary to work the soil with some instrument in order to prevent the moisture from evaporating too quickly. This is on the same theory as mulching. The moisture rises to the surface in compact soil like the oil in a lamp wick and from there it evaporates by the heat of the sun. In order to prevent this the surface is broken and pulverized, the moisture then rises to the pulverized layer and no further. To receive the practical benefit from cultivation a planter must plow his soil just before the rainy season is over and harrow it before it dries out. After that he must harrow or cultivate 2 to 4 inches deep after every sprinkle of rain, or frequently enough to prevent the pulverized layer from coming in solid contact with the underlying uncultivated portion. Many planters know

this and practice it, but they frequently start plowing and harrowing too late, trying to save the moisture after it has already escaped.

Irrigation is usually of great benefit and frequently it is absolutely necessary for several months in the winter. In some soils it is possible to preserve enough moisture to keep the trees in growing condition by constant cultivation, but a great many soils become too dry and the bloom which should form in February does not appear before June. With irrigation not only the time of blooming may be controlled but many insect pests as well.

The method of irrigating will depend on local conditions. Water can seldom be obtained from streams by gravitation, and it must therefore be pumped from natural supplies or from wells. The cost of this will vary according to the height the water has to be lifted and the machinery used. If there is an unlimited supply of water within 15 to 20 feet of the surface the problem is a simple one because under such conditions a 4 inch centrifugal pump driven by a 4 horse power engine, supplying about 300 gallons per minute would be enough for a ten acre grove. Where the lift is 100 to 200 feet like in some groves in Cuba, the problem becomes more complicated and the cost very great. In connection with pumping plants it is usually necessary to have some kind of storage tank unless the capacity of the plant is large, because the main thing is to be able to supply large volumes of water when needed. A windmill can not be relied on where much water is needed, but for a small supply a windmill in connection with a storage reservoir may sometimes be satisfactory. The water may

be distributed in open ditches or by means of pipes. The open ditches can be used only where the soil is firm enough to prevent too great loss by seepage. The lay of the land must also be considered. Too much fall will cause washing, on the other hand there must be enough fall for a rapid flow of water. A pipe system may be confined to a main line only, from which the water can be led out through valves into short lateral ditches between the rows of trees. Or the system may be more elaborate with pipes all through the field and stand-pipes for overhead springling, which latter method has much to recommend it and it is one that will undoubtedly be used extensively in the future. (*)

Fertilizing (**) is undoubtedly the most important of all the problems confronting the orange grower. It is unquestionable that moisture, physical condition of the soil, fungus and insect pests, etc., are important problems, but the **underlying** cause of the most serious troubles in citrus groves is to be found in the chemical make up of the soil. A citrus tree like all other plants needs a number of chemical elements for wood making and fruit producing. As these elements are taken up and digested by the plant they are called plant foods which is a more intelligible term than fertilizer, because anything that a fertilizer may contain beside plant food is valueless as such. Many of the elements entering into a plant are found in abundance in all soils and there are but four that need to be considered by the practical

(*) For information on irrigation see Reports Fla. State Horticultural Society 1910 and Cuban Nat. Horticultural Society 1911.

(**) For more information on fertilizers our book "Fertilizer Guide" can be obtained free of charge.

Young Grove Isle of Pines, showing heavy growth of grass around the trees where fertilizer was applied and none in the unfertilized soil between the trees.

planter. These are Lime, Potash, Phosphorous and Nitrogen.

Lime is usually present in the soil in fairly large quantity, and it is seldom necessary to apply it as a fertilizer, because all fertilizers containing phosphate contain enough lime to satisfy the needs of the average plant. But lime is also used to correct acidity in sour soils and is often needed for that purpose. A soil that is sour is not suitable for citrus trees and such soils are always benefited by 1000 pounds or more of burnt lime or several tons of finely ground limestone according to the amount of acid it contains.

Potash is the one plantfood of especial importance to citrus trees, not alone to the growth of the trees, but more so to the health and especially to the quality of the fruit. Potash is present in the different soils in varying quantities. For instance sandy soils are usually poor in potash and clayey soils contain comparatively large quantities, but the actual amount present is not what the planter is so much interested in because often the clay soils showing large quantities when analysed are most benefitted by an application of potash salts. This is due to the fact that the potash in the soil is usually present in insoluble forms which are not available to cultivated plants. The potash used for fertilizers comes from the potash mines in Germany. It is found there in several forms and sold as Sulphate of Potash, Sulphate of Potash Magnesia, Kainit, etc.

Phosphorous or phosphoric acid is another plant food. It is used for fertilizer in the form of phosphate of lime, that is, a combination of phosphoric acid and lime. It is sold as acid phosphate or superphosphate also as double

acid phosphate and as basic slag, all of which contain various amounts of phosphoric acid. Bone meal is also a phosphate fertilizer and so are the various bat guanos.

Nitrogen or ammonia is the third element contained in fertilizers. It is present in all organic matter such as the vegetable matter in the soil, and also in the air, from which latter source leguminous plants can supply themselves. New land recently cleared from forest and containing much vegetable matter is rich in nitrogen, and frequently trees planted on such land suffer from gumming, die-back, etc., because they take up more of that than of potash and phosphoric acid. For the same reason, stable manure can not be used in large quantities for citrus trees. For fertilizing purposes, nitrogen is sold in the forms of nitrate of soda, sulphate of ammonia, dried blood and the various slaughter house and fish scrap offals as well as nitrate of lime which is the new product made where nitrogen is obtained from the air.

Cover Crops is a name applied to plants grown in the grove for the purpose of benefiting the soil. A cover crop may or may not be a fertilizer according to what kind of plant it is. Common weeds such as grasses usually found in uncultivated groves do not add plant food to the soil. They feed on whatever plant foods there are in the soil and when they decay there is nothing more nor less left than what there was before. There is a gain however because the humus formed improves the physical condition of the soil and the plant foods in dead vegetation are in a form readily available for other plants.

Leguminous plants, such as beans or peas do not alone improve the physical condition of the soil but they also add something to it. They feed on nitrogen, taking it

directly from the air, and for this reason they are called green manure crops. In citrus groves a number of legumes are used such as cowpeas, velvet-beans, lablab beans, canavalia, etc. The practice is to plant a crop between the trees at the beginning of the rainy season, and stop all cultivation as long as the rains last, except keeping the vines from climbing up the trees. In this way the soil is greatly benefited by the large amount of humus and the fertilizer bill is greatly reduced, because it is necessary to apply only potash and phosphorous, and these ingredients are cheap compared to the nitrogen which is in this case supplied by the cover crops.

INFLUENCE OF FERTILIZERS ON THE TREES

In sandy soils there is no question about fertilizers being needed, because experience has shown very thoroughly that a citrus tree will not make satisfactory growth without it. Those who start groves on heavier soils such as clay and clay loam often do so because they figure that they will not have to use fertilizers. This is unfortunately a mistake, and the sooner a planter realizes the fact the better it is for him. It is undeniable that citrus trees often grow faster on such soils without fertilizer than on lighter soils with fertilizer, but too frequently the loss is very great from gumming. On light soil a tree will begin to bear when three years old if properly fertilized, but on many of the heavier soils trees 5 to 6 years old do not even bloom. Such soils do respond to fertilizers however, and a medium heavy soil if well drained can be handled very satisfactorily and usually cheaper than a light sandy soil. The thing to remember is that virgin soil containing

Orange tree 4 years old. Average tree in unfertilized plat,
Dr. H. C. Leonhart's grove, Bayamon, Porto Rico.

Orange tree 4 years old. Average tree in fertilized plat,
Dr. H. C. Leonhart's grove, Bayamon, Porto Rico.

much humus is rich in available nitrogen and unless potash and phosphate is supplied artificially the growth is liable to go to wood instead of fruit. Also the growth is liable to be unhealthy, the wood soft and the tree attacked with gummosis.

What fertilizing will do in different soils and under varying conditions is best illustrated by photos, and those shown in this pamphlet are absolutely true. Any planter can duplicate the experiments by trying for himself, using the formulas recommended here.

INFLUENCE OF FERTILIZERS ON THE FRUIT

While it is fairly well known what fertilizers will do in regard to tree growth and fruit production, the tremendous influence on the quality of the fruit and time of ripening is not generally recognized. This is probably due to the fact that experiments have been conducted principally on light soil and the influence seems to be more noticeable on heavy soil.

Following we give some results: At the exhibition held by the Cuban National Horticultural Society, January 1909, some oranges were exhibited by Mr. D. H. Howell of Ceiba Mocha, which were commented on as being insipid and **stale**. It was decided to try and remedy this and in June, 4 lbs. high grade sulphate of potash was applied per tree on 40 trees. The soil was a red clay, fairly deep, underlaid with coral rock. The grove was not well cared for that summer on account of the owner's absence, and it was found necessary to pick the fruit early to prevent it from being stolen. The data was taken in December and several people were asked to give their opinion on the quality of the fruit. All who tasted it

Orange tree from unfertilized part of Fiske grove, Ceballos, Cuba. This part of the grove bore much less fruit than the fertilized part, and it was over ripe in January.

Orange tree from fertilized part Fiske grove, Cebalos, Cuba. This part of the grove was well loaded with fruit, and the fruit was not nearly ripe in January.

remarked on the great difference in flavor and acidity, the unfertilized fruit being even at that early date insipid and inclined to be stale, while the fertilized fruit was of good flavor but decidedly acid, indicating that it lacked 4 to 8 weeks in being ripe.

In 1910 part of a five hundred acre grove at Ceballos, Cuba, was fertilized with a regular fruit fertilizer 3% Nitrogen 7% Phosphoric Acid and 12% Potash and the results were noted in January 1911 by the writer in company with Mr. Gustschow, the manager of the grove. At that time nearly all the fruit had been picked from the unfertilized part and what was left was over ripe, while on the fertilized part the fruit was still hanging on the trees and not too ripe for hanging 3 to 4 weeks longer.

In April 1910 an experiment was started in the grove of the Cuban Development Co., Victoria de las Tunas, Cuba, and the result noted January 1911. There were several plots with various fertilizer combinations from which the results were not very clear cut, but the difference in flavor and acidity due to different stages of ripeness was observed to be as in the above mentioned experiments although less pronounced.

In Porto Rico the same results were obtained in the Southern Cross grove one year, but it was seemingly not duplicated the next year. From Dr. H. C. Leonhardt's grove, Bayamon, Porto Rico, ten boxes of fruit from plats fertilized with different combinations of fertilizer were shipped to New York and from there to Havana in January 1911. This fruit was carefully noted and showed great variation in flavor and shipping quality due to fertilizing.

More observations might be given but this will serve

Grapefruit tree 3 years old well fertilized. Mr. H. A. Brigg's grove,
Santa Fe, Isle of Pines.

to show that a fruit fertilizer like formula No. 2 or No. 3 in connection with leguminous cover crops has a very beneficial influence. A fertilizer high in potash content or even potash alone where nothing else is required seems to have a decided retarding influence on citrus fruit. This is very important in the West Indies where it is advisable to hold the fruit until the bulk of the Florida and California fruit has been marketed. Of course the flavor and carrying quality is improved also, but this may, as shown in some of the experiments cited, be due to the difference in time of ripening. A fruit when just ripe is of good flavor, but when it is over ripe it becomes stale and loses its good flavor.

Also a fruit may be just ripe when picked, but if it is mussy when it reaches the market, what is the use of shipping it. On asking a number of fruit venders "Why do you handle the California Navel when there are other better quality oranges in the market?", The substance of their answers was because it looks good and it keeps. That is the keynote of the marketing problem, it must look good and keep well after reaching the market, and proper fertilizing is the only thing that will accomplish it.

FERTILIZER FORMULAS

The following formulas will be found referred to in the text by numbers.

Formula No. 1 for young trees on light soil:

3% Nitrogen

5% Phosphoric Acid

5% Potash.

This can be obtained from reputable fertilizer dealers,

or it can be mixed by the planter himself from the following materials:

400 lbs.	Nitrate of Soda	containing 15%	Nitrogen
500 "	Basic Slag	" 20%	Phos. Acid
200 "	Sulphate of Potash	" 50%	Potash
900 "	Dry earth as filler.		
<hr/>			
2000 lbs.			

OR

300 lbs.	Sulphate of Ammonia	containing 20%	Nitrogen
250 "	Acid Phosphate	" 40%	Phos. Acid
375 "	Sulphate of Potash Magnesia	" 27%	Potash
1075 "	Dry earth as filler.		

OR

430 lbs.	Dried Blood	containing 14%	Nitrogen
670 "	Acid Phosphate	" 15%	Phos. Acid
200 "	Muriate of Potash	" 50%	Potash
700 "	Dry earth as filler.		
<hr/>			
2000 lbs.			

Neither the dried blood nor the muriate of potash as used in the last mixture is considered well suited for citrus fertilizer. Dried blood and other organic forms should not be used where the trees are subject to gummosis. Muriate of potash may be used for young trees and probably for bearing trees as well, but the influence of chlorine on the fruit has not yet been determined.

Any of the above mixtures contain approximately 3% nitrogen, 5% phosphoric acid and 5% potash when mixed with filler to bring the weight up to 2000 lbs. Filler is not strictly necessary especially if the fertilizer is used shortly after mixing, but in that case corresponding smaller amounts should be applied. For instance of the

Grapefruit tree 4 years old. Average tree from unfertilized part of grove belonging to the Cuban Development Co., Las Tunas, Cuba.

Grapefruit tree 4 years old. Average tree from fertilized part of grove
belonging to the Cuban Development Co., Las Tunas, Cuba.

second mixture without filler 1 to 3 lbs. should be applied per year to trees under six years old instead of 2 to 6 lbs. of the mixture containing filler.

Formula No. 2, for bearing trees on light soil:

3% Nitrogen
6% Phosphoric Acid
12% Potash.

If a formula like this can not be obtained it may be made by mixing the following materials:

400 lbs. Nitrate of Soda	containing 15%	Nitrogen
600 „ Basic Slag	„ 20%	Phos. Acid
480 „ Sulphate of Potash	„ 50%	Potash
520 „ Dry earth as filler.		

2000 lbs.

or

300 lbs. Sulphate of Ammonia	containing 20%	Nitrogen
300 „ Acid Phosphate	„ 40%	Phos. Acid
900 „ Sulphate of Potash Magnesia	„ 27%	Potash
500 „ Dry earth as filler.		

2000 lbs.

This like the former may be used with or without filler. With filler a fair application is 9 lbs. the fourth year, 12 lbs. the 5th. year, 15 lbs. the 6th. year, and up to 30 lbs. or more later according to the amount of fruit produced.

Formula No. 3 for trees on land containing considerable humus and nitrogen, the same amounts of potash and phosphoric acid as used in formula Nos. 1 and 2 may be used in this mixture but half or all of the nitrogen can be omitted according to the character of

the soil. A rich dark loam or a soil producing leguminous crops continually for any length of time will need but little nitrogen and in many cases none at all. Frequently such soils contain so much nitrogen as to cause gummosis in young trees and potash and phosphate must be applied to rectify the trouble.

Formula No. 4 for leguminous plants: Leguminous plants can supply themselves with nitrogen from the air by means of bacteria which live in **nodules** on the roots, but as they can obtain no nitrogen before such nodules have formed it may be necessary to apply some for the first crop on poor sandy soils. A good formula for such is as follows:

250 lbs.	Nitrate of Soda
600 „	Basic Slag
300 „	Muriate of Potash
850 „	Dry earth as filler.

2000 lbs.

or

200 lbs.	Sulphate of Ammonia
300 „	Double Acid Phosphate
300 „	Muriate or Sulphate of Potash
1200 „	Dry earth as filler.

2000 lbs.

This can be applied to three acres either with the filler or without. That is, the application should be about 633 lbs. per acre of the last formula with filler, or 266 lbs. without filler. For the second and successive crops the nitrogen may be omitted entirely.

HOW TO TAKE CARE OF THE GROVE

The young trees should preferably be mulched as soon as planted. If no mulch is available the soil should be

Orange tree 5 years old not fertilized. Average tree in unfertilized part,
Mr. John Boston's grove, La Gloria, Cuba.

Orange tree 5 years old. Average tree from part of grove fertilized with wood ashes and cover crops. (Wood ashes contain potash and lime).

stirred and kept well pulverized on top in order to prevent drying out. In light soil an application of fertilizer should be made two or three months after planting, using $\frac{1}{2}$ lb. per tree of formula No. 1. This should be scattered around the trees not closer than 6 inches to the trunk, and the application should be repeated every three months during the first year. If plants are used for wind protection they should be cultivated and if cover crops are planted, strips on each side of the citrus rows or circles around each tree, should be kept clean. At the end of the rainy season the cover crops should be cut and used as a mulch immediately around the trees, and the middles should be plowed and cultivated often enough to prevent the moisture from escaping.

During the first three years this treatment may be kept up. Fertilizers like formula No. 1 should be applied from two to four times per year gradually increasing the amount from 2 to 6 lbs. and gradually increasing the width of the circle and distance from the trunk of the tree. On light soil the first cover crop should be fertilized with 300 lbs. to 600 lbs. of formula No. 4, and if a large portion of the crop is used for mulching the trees less fertilizer will be needed for them. With heavy growth of cover crops plowed in every year, formula No. 3 should be used for the trees after the second year. The kind of formula to use can largely be determined by the color of the leaves, a dark green color and vigorous growth indicates an abundance of nitrogen, whereas a yellowish color indicates lack of nitrogen. On heavier soil where ordinarily no fertilizer is needed during the first 3 to 4 years, formula No. 3 can profitably be used if the trees show a tendency towards gumming.

View of grove, Victoria de las Tunas, Cuba. Row to the right not fertilized.
Row to left fertilized.

Bearing trees should receive essentially the same treatment as young trees during the first few years, with the exception that fertilizer formula No. 2 should be used and the quantity should be gradually increased reaching 15 lbs. per tree on light soil by the time the tree is 6 years old. On soil naturally rich in vegetable matter or one producing large crops of legumes, formula No. 3 should be used.

It is usually understood that a citrus tree will begin to bear when three years old, and it will do so if treated as here recommended. If however the tree is neglected a planter is lucky if he can harvest fruit in quantity at the end of 5 years.

The fruit from a three year old tree may be of good quality, but it usually is not so unless the tree has been properly fertilized. The quality depends somewhat on the variety but most grapefruit varieties are apt to be pearshaped and pithy and orange varieties such as the navel for instance are seldom at their best before the second or third or fourth crop although with irrigation and potash fertilizer they may be all right the first crop. If the tree is healthy and vigorous it will be able to carry all the fruit it sets. If it is in any way injured, it may set more fruit than it can mature without injury to itself, and in such cases the fruit should of course be thinned out or all taken off.

If local wind protection has been used it should be cut out when the trees reach the age of 4 to 5 years, because then they should be large enough to protect one another. After 6 years the method of cultivation may in some cases be changed. Large trees of that age will have the root system extending all through the soil and the two

View of two groves, Herradura, Cuba. To right poorly cultivated and not fertilized.
To left well cultivated and fertilized.

essential needs are water and plant food. If irrigation is used the soil can be kept moist without cultivating and if the overhead system is used it will not be packed from the application of water. Therefore all that is necessary is to irrigate when needed and apply the fertilizer broadcast without working it into the soil. The weeds or leguminous crops that can be made to grow under the trees should be cut frequently and left on the ground preferably each time after an application of fertilizer, so as to serve as a temporary cover for that.

Of course this system is practical only in connection with irrigation or with mulching where plenty of that can be had. Ordinarily cultivation must be continued in order to keep the soil in good condition. Fertilizers should be applied as before with the exception that the quantity should be gradually increased according to the crop taken off. Thirty to fifty pounds per tree per year scattered broadcast in 2 to 4 applications will pay under most conditions.

CITRUS ENEMIES AND REMEDIES

In the natural state the citrus tree and fruit is not very much subject to diseases, but under the extremely artificial conditions prevailing in the majority of planted groves, remedies have to be used in order to make the business a success. There are three distinct troubles to combat, viz: physiological, insect and fungus.

Physiological Troubles are usually caused by unfavorable soil conditons, such as heavy soil that is too wet or too rich in nitrogen. The symptoms are an outflow of gum from any part of the tree, usually from the lower part of the trunk on young trees, and often from crotches

Tree nearly girdled from gummosis. Note method of inarching watersprouts to keep up sap circulation. Gummosis can often be remedied by an application of potash.

of the limbs on older trees. It is called gummosis (*) or gumming, or on older trees, when occurring at the base and on the crown roots it is called foot-rot.

The remedies are drainage, an application of potash and cutting away the dead bark and painting the wound with some antiseptic like a very weak solution of carbolic acid in water, or even Bordeaux mixture. Carbolineum is also used and gives good results, especially when mixed with a soap solution. The pure carbolineum may burn the live bark.

Insects are treated according to the kind of damage they do. Some are biting, that is they eat the foliage and they can be killed with poisons. Others are sucking, that is they feed on the juices of the tree or the fruit, and they must be killed with contact remedies. That is soap or other sticky solutions that stop up the breathing pores.

Of leaf-eating insects there are various beetles all of which are more or less difficult to kill. Often the only remedy is shaking the tree and collecting the beetles on a sheet underneath. Of sprays, the only practical one seems to be Arsenate of Lead. That can be bought ready prepared and mixed with water according to directions. The greatest damage is caused by beetles in the larval stage. The larva, which is commonly referred to as the grub, lives in the ground and feeds upon the roots. It is difficult to kill, because anything that will kill it, will

(*) Since this was written Prof. Fawcett of the Florida Experiment Station has found a fungus connected with gummosis and it may in the future be classed among fungus diseases. That however does not alter the treatment here recommended, but pruning tools should be cleaned and washed in a 5% solution of carbolic acid before using on healthy trees.

kill the tree also. Sometimes digging out is resorted to and again repellants are used, such as the Kerosene Carbolic Acid Emulsion used for scale or Carbolineum. Kainit spread around the tree is also of some use, but the best remedy is plenty of fertilizer and irrigation, which causes the tree to outgrow the damage done by the grub.

Of sucking Insects various scales, a white fly and two species of red spider attack the tree and a rust mite attacks the fruit. Some of the scale insects like the Wax Scale and Hemispherical Scale are called soft scales, and they are readily kept in check by spraying with whale oil soap or kerosene emulsions. Other scales having a hard covering are much more difficult to destroy. Those giving the greatest trouble are the Purple Scale, the White or Orange Snow Scale and the Florida Red Scale. All of these are rather difficult to kill, and the Purple Scale especially so. An ounce of preventive is in this case worth more than the proverbial pound of cure, and windbreaks are the preventive. If it becomes necessary to spray, several mixtures may be used, but the following recommended in the Porto Rico Experiment Station Bulletin No. 10, is probably as effective as any and practical enough for any planter to use.

Kerosene and Crude Carbolic Acid Emulsion:

Stock Solution.

Kerosene	2 gallons
Water	1 „
Soft Soap	1 quart
or	
Common Washing Soap	½ pound
Crude Carbolic Acid	1 pint

Spraying Solution

1 Part of the Emulsion

15 Parts of Water.

For making this a hand spray pump and a pail may be used. Pour the kerosene into the pail and dissolve the soap in boiling water, after which pour this into the pail, together with the carbolic acid. Then mix the liquid by pumping vigorously for 10 to 20 minutes, directing the stream back into the pail. The mixture is safe to use if the oil does not separate from the soap after standing a few days. If there is free oil present it is liable to do serious damage to the trees.

This spray may be used for Scale, White Fly and Ants, and even for Red Spider, but Rust Mite and Red Spider may be kept in check by

Sulphur Wash:

Flowers of Sulphur	2 pounds
Caustic Soda 98%	1 „
Water	1½ pint

Place the sulphur in a half barrel and moisten it with water, then add the soda and stir vigorously until it starts to boil. Strain the brown liquid and cork it in bottles. For spraying use one gallon of the solution with 25 gallons of water.

Fungus Pests are not very troublesome on citrus trees in the West Indies, and practically the only one that has so far done damage enough to justify spraying for is the Scab. This attacks Sour Orange, Lemon and often Grape-Fruit severely. Where it is destructive to the fruit it may be necessary to spray and practically the only remedy is a solution of copper

Bordeaux Mixture

Copper Sulphate	4 pounds
Quicklime	6 „
Water	50 gallons

Place the copper sulphate in a grain bag and suspend it in a barrel containing about 25 gallons of water, in which way it will dissolve over night. Slake a quantity of freshly burned lime in as much water as it will take up and no more. Such slaked lime can conveniently be kept in a barrel sunk in the ground and the lime kept covered with water to prevent drying. Of this slaked lime take about 10 pounds and stir it up in about 25 gallons of water. Then strain it and pour it into another barrel simultaneously with the copper sulphate solution while stirring vigorously. The barrel should be filled up to the 50 gallon mark and the mixture is ready for use.

In spraying with Bordeaux mixture the beneficial fungi that keep the scale insects in check are destroyed as well as that which cause injury to the tree. For this reason scale insects multiply fast after spraying with Bordeaux mixture and it should never be used except in connection with scale sprays or closely followed by an application of such. It should be well understood that Bordeaux mixture does not destroy scale nor any other kind of insect, neither do soaps or oil emulsions destroy fungi.

CAGE

7 DAY USE

RETURN TO DESK FROM WHICH BORROWED

Agric. Reference Service

This publication is due on the LAST DATE
stamped below.

MAY 15 1962

RB 17-60m-8,'61
(01641s10)4188

General Library
University of California
Berkeley

259951

SB369

G32

UNIVERSITY OF CALIFORNIA LIBRARY