UC-NRLF

\$B 71 694

LABORATORY MANUAL OF HORTICULTURE

HOOD

GIFT OF
F. L. Griffin

EX LIBRIS

THE 1880 POINT

100

IN THE CITY OF NEW YORK

LABORATORY MANUAL OF HORTICULTURE

WITH ILLUSTRATIONS OF
METHODS, EQUIPMENT, AND APPARATUS

BY

GEORGE W. HOOD

ASSOCIATE PROFESSOR OF HORTICULTURE
UNIVERSITY OF NEBRASKA

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

GINN AND COMPANY

BOSTON • NEW YORK • CHICAGO • LONDON
ATLANTA • DALLAS • COLUMBUS • SAN FRANCISCO

SB 365
H7

Gift
F. L. Griffin.

9/7/21

COPYRIGHT, 1915, BY

GEORGE W. HOOD .

ALL RIGHTS RESERVED

215.8

TO THE
LIBRARY OF
THE
ATHENÆUM PRESS
GINN AND COMPANY
BOSTON U.S.A.

The Athenæum Press
GINN AND COMPANY • PROPRIETORS • BOSTON • U.S.A.

PREFACE

This manual contains the exercises that have been given in connection with the work in General Horticulture at the Ohio State University, Michigan Agricultural College, and University of Nebraska.

It is a modest attempt to fill a place that, for a long time, has seemed to be vacant. The aim is to give the student an outline by which he can better grasp some of the fundamental principles of horticulture, and an opportunity to learn by actual experimentation and observation the reasons for certain necessary operations in horticulture.

It is the object of this manual to give explicit directions for every detail of the work. In certain exercises it may be impossible for the student to do all the tasks outlined, but only such parts as are suited to the conditions under which the student is working should be attempted. This must be determined by the instructor, who, it is hoped, will select those exercises that are most practicable for his class.

The drawings asked for and the questions raised in the several exercises are designed to bring out fundamental principles and important points as well as to fix firmly in the mind of the student the necessity of careful and systematic work.

The author has made use of suggestions found in other works, and has endeavored to bring together the information that will be the most helpful to the student, and to present it in such a manner that it will be of value to the beginner. He makes no claim of completeness for this outline: only such exercises have been included as seem best adapted to a general course in horticulture. Most of these exercises can be done in the laboratory, but after the fundamental principles have been thoroughly mastered field practice should be given in pruning, spraying, etc.

The exercises are arranged in groups which naturally fall under a common head. It is not likely that the instructor can begin with the first exercise and continue uninterruptedly to the

LABORATORY MANUAL OF HORTICULTURE

last; he will probably find it desirable to make the order in which the exercises are used conform to the work in hand. For instance, if the discussion is on buds and pruning, it will be advisable to turn to that section and work out those exercises. If the discussion is on seeds, then the laboratory exercises dealing with germination of seeds should be assigned, and so on through the book. Care should be taken, however, not to assign tasks based on preceding exercises that have not been already worked out or discussed.

The author wishes to express his appreciation to Professor Wendell Paddock of the Ohio State University, Professors R. A. Emerson and J. R. Cooper of the University of Nebraska, and others for valuable suggestions offered in connection with this work.

GEORGE W. HOOD

CONTENTS

EXERCISES	PAGE
I. Comparative study of seed-testers	2
II. Seed-testing	6
III. Seed vitality limited by age	10
IV. Absorption of water by seeds	12
V. Germination hastened by soaking seeds	16
VI. The effect of drought on germinating seeds	18
VII. Seeds treated with chemicals to hasten germination	20
VIII. The effect of commercial-fertilizer solutions on seeds	24
IX. The action of fungicides on seeds	26
X. A study of bulbs	28
XI. A study of corms	30
XII. A study of soft or green wood cuttings	32
XIII. A study of leaf cuttings	36
XIV. A study of root cuttings	38
XV. A study of hardwood, or dormant, cuttings	40
XVI. A study of one-eye, two-eye, three-eye, mallet, and heel cuttings	44
XVII-XVIII. A study of buds (<i>Prunus</i> sp.)	
Plums	
<i>Prunus domestica</i>	48
<i>Prunus triflora</i>	50
<i>Prunus americana</i>	50
<i>Prunus hortulana</i>	52
<i>Prunus munsoniana</i>	52
The cherry	
<i>Prunus avium</i> (sweet cherry)	54
<i>Prunus cerasus</i> (sour cherry)	58
XIX. A study of buds (<i>Amygdalus</i> sp.)	
The peach	
<i>Amygdalus persica</i>	60
North China race	60
Persian race	62
South China race	62
Spanish race	64
<i>Amygdalus platycarpa</i>	64
Peen-to race	64
XX. A study of buds — pomaceous fruits	
The apple	
<i>Malus sylvestris</i>	66
The pear	
<i>Pyrus communis</i> and <i>Pyrus sinensis</i>	68
The quince	
<i>Cydonia vulgaris</i>	70

LABORATORY MANUAL OF HORTICULTURE

EXERCISES

PAGE

XXI. A study of buds — vine fruits	
The grape	
<i>Vitis labrusca</i>	72
<i>Vitis vinifera</i>	74
<i>Vitis labrusca</i> × <i>vinifera</i>	74
XXII. Pruning	76
XXIII. The pruning of the apple	80
XXIV. The pruning of the pear	82
XXV. The pruning of the plum and the cherry	84
XXVI. The pruning of the peach	86
XXVII. The pruning of the grape	88
XXVIII. The pruning of the brambles	90
XXIX. The pruning of the bush fruits	92
XXX. The study of shield and prong budding	94
XXXI. The study of plate budding and H-budding	98
XXXII. The study of flute budding, ring budding, and chip budding	102
XXXIII. The study of grafting-waxes	
Beeswax and resin waxes	108
Alcoholic waxes	110
Pitch wax	110
XXXIV. The study of splice, whip, and double-whip grafting	112
XXXV. The study of cleft grafting and bark grafting	118
XXXVI. The study of bridge grafting	126
XXXVII. The study of side grafting	128
XXXVIII. The study of veneer grafting	130
XXXIX. The study of Bordeaux mixture	132
XL. The study of lime-sulphur	138
XLI. The study of self-boiled lime-sulphur	144
XLII. The study of ammoniacal solution of copper carbonate	146
XLIII. The study of Paris green and arsenate of lead	148
XLIV. The study of kerosene emulsion	152
XLV-LIV. A study of the apple	154
LV-LX. A study of the grape	174
LXI-LXVI. A study of the peach	186
LXVII-LXXII. A study of the pear	198
LXXIII-LXXVIII. A study of the plum	210
GLOSSARY	223
INDEX	233

UNIVERSITY OF CALIFORNIA
LIBRARY

LABORATORY MANUAL OF
HORTICULTURE

EXERCISE I

COMPARATIVE STUDY OF SEED-TESTERS

Material. Dinner plates, blotters, canton flannel, germinating cups, tile germinator, Geneva seed-tester, germinating chamber.

Seed-testers are used for the purpose of testing the vitality of seeds. They differ greatly, and their efficiency varies with the kind. Study carefully four kinds.

1. The dinner-plate germinator consists of two large-sized dinner plates, one turned over the other, between which are two or three blotters covered with two pieces of canton flannel. Make a drawing showing the two plates and the inside material. The drawings should be at least three inches in diameter. Describe fully. Label all parts and describe the materials of which it is made. What can you say about the efficiency of this tester?

2. Germinating cups are small earthen cups three inches in diameter or three inches square and one and one-half inches deep, covered with a lid of the same size and shape as the top of the cup, in which are a number of small holes. Make a drawing, natural size, of a germinating cup and lid. Label, and give the dimensions on the drawings. Describe how the seeds can be germinated in this vessel. The cups must be placed in a shallow pan of water. Discuss the purpose of this procedure.

3. The tile germinator is a large tile twelve inches wide by fifteen inches long and two inches deep, in which have been molded pockets of varying sizes from one inch to three inches in diameter. This tester is placed in a shallow pan of water the same as the germinating cups. Make a drawing of the top view and a cross section, showing the number, size, and the depth of pockets. What advantage, if any, has this tester over the previous one? Give the exact dimensions on your drawing. Make the drawing two by three inches. How does the moisture get to the seeds? Discuss fully how you would germinate seeds in this tester. A pane of glass is usually placed over the top of the tester. What is the purpose of the glass? Give your opinion as to the value derived from its use.

COMPARATIVE STUDY OF SEED-TESTERS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

4. The Geneva seed-tester is a galvanized iron pan ten inches wide, fourteen inches long, and three and one-half inches deep, with a ledge three eighths of an inch wide along the sides. The seeds are held in folds of cloth suspended on rods which rest on this ledge. Describe how the water is carried to the seeds.

FIG. 1. Tile germinator, showing the different-sized holes in which the seeds are placed for testing

Make a drawing of the pan, three inches in length and of proportional width, showing the ledge. Draw an end view of the cloth. Give the depth of the pocket in inches, and measure the length of each end of the cloth from the supporting-rod. Why should the end pieces of the cloth be longer than the depth of the pocket?

FIG. 2. Geneva seed-tester

A, supporting-rod for the cloth ; B, Canton flannel pocket ; C, opening through which the supporting-rod passes

COMPARATIVE STUDY OF SEED-TESTERS

STUDENT'S NOTES AND REPORT

EXERCISE II

SEED-TESTING

Material. Seeds of clover, celery, onion, cucumber, tomato, wheat, cabbage, oats, eggplant, beet, peas, rye, beans.

Apparatus. Sprouting-chamber, Geneva seed-tester, pane of glass the size of the tester.

Use the Geneva seed-tester to familiarize the student with this method, which can be applied on a large or a small scale. Each seed-tester should be thoroughly sterilized by placing about one inch of water in the pan, allowing the cloth to remain in the folds suspended on the rods, and boiling the water from fifteen minutes to a half hour, or until everything is thoroughly steamed. A piece of glass should be placed over the top of the tester to hold the steam and to give a more thorough sterilization.

FIG. 3. Sprouting-chamber, or germinating-oven, showing the Geneva seed-testers on the shelves

1. Count out one hundred seeds of clover, celery, onion, cucumber, tomato, wheat, cabbage, oats, eggplant, beets, rye, and fifty seeds of peas and beans. Place the seeds in the tester, which has previously been thoroughly sterilized with steam. Discuss why we sterilize the tester. The tester should be placed in a sprouting-chamber

where the temperature can be maintained around 80° F. Why? Discuss fully what would happen to the seeds if the temperature were to fall as low as 50° F. Why is it important that seeds should not be planted until the soil is thoroughly warmed up? Will any seed germinate in the presence of oxygen and moisture

SEED-TESTING

STUDENT'S NOTES AND REPORT

TABLE OF RESULTS

NAME OF SEED	DATE GERMI- NATION BEGAN	NO. OF SEEDS	NO. OF SEEDS GERMINATING, BY DAYS							TOTAL NO. GERMI- NATING	PER CENT GERMI- NATION
			1st	2d	3d	4th	5th	6th	7th		

LABORATORY MANUAL OF HORTICULTURE

without heat? In testing seeds do all of them germinate at the same time? Discuss the irregularity of seeds in germinating when the same external conditions are given. Is a sprouting-chamber necessary in order to germinate seeds? Discuss the advantages gained by the rapid germination of seeds.

2. Tabulate your results, and figure out the per cent of germination. This test should continue from a week to ten days, but a week is usually sufficient for most seeds. Remove all of the seeds as they germinate after the count has been tabulated. Discuss fully what is gained by seed-testing. Discuss the three essentials necessary for the germination of seeds.

NOTE. I have found that the most convenient size of the Geneva seed-tester for class use is eight inches wide, fourteen inches long, and three inches deep. No. 12 wire makes a good size to support the cloth, and the length of the cloth most convenient is four feet, containing ten pockets. In making the pockets, enough Canton flannel should be allowed for the making of the seam, so that the pocket may be at least one and one-half inches deep. Two or three such cloths may be placed in each pan if necessary.

REFERENCES :

Bailey, "The Nursery Book," chap. i.

Bulletin No. 35, Rhode Island Agricultural Experiment Station.

Goff, "Principles of Plant Culture," pp. 24-32.

SEED-TESTING

STUDENT'S NOTES AND REPORT

SEED VITALITY LIMITED BY AGE

Apparatus. Sprouting-chamber, Geneva seed-tester.

1. Secure one hundred seeds each of the cucumber, tomato, and cabbage, one year old, two years old, three years old, four years old, five years old, six years old, seven years old, eight years old, nine years old, and ten years old, and place them in a Geneva seed-tester, the same as in Exercise II. Carry the germination test seven days and tabulate your results.

2. From your results, plot a curve showing the decrease in the vitality due to the age of the seed. At what age does the greatest vitality occur in each kind of seed? How do you account for this? Discuss the age limit of the different seeds.

TABLE OF RESULTS

[illegible]

[10]

SEED VITALITY LIMITED BY AGE

STUDENT'S NOTES AND REPORT

HOW TO PLOT THE CURVE

EXERCISE IV

ABSORPTION OF WATER BY SEEDS

Material. Seeds of beans, peas, corn, canna, honey locust, and field beans.

Apparatus. Scale, 250-cc. graduated cylinder, 100-cc. wide-mouthed bottles.

1. *To determine volume.* Partly fill the graduated cylinder with water and read the height to which the column of water stands. Now count out fifty seeds of the bean, pea, and corn and drop each lot separately into the water in the cylinder. Remove the air bubbles each time by slightly shaking the cylinder. The height of the column of water is again read and the difference in the two readings indicates the volume of each sample in cubic centimeters. How does this vary with the different seeds? Explain.

2. *To determine the amount of water absorbed by seeds.* Weigh fifty seeds of peas, beans, and corn and tabulate the weight of each lot. Place the seeds in a wide-mouthed bottle of about 100 cc. and cover them with water. Allow them to stand for twenty-four hours in a room where the temperature is about 70° F. The following day remove the seeds, wipe dry, and weigh. Tabulate your results. Discuss. Allow the same seeds to soak in water for another day and note whether there has been any increase in weight. At what time was the greatest amount of water taken up? Discuss.

3. Now take fifty more seeds, the same as in 2, and treat in the same manner, only place them in a temperature of about 40° F. Note your results and explain.

4. Secure fifty seeds of the canna or honey locust and treat them in the same manner as in 1 and 2. Note the difference, if any, and discuss. How do you account for the change?

5. Weigh out two samples of field beans and place one lot in water and the second sample in damp sand. Weigh each lot the two following days and compute the per cent of water absorbed by each.

ABSORPTION OF WATER BY SEEDS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

6. Weigh out two samples of field beans and place one sample in loose, moist sand and the other in the same kind of sand, but well packed and pressed over the seed. Weigh the seeds the two following days. Discuss and give your reason for the results obtained.

REFERENCE :

Goff, "Principles of Plant Culture," pp. 22-25.

ABSORPTION OF WATER BY SEEDS

STUDENT'S NOTES AND REPORT

EXERCISE V

GERMINATION HASTENED BY SOAKING SEEDS

Material. Seeds of beans, peas, corn, lettuce, beet, cabbage, salsify, parsnip, carrot, wheat, and oats.

Apparatus. 100-cc. wide-mouthed bottles, Geneva seed-tester, sprouting-chamber.

1. Count out fifty seeds of beans, peas, and corn. Divide these into two lots of twenty-five seeds each and allow the first lot to soak for two hours. Treat the second lot in exactly the same way, but allow them to soak for twelve hours. Now place both sets in a Geneva seed-tester and allow them to germinate. What differences do you observe? How do you account for your results? Explain.

2. Count out one hundred seeds each of lettuce, beet, cabbage, salsify, parsnip, carrot, wheat, and oats and treat the same as in 1.

Now compare these results with those obtained in Exercise II, and plot a curve for beans, cabbage, and salsify, showing the difference, if any, in time of germination. Explain fully your results. What is gained by the soaking of seeds? Do all seeds respond alike? If not, why?

GERMINATION HASTENED BY SOAKING SEEDS

STUDENT'S NOTES AND REPORT

NOTE. Use different-colored inks ; for example, red = bean, black = cabbage, green = salsify.

EXERCISE VI

THE EFFECT OF DROUGHT ON GERMINATING SEEDS

Material. Seeds of clover, peas, beans, alfalfa, cowpeas, corn, wheat, rye, and oats.

Apparatus. Sprouting-chamber, Geneva seed-tester.

Germination is often interfered with by the drying out of the soil. The seeds of all dicotyledons suffer considerably from drying out, while the cereals and, in general, the seeds of all monocotyledons suffer but little. There are, however, a few exceptions, as corn, clover, beans, peas, etc.

1. Count out one hundred seeds each of clover, peas, beans, alfalfa, cowpeas, corn, wheat, rye, and oats. Place them in a Geneva seed-tester and allow them to germinate from twenty-four to forty-eight hours. Now remove the tester from the germinating-chamber, pour out the water, and spread the cloths out so that the seeds will dry for twenty-four hours, or until they are well dried, then pour water in the tester, and again wet the seeds thoroughly and place them in the germinating-chamber. Do this several times and note the results. At the same time carry on a check for each seed which has been treated in the normal way to compare with the others. What difference is seen in the germination? Do all the seeds respond alike? If not, why? What seeds are injured the most by drying out? What ones are not injured? Study carefully each sample and determine why some seeds are injured and some are not. Has the construction of the seed anything to do with the drying out? Tabulate and discuss your results fully.

THE EFFECT OF DROUGHT ON GERMINATING SEEDS

STUDENT'S NOTES AND REPORT

TABLE OF RESULTS

[illegible]

EXERCISE VII

SEEDS TREATED WITH CHEMICALS TO HASTEN GERMINATION

Material. Seeds of the honey locust, asparagus, wheat, barley, oats, rye, radish, cabbage. 5, 10, and 50 per cent and concentrated sulphuric acid, saturated solution of limewater, 2 per cent solution of iodine, saturated solution of chlorine water, 1 per cent monobromide of camphor.

Apparatus. 100-cc. wide-mouthed bottles, Geneva seed-tester, sprouting-chamber.

The treatment of seeds with chemicals has long been known, but definite information concerning the exact effect of such treatment on germination is meager and based on three hypotheses.

Hypothesis 1. If seeds are treated with acids, the hard seed coat is loosened, which makes possible the more rapid imbibition of water.

Count out five lots of ten seeds each of the honey locust and asparagus and proceed as follows: Place the seeds of each lot in a small, wide-mouthed bottle and immerse four of them each respectively in 5, 10, and 50 per cent dilutions of sulphuric acid and in the concentrated acid for thirty minutes. Place them in the Geneva seed-tester and germinate. Use one lot untreated to serve as a check. Tabulate the results every twenty-four hours on both treated and untreated seeds. Describe what action the acid of different strengths has on the seeds. What per cent of acid gives the best result? What per cent solution would you advise? Are both kinds of seed acted upon alike by the same solution? Plot a curve showing the increase in the time of germination over the nontreated seed. Use colored inks to show the difference in the time.

Hypothesis 2. If the evolution of carbonic acid gas can be hastened by the use of chemicals, germination will be facilitated.

Count out ten lots of one hundred seeds each of wheat, barley, oats, and rye. Place one lot of each seed in a 100-cc. wide-mouthed bottle, cover the seeds with a saturated solution of limewater, and allow them to remain immersed for thirty minutes.

SEEDS TREATED WITH CHEMICALS TO HASTEN GERMINATION

STUDENT'S NOTES AND REPORT

CURVE FOR ACID TREATMENT

CURVE FOR LIMEWATER TREATMENT

NOTE. Use colored inks, as, for example, red ink = canna seed, black ink = asparagus, blue = check ; or use the following symbols: - - - = canna seed, — — — = asparagus, — — — — = check, etc.

LABORATORY MANUAL OF HORTICULTURE

Now place the seeds in a Geneva seed-tester and germinate. The second lot of seeds should be germinated untreated in the same tester, under the same conditions, to serve as a check. Tabulate your results and describe fully what takes place. Plot a curve to show the increase in germination of seed that has been treated over seed that has not been treated.

Hypothesis 3. If you can set oxygen free, to unite with the carbon of the seed and thus form carbonic acid gas, which must pass off, germination will be hastened.

a. Count out one hundred seeds each of the radish, cabbage, and onion, and place each in a wide-mouthed bottle. Now cover them with a 60 per cent alcoholic solution of iodine (approximately 1.1 per cent solution) and allow the seed to soak or be kept moist with the solution until germination begins. Note the time of sprouting. Tabulate your results to show the increase in time. Discuss.

b. Count out four lots of one hundred seeds each of the radish. Place each lot in a bottle and keep the first moist with pure water, the second with a 60 per cent alcoholic solution of iodine, the third with a saturated solution of chlorine water (approximately 1.8 per cent), and the fourth with a 1 per cent solution of monobromide of camphor. Describe fully what takes place and tabulate your results. Plot a curve to show what effect the different chemicals have on the germination of the radish seeds.

SEEDS TREATED WITH CHEMICALS TO HASTEN GERMINATION

STUDENT'S NOTES AND REPORT

CURVE TO SHOW RADISH GERMINATION

No. of seeds

NOTE. Use different-colored inks or different symbols to represent the different solutions.

EXERCISE VIII

THE EFFECT OF COMMERCIAL-FERTILIZER SOLUTIONS ON SEEDS

Material. Clover seed, 1 per cent solution of sodium nitrate, 1 per cent solution of muriate of potash, 1 per cent solution of phosphoric acid, and a saturated solution of limewater.

Apparatus. 100-cc. wide-mouthed bottles, Geneva seed-testers. It has long been known that certain chemical fertilizers, as well as certain forms of other chemicals, have some influence on the seed, in the soil solution as well as in an aqueous solution.

*1. Count out four lots of one hundred seeds of the clover. Place each lot in a wide-mouthed bottle and cover them respectively with a 1 per cent solution of sodium nitrate, a 1 per cent solution of muriate of potash, and a 1 per cent solution of phosphoric acid, allowing each lot to remain in its respective solution for twenty-four hours. Remove each sample, and germinate in the Geneva seed-tester. Place the fourth batch untreated in the seed-tester to serve as a check. Tabulate and discuss your results in each case. Plot a curve showing the per cent of germination with each of the fertilizers and also when no fertilizer is used. Does it pay to allow a commercial fertilizer to come in direct contact with the seed? Discuss.

2. Mix in equal proportions a saturated solution of limewater with a 1 per cent solution of each of the following: sodium nitrate, muriate of potash, and phosphoric acid. Now with each solution treat such a batch of seeds as in 1, and germinate. Does the addition of the limewater to the seed aid in preventing any detrimental effect of the commercial fertilizers?

Do the fertilizers all act alike on the different seeds? Which fertilizer shows the most injurious effect? Are the seeds hastened or retarded in germinating by the use of fertilizers? Discuss. Tabulate your results, and plot a curve showing what effect the limewater has when added to each of the fertilizers.

THE EFFECT OF COMMERCIAL-FERTILIZER SOLUTIONS

STUDENT'S NOTES AND REPORT

NOTE. Use different-colored inks or different symbols to illustrate each fertilizer.

RECORD

NAME OF FERTILIZER	NO. OF SEEDS GERMINATING, BY DAYS							TOTAL NUMBER GERMINATING	PER CENT GERMINATION
	1st	2d	3d	4th	5th	6th	7th		
Nitrate									
Potash									
Phosphoric acid . . .									
Nitrate + lime . . .									
Potash + lime . . .									
Phosphoric acid + lime									
Check									

EXERCISE IX

THE ACTION OF FUNGICIDES ON SEEDS

Material. Seeds of the wheat, cabbage, and corn, 1 per cent solution of copper sulphate, 25 per cent salt solution.

Apparatus. 100-cc. wide-mouthed bottles, Geneva seed-tester.

Seeds are often treated with fungicides to destroy the spores of some diseases. This treatment, however, is dangerous and must be applied with extreme care and precision, in order that no damage may result to the seed.

1. Count out two lots, of one hundred seeds each, of the wheat, of the cabbage, and of the sweet corn. Place the samples of each variety in separate bottles. Now cover one lot of seeds with a 1 per cent solution of copper sulphate. Place the second lot in pure water to serve as a check. Allow both batches to soak for two hours, and then germinate. Tabulate your results, and explain. What strength of solution would you recommend for the treatment of seeds in order to insure the best results? Compare the treated seeds with the check. What is gained by the treatment? Did you note any fungus growing? What seeds, if any, were affected? Discuss.

2. Count out four lots of one hundred seeds each, as in 1. Place each lot in a 100-cc. bottle, and keep one as a check. Pour enough of the salt solution over the first batch to cover the seeds, and allow them to remain immersed one minute. Soak the second batch of seeds for five minutes and the third batch for fifteen minutes. Germinate as in 1, and note the results. Describe fully. Tabulate the results as in 1, and discuss fully.

THE ACTION OF FUNGICIDES ON SEEDS

STUDENT'S NOTES AND REPORT

COPPER SULPHATE SOLUTION

NAME OF SEED	NO. OF SEEDS GERMINATING, BY DAYS							TOTAL NUMBER GERMI- NATING	PER CENT GERMINATION
	1st	2d	3d	4th	5th	6th	7th		
Wheat									
Cabbage									
Corn									

SALT SOLUTION

NAME OF SEED	LENGTH OF TIME IMMERSED	NO. OF SEEDS GERMINATING, BY DAYS							TOTAL NUMBER GERMI- NATING	PER CENT GERMINATION
		1st	2d	3d	4th	5th	6th	7th		
Wheat	1 min.									
	5 min.									
	15 min.									
	Check									
Cabbage	1 min.									
	5 min.									
	15 min.									
	Check									
Corn	1 min.									
	5 min.									
	15 min.									
	Check									

REFERENCE :

Bulletin No. 10, United States Department of Agriculture, Division of Vegetable Physiology and Pathology.

EXERCISE X

A STUDY OF BULBS

Material and apparatus. Bulbs of the hyacinth, tulip, onion, and lily; sharp knife.

Bulbs of all kinds are specialized buds. They are made up of a short and rudimentary axis closely incased in transformed and thickened leaves called bulb scales. Bulbs occur only in plants which are accustomed to long periods of inactivity, and are divided into two classes determined by the scales.

FIG. 4. Bulb of Easter lily

1. Study bulbs of the hyacinth, onion, tulip, and tiger lily or Easter lily. Make a drawing of each one, natural size, showing the external characteristics. Describe fully the differences in each bulb. How does the lily differ from the hyacinth?

2. Keep a close watch for bulbils, which are little bulbs found on the parent bulb. Where do you find these located? Are they in essentially the same place in each bulb? Make a drawing of the bulbil, natural size, showing the attachment to the parent.

3. Draw a cross section, natural size, of each bulb. What differences do you observe? Carefully examine and describe the bulb scales of the hyacinth and the tulip. How does the scale of the lily differ from that of the tulip and the hyacinth?

4. Draw a cross section of each scale. Describe in detail the differences. Measure the thickness of each bulb scale. What significance does the thickness of the bulb scale have? What is contained in the scales of all bulbs? How do the scales of the lily differ in arrangement and shape from those of the hyacinth and the tulip?

REFERENCE :

Bailey, "The Nursery Book," chap. ii.

BULBS

STUDENT'S NOTES AND REPORT

EXERCISE NO. FIFTY-4

A student of the University of California, Berkeley, California, has been studying the habits of the California Bulb, and has been able to determine the following facts:

The student has also been able to determine the following facts:

EXERCISE XI

A STUDY OF CORMS

Material and apparatus. Corms of the gladiolus and the crocus; sharp knife.

A corm is a bulblike organ, differing from a bulb in being solid throughout and not composed of fleshy scales. Cormous plants are multiplied in essentially the same way as are bulbous plants.

FIG. 5. Corms of the gladiolus, showing the cormels, or the little corms, about the parent

1. Study the corm of the gladiolus and the crocus. Make a natural-size drawing of each of the corms. What external differences do you note? Remove the outer husk and note the lines surrounding the corm. Discuss how a corm grows.

2. Draw a cross section, natural size, of both of the corms. How do they differ from the bulbs in Exercise X? Is the shape of the corm the same as that of the bulb? Look for the central axis. Is this differentiated from the main body of the corm? Carefully examine the corm for eyes. Discuss fully.

THE STUDY OF CORMS

STUDENT'S NOTES AND REPORT

100

THE STUDY OF CORMS
CORMS ARE THE THICKENED PARTS OF THE ROOTS OF PLANTS
WHICH STORE FOOD FOR THE PLANT. THEY ARE FOUND IN
MANY PLANTS, SUCH AS IRIS, GLADIOLUS, AND
HYACINTH. THE CORMS OF THESE PLANTS ARE
USED FOR PROPAGATION AND FOR THE
PREPARATION OF MEDICINES.

THE STUDY OF CORMS
CORMS ARE THE THICKENED PARTS OF THE ROOTS OF PLANTS
WHICH STORE FOOD FOR THE PLANT. THEY ARE FOUND IN
MANY PLANTS, SUCH AS IRIS, GLADIOLUS, AND
HYACINTH. THE CORMS OF THESE PLANTS ARE
USED FOR PROPAGATION AND FOR THE
PREPARATION OF MEDICINES.

EXERCISE XII

A STUDY OF SOFT OR GREEN WOOD CUTTINGS

Material. Stock plants of the geranium, coleus, chrysanthemum, carnation, cactus, oleander, hydrangea, and rose, and rooted cuttings of each.

Soft or green wood cuttings are more commonly employed than those from mature wood because they root more quickly and can

FIG. 6. Soft wood cuttings of different plants

1, rose geranium; 2, geranium; 3, variegated geranium; 4, 5, coleus; 6, rose; 7, carnation; 8, chrysanthemum; 9, oleander; 10, hibiscus; 11, tomato

be handled under glass in winter, and also because more species are capable of propagation by this method. All soft-wooded and many ornamental plants are propagated by green wood cuttings.

There are two general classes of green wood cuttings: first, those that are made from the soft and still-growing wood, and second, those that are made from the green but nearly ripened wood.

SOFT OR GREEN WOOD CUTTINGS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

1. Select several plants, such as the geranium, coleus, carnation, chrysanthemum, cactus, etc., and study. Make a number of cuttings from each kind of plant. Draw a cutting, natural size, from each of the plants, and label. It is always best to remove about one half of the leaves before planting the cutting. Why is this procedure necessary? Explain. Now secure some cuttings from each of the above plants that have been rooted, and examine carefully. Where do the roots start? Is there any callus tissue present? If so, describe. Make a drawing showing the callus tissue, and the roots just starting. Label all of the parts.

2. Secure several plants with semihard wood, such as the oleander, rose, hydrangea, etc., and compare these plants with those in 1. What difference do you find in the texture of the wood? Describe any other unlikeness or differences. Make a natural-size drawing of each cutting, four inches in length. Differentiate between the still-growing and the nearly ripened wood cutting.

SOFT OR GREEN WOOD CUTTINGS

STUDENT'S NOTES AND REPORT

THEORY OF THE EXPERIMENT

THEORY OF THE EXPERIMENT

The purpose of this experiment is to determine the effect of the temperature of the water on the rate of diffusion of a substance into a soft wood cutting. The experiment is performed by placing a soft wood cutting in a beaker of water and measuring the amount of substance that diffuses into the wood over a period of time.

The results of the experiment show that the rate of diffusion increases with the temperature of the water. This is because the molecules of the substance have more kinetic energy at higher temperatures, and therefore move more rapidly through the wood. The experiment also shows that the rate of diffusion is affected by the thickness of the wood cutting, with thinner cuttings showing a faster rate of diffusion.

EXERCISE XIII

A STUDY OF LEAF CUTTINGS

Material. Stock plants of the begonia, bryophyllum calycinum, and gloxinia, and rooted leaf cuttings of each.

Many plants having thick, fleshy leaves can easily be propagated by leaf cuttings. In some plants the entire leaf is used ;

in others the leaf is cut half in two and placed in the ground on its edge. In some begonias the leaf is cut into several fan-shaped sections, each cutting having a portion of the petiole attached.

Study the begonia, bryophyllum, and gloxinia. Each plant represents a different form of leaf cutting.

1. *Begonia*. Draw a leaf of one of the large, thick-leaved types. Indicate by dotted lines where the cuts should be made to produce the fan-shaped cuttings. Draw a single cutting.

FIG. 7. Leaf of the bryophyllum, showing the starting of the young plants

How is it planted ? Where does the young plant start ? Discuss.

2. *Bryophyllum*. Draw a leaf natural size, and note the growing points. Describe. This plant is usually propagated by laying the leaf flat on the surface of the ground, where several small plants will start. Study a leaf from which the young plants have started. Where is the origin of each plant ?

3. *Gloxinia*. Draw and describe a rooted leaf cutting.

A STUDY OF LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

THE STUDENT

DATE

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

LEAF CUTTINGS

STUDENT'S NOTES AND REPORT

EXERCISE XIV

A STUDY OF ROOT CUTTINGS

Material. Blackberry and horse-radish roots and planted cuttings of each.

Many plants can be multiplied with ease by means of short cuttings of the roots. True root cuttings possess no buds at all,

FIG. 8. Root cuttings of the horse-radish

but buds develop after the cutting is planted. The roots are cut into pieces from one inch to three inches in length and planted horizontally.

1. Secure roots of the blackberry and horse-radish, both of which are easily propagated by root cuttings. Practice making a number of these cuttings. Make a natural-size drawing of a root cutting from each of the plants. Look carefully for buds. Can any be found? Now secure cuttings of the above that have been planted for some little time. Examine these for buds. Where are the buds found? Make a drawing showing the bud, and differentiate between the two cuttings. Discuss.

A STUDY OF ROOT CUTTINGS

STUDENT'S NOTES AND REPORT

THE STUDENT

DATE

The student has been studying the effects of root cuttings on the growth of plants. The student has found that root cuttings can be used to propagate many plants. The student has also found that root cuttings can be used to study the effects of various factors on plant growth. The student has found that root cuttings can be used to study the effects of light, temperature, and humidity on plant growth. The student has found that root cuttings can be used to study the effects of various nutrients on plant growth. The student has found that root cuttings can be used to study the effects of various hormones on plant growth. The student has found that root cuttings can be used to study the effects of various diseases on plant growth. The student has found that root cuttings can be used to study the effects of various insects on plant growth. The student has found that root cuttings can be used to study the effects of various fungi on plant growth. The student has found that root cuttings can be used to study the effects of various bacteria on plant growth. The student has found that root cuttings can be used to study the effects of various viruses on plant growth. The student has found that root cuttings can be used to study the effects of various parasites on plant growth. The student has found that root cuttings can be used to study the effects of various predators on plant growth. The student has found that root cuttings can be used to study the effects of various competitors on plant growth. The student has found that root cuttings can be used to study the effects of various mutualists on plant growth. The student has found that root cuttings can be used to study the effects of various commensals on plant growth. The student has found that root cuttings can be used to study the effects of various symbionts on plant growth. The student has found that root cuttings can be used to study the effects of various pathogens on plant growth. The student has found that root cuttings can be used to study the effects of various herbivores on plant growth. The student has found that root cuttings can be used to study the effects of various carnivores on plant growth. The student has found that root cuttings can be used to study the effects of various omnivores on plant growth. The student has found that root cuttings can be used to study the effects of various detritivores on plant growth. The student has found that root cuttings can be used to study the effects of various decomposers on plant growth. The student has found that root cuttings can be used to study the effects of various scavengers on plant growth. The student has found that root cuttings can be used to study the effects of various predators on plant growth. The student has found that root cuttings can be used to study the effects of various competitors on plant growth. The student has found that root cuttings can be used to study the effects of various mutualists on plant growth. The student has found that root cuttings can be used to study the effects of various commensals on plant growth. The student has found that root cuttings can be used to study the effects of various symbionts on plant growth. The student has found that root cuttings can be used to study the effects of various pathogens on plant growth. The student has found that root cuttings can be used to study the effects of various herbivores on plant growth. The student has found that root cuttings can be used to study the effects of various carnivores on plant growth. The student has found that root cuttings can be used to study the effects of various omnivores on plant growth. The student has found that root cuttings can be used to study the effects of various detritivores on plant growth. The student has found that root cuttings can be used to study the effects of various decomposers on plant growth. The student has found that root cuttings can be used to study the effects of various scavengers on plant growth.

EXERCISE XV

A STUDY OF HARDWOOD, OR DORMANT, CUTTINGS

Material and apparatus. Currant and gooseberry branches and rooted cuttings of each; sharp knife.

Hardwood cuttings are made from the dormant, or mature, wood of dicotyledonous plants, and should be cut just below a node because they form roots more promptly if cut at this point. They are usually taken in the fall or the early winter and stored in a cool place in sand so that they will callus over before being planted the following spring.

1. Study hardwood cuttings of the currant and the gooseberry. These cuttings should be about six inches in length, and, if the weather permits, the students should be allowed to collect their own cuttings from the parent bush.

Make a natural-size drawing of each of the cuttings. How many buds are usually found on each? How does the cutting of the currant differ from that of the gooseberry? What is the age of the wood from which cuttings are taken? Describe.

2. Study some cuttings of the currant and the gooseberry that have been stored in damp sand for two months. How do they differ from cuttings just taken? Make a drawing of the lower end of the cutting, showing the callus tissue. What is the purpose of this tissue? Label all parts, and discuss.

3. Study some cuttings that have just started into growth. From what position on the cuttings do the roots come? Explain. What indications are there on the cuttings that roots are forming? Discuss.

REFERENCES:

Bailey, "The Nursery Book," chap. iv.

Goff, "Principles of Plant Culture," pp. 207-218.

A STUDY OF HARDWOOD, OR DORMANT, CUTTINGS

STUDENT'S NOTES AND REPORT

FIG. 9. Hardwood cuttings of the gooseberry and the currant
Natural size

A STUDY OF HARDWOOD, OR DORMANT, CUTTINGS

STUDENT'S NOTES AND REPORT

EXERCISE XVI

A STUDY OF ONE-EYE, TWO-EYE, THREE-EYE, MALLET, AND HEEL CUTTINGS

Material. Grape canes, stored and rooted cuttings of the grape.

These cuttings fall under the general head of hardwood cuttings, but they represent different forms which are best suited

to certain plants. The plant which lends itself best to the study of these forms of hardwood cuttings is the grape.

1. A one-eye cutting is a hardwood cutting containing one well-developed and mature bud, with a small piece of the branch on each side of the bud. These cuttings are laid flat when planted. Make a natural-size drawing of a one-eye cutting of the grape. How long is this cutting? Label, and describe.

FIG. 10. Hardwood cuttings of the grape, showing the three-eye, the two-eye, the one-eye, the heel, and the mallet cutting

2. A two-eye and a three-eye cutting are cuttings containing two and three buds respectively, with a small portion of the branch at each end. Make a natural-size drawing of each cutting. How long are

A STUDY OF MALLET AND HEEL CUTTINGS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

the internodes? Study from a stored cutting the callus tissue which has formed. Does this differ from the callus tissue in the currant and the gooseberry?

3. A *mallet cutting* is a cutting that contains a small section of the parent branch forming a T, or *mallet*, at its base (Fig. 10). Some varieties of grapes, as well as many ornamentals, root easily from this kind of a cutting. It should usually have two buds. Make a natural-size drawing showing the *mallet*. The *mallet* portion should be about one inch in length. Label all parts. These cuttings are treated in the same way as single-stem cuttings.

4. A *heel cutting* is a cutting with a small portion of the wood of the previous year at its base (Fig. 10). This is similar to the *mallet cutting*, but the *heel* is only a small piece shaved from one side of the parent branch.

Make a drawing showing the cutting attached to the parent branch, and indicate by dotted lines where the cut should be made. This cutting should contain two buds.

Compare this with the drawing made in 3 and give the significance of both forms. What advantages have *mallet cuttings* and *heel cuttings* over one-eye, two-eye, and three-eye cuttings? Discuss *hardwood cuttings* in general.

REFERENCE:

Farmers' Bulletin No. 157, pp. 10-12.

A STUDY OF MALLET AND HEEL CUTTINGS

STUDENT'S NOTES AND REPORT

EXERCISE XVII

A STUDY OF BUDS (*Prunus* sp.)

Material and apparatus. Branches of several of the commercial groups of plums, bearing fruit buds and branch buds; razor, dissecting lens.

FIG. 11. Branch buds and fruit buds of plum
Note the clusters of buds that are borne on the
short spurs

The genus *Prunus* includes the plum and the cherry, and is widely distributed. These species, cultivated for their edible fruit, are found only in the temperate zone of the northern hemisphere. The flowers, twigs, and buds in each group will be found to differ, as may be verified after studying them.

PLUMS

The plums are divided into several important, more or less distinct, commercial groups, with many hybrids between the groups. The following groups are a few that admit of careful study.

1. *Prunus domestica* includes the following varieties: Bavay, Lombard, Gold Drop, Arch Duke, Diamond, Monarch, Italian Prune, Yellow Gage, and several others.

a. Secure branches, including both fruit buds and branch buds, of one of the above varieties. Draw a twig, natural size, and at

A STUDY OF BUDS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

least four inches in length. Measure the distance between the internodes and count the number of buds in a cycle. Describe the color of the bark. Are the lenticels numerous or few? Describe.

b. Draw a longitudinal and a cross section of both a flower bud and a branch bud. Describe the differences. How many ovules do you find in a bud? Is the number constant or variable? Discuss. Does the fruit bud differ from the branch bud in position on the stem? Discuss.

c. Make a drawing, at least one inch in length, of a branch bud and a flower bud. Remove several of the bud scales, and make a drawing of a scale from both the branch bud and the flower bud. Are there any differences? If so, describe.

2. *Prunus triflora* includes the following varieties: Ogon, Occident, Red June, Willard, Abundance, and several others.

a. Secure branches from one or more of the varieties named or others of the *Prunus triflora* that are available. Make a natural-size drawing of a branch four inches in length, showing the number and the position of the buds. Count from external appearance the number of fruit buds and branch buds. Measure the distance between the buds, and count the number in a cycle. Is the number constant? Discuss.

b. Make a drawing of a longitudinal and a cross section of both a branch bud and a flower bud. This drawing should show detail and be at least one inch in length. How many ovules do you find in the flower bud. Label all parts. Describe.

c. After a branch bud and a flower bud have been definitely determined, make a drawing showing each bud enlarged to one inch. What internal and external differences do you note? How can you distinguish the flower bud from the branch bud by external appearance?

d. Describe fully the difference between these buds and those of the *Prunus domestica* and the *Prunus triflora* in gross appearance, in size, in location and arrangement, and in any other points noted by you.

3. *Prunus americana* includes the following varieties: Hawkeye, De Soto, Weaver, Wood, Wyant, and some others.

a. Secure branches from one of the above varieties, including both fruit buds and branch buds, and make a drawing, natural size, and four inches in length. Note the color of the bark and

A STUDY OF BUDS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

the number of the lenticels. How does the *Prunus americana* differ from the *Prunus domestica* and the *Prunus triflora* in these respects? Describe fully.

b. Make a drawing, at least one inch in length, of both the branch bud and the flower bud. Are the buds arranged on the stem singly, or in clusters at the apex of a spur? Count the buds in each cluster, and note whether the number is constant or variable. Describe a fruit spur, and note how it grows. What is the direction of the growth? Why is this? Where do you find the branch bud? Has the position of the branch bud anything to do with the direction of the growth in the spur?

4. *Prunus hortulana* includes the following varieties: Moreman, Wayland, World Beater, Golden Beauty, and several others.

a. Secure branches from one of the above varieties, including both fruit buds and branch buds. Make a drawing, four inches in length, and natural size, to show a twig which has both kinds of buds. Describe the color of the bark and the general appearance of the lenticels. Note any differences from the preceding groups.

b. Make an enlarged drawing of a fruit bud and a branch bud. Describe how the buds are arranged. Are they larger or smaller than those of the preceding groups? Discuss.

5. *Prunus munsoniana* includes the following varieties: Poole Pride, Pottawattamie, Newman, Robinson, Sophie, Wild Goose.

a. Make a drawing, natural size, and four inches in length, of a twig, including flower buds and branch buds. How do these buds differ from those of the preceding groups? Describe the buds, the lenticels, and the color of the bark.

b. Make an enlarged drawing, one inch in length, of a branch bud and a flower bud. Draw a separate scale of each. How do the scales differ? Are the buds arranged singly or in groups?

Comparative study of the plums. Describe in detail the points of difference between the branch buds and the flower buds in each group. Which are arranged singly, and which are in clusters? Count the ovules in at least ten different buds. Is the number constant? Describe any difference in the longitudinal and the cross sections of the buds in the different groups.

REFERENCES:

Bailey, "The Pruning Book," pp. 21-29, 46-50.

Goff, "Principles of Plant Culture," pp. 86-95.

Paddock and Whipple, "Fruit Growing in Arid Regions," pp. 66-80.

A STUDY OF BUDS
STUDENT'S NOTES AND REPORT

EXERCISE XVIII

A STUDY OF BUDS (*Prunus* sp.)

Material and apparatus. Branches of the sour and the sweet cherry, bearing both branch buds and fruit buds; razor, dissecting lens.

THE CHERRY

There are two large groups of cherries — *Prunus cerasus* and *Prunus avium* — which probably sprang from the European species. They differ in habit of growth, in shape, and in the arrangement of buds, as well as in other characteristics. The domestic forms of *Prunus avium* are characterized by a tall, erect growth, by reddish-brown bark, and by flowers generally in clusters on lateral spurs. The fruit is red, yellow, or black, and sweet. *Prunus avium* has four representatives in the United States — mazzards, hearts, bigarreaus, and dukes. *Prunus cerasus* is characterized by low-headed and spreading forms of trees, by flowers in clusters on lateral spurs, and by thin, willowy branches. There are two representatives in the United States — amarells and morellos. For convenience all cherries are divided into the bigarreaus, or sweet cherries, and the morellos, or sour cherries. These are subdivided according to color and form of fruit, but the subdivisions are of minor importance for consideration at this time.

1. *Bigarreau group.* The bigarreaus, or sweet cherries (*Prunus avium*), include the following varieties: Black Eagle, Black Heart, Black Tartarian, American Heart, Cleveland, Coe's Transparent, Sweet Montmorency, Windsor, Bing, Lambert, Royal Ann, and many others.

a. Secure branches from one or more of the varieties named above, and study thoroughly. Make a drawing, natural size, and at least four inches in length, showing both the branch buds and the fruit buds. Where are they located, and how are they arranged? What external differences do you find between the fruit bud and the branch bud? What bud predominates, and

A STUDY OF BUDS

STUDENT'S NOTES AND REPORT

FIG. 12. Branches showing the branch buds and the flower buds of the sour and the sweet cherry

why? Does the position of the fruit bud differ from that of the branch bud? Is the relative position of the two kinds of buds always the same? Discuss.

b. Make a drawing, at least one inch in length, of a branch bud and a flower bud. Remove a scale from each bud, and draw. Are there any differences between the branch-bud scales and the fruit-bud scales? Describe.

A STUDY OF BUDS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

c. Make a drawing of a longitudinal and a cross section of a branch bud and a fruit bud. Describe the internal differences. Make a dozen or more sections of the flower buds and count the ovules in each. Is the number constant? Can you give any reason for variation? Is there any ratio between the number of ovules in the buds and the number of clusters in the group? Do you find the buds arranged singly, or in groups on short spurs? Where do you find the greatest number of buds, and how old is the wood on which they are borne? Describe the relative position of the branch bud and the fruit bud.

2. *Morello group*. The morellos, or sour cherries (*Prunus cerasus*), include the following varieties: Early Richmond, May Duke, Montmorency, Morello, Royal Duke, Wragg, and several more.

a. Secure branches from one or two of the above-named varieties, or from any other of the sour cherries, and study carefully. Make a drawing of a branch, natural size, and at least four inches in length. Describe the differences between the branches of the sweet and the sour cherry.

b. Make a drawing, at least one inch in length, of a fruit bud and a branch bud. How do these buds differ from those of the sweet cherry? Are the buds arranged singly or in clusters? Do you find spurs on the sour cherries? Describe fully. Now cut cross sections of several fruit buds. How many ovules do you find? Is the number constant, and does it differ from the number in the sweet cherry?

c. Describe fully the differences between the buds of the sweet cherry and those of the sour cherry. Describe the general appearance of a branch of each. Which branch gives the impression of a rigid, stiff, and upright growth? Describe the difference in the bark of the two cherries.

A STUDY OF BUDS

STUDENT'S NOTES AND REPORT

EXERCISE XIX

A STUDY OF BUDS (*Amygdalus* sp.)

Material and apparatus. Branches of the several races of the peach, bearing fruit buds and branch buds ; razor, dissecting lens.

THE PEACH

The peach is divided into five somewhat distinct races. The characteristics which separate each race are more or less pronounced. Each one includes a number of different varieties, some of which are of commercial importance, while others do not rank so high. A comparative study of the buds and branches of each type will aid in bringing out the differences.

1. *North China race.* The North China race (*Amygdalus persica*) includes the following varieties: Albert, Chinese Cling, Elberta, Smock, Spottswood, Thurber, Ford, Family Favorite, and others. The original parent of this race is almost a dwarf. The trees bloom later than the Persian types, and the flowers are nearly always larger. The foliage hangs on well during drought, but sheds off early in the fall. This race is supposed to have originated in northern China, and is adapted to zones north of those suited to the Spanish race.

a. Secure branches from several varieties of this race and make a drawing, natural size, and four inches in length, showing both the flower bud and the branch bud. How are the buds arranged? Describe.

b. Select a fruit bud and a branch bud, taking care that both are mature and perfect, and make a drawing, one inch in length,

FIG. 13. Fruit buds and branch buds of the peach

Note the cluster of buds

A STUDY OF BUDS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

of each. What difference is observed between the two? How are they arranged, and where do you find the branch buds with reference to the fruit buds? Describe the scales and pubescence of the buds?

c. Make a longitudinal and a cross section of both a flower bud and a branch bud. How do the flower buds and the branch buds differ in internal and in external structure? How many ovules are present in a flower bud? Is the number constant or variable? How old is the wood on which the flower buds are found? Does the age of the wood on which the flower buds are borne differ from that of the other fruits studied?

2. *Persian race.* The Persian race (*Amygdalus persica*) includes the following varieties: Lemon, Alexander, Conkling, Crawford's Late, Golden Drop, Old Mixon Free, St. John, Salway, Scott, Mountain Rose, Hale Early, and several others.

The trees of this race are of medium size, with short, thick limbs. The bark on the young wood is usually a rich purplish red. The bloom is variable, and the foliage is crumpled and has a purplish tinge. The trees require a long period of rest. The fruit is highly colored and of good flavor. This race is best adapted to northern sections and is not recommended for Southern planting.

a. Study branches from several of the above-named varieties. Make a drawing, natural size, and four inches in length, showing the details of the branch and of the fruit buds and branch buds. How are the buds arranged, and does this arrangement differ from that of the North China race? What is the color of the bark on the small branches? Is the color constant or variable?

b. Make a drawing, at least one inch in length, of both a flower bud and a branch bud. Compare the buds with those of the North China race in respect to size. Describe.

3. *South China race.* The South China race (*Amygdalus persica*) includes the following varieties: Honey, Pallas, Climax, Coleman, Early China, and others.

The trees are of medium size, and the branches come out at an angle of about fifty degrees and are less willowlike than the Peen-to. The bloom is large and profuse. This race is valuable for Southern planting.

Secure branches from one or more of the above varieties and study carefully. Make a drawing, natural size, and four

A STUDY OF BUDS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

inches in length, of a branch, including both fruit buds and branch buds. Describe the arrangement of the buds. What color is the young wood?

4. *Spanish race*. The Spanish race (*Amygdalus persica*) includes the following varieties: Cobler, Columbia, Victoria, Texas, Sanders, and several others. The trees are very large; the limbs large, long, and spreading; the branches low; the flowers usually large; the foliage small and flat. The fruit is very decided in character, very late, and nearly always yellow. This race is adapted to isothermal lines north of those in which the South China race flourishes.

Secure branches from one or more varieties of this race. Study carefully, and compare with the races already studied. Make a drawing, natural-size, and four inches in length, of a branch, showing the position and arrangement of the buds. Describe any differences in this respect between this race and the others studied. What is the color of the bark? Where do you find the fruit buds?

5. *Peen-to race* (*Amygdalus platycarpa*). The Angel and the Waldo are two varieties which are seedlings of the Peen-to. This is probably the least important race, and the varieties are restricted as to the location in which they will grow. The trees are rather large and vigorous, with willowlike branches and large flowers.

Secure branches from one-year-old and two-year-old wood of the above varieties. Make a drawing, natural size, and four inches in length, of a branch, showing both leaf and branch buds. Describe fully. Compare this race with the South China race, and describe any difference.

Comparative study of the peach. Describe in detail the differences between the races — in the twigs, in the leaf and flower buds, in the color of the branches, in the position and arrangement of the buds, etc. — as seen by your study.

REFERENCES:

Bailey, "The Pruning Book," pp. 50-54.

Bulletin No. 39, Texas Agricultural Experiment Station.

A STUDY OF BUDS

STUDENT'S NOTES AND REPORT

EXERCISE XX

A STUDY OF BUDS—POMACEOUS FRUITS

Material. Branches of several varieties each, of the apple, pear, and quince, bearing both leaf and fruit spurs.

Apparatus. Razor and lens.

Apples and pears are classed with a natural group of plants in which the fruit is more or less fleshy and contains seed cells inclosed in parchmentlike carpels. The apple, pear, and quince, so similar in many respects, belong to this class. This genus is called *Malus*, and includes many species of apples and crab apples. *Malus sylvestris*, the apple, includes many varieties. *Pyrus communis* and *Pyrus sinensis*, the pear, also include many varieties. *Cydonia vulgaris*, the quince, is not so important, but for convenience it is placed with the apple and the pear.

The varieties of all three of these fruits are numerous, and the difference in relationship is very slight.

FIG. 14. Fruit buds and branch buds of the apple

Note the fruit spurs and branch spurs and compare them with those of the pear (Fig. 15)

THE APPLE (*Malus sylvestris*)

Branches of the following varieties are suggested: Northern Spy, Ben Davis, Wolf River, Early Harvest, Fameuse, Jonathan, Lady, Spitzenburg, Duchess of Oldenburg, Pippin, Greening, Russet, Bellflower.

1. Make a drawing, natural size, and four inches in length, of a twig, including the branch buds and leaf spurs, of at

A STUDY OF BUDS—POMACEOUS FRUITS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

least four varieties. Describe the difference between a branch bud and a leaf spur.

2. Draw at least three fruit spurs, including the branch, natural size, and four inches in length. What difference do you note externally between the branch bud and the fruit bud? Is

FIG. 15. Branch buds and fruit buds of the pear

Note the fruit spurs and leaf spurs

there any difference in size, shape, or position of the branch bud and the fruit bud? Describe fully. How are the buds constructed? Study carefully the differences between the leaf spur and the fruit spur. How does the bud on the leaf spur differ from the bud found on one-year-old wood?

3. Tear the buds apart, and draw a separate scale from both the fruit bud and the branch bud. Describe any difference between the scale found on the branch bud and that found on the fruit bud.

4. Now with a very sharp knife make a longitudinal and a cross section of a branch bud and of a fruit bud. Make a separate drawing, at least one inch in length, of each. Can you detect any flowers or embryonic fruit? Describe the difference as seen in a cross

section of both the fruit bud and the branch bud. Describe any varietal differences observed from your study.

THE PEAR (*Pyrus communis* and *Pyrus sinensis*)

Study the pear in the same way as the apple. Varieties suggested: Bartlett, Seckel, Sand pear, President, Kieffer, Anjou, Angoulême.

1. Make drawings, natural size, and four inches in length, showing both the fruit spurs and the leaf spurs of at least four

A STUDY OF BUDS—POMACEOUS FRUITS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

varieties. Is there any difference between the fruit spurs of the pear and of the apple? Describe fully. What differences do you note in the fruit bud and in the branch bud of each. Describe the difference in the color of the bark, in the pubescence, and in the lenticels, etc., as seen in the different varieties.

FIG. 16. Branches of the quince, showing the buds

2. Make a drawing, two inches in length, showing the leaf scars. Explain the cause of these scars.

3. Draw a cross section, at least one inch in length, of a branch bud and a fruit bud. Cut a longitudinal and a cross section of both the fruit bud and the branch bud and describe any differences between the pear and the apple. Note and describe any varietal differences. How do the fruit spurs of the apple and pear differ from those of the cherry and plum?

THE QUINCE (*Cydonia vulgaris*)

The quince is a fruit similar in structure to the apple and the pear. It is borne upon small, irregular-growing trees from ten to twelve feet high, and is a most excellent fruit for preserves.

Study the quince in the same way as the apple and the pear. The following varieties are suggested: Orange, Chinese, Portugal, Champion, Angers.

1. Make a drawing of a branch, four inches in length. How does the branch of the quince differ from that of the apple and the pear?

2. Make a longitudinal and a cross section of a bud, one inch in length. Where are the fruit buds found, and how do they differ in position from those of the apple and the pear? Where are the branch buds found? Describe.

3. Describe fully the difference in the buds of the three fruits in respect to position, structure, size, and shape.

REFERENCE:

Bailey, "The Pruning Book," pp. 30-46.

A STUDY OF BUDS—POMACEOUS FRUITS

STUDENT'S NOTES AND REPORT

EXERCISE XXI

A STUDY OF BUDS — VINE FRUITS

Material and apparatus. Canes of the grape, showing well-developed buds; razor, lens.

THE GRAPE (*Vitis* sp.)

The grape is one of our oldest fruits. The history of the Old World grape goes back to prehistoric times. Grape seeds have been found in the remains of the Swiss lake dwellings of the bronze age, and the original habitat of the grape is supposed to have been about the Caspian Sea. Several species of grapes are recognized, but the two of greatest commercial importance are the *Vitis vinifera*, or the Old World grape, and the *Vitis labrusca*, or the New World grape. There are many hybrids between these two.

FIG. 17. Grape canes, showing the buds

Vitis labrusca. Varieties suggested for study: Black Hawk, Champion, Chautauqua, Colerain, Concord, Cottage, Eaton, Glenfeld, Hosford, Moore Early, Lutie, Lucile, Wyoming, Worden, Vergennes.

1. Secure canes of one or more of the above varieties, and study carefully. Make a drawing of a cane, natural size, including at least two buds. Describe the position and the arrangement of the buds. What is the color of the bark? Describe. Make a drawing, at least one inch in length, of a bud. How does this bud differ from the bud of the cherry and of the apple? Describe the bud scales.
2. Make a longitudinal and a cross section of a bud. Describe the internal structure. Is this a fruit bud or a leaf bud? Where is the fruit borne on the grape? Describe fully.

A STUDY OF BUDS—VINE FRUITS

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

3. Make a longitudinal section of a cane, through a node, and show by a drawing the internal structure. Label the pith, bark, and wood. What proportion does the bark bear to the pith? Can you locate the diaphragm which is found at the node, after a longitudinal section of the cane is made? Label. Describe.

Vitis vinifera. Varieties suggested for study: Black Hamburg, Tokay, and Malaga.

1. Secure canes from one of the above varieties, and study them critically. Make a drawing, natural size, of a cane, including at least two buds. Describe the buds and the bark. Make a drawing, at least one inch in length, of a bud. Describe the difference between the *labrusca* and the *vinifera*. What differences do you observe in the bud and the cane of each? Where are the *vinifera* types of grapes grown?

2. Make a longitudinal section of a cane, showing the diaphragm. Does the diaphragm of the *vinifera* differ from that of the *labrusca*? Describe.

Vitis labrusca $\times$ *vinifera*. Varieties suggested for study: Brighton, Campbell Early, Catawba, Diamond, Essex, Goethe, Hartford, Herbert, Iona, Isabella, Lady Washington, Jefferson, Woodruff.

This group includes many good varieties of grapes which have originated by the crossing of the *vinifera* with the *labrusca*. The characteristics are derived from both of the parents, but some varieties show more characteristics from the male, others more from the female.

1. Secure canes from one or more of the above varieties, and study carefully. Make a drawing of a cane, including at least two buds. How do these buds differ from those of *Vitis labrusca* and *Vitis vinifera*? Can you see any resemblance to either parent? Study carefully, and note which parent they seem to resemble the most. Make a drawing, at least one inch in length, of a bud. Describe.

2. Draw a longitudinal section of a cane, showing the diaphragm. Describe. Describe in detail the differences between the *Vitis labrusca*, the *Vitis vinifera*, and the cross between the two.

REFERENCES:

- Bulletin No. 193*, California Agricultural Experiment Station.
Circular No. 16, New York Agricultural Experiment Station.

A STUDY OF BUDS—VINE FRUITS

STUDENT'S NOTES AND REPORT

EXERCISE XXII

PRUNING

Material and apparatus. Small orchard; pruning-saw, pruning-knife, pruning-shears, long-handled pruner.

After the student has become thoroughly familiar with the different kinds of buds of the several fruits and is able to distinguish a fruit bud from a branch bud, or a fruit spur from a leaf spur, several exercises should be given in the general practice of pruning. It is advisable to have at least one exercise each on the pruning of the apple, the cherry, the peach, and the grape. Study carefully each kind of plant pruned. Discuss fully what is the essential thing for which to look. Discuss the differences between the fruit bud and the branch bud as seen from your practice in pruning. Discuss fully what kind of a cut should be made in every case. Discuss why it is essential, when cutting a large limb, to make a cut first on the underside. Discuss why the peach must be pruned differently from the apple. Tell why the grape should be pruned back to three or four buds. On what age of wood are the fruit buds of the peach, the pear, the grape, and the cherry usually found. Compare several of the different kinds of pruning-saws. Give the advantages and the disadvantages of each. Tell where the pruning-shears should be used and where they should not be used.

REFERENCES :

Bailey, "The Pruning Book," chap. iv.

Farmers' Bulletin No. 181, United States Department of Agriculture.

Paddock and Whipple, "Fruit Growing in Arid Regions," pp. 81-100.

Thomas, "The American Fruit Culturist," chap. vii.

PRUNING

STUDENT'S NOTES AND REPORT

FIG. 18. Pruning tools

1, 2, 3, 4, pruning-saws; 5, 6, pruning-knives; 7, 8, 9, 10, pruning-shears

PRUNING

STUDENT'S NOTES AND REPORT

EXERCISE XXIII

THE PRUNING OF THE APPLE

Material and apparatus. Small orchard; pruning-saw, pruning-knife.

Discuss the pruning of the apple and give the reason for each step. Discuss how a one-year-old tree must be pruned; a two-year-old tree; a five-year-old tree; a mature tree. How is a low-headed tree formed? When should the head be started? Discuss. Give the advantages and the disadvantages of a low-headed and a high-headed tree.

• REFERENCES :

Thomas, "The American Fruit Culturist," pp. 98-109.

Waugh, "The American Apple Orchard," chap. ix.

THE PRUNING OF THE APPLE

STUDENT'S NOTES AND REPORT

EXERCISE XXIV

THE PRUNING OF THE PEAR

Material and apparatus. Small pear orchard; pruning-saw, pruning-knife.

Discuss the pruning of a one-year-old tree, a two-year-old tree, a ten-year-old tree, and a mature tree. How does the pruning of the pear differ from that of the apple? Discuss in detail.

REFERENCES:

Bailey, "The Pruning Book," pp. 269-278.

Paddock and Whipple, "Fruit Growing in Arid Regions," pp. 133-138.

THE PRUNING OF THE PEAR

STUDENT'S NOTES AND REPORT

EXERCISE XXV

THE PRUNING OF THE PLUM AND THE CHERRY

Material. Small plum and cherry orchard.

Apparatus. Pruning-saw, pruning-knife.

Discuss the pruning of the plum and the cherry. What different methods are practiced on the two kinds of trees? How do these methods of pruning differ from those of the pear and the apple?

REFERENCES :

Paddock and Whipple, "Fruit Growing in Arid Regions," pp. 139-141.
Thomas, "The American Fruit Culturist," p. 109.

THE PRUNING OF THE PLUM AND THE CHERRY

STUDENT'S NOTES AND REPORT

EXERCISE XXVI

THE PRUNING OF THE PEACH

Material and apparatus. Small peach orchard; pruning-saw, pruning-shears.

Give in detail the proper method to follow in the pruning of the peach. How does this differ from the method used for the apple and the cherry? Where is the fruit-bearing wood found? Discuss.

REFERENCES:

Bulletin No. 159, Maryland Agricultural Experiment Station, pp. 121-136.

Bulletin No. 55, Missouri Agricultural Experiment Station.

Paddock and Whipple, "Fruit Growing in Arid Regions," pp. 102-110, 127-132.

THE PRUNING OF THE PEACH

STUDENT'S NOTES AND REPORT

EXERCISE XXVII

THE PRUNING OF THE GRAPE

Material and apparatus. Small vineyard; pruning-shears, pruning-knife.

Discuss in detail the pruning of the grape. Where are the fruit buds found? Do they occur on the present season's wood or later? Tell why it is advisable to prune the grape to three or four buds. Does this method of pruning reduce the yield of the crop? Is it an advantage or a disadvantage to use this method? Discuss. Describe in detail each of the following methods:

1. Renewal system of pruning.
2. Marvin's system of fan pruning.
3. Kniffin system of pruning.
4. Y-trunk Kniffin system of pruning.
5. Double Kniffin system of pruning.
6. Overhead Kniffin system of pruning.

REFERENCES:

Bailey, "The Pruning Book," chap. viii.

Bulletin No. 193, California Agricultural Experiment Station.

Circular No. 16, New York Agricultural Experiment Station.

Year Book of the Department of Agriculture, 1896, pp. 499-542.

THE PRUNING OF THE GRAPE

STUDENT'S NOTES AND REPORT

EXERCISE XXVIII

THE PRUNING OF THE BRAMBLES

Material and apparatus. Small patch of brambles; bramble hooks and pruning-shears.

The blackberry, raspberry, and dewberry are classed as brambles. The method of pruning such plants is very different from the method used for tree fruits. Discuss the best method of procedure. What canes should be removed? Is there any disadvantage in using the bramble hook? If so, discuss. When should the brambles be pruned, and what canes should be removed?

REFERENCES:

Bailey, "The Pruning Book," pp. 323-327.

Farmers' Bulletin No. 213, pp. 15-16.

THE PRUNING OF THE BRAMBLES

STUDENT'S NOTES AND REPORT

ALL RIGHTS RESERVED

PRINTED BY THE AUTHOR

THE PRUNING OF THE BRAMBLES
A STUDENT'S NOTES AND REPORT
ON THE HISTORY AND DEVELOPMENT OF THE
BRAMBLE PRUNING INDUSTRY
IN THE UNITED STATES
BY
J. H. BROWN
B. S. AGRICULTURE
UNIVERSITY OF CALIFORNIA
DAVIS, CALIF. 1914

THE PRUNING OF THE BRAMBLES
A STUDENT'S NOTES AND REPORT
ON THE HISTORY AND DEVELOPMENT OF THE
BRAMBLE PRUNING INDUSTRY
IN THE UNITED STATES
BY
J. H. BROWN
B. S. AGRICULTURE
UNIVERSITY OF CALIFORNIA
DAVIS, CALIF. 1914

EXERCISE XXIX

THE PRUNING OF THE BUSH FRUITS

Material and apparatus. Small patch of currants and gooseberries ; pruning-knife and pruning-shears.

The currant and the gooseberry are two of the most important of the bush fruits. The method of pruning these fruits is different from that used for tree fruits. After some practice in the field discuss the best ways in which the bush fruits are pruned. What wood should be removed each year? Should the plants be severely headed back, or thinned out by the removal of the older canes? Is it wise to keep approximately a definite number of canes in each clump? Discuss.

REFERENCES :

Bailey, "The Pruning Book," pp. 327-333.

Bulletin No. 357, New York Agricultural Experiment Station.

THE PRUNING OF THE BUSH FRUITS

STUDENT'S NOTES AND REPORT

EXERCISE XXX

THE STUDY OF SHIELD AND PRONG BUDDING

Material. Bud sticks, stocks of the peach or willow (preferably steamed, if the exercise is practiced in the laboratory).

Apparatus. Raffia or grafting-cord, budding-knife.

Budding is the operation of applying a single bud, bearing little or no wood, to the surface of the growing wood of the stock. The bud is applied directly to the cambium layer of the stock. It is usually inserted under the bark, but in some forms of budding the bud is used to cover the wound made on the stock. Plants with thin bark and plenty of sap are likely to do better when budded than plants with thick, heavy bark.

FIG. 19. Shield budding

A, the stock, showing the longitudinal and the transverse cut; B, the bud removed from the bud stick and ready to be inserted into the stock; C, the stock, showing the bud in position ready for tying

SHIELD BUDDING

This is the style of budding that is in general use, and the one that

is the most often practiced. It is known as shield budding, from the shieldlike shape of the portion of bark bearing the bud. Secure small branches, about three eighths of an inch in diameter, of either the peach or the willow. If the exercise is

THE STUDY OF SHIELD AND PRONG BUDDING

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

desired in the winter, the steaming of the wood will be found to loosen the bark so that it will slip. Now hold the branch firmly with the left hand and with the right make a clean, sharp cut, with a budding-knife, through to the wood and one third of the distance around the twig. Now, by another operation, make a straight cut, about one inch in length, at right angles to the one just finished. The two cuts now form a letter T (see Fig. 19). With the budding-knife carefully peel the bark back, down to the wood, thus breaking the cambium layer. Take a bud stick

FIG. 20. Prong bud ready to be inserted into the stock

which has previously been procured and, holding it in the left hand, carefully cut a bud from the stick. The bark, including the bud, should be about one inch in length. It should contain as little as possible of the wood of the stick. Now insert the bud into the cut made in the stock, place the bark over the bud, and tie the whole firmly with raffia or grafting-cord.

Make three drawings, four inches in length, to show first, the longitudinal and the cross cut; second, the bud in place; and third, the bud wrapped. Make two natural-size drawings of the bud after it has been removed from the bud stick, the first a front view, showing the shieldlike shape of the portion of the bark; the second a side view, showing the width of the bark bearing the bud. Label all drawings.

PRONG, OR TWIG, BUDDING

This is a modification of shield budding in which a short prong, or spur is used in the place of a simple bud. The bud is cut in essentially the same manner as for shield budding.

Secure for your bud stick small branches of the walnut tree, or any other tree having short prongs, or spurs. By the same operation as in shield budding remove the prong bud. The incision into which the prong bud is placed is identical with that of the shield bud. Make two drawings, natural size, of the prong bud, showing the front and side views. Label. Discuss.

REFERENCES:

Bailey, "The Nursery Book," pp. 94-107.

Goff, "Principles of Plant Culture," pp. 230-234.

Farmers' Bulletin No. 157, pp. 21-23.

THE STUDY OF SHIELD AND PRONG BUDDING

STUDENT'S NOTES AND REPORT

U.S. GEOLOGICAL SURVEY

Geological Survey of the United States

Washington, D.C.

1900

By [Name]

[Faint, illegible text covering the main body of the page, likely containing student notes and a report.]

EXERCISE XXXI

THE STUDY OF PLATE BUDDING AND H-BUDDING

Material. Bud sticks, stocks of the peach or willow.

Apparatus. Budding-knife, raffia or grafting-cord.

PLATE BUDDING

This form of budding is used on certain plants where the shield bud is not successful.

FIG. 21. Plate bud, showing the stock with the bark turned back and the bud ready for insertion

Secure branches of either the peach or the willow, about one-half inch in diameter or a trifle larger and of any convenient length. With the budding-knife make a rectangular incision, three fourths of an inch in length and one fourth to three eighths of an inch in width, in the stock through the bark to the wood (Fig. 21), and turn down the flap of bark. Now cut a section of bark including a well-formed bud, similar in shape to the rectangular section in the stock. The bud on the bark should be about twice as far from the bottom as from the top of the bark. No wood should be attached to the bark containing the bud. Now place the bud under the flap of bark so that the cambium layer of the bud and the stock are in close apposition, and turn the flap over the bud and tie firmly. The operation is now complete.

Make a drawing, four inches in length, showing the stock with the flap of bark cut and turned down. Label all parts. Make a drawing of the bud twice its natural size, and label. Make another drawing, four inches in length, showing the stock with the bud in place and wrapped. Describe fully each operation. On what kind of plants is this form of budding used? Discuss.

THE STUDY OF PLATE BUDDING AND H-BUDDING

STUDENT'S NOTES AND REPORT

H-BUDDING

This form of budding is a modification of plate budding. In H-budding a flap is formed both above and below the horizontal cut on the stock, which covers the bud from both ends. This allows the more perfect fitting of the bark around the bud.

FIG. 22. H-Budding, showing the stock and the bud

Secure a small branch, similar to that used in plate budding, and make the same cut as in plate budding, but, instead of forming one flap, continue your longitudinal cuts equal distances on each side of the transverse cut. Now with the budding-knife peel the bark in opposite directions from the transverse cut, forming the letter H (Fig. 22). Proceed to cut your bud, including a piece of bark the size of the opening made by turning back both flaps of the bark on the stock. Cut the bark including the bud in such a manner that the bud is in the center of the bark. Now place the cambium layer of the bud and of the stock together and tie the two flaps of bark on the stock firmly around the bud. At least a dozen trials should be made to insure good technique and ease of operation. Make a

drawing of the stock, four inches in length, showing the cut, with both flaps of bark turned back. Make a drawing of the bud, showing the front and the side view. Label. Describe fully each operation.

THE STUDY OF PLATE BUDDING AND H-BUDDING

STUDENT'S NOTES AND REPORT

EXERCISE XXXII

THE STUDY OF FLUTE BUDDING, RING BUDDING, AND CHIP BUDDING

Material. Bud sticks, stocks of the peach or willow.

Apparatus. Raffia or grafting-cord, budding-knife.

FLUTE BUDDING

Flute budding is a form of budding occasionally employed on certain plants. It differs from the other forms in that a portion of the bark is removed entirely from the stock and a similar piece is fitted into its place. The section of bark removed extends only part of the way around the stem (Fig. 23).

FIG. 23. Flute budding, showing the stock and the bud

Secure a small branch about one half to five eighths of an inch in diameter, and with the budding-knife remove a section of the bark about one inch in length and about one third of the diameter of the stock in width. The cut necessarily varies somewhat in size, according to the size of the stock which is to be budded, but the usual size is about one inch in length and from one third to one half of the distance around the stock. Now remove a piece of bark containing a healthy, vigorous bud and cut it to the exact size of the wound on the stock. Do not remove any wood with the bark containing the bud. Place the bud over the wounded surface on the stock and tie it firmly. The operation is now complete. Make a drawing of the stock,

four inches in length, showing the front and side views. Label all parts, and give the dimensions of the cut surface. Make a natural-size drawing of the bud. Label and describe.

FLUTE BUDDING, RING BUDDING, AND CHIP BUDDING

STUDENT'S NOTES AND REPORT

[Faint, illegible text, likely bleed-through from the reverse side of the page.]

LABORATORY MANUAL OF HORTICULTURE

RING, OR ANNULAR, BUDDING

This is a modification of flute budding, the difference being that the portion of bark removed extends entirely around the stock, forming a ring; hence the name. The same size stocks should be used as for flute budding.

Secure stocks of the same size as in flute budding, and with the budding-knife remove a ring of bark, about one inch in length, from the stock. Secure the plant from which you wish to bud, and remove a piece of the bark bearing a healthy bud, the same size as the wound on the stock. The bark bearing the bud should be split lengthwise on the side opposite the bud. The bud should occupy the central portion of the bark. Why? Make a drawing of the stock, natural size, and four inches in length, showing the portion of bark removed. Label, and give the dimensions of the bud. Make a natural-size drawing of the bud, showing the ring character. Label. Discuss the advantages and the disadvantages of this method.

FIG. 24. Ring budding
A, stock, showing the
ring of bark removed;
B, bud ready to be placed
on the stock

CHIP BUDDING

This is a form of budding that is used when the stock is dormant and the bark will not slip. The chip bearing the bud is held in place by tying or, still better, by the use of grafting-wax.

Secure small stocks the same as those employed in flute budding and make an incision into the stock similar to that in Fig. 25. The cut in the stock should be smooth, clean, and even,

FIG. 25. Chip budding, showing the cut in the stock and the bud ready to be inserted

FLUTE BUDDING, RING BUDDING, AND CHIP BUDDING

STUDENT'S NOTES AND REPORT

and about one inch in length and from three eighths to one half of an inch in width. The depth of the cut should not exceed one eighth of an inch, and the bud when inserted should be flush with the bark of the stock. Now with the budding-knife remove the bud with sufficient wood so that it will fit snugly into the cut made in the stock. It is important that at least two sides of the bud should be in close apposition with the sides of the stock and that the cambium layers of the bud and the stock should meet at several points.

Make a drawing of the stock, four inches in length, showing the front and the side views. Draw a front and a side view of the bud, give the dimensions, and label. Discuss the operation and tell what kind of plants are propagated by this method.

FLUTE BUDDING, RING BUDDING, AND CHIP BUDDING

STUDENT'S NOTES AND REPORT

FLUTE BUDDING

STUDENT'S NOTES AND REPORT

The first step in flute budding is to select a healthy, well-developed scion. The scion should be cut from a branch that is free from disease and insect damage. It should be about 12 inches long and 1/2 inch in diameter. The scion should be cut at an angle of 45 degrees at both ends. The bottom end should be cut with a sharp knife, and the top end should be cut with a pruning shears. The scion should be kept in a cool, moist place until it is ready to be budded.

STUDENT'S NOTES AND REPORT

The next step in flute budding is to prepare the stock. The stock should be a healthy, well-developed branch that is free from disease and insect damage. It should be about 12 inches long and 1/2 inch in diameter. The stock should be cut at an angle of 45 degrees at both ends. The bottom end should be cut with a sharp knife, and the top end should be cut with a pruning shears. The stock should be kept in a cool, moist place until it is ready to be budded.

STUDENT'S NOTES AND REPORT

The final step in flute budding is to attach the scion to the stock. The scion should be inserted into the stock, and the two should be secured with a rubber band. The rubber band should be placed around the scion and the stock, and it should be tightened so that it is secure. The rubber band should be removed after the scion has been attached to the stock.

EXERCISE XXXIII

THE STUDY OF GRAFTING-WAXES

Material. Resin, beeswax, rendered beef tallow, raw linseed oil, white resin, alcohol, turpentine, Burgundy pitch, red ocher, white pitch, Venetian red, No. 16 white knitting-cotton.

Apparatus. Granite kettle, small pail, cooking-burner.

Grafting-waxes are used for the protection of wounds and for the covering of wounded surfaces when certain kinds of grafts are made. Grafting-waxes are made by the following methods:

BEESWAX AND RESIN WAXES

Formula 1. Weigh out one pound of resin, one half of a pound of beeswax, and one fourth of a pound of rendered beef tallow. Place the ingredients in a granite or an iron kettle and allow them to melt, then boil for about one minute, or until the mass is thoroughly mixed. Remove from the fire, and pour the melted mass directly into a pail of cold water. When the mass is cold enough to handle, gather it into a ball and work it with the hands until it assumes a light-brown color. Grease the hands thoroughly with tallow before working, to prevent the wax from sticking to them. Why is the melted wax poured into the water? What is the purpose of the molding and working with the hands? It is important that pure beeswax be used and not wax which has been adulterated with paraffin. Why? Does the wax aid in the healing of a wound? Discuss.

Formula 2. Weigh out one pound of resin, one fourth of a pound of beeswax, one fourth of a pint of raw linseed oil. Place the ingredients in a granite kettle and allow them to melt gradually. When the mass is thoroughly melted and incorporated, pour it into a bucket containing cold water, and treat as above. After twenty-four hours examine both batches and describe fully the differences, if any. Explain.

Grafting-cord. Often in the making of root grafts and occasionally in budding, grafting-cord is needed. This is made

THE STUDY OF GRAFTING-WAXES

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

by securing a ball of No. 16 knitting-cotton and placing it in the kettle of melted wax of Formula 1. Allow the ball of cotton to remain in the melted wax for at least five minutes, or until it is completely saturated. The ball should be turned frequently, in order to insure complete saturation. When finished, allow it to cool, and wrap in oiled paper for future use.

Why is cord prepared in this way superior for grafting purposes to the ordinary wrapping-twine?

ALCOHOLIC WAXES

Formula 3. Weigh out one and one-half pounds of white resin and one fourth of a pound of beeswax. Place the ingredients in a granite kettle and heat them until they are thoroughly incorporated. Remove the mass from the fire and stir until partially cooled. Now gradually add enough alcohol to give to the mass, when cooled, the consistency of thick paint. Describe the color and the appearance of this wax and tell where it can best be used.

Formula 4. Weigh out one-half pound of white resin and one-half ounce of rendered beef tallow. Heat the two substances together until they are incorporated. Remove from the fire, cool, and add one teaspoonful of turpentine. Stir the mass until thoroughly mixed, and gradually add enough alcohol to make the mass assume the consistency of thick honey. (From six to eight fluid ounces of 95 per cent alcohol will usually be sufficient.)

What are the differences between formulas 3 and 4? Describe the appearance of the two waxes, and give your opinion as to which is the best.

PITCH WAX

Formula 5. Weigh out twelve ounces of resin and six ounces of Burgundy pitch and melt them together in a granite kettle, stirring thoroughly. At the same time melt three ounces of rendered beef tallow in another vessel. Pour both liquids together while hot, and stir thoroughly. To this mass add four or five ounces of red ocher, and stir the mixture until it is smooth and until the different ingredients are completely incorporated. Describe the mass and tell when and where it should be employed.

NOTE. The linseed oil should be entirely free from cottonseed oil. These grafting-waxes may be made by groups of from two to four students.

STUDENT'S NOTES AND REPORT

EXERCISE XXXIV

THE STUDY OF SPLICE, WHIP, AND DOUBLE-WHIP GRAFTING

Material. One-year-old apple-seedling roots, scions from apple trees.

Apparatus. Sharp budding-knife, raffia or grafting-cord.

Grafting is the operation of inserting into a stock a scion or twig comprising one or more buds. It is classified in various ways, chiefly with reference to the method in which the scion and the stock are joined.

The union of the scion with the stock depends upon the growing together of the cambium tissue of the two plants. This tissue is the living active cells found between the bark and the wood. It is composed of cells which are actively dividing. The layer of cambium tissue is very narrow, being only four or five cells in thickness. It is very essential that careful and painstaking work be done and that the student be thoroughly familiar with the position the cambium layer occupies in both the stock and the scion.

The scions are cut in the late fall, in winter, or at any time before the buds swell in the spring. It is thought, however, that the later the scions are taken from the tree before the buds swell, the greater will be the success in grafting, other things being equal. Each scion should contain from three to five well-developed buds, and it should be cut about one-half inch above and below a bud. Only the previous year's growth is used in all ordinary cases, but an exception to this is found in the maples and several of the ornamentals in which the older wood may be used.

The scions should be stored in moss, sand, or sawdust and kept in a cool place until they are used to prevent drying out or swelling of the buds. This is very important for successful grafting. Only well-formed branch buds should be used.

SPLICE GRAFTING

Splice grafting represents the simplest form of grafting. In this form the scion and the stock should be as nearly the same size as it is possible to get. In splice grafting it is well to use

SPLICE, WHIP, AND DOUBLE-WHIP GRAFTING

STUDENT'S NOTES AND REPORT

one-year-old apple seedlings, and scions taken from apple trees. The two parts, namely, the stock and the scion (Fig. 26), are simply cut diagonally across and the two firmly tied together.

FIG. 26. Splice grafting, showing the relative size of the stock and the scion

The cut should be one inch to one and one-half inches across. This form of grafting is frequently employed upon plants with soft or tender wood which will not admit of splitting.

Make a drawing, natural size, and three inches in length, of both the stock and the scion, showing the parts in position ready for tying. How long did you make the diagonal cut? Place the dimensions on your drawings. To what kind of stock is this method of grafting best adapted? Practice by making a number of these grafts. Care must be taken that the cambium layers are in apposition. Why is this important? Discuss.

WHIP, OR TONGUE, GRAFTING

This form of grafting is employed only on small stocks, one or two years old. The stocks are usually seedling roots, and the grafting is usually performed in the winter. Both the scion and the stock (Fig. 27) are cut diagonally across, as in splice grafting, and the cut should be at least one to one and one-half inches across, according to the size of the stock. A vertical cleft is then made in both the stock and the scion, and the two are joined by shoving the tongue of the scion into the cleft of the stock. The two parts are held together by tying with raffia or grafting-cord. If the wound stands above the ground, it must be protected by applying grafting-wax over the bandage.

FIG. 27. Whip grafting
A, scion, showing the diagonal cut and the split; B, scion inserted into the stock

SPLICE, WHIP, AND DOUBLE-WHIP GRAFTING

STUDENT'S NOTES AND REPORT

 

LABORATORY MANUAL OF HORTICULTURE

1. Make a natural-size drawing of the root, showing the diagonal cut and the cleft. How long should this cut be? Place the dimensions on the drawing, and describe how the cut, which should be clean and even, is best made. Discuss.

2. Make a natural-size drawing of the scion, showing the cut and the cleft. The scion should contain three or four well-developed and plump buds. How long is the scion? Label all parts, and place the dimensions on your drawings.

3. Show by a natural-size drawing the stock and the scion tied in position. What size of stocks is best suited to this form of

grafting? What commercial operation depends upon this method of propagation? What superiority, if any, does a tree produced by this method have over a tree produced by budding? When is the best time of the year to do root grafting, and why? Why is it essential to have at least one side of the stock and the scion flush with each other? Where do you find the cambium layer on the stock and the scion? Indicate on your drawing by an arrow the location of the cambium tissue.

FIG. 28. Double-whip grafting

A, scion; *B*, stock, showing the double split; *C*, scion inserted into the stock

DOUBLE-WHIP GRAFTING

This is a modified form of whip grafting in which two clefts are made instead of one (Fig. 28). The scion, having two supports in the stock, forms a more intimate contact with it. This method is usually too slow for general practice, but is good in special cases, particularly with some plants that are a little hard to graft.

Make a natural-size drawing of the stock and the scion.

Make a drawing of the stock with scion in position. Describe.

REFERENCES :

Bailey, "The Nursery Book," chap. v.

Goff, "Principles of Plant Culture," pp. 219-230.

SPLICE, WHIP, AND DOUBLE-WHIP GRAFTING

STUDENT'S NOTES AND REPORT

EXERCISE XXXV

THE STUDY OF CLEFT GRAFTING AND BARK GRAFTING

Material. Stocks cut from apple or pear trees from one to two inches in diameter and from twelve to fifteen inches in length.

Apparatus. Grafting-chisel, sharp knife, grafting-wax.

These two forms of grafting are used principally for the top-working of old trees or for changing the variety of a large tree of

inferior flavor. Occasionally they are used to replace large broken limbs.

Cleft grafting is the form universally used in top-working or grafting in the limbs of a tree. It is limited to branches less than two inches in diameter, and is best suited to

those between one inch and one and one-half inches in diameter.

The two essential parts in this form of grafting are the *stub* (Fig. 29), or that part of the branch on which the grafting is performed, and the *scion* (Fig. 33). The scion is a twig taken from one-year-old wood, and should contain three or four strong, healthy buds. It is usually from four to six inches in length.

The scion is cut wedge-shaped, and the length of the wedge should be not less than one inch, but the cut can vary from one to one and one-half inches. The cut must be smooth, and made by two draws of the knife. The outer edge is usually a little thicker than the inner one, to insure the scion being held firmly.

FIG. 29. A very good type of grafting-chisel, showing the first position in cleft grafting

FIG. 30. The second position of the grafting-chisel in cleft grafting

CLEFT GRAFTING AND BARK GRAFTING

STUDENT'S NOTES AND REPORT

CHAPTER I

The first step in grafting is the selection of the stock and the scion. The stock should be a healthy, well-developed plant of the species to be grafted. The scion should be a young, healthy shoot of the same species, or of a different species, which is to be grafted onto the stock. The scion should be cut in the winter, before the buds have opened, and should be kept in a cool, moist place until it is ready to be grafted.

The next step is to make a cleft in the bark of the stock. This is done by using a cleft grafting knife to make a vertical incision in the bark, about 2-3 inches long. The incision should be made in a place where the bark is thin and the wood is not too hard. The cleft should be made deep enough to reach the cambium layer of the wood.

After the cleft has been made, the scion is inserted into it. The scion should be cut at an angle, so that it fits snugly into the cleft. The scion should be inserted so that its cambium layer is in contact with the cambium layer of the stock. The graft is then secured by wrapping the cleft with a grafting tape or bandage. The graft should be kept in a cool, moist place for a few days before it is planted in the soil.

LABORATORY MANUAL OF HORTICULTURE

The limb, or stub, is split either by a grafting-chisel (Fig. 30), of which two kinds are in common use, or by the ordinary wood chisel. The essential requisite is to have the knife equipped with a curved edge so as not to split the bark but rather to cut it, and this is best accomplished by a curved blade.

CLEFT GRAFTING

1. Practice making several clefts and inserting the scions.

Make a drawing, four inches in length and one and one-half inches in diameter, showing a stub and cut with the chisel in

FIG. 31. Cleft graft, showing the scions in position in the stub before it is waxed over

position. Show the length of the split along the side. Does this vary with each graft? What is the reason for this variation? Where is the cambium layer in the stub found?

Cut your scions, and insert two in each stub, slanting them outward at an angle of four or five degrees (Fig. 31). Now wax the stub over with grafting-wax. The wax should extend over the edge about a quarter of an inch and completely fill the split on both sides. Why?

2. Make a natural-size drawing of the scion. What is the distance to the first bud? Why should a small

portion of the wood be left above the last bud? The top of the wedge-shaped cut should be near a bud. Why this precaution? Where is the cambium layer in the scion? How do you insure the cambium layer of the scion coinciding with that of the stock? Discuss.

3. Make a drawing, four inches in length, showing the scions in the stub after they have been waxed over (Fig. 32). The two scions should slant outward at an angle of about five degrees. Why? What is the purpose of the wax?

CLEFT GRAFTING AND BARK GRAFTING

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

BARK GRAFTING

Bark grafting is employed on limbs that are too large for cleft-grafting. Limbs over two inches in diameter are best grafted in this way. Bark grafting is often employed to replace large limbs which have been destroyed.

The essential principle in this form of grafting is the same as that involved in cleft grafting. From three to twelve scions are placed in each stub, according to its size.

FIG. 32. Cleft graft, showing the stub after it has been waxed over

FIG. 33. Scion for bark and cleft grafting

The scions are cut very thin, with a shoulder on each side (Fig. 33). The cut should be at least one inch in length and not much more than one sixteenth of an inch in width. Why should this cut be so thin? Make a natural-size drawing of the scion, showing the shoulder. The scion should contain three or four well-developed buds.

1. Make a drawing, at least four inches in length and two and one-half inches in diameter, showing the stub and the location of each scion. In this method the scions are placed between the bark and the wood, and no cleft is made in the stub. A small penknife

CLEFT GRAFTING AND BARK GRAFTING

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

is convenient in making an opening into which the scion can be inserted. Why are the scions placed between the bark and the wood instead of being placed in a cleft in the stock? After the scions have been placed in the stock, wax the whole surface over with grafting-wax, allowing the wax to extend over the edge of the stub about one fourth of an inch.

2. Make a drawing, four inches in length, showing the scions in place, after the whole surface has been waxed over with grafting-wax. Label all parts. When is this form of grafting used? Discuss.

CLEFT GRAFTING AND BARK GRAFTING

STUDENT'S NOTES AND REPORT

EXERCISE XXXVI

THE STUDY OF BRIDGE GRAFTING

Material. Stocks of the apple, three inches in diameter and twelve inches in length, scions from the same kind of trees.

Apparatus. Sharp knife, grafting-wax.

Bridge grafting is used to save girdled trees and for the healing over of wounds that have been made by cultivating-implements.

The wounded part is first thoroughly cleaned out (Fig. 34, *A*) and the bark cut back until it is smooth. The scions are cut wedge-shaped at both ends (Fig. 34, *B*) and varying in length according to the size of the wounded area, and are then pushed up firmly under the bark. They should be about an inch longer than the wound. From three to six scions are used, according to the size of the wound.

FIG. 34. Bridge grafting

A, stock, showing scions in position ready for waxing; *B*, scion, showing wedge at each end

1. Make a drawing, two inches in diameter, showing the injury to the limb. Show by dotted lines the position of the cut which must be made in order to smooth up the edges. Why is it important to cut this bruised tissue back to firm bark? Discuss.

2. Make another drawing, the same size as the above, showing the scions set in place and the whole surface waxed over. Label and describe.

Make a drawing of a scion. How many buds are found?

THE STUDY OF BRIDGE GRAFTING

STUDENT'S NOTES AND REPORT

EXERCISE XXXVII

THE STUDY OF SIDE GRAFTING

Material. Stocks of the apple or pear, one inch in diameter and twelve inches in length, scions from the same kind of trees.

Apparatus. Sharp, narrow chisel and sharp knife.

Side grafting is the method of grafting by inserting the scion into the stock without the cutting off of the stock (Fig. 35). It

FIG. 35. Side grafting

A, stock, showing the chisel in position; B, stock, showing the scion inserted

is often used to place a limb in a certain region where one is lacking, or for the grafting of some difficult subject, such as the mulberry or some of the beeches. The scion is placed in the side of the stock, set into an oblique incision made for the purpose. This cut can best be made by a thin-bladed chisel with a bent shank; however, any ordinary knife or chisel may be used. The angle at which the scion is placed is usually about 30 degrees. The incision should be made about an inch deep, and the scion should be cut wedge-shaped. As in bark grafting, the

length of the wedge-shaped cut should correspond to the depth of the incision made in the stock.

1. Secure stocks about one inch in diameter and ordinary scions containing from two to four buds. Practice by making a number of these grafts.

Make a natural-size drawing of the stock, four inches in length, showing the chisel or the knife in position.

2. Make a drawing the same size as in 1, showing the scion in position. How long is the wedge-shaped cut on the scion? Does the entire wedge of the scion fit into the incision made in the stock? Make a natural-size drawing of the scion. Discuss.

THE STUDY OF SIDE GRAFTING

STUDENT'S NOTES AND REPORT

EXERCISE XXXVIII

THE STUDY OF VENEER GRAFTING

Material. Stocks of the apple or pear from one half to three fourths of an inch in diameter.

Apparatus. Grafting-cord or raffia, sharp knife or chisel.

This form of grafting is adapted to ornamentals and rare stocks which are usually grown in pots. In this method of grafting an

incision is made into the wood, and all of the wounded surfaces are completely covered by the matching of the scion and the stock. No grafting-wax is used. This method is employed from November until March, and the scions are applied an inch or two above the surface of the soil. Both dormant and growing scions are used.

To make a veneer graft, an incision from one to one and one-half inches in length is made upon the stock, just through the bark, and the small piece of bark is removed by a downward-sloping cut at the base (Fig. 36). The base of the scion is cut off obliquely, and upon the longest side a portion of the

FIG. 36. Veneer grafting

A, stock, showing the cut; *B*, scion, showing the cut at the base; *C*, scion and stock united and tied

bark is removed, corresponding to the part taken from the stock. The scion is then tied tightly to the stock.

1. Secure branches of the apple or the pear, from about one half to three fourths of an inch in diameter, and practice by making several of these grafts. Make all cuts clean and sharp. Make a natural-size drawing of the stock and the scion, at least three inches in length. Show the cut surfaces separately, and give the size of each.

2. Make a natural-size drawing, four inches in length, after the stock and the scion have been placed together and firmly tied.

THE STUDY OF VENEER GRAFTING

STUDENT'S NOTES AND REPORT

EXERCISE XXXIX

THE STUDY OF BORDEAUX MIXTURE

Material. Copper sulphate, stone lime, hydrated lime.

Apparatus. Quart fruit jar, granite kettle, glass stirring-rod.

Fungicides are chemicals that are used for the prevention or the killing of plant diseases. In some cases, where the fungus, or parasitic growth, is on the outside of the host, the external application of the fungicide will kill the disease. In other cases, where the disease is on the inside of the plant tissue, the host is covered with a fungicide, so that the spores, or reproductive bodies of the disease, are killed before they have a chance to enter the host plant.

There are several important fungicides. Bordeaux mixture is probably the best-known one, and it is made in different strengths for different purposes. It is a mixture of copper sulphate and lime, and the different strengths depend upon the amount of the chemicals used. The regular standard formulas for commercial spraying are as follows:

NO. 1, STANDARD BORDEAUX
MIXTURE

4 lb. copper sulphate
4 lb. stone lime
50 gal. water

NO. 2, HALF-STRENGTH BORDEAUX
MIXTURE

2 lb. copper sulphate
2 lb. stone lime
50 gal. water

Laboratory formula for standard Bordeaux mixture. Dissolve one ounce of copper sulphate in two pints of water. Place one ounce of stone lime in a vessel, and slake in two pints of water and stir while slaking. Describe fully what takes place. Use these as stock solutions.

1. Now take one fourth of the copper sulphate solution and dilute it with the same quantity of cold water. Also take one fourth of the lime solution and dilute it with the same quantity of cold water. Pour the two diluted solutions together, fill a quart jar, and label. Describe what takes place and give the color of the solution.

THE STUDY OF BORDEAUX MIXTURE

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

2. Now repeat the same operation as in 1, but use hot water to dilute each of the stock solutions. Place in a second glass jar, and label. Describe, and give your results.

3. Take one ounce of each of the stock solutions and pour into a glass jar. Dilute the mixture to eight ounces with cold water, and label.

4. Shake all three samples of Bordeaux and allow them to settle. After half an hour examine each sample carefully, note the differences, and record all data. Now permit the solutions to remain for three or four days and again agitate them and describe fully what takes place.

5. Make a second set of solutions as in Laboratory Formula 1, but instead of using stone lime use one and one-third ounces of hydrated lime. What is the difference between the hydrated lime and the stone lime? Describe. What result do you get from pouring water on the hydrated lime? Does this differ any from Laboratory Formula 1? If so, how and why? Describe the Bordeaux made from the hydrated lime. Does this differ any from that made from the stone lime? Describe.

Describe Bordeaux mixture. What, in your opinion, is the best sample? Why? After Bordeaux has stood for some time is it as good as when it is first made? Discuss. Is it advisable to keep a stock solution of Bordeaux? Why? Tell which is the most important chemical in the mixture and give the reason for your answer. Why is it necessary to make Bordeaux in either glass or wooden vessels? Tell the differences between the combined solutions and the separate ingredients. Why is it important that the diluted solutions be mixed rather than the concentrated? Discuss.

Laboratory formula for half-strength Bordeaux mixture. Now take one pint of the standard Bordeaux mixture and add one pint of water. What does this dilution of the mixture make? Describe the difference between the half-strength Bordeaux and the standard Bordeaux. When and on what plants is half-strength Bordeaux mixture used?

Now take one pint of the standard Bordeaux made from hydrated lime, and add one pint of water. Does this differ from that made from the stone lime? If so, describe.

Problem 1. If it takes seven and one-half gallons of standard Bordeaux mixture to spray a fair-sized apple tree once, how

THE STUDY OF BORDEAUX MIXTURE

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

much will be required to spray fifty trees three times? If copper sulphate costs five and one-half cents a pound and lime a cent a pound, what will be the cost of the material?

Problem 2. I have a ten-acre apple orchard. There are forty-nine trees to the acre. I wish to spray this orchard four times. How much Bordeaux mixture will be required, and what will be the cost of the chemicals?

REFERENCES:

Bulletin No. 135, Illinois Agricultural Experiment Station.

Farmers' Bulletin No. 243, United States Department of Agriculture,
pp. 5-14.

Lodeman, "The Spraying of Plants," pp. 19-45.

THE STUDY OF BORDEAUX MIXTURE

STUDENT'S NOTES AND REPORT

EXERCISE XL

THE STUDY OF LIME-SULPHUR

Material. Stone lime, flowers of sulphur, commercial lime-sulphur.

Apparatus. One-gallon granite kettle, hydrometer, large glass cylinders, cooking burner.

It has been learned that lime-sulphur can be used as a fungicide and in some cases is superior to Bordeaux mixture for the spraying of certain plants and the controlling of certain diseases. The point of difference between lime-sulphur as a fungicide and as an insecticide is the degree of dilution, or the strength of the solution. The strength can best be determined by the use of a hydrometer (Fig. 37), which tests the density of the solution.

There are several formulas from which the lime-sulphur is made. A good commercial formula to follow is

Stone lime	40 lb.
Flowers of sulphur	80 lb.
Water	50 gal.

FIG. 37. Cylinder, showing hydrometer for testing the density of lime-sulphur solution

Laboratory formula. Weigh out four ounces of stone lime and eight ounces of flowers of sulphur. Place the lime in the granite vessel and slake with water, adding the water gradually and being careful not to smother the lime with too much water. Stir until the lime is thoroughly slaked and forms a uniform paste.

Now add enough water to make a half gallon, after which place over the fire and boil. Now mix the flowers of sulphur to a thin paste, and add gradually to the lime when boiling, and stir constantly. Allow the mixture to boil for about an hour, stirring at frequent intervals. Keep the water in the

THE STUDY OF LIME-SULPHUR

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

vessel at a constant level by adding hot water from time to time. This level can be determined by measuring the depth of the liquid with a stick that has previously been notched.

As soon as this solution is cool enough, allow it to settle, pour off the liquid into a cylinder, and test with the hydrometer. Note the reading, and tabulate. Now dilute a portion of the solution with water according to the following scale, until it is the proper strength for a summer spray, and dilute another portion until it is the proper strength for a dormant spray.

DEGREE, BAUMÉ	SPECIFIC GRAVITY	DILUTION FOR DORMANT SPRAY	DILUTION FOR SUMMER SPRAY
35	1.318	1 to 9	1 to 45
34	1.306	1 to $8\frac{1}{2}$	1 to 43
33	1.294	1 to 8	1 to 41
32	1.283	1 to $7\frac{1}{2}$	1 to 40
31	1.271	1 to $7\frac{1}{2}$	1 to 38
30	1.260	1 to 7	1 to 36
29	1.250	1 to $6\frac{1}{2}$	1 to 34
28	1.239	1 to $6\frac{1}{2}$	1 to 32
27	1.228	1 to 6	1 to 31
26	1.218	1 to $5\frac{1}{2}$	1 to 29
25	1.208	1 to $5\frac{1}{2}$	1 to 28
24	1.198	1 to $5\frac{1}{4}$	1 to 26
23	1.188	1 to $4\frac{3}{4}$	1 to 24
22	1.178	1 to $4\frac{1}{2}$	1 to 23
21	1.169	1 to $4\frac{1}{4}$	1 to 21
20	1.160	1 to 4	1 to 20
19	1.150	1 to $3\frac{1}{2}$	1 to 18
18	1.141	1 to $3\frac{1}{4}$	1 to 17
17	1.132	1 to 3	1 to 16
16	1.124	1 to $2\frac{3}{4}$	1 to 15
15	1.115	1 to $2\frac{1}{2}$	1 to 14

The above table is for diluting the concentrated lime-and-sulphur solution for the dormant as well as for the summer sprays. The dilution for the summer sprays is usually from thirty-five to forty-five gallons of water when the Baumé test is 32° or 33°. Now figure out from this table the degree of the dilution of your sample made in the laboratory.

Example 1. The concentrated lime-sulphur solution, test 35° Baumé, by the table has a specific gravity of 1.318. If it is desired to spray trees with lime-sulphur at 20° Baumé, which has a specific gravity of 1.160, calculate the number of dilutions required as follows: the decimal of concentration of the

THE STUDY OF LIME-SULPHUR

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

Baumé 35° is .318; this, divided by the decimal for Baumé 20°, which is .160, will give 19.8, or the number of dilutions required, this result being obtained by adding 19.8 volumes of water to 1 volume of the concentrated lime and sulphur. This rule is based upon the general fact that the density of a solution heavier than water varies inversely with the number of dilutions.

When the density of the lime-sulphur solution or the home-made wash has been first determined by the use of a hydrometer, sprays of any desired density may be calculated by using the table. The rule for determining the number of dilutions is as follows: divide the decimal of the concentrate by the decimal of the spray desired; the quotient will be the number of dilutions.

$$\frac{\text{Decimal of the concentrate}}{\text{Decimal of the spray desired}} = \text{the number of dilutions}$$

$$\frac{.318}{.160} = 19.8$$

Example 2. Assuming that the decimal of the concentrate is known and that this concentrate is diluted by a certain number of volumes of water, what is the decimal of the spray? Taking the same example as in Ex. 1, we have the decimal of the concentrate, .318, and the number of dilutions, 19.8. Dividing .318 by 19.8 equals .160, which is the decimal of the spray and corresponds to 20° Baumé. Now figure out the degree of dilution for each sample you make in the laboratory.

$$\frac{\text{Decimal of the concentrate}}{\text{Number of dilutions}} = \text{decimal of the spray}$$

$$\frac{.318}{19.8} = .160$$

Secure some concentrated lime-sulphur solution. How does this differ in color and looks from the home-made lime-sulphur solution? Test each of the solutions with the hydrometer. Note the reading in each case. What amount of water was necessary to bring both of the solutions to the same reading? How do you account for the difference, if any? Discuss fully.

REFERENCES:

- Bulletin No. 288*, Cornell Agricultural Experiment Station.
- Bulletin No. 289*, Cornell Agricultural Experiment Station.
- Circular No. 10*, Michigan Agricultural Experiment Station.
- Bulletin No. 330*, New York Agricultural Experiment Station.

THE STUDY OF LIME-SULPHUR
STUDENT'S NOTES AND REPORT

EXERCISE XLI

THE STUDY OF SELF-BOILED LIME-SULPHUR

Material. Stone lime, flowers of sulphur.

Apparatus. Granite kettle, stirring-rod.

Self-boiled lime-sulphur differs from the lime-sulphur wash in Exercise XL chiefly in that there is no artificial heat applied and all of the cooking that the solution undergoes is done by the heat generated by the stone lime in slaking. This is used only as a summer spray, and is proving very satisfactory for use on the peach against the scab and brown rot. The formula that is most often used is

Flowers of sulphur	8 lb.
Stone lime	8 lb.
Water	50 gal.

Laboratory formula. Weigh out two and one-half ounces of the flowers of sulphur and two and one-half ounces of stone lime. Place the lime and the sulphur in a granite vessel and add enough cold water to slake the lime, stirring constantly to prevent burning. Cover the granite kettle, so as to retain as much of the heat as possible. Watch the mixture closely, and as soon as an orange-colored liquid begins to gather on the surface add enough more water to make a gallon. Do not use hot water or allow the mixture to stand after the lime is slaked or before dilution. In this spray we do not want soluble sulphides to form, because these will injure the foliage as well as the fruit. How does this solution differ from the one in Exercise XL? Describe the color of each solution and note any differences. Why should cold water be used instead of hot water? Why is the prevention of the formation of soluble sulphides especially important? Discuss.

REFERENCE:

Special Bulletin No. 61, Michigan Agricultural Experiment Station.

THE STUDY OF SELF-BOILED LIME-SULPHUR

STUDENT'S NOTES AND REPORT

EXERCISE XLII

THE STUDY OF AMMONIACAL SOLUTION OF COPPER CARBONATE

Material. Copper sulphate, carbonate of soda, ammonia.

Apparatus. Quart glass jar, stirring-rod, granite kettle.

To make this fungicide it is necessary first to make the copper carbonate. The common formula for copper carbonate is

Copper sulphate	10 lb.
Water	10 gal.
Carbonate of soda	12 lb.
Water	10 gal.

Laboratory formula. Weigh out two ounces of copper sulphate and dissolve in one pint of water, and in a separate vessel dissolve two ounces of carbonate of soda in one pint of water. When cool, mix the two solutions slowly, stirring well. Allow the solution to stand twelve hours to settle, after which pour off the liquid. Add one quart of water and stir a second time, again allowing the solution to stand for twelve hours. Repeat this operation several times and then filter off the residue. The blue powder thus obtained is copper carbonate. Describe fully the operation and explain each step. What is the difference between the finished product and the copper sulphate? Write out the chemical formula and give the resulting product.

The common formula for ammoniacal solution of copper carbonate is

Copper carbonate	6 oz.
Ammonia	3 pt.
Water	50 gal.

Laboratory formula. Weigh out one fourth of an ounce of the copper carbonate just prepared and dissolve this in one eighth of a pint of ammonia. When thoroughly dissolved, add enough water to make two gallons. What is the function of the ammonia? Where is this fungicide used? How does it differ in this respect from Bordeaux mixture? Give the chemical formula of copper carbonate when mixed with the ammonia.

AMMONIACAL SOLUTION OF COPPER CARBONATE

STUDENT'S NOTES AND REPORT

EXERCISE XLIII

THE STUDY OF PARIS GREEN AND ARSENATE OF LEAD

Material. Paris green, powdered arsenate of lead, paste arsenate of lead.

Apparatus. Quart glass jars, stirring-rod.

Insecticides differ from fungicides in that they are chemicals used for the destruction of insects that prey upon cultivated plants. There are two kinds of insecticides: namely, stomach poisons, or insecticides that are eaten by the insects and kill by being taken into the stomach; and contact insecticides, or those which are sprayed on the insects and kill by coming in contact with their bodies.

PARIS GREEN

Paris green, one of our important insecticides, is an aceto-arsenite of copper, and Ehrmann has given the composition of pure Paris green as follows:

Copper oxide	31.29%
Arsenious acid	58.65%
Acetic acid	10.06%

However, most samples show some variation from the above. The free arsenic in the Paris green is dangerous to foliage, and when Paris green is used alone, lime should be added to avoid burning the leaves.

The usual formula for spraying is

Paris green	1 lb.
Water	100 to 300 gal.

Laboratory formula. Weigh out one fourth of an ounce of Paris green and place it in a quart fruit jar of water. Study carefully the action of the poison when placed in the water. Describe what takes place. Now with a stirring-rod vigorously agitate the water. Does it mix with the water? If not, why? Describe. Is Paris green in water a solution or a mixture?

PARIS GREEN AND ARSENATE OF LEAD

STUDENT'S NOTES AND REPORT

LABORATORY MANUAL OF HORTICULTURE

ARSENATE OF LEAD

Arsenate of lead is another of our important insecticides and to-day probably ranks first. It can be purchased in two forms — as a powder or as a paste. The paste form is the most widely used. The usual formula for arsenate of lead is from two to three pounds to fifty gallons of water, but the proportion varies with the crop to be sprayed.

Laboratory formula. Weigh out one-half ounce each of the powdered and the paste form of arsenate of lead, and dissolve each in a separate quart glass fruit jar of water. Study carefully what takes place. Now place all three of the jars containing the insecticides where the light will pass through them. Stir each vigorously for a minute and note the results. Watch the solutions for fifteen minutes, noting what takes place in each case. Are the insecticides completely dissolved, forming a chemical union, or are they held in suspension, with only a slight chemical action taking place? Discuss. Note accurately the length of time required for each solution to settle. Discuss in detail why it is absolutely necessary to have the spray mixture thoroughly agitated during the whole time of spraying.

PARIS GREEN AND ARSENATE OF LEAD

STUDENT'S NOTES AND REPORT

EXERCISE XLIV

THE STUDY OF KEROSENE EMULSION

Material. Laundry soap, kerosene.

Apparatus. Granite kettle, cooking-burner, stirring-rod.

This insecticide is used for many scale insects, and also for some other of the sucking insects, such as the plant lice, etc. The emulsion is usually made in the form of a stock solution and diluted to the required strength when needed. The regular formula is

Laundry soap	$\frac{1}{2}$ lb.
Kerosene	2 gal.
Soft water	1 gal.

Laboratory formula. Weigh out one ounce of laundry soap, cut into small pieces, and dissolve it in one pint of water. Allow this solution to boil, *remove it from the fire*, and add one quart of kerosene. Stir the mixture violently until it becomes a creamy mass that will not separate. This usually requires from five to ten minutes. The emulsion is now finished. Describe kerosene emulsion. Name several kinds of insects upon which it is used. What is an emulsion?

For use, dilute one part of the emulsion with eight or ten parts of water for hard-bodied insects and with fifteen or twenty parts of water for soft-bodied insects. Why is it detrimental to use the kerosene alone, and what is the destructive principle in the emulsion?

THE STUDY OF KEROSENE EMULSION

STUDENT'S NOTES AND REPORT

EXERCISE XLV

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small	CALYX LOBES	Submerged	Medial
FORM	Acuminate	Gray	CARPELS
Conical	Acute	Red	Cordate
Irregular	Obtuse	Russet	Emarginate
Oblate	Long	White	Ovate
Oblong	Medium	CORE	Round
Oblique	Short	Abaxile	Tufted
Ovate	Narrow	Axile	SEED
Ribbed	Wide	Large	Acuminate
Roundish	Separated at base	Medium	Acute
Symmetrical	BASIN	Small	Flat
STEM	Abrupt	Closed	Large
Long	Corrugated	Open	Long
Medium	Furrowed	Distant	Medium
Short	Obtuse	Median	Narrow
Slender	Symmetrical	Sessile	Obtuse
Thick	Deep	CORE LINE	Plump
CAVITY	Medium	Clasping	Small
Acuminate	Shallow	Meeting	Tufted
Acute	Narrow	CALYX TUBE	Wide
Lipped	Wide	Conical	FLAVOR
Obtuse	SKIN	Funnel-shaped	Aromatic
Russeted	Dull	Urn-shaped	Flat
Symmetrical	Glossy	Long	Sour
Narrow	Rough	Short	Sprightly
Wide	Russet	Narrow	Sub-acid
Deep	Smooth	Wide	QUALITY
Medium	Tender	FLESH	Best
Shallow	Thin	Color	Good
COLOR	Tough	TEXTURE	Very good
Ground	Waxen	Coarse	Medium
Overlying	DOTS	Crisp	Poor
Blushed	Areolar	Dry	Very poor
Mottled	Elongated	Fine	MARKET
Striped	Stellate	Firm	Distant
Polish	Few	Juicy	Local
Prevailing effect..	Many	Medium	USE
CALYX	Large	Sweet	Cooking
Closed	Medium	Tender	Dessert
Open	Small	Tough	Market

A STUDY OF THE APPLE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE XLVI

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small	CALYX LOBES	Submerged	Medial
FORM	Acuminate	Gray	CARPELS
Conical	Acute	Red	Cordate
Irregular	Obtuse	Russet	Emarginate
Oblate	Long	White	Ovate
Oblong	Medium	CORE	Round
Oblique	Short	Abaxile	Tufted
Ovate	Narrow	Axile	SEED
Ribbed	Wide	Large	Acuminate
Roundish	Separated at base	Medium	Acute
Symmetrical	BASIN	Small	Flat
STEM	Abrupt	Closed	Large
Long	Corrugated	Open	Long
Medium	Furrowed	Distant	Medium
Short	Obtuse	Median	Narrow
Slender	Symmetrical	Sessile	Obtuse
Thick	Deep	CORE LINE	Plump
CAVITY	Medium	Clasping	Small
Acuminate	Shallow	Meeting	Tufted
Acute	Narrow	CALYX TUBE	Wide
Lipped	Wide	Conical	FLAVOR
Obtuse	SKIN	Funnel-shaped	Aromatic
Russeted	Dull	Urn-shaped	Flat
Symmetrical	Glossy	Long	Sour
Narrow	Rough	Short	Sprightly
Wide	Russet	Narrow	Sub-acid
Deep	Smooth	Wide	QUALITY
Medium	Tender	FLESH	Best
Shallow	Thin	Color.....	Good
COLOR	Tough	TEXTURE	Very good
Ground.....	Waxed	Coarse	Medium
Overlying	Dors	Crisp	Poor
Blushed	Areolar	Dry	Very poor
Mottled	Elongated	Fine	MARKET
Striped	Stellate	Firm	Distant
Polish.....	Few	Juicy	Local
Prevailing effect..	Many	Medium	USE
CALYX	Large	Sweet	Cooking
Closed	Medium	Tender	Dessert
Open	Small	Tough	Market

A STUDY OF THE APPLE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE XLVII

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small		Submerged	Medial
FORM	CALYX LOBES	Gray	CARPELS
Conical	Acuminate	Red	Cordate
Irregular	Acute	Russet	Emarginate
Oblate	Obtuse	White	Ovate
Oblong	Long	CORE	Round
Oblique	Medium	Abaxile	Tufted
Ovate	Short	Axile	SEED
Ribbed	Narrow	Large	Acuminate
Roundish	Wide	Medium	Acute
Symmetrical	Separated at base	Small	Flat
STEM	BASIN	Closed	Large
Long	Abrupt	Open	Long
Medium	Corrugated	Distant	Medium
Short	Furrowed	Median	Narrow
Slender	Obtuse	Sessile	Obtuse
Thick	Symmetrical	CORE LINE	Plump
CAVITY	Deep	Clasping	Small
Acuminate	Medium	Meeting	Tufted
Acute	Shallow		Wide
Lipped	Narrow	CALYX TUBE	FLAVOR
Obtuse	Wide	Conical	Aromatic
Russeted	SKIN	Funnel-shaped	Flat
Symmetrical	Dull	Urn-shaped	Sour
Narrow	Glossy	Long	Sprightly
Wide	Rough	Short	Sub-acid
Deep	Russet	Narrow	QUALITY
Medium	Smooth	Wide	Best
Shallow	Tender	FLESH	Good
COLOR	Thin	Color	Very good
Ground	Tough	TEXTURE	Medium
Overlying	Waxen	Coarse	Poor
Blushed	DOTS	Crisp	Very poor
Mottled	Areolar	Dry	MARKET
Striped	Elongated	Fine	Distant
Polish	Stellate	Firm	Local
Prevailing effect..	Few	Juicy	USE
CALYX	Many	Medium	Cooking
Closed	Large	Sweet	Dessert
Open	Medium	Tender	Market
	Small	Tough	

A STUDY OF THE APPLE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE XLVIII

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY_____DATE_____

GROUP_____

SYNONYMS_____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small	CALYX LOBES	Submerged	Medial
FORM	Acuminate	Gray	CARPELS
Conical	Acute	Red	Cordate
Irregular	Obtuse	Russet	Emarginate
Oblate	Long	White	Ovate
Oblong	Medium	CORE	Round
Oblique	Short	Abaxile	Tufted
Ovate	Narrow	Axile	SEED
Ribbed	Wide	Large	Acuminate
Roundish	Separated at base	Medium	Acute
Symmetrical	BASIN	Small	Flat
STEM	Abrupt	Closed	Large
Long	Corrugated	Open	Long
Medium	Furrowed	Distant	Medium
Short	Obtuse	Median	Narrow
Slender	Symmetrical	Sessile	Obtuse
Thick	Deep	CORE LINE	Plump
CAVITY	Medium	Clasping	Small
Acuminate	Shallow	Meeting	Tufted
Acute	Narrow	CALYX TUBE	Wide
Lipped	Wide	Conical	FLAVOR
Obtuse	SKIN	Funnel-shaped	Aromatic
Russeted	Dull	Urn-shaped	Flat
Symmetrical	Glossy	Long	Sour
Narrow	Rough	Short	Sprightly
Wide	Russet	Narrow	Sub-acid
Deep	Smooth	Wide	QUALITY
Medium	Tender	FLESH	Best
Shallow	Thin	Color.....	Good
COLOR	Tough	TEXTURE	Very good
Ground.....	Waxen	Coarse	Medium
Overlying	DOTS	Crisp	Poor
Blushed	Areolar	Dry	Very poor
Mottled	Elongated	Fine	MARKET
Striped	Stellate	Firm	Distant
Polish.....	Few	Juicy	Local
Prevailing effect..	Many	Medium	USE
CALYX	Large	Sweet	Cooking
Closed	Medium	Tender	Dessert
Open	Small	Tough	Market

A STUDY OF THE APPLE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE XLIX

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small		Submerged	Medial
FORM	CALYX LOBES	Gray	CARPELS
Conical	Acuminate	Red	Cordate
Irregular	Acute	Russet	Emarginate
Oblate	Obtuse	White	Ovate
Oblong	Long	CORE	Round
Oblique	Medium	Abaxile	Tufted
Ovate	Short	Axile	SEED
Ribbed	Narrow	Large	Acuminate
Roundish	Wide	Medium	Acute
Symmetrical	Separated at base	Small	Flat
STEM	BASIN	Closed	Large
Long	Abrupt	Open	Long
Medium	Corrugated	Distant	Medium
Short	Furrowed	Median	Narrow
Slender	Obtuse	Sessile	Obtuse
Thick	Symmetrical	CORE LINE	Plump
CAVITY	Deep	Clasping	Small
Acuminate	Medium	Meeting	Tufted
Acute	Shallow	CALYX TUBE	Wide
Lipped	Narrow	Conical	FLAVOR
Obtuse	Wide	Funnel-shaped	Aromatic
Russeted	SKIN	Urn-shaped	Flat
Symmetrical	Dull	Long	Sour
Narrow	Glossy	Short	Sprightly
Wide	Rough	Narrow	Sub-acid
Deep	Russet	Wide	QUALITY
Medium	Smooth	FLESH	Best
Shallow	Tender	Color	Good
COLOR	Thin	TEXTURE	Very good
Ground	Tough	Coarse	Medium
Overlying	Waxen	Crisp	Poor
Blushed	DOTS	Dry	Very poor
Mottled	Areolar	Fine	MARKET
Striped	Elongated	Firm	Distant
Polish	Stellate	Juicy	Local
Prevailing effect..	Few	Medium	USE
CALYX	Many	Sweet	Cooking
Closed	Large	Tender	Dessert
Open	Medium	Tough	Market
	Small		

A STUDY OF THE APPLE

SEASON _____

LOCALITY _____

PRODUCTIVENESS _____

GENERAL APPEARANCE _____

DESIRABILITY _____

HISTORY _____

REMARKS _____

Draw a longitudinal section of the fruit.

EXERCISE L

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small	CALYX LOBES	Submerged	Medial
FORM	Acuminate	Gray	CARPELS
Conical	Acute	Red	Cordate
Irregular	Obtuse	Russet	Emarginate
Oblate	Long	White	Ovate
Oblong	Medium	CORE	Round
Oblique	Short	Abaxile	Tufted
Ovate	Narrow	Axile	SEED
Ribbed	Wide	Large	Acuminate
Roundish	Separated at base	Medium	Acute
Symmetrical	BASIN	Small	Flat
STEM	Abrupt	Closed	Large
Long	Corrugated	Open	Long
Medium	Furrowed	Distant	Medium
Short	Obtuse	Median	Narrow
Slender	Symmetrical	Sessile	Obtuse
Thick	Deep	CORE LINE	Plump
CAVITY	Medium	Clasping	Small
Acuminate	Shallow	Meeting	Tufted
Acute	Narrow	CALYX TUBE	Wide
Lipped	Wide	Conical	FLAVOR
Obtuse	SKIN	Funnel-shaped	Aromatic
Russeted	Dull	Urn-shaped	Flat
Symmetrical	Glossy	Long	Sour
Narrow	Rough	Short	Sprightly
Wide	Russet	Narrow	Sub-acid
Deep	Smooth	Wide	QUALITY
Medium	Tender	FLESH	Best
Shallow	Thin	Color	Good
COLOR	Tough	TEXTURE	Very good
Ground	Waxen	Coarse	Medium
Overlying	DOTS	Crisp	Poor
Blushed	Areolar	Dry	Very poor
Mottled	Elongated	Fine	MARKET
Striped	Stellate	Firm	Distant
Polish	Few	Juicy	Local
Prevailing effect..	Many	Medium	USE
CALYX	Large	Sweet	Cooking
Closed	Medium	Tender	Dessert
Open	Small	Tough	Market

A STUDY OF THE APPLE

SEASON _____

LOCALITY _____

PRODUCTIVENESS _____

GENERAL APPEARANCE _____

DESIRABILITY _____

HISTORY _____

REMARKS _____

Draw a longitudinal section of the fruit.

EXERCISE LI

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small		Submerged	Medial
FORM	CALYX LOBES	Gray	CARPELS
Conical	Acuminate	Red	Cordate
Irregular	Acute	Russet	Emarginate
Oblate	Obtuse	White	Ovate
Oblong	Long	CORE	Round
Oblique	Medium	Abaxile	Tufted
Ovate	Short	Axile	SEED
Ribbed	Narrow	Large	Acuminate
Roundish	Wide	Medium	Acute
Symmetrical	Separated at base	Small	Flat
STEM	BASIN	Closed	Large
Long	Abrupt	Open	Long
Medium	Corrugated	Distant	Medium
Short	Furrowed	Median	Narrow
Slender	Obtuse	Sessile	Obtuse
Thick	Symmetrical	CORE LINE	Plump
CAVITY	Deep	Clasping	Small
Acuminate	Medium	Meeting	Tufted
Acute	Shallow	CALYX TUBE	Wide
Lipped	Narrow	Conical	FLAVOR
Obtuse	Wide	Funnel-shaped	Aromatic
Russeted	SKIN	Urn-shaped	Flat
Symmetrical	Dull	Long	Sour
Narrow	Glossy	Short	Sprightly
Wide	Rough	Narrow	Sub-acid
Deep	Russet	Wide	QUALITY
Medium	Smooth	FLESH	Best
Shallow	Tender	Color	Good
COLOR	Thin	TEXTURE	Very good
Ground	Tough	Coarse	Medium
Overlying	Waxen	Crisp	Poor
Blushed	DORS	Dry	Very poor
Mottled	Areolar	Fine	MARKET
Striped	Elongated	Firm	Distant
Polish	Stellate	Juicy	Local
Prevailing effect..	Few	Medium	USE
CALYX	Many	Sweet	Cooking
Closed	Large	Tender	Dessert
Open	Medium	Tough	Market
	Small		

A STUDY OF THE APPLE

SEASON _____

LOCALITY _____

PRODUCTIVENESS _____

GENERAL APPEARANCE _____

DESIRABILITY _____

HISTORY _____

REMARKS _____

Draw a longitudinal section of the fruit.

EXERCISE LII

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small	CALYX LOBES	Submerged	Medial
FORM	Acuminate	Gray	CARPELS
Conical	Acute	Red	Cordate
Irregular	Obtuse	Russet	Emarginate
Oblate	Long	White	Ovate
Oblong	Medium	CORE	Round
Oblique	Short	Abaxile	Tufted
Ovate	Narrow	Axile	SEED
Ribbed	Wide	Large	Acuminate
Roundish	Separated at base	Medium	Acute
Symmetrical	BASIN	Small	Flat
STEM	Abrupt	Closed	Large
Long	Corrugated	Open	Long
Medium	Furrowed	Distant	Medium
Short	Obtuse	Median	Narrow
Slender	Symmetrical	Sessile	Obtuse
Thick	Deep	CORE LINE	Plump
CAVITY	Medium	Clasping	Small
Acuminate	Shallow	Meeting	Tufted
Acute	Narrow	CALYX TUBE	Wide
Lipped	Wide	Conical	FLAVOR
Obtuse	SKIN	Funnel-shaped	Aromatic
Russeted	Dull	Urn-shaped	Flat
Symmetrical	Glossy	Long	Sour
Narrow	Rough	Short	Sprightly
Wide	Russet	Narrow	Sub-acid
Deep	Smooth	Wide	QUALITY
Medium	Tender	FLESH	Best
Shallow	Thin	Color	Good
COLOR	Tough	TEXTURE	Very good
Ground	Waxen	Coarse	Medium
Overlying	DOTS	Crisp	Poor
Blushed	Areolar	Dry	Very poor
Mottled	Elongated	Fine	MARKET
Striped	Stellate	Firm	Distant
Polish	Few	Juicy	Local
Prevailing effect..	Many	Medium	USE
CALYX	Large	Sweet	Cooking
Closed	Medium	Tender	Dessert
Open	Small	Tough	Market

A STUDY OF THE APPLE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE LIII

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small		*Submerged	Medial
FORM	CALYX LOBES	Gray	CARPELS
Conical	Acuminate	Red	Cordate
Irregular	Acute	Russet	Emarginate
Oblate	Obtuse	White	Ovate
Oblong	Long	CORE	Round
Oblique	Medium	Abaxile	Tufted
Ovate	Short	Axile	SEED
Ribbed	Narrow	Large	Acuminate
Roundish	Wide	Medium	Acute
Symmetrical	Separated at base	Small	Flat
STEM	BASIN	Closed	Large
Long	Abrupt	Open	Long
Medium	Corrugated	Distant	Medium
Short	Furrowed	Median	Narrow
Slender	Obtuse	Sessile	Obtuse
Thick	Symmetrical	CORE LINE	Plump
CAVITY	Deep	Clasping	Small
Acuminate	Medium	Meeting	Tufted
Acute	Shallow	CALYX TUBE	Wide
Lipped	Narrow	Conical	FLAVOR
Obtuse	Wide	Funnel-shaped	Aromatic
Russeted	SKIN	Urn-shaped	Flat
Symmetrical	Dull	Long	Sour
Narrow	Glossy	Short	Sprightly
Wide	Rough	Narrow	Sub-acid
Deep	Russet	Wide	QUALITY
Medium	Smooth	FLESH	Best
Shallow	Tender	Color	Good
COLOR	Thin	TEXTURE	Very good
Ground	Tough	Coarse	Medium
Overlying	Waxed	Crisp	Poor
Blushed	DOTS	Dry	Very poor
Mottled	Areolar	Fine	MARKET
Striped	Elongated	Firm	Distant
Polish	Stellate	Juicy	Local
Prevailing effect..	Few	Medium	USE
CALYX	Many	Sweet	Cooking
Closed	Large	Tender	Dessert
Open	Medium	Tough	Market
	Small		

A STUDY OF THE APPLE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE LIV

A STUDY OF THE APPLE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Large	Compressed	STAMENS
Large	Medium	Raised	Basal
Medium	Small	Rough	Marginal
Small		Submerged	Medial
FORM	CALYX LOBES	Gray	CARPELS
Conical	Acuminate	Red	Cordate
Irregular	Acute	Russet	Emarginate
Oblate	Obtuse	White	Ovate
Oblong	Medium	CORE	Round
Oblique	Short	Abaxile	Tufted
Ovate	Narrow	Axile	SEED
Ribbed	Wide	Large	Acuminate
Roundish	Separated at base	Medium	Acute
Symmetrical		Small	Flat
STEM	BASIN	Closed	Large
Long	Abrupt	Open	Long
Medium	Corrugated	Distant	Medium
Short	Furrowed	Median	Narrow
Slender	Obtuse	Sessile	Obtuse
Thick	Symmetrical	CORE LINE	Plump
CAVITY	Deep	Clasping	Small
Acuminate	Medium	Meeting	Tufted
Acute	Shallow	CALYX TUBE	Wide
Lipped	Narrow	Conical	FLAVOR
Obtuse	Wide	Funnel-shaped	Aromatic
Russeted	SKIN	Urn-shaped	Flat
Symmetrical	Dull	Long	Sour
Narrow	Glossy	Short	Sprightly
Wide	Rough	Narrow	Sub-acid
Deep	Russet	Wide	QUALITY
Medium	Smooth	FLESH	Best
Shallow	Tender	Color	Good
COLOR	Thin	TEXTURE	Very good
Ground	Tough	Coarse	Medium
Overlying	Waxy	Crisp	Poor
Blushed	DORS	Dry	Very poor
Mottled	Areolar	Fine	MARKET
Striped	Elongated	Firm	Distant
Polish	Stellate	Juicy	Local
Prevailing effect..	Few	Medium	USE
CALYX	Many	Sweet	Cooking
Closed	Large	Tender	Dessert
Open	Medium	Tough	Market
	Small		

A STUDY OF THE APPLE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE LV

A STUDY OF THE GRAPE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

BERRY	Wide	JUICINESS	PIGMENT
SIZE	Winged	Medium	FLAVOR
Large	Irregular	Dry	Astringent
Medium	Regular	Juicy	Foxy
Small	Long	Clear	Neutral
Uniform	Medium	Pink	SEED
Variable	Short	Purple	Number.....
FORM	Double-shouldered	Thick	SIZE
Oblate	Single-shouldered	Thin	Large
Oval			Medium
Roundish	NUMBER PER SHOOT	TEXTURE	Small
	COMPACTNESS	Coarse	SHAPE
COLOR	Compact	Melting	Notched
Amber	Loose	Slippery	Oblong
Black	Medium	Solid	Plump
Blue	PEDUNCLE	Stringy	Blunt
Green	Long	Tender	Pointed
Purple	Medium	Tough	Rounded
Red	Short	AROMA	Narrow
Violet	Slender	Aromatic	Wide
White	Thick	Foxy	Long
Yellow		Spicy	Medium
Glossy	PEDICEL	FLAVOR	Short
Translucent	Long	Acid	COLOR
BLOOM	Medium	Foxy	Brown
Abundance.....	Short	Mild	Yellow
Color.....	Slender	Spicy	RAPHE
APEX	Thick	Sweet	Cordlike
Adherence of	Swellings	Tart	Large
stigma	Warts	Vinous	Small
ADHESION	BRUSH	QUALITY	Obscure
Drops	Color.....	Best	Prominent
Persistent	Length.....	Good	CHALAZA
CLUSTER	FLESH	Very good	Above center
SIZE	COLOR	Medium	Below center
Large	Greenish	Poor	Circular
Medium	Red	Very poor	Distinct
Small	White	SEPARATION FROM	Obscure
FORM	Translucent	SEED	Oval
Cylindrical	Transparent	With difficulty	USE
Ovate		Easily	Dessert
Roundish	FIRMNESS	SKIN	Kitchen
Slender	Firm	Medium	Market
Tapering	Medium	Thick	Raisin
	Soft	Thin	Shipping
		Tender	Wine
		Tough	

A STUDY OF THE GRAPE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Make drawings of several different seeds, showing the position and the shape of the chalaza.

EXERCISE LVI

A STUDY OF THE GRAPE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

BERRY	Wide	JUICINESS	PIGMENT.....
SIZE	Winged	Medium	FLAVOR
Large	Irregular	Dry	Astringent
Medium	Regular	Juicy	Foxy
Small	Long	Clear	Neutral
Uniform	Medium	Pink	SEED
Variable	Short	Purple	Number.....
FORM	Double-shouldered	Thick	SIZE
Oblate	Single-shouldered	Thin	Large
Oval	NUMBER PER SHOOT	TEXTURE	Medium
Roundish	COMPACTNESS	Coarse	Small
COLOR	Compact	Melting	SHAPE
Amber	Loose	Slippery	Notched
Black	Medium	Solid	Oblong
Blue	PEDUNCLE	Stringy	Plump
Green	Long	Tender	Blunt
Purple	Medium	Tough	Pointed
Red	Short	AROMA	Rounded
Violet	Slender	Aromatic	Narrow
White	Thick	Foxy	Wide
Yellow	PEDICEL	Spicy	Long
Glossy	Long	FLAVOR	Medium
Translucent	Medium	Acid	Short
BLOOM	Short	Foxy	COLOR
Abundance.....	Slender	Mild	Brown
Color.....	Thick	Spicy	Yellow
APEX	Swellings	Tart	RAPHE
Adherence of stigma	Warts	Vinous	Cordlike
ADHESION	BRUSH	QUALITY	Large
Drops	Color.....	Best	Small
Persistent	Length.....	Good	Obscure
CLUSTER	FLESH	Very good	Prominent
SIZE	COLOR	Medium	CHALAZA
Large	Greenish	Poor	Above center
Medium	Red	Very poor	Below center
Small	White	SEPARATION FROM SEED	Circular
FORM	Translucent	With difficulty	Distinct
Cylindrical	Transparent	Easily	Obscure
Ovate	FIRMNESS	SKIN	Oval
Roundish	Firm	Medium	USE
Slender	Medium	Thick	Dessert
Tapering	Soft	Thin	Kitchen
		Tender	Market
		Tough	Raisin
			Shipping
			Wine

A STUDY OF THE GRAPE

SEASON _____

LOCALITY _____

PRODUCTIVENESS _____

GENERAL APPEARANCE _____

DESIRABILITY _____

HISTORY _____

REMARKS _____

Make drawings of several different seeds, showing the position and the shape of the chalaza.

EXERCISE LVII

A STUDY OF THE GRAPE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

BERRY	Wide	JUICINESS	PIGMENT
SIZE	Winged	Medium	FLAVOR
Large	Irregular	Dry	Astringent
Medium	Regular	Juicy	Foxy
Small	Long	Clear	Neutral
Uniform	Medium	Pink	SEED
Variable	Short	Purple	Number.....
FORM	Double-shouldered	Thick	SIZE
Oblate	Single-shouldered	Thin	Large
Oval	NUMBER PER SHOOT	TEXTURE	Medium
Roundish	COMPACTNESS	Coarse	Small
COLOR	Compact	Melting	SHAPE
Amber	Loose	Slippery	Notched
Black	Medium	Solid	Oblong
Blue	PEDUNCLE	Stringy	Plump
Green	Long	Tender	Blunt
Purple	Medium	Tough	Pointed
Red	Short	AROMA	Rounded
Violet	Slender	Aromatic	Narrow
White	Thick	Foxy	Wide
Yellow	PEDICEL	Spicy	Long
Glossy	Long	FLAVOR	Medium
Translucent	Medium	Acid	Short
BLOOM	Short	Foxy	COLOR
Abundance.....	Slender	Mild	Brown
Color.....	Thick	Spicy	Yellow
APEX	Swellings	Sweet	RAPHE
Adherence of stigma	Warts	Tart	Cordlike
ADHESION	BRUSH	Vinous	Large
Drops	Color.....	QUALITY	Small
Persistent	Length.....	Best	Obscure
CLUSTER	FLESH	Good	Prominent
SIZE	COLOR	Very good	CHALAZA
Large	Greenish	Medium	Above center
Medium	Red	Poor	Below center
Small	White	Very poor	Circular
FORM	Translucent	SEPARATION FROM SEED	Distinct
Cylindrical	Transparent	With difficulty	Obscure
Ovate	FIRMNESS	Easily	Oval
Roundish	Firm	SKIN	USE
Slender	Medium	Medium	Dessert
Tapering	Soft	Thick	Kitchen
		Thin	Market
		Tender	Raisin
		Tough	Shipping
			Wine

A STUDY OF THE GRAPE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Make drawings of several different seeds, showing the position and the shape of the chalaza.

EXERCISE LVIII

A STUDY OF THE GRAPE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY_____ DATE_____

GROUP_____

SYNONYMS_____

BERRY	Wide	JUICINESS	PIGMENT.....
SIZE	Winged	Medium	FLAVOR
Large	Irregular	Dry	Astringent
Medium	Regular	Juicy	Foxy
Small	Long	Clear	Neutral
Uniform	Medium	Pink	SEED
Variable	Short	Purple	Number.....
FORM	Double-shouldered	Thick	SIZE
Oblate	Single-shouldered	Thin	Large
Oval	NUMBER PER SHOOT	TEXTURE	Medium
Roundish	COMPACTNESS	Coarse	Small
COLOR	Compact	Melting	SHAPE
Amber	Loose	Slippery	Notched
Black	Medium	Solid	Oblong
Blue	PEDUNCLE	Stringy	Plump
Green	Long	Tender	Blunt
Purple	Medium	Tough	Pointed
Red	Short	AROMA	Rounded
Violet	Slender	Aromatic	Narrow
White	Thick	Foxy	Wide
Yellow	PEDICEL	Spicy	Long
Glossy	Long	FLAVOR	Medium
Translucent	Medium	Acid	Short
BLOOM	Short	Foxy	COLOR
Abundance.....	Slender	Mild	Brown
Color.....	Thick	Spicy	Yellow
APEX	Swellings	Sweet	RAPHE
Adherence of	Warts	Tart	Cordlike
stigma	BRUSH	Vinous	Large
ADHESION	Color.....	QUALITY	Small
Drops	Length.....	Best	Obscure
Persistent	FLESH	Good	Prominent
CLUSTER	COLOR	Very good	CHALAZA
SIZE	Greenish	Medium	Above center
Large	Red	Poor	Below center
Medium	White	Very poor	Circular
Small	Translucent	SEPARATION FROM	Distinct
FORM	Transparent	SEED	Obscure
Cylindrical	FIRMNESS	With difficulty	Oval
Ovate	Firm	Easily	USE
Roundish	Medium	SKIN	Dessert
Slender	Soft	Medium	Kitchen
Tapering		Thick	Market
		Thin	Raisin
		Tender	Shipping
		Tough	Wine

A STUDY OF THE GRAPE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Make drawings of several different seeds, showing the position and the shape of the chalaza.

EXERCISE LIX

A STUDY OF THE GRAPE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

BERRY	Wide	JUICINESS	PIGMENT
Size	Winged	Medium	FLAVOR
Large	Irregular	Dry	Astringent
Medium	Regular	Juicy	Foxy
Small	Long	Clear	Neutral
Uniform	Medium	Pink	SEED
Variable	Short	Purple	Number.....
FORM	Double-shouldered	Thick	SIZE
Oblate	Single-shouldered	Thin	Large
Oval	NUMBER PER SHOOT	TEXTURE	Medium
Roundish	COMPACTNESS	Coarse	Small
COLOR	Compact	Melting	SHAPE
Amber	Loose	Slippery	Notched
Black	Medium	Solid	Oblong
Blue	PEDUNCLE	Stringy	Plump
Green	Long	Tender	Blunt
Purple	Medium	Tough	Pointed
Red	Short	AROMA	Rounded
Violet	Slender	Aromatic	Narrow
White	Thick	Foxy	Wide
Yellow	PEDICEL	Spicy	Long
Glossy	Long	FLAVOR	Medium
Translucent	Medium	Acid	Short
BLOOM	Short	Foxy	COLOR
Abundance.....	Slender	Mild	Brown
Color.....	Thick	Spicy	Yellow
APEX	Swellings	Sweet	RAPHE
Adherence of stigma	Warts	Tart	Cordlike
ADHESION	BRUSH	Vinous	Large
Drops	Color.....	QUALITY	Small
Persistent	Length.....	Best	Obscure
CLUSTER	FLESH	Good	Prominent
Size	Color	Very good	CHALAZA
Large	Greenish	Medium	Above center
Medium	Red	Poor	Below center
Small	White	Very poor	Circular
FORM	Translucent	SEPARATION FROM SEED	Distinct
Cylindrical	Transparent	With difficulty	Obscure
Ovate	FIRMNESS	Easily	Oval
Roundish	Firm	SKIN	USE
Slender	Medium	Medium	Dessert
Tapering	Soft	Thick	Kitchen
		Thin	Market
		Tender	Raisin
		Tough	Shipping
			Wine

A STUDY OF THE GRAPE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Make drawings of several different seeds, showing the position and the shape of the chalaza.

EXERCISE LX

A STUDY OF THE GRAPE

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

BERRY	Wide	JUICINESS	PIGMENT.....
SIZE	Winged	Medium	FLAVOR
Large	Irregular	Dry	Astringent
Medium	Regular	Juicy	Foxy
Small	Long	Clear	Neutral
Uniform	Medium	Pink	SEED
Variable	Short	Purple	Number.....
FORM	Double-shouldered	Thick	SIZE
Oblate	Single-shouldered	Thin	Large
Oval	NUMBER PER SHOOT	TEXTURE	Medium
Roundish	COMPACTNESS	Coarse	Small
COLOR	Compact	Melting	SHAPE
Amber	Loose	Slippery	Notched
Black	Medium	Solid	Oblong
Blue	PEDUNCLE	Stringy	Plump
Green	Long	Tender	Blunt
Purple	Medium	Tough	Pointed
Red	Short	AROMA	Rounded
Violet	Slender	Aromatic	Narrow
White	Thick	Foxy	Wide
Yellow	PEDICEL	Spicy	Long
Glossy	Long	FLAVOR	Medium
Translucent	Medium	Acid	Short
BLOOM	Short	Foxy	COLOR
Abundance.....	Slender	Mild	Brown
Color.....	Thick	Spicy	Yellow
APEX	Swellings	Sweet	RAPHE
Adherence of stigma	Warts	Tart	Cordlike
ADHESION	BRUSH	Vinous	Large
Drops	Color.....	QUALITY	Small
Persistent	Length.....	Best	Obscure
CLUSTER	FLESH	Good	Prominent
SIZE	COLOR	Very good	CHALAZA
Large	Greenish	Medium	Above center
Medium	Red	Poor	Below center
Small	White	Very poor	Circular
FORM	Translucent	SEPARATION FROM SEED	Distinct
Cylindrical	Transparent	With difficulty	Obscure
Ovate	FIRMNESS	Easily	Oval
Roundish	Firm	SKIN	USE
Slender	Medium	Medium	Dessert
Tapering	Soft	Thick	Kitchen
		Thin	Market
		Tender	Raisin
		Tough	Shipping
			Wine

A STUDY OF THE GRAPE

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Make drawings of several different seeds, showing the position and the shape of the chalaza.

EXERCISE LXI

A STUDY OF THE PEACH

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Prolonged tip	QUALITY
Large	Roundish	Best
Medium	Dots	Very good
Small	Color.....	Good
FORM	Conspicuous	Fair
Compressed	Inconspicuous	Poor
Conical	Few	Very poor
Oblate	Numerous	USE
Oblong	Size.....	Dessert
Oval	FLESH	Kitchen
Round	Green	Market
Halves equal	Red	PIT
Halves unequal	White	ADHESION
COLOR	Yellow	Cling
Creamy	JUICE	Free
Green	Dry	Half free
Red	Juicy	SIZE
White	Color	Large
Yellow	Clear	Medium
Mottled	Dark	Small
CAVITY	Light	SHAPE
Deep	Pink	Angular
Medium	TEXTURE	Flattened
Shallow	Coarse	Necked
Narrow	Crisp	Oblique
Wide	Firm	Obovate
SUTURE	Fine	Oval
A mere line	Melting	Ovate
Indistinct	Stringy	Plump
Lacking	Tender	Blunt
Prominent	Medium	Pointed
Continuous	Tough	Prolonged
Deep	FLAVOR	Roundish
Shallow	Flat	Winged
APEX	Mild	SURFACE
Depressed	Sour	Pitted
Flattened	Sweet	Rough
Pointed	Sub-acid	Smooth

A STUDY OF THE PEACH

SEASON _____

LOCALITY _____

PRODUCTIVENESS _____

GENERAL APPEARANCE _____

DESIRABILITY _____

HISTORY _____

REMARKS _____

Draw a longitudinal section of the fruit and make a drawing of the seed.

EXERCISE LXII

A STUDY OF THE PEACH

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Prolonged tip Roundish	QUALITY
Large		Best
Medium		Very good
Small	DORS	Good
	Color.....	Fair
FORM	Conspicuous	Poor
Compressed	Inconspicuous	Very poor
Conical	Few	
Oblate	Numerous	USE
Oblong	Size.....	Dessert
Oval		Kitchen
Round	FLESH	Market
Halves equal	Green	
Halves unequal	Red	PIT
	White	ADHESION
COLOR	Yellow	Cling
Creamy		Free
Green	JUICE	Half free
Red	Dry	
White	Juicy	SIZE
Yellow	Color	Large
Mottled	Clear	Medium
	Dark	Small
CAVITY	Light	
Deep	Pink	SHAPE
Medium		Angular
Shallow	TEXTURE	Flattened
Narrow	Coarse	Necked
Wide	Crisp	Oblique
	Firm	Obovate
SUTURE	Fine	Oval
A mere line	Melting	Ovate
Indistinct	Stringy	Plump
Lacking	Tender	Blunt
Prominent	Medium	Pointed
Continuous	Tough	Prolonged
Deep		Roundish
Shallow	FLAVOR	Winged
	Flat	
APEX	Mild	SURFACE
Depressed	Sour	Pitted
Flattened	Sweet	Rough
Pointed	Sub-acid	Smooth

A STUDY OF THE PEACH

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and make a drawing of the seed.

EXERCISE LXIII

A STUDY OF THE PEACH

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Prolonged tip	QUALITY
Large	Roundish	Best
Medium	Dots	Very good
Small	Color.....	Good
FORM	Conspicuous	Fair
Compressed	Inconspicuous	Poor
Conical	Few	Very poor
Oblate	Numerous	USE
Oblong	Size.....	Dessert
Oval	FLESH	Kitchen
Round	Green	Market
Halves equal	Red	PIT
Halves unequal	White	ADHESION
COLOR	Yellow	Cling
Creamy	JUICE	Free
Green	Dry	Half free
Red	Juicy	SIZE
White	Color	Large
Yellow	Clear	Medium
Mottled	Dark	Small
CAVITY	Light	SHAPE
Deep	Pink	Angular
Medium	TEXTURE	Flattened
Shallow	Coarse	Necked
Narrow	Crisp	Oblique
Wide	Firm	Obovate
SUTURE	Fine	Oval
A mere line	Melting	Ovate
Indistinct	Stringy	Plump
Lacking	Tender	Blunt
Prominent	Medium	Pointed
Continuous	Tough	Prolonged
Deep	FLAVOR	Roundish
Shallow	Flat	Winged
APEX	Mild	SURFACE
Depressed	Sour	Pitted
Flattened	Sweet	Rough
Pointed	Sub-acid	Smooth

A STUDY OF THE PEACH

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and make a drawing of the seed.

EXERCISE LXIV

A STUDY OF THE PEACH

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Prolonged tip	QUALITY
Large	Roundish	Best
Medium	DOTS	Very good
Small	Color.....	Good
FORM	Conspicuous	Fair
Compressed	Inconspicuous	Poor
Conical	Few	Very poor
Oblate	Numerous	USE
Oblong	Size.....	Dessert
Oval	FLESH	Kitchen
Round	Green	Market
Halves equal	Red	PIT
Halves unequal	White	ADHESION
COLOR	Yellow	Cling
Creamy	JUICE	Free
Green	Dry	Half free
Red	Juicy	SIZE
White	Color	Large
Yellow	Clear	Medium
Mottled	Dark	Small
CAVITY	Light	SHAPE
Deep	Pink	Angular
Medium	TEXTURE	Flattened
Shallow	Coarse	Necked
Narrow	Crisp	Oblique
Wide	Firm	Obovate
SUTURE	Fine	Oval
A mere line	Melting	Ovate
Indistinct	Stringy	Plump
Lacking	Tender	Blunt
Prominent	Medium	Pointed
Continuous	Tough	Prolonged
Deep	FLAVOR	Roundish
Shallow	Flat	Winged
APEX	Mild	SURFACE
Depressed	Sour	Pitted
Flattened	Sweet	Rough
Pointed	Sub-acid	Smooth

A STUDY OF THE PEACH

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and make a drawing of the seed.

EXERCISE LXV

A STUDY OF THE PEACH

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Prolonged tip	QUALITY
Large	Roundish	Best
Medium		Very good
Small	DOTS	Good
	Color.....	Fair
FORM	Conspicuous	Poor
Compressed	Inconspicuous	Very poor
Conical	Few	
Oblate	Numerous	USE
Oblong	Size.....	Dessert
Oval		Kitchen
Round	FLESH	Market
Halves equal	Green	
Halves unequal	Red	PIT
	White	ADHESION
	Yellow	Cling
COLOR		Free
Creamy	JUICE	Half free
Green	Dry	
Red	Juicy	SIZE
White	Color	Large
Yellow	Clear	Medium
Mottled	Dark	Small
CAVITY	Light	
Deep	Pink	SHAPE
Medium		Angular
Shallow	TEXTURE	Flattened
Narrow	Coarse	Necked
Wide	Crisp	Oblique
	Firm	Obovate
SUTURE	Fine	Oval
A mere line	Melting	Ovate
Indistinct	Stringy	Plump
Lacking	Tender	Blunt
Prominent	Medium	Pointed
Continuous	Tough	Prolonged
Deep		Roundish
Shallow	FLAVOR	Winged
	Flat	
APEX	Mild	SURFACE
Depressed	Sour	Pitted
Flattened	Sweet	Rough
Pointed	Sub-acid	Smooth

A STUDY OF THE PEACH

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and make a drawing of the seed.

EXERCISE LXVI

A STUDY OF THE PEACH

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	Prolonged tip	QUALITY
Large	Roundish	Best
Medium	DOTS	Very good
Small	Color.....	Good
FORM	Conspicuous	Fair
Compressed	Inconspicuous	Poor
Conical	Few	Very poor
Oblate	Numerous	USE
Oblong	Size.....	Dessert
Oval	FLESH	Kitchen
Round	Green	Market
Halves equal	Red	PIT
Halves unequal	White	ADHESION
COLOR	Yellow	Cling
Creamy	JUICE	Free
Green	Dry	Half free
Red	Juicy	SIZE
White	Color	Large
Yellow	Clear	Medium
Mottled	Dark	Small
CAVITY	Light	SHAPE
Deep	Pink	Angular
Medium	TEXTURE	Flattened
Shallow	Coarse	Necked
Narrow	Crisp	Oblique
Wide	Firm	Obovate
SUTURE	Fine	Oval
A mere line	Melting	Ovate
Indistinct	Stringy	Plump
Lacking	Tender	Blunt
Prominent	Medium	Pointed
Continuous	Tough	Prolonged
Deep	FLAVOR	Roundish
Shallow	Flat	Winged
APEX	Mild	SURFACE
Depressed	Sour	Pitted
Flattened	Sweet	Rough
Pointed	Sub-acid	Smooth

A STUDY OF THE PEACH

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and make a drawing of the seed.

EXERCISE LXVII

A STUDY OF THE PEAR

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	COLOR	Tough	STAMENS
Large	Ground.....	Waxen	Basal
Medium	Overlying	CORE	Marginal
Small	Blushed	Abaxile	Medial
FORM	Mottled	Axile	CARPELS
Acute	Striped	Large	Cordate
Irregular	Polished.....	Medium	Emarginate
Oblong	Prevailing effect.....	Small	Ovate
Obovate	CALYX	Closed	Round
Ovate	Closed	Open	Tufted
Pyriform	Open	Distant	SEED
Ribbed	Large	Medial	Acuminate
Roundish	Medium	Sessile	Acute
Sides unequal	Small	CORE LINE	Flat
Symmetrical	CALYX LOBES	Clasping	Large
Turbinate	Acuminate	Meeting	Long
NECK	Acute	CALYX TUBE	Medium
Acute	Obtuse	Conical	Narrow
Distinct	Long	Funnel-shaped	Obtuse
Narrow	Medium	Urn-shaped	Plump
Obscure	Short	Long	Small
Obtuse	Narrow	Short	Tufted
Short	Wide	Narrow	Wide
STEM	Separated at base	Wide	FLAVOR
Long	BASIN	DOTS	Aromatic
Medium	Abrupt	Absent	Astringent
Short	Angular	Present	Flat
Slender	Corrugated	FLESH	Neutral
Thick	Furrowed	Color	Sprightly
CAVITY	Obtuse	TEXTURE	Sour
Acuminate	Symmetrical	Buttery	Sub-acid
Acute	Wrinkled	Coarse	Sweet
Furrowed	Deep	Crisp	QUALITY
Lipped	Medium	Dry	Best
Obtuse	Shallow	Granular	Very good
Russeted	Narrow	Fibrous	Good
Symmetrical	Wide	Fine	Medium
Narrow	SKIN	Firm	Poor
Wide	Dull	Juicy	Very poor
Deep	Glossy	Medium	MARKET
Medium	Rough	Sweet	Distant
Shallow	Russet	Tender	Local
	Smooth	Tough	USE
	Tender		Cooking
	Thin		Dessert
			Market

A STUDY OF THE PEAR

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE LXVIII

A STUDY OF THE PEAR

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	COLOR	Tough	STAMENS
Large	Ground.....	Waxen	Basal
Medium	Overlying	CORE	Marginal
Small	Blushed	Abaxile	Medial
FORM	Mottled	Axile	CARPELS
Acute	Striped	Large	Cordate
Irregular	Polished.....	Medium	Emarginate
Oblong	Prevailing effect.....	Small	Ovate
Obovate	CALYX	Closed	Round
Ovate	Open	Open	Tufted
Pyriform	Large	Distant	SEED
Ribbed	Medium	Medial	Acuminate
Roundish	Small	Sessile	Acute
Sides unequal	CALYX LOBES	CORE LINE	Flat
Symmetrical	Acuminate	Clasping	Large
Turbinate	Acute	Meeting	Long
	Obtuse	CALYX TUBE	Medium
NECK	Long	Conical	Narrow
Acute	Medium	Funnel-shaped	Obtuse
Distinct	Short	Urn-shaped	Plump
Narrow	Narrow	Long	Small
Obscure	Wide	Short	Tufted
Obtuse	Separated at base	Narrow	Wide
Short	BASIN	Wide	FLAVOR
STEM	Abrupt	DOTS	Aromatic
Long	Angular	Absent	Astringent
Medium	Corrugated	Present	Flat
Short	Furrowed	FLESH	Neutral
Slender	Obtuse	Color.....	Sprightly
Thick	Symmetrical	TEXTURE	Sour
CAVITY	Wrinkled	Buttery	Sub-acid
Acuminate	Deep	Coarse	Sweet
Acute	Medium	Crisp	QUALITY
Furrowed	Shallow	Dry	Best
Lipped	Narrow	Granular	Very good
Obtuse	Wide	Fibrous	Good
Russeted	SKIN	Fine	Medium
Symmetrical	Dull	Firm	Poor
Narrow	Glossy	Juicy	Very poor
Wide	Rough	Medium	MARKET
Deep	Russet	Sweet	Distant
Medium	Smooth	Tender	Local
Shallow	Tender	Tough	USE
	Thick		Cooking
	Thin		Dessert
			Market

A STUDY OF THE PEAR

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE LXIX

A STUDY OF THE PEAR

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	COLOR	Tough	STAMENS
Large	Ground	Waxen	Basal
Medium	Overlying	CORE	Marginal
Small	Blushed	Abaxile	Medial
FORM	Mottled	Axile	CARPELS
Acute	Striped	Large	Cordate
Irregular	Polished	Medium	Emarginate
Oblong	Prevailing effect	Small	Ovate
Obovate	CALYX	Closed	Round
Ovate	Closed	Open	Tufted
Pyriform	Open	Distant	SEED
Ribbed	Large	Medial	Acuminate
Roundish	Medium	Sessile	Acute
Sides unequal	Small	CORE LINE	Flat
Symmetrical	CALYX LOBES	Clasping	Large
Turbinate	Acuminate	Meeting	Long
NECK	Acute	CALYX TUBE	Medium
Acute	Obtuse	Conical	Narrow
Distinct	Long	Funnel-shaped	Obtuse
Narrow	Medium	Urn-shaped	Plump
Obscure	Short	Long	Small
Obtuse	Narrow	Short	Tufted
Short	Wide	Narrow	Wide
STEM	Separated at base	Wide	FLAVOR
Long	BASIN	DOTS	Aromatic
Medium	Abrupt	Absent	Astringent
Short	Angular	Present	Flat
Slender	Corrugated	FLESH	Neutral
Thick	Furrowed	Color	Sprightly
CAVITY	Obtuse	TEXTURE	Sour
Acuminate	Symmetrical	Buttery	Sub-acid
Acute	Wrinkled	Coarse	Sweet
Furrowed	Deep	Crisp	QUALITY
Lipped	Medium	Dry	Best
Obtuse	Shallow	Granular	Very good
Russeted	Narrow	Fibrous	Good
Symmetrical	Wide	Fine	Medium
Narrow	SKIN	Firm	Poor
Wide	Dull	Juicy	Very poor
Deep	Glossy	Medium	MARKET
Medium	Russet	Sweet	Distant
Shallow	Smooth	Tender	Local
	Thick	Tough	USE
	Thin		Cooking
			Dessert
			Market

A STUDY OF THE PEAR

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE LXX

A STUDY OF THE PEAR

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	COLOR	Tough	STAMENS
Large	Ground.....	Waxen	Basal
Medium	Overlying	CORE	Marginal
Small	Blushed	Abaxile	Medial
FORM	Mottled	Axile	CARPELS
Acute	Striped	Large	Cordate
Irregular	Polished.....	Medium	Emarginate
Oblong	Prevailing effect.....	Small	Ovate
Obovate	CALYX	Closed	Round
Ovate	Closed	Open	Tufted
Pyriform	Open	Distant	SEED
Ribbed	Large	Medial	Acuminate
Roundish	Medium	Sessile	Acute
Sides unequal	Small	CORE LINE	Flat
Symmetrical	CALYX LOBES	Clasping	Large
Turbinate	Acuminate	Meeting	Long
NECK	Acute	CALYX TUBE	Medium
Acute	Obtuse	Conical	Narrow
Distinct	Long	Funnel-shaped	Obtuse
Narrow	Medium	Urn-shaped	Plump
Obscure	Short	Long	Small
Obtuse	Narrow	Short	Tufted
Short	Wide	Narrow	Wide
STEM	Separated at base	Wide	FLAVOR
Long	BASIN	Dots	Aromatic
Medium	Abrupt	Absent	Astringent
Short	Angular	Present	Flat
Slender	Corrugated	FLESH	Neutral
Thick	Furrowed	Color.....	Sprightly
CAVITY	Obtuse	Texture	Sour
Acuminate	Symmetrical	Buttery	Sub-acid
Acute	Wrinkled	Coarse	Sweet
Furrowed	Deep	Crisp	QUALITY
Lipped	Medium	Dry	Best
Obtuse	Shallow	Granular	Very good
Russeted	Narrow	Fibrous	Good
Symmetrical	Wide	Fine	Medium
Narrow	SKIN	Firm	Poor
Wide	Dull	Juicy	Very poor
Deep	Glossy	Medium	MARKET
Medium	Rough	Sweet	Distant
Shallow	Russet	Tender	Local
	Smooth	Tough	USE
	Tender		Cooking
	Thick		Dessert
	Thin		Market

A STUDY OF THE PEAR

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit.

EXERCISE LXXI

A STUDY OF THE PEAR

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	COLOR	Tough	STAMENS
Large	Ground.....	Waxen	Basal
Medium	Overlying	CORE	Marginal
Small	Blushed	Abaxile	Medial
FORM	Mottled	Axile	CARPELS
Acute	Striped	Large	Cordate
Irregular	Polished.....	Medium	Emarginate
Oblong	Prevailing effect.....	Small	Ovate
Obovate	CALYX	Closed	Round
Ovate	Closed	Open	Tufted
Pyriform	Open	Distant	SEED
Ribbed	Large	Medial	Acuminate
Roundish	Medium	Sessile	Acute
Sides unequal	Small	CORE LINE	Flat
Symmetrical	CALYX LOBES	Clasping	Large
Turbinate	Acuminate	Meeting	Long
	Acute	CALYX TUBE	Medium
	Obtuse	Conical	Narrow
NECK	Long	Funnel-shaped	Obtuse
Acute	Medium	Urn-shaped	Plump
Distinct	Short	Long	Small
Narrow	Narrow	Short	Tufted
Obscure	Wide	Narrow	Wide
Obtuse	Separated at base	Wide	FLAVOR
Short	BASIN	DOTS	Aromatic
STEM	Abrupt	Absent	Astringent
Long	Angular	Present	Flat
Medium	Corrugated	FLESH	Neutral
Short	Furrowed	Color.....	Sprightly
Slender	Obtuse	TEXTURE	Sour
Thick	Symmetrical	Buttery	Sub-acid
CAVITY	Wrinkled	Coarse	Sweet
Acuminate	Deep	Crisp	QUALITY
Acute	Medium	Dry	Best
Furrowed	Shallow	Granular	Very good
Lipped	Narrow	Fibrous	Good
Obtuse	Wide	Firm	Medium
Russeted	SKIN	Juicy	Poor
Symmetrical	Dull	Medium	Very poor
Narrow	Glossy	Sweet	MARKET
Wide	Rough	Tender	Distant
Deep	Russet	Tough	Local
Medium	Smooth		USE
Shallow	Tender		Cooking
	Thick		Dessert
	Thin		Market

A STUDY OF THE PEAR

SEASON _____

LOCALITY _____

PRODUCTIVENESS _____

GENERAL APPEARANCE _____

DESIRABILITY _____

HISTORY _____

REMARKS _____

Draw a longitudinal section of the fruit.

EXERCISE LXXII

A STUDY OF THE PEAR

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	COLOR	Tough	STAMENS		
Large	Ground.....	Waxen	Basal		
Medium	Overlying	CORE	Marginal		
Small	Blushed		Medial		
FORM	Mottled		Abaxile	CARPELS	
	Striped	Axile	Cordate		
	Polished.....	Large	Emarginate		
	Prevailing effect.....	Medium	Ovate		
Acute	CALYX	Small	Round		
Irregular		Closed	Tufted		
Oblong		Open	SEED		
Obovate	Distant				
Ovate	Large	Acuminate			
Pyriform	Medium	Acute			
Ribbed	Small	Sessile	Flat		
Roundish	CALYX LOBES	CORE LINE	Large		
Sides unequal			Clasping	Long	
Symmetrical			Meeting	Medium	
Turbinate		Obtuse	Narrow		
NECK	Long	CALYX TUBE	Obtuse		
	Medium	Conical	Plump		
	Short	Funnel-shaped	Small		
	Narrow	Urn-shaped	Tufted		
Obscure	Narrow	Long	Wide		
Obtuse	Separated at base	Short	FLAVOR		
Short	BASIN	Narrow		Aromatic	
STEM		Wide		DOTS	Astringent
		Abrupt	Absent	Flat	
	Angular	Present	Neutral		
	Long	Corrugated	FLESH	Sprightly	
Medium	Furrowed	Sour			
Short	Obtuse	Sub-acid			
Slender	Symmetrical	COLOR.....	Sweet		
Thick	Wrinkled		TEXTURE	QUALITY	
CAVITY	Deep	Buttery			Best
	Medium	Coarse			Very good
	Shallow	Crisp	Good		
	Acuminate	Narrow	Dry	Medium	
Acute	Wide	Granular	Poor		
Furrowed	SKIN	Fibrous	Very poor		
Lipped		Dull	MARKET	USE	
Obtuse		Glossy			Distant
Russeted	Rough	Local			
Symmetrical	Russet	Juicy	Cooking		
Narrow	Smooth	Medium		Dessert	
Wide	Tender	Sweet		Market	
Deep	Thick	Tender			
Medium	Thin	Tough			
Shallow					

A STUDY OF THE PEAR

SEASON _____

LOCALITY _____

PRODUCTIVENESS _____

GENERAL APPEARANCE _____

DESIRABILITY _____

HISTORY _____

REMARKS _____

Draw a longitudinal section of the fruit.

EXERCISE LXXIII

A STUDY OF THE PLUM

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	PUBESCENT.....	DOTS	QUALITY
Large		Few	Best
Medium	ANGLE OF STEM.....	Many	Very good
Small		Large	Good
	CAVITY	Small	Fair
FORM	Acuminate	Color	Poor
Compressed	Acute	Conspicuous.....	Very poor
Conical	Lipped		
Cordate	Symmetrical	FLESH	USE
Halves equal	Narrow	Colorless	Cooking
Halves unequal	Wide	Green	Dessert
Necked	Deep	Purple	Market
Oblate	Medium	Red	
Oblong	Shallow	White	PIT
Oval		Yellow	ADHESION
Ovate			Cling
Pyriform	SETURE	JUICE	Free
Roundish	A mere line	Dry	Half free
Truncate	Continuous	Juicy	
	Indistinct	COLOR	SIZE
COLOR	Lacking	Clear	Large
Amber	Prominent	Dark-colored	Medium
Black	Deep	Light-colored	Small
Blue	Shallow	Pink	
Crimson		TEXTURE	SHAPE
Mottled	APEX	Coarse	Angular
Purple	Depressed	Crisp	Blunt
Red	Flattened	Fine-grained	Flattened
White	Pointed	Firm	Necked
Yellow	Prolonged tip	Melting	Oblique
	Roundish	Stringy	Obovate
BLOOM		Tender	Oval
Blue	SKIN	Medium	Ovate
Gray	Astringent	Tough	Plump
Lilac	Medium		Pointed
White	Thick	FLAVOR	Prolonged at ends
Thick	Thin	Aromatic	Roundish
Thin	Tender	Astringent	Winged
STEM	Tough	Flat	
Slender	Translucent	Mild	
Medium	Adherence to	Sour	SURFACE
Thick	flesh	Sprightly	Pitted
Long	Adherent	Sub-acid	Rough
Short	Free	Sweet	Smooth
		Tart	

A STUDY OF THE PLUM

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and a side view of the seed.

EXERCISE LXXIV

A STUDY OF THE PLUM

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	PUBESCENT.....	DOTS	QUALITY
Large		Few	Best
Medium	ANGLE OF STEM.....	Many	Very good
Small		Large	Good
	CAVITY	Small	Fair
FORM	Acuminate	Color.....	Poor
Compressed	Acute	Conspicuous.....	Very poor
Conical	Lipped		
Cordate	Symmetrical	FLESH	USE
Halves equal	Narrow	Colorless	Cooking
Halves unequal	Wide	Green	Dessert
Necked	Deep	Purple	Market
Oblate	Medium	Red	
Oblong	Shallow	White	PIT
Oval		Yellow	ADHESION
Ovate			Cling
Pyriform	SUTURE	JUICE	Free
Roundish	A mere line	Dry	Half free
Truncate	Continuous	Juicy	
	Indistinct	COLOR	SIZE
COLOR	Lacking	Clear	Large
Amber	Prominent	Dark-colored	Medium
Black	Deep	Light-colored	Small
Blue	Shallow	Pink	
Crimson		TEXTURE	SHAPE
Mottled	APEX	Coarse	Angular
Purple	Depressed	Crisp	Blunt
Red	Flattened	Fine-grained	Flattened
White	Pointed	Firm	Necked
Yellow	Prolonged tip	Melting	Oblique
	Roundish	Stringy	Obovate
BLOOM		Tender	Oval
Blue	SKIN	Medium	Ovate
Gray	Astringent	Tough	Plump
Lilac	Medium	FLAVOR	Pointed
White	Thick	Aromatic	Prolonged at ends
Thick	Thin	Astringent	Roundish
Thin	Tender	Flat	Winged
STEM	Tough	Mild	
Slender	Translucent	Sour	
Medium	Adherence to flesh	Sprightly	SURFACE
Thick	Adherent	Sub-acid	Pitted
Long	Free	Sweet	Rough
Short		Tart	Smooth

A STUDY OF THE PLUM

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and a side view of the seed.

EXERCISE LXXV

A STUDY OF THE PLUM

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	PUBESCENT.....	DOTS	QUALITY
Large		Few	Best
Medium	ANGLE OF STEM.....	Many	Very good
Small		Large	Good
	CAVITY	Small	Fair
FORM	Acuminate	Color	Poor
Compressed	Acute	Conspicuous.....	Very poor
Conical	Lipped		
Cordate	Symmetrical	FLESH	USE
Halves equal	Narrow	Colorless	
Halves unequal	Wide	Green	Cooking
Necked	Deep	Purple	Dessert
Oblate	Medium	Red	Market
Oblong	Shallow	White	
Oval		Yellow	PIT
Ovate			ADHESION
Pyriform	SUTURE	JUICE	
Roundish	A mere line	Dry	Cling
Truncate	Continuous	Juicy	Free
	Indistinct		Half free
COLOR	Lacking	COLOR	SIZE
Amber	Prominent	Clear	
Black	Deep	Dark-colored	Large
Blue	Shallow	Light-colored	Medium
Crimson		Pink	Small
Mottled			
Purple	APEX	TEXTURE	SHAPE
Red	Depressed	Coarse	
White	Flattened	Crisp	Angular
Yellow	Pointed	Fine-grained	Blunt
	Prolonged tip	Firm	Flattened
	Roundish	Melting	Necked
		Stringy	Oblique
BLOOM		Tender	Obovate
Blue	SKIN	Medium	Oval
Gray	Astringent	Tough	Ovate
Lilac	Medium		Pump
White	Thick	FLAVOR	Pointed
Thick	Thin	Aromatic	Prolonged at
Thin	Tender	Astringent	ends
STEM	Tough	Flat	Roundish
Slender	Translucent	Mild	Winged
Medium	Adherence to	Sour	
Thick	flesh	Sprightly	SURFACE
Long	Adherent	Sub-acid	
Short	Free	Sweet	Pitted
		Tart	Rough
			Smooth

A STUDY OF THE PLUM

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and a side view of the seed.

EXERCISE LXXVI

A STUDY OF THE PLUM

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	PUBESCENT.....	DOTS	QUALITY
Large		Few	Best
Medium	ANGLE OF STEM.....	Many	Very good
Small		Large	Good
	CAVITY	Small	Fair
FORM	Acuminate	Color	Poor
Compressed	Acute	Conspicuous.....	Very poor
Conical	Lipped	FLESH	USE
Cordate	Symmetrical	Colorless	
Halves equal	Narrow	Green	Cooking
Halves unequal	Wide	Purple	Dessert
Necked	Deep	Red	Market
Oblate	Medium	White	PIT
Oblong	Shallow	Yellow	ADHESION
Oval		JUICE	
Ovate	SUTURE	Dry	Cling
Pyriform	A mere line	Juicy	Free
Roundish	Continuous	COLOR	Half free
Truncate	Indistinct	Clear	SIZE
COLOR	Lacking	Dark-colored	Large
Amber	Prominent	Light-colored	Medium
Black	Deep	Pink	Small
Blue	Shallow	TEXTURE	SHAPE
Crimson	APEX	Coarse	
Mottled	Depressed	Crisp	Angular
Purple	Flattened	Fine-grained	Blunt
Red	Pointed	Firm	Flattened
White	Prolonged tip	Melting	Necked
Yellow	Roundish	Stringy	Oblique
BLOOM	SKIN	Tender	Obovate
Blue	Astringent	Medium	Oval
Gray	Medium	Tough	Ovate
Lilac	Thick	FLAVOR	Plump
White	Thin	Aromatic	Pointed
Thick	Tender	Astringent	Prolonged at ends
Thin	Tough	Flat	Roundish
STEM	Translucent	Mild	Winged
Slender	Adherence to flesh	Sour	
Medium	Adherent	Sprightly	SURFACE
Thick	Free	Sub-acid	
Long		Sweet	Pitted
Short		Tart	Rough
			Smooth

A STUDY OF THE PLUM

SEASON _____

LOCALITY _____

PRODUCTIVENESS _____

GENERAL APPEARANCE _____

DESIRABILITY _____

HISTORY _____

REMARKS _____

Draw a longitudinal section of the fruit and a side view of the seed.

EXERCISE LXXVII

A STUDY OF THE PLUM

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	PUBESCENT.....	DOTS	QUALITY
Large		Few	Best
Medium	ANGLE OF STEM.....	Many	Very good
Small		Large	Good
	CAVITY	Small	Fair
FORM	Acuminate	Color	Poor
Compressed	Acute	Conspicuous.....	Very poor
Conical	Lipped	FLESH	USE
Cordate	Symmetrical	Colorless	
Halves equal	Narrow	Green	Cooking
Halves unequal	Wide	Purple	Dessert
Necked	Deep	Red	Market
Oblate	Medium	White	PIT
Oblong	Shallow	Yellow	ADHESION
Oval		JUICE	
Ovate	SUTURE	Dry	Cling
Pyriform	A mere line	Juicy	Free
Roundish	Continuous		Half free
Truncate	Indistinct	COLOR	SIZE
	Lacking	Clear	
COLOR	Prominent	Dark-colored	Large
Amber	Deep	Light-colored	Medium
Black	Shallow	Pink	Small
Blue		TEXTURE	SHAPE
Crimson	APEX	Coarse	
Mottled	Depressed	Crisp	Angular
Purple	Flattened	Fine-grained	Blunt
Red	Pointed	Firm	Flattened
White	Prolonged tip	Melting	Necked
Yellow	Roundish	Stringy	Oblique
		Tender	Obovate
BLOOM	SKIN	Medium	Oval
Blue	Astringent	Tough	Ovate
Gray	Medium	FLAVOR	Plump
Lilac	Thick	Aromatic	Pointed
White	Thin	Astringent	Prolonged at ends
Thick	Tender	Flat	Roundish
Thin	Tough	Mild	Winged
STEM	Translucent	Sour	
Slender	Adherence to flesh	Sprightly	SURFACE
Medium	Adherent	Sub-acid	Pitted
Thick	Free	Sweet	Rough
Long		Tart	Smooth
Short			

A STUDY OF THE PLUM

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and a side view of the seed.

EXERCISE LXXVIII

A STUDY OF THE PLUM

(Underscore or star the terms that apply to the variety of fruit being studied)

VARIETY _____ DATE _____

GROUP _____

SYNONYMS _____

SIZE	PUBESCENT.....	DOTS	QUALITY
Large		Few	Best
Medium	ANGLE OF STEM.....	Many	Very good
Small		Large	Good
	CAVITY	Small	Fair
FORM	Acuminate	Color	Poor
Compressed	Acute	Conspicuous.....	Very poor
Conical	Lipped	FLESH	USE
Cordate	Symmetrical	Colorless	
Halves equal	Narrow	Green	Cooking
Halves unequal	Wide	Purple	Dessert
Necked	Deep	Red	Market
Oblate	Medium	White	PIT
Oblong	Shallow	Yellow	ADHESION
Oval		JUICE	
Ovate	SUTURE	Dry	Cling
Pyriform	A mere line	Juicy	Free
Roundish	Continuous	COLOR	Half free
Truncate	Indistinct	Clear	SIZE
COLOR	Lacking	Dark-colored	Large
Amber	Prominent	Light-colored	Medium
Black	Deep	Pink	Small
Blue	Shallow	TEXTURE	SHAPE
Crimson		Coarse	
Mottled	APEX	Crisp	Angular
Purple	Depressed	Fine-grained	Blunt
Red	Flattened	Firm	Flattened
White	Pointed	Melting	Necked
Yellow	Prolonged tip	Stringy	Oblique
BLOOM	Roundish	Tender	Obovate
Blue	SKIN	Medium	Oval
Gray	Astringent	Tough	Ovate
Lilac	Medium	FLAVOR	Plump
White	Thick	Aromatic	Pointed
Thick	Thin	Astringent	Prolonged at ends
Thin	Tender	Flat	Roundish
STEM	Tough	Mild	Winged
Slender	Translucent	Sour	
Medium	Adherence to	Sprightly	SURFACE
Thick	flesh	Sub-acid	
Long	Adherent	Sweet	Pitted
Short	Free	Tart	Rough
			Smooth

A STUDY OF THE PLUM

SEASON_____

LOCALITY_____

PRODUCTIVENESS_____

GENERAL APPEARANCE_____

DESIRABILITY_____

HISTORY_____

REMARKS_____

Draw a longitudinal section of the fruit and a side view of the seed.

GLOSSARY

DESCRIPTIVE TERMS

This list is intended to cover most of the important terms which might be confusing to the student. Several general terms used in describing fruit are listed first, and the remaining terms follow in alphabetical order under the various headings.

GENERAL TERMS

Acute. Tapering nearly to a point.

Apex. A term understood as applying to the part most remote from the branch or root. (In pears, this part is usually spoken of as the crown.)

Axis. A line connecting the base and the apex.

A longitudinal section of a fruit is made by cutting it from base to apex; a cross, or transverse, section is made by cutting it at right angles to the axis.

Base (of a fruit). The portion toward the branch or root.

Compressed. With the sides flattened, as in some plums.

Depressed. Sunken, as the apex of peaches.

Furrowed (of fruit). More or less regularly channelled.

Irregular. A term applied to an outline that is elliptical or angular.

Obtuse. Blunt; approximating an obtuse angle.

Pit. A term applied to the seed found in drupaceous fruits.

Regular. When a section through the equatorial diameter shows nearly a circular outline, the fruit is said to be regular.

Ribbed. Having large obtuse and rounded ridges.

Symmetrical. Having parts mutually well proportioned; regular.

Truncate. Abruptly flattened at the end, or (of fruit) appearing as though cut squarely across.

FIG. 38. A, apex; B, base

BASIN

The depression in which the calyx is set is technically called the *basin*, and several terms are used in describing it.

Abrupt basin. A basin in which the sides show a sudden slope.

Corrugated basin. A basin in which the furrows are wrinkled or folded about the calyx lobes.

Mammiform basin. A basin in which peculiar fleshy protuberances occur about the base of the calyx lobes.

LABORATORY MANUAL OF HORTICULTURE

Obtuse basin. A basin in which the sides show a very gradual slope.

Regular basin. A basin in which the sides are smooth.

Ridged, ribbed, and furrowed. Terms expressing slightly different degrees of the condition in which depressed lines extend up the sides of the basin.

FIG. 39. A, narrow basin; B, wide basin; C, obtuse basin;
D, abrupt basin

Symmetrical basin. A basin that is nearly round, or uniform in shape.

Wavy basin. A basin in which the furrows are indistinct.

CALYX

The lower part of the flower is called the *calyx*. The outer green coverings of the flower bud are called the *calyx lobes*. Calyx lobes remain visible in the mature fruit of some apples, pears, and quinces.

Closed calyx. A calyx whose lobes fold over each other in such a way that the center is completely obscured.

Connivent. A term applied to calyx lobes which are upright and have the tip inclined toward the axis.

Convergent. A term applied to calyx lobes which are flat and inclined toward the axis.

Deciduous. A term applied to calyx lobes which fall away, like those of the Sand Pears and of some pure-Siberian-crab species.

Divergent or reflexed. Terms applied to calyx lobes which turn backward.

Open calyx. A calyx whose lobes flare out in such a way that the center is clearly exposed.

Partly open calyx. A calyx whose condition is intermediate between the open and the closed state.

Persistent calyx. A calyx whose lobes remain attached to the fruit, as in some apples, pears, and quinces.

FIG. 40. A, basin; B, calyx;
C, axis; D, cavity; E, stem

In examining different specimens of the same variety of a species we often find great variation in the above features.

GLOSSARY

CALYX TUBE

We recognize three kinds, or shapes, of calyx tubes; namely, the cone-shaped, the funnel-shaped, and the urn-shaped.

Cone-shaped calyx tube. A calyx tube resembling, or conforming to, the general shape of a cone.

Funnel-shaped calyx tube. A calyx tube whose general shape resembles that of a funnel, or which has a little tube attached to the base of the cone-shaped tube.

Urn-shaped calyx tube. A calyx tube resembling an urn in shape, or having sides which are neither straight nor funnel-form, but which have a double curve and are comparatively broad and rounded toward the base.

CAVITY

The depression around the stem of a fruit is technically called the *cavity*.

Acuminate cavity. A cavity in which the stem meets the side of the cavity in a sharp, narrow angle.

FIG. 41. A, obtuse cavity ; B, acuminate cavity ;
C, acute cavity

Acute cavity. A cavity intermediate between an acuminate and an obtuse cavity, as in the Canada Red apple.

Obtuse cavity. A cavity in which the angle between the stem and the side is wide and blunt, as in the Rome Beauty apple.

In addition to the above we speak of the cavity as being *wide* or *narrow*, and *deep*, *medium*, or *shallow*.

COLOR

The *color* of fruit is quite varied and ranges from the light yellow of the white fruits, through the various shades of red, into deep purple. This condition can better be explained by several examples. Thus, we have the white flesh of the snow peach, followed by pale yellows shading gradually into the deeper colors. Again, we find cherries and plums gradually shading from the yellowish tints through to orange and the deeper reds; and the grapes follow with blues, violets, and purples. Amber and fawn with slight mixtures of grays are also found. In addition to solid color in fruits, we find various striped and mottled effects.

LABORATORY MANUAL OF HORTICULTURE

Blotched. Having different abrupt shades more or less blended together without any order or regularity.

Blushed. Distinguished from mottled, striped, or splashed by having the surface overspread with a red tint that is not much broken.

Clouded. Having blotches broader and more softly shaded and blended together.

Dotted. Having small, distinct dots.

Marbled. Having wide, faint, irregular, or wavy stripes.

Mottled. Covered with dots that run together without any marked lines of distinction.

Scarf skin. A term descriptive of a condition found on fruit (principally on apples), which gives a characteristic appearance to the fruit. It usually extends outward from the base in whitish lines, or stripes, readily distinguished by contrast with the green or yellow color.

Splashed. Having stripes of color which are much broken and of varying widths.

Spotted. Having larger dots than those characterizing a fruit as "dotted."

Stained. Having the lighter shades of the blotched or clouded effect.

Streaked. Having long and narrow color bands.

Striped. Having one or more shades of color alternating in broad lines.

FIG. 42. Abaxile core

CORE

In describing the *core*, its location in the fruit and the size and character of the carpels and seeds are all considered.

Abaxile. A term used to describe the core when it lies to one side of the axis of the fruit.

Axile. A term used to describe the core when it is equidistant on either side of the axis of the fruit.

Distant. A term used to describe a core which is comparatively far from the stem.

Median. A term used to describe a core which is located at about the center of the fruit.

Sessile. A term used to describe a core which is very close to the stem.

CARPELS

Carpels are the parchmentlike walls of the seed cells. They vary in form somewhat as leaves do, and the terms used in describing leaves may well be applied in describing the carpels. The two sides of the

GLOSSARY

seed cells correspond to the two halves of a leaf, with the axis of the fruit representing the midrib of the leaf.

Cordate. A term used to describe a carpel whose two halves are heart-shaped.

Emarginate. A term used to describe carpels which are indented at the top or bottom, but too narrow to be cordate.

Mucronate. A term used to describe carpels which come to a sharp point.

Obcordate. Reversed cordate.

Round. A term used to describe a carpel whose two halves are nearly circular.

CORE LINE

The *core line* is that portion of the fleshy part of the apple surrounding the seed cavities. It constitutes part of the core and is limited by visible lines.

Clasping core line. A core line which appears to join the calyx tube somewhat above the base.

Meeting core line. A core line which joins the calyx tube near or at its base.

Dots

Dots are small, irregular specks found on some fruits. They are described by the following terms:

Areolar. A term used to describe dots which are surrounded by a halo of a paler or brighter color. The halo is decidedly conspicuous on some varieties and inconspicuous on others.

Depressed. A term used to describe dots which are sunken as indicated by touch.

Elongate. A term used to describe dots which are stretched out, or approach the general shape of a rectangle. These dots are often seen on the base of the fruit, along lines radiating from the cavity.

Stellate. A term used to describe dots which approach the form of a star.

Submerged. A term used to describe dots which are visible under the epidermis.

FLAVOR

The *flavor* of fruit is determined by the acids, sugars, aromatic oils, and other constituents found in the fruit, and it varies slightly with each fruit.

Acid. Possessing a sour taste.

Astringent. Possessing the power of contracting the muscles; puckery.

Neutral. Neither sweet nor sour, but intermediate between the two.

Perfumed. Possessing pleasing odors.

Spicy. Having an aromatic or fragrant flavor resembling spices.

Vinous. Rich and highly flavored, bordering on acid.

In addition to the above we have *sweet*, *sour*, *mild*, and several other terms that are self-explanatory.

FIG. 43. Axile core

A, core line

LABORATORY MANUAL OF HORTICULTURE

FORM

The *form* of a fruit means the shape and is described by the following terms:

Angular. Having sides more or less ribbed.

Conical. Tapering gradually from the base to the apex.

Globose, globular, or round. Terms used to describe fruit which is nearly spherical, as the Fameuse.

Oblate. Flattened, having the height less than the breadth.

Oblique. Having an axis which slants obliquely. This form is sometimes called lopsided.

Oblong. A term used to describe a fruit whose length exceeds its breadth and whose sides are nearly parallel.

Obovate. Reversed ovate; a term used to describe an egg-shaped fruit whose smaller end is at the base instead of at the apex.

Ovate, egg-shaped. Terms used to describe fruit whose length somewhat exceeds its breadth, and which has a slightly rounded taper from the base to the apex.

Roundish. Varying slightly from round, or having the length and the breadth nearly equal.

FIG. 44. Oblong

SKIN

The *skin* of the different fruits becomes much modified because of the varying conditions under which they are grown. The texture, form, color, and any other characteristics should be noted.

Bloom. A delicate powdery substance, varying in color and found on many fruits. It is seen most frequently on grapes and plums, and occasionally on apples.

Pubescence. Hairiness; a condition sometimes found on the skin of certain fruit. This condition is particularly noticeable on certain portions of the fruit; for example, around the calyx.

Rough skin. A term applied to the surface of fruit when minute capillary russet netted veins or russet dots or both are scattered over the skin.

Russet skin. A condition found on some apples in which all or most of the surface is covered with russet, as seen on the Golden Russet.

Waxy, or oily, skin. Found on some fruits, particularly the apple. This character can best be determined by the sense of touch, the feeling of such a skin much resembling that of a greasy surface.

SIZE

Fruits are usually graded into seven sizes as follows: *very small, small, medium, below medium, above medium, large, very large.*

It is apparent that, owing to the great variation in fruits, no definite line can be drawn between these terms, and they must of necessity be only

FIG. 45. Conical

FIG. 46. Oblate-conical

FIG. 47. Oblate

FIG. 48. Obovate

FIG. 49. Ovate

FIG. 50. Round-ovate

LABORATORY MANUAL OF HORTICULTURE

comparative. Among apples the crabs are usually referred to as being small or very small, the Jonathan or Baldwin as medium, and the Wolf River as large or very large. The other varieties of apples can be gauged according to these already mentioned.

STAMENS

Basal. A term applied to the stamens when they are located near the base of the calyx tube.

Marginal. A term applied to the stamens when they are located near the outer margin of the calyx tube.

Medial. A term applied to the stamens when they are located about the middle of the calyx tube.

STEM

The character of the *stem* is of some importance, although it varies much on different individual fruits of the same variety. Unfortunately, however, there are comparatively few varieties which show a stem so characteristic that the fruit may be recognized by this character alone. In this case, as with the size, the descriptive terms must of necessity only be comparative. We speak of a stem as being *long* or *short*, *medium*, *slender*, *thick*, *fleshy*, or *warty*. *Lipped* is a term which signifies that the flesh forms a protuberance, or lip, under which the stem is inserted; for example, the Pewaukee apple.

SUTURE

Suture (meaning "seam") is more conspicuous on the stone fruits than on any other class. It is a furrow, or depression, extending from the base to the apex on either one or both sides of the fruit.

Distinct. A term used to describe a suture in which the seam is clear and well defined.

Large. A term used to describe a suture in which the seam is wide and deep.

Obscure. A term used to describe a suture in which the seam is faint or indistinct.

Obsolete. A term used to describe fruit in which the suture either does not exist or exists only as a faint line on the surface.

SPECIAL TERMS FOR THE GRAPE

Foxy. Possessing a sour, unpleasant taste, as found in some grapes which have a coarse flavor, like the fox grapes.

Brush. The bristlelike portion left attached to the pedicel when the berry is removed; the central portion of the berry, which usually pulls out of the flesh and remains attached to the pedicel.

Pedicel. The divisions or branches of the peduncle.

Peduncle. A flower stalk; the stem that supports the flowers or fruit.

Pigment. Opaque coloring matter deposited in the plant cell or tissue.

Raphe. The seamlike union of two halves of an organ; in anatropous ovules, that part of the funicle adnate to the integument forming a ridge

GLOSSARY

along the body of the ovule. A medium line, or suture, between two pericarps. The line found on the seed of the grape.

Chalaza. The depression found near the center of the seed. The point at which the nucellus and integuments of the ovule are united.

SPECIAL TERMS FOR THE PEAR

Neck. The narrow part toward the stalk. The neck may be *long*, *narrow*, *short*, *obtuse*, *acute*, *distinct*, or *obscure*.

Body. The larger part of the fruit toward the crown. The body may be *small*, *large*, *oblate*, *round*, *conical*, or *ovate*.

FIG. 51. A, neck; B, body

FIG. 52. Acute

Pyriform. A term used to describe the shape of a pear when the sides formed by the body and the neck are more or less concave, or hollowed out.

Turbinate. Top-shaped, or having the body nearly round.

INDEX

- Absorption of water by seeds, 12
 Alcoholic grafting-wax, 110
 Ammoniacal solution of copper carbonate, 146
Amygdalus persica, 60, 62, 64
Amygdalus platycarpa, 64
 Annular budding, 104
 Apple, buds of, 66; pruning of, 80; systematic study of, 154-173
 Arsenate of lead, 150
- Bark grafting, 118, 122
 Beeswax and resin grafting-wax, 108
 Begonia, 36
 Bigarreau, 54
 Blackberry, 38
 Brambles, pruning of, 90
 Bridge grafting, 126
 Bryophyllum, 36
 Budding, shield, 94; prong, or twig, 96; plate, 98; H-, 100; flute, 102; ring, or annular, 104; chip, 104
 Buds: *Prunus domestica*, 48; *Prunus americana*, 50; *Prunus triflora*, 50; *Prunus hortulana*, 52; *Prunus munsoniana*, 52; *Prunus avium*, 54, 56, 58; *Prunus cerasus*, 54, 58; *Amygdalus persica*, 60, 62, 64; *Amygdalus platycarpa*, 64; *Malus sylvestris*, 66; *Pyrus communis*, 68; *Pyrus sinensis*, 68; *Cydonia vulgaris*, 70; *Vitis labrusca*, 72; *Vitis vinifera*, 74
 Bulbil, 28
 Bulbs, 28
 Bush fruits, pruning of, 92
- Cactus, 32
 Carnation, 32
 Chemicals, effect on germination of seeds, 20
 Cherry, buds of, 54, 56, 58; pruning of, 84
 Chip budding, 104
 Chrysanthemum, 32
 Cleft grafting, 118, 120
 Coleus, 32
 Commercial fertilizers, effect on seeds, 24
 Corms, 30
 Crocus, 30
 Currant, 40, 42
 Cuttings, soft or green wood, 32, 34; leaf, 36; root, 38; hardwood, or dormant, 40; one-eye, two-eye, three-eye, 44; mallet, 44, 46; heel, 44, 46
- Dicotyledons, 18
 Dinner-plate seed-tester, 2
 Dormant cuttings, 40
 Double-whip grafting, 116
 Drought, effect on germinating seeds, 18
- Flute budding, 102
 Fungicides: effect on seeds, 26; Bordeaux mixture, 132; lime-sulphur, 138, 140, 142; self-boiled lime-sulphur, 144; ammoniacal solution of copper carbonate, 146
- Geneva seed-tester, 2, 4, 8
 Geranium, 32
 Germinating chamber, 2
 Germinating cup, 2
 Germination, hastened by soaking, 16
 Gladiolus, 30
 Gloxinia, 36
 Gooseberry, 40, 42
 Grafting, splice, 112; whip, or tongue, 114; double-whip, 116; cleft, 118, 120; bark, 118, 122; bridge, 126; side, 128; veneer, 130
 Grafting-cord, 108
 Grafting-wax, 108, 110
 Grape, cuttings of, 44, 46; buds of, 72, 74; pruning of, 88; systematic study of, 174-185
 Green wood cuttings, 32, 34
- Hardwood cuttings, 40, 42, 44
 H-budding, 100
 Heel cutting, 44, 46
 Horse radish, 38
 Hyacinth, 28
 Hydrangea, 34
 Hypothesis, 20, 21
- Insecticides: Paris green, 148; arsenate of lead, 150; lime-sulphur, 138, 140, 142; self-boiled lime-sulphur, 144; kerosene emulsion, 152
- Kerosene emulsion, 152
- Leaf cuttings, 36
 Lily, 28
 Lime-sulphur, 138; self-boiled, 144
- Mallet cutting, 44, 46
 Monocotyledons, 18
 Morello, 58

LABORATORY MANUAL OF HORTICULTURE

- Oleander, 32
- One-eye cutting, 44
- Onion, 28
- Paris green, 148
- Peach, buds of, 60, 62, 64; pruning of, 86; systematic study of, 186-197
- Pear, buds of, 68; pruning of, 82; systematic study of, 198-209
- Pitch wax, 110
- Plate budding, 98
- Plum, buds of, 48, 50, 52; pruning of, 84; systematic study of, 210, 221
- Pomaceous fruits, 66
- Prong budding, 96
- Quince, buds, 70
- Ring budding, 104
- Root cuttings, 38
- Rose, 32
- Scion, 122
- Seed vitality limited by age, 10
- Seed-testers, 2
- Seed-testing, 6
- Seeds, amount of water absorbed by, 12; determining volume of, 12; germination of, 16; of dicotyledons, 18; effect of drought on, 18; of monocotyledons, 18; treated with chemicals, 20; effect of commercial fertilizers on, 24
- Shield budding, 94
- Side grafting, 128
- Soaking of seeds, 16
- Soft wood cuttings, 32
- Splice grafting, 112
- Sprouting-chamber, 6
- Systematic study, of apple, 154-173; of grape, 174-185; of peach, 186-197; of pear, 198-209; of plum, 210-221
- Three-eye cutting, 44
- Tile germinator, 2, 4
- Tongue grafting, 114
- Tulip, 28
- Twig budding, 96
- Two-eye cutting, 44
- Veneer grafting, 130
- Vine fruits, 72
- Vitis labrusca*, 72
- Vitis vinifera*, 74
- Volume of seeds, 12
- Whip grafting, 114

UNIVERSITY OF CALIFORNIA LIBRARY

This book is DUE on the last date stamped below.

Fine schedule: 25 cents on first day overdue
50 cents on fourth day overdue
One dollar on seventh day overdue.

OCT 20 1947

LD 21-100m-12,'46(A2012s16)4120

YC 62059

445125

SB 355

H7

UNIVERSITY OF CALIFORNIA LIBRARY