IMAGE EVALUATION TEST TARGET (MT-3)

**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1981

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured covers/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- ☐ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☐ Bound with other material/
Relié avec d'autres documents
- ☒ Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- ☐ Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- ☐ Additional comments:/
Commentaires supplémentaires:

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☐ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☐ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscuries par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X	32X
☐	☐	☐	☐	☐	☐	☒
12X	16X	20X	24X	28X		

med here has been reproduced thanks
osity of:

National Library of Canada

appearing here are the best quality
nsidering the condition and legibility
nal copy and in keeping with the
tract specifications.

ies in printed paper covers are filmed
with the front cover and ending on
e with a printed or illustrated impres-
back cover when appropriate. All
al copias are filmed beginning on the
with a printed or illustrated impres-
ading on the last page with a printed
d impression.

ordred frame on each microfiche
n the symbol $\rightarrow$ (meaning "CON-
or the symbol ∇ (meaning "END"),
applies.

s, charts, etc., may be filmed at
duction ratios. Those too large to be
uded in one exposure are filmed
n the upper left hand corner, left to
p to bottom, as many frames as
ne following diagrams illustrate the

1	2	3
---	---	---

1	2	3
4	5	6

L'exemplaire filmé fut reproduit grâce à la
générosité de:

Bibliothèque nationale du Canada

Les images suivantes ont été reproduites avec le
plus grand soin, compte tenu de la condition et
de la netteté de l'exemplaire filmé, et en
conformité avec les conditions du contrat de
filmage.

Les exemplaires originaux dont la couverture en
papier est imprimée sont filmés en commençant
par le premier plat et en terminant soit par la
dernière page qui comporte une empreinte
d'impression ou d'illustration, soit par le second
plat, selon le cas. Tous les autres exemplaires
originaux sont filmés en commençant par la
première page qui comporte une empreinte
d'impression ou d'illustration et en terminant par
la dernière page qui comporte une telle
empreinte.

Un des symboles suivants apparaîtra sur la
dernière image de chaque microfiche, selon le
cas: le symbole $\rightarrow$ signifie "A SUIVRE", le
symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être
filmés à des taux de réduction différents.
Lorsque le document est trop grand pour être
reproduit en un seul cliché, il est filmé à partir
de l'angle supérieur gauche, de gauche à droite,
et de haut en bas, en prenant le nombre
d'images nécessaire. Les diagrammes suivants
illustrent la méthode.

1
2
3

21" 645
/

OUTLINES
OF
CLASSIFICATION OF PLANTS,

BY
D. P. PENHALLOW, B.Sc., F.R.S.C.,
Professor of Botany, McGill University,

MONTREAL:
E M RENOUF, Publisher.
1895.

Entered according to Act of Parliament of Canada in the Year one thousand
eight hundred and ninety five by D. P. PENHALLOW, in the Office of the Minister
of Agriculture.

OUTLINES OF CLASSIFICATION.

Synopsis.

Branch I.—Protophyta.

CLASS 1.—MYXOGASTRES, (Slime Moulds.)

Orders—Peritrichae.
Columelliferae.
Lithodermiae.
Calotrichae.

CLASS 2.—SCHIZOPHYCEAE, (Fission Fungi)

Orders—Myxobacteriaceae.
Cystiphorae.
Nematogenerae.

From the nature of the organisms included in the protophyta, is improbable that many of them will ever be found in the fossil state, although Renault claims to have recognized several species of bacteria in the tissues of carboniferous plants.

Branch II.—Thallophyta.

CLASS 3.—ASCOMYCETES, (Sac Fungi)

Orders—Perisporiaceae.
Tuberaceae.
Pyrenomycetaceae.
Discomycetaceae.
Uredineae.
Ustilagineae.
Sphaeropsidaceae.
Melanconiceae.
Hyphomycetaceae.

Fossil representatives occur in the carboniferous and later formations, chiefly upon leaves and in lignite.

CLASS 4.—BASIDIOMYCETES, (Higher Fungi)

Orders—Gasteromycetaceae.
Hymenomycetaceae.

Fossil representatives occur in the carboniferous, *Grilletta macrospora*, *Gasteromyces farinatus*, etc.

CLASS 5.—CHLOROPHYCEAE, (Green Algae.)

Orders—Protozoocoidae.
Congugatae.
Siphonae.
Conferoideae.

Fossil representatives, in the form of siliceous diatom valves, are especially abundant in the Tertiary and Quaternary, often forming extensive deposits known under the name of infusorial earth.

CLASS 6.—PHLOPHYCEAE, (Brown Algae.)

Orders—Phaeosporae.
Dictyotae.
Fucoideae.

The earliest fossil representatives of the brown algae occur in the Upper Silurian whence they pass into the Lower Devonian. They appear in these early formations as plants of great size (*Acmetophyton*), indicating that they must also have flourished at much earlier periods, but the remains of the vegetation which existed prior to the Upper Silurian are now represented only by a carbonaceous residue in the form of graphite. Phaeophyceae are found more or less abundantly in all the later formations, and the genus *Fucus* is well defined in the Cretaceous.

CLASS 7.—COLEOCHLITACEAE.

Order—Coleochaetaceae.

No fossil representatives known.

CLASS 8.—RHODOPHYCEAE.

Order—Florideae.

Fossil representatives from the Silurian upward, and especially abundant in the Cretaceous.

CLASS 9.—CHAROPHYCEAE.

Order—Characeae.

Represented by many species in the lower Cretaceous, in the Tertiary and Quaternary.

Synopsis.—(Continued.)

NOTES.

Branch III.—Bryophyta.

CLASS 10.—HEPATICA.

- Orders*—Jungermanniaceae,
Ricciaceae,
Anthocerotaceae,
Marchantiaceae.

Found only in recent formations—Tertiary and Quaternary.
Larchantia.

CLASS 11.—MUSCINE.

- Orders*—Sphagnaceae,
Andreaeaceae,
Phaeaceae,
Bryaceae.

Fossils found only in the Tertiary and Quaternary. *Gymnomium*, *Sphagnum*, *Hypnum*.

Branch IV.—Pteridophyta.

CLASS 12.—FILICES.

- Orders*—Filices,
Salvinaceae,
Marsiliaceae,
Ophioglossaceae,
Marattiaceae.

Fossil representatives numerous from the Devonian (*Parkia*, *Botryopteris*, *Neuropteris*, *Sphenopteris*, etc.) through the carboniferous to recent formations.

CLASS 13.—EQUISETINE.

- Orders*—Equisetaceae,
Calamitae,
Annulatifer,
Astrophyllitae.

Fossil representatives numerous from the Devonian (*Calamites*, *Astrophyllites*, etc.) and throughout the carboniferous.

CLASS 14.—SPHENOPHYLLACE.

- Order*—Sphenophyllaceae.

Fossil representatives in the Siluro-Cambrian and upward.

CLASS 15.—LYCOPODINE.

- Orders*—Lycopodiaceae,
Psilotaceae,
Selaginellaceae,
Isoetaceae.

Fossil representatives numerous and often very large, from the Upper Silurian (*Psilophyton*, etc.) through the Devonian (*Lepidodendron*) and Carboniferous.

Branch V.—Spermaphyta.

CLASS 16.—GYMNOSPERME.

- Orders*—Cycadaceae,
Coniferae,
Taxaceae,
Gnetaceae.

Fossil representatives very numerous from the Devonian (*Dadoxylon*, *Conioides*, etc.) and through the Carboniferous and more recent formations.

CLASS 17.—ANGIOSPERME.

- Sub-Class 1.*—Monocotyledons.
Sub-Class 2.—Dicotyledons.

Orders numerous.

Fossil representatives are very numerous as leaves, lignite, &c., from the Permian upward, becoming more numerous in recent formations.

BRANCH I.—Protophyta.

NOTES.

Unicellular Plants often forming more or less extensive gelatinous plasmodia by coalescence.

CLASS 1.—MYXOGASTRES,
(Slime Moulds.)

Orders.

1. Peritrichen.	2. Columellifera.
3. Lathodermene.	4. Calotrichen.

Asexual reproduction predominant.

Saprophytic plants devoid of a cell wall except in the spores. Reproduction through amoeboid-swarm spores, asexual by simple fission, or sexual (?) (Goebel) by conjugation and the formation of plasmodia, from which asexual spores arise.

CLASS II.—SCHIZOPHYCEAE.
(Fission Fungi.)

Index

1. Myxobacteriacea, 2. Cystiphora,
3. Nematogenceae

Asexual reproduction predominant.

Parasitic or saprophytic plants sometimes forming gelatinous plasmodia. Reproduction wholly asexual by simple division of a mother cell, more rarely by endogenous spore formation.

place of death.

BRANCH II.—Thallophyta.

NOTES.

Unicellular or multicellular plants, the latter often with a stemlike habit of growth.

Sexual generation predominant.

III. ASCOMYCETES. (Sac Fungi.)	IV. BASIDIOMYCETES. (Higher Fungi)	V. CHLOROPHYCEAE. (Green Algae)	VI. PHAEOPHYCEAE. (Brown Algae)	VII. COLLOIDIUM.
<i>Orders.</i>	<i>Orders.</i>	<i>Orders.</i>	<i>Orders.</i>	<i>Order.</i>
1. Perisporiaceae. 2. Tuberoideae. 3. Pyrenomycetaceae. 4. Discomycetaceae. 5. Uredineae. 6. Ustilagineae. 7. Sphaeropsidaceae. 8. Melanconiaceae. 9. Hyphomycetaceae.	1. Gasteromycetaceae. 2. Hymenomycetaceae.	1. Protococcoidae. 2. Conjugatae. 3. Siphonae. 4. Confervoidae.	1. Phaeosporaceae. 2. Dictyotaceae. 3. Fucoideae.	1. Coleochaetaceae.
Asexual reproduction commonly predominant.	Asexual reproduction predominant.	Sexual reproduction often by conjugation with the formation of Zygospores. Asexual reproduction predominant.	Reproduction wholly sexual in Fucoaceae, asexual in Phaeosporaceae and Dictyotaceae.	Asexual reproduction by swarm-spores often predominant.
1. Perfect plant ♀. 2. Antheridia ♂.		1. Perfect plant ♀. 2. Antheridia ♂. 3. Spermatozoids.	1. Perfect plant ♀ or ♂. 2. Antheridia ♂. 3. Spermatozoids.	1. Perfect plant ♀. 2. Antheridia ♂. 3. Spermatozoids.
1. Perfect plant ♀. II. Archicarpus ♂.		1. Perfect plant ♀. II. Oogonium.	1. Perfect plant ♀ or ♂. II. Oogonia.	1. Perfect plant ♀. II. Oogonium.
	No clearly defined sexual reproduction known.	III. Oospere (Ovum).	III. Oospere (Ova).	III. Oospere (Ovum).
3 x III. Ascus.		4 x IV. Oospore.	4 x IV. Oospore.	4 x IV. Oospore.
4 x IV. Spores.		5 x V. Swarm-spores.		5 x V. Swarm-spores.

Alternation of generations often well defined.

BRANCH III—Bryophyta.

Cellular plants with roots, epidermis and stomata,
often with leafy stems,
Sexual generation predominant.

Thallophyta.—(Continued.)

II. RHODOPHYCEÆ. (Red Seaweeds.)	IX. CHAROPHYCEÆ	X. HEPATICA. (Liverworts)	XI. MUSCINEÆ. (Mosses)
<i>Order.</i> Florideæ.	<i>Order.</i> 1. Characeæ.	<i>Order.</i> 1. Jungermanniaceæ. 2. Ricciaceæ. 3. Anthocerotaceæ. 4. Marchantiaceæ.	<i>Order.</i> 1. Sphagnaceæ. 2. Andreaceæ. 3. Phascaceæ. 4. Bryaceæ.
Asexual reproduction often predominant by means of tetrahedria.	Asexual reproduction often predominant by means of bulbils, or tubers, or special vegetative branches.	Asexual reproduction often predominant, by means of simple separation of the vegetative axis, by adventitious shoots or by gemmæ.	Asexual reproduction predominant, by the development of a protonema, more rarely through gemmæ.
Perfect plant ♂ or ♀ Spermatogonia. Spermatia ♀	1. Perfect plant ♂ 2. Antheridium ♀ 3. Spermatozoids motile	1. Perfect plant ♂ or ♀ 2. Antheridia ♀ 3. Spermatozoids motile	1. Perfect plant ♂ or ♀ 2. Antheridia ♀ 3. Spermatozoids motile.
Perfect plant ♂ or ♀ Procarp. Trichogyne.	1. Perfect plant ♂ II. Oogonium. III. Oosphere.	1. Perfect plant ♂ or ♀ II. Archegonium. III. Oosphere (Ovum).	1. Perfect plant ♂ or ♀ II. Archegonium. III. Oosphere (Ovum).
III. Carpospores. IV. Protonema.	1 x IV. Oospore. 5 x V. Pro-embryo.	4 x IV. Oospore. 5 x V. Embryo rudimentary. 6 x VI. Sporogonium parasitic upon the sexual generation. 7 x VII. Spores with elaters. 8 x VIII. Protonema.	4 x IV. Oospore. 5 x V. Embryo rudimentary. 6 x VI. Sporogonium parasitic upon the sexual generation. 7 x VII. Spores 8 x VIII. Protonema.

Oöphyte Generation.

(sexual Phase.)

Alternation of generations well defined.

Sporophyte Generation.

(Asexual Phase)

Oophyte Generation.
 (Sexual Phase.)
 Sporophyte Generation
 (Asexual Phase.)
 Alternation of generations often well defined.

XII.—FILICES L.
(Ferns.)

XIII.—EQUISETACE.
(Horse-tails)

A. Leptosporangiate.		B. Eusporangiate.		
1. Homosporous.	2. Heterosporous.	3. Homosporous.	Homosporous.	Heterosporous.
<i>Orders.</i>	<i>Orders.</i>	<i>Orders.</i>	<i>Orders.</i>	<i>Orders.</i>
1. Filices. Asexual reproduction wholly subordinate; when present, chiefly by extension and division of the main axis, more rarely by bulbs, or by stoloniferous fronds, or even by apogamy.	1. Salviniaceae. 2. Marsiliaceae. Asexual reproduction wholly subordinate.	1. Ophioglossaceae. 2. Marattiaceae. Asexual reproduction subordinate, by means of adventitious buds on the roots.	1. Equisetaceae. 2. Calamitae (?). Asexual reproduction subordinate by means of tubers.	1. Annulariaceae. 2. Asterophyllitaceae. Represented by sil forms only, in succession as in heterosporous Lycopodium.
1. Prothallus ♂ well formed.	1. Prothallus ♂ well formed.	1. Prothallus ♂ well formed.	1. Prothallus ♂ well developed.	
2. Antheridium ♂	2. Antheridium.	2. Antheridium ♂	2. Antheridium ♂	
3. Spermatozoids motile.	3. Spermatozoids.	3. Spermatozoids motile.	3. Spermatozoids motile.	
1. Prothallus ♂ well formed.	1. Prothallus well formed.	1. Prothallus ♂ well formed.	1. Prothallus ♂ well developed.	
II. Archegonia	II. Archegonia.	II. Archegonia	II. Archegonium	
III. Oosphere (Ovum.)	III. Oosphere (Ovum)	III. Oosphere (Ovum)	III. Oosphere (Ovum)	
4 x IV. Oospore.	4 x IV. Oospore	4 x IV. Oospore.	4 x IV. Oospore.	
5 x V. Embryo rudimentary.	5 x V. Embryo rudimentary.	5 x V. Embryo rudimentary.	5 x V. Embryo rudimentary.	
6 x VI. Normal plant.	6 x VI. Normal plant	6 x VI. Normal plant.	6 x VI. Normal plant	
7 x VII. Sporophyll.	7 x VII. Sporocarp.	7 x VII. Sporophyll.	7 x VII. Scales of ter	
8 x VIII. Sporangia.	8 x VIII. Sporangia	8 x VIII. Sporangia.	8 x VIII. Sporangia.	
9 x IX. Spores.	9 x IX. Spores and ♂	9 x IX. Spores.	9 x IX. Spores with elaters.	

phyta.

inct Vascular System.
(ptogams.)
ordinate.

NOTES.

XIV. SPHENOPHYLLACE.		XV. LYCOPODINE.	
(Club Mosses.)			
Heterosporous.	Homosporous or Heterosporous, the latter fossil only.	Homosporous.	Heterosporous.
Order.	Order.	Order.	Order.
Sphenophyllaceae.	1. Lycopodiaceae	1. Psilotaceae.	1. Selaginellaceae. 2. Isoetaceae.
	Asexual reproduction subordinate, by means of axillary bulbils, or by lateral budding of underground tubers.	Sexual reproduction predominant.	Asexual propagation subordinate, by division of the main axis, rarely by apogamy.
	1. Prothallus ♂ or ♀ rudimentary. 2. Antheridium ♀ 3. Spermatozooids motile.	1. Prothallus ♀ rudimentary. 2. Antheridium ♀ 3. Spermatozooids motile.	1. Prothallus ♀ rudimentary. 2. Antheridium. 3. Spermatozooids motile.
	1. Prothallus ♂ or ♀ rudimentary. II. Archegonium III. Oosphere (Ovum).	1. Prothallus rudimentary. II. Archegonium III. Oosphere (Ovum).	1. Prothallus rudimentary. II. Archegonium. III. Oosphere (Ovum).
	4 x IV. Oospore. 5 x V. Suspensor. 6 x VI. Embryo rudimentary. 7 x VII. Normal plant. 8 x VIII. Sporangiferous leaf. 9 x IX. Sporangium. 10 x X. Spores.	4 x IV. Oospore. 5 x V. Suspensor 6 x VI. Embryo rudimentary. 7 x VII. Normal plant. 8 x VIII. Sporangiferous leaf. 9 x IX. Sporangium. 10 x X. Spores.	4. Oospore. 5. Suspensor. 6. Embryo rudimentary. 7. Normal plant. 8. Sporangiferous leaf. 9. Microsporangia. 10. Microspores. IV. Oospore. V. Suspensor. VI. Embryo rudimentary. VII. Sporangiferous leaf. VIII. Normal plant. IX. Macrosporangia. X. Macrospores.

Represented by fossil forms only. The succession as in the heterosporous Lycopodinae.

BRANCH V.—Spermatophyta.

True seed plants.

Sexual generation very subordinate, in the Dicotyledons becoming almost completely suppressed with respect to the general structure.

NOTES.

XVI.—GYMOSPERMÆ.

Carpels open, seeds naked.

XVII.—ANGIOSPERMÆ.

Carpels forming closed seed vessels.

Heterosporous; prothalli two $\times$ τ .

Heterosporous, the rudimentary prothalli often wanting.

Orders.

1. Cycadaceæ. 2. Coniferae.
3. Taxaceæ. 4. Gnetaeæ.

Orders numerous.

Order: numerous.

Asexual propagation subordinate, rarely wholly replacing the sexual; varied, by means of runners, stolons, offsets, bulbs, suckers, tubers, and rarely by parthenogenesis.

Asexual propagation subordinate; sometimes by tubers.

1. Prothallus rudimentary. τ
2. Antheridium rudimentary. (Pollen tube.)
3. Spermatozooids none. (Protoplasm of the antheridium.)

1. Prothallus not developed.
2. Antheridium rudimentary. (Pollen tube.)
3. Spermatozooids none. (Protoplasm of the antheridium.)

1. Prothallus not developed. τ
2. Antheridium rudimentary. (Pollen tube.)
3. Spermatozooids none. (Protoplasm of the antheridium.)

- I. Prothallus (endosperm), formed before impregnation; parasitic upon the asexual generation. (Seeds albuminous.)
- II. Archegonium well formed.
- III. Oosphere (Ovum.)

- I. Prothallus (endosperm), formed only after impregnation of the ovum; parasitic upon the asexual generation. (Seeds albuminous.)
- II. Archegonium not formed.
- III. Oosphere (Ovum. Germ cell.)

- I. Prothallus (endosperm) formed only after impregnation of the ovum, generally obliterated. (Seeds exalbuminous.)
- II. Archegonium not formed.
- III. Oosphere (Ovum or Germ cell.)

- I. Oospore.
5. Suspensor.
6. Embryo with 2 to several cotyledons.
7. Normal plant.
8. Anthophylla.
9. Microsporangia (Anthers.)
10. Microspores (Pollen.)

- I. Oospore.
5. Suspensor.
6. Embryo with one cotyledon.
7. Normal plant.
8. Stamens.
9. Microsporangia (Anthers.)
10. Microspores (Pollen.)

- I. Oospore.
5. Suspensor.
6. Embryo with two cotyledons.
7. Normal plant.
8. Stamens.
9. Microsporangia (Anthers.)
10. Microspores (Pollen.)

- IV. Oospore.
- V. Suspensor.
- VI. Embryo with 2 to several cotyledons.
- VII. Normal plant.
- VIII. Carpophyllum.
- IX. Macrosporangia (Ovules.)
- X. Macrospore (Embryo sac.)

- IV. Oospore.
- V. Suspensor.
- VI. Embryo with one cotyledon.
- VII. Normal plant.
- VIII. Pistil.
- IX. Macrosporangia (Ovules.)
- X. Macrospore (Embryo sac.)

- IV. Oospore.
- V. Suspensor.
- VI. Embryo with two cotyledons.
- VII. Normal plant.
- VIII. Pistil.
- IX. Macrosporangia (Ovules.)
- X. Macrospore (Embryo sac.)