IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions

Institut canadien de microreproductions historiques

1980

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

☒ Coloured covers/
Couverture de couleur

☐ Covers damaged/
Couverture endommagée

☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée

☐ Cover title missing/
Le titre de couverture manque

☐ Coloured maps/
Cartes géographiques en couleur

☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)

☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur

☐ Bound with other material/
Relié avec d'autres documents

☐ Tight binding may cause shadows or distortion along interior margin/
La reliure serrée peut causer de l'ombre ou de la distortion le long de la marge intérieure

☐ Blank leaves added during restoration may appear within the text. Whenever possible, these have been omitted from filming/
Il se peut que certaines pages blanches ajoutées lors d'une restauration apparaissent dans le texte, mais, lorsque cela était possible, ces pages n'ont pas été filmées.

☐ Additional comments:/
Commentaires supplémentaires:

☐ Coloured pages/
Pages de couleur

☐ Pages damaged/
Pages endommagées

☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées

☐ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées

☐ Pages detached/
Pages détachées

☒ Showthrough/
Transparence

☐ Quality of print varies/
Qualité inégale de l'impression

☐ Includes supplementary material/
Comprend du matériel supplémentaire

☐ Only edition available/
Seule édition disponible

☐ Pages wholly or partially obscured by errata slips, tissues, etc., have been refilmed to ensure the best possible image/
Les pages totalement ou partiellement obscurcies par un feuillet d'errata, une pelure, etc., ont été filmées à nouveau de façon à obtenir la meilleure image possible.

☒ This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
					☒						

The copy filmed here has been reproduced thanks to the generosity of:

Library of the Public
Archives of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

La bibliothèque des Archives
publiques du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

errata
to

la pelure,
on à

CONTRIBUTIONS

TO THE

du ho

NATURAL HISTORY

OF

NOVA SCOTIA.

BY J. MATTHEW JONES, F. L. S.

INSECTA

COLEOPTERA.—PART I.

HALIFAX, NOVA SCOTIA.

PRINTED BY JAMES BOWES & SONS, BEDFORD ROW.

1870.

1870
(83)

8
18

t
c
g
P
g
a
in
h
m
n
A
th
to

in

se
co
no
D

Proc. & Trans. Nova Scotia Inst. Scien.
1870, Halifax, Vol. II, Part 3, pp. 141-55.)

NOVA SCOTIAN COLEOPTERA.

To form a complete list of any particular order of insects inhabiting a country of some extent, requires the united labour of divers entomologists, and that continuous for several years; but in a new, and I may almost say, untouched field * like the province of Nova Scotia presents, the isolated entomologist can look forward to nothing better than a meagre addition from time to time to the small collection he is able to form. It is to be hoped, however, that our Institute is gradually maturing the work it was designed to effect, viz.: the engendering in the minds of our people a love of nature; and that ere a few years have elapsed, some of its members will take up the study of entomology, and render service in the cause of promoting a knowledge of that interesting branch of Natural History.

The services of an entomologist, who can set before the inhabitants of the land in which he lives the various injuries done by certain species, and the direct benefits conferred by others, are of great value; and as there is no country in the world in which the people do not suffer in a greater or lesser degree from damage to growing crops, produce in bulk, clothing materials, and in fact almost every thing capable of being injured by the myriad forms of insect life which everywhere exist, a knowledge of insects, their habits, haunts, and economy, is very desirable, as it may be the means of saving much valuable property, even to the extent of many thousand pounds per annum. Indeed, a writer in a recent American magazine asserts that "*taking one year with another the United States suffer from the depredations of noxious insects to the annual amount of \$300,000,000.*"

Under these circumstances, therefore, it is a matter of some importance to ascertain whether injurious insects are to be found

* I am not aware of any collection ever having been made in this country and scientifically described, save that of Lieut. Redman, who some 40 years ago collected for the British Museum, and in the cabinet of which institution it may now be seen still in an excellent state of preservation. It is chiefly composed of *Diptera*.

abundantly or otherwise in the country we live in; and hence, the necessity of publishing lists of native insects as often as the diligence of collectors will allow of such publications. If for this and no other reason this present incomplete list of the Coleoptera of the province is offered to the Institute I trust it will be received as a small contribution to our monthly papers, and that (D. V.) I may be allowed to continue the subject at some future time.

I must not omit to express the obligation I feel myself under to the Rev. C. J. S. Bethune, of Credit, Ontario (Secretary to the Entomological Society of Canada), who very kindly identified nearly the whole of the species comprised in this list; and to Melsheimer's Catalogue of United States Coleoptera, and Mr. Ritchie's excellent Catalogue of Montreal Coleoptera, published in the 'Canadian Naturalist,' I owe my synonyma.

Fam. CICINDELIDÆ.

Gen. CICINDELA.

1. *Cicindela vulgaris*, Say.
C. obliquata, Dej.
C. Tranquebarica, Herbst.
2. *Cicindela albilabris*, Kirby.
C. longilabris, Say.

Common in the interior of the country, particularly on the highroads about Truro. It is not so common, however, in the neighborhood of the Atlantic coast, where *C. 12 guttata* is more abundant.

This species, which seems to be rare in the Northern United States and Canada West, is of frequent occurrence in the neighborhood of Halifax: indeed, I may almost say more abundant than *C. vulgaris*. There can be no mistake as to the insect, for the lengthened light coloured labrum is peculiar to this species. From the fact of its being, as far as I have observed, only common over the more barren portions of this province, coupled with the knowledge of its also frequenting the barren shores of the Arctic Sea, in lat. 70° N., where it was taken by Sir John Richardson, as also Fort Simpson, on the Mackenzie River; it would appear that the rocky and uncultivated districts of North America, in the vicinity of the sea,

sparsely clothed with vegetation, are more suited to its habits. Of eight specimens now before me, some have the white dots and angular medial marking on the elytra much more distinct than others. It makes its appearance in spring earlier than any of our *Cicindelas*, as I have taken it at the end of April.

3. *Cicindela* 12, *guttata*, Dej.

C. Proteus, Kirby.

This may be considered the most common species in the neighborhood of Halifax, where, in open spots, and more especially "Halifax Common," it may be taken in hundreds. On that portion of this open plain called "Camp Hill," those parts which are destitute of grass, are perfectly riddled with the larval holes of this insect. In damp, foggy, or rainy weather, I have observed it lying partially torpid beneath surface stones.

4. *Cicindela* ———? Of this rare species I have only three specimens, which I have not yet identified.

5. *Cicindela* ———? Another species, rare, and undetermined.

Fam. CARABIDÆ.

Gen. ELAPHIRUS.

Elaphrus ruscarius, Say.

E. Americanus, Dej.

The only specimen I have seen, is one in the possession of Mr. Arthur Silver, by whom it was taken at River Bank, Preston.

Gen. CALOSOMA.

Calosoma calidum, Fabr.

C. lepidum, Lec.

This carab cannot be considered as abundant in the neighborhood of Halifax, although fine specimens may be obtained in the heat of summer on "the Common" hunting amid the scanty herbage for prey. I have never observed it in the forest. It appears to be partial to the colder regions of North America, and specimens have been taken as far north as the Mackenzie and Slave Rivers, between lat. 58° and 65° N.

Gen. CYCHRUS.

Cychnus Lecontei, Dej. Rare. I have only taken one specimen.

Gen. PTEROSTICHUS.

1. *Pterostichus lucublandus*, Lec.

Feronia lucublandus, Say.

Poecilus lucublandus, Kirby.

Common under stones; Halifax common. A pungent odour emanates from it when handled.

2. *P. chalcites*, Lec.

Feronia chalcites, Say.

Poecilus Sayi, Brulle.

Poecilus chalcites, Kirby.

Poecilus micans, Chand.

Very common in gardens during the hottest days of summer.

3. *Pterostichus* ———? A large species; rare.

4. *Pterostichus* ———? Rare.

Gen. PLATYNUS.

1. *Platynus extensicollis*, Lec.

Feronia extensicollis, Say.

Anchomenus extensicollis, Ibid.

A. Lecontei, Dej.

—var. *proximus*, Harris, New. Eng. Farmer.

Common under stones on the shores of Halifax harbour, near tidal marks. This insect must be able to lead a partially amphibious life, for I find the places it frequents perfectly wet with the salt water.

2. *Platynus cupripennis*, Lec.

Feronia cupripennis, Say.

Agonum cupripenné, Dej.

Very common in gardens, where it may be found all day and also in the darkness of night, busily searching for food.

3. *Platynus* ———? This, and two or three other species in my collection, yet undetermined.

Gen. AMARA.

Amara obesa, Say.

Percosia obesa, Lec.

Of this carab I have only obtained one specimen, and consider it rare about Halifax.

Fam. DYTISCIDÆ.

Gen. DYTISCUS.

Dytiscus marginalis, Linn.

Not uncommon in the lakes with which this province abounds. It answers in every particular the characteristics of the European species.

Gen. HYDATICUS.

1. *Hydaticus* —? Species undetermined. It appears early, as I have observed it moving about in a lake on the 4th of April.
2. *Hydaticus* —? Species undetermined.

Fam. GYRINIDÆ.

Gen. GYRINUS.

Gyrinus —? Very common on all our lakes and streams, appearing about the last week of April.

Fam. SILPHIDÆ.

Gen. NECROPHORUS.

1. *Necrophorus Americanus*, Oliv.

N. grandis, Fabr.

N. virginicus, Frölich Natf.

Common in putrid carcasses of animals.

2. *Necrophorus orbicollis*, Say.

N. Hallü, Kirby.

N. 4 maculatus, Dej.

var. *tibialis*, Lec.

Not quite so common as the preceding species, in similar situations.

3. *Necrophorus velutinus*, Fabr.

N. tomentosus, Web.

Not common.

4. *Necrophorus* —? A fine species somewhat larger than *N. orbicollis*; rare.

5. *Necrophorus* —? A small species; rare.

Gen. SILPHA.

Silpha Surinamensis, Fabr.*Necrodes Surinamensis* Oliv, et Herbst.

Common in putrid flesh, with the *Necrophori*. Some specimens are of a light chesnut colour and smaller than the darker ones.

2. *Silpha Americana*, Linn.*Oiceptoma affine*, Kirby.*Necrophila peltata*, Lec.*Scarabeus peltatus*, Cates.By far the most common of our *Silphæ*.3. *Silpha marginalis*, Fab.*Oiceptoma marginata*, Kirby.*O. noveboracensis*, Forst.

Common.

4. *Silpha lapponica*, Linn.*Thanatophilus caudatus*, Say.*T. tuberculata*, Germ.*T. Californica*, Mann.*T. granigera*, Chev.

Rare about Halifax. This species appears to have a wide range in North America, having been taken at Lake Superior, and as far north as the Mackenzie and Slave rivers.

Fam. STAPHYLINIDÆ.

Gen. CREOPHILUS.

Creophilus villosus, Grav.*Staphylinus villosus*, Kirby.*S. fasciatus*, Lap.

Not uncommon in decayed animal, and more particularly, vegetable matter, cattle droppings, &c. This species is also known in the Hudson's Bay Territory.

Gen. LEISTOTROPHIUS.

Leistotrophus cingulatus, Perty.*Staphylinus cingulatus*, Grv.*S. speciosus*, Mann.*S. chrysurus*, Kirby.

Not common.

NOTE.—Two or three other Staphylinæ undetermined.

Fam. HISTERIDÆ.

Gen. HISTER.

1. *Hister* ———? Undetermined; rare.
2. *Hister* ———? Another species; rare.

Fam. DERMESTIDÆ.

Gen. DERMESTES.

Dermestes lardarius, Linn.

Very common. A perfect nuisance to the collector, whose zoological specimens are never safe from its ravages.

Fam. BYRRHIDÆ.

Gen. BYRRHUS.

- Byrrhus* ———? Not determined.
Byrrhus ———? Not determined.

Fam. SCARABÆIDÆ.

Gen. APHODIUS.

1. *Aphodius fossor*, Fabr.
Scarabæus fossor, Linn.
Common in cattle droppings.
2. *Aphodius fimetarius*, Hald.
A. nodifrons, Rand.
Scarabæus fimetarius, Linn.

Very common. It is one of the first insects to appear in spring as soon as the frost disappears, frequenting the highroads, where it alights upon the cattle droppings. It is identical with the British species, being of similar size to those I have taken in North Wales; but smaller than those taken in Dorsetshire.

3. *Aphodius scybalarius*, Fabr.

This European form is not uncommon about Halifax in company with the preceding species. There can be no doubt as to its identity, for I have English specimens of my own collecting for comparison. I do not find it in any of the American catalogues.

Gen. SERICA.

Serica vespertina, Lec.*Melolontha vespertina*, Schon.*Omaloplia vespertina*, Har.*Camptorhina atricapilla*, Kirby.

Very common, It hibernates in the ground, and many specimens may be procured where ploughing is going on late in the autumn.

Gen. DIPLLOTAXIS.

Diplotaxis liberta, Germ.*D. moesta*, Say.*D. tristis*, Kirby.

Not common.

Gen. LACHNOSTERNA.

Lachnosterna fusca, Lec.*Melolontha fusca*, Frohl.*M. quercina*, Kroch.*M. fervens*, Gyll.

Gen. OSMODERMA.

Osmoderma scabra, Dej.*Trichius scabra*, Beauv.*Gymnodus foveata*, Kirby.

Rare ; taken by Mr. Arthur Silver, at River Bank, Preston.

Gen. TRICHIUS.

Trichius piger, Fabr.*T. rotundicollis*, Kirby.*T. Drummondii*, Gor et Perh.

Common ; on the wild rose and other flowering shrubs.

Fam. BUPRESTIDÆ.

Gen. DICERCA.

Dicerca divaricata, Lec.*D. dubia*, Melsh.*Buprestis divaricata*, Say.*Stenuris divaricata*, Kirby.

Not uncommon.

Gen. ANCYLOCHEIRA.

1. *Ancylocheira maculiventris*, Lec.*Buprestis maculiventris*, Say.*B. sexnotata*, Lap.*Anoplis rusticorum*, Kirby.

Not uncommon.

2. *Ancylocheira fasciata*, Dej.*A. sexmaculata*, Herbst.

Rare; taken by Mr. Arthur Silver, at River Bank, Preston.

3. *Ancylocheira* ———? Rare; a very beautiful species about the size of *A. maculiventris*. Elytra grooved longitudinally between five ridges, including suturals. Outer margin deeply indented, cupreous. Suture broadly margined with cupreous, expanding anteriorly. The two central grooves metallic green. Thorax punctured, cupreous, metallic green.

Gen. CHALCOPHORA.

Chalcophora virginica, Drury.*C. virginensis*, Herbst.*Buprestis mariana*, Linn.

Common.

Fam. ELATERIDÆ.

Gen. ELATER.

1. *Elater nigricollis*, Herbst.

Not uncommon.

2. *Elater* ———? A species having two buff patches on outer margin of elytra. Taken under dead spruce bark in Point Pleasant Woods.

Gen. MELANOTUS.

Melanotus scrobicollis, Lec.

Not common.

Gen. CORYMBITES.

Corymbites hieroglyphicus, Say.*Selatosomus hieroglyphicus*, Steph.

Not uncommon; taken by Mr. Arthur Silver, at River Bank, Preston.

2. *Corymbites aeripennis*, Lec.

Elater aeripennis, Kirby.

E. appropinquans, Rand.

3. *Corymbites* ———? Species undetermined.

NOTE.—Four or five other species of small size, undetermined.

Gen. ASAPHES.

Asaphes memnonius, Herbst.

Not uncommon.

Fam. LAMPYRIDÆ.

Gen. PHOTINUS.

Photinus corruscus, Lac.

Lampyris corrusca, Linn.

Ellychnia latipennis, Motsch.

E. corrusca. White. (Brit. Mus. Cat.)

Pyractomena fenestralis, Melsh.

Very common. This species is widely distributed over the northern parts of America, and is found as far west as Lake Winnipeg.

Gen. PHOTURIS.

1. *Photuris Pennsylvanica*, Lec.

Lampyris versicolor, Fabr.

L. Pennsylvanica, De Geer.

Our common fire fly. The light emitted is sufficient to enable the collector to read the lines of his note book when the insect is held close over it. It generally appears about the end of June, but I have seen it much earlier. It seems to delight in dark nights with a close humid atmosphere.

2. *Photuris* ———? A small species, undetermined.

Fam. TENEBRIONIDÆ.

Gen. UPIS.

Upis ceramboides, Fabr.

U. reticulata, Say.

Tenebrio ceramboides, Linn.

Not common. This species extends as far north as lat. 63°, near the southern confine of the Arctic Circle, and as far west as the Mackenzie River.

Gen. IPHITHINUS.

Iphthinus serratus, Mann.*Nyctobates serratus*, Beit.

Not common.

Gen. TENEBRIO.

Tenebrio obscurus, Fabr.

Not common.

Tenebrio molitor, Linn.

Not common.

Fam. MELOIDÆ.

Gen. MELOE.

Meloe angusticollis, Say.

Halifax Common; abundant about the first week in June, when the sexes pair.

Fam. OEDEMERIDÆ.

Gen. NACERDES.

Nacerdes melanura, Schmdt.*N. notata*, Fabr.*N. analis*, Oliv.*N. apicalis*, Say.*Cantharis melanura*, Linn.

Rare.

Gen. CANTHARIS.

1. *Cantharis* ———? Rare; Taken by Mr. Arthur Silver, at River Bank, Preston.2. *Cantharis* ———? Rare; Taken in the same locality.3. *Cantharis* ———? Rare; Taken in the same locality.

Fam. CURCULIONIDÆ.

Gen. PISSODES.

Pissodes nemorensis, Germ.

Rare. Taken by Mr. Silver, at River Bank, Preston.

Gen. HYLOBIUS.

1. *Hylobius pales*, Herbst.*H. macellus*, Germ.

Very common, frequenting wooden dwellings and out-houses. This beetle in the larva state is probably more destructive to our pine trees than any other in the province. Its ravages in certain parts of the United States are very great, and the observant naturalist Wilson many years ago, endeavoured to befriend the family of woodpeckers, by proving the vast amount of good they did in consuming numberless larvæ of this insect.

1. *Hylobius* ———? Species undetermined.2. *Hylobius* ———? Species undetermined.

Fam. CERAMBYCIDÆ.

Gen. CRIOCEPHALUS.

1. *Criocephalus agrestis*, Kirby.

Common.

2. *Criocephalus* ———? A smaller species undetermined.

Not common.

Gen. CALLIDIUM.

Callidium violaceum, Fabr.*Cerambyx violaceus*, Linn.

Rare. This is another insect destructive to the pine trees, and is common to Europe and America.

Gen. CLYTUS.

1. *Clytus caprea*, Say.*C. elevatus*, Lap.*C. gibbicollis*, Ibid.

Not common.

2. *Clytus* ———? Another species undetermined, rare; taken by my eldest boy W. Salton Jones.

Gen. MONOHAMMUS.

Monohammus titillator, Fabr.

Very common. Numbers of these insects may be obtained in the forest about August, where the Weymouth Pine (*Pinus strobus*)

has been cut down. I have never, however, met with them on any species of *Abies*, although some lay near the pines. From this circumstance I am inclined to believe that the larva lives in our common pine.

2. *Monohammus scutellatus*, Say.

M. resutor, Kirby.

Very common. Also taken on the newly fallen Weymouth Pine. This species has an extensive range in British America, being found as far north as lat 63°, and as far west as the Mackenzie River.

Monohammus ——? A very handsome species, rather larger than *M. titillator*, with elytra and thorax prettily dappled with patches of white and yellow hairs, and the whole shining with a gloss as if varnished. I have only one specimen in my collection taken at Mount Uniacke, and kindly given me by the hospitable owner of that charming retreat, the Rev. Fitzgerald Uniacke.

Gen. SAPERDA.

Superda candida, Fabr.

S. bivittata, Say.

Rare. In the larval state this is the well known apple tree borer of the northern United States, which commits such sad ravages in the orchards. The only specimen I possess was taken at the foot of an apple tree at Ashbourne. In the apple growing districts of Annapolis and Cornwallis it may be more common than it is near Halifax, where these fruit trees are not much cultivated.

Gen. RHAGIUM.

Rhagium lineatum, Kirby.

Stenocorus lineatus, Oliv.

Common.

Gen. TYPOCERUS.

Typocerus fugax, Fabr.

Strangalia fugax, Sch.

Leptura tenuior, Kirby.

Extremely common; frequents the yellow blossoms of the Golden Rod (*Solidago*) and other flowers.

Gen. LEPTURA.

1. *Leptura elegans*, Lec.
Strangalia elegans, Dej.
Rare.
2. *Leptura vittata*, Oliv.
L. abbreviata, Germ.
L. semivittata, Kirby.
Rare.
3. *Leptura auripilis*, Lec.
Not uncommon.
4. *Leptura Canadensis*, Oliv.
L. tenuicornis, Hald.
Common.
5. *Leptura erythroptera*, Kirby.
L. cinnamoptera, Hald.
Rare.
6. *Leptura chrysoma*, Kirby.
Rare.
7. *Leptura* ———? Species undetermined; rare.
8. *Leptura* ———? Small species undetermined; rare.

Fam. CHRYSOMELIDÆ.

Gen. CHRYSOMELA.

1. *Chrysomela Philadelphica*, Linn.
Calligrapha Philadelphica, Chev.
Rare.
2. *Chrysomela polygoni*, Linn.
This European form is very common in the neighbourhood of
Halifax on grass stalks in waste ground.
3. *Chrysomela elegans*, Oliv.
Rare. Taken by Mr. A. Silver at River Bank, Preston. The
vertical yellow stripes on the elytra and the marginal band of the
same colour around the thorax, are sometimes deep orange.
4. *Chrysomela* ———? A species of the size of *C. Philadel-*
phica, undetermined.
5. *Chrysomela* ———? Undetermined; rare.

Fam. COCCINELLIDÆ.

Gen. COCCINELLA.

1. *Coccinella trifasciata*, Linn.

C. perplexa, Muls.

Common.

2. *Coccinella bipunctata*, Linn.

Adalia bioculata, Say.

Extremely abundant.

Gen. HIPPODAMIA.

Hippodamia 13 punctata, Linn.

H. tibialis, Say.

Rare. Taken by Mr. A. Silver at River Bank, Preston. This species has a wide range over British America, being found at Lake Superior, and as far to the north-west as Great Bear Lake in lat 69° N. long 120° W.

Gen. MYSIA.

Mysia 15 punctata, Muls.

Coccinella 15 punctata, Oliv.

Anatis mali, Say.

A. labiculata, Ibid.

I have only taken one specimen of this insect.