IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1981

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- ☒ Coloured covers/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☒ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- ☐ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☐ Bound with other material/
Relié avec d'autres documents
- ☒ Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- ☐ Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- ☐ Additional comments:/
Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☒ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
☐	☐	☐	☒	☐	☐
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:

Library,
Geological Survey of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque,
Commission Géologique du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

rata
o

elure,
à

MEFL
D27

[FROM THE AMERICAN JOURNAL OF SCIENCE, VOL. XX, NOVEMBER, 1880.]

REVISION OF THE LAND SNAILS OF THE
PALEOZOIC ERA, WITH DESCRIPTIONS
OF NEW SPECIES.

By J. W. DAWSON.

[FROM

REVI

[FROM THE AMERICAN JOURNAL OF SCIENCE, VOL. XX, NOVEMBER, 1880.]

REVISION OF THE LAND SNAILS OF THE
PALEOZOIC ERA, WITH DESCRIPTIONS
OF NEW SPECIES.

By J. W. DAWSON.

[FROM

REVI

THE
brian,
tion of
instead
any for
bonifer

[FROM THE AMERICAN JOURNAL OF SCIENCE, VOL. XX, NOVEMBER, 1880.]

REVISION OF THE LAND SNAILS OF THE
PALEOZOIC ERA, WITH DESCRIPTIONS
OF NEW SPECIES.

By J. W. DAWSON.

THE Gasteropods as a class occur as early as the Upper Cambrian, but all the earlier known types are marine. That portion of the group distinguished by the possession of air sacs instead of gills (Pulmonifera) has not hitherto been found in any formation older than the Carboniferous, and only four Carboniferous species have been described. In the present paper

I propose to state some additional facts respecting the species already known, to discuss their affinities, and to describe two additional species, making six in all from the Paleozoic rocks, including one from the Erian or Devonian. For reasons to be mentioned in the sequel, I do not admit the genus *Palæorbis* founded, by some German naturalists, on fossils which I believe to be tubes of Annelids.

It may be useful to premise that of the two leading subdivisions of the group of Pulmonifera, the Operculate and Inoperculate, the first has been traced no farther back than the Eocene. The second, or Inoperculate division, includes some genera that are aquatic and some that are terrestrial. Of the aquatic genera no representatives are known in formations older than the Wealden and Purbeck, and these only in Europe. The terrestrial group or the family of the *Helicidæ*, which, singularly enough, is that which diverges farthest from the ordinary gill-bearing Gasteropods, is the one which has been traced farthest back, and includes the Paleozoic species. It is further remarkable that a very great gap exists in the geological history of this family. No species are known between the Carboniferous and the early Tertiary, though in the intervening formations there are many fresh-water and estuarine deposits in which such remains might be expected to occur. There is perhaps no reason to doubt the continuance of the *Helicidæ* through this long portion of geological time, though it is probable that during the interval the family did not increase much in the number of its species, more especially as it seems certain that it has its culmination in the modern period, when it is represented by very many and large species, which are dispersed over nearly all parts of our continents.

The mode of occurrence of the Paleozoic Pulmonifera in the few localities where they have been found is characteristic. The earliest known species, *Pupa vetusta*, was found by Sir Charles Lyell and the writer, in the material filling the once hollow stem of a *Sigillaria* at the South Joggins in Nova Scotia, and many additional specimens have subsequently been obtained from similar repositories in the same locality, where they are associated with bones of Batrachians and remains of Millipedes. Other specimens, and also the species *Zonites priscus*, have been found in a thin, shaly layer, containing debris of plants and crusts of Cyprids, and which was probably deposited at the outlet of a small stream flowing through the coal-formation forest. The two species found in Illinois occur, according to Bradley, in an underclay or fossil soil which may have been the bed of a pond or estuary, and subsequently became a forest sub-soil. The Erian species occurs in shales charged with remains of land plants, and which must consequently have

receive
in such
pected
Pulmon
should
minous
Cyprid
fossil p

With
present
sembla
belong
to sub
nent a
ble fro
though
small

All the
moder

I sh
of the
known
explor
of the
of exc
anima
ment
the R
three

[Sir C
the Sou
1832 (6
Air-bre
184, 18

De

at th
eter i
adult
ter o
not
teeth
ing,
spac
in w
spec
roun

received abundant drainage from neighboring land. It is only in such deposits that remains of true land-snails can be expected to occur; though, had fresh-water or brackish water Pulmonates abounded in the Carboniferous age, their remains should have occurred in those bituminous and calcareo-bituminous shales which contain such vast quantities of debris of Cyprids, Lamellibranchs and fishes of the period, mixed with fossil plants.

With reference to their affinities, the Paleozoic land snails present no very remarkable peculiarity except their close resemblance to some modern forms. Of the known species, four belong to the genus *Pupa* in its wider sense, and are very near to sub-generic types still represented on the American continent and its islands. One is a small helicoid shell not separable from the modern genus *Zonites*, and the remaining one, though it has been placed in a new genus, is very near to some small American snails of the present day (*Stenotrema*, etc.) All the species are of small size, though not smaller than some modern shells of the same types.

I shall now proceed to give the characters and descriptions of the several species, adding to the account of those previously known, such new facts as have occurred in my more recent explorations and examinations. I should state here that many of the new facts detailed have been obtained in the course of excavations for the extraction of erect trees holding land animals, undertaken with the aid of a grant from the Government fund for aiding original researches, at the disposal of the Royal Society of London, and carried on within the past three years.

1. *Pupa vetusta* Dawson. (Figs. 1 to 4, and 14, *a*, *b*.)

[Sir C. Lyell and Dr. Dawson on Remains of Reptiles and a Land shell from the South Joggins in Nova Scotia, *Journal of Geological Society of London*, vol. ix, 1832 (figured but not named). Dawson's *Acadian Geology*, 1855, p. 160. Dawson's *Air-breathers of the Coal Period*, 1863. *Acadian Geology*, 2d and 3d editions, p. 384, 1868 and 1879.]

Description.—Shell cylindrical, somewhat abruptly conical at the apex, in some specimens tending to diminish in diameter in the later turns or whorls of the shell. Whorls nine in adult shells, slightly convex, in width equal to half the diameter of the shell. Suture impressed. Aperture evenly rounded, not continuous above, rather longer than broad, destitute of teeth; peristome slightly reflected and smooth. Surface shining, marked with longitudinal smooth ridges, separated by spaces a little wider than the ridges; spaces about $\frac{1}{10}$ th inch in width. Shell calcareous, thin, prismatic in structure. Young specimens abruptly conical and helicoid in form. Nucleus round, smooth, the first turn below the nucleus marked with

rows of little pits which gradually pass into the continuous striae. The last whorl of the adult presents irregular lines of growth, instead of the regular microscopic ribs of the middle turns. Mature ovum membranous, or so slightly calcareous that it can be compressed without breaking: the embryo shell sometimes visible within. Length of adult shell rather less than 1 centimeter, breadth in middle 4 millimeters.

Variety tenuistriata.—Along with the ordinary form there are others of similar size and general structure, but with the apex less obtuse and a somewhat greater tendency to diminish in diameter in the later whorls. They have also the microscopic ridges in the shell about half as far apart as those of the ordinary form. This form I was at first disposed to regard as specifically distinct, but there seems to be a gradual transition from one to the other, and the two forms seem to accompany each other throughout the entire range of the species.

State of preservation.—The shells are usually entire, but often somewhat flattened, and cracked or distorted in the process. Many fragments of shells, however, occur with the entire specimens, and some of these have a whitened or bleached appearance like that of modern land shells after having been exposed to the weather. In one layer I found impressions of several flattened shells, the substance of the shell having been altogether removed. Ordinarily the shell remains in such a state as to show its structure, and the more perfect specimens found in the erect trees have a grayish brown color, like that of some modern Pupæ.

The habitat of this species was in forests of the Coal-formation period, composed of *Sigillaria*, *Calamites*, *Lepidophloios* and *Ferns*. The only known locality is the South Joggins, Nova Scotia. At this place the shells have been obtained in considerable numbers, though perfect specimens which can be disengaged from the matrix, are comparatively few. They have been found in erect *Sigillariæ* and also in a bed of shale. The lowest and highest beds in which they occur are separated by 2,000 feet of vertical thickness of strata including no less than thirty-five beds of coal and many underclays supporting erect trees, so that the species must have inhabited this locality for a very long time and must have survived many physical vicissitudes.

The first specimen, which was also the first known Paleozoic land shell, was found by Sir Charles Lyell and the writer in 1851, in breaking up the contents of an erect tree holding reptilian bones. The specimens obtained from this tree having been taken by Sir Charles to Cambridge and submitted to the late Prof. Jeffries Wyman, the shell in question was recognized by him and the late Dr. Gould, of Boston, as a land shell. It

5.

Fig.
× 8;
5, 6,
ture
section
crush

Fig. 1, *Pupa retusta*, magnified 8 times lineally; 2, same, showing the aperture, $\times 8$; 3, same, nuclear whorl, $\times 25$; 4, same, mature egg and embryo shell, $\times 25$. 5, 6, *Pupa Bigsbyi*, $\times 8$. 7, *Pupa Vermilionensis*, $\times 8$; 8, same, showing aperture $\times 8$, the small tooth on the columella somewhat exaggerated; 9, same, section of aperture, showing tooth $\times 16$. 10, *Zonites priscus*, $\times 8$; 11, same, crushed specimen, showing aperture $\times 20$.

was subsequently examined by M. Deshayes and Mr. Gwynd is b Jeffries, who concurred in this determination; and its micr about 4 scopic structure was described by the late Prof. Quekett, which t London, as similar to that of modern land shells. The singhickne specimen obtained on this occasion was somewhat crushed and he she did not show the aperture. Hence the hesitation as to its Loga nature, and the delay in naming it, though it was figure Division and described in the paper above cited in 1852. Better specimells o mens showing the aperture were afterward obtained by th In th writer, and it was named and described by him in his "Ains alrea breathers of the Coal Period," in 1863. Prof. Owen, in his gray co 'Palæontology,' subsequently proposed the generic name *DenZonites dropupa*. This I have hesitated to accept, as expressing a few v generic distinction not warranted by the facts; but shouldaceous the shell be considered to require a generic or sub-generic disare mos tion, Owen's name should be adopted for it. There seems as frag however, nothing to prevent it from being placed in one of the mud de modern sub-genera of simple-lipped Pupæ. With regard to the stream. form of its aperture, I may explain that some currency has occur it been given to an incorrect representation of it, through an unonly g fortunate accident. In the case of delicate shells like this, though imbedded in a hard matrix, it is of course difficult to work out would the aperture perfectly; and in my published figure in the "Air a sandy breathers," I had to restore somewhat the broken specimens a sandy in my possession. This restoration, specimens subsequently or in a found have shown to be very exact. Nevertheless it was Except ries are criticised by some English conchologists, and when Sir Charles Lyell was about to publish his Student's Manual, he asked me that m to give him one of my best specimens to be figured. This I they co sent with micro-photographs of others. It seems, however, ble tha that the artist or engraver mistook the form of the aperture habit c and gave it an entirely unnatural appearance in the Student's dence Manual. That now given is taken from a photograph of the ology the most perfect and least compressed specimen in my possession. They anima

As already stated, this shell seems closely allied to some modern Pupæ. Perhaps the modern species which approaches most nearly to it in form, markings and size, is *Macrocheilus Gossei* from the West Indies, specimens of which were sent to me some years ago by Mr. Bland, of New York, with the remark that they must be very near to my Carboniferous species. Such edentulous species as *Pupa (Leucochila) fallax* of Eastern America very closely resemble it; and it was regarded by the late Dr. Carpenter as probably a near ally of those species which are placed by some European conchologists in the genus *Pupilla*.

The lowest bed in which *Pupa vetusta* occurs belongs to group VIII of Division 4 of my section of the South Joggins,

and Mr. Gwynd is between Coal 37 and Coal 38 of Logan's section, being about 42 feet below Coal 37. The next horizon, and that in which the shell was first discovered, is 1217 feet of vertical distance higher, in group XV of Division 4 of my section. The shells occur here in erect *Sigillariae*, standing on Coal 15 of Logan's section. The third horizon is in group XXVI of Division 4, about 800 feet higher than the last. Here also the shells occurred in an erect *Sigillaria*.

In the lowest of these three horizons, the shells are found, in a thin bed of concretionary clay of dark gray color, though associated with reddish beds. It contains *Denizonites priscus* as well, though this is very rare, and there are also a few valves of *Cythere* and shells of *Naiadites* as well as carbonaceous fragments, fronds of ferns, *Trigonocarpa*, etc. The *Pupæ* are mostly adult, but many very young shells also occur, as well as fragments of broken shells. The bed is evidently a layer of mud deposited in a pond or creek, or at the mouth of a small stream. In modern swamps, multitudes of fresh-water shells occur in such places, and it is remarkable that in this case the only gasteropods are land shells, and these very plentiful, though only in one bed about an inch in thickness. This would seem to imply an absence of fresh-water Pulmonifera. In the erect *Sigillariae* of group XV, the shells occur either in a sandy matrix, more or less darkened with vegetable matter, or in a carbonaceous mass composed mainly of vegetable debris. Except when crushed or flattened, the shells in these repositories are usually filled with brownish calcite. From this I infer that most of them were alive when imbedded, or at least that they contained the bodies of the animals; and it is not improbable that they sheltered themselves in the hollow trees, as is the habit of many similar animals in modern forests. Their residence in these trees as well as the characters of their embryology are illustrated by the occurrence of their mature ova. They may also have formed part of the food of the reptilian animals whose remains occur with them. In illustration of this I have elsewhere stated that I have found as many as eleven unbroken shells of *Physa heterostrophæ* in the stomach of a modern *Menobanchus*. I think it certain, however, that both the shells and the reptiles occurring in these trees must have been strictly terrestrial in their habits, as they could not have found admission to the erect trees unless the ground had been sufficiently dry to allow several feet of the imbedded hollow trunks to be free from water. In the highest of the three horizons the shells occurred in an erect tree, but without any other fossils, and they had apparently been washed in along with a grayish mud.*

* The discovery of the shells in this tree was made by Albert I. Hill, C.E.

2. *Pupa Bigsbyi* s. n. (Figs. 5 and 6.)

Description.—Shell half the size of *Pupa vetusta*, or between three and four millimeters in length and one and five-tenths millimeters in breadth. Form, long conical. Body whorl about one-third of the entire length, giving the shell a somewhat bulboid form. Whorls five in the largest specimens found. Tumid, suture much impressed. Surface smooth. Aperture apparently oval in form, but not perfectly known, as the body whorl is crushed in all the specimens.

A few specimens, none of them quite perfect, were found in the erect trees of group XV at the Joggins, along with *Pupa vetusta*. They differ from that species in smaller size, different form and absence of sculpture. The specimens do not show whether the aperture was toothed or simple, but it was probably the latter, as the lip is evidently very thin and delicate. From its form it is probable that it belongs to a different subgenus from *P. vetusta*. It is very much more rare than that species in the erect trees, and has not been found elsewhere.

I dedicate it to my venerable and dear friend Dr. Bigsby F.R.S., of London, a pioneer in American geology, and still an indefatigable worker in the science.

3. *Pupa Vermilionensis* Bradley. (Figs. 8 and 9, and 14c.)

[Bradley in Report of Geological Survey of Illinois, vol. iv, p. 254. Id. in Am Journ. Sci., III, vol. iv, p. 87.]

*Description.**—Shell spindle-shaped, tapering to an obtuse apex, covered with microscopic ridges (25 to 30 in a millimeter) parallel to the lines of growth. Aperture oblique, oval. Outer lip thin, slightly reflexed. Columella lip reflexed, thickened; furnished with a single central curved tooth, projecting nearly half way across the aperture. Junction of columella and outer lip somewhat angular and dentiform. In old individuals the columella tooth is often continuous through an entire turn or farther. It is not seen on shells having less than three turns. The last turn forms nearly half the length of the shell. Whorls rounded. Suture impressed. Surface glossy. Color black or gray. Length three and six-tenths millimeters. Width two millimeters. Some individuals are smooth or destitute of the fine microscopic ridges, but whether this is a natural peculiarity or a result of injury to the outer surface, is not certain.

As compared with *Pupa vetusta* this shell is less than half the size, of a less cylindrical form, its whorls more rounded, and its body whorl much larger in proportion. Its sculpture is much finer. The conspicuous tooth in the aperture is of

* Slightly modified from Bradley.

course al
rom its
microsc
ous te
ight col
areous.
As co
sis is v
the aper
to *P. (L*
of *P. ru*
species
This s
1869, in
of Coal
ion Rive
Report o
Pupa ve
in the
cited.

I am
of the G
of Danv

4. *Zonite*

[Quarter
ogy, 2d ed

Descr
tenths
Spire li
flattene
excavat
somewh
with u
growth
probab

This
referre
Pupa
this b
hold c
some
materi
contai
patche

6.) course also a strong mark of distinction. The shell is thin, and from its black color and failure to show structure under the microscope, I infer that it must have been of a horny or cor- and five-tenths of an inch in diameter. The matrix is a somewhat light colored and concretionary, and somewhat hard and calcareous. The specimens found are as follows:

Apertures. As compared with modern American species, *P. Vermilionensis* is very near to several of the smaller forms with teeth in the aperture. In its form and aperture it approaches closely to *P. (Leucochila) corticaria* of Say, or to the immature shell of *P. rupicola*. It has also some resemblance to the western species *P. hordeacea* Gabb, from Arizona.

This shell was discovered by the late Mr. F. H. Bradley in 1869, in concretionary limestone accompanying the underlay of Coal No. 6, Wabash Valley Section, at Pelly's Fort, Vermilion River, Illinois. In the first notice, which appeared in the Report of the Geological Survey of Illinois, it was referred to *Pupa vetusta*, but was subsequently described by Mr. Bradley and Dr. Bigsby in the American Journal of Science, under the name above cited.

I am indebted for specimens of this shell to Mr. John Collett, of the Geological Survey of Indiana, and also to Mr. W. Gurley, of Danville, Illinois.

9, and 14c.)

254. Id. in Am.

to an obtuse

n a millime

oblique, oval

flexed, thick

h, projecting

of columella

In old indi

through an

ng less than

length of the

face glossy.

millimeters.

th or desti

is a natural

is not cer-

s than half

e rounded,

s sculpture

ture is of

4. *Zonites (Conulus) priscus* Carpenter. (Figs. 10 and 11, and 14d.)

[Quarterly Journal of Geological Society of London, Nov. 1867. *Acadian Geology*, 2d edition, 1868, p. 385.]

*Description.**—Shell small, helicoid. Length two and five-tenths millimeters, width two and eight-tenths millimeters. Spire little elevated. Nucleus small. Whorls four, somewhat flattened, with the suture little impressed. Base somewhat excavated with large umbilicus. Aperture oblique, suboval, somewhat regularly rounded. Lip simple. Surface marked with uneven striæ and somewhat more conspicuous ridges of growth. Angle of divergence about 130°. Shell thin and probably horny.

This little shell was discovered in 1866, in the bed already referred to as the lowest of those at the South Joggins in which *Pupa vetusta* has been found. Shortly after I had discovered this bed, being impressed with the probability that it might hold other remains of land animals beside the *Pupa*, I had some excavations made in it, and a considerable quantity of material taken out. I found, however, that the thin layer containing the land shells was not continuous but in limited patches, and was rewarded only by the discovery of a few

* Slightly modified from Carpenter.

specimens of *Zonites priscus* and a small and not determined fragment of bone, in addition to specimens of *Pupa vetusta*.

The specimens found at this time were submitted to the Dr. P. P. Carpenter, by whom the species was named and described. One or two crushed specimens have been subsequently found in the erect trees holding *Pupa vetusta* in growth XV, but the species is extremely rare in comparison. This however have depended on some difference in habitat or mode of life, rendering it less likely to be imbedded in the deposit in process of formation. It is also to be observed that the shell is much more delicate than that of *Pupa vetusta*, and therefore less likely to be preserved.

Fig. 12, *Dawsonella Meeki*, $\times 8$; 13, same, section of aperture, $\times 16$; 14, Markings of surface $\times 100$: (a) *Pupa vetusta*; (b) *Pupa vetusta* var. *tenuistriata*; (c) *Pupa Vermilionensis*; (d) *Zonites priscus*; 15, *Strophites grandeva*, natural size and magnified 8 diameters.

With regard to its affinities, it was compared by Dr. Carpenter with the African species *Paryphanta Caffra* Fer., "on an extremely small scale." Dr. Carpenter also compared it with *Hygromia*, and stated that it might well be ranked under *Pseudosuccinea*.

not determined by *hyalina* of Morse, with the living species *minuscule* and *exigua*. *Pupa vetusta*, I thought it best, however, to place it in the subgenus *Conulus* attributed to the genus *Zonites*, as defined by Messrs. Adams. With regard to the subgeneric name, Dr. Carpenter explained that the subgenus *Conulus* of Fitz, 1833, appears to be synonymous with *Trochiscus* Held, 1837 (non Sby.); also with *Petasia* Beck, 1837; and with *Perforatella* Schlütt.; and according to Adams the habitat or me a subgenus of *Zonites* Montf. (non Leach, Gray). Those who do not care to enter into these subgeneric distinctions, may designate the species as a *Zonites*, or even, speaking loosely, as a *Helix*. There seems nothing in its characters to separate it more than specifically, from many of our smaller helicoid shells with thin shells and simple aperture.

15.

5. *Dawsonella Meeki* Bradley. (Figs. 12 and 13.)

[Report of Geological Survey of Illinois, vol. iv, p. 254. Am. Journ. of Sci., III, n. s. iv, p. 88. Ibid, vol. vii, p. 157.]

Description.—Shell broad, depressed, helicoid. Spire obtuse, consisting of three to three and one-half turns. Length three and two-tenths millimeters, width four millimeters. Surface smooth, but with fine microscopic lines of growth, about fifteen in a millimeter. Aperture oblique, oval, greatly contracted by a broad lamellar expansion of the columella, extending more than half way across, even in small individuals. Outer lip thickened, slightly reflexed. Suture little impressed, imperforate, but last turn slightly excavated in the umbilical region. The shell is usually black in color, and under the microscope shows no distinct structure, from which it may be inferred that it was corneous in texture. It is thicker than the shell of *Zonites priscus*.

This species is found along with *Pupa Vermilionensis*, and was discovered by Bradley, who was, however, at first disposed to refer it to genus *Anomphalus* of Meek; but subsequently, and with good reason, regarded it as distinct and as a land shell. In size and general form it resembles *Zonites priscus*, though expanding less rapidly and with rounder whorls; but it is at once distinguished by its want of the somewhat coarse sculpture of that species, and by the plate which partially covers its aperture. Its nearest modern allies in eastern America would seem to be such shells as *Helix (Triodopsis) palliata*, and *H. (Senotrema) monodon*.

For specimens of this shell I am indebted to the persons above named as having furnished specimens of *Pupa Vermilionensis*.

6. *Strophites grandæva*, s. n. (Fig. 15.)

Description.—Shell cylindrical, with obtuse apex. Whorls four or more. Surface covered with sharp vertical ridges,

* Modified from Bradley.

separated by spaces three times as wide. The body whorl about 4 millimeters in diameter, with about thirteen vertical ridges visible on one side. Length of a specimen probably not quite perfect, about 8 millimeters. The shell, which has disappeared, must have been very thin, and the surface remaining is smooth and shining. In general form, so far as can be ascertained from a very imperfect specimen, this shell must have closely resembled the modern Pupæ of the genus *Strophia* of Albers.

The only specimen known is from the Erian (Devonian) plant-beds of St. John, New Brunswick, which, besides affording great numbers of remains of land plants, have produced the only Erian insects as yet known. It was sent to me by Mr. G. F. Matthew, of St. John, along with specimens of fossil plants, several years ago, but I hesitated to describe it, waiting in hope of additional specimens. As these have not occurred, and I have now carefully examined the whole of the material from these beds to which I have been able to obtain access, I venture to name it as probably the oldest known land shell, the beds in which it is found being either middle or upper Erian.

If a land snail, it is larger in size and probably of higher type than any of those known from the Coal-formation. This would not be wonderful, when we consider the greater variety of surface and the high character of the vegetation, which, as I have elsewhere endeavored to show, distinguished the later Erian age in Northeastern America.

Concluding Remarks.

It may be proper to mention here the alleged Pulmonifera of the genus *Pulvorbis* described by some German naturalists. These I believe to be worm-tubes of the genus *Spirorbis*, and in fact to be nothing else than the common *S. carbonarius* or *S. pusillus* of the Coal-formation. The history of this error may be stated thus. The eminent paleobotanists Germar, Göppert and Geinitz have referred the *Spirorbis*, so common in the Coal-measures to the fungi, under the name *Gyromyces*, and in this they have been followed by other naturalists, though as long ago as 1868 I had shown that this little organism is not only a calcareous shell, attached by one side to vegetable matters and shells of mollusks, but that it has the microscopic structure characteristic of modern shells of this type.* More recently Van Beneden, Cænius and Goldenberg, perceiving that the fossil is really a calcareous shell, but apparently unaware of the observations made in this country by myself and Mr. Lesquereux, have held the *Spirorbis* to be a pulmonate mollusk allied to *Planorbis*, and have supposed that its presence on fossil

* Acadian Geology, 2d edition, p. 205.

plants
attache
found a
R. Eth
has rec
these sh
of artic

If we
Paleozo
howeve
Insects
likely t
announ
are in
John, a
of the
other G
holding
of the
in so fa
rated n
success

It is p
erous w
faunæ o
consider
have be
contain
any larg
occurred
Paleozoi
in those
most co
that we
Carbonif
country
revelatio
Coal-fo
snails an
of reptil
perfect t
of this k
speculate
with anir
or later f
of form
time whi

plants is confirmatory of this view, though the shells are attached by a flattened side to these plants, and are also found attached to shells of bivalves of the genus *Naiadites*. Mr. R. Etheridge, Jr., of the Geological Survey of Great Britain, has recently summed up the evidence as to the true nature of these shells, and has revised and added to the species, in a series of articles in the Geological Magazine of London, vol. viii.

If we exclude the alleged *Palaeorbis* above referred to, all the Paleozoic Pulmonifera hitherto found are American. Since, however, in the Carboniferous age, Batrachians, Arachnidans, Insects and Millipedes occur on both continents, it is not unlikely that ere long European species of land snails will be announced. The species hitherto found in Eastern America, are in every way strangely isolated. In the plant-beds of St. John, about 9,000 feet in thickness, and in the Coal-formation of the South Joggins, more than 7,000 feet in thickness, no other Gasteropods occur, nor, I believe, do any occur in the beds holding land snails in Illinois. Nor, as already stated, are any of the aquatic Pulmonifera known in the Paleozoic. Thus, in so far as at present known, these Paleozoic snails are separated not only from any predecessors, if there were any, or successors, but from any contemporary animals allied to them.

It is probable that the land snails of the Erian and Carboniferous were neither numerous nor important members of the faunæ of those periods. Had other species existed in any considerable numbers, there is no reason why they should not have been found in the erect trees, or in those shales which contain land plants. More especially would the discovery of any larger species, had they existed, been likely to have occurred. Further, what we know of the vegetation of the Paleozoic Period would lead us to infer that it did not abound in those succulent and nutritious leaves and fruits which are most congenial to land snails. It is to be observed, however, that we know little as yet of the upland life of the Erian or Carboniferous. The animal life of the drier parts of the low country is indeed as yet very little known; and but for the revelations in this respect of the erect trees in one bed in the Coal-formation of Nova Scotia, our knowledge of the land snails and Millipedes, and also of an eminently terrestrial group of reptiles, the *Microsauria*, would have been much more imperfect than it is. We may hope for still further revelations of this kind, and in the meantime, it would be premature to speculate as to the affinities of our little group of land snails with animals either their contemporaries or belonging to earlier or later formations, except to note the fact of the little change of form or structure in this type of life in that vast interval of time which separates the Erian Period from the present day.