F7
7295
M46
C.2
HARVARD UNIVERSITY HERBARIUM.

MH-G

Bought

LIBRARY OF THE GRAY HERBARIUM
HARVARD UNIVERSITY

FLORA BARBADENSIS.

To Charles Mackenzie Esq
from his sincere Friend.
The Author.

FLORA HARRADEZIS:

A CATALOGUE

PLANTARUM

INDICUM NATURALIS ET CULTIVATAE

HARRADON

A GEOLOGICAL DESCRIPTION

OF THE

JAMES DOUGLAS MAYCOCK M.A. F.R.S.

A BIRTHDAY PRESENT TO THE SOCIETY OF FRIENDS

THE SOCIETY OF FRIENDS OF THE ARTS AND SCIENCES
HAS THE HONOUR TO ANNOUNCE THAT THE
FOLLOWING WORKS, BY JAMES DOUGLAS MAYCOCK, M.A. F.R.S.
AND WILLIAM DOUGLAS, ESQ. F.R.S. &c. &c.
ARE NOW IN THE PRESS.

Vol. I.

LONDON

PUBLISHED BY JAMES DOUGLAS, 10, PATERNOSTER ROW.

1830.

FLORA BARBADENSIS:

A CATALOGUE

OF

PLANTS,

INDIGENOUS, NATURALIZED, AND CULTIVATED,

IN

BARBADOS.

TO WHICH IS PREFIXED,

A GEOLOGICAL DESCRIPTION

OF THE

ISLAND.

BY

JAMES DOTTIN MAYCOCK, M.D. F.L.S.

A MEMBER OF HIS MAJESTY'S COUNCIL IN BARBADOS.

*"Est enim animorum ingeniorumque naturale quoddam quasi pabulum
consideratio contemplatioque naturæ. Erigimur; latiores fieri videmur;
humana despiciamus: cogitantesque supera atque cælestia, hæc nostra,
ut exigua et minima contemnimus."*

Cic. Acad.

LONDON:

PUBLISHED BY JAMES RIDGWAY, 169, PICCADILLY.

1830.

PRINTED BY JOHN AND JAMES KEENE,
KINGSMEAD STREET, BATH.

PREFACE.

HAVING devoted much of my leisure, during a long professional residence in Barbados, to preparing a Flora of that island, which will, in a great measure, answer for the whole Caribbean Archipelago, I am induced to lay it before the public, hoping that it may be found useful to botanical students in general, and more especially adapted to the wants of those persons who are resident in our valuable colonies in the West. As, however, there are several works which might seem to supersede the necessity of that I now offer, it may be allowed me to remark, that whatever may be expected from their title-pages, or the abilities of their authors, there is none of them which could be adopted as a Flora of Barbados and the neighbouring islands.

It would be impossible for any one, who is a lover of botany, to contemplate detracting from

favourite pursuits. But since the year 1756, when the Civil and Natural History of Jamaica was published, great alterations have been made in, and abundant matter added to, the science of botany. Swartz, Aublet, Jacquin, Plumeir, and others, have largely increased our knowledge of intertropical plants; nevertheless the works of these authors, whether from being descriptive of plants of other places, including comparatively few of those found in the islands, or from their being confined to a particular class of plants, or to plants little or not at all previously known, are none of them of such a nature as to supercede the necessity or usefulness of the Flora I have endeavoured to furnish.

The Natural History of Barbados, published in 1750, by the Rev. Griffith Hughes, A.M. is adorned with many characteristic and well executed plates. It must be admitted, however, that these are the most valuable part of his book, the descriptions in the text being so loose, imperfect, and inaccurate, and so evidently behind the state of the science at the time they were written, that it would hardly be possible to recognize any of his plants, not represented in a plate, if it were not for the agreement of the title in the Natural History with the name at present in current use.

Mr. Hughes was, in fact, altogether unacquainted with botany, and unequal to the task in which he engaged: his descriptions appear to have been frequently taken from the information of superficial observers, which will account for their inaccuracy as far as they go; and his book, being scarcely more than a list of vernacular names, is of little, or indeed of no value to the student. Mr. Hughes himself seems to have been aware of this, for he observes, "As this treatise, with regard to vegetables, is confined within a narrow sphere, it cannot be supposed to comprise rules for a regular system of botany; nor does it, therefore, require me to descend to every minute circumstance, necessary, and expected only, to be expatiated upon by those who undertake to write botanic systems."*

My object in the present attempt has been to exhibit a correct scientific arrangement of all the plants I have met with in Barbados, according to

* I cannot here omit to notice a manuscript description of many of the plants of Barbados, prepared some years since by the late Dr. Collins. This Gentleman's access to books was very limited, but he was possessed of considerable botanical knowledge; and the manuscript, although left imperfect and uncorrected, bears abundant testimony of the ability and science of the author.

the Linnæan system, and conforming to the most recent and improved alterations and amendments. I have added the vernacular name assigned to each of these plants in the Gardener's Dictionary, the Hortus Kewensis, the Hortus Britannicus, the English Flora, and in some other popular works, as marking the appellation by which they are more familiarly known in Britain; also those in use in Jamaica, as far as I have been able to ascertain them from Browne and Long. I have carefully endeavoured to identify the plants mentioned in the Natural History of Barbados with their proper originals; although, from the looseness and inaccuracy of description already alluded to, it has generally been difficult, and frequently impossible to effect this with any thing like certainty. I have also added the names now currently prevalent in Barbados, when different from those adopted by Mr. Hughes; but many plants are not distinguished by any particular English name, being simply included by the vulgar under the general denomination of wild plants.

Although the Linnæan artificial method has been adopted, I have thought it expedient to mark the order to which each genus belongs in the natural arrangement proposed by Linnæus, and

in the more extended and very erudite system of the younger Jussieu; and without particularly tracing the alterations and improvements, from time to time made by the latter author, in the termination of the names of his orders, and in the disposition of some of his genera, I have endeavoured to keep pace with our advancing knowledge by very frequent references to recent sources of information; and herein, more particularly, to the labours of M. De Candolle, and Mr. Robert Brown. In these authors most ample details of generic distinctions will be found, to which I think it sufficient to direct the attention of the botanical student.

That very considerable difficulties obstruct the formation of a natural arrangement of plants was perceived and explained by Linnæus; and it is certain, that all attempts hitherto made have been clogged with imperfections and inconsistencies. The plan itself of Jussieu is indeed not entirely natural: the formation of his classes, from the cotyledons and the position of the stamens, is beyond doubt artificial; and the classes of De Candolle, on the same principle, are not proposed by that author as natural. The distinction of some orders is, moreover, entirely dependant on definitions, in a great degree arbitrary. Many other orders, however, are so purely natural, that we may rea-

sonably conclude that the natural affinities of plants are real, and not fancied, and that our attempts to arrange them accordingly will attain a high degree of perfection, as we become more familiar with the objects of the vegetable kingdom. It is not, however, to be contemplated, that natural arrangement will ever be rendered so complete as to supercede the artificial plan, founded by Linnæus on the number, situation, proportion, and connection of the stamens and pistils; which, when it has received from the process of science, those few amendments, of which it at present stands more or less in need, will become perfect as an *artificial* system, which is all it professes to be.* The aim, indeed, of natural and artificial arrangement is clearly different; and it is altogether unnecessary, and improper, to consider these methods as opposed; or to urge that the latter separates genera, between which there is a strong relationship and analogy, and involves unnatural assemblages. In answer to this it is sufficient to admit, that the artificial plan, as its distinctive appellation unequivocally explains, does not profess to be a natural one; but is intended to assist the memory,

* See an able defence of the Linnæan plan, by Mr. Roscoe, in the Transactions of the Linnæan Society.

and to afford the means of arranging plants where they may be found with the greatest degree of facility: and it may, therefore be contended that whatever it attains beyond this is so much more than it promises. Natural orders, on the other hand, depend on the more obvious and general habits, characters, and relations of plants; they appear indicated by Nature herself, and in some instances are peculiarly evident and perfect, as in the grasses, the palms, and the ferns, compound flowers, umbelliferous and cruciform plants, &c. &c.

Natural arrangement has been long the favourite of botanists of the French school; but has recently been cultivated with great success in Britain. Although, therefore, for the purposes already stated, the artificial system is pre-eminently useful, natural affinities must be studied by all who wish to become philosophically acquainted with botanical science. The many points of connexion and distinction between the orders cannot be well understood, without greatly facilitating our discrimination of plants; and not unfrequently, in the investigation of unknown plants, a knowledge of such circumstances will at once direct us into the right tract.

The specific character has generally been made to rest on the authority of some standard author;

but in the case of a few plants indigenous in Barbados, I have felt constrained to depart from this rule.* Evanescent varieties have not been noticed, and those which, on account of their greater permanence, it has been thought necessary to introduce, are numbered as they occur in Barbados, without regard to the more extensive enumeration to be found in systematic works. I have been careful to select synonyms from the best authority, and at the same time not to increase the bulk of the volume by unnecessarily multiplying them. A reference to Sloane and Browne has never been intentionally omitted; and I have seldom neglected to quote Plumier, Jacquin, Aublet, and Swartz. The older authors, Dillenius, Plukenet, Ray, Bauhin, Tournefort, &c. have more seldom been introduced. I have, however, too much consulted the gratification of those who may look into his book, not to direct their attention to the splendid specimens of British and foreign art,

* It may be necessary to remind the classical reader that the barbarous words, which too frequently occur in the definitions, are in general use with botanists, and constitute a portion of the technical language of the science. To account for some apparent inconsistencies in orthography, the author would have it understood that it has been his wish to quote *literally* the synonyms admitted.

with which the present age so fruitfully abounds. Thus has the addition of plates to this work been rendered in a great degree unnecessary. They would, indeed, have been highly ornamental; but they must have considerably increased the expence of the volume, without, in an adequate degree, extending its usefulness.

The following observations on the subject of references, from the preface of Sir James E. Smith to his English Flora, have made on my mind their proper impression. "I cannot," says this much lamented botanist, "insist too often on the mischief of copying synonyms from other works. The value of synonyms consists entirely in their collector being answerable for their exactness; by which he enriches his own work with the accumulated treasures of the botanical library, as far as his readers think proper to follow up the pursuit. But if he blindly transcribes them, such surreptitious references are sure to be soon discovered by the frequent errors of the press, respecting pages or figures, which occur in most books. At a single discovery of this kind, all our respect and confidence vanish; for we know not that every thing else of the same nature is not equally fallacious; and an author who has appeared very learned and deeply read, proves a

copyist of the lowest rank, decked out with imposing, but borrowed feathers." With the full force of these truths constantly operating, I have spent much time and labour in consulting the authors quoted; abundant proofs of which are scattered through the following pages. Whatever errors, therefore, I may have fallen into, I confidently trust I shall be acquitted of those of a copyist.

I will not pretend that every plant in Barbados has been described. Some, very many perhaps, from the infrequency of their occurrence in ordinary situations, and from their minuteness, I can readily believe have escaped observation. Several have been purposely omitted, as I had not been able to see and examine them in their flowering and fruiting states, when I was hurried from the island by ill health. Some, which I was led to expect to find, were searched for in vain, and have probably been exterminated by the almost universal cultivation of the soil, or remain only in the most secluded and inaccessible spots. When any doubt has remained on my mind respecting a plant from insufficient opportunity of examining it, which, in more than one instance, has been the case, I have thought it incumbent on me to mark that doubt by a note of interro-

gation, placed after the supposed specific name; that others may be guided to a careful examination, before admitting as certain what I only propose as probable.

The Introductory Essay on the Geological Features of the island, and the map and sections, will probably be interesting to some persons, and I hope may not be deemed altogether foreign to the principal object of the publication.

The addition of the vernacular names is confessedly intended for the benefit of those who are attached to botanical pursuits, but who might find much difficulty and trouble in recognizing individual plants by their generic and specific characters alone. Anxious to promote the study of botany in the West Indies, I have in this way endeavoured to afford that assistance, which has been so amply supplied to the student in Britain.

Botany has not unfrequently been represented as a mere detail of verbal distinctions; but how falsely has, I believe, been experienced by all who have given much consideration to the subject. It should rather be considered as a science, which, while it habituates the mind to close observation and minute arrangement, tends to meliorate the dispositions, and to fill the soul with a grateful sense of the rational delight to

be derived from the boundless charms of nature. Hard, indeed, must be the heart of that man, who can enjoy a benefit without gratitude to the giver. The evil of our mind is, that we possess, and appropriate, but do not enjoy. To a large class of mankind creation hourly displays its perfection in vain: their taste for enjoyment, unimproved by neglect, or vitiated by habit, is confined to objects purely sensual, and raises not the mind above subordinate agents; they are strangers to the contemplation of nature, which is attended with intellectual delight, exhorts the thoughts, purifies the desires, and displays to the mind overwhelmed with admiration the infinite attributes of the Creator of the Universe.

It is not my intention to state, that the beauties of the vegetable kingdom are exhibited for the botanist alone; but that his pursuits afford a fertile source of innocent amusement, remove him, for a time at least, from the cares and contentions of the world, and lead him to scenes of solitude and of peace, where feelings are naturally called forth, far more precious than that knowledge of classification and arrangement, which is supposed to be the sole object of his science. Enjoyment such as this is thankfulness, and thankfulness is praise. From our moral and physical constitution, we

require that the serious business of life should alternate with periods of pleasurable relaxation; but as there is one point, to which all our pursuits, as rational and responsible agents should tend, it is of the greatest consequence to our present and future well-being that our sources of amusement be wisely chosen. "Some studies seem to contract the mind, but such is not the character of natural science, which enlarges the understanding by a perpetual display of the power and wisdom of God, and encourages our hopes by sure testimonies of his goodness. He who feeds the sparrows, and clothes the golden lily of the fields in a splendour beyond that of Solomon himself, invites us, his rational creatures, to confide in his promises of eternal life. The simplest blade of grass, and the grain of corn, to which 'he gives its own body,' are sufficient to convince us that our trust cannot be in vain."*

It is indeed true, that the works of nature, eloquent as they are, do not to every one plead with irresistible force. There have been some persons, intimately familiar with natural science, who, from causes too serious to be here alluded to in a superficial manner, have had their hearts little or not at all affected by the stupendous objects of

* Preface to Smith's English Flora.

their continued research. It is with the works, as it is with the words of God : they are not unfrequently so studied, that the knowledge obtained has nothing of the character of true wisdom, and is in some instances, even turned against the most sacred and immutable truths. Such aberration of cultivated genius, when it occurs, exhibits in the strongest light the imperfection, and insufficiency of unassisted human reason; is a becoming lesson of humiliation to the pride of philosophy, and, could we even confine our views to this perishable world, is well calculated to excite our commiseration. But we must not measure the excellence of nature by the effects it has on the obdurate heart of man. Creation is always the same, not more astonishing as a whole, from the perfection of its harmony, and the immensity of its grandeur, than in its minutest parts, from their variety, beauty, and adaptation. Surely he who is best acquainted with such wonders—he who contemplates in detail the innumerable objects of admiration, which present themselves in a blended mass to the multitude—surely he ought to be most deeply and sensibly affected by them; and such, I am confident, is the general fact as regards the student of natural history. If, indeed, the apothegm, “the proper study of

mankind is man," be literally and exclusively admitted, natural science in general must yield much of its claim to the estimation of the wise; but if the noblest study of mankind be God, then may natural science firmly maintain its ground, independent of the many applications to the necessities, the comforts, and the elegancies of human life, to which it is subservient. By it the book of creation is laid open before us, and in it the indubitable proofs of the Divine Author—in it "his glorious attributes are written in large and legible characters." As relates to botany, the endless number of plants with which the Deity has covered the earth, are, as Linnæus has somewhere piously observed, so many monuments of his wisdom and his power—from effects we instinctively pass to the cause—from the creature to the Creator. It is impossible to commune with nature, and be unmindful of God: and to conclude with the words of the excellent Paley,* "if one train of thinking be more desirable than another, it is that which regards the phenomena of nature, with a constant reference to a supreme, intelligent author. To have made this the ruling, the habitual sentiment of our minds, is to have laid the foundation of

* Natural Theology.

every thing which is religious. The world thenceforth becomes a temple ; and life one continued act of adoration."

INTRODUCTORY ESSAY.

**GEOLOGICAL
MAP OF
BARBADOES.**

GEOLOGICAL DESCRIPTION OF BARBADOS.

*"Obiuerat tumulos immensa licentia ponti,
"Pulsabantque novi montana cacumina fluctus."*
OVID.

It would be impossible by any similitude to illustrate sufficiently the irregular figure of Barbados. It is twenty-one miles in length; and its greatest breadth does not exceed fourteen. It is situate in the thirteenth degree of North latitude, and the fifty-ninth of West longitude.

That portion of the coast, the aspect of which is to the West, and to the South, is generally shelving to the sea, with a flat shallow beach; the South-Eastern, and the Northern coasts are, on the contrary, perpendicularly precipitous from thirty to sixty feet, and the water immediately becomes deep, except in some of the small creeks, where steep sandy beaches occur under the rocky cliffs. The windward, or North-Eastern coast, to the extent of fifteen miles or more, exhibits a mixed character, the low land sinking very gradually under the sea, and the rugged conical hills terminating not in mural precipices, but sloping abruptly

to a flat extended beach. The island is nearly encircled with rocks, many of which are rolled masses at a considerable distance from their original situation. The greater part, however, of this rocky belt consists of the substance of the island, extended under the surface of the water in tables, and rising in reefs, or insulated rocks, at no great distance from the shore.

The surface of this island may properly be distinguished into the flat, and the hilly. The low land occupies the northern, southern, and western parts; and rises by precipitous, broken, acclivities, running parallel to the coast, in terraces of flat, open country to the highest land, situate something to the North of the centre of the island. This progressive rise is, indeed, sometimes interrupted by the occurrence of valleys; one only of which, termed *THE VALLEY*, is deserving particular notice. This tract of low land passes from the windward coast of the *THICKETS* between two elevated ridges, denominated *The Ridge* and *The Cliff*, through the parishes of Saint Philip, and Saint George, to Bridge Town; forming the only general interruption to the regular terraced rise from the sea to the highest land. If the sea were fifty or sixty feet above its present level, Barbados would be divided into two islets, of unequal size, by a narrow strait occupying the site of what is now *THE VALLEY*.

MOUNT HILLABY is the highest land, its altitude being eleven hundred and forty-seven feet above the level of the sea; from which it is distant, in a direct line, about four miles. From this point the high land branches off in steep precipitous ridges, in two directions, northerly and easterly, and southerly and easterly, towards the sea on the windward coast; suddenly diminishing in height as they approach it. These two ridges of high land embrace a country, the appearance of which is altogether different from the flat, open scenery of that which has just been described. This portion of the island is distinguished by the appellations, SCOTLAND, and BELOW THE CLIFF. The hills in this district are numerous; they are lofty, conical, and steep, and project irregularly from the ridges of high land, or rise in small groups from the plain, which is little above the level of the sea. The deep valleys intersecting the hills are covered with the most luxuriant vegetation, the hills themselves appearing naked and barren, or richly clothed with timber. The scenery is every where wild, irregular, and picturesque, and displays in miniature all the beauties of a mountainous country.

Such is the striking dissimilarity in the general appearance of the two districts. Attentive observation points out an essential difference in the immediate substratum of the soil; that of the flat country being entirely calcareous, the soil of the

hilly district resting almost exclusively on mineral substances belonging to the clay genus.

Upon examining the structure of the calcareous formation, we find it to consist of the spoils of zoophites; of which several species of madreporæ, milleporæ, corallinæ, and alcyoniæ, are strikingly evident. These are cemented together by carbonate of lime, containing an abundance, and great variety, of shells. The cement may be said to vary from marl, more or less indurated, to a hard compact limestone, with conchoidal fracture, and translucency on the edges. In some places the organic remains constitute the principal, in all a very considerable portion of this formation; and although these remains are intimately blended in the common structure, they appear to be arranged in families; in some situations the madreporæ, in others the alcyoniæ, being most conspicuous. This structure of the calcareous formation prevails throughout its whole extent, even at its highest elevation, which reaches nearly eleven hundred and fifty feet; and in some places, as in the neighbourhood of Hackleton's Cliff, at Ape's Hill Gully, and in many other similar situations, mural precipices of this coralline aggregate are exposed, between one and two hundred feet in height. Upon this coralline mass is occasionally found beds of white shelly sandstone, similar to that of Guadaloupe, in which the reliquiæ of human

bodies have recently been discovered. This sandstone is quarried for the purposes of building, and, being sufficiently porous, is employed for the filtration of water. It sometimes seems disposed to assume the slaty structure, and when the beds are of considerable thickness they simulate the appearance of stratification. Calcareous spar also, and calc sinter, occur abundantly; and I have seen small specimens of white granular limestone. These are found attached to the common calcareous rock, and, like the spar, have been deposited in accidental cavities at a comparatively recent period.

The whole of the calcareous portion of the island presents numerous rents and fissures; the smaller are filled with chrySTALLIZED and other modifications of carbonate of lime; the larger remain open, and are the deep precipitous ravines, or gullies, which are so very frequent in the higher parts of this district, and which become during the rainy season the conducting channels of temporary torrents. This calcareous formation is extremely cavernous; dislocations, and sinking of the surface, occasionally take place at the present time, and from general appearances, we must conclude that they happened very frequently, and to a considerable extent, at former periods. It is to this cause that the island is plentifully supplied with fissures, through which the water lodged on the surface is drawn off, and conducted to the ocean by means of subter-

anean channels; and from this cause, together with the breaking away of the face of the ridges, has arisen the precipitous mural cliffs, so common on the coast, and in the interior of the calcareous or coralline district.

The hilly portion, or that part of the island which has been denominated *Scotland*, and *Below the Cliff*, is principally composed of mineral substances belonging to the clay genus; particularly loam, potter's clay, and slate clay. There is also found here a fine-grained friable brown sandstone, which is for the most part micaceous; and beds of very loosely cohering black sandstone: also Gypsum, in fragments, crystals, and distinct concretions, from a very small size to such as weigh several ounces, disseminated through beds of sandy clay: and there occur, imbedded in the same, or scattered on the surface, fragments of ferruginous conglomerate, and of quartz sandstone, and balls of flint and iron flint: also a variety of other ores of iron, such as clay ironstone, compact black ironstone, compact and ochery brown ironstone, and the like. I have here seen in a hill, from the white appearance of its broken cliffs, termed CHALKEY MOUNT, a bed of porphyritic slate, or clinkstone porphyry, about eighteen inches in thickness, lying between beds of very loosely cohering sandstone, and dipping to the N. E. at an angle of 30° : also a bed, a few inches thick, composed of alternate laminæ of

fuller's earth, and yellow earth. Beds of bituminous shale are frequent, and petroleum or mineral oil, and asphaltum are more or less abundant. The mineral oil exudes from the base of clay hills, and is collected on the surface of natural or artificial reservoirs of water; or it is found penetrating the substance of sandy rocks. In the island it is generally known by the appellation of Green Tar, and in England by that of Barbados Tar.*

* It may be proper to add a few words relative to the general application of this tar in Barbados. It is principally there used instead of other oils, for burning, and painting the roofs of out-buildings; and it is freely and beneficially employed as a medicine internally, and externally for horses, and horned cattle. As a remedy against the diseases of man its reputation has greatly fluctuated, which is readily accounted for by the fact, that at different times it has had the character of being efficacious in the cure of such diseases as cancer, pulmonary consumption, the lepra Arabum or elephantiasis, and tetanus. It is in this latter most dreadful, and frequently fatal affection, that, during a long practice, I have had most opportunity of seeing it employed. It may seem foreign to my present purpose to state that tetanus is decidedly epidemic in Barbados; yet I am so fully convinced that this is the fact, and that a knowledge of it will materially affect the result of surgical operations in the West-Indies, that I will not omit to remark it in this place: the more particularly as I have not seen it mentioned in any books, although I cannot suppose that it has escaped the observation of those practitioners, who have had extensive opportunities of seeing the disease. In the cure of tetanus the mineral oil, externally employed, is at least equal to

The clay, which is plentifully distributed through the hilly district, is not very pure, being generally charged with iron, bituminous, or calcareous matter; but in most places it answers sufficiently well for the manufacture of coarse ware and bricks, and accordingly there are several pot-kilns in the parishes of St. John, St. Joseph, and St. Andrew. Furnaces are here frequently constructed of unburnt bricks, the cement on such occasions being a

the oil of amber. Of its powers, administered internally, I can say nothing decisive; for in an affection so frequently, and so quickly fatal, I have never been induced, for the purpose of an experiment, to trust to a remedy, the powers of which are not fully ascertained; and as the means, with which it was associated, were of the most active kind, and employed with the boldness called for by the danger of the patient, it would be quite problematical to conjecture how much of the benefit was to be attributed to the oil, in those cases which have recovered, and in which it was used. The medicinal qualities of the mineral oil, or Green Tar, are externally stimulant, internally aperient, diaphoretic, and diuretic. It might be employed, with much promise of advantage, in several diseases of a chronic nature, both internally and externally, particularly in conjunction with medicated baths. Its disagreeable appearance has, however, kept it out of general use; but this may, in a great degree, be surmounted by pharmaceutical preparation; and it is certainly desirable that it should be submitted to the test of unprejudiced experiment, that its medicinal virtues may be fully, and accurately ascertained. Its sensible qualities are conspicuous, and it deserves a trial in those cases, to which it seems applicable; the more especially, as some of them do not readily yield to the means at present in general use.

paste of the same kind of clay as that of which the bricks are made. Upon the application of heat, the whole becomes consolidated into one mass; and furnaces of this description, if the bricks be previously well cured by exposure to the air, will last many years, although subjected to the very strong fires employed in the manufacture of sugar.

Masses of the calcareous formation, some of considerable magnitude, are to be seen in SCOTLAND. These are either projections from the high ridges, which have never been covered by the clay formation; or they are rolled fragments, of which there is an endless number. Some of these are at a great distance from their original situation, to which, however, they can generally be traced. I have diligently searched for the remains of animal or vegetable substances, but have never been able to find any in the minerals of this district.

I cannot omit taking notice in this place of an extinguished pseudo-volcanic hill, situate on the windward coast, in one of the estates belonging to the Society for the propagation of Christian knowledge. It is to this day very properly denominated the *Burnt Hill*, and is mentioned by Hughes, as having been accidentally set on fire by a slave, and as having continued to burn for the space of five years. It consists entirely of highly burnt clay, and earth slag; and the neighbourhood abounds in bituminous shale, and mineral oil.

The natural superficial springs of Barbados are not very numerous. The inhabitants of the flat country are supplied with water principally from wells, which are frequently of very considerable depth, sunk entirely through the coralline rock; but running streams are abundant in the hilly district, in which occur several saline, and chalybeate springs. There is also a spring in Scotland, called the *Burning Spring*, which generally attracts the notice of the traveller. This little streamlet rises in a deep sequestered ravine, at the foot of a hill richly clothed with timber; and on its first appearance forms for itself a little basin, in which the water is in a continued state of ebullition, from the passage of inflammable gas through it. This readily inflaming on the application of a lighted taper, gives to the spring its characteristic appellation. The gas does not, indeed, rise in great quantity, but the scenery in the approach to the spot is beautiful and imposing; and one can hardly view it without fancying what might have been its celebrity, and importance, had it been known to a people prone to attach superstitious veneration to unusual phenomena.

The saline springs make their appearance at an inconsiderable height above the level of the sea, through the sides, and very near the base of clay hills, abounding in gypsum; and it is quite evident that the saline beds, over which they flow, and from

which they derive their impregnation, are subjacent to those minerals which appear as the external crust of SCOTLAND. The waters have not been carefully analyzed: in taste, and other qualities, they resemble those of Cheltenham, and they are occasionally employed with effect to answer the same medicinal purposes.

From what has been said, it will appear that the great mass of the island consists of calcareous matter combined with carbonic acid; but that the hilly district is composed principally of argil, or of argil and silex, frequently blended with ferruginous or bituminous matter, and containing fragments of rocks of apparently much older formation. After careful inspection, I am quite satisfied that the argillaceous substances rest on the calcareous or coralline mass, which constitutes the exterior crust of the greater part of the island.

It cannot be doubted that the calcareous formation, which has so frequently been mentioned, has originated in the submarine operations of insects belonging to the order of Zoophytes; and that the various modifications of carbonate of lime, by which the corallines are cemented, have been derived from these substances acted on by water. This island, which was once undeniably under the surface of the ocean, now rises considerably above it, the coralline rock, of which its highest elevation consists, reaching between eleven and twelve hun-

dred feet above the level of the sea; and forming one of the many indubitable proofs, some found at very high elevations, that the dry land of our present habitable world has once been under the surface of the ocean—that the land has been elevated, or the waters have subsided. Although it is interesting to inquire how far one, or both of these causes may have operated in far remote ages, it is entirely foreign to my present object to enter into the wide field of inquiry, to which such phenomena would lead. I would rather call the attention of Geologists to the following points.

The first is the high elevation, which the coralline aggregate attains in Barbados; and the fact, that in many situations precipitous walls are exposed of considerably more than one hundred feet in perpendicular height, entirely composed of coral. How are these circumstances to be reconciled to the position of French, and indeed of English, naturalists, that madrepores, in the formation of coral rocks, do not commence their labours at greater depths than twenty-five or thirty feet below the surface of the sea; resting their habitations on the summit of submarine rocks, bringing them nearer to the surface, but not forming them entirely from the bottom of the ocean? I question not the fact that these polypi commence their operations on previously existing rocky summits; nor do I doubt that the nucleus, on which they have

rested their labours in forming the external crust of the greater part of Barbados, is a lower elevation of the same primary rock, which appears as mountains in some of the other islands. But if the range of action of the madreporæ, &c. be limited to twenty-five or thirty feet, in what manner has the coralline crust been formed of so much greater thickness in Barbados? That it is of greater thickness is quite certain; for the formation extends uninterruptedly for many miles, intersected in every direction by deep ravines or gullies, the precipitous walls of which are exposed to a much greater depth than twenty-five or thirty feet. Whether we assume that the land has been elevated, or that the waters have subsided, this fact appears to me at variance, and indeed irreconcilable, with the position laid down by M. M. Quoi and Gaimard; for if the range of action of the submarine zoophytes be circumscribed, as stated by these naturalists and others, no supposed elevation, no tilt or dislocation of the coralline mass, can in any way account for the phenomena, which in numberless places present themselves to observation in Barbados.

I would next observe, that the argillaceous minerals are constantly superimposed on the calcareous, into which they pass abruptly. Also that the argillaceous minerals are found only on the North-Eastern portion of the island, in a deep hollow,

protected to the West, North-West, and South-West, by high ridges of coralline structure; and that they are every where to be found on the North-eastern coast, extending more or less into the body of the island, according to the depth of the indentations of the coast; and attaining elevations proportioned to the height of the coral ridges against which they rest. To illustrate the latter part of this statement, I will instance only a single example to be found at Skeet's Bay, on the North-East of the THICKETS, where, notwithstanding the land is low, the minerals of Scotland, clay and gypsum, are thrown up against the calcareous cliffs; but immediately round the point, these are wanting, and the South-eastern coast appears exposed precipitous, and undermined by the waters of the ocean.

This disposition of the argillaceous minerals seems to me to indicate that they were deposited in their present situation after the coral rock, on which they lie, had attained its present elevation; and the occurrence of saline springs in this part of the island rather confirms me in this opinion. The argillaceous deposition of Scotland, I therefore think should be referred to the rising of the waters of the ocean, at some period subsequent to the formation of the coral; and should be considered as of diluvial origin; a view of the subject well according with the appearance of petroleum,

and the spiracle of inflammable air. It is also apparent, that these must have been deposited under the influence of a current setting from a North-East point, which, while it furnished the argillaceous and other materials to be accumulated in the protected hollow and similar situations, would wash freely down the inclined surface of the other parts of the island the loose substances, which may antecedently have been collected thereon. It is scarcely necessary to add, that a current setting from the North-East at present exists, and is extensively influential; for being stopped in its progress by the Isthmus of Darien, it is reflected through the Gulf of Mexico, and passing between the shores of Florida and the Bahama bank, it constitutes the gulf stream, which so powerfully affects the navigation of the Atlantic. This current being dependent on the North-easterly trade wind, must have been coeval with the present direction of our terrestrial poles; but for the transportation and deposition of the substances forming Scotland, it would seem necessary that its force should have been increased by causes, which may be presumed to have operated during such a catastrophe as the tumultuous rising of the waters of a deluge.

In what manner the inequalities under the ocean, one eminence of which constitutes the foundation of Barbados, were produced, I do not contemplate in this place inquiring; neither shall I attempt to

decide the question, whether after the coral was formed, the waters subsided or the rock was elevated. The solution of this problem must form a part of a general theory, and is left for those, who, from extensive acquaintance with geological facts, may be sufficiently qualified for the task of generalization.

FLORA BARBADENSIS.

BOOKS QUOTED IN THE FLORA.

- An. du Mus.* Annales du Museum National d'Histoire Naturelle. *Paris.* 1802—
- Aubl.* Histoire des Plantes de la Guiane François, par M. Fusée Aublet. *Lond. et Paris.* 1775.
- Bauh. Hist.* Historia Plantarum Universalis, auctore Johanne Bauhino. *Ebroduni.* 1750—51.
- Bauh. Pin.* Caspari Bauhini Πινάξ Theatri Botanici. *Basilicæ.* 1771.
- Blackw.* Herbarium Blackwellianum. *Norimbergæ.* 1757—65.
- Bot. Cab.* The Botanical Cabinet, by Conrade Loddiges and Sons. *Lond.* 1817—
- Bot. Mag.* Curtis' Botanical Magazine, continued by James Sims, M.D. *Lond.* 1787,—New Series, by Samuel Curtis, F.L.S. and William Hooker, L.L.D. *Lond.* 1827—
- Bot. Reg.* The Botanical Register; the designs by Sydenham Edwards. *Lon.* 1815—

- Bot. Repos.* The Botanist's Repository, by H. Andrews.
- Brown.* Robert Brown; author of Prodrômus Floræ Novæ Hollandiæ, General Remarks on the Botany of Terra Australis, &c. &c. &c.
- Br. Browne.* The Civil and Natural History of Jamaica, by Patrick Browne, M.D. Lond. 1756.
- Catesb. Car.* The Natural History of Carolina, Florida, and the Bahama Isles; by Mark Catesby, F.R.S. Lond. 1743.
- Cav. Diss.* Monadelphici Classis Dissertationes Decem. Auctore Ant. Jos. Cavanilles. *Matriti.* 1790.
- Cav. Ic.* Icones et Descriptiones Plantarum. Auct. Ant. Jos. Cavanilles. *Matriti.* 1798.
- Com. Præl.* Caspari Commelini M. D. Præludia Botanica. *Lugd. Bat.* 1703.
- De Cand.* A. P. De Candolle; author of Théorie Élémentaire de la Botanique. *Par.* 1813. Regni Vegetabilis Systema Naturale. *Par.* 1818. Prodrômus Systematis Regni Vegetabilis. *Par.* 1824. &c. &c. &c.
- Dill. Elth.* Hortus Elthamensis. Auctore Joanne Jac. Dillenio, M.D. Lond. 1732.

- Engl. Bot.* English Botany, by James Sowerby.
Lond. 1790—
- Engl. Fl.* English Flora, by Sir James E. Smith.
Lond.—1829.
- Exot. Bot.* Exotic Botany, by J. E. Smith, M.D.
the figures by James Sowerby,
F. L. S. *Lond.* 1804—
- Gaert. Fruct.* Josephus Gaertner de Fructibus et
Seminibus Plantarum. *Lepsiæ.*
1801—2.
- Ham. Prod.* Gulielmi Hamilton, M. B. Prodro-
mus Plantarum Indiæ Occidentalis.
Lond. 1825.
- Fl. Br.* Flora Brittanica. Auct. Jacobo Ed-
vardo Smith, M. D. *Turici.* 1804.
- Haw. Succ.* Synopsis Plantarum Succulentarum.
Auct. A. H. Haworth, F. L. S.
Lond. 1812.
- Herm. Par.* Pauli Hermanni, M. D. Paradisus
Batavus. *Lugd. Bat.* 1705.
- Hort. Br.* Sweet's Hortus Britannicus. *Lond.*
1827.
- Hort. Gram.* Hortus Gramineus Woburnensis, by
George Sinclair, F. R. S. F. H. S.
Lond. 1824.
- Hort. Kew.* Hortus Kewensis, or a Catalogue of
Plants cultivated in the Royal Bo-
tanic Garden at Kew; by William
Townsend Aiton. *Lond.* 1811.

- Hort. Mal.* Hortus Indicus Malabaricus. Auct.
Henrico Von Rheede. *Amstel.* 1678.
- Hughes.* The Natural History of Barbados;
by the Rev. Griffith Hughes, A. M.
Lond. 1750.
- Jacq. Amer.* Nic. Jos. Jacquin Selectarum Stri-
pium Americanarum Historia.
Vind. 1763.
- Jacq. Austr.* Floræ Austriacæ, &c. Icones Nicolai
Jos. Jacquin. *Viennæ.* 1773—5.
- Jacq. Coll.* Nicolai Jos. Jacquin Collectanea ad
Botanica, &c. Spectantia. *Vind.*
1786—90.
- Jacq. Ic.* Icones Plantarum Rariorum. Nica-
lao Jos. Jacquin. *Vind.* 1781—93.
- Jacq. Obs.* Nicolai Josephi Jacquin Observa-
tionum Botanicarum Partes Qua-
tuor. *Vind.* 1764—71.
- Jacq. Schæn.* Plantarum Rariorum Horti Cæsari
Schœnbrenensis. *Viennæ,* &c. 1797.
- Jacq. Vind.* Hortus Botanicus Vindobonensis curâ
Nicolai Jos. Jacquin. *Vind.* 1770—3.
- Juss.* Antonii Laurentii de Jussieu Genera
Plantarum. *Turici.* 1791.
- Kunth.* Synopsis Plantarum Æquinoctialium.
Auct. Carolo S. Kunth. *Paris.* 1825.
- Lawr. Ros.* A Collection of the Roses from Nature,
by Miss Lawrance.

- Ligon.* A True and exact History of the
Island of Barbados; by Richard
Ligon, gent. *Lond.* 1673.
- Lin.* Caroli Linnæi Genera Plantarum.
Holmiæ. 1764
- Lin. Cliff.* Hortus Cliffortianus. Auct. Carolo
Linnæo. *Amstel.* 1737.
- Lin. Sp. Pl.* Caroli Linnæi Species Plantarum.
Holmiæ. 1762—3.
- Lin. Suppl.* Supplementum Plantarum Carolo à
Linnée. *Brunsv.* 1781.
- Lin. Tr.* Transactions of the Linnæan Society
of London.
- Lin. Zeyl.* Caroli Linnæi Flora Zeylanica. *Hol-
miæ.* 1747.
- Long.* The History of Jamaica. *Lond.* 1774.
- Lour. Cochin.* Flora Cochinchinensis Johanni de
Loureiro. *Ulyss.* 1790.
- Mart. Palm.* Genera et Species Palmarum. C. J.
P. de Martius, M. D. *Monachii.*
1823—26.
- Menth. Br.* Menthæ Britannicæ; by William
Sole. *Bath.* 1798.
- Mich. Amer.* Andreas Michaux. Flora Boreali-
Americana. *Paris.* 1803.
- Miller.* The Gardener's and Botanist's Dic-
tionary, by Phillip Miller, F.R.S.;
new edition by Thomas Martyn,
B.D. F.R.S. *Lond.* 1807.

- Mor. Hist.* Roberti Morison, M. D. Historia Plantarum Universalis Oxeniensis. Oxon. 1680.
- Pluk. Alm.* Almagestum Botanicum. Lond. 1696.
- Pluk. Phyt.* Leonardi Plukenetii Phytographia. Lond. 1691.
- Plum. Amer.* Descriptions des Plantes de l'Amerique; par le R. P. Charles Plumier. Paris. 1693.
- Plum. Fil.* Traité des Fougères de l'Amerique; par le R. P. Charles Plumier. Paris. 1705.
- Plum. Gen.* Nova Plantarum Americanarum Genera. Auct. P. Carolo Plumier. Paris. 1703.
- Plum. Ic.* Caroli Plumier Plantarum Americanarum Fasciculi Decem. Amstel. 1750.
- Raj. Hist.* Historia Plantarum Joanne Rajo. Lond. 1686.
- Retz. Obs.* Andreæ Joannis Retzii Fasciculus Observationum Botanicarum Tertius. Lipsiæ. 1783.
- Rottböll. Gram.* Descriptiones et Icones, &c. Auct. Christophoro Rottböll. Havnæ. 1786.
- Roem. Syst.* Caroli à Linnée Systema Vegetabilium. Auct. Joanne J. Roemer, M. D. et Jos. Aug. Schultes, &c. Stuttgart. 1817—20.

- Roxb. Cor.* Plants of the Coast of Coromandel;
by William Roxburgh. *Lond.* 1798.
- Rumph. Amb.* Georgii Everhard Rumphii Herbarium Amboinense. Ed. J. Burmanno, M.D. *Amstel.* 1741.
- Sl.* A Voyage to the Islands of Madera, Barbados, Neives, St. Christopher, and Jamaica, &c. &c.; by Sir Hans Sloane, bart. *Lond.* 1707—25.
- Spreng. Syst.* Coroli Linnæi Systema Vegetabilium Curante Curtio Sprengel. *Gottingæ.* 1825—8.
- Swtz. Fl.* Olavi Swartz, M.D. Flora Indiæ Occidentalis. *Erlangæ.* 1797—1800—1806.
- Swtz. Obs.* Observationes Botanicae Olavi Swartz, M.D. *Erlangæ.* 1791.
- Swtz. Prod.* Nova Genera et Species Plantarum Olavi Swartz, M.D. *Holmiæ.* 1788.
- Sweet Ger.* Geraniaceæ. The Natural Order of Gerania; by Robert Sweet, F.L.S. *Lond.* 1820—8.
- Thumb. Jap.* Caroli Petri Thumberg Flora Japonica. *Lipsiæ.* 1784.
- Thumb. Prod.* Prodomus Plantarum Capensium Carolo Petro Thumberg. *Upsaliæ.* 1794.
- Tourn. Inst.* Josephi Pitton Tournefort Institutiones Rei Herbariæ. *Paris.* 1700.

Willd. Berol. Caroli Ludovici Willdenow Hortus
Berolinensis. *Berolini.* 1803—6.

Willd. Sp. Pl. Caroli à Linnée Species Plantarum,
Curante Carolo Ludovico Willde-
now. *Berolini.* 1797—

The reader is requested to bear in mind that the genus
Triphasia should have been placed in the class Hexandria, order
Monogynia; and to make the following necessary corrections:

P. 32, l. 20, for *vegetabilium*, read *vegetabilium*.

P. 56, l. 20, for *f. 1*, read *f. 2*.

P. 224, l. 29, for *lay*, read *lies*.

P. 272, l. 10 and 18, for *Monkies*, read *Monkeys*.

P. 385, l. 16, for *calice*, read *calyce*.

TO THE BINDER.—The plates are to be placed immediately
before the Introductory Essay.

MONANDRIA MONOGYNIA.

CANNA.

Scitamineæ. *Lin.*—Cannæ. *Juss.*—Canneæ. *Brown.*

indica. C. corollæ limbo interiore trifido: laciniis lanceolatis acuminatis strictis. *Roscoe*
Lin. Tr. 8. p. 338. Hort. Kew. 1. p. 1.
Bot. Reg. t. 776. 2

C. *indica.* α . *rubra.* *Willd. Sp. Pl. 1. p. 3.*
Canacorus. *Rumph. Amb. 5. p. 177. t. 71. f. 2.*
Common Indian Shot. *Sweet.*

coccinea. C. corollæ limbo interiore trifido: laciniis emarginatis strictis. *Roscoe Lin. Tr. 8.*
p. 338. Hort. Kew. 1. p. 1. 2

C. *indica.* γ . *coccinea.* *Willd. Sp. Pl. 1. p. 3.*
C. *indica.* *Bot. Mag. t. 454.*
Scarlet Indian Shot. *Sweet.*

lutæa. C. corollæ limbo interiore bifido. *Roscoe*
Lin. Tr. 8. p. 338. Hort. Kew. 1. p. 2.
Bot. Reg. t. 773.

- C. indica.* β . *lutea.* *Willd. Sp. Pl.* 1. p. 3.
 Yellow Indian Shot. *Sweet.*
- occiden-* *C. corollæ limbi interioris labio superiore*
talis. bipartito: laciniis integris ovatis inæqua-
 libus: labio inferiore declinato (vel rec-
 tius revoluto). *Roscoe MS. in Bot. Reg.*
t. 772. 4
- C. indica.* *Sl.* 1. p. 253.
 Plantain Shot. *Hughes.* 168.
 West Indian Shot. *Sweet.*

MARANTA.

Scitamineæ. *Lin.*—Cannæ. *Juss.*—Cannæ. *Brown.*

- arundi-* *M. culmo ramoso herbaceo, foliis ovato-*
nacea. lanceolatis subtus pilosiusculis. *Roscoe*
Lin. Tr. 8. p. 339. *Hort. Kew.* 1. p. 2.
Willd. Sp. Pl. 1. p. 13. *Aubl.* 3. *Bot.*
Mag. t. 2307. 4
- M. foliis lanceolato-ovatis.* *Br.* 112.
Canna indica radice alba alexipharmaca.
Sl. 1. p. 253. *t.* 149. *f.* 2.
 Arrow-Root. *Hughes.* 221.
 Indian Arrow-Root. *Long.* 2. p. 759.
Miller.

ALPINIA.

Scitamineæ. *Lin. Brown.*—Cannæ. *Juss.*

- nutans.* A. nectario obsolete trilobato margine involuto, foliis lanceolatis, capsula membranacea lanuginosa. *Roscoe Lin. Tr.* 8. p. 346. *Hort. Kew.* 1. p. 4. *Bot. Mag.* t. 1903. *Exot. Bot.* t. 106. 4
- Globba nutans. *Willd. Sp. Pl.* 1. p. 153.
- Renealmia nutans. *Bot. Repos.* t. 360.
- Drooping Alpinia. *Aiton.*

ZINGIBER.

Scitamineæ. *Brown. (Amomum) Lin.*—Cannæ. *Juss.*

- officinale.* Z. bracteis ovato-lanceolatis, laciniis corollæ revolutis, nectario trilobato. *Roscoe Lin. Tr.* 8. p. 348. *Hort. Kew.* 1. p. 5. 4
- Zinziber foliis lanceolatis. *Br.* 119.
- Amomum zingiber. *Willd. Sp. Pl.* 1. p. 6. *Aubl.* 2.
- Ginger. *Hughes.* 233. *Long.* 2. p. 700.
- Narrow-leaved Ginger. *Miller.*

CURCUMA.

Scitamineæ. *Lin. Brown.*—Cannæ. *Juss*

- longa.* C. foliis ovato-lanceolatis, bracteis spatulatis. *Roscoe Lin. Tr. 8. p. 355. Hort. Kew. 1. p. 9. Willd. Sp. Pl. 1. p. 14. Blackw. t. 396. Bot. Reg. t. 886. 2*
Long-rooted Turmeric. *Miller.*

BOERHAAVIA.

Aggregatæ. *Lin.*—Nyctagines. *Juss. Brown.*

- diffusa.** B. caule tereti pubescente, floribus capitato-corymbosis. *Willd. Sp. Pl. 1. p. 20. Hort. Kew. 1. p. 11. Br. 123. 2*
Valerianella folio subrotundo. *Sl. 1. p. 210.*
Hog-weed. *Long. 2. p. 760.*
Spreading Hog-weed. *Miller.*

* This genus is by many considered as properly belonging to the class Diandria. "Genus potius ad diandriam referri debet, plurimæ enim species diandræ, rarissime filamentis unico gaudent."—*Swtz. Obs. 8.* It is placed in the class Diandria by Roemer and Shultes, in their edition of the *Systema Vegetabilium*, now in the progress of publication. Jussieu (*Gen. Pl. 102.*) affixes to the character of the genus "stamina 1. aut 2." Willde-

DIANDRIA MONOGYNIA.

JASMINUM.

Sepiariæ. *Lin.*—Jasmineæ. *Juss. Brown.*

- sambac.* *J. foliis oppositis simplicibus ellipticis ovatis subcordatisque membranaceis opacis, ramulis petiolisque pubescentibus, laciniis calycinis subulatis.* *Willd. Sp. Pl.* 1. p. 35. *Hort. Kew.* 1. p. 15. $\frac{1}{2}$
 α . floribus simplicibus. *Bot. Reg.* t. 1.
 Arabian Jessammy. *Hughes.* 165.
 Single-flowered Arabian Jasmine.
 β . floribus multiplicatis. *Bot. Repos.* t. 497.
 Common double Arabian Jasmine.
 γ . floribus plenis. *Bot. Mag.* t. 1785.
 Great flowered double Arabian or
 Tuscan Jasmine.

now says, of the species *diffusa*, “flores purpurei monandri;” but after repeated and careful examination, I am quite certain that the flowers of *B. diffusa* are monandrous or diandrous on the same plant. What then is to be considered as the specific distinction of *B. husuta*?

- hirsutum.* *J. foliis oppositis ovatis cordatisve ramisque hirsutis, foliolis calycinis linearibus hirsutis.* *Hort. Kew.* 1. p. 16. *Bot. Mag. t.* 1991. *Bot. Reg. t.* 15. *Exot. Bot. t.* 118. h
- J. hirsutum.* *Willd. Sp. Pl.* 1. p. 36.
- J. pubescens.* *Willd. Sp. Pl.* 1. p. 37.
- J. multiflorum.* *Bot. Repos. t.* 496.
- Hairy Indian Jasmine. *Aiton.*
- azoricum.* *J. foliis oppositis ternatis, foliolis ovatis subcordatisque undulatis, ramis glabris teretibus, corollæ laciniis tubo æqualibus.* *Willd. Sp. Pl.* 1. p. 39. *Hort. Kew.* 1. p. 17. *Bot. Mag. t.* 1889. *Bot. Reg. t.* 89. h
- J. album trifoliatum.* *Pluk. Alm.* 195. *Phyt. t.* 303. f. 2.
- Azorian Jasmine. *Aiton.*
- revolutum.* *J. foliis omnibus pinnatis, paniculis oppositifoliis, laciniis corollæ revolutis.* *Bot. Mag. t.* 1731. *Bot. Reg. t.* 178. h
- Yellow Nepaul Jasmine. *Sweet.*
- officinale.* *J. foliis oppositis pinnatis, foliolis acuminatis, gemmulis erectiusculis.* *Willd. Sp. Pl.* 1. p. 40. *Hort. Kew.* 1. p. 18. *Bot. Mag. t.* 31. h
- J. volubile.* *Br.* 115.
- Common white Jasmine. *Miller.*

JUSTICIA.

Personatæ. *Lin.*—Acanthi. *Juss.*—Acanthaceæ. *Brown.*

sexangu- *J. pedunculis trifloris, bracteis cuneiformibus foliis ovatis. Willd. Sp. Pl. 1. p. 83.*
laris.

Hort. Kew. 1. p. 35. ☉

Dicliptera sexangularis. Roem. Syst. 1. p. 171.

Chickweed-leaved Justicia. *Miller.*

coccinea. *J. spicis terminalibus, foliis bracteisque ellipticis margine nudis, corollarum galea lanceolata apice reflexa. Willd. Sp. Pl. 1.*

p. 86. Hort. Kew. 1 p. 36. Aubl. 10.

t. 3. Bot. Mag. t. 432. h

J. lucida. Bot. Repos. t. 313.

Scarlet-flowered Justicia. *Miller.*

picta. *J. racemis axillaribus terminalibusque, floribus verticillatis, foliis ovatis pictis.*

Willd. Sp. Pl. 1. p. 88. Hort. Kew. 1.

p. 37. h

α. foliis macula alba pictis.

β. foliis macula rubente pictis.

*γ. foliis viridibus.**

* Not only does the first variety frequently exhibit branches furnished entirely with plain leaves, but I have known an old shrub raised from seeds of the first variety, which never had a

♂. foliis lurido-sanguineis. *Bot. Mag.*
t. 1870.

Painted Justicia. *Aiton.*

nitida. J. racemis terminalibus* subramosis, floribus subverticillatis, foliis lanceolato-ellipticis utrinque attenuatis. *Willd. Sp. Pl.* 1. p. 88. *Hort. Kew.* 1. p. 36. *Swtz. fl.* 26. *Jacq. Amer.* 5. h

Teucroides siliculosum. *Sl.* 1. p. 37. t. 10. f. 2.

Glossy Justicia. *Aiton.*

secunda. J. racemis terminalibus compositis: racemulis secundis, foliis ovato-lanceolatis acuminatis. *Willd. Sp. Pl.* 1. p. 91. *Hort. Kew.* 1. p. 37. *Bot. Mag.* t. 2060. h

Side-flowering Justicia. *Aiton.*

retusa. J. spicis terminalibus, bracteis obovatis subretusis imbricatis, foliis ovatis acutis. *Willd. Sp. Pl.* 1. p. 92. *Bot. Cab.* t. 724. h

Dicliptera retusa. *Roem. Syst.* 1. p. 169.

Retuse Justicia. *Sweet.*

pectoralis. J. panicula terminali dichotoma, floribus spicatis remotis. *Hort. Kew.* 1. p. 40. *Willd. Sp. Pl.* 1. p. 93. *Jacq. Amer.* 3. t. 3. *Bot. Reg.* t. 796. u

painted leaf; but in all other respects was perfectly marked by the specific characters of its parent plant.

* The racemes are also axillary at the top of the branches.

Dianthera foliis lanceolatis. *Br.* 118.

Antigua Balsam. *Hughes.* 222.

Jamaica Garden Balsam. *Aiton.*

Adhatoda. *J. spicis axillaribus oppositis, bracteis ovatis acutis nervosis.* *Willd. Sp. Pl.* 1. p. 96. *Hort. Kew.* 1. p. 41. *Bot. Mag.* t. 861. h

Adhatode zeylonensium. *Pluk. Alm.* 9. *Phyt. t.* 173. *f.* 3.

Malabar Nut. *Aiton.*

nervosa. *J. spicis axillaribus terminalibusque imbricatis, bracteis oblongis venosis foliisque ovatis acuminatis.* *Hort. Kew.* 1. p. 42. *Bot. Mag. t.* 1358. h

J. pulchella. *Roxb. Cor.* 2. p. 41. *t.* 177.

Eranthemum nervosum. *Roem. Syst.* 1. p. 174.

Eranthemum pulchellum. *Bot. Repos. t.* 88.

Blue-flowered Justacia. *Aiton.*

bicolor. *J. pedunculis axillaribus trifloris petiolo brevioribus, foliis ovalibus acuminatis glabris, limbo corollæ plano, tubo longissimo.* *Bot. Mag. t.* 1423. h

Eranthemum bicolor. *Spreng. Syst.* 1. p. 88.

Eranthemum hypocrateriforme. *Roem. Syst.* 1. p. 175.

Two-coloured Eranthemum. *Sweet.*

ROSMARINUS.

Verticillatæ. *Lin.*—*Labiatae. Juss. Brown.*

- officinalis.* *R. foliis sessilibus. Willd. Sp. Pl. 1. p.*
 126. *Hort. Kew. 1. p. 52.* 4
R. fruticosus incanis. Br. 117.
 Officinal Rosemary. *Miller.*

SALVIA.

Verticillatæ. *Lin.*—*Labiatae. Juss. Brown.*

- officinalis.* *S. foliis lanceolato-ovatis crenulatis, ver-*
 ticillis paucifloris, calycibus mucronatis.
 Willd. Sp. Pl. 1. p. 129. Hort. Kew. 1.
 p. 52. Br. 117. 4
 Garden Sage. *Miller.*
- occiden-* *S. foliis ovatis serratis, spicis laxis, brac-*
 talis. teis cordatis subtrifloris. Willd. Sp. Pl.
 1. p. 131. *Swtz. fl. 43.* 4
 S. spicata repens. Br. 117.
 Hop-weed (*red*). *Hughes. 169.*
 West Indian Sage. *Miller.*
 Creeping sweet-scented Sage. *Browne.*

DIANDRIA TRIGYNIA.

PIPER.

Piperitæ. *Lin.*—Urticis aff. *Juss.*—Piperaciæ. *Kunth.*

Amalago. P. foliis lanceolato-ovatis quinquenerviis rugosis. *Willd. Sp. Pl.* 1. p. 160. *Hort. Kew.* 1. p. 69. h

P. frutescens diffusum. *Br.* 121.

P. longum arboreum altius. *Sl.* 1. p. 134. t. 87. f. 1.

Rough-leaved Pepper. *Miller.*

Small-grained Black Pepper. *Browne.*

aduncum. P. foliis oblongo-ovatis acuminatis basi inæqualibus venosis, spicis solitariis axillaribus uncinatis. *Willd. Sp. Pl.* 1. p. 162. *Hort. Kew.* 1. p. 70. *Aubl.* 21. *Jacq. Ic.* 2. t. 210. h

P. frutescens diffusum. *Br.* 122.

P. longum folio nervoso. *Sl.* 1. p. 135. t. 87. f. 2.

Saururus arborescens fructu adunco. *Plum. Amer.* 58. t. 77.

Rock-Bush. *Hughes.* 219.

Hooked-spiked Pepper. *Miller.*

- pellucidum.* P. foliis cordatis petiolatis, caule herbaceo. *Willd. Sp. Pl.* 1. p. 163. *Hort. Kew.* 1. p. 71. *Aubl.* 21. ☉
Piperomia pellucida. *Hort. Br.* 380.
Saururus minor procumbens. *Plum. Amer.* 54. t. 72.
 Pellucid-leaved Pepper. *Miller.*
- obtusifolium.* P. foliis obovatis enerviis carnosiss, caule prostrato. M—. *Willd. Sp. Pl.* 1. p. 165. *Hort. Kew.* 1. p. 71. *Aubl.* 22. *Jacq. Ic.* 1. t. 9. 2
 P. longum humilius. *Sl.* 1. p. 136.*
Piperomia obtusifolia. *Hort. Br.* 380.
Saururus major repens. *Br.* 204.
Saururus humilis. *Plum. Amer.* 53. t. 70.
 Rock-Balsam. *Hughes.* 221.
 Blunt-leaved Pepper. *Miller.*
- acuminatum.* P. herbaceum, foliis lanceolato-ovatis nervosis carnosiss, caule erectiusculo. *Willd. Sp. Pl.* 1. p. 164. *Aubl.* 21. 2
Piperomia acuminata. *Hort. Br.* 380.

* The leaves of the Rock Balsam are in general smaller and more pointed than those of the "Piper longum humilius" of Sloane, which are represented as being "four inches long and two broad;" but Sloane observes, "there is a variety of this with smaller and more pointed leaves, which I gathered in Barbadoes," "on rocky ground." The leaves of this kind of Piper vary much in Barbados in size and form: they occur not only round-pointed, as represented in Plumier's plate, but even a little emarginate, as remarked by Sloane.

Saururus alius humilis. *Plum. Amer.* 54.
t. 71.

Rock Balsam. *Hughes.* 221.

peltatum. P. foliis peltatis orbiculato-cordatis obtusis repandis, spicis umbellatis. *Willd. Sp. Pl.* 1. p. 166. *Hort. Kew.* 1. p. 72. *Aubl.* 22. 24

Saururus foliis amplis. *Br.* 203. No. 2.

Saururus arborescens foliis amplis rotundis. *Plum. Amer.* 56. t. 74.

Peltated Pepper. *Aiton.*

Monkey's Hand.

umbellatum. P. foliis orbiculato-cordatis acuminatis venosis, spicis umbellatis, caule erecto sulcato pubescente. *Willd. Sp. Pl.* 1. p. 167. *Hort. Kew.* 1. p. 72. *Aubl.* 22. *Jacq. Ic.* 2. t. 204. 1/2

P. longum racemosum malvaceum. *Sl.* 1. p. 136.

Saururus foliis amplis. *Br.* 203. No. 1.

Saururus arborescens foliis amplis cordatis. *Plum. Amer.* 53. t. 73.

Umbelled-Pepper, or Santa Maria Leaf. *Miller.*

Wild Elder Bush.

TRIANDRIA MONOGYNIA.

MELOTHRIA.

Cucurbitaceæ. *Lin. Juss. De Cand.*

pendula. M. foliis cordatis quinquelobis dentatis, cirris simplicibus, floribus femineis solitariis longe pedunculatis, corolla subpilosa denliculata, fructibus ovato-subglobo-
sis pendulis. *De Cand. Prod.* 3. p. 313.
Willd. Sp. Pl. 1. p. 189. *Hort. Kew.* 1.
p. 78. ☉

M. scandens. *Br.* 124.

Cucumis minima. *Sl.* 1. p. 227. t. 142. f. 1.

Small Wild Cucumber. *Hughes*, 166.

Pendulous Melothria. *Aiton.*

MORÆA.

Ensatae. *Lin.*—Irides. *Juss.*—Irideæ. *Brown.*

plicata. M. scapo tereti, foliis petiolatis oblongis nervosis plicatis, racemis bifidis. *Willd.*
Sp. Pl. 1. p. 243. *Swtz. fl.* 82. 2

- M. palmifolia.* *Jacq. Ic. 2. t. 227.*
Marica plicata. *Bot. Mag. t. 655.*
Sisyrinchium latifolium. *Swtz. Prod. 17.*
Hort. Kew. 4. p. 135.
Ixia americana. *Aubl. 33.*
Bermudiana palmæ folio. *Tourn. Inst. 388.*
 Blood-worth. *Hughes, 170.*
 Broad-leaved *Sisyrinchium.* *Sweet.*

COMMELINA.

Ensatae. Lin.—Junci. Juss.—Commelineæ. Brown.

- communis.* *C. corollis inæqualibus, foliis ovato-lanceolatis acutis, caule repente glabro.*
Willd. Sp. Pl. 1. p. 249. Hort. Kew. 1.
p. 123. Aubl. 37. ☉
 Pond-grass or Canker-weed. *Hughes, 239.*
 Common American Commelina. *Miller.*
- erecta.* *C. corollis inæqualibus, foliis ovato-lanceolatis, caule erecto scabro simplissimo.*
Willd. Sp. Pl. 1. p. 250. Hort. Kew. 1.
p. 124. ☉
 Pond-grass or Canker-weed. *Hughes, 239.*
 Upright Virginian Commelina. *Miller.*

CALLISIA.

Ensatae. *Lin.*—Junci. *Juss.*—Commelineae. *Brown.*

- repens.* C. foliis ovato-lanceolatis sessilibus, caule procumbente, floribus axillaribus sessilibus. *Roem. Syst.* 1. p. 527. *Willd. Sp. Pl.* 1. p. 254. *Hort. Kew.* 1. p. 125. *Aubl.* 39. 2
Hapalanthus repens. *Jacq. Amer.* 11. t. 11.
 Creeping Callisia. *Aiton.*

KYLLINGIA.

Cyperoideae. *Juss.*—Cyperaceae. *De Cand. Brown.*

- monocephala.* K. spiculis unifloris, squamis pilosociliatis tenuissime nervosis, capitulo indiviso vel trilobo, involucri 3—4-phyllo elongato foliisque linearibus. *Brown. Prod.* 75. *Willd. Sp. Pl.* 1. p. 256. *Hort. Kew.* 1. p. 125. *Rottb. Gr. t.* 4. f. 4. 2
Cyperus minimus pratensis. *Br.* 127.
 One-headed Kyllingia. *Aiton.*

SCHCENUS.

Calamariæ. *Lin.*—Cyperoideæ. *Juss.*

Cyperaceæ. *De Cand. Brown.*

stellatus. S. culmo triquetro basi folioso, capitulo glomerato parvo, involucri stellato colorato pentaphyllo. *Willd. Sp. Pl. 1. p. 264.*

Swtz. fl. 102.

4

Dichroma ciliatum. *Ham. Prod. 15.*

Gramen cyperoides spica compacta alba.

Sl. 1. p. 119. t. 78. f. 1.

Star Grass. *Hughes. 238.*

Starry Bog-Rush. *Miller.*

CYPERUS.

Calamariæ. *Lin.*—Cyperoideæ. *Juss.*

Cyperaceæ. *De Cand. Brown.*

ligularis. C. culmo triquetro, umbella composita, spiculis paucifloris spicatis, glumis obtuse mucronatis, involucri polyphylo longissimo. *Willd. Sp. Pl. 1. p. 277. Aubl. 47.*

Rottb. Gr. t. 11. f. 2.

4

C. major subtriquetrus. *Br. 128.*

C. maximus paucula minus sparsa. *Sl.* 1.
p. 36. *t.* 9.

Dwrah, or Ammadwrah. *Hughes.* 240.

imbrica-
tus.

C. culmo triquetro nudo, umbella decomposita, umbellarum radiis imbricatis, involucri hexaphylli foliis tribus longissimis tribus brevissimis. *Willd. Sp. Pl.* 1 *p.* 278. 4

elegans.

C. culmo triquetro, umbella decomposita, spiculis ovatis, glumis mucronatis patulis, involucri tetraphyllo umbella longiore. *Willd. Sp. Pl.* 1. *p.* 278. 4

C. major umbellatus. *Br.* 128.

C. panicula maxime sparsa ferruginea. *Sl.* 1. *p.* 117. *t.* 75. *f.* 1.

Elegant Cyperus. *Sweet.*

strigosus.

C. culmo triquetro nudo, umbella simplici, spiculis linearibus confertissimis horizontalibus. *Willd. Sp. Pl.* 1. *p.* 281. *Hort. Kew.* 1. *p.* 129. 4

C. rotundus panicula sparsa. *Sl.* 1. *p.* 116. *t.* 74. *f.* 2. 3.

Wild Dwrah. *Hughes.* 241.

Bristled-spiked Cyperus. *Aiton.*

tenuis.

C. culmo triquetro nudo, umbella simplici, spiculis subulatis confertis horizontali-reflexis, foliis involucri lineari-setaceis. *Willd. Sp. Pl.* 1. *p.* 281. *Swtz. fl.* 119. *Rottb. Gr.* *t.* 9. *f.* 3. 4

compres- C. culmo triquetro nudo, umbella uni-
sus. versali triphylla, glumis mucronatis late-
 ribus membranaceis. *Willd. Sp. Pl.* 1.
p. 282. 4

C. rotundus gramineus. *Sl.* 1. *p.* 117.
t. 76. *f.* 1.

Compressed Cyperus. *Sweet.*

odoratus. C. culmo triquetro nudo, umbella decom-
 posita simpliciter foliosa, pedicellis dis-
 tiche spicatis. *Willd. Sp. Pl.* 1. *p.* 284. 4

C. longus odoratus. *Sl.* 1. *p.* 116. *t.* 8.
et t. 74. *f.* 1.

esculentus. C. culmo triquetro nudo, umbella foliosa,
 radicum tuberibus ovatis zonis imbri-
 catis. *Willd. Sp. Pl.* 1. *p.* 284. *Hort.*
Kew. 1. *p.* 129. 4

C. rotundus esculentus. *Mor. Hist.* 236.
t. 11. *f.* 10.

Nut Grass. *Hughes.* 241.

Eatable Cyperus—Rush-nut. *Aiton.*

SCIRPUS.

Calamanæ. *Lin.*—Cyperoideæ. *Juss.*

Cyperaceæ. *De Cand. Brown.*

mutatus. S. culmo triquetro nudo, spica cylin-
 drica terminali. *Willd. Sp. Pl.* 1. *p.* 290.
Aubl. 49. 4

- S. culmo triquetro, spica oblonga. Br. 126.*
Common Rush.
- geniculatus.* *S. culmo tereti nudo, spica oblonga terminali. Willd. Sp. Pl. 1. p. 291. Aubl. 49.*
S. culmo rotundo. Br. 126. 2
Juncus aquaticus geniculatus. Sl. 1. p. 122. t. 75. f. 2.
Knee jointed Club-Rush. *Miller.*
- ferrugineus.* *S. culmo triquetro subnudo, involucris longitudine paniculæ ciliatis. Willd. Sp. Pl. 1. p. 304. Aubl. 49. 2*
Gramen cyperoides majus. Sl. 1. p. 118. t. 77. f. 2.

CENCHRUS.

Gramina, *Lin.*—Gramineæ. *Juss. Brown. De Cand.*

- echinatus.* *C. spica spiculis approximatis, involucris decempartitis villosis. Roem. Syst. 2. p. 253. Willd. Sp. Pl. 1. p. 317. Hort. Kew. 1. p. 135. Aubl. 933. Cav. Ic. 5. p. 39. t. 462. ☉*
C. spica oblonga simplici echinata. Br. 367.
Gramen echinatum maximum. Sl. 1. p. 108.
Rough-spiked Cenchrus. *Sweet.*
Bur Grass.

tribuloides. C. spica spiculis alternis, involucris integris spinosis. *Roem. Syst.* 2. p. 253.

Willd. Sp. Pl. 1. p. 317. ☉

Gramen maritimum echinatum procumbens. *Sl.* 1. p. 108. t. 65. f. 1.

Spinous Cenchrus. *Sweet.*

Bur Grass.

TRIANDRIA DIGYNIA.

SACCHARUM.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

officina- S. floribus paniculatis, foliis planis.
rum. *Roem. Syst.* 2. p. 285. *Willd. Sp. Pl.* 1.
p. 321. *Hort. Kew.* 1. p. 136. *Aubl.*
50. 4

α. commune.

Native Cane.

*β. otaheitense.**

Bourbon Cane.

S. geniculatum et succulentum. *Br.* 129.
Arundo saccharifera. *Sl.* 1. p. 108. t. 66.
Sugar Cane. *Hughes.* 244. *Long.* 2. p. 674.
Common Sugar Cane. *Sweet.*

* Var. *β.* per Gallos et serius per Anglos in Antillas delata differt statura majore robustiore, internodiis longioribus, pilis spicula duplo triplo longioribus, glumis binerviis nec obsolete uninerviis, apice minus ciliatis. An species distincta?—(*Roem. et Shultes Syst. l. c.*) I have little doubt that the Bourbon and Native Canes are distinct species, although I cannot more satisfactorily point out their specific difference.

*viola-
ceum.**

S. culmo foliisque violaceo-coloratis, panicula effusa, spiculis triandris, glumis quadrinerviis dorso longissime pilosis.

Roem. Syst. 2. p. 285.

4

Violet-coloured Sugar-Cane. *Sweet.*

ERIANTHUS.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

*saccha-
roides.*

E. culmo altissimo, vaginis foliorum ad collum confertissima lana barbatis, panicula stricta, gluma villis involucrantibus multo brevioribus: valvis exterioribus juxta margines longe villosis. *Roem. Syst. 2.*

p. 323.

4

Wild Cane. *Hughes. addenda.*

Sugar-cane-like Erianthus. *Sweet.*

* This species, commonly considered as a variety, is frequently found in fields of Canes, and is readily to be distinguished.

ANDROPOGON.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

angusti- A. foliis linearibus canaliculatis pilosis,
folius. spicis digitatis corymbosis plurimis, caly-
cibus striatis pedicellisque hirsutis.

Roem. Syst. 2. p. 822. 4

A. Ischæmum. *Spreng. Syst. 1. p. 286.*

Jacq. Austr. 4. p. 43. t. 384.

Red Flag, or Plush-Grass. *Hughes. 241.*

Narrow-leaved Andropogon. *Sweet.*

Narrow-leaved Sour-Grass.

alopecu- A. panicula simplici oblonga elongata,
roides. spicis verticillatis simplicibus, floribus
aristatis geminatis altero sessili altero
pedicellato, rachi lanata, lana floribus
longiore. *Roem. Syst. 2. p. 815. Willd.*

Sp. Pl. 4. p. 911. 4

Gramen dactylon Alopecuroidis facie.

Sl. 1. p. 113. t. 70. f. 1.

Flag-Grass. *Hughes. 240.*

CYMBOPOGON.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- Schoenanthus.* C. bractea foliacea striata, flosculis subspicatis: flosculis hermaphroditis quaternis. *Roem. Syst.* 2. p. 833. *Spreng. Syst.* 1. p. 289. 4
Andropogon Schoenanthus. *Willd. Sp. Pl.* 4. p. 915. *Hort. Kew.* 5. p. 427.
Schoenanthum amboinicum. *Rumph. Amb.* 5. p. 181. t. 72. f. 2.
 Lemon-Grass. *Sweet,*

CHLORIS.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- radiata.* C. spicis plurimis fasciculatis erectiusculis, flosculis subulatis glabris. *Willd. Sp. Pl.* 4. p. 923. *Hort. Kew.* 5. p. 429. 4
Swtz. Fl. 201.
 Plush-Grass. *Hughes.* 239.
 Many-spiked Chloris. *Aiton.*

RHABDOCHLOA.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- mucronata.* R. culmo repente, spicis plerumque quatuor, rachi triangulari mucronata, spiculis subquadrifloris, gluma exterior mucronata. *Roem. Syst. 2. p. 618.* 4
Chloris mucronata. Mich. Amer. 1. p. 59.
Dactyloctenium meridionale. Ham. Prod. 6.
 Gramen dactylon spicis gracilioribus.
*Sl. 1. p. 111. t. 68, f. 3.**
 Mucronate Rhabdochloa, *Sweet.*
- cruciata.* R. spicis subquaternis cruciatis, calycinis valvulis acuminatis, oris vaginarum glabris. *Roem. Syst. 2. p. 617.* 4
Chloris cruciata. Willd. Sp. Pl. 4. p. 923.
Swtz. Fl. 192.
 Cross-spiked Rhabdochloa, *Sweet.*

* This very characteristic plate of a grass growing in Barbados is, by all the authors I have met with, referred to *Chloris radiata*, with the epithet *pessima* affixed to it; an example of the evil of following authority in the quotation of synonyms. Sloane's plate of a grass with four spikes disposed crosswise would, indeed, be a very bad representation of *Chloris radiata*, with very many fascicled spikes, had it ever been intended as a figure of that grass.

SORGHUM.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- vulgare.* S. calycibus obovatis nitidis pilosiusculis apice pubescentibus, seminibus compressis, panicula coarctata oblonga. *Roem. Syst. 2. p. 836.* 4
 Holcus Sorghum. *Willd. Sp. Pl. 4. p. 929. Hort. Kew. 5. p. 430.*
 Panicum erectum maximum. *Br. 135.*
 Milium indicum arundinaceo caule. *Sl. 1. p. 104.*
 Guinea Corn. *Hughes. 254.*
 Indian Millet. *Sweet.*
- bicolor.* S. calycibus apice strigoso-pubescentibus nigris, seminibus albis, panicula coarctata ovata. *Roem. Syst. 2. p. 837.* 4
 Holcus bicolor. *Willd. Sp. Pl. 4. p. 929.*
 Guinea Corn. *Hughes. 254.*
 Two-coloured Sorghum. *Sweet.*
- cernuum.* S. calycibus villosis, seminibus globosis, panicula coarctata ovata pendula. *Roem. Syst. 2. p. 837.* 4
 Holcus cernuus. *Willd. Sp. Pl. 4. p. 930.*
 Guinea Corn. *Hughes. 254.*
 Nodding Sorghum. *Sweet.*

- saccharatum.* S. calycibus oblongis acutis pubescentibus, seminibus compressis, panícula effusa patentissima. *Roem. Syst.* 2. p. 837. 4
Holcus saccharatus. *Willd. Sp. Pl.* 4. p. 930.
 Guinea Corn. *Hughes.* 254.
 Yellow-seeded Sorghum. *Sweet.*

ZEA.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- Mays.* Z. foliis integerrimis. *Willd. Sp. Pl.* 4. p. 200. *Hort. Kew.* 5. p. 235. *Aubl.* 847. ☉
 Z. seminibus subcompressis obovatis. *Br.* 335.
Mays. Frumentum turcicum. *Blackw.* t. 547. a. b.
 Frumentum indicum *Mays* dictum. *Sl.* 1. p. 105.
 Indian Corn. *Hughes.* 253. t. 23. f. 1.
 Great Corn, or Indian Maize. *Long.* 2. p. 762.
 Indian Corn, or Maize. *Miller.*

COIX.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

Lacryma. C. culmo superne semitereti obtuso,
floribus nudis, fructibus ovatis. *Willd.*
Sp. Pl. 4. p. 202. *Hort. Kew.* 5. p. 236.
Aubl. 847. *Bot. Mag.* t. 2479. 2

Lacryma Jobi. *Rumph. Amb.* 5. p. 193.
t. 75. f. 2.

Lithospermum arundinaceum. *Bauh.*
Pin. 258.

Job's Tears. *Long.* 2. p. 831.

Common Job's Tears. *Sweet.*

SCLERIA.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

Flagel- S. culmo triquetro scaberrimo, foliis
lum. margine et costa media aculeato-his-
pidis, paniculis terminalibus et axillari-
bus corymbosis, pedunculis communibus
hirsutis, nucibus lævibus nitidis apice
puncto opaco instructis. *Willd. Sp. Pl.*
4. p. 312. *Swtz. Fl.* 88. 2

Gramen cyperoides sylvaticum maximum. *Sl.* 1. p. 118. t. 77. f. 1.

- hirtella.* S. culmo simplissimo triquetro foliisque pubescentibus, spica glomerata nuda: glomerulis alternis distantibus, glumis setosis, nucibus elliptico-globosis mucronatis lævibus nitidis. *Willd. Sp. Pl.* 4. p. 318. *Swtz. Fl.* 93. 4
Hairy Scleria. *Sweet.*

PASPALUM.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- virgatum.* P. spicis paniculatis alternis basi villosis, floribus geminis. *Willd. Sp. Pl.* 1. p. 331. 4

Gramen dactylon majus. *Sl.* 1. p. 112. t. 69. f. 2.

Rod-like Paspalum. *Miller.*

- distichum.* P. spicis duabus erectiusculis altera sessili, floribus oblongis glabris, culmo ascendente. *Willd. Sp. Pl.* 1. p. 332. *Hort. Kew.* 1. p. 138. *Swtz. Obs.* 35. t. 2. f. 1. 4

Gramen dactylon bicornis, spicis purpurascens majus. *Sl.* 1. p. 112. t. 65. f. 3.

Savannah Grass. *Hughes. 239.*

Two-spiked Paspalum. *Miller.*

conjugatum. P. spicis duabus horizontalibus conjugatis, spiculis ovatis, culmo erecto, foliis involutis. *Willd. Sp. Pl. 1. p. 333. Swtz. Fl. 133.* 2

P. tenue. *Gaert. Fruct. 2. p. 2. t. 80. f. 4.*

Gramen dactylon bicornis repens. *Sl. 1. p. 112.*

Conjugate-spiked Paspalum. *Miller.*

Broad-leaved Sour Grass.

PANICUM.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

colonum. P. spicis alternis secundis muticis ovatis scabris, rachi teretiuscula. *Willd. Sp. Pl. 1. p. 338. Hort. Kew. 1. p. 141. ☉*
Gramen panicum minimum. *Sl. 1. p. 107. t. 64. f. 3.*

Purple Panic Grass. *Miller.*

pilosum? P. spicis paniculatis alternis secundis, spiculis geminis flosculo altero minore acuminatis lævibus, rachi compressa pilosa, culmo divaricato geniculato. *Roem. Syst. 1. p. 427. Willd. Sp. Pl. 1. p. 340. Swtz. Fl. 141.* 2

Scotch Grass. *Hughes*. 240.

molle. P. spicis paniculatis alternis secundis patentibus, spiculis approximatis pedicellatis secundis muticis. *Willd. Sp. Pl.* 1. p. 340. *Swtz. Fl.* 143. 4

P. silvestris. *Br.* 133.

Rice Grass. *Hughes*. 239.

Soft Panic Grass. *Miller*.

fasciculatum. P. spicis paniculatis alternis erectis subfastigiatis, spiculis secundis subrotundis. *Willd. Sp. Pl.* 1. p. 341. *Swtz. Fl.* 145. ☉

Fascicled Panic Grass. *Miller*.

carthaginense. P. spicis paniculatis foliis brevioribus, spiculis secundis ovatis acutis, culmis prostrato-ramosis, vaginis pubescentibus. *Roem. Syst.* 2. p. 427. *Willd. Sp. Pl.* 1. p. 341. *Swtz. Fl.* 148. 4

Carthagenâ Panic Grass. *Sweet*.

lineare. P. spicis digitatis subquaternis linearibus, flosculis solitariis secundis muticis. *Willd. Sp. Pl.* 1. p. 344. 4

Gramen dactylon panicula longa. *Sl.* 1. p. 113. t. 70. f. 3.

Linear-spiked Panic Grass. *Miller*.

distachyon. P. Spicis geminis secundis lævibus. *Willd. Sp. Pl.* 1. p. 345. ☉

Digitaria distachya. *Ham. Prod.* 6.

Distich-spiked Panic Grass. *Miller*.

Short Grass.

repens. P. panicula virgata, foliis divaricatis.
Willd. Sp. Pl. 1. p. 347. Hort. Kew. 1.
p. 143. ☉

Creeping Panic Grass. *Miller.*

latifolium. P. panicula racemis lateralibus simplici-
 bus, foliis ovato-lanceolatis collo pilosis.
Willd. Sp. Pl. 1. p. 350. Hort. Kew. 1.
p. 144. Aubl. 50. 4

Ginger Grass. *Hughes. 241.*

Broad-leaved Panic Grass. *Miller.*

May Grass.

arundina- P. panicula composita patente: ramis ra-
ceum. mulisque strictis capillaribus, spiculis
 subrotundis, culmo subdiviso geniculato,
 foliis lato-lanceolatis acuminatis rigidis.

Willd. Sp. Pl. 1. p. 353. Swtz. Fl. 166. 4

Arundo silvestris ramosa tenuis. *Br. 138.*

Gramen miliaceum sylvaticum maxi-
 mum. *Sl. 1. p. 114. t. 71. f. 3?*

Reed: *Hughes. 252.*

Reedy Panic Grass. *Miller.*

polyga- P. panicula composita capillari patente,
mum. flosculis polygamis, culmo articulato hir-
 suto. *Willd. Sp. Pl. 1. p. 353. Swtz.*
Prod. 24. 4

P. maximum. Swtz. Fl. 170. Jacq. Ic. 1.
t. 13.

Holcus major assurgens. *Br. 366.*

Guinea Grass. *Sweet.*

CYNODON.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

dactylon. C. spicis 3—5—digitatis, perianthiis
glabris subciliatis gluma longioribus,
setula imberbi ad basin valvulæ inte-
rioris. *Brown. Prod.* 43. *Hort. Gram.*
290. *et Ic. Engl. Fl.* 95. 24
Panicum dactylon. Willd. Sp. Pl. 1. p.
342. *Hort. Kew.* 1. p. 142. *Fl. Br.* 67.
Engl. Bot. t. 850.
Creeping Dog's-tooth Grass. *Smith.*
Devil's Grass.*

* This is the Durva, Dub, or Doob-grass of the Hindoos.
vid. Lambert, in Lin. Tr. 6. Its habitat in Britain is about
Penzance and Marketjue, in Cornwall.—*Fl. Br.* It is there-
fore a native of at least three quarters of the globe. It is a most
injurious weed if allowed to grow with cultivated plants, but
is highly valuable as fodder, and can easily be exterminated.

AGROSTIS.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- purpurascens.* A. panicula coarctata elongata ramis appressis erectis, flosculis inæqualibus acuminatis. *Roem. Syst.* 2. p. 368. *Willd. Sp. Pl.* 1. p. 375. *Swtz. Fl.* 185. ☉
Sporobolus purpurascens. *Ham. Pr.* 5.
Purple Bent Grass. *Miller.*
- indica.* A. panicula contracta mutica, racemis lateralibus erectis alternis. *Roem. Syst.* 2. p. 368. *Willd. Sp. Pl.* 1. p. 375. *Swtz. Obs.* 38. 2
Sporobolus indicus. *Brown. Prod.* 26.
Sporobolus Lamarckii. *Ham. Prod.* 4?
Gramen pratense panicula et foliis longissimis. *Sl.* 1. p. 115. t. 73. f. 1.*
Indian Bent Grass. *Miller.*

* By a strange mistake, Linnæus made this "Gramen pratense" of Sloane a synonym both of "Agrostis purpurascens" and of "Poa ciliaris," in which he has been followed by Miller and Martyn in the Gardener's Dictionary, and by Willdenow in his recent edition of the Species Plantarum. The error seems to have originated in Browne having identified this Gramen of Sloane with his Briza. Even in the recent work, now in the course of publication, of Roemer, and Shultes, this plate, 73. f. 1. is twice referred to—once as a representation of Agrostis

OPLISMENUS.

Gramina. *Lin.*—Gramineæ *Juss. Brown.*

hirtellus. O. spica composita spiculis appressis alternis, calycibus geminis, valvulis omnibus aristatis, extima longissima.

Roem. Syst. 2. p. 481. 2

Panicum hirtellum. *Willd. Sp. Pl.* 1. p. 340. *Hort. Kew.* 1. p. 142.

Orthopogon hirtellus. *Spreng. Syst.* 1. p. 306.

Panicum majus panicula rariori. *Br.* 133.

Gramen panicum maximum. *Sl.* 1. p. 106.

Scotch Grass.* *Long.* 2. p. 761. *Miller.*

Pagister Grass.

indica, and again of *Megastachya ciliaris*, identical with *Poa ciliaris*.—*Willd.* An example, this, of the continuance of error by the copying of synonyms. The *Gramen pratense* of Sloane is certainly an *Agrostis*; and is recognized as such by Swartz.

* Sloane informs us, that this grass takes its familiar name “from that part of Barbados called Scotland,” from which place it was transported into Jamaica. The Scotch grass of Jamaica is different from the grass bearing that name in Barbados. If the references above given on the authority of Willdenow, Roemer, Swartz, Martyn, &c. to Sloane and Browne be correct, the Scotch grass of Jamaica is identical with the Pagister grass of Barbados.

POA.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- ciliaris.* P. panicula contracta, glumarum valvulis interioribus piloso-ciliatis. *Willd. Sp. Pl. 1. p. 402. Hort. Kew. 1. p. 158. Aubl. 51. Jacq. Ic. 2. t. 304.* ☉
 Megastachya ciliaris. *Roem. Syst. 2. p. 592.*
 Briza tenuissima. *Br. 135.*
 Dog's Grass. *Hughes. 241.*
 Ciliated Meadow Grass. *Miller.*

CYNOSURUS.

Gramina. *Lin.*—Gramineæ. *Juss. Brown.*

- indicus.* C. spicis digitatis linearibus, culmo compresso declinato basi nodoso, foliis alternis. *Willd. Sp. Pl. 1. p. 417. Hort. Kew. 1. p. 162. Aubl. 51.* ☉
 Eleusine indica. *Roem. Syst. 2. p. 582.*
 Gramen majusculmo compresso. *Br. 137.**

* This synonym is applied by Swartz (*Fl. 143*) to *Panicum molle*, which, according to him, "Jamaicensibus dicitur Dutch Grass," and for which he refers inaccurately to p. 113. of

Gramen dactylon procumbens. *Sl.* 1.

p. 111.

Dutch Grass. *Hughes.* 239.

Indian Dog's-tail Grass. *Miller.*

DONAX.

Gramina. *Lin.*—Gramineæ (*Arundo*). *Juss.* *Brown.*

arundina- D. calycibus sub-3—5-floris flosculis
ceus. calycem æquantibus, culmo basi lignoso.

Roem. Syst. 2. *p.* 600. 4

Arundo Donax. *Willd. Sp. Pl.* 1. *p.* 454.

Hort. Kew. 1. *p.* 173.

Arundo Sativa. *Donax Dioscoridis.*

Bauh. Pin. 17.

Cultivated Reed. *Miller.*

Browne instead *p.* 137. But in the “Gramen Dactylon Procumbens”—the Dutch Grass of Jamaica known to Sloane—the “spikes are usually three or four, very green, broad and large, all coming from the top of the stalks.” The *Panicum molle* cannot be confounded with this gramen. It is most probable that Sloane and Browne applied the vernacular name, Dutch Grass, to the same plant, and Sloane’s description applies very perfectly to the Dutch Grass of Barbados. Browne’s description is unusually imperfect; as far however as it goes, it corresponds very well with *Cynosurus indicus*; and, for the reasons stated, I have no doubt his “Gramen Majus Culmo Compresso” and *Cynosurus Indicus* are identical.

TRIANDRIA TRIGYNIA.

DRYMARIA.

Caryophyllææ. *De Cand. (Holosteum) Lin. Juss.*

cordata. D. caule foliisque glabris, foliis breviter petiolatis ovato subrotundis acuto-mucronatis basi rotundatis vel obsolete cordatis, pedunculis dichotomis multifloris, calyce glabro petala superante, ovariis 7—10-spermis. *De Cand. Prod. 1. p. 395.* ☉

• *Holosteum cordatum. Willd. Sp. Pl. 1. p. 488.*

Holosteum foliis orbiculatis. Br. 139.

• *Alsine americana. Sl. 1. p. 203.*

Heart-leaved Drymaria. *Sweet.*

TETRANDRIA MONOGYNIA.

SPERMACOCE.

Stellatæ. *Lin.*—*Rubiaceæ. Juss.*—*Spermacocæ. Kunth.*

- tenuior.* *S. glabra*, foliis lanceolatis, staminibus inclusis, floribus verticillatis, seminibus hirtis. *Willd. Sp. Pl.* 1. p. 568. *Hort. Kew.* 1. p. 232. ☉
- S. scandens.* *Br.* 141.
- Anonymos americana.* *Pluk. Alm.* 33. *Phyt. t.* 136. f. 4.
- Slender Button Weed. *Miller.*
- Iron Grass, or Climbing Spermacoce. *Browne.*
- radicans.* *S. glabra*, foliis subsessilibus lanceolatis, floribus verticillatis, caule procumbente radicante. *Willd. Sp. Pl.* 1. p. 570. *Hort. Kew.* 1. p. 233. *Aubl.* 58. t. 20. f. 4. 2
- Rooting Button Weed. *Miller.*
- hirta.* *S. scabra ramosa*, foliis ovato-lanceolatis, floribus confertis axillaribus, staminibus exsertis. *Willd. Sp. Pl.* 1. p. 571. *Jacq. Ic.* 2. t. 308. ☉

S. erecta subhirsuta. . *Br.* 141.

Rough-haired Button Weed. *Miller.*

Purple Eye-bright.

spinosa.

S. suffruticosa, foliis linearibus spinulis
ciliatis. *Willd. Sp. Pl.* 1. p. 573. *Jacq.*

Amer. 21.

2

S. erecta simplex. *Br.* 140.

Thorny Button Weed. *Miller.*

IXORA.

Stellatæ. *Lin.*—*Rubiaceæ.* *Juss.* *Coffeaceæ.* *Kunth.*

coccinea. *I. foliis sessilibus ellipticis obtusis mucronatis, corymbis terminalibus.* *Willd.*

Sp. Pl. 1. p. 609. *Hort. Kew.* 1. p. 244.

Bot. Mag. t. 169?

h

I. grandiflora. *Bot. Reg. t.* 154.

Flamma sylvarum. *Rumph. Amb.* 4. p.
105. *t.* 46 ?

Scarlet Ixora. *Miller.*

*fascicu-
lata.*

*I. foliis ovato-ellipticis; ramulorum sub-
fasciculatis pedunculis filiformibus sub-
trifloris.* *Willd.* 1. p. 610.

h

Chomelia fasciculata. *Swtz. Fl.* 238.

Dart Wood. *Hughes.* 155 ?

PAVETTA.

Stellatæ. *Lin.*—Rubiaceæ. *Juss.*—Coffeaceæ. *Kunth.*

pentandra. *P.* foliis oblongo-lanceolatis acuminatis, panicula trichotoma axillari, floribus pentandris. *Willd. Sp. Pl.* 1. p. 611. *Swtz. Fl.* 233. h

Psychotria Pavetta. *Swtz. Pr.* 45.
Wild Coffee.*

ÆGIPHILA.

Stellatæ. *Lin.*—Vitices. *Juss.*

Verbenaceæ. *Juss. An. du Mus. Brown.*

martini- *Æ.* foliis ovato-lanceolatis acuminatis
censis. glabris, ramis diffusis, paniculis terminalibus axillaribusve, calycibus glabris.

Willd. Sp. Pl. 1. p. 615. *Hort. Kew.* 1. p. 264. *Bot. Cab. t.* 132. h

Spirit Weed. *Hughes.* 156.†
Martinico Ægiphila. *Sweet.*

* Several different plants are known in Barbados by this familiar appellation.

† Ægiphila Martinicensis is doubtless the Spirit Weed of Hughes, although he says "the flowers are yellow"—a state-

elata. *Æ.* foliis elliptico-acuminatis membrana-
ceis, paniculis terminalibus, calycibus
pubescentibus. *Willd. Sp. Pl.* 1. p.
616. h

BUDDLEIA.

Scrophulariæ, *Juss.*—Scrophularinæ, *Brown.*

- occiden-* *B. foliis lanceolatis. Willd. Sp. Pl. 1.*
talis. *p. 632.* h
 Ophyoxylon americanum. Pluck. alm.
 270. Phyt. t. 210. f. 1.
 Spear-leaved Buddleia. Miller.

the genus *Manettia* is “Calyx octophyllus superus.” Browne, however, describes the calyx of *Lygistum* as “monophyllus conico-campanulatus ore quadricrenato,” and the seed vessel as “calyci imposita;” and the “twigs or boughs” as “furnished with moderate bunches of flowers towards the top.” It is, therefore, quite evident that the *Lygistum* of Browne cannot be the *Manettia* of Willdenow and Swartz: nor is it less certain that it should be assumed as a synonym to *Ægiphila elata* to the exclusion of *Knoxia scandens*. The want of accordance between Browne’s description and the plate referred to did not escape Swartz, but he did not trace the error to its source, and could not correct it. Speaking of *Ægiphila elata*, he says, “Folia ovato-cordata descripit Browne, elliptico-lanceolata tamen illa delineari curavit.”—(*Swtz. Fl. 255*).

PETITIA.

Vitices. *Juss.*—Verbenaceæ. *Juss. An. du Mus. Brown.*

domingen- *P. domingensis. Willd. Sp. Pl. 1. p. 614.*
sis. Jacq. Amer. 14. t. 182. f. 6. h

SIDERODENDRUM.

Rubiaceæ. *Juss.*—Coffeaceæ. *Kunth.*

triflorum. S. triflorum. Willd. Sp. Pl. 1. p. 612.
Hort. Kew. 1. p. 245. h

Sideroxylum americanum. Pluk. Alm.
346. Phyt. t. 224. f. 2.

Sideroxyloides ferreum. Jacq. Amer. 19.
t. 175. f. 9.

Iron-wood. *Hughes. 142.*

Three-flowered Iron Tree. *Aiton.*

CISSUS.

Hederaceæ. *Lin.*—Vites. *Juss.*—Ampelideæ. *De Cand.*

sicyoides.* *C. foliis cordatis ovatis glabris crassiusculis setaceo-serratis: serraturis appressis, ramulis teretibus. De Cand. Prod. 1. p. 628. Willd. Sp. Pl. 1. p. 656. Hort. Kew. 1. p. 259. Aubl. 77. Jacq. Amer. 22. t. 15.* h

Irsiola scandens. Br. 147. t. 4.†

Bryonia alba geniculata. Sl. 1. p. 233. t. 144. f. 1.

Poison Wyth. Hughes. 166.

Heart-leaved Cissus. Aiton.

Bastard Bryony. Browne.

Scratch-Wyth.

* “Flores sunt parvi, numerosi, inodori, in variis locis vel rubri, vel flavi, vel virides.”—*Jacq. l. c.*

† “Valde accidit” (that is, *Cissus ovata*) “ad *sicyoidem*, sed folia ovata nec cordata.”—*De Cand.* I altogether doubt whether the *C. ovata* be sufficiently distinct from *C. sicyoides*; but at all events I am pretty confident that Browne’s *Irsiola* is identical with the latter, and with Sloane’s *Bryonia*. I however hold this opinion subject to correction, for it is in opposition to M. De Candolle.

AMMANNIA.

Calycanthemæ. *Lin.*—Salicariæ. *Juss.*Lythraræ. *De Cand.*

- latifolia*.* *A. caule tetragono erecto ramoso, foliis lineari-lanceolatis basi obtuse cordato-auriculatis, floribus axillaribus 3—5 sessilibus, petalis?† staminibusque quatuor. De Cand. Prod. 3. p. 78. Willd. Sp. Pl. 1. p. 678. Hort. Kew. 1. p. 260. ☉ Isnardia foliis sessilibus. Br. 148. Aparines folio anomola. Sl. 1. p. 44. t. 7. f. 4. Broad-leaved Ammannia. Miller.*

* The plant described by Sloane and Browne, beyond doubt the species of *Ammannia* growing in Barbados, hardly deserves the epithet *latifolia*, even in reference to the other species.

† De Candolle places this species in the tetrapetalous subdivision of the genus, but questions “an potius apetala?” The flower is certainly apetalous, and greatly resembles that of *A. ægyptiaca*, represented in *Willd. Hort. Berolinensis*, from which this species is principally to be distinguished by the base of the leaf.

POTHOS.

Piperitæ. *Lin.*—Aroideæ. *Juss. Brown.*

- acaulis.* P. foliis lanceolatis integerrimis enerviis.
Willd. Sp. Pl. 1. p. 684. *Hort. Kew.* 1.
 p. 268. *Jacq. Amer.* 240. t. 153. 4
 Narrow-leaved Mistletoe. *Hughes.* 156.
 Stemless Pothos. *Aiton.*
- lanceolata.* P. foliis lanceolatis integerrimis trinerviis, scapo apice triquetro. *Willd. Sp. Pl.* 1. p. 684. *Hort. Kew.* 1. p. 268. 4
 Arum foliis rigidis angustis et acuminatis *Plum. Amer.* 47. t. 62.
 Long-leaved Mistletoe. *Hughes.* 222.
 Lance-leaved Barbados Pothos. *Aiton.*

RIVINA.

Holeraceæ. *Lin.*—Atriplices. *Juss.*—Phytolaceæ. *Brown.*

- humilis.* R. racemis simplicibus, floribus tetrandris, foliis pubescentibus. *Willd. Sp. Pl.* 1. p. 694. *Hort. Kew.* 1. p. 273.
Aubl. 90. *Bot. Mag.* t. 1781. 2

R. dichotoma erecta. *Br.* 148.

Solanum barbadense. *Pluk. Alm.* 353.

Phyt. t. 112. *f.* 2.

Cat's Blood. *Hughes.* 160.

Downy Rivina. *Miller.*

octandra. *R. racemis simplicibus, floribus octan-*
dris dodecandrisve. Willd. Sp. Pl.* 1. *p.*

695. *Hort. Kew.* 1. *p.* 273. *Aubl.* 90. $\frac{1}{2}$

R. sarmentosa. *Br.* 149. *t.* 23. *f.* 2.

Black Basket-Wyth. *Hughes.* 242.

Hoop-Wyth. *Long.* 2. *p.* 755.

Climbing Rivina. *Miller.*

Wild Hicory.

* "Stamina quatuor intra calycis folia; reliqua octo alternantia."—*Willd.* All the flowers I have examined are dodecandrous. The filaments come from eight points at the base of the germen, the alternate points bearing two filaments, as described by Browne.

TETRANDRIA TETRAGYNIA.

MYGINDA.

Dubii ord. *Lin.*—Rhamni. *Juss.*—Celastrineæ. *De Cand.*

latifolia. M. foliis ellipticis crenatis lævibus breviter petiolatis coriaceis, pedunculis trifidis paucifloris, stigmatibus 2—4 subsesilibus. *De Cand. Prod.* 2. p. 13.

Willd. Sp. Pl. 1. p. 722. *Hort. Kew.* 1. p. 283. h

Broad-leaved Myginda. *Aiton.*

PENTANDRIA MONOGYNIA.

HELIOTROPIUM.

Asperifoliæ. *Lin.*—Boragineæ. *Juss. Brown.*

indicum. H. foliis cordato-ovatis acutis scabriusculis, spicis solitariis, fructibus bifidis.
Willd. Sp. Pl. 1. p. 740. *Hort. Kew.* 1. p. 284. *Aubl.* 116. *Bot. Mag. t.* 1837. ☉

H. americanum cæruleum. *Sl.* 1. p. 213.

Tiaridium indicum. *Spreng. Syst.* 1. p. 568.

Indian Turnsole, or Heliotrope. *Miller.*
Wild Clary.

parviflorum. H. foliis ovatis rugosis scabris oppositis alternisque, spicis conjugatis. *Willd. Sp.*

Pl. 1. p. 741. *Hort. Kew.* 1. p. 284. ☉

H. barbadense. *Dill. Elth.* 178. *t.* 146. *f.* 175.

H. hirsutum læte virens. *Br.* 151.

Wild Clary. *Hughes.* 169.

Small-flowered Turnsole. *Miller.*

curassavi- H. foliis lanceolato-linearibus glabris
cum. aveniis, spicis conjugatis. *Willd. Sp. Pl.*
 1. p. 743. *Hort. Kew.* 1. p. 285. *Aubl.*
 117. *Bot. Mag.* t. 2669. ☉

H. supinum leucophæum. *Br.* 151.

H. maritimum minus. *Sl.* 1. p. 213. t.
 132. f. 3.

Wild Lavender. *Hughes.* 220.

Glancous Turnsole. *Miller.*

gnapha- H. foliis linearibus alternis tomentosis,
lodes. pedunculis dichotomis, spicarum floribus
 quaternis, caule frutescente. *Willd. Sp.*
Pl. 1. p. 745. *Aubl.* 117. *Jacq. Amer.*
 25. t. 173. f. 11. ½

H. arboreum maritimum. *Sl.* 1. p. 213.

Tournefortia gnaphalodes. *Roem. Syst.*
 4. p. 538.

Crab Bush, or Sea-side Laurel. *Hughes.*
 155.

Sea-side Lavender.

TOURNEFORTIA.

Asperifoliæ. Lin.—Boragineæ. Juss. Brown.

volubilis. T. foliis ovatis acuminatis glabris, petiolis
 reflexis, caule volubili. *Willd. Sp. Pl.* 1.
 p. 791. *Hort. Kew.* 1. p. 303. *Aubl.* 117. ½

T. fruticosa scandens. *Br.* 170.

Messerschmidia volubilis. *Roem. Syst.* 4.
p. 541.

Bryonia nigra fruticosa. *Sl.* 1. *p.* 234.
t. 143. *f.* 2.

Chigery-Bush. *Hughes.* 160.

Soldier-Bush. *Hughes.* 161.

Climbing Tournefortia. *Miller.*

bicolor. *T. foliis ovatis acuminatis glabris superne
rugosiusculis, spicis cymosis erectis race-
mosis.* *Willd. Sp. Pl.* 1. *p.* 793. *Swtz.*
Fl. 344. h

T. reclinata diffusa. *Br.* 169.

Basket-Wyth. *Long.* 2. *p.* 834.

Two-coloured Tournefortia. *Sweet.*

White Hoop.

CORDIA.

Asperifoliæ. Lin.—Boragineæ. Juss. Brown.

Myxa. *C. foliis ovalibus sub repando-dentatis
sub-coriaceis, paniculis lateralibus dif-
fuis. M—.* *Willd. Sp. Pl.* 1. *p.* 1027.
Hort. Kew. 2. *p.* 8. h

Sebestena. Myxa. Blackw. t. 398.

Sebestena officinalis. *Gaert. Fruct.* 1. p. 364. t. 76. f. 1.

Prunus Sebestena. *Pluk. Alm.* 306. *Phyt.* t. 217. f. 3.

Smooth-leaved Cordia, *Sebestena*, or African Plum. *Miller.*

Sebestena. C. foliis oblongo-ovatis repandis scabris. *Willd. Sp. Pl.* 1073. *Hort. Kew.* 2. p. 8. *Jacq. Amer.* 42. h

Sebestena scabra. *Dill. Elth.* 340. t. 255. f. 331.

α. floribus luteis. *Bot. Repos.* t. 157. Yellow Cordia.

β. floribus fulvo-aureis. *Bot. Mag.* t. 794. Orange Cordia.

γ. floribus coccineis.

C. foliis amplioribus. *Br.* 202.

Caryophyllus spurius. *Sl.* 2. p. 20. t. 164.

Scarlet Cordia.

Rough-leaved Cordia. *Miller.*

Collococca. C. foliis oblongo-ovatis integerrimis, floribus corymbosis, calycibus interne tomentosis. *Willd. Sp. Pl.* 1. p. 1075. *Hort. Kew.* 2. p. 9. *Aubl.* 219. h

Collococcus foliis rugosis. *Br.* 167.

Ceraso affinis arbor baccifera. *Sl.* 2. p. 95. t. 203. f. 2.

Clammy Cherry Tree. *Hughes.* 151.

Clammy Cherry, or Turkey-Berry Tree.

Browne.

Long-leaved Cordia. *Miller.*

VARRONIA.

Asperifoliæ. Lin.—Boragineæ. Juss. Brown.

curassa- V. foliis lanceolatis, spicis oblongis.
vica. *Willd. Sp. Pl. 1. p. 1080. Hort. Kew. 2.*
p. 11. Jacq. Amer. 40. h

V. assurgens sarmentosa. *Br. 172.*

Cordia curassavica. *Roem. Syst. 4. p. 460.*

Periclymenum rectum salviæ folio rugoso
 majore. *Sl. 2. p. 81.*

Black Sage Bush. *Hughes. 159.*

Long-spiked Varronia. *Miller.*

alba. V. foliis cordatis, floribus cymosis. *Willd.*
Sp. Pl. 1. p. 1081. Jacq. Amer. 41. h

Loblolly Tree. *Hughes. 143.*

White-fruited Varronia. *Miller.*

SPIGELIA.

Stellatæ. *Lin.*—Gentianeæ. *Juss. Brown.*

Anthelmia. S. caule herbaceo, foliis summis quaternis. *Willd. Sp. Pl.* 1. p. 824. *Hort. Kew.* 1. p. 317. *Aubl.* 126. *Bot. Mag.* t. 2359. ☉

S. quadrifolia. *Br.* 156. t. 37. f. 3.

Arapabaca quadrifolia. *Plum. Gen.* 11.

Loggerhead Weed. *Hughes.* 230.

Worm-Grass. *Long.* 2. p. 766.

Annual Worm-Grass. *Miller,*

Water-Weed.

PLUMBAGO.

Dubii ord. *Lin.*—Plumbagines. *Juss.*

Plumbagineæ. *Brown.*

rosea.

P. foliis petiolatis ovatis glabris subdenticulatis, caule geniculis gibbosis. *Willd.*

Sp. Pl. 1. p. 858. *Hort. Kew.* 1. p. 324.

Bot. Mag. t. 230. ♀

Rose-coloured Lead-Wort. *Miller.*

scandens. P. foliis petiolatis ovatis glabris, caule flexuoso scandente. *Willd. Sp. Pl.* 1. p. 858. *Hort. Kew.* 1. p. 324. *Aubl.* 136. h

P. spicis ramosis. *Br.* 158.

Dentillaria lychnoides, &c. flore albo. *Sl.* 1. p. 211. *t.* 133. *f.* 1.

Climbing Lead-Wort. *Miller.*

CONVOLVULUS.*

Campanacæ. *Lin.*—Colvolvuli. *Juss.*

Convolvulacæ. *Brown.*

hedera- C. foliis cordatis integris trilobisve,
ceus. corollis indivisis, fructibus erectis. *Willd.*
Sp. Pl. 1. p. 851. *Hort. Kew.* 1. p. 329.
Aubl. 140. ⊙

Ipomœa hederacea. *Bot. Reg.* t. 85.

Ipomœa punctata. *Roem. Syst.* 4. p. 227.

Wild Slip. *Hughes.* 172.

Ivy-leaved Bind-Weed. *Miller.*

* The generic arrangement of convolvulaceous plants is so continually being changed, that I have been content to follow the species Plantarum of Willdenow, referring to some of the principal recent alterations as synonyms.

- Nil.* C. foliis cordatis trilobis, corollis semiquinquefidis, pedunculis petiolo brevioribus. *Willd. Sp. Pl.* 1. p. 851. *Hort. Kew.* 1. p. 330. *Bot. Mag.* t. 188. ☉
Ipomœa hederacea. *Br. Prod.* 342.*
 Monkey Vine. *Hughes.* 168.
 Blue, or Azure Bind-Weed. *Miller.*
- purpureus.* C. foliis cordatis indivisis, fructibus cernuis, pedicellis incrassatis. *Willd. Sp. Pl.* 1. p. 852. *Hort. Kew.* 1. p. 330. *Bot. Mag.* t. 113. ☉
Ipomœa purpurea. *Hort. Br.* 288.
 Purple Bind-Weed. *Miller.*
- Batatas.* C. foliis cordatis hastatis (*trilobis quinquelobisque*) quinquenerviis, caule repente hispido tuberifero. *Willd. Sp. Pl.* 1. p. 853. *Hort. Kew.* 1. p. 331. *Aubl.* 140. 2
 C. edulis? † *Willd. Sp. Pl.* 1. p. 875.

* Mr. Brown considers the *Convolvulus Nil. L.* and the *Ipomœa hederacea. L.* as identical, and refers to them both as synonyms of his *Ipomœa hederacea.*

† Many varieties of the sweet Potatoe are cultivated in Barbados, propagated by the slip or by the tuberosc root. I am not certain that the *C. Batatas*, and *C. edulis*, are specifically distinct. Thunberg, describing the *C. edulis*, says “Radices magnitudine sæpe pugni, tuberculatæ, carnosæ, uti *Batatas*, esculentæ, mollissimæ et sapidissimæ.” “Differt a *C. Batatas* foliis cordatis integris trilobis et quinquelobis, nec angustatis in medio ut sagittata evadant.”—*Thunb. Jap.* 85. The leaves of the Potatoe are very

C. repens, &c. radice crassa carnosâ alba.
Br. 154.

C. repens, &c. radice crassa carnosâ lutea.
Id. ibid.

C. radice tuberosa esculenta. *Sl.* 1. p. 150.

Ipomœa Batatas. *Roem. Syst.* 4. p. 218.

Potatoes. *Hughes.* 228.

Sweet, or Bermudas Potatoe. *Long.* 2.
p. 774.

Tuberose-rooted Bind-Weed, or Spanish
Potatoe. *Miller.*

maximus. *C. foliis cordatis ovatis acuminatis integerrimis glaberrimis, caule petiolisque glabris.* *Willd. Sp. Pl.* 1. p. 853. *Hort. Kew.* 1. p. 331. h

Great Ceylon Bind-Weed. *Aiton.*

verticillatus. *C. foliis cordatis oblongis nudis, pedunculis umbellatis bifidis multifloris.* *Willd. Sp. Pl.* 1. p. 856. ⊙

C. minor scandens. *Br.* 153.

Wild Vine. *Hughes.* 242.

umbellatus. *C. foliis cordatis, caule volubili, pedunculis umbellatis.* *Willd. Sp. Pl.* 1. p. 857.
Hort. Kew. 1. p. 331. *Aubl.* 140. u

C. foliis cordato-acuminatis. *Br.* 154.

C. polyanthos. *Sl.* 1. p. 154.

various, and some agree best with those assigned to *C. edulis*. It is not improbable, however, that some of the supposed varieties of the sweet Potatoe will be found to be distinct species.

C. luteus polyanthos. *Plum. Amer.* 88.
t. 102.

Ipomœa polyanthes. *Roem. Syst.* 4. p. 234.

Hog-Slip. *Hughes.* 171. Hog-Vine.
Id. 243.

Umbelled Bind-Weed. *Miller.*

speciosus. *C. foliis cordatis subtus tomentoso-sericeis, pedunculis petiolo longioribus umbelliferis, calycibus acutis, caule volubili.*

Willd. Sp. Pl. 1. p. 859. *Hort. Kew.* 1.
p. 332. h

Ipomœa speciosa. *Roem. Syst.* 4. p. 239.

Bot. Mag. t. 2446.

Argyreia speciosa. *Hort. Br.* 289.

Broad-leaved Bind-Weed. *Miller.*

Cephalic Vine.

dissectus. *C. foliis palmatis septempartitis dentato-sinuatis glabris, caule piloso, pedunculis unifloris.* *Willd. Sp. Pl.* 1. p. 864. *Jacq.*

Vind. 2. p. 74. t. 159. u

Ipomœa sinuata. *Roem. Syst.* 4. p. 207.

Noyeau Vine.

*brasiliensis.** *C. foliis emarginatis basi biglandulosis, pedunculis trifloris.* *Willd. Sp. Pl.* 1. p.

877. *Hort. Kew.* 1. p. 338. *Aubl.* 141. u

C. maritimus. *Br.* 153.

* Mr. Brown considers the *Convolvulus braziliensis*, L. and the *C. pes carpæ*, L. as identical.—*Br. Prod.* 342.

C. marinus catharticus. *Plum. Amer.* 89.
t. 104.

Ipomœa maritima. *Roem. Syst.* 4. *p.* 249.
Bot. Reg. t. 319.

Sea-side Vine. *Hughes.* 243.

Purging Sea Bind-Weed, or Scammomy.
Long. 2. *p.* 833.

Brazilian Bind-Weed. *Miller.*

IPOMŒA

Campanacææ. *Lin.*—Convolvuli. *Juss.*

Convolvulacææ. *Brown.*

Quamoclit. *I. foliis pinnatifidis linearibus, floribus subsolitariis.* *Willd. Sp. Pl.* 1. *p.* 879.

Hort. Kew. 1. *p.* 338. *Aubl.* 144. ☉

I. foliis capillaceis pinnatis. *Br.* 155.

α. floribus coccineis. *Bot. Mag. t.* 244.

Barbabos Pink. *Hughes.* 168.

β. floribus albis.

Winged-leaved *Ipomœa.* *Miller.*

American Jessamine. *Browne.*

Indian Creeper—Indian Pink.—

coccinea. I. foliis cordatis acuminatis basi angulatis, pedunculis multifloris. *Willd. Sp. Pl.* 1. p. 880. *Hort. Kew.* 1. p. 338. *Aubl.* 144. *Bot. Mag. t.* 221. *Bot. Repos. t.* 499. ☉

Convolvulus coccineus. *Plum. Amer.* 89. t. 103.

Scarlet-flowered Ipomœa. *Miller.*

bona-nox. I. foliis cordatis acutis integerrimis (*subangulato-dentatis*) caule aculeato, floribus ternis corollis indivisis. *Willd. Sp. Pl.* 1. p. 882. *Hort. Kew.* 1. p. 339. *Bot. Mag. t.* 752. ½

I. sylvestris. *Br.* 155.

I. grandiflora. *Bot. Repos. t.* 403?

Convolvulus maximus, caule spinulis obtusis obsito. *Sl.* 1. p. 151. t. 96. f. 1.

Argyreia bona-nox. *Hort. Br.* 289.

Prickly Ipomœa. *Miller.**

Large white-flowered Ipomœa. *Browne.*

Large night-blowing *Convolvulus.*

* The aculei are properly represented in the Gardener's Dictionary as "blunt, herbaceous, short, variously shaped." They are, in fact, little more than protuberances, and afford no ground for the trivial appellation applied to the plant in that work. The epithet night-flowering, assigned by Sweet to *Argyreia bona-nox*, is certainly much more characteristic.

PSYCHOTRIA.

Stellatæ. *Lin.*—*Rubiaceæ. Juss.*—*Coffeaceæ. Kunth.*

- nervosa.* P. stipulis oblongis emarginatis deciduis, foliis ovatis utrinque acuminatis nervosis subundulatis, paniculis sessilibus erectiusculis. *Willd. Sp. Pl. 1. p. 964. Swtz. Fl. 403.* h
St. John's Bush.
- tenuifolia.* P. stipulis ovatis emarginatis (*apice bifidis*) deciduis, foliis oblongis tenuioribus glabris, paniculis erectis subsessilibus. *Willd. Sp. Pl. 1. p. 964. Swtz. Fl. 402.* h
Wild Coffee.
- horizontalis.* P. stipulis ovatis, foliis lanceolato-ovatis acutis, ramis foliis ramulisque panicularum horizontalibus. *Willd. Sp. Pl. 1. p. 965. Swtz. Fl. 410.* h

COFFEA.

Stellatæ. *Lin.*—*Rubiaceæ. Juss.*—*Coffeaceæ. Kunth.*

- arabica.* C. foliis oblongo-ovatis, pedunculis axillaribus aggregatis, corollis quinquefidis.
Willd. Sp. Pl. 1. p. 973. Hort. Kew. 1. p. 374. Aubl. 154. Bot. Mag. t. 1303. h
 C. fruticosa foliis oppositis. *Br. 161.*
 Coffee Tree. *Hughes. 158. Browne.*
 Eastern Coffee Tree. *Miller.*
- occidentalis.* C. foliis oblongo-lanceolatis acuminatis, panicula pauciflora trifida terminali, corollis quadrifidis, baccis monospermis, ramulis tetragonis. *Willd. Sp. Pl. 1. p. 974. Hort. Kew. 1. p. 374. Aubl. 154. Jacq. Amer. 67. t. 47. h*
 Pavetta foliis oblongo-ovatis. *Br. 142. t. 6. f. 2.**
 Pavetta coffeoides. *Ham. Prod. 18.*
 Jasminum forte arboreum. *Sl. 2. p. 97. t. 205. f. 1? †*

* The reference to the plate in Browne is erroneously marked fig. 1.

† Although I have followed Willdenow in giving this synonym from Sloane, I must add that it certainly does not seem to apply to *Coffea occidentalis*, nor does Browne consider Sloane's

Tetramerium odoratissimum. Spreng.

Syst. 1. p. 409.

Western Coffee Tree. Miller.

Wild Jessamine. Browne.

Wild Coffee.

CHIOCOCCA.

Aggregatæ. Lin.—Rubiaceæ. Juss.—Coffeaceæ. Kunth.

racemosa. C. foliis ovatis acuminatis, racemis subdivisis axillaribus secundis subnutantibus. Willd. *Sp. Pl.* 1. p. 975. Hort. Kew. 1. p. 374. Jacq. *Amer.* 68. Bot. Repos. t. 284. h

Jasminum folio myrtino. Sl. 2. p. 97. t. 188. f. 3.

α. C. sarmentosa. Br. 164.

β. C. scandens. Br. 164.

Briny Roots. Hughes. 225.

Climbing Snow-Berry Tree, or David's Root. Miller.

Candle Wood.

Jasminum as identical with his *Pavetta*. Sloane represents the corolla of his plant as "divided towards the edges into five points." Browne describes the corolla of his *Pavetta* as "tubus longus cylindraceus, limbus in quatuor lacinias lanceolatas patentes sectus;" from whence, in the species plantarum, the *Coffea occidentalis* is properly characterized "corollis quadrifidis."

CONOCARPUS.

Aggregatæ. *Lin.*—*Elæagni. Juss.*Combretaceæ. *De Cand. Brown.**erecta.*

C. foliis lanceolatis. Willd. Sp. Pl. 1. p. 994. Hort. Kew. 1. p. 381. Jacq. Amer. 78. t. 52. f. 1. h

C. erecta. α. arborea. De Cand. Prod. 3. p. 16.

C. foliis oblongis. Br. 159.

Manghala arbor curassavica. Catesb. Car. 2. p. 33. t. 33.

Alni fructu laurifolia arbor maritima. Sl. 2. p. 18. t. 161. f. 2.

Jamaica Button Tree. Miller.

MIRABILIS.

Dubii ord. *Lin.*—*Nyctagines. Juss. Brown.**Jalapa.*

M. floribus congestis pedunculatis, foliis glabris. Willd. Sp. Pl. 1. p. 999. Hort. Kew. 1. p. 382. u

Admirabilis peruviana. Sl. p. 211.

Solanum mexicanum. *Bauh. Pin.* 168.

α. Jalapa flore purpureo. *Tourn. Inst.* 129. *Bot. Mag. t.* 371.

Common Marvel of Peru. *Aiton.*

β. Jalapa flore flavo. *Tourn. Inst.* 129.

Yellow Marvel of Peru. *Aiton.*

γ. Jalapa flore exalbido. *Tourn. Inst.* 129.

White Marvel of Peru. *Aiton.*

δ. Jalapa flore ex purpureo et luteo mixto. *Tourn. Inst.* 129.

Yellow-striped Marvel of Peru. *Aiton.*

Jallop, or Four-o'Clock Flower. *Hughes.* 203. *Long. 2. p.* 814.

DATURA.

Luridæ. Lin.—Solaneæ. Juss. Brown.

Strammonium. *D. pericarpiis spinosis erectis ovatis, foliis ovatis (sinuatis. Engl. Fl.) glabris. Fl. Br.* 254. *Willd. Sp. Pl. 1. p.* 1008. *Hort. Kew. 1. p.* 387. *Aubl.* 212. *Engl. Bot. t.* 1288. ☉

D. foliis profunde crenatis. Br. 167.

Strammonium. Blackw. t. 313.

Solanum fætidum, pomo spinoso. Bauh. Pin. 168.

Night-Shade. *Hughes.* 211.

Common Thorn-Apple. *Miller.*

Thorn-Apple, or Burn-Weed. *Browne.*

arborea. *D. pericarpiis glabris inermibus nutantibus, caule arboreo. Willd. Sp. Pl. 1. p. 1009. Hort. Kew. 1. p. 388.* ^h

Brugmansia candida. Hort. Br. 297.

Tree Thorn-Apple. *Miller.*

NICOTIANA.

Luridæ. Lin.—Solaneæ. Juss. Brown.

Tabacum. *N. foliis lanceolato-ovatis sessilibus decurrentibus, floribus acutis. Willd. Sp. Pl. 1. p. 1014. Hort. Kew. 1. p. 394. Aubl. 212. Blackw. t. 146.* [⊙]

Virginian Tobacco. *Miller.*

PHYSALIS.

Luridæ. Lin.—Solaneæ. Juss. Brown.

angulata. *P. ramosissima, ramis angulatis glabris, foliis ovatis dentatis. Willd. Sp. Pl. 1. p. 1022. Hort. Kew. 1. p. 394. Aubl. 212.* [⊙]

Pops. *Hughes*. 161.

Tooth-leaved Winter Cherry. *Miller*.

barbaden- *P. ramosissima*, foliis ovato-cordatis pu-
sis. bescentibus, floribus pendulis, calycibus
fructiferis ovato-acuminatis angulatis.

Willd. Sp. Pl. 1. p. 1023. *Jacq. Ic.* 1.
p. 5. t. 39. ☉

Pop Vine. *Hughes*. 161.

Barbados Winter Cherry. *Miller*.

SOLANUM.

Luridæ. *Lin.*—Solaneæ. *Juss. Brown.*

tuberosum. *S. caule inermi herbaceo, foliis pinnatis*
integerrimis, pedunculis subdivisis.

Willd. Sp. Pl. 1. p. 1033. *Hort. Kew.* 1.
p. 399. *Blackw. t.* 523. a. b. 2

Irish Potatoe. *Long.* 2. p. 774.

Tuberose-rooted Nightshade, or Com-
mon Potatoe. *Miller*.

Lycoper- *S. caule inermi herbaceo, foliis pinnatis*
sicum. incisiss, racemis bipartitis aphyllis, fruc-
tibus glabris. *Willd. Sp. Pl.* 1. p. 1033.

Hort. Kew. 1. p. 399. *Aubl.* 215. ☉

Lycopersicum esculentum. *Roem. Syst.*
4. p. 568.

Lycopersicon subhirsutum. *Br.* 175.

Pomum amoris. *Blackw. t.* 133.

Tammatas Shrub. *Hughes*. 148.

Tomato. *Long*. 2. p. 773.

Love Apple, or Tomato. *Miller*.

nigrum. *S. caule inermi herbaceo, foliis ovatis dentato-angulatis, umbellis lateralibus nutantibus. Fl. Br.* 256. *Willd. Sp. Pl.* 1. p. 1035. *Hort. Kew.* 1. p. 399. *Aubl.* 215. *Engl. Bot. t.* 566. ☉

S. humilius diffusum. Br. 174.

S. Desvauxii. Ham. Prod. 26? *

Gumma Bush. *Hughes*. 164.

Branched Calalue. *Long*. 2. p. 771.

Small Black-berried Nightshade. *Miller*.

Seafortthianum. S. caule inermi fruticoso scandente, foliis ternatis simplicibus lanceolato-ovatis acuminatis undulatis, racemis paniculatis, pedicellis incurvis. Bot. Mag. t. 1982. *Bot. Reg. t.* 969. *Bot. Repos.* t. 504. ♀

* I would not be understood to insist that the *Solanum Desvauxii* is not a distinct species, only that its characters have not as yet been sufficiently marked to separate it specifically from *Solanum nigrum*. They are both described with unarmed, herbaceous stems, and small, round, black berries; the only mark of distinction pointed out being in the leaves, and that by no means a strong one. In *Solanum Desvauxii. Ham.* these are "ovato-lanceolatis integerrimis subrepandis;" in *S. nigrum. Willd.* "ovatis dentato-angulatis." The plant I have seen in Barbados I cannot but consider as *S. nigrum. Willd.* *S. humilius diffusum. Br.* and the Gumma Bush. *Hughes*.

Seaforth's Nightshade. *Sweet.*

St. Vincent Lilac.

*Melon-
gena.*

S. caule inermi herbaceo, foliis ovatis tomentosis, pedunculis incrassatis, calycibus inermibus (vel aculeatis. M.—)

Willd. Sp. Pl. 1. p. 1036. Hort. Kew. 1. p. 400. Aubl. 215. ©

*S. insanum. Willd. Sp. Pl. 1. p. 1037.**

S. hirsutum et spinosum. Br. 173.

S. esculentum. Roem. Syst. 4. p. 638.

Montabba, *alias* Trubba. *Hughes. 147.*

Brown Jolly, Valinghanna, or Mad Apple.

Long. 2. p. 772.

Large-fruited Nightshade, or Egg Plant.

Miller.

torvum. S. caule aculeato fruticoso, aculeis curvis, foliis subcordatis ovatis sinuatis tomen-

* I entirely adopt the opinion of Loureiro, that the species *Melongena* and *insanum* are not specifically distinct. In the definitions given by Willdenow the aculeated stem and calyx of *S. insanum* is the only distinguishing character. It cannot however be admitted as such. In the Gardener's Dictionary the *Melongena* is represented as having a calyx "armed with very strong prickles on the outside;" and one variety as also having "the stalk and leaves armed with very strong thorns." Mr. Miller adds, that in the same garden, and from the same seeds, he has raised plants, some with and some without prickles, a fact confirmed by Loureiro—"Solanum insanum non est specie diversum a *Melongena*; sequidem aculeos illius, per quos videbatur distingui, ego sæpe observavi in ista ex iisdem seminibus et in eodem horto nata."—*Lour. Fl. Cochin. 130.*

toxis, rachibus aculeatis,* calycibus inermibus. *Willd. Sp. Pl. 1. p. 1038. Swtz. Fl. 456.* h

Small Red Trubba. *Hughes. 148.*

Curved Prickled Nightshade. *Sweet.*

mammosum. *S. caule aculeato herbaceo, foliis cordatis angulato-lobatis utrinque villosis aculeatis. Willd. Sp. Pl. 1. p. 1040. Hort. Kew. 1 p. 401. Aubl. 216.* ⊙

S. barbadense spinosum. Pluk. Alm. 350. Phyt. t. 226. f. 1.

S. pomiferum tomentosum. Sl. 1. p. 38. t. 12. f. 1.

Bachelor's Pears. *Hughes. 139.*

Yellow-spined Nightshade. *Miller.*

xantocarpum. *S. caule decumbente diffuso aculeato, foliis pinnatifidis aculeatis stellato-pubescentibus: laciniis sinuatis acutis margine nudis, calycibus aculeatis. Willd. Sp. Pl. 1. p. 1041.* ⊙

Bachelor's Pears. *Hughes. 139.*

Yellow-fruited Nightshade. *Miller.*

stramonifolium. *S. caule aculeato fruticoso, foliis cordatis angulato-lobatis integris subinermibus subtus tomentosiusculis. Willd. Sp. Pl. 1. p. 1044. Hort. Kew. 1. p. 403. Jacq. Ic. 1. p. 5. t. 44.* h

* "Variat foliis inermibus."—*Swtz.*

Broad-leaved Nightshade. *Miller.*

bahamense. *S. caule aculeato fruticoso, foliis lanceo-*
latis repandis obtusis margine reflexis,
racemis simplicibus. Willd. Sp. Pl. 1. p.
1046. Hort. Kew. 1. p. 404. h

S. fruticosum bacciferum. Sl. 1. p. 236.*
t. 145. f. 3.

Bahama Nightshade. *Miller.*

igneum. *S. caule aculeato fruticoso, foliis lanceo-*
latis acuminatis basi utrinque revolutis,
racemis simplicibus. Willd. Sp. Pl. 1.
p. 1048. Hort. Kew. 1. p. 405. Aubl.
216. h

S. bacciferum fruticosum. Sl. 1. p. 38.
t. 11. f. 3.

S. spiniferum frutescens. Pluk. Alm.
350. t. 225. f. 5.

Berry Bush, or Hedge Bush. *Hughes.*

Red-spined Nightshade. *Miller.*

* It is this *Solanum fruticosum bacciferum*, and not *S. bacciferum fruticosum*, *Sl. 1. p. 38.* which is the synonym to *S. bahamense*. The *Solanum* of which Sloane, *p. 38.* says "the prickles are very red," is obviously the Berry Bush of Hughes.—The *Solanum igneum. Willd.* By comparing *t. 11.* with *t. 145.* the difference will readily be perceived,

CAPSICUM.

Luridæ. *Lin.*—Solaneæ. *Juss.*

- annuum.* C. fructibus oblongis pendulis ovatisque glabris, caule herbaceo. *Roem. Syst.* 4. p. 559. ☉
 Bell Pepper. *Hughes.* 213.
 Annual Capsicum. *Sweet.*
- longum.* C. fructibus oblongo-acuminatis subincurvis, petiolis glabris, caule herbaceo. *Roem. Syst.* 4. p. 560. ☉
 Long Pepper. *Hughes.* 213.
 Long-fruited Capsicum. *Sweet.*
- angulosum.* C. caule herbaceo, fructu cordiformi anguloso. *Roem. Syst.* 4. p. 561. ☉
 Bonnet-Pepper (*ribbed*). *Hughes.* 213.
 Angular-fruited Capsicum. *Sweet.*
- tetragonum.* C. caule herbaceo, fructu maximo anguloso obtuso. *Roem. Syst.* 4. p. 561. ☉
 Bonnet-Pepper (*plain*). *Hughes.* 213.
 Four-sided Capsicum. *Sweet.*
- grossum.** C. fructibus oblongo-ovatis subcompressis erectis deflexisque, petiolis glabris,

* “Fructus relatione ad plantam maximus, mali fere magnitudine, varia in variis individuis forma.”—*Willd.* 1. p. 1051.

caule herbaceo vel potius perennante.

Roem. Syst. 4 p. 562. 2

C. fructu maximo. *Br. 176.*

Turbilo Pepper. *Hughes. 213 ?*

Bell-Pepper. *Browne.*

Various-fruited Capsicum. *Sweet.*

conoides. C. caule suffrutescente, fructibus ovato-conicis surrectis. *Roem. Syst. 4. p. 562.* 2

C. fructu minimo conico rubro. *Br. 176.*

C. minus fructu parvo pyramidali. *Sl. 1. p. 240 t. 146. f. 2.* 2

Negro-Pepper. *Hughes. 213.*

Bird Pepper.* *Browne.*

Conical-fruited Capsicum. *Sweet.*

cerasi-forme. C. fructibus globosis, petiolis glabris, caule fruticoso. *Roem. Syst. 4. p. 563.* 2

Cherry-Pepper. *Hughes. 213.*

Cherry-fruited Capsicum. *Sweet.*

frutescens. C. fructibus oblongis obtusis, petiolis glabris, caule erecto. *Roem. Syst. 4. p. 563.* 2

Spur-Pepper. *Hughes. 213.*

* The Bird Pepper of Jamaica is a different species from the Bird Pepper of Barbados. The Jamaica name for this species of Capsicum prevails very much in England, and is sometimes changed for Bird's-Eye Pepper. Mr. Sweet has, however, in the *Hort. Brit.* properly applied it to *C. baccatum*; but he has made an erroneous reference to Sloane for a plate of the Bird Pepper, having been misled, probably by the authority of Willdenow.

Shrubby Capsicum. *Sweet.*

bicolor. C. fructibus oblongis mucronatis, petiolis glabris, caule fruticoso. *Roem. Syst. 4. p. 563. Bot. Mag. t. 1835.* h

Purple-flowered Capsicum. *Sweet.*

baccatum. C. pedunculis geminis solitariisque, fructibus globoso-ovatis erectis, petiolis glabris, caule frutescente. *Roem. Syst. 4. p. 564.* h

C. minus fructu rotundo. *Sl. 1. p. 240.*

Bird-Pepper. *Hughes. 213. Sweet.*

sinense. C. fructibus ovatis pendulis, petiolis pubescentibus, caule frutescente. *Roem. Syst. 4. p. 564.* h

Olive-Pepper. *Hughes. 213.*

Oval Chinese Capsicum. *Sweet.*

CESTRUM.

Luridæ. *Lin.*—Solaneæ. *Juss. Brown.*

laurifolium. C. filamentis denticulatis nudisve,* foliis ellipticis coriaceis nitidissimis, pedunculis petiolo brevioribus. *Willd. Sp. Pl. 1. p. 1054. Hort. Kew. 2. p. 1.* h

* Of all the flowers I have examined, the filaments were denticulated. Some few had six segments of the corolla, and six stamens.

Laureola sempervirens americana. *Pluk.*

Alm. 209. *Phyt. t.* 95. *f.* 1.

Wild Calabash. *Hughes.* 160.

Laurel-leaved Cestrum. *Miller.*

noctur-
num.

*C. filamentis edentatis, corollæ tubo fili-
formi: laciniis acuminatis, floribus sub-
racemosis, pedunculis axillaribus brevis-
simis, foliis ovatis acuminatis. M.—*

Willd. Sp. Pl. 1. *p.* 1055. *Hort. Kew.*

2. *p.* 1.

h

Jasminoides, &c. flore virescente noctu
odoratissimo. *Dill. Elth* 183. *t.* 153. *f.* 185.

Night-smelling Cestrum. *Miller.*

Lady of the Night.*

* It is impossible to doubt the identity of the Lady of the Night of Barbados with the *Cestrum nocturnum*; yet the definition given by Willdenow, Aiton, and others, of *C. vespertinum*, *i. e.* “*C. filamentis edentatis, tubo filiformi, pedunculis brevissimis*,” is more descriptive of this “night-smelling Cestrum” than that affixed to *C. nocturnum*, which represents it “*pedunculis subracemosis folio æquantibus*.” Martyn, between his respect for authority and his personal experience, has fallen into a strange inconsistency. He follows the definition of Linnæus, “peduncles subracemed, *equal* to the leaf;” yet he describes the leaves and inflorescence as he actually saw them: the leaves as “near four inches long,” and the flowers as “standing on short peduncles.” Those peduncles cannot surely be short, which are equal to a leaf near four inches long.—See *Gard. Dict. sub verb.*

ARDISIA.

Myrsineæ. *Brown.*

coriacea. A. floribus paniculatis, foliis oblongis
integrifoliis coriaceis. *Willd. Sp. Pl.*
1. p. 1062. *Swtz. Fl.* 470. h

JACQUINIA.

Dubii ord. *Lin.*—Sapotæ. *Juss.*—Myrsineæ. *Brown.*

armillaris. J. foliis cuneiformibus, ramis ad ramifi-
cationes nodosis verticillatis. *Willd. Sp.*
Pl. 1. p. 1064. *Hort. Kew.* 2. p. 5.
Jacq. Amer. 53. t. 39. h
Xyloceste fruticosum. *Br.* 372.
Arbor baccifera laurifolia. *Sl.* 2. p. 89.
t. 190. f. 2.
Pye-Crust. *Hughes.* 147.
Obtuse-leaved Jacquinia. *Miller.*

CHRYSOPHYLLUM.

Dumosæ. *Lin.*—Sapotæ. *Juss.*—Sapoteæ. *Brown.*

- Cainito.* C. foliis ovatis parallele striatis subtus tomentoso-nitidis. *Willd. Sp. Pl.* 1. p. 1083. *Hort. Kew.* 2. p. 12. *Aubl.* 234. $\frac{1}{2}$
 α . C. *Cainito.* *Jacq. Amer.* 51. t. 37. f. 1.
 β . C. *jamaicense.* *Jacq. Amer.* 52.*
 C fructu majori globoso. *Br.* 171. t. 14. f. 2.
Anona, &c. fructu rotundo majori. *Sl.* 2. p. 176. f. 229.
 γ . C. *cæruleum.* *Jacq. Amer.* 52. t. 37.
 Broad-leaved Star-Apple. *Miller.*
- monopyrenum.* C. foliis oblongis acuminatis subtus tomentoso-aureis, fructu ovato monopyreno.† *Willd. Sp. Pl.* 1. p. 1083. *Swtz. Fl.* 480. $\frac{1}{2}$
 C. fructu minori glabro. *Br.* 171.

* It is this variety which is generally brought to the desert in Barbados; but I have seen the others.

† Fructus qui jamaicensibus dicitur Damson-Plum constanter monospermus invenitur.—*Swtz. Fl.* 481. This fruit is generally single-seeded, but it occurs with two seeds.

Damson-Plum. *Browne.*

Star-Plum.

glabrum. C. foliis ovato-oblongis acutis utrinque
glabris, fructu elliptico glabro. *Willd.*
Sp. Pl. 1. p. 1084. h

C. foliis utrinque glaberrimis. *Jacq.*
Amer. 53. t. 38. f. 2.

Damacen, or Callimato Tree. *Hughes.*
179.

Smooth Star-Apple. *Miller.*

BUMELIA.

Dumosæ. *Lin.*—Sapotæ. *Juss.*—Sapoteæ. *Brown.*

nigra. B. foliis terminalibus oblongo-lanceolatis
glabris margine undulatis, ramis laxis,
ramulis virgatis floriferis. *Willd. Sp. Pl.*
1. p. 1085. *Swtz. Fl.* 487. h

Achras fructibus minoribus. *Br.* 201.

Bully-Berry Tree. *Hughes.* 148. *Id.* 177.

Black Bumelia. *Miller.*

Bastard Bully Tree. *Browne.*

Bully Tree.

salicifolia.* B. foliis lanceolato-ovatis acuminatis, pedunculis confertis axillaribus et lateralibus. *Willd. Sp. Pl.* 1. p. 1086. *Swtz. Fl.* 491. . h

Achras salicifolia. *Lin. Sp. pl.* 470.

Achras foliis oblongis. *Br.* 201. t. 17. f. 4.

Salicis folio lato splendente arbor. *Sl.* 2. p. 98. t. 206. f. 2.

White Bully Tree, or Galimeta-Wood.
Long. 2. p. 748.

Willow-leaved Bumelia. *Miller.*

* I have been obliged to exclude as synonyms to *B. salicifolia* the *Cornus* t. 75. of Catesby, and Jacquin's *Sideroxylum Mastichodendron*, which are unadvisedly admitted by Willdenow. Brown describes the berries of his *Achras* 201. t. 17. f. 4. which is the *B. salicifolia* of Willdenow, as "black, smooth, and very small." Catesby and Jacquin represent the fruit of their tree as yellow, "and of the size of a nutmeg." It is, therefore, impossible that their tree should be identical with Browne's. Swartz says nothing of the colour of the fruit of *B. salicifolia*, but simply, "drupa parva, oblonga, extus succosa, sæpe disperma." The synonyms in question have been removed to *Sideroxylon Mastichodendron* of Sprengel's *Systema*, to which they seem unequivocally to belong.

SIDEROXYLON.

Dumosæ. *Lin.*—Sapotæ. *Juss.*—Sapoteæ. *Brown.*

Masticho- S. foliis oblongo-lanceolatis utrinque at-
*dendron.** tenuatis sempervirentibus supra nitidis
 subtus parallele venosis, pedunculis axil-
 laribus confertis abbreviatis calycibusque
 splendido-tomentosis. *Spreng. Syst.* 1.
p. 666. *Jacq. Coll.* 2. *p.* 253. *t.* 17.
f. 5. h

Cornus foliis laurinis. *Catesb. Car.* *p.*
 75. *t.* 75.

Prunifera vel nucifera, sen potius nuci-
 prunifera. *Sl.* 1. *p.* 40. *Pluk. Alm.* 307.
Phyt. t. 217. *f.* 5.

Mastick Tree. *Hughes.* 149.

* This species of Sideroxylon, the hard wood represented by Sloane, Ligon, Hughes, and others, under the name of Mastick, is confounded by Willdenow with Bumelia salicifolia. This valuable tree, which was once so common in Barbados, is now very rarely to be seen.

TECTONA.

Vitices. *Juss.*—Verbenaceæ. *Juss. An. du Mus. Brown.*

grandis. T. *grandis.* *Willd. Sp. Pl.* 1. p. 1088.
Hort. Kew. 2. p. 12. *Roxb. Cor.* 1. p.
 10. t. 6. h

Jatus. *Rumph. Amb.* 3. p. 34. t. 18.

Indian Oak, or Teak-Wood. *Aiton.*

CEANOTHUS.

Dumosæ. *Lin.*—Rhamni. *Juss.*—Rhamneæ. *De Cand. Brown.*

colubrinus. C. foliis ovatis subacuminatis integris
 subtus ramulis floribusque ferrugineo-
 villosis, floribus axillaribus corymboso-
 aggregatis. *De Cand. Prod.* 2. p. 31. h
Rhamnus colubrinus. *Willd. Sp. Pl.* 1.
 p. 1096. *Hort. Kew.* 2. p. 15. *Jacq.*
Amer. 74.

Rhamnus arboreus, foliis ovatis venosis.
Br. 172.

Bahama Red-Wood. *Sweet.*

Black Bead Tree.

ZIZYPHUS.

Dumosæ. *Lin.*—Rhamni. *Juss.*—Rhamneæ. *De Cand.*

- Jujuba.* *Z.* foliis subrotundo-ovatis obtusis subintegris subtus petiolis ramulisque tomento brevissimo canis, aculeis subsolitariis recurvis, corymbis axillaribus tomentosis.
De Cand. Prod. 2. p. 21. *Willd. Sp. Pl.* 1. p. 1104. *Hort. Kew.* 2. p. 18. ½
Jujuba indica spinosa. *Pluk. Alm.* 199.
Rhamnus Jujuba. *Lin. Sp. Pl.* 282.
 Dunk Tree, or Mangustine. *Hughes.* 134.
 Blunt-leaved Zizyphus. *Aiton.*

CEDRELA.

Miscellaneæ. *Lin.*—Meliis aff. *Juss.*—Meliaceæ. *De Cand.*

- odorata.* *C.* calycibus corollisque nudis. *Roem. Syst.* 5. p. 464. *De Cand. Prod.* 1. p. 624. *Willd. Sp. Pl.* 1. p. 1147. *Hort. Kew.* 2. p. 38. *Aubl.* 246. ½
C. foliis pinnatis, ligno odorato. *Br.* 158.
t. 10. *f.* 1.

Cedrus barbadensium. *Pluk. Alm.* 92.

Phyt. t. 157. *f.* 1.

Pruno forte affinis arbor. *Sl.* 2. *p.* 128.

t. 220. *f.* 2.

Cedar. *Hughes.* 115.

Barbadoes Cedar. *Long.* 2. *p.* 835.

Barbadoes Bastard Cedar. *Miller.*

MANGIFERA.

Dubii ord. *Lin.*—Terebinthaceæ. *Juss. De Cand.*

indica. *M. foliis oblongo-lanceolatis petiolatis, paniculis erectis, petalis apice patulis, stamine unico fertili, drupa subreniformi glabra.* *De Cand. Prod.* 2. *p.* 63. *Willd.*

Sp. Pl. 1. *p.* 1150. *Hort. Kew.* 2. *p.* 39.

Jacq. Ic. 2. *t.* 337. *Bot. Repos. t.* 425. *h*

Manga domestica. *Rumph. Amb.* 1. *p.* 93. *t.* 25.

Persicæ similis. *Bauh. Pin.* 440.

East India Mango Tree. *Hughes.* 177.

Mango Tree. *Aiton.*

BALSAMINA.

Corydales (*Impatiens*) *Lin.*—*Gerania. Juss.*Balsamineæ. *De Cand.*

hortensis.* *B. pedicellis aggregatis, foliis lanceolatis; superioribus alternis, calcaribus flore brevioribus. De Cand. Prod. 1. p. 685. ☉ Impatiens Balsamina. Willd. Sp. Pl. 1176. Hort. Kew. 2. p. 49. Garden Balsam. Miller.*

VITIS

Hederaceæ. *Lin.*—*Vites Juss.*—*Ampelideæ. De Cand.*

vinifera. *V. foliis lobatis sinuato-dentatis nudis aut tomentosis. De Cand. Prod. 1. p. 633. Willd. Sp. Pl. 1. p. 1180. Hort. Kew. 2. p. 51. h*
α. vulgaris.
 Common Grape.

* “Variat in hortis flore pleno albo, roseo, rubro, et ex his colore vario.”—*De Cand.*

β. apyrena.

Corynth, or Currant Grape.

Common Vine. *Aiton.*

CELOSIA.

Miscellaneæ. *Lin.*—Amaranthi. *Juss.*—Amaranthaceæ. *Brown.*

argentea. C. foliis lanceolatis, stipulis subfalcatis, pedunculis angulatis, spicis scariosis.

Willd. Sp. Pl. 1. p. 1197. Hort. Kew.

2. p. 58.

⊙

Silvery-spiked Celosia. *Miller.*

cristata. C. foliis oblongo-ovatis, pedunculis teretibus substriatis, spicis oblongis. *Willd.*

Sp. Pl. 1. p. 1198. Hort. Kew. 2. p. 58.

⊙

Crested Amaranth, or Cock's Comb.

Miller.

coccinea. C. foliis ovatis strictis inauriculatis, caule sulcato, spicis multiplicibus cristatis.

Willd. Sp. Pl. 1. p. 1199. Hort. Kew.

2. p. 58.

⊙

Scarlet Celosia, or Chinese Cock's Comb.

Miller.

GARDENIA.

Contortæ. *Lin*—*Rubiaceæ*. *Juss.* *Brown.*

- florida.* *G. inermis*, foliis ellipticis, corollis hypocrateriformibus, laciniis calycinis verticalibus lanceolato-subulatis. *Willd. Sp. Pl.* 1. p. 1225. *Hort. Kew.* 1. p. 368. h
 α . floribus simplicibus. *Bot. Reg.* t. 449.
 β . floribus plenis. *Bot. Mag.* t. 2627.
 Fragrant Gardenia, or Cape Jasmine.
Miller.
- armata.* *G. spinosa*, spinis ramulorum terminalibus quaternis, laciniis calycinis lineari-cuneiformibus, floribus aggregatis. *Willd. Sp. Pl.* 1. p. 1229. *Swtz. Fl.* 524. h
Mussænda spinosa. *Jacq. Amer.* 70. t. 49.
 Armed Gardenia. *Sweet.*
- Randia.* *G. spinosa*, spinis ramulorum terminalibus binis, foliis ovato-subrotundis, corollis hypocrateriformibus. *Willd. Sp. Pl.* 1. p. 1230. *Hort. Kew.* 1. p. 370.
Swtz. Fl. 526. *Bot. Mag.* t. 1841. h
Randia aculeata. *Roem. Syst.* 5. p. 246.
Randia foliis subrotundis. *Br.* 143. t. 8. f. 1.

Lycium forte foliis subrotundis. *Sl.* 1.

p. 40. *t.* 11. *f.* 4.

Indigo-Berry. *Long.* 2. *p.* 736.

Round-leaved Gardenia. *Miller.*

Dog-Wood.

CERBERA.

Contortæ. *Lin.*—Apocynæ. *Juss.* *Brown.*

Thevetia. *C.* foliis linearibus longissimis confertis.

Willd. Sp. Pl. 1. *p.* 1223. *Hort. Kew.* 2.

p. 66. *Jacq. Amer.* 48. *t.* 34. *Bot.*

Mag. t. 2309. h

Plumeria arborescens, foliis lanceolatis.

Br. 181.

Nerio affinis, &c. flore luteo. *Pluk. Alm.*

268. *Phyt. t.* 207. *f.* 3.

French Willow. *Hughes.* 133. *t.* 8.

Linear-leaved Cerbera. *Miller.*

ALLAMANDA.

Contortæ. *Lin.*—Apocynæ. *Juss.* *Brown.*

cathartica. *A. cathartica.* *Willd. Sp. Pl.* 1. *p.* 1231.

Hort. Kew. 2. *p.* 66. *Bot. Mag. t.* 338. h

Orelia grandiflora. *Aubl.* 271. *t.* 106.

Willow-leaved Allamanda. *Aiton.*

VINCA.

Contortæ. *Lin.*—Apocynæ. *Juss.* *Brown.*

rosea.

V. caule suffrutescente erecto, floribus geminis sessilibus, foliis ovato-oblongis, petiolis basi bidentatis. *Willd. Sp. Pl.* 1. *p.*

1233. *Hort. Kew.* 2. *p.* 66. h

α . flore carneo. *Bot. Mag.* *t.* 248.

Madagascar Red Periwinkle. *Aiton.*

β . flore albo, unbone carneo.

Madagascar White Periwinkle. *Aiton.*

Old Maid, or Cayenne Jasmine.

NERIUM.

Contortæ. *Lin.*—Apocynæ. *Juss.* *Brown.*

odorum.

N. foliis lineari-lanceolatis ternis subtus costatis, laciniis calycinis erectis, nectariis multipartitis: laciniis filiformibus.

Willd. Sp. Pl. 1. *p.* 1235. *Hort. Kew.* 2. *p.* 67. h

N. foliis lanceolatis. *Br.* 181.

N. Oleander. *Lour. Cochin.* 151.

α. floribus simplicibus.

Single-flowered sweet-scented Rose-Bay, or Oleander.

β. floribus plenis. *Bot. Reg. t.* 74. *Bot.*

Mag. t. 1799. *β.*

Full-flowered sweet-scented Rose-Bay, or Oleander.

South-Sea Rose. *Hughes.* 218.

*coronari-
um.*

N. foliis ellipticis, pedunculis ex dichotomia ramorum geminis bifloris. *Willd.*

Sp. Pl. 1. *p.* 1236.

h

Tabernæmontana coronaria. *Hort. Kew.* 2. *p.* 72.

α. floribus simplicibus.

Tabernæmontana coronaria. *Bot. Reg. t.* 1064.

β. floribus plenis.

N. coronarium. *Bot. Mag. t.* 1865.

Broad-leaved Rose-Bay. *Miller.*

PLUMERIA.

Contortæ, *Lin.*—Apocynææ, *Juss. Brown.*

rubra.

P. foliis ovato-oblongis, petiolis biglandulosus. Willd. Sp. Pl. 1. p. 1242. Hort. Kew. 2. p. 70. Aubl. 258. Jacq. Amer. 35. Bot. Mag. t. 279. Bot. Reg. t. 780. h

P. arborescens. Br. 181. No. 2.

P. flore roseo odoratissimo. Catesb. Car. 1. p. 92. t. 92.

Nerium arboreum, folio maximo. Sl. 2. p. 61. t. 185—186.

Tree Jasmine. Hughes. 189. t. 17.

Jasmin Tree. Browne.

Red Plumeria. Miller.

alba.

P. foliis lanceolatis revolutis, pedunculis superne tuberosis. Willd. Sp. Pl. 1. p. 1242. Hort. Kew. 2. p. 70. Aubl. 259. Jacq. Amer. 36. t. 174. f. 12. h

P. arborescens. Br. 181. No. 3.

Nerium arboreum, &c. folio angusto. Sl. 2. p. 62.

White-flowered Jasmin Tree. Browne.

White Plumeria. Miller.

PENTANDRIA DIGYNIA.

CALOTROPIS.

Contortæ (*Asclepias*). *Lin.*—Apocynæ (*Asclepias*). *Juss.*
Asclepiadeæ. Brown.

- procera.* C. corollæ laciniis patulis. *Hort. Kew.* 2.
p. 78. *Roem. Syst.* 6. p. 90. h
Asclepias procera. Willd. Sp. Pl. 1.
p. 1263.
Asclepias frutescens incana. Br. 182.
Apocynum erectum majus. Pluk. Alm.
35. *Phyt. t.* 175. f. 3.
Bell-flowered Gigantic Swallow-Wort.
Miller.
- gigantea.* C. corollæ laciniis reflexis involutis.
Hort. Kew. 2. p. 78. *Roem. Syst.* 6.
p. 91. h
Asclepias gigantea. Willd. Sp. Pl. 1.
p. 1264.
Curl-flowered Gigantic Swallow-Wort.
Miller.

HOYA.

Contortæ (*Asclepias*). *Lin.*—Apocynæ (*Asclepias*). *Juss.*

Asclepiadæ. *Brown.*

- carnosa.* H. foliis ovali-oblongis carnosis, corollis
barbatis, coronæ foliolis subtus sulcatis.
Brown. Prod. 316. *Hort. Kew.* 2. p. 84.
Roem. Syst. 6. p. 50. h
Asclepias carnos. *Willd. Sp. Pl.* 1. p.
1264. *Bot. Mag. t.* 788. *Ex. Bot. t.* 70.
Fleshy-leaved Hoya. *Sweet.*

ASCLEPIAS.

Contortæ. *Lin.*—Apocynæ. *Juss.*—Asclepiadæ. *Brown.*

- curassa-* A. foliis lanceolatis petiolatis glabris
vica. nitidis, caule simplici, umbellis erectis
solitariis lateralibus. *Willd. Sp. Pl.* 1.
p. 1266. *Hort. Kew.* 2. p. 81. *Bot.*
Reg. t. 81. h
A. erecta, foliis angustis. *Br.* 183.
Apocynum erectum, folio oblongo. *Sl.* 1.
p. 206. t. 129. f. 4, 5.

Indian Root. *Hughes*. 209.

Wild, or Bastard Ipecacuanha. *Browne*.

Curassoa Swallow-Wort. Bastard Ipecacuanha. *Miller*.

CYNANCHUM.

Contortæ. *Lin.*—Apocynæ. *Juss.*—Asclepiadæ. *Brown*.

mucronatum. C. foliis ovali-oblongis mucronatis membranaceis subpubescentibus, umbellis axillaribus. *M.*— *Bot. Repos. t. 515.* $\frac{1}{2}$ *Asclepias mucronata. Willd. Sp. Pl. 1. p. 1263. Thumb. Jap. 47.*
Sharp-pointed Cynanchum. *Sweet.*

SARCOSTEMMA.

Contortæ (*Asclepias*). *Lin.*—Apocynæ (*Asclepias*). *Juss.*
Asclepiadæ. *Brown*.

Swartzia- S. caule suffrutescente volubili filiformi,
num. foliis oppositis lanceolatis glabris, umbellis lateralibus multifloris. *Roem. Syst. 6. p. 115.* $\frac{1}{2}$

Asclepias viminalis Willd. *Sp. Pl.* 1. p. 1270. *Swtz. Fl.* 539.

Asclepias funiculacea. Br. 183.

Apocynum fruticosum scandens. Sl. 1. p. 207. t. 131. f. 1.

Down Vine. Hughes. 214.

Swartz's *Sarcostemma.* Sweet.

CHENOPODIUM.

Holeraceæ. Lin.—Atriplices. Juss.—Chenopodeæ. Brown.

anthelmin- C. foliis ovato-oblongis dentatis, racemis
ticum. aphyllis. Willd. *Sp. Pl.* 1. p. 1304. *Hort.*
Kew. 2. p. 100. h

Worm-seed Weed. Hughes. 170.

Worm-seed Goosefoot. Miller.

BETA.

Holeraceæ. Lin.—Atriplices. Juss.—Chenopodeæ. Brown.

vulgaris. B. floribus congestis, foliis inferioribus
ovatis. Willd. *Sp. Pl.* 1. p. 1308. *Hort.*
Kew. 2. p. 101. δ

α. viridis.

Green Beet. *Sweet.*

β. rubra.

Red Beet. *Sweet.*

γ. lutea.

Yellow Beet. *Sweet.*

δ. macrorhiza.

Mangel-Wurzel. *Sweet.*

GOMPHRENA.

Miscellaneæ. *Lin.*—*Amaranthi. Juss.*

Amaranthaceæ. Brown.

globosa. G. caule erecto, foliis ovato-lanceolatis,
capitulis solitariis, pedunculis diphyllis.
Willd. Sp. Pl. 1. p. 1321. . Hort. Kew.
2. p. 105. Aubl. 280. ☉

G. erecta hirsuta. *Br. 184.*

α. floribus purpureis.

Amaranthoides, &c. capitulis purpureis. *Tourn. Inst. 654.*

Purple Bachelor's Button.

β. floribus albis.

Amaranthoides, &c. capitulis argenteis.
Tourn. Inst. 654.

White Bachelor's Button.

Bachelor's Button, or Everlasting.

Hughes. 172.

Annual Globe Amaranth. *Miller.*

brasiliensis.

G. caule erecto, foliis ovato-oblongis, capitulis pedunculatis globosis aphyllis.

Willd. Sp. Pl. 1. 1322. *Hort. Kew.*

2 p. 106. *Jacq. Ic.* 2. t. 346. ☉

G. brasiliana. *Aubl.* 281.

Philoxerus brasiliensis. *Roem. Syst.* 5.

p. 542.

Brasilian Globe Amaranth. *Miller.*

DAUCUS.

Umbellatæ. *Lin.*—Umbelliferæ. *Juss.*

Carota.

D. fructibus hispidis, petiolis subtus nervosis. *Fl. Br.* 300. *Willd. Sp. Pl.* 1.

p. 1389. *Hort. Kew.* 2. p. 128. *Engl.*

Bot. t. 1174. ♂

α. radice lutea.

Pastinaca radice lutea. *Bauh. Pin.* 151.

Yellow Garden Carrot. *Aiton.*

β. radice atrorubente.

Pastinaca radice atrorubente. *Bauh.*

Pin. 151.

Red Garden Carrot. *Aiton.*

ERYNGIUM.

Umbellatæ. *Lin.*—Umbelliferæ. *Juss.*

fœtidum. E. foliis radicalibus lanceolatis serratis;
floralibus multifidis, caule dichotomo.

Willd. Sp. Pl. 1. p. 1356. *Hort. Kew.*

2. p. 115. *Aubl.* 284. *Br.* 155. 4

E. foliis angustis serratis. *Sl.* 1. p. 264.
t. 156. *f.* 3. 4.

Fit-Weed. *Hughes.* 229.

Stinking Eryngo, or Fit Weed. *Long.*
2. p. 846.

Stinking Eryngo. *Miller.*

ANETHUM.

Umbellatæ. *Lin.*—Umbelliferæ. *Juss.*

graveo- A. fructibus compressis. *Willd. Sp. Pl.*
lens. 1. p. 1469. *Hort. Kew.* 2. p. 158. ☉

A. hortense. *Bauh. Pin.* 147.

Common Dill. *Miller.*

- Fœniculum.* A. fructibus ovatis. *Willd. Sp. Pl.* 1. p. 1469. *Hort. Kew.* 2. p. 159. *Fl. Br.* 329. *Engl. Bot. t.* 1208. 2
Fœniculum dulce. *Bauh. Pin.* 147.
Fœniculum vulgare. *Hort. Br.* 187.
 Fennel, or Finckle. *Miller.*

APIUM.

Umbellatæ. *Lin.*—Umbelliferæ. *Juss.*

- Petroselinon.* A. foliolis caulinis linearibus, involucellis minutis. *Willd. Sp. Pl.* 1. p. 1475. *Hort. Kew.* 2. p. 161. ♂
 α. A. hortense, *Petroselinum* vulgo. *Tourn. Inst.* 305.
 Common Parsley. *Aiton.*
 β. A. vel *Petroselinum crispum.* *Tourn. Inst.* 305.
 Curled Parsley. *Aiton.*
graveolens. A. foliis caulinis cuneiformibus. *Willd. Sp. Pl.* 1. p. 1475. *Hort. Kew.* 2. p. 161. *Engl. Bot. t.* 1210. ♂
 A. dulce, *Celleri Italorum.* *Tourn. Inst.* 305.
 Celery. *Aiton.*

PENTANDRIA TRIGYNIA.

XYLOPHYLLA.

Tricoccæ. (*Phyllanthus*) *Lin.*—Euphorbiæ. *Juss.*

Euphorbiaceæ. *De Cand.*

- falcata.* X. foliis sparsis lineari-lanceolatis subfalcatis crenatis, floribus subsessilibus triandris monoicis. *Willd. Sp. Pl.* 1. p. 1501.
Hort. Kew. 2. p. 336. *Swtz. Prod.* 28.
Bot. Reg. t. 373. *Bot. Repos.* t. 331. ½
Phyllanthus falcata. *Swtz. Fl.* 1115.
Filicifolia hemionitidi affinis. *Pluk. Alm.* 154. *Phyt.* t. 247. f. 4.
Sea-side Laurel. *Hughes.* 233. t. 22.
Sickle-leaved Sea-side Laurel. *Miller.*

BASELLA.

Holeraceæ. *Lin.*—Atriplices. *Juss.*

Chenopodeæ. *Brown.*

cordifolia. B. foliis cordato-subrotundis, pedunculis simplicibus folio brevioribus. *Willd. Sp.*

Pl. 1. p. 1514.

Heart-leaved Malabar Nightshade. *Sweet.*
Calalue.

PENTANDRIA TETRAGYNIA.

EVOLVULUS.

Campanceæ. *Lin.*—Convolvuli. *Juss.*

Convolvulaceæ. *Brown.*

nummularis. E. foliis subrotundis, caule repente, floribus subsessilibus. *Willd. Sp. Pl.* 1. p. 1516. ☉

Convolvulus herbaceus repens. *Br.* 153.

Convolvulus minor repens. *Sl.* 1. p. 157. t. 99. f. 2.

Small Creeping Convolvulus. *Browne.*

Round-leaved Evolvulus. *Sweet.*

HEXANDRIA MONOGYNIA.

AGAVE.

Coronariæ *Lin.*—Bromeliæ. *Juss.*—Bromeliaceæ. *Brown.*

americana. A. acaulis, foliis dentato-spinosis, scapo ramoso, tubo corollæ medio angustato, staminibus corolla longioribus, stylo staminibus longiore. *Willd. Sp. Pl.* 2. p. 192. *Hort. Kew.* 2. p. 301. *Bot. Repos. t.* 438. h

α. foliorum margine concolore.

A. foliis subcompressis. *Br.* 199.

May-Pole. *Hughes.* 223.

Great American Aloes, or Caratœ.

Long. 2. p. 710.

Common American Agave. *Miller.*

Silk-Grass.

β. foliorum margine luteo.*

* I am much inclined to think that this is a distinct species; it however stands as a variety in Willdenow, and I recently saw

vivipara.* *A. acaulis*, foliis dentatis (*dentato-spinosis*) scapo ramoso, tubo corollæ medio angustato, staminibus corollam æquantibus. *Willd. Sp. Pl. 2. p. 193. Hort. Kew. 2 p. 302.* 24

Silk-Grass. *Hughes. 224.†*

Viviparous or Childing Agave. *Miller.*

BROMELIA.

Coronariæ. *Lin.*—Bromeliæ. *Juss.*—Bromeliaceæ. *De Cand.*

Ananas. *B. foliis ciliato-spinosis mucronatis, spica comosa. Willd. Sp. Pl. 2. p. 7. Hort. Kew. 2. p. 200. Aubl. 301.* 24

it marked as *Agave americana* in the Botanic Garden at Geneva, which is under the immediate superintendence of M. De Candolle.

* In the *Agave americana* the seeds vegetate, and the young plants grow to be several inches long before they are separated from the parent stem, so that it is as strictly viviparous as this species, which has been distinguished specifically by the appellation *vivipara*. The *A. americana* is very generally called Silk-Grass in Barbados, and is the species principally planted as hedges: the *A. vivipara* being mostly found in gullies, and of accidental growth.

† Not the Silk-Grass of Jamaica *vid.—Long. 2. p. 814.*

- Anassa. *Rumph. Amb.* 5 p. 227. t. 81.
 Pine-Apple. *Hughes.* 230. t. 21.*
 Common Pine-Apple. *Sweet.*
- Karatas.* B. foliis erectis, floribus acaulibus sessilibus aggregatis. *Willd. Sp. Pl.* 2. p. 8.
Hort. Kew. 3. p. 200. *Jacq. Amer.* 90.
Jacq. Vind. 1. p. 11. t. 31—32. $\frac{1}{2}$
 Pen-Gwyn. *Hughes.* 232.
 Upright-leaved Pine-Apple. *Sweet.*
- Pinguin.* B. foliis ciliato-spinosis mucronatis, racemo terminali. *Willd. Sp. Pl.* 2. p. 8.
Hort. Kew. 2. p. 200. *Aubl.* 302
Jacq. Amer. 91. $\frac{1}{2}$
 B. caule assurgente. *Br.* 193.
Pinguin. *Long.* 2. p. 738.
 Large Wild Barren Pine. *Hughes.* 231.
 Broad-leaved Pine-Apple. *Sweet.*
- panicul-
gera.* B. foliis serrato-spinosis, bracteis lanceo-
 latis, racemo composito, racemulis sub-
 divisis, floribus pedunculatis. *Willd. Sp.*
Pl. 2. p. 9. *Swtz. Fl.* 585. $\frac{1}{2}$
 Small Barren Pine. *Hughes.* 231.
 Panicked Pine-Apple. *Sweet.*

* The Wild Pine and the Button Pine (*Hughes.* 231.) are doubtless only varieties of *Bromelia Ananas*.

PITCAIRNIA.

Coronariæ. *Lin.*—Bromeliæ. *Juss.*—Bromeliaceæ. *De Cand.*

*bromeliæ-
folia.* P. foliis ciliato-spinosis, pedunculis germinibusque glaberrimis. *Willd. Sp. Pl.* 2. p. 10. *Hort. Kew.* 2. p. 201. *Swtz. Fl.* 580. *Bot. Mag. t.* 824. *Bot. Reg. t.* 1011. h

Hepetis angustifolia. *Swtz. Prod.* 56.

Scarlet Pitcairnia. *Miller.*

SANSEVERIA.

Hemerocallideæ. *Brown.*

spicata. S. angustior, foliis subundecim subrectis prærigidis fragilibus lanceolato-ensiformibus, fasciis valde obsoletis. *Haw. Succ.* 66. 4

S. zeylanica. *Rox. Cor.* 2. p. 43. t. 184.

Salmia spicata. *Cav. Ic.* 3. p. 24. t. 264.

Spiked Sanseveria. *Sweet.*

TRADESCANTIA.

Ensatae. *Lin.*—Junci. *Juss.*—Commelineae. *Brown.*

Zanonia. T. erecta, foliis lato-lanceolatis, pedunculis lateralibus solitariis medio geniculatis multifloris, bracteis geminis. *Willd.*

Sp. Pl. 2. p. 17. *Hort. Kew.* 2. p. 205.

Swtz. Fl. 604. 4

Zanonia graminea perfoliata. *Plum. Gen.* 38.

Campelia Zanonia. *Kunth. Syn.* 1. p. 272.

Commelina Zanonia. *Aubl.* 37.

Commelina erecta major. *Br.* 125.

Periclymenum rectum herbaceum. *Sl.* 1. p. 243.

Gentian-leaved Spider's-Wort. *Aiton.*

discolor. T. acaulis lævis, bracteis equitantibus compressis, foliis lanceolatis subtus coloratis. *Willd. Sp. Pl.* 2. p. 18. *Hort. Kew.*

2. p. 205. *Swtz. Fl.* 607. *Bot. Mag.* t. 1192. 4

Purple-leaved Spider's-Wort. *Aiton.*

geniculata. T. procumbens hirsuta. *Willd. Sp. Pl.* 2. p. 19. *Hort. Kew.* 2. p. 205. *Jacq. Amer.* 94. t. 64. 4

Knotted Spider's-Wort. *Aiton.*

PONTEDERIA.

Ensatæ. *Lin.*—Narcissi. *Juss.*—Pontidereæ. *Kunth.*

- azurea.* P. foliis subcordatis ellipticis basi petio-
lisque incrassatis, floribus spicatis. *Willd.*
Sp. Pl. 2. p. 28. *Swtz.* 609. 2
P. aquatica caulescens. *Br.* 195.
Blue-flower'd Pontederia. *Miller.*
Round-leaved Water Plantain. *Browne.*
Water Lily.

PANCRATIUM.

Spathaceæ. *Lin.*—Narcissi. *Juss.*—Amaryllideæ. *Brown.*

- fragrans.* P. spatha multiflora, foliis ellipticis, nec-
tarii dentibus sex staminiferis: sinubus
integræ aut varie dentatis. *M.*— *Willd.*
Sp. Pl. 2. p. 43. *Hort. Kew.* 2. p. 219.
Bot. Cab. t. 834. *Salisb. Lin. Tr.* 2. p.
72. t. 11. 2

P. caribæum.* *Willd. Sp. Pl. 2. p. 42.*

Aubl. 303. Bot. Mag. t 826.

P. foliis compressis obtusis. Br. 194.

Narcissus totus albus. Sl. 1. p. 244.

White Lily. Hughes. 228. Browne.

Fragrant Pancratium. Aiton.

AMARYLLIS.

Spathaceæ. Lin.—Narcissi. Juss.—Amaryllideæ. Brown.

*equestris. A. spatha subbiflora, pedicellis erectis
sparsis brevioribus, tubo filiformi hori-*

* The definition of *Pancratium caribæum* in the species *Plantarum* is peculiarly imperfect; and I question its claims to be received as a species distinct from *P. fragrans*. The Jamaica plant, mentioned by Browne and Sloane, is doubtless identical with the Barbados plant, so carefully examined, and so fully described, by Mr. Salisbury, in the *Transactions of the Linnæan Society*. Contrary however to the terms of Mr. Salisbury's definition, the sinus of the nectary is, according to my observation, generally furnished with teeth, which are frequently bifid. The Editor of the *Botanical Magazine* observes, "little stress in specific distinction can be laid on the sinus of the crown being entire or toothleted, as we have frequently seen them vary infinitely in the same plant." In accordance with this statement, and my own observation, I have taken upon me to alter Mr. Salisbury's definition.

zontali, limbo oblique patulo sursum
curvo, fauce piloso. *Willd. Sp. Pl.* 2. p.
53. *Hort. Kew.* 2. p. 224. *Bot. Mag.*
t. 305. *Bot. Reg. t.* 234. 4

A. brasiliensis. *Bot. Repos. t.* 358.

Red Lily. *Hughes.* 229.

Barbados Lily. *Aiton.*

tubispa- *A. spatha monophylla tubulosa bifida*
tha. uniflora, pedunculo spatha duplo lon-
giore. *Willd. Sp. Pl.* 2. p. 51. *Bot. Mag.*
t. 1586. 4

Zephyranthes tubispatha. *Hort. Br.* 401.

Barbados Snow-Drop.

zeylanica. *A. spatha multiflora, corollis reclinatis,*
tubo filiformi longissimo, laciniis unci-
natis. *Willd. Sp. Pl.* 2. p. 56. 4

A. ornata. β. *Hort. Kew.* 2. p. 226.
Bot. Mag. t. 1171.

Crinum latifolium. *Bot. Repos. t.* 478.

Tulipa Javana. *Rumph. Amb.* 5. p. 306.
t. 105.

Wild Tulip. *Hughes.* 229.*

Crimson and White Amaryllis. *Aiton.*

* The description Hughes gives of the Wild Tulip sufficiently identifies it with the second variety of the *Amaryllis ornata* of the Hortus Kewensis. I have made repeated and diligent search for the *Amaryllis Belladonna*, the *habitat* of which is said to be in Barbados, but I have never been able to find it.

CRINUM.

Spathaceæ. *Lin.*—Narcissi. *Juss.*—Amaryllideæ. *Brown.*

- america-* C. foliis oblongo-lanceolatis margine
num. glaberrimis apice coarctato-unguiculatis,
floribus pedicellatis limbo brevioribus.
Willd. Sp. Pl. 2. p. 46. Hort. Kew. 2.
p. 221. Aubl. 303. Bot. Mag. t. 1034. ♀
American Crinum. *Aiton.*

ALLIUM.

Spathaceæ. *Lin.*—Asphodeli. *Juss.*—Asphodeleæ. *Brown.*

- Porrum.* A. caule planifolio umbellifero, stamini-
bus tricuspidatis, radice tunicata. *Willd.*
Sp. Pl. 2. p. 64. Hort. Kew. 2. p.
232. ♂
Common Leek. *Miller.*
- sativum.* A. caule planifolio bulbifero, bulbo com-
posito, staminibus tricuspidatis. *Willd.*
Sp. Pl. 2. p. 68. Hort. Kew. 2. p.
234. ♂
Common Garlick. *Miller.*

- Scorodo-
prasum.* A. caule planifolio bulbifero, foliis cre-
nulatis vaginis ancipitibus, staminibus
tricuspidatis. *Willd. Sp. Pl. 2. p. 69.*
Hort. Kew. 2. p. 230. ½
Rocambole. *Miller.*
- Ascalo-
nium.* A. scapo nudo tereti, foliis subulatis, um-
bella globosa, staminibus tricuspidatis.
Willd. Sp. Pl. 2. p. 75. Hort. Kew. 2.
p. 236. ¼
α. vulgaris.
Shallot. *Sweet.*
β. chinense.
Chinese Shallot. *Sweet.*
γ. Majus.
Scallion. *Sweet.*
- Cepa.* A. scapo nudo inferne ventricosso longi-
ore, foliis teretibus. *Willd. Sp. Pl. 2.*
p. 80. Hort. Kew. 2. p. 238. ¼
Common Onion. *Miller.*

ASPARAGUS.

Sarmentaceæ. *Lin.*—Asparagi. *Juss.*—Asphodeleæ. *Brown.*

- officinalis* A. caule herbaceo tereti inermi sube-
recto, foliis setaceis mollibus, stipulis

subsolitariis. *Fl. Br.* 369. *Willd. Sp. Pl.* 2. p. 150. *Hort. Kew.* 2. p. 273. *Engl. Bot. t.* 339. 2

Common Asparagus, or Sperage. *Miller.*

DRACÆNA.

Coronariæ (*Aletris*). *Lin.*—Asparagi (*Dianella*). *Juss.*

Asphodeliæ. *Brown.*

ferrea. D. arborea, foliis lanceolatis acutis.
Willd. Sp. Pl. 2. p. 157. *Hort. Kew.* 2. p. 277 2

D. terminalis. *Jacq. Ic.* 2. t. 448.

Purple Dracæna. *Aiton.*

Dragon's Blood.

fragans. D. arborea, foliis lanceolatis reflexis, paniculæ ramis terminatis capitulo florum.
Hort. Kew. 2. p. 277. *Bot. Mag. t.* 1081. 2

Aletris fragrans. *Willd. Sp. Pl.* 2. p. 183.
Bot. Repos. t. 306.

Sweet-scented Dracæna. *Aiton*

POLIANTHES.*

Coronariæ. *Lin.*—Narcissi. *Juss.*—Hemerocallideæ. *Brown.*

- tuberosa.* P. foliis linearibus scapo brevioribus.
Willd. Sp. Pl. 2. p. 164. Hort. Kew. 2.
p. 281. 2
 α . floribus simplicibus. *Bot. Reg. t. 63.*
 Common Polianthes. *Sweet.*
 β . floribus plenis. *Bot. Mag. t. 1817.*
 Double-flowered Polianthes. *Sweet.*
 Tuberose. *Hughes. 172. Miller.*

ALETRIS.

Coronariæ. *Lin.*—Asphodeli. *Juss.*—Hemerocallideæ. *Brown.*

- farinosa.* A. acaulis, foliis lanceolatis membra-
 naceis, floribus alternis. *Willd. Sp. Pl. 2.*
p. 183. Hort. Kew. 2. p. 290. Bot.
Mag. t. 1418. 2
 Common Colic-Root. *Sweet.*

* There is a difference of opinion as to the correct manner of spelling this generic title, arising from a doubt respecting its etymology. Authority is strongly in favour of the orthography above adopted.

ALOE.

Coronariæ. *Lin.*—Asphodeli. *Juss.*—Hemerocallideæ. *Brown.*

vulgaris. A. foliis patulis ascendentibus margine spinosis, pedunculo ramoso, ramis bractea duplici amplexis. *Hort. Kew. 2. p. 292.* h

A. sinuata. *Willd. Sp. Pl. 2. p. 187.*

A. barbadensis. *Haw. Lin. Tr. 7. p. 19.*

Aloes Plant. *Hughes. 153.*

Yellow-flowered Aloe. *Aiton.*

YUCCA.

Coronariæ. *Lin.*—Lilia. *Juss.*—Tulipaceæ. *Kunth.*

Draconis. Y. foliis crenatis nutantibus. *Willd. Sp. Pl. 2. p. 184. Hort. Kew. 2. p. 291.* h

Drooping-leaved Adam's Needle. *Miller.*

Wild Pine. Spanish Needles.

THRINAX.

Palmæ. *Lin. Juss. Brown.*

parviflora. T. *parviflora.* *Willd. Sp. Pl. 2. p. 202.*

Hort. Kew. 2. p. 307. Swtz. Fl. 614. h

Corypha palmacea. *Br. 190.*

Palma brasiliensis. *Sl. 2. p. 121. t. 213. f. 2.*

Palmetto Royal. *Long. 2. p. 741. Miller.*

Fan-Palm.

ACHRAS.

Dumosæ. *Lin.*—Sapotæ. *Juss.*—Sapoteæ. *Brown.*

mammosa. A. *floribus solitariis, foliis cuneiformi-lanceolatis.* *Willd. Sp. Pl. 2. p. 224.*

Hort. Kew. 2. p. 312. Aubl. 307. h

A. *fructu maximo ovato.* *Br. 201.*

Lucuma mammosum. *Hort. Br. 269.*

Mammee Sapota. *Miller.*

Mammee Sapote Tree. *Browne.*

- Sapota.* A. floribus solitariis, foliis lanceolato-ovatis. *Willd. Sp. Pl.* 2. p. 224. *Hort. Kew.* 2. p. 312. *Aubl.* 307. $\frac{1}{2}$
- α . A. Zapota. *Jacq. Amer.* 57. t. 41.
A. fructu elliptico scabro majore. *Br.* 200. t. 19. f. 3.
Sapota fructu ovato majore. *Plum. Gen.* 43.
Sapodella Tree. *Browne.*
- β . A. Zapotilla. *Jacq. Amer.* 57. t. 41.
A. brachiatus diffusus, fructu subrotundo. *Br.* 200.
Sapota fructu turbinato minore. *Plum. Gen.* 43.
Nisberry Tree. *Browne.**
Sappadilla Tree. *Hughes.* 132.
Common Sapota. *Miller.*

* The second variety is the *Achras* No. 2, or Nisberry Tree, of Browne. Mr. Aiton confounds the Sapodella Tree with the Nisberry Bully Tree, the latter of which is the *Achras* No. 3. of Browne. The Sapodella fruit is in great estimation; but, like the medlar, it is not eaten until it is over ripe. "Fructus edulis solummodo est, quando jam putrescere coepit."—*Jacq. Amer.* 58.

BAMBUSA.

Gramina (*arundo*). *Lin.*—Gramineæ. *Juss. Brown.*

arundi- B. panicula* ramosa divaricata. *Willd.*
nacea. *Sp. Pl.* 2. p. 245. *Hort. Kew.* 2. p. 316. $\frac{1}{2}$
 Bambos arundinacea. *Roxb. Cor.* 1. p.
 56. t. 79.

Arundo Bambos. *Sp. Pl.* 120.

Bambu, or Bamboo Cane. *Miller.*

* The Bamboo flowered for the first time in Barbados about seven or eight years since, having been introduced into the island nearly half a century previously, and extensively propagated by cuttings of the stem. The year was generally dry, and the period of flowering particularly so; but the plants which flowered were in an elevated situation, not far from Mount Hilaby, and in a cool and moist spot, termed Pory's Spring. The plants, although of considerable growth, were by no means the oldest in the island, the first planted roots being at an estate situate in the valley.

HEXANDRIA POLYGYNIA.

ALISMA.

Tripetaloideæ. *Lin.*—Junci. *Juss.*—Alismaceæ. *Brown.*

cordifolia. A. foliis cordatis obtusis, floribus dodecandris, seminibus uncinato-muricatis.

Willd. Sp. Pl. 2. p. 277.

24

A. foliis lunato-sagittatis. *Br. 204.*

Great Water-Plantain. *Browne.*

Pond-Weed.*

* Not the Pond-Weed of Hughes. 171. and 240.

HEPTANDRIA MONOGYNIA

PISONIA.

Dubii ord. *Lin.*—Nyctagines. *Juss.*

Nyctagineæ. *De Cand.*

aculeata. P. spinis axillaribus horizontalibus, foliis ovatis utrinque attenuatis, corymbis axillaribus. *Willd. Sp. Pl. 2. p. 283. Hort. Kew. 2. p. 333. Jacq. Amer. 274. Aubl. 948.* h

P. assurgens. *Br. 358.*

Paliuro affinis arbor spinosa. *Sl. 2. p. 25. t. 167.*

Black Thorn. *Hughes. 222.*

Fingrigo, or Cockspur. *Long. 2. p. 755.*

Prickly Pisonia. *Miller.*

nigricans. P. inermis; foliis ovatis acuminatis, floribus cymosis erectis, fructibus baccatis. *Willd. Sp. Pl. 2. p. 283. Hort. Kew. 2. p. 334. Swtz. Fl. 643.* h

P. inermis. *Jacq. Amer.* 275.

Beef-wood Tree. *Hughes.* 146.

Black Pisonia. *Aiton.*

PETIVERIA.

Holeraceæ. *Lin.*—Atriplices. *Juss.*—Phytolaceæ. *Brown.*

alliacea. *P. aliacea.* *Willd. Sp. Pl.* 2. p. 284. *Hort.*
Kew. 2. p. 334. h

α. floribus hexandris.

P. aliacea. *Hort. Br.* 338. *Bot. Cab.* 148.

P. foliis oblongo-ovatis. *Br.* 274.

Verbenæ aut Scorodoniæ affinis. *Sl.*
1. p. 172.

Common Guinea-Hen Weed. *Aiton.*

β. floribus octandris.

P. octandra. *Jacq. Amer.* 101. *Aubl.*
323. Hort. Br. 338.

P. solani foliis. *Plum. Ic.* 213. *t.* 219.

Dwarf Guinea-Hen Weed. *Aiton.*

Gully Root. *Hughes.* 205.

OCTANDRIA MONOGYNIA.

MELICOCCA.

Sapindi. *Juss.*—Sapindaceæ. *De Cand.*

bijuga.* M. foliis bijugis, rachi subulata, racemis simplicibus spiciformibus, floribus tetrapetalis, drupis ovatis. *De Cand. Prod.* 1. p. 615. *Willd. Sp. Pl.* 2. p. 330. *Hort. Kew.* 2. p. 350. ½

Melicoccus bijugatus. *Jacq. Amer.* 108. t. 72.

Melicoccus foliis ut plurimum bijugatis. *Br.* 210.

Wing-leaved Melicocca, or Honey Berry. *Aiton.*

* This is the Genip Tree of Jamaica, but must not be confounded with the Genip Tree of America, which is a Gardenia.

DODONÆA.

Terebinthaceis aff. *Juss.*—Sapindaceæ. *De Cand.*

- viscosa.* D. foliis obovato-oblongis basi cuneatis viscosis, floribus racemosis, fructibus 2—3—alatis pedicello longioribus. *De Cand. Prod.* 1. p. 616. *Willd. Sp. Pl.* 2. p. 343. *Hort. Kew.* 2. p. 358. *Jacq. Amer.* 109. h
 Aceri vel Paliuro affinis. *Sl.* 2. p. 27. *t.* 162. *f.* 3.
 Ptelea folio simplici. *Plum. Ic.* 245. *t.* 247. *f.* 2.
 Hop-Shrub. *Hughes.* 168.
 Clammy Dodonæa. *Sweet.*

LAWSONIA.

Dubii ord. *Lin.*—Salicariæ. *Juss.*—Lythrarieæ. *De Cand.*

- alba.* L. foliis oppositis ovali-lanceolatis integerrimis, floribus paniculatis. *De Cand. Prod.* 3. p. 91. h

L. inermis. *Willd. Sp. Pl. 2. p. 344.*

Hort. Kew. 2. p. 354.

*L. spinosa.** *Willd. Sp. Pl. 2. p. 345.*

Hort. Kew. 2. p. 354.

Smooth Lawsonia. Prickly Lawsonia.

Sweet.

AMYRIS.

Dubii ord. *Lin.*—Terebinthaceis. *Juss. De Cand.*

toxifera. *A. foliolis 5—7 petiolatis ovatis subcordatis acuminatis, racemis simplicibus petioli circiter longitudine. De Cand.*

Prod. 2. p. 81.

h

A. balsamifera. Willd. Sp. Pl. 2. p. 338.

A. arboreus. Br. 208.

Gum-Elemi Tree. *Hughes. 224.*

Sweet-Wood, or Shrubby Sweet-Wood.

Long. 2. p. 750.

White Candle-Wood, or Rose-Wood.

Miller.

* “Junior inermis, vetusta ramis induratis spinescit.”—*De Cand.*

OCTANDRIA TRIGYNIA.

COCCOLOBA.

Holeraceæ. *Lin.*—Polygoneæ. *Juss. Brown.*

uvifera. C. foliis cordato-subrotundis nitidis
Willd. Sp. Pl. 2. p. 457. Hort. Kew. 2.
p. 421. Jacq. Amer. 112. t. 73. h

Coccolobis foliis crassis. *Br. 209.*

Uvifera litorea. *Pluk. Alm. 394. Phyt.*
t. 236. f. 7.

Prunus maritima racemosa. *Sl. 2. p. 129.*
t. 220. f. 3. Catesb. Car. 2. p. 96. t. 96.

Bay Grape Tree. *Hughes. 180.*

Seaside Grape. *Long. 2. p. 790.*

Round-leaved Seaside Grape, or Man-
grove Grape. *Miller.*

pubescens. C. foliis orbiculatis pubescentibus. *Willd.*
Sp. Pl. 2. p. 457. Hort. Kew. 2. p.
421. h

- C. grandifolia*.* *Jacq. Amer.* 113.
Coccolobis arborea. *Br.* 210.
Scortea arbor americana. *Pluk. Phyt. t.*
 222. *f.* 8.
 Leather-Coat Tree. *Hughes.* 151.
 Great-leaved Seaside Grape. *Miller.*
nivea. *C. foliis oblongis acuminatis basi attenu-*
atis, racemis erectiusculis. *Willd. Sp.*
Pl. 2. p. 458. *Jacq. Amer.* 115. *t.* 78.
Swtz. Fl. 693. h
Coccolobis foliis oblongo-ovatis. *Br.*
 210.†
 Chigery-Grape Tree. *Hughes.* 179.
 Chequered Grape Tree. *Browne.*
barbaden- *C. foliis cordato-ovatis undulatis.* *Willd.*
sis. *Sp. Pl. 2. p.* 459. *Hort. Kew. 2. p.*
 422. *Jacq. Obs. 1. p.* 18. *t.* 8. h
 Rod-Wood. *Hughes.* 144.‡

* “*Folia diametro bipedali.*” The wood very hard. “*Per secula enim incorrupta perstat, et tandem in lapidem permutatur durissimum ista illius pars quæ sub terra recondita fuit.*” *Jacq.*

† I cannot persuade myself that Willdenow is not in error in making the *Coccolobis*, No. 3. of Browne, different from the *Coccoloba nivea* of Jacquin, which is certainly the tree commonly called Chigery-Grape, in Barbados. I have, therefore, referred to the *Coccolobis*, No. 3. of Browne, as a synonym to the *Coccoloba nivea* of Jacquin, and Willdenow.

‡ I cannot identify the Red-Wood of Hughes. 143. with any tree in Barbados. His Rod-wood. 144. is unquestionably the *Coccoloba barbadensis*, which commonly goes by the name of

Barbados Seaside Grape. *Aiton.*
Red Wood.

PAULLINIA.

Trihilatæ. Lin.—Sapindi. Juss.—Sapindaceæ. Brown.

Cururu. *P. capsulis pyriformibus obtusis, foliis ternatis: foliolis oblongis dentato-serratis subacuminatis, petiolis alatis. Willd. Sp. Pl. 2. p. 460. De Cand. Prod. 1. p. 604. Jacq. Obs. 3. p. 11. t. 61. f. 4. ½*
P. foliis ternatis. Plum. Ic. 102. t. 111. f. 2.
Cururu scandens triphylla. Plum. Gen. 34.

Bread and Cheese, or Sucking-Bottle.
Hughes. 139.

Supple Jack.

barbaden- *P. valvulis capsularum semiovatis villosi-*
sis. *usculis, foliis biternatis: foliolis ovalibus integerrimis serratisque coriaceis, petiolis submarginatis. Willd. Sp. Pl. 2. p. 461. De Cand. Prod. 1. p. 605. Hort. Kew. 2. p. 423. Jacq. Obs. 3. p. 12. t. 62. f. 9. ½*

Red-Wood amongst persons frequenting the woods. The young vigorous shoots of this tree present a very different appearance, especially as regards the leaf, to that of the older branches.

Barbados Paullinia. *Miller.*
 Supple Jack.*

CARDIOSPERMUM.

Trihilatæ. *Lin.*—Sapindi. *Juss.*—Sapindaceæ. *De Cand.*

Halicacabum. C. caule petiolis foliisque glabris, foliis
 biternatim sectis; segmentis petiolatis in-
 ciso-dentatis. *De Cand. Prod.* 1. p. 601.
Willd. Sp. Pl. 2. p. 467. *Hort. Kew.*
 2. p. 424 *Aubl.* 355. *Bot. Mag.*
 1049. ☉

C. scandens. *Br.* 213.

Pisum vesicarium. *Sl.* 1. p. 238.

Wild Parsley. *Hughes.* 212.

Parsley-leaved Heart-Pea, or Heart-Seed.
Miller.

* Neither this nor the preceding is the real Supple Jack, but
 Paullinia polyphylla. *Willd.* Serjana triternatea. *De Cand.*

SAPINDUS.

Trihilatæ. *Lin.*—Sapindi. *Juss.*—Sapindaceæ. *De Cand.*

saponaria. *S. rachi foliorum decurrente late alata, foliolis integerrimis lanceolatis tri-vel quadrijugis: terminalibus longe acuminatis. De Cand. Prod. 1. p. 607. Willd. Sp. Pl. 2. p. 468. Hort. Kew. 2. p. 468. Aubl. 359.* h

S. foliis oblongis. Br. 206.

Prunifera racemosa. Sl. 2. p. 131.

Black Nicker Tree. *Hughes. 118.**

Soap-Tree, or Soap-Berry. *Long. 2. p. 823.*

Common Soap-Berry Tree. *Miller.*

* This cannot be confounded with the Soap-Berry-Bush, or Fire-Burn-Leaf, of Hughes, 149. which I have not been able to identify with any plant I have met with.

OCTANDRIA TETRAGYNIA.

BRYOPHYLLUM.

Crassulaceæ. *De Cand.*

calycinum. B. calycinum. *De Cand. Prod.* 3. p. 396.

Bot. Mag. t. 1409.* h

Cotyledon pinnata. *Hort. Kew.* 3. p. 111.

Large-Cupped Bryophyllum. *Sweet.*

Germinating-Leaf Plant.

* This plant is placed, surely improperly, by the Editor of the Botanical Magazine, in the order Monogynia.

ENNEANDRIA MONOGYNIA.

PERSEA.

Holeraceæ (*Laurus*). *Lin.*—Lauri (*Laurus*). *Juss.*

Laurinæ. *Brown.*

*gratis-
sima.*

*P. foliis elliptico-oblongis obtusiusculis
subtus pubescentibus glaucescentibus,
corymbis axillaribus, calycibus tomen-
tosis, fructu pyriformi. Spreng. Syst. 2.
p. 268.*

h

Laurus Persea. Willd. Sp. Pl. 2. p. 480.

Hort. Kew. 2. p. 429. Aubl. 364.

Laurus foliis oblongo-ovatis. Br. 214.

*Prunifera arbor fructu maximo. Sl. 2.
p. 132. t. 222. f. 2.*

Avigato Pear Tree. Hughes. 130.

Avocado Pear. Long. 2. p. 808.

Alligator Pear. Miller.

LAURUS.

Holeraceæ. *Lin.*—Lauri. *Juss.*—Laurinæ. *Brown.*

- Cinnamomum.* L. foliis trinerviis ovato-oblongis nervis versus apicem evanescentibus. *Willd. Sp. Pl.* 2. p. 477. *Hort. Kew.* 2. p. 427. *Aubl.* 362. *Jacq. Amer.* 117. *Bot. Repos.* t. 596. h
- L. Cassia. *Bot. Mag.* t. 1636?
- Cinnamomum.* *Blackw.* t. 354.
- Cinnamomum verum.* Hort. Br.* 344.
- Cinnamomum*; sive *Canella zeylanica.* *Bauh. Pin.* 408.
- Cinnamon Tree. *Miller.*
- carolinensis.* L. foliis perennantibus ovali-lanceolatis subtus subglaucis, pedunculis simplicibus fasciculo subcapitato paucifloro terminatis, calycis laciniis exterioribus duplo brevioribus. *Mich. Amer.* 1. p. 245. h

* I will not be certain that the sickly exotic, now and then found in Barbados, under the name, Cinnamon Tree, is really the plant producing the true cinnamon. The similitude between the *L. Cinnamomum* and *L. Cassia* is so great, that they have by many been considered as varieties, although it seems to be at present settled by good authority that they are distinct species.

L. borbonica. *Willd. Sp. Pl. 2. p. 481?*

Hort. Kew. 2. p. 429?

L. foliis acuminatis. *Catesb. Car. 1. p. 63. t. 63.*

Sweet-Wood Tree. *Hughes. 157.*

Broad-leaved Carolina Bay—Red Bay.
Miller.

Timber Sweet-Wood.

parviflora. *L. foliis oblongo-ellipticis acuminatis planis membranaceis, racemis terminalibus axillaribusve folio brevioribus erectis confertis subpaniculatis.* *Willd.*

Sp. Pl. 2. p. 483. Swtz. Fl. 717. h

L. foliis oblongo-ovatis alternis venosis.
Br. 213?

L. folio longiore. *Sl. 2. p. 21. t. 165?*

Sweet-Wood Tree. *Hughes. 157.**

Water Sweet-Wood.

* Hughes represents the Sweet-Wood Tree as “of two sorts,” one affording “the best and most solid timber,” the other “of less value;” and doubtless alluded to the two species of *Laurus* above described.

ANACARDIUM.

Holeraceæ *Lin.*—Terebinthaceæ. *Juss. De Cand.*

- occidentale.* A. foliis ovalibus obtusissimis subemarginatis basi vix subattenuatis latitudine paulo longioribus. *De Cand. Prod.* 2. p. 62. *Willd. Sp. Pl.* 2. p. 486. *Hort. Kew.* 2. p. 430. *Jacq. Amer.* 124. t. 181. f. 35. *Aubl.* 392. h
- A. fructu obverse ovato. *Br.* 226.
- Acajouba occidentalis. *Gaert. Fruct.* 1. p. 192. t. 40.
- Pomifera seu potius prunifera. *Sl.* 2. p. 136. *Catesb. Car. App.* 9. t. 9.
- Cassuvium. *Rumph. Amb.* 1. p. 177. t. 69.
- Cashew Tree. *Hughes.* 135.
- Cashew, or Cashou Tree. *Long.* 2. p. 725.
- Cashew Nut, Cassu, or Acajou. *Miller.*

DECANDRIA MONOGYNIA.

BAUHINIA.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

tomentosa. B. foliis basi ovatis subrotundisve subtus petiolis ramulis stipulis pedunculis bracteis calycibusque subvillosis: foliolis ovalibus obtusis 3—4—nerviis ultra medium concretis, pedunculis 1—3—floris, petalis obovatis subobtusis, staminibus decem fertilibus inæqualibus. *De Cand. Prod.*

2. p. 514. *Willd. Sp. Pl.* 2. p. 511. ½
Mandaru maderaspatense. *Pluk. Alm.*
240. *Phyt. t.* 44. *f.* 6.

Cancshena-Pou. *Hort. Mal.* 63. *t.* 35.

Downy Mountain Ebony. *Miller.*

HYMENÆA.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss De Cand.*

- Courbaril.* *H.* foliolis coriaceis subaveniis basi inæqualibus oblongis brevissime et obtuse acuminatis, paniculæ floribus pedunculatis, leguminibus non tuberculatis. *De Cand. Prod.* 2. p. 511. *Willd. Sp. Pl.* 2. p. 512. *Hort. Kew.* 3. p. 24. *Aubl.* 378. ½
H. foliis geminatis. *Br.* 221.
Courbaril bifolia. *Plum. Gen.* 49.
Ceratia diphyllus. *Pluk. Alm.* 96. *Phyt.* t. 82. f. 3.
Locust Tree. *Hughes.* 192. *Long.* 2. p. 728. *Miller.*

PARKINSONIA.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

- aculeata.* *P. aculeata.* *De Cand. Prod.* 2. p. 486. *Willd. Sp. Pl.* 2. p. 513. *Hort. Kew.* 3. p. 24. *Jacq. Amer.* 121. t. 80. *Aubl.* 379. ½
P. foliis minutissimis pinnatis. *Br.* 222.

Holy Thorn, or Royal Cashiauw.

Hughes. 194.

Jerusalem Thorn. *Long.* 2. p. 845.

Prickly Parkinsonia. *Miller.*

CASSIA.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

Fistula. C. foliis 4—6- jugis ovatis subacuminatis glabris, petiolis eglandulosis, racemis laxis ebracteatis, leguminibus teretibus rectis subobtusis lævibus. *De Cand.*

Prod. 2. p. 490. *Willd. Sp. Pl.* 2. p.

518. *Hort. Kew.* 3. p. 27. *Aubl.* 381.

Rumph. Amb. 2. p. 83. t. 21. h

C. arborea, &c. siliqua maxima cylindracea. *Br.* 222.

C. major, seu *Fistula prima.* *Sl.* 2. p. 44.

Cassia *Fistula Tree.* *Hughes.* 192.

Alexandrian Purgine Cassia, &c. *Miller.*

alata.

C. foliolis 8—12—jugis obovato-oblongis utrinque glabris; extimis majoribus; inferioribus axilla proximis, petiolis eglandulosis. *De Cand. Prod.* 2. p. 492. *Willd.*

Sp. Pl. 2. p. 523. *Hort. Kew.* 3. p. 29. u

C. herpetica. *Jacq. Obs.* 2. p. 24. t. 45. f. 2.

C. siliquis quadrialatis. *Br.* 224.

Juglandis folio, &c. siliqua quadrialata.

Sl. 2. *p.* 59. *t.* 175. *f.* 2.

French Guava. *Hughes.* 202.

Ring-Worm Bush. *Long.* 2. *p.* 845.

Wild, or Bastard French Guava.

obtusifolia. *C. foliolis* 3—jugis obovatis obtusis sub-
tus pubescentibus, glandula oblongo-
cylindracea acuta inter infima, stipulis
lineari-subulatis, leguminibus longis gra-
cilibus recurvis. *De Cand. Prod.* 2. *p.*
493. *Willd. Sp. Pl.* 2. *p.* 516. *Hort.*
Kew. 3. *p.* 26. ☉

C. foetida, foliis Sennæ Italicæ. *Dill.*
Elth. 71. *t.* 62. *f.* 72.

Senna minor herbacea. *Sl.* 2. *p.* 47. *t.*
180. *f.* 5.

Italian Senna. *Hughes.* 207.

Round-leaved Cassia. *Miller.*

sulphurea. *C. foliolis* 5—6—jugis ovali-oblongis
subtus glaucis; adultis glabris; juniori-
bus ramulis petiolisque subpuberulis,
glandula ovata sessili inter 3—4—paria
inferiora, stipulis lineari-subulatis erecto-
falcatis, racemis axillaribus erectis folio
brevioribus. *De Cand. Prod.* 2. *p.* 495. ♀
C. arborescens. *Willd. Sp. Pl.* 2. *p.* 520.
Hort. Kew. 3. *p.* 27.

Tree Cassia. *Aiton.*

*occiden-
talis.*

C. foliolis 4—6—jugis ovato-lanceolatis margine pubescentibus, glandula crassa ad basin petioli, pedunculis brevibus 2—4—floris; inferioribus axillaribus; cæteris in racemum terminalem dispositis, leguminibus plano-compressis suturis calloso-tumidulis. De Cand. Prod. 2. p. 497. Willd. Sp. Pl. 2. p. 518. Hort. Kew. 3. p. 26. Aubl. 381. Bot. Reg. t. 83. ☉

C. herbacea major erecta. Br. 224.

Senna occidentalis. Sl. 2. p. 48. t. 175. f. 3, 4.

Wild French Guava, or Stinking Bush. Hughes. 202.

Occidental Cassia. Miller.

Stinking Weed. Browne.

*emargi-
nata.*

C. foliolis 4—jugis ovatis obtusis aut subemarginatis subtus petiolis ramulisque pubescenti-hirsutis, petiolis eglandulosis, racemis axillaribus confertis.

De Cand. Prod. 2. p. 499. Willd. Sp. Pl. 2. p. 516. ½

C. arborescens diffusa. Br. 223.

C. minor fruticosa. Sl. 2. p. 44. t. 180. f. 1—4.

*Wild Senna, or Wild Cassia Fistula. Hughes. 204. t. 20.**

* The Wild Senna of Hughes is certainly the *Cassia minor* of Sloane, the *C. emarginata* of modern botanists. Martyn is in

Notch-leaved Cassia. *Sweet.*

Senna Tree. *Browne.*

*Chamæ-
crista.*

C. erecta, foliolis 10—12—jugis oblongo-linearibus mucronatis, glandula sessili infra par infimum, pedicellis 2—3 supra-axillaribus fasciculatis petiolo brevioribus, leguminibus hispidulis. *De Cand. Prod. 2. p. 503. Willd. Sp. Pl. 2. p. 528.*

Hort Kew. 3. p. 30. Aubl. 382. Bot. Mag. t. 107. h

C. suffruticosa erecta. Br. 225.

Wild Tamarinds. *Hughes, 215.*

Dwarf Cassia. *Miller.*

error in describing the seed vessel as “flat, broad,” it being more correctly represented by Hughes as “five inch long, brownish, round pods.” Sloane’s plate, above referred to, affords a good representation of the lomentum. In this plate the leaflets all appear emarginate; in Hughes’s plate they are simply rounded or slightly mucronate. In the same plant I have seen all these varieties; but although the leaf is generally emarginate, it is seldom so much so as in Sloane’s plate. In the definition of this species, M. De Candolle has with his usual precision employed the term subemarginate.

POINCIANA,

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

*pulcher-
rima.*

P. aculeata, foliolis obovatis, calycibus glabris, petalis fimbriatis longe stipitatis.

De Cand. Prod. 2. p. 484. *Hort. Kew.* 3.

p. 30. *Aubl.* 386. *Jacq. Amer.* 122. ♀

Cæsalpinia pulcherrima. *Willd. Sp. Pl.*

2. p. 531.

α. Corymbis floribus luteis. *Bot. Mag.* t. 995.

P. aculeata, &c. floribus croceis. *Br.* 225.

Flower Fence, or Spanish Carnation (yellow). *Hughes.* 201.

β. Corymbis floribus variegatis.

Senna spuria, &c. flore ex luteo et rubro speciosa. *Sl.* 2. p. 49.

Flower Fence, or Spanish Carnation (red and yellow). *Hughes.* 201.

Barbadoes Pride, or Flower Fence.

Long. 2. p. 815.

Prickly Poinciana, or Barbadoes Flower Fence. *Aiton.*

GUILANDINA.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

Bonduc. G. foliis pubescentibus aut villosis-subvelutinis. *De Cand. Prod.* 2. p. 480.
Hort. Kew. 3. p. 32. h

α. Seminibus flavicantibus.

G. Bonduc. *Willd. Sp. Pl.* 2. p. 534.

Bonduc vulgare majus. *Plum Gen.* 25.

Lobus echinatus fructu flavo. *Sl.* 2.
p. 40.

Yellow Nicker. *Long.* 2. p. 824.

Yellow Bonduc, or Nicker Tree.
Miller.

β. Seminibus griseis.

G. Bonducella. *Willd. Sp. Pl.* 2. p.
534.

Bonduc vulgare minus. *Plum Gen.* 25.

Lobus echinatus fructu cæsius. *Sl.* 2.
p. 41.

Horse-Nicker, or Chick-Stone Tree.
Hughes. 195. t. 18.

Grey Nicker. *Long.* 2. p. 824.

Grey Bonduc. *Miller.*

MORINGA.

Lomentaceæ (*Guilandina*). *Lin.*—Leguminosæ. *Juss. De Cand.*

pterygos- *M. leguminibus trigonis vel tetragonis,*
perma. *seminibus trigonis angulis in alas ex-*

pansis, floribus semidecandris. M.—

De Cand. Prod. 2. p. 478. ½

Hyperanthera Moringa. Willd. Sp. Pl.

2. p. 536. Hort. Kew. 3. p. 32.*

Guilandina Moringa. Jac. Ic. 3. t. 461.

Lignum peregrinum. Bauh. Pin. 416.

Wing-seeded Horse-Radish Tree. Sweet.

Moringa Tree.

* Willdenow questions whether the *Hyperanthera decandra*, “*leguminibus suboctogonis*,” be a species distinct from *H. Moringa*, “*leguminibus trigonis*.” De Candolle also doubts whether the *Moringa polygona* be not a variety of *M. pterygosperma*. On this subject I would remark, that the seed vessel, in its green state, is either subhexangular or suboctangular; when dry, either triangular or quadrangular. This variation occurs not only on different plants, but also on the same plant; and I have frequently seen on the same common peduncle the legumen triangular and trivalved, and quadrangular quadrivalved. After repeated and diligent search, I have however never been able to find a single flower decandrous in Barbados, five of the ten stamens being always imperfect. On this ground, unless the observations of others shall prove that I have been

HÆMATOXYLON.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

campechi- *H. campechianum. De Cand. Prod. 2. p.*
anum. 485. Willd. Sp. Pl. 2. p. 547. Hort.
Kew. 3. p. 36. Aubl. 393. h

H. spinosum. Br. 221.

Lignum campechianum. Catesb. Car.
*2. p. 66. t. 66.**

Logwood. Hughes. 169. Long. 2. p. 754.

Logwood, Blood Wood, or Campechi
Wood. Miller.

unfortunate in the objects of my examination, I think the species in question must be kept distinct. I have thought it necessary not only to correct the definition of M. De Candolle as regards the legumen, but also to insert the semidecandrous character of the flower, as affording an additional, and necessary specific distinction.

* This plate very well represents the raceme of flowers; but the foliage is not good: and the figure of the legumen or pod is altogether an error.

ADENANTHERA.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

pavonina. A. foliolis ovalibus obtusis utrinque glabris, legumine subfalcato. M.— *De Cand. Prod.* 2. p. 446. *Willd. Sp. Pl.* 2. p. 550. *Hort. Kew.* 3. p. 37. *Jacq. Coll.* 4. p. 212. t. 23. h

Poinciana foliis duplicato-pinnatis. *Lin. Cliff.* 158. No. 2.

Bastard Flower Fence. *Aiton.*

Circassian Bead Tree.

TRIBULUS.

Gruinales. *Lin.*—Rutaceæ. *Juss.*—Zygophylleæ. *De Cand.*

cistoides. T. foliis 8—jugis : foliolis subæqualibus subtus sericeis, pedicellis petioli longitudine. *De Cand. Prod.* 1. p. 703. *Willd. Sp. Pl.* 2. p. 567. *Hort. Kew.* 3. p. 42. *Aubl.* 398. *Bot. Reg.* t. 791. u

T. terrestris americanus. *Pluk. Alm.*
 374. *Phyt. t.* 67. *f.* 4.
Cistus-like Caltrops. *Aiton.*

GUAIACUM.

Gruinales. *Lin.*—Rutaceæ. *Juss.*—Zygophylleæ. *De Cand.*

officinale. *G. foliis 2—3—jugis: foliolis obovatis ovalibusve obtusis, capsulis bilocularibus.*
M.— Willd. Sp. Pl. 2. p. 538. De Cand.
Prod. 1. p. 707. Hort. Kew. 3. p. 33. ½
G. foliis fere impetiolatis. Br. 225.
G. flore cæruleo. Plum. Gen. 39.
G. jamaicense. Pluk. Alm. 180. Phyt.
t. 35. f. 3, 4.
Pruno vel evonymo affinis arbor. Sl. 2.
p. 133. t. 222. f. 3—6.
Lignum Vitæ. Hughes. 142. t. 9. Browne.
Officinal Guaiacum. Miller.

SWIETENIA.

Miscellaneæ. *Lin.*—Meliis aff. *Juss.*—Meliaceæ. *De Cand.*

Mahogoni. S. foliis sub—4—jugis: foliolis ovato-lanceolatis basi inæqualibus apice acuminatis, racemis axillaribus paniculatis.

De Cand. Prod. 1. p. 625. *Willd. Sp. Pl.*

2, p. 557. *Hort. Kew.* 3. p. 38. *Jacq.*

Amer. 127. *Cav. Diss.* 365. t. 209. ½

Cedrela foliis pinnatis, &c. ligno graviore. *Br.* 158.

Mahogany Tree. *Long.* 2. p. 842. *Miller.*

QUASSIA.

Gruinales. *Lin.*—Magnoliis aff. *Juss.*

Simaroubeæ. *De Cand.*

*amara.** Q. floribus hermaphroditis, foliis imparipinnatis: foliolis oppositis sessilibus, pe-

* “Lignum Quassiæ venale non ex hoc frutice venit; colligitur a Quassia excelsa quæ minus amara.”—*Willd.*

tiolo articulato alato, floribus racemosis.

Willd. Sp. Pl. 2. p. 567. Hort. Kew. 2. p. 42. Bot. Mag. t. 497. De Cand. Prod. 1. p. 733. h

Bitter Quassia. *Miller.*

excelsa. Q. floribus polygamis pentandris paniculatis, foliis impari-pinnatis: foliolis oppositis petiolatis, petiolo nudo. *Willd. Sp.*

Pl. 2. p. 569. Swtz. Fl. 742. h

Simarouba? *excelsa. De Cand. Prod. 1. p. 733.*

Gall Tree. *Hughes. 116.*

Lofty Quassia. *Miller.*

Bitter Wood.

MELIA.

Trihilatæ. *Lin.*—Melix. *Juss.*—Meliaceæ. *De Cand.*

Azedarach. M. foliis bipinnatis: foliolis lævibus incisis subquinatis. *De Cand. Prod. 1. p. 621. Willd. Sp. Pl. 2. p. 558. Hort. Kew. 3. p. 39. Aubl. 393. Cav. Diss. 363. t. 207. Bot. Mag. t. 1066.* h

Common Bead-Tree. *Sweet.*

Barbados Lilac.

TRIPHASIA.

Dubii ord. (*Limonia*.) *Lin.*—*Aurantiaceæ*. *Juss. De Cand.*

trifoliata. *T. foliis 3—foliatis. De Cand. Prod. 1. p. 536.* h

Limonia trifoliata. Willd. Sp. Pl. 2. p. 571. Hort. Kew. 3. p. 43. Jacq. Ic. 3. t. 463. Bot. Repos. t. 143.

Three-leaved Triphasia. *Sweet.*

Limonia.—Myrtle Lime.*

* “*Limoniam trifoliatam vidi 3—petalam 5—andram.*”—*Juss. Gen. 290.* Jacquin describes the flower as tripetalous hexandrous, to which he adds, “*In floribus paucis vidi petala quatuor cum staminibus octo.*”—*Collect. 3. p. 177.* The most general character of the flower is certainly tripetalous hexandrous; but I have seen on the same plant, flowers 3—petaled 6—androus, 4—petaled 6—androus, 4—petaled 7—androus, 4—petaled 8—androus, and 5—petaled 6—androus; but even on this plant the 3—petaled 6—androus flowers greatly prevailed, and this is to be considered as the general character of the flower, and agrees perfectly with that assigned to the genus by De Candolle—“*Flores proportionem partium ternaria.*”

JUSSIÆA.

Calycanthemæ. *Lin.*—Onagræ. *Juss.*

Onagrariæ. *De Cand.*

octonervia. *J.* herbacea erecta glabra, foliis angustolanceolatis acuminatis, floribus sessilibus, calycis lobis quatuor lanceolatis acutis: tubo cylindrico 8—nervio striato folio florali brevior, petalis obovatis emarginatis lobos calycis superantibus. *De Cand. Prod.* 3. p. 57. ☉

J. octovalvis. *Willd. Sp. Pl.* 2. p. 576.

œnothera octovalvis. *Lin. Sp. Pl.* 492.

Jacq. Amer. 102. t. 70.

œnothera assurgens glabra. *Br.* 208.

Many Seed. *Hughes.* 212.

Spear-leaved Jussiaea. *Sweet.*

Smooth Primrose Willow. *Browne.*

TETRAZYGIA.

Calycanthemæ (*Melastoma*). *Lin.*—*Melastomæ. Juss.*

Melastomaceæ. De Cand.

discolor. T. ramis obtuse tetragonis, petiolis paniculis foliisque junioribus subtus pubescentibus stellata brevi subfloccosa albidis, foliis repando-subdenticulatis ovali-acuminatis 5-nerviis supra glabris, cyma paniculato-corymbosa trichotoma terminali, calycis dentibus quatuor brevibus obtusis, petalis ovalibus retusis. *De Cand. Prod.* 3. p. 172. h

Melastoma discolor. Willd. Sp. Pl. 2. p. 599. *Hort. Kew.* 3. p. 46. *Jacq. Amer.* 130. t. 84.

Grossulariæ fructu arbor, &c. Sl. 1. p. 39? p. 40?

Two-coloured *Melastoma. Aiton.*

CREMANIUM.

Calycanthemæ (*Melastoma*). *Lin.*—*Melastomæ. Juss.*

Melastomaceæ. De Cand.

quadran- C. ramis tetragonis glabriusculis, foliis
gulare. petiolatis ovato-lanceolatis acuminatis
integris 3—nerviis rigidis supra glabris
subtus ferrugineis, thyrso paniculato ter-
minali stricto: ramulis oppositis angu-
latis, calycis ovati pube stellata cinerei
lobis obtusis ovatis membranaceis, stig-
mate capitato. *De Cand. Prod. 3. p.*
192. h

Melastoma quadrangularis. Willd. Sp.
Pl. 2. p. 585. Swtz. Fl. 770.

CASEARIA.

Incertæ sedis. Juss.—*Samydeæ. De Cand.*

ramiflora. C. floribus 8—antheris 5—partitis, pedi-
cellis 1—floris infra folia secus ramulos

fasciculatis, foliis ellipticis acutis serratis
utrinque glabris. *De Cand. Prod.* 2. p.
49. *Willd. Sp. Pl.* 2. p. 627. $\frac{1}{2}$

Iroucana guianensis. *Aubl.* 329. t. 127.
Branch-flowered Casearia. *Sweet.*

parviflora. C. floribus 10—antheris 5—partitis mi-
nimis, pedicellis confertis axillaribus
1—floris, foliis oblongis acuminatis cre-
nulatis utrinque glabris nitidis evidenter
pellucido-punctatis. *De Cand. Prod.* 2.
p. 49. *Willd. Sp. Pl.* 2. p. 627. $\frac{1}{2}$

Samyda parviflora. *Lin. Sp. Pl.* 557.

Samyda foliis ovatis cum acumine. *Br.*
217.

Arbor baccifera, foliis oblongis acumi-
natis *Sl.* 2. p. 108. t. 211. f. 2.

Wild Honey Tree. *Hughes.* 159.

Smaller Cloven-Berry Bush. *Browne.*

Small-flowered Casearia. *Sweet.*

CLETHRA.

Bicornes. *Lin.*—*Ericæ.* *Juss.*

tinifolia. C. foliis oblongis integerrimis subtus
incanis, racemis paniculatis florentibus

ebracteatis, pedunculis tomentosis. *Willd.*

Sp. Pl. 2. p. 620. *Swtz. Fl.* 845. b

Tinus occidentalis. *Lin. Sp. Pl.* 530.

Volkameria arborea. *Br.* 214. t. 21. f. 1.

Baccifera arbor calyculata. *Sl.* 2. p. 86.

t. 198. f. 2.

Bastard Locust, or Forest Tree. *Hughes.*

149.

Forest Bark, or Bastard Locust. *Hughes.*

224.

DECANDRIA DIGYNIA

HYDRANGEA.

Succulentæ. *Lin.*—Saxifragæ. *Juss.*

hortensis. *H.* cymis radiatis, foliis ellipticis utrinque attenuatis dentatis glabris. *Willd.*
Sp. Pl. 2. p. 633. *Hort. Kew.* 3 p. 63.
Bot. Mag. t. 438. h

Garden Hydrangea, or Chinese Guelder
Rose. *Miller.*

DIANTHUS.

Caryophylli. *Lin.*—Caryophylleæ. *Juss. De Cand.*

barbatus. *D.* floribus aggregatis fasciculatis, squamis calycinis ovato-subulatis tubum æquantibus, foliis lanceolatis. *Willd.*
Sp. Pl. 2. p. 671. *Hort. Kew.* 3. p. 78.
De Cand. Prod. 1. p. 355. 2

α . flore simplici. *Bot. Mag. t. 207.*

Caryophyllus barbatus hortensis.

Bauh. Pin. 208.

β . flore pleno.

Caryophyllus barbatus flore multiplici. Bauh. Pin. 208.

Bearded Pink, commonly called Sweet William. *Miller.*

chinensis. *D. floribus solitariis, squamis calycinis subulatis patulis foliaceis tubum æquantibus, petalis crenatis, foliis lanceolatis.*

Willd. Sp. Pl. 2. p. 677. Hort. Kew. 3.

p. 80. De Cand. Prod. 1. p. 359. Bot.

Mag. t. 25.

⊙

Chinese Pink. *Miller.*

virgineus. *D. floribus solitariis, caulibus paucifloris, squamis calycinis brevissimis obtusissimis binis, petalis crenatis. Willd. Sp.*

Pl. 2. p. 680. De Cand. Prod. 1. p. 361.

Bot. Mag. t. 1740.

2

Upright Pink. *Miller.*

Caryophyllus. *D. floribus solitariis, squamis calycinis subrhombeis brevissimis, petalis crenatis imberbibus. Fl. Br. 461. Willd. Sp. Pl.*

2. p. 674. Hort. Kew. 3. p. 79. De

Cand. Prod. 1. p. 359.

2

α . flore simplici. *Engl. Bot. t. 214.*

Clove Pink. *Aiton.*

β . flore pleno. *Bot. Mag. t. 39.*

Carnation. *Aiton.*

DECANDRIA TRIGYNIA.

MALPIGHIA.

Trihilatæ. *Lin.*—Malpighiæ. *Juss.*—Malpighiaceæ. *De Cand.*

urens. M. ramis glabris, foliis oblongo-ovatis
subtus decumbenti-hispidis superne gla-
bris, pedunculis unifloris corymbiferisve
folio duplo brevioribus, petalis æqualibus.

De Cand. Prod. 1. p. 577. *Willd. Sp.*

Pl. 2. p. 734. *Hort. Kew.* 3. p. 104.

Cav. Diss. 407. t. 235. f. 1. *Aubl.* 462.

Bot. Reg. t. 96.

h

M. viminea. *Br.* 229.

Arbor baccifera folio oblongo. *Sl.* 2. p.

106. t. 207. f. 3.

Stinging Barbadoes Cherry. *Miller.*

Cowhage Cherry. *Browne.*

coccifera. M. foliis subovatis dentato-spinosis gla-
bris. *De Cand. Prod.* 1. p. 578. *Willd.*

Sp. Pl. 2. p. 737. *Hort. Kew.* 3. p. 105.

Cav. Diss. 408. t. 235. f. 2. *Bot. Reg.*

t. 568.

h

M. coccigera. *Jacq. Ic. 3. t. 470.*

M. humilis ilicis cocciglandiferæ foliis.
Plum. Gen. 46.

Kermes-Oak-leaved Barbadoes Cherry.
Aiton.

glabra. *M. foliis ovatis integerrimis glabris, pedunculis axillaribus umbellatis. De Cand.*
Prod. 1. p. 578. Willd. Sp. Pl. 2. p. 731.
Hort. Kew. 3. p. 102. Aubl. 461. Cav.
Diss. 406. t. 234. f. 1. h

M. fruticosa erecta, foliis nitidis ovato-acuminatis. Br. 230.

Arbor baccifera folio subrotundo. *Sl. 2.*
p. 106. t. 207. f. 2.

Red Cherry-Tree. *Hughes. 151.*

Smooth-leaved Barbadoes Cherry. *Miller.*

Shrubby erect Malpighia. *Browne.*

puniceifolia. *M. foliis ovatis integerrimis glabris, pedunculis axillaribus unifloris. De Cand.*
Prod. 1. p. 578. Willd. Sp. Pl. 2. p. 732. Hort. Kew. 3. p. 103. Aubl. 462.
Cav. Diss. 406. t. 234. f. 2. h

M. fruticosa erecta ramulis gracilibus patentibus. Br. 230.

Pomegranate-leaved Barbadoes Cherry.
Miller.

Cherecze, or Barbadoes Cherry *Browne.*

BYRSONIMA.

Trihilatæ. (*Malpighia*). *Lin.*—Malpighiæ (*Malpighia*). *Juss.*
 Malpighiaceæ. *De Cand.*

coriacea. B. foliis ovatis acutis integerrimis utrinque glabris, racemis dense spicatis pubescentibus erectis. *De Cand. Prod.* 1. p. 580. h

Malpighia coriacea. *Willd. Sp. Pl.* 2. p. 736. *Swtz. Fl.* 854.

Malpighia arborea floribus spicatis. *Br.* 230.

Tiliæ affinis laurifolia. *Sl.* 2. p. 20. t. 163. f. 1.

Leathery-leaved Byrsonima. *Sweet.*

Locus-Berry Tree. *Browne.*

BUNCHOSIA.

Trihilatæ (*Malpighia*). *Lin.*—Malpighiæ (*Malpighia*). *Juss.*
 Malpighiaceæ. *De Cand.*

glandulosa. B. foliis ovato-ellipticis acuminatis glabris, petiolis apice biglandulosis, racemis

simplicibus folio brevioribus. *De Cand.*

Prod. 1. p. 581. h

Malpighia glandulosa. *Willd. Sp. Pl.* 2.

p. 733. *Cav. Diss.* 411. t. 239. f. 2.

Two-glanded *Bunchosia.* *Sweet.*

TRIOPTERIS.

Trihilatæ. *Lin.*—*Malpighiæ.* *Juss.*—*Malpighiaceæ.* *De Cand.*

jamaicensis. *T. foliis oblongis acuminatis venosis nitidis, racemis compositis terminalibus*

laxis, fructibus trialatis. *Willd. Sp. Pl.*

2. p. 743. *De Cand. Prod.* 1. p. 586.

Aubl. 467. h

Banisteria seminibus trialatis. *Br.* 231.

Fire-Burn Bush. *Hughes.* 172.

Jamaica *Triopteris.* *Sweet.*

DECANDRIA PENTAGYNIA.

SPONDIAS.

Dubii ord. *Lin.*—Terebinthaceæ. *Juss. De Cand.*

purpurea. *S. foliis impari-pinnatis, petiolo compresso. De Cand. Prod. 2. p. 75.* h

S. Mombin. Willd. Sp. Pl. 2. p. 750.

S. Myrobalanus. Jacq. Amer. 139. t. 88.

S. diffusa. Br. 228.

Myrobalanus, &c. fructu purpureo. Sl. 2. p. 126. t. 219. f. 3—5.

Jamaica Plum Tree. Hughes. 179.

Top-knot Plum. Long. 2. p. 827.

Purple Hog-Plum, or Spanish Plum. Miller.

lutea. *S. foliis impari-pinnatis, petiolo tereti, foliolis 7—9 ovato-oblongis nitidis sensim acuminatis. De Cand. Prod. 2. p. 75.*

Aubl. 496. h

S. Myrobalanus. Willd. Sp. Pl. 2. p. 751. Hort Kew. 3. p. 108.

S. Mombin. *Jacq. Amer.* 138.

S. foliolis plurimis. *Br.* 229.

Myrobalanus, &c. fructu luteo. *Sl.* 2. *p.* 125. *t.* 219. *f.* 1, 2.

Gully Plum Tree. *Hughes.* 178. *t.* 13.

Yellow, or Jamaica Plum. *Long.* 2. *p.* 827.

Yellow Hog-Plum, or Jamaica Plum.
Miller.

Hog-Plum Tree.

dulcis.

S. foliis impari-pinnatis, petiolis tere-
tibus: foliolis 11—13 ovali-oblongis acu-
minatis costatis serratis. *De Cand. Prod.*

2. *p.* 75. *Willd. Sp. Pl.* 2. *p.* 752.

Hort. Kew. 3. *p.* 108.

h

S. cytherea. *Gaert. Fruct.* 2. *p.* 101. *t.* 103.

Otaheite Apple. *Aiton.*

Golden Apple.

DODECANDRIA MONOGYNIA.

RHIZOPHORA.

Holeraceæ. *Lin.*—Caprifoliæ. *Juss.*

Rhizophoreæ. *De Cand. Brown.*

- Mangle.* *R. foliis obovato-oblongis obtusis, pedunculis 2—3—floris petiolo longioribus, fructibus subulato-clavatis. De Cand. Prod. 3. p. 32. Willd. Sp. Pl. 2. p. 843. Jacq. Amer. 141. t. 89.* b
- R. utrinque brachiata. Br. 211.*
- Mangle Pyri foliis. Sl. 2. p. 63.*
- Candela americana. Catesb. Car. 1. p. 63. t. 63.*
- Red Mangrove Tree. Hughes. 194. Long. 2. p. 811.*
- Mangrove Tree. Miller.*

GARCINIA.

Guttiferæ. *Juss. De Cand.*

- Mangos-* G. foliis ovatis acutis venosis, floribus
tana. terminalibus solitariis, corollis rubris,
 stigmatibus 6—8—lobo, baccis speciosissimis
 edulibus, pericarpio fungoso. *De Cand.*
Prod. 1. p. 560. *Willd. Sp. Pl.* 2. p. 848.
Hort. Kew. 3. p. 143. h
Mangostana. *Rumph. Amb.* 1. p. 132. t. 43.
Laurifolia javanensis. *Raj. Hist.* 1662.
Mangostan, or Mangosteen. *Miller.*

CANELLA.

Melicæ. *Juss.*—Guttiferæ. *De Cand.*

- alba.** C. alba. *Willd. Sp. Pl.* 2. p. 851. *De*
Cand. Prod. 1. p. 563. *Hort. Kew.* 3.
 p. 144. *Swtz. Lin. Tr.* 1. p. 96. t. 8. h

* On the authority of recent Pharmacopœias the Canella has
 entirely superceded the medicinal use of the true Winter's Bark—
Wintera aromatica. *Willd. Sp. Pl.* 2. p. 1239.

C. foliis oblongis obtusis nitidis. *Br.* 275.

t. 27. *f.* 2, 3.*

Arbor baccifera laurifolia aromatica.

Sl. 2. *p.* 87. *t.* 191. *f.* 2. *Catesb. Car.* 1.
p. 50. *t.* 50.

Cortex Winteranus. *Elackw. t.* 206.

Wild Cinnamon. *Hughes.* 143. *Long.*
2. p. 705.

Laurel-leaved Canella. *Miller.*

Canella, or Winter's Bark. *Browne.*

CRATÆVA.

Putamineæ. Lin.—*Capparides. Juss.*—*Capparideæ. De Cand.*

gynandra. C. staminibus 20--24 toro cylindraceo
insertis petalis longioribus, bacca ovata,
foliolis ovatis acutis. *De Cand. Prod.* 1.

p. 243. *Willd. Sp. Pl.* 2. *p.* 852. *Hort.*
Kew. 3. *p.* 144. *Aubl.* 474. h

C. arborea triphylla, foliis ovatis glabris.
Br. 246.

* Plate 27. f. 2. of Browne certainly represents *Canella alba*, and not *Breynia fruticosa*, as it is by mistake made to do by Browne. *vid. Swtz. Lin. Tr.* 1. *p.* 100.

Anona trifolia. *Sl.* 2. *p.* 169.

Garlick-Pear Tree. *Hughes.* 134.

Thin-leaved *Cratæva*, or Garlick-Pear.
Browne.

Thin-leaved Garlick-Pear. *Aiton.*

TRIUMFETTA.

Calumniferæ. *Lin.*—Tiliaceæ. *Juss. De Cand.*

*semitri-
loba.*

T. foliis basi ovatis trilobis callososerratis utrinque velutinis; inferioribus sub—5—lobis; superioribus oblongis indivisis, calycibus pubescentibus subapiculatis, fructibus globosis glabris echinatis, setis retrorsum pilosulis. *De Cand.*

Prod. 1. *p.* 507. *Willd. Sp. Pl.* 2. *p.* 855. *Hort. Kew.* 3. *p.* 146. *Jacq. Amer.* 147. h

T. villosa. *Br.* 233? *

* I have questioned whether the *Triumfetta* of *Browne* be properly a synonym of *T. semitriloba*, *Willd.* partly on the authority of *Swartz*, who thinks it more likely to be identical with *T. rhombæfolia*, but principally from *Browne's* statement that "one half the capsule is echinated, the other smooth."

Mallow-leaved Triumfetta. *Miller.*

Bur-Bark. *Browne.*

rhomboidea.

T. foliis rhomboideis; inferioribus sub—5—lobis; mediis sub—3—lobis; summis oblongis; omnibus calloso-serratis utrinque velutinis basi subcuneatis, floribus axillaribus, calycibus apiculatis, fructibus echinatis subpilosis. De Cand. Prod. 1.

p. 507. Jacq. Amer. 147. t. 90. ½

T. rhombeæfolia. Willd. Sp. Pl. 2. p. 856.

Swtz. Fl. 863.

Rhomb-leaved Triumfetta. *Miller.*

PORTULACA.

Succulentæ. *Lin.*—Portulacææ. *Juss. De Cand.*

oleracea. P. foliis cuneiformibus carnosis, axillis geniculisque nudis, floribus sessilibus.

De Cand. Prod. 3. p. 353. Willd. Sp.

Pl. 2. p. 859. Hort. Kew. 3. p. 147.

Aubl. 475. Br. 233.

⊙

α. sylvestris.

Small Purslane. *Sweet.*

β. sativa.

Garden Purslane. *Sweet.*

Purslane (*red and white*). *Hughes*. 165.
parvifolia. *P. foliis cuneiformibus minutis carnosis, caule ramosissimo prostrato, floribus alte pedunculatis sessilibusque. De Cand.*

Prod. 3. p. 353. ☉

Wild Purslane (*white*). *Hughes*. 243.

Small-leaved Purslane. *Sweet*.

pilosa. *P. caule diffuso, geniculis longe pilosis, foliis alternis lineari-lanceolatis; floribus subverticillatis, floribus ad apices ramorum congestis sessilibus villo longo circumdatis. De Cand. Prod.* 3. p. 354.

Willd. Sp. Pl. 2. p. 860. *Hort. Kew.* 3.

p. 147. *Bot. Reg.* t. 792. ☉

Wild Purslane (*red*). *Hughes*. 243.

Hairy Purslane. *Sweet*.

TALINUM.

Succulentæ (*Portulaca*). *Lin.*—*Portulacæ. Juss. De Cand.*

triangulare. *T. caule fruticoso erecto, foliis planis canaliculatis cuneiformibus emarginatis mucronatis, racemo simplici, rachi triquetra. De Cand. Prod.* 3. p. 357. *Willd.*

Sp. Pl. 2. p. 862. *Hort. Kew.* 3. p. 148. ½

Portulaca triangularis. *Jacq. Amer.* 147.

Triangular Talinum. *Aiton.*

patens. T. caule suffruticoso erecto, foliis planis ovalibus mucronatis; infimis obovatis, panicula terminali, pedunculis alternis dichotomis ebracteatis. *De Cand. Prod.*

3. p. 357. *Willd. Sp. Pl.* 2. p. 863.

Hort. Kew. 3. p. 148. h

T. paniculatum. *Gaert. Fr.* 2. p. 219. t. 128.

Portulaca paniculata. *Jacq. Amer.* 148.

Jacq. Vind. 2. p. 71. t. 151.

Panicked Talinum. *Aiton.*

RESEDA.

Miscellanææ. *Lin.*—Capparideææ. *Juss.*

Resedaceææ. *De Cand.*

odorata. R. foliis integris trilobisque, calycibus florem æquantibus. *Willd. Sp. Pl.* 2. p. 881. *Hort. Kew.* 3. p. 155. *Bot. Mag. t.* 29. ☉

Sweet Reseda, or Mignonette. *Miller.*

ICOSANDRIA MONOGYNIA

MELOCACTUS.

Succulentæ (*Cactus*). *Lin.*—Cacti (*Cactus*). *Juss.*

Cactææ. *De Cand.*—Opuntiaceæ. *Kunth.*

communis. M. ovato-subrotundus atro-virens 12—
18—angulatus, costis rectis, spinis fasci-
culatis fuscis subæqualibus. *De Cand.*
Prod. 3. p. 460. h

Cactus melocactus. *Willd. Sp. Pl.* 2. p.
938. *Hort. Kew.* 3. p. 175.

Cactus humilis subrotundus. *Br.* 238.

Turk's Head. *Hughes.* 226.

Great Melon-Thistle, or Turk's Cap.
Miller.

Turk's Head, or Pope's Head Indian Fig.
Browne.

Cactus trigonus repens. *Plum. Ic.* 193.
t. 200. *f.* 1.

Great triangular *Cereus.* *Sweet.*

trigonus. *C. repens triqueter, angulis vix canalicu-*
latis, aculeis 5—7 stellatis. *De Cand.*
Prod. 3. *p.* 468. h

Cactus triangularis. β . *Willd. Sp. Pl.*
2. p. 942. *Hort. Kew.* 3. *p.* 178.
Aubl. 482.

Cactus triangularis foliosus. *Jacq. Amer.*
152. t. 181. *f.* 65.

Cactus caule triangulari. *Plum. Ic.* 193.
t. 200. *f.* 2.

Cactus debilis brachiatus. *Br.* 238.

Ficus indica folio triangulari. *Sl.* 2. *p.* 155.

Prickly Pear Vine. *Hughes.* 186.

Small triangular *Cereus.* *Sweet.*

Strawberry-Pear. *Browne.*

OPUNTIA.

Succulentæ (*Cactus*). *Lin.*—*Cacti* (*Cactus*). *Juss.*

Cactæ. *De Cand.*—*Opuntiaceæ.* *Kunth.*

Tuna. *O. erecta, articulis late ovato-oblongis,*
aculeis subulatis longis flavicantibus,
lana brevi. *De Cand. Prod.* 3. *p.* 472. h

Tuna major, spinis validis flavicantibus.

Dill. Elth. 396. *t.* 295. *f.* 380.

Cactus Tuna. α . *Willd. Sp. Pl.* 2. *p.* 944. *Hort. Kew.* 3. *p.* 179.

Pimploes. *Hughes.* 185.

Great Yellow-spined Indian Fig. *Sweet.*

coccinelli- *O. erecta*, articulis ovato-oblongis sub-
fera. inermibus, floribus conniventibus, stami-

nibus styloque exsertis. *De Cand. Prod.* 3. *p.* 473. h

O. spinulis obtusis mollibus et innocentibus. *Sl.* 2. *p.* 152.

Cactus coccinellifer. *Willd. Sp. Pl.* 2. *p.* 944. *Hort. Kew.* 3 *p.* 179. *Bot. Mag.* *t.* 2741, *t.* 2742. *Bot. Repos.* *t.* 533.

Cactus brachiatus, &c. subinermis major. *Br.* 237.

Tuna mitior, flore sanguineo. *Dill. Elth.* 399. *t.* 297. *f.* 383.

Cochineel Shrub. *Hughes.* 135.

Cochineal Indian Fig. *Miller.*

PERESKIA.

Succulentæ (*Cactus*). *Lin.*—Cacti (*Cactus*). *Juss.*

Cactææ. *De Cand.*

aculeata. P. foliis ellipticis, aculeis in axilla solitariis
demum in caule fasciculatis, floribus sub-
paniculatis, fructu globoso sepala foliacea
gerente. *De Cand. Prod.* 3. p. 474. *Dill.*
Elth. 305. t. 227. f. 294. ½

Cactus Pereskia. *Willd. Sp. Pl.* 2. p.
946. *Hort. Kew.* 3. p. 180.

Cactus sarmentosus foliatus. *Br.* 237.

Grossulariæ fructu majore arbor spinosa.
Sl. 2. p. 86.

Gooseberry Shrub. *Hughes.* 162.

Barbadoes Gooseberry. *Miller.*

PSIDIUM.

Hesperidæ. *Lin.*—Myrti. *Juss.*—Myrtacææ. *De Cand.* *Brown.*

pyriferum. P. ramulis tetragonis, foliis ellipticis
acutis, nervis prominulis lineatis subtus

pubescenti-velutinis, pedicellis unifloris, fructibus pyriformibus. *De Cand. Prod.*

3. p. 233. *Willd. Sp. Pl.* 2. p. 957.

Hort. Kew. 3. p. 185. *Aubl.* 487. h

Cujavus domestica. *Rumph. Amb.* 1. p. 140. t. 47.

Malo punicæ affinis, &c. fructu majore albo. *Sl.* 2. p. 163?

Guava Tree (*pear fashioned or perfumed*).
Hughes. 132.

Pear-fruited Guava. *Sweet.*

French Guava.

*pomife-
ram.*

P. ramulis tetragonis, foliis ovali-aut-oblongo-lanceolatis subtus pubescentibus, pedunculis tri-aut-multifloris, fructibus globosis. *De Cand. Prod.* 3. p. 234. *Willd.*

Sp. Pl. 2. p. 958. *Hort. Kew.* 3. p. 185.

Aubl. 487. h

Cujavus agrestis. *Rumph. Amb.* 1. p. 142. t. 48.

P. fruticosum. *Br.* 238.

Malo punicæ affinis, &c. fructu rubro majore. *Sl.* 2. p. 161.*

Malo punicæ affinis, &c. fructu minore albo. *Sl.* 2. p. 163.

Guava Tree (*round*). *Hughes.* 132.

* The large red Guava of Sloane improved by cultivation becomes the Garden Guava.

Apple-fruited Guava. *Sweet.**
Common and Garden Guava.

MYRTUS.

Hesperidæ. *Lin.*—Myrti. *Juss.*—Myrtaceæ. *De Cand. Brown.*

communis. *M. pedicellis solitariis unifloris folii circiter longitudine, bracteolis 2 linearibus deciduis sub flore, calycibus quinquefidis, foliis ovatis lanceolatisve acutis.*

De Cand. Prod. 3. p. 239. Willd. Sp.

Pl. 2. p. 967. Hort. Kew. 3. p. 188. ½

α. romana.

M. latifolia romana. Bauh. Pin. 468.

Common broad-leaved Myrtle. *Aiton.*

β. tarentina.

M. minor vulgaris. Bauh. Pin. 469.

Box-leaved Myrtle. *Aiton.*

* The vernacular epithets, red and white, applied in the Gardener's Dictionary, and in the Hortus Kewensis, to the apple-fruited and pear-fruited Guava, should be discarded. The editor of the former work very properly states, although the statement involves a contradiction, that the berry of his red Guava is within "either white or red."

γ. italica.

M. communis italica. *Bauh. Pin.* 468.

Italian, or Upright Myrtle. *Aiton.*

δ. boetica.

M. latifolia boetica. *Bauh. Pin.* 469.

Orange-leaved Myrtle. *Aiton.*

MYRCIA.

Hesperidæ (*Myrtus*). *Lin.*—Myrti (*Myrtus*). *Juss.*

Myrtaceæ. *De Cand.*

coriacea.

M. pedunculis terminalibus paniculatis folio longioribus glabris, ramulis distantibus oppositis apice 1—3 floris, floribus quinquelobis subnudis, foliis obovato-aut elliptico-subrotundis obtusis coriaceis opacis margine subrevolutis utrinque lucidis superne impresso punctatis ramisque glabris. *De Cand. Prod.* 3. p. 243. h

Myrtus coriacea. *Willd. Sp. Pl.* 2. p. 973.

Hort. Kew. 3. p. 190. *Swtz. Fl.* 909.

Myrtus acris. β. *Swtz. Prod.* 79.

Wild Bay-Berry Tree. *Hughes.* 145.

Sumach-leaved Myrtle. *Miller.*

acris.

M. pedunculis axillaribus et terminalibus trichotomis corymbosis folio longioribus compressis, floribus quinquefidis, foliis ellipticis obtusis convexis coriaceis glaberrimis superne venis elevatis reticulatis subtilissime pellucido-punctatis.

De Cand. Prod. 3. p. 243.

h

Myrtus acris. Willd. Sp. Pl. 2. p. 973.

Hort. Kew. 3. p. 190. Swtz. Fl. 909.

Caryophyllus foliis oblongo-ovatis oppositis. Br. 247.

*Bay-Berry Tree. Hughes. 145. t. 10.**

Wild Clove, or Bay-Berry Myrtle. Miller.

Bay-Berry Tree. Wild Cinnamon. Wild Clove. Browne.

* In differing from so great authority as M. De Candolle, by making the Bay-Berry of Hughes a synonym to *Myrcia acris*, and not to *Eugenia Pimenta*, to which it indeed has a near resemblance, I may be allowed to suspect that his mistake may have arisen from Hughes having stated of his Bay-berry Tree that "its berries are well known by the name of Jamaica pepper"; whereas the true Pimento, Jamaica pepper, or Allspice is no part of the produce of Barbados; the *Eugenia Pimenta* being a rare exotic, cultivated only by a few who are curious in plants; nor could the berries of the Barbados Bay-Berry be passed in commerce as Pimenta. Of *Eugenia Pimenta*, De Candolle says "bacca globosa 1—sperma." The Bay-Berry of Hughes is oblong, and generally four-seeded, agreeing with the description given by Swartz of *Myrtus acris*, "bacca oblonga 2—4 sperma." The berry of the *Eugenia Pimenta*, however, also varies much in the number of its seeds.

CARYOPHYLLUS.

Hesperidæ (*Eugenia*). *Lin.*—Myrti (*Eugenia*). *Juss.*

Myrtaceæ. *De Cand.*

*aroma-
ticus.*

C. foliis ovato-oblongis utrinque acuminatis, cymis multifloris. De Cand. Prod. 3. p. 262. Blackw. t. 338. Bot. Mag. t. 2749, t. 2750. h

Myrtus Caryophyllus. Spreng. Syst. 2. p. 485.

Eugenia Caryophyllata. Willd. Sp. Pl. 3. p. 965. Hort. Kew. 3. p. 188.

Clove Tree. Miller.

EUGENIA.

Hesperidæ. *Lin.*—Myrti. *Juss.*—Myrtaceæ. *De Cand.*

ligustrina. E. pedicellis unifloris subsolitariis ex axillis foliorum aut bractearum elongatis subflore ebracteolatis, foliis lanceolatis obtusis subconcavis superne lucidis utrinque ramis floribusque glaberrimis. De Cand. Prod. 3. p. 263. Willd. Sp. Pl. 2. p. 962. h

Myrtus ligustrina. *Swtz. Fl.* 885.

Myrtus cerasina. *Willd. Sp. Pl.* 2. p. 968.

Arbor baccifera myrtifolia. *Sl.* 2. p. 107.

Black Cherry Tree. *Hughes.* 151?

Privet-leaved *Eugenia.* *Sweet.*

procera. *E. pedicellis unifloris axillaribus 2—4 confertis folio brevioribus sub flore bibracteolatis, foliis ovatis obtuse acuminatis ramisque glabris.* *De Cand. Prod.* 3. p. 268. h

Myrtus procera. *Willd. Sp. Pl.* 2. p. 968.
Swtz. Fl. 887.

Tall Myrtle. *Miller.*

triflora. *E. pedunculis axillaribus subternis trifloris, floribus ebracteatis sessilibus, foliis alternis* petiolatis oblongis basi attenuatis nitidis subtus viridi-cinereis.* *De Cand. Prod.* 3. p. 274. *Ham. Prod.* 44. h

Myrtus triflora. *Jacq. Amer.* 153. t. 193. f. 59.

Black Wood. *Hughes.* 180.

virgultosa. *E. racemis axillaribus et terminalibus puberulis subcompressis folio brevioribus, pedicellis subremotis decussatis sub flore bibracteolatis, foliis ovali-oblongis utrinque acuminatis subopacis*

* Ab omnibus differt ob folia alterna dicta! forsan tantum in summitate ramorum.—*De Cand.*

ramisque glabris, superne lucidis punctis
venisque impressis notatis. *De Cand.*

Prod. 3. p. 280. h

Myrtus virgultosa. *Willd. Sp. Pl.* 2. p.
972. *Hort. Kew.* 3. p. 190. *Swtz. Fl.* 905.

Black-Berry-Hunters. *Hughes.* 145.

Twiggy Myrtle. *Aiton.*

Rod-Wood.

Pimenta. E. pedunculis axillaribus et terminalibus
trichotomo-paniculatis, floribus 4—fidis
in dichotomiis subsessilibus; cæteris pa-
niculatis, foliis oblongis ovalibusve pellu-
cido-punctatis subopacis glabris, ramis
teretibus: ramulis compressis; junioribus
pedicellisque pubescentibus. *De Cand.*

Prod. 3. p. 285. h

Myrtus Pimenta. *Willd. Sp. Pl.* 2. p. 973.

Hort. Kew. 3. p. 191. *Aubl.* 503. *Bot.*

Mag. t. 1236.

Myrtus arborea aromatica. *Sl.* 2. p. 76.
t. 191.

Caryophyllus foliis oblongo-ovatis alter-
nis. *Br.* 247.

Pimento, Jamaica Pepper, or All-spice.

Long. 2. p. 702. *Miller.*

JAMBOSA.

Hesperidæ (*Eugenia*). *Lin.*—Myrti (*Eugenia*). *Juss.*

Myrtaceæ. *De Cand.*

macro- *J. cymis* lateralibus fasciculatis, foliis
phylla. ovato-lanceolatis basi obtusis apice acu-
minatis. *De Cand. Prod.* 3. p. 287. 6
J. sylvestris alba. *Rumph. Amb.* 1. p.
127. t. 39.*
Malacca Shambu. *Hort. Mal.* 27. t. 17.
Rose-Apple. Pomme-Rose.

* This synonym has been removed from *Eugenia Jambos*. *Willd.* to which it certainly cannot belong. Very great alterations have recently been made in the disposition of the members of the Myrtaceous family, and in the synonyms assigned to individuals. Rumphius describes the fruit of *Jambosa domestica*, which has by some been identified with *Eugenia Jambos*, by others with *E. malaccensis*, but which is considered by De Candolle as a distinct species, as follows:—"Caro ejus est succosa grata et tam jucunde spirans, ac si in aqua rosarum foret infusa, cujus sapor ex dulci et acido adeo grate mixtus est, ac si pomum esset cum aqua rosarum mixtum."—*Herb. Amb.* 1. p. 121. This description seems sufficiently applicable to the Rose-Apple of Barbados, but the fruit is oblong. Of the *Jambosa sylvestris* alba, he adds, "hujus fructus substantia durior est quam domesticæ Jambosæ, neque ita succosa, sapore cæterum conveniens sed paulo magis fatua."—*Herb. Amb.* 1. p. 128. This

PUNICA.

Pomaceæ. *Lin.*—Myrti. *Juss.*—Granateæ. *De Cand.*

Granatum. *P. foliis lanceolatis, caule arboreo.* *Willd.*
Sp. Pl. 2. p. 981. *Hort. Kew.* 3. p. 194.
De Cand. Prod. 3. p. 3. *Bot. Mag. t.*
 1832. h

P. fruticosa major. *Br.* 239.

Malus punica sativa. *Sl.* 2. p. 163.

Pomegranate Tree. *Hughes.* 130.
Browne.

Common Pomegranate Tree. *Miller.*

nana. *P. foliis linearibus, caule fruticoso.* *Willd.*
Sp. Pl. 2. p. 981. *Hort. Kew.* 3. p. 194.
De Cand. Prod. 3. p. 4. *Aubl.* 513.
Bot. Mag. t. 634. h

P. fruticosa humilior. *Br.* 239.

Small Pomegranate Tree. *Hughes.* 130.

Dwarf Pomegranate Tree. *Miller.*
Browne.

description and the plate of *Jambosa sylvestris alba* identify it with the Rose-Apple, and I unhesitatingly follow De Candolle in assigning it as a synonym to *Jambosa macrophylla*. The Malacca Schambu of Rheede would seem to apply to this species rather than to the *Jambosa vulgaris*, *De Cand.* "racemis cymosis terminalibus."

CHRYSOBALANUS.

Pomaceæ. *Lin.*—Rosaceæ. *Juss. De Cand.*

Chrysobalanæ. *Brown.*

- Icaco.* C. foliis subrotundis obovatisve emarginatis, racemis axillaribus dichotomis, staminibus hirsutis. *De Cand. Prod.* 2. p. 525. *Willd. Sp. Pl.* 2. p. 998. *Hort. Kew.* 3. p. 200. *Aubl.* 513. *Jacq. Amer.* 154. t. 94. h
- C. fruticosus. *Br.* 250.
- Frutex cotini fere folio. *Catesb. Car.* 1. p. 25. t. 25.
- Fat-Pork Tree. *Hughes.* 180.
- Cocco-Plum. *Long.* 2. p. 827.
- Cocoa-Plum. *Miller.*

CERASUS.

Hesperidæ. *Lin.*—Rosaceæ. *Juss. De Cand.*

- occidentalis.* C. racemis lateralibus, foliis eglandulosis oblongis acuminatis integerrimis utrinque glabris. *De Cand. Prod.* 2. p. 540. h

Prunus occidentalis. Willd. *Sp. Pl.* 2.
p. 987. Hort. Kew. 3. p. 196. Swtz.
Fl. 925.

Black Cherry. Hughes. 159 ?*

West Indian Cherry. Sweet.

sphæro- C. racemis axillaribus erectis minimis
carpa? folio brevioribus, foliis eglandulosis in-
tegerrimis nitidis, floribus distantibus,
fructibus subglobosis. De Cand. *Prod.*
2. p. 540. h

Prunus sphærocarpa. Willd. *Sp. Pl.* 2.
p. 987. Swtz. *Fl.* 927.

Myrtifolia Arbor. Sl. 2. p. 79. t. 193. f. 1.

Stopper-Berry Tree. Hughes. 176.

Round-fruited Cherry. Sweet.

* I am not certain that the Black Cherry Tree, Hughes. 159. be distinct from his Stopper-Berry Tree, the fruit of which is often called Black Cherries, from their resemblance to the European fruit of the same name. If it be distinct, it is probably the *Cerasus occidentalis*. De Cand. Of the fruit of this tree Swartz observes, "drupa ovalis pulposa monosperma, matura nigra." "Jamaicensibus Prune Tree."—*Fl.* 925. A comparison of the above description, with the representation in Catesby (*Car.* 2. t. 94.) would, I think, indicate that the trees respectively described by these authors cannot be identical, although M. De Candolle, under the head *Cerasus occidentalis*, refers, doubtingly however, to this plate of Catesby.—My notes of the Stopper-Berry Tree are, I am sorry to find, not quite so complete as I could have wished.

ICOSANDRIA TRIGYNIA.

SESUVIUM.

Succulentæ. *Lin.*—Ficoideæ. *Juss. De Cand.*

- Portulaca-* S. foliis lineari-aut-lanceolato-oblongis
castrum. planis, floribus pedicellatis. *De Cand.*
Prod. 3. p. 453. *Willd. Sp. Pl.* 2. p.
1009. *Hort. Kew.* 3. p. 205. *Jacq.*
Amer. 155. t. 95. *Bot. Mag. t.* 1701. 4
Portulaca aizoides maritima procum-
bens. *Sl.* 1. p. 205.
Aizoon repens. *Br.* 241.
Seaside Samphire. *Hughes.* 220.
Purslane-leaved Sesuvium. *Aiton.*

ICOSANDRIA PENTAGYNIA.

KERRIA.

Rosaceæ. *De Cand.*

japonica. *K. japonica.* *De Cand. Prod.* 3. p. 541. $\frac{1}{2}$

α . floribus simplicibus.

β . floribus plenis.

Corchorus japonicus. *Willd. Sp. Pl.* 2.

p. 1218. *Hort. Kew.* 3. p. 314. (*fl. pl.*)

Bot. Mag. t. 1296.

Japanese Kerria. *Sweet.*

ICOSANDRIA POLYGYNIA.

ROSA.*

Senticosæ. *Lin.*—Rosaceæ. *Juss. De Cand.*

*damas-
cena.*

R. calycibus semipinnatis, germinibus ovatis turgidis pedunculisque hispidis, caule petiolisque aculeatis, foliolis ovatis acuminatis subtus villosis. Willd. Sp. Pl. 2. p. 1072. Hort. Kew. 3. p. 263. De Cand. Prod. 2. p. 620. Lawr. Ros. t. 5. t. 18. t. 52. t. 70.

h

Damask Rose. *Miller.*

* "This is the most favorite genus of the whole vegetable kingdom, on account of its beauty and fragrance; but one most difficult with respect to the determination of its species."—*Sm. Engl. Fl. sub. verb.* The difficulties alluded to by this very able botanist have, I am afraid, not been yet successfully encountered; I have therefore been less solicitous to follow the alterations recently made in the arrangement of species and varieties. A few species introduced into Barbados from America, a short time before I quitted the island, have not been inserted, as I had no opportunity of examining them.

- sempervirens.* R. germinibus ovatis calycibus pedunculisque hispidis, caule petiolisque aculeatis, floribus subumbellatis, bracteis lanceolatis reflexiis. *Willd. Sp. Pl.* 2. p. 1072. *Hort. Kew.* 3. p. 263. *De Cand. Prod.* 2. p. 597. *Bot. Reg.* t. 465. *Lawr. Ros.* t. 45. h
 Evergreen Rose. *Miller.*
- rubiginosa.* R. fructibus ovatis pedunculisque hispidis, aculeis aduncis, foliolis ellipticis subtus glanduloso-rubiginosis. *Fl. Br.* 540. *Willd. Sp. Pl.* 2. p. 1073. *Hort. Kew.* 3 p. 264. *De Cand. Prod.* 2. p. 615. *Engl. Bot.* t. 992. *Lawr. Ros.* t. 56. h
 Sweet-Briar Rose. *Miller.*
- moschata.* R. fructibus ovatis pedunculisque villosis, caule petiolisque aculeatis, foliolis oblongis acuminatis glabris, pedunculis multifloris. *Willd. Sp. Pl.* 2. p. 1074. *Hort. Kew.* 3. p. 264. *De Cand. Prod.* 2. p. 598. h
 α . floribus simplicibus. *Lawr. Ros.* t. 64.
 Single Musk Rose. *Aiton.*
 β . floribus multiplicatis et plenis. *Lawr. Ros.* t. 53.
 Double Musk rose. *Aiton.*
- semperflorens* R. germinibus oblongis pedunculisque hispidis, caule petiolisque aculeato-his-

pidis, foliis subternatis aculeatis. *Willd.*
Sp. Pl. 2. p. 1078. *Hort. Kew.* 3. p.
 266. h

R. indica θ . *semperflorens.* *De Cand.*
Prod. 2. p. 601.

α . floribus simplicibus. *Ex. Bot.* t. 91.

β . floribus plenis. *Bot. Mag.* t. 284.
Lawr. Ros. t. 23.

Ever-blowing Rose. Dark Chinese
 Rose. *Aiton.*

Crimson ever-bearing Rose.

chinensis. *R. germinibus ovatis pedunculisque gla-*
bris, petiolis cauleque aculeatis, foliolis
ovato-lanceolatis subternatis serrulatis
glabris. *Willd. Sp. Pl.* 2. p. 1078. h

R. odorata. *Hort. Brit.* 138.

R. Indica β . *odorata.* *Bot. Reg.* t. 804.

R. Indica δ . *fragrans.* *De Cand. Prod.*
 2. p. 600.

Sweet-scented China Rose. *Sweet.*

indica. *R. germinibus ovatis pedunculisque gla-*
bris, caule subinermi, petiolis aculeatis.
Willd. Sp. Pl. 2. p. 1079. *Hort. Kew.*
 3. p. 266. *Lawr. Ros.* t. 26. h

R. indica. α . *vulgaris.* *De Cand. Prod.*
 2. p. 600.

Blush Chinese Rose. *Aiton.*

Common ever-bearing, or China Rose.

bracteata. R. germinibus obovatis pedunculis bracteatis ramulisque villosis, caule petiolisque aculeatis, foliolis glabris subrotundis crenatis subaculeatis. *Willd. Sp.*

Pl. 2. p. 10. 1079. *Hort. Kew.* 3. p. 267. *De Cand. Prod.* 2. p. 602. h

R. lucida. *Lawr. Ros. t.* 84.

Macartney Rose. *Sweet.*

Austrian Rose.

POLYANDRIA MONOGYNIA.

CAPPARIS.

Putamineæ. *Lin.*—Capparides. *Juss.*

cynophal- C. foliis glabris coriaceis oblongis brevi-
lophora. ter petiolatis, glandula axillari ovato-
oblonga, pedunculis paucifloris folio
brevioribus. *De Cand. Prod.* 1. p. 249.

Willd. Sp. Pl. 2. p. 1136. *Hort. Kew.* 3.
p. 285. h

C. cynallophora. *Jacq. Amer.* 158. t. 98.
Breynia fruticosa foliis oblongis. *Br.* 246.
t. 27. f. 1.

Acaciæ affinis arbor siliquosa. *Sl.* 2. p. 59.
Dog-Willow, Rat's-Bane, or Raw Head
and Bloody Bones. *Hughes.* 201.

Bay-leaved Capparitis. *Aiton.*

torulosa. C. foliis* ovato-lanceolatis coriaceis pe-
tiolatis superne glabris subtus ramulisque

* The leaves on the adult tree are ovate-lanceolate as repre-
sented; but on the young tree, when only 2—4 feet high, they

lepidotis; pedunculis paucifloris racemoso-corymbosis, staminibus 28, siliqua longissima. *De Cand. Prod.* 1. p. 252.

Willd. Sp. Pl. 2. p. 1134. *Swtz. Fl.* 932. h

Breynia arborescens. *Br.* 246.

Black Willow. *Hughes.* 196.*

Twisting Capparis. *Sweet.*

Breynia. C. foliis ellipticis coriaceis petiolatis superne glabris subtus ramulisque lepidotis, pedunculis angulatis racemoso-corymbosis, staminibus 16,† siliqua longissima. *De Cand. Prod.* 1. p. 252. *Willd. Sp. Pl.* 2. p. 1138. *Hort. Kew.* 3. p. 285. *Jacq. Amer.* 161. t. 103. h

are perfectly linear, the breadth of the middle increasing as the age of the plant advances.

* I cannot say that I know the Red Willow of Hughes, p. 196. as distinct from his Black Willow. If they be really different, and the *Capparis ferruginea* be growing in Barbados, it would doubtless be the Red Willow; but I have never seen this species in the island, nor the *Capparis longifolia*, the habitat of which latter is especially fixed by De Candolle in Barbados. The *C. torulosa* is generally known as the Black Willow, but I have frequently also heard that name applied to the *C. Breynia*.

† "Floribus octandris." *Willd.* "Stamina numerosa." *Jacq.* Previously to seeing the definition of M. De Candolle, and with the view of ascertaining where the error indicated by the above incongruity lay, I examined a great many flowers, and always found the number of stamens to be sixteen.

White Willow. *Hughes*. 196.

Oleaster-leaved Capparis. *Sweet*.

ARGEMONE.

Rheadæ. *Lin.*—Papaveraceæ. *Juss. De Cand.*

mexicana. *A. mexicana*. *De Cand. Prod.* 1. p. 120.

Willd. Sp. Pl. 2. p. 1148. *Hort. Kew.* 3.

p. 290. *Aubl.* 532. *Bot. Mag. t.* 243. ☉

A. spinosum. *Br.* 244.

Papaver spinosum. *Bauh. Pin.* 171.

Holly-Hock. *Hughes*. 208.

Yellow Thistle. *Long.* 2. p. 845.

Prickly Argemone, or Poppy. *Miller*.

NYMPHÆA.

Hydrocharides. *Juss.*—Nymphæaceæ. *De Cand.*

odorata. *N. foliis cordatis integerrimis subtus
nervis venisque prominentibus, stigmatibus*

16—20—radiato : radiis erectis apice inflexis. *De Cand. Prod.* 1. p. 116. *Willd.*

Sp. Pl. 2. p. 1153. *Hort. Kew.* 3. p. 292.

Bot. Mag. t. 819. *Bot. Repos.* t. 297. 2

N. alba major. *Sl.* 1. p. 252.

Large Duck-Weed. *Hughes.* 233.

Sweet-scented Water-Lily. *Aiton.*

BIXA.

Columniferæ. *Lin.*—Tiliaceæ. *Juss.*—Bixineæ. *De Cand.*

Orellana. *B. foliis utrinque glabris.* *De Cand. Prod.*

1. p. 259. *Willd. Sp. Pl.* 2. p. 1154.

Hort. Kew. 3. p. 296. *Aubl.* 533. *Bot.*

Mag. t. 1456. h

B. foliis cordatis cum acumine. *Br.* 254.

Arnotto Shrub. *Hughes.* 202.

Arnotto, or Roucou. *Long.* 2. p. 714.

Arnotto, or Arnotta. *Miller.*

MAMMEA.

Dubii ord. *Lin.*—Guttiferæ. *Juss. De Cand.*

americana. *M.* foliis obovatis obtusissimis integerimis, fructibus maximis tetraspermis.

De Cand. Prod. 1. p. 561. *Willd. Sp.*

Pl. 2. p. 1157. *Hort. Kew.* 3. p. 297.

Aubl. 917. *Jacq. Amer.* 268. t. 181.

f. 82.

h

M. foliis ovalibus nitidis. *Br.* 249.

Malus persica maxima. *Sl.* 2. p. 123.

t. 217. f. 3.

Mammee Tree. *Hughes.* 133. *Long.* 2.

p. 836.

American Mammee. *Miller.*

LAGERSTRÆMIA.

Dubii ord. *Lin.*—Salicariæ. *Juss.*—Lythrarieæ. *De Cand.*

indica. *L.* foliis subrotundo-ovatis acutis glabris,
panicula multiflora terminali, petalis

crispis longe unguiculatis. *De Cand.*
Prod. 3. p. 93. *Willd. Sp. Pl.* 2. p. 1178.
Hort. Kew. 3. p. 302. *Bot. Mag. t.* 405. $\frac{1}{2}$
 Indian Lagerstroemia. *Aiton.*
 King of Flowers.

Regina. L. foliis oblongis glabris, panícula termi-
 nali, calycibus tomentosis, petalis orbi-
 culatis undulatis breviter unguiculatis.
De Cand. Prod. 3. p. 93. *Willd. Sp. Pl.*
 2. p. 1178. *Hort. Kew.* 3. p. 302. *Roxb.*
Cor. 1. p. 46. t. 65. $\frac{1}{2}$
 Oblong-leaved Lagerstroemia. *Aiton.*
 Queen of Flowers.

CALOPHYLLUM.

Dubii ord. *Lin.*—*Aurantiae. Juss.*—*Guttiferæ. De Cand.*

Calaba. C. foliis ovatis obtusis, floribus her-
 maphroditis aut masculis, racemis late-
 ralibus brevioribus. *De Cand. Prod.* 1. p.
 562. *Willd. Sp. Pl.* 2. p. 1160. *Hort.*
Kew. 3. p. 298. *Jacq. Amer.* 269. t.
 165. $\frac{1}{2}$
 Arbor altissima. *Br.* 372.
 Mali Persicæ, &c. folio arbor maxima.
Sl. 2. p. 124.

Bastard Mammee, or Santa Maria.

Long. 2. p. 835.

Calaba Tree, or Santa Maria. *Aiton.*

CORCHORUS.

Columniferæ. *Lin.*—Tiliaceæ. *Juss. De Cand.*

siliquosus. C. capsulis linearibus compressis bivalvibus glabris, foliis lanceolatis æqualiter serratis, caule ramosissimo glabriusculo.

De Cand. Prod. 1. p. 504. Willd. Sp.

Pl. 2. p. 1218. Hort. Kew. 3. p. 314.

Aubl. 578. Plum. Ic. t. 103. f. 1. h

Corchoro affinis chamædryos folio. *Sl.*

1. p. 145. t. 94. f. 1.

Coreta foliis minoribus. *Br. 147.*

Broom-Weed.* *Hughes. 171. Browne.*

Germander-leaved Corchorus. *Miller.*

hirtus. C. capsulis linearibus compressis bivalvibus cauleque pilosis, foliis oblongis æqualiter serratis. *De Cand. Prod. 1. p.*

504. Willd. Sp. Pl. 2. p. 1218. Aubl.

579. Plum. Ic. t. 103. f. 2. h

* This plant, as well as the Sidas, hereafter to be noticed, obtains the name of Broom in Barbados and Jamaica, being commonly employed as besoms.

Rata-Pepper, or Rata-Ockro, *alias* Hoho.

Hughes. 211.

Hairy Corchorus. *Sweet.*

æstuans.

C. capsulis oblongis 3—valvibus 6—sulcatis, foliis subcordatis ovato-acuminatis serratis: serraturis infimis longe acuminatis. De Cand. Prod. 1. p. 504. Willd.

Sp. Pl. 2. p. 1215. Hort. Kew. 3. p. 313. ☉

Alcea cibaria, seu corchorus americana.

Pluk. Phyt. t. 127. f. 3.

Triumfetta subvillosa. Br. 232. t. 25. f. 1.

Papaw-Ockroe. Hughes. 197.

Hornbean-leaved Corchorus. *Miller.*

tridens.

C. capsulis gracilibus 2—3—valvibus 2—3—angulatis 2—3—cornutis, foliis oblongis serratis: serraturis infimis setaceis. De Cand. Prod. 1. p. 505. Willd.

Sp. Pl. 2. p. 1215. ☉

C. americanus, angusto barbato folio.

Pluk. Phyt. t. 127. f. 4.

POLYANDRIA POLYGYNIA.

NELUMBIUM.

Hydrocharides. *Juss.*—*Nymphæaceæ. De Cand.*

- jamai-* *N. corolla polypetala, antheris , fo-*
cense. *liis peltatis subtus radiatis. M—. De*
 Cand. Prod. 1. p. 114. 24
N. speciosum. γ. Willd. Sp. Pl. 2. p.
 1258. Hort. Kew. 3. p. 332.
Nymphæa foliis orbiculatis. Br. 243.
Broad Pond Duck-Weed. Hughes. 222.
Ægyptian Bean, or Great Water-Lily.
 Browne.
Jamaica Sacred Bean. Sweet.

ANONA.

Coadunatæ. *Lin.*—Anonæ. *Juss.*—Anonaceæ. *De Cand.*

muricata. A. foliis ovato-lanceolatis glabris subnitidis, pedunculis solitariis unifloris, petalis exterioribus cordatis acuminatis; interioribus obtusis, fructibus muricatis mucronibus carnosus. *De Cand. Prod.* 1.

p. 84. *Willd. Sp. Pl.* 2. *p.* 1264. *Hort.*

Kew. 3. *p.* 333, *Aubl.* 617. h

A. foliis oblongo-ovatis nitidis. *Br.* 255.

A. maxima foliis latis splendentibus. *Sl.*

2. *p.* 166. *t.* 225.

Sour-sop Tree. *Hughes.* 131. *Browne.*

Rough-fruited Custard Apple, or Sour-sop. *Miller.*

palustris. A. foliis ovato-oblongis coriaceis glaberrimis, floribus solitariis pedunculatis, petalis omnibus acutis, fructibus subareolatis. *De Cand. Prod.* 1. *p.* 84. *Willd.*

Sp. Pl. 2. *p.* 1267. *Hort. Kew.* 3.

p. 335. h

A. religiosa. *Br.* 256.

A. aquatica. *Sl.* 2. *p.* 169. *t.* 228. *f.* 1.

Monkey-Apple Tree. *Hughes.* 129.

Aligator-Apple, or Cork-Wood. *Long.*
2. p. 832.

Shining-leaved Custard-Apple. *Miller.*
squamosa. A. foliis lanceolatis glabris pellucido-
punctatis, petalis exterioribus subclausis,
fructibus ovoideis squamosis. *De Cand.*

Prod. 1. p. 85. *Willd. Sp. Pl.* 2. p. 1265.

Hort. Kew. 3. p. 334. *Aubl.* 617. h

A. foliis ovatis undulatis. *Br.* 256.

A. foliis odoratis minoribus. *Sl.* 2. p. 168.
t. 227.

Sugar-Apple Tree. *Hughes.* 129.

Undulated Custard-Apple, or Sweet Sop.
Miller.

Sweet Sop, or Sugar-Apple Tree.
Browne.

reticulata. A. foliis oblongo-lanceolatis acutis gla-
bris subpunctatis, petalis exterioribus
oblongis subclausis, fructibus ovato-glo-
bosis reticulato-areolatis. *De Cand. Prod.*

1. p. 85. *Willd. Sp. Pl.* 2. p. 1265.

Hort. Kew. 3. p. 334. *Aubl.* 617. h

A. foliis oblongis undulatis venosis.
Br. 256.

A. maxima foliis oblongis angustis. *Sl.*
2. p. 167. *t.* 226.

Custard-Apple Tree. *Hughes.* 140.
Browne.

Netted Custard-Apple. *Miller.*

DIDYNAMIA GYMNOSPERMIA.

SATUREJA.

Verticillatæ. *Lin.*—Labiatae. *Juss. Brown.*

hortensis. S. pedunculis axillaribus subcymosis,
foliis lanceolatis integerrimis, caule bra-
chiato. *Willd. Sp. Pl. 3. p. 45. Hort.*
Kew. 3. p. 374. ☉
Summer Savoury. *Miller.*

NEPETA.

Verticillatæ. *Lin.*—Labiatae. *Juss. Brown.*

Cataria. N. floribus spicatis, verticillis subpedi-
cellatis, foliis petiolatis cordatis dentato-
serratis. *Fl. Br. 608. Willd. Sp. Pl. 3.*
p. 49. Hort. Kew. 3. p. 377. Engl.
Bot. t. 137. 24

Mentha Cataria. *Bauh. Pin.* 228
Common Catmint. *Miller.*

LAVANDULA.

Verticillatæ. *Lin.*—*Labiatae.* *Juss. Brown.*

Stæchas. *L. foliis sessilibus linearibus tomentosis
 margine revolutis, spica coarctata co-
 mosa, bracteis subtrilobis.* *Willd. Sp.*
Pl. 3. p. 60. Hort. Kew. 3. p. 382. ½
French Lavender. *Miller.*

BYSTROPOGON.

Verticillatæ (Clinopodium). *Lin.*—*Labiatae.* *Juss. Brown.*

pectina- *B. paniculis compactis floribus secundis,*
tum. *foliis ovatis.* *Willd. Sp. Pl. 3. p. 72. ☉*
Hyptis pectinata. *Hort. Kew. 3. p. 392.*
Poiteau An. du Mus. 7. p. 474. t. 30,
Balm-leaved Bystropogon. *Miller.*
Wild Spikenard. *Browne.*

- suaveolens.* B. pedunculis axillaribus solitariis, calycibus truncatis aristatis, foliis cordatis.
Willd. Sp. Pl. 3. p. 72. ☉
 Ballota suaveolens. *Jacq. Amer. 172.*
 Hyptis suaveolens. *Poiteau An. du Mus. 7. p. 472. t. 29. f. 2.*
 Hyptis ebracteata. *Hort. Kew. 3 p. 391.*
 Mesosphærum hirsutum. *Br. 257. t. 18. f. 3.*
 Mentastrum maximum. *Sl. 1. p. 171. t. 102. f. 2.*
 Musketo-Bush, or White Hop-Weed.
Hughes. 169.
 Sweet-scented Bystropogon. *Miller.*
 Spikenard. *Browne.*

MENTHA.

Verticillatæ. *Lin.*—Labiatae. *Juss. Brown.*

- viridis.* M. spicis interruptis, foliis sessilibus lanceolatis acutis nudis, bracteis setaceis dentibusque calycinis subhirsutis. *Fl. Br. 612. Hort. Kew. 3. p. 388. Willd. Sp. Pl. 3. p. 76. Menth. Br. 11. t. 5. 4*
 Spear-Mint. *Miller.*

- piperita.* M. spicis obtusis inferne interruptis, foliis petiolatis subovatis glabriusculis, calyce basi glaberrimo. *Fl. Br.* 613.
Hort. Kew. 3. p. 388. *Willd. Sp. Pl.* 3. p. 79. *Engl. Bot. t.* 687. 4
 M. *piperita* officinalis. *Menth. Br.* 15. t. 7.
 Pepper-Mint. *Miller.*
- Pulegium.* M. floribus verticillatis, foliis ovatis, caule prostrato, pedicellis calycibusque undique tomentosis, dentibus ciliatis. *Fl. Br.* 614. *Hort. Kew.* 3. p. 390. *Willd. Sp. Pl.* 3. p. 82. *Engl. Bot. t.* 1026.
Menth. Br. 51. t. 23. 4
 Pennyroyal. *Miller.*

HYPTIS.

Verticillatæ (*Clinopodium*). Labiatæ. *Juss. Brown.*

- capitata.* H. capitulis oppositis pedunculis internodii longitudine, bracteis lanceolatis calyce frugifero brevioribus, foliis oblongis utrinque attenuatis (*subangulatis*) inæqualiter serratis. *Poiteau An. du Mus.* 7. p. 464. t. 27. f. 1. *Hort. Kew.* 3. p. 391. *Willd. Sp. Pl.* 3. p. 84. *Jacq. Ic.* 1. t. 114. 3

Clinopodium capitatum. *Swtz. Prod.* 88.

Clinopodium subhirsutum. *Br.* 259.

Sideritis spicata. *Sl.* 1. *p.* 174. *t.* 109. *f.* 2.

Jamaica Hyptis. *Aiton.*

Wild Hops. *Browne.*

STACHYS.

Verticillatæ. Lin.—Labiatae. Juss. Brown.

circinata. *S.* verticillis spicatis sexfloris, bracteis cordatis, foliis cordatis petiolatis obtusis crenato-dentatis. *Willd. Sp. Pl.* 3. *p.* 97.

Hort. Kew. 3. *p.* 398. 24

S. canariensis. *Jacq. Ic.* 1. *t.* 108.

Blunt-leaved Stachys. *Aiton.*

LEONURUS.

Verticillatæ. Lin.—Labiatae. Juss. Brown.

sibiricus. *L.* foliis tripartitis: laciniis subincisis, calycibus inermibus, labio corollæ su-

periore concavo recto. *Willd. Sp. Pl.* 3.
p. 117. *Hort. Kew.* 3. *p.* 406. *Ex.*
Bot. t. 94. 2
 Siberian Motherwort. *Miller.*

LEUCAS.

Verticillatæ (*Phlomis*). *Lin.*—*Labiatae*. *Juss. Brown.*

martini-
censis.

L. foliis oblongis dentatis subtus pubes-
centibus, verticillis multifloris globosis,
calycibus incurvis decemdentatis: dente
supremo longissimo. Hort. Kew. 3. *p.*
 409. ☉

Phlomis martinicensis. Willd. Sp. Pl. 3.
p. 123.

Phlomis caribæa. Swtz. 1009. *Jacq. Ic.*
 1. *t.* 110.

Clinopodium martinicense. Jacq. Amer.
 173. *t.* 177. *f.* 75.

West Indian Leucas. *Aiton.*

LEONOTIS.

Verticillatæ (*Phlomis*). *Lin.*—*Labiatae*. *Juss.* *Brown.*

nepetifolia. *L.* foliis cordatis acutis inciso-crenatis, calycibus aristatis octodentatis: dente supremo maximo, caule herbaceo. *Hort.*
Kew. 3. p. 409. *Bot. Reg.* t. 281. ☉
Phlomis nepetifolia. *Willd. Sp. Pl.* 3. p. 126.
 Catmint-leaved Leonotis. *Aiton.*

ORIGANUM.

21

Verticillatæ. *Lin.*—*Labiatae*. *Juss.* *Brown.*

Majorana. *O.* spicis subrotundis ternis compactis pedunculatis, foliis petiolatis ellipticis obtusis glabriusculis, radice annua.
Willd. Sp. Pl. 3. p. 137. *Hort. Kew.* 3. p. 413. ☉
Majorana vulgaris. *Bauh. Pin.* 224.
 Sweet, or Knotted Marjoram. *Miller.*

THYMUS.

Verticillatæ. *Lin.*—*Labiatae. Juss. Brown.*

vulgaris. T. erectus, foliis revolutis ovatis, floribus
verticillato-spicatis. *Willd. Sp. Pl. 3. p.*
139. *Hort. Kew. 3. p. 414.* h
Garden Thyme. *Miller.*

OCYUM.

Verticillatæ. *Lin.*—*Labiatae. Juss. Brown.*

Basilicum. O. foliis ovatis glabris, calycibus ciliatis.
Willd. Sp. Pl. 3. p. 161. Hort. Kew. 3.
p. 423. Aubl. 624. ⊙
Common Sweet Basil. *Miller.*
Garden Basil.

americana- P. foliis sublanceolatis acuminatis sub-
num. serratis, racemis teretibus, caule subher-
baceo. *Willd. Sp. Pl. 3. p. 163. Hort.*
Kew. 3. p. 423. Aubl. 625. Jacq. Vind.
3. p. 45. t. 86. ⊙
American Basil. *Miller.*
Wild Basil.

DIDYNAMIA ANGIOSPERMIA.

GLOXINIA.

Campanulaceæ. *Juss.*—Gesnereæ. *Juss. An. du Mus.*

- speciosa.* G. subacaulis, foliis hispidis crenatis, pedunculis erectis flore longioribus. *Bot. Mag. t. 1937. Bot. Reg. t. 213. 4*
Many-flowered Gloxinia. *Sweet.*

BESLERIA.

Personatæ. *Lin.*—Scrophulariæ. *Juss.*

Gesnereæ. *Juss. An. du Mus.*

- pulchella.* B. foliis oblongo-ovatis rugosis crenatis in petiolum decurrentibus, petiolis connatis, pedunculis axillaribus corymbi-

feris, involucro pedicellis brevior, calycibus pentagonis serrulatis coloratis.

Hort. Kew. 4. p. 21. *Bot. Mag.* t. 1146. $\mathfrak{h}$
Large-flowered Besleria. *Sweet.*

BIGNONIA.

Personatæ. *Lin.*—Bignoniæ *Juss.*—Bignoniaceæ. *Brown.*

Unguis. B. foliis conjugatis cirrosis: foliolis ovatis acuminatis, pedunculis axillaribus unifloris. *Willd. Sp. Pl.* 3. p. 292. *Hort.*

Kew. 4. p. 31. *Aubl.* 657. $\mathfrak{h}$

Clematis quadrifolia. *Plum. Amer.* 80. t. 94.

Apocyno affine gelsimum. *Sl.* 1. p. 208.

Right-Wyth, or Cat's-Claw Wyth.

Hughes. 244.

Barbados Trumpet-Flower. *Aiton.*

Leucoxy- B. foliis digitatis: foliolis lanceolatis acu-
lon. minatis integerrimis glabris, floribus terminalibus solitariis. *Willd. Sp. Pl.* 3.

p. 300. *Hort. Kew.* 4. p. 33. *Aubl.* 658.

Bot. Repos. t. 43. $\mathfrak{h}$

B. pentaphylla arborea. *Br.* 263.

Nerio affinis arbor. *Sl.* 2, p. 62.

White-wood Tree. *Hughes.* 192.

Smooth five-leaved Trumpet-Flower, &c.
Miller.

stans. White Cedar, or White-Wood. *Browne.*

B. foliis pinnatis: foliolis oblongo-lanceolatis serratis, racemo simplici terminali, caule erecto. Willd. Sp. Pl. 3. p. 302.

Hort. Kew. 4. p. 34. h

B. fraxini foliis. Catesb. Car. 1. p. 65. t. 65.

B. fruticosa, foliis pinnatis. Br. 264.

Apocyno affine gelsiminum. *Sl. 2. p. 63.*

Branching - flowered Trumpet-Flower.
Miller.

Ash-leaved Shrubby Bignonia. *Browne.*

CRESCENTIA.

Putamineæ. *Lin.*—Solaneis aff. *Juss.*

Cujete. *C. foliis cuneato-lanceolatis confertis.*

Willd. Sp. Pl. 3. p. 311. Hort. Kew. 4. p. 37. Aubl. 664. Jacq. Amer. 175. t. 111. h

C. arborescens, foliis confertis. Br. 265.

Calabash Tree. *Hughes. 116. Browne.*

Narrow-leaved Calabash. *Miller.*

CITHAREXYLUM.

Personatæ. *Lin.*—Vitices. *Juss.*

Verbenaceæ. *Juss. An. du Mus. Brown.*

- cinereum.* C. ramis teretibus, foliis oblongis acuminatis integerrimis,* racemis pendulis, calycibus dentatis. *Willd. Sp. Pl. 3. p. 308. Hort. Kew. 4. p. 37.* h
C. teres. *Jacq. Amer. 185. t. 118.*
C. fruticosum, cortice cinereo. *Br. 264.*
Fiddle-Wood Tree.† *Hughes. 146.*
Ash-coloured Fiddle-Wood. *Miller.*
Old Woman's Bitter. *Browne.*

* The young branches are generally quadrangular, and the leaves, when of luxuriant growth, are very frequently dentated.

† Hughes states the Fiddle-Wood to be "distinguished into the white, the red, and the black sort." His description, however, applies exclusively to the *Citharexylum cinereum*.

PETRÆA.

Personatæ. *Lin.*—Vitices. *Juss.*

Verbenaceæ. *Juss. An. du Mus. Brown.*

volubilis. P. caule fruticoso volubili, foliis elliptico-oblongis coriaceis utrinque scabris subtus venoso-reticulatis, racemis laxis elongatis, laciniis calycinis exterioris spathulatis obtusis reticulatis. *Spreng. Syst. 2. p. 761. Willd. Sp. Pl. 3. p. 313. Hort. Kew. 4. p. 38. Aubl. 664. Jacq. Amer. 180. t. 114. Bot. Mag. t. 628. ½ Climbing Blue-flowered Petræa. Aiton.*

LANTANA.

Personatæ. *Lin.*—Vitices. *Juss.*

Verbenaceæ. *Juss. An. du Mus. Brown.*

mista. L. foliis oppositis ovatis acutis pilosis, caule inferne aculeato, floribus capitatis: capitulis subrotundis, bracteis lanceolatis. *Willd. Sp. Pl. 3. p. 315. Hort. Kew. 4. p. 42. Bot. Cab. t. 68. ½*

Camara, &c. flore misto. *Dill. Elth.* 64.
t. 56. f. 64.

Various-flowered Lantana. *Miller.*

*involu-
crata.*

*L. foliis oppositis ternisque rhombeo-
ovatis obtusis tomentosis, caule inermi,
capitulis squamosis, bracteis ovatis.*

Willd. Sp. Pl. 3. p. 318. *Hort. Kew.*
4. p. 43. *Aubl.* 665. h

L. frutescens, &c. floribus subcarneis.
Br. 268.

Periclimenum rectum salviæ folio. *Sl.* 1.
p. 39.

White Sage. *Hughes.* 155.

Round-leaved Lantana. *Miller.*

Wild Sage with white* Flowers. *Browne.*

Rock-Sage. Button-Sage.

aculeata.

*L. foliis oppositis ovatis subcordatis
subtus molliusculis, caule aculeato, ca-
pitulorum bracteis lineari-cuneiformi-
bus.* *Willd. Sp. Pl.* 3. p. 320. *Hort. Kew.*

4. p. 44. *Aubl.* 665. *Bot. Mag.* t. 96. h

L. frutescens spinosa. *Br.* 269.

Prickly Lantana. *Miller. Browne.*

* Although Browne here says with "white flowers," in his definition he more properly describes the plant "*floribus subcarneis.*" Miller observes, "the yellow colour of the throat soon changes to white, and hence the flower is whitish, with a pale flesh-coloured margin."

CAPRARIA.

Personatæ. *Lin.*—Scrophulariæ. *Juss.*—Scrophularinæ. *Brown.*

- biflora.* *C. glabra*, foliis alternis lanceolatis serratis, floribus geminis, calycis laciniis sububulatis, corollæ fauce imberbi. *Hort. Kew.* 4. p. 45. *Willd. Sp. Pl.* 3. p. 323. *Jacq. Amer.* 182. t. 115. h
- C. erecta ramosa.* *Br.* 268.
- West-India Tea. *Long.* 2. p. 806.
- Shrubby Goat-Weed, or Sweet-Weed. *Miller.*

BROWALLIA.

Luridæ. *Lin.*—Scrophulariæ. *Juss.*—Scrophularinæ. *Brown.*

- elata.* *B. peduculis unifloris multiflorisque.*
Willd. Sp. Pl. 3. p. 339. *Hort. Kew.* 4.
 p. 51. *Bot. Mag.* t. 34 ⊙
- Upright Browallia. *Miller.*

APHELANDRA.

Personatæ (*Justicia*). *Lin.*—Acanthi (*Justicia*). *Juss.*

Acanthaceæ. *Brown.*

- cristata.* A. foliis ellipticis oblongisve acuminatis, spicis tetragonis, bracteis ovatis integerrimis, corollis glabris. *Hort. Kew.* 4. p. 55. *Bot. Mag.* t. 1578. h
Ruellia cristata. *Bot. Repos.* t. 506.
Justicia pulcherrima. *Willd. Sp. Pl.* 1. p. 86. *Aubl.* 15. *Jacq. Amer.* 6. t. 2. f. 4. *Jacq. Ic.* 2. t. 204.
Justicia tetragona. *Willd. Sp. Pl.* 1. p. 85.
 Dense-spiked *Aphelandra.* *Sweet.*

RUELLIA.

Personatæ. *Lin.*—Acanthi. *Juss.*—Acanthaceæ. *Brown.*

- clandestina.* R. foliis petiolatis oblongis obtusis basi attenuatis subdentatis, pedunculis trifloris foliis brevioribus. *Willd. Sp. Pl.* 3. p. 365. *Hort. Kew.* 4. p. 57. u
 Three-flowered *Ruellia* *Sweet.*

- tuberosa.* R. foliis cuneato-ovatis crenatis, pedunculis tripartitis, caule simplici. *Willd. Sp. Pl.* 3. p. 368. *Hort. Kew.* 4. p. 57. *Aubl.* 670. 2
- R. erecta, asphodeli radice. *Br.* 268.
- Gentianella flore cæruleo. *Sl.* 1. p. 149. t. 95. f. 1.
- Many-Roots. *Hughes.* 210.
- Tuberose-rooted Ruellia. *Sweet.*
- Menow-Weed, Spirit-Weed, and Snapdragon. *Browne.*
- formosa.* R. foliis petiolatis integerrimis ovatis utrinque pubescentibus, pedunculis axillaribus alternis longissimis paucifloris, corollis subringentibus. *Hort. Kew.* 4. p. 58. *Bot. Mag. t.* 1400. 2
- Splendid Ruellia. *Sweet.*

DURANTA.

Personatæ. *Lin.*—Vitices. *Juss.*

Verbenacæ. *Juss. An. du Mus. Brown.*

- Ellisia.* D. calycibus frutescentibus erectis. *Willd. Sp. Pl.* 3. p. 380. *Hort. Kew.* 4. p. 59. *Jacq. Amer.* 187. t. 170. f. 77. *Bot. Mag. t.* 1759. 2

- D. Plumieri.* *Bot. Reg. t. 244?*
Ellisia frutescens. *Br. 262. t. 29. f. 1.*
Jasminum folio integro. *Sl. 2. p. 97.*
Prickly Duranta. Aiton.*

CLERODENDRUM.

Personatæ. *Lin.*—Vitices. *Juss.*

Verbenacæ. *Juss. An. du Mus. Brown.*

Siphonanthus. *C. foliis verticillatis elongato-lanceolatis
 glabris, pedunculis axillaribus corym-
 bosis, tubo corollæ longissimo. Spreng.
 Syst. 2. p. 758.* h

* This plant, although commonly called Prickly Duranta, is not always found so. Browne describes it “quandoque spinosa,” although in his plate he represents it with thorns; and, according to Willdenow, “variat caule spinoso et inermi, foliis serratis et integerrimis.” The Duranta in Barbados more commonly occurs unarmed, and with a leaf quite entire; but I have seen it with pretty long thorns, which are hardened flower stalks, and on the same plant leaves entire and serrate. The Duranta *Ellisia* is exhibited in the Bot. Mag. with spines and a serrated leaf, and is certainly less like the Duranta as it occurs in Barbados than the plate in Bot. Reg. of *Duranta Plumieri*. I cannot, however, persuade myself that *D. Ellisia* and *D. Plumieri* are distinct species.

Siphonanthus angustifolia. Willd. *Sp. Pl.* 1. p. 606.

Whorl-leaved Clerodendrum. Sweet.

fragrans. C. foliis latis subcordato-ovatis repandodentatis basi biglandulosis subtus tomentosis, corymbis terminalibus densis hemisphæricis, tubo corollæ calycem duplo excedente. Spreng. *Syst.* 2. p. 760. Hort. Kew. 4. p. 63. (*fl. plenis.*) Bot. Mag. t. 1834. h

Fragrant Clerodendrum. Sweet.

VOLKAMERIA.

Personatæ. Lin.—Vitices. Juss.

Verbenacæ. Juss. An. du Mus. Brown.

aculeata. V. foliis oblongis acutis integerrimis, spinis e petiolorum rudimentis. Willd. *Sp. Pl.* 3. p. 383. Hort. Kew. 4. p. 62. *Aubl.* 673. h

Clerodendrum fruticosum. Br. 262. t. 30. f. 2.

Paliuro affinis ligustrifolia spinosa. Sl. 2. p. 25. t. 166. f. 2, 3.

Prickly Volkameria. Miller.

Wild Pomegranate.

AVICENNIA.

Personatæ. *Lin.*—Vitices. *Juss.*—Myoporinæ. *Brown.*

tomentosa. A. foliis oblongis obtusis subtus tomentosis. *Willd. Sp. Pl. 3. p. 395. Hort. Kew. 4. p. 62. Jacq. Amer. 178. t. 112. f. 2.* h

Bontia foliis integris. *Br. 263.*

Mangle laurocerasi foliis. *Sl. 2. p. 66.*

Dwarf Mangrove Tree. *Hughes. 147 ?*

Downy-leaved Avicennia. *Sweet.*

Olive Mangrove. *Browne.*

BONTIA.

Personatæ. *Lin.*—Solaneis aff. *Juss.*—Myoporinæ. *Brown.*

daphnoides. B. daphnoides. *Willd. Sp. Pl. 3. p. 394. Hort. Kew. 4. p. 61. Aubl. 673.* h

Olea sylvestris barbadensis. *Pluk. Alm. 269. Phyt. t. 209. f. 3.*

Wild Olive. *Hughes. 177. t. 12.*

Barbados Wild Olive. *Miller.*

THUMBERGIA.

Acanthi. *Juss.*—Acanthaceæ. *Brown.*

fragrans.* *T. foliis cordatis acuminatis basi subangulato-dentatis, caule scandente. Willd.*

Sp. Pl. 3. p. 388. Hort. Kew. 4. p. 66.

Roxb. Cor. 1. p. 47. t. 67. Bot. Mag. t. 1881. Bot. Repos. t. 123. h

Twining Thumbergia. *Aiton.*

* “The flowers are quite scentless, nor have we observed any fragrance in any part of the plant.” “Perhaps the heat of a tropical sun is necessary to bring forth its odour.”—*Sims in the Bot. Mag.* The odour is certainly not elicited by the great heat sometimes prevailing in Barbados, and the stem is not simply scandent, but twining. For these reasons the specific appellation should be changed from *fragrans* to *volubilis*.

TETRADYNAMIA SILICULOSA.

LEPIDIUM.

Siliquosæ. *Lin.*—Cruciferæ. *Juss. De Cand.*

- virginicum.* L. siliculis orbiculatis emarginatis pedicello brevioribus, floribus 2—4—andris, foliis caulinis lineari-lanceolatis inciso-serratis glabris. *De Cand. Prod.* 1. p. 205.
Willd. Sp. Pl. 3. p. 440. *Hort. Kew.* 4 p. 89. *Aubl.* 675. ☉
- L. erectum ramosum. *Br.* 272.
- Iberis humilior annua virginica. *Sl.* 1. p. 195. *t.* 123. *f.* 3.
- Wild Pepper-Grass. *Hughes.* 162.
- Virginian Pepper-Wort. *Miller.*
- Upright branching Pepper - Grass.
Browne.

COCHLEARIA.

Siliquosæ. *Lin.*—Cruciferæ. *Juss. De Cand.*

- Armoracia.* C. siliculis ellipsoideis, foliis radicalibus oblongis crenatis; caulinis elongato-lanceolatis dentatis incisive, radice carnosa maxima. *De Cand. Prod.* 1. p. 173. *Fl. Br.* 690. *Willd. Sp. Pl.* 3. p. 451. *Hort. Kew.* 4. p. 90. *Engl. Bot. t.* 2323. 4
Raphanus rusticus. *Bauh. Pin.* 96.
 Horse-Radish. *Miller.*

NASTURTIIUM.

Siliquosæ. *Lin.*—Cruciferæ. *Juss. De Cand.*

- officinale.* N. foliis pinnatisectis: segmentis ovatis subcordatis repandis. *De Cand. Prod.* 1. p. 137. *Hort. Kew.* 4. p. 110. *Engl. Fl.* 3. p. 193. 4
 N. aquaticum vulgare. *Sl.* 1. p. 193.
Sisymbrium Nasturtium. *Willd. Sp. Pl.* 3. p. 489. *Fl. Br.* 700. *Engl. Bot. t.* 855.
Sisymbrium aquaticum. *Br.* 272.
 Water-Cress. *Sweet.*

TETRADYNAMIA SILIQUOSA.

BRASSICA.

Siliquosæ, *Lin.*—Cruciferae, *Juss. De Cand.*

oleracea.* *B.* foliis polline glaucis subcarnosis repandis lobatisve etiam junioribus glaberrimis. *De Cand. Prod.* 1. p. 213. *Fl Br.* 720. *Willd. Sp. Pl.* 3. p. 548. *Hort. Kew.* 4. p. 123. *Engl. Bot. t.* 637. ♂

α. *B. capitata alba.* *Bauh. Pin.* 111.
Heading White Cabbage. *Miller.*

β. *B. capitata rubra.* *Bauh. Pin.* 111.
Red Cabbage. *Miller.*

* “Species a vetustissimis temporibus culta, et ideo valde polymorpha.”—*De Cand.* So few varieties are found, in Barbados, that I have not thought it necessary to follow the arrangement of varieties given by M. De Candolle.

γ. *B. alba vel viridis. Bauh. Pin. 111.*
 Indian Cale, or Seven Year's Cab-
 bage. *Hughes. 221.*
 Green Colewort. *Miller.*

Rapa. *B. foliis radicalibus lyratis polline glauco*
destitutis setoso-scabris; caulinis mediis
incisis; summis integerrimis lævibus.
De Cand. Prod. 1. p. 214. Fl. Br. 719.
Willd. Sp. Pl. 3. p. 548. Hort. Kew.
4. p. 123. Engl. Bot. t. 2176. ♂
 Turnip. *Miller.*

SINAPIS.

Siliquosæ. Lin.—Cruciferae. Juss. De Cand.

nigra. *S. siliquis glabris lævibus subtetragonis*
pedunculo appressis, foliis infimis lyratis;
summis lanceolatis integerrimis petio-
latis. De Cand. Prod. 1. p. 218. Fl. Br.
722. Willd. Sp. Pl. 3. p. 555. Hort.
Kew. 4. p. 127. Engl. Bot. t. 969. ☉
Sinapi Rapi folio. Bauh. Pin. 99.
 Common, or Black Mustard. *Miller.*

alba. *S. siliquis hispidis patentibus rostro en-*
siformi subangustioribus, foliis lyratis

caulique subglabris. *De Cand. Prod.* 1.
p. 220. *Fl. Br.* 721 *Willd. Sp. Pl.* 3.
p. 555. *Hort. Kew.* 4. *p.* 126. *Engl.*
Bot. t. 1677. ☉

Sinapi Apii folio. *Bauh. Pin.* 99.

White Mustard. *Miller* }

RAPHANUS.

Siliquosæ. *Lin.*—Cruciferæ. *Juss. De Cand.*

sativus. R. siliquis teretibus torosis acuminatis
vix pedicello longioribus. *De Cand.*
Prod. 1. *p.* 228. *Willd. Sp. Pl.* 3. *p.* 560.
Hort. Kew. 4. *p.* 129. ☉

α. *rotunda.*

R. major orbicularis vel rotundus.
Bauh. Pin. 96.

Turnip Radish. *Aiton.*

β. *oblonga.*

R. minor oblongus. *Bauh. Pin.* 96.

Common Garden Radish. *Aiton.*

GYNANDROPSIS.

Capparides. *Juss.*—Capparideæ. *De Cand.*

*penta-
phylla.*

G. glabriuscula, foliis mediis 5—foliolatis; infimis floralibusque 3—foliolatis: foliolis integris subserrulatisve. *De Cand.*

Prod. 1. p. 238.

⊙

Cleome pentaphylla. *Willd. Sp. Pl.* 3. p. 564. *Hort. Kew.* 4. p. 130. *Aubl.* 677. *Bot. Mag. t.* 1681.

Cleome procumbens pentaphylla. *Br.* 273.

Sinapistrum indicum pentaphyllum. *Sl.* 1. p. 194.

Coyers. *Hughes.* 210.

Five-leaved Gynandropsis. *Sweet.*

Sambo. *Browne.*

MONADELPHIA TRIANDRIA.

TAMARINDUS.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

- indica.* *T. indica.* *Willd. Sp. Pl.* 3. p. 577. *Hort.*
 Kew. 4. p. 134. *Aubl.* 24. *Jacq. Amer.*
 10. t. 10. h
T. indica. *De Cand. Prod.* 2. p. 488? *
T. occidentalis. *De Cand. Prod.* 2. p. 489.
T. diffusus. *Br.* 125.
T. Gerhardi. *Sl.* 2. p. 45.
Tamarind Tree. *Hughes.* 189. *Long.* 2.
 p. 729. *Aiton.*

* De Candolle has introduced two species of the *Tamarindus*, founded on the length of the pod, and the number of its seeds. In this I cannot follow him, as I have seen the seed-vessel to vary very much in length and number of seeds on the same tree. On the West-Indian *Tamarind Tree* I have commonly seen the pod, or lomentiform fruit, fully six times as long as broad, and the seeds more than four in number, although, perhaps, never twelve. It is sometimes single seeded, and nearly round.

MONADELPHIA PENTANDRIA.

WALTHERIA.

Calumniferæ. *Lin.*—Tiliaceæ. *Juss.*

Byttneriaceæ. *De Cand.*

americana. *W.* foliis ovalibus plicatis acute inæqualiter dentatis tomentosis, capitulis pedunculatis. *Willd. Sp. Pl.* 3. p. 586. *Hort.*

Kew. 4. p. 138. *Aubl.* 678. h

W. fruticosa subhirsuta. *Br.* 276.

W. arborescens. *Cav. Diss.* 316. t. 170.
f. 1.

W. indica. *Jacq. Ic.* 1. t. 130.

Malva americana abutili folio. *Sl.* 1.
p. 218. t. 138. f. 2, 3.*

* Sloane's plate, 138. f. 2. 3. certainly applies to his *Malva americana* above, rather than to his *Abutilon arboreum*, with which he connects it. The leaves of the latter he describes as "standing on an eighth of an inch long footstalks," and says "they are three quarters of an inch long, and one third part

Dialthæa. *Hughes*. 172.

American Waltheria. *Aiton*.

Buff-Coat.

RIEDLEIA.

Byttneriaceæ. *De Cand.*—Columniferæ. (*Melochia*). *Lin.*

Malvaceæ. (*Melochia*). *Juss.*

nodiflora. *R.* foliis ovatis acuminatis serratis glabris; junioribus appresse pilosis, floribus axillaribus conglomeratis sessilibus. *De*

Prod. 1. p. 491 h

Melochia nodiflora. *Willd. Sp. Pl.* 3. p. 603. *Swtz. Fl.* 1139.

Abutilon fruticosum. *Sl.* 1. p. 219. t. 135. f. 2.

Red Dialthæa. *Hughes*. 212,

Knot-flowered Riedleia. *Sweet*.

of an inch broad near the base, where round and broadest." In the plate the footstalks are represented an inch long, and the leaves more than two inches long, and more than an inch broad. Compare Sloane's plate with those of Jacquin and Cavanilles.

OCHROMA.

Columniferæ. *Lin* —Malvaceæ. *Juss.*—Bombaceæ. *De Cand.*

Lagopus. *O.* foliis cordatis 5—angulato-sublobatis denticulatis subtus pubescentibus. *De Cand. Prod.* 1. p. 480. *Willd. Sp. Pl.* 3. p. 605. *Hort. Kew.* 4. p. 147. *Swtz. Fl.* 1144. t. 23. h

Hibiscus arborescens. *Br.* 286.

Bombax pyramidale. *Cav. Diss.* 294. t. 153.

Gossypium vel Xylon arboreum. *Pluk. Alm.* 173. *Phyt. t.* 189. f. 2.

Down Tree. *Hughes.* 194.

Pyramidal Ochroma. *Aiton.*

Bombast-Mahoe. *Browne.*

PASSIFLORA.

Cucurbitaceæ. *Lin.*—Cucurbitaceis aff. *Juss.*

Passifloreæ. *De Cand.*

malifor- *P.* foliis oblongo-ovatis cordatis tripli-
mis. nerviis venosis integerrimis, petiolis

biglandulosis, involucro triphylo flore
 majore. *Willd. Sp. Pl. 3. p. 608. Hort.*
Kew. 4. p. 148. Aubl. 831. Bot. Reg.
t. 94. Bot. Repos. t. 217. h

P. foliis cordatis productis. Br. 328.

Clematitis indica latifolia. Plum. Amer.
67. t. 82.

Apple-fruited Passion-Flower. *Miller.*

Water-Lemon. *Browne.*

Conch-Apple. Conch-Nut.

quadran- *P. foliis oblongo-ovatis subcordatis inte-*
gularis. *gerrimis venosis, petiolis sexglandulosis,*
stipulis subrotundo-ovatis, involucro tri-
phylo, caule membranaceo-tetragono.

Willd. Sp. Pl. 3. p. 608. Hort. Kew. 4.
p. 148. Jacq. Amer. 231. t. 143. Cav.
Diss. 453. t. 283. Bot. Reg. t. 14. h

P. foliis amplioribus. Br. 327.

Granadilla Vine. *Hughes. 187. Long.*
2. p. 804.

Square-stalked Passion-Flower. *Miller.*

laurifolia. *P. foliis oblongis integerrimis venosis,*
petiolis biglandulosis, involucro triphylo
apice dentato. Willd. Sp. Pl. 3. p. 609.

Hort. Kew. 4. p. 148. Aubl. 831. Cav.
Diss. 454. t. 284. Bot. Reg. t. 13. h

P. foliis ovatis. Br. 327.

Clematitis indica, fructu citriformi. Plum.
Amer. 64. t. 84.

Water-Lemon Vine. *Hughes*. 184.

Laurel-leaved Passion-Flower. Water-Lemon. *Miller*.

Honey-Suckle. *Browne*.

suberosa. *P. foliis trilobis glabris: lobis oblongis: lateralibus brevissimis, petiolis biglandulosus, pedunculis subgeminis petiolo longioribus, floribus apetalis, caule, basi suberoso. Willd. Sp. Pl. 3. p. 617. Hort. Kew. 4. p. 151. Aubl. 832. Cav. Diss. 442. t. 265. Ex. Bot. t. 28.* h

Clematitis indica, flore minore pallido. Plum. Amer. 73. t. 88.

Ink Vine. *Hughes*. 160.

Cork-barked Passion-Flower. *Miller*.

foetida. *P. foliis trilobis cordatis pilosis, involucri multifido-capillaribus. Willd. Sp. Pl. 3. p. 619. Hort. Kew. 4. p. 153. Cav. Diss. 458. t. 289. Bot. Reg. t. 321.* u

P. vesicaria. Br. 327.

Clematitis indica hirsuta foetida. Plum. Amer. 71. t. 86.

Wild Water-Lemon Vine, or Love-in-a-Mist. *Hughes*. 184.

Stinking Passion-Flower. *Miller*.

MONADELPHIA HEPTANDRIA.

PELARGONIUM.

Gruinales (*Geranium*). *Lin.*—*Gerania*. *Juss.*

Geraniaceæ. *De Cand.*

humifusum. *P. multicaule procumbens, foliis cordatis sub—3—partitis 5—lobisve dentatis, pedunculis 3—5—floris, floribus 4—5—antheris. De Cand. Prod. 1. p. 654.*

Sweet. Ger. t. 42.

⊙

Trailing Stork's-Bill. *Sweet.*

zonale. *P. foliis cordato-orbiculatis obsolete lobatis dentatis superne zonatis, pedunculis multifloris, petalis cuneatis. De Cand.*

Prod. 1. p. 659. Willd. Sp. Pl. 3. p.

667. Hort. Kew. 4. p. 172.

h

α. foliis zonatis.

β. foliis albo-marginatis.

Ciconium zonale. Hort. Br. 84.

Horse - shoe (*Stork's - Bill*) *Ciconium.*

Sweet.

Monstrum. P. foliis orbiculato-reniformibus obsolete lobatis subzonatis complicatis crispis utrinque pubescentibus, pedunculis multifloris, floribus aggregatis: petalis lineari-cuneatis. *De Cand. Prod.* 1. p. 659. $\frac{1}{2}$
P. monstrosum. *Willd. Sp. Pl.* 3. p. 668.
Hort. Kew. 4. p. 172. *Sweet. Ger. t.* 13.
Ciconium Monstrum. *Hort. Brit.* 84.
Cluster-leaved (*Stork's-Bill*) Ciconium.
Sweet.

pyrethri- P. ramosissimum, foliis cordatis palma-
folium. to-bipinnatifidis scabris margine planis: segmentis oblongo-lanceolatis obtusis dentatis, pedunculis 2—3—floris, petalis inferis ligulatis, tubo nectarifero longitudine calycis. *De Cand. Prod.* 1. p. 663.
Sweet. Ger. t. 153. $\frac{1}{2}$
Feverfew-leaved Stork's-Bill. *Sweet.*

quinque- P. caule suffruticoso, foliis hispidis tri-
vulnerum. sectis: segmentis multifidis: laciniis lineari-lanceolatis serratis, umbellis multifloris. *De Cand. Prod.* 1. p. 664. *Sweet.*
Ger. t. 161. $\frac{1}{2}$
Dark-flowered Stork's-Bill. *Sweet.*

peltatum. P. caule fruticoso, ramis carnosis angulatis, foliis peltatis 5—lobis integerrimis carnosissimis, umbellis paucifloris. *De Cand. Prod.* 1. p. 666. *Willd. Sp. Pl.* 3. p. 669.
Hort. Kew. 4. p. 173. *Bot. Mag. t.* 20. $\frac{1}{2}$

Peltated-leaved Stork's-Bill. *Sweet.*

capitatum. P. foliis cordatis trilobis undatis molli-
ter villosis dentatis, stipulis latis cordatis,
caulibus diffusis, umbellis multifloris ca-
pitatis, tubo nectarifero calyce triplo
breviore. *De Cand. Prod.* 1. p. 674.

Willd. Sp. Sp. Pl. 3. p. 676. *Hort.*

Kew. 4. p. 176.

h

Rose-scented Stork's-Bill. *Sweet.*

dissectum. P. caule fruticoso ramoso diffuso, foliis
cordatis 5—7—partitis canescenti-pu-
bescentibus: segmentis flabelliformibus
imbricatis inciso-dentatis obtusis, um-
bellis multifloris paniculatis, petalis sub-
æqualibus, tubo nectarifero calyce sub-
æquali aut paulo brevior. *Sweet. Ger.*

t. 247.

h

Dissected-leaved Stork's-Bill. *Sweet.*

MONADELPHIA ENDECANDRIA.

BROWNEA.

Dubii ord. *Lin.*—Leguminosæ. *Juss. De Cand.*

coccinea. B. foliolis 2—3—jugis ovali-oblongis
acuminatis, floribus subfasciculatis, ramis
petiolisque glabris. *De Cand. Prod.* 2.
p. 477. *Willd. Sp. Pl.* 3. *p.* 715. *Hort.*
Kew. 4. *p.* 191. *Jacq. Amer.* 194.
t. 121. h

Scarlet-flowered Brownea. *Aiton.*

MONADELPHIA POLYANDRIA.

CAROLINEA.

Malvaceæ (*Pachira*). *Juss.*—Bombaceæ. *De Cand.*

insignis.* *C. foliolis* 5—7 obovato-oblongis, calyce sinuato, petalis erectis apice summo patulis. *De Cand. Prod.* 1. p. 478. *Willd. Sp. Pl.* 3. p. 730. *Hort. Kew.* 4. p. 195. *Swtz. Fl.* 1202. h
Bombax grandiflorum. *Cav. Diss.* 295. t. 154.
Great-flowered Carolina. *Aiton.*

* Commonly known in Barbados as Carolina princeps.

ADANSONIA.

Columniferae. *Lin.*—Malvaceae. *Juss.*—Bombaceae. *De Cand.*

- digitata.* *A. digitata.* *Willd. Sp. Pl.* 3. p. 730.
Hort. Kew. 4. p. 195. *De Cand. Prod.*
 1. p. 478. *Cav. Diss.* 298. t. 157. $\frac{1}{2}$
A. Baobab. *Gaert. Fr.* 2. p. 253. t. 135.
Baobab sive Abavi. *Bauh. Hist.* 1. p. 110.
Abavo arbor. *Bauh. Pin.* 434.
 Corn Tree. *Hughes.* 218? *
 Ethiopian Sour Gourd, or Monkies'
 Bread. *Miller.*

* There is certainly no tree in Barbados that bears for its fruit a "long cylindrical ear, not ill resembling a large mould-candle, round which stalk the grains (of corn) are disposed," as stated by Hughes. From his general description, however, of the corn tree, and from his account of the period of its introduction into the island, I have little doubt of his meaning what is now generally termed the Baobab, or Monkies' Bread-fruit Tree.

BOMBAX.

Columniferæ. *Lin.*—Malvaceæ. *Juss.*—Bombaceæ. *De Cand.*

- Ceiba.* B. caudice aculeato, foliis palmatis 5—
foliolatis, fructu turbinato apice concavo.
De Cand. Prod. 1. p. 478. *Willd. Sp.*
Pl. 3. p. 732. *Hort. Kew.* 4. p. 196.
Jacq. Amer. 192. t. 176. f. 71. *Aubl.*
702. *Cav. Diss.* 294. t. 152. h
B. foliis digitatis. *Br.* 277.
Gossypium arboreum maximum. *Sl.* 2.
p. 72.
Silk-Cotton Tree. *Hughes.* 190. *Long.*
2. p. 736.
Five-leaved Silk-Cotton Tree. *Aiton.*

SIDA.

Columniferæ. *Lin.*—Malvaceæ. *Juss.* *De Cand.*

- angusti-* S. foliis lineari-lanceolatis dentatis, pe-
*folia.** dunculis subsolitariis axillaribus, capsulis

* The authors to whom Willdenow refers in general represent this species as spinous. He himself is silent on this head, and

bicuspidatis. *Willd. Sp. Pl. 3. p. 734.*

Hort. Kew. 4. p. 197. De Cand. Prod.

1. p. 459. Cav. Diss. 14. t. 2. f. 2. h

S. ulmifolia. Retz. Obs. 3. p. 37.

Broom-Weed (*white*). *Hughes. 171.*

Narrow-leaved Sida. *Sweet.*

jamaicensis. *S. foliis ovatis serratis tomentosis, floribus axillaribus subpedunculatis, capsulis*

quinis bicornibus. Willd. Sp. Pl. 3. p.

737. De Cand. Prod. 1. p. 460. Cav

Diss. 17. t. 2. f. 3. h

S. humilior foliis ovatis serratis. Br. 280.

Broom-Weed (*crumple-leaved*). *Hughes. 171.*

Broad-leaved Broom-Weed. *Long. 2. p 831.*

Jamaica Sida. *Sweet.*

populifolia. *S. foliis subrotundo-cordatis acuminatis inæqualiter repando-dentatis, pedunculis solitariis petiolo longioribus, capsulis acutis truncatis calyce longioribus.*

Willd. Sp. Pl. 3. p. 751. De Cand

Prod. 1. p. 470. Hort. Kew. 4. p. 202.

Cav. Diss. 32. (t. 7. f. 9. mala) t. 128.

f. 2. bona.

⊙

simply says, "axillæ foliorum 1—2—floræ." According to Cavanilles, "sub petiolo trium spinularum rudimenta;" but Retzius' statement, "stipulæ setacæ, spinæ nullæ," entirely accords with the Barbados narrow-leaved Sida.

S. major assurgens. *Br.* 281.

Poplar-leaved Sida. *Miller.*

althæifolia. *S. foliis cordatis subangulatis obtusis serrato-crenatis utrinque tomentosis, rostris seminum calyce brevioribus.* *Willd. Sp.*

Pl. 3. p. 753. De Cand. Prod. 1. p. 464.

Swtz. Fl. 1207.

h

Althæa flore luteo. Sl. 1. p. 218. t. 136.

f. 2.

Marsh-Mallow-leaved Sida. *Miller.*

urens.

S. foliis ovato-cordatis acuminatis dentatis, pedunculis axillaribus multifloris glomeratis, capsulis muticis. *Willd. Sp.*

Pl. 3. p. 759. De Cand. Prod. 1. p. 465.

Hort. Kew. 4. p. 204.

h

S. urticata foliis cordatis serratis. Br. 280.

Althæa spicata betonicæ folio. Sl. 1. p.

44. t. 14. f. 3.

Stinging Sida. *Miller.*

Nettle Sida. *Browne.*

MALACHRA.

Columniferæ (*Sida*). *Lin* — Malvaceæ. *Juss. De Cand.*

capitata. *M. capitulis pedunculatis triphyllis** septemfloris, foliis subcordatis obsolete lobatis, caule scabro. *Willd. Sp. Pl.* 3. p. 768. *De Cand. Prod.* 1. p. 440. *Hort. Kew.* 4. p. 207. ☉

Sida capitata. *Aubl.* 704.

Sida hirta assurgens. *Br.* 281.

Malva aspera major aquatica. *Sl.* 1. p. 217. *t.* 137. *f.* 1.

Wild Hemp (*white*). *Hughes.* 167.

Headed Malachra. *Sweet.*

Bastard Ockro. *Browne.*

Wild Ockro.

fasciata. *M. capitulis pedunculatis triphyllis* subquinquefloris, foliis subrotundis obsolete lobatis, caule villosa. *Willd. Sp. Pl.* 3. p. 769. *De Cand. Prod.* 1. p. 440. *Jacq. Ic.* 3. *t.* 548. ☉

Wild Hemp (*red*). *Hughes.* 167.

Fasciate Malachra. *Sweet.*

* In every head that I have examined in this and the following species there is a floral leaf with two setaceous apendiculæ to each flower. The "capitulis triphyllis" is taken from the three larger and more mature external leaves.

ALTHÆA.

Columniferæ. *Lin.*—Malvaceæ. *Juss. De Cand.*

- rosea.* *A. foliis quinque—septemlobo-angulatis.*
 Willd. Sp. Pl. 3. p. 773. Hort. Kew. 4.
 p. 209. ♂
Malva rosea. Bauh. Pin. 315.
Common Holly-Hock. Aiton.

GOSSYPIUM.*

Columniferæ. *Lin.*—Malvaceæ. *Juss. De Cand.*

- herba-* *G. foliis quinquelobis subtus uniglandu-*
ceum. *losis : lobis rotundatis mucronatis, invo-*
 lucello serrato, caule lævi. De Cand.
 Prod. 1. p. 456. Willd. Sp. Pl. 3. p.
 803. Hort. Kew. 4. p. 222. Cav. Diss.
 310. t. 164. f. 2. ♀
 Small Cotton Tree (Flying-Fish sort).
 Hughes. 191.
 Common Cotton. Aiton.

* "Species omnes incertæ ex characteribus mancis stabilitatæ."—*De Cand.*

Pollard, or White Wool Cotton.

arboreum? G. foliis quinquelobo-palmatis: lobis lanceolatis obtusis setula brevi mucronatis subtus uniglandulosus, involucellis subintegris. *De Cand. Prod.* 1. p. 456.

Willd. Sp. Pl. 3. p. 804. *Hort. Kew.* 4. p. 222. *Aubl.* 705. *Cav. Diss.* 311. t. 165. h

Tree Cotton. *Aiton.*

Persian Cotton.

vitifolium. G. foliis inferioribus quinquelobo-palmatis; superioribus trilobis subtus uniglandulosus, involucellis laciniatis, calycibus basi triglandulosus, caule punctato glabro.

De Cand. Prod. 1. p. 456. *Willd. Sp. Pl.* 3. p. 804. *Hort. Kew.* 4. p. 223. *Cav. Diss.* 311. t. 166. h

Small Cotton Tree (*Vine sort*). *Hughes.* 191.

Vine-leaved Cotton Tree. *Aiton.*

Vine Cotton.

hirsutum. G. foliis superioribus indivisis cordatis; inferioribus tri-quinquelobis subtus uniglandulosus, ramulis petiolisque hirsutis, involucellis apice tridentatis. (*seminibus viridibus adhærentibus. W.*) *De Cand.*

Prod. 1. p. 456. *Willd. Sp. Pl.* 3. p. 805. *Hort. Kew.* 4. p. 223. *Aubl.* 705. *Cav. Diss.* 312. t. 167. 4

Hairy Cotton. *Aiton.*

Sea-Island Cotton.

barba-
dense.

G. foliis superioribus trilobis; inferioribus quinquelobis subtus triglandulosis, caule læviori, seminibus liberis. De Cand.

Prod. 1. p. 456. Willd. Sp. Pl. 3. p. 806.

Hort. Kew. 4. p. 224. Aubl. 705. 2

Small Cotton Tree (*great sort*). *Hughes.*
191.

Barbadoes Cotton. *Aiton.*

Common brown wool Cotton.

peruvia-
num.

G. foliis quinquelobis triglandulosis; inferioribus indivisis, involucellis laciniatis basi triglandulosis. De Cand. Prod. 1.

p. 456. Cav. Diss. 313. t. 168. 2

Peruvian Cotton.

THESPESIA.

Columniferæ (*Hibiscus*). *Lin.*—*Malvaceæ. Juss. De Cand.*

populnea. *T. foliis subrotundo-cordatis acuminatis. De Cand. Prod. 1. p. 456. 2*

Hibiscus populneus. Willd. Sp. Pl. 3. p. 809. Hort. Kew. 4. p. 224. Aubl. 705. Cav. Diss. 152. t. 56. f. 1.

Alcæa malabariensis. Raj. Hist. 1069.

Novella litorea. *Rumph. Amb. 2. p. 224.*
t. 74.

Poplar-leaved Thespesia. *Sweet.*

Poplar Tree. Poppy Tree.

HIBISCUS.

Columniferæ. *Lin.*—Malvaceæ. *Juss. De Cand.*

arboreus. H. caule arboreo, ramis pubescentibus, foliis subrotundo-cordatis acuminatis apice obscure dentatis supra punctatis subtus cinereo tomentosus, calyce exteriori decemdentato: dentibus subbrevis, floribus amplis luteis. *Ham. Prod.*
 59. h

H. abutiloides. *De Cand. Prod. 1. p. 454.*

Althæa maritima arborescens. *Br. 284.*

Malva arborea maritima *Sl. 1. p. 215.*
t. 134. f. 4.

Rope Mangrove. *Hughes. 199.*

Rosa- H. foliis cordatis acuminatis dentatis
sinensis. glabris basi integerrimis, caule arboreo, calyce exteriori suboctophyllo. *Willd.*
Sp. Pl. 3. p. 812. Hort. Kew 4. p.
225. De Cand. Prod. 1. p. 448. h

α . flore simplici. *Cav. Diss.* 158. *t.* 69.

f. 2. *Bot. Mag.* *t.* 158.

Chinese Rose-Hibiscus. *Aiton.*

Common Scarlet Hibiscus.

β . flore pleno.

Flos festalis. *Rumph. Amb.* 4. *p.* 24.
t. 8.

Double Chinese Rose-Hibiscus. *Aiton.*

Rose of Sharon.

phœni-
*ceus.**

H. foliis ovatis acuminatis serratis; inferioribus subcordatis tricuspidatis, pedunculis articulatis, seminibus lanatis.

Willd. Sp. Pl. 3. *p.* 813. *Hort. Kew.*
4. *p.* 225. *Bot. Reg.* *t.* 230. $\frac{1}{2}$

H. phœniceus. β . involucro calycem æquante. *De Cand. Prod.* 1. *p.* 452.

H. hirtus. *Cav. Diss.* 156. *t.* 67. *f.* 3.

Small purple-flowered Hibiscus. *Aiton.*
Dwarf Hibiscus.

mutabilis. *H. foliis cordatis angulatis quinquelobis acuminatis dentatis, calyce exteriori oc-*

* “*Hibiscus hirtus* hujus solummodo varietas, quæ forte a sola cultura dependet.”—*Lin. Suppl.* 310. The plate of *Hibiscus phœniceus* of Cavanilles (*t.* 67. *f.* 2.) is by no means so characteristic of the Barbados plant as that of *H. hirtus*, referred to above. The small Hibiscus so common in the gardens is the β . variety of De Candolle, the characters of which should rather, I think, be considered as marking a distinct species.

tophylo, capsula villosa, caule arboreo.

Willd. Sp. Pl. 3. p. 817. *Hort. Kew.* 4.
p. 226. *De Cand. Prod.* 1. p. 452.
Aubl. 706. *Cav. Diss.* 165. t. 62. f. 1.
Bot. Repos. t. 228. h

H. fruticosus brachiatus. *Br.* 286.

Flos horarius. *Rumph. Amb.* 4. p. 27. t. 9.

French Rose Tree. *Hughes.* 218.

Chinoise Rose. *Browne.*

Changeable Rose-Hibiscus. *Miller.*

pruriens. *H. caule herbaceo piloso, foliis longe
petiolatis ovatis subtrilobis serratis mem-
branaceis glabriusculis, pedicellis brevis-
simis, involucello et calyce basi pilosis-
simis corollam æquantibus: utroque
5—partito acuminato.* *De Cand. Prod.*
1. p. 448. *Bot. Repos.* t. 498. ⊙

Nettle Hibiscus. *Andrews.*

Sabdariffa. *H. foliis dentatis; inferioribus ovatis in-
divisis; superioribus trilobis basi cune-
atis, calyce exteriori subduodecimden-
tato, caule inermi, floribus subsessilibus.*

Willd. Sp. Pl. 3. p. 821. *Hort. Kew.* 4.
p. 227. *De Cand. Prod.* 1. p. 453. *Aubl.*
706. *Cav. Diss.* 170 & 351. t. 198. f. 1. ⊙

α. caule rufescente.

H. rufescens acetosus. *Br.* 285.

Sorrel (red). *Hughes.* 204. *Long.* 2.
p. 805.

β. caule virescente.*

Sorrel (*white*). *Hughes*. 204. *Long*.
2. p. 805.

Various-leaved Hibiscus. *Miller*

*cannabi-
nus.*

H. foliis superioribus palmatis quinque-
partitis subtus uniglandulosis, caule acu-
leato, floribus sessilibus. *Willd. Sp. Pl.*
3. p. 822. *Hort. Kew.* 4. p. 228. *De*
Cand. Prod. 1. p. 450. *Aubl.* 707. *Roxb.*
Cor. 2. p. 43. t. 190. *Cav. Diss.* 148. t.
52. f. 1. ☉

Hemp-leaved Hibiscus. *Miller.*

vitifolius.

H. foliis glabriusculis dentatis quinquan-
gularibus acuminatis, floribus cernuis.
Willd. Sp. Pl. 3. p. 829. *Hort. Kew.* 4.
p. 230. *De Cand. Prod.* 1. p. 450. *Cav.*
Diss. 145. t. 518. f. 2. ☉

Vine-leaved Hibiscus. *Miller.*

*Abelmos-
chus.*

H. foliis subpellato-cordatis septemangu-
laribus acuminatis serratis, calyce ex-
teriore suboctophyllo, caule hispido.
Willd. Sp. Pl. 3. p. 826. *Hort. Kew.* 4.
p. 229. *De Cand. Prod.* 1. p. 452.
Aubl. 707. *Cav. Diss.* 167. t. 62. f. 2. ♂
H. hispidus. *Br.* 285.

Alcæa ægyptiaca villosa. *Bauh. Pin.* 317.

Musk-Bush, or Wild Ockro. *Hughes.* 207.

* I am not confident that the white sorrel is not a distinct species.

Musk Ockro. *Long.* 2. p. 776.

Target-leaved Hibiscus. *Miller.*

esculentus. *H.* foliis cordatis quinquelobis obtusiusculis dentatis, petiolis flore longioribus, calycibus subdecaphyllis deciduis: interioribus longitudinaliter rumpentibus.

Willd. Sp. Pl. 3. p. 827. *Hort. Kew.* 4.

p. 229. *De Cand. Prod.* 1. p. 450.

Aubl. 707. *Cav. Diss.* 168 & 351. t.

61. f. 2. ☉

H. ramosus hirsutus. *Br.* 285.

Alcæa maxima malvæ rosæ folio. *Sl.* 1.

p. 223. t. 133. f. 3.

Ockra—Ockro. *Hughes.* 210. *Long.* 2.

p. 775.

Eatable Hibiscus. *Miller.*

MYRODIA.

Bombaceæ. *De Cand.*

turbinata. *M.* foliis ovato-oblongis, calycibus turbinatis, columna staminum petalis brevior. *Willd. Sp. Pl.* 3. p. 843. *Hort.*

Kew. 4. p. 236. *Swtz. Fl.* 1227. h

Five-sprig-Tree. *Hughes.* 217.

Short-flowered Myrodia. *Aiton.*

DIADELPHIA OCTANDRIA.

BADIERA.

Lomentaceæ (*Polygala*). *Lin.*—Pediculares (*Polygala*). *Juss.*

Polygaleæ. *De Cand.*

*diversi-
folia.*

B. racemis brevibus axillaribus, foliis ovali-oblongis utrinque acuminatis. De Cand. Prod. 1. p. 334. h

Polygala diversifolia. Willd. Sp. Pl. 3. p. 887. Aubl. 739.

Polygala fruticosa. Br. 287. t. 5. f. 3, 4.

Genista affinis anonyma arbor. Sl. 2. p. 32. t. 170. f. 2.

Bastard Lignum Vitæ. Hughes. 142. Long. 2. p. 825.

SECURIDACA.

Leguminosæ. *Juss.*—Polygaleæ. *De Cand.*

- volubilis.* S. caule scandente, ramulis subpubescentibus, foliis ovali-oblongis acutis, racemis lateralibus, capsulæ ala obovata dorso ad basin apendiculata. *De Cand. Prod.* 1. p. 340. *Willd. Sp. Pl.* 3. p. 899. *Hort. Kew.* 4. p. 247. h
 S. scandens. *Br.* 288. *Jacq. Amer.* 197. *t.* 183. *f.* 83. *Aubl.* 739.
 S. caule volubili. *Plum. Ic.* 244. *t.* 247. *f.* 1.

Climbing Securidaca. *Miller.*

DIADELPHIA DECANDRIA.

ECASTAPHYLLUM.

Papilionaceæ (*Pterocarpus*). *Lin.*

Leguminosæ. *De Cand.* (*Pterocarpus*.) *Juss.*

Brownei. *E.* foliis 1—foliolatis, folio nempe unico terminali ovato basi subrotundo subcordato apice acuminato subtus pubescente.

De Cand. Prod. 2. p. 420. h

E. frutescens. *Br.* 299. t. 32. f. 1.

Pterocarpus Ecastaphyllum. *Willd. Sp.*

Pl. 3. p. 907. *Hort. Kew.* 4. p. 249.

Hedysarum Ecastaphyllum. *Aubl.* 773.

Oval-leaved *Ecastaphyllum.* *Sweet.*

ABRUS.

Papilionaceæ (*Orobus*). *Lin.*—Leguminosæ. *Juss. De Cand.*

- precato- A. precatorius. Willd. Sp. Pl. 3. p. 911.*
rius. Hort. Kew. 4. p. 250. De Cand. Prod.
2. p. 381. h
Glycine scandens. Br. 297.
Phaseolus Glycyrrhizites. Sl. 1. p. 180.
t. 112. f. 4—6.
Crab's-Eye Vine. Hughes. 198.
Wild Liquorice, or Red-bead Vine.
Long. 2. p. 799.
Jamaica Wild Liquorice. Miller.

ERYTHRINA.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

- Corallo- E. caule arboreo aculeato, petiolis iner-*
dendron. mibus, foliolis late ovato-rhombeis acutis
glabris, calyce truncato 5—dentato, vex-
illo oblongo, stamine decimo libero aliis
subæquali. De Cand. Prod. 2. p. 411.
Willd. Sp. Pl. 3. p. 913. Hort. Kew.
4. p. 251. h

E. arborea. *Br.* 288.

Coral Arbor. *Sl.* 2. *p.* 38.

Bean Tree, or Throve Tuesday. *Hughes.*
194.

Coral, or Red Bean Tree. *Long.* 2. *p.* 788.

Smooth-leaved Coral Tree. *Miller.*

TERAMNUS.

Papilionaceæ (*Dolichos*). *Lin.*—Leguminosæ. *Juss. De Cand.*

uncinatus. *T. ramis retrorsum sericeis, foliis subtus sericeis superne pubescentibus: foliolis ovatis aut oblongis.* *De Cand. Prod.* 2. *p.* 382. h

T. volubilis. *Willd. Sp. Pl.* 3. *p.* 971,
Swtz. Fl. 1241.

T. triphyllus subhirsutus. *Br.* 290.

Dolichos uncinatus. *Lin. Sp. Pl.* 1019.

Phaseolus sylvestris minor. *Sl.* 1. *p.* 182.

Horse Vine. *Hughes.* 208.

Hooked Teramnus. *Sweet.*

Rabbit Vine.

CROTALARIA.

Papilionaceæ, *Lin.*—Leguminosæ. *Juss. De Cand.*

- retusa.* C. foliis simplicibus oblongis cuneiformibus retusis, racemo terminali. *Willd. Sp. Pl. 3. p. 976. Hort. Kew. 4. p. 272. De Cand. Prod. 2. p. 125. Bot. Mag. t. 2561. Bot. Reg. t. 253.* ☉

Wedge-leaved Crotalaria. *Miller.*

Yellow Lupin.

- verucosa.* C. foliis simplicibus ovatis retusis, stipulis lunatis declinatis, racemo terminali, ramis tetragonis. *Willd. Sp. Pl. 3. p. 977. Hort. Kew. 4. p. 272. Aubl. 763. Bot. Repos. t. 308.* ☉

C. *verucosa.* α . *obtusa.* *De Cand. Prod. 2. p. 125.*

Blue-flowered Crotalaria. *Miller.*

Blue Shake-Shake.

- lotifolia.* C. foliis ternatis: foliolis cuneiformibus emarginatis subtus villosis-subsericeis, pedunculis axillaribus solitariis unifloris. *Willd. Sp. Pl. 3. p. 983. Hort. Kew. 4. p. 273. De Cand. Prod. 2. p. 134. Aubl. 763.* ☉

C. *trifolia fruticosa.* *Sl. 2. p. 33. t. 176. f. 1, 2.*

Coot-Weed. *Hughes*. 212.

Lotus-leaved Crotalaria. *Miller*.

incana.

C. foliis ternatis ovalibus subtus villosis, racemis spiciformibus, carina margine tomentosa, leguminibus sessilibus hirsutis. *Willd. Sp. Pl.* 3. p. 985. *Hort. Kew.*

4. p. 274. *De Cand. Prod.* 2. p. 132. ☉

C. trifolia fruticosa. *Sl.* 2. p. 34. t. 179. f. 1.

Rattle-Bush, or Shake-Shake. *Hughes*. 212.

Hoary Crotalaria. *Miller*.

ARACHIS.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

hypogæa. *A. hypogæa.* *Willd. Sp. Pl.* 3. p. 1021.

Hort. Kew. 4. p. 285. *De Cand. Prod.*

2. p. 474. *Aubl.* 765. ☉

A. tetraphylla. *Br.* 295.

Arachnida Indiæ utriusque. *Sl.* 1. p. 184.

Ground-Nut. *Hughes*. 232.

Pindals, or Ground-Nuts. *Long.* 2. p. 788.

Common Earth or Ground Nuts. *Miller*.

PHASEOLUS.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

perennis. *P. volubilis pubescens, foliolis ovatis acuminatis triplinerviis, racemis 1—3 in axilla folio longioribus paniculatis, bracteolis minimis, leguminibus pendulis latis falcatis mucronatis. De Cand.*

Prod. 2. p. 391. Willd. Sp. Pl. 3. p. 1031. Br. 292. 4

Lima Bean. *Long. 2. p. 786.*

Perennial Kidney-Bean. *Sweet.*

multiflorus. *P. volubilis glabriusculus, foliolis ovatis acuminatis, racemis pedunculatis folio longioribus, pedicellis geminatis, bracteolis calyce subbrevioribus appressis, leguminibus pendulis subfalcatis torulosis scabris. De Cand. Prod. 2. p. 392.*

Willd. Sp. Pl. 3. p. 1030. Hort. Kew. 4. p. 288. ☉

Scarlet-Runner Kidney-Bean. *Sweet.*

French Bean.

vulgaris. *P. volubilis glabriusculus, foliolis ovatis acuminatis, racemis pedunculatis folio brevioribus, pedicellis geminis, leguminibus pendulis rectiusculis subtorulosis*

longe mucronatis, semine ovato-subcompresso. *De Cand. Prod.* 2. p. 392.

Willd. Sp. Pl. 3. p. 1030. *Hort. Kew.* 4. p. 288. ☉

Common Kidney-Bean. *Sweet.*

French Bean.

lunatus. *P. volubilis* glabriusculus, foliolis ovatis acuminatis, racemis subpedunculatis folio brevioribus, pedicellis geminis, bracteolis minimis appressis, vexillo concavo subrotundo, legumine acinaciformi sublunato lævi 2—4—spermo. *De Cand.*

Prod. 2. p. 393. *Willd. Sp. Pl.* 3. p. 1031. *Hort. Kew.* 4. p. 288. ☉

P. scandens. *Br.* 292.

Cimeter-podded Kydney-Bean. *Sweet.*

Sugar Bean.

vexillatus. *P. caule* prostrato volubili pilosiusculo, foliolis oblongo-ovatis, pedunculis longissimis, floribus 5—7. capitatis: vexillo majusculo emarginato: alis parvulis, leguminibus teretibus pilosiusculis, seminibus lanatis. *De Cand. Prod.* 2. p. 395.

Willd. Sp. Pl. 3. p. 1032. *Hort. Kew.* 4. p. 289. *Jacq. Vind.* 3. p. 45. t. 102. ☉

Sweet-scented Kidney-Bean. *Sweet.*

Mungo. *P. caule* flexuoso tereti hispido, foliolis ovatis acutis subrepandis, pedunculis

folio paulo longioribus, floribus 6—7.
capitatis: carina sinistrorsum cornuta,
leguminibus teretibus hirtis subtorulosis,
seminibus cylindraceis truncatis. *De*
Cand. Prod. 2. p. 395. *Willd. Sp. Pl.*
3. p. 1036. ☉

P. hirsutus flexicaulis. Pluk. Alm. 290.
Small-fruited Kidney-Bean. *Sweet.*
Woolly-Pyroe.

semierectus. *P. caule demum volubili tereti pubes-*
cente, foliolis ovato-lanceolatis acutis
glabris, pedunculis elongatis, floribus
spicatis geminatis, carina mutica unci-
nata dextrorsum versa, leguminibus sub-
patentibus rectis subulato-compressis
rostro acuminatis, seminibus oblongis.

De Cand. Prod. 2. p. 396. *Willd. Sp.*

Pl. 3. p. 1033. *Hort. Kew.* 4. p. 289.

Jacq. Ic. 3. t. 558. *Bot. Reg.* t. 743. ☉

P. barbadensis erectior. Dill. Elth. 312.
t. 233. f. 301.

Wild Dolly. *Hughes.* 214.

Dark Red Kidney-Bean. *Sweet.*

CLITORIA.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

ternatea. *C.* caule volubili subpubescente, foliolis 2—3—jugis ovalibus ovatisve, stipulis subulatis, pedicellis solitariis 1—floris, bracteolis magnis subrotundis, leguminibus glabriusculis. *De Cand. Prod.* 2.

p. 233. *Willd. Sp. Pl.* 3. *p.* 1068. *Hort.*

Kew. 4. *p.* 300. *Bot. Mag. t.* 1542. 2

Winged-leaved Clitoria. *Miller.*

Blue Vine.

*virginiana.** *C.* caule scandente foliisque glabris aut subpuberulis, pedunculis 1—4—floris, bracteolis lanceolatis circiter calycis longitudine, leguminibus linearibus compressis. *De Cand. Prod.* 2. *p.* 234. *Willd.*

Sp. Pl. 3. *p.* 1069. *Hort. Kew.* 4. *p.*

301. *Aubl.* 765. *Bot. Reg. t.* 1047. 2

C. major scandens. *Br.* 298.

Wild Pea. *Hughes.* 215.

Small-flowered Clitoria. *Miller.*

* “Flores purpurascentes aut cærulei.”—*De Cand.* “Vidi plures varietates foliis angustioribus latioribusque corollis cæruleis et albis.”—*Willd.* In the plant mentioned by Hughes “the flowers are white, except that near their centre; they are slightly stained with red.”

PISUM.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

sativum. P. petiolis teretibus trijugis, foliolis ovatis integris margine undulatis sæpe oppositis mucronulatis, stipulis ovato-subsemicordatis crenatis, pedunculis bivul multifloris, leguminibus subcarnosis.

De Cand. Prod. 2. p. 368. *Willd. Sp.*

Pl. 3. p. 1070. *Hort. Kew.* 4. 302. ☉

Common Pea. *Miller.*

MUCUNA.

Papilionaceæ (*Dolichos*). *Lin.*—Leguminosæ. *Juss. De Cand.*

urens. M. floribus racemosis, leguminibus urentibus sulcis transversim lamellosis, foliolis subtus tomentoso-nitidis. *De Cand.*

Prod. 2. p. 405. h

Dolichos urens. *Willd. Sp. Pl.* 3. p. 1042.

Hort. Kew. 4. p. 293. *Jacq. Amer.* 202.

t. 182. *f.* 84. *Aubl.* 764.

Zoophthalmum siliquis majoribus. *Br.*
295.

Phaseolus brasiliensis frutescens. *Sl.* 1.
p. 178.

Horse-Eye Bean Vine. *Hughes.* 215.

Horse-Eye Bean. *Aiton.*

Ox-Eye Bean. *Browne.*

pruriens. M. floribus racemosis, leguminibus uren-
tibus valvulis subcarinatis, foliolis subtus
hirsutis acuminatis: medio rhomboideo:
lateralibus extus dilalatis. *De Cand.*
Prod. 2. *p.* 405. h

Dolichos pruriens. *Willd. Sp. Pl.* 3. *p.*
1041. *Hort. Kew.* 4. *p.* 293. *Jacq.*
Amer. 201. *t.* 122. *Aubl.* 764.

Stizolobium spicis multifloris. *Br.* 290.
t. 31. *f.* 4.

Phaseolus utriusque Indiæ. *Sl.* 1. *p.* 37.

Cow-Itch Vine. *Hughes.* 206.

Cow-Itch, or Cowhage. *Long.* 2. *p.* 825.

Common Cow-Itch. *Sweet.*

CAJANUS.

Papilionaceæ (*Cytisus*). *Lin.*—Leguminosæ. *De Cand.*
(Cytisus). *Juss.*

flavus. C. vexillo extus concolore, leguminibus
 2—3—spermis calycibusque immaculatis,
 stipellis foliolorum lateralium petiolulo
 dimidio brevioribus. *De Cand. Prod.* 2.
p. 406. h

Cytisus Cajan. *Willd. Sp. Pl.* 3. *p.* 1121.
Hort. Kew. 4. *p.* 320. *Aubl.* 768.

Cytisus fruticosus erectus. *Br.* 296.

Phaseolus erectus incanus. *Pluk. Alm.*
 293. *Phyt.* 213. *f.* 3.

Laburnum humilius. *Sl.* 2. *p.* 31.

Pigeon Pea Tree. *Hughes.* 199. *t.* 19.

Pigeon, or Angola Peas. *Long.* 2. *p.* 787.

Yellow-flowered Pigeon Pea. *Sweet.*

LONCHOCARPUS.

Papilionaceæ (*Robinia*). *Lin.*—Leguminosæ. *De Cand.*
(*Robinia*). *Juss.*

- violaceus*. *L.* foliolis 7—11 ovatis obtusis subemarginatis glabris pellucido-punctatis membranaceis nervo medio subtus subprominente, calycibus glabris, racemi pedunculis bifloris. *De Cand. Prod.* 2. p. 259. $\frac{1}{2}$
Robinia violacea. *Willd. Sp. Pl.* 3. p. 1131. *Hort. Kew.* 4. p. 323. *Jacq. Amer.* 210. t. 177. f. 49.
Spanish Ash. *Hughes.* 189.
White Spanish Ash. *Hughes.* 190 ?
Violet Lonchocarpus. *Sweet.*

ÆSCHYNOMENE.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

- sensitiva*. *Æ.* caule fruticoso lævi tereti, foliolis linearibus 16—20—jugis, leguminibus racemisque glabris, pedunculis ramosis

paucifloris, legumis articulis 8—10 fere quadratis medio lævibus ad saturam superiorem subpilosiusculis. *De Cand.*

Prod. 2. p. 320. *Willd. Sp. Pl.* 3. p. 1162. *Hort. Kew.* 4. p. 336. *Swtz. Fl.* 1276. h

Æ. mitis. *Plum. Ic.* 140. t. 149. f. 2.

West India Shrubby *Æschynomene*.
Aiton.

americana. *Æ.* caule erecto tereti hispido, foliolis linearibus mucronatis 10—20—jugis subciliatis, racemis simplicibus paucifloris, leguminis glabri impunctati articulis 4—8 hinc rectis illinc subrotundis. *De*

Cand. Prod. 2. p. 320. *Willd Sp. Pl.* 3. p. 1163. *Hort. Kew.* 4. p. 337. *Aubl.* 774. ⊙

Hedysarum caule hirsuto. *Sl.* 1. p. 186. t. 118. f. 3.

Hairy *Æschynomene.* *Aiton.*

STYLOSANTHES.

Papilionaceæ (*Hedysarum*). *Lin.*

Leguminosæ. *Juss. De Cand.*

*procum-
bens.*

S. caule suffrutescente procumbente apice appresse pubescente, foliolis oblongis acutis glabris lineatis, spicis multifloris.

De Cand. Prod. 2. p. 317. Willd. Sp.

Pl. 3. p. 1165. Swtz. Fl. 1282. t. 25. ½

Trifolium procumbens. Br. 298.

Hedysarum hamatum. α. Lin. Sp. Pl. 1056.

Anonis non spinosa. Sl. 1. p. 187. t. 119. f. 2.

Procumbent Stylosanthes. Sweet.

Small Creeping Trefoil. Browne.

LOUREA.

Papilionaceæ (*Hedysarum*). *Lin.*

Leguminosæ. *De Cand. (Hedysarum). Juss.*

*Vesperti-
lionis.*

L. foliolis lateralibus nullis aut minimis : terminali transversim et falcatis oblongo-

subemarginato longitudine decies latiore.

De Cand. Prod. 2. p. 323. ♂

Hedysarum Vespertilionis. Willd. Sp.

Pl. 3. p. 1177. *Hort. Kew.* 4. p. 340.

Jacq. Ic. 3. t. 566.

Bat-winged Lourea. *Sweet.*

DESMODIUM

Papilionaceæ (*Hedysarum*). *Lin.*—Leguminosæ. *Juss. De Cand.*

supinum. *D.* caule frutescente declinato superne pubescente, foliolis ovatis subincano-villosis, stipulis lanceolatis acuminatis puberulis, racemis terminalibus, pedicellis solitariis, leguminis erectiusculi articulis semiovatis pubescenti-hirtis. *De Cand.*

Prod. 2. p. 332. ♀

Hedysarum supinum. Willd. Sp. Pl. 3.

1190. *Swtz. Fl.* 1264.

Hedysarum triphyllum fruticosum supinum. Sl. 1. p. 185. t. 118. f. 2.

Sweet Heart. *Hughes.* 213.

Trailing Desmodium. *Sweet.*

spirale. *D.* caule herbaceo scandente basi radicante tereti glabro, foliolis ovatis glabris, stipulis subulatis, racemis terminalibus

elongatis, pedicellis binis ternisve, leguminibus pubescentibus spirali-tortuosis.

De Cand. Prod. 2. p. 332. ½

Hedysarum spirale. *Willd. Sp. Pl. 3. p. 1199. Swt. Fl. 1273.*

Iron Vine. *Hughes. 214.*

Spiral Desmodium. *Sweet.*

repens. D. caulibus procumbentibus, foliolis obcordatis, racemis lateralibus. *De Cand.*

Prod. 2. p. 338. ¼

Hedysarum repens. *Willd. Sp. Pl. 3. p. 1201.*

Hedysarum procumbens. *Dill. Elth. 172. t. 142. f. 169.*

Creeping Desmodium (*Hedysarum*). *Miller.*

Poverty-Weed.

HEDYSARUM.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

volubile. H. foliis ternatis, caule volubili, racemis axillaribus. *Willd. Sp. Pl. 3. p. 1204.*

Hort. Kew. 4. p. 345. ¼

H. trifoliatum scandens. *Dill. Elth. 173. t. 143. f. 170.*

Silver Vine. *Hughes*. 243.

Twining Hedysarum. *Miller*.

INDIGOFERA.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

Anil. I. caule suffruticoso erecto, foliis pinnatis
3—7—jugis: foliolis ovalibus subtus vix
pubescentibus, racemis axillaribus folio
brevioribus, leguminibus compressis non
torulosis deflexis arcuatis sutura utraque
calloso-prominula. *De Cand. Prod.* 2. p.
225. *Willd. Sp. Pl.* 3. p. 1236. *Hort.*
Kew. 4. p. 354. 24

I. assurgens. *Br.* 302.

Coluteæ affinis, siliquis falcatis. *Sl.* 2.
p. 37. t. 176. f. 3.

Indigo Weed. *Hughes*. 203.

Wild Indigo. *Miller. Browne.*

LABLAB.

Papilionaceæ (*Dolichos*). *Lin.*—Leguminosæ. *De Cand.*
(*Dolichos*). *Juss.*

vulgaris. L. leguminibus oblongo-ventricosus acinaciformibus, pericarpio facile detractili, seminibus ovatis subcompressis glandula basilari hemisphærica sulcata. *De Cand. Prod.* 2. p. 401. ☉

α. floribus violaceis, seminibus nigris.

Dolichos Lablab. *Willd. Sp. Pl.* 3. p. 1037. *Hort. Kew.* 4. p. 291. *Bot. Mag.* t. 896.

Phaseolus ægyptiacus. *Bauh. Pin.* 341.

Common Lablab. *Sweet.*

β. floribus purpureis, seminibus nigro-purpureis.

Dolichos purpureus. *Willd. Sp. Pl.* 3. p. 1048. *Hort. Kew.* 4. p. 295. *Bot. Reg.* t. 830. *Ex. Bot.* t. 74.

Phaseolus maximus perennis. *Sl.* 1. p. 177. t. 113. f. 4.

Buona Vista. Bonny-vis. (*Red and black.*) *Hughes.* 216.

Purple Lablab. *Sweet.*

γ. floribus albis, seminibus pallide ferrugineis.

Dolichos bengalensis. *Willd. Sp. Pl.*

3. p. 1038. *Jacq. Vind.* 2. p. 57. t. 124.

Phaseolus maximus perennis. *Sl.* 1.

p. 177. t. 113. f. 1—3.

Phaseolus jamaicensis et barbadensis.

Pluk. Alm. 291.

Buona Vista. Bonny-vis. (*White.*)

Hughes. 216.

Bengal Lablab. *Sweet.*

perennans. L. foliis pubescentibus, seminibus subrotundis. *De Cand. Prod.* 2. p. 402. 4

Dolichos albus. *Lour. Cochin.* 2. p. 534.

Rumph. Amb. 5. t. 137.

Phaseolus barbadensis septenii durationis.

Pluk. Alm. 291.

Buona Vista. Bonny-vis. (*Old Sort.*)

Hughes. 216.

Perennial Lablab. *Sweet.*

nankinicus. L. leguminibus oblongo-ventricosus acinaciformibus, pericarpio facile detractili, seminibus ovato-turgidiusculis glandula basilari hemisphærica sulcata deorsum acuminata. *De Cand. Prod.* 2. p. 402. ☉

Buona Vista. Bonny-vis. (*Green.*)

Hughes. 216.

Chinese Lablab. *Sweet.*

- cultratus.* L. leguminibus apice uncinatis. *De Cand. Prod.* 2. p. 402. ☉
Dolichos cultratus. Willd. *Sp. Pl.* 3. p. 1039.
Dolichos ensiformis. Thumb. *Jap.* 279.
 Buona Vista. Bonny-vis. (*Small sort.*)
Hughes. 216.
 Sharp-edged Lablab. *Sweet.*
- leucocarpus.* L. leguminibus sublanceolatis compressis torosis, pericarpio crispo undulato difficile detractili, semine subgloboso glandula basilari hemisphærica obtusa. *De Cand. Prod.* 2. p. 402. ☉
 Buona Vista. Bonny-vis. (*Moonshine.*)
Hughes. 216.
 White-podded Lablab. *Sweet.*

VIGNA.

Papilionaceæ (*Dolichos*). Lin.—Leguminosæ. *De Cand.*
 (*Dolichos*). *Juss.*

- glabra.* V. volubilis glabra, calycis labio superiore obtuso. *De Cand. Prod.* 2. p. 401. ☉
Dolichos luteolus. Willd. *Sp. Pl.* 3. p. 1038. *Jacq. Vind.* 1. p. 39. t. 90.

Phaseolus, &c. siliqua longa ventricosa.

Sl. 1. *p.* 183.

Yellow Vigna. *Sweet.*

Rouncivall Peas.

DOLICHOS.

Papilionaceæ, *Lin.*—Leguminosæ. *Juss. De Cand.*

luteus. D. caule volubili repente herbaceo glabro, foliolis subrotundo-ovato-rhombeis glabris, racemis spicatis petiolo longioribus, leguminibus subcylindricis glabris. *De Cand. Prod.* 2. *p.* 398. *Willd. Sp. Pl.* 3. *p.* 1050. *Swtz. Fl.* 1246. ☉

Eatable Wild Pea. *Hughes.* 216.

Yellow-flowered Dolichos. *Sweet.*

Monkey's Peas.

unguiculatus. D. caule volubili foliisque glabris, pedunculis folii longitudine apice flores 2—3 subumbellatos gerentibus, leguminis rostro recurvo. *De Cand. Prod.* 2. *p.* 400. *Willd. Sp. Pl.* 3. *p.* 1039. *Hort. Kew.* 4. *p.* 292. ☉

Phaseolus erectus, &c. seminibus oblongis albis. *Br.* 292.

Phaseolus erectus minor, &c. semine albo.

Sl. 1. *p.* 183.

Cuckold's Increase (*small*). *Hughes.* 216.

Long. 2. *p.* 787.

Bird's-foot *Dolichos.* *Sweet.*

barba-
densis.

D. caule procumbente, foliolis lateralibus angulato-ovatis: terminali hastato-subtrilobo, stipulis falcatis acutis, leguminibus capitatis cylindraceis polyspermis, rostro cochleariformi. M—. ☉

Galivaughn* Peas.

sinensis.

D. caule subvolubili herbaceo glabro, foliolis ovatis acuminatis, pedunculis bifloris folio brevioribus, leguminibus teretibus torulosis pendulis. *De Cand.*

Prod. 2. *p.* 400. *Willd. Sp. Pl.* 3. *p.*

1038. *Bot. Mag. t.* 2232. *Jacq. Vind.*

3. *p.* 39. *t.* 71. ☉

Phaseolus erectus, &c. seminibus rufescentibus oblongis. *Br.* 292.

Phaseolus erectus major, &c. semine rubro. *Sl.* 1. *p.* 183. *t.* 115. *f.* 2, 3.

Calavances, or Red Bean. *Long.* 2. *p.* 786.

Chinese *Dolichos.* *Sweet.*

Clay Peas.

sesquipe-
dalis.

D. caule volubili glabro, foliolis late ovatis, leguminibus subcylindraceis apice

* Probably a corruption of the word Calavance, applied in the West-India Islands to several kinds of pulse.

mucronato-uncinatis lævibus longissimis
ad semina torulosis. *De Cand. Prod.* 2. p.
400. *Willd. Sp. Pl.* 3. p. 1040. *Hort. Kew.*
4. p. 293. *Jacq. Vind.* 1. p. 27. t. 67. ☉
Cockold's Increase (*large*). *Hughes.* 216.
Long-podded Dolichos. *Sweet.*
Halifax Peas.

sphæros- *D. caule erecto ramoso glabro, foliolis ova-*
permus. *tis subacutis, pedunculis elongatis apice*
paucifloris, leguminibus rectis, seminibus
sphæricis. *De Cand. Prod.* 2. p. 400. ☉
Phaseolus sphærospermus. *Willd. Sp.*
Pl. 3. p. 1037.

Phaseolus erectus siliquis gracilibus.
Br. 292.

Phaseolus erectus minor, &c. *Sl.* 1. p.
184. t. 117.

Black-eyed Peas. *Long.* 2. p. 787.

CANAVALIA.

Papilionaceæ (Dolichos). Lin.—Leguminosæ. De Cand.
(Dolichos.) Juss.

ensiformis. *C. foliolis ovatis acutis, leguminibus lati-*
tudine quintuplo et ultra longioribus.
De Cand. Prod. 2. p. 404. ☉

Dolichos ensiformis. Willd. *Sp. Pl.* 3. p.

1050. *Hort. Kew.* 4. p. 295. *Aubl.* 764.

Dolichos acinaciformis. Jacq. *Ic.* 3. t. 559.

Phaseolus suberectus major. Br. 291.

Phaseolus maximus siliqua ensiformi.

Sl. 1. p. 177. t. 114. f. 1—3.

Horse-Bean. *Hughes.* 215. *Browne.*

Cimeter-podded Canavalia. *Sweet.*

RHYNCHOSIA.

Papilionaceæ (*Dolichos*). Lin — Leguminosæ. *De Cand.*

(*Dolichos*.) *Juss.*

minima. R. caule volubili gracili angulato subpubescente, foliolis rhombeis acutis glabriusculis subtus punctatis, racemis folio longioribus, floribus reflexis, leguminibus oblongis basi attenuatis tenuissime subvelutinis. *De Cand. Prod.* 2. p. 385. ☉

Dolichos minimus. Willd. *Sp. Pl.* 3. p.

1044. *Hort. Kew.* 4. p. 294. Br. 294.

Aubl. 764.

Phaseolus minimus foetidus. *Sl.* 1. p.

182. t. 115. f. 1.

Least Wild Pea Vine. *Hughes.* 198.

LATHYRUS.

Papilionaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

- odoratus.* *L. hirsutus*, caulibus diffusis alatis, foliolis ovatis mucronulatis, stipulis semisagittato-lanceolatis petiolo multo brevioribus, pedunculis 2—3—floris folio multo longioribus, leguminibus oblongo-linearibus compressis hirsutis. *De Cand. Prod.* 2. p. 374. *Willd. Sp. Pl.* 3. p. 1083. *Hort. Kew.* 4. p. 307. ☉
- α. florum vexillis purpureis. *Bot. Mag.* t. 60.
- β. florum vexillis roseis.
- Sweet Pea. *Sweet.*

SESBANIA.

Papilionaceæ (*Coronilla*). *Lin.*—Leguminosæ. *Juss. De Cand.*

- occidentalis.* *S. fruticosa* glabra, foliolis ellipticis 12—15—jugis, racemis paucifloris, leguminibus teretibus rectis petiolo triplo longioribus. *De Cand. Prod.* 2. p. 264. ½

Coronilla occidentalis. *Willd. Sp. Pl.* 3.
p. 1147. *Plum. Ic. t.* 125. *f.* 1.
 West Indian *Sesbania.* *Sweet.*

AGATI.

Papilionaceæ (Coronilla). *Lin.*—*Leguminosæ.* *De Cand.*
(Sesbania.) Juss.

*grandi-
 flora.*

*A. foliolis glabris, leguminibus evidenter
 compressis.* *De Cand. Prod.* 2. *p.* 266. $\mathfrak{h}$
Sesbania grandiflora. *Hort. Kew.* 4. *p.*
 330.

Coronilla grandiflora. *Willd. Sp. Pl.* 3.
p. 1145.

Æschynomene grandiflora. *Lin. Sp. Pl.*
 1050.

α . floribus albis.

β . floribus rubris.

Great-flowered Agati. *Sweet.*

POLYADELPHIA DECANDRIA.

THEOBROMA.

Columniferæ. *Lin.*—Malvaceæ. *Juss.*—Byttneriaceæ. *De Cand.*

Cacao. T. foliis integerrimis ovato-oblongis
acuminatis utrinque glaberrimis con-
coloribus. *De Cand. Prod.* 1. p. 484.

Willd. Sp. Pl. 3. p. 1422. *Hort. Kew.*
4. p. 408. h

T. fructu subrotundo. *Br.* 306.

T. fructu ovato acuminato. *Id. ibid.*

Cacao. *Sl.* 2. p. 15. t. 160. *Catesb. Car.*

App. 6. t. 6.

Cocoa Tree. *Hughes.* 195.

Chocolate-Nut Tree. *Miller.*

POLYADELPHIA ICOSANDRIA.

CITRUS.*

Bicornes. *Lin.*—Aurantia. *Juss.*—Aurantiaceæ. *De Cand.*

- medica.* C. petiolis nudis, foliis oblongis acutis, floribus 40—andris sæpe agynis, fructuum oblongorum rugosorum cortice crasso, pulpa acidula. *De Cand. Prod.* 1. p. 539. *Willd. Sp. Pl.* 3. p. 1426. *Hort. Kew.* 4. p. 419. b
- α. fructu ovato acido. *Br.* 308.
Limo arbor. *Sl.* 2. p. 178.
St. Helena Lemon Tree. *Hughes.* 128.
Spanish Lemon Tree. *Id. ibid.*

* “It is very difficult to determine what is a variety and what a species in this genus,”—*Martyn in Gard. Dict.* I am sorry to feel that this difficulty has not yet been overcome. The whole genus requires revision, and a correct monograph of it would be highly acceptable.

β . fructu majori acido. *Br.* 308.

Wild Shaddock. Large Lime Tree.

Hughes. 125.

γ . fructu oblongo majori. *Br.* 309.

Citrus arbor. *Sl.* 2. p. 176.

Citron Tree. *Hughes.* 127.

Lemon Tree. *Sweet.*

Limetta. C. petiolis nudis, foliis ovatis rotundatis serratis, floribus 30—andris, fructuum globosorum umbone obtuso coronatorum cortice firmo, pulpa dulci. *De Cand. Prod.* 1. p. 539. h

Burgamot Tree. *Hughes.* 127.*

Bergamotte. *Sweet.*

Sweet Lime?

Limonum. C. petiolis subalatis, foliis oblongis acutis dentatis, floribus 35—andris sæpe agynis, fructuum globosorum oblongorum cortice tenuissimo, pulpa acidissima. *De Cand. Prod.* 1. p. 539. h

C. medica. β . *Willd. Sp. Pl.* 3. p. 1426.
Hort. Kew. 4. p. 419.

C. fructu sphærico-ovato. *Br.* 308.

Malus Aurantia fructu Limonis pusillo.
Sl. 2. p. 182.

Lime Tree. *Hughes.* 129. *Sweet.*

* The characters assigned to this species, commonly known as the Bergamotta and Limo dulcis, does not well accord with Hughes's description of his Burgamot Tree.

*Auran-
tium.*

C. petiolis subnudis, foliis ovato-oblongis acutisque, floribus 20—andris, fructuum globosorum cortice tenui, pulpa dulci.

De Cand. Prod. 1. p. 539. h

C. Aurantium. β . *Willd. Sp. Pl.* 3. p. 1427. *Hort. Kew.* 4. p. 420.

Malus Aurantia sinensis. *Sl.* 2. p. 181.

China Orange Tree. *Hughes.* 126. *Aiton.*

vulgaris.

C. petiolis alatis, foliis ellipticis acutis crenulatis, floribus 20—andris, fructuum globosorum cortice tenui scabroso, pulpa acra amara.

De Cand. Prod. 1. p. 539. h

C. Aurantium. α . *Willd. Sp. Pl.* 3. p. 1427. *Hort. Kew.* 4. p. 420.

C. Aurantium. γ . *myrtifolius.* *Bot. Reg.* t. 346.

C. fructu sphærico acido. *Br.* 308.

Malus Aurantia vulgaris. *Sl.* 2. p. 179.

Sour Orange Tree. *Hughes.* 125.

Golden Orange Tree. *Id.* 126.

Seville Orange. *Aiton.*

*Decu-
mana.*

C. petiolis alatis, foliis obtusis emarginatis, fructibus maximis turbinato-subrotundis cortice crasso scabro. M—. *De*

Cand. Prod. 1. p. 539. h

C. fructu sphærico-obovato maximo, *Br.* 309.

Malus Aurantia fructu rotundo maximo.

Sl. 1. p. 41. t. 12. f. 2—4.

Shaddock Tree. *Hughes*. 124.

Lesser Shaddock Tree. *Id.* 125.

Pumple-Nose Tree. *Id.* 126.

Shaddock. *Sweet*.

Hystrix? C. petiolis alatis, foliis ovatis obtusis,
floribus racemosis, fructibus sphæricis
cortice subcrasso lævi. M—. *De Cand.*
Prod. 1. p. 539. h

Grape-Fruit Tree.*

buxifolia? C. petiolis lineari-alatis, foliis ovatis re-
tusis, floribus racemosis, fructibus ovatis
cortice subcrasso lævi. M—, *De Cand.*
Prod. 1. p. 540. h

C. fructu sphærico-ovato minori. *Br.*
309.

Forbidden-Fruit Tree. *Hughes*. 127.

* My notes of the Grape-Fruit and Forbidden-Fruit Trees, I am sorry to find, are too imperfect to enable me to say with certainty that they are specifically distinct, although I am inclined to think they are. I feel quite certain that they are not varieties of C *Decumana*, as they have by some been supposed to be.

SYNGENESIA POLYGAMIA ÆQUALIS.

SONCHUS.

Compositæ. *Lin.*—Cichoracæ. *Juss.*

agrestis. S. pedunculis tomentosis multifloris, calycibus glabris, caule striato, foliis incisoserratis sessilibus. *Willd. Sp. Pl. 3. p. 1513. Swtz. 1290.* ☉

S. lævis. *Sl. 1. p. 255.*

Rabbit-Weed. *Hughes. 209.*

oleraceus. S. pedunculis subtomentosis umbellatis, calycibus glabris, foliis oblongo-lanceolatis, amplexicaulibus denticulatis subsinuatis. *Willd. Sp. Pl. 3. p. 1514. Hort. Kew. 4. p. 437. Fl. Br. 817. Engl. Bot. t. 843.* ☉

Thistle, or Rabbit-Weed. *Hughes. 172.*

Common Sow-Thistle. *Miller.*

LACTUCA.

Compositæ. *Lin.*—Cichoraceæ. *Juss.*

- sativa.* L. foliis rotundatis; caulinis cordatis, caule corymboso. *Willd. Sp. Pl.* 3. p. 1523. *Hort. Kew.* 4. p. 440. ☉
Garden Lettuce. *Miller.*

BIDENS.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

- leucantha.* B. floribus radiatis calyce exteriori longitudine interioris, foliis inferioribus pinnatis; superioribus ternatis, foliolis ovatis serratis basi æqualibus. *Willd. Sp. Pl.* 3. p. 1719. ☉
Corcopsis leucantha. *Lin. Sp. Pl.* 1282.
Spanish Needle. *Hughes.* 214.*
White-flowered Bidens. *Sweet.*

* This vernacular name, derived from the seed, is frequently applied to all the species of *Bidens* herein enumerated.

pilosa. B. floribus discoideis, calyce exteriori longitudine interioris, foliis inferioribus pinnatis; superioribus ternatis: foliolis oblongis: terminali lanceolato reliquis duplo longiore. *Willd. Sp. Pl. 3. p. 1719.*

Hort. Kew. 4. p. 495. Aubl. 794. ☉

Hairy Bidens. *Miller.*

bipinnata. B. floribus subradiatis, calyce exteriori longitudine interioris, foliis pinnatis: foliolis lanceolatis pinnatifidis. *Willd.*

Sp. Pl. 3. p. 1721. Hort. Kew. 4. p. 496. ☉

Hemlock-leaved Bidens. *Miller.*

EUPATORIUM.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

odoratum. E. foliis petiolatis triangulari-ovatis serratis apice integerrimis subtus pubescentibus, corymbis divaricatis terminalibus sessilibus. *Willd. Sp. Pl. 3. p. 1767.*

Hort. Kew. 4. p. 509. Br. 313. ♀

Eupatoria conyzoides. *Pluk. Phyt. t. 177. f. 3.*

Christmas Bush. *Hughes. 205.*

Sweet-scented Hemp Agrimony. *Miller.*
Archangel. *Browne.*

AGERATUM.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

conyzoides.

A. foliis ovatis subcordatis, caule piloso,
paleis pappi aristatis denticulatis. *Willd.*

Sp. Pl. 3. p. 1773. Hort. Kew. 4. p. 509.

Aubl. 799.

⊙

Conyza urticæ folio. Sl. 1. p. 258. t.
152. f. 2.

Hairy Ageratum. *Miller.*

SYNGENESIA POLYGAMIA SUPERFLUA.

ARTEMISIA.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

gallica. A. foliis incanis; radicalibus bipinnatis lineari-filiformibus, caule erecto ramoso tomentoso, ramis erectis, floribus oblongis subsessilibus erectis. *Willd. Sp. Pl. 3. p. 1834. Hort. Kew. 5. p. 7. Engl. Bot. t. 1001.* 2
Upright-flowered Sea-Wormwood. *Aiton.*
French Wormwood.

Absinthium. A. foliis incanis; radicalibus triplicatopinnatifidis: laciniis lanceolatis dentatis obtusis; caulinis bipinnatifidis pinnatifidisve: laciniis lanceolatis acutiusculis; floralibus indivisis lanceolatis, floribus globosis pedunculatis nutantibus. *Willd. Sp. Pl. 3. p. 1844. Hort. Kew. 5. p. 8. Fl. Br. 864. Engl. Bot. 1230.* 2

Common Wormwood. *Aiton.*
English Wormwood.

BACCHARIS.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

odorata. B. fruticosa, ramis tomentosis, foliis ovatis subacutis obscure serratis subtus canis, corymbo terminali dense composito. *Ham. Prod.* 52. h

Conyza odorata. *Willd. Sp. Pl.* 3. p. 1944. *Hort. Kew.* 5. p. 30.

Conyza foliis lanceolatis ovatis. *Plum. Ic.* 85. t. 97.

Sweet-scented Flea-Bane. *Aiton.*
Wild Tobacco.

TUSSILAGO.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

nutans. T. scapo unifloro nudo, flore radiato nutante, foliis petiolatis obovatis dentatis

basi sinuatis subtus tomentosis. *Willd.*

Sp. Pl. 3. p. 1965. *Hort. Kew.* 5. p. 34. ☉

Leontodon foliis radicalibus oblongis.

Br. 310.

Dens Leonis folio subtus incano. Sl. 1.

p. 255. *t.* 150. *f.* 2.

Silver-leaved Plant. *Hughes.* 222.

Drooping-flowered Colt's-Foot. *Aiton.*

ASTER.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

dumosus. A. foliis linearibus integerrimis glabris;
ramulorum brevissimis, ramis panicu-
latis, calycibus cylindraceis arcte imbri-
catis. *Willd. Sp. Pl.* 3. p. 2026. *Hort.*
Kew. 5. p. 52. 4

α. radio pallide violaceo.

β. radio albo.

Bushy Starwort. *Miller.*

chinensis. A. foliis ovatis grosse dentatis petiolatis;
caulinis sessilibus basi cuneatis; flora-
libus lanceolatis integerrimis, caule his-
pido, ramis unifloris, calycibus foliaceis.
Willd. Sp. Pl. 3. p. 2038. *Hort. Kew.* 5.
p. 58. 4

Callistema hortensis. *Hort. Br.* 226.
Chinese Starwort, or China-Aster. *Miller.*

SOLIDAGO.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

cambrica. *S.* caule simplissimo pubescente, foliis
cuneiformi-lanceolatis serratis pilosius-
culis, racemis erectis, ligulis elongatis.

Willd. Sp. Pl. 3. p. 2066. *Hort. Kew.*
5. p. 70. 24

S. virgaurea. γ . *cambrica.* *Fl. Br.* 889,
Welch Golden Rod. *Miller.*

DAHLIA.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

*superflua,** *D.* caule non pruinoso, ligulis foemineis.

Hort. Kew. 5. p. 87. *Bot. Mag. t.* 1885.

A. & B. Bot. Reg. t. 55. 24

* The varieties of this species are very readily multiplied by culture, both as to colour and fulness. The only varieties I have seen in Barbados are the single and double-flowered purple and the red-flowered.

D. pinnata. *Bot. Repos. t.* 408.

Georgina purpurea. *Willd. Sp. Pl. 3.*
p. 2124.

Georgina rosea. *Willd. Sp. Pl. 3. p.* 2124.

Fertile-rayed Dahlia. *Aiton.*

TAGETES.

Compositæ. Lin.—Corymbiferae. Juss.

patula.

T. foliis pinnatis: foliolis lanceolatis ciliato-serratis, pedunculis unifloris sub-incrassatis, calycibus lævibus, caule patulo. *Willd. Sp. Pl. 3. p.* 2126. *Hort. Kew. 5. p.* 88. ☉

α. floribus simplicibus. *Bot. Mag. t.* 150.
Single - flowered French Marigold.
Aiton.

β. floribus multiplicatis.

Double - flowered French Marigold.
Aiton.

erecta.

T. foliis pinnatis: foliolis lanceolatis ciliato-serratis, pedunculis unifloris ventricosο-incrassatis, calycibus angulatis, caule erecto. *Willd. Sp. Pl. 3. p.* 2127.
Hort. Kew. 5. p. 89. *Aubl.* 802. ☉

α . floribus simplicibus.

Single-flowered African Marigold.

Aiton.

β . floribus multiplicatis.

Double-flowered African Marigold.

Aiton.

tenuifolia. T. foliis pinnatis: foliolis linearibus serratis: serraturis inferioribus elongatis, caule paniculato, pedunculis alternis unifloris, calycibus clavatis. *Willd. Sp. Pl.*

3. p. 2128. *Hort. Kew.* 5. p. 89. *Bot.*

Mag. t. 2045.

⊙

Fine-leaved Tagetes, or Peruvian Marigold. *Aiton.*

PECTIS.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

humifusa. P. foliis oblongo-ovatis subtus punctatis, basi utrinque ciliatis, caulibus procumbentibus. *Willd. Sp. Pl.* 3. p. 2130.

Swtz. 1362.

⊙

Melampodium foliis lanceolatis. *Plum.*

Ic. t. 95. f. 2.

Trailing Pectis. *Sweet.*

HETEROSPERMUM.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

ovatum. H. caule villosa, foliis ovatis apice serratis. *Willd. Sp. Pl. 3. p. 2129.* ☉

ZINNIA.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

elegans. Z. floribus pedunculatis, foliis oppositis cordato-ovatis sessilibus amplexicaulibus, caule hirsuto, paleis corollæ serratis.
Willd. Sp. Pl. 3. p. 2140. Hort. Kew. 5. p. 92. Bot. Mag. t. 527. ☉
 Z. violacea. *Bot. Repos. t. 55.*
 Purple Zinnia. *Miller.*

CHRYSANTHEMUM.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

- sinense.* C. foliis coriaceis petiolatis sinuato-pin-
natifidis dentatis glaucescentibus, radio
longissimo, caule fruticoso. *Sabine Lin.*
Tr. 14. p. 145. h
C. indicum. *Hort. Kew.* 5. p. 95. *Bot.*
Mag. t. 327. *Bot. Reg.* t. 455. t. 616.
Anthemis artimesiæfolia. *Willd. Sp. Pl.*
3. p. 2184.
Variable Chinese Chrysanthemum. *Sweet.*

ECLIPTA.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

- punctata.* E. caule erecto scabro punctato, foliis
lanceolatis sessilibus serratis. *Willd. Sp.*
Pl. 3. p. 2217. ⊙
Bellis ramosa. *Jacq. Amer.* 216. t. 129.
Dotted-stalked Eclipta. *Miller.*

ANTHEMIS.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

- bupthal-*
moides. *A.* foliis oppositis petiolatis ovatis serratis triplinerviis subtus pubescentibus supra scabris, radio multifloro. *Willd. Sp. Pl.* 3. p. 2185. *Hort. Kew.* 5. p. 108. h
Bupthalmum scabrum. *Cav. Ic.* 2. p. 53. t. 167.
Wild Marigold. *Hughes.* 171.
Oval-leaved Peruvian Chamomile. *Aiton.*

VERBESINA.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

- nodiflora.* *V.* foliis oppositis ovatis serratis triplinerviis, floribus axillaribus subsessilibus, calycibus simplicibus, caule trichotomo. *Willd. Sp. Pl.* 3. p. 2226. *Hort. Kew.* 5. p. 121. *Aubl.* 803. ⊙
V. erecta hirsuta. *Br.* 319.

332 SYNGENESIA POLYGAMIA SUPERFLUA.

Bidens nodiflora. *Dill. Elth.* 53. *t.* 45.
f. 53.

Synedrella nodiflora. *Hort. Br.* 235.

Chrysanthemum conyzoides nodiflorum.
Sl. 1. *p.* 262. *t.* 154. *f.* 4.

Single-flowered *Verbesina.* *Miller.*

SYNGENESIA POLYGAMIA FRUSTRANEA.

HELIANTHUS.

Compositæ. *Lin.*—Corymbiferæ. *Juss.*

- annuus.* H. foliis omnibus cordatis trinervatis,
pedunculis incrassatis, floribus cernuis.
Willd. Sp. Pl. 3. p. 2237. Hort. Kew. 5.
p. 126 ☉
Helenium indicum maximum. *Bauh.*
Pin. 276.
Annual Sun-Flower. *Miller.*

COSMEA.

Compositæ. *Lin.*—Corymbiferæ. *Juss.*

- sulphurea.* C. foliis bipinnatifidis: laciniis lanceo-
latis, calycis exterioris laciniis lan-

ceolatis. *Willd. Sp. Pl. 3. p. 2250. Hort.*

Kew. 5. p. 132. ☉

Coreopsis artemisiæfolia. Jacq. Ic. 3. t. 595.

Southern-Wood-leaved Cosmea: *Aiton.*

bipinnata. C. foliis bipinnatis: foliolis lineari-subulatis, calycis exterioris laciniis ovatis (lineari-lanceolatis). *Willd. Sp. Pl. 3. p. 2250. Hort. Kew. 5. p. 130. Bot. Mag. t. 1535.* ☉

Cosmos bipinnatus. Cav. Ic. 1. p. 10. t. 14.

Double Wing-leaved Cosmea. *Aiton.*

COREOPSIS.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

reptans. C. foliis serratis ovatis; summis ternatis, caule repente. *Willd. Sp. Pl. 3. p. 2255.*

Hort. Kew. 5. p. 135. Aubl. 804. 4

Verbesina foliis inferioribus simplicibus. Br. 319.

Chrysanthemum trifoliatum scandens.

Sl. 1. p. 261. t. 154. f. 2, 3.

Trailing Coreopsis. *Miller.*

SYNGENESIA POLYGAMIA NECESSARIA.

CALENDULA.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

officinalis. C. seminibus cymbiformibus muricatis incurvatis omnibus. *Willd. Sp. Pl. 3. p. 2340. Hort. Kew. 5. p. 166.* ☉
Garden Marigold. *Miller.*

PARTHENIUM.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

Hystérophorus. P. foliis bipinnatis. *Willd. Sp. Pl. 3. p. 2385. Hort. Kew. 5. p. 181. Aubl. 854. Bot. Mag. t. 2275.* ☉

336 SYNGENESIA POLYGAMIA NECESSARIA.

P. subhirsutum ramosum. *Br.* 340.

Absinthium erysimi folio. *Pluk. Alm.* 2.

Phyt. t. 45. *f.* 3.

Cut-leaved Parthemium, or Bastard
Feverfew. *Miller.*

Wild Wormwood. *Browne.*

SYNGENESIA POLYGAMIA SEGREGATA.

ELEPHANTOPUS.

Compositæ. *Lin.*—Corymbiferae. *Juss.*

- spicatus.* *E. foliis ovato-lanceolatis serratis scabris, fasciculis florum sessilibus lateralibus, caule ramoso. Willd. Sp. Pl. 3. p. 2390. Swtz. Fl. 1380. Aubl. 808. 2*
E. erectus hirsutus. Br. 311.
Conyza major inodora. Sl. 1. p. 256. t. 150. f. 3, 4.
Spiked Elephant's-Foot. Sweet.

GYNANDRIA MONANDRIA.

EPIDENDRUM.

Orchideæ. *Lin. Juss. Brown.*

- ciliare.* E. caule simplici, foliis binis oblongis
aveniis, lamina labelli tripartita: lacinia
intermedia lineari. *Willd. Sp. Pl.* 4. p.
119. *Hort. Kew.* 5. p. 219. *Aubl.* 822.
Jacq. Amer. 224. t. 179. f. 89. *Bot.*
Mag. t. 463. *Bot. Reg.* t. 784. 2
Wild Lily. *Hughes.* 229.
Fringed Epidendrum. *Sweet.*

ONCIDIUM.

Orchideæ. *Lin. Juss. Brown.*

- Papilio.* O. foliis solitariis ovalibus pictis paten-
tibus, scapo articulato ancipiti paucifloro,

sepalis superioribus linearibus longis-
simis: inferioribus distinctis ovato-lan-
ceolatis undulatis: columna bicorni: alis
fimbriatis. *Bot. Reg. t. 910.* 2

Butterfly-Oncidium. *Sweet.*

Trinidad Butterfly-Plant.

GYNANDRIA HEXANDRIA.

ARISTOLOCHIA.

Sarmentaceæ. *Lin.*—Aristolochiæ. *Juss. Brown.*

Asarinæ. *Kunth.*

odoratissima. A. foliis cordato-ovatis, caule volubili, pedunculis folio longioribus unifloris, labio cordato-lanceolato corolla longiore.

Willd. Sp. Pl. 4. p. 156. Hort. Kew. 5. p. 224. Aubl. 833. h

A. scandens, foliis cordatis acuminatis. *Br. 329.*

A. scandens odoratissima. *Sl. 1. p. 162. t. 104. f. 1.*

Junction Vine. *Hughes. 199.*

Sweet-scented Birthwort. *Miller.*

MONŒCIA MONANDRIA.

EUPHORBIA.*

Tricoccæ. *Lin.*—Euphorbiæ. *Juss.*—Euphorbiaceæ. *Brown.*

*antiquo-
rum.*

E. aculeata subnuda triangularis articulata, ramis patentibus. Willd. Sp. Pl. 2. p. 881. Hort. Kew. 3. p. 155. h

Tithymalus azoides triangularis et quadrangularis. Com. Præl. 55. t. 5.

Spreading Three-sided Spurge. Sweet.

* “Euphorbia, placed by Linnæus in the class Dodecandria, is now understood by Jussieu (*Gen. Pl.*) and Brown (*Terr. Austr.*) to have separated flowers, conformably to the general character of its natural order. What Linnæus and others have taken for stamens appear to be distinct monandrous barren flowers;” “consequently Euphorbia is to be removed to Monœcia Monandria.”—*Sm. Engl. Fl. 2. p. 349. See also Gram. of Bot. p. 185.* The successive manner in which the supposed stamens are evolved is a strong argument in favour of their being flowers; and I have had no hesitation in adopting the above arrangement.

neriefolia. *E. aculeata* seminuda, aculeis geminatis, angulis oblique tuberculatis superne foliosis, foliis oblongis. *Willd. Sp. Pl.* 2. p. 884. *Hort. Kew.* 3. p. 157. h

Tithymalus azoides arborescens. *Com. Præl.* 56. t. 6.

Oleander-leaved Spurge. *Sweet.*

Tithyma-
loides. *E. inermis* fruticosa, foliis distichis alternis ovatis. *Willd. Sp. Pl.* 2 p. 890. *Hort. Kew.* 3. p. 159. h

α . foliis acutis. *Jacq. Amer.* 149. t. 92.

Pedilanthus Tithymaloides. *Ham. Prod.* 43. *Bot. Reg.* t. 837.

Tithymalus curassavicus myrtifolius.

Pluk. Alm. 369. *Phyt. t.* 230. f. 2.

Crepidaria myrtifolia. *Haw. Succ.* 136.

Myrtle-leaved Spurge. *Aiton.*

β . foliis obtusis vel emarginatis.

Tithymaloides lauro-cerasi folio. *Dill. Elth.* 383. t. 288. f. 372.

Pedilanthus padifolius. *Ham. Prod.* 42.

Crepidaria padifolia. *Haw. Succ.* 136.

Laurel-leaved Spurge. *Aiton.*

Tithymaloides. *Hughes.* 165.

glabrata. *E. frutescens* ramosa, foliis oppositis cordato-ovatis acutis glabris integerrimis.

Willd. Sp. Pl. 2. p. 893. *Swtz. Fl.* 870. h

Peplis fruticosa maritima geniculata. *St. l.* p. 198.

Smooth Spurge. *Miller.*

Sea-Side Milk-Weed.

atropur- *E. caule trichotomo, ramis geniculatis,*
purea. *foliis ternis ovato-lanceolatis subcordatis*
longe petiolatis coloratis subtus glaucis,
bracteis linearibus subquaternis deciduis,
umbella multifida radiis subquadrifidis,
involucris obtusis basi connatis. M—. *h*

E. atropurpurea. Spreng. Syst. 3. p. 790?

Dark Purple Spurge. *Sweet.*

Indian Rubber Tree.*

hyperici- *E. dichotoma, foliis serratis ovali-ob-*
folia.† *longis glabris, corymbis terminalibus*
ramis divaricatis. Willd. Sp. Pl. 2. p.
895. Hort. Kew. 3. p. 161. Aubl.
479.

⊙

* This shrubby tree generally goes by the name of the Indian Rubber in Barbados, but the appellation is also applied to the *Ficus elastica*, lately introduced into the island, and at present a rare exotic. I cannot be confident that I am right in making it identical with the *E. atropurpurea* of Sprengel. By altogether excluding from his plan a reference to authors, Sprengel has greatly increased the difficulty of recognising his species, and thereby circumscribed the usefulness of his work.

† I have excluded Browne's synonyme, admitted by Willdenow, as it undoubtedly must belong to some other species. Sloane's *Tithymalus* is certainly the Barbados plant in question. He describes it as having "one red stalk dividing itself into several green round branches, rising ten inches high." This

Tithymalus erectus acris. *Sl.* 1. *p.* 197?
t. 126.

Milk-Weed (*red*). *Hughes.* 163.

St. John's-Wort-leaved Spurge. *Miller.*

maculata. *E. dichotoma*, foliis serratis oblongis pilosis, floribus axillaribus solitariis, ramis patulis. *Willd. Sp. Pl.* 2. *p.* 896.
Hort. Kew. 3. *p.* 162. *Aubl.* 497. *Jacq. Vind.* 2. *p.* 87. *t.* 186. ☉

Tithymalus seu Chamæsyce. *Pluk. Alm.* 372. *t.* 65. *f.* 8.

Chamæsyce. *Sl.* 1. *p.* 198.

Dove-Weed. *Hughes.* 163.

Spotted Spurge. *Miller.*

pilulifera. *E. dichotoma*, foliis serratis ovali-oblongis, pedunculis bicapitatis axillaribus caule erecto. *Willd. Sp. Pl.* 2. *p.* 897.
Aubl. 479. *Jacq. Ic.* 3. *t.* 478. ☉

Milk-Weed (*white*). *Hughes.* 163.

Globular Spurge. *Sweet.*

must surely be different from Browne's *Euphorbia minima reclinata*—"a slender, weakly creeper, which seldom rises above three or four inches from the root." I should almost doubt Sloane's *Tithymalus* being identical with *Euphorbia hypericifolia*, had he not given a plate of it; for a plant, "floribus ad caulium nodos conglomeratis," very ill accords with the "corymbis terminalibus" of the definition of this species; nor does it, indeed, accord with the character of the inflorescence represented in the plate.

obliterata. E. umbella trifida, foliis oblongis trapezoides serratis pubescentibus altera baseos parte obliteratis. *Willd. Sp. Pl.* 2. p. 904. *Swtz. Fl.* 872. *Jacq. Amer.* 151. ☉

Milk-Weed (*red*). *Hughes.* 163.

ARTOCARPUS.

Urticæ. *Juss* —Urticæ. *De Cand.*

incisa. A. foliis pinnatifido-sinuatis scabris subtus pubescentibus, amentis masculis cernuis. *Willd. Sp. Pl.* 4. p. 188. *Hort. Kew.* 5. p. 231. h

α. nucifera.

Soccus granosus. *Rumph. Amb.* 1. p. 112. *t.* 33.

Nut-bearing Bread-Fruit. *Sweet.*

Bread-Nut Tree.

β. apyrena.

Soccus lanosus. *Rumph. Amb.* 1. p. 110. *t.* 32.

Seedless Bread-Fruit. *Sweet.*

Bread-Fruit Tree.

*integri-
folia.*

A. foliis oblongis indivisis basi attenuatis
subtus scabris. *Willd. Sp. Pl.* 4. p. 189.

Hort. Kew. 5. p. 231.

h

Jaca indica. *Bauh. Hist.* 1. p. 115.

Jaca Tree. *Sweet.*

Jack-Fruit Tree.

MONŒCIA TRIANDRIA.

TRAGIA.

Tricoccæ. *Lin.*—Euphorbiæ. *Juss.*—Euphorbiacæ. *De Cand.*

volubilis. T. foliis cordatis ovatis acuminatis serratis glabriusculis, petiolis ciliatis, foliolis calycinis fœmineis indivisis hirtis, caule scandente. *Willd. Sp. Pl.* 4. p. 322.

Hort. Kew. 5. p. 255. *Aubl.* 847. ½

T. scandens. *Br.* 336.

Urtica racemosa scandens. *Sl.* 1. p. 123.
t. 82. *f.* 1.

Vine-Nettle. *Hughes.* 220.

Twining Tragia. *Miller.*

Creeping Cowhage. *Browne.*

HERNANDIA.

Tricoccæ. *Lin.*—*Lauris* aff. *Juss.*

sonora. *H. foliis peltatis.* *Willd. Sp. Pl.* 4. p. 327.
Hort. Kew. 5. p. 256. *Jacq. Amer.* 245.
Aubl. 852. h

H. arborea. *Br.* 373.

Jack-in-a-Box. *Hughes.* 119. *Long.* 2.
p. 810.

Whistling Hernandia. *Miller.*

MONŒCIA TETRANDRIA.

CICCA.*

Euphorbiæ. *Juss.*—Euphorbiacæ. *De Cand.*

- disticha.* C. foliis pinnatis: foliolis oblongis, racemis lateralibus. *Willd. Sp. Pl.* 4. p. 332.
Hort. Kew. 5. p. 258. h.
Phyllanthus longifolia. *Jacq. Schæn.* 2.
p. 36. t. 194.
Averrhoa acida. *Lin. Sp. Pl.* 613.
Long-leaved Cicca. *Aiton.*
Otaheite Gooseberry.

* The seed vessel of the Cicca has been variously described: by some as an apple, by others as a berry. Willdenow states it to be a capsule,—“capsula tetracocca, non dehiscens subcarnosa.” It is, however, a subrotund monospermous drupe, eight furrowed, the alternate furrows deeper. The nut is subrotund, flattened at the ends, and furrowed to correspond with the drupe, four-celled, each cell single-seeded.

URTICA.

Scabridæ. *Lin.*—Urticæ. *Juss.*—Urticeæ. *De Cand.*

*grandi-
folia.*

U. foliis oppositis subrotundo-ovatis grosse serratis, stipulis ellipticis integerrimis, corymbis pedunculatis ramossissimis longitudine foliorum. *Willd. Sp. Pl.* 4. p. 349. *Hort. Kew.* 5. p. 262. *Aubl.* 853. h

U. erecta, foliis ovato-acuminatis. *Br.* 337.

U. iners racemosa sylvatica. *Sl.* 1. p. 124. t. 83. f. 2.

Nettles (*white sort*). *Hughes.* 220.

Great-leaved Nettle. *Aiton.*

*micro-
phylla.*

U. foliis ovatis acutis integerrimis minoribus oppositis et intermixtis, floribus monoicis,* caulibus simpliciusculis ascendentibus. *M—, Willd. Sp. Pl.* 4. p. 359.

Hort. Kew. 5. p. 264. *Swtz. Fl.* 305. u

U. humilior disticha. *Br.* 336.

Parietaria microphylla. Lin. Sp. Pl. 1492.

* This plant is represented by Swartz as dioecious; but in every plant which I have examined the male and female flowers were mixed. Nor have I mistaken the *U. herniarifolia* for this plant. If therefore Swartz' plant be really dioecious, this is a new species.

Herniaria lucida aquatica. *Sl.* 1. *p.* 145.
t. 93. *f.* 2.

Small-leaved Nettle. *Aiton.*

æstuans. *U. foliis alternis cordatis, racemis dichotomis, fructibus orbiculato-corymbosis.*
Willd. Sp. Pl. 4. *p.* 362. *Hort. Kew.* 5.
p. 265. *Aubl.* 854. *Jacq. Ic.* 3. *p.* 72.
t. 388. ☉

Nettles (*red sort*). *Hughes.* 220.

Surinam Nettle. *Aiton.*

MORUS.

Scabridæ. *Lin.*—Urticæ. *Juss.*—Urticeæ. *De Cand.*

tinctoria. *M. foliis oblongis basi hinc productionibus, spinis axillaribus solitariis.* *Willd.*
Sp. Pl. 4. *p.* 371. *Hort. Kew.* 5. *p.*
 267. h

M. Zanthozylum. *Jacq. Amer.* 247.

M. lactescens. *Br.* 339.

M. fructu viridi. *Sl.* 2. *p.* 3. *t.* 158. *f.* 1.

Maclura tinctoria. *Hort. Br.* 362.

Broussonetia tinctoria. *Kunth. Syn.* 1.
p. 370.

Fustic Tree. *Hughes.* 178. *Long.* 2.
p. 829.

Dyer's Mulberry, or Fustic Wood. *Miller.*

MONŒCIA PENTANDRIA.

AMARANTHUS.

Miscellanææ. *Lin.*—Amaranthi. *Juss.*—Amaranthaceæ. *Brown.*

tricolor. A. glomerulis triandris sessilibus, foliis oblongo-lanceolatis coloratis. *Willd. Sp. Pl.* 4. p. 385. *Hort. Kew.* 5. p. 272. *Aubl.* 855. ☉

A. folio variegato. *Bauh. Pin.* 121.
Three-coloured Amaranth. *Miller.*

viridis. A. glomerulis axillaribus geminatis triandris, floribus masculis triphyllis, foliis ellipticis emarginatis margine undulatis. *Willd. Sp. Pl.* 4. p. 388. *Hort. Kew.* 5. p. 274. ☉

Blitum minus album. *Sl.* 1. p. 143. t. 92. f. 1.

Caterpillar (*White*). *Hughes.* 170.
Green Amaranth. *Miller.*

- polygono-*
noides. A. glomerulis triandris triphyllis, floribus fœmineis infundibuliformibus, foliis rhombeo-ovatis emarginatis. *Willd. Sp. Pl.* 4. p. 389, *Hort. Kew.* 5. p. 275. *Aubl.* 855. ☉
Blitum polygonoides. *Sl.* 1. p. 144. t. 92. f. 2.
Chenopodium humile. *Br.* 184.
Caterpillars (*red*). *Hughes.* 170.
Spotted-leaved Amaranth. *Miller.*
- sanguineus.* A. racemis pentandris supra decompositis erectis, ramis patentibus glabris, foliis oblongis acutis. *Willd. Sp. Pl.* 4. p. 390. *Hort. Kew.* 5. p. 275. ☉
Spreading, or Bloody Amaranth. *Miller.*
- spinosus.* A. racemis pentandris terminalibus compositis, axillis spinosis. *Willd. Sp. Pl.* 4. p. 393. *Hort. Kew.* 5. p. 277. *Aubl.* 856. ☉
Blitum americanum spinosum. *Sl.* 1. p. 143.
Caterpillars (*prickly*). *Hughes.* 170.
Prickly Amaranth. *Miller.*

MONŒCIA HEXANDRIA.

GUETTARDA.

Tricoccæ. *Lin.* — Rubiaceæ. *Juss.*

argentea. G. foliis subrotundo-ovatis acutis subtus tomentosis floribus pentandris quinquefidis. *Willd. Sp. Pl.* 4. p. 399. h

Halesia arborescens. *Br.* 205. t. 20. f. 1.

scabra. G. foliis obovatis breviter et obtuse acuminatis scabris. M—. *Vent. Choix.* t. 1. h

Mathiola scabra. *Willd. Sp. Pl.* 1. p. 998.

Mathiola folio aspero subrotundo. *Plum.*

Ic. t. 173. f. 2.

Rough Guettarda. *Sweet.*

Buff-Coat Tree.

COCOS.

Palmæ. *Lin. Juss. Brown.*

- nucifera.* C. inermis, frondibus pinnatis: foliolis replicatis ensiformibus. *Willd. Sp. Pl.* 4. p. 400. *Hort. Kew.* 5. p. 279. *Aubl.* 973. *Mart. Palm.* t. 62. t. 75, t. 88. f. 3—6. h
 C. nucifer. *Jacq. Amer.* 277. t. 168.
 Palma spadicebus alaribus. *Br.* 341.
 Palma indica nucifera. *Sl.* 2. p. 8.
 Coco-Nut Tree. *Hughes.* 103. t. 3.
 Cocoa-Nut Tree. *Long.* 2. p. 738. *Miller.*

ACROCOMIA.

Palmæ. *Brown. (Cocos.) Lin. Juss.*

- fusiformis.* A. caudice fusiformi superne aculeatis, frondibus pinnatis: foliolis linearibus apice præmorsis, stipitibus spathisque aculeatis. M—. *Hort. Br.* 432. h
 Cocos fusiformis. *Willd. Sp. Pl.* 4. p. 401.
Hort. Kew. 5. p. 280. *Swtz. Fl.* 616.

Cocos aculeata. *Swtz. Prod.* 151.

Palma pinnis, &c. aculeatissimis. *Br.* 344.

Palma tota spinosa major. *Sl.* 2. p. 119.

Mackaw Tree. *Hughes.* 114.

Great Mackaw Tree. *Long.* 2. p. 740.

Miller.

MONŒCIA POLYANDRIA.

CERATOPHYLLUM.

Inundatæ. *Lin.*—Naides. *Juss.*—Fluviales. *Kunth.*

- demersum.* C. foliis dichotomo-bigeminis, fructibus
trispinosis. *Willd. Sp. Pl.* 4. p. 405.
Hort. Kew. 5. p. 281. *Fl. Br.* 1020.
Engl. Bot. t. 947. 24
C. foliis verticillatis. *Br.* 345.
Small Duck Weed. *Hughes.* 234.
Prickly-seeded Hornwort. *Miller.*
Morass Weed. *Browne.*

BEGONIA.

Holeraceæ. *Lin.*—Incertæ sedis. *Juss.*—Begoneaceæ. *Brown.*

- crenata.* B. caulescens, foliis inæqualiter cordatis
subrotundis obtusis crenato-dentatis,

capsulis bilocularibus. *Willd. Sp. Pl.* 4. p. 414. *Dryander Lin. Tr.* 1. p. 162. t. 14. f. 3. h

CALADIUM.

Piperitæ (*Arum*), *Lin.*—Aroideæ. *Juss. Brown.*

macrorhi- C. subacaule* foliis profunde cordatis
zon. repandis. *Brown. Prod.* 192. u

Arum macrorhizon, Willd. Sp. Pl. 4. p. 482.

Arum peregrinum. Willd. Sp. Pl. 4. p. 482.

Arum maximum. Macrorhizon zeylanicum. Herm. Par. 73. t. 73.

Arum indicum sativum. Rumph. Amb. 5. p. 308. t. 106.

Arum acaule maximum. Br. 332. No. 5.

Roasting Eddas. *Hughes.* 227.†

* The cultivated plant in Barbados is stemless: the root not being allowed to remain in the ground long enough to become caulescent.

† The roasting Eddas (*Eddoes*) of Hughes is certainly not identical with the *Arum maximum ægyptiacum* of Caspar Bauhin, *Arum Colocasia* (*Willd.*) as has been concluded on the autho-

Long-rooted Caladium (*Arum*). *Sweet*.

White Cocco and Tyre. *Browne*.

nymphæi- C. acaule, foliis peltatis cordato-sagit-
folium. tatis, spadice spatha cylindracea apice
lanceolata longiore. *Willd. Sp. Pl.* 4. p.
488. *Hort. Kew.* 5. p. 311. 2

Arum acaule medium. *Br.* 332. *No.* 7.*

Water-Lilly-leaved Caladium. *Aiton*.

Indian Kale. *Browne*.

Wild, or Monkey Eddoe.

sagitti- C. acaule, foliis sagittatis acuminatis,
folium. spadice spatha ovata cucullata brevior.
Willd. Sp. Pl. 4. p. 489. *Hort. Kew.* 5.
p. 311 2

rity of Sloane and Browne. The leaf of the *Arum Colocasia* is peltated; that of the roasting Eddoe is not. Mr. Miller considered the *A. peregrinum* as the species "most commonly cultivated in the West Indies, under the name of Edder." I am confident that the *Arum peregrinum*, and the *A. macrorhizon* are not distinct species, but are both identical with the *Arum max. &c.* of Hermannus. The definition therefore of *Arum macrorhizon*, representing the plant with a peltated leaf, in opposition to Hermannus from whom the specific title is taken, should be excluded; the two species in question should be considered as one, the title *macrorhizon* adopted, and the species thus arranged be carried to the genus *Caladium*, to which it undoubtedly belongs.

* Browne is certainly in error; making his *Arum*, *No.* 7, the same as Sloane's *A. Nymphææ foliis*, which is properly recognised by Willdenow as *Caladium esculentum* (*Arum esculentum*), and which I have no doubt is Browne's *Arum*, *No.* 6.

Arum acaule purpureum. *Br.* 332.

Arum minus esculentum. *Sl.* 1. p. 167.
t. 106. *f.* 2.

Blue Eddas. *Hughes.* 227.

Arrow-leaved Caladium. *Aiton.*

Purple Cocco and Taniers. *Browne.*

Nut Eddoes. Taniers.

seguinum. C. caulescens suberectum, foliis oblongis
cuspidatis, spadice spatha oblonga bre-
vior. *Willd. Sp. Pl.* 4. p. 490. *Hort.*

Kew. 5. p. 312. *Bot. Mag. t.* 2606. *h.*

Arum seguinum. *Jacq. Amer.* 239. *t.* 151.

Arum caule erecto geniculato. *Br.* 331.

Arum caule geniculato. *Sl.* 1. p. 168.

Dumb Cane. *Hughes.* 252. *Browne.*

Dumb Cane or Caladium. *Aiton.*

ARUM.

Piperitæ. *Lin.*—Aroideæ. *Juss. Brown.*

divarica- A. acaule foliis cordato-hastatis, spadice
*tum.** subulato, spatha ovato-lanceolata reflexa
longiore. *Willd. Sp. Pl.* 4. p. 482. *Hort.*
Kew. 5. p. 308. 24

* I will not be certain that this is an Arum, and not a Caladium, as I have not seen the fructification. It is an Arum in Willdenow.

A. acaule, foliis triangularibus. *Br.* 332.

No. 8.*

Divaricated Arum. *Aiton.*

Smaller Indian Kale. *Browne.*

esculentum.†

A. acaule, foliis peltatis cordatis, spadice spatha ovato-lanceolata brevior. 2

Caladium esculentum. *Willd. Sp. Pl.* 4. p. 489. *Hort. Kew.* 5. p. 311.

A. ægyptium, seu *Caladium sativum.*

Rumph. Amb. 5. p. 313. t. 109. t. 110.

f. 1.‡

* Browne considers his Arum, *No.* 8. as the plant described by Sloane, as Arum minus esculentum; and in this he has been followed by Willdenow. Nevertheless they are distinct. Browne describes his Arum as edible only as a green; Sloane says "the roots are boiled, and taste musked or sweet." Sloane's plant is the Nut Eddoe of Barbados, the roots of which are so conspicuous and valuable as food, that in this light they could not have escaped the notice of Browne. The Nut Eddoe, as already stated, is the Arum, *No.* 4. of Browne.

† This plant is certainly not a Caladium. The male flowers are not disposed at the top of the spadix, agreeably to the generic character of Caladium; but the spadix ends in a 1—3 inch long nudate process. It is therefore generically distinct from Mr. Brown's Caladium acre, which is undoubtedly a Caladium. That profound botanist had an opportunity of seeing the fructification, and represents it in these words, "spadice usque ad apicem obtusiusculum antherifero mucronulove brevissimo nudo."—*Prod.* 192.

‡ I do not exactly perceive how the Arum Colocasia is to be distinguished from the above Arum—*See Brown. Prod.* 192.

A. minus nymphaeæ foliis. *Sl.* 1. *p.* 167.

t. 106. *f.* 1.

Scratching Eddas. *Hughes.* 227.

Esculent Arum. *Miller.*

Scratch Cocco. *Edyes.* *Browne.*

I am inclined to think that these two species, as they stand in Willdenow, are in fact only one; to which, whether it be termed *Colocasia* or *esculentum*, apply both the plates of Rumphius, *t.* 109. and *t.* 110. *f.* 1. Much confusion has arisen from the relative terms, *minus*, *majus*, and *maximum*, employed by the older authors; and both the genus *Arum* and *Caladium* require much revision to ascertain the limits of each, and the appropriate synonyms of the species.

MONÆCIA MONADELPHIA.

CROTON.

Tricoccæ. *Lin.*—Euphorbiæ. *Juss.*—Euphorbiaceæ. *Brown.*

palustre. C. foliis ovato-lanceolatis plicatis serratis
scabris. *Willd. Sp. Pl.* 4. p. 535. *Hort.*
Kew. 5. p. 326. *Aubl.* 883. ☉
Pond-Weed. *Hughes.* 171.—240.
Marshy Croton. *Miller.*

balsami- C. foliis ovato-lanceolatis integerrimis
ferum. scabris subtus tomentosis basique bi-
glandulosis. *Willd. Sp. Pl.* 4. p. 548. ½
C. foliis lanceolatis acuminatis integris.
Jacq. Amer. 255. t. 162. f. 3.
C. fruticosum erectum. *Br.* 347.*

* I have removed the above synonym of Browne from Croton Eluteria, with which it does not agree. The "Croton Eluteria" of Willdenow, the "Mali folio arbor" of Sloane, is a tree; the "Croton fruticosum erectum" of Browne is an undershrub,

Balsam, or Sea-side Sage. *Hughes*. 143.

t. 10. *f.* 1.

Balsam Croton. *Miller*.

Sea-side Balsam. *Browne*.

JATROPHA.

Tricoccæ. *Lin.*—Euphorbiæ. *Juss.*—Euphorbiacæ. *Brown*.

gossipi- J. foliis cordatis quinquelobis serratis
folia. glanduloso-ciliatis, pilis ramosis glandu-
losis in foliorum axillis et petiolis. *Willd.*
Sp. Pl. 4. p. 557. *Hort. Kew.* 5. p. 329.
Aubl. 883. *Jacq. Ic.* 3. t. 623. *Bot.*
Reg. t. 746. h

J. humilior. *Br.* 348.

Ricinus minor staphisagriæ folio. *Sl.* 1.
p. 129. *t.* 84.

Belly-Ach. *Hughes*. 152. *t.* 11. *f.* 1.

Wild Cassada. *Long.* 2. p. 776.

Cotton-leaved Physic-Nut, or Wild
Casava. *Miller*.

and “grows only to the height of four or five feet.” The inflo-
rence of Croton Eluteria, “racemis compositis axillaribus,”
that of Croton fruticosum erectum, “spicis terminalibus.”

Curcas. J. foliis cordatis angulatis. *Willd. Sp. Pl.* 4. p. 560. *Hort. Kew.* 5. p. 330. *Aubl.* 884. h

J. assurgens ficus folio. *Br.* 348.

Ricinus ficus folio. *Sl.* 1. p. 127.

Physic-Nut Tree. *Hughes.* 115. *Browne.*

Angular-leaved Physic-Nut. *Miller.*

multifida. J. foliis palmatis undecimlobis lævibus: lobis pinnatifidis crenatis, stipulis setaceis multifidis. *Willd. Sp. Pl.* 4. p. 561. *Hort. Kew.* 5. p. 330. *Aubl.* 884. h

J. assurgens foliis digitatis. *Br.* 348.

Ricinus americanus tenuiter diviso folio. *Sl.* 1. p. 36.

French Physic-Nut. *Hughes.* 115. t. 6. *Browne.* *Miller.*

JANIPHA.

Tricoccæ (*Jatropha*). *Lin.*—Euphorbiæ. *Juss.*

Euphorbiaceæ. *Brown.*

Löflingii. J. foliis cordatis 5—partitis: laciniis acuminatis integerrimis: intermedia panduriformi, floribus racemosis. *Spreng. Syst.* 3. p. 77. h

Jatropha Janipha. Willd. *Sp. Pl.* 4. p. 562.

Jatropha carthaginensis. Jacq. *Amer.*
256. t. 162. f. 1.

Jatropha foliis palmatis, &c. carne nivea.
Br. 350.

Cassado Tree (*white sort*). *Hughes.* 150.

Sweet Cassada. *Long.* 2. p. 781.

Löflings Janipha. *Sweet.*

Roasting Cassava.

Manihot. *J. foliis palmatis 5—7—partitis glabris*
subtus glaucis: laciniis lanceolatis inte-
gerrimis, floribus racemosis. *Spreng.*
Syst. 3. p. 77. h

Jatropha Manihot. Willd. *Sp. Pl.* 4. p.
562. *Hort. Kew.* 5. p. 330. *Aubl.* 884.

Jatropha foliis palmatis, &c. carne sub-
latea. *Br.* 349.

Ricinus minor viticis obtuso folio. *Sl.* 1.
p. 130. t. 85.

Cassado Tree (*old sort*). *Hughes.* 150.

Cassada. *Long.* 2. p. 777.

Cassava. *Sweet.*

RICINUS.

Tricoccæ. *Lin.*—Euphorbiæ. *Juss.*—Euphorbiaceæ. *Brown.*

communis. *R.* caule herbaceo pruinoso, foliis peltato-palmatis: lobis lanceolatis dentatis, stigmatibus tribus apice bifidis, capsulis echinatis. *M—.* *Willd. Sp. Pl.* 4. p. 564.

Hort. Kew. 5. p. 331. *Aubl.* 884. ☉

R. fruticosus assurgens. *Br.* 350.

R. americanus fructu racemoso hispido.

Sl. 1. p. 126.

R. ruber. *Rumph. Amb.* 4. p. 97. t. 41.

Negro-Oil Bush (*red sort*). *Hughes.* 116.

Common Palma Christi, or Castor-Oil Plant. *Aiton.*

Oil-Nut Tree. *Browne.*

*viridis.** *R.* caule herbaceo pruinoso, foliis peltato-palmatis: lobis lanceolatis dentatis, stig-

* This species, although much like the former, is distinct. Hughes observes, “the oil extracted from the berries (*seeds*) of the red is less rank than that of the other.” That prepared from the seeds of the *R. viridis* is indeed more drastic, nauseating, and uncertain in its operation; and it is greatly to be desired that the oil employed in medicine—a most excellent article when genuine, and prepared as directed by the College of Physicians—should be obtained exclusively from the *R. communis*. It is

matibus sex simplicibus, capsulis echinatis. M—. *Willd. Sp. Pl.* 4. p. 564.
Hort. Kew. 5. p. 331. *Willd. Berol.*
t. 49. ☉

R. communis. *Bot. Mag. t.* 2209.

R. americanus major caule virescente.
Sl. 1. p. 127.

R. albus. *Rumph. Amb.* 4. p. 92.

Negro-Oil Bush (*white sort*). *Hughes.* 116.

Green Palma Christi. *Aiton.*

HIPPOMANE.

Tricoccæ. *Lin.*—Euphorbiæ. *Juss.*—Euphorbiacæ. *Brown.*

Mancinella. *H. Mancinella.* *Willd. Sp. Pl.* 4. p. 571.
Hort. Kew. 5. p. 333. *Aubl.* 885. *h*
H. Mancanilla. *Jacq. Amer.* 250. *t.* 159.

however fruitless to expect this, so long as the preparation is made abroad, and is in the hands of uninformed persons, who do not think it of importance to select the seeds from a particular species. It is, moreover, certain that the unprincipled do not hesitate to mix with the *Ricinus* seeds, those obtained from other oleaginous plants, such as species of the *Jatropha*, *Croton*, &c. which abound in warm climates. This will account for the disappointment experienced in the medicinal use of Castor-Oil, without resorting to constitutional idiosyncrasies.

H. arboreum lactescens. *Br.* 351.

Mancanilla Pyri facie. *Plum. Gen.* 50.

Juglandi affinis arbor. *Sl.* 2. *p.* 3. *t.* 159.

Manchaneel Tree. *Hughes.* 122.

Manchineel. *Long.* 2. *p.* 838.

Mancheneel Tree. *Miller.*

SAPIUM.

Tricoccæ (Hippomane). *Lin.*—*Euphorbiæ.* *Juss.*

Euphorbiacæ. *Brown.*

aucupato- *S. foliis oblongis acuminatis serrulatis,*
rium. *petiolis apice biglandulosis.* *Willd. Sp.*
Pl. 4. *p.* 572. *Hort. Kew.* 5. *p.* 333.
Jacq. Amer. 249. *t.* 158. h

S. arboreum. *Br.* 338.

Hippomane biglandulosa. *Aubl.* 885.

Mancanilla lauri foliis. *Plum. Gen.* 50.

Laurifolia venenata. *Sl.* 1. *p.* 39.

Poison Tree. *Hughes.* 144.

Gum Tree. *Long.* 2. *p.* 278. *Miller.*

PHYLLANTHUS.

Tricoccæ. *Lin.*—Euphorbiæ. *Juss.*—Euphorbiaceæ. *Brown.*

- nutans.* P. foliis elliptico-ovatis, racemo terminali nutante, caule ramoso. *Willd. Sp. Pl. 4. p. 577. Swtz. Fl. 1203.* $\frac{1}{2}$
 Tilia forte arbor racemosa. *Sl. 2. p. 19. t. 158. f. 3.*
 Box-Wood. *Hughes. 159?*
 Pendulous-flowered Phyllanthus. *Miller.*
 Rod-Wood.
- Niruri.* P. foliis pinnatis floriferis: foliolis ellipticis obtusis, pedunculis inferioribus subgeminis masculis; superioribus solitariis foemineis, petiolis cauleque erecto teretibus. *Willd. Sp. Pl. 4. p. 583. Hort. Kew. 5. p. 335.* ☉
 Herba mœroris alba. *Rumph. Amb. 6. p. 41. t. 17. f. 1.*
 Annual Phyllanthus. *Miller.*

HURA.

Tricoccæ. *Lin.*—Euphorbiæ. *Juss.*—Euphorbiaceæ. *Brown.*

crepitans. *H. crepitans.* *Willd. Sp. Pl. 4. p. 592.*

Hort. Kew. 5. p. 338. Aubl. 885. h

Hippomane arboreum, &c. foliis cordatis
crenatis. *Br. 350.*

Sand - Box Tree. *Hughes. 114. t. 5.*

Miller. Browne.

ARECA.

Palmæ. *Lin. Juss. Brown.*

oleracea. *A. frondibus pinnatis: foliolis linearibus
acutis, stipitibus spathis lanceolatis in-
curvatis spadicebusque ramosis lævibus,
fructibus oblongis. M.— Willd. Sp. Pl.*

*4. p. 596. Hort. Kew. 5. p. 340. Jacq.
Amer. 278. t. 170. h*

Palma caudice altissimo. *Br. 342.*

Palma pinnis inferne vaginantibus. *Br.
343.*

Palma altissima non spinosa. *Sl.* 2. p. 115. t. 215.

Cabbage Tree. *Hughes.* 107. t. 4. *Miller.*
Barbados Cabbage Tree. Jamaica Cabbage Tree. *Long.** 2. p. 744.

LAGENARIA.

Cucurbitaceæ. *De Cand.* (*Cucurbita*). *Lin.* *Juss.*

vulgaris. L. molliter pubescens moschata, caule scandente, cirris 3—4—fidis, foliis cordatis integriusculis piloso-subglaucescens, tibus basi biglandulosis, floribus monoi-cis stellatis patentissimis fasciculatis connectivo papillis oblongo-ovatis acutis

* Long justly states the Barbados and Jamaica Cabbage Trees to be only accidental varieties. Browne represents them as specifically distinct. Hughes, p. 111. mentions a small Cabbage Tree, which he thinks different from that generally growing in Barbados, and noticed at p. 107. of his history. "This," he says, "even at its greatest growth, is less than the Cabbage Tree already described, having likewise the green part near the top much less in proportion to its bulk." This I believe to be only the *Areca oleracea* impeded in its growth by sterility of soil, or by being crowded by other trees.

obsito, fructibus pubescentibus maturitate glabratis lævissimis, carne albida (eduli?) *De Cand. Prod.* 3. p. 299.* ☉

Cucurbita Lagenaria. *Willd. Sp. Pl.* 4. p. 606. *Hort. Kew.* 5. p. 343. *Aubl.* 887.

Cucurbita fructu pyriformi. *Br.* 354.

Cucurbita fructu maximo subrotundo.
Br. 354.

Small Gourd Vine. *Hughes.* 138? *Long.* 2. p. 801.

Large Gourd Vine. *Hughes.* 138. *Long.* 2. p. 801.

Bottle Gourd. *Sweet.*

CUCUMIS.

Cucurbitaceæ. *Lin. Juss. De Cand.*

Melo.† *C. caule humifuso scabro cirrifero, foliis rotundatis angulatis petiolatis, floribus masculis tubo calycis basi subventricosos*

* De Candolle enumerates four varieties depending on the difference of shape in the fruit; but as these pass into each other by an indefinite series I have not thought it proper to introduce them. I am not informed that the pulp is in any instance edible.

† De Candolle enumerates three principal varieties.

apice dilalato staminibus inclusis antheris connectivo brevioribus; floribus hermaphroditis antheris ut in masculinis, stigmatibus 3—4 breve bilobis, fructu ovato vel subgloboso 8—12 sulcato, carne saccharata flava vel alba. *De Cand.*

Prod. 3. p. 300. *Willd. Sp. Pl.* 4. p.

613. *Hort. Kew.* 5. p. 345. *Aubl.* 887. ☉

C. fructu oblongo sulcato. *Br.* 353.

Melo vulgaris. *Bauh. Pin.* 310.

Musk-Melon Vine. *Hughes.* 136.

Melon. *Long.* 2. p. 800.

Common or Musk-Melon. *Miller.*

sativus.

C. caule scabro cirrifero, foliis cordatis obscure quinquelobis petiolatis lobo terminali, floribus breve pedunculatis subternis majusculis, florum masculinorum tubo calycis tubuloso-campanulato: limbo patente deflexo, petalis acutiusculis, fructibus oblongis subtriquetris per maturitatem sublævibus sæpe nitidis, carpellis distinctis separabilibus. *De Cand. Prod.*

3. p. 300. *Willd. Sp. Pl.* 4. p. 615. *Hort.*

Kew. 5. p. 346. *Aubl.* 887. *Br.* 353.

Blackw. t. 4.

☉

C. sativus vulgaris. *Bauh. Pin.* 310.

Common Cucumber. *Miller.*

flexuosus. *C.* caule humifuso scabro flexuoso cirrifero, foliis cordato-ovatis sublobatis den-

ticulatis petiolatis, floribus fasciculatis in axillis foliorum, calyce valde piloso, fructibus elongatis cylindrico-clavatis sulcatis flexuosis replicatis albidis vel luteis.

De Cand. Prod. 3. p. 300. Willd. Sp.

Pl. 4. p. 615. Hort. Kew. 5. p. 346. ☉

Flexuose Snake Cucumber. *Sweet.*

Anguria. C. caule subfiliformi cirrifero, foliis palmato-sinuatis basi cordatis scabris, floribus subsolitariis magnitudine Bryoniæ dioicæ, fructibus globosis echinatis. *De*

Cand. Prod. 3. p. 301. Willd. Sp. Pl.

4. p. 612. Hort. Kew. 5. p. 345. Aubl. 887. ☉

C. subhirsutus minor. *Br. 353.*

C. Anguriæ folio latiore. *Sl. 1. p. 227.*

Wild Cucumber Vine. *Hughes. 186.*

Long. 2. p. 801.

Round Prickly-fruited Cucumber. *Miller.*

Citrullus. C. valde pilosus, caule humifuso cirrifero, foliis obtuse pinnatisectis subglaucescens, floribus solitariis unibracteatis: bractea oblonga, fructibus ellipticis oblongisve (*subglobosis D.C.*) glabris stellato-maculatis. *De Cand. Prod. 3. p. 301. ☉*

C. foliis multipartitis. *Br. 353.*

Cucurbita Citrullus. *Willd. Sp. Pl. 4. p.*

610. Hort. Kew. 5. p. 344. Aubl. 887.

Anguria prima Citrullus dicta. *Sl.* 1. p. 226.

Water Melons. *Hughes.* 136. *Long.* 2. p. 300. *Miller.*

Dudaim. *C. hispidus*, foliis inferioribus rotundatis; superioribus subquinelobis basi cordatis denticulatis, cirris simplicibus, petalis ovato-rotundatis, florum masculinorum calyce basi rotundato fauce dilatato connectivo antheris longiore; florum hermaphroditorum tubo calycis ovato piloso, stigmatibus 4—6, fructibus globosis læviusculis variegatis raro verrucosis, carne albida odoratissima sed insipida.

De Cand Prod. 3. p. 301. *Willd. Sp.*

Pl. 4. p. 613. *Hort. Kew.* 5. p. 345.

Bot. Repos. t. 548.

⊙

Apple-shaped Cucumber. *Miller.*

BRYONIA.

Cucurbitaceæ. *Lin. Juss. De Cand.*

racemosa. *B.* foliis inferioribus subpalmatis; superioribus trilobis indivisisque; laciniis ovatis,

floribus racemosis, pedunculis subsecundis, fructibus ovalibus. *De Cand. Prod.* 3. p. 308. *Willd. Sp. Pl.* 4. p. 623. *Swtz. Fl.* 1148. *Plum. Amer.* 83. t. 97. 4

MOMORDICA.

Cucurbitaceæ. *Lin. Juss. De Cand.*

Charantia. M. foliis septemlobo-palmatis dentatis subhirsutis, cirris pubescentibus, fructibus oblongis acuminatis angulatis tuberculatis croceis vel rubris, pulpa lutea molli, bractea cordata integerrima infra medium pedicelli, seminibus oblongis tuberculatis arilla rubro-sanguinea. *De Cand. Prod.* 3. p. 311. *Willd. Sp. Pl.* 4. p. 602. *Hort. Kew.* 5. p. 342. *Aubl.* 886. *Bot. Mag.* t. 2455. ☉

M. subhirsuta. *Br.* 353.

Hairy Momordica. *Sweet.*

Hairy Cerasee. *Browne.*

operculata. M. foliis quinquelobis dentatis, fructibus ellipticis angulato-tuberculatis rostratis

rostro deciduo operculatis. *De Cand.*
Prod. 3. p. 311. *Willd. Sp. Pl.* 4. p. 603.
Hort. Kew. 5. p. 342. *Aubl.* 886. ☉
 Rough-fruited Momordica. *Sweet.*

SECHIUM.

Cucurbitaceæ. *Juss. De Cand. (Sicyos.) Lin.*

edule. S. caule teretiusculo striato lævi, foliis
 cordato-angulatis subtus rugosiusculis,
 lobis basi conniventibus dentatis termi-
 nali longiori acuminatis, cirris 4—5—fidis,
 floribus masculis racemosis; foemineis so-
 litariis in eadem axilla, fructibus magnis
 obovatis 5—sulcatis apice gibbis piloso-
 echinatis. *De Cand. Prod.* 3. p. 313.
Willd. Sp. Pl. 4. p. 627. *Swtz. Fl.*
 1150. ☉
 S. foliis cordato-angulatis. *Br.* 355.
Sicyos edulis. Jacq. Amer. 258. t. 163.
Swtz. Prod. 116.
 Chocho Vine. *Long.* 2. p. 802. *Miller.*
 Christophine.

TRICHOSANTHES.

Cucurbitaceæ. *Lin. Juss. De Cand.*

anguina? T. caule pentagono, foliis cordatis trilobis repando-dentatis puberulis, cirris bifidis longissimis, floribus masculis racemosis: racemo longe pedunculato, lobis calycinis brevissimis subdentiformibus, lobis corollæ ovatis lacinioso-ciliatis, fructibus tereti-oblongis longe rostratis piloso-hispidulis rumpentibus, seminibus *De Cand. Prod. 3. p. 314.*

Willd. Sp. Pl. 4. p. 598. Hort. Kew. 5. p. 341. Bot. Mag. t. 722. ☉

Cucumis anguinus. *Willd. Sp. Pl. 4. p. 615?*

Sweet Gourd. *Hughes. 138.*

Slender Winding Gourd, or Sweet Gourd.
Long. 2. p. 801.

Snake Gourd. *Aiton.*

CUCURBITA.

Cucurbitaceæ. *Lin. Juss. De Cand.*

- Pepo.* C. foliis cordato-obtusis subquinelobis denticulatis, calycibus in collum infra limbum desinentibus, fructibus subrotundis oblongisve lævibus. *De Cand. Prod. 3. p. 317. Willd. Sp. Pl. 4. p. 609. Hort. Kew. 5. p. 343. Aubl. 887. ☉*
Pepo maximus indicus. Sl. 1. p. 226.
Pumkins. Hughes. 137.
Pumpion, or Pumpkin Gourd. Miller.
- Melopepo.* C. foliis cordato-obtusis subquinelobis denticulatis, cirris in foliis valde imperfectis sæpe transformatis, calyce hemisphærico-campanulato brevi, fauce valde dilatata, fructibus depressis carpellis ultra faucem calycis prominentibus irregularibus, carne sicca spongiosa albida, loculis 4—5, seminibus *De Cand. Prod. 3. p. 317. Willd. Sp. Pl. 4. p. 610. Hort. Kew. 5. p. 344. ☉*
Melopepo Clypieformis. Bauh. Pin. 312.
Squashes. Hughes. 137.
Squash-Gourd. Miller.

DICECIA MONANDRIA

PANDANUS.

Palmæ. *Juss.*—Pandaneæ. *Brown.*

odoratis- P. foliis dorso margineque spinoso-den-
simus. tatis, fructu globoso solitario. *Willd.*
Sp. Pl. 4. p. 645. *Hort. Kew.* 5. p. 351.
Roxb. Cor. 1. p. 65. t. 94. 95, 96. ½
Sweet-scented Pandanus, or Skrew Pine.
Miller.

DICECIA DIANDRIA.

CECROPIA.

Scabridæ. *Lin.*—Urticæ. *Juss.*—Urticæ. *De Cand.*

- peltata.* C. foliis novemlobis : lobis oblongis obtusiusculis supra hispido-scabris subtus tomentoso-albis. *Willd. Sp. Pl.* 4. p. 651.
Hort. Kew. 5. p. 352. *Aubl.* 894. $\frac{1}{2}$
Corlotapalus ramis excavatis. *Br.* 111.
Yaruma de Oviedo. *Sl.* 1. p. 137. t. 88.
f. 2.
Ficus surinamensis. *Pluk. Alm.* 146.
Phyt. t. 243. f. 5.
Snake-Wood. *Hughes.* 223. t. 10. f. 2.
Trumpet-Tree, or Snake-Wood, *Long.*
2. p. 757. *Miller.*

DICECIA TRIANDRIA.

PHŒNIX.

Palmæ. *Lin. Juss. Brown.*

*dactyli-
fera.*

P. frondibus pinnatis inermibus: foliolis complicatis lineari-lanceolatis strictis.

Willd. Sp. Pl. 4. p. 730. Hort. Kew. 5. p. 369. Aubl. 974.

h

Palma dactylifera major. Sl. 2. p. 111.

Palma subcinerea. Br. 344.

Date Tree. Hughes. 102. Long. 2. p. 742.

Date-Palm Tree. Miller.

DICECIA TETRANDRIA.

VISCUM.

Aggregatæ *Lin.*—Caprifolia. *Juss.*

Loranthæ. *Juss. Ann. du Mus.*

verticilla- V. caule verticillato, foliis ovatis triner-
tum. viis obtusis. *Willd. Sp. Pl.* 4. p. 741. ½

V. parasiticum ramulis verticillatis. *Br.*
356.

V. ramulis et foliis longis. *Sl.* 2. p. 93.
t. 201. *f.* 2.

Mistletoe. *Hughes.* 156.

Whorled Mistletoe. *Miller.*

KŒLERA?*

* * * * *

serrata. K. foliis alternis ovato-lanceolatis acuminatis serratis basi inæqualibus, pedunculis unifloris axillaribus confertis. M.—
Silver-Wood Tree.

* I cannot identify the Silver Wood Tree with any known species, nor are my notes sufficient to enable me to ascertain the natural order to which it should be referred. It would seem to belong to the genus *Kœlera*; but differs from *K. laurifolia*, in having an unarmed trunk, and a serrated leaf. It is a large and elegant tree, growing in the neighbourhood of Joes River.

MAS.	FEM.
Calyx. 4-partitus.	Calyx et Cor. maris.
Cor. o.	Pist. germen ovatum superum, stylus nullus, stigma capitatum.
Stam. 4.* calice longiora.	Nect. annulare basi germen cingens.
Nect. capitato-depressum.	Pericarp. Capsula monosperma, stigmate persistente uno latere coronata.

* The male flowers have sometimes five, more rarely six stamens, the segments of the Calyx varying in like manner.

DICECIA PENTANDRIA.

ZANTHOXYLUM.

Hederaceæ. *Lin.*—Terebinthaceæ. *Juss.*—Rutaceæ. *De Cand.*

Ochroxylum. Z. trunco aculeato, foliis ovatis punctato-pellucidis, floribus 5--andris 3--gynis. *De Cand. Prod.* 1. p. 725. $\frac{1}{2}$
Yellow Hercules. *Hughes.* 146.

tragodes. Z. aculeatum, foliis impari pinnatis: foliolis cuneiformibus emarginatis, petiolo alato in articulos foliola fere referentes solubili. *De Cand. Prod.* 1. p. 725. $\frac{1}{2}$
Fagara tragodes. *Willd. Sp. Pl.* 1. p. 667. *Hort. Kew.* 1. p. 263. *Jacq. Amer.* 21. t. 14.

Fingrigo, or Savine Tree. *Hughes.* 197.
Prickly-leaved Zanthoxylum. *Sweet.*

Pterota. Z. aculeatum, foliis impari-pinnatis: foliolis obovatis subcrenatis, petiolo anguste

marginato, aculeis duobus stipularibus uncinatis. *De Cand. Prod.* 1. p. 725. $\frac{1}{2}$
 Pterota subspinoso. *Br.* 146. t. 5. f. 1.
 Fagara Pterota. *Willd. Sp. Pl.* 1. p. 666.
Hort. Kew. 1. p. 263.

Lauro affinis Jasmini folio. *Sl.* 2. p. 25.
 t. 162. f. 1.

Lentiscus-leaved Zanthoxylum. *Sweet.*
 Saven Tree, or Bastard Iron-Wood.
Browne.

emarginatum. Z. inerme, foliis pinnatis 2—3—jugis :
 foliolis ovatis emarginatis villosis, racemis terminalibus. *De Cand. Prod.* 1. p.
 726. *Willd. Sp. Pl.* 4. p. 753. *Hort.*
Kew. 5. p. 382. *Swtz. Fl.* 572. $\frac{1}{2}$

Fagara emarginata. *Swtz. Prod.* 33.
 Lauro affinis Terebinthi folio alato.
Sl. 2. p. 24. t. 168. f. 4.

Notch-leaved Zanthoxylum. *Sweet.*

Clava Z. aculeatum, foliis pinnatis 4—jugis :
Herculis. foliolis ovatis repando-dentatis basi inæqualibus sessilibus pellucido-punctatis glaberrimis, paniculis terminalibus. *De*
Cand. Prod. 1. p. 727. *Willd. Sp. Pl.*
 4. p. 754. *Hort. Kew.* 5. p. 382. $\frac{1}{2}$

Z. foliis oblongo-ovatis pinnatis. *Br.* 189.
 Prickle Yellow Wood. *Hughes.* 167.
 Clubbed Zanthoxylum. *Sweet.*

Prickly Yellow Wood, or Yellow Hercules. *Browne*.

Yellow Hercules.

fraxineum. *Z.* foliis impari pinnatis 4—5—jugis: foliolis ovatis obsolete serrulatis basi æqualibus, petiolis teretibus inermibus, aculeis stipularibus, umbellis axillaribus.

De Cand. Prod. 1. p. 726. *Willd. Sp.*

Pl. 4. p. 757. *Hort. Kew.* 5. p. 383. $\frac{1}{2}$

Z. Clava Herculis. β . *Lin. Sp. Pl.* 1455.

White Hercules. *Hughes.* 164.

Prickly White Wood. *Long.* 2. p. 828.

Common *Zanthoxylum*. *Sweet.*

DIOECIA HEXANDRIA

DIOSCOREA.

Sarmentaceæ. *Lin.*—Asparagi. *Juss.*—Dioscoreæ. *Brown.*

aculeata. D. foliis alternis subrotundo-cordatis
acuminatis septemnerviis, caule aculeato
bulbifero. *Willd. Sp. Pl.* 4. p. 792. *Hort.*
Kew. 5. p. 392. *Aubl.* 903. 4
D. foliis cordatis, caule tereti aculeato.
Br. 359.*

Prickly Yam Vine. *Hughes.* 227.

Prickly-stem'd Dioscorea. *Aiton.*

* There is indeed, as Dr. Martyn observes, "much confusion in the synonyms" to the different species of Dioscorea. I cannot however agree with him, that the Dioscorea No. 1. of Browne, with an aculeated stem, is identical with *D. sativa*. Browne is in error in considering his Dioscorea No. 1. as the *Volubilis nigra radice tuberosa compressa* of Sloane, termed by him Negro-Country Yam, in reference to the part of the

- alata.* D. foliis oppositis ovatis cordato-sagittatis cuspidatis septemnerviis, caule alato bulbifero. *Willd. Sp. Pl.* 4. p. 792. *Hort. Kew.* 5. p. 392. *Aubl.* 903. 4
D. foliis cordatis, caule alato. *Br.* 359.
Volubilis nigra, radice alba aut purpurea. *Sl.* 1. p. 139.
Winged-stalked Dioscorea. *Aiton.*
Red Yam.
- bulbifera?* D. foliis alternis cordatis subrotundo-ovatis acuminatis subnovemnerviis, caule lævi bulbifero. *Willd. Sp. Pl.* 4. p. 793.
Hort. Kew. 5. p. 392. *Aubl.* 903. 4
Rhizophora, &c. radice rotunda. *Herm. Par.* 217. t. 217.
Black-Bryony-leaved Dioscorea. *Aiton.*
Grenada Yam.
- sativa.** D. foliis alternis cordatis subrotundo-ovatis cuspidatis subnovemnerviis lobis

world from which it had been introduced into Jamaica; and which I think is plainly the *D. sativa*. Long, by the term Negro-Yam, *Dioscorea radice tuberosa lutea*, evidently means the prickly Yam; and Browne's No. 1. is without doubt the same plant, the *Dioscorea aculeata*.

* This is certainly not, as supposed by Willdenow, the *Dioscorea*, No. 3. or Wild Yam of Browne, which he says is "not put to any use in Jamaica." I have therefore omitted the synonym from Browne, referred to in the *Species Plantarum*.

baseos approximatis, capsulis obovatis,
 caule lævi tereti. *Willd. Sp. Pl.* 4. p.
 795. *Hort. Kew.* 5. p. 392. *Aubl.*
 904. 2

Volubilis nigra, radice tuberosa com-
 pressa. *Sl.* 1. p. 140.

Yams. *Hughes.* 226.*

Cultivated Dioscorea, or Yam. *Aiton.*

ELAIS.

Palmæ. Lin. Juss. Brown.

guineensis. E. frondibus pinnatis, stipitibus dentato-
 spinosis divergentibus, denticulis su-
 premis recurvatis. *Willd. Sp. Pl.* 4. p.
 799. *Hort. Kew.* 5. p. 393. *Jacq. Amer.*
 280. t. 172. *Mart. Palm.* t. 54—56. 2
 Palma foliorum pediculis spinosis. *Sl.*
 2. p. 113. t. 214.†

* Several kinds of Yam are cultivated in Barbados. I have not had an opportunity of informing myself how far they are distinct species or only varieties.

† Willdenow, by an oversight, has made this Elais and the Cocos (*Acrocomia*) fusiformis both identical with the Palma tota

Palm Oil Tree. *Hughes.* 111.

Oily Palm Tree. *Aiton.*

occiden- E. frondibus pinnatis: foliolis vaginatis
talıs. stipitibus caudiceque inermi. *Willd.*

Sp. Pl. 4. p. 799. *Swtz. Fl.* 619. ½

Cordage Palm.

MAURITIA.

Palmæ. Lin. Juss. Brown.

vinifera. M. cadice elato stricto inermi, frondibus
flabelliformi-pinnatifidis concoloribus, pe-
tiolis semiteretibus canaliculatis, squa-
mis baccarum ovatarum subæqualiter
rhombeis. *Mart. Palm.* 42. t. 38, 39.

f. 1, 2. ½

Straight-stem'd Mauritiam. *Sweet.*

spinosa major of Sloane. I should have had no hesitation in adopting the above synonym even without the authority of Swartz, from which however I derive additional confidence. I have felt obliged to exclude the synonym from Browne admitted in the species *Plantarum*, for his *Palma caudice aculeatissimo*, and Sloane's *Palma spinosa minor*, refer to the small Macaw Tree—a prickly Palm growing in Jamaica, which I have not seen in Barbados.

DICECIA DECANDRIA.

CARICA.

Tricoccæ. *Lin.*—Cucurbitaceis aff. *Juss.*

Papaya. C. foliis palmatis septemlobis : lobo inter-
medio sinuato : laciniis oblongis acutis,
floribus masculis corymbosis. *Willd. Sp.*
Pl. 4. *p.* 814. *Hort. Kew.* 5. *p.* 399.
Aubl. 909. h

C. fronde comosa. *Br.* 360.

Papaya major. *Sl.* 2. *p.* 164--166.

Papaja, mas. et foem. *Rumph. Amb.* 1.
p. 145. *t.* 50, 51.

Papaw Tree. *Hughes.* 181. *t.* 14, 15.
Long. 2. *p.* 802.

Common Papaw Tree. *Miller.*

DICÆCIA MONADELPHIA.

JUNIPERUS.

Coniferæ. *Lin. Juss.*

- barbaden-* J. foliis omnibus quadrifariam imbrica-
sis. tis; junioribus ovatis; senioribus acutis.
Willd. Sp. Pl. 4. p. 851. Pluk. Alm.
201. Phyt. t. 197. f. 4. h
Barbadoes Juniper. *Miller.*
- bermu-* J. foliis inferioribus ternis; superioribus
diana. binis decurrentibus subulatis patulis acu-
tis. *Willd. Sp. Pl. 4. p. 851. Hort. Kew.*
5. p. 413. h
Bermudas Juniper. *Miller.*
Bermudas Cedar.*

* I cannot be certain whether the plant named by Browne and Long Bermudas Cedar be the *J. bermudiana*, or *J. barbadensis*. Browne refers to both these Linnæan species. Sloane questions whether it be not the *Juniperus barbadensis* of Pluke-

MYRISTICA.

Lauri. *Juss.*—Myristicæ. *Brown*,

moschata. *M.* foliis oblongis acuminatis glabris,
venis simplicibus, fructu solitario glabro.

Willd. Sp. Pl. 4. p. 869. *Hort. Kew.*
5. p. 417. h

M. officinalis. *Bot. Mag. t.* 2756, 2757.

Nux moschata fructu rotundo. *Pluk.*

Alm. 265. *Phyt. t.* 209. *Blackw. t.* 353.

Aromatic, or True Nutmeg Tree. *Aiton.*

net; but Long seems quite confident that it is *J. bermudiana*. Not having seen the tree indigenous in Jamaica, I cannot attempt to decide this question; I have therefore omitted all reference to Browne, Sloane, and Long. From Sloane's plate I should rather take the Juniper of Jamaica to be identical with the *J. barbadensis*, a tree by no means common at this time in the island from which it takes its specific name.

POLYGAMIA MONŒCIA.

MUSA.

Scitamineæ. *Lin.*—Musæ. *Juss.*—Musaceæ. *De Cand.*

- paradi-* M. spadice nutante, floribus masculis
siaca. persistentibus, spathis oblongis acutis,
fructu cylindraceo trigono. *Willd. Sp.*
Pl. 4. p. 893. Hort. Kew. 5. p. 424.
Aubl. 930. 4
- M. spadice nutante, fructu triquetro.
Br. 363.
- M. caudice viridi. *Sl. 2. p. 141.*
Plantain Tree. *Hughes. 182.* Long. 2.*
p. 782. Miller.
- sapientum.* M. spadice nutante, floribus masculis

* The Wild Plantain Tree, of which Hughes states, "that though it blossoms it never bears any fruit," is not specifically distinct from this.

deciduis, spathis oblongis obtusiusculis,
fructu elliptico. *Willd. Sp. Pl.* 4. p. 894.

Hort. Kew. 5. p. 424. *Aubl.* 930. 2

M. spadice nutante fructu brevior. *Br.*
363.

M. caudice maculato. *Sl.* 2. p. 147.

α . fructu flavicante.

β . fructu purpureo.

Banana Tree. *Hughes.* 184. t. 16. *Long.*

2. p. 783. *Miller.*

TERMINALIA.

Elæagni. *Juss.*—*Combretaceæ.* *De Cand. Brown.*

Catappa. *T. foliis obovatis basi attenuatis subtus molliter pubescentibus, glandulis minimis subtus in basi folii ad latus nervii medii.* *De Cand. Prod.* 3. p. 11. *Willd.*

Sp. Pl. 4. p. 976. *Hort. Kew.* 5. p. 441.

Jacq. Ic. 1. p. 19. t. 197. 1/2

Catappa. *Rumph. Amb.* 1. p. 174. t. 68.

Amygdalus indica. *Raj. Hist.* 1521.

Large-leaved Terminalia. *Sweet.*

Almond Tree.

CLUSIA.

Dubii ord. *Lin.*—Guttiferæ. *Juss. De Cand.*

flava.

C. floribus polygamis, calyce polyphyllo, corolla 4—petala, staminibus numerosis brevibus, stigmatibus circiter duodecim, foliis obovatis obtusis aveniis carnosis.

M.— *De Cand. Prod.* 1. p. 559. *Willd. Sp. Pl.* 4. p. 977. *Hort. Kew.* 5. p. 444. *Jacq. Amer.* 272. *t.* 167. *Bot. Repos.* *t.* 223. ½

C. arborea, foliis crassis. *Br.* 236.

Terebinthus folio singulari. *Sl.* 2. p. 91. *t.* 200. *f.* 1.

Balsam Tree. *Hughes.* 157. *Long.* 2. p. 723.

Yellow-flowered Balsam. *Miller.*

GOUANIA.

Rhamnis aff. *Juss.*—Rhamneæ. *De Cand.*

- domingensis.* G. foliis ovatis acuminatis hinc inde serratis glabriusculis. *De Cand. Prod.* 2. p. 39. *Willd. Sp. Pl.* 4. p. 999. *Hort. Kew.* 5. p. 450. *Aubl.* 934. h
 G. glabra. *Jacq. Amer.* 264. t. 179. f. 40.
 Lupulus sylvestris americana. *Pluk. Alm.* 229. *Phyt. t.* 201. f. 4.
 Chaw-Stick. *Miller.*

MIMOSA.

Lomentaceæ. *Lin.*—Leguminosæ. *Juss. De Cand.*

- pudica.* M. caule herbaceo aculeato plus minus petiolis pedunculisque piloso-hispidis, foliis subdigitato-pinnatis: pinnis 4—multijugis: foliolis linearibus. *De Cand. Prod.* 2. p. 426. *Willd. Sp. Pl.* 4. p. 1031. *Hort. Kew.* 5. p. 455. *Aubl.* 944. *Bot. Repos. t.* 544. ⊙

Sensitive Plant. *Hughes*. 209.

Humble Plant. *Miller*.

INGA.

Lomentaceæ (*Mimosa*). *Lin.*—Leguminosæ. *Juss. De Cand.*

Bourgoni. *I.* foliolis 2—3—jugis ovatis nitidis glabris, glandulis petioli ad apicem articulationum subalati distinctis, spicis axillaribus subquaternis brevibus. *De Cand.*

Prod. 2. p. 434. ½

I. marginata. *Willd. Sp. Pl.* 4. p. 1015.

Hort. Kew. 5. p. 453.

Mimosa Bourgoni. *Aubl.* 941. t. 358.

Mimosa fagifolia. *Jacq. Amer.* 264. t. 164.

Spanish-Oak. *Hughes*. 196.

Beech-leaved Inga. Spanish-Oak. *Aiton*.

Unguis Cati. *I.* spinis stipularibus rectis, foliolis subrotundo ellipticis subdimidiatis emarginatis membranaceis glabris, glandula in dichotomia petioli glabri et inter foliola, florum capitulis globosis in racemum terminalem dispositis, legumine torto.

De Cand. Prod. 2. p. 436. *Willd. Sp.*

Pl. 4. p. 1006. *Hort. Kew.* 5. p. 452. ½

Mimosa Unguis Cati. *Aubl.* 944.

Mimosa fruticosa. *Br.* 252.

Mimosa spinis geminis. *Plum. Ic.* 2. t. 4.

Acacia arborea major spinosa. *Sl.* 2.
p. 56.

Moabite, *alias* Mangrove - Bead Tree.
Hughes. 193.

Four-leaved Inga. *Aiton.*

Bread and Cheese.

purpurea. *I. inermis*, foliis conjugato-pinnatis: pin-
narum foliolis 3—7—jugis oblongis ob-
tusis, capitulis pedunculatis axillaribus.

De Cand. Prod. 2. p. 439. *Willd. Sp.*

Pl. 4. p. 1021. *Hort. Kew.* 5. p. 454.

Bot. Reg. t. 129. h

Mimosa purpurea. *Bot. Repos.* t. 372.

Purple Inga, or Soldier-Wood. *Aiton.*

DESMANTHUS.

Lomentaceæ (*Mimosa*). *Lin.*—Leguminosæ. *Juss. De Cand.*

*virgatus.** *D.* caule suffruticoso erecto angulato,
foliis pinnis 3—4—jugis, glandula infra
jugum inferius, pedunculis nudis, spicis
paucifloris capitatis, floribus decandris

* *Desmanthus strictus*, *D. C.* the habitat of which is said to
be in the Caribee Islands, seems to be only a variety of this.

legumine anguste lineari 25—30—spermo. *De Cand. Prod. 2. p. 445. Willd. Sp. Pl. 4. p. 1047. Hort. Kew. 5. p. 458. Bot. Mag. t. 2454.* h

Mimosa virgata. Jacq. Vind. 1. p. 34. t. 80.

Largest Sensitive Plant. *Hughes. 198.*

Long-twiggèd Desmanthus. *Aiton.*

ACACIA.

Lomentaceæ (*Mimosa*). *Lin.*—Leguminosæ. *Juss. De Cand.*

horrida. A. spinis geminis foliis subæquantibus, ramis foliisque glabris, pinnis 2—3 jugis: foliolis 9—12—jugis oblongis obtusis costa biglandulosa, capitulis pedunculatis axillaribus sæpius aggregatis. *De Cand. Prod. 2. p. 460. Willd. Sp. Pl. 4. p. 1082.* h

Nemnem, or Toothache Tree. *Hughes. 200.*

Long-spined Acacia. *Sweet.*

tortuosa. A. spinis geminis connatis, ramis petiolisque pubescentibus, pinnis 2—6—jugis: foliolis 12—16—jugis, glandula infra in-

fimum par et sæpe inter pinnas extremas, capitulis solitariis axillaribus, legumine compresso crassiusculo pubescente subcorniculato. *De Cand. Prod.* 2. p. 461.

Willd. Sp. Pl. 4. p. 1083. h

Akasee, or Sweet Briar. *Hughes.* 200. t. 10. f. 2.

Twisted Acacia. *Sweet.*

latisiliqua. A. inermis glabra, pinnis 5—jugis: foliolis 10—jugis ellipticis obtusis, stipulis bracteiformibus dimidiato-cordatis, capitulis pedunculatis aggregatis in paniculam terminalem subdispositis. *De Cand.*

Prod. 2. p. 467. *Willd. Sp. Pl.* 4. p. 1067. *Hort. Kew.* 5. p. 467. h

Mimosa latisiliqua. *Aubl.* 945.

Broad-podded Acacia. *Sweet.*

frondosa? A. inermis, pinnis 6—jugis: foliolis 14—jugis lanceolatis distantibus subtus ramis petiolisque pubescentibus, glandula inter pinnas inferiores, capitulis axillaribus pedunculatis solitariis. *De*

Cand. Prod. 2. p. 468. *Willd. Sp. Pl.* 4. p. 1076. h

Frondose Acacia. *Sweet.*

RHAPIS.

Palmae. *Lin. Juss. Brown.*

*flabelli-
formis.*

R. frondibus palmatis plicatis: plicis
marginibusque aculeato – denticulatis.

Willd. Sp. Pl. 4. p. 1093. Hort. Kew.

5. p. 473. Bot. Mag. t. 1371. h

Creeping-rooted Rhapsis, or Ground-
Ratan. *Aiton.*

Dwarf Palm.

POLYGAMIA DICECIA.

BURSERA.

Dubii ord. *Lin.*—Terebinthaceæ. *Juss. De Cand.*

*gummi-
fera.*

B. foliis deciduis sæpius impari-pinnatis: foliolis ovatis acutis membranaceis, racemis axillaribus. De Cand. Prod. 2. p. 78.

Willd. Sp. Pl. 4. p. 1119. Hort. Kew. 5. p. 481.

h

Burseria gummifera. Jacq. 94. t. 65.

Terebinthus foliis cordato-ovatis pinnatis. Br. 345.

Terebinthus major. Sl. 2. p. 89. t. 199. f. 2.

Betula arbor americana. Pluk. Alm. 67. Phyt. t. 151. f. 1.

Birch-Gum Tree. Hughes. 144.

Jamaica Birch Tree. Miller.

Birch, or Turpentine Tree. Browne.

FICUS.

Scabridæ. *Lin.*—Urticeæ. *Juss.*

- Carica.* F. foliis cordatis tri-quinquelobisve repando-dentatis: lobis obtusis supra scabris subtus pubescentibus, receptaculis pyriformibus glabris. *Willd. Sp. Pl.* 4. p. 1131. *Hort. Kew.* 5. p. 483, *Aubl.* 951. *Blackw. t.* 125. h

Common Fig Tree. *Sweet.*

- martinicensis.* F. foliis oblongo-lanceolatis integerrimis apice attenuatis acutis basi rotundatis supra albo-punctatis. *Willd. Sp. Pl.* 4. p. 1137. *Hort. Kew.* 5. p. 485. h

F. laurifolia. *Roem. Syst.* 1. p. 511.F. indica. *Aubl.* 951.F. indica maxima folio oblongo. *Sl.* 2. p. 140. *t.* 223.F. arborea assurgens. *Br.* 110?Bearded Fig Tree. *Hughes.* 175.

- nitida.* F. foliis ellipticis acutis glabris basi attenuatis subtrinerviis parallele venosis, receptaculis globosis involucratis in apicibus romulorum sessilibus. M.—*Roem. Syst.* 1. p. 506. h

F. nitida. *Exot. Fl. t.* 111?*

F. pertusa. *Willd. Sp. Pl.* 4. *p.* 1144.
Hort. Kew. 5. *p.* 487. *Spreng. Syst.* 3.
p. 781.†

* Mr. Sweet considers the *Ficus nitida* of the Exotic Flora as a distinct species, under the title *F. Hookeri*. I had the satisfaction of conversing with him on this subject; and I readily admit that the *F. nitida*, represented in the Exotic Flora, must, from the figure and definition there given, be considered as a species distinct from the *F. nitida* of Roemer. Yet it is said, that the plant from which Dr. Hooker's plate was taken was brought from the West-Indies; and in some respects it is exceedingly like the Garden Mangrove of Barbados. The dissimilarity is principally in the form of the leaf, which seems to have varied very much even on the specimen from which the drawing was taken. The colour of the leaf is also much paler, and the disposition to be 3-nerved is not exhibited. These points of difference may in a great degree be accounted for from the original of Dr. Hooker's plate having been a hot-house exotic; and I am therefore induced to pause before admitting the *F. Hookeri* as a distinct species.

† The *F. pertusa* of Willdenow, Aiton, and Sprengel, "*receptaculis globosis sessilibus*," cannot be the same species as the *F. pertusa* of Roemer, "*receptaculis globosis pedunculatis*." The *pertusa* of Willdenow seems to be the same as the *nitida* of Roemer, and both identical with the *Itti Arealou* of Rheede and the Garden Mangrove of Hughes. This species of *Ficus*, although very common in Barbados, is seldom or never of accidental growth, being propagated principally from layers or cuttings of the stem, very rarely from the seed, and planted in the neighbourhood of dwelling-houses on account of its great beauty and evergreen

Itti Arealou. *Hort. Mal.* 3. p. 69. t. 55.

Garden Mangrove. *Hughes.* 193.

Glossy-leaved Fig Tree. *Sweet.*

Mangrove Evergreen Tree.

elastica. F. foliis ellipticis acutiusculis coriaceis
splendidis margine cartilagineis paral-
lele venosis. *Spreng. Syst.* 3. p. 781. ½
Indian Rubber Fig-Tree. *Sweet.*

shade. It was probably at an early period introduced into the island, as has more recently been the Bamboo, the Bread-Fruit Tree, the Shaddock, the Hibiscus Rosa-sinensis, &c. &c. now very generally met with. Its native soil is the East-Indies and China.

CRYPTOGAMIA STACHYOPTERIDES.

LYCOPodium.

Musci. *Lin. Juss.*—Lycopodiaceæ. *Kunth.*

Lycopodineæ. *Brown.*

cernuum. L. foliis sparsis curvatis, caule ramosissimo, spicis cernuis, squamis appressis membranaceis serrato-ciliatis. *Willd.*

Sp. Pl. 5. p. 30. 4

Muscus maximus coridis folio. *Plum.*

Fil. 144. t. 165. A.

CRYPTOGAMIA FILICES.

ACROSTICUM.

Filices. *Lin. Juss. Brown.*

*longifo-
lium.*

A. frondibus oblongo-lanceolatis marginatis basi apiceque attenuatis utrinque glabris, stipitibus frondis fertilis lanceolatae apice spiraliter tortae paleaceis.

Willd. Sp. Pl. 5, p. 105. 2

Lingua cervina rigida et glabra. *Plum. Fil. 118. t. 135.*

*Calome-
lanos.*

A. frondibus bipinnatis: pinnulis oblongis subtus albo-farinosus: inferioribus subpinnatifidis basi sursum auritis: superioribus serratis: summis confluentibus. *Willd. Sp. Pl. 5. p. 123. Hort. Kew.*

5. p. 501. 2

α . Filix non ramosa major, caule nigro.

Sl. 1. *p.* 92. *t.* 30. *f.* 2.

Filix albissimo pulvere conspersa.

Plum. Fil. 30. *t.* 40.

Adiantum nigro simile. *Plum. Amer.*

30. *t.* 44.

β . Filix non ramosa minor, caule nigro.

Sl. 1. *p.* 3. *t.* 7. *f.* 1.

Filix non ramosa minima, caule nigro.

Sl. 1. *p.* 92. *t.* 53. *f.* 1.

Maiden-Hair. *Hughes.* 237.

Mealy Acrosticum. *Aiton.*

POLYPODIUM.

Filices, *Lin.* *Juss.* *Brown.*

Phylli-
tidis.

P. frondibus lanceolatis marginatis acutis
basi attenuatis glabris, soris biserialibus.

Willd. Sp. Pl. 5. *p.* 157. *Hort. Kew.* 5.

p. 504. 4

P. foliis linguæ cervinæ. *Plum. Amer.*

26. *t.* 38.

Lingua cervina longis angustis et undu-
latis foliis. *Plum. Fil.* 114. *t.* 130.

Phyllitis arboribus innascens. *Sl.* 1. *p.* 72.

Hart's-tongue Polypody. *Aiton.*

- aureum.* P. frondibus profunde pinnatifidis glaucescentibus: laciniis lanceolatis acuminatis integerrimis: terminali elongata, soris laciniarum inferiorum sparsis: laciniæ terminalis solitariis. *Willd. Sp. Pl.* 5. p. 169. *Hort. Kew.* 5. p. 504. 2
- P. majus aureum. *Plum. Amer.* 25. t. 35. *Plum. Fil.* 59. t. 76.
- Agnus Scithycus. *Hughes.* 235.
- Golden Polypody. *Aiton.*

ASPIDIUM.

Filices. *Lin. Juss. Brown.*

- heracleifolium.* A. frondibus pinnatis: pinnis oblongo-lanceolatis trilobis cordatis subrepandis acuminatis, soris biserialibus. *Willd. Sp. Pl.* 5. p. 217. 2
- Hemionitis altera pentaphylla. *Plum. Fil.* 126. t. 147.
- macrophyllum.* A. frondibus pinnatis: pinnis oblongo-lanceolatis acuminatis: terminali tripartita serrata: lateralibus subsinuato-repandis: infimis bi-trilobisve subsinuatis, soris biserialibus. *Willd. Sp. Pl.* 5. p. 217. 2

Hemionitis maxima linguæ cervinæ
affinis. *Plum. Fil.* 124. *t.* 145.

patens.

A. frondibus pinnatis: pinnis pinnati-
fidis: laciniis lanceolatis acutis: infimis
pinnarum ultimarum longioribus pinna-
tifido-incisis, venis subtus hirtis. *Willd.*

Sp. Pl. 5. *p.* 244. *Hort. Kew.* 5. *p.* 508. 4

Polypodium patens. *Swtz. Fl.* 1673.

Filix non ramosa minor. *Sl.* 1. *p.* 91. *t.*
52. *f.* 1.

Fern-like Plant. *Hughes.* 237.

Downy Shield-Fern. *Aiton.*

PTERIS.

Filices. *Lin. Juss. Brown.*

lanceolata. P. frondibus simplicibus lanceolatis utrin-
que acutis subintegerrimis apice fructi-
ficantibus. *Willd. Sp. Pl.* 5. *p.* 356. 4

Lingua cervina foliis acutis et ad oras
summitatum pulverulentibus. *Plum. Fil.*

116. *t.* 132. *Plum. Amer.* 28. *t.* 40.

ADIANTUM.

Filices. *Lin. Juss. Brown.*

- pumillum.* A. frondibus pinnatis: pinnis subrotundis cuneatis obtusis serrulatis: terminali rhombeo-obovata, soris margine superiori subsolitariis. *Willd. Sp. Pl. 5. p. 431. Swtz. Fl. 1703.* 2
- A. minimum erectum. *Br. 87.*
- Black Maiden-Hair. *Hughes. 237.*
- Dwarf Maiden-Hair. *Miller.*

CYATHEA.

Filices. *Lin. Juss. Brown.*

- arborea.* C. frondibus bipinnatis: pinnulis lanceolatis serratis acutiusculis: summis confluentibus, stipiti lævi, caudice arboreo. *Willd. Sp. Pl. 5. p. 491. Hort. Kew. 5. p. 529.* 2

Polypodium arboreum maximum. *Br.*
104.

Filix arborescens pinnulis dentatis.
Plum. Fil. 1. t. 1, 2. Plum. Amer. 1.
t. 1, 2.

Mountain-Fern. *Hughes. 237.*

Tree-Cyathea, or Cup-Fern. *Aiton.*

APPENDIX I.

PLANTS MENTIONED IN THE NATURAL HISTORY
OF BARBADOS, BUT NOT IDENTIFIED WITH
ANY OF THE SPECIES ENUMERATED
IN THE PRECEDING PAGES.*

Anchovie-Apple Tree	<i>Hughes.</i>	131.
Antigua Rose Tree	<i>id.</i>	220.
Bumbo-Bush	<i>id.</i>	164.
Cow Wyth	<i>id.</i>	242.
Crab Tree	<i>id.</i>	203.
Fire-Burn Weed	<i>id.</i>	162.
Forster's Plant	<i>id.</i>	206.

* Several of the plants, inserted in this Appendix, have been excluded from the Flora, in consequence of the author not having had an opportunity of examining them in their flowering and fruiting states, so as satisfactorily to ascertain their generic and specific characters. Others, for the reasons explained in the preface, he cannot identify with any of the species noticed in the Flora. The Olive and the Lucern are probably at this time no where to be found in the island.

Goat's Rue	<i>Hughes.</i>	219.
Grapple Wood	<i>id.</i>	217.
Honey Wyth	<i>id.</i>	242.
Jallop Vine	<i>id.</i>	204.
Lignum Rorum	<i>id.</i>	158.
Lucern	<i>id.</i>	202.
Olive Tree	<i>id.</i>	177.
Patience	<i>id.</i>	219.
Pigeon Wood	<i>id.</i>	157.
Prickly Hoop, or White Thorn	<i>id.</i>	153.
Pudding Wyth	<i>id.</i>	242.
Red Willow	<i>id.</i>	196.
Soap Berry Bush, or Fire Burn Leaf	<i>id.</i>	149.
Sober's Plant	<i>id.</i>	218.
Sweet Vine	<i>id.</i>	243.
Turky Weed	<i>id.</i>	219.
White Plum Tree	<i>id.</i>	179.
Wild Clove	<i>id.</i>	221.
Wild Pennyroyal	<i>id.</i>	170.
Wild Potatoe Vine	<i>id.</i>	198.
Wild Yam Vine	<i>id.</i>	242.
Wood Sorrel	<i>id.</i>	203.

APPENDIX II.

SOME OF THE PRINCIPAL VERNACULAR NAMES
OCCURRING IN THE FOREGOING PAGES,
WITH THEIR SYSTEMATIC
SYNONYMS.

Ægyptian Bean	<i>Nelumbium jamaicense.</i>
African Marigold	<i>Tagetes erecta.</i>
African Plum	<i>Cordia Myxa.</i>
Agnus Scythicus	<i>Polypodium aureum.</i>
Aligator Apple	<i>Anona palustris.</i>
Almond Tree	<i>Terminalia Catappa.</i>
Aloes Plant	<i>Aloe vulgaris.</i>
American Jasmine	<i>Ipomœa Quamoclit.</i>
American Torch	<i>Cereus monoclonos.</i>
Annual Sun Flower	<i>Helianthus annuus.</i>
Antigua Balsam	<i>Justicia pectoralis.</i>
Arabian Jasmine	<i>Jasminum Sambac.</i>
Archangel	<i>Eupatorium odoratum.</i>
Arnotta	<i>Bixa Orellana.</i>
Arrow-Root	<i>Maranta arundinacea.</i>
Austrian Rose	<i>Rosa bracteata.</i>
Avigato Pear Tree	<i>Persea gratissima.</i>

Bachelor Button	Gomphræna globosa.
Bachelor's Pears	Solanum mammosum.
Bachelor's Pears	Solanum xantocarpum.
Bahama Red Wood	Ceanothus colubrinus.
Balsam, or Sea-side Sage,	Croton balsamiferum.
Balsam Tree	Clusia flava.
Bambu, or Bamboo	Bambusa arundinacea.
Banana Tree	Musa sapientum.
Barbados Cedar	Cedrela odorata.
Barbados Lilac	Melia Azedarach.
Barbados Pride, or	
Flower Fence	Poinciana pulcherrima.
Barbados Trumpet	
Flower	Bignonia Unguis.
Basket Wyth	Tournefortia bicolor.
Bastard Briony	Cissus sicyoides.
Bastard Bully Tree	Bumelia nigra.
Bastard Feverfew	Parthenium Hystero-
	phorus.
Bastard Flower-Fence ...	Adenanthera pavonina.
Bastard Ipecacuahna ...	Asclepias curassavica.
Bastard Lignum Vitæ ...	Badiera diversifolia.
Bastard Locust, or Forest	
Tree	Clethra tinifolia.
Bastard Mammee, or	
Santa Maria	Calophyllum Calaba.
Bastard Ockro	Malachra capitata.
Bay-Berry Tree	Myrcia acris.
Bay-Grape Tree	Coccoloba uvifera.

Bean Tree, or Throve	} Erythrina Coralloden-
Tuesday	
Bearded-Fig Tree	Ficus martinicensis.
Beef-Wood Tree	Pisonia nigricans.
Beet	Beta vulgaris
Bell Pepper	Capsicum annuum.
Bell Pepper	Capsicum grossum.
Belly-Ach	Jatropha gossipifolia.
Bermudas Cedar	Juniperus bermudiana.
Birch Gum Tree	Bursera gummifera.
Birch or Turpentine	
Tree	Bursera gummifera.
Bird Pepper	Capsicum baccatum.
Bird Pepper	Capsicum conoides.
Bitter Wood	Quassia excelsa.
Black Basket Wyth	Rivina octandra.
Black Bead Tree	Ceanothus colubrinus.
Black Berry Hunters ...	Eugenia virgultosa.
Black Cherry	Eugenia ligustrina.
Black Cherry	Cerasus occidentalis.
Black-eyed Peas	Dolichos sphærospermus.
Black Maiden Hair	Adiantum pumillum.
Black Nicker	Sapindus saponaria.
Black Sage Bush	Varronia currasavica.
Black Thorn	Pisonia aculeata.
Black Willow	Capparis torulosa.
Black Wood	Eugenia triflora.
Bloodworth	Moræa plicata.
Blue Eddas	Caladium sagittifolium.

Blue Shake Shake	<i>Crotalaria verucosa</i> .
Blue Vine	<i>Clitoria ternatea</i> .
Bombast Mahoe	<i>Ochroma Lagopus</i> .
Bonnet or Bonny Pepper.	<i>Capsicum tetragonum</i> .
Bonnet or Bonny Pepper.	<i>Capsicum angulosum</i> .
Box Wood	<i>Phyllanthus nutans</i> .
Branched Calalue	<i>Solanum nigrum</i> .
Bread and Cheese	<i>Inga Unguis Cati</i> .
Bread and Cheese, or	
Sucking Bottle	<i>Paullinia Cururu</i> .
Bread Fruit Tree	<i>Artocarpus incisa</i> . β .
Bread Nut Tree	<i>Artocarpus incisa</i> . α .
Broad-leaved Rose Bay.	<i>Nerium coronarium</i> .
Broad-leaved Sour Grass.	<i>Paspalum conjugatum</i> .
Broad Pond Duck-Weed.	<i>Nelumbium jamaicense</i> .
Broom Weed	<i>Corchorus siliquosus</i> .
Broom Weed	<i>Sida jamaicensis</i> .
Broom Weed	<i>Sida angustifolia</i> .
Brown Jolly, &c.	<i>Solanum Melongena</i> .
Brown Wool Cotton	<i>Gossypium barbadense</i> .
Buff-Coat	<i>Waltheria americana</i> .
Buff Coat Tree	<i>Guettarda scabra</i> .
Bully-Berry Tree	<i>Bumelia nigra</i> .
Bully Tree	<i>Bumelia nigra</i> .
Buona Vista. Bonny-vis.	<i>Lablab nankinicus</i> .
Buona Vista. Bonny-vis.	<i>Lablab leucocarpus</i> .
Buona Vista. Bonny-vis.	<i>Lablab perennans</i> .
Buona Vista. Bonny-vis.	<i>Lablab vulgaris</i> . β .
Buona Vista. Bonny-vis.	<i>Lablab cultratus</i> .

Buona Vista. Bonny-vis.	Lablab vulgaris. γ .
Burgamot Tree	Citrus Limetta.
Bur Grass	Cenchrus echinatus.
Bur Grass	Cenchrus tribuloides.
Butterfly Oncidium	Oncidium Papilio.
Button Sage	Lantana involucrata.
Cabbage	Brassica oleracea.
Cabbage Tree ..,	Areca oleracea.
Calaba Tree, or Santa	
Maria	Calophyllum Calaba.
Calabash Tree	Crescentia Cujete.
Calalue	Basella cordifolia.
Calavances, or Red Bean.	Dolichos sinensis.
Candle Wood	Chiococca racemosa.
Canella, or Winter's	
Bark	Canella alba.
Cape Jasmine	Gardenia fragrans.
Carnation	Dianthus Caryophyllus. β .
Cashew Tree	Anacardium occidentale.
Cassava	Janipha Manihot.
Cassia Fistula Tree	Cassia Fistula.
Castor-Oil Plant	Ricinus communis.
Caterpillars	Amaranthus viridis.
Caterpillars	Amaranthus polygo- noides.
Cedar	Cedrela odorata.
Celery	Apium graveolens.
Cephalic Vine	Convolvulus speciosus.

Changeable Rose Hibis-

cus	<i>Hibiscus mutabilis</i> .
Chaw-Stick	<i>Gouania domingensis</i> .
Checquered Grape Tree.	<i>Coccoloba nivea</i> .
Chereese, or Barbados	
Cherry	<i>Malpighia puniceifolia</i> .
Cherry Pepper	<i>Capsicum cerasiforme</i> .
Chigery Bush	<i>Tournefortia volubilis</i> .
Chigery Grape Tree	<i>Coccoloba nivea</i> .
China Aster	<i>Aster chinensis</i> .
China Orange Tree	<i>Citrus Aurantium</i> .
Chinaise Rose	<i>Hibiscus mutabilis</i> .
Chinese Cock's-Comb ...	<i>Celosia coccinea</i> .
Chinese Pink	<i>Dianthus chinensis</i> .
Chinese Rose Hibiscus,	<i>Hibiscus Rosa sinensis</i> .
Choco Vine	<i>Sechium edule</i> .
Chocolate-Nut Tree	<i>Theobroma Cacao</i> .
Christmas Bush	<i>Eupatorium odoratum</i> .
Christophine	<i>Sechium edule</i> .
Cinnamon Tree	<i>Laurus Cinnamomum</i> .
Circassian Bead Tree ...	<i>Adenãthera pavonina</i> .
Citron Tree	<i>Citrus medica</i> .
Clay Peas	<i>Dolichos sinensis</i> .
Climbing Snowberry	
Tree	<i>Chiococca racemosa</i> .
Clove Pink	<i>Dianthus caryophyllus</i> .
Clove Tree	<i>Caryophyllus aromaticus</i> .
Cocco Plum	<i>Chrysobalanus Icaco</i> .
Cochineel Shrub	<i>Opuntia coccinellifera</i> .

Cocoa Nut Tree	<i>Cocos nucifera</i> .
Cocoa Plum	<i>Chrysobalanus Icaco</i> .
Cocoa Tree	<i>Theobroma Cacao</i> .
Coffee Tree	<i>Coffea arabica</i> .
Common Cucumber	<i>Cucumis sativus</i> .
Common Dill	<i>Anethum graveolens</i> .
Common Everbearing	
Rose	<i>Rosa indica</i> .
Common Fig Tree	<i>Ficus Carica</i> .
Common Garlick	<i>Allium sativum</i> .
Common Grape	<i>Vitis vinifera</i> . α .
Common Kidney Bean.	<i>Phaseolus vulgaris</i> .
Common Leek	<i>Allium Porrum</i> .
Common Pea	<i>Pisum sativum</i> .
Common Soap Tree	<i>Sapindus saponaria</i> .
Common Sow Thistle ...	<i>Sonchus oleraceus</i> .
Common Sweet Basil ...	<i>Ocimum Basilicum</i> .
Common Vine	<i>Vitis vinifera</i> .
Common Wormwood ...	<i>Artemisia Absinthium</i> .
Conch-Apple. Conch-Nut.	<i>Passiflora maliformis</i> .
Coot Weed	<i>Crotalaria lotifolia</i> .
Coral, or Red Bean Tree.	<i>Erythrina Coralloden-</i> <i>dron</i> .
Cork Wood	<i>Anona palustris</i> .
Corn Tree	<i>Adansonia digitata</i> .
Corynth, or Currant	
Grape	<i>Vitis vinifera</i> . β .
Cowhage Cherry	<i>Malpighia urens</i> .
Cow-Itch Vine	<i>Mucuna urens</i> .

Crab Bush, or Sea-side	} Heliotropium gnapha-
Laurel	
	lodes.
Crab's-Eye Vine	Abrus precatorius.
Creeping Cowhage	Tragia volubilis.
Creeping-rooted Rhaps,	
or Ground Ratan	Rhapis flabelliformis.
Crested Amaranth, or	
Cock's-Comb	Celosia cristata.
Crimson Everbearing	
Rose	Rosa semperflorens.
Cuckold's Increase	Dolichos unguiculatus.
Cuckold's Increase	Dolichos sesquipedalis.
Custard Apple	Anona reticulata.
Damacen or Callimato	
Tree	Chrysophyllum glabrum.
Damask Rose	Rosa damascena.
Damson Plum	Chrysophyllum monopy-
	renum.
Dark Chinese Rose	Rosa semperflorens.
Dart Wood	Ixora fasciculata.
Devil's Grass	Cynodon dactylon.
Date Tree	Phoenix dactylifera.
David's Root	Chiococca racemosa.
Dialthæa	Waltheria americana.
Dog's Grass	Poa ciliaris.
Dog Wood	Gardenia Randia.
Dove Weed	Euphorbia maculata.
Down Tree	Ochroma Lagopus.

- Down Vine *Sarcostemma Swartzianum*.
- Downy Mountain Ebony. *Bauhinia tomentosa*.
- Dragon's Blood *Dracæna ferrea*.
- Drooping-leaved Adam's
Needle *Yucca Draconis*.
- Dwarf Hibiscus *Hibiscus phoeniceus*.
- Dwarf Maiden Hair *Adiantum pumillum*.
- Dwarf Palm *Rhapis flabelliformis*.
- Dwarf Pomegranate
Tree *Punica nana*.
- Dwrah, or Amadwrah ... *Cyperus ligularis*.
- Dumb Cane *Caladium seguinum*.
- Dunk Tree, or Man-
gostine *Ziziphus Jujuba*.
- Dutch Grass *Cynosurus indicus*.
- Dyer's Mulberry, or
Fustic Wood *Morus tinctoria*.
- East India Mango Tree. *Mangifera indica*.
- Eatable Wild Pea *Dolichos luteus*.
- Egg Plant *Solanum Melongena*.
- English Plantain *Plantago media*.
- English Wormwood *Artemisia Absinthium*.
- Ethiopian Sour Gourd,
or Monkey's Bread ... *Adansonia digitata*.
- Fan Palm *Thrinax parviflora*.
- Fat Pork *Chrysobalanus Icaco*.

Fennel, or Finkle	<i>Anethum Fœniculum.</i>
Fern-like Plant	<i>Aspidium patens.</i>
Fiddle-Wood Tree	<i>Cytharexylum cinereum.</i>
Fingrigo, or Cockspur ...	<i>Pisonia aculeata.</i>
Fingrigo, or Savine Tree.	<i>Zanthoxylum tragodes.</i>
Fit-Weed	<i>Eryngium foetidum.</i>
Fire Burn Bush	<i>Triopteris jamaicensis.</i>
Five Sprig Tree	<i>Myrodia turbinata.</i>
Flexuose Snake Cucum- ber	<i>Cucumis flexuosus.</i>
Flower Fence, or Spanish Carnation	<i>Poinciana pulcherrima.</i>
Forbidden Fruit Tree ...	<i>Citrus buxifolia ?</i>
Forest Bark, or Bastard Locust	<i>Clethra tinifolia.</i>
French Bean	<i>Phaseolus vulgaris.</i>
French Bean	<i>Phaseolus multiflorus.</i>
French Guava	<i>Cassia alata.</i>
French Guava	<i>Psidium pyrifera.</i>
French Lavender	<i>Lavandula Stœchas.</i>
French Marigold	<i>Tagetes patula.</i>
French Physic-Nut Tree.	<i>Jatropha multifida.</i>
French Rose Tree	<i>Hibiscus mutabilis.</i>
French Willow	<i>Cerbera Thevetia.</i>
French Wormwood	<i>Artemisia gallica.</i>
Fustic Tree	<i>Morus tinctoria.</i>
Galimeta Wood	<i>Bumelia salicifolia.</i>
Galivaughn Peas	<i>Dolichos barbādensis.</i>

Gall Tree	<i>Quassia excelsa.</i>
Garden Balsam	<i>Balsamina hortensis.</i>
Garden Balsam	<i>Justicia pectoralis.</i>
Garden Basil	<i>Ocimum Basilicum.</i>
Garden Carrot	<i>Daucus Carota.</i>
Garden Hydrangea, or Chinese Guelder Rose.	<i>Hydrangea hortensis.</i>
Garden Lettuce	<i>Lactuca sativa.</i>
Garden Mangrove	<i>Ficus nitida.</i>
Garden Marigold	<i>Calendula officinalis.</i>
Garlick Pear Tree	<i>Cratæva gynandra.</i>
Germinating-leaf Plant.	<i>Bryophyllum Calycinum.</i>
Ginger	<i>Zingiber officinalis.</i>
Ginger Grass	<i>Panicum latifolium.</i>
Golden Apple	<i>Spondias dulcis.</i>
Golden Orange Tree	<i>Citrus vulgaris.</i>
Gooseberry Shrub	<i>Pereskia aculeata.</i>
Gourd Vine	<i>Lagenaria vulgaris.</i>
Granada Yam	<i>Dioscorea bulbifera?</i>
Granadilla Vine	<i>Passiflora quadrangularis</i>
Grape-Fruit Tree	<i>Citrus Hystrix?</i>
Great Corn, or Indian Maize	<i>Zea Mays.</i>
Great Macaw Tree	<i>Acrocomia fusiformis.</i>
Great Water Lily	<i>Nelumbium jamaicense.</i>
Grey Nicker	<i>Guilandina Bonduc. β.</i>
Ground Nut	<i>Arachis hypogœa.</i>
Guava Tree	<i>Psidium pomiferum.</i>
Gully Plum Tree	<i>Spondias lutea.</i>

Gully-Root	<i>Petiveria alliacea.</i>
Gum Elemi Tree	<i>Amyris toxifera.</i>
Gum Tree	<i>Sapium aucupatorium.</i>
Gumma Bush	<i>Solanum nigrum.</i>
Guinea Corn	<i>Sorghum vulgare.</i>
Guinea Corn	<i>Sorghum cernuum.</i>
Guinea Corn	<i>Sorghum bicolor.</i>
Guinea Corn	<i>Sorghum saccharatum.</i>
Guinea Grass	<i>Panicum polygamum.</i>
Hairy Cerasee	<i>Momordica Charantia.</i>
Halifax Peas	<i>Dolichos sesquipedalis.</i>
Hog Plum Tree	<i>Spondias lutea.</i>
Hog Slip. Hog Vine ...	<i>Convolvulus umbellatus.</i>
Hog Weed	<i>Boerhaavia diffusa.</i>
Holly-Hock	<i>Argemone mexicana.</i>
Holly-Hock	<i>Althæa rosea.</i>
Holy Thorn, or Royal	
Cashaw	<i>Parkinsonia aculeata.</i>
Honey-Berry	<i>Melicocca bijuga.</i>
Honey-Suckle	<i>Passiflora laurifolia.</i>
Hoop Wyth	<i>Rivina octandra.</i>
Hop Shrub	<i>Dodonæa viscosa.</i>
Hop Weed	<i>Bystropogon suaveolens.</i>
Hop Weed	<i>Salvia occidentalis.</i>
Horse Bean	<i>Canavalia ensiformis.</i>
Horse-Eye Bean Vine ...	<i>Mucuna urens.</i>
Horse Nicker	<i>Guilandina Bonduc. β.</i>
Horse Radish	<i>Cochlearia Armoracia.</i>

Horse Vine	<i>Teramnus uncinatus.</i>
Jaca Tree	<i>Artocarpus integrifolia.</i>
Jack Fruit Tree	<i>Artocarpus integrifolia.</i>
Jack-in-a-Box	<i>Hernandia sonora.</i>
Jallop, or Four o'Clock	
Flower	<i>Mirabilis Jalapa.</i>
Jamaica Birch Tree	<i>Bursera gummifera.</i>
Jamaica Plum Tree	<i>Spondias purpurea.</i>
Jasmine Tree	<i>Plumeria rubra.</i>
Jerusalem Thorn	<i>Parkinsonia aculeata.</i>
Indian Cale, or Seven	
Year's Cabbage	<i>Brassica oleracea. γ.</i>
Indian Corn	<i>Zea Mays.</i>
Indian Creeper. Indian	
Pink	<i>Ipomœa Quamoclit.</i>
Indian Kale	<i>Caladium nymphæifo-</i> <i>lium.</i>
Indian Oak, or Teak	
Wood	<i>Tectona grandis.</i>
Indian Root	<i>Asclepias curassavica.</i>
Indian Rubber	<i>Ficus elastica.</i>
Indian Rubber Tree	<i>Euphorbia atropurpurea.</i>
Indigo Berry	<i>Gardenia Randia.</i>
Indigo Weed	<i>Indigofera Anil.</i>
Ink Vine	<i>Passiflora suberosa.</i>
Job's Tears	<i>Coix Lacryma.</i>
Irish Potatoe	<i>Solanum tuberosum.</i>
Iron Grass	<i>Spermacoce tenuoir.</i>

Iron Wood *Siderodendrum triflorum*.
 Iron Vine *Desmodium spirale*.
 Italian Senna *Cassia obtusifolia*.
 Junction Vine *Aristolochia odoratissima*

King of Flowers *Lagerstroemia indica*.

Lady of the Night *Cestrum nocturnum*.
 Large Duck Weed *Nymphæa odorata*.
 Large Lime Tree *Citrus medica*. β .
 Large Wild Barren Pine. *Bromelia Pinguin*.
 Largest Sensative Plant. *Desmanthus virgatus*.
 Least Wild Pea Vine *Rhynchosia minima*.
 Leather Coat Tree *Coccoloba pubescens*.
 Lemon Grass *Cymbopogon Schoenanthus*.

Lesser Shaddock Tree ... *Citrus Decumana*.
 Lignum Vitæ *Guaiacum officinale*.
 Lily *Amaryllis equestris*.
 Lily *Pancratium fragrans*.
 Lily *Epidendrum ciliare*.
 Lime Tree *Citrus Limonum*.
 Lima Bean *Phaseolus perennis*.
 Limonia *Triphasia trifoliata*.^{*}
 Loblolly Tree *Varronia alba*.
 Locus-Berry Tree *Byrsonima coriacea*.

* This genus, inserted by mistake in the class Decandria, properly belongs to the class Hexandria, order monogynia.

Locust Tree	<i>Hymenæa Courbaril.</i>
Loggerhead Weed	<i>Spigelia Anthelmia.</i>
Logwood	<i>Hæmatoxylon campe-</i> <i>chianum.</i>
Long-leaved Mistletoe ...	<i>Pothos lanceolata.</i>
Long Pepper	<i>Capsicum longum.</i>
Love-in-a-Mist	<i>Passiflora foetida.</i>
Macartney Rose	<i>Rosa bracteata.</i>
Madagascar Periwinkle.	<i>Vinca rosea.</i>
Macaw Tree	<i>Acrocomia fusiformis.</i>
Mahogany Tree	<i>Swietenia Mahogoni.</i>
Maiden Hair	<i>Acrosticum Calomelanos.</i>
Malabar Nut	<i>Justicia Adhatoda.</i>
Mammee Sapota	<i>Achras mammosa.</i>
Mammee Tree	<i>Mammea americana.</i>
Manchineel Tree	<i>Hippomane Mancinella.</i>
Mangel-Wurzel	<i>Beta vulgaris. δ.</i>
Mango Tree	<i>Mangifera indica.</i>
Mangostan, or Mangos-	
teen	<i>Garcinia Mangostana.</i>
Many Roots	<i>Ruellia tuberosa.</i>
Many Seed	<i>Jussiaea octonervia.</i>
Marvel of Peru	<i>Mirabilis Jalapa.</i>
Mastick Tree	<i>Sideroxylon Masticho-</i> <i>dendron.</i>
May Grass	<i>Panicum latifolium.</i>
May Pole	<i>Agave americana.</i>
Melon	<i>Cucumis Melo.</i>

Menow Weed	Ruellia tuberosa.
Milk Weed	Euphorbia obliterata.
Milk Weed	Euphorbia glabrata.
Milk Weed	Euphorbia pilulifera.
Mistletoe	Viscum verticillatum.
Mignonette	Reseda odorata.
Moabite <i>alias</i> Mangrove	
Bead Tree	Inga Unguis Cati.
Monkey Apple Tree	Anona palustris.
Monkey's Bread	Adansonia digitata.
Monkey's Hand	Piper peltatum.
Monkey's Peas	Dolichos luteus.
Montabba <i>alias</i> Trubba.	Solanum Melongena.
Morass Weed	Ceratophyllum demersum.
Mountain Fern	Cyathæa arborea.
Musk Bush, or Wild	
Ockro	Hibiscus Abelmoschus.
Musk Melon	Cucumis Melo.
Musk Ockro	Hibiscus Abelmoschus.
Musk Rose	Rosa moschata.
Muskito Bush	Bystropogon suaveolens.
Myrtle Lime	Triphasia trifoliata.
Narrow-leaved Sour	} Andropogon angustifolius.
Grass	
Negro-Oil Bush	Ricinus communis.
Negro-Oil Bush	Ricinus viridis.
Negro Pepper	Capsicum conoides.

Nem Nem, or Toothach

Tree	<i>Acacia horrida</i> .
Nettles	<i>Urtica æstuans</i> .
Nettles	<i>Urtica grandifolia</i> .
Nettle Hibiscus	<i>Hibiscus pruriens</i> .
Nettle Sida	<i>Sida urens</i> .
Night-blowing Convol- vulus	<i>Ipomœa bona-nox</i> .
Night Shade	<i>Datura Strommonium</i> .
Noyeau Vine	<i>Convolvulus dissectus</i> .
Nut Eddoe. Taniers	<i>Caladium sagittifolium</i> .
Nut Grass	<i>Cyperus esculentus</i> .
Nutmeg Tree	<i>Myristica moschata</i> .

Ockra. Ockro *Hibiscus esculentus*.

Oil-Nut Tree *Ricinus communis*.

Old Maid, or Cayenne

Jasmine *Vinca rosea*.

Old Woman's Bitter *Citharexylum cinereum*.

Oleander, or Sweet-

scented Rose Bay *Nerium odorum*.

Olive Mangrove *Avicennia tomentosa*.

Olive Pepper *Capsicum cerasiforme*.

Otaheite Apple *Spondias dulcis*.

Otaheite Gooseberry *Cicca disticha*.

Ox-Eye Bean *Mucuna urens*.

Pagister Grass *Oplismenus hirtellus*.

Palm Oil Tree *Elais guineensis*.

Parsley	<i>Apium Petroselinon.</i>
Pawpaw Ockroe	<i>Corchorus æstuans.</i>
Pawpaw Tree	<i>Carica Papaya.</i>
Pen-Gwyn	<i>Bromelia Karatas.</i>
Pepper-Mint	<i>Mentha piperita.</i>
Pennyroyal	<i>Mentha Pulegium.</i>
Peruvian Cotton	<i>Gossypium peruvianum.</i>
Peruvian Marigold	<i>Tagetes tenuifolia.</i>
Physic Nut Tree	<i>Jatropha Curcas.</i>
Pigeon Pea Tree	<i>Cajanus flavus.</i>
Pimento, Jamaica Pepper,	
• or Allspice	<i>Eugenia Pimenta.</i>
Pimploes	<i>Opuntia Tuna.</i>
Pindals, or Ground Nuts.	<i>Arachis hypogœa.</i>
Pine-Apple	<i>Bromelia Ananas.</i>
Pinguin	<i>Bromelia Pinguin.</i>
Plantain Tree	<i>Musa paradisiaca.</i>
Poison Tree	<i>Sapium aucupatorium.</i>
Poison Wyth	<i>Cissus sicyoides.</i>
Pollard, or White Wool	
Cotton	<i>Gossypium herbaceum.</i>
Pomegranate Tree	<i>Punica Granatum.</i>
Pond Grass, or Canker	
Weed	<i>Commelina communis.</i>
Pond Grass, or Canker	
Weed	<i>Commelina erecta.</i>
Pond Weed	<i>Croton palustre.</i>
Pond Weed	<i>Alisma cordifolia.</i>
Poplar, or Poppy Tree.	<i>Thespesia populnea.</i>

Pops	Physalis angulata.
Pop Vine	Physalis barbadensis.
Potatoes	Convolvulus Batatas.
Poverty Weed	Desmodium repens.
Prickle Yellow Wood ...	Zanthoxylum Clava Her- culis.
Prickly Argemone, or	
Poppy	Argemone mexicana.
Prickly Pear Vine	Cereus trigonus.
Prickly Yam Vine	Dioscorea aculeata.
Prickly Yellow Wood, } or Yellow Hercules. }	Zanthoxylum Clava Her- culis.
Pumkins	Cucurbita Pepo.
Pumpion, or Pumpkin	
Gourd	Cucurbita Pepo.
Pumple Nose Tree	Citrus Decumana ?
Purple Cocco, and	
Taniers	Caladium sagittifolium.
Purple Eye Bright	Spermacoce hirta.
Purple Inga, or Soldier	
Wood	Inga purpurea.
Purslane	Portulaca oleracea.
Pye-Crust	Jacquinia armillaris.
Queen of Flowers	Lagerstroemia Regina.
Rabbit Vine	Teramnus uncinatus.
Rabbit Weed	Sonchus agrestis.

Rata Pepper. Rata Ockro	
<i>alias</i> Hoho	Corchorus hirtus.
Rattle Bush, or Shake	
Shake	Crotalaria incana.
Red Cherry Tree	Malpighia glabra.
Red Dialthæa	Reidleia nodiflora.
Red Flag or Plush Grass.	Andropogon angustifolius.
Red Mangrove Tree	Rhizophora Mangle.
Red Wood	Coccoloba barbadensis.
Red Yam	Dioscorea alata.
Reed	Panicum arundinaceum.
Reed	Donax arundinaceus.
Rice Grass	Panicum molle.
Right Wyth, or Cat's-	
Claw Wyth	Bignonia Unguis.
Ring Worm Bush	Cassia alata.
Roasting Cassava	Janipha Löflingii.
Roasting Eddoes	Caladium Macrorhizon.
Rocambole	Allium Scorodoprasum.
Rock Balsam	Piper obtusifolium.
Rock Balsam	Piper acuminatum.
Rock Bush	Piper aduncum.
Rock Sage	Lantana involucrata.
Rod Wood	Coccoloba barbadensis.
Rod Wood	Eugenia virgultosa.
Rope Mangrove	Hibiscus arboreus.
Rose of Sharon	Hibiscus Rosa-sinensis.

Raucou	Bixa Orellana.
Rouncivall Peas	Vigna glabra.
Rush	Scirpus mutatus.
Rush	Scirpus geniculatus
St. Helena Lemon Tree.	Citrus medica.
St. John's Bush	Psychotria nervosa.
St. Vincent Lilac	Solanum Seaforthianum.
Sambo	Gynandropsis penta- phylla.
Sandbox Tree	Hura crepitans.
Santa Maria Leaf	Piper umbellatum.
Sappadilla Tree	Achras sapota. x
Savanna Grass	Paspalum distichum.
Scarlet Runner Kidney	
Bean	Phaseolus multiflorus.
Saven Tree, or Bastard	
Iron Wood	Zanthoxylum Pterota.
Scallion	Allium Ascalonium.
Scotch Grass	Panicum pilosum.
Scotch Grass	Oplismenus hirtellus.
Scratch Cocco. Eddyes.	Arum esculentum.
Scratch Wyth	Cissus sicyoides.
Scratching Eddas	Arum esculentum.
Screw-Pine	Pandanus odoratissimus.
Sea Island Cotton	Gossypium hirsutum.
Seaside Balsam	Croton balsamiferum.
Seaside Grape	Coccoloba uvifera.
Seaside Lavender	Heliotropium gnapha- lodes.

Seaside Laurel	<i>Xylophylla falcata.</i>
Seaside Laurel	<i>Heliotropium gnaphalodes.</i>
Seaside Milk Weed	<i>Euphorbia glabrata.</i>
Seaside Samphire	<i>Sesuvium Portulacastrum</i>
Seaside Vine	<i>Convolvulus brasiliensis.</i>
Senna Tree	<i>Cassia emarginata.</i>
Sensitive Plant	<i>Mimosa pudica.</i>
Seville Orange Tree	<i>Citrus vulgaris.</i>
Shaddock Tree	<i>Citrus Decumana.</i>
Shallot	<i>Allium Ascalonium.</i>
Short Grass	<i>Panicum distachyon.</i>
Shrubby Goat Weed, or Sweet Weed	<i>Capraria biflora.</i>
Siberian Motherwort	<i>Leonurus sibiricus.</i>
Silk Cotton Tree	<i>Bombax Ceiba.</i>
Silk Grass	<i>Agave americana.</i>
Silk Grass	<i>Agave vivipara.</i>
Silver-leaved Plant	<i>Tussilago nutans.</i>
Silver Wood	<i>Koelera serrata.</i>
Silver Vine	<i>Hedysarum volubile.</i>
Small Cotton Tree	<i>Gossypium barbadense.</i>
Small Cotton Tree	<i>Gossypium herbaceum.</i>
Small Cotton Tree	<i>Gossypium vitifolium.</i>
Small Barren Pine	<i>Bromelia paniculegera.</i>
Small Duck Weed	<i>Ceratophyllum demersum.</i>
Small-grained Black Pepper	<i>Piper Amalago.</i>
Small Red Thrubba	<i>Solanum torvum.</i>

Small Pomegranate Tree.	<i>Punica nana</i> .
Small Wild Cucumber.	<i>Melothria pendula</i> .
Smaller Cloven-Berry	
Bush	<i>Casearia parviflora</i> .
Smaller Indian Kale	<i>Arum divaricatum</i> .
Smooth and Prickly	
Lawsonia	<i>Lawsonia alba</i> .
Smooth Primrose Willow.	<i>Jussiaea octonervia</i> .
Snake Gourd	<i>Trichosanthes anguina</i> .
Snake Wood	<i>Cecropia peltata</i> .
Soap Tree, or Soap-Berry.	<i>Sapindus saponaria</i> .
Sorrel	<i>Hibiscus Sabdariffa</i> .
Sour Orange Tree	<i>Citrus vulgaris</i> .
Soursop Tree	<i>Anona muricata</i> .
South Sea Rose	<i>Nerium odorum</i> .
Spanish Ash	<i>Lonchocarpus violaceus</i> .
Spanish Lemon Tree	<i>Citrus medica</i> .
Spanish Needle	<i>Bidens leucantha</i> .
Spanish Needles	<i>Yucca Draconis</i> .
Spanish Oak	<i>Inga Bourgoni</i> .
Spear-Mint	<i>Mentha viridis</i> .
Spirit Weed	<i>Ægiphila martinicensis</i> .
Spirit Weed	<i>Ægiphila elata</i> .
Spikenard	<i>Byrtropogon suaveolens</i> .
Spur Pepper	<i>Capsicum frutescens</i> .
Squashes	<i>Cucurbita Melopepo</i> .
Stinking Weed	<i>Cassia occidentalis</i> .
Star Apple	<i>Chrysophyllum Cainito</i> .
Star Grass	<i>Schoenus stellatus</i> .

Star Plum	<i>Chrysophyllum mono-</i> <i>pyrenum.</i>
Stopper-Berry Tree	<i>Cerasus sphærocarpa.</i>
Strawberry Pear	<i>Cereus trigonus.</i>
Sugar Apple Tree	<i>Anona squamosa.</i>
Sugar Bean	<i>Phaseolus lunatus.</i>
Sugar Cane	<i>Saccharum officinarum.</i>
Supple Jack	<i>Paullinia Cururu.</i>
Supple Jack	<i>Paullinia barbadensis.</i>
Sweet Briar	<i>Acacia tortuosa.</i>
Sweet Briar Rose	<i>Rosa rubiginosa.</i>
Sweet Cassava	<i>Janipha Löflingii.</i>
Sweet Gourd	<i>Trichosanthes anguina.</i>
Sweet Heart	<i>Desmodium supinum.</i>
Sweet Marjoram	<i>Origanum Majorana.</i>
Sweet Lime	<i>Triphasia trifoliata.</i>
Sweet Lime	<i>Citrus Limetta.</i>
Sweet Pea	<i>Lathyrus odoratus.</i>
Sweet-scented Birthwort.	<i>Aristolochia odoratissima</i>
Sweet-scented China	
Rose	<i>Rosa chinensis..</i>
Sweet-scented Flea Bane.	<i>Baccharis odorata.</i>
Sweet-scented Hemp	
Agrimony	<i>Eupatorium odoratum.</i>
Sweet-scented Water	
Lily	<i>Nymphæa odorata.</i>
Sweet Sop, or Sugar	
Apple Tree	<i>Anona squamosa.</i>
Sweet Wood Tree	<i>Laurus caroliniensis.</i>

Sweet Wood Tree	<i>Laurus parviflora.</i>
Tamarind Tree	<i>Tamarindus indica.</i>
Tammatas	<i>Solanum Lycopersicon.</i>
Teak Wood, or Indian	
Oak	<i>Tectona grandis.</i>
Thistle, or Rabbit Weed.	<i>Sonchus oleraceus.</i>
Timber Sweet Wood	<i>Laurus caroliniensis.</i>
Tithymaloides	<i>Euphorbia Tithymaloides</i>
Top-Knot Plum	<i>Spondias purpurea.</i>
Trinidad Butterfly Plant.	<i>Oncidium Papilio.</i>
Trumpet Tree, or Snake	
Wood	<i>Cecropia peltata.</i>
Tuberoze	<i>Polianthes tuberosa.</i>
Turmerick	<i>Curcuma longa.</i>
Turkey Berry Tree	<i>Cordia Collococca.</i>
Turk's Head	<i>Melocactus communis.</i>
Turnip	<i>Brassica Rapa.</i>
Vine Nettle	<i>Tragia volubilis.</i>
Virginian Pepper Wort.	<i>Lepidium virginicum.</i>
Upright Pink	<i>Dianthus virgineus.</i>
Water Cress	<i>Nasturtium officinale.</i>
Water Lemon	<i>Passiflora laurifolia.</i>
Water Lemon	<i>Passiflora maliformis.</i>
Water Sweet Wood	<i>Laurus parviflora.</i>
Water Weed	<i>Spigelia Anthelmia.</i>
West India Tea	<i>Capraria biflora.</i>

Welch Golden Rod	<i>Solidago cambrica.</i>
White Bully Tree	<i>Bumelia salicifolia.</i>
White Cedar Tree	<i>Bignonia Leucoxylon.</i>
White Cocco, or Tyre ...	<i>Caladium Macrorhizon.</i>
White-flowered Jasmine	
Tree	<i>Plumeria alba.</i>
White Hercules	<i>Zanthoxylum fraxineum.</i>
White Hoop	<i>Tournefortia bicolor.</i>
White Sage	<i>Lantana involucrata.</i>
White Willow	<i>Capparis Breynia.</i>
White Wood Tree	<i>Bignonia Leucoxylon.</i>
Wild Basil	<i>Ocymum americanum.</i>
Wild or Bastard Ipeca-	
cuanha	<i>Asclepias curassavica.</i>
Wild or Bastard French	
Guava	<i>Cassia alata.</i>
Wild Bay Berry Tree ...	<i>Eugenia coriacea.</i>
Wild Cane	<i>Erianthus arundinaceus.</i>
Wild Calabash	<i>Cestrum laurifolium.</i>
Wild Cassava	<i>Jatropha Gossypifolia.</i>
Wild Cinnamon	<i>Canella alba.</i>
Wild Clary	<i>Heliotropium indicum.</i>
Wild Clary	<i>Heliotropium parviflo-</i>
	<i>rum.</i>
Wild Clove, or Bay-	
Berry Myrtle	<i>Myrcia acris.</i>
Wild Coffee	<i>Coffea occidentalis.</i>
Wild Coffee	<i>Pavetta pentandra.</i>
Wild Coffee	<i>Psychotria tenuifolia.</i>

Wild Dolly	<i>Phaseolus semierectus.</i>
Wild Dwrah	<i>Cyperus strigosus.</i>
Wild Elder Bush	<i>Piper umbellatum.</i>
Wild French Guava	<i>Cassia occidentalis.</i>
Wild Hemp	<i>Malachra capitata.</i>
Wild Hemp	<i>Malachra fasciata.</i>
Wild Hiccory	<i>Rivina octandra.</i>
Wild Honey Tree	<i>Casearia parviflora.</i>
Wild Hops	<i>Hyptis capitata.</i>
Wild Indigo	<i>Indigofera Anil.</i>
Wild Lavender	<i>Heliotropium curassavi-</i> <i>cum.</i>
Wild Lily	<i>Epidendrum ciliare.</i>
Wild Liquorice	<i>Abrus precatorius.</i>
Wild, or Monkey Eddoe.	<i>Caladium nymphæifo-</i> <i>lium.</i>
Wild Ockro	<i>Malacra capitata.</i>
Wild Olive	<i>Bontia daphnoides.</i>
Wild Parsley	<i>Cardiospermum Helica-</i> <i>cabum.</i>
Wild Pepper Grass	<i>Lepidium virginicum.</i>
Wild Pines	<i>Yucca Draconis.</i>
Wild Pomegranate	<i>Volkameria aculeata.</i>
Wild Purslane	<i>Portulaca parvifolia.</i>
Wild Purslane	<i>Portulaca pilosa.</i>
Wild Sage	<i>Lantana involucrata.</i>
Wild Senna, or Wild	<i>Cassia emarginata.</i>
Cassia Fistula	
Wild Shaddock	<i>Citrus medica.</i>

Wild Spikenard	<i>Bystropogon pectinatum.</i>
Wild Slip	<i>Convolvulus hederaceus.</i>
Wild Tamarinds	<i>Cassia Chamæcrista.</i>
Wild Tobacco	<i>Baccharis odorata.</i>
Wild Tulip	<i>Amaryllis zeylanica.</i>
Wild Water Lemon Vine.	<i>Passiflora foetida.</i>
Wild Wormwood	<i>Parthenium Hysterophorus.</i>
Winged - seeded Horse- Radish Tree	<i>Moringa pterygosperma.</i>
Woolly-Pyroe	<i>Phaseolus Mungo.</i>
Worm Seed Weed	<i>Chenopodium anthelmin- ticum.</i>
Yams	<i>Dioscorea sativa.</i>
Yellow Hercules	<i>Zanthoxylum Ochroxyl- lum.</i>
Yellow Hercules	<i>Zanthoxylum Clava Her- culis.</i>
Yellow, or Jamaica Plum.	<i>Spondias lutea.</i>
Yellow Lupin	<i>Crotalaria retusa.</i>
Yellow Nicker	<i>Guilandina Bonduc. α.</i>
Yellow Thistle	<i>Argemone mexicana.</i>

FINIS.

INDEX.

	Page		Page		Page
ABRUS	288	Avicennia	253	Cardiospermum ..	158
Acacia	402	Baccharis	324	Carica	293
Acrosticum	410	Badiera	285	Carolinea	271
Adenanthera	175	Balsamina	115	Caryophyllus	210
Adiantum	414	Bambusa	148	Casearia	182
Acrocomia	355	Banisteria	191	Cassia	167
Adansonia	272	Basella	131	Ceanothus	112
Agave	133	Bauhinia	165	Cecropia	382
Ægiphila	70	Begonia	357	Cedrela	113
Æschynomene ...	299	Besleria	242	Celosia	116
Agati	313	Beta	125	Cenchrus	48
Ageratum	322	Bidens	320	Cerasus	215
Agrostis	63	Bignonia	243	Ceratophyllum ...	357
Aletris	144	Bixa	226	Cerbera	118
Alisma	149	Boerhaavia	32	Cereus	202
Allamanda	118	Bombax	273	Cestrum	105
Allium	141	Bontia	253	Chenopodium	125
Aloe	145	Brassica	257	Chiococca .. .	94
Alpinia	31	Bromelia	134	Chloris	53
Althæa	277	Browallia	248	Chrysanthemum ..	330
Amaranthus	352	Brownea	270	Chrysobalanus ...	215
Amaryllis	139	Bryonia	376	Chrysophyllum ..	108
Ammannia	76	Bryophyllum	160	Cicca	349
Amyris	154	Budleia	72	Cissus	74
Anacardium	164	Bumelia	109	Citharexylum ...	245
Andropogon	52	Bunchosia	189	Citrus	315
Anethum	128	Bursera	405	Clethra	183
Anona	232	Byrsonima	189	Clerodendrum ...	251
Anthemis	331	Bytropogon	235	Clitoria	295
Aphelandra	249			Clusia	398
Apium	129			Coccoloba	155
Arachis	291	Cajanus	298	Cochlearia	256
Ardisia	107	Caladium	358	Cocos	355
Areca	371	Calendula	335	Coffea	93
Argemone	225	Callisia	44	Coix	57
Aristolochia	340	Calophyllum	228	Commelina	43
Artemisia	323	Calotropis	122	Conocarpus	95
Artocarpus	345	Canavalia	310	Convolvulus	86
Arum	360	Canella	195	Corchorus	229
Asclepias	123	Canna	29	Coreopsis	334
Asparagus	142	Capparis	223	Cordia	82
Aspidium	412	Capraria	248	Cosmea	333
Aster	325	Capsicum	103	Crataeva	196

INDEX.

	Page		Page		Page
Cremanium	182	Guilandina	172	Malachra	276
Crescentia	244	Gynandropsis	260	Malpighia	187
Crinum	141			Mammea	227
Crotalaria	290	Haematoxylon ...	174	Mangifera	114
Croton	363	Hedysarum	303	Maranta	30
Cucumis	373	Helianthus	333	Mauritia	392
Cucurbita	380	Heliotropium	80	Melia	178
Curcuma	32	Hernandia	348	Melicocca	152
Cyathæa	414	Heterospermum ..	329	Melocactus	201
Cymbopogon	53	Hibiscus	280	Melothria	42
Cynanchum	124	Hippomane	368	Mentha	236
Cynodon	62	Hoya	123	Mimosa	399
Cynosurus	65	Hura	371	Mirabilis	95
Cyperus	45	Hydrangea	185	Momordica	377
		Hymenæa	166	Moræa	42
Dahlia	326	Hyptis	237	Moringa	173
Datura	96			Morus	351
Daucus	127	Jacquinia	107	Mucuna	296
Desmanthus	401	Jambosa	213	Musa	396
Desmodium	302	Janipha	365	Myginda	79
Dianthus	185	Jasminum	33	Myrcia	208
Dioscorea	389	Jatropha	364	Myristica	395
Dodonæa	153	Indigofera	304	Myrodia	284
Dolichos	308	Inga	400	Myrtus	207
Donax	66	Ipomœa	90		
Dracæna	143	Juniperus	394	Nasturtium	256
Drymaria	67	Jussiaea	180	Nelumbium	231
Duranta	250	Justicia	35	Nerium	119
		Ixora	69	Nicotiana	97
Ecastaphyllum ..	287			Nymphæa	225
Elais	391	Kerria	218		
Elephantopus	337	Kœlera	385	Ochroma	264
Epidendrum	338	Kyllingia.....	44	Ocymum	241
Erianthus	51			Oncidium	338
Eringium	128	Lablab	305	Oplismenus	64
Erythrina	288	Lactuca	320	Opuntia	203
Eugenia	210	Lagenaria	372	Origanum	240
Evolvulus	132	Lagerstroemia ...	227		
Eupatorium	321	Lantana	246	Pancratium	138
Euphorbia	341	Lathyrus	312	Pandanus	381
		Laurus	162	Panicum	59
Ficus	406	Lavandula	235	Parkinsonia	166
		Lawsonia	153	Parthenium	335
Garcinia	195	Leonotis	240	Paspalum	58
Gardenia	117	Leonurus	238	Passiflora	264
Gloxinia	242	Lepidium	255	Pavetta	70
Gomphræna	126	Leucas	239	Paullinia	157
Gossypium	277	Lonchocarpus ...	299	Pectis	328
Gouania	399	Lourea	301	Pelargonium	207
Guaiacum	176	Lycopodium	409	Pereskia	205
Guettarda	354			Persea	161

INDEX.

	Page		Page		Page
Petitia	73	Rosmarinus	38	Terminalia	397
Petiveria	151	Ruellia	249	Tetrazygia	181
Petræa ..	246			Theobroma	314
Phaseolus	292	Saccharum	50	Thespesia	279
Phoenix	383	Salvia	38	Thrinax	140
Phyllanthus	370	Sanseveria	136	Thumbergia	254
Physalis	97	Sapindus	159	Thymus	241
Piper	39	Sapium	369	Tournefortia	81
Pisonia	150	Sarcostemma	124	Tragia	347
Pisum	296	Satureja	234	Tradescantia	137
Pitcairnia	136	Schoenus	45	Tribulus	175
Plantago	75	Scirpus	47	Trichosanthos	379
Plumbago	85	Scleria	57	Triopteris	190
Plumeria	121	Scoparia	75	Triphasia	179
Poa	65	Sechium	378	Triumfetta	197
Poinciana	171	Securidaca	286	Tussilago	324
Polianthes	144	Sesbania	312		
Polypodium	411	Sesuvium	217	Varronia	84
Pontederia	138	Sida	273	Verbesina	331
Portulaca	198	Siderodendrum	73	Vigna	312
Pothos	77	Sideroxylon	111	Vinca	119
Psidium	205	Sinapis	258	Viscum	384
Psychotria	92	Solanum	98	Vitis	115
Pteris	413	Solidago	326	Volkameria	252
Punica	214	Sonchus	319		
		Sorghum	55	Urtica	350
Quassia	177	Spermacoce	68		
		Spigelia	85	Waltheria	262
Raphanus	259	Spondias	192		
Reseda	200	Stachys	238	Xylophylla	130
Rhabdochloa	54	Stylosanthos	301		
Rhapis	404	Swietenia	177	Yucca	145
Rhizophora	194				
Rhynchosia	311	Tagetes	327	Zanthoxylum	386
Ricinus	367	Talinum	179	Zea	56
Riedleia	263	Tamarindus	261	Zingiber	31
Rivina	77	Tectona	112	Zinnia	329
Rosa	219	Teramnus	289	Zizyphus	113