IMAGE EVALUATION TEST TARGET (MT-3)

Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1983

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- ☐ Coloured covers/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- ☐ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☒ Bound with other material/
Relié avec d'autres documents
- ☐ Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- ☐ Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- ☐ Additional comments:/
Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☒ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
				✓							

The copy filmed here has been reproduced thanks to the generosity of:

Library
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ONTARIO AGRICULTURAL COLLEGE
EXPERIMENT STATION

BULLETIN LXVII

WINTER WHEAT EXPERIMENTS.

BY THOMAS SHAW, PROFESSOR OF AGRICULTURE, AND
C. A. ZAVITZ, ASSISTANT SUPERINTENDENT
OF EXPERIMENTS.

PUBLISHED BY THE DEPARTMENT OF AGRICULTURE
Aug. 12, 1891

TORONTO

PRINTED BY WARWICK & SONS

MINISTER OF AGRICULTURE

HON. JOHN DRYDEN, TORONTO.

Ontario Agricultural College and Experimental Farm, Guelph,
under control of the Minister of Agriculture.

JAMES MILLS, M.A.		President.
THOMAS SHAW		Professor of Agriculture and Farm Superintendent.
A. E. SHUTTLEWORTH, B.A. Sc.		Professor of Chemistry
J. HOYES PANTON, M.A., F.G.S.		Professor of Natural History and Geology.
F. C. GRENSIDE, V.S.		Professor of Veterinary Science.
H. H. DEAN, B.S.A.		Professor of Dairy Husbandry.
E. LAWRENCE HUNT, B.A.		Assistant Resident and Mathematical Master.
CAPTAIN WALTER CLARKE		Instructor in Drill and Gymnastics.
C. A. ZAVITZ, B.S.A.		Assistant Superintendent of Experiments.
GEORGE HARCOURT, B.S.A.		Assistant Chemist.
A. MCCALLUM		..Bursar.

ADVISORY BOARD.

G. C. JAMES, M.A., <i>Secretary</i>		Deputy Minister of Agriculture, Toronto.
JOHN I. HOBSON, <i>Chairman</i>		Mosborough, County of Wellington.

BULLETIN LXVII.

EXPERIMENTS WITH WINTER WHEATS.

The winter wheat crop will always be one of considerable importance to Ontario notwithstanding the facilities for growing it in Manitoba and the Northwest. This view is supported by the following reasons along with other others which might be given :

1. A large portion of the province is well adapted to the growth of wheat. In view of this fact the advantage of growing a less amount than will suffice for home consumption is more than doubtful. Adaptability should always be a prime factor in determining the crops that should be grown. That winter wheat may still be grown in western Ontario in as fine form as in any country in the world has been amply demonstrated by the yields of the present year.

2. The winter wheat crop may with much advantage be made to form a prominent feature of a regular rotation.

3. This crop provides bedding for live stock more abundantly and more cheaply than any crop now grown. This feature alone adds much more to the value of the crop than is generally supposed.

4. It furnishes a crop suitable for sowing grasses along with it, a fact of no little significance in view of the probable decrease in the acreage of barley. It also admits of the autumn sowing of timothy which generally ensures a catch of the seed in any season.

5. The increased attention to stock production will tend to improve the yield per acre owing to the increase of fertilising materials which are thus made available to the farmer.

6. The growth of winter wheat tends to a more even distribution of labor throughout the year.

7. It encourages the manufacture of flour in the country, and this furnishes a plentiful supply of bran, which is almost indispensable for certain feeding purposes.

8. In growing winter wheat for home consumption the farmers always will have protection to the extent of the cost of conveyance from other provinces.

Our farmers should therefore continue to give careful attention to the growth of winter wheat in the parts of the province favorable to the same. The aim should be not so much to grow a large acreage

as to adopt such methods of cultivation and to secure such varieties as are likely to give a large yield per acre. With this end in view the importance of good drainage, sowing at the right time and suitable conditions of soil in relation to rotation, fertility and natural adaptability cannot easily be over-estimated. All things considered it would be better not to sow winter wheat at all than to sow it under conditions not likely to produce a good crop.

SELECTION OF VARIETIES. In selecting varieties to sow, a careful regard must also be given to the wants of the millers. Those good yielding varieties only should be grown which are well adapted to milling purposes. Some years ago quantity was the great consideration with the grower. At present quality is at least equally important with quantity. This statement finds ample confirmation in the low price paid for the Wild Goose wheat notwithstanding its ample yields. The choice of the millers is not a mere arbitrary one. As in this fastidious age the tastes of the consumer rule, the miller must cater to those tastes or he will lose his custom. The consumer demands a beautifully appearing snowy white bread, and the miller must furnish flour that will produce it. In this we find one of the principal reasons for his preference for white wheats. The baker wants a strong flour, that is, one rich in gluten, and the miller must give it to him or he will cease to buy. Because of this the Manitoba Fyfe spring wheat stands high in favor with the millers. A heavy weighing wheat gives a larger percentage of flour than a light weighing one, hence the preference of the millers for the varieties which weigh the heaviest. The farmers therefore will do well to give diligent heed to the recommendations of the millers as to the varieties which stand high in favor with them, and in making their selections for sowing they should choose accordingly.

THE VARIETIES GROWN. There were in all some 70 plots of winter wheat grown at this station during the present year, including 51 varieties. A few of the varieties were grown in duplicate plots, and in another field we grew 10 varieties in acre and half acre plots. These larger plots, however, were simply duplicates of varieties grown in the smaller plots, with the exception of the Longberry Red, a variety introduced last year from Indiana, and which through some unfortunate oversight was not grown in the small plots. The details relating to the larger plots will be given in the annual report. Of the 51 varieties 24 were Canadian and American, and 27 were foreign. The foreign varieties came chiefly from Germany, England, France and Russia. As none of the foreign varieties have as yet proved equal to some of the Canadian and American kinds we do not feel justified at present in recommending them to be sown. This bulletin, therefore, gives the particulars relating to 23 Canadian and American varieties grown under exactly the same conditions.

GENERAL CLASSIFICATION OF VARIETIES OF FALL WHEATS.

Bald Head.	White Chaff.	White Grain.	{	(2) Garfield or Natural Cross.
			{	(3) Surprise.
	Red Grain.	Red Grain.	{	(4) Canadian Velvet Chaff.
			{	(5) Bonnell or Landreth.
Bearded Head.	White Chaff.	White Grain.	{	(11) Winter Pearl.
			{	(14) Martin Amber.
	Red Chaff.	Red Grain.	{	(1) American Bronze.
			{	(9) Red Velvet Chaff.
	Red Chaff.	Red Grain.	{	(10) Jones' Winter Fyfe.
			{	(12) Rogers.
	Red Chaff.	Red Grain.	{	(19) Golden Drop.
			{	(23) New Monarch.
	White Chaff.	White Grain.	{	(6) Seneca or Clawson.
			{	(7) Standard.
Bearded Head.	Red Chaff.	Red Grain.	{	(8) Early Red Clawson.
			{	(16) Manchester.
	White Chaff.	White Grain.	{	(17) Bulgarian.
			{	(22) Democrat.
	Red Chaff.	Red Grain.	{	(13) Egyptian.
			{	(15) Golden Cross or Volunteer.
	Red Chaff.	Red Grain.	{	(18) Red Lyon.
			{	(20) Hybrid Mediterranean.
	Red Chaff.	Red Grain.	{	(21) Lancaster.
			{	(21) Lancaster.

The above classification is based upon the recognition of all winter wheats as belonging to the bald or bearded varieties. But observe that those varieties with very short beards, often found only on the upper portion of the head, are classified as bald. The wheats in each of these general divisions are again classified according to the color of the grain. It should be borne in mind that sometimes we have white chaff with red wheat, and red chaff with white wheat. The finer distinctions of shade in color, as amber, bronze, etc., are not given in this classification, as they are liable to vary with a change of soil and climate. The number standing before the name of each variety gives the order of the yield of the said variety.

LOCATION AND SOIL. All the varieties of winter wheat both native and foreign were grown in plots side by side in the same range. These plots contain exactly the one-fiftieth of an acre each. The aspect of the land is north-easterly, but the slope is very gentle. The soil may be termed a clay loam with good natural drainage.

PREPARATION OF THE SOIL AND SOWING. The soil was prepared on the bare fallow system as it was felt that a test of this nature should be made under the most favorable conditions

attainable, to bring out to the fullest extent the capabilities of the grains grown. This is the only bare fallow that we had on the farm. The ground was plowed twice the previous summer. Additional surface cultivation was also given. Farmyard manure was applied at the rate of 15 tons to the acre.

MANNER AND TIME OF SEEDING. The seed was sown by hand, as we have no machines suitable for sowing in drills in plots of the size mentioned. The plots were all sown September 4th with the exception of the Winter Pearl variety, which was sown on September 8th. The same amount of seed by weight was sown upon each plot, and it was sown at the rate of $1\frac{3}{4}$ bushels per acre. As the location was favorable, the soil well prepared, the weather favorable from time of sowing until time of reaping and especially so during the ripening period, as there was no rust or blight to interfere with the filling of the grain, and as all the conditions were exactly similar and favorable to the most complete development, we may safely conclude that a more favorable opportunity for testing the full capabilities of these wheats when at their best could not easily have been furnished.

Table 1 shows the average yields of twenty-three varieties for 1891.

No.	Varieties.	Weight of grain per measured bushel.	Amount of straw per acre.	Yield of grain per acre (bush. 60 lb.)
		lb.	tons.	
1	American Bronze.....	63.00	2.16	65.2
2	Garfield or Natural Cross.	62.00	2.52	64.4
3	Surprise.....	63.25	1.99	63.8
4	Canadian Velvet Chaff...	62.25	1.82	59.9
5	Bonnell or Landreth.....	62.75	2.26	59.5
6	Seneca or Clawson.....	62.75	2.07	58.9
7	Standard.....	63.50	2.02	58.3
8	Early Red Clawson.....	62.50	1.93	58.3
9	Red Velvet Chaff.....	64.00	1.92	56.9
10	Jones Winter Fyfe.....	64.25	1.77	56.3
11	Winter Pearl.....	61.00	2.51	55.5
12	Rogers.....	62.50	1.83	54.6
13	Egyptian.....	64.00	1.94	53.4
14	Martin Amber.....	61.00	2.13	50.4
15	Golden Cross or Volunteer.	64.00	2.17	49.9
16	Manchester.....	64.50	1.53	47.7
17	Bulgarian.....	64.50	1.60	46.3
18	Red Lyon.....	64.00	1.67	44.5
19	Golden Drop.....	63.00	1.61	44.3
20	Hybrid Mediterranean.....	64.25	1.92	44.0
21	Hamaker.....	64.50	2.11	42.8
22	Democrat.....	64.50	1.69	41.6
23	New Monarch.....	63.50	1.72	40.8
	Average.....	63.28	1.95	52.93

It will be noticed that in the above table the 23 varieties of wheat mentioned have given most extraordinary yields, and produced wheat of an uncommonly good quality. The highest yield is from the American Bronze, which produced at the rate of 65.2 bush. per acre, and the lowest is from the New Monarch, which produced at the rate of 40.8 bush. per acre. The average yield of the 23 varieties is at the rate of 52.93 bush. per acre. The highest weight per bush. was $64\frac{1}{2}$ lb., reached by several of the varieties; the lowest weight was 61 lb. per bush., and the average weight of the 23 varieties was 63.28 lb. per bush. In view of the extraordinary yields thus obtained, and the even more extraordinary weights of the grain, the curiosity is not unnatural which would desire to know whether these yields have a parallel on the continent of America.

It should be stated here that the weights of the different varieties of wheat, as given in the above table, were obtained from tests conducted at our request by Mr. James Goldie, Guelph, of James Goldie & Sons. When the weights were taken Mr. Goldie considered all the varieties sufficiently dry to grind, with but two or three exceptions.

DISCOUNTING THE YIELD. It may be well, also, to remark that yields such as the above would not have been obtained from whole fields of these respective varieties, nor could they be looked for from cultivation given under average conditions. Small plots yield more relatively than large ones, owing to cultivation around the borders and probably to other causes, and from this extra yield it has been thought that if one-fifth were deducted, the result would be a fair average to expect from a whole field under like conditions in other respects.

Table II gives comparative summary results :

Class.	Number varieties.	Weight per measured bushel.	Yield per acre. (Standard bushel by weight).
{ Bald	16	62.88	55.93
{ Bearded.....	7	64.25	46.07
{ White Chaff.....	15	63.03	54.19
{ Red Chaff.	8	63.75	50.55
{ White Wheat....	10	62.75	55.86
{ Red Wheat.....	13	63.71	50.68

From this table it is apparent that when wheat is grown under favorable conditions the bald varieties yield considerably more than the bearded, but that the bearded varieties weigh heavier on an average. It would be interesting to know how far these comparative

results would be modified on soils less fertile and in seasons less propitious. The white chaff varieties were also the best yielders, but they did not quite equal the red chaff varieties in weight, and the white grain varieties also gave the highest yield per acre, but the red grain varieties were ahead in weight.

TABLE III gives many interesting particulars relating to the different varieties :

Number.	Variety.	Date of maturity.	Length of plant.	Width of 25 straws side by side.	Average length of 10 heads.	Average No. of grains per head.	Weight of 1,000 grains. in grammes.
		July.	inches	mm.	inches.		
1	American Bronze	25	39	97	3.1	29.2	47.4
2	Garfield or Natural Cross	23	39	87	2.9	29.3	45.2
3	Surprise	21	38	87	3.0	33.1	41.8
4	Canadian Velvet Chaff	21	35	81	2.9	31.4	43.0
5	Bonnell or Landreth ..	23	39	85	3.2	34.2	48.9
6	Seneca or Clawson	23	39	78	3.1	30.9	42.4
7	Standard	22	36	91	2.7	35.1	39.3
8	Early Red Clawson	20	37	91	2.7	26.4	48.3
9	Red Velvet Chaff	23	39	87	2.9	30.1	43.1
10	Jones' Winter Fyfe	22	38	83	2.8	34.1	39.6
11	Winter Pearl	24	40	78	3.7	30.5	49.7
12	Rogers	23	36	74	3.1	30.6	43.5
13	Egyptian	21	44	83	2.8	29.4	40.5
14	Martin Amber	23	38	78	3.3	37.4	47.2
15	Golden Cross or Volunteer	21	39	85	2.5	27.1	43.3
16	Manchester	22	37	73	2.8	28.8	41.1
17	Bulgarian	21	40	70	2.9	30.8	40.1
18	Red Lyon	21	38	70	2.5	22.6	44.2
19	Golden Drop	23	39	74	3.1	34.9	44.3
20	Hybrid Mediterranean	23	38	80	2.5	25.7	44.7
21	Lancaster	21	38	75	2.6	21.3	43.8
22	Democrat	22	40	72	2.7	27.2	40.1
23	New Monarch	22	37	80	3.2	32.2	41.4

From the foregoing table we learn that there were only four days of difference in the ripening of the 23 varieties. The greatest difference in the average length of the plant in the plots was only seven inches, and the average length of the plant in all the plots was 38.4 inches. The width of the straw is given in millimeters to give an idea of size and to some extent of its strength. As the straw in all the varieties stood fairly erect until the time of cutting, it would not be so easy perhaps to convey a correct idea in any other way. The average length of head was 2.9 inches, and the average number of grains per head was 30.1. From the column giving the weight in grammes of 1,000 grains of each variety, a somewhat accurate idea of the comparative size of the grains can be obtained.

The seed in 15 varieties was obtained from the College, and the balance as follows : Nos. 1, 8, 10, and 15 (Steele) ; Nos. 4 and 16 (Pearce) ; No. 11 (Wilkinson) and No. 13 (Gordon).

The color of the straw was white in all varieties, except Nos. 15 and 21, which were a deep slate, and Nos. 16, 18 and 20, which were a light slate.

The surface of chaff was smooth with the exception of the three velvet varieties, Nos. 4, 9 and 10.

In table iv. the yields of the 15 leading varieties for 1890 are given in comparison with the yields of the same for 1891 :

Varieties.	Weight of grain per measured bushel.			Yield of grain per acre in bushels of 60 lb.		
	1891.	1890.	Average 1890-1.	1891.	1890.	Average 1890-1.
Surprise.	63.3	58.7	61.0	63.8	29.6	46.7
Red Velvet Chaff.	64.0	61.0	62.5	56.9	35.8	46.3
Bonnell or Landreth.	62.8	61.0	61.9	59.5	32.1	45.8
Early Red Clawson.	62.5	58.0	60.3	58.3	32.1	45.2
Standard.	63.5	59.3	61.4	58.3	31.7	45.0
Rogers.	62.5	60.8	61.7	54.6	34.6	44.6
Seneca or Clawson.	62.8	59.0	60.9	58.9	25.4	42.1
Martin Amber.	61.0	62.5	61.8	50.4	32.5	41.5
Golden Gross or Volunteer.	64.0	58.3	61.2	49.9	32.1	41.0
Manchester.	64.5	61.0	62.8	47.7	30.4	39.1
Golden Drop.	63.0	61.2	62.1	44.3	32.5	38.4
Lancaster.	64.5	61.7	63.1	42.8	33.3	38.1
Hybrid Mediterranean.	64.3	60.2	62.2	44.0	32.1	38.1
Red Lyon.	64.0	60.0	62.0	44.5	29.2	36.9
New Monarch.	63.5	56.7	60.1	40.8	20.0	30.4
Averages.	63.3	60.0	61.7	51.6	30.9	41.3

It will be observed that the Surprise heads the list, while as has been already stated it ranks high as a milling wheat. The Red Velvet Chaff which stood first last year for yield and weight combined, is second in comparison for the two years, but drops down to ninth place in the list of 1891. It may also be observed that on the whole, the different varieties have given yields in much the same order for the two years. The average yield per acre for the two years, and the average weight per bushel, are certainly satisfactory. The former was no less than 43.1 bushels, and the latter 61.7 lb.

REPORT OF THE DOMINION MILLERS' ASSOCIATION.

A deputation was appointed by the Dominion Millers' Association to visit this station, examine the different kinds of winter wheat and report upon the same. The deputation consisted of ten persons, in-

cluding the President and Secretary of the Association. The visit was made on August 5th and the following is a summary of the report :

"The Committee are anxious to extend the growth of white wheat. They think very highly of the Surprise variety and advise farmers to sow it. They highly recommend the Canadian Velvet Chaff and regard it is a good milling wheat. They also consider the Bulgarian an extra good milling wheat.

"In regard to the red varieties they advise that the Rogers should not be sown as it is inclined to smut and contains no good milling properties. They also fear that the Manchester is a weak wheat for milling purposes, but this conclusion is only based on the grinding of small quantities. They consider the Jones' Winter Fyfe an extraordinarily good wheat for grinding, and they are of the opinion that if the farmers could raise it plentifully it would obviate the necessity of using so much Manitoba wheat. They very highly recommend the Hybrid Mediterranean, looking upon it as a good wheat, and they also highly recommend the Longberry Red."

More complete particulars regarding these wheats may be obtained from the report of the Millers' Association, which will doubtless have reached the public through the press before the issuing of this bulletin.

CONCLUSIONS.

The results of the experiments may be thus summarised :

1. The splendid returns obtained in this experiment prove that the capabilities of Ontario, as a wheat producing country, are still of a high order.

2. The bald wheats have on an average given 9.86 bushels more per acre, or 21.42 per cent. than the bearded varieties, but the latter have weighed on an average 1.37 lb. more per bushel.

3. The white wheats have given an average of 5.18 bushels more per acre than the red wheats, and they also stand higher in the estimation of the millers than the latter.

4. The bald white chaff white wheats gave an average of 13.6 bushels per acre more than the bearded red chaff red wheats.

5. The seven leading varieties in point of yield were all white wheats except the American Bronze.

6. These experiments tend to confirm the popular opinion that white wheats under favorable condition will give more bountiful returns than red wheats, and that the same may be said of the bald varieties as contrasted with the bearded.

7. The four best yielding white wheats for 1891 were the Garfield, Surprise, Canadian Velvet Chaff and Bonnell; and the four best yielding varieties of red wheat were the American Bronze,

Early Red Clawson, Red Velvet Chaff and Jones' Winter Fyfe, in the order named in both instances.

8. The best four weighing varieties were the Manchester, Bulgarian, Lancaster and Democrat, each of which gave $64\frac{1}{2}$ lb. per bushel.

9. The three velvet chaff varieties gave an average yield of 4.77 bushels per acre in excess of the mean average of the 23 varieties, and weighed .22 lb. more per bushel, and they are also included in the leading varieties mentioned in conclusion 7.

10. Of the varieties enumerated in this bulletin, the Dominion Millers' Association recommend the following as the most serviceable for milling purposes, viz., of the white wheats, the Surprise, Canadian Velvet Chaff and Bulgarian; and of the red wheats, the Jones' Winter Fyfe, the Hybrid Mediterranean and the Longberry Red.

DISTRIBUTION OF SEED.

As we have received many enquiries regarding seed, we append the following in reference to the mode of distribution that we have decided to adopt: We will supply any of the following varieties, viz., the Surprise, Red Velvet Chaff, Bonnell or Landreth, Early Red Clawson, Bulgarian, Garfield, American Bronze and Canadian Velvet Chaff in one bushel and half bushel lots. As we have only a limited quantity of the seed of each of these varieties, we can only furnish seed while the supply lasts and in the order in which we receive the applications. The prices charged will be moderate. For further information apply to the Professor of Agriculture.

The varieties will be distributed in smaller lots through the medium of the Ontario Agricultural and Experimental Union. The Experimental Union, which meets annually at the Agricultural College, is composed of the officers, ex-students and students of the college, and all farmers throughout the province are invited to co-operate in the work that is being carried on by the Association. This work consists of the testing of seeds and fertilisers, under conditions as nearly similar as can be found practicable. The seeds are furnished by the Union free to the farmers, and full instructions regarding the mode of conducting the tests are also furnished at the same time. The only return asked of the farmer is a report of the results, to be sent after harvest by a time fixed upon as mentioned in the instruction sheet. These reports are made upon blank forms furnished to each experimenter along with the instructions.

At the present time there are no less than 1,642 plots under experiment, which are conducted by ex-students and other farmers

throughout the province. These plots comprise the following:—
70 with fertilisers, 196 with lucerne and corn, 350 with roots,
1,026 with spring wheats.

The results of this work, which are published annually, cannot fail to be of immense service to the farmers.

In the subjoined table will be found the different sets of varieties of wheats, which will be furnished by mail in half pound lots of each variety, to farmers applying for them, and in the order of the applications so long as the supply lasts.

FIVE SETS OF FALL WHEAT FOR CO-OPERATIVE TESTS

(1)	(2)	(3)	(4)	(5)
American Bronze. Canadian Vel- vet Chaff. Surprise. Manchester. Bulgarian.	American Bronze. Canadian Vel- vet Chaff. Garfield. Golden Cross. Standard.	American Bronze. Canadian Vel- vet Chaff. Bonnell. Early Red Clawson. Lancaster.	American Bronze. Canadian Vel- vet Chaff. Winter Pearl. Red Velvet Chaff. Hybrid Medi- terranean.	American Bronze. Canadian Velvet Chaff. Longberry Red. Jones' Winter Fyfe. Democrat.

Each farmer wishing one of these sets will please address to the Secretary, O. A. Zavitz, Experiment Station, Guelph, *mentioning which set he desires*, and the grain, instructions for testing and blank forms on which to report will be forwarded to his address.