Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1981

Tt
to

**The
po
of
fil**

O
be
th
si
of
fin
si
of

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☒ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refiled to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

**T
s
T
W**

M
d
e
b
r
i
r
e
m

The copy filmed here has been reproduced thanks to the generosity of:

Library,
National Museums of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol ➡ (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque
Musées Nationaux du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole ➡ signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ails
du
modifier
une
image

rrata
o

pelure,
n à

THE
RAY SOCIETY.

INSTITUTED MDCCCXLIV.

*This volume is issued to the Subscribers to the RAY SOCIETY for
the Year 1867.*

LONDON:

MDCCCLXVII.

THE
MISCELLANEOUS
BOTANICAL WORKS

OF

ROBERT BROWN, ESQ., D.C.L., F.R.S.,
FOREIGN ASSOCIATE OF THE ACADEMY OF SCIENCES OF THE
INSTITUTE OF FRANCE, ETC., ETC., ETC.

VOL. II.
CONTAINING
III. SYSTEMATIC MEMOIRS.
AND
IV. CONTRIBUTIONS TO SYSTEMATIC WORKS.

LIBRARY
NATIONAL MUSEUM
OF CANADA
LONDON:

PUBLISHED FOR THE RAY SOCIETY BY
ROBERT HARDWICKE, 192, PICCADILLY.

MDCCCLXVII.

003402

PRINTED BY J. E. ADLARD,
BARTHOLOMEW CLOSE.

PREFACE TO THE SECOND VOLUME.

(BY THE EDITOR.)

THE present volume contains, as announced in the Preface to the first, the Systematic Memoirs and Miscellaneous Descriptions of Plants; and to these have been added a number of short contributions to different publications, together with notices of the assistance acknowledged to have been afforded by the Author to various writers, wherever the nature of the assistance is distinctly pointed out.

In Pritzel's '*Thesaurus Literaturæ Botanicae*,' p. 35, at the end of the enumeration of Mr. Brown's works, it is stated, "*Robertus Brown dedit præterea descriptiones ultimi (an etiam tertii?) Fasciculi Plantarum Cryptogamicarum Dicksonianarum, necnon eas tertii voluminis Plant. Coromand. Roxburghii, et curavit Editionem alteram Horti Kewensis a classe XII ad XXIV.*" To all the works here mentioned there is no doubt that Mr. Brown contributed valuable assistance, but of different kinds and in different degrees. The fourth fasciculus of Dickson's '*Plantæ Cryptogamicæ*,' but not, I believe, the third, was largely indebted to him; but it would be difficult, if not impossible, to determine what portions of the text were actually furnished by him. To the publication of the third volume of Roxburgh's '*Coromandel Plants*' he gave a general superin-

tendence, but in dealing with the text confined himself to slight verbal corrections. One generic name, that of *Gynocardia*, was substituted by him for the barbarous name, *Chaulmoogra*, adopted by Roxburgh, and this alteration has met with general acquiescence. On the portion of the second edition of the 'Hortus Kewensis' printed after the death of Dryander, in 1810, he bestowed the same attention which had been devoted by Dryander to the earlier portion, and by Solander to the first edition, adding largely to the elucidation of certain Families and Genera. These contributions will be found extracted in the present volume.

In the list of Mr. Brown's works given by Pritzel there occur two: "No. 1365†, Of three species of the natural order Orchideæ, Lond. 1817, 4, 8 pp. 2 tab. col.;" and No. 1366†, Select Orchideæ, ib. 4, 2 pp." both extracted from 'The Journal of Science and the Arts,' which do not properly belong to Mr. Brown, but formed part of a series of papers on Cape *Orchideæ*, by Mr. Bellenden Ker. These, of course, have not been admitted into the present edition.

I have only, in addition, to return my best thanks to my friends, Mr. Carruthers, and the Rev. W. W. Newbould, for much valuable assistance in reading and comparing the proof-sheets; and to Mr. Carruthers, in particular, for the preparation of the elaborate Index of Names of Plants appended to the present volume.

JOHN J. BENNETT.

NOVEMBER 30TH, 1867.

CONTENTS.

PART III.

SYSTEMATIC MEMOIRS.

	PAGE
On the Natural Order of Plants called Proteaceæ	3—192
On the Asclepiadæ, a Natural Order of Plants separated from the Apocinæ of Jussieu	193—247
On Woodsia, a New Genus of Ferns	249—255
Observations on the Natural Family of Plants called Compositæ	257—318
Characters and Descriptions of Three New Species of Plants, found in China by Clarke Abel, Esq.	319—328
Characters and Description of Lyellia, a New Genus of Mosses, with Observations on the Section of the Order to which it belongs; and some Remarks on Leptostomum and Buxbaumia	329—351
Remarks on the Structure and Affinities of Cephalotus	353—359
Characters and Description of Limnanthes, a New Genus of Plants allied to Floerkea	361—364

PART IV.

CONTRIBUTIONS TO SYSTEMATIC WORKS.

Genera et Species Plantarum Cruciferarum, necnon Generis Cleome, quæ in Horto Kewensi coluntur	367—415
Genera et Species quædam Plantarum Leguminosarum, quæ in Horto Kewensi coluntur	417—447
Genera et Species quædam Plantarum Myrtacearum, quæ in Horto Kewensi coluntur	449—457

	PAGE
Genera et Species quædam Plantarum Compositarum, quæ in Horto Kewensi coluntur	459—470
Genera et Species Plantarum Orchidearum, quæ in Horto Kewensi coluntur	471—499
Genera et Species Plantarum e variis Familiis, quæ in Horto Kewensi coluntur	501—510
Extracts from Dr. Richardson's Botanical Appendix to the "Narrative of a Journey to the Polar Sea, by Captain Franklin;" with Addenda	511—527
Names of and Notes on Indian Plants; extracted from Dr. Wallich's "Numerical List"	529—538
Notes and Observations on Indian Plants; extracted from Dr. Wallich's "Plantæ Asiaticæ Rariores"	539—556
Extracts from "Plantæ Javanicæ Rariores"	557—666
Contributions to the "Botanical Magazine"	667—679
Contributions to the "Botanical Register"	681—699
Names, &c., of Plants contributed to various publications, between 1818 and 1826; and Names quoted from R. Brown's Herbarium in Mr. Bentham's "Flora Australiensis"	701—711
Short Contributions to various Works, and occasional Notices, Botanical and Geological	713—727
Index of Subjects to Vol. II	729—735
Index of Names of Plants to both Volumes	737—786

PAGE
 n Horto
 . 459—470
 Kewensi
 . 471—499
 Kewensi
 . 501—510
 ative of
 enda
 511—527
 ullich's
 529—538
 ullich's
 539—556
 557—606
 667—679
 681—699
 n 1818
 um in
 701—711
 tanical
 713—727
 729—735
 737—786

PART III.

SYSTEMATIC MEMOIRS.

we

ON THE
NATURAL ORDER OF PLANTS

CALLED

PROTEACEÆ.

BY

ROBERT BROWN, LIB. L.S.

(READ JAN. 17TH, 1809.)

[*Extracted from the 'Transactions' of the Linnean Society of London,
Vol. X, pp. 15—226.*]

LONDON.

1810.

ON THE
PROTEACEÆ OF JUSSIEU.

THE Linnean system of botany, though confessedly ¹¹⁵ artificial, has not only contributed more than all others to facilitate the knowledge of species, but, by constantly directing the attention to those essential parts of the flower on which it is founded, has made us acquainted with more of their important modifications than we probably should have known, had it not been generally adopted, and has thus laid a more solid foundation for the establishment of a natural arrangement, the superior importance of which no one has been more fully impressed with than Linnæus himself.

There are still, however, certain circumstances respecting the stamina and pistilla, which appear to me to have been much less attended to than they deserve, both by Linnæus and succeeding botanists. What I chiefly allude to is the state of these organs before the expansion of the flower. The utility of ascertaining the internal condition of the ovarium before fœcundation will hardly be called in question, now that the immortal works of Gærtner and Jussieu have demonstrated the necessity of minutely studying the fruits of plants in attempting to arrange them according to the sum of their affinities, as in many cases the true nature of the ripe fruit, especially with respect to the placentation of the seeds, can only be determined by this means. Its importance is indeed expressly inculcated by many botanists, who, however, have frequently neglected it in practice: ¹¹⁶ nor do I find any one who has steadily kept it in view, except Aubert Du Petit-Thouars in his excellent work on

the plants of Madagascar and the Isles of France and Bourbon.

The bursting of the antheræ has, it is true, been generally observed, and many of its most unusual modes have been introduced into the characters of genera; but the examination of these organs, at a still earlier period, has been universally neglected; and hence the very imperfect knowledge which, even now, is possessed of their real nature in two of the most remarkable families of plants, the Orchideæ and Asclepiadeæ.

Examples of the great advantage of observing the antheræ in this early stage will hereafter be given in my general remarks on the order which is the proper subject of this essay. But I trust I shall be pardoned for here introducing some account of their structure in Asclepiadeæ, as it will enable me not only to bring forward the most striking proof of the importance of this consideration with which I am acquainted, but also, as I apprehend, to decide a question which has long occupied, and continues to divide, the most celebrated botanists.

The point in dispute is whether this order, comprehending *Asclepias*, *Cynanchum*, *Pergularia*, *Stapelia*, and several genera, at present confounded with these, ought to be referred to Pentandria or Gynandria, and, if to the latter, whether the antheræ are to be considered as five or ten; all of which opinions have had advocates of the greatest name in the science.

According to Linnæus, Jussieu, and Richard, they belong to Pentandria.

Linnæus has assigned no reason for his opinion, which, however, it appears he retained after he became acquainted with the observations of Jacquin and Rottboell; but it is ¹⁷³ probable he was induced to adopt it more from the consideration of the close analogy these plants have with the manifestly pentandrous Apocineæ, than from regarding them as strictly referable to this class; for, in his natural generic characters of *Asclepias* and *Pergularia*, he very clearly describes both these genera as gynandrous.

Jussieu has entered more fully into the subject, but seems

also to have been chiefly guided by this analogy and the observations of others; as he concludes by expressing his doubts, respecting both the origin and use of the parts.

Richard, whose description of these organs I find in Persoon's synopsis, has indeed come nearer to the solution of the question; his account, however, of the origin of the lateral processes hereafter mentioned, proves that this description was not altogether formed on actual observation.

Jacquin, the first botanist that submitted these plants to minute examination, and whose figures well illustrate most points of their structure, has adopted a very different opinion, referring them to Gynandria, in which he is followed by Koelreuter, Rottboell, and Cavanilles, all of whom likewise agree with him in considering them as decandrous; while Dr. Smith, in his late valuable Introduction to Botany, who conceives that "no plants can be more truly gynandrous," regards them as having only five antheræ. And lastly Desfontaines supposes the five glands of the stigma to be the true antheræ, considering the attached masses of pollen as mere appendages to these.

All the authors who thus refer them to Gynandria seem quite confident in the justness of their views; and yet the inspection of a single flower bud overturns, as it appears to me, with irresistible evidence, the conclusion they had formed from premises apparently so satisfactory.

My attention, while in New Holland, having been much engaged by the plants of this family, the species in that continent being both numerous and with difficulty reducible to established genera: I there observed the following facts concerning them, all of which I have, since my return to England, confirmed by the examination of different species of the same tribe.

The observations of Jacquin on this subject being generally known, it must be unnecessary to enter into a minute description of those organs which are well exhibited by his figures in every respect, except as to the origin of the supposed antheræ.

If a flower bud of any plant of this family, while scarcely

half the size it attains immediately before expansion, be carefully examined, it will be found that the polleniferous sacs, as they are termed by Jacquin and his followers, in which they suppose the antheræ to be merely immersed, are really the organs by which the fœcundating matter is secreted: for at this period they are perfectly closed, and consequently all communication cut off between the stigma and their contents now consisting of a turbid fluid or pulpy mass.* If the stigma be at the same time observed, the gland-like bodies which originate in its grooved angles are already visible; but, instead of having the cartilaginous or horny texture which they at length acquire, are as yet semi-fluid, and of hardly a determinate form. Near the base of each side of these grooves a more superficial depression is observable, which, though in some cases extremely short, is in others of considerable length, and generally forms a right angle with the corresponding groove. In these depressions, the processes by which, at a more advanced stage, the contents of the antheræ are connected with the stigma, are immersed, and at this period they are found to be semi-fluid. By degrees the glands, as well as their lateral processes, acquire a firmer consistence, and the inferior or outer extremity of each of the processes, being extended beyond its depression or furrow, on the bursting of the opposite cell of the corresponding anthera, firmly attaches itself to its contents, now become a regular mass of a waxy consistence.

If the accuracy of this statement be admitted, it will probably be allowed that the *Asclepiadeæ* cannot be regarded as gynandrous, especially in the sense in which they are so considered by botanists; but lest it should not be thought completely satisfactory, it may be added, that in a still earlier stage of the flower bud I have found the fœcundating matter already secreted in the cells of the antheræ, while the glands of the stigma, as well as their processes, were absolutely invisible.

As to the question of their being pentandrous or decandrous, every analogy must lead us to refer them to the former class; nor, indeed, have they, when not considered

as gynandrous, been ever supposed to belong to Decandria.

An œconomy, in many respects similar to that now described, obtains also in Orchideæ; in which, however, the processes connecting the antheræ with the stigma, where they exist, are in many cases derived from the masses of pollen themselves; but in others they as certainly originate from the stigma, or its glandular appendage.

The result of my examination of these two interesting orders of plants I hope hereafter to submit to the Society; and I now proceed to the proper subject of the present paper.

The natural order of PROTEÆ, or, as it is less exceptionably called, PROTEACEÆ, was first established in the Genera Plantarum of the celebrated Jussieu; and the description there prefixed to it will, with a few alterations, still apply to the order, now that it has received so many additions, not only in species, but in very distinct genera, several of which were first published by Dr. Smith, in the 4th vol. of the Society's Transactions, and others are in the present paper submitted to the consideration of botanists.

The general description and definition of the order will be most advantageously placed at the head of its systematic arrangement; before entering upon which, I shall offer some remarks on its geographical distribution, and likewise on such modifications of structure in the different organs as appear to be of the greatest importance in indicating or characterising genera.

The geography of plants being as yet in its infancy, the smallest addition to our knowledge of a subject which promises to become of considerable importance, will probably be received with indulgence; and in this persuasion I venture to make the following observations on the order before us. In the first place, it is remarkable that the PROTEACEÆ are almost entirely confined to the southern hemisphere. This observation originated with Mr. Dryander, and the few exceptions hitherto known to it, occur considerably within the tropic. The fact is the more

deserving of notice, as their diffusion is very extensive in the southern hemisphere, not merely in latitude and longitude, but also in elevation; for they are not only found to exist in all the great southern continents, but seem to be generally, though very unequally, spread over their different regions: they have been observed also in the larger islands of New Zealand and New Caledonia; but hitherto neither in any of the lesser ones, nor in Madagascar. As in America, they have been found in Terra del Fuego, in Chili, Peru, and even Guiana, it is reasonable to conclude that the intermediate regions are not entirely destitute of them. But with respect to this continent, it may be observed, that the number of species seems to be comparatively small, their organization but little varied; and further, that they have a much greater affinity with those²¹ of New Holland than of Africa.

Of the botany of South Africa, scarce anything is known, except that of the Cape of Good Hope, where this family occurs in the greatest abundance and variety; but even from the single fact of a genuine species of *Protea* having been found in Abyssinia by Bruce, it may be presumed, that in some degree they are also spread over this continent.

With the shores, at least, of New Holland, under which I include Van Diemen's Island, we are now somewhat better acquainted, and in every known part of these, Proteaceæ have been met with.

But it appears that, both in Africa and New Holland, the great mass of the order exists about the latitude of the Cape of Good Hope; in which parallel it forms a striking feature in the vegetation of both continents.

What I am about to advance respecting the probable distribution of this family in New Holland, must be very cautiously received; as it is in fact chiefly deduced from the remarks I have myself made in Captain Flinders's Voyage, and subsequently during my short stay in the settlements of New South Wales and Van Diemen's Island, aided by what was long ago ascertained by Sir Joseph Banks, and by a very transitory inspection of an herbarium collected on the west coast, chiefly in the neighbourhood

of Shark's Bay, by the botanists attached to the expedition of Captain Baudin.

From knowledge so acquired I am inclined to hazard the following observations.

The mass of the order, though extending through the whole of the parallel already mentioned, is by no means equal in every part of it; but on the south-west coast forms a more decided feature in the vegetation of the country, and contains a far greater number of species than on the east:—and in that part of the south coast, which was first examined by Captain Flinders, it seems to be more scanty than at either of the extremes.

On the west coast also, the species upon the whole are more similar to those of Africa than on the east, where they bear a somewhat greater resemblance to the American portion of the order.

From the parallel of the mass, the order diminishes in both directions; but the diminution towards the north is probably more rapid on the east than on the west coast.

Within the tropic, on the east coast, no genera have hitherto been observed, which are not also found beyond it; unless that section of *Grevillea*, which I have called *Cycloptera*, be considered as a genus. Whereas at the southern limit of the order several genera make their appearance, which do not occur in its chief parallel.

The most numerous genera are also the most widely diffused. Thus *Grevillea*, *Hakea*, *Banksia*, and *Persoonia*, extensive in species in the order in which they are here mentioned, are spread nearly in the same proportion; and they are likewise the only genera that have as yet been observed within the tropic.

Of such of the remaining genera, as consist of several species, some, as *Isopogon*, *Petrophila*, *Conospermum*, and *Lambertia*, are found in every part of the principal parallel, but hardly exist beyond it. Others, as *Josephia*¹ and *Synaphea*, equally limited to this parallel, have been observed only towards its western extremity; while *Embothrium*

¹ Subsequently in this paper altered to *Dryandra*. Confer *Prodr. Fl. Nov. Holl.* 1, p. 396.—EDIT.

(comprehending for the present under this name all the many-seeded plants of the order),¹ which is chiefly found on the east coast, and makes very little progress towards the west, advances to the utmost limit of south latitude, and there ascends to the summits of the highest mountains.

^{23]} Genera consisting of one or very few species, and which exhibit generally the most remarkable deviations from the usual structure of the order, are the most local, and are found either in the principal parallel, or in the highest latitude.

The range of species in the whole of the order seems to be very limited; and the few cases which may be considered as exceptions to this, occur in the most extensive genera, and in such of their species as are most strictly natives of the shores. Thus *Banksia integrifolia*, which grows more within the influence of the sea than any plant of the order, is probably also the most widely extended, at least in one direction, being found within the tropic, and in as high a latitude as 40°. It is remarkable, however, that with so considerable a range in latitude, its extension in longitude is comparatively small: and it is still more worthy of notice, that no species of this family has been found common to the eastern and western shores of New Holland.

The celebrated traveller Humboldt is the first who has expressly pointed out a remarkable difference in the distribution of the species of plants.

He observes that, while the greater number grow irregularly scattered and mixed with each other, there are some which form considerable masses, or even extensive tracts, to the nearly absolute exclusion of other species. Of plants growing thus in society, the greater number occur in the temperate zones; and of these, the most decided instances will readily present themselves to every botanist. I venture to add, that such as exist within the tropic, are found, either at considerable heights or on the sea-shores.

To this class very few of the Proteaceæ can be said to

¹ In the systematic part of the present Memoir, the Australian species of *Embothrium* are separated under the generic names of *Knightia*, *Lomatia*, and *Stenocarpus*.—EDIT.

belong. *Protea argentea* of Linnæus is the most striking example among the African species; and my friend Mr. ^[24] Ferdinand Bauer has observed a similar tendency in *Protea mellifera*.

Among the New Holland species, *Banksia speciosa* is the sole instance, and even that only in certain circumstances, of this manner of growth.

The favourite station of Proteacæ is in dry stony exposed places, especially near the shores, where they occur also, though more rarely, in loose sand. Scarcely any of them require shelter, and none a good soil. A few are found in wet bogs, or even in shallow pools of fresh water; and one, the *Embothrium ferrugineum* of Cavanilles, grows, according to him, in salt marshes.

Respecting the height to which plants of this order ascend, a few facts are already known. The authors of the Flora Peruviana mention, in general terms, several species as being alpine; and Humboldt, in his valuable Chart of Æquinoctial Botany, has given the mean height of *Embothrium emarginatum* about 9300 feet, assigning it a range of only 300 feet. On the summits of the mountains of Van Diemen's Island, in about 43° south latitude, at the computed height of about 4000 feet, I have found species of *Embothrium*, as well as other genera hitherto observed in no other situation. *Embothrium*, however, as it is the most southern genus of any extent, so it is also, as might have been presumed, the most alpine of the family.

Two genera only of this order are found in more than one continent: *Rhopala*, the most northern genus, which, though chiefly occurring in America, is to be met with also in Cochin-China and in the Malay Archipelago; and *Embothrium*, the most southern genus of any extent, is common to New Holland and America.

From this account of the geographical distribution of ^[25] the Proteacæ, I proceed to make some general remarks on the structure and modifications of their different parts. The order, which consists of shrubs of the most rigid nature, or of trees of moderate size, contains also one herbaceous plant, my *Symphionema paludosum*, which, how-

all the
y found
ards the
ude, and
ains.

ies, and
ons from
and are
latitude.
seems to
be con-
most ex-
are most
egrifolia,
than any
idely ex-
within the
markable,
itude, its
and it is
is family
rn shores

who has
ne distri-

row irre-
are some
e tracts,
Of plants
r in the
instances
venture
ad, either

said to

species of
omatia, and

ever, except in this respect and in the union of the tops of its filaments, does not remarkably differ from the usual structure of the family.

The pubescence, which is very general in the order, consists either of a short and in many cases nearly impalpable tomentum, or of soft hairs which are either spreading, close pressed, or somewhat crisped, generally simple, but in some genera fixed by the middle, and in a very few cases glandular.

The existence or absence of pubescence in the adult leaves cannot always be depended upon in distinguishing species; but the short tomentum, especially of their under surface, is of greater consequence than the spreading hairs. In the bractæ more reliance may be placed on it, and in the different parts of the flower I have never hesitated to employ it in my specific characters. In the calyx I have even derived the greatest advantage in some difficult genera, especially *Serruria*, from attending to its differences in direction.

Mr. Salisbury has introduced the pubescence of fruit into several of his generic characters, and in some I think with evident advantage, but in such only as where from its abundance and length it performs a function of manifest importance in assisting dissemination: hence I conceive it may be safely admitted into the characters of *Protea* and *Isopogon*; but I can perceive no advantage whatever in employing it in those of *Serruria* and *Spatalla*. For this reason too it ought not to be used in the capsular or drupaceous genera, in which indeed experience proves it to be of no further moment than in distinguishing species.

Dr. Smith has given it as his opinion, that from the disposition of leaves in New Holland plants no conclusion can safely be drawn as to their genera. This remark however appears to me only applicable to certain families, or rather genera; for in many tribes the plants of that country are altogether as constant in their leaves as in any other part of the world. In proof of this, it may be sufficient to mention the order Rubiaceæ; and there are many others in which I find nothing at all remarkable in this respect.

As to Proteaceæ, it must be acknowledged that in *Banksia* both verticillated and scattered leaves occur; but the leaves constantly in threes in *Lambertia* seems to me a circumstance of even greater importance than the number of flowers in the involucre; and the opposite leaves of *Xylomelum* distinguish it at once both from *Rhopala* and *Hakea*.

Although the form and divisions of leaves in the order are variable in no common degree, yet there are certain genera, both among those of Africa and New Holland, which the leaves even in these respects assist in indicating. Thus, in that genus to which I have applied the name of *Protea* (the *Erodendrum* of Mr. Salisbury), and I believe also in my *Leucadendron*, there is no instance of a divided or toothed leaf; thus also the leaves of *Spatalla* are filiform and undivided, and those of *Serruria* filiform and almost always pinnatifid. Their dichotomous divisions in *Simsia* and *Franklandia* are still more characteristic; and their division and remarkable reticulation readily distinguish *Synaphea* from *Conospermum*.

The inflorescence in Proteaceæ, whatever use botanists may think proper to make of it in their generic characters, is of undoubted importance in determining genera, and even in the primary division of the order it appears to be of nearly equal consequence with the fruit itself; for, in dividing the order into two sections from the structure of the ovary, it will be found that while all the single-seeded genera have each flower subtended by a proper bractea, or more rarely are without one, those with two or more seeds have, with very few exceptions, the flowers of their spikes or clusters disposed in pairs, each pair being furnished with only one bractea common to both flowers: it may also be observed that all the American and two thirds of the New Holland species have this mode of inflorescence, while only one instance of it occurs in Africa.

The single envelope of the stamina and pistillum in Proteaceæ I have, with Jussieu, denominated CALYX, chiefly because the stamina, of equal number with its laciniae, are constantly opposite to them, and from the close analogy

subsisting between this family and that of Thymeleæ, in which I believe the greater number of botanists will allow that this envelope is really calyx: and as this latter argument may be considered as the stronger, I shall endeavour to establish the identity of this organ in these two families. In several of the Thymeleæ, especially in *Pimelea*, the lower part of the tube of the calyx is, as it were, jointed with the upper; after the falling off of which, it remains surrounding the fruit: this is also the case in several genera of Proteaceæ, as in *Adenanthos* of Labillardière, in *Isopogon*, in *Grevillea* *Chrysodendron*, and still more remarkably in *Franklandia*, in which the persistent tube becomes indurated and even nearly woody, a change surely not likely to take place in a genuine corolla. But though I have thus adopted the language of Jussieu, I am decidedly of opinion that, in all families having a single ^{28]} envelope, it will be still better to call it perianthium or perigonium, which latter term was proposed by Ehrhart, and is adopted by Decandolle.

A circumstance meriting the attention of the theoretical botanist, respecting the calyx in this order, is its invariable division into four leaves or segments; for the single exception noted by Linnæus in his description of the male flowers of *Brabejum*, he himself seems afterwards to have distrusted, from the manner in which he has introduced it into the amended generic character given in the Mantissa; and I may add, that in nearly 400 species of the order, which I have examined, I have not met with a single exception to this rule.

With this uncommon constancy in point of number, it is remarkable that there is, in the whole order, a strong tendency to irregularity in form, the various kinds of which are of great importance in characterising genera.

Before the expansion of the calyx the margins of its segments are applied to each other; and from the unequal degrees of cohesion in many cases subsisting among them after expansion, several kinds of irregularity arise. I am not sure that any term has been contrived for this manner of æstivation, except it be the *æstivatio valvata* of Linnæus;

but as he has not defined it, and as his commentator Reuss has given the very different æstivation of grasses as an example, I have, in introducing this circumstance into the general description of the order, specified it at length.

From the colour of the calyx, many genera of Proteaceæ are indicated with tolerable certainty. Thus *Synaphea* is distinguished from *Conospermum* by its yellow flowers; and no instance of yellow flowers has been met with in the numerous genera *Serruria* and *Spatalla*, nor any of purple in *Leucadendron*. In some genera, however, as in *Banksia*,^[29] and *Isopogon*, it is evidently of very little importance.

The fleshy or scale-like bodies, which surround the ovarium in the greater number of plants of this family, are in many cases so manifestly secreting organs, that it is surprising Mr. Salisbury should hesitate in considering them as nectaria, and denominate them calli; a term which excludes the idea of secretion. But whatever their functions may be, great assistance may certainly be derived from their various modifications, in distinguishing genera. Their importance however in this respect, like that of all other parts, not only in this, but, as I apprehend, in every natural family, is very unequal, and in some cases seems to be entirely lost. Thus, in the genus *Leucadendron* as it is here constituted, they are wanting in several species, and in some I am inclined to think exist only in the males.

In most of the regular-flowered genera they are four in number, and alternate with the leaves or laciniae of the calyx. In these genera they are also generally in the form of succulent scales, distinct, or more rarely cohering at their base, and in a very few instances adhering to the calyx; but in *Persoonia* they are nearly round and fleshy, and in *Bellendena*, *Symphionema*, *Simsia*, *Agastachya*, *Petrophila*, and *Isopogon*, they are entirely wanting.

In the irregular-flowered genera with two or many seeds their number is less than four, in most cases only one exists, in a few others three, and in some none.

Varieties in the structure or apparent origin of the STAMINA, afford, as might be expected, important generic characters. Their usual insertion in the order is in the

concave tops of the laciniae of the calyx; all considerable deviations from which may safely be employed in characterising genera. In this way *Rhopala*, *Xylomelum*, and *Lambertia* are readily distinguished from *Embothrium*, *Grevillea*, and *Hakea*; and thus also *Persoonia* and *Brabejum* remarkably differ from *Guevina*; while *Bellendena* differs from all others in having its stamina distinct from the calyx, affording however an indication of the real origin of these organs in the whole family.

The deviations from the usual structure of ANTHERÆ in this order are not many; but some of them are of so singular a nature as to constitute the essential characters of the genera in which they take place. These genera are *Simsia*, *Conospermum*, and *Synaphea*, all of which are most truly syngenesious; for not only do their antheræ firmly cohere together, but the corresponding lobes of these being, when considered separately, entirely open, are so applied to each other as to form but one cell, without a trace of any intermediate membrane. In *Simsia* the four antheræ are perfect, each consisting, as in the rest of the order, of two lobes, and therefore the whole before bursting constitute four cells. Whereas in *Conospermum* and *Synaphea* one filament is entirely barren, the two lateral ones have each a single-lobed anthera, and the fourth alone is perfect: hence before bursting the whole form only two cells.

This remarkable structure, which can only be ascertained before the opening of the calyx, necessarily escaped Dr. Smith in describing his *Conospermum*, for I conclude he had only the expanded flower before him, and the appearance of the antheræ in this state after their separation justifies him in referring the genus to Tetrandria: but according to the view now given of its structure, it can have no other pretension to a place in this class than its belonging to Proteaceæ; and the order Syngenesia Monogamia being abolished, it must be referred to Triandria.

The only remaining anomaly in these parts occurs in ³⁰¹ *Franklandia*, and consists in the anthera, or rather that portion of the filament on which it is fixed, adhering to the calyx through its whole length.

considerable
in charac-
Xylomelum,
Embothrium,
and *Bra-*
Bellendena
distinct from
real origin

ANTHERÆ in
e of so sin-
acters of the
are *Simsia*,
most truly
rmly cohere
eing, when
died to each
e of any in-
antheræ are
rder, of two
g constitute
naphea one
have each a
fect: hence

y be ascer-
ily escaped
I conclude
n, and the
heir separa-
ndria: but
ure, it can
ass than its
esia Mono-
iandria.

occurs in
rather that
adhering to

The figure of the POLLEN has been attended to by a few theoretical, but by hardly any practical botanists; yet I am inclined to think, not only from its consideration in this family, but in many others, that it may be consulted with advantage in fixing our notions of the limits of genera: and though its minuteness may perhaps always exclude it from a place in generic characters, yet it well deserves, to use the words of Linnæus when speaking of habit, to be "occulte consulendus."

Its usual figure in the order is triangular with secreting angles, a beautiful contrivance for insuring impregnation in a tribe, in which, from the very scanty, or in many cases apparent want of secretion by the stigma, it must otherwise have been very uncertain; for by this form and secretion, as well as by the singular œconomy of the calyx, it remains so long in contact with the stigma, as probably to compensate for the somewhat defective structure of that organ.

From this figure the principal deviation is in the extensive genera *Banksia* and *Josephia*, in all of which it is elliptical or oblong, and either straight or bent into a semilunar form; and in *Franklandia* and *Aulax*, where it is spherical. The only remaining exception with which I am acquainted is the original *Embothrium* of Forster, his *E. coccineum*, in which, as in *Banksia*, it is oblong; a circumstance that, together with the more important character of a regular club-shaped stigma, and some other differences, has determined me to separate it from all the other species of *Embothrium*, except *E. lanceolatum* of Flora Peruviana, whose pollen however remains to be examined.

The external modifications of the OVARIUM must be very cautiously used in the generic characters of this family; even its being sessile or pedicellated is not always of sufficient importance, though I think Mr. Salisbury has done well in introducing it into his characters of *Serruria* and *Spatalla*, in both which genera I had overlooked it before the publication of his Essay.

Its internal structure, which ought always to be ascertained, will be found of the greatest importance in most

cases, but fails in *Persoonia*, the species of which differ in having one or two seeds: it would seem however, in this case, that an irregularity in a point of such importance could not take place unaccompanied with other anomalies in the same organ, and accordingly such are found to exist in this genus, and will be mentioned when treating of the fruit.

Besides number, the insertion of the ovula is also to be attended to; for though this may generally be presumed from the situation of the radicula in the ripe seed, yet to this criterion there are several exceptions, even in the present order: thus, while the radicula constantly points downward in the whole of the order, the insertion of the ovulum is in many cases at the top or side of the cell of the ovarium. My observations on this subject are as yet incomplete; but, from those that I have made, I am inclined to think such differences will be connected with genera, or rather perhaps with particular kinds of fruit. Thus I conjecture, in *Leucospermum*, *Mimetes*, *Nivenia*, and *Spatalla*, the insertion to be uniformly lateral.

The STYLE, though not subject to much variety in this family, will be found in a few cases to furnish generic characters. Thus in *Protea*, strictly so called, the persistent subulate style forms an important part of its character: and the persistency of the whole of the style in the greater number of species of *Grevillea* will probably be used by ^{33]} future botanists in distinguishing them from that remarkable section of the genus, which I have at present united with them and called *Cycloptera*. Its length also, when compared with that of the calyx, seems in some cases to be of importance, as in distinguishing *Adenanthos* from *Spatalla*; but in general this circumstance can hardly be had recourse to except in specific characters.

The form of the STIGMA is in many cases of considerable importance in characterising genera, a fact which could not escape the penetration of Dr. Smith when establishing his new genera of this order: thus its conical papilla in his *Conchium* (the *Hakea* of Schrader) will in many, though certainly not in all cases, distinguish it from *Grevillea*: but

its form in both these genera will readily serve to separate them from *Xylomelum* and *Rhopala*; and thus also *Spatalla* remarkably differs from *Adenanthos*. Upon the whole, however, it seems that its obliquity is of greater importance than its form; for this, when existing in any great degree, is generally accompanied with a corresponding irregularity in the calyx: but as this irregularity is produced for the purpose of bringing all the antheræ into contact with the stigma, so its obliquity in the dioiceous genera *Leucadendron* and *Aulax* is not attended with so great a degree of irregularity, which would here serve no end, impregnation depending on the pollen of different individuals, to insure which the surface of the stigma in these genera is rough with papulæ; a circumstance that, together with its form, readily distinguishes them from all others of the order.

In *Synaphea*, the stigma or summit of the style inoscillates with the divisions of the barren filament, which in some species appear beyond it in horn-like processes, but in others are entirely lost in its substance. I am acquainted with nothing like this in the whole vegetable kingdom; and such a singularity alone, when occurring in several species, would have determined me to separate these plants from *Conospermum*: but being also accompanied by other remarkable differences, both of structure and appearance, no genus, I apprehend, can be better founded than this.

That the opinion of Christian Knaut and Vaillant respecting the non-existence of naked seeds is correct when anatomically considered, there can be no doubt; but the practical utility of deviating in this subject from the common language of botanists may still be questioned: and accordingly Gærtner, who was fully aware of the truth of their position, has nevertheless continued to describe the seeds of many plants as naked. I confess however I am inclined to adopt the opposite decision of the French botanists, at the head of whom is Richard, who has also proposed terms for distinguishing the various species hitherto confounded under the name of naked seeds. The fruit of the monospermous genera of Proteaceæ might probably be with

advantage referred to that which he has termed *Akenium*; but as I am unwilling in the present paper to adopt any term not more generally sanctioned and understood than this, I shall content myself with calling those *nucæ*, which are either not at all or but slightly compressed and not bordered; and apply the term *samara* to such as are either very much compressed, or with a less remarkable compression are surrounded or terminated by a membranaceous border: that I regard these distinctions however as in some cases of very little importance, may be inferred from this, that my genus *Leucadendron* includes both these kinds of fruit.

The first observation I have to offer on the fruits of Proteaceæ is, that there is no really bivalvular capsule in the order; a truth which was not perceived by Gärtner in ^{35]} describing his *Banksia dactyloides* (the *Conchium dactyloides* of Dr. Smith), and which has equally escaped Cavanilles and Labillardière in their characters of *Hakea*. Dr. Smith has more cautiously omitted this consideration in his character of that genus, and Professor Schrader has accurately described the suture as only existing on one side: such fruits then are as truly folliculi as those of *Grevillea*, *Rhopala*, or *Embothrium*; and that the existence of a distinct placenta is by no means necessary to constitute this kind of fruit, is proved even by some genera of Apocineæ, to which family this term was first applied.

A circumstance occurs in some species of *Persoonia* to which I have met with nothing similar in any other plant: the ovarium in this genus, whether it contain one or two ovula, has never more than one cell; but in several of the two-seeded species a cellular substance is after fecundation interposed between the ovula; and this gradually indurating acquires in the ripe fruit the same consistence as the putamen itself, from whose substance it cannot be distinguished; and thus a fruit originally of one cell becomes bilocular: the cells however are not parallel, as in all those cases where they exist in the unimpregnated ovarium, but diverge more or less upwards.

In all the seeds of this order there is a very manifest

CHALAZA, which, whatever may be the point of insertion of the seed, is always situated at its upper extremity ; and I have not been able to observe any fasciculus of vessels connecting it with the umbilicus in cases where this latter is placed in a different part of the seed.

I am not aware of any function being ascribed to the CHALAZA of seeds, except the nutrition of their proper membrane : but it appears to me too remarkable a part to be destined for this purpose only ; and some observations I have made induce me to suppose that it is the organ ³⁶ secreting the liquor amnios. This opinion I was first led to form by observing in some species of *Persoonia*, in which the inspissated remains of this fluid are visible in the ripe fruit, that it evidently originated in the *chalaza* and continued to adhere to it : nothing has hitherto occurred to invalidate this opinion, which is here however hazarded merely as a conjecture, requiring for its confirmation more numerous and decisive facts than I can at present adduce.

That the ALBUMEN of seeds is merely that condensed portion of the liquor amnios which remains unabsorbed by the embryo, seems to me very satisfactorily established ; and as this fluid is in the early stage never wanting, all seeds may in one sense be said to have albumen : but while in some tribes this unabsorbed part in the ripe seed many times exceeds the size of the embryo, so there are others in which not a vestige of it remains ; and such has hitherto been supposed to be the case with Proteaceæ : nor are the few exceptions with which I am at present acquainted of so decisive a nature as to invalidate this character of the order ; for they occur only in some species of *Persoonia*, where the semi-fluid remains of this substance are observable between the cotyledons ; and in *Bellendena*, in which it continues to form a thin fleshy coat on the inner surface of the proper membrane of the seed. From such instances, however, we may expect to find plants with a more copious albumen, which nevertheless it may be necessary from the whole of their organisation to refer to this family.

The RADICULA pointing towards the base of the fruit in

all Proteaceæ is a circumstance of the greatest importance in distinguishing the order from the most nearly related tribes; and its constancy is more remarkable, as it is not accompanied by the usual position or even uniformity in the situation of the external umbilicus.

^{37]} If Gärtner had not described the PLUMULA of *Protea argentea*, I should not have hesitated to assert that it was inconspicuous in the whole order.

The number of COTYLEDONS when more than two is a circumstance of little importance. In *Persoonia*, the only genus of the order in which a plurality of cotyledons has been observed, I am not even certain that their number is constant in those species in which this anomaly occurs.

In the following part of this essay it may be observed, that the genera into which I have subdivided the great African family *Protea*, are in most cases similar to those already proposed by Mr. Salisbury in the *Paradisus Londinensis*: from that essay however they are certainly not derived, but before its publication were formed and submitted to the judgment of Mr. Dryander, at whose suggestion they are now offered to the Society. That the results of an examination conducted by two observers wholly independent of each other, are so similar, will probably be considered as some proof of their correctness.

As Mr. Salisbury's generic names have the unquestionable right of priority of publication, I have in most cases adopted them, though I wish some of them had been differently constructed. But as I cannot accede to his application of the Linnean names *Protea* and *Leucadendron*, I shall here, that I may not disturb the following arrangement, assign my reasons for differing from him in this respect; and as in so doing I am obliged to trace the progress of Linnæus's knowledge of the family, I persuade myself that this will in some degree compensate for the otherwise unwarrantable length of the discussion.

The name PROTEA, which originated with Linnæus, first occurs in the folio edition of his *Systema Naturæ* published in 1735; no generic characters are there given,

but from the references to Boerhaave's figures it is evident that the genus is to be understood in the same extensive sense which he at length gave it in the second Mantissa. In 1737 appeared the *Genera Plantarum*, and in it for the first time the natural generic character of *Protea*: as in this work he only cites *Lepidocarpodendron* and *Hypophyllocarpodendron* of Boerhaave, it follows that here the genus is more limited, though its character is not peculiarly applicable to either of Boerhaave's genera referred to; and the description of antheræ and germen is not reconcilable to any plant whatever of the family. In the same year *Hortus Cliffortianus* was published, in which he resumes his first opinion of *Protea*, reducing to it all Boerhaave's genera, but referring to the character given in his own *Genera Plantarum*. It does not appear on what ground this change of opinion was formed; for in Clifford's garden, according to *Viridarium Cliffortianum*, there had only been two species, *Protea argentea* and *saligna*, neither of which had flowered, and the former was already lost; while in his Herbarium, now in the collection of Sir Joseph Banks, the specimens of all the three species given in the body of the work are without fructification, and of *Protea racemosa* added in the appendix there is no specimen whatever.

If Linnæus is to be considered in a great degree the author of the *Prodromus Floræ Leydensis*, published by A. Van Royen in 1740, as has been asserted by some of his pupils, and may be inferred from a passage in his Diary published by Dr. Maton, it must be noticed as his next work in the order of time; for from the same Diary it appears that he could only have been employed in its composition in 1738. In this work the genus *Protea* is given in the same extensive sense as in *Hortus Cliffortianus*, and no fewer than twenty-one species are characterised, of which however only two were in the Leyden garden, the rest being described from specimens in Van Royen's Herbarium.

In 1738 he also published his *Classes Plantarum*, in which, notwithstanding he appears to have composed it

while engaged in the arrangement of Van Royen's collection, another fluctuation of opinion occurs, *Protea* being limited as in the first edition of the *Genera Plantarum*, and to *Leucadendros*, which here for the first time occurs, he refers the *Conocarpodendron* of Boerhaave.

In 1740 he published the second edition of *Systema Naturæ*, where the names *Protea* and *Leucadendron* are both given; but the references to Boerhaave are reversed, *Protea* being confined to his *Conocarpodendron*, and *Leucadendron* comprehending his other two genera. In this sense they also appear in the second edition of the *Genera Plantarum* published in 1742, in which the character of *Leucadendron* is first given, some of whose species he must, from the annexed asterisk, have seen recent: his description of corolla and pistillum is only applicable to *Lepidocarpodendron*.

In 1745 Linnæus received the Herbarium of Hermann, from which he composed his *Flora Zeylanica*: the fourth volume of this collection containing a mixture of Ceylon and African plants, the latter are not noticed in this work; but from an inspection of the Herbarium itself, now in the Banksian collection, it appears that he had added generic names to most of them: of *Proteæ* only three species exist in the volume, of which *Protea conocarpa* is one: of this there are on the same page two specimens, whose heads of flowers are separately pasted; under one of these specimens he has written *Leucadendron*, and under the second *Protea*; to a specimen of *Protea Serraria* on a different ^{40]} page he has given the name of *Santolina*. These facts are mentioned to prove, that at this period his knowledge of the family must have been chiefly derived from Boerhaave's figures, and perhaps from specimens which he had casually seen:

In 1748 the sixth edition of *Systema Naturæ* appeared, where the essential characters of *Protea* and *Leucadendron* first occur, both of them evidently derived from the natural characters previously given.

In 1753 the *Species Plantarum*, the most accurate of all his works, was given to the world; both genera are found

in it, their species characterised, and trivial names for the first time applied to them: c^t Protea there are only two species, *P. argentea* and *fusca*; to the former however he referred as varieties *P. saligna*, *conifera*, and three others; to the whole adding the following observation, which may be supposed to contain his chief reason for applying his name Protea to this genus rather than to that for which in his Classes Plantarum he had first intended it. "*Planta naturalis in patria argentea excellit fronde inter arbores nitidissima omnium; at culta et captiva extra patriam exiit decus; variat dein etiam domi mille modis verè Protea.*"

At this time he had in his Herbarium a specimen without fructification of *Protea argentea* properly so called; but of its supposed varieties or of *P. fusca* none whatever. Of his genus *Leucadendron* he had only one species, *L. proteoides*, afterwards called *Protea purpurea*, a plant differing in many respects from the tribe to which he had, though not without hesitation, referred it.

In 1754 the fifth edition of *Genera Plantarum* appeared, in which the characters of both genera remain exactly as in the second.

In 1759 was published the tenth edition of *Systema Naturæ*, where the essential generic characters are nearly the same as in the sixth, and the specific characters are copied from the *Species Plantarum*.

Of this latter work the second edition appeared in 1762: it contains two additional species of *Leucadendron* described from Burmannus's Collection and *Plantæ Africanæ*; *Protea argentea* of the first edition is here divided into two species; the first *Protea argentea* now so called, the second comprehending *P. saligna*, *conifera*, and three other nearly related species: to this latter the greater part of the observation added to *P. argentea* of the first edition is annexed, though evidently less applicable to the species thus divided.

In the sixth edition of *Genera Plantarum* printed in 1764 no alterations are made in the characters of these two genera.

In *Mantissa prima* published in 1767, two new species of *Leucadendron* are described: neither of these, how-

ever, he had in his Herbarium: the first, *Leucadendron speciosum*, he had probably accidentally seen, the antheræ of which are described as filaments, and their callous apices alone as true antheræ: the description of the second, *L. pinifolium*, is by Van Royen.

In the twelfth edition of *Systema Naturæ* published in the same year, the species of *Leucadendron* are arranged in a different, and, as the author intended, a more natural order; from which it may be concluded that at this time considerable additions had been made to his Herbarium: but *L. glomeratum* is unaccountably omitted. *Protea* here receives again *P. Levisanus*, the *P. fusca* of the first edition of the *Species Plantarum*, which in the second had been referred to *Brunia*.

In *Mantissa altera* published in 1771, the two genera are united under the name of *Protea*; new characters are ⁴²¹ given to the species, and most of them are described from specimens then in his Herbarium; five species are added which had already been published by the accurate Bergius; and three, *P. totta*, *strobilina* and *parviflora*, are here first met with: in his description of the last he seems to suspect it to be a male plant, which we now certainly know to be the case. *P. glomerata* is here again taken up; but *Protea acaulis*, *cancellata* and *conocarpa* are omitted; and *Protea conifera* of the second edition of the *Species Plantarum* is subdivided into three species, *P. conifera*, *pallens* and *saligna*.

In the thirteenth edition of the *Systema Vegetabilium* published in 1774, the essential character of the genus is adapted to its present state, and no alteration occurs among the species, except that *P. speciosa* is considered as a variety of *P. Lepidocarpodendron*.

From this statement it appears, that Linnæus in his earlier works had not sufficient materials for obtaining an accurate notion of this family; and hence that perpetual fluctuation of opinion concerning it, which has been now pointed out, and may in few words be recapitulated.

1st, He gave the genus *Protea* the same extent which he at length assigned to it in the *Mantissa*.

2dly, He limited it, leaving unnoticed that part to which at a latter period he exclusively applied the name.

3rdly, He resumed his first opinion.

4thly, He subdivided it into two genera, giving them the same names which are adopted in the present paper.

5thly, He continued the subdivision but reversed the names, and for a reason, as it would seem, which is now known to be founded in error.

And lastly, Having acquired more perfect materials and perceiving the insufficiency of his characters, he united them together, thus ending exactly where he commenced.

But, as in this he has been universally followed for nearly forty years, *Protea* can no longer be considered as more strongly associated with any one species of the genus than another; and therefore this name so familiar to botanists, if the necessity of again subdividing the genus be allowed, ought certainly to be given to that part which is best known, and which contains the greatest number of published species, especially if the name be at least as applicable to this as to any other subdivision: now this part unquestionably is the *Lepidocarpodendron* of Boerhaave, the *Protea* of the first edition of the *Genera Plantarum* and *Classes Plantarum*, and of the present Essay.

The question respecting the application of the name *Leucadendron* is reducible to a smaller compass. Mr. Salisbury is aware that the Linnean character of the genus is only applicable to *Lepidocarpodendron* of Boerhaave; and therefore, consistently with the reasons which determined him in his application of the name *Protea*, *Leucadendron* ought to have been retained for that which he has called *Erodendrum* in *Paradisus Londinensis*; and this it seems he would have done, had it not been differently used by Plukenet, whom he professes to follow in this respect. But as rejecting Linnean names when accompanied by characters, for those of Plukenet who never published a single character, is somewhat unusual, it must be supposed to have arisen from the latter author's more appropriate use of this significant name, while it may also be presumed that Linnæus's application of it is wholly unsuitable;

and it is at least to be expected that in his own application he is consistent with Plukenet, whom he means to follow.

^{44]} To determine how far this is the case, I have examined the figures published by Plukenet under the name of *Leucadendros*, and also his Herbarium, which forms part of the Sloanean collection in the British Museum. Of his three species so named the first is *Protea argentea*, his "*Leucadendros africana arbor. tota argentea sericea foliis integris, Atlas Tree, D. Herman*," of which the figure represents a branch without fructification, and a separate fruit possibly of the same plant, but rather, as I suspect, belonging to a different species of the same genus.

On the same plate is figured a single leaf, in all probability belonging to *P. conocarpa*, with the following name, "*Leucadendro similis africana arbor argentea folio symmo crenaturis florida, an Leucadendros africana foliis serratis D. Herman?*" The separate fruit accompanying this, probably does not belong to it, but to some species of that division of *Leucadendron* which Mr. Salisbury has called *Euryspermum*.

The third species, his "*Leucadendros africana, seu Scolymocephalus angustiori folio apicibus tridentatis*," is a good figure of a flowering branch of *Protea cucullata*.

It could not certainly from his publications alone be understood why the name *Leucadendros* is applied to these three plants so little alike, while different names are given to species much more nearly related to some of them than they are to each other: of this however the solution is to be found in his Herbarium; on consulting which I find, that after the publication of *Protea argentea*, with whose flowers he was unacquainted, he had acquired flowering specimens of *Protea hirta*, and had supposed these two species to be the same, pasting between two leaves of *argentea* four loose heads of *hirta*, and under the whole copying in his own hand the name *Leucadendros*, &c. at full length from his *Phytographia*. This satisfactorily ^{45]} explains why he referred *P. cucullata* to *Leucadendros*, its flowers being very similar to those of *Protea hirta*. As

to his application of this name to *P. conocarpa*, it could only proceed from his total ignorance of its fructification; for, as he has figured a nearly related species, *P. hypophylla*, under the very different name of *Thymelea*, &c., it is reasonable to conclude, that had he seen the flower of *P. conocarpa* he would have given it the same generic name. This *P. conocarpa* however, of which it may truly be said he knew nothing, and concerning which at least no information is to be derived from his works, is the only species of the three which belongs to Mr. Salisbury's genus *Leucadendron*.

But the original *Leucadendros* of Hermann, of Plukenet, and of Linnaeus himself, is *Protea argentea*, the only plant of the family to which the name can properly be applied; to this therefore I have assigned it in the following arrangement.

Before proceeding to this arrangement, I am happy in having an opportunity of acknowledging that assistance which has so liberally been afforded me.

To the invaluable Herbarium and Library of Sir Joseph Banks I have on this, as on all other occasions, enjoyed the freest access; an advantage which has been greatly enhanced by the opportunity it has given me of consulting my friend Mr. Dryander, both as to the formation of genera and respecting synonyms, on which points his sound judgment and unrivalled erudition so well enable him to decide.

To Dr. Smith I am indebted for the permission of inspecting the Linnean Collection, and for the most friendly and satisfactory answers to the queries on this subject which he allowed me to put to him.

Mr. Lambert, whose Herbarium in this tribe is only surpassed by that of Sir Joseph Banks, has, with his accustomed liberality, submitted it without reserve to my examination.

Mr. Hibbert, who for many years possessed the most extensive collection of living Proteas that has ever been formed, and who also received from his intelligent collector Mr. Niven a valuable Herbarium of native specimens, most

obligingly permitted me to examine these, and even to dissect such as were new. For the like privilege I am indebted to the friendship of Mr. Aiton of Kew, who sent me his whole collection, peculiarly valuable as containing many of the original specimens of Mr. Masson : and lastly, I have to acknowledge the great assistance I have derived from the extensive collection presented to this Society by my friend Dr. Roxburgh, who during his short residence at the Cape appears to have paid particular attention to this tribe of plants, and who, besides the many new species discovered by him, has given a greater value to his Herbarium by numerous observations on the sexes, the size, and places of growth, which I have everywhere inserted on his authority.

PROTEACEÆ.

DIAGNOSIS.

Calyx tetraphyllus v. quadrifidus, æstivatione valvatâ.

Corolla nulla.

Stamina quatuor, (altero nunc sterili,) laciniis calycis opposita.

Ovarium unicum, liberum. *Stylus* simplex.

Stigma subindivisum.

Semen (pericarpium varii) exalbuminosum.

Embryo dicotyledoneus, (raro polycotyledoneus,) rectus.

Radicula infera.

47]

DESCRIPTIO.

Frutices v. *Arbores* vix excelsæ ; rarissimè *Herbæ*.

Rami in plerisque annotino-umbellati.

Folia sparsa, nunc verticillata v. opposita, persistentia, exstipulata, indivisa v. variè dentata, seu incisa profundiusve laciniata, rarissimè verè composita.

Inflorescentia subspicata, modò laxius, in racemum v. corymbum floribus sæpè geminatis, nunc densiùs congesta

in capitulum, vel aggregata supra receptaculum planiusculum, involucri persistenti, sæpius imbricato, subtensum: in quibusdam quasi abortione, uniflorum, indicante involucri calyculum tunc æmulante. *Bractæ* dum flores geminati singulis paribus communes; in capitatis persistentes, sæpiusque auctæ et induratæ, rarò connatæ; in aggregatis nanæ, plerumque deciduæ, quandoque nullæ.

Flores in plerisque hermaphroditi perfecti, nunc organorum vitio dielines.

Calyx tetraphyllus, foliolis distinctis v. sæpius plius minùs arcè cohærentibus tubulosus; *limbo* quadrifido, æquali, laciniis subspathulatis; nunc irregulari sive ex earum cohæsione rariùsve inæqualitate: coloratus, subcoriaceus, aveniùs, extùs sæpè pubescens, intùs glaber rariùsve barbâ ut plurimùm partiali instructus, valvatim aperiens, ante expansionem marginibus subtruncatis mutuò cohærentibus: deciduus v. marcescens, dum tubulosus sæpius a basi tandem quadrifidâ abscedens, quandoque basi integrâ diutiùs persistente.

Corolla nulla.

Stamina quatuor, (altero nunc sterili,) foliolis calycis opposita, iisdemque sæpissimè inserta, in plerisque juxta apicem, quandoque prope medium v. basin; rarò hypogyna; calycem nunquam superantia.

Filamenta brevissima v. mediocria, rarissimè partim cohærentia.

Antheræ adnatæ, biloculares, lineares, loculis per axin longitudinaliter dehiscentibus; rarò bipartitæ lobis respondentibus vicinarum connatis loculumque unicum tandem bivalvem efformantibus, altero lobo in quibusdam deficiente.

Pollen triangulare, angulis subsecernentibus, quandoque ellipticum v. lunatum, rarò sphericum.

Squamulæ Glandulæve hypogynæ v. quatuor foliolis calycis alternantes, distinctæ seu connatæ; v. pauciores et intus secundæ, interdum nullæ, rarissimè staminuliformes.

Ovarium unicum, liberum, sæpè pedicellatum, pedicello rarissimè articulado, mono-di-poly-spermum quandoque biloculare: ovulis apice, basi v. latere ovarii affixis.

Stylus simplex, terminalis.

Stigma in plerisque indivisum, modò emarginatum rariùsve bifidum; sæpè obliquum, figurâ varium, plerumque glabrum, quandoque papulosum, hispidulum v. tomentosum.

Pericarpium, *Nux*, *Samara* v. *Drupa* monosperma rarò disperma, vel *Folliculus* coriaceus seu ligneus, di-poly-spermus basi, marginibusve suturæ seminifer; rarò bilocularis, dissepimento libero parallelo bipartibili!

Semen sessile, ventricosum, v. sæpiùs compressum, in folliculatis sæpè alatum; exalbuminosum, apice *Chalazá* venosâ insignitum, *Rhaphi* nullâ.

Embryo dicotyledoneus, rarò polycotyledoneus, rectus, albus.

Radicula infera, brevis.

Plumula vix conspicua.

(Table facing page 48.)

PROTEACEÆ.

I. FRUCTUS CLAUSUS.

A. ANTHERÆ DISTINCTÆ.

a. ANTHERÆ A CALICE LIBERÆ.

† *Flores dioici*, (organis imperfectis). *Stigma* femineorum obliquum, emarginatum, papulosum.1. *Nux exserta*, (barbata) : *squamis* capituli feminei subulatis. *Masculi flores* racemosi.2. *Nux* v. *Samara squamis* dilatatis strobili inclusa. *Masculi flores* capitati†† *Flores hermaphroditi*, raro polygami dioici, *stigmate* tunc verticali.a. *Antheræ* apicibus concavis calycis immersæ.§ *Ovarium monospermum*. *Nux* subcrustacea v. *Samara*.a. *Squamulae* glanduleve nullæ hypogynæ.1. *Nux* lenticularis, hinc barbata v. *Samara* glabra. *Calyx* totus simul deciduus2. *Nux* ventricosa, undique æqualiter barbata. *Calyx* tubo gracili diutius persistentib. *Squamulae* v. *Glandulae* quatuor hypogynæ.|| *Calyx* irregularis, labiatus, laciniis tribus (raro omnibus) coherentibus.1. *Calyx* bipartibilis. Labii majoris laminae staminiferæ coherentes. *Stylus* persistens, subulatus.2. *Calyx* tubulosus. *Laminae* staminiferæ distinctæ. *Stylus* deciduus, filiformisPag. Pag.
39 (49)
40 (50)AULAX.....
LEUCADENDRON

PETROPHILA ... 55 (67)

ISOPOGON 58 (71)

PROTEA 60 (74)

LEUCOSPERMUM 78 (95)

LIBRARY
NATIONAL MUSEUM
OF CANADA

A. ANTHERÆ DISTINCTÆ—(contin.)

a. ANTHERÆ A CALYCE LIBERÆ—(contin.)

|| || Calyx lacinii quatuor distinctis (sepius æquidistantibus).

a. Capitulum indefinite multiflorum, subpaleaceum. *Involucrum* dum adsit imbricatum.

1. <i>Nux</i> brevissimè pedicellata. <i>Paleæ</i> persistentes. <i>Receptaculum</i> convexum	<i>SERRURIA</i>	Pag. 92 (112)
2. <i>Nux</i> sessilis. <i>Paleæ</i> deciduæ, angustissimæ v. nullæ. <i>Receptaculum</i> planum	<i>MIMETES</i>	86 (105)

β. *Involucrum* uniflorum v. definitè pauciflorum. *Paleæ* nullæ.

† *Squamulæ hypogynæ* a calyce toto deciduo liberæ.

* *Stigma* verticale. *Calyx* regularis.

1. <i>Nux</i> sessilis, nitens, basi integrâ. <i>Involucrum</i> fructiferum induratum, 4-pl., 4-flor.	<i>NIVENIA</i>	110 (133)
2. <i>Nux</i> pedicellata v. basi emarginata. <i>Involucrum</i> fructiferum non mutatum	<i>SOROCEPHALUS</i>	115 (139)

** *Stigma* obliquum, dilatatum. *Calycis* lacinia quartâ sepiè majore

SPATALLA

119 (143)

†† *Squamulæ hypogynæ* infra adnatæ basi persistenti calycis

ADENANTHOS

125 (151)

§ § *Ovarium dispersum*. *Drupa* baccata, putamine osseo. *Stigma* obliquum. *Calyx* irregularis

GUEVINA

138 (165)

β *Anthere* exsertæ. *Stamina* medio v. basi calycis inserta v. hypogyna.

† *Glandulæ* hypogynæ distinctæ s. connatæ. v. *Stamina* quatuor sterilia. *Drupa* putamine osseo.

* *Drupa* exsucca, tomentosa. *Filamenta* basi calycis inserta. *Vagina* hypogyna. *Flores* polygami

BRABEUM

137 (164)

- β *Antheræ* exsertæ. *Stamina* medio v. basi calycis inserta v. hypogynæ.
- + *Glandula* hypogynæ distinctæ s. connatæ. v. *Stamina* quatuor sterilia. *Drupa* putamine osseo.
- * *Drupa* exsucca, tomentosa. *Filamenta* basi calycis inserta. *Vagina* hypogyna. *Flores* polygami *BRABEUM* 137 (164)

- ** *Drupa* baccata. *Flores* hermaphroditi.
1. *Glandula* hypogynæ carnosæ. *Stamina* medio calycis supra recurvi inserta. *Ovarium* pedicellatum *PERSOONIA* 133 (159)
2. *Glandula* hypogynæ staminuliformes. *Stamina* basi calycis supra angustati inserta. *Ovarium* sessile *CENARRHENS* 132 (158)
- ++ *Glandula* nullæ hypogynæ.
- § *Stamina* calyci inserta.
1. *Filamenta* distincta. *Stigma*, unilaterale. *Ovarium* monospermum, trigonum (imberbe) *AGASTACHYS* ... 132 (158)
2. *Filamenta* apice coherentia. *Stigma* subtruncatum. *Ovarium* dispermum, teres *SYMPHONEMA* 131 (157)
- § § *Stamina* receptaculo inserta. *Samaræ* apteræ, 1-2-sperma. *Stigma* simplex... *BELLENDENA* ... 139 (166)
- b. *ANTHERÆ ADNATÆ* TUBO CALYCIS HYPOCRATERIFORMIS! *Næz* fusiformis, pedicellata; apice dilatato papposo FRANKLANDIA... 130 (157)
- b. *ANTHERÆ COHERENTES*, vicinarum lobis proximis loculum unicum constituentibus! tandem distinctæ.

- a. *Calycis* regularis. *Antheræ* exsertæ, omnes bilobæ *SIMSIA* 126 (152)
- β. *Calycis* ringens. *Antheræ* inclusæ, laterales dimidiatæ; *Stamine* quarto sterili.
1. *Stigma* liberum. *Anthera* labii superioris biloba *CONOSPERMUM* 127 (153)
2. *Stigma* filamento sterili coherentis. *Anthera* media labii inferioris biloba *SYNAPHEA* 129 (155)

II. FRUCTUS DEHISCENS.

A. UNILOCULARIS.

- a. *OVARIUM* DISPERSUM. *Fructus* quandoque monospermus.

† *Antheræ* apicibus concavis calycis immerse. *Glandula* hypogyna unica dimidiata, v. nulla.

- * *Glandula* nulla hypogyna. *Stigma* conicum. *Semen* apterum *ANADENIA* 139 (166)
- ** *Glandula* hypogyna dimidiata, quandoque lobata.
1. *Folliculus* (coriaceus v. ligneus) loculo centrali. *Semina* alia apicis dum adsit nucleo breviora *GREVILLEA* 140 (167)
2. *Folliculus* incrassato-ligneus, loculo excentrico. *Semina* alia apicis nucleo longiora... *HAKEA* 149 (178)

A. UNILOCULARIS—(contin.)

a. OVARIUM DISPERMUM—(contin.)		Pag.	Pag.
†† <i>Antheræ</i> exsertæ, apicibus calycis revolutis. <i>Glandulæ</i> hypogynæ quatuor, distinctæ v. connatæ.			
‡ <i>Involucrum</i> coloratum, deciduum, uni-multiflorum, receptaculo plano. <i>Semina</i> marginata.			
<i>Stigma</i> subulatum.....	LAMBERTIA.....	157	(187)
†† <i>Involucrum</i> nullum. <i>Flores</i> spicati.			
a. <i>Semina</i> apice (solum) alata.			
1. <i>Folliculus</i> incrassato-ligneus, loculo excentrico. <i>Stigma</i> clavatum. <i>Flores</i>	XYLOMELUM	159	(189)
polygami	ORITES	159	(189)
2. <i>Folliculus</i> coriaceus, loculo centrali. <i>Stigma</i> subcylindraceum			
β. <i>Semina</i> utrinque alata, marginata. <i>Stigma</i> clavatum. <i>Folliculus</i> ligneo-coriaceus	RHOEPA.....	160	(190)
b. OVARIUM TETRASPERMUM! <i>Calyx</i> regularis. <i>Antheræ</i> exsertæ. <i>Semina</i> apice alata. <i>Glandulæ</i> quatuor	KNIGHTIA	163	(193)
hypogynæ ..			
c. OVARIUM POLYSPERMUM. <i>Calyx</i> irregularis, apicibus concavis staminiferis.			
§ <i>Semina</i> apice alata.			
* <i>Stigma</i> verticale, clavatum. <i>Glandula</i> hypogyna unica, semiannularis	EMBOTHRIUM	165	(195)
** <i>Stigma</i> obliquum, unilaterale.			
1. <i>Glandula</i> nulla hypogyna. <i>Stigma</i> dilatatum, concavum. <i>Involucrum</i> (racemi) nullum	OREOCALLIS	165	(196)
2. <i>Glandula</i> unica hypogyna, subannularis. <i>Stigma</i> clavatum, convexum. <i>Involucrum</i>	TELOPEA.....	166	(197)
(racemi) deciduum			
3. <i>Glandulæ</i> tres hypogynæ, secundæ. <i>Stigma</i> dilatatum, concavum. <i>Calyx</i> foliola	LOMATIA.....	168	(199)
distincta	STENOCARPUS.....	170	(201)
§ § <i>Semina</i> basi alata! <i>Glandula</i> hypogyna unica dimidiata. <i>Stigma</i> dilatatum, concavum			

B. BILOCULARIS, dissepimento libero, bifido.

1. <i>Amentum</i> paribus flosculorum tribracteatis	BANESIA	171	(202)
2. <i>Receptaculum commune planum</i> ; involucri imbricato; flosculis indeterminatim confertis, paleis solitariis			
v. nullis	DRYANDRA	179	(211)

1. AULAX.

[49]

*Berg. Cap. 33. Salisb. Parad. 67.*CHAR. GEN. *Flores* dioici, organis imperfectis.Masc. racemosi: *Calyx* tetraphyllus foliolis medio staminiferis.Fem. *Stigma* obliquum, clavatum, hispidulum, emarginatum. *Nux* exserta, ventricosa, barbata, squamis capituli subulatis.HABITUS. Frutices *glaberrimi*. Folia *integerrima*. Flores *terminales, unibracteati*. Masculi *in racemis aggregatis, nudis*. Feminei *in capitulo solitario, cincto foliolis intus auctis appendiculâ aceroso-multifidâ, capitulo quasi abortivo racemis exterioribus maris analogo, (interdum florifero, fide Cel. Salisburii.)*1. *A. pinifolia*, foliis filiformibus canaliculatis.Masc. Pini foliis planta Africana Cyperi capitulis. *Herm. Afr. 18. Burm. Afr. 193, t. 70, f. 3.*Pini folio planta Capensis floribus spicatis. *Raj. Hist. 3, App. p. 247, n. 47.*Pini facie frutex africanus, Cyperi capitulis umbellatis. *Sher. in Raj. Hist. 3, Dendr. p. 130.**Aulax pinifolia. Berg. Cap. 33.***Leucadendron pinifolium. D. Van Royen in Linn. Mant. 36.* Syst. Nat. ed. xii, t. 2, p. 110.**Protea pinifolia. Linn. Mant. 187. (sed specimen maris A. umbellatæ habebat in Herb.) Syst. Nat. ed. xiii, t. 2, p. 117. Thunb. Diss. n. 20.* Prod. 26. Willd. Sp. Pl. 1, p. 515. Lam. Illust. Gen. 1, p. 237, n. 1244. And. Repos. 76, bona. Poiret, Encyc. Botan. 5, p. 651.*Fem. *Scolymocephalus africanus* foliis *Rorismarini* acutis. *Herm. Afr. 20. Raj. Hist. 3, Dendr. p. 10.**Conophorus capensis* pini folio. *Petiv. Gazoph. 3, n. 458, t. 25, f. 7, 8vo, p. 40.**Lepidocarpodendron*; foliis angustissimis, gramineis; fructu cancellato; semine coronato. *Boerh. Lugd. Bat. 2, p. 193, c. tab.*

Leucadendron cancellatum. *Linn. Sp. Pl. ed. 1, p. 91; ed. 2, p. 134; omiss. in Mant. et Syst. Veg. ed. xiii.*

Protea bracteata. *Thunb. Diss. n. 24,* tab. 1. Prod. 26. Linn. Suppl. 118. Willd. Sp. Pl. 1, p. 517. Lam. Illust. Gen. 1, p. 238, n. 1245. Poiret, Encyc. Botan. 5, p. 652.*

HAB. In Africæ Australis montibus; prope Plattekloof, Hottentots-Holland, et alibi. (v. s. in Herb. plur.)

Obs. Pollen globosum.

2. *A. umbellata*, foliis planis spathulato-linearibus.

MASC. *Protea aulacea*. *Thunb. Diss. n. 33,* tab. 2, bona. Prod. 26. Willd. Sp. Pl. 1, p. 520. Lam. Illust. Gen. 1, p. 237, n. 1243. Poiret, Encyc. Botan. 5, p. 651.*

FEM. *Protea umbellata*. *Thunb. Diss. n. 34.* Prod. 26. Linn. Suppl. 118. Willd. Sp. Pl. 1, p. 520. Lam. Illust. Gen. 1, p. 237, n. 1242. And. Repos. 248. Poiret, Encyc. Botan. 5, p. 650.*

HAB. In Africæ Australis montibus, prope Prom. B. Spei; Taffelberg, Platte-kloof, &c. (v. s. in Herb. Banks.)

Obs. Pollen subglobosum, obtusissimè trigonum.

2. LEUCADENDRON.

Herm. Pluk. Linn. in Class. Plant. Conocarpodendra, t. 195, 197, 200, 202, 203, 204. Boerh. Protea. Linn. Gen. Pl. ed. 2, 5, 6. Conocarpos. Adans. Famill. Protea, Euryspermum, Chasme. Salisb. Parad. Lond.

CHAR. GEN. Flores dioici, organis imperfectis; capitati.

Fem. *Stigma* obliquum, clavatum, emarginatum, hispidulum. *Nux* v. *Samara* monosperma, squamis (quandoque cohærentibus) strobili inclusa.

HABITUS. Frutices, rarò Arbores, sæpe sericeo-tomentosi. Folia integerrima. Capitula terminalia, solitaria; bracteis imbricatis foliisve verticillatis et subcoloratis plerumque cincta.

Obs. The separation of sexes in the genus *Protea* of authors, obscurely suspected by Linnæus himself in his *Protea parviflora*, and afterwards more expressly by La-

mark in *P. pinifolia*, was first ascertained in *Aulax* and the present genus (as Mr. Dryander informs me) by our countryman Masson, during his last residence at the Cape of Good Hope, and is beautifully illustrated by that eminent botanical painter Mr. Francis Bauer, in his unpublished drawings preserved in the Banksian collection. Numerous observations on the same subject have also more recently been made by Dr. Roxburgh and Mr. Niven, who have bestowed much pains in ascertaining its limits, of which, as far as regards the African part of the family, Mr. Salisbury has given an accurate account in his Essay already quoted. The Dissertation of Thunberg, who was wholly unacquainted with this separation of sexes in these plants, is necessarily imperfect, and he has, in several cases, described the different sexes as distinct species; and thus also Bergius has founded his genus *Aulax* on the male of a species, whose female he had previously published as a *Leucadendron*. On the other hand, Jussieu, deceived by the resemblance in inflorescence, between *Brabejum* and the spiked species of *Protea*, has erroneously suspected these to be monoicous, while he has totally overlooked the truly dioicous nature of the present genus.

† *Nux ventricosa*, stylo toto calyceque persistentibus.

[52]

1. *L. argenteum*, arboreum, foliis lanceolatis argenteis: marginibus ramisque villosis, bracteis involucrentibus abbreviatis tomentosis, calycibus masculis sericeis.

Scolymocephalus Africana, foliis sericeis argenteis longis acutis. *Herm. Cat.*

Leucadendros Africana, arbor tota argentea, sericea, foliis integris. Atlas Tree. D. Herman. *Pluken. Phyt. t.* 200, f. 1, ramulus sine floribus nuce separata; fortè speciei distinctæ.

Conifera salicis facie, folio et fructu tomento sericeo candicante obductis, semine pennato. *Sloane in Philos. Trans.* 17, p. 664, strobilus cum fructu separato.

Frutex *Æthiopicus* conifer, foliis lanuginosis omnium maximis. *Breyn. Prod.* 2, p. 66.

Argyrodendros africana foliis sericeis et argenteis. *Com. Hort.* 2, p. 51, t. 26. *Raj. Hist.* 3, *Dendr.* p. 9.

Globularia Africana frutescens *Thymeleæ* folio lanuginoso. *Tournef. Inst.* 467?

Conocarpodendron; foliis argenteis, sericeis, latissimis. *Boerh. Lugd. Bat.* 2, p. 195, c. tab.

Scolymocephalus africanus, folio crasso nervoso sericeo. *Weinm. Phyt.* 4, p. 293, t. 900.

Brückm. Epist. Itin. 2, p. 8, t. 4, strobilus.

Protea foliis lanceolatis integerrimis acutis hirsutis nitidis. *Hort. Cliff.* 29. *Virid. Cliff.* 8. *Roy. Lugd. Bat.* 184. *Wachend. Ultraj.* 201.

Protea argentea a. *Sp. Pl. ed.* 1, p. 94.

Protea argentea. *Sp. Pl. ed.* 2, p. 137. *Mant.* 194. *Thunb. Diss. n.* 48.* *Prod.* 27. *Gært. Sem.* 1, p. 239, t. 51. *Willd. Sp. Pl.* 1, p. 529. *Lam. Illust. Gen.* 1, p. 237, n. 1236, t. 53, f. 1. *Poiret, Encyc. Botan.* 5, p. 648.

⁵⁵³ HAB. In Africâ Australi, ad radices lateraque montium, prope *Pro* 3. *Speci.* (v. s. in *Herb. Banks. &c.*)

OBS. Squamulæ quatuor hypogynas, longas, angustolineares, in floribus masculis observavi: femineos nondum vidi.

2. *L. plumosum*, fruticosum, foliis lineari-lanceolatis muticis glabris subsericeisve: basi attenuatâ tortâ, involucris calycibusque masculis glabris: femineis persistentibus plumosis quadrifidis, nucibus cuneato-oblongis villosis.

MASC. *Protea parviflora*. *Linn. Mant.* 195.* *Syst. Veg. ed.* xiii, p. 119. *Thunb. Diss. n.* 40,* tab. 4, bona. *Prod.* 27. *Willd. Sp. Pl.* 1, p. 524. *Lam. Illust. Gen.* 1, p. 235, n. 1220. *Poiret, Encyc. Botan.* 5, p. 643.

FEM. *Protea obliqua*. *Thunb. Diss. n.* 39.* *Prod.* 27. *Linn. Suppl.* 117,* fide descriptionis, nullum enim specimen in herbario, monente Cel. Smith. *Willd. Sp. Pl.* 1, p. 524.

Protea plumosa. *Hort. Kew.* 1, p. 127.

HAB. In Africâ Australi, prope *Fransche Hoek* et alibi haud infrequens. (v. s. in *Herb. Banks.*)

Obs. Squamulæ nullæ hypogynæ, in mare: femina ad anthesin haud observata.

*3. *L. retusum*, fruticosum, foliis oblongo-spathulatis glabris: basi attenuatis; callo apicis subretuso, ramis tomentosis, involucris pubescentibus, calycibus masculis glabris, femineis plumosis quadripartitis, nucibus glabris ovato-orbiculatis.

HAB. In Africæ Australis montibus prope Prom. B. Spei; Picket-Berg. (v. s. in Herb. Soc. Linn.)

Obs. Strobilus cum Nucibus et Calyce ad basin tabulæ 199. *Boerh. Lugd. Bat.* vel ad hanc vel ad sequentem speciem pertinere videntur.

*4. *L. spathulatum*, fruticosum, foliis oblongo-spathulatis: basi attenuatis; callo apicis acutiusculo recurvo ramisque glabris, calycis feminei tardiùs decidui laminis nudis, nucibus glabris latioribus quàm longis demum muticis.

HAB. In Africæ Australis planitiis elevatioribus arenosis, prope Promont. B. Spei. (v. s. in Herb. Hibbert.)

Obs. In Mare? squamulas quatuor longissimas hypogynas observavi.

*5. *L. sessile*, fruticosum, foliis lanceolato-oblongis glabris: basi obtusâ.

HAB. In Africâ Australi prope Promont. B. Spei. *D. Masson.* (v. s. in Herb. Banks.)

Obs. Squamulæ hypogynæ longissimæ, persistentes.

†† *Nux ventricosa* v. *lenticularis aptera*, undique marginibusve pilosa. Stylus totus deciduus, basive solum remanenti. Calyx diù persistens quadripartitus.

*6. *L. angustatum*, foliis lineari-spathulatis (passim angustatis) obtusis muticis concaviusculis ramisque glabris, strobili squamis conniventibus nucibusque pubescentibus subglobois muticis, calycibus plumosis.

HAB. In Africâ Australi, prope Promont. B. Spei. *D. Masson.* (in Herb. D. Aiton vidi.)

Desc. *Frutex*, ramis strictis, divisis. *Folia* sparsa, fre-

quentia, erecta, 8—9 lineas longa, obtusissima, ad apicem (vix callosum) sesquilineam lata, in eodem ramo passim dimidio angustiora. *Strobilus* subglobosus, magnitudine globuli sclopi minoris: *Squamis* ovatis exterioribus latioribus. *Nux* magnitudine seminis Viciæ, levissimè compressa, undique pube brevi induta, cincta calyce ad basin quadripartito vix longiore.

^{55]} *7. *L. imbricatum*, foliis lanceolato-linearibus glabris imbricatis basi obtusis, squamis strobili dilatato-cuneatis sericeis, nucibus undique comosis basi styli cuspidatis.

Protea Levisanus. *Herb. Linn.*

HAB. In Africâ Australi, prope Prom. B. Spei; Rode Zant. (v. s. in *Herb. Banks. et Soc. Linn.*)

OBS. Squamæ nullæ hypogynæ.

*8. *L. buxifolium*, foliis ovali-lanceolatis subimbricatis: adultis glabris, squamis strobili dilatato-cuneatis sericeis, nucibus undique comosis basi styli mucronatis.

Protea imbricata. *Wend. Hort. Herenh. tab. 14?* excl. syn.

HAB. In Africâ Australi. *D. Masson.* (v. s. in *Herb. Banks.*)

DESC. *Frutex* erectus, ramosissimus. *Rami* umbellati, tenuissimè tomentosi. *Folia* frequentia, erecto-imbricata, sessilia, concaviuscula, avenia, opaca, semiunguicularia; superiora tenuissimè pubescentia; floralia angustiora. MASC. *Capitulum* sessile magnitudine pisi. *Calyx* tubo unguibusque sericeis, laminiis glabriusculis. *Squamulæ* quatuor hypogynæ longæ, lineares. FEM. *Capitulum* paulo majus. *Calycis* unguis laminæque sericeæ. *Squamulæ* nullæ hypogynæ. *Nux* ovata, calyce diù cincta.

OBS. I. Nimis affine *L. imbricato*, figurâ foliorum præsertim distinguendum.

OBS. II. Icon *Wend.* suprâ citata forsân diversæ speciei; foliis oblongis semuncialibus pilosis, strobilis longioribus, squamulis hypogynis: an potiùs ad priorem referenda?

9. *L. Levisanus*, foliis obovato-spathulatis obtusissimis : adultis glabris, ramulis pilosis, capitulis masculis sessilibus, nucibus undique comosis muticis.

Levisanus capensis serpylli folio. Petiv. Gazoph. 9, [56 t. 5, f. 7 ? mala.

Chrysanthemum Conyzoides Æthiopicum, capitulo aphylo, Tithymali paralii foliis subrotundis, hiuleculis in superficie conspicuis. *Pluk. Mant. 47, t. 343, f. 9, bona.*

Conocarpodendron ; foliis subrotundis, brevissimis, capituli immaturi globosi parte inferiore fuscâ, mediâ aureâ, supremâ viridi. *Boerh. Lugd. Bat. 2, p. 202, c. tab.*

Brunia foliis oblongis incanis, florum capitulo ramulum terminante. *Burm. Afr. 267,* t. 100, f. 2, Mas. bona.*

Scolymocephalus seu *Conocarpodendron*, foliis brevissimis. *Weinm. Phyt. 4, p. 296, t. 904, a, pessima quoad colores.*

Protea fusca. Linn. Sp. Pl. ed. 1, p. 95.

Brunia Levisanus. Linn. Sp. Pl. ed. 2, p. 289.

Leucadendron Levisanus. Berg. Act. Stockh. 1766, p. 324. Berg. Cap. 20,* Mas.*

Protea Levisanus. Linn. Syst. Nat. ed. xii, t. 2, p. 111. Mant. 194, quoad descriptionem, sed specimen in Herbario est feminæ L. imbricati. Thunb. Diss. n. 43.* Prod. 27. Willd. Sp. Pl. 1, p. 526. Lam. Illust. Gen. 1, p. 235, n. 1223. Poiret, Encyc. Botan. 5, p. 644. Wend. Hort. Herenh. t. 1, Mas.*

HAB. In Africæ Australis campis sabulosis ericetisque, prope Prom. B. Spei. (v. s. in Herb. Banks.)

Obs. *Squamulæ nullæ hypogynæ.*

10. *L. tortum*, foliis spathulato-linearibus obtusis basi tortis : adultis glabris ; junioribus ramulisque subsericeis, capitulis masculis pedunculatis, calycis laminis sericeis, nucibus undique comosis muticis.

Protea torta. Thunb. Diss. n. 31? Prod. 26? Willd. Sp. Pl. 1, p. 519?*

Protea cinerea. Willd. Sp. Pl. p. 521. Fem. sec. [57 desc. exclus. syn.*

HAB. In Africæ Australis depressis, prope Prom. B. Spei. (v. s. in Herb. Banks. et Soc. Linn.)

11. *L. cinereum*, foliis spathulato-linearibus argenteis, capitulis masculis sessilibus, nucibus obovato-cuneatis villosiusculis muticis.

Protea alba. *Thunb. Diss. n. 32,* sec. desc. Prod. 26. Willd. Sp. Pl. 1, p. 520. Lam. Illust. Gen. 1, p. 236, n. 1233. Poiret, Encyc. Botan. 5, p. 647.*

Protea cinerea. *Hort. Kew. 1, p. 127.* Fem. fid. spec. descript. in Herb. Banks.

HAB. In Africâ Australi, prope Prom. B. Spei. (v. s. in Herb. Banks.)

OBS. Squamulæ quatuor hypogynæ in utroque sexu.

12. *L. corymbosum*, foliis lineari-subulatis imbricatis glabris, strobili squamis acutis apice recurvis, nucibus subcompressis obcordatis margine pilosis.

Leucadendron corymbosum. *Berg. Act. Stockh. 1766, p. 325.* Berg. Cap. 21,* Mas.*

Protea corymbosa. *Thunb. Diss. n. 28,* (desc. e mare præcipuè,) tab. 2, Mas. Thunb. Prod. 26. Willd. Sp. Pl. 1, p. 518. Lam. Illust. Gen. 1, p. 238, n. 1250. Poiret, Encyc. Botan. 5, p. 653.*

Protea bruniades. *Linn. Suppl. 117.** Mas. fid. spec. in illius Herb.

Protea corymbosa. *And. Repos. 495, Fem.*

HAB. In Africâ Australi, Drakenstein, Swartland, Rode Zant. (v. s. in Herb. Soc. Linn.)

OBS. Squamulæ quatuor hypogynæ in utroque sexu.

58] †† *Samara glabra alata v. aptera*. Stylo (fere toto) calyceque deciduis.
Squamæ strobili distinctæ.

13. *L. decorum*, foliis oblongis venosis callo recurvo: adultis glabris; junioribus ramisque sericeis; floralibus coloratis semiscariosis, strobili squamis extus tomentosis: apice retuso parùm coarctato glabro, nucibus marginatis impresso-punctatis.

Prom. B.

Protea laureola. *Lam. Illustr. Gen.* 1, p. 234, n. 1214.
Poiret, Encyc. Botan. 5, p. 641.* Fem. exclus. syn.
Linnei, Thunbergii, Schraderi.

HAB. In Africâ Australi, prope Prom. B. Spei. (v. s. in
Herb. Banks.)

c. *Prod.*
Gen. 1, p.

. fid. spec.

i. (v. s. in

e sexu.

imbricatis
cibus sub-

ckh. 1766,

sc. e mare

lld. *Sp. Pl.*

. *Poiret,*

. fid. spec.

land, Rode

e sexu.

ue deciduis.

o recurvo :

floralibus

omentosis :

marginatis

14. *L. squarrosus*, foliis lanceolato-oblongis : callo recurvo anticè sulco lineari ; adultis ramisque glaberrimis, amenti feminei oblongo-ovati squamis sursum glabris dilatatis integris ciliatis : strobili recurvis undulatis.

MASC. *Protea arcuata*. *Lam. Illustr. Gen.* 1, p. 234, n. 1215 ?

Protea obliqua α. *Poiret, Encyc. Botan.* 5, p. 642* ? exclus. syn. Thunber. Linnei et Boerhaav.

FEM. *Protea strobilina*. *Linn. Mant.* 192.*

β. *Conocarpodendron* ; folio rigido, crasso, angusto, cono laricis parvo. *Boerh. Lugd. Bat.* 2, p. 197, c. tab. Fem. ?

HAB. In Africâ Australi, prope Prom. B. Spei. (v. s. in Herb. Soc. Linn.)

15. *L. concolor*, foliis spathulato-oblongis : callo anticè subrotundo ; adultis glabris ; floralibus masculis concoloribus, ramis pubescentibus, amenti feminei squamis retusis basi tomentosis, margine ciliatis.

MASC. *Protea arcuata* β. *Lam. Illustr. Gen.* 1, p. 234, n. 1215 ?

Protea globosa. *And. Repos.* 307, bona. *Sims in Bot. Mag.* 878.

Protea obliqua β. *Poiret, Encyc. Botan.* 5, p. 642 ? exclus. syn. Boerh. Thunb. Linnei, Willd.

FEM. *Protea strobilina*. *Schrad. Sert. Hanov.* 1, p. 7, t. 1.

HAB. In Africâ Australi, prope Prom. B. Spei. (v. s. in Herb. Soc. Linn.)

16. *L. grandiflorum*, foliis lanceolato-oblongis : callo apicis anticè subrotundo ; adultis glabris ; floralibus coloratis, ramis tenuissimè tomentosis, squamis amenti utriusque sexûs ovatis obtusiusculis glabris fucatis.

MASC. *Euryspermum grandiflorum*. *Salisb. Parad.* 105.
HAB. In Africâ Australi, prope Prom. B. Spei. (v. s.)

*17. *L. ovale*, foliis oblongo-ovalibus subaveniis: callo obtuso; adultis utrinque glabris margine tomentosis, strobili squamis lanceolato-ovatis acutis glabris, samaris apteris impunctatis extûs ventricosus.

Protea strobilina. *Thunb. Diss. n. 44,** secund. descrip.
HAB. In Africâ Australi. *D. Masson*. Palmetta River.
Gul. Roxburgh M.D. (v. s. in Herb. Banks.)

*18. *L. venosum*, foliis oblongo-lanceolatis venosis glabris: callo acuto, strobili squamis ovato-lanceolatis acutis ciliatis extra medium glabris, calycibus persistentibus, nucibus apteris.

HAB. In Africâ Australi. *Gul. Roxburgh M.D.* (v. s. in Herb. Soc. Linn.)

19. *L. decurrens*, foliis lanceolato-spathulatis basi attenuatis subdecurrentibus concaviusculis ramisque glabris, calycis feminei tubo hirsuto, strobili squamis subrotundis demùm glabriusculis, samaris obcordatis alatis cinereis utrinque convexis.

Protea pallens. *Thunb. Diss. n. 41,** secund. descrip. exclus. omnibus synonymis.

Protea chamelæa. *Lam. Illust. Gen. 1, p. 237, n. 1240?* exclus. syn. *Poiret, Encyc. Botan. 5, p. 650*?*

HAB. In Africâ Australi, prope Prom. B. Spei. (v. s. in Herb. Banks. et Soc. Linn.)

*20. *L. glabrum*, foliis lineari-lanceolatis aveniis: adultis ramisque glabris, strobili squamis obtusissimis subretusis tuboque calycis nudis, samaris alatis nigris planiusculis dilatatis.

HAB. In Africâ Australi, prope Prom. B. Spei. *Gul. Roxburgh M.D.* (v. s. in Herb. Soc. Linn.)

21. *L. strictum*, foliis linearibus mucrone subulato glabris, bracteis involucrantibus ovatis acutis capitulo florido longio-

ribus, strobili squamis dilatatis rotundatis glabris, samaris apteris impresso-punctatis.

MASC. *Euryspermum salicifolium*. *Salisb. Parad.* 75, bona.

Protea conifera. *And. Repos.* 541.

Protea conica. *Lam. Illust. Gen.* 2, p. 237, n. 1237?

Protea conifera A. *Poiret, Encyc. Botan.* 5, p. 649?

HAB. In Africâ Australi, ad ripas fluviorum et in humidis inter saxa, prope Stellenbosch et Rode Zant. *Gul. Roxburgh M. D.* (v. s. in Herb. Banks. et Soc. Linn.)

*22. *L. virgatum*, foliis linearibus acutis pellucido-marginatis ramisque glaberrimis: floralibus lineari-lanceolatis elongatis, strobili squamis ovatis integris incanotomentosis, samaris alatis emarginatis.

HAB. In Africâ Australi, prope Promont. B. Spei. (v. s. in Herb. Soc. Linn.)

Obs. Folia ramea basi torta; floralia intus colorata ramis strobilisque duplo longiora; bracteæ involucrantes capitulo florido breviores. Strobilus ovatus magnitudine Avellanæ, squamis tomento vix nitente, superioribus apice sub-coarctatis.

*23. *L. adscendens*, foliis lineari-lanceolatis acutis: floralibus lanceolatis apice coarctatis coloratis corcavis, strobili squamis ovatis integris incano tomentosis, samaris alatis emarginatis, ramis adscendentibus, caulibus subdepressis.

Thymelæa capitata seu *julifera* angusto salicis folio promontorii Bonæ Spei. *Pluk. Mant.* 181, t. 229, f. 6, Mas. fide specim. in Herb. Pluk.

Conocarpodendron; folio angusto, rigido, brevior; cono parvo aureo, coronâ foliaceâ succinto. *Boerh. Lugd. Bat.* 2. p. 200. c. tab.?

Scolymocephalus minor. *Wein. Phyt.* 4, p. 295, t. 903. a.?

Protea pallens. *Linn. Mant.* 193.* Mas. fid. spec. in ejusd. Herb.

Protea conifera. *Linn. Mant.* 193.* Fem. fid. spec. in ejusd. Herb.

HAB. In Africæ Australis montibus, prope Promont. B. Spei. (v. s. in Herb. Banks.)

OBS. I. Strobilus ad basin tab. 9, Breyn. Cent. huc pertinet.

OBS. II. *Protea pallens* et conifera auctorum recentiorum ex eorum descriptionibus incompletis et ab altero solùm sexu desumptis vix extricandæ et æquo jure ad hanc speciem vel ad *L. virgatum* v. *glabrum* citari possint.

*24. *L. concinnum*, foliis lanceolato-oblongis obtusiusculis ⁶²⁾aveniis ramisque glaberrimis: floralibus subconformibus semicoloratis, strobili squamis ovatis integris argentecomentosis, samaris alatis emarginatis.

HAB. In Africæ Australis montibus. *D. Niven.* (in Herb. Hibb. vidi.)

DESC. *Frutex* decempedalis (*Niven.*) *Rami* stricti, glaberrimi. *Folia* frequentia, erecta, subimbricata, uncialia, marginibus angustissimis, semipellucidis, scabriusculis, parùm concava, callo apicis obtusiusculo: *floralia* sesquiuncialia, strobilo maturo vix duplo longiora.

25. *L. salignum*, foliis lanceolato-linearibus acutissimè mucronatis subsericeis: floralibus lanceolatis coloratis, strobili squamis tomentosis sursùm dilatatis retuso-bilobis margine glabris, samaris apice latiusculè margine angustissimè alatis.

Conocarpodendron; folio tenui, angusto, saligno; conocalyculato, coronâ foliaceâ succincto. *Boerh. Lugd. Bat. 2, p. 204, c. tab.*

Protea foliis lineari-lanceolatis integerrimis acutis. *Jort. Cliff. 29*, secund. specim. in Herb. Cliff. quod ramulus absque fructificatione.

Protea foliis lineari-lanceolatis integerrimis superioribus hirsutis nitidis. *Roy. Lugd. Bat. 184.*

Protea argentea β. *Linn. Sp. Pl. ed. i, p. 94*, exclus. syn. Breynii et Tournefortii.

Protea conifera α. *Linn. Sp. Pl. ed. ii, p. 138*, excl. syn.

Protea saligna. *Linn. Mant. 194*, Mas. exclus. syn.

Bergii et Breynii. *Lam. Illustr. Gen.* 1, p. 236, n. 1235.
Poiret, Encyc. Botan. 5, p. 648,* Fem.

HAB. In Africâ Australi, prope Promont. B. Spei, in montosis. (v. s. in Herb. Soc. Linn.)

26. *L. uliginosum*, foliis lanceolato-linearibus utrinque argenteis tomento arcuè adpresso: callo apicis acuto, ramis tomentosis, calycis feminei tubo hirsuto, squamis strobili sericeis dilatatis subundulatis obsolete retusis, samaris apteris.

Frutex æthiopicus conifer foliis cneori salici æmulis. *Breyn. Cent.* 21, t. 9, excepto strobilo ad basin tabulæ, qui ad *L. adscendens* pertinet.

Protea saligna. *Thunb. Diss.* n. 47,* secund. descrip.

HAB. In Africæ Australis uliginosis, prope Prom. B. Spei. (v. s.)

27. *L. floridum*, foliis lanceolato-linearibus sericeis supra villosis: callo apicis acuto; floralibus subtus ramisque hirsutis, calycibus masculis longitudinaliter pilosis, strobili squamis tomentosis apice dilatatis integris, samaris apteris.

Thyridæa capensis sericeis longioribus et acutis foliis caule geniculato piloso. *Pluk. Phyt.* 181, t. 229, f. 4, fide specim. in ejus Herb.

Protea saligna mas et fem. *And. Rep.* 572?

HAB. In Africâ Australi, prope Promont. B. Spei, in paludosis ad radicem Montis Wynberg. *Gul. Roxburgh M. D.* (v. s. in Herb. Banks. et Mus. Brit.)

+++ Squamæ strobili connatæ. Samara foliaceo-compressa, glabra. Folia aliqua vel omnia filiformia.

*28. *L. platyspermum*, foliis superioribus lineari-spathulatis: callo obtuso, strobili squamis duplicatis longitudinaliter accretis: rimis semicircularibus, samaris duplo latioribus quàm longis.

HAB. In Africâ Australi, prope Promont. B. Spei. Hout Hoek. *And. Auge.* (v. s. in Herb. Banks.)

DESC. *Frutex* glaber. *Folia* inferiora filiformia, canaliculata, sesquiunciolia; superiora plana, vix longiora.

⁶⁴ MASC. *Amentum* ovatum, bracteis lanceolatis, sericeis subtensum. *Laminæ* calycis glabræ. *Stigma* clavatum.

FEM. *Amentum* oblongum: *Squamis* glabris connatis. *Ungues* calycis villosæ. *Laminæ* glabræ. *Stigma* dilatatum, obliquum, papulosum. *Strobilus* oblongus, quandoque ovatus, rimis transversis semicirculum sub-æquantibus. *Samara* cinerea, lævis.

29. *L. comosum*, foliis superioribus spathulato-lanceolatis obtusis mucronatis rugoso-striatis, strobilis oblongis: squamis basi connatis suprâ distinctis marginibus inflexis sub-barbatis, samaris subrotundis nigris.

Protea comosa. *Thunb. Diss. n. 25,** secund. descript. folior. *Prod. 26. Willd. Sp. Pl. 1, p. 517. Lam. Illustr. Gen. 1, p. 238, n. 1254. Poiret, Encyc. Botan. 5, p. 655.*

HAB. In Africâ Australi, pone Montes Swellendam. *D. Masson.* (v. s. in Herb. Banks. et D. Aiton.)

*30. *L. æmulum*, foliis superioribus lanceolato-spathulatis acutis rugoso-striatis, strobilis ovatis: squamis basi cohærentibus suprâ distinctis marginibus recurvis imberbibus, samaris subrotundis nigris.

Protea incurva. *And. Repos. 429*, fortè *Mas* hujus speciei, quamvis folia superiora vix duplo latiora.

HAB. In Africâ Australi. *Gul. Roxburgh M. D.* (v. s. in Herb. Soc. Linn.)

31. *L. abietinum*, foliis omnibus filiformibus canaliculatis obtusiusculis lævibus patulis arcuato-incurvis, strobili squamis marginibus axibusque infrâ connatis suprâ distinctis bilobis.

⁶⁵ *Protea teretifolia*. *And. Repos. 461*, femina et ramulus ad 4. mas.

HAB. In Africâ Australi, prope Prom. B. Spei, frequens. (v. s. in Herb. Banks. et Soc. Linn.)

*32. *L. scabrum*, foliis omnibus filiformibus canaliculatis acutis imbricatis rectiusculis margine scabris pilosisve, strobili squamis marginibus axibusque infra connatis apicibus distinctis bilobis.

HAB. In Africâ Australi. (v. s. in Herb. Hibbert.)

++++ Dubiæ tribus. Feminis adhucdum incognitis.

33. *L. sericeum*, foliis lanceolatis sericeis semiunguicularibus, caule erecto, capitulis sessilibus solitariis aggregatisve turbinatis, calycibus masculis longitudinaliter pubescentibus: tubo gracili infernè stylo cohærente.

Protea sericea. Linn. *Suppl.* 118,* fide specim. in ejus Herb.

Protea sericea. Thunb. *Diss. n.* 46,* sed caulis erectus videtur.

HAB. In Africâ Australi, prope Promont. B. Spei. (v. s. in Herb. Banks.)

34. *L. Globularia*, foliis lineari-spathulatis glabris aveniis: callo obtusissimo; basi attenuatâ tortâ, capitulis sessilibus depresso-globosis: bracteis tomentosis, calycibus masculis pubescentibus, stigmatibus clavato.

Protea globularia. Lam. *Illustr. Gen.* 1, p. 236, n. 1232, t. 53, f. 2. *Poirer, Encyc. Botan.* 5, p. 647? exclus. syn. Thunbergii.

Desc. *Fruticulus* erectus ramosissimus, ramis strictis, ultimis sericeis. *Folia* sparsa, 8—9 lineas longâ, inferiora ramorum breviora, capitulum subtendentia confertiora. *Bracteæ* involucrantès capitulo dimidio breviores, ovatæ, artè imbricatæ. *Calyx* tubo gracili unguibusque villosis, lamina glabris. *Ovarium* nullum. *Stylus* villosus. *Squamulæ* quatuor, lineares, longæ, basi styli infra adnatæ.

*35. *L. pubescens*, foliis spathulato-linearibus obtusis obliquis: adultis pubescentibus; junioribus sericeis, ramis villosis, involucris capitulo globoso sessili brevioribus tomentosis, calycibus stylisque pubescentibus.

HAB. In Africâ Australi. *Gul. Roxburgh M. D.* (v. s. in Herb. Soc. Linn.)

Obs. Quam maximè affine præcedenti.

*36 *L. ericifolium*, foliis acerosis glabris semiunguiculatis, capitulis corymbosis paucifloris, calycibus tomentosis.

HAB. In Africâ Australi. *Dom. J. Roxburgh.* (v. s. in Herb. Lambert. et Soc. Linn.)

DESC. *Frutex* erectus, ramosissimus, ramis rubicundis, ramulis tenuissimè tomentosis. *Folia* frequentia, erecta, imbricata, mutica, concaviuscula. *Capitula* breviter pedunculata, *Involucro* breviori sericeo subtensa. *Calyx* tubo gracili. *Ovarium* nullum. *Stylus* glaber. *Stigma* clavatum. *Squamulæ* nullæ intra calycem.

*37. *L. crassifolium*, foliis cuneato-obovatis obtusissimis glaberrimis crassis aveniis (3-uncialibus) basi attenuatis, capitulis globosis, bracteis propriis lanatis, calycibus glabris.

HAB. In Africâ Australi. *D. Masson.* (v. s. in Herb. Banks.)

DESC. *Rami* glaberrimi, glauci, crassitie digiti minimi. *Folia* frequentia, glauca, rigida, sesquiunciam lata, callo subrotundo, acutiusculo, marginibus subsimplicibus per lentem minutè crenulatis, novellis ciliatis. *Capitulum* magnitudine cerasi maximi. *Calycis Tubus* cylindraceus. *Laminæ* lineares, planiusculæ, unguisque recurvæ. *Stylus* glaber. *Stigma* oblongo-clavatum.

An hujus generis ?

*38. *L. cartilagineum*, foliis ovali-spathulatis obtusissimis: callo subtruncato: aveniis cartilagineis glaberrimis; basi attenuatâ lineari, capitulis globosis subpedunculatis, bracteis calycibusque tomentosis.

HAB. In Africâ Australi. (v. s. in Herb. Soc. Linn. et D. Hibbert.)

DESC. *Frutex* erectus, ramulis tenuissimè tomentosis. *Folia* vix uncialia, siccatione venis obsoletis depressis. *Capitulum* magnitudine cerasi minoris, pedunculo brevissimo

bracteisque villosis. *Calyx* quadrifidus. *Stylus* glaber.
Stigma oblongo-clavatum.
 An hujus generis?

3. PETROPHILA.

ATYLI species. *Salisb. Parad.*

GEN. CHAR. *Calyx* quadrifidus, totus simul deciduus. *Stylus* basi persistenti. *Stigma* fusiforme, apice attenuato. *Squamulæ* nullæ hypogynæ. *Strobilus* ovatus. *Nux* lenticularis, hinc comosa, v. *Samara* basi barbata.

HABITUS. Frutices rigidi. Folia glabra, varia, filiformia v. plana, indivisa, lobata v. pinnatifida, quandoque in eodem frutice diversiformia. Amenta ovata v. oblonga, terminalia et axillaria, nunc aggregata. Genus, posthac, speciebus incrementibus, dividendum, phalangibus infra propositis genera futura indicantibus.

ETYM. Πέτρος et φίλεω. Hi frutices enim semper in saxosis apricis proveniunt.

OBS. Mr. Salisbury has united such of the New Holland Proteæ as he had seen, into one genus, which he calls *Atylus*; a name meant to express the want of those bodies that usually surround the ovarium, in this order, and which he chooses to term *calli*: but as I conceive they are certainly secreting organs, the name on this ground would be exceptionable: my chief reason however for not adopting it, either for the present or the following genus, is, that the whole of his essential character does not apply to either of them. In his secondary character he has also considered them as monoicous, a mistake into which he has probably been led, not only from the striking similarity between the strobili of *Petrophila* and *Leucadendron*, but also from the style of the former remaining for some time unwithered after the calyx has fallen off. In one species he has even described the relative situation of the sexes; regarding the terminating amentum of *P. pulchella* as female, and the lesser ones, which frequently though not always surround

it, as male; but that this is not the case is proved by Cavanilles' figure of the species, in which all the amenta are in fruit, and a specimen in the same state may be seen in Sir Joseph Banks's Herbarium.

† *Stigma articulatum, articulo inferiore angulato, glabro, superiore tomentoso. Nux lenticulari-compressa, intus marginibusque comosa. Folia filiformia indivisa.*

*1. *P. teretifolia*, foliis teretibus exsulcis, squamis strobili enervibus, stigmatis articulo superiore stuposo triplo longiore.

⁶⁹⁾ HAB. In Novæ Hollandæ orâ australi, Lewin's Land. (ubi v. v.)

*2 *P. filifolia*, foliis teretibus exsulcis, squamis strobili nervosis orbiculatis, stigmatis articulo superiore barbato vix duplo longiore.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land; in collibus saxosis. (ubi v. v.)

*3. *P. acicularis*, foliis filiformibus suprâ obsoletè sulcatis, squamis strobili nervosis ovatis.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land; in campis collibusque sterilibus. (ubi v. v.)

†† *Stigma inarticulatum, hispidiusculum. Nux lenticulari-compressa, intus marginibusque comosa. Strobilus squamis distinctis. Folia filiformia bipinnatifida.*

*4. *P. rigida*, foliis triternatis: laciniis divaricatis, calycibus barbatis: laminarum apiculis glabris.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land; in collibus saxosis. (ubi v. v.)

5. *P. pulchella*, foliis trifido-bipinnatis: laciniis erectis, calycibus sericeis: laminarum apicibus tomentosis.

Protea pulchella. Schrad. Sert. Hanov. ii. p. 15. t. 7. Willd. Sp. Pl. 1, p. 507. Cavan. Anal. 1, p. 237.* Ic. 6, p. 33, t. 550. Sims, Bot. Mag. 796.*

Protea fucifolia. Salisb. Prod. 48.

Protea dichotoma. *Cavan. Anal.* 1, p. 239.* *Ic.* 6, p. 34,* t. 551.

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson; in arenosis inter saxa. (ubi. v. v.)

*6. *P. fastigiata*, foliis trifido-bipinnatis: laciniis erectis ⁽⁷⁰⁾ fastigiatis teretiusculis muticis, calycibus glabris, strobilis terminalibus sessilibus: squamis lanatis.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land; in ericetis aridis elevatioribus. (ubi. v. v.)

*7. *P. pedunculata*, foliis tripinnatifidis: laciniis canaliculatis divaricato-patulis, calycibus glabriusculis, strobilis pedunculatis: squamis glabris.

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson; in montibus saxosis. (ubi v. v.)

†† Strobilus *squamis connatis*. *Samara foliacea, dilatata*. *Stigma inarticulatum, hispidiusculum*. *Folia plana, bipinnatifida*.

*8. *P. diversifolia*, foliis bi-tripinnatifidis planis: laciniis mucronatis, calycibus barbatis, strobilis axillaribus pedunculatis: squamis lanatis cohærentibus.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land; in collibus saxiosis. (ubi v. v.)

††† Strobilus *squamis distinctis*. *Samara planiuscula*. *Folia plana, ternatim divisa*.

*9. *P. squamata*, foliis trifidis: lobis lineari-lanceolatis; lateralibus sæpiùs bi-trifidis, strobilis axillaribus sessilibus: squamis apice scariosis glabris.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land; in ericetis collibusque aridis. (ubi v. v.)

*10. *P. trifida*, foliis trifidis: lobis spathulato-lanceolatis sæpissimè indivisis, strobilis axillaribus sessilibus: squamis apice sericeis.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land; ⁽⁷¹⁾ in collibus apricis sterilibus. (ubi v. v.)

4. ISOPOGON.

ATYLI species. *Salisb. Parad.*

GEN CHAR. *Calyx* quadrifidus, tubo gracili, diutius persistente. *Stylus* totus deciduus. *Stigma* fusiforme, v. cylindraceum. *Squamæ* nullæ hypogynæ. *Nux* sessilis, ventricosa, undique comosa.

HABITUS. Frutices *rigidi*. Folia *glabra, plana v. filiformia, divisa v. integerrima*. Capitula *terminalia, rarò axillaria*. Floribus *modò densissimè imbricatis strobilo globoso; modò fastigiatis receptaculo communi planiusculo subinvolucrato, paleis deciduis congestis*. Genus distinctum, præcedenti proximum, posthac forsàn in duo dividendum, ratione inflorescentiæ secundum quam species infrà distributæ sunt, in duas phalanges habitu parùm diversas.

ETYM. *Ισος* et *πογον*, ob nuces undique æqualiter barbatus; quâ notâ a *Petrophilâ* faciliè distinguendus.

† *Strobilus globosus; squamis densissimè imbricatis, tardiùs deciduis.*

*1. *I. teretifolius*, foliis bi- v. triternatis filiformibus exsulcis, ramulis tomentosis, calycis tubo sericeo: laminis longitudinaliter barbatis.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land; in ericetis collibusque saxosis. (ubi v. v.)

2. *I. anethifolius*, foliis pinnatifidis bipinnatifidisque filiformibus suprâ sulcatis: laciniis erectiusculis, ramis glabris, calycis tubo pubescente: laminis infrâ glabris apice barbato.

⁷² *Protea anethifolia. Salisb. Prod.* 48.

Protea acufera. Cavan. Anal. 1, p. 236.* *Ic.* 6, p. 33, t. 549.

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson; in ericetis. (ubi v. v.)

*3. *I. formosus*, foliis bipinnatifidis subtriternatis fili-

formibus suprâ canaliculatis : laciniis divaricatis, ramulis tomentosis, calycibus glabris : laminis apice pilosiusculis.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land ; in collibus saxiosis. (ubi v. v.)

4. *I. anemonifolius*, foliis trifido-pinnatifidis bipinnatifidisve : laciniis linearibus planis patenti-erectis subtus lævibus, strobili squamis stuposis.

Protea anemonifolia. *Salisb. Prod.* 48. *Sims, Bot. Mag.* 697. *And. Repos.* 332.

Protea tridactylites. *Cavarr. Anal.* 1, p. 235.* *Ic.* 6, p. 33,* t. 548.

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson ; in ericetis saxiosis. (ubi v. v.)

*5. *I. ceratophyllus*, foliis trifido-bipinnatifidis : laciniis linearibus planis divaricato-patulis utrinque striatis mucronatis ; floralibus basi dilatatis, strobili squamis glabratiss.

HAB. In Novæ Hollandiæ orâ australi, prope Port Phillip ; in campis et collibus. (ubi v. v.)

*6. *I. trilobus*, foliis cuneatis planis trilobis : basi attenuatis petiolatis ; lobis integerrimis, ramulis tomentosis.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land ; in ericetis siccis. (ubi v. v.)

*7. *I. longifolius*, foliis lineari-lingulatis : superioribus integerrimis ; inferioribus passim trifidis, calycibus sericeis, stigmate glabro.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land ; in collibus saxiosis. (ubi v. v.)

†† *Receptaculum commune planum v. convexiusculum, paleis deciduis.*

*8. *I. cuneatus*, foliis oblongo-cuneatis obtusissimis, involucri bracteis tomentosis, calycibus glabris, stigmate fusiformi.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land. *D. A. Menzies.* (v. s. in Herb. Banks.)

*9. *I. attenuatus*, foliis elongato-oblongis mucronulatis basi attenuatis, rariis bracteisque involucrantibus glabris, capitulis solitariis, calycis laminis apice barbatis, stigmatè cylindræo.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land ; in collibus saxosis. (ubi v. v.)

*10. *I. polycephalus*, foliis lineari-oblongis mucronulatis, ramulis tomentosis, capitulis subaggregatis : bracteis omnibus lanatis, stigmatè cylindræo.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land ; in collibus saxosis. (ubi v. v.)

*11. *I. buxifolius*, foliis ovatis sessilibus acutis : apicibus recurvis, caulibus proliferis, capitulis solitariis foliis obvallatis ; bracteis subulatis, stigmatè fusiformi.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land ; in ericetis elevatioribus subhumidis. (ubi v. v.)

*12. *I. axillaris*, foliis cuneato-lingulatis mucronulatis, capitulis axillaribus paucifoliis ; bracteis involucrantibus ovatis imbricatis, calycis laminis longitudinaliter barbatis, stigmatè fusiformi.

HAB. In Novæ Hollandiæ orâ australi, Lewin's Land ; in collibus saxosis. (ubi v. v.)

5. PROTEA.

Linn. Gen. Pl. ed. i. Class. Plant. Leucadendron. Linn. Gen. Pl. ed. ii, v, et vi. Proteæ sp. Linn. Syst. Veg. xiii. Thunb. Diss. Juss. Gen. Erodendrum. Salisb. Parad.

GEN. CHAR. *Calyx* bipartibilis, inæqualis, labii latoris laminis staminiferis cohærentibus. *Stylus* subulatus. *Stigma* angustius, cylindræum. *Nux* undique barbata, stylo persistenti caudata. *Receptaculum* commune, paleis abbreviatis persistentibus. *Involucrum* imbricatum, persistens.

HABITUS. Frutices modò proceriores et quandòque arbo-

rescentes, modò subacaules. Folia integerrima. Capitula terminalia, rariusve lateralialia: Receptaculo planiusculo, nunc convexo, sæpissimè glabro, paleis quandòque connatis alveolato: Involucro magno, colorato, turbinato v. hemisphærico: Calycis labio latiore sæpè 2—3-aristato.

† *Capitula terminalia.* [73]

1. *P. Cynaroides*, foliis subrotundis petiolatis, involucris sericeis; bracteis interioribus acutis imberbibus, stylo infra medium pubescenti.

Scolymocephalos Africana lato rotundo glabro folio, cono maximo sericeo candido. *Herm. Cat. Mt. Raj. Hist.* 3, *Dendr.* p. 9.

Cinaroides frutex folio subrotundo rigido e Monte Tabulari. *Petiv. Mus.* 374.

Lepidocarpodendron; folio subrotundo, rigido, in pedunculo longo, crasso; flore maximo, purpureo. *Boerh. Lugd. Bat.* 2, p. 184,* c. tab. bona.

Scolymocephalos africanus folio lato rotundo. *Weinm. Phyt.* 4, p. 287, t. 892.

Leucadendron foliis subrotundis patentissimis petiolatis, foliolis calycinis carinatis. *Wachen. Ultraj.* 204.

Leucadendron cinaroides a. *Linn. Sp. Pl. ed.* i, p. 92, ed. ii, p. 135, *Berg. Act. Stockh.* 1766, p. 319.

Protea cynaroides. *Linn. Mant.* p. 190.* *Syst. Veg.* xiii, p. 118. *Thunb. Diss. n.* 59.* (exclus. syn. *Lepidocarpodendron*, &c. *Boerh. Lugd. Bat.* 2, p. 199, c. tab.) *Thunb. Prod.* 28. *Lam. Illust. Gen.* 1, p. 234, n. 1209. *Willd. Sp. Pl.* 1, p. 534. *And. Repos.* 288. bona. *Poiret, Encyc. Botan.* 5, p. 639. *Sims, Bot. Mag.* 770.

HAB. In Africâ Australi, ad latera montium prope Promont. B. Spei. (v. v. in Hort. var.)

*2. *P. latifolia*, foliis late-ovatis semicordatis sessilibus, involucro sericeo-tomentoso; bracteis interioribus angustatis apice dilatato barbato, calyce tomentoso: aristis hirsutis longitudine laminarum, stylo pubescenti.

HAB. In Africâ Australi. *Gul. Roxburgh M. D.* in 176

arenosis Zwartberg. *D. Niven.* (v. s. in Herb. Hibbert. et Lambert.)

DESC. *Frutex* 6—8 pedes altus. *Rami* tenuissimè tomentosi. *Folia* obtusissima, callo nullo prominulo, marginata, venosa, 3—4 uncias longa, 2—3 uncias lata, adulta glabra marginibus quandòque lanatis. *Involucrum* folia superiora superans, turbinatum, pugno majus: *Bracteis* obtusis, ciliatis; exterioribus latè ovatis; mediis oblongis; intimis elongatis, ungue lineari, laminâ oblongâ. *Calyx* ferè triuncialis, sericeo-tomentosus, labio latiore triaristato, aristis hirsutis, villis patulis, terminalibus præsertim purpureis. *Stylus* angulato-compressus, pube brevi adpressa subsericeus, apice glabro, curvato. *Stigma* subulatum, apice obtusiusculo.

*3. *P. compacta*, foliis oblongo-ovatis cordatis marginatis: callo apicis prominulo, involucro sericeo-tomentoso ciliato imberbi, calycis aristis longitudine laminarum, stylo glabro apice curvato, stigmatis apice conico-incrassato.

HAB. In Africæ Australis montosis, Hout Hoek. *D. Masson.* (v. s. in Herb. Banks.)

4. *P. longiflora*, foliis ovato-oblongis sessilibus basi subcordatis simplicibusve, ramis tomentosis, involucro sericeo; bracteis intimis elongatis sericeo-ciliatis, calycis aristis brevissimis, stylo glabro involucro longiore.

Conocarpodendron; folio subrotundo, crasso, rigido, valdè nervoso; cono longo, variegato, ex rubro et flavo; flore aureo. *Boerh. Lugd. Bat.* 2, p. 199, c. tab. bona respectu capituli, foliis vix convenientibus et potiùs ad *P. compactam* v. *latifoliam* pertinentibus: strobilo nucibus et flosculo ad calcem tabulæ jamjam ad *Leucadendron retusum* relatis.

77 Scolymocephalus foliis subrotundis glabris. *Weinm. Phyt.* 4, p. 294, tab. 902, b. a Boerh. icone mutuata omissio tamen strobilo.

Leucadendron, foliis subsessilibus cordato-ovatis imbricatis glabris. *Wachend. Ultraj.* 204. caractere ab icone Boerh. desumpto.

Protea longiflora. *Lam. Illust. Gen.* 1, p. 234, n. 1211, *Poiret, Encyc. Botan.* 5, p. 640.*

Protea lacticolor. *Salisb. Parad.* 27.

Protea ochroleuca. *Smith, Exot. Bot.* 2, p. 43, t. 81.

HAB. In Africæ Australis montibus, prope Prom. B. Spei. (v. v. in Hort. Reg. Kew.)

*5. *P. coccinea*, foliis obovatis obtusissimis sessilibus venosis ramisque glabris, involucris bracteis interioribus spathulatis apice barbatis, stylo glabro, calycis aristis ferè longitudine laminarum : margine pilosis ; apice imberbibus.

HAB. In Africæ Australis montibus, prope Promont. B. Spei, Devil's Head : solo fertiliori. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* 4—5 pedes altus. *Rami* crassitie digiti. *Folia* lævia, glauca, per lentem punctis minutissimis depressiusculis conspersa ; dum 4 uncias longa, 2—3 uncias lata ; superiora basi quandòque semicordata ; summa capitulum æquantia. *Involucrum* sessile, solitarium, turbinatum, 4—5-unciale, bracteis extùs demum glabriusculis ; interiorum barbâ marginali, copiosâ, longâ, persistenti. *Calyx* inclusus, 2½ uncias longus ; *Unguibus* hirsutis ; *Laminis* dorso glabris, margine pilosis ; *Aristis* vix longitudine laminarum. *Stylus* compressus. *Stigma* subulatum ; inde exsulcum a stylo absque manifestâ curvaturâ continuum.

6. *P. speciosa*, foliis obovato-oblongis basi angustatis 78 ramisque glabris, involucris bracteis omnibus sericeis : interioribus apice subdilatatis mediisque barbatis, stylo pubescenti, aristis calycis apice lanatis.

Scolymocephalus Africana foliis longis glabris, cono sericeo ; squamis rubiginèâ villosâ cristâ ornatis. *Herm. Cat. Mt. Raj. Hist.* 3, *Dendr.* p. 9.

Lepidocarpodendron ; folio oblongo, viridi, limbo rubro ornato ; squamarum apice, et margine, lanuginosis. *Boerh. Lugd. Bat.* 2, p. 185. c. tab.

Scolymocephalus foliis longis, seu *Tulipifer latifolius*. *Weinm. Phyt.* 4, p. 288, t. 893, a. bona.

Scolymocephalus Africanus foliis angustis villosis. *Weinm. Phyt.* 4, p. 289, t. 894?

Bruckm. Epist. Itin. 2, p. 8, t. 3, capitulum.

Leucadendron speciosum. *Linn. Mant.* p. 36.* excl. syn. *Clusii.*

Protea speciosa. *Linn. Mant.* p. 191.

Protea Lepidocarpodendron β. *Linn. Syst. Veg.* xiii, p. 118.

Protea barbata. *Lam. Illust. Gen.* 1, p. 236, n. 1228.

Protea speciosa latifolia. *And. Repos.* 110, fortè huc pertinet monente D. Bellenden Ker; at pessima figura.

Protea speciosa. *Sims, Bot. Magaz.* 1183.

HAB. In Africæ Australis montibus, prope Prom. B. Spei. (v. v. in Monte Tabulari.)

*7. *P. macrophylla*, foliis elongato-oblongis marginatis venosis glabris basi subattenuatis, involucri bracteis omnibus tomentosis; intimis lingulatis imberbibus, calycis aristis hirsutis, stylo exsulco infra medium pubescente: apice curvato.

⁷⁹¹ HAB. In Africâ Australi, ad latera Montium Attaquas Kloof. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* validus, 8—10 pedes altus. (*Niven.*) *Rami* glabri apice tomento brevissimo quasi rore canescenti obducti. *Folia* basi parùm attenuatâ tortâ; superiora longiora, involucri longè superantia, spithamea, ultra pollicem lata. *Involucrum* bracteis omnibus obtusis incanis; extimis ovatis; mediis oblongis; intimis apice haud dilatato. *Calyx* involucri parùm longior; unguibus laminisque tomento albo villisque concoloribus patulis: *Aristis* longitudine laminarum, tomento albo villisque longis, patulis, nigro-purpureis, terminalibus subcrispatis.

8. *P. formosa*, foliis angusto-oblongis venosis obliquis: basi simplicibus; marginibus ramisque tomentosis, involucri bracteis ciliatis; intimis lingulatis imberbibus, calycibus aristisque tomentosis, stylo glabro apice curvato, stigmate apice incrassato.

Protea coronata. *And. Repos.* 469.

- Erodendrum formosum*. *Salisb. Parad.* 76.
 HAB. In Africâ Australi. *D. Masson*. (v. s. in Herb. D. Aiton, e Hort. Reg. Kew.)
 OBS. Affinitate proxima *P. compactæ*, foliis præsertim diversa.

9. *P. melaleuca*, foliis lineari-lingulatis marginatis ciliatis, ramis pilosiusculis, involucris elongato-turbinatis: bracteis albociliatis; exterioribus squarrosis; interioribus conniventibus spathulatis dorso nigro-tomentosis.

Lepidocarpodendron; folio saligno, viridi; nervo et margine flavo; cono longo, superiore parte maximè clauso. [80] *Boerh. Lugd. Bat.* 2, p. 189,* c. tab.?

Scolymocephalus seu *Lepidocarpodendron* frutice conifero. *Weinm. Phyl.* 4, p. 291, t. 898? diversa tamen bracteis intimis viridibus fortè e descriptione in Cod. Witsen. pictis.

Protea coronata. *Lam. Illust. Gen.* 1, p. 236, n. 1227? exclus. syn. priore Boerhaavii. *Poiret, Encyc. Botan.* 5, p. 645? desc. conveniente.

Protea speciosa nigra. *And. Repos.* 103.

Protea Lepidocarpon. *Ker in Bot. Mag.* 674.

HAB. In Africâ Australi, prope Prom. B. Spei.

OBS. Species, ex figuris recentioribus huc citatis, nec non e pulcherrimâ ineditâ D. Franc. Bauer, quæ omnes inter se exactè conveniunt, distincta videtur, at quoniam specimina his respondentia nondum vidi, haud sine hesitatione a sequente separavi.

10. *P. Lepidocarpon*, foliis lineari-lingulatis marginatis scabriusculis nitentibus ramisque glabris, involucris bracteis interioribus spathulatis dorso marginibusque nigro-barbatis, calycis aristis intus pennatis, stylo pubescenti.

Scolymocephalus Africana, foliis longis glabris, cono variegato resinifero. *Herm. Cat. Mt.*

Scolymocephalus Africana, cono variegato resinifero. *Raj. Hist.* 3, *Dendr.* p. 9.

Lepidocarpodendron; foliis angustis, longioribus, saliginis; calycis squamis elegantissimè ex flavo fusco albo nigro

variegatis ; florum plumulis atro-purpureis. *Boerh. Lugd. Bat. 2, p. 188, c. tab.*

Scolymocephalus Africana, foliis longis, cono variegato. *Weinm. Phyt. 4, p. 289, t. 895.*

^{81]} *Protea* foliis lanceolatis integerrimis glabris calycinis supernè villosis. *Roy. Lugd. Bat. 186.*

Leucadendron *Lepidocarpodendron*. a. *Linn. Sp. Pl. ed. i, p. 91, ed. ii, p. 134. Berg. Act. Stockh. 1766, p. 322.*

Protea *Lepidocarpodendron*. *Linn. Mant. 190,* desc. opt. nullo tamen specimine in Herb.*

Protea *Lepidocarpodendron* a. *Linn. Syst. Veg. xiii. p. 118.*

Protea speciosa. *Thunb. Diss. n. 53,* Prod. 27. Willd. Sp. Pl. 1, p. 531.*

Protea cristata, *Lam. Illustr. Gen. 1, p. 235, n. 1226. Poiret, Encyc. Botan. 5, p. 644, exclus. syn. Roy. Linn. et Andr.*

Protea grandiflora var. foliis undulatis. *And. Repos. 301?*

HAB. In Africæ Australis montibus, prope Prom. B. Spei. (v. v. in Monte Tabul.)

*11. *P. neriifolia*, foliis lineari-lingulatis lævibus opacis margine subsimplicibus basi extùs ramisque tomentosis, involucri bracteis interioribus apice parùm latioribus dorso argenteo-sericeo margine nigro-barbato, calycis aristis laminas superantibus intùs pennatis, stylo pubescenti.

Cardui generis elegantissimi cujusdam caput. *Clus. Exot. 38.* fig. xv.*

HAB. In Africâ Australi, ad radices montium prope Prom. B. Spei. (v. s. in Herb. Soc. Linn.)

OBS. I. Quam maximè affinis *P. Lepidocarpo*, at distincta videtur.

OBS. II. Synonymon Clusii huc retuli ob descriptionem optimè convenientem.

12. *P. pulchella*, foliis lineari-lingulatis marginatis niten-
^{82]} tibus scabriusculis, ramis parùm tomentosis, involucri bracteis interioribus apice lanceolato-dilatato sericeo marginibus nigro-barbatis, calycis aristis vix longitudine laminarum, stylo pubescenti.

Protea pulchella. *And. Repos.* 270, bona quoad capitulum, sed folia opaca margine ciliata.

Protea speciosa, var. foliis glabris. *And. Repos.* 277, optima respectu capituli et foliorum nitore quæ autem margine concolori diversa.

Protea pulchella var. *speciosa*. *And. Repos.* 442, differt figura bractearum interiorum aristisque calycis laminâ longioribus.

HAB. In Africæ Australis montibus, prope Stellenbosch. *Gul. Roxburgh, M. D.* (v. s. in Herb. Banks. et Soc. Linn.)

13. *P. patens*, foliis angusto-oblongis subundulatis marginatis basi subattenuatis, ramisque villosis procumbentibus, involucri hemisphærico: bracteis sericeis; interiorum barba nigro-purpurea, stylo infra pubescenti, calycis aristis longitudine laminarum.

Protea speciosa patens. *And. Repos.* 543.

HAB. In Africæ Australis montibus saxosis, prope Wilde River. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* procumbens. (*Niven.*) *Rami* tomentosi et villis patulis brevibus incani. *Folia* secunda, frequentia, obtusiuscula, venosa, 4—5 uncias longa, 7—9 lineas lata. *Involucri* sessile, magnitudine pugni minoris: *Bracteis* obtusis, albo-sericeis, concaviusculis, interioribus nec dilatatis nec angustatis mediisque barbâ nigro-purpureâ instructis. *Calyx* sesquiuncialis albo-lanatus, aristis apice purpureis. *Stylus* basi compressâ tomentosâ suprâ subulatus et infra medium pube rarâ, suprâ glaber, apice curvato. *Stigma* acutiusculum.

*14. *P. incompta*, foliis lingulato-oblongis: summis ramisque hirsutis, involucri bracteis interioribus apice orbiculato-dilatato margine barbato, calycis lanati aristis longitudine laminarum, stylo glabro apice simplici.

Protea foliis lanceolatis integerrimis glabris calycem succingentibus hirsutis. *Roy. Lugd. Bat.* 186? exclus. syn. Boerh. *t.* 189.

HAB. In Africâ Australi. *Oldenburgh*: prope Wynberg.

Gul. Roxburgh, M. D. (v. s. sub eodem nomine in Herb. Banks.)

DESC. *Frutex* erectus. *Rami* hirsutissimi villis longis patulis. *Folia* frequentia, modicè patentia, 4 uncias longa, 1 unciam lata, venosa, basi obtusa, marginibus simplicibus; callo apicis acuto, recurvo; inferiora glabra; summa angustiora, capitulum paulò superantia. *Involucrum* turbinatum, 4 uncias longum, bracteis tomentosis; exterioribus mediisque oblongis, imberbibus; interioribus barbâ marginali, albâ. *Calyx* lanâ albâ, implexâ.

15. *P. longifolia*, foliis elongato-linearibus basi attenuata, involucri turbinati bracteis glabris acutis imberbibus, calycis aristis laminâ longioribus, stylo pubescenti apice curvato.

Lepidocarpodendron; foliis angustis, longis, salignis nervo rubro; florum plumis violaceo-purpureis. *Boerh. Lugd. Bat. 2, p. 186,* c. tab.*

Protea longifolia nigra. And. Repos. 132.

Protea longifolia var. cono turbinato. *And. Repos. 144.*

Protea longifolia ferrugineso-purpurea. And. Repos. 133.

^{84]} *Protea Lepidocarpodendron. Herb. Linn.*

HAB. In Africâ Australi, prope Prom. B. Spei. *Oldenburgh. (v. s. in Herb. Banks.)*

16. *P. mellifera*, foliis lanceolato-lingulatis basi attenuatis, involucro turbinato: bracteis glabriusculis imberbibus viscidis, aristis calycis albo-lanatis longitudine laminarum, stylo glabro: apice simplici.

Scolymocephalus Africana, foliis longis acutioribus hirsutis, cono mellifero. *Herm. Cat. Mt.*

Conifera Alypi folio seminibus pennatis, pluribus in medio coni conglomeratis, et non inter squamas aliorum conorum more nascentibus! *Sloane in Philos. Trans. 17, p. 666,* c. tab.*

Scolymocephalus Africana, foliis angustis villosis, cono mellifero. *Raj. Hist. 3. Dendr. p. 9¹*

Lepidocarpodendron; foliis angustis, brevioribus, salignis; calycis squamis elegantissimè ex roseo aureo albo atro-

¹ The point of interrogation is added in M.S. by Mr. Brown in his own separate copy.—EDIT.

rubro variegatis; florum plumis albis. *Boerh. Lugd. Bat. 2, p. 187, c. tab.*

Scolymocephalus seu *Lepidocarpodendron* folio saligno. *Weinm. Phyt. 4, p. 289, t. 896.*

Protea caule multifloro calycibus oblongis foliis lanceolatis integerrimis. *Roy. Lugd. Bat. 185.*

Leucadendron repens a. *Linn. Sp. Pl. ed. i, p. 91, ed. ii, p. 135.*

Leucadendron repens. *Berg. Act. Stockh. 1766, p. 322.*

Protea repens. *Linn. Mant. 189,* Syst. Veg. xiii, p. 118.*

Protea mellifera. *Thunb. Diss. n. 37.* Prod. 26. Lam. Illust. Gen. 1, p. 236, n. 1229. Salisb. Prod. 49. Willd. Sp. Pl. 1, p. 522. Poiret, Encyc. Botan. 5, p. 646. Curt. Mag. 346. Wend. Hort. Herrenh. 13.*

HAB. In Africæ Australis collibus et campis, prope Prom. B. Spei, gregatim quandoque crescens. (v. v. prope Constantiam.)

17. *P. grandiflora*, foliis oblongis sessilibus ramisque glabris, involucri hemisphærico imberbi nudiusculo, calyce tomentoso; unguibus dorso glabriusculis; aristis brevissimis, stylo glabro.

Scolymocephalus foliis oblongis glabris crassioribus latioribus. *Herm. Cat. Mt.*

Lepidocarpodendron; folio saligno lato; caule purpureo. *Boerh. Lugd. Bat. 2, p. 183,* c. tab.*

Scolymocephalus foliis oblongis. *Weinm. Phyt. 4, p. 28, t. 891.*

Protea foliis lanceolatis integerrimis flore patente glabro stylis longissimis. *Roy. Lugd. Bat. 186.*

Protea cinaroides β. *Linn. Sp. Pl. ed. i, p. 92, ed. ii, p. 136.*

Protea grandiflora. *Thunb. Diss. n. 51.* Prod. 27. Willd. Sp. Pl. 1, p. 530. Lam. Illust. Gen. 1, p. 234, n. 1210? Poiret, Encyc. Botan. 5, p. 640?*

β. *Protea marginata.* *Lam. Illust. Gen. 1, p. 235, n. 1225.*

HAB. In Africæ Australis collibus et montibus, frequens. (v. v. in Monte Tabulari.)

OBS. Variat involucri penitus glabro bracteisque exteri-

oribus albo-tomentosis. *Folia* quandoque lineari-oblonga et tunc ab icone *P. abyssinica* haud distinguenda.

18. *P. Abyssinica*, foliis lanceolato-lingulatis obtusiusculis basi subangustatis, involucri hemisphærico: bracteis obtusis imberbibus, calyce tomentoso; aristis brevissimis, receptaculo villosa? caule arborescenti.

Gaguedi. *Bruce. Abyss.* 5, p. 52, c. tab. duab.

⁸⁶¹ *Protea abyssinica*. *Willd. Sp. Pl.* 1, p. 522.

HAB. In Abyssiniâ, Lamalmon. *Bruce l. c.*

19. *P. Scolymus*, foliis lineari-lanceolatis acutis submucronatis basi attenuatis, involucri hemisphærico; bracteis glabris obtusis, calycibus muticis, receptaculo villosa, caule ramoso multifloro.

Thymelæa capitata rapunculoides æthiopica saligneis foliis perianthio magno conformi squamoso. *Pluk. Mant.* 181, t. 440, f. 1, mala.

Scolymocephalus fruticis Æthiopici coniferi Breynii foliis; capite majore squamato. *Raj. Hist.* 3. *Dendr.* p. 10.

Lepidocarpodendron; acaulon; ramis numerosis e terrâ excrescens, calyce floris immaturo extus ex rubro et flavo variegato intus flavo. *Boerh. Lugd. Bat.* 2, p. 192, c. tab.

Scolymocephalus foliis angustis longis. *Weinm. Phyt.* 4. p. 288, t. 893, f. b.

Leucadendron Scolymocephalum. *Linn. Sp. Pl. ed. i,* p. 92, ed. ii, p. 135. *Berg. Act. Stockh.* 1766, p. 323.

Protea Scolymus. *Thunb. Diss. n. 36.* Prod.* 26. *Lam. Illustr. Gen.* 1, p. 236, n. 1231. *Willd. Sp. Pl.* 1, p. 522. *Poiret, Encyc. Botan.* 5, p. 647. *And. Repos.* 409. *Wend. Sert. Hanov. t. 20.* *Sims, Bot. Mag.* 698.

Protea angustifolia. *Salis. Prod.* 49.

HAB. In Africae Australis ericetis elevationibus, prope Prom. B. Spei. (v. v. in Hort. Angl.)

20. *P. mucronifolia*, foliis lanceolato-linearibus mucronatis pungentibus basi obtusa, bracteis involucri lanceolatis mucronatis glabris, caule erecto multifloro.

Protea mucronifolia. *Salisb. Parad.* 24. *Sims, Bot.* 187 *Mag.* 933. *And. Repos.* 500.

Protea odoratissima. *Masson, in Herb. Ait.*

HAB. In Africâ Australi. *Masson*. In arenosis prope Berg River. *Niven*. (v. v. in Hort. Hibbert.)

21. *P. nana*, foliis subulatis mucronatis, involucris nantibus hemisphæricis; bracteis glabris obtusis.

Thymelæa æthiopica abietiformis floribus phœniceis. *Pluk. Mant.* 180.

Leucadendron nanum. *Berg. Act. Stockh.* 1766, p. 325.* *Berg. Cap.* 22,* exclus. syn. Petiv. ad feminam *Aulacis pinifoliae* jam citato.

Protea rosacea. *Linn. Mant.* p. 189,* *Syst. Veg.* xiii, p. 118.

Protea nana. *Thunb. Diss. n.* 29.* *Prod.* 26. *Hort. Kew.* 3, p. 484. *Willd. Sp. Pl.* 1, p. 519.

Protea rosacea. *Lam. Illust. Gen.* 1, p. 238, n. 1251. *Poiret, Encyc. Botan.* 5, p. 653. *Smith, Exot. Bot.* 1, p. 85, t. 44.

Protea acutifolia. *Salisb. Parad.* 2.

HAB. In Africæ Australis montosis; prope Roode Zant Cascade. (v. s. in Herb. Linn. a Bergio.)

OBS. Nomen Cel. Bergii utpotè primum, nec ineptum et a Thunbergio, Dryandro et Willdenovio receptum, prætuli.

*22. *P. pendula*, foliis lineari-lanceolatis mucronulatis: terminalibus ramorum floriferorum recurvorum reclinatis, bracteis involucris obtusis demùm glabriusculis.

HAB. In Africâ Australi. *Masson*. (v. s. in Herb. Banks.)

DESC. *Frutex* erectus. *Rami* teretes, glabri; ultimi tenuissimè tomentosi: floriferi supra medium recurvi. *Folia* sparsa, passim subopposita, frequentia, modicè patentia; extra medium parùm latiora, obtusa, mucronulo patenti, marginibus subrecurvis, glauca, compacta, adulta glaberrima, sesquiunciam longa. *Involucra* pendula, solitaria, hemisphærica, magnitudine pruni: *Bracteis* arcuè imbricatis, imberbibus, exutâ pube tenuissimâ sericâ demùm glabriusculis; interioribus sensim longioribus. *Calyces*

inclusi, submutici, laminis barbatis. *Stylus* glaber, vix uncialis, apice simplici.

23. *P. tenax*, foliis lineari-lanceolatis planis: basi attenuatis; margine scabriusculis, ramis decumbentibus, involucri hemisphærico sericeo obtuso, calycis (uncialis) unguibus glabriusculis: aristis lanatis laminâ dimidio brevioribus.

Erodendrum tenax. *Salisb. Parad.* 70.

HAB. In Africæ Australis depressis, Lange Kloof. *D. Niven.* (v. s. in *Herb. Soc. Linn.*)

DESC. *Frutex* diffusus. *Rami* glabri v. hirsuti. *Folia* 4—6 uncias longa, 4—6 lineas lata, acuta, uninervia, venis obsolete, minute punctulata, scabriuscula; ramorum subsecunda. *Involucrum* bracteis concavis, tenuissimè ciliatis, exterioribus ovatis; interioribus oblongis. *Calyx* unguibus suprà pilosiusculis; laminis dorso nudiusculis. *Stylus* glaber, apice simplici.

24. *P. canaliculata*, foliis linearibus aveniis lævibus: suprà concaviusculis; ramisque glabris decumbentibus, involucri obtuso: bracteis interioribus subsericeis, calycis unguibus glabris: aristis pectinatis laminæ dimidio brevioribus.

Protea canaliculata. *And. Repos.* 437.

HAB. In Africæ Australis depressis arenosis, Lange Kloof. *D. Niven.* (v. s. in *Herb. Lambert.*)

^{89]} DESC. *Frutex* subdecumbens. (*Niven.*) *Folia* frequentia, 4—6 uncias longa, vix duas lineas lata, acuta. *Involucrum* magnitudine pruni, *Bracteis* concavis; exterioribus glabris; interioribus brevissimè ciliatis. *Calyx* uncialis, unguibus laminisque glabris: Aristis albo-barbatis. *Stylus* glaber apice simplici.

25. *P. acaulis*, caulibus abbreviatis, ramis depressis, foliis obovato-oblongis marginatis venosis basi attenuatis, involucris hemisphæricis inclinatissimis; bracteis obtusis glabris, calycibus muticis.

Scolymocephalus Africana foliis oblongis glabris humilis et procumbens. *Herm. Cat.* 19.

Scolymocephalus Africanus Lauri folio humilis et procumbens. *Raj. Hist.* 3. *Dendr.* p. 9.

Lepidocarpodendron; acaulon; foliis paucis, latis, nervo et marginibus rubris ornatis; fructu parvo. *Boerh. Lugd. Bat.* 2, p. 191,* c. tab.

Scolymocephalus s. *Lepidocarpodendron* acaulon. *Weinm. Phyt.* 4. p. 291, t. 898, h. bona.

Protea caule unifloro foliis lanceolatis. *Roy. Lugd. Bat.* 186.

Leucadendron acaulon. *Wachend. Ultraj.* 204. *Linn. Sp. Pl. ed. i, p. 92, ed. ii, p. 135.* Syst. Nat. xii, t. 1, p. 110, omitta in Mant. et Syst. Veg. xiii.*

Protea acaulis. *Thunb. Diss. n. 49.* Prod.* 27. *Willd. Sp. Pl.* 1, p. 529.

Protea rana. *Lam. Illust. Gen.* 1, p. 233, n. 1208. *Poiret, Encyc. Botan.* 5, p. 639.

Protea glaucophylla. *Salisb. Parad.* 11.

HAB. In Africæ Australis collibus, prope Promont. B. Spei. (v. v. juxta Simons Bay.)

*26. *P. elongata*, caulibus nanis, foliis elongato-lanceolatis (pedalibus) planis marginatis venosis lævibus; basi valdè attenuata lineari, involucri hemisphærico inclinato; bracteis glabris obtusis, calycibus brevissimè aristatis.

HAB. In Africæ Australis humidis elevationibus. Roode Zant Cascade. *D. Niven.* (v. s. in Herb. Hibbert.)

Obs. Nimis affinis *P. acauli*.

*27. *P. angustata*, caulibus nanis, foliis lanceolato-lineari-bus planis marginatis venosis lævibus, involucri hemisphærico inclinato; bracteis glabris obtusis, calycibus muticis; unguibus extùs glabris margine lanatis.

HAB. In Africæ Australis montosis solo fertiliori; Hout Hoek. *D. Niven.* (v. s. in Herb. Hibbert.)

Obs. An species distincta a *P. acaule*?

*28. *P. revoluta*, caulibus nanis, foliis canaliculato-semi-teretibus lævibus, involucri hemisphærico inclinato: bracteis

glabris obtusis, calycibus muticis : unguibus extùs glabris margine lanatis.

HAB. In Africæ Australis montibus aridis. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* humilis, basi divisus. *Rami* adscendentes, glabri, vix longitudine foliorum. *Folia* 6—9 uncias longa, acuta, impunctata, marginibus recurvis, simplicibus, canaliculata, infra medium teretiuscula parùmque attenuata. *Involucrum* brevissimè pedunculatum magnitudine pruni minoris. *Calycis* laminæ sericæ. *Stylus* glaber, apice simplici.

*29. *P. tenuifolia*, caulibus nanis, foliis canaliculato-
 91] semiteretibus scabris, involucri hemisphærico : bracteis tomentosis, calycis unguibus laminisque hirsutis : aristis brevissimis.

HAB. In Africæ Australis montibus sterilibus. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Folia* numerosa, punctis elevatis utrinque scabra, marginibus revolutis canaliculata, basi planiuscula, spithamea v. dodrantalia. *Involucrum* erectum, sessile, magnitudine pomi minoris, tomento ferrugineo tardiùs deciduo. *Calyx* sesquiuncialis laminarum villis brevioribus, aristis (mucronibus potiùs) duabus lamina quadruplo brevioribus. *Stylus* glaber, apice simplici.

*30. *P. lævis*, caulibus nanis decumbentibus, foliis elongato-linearibus lævibus aveniis marginibus recurvis, involucri hemisphærico : bracteis obtusis subsericeis, calycibus subuncialibus muticis.

HAB. In Africâ Australi. *D. Masson.* (v. s. in Herb. Banks.)

DESC. *Caulis* brevissimus, decumbens (*Masson.*) glaber. *Folia* secunda, glauca, spithamea, acuta, marginibus lævibus non incrassatis, basi attenuata plana. *Involucrum* sessile, erectum, magnitudine pomi minoris ; *Bracteis* primùm subsericeis, demùm glabriusculis, marginibus brevissimè ciliatis. *Calyx unguibus* extùs glabriusculis, margine lanatis ; *Laminis* villosis.

*31. *P. scabra*, caulibus nanis, foliis elongato-linearibus

scabris obsoletè venosis margine subrecurvis, involucrio turbinato-hemisphærico: bracteis obtusis tomentosis, unguibus calycis hirsutis: aristis laminâ dimidio brevioribus.

HAB. In Africâ Australi, prope Promont. B. Spei. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Caulis* semisepultus, divisus, ramis adscendenti-⁹²bus, folio brevioribus. *Folia* conferta, erecta, spithamea, vix pedalia, 3—4 lineas lata, apice acuto sphacclato, uninervia, obsoletè venosa, utrinque tuberculis pustuliformibus scabra, aliisque minutissimis conspersa, basi attenuatâ petioliformi lævi. *Involucrum* sessile, erectum, magnitudine pomi minoris; *Bracteis* tomento ferrugineo demùm subdeciduo. *Calyx* vix semuncialis; laminis hirsutis; aristis villis flexuosis cinereis ferrugineisve.

OBS. In Herbario D. Hibbert plantam vidi Foliis planis elongato-lanceolatis; Involucris turbinatis; Calycibus albolanatis aristarum lana longiore magisque implexa; Stylo biunciali, vix arcuato: an distincta species?

32. *P. repens*, caulibus nanis, foliis elongato-linearibus scabriusculis margine revolutis, involucrio turbinato: bracteis obtusis tomentosis: interioribus margine lanatis, calycibus biuncialibus; unguibus hirsutis; aristis laminâ brevioribus, stylo apice simplici.

Lepidocarpodendron; foliis longissimis, angustissimis, fructum elegantissimè ex rubro flavo et albo variegatum instar coronæ succingentibus; radice repente. *Boerh. Lugd. Bat.* 2, p. 190,* c. tab.

Scolymocephalus s. *Lepidocarpodendron* foliis longissimis. *Weinm. Phyt.* 4, p. 290, t. 897, a.

Protea caule unifloro calyce oblongo foliis linearibus longissimis. *Roy. Lugd. Bat.* 185.

Leucadendron, foliis longissimis obtusè trigonis longitudine florem superantibus. *Wachend. Ultraj.* 204.

Leucadendron repens β. *Linn. Sp. Pl. ed. i*, p. 92, ed. ii, p. 135.

Protea repens. *Thunb. Diss. n.* 38.* *Prod.* 26. *Lam.* ¹⁹³*Illust.* 1, p. 236, n. 1230. *Willd. Sp. Pl.* 1, p. 523. *Poiret, Encyc. Botan.* 5, p. 646.

HAB. In Africæ Australis campis arenosis prope Prom.
B. Spei. (v. s. in Herb. Banks.)

Obs. Varietas? foliis vix punctatis, sesquipedalibus.

*33. *P. lorea*, caulibus nanis, foliis teretibus elongatis lævibus, involucro turbinato sub-pedunculato: bracteis acutiusculis sericeis, calycis unguibus extus glabris: aristis laminâ brevioribus, stylo apice curvato.

HAB. In Africâ Australi, prope Promont. B. Spei. *D. Masson.* (v. s. in Herb. Banks.)

DESC. Caulis brevissimus, semisepultus. *Folia* numerosa, pedalia, crassitie fili ligaterii. *Involucrum* pedunculo brevi squamis arcuè imbricatis tecto: *Bracteis* exterioribus ovatis acutiusculis, interioribus oblongo-linearibus. *Calyx* Unguibus Laminisque extus *Aristis* undique lana brevi densa alba crispata. *Stylus* glaber.

34. *P. turbiniflora*, caulibus nanis, foliis elongato-lanceolatis marginatis subundulatis lævibus, involucro subturbinato: bracteis tomentosis obtusis, calycis aristis longitudine laminarum: lana apicis longiore crispa.

Erodendrum turbiniflorum. *Salisb. Parad.* 108.

Protea cæspitosa. *And. Repos.* 526.

HAB. In Africæ Australis montibus; in humidis solo fertiliori. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Caules* cæspitosi, abbreviati, divisi, erecti. *Folia* uninervia, venosa, juniora villosa, adulta glabra, nitida, minutissimè punctata, acutissima, basi valdè attenuatâ petioliformi, spithamea, vix pedalia, unciam circiter lata; extima ramorum nana, biuncialia, basi vix attenuata membranacea, subscariosa. *Involucrum* sessile, vix biunciale: *Bracteis* subincanis, ciliatis, interioribus apice lanatis. *Calyx* unguibus laminisque lanatis: *Aristis* curvatis, albo-lanatis, lanâ terminali fulvâ. *Stylus* glaber, apice levissimè curvato.

*35. *P. Scolopendrium*, caulibus nanis, foliis elongato-lanceolatis marginatis lævibus, involucro turbinato: bracteis

lanceolatis acuminatis apice tomentosis, aristis calycis lamina dimidio brevioribus.

HAB. In Africâ Australi, Wintershoek. *D. Joh. Roxburgh.* (v. s. in Herb. Lambert.)

DESC. *Caulis* foliis aliquoties brevior. *Folia* pedalia, sesquiunciam vix duas uncias lata, costâ subtùs eminente, venis ramosis minutissimè punctata, basi valdè attenuata. *Involucra* subsessilia, solitaria v. bina, quandoque tres uncias longa: *Bracteis* interioribus apice tomento persistente cinereis. *Calyx* lanatus. *Stylus* glaber, infra medium dilatatus. *Ovarii* barba alba.

†† *Flores laterales.*

36. *P. cordata*, floribus lateralibus, foliis cordatis subrotundis nervosis, bracteis involucri glabris.

Protea cordata. *Thunb. Diss. n. 60,* tab. 5, bona. Prod. 28. Lam. Illust. Gen. 1, p. 233, n. 1207. Willd. Sp. Pl. 1, p. 534. Poiret, Encyc. Botan. 5, p. 639. And. Repos. 289.*

Protea cordifolia. *Sims, Bot. Mag. 649.*

HAB. In Africæ Australis montibus, Hottentots Holland et prope Fluvium Zonder End. *Thunb. l. c.* (v. s. in Herb. Banks.)

37. *P. amplexicaulis*, floribus lateralibus, foliis cordatis ovatis amplexicaulibus divaricatis apice recurvis, bracteis involucri pubescentibus.

Erodendrum amplexicaule. *Salisb. Parad. 67.*

Protea repens. *And. Repos. 453.*

HAB. In Africâ Australi. *D. Masson.* (v. s. in Herb. Banks.)

38. *P. humilis*, floribus lateralibus, foliis linearibus acutis, (biuncialibus,) receptaculo conico: paleis acutis.

Protea humiflora. *And. Repos. 532.*

HAB. In Africâ Australi. *D. Masson.* (v. s. in Herb. Banks.)

DESC. *Caulis* nanus. *Rami* glabri. *Folia* plana, fere

3 uncias longa. *Involucra* hemisphærica, bracteis obtusis, interioribus apice pube adpressâ ferrugineâ.

*39. *P. acerosa*, floribus lateralibus, foliis subulatis, receptaculo convexiusculo : paleis obtusis.

HAB. In Africâ Australi. *D. Masson.* (v. s. in Herb. Banks.)

DESC. *Caulis* brevis. *Rami* erecti, glabri. *Folia* lævia. *Involucra* ramea, subaggregata, breviter pedunculata; *Bracteis* obtusis, interioribus pube diutius persistenti subsericeis. *Calyx* muticus, apice barbato. *Receptaculi* paleæ connatæ. *Squamulæ hypogynæ* subulatæ.

Obs. Varietas? Foliis longioribus (sesquiuncialibus) semiteretibus in Herbario et Hort. D. Hibbert vidi, quæ secundum *D. Niven* 3—4 pedes alta in montosis solo fertiliori prope Zonder End lecta. Hæc *Protea virgata*. *And. Repos.* 577.

6. LEUCOSPERMUM.

LEUCADENDRUM. *Salisb. Parad.* Proteæ sect. 3. *Linn.*

Mant. CONOCARPODENDRA (spuria 196 et 198). *Boerh. Lugd.*

96] CHAR. GEN. *Calyx* irregularis, labiatus, unguibus tribus (rarò omnibus) cohærentibus, laminis staminiferis distinctis. *Stylus* filiformis, deciduus. *Stigma* incrassatum, glabrum (nunc inæquilaterale). *Nux* ventricosa, sessilis, lævis. *Capitulum* indefinitè multiflorum : *Involucro* polyphyllo imbricato.

HABITUS. Frutices sæpe humiles, quandoque arborescentes, plerique tomentosi v. hirsuti. *Folia* integra v. apice callosodentata. *Capitula* terminalia; *Floribus* flavis, modò imbricatis bracteis distinguendis persistentibus induratis; modò fastigiatis receptaculo planiusculo, paleis angustis, non mutatis, subdeciduis.

† *Capitulum amentaceum*; *Bracteis propriis persistentibus subinduratis.*

1. *L. linearis*, stylo calycem hirsutum superante, stigmatē hinc gibboso, involucro tomentoso, foliis linearibus integris; callo apicis subbarbato, ramis glabris.

Protea linearis. *Thunb. Diss. n. 35,* tab. 4*, pedunculo insolitè elongato stylisque apice nimis arcuatis. *Thunb. Prod. 26. Lam. Illust. Gen. 1, p. 237, n. 1241. Willd. Sp. Pl. 1, p. 521. Poiret, Encyc. Botan. 5, p. 650.*

HAB. In Africæ Australis arenosis, Paarl, Drakenstein, Stellenbosch. (v. s. in Herb. Banks. Lambert. Soc. Linn.)

Obs. Folia sæpius canaliculata marginibus inflexis, nunquam reflexis, callo apicis villis albis diu tecto; dum plana obsoletè striata marginibus scabriusculis; rarissimè 2—3-dentata.

*2. *L. attenuatum* stylo calycem hirsutum superante, stigmatē subæquilaterali, foliis cuneato-linearibus tridentatis aveniis basi attenuata, involucris ramisque tomentosis.

HAB. In Africæ Australis arenosis elevationibus inter saxa; Zwellendam. *D. Niven.* (v. s. in Herb. Banks. Lambert. Hibbert.)

DESC. *Frutex* erectus, tripedalis. *Rami* stricti, crassitie pennæ anserinæ, incani. *Folia* glaberrima, lævia, basi uninervi vix tortâ, crassa, rarè 5-dentata, sesquiuncialia, biuncialia. *Capitula* solitaria v. gemina, breviter pedunculata, obovata, magnitudine pruni majoris. *Bracteæ involucri* ovatæ, acuminatæ, arctè imbricatæ; *pedunculi* patulæ. *Stylus* calyce unam quartam longior. *Stigma* conico-ovatum.

Obs. Hujus Varietas? insignis. Foliis latoribus, apice profundè tridentatis, dentibus lateralibus sæpissimè bi-intermedio tri-dentatis. Ramulis præter tomentum incanum villis patulis brevibus. (v. s. in Herb. Hibbert.)

3. *L. Tottum*, stylo calycem hirsutum $\frac{1}{4}$ superante, stigmatē hinc gibboso, foliis lineari-oblongis sub-integris venosis basi obtusâ, bracteis involucri glabris ciliatis.

Protea Totta. *Linn. Mant. 191,* fide spec. in illius Herb.*

Thunb. Diss. n. 54. Prod. 27. Lam. Illust. Gen. 1, p. 235, n. 1224. Willd. Sp. Pl. 1, p. 532. Poiret, Encyc. Botan. 5, p. 644.*

HAB. In Africæ Australis montosis; Roode Zant Cascade. (v. s. in Herb. Linn. Banks., &c.)

Obs. *Frutex* subdecumbens (secund. *D. Niven*). *Rami* sæpiùs hirsuti, quandoque glabri. *Folia* interdum 2—3-dentata, venis obsoletis. *Calyces* bracteis triplo longiores. *Stigma* indivisum.

*4. *L. medium*, stylo calycem hirsutum ferè bis superante, stigmatè hinc gibboso, foliis lineari-oblongis integris passimque 2—3-dentatis: cullis acutis; basi obtusâ, bracteis involucri tenuissimè pubescentibus ciliatis, capitulis cernuis. ^{98]} *Protea formosa. And. Repos. 17?* quæ differt tamen, Foliis longioribus, Calycibus unilabiatis unguibus omnibus longitudinaliter cohærentibus, Bracteis involucri sphacelatis, Stigmatè ovato-oblongo vix gibboso.

HAB. In Africæ Australis montibus. (v. s. in Herb. Soc. Linn.)

Obs. Species inter *L. Totum* et *ellipticum* media, illo foliis, hoc floribus ferè exactè conveniens.

5. *L. ellipticum*, stylo calycem hirsutum ferè bis superante, stigmatè conico-ovato hinc gibboso, foliis oblongis 3—4-dentatis; basi obtusis; biuncialibus: bracteis involucri tenuissimè pubescentibus ciliatis, capitulis erectis.

Protea elliptica. Thunb. Diss. n. 15. Prod. 26. Willd. Sp. Pl. 1, p. 512.*

Protea vestita. Lam. Illust. Gen. 1, p. 239, n. 1259?

Protea conocarpa A. Poiret, Encyc. Botan. 5, p. 657?

HAB. In Africæ Australis montibus. (v. s.)

Obs. Calli apicis foliorum obtusiusculi.

*6. *L. nutans*, stylo calycem supra sericeum bis superante, stigmatè obliquo turbinato! involucri bracteis tomentosis incanis, capitulis nutantibus, foliis ovatis oblongisve 3—5-dentatis; basi obtusis.

a. Foliis subovatis cordatis vix sesquiuncialibus.

β. Foliis lineari-oblongis basi simplicibus, 2—3 uncias longis.

HAB. In Africæ Australis montibus. *Masson.* (v. s. a. in Herb. Banks., β. in Herb. Lambert.)

Obs. Distincta stigmatē obliquo, apice depresso, axi longitudinali elevata.

Variat ramis tomentosis et hirsutis.

7. L. *Conocarpum*, stylo calycem villosissimum superante, stigmatē subæquilaterali oblongo-conico, foliis ovalibus 3—9-dentatis, ramis bracteisque hirsutissimis.

Scolymocephalus africanus latifolius lanuginosus foliis in summitate crenatis. *Herm. Cat.* 20.

Leucadendron similis *Africana* arbor argentea folio summo crenaturis florida. *Plukn. Phyt. t.* 200, *f.* 2, folium, sed nux vix hujus generis.

Leucadendron, *africana* arbor argentea summo folio crenato. *Plukn. Alm.* 212.

Conophoros capitis Bonæ Spei, folio in summo dentato. *Raj. Hist. 3. App.* 240. *Petiv. Mus.* 172, fide spec. in Herb. *Petiv.*

Conocarpodendron; folio crasso, nervoso, lanuginoso, suprâ crenato, ibique limbo rubro; flore aureo; cono facilè deciduo. *Boerh. Lugd. Bat. 2, p.* 196, *c. tab.* bona.

Scolymocephalus africanus folio crasso nervoso. *Weinm. Phyt.* 4, *p.* 292, *t.* 899, *f.* 6.

Protea foliis oblongo-ovatis apice quinquedentato-callosis. *Roy. Lugd. Bat.* 184.

Leucadendron foliis ovatis obversis oblongis, margine calloso fimbriatis ad apicem crenatis. *Wachend. Ultraj.* 203.

Leucadendron Conocarpodendron. *Linn. Sp. Pl. ed. i, p.* 93; *ed. ii, p.* 136. *Syst. Nat. xii, t. 2, p.* 110. *Berg. Act. Stockh.* 1766, *p.* 321. Omiss. in *Linn. Mant. et Syst. Veg. xiii.*

Protea conocarpa. *Thunb. Diss. n.* 14,* desc. partim a *L. grandiflora* desumptâ. *Thunb. Prod.* 25. *Willd. Sp. Pl. 1, p.* 512. *Lam. Illustr. Gen. 1, p.* 239, *n.* 1260, *tab.* 53, *f.* 3, mala, præcipuè floribus separatis. *Poiret, Encyc. Botan. 5, p.* 656.

HAB. In Africæ Australis campis et collibus sterilibus, prope Promont. B. Spei. (v. v. ad littora Simon's Bay.)

100] 8. *L. grandiflorum*, stylo calycem villosissimum superante, stigmatæ æquilaterali oblongo-cylindraco, foliis oblongo-lanceolatis tridentatis integrisque, ramis hirsutissimis, bracteis involucri glabris ciliatis.

Leucadendron grandiflorum. *Salisb. Parad.* 116.

HAB. In Africæ Australis montosis. (v. s. in Herb. Banks. sub nomine *Protea villosiusculæ*.)

9. *L. puberum*, stylo calycem hirsutum superante, stigmatæ æquilaterali ovato, foliis lanceolatis ellipticisve integris uncia brevioribus pubescentibus, ramis hirsutis, bracteis involucri incano-villosis ellipticis longè acuminatis.

Protea pubera. *Linn. Mant.* 192,* fide spec. in illius Herb., exclusis synonymis. *Thunb. Diss.* n. 56. *Prod.* 27. *Lam. Illust. Gen.* 1, p. 234, n. 1216. *Willd. Sp. Pl.* 1, p. 533, excl. syn. Bergii. *Poiret, Encyc. Botan.* 5, p. 642.

HAB. In Africæ Australis summis montibus; Hottentot's Holland. (v. s. in Herb. Banks., Lambert.)

OBS. Variat foliis angusto-lanceolatis.

*10. *L. buxifolium*, stylo calycem hirsutum superante, stigmatæ æquilaterali ovato, foliis ovalibus obtusis integris unguicularibus pubescentibus, ramis hirsutis, bracteis involucri orbiculato-ovatis breviter acuminatis glabriusculis ciliatis.

HAB. In Africæ Australis montibus. *Masson*. (v. s. in Herb. Banks.)

OBS. Proximum priori et fortè cum eo a Thunbergio confusum.

*11. *L. patulum*, stylo calycem tomentoso-villosum superante, ante, stigmatæ æquilaterali ovato, foliis spathulato-linearibus integris: adultis glabris, ramis divaricatis tomentosis, capitulis pedunculatis.

HAB. In Africâ Australi. *Masson*. (v. s. in Herb. Banks.)

DESC. *Frutex* humilis, ramosissimus. *Folia* conferta, uncia breviora, basi angustata, callo apicis acutiusculo, summa tomentosa. *Capitula* magnitudine avellanæ; pedunculo tomentoso, bracteis lanceolatis; *Bractea* involucrantes ovatæ, acuminatæ, tomentosæ, incanæ. *Calyx* tubulosus, bilabiatus, tomentosus, villisque brevibus patulis suprà frequentioribus. *Stylus* 9 lineas longus. *Stigma* breve.

OBS. Valdè affinis *L. pubero*.

*12. *L. spathulatum*, stylo calycem villosotomentosum superante, stigmatæ æquilaterali, foliis spathulatis basi lineari: adultis glabris uncialibus, ramis hirsutis patulis, capitulis pedunculatis, bracteis tomentosis acuminatis.

HAB. In Africâ Australi. *D. Niven*. (v. s. in Herb. Hibbert.)

DESC. *Frutex* humilis, ramosissimus. *Rami* villis brevibus patulis tomentoque cinereo instructi. *Folia* elliptico-spathulata, basi attenuata, lineari, tortâ: callo apicis obtuso; obsoletè venosa. *Capitulum* magnitudine juglandis minoris; *Bracteis* involucrantes ovatis, acuminatis. *Calyces* villis brevibus patulis densè tecti, laminarum decumbentibus, brevissimis. *Stylus* uncialis.

13. *L. tomentosum*, stylo sublongitudine calycis, caule erecto, foliis linearibus cuneatisve tridentatis tomentosis, bracteis lanceolatis tubum calycis subæquantibus.

Protea tomentosa. *Thunb. Diss. n. 18.* Prod. 26. Linn. Suppl. 118. Lam. Illustr. Gen. 1, p. 239, n. 1257. Willd. Sp. Pl. 1, p. 514. Poiret, Encyc. Botan. 5, p. 656.*

a. foliis linearibus canaliculatis aveniis, ramis bracteisque tomentosis, calycis laminis barbatis.

β. foliis lineari-cuneatis planis subvenosis 3—5-dentatis, ramis hirsutis, bracteis calycisque laminis tomentosis.

Protea candicans. And. Repos. 294.

γ. foliis linearibus planis, ramis hirsutis, bracteis glabris ciliatis.

HAB. In Africæ Australis montibus, prope Promont.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 272-4503

25
28
32
22
20

10
01

B. Spei. (v. s. in Herb. Banks., Lambert., et Soc. Linn.)

OBS. Plantæ pro varietatibus suprâ habitæ fortè species distinctæ.

14. L. *Hypophyllum*, stylo longitudine calycis, caule procumbente, foliis linearibus tridentatis, bracteis orbiculato-ovatis tomentosis tubo calycis dimidio brevioribus.

Thymelæa capitata Rapunculoides Nerii crassioribus foliis summo apice tridentatis æthiopica coniformi calyce squamato. *Plukn. Mant.* 181, t. 440, f. 3.

Conophoros capensis folio angusto summo dentato *Petiv. Mus.* 900, fide spec. in illius Herbar.

Scolymocephalos foliis angustis in summitate tridentatis. *Raj. Hist.* 3, *Dendr.* p. 9.

Conocarpodendron; folio rigido, angusto, apice tridentato rubro; flore aureo. *Boerh. Lugd. Bat.* 2, p. 198.* c. tab.

Scolymocephalus seu Conocarpodendron folio angusto. *Weinm. Phyt.* 4, p. 294, t. 902, f. a.

Protea foliis lanceolato-linearibus apice tridentato-callosis. *Linn. Hort. Cliff.* 29. Herb. Cliff. absque fructificatione.

¹⁰⁸⁷ Protea foliis lanceolatis linearibus apice tridentato callosis capitulis aphyllis. *Roy. Lugd. Bat.* 184. *Wachend. Ultraj.* 202.

Leucadendron Hypophyllocarpodendron. *Linn. Sp. Pl.* ed. i, p. 93; ed. ii, p. 136. *Berg. Act. Stockh.* 1766, p. 321.* *Berg. Cap.* 16.*

Protea Hypophyllocarpodendron. *Linn. Mant.* 191.* desc. opt.

Protea Hypophylla. *Thunb. Diss. n.* 16.* *Prod.* 26. *Lam. Illust. Gen.* 1, p. 239, n. 1256. *Willd. Sp. Pl.* 1, p. 513. *Poiret, Encyc. Botan.* 5, p. 655.

HAB. In Africæ Australis sabulosis depressis prope Prom. B. Spei. (v. v. in collibus juxta Simon's Bay.)

OBS. Variat foliis glabris, pubescentibus et incanotomentosis, 3 — 5-dentatis passimque integris, planis canaliculatisve, ramis nudiusculis, villosis v. tomentosis;

Capitulis subsessilibus pedunculatisque ; Bracteis latè ovatis, acutis orbiculatisve.

†† *Receptaculum planiusculum ; Bracteis propriis angustis deciduis.*

*15. *L. molle*, foliis ellipticis acutis 2—3-dentatis integrisve subsericeo-pubescentibus mollibus, bracteis exterioribus glabriusculis, stigmatè ovato.

HAB. In Africæ Australis montibus. (v. s.)

OBS. Proximum *L. crinito*, diversum figura foliorum et fortè caule procumbenti.

16. *L. crinitum*, foliis obovato-oblongis obtusis 3—5-dentatis integrisve ; basi angustatis ; pubescentibus demum glabris scabriusculis, bracteis omnibus villosis.

Protea crinita. *Thunb. Diss. n. 13?** *Prod. 25. Willd. Sp. Pl. 1, p. 511?* *Poiret, Encyc. Botan. 5, p. 657.*

HAB. In Africâ Australi. (v. s. in Herb. Soc. Linn.)

17. *L. oleæfolium*, foliis ovali-oblongis sublanceo-¹⁰⁴ latisve tridentatis et integris : adultis glabris, bracteis omnibus villosis, stigmatè oblongo.

Leucadendron oleæfolium. *Berg. Act. Stockh. 1766, p. 320.* Berg. Cap. 15.**

Protea criniflora. *Linn. Suppl. 117.**

HAB. In Africâ Australi. (v. s. in Herb. Banks.)

OBS. Duplex varietas, altera foliis ovali-oblongis obtusis ; bracteis exterioribus glabriusculis apice barbatis : altera foliis lineari-oblongis acutiusculis bracteis omnibus villosis. Ambæ à *L. crinito* diversæ foliis basi haud angustatâ.

18. *L. diffusum*, foliis cuneato-linearibus integris 2—3-dentatisve basi angustatis : adultis glabris, ramis procumbentibus, bracteis tomentosis lanceolatis acuminatis calyce dimidio brevioribus.

Protea heterophylla. *Thunb. Diss. n. 19?** *Prod. 26? Willd. Sp. Pl. 1, p. 515.*

HAB. In Africâ Australi. *Gul. Roxburgh, M.D.* (v. s. in Herb. Banks. et Soc. Linn.)

DESC. *Frutex* prostratus? *Rami* longi, glabri v. hirsuti, quandoque adscendentes. *Folia* uncialia, plana v. marginibus leviter inflexis concaviuscula, obsolete venosa, in ramis prostratis secunda. *Capitula* solitaria, breviter pedunculata, turbinata, magnitudine avellanæ; *Bracteæ* involucri incanæ, calyce hirsuto dimidio breviores. *Pistillum* calyce sesquialongius. *Stigma* clavatum *stylo* capillari parum crassius.

OBS. Species affinis *L. patulo*.

105]

7. MIMETES.

Salisb. Parad. HYPOPHYLLOCARPODENDRON. Boerh. Lugd. Proteæ Sp. 9—10. Linn. Mant.

CHAR. GEN. *Calyx* quadripartitus, æqualis, laciniis distinctis. *Stylus* filiformis, deciduus. *Stigma* cylindraceum, gracile. *Nux* ventricosa, sessilis, lævis. *Receptaculum* commune planum, paleis angustis, deciduis. *Involucrum* indefinitè polyphyllum, imbricatum.

HABITUS. Frutices. *Folia integra v. calloso-dentata. Capitula axillaria, in quibusdam folio superiori cucullato amplexa! quandoque terminalia. Involucra membranacea, rarè coriacea, nunc dimidiata! Pistilla calyce post expansionem flaccido longiora. Stigma sæpissimè acutum.*

† *Capitula axillaria.*

1. *M. hirta*, involucris æquilateralibus coloratis acuminatis semi-exsertis 8—10-floris, stigmatibus subulato, laminis calycis plumosis, foliis acutis integerrimis.

Scolymocephalus Africanus argenteus foliis Dorycnii Plateau. Herman. Cat. Mt.

Conophoros capensis foliis pilosis apice nigricante. Petiv. Mus. 62, fid. spec. in illius Herb.

Lepidocarpodendron; foliis sericeis, brevibus, confertissimè natis; fructu gracili, longo. *Boerh. Lugd. Bat. 2, p. 194, c. tab.*

Scolymocephalus africanus argenteus foliis Dorycnii.
Weinm. Phyt. 4, p. 292, t. 899, bona.

Leucadendron hirtum. *Amœn. Acad.* 6, p. 83.* *Sp. Pl. ed. ii, p. 136.*

Protea hirta. *Linn. Mant. p. 188.** (Herb. Linn.) ¹¹⁰⁶
*Thurb. Diss. n. 55,** exclus. syn. Boerh. *Lugd.* 2, p. 205.
Thunb. Prod. 27. *Lam. Illustr. Gen.* 1, p. 234, n. 1213.
Willd. Sp. Pl. 1, p. 532. *Poiret, Encyc. Botan.* 5, p. 641.

HAB. In Africæ Australis campis collibusque, in locis humidis. (v. v. in collibus humidis prope Simon's Bay.)

*2. *M. capitulata*, involucris æquilateralibus coloratis acutis semi-exsertis pubescentibus 8—10-floris, stigmatè apice conico-incrassato! laminis calycis plumosis, foliis acutis integerrimis.

HAB. In Africâ Australi. *Gul. Roxburgh, M.D.* (v. s. in Herb. Banks.)

DESC. *Frutex* erectus. *Rami* pubescentes. *Folia* elliptico-lanceolata, vix uncialia, pubescentia, haud sericea, ciliata, floralia parùm latiora; *Involucra* foliis paulò longiora; *Bracteis* ellipticis, acutis, rubris, tenuissimè pubescentibus. *Calyces* involucri vix longiores. *Styli* calycibus ferè duplo longiores apice parùm incrassato tetragono subfusiformi. *Stigma* stylo nodulo articuliformi connexum, cylindraceum, sulcatum, apice duplo crassiore conico-capitato.

*3. *M. pauciflora*, involucris subæquilateralibus coloratis acutis villosiusculis subquadrifloris, calycibus pistilla æquantibus! laminis nudiusculis, stigmatè cylindraceo, foliis obtusis integerrimis sericeis.

HAB. In Africâ Australi. *Gul. Roxburgh, M.D.* (v. s. in Herb. Lambert.)

DESC. *Frutex* erectus. *Rami* villosi, vestiti. *Folia* imbricata, frequentia, ovalia, plèna, uncialia, venis altè immersis adversus lucem tantummodò obviis. *Involucra* ¹¹⁰⁷ cylindracea, foliis sesquilogiora. *Bracteis* membranaceis, rubris, extimis obtusis ter brevioribus. *Calyx* involucri ferè unam quartam longior; *Unguibus* hirsutis; *Laminis*

glabriusculis, pube brevi adpressâ. *Stylus* calycem vix superans, extra medium angulatus. *Stigma* cylindraceum, sub-emarginatum, crassitie styli, quo cum nodulo connexum.

4. *M. cucullata*, involucris inæquilateralibus subdimidiatis acuminatis glabriusculis, foliis lineari-oblongis tridentatis glabris: floralibus infra dilatatis marginibus recurvis, stigmatе subulato acutissimo.

Scolymocephalus africana, foliis angustis brevioribus, tribus in summitate denticulis, capitulis foliosis interceptis. *Herm. Afr.* 20.

Leucadendros africana s. *Scolymocephalus angustiori* folio apicibus tridentatis. *Plukn. Alm.* 212, t. 304, f. 6, bona.

Hypophyllocarpodendron foliis inferioribus apice trifido rubro superioribus penitus rubris glabris. *Boerh. Lugd. Bat.* 2, p. 206, c. tab.

Scolymocephalus seu *Hypophyllocarpodendron* foliis tribus in summitate. *Weinm. Phyt.* 4, p. 297, t. 905.

Protea foliis lanceolatis obtusis foliis involventibus apice tridentato-callosis. *Roy. Lugd. Bat.* 184.

Leucadendron foliis cuneiformibus apice tridentato-callosis summis ultra florem protensis. *Wachend. Ultraj.* 203.

Leucadendron cucullatum. *Linn. Sp. Pl. ed.* i, p. 93; ed. ii. p. 136. *Berg. Act. Stockh.* 1766, p. 320.* *Berg. Cap.* 14.*

Protea cucullata. *Linn. Mant.* 189.* *Thunb. Diss. n.* 17.* *Prod.* 26. *Lam. Illust. Gen.* 1, p. 239, n. 1258. *Willd. Sp. Pl.* 1, p. 514. *Poiret, Encyc. Botan.* 5, p. 656.*

^{108]} *HAB.* In Africæ Australis uliginosis prope Prom. B. Spei. (v. v. juxta Simon's Bay et Constantiam.)

OBS. *Frutex* 2—3 pedes altus. *Folia* vix sesquiuncialia, subavenia; floralia suprâ glabriuscula. *Stigma* infra apicem non incrassatum. Varietas foliis unciâ brevioribus sub-linearibus.

5. *M. Hartogii*, involucris inæquilateralibus subdimidiatis: bracteis acuminatis pubescentibus: interioribus

tomentosis incanis, foliis lineari-oblongis tridentatis: adultis glabris marginibus niveo-lanatis; floralium apice angustato suprâ sericeo, stigmatе extra medium fusiformi: acumine setaceo.

Hypophyllocarpodendron; foliis lanuginosis, in apice trifido rubro quasi florescens. *Boerh. Lugd. Bat. 2, p. 205, c. tab.*

Scolymocephalus seu Hypophyllocarpodendron foliis lanuginosis. *Weinm. Phyt. 4, p. 297, t. 906, a.*

Protea cucullata β . *Lam. Illust. Gen. 1, p. 239, n. 1258.*

HAB. In Africæ Australis collibus, prope Prom. B. Spei. (v. v. in montibus juxta False Bay.)

DESC. *Arbuscula* orgyalis. *Rami* patentes, tomentosi. *Folia* frequentia, imbricata, plana, biuncialia et ultra, 8 lineas lata, subvenosa, utrinque tenuissimè pubescentia, pube demùm deciduâ, lanâ marginis persistenti; floralia dimidio inferiore dilatato, oblongo, marginibus reflexis cucullato, capitulum proximè inferius amplexante; superiore breviorе, lineari, suprâ sericeo, marginibus planis. *Calyx* sesquiuncialis, plumoso-barbatus. *Stylus* calycem superans, sulcato-angulatus. *Stigma* sulcato-quadrangulum sulcis striâ parùm elevatâ. *Receptaculum* paleis subulatis, lanatis.

*6. *M. Hibbertii*, involucris inæquilateralibus subdimidiatis: bracteis obtusis: exterioribus glabris, foliis argenteis oblongo-ellipticis tridentatis integerrimisve.

HAB. In Africæ Australis alpinis humidis, prope Barbiers Kraal. *D. Niven.* (v. s. in Herb. Hibbert., Banks., Lambert.)

DESC. *Frutex* 5—6 pedes altus. *Rami* tomentosi, cinerei. *Folia* imbricata, sessilia, plana, obsoletè venosa, dum duas uncias longa, vix 8 lineas lata. *Involucra* foliis breviora, turbinato-ovata, 7—8-flora. *Bracteis* latè ovatis, exterioribus ciliatis, interioribus sericeis. *Calyx* villosissimus. *Stylus* calyce longior. *Stigma* filiforme, striatum, acutiusculum.

*7. *M. Massoni*, involucris æquilateralibus calyce dimidio brevioribus: bracteis subrotundis obtusis coriaceis folisque argenteis ovatis integris.

HAB. In Africæ Australis montibus prope Franche Hoek. *Masson*. (v. s. in Herb. Banks.)

DESC. *Frutex* erectus. *Rami* sericei. *Folia* imbricata, frequentia, plana, holosericea, obsoletè venosa, biuncialia, sesquiunciam lata, callo apicis nudiusculo. *Involucra* vix semuncialia, globosè ovata, circiter octo-flora; *Bracteis* fructiferis induratis. *Calyx* villis longis sub-adpressis incanus. *Stylus* calyce longior. *Stigma* filiforme, acutum, striatum, vix crassitie styli. *Receptaculum* villosum, angustum, cpaletatum.

†† *Capitula* terminalia. *Mimetes spuræ*.

8. *M. thymelæoides*, caule erecto, foliis ovalibus obtusis pubescentibus semunciâ brevioribus, capitulis subaggregatis, stylis infra medium pubescentibus.

Leucadendron thymelæoides. *Berg. Act. Stockh.* 1766, p. 324.* *Berg. Cap.* 19.*

¹¹⁰¹ HAB. In Africâ Australi, prope Promont. B. Spei. (v. s. in Herb. Banks.)

DESC. *Frutex* ramosissimus. *Rami* stricti, vestiti. *Folia* imbricata, vix unguicularia, subavenia, inferiora glabra. *Capitula* sessilia, globosa, magnitudine vix cerasi nigri. *Bracteæ* involucris lanceolato-ellipticæ. *Palcæ* undique densè lanatæ. *Calyx* sericeo-lanatus. *Stylus* calyce longior. *Stigma* acutiusculum.

9. *M. myrtifolia*, caule erecto, foliis lineari-oblongis obliquis integris v. 2—3-dentatis unciâ brevioribus, stylo glabro, capitulis subsolitariis.

a. foliis tomentosis, passim 2—3-dentatis, bracteis acuminatis.

β. foliis glabriusculis, summis capitulo parùm longioribus, bracteis obtusiusculis.

Protea myrtifolia. *Thunb. Diss. n.* 50*? *Prod.* 27. *Willd. Sp. Pl.* 1, p. 530. *Poiret, Encyc. Botan.* 5, p. 641.

HAB. In Africâ Australi. (v. s. in Herb. Banks. et Soc. Linn.)

DESC. a. *Frutex* parvus. *Rami* brunnei, adulti glabri, juniores villosi. *Folia* avenia, tenuissimè pubescentia v.

glabra. *Capitula* turbinata, sessilia, solitaria v. pauca aggregata, piso vix duplò majora, multiflora. *Bracteæ* involucri pubescentes, ciliatæ; exteriores ovato-lanceolatae, acumine brevi; interiores oblongo-ellipticæ, obtusiusculæ. *Calyx* tetraphyllus, plumoso-villosus. *Pistillum* calyce longius. *Stigma* crassitie styli. *Squamulæ* hypogynæ subulatæ, persistentes. *Nux* elliptica, vix compressa, tenuissimè pubescens, basi styli terminata: cortex membranaceus, tenuis, albus, separabilis apice rugoso, putamen crustaceum, nigro-fuscum. *Nucleus* integumento simplici, tenuissimo. *Chalaza* apicis lata, venis radiantibus. *Receptaculum* plumum, villosum, epaleatum.

10. *M. divaricata*, caule procumbente, foliis ovalibus obtusis pubescentibus, stylo glabro.

a. bracteis oblongo-linearibus obtusis semifoliacis, laminis calycis sericeis.

Scolymocephalos africanus argenteus, foliis brevioribus, myrtiformibus, capitulis rarioribus. *Herm. Afr.* 20.

Leucadendron divaricatum. *Berg. Act. Stockh.* 1766, p. 324.* *Berg. Cap.* p. 19.*

Protea divaricata. *Linn. Mant.* 194.* *Thunb. Diss.* n. 57.* *Prod.* 27. *Lam. Illust. Gen.* 1, p. 235, n. 1221. *Poiret, Encyc. Botan.* 5, p. 643. *Willd. Sp. Pl.* 1, p. 533.

β. bracteis lanceolatis acutiusculis subscaricis.

HAB. In Africæ Australis campis et collibus, ubique prope Promont. B. Spei. (v. v. ad latera montium, juxta Simon's Bay.)

OBS. *Calyx* tetraphyllus. *Receptaculum* epaleatum.

11. *M. purpurea*, caule procumbenti, ramis adscendentibus, foliis lineari-subulatis canaliculatis, laminis calycis glabris.

Protea foliis linearibus simplicissimis ramis determinatis floribus terminatricibus. *Roy. Ludg. Bat.* 186.

Leucadendron proteoides. *Linn. Sp. Pl. ed. i*, p. 91.* (fid. spec. tunc in Herb.) *ed. ii*, p. 134.* *Berg. Act. Stockh.* 1766, p. 326.* *Berg. Cap.* 24.*

Protea purpurea. *Linn. Mant.* 195.* *Thunb. Diss.*

n. 26.* *Prod.* 26. *Lam. Illust. Gen.* 1, p. 238, *n.* 1252. *Willd. Sp. Pl.* 1, p. 518. *Poiret, Encyc. Botan.* 5, p. 654.

HAB. In *Africæ Australis* collibus, prope Promont. B. Spei; frequens. (v. v. ad latera montium, prope Simon's Bay.)

Obs. I. Receptaculum epaleatum.

Obs. II. Variat Caule erectiusculo; Foliis undique versis ^{112]} et secundis; Bracteis acumine subulato, longo, brevissimo, vel nullo.

8. SERRURIA.

Salisb. Parad. SERRARIA. Burm. Afr. Adans. Fam.

GEN. CHAR. *Calyx* quadrifidus, subæqualis, unguibus distinctis. *Stigma* verticale, glabrum. *Squamulæ* quatuor hypogynæ. *Nux* brevissimè pedicellata, ventricosa. *Capitulum* indefinitè multiflorum; *paleis* persistentibus, imbricatis.

HABITUS. Frutices. Folia *filiformia*, *trifido-pinnatifida*, rarò *indivisa*. *Capitula terminalia* v. *c. summis alis, simplicia*, nunc *composita partialibus congestis* v. *pedunculo communi diviso corymbosa*. *Involucrum imbricatum, membranaceum, floribus sæpissimè brevius, in paucis longius, quandoque nullum.* Flores semper sessiles, purpurei. *Pistillum longitudine calycis. Stigma clavatum, rariusve cylindraceum. Nux ovalis, tenuiter pubescens, modò barbata, aliquando glabriuscula.*

Obs. Secundum Cl. *Salisburium*, "Flores interdum pedicellati," quod nunquam observare licuit.

† *Capitula simplicia; Pedunculi indivisi* v. *nulli.*

*1. *S. glaberrima*, capitulis axillaribus pedunculatis, bracteis laminisque calycis glabris, foliis indivisis passimque trifidis, caule procumbente.

HAB. In *Africæ Australis* umbrosis montium. *Masson. Kleine Hoot. Hoek. Gul. Roxburgh, M.D.* (v. s. in *Herb. Soc. Linn. et Banks.*)

Desc. *Frutex* prostratus, glaber. *Rami* filiformes, sub-

flexuosi. *Folia* alterna, remotiuscula, ramis parùm graciliora, 2—3 uncias longa. *Capitula*, erecta, sub-octoflora, pedunculo bracteato parùm breviora. *Bracteæ* propriæ subrotundæ, mucronatæ, cucullatæ, glabræ, scariosæ. *Calyx* strictus, unguibus villosiusculis.

*2. *S. cygnea*, capitulis axillaribus terminalibusque pedunculatis, bracteis glabris subciliatis, calycibus curvatis sericeis, foliis bipinnatis, caule procumbente.

α. Capitula floribus viginti pluribusve : bracteis involu-
crantibus nullis.

β. Capitula floribus viginti paucioribus : bracteis involu-
crantibus nonnullis, lanceolato-ovatis.

HAB. In Africâ Australi prope Winterhoek et alibi. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn., β in Herb. Banks.)

DESC. *Frutex* procumbens, ramosus, glabriusculus. *Folia* sesquiunciam longa, quandoque biuncialia, superiora interdum breviora. *Pedunculi* capitulo longiores, bracteis distantibus, sæpius curvati. *Capitula* globosa, magnitudine cerasi ; *Bracteis* propriis latè ovatis, acuminatis. *Calyx* unguibus sigmoideo-curvatis ; *Laminis* nutantibus. *Stylus* pariter arcuatus. *Stigma* pendulum.

*3. *S. acrocarpa*, capitulis axillaribus pedunculatis, bracteis tomentosis, calycibus curvatis sericeis, nucibus basi pubescenti styli mucronatis, foliis bipinnatifidis, caule erecto.

HAB. In Africâ Australi, Brant-fly plain. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Frutex* bipedalis et ultrà. *Ramuli* pubescentes. *Folia* biuncialia, adulta glabra. *Pedunculi* capitulo longiores, sæpè curvati, bracteis glabris distantibus, apice tenuissimè pubescentes. *Capitulum* magnitudine cerasi : *Bracteis* propriis ovato-subrotundis, breviter acuminatis, involu-
crantibus paucis similibus. *Stylus* basi incrassatâ apice arcuato. *Stigma* pendulum. *Nux barbata* pilis strictis patulis.

*4. *S. elevata*, capitulis axillaribus pedunculo brevioribus, bracteis cuneato-orbiculatis tomentosis, calycibus breviter barbatis curvatis, nucibus submuticis, foliis bipinnatis unciâ longioribus, caule erecto.

HAB. In Africæ Australis arenosis. *Masson*. Picket Berg. *Gul. Roxburgh, M.D.* (v. s. in Herb. Banks., et Soc. Linn.)

DESC. *Frutex* orgyalis. *Rami* tomentosi, cinerei. *Folia* frequentia, pilosiuscula, viridia, inferiora glabra, sesquiuncialia, callis obtusiusculis. *Pedunculi* folia sæpissimè superantes, quandoque 3-unciales, tomentosi, cinerei, bracteis alternis, lanceolatis, patentibus. *Capitulum* magnitudine cerasi, floribus viginti pluribus, semunciâ brevioribus. *Bracteæ* omnes extûs sericeo-tomentosæ. *Nux* submutica, mucronulo vix manifesto, barbata.

Obs. Descriptio e plantâ Massoni : Roxburgiana paulò diversa, *Calycibus* quandoque sericeis ; *bracteis* mucrone longiore ; *pedunculis* brevioribus, paucioribus ; *foliis* recentioribus magis hirsutis : fortè species distincta.

*5. *S. Aitoni*, capitulis axillaribus subterminalibus pedunculo brevioribus, bracteis cuneato-subrotundis mucronatis glabriusculis, calycibus plumosis, nucibus mucronatis, foliis tripartito-bipinnatis sericeis unciâ brevioribus, caule erecto.

HAB. In Africâ Australi. *D. Masson*. (v. s. in Herb. D. Aiton.)

DESC. *Rami* stricti, pedales, tomentosi, vestiti. *Folia* erecta, frequentia, 8—10 lineas longa, subargentea tomento ¹¹⁵¹ arcè adpresso, profundè trifida, laciniis bipinnatifidis, intermediâ parùm longiore magisque divisâ, lacinulis intûs sulco tenui, apiculis subrecurvis, callo obtusiusculo. *Pedunculi* e summis alis et terminalibus, corymbosi, unciales et ultrâ, tomento brevissimo cinerei, bracteis alternis e basi erectâ lanceolatâ subulatis, recurvis. *Capitula* globosa, magnitudine ferè juglandis, floribus viginti pluribus. *Bracteæ* exteriores acumine longiore, interiores latiores, omnes glabriusculæ, subciliatæ. *Calyx* 7—8 lineas longus, unguibus laminisque plumoso-barbatis. *Stigma* clavatum,

oblongum. *Nux* villis strictis sericeis barbata, basi styli mucronata. *Squamulæ* hypogynæ quatuor, subulatæ, persistentes.

*6. *S. simplicifolia*, capitulis terminalibus pedunculatis, bracteis villosis, calycibus barbatis, foliis indivisis rariùsve trifidis, caule erecto.

HAB. In Africæ Australis arenosis : Roode Zant Cascade. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Fruticulus* pedalis, sesquipedalis, simplex v. subramosus, glaber, apicem versus tenuissimè pubescens. *Folia* uncialia sesquiuncialia, canaliculata, pleraque indivisa, aliqua passim trifida, juniora hirsuta ; radicalia elongata, crassiora, canali latiore. *Pedunculi* solitarii, capitulo longiores, incano-tomentosi ; *bracteis* glabriusculis, lanceolatis, distantibus. *Capitulum* magnitudine cerasi, floribus circiter viginti. *Bracteæ* subrotundæ, breviter acuminatæ, tomentosæ, subincanæ. *Calyx* densè plumosus, niveus. *Stigma* subcylindraceum.

*7. *S. diffusa*, capitulis terminalibus pedunculatis, bracteis lanceolato-ovatis acuminatis, calycibus barbatis, foliis trifidis v. pinnatifidis subindè simplicibus uncialibus ramisque glabris, caule procumbente.

HAB. In Africæ Australis arenosis saxosis ; in elevationibus prope Roode Zant. *Gul. Roxburgh, M.D.* Prope Wilde River. *D. Niven.* (v. s. in Herb. Soc. Linn., et Hibbert.)

DESC. *Frutex* diffusus, pedalis bipedalis. *Folia* vix sesquiuncialia, dum pinnatifida laciniis quinque indivisis. *Pedunculi* solitarii, tomentosi, capitulo vix longiores ; *bracteis* angustè lanceolatis, concavis, patulis, glabris. *Capitulum* magnitudine cerasi, floribus circiter viginti. *Bracteæ* lanceolato-ovatæ, acuminatæ, villosæ, scariosæ ; extimæ angustiores, glabriusculæ. *Calyx* unguibus laminisque plumosis. *Stigma* clavato-cylindraceum.

8. *S. pinnata*, capitulis terminalibus axillaribusque pedunculatis subaggregatis, bracteis lanceolatis acuminatis villosis dimidio calyce longioribus, calycis unguibus sub-

sericeis : laminis apice barbatis, foliis pinnatifidis trifidisve unciâ longioribus, caule procumbente piloso.

Protea pinnata. *And. Repos.* 512? sed folia nimis longa.

HAB. In Africæ Australis montibus aridis ; in ascensu Paarl Berg. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* totus prostratus, basi divisus, ramis pedalis, pubescentibus. *Folia* secunda, erecta, subsesquiuncialia, sæpiùs pinnatifida, laciniis quinque, passim trifida, pilosiuscula, callis acutis. *Pedunculi* e summis alis et terminales, capitula subæquantes, adscendentes, tomentosi, bracteis alternis, ovato-lanceolatis, acuminatis, glabriusculis. *Capitula* globosa, magnitudine ferè juglandis, multiflora. *Calycis* laminæ infra sericeæ, apice penicillatim barbatae. *Stigma* erectiusculum, subclavatum, apice dilatato cavo.

117 *9. *S. arenaria*, capitulis terminalibus pedunculo longioribus, bracteis lanceolato-ovatis acuminatis villosis, calycis laminis tribus plumoso-barbatis quartâ subimberbi ; unguibus nudiusculis, foliis trifidis pinnatifidisve unciâ brevioribus, caule pubescenti.

HAB. In Africæ Australis arenosis montium. 'Tygerhoek Hill, Blue berg, &c. *Gul. Roxburgh, M.D. & D. Niven.* (v. s. in Herb. Soc. Linn. et D. Hibbert.)

DESC. *Frutex* erectus v. decumbens, pedalis, parùm ramosus. *Folia* frequentia, laciniis indivisis, sæpè secunda. *Pedunculi* solitarii, capitulo dimidio breviores. *Ungues* calycis glabri v. pilis raris patentibus.

10. *S. cyanoides*, capitulis terminalibus pedunculo longioribus, bracteis orbiculato-ovatis acuminatis villosis, calycis laminis tribus longitudinaliter plumoso-barbatis quartâ nudiusculâ, foliis patulis : superioribus subbipinnatifidis vix uncialibus ; inferioribus brevioribus trifidis, caule erectiusculo.

Cyanus æthiopicus rigidis capillaceis tenuissimis foliis trifidis ex Prom. B. Spei. *Plukn. Mant.* 61, t. 345, f. 6, fid. spec. in ejus Herb.

Protea foliis linearibus ramosis. Roy. Lugd. Bat. 186. Wachend. Ultraj. 202.

*Leucadendron cyanoides. Linn. Sp. Pl. ed. i, p. 93; ed. ii, p. 137. Berg. Act. Stockh. 1766, p. 326. Berg. Cap. 27.**

Protea cyanoides. Linn. Mant. 188. Herb. Linn.*

Protea cyanoides. Lam. Illustr. Gen. 1, p. 239, n. 1263. Poiret, Encyc. Botan. 5, p. 658, fortè diversa species.

HAB. In Africæ Australis collibus, prope Promont. B. Spei. (v. v. ad latera montium juxta Simon's Bay.)

DESC. Frutex humilis. Ramuli glabriusculi v. tenuissimè pubescentes. Folia vix uncialia, pilosiuscula, demum glabra. Pedunculi solitarii, tomentosi, capituli sæpiùs dimidio breviores, nunc subæquantes, bracteis alternis. Capitulum magnitudine cerasi majoris, folia superiora superans. Bractee scariosæ, villis adpressis, acumine subulato brevior. Stigma clavato-cylindraceum.

*11. *S. furcellata*, capitulis terminalibus pedunculatis, bracteis lanceolatis: exterioribus pedunculisque glabris; interioribus villosis, calycibus barbatis, foliis unciâ longioribus trifidis: laciniis 2—3-fidisve fastigiatis ramisque glabris, caule erecto.

HAB. In Africâ Australi. Gul. Roxburgh, M.D. (v. s.)

DESC. Rami virgati. Folia alterna, sesquiuncialia, ad medium trifida, laciniis modicè patentibus, vix calcatis, callo brevi acuto, lateralibus bifidis, intermediâ sæpiùs trifidâ. Pedunculi capitula subæquantes, bracteis linearilanceolatis, imbricatis, glabris, vestiti. Capitulum magnitudine cerasi nigri. Stigma cylindraceo-clavatum.

Obs. Valdè affinis sequenti.

*12. *S. scariosa*, capitulis terminalibus pedunculatis, bracteis lanceolatis glabriusculis calyces sericeos æquantibus apice patulis, pedunculis squarrosis, foliis bipinnatis laciniisque divaricatis ramisque glabris, caule erecto.

Protea sphærocephala. Poiret, Encyc. Botan. 5, p. 658, secund. descript. synonyma autem omnia excludenda.*

HAB. In Africâ Australi; in depressis, rariùs. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Rami* rubicundi, parùm flexuosi. *Folia* sesquiunciam longa, pinnarum lacinulis paucis, subfastigiatis, callo acuto; superiora modicè patentia. *Pedunculi* sub-
119) umbellati, capitulo paulo longiores, pilosiusculi, bracteis lanceolatis, glabris, divaricatis. *Capitulum* globosè-ovatum, magnitudine cerasi minoris; *Bracteis* omnibus scariosis, glabriusculis, carinatis, apice acuto, patulo. *Calyx* villis arcè adpressis sericeus. *Stigma* subcylindraceum.

13. *S. pedunculata*, capitulis terminalibus pedunculatis, bracteis latè ovatis tomentosis, calycibus curvatis plumoso-barbatis: laminâ interiori villis adpressis sericeâ, foliis bitripinnatifidis cauleque erecto hirsutis.

Protea pedunculata. *Lam. Illustr. Gen.* 1, p. 240, n. 1264.

Protea sphærocephala A. *Poiret, Encyc. Botan.* 5, p. 658.

Protea glomerata. *And. Repos.* 264, bona quoad faciem sed stigma nimis inclinans.

HAB. In Africæ Australis montosis; solo fertiliori; Roode Zant Cascade. *Gul. Roxburgh, M.D.* (v. s. in Herb. Banks., Lambert., Hibbert., et Soc. Linn.)

DESC. *Frutex* quandoque orgyalis. *Rami* stricti, pubescentes. *Folia* frequentia, sesquiunciam longa, pube decumbenti v. patulâ, annotino-interrupta. *Pedunculus* terminalis, dum plures aliqui axillares, capitulo sæpiùs longiores, rarè nulli, ramulis floriferis tunc foliis nanis instructis. *Capitulum* magnitudine ferè juglandis. *Calyx* densissimè barbatus, villis strictis, patulis. *Stigma* cylindræo-clavatum.

*14. *S. scoparia*, capitulis terminalibus pedunculatis, bracteis latè-ovatis villosis, calycibus barbatis, foliis triter-natis patulis unciâ brevioribus ramisque hirsutis, caule decumbente.

HAB. In Africæ Australis depressis arenosis et saxosis;

inter 24 Rivers et Fontainage Flat. *Gul. Roxburgh*, [120
M.D. (v. s. in Herb. Soc. Linn. et D. Hibbert.)

Desc. *Rami* adscendentes, demùm glabriusculi. *Folia* 8—10 lineas longa, subdivaricata, ad medium trifida, laciniis subbipinnatis, lateralibus intermediam æquantibus. *Calycis* unguis hirsuti; laminæ densiùs barbatae, interioris barbâ brevior. *Stigma* clavatum.

15. *S. hirsuta*, capitulis terminalibus pedunculo longioribus, bracteis lineari-lanceolatis hirsutis, calycibus plumoso-barbatis, foliis bipinnatis uncialibus, ramis hirsutis, caule erecto.

Protea phyllicoides. *Thunb. Diss. n. 9.* Prod. 25. Willd. Sp. Pl. 1, p. 510*, excluso synonymo Bergii.

HAB. In Africæ Australis collibus saxosis, prope Prom. B. Spei. (v. v. juxta Simon's Bay.)

Desc. *Frutex* 2—3 pedes altus. *Rami* umbellati, stricti, villis patulis, persistentibus, hirsuti. *Folia* frequentia, quandoque sesquiuncialia, modicè patentia, juniora hirsuta, adulta glabra, laciniis acutissimis. *Pedunculi* solitarii v. sæpè uno plures, capitulo dimidio breviores, bracteis lanceolato-subulatis, divaricatis. *Capitulum* magnitudine ferè juglandis, folia superiora superans. *Calyx* leviter arcuatus, barbâ laminæ interioris brevior. *Stigma* clavato-cylindraceum.

*16. *S. stilbe*, capitulis terminalibus subsessilibus ovatis, bracteis hirsutis ovatis acumine recurvo, calycibus barbatis, foliis 2—3-ternatis unciâ brevioribus, ramis pubescentibus, caule erecto.

α. folia subbitermata, semunciâ breviora, imbricata, [121
 adulta glabra; bractæ pilosiusculæ; nuces glabriusculæ.

β. folia bitermata, ferè semuncialia, subimbricata ramique hirsuta; bractæ nucesque barbatae.

γ. folia subtritermata, semunciâ longiora, patula, ramulorum floriferorum nana; bractæ nucesque hirsutæ.

HAB. In Africæ Australis montibus saxosis. *Masson*, et *Gul. Roxburgh*, *M.D.* (v. s. α. in Herb. Banks., β. et γ. in Herb. Soc. Linn.)

Obs. Plantæ hic ut varietates propositæ, fortè species distinctæ.

*17. *S. Niveni*, capitulis terminalibus sessilibus, bracteis lanceolatis: extimis glabris; interioribus sericeis, calycibus barbatis, foliis bipinnatifidis subuncialibus: summis capitulum superantibus ramisque glaberrimis, caule decumbente.

Protea decumbens. *And. Repos.* 349.

HAB. In Africæ Australis montibus saxosis. Swartberg. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Fruticulus* diffusus, spithameus, ramosissimus. *Rami* ramulique teretes, glaberrimi, rubicundi. *Folia* bitermata et bipinnatifida, intus canaliculata, mucronibus laciniarum acutissimis, semipellucidis, innocuis; modicè patentia; ramorum procumbentium secunda. *Capitula* solitaria, subsessilia, globosa, magnitudine cerasi nigri. *Bracteæ* extimæ breviter acuminatæ, extus glaberrimæ, marginibus tenuissimè ciliatis, dimidio capitulo parum longiores; reliquæ sericæ, apicibus glabriusculis. *Calyx* densè barbatus, laminâ interiori villis adpressis sericæ. *Stigma* cylindraceum, stylo vix crassius.

122] 18. *S. villosa*, capitulis terminalibus sessilibus, bracteis lanceolatis acuminatis tomentosis, calycis laminis barbatis: unguibus tomentosis, foliis subbiternatis: superioribus capitulum superantibus, ramis hirsutis, caule erecto.

Protea villosa. *Lam. Illust. Gen.* 1, p. 240, n. 1265.

Protea phyllicoides. *Poiret, Encyc. Botan.* 5, p. 659,* exclusis synonymis Bergii et Thunbergii.

HAB. In Africâ Australi, prope Promont. B. Spei; in montibus prope Simon's Bay. *Gul. Roxburgh, M.D.* In depressis prope Constantiam legi. (v. v.)

DESC. *Frutex* bipedalis et ultra. *Rami* umbellati, stricti, divisi, ultimi hirsuti. *Folia* vix uncialia, trifida; laciniis lateralibus bifidis trifidisve; intermediâ trifidâ, quandoque pinnatâ, mucronibus lacinularum acutissimis, subincurvis: modicè patentia, adulta glabra. *Capitula* solitaria, magnitudine cerasi. *Calycis unguis* tomento arcuè adpresso; *Laminæ* penicillatim barbata. *Stigma* cylindraceo-clavatum.

*19. *S. faniculacea*, capitulis terminalibus subsessilibus, bracteis glabris ovatis acuminatis, calycibus sericeis, foliis bipinnatis sesquiuncialibus: superioribus capitulum superantibus; ramisque glabris, caule erecto.

HAB. In Africæ Australis depressis, prope Constantiam, (ubi v. v.)

DESC. *Frutex* bipedalis, ramis umbellatis, rubicundis. *Folia* modicè patentia, laciniis gracili-filiformibus, acutissimis. *Capitula* solitaria, magnitudine cerasi; *pedunculo* brevissimo, bracteis imbricatis tecto, v. nullo. *Bracteæ* breviter ciliatæ. *Calyx* leviter arcuatus, unguibus laminisque argenteo-sericeis villis arcuè adpressis. *Stigma* oblongo-clavatum.

OBS. Facie, foliis, bracteis, calycibusque affinitatem quandam cum *S. glomeratâ* habet; sed capitulis semper solitariis distincta.

*20. *S. ciliata*, capitulis terminalibus pedunculo longioribus, bracteis subulatis glabris margine hirsutis dimidio capituli longioribus, calycibus sericeis, foliis subbipinnatis ramisque glabris, caule erecto.

HAB. In Africæ Australis depressis arenosis prope Physers-Hoek. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Frutex* ramosissimus. *Rami* rubicundi, ultimi tenuissimè pubescentes. *Folia* vix uncialia, modicè patentia, biternata v. subbipinnatifida; superiora capitulum vix æquantia. *Pedunculi* solitarii v. sæpè aggregati, bracteis subulatis squarrosi. *Capitula* turbinato-obovata, ceraso nigro minora. *Bracteæ* extus glabriusculæ, punctis elevatis scabriusculæ. *Calyx* arcuatus. *Stigma* cylindraceo-clavatum.

*21. *S. congesta*, capitulis terminalibus sessilibus, bracteis subulatis margine hirsutissimis dimidio capituli longioribus, calycibus barbatis, foliis subbiternatis semuncialibus, ramis pilosiusculis, caule erecto.

HAB. In Africæ Australis arenosis, inter Roode Zant et Urbem Cap. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Frutex* ramosissimus. *Rami* sparsi, adulti glabri.

Folia erecta, quandoque pinnatifida, laciniis indivisis. *Capitula* turbinata, vix magnitudine cerasi nigri, sæpiùs aggregata. *Bracteæ* extùs punctis elevatis, crebris, junioribus piliferis. *Calyx* densè barbatus, villis patulis, parallelis. *Stigma* cylindraceo-clavatum.

^{124]} *22. *S. nitida*, capitulis terminalibus pedunculo squaroso duplo longioribus, bracteis capitulo parùm brevioribus: exterioribus subulatis glabris; interioribus villosissimis sericeis, calycis laminis plumoso-barbatis: interiori unguibusque nudiusculis, foliis uncia longioribus.

Protea cyanoides. *Thunb. Diss. n. 3**? *Prod. 25*? *Willd. Sp. Pl. 1, p. 507*?

HAB. In Africæ Australis montibus. Hottentots-Holland-Kloof. *Gul. Roxburgh, M.D.* (v. s. in *Herb. Soc. Linn.*)

DESC. *Frutex* glaberrimus, ramis rubicundis. *Folia* pinnatifida et subbipinnatifida, ferè sesquiuncialia. *Capitula* solitaria, globosa, magnitudine avellanæ. *Calyx* strictus, unguibus perangustis, pilosiusculis; lamina interiori apice barbâ brevi rarâ, reliquis longitudinaliter barbatis, villis terminalibus dimidio laminæ longioribus. *Stigma* cylindraceum.

*23. *S. squarrosa*, capitulis terminalibus axillaribusque, pedunculis ramuliformibus squarrosis, bracteis dimidium capituli superantibus: exterioribus linearibus glabris; interioribus lineari-lanceolatis pilosis, calycis laminis penicillatim barbatis: interiori unguibusque nudiusculis, aliis subbiuncialibus.

HAB. In Africâ Australi. *Gul. Roxburgh, M.D.* (v. s. in *Herb. Lambert.*)

DESC. *Frutex* erectus, glaberrimus, ramosissimus, ramulis rubicundis. *Folia* bipinnatifida, patentia. *Pedunculi* capitulis parùm longiores; *bracteis* numerosis, divaricatis, inferioribus teretiusculis, foliaceis, superioribus longioribus, linearibus, confertissimis. *Bracteæ* interiores capituli pilis sparsis, patulis, hirsutæ. *Stigma* cylindraceum.

24. *S. phyllicoides*, capitulis terminalibus axillari-¹²⁵ busque, pedunculis ramuliformibus squarrosis, bracteis dimidium capituli superantibus: extimis lineari-subulatis; interioribus lanceolatis; utrisque glabris, calycis laminis penicillato-barbatis; interiori nudiusculâ; unguibus glabris, foliis sesquiuncialibus.

Leucadendron phyllicoides. *Berg. Act. Stockh.* 1766, p. 328.* *Berg. Cap.* 29,* desc. opt.

Protea sphærocephala. *Linn. Mant.* 188.* (*Herb. Linn.*) exclus. syn. *Bergii*.

Protea abrotanifolia. *And. Repos. t.* 507.

HAB. In Africâ Australi. (v. s. in *Herb. Linn. et Soc. Linn.*)

DESC. *Frutex* erectus, glaberrimus, ramulis rubicundis. *Folia* bipinnatifida, passimque pinnatifida, modicè patentia, quandoque biuncialia. *Pedunculi* (si placeas ramuli floreri) axillares et terminales, subcorymbosi, capitulis longiores, bracteis foliaceis, subulatis, indivisis, squarrosi. *Capitula* magnitudine avellanæ. *Bracteæ* extimæ punctis elevatis, interiores læves, marginibus nudis rariùsve ciliatis. *Calyx* strictus, unguibus glaberrimis, laminis exterioribus niveo-barbatis, villis terminalibus longitudine antherarum; interiori glabriusculâ. *Stigma* cylindraceum.

*25. *S. æmula*, bracteis capitulo terminali subsessili parùm brevioribus: exterioribus lanceolatis tomentosis ciliatis; interioribus minoribus villosis, calycis laminis omnibus plumoso-barbatis, foliis bipinnatifidis.

HAB. In Africæ Australis montibus prope Franche Hoek. *Gul. Roxburgh, M.D.* (v. s. in *Herb. Soc. Linn. et D. Hibbert.*)

DESC. *Frutex* 3—4 pedes altus. (*Niven.*) *Rami* ultimi tomento tenuissimo cinerascents. *Folia* sesquiuncialia, modicè patentia, glabra, laciniis acutissimis. *Pedunculi*¹²⁶ capitulo breviores, quandoque brevissimi; *bracteis* subulatis, tomentosis, divaricatis, squarrosi. *Capitula* magnitudine avellanæ majoris. *Bracteæ* membranaceæ. *Calyx* strictus, unguibus nudiusculis. *Stigma* cylindraceum.

26. *S. florida*, bracteis capitulo pedunculato longioribus : exterioribus glabris oblongo-lanceolatis acuminatis ; interioribus inclusis lineari-lanceolatis ciliatis, foliis pinnatifidis bipinnatifidisve.

Protea florida. *Thunb. Diss. n. 2,* tab. 1, bona. Lam. Illust. Gen. 1, p. 240, n. 1271. Willd. Sp. Pl. 1, p. 506. Poiret, Encyc. Botan. 5, p. 662.*

HAB. In Africæ Australis montibus prope Franche Hoek. *Masson.* (v. s. in Herb. Banks.)

†† *Capitula composita ; partialibus congestis.*

*27. *S. decumbens*, caule prostrato foliisque glabris trifidis : laciniis indivisis, capitulis partialibus subquadrifloris.

Protea decumbens. Thunb. Diss. n. 1, tab. 1. Prod. 25. Willd. Sp. Pl. 1, p. 506. Lam. Illust. Gen. 1, p. 239, n. 1261. Poiret, Encyc. Botan. 5, p. 657.*

*Protea procumbens. Linn. Suppl. 116.**

HAB. In Africæ Australis lateribus saxosis montium, prope Promont. B. Spei. (v. v. juxta Simon's Bay.)

DESC. *Frutex* prostratus, glaber, basi divisus. *Rami* elongati, rubicundi, parum flexuosi, sæpè annotino-articulati. *Folia* alterna, erecta, secunda, biuncialia, infra medium trifida, laciniis subæqualibus. *Pedunculi* terminales et sæpè e summis alis, adscendentes, graciles ; bracteis nonnullis, parvis, glabris. *Capitulum* commune sub-^{127]} conicum, magnitudine ferè juglandis, e quatuor ad sex partialibus imbricatis, breviter pedunculatis, 3—4-floris, quandoque abortione simplex. *Bractee* capitulorum partialium orbiculato-ovatae, acumine brevi, subsericeae, passimque glabriusculae. *Calyx* levissimè arcuatus, subsericeus, villis arcè adpressis. *Stigma* cylindraceum.

28. *S. ascendens*, caule procumbente foliisque glabris pinnatifidis bipinnatifidisque, pedunculis partialibus incanotomentosis, calycibus curvatis.

Protea ascendens. Lam. Illust. Gen. 1, p. 239, n. 1262. Poiret, Encyc. Botan. 5, p. 658?*

HAB. In Africæ Australis montibus. *Kleine-Hoot-Hoek. Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Frutex* glaber. *Rami* rubicundi, quandoque adscendentes. *Folia* sæpius bipinnatifida, passim pinnatifida, sesquiuncialia, biuncialia. *Pedunculi* communes terminales et interdum e summis alis, capituli dimidio breviores. *Capitulum* obtusè conicum, magnitudine ferè juglandis, compositum partialibus quinque ad septem, imbricatis, breviter pedunculatis, 6-7-floris. *Bracteæ* ovato-lanceolatae, acumine patulo, glabræ, basi tomentosâ subincanâ. *Calyx* villis adpressis, argenteis sericeus. *Stigma* subcylindraceum.

*29. *S. flagellaris*, caule procumbente foliisque pilosis bipinnatifidis, pedunculis partialibus subtomentosis, calycibus strictis.

HAB. In Africæ Australis campis arenosis lateribusque montium; prope Simon's Bay, (ubi v. v.)

DESC. *Frutex* prostratus, basi divisus. *Rami* elongati, apice adscendentes, adulti glabriusculi. *Folia* erecta, secunda, circiter biuncialia, laciniis patentibus, fastigiatis, hirsutis, pilis patulis, tardiùs deciduis. *Pedunculi* communes terminales; bracteis alternis, subulatis, vix longitudine capituli; quandoque recurvi. *Capitulum* magnitudine juglandis, e partialibus 5-8, racemoso-congestis, 8-10-floris. *Pedunculi* partiales capitulis suis breviores, tomento raricre cinerascetes. *Bracteæ* ovatae, acuminatae, pube rarâ appressâ conspersæ, ciliatae. *Calyx* sericeus, villis adpressis imbricatis. *Stigma* subcylindraceum.

30. *S. rubricaulis*, caule erecto foliisque glabriusculis subbipinnatifidis uncialibus, capitulis partialibus paucifloris, bracteis ovatis acuminatis glabris, pedunculis partialibus pilosiusculis, stigmate cylindraceo.

Protea sphærocephala. Thunb. Diss. n. 5? exclus. syn. omn.*

HAB. In Africâ Australi. *Gul. Roxburgh, M.D.* (v. s.)

DESC. *Rami* stricti, rubicundi, glabri, pilisve paucis patulis. *Folia* biternata et subbipinnatifida, erecta, vix

sesquiuncialia. *Pedunculus* communis terminalis, capitulo brevior, glaber, bracteis alternis; partiales capitulis suis dimidio breviores, pilosi, quandoque glabriusculi. *Bractee* ovatae, acumine recurvo, glabrae, ciliatae, scariosae. *Calyx* sericeus, villis adpressis.

Obs. Valdè affinis *S. adscendenti*.

31. *S. glomerata*, caule erecto foliisque glabris bipinnatifidis unciâ longioribus, capitulis partialibus multifloris, bracteis exterioribus glabris: interioribus subsericeis, pedunculo communi squarroso, stigmatè clavato.

Serraria foliis tenuissimè divisis capitulis tomentosis. *Burm. Afr. p.* 265, *t.* 99, *f.* 2, mala.

Leucadendron Serraria. *β. Linn. Sp. Pl. ed. i, p.* 94.

^{129]} Leucadendron glomeratum. *Linn. Sp. Pl. ed. ii, p.* 137, (omissum in *Syst. Nat. ed. xii.*) *Berg. Act. Stockh.* 1766, *p.* 328.

Protea glomerata. Linn. Mant. 187. Herb. Linn.*

Protea patula. Thunb. Diss. n. 4?*

HAB. In Africæ Australis collibus saxosis, prope Promont. B. Spei. (v. s. in *Herb. Linn., Banks., Soc. Linn.*)

DESC. *Frutex* ramis rubicundis. *Folia* modicè patentia, quandoque biuncialia, glaberrima. *Pedunculi communes* sæpè aggregati, bracteis patulis, latè-ovatis, acuminatis, glabris squarrosi, capitula subæquantes; *partiales* capitulis suis breviores; utrique pubescentes. *Capitula* partialia magnitudine pisi majoris, bracteis densissimè imbricatis, subrotundis, acuminatis. *Calyx* sericeus, villis adpressis.

32. *S. decipiens*, caule erecto ramulis pubescentibus, foliis bipinnatifidis uncialibus et ultra, capitulis partialibus paucifloris communique breviter pedunculatis, bracteis omnibus villosissimis, calycibus sericeis.

a. *Frutex* 4—5-pedalis, foliis sesquiuncialibus biuncialibusque.

β. *Frutex* 1—2-pedalis, foliis uncialibus, bractearum acumine glabro.

HAB. In Africæ Australis planitiis elevationibus arenosis. *Gul. Roxburgh, M.D.* (v. s. in *Herb. Soc. Linn.*)

DESC. *Frutex* ramosissimus, ramis tenuissimè pubescentibus. *Folia* patentia, superiora capitula superantia. *Capitula communia* sæpè aggregata; *partialia* 5—6-flora; *Bracteæ* ovatæ, villis longis, decumbentibus incanæ, acumine subulato, nunc glabro. *Calyx* curvatus.

33. *S. compar*, caule erecto ramis glabris, foliis bipinnatifidis unciâ longioribus, capitulis partialibus paucifloris ^[130] communique breviter pedunculatis, bracteis tomentosis, calycibus barbatis.

HAB. In Africâ Australi. (v. s.)

Obs. Nimis affinis *S. decipienti*. Differt præsertim ramis glabris, calycibus barbatis villis brevissimis patulis, bracteis exterioribus tenuissimè tomentosis, acumine recurvo.

34. *S. Roxburghii*, caule erecto, foliis triternatis fastigiatis semunciâ brevioribus, capitulo communi partialibusque sessilibus paucifloris.

HAB. In Africâ Australi, prope Pardberg in Swartland. *Gul. Roxburgh*, M.D. (v. s. in Herb. Soc. Linn.)

DESC. *Frutex* 3—4-pedalis, ramosissimus. *Rami* umbellati, spithamei, pubescentes. *Folia* adulta glabra, patula, flabelliformia, lacinulis acutissimis. *Capitulum* terminale, sæpè magnitudine juglandis minoris, quandoque vix cerasi. *Bracteæ* lanceolato-ovatæ, acuminatæ, villosissimæ, incanæ, acumine nudiusculo. *Caly.* argenteo-sericeus, villis laxiùs adpressis. *Stigma* cylindraceo-clavatuni.

†† *Pedunculi divisi. Capitulis distinctis, corymbosis v. racemosis.*

35. *S. candicans*, capitulis racemosis paucifloris, pedunculis partialibus calyce barbato brevioribus, foliis bipinnatifidis ramulisque incanis.

HAB. In Africâ Australi. (v. s.)

Obs. Facies *S. Burmanni* β, eique quam maximè affinis.

36. *S. Burmanni*, capitulis corymbosis subdecemfloris, calycibus fastigiatis sericeis apiceve nudiusculis pedunculo

partiali brevioribus, foliis bipinnatifidis setaceis vix biuncialibus.

^{131]} *a.* Ramis foliisque pilosiusculis; capitulis turbinatis, bracteis acumine glabriusculo; calycis laminis demùm nudiusculis.

Abrotanoides arboreum monamotapense floribus in ramulorum cymis. *Plukn. Mant.* 1, t. 329, f. 1, fide specim. in illius Herb.

Serraria foliis tenuissimè divisis floribus rubris apetalis. *Burm. Afr. p.* 264, t. 99, f. 1, mala, nisi quoad figuram capitulorum.

Leucadendron Serraria a. *Linn. Sp. Pl. ed. i,* p. 93; *ed. ii,* p. 137.

Protea Serraria. *Linn. Mant.* 188.* *Herb. Linn. Thunb. Diss. n.* 6.* *Prod.* 25. *Willd. Sp. Pl.* 1, p. 508. *Lam. Illust. Gen.* 1, p. 240, n. 1268. *Poiret, Encyc. Botan.* 5, p. 660.

β. Ramis foliisque subsericeis; capitulis basi obtusis, bracteis totis calycibusque sericeis.

HAB. In Africæ Australis depressis sterilibus, et ad latera montium. *a.* ubique. *β.* rariùs; fortè distincta species: (*a.* v. v. juxta Simon's Bay. *β.* v. s. in Herb. Soc. Linn. et D. Hibbert.)

37. *S. triternata*, corymbis compositis, capitulis globosis; floribus viginti pluribus imbricatis, bracteis pedunculisque partialibus sericeis, foliis triternatis digitalibus cauleque glaberrimis.

Protea triternata. *Thunb. Diss. n.* 7.* *Prod.* 25. *Willd. Sp. Pl.* 1, p. 509. *Poiret, Encyc. Botan.* 5, p. 660.

Protea argentiflora. *And. Repos.* 447, bona.

HAB. In Africâ Australi, prope fluvium ad Roode Zant. *D. Niven.* (v. s. in Herb. Banks., Hibbert., et Soc. Linn.)

DESC. *Frutex* erectus, orgyalis. *Rami* rubicundi crassitie pennæ anserinæ. *Folia* patentia. *Corymbus* ^{132]} paniculatus, foliis sæpiùs longior, ramis glabris, ramulis tomentosus, incanis, subangulatis. *Bracteæ* ad divisuras glabriusculæ, acutæ, patentæ. *Capitula* magnitudine

cerasi nigri. *Bractea* ovata, acuminata. *Calyx* argenteo-sericeus, villis laxiùs decumbentibus. *Stigma* ovale.

38. *S. elongata*, corymbis simplicibus subcompositisve, pedunculo communi elongato: partialibus bracteisque glabris: acumine subulato recurvo dimidium baseos ovatae superante, foliis 2—3-pinnatifidis digitalibus.

Leucadendron elongatum. *Berg. Act. Stockh.* 1766, p. 327.* *Berg. Cap.* 27.*

Protea glomerata. *Thunb. Diss. n.* 8,* exclus. synon. Linnæi et fortè Burmanni. *Thunb. Prod.* 25. *Willd. Sp. Pl.* 1, p. 509, sec. descrip. a Thunb. mutato.

Protea thyrsoidea. *Lam. Illustr. Gen.* 1, p. 240, n. 1267. *Poiret, Encyc. Botan.* 5, p. 660.*

HAB. In Africæ Australis montibus. Hottentots-Holland-Kloof. Kleine-hoot-Hoek. *Gul. Roxburgh, M.D.* (v. s. in Herb. Banks. et Soc. Linn.)

DESC. *Frutex* erectus, subramosus, sesquipedalis, glaber. *Folia* (*Crithmi*) ad apicem rami articulive annotini conferta, infra nulla. *Pedunculus* communis 3—10-uncialis, infra bracteis distantibus, apice corymbosus. *Capitula* globosa, 16—20-flora, superiora præcociora. *Bractea* scariosa, latè ovata. *Calyx* sericeus. *Stigma* clavato-oblongum.

39. *S. crithmifolia*, racemis simplicibus, pedunculo communi elongato partialibusque glabris, capitulis subtrigintifloris, bracteis glaberrimis latioribus quàm longis: mucrone brevissimo obtuso erecto, foliis bi-tripinnatifidis digitalibus.

HAB. In Africâ Australi. *D. Niven.* (v. s. in Herb. D. Hibbert.)

DESC. *Frutex* erectus, simplex? *Folia* 3—4-uncialia, laciniis teretiusculis, callo apicis obtuso. *Pedunculus* terminalis, scapiformis, sæpè infra racemum 8—10 uncias æquans, bracteis paucissimis. *Racemus* scapo plerumque brevior, 8—10-florus. *Pedunculi* partiales, capitulo longiores, basi dilatata, cum processu scutelliformi racheos articulati. *Capitula* magnitudine avellanæ, globosa. *Calyx*

semuncialis. *Nux* undique pubescens, pedicello brevissimo, glabro, rugoso.

9. NIVENIA.

Paranomus. *Salisb. Parad.*

CHAR. GEN. *Calyx* quadrifidus, æqualis, totus deciduus. *Stigma* clavatum, verticale. *Nux* ventricosa, nitens, sessilis, basi integrâ. *Involucrum* simplici serie tetraphyllum, quadriflorum, fructiferum induratum; *Receptaculo* plano epaleato.

HABITUS. Frutices. *Folia sparsa, inferiora bipinnatifida filiformia; superiora, in quibusdam, indivisa, plana. Involucra in spicam variûsve capitulum terminale digesta, sessilia, bractea unicâ subtensa. Flores purpurascens.*

This genus is published by Mr. Salisbury: his primary generic character does not indeed at all differ from that which he has given to *Mimetes*; in his account of Inflorescence, however, it is evident he understood the genus nearly as I have here proposed it: I should therefore have adopted his name had it appeared to me tenable; but I am disposed to believe that he will, on reconsidering the subject, see the propriety of relinquishing it; for the irregularity or unusual structure, which (if I understand him) he says exists "tot partibus diversis," only takes ¹³⁴ place in the leaves of a small number of species; on the other hand, the flowers of all are perfectly regular, and that too in opposition to some of the most nearly related genera, while the great uniformity and regularity of inflorescence forms an essential part of its character. I have therefore named it in honour of Mr. James Niven, an intelligent observer and indefatigable collector, to whom botanists are indebted for the discovery of many new species, especially in the two extensive South-African families of *Erica* and *Proteaceæ*.

† *Folia superiora indivisa, latiora.*

1. *N. Sceptrum*, foliis obovatis lanceolatisve planiusculis margine simplicibus, calyce sericeo villis adpressis.

Protea Sceptrum Gustavianum. *Sparm. in Act. Stockh.* 1777, p. 55, t. 1, bona. *Linn. Suppl.* 116. (*Herb. Linn.*)

Protea Sceptrum. *Thunb. Diss. n.* 12.* *Prod.* 25. *Willd. Sp. Pl.* 1, p. 511. *Poiret, Encyc. Botan.* 5, p. 662.

Protea alopecuroides. *Lam. Illustr. Gen.* 1, p. 240, n. 1272.

HAB. In Africæ Australis summis montibus Hottentots-Holland. (v. s. in *Herb. Banks.*)

Obs. Involucri fructiferi foliola aucta, indurata.

*2. *N. marginata*, foliis latioribus quàm longis cucullatis marginatis, calyce sericeo villis adpressis, involucri foliolis acutis apice glabriusculis.

HAB. In Africæ Australis montibus. *Gul. Roxburgh, M.D.* (v. s. in *Herb. Soc. Linn.*)

DESC. *Frutex*. Rami umbellati, stricti, glabri, rubicundi. *Folia* subrotunda, parùm latiora quàm longa, diametro 8—10-lineari, glauca, margine cartilagineo, latiusculo, semi-pellucido (infima nondum visa). *Spica* subsessilis, sesquiuncialis. *Bracteæ* subulatæ, concavæ, glabriusculæ. *Stylus* glaber. *Stigma* clavâ oblongâ.

3. *N. spathulata*, foliis latioribus quàm longis cucullatis marginatis, involucri foliolis obtusis, calyce barbato, stylo glabro, stigmatibus clavato-oblongo.

Protea spathulata. *Thunb. Diss. n.* 58,* t. 5. *Prod.* 28. *Lam. Illustr. Gen.* 1, p. 235, n. 1218. *Willd. Sp. Pl.* 1, p. 533. *Poiret, Encyc. Botan.* 5, p. 642.

HAB. In Africæ Australis montibus, Platte-Kloof. *D. Masson.* (v. s. in *Herb. Banks.*)

Obs. *Folia* infima 2—3-pinnatifida, filiformia, canaliculata.

*4. *N. parvifolia*, foliis latioribus quàm longis cucullatis, calycibus barbatis, stylo lanato, stigmate conico-capitato.

Protea Sceptum. *Lam. Illust. Gen.* 1, p. 241, n. 1273?

Protea Gustaviana. *Poiret, Encyc. Botan.* 5, p. 663? exclus. syn. *Sparm.* et *Linnei*.

Protea spathulata. *Thunb. Diss. tab.* 5, quoad figuram.

HAB. In Africæ Australis montibus. *D. Masson.* (v. s. in *Herb. Banks.*, *Soc. Linn.*, *Hibbert.*)

DESC. *Frutex* ramosissimus. *Rami* umbellati, patentes; ramuli tenuissimè pubescentes. *Folia inferiora* bipinnatifida, filiformia, canaliculata; *religua* orbiculato-rhombea, frequentia, glaberrima, diametro vix unguiculari, margine cartilagineo, angusto, crenulato. *Petiolì* adpressi, foliis breviores. *Spicæ* terminales, solitariae, v. aggregatae, sesquiunciales—biunciales, dum solitariae sessiles, dum aggregatae sæpè pedunculatae. *Involucrum* foliolis subrotundis, fructiferis auctis, induratis. *Stylus* angulatus, dimidio inferiore longiore, lanato. *Stigma* magnum, apice styli duplo crassius, rugosiusculum.

136]

†† *Folia omnia bipinnatifida.*

5. *N. spicata*, pedunculis subumbellatis dimidio spicæ cylindraceæ longioribus, bracteis subtendentibus pedunculi ovatis, involucris inferioribus distinctis, stylis ad duas tertias villosissimis, foliis glabris, ramis tomentosis.

Leucadendron spicatum. *Berg. Act. Stockh.* 1766, p. 327.* *Berg. Cap.* 25.*

Protea spicata. *Linn. Mant.* 187.* (*Herb. Linn.*) *Thunb. Diss. n.* 11.* *Prod.* 25. *Willd. Sp. Pl.* 1, p. 511.

HAB. In Africæ Australis montibus. *Hottentots-Holland-Kloof.* (v. s. in *Herb. Banks.*)

DESC. *Frutex* erectus, ramis tenuissimè tomentosis, villis præterea nullis. *Folia* subtriterinata, biuncialia, canaliculata, callis obtusis. *Pedunculi* terminales, quandoque solitarii, sæpius 3—5 umbellati, tomento villisque brevibus patulis incani; bracteis alternis, numerosis, adpressis; ses-

quiunciales—biunciales. *Spicæ* sesquiunciales, usque $2\frac{1}{2}$ uncias æquantes. *Involucra* superiora conferta, inferiora distincta; bracteis subtendentibus ovatis, acumine brevissimo; *foliolis* ovatis, acutis, fractiferis auctis, induratis. *Calyx* basi villosus, unguis tomentosi, laminis breviter barbatis. *Stylus* ipsâ basi et tertiâ parte superiore glabris. *Stigma* clavato-ovale. *Nux* ovata, cortice albo nitente tenuissimo; denudata fusca, basi parum incrassatâ, stylo diu terminata.

6. *N. crithmifolia*, pedunculis umbellatis spicas conico-cylindræas subæquantibus, bracteis subtendentibus ovatis acuminatis, involucris alternis: foliolis obtusis, stylis ad medium villosis, foliis divaricatis glabris.

Protea Lagopus. *And. Repos.* 243.

HAB. In Africæ Australis montibus. *D. Niven.* [137 (v. s. in Herb. Hibbert.)]

Obs. Nimis affinis *P. spicato*, et fortè haud distincta species: differt tamen foliis divaricatis, lacinulis latioribus, sursum paulò dilatatis; bracteis pedunculi paucioribus parùmque angustioribus; spicis pedunculo vix longioribus; involucris magis distinctis, foliolis obtusioribus tomento arcè adpresso; styli dimidio superiore glabro.

7. *N. media*, spicis cylindræis pedunculo quater longioribus, bracteis subtendentibus capitulorum lanceolato-subulatis, involucris inferioribus subdistinctis: foliolis ovatis acutis apice imberbibus, stylo infra medium pubescenti, foliis glabris, ramis tomentosis.

Protea spicata. *And. Repos.* 234?

HAB. In Africæ Australis montibus, frequens. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* 6—8 pedes altus (*Niven*). *Rami* umbellati, stricti, tomento tenuissimo cinerascetes. *Folia* erecta, sesquiuncialia; inferiora biternata et subtrternata; superiora trifida, laciniiis lateralibus subsimplicibus. *Pedunculi* terminales, solitarii, vix unciales, villosi, bracteis lanceolatis, sparsis, erectis, tomentosis. *Spicæ* 3—5 uncias longæ, involucris distinctis, tamen approximatis, foliolis

acutissimis, tomento arcè adpresso. *Calyx* tubo tomentoso, involucri ferè ter longiore; laminis villis brevibus, sericeis, subdecumbentibus, barbatis. *Stylus* vix ultra unam tertiam a basi pubescens. *Stigma* gracile, clavatum.

8. *N. Lagopus*, spicis subsessilibus cylindraceis, capitulis imbricatis: bracteis subtendentibus lanceolato-subulatis: ^{138]} involucri subrotundis apice acuto barbato, stylo infra medium pubescente, foliis adultis glabris: junioribus ramulisque pilosis.

Protea Lagopus. *Thunb. Diss. n. 10.* Willd. Sp. Pl. 1, p. 510.*

HAB. In Africæ Australis montibus. *Gul. Roxburgh, M.D. (v. s.)*

DESC. Frutex erectus. *Rami* umbellati. *Folia* vix sesquiuncialia, modicè patentia, biternata. *Spicæ* solitariae, densæ, 2—4 unciales, pedunculo quandoque semunciali, sæpè brevissimo v. nullo. *Bracteæ* subtendentes apice barbatae. *Calyx* unguibus tomentosus, laminis barbatis, villis longis, numerosis, patulis. *Stylus* vix ad medium pubescens. *Stigma* ovali-clavatum.

*9. *N. mollissima*, spicis pedunculosis vix æquantibus, foliis sericeis triternatis (uncialibus), calycis unguibus tomentosus: laminis barbatis.

HAB. In Africæ Australis montibus. *D. Joh. Roxburgh. (v. s. in Herb. Banks., Lambert., Soc. Linn.)*

DESC. Frutex erectus, tomentosus, incanus. *Rami* ramulique tomento arcè adpresso. *Folia* mollissima, profundè trifida, lacinulis fastigiatis. *Pedunculi* terminales, subsolitarii, foliis breviores. *Spicæ* subovatae, capitulis inferioribus distinctis, bracteis ovatis acutis, involucri similibus, utrisque tomentosus, imberbibus. *Calycis* unguis involucri ferè ter longiores. *Stylus* infra medium pubescens. *Stigma* gracile. *Nux* ovata, cuticulâ albâ nitente tenuissimè pubescente, basi incrassatâ styli diu coronata; involucri foliolis coriaceo-induratis, parùmque auctis, demùm patulis cincta.

*10. *N. capitata*, capitulo communi globoso subsessili, unguibus laminisque calycis barbatis, foliis semuncialibus : ramulorum inferioribus glabris.

HAB. In Africæ Australis montosis, near Brant-fly's Hill. *Gul. Roxburgh, M.D.* (v. s. in Herb. Banks., Lambert., Soc. Linn.)

DESC. *Frutex* erectus, tripedalis et ultrà. *Rami* umbellati, ultimi tomentosi. *Folia* biternata, canaliculata, superiora ramulorum sericea. *Capitula* communia vix magnitudine cerasi nigri, pauciflora, quandoque aggregata breviterque pedunculata. *Involucrum* foliolis lanceolato-ovatis, acutis. *Stylus* medio pubescens, utroque fine glaber. *Stigma* ovali-clavatum.

10. SOROCEPHALUS.

Spatallæ sp. *Salisb. Parad.*

CHAR. GEN. *Calyx* quadrifidus, æqualis, totus deciduus. *Stigma* verticale, clavatum. *Nux* ventricosa, brevissimè pedicellata v. basi emarginata. *Involucrum* subsimplici serie 3—6-phyllum, definitè pauciflorum v. uniflorum : fructiferum non mutatum. *Receptaculum* epaleatum.

HABITUS. Frutices. *Ramis virgatis*. *Folia sparsa, filiformia v. plana, indivisa, infima rariùs bipinnatifida*. *Involucra subsessilia, unibracteata, in spicam capituliformem basi nunc bracteis imbricatis subtensam, congesta*. Flores *purpurascentes*.

ETYM. *σωρος* cumulus, et *κεφαλή* caput; ob capitula congesta.

Obs. Genus complectens phalanges duas facie et structurâ parùm diversas, quarum *prima* habitu et inflorescentiâ *Spatallæ* proxima, diversa tamen stigmate verticali, calyceque semper regulari: *secunda* e speciebus inter se convenientibus capitulo communi involucrato, sed discrepantibus numero florum foliolorumque involucri partialis, nec non foliis in quibusdam filiformibus, in aliis planis, et in unicâ dimorphis instar *Niveniæ*: fructus in hujus

sectionis duabus speciebus tantummodo observatus, in alterâ (foliis filiformibus) brevissimè pedicellatus, basi obsoletè emarginatâ, tenuissimè pubescens; in alterâ (foliis planis) glaberrimus, sessilis, basi angustatâ, profundè emarginatâ.

† *Spica nudiuscula*. Involucra 1—3-flora. Nux brevissimè pedicellata, basi integrâ. Folia filiformia, indivisa.

*1. *S. setaceus*, involucris unifloris, foliis setaceis incurvis (uncialibus) ramulisque hirsutis.

HAB. In Africâ Australi. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Frutex* erectus. *Rami* virgati, stricti, umbellati. *Folia* frequentia, vix sesquiuncialia, mucrone setaceo, spha-celato: inferiora minùs incurva. *Capitulum* terminale, sessile, ovatum, magnitudine cerasi nigri. *Calyx* unguibus laxiùs tomentosus; laminis barbatis. *Stigma* conico-ovatum.

*2. *S. salsoloides*, involucris unifloris, foliis triquetro-filiformibus incurvis (semuncialibus) glabris.

HAB. In Africâ Australi. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Frutex* erectus, ramosissimus. *Rami* glabri, ramuli tenuissimè pubescentes. *Folia* frequentia, semiteretia, suprâ sulcata, mucrone acuto subconcolori. *Capitulum* terminale, sessile, ovatum, vix magnitudine cerasi nigri, bracteolis paucis, brevissimis, lanceato-linearibus, subtensum. *Calyx* barbatus, villis brevibus. *Stigma* erectum v. parùm inclinans.

*3. *S. imberbis*, involucris trifloris, laminis calycis acuminibusque bractearum glabris.

^{141]} HAB. In Africâ Australi. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* erectus, ramosissimus. *Ramuli* pubescentes. *Folia* glabra, uncialia, modicè patentia, parùm incurva, suprâ sulcata, acutè mucronata. *Capitulum*

terminale, breviter pedunculatum, subglobosum, magnitudine cerasi nigri. *Bractea* lanceolata, ciliatæ, acumine subulato, glabro. *Calycis* unguis barbati. *Stylus* strictus. *Stigma* ovato-clavatum; æquale.

*4. *S. spatuloides*, involucris trifloris subpedicellatis, calycis laminis barbatis.

HAB. In Africâ Australi; prope Franche-hoek. *D. Niven.* (v. s. in Herb. Soc. Linn., et *D. Hibbert.*)

DESC. *Frutex* erectus. *Rami* umbellati, tenuissimè pubescentes. *Folia* modicè patentia, parùm incurva, vix uncialia, juniora pilosa. *Capitula* solitaria v. 2—3 aggregata, breviter pedunculata, ovata v. oblonga, magnitudine avellanæ. *Bractea* lanceolata, acutæ, pubescentes, apice quandoque glabriusculo. *Calycis* laminæ longiùs barbatae. *Stylus* apice sæpiùs incurvo, modò rectiusculo. *Stigma* styli hamati parùm inæquale; rectiusculi æquilaterale, ovatum.

†† *Spica subinvolucrata.* Involucra 4—6-flora. *Nux basi emarginatâ.*

*5. *S. tenuifolia*, foliis filiformibus (semunciâ brevioribus), capitulis paucifloris, calycis laminis plumoso-barbatis: interiori nudiusculâ.

HAB. In Africâ Australis montosis; in humidis prope Breede River. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* 3—4 pedes altus (*Niven*), facie *Spatallæ proliferae*. *Rami* glabri, rubicundi, vestiti; ramuli villosiusculi. *Folia* imbricata, scabriuscula, mucrone acuto; juniora hirsuta. *Capitulum commune* terminale, sessile, magnitudine pisi, e partialibus 2—4 compositum. *Involucra partialia* subimbricata, foliolis lanceolatis, barbatis, apice glabriusculo. *Calyx* profundè quadrifidus, æqualis. *Stylus* strictus. *Stigma* æquilaterale, erectum, ovatum.

6. *S. lanatus*, foliis triquetro-filiformibus (semunciâ longioribus) suprâ sulcatis, capitulis multifloris, calycis laminis omnibus plumoso-barbatis.

Protea lanata. *Thunb. Diss. n. 30,* t. 3. Prod. 26.*

Willd. Sp. Pl. 1, p. 519. *Poiret, Encyc. Botan.* 5, p. 653.

HAB. In Africæ Australis montosis. Swartland. (v. s. in Herb. Banks.)

DESC. *Frutex* erectus. *Rami* subumbellati, stricti, vestiti, tenuissimè pubescentes. *Folia* imbricata, 5—8 lineas longa. *Capitulum* terminale, solitarium, sessile, globosum, magnitudine avellanæ majoris: *partialia* densissimè congesta, 5—8-flora: *Involucris* 5—7-phyllis, foliolis augusto-lanceolatis, barbatis. *Calyx* profundè 4-fidus, æqualis. *Stylus* strictus. *Stigma* ovatum, æquilaterale, stylo ferè duplò crassius. *Nux* brevissimè pedicellata basique leviter emarginata, tenuissimè pubescens, cortice tenui, rugosiusculo, fusco.

OBS. Variat foliis subtùs triquetris teretibusque, scabriusculis et lævitus.

7. *S. imbricatus*, foliis lanceolatis subtùs scabris, unguibus calycis glanduloso-pilosis, stigmatè clavato.

Protea imbricata. *Thunb. Diss.* n. 45, t. 5. *Prod.* 27. *Linn. Suppl.* 116.* *Lam. Illust. Gen.* 1, p. 235, n. 1222. *Willd. Sp. Pl.* 1, p. 527. *Poiret, Encyc. Botan.* 5, p. 643. *And. Repos.* 527.

¹⁴⁸⁷ HAB. In Africæ Australis montibus. (v. s. in Herb. var. et v. in Hort. D. Hibbert.)

DESC. *Frutex* erectus. *Rami* elongati, stricti. *Folia* imbricata, subtùs convexiuscula venoso-striata, suprà concaviuscula lævia, unguicularia, mucrone incurvo. *Capitulum* terminale, sessile, subovatum, solitarium, v. e. 2—3 aggregatis compositum. *Involucrum commune* polyphyllum, imbricatum, capitulo brevius; foliolis lanceolatis, membranaceis, coloratis, scabriusculis. *Involucra partialia* sæpiùs quadriflora, tetraphylla; foliolis lanceolatis, hirsutis. *Calyx* tubo gracili, laminis barbatis. *Ovarium* barbatum. *Stylus* strictus. *Stigma* elliptico-clavatum, hinc gibbosiusculum. *Nux* glaberrima, nitens, fusca, oblonga, basi angustatâ concolori emarginatâ.

*8. *S. diversifolius*, foliis spathulato-lanceolatis subtùs

lævibus : infimis bipinnatifidis, unguibus laminisque calycis barbatis, stigmatè cylindræo.

HAB. In Africæ Australis montibus saxosis, prope Goud Rivier. *D. Niven.* (v. s. in Herb. Banks., Lambert., et Hibbert.)

DESC. *Frutex* erectus, glaber, bipedalis usque orgyalis, indivisus, v. bifidus, strictus, crassitie pennæ olorinæ, suprâ pubescens. *Folia infima* trifido-bipinnatifida, canaliculata, biuncialia ; *reliqua* imbricata, obtusiuscula, parùm concava, vix semuncialia. *Capitulum* terminale, solitarium, sessile, ovatum, obtusum, magnitudine pruni minoris.

11. SPATALLA.

Salisb. Parad.

CHAR. GEN. *Calyx* quadrifidus, laciniâ inferiore (in plerisque) majore, totus deciduus. *Stigma* obliquum, dilatatum. *Nux* ventricosa, brevissimè pedicellata. *Involucrum* simplici serie 2—4-phyllum, uniflorum v. definitè pauciflorum. *Receptaculum* epaleatum.

HABITUS. Frutices. *Folia sparsa, filiformia, indivisa.* *Involucra terminalia, spicata v. racemosa, unibracteata, fructifera haud mutata.* Flores *purpurascentes.* *Anthera lacinie majoris calycis proportionatim major, et in quibusdam unica fertilis.*

† *Involucra uniflora. Stigma concavum, cochleariforme. Calyx inæqualis.*

*1. *S. mollis*, involucro diphylo : foliolis integerrimis, foliis strictis ramulisque villosis.

HAB. In Africæ Australis montibus. *D. Joh. Roxburgh.* (v. s. in Herb. Lambert.)

DESC. *Frutex* erectus, ramosissimus. *Rami* rubicundi, ramuli graciles, erecti. *Folia* erecto-patentia, 7—8 lineas longa, callo acutissimo, villis modicè patentibus sericea. *Spica* sessilis, erecta, solitaria, oblongo-cylindræa, densa, racemosa, vix uncialis. *Bracteæ* foliaceæ, pedicellis duplo

longiores. *Involucrum* foliolis ovatis, villosis, exteriore latiore. *Calyx* densè barbatus, laminâ lacinia majoris villis marginalibus inflexis. *Squamulae hypogynae* quatuor, lineares, persistentes.

*2. *S. pedunculata*, involucro diphylo: foliolo latiore tridentato, spicâ imbricatâ, pedunculo foliis longiore triquetris incurvis basi attenuatis, bracteis sericeis involucro brevioribus.

HAB. In Africae Australis montibus. Kleine-Hoot-Hoek. *Gul. Roxburgh, M.D.* (v. s. in Herb. Soc. Linn.)

DESC. *Frutex* erectus, ramosissimus, foliis ramisque adultis glabris, junioribus sericeis. *Folia* frequentia, ferè uncialia, basi attenuatâ, erectâ, suprâ patentia, falcato-incurva, ^{145]} callo apicis obtusiusculo. *Pedunculi* sesquiunciales, solitarii, sericei; bracteis alternis, subulatis. *Spica* cylindracea, pedunculo vix longior; pedicellis, involucris, calycibusque sericeis.

*3. *S. nivea*, involucro diphylo: foliolo latiore tridentato, spicâ imbricatâ, pedunculo foliis rectiusculis acutissimis brevior, bracteis foliaceis villosiusculis involucra æquantibus.

HAB. In Africae Australis montibus. *D. Niven.* (v. s. in Herb. D. Hibbert.)

DESC. *Frutex* erectus, ramosissimus, ramis folisque adultis glabris, novellis sericeis. *Folia* uncialia, leviter incurva, v. rectiuscula, basi parùm attenuata. *Pedunculi* solitarii, subsericei, bracteis alternis subulatis. *Spica* sesquiuncialis, pedunculo duplò longior. *Involucrum* foliolium exterius profundè tridentatum, dente intermedio angustiore. *Calycis* laminæ villis brevibus, patulis, niveis barbatae.

*4. *S. ramulosa*, involucro diphylo: foliolo latiore trifido, spicâ sessili imbricatâ, bracteis superioribus longitudine pedicellorum, foliis acutè mucronatis.

Protea foliis setaceis, floribus racemosis. *Hort. Cliff.* 496 ?

Leucadendron racemosum. *Linn. Sp. Pl. ed. i, p. 91 ?*

ed. ii, p. 194? *Berg. Act. Stockh.* 1766, p. 325.* *Perg. Cap.* p. 23.*

Protea racemosa. *Thunb. Diss. n.* 21*? *Prod.* 26?

HAB. In Africæ Australis montibus. Roode Zant Kloof. (v. s. in *Herb. Banks., Soc. Linn.*)

DESC. *Frutex* erectus, ramosissimus; ramis virgatis filiformibus, foliisque adultis glabris, novellis sericeis. *Folia* frequentia, modicè patentia, parùmque incurva, basi attenuata, vix uncialia, suprâ canaliculata, subtùs convexa, callo acuto mucroniformi. *Spica* terminalis, breviter pedunculata, cylindræa, uncialis, sesquiuncialis, densa, subracemosa, floribus omnibus imbricatis, ramulo uno alterove sericeo brevi, sæpissimè stipata. *Bracteæ* omnes pedicellos pariter tomentosos æquantes. *Involucrum* laciniâ mediâ labii majoris angustiore. *Calyx* breviter densèque barbatus, villis marginalibus, laminæ majoris arcuè inflexis. *Stigma* cochleariforme, papillâ centrali.

*5. *S. laxa*, involucri diphyllo: foliolo latiore trifido, racemo subpedunculato laxiusculo, bracteis superioribus pedicello brevioribus.

HAB. In Africæ Australis montibus. Kleine-Hoot-Hoek. *Gul. Roxburgh, M.D.* (v. s. in *Herb. Banks., Lambert., Soc. Linn.*)

DESC. *Frutex* erectus, 4—6 pedalis, (*Niven*) ramosus. *Rami* graciles, virgati, rubicundi, ramuli subsericei. *Folia* patenti-erecta, leviter incurva, v. rectiuscula, basi attenuata, callo apicis acutiusculo, v. obtusiusculo, uncialia, inferiora glabra, superiora sericea. *Racemi* breviter pedunculati, solitarii, erecti, sesquiunciales, ramulo brevi quandoque stipati. *Bracteæ* tomentosæ, pedicellis fructiferis breviores; inferiores floriferorum subæquantes. *Involucra* vix longitudine pedicellorum, sericea, fructifera labio majore tripartito, laciniâ intermediâ angustissimâ. *Nux* ovata, subsessilis, sericea, involucri persistenti duplò longior *stilo* curvato diù coronata, basi barbata pilis strictis.

*6. *S. bracteata*, involucri diphyllo: foliolo latiore profundè trifido, spicâ pedunculatâ imbricatâ, bracteis teretibus

involucra pedicellata superantibus, foliis incurvis (uncialibus) glabriusculis.

^{147]} *Protea racemosa*. Linn. Mant. 187? (*Herb. Linn.*)

HAB. In Africæ Australis montibus. Franche Hoek. (v. s. in *Herb. Banks., Lambert., Soc. Linn.*)

DESC. *Frutex* erectus, 6—7 pedalis, (*Niven*) ramosissimus, ramulis ultimis sericeis. *Folia* e basi attenuatâ, adpressâ, suprâ patentia, et falcato- v. sigmoideo-curvata, callo obtusiusculo, adulta glabra, recentiora sericea, quandoque sesquiuncialia. *Pedunculi* terminales, solitarii, spicâ sesquiunciali breviores. *Pedicelli* imbricati, inferiores involucra æquantés, superiores iisdem parùm breviores. *Involucra* sericea, labio majore sæpè tripartito. *Calyx* unguibus tomento adpresso; laminis barbatis villis modicè patentibus, marginalibus haud inflexis.

*7. *S. sericea*, involucro diphylo: foliolo latiore tripartito, spicâ sessili imbricatâ: bracteis involucra subsessilia æquantibus, foliis semuncialibus ramulisque sericeis.

HAB. In Africæ Australis montibus. *Gul. Roxburgh, M.D.* (v. s. in *Herb. Soc. Linn.*)

DESC. *Frutex* erectus, ramosissimus. *Rami* ramulique virgati, stricti, hi sericei, illi glabri. *Folia* frequentia, imbricata, patent-erecta, rectiuscula v. leviter incurva, suprâ obsoletissimè sulcata. *Spicæ* solitariae, vix unciales. *Involucra* labio majore laciniis subulatis, mediâ augustiore. *Calyx* unguibus tomentosis, laminis barbatis.

8. *S. prolifera*, involucro tetraphyllo: foliolis apice sphacelatis, spicâ conico-capitatâ: floribus subsessilibus.

Protea prolifera. *Thunb. Diss. n. 27,* tab. 4. Prod. 26. Linn. Suppl. 118. Lam. Illust. Gen. 1, p. 238, n. 1233. Willd. Sp. Pl. 1, p. 518. Poiret, Encyc. Botan. 5, p. 654.*

^{148]} HAB. In Africæ Australis montibus, Hottentots-Holland: Roode Zant. (v. s. in *Herb. Banks., Lambert., Soc. Linn.*)

DESC. *Frutex* erectus, sesquipedalis, bipedalis, ramosissimus. *Rami* ramulique umbellati, hi subsericei, illi

rubicundi glabriusculi. *Folia* imbricata, conferta, vix unguicularia, ramulorum recentiorum sericea. *Spica* sessilis. *Bracteæ* foliaceæ. *Involucri* foliola subulata, demùm glabriuscula. *Calyx* densissimè barbatus, villis brevibus sericeis; laminâ interiori duplò majori, villis marginalibus arcuè inflexis. *Stigma* planiusculum, papillâ centrali. *Squamulæ hypogynæ* quatuor, lineari-subulatæ.

*9. *S. pyramidalis*, involucro tetraphyllo: foliolis acuminatis pedicellos subæquantibus, spicâ erectâ solitariâ sessili oblongo-pyramidali foliis semuncialibus duplò longiore.

HAB. In Africæ Australis montibus, prope Zwellendam. *Gul. Roxburgh, M.D.* (v. s. in Herb. Lambert. et Soc. Linn.)

DESC. *Frutex* erectus, ramosissimus, ramis ramulisque umbellatis, pubescentibus. *Folia* confertissima, modicè patentia, stricta v. parùm incurva, villosiuscula, callo acuto, mucroniformi. *Spica* densa, subuncialis. *Bracteæ* foliaceæ, involucra æquantés. *Involucra* pubescentia, foliolis e latiore basi subulatis, apice patulis, exteriori parùm angustiore. *Calyx* laminâ interiori parùm majori, villis marginalibus simplicibus. *Stigma* concavum, papillâ centrali. *Squamulæ hypogynæ* lineari-subulatæ. *Receptaculum* barbatum.

*10. *S. polystachya*, involucro tetraphyllo: foliolis apice patulis, spicis nutantibus aggregatis pedunculatis, foliis uncialibus curvatis.

HAB. In Africæ Australis montibus. *Gul. Roxburgh, M.D.* (v. s. in Herb. Lambert. et Soc. Linn.)

DESC. *Frutex* erectus, ramosissimus. *Rami* ramulique umbellati, rubicundi, ultimi pubescentes. *Folia* conferta, patula, subsigmoideo-curveda, villosa, mucrone acutissimo, novella sericea. *Spicæ* 4—6, reflexæ, sesquiunciales, breviter pedunculatæ, ramulis umbellatis longioribus stipatæ. *Bracteæ* pedicellis ter longiores. *Involucra* foliolis subæqualibus, concavis, lanceolato-subulatis, acuminatis. *Calyx* subæqualis. *Stigma* planiusculum, papillâ centrali. *Nux* brevissimè pedicellata, tenuissimè pubescens.

† Involucra 3—4-flora. Stigma convexiusculum. Calyx subæqualis.

11. *S. incurva*, spicis racemosis subpedunculatis, bracteis involucro tomentoso (sub-4-floro) brevioribus, foliis incurvis, calycibus inæqualibus.

a. Spicæ sæpè aggregatæ. Bracteæ pedicellos subæquantes. Folia ferè uncialia, inferiora ramulorum glabra.

Protea incurva. *Thunb. Diss. n. 22,* tab. 3*, bona. *Willd. Sp. Pl. 1, p. 516*. *Poiret, Encyc. Botan. 5, p. 652*.

β. Spicæ solitariae. Bracteæ pedicellos superantes. Folia semuncialia, ferè omnia ramulorum sericea.

HAB. In Africæ Australis arenosis humidis subumbrosis; Roode Zant Cascade. (v. s. a. in Herb. Banks., Lambert., Soc. Linn.; β. in Herb. Hibbert.)

OBS. I. Calyx inæqualis. Stigma planiusculum, papillâ centrali.

OBS. II. β. Forsan distincta species: Foliis confertissimis, pedicellis involucro ferè dimidio brevioribus.

1607 *12. *S. propinqua*, spicâ subpedunculatâ, bracteis subulatis foliaceis involucra subsessilia tomentosa subbiflora æquantibus, foliis semuncialibus strictis ramulisque villosis, calycibus subæqualibus.

HAB. In Africâ Australi. *A. Auge*. (v. s. in Herb. Banks.)

OBS. Spica biuncialis. Pedicelli brevissimi. Nux pedicello manifesto, glabro, tenuissimè pubescens.

13. *S. caudata*, spicâ sessili, bracteis involucrisque ovato-lanceolatis glabriusculis ciliatis, foliis glabris acutis.

Protea caudata. *Thunb. Diss. sec. ic. tab. 2*.

HAB. In Africâ Australi; prope Palmetta River. *D. Masson*. (v. s. in Herb. Banks.)

DESC. *Frutex* erectus, ramosissimus; ramis umbellatis glabriusculis. *Folia* vix semuncialia, suprâ canaliculata, acuta, stricta. *Spicæ* sæpè aggregatæ, cylindraceæ, densæ, unciales, quandoque biunciales. *Involucra* subsessilia,

sæpiùs triflora. *Calyx* subæqualis, barbatus. *Stigma* convexum. *Nux* tenuissimè pubescens.

14. *S. Thunbergii*, spicâ sessili, bracteis involucrisque ovato-lanceolatis villosis, foliis calyce longioribus acutis canaliculatis ramisque pilosis.

Protea caudata. *Thunb. Diss. n. 23,** secund. descript.

HAB. In Africæ Australis montosis. *D. Niven.* (v. s. in Herb. Hibbert.)

DESC. *Frutex* erectus, ramosissimus. *Folia* vix semuncialia, conferta, imbricata, stricta v. parùm incurvâ. *Spica* cylindracea, densa, uncialis, sesquiuncialis. *Involucra* brevissimè pedicellata, bracteis parùm longiora, villis persistentibus. *Calyx* subæqualis, laminis brevissimè barbatis, subsericeis. *Stigma* convexum. *Nux* tenuissimè pubescens, pedicello brevissimo, crasso, glabro.

*15. *S. brevifolia*, foliis calyce brevioribus obtusiusculis subsericeis triquetris, spicis densis, bracteis involucrisque pubescentibus.

HAB. In Africæ Australis montosis. *D. Masson.* (v. s. in Herb. Banks. et D. Aiton.)

DESC. *Frutex* erectus, ramis umbellatis, virgatis, pubescentibus. *Folia* subtriquetra, suprâ canaliculata, patentia erecta, villosiuscula, subtrilinearia. *Spica* solitaria, sessilis, uncialis, sesquiuncialis, rachi pedicellis bracteisque pubescentibus. *Bracteæ* e basi membranaceâ, lanceolatâ, subulatâ. *Involucra* brevissimè pedicellata, 2—3-flora. *Calyx* æqualis. *Stigma* convexum, papillâ elevatiore. *Squamulæ hypogynæ* quatuor subulatæ.

12. ADENANTHOS.

Labill. Nov. Holl. 1, p. 28.

CHAR. GEN. *Calyx* quadrifidus, infrâ circumscissus. *Squamulæ* quatuor hypogynæ, basi persistenti calycis adnatæ.

Pistillum calyce longius. *Stigma* verticale. *Nux* ventricosa. *Involucrum* uniflorum, imbricatum, 4—8-phyllum.

HABITUS. Frutices. Folia *sparsa, in diversis varia*. Flores *axillares, solitarii, rubicundi; raro terminales, subaggregati, lutescentes*.

1. *A. obovata*, foliis obovatis integerrimis glabris.

Adenanthos obovata. *Labillard. Nov. Holl. 1, p. 29,* tab. 37.*

HAB. In collibus saxosis oræ australis Novæ Hollandiæ; Lewins Land. (ubi v. v.)

152] 2. *A. cuneata*, foliis cuneatis sericeis apice dentato-crenato.

Adenanthos cuneata. *Labillard. Nov. Holl. 1, p. 28,* tab. 36.*

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: prope littora. (ubi v. v.)

3. *A. sericea*, foliis filiformibus biternatis sericeis, floribus axillaribus solitariis, stylo glabro.

Adenanthos sericea. *Labillard. Nov. Holl. 1, p. 29,* tab. 38.*

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in arenosis prope littora. (ubi v. v.)

4. *A. terminalis*, foliis filiformibus trifidis: lacinâs lateralibus bifidis intermediâ indivisâ, floribus terminalibus solitariis ternisve, stylo villosa.

HAB. In Novæ Hollandiæ orâ australi. Flinders' Land: in depressis prope littora. (ubi v. v.)

13. SIMSIA.

CHAR. GEN. *Calyx* tetraphyllus, regularis, laminis reflexis. *Stamina* exserta. *Antheræ* tandem liberæ, primò cohærentes, lobis proximis vicinarum loculum constituentibus. *Stigma* dilatatum, concavum. *Nux* obconica.

HABITUS. Frutices *humiles, glabri*. Folia *alterna, filiformia, dichotoma, petioli basi dilatata*. Capitula *globosa, parva, terminalia, racemosa, v. paniculata, involucri brevis v. nullo*. Flosculi *flavi, glabri, unibracteati*.

I have named this genus in honour of Dr. John Sims, the respectable editor of the 'Botanical Magazine.'

*1. *S. tenuifolia*, capitulis nudis, paniculæ ramis sub-unifloris bracteolatis.

HAB. In Novæ Hollandiæ orâ australi: Lewins Land; ad latera saxosa collium. (ubi v. v.)

*2. *S. anethifolia*, capitulis involucriatis bracteolis imbricatis, paniculæ ramis multifloris: ramulis capitula subæquantibus.

HAB. In Novæ Hollandiæ orâ australi: Lewins Land; in arenosis prope littora. (ubi v. v.)

14. CONOSPERMUM.

Smith, Linn. Trans. vol. 4. Exot. Bot. Gart. Carp. 3, p. 198, t. 215.

CHAR. GEN. Calyx tubulosus, ringens, laciniâ supremâ basi fornicatâ. Antheræ tres, inclusæ, laterales dimidiatæ, superior biloba: primò cohærentes, lobis proximis vicinarum loculum constituentibus. Stigma liberum. Nuæ obconica, papposa.

HABITUS. Frutices. Folia *sparsa, integerrima, plana, rariusve filiformia*. Spicæ *axillares v. terminales, compositæ, sensim florentes, hinc corymbosæ*. Flores *solitarii, sessiles, unibracteati, albi v. cærulescentes*; Calyce *deciduo*; Bractea *cucullatâ persistenti*.

OBS. Jussieu and Ventenat have referred this genus to the natural order Thymeleæ; but that it is a genuine Proteacea, as Dr. Smith has considered it, is proved by the erect embryo, the terminal style, and the æstivation of the Calyx; and is rendered evident by its affinity to Simsia,

which, with the more usual appearance of this order, agrees with *Conospermum* in the structure of its *Antheræ*.

† *Calycis lacinie acutæ, tubo vix longiores. Conosperma vera.*

1. *C. ellipticum*, foliis ovali-oblongis obtusis mucronulatis aveniis, pedunculis axillaribus.

¹⁸⁴³ *Conospermum ellipticum. Smith in Rees, Cyclop.*

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson; in ericetis aridis. (ubi v. v.)

2. *C. taxifolium*, foliis lanceolato-linearibus acutis mucronatis tenuissimè pubescentibus verticalibus, basi tortis, pedunculis axillaribus.

Conospermum taxifolium. Smith in Rees, Cyclop.

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson; in ericetis. (ubi v. v.)

3. *C. ericifolium*, foliis subulato-filiformibus imbricatis, spicis axillaribus pedunculo brevioribus.

Conospermum ericifolium. Smith in Rees, Cyclop.

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson; in ericetis. (ubi v. v.)

4. *C. longifolium*, foliis oblongis linearibusve planis venosis, pedunculisque elongatis scapiformibus, corymbis decompositis, calycis limbo extûs pubescenti tubum vix æquante.

Conospermum longifolium. Smith, Exot. Bot. 2, p. 45, t. 82.

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson; in ericetis collibusque saxosis. (ubi v. v.)

*5. *C. tenuifolium*, foliis lineari-filiformibus subcanaliculatis aveniis, pedunculisque elongatis scapiformibus, corymbis subsimplicibus, calycis limbo extûs pubescenti tubo longiore.

HAB. In Novæ Hollandiæ orâ orientali, prope Port Jackson; in collibus arenosis prope littora. (ubi v. v.)

*6. *C. cæruleum*, foliis oblongis lanceolatisve planis venosis, pedunculisque elongatis scapiformibus, corymbis compositis, calycis limbo glaberrimo tubo longiore.

HAB. In Novæ Hollandiæ orâ australi : Lewins Land. (ubi v. v.)

†† *Calycis lacinia caudatæ*. Chilurus.

*7. *C. teretifolium*, foliis teretibus pedunculisque elongatis, corymbis compositis.

HAB. In Novæ Hollandiæ orâ australi ; Lewins Land : in collibus saxosis. (ubi v. v.)

*8. *C. capitatum*, foliis linearibus elongatis tortilibus, capitulis sessilibus e spiculis paucifloris congestis.

HAB. In Novæ Hollandiæ orâ australi ; Lewins Land : in collibus apricis graminosis. (ubi v. v.)

††† *Incertæ tribus*.

*9. *C. distichum*, foliis filiformibus subdistichis curvatis, spicis axillaribus indivisis.

HAB. In Novæ Hollandiæ orâ australi ; Lewins Land : in ericetis. (ubi v. v. flor. delaps.)

15. SYNAPHEA.

CHAR. GEN. *Calyx* tubulosus, ringens, laciniaâ supremâ latiore. *Antheræ* tres, inclusæ, laterales dimidiatæ, inferior biloba : primò cohærentes, lobis proximis vicinarum loculum constituentibus ! *Stigma* filamento superiori sterili connatum ! *Nux* obovata.

HABITUS. Frutices humiles. Folia sparsa, plana, pulcherrimè reticulata, circumscriptione cuneiformia, lobata, inferiora ejusdem fruticis sæpiùs indivisa : petioli basi dilatata semivaginant. Spicæ axillares v. terminales, simplices v. ramosæ. Flores alterni, solitarii, sessiles, uni-

bracteati. Calyx flavus, deciduus, quadripartibilis. Bractea cucullata, persistens.

ETYM. *συναφή* connectio, ob peculiarem cohærentiam stigmatis v. apicis styli cum filamento sterili.

*1. *S. favosa*, foliis oblongo-cuneiformibus indivisis trilobisque: lobis integris, petiolis spicisque glabris, stigmatе bicorni.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

2. *S. dilatata*, foliis apice dilatatis trilobis: lobis incindentatis, petiolis spicisque villosis, stigmatе bicorni.

Conospermum reticulatum. *Smith in Rees, Cyclop.*

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*3. *S. petiolaris*, foliis rameis petiolos subæquantibus tripartitis: lobis divisis planis; infimis trilobis integrisque, spicis elongatis ramosis, stigmatе acuto.

Polypodium spinulosum. Burm. Ind. p. 233, t. 67, f. 1, vel ad hanc v. ad plantam congenerem pertinere videtur.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*4. *S. polymorpha*, foliis rameis brevissimè petiolatis tripartitis canaliculatis: lobis subdivisis; infimis indivisis trilobisque, spicis simplicibus pedunculo longioribus, stigmatе acuto.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

16. FRANKLANDIA.

CHAR. GEN. *Calyx* hypocrateriformis, limbo quadripartito, plano, deciduo, tubo persistenti. *Antheræ* inclusæ, calyci adnatæ! *Squamæ* hypogynæ, in vaginam connatæ. *Nux* fusiformis, pedicellata, apice dilatato papposo.

HABITUS. Frutex *glaber*. Folia *alterna, filiformia, dichotoma*. Spicæ *axillares, indivisæ, floribus alternis, unibracteatis, sordidè flavis*. Pollen *sphæricum*. Cotyledones *brevissimæ*!

This genus is named in honour of Sir Thomas Frankland, Baronet, to whom English botany is much indebted, and whose valuable observations and excellent figures of submarine plants it is hoped he may be induced to communicate to the public.

*FRANKLANDIA *fucifolia*.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis humidis. (ubi v. v.)

17. SYMPHIONEMA.

CHAR. GEN. *Calyx* regularis, tetraphyllus, basi cohærens, medio staminifer. *Filamenta* apice cohærentia! *Antheræ* distinctæ. *Glandulæ* nullæ hypogynæ. *Ovarium* dispersum. *Stigma* subtruncatum. *Nux* monosperma, cylindræa.

HABITUS. Suffrutices v. Herbæ *glabræ, pilisve raris glandulosis*. Folia *tripartita, lobis divisis; inferiora opposita*! Spicæ *terminales et e summis alis, simplices*. Flores *alterni, sessiles, unibracteati*. Calyx *flavus, deciduus*. Bractæ *cucullatæ, persistentes*.

*1. *S. paludosum*, laciniis foliorum subulatis semi-teretibus, rachibus bracteisque glaberrimis.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis paludosis. (ubi v. v.)

*2. *S. montanum*, laciniis foliorum planis linearibus uninerviis, rachibus bracteisque pubescentibus pilis glandulosis brevissimis.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in rupibus humidis. (ubi v. v.)

18. AGASTACHYS.

CHAR. GEN. *Calyx* regularis, tetraphyllus, basi cohærens, medio staminifer. *Filamenta* distincta. *Glandulæ* nullæ hypogynæ. *Ovarium* sessile, monospermum, trigonum. *Stigma* unilaterale.

HABITUS. Frutex *glaberrimus*. Folia *sparsa, integerrima, plana*. Spicæ *numerosæ, terminales et e summis alis, simplices*. Flores *alterni, sessiles, unibracteati*. Calyx *flavescens, deciduus*. Pistillum *staminibus brevius*. Bractææ *encullatæ, persistentes*.

ETYM. ἀγασταχὺς spicis abundans.

*AGASTACHYS *odorata*.

HAB. In Insulæ Diemen plagis australioribus; prope Adventure Bay: ubi primum a *D. Nelson* detecta, nuperius lecta a *D. G. Caley*. (v. s. in Herb. Banks.)

19. CENARRHENES. *Labill. Nov. Holl.* 1, p. 36, t. 50.

CHAR. GEN. *Calyx* tetraphyllus, regularis, foliolis supra angustatis, deciduus. *Stamina* basi calycis inserta. *Glandulæ* ^{159]} quatuor hypogynæ, staminiformes. *Ovarium* sessile, monospermum. *Stigma* simplex. *Drupa* baccata.

HABITUS. Arbor *glabra*. Folia *alternata; plana, dentato-serrata, nitida*. Spicæ *axillares, simplices*. Flores *alterni, sessiles, unibracteati*.

OBS. Labillardière considers this genus as most nearly related to *Lauri*. Jussieu, however, has (in *Annales du Muséum*, v. 5, p. 224) stated sufficient reasons for excluding it from that order, but has not attempted to determine its affinity. I have ventured to place it in Proteaceæ, from the structure of its fruit, stamens and calyx, and the only circumstance in which it differs from them, consists in its having (according to Labillardière) four barren

stamina; but even these occupy the place of the glands or scales usually found in the order, and the resemblance they bear to stamina in this genus, may assist in explaining their nature in all: nor does their being in most cases secreting organs render this view of their origin improbable; for the function of secretion, which, as it is far from universal, must be considered as only of secondary importance in assisting impregnation, is more frequently accomplished by the modification of some of the usual parts of the flower than by the production of an additional organ.

CENARRHENES nitida. Labill. Nov. Holl. 1, p. 36,* t. 50.

HAB. In Insulæ Diemen plagis australioribus. *Labillardière.* (v. s. cum fructu sed floribus delapsis in Herb. D. Lambert.)

20. *PERSOONIA.* Smith in Linn. Trans. iv. Gært. Carp. 3, p. 218, t. 220. Pentadactylon. Gært. l. c., p. 219, t. 220. Linkia. Cav. Ic. 4.

CHAR. GEN. *Calyx* tetraphyllus, regularis, foliolis medio staminiferis, suprâ recurvis, deciduus. *Stamina* exserta. *Glandulæ* quatuor hypogynæ. *Ovarium* pedicellatum, 1-loculare, 1—2-spermum. *Stigma* obtusum. *Drupa* baccata; *Nuce* 1—2-loculari!

HABITUS. Frutices v. Arbusculæ, cortice in quibusdam scarioso-lamellosa. Folia sparsa, integerrima, sæpius plana. Pedunculi axillares, solitarii, ebracteati, v. racemosi, unibracteati. Flores flavi. Pedicellus ovarii in quibusdam articulatus! Cotyledones sæpius plures!

*1. *P. teretifolia*, foliis filiformibus exsulcis, pedunculis unifloris solitariis, antheris acuminatis, styli ovario brevioribus.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*2. *P. microcarpa*, foliis filiformibus canaliculatis, pedunculis solitariis geminis ternisve, antheris muticis, stylis ovario aliquoties longioribus, stigmatibus cernuo.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis paludosis. (ubi v. v.)

*3. *P. pinifolia*, foliis filiformibus laxis, spicâ foliatâ elongatâ pyramidalis: foliis floralibus abbreviatis, ovario monospermo.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis et ad ripas rivulorum. (ubi v. v.)

4. *P. juniperina*, foliis subulatis strictis pungentibus, pedunculis axillaribus sparsis spicisve foliatis abbreviatis, ovario dispermis glabris.

Persoonia juniperina. *Labill. Nov. Holl.* 1, p. 33,* tab. 45.

HAB. In Insulâ Diemen: et Novæ Hollandiæ orâ australi, prope Port Phillip: in ericetis aridis lateribusque collium. (ubi v. v.)

161] 5. *P. hirsuta*, foliis linearibus hirsutis scabris margine recurvis, pedunculis axillaribus, ovario monospermo sericeis.

Persoonia hirsuta. *Pers. Syn.* 1, p. 118.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis humidis. (ubi v. v.)

*6. *P. mollis*, foliis longo-lanceolatis villosis subtus mollissimis, calycibus barbatis, ovario dispermis glabris.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: ad ripas arenosas fluviorum. (ubi v. v.)

7. *P. linearis*, foliis angusto-linearibus elongatis glabris, pedunculis erectis calycibusque pubescentibus, pedicello ovario inarticulato, caule arborescenti: cortice lævi.

Persoonia linearis. *And. Repos.* 77. *Vent. Malmais.* 32. *Sims, Bot. Mag.* 760. *Pers. Syn.* 1, p. 118.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in campis et collibus. (ubi v. v.)

*8. *P. lucida*, foliis lanceolato-linearibus elongatis glabris, pedunculis erectis calycibusque pubescentibus, pedicello ovarii inarticulato, caule arborescenti: cortice scarioso-lamellosa.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in montosis ad ripas fluviorum. *D. Fer. Bauer.* (v. s.)

*9. *P. virgata*, foliis linearibus oblongo-linearibusve sparsis verticalibus glaberrimis margine lævibus, pedunculis erectis calycibusque glabris, caule arborescenti: cortice lævi.

HAB. In Novæ Hollandiæ orâ orientali; prope Sandy Cape: in arenosis prope littora. (ubi v. v.)

*10. *P. flexifolia*, foliis lanceolato-linearibus mucronatis confertis basi tortis utrinque lævibus punctis crystallinis micantibus; marginibus scabris, calycibus glabris, caule fruticoso.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: ad latera saxosa collium. (ubi v. v.)

*11. *P. scabra*, foliis lineari-lanceolatis mucronatis utrinque scabris punctis crystallinis aliisque minutissimis opacis conspersis, calycibus pubescentibus.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*12. *P. spathulata*, foliis lanceolato-spathulatis mucronatis concaviusculis utrinque scaberrimis punctis crystallinis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*13. *P. nutans*, foliis linearibus lævibus, pedunculis axillaribus recurvis calycibusque glabris.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in sylvis solo arenoso, ad radices montium. (ubi v. v.)

LIBRARY
NATIONAL MUSEUM
OF CANADA

*14. *P. falcata*, foliis elongato-lanceolatis basi attenuatis subpetiolatis falcatis aversis coriaceis, antheris acuminatis, caule arborescenti: cortice lamelloso.

HAB. In Novæ Hollandiæ orâ orientali; Endeavour River: *Jos. Banks, Bart.*: septentrionali, Carpentaria; prope littora. (ubi v. v. cum fruct. matur. flor. delaps.)

15. *P. lanceolata*, foliis lanceolatis ellipticisve mucro-
1637 natis glabris lævibus, pedunculis axillaribus unifloris, calycibus pube adpressâ subsericeis, pedicello ovarii inarticulato.

Persoonia lanceolata. *And. Repos.* 74. *Pers. Syn.* 1, p. 118.

β. *Persoonia latifolia.* *And. Repos.* 280?

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in campis ericetisque, prope littora. (ubi v. v.)

16. *P. salicina*, foliis lanceolato-oblongis in lateralibus aversis, racemis lateralibus pedunculis axillaribus unifloris, calycibus glabriusculis, caule arborescenti: cortice scarioso-lamelloso.

Linkia lævis. *Cavan. Ic.* 4, p. 61, t. 389? an varietas *P. lanceolata*?

Persoonia salicina. *Pers. Syn.* 1, p. 118.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in campis collibus et sylvis. (ubi v. v.)

17. *P. ferruginea*, foliis ellipticis æquilateralibus venosis adversis, pedunculis axillaribus multifloris calycibusque ferrugineo-tomentosis, caule erecto.

Persoonia laurina. *Pers. Syn.* 1, p. 118.

Persoonia ferruginea. *Smith, Exot. Bot.* 2, p. 47, t. 83.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in campis. (ubi v. v.)

*18. *P. prostrata*, foliis ovalibus obtusis margine pubescentibus, pedunculis axillaribus uni- v. paucifloris, caule procumbente.

HAB. In Novæ Hollandiæ orâ orientali; prope Sandy

Cape: in arenosis prope littora. (ubi v. v. cum fruct. matur., flor. delaps.)

*19. *P. elliptica*, foliis ellipticis venosis, racemis lateralibus, calycibus glabris, pedicello ovarii articulado.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: ad litora saxosa collium. (ubi v. v.)

*20. *P. articulata*, foliis elongato-lanceolatis æquilateralibus glabris, racemis lateralibus pedunculisve unifloris, calycibus glabriusculis, ovarii pedicelli articulo inferiore glandulas hypogynas æquante.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*21. *P. longifolia*, foliis elongato-linearibus falcatis, racemis lateralibus pedunculisve unifloris, calycibus pube adpressâ tectis, ovarii pedicelli articulo inferiore glandulis hypogynis longiore.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*22. *P. graminea*, foliis rameis linearibus longissimis margine recurvis, racemis secundis multifloris, calycibus glabris, caule suffruticoso abbreviato.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: ad ripas arenosas stagnorum. (ubi v. v.)

21. BRABEIUM.

Linn. Gen. Pl. 1, n. 85. *Mant.* 168; *ed. Schreb. n.* 1580.

CHAR. GEN. *Calyx* tetraphyllus, regularis. *Stamina* basi calycis inserta. *Vaginula* hypogyna. *Ovarium* sessile. *Stigma* verticale. *Drupa* exsucca, monosperma, putamine osseo.

HABITUS. Arbor. Folia (Theophrasti,) *verticillata*, serrato-dentata. Spicæ axillares, floribus fasciculatis, ternis

pluribusve, bractea communi subtensis, plerisque masculis pistillo imperfecto.

BRABEIUM stellatifolium.

Arbor hexaphylla æthiopica, foliis circa caulem ad intervalla senis. *Pluk. Alm.* 47, t. 265, f. 3.

Amygdalus æthiopica fructu holosericeo. *Breyn. cent.* 1, t. 1.

Brabejum. *Hort. Cliff.* 36. *Roy. Lugd. Bat.* 400.

Brabejum stellatifolium. *Linn. Sp. Pl. ed. i,* p. 121; *ed. ii,* p. 177. *Mant. p.* 332.*

Brabya. *Mant. p.* 137.*

Brabeium stellulifolium. *Linn. Syst. Veg.* xiii, p. 764. *Houtt. Nat. Hist. par.* 2, t. 6, p. 424, tab. 37; *ed. Germ.* t. 4, p. 647, t. 37, f. 1. *Lam. Encyc. Botan.* 1, p. 459*? *Illust. Gen. tab.* 847. *Willd. Sp. Pl.* 4, p. 972.

Brabeium stellatum. *Thunb. Prod.* 31.

HAB. In Africâ Australi, prope Prom. B. Spei. (v. s. in Herb. Banks., Lambert.)

22. GUEVINA.

Molin. Chil. 198. *Juss. Gen.* 424.

Quadria. *Gen. Flor. Peruv. et Chil.* 16, tab. 33. *Gært. Carp.* 3, p. 220, tab. 220.

CHAR. GEN. *Calyx* tetraphyllus, irregularis, foliolis tribus revolutis, quarto erecto. *Antheræ* apicibus concavis calycis immersæ. *Glandulæ* duæ hypogynæ, anticæ. *Ovarium* dispermium. *Stigma* obliquum. *Drupa* putamine osseo, monospermo.

HABITUS. Arbor. Folia alterna, pinnata. Racemi axillares, floribus geminis, pedicellatis, paribus unibracteatis. Calyx tomentosus, deciduus. *Drupa* parùm carnosa, nucleo amygdalino.

[156] *GUEVINA Avellana.* *Molin. Chil.* 198.*

Nebu subrotundo fraxini folio. *Feuill.* 3, p. 46, t. 33.

Quadria heterophylla. *Flor. Peruv. et Chil.* 1, p. 63, t. 99, f. b.

HAB. In sylvis et ad radices montium Chilensium. (v. s. in Herb. Banks. a Dombey.)

23. BELLENDENA.

CHAR. GEN. *Calyx* tetraphyllus, regularis, patens. *Stamina* hypogyna. *Glandulæ* nullæ hypogynæ. *Ovarium* dispermium. *Stigma* simplex. *Samara*? aptera, 1—2-sperma.

HABITUS. Frutex *glaberrimus*. *Folia sparsa, plana, apice trifida*. *Spica racemosa, terminalis; floribus sparsis, rarò geminatis*. *Calyx albus citò deciduus*. *Ovarium cum pedicello suo articulatum*. *Samara colorata margine altero sulcato*.

This genus is named in honour of JOHN BELLENDEN KER, Esq., whose botanical merits are established by an excellent Essay on *Ensatæ*, published in the 'Annals of Botany,' and by his elaborate disquisitions on the Genera of that and other monocotyledonous families, in the latter volumes of the 'Botanical Magazine.'

BELLENDENA *montana*.

HAB. In Insulâ Dicmen: in summis montibus. (ubi v. v.)

24. ANADENIA.

CHAR. GEN. *Calyx* tetraphyllus, apicibus concavis staminiferis. *Antheræ* immersæ. *Glandulæ* nullæ hypogynæ. *Ovarium* dispermium. *Stigma* conicum. *Folliculus* unilocularis, abortione monospermus. *Semen* apterum.

HABITUS. Frutices. (*Grevilleis affines* :) *pube dum adsit medio affixâ*. *Folia pinnatifida v. lobata, circumscriptione cuneiformia*. *Spicæ terminales, v. laterales, floribus geminatis, paribus unibracteatis, summis quandoque præcioribus!*

ETYM. α priv. et αδην glandula.

*1. *A. pulchella*, foliis pinnatifidis pilosiusculis: lobis cuneiformibus apice trifidis v. inciso-pinnatifidis, folliculis viscidis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*2. *A. trifida*, foliis cuneiformibus triplinerviis aveniis trifidis (unguicularibus) subtus argenteis: lobis integerimis lateralibusve 2—3-dentatis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in sylvis solo arenoso. (ubi v. v.)

Obs. Fortè generis distincti, ob calycem irregularem, stigma paullò diversum, et folliculum ligneum bipartibilem.

*3. *A. ilicifolia*, foliis cuneiformibus (uncialibus) venosis subtus argenteis basi attenuatis extra medium pinnatifido-incisis.

HAB. In Novæ Hollandiæ orâ australi; Flinders' Land: in areiosis prope littora. (ubi v. v. floribus inexpansis absque fructu.)

25. GREVILLEA.

CHAR. GEN. *Calyx* irregularis foliolis laciniisve secundis, apicibus cavis staminiferis. *Antheræ* immersæ. *Glandula* unica hypogyna, dimidiata. *Ovarium* dispermum. *Stigma* obliquum, depressum (rarò subverticale, conicum). *Folliculus* unilocularis, dispermus, loculo centrali. *Semina* marginata v. apice brevissimè alata.

HABITUS. Frutices rarò Arbores, *pube dum adsit medio affixâ*. Folia alterna, indivisa v. pinnatifida. Spicæ modò elongatæ racemosæ, modò abbreviatæ corymbosæ v. fasciculiformes, involucro nullo, pedicellis geminatis, rarò pluribus fasciculisve unibracteatis. Calyces sæpissimè rubicundi, nunc flavi; in quibusdam obliquè inserti. Folliculi vel coriacei, ovati, stylo toto coronati; seminibus ovalibus angus-

tissimè marginatis et apice brevissimè alatis: vel lignei, subrotundi, pseudo-bivalves basi tantum styli mucronati; seminibus undique alatis.

This extensive genus, of which a few of the least remarkable species have been already published as Embotriums by Dr. Smith, Cavanilles, and others, I have dedicated to the Right Honourable CHARLES FRANCIS GREVILLE, one of the Vice-Presidents of the Royal Society; a gentleman eminently distinguished for his acquirements in natural history, and to whom the botanists of this country are indebted for the introduction and successful cultivation of many rare and interesting plants.

Grevillea is probably the most extensive genus of Proteaceæ in New Holland, and admits of division into several very natural sections, most of which are readily distinguishable by more than one character, existing either in the parts of fructification or in habit; notwithstanding which, I have not ventured to separate them into distinct genera, as I probably should have done, had I been acquainted with fewer species; but have given to each section a proper name, a practice that may perhaps be advantageously adopted in all large genera where they are thus capable of natural subdivision. It must be unnecessary to add that proper names can in this manner be given only where the sections are perfectly natural, and not in those cases where genera have been subdivided from single characters, and those too of but little importance, as in Thunberg's division of Protea, from the form and division of the leaves; to which may be opposed the masterly ^[169] subdivision of the same genus previously given by Linnæus in the Mantissa, whose sections, though apparently depending on single characters, are evidently formed from a contemplation of the whole structure, as far as it was then understood; and it is remarkable that, with the exception of the first species, with whose real structure he was necessarily unacquainted, the rest are arranged, and even divided into sections, in most cases corresponding with the genera proposed in the present essay.

† Folliculi coriacei, stylo toto stigmatæque depresso coronati. Semina ovalia, angustissimè marginata, apicæque brevissimè alata.

A. LYSSOSTYLIS.

Folia omnia integerrima (in plerisque marginibus refractis v. replicatis pseudo-3-nervia). Flores fusciculati v. in racemo abbreviato. Stylus glaber. Folliculus ecostatus.

1. *G. punicea*, foliis elliptico-oblongis basi subattenuatis marginibus refractis, ramulis floriferis racemoque abbreviato recurvis, pistillis uncialibus, barbâ interiore calycis oblongâ dimidium inferiorem unguium æquante.

Embothrium sericeum β . *Smith, New Holl.* 27, t. 9, f. 5, β .

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis subhumidis. (ubi v. v.)

*2. *G. dubia*, foliis ellipticis marginibus refractis, ramis ramulisque tomentosis, floriferis racemoque abbreviato recurvis, pistillis unciâ brevioribus.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in saxosis subhumidis prope littora. (ubi v. v.)

OBS. Nimis affinis præcedenti.

170] 3. *G. sericea*, foliis ellipticis oblongisve obtusis mucronatis marginibus refractis, ramulis floriferis erectis, racemis abbreviatis recurvis, pistillis semuncialibus, barbâ interiori calycis dimidio inferiore unguium brevior.

Embothrium sericeum. *Smith, New Holl.* 25, t. 9, f. 1, 2, 3, 4. *Willd. Sp. Pl.* 1, p. 539. *And. Repos.* 100. *Sims, in Bot. Mag.* 862.

Embothrium cytisoides. *Cav. Ic.* 4, p. 60, t. 386, f. 2.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in saxosis prope littora marina et ad rivulorum ripas. (ubi v. v.)

4. *G. linearis*, foliis lineari-lanceolatis acutis mucronatis marginibus refractis, racemis abbreviatis erectiusculis, stylis apice glaberrimis.

Embothrium linearifolium. *Cavan. Ic.* 4, *p.* 59, *t.* 386, *f.* 1.

Embothrium lineare. *And. Repos.* 272.

Embothrium sericeum γ. *Smith, New Holl.* 27, *t.* 9, *f.* 6.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in saxosis præsertim prope littora. (ubi v. v.)

*5. *G. stricta*, foliis lanceolato-linearibus acutis mucronatis marginibus refractis costâque denticulato-scabris, stylis apice sericeis.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: ad ripas saxosas fluviorum. (ubi v. v.)

*6. *G. riparia*, foliis elongato-linearibus marginibus refractis costâque lævibus, stylis apice glaberrimis, pistillis quadrilinearibus: pedicello ovarium superante, barbâ interiori calycis densâ.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: ad ripas fluviorum. (ubi v. v.)

*7. *G. parviflora*, foliis subulato-linearibus marginibus refractis costâque lævibus, ramulis glabriusculis, calycibus ferrugineis barbâ interiori obsoletâ, pistillis bilinearibus: pedicello ovarium vix æquante.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in fruticetis a littore remotis. (ubi v. v.)

*8. *G. juniperina*, foliis subulatis fasciculatis divaricatis marginibus refractis, ramulis villosis teretiusculis, pistillis semuncialibus pedunculo partiali quadruplò longioribus.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis rariùs. *D. G. Caley, & A. Gordon.* (v. s.)

*9. *G. australis*, foliis lanceolato-subulatis unciâ brevioribus margine subrecurvis, suprâ pube deciduâ conspersis subtùs sericeis, ramis ramulisque tomentosius teretibus.

HAB. In Insulæ Diemen plagis australioribus: ad fluviorum ripas. (v. v. absque flor. v. fruct.)

*10. *G. tenuifolia*, foliis subulatis margine revolutis unciâ brevioribus, fasciculis sessilibus, pistillis bilinearibus.

HAB. In Insulâ Diemen; prope Port Dalrymple: ad ripas saxosas fluviorum. (ubi v. v.)

*11. *G. pauciflora*, foliis lineari-oblongis planiusculis obtusis mucronulatis suprâ lævibus subtus subsericeis: inferioribus glabriusculis, fasciculis 2-4-floris erectis, calycibus nudiusculis pistillum subæquantibus.

HAB. In Novæ Hollandiæ orâ australi; Flinders' Land: in depressis apricis prope littora. (ubi v. v.)

*12. *G. aspera*, foliis lineari-oblongis obtusis mucronulatis suprâ punctato-asperis subtus argenteis, racemis abbreviatis recurvis, stylis brevissimis, stigmate cochleariformi.

HAB. In Novæ Hollandiæ orâ australi; Flinders' Land: in ericetis aridis. (v. v. flor. delaps. fruct. matur.)

*13. *G. concinna*, foliis linearibus margine revolutis lævibus erectis, racemis recurvis secundis multifloris, ovariis lanatis, stylis glaberrimis calyce subsericeo duplò longioribus.

OBS. A reliquis sectionis facie differt.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in sterilibus prope littora marina. (ubi v. v.)

† B. PTYCHOCARPA.

Folia omnia integerrima. Flores fasciculati v. in racemo abbreviato, floribus superioribus præcocioribus! Stylus hirsutus v. tomentosus. Ovarium subsessile. Folliculus costatus!

*14. *G. arenaria*, foliis oblongis obtusis mucronulatis, racemis recurvis paucifloris, pistillis tomentosis.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: ad ripas arenosas fluviorum. (ubi v. v.)

*15. *G. montana*, foliis lanceolatis acutis suprâ læviusculis subtus sericeis, floribus geminatis, pedunculis glabris

calyces nudiusculos subæquantibus, pistillis hirsutis, tomento ramulorum arcuè adpresso.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in montosis. (v. s.)

*16. *G. acuminata*, foliis lanceolatis subacuminatis mucronatis suprà punctato-scabris subtùs cinereo-tomentosis, racemis paucifloris porrectis recurvisve, pistillis hirsutis, calycibus demùm glabriusculis, ramulis pubescentibus.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in montosis. (v. s.)

17. *G. cinerea*, foliis ellipticis obovatisve mucronatis suprà scabriusculis subtùs cinereo-tomentosis, racemis paucifloris recurvis, pistillis hirsutis, calycibus pedunculisque lanatis.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in montosis ad ripas saxosas fluviorum. (ubi v. v.)

*18. *G. mucronulata*, foliis obovatis obtusis mucronulatis suprà scabris nitentibus subtùs parùm sericeis, racemis abbreviatis, pistillis hirsutis, calycibus pilosiusculis pube adpressâ.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis. (ubi v. v.)

*19. *G. Baueri*, foliis oblongis obtusis mucronulatis utrinque glabris lævibus, racemis abbreviatis, pistillis hirsutis, calycibus pedunculisque glaberrimis.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in depressis a littore remotis. (ubi v. v.)

† C. ERIOSTYLIS.

Folia omnia integerrima. Flores fasciculati, subumbellati. Pistillum lanatum, pedicellatum. Folliculus ecosatus.

*20. *G. occidentalis*, foliis lanceolatis suprà punctatis scabris subtùs sericeis, fasciculis axillaribus terminalibusque, calycibus utrinque stylisque lanâ patulâ cinereis, stigmate mutico.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in sylvis solo sterili. (ubi v. v.)

*21. *G. sphacelata*, foliis oblongis lanceolatisve suprâ punctis minutis scabriusculis subtus sericis, fasciculis terminalibus, calycibus extus ferrugineo-tomentosis intus stylisque cinereo lanatis, stigmatibus mutico.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in saxosis. (ubi v. v.)

*22. *G. phyllicoides*, foliis lineari-lanceolatis: suprâ punctato-scabris superioribus villosis; subtus pubescentibus cinereis, stigmatibus ovalibus appendice duplò longioribus.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in montibus saxosis. (ubi v. v.)

23. *G. buxifolia*, foliis ellipticis suprâ punctatis scabris subtus tomento adpresso cinereis, stigmatibus orbiculatis appendicem curvam vix æquantibus.

Emboth. . . buxifolium. *Smith, New Holl.* 29, t. 10. *Willd. Sp. Pl.* 1, p. 538. *And. Repos.* 218.

Embothrium genianthum. *Cav. Ic.* 4, p. 60, t. 387.

HAB. In Novæ Hollandiæ orâ australi; prope Port Jackson: in ericetis saxosis. (ubi v. v.)

† D. PLAGIOPODA.

Folia integerrima v. divisa. Racemus thyrsiformis. Pedicellus ovarii accretus apici obliquo pedunculi, cui utrinque foliola duo calycis unum supra alterum inserta!

*24. *G. Goodii*, foliis integerrimis oblongis undulatis ^{176]}venosis utrinque glabris, racemis elongatis pedunculatis, caulibus prostratis.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria, et Arnhem's Land: in depressis arenosis, prope littora. (ubi v. v.)

*25. *G. venusta*, foliis pinnatifidis v. trifidis passimque

indivisis subtùs sericeis, racemis erectis, calycibus glaberrimis, stylis hirsutissimis.

HAB. In Novæ Hollandiæ orâ orientali; prope Cape Townsend: in umbrosis, ad radices montium. (ubi v. v.)

† E. CALOTHYRSUS. (GREVILLIA strictè sic dicta.)

Racemus *thyrsiformis*. Folia *pinnatifida* (raro passim indivisa).

*26. *G. pungens*, foliis pinnatifidis suprâ glabris subtùs argenteis: laciniis subulato-linearibus mucronatis pungentibus, racemis refractis, calycibus pistillisque glaberrimis.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria: prope littora. (ubi v. v.)

*27. *G. Dryandri*, foliis pinnatis subtùs sericeis: foliolis elongato-linearibus, racemis pedunculatis porrectis longissimis, calycibus insertione subobliquis pistillisque glaberrimis, caule patulo.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria, Arnhem's Land: prope littora. (ubi v. v.)

*28. *G. aspleniifolia*, foliis elongatis linearibus pinnatifido-incisis integerrimisque subtùs tomentosis, racemis folio ter brevioribus, calycibus pubescentibus, stylis glabris.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: rariùs. (v. s. in Herb. Banks.)

*29. *G. Banksii*, foliis pinnatifidis subtùs sericeis: laciniis elongato-lanceolatis, racemis erectis æqualibus, calycibus tomentosis, stylis glabris, ovariis sessilibus.

HAB. In Novæ Hollandiæ orâ orientali; Keppel Bay, Pine Port, &c.: in collibus saxosis. (ubi v. v.)

*30. *G. Chrysodendrum*, foliis pinnatifidis bipinnatifidisque: laciniis angusto-linearibus elongatis, racemis cylindraceis: floribus semiverticillatis, calycibus tomentosis basi persistenti! ovariis subsessilibus, stylis glabris.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria: prope littora. (ubi v. v.)

†† CYCLOPTERA.

Folliculi *lignei, subrotundi, basi styli mucronati. Semina undique alata.*

*31. *G. heliosperma*, foliis pinnatis subbipinnatisque glabris: pinnis oblongo-linearibus v. oblongis: inferioribus petiolatis, racemis divisis erectis, calycibus pistillisque glaberrimis.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria: prope littora. (ubi v. v.)

*32. *G. refracta*, foliis pinnatis passim indivisis: foliolis elongato-linearibus subtùs argenteis, racemis refractis divisis, calycibus sericeis, pistillis glaberrimis.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria: prope littora. (ubi v. v.)

177 *33. *G. ceratophylla*, foliis 2-3-fidis indivisisque subtùs nervosis sericeis: laciniis elongato-linearibus, folliculis glaberrimis ovalibus.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria: prope littora. (ubi v. v. sine flor.)

*34. *G. mimosoides*, foliis integerrimis ensiformibus planis nervosis ramisque glabris, folliculis obovatis viscidis.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria: prope littora. (ubi v. v. sine flor.)

*35. *G. polystachya*, foliis lineari-ensiformibus integerrimis laxis subtùs nervosis sericeis, racemis terminalibus alternis, pistillis semunciâ longioribus, stigmatibus obliquo concavo papillâ centrali.

HAB. In Novæ Hollandiæ orâ orientali, intra tropicum. (ubi v. v.)

*36. *G. striata*, foliis lineari-ensiformibus integerrimis strictis subtùs multinervibus sericeis, racemis terminalibus

alternis, pistillis semunciâ brevioribus, stigmate verticali depresso-conico.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria: prope littora. (ubi v. v. sine fructu.)

*37. *G. lorea*, foliis teretibus! pendulis longissimis, stigmate truncato-pyramidato.

HAB. In Novæ Hollandiæ orâ orientali, prope littora; Shoalwater Bay. (ubi v. v. sine fructu.)

*38. *G. gibbosa*, foliis elongato-lanceolatis integerrimis pubescentulis uninerviis venosis, racemis elongatis, ¹¹⁷⁸ stigmate conico, folliculis gibboso-incrassatis.

HAB. In Novæ Hollandiæ orâ orientali, intra tropicum; prope Endeavour River. *J. Banks, Bart.* (v. s.)

26. HAKEA.

Schrad. Sert. Hanov. Cavan. Ic. 6. Labill. Nov. Holl. 1, p. 30. Pers. Syn. 117. Conchium. Smith, Linn. Trans. iv, p. 215. Vent. Malmais. 110. Gært. Carp. 3, p. 216.

CHAR. GEN. *Calyx* tetraphyllus, irregularis, foliolis secundis. *Stamina* apicibus concavis calycis immersa. *Glandula* hypogyna unica, dimidiata (rard biloba). *Ovarium* pedicellatum, dispermum. *Stigma* subobliquum, c basi dilatata conico-mucronatum. *Folliculus* unilocularis, ligneus, loculo excentrico, pseudobivalvis. *Semina* alâ apicis nucleo longiore.

HABITUS. Frutices rigidi, quandoque Arbores mediocres; pube dum adsit medio affixâ. Folia sparsa, in variis varia, nunc in eodem frutice diversiformia. Fasciculi v. Racemuli, sæpius axillares, in plerisque involucrati, squamis imbricatis, scariosis, caducis, rudimenta ramulorum quandoque simul includentibus, ideoque potius pro gemmâ habendis, sed genus, unicâ exceptâ specie, a confinibus, optimè distinguuntibus, aliis notis in quibusdam vacillantibus. Pedicelli colorati, in

racemosis geminati, paribus unibracteatis. Flores parvi, albi v. ochroleuci. Pistillum glaberrimum, stylo subdeciduo. Capsula parietibus incrassatis. Semina nigra, rarò cinerea.

† *Folia omnia filiformia.*

Δ. *Capsulæ juxta apicem ecalcaratæ.*

1. *H. pugioniformis*, foliis filiformibus indivisis glabris, calycibus sericeis hirsutisque, capsulis lanceolatis acuminatis rectis utrinque infra medium transversim cristatis.

¹⁷⁹¹ a. Calyces sericei.

Banksia teretifolia. Salisb. Prod. 51.

Hakea glabra. Schrad. Sert. Hanov. 27, t. 17.

Hakea pugioniformis. Cavan. Anal. de Hist. Natur. 1, p. 213. Ic. 6, p. 24,* tab. 533.*

*Conchium pugioniforme. Smith, Linn. Trans. 9, p. 122.**

*Conchium longifolium. Smith, Linn. Trans. 9, p. 121.**

Lambertia teretifolia. Gært. Carp. 3, p. 213, t. 217.

β. Calyces hirsuti. Ramuli ultimi tomentosi.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in cricetis aridis, collibusque saxosis. β fortè distincta species. In Insulâ Diemen. (ubi v. v.)

*2. *H. rugosa*, foliis filiformibus indivisis glabris fructu parùm longioribus, capsulis obovatis curvatis refractis utrinque cristatis rugosis; acumine subulato lævi adscendenti, caule diffuso.

HAB. In Novæ Hollandiæ orâ australi; Flinders' Land: in campis sterilibus prope littora. (ubi v. v. absque flor.)

3. *H. epiglottis*, foliis filiformibus indivisis glabris fructu duplò longioribus, capsulis curvatis refractis utrinque rugosis ecristatis: mucrone adscendenti subulato carinato, seminum alâ obovatâ, caule erecto.

Hakea epiglottis. Labill. Nov. Holl. 1, p. 30, tab. 40.

Conchium teretifolium. Gært. Carp. 3, p. 217, t. 219.

HAB. In Insulâ Diemen; ad fluviorum rivulorumque ripas. (ubi v. v.)

*4. *H. nodosa*, foliis filiformibus indivisis compressiusculis, capsulis gibbosis obtusis nodosis seminumque ^[180] alâ obovatis, calycibus glabris, pedunculis pubescentibus.

HAB. In Novæ Hollandiæ orâ australi, prope Port Phillip; ad latera montium. (ubi v. v.)

Obs. Sequenti nimis affinis, an species distincta?

*5. *H. flexilis*, foliis filiformibus indivisis parùm compressis, capsulis ellipticis acutiusculis modicè convexis lævibus.

HAB. In Novæ Hollandiæ orâ australi, prope Port Phillip; ad latera montium. (ubi v. v.)

*6. *H. leucoptera*, foliis teretibus indivisis fructu duplò longioribus, ramis erectis virgatis subflexuosis, capsulis ovatis infrâ gibbosis suprâ compressis, seminibus albinereis!

HAB. In Novæ Hollandiæ orâ australi; Flinders' Land: ad margines sylvarum prope radices montium. (ubi v. v. sine flor.)

*7. *H. obliqua*, foliis teretibus indivisis, ramis tomentosis, glandulâ hypogynâ adnatâ apici obliquo pedunculi, calycibus sericeis, capsulis gibbosis subnodosis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis aridis. (ubi v. v.)

*8. *H. sulcata*, foliis filiformibus indivisis undique sulcatis divaricatis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis aridis. (ubi v. v. sine flor. caps. immat.)

† B. *Capsulæ juxta apicem bicalcaratæ.*

*9. *H. lissosperma*, foliis filiformibus indivisis undique exsulcatis glabris fructu duplò longioribus, capsulis gibbosis intus lævibus: calcaribus brevissimis, seminis alâ obovatâ: nucleo lævi basi immarginato.

HAB. In Insulæ Diemen montibus australioribus; inter fluvia Derwent et Huon. (ubi v. v. sine flor.)

10. *H. gibbosa*, foliis filiformibus indivisis subtùs basi obsoletissimè sulcatis ramisque subpubescentibus, ramulis pedunculisque hirsutis, calycibus glabriusculis, capsulis gibbosis intùs lacunosis, seminis alâ semiellipticâ, nucleo lacunoso basi marginato.

Banksia gibbosa. *Smith in White's Voy.* 224, t. 22, f. 2. *Willd. Sp. Pl.* 1, p. 536.

Banksia pinifolia. *Salisb. Prod.* 51.

Hakea pubescens. *Schrad. Sert. Hanov.* 27.

Hakea gibbosa. *Cavan. Anal. de Hist. Nat.* 1, p. 214.*
lc. 6, p. 24,* t. 534.

Conchium gibbosum. *Smith in Linn. Trans.* 9, p. 119.*

Conchium sphæroideum. *Smith in Linn. Trans.* 9, p. 120 P*

Conchium cornutum. *Gært. Carp.* 3, p. 216, t. 219.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis. (ubi v. v.)

Obs. Calyces non penitus glabri, sed pilis paucis longiusculis decumbentibus, sæpiùs deciduis, conspersi.

11. *H. acicularis*, foliis filiformibus indivisis glabris subtùs infra medium obsoletè sulcatis longitudine fructùs, ramulis ultimis subsericeis, pedunculis hirsutis calyces glaberrimos subæquantibus, capsulis gibbosis subrugosis intùs lacunosis.

Banksia tenuifolia. *Salisb. Prod.* 51.

Hakea sericea. *Schrad. Sert. Hanov.* 27.

¹⁸²⁷ *Conchium aciculare*. *Vent. Malm. t.* 111. *Smith in Linn. Trans.* 9, p. 121.

β. *Conchium compressum*. *Smith in Linn. Trans.* 9, p. 121.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis saxosis. (ubi v. v.)

*12. *H. vittata*, foliis filiformibus indivisis exsulcis glabris fructu duplò longioribus, capsulis ovatis convexi-

usculis æquilateralibus basi citiùs dehiscentibus intùs lacunosis, seminis alâ obovatâ, ramulis tomentosus.

HAB. In Novæ Hollandiæ orâ australi; Flinders' Land : in campis sterilibus, prope littora. (ubi v. v. sine flor.)

*13. *H. cycloptera*, foliis filiformibus indivisis fructu duplò longioribus ramulisque glaberrimis, capsulis gibbosis intùs lacunosis, seminibus utrinque alatis; alâ inferiore nucleum subæquante!

HAB. In Novæ Hollandiæ orâ australi; Flinders' Land : in campis prope littora. (ubi v. v. sine flor.)

*14. *H. suaveolens*, foliis filiformibus pinnatifidis passimque indivisis suprâ sulcatis, floribus racemosis glabris : rachi tomentosâ, capsulis gibbosis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land : in saxosis prope littora. (ubi v. v.)

†† *Folia pleraque filiformia, aliqua plana.*

*15. *H. microcarpa*, foliis integerrimis glabris : rameis teretibus; infimis planis, calycibus pedunculisque glaberrimis, capsulis bicalcaratis umbellatis folio multoties brevioribus.

HAB. In Insulâ Diemen; ad ripas saxosas fluviorum. (ubi v. v.)

16. *H. trifurcata*, foliis filiformibus 2-3-fidis indivisisve subtùs sulcatis : passim planis ovalibus integerrimis, calycibus hirsutis, capsulis compressis ecalcaratis.

Conchium trifurcatum. Smith in Linn. Trans. 9, p. 122.*

HAB. In Novæ Hollandiæ orâ australi; Lewins Land : in campis sterilibus. (ubi v. v.)

*17. *H. varia*, foliis superioribus filiformibus divisissimicibusque : inferioribus planis pinnatifidis laciniis linearibus subulatisve, capsulis bicalcaratis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in campis sterilibus. (ubi v. v. sine flor.)

††† *Folia omnia plana.*

A. *Folia aliqua v. omnia dentata v. incisa.*

*18. *H. attenuata*, foliis cuneatis apice dentatis pinnatifidisve: passim lanceolatis integerrimis basi attenuatis, capsulis bicalcaratis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in collibus saxosis. (ubi v. v. sine flor.)

*19. *H. linearis*, foliis lanceolato-linearibus spinuloso-paucidentatis integerrimisque aveniis impunctatis, ramulis pedunculoque communi glabris, fasciculis terminalibus axillaribusque, capsulis bicalcaratis compressiusculis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in campis sterilibus. (ubi v. v.)

*20. *H. florida*, foliis angusto-lanceolatis spinuloso-dentatis, minutissimè punctatis marginibus scabriusculis, ramulis pedunculoque communi brevissimo pubescentibus, capsulis bicalcaratis convexiusculis.

184] HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
ad latera collium. (ubi v. v.)

*21. *H. ilicifolia*, foliis circumscriptione ovalibus opacis sinuato-dentatis spinulosis subpetiolatis, ramis tomentosis, capsulis bicalcaratis ovatis gibbosis apice compressis intùs scrobiculatis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in collibus sterilibus. (ubi v. v.)

*22. *H. nitida*, foliis lanceolatis oblongisve basi attenuatis spinuloso-paucidentatis integrisque nitidis subvenosis ramulisque glaberrimis, capsulis bicalcaratis gibbosiusculis intùs læviusculis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in collibus saxosis. (ubi v. v. absque flor.)

*23. *H. amplexicaulis*, foliis sinuato-dentatis nitidis subvenosis: basi dilatatâ cordatâ amplexicauli, caule prostrato, ramis glabris, capsulis ecalcaratis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus sterilibus. (ubi v. v. sine flor.)

*24. *H. prostrata*, foliis angulato-dentatis apice dilatatis cuneatis: basi cordatâ amplexicauli, caule prostrato, ramis pubescentibus, capsulis ecalcaratis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus sterilibus. (ubi v. v.)

25. *H. ceratophylla*, foliis pinnatifidis bipinnatifidisve linearibus planis, calycibus ferrugineo-tomentosis, capsulis ecalcaratis.

Conchium ceratophyllum. *Smith, Linn. Trans.* 9, [185 p. 124.*

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in campis collibusque. (ubi v. v.)

*26. *H. undulata*, foliis obovatis trinerviis reticulato-venosis undulatis spinoso-dentatis, capsulis ecalcaratis tumidis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v. sine flor.)

†††B. *Folia omnia integerrima.*

27. *H. oleifolia*, foliis lanceolatis integerrimis uninerviis obsoletè venosis mucronulo spinoso: superioribus pubescentibus, ramulis tomentosis, capsulis terminalibus bicalcaratis gibbosis.

Conchium oleifolium. *Smith, Linn. Trans.* 9, p. 124.*

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in campis et collibus. (ubi v. v.)

28. *H. saligna*, foliis elongato-lanceolatis integerrimis uninerviis acutis apiculo sphacelato; omnibus ramulisque glaberrimis, capsulis axillaribus gibbosis: apice compresso utrinque carinato.

Embothrium salignum. *And. Repos. t.* 215.

Conchium salignum. *Smith, Linn. Trans.* 9, p. 124.*

Conchium salicifolium. *Gært. Carp.* 3, p. 217, t. 219.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis elevationibus. (ubi v. v. sine flor.)

*29. *H. marginata*, foliis lanceolatis integerrimis marginatis uninerviis (unciâ brevioribus) mucrone spinoso: summis pubescentibus, capsulis ecalcaratis acuminatis nitidis subsessilibus.

¹⁸⁰³ HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis elevationibus. (ubi v. v. sine flor.)

30. *H. ruscifolia*, foliis ellipticis obovatisve petiolatis integerrimis spinoso-cuspidatis suprâ punctato-scabris subtus tomentosis, ramulis hirsutis, capsulis ecalcaratis punctatis scabriusculis.

β. *Hakea ruscifolia.* *Labill. Nov. Holl.* 1, p. 30,* t. 39.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: ad latera collum. (ubi v. v. sine flor.)

*31. *H. cinerea*, foliis lineari-lanceolatis elongatis integerrimis trinerviis obsoletè venosis scabriusculis apiculo sphacelato, ramulis squamisque involucri tomentosis, capsulis lanceolatis acuminatis subcompressis ecalcaratis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in arenosis prope littora. (ubi v. v. sine flor.)

32. *H. dactyloides*, foliis integerrimis triplinerviis venosis obovato-oblongis v. lineari-lanceolatis aversis, ramulis angulatis, pedicellis pilosis, calycibus glabris, capsulis ecalcaratis: cortice verrucoso.

α. Folia obovato-oblonga, passim lanceolata, venis anastomozantibus.

Banksia dactyloides. *Gært. Sem.* 1, p. 221, t. 47, f. 2. *Lam. Illust. Gen.* 1, p. 242, n. 1279, t. 54, f. 3, a Gært. mutuât.

Banksia oleæfolia. *Salisb. Prod.* 54.

Hakea dactyloides. *Cavan. Anal. de Hist. Nat.* 1, p. 215. *Ic.* 6, p. 25, t. 535.

Conchium dactyloides. *Vent. Malm. t.* 110. *Smith, Linn. Trans.* 9, *p.* 123.

Conchium nervosum. *Gært. Carp.* 3, *p.* 217, *t.* 219. ^[187]

β. *Folia lineari-lanceolata, venis obsoletis.*

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: α. in saxosis prope littora. β. ad ripas fluviorum in regione montanâ. (ubi v. v.)

33. *H. elliptica*, foliis integerrimis quinquenerviis reticulato-venosis ellipticis ovalibusve muticis, pedicellis calycibusque glabris, capsulis ecalcaratis acutis gibbosis: cortice nitido.

Conchium ellipticum. *Smith, Linn. Trans.* 9, *p.* 123.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

34. *H. clavata*, foliis integerrimis lingulatis cartilagineo-carnosis mucronatis enerviis, floribus racemosis glabris, capsulis bicalcaratis.

Hakea clavata. *Labill. Nov. Holl.* 1, *p.* 31,* *t.* 41.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis prope littora. (ubi v. v.)

*35. *H. arborescens*, foliis integerrimis lingulatis linearibusve obsoletè nervosis muticis, involucris nullis! umbellis pedunculatis, pedicellis calycibusque tomentosis, capsulis ecalcaratis.

HAB. In Novæ Hollandiæ orâ septentrionali; Carpentaria: in apricis prope littora. (ubi v. v.)

Obs. Species unica tropica et gemmis floralibus nudis.

27. LAMBERTIA.

Smith, Linn. Trans. 4, *p.* 214. *Cavan. Ic.* 6, *p.* 31.

CHAR. GEN. *Calyx* tubulosus, quadrifidus, laciniis spiraliter revolutis. *Stamina* laciniis inserta. *Squamulæ* ^[188] hypogynæ 4, distinctæ v. in vaginulam connatæ. *Ovarium*

dispermum. *Stigma* subulatum. *Folliculus* unilocularis, coriaceo-ligneus. *Semina* marginata. *Involucrum* 1—7-florum, imbricatum, deciduum. *Receptaculum* planum, epaleatum.

HABITUS. Frutices *pulcherrimi, ramis verticillatis*. Folia *terna, sæpius integerrima*. Involucra *terminalia, solitaria, colorata, in plerisque septemflora, rarò uniflora*. Folliculi *subcuneati, apice hinc cuspidati, inde bicornes v. mutici, quandoque echinati*.

*1. *L. uniflora*, involucris unifloris, foliis obovatis mucronatis glabris reticulatis, folliculis hinc cuspidatis inde ecornibus.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: prope littora saxosa sinuum. (ubi v. v.)

*2. *L. inermis*, involucris septemfloris: foliolis interioribus calycis dimidio brevioribus, stylis glabris, folliculis hinc cuspidatis inde ecornibus, foliis oblanceolatis obovatisque muticis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: ad latera saxosa collium. (ubi v. v.)

3. *L. formosa*, involucris septemfloris: foliolis interioribus calycem æquantibus, stylis pilosis, folliculis hinc cuspidatis inde bicornibus, foliis lineari-lanceolatis cuspidato-mucronatis margine revolutis.

Lambertia formosa. *Smith, Linn. Trans. tab. 20. And. Repos. 69. Cavan. Anal. de Hist. Nat. 1, p. 233.* Ic. 6, p. 31,* t. 547.*

Protea nectarina. *Wendl. Sert. Hanov. fasc. 4, p. 5, t. 21.*

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis saxosis. (v. v.)

189] *4. *L. ? echinata*, foliis linearibus glabris reticulatis apice dilatato-lobato! mucronato, folliculis bicornibus undique echinatis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
ad latera saxosa collium. (ubi v. v. absque flor.)

28. XYLOMELUM.

Smith, Linn. Trans. 4, p. 214.

CHAR. GEN. *Calyx* tetraphyllus, regularis, foliolis apice revolutis. *Stamina* inserta supra medium foliolorum, iisque recurvatis exserta. *Glandulæ* quatuor hypogynæ. *Ovarium* dispermum. *Stylus* deciduus. *Stigma* verticale, clavatum, obtusum. *Folliculus* incrassato-ligneus, unilocularis, loculo excentrico. *Semina* apice alata.

HABITUS. Arbor. Folia *opposita*, *adultæ integerrima*, *plantæ juvenilis dentata*. Spicæ *axillares*, *oppositæ*, *amentaceæ*, *florum paribus unibracteatis*, *infimis solùm perfectis*, *reliquis ovario destitutis stigmatæque minore abortientibus*. *Folliculus unicus tantum maturescens, obpyriformis, tomentosus, crassissimus, intus suturâ dehiscens, inde siccatione partibilis*.

XYLOMELUM *pyriforme*.

Banksia pyriformis. Gært. Sem. 1, p. 220, t. 47, f. 1, fructus. Lam. Illust. Gen. 1, p. 242, n. 1278, t. 54, f. 4, a Gært. mutuât. White, Voy. 224.

Hakea piriformis. Cavan. Anal. de Hist. Nat. 1, p. 217.* Ic. 6, p. 25,* t. 536.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in campis et collibus saxosis. (ubi v. v.)

29. ORITES.

CHAR. GEN. *Calyx* tetraphyllus, regularis, foliolis apice recurvis. *Stamina* inserta supra medium foliolorum, iisque recurvatis exserta. *Glandulæ* quatuor hypogynæ. *Ova-*^[190]
rium sessile, dispermum. Stylus strictus. Stigma ob-

tusum, verticale. *Folliculus* coriaceus, unilocularis, loculo subcentrali. *Semina* apice alata.

HABITUS. Frutices. Folia *alterna*, *integerrima* v. *dentata*. Spicæ *axillares* v. *terminales*, *breves*, *florum paribus unibracteatis omnibus hermaphroditis*.

ETYM. Οπισθης monticola. Hi Frutices enim in summis montibus crescunt.

*1. *O. diversifolia*, foliis planis lanceolatis dentatis integerrimisve subtùs tomentosiusculis, folliculis suture truncatâ leviterve excisâ.

HAB. In Insulæ Diemen summis montibus. (ubi v. v.)

*2. *O. revoluta*, foliis margine revolutis linearibus integerrimis subtùs incano-tomentosis, folliculis suture rotundatâ.

HAB. In Insulæ Diemen summis montibus. (ubi v. v. absque flor.)

30. RHOPALA.

Schreb. *Gen. Pl.* 144. Roupala. *Aubl. Guian.* 1, p. 83, t. 32. *Gært. Carp.* 3, p. 212, t. 217.

CHAR. GEN. *Calyx* tetraphyllus, regularis, foliolis apice recurvis. *Stamina* supra medium foliolorum inserta, iisque recurvatis exserta. *Squamulæ* hypogynæ quatuor, distinctæ v. connatæ. *Ovarium* dispermium. *Stylus* persistens. *Stigma* verticale, clavatum. *Folliculus* unilocularis, ligneo-coriaceus. *Semina* utrinque alata, marginata, nucleo centrali.

HABITUS. Arbores. Folia *alterna*, rarò *verticillata*, *simplicia integerrima* v. *dentata*, rariùs *pinnata* v. *ternata*, in eodem ramo. Spicæ *axillares*, quandoque *terminales*, ^[191] *racemosæ*, *floribus geminis paribus unibracteatis*.

1. *R. montana*, foliis alternis integerrimis ovatis complicatis breviter acuminatis reticulato-venosis racemo axillari

brevioribus, pedunculis cum calycibus ovariisque ferrugineo-tomentosis.

Roupala montana. Aublet, *Guian.* 1, p. 83, t. 32. *Lam. Illust. Gen.* 1, p. 243, t. 55. *Poiret, Encyc. Botan.* 6, p. 316.*

Rupala montana. Vahl, *Symb.* 3, p. 20. *Willd. Sp. Pl.* 1, p. 536. *Gært. Carp.* 3, p. 212, t. 217.

HAB. In Americæ æquinoctialis Guianâ Gallicâ. Aublet. (v. s. in Herb. Aubl., nunc in Mus. Banks.)

*2. *R. media*, foliis alternis integerrimis ovatis planis acuminatis petiolum decurrentibus immersè venulosis racemo axillari brevioribus, pedicellis calycibusque pubescentibus, ovariis tomentosis.

HAB. In Americæ æquinoctialis Guianâ Gallicâ. *Jul. V. Rohr.* (v. s. in Herb. Banks.)

3. *R. nitida*, foliis alternis integerrimis ellipticis breviter acuminatis planis racemum axillarem subæquantibus, pedicellis cum calycibus ovariisque glabris.

Ropala nitida. *Rudge, Guian.* 1, p. 26, t. 39.

HAB. In Americæ æquinoctialis Guianâ Gallicâ. *Jos. Martin.* (v. s. in Herb. Banks. et Lambert.)

*4. *R. moluccana*, foliis alternis integerrimis ellipticis planis venulosis subreticulatis spicâ longioribus, pedicellis calycibusque glabris.

HAB. In Insulis Moluccanis. *D. Christoph. Smith.* (v. s. in Herb. Banks.)

5. *R. cochinchinensis*, foliis alternis ovato-ellipticis breviter acuminatis planis extra medium subserratis racemum axillarem subæquantibus, pedicellis cum calycibus ovariisque glabris.

Helicia cochinchinensis. *Lour. Cochin.* 83,* fide specimenis ab auctore, in Herb. Banks.

HAB. In sylvis Cochinchinæ. *Loureiro, l. c.* (v. s. absque fructu.)

DESC. Rami glaberrimi, teretes. Folia petiolata, gla-

berrima, uninervia, 2-2½ uncias longa; quandoque integerrima. *Racemi* solitarii. *Calyx* ante expansionem clavatus, clavâ ovali tubi dimidio brevior et duplò crassior. *Antheræ* foliolis calycis spiraliter revolutis exsertæ. *Ovarium* brevissimè pedicellatum. *Stylus* filiformis, strictus. *Stigma* clavatum, striatum, oblongum, æquilaterale. *Squamulæ* quatuor hypogynæ, breves, ad medium connatæ, persistentes (a Loureiro post lapsum calycis visæ et uti calyculus quadrifidus descriptæ).

*6. *R. serrata*, foliis alternis latè ellipticis parùm acuminatis serratis racemo axillari longioribus: basi subattenuatâ integerrimâ; paginis discoloribus, pedicellis cum calycibus ovariisque tomentosis.

HAB. In Insulis Moluccanis. *D. Christoph. Smith.* (v. s. in Herb. Bank^e et Roxb.)

*7. *R. dentata*, foliis alternis ovato-lanceolatis complicatis dentatis utrinque attenuatis racemo axillari parùm brevioribus: acumine lineari, calycibus ovariisque tomentosis.

HAB. In Americæ æquinoctialis Guianâ Gallicâ. *D. Alex. Anderson.* (v. s. in Herb. Banks.)

8. *R. peruviana*, foliis alternis ovatis serratis lanuginosis subtùs ferrugineis racemo axillari brevioribus.

183] Embothrium monospermum. *Flor. Peruv. et Chil.* 1, p. 63,* t. 98.

HAB. In Peruviæ montibus frigidis; prope Panoa, vicum ad Portachuelo declivia. *Flor. Peruv. l. c.*

9. *R. diversifolia*, foliis alternis simplicibus pinnatisque venosissimis subtùs pubescentibus racemo axillari brevioribus, folliculis acinaciformibus tomentosis.

Embothrium pinnatum. *Fl. Peruv. et Chil.* 1, p. 63,* t. 99.

HAB. In Peruviâ; in Muna ruderatis et versuris. *Flor. Peruv. l. c.*

10. *R. sessilifolia*, foliis quaternis subsessilibus cuneato-oblongis subacuminatis integerrimis, racemis terminalibus verticillatis umbellatisve.

Roupala sessilifolia. *Rich. in Act. Soc. Hist. Nat.* 1, p. 106. *Poiret, Encyc. Botan.* 6, p. 316.* *Willd. Sp. Pl.* 1, p. 537.

Roupala hameliæfolia. *Rudge, Guian.* 1, p. 22,* t. 31.

HAB. In Americæ æquinoctialis Guianâ Gallicâ. (v. s. in Herb. Banks. et Lamb.)

31. KNIGHTIA.

CHAR. GEN. *Calyx* tetraphyllus, regularis, foliolis revolutis. *Stamina* calyci extra medium inserta. *Glandulæ* hypogynæ quatuor. *Ovarium* tetraspermum, sessile. *Stigma* verticale subclavatum. *Folliculus* coriaceus, stylogerus, unilocularis. *Semina* apice alata.

HABITUS. *Arbor excelsa*. *Folia sparsa, serrata*. *Racemi axillares, floribus geminatis, paribus unibracteatis*. *Folliculi oblongi, tomentosi*.

Genus proximum *Rhopalæ*, distinctum, Seminibus quaternis, apice solùm alatis.

This genus, which was discovered by Sir Joseph Banks, is, with his approbation, named in honour of his friend Thomas Andrew Knight, Esq., the author of many valuable essays on Vegetable Physiology, published in the Philosophical Transactions.

For the figure here given I am also indebted to the liberality of the illustrious President of the Royal Society, who has enabled me to complete the account of this remarkable plant, by permitting me to copy Dr. Solander's description, which I was the more desirous to give, as it exhibits a specimen of the accuracy with which subjects of natural history were investigated in that celebrated voyage; of whose important results it is to be lamented so little is known to foreign naturalists, though in this country they have ever been open to the public, and in the most advantageous manner.

KNIGHTIA excelsa. Tab. 36 (II).

HAB. In Novâ Zelandiâ; prope Tolaga et Oपुरagi. *Josephus Banks*, Baronetus. (v. s. folliculis vacuis sed impressionibus seminum insignitis.)

DESC. *Arbor* sylvestris, magna, sæpè 80 pedalis. *Caudex* strictissimus. *Rami* erecti, teretes, glabri. *Ramuli* ultimi parùm compressi, villosiusculi. *Coma* pyramidalis. *Folia* numerosa, densè sparsa, erecta, petiolata, lanceolato-oblonga (v. oblonga), acutiuscula, profundè serrata, serraturis remotis obtusis, plana, coriacea, rigida: supernè glabra, nitida, lævia, subtùs venosa venulisque numerosissimis subreticulata, villis copiosissimis brevissimis densissimis cinerascens: 4-5 uncias longa. *Petoli* foliis sexies breviores. *Racemi* sessiles, axillares, simplices, multiflori, ovato-oblongi, foliis duplò breviores, sæpè in ramis nudis collocati, ubi ante decessum foliorum axillares fuerunt, unde primo intuitu videntur quasi lateribus ramorum erumpentes. *Rachis* ruberrima. *Pedicelli* holosericei, ¹⁹⁶³ ruberrimi, crassiusculi, patentissimi, semunciales, bipartiti, unde biflori. *Calyx* tetraphyllus, foliola ante expansionem arcuè in tubum connata, linearia, acutiuscula, sesquiuncialia, coriacea, extùs villosa holosericea, ruberrima, usque ad basin revoluta, æqualia, in medio paulò angustiora. *Glandulæ* quatuor, receptaculo inter basin foliorum insertæ, e latâ basi acutæ, virescentes, apice rubicundæ, semilineam longæ. *Filamenta* quatuor, unguibus petalorum adnata, suprâ medium per spatium lineare libera, filiformia, plana, erecta, rubicunda. *Antheræ* lineares, longitudine foliolorum, supernè parùm incurvæ, flavæ, ipso apice casso lanceolato. *Ovarium* conicum, subangulatum, villosiusculum, rubrum. *Stylus* filiformis, crassiusculus, strictus, persistens, longitudine filamentorum, rubicundus. *Stigma* cylindraceo-angulatum, incrassatum, apice attenuatum, longitudine antherarum, virescens. *Folliculus* oblongo-lanceolatus, stylo persistenti coronatus, coriaceus, crassus, durus, unilocularis, sesquiuncialis v. paulò longior, extùs sericeus. *Hactenus* SOLANDER.

Obs. Pollen triangulare, angulis per lentem pellucen-
tioribus, flavum. *Ovarium* tetraspermum, ovulis apice
alatis.

32. EMBOTHRIMUM.

Embothrii species. *Forst. Gen.* 15, t. 8, litt. g. et seq.

CHAR. GEN. *Calyx* irregularis, hinc longitudinaliter fissus, inde quadrifidus. *Stamina* apicibus concavis calycis immersa. *Glandula* hypogyna unica, semiannularis. *Ovarium* pedicellatum, polyspermum. *Stylus* persistens. *Stigma* verticale, clavatum. *Folliculus* oblongus. *Semina* apice alata.

HABITUS. Frutices v. *Arbusculæ glabræ*. Ramuli squamis persistentibus gemmarum quandoque obsiti. Folia sparsa, integerrima. Racemi terminales, corymbosi, paribus pedicellorum unibracteatis: Involucro communi nullo. [196] Flores coccinei, glaberrimi.

1. *E. coccineum*, foliis ovali-oblongis obtusis mucronulatis: paginis discoloribus, ramulis squamatis.

Embothrium coccineum. *Forst. Gen.* p. 16, t. 8, litt. g. —m. *Linn. Suppl.* 128. *Forst. Com. Soc. Reg. Goett.* 9, p. 24. *Lam. Encyc. Botan.* 2, p. 351.* *Illust. Gen.* 1, p. 244, n. 1284, t. 55, f. 2. *Willd. Sp. Pl.* 1, p. 537.

HAB. In Americâ Australi ad littora freti Magellanici, et in Terra del Fuego. (v. s. in Herb. Banks.)

Obs. Pollen ellipticum, levissimè arcuatum, extremitate utrâque pellucentiore; fovillâ majusculâ globosâ.

2. *E. lanceolatum*, foliis lanceolato-linearibus, ramis esquamatis.†

Embothrium lanceolatum. *Flor. Peruv. et Chil.* 1, p. 62, t. 96.

HAB. In Chili collibus et montibus altis, inter Conceptionis urbem et Arauci arcem. *Flor. Peruv. l. c.*

33. OREOCALLIS.

Embothrii species. *Flor. Peruv. et Chil.*

CHAR. GEN. *Calyx* irregularis, hinc longitudinaliter

fissus, inde quadridentatus. *Stamina* apicibus concavis calycis immersa. *Glandula* nulla hypogyna. *Ovarium* pedicellatum, polyspermum. *Stigma* obliquum, orbiculato-dilatatum, concaviusculum. *Folliculus* cylindraceus. *Semina* apice alata. *Involucrum* (racemi) nullum.

HABITUS. Frutex *speciosus*. Folia *sparsa, integra, paginis discoloribus*. Racemus *thyrsoides, terminalis, paribus pedicellorum unibracteatis*. Flores *coccinei, glaberrimi*.

ETYM. Ορος mons, et καλος formosus.

197] *OREOCALLIS grandiflora*.†

Embothrium grandiflorum. *Lam. Encyc. Botan.* 2, p. 354.* *Illust. Gen.* 1, p. 244, n. 1283. *Willd. Sp. Pl.* 1, p. 538.

Embothrium emarginatum. *Flor. Peruv. et Chil.* p. 62, t. 95.

HAB. In Peruviae montibus; in collibus frigidis Provinciae Tarmæ. *Flor. Peruv.* l. c.

34. TELOPEA.

Embothrii species. Smith, Salisb.

CHAR. GEN. *Calyx* irregularis, hinc longitudinaliter fissus, inde quadrifidus. *Stamina* apicibus concavis calycis immersa. *Glandula* hypogyna unica, subannularis. *Ovarium* polyspermum, pedicellatum. *Stylus* persistens. *Stigma* obliquum, clavatum, convexum. *Folliculus* unilocularis, cylindraceus. *Semina* apice alata, alâ hinc immarginatâ inde vasculosâ nervo obliquè recurrenti. *Involucrum* (racemi v. corymbi) imbricatum, deciduum.

HABITUS. Frutices *ramis determinatis*. Folia *sparsa, dentata v. integra*. Racemi *terminales, corymbosi, paribus pedicellorum unibracteatis*. Flores *coccinei*.

ETYM. τηλωπος qui e longinquo cernitur, quod de his fruticibus, floribus coccineis speciosis, valet.

In this genus, as well as in *Lomatia*, and perhaps in all those with an indefinite number of seeds, an extremely

thin black-brown crust is interposed between the ripe seeds, exactly corresponding with them in size and form, and which is probably the remains of a fluid matter that had separated them in the unripe state.

The most important characters distinguishing this genus from *Lomatia*, seem to be the single semiannular or nearly circular gland, the cohering calyx, and the vascular wing of the seed; for the Involucrum, which at first seems to afford so excellent a distinction, considerably loses its importance in *Telopea truncata*, in which it almost always includes the rudiments of branches, as in *Hakea*. In natural affinity *Telopea* approaches much more nearly to *Oreocallis*, which differs principally in having no gland at the base of the footstalk of its ovary, and in the want of an Involucrum: the wing of the seed seems (from the figure in the *Flora Peruviana*) to be in like manner vascular. Embothrium itself, which is also very near akin to *Telopea*, is distinguishable by its vertical stigma, oval pollen, and naked corymbi.

1. *T. speciosissima*, foliis cuneato-oblongis inciso-dentatis venosis cum ramulis involucrisque glaberrimis.

Embothrium speciosissimum. *Smith, New Holl.* 19, t. 7. *Sims, Bot. Mag.* 1128.

Embothrium speciosum. *Salisb. Parad.* 111.

Embothrium spathulatum. *Cav. Ic.* 4, p. 60, t. 388. *Gært. Carp.* 3, p. 214, t. 218.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: locis saxosis, præsertim subumbrosis. (ubi v. v.)

2. *T. truncata*, foliis lanceolato-oblongis integerrimis passimque paucidentatis subtùs ramulisque pubescentulis, involucris extùs tomentosis.

Embothrium truncatum. *Labill. Nov. Holl.* 1, p. 32, t. 44.

OBS. Alam seminis in hâc apice semper rotundatam in præcedenti sæpiùs truncatam observavimus.

HAB. In Insulæ Diemen montibus australioribus. (ubi v. v.)

35. LOMATIA.

Embothrii species. *Smith, Cavan.*

CHAR. GEN. *Calyx* irregularis, foliolis distinctis secundis. *Stamina* apicibus concavis calycis immersa. *Glandulæ* hypogynæ tres, secundæ. *Ovarium* pedicellatum, polyspermum. *Stylus* persistens. *Stigma* obliquum, dilatatum, subrotundum, planiusculum. *Folliculus* ovali-oblongus. *Semina* apice alata; alâ marginatâ disco evasculoso.

HABITUS. Frutices. Folia alterna, in plerisque divisa, v. dentata, rarius integerrima, quandoque in eodem frutice varia. Racemi terminales, interdum axillares, elongati, laxi, nunc abbreviati, corymbosi, paribus pedicellorum unibracteatis. Flores ochroleuci. Involucrum nullum. Seminis nucleus farinâ sulphurâ conspersus.

ETYM. λωμα, margo, ob seminum alam marginatam.

1. *L. silaifolia*, foliis bipinnatifidis glaberrimis: pinnulis cuneato-linearibus lanceolatisve incisis acutis mucronatis reticulato-venosis, racemis glaberrimis elongatis divisis simplicibusve.

Embothrium silaifolium. *Smith, New Holl.* 23, t. 8. *Willd. Sp. Pl.* 1, p. 537.

Embothrium herbaceum. *Cav. Ic.* 4, p. 60, t. 388.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in campis et ericetis. (ubi v. v.)

2. *L. tinctoria*, foliis pinnatifidis bipinnatifidisve (raro indivisis) glabris: pinnulis linearibus distichis uninerviis subaveniis obtusiusculis mucronulatis, racemis elongatis glabris indivisis.

Embothrium tinctorium. *Labill. Nov. Holl.* 1, p. 31, tab. 42 et 43.

HAB. In Insulæ Diemen campis et collibus. (ubi v. v.)

200] 3. *L. ferruginea*, foliis bipinnatifidis tomentosis:

pinnulis ovatis lanceolatisve, racemo terminali foliis brevioribus.

Embothrium ferrugineum. *Cavan. Ic. 4, p. 50,* t. 385.*

HAB. In Americæ Australis "San Carlos de Chiloe in solo aquâ marinâ quandoque inundato." *Cavan. l. c.*

*4. *L. polymorpha*, foliis lineari-lanceolatis integerrimis v. dentatis pinnatifidisve subtus cum ramulis pedicellisque tomentosis, racemis terminalibus corymbosis, calycibus pilosiusculis, pistillis glaberrimis.

a. *cinerea*. Folia lineari-lanceolata integerrima, marginibus recurvis, subtus cinereo-tomentosa; folliculi semunciales.

β. *rufa*. Folia lanceolata v. lineari-lanceolata, incisa v. pinnatifida, passim integerrima, subtus ferrugineo-tomentosa; folliculi subunciales.

Embothrii tinctorii var. *Labill. Nov. Holl. l. c.*

HAB. In Insulæ Diemen montibus australioribus. (ubi v. v.)

*5. *L. ilicifolia*, foliis oblongo-ovatis acutis spinuloso-dentatis reticulatis petiolisque glaberrimis, racemis terminalibus elongatis.

HAB. In Novæ Hollandiæ orâ australi; prope Port Phillip: in campis sterilibus lateribusque montium: (ubi v. v. flor. delaps.)

*6. *L. longifolia*, foliis lineari-lanceolatis elongatis glabris remotè serratis, racemis axillaribus, pedunculis calycibusque pilosiusculis, pistillis glaberrimis.

Embothrium myricoides. *Gært. Carp. 3, p. 215, t. 218?*

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: ad ripas saxosas fluviorum et rivulorum. (ubi v. v.)

7. *L. dentata*, foliis ovalibus serrato-dentatis petiolisque glaberrimis, racemis lateralibus abbreviatis, calycibus pilosis, ovario tomentoso.†

Embothrium dentatum. *Flor. Peruv. et Chil.* 1, p. 62, t. 94 a.

HAB. In nemoribus et sylvis regni Chilensis. *Flor. Peruv. l. c.*

8. *L. obliqua*, foliis ovatis serratis glabris, racemis axillaribus, pedicellis calycibusque pilosis, stigmate deciduo.

Embothrium obliquum. *Flor. Peruv. et Chil.* 1, p. 63, t. 97.

Embothrium hirsutum. *Lam. Encyc. Botan.* 2, p. 355. *Illust. Gen.* 1, p. 245, n. 1286.

HAB. In Conceptionis Chili et Puchacay provinciarum montibus. *Flor. Peruv. l. c.*

Obs. Ala seminis hujus et præcedentis examinanda.

36. STENOCARPUS.

EMBOTHRII species. *Forst. Gen.*

CHAR. GEN. *Calyx* irregularis, foliolis distinctis, secundis. *Stamina* apicibus concavis foliolorum immersa. *Glandula* hypogyna unica, semiannularis. *Ovarium* pedicellatum, polyspermum. *Stylus* deciduus. *Stigma* obliquum, orbiculato-dilatatum, planiusculum. *Folliculus* linearis. *Semina* basi alata!

HABITUS. Frutices *glaberrimi*. Folia *alterna, integerrima*. Umbellæ *axillares, v. terminales, pedunculatæ*. Flores *ochroleuci*.

ETYM. στενος angustus, et καρπος fructus.

1. *S. Forsteri*, foliis oblongis obtusis enervibus:

Embothrium umbellatum. *Forst. Gen.* 16, t. 8, f. a.—*f. Forst. Aust. n.* 60. *Linn. Suppl.* 228. *Lam. Encyc. Botan.* 2, p. 352. *Illust. Gen.* 1, p. 245, n. 1285, t. 55, f. 1. *Willd. Sp. Pl.* 1, p. 538.

^{202]} HAB. In Novâ Caledoniâ. *J. R. et G. Forster.* (v. s. sine fructu in Herb. Banks. et Lambert.)

*2. *S. salignus*, foliis elongato-lanceolatis basi trinerviis.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: ad ripas saxosas fluviorum et rivulorum. (ubi v. v.)

37. BANKSIA.

Linn. fil. Suppl.

CHAR. GEN. *Calyx* quadripartitus (rarò quadrifidus). *Stamina* apicibus concavis laciniarum immersa. *Squamulæ* hypogynæ quatuor. *Ovarium* biloculare, loculis monospermis. *Folliculus* bilocularis, ligneus: *Dissepimento* libero, bifido. *Amentum* flosculorum paribus tribracteatis!

HABITUS. Frutices v. Arbores, *vix excelsæ*. Rami *umbellati*. Folia *sparsa*, rarò *verticillata*, *integra*, *serrata*, v. *pinnatifido-incisa*, in eodem stirpe quandoque varia; in plantâ juvenili v. *mutilatâ sæpè serrata*, v. *incisa*, dum in adultâ et *illæsâ integerrima*. Amenta *solitaria*, *terminalia* v. *e dichotomiis*, rarò *lateralia*, *bracteolis nonnullis*, *brevibus*, *angustis subtensa*, *cylindracea*, in quibusdam *abbreviata*. Bracteæ *flosculorum persistentes*, *maiores solitariae*; *minores geminatae*, *collaterales*, *interiores*. Amenti *fructiferi rachis ut plurimum incrassata*, et cum *folliculorum basibus conferruminata*. Semina *nigra*, *apice cuneato-alata*, *nucleo in lacunâ respondente dissepimenti lignei semiimmerso*.

*1. *B. pulchella*, foliis acerosis integerrimis muticis (unguicularibus), calycis unguibus lanatis: laminis glabris, stigmate depresso-capitato.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land; [203 in ericetis aridis prope littora. (ubi v. v.)

*2. *B. sphaerocarpa*, foliis acerosis integerrimis mucronulatis (uncialibus), calycis unguibus laminisque hirsutis, stigmate subulato, strobilis globosis, folliculis ventricosos apice compressiusculis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis depressis. (ubi v. v.)

*3. *B. nutans*, foliis acerosis integerrimis mucronulatis, amentis nutantibus, calycibus sericeis, folliculis apice dilatatis depressis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis aridis prope littora. (ubi v. v.)

4. *B. ericifolia*, foliis accrosis emarginato-bidentatis (unguicularibus): marginibus integerrimis, amentis elongatis, calycibus sericeis, stigmatē capitato.

Banksia ericæfolia. *Linn. Suppl.* 127. *Lam. Encyc. Botan.* 1, p. 369. *Illustr. Gen.* 1, p. 242, n. 1276. *Willd. Sp. Pl.* 1, p. 536. *And. Repos.* 156. *Cavan. Anal. de Hist. Nat.* 1, p. 221.* *Ic.* 6, p. 27, t. 538. *Pers. Synop.* 1, p. 117.

Banksia. *White's Voy. tab. ad p.* 225, fig. 1, strobilus.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis saxosis. (ubi v. v.)

5. *B. spinulosa*, foliis (adultis) acerosis (1—3-uncialibus) apice tridentatis dente intermedio longiore: marginibus spinuloso-dentatis integerrimisve, calycibus basi intus imberbibus, stigmatē subulato.

Banksia spinulosa. *Smith, New Holl.* 1, p. 13,* t. 4.
204] *Willd. Sp. Pl.* 1, p. 536. *Cavan. Anal. de Hist. Nat.* 1, p. 219.* *Ic.* 6, p. 26,* t. 537. *Pers. Synop.* 1, p. 17.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis aridis. (ubi v. v.)

Obs. Frutex est et sæpius humilis, nec Arbor decem-pedalis, &c. ut habet Cavanilles. l. c.

*6. *B. collina*, foliis linearibus spinuloso-dentatis denticulo terminali breviorē subtus venosis, bracteis amenti obtusis apice tomentosis, calycibus basi intus imberbibus, caule fruticoso.

HAB. In Novæ Hollandiæ orâ orientali; in collibus apricis prope littora. Hunter's River. (ubi v. v.)

*7. *B. occidentalis*, foliis linearibus extra medium spinuloso-dentatis subtus aveniis, bracteis amenti apice glabris,

calycibus marcescentibus: unguibus basi intùs barbatis, folliculis ventricosus tomentosus: apice compressiusculo nudo, caule fruticoso, ramulis glabris.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis. (ubi v. v.)

*8. *B. littoralis*, foliis elongato-linearibus spinuloso-dentatis basi attenuatis subtùs aveniis, calycibus deciduis, folliculis compressis bracteisque strobili apice tomentosus, caule arboreo, ramulis tomentosus.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: ad littora arenosa sinuum. (ubi v. v. flor. delaps.)

9. *B. marginata*, foliis linearibus truncatis mucronulatis integerrimis v. dentatis: venulis subtùs inconspicuis, ramis ultimis hirsutis, bracteis omnibus amenti apice glabris: (205 majoribus acutis, caule fruticoso.

a. Frutex erectus, orgyalis. Amentum foliis plerumque integris longius.

Banksia marginata. Cavan. *Anal. de Hist. Nat.* 1, p. 227. *Ic.* 6, p. 29,* t. 544.

β. Frutex erectus, orgyalis. Folia spinuloso-dentata, planiuscula, amento quandoque longiora.

Banksia microstachya. Cavan. *Anal. de Hist. Nat.* 1, p. 224. *Ic.* 6, p. 28,* t. 541, exclus. syn. Linnei.

γ. Frutex humilis, diffusus. Folia spinuloso-dentata, planiuscula, cuneata, amento longiora.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis. (ubi v. v.)

*10. *B. depressa*, foliis elongato-cuneatis truncatis mucronulatis spinuloso-dentatis: subtùs obsolete costatis venulis inconspicuis, bracteis omnibus amenti (folia vix æquantis) tomentosus obtusis, caule prostrato, ramulis ultimis hirsutis.

HAB. In Insulæ Diemen plagis australioribus; in saxosis ad radices montium. (ubi v. v.)

*11. *B. patula*, foliis cuneato-linearibus truncatis mucro-

nulatis integris v. paucidentatis (uncialibus) subtùs reticulato-venosis, bracteis amenti apice tomentosis obtusis, calycis laminis carinâ glabrâ, caule diffuso, ramulis ultimis tomentosis.

HAB. In Novæ Hollandiæ orâ australi; Flinders' Land: inter frutices, in sterilibus elevationibus. (ubi v. v.)

*12. *B. australis*, foliis linearibus truncatis mucronulatis margine recurvis integris subtùs reticulato-venosis, ramulis ultimis tomentosis, bracteis amenti obtusis subæqualibus apice tomentosis, calycis laminis carinâ obsoletissimâ sericeâ, caule arboreo.

HAB. In Insulâ Diemen, ubique in campis et prope littora, necnon in orâ australi Novæ Hollandiæ prope Port Phillip. (ubi v. v.)

*13. *B. insularis*, foliis lineari- v. cuneato-oblongis subrotundatis cum mucronulo sparsis verticillatisve subtùs reticulato-venosis, bracteis amenti obtusis extrorsum tomentosis, folliculis compressis apice glabris.

HAB. In Insulis Freti Bass, et in Insulâ Diemen, prope littora. (ubi v. v.)

14. *B. integrifolia*, foliis verticillatis oblongo-lanceolatis integris mucronulatis: subtùs venulis reticulantibus conspicuis, folliculis tomentosis, caule arboreo.

a. Arbor parva v. mediocris. Folia oblanceolata, sæpiùs acuta, basi attenuata. Bracteæ geminatæ obtusæ, solitariis acutis dimidio minores.

Banksia integrifolia. Linn. *Suppl.* 127. *Lam. Encyc. Botan.* 1, p. 369. *Illust. Gen.* 1, p. 242, n. 1275. *Willd. Sp. Pl.* 1, p. 535. *Cavan. Anal. de Hist. Nat.* 1, p. 229. *Ic.* 6, p. 30, tab. 546.

Banksia spicata. *Gært. Sem.* 1, p. 221, t. 48.

Banksia oleæfolia. *Cavan. Anal. de Hist. Nat.* 1, p. 228. *Ic.* 6, p. 30, t. 545.

Banksia glauca. *Cavan. Anal. de Hist. Nat.* 1, p. 230. *Ic.* 6, p. 31.*

β. Arbor magna. Folia lanceolato-oblonga, sæpiùs ob-

tusiuscula, basi acuta. Bracteæ geminatae obtusæ, solitariis acutiusculis haud dimidio minores.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: juxta littora marina. β . In orâ australi, prope Port Phillip. (v. v.)

ONS. Species polymorpha, cui nimis affines sunt *B. insularis* et *compar*.

*15. *B. compar*, foliis sparsis lingulato-oblongis emarginatis muticis dentatis integrisve: subtus reticulato-venosis niveis, ramulis bracteisque tomentosis, calycibus sericeis, caule arboreo.

HAB. In Novæ Hollandiæ orâ orientali; prope Keppel Bay: juxta littora. (ubi v. v. absque fructu.)

ONS. Præcedenti proxima; an distincta species?

*16. *B. verticillata*, foliis verticillatis lingulato-oblongis obtusis muticis: subtus aveniis niveis, bracteis amentitomentosis obtusis; involucrantibus hirsutis, caule arboreo.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: prope littora. (ubi v. v.)

*17. *B. coccinea*, foliis alternis cuneato-obovatis oblongisve dentatis truncatis costatis reticulato-venosis basi transversis, bracteis subulatis calycibusque lanatis, stigmatē pyramidali.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in campis prope littora. (ubi v. v.)

*18. *B. paludosa*, foliis subverticillatis cuneato-oblongis subtruncatis basi attenuatis extra medium dentato-serratis margine subrecurvis: subtus costatis reticulato-venosis, petiolis ramulisque glabris, calycibus sericeis, caule fruticoso.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in paludosis. (ubi v. v.)

19. *B. oblongifolia*, foliis sparsis angusto-oblongis truncatis dentato-serratis basi acutiusculis: subtus costatis

reticulato-venosis, petiolis ramulisque tomentosis, bracteis majoribus amenti acuminatis, calycibus sericeis, caule fruticoso.

Banksia oblongifolia. *Cavan. Anal. de Hist. Nat.* 1, p. 225.* *Ic.* 6, p. 28,* t. 542.

Banksia asplenifolia. *Salisb. Prod.* 51?

Banksia salicifolia. *Cavan. Anal. de Hist. Nat.* 1, p. 231. *Ic.* 6, p. 31,*? folia enim in hâc specie quandoque integra.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in ericetis. (ubi v. v.)

20. *B. latifolia*, foliis obovato-oblongis spinuloso-serratis basi acutis: subtùs costatis reticulatis cinereo-tomentosis, calycis unguibus sericeis: laminis glabris, caule fruticoso.

Banksia robur. *Cavan. Anal. de Hist. Nat.* 1, p. 226.* *Ic.* 6, p. 29,* t. 543.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in paludosis. (ubi v. v.)

Obs. Hujus speciei nomen Cavanillesii mutare coactus sum, quoniam nunquam arborescit sed frutex humilis est.

21. *B. marcescens*, foliis cuneiformibus planis sparsis truncatis extra medium dentato-serratis: basi acutiusculâ, ramis tomentosis, calycibus persistentibus folliculisque glabris.

Banksia præmorsa. *And. Repos.* 258.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: prope littora. (ubi v. v.)

Obs. Cùm folia minimè præmorsa falsum nomen mutare non hesitavi.

200] *22. *B. attenuata*, foliis elongato-linearibus truncatis basi attenuatis extra medium serratis: subtùs costatis reticulatis areolis tomentosis, bracteis apice hirsutis, calycibus glabris, folliculis tomentosis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: prope littora. (ubi v. v.)

*23. *B. elatior*, foliis elongato-linearibus subtruncatis serratis subtùs reticulatis: adultis glabriusculis, bracteis imberbibus calycibusque tomentosis, stylo glaberrimo, stigmathe ovali-clavato, caule arboreo.

HAB. In Novæ Hollandiæ orâ orientali; prope Sandy Cape: prope littora. (ubi v. v.)

24. *B. serrata*, foliis lato-linearibus elongatis truncatis serratis: subtùs reticulatis glabriusculis: basi attenuatâ, stylo imo pulvereo-pubescenti, stigmathe cylindraceo sulcato: basi oblique incrassatâ, caule arboreo.

Banksia serrata. *Linn. Suppl.* 126.* *Lam. Encyc. Botan.* 1, p. 369. *Illust. Gen.* 1, p. 242, t. 54, f. 1. *White's Voy.* 222, cum tab. 2 prioribus. *Willd. Sp. Pl.* 1, p. 535. *And. Repos.* 82.

Banksia conchifera. *Gært. Sem.* 1, p. 221, t. 48, f. 1.

Banksia serrata. *Cavan. Anal. de Hist. Nat.* 1, p. 222. *Ic.* 6, p. 27, t. 540. (forsan ad sequentem pertinet.)

Banksia dentata. *Wend. Hort. Herrenh. tab.* 8. ? vel ad sequentem pertinens.

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in campis prope littora. (v. v.)

*25. *B. æmula*, foliis lato-linearibus elongatis truncatis profundè serratis: subtùs reticulatis glabriusculis, calycibus sericeis, stigmathe capitato exsulco nitido apice (quadragulo) styli duplò crassiore, caule fruticoso.

Banksia serratifolia. *Salisb. Prod.* 51. ?

Banksia serrata. *White's Voy.* 222, tab. tertia ?

HAB. In Novæ Hollandiæ orâ orientali; prope Port Jackson: in campis arenosis ericetisque. (ubi v. v.)

OBS. *B. serrata* *Cavan.* et *dentata* *Wend.* suprâ ad *B. serratam* citatæ, fortè ad hanc, valdè affinem, pertinent.

26. *B. dentata*, foliis cuneato-oblongis truncatis sinuato-dentatis undulatis basi acutis: subtùs costatis venulosis niveis, calycibus sericeis, folliculis tomentosis.

Banksia dentata. *Linn. Suppl.* 127. *Willd. Sp. Pl.* 1, p. 536.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

15 28 25
32 22
20
9

10

HAB. In Novæ Hollandiæ orâ orientali, prope Endeavour River; et in septentrionali, Arnheims Land: prope littora. (ubi v. v.)

*27. *B. quercifolia*, foliis oblongo-cuneatis subtruncatis glabris serrato-incisis: incisuris mucronatis, calycis laminis aristatis! folliculis glabriusculis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in campis prope littora. (ubi v. v.)

*28. *B. speciosa*, foliis linearibus pinnatifidis: lobis triangulari-semiovatis mucronatis subtus niveis obsolete nervosis, calycis laminis lanatis, stylo pubescenti, folliculis tomentosis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in saxosis prope littora. (ubi v. v.)

29. *B. grandis*, foliis pinnatifidis: lobis triangulari-ovatis acutis planis subtus nervosis glabriusculis, calycis laminis folliculisque glabris.

an Banksia grandis. Willd. Sp. Pl. 1, p. 535.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

30. *B. repens*, foliis pinnatifidis: lobis sinuatis v. dentatis, caule prostrato.

Banksia repens. Labill. Voy. 1, p. 412, t. 23. Nov. Holl. 2, p. 118.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in campis collibusque saxosis. (ubi v. v.)

*31. *B. ilicifolia*, foliis cuneatis inciso-serratis subtus glabriusculis, amentis brevissimis, calycis unguibus diu cohærentibus stylum æquantibus: laminis citius dehiscentibus!

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in campis collibusque prope littora. (ubi v. v.)

Obs. Species tam singularis ut ferè proprii generis, transitum ad Dryandras facilem reddit.

38. DRYANDRA.

CHAR. GEN. *Calyx* quadripartitus v. quadrifidus. *Stamina* apicibus concavis laciniarum immersa. *Squamulæ* hypogynæ quatuor. *Ovarium* biloculare, loculis monospermis. *Folliculus* bilocularis, ligneus: *Dissepimento* libero, bifido. *Receptaculum commune* planum, floribus indeterminatim confertis, paleis angustis, raro nullis. *Involucrum commune* imbricatum.

HABITUS. Frutices *plerumque humiles*. Rami dum adsint sparsi vel umbellati. Folia sparsa, pinnatifida v. incisa, plantæ juvenilis conformia. Involucra solitaria, terminalia, raro lateralibus sessilibus, foliis confertis interioribus quandoque nanis obvallata, hemisphærica, bracteis adpressis, in quibusdam apice appendiculatis.

Obs. Dryandra of Thunberg, first published in Flora Japonica, being not generically different from Aleurites, which was previously established by Forster, I have peculiar satisfaction in giving the name of my respected friend, Mr. DRYANDER, to a genus so nearly related to Banksia, from which indeed it differs chiefly in Inflorescence, but in that respect so widely as to be at once distinguishable: there is also something in the habit, especially in the leaves of the greater number of species, by which, independent of the parts of fructification, the genus is pretty certainly indicated; and it is worthy of notice, that, while Banksia is generally spread over all the coasts of New Holland and of Van Diemen's Island, Dryandra has hitherto been observed only on that part of the south coast called Lewins Land, where, however, its species are nearly as numerous and abundant as those of Banksia itself.

*1. *D. floribunda*, foliis cuneiformibus inciso-serratis, involucri bracteis exterioribus glabriusculis, calycis laminis glabris, stigmate subclavato obtuso.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in collibus saxosis. (v. v.)

Variat receptaculo epaleato.

*2. *D. cuneata*, foliis cuneiformibus sinuato-dentatis spinosis petiolatis, involucri bracteis omnibus sericeis, calycis laminis barbatis, stigmate subulato-filiformi acuto.

α. Folia vix sesquiunciam longa, dentibus terminalibus subequalibus.

β. Folia biuncialia, apicis dilatati denticulo medio brevioribus sinibus latioribus. Forsan species distincta.

²¹³ HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in collibus saxosis. (ubi v. v.)

*3. *D. armata*, foliis pinnatifidis : lobis triangularibus planis divaricatis rectis spinoso-mucronatis : terminali proximis longiore ; subtus reticulatis venulis nudis, ramis calycisque laminis glabris, stylo basi pubescenti, stigmate subulato sulcato.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in collibus saxosis. (ubi v. v.)

*4. *D. falcata*, foliis pinnatifidis : lobis subulato-triangularibus divaricatis falcato-recurvis spinoso-mucronatis : terminali proximis brevioribus ; subtus reticulatis venulis nudis, ramis pubescentibus, laminis calycis styloque longitudinaliter glabris, stigmate clavato exsulco.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in collibus saxosis. (ubi v. v.)

*5. *D. formosa*, foliis elongato-linearibus pinnatifidis : lobis scaleno-triangularibus muticis planis subtus niveis, involucri tomentosis : foliolis interioribus lineari-oblongis, receptaculo paleaceo. *Tab. 37 (III).*

HAB. In Novæ Hollandiæ orâ australi; Lewins Land :
in sterilibus prope littora. (ubi v. v.)

*6. *D. mucronulata*, foliis elongato-linearibus pinnatifidis : lobis isoscelo-triangularibus mucronulatis planis subtus

niveis, involucris tomentosis : foliolis interioribus linearibus mucronatis, receptaculo paleaceo, caule subsimplici.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land : [214] in depressis saxosis. (ubi v. v.)

*7. *D. plumosa*, foliis elongato-linearibus pinnatifidis : lobis isoscelo-triangularibus mucronulatis margine subrecurvis subtùs niveis, involucris foliolis interioribus plumoso-aristatis, receptaculo epaleato.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land : in lateribus saxosis collium. (ubi v. v.)

*8. *D. obtusa*, foliis linearibus pinnatifidis caule decumbenti tomentoso longioribus : lobis triangularibus obtusis subtùs niveis margine incrassato-recurvis, involucris bracteis exterioribus ovatis : interioribus lineari-oblongis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land : in apricis prope littora. (ubi v. v.)

9. *D. nivea*, foliis linearibus pinnatifidis caulem glabrum subæquantibus : lobis scaleno-triangularibus acutis mucronulatis subtùs niveis margine recurvis, involucris bracteis lineari-lanceolatis glabris ciliatis, calyce quadrifido, unguibus laminisque hirsutis.

a. Folia lobis adscendentibus, mucronatis, subtùs venosis. Stigma stylo parùm crassius.

Banksia nivea. *Labill. Voy.* 1, p. 413, t. 24. *Nov. Holl.* 2, p. 118.

β. Folia lobis divaricatis, uninerviis, subaveniis. Stigma stylo vix crassius.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land : in saxosis prope littora. (ubi v. v.)

*10. *D. longifolia*, foliis linearibus pinnatifidis longis- [215] simis acutis subtùs cinereo-tomentosis : basi attenuatâ integerrimâ ; lobis triangularibus adscendentibus decurrentibus margine recurvis, involucris bracteis elongato-linearibus margine barbatis extùs glabris, calycis unguibus basi lanatis supra pubescentibus : laminis pilosiusculis, caule tomentoso.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in collibus saxosis. (ubi v. v.)

*11. *D. tenuifolia*, foliis elongato-linearibus pinnatifidis subtruncatis subtùs niveis: basi attenuatâ integerrimâ petioliformi; lobis triangularibus decurrentibus divaricatis margine recurvis, involucri bracteis tomentosis: exterioribus ovato-lanceolatis, calycis unguibus basi lanatis suprâ cauleque glabris.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: in ericetis. (ubi v. v.)

*12. *D. pteridifolia*, foliis pinnatifidis caule tomentoso longioribus: lobis linearibus acutis mucronatis margine revolutis basi dilatatis, involucri bracteis tomentosis ovatis.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: ad latera saxose collium. (ubi v. v.)

*13. *D. blechnifolia*, foliis pinnatifidis caule tomentoso longioribus: lobis linearibus obtusis mucronulatis trinerviis margine recurvis basi simplicibus.

HAB. In Novæ Hollandiæ orâ australi; Lewins Land: prope King George's Sound. *D. Archibald Menzies.* (v. s. absque fructificatione.)

Obs. Ad hoc genus retuli, ob summam affinitatem cum *Dryandra pteridifoliâ*, cujus vix varietas.

216] TO RENDER this essay as complete as I am able, I proceed to notice such plants, as either belong or have been referred to Proteaceæ, but from my imperfect acquaintance with which, or from the unsatisfactory accounts hitherto given of them, could not with certainty be referred to any of the genera described, or, if referable to any of them, I could not with confidence propose as distinct species; and shall conclude with the addition of a few synonyms to the species described, from Ray's *Historia Plantarum*, which

had escaped me when the paper was first read to the Society.

LEUCADENDRON linifolium, foliis lineari-spathulatis aversis basi attenuatis ramisque glabris, capitulo masculo sessili foliis circumvallantibus longiore, calycis tubo barbato: laminis stylisque imberbibus.

Protea linifolia. *Jacq. Hort. Schænb.* 1, p. 11, t. 26.

Obs. There can be no doubt of the genus of this plant, or of the individual figured by Jacquin being a male. From the same figure, by which alone I am acquainted with it, it seems to be very nearly related to *Leucadendron tortum*, from which it differs in having the male heads sessile, and in the laminæ of the calyx being quite smooth.

LEUCADENDRON fusciflorum, foliis lineari-lanceolatis glabris junioribus rectis basi attenuatis, capitulo femineo foliis circumvallantibus brevioribus, calycis laminis plumoso-barbatis: tubo pilosiusculo.

Protea fusciflora. *Jacq. Hort. Schænb.* 1, p. 11, t. 27.

This also is known to me only from Jacquin's figure, ^[217] from which it is unquestionably a *Leucadendron*, and a female plant; it can hardly however be supposed the female of the preceding species; and though I have constructed a specific character for it, I think it is not improbably a variety of *Leucadendron angustatum*.

LEUCADENDRON.

Protea linearis. *Houtt. Nat. Hist. par.* 2, vol. 4, p. 116, t. 19, f. 2, ed. *Germ.* vol. 3, p. 84, t. 19.

This is undoubtedly a *Leucadendron*, and probably a female plant; but from the figure alone its species cannot be determined.

LEUCADENDRON.

Protea stellaris. *Sims, Bot. Mag.* 881.

Seems to be a male plant, and apparently different from anything I have seen. From the form of the leaves and the length of those surrounding the capitulum, I am

inclined to consider it as the male of Jacquin's *Protea fusci-flora* already noticed.

LEUCADENDRON.

Conocarpodendron; folio tenuissimo, angustissimo, saligno; cono calyculato. *Boerh. Lugd. Bat. 2, p. 203, c. tab.*

This is probably a male plant, notwithstanding the figure of a ripe cone is given at the bottom of the plate; the separate fruits of some of Boerhaave's figures belonging decidedly to very different species. It may be the male of *Leucadendron adscendens*.

LEUCADENDRON ?

Protea odorata. Thunb. Prod. Append. 187.

²¹⁸ There is no means of determining the genus of this plant, but it is rather more probably a *Leucadendron* than belonging to any other.

LEUCADENDRON ? ?

Conocarpodendron; acaulon; folio rigido, nervoso, oblongo, latiori; cono fusco; semine oblongo, in medio quasi excavato. *Boerh. Lugd. Bat. 2, p. 201, c. tab.*

I know not what to make of this. If the strobilus and nucs at the bottom of the plate really belong to it, it must be referred to *Leucadendron*, and will stand near *L. retusum* or *L. plumosum*; but there are some circumstances both in the figure and description which render this very doubtful. Thunberg refers it to his *Protea strobilina*, but the descriptions by no means agree.

LEUCADENDRON ?

Scolymocephalus Oleæ folio. Sherard, in Raj. Hist. 3, Dendr. p. 10.

This, according to Boerhaave, is his *Conocarpodendron*, &c. 2, p. 197, c. tab. which I have considered as the female of *Leucadendron squarrosus*.

LEUCADENDRON ? ?

Protea glabra. Thunb. Diss. n. 52.

From the very short and unsatisfactory description of Thunberg, the genus of this plant cannot be determined, or even with much probability guessed at.

ISOPOGON.

Protea divaricata. *And. Repos.* 465.

Can this be a variety of *Isopogon anemonifolius*? The yellow flowers satisfy me that it is not a species of ⁽²¹⁹⁾ *Serruria*, and prevent me at the same time from referring it to *Isopogon anethifolius*, whose leaves are not unlike, but whose flowers are of a very different colour.

PROTEA.

Protea venosa. *Lam. Illust. Gen.* 1, p. 234, n. 1212. *Poiret, Encyc. Botan.* 5, p. 640.

Said by Poiret to resemble in most respects *Protea longiflora*; it must therefore be a genuine *Protea*.

PROTEA.

Scolymodendros Africanus ex Monte Tabulari. *Pluk. Mant.* 168, t. 442, f. 4.

This is manifestly a *Protea*, which it appears Plukenet had seen only in the possession of Woodward. The head, especially in the form of the bractæ, bears a great resemblance to that of *Protea cynaroides*; but the leaves are so very different, that, unless we suppose they were drawn from memory and disproportionately reduced, it cannot be referred to this species. It is probably however one of the more common kinds, and I know not what else to suppose it may be, except *Protea grandiflora*. The figure itself has never, so far as I know, been noticed by any author.

LEUCOSPERMUM.

Scolymocephalus Africanus, foliis in summitate profundius crenatis, intercrenatis majoribus, florum staminulis longis recurvis. *Raj. Hist.* 3, *Dendr.* p. 10.

This is probably a *Leucospermum*, and perhaps *L. ellipticum*.

220] MIMETES ?

Protea dichotoma. *Lam. Illust. Gen.* 1, p. 235, n. 1219.
Poiret, Encyc. Botan. 5, p. 643.

Probably a spurious *Mimetes*.

SERRURIA Bergii, capitulis simplicibus solitariis subpedunculatis, bracteis cuneiformibus truncatis cum acumine villosis: inferioribus glabris, calycibus curvatis sericeis, stigmate turbinato-capitato, ramulis foliisque glabris.

Leucadendron sphærocephalum. *Berg. Cap.* 26.*

This I have no hesitation in referring to *Serruria*; and from the description of the accurate *Bergius* I am disposed to think it distinct from any that I am acquainted with. It seems most nearly related to *Serruria acrocarpa*, differing chiefly in the smoothness of its branches, and in having terminal heads.

SERRURIA.

Protea sphærocephala. *Houtt. Nat. Hist. par.* 2, vol. 4, p. 99, t. 19, f. 1, ed. *Germ.* vol. 3, p. 72, t. 19.

Unquestionably a *Serruria*, and probably referable either to *S. hirsuta* or *pedunculata*.

SERRURIA.

Protea villosa. *Thunb. Prod. Append.* 186.

A *Serruria* whose characters cannot be made out from the specific difference given by *Thunberg*.

SERRURIA.

Protea triternata. *And. Repos.* 337.

This may be intended for *S. congesta*, but I cannot with confidence refer to it as such.

SERRURIÆ.

221] *Protea abrotanifolia minor*. *And. Repos.* 536.

Protea abrotanifolia hirta. *And. Repos.* 522.

Protea abrotanifolia odorata. *And. Repos.* 545.

These are manifestly *Serruriæ*, but I do not venture to refer them to any of the species I have described; nor are

there sufficient materials from which they may be characterized as distinct species.

NIVENIA.

Protea concava. *Lam. Illustr. Gen.* 1, p. 234, n. 1217.
Poiret, Encyc. Botan. 5, p. 642.

A species of *Nivenia*, and perhaps one of those described.

NIVENIA ?

Protea candicans. *Thunb. Prod. Append.* 186.

Probably a *Nivenia*, and perhaps not different from *N. mollissima*: it may however be a species of *Serruria*, in which case it is probably *S. candicans*.

PROTEA prostrata. *Thunb. Prod.* 27.

I know not to what genus this may belong; but from the species near which Thunberg has placed it, it may be supposed to be either a *Protea* or a *Leucadendron*: if the latter, it is probably not very different from *L. retusum*.

HAKEA.

Conchium drupaceum. *Gært. Carp.* 23, p. 217, t. 219.

I cannot refer this fruit to any of the species I have described.

EMBOTHRIUM chaparro. *Humb. Æquin. Bot.*

Of this I know nothing but the name, which occurs in Humboldt's Chart of Æquinoctial Botany, and is placed there at the height of about 1600 feet.

EMBOTHRIUM strobilinum. *Labill. Nov. Holl.* 2, p. 116, t. 265.

The seeds of this remarkable plant, which I am acquainted with only from Labillardiere's figure and description, being unknown, and the internal structure of its ovary not having been examined, its genus cannot be determined. Its regular and deeply divided calyx, the four glands at the base of the ovary, and its vertical equilateral stigma, point out its near affinity to *Knightia*, from which it differs

in the style being deciduous, and perhaps also in the number and form of its seeds. If these are but two in number, it would be still more nearly related to *Orites*; but something in its whole appearance, and especially its uncommonly large bractæ, indicates its being a distinct genus.

According to Labillardiere, it is a native both of New Caledonia and the south-west coast of New Holland: but as I am acquainted with no plant of the order, which has so wide a range as this, and as it may be presumed the specimens from New Holland were very imperfect, otherwise so remarkable a plant would surely have found a place in the body of his work, I may be permitted to question the accuracy of the statement. I confess however that I know no plant of Lewins Land with which this could be confounded.

ROUPALA pinnata. *Lam. Illustr. Gen.* 1, p. 243, n. 1282. *Poiret, Encyc. Botan.* 6, p. 317. *Rudge, Pl. Guian.* 25, t. 38.

There can be little doubt of this plant constituting a distinct genus; but its fruit being entirely unknown, it is better to place it among those which require a further examination. It was referred to *Rhopala* at a time when that genus was not at all understood. In its compound ²²³⁷ leaves, its irregular calyx, and even in some degree in the glands subtending the ovarium, it seems to approach more nearly to *Gevuina*; and I am therefore inclined to think its fruit will be found to be a drupa, and not a folliculus as that of *Rhopala*. The whole plant however is so remarkable, that I here add a description taken from an excellent specimen, in Mr. Lambert's Herbarium, collected by the unfortunate Martin in Guiana, where it seems to have been first found by Richard.

Frutex? v. *Arbor.* *Ramuli* teretes, tomento minuto cinerascens. *Folia* alterna, abruptè pinnata, 3-4-juga. *Foliola* opposita, petiolata, latè ovata, obtusa quandoque acutiuscula, integerrima, glaberrima, super nitida, subter ferè opaca, venulis anastomozantibus parùm emersis reticulata: dum $3\frac{1}{2}$ uncias longa 2 uncias lata. *Petioli* partiales

semunciales, semiteretes, cum rachi teretiusculâ articulati. *Spica* terminalis, pedunculata, erecta, folio brevior, pedunculo longior, racemosa: *Pedunculo* rachique teretibus, pube brevissimâ cinereo-ferrugineis (in sicco). *Pedicelli* geminati, teretes, calyce breviores. *Calyx* tetraphyllus. *Foliola* ante expansionem in tubum curvatum cylindraceum clausum utrinque ampliaturum coherentia, mox ad basin distincta, decidua, linearia, extûs pube tenuissimâ arcuè appressâ (in sicco) cinereo-ferrugineâ; intûs glabra: *Unguib* linearibus, basi dilatatis: *Laminis* ovatis, acutis, concavis. *Stamina* 4. *Filamenta* brevissima, basi laminarum imposita. *Antherarum* lobi (connectivo) adnati, distincti, basi parium divergentes, longitudinaliter dehiscentes. *Pollen* flavum. *Ovarium* brevè pedicellatum, parvum, uniloculare, dispernum, ovulis collateralibus: *Pedicello* basi cincto *Squamâ* latâ, glabrâ, adnatâ, (in sicco) corrugatâ, posticè subdeficiente, intersticiâ angustissimâ. *Stylus* cylindraceus, crassiusculus, glaber, longitudine unguum calycis. *Stigma* obliquum, convexum, stylo crassius, papillâ centrali.

Obs. Singularis, Foliis verè compositis, petiolellis cum rachis articulatis; et *Squamâ* hypogynâ pedicello ovarii adnatâ, nec ipso receptaculo connexâ.

ORITINA acicularis. *Append. Flor. Nov. Holl. ined.*

This is a perfectly smooth erect shrub; with alternate cylindrical leaves, furrowed on the upper surface and terminated by a pungent mucro. I observed it only on the summit of the Table Mountain, at the southern extremity of Van Dienten's Island. The perfect flowers I have not seen, but have examined the ovary soon after fecundation, that I have no doubt of its containing originally only two ovula; and as its base is surrounded by four glands, the calyx is probably regular. Hence its near affinity to *Orites*, with which it also agrees in inflorescence and apparently in stigma. The fruit is a smooth compressed coriaceous follicle, containing two seeds, which are winged at both ends; on which account I have not absolutely referred it to *Orites*, but, until its flowers are discovered, have given it a temporary name, indicating its affinity to that genus.

BANKSIA musculiformis. Gært. Sem. 1, p. 221. Lam. *Illust. Gen.* 1, p. 242, n. 1280.

Fructus musculiformis. Rumph. Amb. 2, p. 184, t. 60.

Gærtner has taken up this plant entirely from Rumpf's figure, and referred it to *Banksia* on account of its fruit containing according to that author two winged seeds. But from Rumpf's description, it appears that the whole plant is lactescent; hence it probably does not belong to this family, but rather to Apocineæ, as Burmannus has already conjectured.

CYLINDRIA. Lour. *Cochin. ed. Willd.* 1, p. 86.

^{225]} Both Willdenow and Ventenat have considered this genus as belonging to Proteaceæ, with whose structure indeed the description of Loureiro in most respects well agrees. Mr. König, however, (*Ann. of Bot.* 1, p. 392,) assures us, on the authority of original specimens, that it is scarcely different from *Olea*, though Loureiro has characterized it as having four bilocular antheræ, included in the concave apices of the segments of the corolla; two circumstances altogether incompatible with Oleinæ, and which render it not improbable that the specimen sent to Sir Joseph Banks by the author was very different from that which he described.

LEUCOSPERMUM *Conocarpum.*

Scolymocephalus Africanus, latifolius, lanuginosus, foliis in summitate crenatis, comâ sericeâ. *Raj. Hist.* 3, *Dend.* p. 9.

MIMETES *Hartogii.*

Scolymocephalus Africanus lanuginosus humilis, foliis in summo tridentatis, flore dilutè purpureo, carinulâ albulâ *Oldenlandii.* *Raj. Hist.* 3, *Dend.* p. 10, fide characteris et descriptionis.

MIMETES *cucullatus.* (*Raj. Hist.* 3, *Dend.* p. 10, n. 10.)

MIMETES hirtus.

Scolymocephalos Africanus, foliis brevioribus acuminatis, floribus rubentibus, summis surculis foliis intermistis. *Raj. Hist. 3, Dend. p. 10.*

Besides the Proteaceæ described or noticed in this ⁽²²⁶⁾ paper, I am acquainted with several very beautiful species, chiefly of *Grevillea* and *Persoonia*, discovered in New Holland by Mr. *George Caley*, a most assiduous and accurate botanist, who, under the patronage of Sir JOSEPH BANKS, has for upwards of eight years been engaged in examining the plants of New South Wales, and whose numerous discoveries will, it is hoped, be soon given to the public, either by himself, or in such a manner as to obtain for him that reputation among botanists to which he is well entitled.

TABULARUM EXPLICATIO.

Tab. 36 (II). KNIGHTIA EXCELSA.

FIG.

1. Flos expansus, parùm auctus.
2. Idem longitudinaliter apertus, magnitudine naturali.
3. Ejusdem basis cum glandulis hypogynis.
4. Pistillum auctum, ovario longitudinaliter secto ovulis quatuor.
5. Ovulorum insertiones et relativas positiones ostendens.
6. Ovulum paulò magis auctum.
7. Pollen plurimùm auctum.

Tab. 37 (III). DRYANDRA FORMOSA.

1. Ramus magnitudine naturali.
2. Flos magnitudine naturali.
3. Idem auctus.
4. Receptaculum commune magnitudine naturali et auctum.
5. Idem verticaliter sectum.
6. Paleæ receptaculi.
7. Folliculus.
8. Dissepimentum cum seminibus.
9. Semina.
10. Dissepimentum.
11. Pollen ad lentem auctum.

ON THE

ASCLEPIADEÆ,

A NATURAL ORDER OF PLANTS SEPARATED
FROM THE APOCINEÆ OF JUSSIEU.

BY

ROBERT BROWN, LIB. LINN. SOC.

(READ 4TH NOV., 1809.)

[*Extracted from the 'Memoirs of the Wernerian Natural History Society,' Vol. I, pp. 12—78.*]

EDINBURGH.

1811.

ASCLEPIADEÆ.

THE vast additions to the number of species which botany is constantly receiving, while they make a natural arrangement absolutely necessary to the general botanist, render it at the same time proportionally difficult. For though there are still many tribes of plants easily distinguishable even by a superficial observer, yet there are others, that hitherto have been thought abundantly distinct, which can no longer be circumscribed by means of characters taken from their organs of reproduction. This is perhaps now the case with the RUBIACEÆ and APOCINEÆ of Jussieu. It is true, that to an experienced observer, it may still be practicable to refer the greater part, perhaps the whole, of these plants to their proper places in a natural series; ^[13] but it is, I apprehend, no longer so, to distinguish the two orders by definitions derived from the usual source. Such at least is the opinion I have been led to form from all that I have seen published respecting them, as well as from what I have lately had an opportunity of observing in New Holland.

As, however, both these families are already too extensive, it becomes expedient rather to attempt their subdivision into smaller groups, which may possibly admit of more accurate limitation, than to unite them into one vast order, the distinguishing characters of which, could they be obtained, must probably be extremely vague, and clogged with numerous exceptions. Such a subdivision, it seems to me, may be easily made of the Apocineæ, by employing

a character at once obvious and important, and which while it preserves the natural series unbroken, has the additional advantage of dividing the order into two nearly equal parts. To one of these which includes the genus *Apocynum*, the name of Apocineæ will of course remain.

The consideration of the other, which from one of its most remarkable genera I propose naming ASCLEPIADEÆ, forms the chief subject of the following essay; but the more completely to illustrate it, I have subjoined new, and I trust amended characters of the genera of the most nearly related section of the Apocineæ strictly so called. The singular structure of the stamina in the Asclepiadeæ has attracted the attention of botanists since the days of Tournefort: it is therefore not a little remarkable, that two opposite opinions should still be held even respecting the origin of these parts, and that between these opinions botanists should be almost equally divided.

In a paper which was some time ago read to the Linnean Society of London,¹ I had occasion, in inculcating the necessity of examining the parts of the flower before expansion, to advert to this tribe of plants; and I there entered at some length, both into the opinions generally received respecting their male organs, and also into that which I had deduced from an examination of these parts before the opening of the corolla: and being unwilling to repeat now, what I then stated, I shall content myself with referring to the figures and descriptions published by Jacquin in the first volume of his "*Miscellanea Austriaca*," which give a correct idea of the state of the organs after expansion; and only add the observations I have made on one species of the family, the *Asclepias Syriaca*, in the earlier stages of the flower.

The flower-bud of this plant I first examined, while the unexpanded corolla was yet green and considerably shorter than the calyx. At this period, the gland-like bodies which afterwards occupy the angles of the stigma were absolutely invisible; the furrows of its angles were ex-

¹ [*Ante*, pp. 6—8.]

tremely slight, and, like the body of the stigma, green; the antheræ, however, were distinctly formed, easily separable from the stigma, and their cells, which were absolutely shut, were filled with a turbid fluid, the parts of which did not so cohere as to separate in a mass; of the cuculli, which in the expanded flower are so remarkable, and constitute the essential character of the genus, there was no appearance.

In the next stage submitted to examination, where the corolla nearly equalled the calyx in length, the gland-like bodies of the stigma were become visible, and consisted of two nearly filiform, light brown, parallel, contiguous and membranaceous substances, secreted by the sides of the furrow, which was now somewhat deeper: instead of the filiform processes, a gelatinous matter occupied an obliquely descending depression proceeding from towards the base of each side of the angular furrow.

In a somewhat more advanced stage, the membranes which afterwards become the glands of the stigma were found to be linear, closely approximated, and to adhere at their upper extremity. At the same time the gelatinous substance in the oblique depression had acquired a nearly membranaceous texture and a light brown colour, and on separating the gland from its furrow, which was then practicable, this membrane followed it. At this period, too, the contents of each cell of the anthera had acquired a certain degree of solidity, a determinate form, and were separable from the cell in one mass; the cuculli were also observable, but still very small and green, nearly scutelliform, having a central papilla, the rudiment of the future horn-like process. Immediately previous to the bursting of the cells of the antheræ, which takes place a little before the expansion of the corolla, the cuculli are completely formed, and between each, a pair of minute light green fleshy teeth are observable, the single teeth of each pair being divided from each other by the descending alæ of the antheræ. The glands of the stigma have acquired a form between elliptical and rhomboidal, a cartilaginous texture, and a brownish-black colour; they are easily separable from the secreting furrow, and on their

under surface there is no appearance of a suture, or any indication of their having originally consisted of two distinct parts: along with them separate also the descending processes, which are compressed, membranous, and light brown; their extremity, which is still unconnected, being more gelatinous but not perceptibly thickened. The pollen has acquired the yellow colour and the degree of consistence which it afterwards retains. On the bursting of the cells, the gelatinous extremity of each descending process becomes firmly united with the upper attenuated end of the corresponding mass of pollen. The parts are then in that condition in which they have been commonly examined, and are exhibited in the figures of Jacquin, who having seen them only in this state, naturally considered these plants as truly gynandrous, regarding the masses of pollen as the antheræ, originating in the glands of the stigma, and merely immersed in the open cells of the genuine antheræ, which he calls antheriferous sacs; an opinion in which he has been followed by Rottboell, Koellreuter, Cavanilles, Smith, and Desfontaines. The conclusion to be drawn from the observations now detailed is sufficiently obvious; but it is necessary to remark, that these observations do not entirely apply to all the plants which I have referred to the Asclepiadeæ, some of them, especially *Periploca*, having a granular pollen, applied in a very different manner to the glands of the stigma: they all, however, agree in having pollen coalescing into masses, which are fixed or applied to processes of the stigma, in a determinate manner; and this is, in fact, the essential character of the order. Dr. Smith, in the second edition of his valuable "Introduction to Botany," has noticed my opinion on this subject; but, probably from an indistinctness in the communication, which took place in conversation, has stated it in a manner somewhat different from what I intended to convey to him: for, according to his statement, the pollen is *projected* on the stigma. The term projection, however, seems to imply some degree of impetus, and at the same time presents the idea of something indeterminate respecting the part to which the body^{18]} so projected may be applied. But nothing can be more

constant than the manner in which the pollen is attached to the processes of the stigma in each species; and as considerable differences in this respect take place in various species, I have with advantage employed these in the new generic divisions of the order which I have attempted to establish, and to which I now proceed.

As I have everywhere mentioned the sources from which my knowledge both of genera and species is derived, it becomes unnecessary particularly to notice here the extent of my obligations to the unrivalled Herbarium of Sir JOSEPH BANKS, who, with his accustomed liberality, has permitted me to examine, and, where necessary, to dissect such specimens as seemed to have any peculiarity of structure, and has thus enabled me to found my genera on a more satisfactory induction than I could have done in any other part of the world.

ASCLEPIADEÆ.

I. POLLINIS MASSE (10-20) CERACEÆ, LÆVES, nec in granulis separandæ.

A CORPUSCULA STIGMATIS (5) SULCO LONGITUDINALI BIPARTIBILI, juxta basin, v. latere utrinque exsertentia processum, Massam Pollinis unicam, affigentem.

a. POLLINIS MASSE RECTÆ v. conniventes, stigmati incumbentes : basi, v. infra medium lateris, affixæ.

† Anthere apice simplices, nec membranâ terminatæ.

* Columna fructificationis tubo corollæ inclusa. Tubus stamineus extus appendiculatus.

1. Corollæ tubus e basi ventricosâ cylindraceus : limbi 5partiti lacinie ligulares, conniventes. Corona staminea 2plex, interioris foliola lobis exterioris opposita
2. Corolla campanulata, limbo 10fido, lacinis accessoris nanis. Corona staminea 2plex ; exterior 5fida lacinis bifidis ; interior 5phylla, foliis lacinis exterioris oppositis,
3. Corolla campanulata, 5fida. Corona staminea simplex, 5phylla, foliis dorso dentato-cristatis

CEROPEGIA.

HUBERIA.

PIRANTHEUS.

** Columna fructificationis exserta v. semi-exserta.

1. Corona staminea duplex : (utriusque figura varia) ; interior (cujus foliola antheris opposita) quandoque obsoleta. Masse pollinis altero margine cartilagineo-pellucido
2. Corona staminea simplex, 10phylla ; foliis 5, antheris oppositis, indivisis, obtusis, 5 reliquis bipartitis, acutis. Masse pollinis utroque margine opaco
3. Corona staminea simplex, 5loba, lobis antheris alternantibus ! denticulis nanis quandoque interjectis. Masse pollinis medio affixæ stigmati incumbentes
4. Corona staminea nulla : Corollæ subrotatæ faux squamis 5 coronata. Masse pollinis basi affixæ, apice coarctato pellucido

STAFELIA.

CARALLUMA.

MICROSTEMMA.

LEPTADENIA.

4. Corona staminea nulla: Corollæ subrotatæ faux squamis 5 coronatâ. Massæ pollinis basi affixæ, apice coarctato pellucido

†† Anthæræ membranâ terminatæ.

† Tubus stamineus extus appendiculatus.

* Corona staminea (simplex, 5phylla), foliolis *depressis*, carnosis. Corolla rotata.

1. Coronæ foliola angulo interiori producto in dentem anthæræ incumbentem
2. Coronæ foliola angulo interiori simplici edentulo

** Corona staminea (5phylla), foliolis *compressis*.

1. Coronæ foliola apice indivisa, intus edentula
2. Coronæ foliola apice indivisa, intus lacinulâ auctâ
3. Coronæ foliola bifida, lacinii apice recurvis

†† Tubus stamineus extus inappendiculatus.

1. Semina comosa. Folliculi graciles, leves. Corolla ureolata, fauce coronatâ, nunc edentulâ

2. Semina calva, marginata. Folliculi ventricosi, carnosi. Corolla subrotata, fauce edentulâ stigmate occultatâ. Corona staminea 1phylla, lobata.

1. Semina comosa. Folliculi sæpius costati v. muricati. (Caulis volubilis)

2. Semina calva, crenata. (Caulis erectus)

c. POLLINIS MASSÆ PENDULÆ, apice v. supra medium lateris affixæ. Anthæræ membranâ terminatæ.

† Tubus stamineus appendiculatus.

† Corona staminea simplex, 5phylla, foliolis antheris oppositis, lacinulis 5 nanis in eadem serie quandoque interjectis.

* Corolla 5partita reflexa. Coronæ foliola subeucullata, apice tubi filamentorum imposita.

1. Coronæ foliola eucullata, e fundo exserentia lacinulam corniformem. Folliculi graciles, leves

2. Coronæ foliola subeucullata, utrinque unidentata, absque lacinulâ interiori. Folliculi inflati, ramentacei

ASCLEPIAS.

GOMPHOCARPUS.

HOYA.

TYLOPHORA.

MARSDENIA.

PERGULARIA.

DISCHIDIA.

GYMNEMA.

SARCOLEPIS.

GONOLOBUS.

MATELEA.

I. POLLINIS MASSÆ (10-20) CEREACEÆ, LEVES, &c.—(contin.).		
A. CORPUSCULA STIGMATIS (5) SULCO LONGITUDINALI BIPARTIBILIA, &c.—(contin.).		
** Corolla rotata, non reflexa. Coronæ foliola acuta v. acuminata, intus edentula		OXYSTELMA.
*** Corollæ subcampanulatæ tubus ventricosus.		
a. Columna fructificationis exserta. Massæ pollinis compressæ.		
1. Coronæ foliola carnosæ, obtusæ, simplicia, denticulis 5 alternantibus nanis. Folliculi ventricosi, ramentacei		XYSMALOBUTUM.
2. Coronæ foliola longitudinaliter adnata, aversa, basi recurva. Folliculi ventricosi, læves		CALOTROPIS.
3. Coronæ foliola carnosæ, obtusæ, solutæ. Massæ pollinis curvaturæ processuum dein adscendentium affixæ. Corollæ lacinia ligulares		OXYPTALUM.
β. Columna fructificationis inclusa. Massæ pollinis ventricosa.		
1. Corolla campanulata. Coronæ foliola e basi dilatata subulata, intus simplicia. Massæ pollinis apice affixæ. Stigma muticum		KANAHIA.
2. Corolla urceolata. Coronæ foliola lacinulâ interiori parallelâ aucta. Massæ pollinis infra apicem affixa. Stigma rostratum		DIPLOLEPIS.
†† Coronæ staminea simplex, monophylla, tubo intus quandoque lacinia carnisve aucto. Corolla subrotata.		
1. Coronæ annularis integerrima. Massæ pollinis compressæ		HOLOSTEMMA.
2. Coronæ ore 5-20fidæ, dum 5fida lobis antheris oppositis		CYNANCHUM.
3. Coronæ 5partita, lobis antheris alternis		METAFLEXIS.
††† * Coronæ staminea composita, seriebus 2 v. 3 insertione distinctis.		
* Coronæ duplex, <i>exterior</i> 1 v. 5phylla, <i>interior</i> 5phylla. Faux corollæ esquamata.		
1. Coronæ <i>exterior</i> 5partita, lacinia acuminatis: <i>interior</i> brevior, exteriori opposita		DITASSA.
2. Coronæ <i>exterior</i> 5partita, cum v. absque lacinulis alternantibus, nanis: <i>interior</i> longior, exteriori alternans, foliolis basi solutis. Folliculi ramentacei		DOEMIA.
3. Coronæ <i>exterior</i> cyathiformis v. annularis, crenulatus: <i>interior</i> longior: <i>utraq</i> ue carnosæ. Folliculi graciles, læves		SARCOSTEMMA.

3. Corona *exterior* cystiformis v. annularis, crenulatus: *interior* longior: *utraq*
ue carnosa. Folliculi graciles, laeves.....

- ** Corona triplex, singula 5phylla, *extima* faucis inserta laciniis opposita; *media* foliolis tripartitis; *intima* indivisis.....
- †† Tubus stamineus nudus.
1. Faux corollæ campanulatæ coronata dentibus 5, tubum decurrentibus.....
 2. Faux corollæ urecolata, nuda. Tubus squamis 5, inclusis, cum fasciculis totidem pilorum alternantibus.....
 3. Faux corollæ sub-urecolata, ruda. Tubus esquamatus, pilosiusculus.....
- B. CORPUSCULA STIGMATIS (5) EXSULCA. Singula apice affigentia Massas pollinis 4, sessiles, subsimiles. Tubus stamineus appendiculatus. Corolla rotata.....

II. POLLINIS MASSÆ (5-20) GRANULOSÆ (granulis e spherulis 4 connatis compositis) 1-4 apici dilatato, soluto, corpusculi singuli stigmati, applicatæ.

1. Filamenta basi connata, supra distincta. Antheræ coherentes, imberbes. Massæ pollinis 4, corpusculo singulo affixæ. Corolla rotata.....
2. Filamenta longitudinaliter distincta. Antheræ coherentes, barbatae. Massa pollinis 1, corpusculo singulo affixa. Corolla rotata, squamis faucis aristatis.....
3. Filamenta longitudinaliter distincta. Antheræ imberbes. Massæ pollinis 4, corpusculo singulo affixæ. Corolla hypocrateriformis.....

EUSTEGIA.

METASTELMA.

MICROLOMA.

ASTEPHANUS.

SECAMONZ.

HEMIDESMUS.

PERIPLOCA.

GYMNANTHERA.

ASCLEPIADEÆ.

Contortarum genera, *Linn.* Apocinearum genera, *Juss.* Apocynarum genera, *Adans.*

CALYX quinquedivisus, persistens.

COROLLA monopetala, hypogyna, quinqueloba, regularis, æstivatione imbricata, rarissime valvata, decidua.

STAMINA 5, epipetala, laciniis limbi alternantia. *Filamenta* sæpius connata. *Antheræ* biloculares, septisve semi-completis nunc subquadriloculares. *Pollen* ad dehiscen-
tiam antherarum coalescens in massas numero loculorum,
rariusve per paria confluentes, et geminatim, v. quaternatim,
v. solitarie processibus quinque stigmatibus affixas.

OVARIA 2. *Styli* 2, arcuè approximati, sæpe brevissimi. *Stigma* ambobus commune, dilatatum, pentagonum, angulis corpusculiferis.

FOLLICULI 2; altero nunc abortiente. *Placenta* suturæ
intus applicata, demum libera.

SEMINA numerosa, imbricata, pendula, ad umbilicum
^{20]} sæpius comosa. *Albumen* tenue. *Embryo* dicotyle-
doneus, rectus. *Cotyledones* foliaceæ. *Radicula* suprema.
Plumula inconspicua.

Frutices, rariusve *Herbæ*, ut plurimum lactescentes et
volubiles. *Folia* integra, opposita, quandoque alterna v.
verticillata, ciliis interpetiolaribus loco stipularum sæpius
instructa. *Flores* subumbellati, fasciculati, v. racemosi,
interpetiulares.

I. ASCLEPIADEÆ VERÆ.

[21]

Massæ Pollinis 10, læves, per paria, (diversis antheris pertinentia), affixæ stigmatis corpusculis, sulco longitudinali, bipartilibus. *Filamenta* connata, extus sæpius appendiculata.

CEROPEGIA.

[Ceropegiae pleræque, Linn.]

CHAR. *Corolla* tubo e basi ventricosâ cylindraceo; limbi laciniis ligularibus.

Columna fructificationis inclusa.

Corona staminea duplex, *exterior* abbreviata, 5loba: *interior* 5phylla, foliolis ligularibus, indivisis, lobis exterioris oppositis.

Antheræ apice simplices.

Massæ pollinis erectæ, basi affixæ, marginibus simplicibus.

Stigma muticum.

Folliculi cylindracei, læves. *Semina* comosa.

HABITUS. *Herbæ* glabræ, volubiles. *Radice* tuberosa.

Umbellæ interpetiolares, nunc paucifloræ.

PATRIA. India Orientalis.

OBS. To this genus belong *C. Candelabrum*, Linn.; *C. biflora*, Linn.; *C. tuberosa*, *bulbosa*, *juncea*, *acuminata*, of Roxburgh, and two undescribed species.

HUERNIA.

[Stapelieæ species, Linn. Juss. Masson.] [22]

CHAR. *Corolla* campanulata, limbo decemfido, laciniis accessoriis nanis, dentiformibus.

Columna fructificationis inclusa.

Corona staminea duplex; *exterior* quinquefida, laciniis bifidis: *interior* 5phylla, foliolis e basi gibbosâ subulatis, indivisis, laciniis exterioris alternantibus.

Antheræ apice simplices.

Massæ pollinis erectæ, basi affixæ, altero margine cartilagineo-pellucido.

Stigma muticum.

Folliculi subcylindracci, læves. *Semina* comosa.

HABITUS *Stapelie*.

PATRIA. Africa Australis.

OBS. The whole of the third section of *Stapelia* in Willdenow's edition of the "Species Plantarum," probably belongs to this genus; but I have only had an opportunity of examining *S. campanulata*, *venusta*, and *guttata*, from which the character is formed. I have named the genus in memory of Justus Huernius, one of the earliest collectors of Cape plants, and from whose drawings the first account of *Stapelia* was taken.

23] PIARANTHUS.

[*Stapelie* sp. *Masson*.

CHAR. *Corolla* campanulata, 5fida, carnosa.

Columna fructificationis inclusa.

Corona staminea simplex, 5phylla, foliolis dorso dentatis.

Antheræ apice simplices.

Massæ pollinis erectæ, basi affixæ, altero margine cartilagineo-pellucido.

Stigma muticum.

Folliculi.

HABITUS *Stapelie*.

PATRIA. Africa Australis.

OBS. The want of the external corona renders it necessary to separate from *Stapelia* and *Huernia*, this genus; of which the only two certain species are *Stapelia punctata* and *pulla* of *Masson*; of both these I have examined specimens collected by *Masson*, and preserved in spirits, in the collection of Sir Joseph Banks.

STAPELIA.

[*Stapelie* plures, *Linn. et Mass.*

CHAR. *Corolla* rotata, 5fida, carnosa.

Columna fructificationis exserta.

Corona staminea duplex, utraque in variis varia; interior quandoque obsoleta.

Antheræ apice simplices.

Massæ pollinis basi affixæ, altero margine cartilagineo-pellucido.

Stigma muticum.

Folliculi subcylindracei, læves. *Semina* comosa.

HABITUS. *Plantæ* carnosæ, aphyllæ, angulatæ, sæpe [24] tuberculatæ. *Flores*, ut plurimum speciosi, odore nauseoso, stercorario.

PATRIA. Africa Australis, præsertim in planitiis desertis, argillaceis, *Karoo* nuncupatis.

Obs. The essential character of this extensive and singular genus, consists, according to Linnæus and all subsequent botanists, in the double corona. But I have already shewn, that certain plants that have been referred to it, and which entirely accord in habit, have a corona of a single series; and it will hereafter appear, that other, and very different genera, agree with *Stapelia* in this part of its structure. The genus, even as it is here limited, is capable of further subdivision; and I have little doubt, that when the species become better known, such a subdivision will be found expedient, and probably from characters like the following, by which, in the mean time, it may be disposed into very natural sections.

I. *Corona exterior 5phylla, foliolis indivisis.*

Stapelia hirsuta, Linn. *sororia*, Mass. *vetula*, Mass. *ambigua*, Mass. *asterias*, Mass. *glandulifera*, Mass. *stellaris*, Jacq. *ined.*

II. *Corona exterior 5partita, laciniis bifidis.*

Stapelia revoluta, Mass. *pedunculata*, Mass. *verru-* [25] *cosa*, Mass. *mixta*, Mass. *variegata*, Linn. *lepida*, Jacq. *ined.*

III. *Corona exterior 1phylla, indivisa vel ciliato-multifida.*

Stapelia articulata, Hort. Kew. et Mass. *geminata*, Mass., *aliæque ineditæ*, in Musco Banksiano, spir. vin. asservatæ.

CARALLUMA.

CHAR. *Corolla* rotata, profundè 5fida.

Columna fructificationis exserta.

Corona staminea simplici serie 10phylla; foliolis quinque antheris oppositis indivisis; reliquis bipartitis, subulatis.

Antheræ apice simplices.

Massæ pollinis erectæ, basi affixæ, marginibus simplicibus.

Stigma muticum.

Folliculi graciles, læves. *Semina* comosa.

HABITUS ferè *Stapelie*.

PATRIA. India Orientalis.

OBS. This genus is the *Stapelia adscendens* of Roxburgh, the *Car-allum* of the Telingas.

MICROSTEMMA.

CHAR. *Corolla* rotata, 5fida.

Columna fructificationis exserta.

Corona staminea monophylla, carnosa, 5loba, lobis cum antheris alternantibus.

²⁶ *Antheræ* apice simplices.

Massæ pollinis medio lateri insertæ, stigmati incumbentes.

Stigma muticum.

Folliculi graciles, læves. *Semina* comosa.

HABITUS. *Herba* glabra, erecta. *Radix* tuberosa. *Caulis* infrà simplex, foliis minutis; suprà ramosus, foliis oppositis linearibus. *Umbellæ* laterales et terminales, subsessiles. *Corollæ* nigro-purpureæ intùs barbatae.

PATRIA. Nova Hollandia tropica.

HOYA.

CHAR. *Corolla* rotata, 5fida.

Corona staminea 5phylla, foliolis depressis, carnis, angulo interiore producto in dentem antheræ incumbentem.

Antheræ membranâ terminatæ.

Massæ pollinis basi affixæ, conniventes, compressæ.

Stigma depressum, papillâ obtusâ.

Folliculi læves. *Semina* comosa.

HABITUS. *Caulis* suffruticosus, volubilis, v. decumbens.

Folia opposita, carnosa, v. membranacea. *Umbellæ* interpetiolares, multifloræ.

PATRIA. India Orientalis, China, et Nova Hollandia tropica.

OBS. I have named this genus in honour of Mr. THOMAS HOY, whose merits as an intelligent and successful cultivator have been long known to the botanists of this country. I have added specific characters of the only two species with which I am acquainted; but *Hoya carnosa* probably includes several species, which can only be determined from living specimens: it is also to be considered as the type of the genus, *Hoya viridiflora* differing in some degree in the structure of its corona, and considerably in habit.

1. *H. carnosa*, foliis ovali-oblongis carnosis, corollis barbatis, coronæ foliolis subtus sulcatis!

Asclepias carnosa, Linn. suppl. 170. Murr. syst. veg. ed. 14, p. 260. Willd. sp. pl. 1, p. 1264. Pers. syn. 1, p. 275. Sims in bot. magaz. t. 788. Smith, exot. bot. 2, p. 21, t. 70.

Stapelia Chinensis, Lour. Coch. 1, p. 205, fide specim. ab auctore missi in Herb. Banks.

HAB. In Asiæ tropicæ variis regionibus, etiam in Nova Hollandia (ubi et in hort. Angl. v. v.)

2. *H. viridiflora*, foliis ovatis acuminatis membranaceis corollisque glabris, coronæ foliolis exsulcatis.

Asclepias volubilis, Linn. suppl. 170*. Willd. sp. pl. 1, p. 1269. Pers. syn. 1, p. 276.

Watta-haka-codi, Rheed. Malab. 9, p. 25, t. 15.

HAB. Inter frutices in nemorosis Zeylonæ, J. G. Kæinig, in Herb. Banks. (ubi v. s.)

28] TYLOPHORA.

CHAR. *Corolla* rotata, 5partita.

Corona staminea 5phylla, foliolis depressis, carnosis, angulo interiori simplici edentulo.

Antheræ membranâ terminatæ.

Massæ pollinis erectæ, basi affixæ, marginibus simplicibus.

Stigma muticum.

Folliculi læves. *Semina* comosa.

HABITUS. *Herbæ* v. *suffrutices* volubiles. *Folia* opposita, membranacea, plana. *Umbellæ* interpetiolares. *Flores* ut plurimum parvi.

PATRIA. Nova Hollandia, præsertim intra tropicum, sed usque ad grad. 33. lat. aust. India Orientalis, et Africa æquinotialis. Octo species nobis cognitæ, quarum nullæ adhucdum editæ sunt.

MARSDENIA.

CHAR. *Corolla* urceolata, 5fida, nunc subrotata.

Corona staminea 5phylla, foliolis compressis, indivisis, intus edentulis.

Antheræ membranâ terminatæ.

Massæ pollinis erectæ, basi affixæ.

Folliculi læves. *Semina* comosa.

HABITUS. *Suffrutices* sæpius volubiles. *Folia* opposita, latiuscula, plana. *Cymæ*, nunc *Thyrsi*, interpetiolares.

Stigma sæpius muticum, quandoque rostratum, rostro indiviso vel bifido.

29] PATRIA. India Orientalis et Nova Hollandia; rarius in America Meridionali, et Syria.

Obs. This genus differs from *Pergularia*, chiefly in the want of the inner lacinia of the corona: it is therefore an arbitrary separation, and made principally to obtain clearer characters for both. The two species, with an elongated stigma, are perhaps not truly of this genus, but if separated from it, must form each a distinct genus.

It is named in honour of WILLIAM MARSDEN, Esq.,

F.R.S., late Secretary to the Admiralty, and author of a very judicious and learned "History of Sumatra," in which, though it is evident that he has not made botany his particular study, he has had the merit of turning the attention of botanists to several valuable plants, among others, to the Camphor-Tree of Sumatra, and to a species of this genus, *Marsdenia tinctoria*, said to afford the best indigo in that island.

† *Stigma muticum*. *Marsdeniæ* veræ.

1. *M. velutina*, caule volubili, foliis cordatis latè ovatis acuminatis tomentosis mollibus, cymis umbelliformibus, fauce nudâ.

HAB. In Nova Hollandia, intra tropicum, (ubi v. v.)

2. *M. tinctoria*, caule volubili, foliis cordatis ovato-oblongis acuminatis glabriusculis basi anticè glandulosis, thyrsis lateralibus, fauce barbatâ.

Tarram akkar. *Mars. Sumat.* 78.

HAB. In insula Sumatra. (v. s. in Herb. Banks.)

3. *M. viridiflora*, caule volubili, foliis oblongo-lanceolatis glabriusculis basi obtusa, tubo intus villosiusculo.

HAB. In Nova Hollandia, intra tropicum, (ubi v. v.)

4. *M. clausa*, caule volubili, foliis lanceolatis utrinque acutis glabris: supra parum rugosis, fauce densè barbatâ.

HAB. In Jamaica. *Swartz.* (in Herb. Banks. ubi v. s.)

5. *M. suaveolens*, caule suberecto, foliis ovali-lanceolatis glabris aveniis, tubo ventricosus, fauce barbatâ.

HAB. In Nova Hollandia, extra tropicum, (ubi v. v.)

6. *M. cinerascens*, caule erecto, foliis ovatis obtusiusculis venosis pube rarâ conspersis, petiolis semuncialibus, corollis subrotatis.

HAB. In Nova Hollandia, intra tropicum, (ubi v. v.)

31]

†† *Stigma rostratum*.

7. *M. erecta*, caule erecto, foliis cordatis ovatis acutis, cymis umbelliformibus, limbi laciniis imberbibus tubo 4-5ies longioribus.

Cynanchum erectum, *Linn.*

8. *M. rostrata*, caule volubili, foliis ovatis subcordatis acuminatis glabris, umbellis multifloris, limbo barbato.

HAB. In Nova Hollandia extra tropicum, (ubi v. v.)

PERGULARIA.

[*Pergulariæ* species, *Linn.*

CHAR. *Corolla* hypocrateriformis, tubo urceolato.

Corona staminea 5phylla, foliolis compressis apice indivisis, intus lacinulâ cuctis.

Antheræ membranâ terminatæ.

Massæ pollinis erectæ, basi affixæ.

Stigma muticum.

Folliculi ventricosi, læves. *Semina* comosa.

HABITUS. *Plantæ* volubiles. *Folia* latiuscula, membranacea. *Cymæ* interpetiolares. *Flores* flavescentes, odoratissimi.

PATRIA ignota: in China et India Orientali ob flores suaveolentes culta.

OBS. Of this genus the only certain species are *Pergularia odoratissima*, *Roxb. et Smith*, and *P. minor*, *And. Repos.* 160. *Pergularia purpurea*, *Vahl.* and *Japonica*, *Thunb.* may belong to it. *P. edulis* of Thunberg, prod. cap. is probably very different. When Linnæus established this genus in his Mantissa, he certainly meant his character to apply to *Pergularia glabra*, of which he had a specimen in his herbarium, and which is the *Flos Pergulanus* of Rumphius; but unfortunately this plant does not belong to the order of Asclepiadæ, but to that section of Apocineæ of which I shall hereafter treat. The character of Linnæus was no doubt chiefly taken from a plant of

P. odoratissima, that had flowered in the Upsal Garden, and which he confounded with the *Asclepias cordata* of Forskael, an error long since pointed out by Dr. Smith in his very accurate and satisfactory account of *Pergularia odoratissima*.

DISCHIDIA.

CHAR. *Corolla* urceolata, 5fida.

Corona staminea 5phylla, foliolis bifidis, laciniis subulatis, patentibus, apice recurvis.

Antheræ membranâ terminatæ.

Massæ pollinis erectæ, basi affixæ.

Stigma muticum.

Folliculi læves. *Semina* comosa.

HABITUS. *Herba* in arboribus parasitica, dependens, perennis, lactescens, farinâ albâ tota conspersa. *Caulis* ad genicula radicans. *Folia* opposita, subrotunda, crassa, carnosa. *Flores* parvi, subumbellati.

PATRIA. India Orientalis, præsertim Insulæ Moluccanæ, necnon Nova Hollandia, ubi prope Endeavour River detexit *Illustr. Banks*.

GYMNEMA.

CHAR. *Corolla* suburceolata 5fida. *Fauce* sæpe coronatâ, squamulis denticulise 5, sinubus insertis.

Corona staminea nulla.

Antheræ membranâ terminatæ.

Massæ pollinis erectæ, basi affixæ.

Folliculi graciles, læves. *Semina* comosa.

HABITUS. *Suffrutices* sæpius volubiles. *Folia* opposita, membranacea, plana. *Umbellæ* interpetiolares, cymosæ.

PATRIA. India Orientalis, Nova Hollandia tropica, et Africa æquinoctialis.

OBS. Of this genus I have examined four species. Two of these are unpublished plants; the third is *Asclepias lactifera* Linn., of which there is no specimen in the

Linnean Herbarium: it therefore rests entirely upon Hermann's specimens, which, though collected 140 years ago, were by maceration in water so far recovered, as to enable me with certainty to determine its genus. The fourth is *Periploca sylvestris*, *Willd. sp. pl. 1, p. 1252*.

34] LEPTADENIA.

CHAR. *Corolla* subrotata, *tubo* brevi, *fauce* coronata, squamis 5 sinibus impositis: *limbo* barbato, æstivatione valvata.

Corona staminea nulla.

Antheræ liberæ, apice simplices.

Massæ pollinis erectæ, basi affixæ, apice coarctato pellucido!

Stigma muticum.

Folliculi

HABITUS. *Herbæ*? perennes? volubiles, tomento impalpabili, pulvereo, cinerascens. *Folia* plana, opposita. *Umbellæ* interpetiolares, quandoque cymosæ. *Corpuscula* stigmatis minuta.

PATRIA. India Orientalis, Africa æquinoctialis et septentrionalis.

OBS. Of this genus I have examined three species in the Banksian Herbarium, none of which are as yet described, though one of them was collected by Forskael; it is unnamed, however, and does not correspond with any of his descriptions.

SARCOLOBUS.

CHAR. *Corolla* subrotata, 5fida. *Fauce* nudâ.

Corona staminea nulla.

Antheræ membranâ terminatæ.

35] *Massæ pollinis* erectæ, basi affixæ.

Stigma muticum.

Folliculi ventricosi, carnosi. *Semina* marginata!

HABITUS. *Frutex* volubilis, glaber. *Folia* opposita, latiuscula. *Umbellæ* interpetiolares, multifloræ.

PATRIA. Java; in Prince's Island, prope Bataviam, dextit *Illust. Banks.*

GONOLOBUS. [*Richard?* in *Mich. Fl. Bor. Amer.*
Cynanchi species, *Linn. Jacq.*

CHAR. *Corolla* subrotata, 5partita.

Corona staminea scutelliformis, lobata.

Antheræ transversim dehiscentes, membranâ terminatæ.

Massæ pollinis extremitati exteriori respectu loculi affixæ stigmatæ tectæ.

Stigma planiusculo-depressum.

Folliculi ventricosi, subcostati. *Semina* comosa.

HABITUS. *Suffrutices* volubiles. *Folia* opposita, latiuscula. *Umbellæ* interpetiolares.

PATRIA. America, præsertim intra tropicos.

Obs. *Cynanchum maritimum* *Linn.* suberosum *Linn.* crispiflorum *Hort. Kew.* belong to this genus; and I suppose also *C. planiflorum*, *grandiflorum*, *rostratum*, *nigrum*, *racemosum*, *Carolinense*, *obliquum*, *hirtum*, *prostratum*, and *undulatum* of Willdenow's *Spec. Plant.*: these, however, I have not determined, and the whole genus requires to be re-examined.

MATELEA. [*Aubl. Gujan. tab.* 109, *Hostea.*
Willd. sp. pl. 1, p. 1274.

CHAR. *Corolla* rotata, 5partita.

Corona staminea scutelliformis, lobata.

Antheræ transversim dehiscentes, membranâ terminatæ.

Massæ pollinis extremitate exteriori respectu loculi affixæ, stigmatæ tectæ.

Stigma planiusculo-depressum.

Folliculi ventricosi, costati. *Semina* calva. (*Aubl.*)

HABITUS. *Frutex* erectus. *Folia* opposita, basi supra biglandulosa. *Flores* racemosi, laterales.

PATRIA. America meridionalis.

ASCLEPIAS.

[*Asclepiadis* sp. *Linn.*

CHAR. *Corolla* 5partita, reflexa.

Corona summo tubo filamentorum imposita, 5phylla, foliolis cucullatis, e fundo exserentibus processum aversum corniformem.

Antheræ membranâ terminatæ.

Massæ pollinis compressæ, apice attenuato affixæ, pendulæ.

Stigma depressum, muticum.

Folliculi læves. *Semina* comosa.

^{37]} HABITUS. *Herbæ* erectæ. *Folia* opposita, nunc alterna! vel verticillata. *Umbellæ* interpetiolares.

PATRIA. America Septentrionalis.

OBS. The species of this genus which I have examined, are *A. Syriaca*, (most improperly so called, as it is a native of North America only), *amæna*, *purpurascens*, *variegata*, *curassavica*, *nivea*, *elevata*, *parviflora*, *incarnata*, *decumbens*, *verticillata*, *tuberosa*, of Willdenow's Spec. Plant., and some unpublished species in the Banksian Herbarium; and there can be little doubt that *A. pulchra*, *citrifolia*, *Mexicana*, *linaria*, *rubra*, and all the species of Michaux, belong to it.

GOMPHOCARPUS.

[*Asclepiadis* sp. *Linn.*

CHAR. *Corolla* 5partita, reflexa.

Corona summo tubo filamentorum imposita, 5phylla, foliolis cucullatis utrinque unidentatis, intus simplicibus.

Antheræ membranâ terminatæ.

Massæ pollinis compressæ, apice attenuato affixæ, pendulæ.

Stigma depressum, muticum.

Folliculi ventricosi, spinis innocuis echinati.

Semina comosa.

HABITUS. *Frutices* v. *suffrutices* erecti. *Folia* opposita, margine sæpe revolutæ. *Umbellæ* interpetiolares.

PATRIA. Africa Australis.

Obs. *Asclepias arborescens*, *fruticosa* and *setosa*, are the three certain species of this genus; but *A. crispa* has probably the same structure. The flowers I have examined, but have not seen the fruit. *A. pubescens* ought probably to be altogether omitted; for, according to the description and specimen in Linnæus's Herbarium, it is *A. arborescens*, while according to the reference to Plukenet, it is *A. crispa*. This observation I have copied from Mr. Dryander's notes.

XYSMALOBIIUM.

[*Asclepiadis* sp. Linn.

CHAR. *Corolla* 5fida, patens.

Corona staminea summo tubo filamentorum imposita, simplici serie decempartita, laciniis 5 antheris oppositis carnosius, subrotundis, intus simplicibus; 5 reliquis nanis.

Antheræ membranâ terminatæ.

Massæ pollinis compressæ, apice affixæ, pendulæ, processibus connectentibus latiusculis.

Stigma muticum.

Folliculi ventricosi, ramentacei. *Semina* comosa.

HABITUS. *Frutices* erecti. *Folia* opposita. *Umbellæ* interpetiolares. *Flores* majusculi, limbo nunc barbato.

PATRIA. Africa Australis.

Obs. This genus at present consists of only two species, *Asclepias undulata* and *grandiflora*.

CALOTROPIS.

[*Asclepiadis* sp. Linn.

CHAR. *Corolla* sub-campanulata, tubo angulato, angulis intus saccatis, limbo 5partito.

Corona staminea 5phylla, foliolis carinæformibus, tubo filamentorum longitudinaliter adnatis, basi recurvâ.

Antheræ membranâ terminatæ.

Massæ pollinis compressæ, apice attenuato affixæ, pendulæ.

Stigma muticum.

Folliculi ventricosi, læves. *Semina* comosa.

HABITUS. *Frutices* erecti, glabri. *Folia* opposita, lata. *Umbellæ* interpetiolares. *Flores* speciosi.

PATRIA. India Orientalis, et Persia.

[¹ SPECIES. *Asclepias gigantea* et *procera*.]

KANAHIA.

CHAR. *Corolla* campanulata, limbo 5partito.

Columna semi-inclusa.

Corona staminea apice tubi filamentorum imposita, 5phylla, foliolis e basi incrassatâ subulatis, indivisis.

Antheræ membranâ terminatæ.

^{40]} *Massæ pollinis* ventricosæ, apice affixæ, pendulæ.

Stigma muticum.

Folliculi graciles, striati. *Semina* comosa?

HABITUS. *Frutex*? erectus. *Folia* opposita, plana. *Pedunculi* interpetiolares, apice fasciculatim multiflori, pedicellis imbricatis, basi unibracteatis.

PATRIA. Arabia, ¹[et Abyssinia.]

OBS. *Asclepias laniflora*, Forsk. *Ægypt. Arab.* p. 51, KANAHH of the Arabs, is the only species of the genus. I have examined the flowers of an original specimen. The account of the fruit is taken from Forskael's description, and Vahl's figure.

OXYSTELMA.

CHAR. *Corolla* rotata, patens.

Columna exserta.

Corona staminea 5phylla, foliolis acutis, indivisis.

Antheræ membranâ terminatæ.

¹ Added in MS. by the author.

Massæ pollinis compressæ, apice attenuato affixæ, pendulæ.

Stigma muticum.

Folliculi læves. *Semina* comosa.

HABITUS. *Suffrutices* volubiles, glabri. *Folia* opposita. *Racemi* v. *umbellæ* interpetiolares.

PATRIA. India Orientalis et Nova Hollandia?

OBS. *Periploca esculenta*, *Linn. Suppl. Roxb. Corom.* [41] 1, p. 13, t. 11, is the type of the genus, and perhaps the only genuine species; for a New Holland plant which I have at present joined with it, differs considerably both in the form of its corolla and that of its corona.

OXYPETALUM.

CHAR. *Corolla* tubo brevi ventricoso; limbo 5partito, laciniis supra ligulatis.

Corona staminea 5phylla, foliolis subrotundis simplicibus, carnosius.

Antheræ membranâ terminatæ.

Massæ pollinis lineares, pendulæ, affixæ curvaturæ processuum, dein adscendentium.

Stigma acumine elongato bipartito.

Folliculi

HABITUS. *Frutex* volubilis. *Folia* opposita, cordata. *Umbellæ* subcorymbosæ, interpetiolares. *Flores* suaveolentes.

PATRIA. America Meridionalis.

OBS. The genus consists of a single species, which was discovered in 1768, near Rio de Janeiro, by Sir Joseph Banks.

DIPLOLEPIS.

CHAR. *Corolla* tubo brevi, urceolato, limbo 5partito.

Corona staminea 5phylla, foliolis obtusis squamulâ [42] interiori auctis.

Antheræ membranâ terminatæ.

Massæ pollinis ventricosæ, juxta apicem affixæ, pendulæ.

Stigma rostro elongato indiviso.

Folliculi

HABITUS. [¹*Frutex* decumbens, glaber. *Folia* oblonga, mucronulata, coriacea. *Cymæ* congestæ, extraalares et terminales.]

PATRIA. America Australis.

Obs. Only one certain species is known, which was discovered by my friend *Mr. Archibald Menzies*, in Chili, near Valparaizo.

Asclepias vomitoria of Kœnig's MSS., of which I have examined specimens in the Banksian Herbarium, agrees with this genus in many respects, especially in the form and insertion of its masses of pollen, but differs considerably in appearance, and in having a blunt stigma.

HOLOSTEMMA.

CHAR. *Corolla* subrotata, 5fida.

Corona imo tubo stamineo inserta, simplex, annularis, integra.

Antheræ membranâ terminatæ.

Massæ pollinis pendulæ, compressæ, apice attenuato affixæ.

Stigma muticum.

Folliculi ventricosi, læves. *Semina* comosa.

HABITUS. *Frutex* volubilis, glaber. *Folia* opposita, lata. *Umbellæ* interpetiolares, subsessiles. *Flores* speciosi.

^{43]} PATRIA. India Orientalis.

Obs. This is Ada-Kodien, *Rheed. Mal.* 9, p. 9, t. 7, in which the leaves are represented alternate, an error that is corrected in the description, which is excellent, and well accords with a specimen in the Banksian Herbarium collected by Dr. Patrick Russell.

¹ Added in MS. by the author.

CYNANCHUM.

[Cynanchi sp. *Linn.*CHAR. *Corolla* subrotata, 5partita.*Corona staminea* monophylla, 5-20loba, cum 5fida lobis antheris oppositis.*Antheræ* membranâ terminatæ.*Massæ pollinis* ventricosæ, pendulæ.*Stigma* apiculatum.*Folliculi* læves. *Semina* comosa.HABITUS. *Plantæ* perennes, vel *suffrutices*, ut plurimum volubiles. *Folia* opposita. *Umbellæ* interpetiolares.

PATRIA varia, a gradu 59° lat. bor. ! usque ad 32° lat. aust.

Obs. Though I have here very much limited the Linnean genus *Cynanchum*, yet it appears still to contain the elements of several genera: I shall therefore add the characters of all the species I am acquainted with, divided into such sections as will probably hereafter be considered distinct genera.

I. Corona staminea tubulosa, columnam includens, ore 14 5-10fido. *Lacinia* 5 interiores antheris exterioribusque oppositæ et parallelæ.

Massæ pollinis infra apicem insertæ.*Stigma* apiculo semibifido.*Folliculi* cylindracei, divaricati.*Caulis* volubilis. *Folia* cordata.

1. *C. acutum*, foliis oblongo-ovatis cordatis acutis, corollæ laciniis oblongis obtusis.

Cynanchum acutum, *Linn.*, cum synonymis.

2. *C. Monspeliacum*, foliis reniformibus: apice coarctato semi-lanceolato, laciniis corollæ lanceolatis obtusiusculis.

Cynanchum Monspeliacum, *Linn.*, cum synonymis.

Obs. Forsan haud distinctum a priori.

3. *C. Chinense*, foliis ovatis cordatis: acuminibus brevi,

laciniis corollæ lanceolato-linearibus acutis, coronæ laciniis 5 exterioribus compresso-filiformibus integris.

HAB. In Chinæ provincia Pechelej, *Georgius Staunton*, *Baronettus*. (v. s. in Herb. Banks.)

^{45]} II. *Corona staminea* tubulosa, columnam includens, ore 5-10fido, carinis decem interioribus simplicibus v. in lacinulas supra productis.

Massæ pollinis infra apicem insertæ.

Stigma apiculo emarginato.

Folliculi ventricosi, angulati, patentes.

Caulis volubilis. *Folia* cordata.

4. *C. pedunculatum*, pedunculo communi foliis glabrisculis longiore, carinis interioribus coronæ in lacinulas productis.

HAB. In Nova Hollandia tropica (ubi v. v.)

5. *C. floribundum*, pedunculo communi foliis glaberrimis brevioribus, carinis interioribus coronæ in lacinulas productis.

HAB. In Nova Hollandia extra tropicum (ubi v. v.)

6. *C. erubescens*, carinis interioribus coronæ 10fidæ corolla brevioris simplicibus, pedunculo communi petiolum æquante, ramulis floribusque extus pubescentibus.

HAB. In Nova Hollandia tropica, ubi a *Josepho Banks*, *Baronetto*, detectum.

7. *C. pauciflorum*, carinis interioribus coronæ 5fidæ corollam æquantis simplicibus, pedunculo communi petiolo brevioribus, ramulis floribusque glabris.

^{46]} *Periploca tunicata*, *Retz. obs. 2, p. 15. Willd. phyt. 1, p. 7, n. 23, t. 5, f. 3. Willd. sp. pl. 1, p. 1252.*

HAB. In India Orientali. (v. s. in Herb. Banks.)

III. *Corona staminea* tubulosa, columnam includens, ore 5-10fido, laciniis carinisve interioribus nullis.

Massæ pollinis apice sæpius affixæ.

Stigma apiculo bifido.

Caulis volubilis. *Folia* subcordata.

8. *C. pilosum*, foliis ovatis acutiusculis calycibusque pilosis, corona 10fida longitudine corollæ.

Periploca Africana, *Linn.*, cum synonymis.

HAB. In Africa Australi prope Promont. B. Spei. (ubi v. v.)

9. *C. crassifolium*, foliis ovatis subcordatis obtusis cum mucronulo carnosus calycibusque glabris, corona 10fida longitudine corollæ.

Cynanchum obtusifolium, *Linn. suppl.* 169?

HAB. In Africa Australi. (v. s. in Herb. Banks.)

10. *C. Capense*, foliis ovatis cordatis obtusis cum mucronulo caulibusque glabris, calycibus pubescentibus, corona 5fida corollâ duplo brevior.

Cynanchum Capense, *Linn. suppl.* 168?

HAB. In Africa Australi. (v. s. in Herb. Banks.)

IV. *Corona staminea* tubulosa, columnam includeus, ore 10fido, intus laciniis carinisve nullis.

Massæ pollinis apice affixæ.

Stigma apiculo integro.

Caulis erectus.

11. *C. roseum*, foliis lanceolato-linearibus undulatis glabrisculis, pedicellis calycibusque pilosis.

Asclepias foliis ex lineari-lanceolatis, floribus umbellatis, umbellis alternis erectis, caule erecto. *Gmel. Sib.* 4, p. 78, t. 42.

Asclepias Davurica, *Willd. sp. pl.* 1, p. 1272?

HAB. In Siberia. *P. S. Pallas*, in Herb. Banks.

V. *Corona scutelliformis*, carnosus, 5-10loba, intus simplex.

Massæ pollinis infra apicem affixæ.

Stigma apiculo brevissimo, integro.

Folliculi læves.

Caules erectiusculi.

12. *C. Vincetoxicum*, caule erecto, corollis imberbibus, pedicellis umbellæ simplicis pedunculo communi triplo longioribus, corona 5loba.

Asclepias Vincetoxicum, *Linn.*

HAB. In Europa: in Suecia, etiam ad 59° 40' Lat.

^{48]} 13. *C. medium*, caule supernè volubili, corollis imberbibus, pedicellis umbellæ sæpe divisæ pedunculo communi vix longioribus, corona 5loba.

HAB. Specimen e Hort. Reg. Paris. in Herb. Banks.

Sequenti proximum.

14. *C. nigrum*, caule supernè volubili, corollis barbatis, pedicellis umbellæ simplicis pedunculo communi vix longioribus, corona semidecemfida.

Asclepias nigra, *Linn.*

HAB. In Europa Australi.

VI. Corona profundè 5fida, laciniis simplicibus.

Massæ pollinis infra apicem affixæ.

Stigma papilla emarginata.

Folliculi ventricosi.

15. *C. Sibiricum*, foliis lanceolato-linearibus oppositis ternisque, caule decumbente.

Asclepias Sibirica, *Linn.*

HAB. In Siberia, etiam in China. (v. s. in Herb. Banks.)

OBS. Vix hujus generis.

METAPLEXIS.

CHAR. *Corolla* subrotata.

Corona staminea 5phylla, foliolis nanis, cucullatis, cum antheris alternantibus.

Antheræ membranâ terminatæ.

Massæ pollinis ventricosæ, pendulæ, latere fixæ. [49]

Stigma rostro elongato indiviso.

Folliculi

HABITUS. *Suffrutex* volubilis, glaber. *Folia* cordata. *Racemi* pedunculati, interpetiolares. *Corollæ* limbus barbatus.

PATRIA. China, in provincia Pecheleiy. *Georgius Staunton*, *Baronettus*.

DITASSA.

CHAR. *Corolla* subrotata.

Corona staminea duplex; *exterior* 5partita, laciniis acuminatis; *interior* 5phylla, brevior, exteriori antherisque opposita.

Antheræ membranâ terminatæ.

Massæ pollinis ventricosæ, infra apicem affixæ, pendulæ.

Stigma apiculo obtuso.

Folliculi

HABITUS. *Suffrutex* volubilis, glaber. *Folia* opposita, plana. *Umbellæ* interpetiolares.

PATRIA. America Meridionalis, Brasilia, ubi prope Rio de Janeiro detexit *Josephus Banks*, *Baronettus*.

DŒMIA.

[50]

CHAR. *Corolla* subrotata, tubo brevi.

Corona staminea duplex, *exterior* brevis, 10partita, laciniis alternis nanis; *interior* 5phylla, foliolis basi solutis, supra subulatis.

Antheræ membranâ terminatæ.

Massæ pollinis compressæ, apice affixæ, pendulæ.

Stigma muticum.

Folliculi ramentacei. *Semina* comosa.

HABITUS. *Caulis* volubilis. *Folia* opposita, cordata. *Flores* umbellati.

PATRIA. India Orientalis, et Africa Æquinoctialis.

OBS. *Asclepias cordata*, *Forsk. Arab. p. 49*, and *Cynanchum extensum*, *Jacq. ic. 1, t. 54*, are the only certain species of the genus: from the latter, *Asclepias scandens*, *Palis. Flor. D'Oware, 1, p. 92, t. 56*, is probably not distinct, and *Cynanchum bicolor*, *And. Repos. 562*, is very nearly related to it.

SARCOSTEMMA.

CHAR. *Corolla* rotata.

Corona staminea duplex; *exterior* cyathiformis v. annularis, crenata; *interior* 5phylla, *exteriorem* superans, foliolis carnosus.

Antheræ membranâ terminatæ.

⁵¹ *Massæ pollinis* apice affixæ, pendulæ.

Stigma submuticum.

Folliculi graciles, læves. *Semina* comosa.

HABITUS. *Caulis* volubilis v. decumbens, aphyllus, articulatus, v. foliis oppositis distantibus. *Flores* umbellati, laterales v. terminales.

PATRIA. India Orientalis, Nova Hollandia, et Africa.

OBS. To this genus belong *Cynanchum viminale*, *Linn.*; a species nearly allied to it from New Holland and New Caledonia; and *Asclepias viminalis*, *Linn.*, all of which I have examined. The probable species are, *Asclepias aphylla*, *Thunb. prod.*; *Asclepias stipitacea*, *Forsk. Arab. 50*; *Cynanchum pyrotechnicum*, and perhaps also *Asclepias aphylla*, of the same author.

EUSTEGIA.

CHAR. *Corolla* rotata.

Corona triplex; singula 5phylla; *extima* fauce inserta laciniis limbi opposita; *reliqua* extimâ alternantia, antheris opposita, foliolis *mediæ* tripartitis, *intimæ* indivisis.

Antheræ membranâ terminatæ.

Massæ pollinis apice attenuato affixæ, pendulæ.

Stigma submuticum.

Folliculi

HABITUS. *Herba* decumbens, humilis. *Folia* opposita, hastata. *Flores* subumbellati.

PATRIA. Africa Australis.

OBS. This remarkable character is taken from *Apocynum hastatum*, *Thunb. Prod.* 47; *A. minutum*, *Linn. Suppl.* 169, and from a very nearly related species found at the Cape of Good Hope, by Mr. Ferdinand Bauer.

METASTELMA.

CHAR. *Corolla* subcampanulata; *Fauce* coronatâ dentibus 5 exsertis, sinibus oppositis, tubum decurrentibus.

Corona staminea nulla.

Antheræ membranâ terminatæ.

Massæ pollinis compressæ, apice attenuato affixæ, pendulæ.

Stigma muticum.

Folliculi

HABITUS. *Planta* perennis, volubilis, glabra. *Folia* opposita, membranacea. *Umbellæ* interpetiolares, subsessiles. *Flores* parvi.

PATRIA. America Meridionalis.

OBS. This is the *Cynanchum parviflorum* of *Swartz*, whose description, however, of the corona, in *Flor. Ind. oc.* 1, p. 537, does not agree with ours, which was taken from excellent specimens in the Banksian Herbarium, collected in the islands of St. Croix and St. Christopher, *Masson* and *Von Rohr*.

MICROLOMA. [Ceropegia sp. *Linn. Jacq.* *Ceropegia* *Lam. Illust. Gen. tab.* 179.

CHAR. *Corolla* urceolata, tubo ventricoso, angulato, fauce nudâ, limbo brevior.

Squamæ 5, inclusæ, medio tubo sub sinubus insertæ, totidem fasciculis villorum alternantes.

Corona staminea nulla.

Antheræ membranâ terminatæ, sagittatæ.

Massæ pollinis compressæ, apice affixæ, pendulæ.

Stigma apiculatum.

Folliculi

HABITUS. *Suffrutices* volubiles. *Folia* opposita. *Umbellæ* interpetiolares.

PATRIA. Africa Australis.

OBS. This genus is formed from the two Cape species of *Ceropegia*, viz. *sagittata* and *tenuiflora*, plants which are widely different from the original species of that genus.

1. *M. sagittatum*, foliis sagittatis pubescentibus, corollæ limbo acutiusculo.

Ceropegia sagittata, Linn. mant.

2. *M. lineare*, foliis linearibus glabris, corollæ limbo obtusissimo.

Ceropegia tenuiflora, Linn.

64] ASTEPHANUS.

[*Apocyni* sp. Lin. fl. et Thun.

CHAR. *Corolla* subcampanulata, fauce tuboque esquamatis.

Corona staminea nulla.

Antheræ membranâ terminatæ.

Massæ pollinis pendulæ.

Stigma caudatum, v. muticum.

Folliculi

HABITUS. *Plantæ* perennes, sæpius volubiles. *Folia* opposita.

Umbellæ interpetiolares. *Flores* parvi.

PATRIA. Africa Australis.

OBS. This generic character is formed from *Apocynum triflorum* and *lineare*, Linn. suppl., and from two new species

in the Banksian collection. *Apocynum cordatum* and *lan-
ceolatum*, *Thunb. prod.*, probably likewise belong to the
genus; and I have modified the character, to admit a very
remarkable plant, found by Mr. Masson in South Africa,
the stem of which is shrubby with spinescent branches, the
leaves extremely minute, opposite, distant, and heartshaped;
the corolla rather urceolate than bellshaped; the orifice of
the tube furnished with deflected hairs; the masses of
pollen are fixed by their attenuated apices; the stigma is
blunt; the folliculi nearly cylindrical and smooth, with seeds
of the usual structure. The whole genus is evidently ^{as}
very near akin to *Microloma*, and differs chiefly in the want
of squamæ within the tube. They might, therefore, be
united; but this would lead to the junction also of *Meta-
stelma*, a native of the West India Islands, which I am
unwilling to join with these South African plants.

II.

Massæ pollinis 20, læves, quaternatim (duabus antheris
pertinentes), affixæ apici corpusculorum exsulcorum stig-
matis.

Filamenta connata, extus appendiculata.

SECAMONE.

[*Periplocæ* sp. *Linn.*

CHAR. *Corolla* rotata.

Corona staminea 5phylla.

Massæ pollinis erectæ quaternatim affixæ apici corpusculi
subsimilis, exsulci stigmatis.

Stigma apice coarctato.

Folliculi

HABITUS. *Frutices* erecti v. volubiles, glabri. *Folia*
opposita. *Cymæ* dichotomæ, interpetiolares. *Flores* minuti.

PATRIA. Africa, India Orientalis, et Nova Hollandia
tropica.

Obs. From the extreme minuteness of the parts, no genus

has been more difficult to determine than this: it is, however, perfectly natural and distinct, and is evidently the ^{56]} connecting link between the true Asclepiadeæ, and the Periploceæ, which follow. I have examined five species, of which two are published plants, viz. *Periploca secamone* Linn. and *emetica* of Retz. and Willdenow. The third is a climber, a native of India, discovered by Dr. Roxburgh; and two are erect shrubs, natives of the tropical part of New Holland.

III. PERIPLOCEÆ.

Massæ pollinis 5-20, granulosæ, (granulis e sphæculis 4 compositis), solitariæ, usque quaternatim apici dilatato corpusculi singuli stigmatis affixæ.

Filamenta partim v. omnino libera.

HEMIDESMUS.

[*Periploceæ* s^r *inn.*

CHAR. *Corolla* rotata, squamis 5, sub sinibus, obtusis.

Filamenta basi connata, supra distincta. *Antheræ* co-hærentes, a stigmati liberæ, imberbes, apice simplices.

Massæ pollinis 20.

Stigma muticum.

Folliculi cylindracei, divaricatissimi, læves. *Semina* comosa.

HABITUS. *Frutices* volubiles, glabri. *Folia* opposita, nitida. *Flores* interpetiolares, congesti, parvi.

PATRIA. India Orientalis.

^{57]} OBS. This genus, whose name is derived from the partial connection of the stamina, is composed of *Periploca Indica*, and two very nearly related unpublished species. Its strict affinity to *Periploca* is unquestionable, but the differences in the structure of its flower appear to me sufficient to justify its separation from that genus.

PERIPLOCA.

[Periplocæ sp. *Linn.*

CHAR. *Corolla* rotata. *Squamis* 5 faucis laciniis alternatibus, aristatis.

Filamenta distincta. *Antheræ* cohærentes, dorso barbatae.

Massæ pollinis apici dilatato corpusculi stigmatis applicitæ, solitariae, e quatuor confluentibus.

Stigma submuticum.

Folliculi cylindracei, divaricatissimi, læves. *Semina* comosa.

HABITUS. *Frutices* sæpius volubiles, glabri. *Folia* opposita, nitida. *Flores* subcorymbosi interpetiolares.

PATRIA. Europa Australis. Syria. Africa Septentrionalis et Æquinoctialis.

Obs. Periploca Græca, and lævigata, are the two principal species of the genus. *P. angustifolia*, *Billard.* no doubt also belongs to it; and I have seen a fourth and very remarkable species brought from Sierra Leone by ¹⁸³⁸ Professor Afzelius, on account of which I have been obliged slightly to modify the character.

GYMNANTHERA.

CHAR. *Corolla* hypocrateriformis. *Corona* faucis 5phylla.

Filamenta distincta, fauci inserta. *Antheræ* imberbes.

Massæ pollinis quaternatim applicitæ apici dilatato corpusculorum.

Stigma apiculo bifido.

Folliculi cylindracei, læves, divaricatissimi. *Semina* comosa.

HABITUS. *Frutex* volubilis, glaberrimus. *Folia* opposita, nitida. *Pedunculi* laterales sub-dichotomi. *Flores* albivirescentes, imberbes.

PATRIA. Nova Hollandia, intra tropicum.

(Table facing page 59.)

APOCINÆÆ.

SECT. I. SEMINA COMOSA.

I. SEMINA AD EXTREMITATEM UMBILICALEM (superiorem) COMOSA. Embryo planus.

A. PAUX COROLLÆ NUDA, nec coronata.

a. Tubus corollæ intus absque squamulis denticulatis.

† Stamina inclusa.

* Anthere medio stigmatē coherentes. Squamulæ 5 hypogynæ distinctæ, raro connatæ.
Corollæ lacinis dimidiatæ

ECHITES.

** Anthere a stigmatē solutæ.

1. Corpuscula 5 hypogyna, filiformia. Anthere sagittatæ, lobis posticis polline vacuis.

ICHCOCARPUS.

Corollæ lacinis falcatæ

2. Corpuscula nulla hypogyna. Anthere lanceolatæ, longitudinaliter polliniferæ.

HOLARRHENA.

Corollæ lacinis æquilateræ

†† Stamina exserta; medio stigmatē coherentes.

* Filamenta fauci inserta.

1. Filamenta apice simplici. Squamæ nullæ hypogynæ. Corolla hypocrateriformis,
limbo spartito

ISONEMA.

2. Filamenta apice extus gibbere carnosio. Squamæ 5 hypogynæ, basi connatæ.

VALLARIS.

Corolla subhypocrateriformis, limbo 5fido

** Filamenta medio v. juxta basin tubi inserta; filiformia, simplicia. Squamæ 5 hypogynæ,
distinctæ v. connatæ.

PARSONSIA.

1. Folliculi duo, placentis liberis

LYONSIA.

2. Capsula bilocularis, septo parallelo placentifero

PAROSSIA.
LYONSIA.

- distinctæ v. connatæ.
1. Folliculi duo, placentis liberis
2. Capsula bilocularis, septo parallelo placentifero

- b. Tubus corollæ intus squamis denticulise 5 inclusis. Stamina inclusa. Styli nulli. Squamæ 5 hypogynæ.
1. Corolla campanulata. Denticuli acuti tubi lacinis limbi oppositi
2. Corolla infundibuliformis. Squamulæ obtusæ tubi lacinis limbi alternantes

APOCYNUM.
CRYPTOLEPIS.

B. FAUX COROLLÆ CORONATA, squamis (tubulove) exsertis.

* Corona monophylla, cum vel absque squamis interioribus.

1. Corona duplex; *exterior* annularis integerrimus; *interior* e squamis 5 lacinis limbi alteruantibus. Urceolus hypogynus
2. Corona simplex, tubulosa, crenulata. Squamæ nullæ hypogynæ, sed 10 basi calycis extra corollam impositæ. Corollæ lacinis æquilateres. Antheræ mucrone brevissimo.

PRESTONIA.
BALFOURIA.

** Corona simplici serie polyphylla.

1. Coronæ foliola divisa. Squamæ nullæ hypogynæ. Corollæ lacinie ecaudatæ inæquilateres. Antheræ elongato-aristatæ
2. Coronæ foliola 10 indivisa. Squamæ 5 hypogynæ. Corollæ lacinie caudatæ

NERIUM.
STROPHANTHUS.

II. SEMINA AD EXTREMITATEM UMBILICO OBVERSAM (inferiorem) COMOSA. Embryo cotyledonibus involutis. Corolla fauce coronata

WRIGHTIA.

III. SEMINA PELTATA, CILIATA, UTRAQUE EXTREMITATE CILIIS ELONGATIS COMOSA. Corolla hypocroteriformis, fauce nuda. Stamina inclusa, antheris liberis. Squamæ 0 hypogynæ

ALSTONIA.

APOCINÆ.

SECT. I. SEMINA COMOSA.

A. *Coma ad extremitatem umbilicalem (superiorem) seminis.*

ECHITES.

[*Brown.* Echitides pleræq. *Jacq.*
Echitidis species, *Linn.*

CHAR. *Corolla* hypocrateriformis, fauce tuboque esquamatis. *Laciniis* limbi 5partiti inæquilateris.

Stamina inclusa. *Antheræ* sagittatæ, medio stigmati cohærentes, lobis posticis polline vacuis.

Ovaria 2. *Stylus* 1, filiformis.

Squamæ 5 hypogynæ.

Folliculi graciles.

HABITUS. *Frutices* volubiles. *Folia* opposita, ciliis interpetiolaribus glandulosis. *Pedunculi* interpetiolares, multiflori. *Flores* ut plurimum speciosi, albi, lutei et purpurei.

PATRIA. America Meridionalis.

Obs. Of Echites I have only examined *E. umbellata*, the original species when the species was established by Brown⁶⁰³ in his "History of Jamaica;" *biflora*, *circinalis*, and *Domingensis*: but, from the descriptions and figures of authors, especially of Jacquin, Swartz, and the authors of the "Flora Peruviana," I without hesitation refer to it the following species: *E. suberecta* *Jacq. repens* *Jacq. agglutinata* *Jacq. asperuginis* *Swartz*, *torulosa* *Jacq. acuminata* *Fl. Peruv. laxa* *Fl. Peruv. hirsuta* *Fl. Peruv. E. quinquangularis*, *Jacq.* and *annularis*, *Linn. Suppl.* are probably not genuine species, on account of the prominent

ring of the faux, and *E. glandulosa* *Fl. Peruv.* which according to the figure has a crown of 5 entire laciniaë and the segments of the corolla equal-sided, must be excluded from this genus. *E. siphilitica*, the specimen of which in the Linnean Herbarium I have seen but not sufficiently examined, is somewhat doubtful. *E. floribunda*, *corymbosa* and *spicata* are removed to another genus. The two remarkable species of South Africa, *E. bispinosa* and *succulenta*, require further examination; for their peculiar habit indicates their being a distinct genus from *Echites*, which it would be desirable to limit to the species of tropical America: hence it will be necessary to re-examine certain plants of India that in many respects agree with this genus, especially *Tsjeria-pupal-valli* of *Rheed, Mal. 7, p. 103, t. 55*, which appears to differ from *Echites* chiefly in having a calyx longer than the tube of the corolla, in the scales surrounding the ovarium being united, (which, however, is also the case in *E. Domingensis*,) and in the greater quantity and density of the albumen, which is between fleshy and cartilaginous.

The authors of the "Flora Peruviana" have reformed the character of *Echites*, but the scales which they describe between the calyx and corolla will certainly not materially assist in distinguishing this genus from those most nearly related to it; and I have observed a nearly similar structure in most of the genera of this family, as well as of *Asclepiadæ*: these scales, however, truly belong to the calyx, and are either five or ten in number, or more rarely consist of an uninterrupted series of ciliaë, not unlike those which so frequently occur within the footstalks of the leaves. My observations are not sufficiently numerous to enable me to determine whether their modifications might not generally assist in characterizing genera, and I have, therefore, very seldom had recourse to them.

ICHNOCARPUS.

CHAR. *Corolla* hypocrateriformis, limbi laciniis dimidiatis, fauce tuboque esquamatis.

Stamina inclusa. *Antheræ* sagittatæ, a stigmatē liberæ.
Ovaria 2. *Stylus* 1, filiformis. *Stigma* ovatum, acuminatum.

^{62]} *Filamenta* 5, hypogyna, staminibus alternantia.

Folliculi graciles.

HABITUS. *Frutex* oppositifolius. *Panicula* terminalis, brachiata. *Flores* parvi.

PATRIA. India Orientalis et Zeylona.

OBS. This is the *Apocynum frutescens* Linn., of which I have examined the original specimen in Hermann's herbarium. The *Quirivelia Zeylanica* of Lamarck and Poiret, in *Encycl. method. botan. vol. 6, p. 42*, considered by them as Linnæus's plant, must, from the description, be widely different, and probably does not belong to the same natural family.

HOLARRHENA.

CHAR. *Corolla* hypocrateriformis, laciniis æquilateris, fauce tuboque esquamatis.

Stamina inclusa, imo tubo inserta. *Antheræ* a stigmatē liberæ, lanceolatæ, integræ, longitudinaliter polliniferæ.

Ovaria 2. *Stylus* brevissimus. *Stigma* cylindraceum.

Squamæ nullæ hypogynæ.

Folliculi graciles.

HABITUS. *Frutices* erecti, glabri. *Folia* membranacea. *Cymæ* terminales et laterales.

PATRIA. India Orientalis et Zeylona.

OBS. This genus consists of two species; one of which is ^{63]} *Carissa mitis*, Vahl, *ymb.* 3, p. 44: a specimen of this, so named by Koenig, I have examined in the Banksian Collection.

ISONEMA.

CHAR. *Corolla* hypocrateriformis; fauce tuboque esquamatis; limbo 5partito.

Stamina exserta. *Filamenta* fauci inserta, apice simplicia. *Antheræ* sagittatæ, medio stigmati cohærentes.

Ovaria 2. *Stylus* 1, filiformis. *Stigma* incrassatum, obtusum.

Squamæ nullæ hypogynæ.

Folliculi

HABITUS. *Frutex* erectus? oppositifolius, pilosus. *Panicula* terminalis, brachiata, floribus corymbosis. *Calycis* foliola basi intus squamâ duplici. *Corollæ* (semuncialis) tubus cylindræus, intus medio barbatus.

PATRIA. Africa Æquinoctialis. (*H. Smeathman*, in *Herb. Banks.*)

VALLARIS. [*Burm. Ind.* 51. *Pergulariæ* sp. *Linn.*

CHAR. *Corolla* hypocrateriformis; fauce tuboque esquamatis, limbo 5fido obtuso.

Stamina exserta. *Filamenta* fauci inserta, brevissima, apice extus gibbere carnosum. *Antheræ* sagittatæ, medio stigmati cohærentes.

Ovarium biloculare. *Stylus* filiformis. *Stigma* conico-ovatum.

Squamæ 5, hypogynæ, basi connatæ, apicibus ciliatis. 164

Folliculi

HABITUS. *Frutex* volubilis, oppositifolius. *Pedunculi* interpetiolares, dichotomi. *Flores* corymbosi, suaveolentes.

PATRIA. India Orientalis.

Obs. This is the *Flos Pergulanus* of Rumphius, which Linnæus considered as the first species of his genus *Pergularia*: it does not, however, belong to the same order with the plant that afforded his generic character, and to which the name has been since generally applied.

PARSONSIA. [*Echitidis* sp. *Jacq. et Swartz.*

CHAR. *Corolla* infundibuliformis, fauce tuboque esquamatis, limbo 5partito, recurvo, laciniis æquilateris.

Stamina exserta. *Filamenta* medio v. juxta basin tubi inserta, filiformia. *Antheræ* sagittatæ, medio stigmati cohærentes, lobis posticis polline destitutis.

Ovaria 2, v. 1, biloculare. *Stylus* 1. *Stigma* dilatatum.

Squamæ 5, hypogynæ, distinctæ, v. connatæ.

Folliculi 2, distincti, v. cohærentes.

HABITUS. *Frutices* volubiles. *Folia* opposita. *Flores* ^{65]} cymosi, v. racemosi, racemis sæpe compositis, terminales v. interpetiolares, parvi, colore in variis vario.

PATRIA. America Meridionalis et Australasia.

Obs. The American species of this genus, viz. *Echites corymbosa* Jacq. floribunda Sw. and spicata Jacq. differ considerably from the rest, which are natives of New Holland and New Zealand. Among these, the only published species is *Periploca capsularis*, Forst. prod. n. 126. When the fruit of all has been examined, they will probably be divided in the following manner:

† *Americanæ*. Ovariis duobus. Folliculis distinctis.

†† *Australasienses*. Ovario biloculari. Folliculis longitudinaliter cohærentibus.

The genus is named in memory of Dr. JAMES PARSONS, the author of a Dissertation on the Analogy between the Propagation of Animals and that of Vegetables, and of an unfinished work, entitled, "The Microscopical Theatre of Seeds." The *Parsonsia* of Brown, which Linnæus reduced to *Lythrum*, is a species of *Cuphea*.

^{66]} LYONSIA.

CHAR. *Corolla* infundibuliformis, fauce tuboque esquamatis, limbo 5partito, recurvo, laciniis æquilateris: æstivatione valvata.

Stamina exserta. *Filamenta* medio tubo inserta, filiformia. *Antheræ* sagittatæ, medio stigmati cohærentes, lobis posticis polline vacuis.

Ovarium biloculare. *Stylus* 1, filiformis, apice dilatato.

Stigma subconicum.

Squamæ 5, hypogynæ, connatæ.

Capsula cylindracea, bilocularis, valvis folliculiformibus, dissepimento parallelo libero utrinque seminifero.

HABITUS. *Frutex* volubilis. *Folia* opposita. *Cymæ* terminales, trichotomæ. *Flores* inter minores, limbo barbato.

PATRIA. Nova Hollandia.

Obs. This genus is perhaps too nearly related to the New Holland portion of *Parsonsia*, from which it differs chiefly in its fruit being capsular. It is named in memory of ISRAEL LYONS, author of "*Fasciculus Plantarum circa Cantabrigiam nascentium*," and from whom Sir Joseph Banks received his earliest instructions in botany.

APOCYNUM.

[*Apocyni* sp. *Linn. et Juss.* 167

CHAR. *Corolla* campanulata. *Tubus* denticulis 5, acutis, inclusis, laciniis limbi oppositis. *Faux* nuda.

Stamina inclusa. *Antheræ* sagittatæ, medio stigmati cohærentes, lobis posticis polline vacuis.

Ovaria 2. *Styli* subnulli. *Stigma* dilatatum, apice conico.

Squamæ 5, hypogynæ.

Folliculi graciles, distincti.

HABITUS. *Herbæ* perennes, erectæ. *Folia* opposita, membranacea. *Flores* cymosi.

PATRIA. America Borealis, et Europa Australis.

Obs. The Linnean genus *Apocynum* at present contains many plants widely different from the species of which it originally consisted. Most of the spurious species I have already referred to different genera, and I here add specific characters of all the genuine species with which I am acquainted.

1. *A. androsæmifolium*, foliis ovatis glabris, cymis terminalibus lateralibusque, tubo corollæ calycem his superante.

Apocynum androsæmifolium, *Linn.*

68] 2. *A. cannabinum*, foliis lanceolatis utrinque acutis glabris, cymis paniculatis, calyce tubum corollæ æquante.
Apocynum cannabinum, Linn.

3. *A. hypericifolium*, foliis oblongis glabris brevissimè petiolatis mucronatis: basi obtusa subcordata, cymis folio brevioribus, calyce tubum corollæ æquante.
Apocynum hypericifolium, Hort. Kew.

4. *A. pubescens*, foliis ovato-oblongis mucronatis; basi obtusis; utrinque cymaque brevioribus pubescentibus, calyce corollam subæquante.

HAB. In Virginia. Mitchell, in Herb. Banks. (ubi v. s.)

5. *A. Sibiricum*, foliis ovato-oblongis mucronatis glabris; basi obtusa, cymis terminalibus pedunculatis pulvereo-pubescentibus, tubo corollæ calycem superante.

HAB. In salsis desertorum Astrachanensium. P. S. Pallas, M.D., in Herb. Banks. (ubi v. s.)

6. *A. Venetum*, foliis oblongo-ellipticis glabris mucronatis; basi subobtusis, cymis paniculatis lateralibus terminalibusque, calyce tubum corollæ æquante.

Apocynum venetum, Linn.

69] CRYPTOLEPIS.

CHAR. *Corolla* infundibuliformis. *Tubus* squamulis 5, obtusis, inclusis, laciniis limbi alternantibus. *Faux* nuda. *Stamina* inclusa, imo tubo inserta. *Antheræ* sagittatæ. *Ovaria* 2. *Styli* 0. *Stigma* dilatatum, apiculo conico. *Squamulæ* 5, hypogynæ. *Folliculi*

HABITUS. *Frutex* volubilis. *Folia* opposita, paginis discoloribus, inferiore venosa. *Corymbi* interpetiolares subsessiles brevissimi.

PATRIA. India Orientalis. (Franciscus Buchanan, M.D., in Herb. Banks.)

PRESTONIA.

CHAR. *Corolla* hypocrateriformis. *Faux* coronata *tubulo* annulari indiviso *squamisque* 5, interioribus laciniis limbi alternantibus.

Antheræ semi-exsertæ, sagittatæ, medio stigmati cohærentes, lobis posticis polline vacuis.

Ovaria 2. *Stylus* 1, filiformis, apice dilatato. *Stigma* turbinatum apiculo angustiore.

Urceolus hypogynus, 1-phyllus.

Folliculi

HABITUS. *Frutex* volubilis, tomentosus. *Folia* opposita, tomentosa. *Corymbi* congesti interpetiolares. *Calyx* 7^o foliaceus, laciniis basi intus squamulâ instructis.

PATRIA. America Meridionalis.

P. tomentosa.

HAB. In sepibus prope Rio de Janeiro, in Brasilia, detexit *Josephus Banks, Baronettus*.

OBS. This genus is named in memory of Dr. CHARLES PRESTON, the correspondent of Ray, and styled by him, "eruditissimus vir, et curiosissimus stirpium observator." Many of his observations, chiefly on the more minute plants, occur in Ray's "Methodus emendatæ." Blair also mentions him as an ingenious and expert botanist.

BALFOURIA.

CHAR. *Corolla* infundibuliformis. *Faux* coronata *tubulo* crenulato. Limbi laciniæ rectæ, æquilateres.

Stamina semi-exserta, faucibus inserta. *Antheræ* sagittatæ, medio stigmati cohærentes, mucronatæ.

Ovarium biloculare. *Stylus* 1, filiformis, apice dilatato.

Stigma angulatum.

Squamulæ 10, basi calycis extra corollam insertæ: hypogynæ nullæ.

Folliculi

HABITUS. *Arbor* inter minores. *Folia* opposita, lanceo-

71) lato-linearia, falcata, denticulis interpetiolaribus. *Cypripæ* trifidæ, laterales et terminales.

PATRIA. Nova Hollandia tropica.

Obs. I have named this genus in memory of Sir ANDREW BALFOUR, the founder of the Edinburgh Botanic Garden and Museum, of whose merits in natural history, especially in botany, an interesting account is given by his friend Sir Robert Sibbald, in a small volume, entitled, "Memoria Balfouriana."

NERIUM.

[*Nerii* sp. *Linn. et Juss.*

CHAR. *Corolla* hypocrateriformis. *Faux* coronata foliolis lacero-multifidis. Læmbs laciniis tortis, inæqualiteris, ecaudatis.

Stamina. *Filamenta* medio tubo inserta. *Antheræ* sagittatæ aristatæ, medio stigmati cohærentes.

Ovaria 2. *Stylus* 1, filiformis, apice dilatato. *Stigma* obtusum.

Squamæ nullæ hypogynæ denticuli in basi calycis, extra corollam.

Folliculi cylindracei.

HABITUS. *Frutices* erecti. *Folia* terna, elongata, coriacea, venis numerosis, parallelis.

PATRIA. India Orientalis.

72) Obs. The only true species of *Nerium* are, *N. Oleander*, *odorum*, and probably *salicinum*, *Forsk.*

Nerium Zeylanicum and antidysentericum, form a very distinct genus, which I have named *Wrightia*. *N. coronarium* is probably a *Tabernæmontana*, and *N. divaricatum*, from an examination of the specimen in Hermann's Herbarium, on which this species entirely rests, I believe to be the same plant. *Nerium obesum*, *Forsk.*, seems to be *sui generis*: it cannot, at least, be a *Nerium*.

STROPHANTHUS, *Decandolle*. [Echitidis sp. *Linn.*

CHAR. *Corolla* infundibuliformis. *Faux* coronata squamulis decem, indivisis. Limbi laciniæ caudatæ.

Stamina medio tubo inserta. *Antheræ* sagittatæ, aristatæ, v. mucronatæ.

Ovaria 2. *Stylus* 1, filiformis, apice dilatato. *Stigma* subcylindraceum.

Squamæ 5, hypogynæ.

Folliculi

HABITUS. *Frutices* sarmentosi. *Folia* opposita.

PATRIA. Africa Æquinoctialis, et India Orientalis.

B. *Coma ad extremitatem umbilico obversam (inferiorem) 173 seminis.*

WRIGHTIA. [Nerii sp. *Linn.*

CHAR. *Corolla* hypocrateriformis. *Faux* coronata squamis decem, divisis.

Stamina exserta. *Filamenta* faucis inserta. *Antheræ* sagittatæ, medio stigmati cohærentes.

Ovaria 2, cohærentia. *Stylus* 1, filiformis, apice dilatato. *Stigma* angustius.

Squamæ 5-10, basi calycis extra corollam insertæ.

Folliculi distincti, v. cohærentes, placentis adnatis.

HABITUS. *Frutices* erecti, arboresce minores. *Folia* opposita. *Corymbi* subterminales. *Flores* albi. *Albumen* 0. *Embryo* cotyledonibus longitudinaliter involutis, albus, aquâ calidâ immersus roseus evadit!

PATRIA. India Orientalis, Zeylona, Archipelago Malaica, et Nova Hollandia tropica.

OBS. Gærtner has given an excellent account of the fruit of this genus, in his description of *Nerium Zeylanicum*, and he no doubt supposed, that the fruit of *Nerium Oleander* was essentially the same. It is, however, very remarkably different. And no genus is more distinct in habit, or more beautifully characterized than this, which I

have dedicated to my much respected friend, WILLIAM 74] WRIGHT, M.D., F.R.S.L. & E., whose ardour in the pursuit of botanical knowledge, even while engaged in extensive medical practice, in the island of Jamaica, has long entitled him to this mark of distinction.

1. *W. antidysenterica*, foliis obovato-oblongis breviter acuminatis glabris, corymbis subterminalibus, tubo corollæ calyce 6ies longiore, folliculis distinctis.

Nerium foliis ovatis acuminatis petiolatis, *Linn. Flor. Zeyl.* 107, fide speciminis in herbario Hermanni.

Nerium antidysentericum, *Linn. sp. pl. ed. 2, p.* 306.

HAB. In Zeylona. *Hermann*, et *J. G. Kænig*, in *Herb. Banks.*

Obs. Codaga pala, *Rheed. mal.* 1, *p.* 85, *t.* 47, vix hujus loci, præsertim ob diversam figuram foliorum, et coronæ defectum; ideoque forsân Holarrhenæ species.

2. *W. Zeylanica*, foliis oblongo-lanceolatis subacuminatis glabris, corymbis terminalibus, tubo corollæ calyce 4-5ies longiore, folliculis distinctis.

Nerium Zeylonicum, *Linn. Amæn. Acad.* 4, *p.* 309*.

HAB. In Zeylona. Exemplar visum ex Herbario fuit *D. Van Royen*, nunc in Museo Banksiano.

Obs. Præcedenti nimis affinis, diversa præsertim figura foliorum, quæ etiam paulo minora sunt.

3. *W. tinctoria*, foliis elliptico-lanceolatis ovatisque acuminatis glabris, ramis corymbisque divaricatis, corollæ tubo calyce duplo longiore, folliculis distinctis.

HAB. In India Orientali. *J. G. Kænig*, *M.D.*, et *Gul. Roxburgh*, *M.D.*, in *Herb. Banks.* (ubi v. s.)

4. *W. pubescens*, foliis elliptico-oblongis acuminatis calycibusque pubescentibus, corymbis erectis, tubo corollæ calyce parum longiore, folliculis cohærentibus.

HAB. In Novæ Hollandiæ orâ septentrionali Arnheims Land, et in insula Timor prope Coepang, (ubi v. v.)

C. Semina peltata, ciliata, utriusque extremitatis ciliis elongatis comosa.

ALSTONIA.

[Echitidis sp. Linn.]

CHAR. *Corolla* hypocrateriformis, fauce tuboque esquamatis.

Stamina inclusa. *Antheræ* lanceolatae, longitudinaliter polliniferae, a stigmatibus liberae.

Ovaria 2. *Stylus* 1, filiformis, apice dilatato. *Stigma* subconicum.

Squamulae nullae hypogynae, nec calycinae.

Folliculi teretes.

HABITUS. *Arbores* saepe proceræ, lactescentes.

Folia verticillata vel opposita, costata, glabra.

Cymæ terminales, paniculatae. *Flores* saepius albi.

Folliculi plerumque longissimi.

PATRIA. India Orientalis, Archipelago Malaica, et 176 Insulae Societatis.

Obs. Had Linnæus examined the fruit of this genus, or even attended to the figure given of it in the "Hortus Malabaricus" (of which work, it is proper to observe, he had not a copy), it is probable he would have distinguished it from Echites, to which it has so little affinity. As it is, I am happy in having it in my power to commemorate the merits of Dr. ALSTON, the predecessor of Dr. Hope, by so distinct and splendid a genus. The *Alstonia* of the Younger Linnæus, is, according to L'Heritier, a species of *Symplocos*.

1. *A. scholaris*, foliis verticillatis 5-7 obovato-oblongis obtusis costatis venaque margini approximata cinctis, cymis breviter pedunculatis, limbo corollae parum barbatis, folliculis longissimis.

Pala, *Rheed. mal.* 1, p. 81, t. 45, optima.

Lignum scholare, *Rumph. amb.* 2, p. 246, t. 82, quoad descriptionem, sed figura potius sequentis.

Echites scholaris, *Linn. mant.* 53.

HAB. In India Orientali et in Insulis Moluccanis. (v. s. in Herb. Banks.)

2. *A. spectabilis*, foliis quaternis elliptico-oblongis sub-
773 acuminatis costatis: margine simplicibus, cymis pedunculatis folio brevioribus, corolla limbo barbato, folliculis longissimis.

HAB. In Insula Timor prope Coepang, cum floribus fructibusque Aprili 1803 observavi.

Obs. Præcedenti quàm maximè affinis, sed reverà distincta, nec malè ab icone cit. Rumphii repræsentata; descriptio autem *A. scholari* melius convenit.

3. *A. venenata*, foliis quaternis oblongo-lanceolatis acuminatis basi attenuatis, cymis dichotomis, corollæ tubo sursum ampliato; limbo imberbi acuto, folliculis utrinque attenuatis folium vix æquantibus.

HAB. In India Orientali, *Gul. Roxburgh, M.D.* (v. s. in Herb. Banks.)

4. *A. costata*, foliis oppositis elliptico-oblongis acuminatis costatis, cymis effusis, limbi laciniis imberbibus lanceolatis tubo longioribus, folliculis longissimis.

Echites costata, *Forst. prod. n.* 123.

HAB. In Insulis Otaheite et Ulaietea, inter juga montium (insularibus Attahé nuncupata). *Josephus Banks, Baronettus.* (v. s. in Herb. Banks.)

Obs. Kametti-valli, *Rheed. mal.* 9, p. 23, t. 14, which is given in Willdenow's *sp. pl.* 1, p. 1240, as a synonyme of this plant, differs in the shortness of the folliculi, and in
783 having winged and naked seeds. Rheede also describes his plant as being a climber. The seeds of the Otaheite plant, which forms a moderate-sized tree, are distinctly

ciliated ; but I am not certain that the ciliæ are so remarkably elongated at each extremity, as in the other species ; and it may, therefore, be a connecting link between this section of Apocineæ, in which I have placed it, and the nearly related genera *Plumeria*, *Cameraria*, and *Vinca*.

ON
WOODSIA,
A NEW GENUS OF FERNS.

BY
ROBERT BROWN, Esq., F.R.S., LIB. LINN. SOC.

(READ NOVEMBER 7TH, 1812.)

[*Extracted from 'The Transactions of the Linnean Society of London,'*
Vol. XI, pp. 170—174.]

ON WOODSIA,

[170

A NEW GENUS OF FERNS.

THERE is perhaps no tribe of cryptogamous plants which since the time of Linnæus has received greater additions to its number of species, or more considerable improvements in its systematic arrangement, than the *Filices*; and certainly no botanist has so essentially contributed to these improvements as the President of this Society; whose ingenious Essay on Dorsiferous Ferns may justly be considered as the groundwork of the more complete dissertations of Professors Swartz and Bernhardi, which have appeared since its publication.¹

Linnæus, in his latest work, the 13th edition of the *Systema Vegetabilium*, enumerates scarcely more than 200 Ferns, which he referred to twelve genera: while the *Species Plantarum* of the late Professor Willdenow contains upwards of a thousand plants of the same order, arranged under forty-three genera. It is however remarkable, that of this vast number of species nearly one half belong to four of the Linnean genera, namely, *Polypodium*, *Acrostichum*, *Asplenium*, and *Pteris*, all of which were first proposed by Ray in his *Methodus Plantarum Emendata*, published in 1703; without names, indeed, but with characters nearly similar to those of Linnæus.

It appears, therefore, that the arrangement of Ferns at present universally followed is not wholly new: and that it has not attained such a degree of perfection as to super-

¹ An. 1793, in *Mém. de l'Académie Royale des Sciences de Turin*, vol. v, p. 401.

sede all changes in nomenclature, may be inferred from the genus *Polypodium* alone, though reduced nearly one half by its present character, still including 157 species, or upwards of a seventh part of the whole order.

The expediency of subdividing *Polypodium*, as well as some of the other genera mentioned, especially *Acrostichum*, is indeed obvious, not merely on account of their great extent, but also from the striking differences in habit existing among the species referred to each.

I have, some time ago,¹ had an opportunity of remarking, that the plants referred to *Polypodium*, *P. ilvense* and *hyperboreum*, form a distinct genus, from the peculiar structure of their involucre, even the existence of which had escaped preceding observers.

This genus I have named in honour of my friend Mr. Joseph Woods, whose merits as an accurate and skilful English botanist are well known to many of the members of this Society: and the object of the present communication is to illustrate it by some additional observations on its structure, and by a very perfect drawing, for which I am indebted to the friendship of Mr. Francis Bauer.

The character distinguishing *Woodsia* from all other genera of Ferns hitherto established, consists in its involucre being inserted under the group of capsules, or, as it is technically called, the *sorus*, which it completely surrounds at the base; while it is in every stage open at the top, having its margin divided into a number of capillary segments, which from their length and incurvation entirely conceal the young capsules, and in a great measure the full-grown.

That so singular a structure should have been hitherto unnoticed, even though both species of the genus have been described and figured since the publication of Dr. Smith's memoir, is not perhaps to be wondered at: for the membranaceous base of the involucre is completely concealed by the capsules, and the marginal hairs, which alone are visible, exactly resembling the pubescence of the frond, have been universally confounded with it.

¹ *Prodr. Fl. Nov. Holl.* i, p. 158, Obs. IV.

The difficulty, too, of separating the membrane entire from the frond, to which, by the pressure of the capsules, it is closely applied, is so considerable, that, since the publication of my remark already quoted, its existence has been doubted by a botanist, whose opinion, especially in whatever regards this order of plants, is of peculiar weight, and in opposition to which I should not retain full confidence in my own observations, though frequently repeated, were they not so distinctly confirmed by Mr. Bauer's excellent drawing.

I first observed the involucre six years ago in living plants of *Woodsia hyperborea*, and have since repeatedly ascertained its existence in dried specimens of the same species, and of *Woodsia ilvensis*. These two plants are indeed so nearly related, that I find myself unable to construct for them clear specific characters; and therefore, in proposing them here as distinct species, I am, from want of sufficient materials to determine the question, rather following the prevailing opinion than my own.

To the characters and synonyms which follow, I have not thought it necessary to add descriptions of the two supposed species, these having been given by several of the authors referred to, and in every respect correctly, except what regards the involucre.

WOODSIA.

[173]

Sori dorsales, subrotundi.

Involucre calyciforme apertum margine crinitum: includens *Capsulas* pedicellatas: receptaculo communi elevato nullo.

Filiculæ, frondibus cæspitosis, pinnatim divisis; pilis simplicibus squamulisque angustis instructæ.

1. *ilvensis*. W. frondibus bipinnatifidis, pinnis oblongis, pinnulis confluentibus multifloris: inferioribus subrepandis: infimis subæqualibus.

Polypodium ilvense, Swartz, *Synop. Fil.* 39. Willden. *Sp. Pl.* 5, p. 198. Schkuhr, *Crypt.* 16, t. 19.

Acrostichum ilvense. Linn. *Sp. Pl. ed. 2, p. 1528.*

Nephrodium lanosum. Michaux, *Amer. 2, p. 198.*

Habitat in rupibus Europæ et Americæ borealis. (v. v.)

2.

2. *hyperborea.* W. frondibus pinnatis, pinnis triangularibus oblongisve inciso-pinnatifidis: lobis integerrimis paucifloris: antico baseos productiore. *Tab. 38 (XI).*

Polypodium hyperboreum. Swartz, *Synop. Fil. 39. Willden. Sp. Pl. 5, p. 197. Eng. Bot. 2023.*

Polypodium arvenicum. Smith, *Fl. Brit. 3, p. 1115.**

Polypodium ilvense. Withering, *Arrang. ed. 3, t. 3, p. 774.*

Acrostichum hyperboreum. Liljeblad in *Act. Stockholm. 1793, p. 201, t. 8*.*

Acrostichum alpinum. Bolton, *Fil. Brit. 76, t. 42.*

Ceterach alpinum. Lamarck et Decandolle, *Fl. Fran. 2, p. 567.*

Habitat in Europæ alpibus (v. v.) 2.

EXPLANATION OF TAB. 38 (XI).

[174]

FIG.

1. A native specimen of *Woodsia hyperborea*, natural size.
2. The stipes and lower part of the frond of the same plant, magnified 3 times in diameter.
3. A pinna of the same plant, magnified 10 diam.
4. A pinna from another specimen, in which the clusters of capsules (sori) are more numerous and confluent, 10 diam.
5. A single cluster of capsules within their involucre, the membranaceous base of which they entirely conceal, magnified 50 diam. (2500 times in superficies).
6. The involucre spread open, with only one capsule left in it, magnified 50 diam.
7. An unripe capsule.
- 8, 9. Side and back views of a ripe capsule.
- 10, 11. Capsule opening and entirely burst, shedding its seeds.
12. A seed magnified 200 diam.
13. A frond of a cultivated plant of the same species, natural size.

OBSERVATIONS
ON
THE NATURAL FAMILY OF PLANTS
CALLED
COMPOSITÆ.

BY
ROBERT BROWN, F.R.S.,
CORRESPONDING MEMBER OF THE ROYAL INSTITUTE OF FRANCE, AND OF THE
ROYAL ACADEMIES OF SCIENCES OF BERLIN AND MUNICH;
MEMBER OF THE WERNERIAN SOCIETY OF
EDINBURGH, LIBB. LINN. SOC.

(READ FEB. 6TH AND 20TH, 1816.)

[*Extracted from 'The Transactions of the Linnean Society of London,'*
Vol. XII, pp. 76—142.]

LONDON.

1817.

OBSERVATIONS
ON
COMPOSITÆ.

THE class *Syngenesia* of the Linnean artificial system, [76] as at present limited, constitutes a family strictly natural, and by far the most extensive in the vegetable kingdom. It is also, with the exception of Grasses only, the most generally diffused, and is almost equally remarkable with that order, for the great apparent uniformity in the structure of its essential parts of fructification.

This class of plants, for which I retain the established name COMPOSITÆ, in preference to any of those recently proposed, has lately become the subject of a minute and accurate examination by Mons. Henri Cassini; two of whose Memoirs on the Style and Stamina of the class, already published in the *Journal de Physique*,¹ are in my opinion models for botanical investigation.

A few years before the publication of M. Cassini's Memoirs on Compositæ I was induced to examine a considerable part of this extensive family, chiefly with a view to the more accurate determination of the New Holland plants belonging to it.

My principal object in the present paper is to communicate such general observations, the results of this investigation, as either have not yet been published by M. Cassini, or respecting which I consider myself to have anticipated that author in my General Remarks on the Botany of New

¹ Of 1813 and 1814.

77) Holland, appended to Captain Flinders's Voyage to Terra Australis.

To these observations I shall add some remarks on certain genera of Compositæ, which occur repeatedly under different names in late systematic works, and whose structure and limits appear to be imperfectly understood.

My *first* observation relates to the peculiar disposition of the nerves or vessels of the corolla of this family of plants.

In the essay already mentioned, which appeared early in the summer of 1814, I have noticed this peculiarity in the following terms:¹

"The whole of Compositæ agree in two remarkable points of structure of their corolla; which, taken together at least, materially assist in determining the limits of the class. The first of these is its valvular æstivation; this, however, it has in common with several other families. The second I believe to be peculiar to the class, and hitherto unnoticed. It consists in the disposition of its fasciculi of vessels or nerves; these, which at their origin are generally equal in number to the divisions of the corolla, instead of being placed opposite to these divisions, and passing through their axes, as in other plants, alternate with them; each of the vessels at the top of the tube dividing into two equal branches, running parallel to and near the margins of the corresponding lacinia, within whose apices they unite. These, as they exist in the whole class and are in great part of it the only vessels observable, may be called primary. In several genera, however, other vessels occur, alternating with the primary, and occupying the axes of the lacinia: in some cases these secondary vessels being most distinctly visible in the lacinia, and becoming gradually fainter as they descend the tube, might be regarded as recurrent; originating from the united apices of the primary
78) branches; but in other cases, where they are equally distinct at the base of the tube, this supposition can hardly be admitted. A monopetalous corolla, not splitting at the

[¹ Vol. i, p. 30.]

base, is necessarily connected with this structure, which seems also peculiarly well adapted to the dense inflorescence of Compositæ; the vessels of the corolla and stamina being united and so disposed as to be least liable to suffer by pressure."

At the date of this publication I certainly had no knowledge of any similar observations having been previously made: but I now see in M. Cuvier's account of the proceedings of the Institute of France for 1815, that M. Cassini is considered to have anticipated me on this subject, and as he says in "termes non équivoques." What these terms are, appears by a letter I have received from M. Cassini himself, in which he states his claim to rest on the following passage:

"Chaque fleur hermaphrodite ou male contient cinq étamines, correspondant aux cinq nervures de la Corolle et par conséquent alternes avec ses lobes."

This passage occurs in a Memoir on the Stamina of Compositæ, which was read to the Institute of France in July 1813, and first appeared with the substance of that Memoir in the Journal de Physique, said to be for April 1814; but the actual date of the publication of which I have reason to believe was somewhat later, and very nearly corresponding with that at which M. de Jussieu was in possession of a copy of my essay containing the observations already quoted. I conclude it is not supposed I could have been acquainted with the passage in the original memoir, unless the report usually made on memoirs read to the Institute should have been printed, and should have actually noticed this passage, or the discovery it is now said to contain.

But independently of the near equality of dates, I cannot consider my observations as either wholly or even in any considerable degree anticipated by the passage in question. My observations notice not only the disposition of the five vessels in the tube of the corolla, but their ramification in the laciniæ, by no means a necessary consequence of that disposition; they notice also the existence, in several genera of Compositæ, of five vessels alternating

with those, and which I considered secondary in this order, though they occupy the place of the primary vessels in other families: and it is this inverted disposition, indicated in the greater part of the class by the primary being the only vessels existing, which I have considered as of material importance in determining the limits of *Compositæ*, though by no means as affording an essential practical character for the whole class.

In the passage quoted from M. Cassini (the only one I can find relative to the subject in the memoir in which it occurs), the existence of five nerves or vessels in the tube of the corolla, alternating with its laciniae, is stated, but their division and disposition in the laciniae are not noticed; it is at the same time to be inferred from the terms of the passage, that no other vessels exist in the tube of the corolla: and it is equally evident that, so far from announcing this disposition of vessels as a discovery, or peculiar to the order, the author rather considers it either as a fact already known, or as the usual structure. That M. Cassini was not then aware of the importance of the fact which he had imperfectly stated, appears likewise from his having, many months after his memoir was read, and at a time when he says he had finished his analysis of the corolla, proposed a name for the class, taken from a supposed peculiarity in the structure of the filament, a name which he is now inclined to abandon for one derived from the disposition of vessels in the corolla.

^{80]} Since my attention has been again turned to the subject, I have endeavoured to collect all that has been observed on the nerves or vessels of the corolla of *Compositæ*, a brief account of which may be not altogether without interest.

The earliest notice I have been able to find is contained in a passage (in page 170) of Grew's *Anatomy of Plants*, where, in speaking of syngenesious flosculi, he says, "they are frequently ridged, or as it were hem'd like the edge of a band." And his figure of a magnified floret of the common Marigold, in tab. 61, gives a tolerable idea of the marginal vessels of its laciniae. Grew however takes no

notice of the trunks from which these branches arise, either in his text or plates.

Van Berkhey, in his Dissertation on Compositæ, published at Leyden in 1760, though he makes no mention of the nerves of the corolla in his text, yet in all the magnified figures he has given both of ligulate and tubular florets, correctly represents the trunks of the primary vessels, without, however, noticing their ramification in the lacinia. I am anticipated therefore by this author's figures exactly in the same degree as by the passage contained in M. Cassini's second memoir.

The accurate Schmidel, in the few Compositæ which occur in his Icones, has correctly represented the trunks of the primary vessels, but has equally omitted their ramifications.

In the Analysis Florum of Batsch, a work published in 1790, the object of which was to give an idea of the structure of the natural families of plants, by a minute description and magnified figures of one or more species selected from each, *Coreopsis tripteris* occurs; and although the vessels of its tubular floret are very indistinctly figured, yet both their trunks and branches are correctly described. The same author however, who in 1802 published an ingenious work on the natural families of plants,¹ takes no notice of the vessels of the corolla in the character of Compositæ which he has there proposed.

In the figures of syngenesious plants given by Schkuhr,² wherever the ligulæ of *Cichoraceæ* are magnified, the trunks of the nerves are correctly represented ending in the sinuses; unless in one plate containing *Lactuca virosa* and *Sonchus sibiricus*, in both of which the vessels are made to pass through the axes of the teeth; but in no case are the marginal branches noticed. It is singular that this generally accurate author, in the many magnified figures he has given of tubular florets, has only in two cases represented the trunks of their vessels, namely in *Echinops Ritro*, where they are correctly placed, and in *Silphium trifoliatum*.

¹ Tabulæ affinitatum regni vegetabilis.

² In Botanisches Handbuch

where, though only five vessels are visible, they are erroneously made to pass through the axes of the laciniae.

The only remaining author that notices these vessels is M. Mirbel, who in the second part of his valuable *Elémens de Physiologie Végétale et de Botanique*, published in 1815, introduces into his character of *Compositæ* the fact of the laciniae of the corolla being furnished with marginal nerves. This observation, if not original, the author may have adopted either from my essay already quoted, of which he was in possession soon after its publication, or from M. Cassini's third memoir, which was read to the Institute of France six months after that essay appeared: but he could not have derived it from the passage in that author's second memoir, on which he rests his claim; no notice being there taken of the disposition of vessels in the laciniae.

In M. Cassini's memoir expressly on the Corolla of *Compositæ*, which was read to the Institute of France in December 1814, and of which an abstract, by the author ⁸²¹ himself, is given in a late number of the *Nouveau Bulletin des Sciences*, the disposition of vessels in the corolla is expressed in the following terms:

"Chacun des cinq pétales dont se compose la corolle est muni de deux nervures très simples qui le bordent d'un bout à l'autre des deux côtés, et confluent par conséquent au sommet."

On this statement I have several remarks to offer. And first, I object to its hypothetical language. Whatever opinion may be formed of the theory here adopted by the author, namely, that every monopetalous corolla is in reality composed of several confluent petals; a theory first proposed by Linnæus himself in his *Prolepsis Plantarum*, and ably supported on different grounds by Mons. Decandolle in his excellent *Théorie Élémentaire de la Botanique*; I can see no advantage in adopting its language in stating a fact of this kind, especially if proposed as a practical character.

For my own part, I consider this opinion as correct in the sense in which it was held by Linnæus, without, how-

ever, connecting with it the ingenious hypothesis of M. Decandolle, namely, that petals are only modified stamina. It remains to be seen on what ground M. Cassini has adopted this theory, as proposed by M. Decandolle, for Compositæ, the only family which seems to present a very important objection to it, in having its principal, and in the greater part of the order its only, vessels occupying the lines of junction of the supposed united petals.

To adapt this disposition of vessels to the theory, M. Cassini is obliged to subdivide their apparently simple trunks; a division, however, which may be regarded as entirely hypothetical. From the observations I have made on the subject, I have no doubt that these trunks are equally simple with the secondary nerves when present, or with the primary in other families. I find them to consist of two kinds of vessels, the spiral and ligneous. Of the spiral vessels there are generally several in the cord: in *Helianthus multiflorus*, however, I have not been able to find more than one, either in the trunk of the nerve above the insertion of stamina, or in the branches of the lacinia. It will be of some interest to verify this fact (which I by no means give with absolute confidence), both on account of the apparently formidable objection it presents to the theory in question, and also that, in following it up by an examination of the point of division, a clearer idea may be obtained of the ramification of spiral vessels than has hitherto been given.

My second objection to M. Cassini's account is, that he describes the nerves as marginal through their whole length. I have formerly, in the passage already quoted, stated them to be parallel and approximated to the margins of the lacinia. Perhaps in no instance can the branches be considered as strictly marginal; in many cases they are manifestly distinct from the margins, and in the genus *Hymenopappus* are further removed from them than from the axis of the lacinia. In *H. scabiosæus* there is also an evident inequality of the two branches in each lacinia, the stronger extending nearly to the apex, while the weaker either entirely disappears before it reaches the stronger, or

unites with it considerably below its termination. In *H. tenuifolius* this irregularity is still greater; one branch being not unfrequently altogether wanting, and even the remaining branch considerably weakened: where this happens a secondary vessel is always produced, though very few flosculi are furnished with five complete middle nerves.

To the fact stated by M. Cassini that the lateral nerves are always simple, I have met¹ with only one apparent exception, in an unpublished species of *Madia*, where they are connected by a few branches with the secondary or ⁸⁴ middle nerve, which in this plant is more strongly marked than the primary, and from which indeed these connecting branches probably originate.

It must, I think, be admitted by M. Cassini, that in many genera of Compositæ five vessels passing through the axes of the segments exist, even ten others are occasionally found, as in *Helianthus*, though these can hardly be traced below the insertion of stamina. But as it has been already shown that the lateral or primary vessels are not strictly marginal through their whole length, and as one instance has been produced in which their branches, if not themselves subdivided, are at least connected by ramifications of the middle nerves,¹ it follows that a monopetalous corolla having in its tube fifteen nerves with distinct origins, three of which are continued through each of its segments, and unite together at the apex, would upon the whole better correspond with the definition M. Cassini has given of the Corolla of Compositæ, than the actual disposition of vessels in that order. Now such a structure exists in the whole of Goodenoviæ,² a family of plants very

¹ M. Cassini himself (in a note to his third memoir published in the Journal de Physique for February 1816, p. 129) has given another instance of the ramification of nerves in *Iva frutescens*.

² I have formerly observed (in Prodr. Flor. Nov. Holl. p. 580, and in General Remarks on the Botany of Terra Australis [vol. i, p. 33]) that *Euthales* and *Velleia*, genera belonging to Goodenoviæ, exhibit the remarkable and nearly peculiar character of a corolla having the lower part of the tube cohering with the ovarium, while the calyx is entirely distinct. I have at the same time remarked that, even in those genera of the same natural family in which the calyx is coherent, the tube of the corolla may be supposed to be continued down to the base of the ovarium; and that this becomes even evident in such species as have the

nearly related to Compositæ. It exists also in *Ernodea*,^[85] in which the lateral nerves, though they give out externally a few branches, observe the same course, and terminate in the same manner in the lacinia as those of Compositæ. A similar disposition is observable in certain genera of Solanaceæ, as *Datura* and *Cestrum*, though in these the lateral nerves are more ramified, and their trunks generally less distinct in the lacinia. It appears therefore that, in adopting M. Cassini's theoretical expression for the vascular structure of the corolla of Compositæ, one peculiarity actually existing is lost.¹

adhering part dilated into nectariferous processes; or in those where, the segments of the calyx not being closely approximated, the coloured corolla is visible in the interstices. In some species of *Goodenia*, particularly *G. decurrens* and *bellidifolia*, I find it practicable to separate not only the adhering calyx, but also the tube of the corolla from the ovarium. In the tube thus separated it appears that the lateral nerves, which preserve their parallelism to the middle nerve nearly to the base of the segment, become more evidently divergent below the point of adhesion, and in such a degree that the corresponding branches of the neighbouring segments unite with each other considerably above the middle of the tube, forming a common trunk, which is continued to the base of the ovarium; the five trunks thus formed uniting internally with those from which the filaments originate, and externally with the axes of the opposite segments of the calyx. The middle nerves of the segments of the corolla are in like manner continued below the point of cohesion to the real base of the tube.

The analogy of this disposition of vessels in the corolla of *Goodenivæ* to that of Compositæ is obvious. To assimilate entirely the two structures, it is only necessary to suppose a deeper division of the five primary vessels of Compositæ, and a continuation of the tube of the corolla below its apparent base to that of the ovarium. That this is its real origin, is rendered not improbable both from the analogous structure now described in the family of *Goodenivæ*, and from the manifestly hypogynous corolla of *Brunonia*; a genus in many respects still more nearly related to Compositæ, though differing in the disposition of the vessels of its corolla.

The more direct proof of this origin, derived from an examination of the surface itself, can hardly, perhaps, be expected where the parts are generally so small, and where, as I conceive, the surface of the pericarpium in many cases depends less on that of the coloring envelopes, than on the proper figure of the ovarium itself, as seems to be likewise the case in Umbellatæ.

There are however a few cases in which this opinion respecting the origin of corolla in Compositæ may derive some additional support from the appearance of the surface of the ovarium, as in *Marshallia* and *Hymenopappus*, in both of which genera, but particularly in the former, it is marked with ten longitudinal striæ, of which the five stronger are continued into the five nerves of the corolla, the remaining five ending abruptly at the apex of the ovarium.

¹ A still stronger objection to M. Cassini's definition is, that while its application to Compositæ is only hypothetical, it very nearly corresponds with the

The principal peculiarity, however, consists in the corolla of a syngenesious plant, when reduced to its smallest number of nerves, having these nerves alternating with its segments in the tube. I am acquainted with no instance of this order of reduction in the nerves of any other monopetalous corolla, but I observe an apparent tendency to it in *Portlandia* and *Catesbæa*. In the tube of the corolla of both these genera there are ten nerves, of which the five that alternate with the segments are manifestly stronger, and seem to furnish the greater part of the vascular system of the upper part of the tube and of the segments; the intermediate nerves being there somewhat like recurrent branches.

I shall conclude this subject by observing, that although the existence of nerves alternating with the segments of a monopetalous corolla, dividing below the sinus and uniting their branches at the apex of the segment, be rare, this disposition is comparatively frequent in a monophyllous calyx, especially where its æstivation is valvular. Labiatae furnish the most striking examples of this structure. I am not however acquainted with any instance of a calyx having five nerves only, and those alternating with its segments.

The æstivation or condition of the corolla before expansion is the subject of my *second* remark on Compositæ. I have, in the observations formerly quoted, stated this to be *valvular*, that is, having the margins of the segments applied to each other and dehiscing like the valves of a capsule. As I have remarked in the same place that this æstivation exists in several other families, it is rather surprising that M. Cassini, in the abstract of his third memoir given in the *Nouveau Bulletin des Sciences* for last October, should seem to consider this character as peculiar to

actual disposition of vessels in certain polypetalous genera. Thus in *Pittosporum revolutum*, each of the petals has three nerves with distinct origins. Of these the two lateral, evidently within the margins, less so, however, than in *Hymenopappus*, are quite simple in the unguis, and ramify more or less in the laminæ, near the top of which they unite with each other and with the middle nerve.

Compositæ.¹ It appears also that he is not aware of any exception to it in the class. I have however, in a different part of the same essay, noticed one exception existing in *Chusqueira*, and I have since found another in *Corymbium*. In both these genera the æstivation is induplicate, that is, the margins of the segments are doubled in, so that in the unexpanded state none of them are visible. I have in the passage referred to observed that the valvular and indu-^[84] plicate modes of æstivation easily pass into each other, merely by an addition or abstraction of the elevated margins of the laciniae: instances of their abstraction, and of the consequent conversion of the induplicate into the valvular mode, occur in several Goodenoviae, and in some Convolvulaceæ and Solanaceæ; while *Chusqueira* and *Corymbium* are examples of their addition in an order where they are generally wanting.

My third remark is entirely borrowed from Schkuhr,² who states that in all *Uchhoraceæ* or *Ligulatae* the pollen is

¹ Since this paper was read, M. Cassini has published his memoir (in the Journal de Physique for February 1816), in which he states the same æstivation to exist in certain other families, namely, Campanulaceæ, Lobeliaceæ, and Rubiaceæ. This observation, if applied to the whole of these families, as is evidently the author's intention, is correct only with respect to Campanulaceæ, from which I have separated Stylidæ as a distinct order, partly, as I have stated, on account of its imbricate æstivation. In a considerable part of the Lobeliaceæ of Jussieu, which includes my Goodenoviae, the æstivation is not valvular but induplicate: and though in Rubiaceæ the valvular mode is very general, there are many remarkable exceptions to it, as *Gardenia*, *Isora*, *Pavetta*, *Coffea*, and several other genera, where it is unilaterally and obliquely imbricate, as in most of the Apocineæ, with which Linnæus united them under the name of Contortæ, derived from this very circumstance. On this subject I may be allowed further to remark, that M. Cassini, who in the memoir now cited has repeatedly asserted his claim to the priority of the observation on the disposition of vessels in the corolla, has in treating of its æstivation omitted to notice what had been already published respecting it in my essay above quoted, where I conclude he must have seen my observation, as he refers to the sentence containing it. The æstivation of corolla in Compositæ is also noticed in the observations on *Brunonia*, contained in my Prodrômus Floræ Novæ Hollandiæ, which I suppose he has not seen: I may therefore, for the general importance of æstivation of calyx and corolla in affording characters both for Orders and Genera, refer him to almost every page of the same work, and to its preface, for an observation on the degree of attention that had been previously paid to this point of structure, which will enable him to correct in some measure his own remark on the subject.

² Botanisches Handbuch 3, p. 8.

angular, and that in *Corymbiferae* and *Carduaceae*, or in all tubular florets, it is spherical or oval.

All the figures which this author has given of pollen in *Cichoraceae* represent it as a regular icosahedron, except that of *Geropogon glabrum*, which is a dodecahedron. I believe neither of these forms of pollen has been observed in any other family of plants.

A *fourth* remark on *Compositae* I do not offer with absolute confidence, as it is opposed to the statement of M. Cassini, on whose general accuracy I have great reliance. It relates to the disposition of the branches of the style or stigmata, which according to M. Cassini are lateral, or right and left with relation to the axis of the common receptacle; whereas, I consider them as anterior and posterior, though in many cases by a slight degree of twisting in the style they acquire what M. Cassini regards as their original position.

This may seem a point of very little consequence to establish. Independent however of the necessity of minute accuracy in every case, it appears to me to have some connexion with my *fifth* remark, which relates to the internal ⁶⁰ structure of the Ovarium of *Compositae*. I am not aware of anything having been yet said on this subject further than that it contains a single erect ovulum, inserted at the base of the cavity. In addition to this, I observe in the greater part of *Compositae*, whose ovarium I have examined, two very slender filiform cords, which, originating from opposite points of the base of the ovulum, or of its short foot-stalk, run up, and are more or less connected with, the lateral parietes of the ovarium, until they unite at the top of its cavity, immediately under the style; between which and the ovulum a connexion is thus formed. In many cases, as in *Liatrix spicata* and *Tussilago odorata*, these cords are easily separable from the ovarium, and have such a degree of tenacity that they may be extracted from it entire, along with the ovulum. In other cases they more firmly cohere with the sides of the cavity: and in those plants in which I have been unable to see them distinctly,

I conclude they are not absolutely wanting, but that their connexion with the parietes is still more intimate.

These cords may be supposed to consist either solely of the vessels through which the ovulum is fecundated, or to contain also the remains or indications of a system of nourishing vessels, or chordæ pistillares, the position of which points out the true nature of the ovarium in this class, or the relation it has to the apparently less simple ovarium of other families. I am inclined to adopt the latter supposition. In order, however, to be understood on this subject, it is necessary to premise that I consider the pistillum or female organ of all phænogamous plants to be formed on the same plan, of which a polyspermous legumen or folliculus whose seeds are disposed in a double series may be taken as the type. A circular series of these pistilla, disposed round an imaginary axis, and whose number corresponds with that of the parts of the calyx or corolla, enters into my notion of a flower complete in all its parts.

But from this type and number of pistilla many deviations take place, arising either from the abstraction of part of the complete series of organs, from their confluence, or from both these causes united; with consequent abortions and obliterations of parts in almost every degree. According to this hypothesis, the ovarium of a syngenesious plant is composed of two confluent ovaria; a structure which is in some degree indicated externally by the division of the style, and internally by the two cords which I consider as occupying the place of two parietal placentæ, each of these being made up of two confluent chordulæ, belonging to different parts of the compound organ. I am well aware how very paradoxical such an hypothesis must seem, especially when applied to a structure apparently so simple as that of the ovarium of Compositæ; and I therefore regret that I am not yet fully prepared to bring forward in its support a series of facts already in my possession, consisting of deviations from the usual structure of organs, and particularly of instances of stamina changed into pistilla.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

1.8 2.0 2.2 2.5 2.8 3.2 3.6 4.0 4.5 5.0 5.6 6.3 7.1 8.0 9.0 10.0 11.2 12.5 14.0 16.0 18.0 20.0 22.4 25.0 28.0 31.5 36.0 40.0 45.0 50.0 56.0 63.0 71.0 80.0 90.0 100.0

1.0 0.1 0.2 0.3 0.5 0.7 1.0 1.5 2.0 3.0 4.0 6.0 8.0 10.0 15.0 20.0 30.0 40.0 60.0 80.0 100.0

In the mean time it may give some plausibility to the hypothesis to remark, that there are families of plants strictly natural in which a series of degradations exist, if I may so speak, from the assumed perfect pistillum, to a structure as simple as that of *Compositæ*.

Thus in *Proteaceæ* we have the type of the perfect pistillum in the many-seeded folliculus of *Embothrium*; the first degree of imperfection in that of *Grevillea*, where only one ovulum of each series remains; a further reduction in the indehiscent monospermous fruit of *Leucospermum*, in which the insertion of the ovulum is lateral; and the simplest form in *Protea* itself, where the single ovulum is inserted at the base of the cavity. *Proteaceæ*, however, exhibit a series of obliterations in the parts of a single pistillum only. An illustration more in point, though somewhat less perfect as a series, may be taken from *Goodenoviæ*, an order of plants very nearly related to the class of which we are treating. In the greater part of *Goodenoviæ*, the ovarium is bilocular, each cell having an indefinite number of seeds; in the greater number of *Scævolaæ*, each cell is reduced to a single ovulum; while in some species of the same genus, and in all the species of *Dampiera*, the ovarium, though retaining its external characters, is reduced to a single monospermous cell, with an erect ovulum, as in *Compositæ*. The natural order *Cruciferaæ* exhibits also obliterations, more obviously analogous to those assumed as taking place in syngenesious plants; namely from a bilocular ovarium with two polyspermous parietal placentæ, which is the usual structure of the order, to that of *Isatis*, where a single ovulum is pendulous from the apex of the unilocular ovarium. And lastly in the genus *Bocconia*, in the original species of which (*B. frutescens*) the insertion of the single erect ovulum has the same relation to its parietal placentæ, as that of *Compositæ* has to its filiform cords, a second species (*B. cordata*) exists in which these placentæ are polyspermous.

My sixth observation on *Compositæ* regards the order in which the florets expand. To understand the relation this order has to that of other families, it may be necessary first

to make a few remarks on the more usual modes of inflorescence.

It is well known that in an absolutely simple spike the expansion of the flowers is ascendent; that is, begins at the base of the spike and proceeds regularly upwards. To this order very few real exceptions occur, several of the apparent deviations being connected with some degree of composition in the spike.

It is also known that in a compound spike, while the expansion of each partial spike is ascendent, that of the spikes, with relation to each other, is descendent; the terminal spike expanding first, and the others in a regular succession downwards. This order, indeed, admits of a greater number of exceptions than that of the simple spike; several of them apparently depending on the density or imperfect composition of the spike; and the more usual deviation consisting in the expansion beginning below the apex, and proceeding in opposite directions from the point of commencement; the upper portion following the order of the simple, the lower that of the compound spike.¹

The simple racemus and corymbus are obviously very slight modifications of the spike, and in their expansion obey the same law.

A syngenesious compound flower, or *capitulum* as it may be termed, is merely a spike with a shortened and generally depressed axis. In cases where this capitulum is unquestionably simple, the expansion of its flowers is uniformly from circumference to centre, or in the order of the simple spike. Where the capitula are disposed in a corymbus, which is their usual mode of combination, the order of the compound spike is observed; their expansion with relation to each other being from centre to circumference. In their denser aggregations, whether forming a

¹ The most remarkable exception to the order of the compound spike exists in the compound umbel of Umbelliferae, of which the outer umbellulæ expand somewhat earlier than the central; and as this order of expansion seems to extend through the whole natural family, *Astrantia*, in which the terminating umbel expands much earlier than those of the lateral branches, cannot be considered as having a compound umbel.

compound spike or head, the same order of expansion obtains, and it continues though the florets in each common calyx or involucrum should be lessened in number, or even reduced to unity, as in *Echinops* and *Rolandra*.

93] The absolute constancy in the order of expansion of the simple capitulum from circumference to centre, and the more or less complete inversion of this order in the compound capitulum, appear to afford tests of the real structure in certain cases where the degree of composition, and consequently the proper names of some of the parts, might otherwise be doubtful.

To illustrate this I select two genera, *Lagasca* and *Cæsulia*.

In *Lagasca* the capitulum, both from its form and the appearance of its involucrum, might at first sight be considered as simple: on examination, however, it is found to differ from all simple capitula, in each floret being furnished with a tubular envelope, exactly resembling a five-toothed perianthium, but which does not in any state cohere with the included ovarium.

Cavanilles, by whom the genus was established, regarded this envelope as a genuine perianthium, and erroneously described its tube as cohering with the ovarium; an error which is copied in Persoon's *Synopsis Plantarum*, where the genus is consequently placed in *Polygamia æqualis*. Jacquin, who has published *Lagasca* under the name of *Noccea mollis*,¹ also describes the envelope of each flower as a proper perianthium, although aware of its tube being distinct from the ovarium. Subsequent writers have, indeed, more correctly referred the genus to *Polygamia segregata*; but the terms involucellum and calyculus, which they apply to the envelope in question, appear to me objectionable, for a reason that will presently be given.

Three suppositions may be formed respecting the nature of this envelope, namely, either that it is an involucrum reduced, as in *Echinops*, to a single flower; secondly, that it is a proper perianthium, which in appearance it very

¹ *Fragm. Bot.* p. 58, tab. 85.

much resembles; or thirdly, that it is more analogous to the outer calyx of *Scabiosa*, which M. Cassini seems to consider different in its nature from both these parts.

But the order of expansion in *Lagasca*, which is, though with some degree of irregularity, from centre to circumference, or that of the compound capitulum, seems to decide the question respecting the envelope of each flower, and to establish its identity with involucre: nor does this involucre differ materially from that of *Echinops*, except in the reduced number and confluence of its component parts.

The real structure of *Cæsulia* is perhaps less obvious.

This genus, which was first published by Dr. Roxburgh,¹ is referred by him to *Polygamia segregata*; the tubular envelope or involucre of each floret being described as distinct from the included ovary.

Kœnig, on the other hand, by whom the genus was discovered, and whose account of it is given in the same work, describes the partial involucre of Roxburgh as the surface of the ovary itself; its segments being, according to him, a pappus of two leaves. And lastly, Willdenow, regarding this involucre as merely paleæ of the receptacle, refers the genus to *Polygamia æqualis*; in which order it is continued, both in Persoon's Synopsis, and in the second edition of Mr. Aiton's Hortus Kewensis.

This last view of the structure seems the most erroneous of any, and was probably adopted by Willdenow, in consequence of his having added to the genus a second species not really belonging to it, and which I shall have occasion to notice in another part of my subject.

An examination of the parts of fructification in different stages reconciles the opposite statements of Kœnig and Roxburgh; for I find that at the time of flowering the envelope of each floret is, as Roxburgh has figured it, distinct from the ovary, with which, however, in a more advanced stage its tube becomes firmly united; a fact that sufficiently accounts for Kœnig's description.

¹ In *Corom. Plants*, i, p. 64, t. 93.

There is here, therefore, a nearer approach to a true perianthium than in the involucre of *Lagasca*; but the expansion of the flowers being, as in that genus, from centre to circumference of the capitulum, I consider the envelope of *Casulia* as unquestionably an involucre, and the genus consequently belonging to *Polygamia segregata*.

I may here remark, that the name *Polygamia segregata*, invented by Linnæus for those genera of *Compositæ* with densely aggregate capitula, is calculated to give an erroneous idea of the nature of the structure; the opposite term *Polygamia congregata* being, according to the view now taken, obviously more proper for those genera, at least, whose involucre contain several flowers. It is not unlikely, indeed, that Linnæus himself was aware of the true nature of the inflorescence of these genera; but the term *Polygamia congregata* would not have suited the artificial arrangement which he adopted in his subdivisions of the class, nor his including in it the order *Monogamia*; for with this order the single-flowered genera of *Polygamia segregata* must then have been confounded.

It is a curious circumstance, that the order of expansion in *Compositæ* does not depend on the number of flowers actually existing, but on the effort, if I may so term it, made to produce them, manifested by the presence of an involucre or common calyx, which is in some cases reduced to a single flower. The fact at the same time contributes to prove, that the whole natural class is formed on ^{96]} that plan of dense aggregation of flowers, for which I have already attempted to show that certain parts of the structure of a syngenesious floret are peculiarly well adapted.

The circumstance, however, is not confined to *Compositæ*, but exists in an equally remarkable degree in *Gramineæ*.

I have formerly¹ considered the gluma, or what Linnæus has termed calyx, in this family of plants, as an involucre.

In those genera where this gluma or involucre con-

¹ [Vol. i. p. 55.]

tains several flowers their expansion is generally ascendent, or in the order of the simple spike. In a spike formed by these many-flowered glumæ, as that of *Triticum* and *Lolium*, the expansion of the partial spikes, with relation to each other, is descendent, or in the order of the compound spike; in most cases, however, with that deviation, which I have already noticed, of the expansion commencing below the apex and proceeding in opposite directions. But as the same descendent expansion takes place in a spike formed of single-flowered glumæ, it may be inferred that the genuine type or most perfect form of a grass is to have several flowers in its gluma or involucre: a view not only consistent with the fact of a great majority of the order having actually this disposition; but also with that peculiarity in the vascular structure of the inner valve of the perianthium; which, whether it be considered as indicating that this part is formed of two confluent valves, an opinion I have elsewhere¹ advanced, or merely as a transposition of vessels in a simple valve, analogous to that in the syngenesious floret, is evidently adapted to the many-flowered spicula, though equally existing in that with a single flower.

The resemblance between the outer calyx of *Dipsacæ* and the single-flowered involucre of *Compositæ* is so striking, that it cannot appear very paradoxical to consider them as both of the same nature.

In *Dipsacæ*, however, there is no instance of the outer calyx containing more than one flower, and the evidence afforded by inflorescence on this subject is not altogether satisfactory.

In *Dipsacus* it has been long noticed that expansion begins about the middle of the spike, and proceeds in opposite directions from the point of commencement: this order is evidently more analogous to that of the compound than of the simple spike; there being several instances of spikes manifestly compound, where the same inversion of the upper part exists.

¹ In General Remarks on the Botany of New Holland [vol. i, p. 55].

But a fact, which I do not find any where observed, respecting the inflorescence of certain species of *Scabiosa*, particularly *succisa* and *atropurpurea*, is not so easily reconcilable with the compound spike: in these, and I have reason to think in many other species of the genus, the expansion begins simultaneously at the base and middle of the capitulum, proceeding regularly upwards from both points. Were this the case in all *Scabiosæ*, the compound nature of the spike in *Dipsacæ*, although by no means proved, might be considered not improbable: there are, however, several species of the genus in which the order of expansion is altogether that of the simple spike.

Connected with the subject of inflorescence, I may remark that priority of development, whether among similar parts in the same flower or the different flowers of the same spike, is generally accompanied with greater perfection of these parts or flowers, and apparently with greater power of resisting the ordinary causes of abortion or obliteration.

I have formerly¹ observed respecting several natural families of plants, in which the stamina are in a determinate number, but a number subject to reduction, that this reduction, where the flower is of a regular form, takes place in the same order in each natural family. Thus in *Junceæ*, which are generally hexandrous, the triandrous species have their stamina constantly placed opposite to the three outer leaves of the perianthium, while in *Restiaceæ*, *Asphodeleæ*, and I believe in a great part of the regular-flowered *Liliaceæ*, in certain species of which a similar reduction occurs, the stamina in the triandrous species are placed opposite to the inner leaves or segments of the perianthium. But in both cases the greater perfection of those stamina that exist in genera or species reduced to the smallest number, is indicated, where there is no reduction, by the earlier bursting of their antheræ; so that from this circumstance the order of reduction or abortion of stamina

¹ In *Prodr. Flor. Nov. Holl.* vol. i, and Appendix to Flinders's Voyage to Terra Australis [vol. i, p. 52].

in any natural family may with some confidence be predicted by an examination of those genera where the number is complete.

Wherever the separation of sexes takes place, it may be assumed that the female flower is the more perfect production. And if this be admitted, where both sexes exist in the same simple spike the female should be found at its base, or where expansion commences, which is almost uniformly the case. For the same reason, in the trifid or trichotomous inflorescence, the female should be placed in the centre, which is also generally the fact.¹

This connexion between præcocity and perfection of development is even more constant than the order of expansion in certain forms of inflorescence; as it is found to extend to several of the exceptions to this order.

Thus in the apparently simple spike of *Poterium*, where the order of expansion is descendent, the female flowers occupy the upper part of the spike; and this relation also exists in the more compound inflorescence of *Ricinus*, *Siphonia*, and *Celtis*, in which the order of expansion is equally inverted.

It may seem rather paradoxical to select *Euphorbia* as an example of the same relation; this genus being considered by Linnæus, and the greater part of the botanists who have adopted his system, as having a dodecandrous hermaphrodite flower. We have already, however, I believe, sufficient evidence that this supposed hermaphrodite flower is in reality formed of several monandrous male flowers surrounding a single female.²

¹ To this order the most remarkable exception occurs in *Begonia*, in which the male flowers are central, and expand long before the lateral female flowers.

² To the arguments I have adduced (in my Remarks on the Botany of New Holland [vol. i, p. 28]) in support of this opinion, I am now enabled to add the more direct proof derived from certain species of *Euphorbia* itself, in which the female flower is furnished with a manifest calyx. I have formerly observed, that in a few cases the footstalk of the ovary is dilated and obscurely lobed at top; but in the species now referred to it terminates in three distinct and equal lobes of considerable length, and which being regularly opposite to the cells of the capsule may be compared to the three outer foliola of the perianthium of *Phyllanthus*, between which and the cells of the capsule the same relation exists. This calyx is most remarkable in an undescribed species of *Euphorbia* from the coast of Patagonia, in the Herbarium of Sir Joseph Banks; but it is observable, though less distinct, in *E. punicea* and several other species.

In conformity with this view of its composition, and with the relation above attempted to be established, the development of the pistillum precedes that of the stamina in many species of the genus.

It is more difficult to determine whether this order of expansion and relative position of sexes in *Euphorbia* be in conformity with the general rule, or an exception to it. For its fasciculus of flowers may be considered as analogous either to the simple spike, and consequently having an inverted order of expansion, as in *Allium descendens*, and certain species of *Grevillea* and *Anadenia*: or it may be assimilated to the compound spike, as in several species of the genus the male flowers appear to be separated into ¹⁰⁰ fasciculi; and according to this view the order of expansion is direct, the central female flower being the representative of the terminal partial spike.

There is even a third species of inflorescence with which the fasciculus of *Euphorbia* may be compared, namely, that consisting of one or more verticilli with a single flower in the centre. In this, which may be considered a modification of the spike or umbel, the usual order of expansion seems to be from centre to circumference. Its simplest form occurs in an unpublished New Holland genus of the same natural family with *Euphorbia*, in which a single verticillus of male flowers surrounds the central female flower. *Lambertia* may be considered as another instance of the same mode, and as far as can be determined, in a case where the flowers are hermaphrodite and their expansion nearly synchronous, following the same order. In all the known species of this genus the leaves are verticillate, and uniformly in threes: in *L. formosa* and *inermis* the involucre constantly contains seven flowers, while in *L. uniflora* it is reduced to one flower. The seven flowers of the two former species I consider as made up of two verticilli, in number of flowers corresponding with that of the leaves, and of a single central or terminal flower; to which terminal flower *L. uniflora* appears to be reduced. From this order of reduction it may be assumed as more probable that species of *Lambertia* should be found with

ten or four flowers in the involucre than with nine, six, or three. But greater permanence being, as has been already remarked, generally connected with greater perfection, it becomes also probable that, if any species of this genus should be discovered with androgynous capitula, the female flower will occupy the centre as in the genus of Euphorbiaceæ above referred to.

It is worthy of remark, and may indeed appear in some degree at variance with the foregoing observations, that although in an assemblage of flowers priority of expansion generally indicates a greater degree of perfection, and consequently a more ready convertibility of the hermaphrodite into the female flower; yet in a hermaphrodite flower the development of stamina usually precedes that of pistilla. The most remarkable exceptions to this order of development which I at present remember, occur in several species of *Plantago*, where the stigmata are fully developed, and often even withered, before the bursting of the antheræ.

I now proceed to make some remarks on certain genera of Compositæ which either occur under different names in late systematic works, or whose structure and limits seem to be imperfectly understood.

SOLIVA

was established in the *Prodromus Floræ Peruvianæ et Chilensis*, and is adopted by Persoon in his *Synopsis Plantarum*.

To this genus *Hippia minuta* of the Linnean Herbarium unquestionably belongs, and it is perhaps not specifically distinct from *Soliva pedicellata*. But on comparing the structure of this plant with the figures and descriptions, given by Mons. de Jussieu (in the fourth volume of the *Annales du Muséum*.) of the different species of his *Gymnostyles*, it appears to me evident that the whole of

this genus is referable to *Soliva*, whose principal characters would consist in the want of corolla or perhaps its accretion with the persistent style in the female florets; in the pericarpia being more or less winged, and presenting their disk instead of their margins to the centre of the capitulum.

^{102]} Sir James Smith has already pointed out the error M. de Jussieu has been led into in referring *Hippia minuta* Linn. to his *Gymnostyles nasturtiifolia*, a plant much more nearly related to *Hippia stolonifera* of Brotero; which, from repeated examination, I can with confidence refer to the same genus.

Gymnostyles anthemifolia is stated by M. de Jussieu to be a native of New South Wales: but as I have observed it only in cultivated ground in the neighbourhood of Sydney, and as it has certainly been found in South America, of which four other species of the genus are unquestionably natives, it has probably been imported into New South Wales, perhaps from Brazil; nor is it altogether improbable that *Hippia stolonifera* of Brotero may have been introduced into Portugal from the same quarter.

GRINDELIA,

described by Willdenow in the Transactions of the Natural History Society of Berlin for 1807, and subsequently in his Enumeratio Plantarum Horti Berolinensis, flowered in Kew Gardens for the first time in 1815, when I had an opportunity of examining it, and of determining its very near affinity with *Donia*, a genus proposed in the second edition of Hortus Kewensis, and adopted by Mr. Pursh in his Flora of North America: the principal distinction between these two genera consisting in a difference in the number of radii of the pappus, which in *Grindelia* is described by Willdenow as of two rays, and according to my observations has more frequently one only. But as even in *Donia* the number of rays, though indefinite, is variable, and the structure of the pappus is very nearly similar in both genera, which in all other respects agree, it

may be perhaps expedient to unite them under the name of *Grindelia*, which was first in order of publication.

TRIDAX

[103]

was first established by Linnæus, in Hortus Cliffortianus, from a specimen found at Vera Cruz by Houston, and sent to Clifford by Miller. As Linnæus had no specimen in his own collection, that in Clifford's Herbarium, now in the possession of Sir Joseph Banks, is the only authority for the genus; and on examining this specimen I find the pappus to be not setaceous, as Linnæus has described it, but distinctly plumose. There is, therefore, no difference whatever between *Tridax* and *Balbisia* of Willdenow; and on comparing *Tridax procumbens* with *Balbisia elongata*, I cannot satisfy myself that they are even specifically distinct.

ANGIANTHUS.

Angianthus tomentosus of Wendland's Collectio Plantarum, (vol. ii, p. 32, tab. 48,) published in 1809, is evidently the same plant as my *Cassinia aurea*, described in the fifth volume of the second edition of Hortus Kewensis, which did not appear till 1813. Wendland neither mentions the native country of his *Angianthus*, nor from whence he received it. He must, no doubt, however, have obtained it from Kew Garden, where it was introduced and flowered from seeds which I collected in 1802, in the island of St. Francis, on the South coast of New Holland.

MEYERA.

This genus, described by Schreber in his edition of the Genera Plantarum, is not adopted by Willdenow. Swartz, however, in his Flora Indiæ Occidentalis, has referred to it, and I have no doubt correctly, *Eclipta sessilis* of his Prodrromus. On comparing this species of *Meyera* with a plant in Sir Joseph Banks's Herbarium, collected in Peru

^{104]} by Dombey, and which exactly agrees with *Sobrega* of the Flora Peruviana, it appears evident that this genus is reducible to *Meyera*. *Enhydra* of Loureiro's Flora Cochinchinensis, though described somewhat differently, and referred to *Polygamia segregata*, I have little doubt, belongs to the same genus; as does unquestionably *Hingstha* of Roxburgh's unpublished Flora Indica, where it is also referred to *Polygamia segregata*. This plant, which I have examined, is scarcely distinct from a species of *Meyera* that grows in New South Wales.

Cryphiospermum of Mons. de Beauvois's interesting Flore d'Oware et Benin, although reduced by him to Cichoraceæ, I have but little hesitation in referring also to *Meyera*. And lastly, *Cæsalia radicans* of Willdenow, likewise a native of æquinoctial Africa, is perhaps not specifically different from *Cryphiospermum repens* of Mons. de Beauvois.

MELAMPODIUM

was established by Linnæus, in the first edition of *Genera Plantarum* and in *Hortus Cliffortianus*, from a specimen found by Houston near Vera Cruz, and communicated by Miller to Clifford, in whose Herbarium, now forming part of the collection of Sir Joseph Banks, it still exists. It does not appear that this plant has been found by any other botanist than Houston; and according to the character given by Linnæus of *Melampodium*, it must be considered the only species of the genus.

In the second edition of *Species Plantarum* he added to it, but with a doubt, *Melampodium australe*, a plant adopted from Læffling, according to whose description the pappus and surface of the seed are widely different from those of the original species. Swartz has referred to the genus a third species, *M. humile*, entirely distinct in these respects from both the former; and more recently a fourth species, *M. longifolium*, with seeds differently modified from all the others, has been annexed to it.

^{105]} But if these four plants, so extremely different from each other in pappus and form of the pericarpium, really

belong to the same genus, as their habit seems strongly to indicate, there can be no reason to separate from them *Alcina* of Cavanilles, erroneously considered by Willdenow as a species of *Wedelia*: and *Dysodium* of Richard, published in Persoon's Synopsis, though differing from all the others in the form of its pericarpium and in that of its receptacle, must also be reduced to this genus. If, however, the part described by Linnæus as pappus in *Melampodium americanum* be really such, and if the pericarpium itself vary so widely both in form and surface, it would be inconsistent with the principles of division generally adopted in Compositæ, to unite all these plants into one genus, notwithstanding their great resemblance in habit as well as in the other parts of fructification; and it would be at least in vain to look for any combining character in this part of their structure.

A careful examination of the female flowers, especially in an early stage, removes this difficulty, by proving that the supposed external coat of the ovarium, with its various inequalities of surface, some of which have been described as pappus, is in reality an involute bractea or foliolum of the involucre, like that of *Micropus*, completely inclosing the ovarium, but from which in several species of the genus it is entirely, and in others in great part, distinct.

CRASPEDIA

first appears in Forster's *Prodromus Florulæ Insularum Australium*, where an essential generic character is given, but no description of the species. The genus is adopted and the character received without remark by Willdenow in his edition of *Species Plantarum*, and by Persoon in his Synopsis. Among George Forster's drawings of subjects of natural history made in Cook's second voyage, and now in the library of Sir Joseph Banks, there is a figure of this plant, from which it appears that he originally referred it to *Stæhelina*; a proof that he had not at that time very carefully examined it. It is not improbable therefore that he afterwards proposed it as a distinct genus, belonging to

Polygamia segregata, from finding that this had been already done by Solander, whose name (*Cartodium*), however, he did not think it necessary to adopt, and with whose generic character he probably was not acquainted. In his own he very erroneously states that there is no partial involucre, and hence perhaps M. Labillardière entirely overlooked *Craspedia* when he established his *Richea* from a nearly related species of the same genus. That such is the case I have long since briefly noticed,¹ and have ascertained by a comparison of the specimen of *Craspedia uniflora* in George Forster's Herbarium with *Richea glauca* of Labillardière, and other species of the same genus which I have observed in New Holland.

M. Labillardière's character of *Richea* is essentially correct. It is well to remark, however, that his general involucre is formed of the bractæ subtending and in equal number with the outer partial capitula; and that the general receptacle has no other paleæ than the analogous bractæ of the inner capitula. It is the more necessary to take this view of the structure, as I have found in New Holland a nearly related genus (*Calocephalus*), which differs from *Craspedia* and *Richea* in the want of these bractæ, as well as in the partial receptacles being without paleæ, and in the rays of the pappus being plumose only in the upper part. I have also another genus of this tribe (*Leucophyta*) from the same country, differing from *Calocephalus* in having a general involucre consisting of a few short bractæ, in the squamæ of its partial involucre being ¹⁰⁷¹ concave and bearded at top, and in the rays of its pappus being plumose through their whole length, as in *Craspedia*, from which it is distinguished by the want of paleæ on the partial receptacles, and very remarkably in habit.

I have selected the foregoing genera as having been either published under different names, or, as it appears to me, unnecessarily subdivided. In this extensive class it

¹ In Prodr. Flor. Nov. Holl. p. 555.

would not be difficult to point out a much greater number consisting of species improperly united. One very remarkable case of this kind is the genus

CALEA,

to which, as I intend to enter fully into the history and affinities of its species, I shall confine myself.

This genus was established by Linnæus in the sixth edition of his *Genera Plantarum*, where the natural character is given: but the following essential character, which is still retained, appears for the first time in the twelfth edition of *Systema Naturæ*, in the third section of *Polygamia æqualis*:

“*Receptaculum* paleaceum, *Pappus* pilosus, *Calyx* imbricatus.”

The species originally referred to *Calea*, in the second edition of *Species Plantarum*, are *C. jamaicensis*, *oppositifolia*, and *Amellus*, described from specimens in Browne's Jamaica Herbarium, which he had received a few years before, and incorporated with his own.

These three plants Linnæus had originally referred to *Santolina*,¹ for which it seems to me rather less difficult to account than for his afterwards uniting them together to form his genus *Calea*; two of them, according to his descriptions,² though in reality one only, being without pappus, and in other respects corresponding with the generic character of *Santolina*; and the third, which Browne had doubtfully referred to the same genus, though furnished with pappus, agreeing with the others in having opposite leaves.

But the difference in habit between all these plants and the original species of *Santolina* is so great, that it probably afterwards determined Linnæus to remove them from that genus; and although he found a sufficient generic character in the pappus of *Calea jamaicensis* only, he united with it the two other species, for a reason perhaps similar

¹ In *Amœnit. Acad.* vol. v, p. 404.

² *Loc. cit.*

to what I have supposed led him to separate all the three from *Santolina*. It is remarkable, however, that not one of these three original species of *Calea* corresponds with his character of the genus; and that they in reality belong to three very distinct genera, on principles which, I conceive, Linnæus himself would have admitted.

The *first* species, *Calea jamaicensis*, is the only one that even seems to agree with the generic character, in having pappus which at first sight (to the naked eye at least) might appear simply capillary, but which on a closer examination proves to be of a very different and nearly peculiar structure. Of this species I have seen only one authentic specimen, received from Browne by Ehret, and now in Sir Joseph Banks's Herbarium. The specimen in question, though incomplete, evidently belongs to the same species with "*Conyza fruticosa* cisti odore, floribus pallide purpureis, summitatibus ramulorum insidentibus," of Sloane,¹ of which I have examined the original very perfect specimens in his Herbarium, preserved in the British Museum,² and am satisfied that its pappus is of the same structure as that of *Calea cordifolia* of Swartz, who has well described it, but who has at the same time given a different ^{109]} account of that of *C. jamaicensis*.³ These two plants are the only published species of this genus, for which the name of *Calea* should be retained, and which may be distinguished by the following characters:

CALEA.

Caleæ species *Linnaei*.

*Involucrum*⁴ imbricatum. *Receptaculum* paleaceum. *Flosculi* tubulosi, uniformes, hermaphroditi. *Antheræ* basi muticæ. *Stigmata* acuta. *Pappus* paleaceus: radiis uninerviis, pinnatifido-striatis.

¹ Hist. Jam. i, p. 257, tab. 151, fig. 3.

² Herb. vci. v, fol. 14 and 15.

³ In Flor. Ind. Occid. vol. iii, p. 1328.

⁴ Calyx communis *Linnaei*.

Frutices (Americæ æquinoctialis,) *pubescentes, scabri*. Folia *opposita, indivisa*. Capitula¹ *corymbosa, v. terminalia, v. axillaria*. Involucri *subovati foliola nervosa, obtusa*. Paleæ *receptaculi convexi distinctæ, figura et textura fere involucris*. Corollæ *luteo-purpureæ* (Swartz), *glæ 2, laciniis dinerviis*. Achenium *subcylindraceum v. obsolete angulatum, glabrum v. pubescens, callo baseos subobliquo*. Pappus *persistens albus, nitens; radiis simplici serie subulatis, indivisis, superne denticulatis*.

OBS. In Sir Joseph Banks's Herbarium there are two plants very nearly related to *Calea*, differing from it merely in having a radius of ligular female florets. If this difference be considered sufficient to constitute a genus, it may be named *Caleacte*. The first of these plants (*C. urticifolia*), with nearly ovate-acute crenated leaves, found by Houston near Vera Cruz, is *Solidago urticæfolia* of Miller, by whom it appears to have been cultivated. The second, with deeply lobed or pinnatifid leaves (*C. pinnatifida*), was lately sent from Brazil by Mr. Sellow.

The second Linnean species, *Calca oppositifolia*, has very little affinity to the first. In attending merely to the technical character of *Santolina*, it might be referred to that genus; but it differs so widely, both in other points of structure and in habit, that there can be no question of the propriety of separating it, which may be done by the following character, and under the name of

ISOCARPHA.

Receptaculum conicum: *paleis distinctis, conformibus: extimis involucrum constituentibus*. *Flosculi* tubulosi, uniformes, hermaphroditi. *Antheræ* basi muticæ. *Stigmata* appendice elongato, hispidulo, acuto. *Achenium* prismaticum: *pappo* nullo.

Herbæ (Americæ æquinoctialis). Folia *opposita (vel*

¹ Corolla communis, Linn.

alterna) *indivisa*. Capitula *ovata*, *terminalia*, *terna* (vel *solitaria*). Paleæ *lanceolatae*. Corollæ *albidae*. Antheræ *basi truncatae*.

Obs. I have so constructed the generic character of *Isocarpa* as to include *Spilanthus atriplicifolius* of Linnæus, which, however, differs very remarkably from *Calea oppositifolia* in having alternate leaves and solitary capitula, as well as in the texture and form of its paleæ.

The pappus, consisting of three or four very minute aristæ, described by Swartz¹ in *Calea oppositifolia*, I have not been able to observe in any of the specimens that I have examined.

The *third* species, *Calea Amellus*, is probably the same plant as *Bidens scandens*, which Linnæus described in Hortus Cliffortianus, but, having no specimen in his own collection, appears to have forgotten. The original specimen in Clifford's Herbarium, now in the possession of Sir Joseph Banks, evidently belongs to the same species, and perhaps to the same individual, with a specimen in Miller's collection, which Mr. Dryander compared, and considered to agree with *Calea Amellus* of the Linnean Herbarium. The true synonym, therefore, of *Calea Amellus* is "*Bidens* *in* suffruticosus vimineus, foliis oblongo-ovatis oppositis, floribus comosis" of Browne;² while Linnæus has quoted and even derived his specific name from the same author's "*Amellus* *ramosus*, foliis remotis terminalibus, fulcris longis divaricatis;"³ which, instead of belonging to *Bidens scandens*, I believe, for the following reasons, to be *Bidens nivea*. 1st, The figure in Burmann's Thesaurus Zeylanicus,⁴ quoted by Browne for his plant, though belonging to *Lavenia erecta*, is at the same time a good representation of *Bidens nivea*, and very unlike *Bidens scandens*. 2dly, Browne's description in most respects very well agrees with

¹ In Obs. Bot. p. 302.

² Browne, Jani. 317.

³ l. c.

⁴ Eupatoriophalacrum scrophulariæ aquaticæ foliis oppositis, Burm. Thesaur. Zeyl. p. 95, t. 42.

the former species, but certainly not with *Bidens scandens*. And 3dly, I infer that *Bidens nivea* was actually in Browne's Herbarium, from finding it in the Flora Jamaicensis published in the 5th volume of Amœnitates Academicæ, and formed chiefly from that Herbarium; though a very erroneous reference for this species is there made to Browne's first *Santolina*, which, from the description, cannot possibly belong to *Bidens nivea*, but is probably *Verbesina gigantea*.

M. Decandolle has lately established a new genus, *Salmea*, consisting of *Bidens scandens*, *Bidens hirsuta*, and a third species which I have not examined. These plants are very properly separated from *Bidens* by this excellent botanist, and well distinguished both from that genus and from *Melananthera*. It is rather remarkable, however, that he has not thought it necessary to compare *Salmea* with *Spilanthus*, from which, according to his description, it differs only in its imbricate involucre. But as in *Spilanthus* the foliola of the involucre are not exactly equal, and are disposed at least in a double series, I have introduced some additional distinctions into the following character of

SALMEA.

Decandolle in Cat. Hort. Monspel. p. 140.

Involucreum imbricatum. *Receptaculum* conicum, paleis persistentibus. *Flosculi* tubulosi, uniformes, hermaphroditi (5-fidi). *Antheræ* sagittatæ. *Achenium* verticaliter compressum, bi-aristatum; *aristis* persistentibus (apteris v. alatis).

Frutices (Americæ æquinoctialis) sæpius decumbentes. Folia opposita, indivisa. Inflorescentia terminalis, subpaniculata, vel corymbosa. Corollæ albidæ. Paleæ receptaculi post lapsum pericarpiorum persistentes.

OBS. Of this genus I have examined specimens of three species in Sir Joseph Banks's Herbarium, differing from each other in several very remarkable characters.

1. *Salmea scandens*, (Decand. l. c.) in which the aristæ are equal and without any membranaceous border : stigmata remarkably dilated, tongue-shaped, obtuse, not hispid, obscurely papulose, and apparently without any terminal appendix : style dilated at the base into a hemispherical bulb which is truncated underneath.

2. *Salmea hirsuta*, (Decand. l. c.) whose aristæ are unequal ; the inner, which is the larger, being furnished with an evident ala ; the outer having a narrow margin only : stigmata sharp and spreading : style dilated into an ovate bulb which has an attenuated base.

3. *Salmea? curviflora* (nob.) differs from both the preceding in the tube of its corolla being remarkably bent outwards. In place of the inner arista there is a broad obtuse wing, of which the inner margin is straight and thickened, the outer continued down nearly to the base of the pericarpium : the outer arista is winged : and besides these, one or two minute processes are generally observable. Stigmata revolute.¹

¹ In the remarkable character of its re-curved florets, as well as in some other respects, this species of *Salmea* agrees with *Spilanthus arbores* of George Forster (in Commentat. Gotting. ix, p. 66), of which he originally formed his genus *Laxmannia* ; from a very erroneous view of its structure, however, having described the Nectarium or glandula epigyna as a "germen superum ;" the real, though imperfect, germen with its two aristæ as a "perianthium bidentatum," and consequently referring the genus to Polygamia segregata.

When he afterwards corrected these errors, and reduced *Laxmannia* to *Spilanthus*, he did not discover that he had only the imperfect hermaphrodite or male plant before him.

That *Spilanthus arbores* is really diœcious, I have ascertained from the examination of numerous specimens collected by Sir Joseph Banks in the Island of St. Helena, where it forms a small tree called by the inhabitants White-wood. It is *Bidens arborea* and perhaps also *Spilanthus tetrandrus* of Dr. Roxburgh's List of Plants appended to General Beatson's Tracts on St. Helena ; the former being probably the female, the latter a starved variety of the male plant.

In re-establishing *Spilanthus arbores* as a genus, sufficiently distinct from *Bidens*, *Spilanthus*, and *Salmea*, it will not, I conclude, be considered expedient to recur to Forster's name *Laxmannia*, which as far as relates to this plant is connected only with a series of blunders, was abandoned by the author himself, and has since been applied to another genus already generally adopted. It may be distinguished by the following character, and named

PETROBIUM.

Involucrum polyphyllum subduplici serie: exteriore breviori, foliis paucioribus. *Receptaculum* paleaceum, planiusculum. *Flosculi* dioici, tubulosi,

In the 12th edition of *Systema Naturæ*, Linnæus added to his genus *Calca* a fourth species, namely *Calea scoparia*; for what reason it would be difficult to discover, as it does not resemble, either in its fructification or habit, any of the three genera of which, as has been shown, *Calea* was originally composed. This fourth species, which he had at first referred to *Chrysocoma*,¹ is now known to be diœcious;—Browne, by whom it was first described and figured, and one of whose specimens I have examined, Linnæus, and even Swartz when he published his *Observationes Botanicae*, being acquainted with the male plant only; which, however, all of them considered hermaphrodite: nor is there any reason to doubt that Gærtner's genus *Sergilus* is also the male of this species; although he has ventured to describe the colour of the embryo, deceived, probably, by the size of the imperfect ovarium, and the colour of its inner surface.

Professor Swartz has since given a more satisfactory account of *Calea scoparia*, and has referred it to *Baccharis*,² to which genus as Richard³ and Jussieu⁴ have proposed to limit it, namely to the diœcious species of America, it unquestionably belongs. This limitation of *Baccharis* it may, upon the whole, be expedient to adopt; by doing so, however, a name of Dioscorides is applied to a genus of plants found only in the new continent; while, notwithstanding the contrary opinion is expressed by M. de Jussieu,⁵ sufficient distinctions exist between those species

4-fidi: *Masculi*: Antheris exsertis; Stigmatibus acutis hispidulis: *Feminei*: Staminibus sterilibus; Stigmatibus acutis recurvis. *Achenium* v. parallelo-compressum v. angulatum; angulis (2-3) aristatis: *aristis* persistentibus, antrorsum denticulatis.

Arbor (Insulæ S^{tæ} Helenæ). Folia opposita, indivisa. Panicula terminalis, brachiata. Involucrum oblongum. Paleæ receptaculi squamis involucri sub-similes. Corollulæ ochroleucæ, tubo arcuato-recurvo (ut capitulum primo intuitu radiatum videatur). Mas. Antheris nigricantibus, basi emarginatis, appendice apicis brevissimo, acuto; loculis vestigio septi longitudinalis instructis. Fem. Staminibus sterilibus distinctis, antheris sagittatis cassis.

¹ Amœn. Acad. v, p. 404, et Syst. Nat. ed. 10, vol. ii, p. 1206.

² Flor. Ind. Occident. iii, p. 1339.

³ Mich. Flor. Bor.-amer. ii, p. 125.

⁴ Annal. du Mus. d'Hist. Natur. vii, p. 385.

⁵ l. c.

of *Baccharis* from which the Linnean character was taken, and *Conyza* when reduced to its original species, *C. squarrosa* and *bifrons*, and a few others since added to the genus: for these differ from *Inula* chiefly in the extreme shortness of their ligulæ.

As no satisfactory character has hitherto been given of *Baccharis*, that will serve to distinguish it, as now limited, from the diœcious *Gnaphalia*, I propose the following.

115]

BACCHARIS.

(Richard) in *Michaux Amer.* ii, p. 125. *Jussieu in Annal. du Mus. d'Hist. Nat.* vii, p. 385. *Molina Ruiz et Pavon Prodr. Flor. Peruv.* 111. *Baccharidis species Linn.*

Involucrum imbricatum. Receptaculum nudum. Flosculi tubulosi, dioici. Masculi: antheris exsertis, basi muticis; stigmatibus appendice acuto hispidulo; pappo subpenicillato. Feminei filiformes; pappo capillari.

Frutices (Americæ æquinoctialis et temperatæ). Folia alterna, raro opposita, in quibusdam minuta vel nulla, ramis tunc foliaceo-alatis. Inflorescentia terminalis rariusve lateralis, corymbosa, nunc fasciculata. Involucris subovati v. oblongi squamæ semiscariosæ, margine simplici. Mas pappo cinereo. Fem. limbo minuto 2-3dentato, staminibus sterilibus nullis: pappo elongato.¹

Willdenow, in his edition of *Species Plantarum*, has retained the four Linnean species of *Culea*, and added to

¹ I have observed another diœcious genus with naked receptacle, capillary pappus, and a habit nearly similar to that of *Baccharis*, of which *Baccharis nereifolia* Linn. is the only published species. It may be named

BRACHYLÆNA.

Involucrum imbricatum, squamis coriaceis. Receptaculum nudum. Flosculi dioici. Masculi: antheris exsertis, basi bisetis. Feminei angustiores, limbo 5-fido; filamentis sterilibus: stigmatibus linguiformibus imberbibus. Pappus utriusque sexus pilosus scaber.

Arbusculæ vel Frutices (Africæ australis) subtomentosi. Folia alterna integerrima vel dentata. Inflorescentia terminalis, subracemosa. Involucra subovata, breviter: squamis ovatis, textura uniformi.

them an equal number, not one of which belongs to any of the genera formed by the original species, but to four others equally distinct.

The *first* of these additional species, taking them in the order in which Willdenow has arranged them, is *Calea aspera*, which he adopted from Jacquin; by whom it is well described and figured, though erroneously referred to *Calea*.

This, and not (as M. Richard has supposed) the nearly related species of North America, is what Linnæus originally intended by his *Bidens nivea*, as appears by the specimen in his Herbarium; by his original reference to Vaillant's "*Ceratocephalus foliis cordatis s. triangularibus flore albo*,"¹ described from a specimen in Surian's Herbarium; and by his afterwards adding as varieties of his species the two plants from Carolina figured in Hortus Elthamensis.

Calea aspera is abundantly distinct from *Bidens*, and has very little affinity with any of the original species of *Calea*, and least of all with *C. jamaicensis*, from which the character was taken. Since its appearance in Willdenow's work, however, it has been continued in this genus, in most of the recent catalogues of Gardens, as those of Desfontaines, Decandolle, and the second edition of Mr. Aiton's Hortus Kewensis; and Lamarck in his Illustrationes Generum has copied Jacquin's figure of it, apparently as the principal example of the genus *Calea*.

It is certainly now too late to rectify to the name of *Amellus*, under which Browne, as I have already attempted to prove, first proposed this plant as a distinct genus; Linnæus having soon after given that generic name to two very different plants, to one of which it is still applied; and the real plant of Browne having till now been mistaken, owing in part to his having entirely overlooked the pappus which is deciduous.

Bidens nivea, however, as long ago as 1784 was described by Von Rohr, and published by him in 1792 in the second

¹ Act. Paris. 1720, p. 327.

volume of the Transactions of the Natural History Society of Copenhagen, as a distinct genus, under the name of *Melanthera*: and in 1803 by Richard, in Michaux's *Flora Boreali-Americana*, where it is called *Melananthera*, and ¹¹⁷ where the two species included by Linnæus in his *Bidens nivea* are for the first time distinguished: and lastly this genus, as named and determined in the work of Michaux, is adopted by Persoon in his Synopsis.

But as both Von Rohr and Richard have given only the natural character of the genus, and the essential character proposed by Persoon is not altogether satisfactory, I have added the following, and adopted the more generally received name of

MELANANTHERA.

(Richard) in Michaux *Amer.* ii, p. 106. *Melanthera* Von Rohr in *Kiobenh. Naturhist. Selskab. bind.* ii, hefte 1, p. 213. Amellus Browne *Jam.* 317. *Bidentis* species Linn. *Calea* species Jacquin.

Involucrum duplici serie polyphyllum, subæquale. *Receptaculum* paleaceum, convexum, paleis foliaceis. *Flosculi* tubulosi, uniformes, hermaphroditi. *Achenium* turbinatum angulatum vertice depresso. *Pappus* e setis (2—18) scabris, distinctis, deciduis.

Herbæ (Americæ æquinoctialis et temperatæ) *pubescentes, scabræ*. *Folia opposita, indivisa v. sublobata*. *Capitula terminalia, pedunculis unifloris, elongatis, ternatis, geminisve*. *Involucrum foliaceum*. *Receptaculi hemisphærici paleæ foliolis involucri subsimiles*. *Corollæ albide*. *Antheræ nigricantes, appendicibus apicis albidis, basi muticæ; paulo post expansionem corollæ exsertæ, dein (contractione filamentorum) fauce inclusæ*. *Stigmata appendice acuto hispidulo, post retractionem tubi antherarum exserta; demum subinclusa*.¹

¹ In the extensive collection of plants made by my lamented friend Dr. Smith, on the banks of the Congo, I have observed a Syngenesious genus, which, though belonging to Polygamia superflua and having yellow flowers, is in other respects so nearly related to *Melananthera*, that had it been found with

Obs. In Von Rohr's natural character of *Melanthera* the Nectarium, or glandular body sheathing the base of the style, is introduced, which is the earliest notice I have yet found of this organ in Compositæ, except in Batsch's Analysis Florum, published in 1790, where it is both described and figured in *Coreopsis tripteris*. The merit, however, of establishing its nearly universal existence in the hermaphrodite florets of this extensive class belongs to M. Cassini.

Both Von Rohr and Richard in their characters of *Melanthera* have described the antheræ as shorter than the corolla, which is indeed the case in a particular state of the flower; immediately after its expansion, however, they project considerably, and again become inclosed in the more advanced stage. This fact has been noticed by Jacquin,¹ who considers the final inclosure of the antheræ to be owing to the elongation of the corolla. But the actual increase in length of the corolla is very slight, and by no means sufficient to account for the appearance; the real cause of which is a considerable, and I believe a gradual, contraction of the filaments. This economy is not unfrequent in Compositæ, especially in the tribe of *Heliantheæ*, to which *Melanthera* belongs.

In M. Cassini's Memoir on the Stamina of Compositæ the retraction of antheræ is not expressly noticed. This

ripe seeds only, it would certainly have been referred to it. The following characters, however, prove it to be sufficiently distinct. It may be named

LIPOTRICHE.

Involucrum duplici serie imbricatum, subæquale. *Receptaculum* convexum, paleis foliaceis, distinctis. *Capitulum* radiatum. *Ligulæ* (simplici serie) femineæ. *Flosculi* hermaphroditi, *stigmatibus* appendice acuto hispidulo. *Achenia* subuniformia, turbinata; *Pappo* setaceo, caduco.

Herbæ (Africæ æquinoctialis) *Folia opposita, indivisa*. *Pedunculi terminales, terni*. *Involucra breviora, foliacea*. *Paleæ receptaculi carinatae, nervosæ, aculeæ*. *Corollulæ flacæ*. *Ligulæ elongatæ, 3-dentatæ*. *Antheræ nigricantes, subinclusæ, basi muticæ*. *Achenium obtusè tetragonum*. *Pappus, in disco verticis depressi, brevis, e setulis simplici serie, numerosis (8—10), distinctis, denticulatis, caducis vel deciduis*.

Melantheræ proximè accedit: affinis quoque *Ecliptæ* Linn. *Wedeliæ* Jacq. et *Diomedæ* Cassini (in Journ. de Phys. tome lxxii, p. 145), sed ab his omnibus satis distincta videtur.

¹ Collect. ii, p. 291, Ic. Rar. iii, t. 583.

appearance, however, can hardly have escaped so accurate an observer; and his opinion respecting its cause may perhaps be inferred from an observation he has made on the stamina of the tribe in which it is most remarkable, namely *Heliantheæ*; whose filaments below the joint, he says, wither very soon after fœcundation.¹ To this withering, which he does not mention as occurring in any other tribe, the phænomenon in question may be supposed to be ascribed.

But it appears to me, that the contraction or collapse of the filaments, from their previous state of extension, is a vital action, and not the effect of withering or decay, which, however, speedily follows it. For the contraction may in great part be prevented by the separation of the floret, when the filaments are in the state of extension: and in many genera of *Compositæ* the antheræ are never retracted, but continue to project till they fall off with the corolla.

This contraction is also analogous to the more evident motion or irritability of the filaments long ago noticed by Borelli and Alexander Camerarius² in *Cinarocephalæ*; and more fully described in the same tribe by Dal Covolo;³ whose observations are confirmed and extended to other subdivisions of *Compositæ* by Koelreuter.⁴ A similar contraction and irritability of the style has been lately described by Mr. Ker in certain species of *Arctotis*.⁵

The second species added to the genus by Willdenow is *Calea lobata*, which Linnæus, from the general appearance, I conclude, rather than from actual examination of the plant in Clifford's Herbarium, had referred to *Conyza*; and having no specimen in his own Herbarium, the twofold error of supposing it to belong to *Polygamia superflua*, and to have a naked receptacle, remained uncorrected in all his subsequent works.

¹ Journal de Physique, tome lxxviii, p. 278.

² Ephemerid. Acad. Nat. Curios. cent. ix et x, p. 194.

³ Discorso della Irritabilita d' alcuni Fiori. Firenze, 1764.

⁴ Von Einigen das Geschlecht der Pflanzen betreffenden versuchen. 3. fortsetz. p. 125.

⁵ Botanical Register, i, 34.

Its real structure was first pointed out by Professor Swartz, who consequently referred it to *Calea*, with the character of which it exactly agrees. This alteration is adopted in the first edition of Hortus Kewensis, where the generic character of *Calea* is modified, to admit those species that are without pappus; and by Gærtner, who limits the genus to *C. lobata* and *C. jamaicensis*, as the only species that correspond with the Linnean character. But as *C. jamaicensis*, the original species of *Calea*, has been shown to have a pappus of a very different kind, it becomes necessary to give a new name to *Calea lobata*; and some additions being also wanting to its generic character, I propose the following, and the name of

NEUROLÆNA.

Calea Gært.

Involucrum imbricatum, foliaceum. *Receptaculum* paleaceum, planiusculum. *Flosculi* tubulosi, uniformes, hermaphroditi. *Antheræ* inclusæ, basi muticæ (emarginatæ). *Stigmata* acuta, recurva. *Pappus* capillaris, denticulatus, persistens.

Frutex (Americæ æquinotialis) erectus. Folia alterna, indivisa, et lobata. Corymbus terminalis, compositus. Involucrum subovati foliola obtusa, nervosa. Paleæ receptaculi involucri subsimiles. Corollulæ flavæ.¹

¹ There are two other genera in many respects agreeing with the character here given of *Neurolæna*, which it is necessary to point out. The first is *Carphophorus* of M. Cassini (in Bulletin des Sciences, 1816, p. 198), sufficiently distinct in having the stigmata of *Eupatorium* or *Liatris* with the habit of the latter, from some species of which it differs only in its receptacle having paleæ. The second, not yet described, may be named

PIPTOCARPHA.

Involucrum imbricatum, turbinatum, scariosum. *Receptaculum*: paleis distinctis. *Flosculi* tubulosi, uniformes, limbo revoluto. *Antheræ* exsertæ, basi bisetæ. *Stigmata* filiformia, acuta, hispidula. *Pappus* pilosus.

Frutex (Brasiliensis) ramosissimus, decumbens? Folia alterna, integerrima, subtus incana. Involucra axillaria et terminalia, fusciculata, glabrata, squamis sessilibus obtusiusculis nerviis, textura uniformi. Paleæ receptaculi squamis

The *third* species, *Calea pinifolia*, is adopted from Forster's *Florulæ Insularum Australium Prodrum*.

The specimen of this plant in George Forster's Herbarium (now forming part of the extensive collection of Mr. Lambert) is very imperfect; it evidently, however, belongs to the same species with a more complete specimen received, without a name, from Forster by Sir Joseph Banks, in whose Herbarium I have examined it, and ascertained that it has a naked receptacle. It therefore cannot be a species of *Calea*, which I have no doubt Forster considered it merely from a certain degree of resemblance to his *Calea leptophylla*. From the structure of its stigmata, antheræ, and involucre, *Calea pinifolia* belongs, indeed, to a very different tribe, and might even be referred to *Gnaphalium* as it at present stands. But this extensive and ill-defined genus evidently requires reformation; and ¹²²¹ if the necessity for its subdivision be admitted, it will also, I believe, be found most expedient to apply the name *Gnaphalium* to that section to which *G. luteo-album*, *sylvaticum*, and *uliginosum* belong, and which is characterised by its naked receptacle, its involucre connivent at top and of equal height with the truncated capitulum, which consists of numerous filiform female florets in the circumference, with a smaller number of hermaphrodite florets in the disk, both of them ripening seeds and having a sessile capillary deciduous pappus.

To *Gnaphalium* so limited *Calea pinifolia*, a shrub with nearly acerose leaves, and in which all or most of the flosculi are hermaphrodite and the radii of the persistent pappus somewhat thickened upwards, cannot be referred.

It seems, however, to approach more nearly to *Antennaria*, a genus separated from *Gnaphalium* by Gærtner, but which, as he has proposed it, consists of three tribes of

intimis involucri subsimiles, et unâ cum iisdem deciduæ. Corollulæ glabræ. antherarum integerrimæ. Pappus albus, radiis simplici serie.

Obs. I have not seen perfect seeds; and as even in the unripe fall off along with the inner squamæ of the involucre, and the antheræ; in a remarkable degree, it is possible the plant here described may be only the male of a diœcious species: it certainly, however, belongs to a genus not before published.

plants sufficiently dissimilar in habit and structure to justify a further subdivision; and, what is remarkable, none of them entirely agreeing with his generic character.

The first tribe consists of herbaceous plants, natives of Europe and North America, having the male and female flosculi in distinct involucri and on different individuals. To this genus the name ANTENNARIA¹ may remain, though

¹ ANTENNARIA.

Antennariæ species. *Gærtner.* Gnaphalii species. *Linn. Jussieu.*

Involuerum imbricatum, scariosum, coloratum. *Receptaculum* epaleatum, scrobiculatum. *Flosculi* dioici. *Masculi*: *antheris* basi bisetis: *stigmatibus* truncatis: *Pappo* vel penicillato v. apice incrassato. *Feminei* filiformes, limbo parvo: *staminum* rudimentis nullis: *Pappo* capillari.

Herbæ perennes, tomentosæ, incanæ. *Folia* plana, adulta sæpe super glabruscula; radicalia in plerisque latiora. *Inflorescentia* corymbosa raro solitaria. *Involueri* turbinati vel quandoque hemisphærici squamæ e basi calycina superne coloratæ (albæ v. purpurascetes). *Coruliulæ* flavæ. *Antheræ* semi-exsertæ. *Pappus* murium niveus, opacus.

Obs. *Gnaphalium margaritaceum*, which I have referred to this genus, was first described by Clusius; from whose account it appears to have been introduced into the English gardens from America towards the end of the sixteenth century.

It has ever since been very generally cultivated, as an ornamental plant, both in this country and on the continent of Europe; and has a place in several of the European Floras, as well as in those of North America. It is surprising, therefore, that hitherto the male plant only should have been observed, uniformly, however, considered as hermaphrodite, except by M. Cassini, who in his first memoir on *Synantheræ* (in *Journal de Physique*, tome lxxvi, p. 200) suspects it to be male, from the imperfect appearance of the ovary.

That this species of *Gnaphalium* is really diœcious, I learned several years ago from an inspection of a specimen of the female plant in the Herbarium of Sir Joseph Banks, who found it on the banks of the Rymney in Glamorgan-shire, where the plant was first observed by Lhwyd. I have since received several specimens of both sexes from Mr. Bicheno, to whom I had mentioned this fact, and who obligingly undertook to observe the different states of the plant in the same place, where it seems to be really indigenous. I have never been able to discover any female florets in the circumference of the capitulum of the male plant; but in the centre of the female capitulum I have always found two or three imperfect male florets, whose antheræ, although cohering and of the usual form, appear to be destitute of pollen.

The separation of sexes in a still more common plant of this class, namely, *Serratula tinctoria*, has been equally overlooked.

All the authors who have noticed this species, which is included in almost every European Flora, as well as in more than one recent Monograph of the genus, have considered it as hermaphrodite, while it really belongs to *Polygamia diœcia*, or has its perfect sexual organs on different plants. The hermaphrodite plant, apparently perfect, but which I believe very seldom ripens seed, is well figured by Schkuhr (in *Botanisches Handbuch*, tab. 234); and the female, whose stigmata are remarkably developed and undulated, while the

^{123]} descriptive of the pappus of the male flower only. Its species are *Gnaphalium dioicum* Linn., *alpinum* L., *carpaticum* Wahlenberg, *plantagineum* L., and *margaritaceum* L.

The second tribe, consisting of *Gnaphalium Leontopodium* ^{124]} and *Leontopodioides*, which may be called LEONTOPIDIUM, is in affinity intermediate between *Antennaria* and *Gnaphalium* as here limited, but has sufficient characters to distinguish it from both.

The third tribe has been found only in South Africa, and consists of shrubs with small rigid heath-like leaves, of which the margins are incurved, the upper surface tomentose, and the under convex and nearly smooth; but by a remarkable twisting they are in most of the species resupinate; a character which seems to have been overlooked in all the described species; namely, *Gnaphalium muricatum*, *mucronatum*, and *seriphioides*. In this tribe, or genus, which may be named METALASIA, the involucre is generally cylindrical, and in most of the species has a short radius formed by the spreading coloured laminæ of the inner scales; the flosculi are few in number, and all hermaphrodite; and the radii of the pappus, which fall off separately, are either thickened or more strongly toothed at top.

Calea pinifolia does not even belong to this genus, though it has a nearly similar habit; but the margins of its leaves are revolute, and their tomentum chiefly on the under surface. In these respects, as well as in the principal characters of fructification, it agrees with several shrubs, chiefly of New Holland and Van Diemen's Island; among which are *Eupatorium ferrugineum*, *Eupatorium rosmarinifolium*, and *Chrysocoma cinerea* of M. Labillardière. Part of these have the inner squamæ of the involucre simple, as seems to be the case in *Calea pinifolia*;

antheræ are evidently imperfect, and which generally produces ripe seeds, is represented in English Botany (tab. 38), in Flora Danica (281), and probably also in Svensk Botanik (170). For my knowledge of this fact respecting *Serratula tinctoria* I am indebted to the Rev. Robert Bree of Camberwell, who pointed out to me both its states, which he was then disposed to consider as distinct species.

while in others, as the two species referred to *Eupatorium* by M. Labillardière, they form a short radius. These I am inclined to consider merely sections of one and the same genus, which may be distinguished by the following character, and named

OZOTHAMNUS.

[125]

Involucrum imbricatum, scariosum, coloratum. *Receptaculum* epaleatum, glabrum. *Flosculi* (pauciores quam 20) tubulosi, vel omnes hermaphroditi, vel paucissimi feminei angustiores in ambitu. *Antheræ* (inclusæ), basi bisetæ. *Stigmata* apice obtuso subtruncato hispidulo. *Pappus* sessilis, pilosus, nunc penicillatus, persistens.

Frutices (Novæ Hollandiæ et Novæ Zelandiæ, vix Africae australis,) *graveolentes, tomentosi*. Folia *sparsa, integerrima, marginibus sæpius recurvis*. Inflorescentia *terminalis, corymbosa v. congesta*. Involucra *alba v. cinerea: squamis intimis nunc conformibus et conniventibus; nunc laminis patulis niveis radium brevem obtusum efformantibus*. Corollulæ *luteæ*. Pappus *albus*.

The fourth species added to *Calea* by Willdenow is *Calea leptophylla* of Forster, whose specimens I have examined in Mr. Lambert's Herbarium. Amongst Forster's drawings, formerly referred to, there is a coloured figure of this plant, by which it appears that he originally considered it to belong to *Gnaphalium*. From this genus he afterwards removed it, probably on finding it referred to *Calea* in the collection of Sir Joseph Banks, by whom it was discovered in New Zealand in a more perfect, at least in a more luxuriant state.

This plant, though agreeing with *Calea* in every part of the Linnean essential character, differs remarkably from it in other points of nearly equal importance, as well as in habit; and along with *Calea aculeata* of M. Labillardière, and several other species also natives of New Holland and Van Diemen's Island, constitutes a genus very nearly

related to *Ozothamnus*, from which it is to be distinguished chiefly by the paleæ of its receptacle.

^{126]} I propose to name this genus in honour of *M. Henri Cassini*, whose well-conducted investigation of Compositæ has already thrown much light on the structure and economy of the more important parts of fructification of this difficult class: and especially of those organs from which the distinguishing characters of *Cassinia* are here derived.

I shall add the characters of the species of this genus, which, like *Ozothamnus*, admits of subdivision into two sections; and I have appended to it *Calea spectabilis* of Labillardière, a plant corresponding with it in character, but differing very much in habit from all the other species.

CASSINIA.

Caleæ sp. *Labillardière*.

Involucrum imbricatum, scariosum, pauciflorum. *Receptaculum*: *paleis* distinctis, squamis intimis involucris subsimilibus. *Flosculi* tubulosi, vel omnes hermaphroditi vel paucissimi feminei angustiores in ambitu. *Antheræ* (inclusæ) basi bisetæ. *Stigmata* apice obtuso subtruncato hispidulo. *Pappus* pilosus v. penicillatus, persistens.

Frutices. *Folia* sparsa, sæpius angustata, marginibus recurvis. Inflorescentia terminalis, corymbosa rariusve paniculata. *Involucra* alba nunc cinerea raro aurea; squamis intimis sæpius apice conniventibus, nunc patulis et radium brevem obtusum efformantibus.

† *Involucrum radiatum* (squamis intimis apice patulis).

1. *C. leptophylla*, foliis lineari-lingulatis subter ramulisque incanis, corymbis terminalibus, involucris turbinatis.

Calea leptophylla. *Forst. Prodr. n.* 287. *Willd. Sp. Pl.* iii, p. 1796. *Persoon, Syn.* ii, p. 406. *Poiret, Encycl. Suppl.* ii, p. 28.

^{127]} *Loc. Nat.* Novæ Zelandiæ campi arenosi prope

Tolaga, &c. *D. Banks.* Prope Queen Charlotte's Sound.
J. R. et G. Forster. (v. s. in Herb. Banks et G. Forster.)

† † *Involucrum connivens.*

A. Fruticosæ.

2. *C. denticulata*, foliis ovalibus oblongisve acutis spinuloso-denticulatis subter tomentosis, corymbis compositis, involucris hemisphæricis.

Loc. Nat. Novæ Hollandiæ ora orientalis prope Port Jackson. *David Burton.* (v. s. in Herb. Banks.)

3. *C. longifolia*, foliis lanceolato-linearibus elongatis lævibus subter tomentosis, corymbis decompositis, involucris turbinatis.

Loc. Nat. Novæ Hollandiæ ora orientalis prope Port Jackson; in dumetis. (v. v.)

4. *C. aurea*, foliis lanceolato-linearibus elongatis lævibus subter glandulosis, corymbis decompositis, involucris ovalibus: squamis apice aureis.

Loc. Nat. Novæ Hollandiæ ora orientalis prope Port Jackson; in sylvis et dumetis. (v. v.)

5. *C. aculeata*, foliis angusto-linearibus margine revolutis super hispidulis subter ramulisque incanis, corymbis compositis decompositisve congestis, involucris turbinatis.

Calea aculeata. *Labill. Nov. Holl.* ii. p. 41, t. 185.
Persoon Syn. ii. p. 406. *Poiret Encycl. Suppl.* ii. p. 28.

Loc. Nat. Insula Van Diemen; in dumetis et ad ripas fluv. (v. v.)

6. *C. affinis*, foliis angustato-linearibus margine revolutis super hispidulis subter concoloribus, corymbis decompositis congestis, involucris turbinatis.

Loc. Nat. Novæ Hollandiæ ora orientalis prope Port Jackson; in dumetis. *D. G. Caley.* (v. s.)

Obs. *C. aculeatæ* nimis affinis.

7. *C. lævis*, foliis angustissime linearibus margine revo-

lutis super lævibus subter ramulisque incano-tomentosis, corymbis compositis, involucris congestis cylindraceis.

Loc. Nat. Novæ Hollandiæ ora australis; in campis ad radices montium prope ortum Spencer's Gulph. (v. v.)

8. *C. arcuata*, foliis angustissime linearibus margine revolutis super lævibus subter ramulisque incano-tomentosis, panicula pyramidata, involucris spicatis cylindraceis arcuatis.

Loc. Nat. Novæ Hollandiæ ora australis; in campis elevatis prope ortum Spencer's Gulph. (v. v.)

9. *C. quinquefaria*, foliis angustissime linearibus super ramulisque glabris, panicula decomposita, involucris turbinatis: squamis 5-fariis.

Loc. Nat. Novæ Hollandiæ ora orientalis prope Port Jackson; in montosis. *D. G. Caley.* (v. s.)

B. *Herbacea.*

10. *C. spectabilis*, panicula decomposita, foliis lanceolatis decurrentibus subter ramisque lanatis.

Calea spectabilis. *Labill. Nov. Holl.* ii. p. 42, t. 186. *Persoon Syn.* ii. p. 406. *Poiret Encycl. Suppl.* ii. p. 28.

Loc. Nat. Novæ Hollandiæ ora australis; in sylvis dumetisque prope Memory Cove, Port Lincoln, &c. legi. In Insula Van Diemen a D. Labillardière detecta. (v. v.)

^{129]} Since the publication of Willdenow's *Species Plantarum* very few alterations have been made in the genus *Calea*.

In Persoon's Synopsis two of the species are excluded; namely, *Calea scoparia*, which, following Swartz, he has referred to *Baccharis*; and *Calea aspera*, adopted from Richard as a species of *Melananthera*. The additional species in the work referred to are *C. cordifolia* of Swartz, already noticed as a genuine *Calea*; *C. aculeata* and *spectabilis* of Labillardière, which belong to *Cassinia*; and *C. cordata*, adopted from Loureiro, of whose plant nothing is known except from the short description in *Flora Cochinchinæ*.

chinensis, which is only sufficient to render it probable that it neither belongs to *Calea* as I have proposed to limit it, nor to any of the genera hitherto confounded with it.

M. Poiret, in the Supplement to the Botanical Dictionary of the Encyclopédie Méthodique, has under the article *Calea* retained all the species of this genus given by Persoon; and also *Calea aspera*; which, however, he has in a subsequent article correctly referred to *Melananthera*.

Connected with the proper subject of this paper, I shall describe and add some observations on a plant lately sent from Brazil by Mr. Sellow; which, though not strictly referable to Compositæ, probably belongs to a genus at present included in this family; and conclude with a few remarks on the structure and affinities of *Brunonia*.

I have named the Brazil plant

ACICARPHA SPATHULATA.

Herba annua? glaberrima, ramosa, diffusa. *Rami* ascendentes, angulati. *Folia* sparsa, petiolata, exstipulata, spathulata mucronulo brevissimo, sesquiuncialia, crassiuscula? glauca? sæpiùs integerrima; inferiora quandoque extra medium dentata. *Petioles* lineares basi parum dilatata^[130] semiamplexicauli; inferiores elongati; superiores plerumque folio aliquoties breviores. *Capitula* solitaria, nunc oppositifolia pedunculata, nunc terminalia subsessilia, basiflora, ovata, flava. *Involucrum* simplici serie pentaphyllum, capitulum floridum superans, foliaceum; foliolis inæqualibus spathulatis sessilibus integerrimis ipsa basi connatis. *Receptaculum* subulato-conicum, paleaceum. *Paleæ* lanceatæ mucronulatæ, inter flosculos hermaphrodito-masculos magis manifestæ, inter hermaphroditos passim abortientes. *Flosculi* tubulosi, uniformes, glabri.

Flosculi ambitus, duplici triplicive serie, hermaphroditi, utroque organo perfecto. *Corollæ Tubus* gracilis cylindraceus, cum ovario continuus, basique stylo accretus, per lentem 10-striatus. *Limbus* infundibuliformis, 5-fidus, æstivatione

valvata; laciniis semilanceolatis, planis, trinerviis; nervis lateralibus margini parallelo-approximatis, indivisis, apice confluentibus, e nervis alternis tubi infra sinus furcatis ortum ducentibus.

Stamina 5 epipetala, limbi laciniis alternantia.

Filamenta inferne cum tubo arcte connata, superne libera fauci quasi inserta, invicem cohærentia in tubulum 5-dentatum ipsis apicibus, subita mutatione texturæ, articulatis; basi intus incrassatum arcis 5 oblongis cum filamentorum axibus alternantibus.

Anthera continuæ, lineares, dimidio inferiore arcte cohærentes, superiore liberæ; biloculares, loculis longitudinaliter dehiscentibus, valvula interiore angustiore, receptaculo pollinis utriusque loculi longitudinali septiformi: basi emarginatæ, lobulis posticis acutiusculis brevibus polliniferis; apice simplices connectivo ultra loculos hand producto. *Pollen* subglobosum, per lentem pluries augeantem obsolete angulatum.

Ovaria connata, singula coronata calyce 5-fido dentibus spinescentibus cum laciniis limbi corollæ alternantibus; ¹³¹⁷ monosperma, ovulo ovato pendulo, paulo infra apicem affixo funiculo crassiusculo ex ipso apice angustato cavitatis orto; chorda vasculari a puncto insertionis ad extremitatem inferiorem ejusdem lateris attingenti. *Stylus* filiformis glaber, inferne cum basi tubi corollæ connatus. *Stigma* simplex obtusum hispidulum.

Flosculi superiores numerosi hermaphrodito-masculi, paulo minores hermaphroditis, calycis laciniis submembranaceis; ovariis (pariter connatis) imperfectis, sæpius absque ovulo.

Pericarpia (flosculorum ambitûs): *Achenia* conferminata, singula coronata calyce aucto 5-spinoso, spinis patulis conico-subulatis e substantia suberosa axi solidiori rigida.

Semen pendulum, ovatum extremitate superiore acuminato: *testa* membranacea: *membrana propria* nucleo adhærens. *Albumen* figura seminis, carnosum, copiosum, album. *Embryo* axilis, subcylindraceus, longitudine fere albuminis; albus, dicotyledoneus. *Cotyledones* lineares,

obtusæ, plano-convexæ, vix longitudine *Radiculæ* cylindraceæ, superæ.

Notwithstanding the great difference between my account of this plant and that given by M. de Jussieu of *Acicarpa tribuloides*, I have very little doubt that they both belong to the same genus; though from the above description it is evident that *Acicarpa spathulata* is not referable to Compositæ. To this plant *Calycera* of Cavanilles, in the seeds of which M. Correa has found albumen, seems to be very nearly related; and a third genus, probably referable to this group, is *Boopis*, described by M. de Jussieu in the same Memoir with *Acicarpa*. The important characters, however, of the pendulous ovulum and inverted embryo remain to be ascertained in all these; and the presence of albumen in *Acicarpa tribuloides* (in *Acicarpa lanata* of Lagasca in Pers. Syn. ii. p. 488, if it really belong to this genus), and in both species of *Boopis*. Another question respecting the latter genus is, whether its capitulum be simple, as it certainly is in *Acicarpa spathulata*; or compound, as Jussieu's figure of *Boopis anthemoides* seems to indicate.

In the mean time, with the necessary knowledge of structure of *Acicarpa spathulata* only, I shall venture to propose this group as a distinct natural family to be placed between Compositæ and Dipsacæ; though upon the whole somewhat more nearly approaching to Compositæ. This family, if my conjectures respecting *Calycera* and *Boopis* should be hereafter verified, may be called CALYCERÆ; *Acicarpa* even as a generic name being barely tenable; provided the original species agrees with that here described: for on this supposition M. de Jussieu has mistaken the laciniae of the perianthium for paleæ of the receptacle, deriving the name of the genus from their form; and has entirely overlooked the real paleæ, which, though they could not have suggested this name, may however sanction its being retained, if it be not still better to change it to *Acicarpa*.

It will be attended with similar advantage to form a separate family of

BRUNONIA,

as a link of equal importance, connecting *Compositæ* with *Goodenoviæ*, but from both of which it is in many respects very distinct. As I have formerly described this genus, and made several observations on its principal affinities,¹ I shall here only state the more important relations and distinctions between it and those families to which it appears to me most nearly to approach.

Brunonia agrees with *Goodenoviæ* in the remarkable indusium of the stigma; in the structure and connexion of ¹³³³ the antheræ; in the seed being erect; and essentially in the æstivation of corolla. It differs from them in having both calyx and corolla distinct from the ovarium; in the disposition of vessels in the corolla; in the filaments being jointed at top; in the seed being without albumen; and in its remarkable inflorescence, compatible, indeed, with the nature of the irregularity in the corolla of *Goodenoviæ*, but which can hardly coexist with that characterizing *Lobeliaceæ*.²

With *Compositæ* it agrees essentially in inflorescence; in the æstivation of corolla; in the remarkable joint or change of texture in the apex of its filaments; and in the structure of the ovarium and seed. It differs from them in having *ovarium liberum* or *superum*; in the want of a glandular disk; in the immediately hypogynous insertion of the filaments; in the indusium of the stigma; and in the vascular structure of the corolla, whose tube has five nerves only, and these continued through the axes of the laciniae, either terminating simply (as is at least frequently the case in *Brunonia sericea*), or (as in *B. australis*) dividing at top into two recurrent branches forming lateral nerves, at first sight resembling those of *Compositæ*, but which hardly reach to the base of the laciniae.

It is a curious circumstance that *Brunonia* should so

¹ Prod. Flor. Nov. Holl. p. 589.

² See Flinders's Voyage to Terra Australis, ii. p. 559 [vol. i, p. 32].

completely differ from Compositæ in the disposition of vessels of the corolla, while both orders agree in the no less remarkable structure of the jointed filament; a character which had been observed in a very few Compositæ¹ only before the publication of M. Cassini's second Dissertation, where it is proved to be nearly universal in the order.

In the opposite parietes of the ovarium of *Brunonia* two nerves or vascular cords are observable, which are continued into the style, where they become approximated and parallel. This structure, so nearly resembling that of Compositæ, seems to strengthen the analogical argument in favour of the hypothesis advanced in the present paper—of the compound nature of the pistillum in that order, and of its type in phænogamous plants generally;—*Brunonia* having an obvious and near affinity to *Goodenovia*, in the greater part of whose genera the ovarium has actually two cells with one or an indefinite number of ovula in each; while in a few genera of the same order, as *Dampiera*, *Diaspasis*, and certain species of *Scævola*, it is equally reduced to one cell and a single ovulum.

Sir James Smith, in establishing *Brunonia* as a genus, is disposed to refer it to *Dipsaceæ*. To certain species of this order it, indeed, bears a striking resemblance in habit; it also very nearly agrees with them in its remarkable inflorescence; and one great objection to its union with them may be supposed to be removed in adopting M. Decandolle's account of their ovarium.

But as *Brunonia* differs from the whole order in the following characters, all of which are of primary importance;—namely, in the origin and æstivation of corolla; in the insertion and whole structure of stamina; in the indusium of the stigma; in the ovulum being inserted at the base of the cavity of the ovarium; in the erect embryo and want of albumen;—I continue to think that its proper place in the natural method is between *Goodenoviæ* and *Compositæ*.

¹ Batsch Anal. Flor. p. 107; et Schkuhr Handb. tab. 236 et 244.

I shall conclude this subject, by proposing a few queries respecting the indusium of *Brunonia* and *Goodenoviæ*.

Is this remarkable covering of stigma in these families merely a process of the apex of the style? or is it a part of distinct origin, though intimately cohering with the pistillum? On the latter supposition, may it not be considered as analogous to the glandular disk surrounding or ¹³⁵⁷ crowning the ovarium in many other families? And, in adopting the hypothesis I have formerly advanced¹ respecting the nature of this disk in certain families,—namely, that it is composed of a series of modified stamina,—has not the part in question a considerable resemblance in apparent origin and division to the stamina of the nearly-related family *Stylideæ*?

To render this supposition somewhat less paradoxical, let the comparison be made especially between the indusium of *Brunonia* and the imperfect antheræ in the female flowers of *Forstera*. Lastly, connected with this view, it becomes of importance to ascertain whether the stamina in *Stylideæ* are opposite to the segments of calyx or of corolla. The latter disposition would be in favour of the hypothesis. This, however, is a point which will not be very easily determined, the stamina being lateral. In the mean time, the existence and division of the *corona faucis* in *Stylidium* render it not altogether improbable that they are opposite to the segments of the corolla.

Since the preceding paper was submitted to the Society, M. Cassini has published² the substance of a Memoir, which he read to the Academy of Sciences of Paris in August last, on a new family of plants named by him *BOOPIDÆ*, and consisting of *Calycera*, *Boopis*, and *Acicarpa*. I have also, through the liberality of Messrs. de Jussieu, Desfontaines, and Baron Delessert, had the opportunity of examining specimens of *Acicarpa tribuloides* in

¹ Linn. Soc. Transact. x. p. 159. [*Ante*, p. 133.]

² Bulletin des Sciences, 1816, p. 160.

flower and fruit, of both species of *Boopis* in flower, and detached flowers and pericarpia of *Calycera*. In all of these I have found the ovulum pendulous; and in *Acicarpha* and *Calycera* an inverted embryo occupying the axis of a fleshy albumen. My conjectures, therefore, on their structure and relation to *Acicarpha spathulata* of the preceding paper, are completely verified by this examination, as well as by the observations of M. Cassini, who with his usual acuteness has detected the principal characters distinguishing *Boopideæ* from *Compositæ* and *Dipsacæ*, between which he has also placed them.

As M. Cassini's Memoir, though read subsequently to mine, is already published, the name *Calycereæ*, which I have proposed for this family, is superseded by that which he has given it.

But as his account of the order is by no means complete, several characters of considerable, though not primary, importance being entirely omitted, I may be allowed to add to my paper some remarks on the more essential points of resemblance and difference between it and the two families to which it is most nearly related.

The principal characters distinguishing *Boopideæ* from the whole of *Compositæ* are the pendulous ovulum and the albumen inclosing the embryo, of which the radicle points to the apex of the pericarpium. It appears to me necessary to state all these characters, and nearly in the terms in which they are here given: for, 1st, A pendulous ovulum most frequently, indeed, is not, however, invariably connected with *radicula supera*, though that direction of radicle might here, as well as in *Compositæ*, with confidence have been inferred from the vascular structure of the ovulum.¹ 2dly, Where the insertion of the ovulum is, as in this family, evidently below the upper extremity, the radicle

¹ Some of the indications in many cases afforded by the structure of the unimpregnated ovulum, of the position and direction of the parts of the future embryo, have hitherto been overlooked: the subject, however, for its elucidation requires details incompatible with the limits of the present communication. I have in another place (Flinders's Voyage to Terra Australis, ii. p. 601 [vol. i, p. 77]) thrown out a similar hint, which has probably attracted no attention, and must reserve the explanation of both for a separate essay.

¹³⁷¹ which points to this extremity cannot in strict propriety be described as directed towards the umbilicus. M. Cassini has not noticed the direction of the radicle; either from supposing it constantly connected with that of the ovulum, or, which is more probable, from not having ascertained it.

These distinctive characters may be considered as fully sufficient to authorise the separation of *Boopideæ* from *Compositæ*; yet the same differences exist between certain genera referred and really belonging to *Rubiaceæ* and the principal part of that order.

There are, however, three other characters unnoticed by M. Cassini, which distinguish the flowers of *Boopideæ* from the hermaphrodite flowers of the whole of *Compositæ*; namely, the accretion of the base of the style with the tube of the corolla; the absence of the epigynous disk or nectarium; and the longitudinal subdivision of each cell of the anthera by a "receptaculum pollinis," as in most other families, and of which, indeed, there seems to be the rudiment in the syngenesious genus *Petrobium*, described in the preceding paper.

In the partial cohesion of the antheræ, in which they resemble *Jasione*, they certainly differ from all known *Compositæ*: but as in certain *Compositæ* the antheræ are very slightly connected or entirely distinct;—this, though a remarkable circumstance, can hardly be employed as a distinguishing character.

The principal characters in which *Boopideæ* differ from the greater part, though not from the whole of *Compositæ*, are the corolla being continuous, or not jointed, with the ovarium; the antheræ having no membranaceous appendix at top; and the undivided stigma.

Boopideæ differ from *Dipsacæ* in the vascular structure and valvular æstivation of corolla; in the æstivation, insertion, and connexion of antheræ; in the absence of the partial involucre; and in having alternate leaves.

In adopting M. Decandolle's description of *Dipsacæ*,¹ they would differ also in the important character of "ova-

¹ Flor. Franc. 3^{me} ed. vol. iv. p. 221.

rium inferum." This distinction, however, is neither universal, nor I believe absolute in any case.

M. Auguste Saint Hilaire in his excellent Memoir on *Primulaceæ*,¹ while he admits the correctness of M. Pecandolle's account with respect to great part of *Dipsacæ*, has at the same time well observed, that in several species of *Scabiosa* the ovarium is entirely united with the tube of the calyx. But neither of these authors has remarked the curious, and I believe peculiar, circumstance, of the base of the style cohering with the narrow apex of the tube of the calyx, even in those species of the order in which the dilated part of the tube is entirely distinct from the ovarium.

This kind of partial cohesion between pistillum and calyx is directly opposite to what usually takes place, namely, the base of the ovarium being coherent, whilst its upper part is distinct. It equally, however, determines the apparent origin or insertion of corolla and stamina, producing the unexpected combination of "flos superus" with "ovarium liberum."

In the vascular structure of the corolla *Boopideæ* may be considered as essentially agreeing with *Compositæ*, in many of whose genera the middle nerves of the tube and segments are equally manifest. In stating the character derived from this source in either of these orders, it is not sufficient to describe the nerves of the laciniae only as M. Mirbel has done in his character of *Compositæ*,² and M. Cassini in that of *Boopideæ*: but it is also necessary to give their disposition in the tube or undivided part of the limb; there being instances in both families where the lateral nerves of the segments do not unite at top; and, as has been formerly remarked, several examples in other families of a nearly similar disposition in the segments, accompanied by a different disposition in the tube. To the examples of this kind formerly given, *Globularia cordifolia* may be added, in the segments of whose lower lip there are three simple nerves, of which the lateral do not unite at top, and con-

¹ Mem. du Mus. d'Hist. Natur. ii. p. 47.

² Elemens de Physiol. Veget. et de Botan. ii. p. 885.

tinue distinct nearly to the base of the tube, where they converge and appear to unite with the middle nerve.

In *Acicarpa* and *Boopis* the filaments appear to me jointed as in *Compositæ*; a character I have not been able to observe in the very few flowers which I have examined of *Calycera*.

In *Acicarpa* the florets of the circumference are hermaphrodite and apparently complete, the antheræ containing pollen and the ovaria producing seed; while those of the disk are male with an incomplete pistillum. Such an arrangement has not hitherto been observed in *Compositæ*, in which, wherever the central florets are male with an imperfect pistillum, those of the circumference are female with or without the rudiments of stamina.

The regularity in the order of expansion of flowers from the base to the top of the capitulum in *Acicarpa tribuloides* and *spathulata*, and the irregularity, approaching to the inverted order, which I have found to exist in both species of *Boopis*, seem to prove the capitulum to be simple in the former genus and compound in the latter, notwithstanding the great resemblance between their involucre. The exact nature of its composition, however, in *Boopis* can only be satisfactorily determined in recent specimens.

140] This irregular expansion in *Boopis*, which renders even the generic name improper, and at present the want of satisfactory characters to distinguish it from *Calycera*, are objections to the name M. Cassini has chosen for this family; while that of *Calyceræ*, which I have proposed, derived from the genus first described, and applicable to all the genera of the order, appears to me unexceptionable: especially as there seems no reason to doubt that the part which I have considered as calyx in *Boopideæ* is really such; its divisions being generally in equal number, and alternating with those of the corolla. It may be observed that a like alternation of the divisions of the pappus with the segments of the corolla obtains in those genera of *Compositæ* where both parts are in equal number. But in some cases, where the division of pappus is still further

reduced, the same alternation does not exist, especially in those genera having vertically compressed pericarpia and two aristæ, as *Spilanthus* and *Salmea*.

The absence of "discus epigynus" in *Boopideæ* is a necessary consequence of the accretion of the base of the style with the tube of the corolla. It seems to me, however, that a modification of the same organ may be traced in the five thickened areolæ observable within and near the base of the tube formed by the filaments in *Acicarpia spatulata*; and much more distinctly in the same situation in *Boopis balsamitifolia*, where they have the appearance of five adnate fleshy bodies alternating with the filaments.

This apparent decomposition of the glandular disk in *Boopideæ*, compared with its state in *Compositæ*, as well as its transposition and the alternation of its parts with the stamina, seem to give some additional support to the conjecture I have formerly hazarded in the paper on *Proteaceæ*, published in the Society's Transactions (vol. x, p. 159¹); namely, that in several families—for the hypothesis is not meant to be extended to all—this part, even in its simplest state, may be considered as formed of a series of modified stamina: Or, merely to state the facts from which the conjecture originates, that there are certain families in some of whose genera this organ exists in its simplest form, that of an undivided fleshy ring; while in other genera of the same families it consists of several distinct bodies alternating with the stamina, and in some cases putting on the appearance of barren filaments.

This hypothesis is chiefly applicable to families in which the number of stamina is equal to the divisions of one floral envelope only, the nectarium being supposed to be formed of the second series: but it receives its principal support from *Scitamineæ*,² where the glandular bodies belong actually to the same series with the perfect stamen.

I am aware at the same time of several objections to its generalisation. Thus, the nectarium or glandular disk exists in families where, though the stamina are definite,

¹ [*Ante*, p. 133.]

² See Flinders's Voyage to Terra Australis, ii. p. 574 [*vol. i*, p. 49].

they are equal in number to the divisions of calyx and corolla united; and moreover, in such families where it consists of distinct parts, these parts are placed where an addition to the number of stamina is least likely to take place, as in *Crassulaceæ*. Here, however, as in many other cases, the divisions of the disk are opposite to the ovaria; they may therefore be supposed more intimately connected with the pistilla than with the stamina; an opinion which I believe held, though not yet published, by the ingenious M. Decandolle with respect to *Ranunculaceæ*. In support of this opinion it may be noticed that in *Pæonia Montan*, where the disk or urceolus is in the state of the greatest development, when a multiplication of the pistilla takes place, which in the double-flowered varieties of this ^{142]} species it not unfrequently does by the addition of one or more inner series, the rudiments of an analogous disk are produced along with each of the additional series.

Yet, in opposition to this view, I have in a single instance found one of the divisions of the urceolus in *Pæonia Montan* changed into an anthera; and the divisions of the apparently analogous organ in *Aquilegia*, which in their usual state resemble barren filaments, have sometimes been observed with perfect antheræ.¹

¹ Schkuhr Ha dbuch, tab. 146.

CHARACTERS AND DESCRIPTIONS

OF

THREE NEW SPECIES OF PLANTS,

FOUND IN CHINA BY CLARKE ABEL, Esq.;

SELECTED FROM

A SMALL COLLECTION OF SPECIMENS,

THE ONLY PART OF HIS HERBARIUM THAT ESCAPED THE WRECK OF
THE ALCESTE.

BY

ROBERT BROWN, F.R.S.

[*Extracted from the 'Narrative of a Journey in the interior of China,'*
by CLARKE ABEL, Esq., pp. 374—379.]

LONDON.

1818.

CHARACTERS AND DESCRIPTIONS

[374

OF

THREE NEW SPECIES OF PLANTS,

&c. &c.

HAMAMELIS. *Linn.*

SYST. LINN. Tetrandria Digynia.

CHAR. GEN. *Petala* 4, elongata, æstivatione valvato-involuta! *Capsula* semiinfera, bilocularis. *Semina* solitaria.

ORD. NAT. HAMAMELIDÆ, *Br.*

CHAR. GEN. *Petala* elongata, æstivatione valvato-involuta!

OBS. Hamamelidæ notis præsertim sequentibus distinguuntur. *Flos* semisuperus, completus, tetrapetalus. *Stamina* antherifera quatuor, petalis alternantia; *Antheris* basi insertis, bilocularibus, loculo singulo dehiscenti valvula unica medio semiseptifera, et vel altera sutura incompleta persistenti, vel utraque solubili decidua. *Ovarium* biloculare, loculis monospermis, ovulis pendulis (v. appensis). *Styli* duo. *Fructus* semiinferus, capsularis. *Embryo* longitudine fere albuminis: *radicula* supera.

Huic ordini, cui referenda Hamamelis, *Linn.*, *Dicoryphe*, *Aub. Du Petit-Thouars*, et *Dahlia*, *Thunb.*, forsan adjicienda, in distincta tamen sectione, *Fothergilla*, *Linn.*,

pericarpio, semine? necnon habitu similis; petalorum defectu, staminibus numerosis et dehiscentia antherarum diversa.

Hamamelideæ hinc affines *Bruniaceis*, (ordo cui pertinent *Brunia*, *Staavia*, *Linconia*, *Thamnea* et *Erasma*.) ab iisdem distinguuntur antherarum insertionem et dehiscentia, ovarii loculis monospermis, capsulæ dehiscentia, floribus quadridis et habitu; inde accedunt *Corno*, *Marleæ*, *Roxb.* (quæ *Stylidium*, *Lour.*, fide exempl. ab ipso auctore,) generibusque affinibus, et quodammodo *Araliaceis*, diversæ structura antherarum, fructu capsulari aliisque notis.

375]

HAMAMELIS CHINENSIS. *Tub.*¹

Hamamelis foliis integerrimis. *Soland. Mss. in Biblioth. Banks.*

Arbuscula sinensis cisti minoris folio rigidiores, flores candicantes in 5 vel 6 lacinias longas angustas ad umbilicum usque divisos binatim plerumque ferens, fructu ovato parvo bivalvi villos glandis ad instar calyculato. *Cunningham in Pluk. Amalth. 32*, tab. 368, fig. 2*, fide specimen in Herb. Pluk.

Loc. NAT. China prope Nan-king, *D. Abel*; Insula Cheusan, *Cunningham*.

DESCRIPTIO. *Frutex* decumbens, ramosissimus, ramis ramulisque teretibus, ultimis cinerascens, furfuraceis pube stellari brevi. *Folia* alterna, brevè petiolata, integerrima, subovata, acutiuscula vel obtusa, basi inæquali, uninervia, supra avenia, subtus venis anastomozantibus reticulata, utrinque pube stellari brevi, subtus copiosiore, cinerea, 8-10 lineas longa. *Stipulæ*? caducæ. *Capitula* 3-5-flora, pedunculata, ramulos breves terminantia v. axillaria. *Calyx* pube stellari cinereus, furfuraceus; limbo supero, quadridido, laciniis ovatis, planis, trinerviis, extus pubescentibus, intus glabris, æstivatione imbricatis, deciduis. *Petala* 4, æqualia, laciniis calycis alternantia, elongata, calycem aliquoties, ter quaterve, superantia, membranacea, glabra,

¹ [See note at vol. i, p. 187.]

plana, linearia, obtusa, integerrima passimque emarginata, trinervia, nervis ad basin usque distinctis, infernè simplicissimis, supernè divisis, medio ramosiore; æstivatione e basi valvata, supra seorsim spiraliter involuta. *Stamina* antherifera quatuor, epigyna, æqualia, calycis laciniis opposita, brevia, æstivatione conniventia. *Filamenta* brevissima, glabra, crassiuscula. *Antheræ* stantes, ovato-quadratae, glabrae, acumine subulato adscendente, ipsa anthera dimidio brevior; biloculares, loculis lateralibus, medio longitudinaliter sulco, septum parziale indicante, insculptis, valvulâ unicâ medio semiseptiferâ, deciduâ. *Squamulae* (stamina sterilia,) quatuor, glabrae, abbreviatæ, latiores quam longæ, emarginatæ v. semibifidæ, staminibus alternantes. *Ovarium* inferum, brevè turbinatum, disco epigyno nullo, biloculare, loculis monospermis, ovulis pendulis (appensis). *Styli* duo, brevissimi, distantes. *Stigmata* simplicia. *Capsula* semiinfera, corticata, dicocca, coccis semibivalvibus, contrariis segmentis corticis bivalvis.

Obs. Hamamelis Chinensis a Virginica differt valvulis antheræ deciduis pauloque habitu; an itaque (sub nomine Loropetali) in proprium genus separanda?

TABULÆ EXPLICATIO.

1876

Ramus florifer et Ramulus fructifer, uterque magnitudine naturali.

Ad alterum latus tabulæ Flores magnitudine aucti.

1. Flos hinc apertus et vi expansus.
2. Anthera ante dehiscentiam: 3. Eadem post dehiscentiam, valvulis jam delapsis.
4. Pars floris cum ovario longitudinaliter secto; ostendens stylos duos et loculos monospermos.
5. Ovulum, cicatrice umbilicali juxta apicem.
6. Capsula, magnitudine naturali.
7. Pubes stellata, aucta.

ABELIA.¹

SYST. LINN. Didynamia Angiospernia, post Linnæam.

CHAR. GEN. *Pericarpium* inferum, (abortione) monospermum, indehiscens, calyce foliaceo coronatum. *Involucrum* 2-multiflorum, 6-polyphyllum.

ORD. NAT. CAPRIFOLIACEÆ, Sect. I. *Juss. Gen.* 211. (Lonicereæ *Br.*) inter Linnæam et Symphoricarpum.

CHAR. GEN. *Calyx* foliaceus. *Corolla* subinfundibuliformis, 5-loba. *Stamina* 4, didynama. *Ovarium* 3-loculare: loculis duobus polyspermis, abortientibus! tertio monospermo fertili. *Pericarpium* monospermum, indehiscens, calycis limbo foliaceo coronatum. *Involucrum* bi-multiflorum, foliolis sex pluribusve.

Frutices decumbentes vel debiles, glabri. Folia opposita petiolata, dentato-crenata. Pedunculi modo axillares, trichotomi vel trifidi; modo terminales indivisi.

PATRIA China et Japonia.

Obs. Linnææ proximum genus, quacum convenit staminibus didynamis et structura ovarii! diversum fructu hinc uninervi inde quinquenervi, coronato calycis limbo foliaceo persistente, necnon inflorescentia et habitu.

³⁷⁷ *Symphoricarpus* habitu paulò magis accedens *Abelie*, ab eadem facile distinguitur, inflorescentia, floribus pentandris, ovarii quadrilocularis loculis duobus monospermis, fertilibus; duobus reliquis polyspermis sterilibus!

Triosteum fructu calyce foliaceo coronato *Abelie* simile, abunde diversum est ovarii trilocularis loculis omnibus monospermis fertilibus, floribus pentandris et inflorescentia.

¹ This account of the genus ABELIA is extracted from a manuscript dissertation on the natural order to which it belongs.

ABELIA CHINENSIS. *Tab.*¹

Abelia involucris bifloris, pedunculis trichotomis, staminibus exsertis.

Loc. Nat. Chinæ provincia Kiang-si; prope lacum Po-Yang. *D. Abel.*

DESCRIPTIO. *Frutex* decumbens. *Ramuli* virgati, cinerascens, pube tenuissima, per lentem solùm obvia. *Folia* opposita, petiolata, exstipulata, vix uncialia, ovata, acutiuscula, plana, crenato-serrata, uninervia, venis anastomozantibus, immersis, paginis subdiscoloribus, superiore pube rarâ brevissimâ simplici conspersâ, marginibus per lentem ciliatis pilis acutis. *Pedunculi* e summis alis et terminales, approximati, thyrsi speciem efformantes, trichotomi, quandoque tantum trifidi, divisuris bibracteolatis. *Involucrum* biflorum, hexaphyllum, subæquale, persistens, abbreviatum ovarii sessilibus aliquoties brevius. *Calyx* superus, limbo quinquepartito, æquali, laciniis spathulato-oblongis, planis, patentibus, uninerviis, venosis. *Corolla* infundibuliformis, calyce duplo longior: *tubo* quinquenervi, nervis quatuor approximatis, quinto distanti: *fauce* imberbi: *limbo* quinquefido, patenti, laciniis ovatis, obtusis, subæqualibus. *Stamina* quatuor (absque rudimento quinti): *Filamenta* dimidio inferiore tubo corollæ adnata, ejusdem nervis quatuor approximatis primo intuitu subopposita, sed reverâ alterna; superiore libera, filiformia, pilosiuscula, parum inæqualia, apicibus exsertis: *Antheræ* incumbentes, oblongolineares, loculis appositis, absque manifesto connectivo, medio longitudinaliter dehiscentibus. *Ovarium* inferum, lineari-oblongum, modice compressum, tenuissime pubescens, hinc uninerve (fig. 2), inde quinquenerve (fig. 1), in collum breve apice angustatum, basi acuta parum attenuata; trilobulare (fig. 1 et 3) (collo e divisione chordæ pistillaris fenestrato), loculis duobus (interioribus?) polyspermis, ovulis simplici serie insertis, supremo erecto! reliquis pendulis, omnibus constanter abortientibus; loculo tertio in latere

¹ [See note at vol. i, p. 187.]

uninervi (exteriore?) ovarii, monospermo, ovulo ovato, reliquis aliquoties majore, a margine inferiore fenestræ chordæ pistillaris pendulo, vasculoso fasciculo dorsali ! 378] *Stylus* filiformis, glaber, longitudine staminum. *Stigma* depresso-capitatum, indivisum, imberbe. *Pericarpium* figura ovarii, eoque vix duplo majus, calycis limbo, proportionatim aucto, coronatum, monospermum, loculis duobus abortientibus ad alterum latus pressis (fig. 4), induriscens, coriaceum, exsuccum. *Semen* (fig. 4 et 5) subcylindraceum, cavitatem ovarii fere replens, prope apicem insertum funiculo brevi. *Integumentum* duplex : *exterius* membranaceum, laxiusculum ; *interius* tenuissimum, albumini arcuè adherens. *Albumen* (fig. 6) figura et magnitudine seminis, dense carnosum, album. *Embryo* (fig. 6 et 7) axilis, rectus, albus, albumine aliquoties (fere quadruplo) brevior : *Cotyledones* breves : *Radicula* supera.

Obs. The genus *Abelia* is named in honour of its discoverer, CLARKE ABEL, Esq., who accompanied the late Embassy to China, as Chief Medical Officer in the suite of Lord Amherst. Nearly the whole of the extensive collections formed in China by this zealous naturalist were, unfortunately for science, lost in the wreck of the *Alceste* ; the only part saved being a small selection of specimens of plants which he had presented, while in China, to Sir George Staunton, by whom they were most liberally returned to him on his arrival in England.

EURYA. *Thunb. Jap.* p. 11.

SYST. LINN. Polygamia Diœcia.

CHAR. GEN. HERMAPH. *Calyx* 5-part. *Corolla* monopetala, 5-partita. *Stamina* 12—15. *Ovarium* superum, 3-loc. *Stylus* 1. *Stigmata* 2—3.

MAS. Cal. Cor. et Stam. ut in Hermaph. *Pistillum* 0.

FEM. Cal. Cor. Hermaph. *Stamina* 0. *Ovar.* 3-loc. *Stylus* 1. *Stigmata* 3, revoluta. *Bacca* polysperma.

ORD. NAT. TERNSTRÖMIACEÆ. *Mirbel, Nouv. Bullet.* 3, p. 381. *De Candolle, Prop. Med. des Plantes*, p. 203.

CHAR. GEN. Flores Polygami-Dioici. *Calyx* 5-part. *Petala* 5, unguibus connatis. *Stamina* 12—15, simplici serie. *Ovarium* 3-loc. polyspermum. *Stylus* 1. *Stigmata* 3. *Bacca* trilocularis, polysperma. *Semina* reticulata.

OBS. Proximum genus *Fresiera*, distinctum floribus polygamis et petalis basi connatis.

EURYA CHINENSIS. *Tab.*¹

379

Eurya foliis cuneato-ovalibus obovatisque, ramulis ultimis pubescentibus.

Loc. Nat. Chinæ provinciæ Kiang-si et Quang-tong; in campis et collibus.

DESC. *Frutex* bipedalis, ramosissimus, erectus, ramis teretiusculis, patentibus, novellis pube simplici, copiosa, subappressa. *Folia* alterna, petiolata, exstipulata, coriacea, sempervirentia, cuneato-ovalia passimque obovatocuneata, serrata, novella subtus pilis raris conspersa, vix uncialia, siccata subtus præsertim flavicantia. *Flores* parvi, axillares, 3—5, fasciculati, brevè pedunculati, pedunculis apice hibracteolatis. *Calyx* foliolis ovatis, obtusiusculis, duobus exterioribus oppositis minoribus, æstivatione imbricatis. *Corolla* monopetala, subcampanulata, alba, calyce sesquilongior, e petalis quinque, unguibus arcè connatis, laminis obovatis apice patulis, formata. *Stamina* circiter 12—15, hypogyna, simplici serie inserta, basi corollæ leviter cohærentia. *Filamenta*, filiformia, glabra, antheris parum longiora. *Antheræ* juxta basin leviter emarginatam insertæ, lineares, mucronatæ, biloculares, loculis absque manifesto connectivo appositis, medio longitudinaliter dehiscentibus. *Ovarium* sessile, disco nullo cinctum, ovatum, glabrum,

¹ [See note at vol. i, p. 187.]

triloculare, loculis polyspermis. *Stylus* unicus, subulatus, glaber. *Stigmata* tria, filiformia, obtusiuscula, stylo breviora. *Masculi flores* absque pistillo. In *femineis* nulla rudimenta staminum.

Obs. Quam maxime affinis *Euryæ Japonicæ Thunb. Jap. 191* quæ ramulis ultimis glaberrimis folisque ellipticis acutis distincta.

CHARACTERS AND DESCRIPTION

OF

2

LYELLIA,

A NEW GENUS OF MOSSES,

WITH

OBSERVATIONS ON THE SECTION OF THE ORDER TO
WHICH IT BELONGS;

AND SOME REMARKS ON

LEPTOSTOMUM AND BUXBAUMIA.

BY

ROBERT BROWN, Esq., F.R.S., LIBR. L.S.

(READ APRIL 6TH, 1819.)

[*Extracted from 'The Transactions of the Linnean Society of London,'*
Vol. XII, pp. 560—583.]

CHARACTERS AND DESCRIPTION

[500]

OF

LYELLIA, &c.

IN the tenth volume of the Society's Transactions, I have given a description of *Dawsonia*,¹ a genus of Mosses having entirely the habit of *Polytrichum*, but whose peristomium, instead of consisting of a single row of short teeth connected by a horizontal membrane, is composed of an indefinite number of capillary cilia. These cilia, originating both from the inner surface of the capsule, immediately within the orifice, and from the columella itself, form a loose pencil, of which the hairs are sufficiently distinct to allow of a gradual discharge of the seeds.

The correctness of this account of *Dawsonia*, especially as to the origin of the peristomium², and the nature of the supposed columella³, has been questioned by some of those authors, who have since adopted the genus.

From a careful re-examination, however, I find no reason to alter in any respect the generic character formerly given; and to the description of the species I have only to add, that the upper surface of the leaves is furnished with parallel lamellæ, like those of *Polytrichum*; and that the inner membrane of the capsule is, in the early stage at least, ^[501] connected by numerous plicæ with those elevations of the

¹ [Vol. i, p. 348.]

² Bridel Meth. Nov. Muscor. p. 205.

³ De la Pylaie in Journal de Botanique appliq. iii. p. 134.

surface of the columella, which are noticed in my description, and well expressed in Mr. Bauer's excellent figure.

My principal object, in the present paper, is to establish another genus of the same family, equally related to *Polytrichum* in habit, which resembles *Dawsonia* in the remarkable form of its capsule, but whose peristomium is of so peculiar a structure as to require its separation from both these, and readily to distinguish it from all other genera of Mosses.

This new genus I shall name *Lyellia*, in honour of Mr. Charles Lyell, an accurate English Botanist, who has particularly studied, and made several important discoveries in, the natural order to which it belongs.

Mr. Hooker has already intimated his intention of establishing a *Lyellia* in the same order, to consist of three species of *Leucodon* that have a mitriform calyptra;¹ but he has readily agreed to transfer the name of our friend from a genus, respecting which there is still some uncertainty, to one so unquestionable as that here proposed; and as I have not a drawing prepared to accompany the present communication, I trust he will give the necessary illustration of this interesting genus, in an early number of his excellent *Musci Exotici*.

LYELLIA.

Stoma edentulum, clausum *epiphragmate* (crasso depresso), cujus *discus circularis* a limbo (latiori) persistenti secedens, cum columellâ remanenti inclusâ cohæret.

Capsula hinc plana inde convexa.

Calyptra cucullata (apice pilosa).

⁵⁶² *Muscus* (Nepalensis) habitu *Polytrichi*; *capsulæ figurâ et structurâ interiori Dawsoniæ similis*; *peristomio ab utroque genere diversissimus*.

¹ *Musci Exotici*, vol. i. 17.

LYELLIA CRISPA.

DESCRIPTIO. *Muscus cæspitosus.*

Caulis erectus, simplicissimus, 3-4-uncialis, ipsa basitomento radicali cinereo copioso tenuissimo tectus, dein, ad $\frac{1}{2}$ circiter longitudinis, basibus emarcidis foliorum squamatus, supra dense foliatus.

Folia undique versa, e basibus dilatatis subcuneatis semivaginantibus membranaceis pallidis imbricatis, subulata, canaliculata, nigro-viridia, opaca; marginibus ab apice ad $\frac{2}{3}$ longitudinis et ipso apice carinæ serratis; disco intus longitudinaliter lamellis numerosis, parallelo-approximatis, tenuissimis, e nervo ipsaque superficie ortis; limbo elamellato minutissime areolato; madoræ patula leviterque intransversum falcata, siccitate contorta.

Masculi Flores non visi.

Fructificatio Feminea terminalis, solitaria.

Vaginula cylindræa, villis implexis instructa, apice truncato simplici.

Seta sesquiuncialis—biuncialis, erecta, teres, lævis, castanea, demum cava.

Calyptra (nonnullæ delapsæ solum a nobis visæ) cucullata, hinc altè fissa inde truncata, lævis, apicè pilis brevibus simplicibus hispidula.

Capsula erectiuscula, circumscriptione ovata, fusca, vix manifeste areolata, hinc plana et ala perangusta cincta, inde convexiuscula, collo brevissimo cylindræo. *Apophysis* nulla nisi *basis* castaneo-fusca et undique instructa punctis parvis sparsis numerosis, pustuliformibus, verticaliter ellipticis, margine paulo incrassatis, disco angusto, porum re-ferente sed laminam tantum exteriorem perforante.

Operculum (unicum tantum visum quod calyptrà delapsâ inclusum) e basi depresso-conicâ desinens in rostrum longitudine circiter ipsius baseos, intus auctum processu cylindræo, centro baseos inserto, et proculdubio disco circulari terminali columellæ applicito.

Peristomium horizontale, nec obliquum, clausum quasi operculo interiori seu *epiphragmate* crasso, opaco, edentulo,

planiusculo, persistente ; limbo latiusculo crassiore castareo ; disco pallido leviter depresso ; ipso centro circulari plano, crassiusculo, fusco, a disco pallido mox separante, arcte cum columella cohærenti, eaque demum abbreviata intia cavitatem retracto.

Membrana interior (s. Theca vera) approximata exteriori, quacum processibus numerosis vasculiformibus connexa, ore coarctato, spongioso-membranaceo, collum breve columellæ arcte amplexante, superficie interiore altè corrugata.

Columella in capsulamatura majuscula, subovalis, lacunosa, rugis elevatis applicitis et forsan connexis plicis respondentibus membranæ interioris.

Semina minutissima, in cumulo viridia, separatim hyalina, lævia.

Lyellia crispa was lately discovered in Nepaul (probably in the vicinity of Kathmandu,) by the botanical collectors sent from the Company's garden at Calcutta, by Dr. Nathaniel Wallich, the worthy successor of Dr. Roxburgh in that establishment.

The specimens here described were received from Dr. Wallich by Sir Joseph Banks ; and I have also seen others sent at the same time to Lambert, part of which he very liberally communicated.

⁵⁶⁴ The whole number of capsules examined does not exceed twenty-five ; but as all of these were ripe and of uniform appearance and structure, they will probably be considered sufficient for the establishment of the genus.

To complete the description of *Lyellia*, male flowers, which, however, probably resemble those of *Polytrichum* and *Dawsonia*, are still wanting ; and although there is no reason to doubt that the calyptra and operculum, both of which I have ventured to describe from fallen specimens, really belong to this species, yet it would be more satisfactory to find them while still attached to the capsule ; for in this state only the form of the operculum and its probable central connection with the orbicular disk of the peristomium can be absolutely determined. It will be necessary also to examine a greater number of specimens, and perhaps in

different states, to ascertain absolutely the economy of this moss in the dispersion of its seeds. It is evident, however, if the description already given be correct, that as far as dispersion takes place, by the mouth of the capsule, it can only be effected by a contraction or shortening of the columella, and a corresponding retraction within the cavity of the capsule, of the central portion of the peristomium which adheres to it; and in this state several of the specimens were actually found.

But it is also manifest, both from the great size of the columella, and its numerous points of connection with the inner membrane, that in this way the whole of the seeds cannot be discharged. It is possible therefore, that in certain circumstances at least, their dispersion may be assisted by the minute pores existing at the base of the capsule. These pores, in the specimens examined, were found to penetrate the outer membrane, or even its external layer only, and being situated below the origin of the seminal bag, their first effect will probably be to accelerate the decay of the internal spongy texture of the base of the capsule. But by thus removing support from the columella and inner membrane, they may contribute to the greater shortening of the former, and consequently in increasing the dispersion by the mouth of the capsule; or, from the same cause, the inner membrane being at length ruptured, the seeds may be in part discharged by the pores themselves.

It seems then not improbable, from what we at present know of the structure of *Lyellia*, that for the dispersion of the seeds in this genus there are two distinct contrivances, both of which, however, in the only species yet known, are apparently imperfect. But hence it is perhaps allowable to conjecture, that either other species of *Lyellia*, or a nearly-related genus may exist, in which while the mouth of the capsule remains absolutely shut, the pores of the base may be sufficiently enlarged for the complete performance of this important function.

Pores exactly resembling those of *Lyellia crispa* have not hitherto been found in any other moss. I have observed,

however, in several specimens of *Polytrichum alpinum* still more minute pustules, not very different in appearance, and similarly situated on the base of the capsule.

In establishing this new genus of Mosses, it is of importance to determine its more intimate affinities in the family to which it belongs. Its place is unquestionably between *Polytrichum* and *Dawsonia*; and it will I believe be admitted, that these three genera, in the natural method, cannot be separated; though they will necessarily form or be referable to distinct sections of an artificial system founded chiefly on modifications of the peristomium.

In attempting to discover characters by which this group of POLYTRICHOIDEÆ may be distinguished from other Mosses, it is in the first place necessary to determine the whole structure of *Polytrichum*; for this genus, though one of the most common of the order, and, from the great size of the capsules in many of its species, most readily admitting of accurate observation, has never yet been thoroughly examined.

One of the most striking characters of *Polytrichum* is the dense texture and consequent opacity of the leaves; in which it agrees with the other two genera of the section. This character, however, is not altogether confined to *Polytrichoidæ*, and is wanting in *Polytrichum undulatum* and *angustatum*. But the lamellæ of the upper surface of the leaves probably exist, though in very different degrees, in all the species of *Polytrichum*; are equally observable in *Lyellia* and *Dawsonia*; and I am not aware that they have been found in any other genera of the order.

These lamellæ, which are represented in several of the species figured in *English Botany*, by Wahlenberg in *P. lævigatum*,¹ and since noticed by Messrs. Hooker and Taylor² as existing in nearly the whole of the genus, do not belong to the nerve only, as the authors of *Muscologia Britannica* seem to suppose, but in several species cover the greater part of the surface of the upper or spreading portion of the leaf; the sheathing base being either entirely

¹ Flora Lappon. tab. 22.

² Muscol. Brit. p. 24.

destitute of them, or having them much less developed and strictly confined to the nerve.

In the form and position of the male flowers there is probably an absolute uniformity in the whole of this group : but the only peculiarities consist in the perigonal leaf or bractea being formed chiefly of the sheathing part of the stem-leaf ; and in the new shoot proceeding from the centre of the star-like cluster. In *P. undulatum*, however, the former character is by no means obvious, and the latter is not perhaps constant ; and both are only presumed to exist in *Lyellia*.

The double calyptra of *Polytrichum*, long considered as the essential character of the genus, equally exists in *Dawsonia*. But this outer or spurious calyptra, formed of densely matted hairs originating from the vaginula and the apex of the inner cucullate calyptra, is wanting in several species of *Polytrichum*, in some of which, and in *Lyellia*, the true calyptra is furnished with a few hairs only, either scattered over the whole of its surface, or confined to its apex ; while in others, as in *Polytrichum undulatum*, it is nearly, and in *P. magellanicum* and *laevigatum*, entirely smooth.

Respecting the teeth of the peristomium of *Polytrichum*, I have very little to add, except that in *P. magellanicum* they appear to be eighty in number, which is a higher multiple of sixteen than has hitherto been noticed in this genus, and the greatest number that has yet been found in a single series in the order.

On the nature of the transverse membrane or tympanum of *Polytrichum*, I have formerly made some remarks in treating of *Dawsonia*, and have there considered it as the remains of the pulpy continuation of the columella, originally occupying the cavity of the operculum. But its uniform texture, as well as its exactly circular form and equal margin remain to be accounted for ; unless this regularity may be supposed to depend on the circular, and apparently corresponding, aperture of the inner membrane.

Most authors have described the tympanum of *Polytrichum* as minutely perforated. These pores I have never

been able to detect; but I observe in many cases an appearance which may perhaps account for the belief in their existence, namely a minute reticulation on the outer surface of the membrane, apparently owing to the corresponding areolæ of the inner surface of the operculum, with which it was originally in contact.

Polytrichum is remarkable for the various forms of its capsule. Those species, indeed, in which it is quadrangular have been considered as a distinct genus by Mons. de Beauvois. But unless this difference of external form should be found connected with others, either in the habit or in the internal structure, which I believe is not the case, it seems hardly sufficient to justify the subdivision of so natural a genus.

The symmetrical quadrangular capsule, admitting of a regular cylindrical inner membrane, is a character of inferior importance to the plano-convex or dimidiate capsule, which almost necessarily implies, and in *Dawsonia* and *Lyellia* at least is found to be connected with, a corresponding irregularity in the figure of the cavity; and hence I have introduced this remarkable form into the characters of both these genera.

On the structure of the inner membrane of the capsule in *Polytrichum* the only observation that I have met with occurs in *English Botany*, where, in the account of *Polytrichum subrotundum* (plate 1624), it is stated that Mr. James D. Sowerby has discovered, in this species as well as in *Polytrichum undulatum*, a real membranaceous peristomium within the teeth, which, according to Sir James Smith, "gives quite a new idea of the generic character."

This inner peristomium, if it may be so termed, which is well represented in the figure referred to, and consists of a horizontal projection of the inner membrane immediately below its apex, will, I believe, be found to exist in all the species of the genus, and in some cases reduces the aperture of the inner capsule to half the size of the outer at the origin of the teeth. It is always, however, quite entire, and, according to my observations, is formed of a doubling of the

inner membrane, with a loose cellular or rather spongy substance interposed between the lamellæ.

Besides this transverse annular projection there are in the inner membrane of all the species of *Polytrichum* that I have examined, *P. undulatum* alone excepted, four longitudinal equidistant processes, extending from the aperture to the base of the capsule; and in many species projecting so far into its cavity as to come in contact either with the salient angles or sides of the columella, and consequently to subdivide it into a determinate number of cells.

The analogy of these longitudinal processes with the more numerous and irregular plicæ in *Dawsonia* and *Lyellia* is obvious; and I have not myself met with similar processes of the inner membrane in any other genus of mosses. They do not, however, afford an absolutely distinguishing character of this group, as they seem to be altogether wanting in *Polytrichum undulatum*; and Hedwig, it must be admitted, has represented an apparently analogous structure in *Gymnostomum pyriforme*.¹

The quadrangular or four-winged columella of *Polytrichum commune* is well figured and described by the accurate Schmidel²; and I have found an equally regular form of this body in most of the species of *Polytrichum* that I have examined; though in many it is less evidently winged than in *P. commune* and the species nearly related to it.

Mons. de Beauvois seems to consider the alæ of the columella as themselves forming complete septa, and also that in this state they exist only in such species as have quadrangular capsules; for to these he limits his genus *Polytrichum*, distinguished by him from *Pogonatum* by its multilocular fruit. The cavity, however, is completely subdivided in several of the species with cylindrical capsules, as in *P. urnigerum*, and in a new species very nearly related to it (*P. microstomum*, nob.) lately received from Dr. Wallich: and the alæ of the columella, as far as I have observed, never form complete septa, at least in the ripe fruit; though

¹ Fundam. Hist. Nat. Muscor. Frond. pars ii. tab. 2, β.

² Icones pl. p. 236, tab. 59, fig. 15.

in those species having quadrangular capsules they nearly reach the parietes opposite to the external angles, towards which they are directed.

As the columella of *Polytrichum* retains its regular form in the ripe capsule, its real structure may even then be in a great measure determined. In this stage I find its wings, or compressed sides, to consist of a double membrane with an intermediate spongy substance, in which there is no appearance of granules; and the internal denser substance of the axis is equally free from granular matter. But as there is no indication of lateral rupture, the sides in several species remaining perfectly smooth, nor of any central cavity, this structure affords a powerful argument in refutation of those hypotheses which assume the existence of two kinds of granules in the capsules of mosses; the one produced in the cavity formed by the internal membrane, the other in the substance or supposed cavity of the columella itself; the latter being considered as seeds in one of the hypotheses,¹ and in the other as pollen.²

This argument, however, is not here advanced to disprove the existence of two kinds of granules in the capsules of mosses, but merely against their production in the distinct cavities assigned to them in the hypotheses referred to.

In the greater number of *Polytricha* as well as in *Lyellia* and *Dawsonia* the seeds are extremely minute; a fact with which the increased surface for their production is probably connected: for in *P. undulatum*, where the seeds are larger than in most other species of the genus, this increased surface does not exist: and in *P. levigatum*, where they are of still greater size, the plicæ of the inner membrane are probably also wanting.

⁵⁷¹¹ Although there is but little resemblance in the structure of the peristomium among the different genera of *Polytrichoïdeæ*, they may still be said essentially to agree in the function of this part: for in all of them the complete separation of the seeds is ensured by the smallness of the apertures for their discharge. It may be remarked, that the

¹ Palisot Beauvois, *Ætheog.* p. 5.

² Keith, *Physiol. Bot.* ii. p. 346.

necessity for this complete dispersion in Mosses seems to be inversely as the size of the seeds. For in those genera of the order in which the capsule either bursts irregularly or has a naked mouth, the seeds are in general larger than in those with a single, and still more manifestly than in those with a double, peristomium. And in conformity with this also, in *Polytrichum undulatum* and *laevigatum* the tympanum is sooner ruptured or removed than in the other species of the genus.

The result of this comparison of *Polytrichum* with *Lyellia* and *Dawsonia*, although it confirms the propriety of their approximation, does not afford any clearly distinguishing mark for the very natural section of the order which these three genera form. In the mean time, however, it may be circumscribed, though not with absolute precision, by a combination of the more general characters which have been now enumerated.

LEPTOSTOMUM.

In defining this genus, which was first proposed in my former paper on Mosses¹, I relied chiefly on the undivided annular projection of the inner membrane of the capsule. I was induced to employ this modification of the peristomium as a character, though certainly far from being obvious, in finding it to exist in several mosses of the southern hemisphere, having a similar and peculiar habit: and which, had it been neglected, I must have referred to *Gymnostomum*, with the greater number of whose species they have hardly anything in common.

Mr. Hooker, however, has since found, as he states, (572 the same structure in several other mosses, generally considered as having a naked peristomium, particularly in *Gymnostomum microstomum*, *Griffithianum*, and *fasciculare*; and as these species have but little resemblance to each other, and still less affinity to *Leptostomum*, he has reduced this genus also to *Gymnostomum*.

If the statement referred to be correct, *Leptostomum*,

¹ [Vol. i, p. 352.]

though it may be a natural genus, must be given up, until other marks shall be found by which it may be distinguished. I have not had specimens sufficiently perfect to enable me to judge of the structure of all the species of *Gymnostomum* mentioned by Mr. Hooker. In one of them, however, *Gymnostomum microstomum*, the peristomium is certainly very different from that of *Leptostomum*. In this species I find, on removing the operculum, that the mouth of the capsule is not only completely covered by a horizontal membrane, but that this covering is derived from the outer membrane of the capsule, and consequently differs in origin as well as in form from the peristomium which it has been said to resemble. Its central portion, however, being extremely thin is soon ruptured and deciduous, and in this state only it has been seen by the authors of the *Muscologia Britannica*.

Gymnostomum microstomum therefore may itself be considered as a distinct genus, to which the name of *Hymenostomum*¹ may be given; and it is worthy of remark, that in its technical character it approaches most nearly to *Lyellia*, though no two mosses differ more widely in almost every other respect.

^{573]} From the account given of *Gymnostomum Griffithianum* it appears that this species also has in the early stage a membrane completely covering the mouth of the capsule. But this membrane probably differs in origin, at least, from that of *Hymenostomum*, as it seems to do both in form and dehiscence from the peristomium of *Leptostomum*.

Of *Gymnostomum fasciculare* I have examined only imperfect specimens, I cannot therefore speak with confidence of its structure. The annular process, however, mentioned by Mr. Hooker is more likely to be the remains of a complete horizontal covering, and probably originating from the inner membrane, than to resemble the peristomium either of

¹ HYMENOSTOMUM.

Fl. Fem. terminalis.

Stoma edentulum, clausum *epiphragmate* (e membrana exteriori orto), disco tenuissimo (a columella libero) mox rupto et evanido; *limbo* persistenti horizontali indiviso.

Calyptra dimidiata, lævis.

Fl. Mas. terminalis, gemmiformis.

Hymenostomum or *Leptostomum*. A membrane of this kind is certainly present in some species of *Gymnostomum*, and perhaps may be found in all those that really belong to that genus. It exists also in *Weissia Templetoni*, which so closely resembles *Gymnostomum fasciculare* as to be with difficulty distinguished from it, unless by the inspection of the peristomium; and, in addition to the erect annular peristomium from which the character of the genus is taken, I have observed a similar membrane in *Leptostomum* itself. It seems even to be not an uncommon process or termination of the inner membrane, though it has been remarked only in some of its more obvious and persistent modifications. Thus the spongy membrane figured and described in the two published species of *Calymperes*, seems to be an analogous structure,¹ as is also the circular disk terminating the columella in several species of *Splachnum*; ¹⁸⁷⁴ and perhaps even the tympanum of *Polytrichum* may be of similar origin.

But these characters of *Leptostomum* and *Hymenostomum*, though they do not appear to have been yet observed in any other mosses, may still perhaps be considered too minute for generic distinctions: and it must be admitted that were nothing to be obtained but the subdivision of an extensive natural genus it could not be necessary to have recourse to them. The divisions in question, however, are certainly not of that kind.

The weakest part indeed of Hedwig's system is its bringing together all those mosses that have a naked peristomium, and even including the greater part of them in the genus *Gymnostomum*; while many of the species so associated are in real affinity much nearer to several other genera of the order having a simple or even a double peristomium.

¹ The circular spongy membrane covering the mouth of the capsule certainly does not form an essential part of the character of *Calymperes*; for, in the only species that I have examined, it is either entirely wanting, or firmly adheres to the inner surface of the operculum, along with which, also, a considerable portion of the columella separates. Nor has Swartz, who established the genus (in *Spreng. Schrad. und Link Jahrb. der Gewäch.* vol. i, p. 1) even noticed this membrane in his description.

Thus *Gymnostomum microstomum*, the *Hymenostomum* of the present paper, has less the habit of the genus in which it is placed than of *Weissia*, to some of whose species, especially *W. affinis* and *trichodes*, it seems to approach even in the structure of its peristomium.

Several species of that section of *Gymnostomum*, to which perhaps the genus should be limited, especially *G. fasciculare*, *Bonplandii*, and *Rottleri*, can hardly be distinguished from *Weissia Templetoni*.¹

^{575]} *Gymnostomum* or *Anictangium pulvinatum* agrees in every other part of its structure with *Grimmia*.

Gymnostomum lapponicum, notwithstanding the difference of calyptra, may be considered as related to *Grimmia Daviesii*, and consequently to *Orthotrichum*, which *G. Daviesii*² resembles in its teeth being approximated in pairs.

Gymnostomum viridissimum has exactly the habit and calyptra of *Zygodon*.

Gymnostomum pennatum (*Schistostega* of Mohr.), in one remarkable character may be compared with *Fissidens*.³

¹ *Weissia Templetoni*, along with a nearly related species found in New Holland, *Funaria minor* of Delile (*Flor. Egypt.*), and perhaps also *Weissia radians*, may form a genus distinct from *Weissia*, and nearly related to *Funaria*, differing chiefly in the irregular bursting and evanescence of the inner peristomium, which in *Funaria* is regularly divided and generally persistent, though in some cases perhaps equally deciduous. In a variety of *Weissia Templetoni*, or a very nearly related species, collected in 1800 in the county of Donegal, I have observed the outer peristomium to be not unfrequently wanting, even before the separation of the operculum; a fact which, if hereafter confirmed, would establish its affinity to *Gymnostomum fasciculare*.

² *Griffithia Daviesii* nob.

³ As Schkuhr (in *Krypt. Gewäch.* ii, p. 31, t. 12) has ascertained that the operculum of *Gymnostomum pennatum* separates entire, the genus *Schistostega* must be again reduced to *Gymnostomum*, until other distinguishing characters are discovered.

Its resemblance to *Fissidens* consists in the somewhat similar disposition of leaves.

In *Fissidens*, as limited by Bridel (*Muscol. Nov.* p. 186) the leaves are universally described as presenting their margin instead of their disk to the stem, and as having a doubling of the lower half of their inner or upper margin, extending as far as the nerve.

On this view Bridel (in *l. cit.*) has formed a separate section of the order, consisting of *Fissidens* and *Octodiceras*; and hence also M. de la Pylae has changed the name of *Fissidens* to *Skitophyllum*. (*Journal de Bot. Appliq.* iv, p. 133.) It seems to me a much simpler explanation of the apparent anomaly to consider the supposed doubling or division of the leaf as its true disk, and

Anictangium aquaticum is evidently related to *Cinclidotus* or *Trichostomum*.

Gymnostomum julaceum and *Hedwigia secunda* of Hooker resemble certain species of *Pterygandrum*, *Neckera* and *Leskea*.

An unpublished moss (*Glyphocarpa capensis*) with a naked peristomium, which I observed on the Table Mountain of the Cape of Good Hope has the spherical ^[576] striated capsule as well as the inflorescence and ramification of *Bartramia*: and with this genus also *Anictangium Humboldtii* agrees in its capsule, though its habit is that of *Leskea* or *Hypnum*.

Drepanophyllum of Richard (*Dicranum? falcifolium* of Hooker,) in form and disposition of leaves is related to *Fissidens* and *Neckera*.

Calymperes approaches to *Orthotrichum*, or rather, perhaps, to *Schlotheimia* or *Macromitrium*.

Lyellia, which belongs to the same division of the artificial system, is evidently allied to *Dawsonia* and *Polytrichum*.

And lastly, *Leptostomum*, the genus more particularly under consideration, appears to me most nearly related to *Bryum*; with which indeed its affinity would be completely established, were Hedwig's account of the peristomium of *Bryum macrocarpum* proved to be correct.

To the observations now made on the various affinities of mosses which agree in having a naked peristomium, it may be added, that the genera with a simple peristomium do not form a strictly natural series, several of them being much more nearly related to those in which the peristomium is double than to each other.

But if the correctness of these statements be admitted, it follows that, in many cases to obtain natural genera in this order either additional sources of distinction must be

the deviation from the usual structure as consequently consisting in the greater compression of the leaf, and in the addition of a dorsal and terminal wing. In support of this view it may be observed, that in the lower leaves of the stem both the additional wings are greatly reduced in size, and in some cases entirely wanting, as they universally are in the perigonal leaves, which have likewise the more ordinary form, being moderately concave and not even navicular.

sought for, or those at present in use more minutely investigated.

Of additional characters, which in some cases may be employed with advantage, I shall merely advert to the membranes of the capsule being distinct or contiguous, and to that more intimate union where there seems to be a single membrane only; to the modifications of internal structure of the inner membrane; the differences in form and duration of the columella, or even its being entirely wanting in the ripe capsule; the presence or absence of ⁵⁷⁷ the annulus; and the insertion, form and relative position of the male flowers, which, though always considered of importance by Hedwig, many of the most distinguished muscologists of the present day entirely exclude from the characters of their genera.

With respect to the principal source of generic distinctions, the Peristomium, in addition to the circumstances generally attended to, namely, the origin, number, direction, form, and actual division of the teeth, it may be of some importance to ascertain their æstivation, which, though very generally, is not always valvular: and especially to mark the existence or want of the *longitudinal striæ* or semi-pellucid lines; for these, if they do not prove the compound nature, at least clearly indicate a tendency to division in the teeth where they are found; division being always in the course of the striæ, and in no instance taking place unless where they are present.

But in considering them, which I am inclined to do, as proving composition or confluence of the teeth, it would appear that there is a much greater uniformity in the structure of the simple or outer peristomium, at least, than is generally admitted; and that the prevailing number of teeth in this series is thirty-two; though by a coalescence, more or less complete, they are frequently reduced to sixteen, in some cases to eight, and in a few even to four.

According to this view, a single longitudinal line in the axis of a tooth indicates the confluence of two teeth; three equidistant lines, one being central, the coalescence of four; and seven lines similarly disposed that of eight.

Nearly the whole of these modifications exist in that natural subdivision of the order, which may be named *Splachneæ*, consisting of *Splachnum*, *Systylium*, *Tayloria* (*Hookeria* of Schwaegrichen), *Splachnum squarrosum* of Hooker, and *Weissia splachnoides*.

[578] The number of teeth in *SPLACHNEÆ* is thirty-two, which, however, are never entirely distinct and at the same time equidistant, but approximated or united in various degrees in the different genera and species of the section.

Thus in *Tayloria* and *Systylium* the thirty-two teeth are distinct and disposed in sixteen pairs.

In *Splachnum rubrum* and *luteum* there are apparently only eight pairs, each tooth, however, having a pellucid and obscurely-perforated axis. In almost all the other genuine species of *Splachnum* there is the same disposition as in *S. rubrum* and *luteum*; but the pellucid axis of each tooth is less distinct and imperforated.

In *Splachnum angustatum*, and I believe also in a second species nearly related to it, the arrangement is somewhat different; for the sixteen apparent teeth are approximated, and at the base even united in fours, the pellucid axis of each tooth being still less obvious. Hence these species in their peristomium very nearly approach to *Tetraphis*, to which they would be absolutely referable were the union complete.

In *Splachnum squarrosum* the apparent number of teeth is eight, without any actual subdivision. But as each tooth has three equidistant pellucid lines, of which the lateral are nearly as distinct as the central, there can be no doubt that the composition is the same here as in the rest of the section.¹

¹ In a late number of *Musci Exotici*, (No. 17, tab. 136,) *Splachnum squarrosum* is transferred to *Octoblepharum*, and, on the authority of M. de Beauvois, is stated to be *Octoblepharum serratum* of Bridel. Mr. Hooker, however, continues to refer it to this genus, on the supposition of its agreeing with the original species in the form of its calyptra: observing that if this should prove not to be the case, it ought to be separated, under the generic name *Orthodon*, formerly given to it by its discoverer M. Bory de St. Vincent.

The calyptra of *O. albidum* is represented as distinctly cucullate, both by Swartz (in *Obs. Bot.* tab. xi, fig. 1) and M. de Beauvois (in *Flore d'Ouave*, i, tab. 31). I have also observed it of the same form in specimens from Madagascar. There seems, therefore, no reason to doubt that these two

579] By these lines also *S. squarrosum* is readily distinguished from *Octoblepharum*, in which the apparent number of teeth is the same: for in *Octoblepharum* each tooth has only a single pellucid line; and hence its affinity to certain species at present referable to *Weissia*, with a nearly similar habit and sixteen distinct teeth, whose axis is not perceptibly pellucid.

Weissia splachnoides differs from the other *Splachnæ* in having sixteen equidistant teeth; but as these teeth, according to the indication of the pellucid axis, are double, the arrangement may be compared with that of *Tayloria* and *Systylium*, in which the separation into thirty-two is complete, and the sixteen pairs equidistant. It agrees, however, also in this respect with *Grimmia* and with several species of *Weissia*: but in other important characters, as well as in habit, it is evidently related to *Splachnum*, and offers perhaps one of the best examples of the importance of the male flowers in distinguishing natural genera.

Even *Tetraphis pellucida* may be cited in proof of the same prevailing number in the peristomium; each of its four teeth, when highly magnified, appearing to have seven longitudinal striæ, which, according to this test, would make the real number thirty-two; a structure contributing to fix the place of *Tetraphis* in the natural series between *Splachnum* and *Orthotrichum*.

580] Better evidence on the same subject is afforded by *Trichostomum*, *Didymodon*, and *Leucodon*, in all of which the thirty-two teeth are distinct, though approximated in pairs; by the sixteen bifid teeth of *Dicranum* and *Fissidens*; and by the like number of teeth with a perforated axis in *Trematodon*, *Weissia nuda*, *Didymodon latifolium*, and several species of *Grimmia*.

mosses differ materially even in this part of their structure; and as other differences, of at least equal importance, also exist, both in the peristomium and male flowers, *Octoblepharum serratum*, whose habit is nearly that of *Splachnum*, may be distinguished both from that genus and from *Octoblepharum* by the following characters.

ORTHODON.

Pl. Fem. terminalis.

Peristomium simplex, octodentatum, dente singulo striis tribus longitudinalibus instructo (ideoque e quatuor coalitis composito).

Calyptra mitriformis (4-fida, pilosa).

Pl. Mas. terminalis, discoideus.

In all the genera having a double peristomium I believe the pellucid axis more or less manifestly exists ; but in these genera there is a great uniformity in the apparent number of teeth in the outer peristomium ; there being no instance of actual division in this series beyond sixteen, or of a further approximation, unless in *Orthotrichum*, in several of whose species the approximation or even union of the double teeth by pairs takes place, while in a few others the sixteen teeth are slightly divided at the apex, and in the whole genus the pellucid axis is remarkably distinct.

The only exceptions to the actual division into thirty-two, or the structure indicating that number, in the simple peristomium of Mosses, occur on the one hand in certain species of *Weissia*, perhaps in *Encalypta* and in *Octoblepharum*, in all of which, I believe, there is a reduction to sixteen : and on the other in *Polytrichum*, where the number is frequently increased, varying in the different species, and chiefly by multiples of sixteen, from thirty-two to eighty. In this genus also, whatever the number may be, the teeth never have a semipellucid, but rather an opaque or thickened axis, and no tendency to union or even approximation is observable. The constant equidistance of the teeth of *Polytrichum* seems to be connected with its peculiar mode of dissemination ; for as this takes place through the interstices of the teeth, and as complete separation of the seeds seems necessary on account of their extreme minuteness, a reduction in number and consequent increase of size of these apertures would probably, in some degree, prevent dispersion, while the unequal distances of the teeth might either produce a deviation from the regular figure, or an early rupture, of the tympanum, which forms an essential part in this economy.

BUXBAUMIA.

In my former paper¹ I have proposed to preserve the genus *Buxbaumia*, as established by Schmidel ; and in construct-

¹ [Vol. i, p. 351.]

ing a character to comprehend both species, I entirely rejected the outer peristomium of Hedwig; and having also adopted his opinion respecting the middle peristomium of *B. aphylla*, which he has termed corona, and considered as analogous to the annulus in many other mosses, it became unnecessary to advert to this part in defining the genus.

Mr. Hooker has since published an excellent analysis of both species, and has followed Ehrhart and Mohr in regarding them as forming distinct genera.

This determination I have now no hesitation in adopting; for, whatever the nature of Hedwig's corona may be, it affords at least an obvious character, and is connected with other differences of sufficient importance to justify the separation; though the two genera must always remain in the same natural section of the order.

The observations that follow belong, therefore, solely to *Buxbaumia aphylla*.

My first remark on this plant relates to its peristomium, on the nature of which, as compared with that of other mosses, at least two different opinions may be formed.

According to one of these, the outer peristomium of Hooker may with Hedwig be considered analogous to the fimbria or annulus existing in many other mosses; and the principal objection to this view would perhaps be obviated by rejecting the outer peristomium of Hedwig, as I have formerly proposed, and which Mr. Hooker has since done; as there would then be nothing either in the origin or texture of this part essentially at variance with the supposition; the principal remaining difference being its greater length; for the cilia of the peristomium of *Buxbaumia* may be compared with the striæ or divisions existing in the annulus, which seem to be equally determinate in number, and in some cases also disposed in a double series.

On the second supposition, the peristomium of *Buxbaumia* originating entirely from the outer membrane, may, though consisting of several and even of dissimilar series, be regarded as analogous to that portion of the pencil of *Dawsonia* which arises from the same part of the capsule. This analogy is suggested by Mr. Hooker, and is confirmed

by a circumstance that he does not seem to have noticed, namely, that his outer peristomium, the corona of Hedwig, consists of a double series of cilia. The number of cilia in each series exceeds sixteen, but hardly amounts to thirty-two; it probably, however, corresponds with that of the plicæ in the membranaceous peristomium.

We have here then a passage from a number still perhaps definite, though disposed in a triple series, to the indefinite number peculiar to, and so striking in, *Dawsonia*.

My second observation relates to the inner membrane of the capsule, of which I find the mouth to be quite entire and open, though before the separation of the operculum it is closed by the terminating process of the columella. Hence *Buxbaumia* has some resemblance to *Polytrichum*, and a still greater to *Lyellia*, in this part of its structure.

Buxbaumia aphylla is the only moss considered as being entirely destitute of leaves; and though it has been oftener and more fully described than any other plant of the ⁶⁸³ order, from the monographs of Linnæus and Schmidel to the excellent illustration recently published by Mr. Hooker, there is no difference of opinion on this point. I have lately ascertained, however, that *Buxbaumia aphylla* is always furnished with perfect leaves, which more nearly resemble, both in texture and division, those of a *Jungermannia* than of any species of moss properly so called; and consequently are widely different from those of *Polytrichoidæ*, to which this genus is in several respects related.

The leaves in the barren plant, where I first observed them, are lanceolate and but slightly divided. Those at the base of the female perichætium are even broader than the former, but more deeply cut, both laterally and at top, into several capillary segments; while the leaves which proceed from the surface of the perichætium are still more deeply divided, and their segments so much elongated that the minute foliaceous base has been universally overlooked, and the perichætium consequently described as covered with hairs.

REMARKS
ON THE
STRUCTURE AND AFFINITIES
OF
CEPHALOTUS.

BY
ROBERT BROWN, Esq., F.R.S., &c.

*[Extracted from the 'London and Edinburgh Philosophical Magazine,'
Vol. I, pp. 314—317.]*

LONDON.

1832.

OF

CEPHALOTUS.

IN the Botanical appendix of Captain Flinders's Voyage to Terra Australis,¹ a figure and description of *Cephalotus follicularis* are given, in some respects more complete than those of M. Labillardière, by whom this remarkable plant, a native of the south-west coast of New Holland, was first published. Both accounts, however, are equally imperfect with regard to the fruit; and my principal object in the present communication is to supply that deficiency.

My earliest knowledge of the ripe fruit of *Cephalotus* was obtained from a single specimen, sent in 1815 by M. Leschenault, who had found the plant in February 1803 near the shores of King George's Sound, where I had gathered it in a less advanced state in the beginning of January 1802.

I have, however, more recently, received numerous specimens with ripe seeds from Mr. William Baxter, who collected them also at King George's Sound in 1829.

Cephalotus was introduced in 1823 from the same place by Capt. King, into His Majesty's Botanic Garden at Kew, where it flowered repeatedly, and ripened seeds from which several plants have been raised. A figure of one of these with expanded flowers, but still without fruit, has lately been published by Dr. Hooker in the Botanical Magazine; and a plant brought also from King George's Sound in 1829 by Mr. William Baxter is now in flower in Mr. Knight's nursery.

The following account of the ripe fruit will serve as a

¹ [Vol. i, p. 76, t. 4.]

supplement to the description of the plant which I have given in the work referred to.

AKENIA membranacea, insecta parva alis conniventibus quodammodò referentia, perianthio parùm aucto staminibusque persistentibus cincta, iisque sesquolongiora, ferè distincta, ipsâ basi, ubi receptaculo communi inserta, post separationem intùs aperta ibique e membranâ simplici crassiusculâ imberbi nitente formata; suprà clausa et e duplici membranâ conflata; harum exterior densè barbata, pilis longis, strictis, acutis, deflexis, stylo persistenti brevi arcuè reflexo rostrata: membrana seu lamella interior tenuis, intùs quandoque dehiscens.

SEMEN unicum (rarissimè duo), basi cavitatis membranæ interioris insertum, oblongo-ovale, teres, funiculo umbilicali brevi juxta basin affixum. *Integumentum* duplex: *Testa* membranacea laxiuscula, *raphe* tenui laterali et apice *chalazâ* parvâ insignita: *Membrana interior* tenuis separabilis. *Albumen* semini conforme, album, carnosum, subfriabile, e materiâ oleosâ cum granulis minutis mixtâ constans.

EMBRYO parvus, in basi axeos albuminis, teretiusculus, albus, rectus, albumine 4—5ies brevior. *Cotyledones* breves, plano-convexæ. *Radicula* teres, basin seminis attingens.

RECEPTACULUM commune fructûs: tuberculum centrale, parvum, brevissimum, subcylindraceum, cujus lateribus bases apertæ akeniorum adnatæ sunt, apice convexiusculo barbato.

From this description, especially of the embryo, it is evident that *Cephalotus* must be removed from *Rosaceæ*, to which it had been referred by M. Labillardière; and also, though not with much confidence, in the account which I published in 1814. M. de Jussieu, indeed, in 1818, ³¹⁶ proposed to exclude it from *Rosaceæ* and append it to *Crassulaceæ*; and the structure of the seed, as well as of the folliculi or akenia, and even their insertion on the minute central receptacle or axis, may seem to confirm the correctness of this approximation.

Cephalotus, however, still appears to me sufficiently remote from every natural order at present established, to entitle it

(like *Philydrum*¹ and *Brunonia*²), now that its structure is completely known, to rank as a distinct family which may be called *CEPHALOTÆÆ*, and which may be placed between *Crassulaceæ* and *Francoaceæ*; differing from both in being apetalous, in the valvate æstivation of the perianthium, and in many characters of inferior importance: from *Crassulaceæ* also in its minute embryo and more copious albumen: and from *Francoaceæ* in the absence of barren stamina and in the pistilla being monospermous and apparently distinct.

The most striking peculiarity of *Cephalotus* consists in the conversion of a portion of its radical leaves into *Ascidia* or pitchers. But as *ascidia* in all cases are manifestly formed from or belong to leaves, and as the various parts of the flower in *Phænogamous* plants are now generally regarded as modifications of the same organs, the question is naturally suggested, how far the form and arrangement of the parts of fructification agree in those plants whose leaves are capable of producing *ascidia* or pitchers. The four principal, and indeed the only genera in which pitchers occur, are *Nepenthes*, *Cephalotus*, *Sarracenia*, and *Dischidia*, and the few other somewhat analogous cases, consisting of the conversion of bractææ or floral leaves into open cuculli, are found in *Marcgravia* and two other genera of the same natural family.

The only thing common to all these plants is, that they are *Dicotyledonous*.

It may also be remarked, that in those genera in which the *Ascidia* have an operculum, namely *Nepenthes*, *Cephalotus*, and *Sarracenia*, they exist in every known species of each genus, and the structure of these genera is so peculiar that they form three distinct natural families; while in *Dischidia*, whose pitchers are formed without opercula, these organs are neither found in every species of the genus, nor in any other genus of the extensive natural order to which it belongs.

The striking resemblance in most points of the *Ascidia* of

¹ Flinders's Voyage, vol. ii, p. 578 [*vol. i, p. 53*].

² Transact. Linn. Soc. vol. xii, p. 132 [*ante, p. 310*].

Cephalotus to those of *Nepenthes*, leads to a comparison in the first place of these two genera. But although both are apetalous, and in the parts of the flower deviate from the quinary or prevailing number in Dicotyledones, yet they ³¹⁷ differ in so many other important characters that they cannot be considered as nearly related.

The place of *Nepenthes* in the natural series I have long since¹, in my account of *Rafflesia*, suggested to be near *Aristolochiæ* or *Asarinæ*, without, however, intending to include it in that family.

This approximation was adopted by M. Ad. Brongniart, who, however, went further, having absolutely referred *Nepenthes* to *Cytinæ*.

The union of plants so utterly unlike in appearance and æconomy, and so different, it may be added, in many of their most important characters, seems to have been generally regarded as somewhat paradoxical; and accordingly Professor Link, in 1829, has established *Nepenthes* as a section or tribe of *Aristolochiæ*, and Dr. Bartling and Mr. Lindley, in 1830, have considered it as forming a distinct natural family.

To the numerous and obvious distinctions between *Cytinæ* and *Nepenthes* may be added the no less important differences in their internal structure. For while *Cytinæ*, like most, perhaps all, other plants parasitical on roots, are destitute of spiral vessels, *Nepenthes* exhibits these vessels in the greatest degree of development and abundance, and also produces them in parts in which they are hardly to be met with in any other dicotyledonous plant.

Thus, in addition to the dense circle or stratum of spiral vessels existing in the stem between the outer parenchyma and the wood, they are found also singly or scattered in the pith, in the loose parenchyma situated between the wood and the bark, if it may be so called, even in the fibres of the root, and everywhere in the substance of the leaves, the pitchers, calyx and capsules. And between these solitary or scattered spiral vessels, which are often of considerable

¹ Transact. Linn. Soc. vol. xiii, p. 219 [vol. i, p. 386].

length, and those forming the stratum or circle externally bounding the wood and existing in the veins of the leaves, no essential difference in structure will I believe be found. In these points there is little resemblance between *Nepenthes* and *Cephalotus*, in the internal structure of which last there is nothing unusual.

Between the parts of fructification of *Nepenthes* or *Cephalotus* and *Sarracenia*, there is still less analogy, and it is obviously unnecessary to compare in this respect any of these genera with *Dischidia*.

September 25th, 1832.

CHARACTERS AND DESCRIPTION
OF
LIMNANTHES,
A NEW GENUS OF PLANTS
ALLIED TO
FLOERKEA.

BY
ROBERT BROWN, Esq., V.P.L.S.

(READ AT THE LINNEAN SOCIETY, JUNE 18TH, 1833.)

LONDON.

1833.

CHARACTERS AND DESCRIPTION

OF

LIMNANTHES.

A PAPER WAS read, entitled "Characters and Description of *Limnanthes*, a new genus of plants allied to *Floerkea*," by Robert Brown, Esq., V.P.L.S.

For specimens of the plant described the writer is indebted to the Horticultural Society, and to Mr. David Douglas, F.L.S., by whom it was recently discovered in California.

Mr. Brown was led more particularly to examine *Limnanthes*, from its resemblance to *Floerkea* of Willdenow, a genus which he had many years since investigated without being able to determine its place in the natural system. Examination proved these two plants to be so nearly akin, that they might perhaps be included in the same genus. They are here, however, separated, and the two genera are considered as forming a family distinct from all those at present known.

The place of this new family (LIMNANTHÆ) is not absolutely determined; but it is suggested that in two remarkable points of its structure, namely, the presence of glands subtending the alternate filaments, and the existence of a gynobase, it more nearly approaches to Hypogynous families than to Perigynous, with which it has hitherto been associated.

The following are the characters of the Natural Order, and of the two Genera forming it.

LIMNANTHÆ.

Flos completus, regularis. *Calyx* 3—5-partitus, æstivatione valvatâ, persistens. *Petala* 3—5, marcescentia. *Stamina* 6—10, insertione ambiguâ (hypo-perigyna), marcescentia. *Filamenta* distincta, 3—5 sepalis opposita basi extus glandulâ munita. *Ovaria* 2—5, sepalis opposita, cum *stylo* communi 2—5-fido mediante gynobasi connexa, monosperma; ovulo erecto, nucleo inverso. *Achenia* subcarnosa. *Semen* exalbuminosum. *Embryo* rectus; *radicula* infera.

Herbæ (Americæ septentrionalis, paludosæ) *glaberrimæ*, *alternifoliæ*, *æstipulatæ*, *foliis divisis*, *pedunculis unifloris*, *ebracteatis*, *apice dilatato basin turbinatam calycis simulantæ*.

Limnanthes.

Calyx 5-partitus. *Petala* 5, calyce longiora, æstivatione contortâ. *Stamina* 10. *Ovaria* 5.

Herba (*Limnanthes Douglasii*, Americæ occidentali-borealis) *foliis bipinnatifidis*, *pinnis suboppositis segmentis alternis*.

Floerkea. Willd.

Calyx 3-partitus. *Petala* 3, calyce breviora. *Stamina* 6. *Ovaria* 2 (raro 3).

Herba (Americæ orientali-borealis) *foliis pinnatifidis*, *segmentis indivisis*.

va-
na
cia.
an-
m-
aa ;
en

ae,
is,
le.

one

li-
tis

6.

lis,

PART IV.

CONTRIBUTIONS TO SYSTEMATIC WORKS.

6"

Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

15 128
16 32 25
17 22
18 20

10

GENERA ET SPECIES
PLANTARUM CRUCIFERARUM,
NECNON GENERIS CLEOME,
QUE
IN HORTO KEWENSI COLUNTUR.

[*Extracted from 'Hortus Kewensis,' 2nd Edition, Vol. IV,*
pp. 71—130.]

LONDON.

1812.

TETRADYNAMIA.

[71]

SILICULOSA.

CAKILE. *Gærtn. sem. 2, p. 291.*

Silicula biarviculata, articulis monospermis: superioris semine erecto sessili; inferioris (quandoque abortientis) pendulo.

1. *C. maritima*, articulis ancipitibus: superiore sagittato, foliis pinnatifidis subdentatis carnosiss.

Cakile maritima. *Willden. sp. pl. 3, p. 416.*

Bunias Cakile. *Eng. bot. 231.*

Nat. of Britain. ☉.

2. *C. perennis*, articulo superiore ovali ancipiti stylo longiore, foliis laciniato-pinnatifidis dentatis acutis.

Myagrum perenne. *Willden. sp. pl. 3, p. 406. Jacqu. austr. 5, p. 7, t. 414.*

Nat. of Germany. ♀.

3. *C. rugosa*, articulo superiore subgloboso rugoso stylo brevior, foliis obtusis dentatis; radicalibus sublyratis.

Myagrum rugosum. *Willden. sp. pl. 3, p. 406.*

Rapistrum rugosum. *Allion. pedem. 1, p. 257, t. 78.*

Nat. of the South of Europe. ☉.

CRAMBE. *Gen. pl. 1071.*

[72]

Silicula articulo superiore subgloboso, semine inverso

funiculo basi loculi inserto; articulo inferiore abortiente pedicelliformi.

1. *C. maritima*, filamentis longioribus bifurcis, silicula mutica, foliis subrotundis sinuatis undulatis dentatis glaucis cauleque glabris.

Crambe maritima. Willden. *sp. pl.* 3, p. 418. *Engl. bot.* 924.

Nat. of Britain. ♀.

2. *C. pinnatifida*, filamentis longioribus bifurcis, silicula mutica, foliis pinnatifidis: lobis oblongis dentatis cauleque glabris.

Crambe orientalis. Jacqu. *ic.* 1, t. 128.

Crambe suecica. Mill. *dict.*

Nat. of Siberia. ♀.

3. *C. orientalis*, filamentis longioribus bifurcis, silicula mutica, foliis sinuato-pinnatifidis dentatis scabris, caule glabro.

Crambe orientalis. Willden. *sp. pl.* 3, p. 418.

Nat. of the Levant. ♀.

4. *C. tatarica*, filamentis longioribus bifurcis, silicula mutica, foliis radicalibus decompositis: pinnulis dentato-incisis; adultis cauleque glabriusculis.

⁷³¹ *Crambe tatarica.* Willden. *sp. pl.* 3, p. 418. Jacqu. *ic.* 1, t. 129.

Nat. of Hungary and Siberia. ♀.

5. *C. hispanica*, filamentis longioribus bifurcis, silicula mutica, foliis pinnato-lyratis scabris: foliolo terminali reniformi obtuso.

Crambe hispanica. Willden. *sp. pl.* 3, p. 419. Schkuhr *handb.* 2, p. 292, t. 189.

Nat. of Spain. ☉.

6. *C. fruticosa*, filamentis longioribus bifurcis, silicula stylo mucronata, foliis lyrato-pinnatifidis dentatis canis, caule fruticoso.

Crambe fruticosa. Willden. *sp. pl.* 3, p. 420.

Nat. of Madeira. Mr. Fr. Masson. ♀.

7. *C. strigosa*, filamentis simplicibus, silicula stylo mucronata, foliis ovatis dentatis basi inæqualibus subauriculatis cauleque fruticoso hispidis.

Crambe strigosa. Willden. *sp. pl.* 3, p. 420.

Myagrum arborescens. Jacqu. *ic.* 1, t. 120.

Nat. of the Canary Islands. Mr. Fr. Masson. ♀.

8. *C. filiformis*, filamentis simplicibus, silicula mutica a pedicello longiore solubili, foliis lyratis pilosis.

Crambe filiformis. Willden. *sp. pl.* 3, p. 419. Jacqu. *ic.* 3, t. 504.

Nat. of Patagonia. ♀.

[74]

MYAGRUM. Tournef. *inst.* 211.

Silicula monosperma evalvis absque suturarum vestigio (quandoque loculis spurii lateralibus). *Semen pendulum*.

1. *M. perfoliatum*, siliculis obcordatis: loculis duobus spuriiis, foliis amplexicaulibus.

Myagrum perfoliatum. Willden. *sp. pl.* 3, p. 407. Schkuhr *handb.* 2, p. 210, t. 178.

Nat. of France and Switzerland. ☉.

EUCLIDIUM.

Silicula ventricosa bilocularis, suturis manifestis valvarum non dehiscentium, loculis 1-spermis. Cotyledones planæ.

1. *E. syriacum*, siliculis scabris: stylo subulato persistenti, foliis lanceolatis petiolatis.

Bunias syriaca. Willden. *sp. pl.* 3, p. 413.

Anastatica syriaca. Jacqu. *austr.* 1, p. 7, t. 6.

Nat. of Austria and the Levant. ☉.

RAPISTRUM. *Gærtn. sem. 2, p. 285.*

Silicula bilocularis non dehiscens nec suturarum vestigio.
Cotyledones planæ. Calyx patens.

1. *R. ægyptiacum*, siliculis angulatis verrucoso-muricatis, foliis runcinatis.

Bunias ægyptiaca. Willden. sp. pl. 3, p. 414. Jacqu. hort. vindob. 2, p. 68, t. 145.

^{75]} *Nat. of Egypt and Greece. ☉.*

2. *R. paniculatum*, siliculis orbiculatis rugosiusculis, foliis lanceolatis subdentatis sagittatis amplexicaulibus.

Myagrum paniculatum. Willden. sp. pl. 3, p. 409. Fl. dan. 304.

Nat. of Europe. ☉.

BUNIAS. *Gen. pl. 1070.*

Silicula nucamentacea evalvis. Cotyledones spirales lineares. Germen biloculare, loculis 1-2 spermis.

1. *B. Erucago*, siliculis tetragonis 4-locularibus: angulis subdentatis.

Bunias Erucago. Willden. sp. pl. 3, p. 411. Jacqu. austr. 4, p. 21, t. 340.

Nat. of Austria and the South of France. ☉.

2. *B. orientalis*, siliculis ovatis gibbis subverrucosis, 1-2-locularibus.

Bunias orientalis. Willden. sp. pl. 3, p. 412. Houtt. nat. hist. 9, tab. 60, f. 2.

Nat. of the Levant. 4.

CORONOPUS.

Gærtn. sem. 2, p. 293. Smith fl. brit. 2, p. 690.

Silicula didyma evalvis aptera, loculis 1-spermis. *Cotyledones* incumbentes, lineares.

1. *C. Ruellii*, siliculis integris cristato-muricatis.
Coronopus Ruellii. *Smith fl. brit.* 2, p. 690. *Engl. bot.* 1660.

Cochlearia Coronopus. *Willden. sp. pl.* 3, p. 450.
Nat. of Britain. ☉.

2. *C. didyma*, siliculis emarginatis didymis rugosis, foliis pinnatifidis: laciniis lineari-lanceolatis indivisis incisive.

Coronopus didyma. *Smith fl. brit.* 2, p. 691.
Lepidium didymum. *Willden. sp. pl.* 3, p. 439. *Engl. bot.* 248.
Nat. of England. ☉.

BISCUTELLA. *Gen. pl.* 1084.

Silicula didyma, segmentis evalvibus foliaceo-compressis monospermis. *Radicula* descendens! *Cotyledones* accum-bentes inversæ.

1. *B. auriculata*, calycibus bisaccatis, siliculæ segmentis in stylum coeuntibus.

Biscutella auriculata. *Willden. sp. pl.* 3, p. 472.
Schkuhr handb. 2, p. 237, t. 182.
Nat. of France and Italy. ☉.

2. *B. apula*, siliculis scabris, foliis cuneato-lanceolatis serrato-dentatis hirsutis.

Biscutella apula. *Willden. sp. pl.* 3, p. 473.
Iondraba alyssoides apula spicata. *Column. ecphr.* 1, p. 283, t. 285, f. 1.
Nat. of Italy. ☉.

3. *B. coronopifolia*, siliculis glabris lævibus, foliis pinnatifido-dentatis hirtis.

Biscutella coronopifolia. *Linn. mant.* 255. *Decand. in annales du museum*, 18, p. 300, t. 14.

Nat. of Spain and Italy. ☉.

4. *B. lævigata*, siliculis glabris, foliis lanceolatis serrato-dentatis.

Biscutella lævigata. *Willden. sp. pl.* 3, p. 474. *Jacqu. austr.* 4, p. 20, t. 339.

Nat. of Austria and Italy. ♀.

5. *B. sempervirens*, siliculis scabriusculis, foliis dentatis tomentosis mollibus.

Biscutella sempervirens. *Willden. sp. pl.* 3, p. 475.

Nat. of Spain. ♀.

PELTARIA. *Gen. pl.* 1083.

Silicula orbiculata evalvis plana. Germen 2-3-spermum. Filamenta edentula. Cotyledones accumbentes.

1. *P. alliacea*, foliis amplexicaulibus oblongis indivisis. *Willden. sp. pl.* 3, p. 471. *Jacqu. austr.* 2, p. 14, t. 123.

^{78]} *Nat. of Austria.* ♀.

CLYPEOLA. *Gen. pl.* 1082.

Silicula evalvis subrotunda plana. Germen monospermum. Filamenta dentata.

1. *C. Ionthlaspi*, foliis integerrimis tomentosis, calycibus persistentibus, siliculis pubescentibus.

Clypeola Ionthlaspi. *Willden. sp. pl.* 3, p. 471. *Calvanill. ic.* 1, p. 22, t. 34, f. 2.

Nat. of France and Italy. ☉.

ISATIS. *Gen. pl.* 1072.

Silicula unilocularis monosperma: valvis navicularibus tardius dehiscentibus. *Filamenta* edentula, distincta.

1. *I. tinctoria*, siliculis obovato-oblongis glabris, foliis caulinis sagittatis glabris.

Isatis tinctoria. *Willden. sp. pl.* 3, p. 420. *Engl. bot.* 97. *Svensk bot.* 35.

Nat. of England. ♂.

2. *I. aleppica*, siliculis lineari-oblongis pubescentibus ciliatis.

Isatis lusitanica. *Linn, sp. pl.* 936.

Isatis aleppica. *Scop. insub.* 2, p. 31, t. 16.

Isatis orientalis. *Willden. enum.* 663.

Nat. of the Levant. ☉.

SUCCOWIA. *Mænoch method.* 265.

[79]

Silicula subglobosa stylo subulato cuspidata, valvis hemisphæricis echinatis: loculis monospermis. *Cotyledones* conduplicatæ.

1. *Succowia balearica*.

Bunias balearica. *Willden. sp. pl.* 3, p. 415. *Jacqu. hort. vindob.* 2, p. 68, t. 144.

Nat. of Minorca. ☉.

VELLA. *Gen. pl.* 1073.

Silicula ventricosa; stylo foliaceo ovato. *Cotyledones* conduplicatæ. *Calyx* clausus.

1. *V. annua*, foliis pinnatifidis, siliculis pendulis. *Willden. sp. pl.* 3, p. 422. *Engl. bot.* 1442.

Nat. of England. ☉.

2. *V. Pseudo-Cytisus*, foliis integris obovatis ciliatis, siliculis erectis. *Willden. sp. pl.* 3, p. 422. *Cavanill. ic.* 1, p. 32, t. 42.

Nat. of Spain. ♀.

ANASTATICA. *Gen. pl.* 1074.

Silicula ventricosa; valvulis juxta apicem auriculâ auctis. *Cotyledones* planæ.

1. *ANASTATICA hierochuntica.* *Willden. sp. pl.* 3, p. 423. *Jacqu. hort. vindob.* 1, p. 23, t. 58.

Nat. of the Levant. γ.

80]

ÆTHIONEMA.

Silicula valvis navicularibus alatis (quandoque non dehiscentibus). *Filamenta* longiora vel connata vel juxta apicem denticulata. *Calyx* insertione inæqualis.

1. *Æ. saxatile*, filamentis longioribus distinctis, foliis oblongis basi attenuatis.

Thlaspi saxatile. *Willden. sp. pl.* 3, p. 444. *Jacqu. austr.* 3, p. 21, t. 236. *Schkuhr handb.* 2, p. 223, t. 180.

Nat. of the South of Europe. ☉.

2. *Æ. monospermum*, siliculis evalvibus monospermis, foliis ovalibus obovatisve.

Nat. of Spain. ♂.

THLASPI. *Gen. pl.* 1078.

Silicula compressa emarginata, valvis navicularibus (sæpius alatis), polysperma. *Filamenta* edentula distincta. *Calyx* insertione æqualis, patens.

1. *T. arvense*, siliculis orbiculatis: ala dilatata longitudinali, seminibus concentrice striatis, foliis oblongis sagittatis dentatis glabris.

Thlaspi arvense. Willden. *sp. pl.* 3, p. 442. Curtis
lond. Engl. bot. 1659. *Fl. dan.* 793. *Svensk bot.* 214.
Nat. of Britain. ☉.

2. *T. alliaceum*, siliculis subrotundis ventricosus: ala
 angustata deorsim obsoleta, foliis oblongis obtusis acute
 sagittatis subdentatis glabris.

Thlaspi alliaceum. Willden. *sp. pl.* 3, p. 443. Jacqu.
ic. 1, t. 121.

Nat. of the South of Europe. ☉.

3. *T. ceratocarpon*, siliculis ventricosus; alis apicis acu-
 tissimis, foliis sagittatis lanceolatis subdentatis glabris.

Thlaspi ceratocarpon. Willden. *sp. pl.* 3, p. 448. Scop.
insub. 1, p. 10, t. 4.

Nat. of Siberia. ☉.

4. *T. Bursa pastoris*, siliculis obcordatis apteris, foliis
 radicalibus pinnatifidis.

Thlaspi Bursa pastoris. Willden. *sp. pl.* 3, p. 447.
Curtis lond. Engl. bot. 1485.

Nat. of Britain. ☉.

5. *T. perfoliatum*, siliculis obcordatis alatis; stylo incluso
 brevissimo, foliis caulinis cordatis subdentatis glabris.

Thlaspi perfoliatum. Willden. *sp. pl.* 3, p. 446. Jacqu.
austr. 4, p. 19, t. 337.

Nat. of England. ♀.

6. *T. alpestre*, siliculis obovatis retusis: loculis 4—6-
 spermis; stylo exserto, staminibus longitudine petalorum,
 foliis caulinis cordato-sagittatis, caule simplici.

Thlaspi alpestre. Willden. *sp. pl.* 3, p. 447. Engl.
bot. 81.

Nat. of England. ♀.

[82]

7. *T. montanum*, siliculis retusis: loculis 2-spermis; stylo
 exserto, staminibus corolla brevioribus, petalis calycem ter
 superantibus, foliis caulinis cordato-sagittatis.

Thlaspi montanum. *Willden. sp. pl.* 3, p. 445. *Jacqu. austr.* 3, p. 22, t. 237.
Nat. of Austria. ♂.

HUTCHINSIA.

Silicula elliptica integra: valvis navicularibus apteris, loculis dispermis. *Filamenta* edentula.

1. *H. rotundifolia*, foliis inferioribus orbiculato-ovatis petiolatis; superioribus cordato-sagittatis integris.

Iberis rotundifolia. *Willden. sp. pl.* 3, p. 454.

Lepidium rotundifolium. *Allion. pedem.* 1, p. 252. t. 55, f. 2.

Nat. of Switzerland and Piedmont. ♀.

2. *H. alpina*, foliis pinnatis integerrimis glabris, petalis calyce deciduo duplo longioribus, siliculis utrinque acutis; stylo brevissimo exserto.

Lepidium alpinum. *Willden. sp. pl.* 3, p. 433. *Jacqu. austr.* 2, p. 23, t. 137.

Nat. of the Alps of Germany, Switzerland, and Italy. ♀.

3. *H. petræa*, foliis pinnatis integerrimis, petalis calyce vix longioribus, siliculis utrinque obtusis; stigmatibus sessilibus.

Lepidium petræum. *Willden. sp. pl.* 3, p. 434. *Engl. bot.* 111.

Nat. of England. ♂.

TEESDALIA.

Silicula emarginata, valvis navicularibus, loculis dispermis. *Filamenta* basi intus squamula aucta.

1. *TEESDALIA nudicaulis*.

Iberis nudicaulis. *Willden. sp. pl.* 3, p. 458. *Curtis lond. Engl. bot.* 327.

Nat. of Britain. ☉.

IBERIS. *Gen. pl.* 1080.

Silicula emarginata : valvis navicularibus alatis ; loculis monospermis. *Petala* inæqualia.

1. *I. semperflorens*, frutescens, foliis spathulatis glabris integerrimis obtusis. *Willden. sp. pl.* 3, p. 452.

Thlaspidium fruticosum Leucoii folio semperflorens. *Zanon. hist.* 214, t. 165.

Nat. of Persia and Sicily. ♀.

2. *I. sempervirens*, frutescens, foliis oblongo-linearibus integerrimis glabris, corymbis racemosis.

Iberis sempervirens. *Willden. sp. pl.* 3, p. 453.

Thlaspidium. *Rivin. tetrapet.* 110.

Nat. of the Island of Candia. ♀.

3. *I. gibraltarica*, frutescens, foliis cuneiformibus obtusis apice dentatis glabris. *Willden. sp. pl.* 3, p. 453.

Curtis magaz. 124.

Nat. of Spain. ♀.

4. *I. saxatilis*, frutescens, foliis integerrimis acutis pubescentibus : inferioribus linearibus ; superioribus lanceolatis.

Willden. sp. pl. 3, p. 453.

Thlaspi saxatile, vermiculato folio. *Garid. prov.* 460, t. 101.

Nat. of France and Italy. ♀.

5. *I. ciliata*, herbacea, foliis subcarnosis ciliatis : radicalibus spathulatis ; caulinis linearibus. *Willden. sp. pl.* 3, p. 455. *Botan. magaz.* 1030.

Nat. of Mount Caucasus. ♀.

6. *I. umbellata*, herbacea, foliis lanceolatis acuminatis ; inferioribus serratis ; superioribus integerrimis. *Willden. sp. pl.* 3, p. 456. *Curtis magaz.* 106.

Nat. of the South of Europe. ☉.

7. *I. amara*, herbacea, foliis lanceolatis acutiusculis subdentatis glabris, floribus racemosis,

⁸⁶¹ *Iberis amara*. *Willden. sp. pl.* 3, p. 456. *Engl. bot.* 52.
Nat. of England. ☉.

8. *I. violacea*, herbacea, foliis petiolatis spathulatis obtusis dentatis integerrimisque ciliatis, corymbo subumbellato.

Nat. ☉.

9. *I. linifolia*, herbacea, foliis linearibus integerrimis glabris, caule ramoso, corymbis hemisphæricis.

Iberis linifolia. *Willden. sp. pl.* 3, p. 457.

Thlaspi lusitanicum umbellatum gramineo folio flore purpurascente. *Garid. prov.* 459, t. 105.

Nat. of Spain and Portugal. ♂.

10. *I. pinnata*, herbacea, foliis pinnatifidis, caule erecto paniculato, petalis calyce triplo longioribus.

Iberis pinnata. *Willden. sp. pl.* 3, p. 458.

Thlaspi alterum minus umbellatum nasturtii hortensis folio, narbonense. *Lobel. ic.* 218.

Nat. of the South of Europe. ☉.

LEPIDIDIUM. *Gen. pl.* 1077.

Silicula loculis monospermis; valvis navicularibus.
Petala æqualia.

1. *L. latifolium*, foliis lanceolatis indivisis serratis integerrimisve, siliculis ovalibus integris.

⁸⁶¹ *Lepidium latifolium*. *Willden. sp. pl.* 3, p. 436. *Engl. bot.* 182.

Nat. of Britain. ♀.

2. *L. lyratum*, foliis lyratis crispis, siliculis ovatis integris.

Lepidium lyratum. *Willden. sp. pl.* 3, p. 435.

Lepidium orientale, nasturtii crispi folio. *Tournef. it.* 2, p. 339, cum tab.

Nat. of the Levant. ☉.

3. *L. Draba*, foliis amplexicaulibus indivisis lanceolatis oblongisve dentatis, siliculis cordatis apice integris stylo acuminatis.

Cochlearia Draba. *Willden. sp. pl.* 3, p. 451. *Jacqu. austr.* 4, p. 8, t. 315.

Nat. of Austria, France, and Italy. ♀.

4. *L. graminifolium*, foliis caulinis linearibus integris, siliculis ovatis acutis integerrimis, floribus hexandris, petalis calyce longioribus.

Lepidium graminifolium. *Linn. sp. pl.* 900.

Lepidium Iberis. *Linn. sp. pl.* 900, secundum synonyma omnia.

Nat. of Germany, France, and Italy. ♀.

5. *L. divaricatum*, foliis inferioribus pinnatifidis; rameis linearibus, caule ramosissimo, siliculis ovalibus obsolete emarginatis imbricatis.

Lepidium divaricatum. *Willden. sp. pl.* 3, p. 441. ¹⁸⁷

Nat. of the Cape of Good Hope. *Mr. Fr. Masson.* 2.

6. *L. ruderale*, floribus diandris apetalis, foliis radicalibus pinnatifidis; rameis linearibus integerrimis, siliculis emarginatis patentibus.

Lepidium ruderale. *Willden. sp. pl.* 3, 440. *Engl. bot.* 1595.

Nat. of Britain. ☉.

7. *L. bonariense*, floribus diandris, petalis minutis, foliis caulinis pinnatifidis incisive cauleque pilosiusculis, siliculis subrotundis emarginatis.

Lepidium bonariense. *Willden. sp. pl.* 3, p. 441.

Thlaspi bonariense multiscissum, flore invisibili. *Dill. elth.* 381, t. 286, f. 370.

Nat. of South America. ☉.

8. *L. Cardamines*, foliis radicalibus pinnatis; caulinis lyratis, siliculis emarginatis.

Lepidium Cardamines. *Willden. sp. pl.* 3, p. 434. *Act. stockholm.* 1755, p. 273, t. 3, 9.

Nat. of Spain. ♂.

9. *L. subulatum*, foliis subulatis integerrimis cauleque suffruticoso pubescentibus, siliculis emarginatis; stigmatē semiexserto.

Lepidium subulatum. *Willden. sp. pl.* 3, p. 438. *D'Asso aray.* 83, t. 6, f. 3.

^{88]} *Nat. of Spain.* 2.

10. *L. piscidium*, foliis ovali-oblongis extrorsum dentatis integerrimisve, siliculis oblongo-obovatis emarginatis; stigmatē exserto.

Lepidium piscidium. *Willden. sp. pl.* 3, p. 437.

Lepidium oleraceum. *Edit. prior.* 2, p. 374, excluso synonymo *Forsteri*.

Lepidium bidentatum. *Montin in nov. act. acad. nat. curios.* 6, p. 324, tab. 5, a.

Nat. of the Society Islands. ☉.

11. *L. perfoliatum*, foliis caulinis pinnatis bipinnatisque; rameis cordatis amplexicaulibus integris.

Lepidium perfoliatum. *Willden. sp. pl.* 3, p. 431. *Jacqu. austr.* 4, p. 24, t. 346.

Nat. of Austria and the Levant. ☉.

12. *L. spinosum*, foliis pinnatis, siliculis cuneatis semi-bifidis.

Lepidium spinosum. *Willden. sp. pl.* 3, p. 434.

Nat. of the Levant. ☉.

13. *L. campestre*, siliculis ovatis emarginatis alatis, foliis caulinis sagittatis dentatis.

Thlaspi campestre. *Willden. sp. pl.* 3, p. 444. *Curt. lond. Engl. bot.* 1385.

Nat. of Britain. ☉.

^{89]} 14. *L. sativum*, cotyledonibus tripartitis.

Lepidium sativum. *Willden. sp. pl.* 3, p. 435. *Zorn. ic.* 16.

α. *Nasturtium hortense vulgatum*. *Bauh. pin.* 103.

β. *Nasturtium hortense crispum*. *Bauh. pin.* 104.
prodr. 44, *cum ic. in p.* 43 *et* 44.

Nat. ☉.

15. *L. virginicum*, floribus 2-4 andris tetrapetalis, foliis
 caulinis lineari-lanceolatis inciso-serratis, siliculis orbiculatis
 emarginatis.

Lepidium virginicum. *Willden. sp. pl.* 3, p. 440.

Lepidium Iberis. *Schkuhr handb.* 2, p. 222, t. 180.

Nat. of North America and Jamaica. ☉.

COCHLEARIA. *Gen. pl.* 1079.

Silicula subovata polysperma : valvis ventricosis. *Semina*
 immarginata. cotyledonibus accumbentibus. *Filamenta*
 breviora edentula. *Calyx* patens.

1. *C. officinalis*, siliculis subrotundis, foliis radicalibus
 reniformibus.

Cochlearia officinalis. *Willden. sp. pl.* 3, p. 448. *Engl.*
bot. 551. *Svensk bot.* 87.

Nat. of Britain. ☉.

2. *C. anglica*, siliculis ellipticis, foliis radicalibus ovatis
 integris.

Cochlearia anglica. *Willden. sp. pl.* 3, p. 449. *Engl.*
bot. 552.

Nat. of Britain. ☉.

[90]

3. *C. danica*, siliculis ellipticis, foliis omnibus deltoideis.
Cochlearia danica. *Willden. sp. pl.* 3, p. 449. *Engl.*
bot. 696.

Nat. of Britain. ☉.

4. *C. glastifolia*, siliculis sphæricis reticulato-venosis,
 foliis caulinis amplexicaulibus sagittato-cordatis acutis.

Cochlearia glastifolia. *Willden. sp. pl.* 3, p. 451.

Lepidium glastifolium perenne. *Moris. hist.* 2, p. 312, s. 3, t. 21, f. 3.

Nat. of Germany. ♂.

5. *C. saxatilis*, siliculis lentiformibus lævibus, foliis radicalibus oblongis dentatis pilosis.

Myagrum saxatile. *Willden. sp. pl.* 3, p. 409. *Jacqu. austr.* 2, p. 17, t. 128.

Nat. of Austria and Switzerland. ♀.

6. *C. Armoracia*, siliculis oblongis; stigmatibus dilatato subsessilibus, foliis radicalibus oblongis crenatis; caulinis elongato-lanceolatis dentatis incisive.

Cochlearia Armoracia. *Willden. sp. pl.* p. 451. *Engl. bot.* 2223. *Schkuhr handb.* 2, p. 229, t. 181.

Nat. of England. ♀.

SUBULARIA. *Gen. pl.* 1075.

Silicula ovalis - mutica polysperma: valvis ventricosis. *Cotyledones* incumbentes lineares biveres.

1. SUBULARIA *aquatica*.

Subularia aquatica. *Willden. sp. pl.* 3, p. 423. *Engl. bot.* 732.

Nat. of Britain. ☉.

DRABA. *Gen. pl.* 1076.

Silicula integra ovalis: valvis planis v. convexiusculis; loculis polyspermis. *Semina* immarginata: cotyledonibus accumbentibus. *Filamenta* edentula.

1. *D. verna*, scapis nudis, petalis bipartitis, foliis lanceolatis subincisis hirtis. *Smith fl. brit.* 2, p. 677. *Engl. bot.* 586. *Curtis lond. fl. dan.* 983.

Nat. of Britain. ☉.

2. *D. rupestris*, scapis nudis monophyllisve; petalis

indivisis, siliculis lanceolatis pubescentibus, foliis planis lanceolatis pilosis.

Draba hirta. *Smith fl. brit.* 2, p. 677 (excluso synonymo Linnæi et Floræ danicæ). *Engl. bot.* 1338.

Nat. of Scotland. ♀.

3. *D. incana*, foliis caulinis numerosis incanis subdentatis, siliculis oblongis glabris contortis.

Draba incana. *Willden. sp. pl.* 3, p. 430. *Engl. bot.* 388.

Nat. of Britain. ♂.

[92]

4. *D. muralis*, caule ramoso, foliis ovatis amplexicaulibus dentatis, siliculis patentibus glabris. *Willden. sp. pl.* 3, p. 429. *Engl. bot.* 912. *Scop. insubr. fasc.* 2, t. 15.

Nat. of England. ☉.

5. *D. nemoralis*, caule ramoso, foliis ovatis amplexicaulibus dentatis, siliculis patentibus pubescentibus. *Willden. sp. pl.* 3, p. 429.

Nat. of Europe. ☉.

6. *D. aizoides*, scapis nudis glabris, foliis lanceolato-linearibus rigidis carinatis ciliatis, staminibus petala æquantibus, stylo longitudine dimidii germinis.

Draba aizoides. *Willden. sp. pl.* 3, p. 424 (exclusis synonymis *Milleri* et *Curtisii*). *Engl. bot.* 1271.

Nat. of Wales. ♀.

7. *D. ciliaris*, scapis nudis glabris, foliis elongato-linearibus ciliatis, staminibus calycem vix æquantibus.

Draba ciliaris. *Willden. sp. pl.* 3, p. 425.

Draba aizoides. *Curtis magaz.* 170.

Alyssum alpinum hirsutum luteum. *Mill. ic.* 1, p. 14, t. 20, f. 2.

Nat. of Switzerland. ♀.

PETROCALLIS.

Silicula integra ovalis: valvis planiusculis; loculis dispermis. *Semina* immarginata: funiculis umbilicalibus dissepimento adnatis! *Filamentu* edentula.

1. PETROCALLIS *pyrenaica*.

Draba pyrenaica. *Willden. sp. pl.* 3, p. 428. *Botan. magaz.* 713.

Nat. of Austria and the Pyrenees. 4.

CAMELINA. *Crantz austr.* 1, p. 18.

Silicula subovata polysperma: valvis ventricosis. *Cotyledones* incumbentes. *Filamenta* edentula.

1. *C. sativa*, siliculis obovatis marginatis, stigmatibus simplicibus, foliis lanceolatis sagittatis.

Myagrum sativum. *Willden. sp. pl.* 3, p. 408. *Fl. dan.* 1038.

Alyssum sativum. *Engl. bot.* 1254.

Nat. of Britain. ☉.

2. *C. austriaca*, siliculis subrotundis immarginatis, stigmatibus capitatis, foliis oblongis sagittatis dentatis glabris.

Alyssum austriacum. *Willden. sp. pl.* 3, p. 408. *Jacq. austr.* 2, p. 7, t. 111.

Nat. of Austria. ☉.

ALYSSUM. *Lamarck illustr. tab.* 559.

Silicula subrotunda: valvis omnino vel limbo planiusculis; loculis 1—4-spermis. *Semina* marginata. *Cotyledones* 94] incumbentes.

* *Herbacea.*

1. *A. calycinum*, siliculis emarginatis pubescentibus, calycibus persistentibus, filamentis edentulis: brevioribus stipatis pari setarum.

Alyssum calycinum. Willden. *sp. pl.* 3, p. 464.

Alyssum campestre. Schkehr *handb.* 2, p. 234, t. 181.

Nat. of Austria and France. ☉.

2. *A. campestre*, siliculis emarginatis pubescentibus stylo 4-plo longioribus, calycibus deciduis, filamentis omnibus appendiculatis, foliis pedunculisque pube stellata sparsa.

Alyssum campestre. Willden. *sp. pl.* 3, p. 457.

Nat. of France. ☉.

3. *A. montanum*, siliculis emarginatis incanis stylo parum longioribus, calycibus deciduis, filamentis omnibus appendiculatis, foliis ramisque incanis.

Alyssum montanum. Willden. *sp. pl.* 3, p. 466. *Curtis magaz.* 419.

Nat. of Germany and Switzerland. ♀.

** *Fruticosa.*

4. *A. saxatile*, siliculis emarginatis glabris, filamentis longioribus simplicibus, foliis oblongis lanceolatisve repandis tomentosis.

Alyssum saxatile. Willden. *sp. pl.* 3, p. 460. *Curtis magaz.* 159.

Nat. of the Island of Candia. ♀.

5. *A. alpestre*, siliculis integris, filamentis omnibus appendiculatis, foliis obovatis spathulatisque incanis, caulibus adscendentibus suffruticosis.

Alyssum alpestre. Willden. *sp. pl.* 3, p. 461. *Jacqu. collect.* 4, p. 227, t. 4, f. 1. *Alli.n. pedem.* 1, p. 241, t. 18, f. 2.

Nat. of Italy. ♀.

6. *A. spinosum*, siliculis integris glabris, filamentis edentulis, ramis floriferis persistentibus spinescentibus, foliis spatulatis incanis; pube stellari minutissima.

Alyssum spinosum. Willden. *sp. pl.* 3, p. 459.

Thlaspi spinosum hispanicum. Barrel. *ic.* 808.

Nat. of the South of Europe. η .

7. *A. maritimum*, siliculis integris glabris: loculis monospermis, filamentis edentulis, foliis lineari-lanceolatis acutis subincanis; pilis bipartitis adpressis.

Alyssum maritimum. Willden. *sp. pl.* 3, p. 459. *Engl. bot.* 1729.

Alyssum halimifolium. Linn. *sp. pl.* 907. *Curtis magaz.* 101.

Alyssum minimum. Linn. *sp. pl.* 908.

Nat. of England. η .

FARSETIA.

Silicula ovali-oblonga polysperma sessilis, valvis planis v. convexiusculis (non inflatis). *Cotyledones* accumbentes. *Semina* marginata vel *Filamenta* quædam denticulata.

* *Semina marginata*. *Valvæ planæ*.

1. *F. Cheiranthoides*, caulibus fruticosis erectis, foliis linearibus incanis, calycibus clausis.

Cheiranthus Farsetia. Willden. *sp. pl.* 3, p. 526. *Desfont. atlant.* 2, p. 89, t. 160.

Nat. of the Levant. η .

2. *F. lunarioides*, caulibus suffruticosis adscendentibus, foliis spatulatis petiolatis siliculisque tomentosis incanis.

Alyssum lunarioides. Willden. *sp. pl.* 3, p. 461.

Lunaria fruticosa perennis incana Leucoii foliis. *Turnef. it.* 1, p. 242, *cum tab.*

Nat. of the Archipelago. γ .

3. *F. clypeata*, caulibus herbaceis erectis, foliis oblongis repandis.

Alyssum clypeatum. Willden. *sp. pl.* 3, p. 468.

Alyssum Dioscoridis. Dod. *pempt.* 89.

Nat. of the South of Europe. ☉.

** *Valvæ convexiusculæ*. *Filamenta breviora denticulata*.

4. *F. mutabilis*, siliculis glabris compressis, seminibus marginatis.

Alyssum mutabile. Venten. *cels.* 85.

[97

Nat. of the Levant. ♀.

5. *F. incana*, siliculis pubescentibus subventricosus, seminibus immarginatis.

Alyssum incanum. Willden. *sp. pl.* 3, p. 463.

Thlaspi incanum machliniense. Clus. *hist.* 2, p. 132.

Nat. of the North of Europe. ♂.

* * *Calyx basi bisaccatus*. *Valvæ convexiusculæ*.

6. *F. deltoidea*, caulibus diffusis, foliis pilosis.

Alyssum deltoideum. Willden. *sp. pl.* 3, p. 470. Curtis *magaz.* 126.

Nat. of the Levant. ♀.

VESICARIA. *Tamarck illustr. tab.* 559.

Silicula inflata, subglobosa, polysperma. *Cotyledones accumbentes*.

1. *V. utriculata*, calycibus bisaccatis, filamentis omnibus edentulis, foliis integerrimis glabriusculis.

Alyssum utriculatum. Willden. *sp. pl.* 3, p. 470. Curtis *magaz.* 130. *Waldst. et Kitaib. hung.* 2, p. 215, t. 196.

Nat. of the Levant and Hungary. ♀.

2. *V. sinuata*, foliis incanis subdentatis, siliculis membranaceis glabris stylo cuspidatis.

Alyssum sinuatum. Willden. *sp. pl.* 3, p. 468.

Eruca peregrina. Clus. *hist.* 2, p. 134.

Nat. of Spain.

LUNARIA. *Gen. pl.* 1085.

Silicula elliptica pedicellata polysperma: valvis planis.
Semina marginata. *Cotyledones* accumbentes.

1. *L. rediviva*, siliculis oblongis utrinque acutis, pedicello germinis stamina breviora superante, dentibus foliorum mucronatis.

Lunaria rediviva. Willden. *sp. pl.* 3, p. 476.

Lunaria græca perennis. Besl. *cyst. vern.* 7, f. 2.

Nat. of Austria and Hungary. ☿.

2. *L. annua*, siliculis ovalibus utrinque obtusis, staminibus brevioribus pedicellum germinis superantibus, dentibus foliorum submuticis.

Lunaria annua. Willden. *sp. pl.* 3, p. 477. J. Miller *illustr.*

Nat. of Germany. ☉.

RICOTIA. *Gen. pl.* 1086.

Silicula oblonga sessilis: valvis planis; septo oblitterato unilocularis. *Cotyledones* accumbentes obcordatæ.

1. *RICOTIA ægyptiaca.* Willden. *sp. pl.* 3, p. 477.

⁹⁹⁾ *Lunaria* foliis supradecompositis: foliolis trifidis, siliquis oblongis pendulis. Mill. *ic.* 2, p. 113, t. 169.

Nat. of Egypt. ☉.

SILICUOSA.

HELIOPHILA. *Gen. pl.* 1092.

Siliqua integra v. moniliformis. *Cotyledones* incumbentes curvatæ (2-3-crures) lineares. *Filamenta breviora* basi extus dentata dum *siliqua* integra.

1. *H. incana*, pubescens, foliis spathulatis integerrimis, siliquis teretiusculis pubescentibus erectis, caule fruticoso.

Heliophila incana. *Willden. sp. pl.* 3, p. 527, excluso synonymo *Burmanni*.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* h.

2. *H. filiformis*, glabra, foliis angusto-linearibus elongatis integerrimis, siliquis teretibus subulatis pendulis.

Heliophila filiformis. *Willden. sp. pl.* 3, p. 529.

Nat. of the Cape of Good Hope. ☉.

3. *H. platysiliqua*, glabra, foliis carnosis semiteretibus integerrimis, siliquis plano-compressis pendulis, caule suffruticoso.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* h.

4. *H. pilosa*, hirsuta: pilis patulis, foliis planis nunc linearibus integerrimis nunc divisis subcuneiformibus, siliquis teretiusculis glabris integerrimis.

Heliophila pilosa. *Lamarck encycl.* 3, p. 90.

a integrifolia. *Heliophila integrifolia*. *Willden. sp. pl.* 3, p. 527.

Heliophila. *Burmman, in nov. act. upsal*, 1, p. 94, t. 7.

β incisa. *Heliophila arabioides*. *Botan. magaz.* 496.

Nat. of the Cape of Good Hope. ☉.

5. *H. amplexicaulis*, glabra, foliis lineari-lanceolatis amplexicaulibus integerrimis suboppositis, siliquis moniliformibus.

Heliophila amplexicaulis. Willden. *sp. pl.* 3, p. 528.
Nat. of the Cape of Good Hope. ☉.

6. *H. fœniculacea*, pilosiuscula, foliis pinnatis bipinnatis-
 que filiformibus, siliquis linearibus, caule erecto angulato.
Nat. of the Cape of Good Hope. Mr. Fr. Masson. ☉.

7. *H. pinnata*, glabra, foliis pinnatis filiformibus, siliquis
 toroso-moniliformibus pendulis.

Heliophila pinnata. Venten. *malmais.* 113, exclusis
 synonymis.

Heliophila pendula. Willden. *sp. pl.* 3, p. 529.
Nat. of the Cape of Good Hope. ☉.

101]

CARDAMINE.

Gen. pl. 1088. *Dentaria.* *Gen. pl.* 1087.

Siliqua linearis marginibus truncatis: valvis planis
 enervibus (elastice sæpius dissilientibus), dissepimento an-
 gustioribus.

1. *C. enneaphylla*, caule simplicissimo apice triphylo,
 foliis ternatis: foliolis lanceolatis inciso-serratis, staminibus
 corollam subæquantibus.

Dentaria enneaphylla. Willden. *sp. pl.* 3, p. 478.
Jacqu. austr. 4, p. 9, t. 316.

Nat. of Austria and Italy. ♂.

2. *C. pentaphylla*, caule simplicissimo apice triphylo, foliis
 quinato-digitatis: foliolis elliptico-lanceolatis inciso-serratis,
 corolla staminibus duplo longiore.

Dentaria pentaphylla. Willden. *sp. pl.* 3, p. 480.

Dentaria pentaphyllos foliis mollibus. *Garid. prov.* 152,
 t. 29.

Nat. of Switzerland and the South of France. ♀.

3. *C. pinnata*, caule simplicissimo apice triphylo, foliis
 pinnatis: foliolis (5—7) elongato-lanceolatis inciso-serratis.

Dentaria pinnata. Willden. *sp. pl.* 3, p. 480.

Dentaria heptaphyllos. Garid. *prov.* 152, t. 28.

Nat. of Switzerland and the South of France. ♀.

4. *C. bulbifera*, caule simplicissimo, foliis inferioribus pinnatis; superioribus simplicibus.

Dentaria bulbifera. Willden. *sp. pl.* 3, p. 479. Eng. ^[102] *bot.* 309. *Select specim.* 7.

Nat. of England. ♀.

5. *C. asarifolia*, foliis simplicibus reniformibus.

Cardamine asarifolia. Willden. *sp. pl.* 3, p. 482.

Nasturtium montanum asari folio. Herm. *parad.* 203, *cum tab.*

Nat. of Italy. ♀.

6. *C. trifolia*, foliis ternatis: foliolis rhombeo-subrotundis denticulatis, caule subnudo simplici.

Cardamine trifolia. Willd. *sp. pl.* 3, p. 483. *Botan. magaz.* 452.

Nat. of Lapland, Austria and Switzerland. ♀.

7. *C. africana*, foliis ternatis acuminatis, caule ramosissimo. Willden. *sp. pl.* 3, p. 483.

Nasturtium africanum, foliis ternis, facie Christophorianæ. Herm. *parad.* 202, *cum. tab.*

Nat. of the Cape of Good Hope. ♀.

8. *C. chelidonia*, foliis pinnatis: foliolis quinis crenato-incisis: lateralibus ternatis auriculatisve, geruine piloso.

Cardamine chelidonia. Willd. *sp. pl.* 3, p. 484. *Waldst. et Kitaib.* 2, p. 149, t. 140.

Nat. of Italy. ♀.

9. *C. græca*, foliis pinnatis: foliolis ovatis lobatis obtusis.

Cardamine græca. Willden. *sp. pl.* 3, p. 484.

Sio minimo Prosperi Alpini affinis, siliquis latis. *Boccon. sic.* 84, t. 44, f. 2.

Nat. of the South of Europe. ☉.

10. *C. amara*, foliis pinnatis: foliolis radicalibus subrotundis; caulinis dentato-angulatis, stylo obliquo, stigmate acutiusculo, caule basi radicante.

Cardamine amara. *Willden. sp. pl.* 3, p. 488. *Curtis lond. Engl. bot.* 1000.

Nat. of Britain. ♀.

11. *C. latifolia*, foliis pinnatis glabris: foliolis (5—7) radicalibus subrotundis; caulinis ovatis dentato-angulatis, stigmate capitato.

Cardamine latifolia. *Willden. sp. pl.* 3, p. 487.

Nasturtium pyrenaicum aquaticum latifolium purpurascente flore. *Herm. paradis.* 203, *cum tab.*

Nat. of Spain. ♂.

12. *C. pratensis*, foliis pinnatis: foliolis radicalibus subrotundis; caulinis lanceolatis, stylo stricto, stigmate capitato.

Cardamine pratensis. *Willden. sp. pl.* 3, p. 487. *Curtis lond. Engl. bot.* 776. *Fl. dan.* 1039.

α floribus simplicibus.

β floribus plenis.

¹⁰⁴⁵ *Nat. of Britain.* ♀.

13. *C. impatiens*, foliis pinnatis: foliolis lanceolatis subincisis, stipulis ciliatis, petalis linearibus nullisve.

Cardamine impatiens. *Willden. sp. pl.* 3, p. 485. *Engl. bot.* 80.

Nat. of Britain. ☉.

14. *C. hirsuta*, foliis omnibus pinnatis exstipulatis: foliolis petiolatis: radicalibus subrotundis, staminibus (4—6) petala subæquantibus, stigmate subsessili.

Cardamine hirsuta. *Willden. sp. pl.* 3, p. 486. *Curtis lond. Engl. bot.* 492.

Nat. of Britain. ☉.

15. *C. resedifolia*, foliis inferioribus indivisis; superioribus tripartitis pinnatisve stipulatis.

- Cardamine resedifolia*. Willden. *sp. pl.* 3, p. 482.
Allion. pedem. 1, p. 261, t. 57, f. 2.
Nat. of Switzerland and Germany. ☉.

ARABIS. *Gen. pl.* 1094.

Siliqua linearis stigmatē subsessili coronata: valvis
 venosis v. nervosis. *Semina* uniseriata. *Cotyledones*
 accumbentes. *Calyx* erectus.

1. *A. cærulea*, foliis obovatis glabris ciliatis apice sub-
 dentatis, caule paucifolio, racemo cernuo.

Arabis cærulea. Willden. *sp. pl.* 3, p. 537.

Turritis cærulea. Allion. *pedem.* 1, p. 270, t. 40, f. 2.

Nat. of Switzerland. ۲. [105]

2. *A. nutans*, foliis subdentatis: radicalibus obovatis
 utrinque pilosis scabris; caulinis ovalibus, racemo subnu-
 tante.

Arabis nutans. Willden. *sp. pl.* 3, p. 537.

Arabis pumila. Jacqu. *austr.* 3, p. 44, t. 281.

Nat. of Switzerland, Austria and Italy. ۲.

3. *A. bellidifolia*, foliis subdentatis glabris: radicalibus
 cuneato-obovatis; caulinis oblongis sessilibus, racemo erecto.

Arabis bellidifolia. Willden. *sp. pl.* 3, p. 537. Jacqu.
austr. 3, p. 44, t. 280.

Nat. of Austria and Switzerland. ۲.

4. *A. alpina*, foliis serrato-dentatis pubescentibus; cau-
 linis amplexicaulibus, pedunculis longitudine calycis, valvis
 planis, stigmatē integro.

Arabis alpina. Willden. *sp. pl.* 3, p. 534. Curtis
magaz. 226.

Nat. of Switzerland. ۲.

5. *A. verna*, foliis pubescentibus dentatis: caulinis cor-
 datis amplexicaulibus, pedunculis calyce duplo brevioribus,
 valvis convexis, stigmatē emarginato.

Hesperis verna. Willden. *sp. pl.* 3, p. 533.

Leucojum maritimum latifolium annuum. Moris. *hist.* 2, p. 241, s. 3, t. 8, f. 5.

1067 *Nat. of the South of France.* ☉.

6. *A. stricta*, foliis dentatis obtusis hispidis: radicalibus sublyratis, caulibus hispidis, petalis erectis. *Smith fl. brit.* 2, p. 712. *Engl. bot.* 614.

Arabis stricta. Willden. *sp. pl.* 3, p. 539.

Nat. of England. ♀.

7. *A. arenosa*, foliis caulinis sinuato-pinnatifidis petiolatis: pube ramosa, caule hispido: pilis simplicibus, petalis calyce triplo longioribus.

Arabis arenosa. Scop. *carn. n.* 837, t. 40.

Sisymbrium arenosum. Willden. *sp. pl.* 3, p. 498.

Nat. of Germany and Switzerland. ☉.

8. *A. hispida*, foliis radicalibus sinuato-lyratis elongato-petiolatis; caulinis plerisque indivisis glabris, pedunculis fructiferis patulis longitudine dimidii siliquæ.

Arabis hispida. *Smith. fl. brit.* 2, p. 713. Willden. *sp. pl.* 3, p. 538.

Cardamine hastulata. *Engl. bot.* 469.

Nat. of Britain. ♀.

9. *A. thaliana*, foliis dentatis pilosis: radicalibus oblongis, staminibus petala subæquantibus siliquis adscendentibus.

Arabis thaliana. Willden. *sp. pl.* 3, p. 535. *Curtis lond. Engl. bot.* 901. *Fl. dan.* 1106.

Nat. of Britain. ☉.

1071 10. *A. ciliata*, foliis subdentatis ovalibus glabris ciliatis: radicalibus subsessilibus obtusis; caulinis semi-amplexicaulibus, caule simplici.

Turritis alpina. *Engl. bot.* 1746. Willden. *sp. pl.* 3, p. 545.

Turritis ciliata. Willden. *sp. pl.* 3, p. 544.

Nat. of Ireland. ♂.

11. *A. hirsuta*, foliis omnibus hispidis dentatis : caulinis semi-amplexicaulibus, siliquis strictis.

Turritis hirsuta. Willden. *sp. pl.* 3, p. 543. *Engl. bot.* 587. *Fl. dan.* 1040. *Schkuhr handb.* 2, p. 256, t. 185.
Nat. of Britain. ♀.

12. *A. pendula*, foliis amplexicaulibus acuminatis, siliquis pendulis ancipitibus, bracteis foliaceis.

Arabis pendula. Willden. *sp. pl.* 3, p. 541. *Jacq. hort. vindob.* 3, p. 20, t. 34.
Nat. of Siberia. ☉.

13. *A. Turrita*, foliis amplexicaulibus, siliquis planis margine incrassatis, bracteis foliaceis.

Arabis Turrita. Willden. *sp. pl.* 3, p. 541. *Engl. bot.* 178.
Nat. of England. ♂.

14. *A. canadensis*, foliis caulinis lanceolatis subdentatis pubescentibus, racemis ebracteatis, siliquis pendulis falcatis.

Arabis canadensis. Willden. *sp. pl.* 3, p. 540. [108]
Arabis falcata. Michaux *amer.* 2, p. 31.
Nat. of North America. ♀.

MACROPODIUM.

Siliqua pedicellata linearis. *Cotyledones* accumbentes.
Calyx erectus.

1. MACROPODIUM nivale.

Cardamine nivalis. Willden. *sp. pl.* 3, p. 482. *Pallas it.* 2, *append. n.* 113, t. U.
Nat. of Siberia. ♀.

TURRITIS. *Gen. pl.* 1095.

Siliqua elongata anceps: valvis nervosis carinatisve.
Semina biseriata. *Cotyledones* accumbentes.

1. *T. arvensis*, glaberrima, calyce clauso, siliqua rostrata, foliis amplexicaulibus repandis; summis cordatis integerrimis.

Brassica arvensis. Willden. *sp. pl.* 3, p. 546.

Brassica sylvestris fabariæ foliis. Boccon. *sic.* 49, t. 25, f. 3, 4.

Nat. of the South of Europe. 4.

^{109]} 2. *T. glabra*, foliis radicalibus dentatis pilosis; caulinis amplexicaulibus integerrimis glabris.

Turritis glabra. Willden. *sp. pl.* 3, p. 542. *Curtis lond. Engl. bot.* 777. *Fl. dan.* 809.

Nat. of England. ☉.

3. *T. Loeselii*, foliis runcinatis dentatis pedunculisque hirtis, caule retrorsum hispido.

Sisymbrium Loeselii. Willden. *sp. pl.* 3, p. 503. *Jacqu. austr.* 4, p. 12, t. 324.

Nat. of Austria and Prussia. ☉.

BARBAREA.

Siliqua tetragono-anceps. *Cotyledones* accumbentes. *Semina* uniseriata. *Calyx* erectus. *Glandulæ* intra filamenta breviora.

1. *B. vulgaris*, foliis inferioribus lyratis: lobo terminali rotundato; superioribus obovatis dentatis.

Erysimum Barbarea. Willden. *sp. pl.* 3, p. 509. *Engl. bot.* 443. *Fl. dan.* 985. *Svensk bot.* 194.

Nat. of Britain. 4.

2. *B. præcox*, foliis inferioribus lyratis; superioribus pinnatifidis: laciniis lineari-oblongis integerrimis.

Erysimum præcox. Willden. *sp. pl.* 3, p. 510. *Engl. bot.* 1129.

Nat. of England. 4.

NASTURTIIUM.

Siliqua teretiuscula (quandoque abbreviata): valvis concavis enervibus, nec carinatis. *Cotyledones* accumbentes. ^[110]
Calyx patens.

1. *N. officinale*, foliis pinnatis: foliolis ovatis subcordatis repandis.

Sisymbrium Nasturtium. *Willden. sp. pl.* 3, p. 489.
Curtis lond. Engl. bot. 855.

Nat. of Britain. 4.

2. *N. sylvestre*, foliis pinnatis: foliolis lanceolatis serratis incisive.

Sisymbrium sylvestre. *Willden. sp. pl.* 3, p. 489. *Curtis lond. Engl. bot.* 2324.

Nat. of Britain. 4

3. *N. terrestre*, foliis pinnatifidis dentatis glabris, radice fusiformi, petalis calycem æquantibus.

Sisymbrium terrestre. *Smith fl. brit.* 2, p. 701. *Curtis lond. Engl. bot.* 1747.

Sisymbrium palustre. *Willden. sp. pl.* 3, p. 490.

Nat. of Britain. ☉.

4. *N. amphibium*, foliis oblongo-lanceolatis pinnatifidis serratisve, petalis calyce longioribus, radice fibrosa.

Sisymbrium amphibium. *Smith fl. brit.* 2, p. 702.
Engl. bot. 1840. *Willden. sp. pl.* 3, p. 491.

Nat. of Britain. 4.

5. *N. pyrenaicum*, siliculis ovalibus, foliis inferioribus lyratis; superioribus pinnatis amplexicaulibus: foliolis linearibus.

Sisymbrium pyrenaicum. *Willden. sp. pl.* 3, p. 491. ^[111]

Sisymbrium foliis pinnatis: imorum pinnis oblongis, su-

periorum linearibus integris, siliquis ovalibus oligospermis.

La Chenal in act. helvet. 4, p. 291, t. 15.

Nat. of Switzerland and the Pyrenees. ۲.

6. *N. sagittatum*, pubescens, foliis oblongis dentatis: radicalibus hastatis; caulinis sagittatis amplexicaulibus.

Sisymbrium sagittatum. Willden. sp. pl. 3, p. 493.

Sisymbrium molle. Jacqu. ic. 1, t. 122.

Nat. of Siberia. ۲.

SISYMBRIUM. *Gen. pl.* 1089.

Siliqua teretiuscula v. angulata. Cotyledones incumbentes (quandoque obliquæ), planæ. Calyx patens, nunc erectiusculus.

1. *S. officinale*, siliquis rachi adpressis subulatis pubescentibus, foliis runcinatis pilosis, caule hispido.

Erysimum officinale. Willden. sp. pl. 3, p. 509. Curtis lond. Engl. bot. 725. Svensk bot. 158.

Nat. of Britain. ☉.

2. *S. Irio*, foliis runcinatis cauleque glabro, siliquis patenti-erectis.

Sisymbrium Irio. Willden. sp. pl. 3, p. 503. Curtis lond. Engl. bot. 1631.

Nat. of England. ☉.

^{112]} 3. *S. Columnæ*, foliis runcinatis dentatis cauleque viloso, siliquis erectis.

Sisymbrium Columnæ. Willden. sp. pl. 3, p. 503. Jacqu. austr. 4, p. 12, t. 323.

Nat. of Germany and Italy. ☉.

4. *S. orientale*, foliis runcinatis tomentosis, caule lævi.

Willden. sp. pl. 3, p. 504.

Nat. of the Levant. ☉.

5. *S. pannonicum*, foliis inferioribus runcinatis dentatis;

superioribus pinnatis : pinnis linearibus integerrimis, siliquis
rectangule patentibus. *Willden. sp. pl.* 3, p. 502.

Sisymbrium pannonicum. *Jacqu. ic.* 1, t. 123.

Nat. of Hungary. ☉.

6. *S. austriacum*, foliis inferioribus lrvato-sinuatis acute
dentatis cauleque glabro, siliquis adscendentibus.

Sisymbrium austriacum. *Jacqu. austr.* 3, p. 35, t. 262.

Nat. of Austria. ☉.

7. *S. sinapoides*, foliis glabris : radicalibus runcinatis ;
caulinis pinnatifidis, siliquis striatis erectis pilosiusculis
pedunculo laxo duplo longioribus.

Sinapis pyrenaica. *Willden. sp. pl.* 3, p. 556. *Jacqu.*
hort. vindob. 3, p. 50, t. 97.

Nat. of the Pyrenees. ♂.

[113]

8. *S. altissimum*, foliis runcinatis flaccidis : foliolis sub-
linearibus integerrimis, pedunculis laxis. *Willden. sp. pl.*
3, p. 501.

Erysimum monspessulanum ; sinapios folio. *Walth.*
hort. 135, t. 22.

Nat. of Siberia. ☉.

9. *S. asperum*, siliquis scabris, foliis pinnatifidis glabris :
laciniis linearibus subdentatis.

Sisymbrium asperum. *Willden. sp. pl.* 3, p. 499.

Sinapi monspessulanum, siliqua aspera hirsuta. *Bauh.*
hist. 2, p. 858.

Nat. of the South of France. ☉.

10. *S. supinum*, siliquis axillaribus sessilibus solitariis,
foliis sinuato-pinnatifidis, caule retrorsum pubescente.

Sisymbrium supinum. *Willden. sp. pl.* 3, p. 494.

Eruca supina alba, siliqua singulari e foliorum alis erum-
pente. *Isnard in act. paris.* 1724, p. 295, t. 18.

Nat. of the South of Europe. ☉.

11. *S. polyceratium*, siliquis axillaribus aggregatis subu-
latis sessilibus, foliis runcinato-sinuatis dentatis acutis.

114] *Sisymbrium polyceratum*. *Willden. sp. pl.* 3, p. 494.
Jacqu. hort. vindob. 1, p. 34, t. 79.

Nat. of France and Italy. ☉.

12. *S. tanacetifolium*, foliis pinnatis: foliolis lanceolatis incis: extimis confluentibus, petalis calyce majoribus.

Sisymbrium tanacetifolium. *Willden. sp. pl.* 3, p. 492.

Eruca tanacetifolia. *Zanon. hist.* 103, t. 72.

Nat. of Italy. ♀.

13. *S. Sophia*, foliis decomposito-pinnatis: pinnulis linearibus: terminali longiore, petalis calyce minoribus.

Sisymbrium Sophia. *Willden. sp. pl.* 3, p. 500. *Engl. bot.* 963. *Svensk bot.* 203.

Nat. of Britain. ☉.

14. *S. millefolium*, foliis supradecompositis tomentosis incanis, petalis calyce majoribus.

Sisymbrium millefolium. *Willden. sp. pl.* 3, p. 500.

Sinapis millefolia. *Jacqu. ic.* 1, t. 127.

Nat. of the Canary Islands. Mr. Fr. Masson. ♀.

15. *S. strictissimum*, foliis lanceolatis petiolatis dentatis pubescentibus, siliquis linearibus.

115] *Sisymbrium strictissimum*. *Willden. sp. pl.* 3, p. 506.
Jacqu. austr. 2, p. 56, t. 194.

Nat. of Switzerland and Italy. ♀.

ERYSIMUM. *Gen. pl.* 1090.

Siliqua tetragona. *Semina* immarginata. *Cotyledones* incumbentes. *Stigma* capitatum, nunc emarginatum lobulis patentibus. *Calyx* clausus.

1. *E. cheiranthoides*, foliis lanceolatis integerrimis nunc paucidentatis: pilis stellato-tripartitis, siliquis erectiusculis: pedunculis patulis; stigmate indiviso subsessili.

Erysimum cheiranthoides. Willden. *sp. pl.* 3, p. 311.
Engl. bot. 942.
Nat. of Britain. ☉.

2. *E. repandum*, foliis lanceolatis repando-dentatis: pilis bipartitis, siliquis patulis torulosis; stigmatē emarginato.
Erysimum repandum. Willden. *sp. pl.* 3, p. 510. *Jacqu. austr.* 1, p. 16, t. 22.
Nat. of Spain and Austria. ☉.

3. *E. diffusum*, foliis lanceolato-linearibus integerrimis v. paucidentatis: pilis bipartitis, unguibus calyce longioribus; laminis obovato-oblongis, siliquis erectiusculis elongatis; stigmatē bilobo; stylo brevissimo.
Cheiranthus alpinus. Linn. *mant.* 93. *Jacqu. austr.* 1, p. 48, t. 75.
Erysimum diffusum. Willden. *sp. pl.* 3, p. 512. (116)
Nat. of the South of Europe. ♀.

4. *E. angustifolium*, foliis linearilanceolatis integris, siliquis erectis brevissime pedunculatis stylo duplo longioribus.
Erysimum angustifolium. Willden. *sp. pl.* 3, p. 513. •
Waldst. et Kitaib. hung. 1, p. 101, t. 98.
Nat. of Hungary. ☉.

5. *E. sessiliflorum*, foliis linearibus integerrimis, siliquis sessilibus stylo paulo longioribus.
Cheiranthus quadrangulus. Willden. *sp. pl.* 3, p. 525.
L'Herit. stirp. 1, p. 91, t. 44.
Nat. of Siberia. ♀.

6. *E. lanceolatum*, foliis lanceolatis: inferioribus dentatis, unguibus calyce longioribus; laminis orbiculato-obovatis, stigmatē sessili.
Cheiranthus erysimoides. Willden. *sp. pl.* 3, p. 514.
Jacqu. austr. 1, p. 48, t. 74.
Nat. of the South of Europe. ♂.

7. *E. helveticum*, foliis lanceolatis integerrimis: pilis

plerisque bipartitis, siliquis erectiusculis; stigmatē pedicellato emarginato.

Cheiranthus helveticus. *Willden. sp. pl.* 3, p. 515. *Jacqu. hort. vindob.* 3, p. 9, t. 9.

¹¹⁷ *Nat. of Switzerland.* ♂.

8. *E. odoratum*, foliis lanceolatis dentatis: pube tripartita, laminis orbiculato-obovatis, siliquis erectis, stigmatē bilobo.

Erysimum odoratum. *Willden. sp. pl.* 3, p. 512.

Erysimum hieracifolium. *Jacqu. austr.* 1, p. 47, t. 73.

Nat. of Austria. ☉.

9. *E. Alliaria*, foliis cordatis petiolatis dentato-incisis.

Erysimum Alliaria. *Willden. sp. pl.* 3, p. 510. *Curtis lond. Engl. bot.* 796. *Fl. dan.* 935. *Svensk bot.* 208.

Nat. of Britain. ♀.

10. *E. orientale*, foliis cordatis amplexicaulibus glabris; radicalibus scabris integerrimis.

Brassica orientalis. *Willden. sp. pl.* 3, p. 545. *Engl. bot.* 1804.

Nat. of England. ☉.

NOTOCERAS.

Siliqua valvis apice cornigeris. *Cotyledones* accumbentes. *Stigma* capitatum. *Calyx* erectiusculus basi æqualis.

1. *N. canariensis*, siliquis tetragonis bicornibus.

Erysimum bicornis. *Willden. sp. pl.* 3, p. 514.

¹¹⁸ *Nat. of the Canary Islands.* *Mr. Fr. Masson.* ☉.

CHEIRANTHUS. *Gen. pl.* 1091.

Siliqua compressa v. anceps. *Cotyledones* accumbentes. *Calyx* clausus, foliolis oppositis basi saccatis. *Stigma* stylo insidens, bilobum, lobis patentibus v. capitatum.

1. *C. Cheiri*, foliis lanceolatis integerrimis: pube (dum adsit) bipartita appressa, siliquis linearibus; stigmatibus lobis recurvis.

Cheiranthus cheiri. Willden. *sp. pl.* 3, p. 516.

α. flore simplici.

β. flore pleno.

Nat. of the South of Europe. ♂. ♀.

2. *C. mutabilis*, foliis lanceolatis argure serratis canescentibus: pube stellari, siliquis linearibus, caule fruticoso.

Cheiranthus mutabilis. Willden. *sp. pl.* 3, p. 517. *Curtis magaz.* 195.

Cheiranthus longifolius. Venten. *malmais.* 83.

Nat. of Madeira. Mr. Francis Masson. ♀.

3. *C. tenuifolius*, foliis filiformi-linearibus integerrimis incanis: pube bipartita adpressa, caule fruticoso.

Cheiranthus tenuifolius. Willden. *sp. pl.* 3, p. 517.

Nat. of Madeira. Mr. Francis Masson. ♀.

MATHIOLA.

[119]

Siliqua (teres vel compressa) coronata *Stigmate* conniventi bilobo, lobis vel dorso incrassatis cotyledonibus tunc incumbenibus; vel basi cuspidate stipatis. *Calyx* clausus. *Filamenta* longiora dilatata.

* *Stigmatibus lobis dorso incrassatis, subsimplicibusve.*

1. *M. incana*, foliis lanceolatis integerrimis, siliquis cylindricis eglandulosis.

Cheiranthus incanus. Willden. *sp. pl.* 3, p. 520. *Engl. bot.* 1935.

α. *Leucojum incanum majus*. *Bauh. pin.* 200.

β. *Cheiranthus coccineus*, foliis lanceolatis undatis, caule erecto indiviso. *Mill. dict.*

γ. *Cheiranthus albus*, foliis lanceolatis integerrimis obtusis incanis, ramis floriferis axillaribus, caule suffruticoso. *Mill. dict.*

3. *Cheiranthus annuus*. *Willden. sp. pl.* 3, p. 520.

ε. *Cheiranthus glabrus*, foliis lanceolatis acutis petiolatis viridibus, caule suffruticoso. *Mill. dict.*

Nat. of England. 2.

2. *M. fenestralis*, foliis confertis obovatis tomentosis, caule indiviso.

Cheiranthus fenestralis. *Willden. sp. pl.* 3, p. 520. *Jacqu. hort. vindob.* 2, p. 84, t. 179.

Nat. ♂.

^{120]} 3. *M. sinuata*, foliis tomentosis: inferioribus sinuatis, siliquis compressis muricatis.

Cheiranthus sinuatus. *Willden. sp. pl.* 3, p. 524. *Engl. bot.* 462.

Nat. of England. ♂.

4. *M. odoratissima*, foliis tomentosis lyrato-sinuatis, siliquis compressis tomentosis lævibus.

Cheiranthus odoratissimus. *Willden. sp. pl.* 3, p. 524.

Nat. of the North of Persia. 2.

5. *M. tristis*, foliis linearibus integerrimis subsinuatisve, floribus sessilibus.

Cheiranthus tristis. *Willden. sp. pl.* 3, p. 522. *Botan. magaz.* 729.

Nat. of the South of Europe. 2.

**** *Siliquis tricuspidatis*.**

6. *M. tricuspidata*, foliis sinuatis, siliquæ cuspidibus acutis subæqualibus.

Cheiranthus tricuspidatus. *Willden. sp. pl.* 3, p. 523. *Schkuhr handb.* 2, p. 250, t. 184.

Nat. of Barbary. ○.

^{121]} 7. *M. parviflora*, foliis lanceolatis repando-dentatis, siliquis sessilibus: cuspidibus lateralibus acutis medio obtuso longioribus.

Cheiranthus parviflorus. Willden. *sp. pl.* 3, p. 519.
Nat. of Morocco. ☉.

MALCOMIA.

Siliqua teres, bivalvis. Stigma acutum. Cotyledones
incumbentes planæ. Calyx clausus.

1. *M. maritima*, foliis obovatis lanceolatisve integerrimis
 v. obsolete dentatis: pube adpressa 2—4-partita, siliquis
 pedunculatis, calycibus deciduis.

Cheiranthus maritimus. Willden. *sp. pl.* 3, p. 518.
Curtis magaz. 166.

Nat. of the South of Europe. ☉.

2. *M. africana*, foliis lanceolatis dentatis: pube stellari
 laxa, siliquis sessilibus, calyce persistenti.

Hesperis africana. Willden. *sp. pl.* 3, p. 532.

Leucojum gallicum, folio halimi. *Bocc. sic.* 77, t. 42,
 f. 1.

Nat. of Africa. ☉.

3. *M. littorea*, foliis lanceolatis integris vel dentatis sili-
 quisque tomentosis incanis.

Cheiranthus littoreus. Willden. *sp. pl.* 3, p. 521.

Leucojum maritimum minus. *Clus. hist.* 1, p. 298.

Nat. of the South of Europe. ☉.

HESPERIS. *Gen. pl.* 1093.

[122]

Siliqua 4-gona v. anceps. Stigma subsessile, lobis con-
niventibus. Cotyledones incumbentes, planæ. Calyx
clausus.

1. *H. tristis*, petalorum laminis oblongis obliquis, siliquis
 divaricatis acipitibus margine incrassatis.

Hesperis tristis. Willden. *sp. pl.* 3, p. 530. *Botan.*
magaz. 730.

Nat. of Austria and Hungary. ♂.

2. *H. matronalis*, petalorum laminis obovatis, siliquis erectis torosis margine simplicibus.

Hesperis matronalis. *Willden. sp. pl.* 3, p. 531.

Hesperis inodora. *Willden. sp. pl.* 3, p. 531. *Engl. bot.* 731.

a. flore simplici.

flore pleno albo.

β. flore pleno purpureo.

Nat. of England. 4.

ERUCARIA. *Gärtn. sem.* 2, p. 298.

Siliqua biarticulata: articulo inferiore bivalvi polyspermo; superiore evalvi oligospermo. *Cotyledones* planæ (rectæ v. curvatae).

1. *ERUCARIA aleppica*.

Erucaria aleppica. *Venten. cels.* 64.

Cordyllocarpus lævigatus. *Willden. sp. pl.* 3, p. 563.

^{123]} *Bunias myagroides*. *Willden. sp. pl.* 3, p. 414.

Nat. of the Levant. ☉.

BRASSICA. *Gen. pl.* 1096.

Siliqua bivalvis (rostro sterili v. monospermo). *Cotyledones* conduplicatae. *Calyx* clausus.

1. *B. Napus*, radice caulescente fusiformi, foliis lævibus: superioribus cordato-lanceolatis amplexicaulibus; inferioribus lyratis dentatis. *Willden. sp. pl.* 3, p. 547. *Engl. bot.* 2146.

Nat. of Britain. ♂.

2. *B. Rapa*, radice caulescente orbiculari depressa carnosa, foliis radicalibus lyratis scabris; caulinis integerrimis lævibus. *Willden. sp. pl.* 3, p. 548. *Engl. bot.* 2176.

Nat. of England. ♂.

3. *B. oleracea*, radice caulescente tereti carnosa, foliis

omnibus glæbris glaucis repandis lobatisve. *Willden. sp. pl. 3, p. 548. Engl. bot. 637. Select. specim. 4, 5.*

a. capitata. Brassica capitata alba. *Bauh. pin. 111.*

β. rubra. Brassica capitata rubræ. *Bauh. pin. 111.*

γ. sabauda. Brassica alba crispa. *Bauh. pin. 111.*

δ. sabellica. Brassica fimbriata. *Bauh. pin. 112.*

ε. botrytis. Brassica cauliflora. *Bauh. pin. 111.*

ζ. Brassica italica broccoli dicta. *Mill. dict.*

η. Napobrassica. Brassica radice napiformi. *Tourn. [124 inst. 219.*

Nat. of the English sea-coasts. ♂.

4. *B. chinensis*, foliis ovalibus subintegerrimis; floralibus amplexicaulibus lanceolatis, calycibus ungue petalorum longioribus. *Willden. sp. pl. 3, p. 550.*

Nat. of China. ☉.

5. *B. Erucastrum*, foliis runcinatis, caule hispido, siliquis lævibus. *Willden. sp. pl. 3, p. 551. Bulliard herb. de la France, 331.*

Nat. of the South of Europe. ☉.

6. *B. Eruca*, foliis lyratis, caule hirsuto, siliquis glabris: rostro compresso articulato.

Brassica Eruca. *Willden. sp. pl. 3, p. 551. Bulliard herb. de la France, 313. Schkuhr handb. 2, p. 263, t. 186.*

Nat. of Switzerland and Austria. ☉.

7. *B. monensis*, foliis pinnatifidis, caule nudiusculo glabro, siliquis lævibus: rostro monospermo.

Brassica monensis. *Huds. angl. 291.*

Sisymbrium monense. *Willden. sp. pl. 3, p. 496. Engl. bot. 962.*

Nat. of Britain. ☉.

8. *B. elongata*, foliis petiolatis: inferioribus sinuato-^[125] pinnatifidis hispidis; superioribus glabris dentatis, siliquis torulosis tetragonis cuspidatis. *Willden. sp. pl. 3, p. 552.*

Brassica elongata. *Waldst. et Kitaib. hung.* 1, p. 26, t. 28.

Nat. of Hungary. ♂.

9. *B. campestris*, radice cauleque tenui, foliis cordatis acuminatis amplexicaulibus: inferioribus lyratis dentatis subhispidis. *Willden. sp. pl.* 3, p. 546. *Engl. bot.* 2234.

Nat. of England. ☉.

10. *B. alpina*, foliis caulinis cordato-sagittatis amplexicaulibus; radicalibus ovatis, petalis erectis. *Willden. sp. pl.* 3, p. 547. *Villars dauph.* 3, p. 330, t. 36.

Nat. of Germany and Switzerland. ♀.

SINAPIS. *Gen. pl.* 1097.

Siliqua bivalvis (nunc biarticulata articulo superiore evalvi). *Cotyledones* conduplicatæ. *Calyx* patens.

1. *S. arvensis*, siliquis multangulis toroso-turgidis rostro ancipiti longioribus. *Willden. sp. pl.* 3, p. 554. *Curtis lond.* *Engl. bot.* 1748.

Nat. of Britain. ☉.

2. *S. orientalis*, siliquis retrorsum hispidis apice subtragonis compressis. *Willden. sp. pl.* 3, p. 554. *Schkuhr handb.* 1, p. 264, t. 186.

^{126]} *Nat. of the Levant.* ☉.

3. *S. alba*, siliquis hispidis: rostro obliquo longissimo ensiformi. *Willden. sp. pl.* 3, p. 555. *Curtis lond.* *Engl. bot.* 1677.

Nat. of Britain. ☉.

4. *S. Allioni*, siliquis ovato-oblongis patulis, foliis pinnatifidis laciniis incisis. *Willden. sp. pl.* 3, p. 557. *Jacqu. hort. vindob.* 2, p. 79, t. 168.

Raphanus crucoides. *Linn. suppl.* 299.

Nat. ☉.

5. *S. chinensis*, siliquis lævibus subarticulatis patulis, foliis lyrato-runcinatis subhirsutis. *Willden. sp. pl.* 3, p. 557.

Sinapis siliquis glabris subarticulatis, &c. *Arduin. spec.* 1, p. 23, t. 10.

Nat. of China. ☉.

6. *S. juncea*, ramis fasciculatis, foliis summis lanceolatis integerrimis. *Willden. sp. pl.* 3, p. 557. *Jacqu. hort. vindob.* 2, p. 80, t. 171.

Nat. of China. ☉.

7. *S. lævigata*, siliquis lævibus patulis, foliis lyratis glabris: summis lanceolatis, caule lævi. *Willden. sp. pl.* 3, p. 559. [127]

Nat. of Spain and Portugal. ♂.

8. *S. nigra*, siliquis adpressis glabris tetragonis: stylo subulato.

Sinapis nigra. *Willden. sp. pl.* 3, p. 555. *Engl. bot.* 969. *Svensk bot.* 83.

Nat. of Britain. ☉.

9. *S. pubescens*, siliquis pubescentibus erectis: rostro compresso, foliis lyratis villosis. *Willden. sp. pl.* 3, p. 556.

Sinapis hirsuta caulibus frutescentibus, &c. *Arduin. spec.* 1, p. 21, t. 9.

Nat. of Sicily. ♀.

10. *S. eruroides*, siliquis lævibus æqualibus, foliis lyratis oblongis glabris, caule glabro. *Willden. sp. pl.* 3, p. 557. *Jacqu. hort. vindob.* 2, p. 80, t. 170.

Nat. of the South of Europe. ☉.

11. *S. frutescens*, siliquis linearibus lævibus, foliis inferioribus oblongis dentatis; superioribus lanceolatis integris, caule glabro frutescente. *Willden. sp. pl.* 3, p. 559.

Nat. of Madeira. *Mr. Fr. Masson.* ♀.

12. *S. tenuifolia*, siliquis linearibus glabris brevissime [128]

rostratis erectis; pedunculis patulis, foliis lanceolatis acutis-
simis pinnatifidis bipinnatifidisve, caule glabro.

Sisymbrium tenuifolium. Willden. *sp. pl.* 3, p. 493.
Engl. bot. 525.

Brassica muralis. Curtis *lond.*

Nat. of England. 4.

13. *S. muralis*, siliquis linearibus glabris brevissime
rostratis erectiusculis; pedunculis patulis, foliis sinuatis
glabris, caule patulo piloso.

Sisymbrium murale. Willden. *sp. pl.* 3, p. 496. *Engl.*
bot. 1090.

Nat. of England. ☉.

14. *S. incana*, siliquis biarticulatis adpressis, foliis sca-
bris: inferioribus lyratis; summis lanceolatis.

Sinapis incana. Willden. *sp. pl.* 3, p. 558. *Jacqu. hort.*
vindob. 2, p. 79, t. 169.

Myagrum hispanicum. Willden. *sp. pl.* 3, p. 407.

Nat. of France, Spain, and Portugal. ♂.

15. *S. hispanica*, siliquis biarticulatis erectis, foliis
bipinnatifidis: laciniis linearibus.

Sinapis hispanica. Willden. *sp. pl.* 3, p. 558.

Nat. of Spain. ☉.

120]

RAPHANUS. *Gen. pl.* 1098.

Siliqua evalvis. *Cotyledones* conduplicatæ. *Calyx* clausus.

1. *R. sativus*, foliis lyratis, siliquis teretibus torosis bilo-
cularibus. Willden. *sp. pl.* 3, p. 560.

α. *Raphanus minor* oblongus. *Bauh. pin.* 96.

β. *Raphanus major* orbicularis vel rotundus. *Bauh.*
pin. 96.

γ. *Raphanus niger.* *Bauh. pin.* 96.

Nat. ☉.

2. *R. Raphanistrum*, foliis simpliciter lyratis, siliquis
unilocularibus articulatis striatis.

Raphanus Raphanistrum. Willden. *sp. pl.* 3, p. 560.
Curtis lond. Engl. bot. 856.
Nat. of Britain. ☉.

3. *R. [maritimus,]* foliis interrupte lyratis, siliquis unilocularibus articulatis striatis.

Raphanus maritimus. *Engl. bot.* 1643.
Nat. of Britain. 4.

CHORISPERMUM.

Silica evalvis bilocularis, in segmenta monosperma clausa discedens. *Cotyledones* planæ, accumbentes. *Stigma* simplex.

1. *C. tenellum*, siliquis foliisque glabris; superioribus lanceolatis dentatis; infimis pinnatifidis.

Raphanus tenellus. Willden. *sp. pl.* 3, p. 561. *Pallas* (180 it. 3, p. 741, t. L. f. 3.
Nat. of Siberia. ☉.

CLEOME. *Gen. pl.* 1099.

Calyx 4-phyllus. *Petala* 4. *Stam.* 4—40. *Silica* unilocularis, bivalvis, polysperma. *Semina* exalbuminosa.

1. *C. Chelidonii*, polyandra: filamentis apice latioribus, foliis 5—9-natis: foliolis utrinque scabris pilis adpressis, racemis terminalibus.

Cleome Chelidonii. Willden. *sp. pl.* 3, p. 567.
Nat. of the East Indies. ☉.

2. *C. viscosa*, staminibus 15—20, foliis quinatis ternatisque, siliquis oblique striatis, seminibus striis transversis elevatis.

Cleome viscosa. Willden. *sp. pl.* 3, p. 566.

Sinapistrum zeylanicum triphyllum et pentaphyllum viscosum flore flavo. *Martyn dec.* 25.

Nat. of Ceylon. ☉.

3. *C. pentaphylla*, floribus gynandris : staminibus medio stipitis insertis, foliis quinatis passim floralibusque ternatis, caule inermi.

Cleome pentaphylla. *Willden. sp. pl.* 3, p. 564. *Jacqu. hort. vindob.* 1, p. 9, t. 24.

Nat. of both Indies. ☉.

^{131]} 4. *C. gigantea*, inermis hexandra : staminibus juxta basin stipitis longissimi insertis, foliis 7—9-natis hirsutis, racemo ebracteato.

Cleome gigantea. *Willden. sp. pl.* 3, p. 567. *Jacqu. obs.* 4, p. 1, t. 76.

Cleome viridiflora. *Schreb. nov. act. ac. nat. cur.* 4, p. 136, t. 3.

Nat. of South America. ♀.

5. *C. spinosa*, aculeata hexandra : staminibus juxta basin stipitis germine aliquoties longioris insertis, foliis 7—5-natisque acuminatis, bracteis racemi simplicibus cordatis.

Cleome spinosa. *Willden. sp. pl.* 3, p. 568 (excluso synonymo *Milleri*).

Cleome Erucago. *Mill. dict.*

Nat. of the West Indies. ☉.

6. *C. Houstoni*, aculeata glanduloso-pubescens hexandra : staminibus basi stipitis germine brevioris insertis, foliis quinatis ternatisque ; floralibus simplicibus, siliquis breviter pedicellatis, stigmatibus dilatato.

Sinapistrum indicum spinosum, flore carneo folio trifido vel quinquefido. *Martyn dec.* 45.

Nat. of the West Indies. ☉.

^{132]} 7. *C. violacea*, hexandra, calyce petalisque inæqualibus, foliis ternatis lanceolato-linearibus ; floralibus omnibus simplicibus, siliquis teretiusculis, seminibus glabriusculis.

Cleome violacea. *Willden. sp. pl.* 3, p. 569. *Schkuhr handb.* 2, p. 292, t. 189, b.

Nat. of Portugal. ☉.

8. *C. Ornithopodioides*, hexandra, foliis ternatis oblongo-

medio
ternatis,
Jacqu.
lanceolatis obtusis; floralibus plerisque simplicibus, siliquis
pendulis sessilibus torulosis, seminibus glabris.

Cleome ornithopodioides. Willden. sp. pl. 3, p. 568.

Sinapistrum orientale triphyllum, ornithopodii siliquis.
Dill. elth. 359, t. 266, f. 345.

Nat. of the Levant. ☉.

9. *C. arabica*, hexandra, foliis ternatis lineari-lanceolatis,
siliquis pendulis, seminibus hirsutissimis.

Cleome arabica. Willden. sp. pl. 3, p. 569. Linn. fl.
fascic. t. 8.

Siliquaria glandulosa. Forsk. descr. 78, ic. t. 16, B.

Nat. of Arabia. ☉.

10. *C. monophylla*, hexandra, foliis simplicibus lanceolatis
pubescentibus: basi obtusa.

Cleome monophylla. Willden. sp. pl. 3, p. 571.

Sinapistrum zeylanicum viscosum, folio solitario flore
flavo siliqua tenui. Burm. zeyl. 217, t. 100, f. 2.

Nat. of the East Indies. ☉.

[133

11. *C. procumbens*, hexandra glaberrima, foliis simplicibus
lanceolatis, caule diffuso.

Cleome procumbens. Willden. sp. pl. 3, p. 571. Jacqu.
amer. 189, t. 120.

Nat. of the West Indies. 4.

medio
ternatis,
Jacqu.

juxta
irsutis,
Jacqu.

4, p.

basin
7—5-
latis.
excluso

andra:
foliis
reviter

trifido

alibus,
unibus
culis.
chkuhr

longo-

GENERA ET SPECIES QUÆDAM

PLANTARUM LEGUMINOSARUM,

QUÆ

IN HORTO KEWENSI COLUNTUR.

[*Extracted from 'Hortus Kewensis,' 2nd Edition, Vol. III, pp. 1—22 ;
Vol. IV, pp. 266—338 ; and Vol. V, pp. 460—468.*]

LONDON.
—
1811-1812-1813.

DECANDRIA.

[iii. 1]

MONOGYNIA.

EDWARDSIA.

Salisb. in linn. soc. transact. 9, p. 298. Brown mss.

*Cal. 5-dentatus. Cor. papilionacea. Legumen tetra-
pterum, polyspermum.*

1. *E. grandiflora*, foliolis (13—19) lanceolato-oblongis.
*Edwardsia grandiflora. Salisb. in linn. soc. transact. 9,
p. 299.*

*Sophora tetraptera. Willden. sp. pl. 2, p. 499. Curtis
magaz. 167.*

*Nat. of New Zealand. The Right Hon. Sir Joseph
Banks, Bart. K.B. h.*

2. *E. microphylla*, foliolis (25—41) obovatis.
*Edwardsia microphylla. Salisb. in linn. soc. transact.
9, p. 299.*

*Sophora microphylla. Willden. sp. pl. 2, p. 499. Jacqu.
hort. schænbr. 3, p. 10, t. 269.*

*Nat. of New Zealand. The Right Hon. Sir Joseph
Banks, Bart. K.B. h.*

SOPHORA. *Gen. pl. 694. Brown mss.* [iii. 2]

*Cal. 5-dentatus. Cor. papilionacea. Legumen monili-
forme, apterum, polyspermum.*

1. *S. tomentosa*, arborea, foliolis subrotundis ovalibusve obtusissimis utrinque calycibusque tomentosis, racemo terminali elongato. *Brown mss.*

Sophora tomentosa. *Willden. sp. pl. 2, p. 500.*

Sophora occidentalis. *Willden. sp. pl. 2, p. 500.* *Trew ehret. 27, t. 59.*

Nat. of both Indies. γ .

2. *S. japonica*, arborea, foliolis oblongo-ovatis acutis leguminibusque glabris. *Brown mss.*

Sophora japonica. *Willden. sp. pl. 2, p. 500.* *Andrews's reposit. 585.*

Nat. of Japan and China. γ .

3. *S. alopecuroides*, herbacea, foliolis oblongis: adultis subtus sericeis. *Brown mss.*

Sophora alopecuroides. *Willden. sp. pl. 2, p. 499.*

Pallas astragal. 117, t. 87.

Nat. of the Levant. γ .

4. *S. flavescens*, herbacea, foliolis ovato-oblongis glabriusculis. *Brown mss.*

Sophora flavescens. *Willden. sp. pl. 2, p. 499.*

Nat. of Siberia. γ .

iii. 3]

ORMOSIA.

Jackson in linn. soc. transact. 10, p. 360. *Brown mss.*

Cal. 5-fidus, bilabiatus. Cor. papilionacea. Stigmata 2, approximata, obtusa: altero unilaterali! Legumen compressum, lignosum, 1—3-spermum.

1. *O. dasycarpa*, foliolis (9—11) acuminatis utrinque glabris, leguminibus tomentosis. *Brown mss.*

Ormosia dasycarpa. *Jackson in linn. soc. transact. 10, p. 362, t. 26.*

Sophora monosperma. *Willden. sp. pl. 2, p. 501.*

Nat. of the West Indies. γ .

ANAGYRIS. *Gen. pl.* 695.

Cal. 5-dent. bilabiatus. *Cor.* papilionacea, *carina* dipetala
alis longior *vexillum* superantibus. *Legumen* compressum,
 polyspermum.

1. *A. fœtida*, *Willden. sp. pl.* 2, 507. *Clus. hist.* 1, p. 93.
Nat. of Spain and Italy. $\frac{1}{2}$.

THERMOPSIS. *Brown mss.*

Cal. oblongus, semi-5-fidus, bilabiatus, postice convexus,
 basi attenuata. *Cor.* papilionacea, petalis longitudine subæ-
 qualibus: *vexillum* lateribus reflexis; *carina* obtusa. *Sta-*
mina persistentia. *Legumen* compressum, lineare, poly-
 spermum.

1. *T. lanceolata*, foliolis oblongo-lanceolatis, stipulis
 lanceolatis petiolo duplo longioribus, pedicellis verti-
 cillatis. *Brown mss.*

Podalyria lupinoides. *Willden. sp. pl.* 2, p. 504. [iii. 4]

Sophora lupinoides. *Pallas astragal.* 119, t. 89.

Nat. of Siberia. $\frac{1}{4}$.

VIRGILIA.

Lamarck illustr. t. 326. *Pers. synops.* 1, p. 453.
Brown mss.

Cal. 5-fidus. *Cor.* papilionacea, petalis longitudine subæ-
 qualibus: *vexillum* lateribus non reflexis. *Stigma* imberbe.
Legumen compressum, oblongum, polyspermum.

1. *V. aurea*, staminibus persistentibus, germinibus to-
 mentosis, foliolis ovalibus obtusis muticis. *Brown mss.*

Podalyria aurea. *Willden. sp. pl.* 2, p. 502.

Robinia subdecandra. *L'Herit. stirp. nov.* 1, p. 157, t. 75.

Nat. of Abyssinia. $\frac{1}{2}$.

2. *V. intrusa*, staminibus persistentibus, germinibus glabris, calycis basi intrusa, foliolis ovalibus obtusis mucronulatis. *Brown mss.*

Nat. of the Cape of Good Hope. Mr. Fr. Masson. h.

3. *V. capensis*, staminibus deciduis basi lanatis, germinibus tomentosis, carina acuminata, foliolis lanceolatis. *Brown mss.*

Podalyria capensis. Willden. sp. pl. 2, p. 501. Andrews's reposit. 347.

Nat. of the Cape of Good Hope. h.

iii. 6]

CYCLOPIA.

Venten. dec. gen. nov. p. 8. Brown mss. Ibettsonia. Sims in botan. magaz. 1259.

Cal. 5-fidus, inæqualis, basi intrusa. Cor. papilionacea: vexillum rugis longitudinalibus! alæ plica transversa! Stamina decidua. Stigma hinc barbatum. Legumen compressum, polyspermum.

1. *C. genistoides*, foliolis subulatis laciniisque calycis muticis, bracteis oblongo-ovatis pedunculo brevioribus, ramulis glabris. *Brown mss.*

Ibettsonia genistoides. Botan. magaz. 1259.

Gompholobium maculatum. Andrews's reposit. 427.

Podalyria genistoides. Willden. sp. pl. 2, p. 502.

Nat. of the Cape of Good Hope. h.

BAPTISIA.

Venten. dec. gen. nov. p. 9. Brown mss. Podalyria. Michaux amer. 1, p. 263. Podalyriæ sp. Lamarck illustr. t. 327, f. 1. Willden. sp. pl. 2, p. 501.

Cal. semi-4—5-fidus, bilabiatus. Cor. papilionacea, petalis longitudine subæqualibus: vexillum lateribus reflexis. Stamina decidua. Legumen ventricosum, pedicellatum, polyspermum.

1. *B. perfoliata*, foliis perfoliatis integerrimis subrotundis. *Brown mss.*

Rafnia perfoliata. *Willden. sp. pl.* 3, p. 949.

Crotalaria Perfoliatæ folio. *Dill. elth.* 122, t. 102, f. 122.

Nat. of Carolina. 2.

[iii. 6]

2. *B. australis*, foliis ternatis petiolatis: foliolis cuneato-lanceolatis, stipulis petiolo longioribus lanceolatis. *Brown mss.*

Podalyria australis. *Willden. sp. pl.* 2, p. 503. *Venten. Cels.* 56.

Sophora australis. *Botan. magaz.* 509.

Sophora cœrulea. *Trew pl. rar.* 6, t. 14.

Nat. of Carolina. 2.

3. *B. tinctoria*, foliis ternatis petiolatis: foliolis subrotundo-obovatis, stipulis setaceis obsoletis. *Brown mss.*

Podalyria tinctoria. *Willden. sp. pl.* 2, p. 503. *Botan. magaz.* 1099.

Nat. of North America. 2.

4. *B. alba*, foliis ternatis petiolatis: foliolis elliptico-oblongis, stipulis deciduis subulatis petiolo brevioribus, germinibus glabris. *Brown mss.*

Podalyria alba. *Willden. sp. pl.* 2, p. 503. *Botan. magaz.* 1177.

Nat. of North America. 2.

PODALYRIA.

Salisb. parad. 7. *Brown mss.* *Podalyriæ* sp. *Lamarck illustr. t.* 327, f. 3, 4.

Cal. 5-fidus, inæqualis, basi intrusa. *Car.* papilionacea, vexillo majore. *Stamina* persistentia, basi connata. [iii. 7] *Legumen* ventricosum, polyspermum.

1. *P. myrtillifolia*, foliis oblongo-obovatis utrinque calycibusque sericeis, pedunculis unifloris folia subæquantibus. *Brown mss.*

Podalyria myrtillifolia. Willden. *sp. pl.* 2, p. 505.
Nat. of the Cape of Good Hope. ʒ.

2. *P. sericea*, foliis oblongo-obovatis utrinque calyceque sericeis pedunculo unifloro aliquoties longioribus. *Brown mss.*

Sophora sericea. Andrews's *reposit.* 440.
Nat. of the Cape of Good Hope. Mr. David Nelson. ʒ.

3. *P. biflora*, foliis ovalibus utrinque sericeis pedunculo bifloro brevioribus, calycibus tomentosis scabris. *Brown mss.*

Podalyria biflora. Willden. *sp. pl.* 2, p. 505. *Botan. magaz.* 753.

Podalyria argentea. Salisb. *parad.* 7.
Nat. of the Cape of Good Hope. Mr. Fr. Masson. ʒ.

4. *P. calyptrata*, foliis ovalibus obovatisque pubescentibus: subtus reticulatis; adultis non sericeis, pedunculis unifloris folia subæquantibus, calycibus villosis: limbo scarioso reflexo. *Brown mss.*

Podalyria calyptrata. Willden. *sp. pl.* 2, p. 504.
Nat. of the Cape of Good Hope. ʒ.

iii. 8] 5. *P. hirsuta*, foliis villosis petiolatis: superioribus ovatis; inferioribus subrotundis, pedunculis unifloris, calycibus villosis: laciniis longitudine alarum. *Brown mss.*

Podalyria hirsuta. Willden. *sp. pl.* 2, p. 505.
Nat. of the Cape of Good Hope. Mr. Fr. Masson. ʒ.

6. *P. cordata*, foliis cordatis subrotundis subsessilibus villosissimis, pedunculis oifloris, calycis villosi laciniis brevioribus alis. *Brown mss.*

Sophora cordata. Thunb. *prod.* 79?
Nat. of the Cape of Good Hope. Mr. Fr. Masson. ʒ.

CHORIZEMA.

Labillard. voy. 1, p. 405. Brown mss.

Cal. semi-5-fidus, bilabiatus. *Cor.* papilionacea, *carina* ventricosa *alis* brevior. *Stylus* brevis, uncinatus. *Stigma* obliquum, obtusum. *Legumen* ventricosum, polyspermum.

1. *C. ilicifolia*, foliis pinnatifido-dentatis spinosis oblongo-lanceolatis; acumine integro dentibus longiore, bracteis ad apicem pedicelli. *Brown mss.*

Chorizema ilicifolia. Labillard. voy. 1, p. 405, t. 21, nov. holl. 2, p. 120.

Chorozema ilicifolium. Smith in linn. soc. transact. 9, p. 251.

Nat. of the South-west coast of New Holland. h.

2. *C. nana*, foliis sinuato-dentatis spinosis oblongis obtusis, bracteis infra apicem pedicelli. *Brown mss.*

Chorizema nana. Sims in botan. magaz. 1032.

Pultenæa nana. Andrews's reposit. 434.

Nat. of the South-west coast of New Holland. (iii.) Robert Brown. h.

3. *C. rhombea*, foliis integerrimis planis mucronatis: inferioribus rhombeo-orbiculatis; summis elliptico-lanceolatis, pedunculis paucifloris. *Brown mss.*

Nat. of the South-west coast of New Holland. Robert Brown. h.

PCDOLOBIUM. *Brown mss.*

Cal. 5-fidus, bilabiatus. *Cor.* papilionacea, *carina* compressa longitudine *alarum* subæquantium *vexillum* explanatum. *Germen* simplici serie 4-spermum. *Stylus* ad-

scendens. *Stigma simplex*. *Legumen* pedicellatum, lineari-oblongum, modice ventricosum, intus læve.

1. *P. trilobatum*, foliis oppositis spinoso-dentatis trilobis: basi transversa; lobis lateralibus terminali dentato multoties brevioribus, germine sericeo. *Brown mss.*

Chorozema trilobatum. *Smith in linn soc. transact.* 9, p. 253.

Pultenaea ilicifolia. *Andrews's reposit.* 320.

Nat. of New South Wales. Mr. *David Burton*. ♀.

OXYLOBIUM.

Andrews's reposit. 492. *Brown mss.*

Cal. profundè 5-fidus, subbilabiatus. *Cor.* papilionacea, *carina* compressa longitudine *alarum* æquantium *vevillum* explanatum. *Stylus* adscendens. *Stigma simplex*. *Legumen* polyspermum, ventricosum, ovatum, acutum.

iii. 10] 1. *O. arborescens*, foliis lineari-lanceolatis, bracteis apicis pedicelli persistentibus, corymbis confertis, leguminibus calyce vix longioribus.

Nat. of Van Diemen's Island. *Robert Brown*. ♀.

2. *O. ellipticum*, foliis ovali-oblongis, bracteis infra apicem pedicelli caducis, corymbis confertis, leguminibus calyce duplo longioribus. *Brown mss.*

Gompholobium ellipticum. *Labillard. nov. holl.* 1, p. 107, t. 135.

Calistachys elliptica. *Venten. malmais.* 115, b. *Smith in linn. soc. transact.* 9, p. 266.

Nat. of Van Diemen's Island. Mr. *David Nelson*. ♀.

3. *O. cordifolium*, foliis ovatis cordatis pilosiusculis, umbellis terminalibus sessilibus. *Brown mss.*

Oxylobium cordifolium. *Andrews's reposit.* 492.

Nat. of New South Wales. The Right Hon. Sir *Joseph Banks*, Bart. K.B. ♀.

BRACHYSEMA. *Brown mss.*

Cal. 5-fidus, parum inæqualis, tubo ventricoso. *Cor.* papilionacea, *vexillum* brevius *carina* compressa *alas* æquante. *Germen* pedicello basi vaginula cincto. *Stylus* filiformis, elongatus. *Legumen* polyspermum, ventricosum.

1. *B. latifolium*, foliis ovatis planis, vexillo oblongo-ovato. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* h. (iii. 11)

GOMPHOLOBIUM.

Smith in linn. soc. transact. 4, p. 220.

Cal. 5-partitus, subæqualis. *Cor.* papilionacea, *vexillo* explanato. *Stigma* simplex. *Legumen* polyspermum, sub-sphæricum, obtusissimum (glabrum).

1. *C. latifolium*, foliis ternatis: foliolis linearibus oblongo-linearibus (uncialibus et ultra), caule erecto, *carina* fimbriata, calyce fructus reflexo. *Brown mss.*

Gompholobium latifolium. *Smith in linn. soc. transact.* 9, p. 249.

Gompholobium fimbriatum. *Smith exot. bot.* 1, p. 113, t. 58.

Gompholobium psoraliaefolium. *Salisb. parad.* 6.

Nat. of New South Wales. Colonel *William Paterson.* h.

2. *G. marginatum*, foliis ternatis: foliolis obovatis marginatis plenis, stipulis petiolum æquantibus, corolla longitudine calycis. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* h.

3. *G. polymorphum*, foliis ternatis quinatisve: foliolis linearibus margine recurvis: apice quandoque dilatato, caule procumbente v. volubili. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* h.

iii. 12] 4. *G. tomentosum*, foliis pinnatis: foliolis subulato-linearibus mucronatis supra scabris, pedunculis subsolitariis, calycibus hirsutis legumine brevioribus, carina sericeo-ciliata. *Brown mss.*

Gompholobium tomentosum. *Labill. nov. holl.* 1, p. 106, t. 134.

Nat. of the South-west coast of New Holland. h.

5. *G. venustum*, foliis pinnatis multijugis: foliolis subulatis venosis margine revolutis calycibusque glabris, corymbo pedunculato multifloro, corollis purpureis. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* h.

BURTONIA. *Brown mss.*

Cal. profunde 5-fidus. *Cor.* papilionacea, decidua, petalis longitudine subæqualibus. *Germen* dispermum. *Stylus* bulatus, basi dilatata. *Stigma* obtusum, imberbe. *Legumen* subrotundum, modice ventricosum. *Strophiola* seminis nulla.

1. *B. scabra*, foliis ternatis, calycibus glabris, stylo extra medium imberbi. *Brown mss.*

Gompholobium scabrum. *Smith in linn. soc. transact.* 9, p. 250.

Nat. of the South-west coast of New Holland. *Archibald Menzies, Esq.* h.

JACKSONIA. *Brown mss.*

Cal. 5-partitus, subæqualis. *Cor.* papilionacea, staminaque decidua. *Germen* dispermum. *Stylus* subulato-iii. 13] filiformis. *Stigma simplex.* *Legumen* modice ventricosum, ovatum v. oblongum, valvis intus pubescentibus. *Strophiola* seminis nulla.

1. *J. scoparia*, arborescens inermis, ramulis angulatis, racemis terminalibus. *Brown mss.*

Nat. of New South Wales. Robert Brown. h.

2. *J. spinosa*, fruticosa, ramulis spinescentibus di-trichotomis patulis angulatis, bracteis apicis pedicelli appressis brevissimis. *Brown mss.*

Gompholobium spinosum. Labill. nov. holl. 1, p. 107, t. 136.

Nat. of the South-west coast of New Holland. h.

VIMINARIA.

Smith in annals of bot. 1, p. 507. Brown mss.

Cal. 5-dentatus, angulatus. Cor. papilionacea. Stylus capillaris, germine dispermo aliquoties longior. Stigma simplex. Legumen evalue, ovatum. Strophiola seminis nulla.

1. *V. denudata, Smith exot. bot. t. 27. Linn. soc. transact. 9, p. 261.*

Daviesia denudata. Venten. choix t. 6.

Sophora juncea. Schrad. sert. hanov. t. 3.

Nat. of New Holland and Van Diemen's Island. h.

SPHÆROLOBIUM.

Smith in annals of bot. 1, p. 509. Brown mss.

Cal. 5-fidus, bilabiatus. Cor. papilionacea. Stylus [iii. 14] apice hinc membrana auctus, inde imberbis. Stigma terminale. Legumen sphæricum.

1. *S. vimineum*, calycis tubo labiis paulo brevior, stylo incluso a basi arcuato, corollis flavis. *Brown mss.*

Sphærolobium vimineum. Smith in linn. soc. transact. 9, p. 261. Botan. magaz. 969.

Sphærolobium minus. *Labillard. nov. holl.* 1, p. 108, t. 138.

Nat. of New Holland and Van Diemen's Island. $\frac{1}{2}$.

2. *S. medium*, calycis tubo labiis dimidio brevior, corollis rubris. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* $\frac{1}{2}$.

AOTUS.

Smith in annals of bot. 1, p. 504. *Brown mss.*

Cal. 5-fidus, bilabiatus. *Cor.* papilionacea. *Stamina* decidua. *Germen* dispermium. *Stylus* filiformis. *Legumen* bivalve. *Strophiola* seminis nulla.

1. *A. villosa*, calycibus sericeis villis appressis, legumine pedicellato, seminibus punctato-rugosis, foliis supra scabris. *Brown. mss.*

Aotus villosa. *Smith in linn. soc. transact.* 9, p. 249. *Botan. magaz.* 949.

Aotus ferruginea. *Labillard. nov. holl.* 1, p. 104, t. 132.

Pultenæa ericoides. *Venten. malmais.* 35.

Nat. of New Holland and Van Diemen's Island. The Right Hon. Sir *Joseph Banks*, Bart. K.B. $\frac{1}{2}$.

DILLWYNIA.

Smith in annals of bot. 1, p. 310. *Brown mss.*

Cal. 5-fidus, bilabiatus, basi attenuata. *Cor.* papilionacea, petalis staminibusque deciduis, medio tubo calycis insertis: *vevilli* lamina duplo et ultra latior quam longa, divaricato-biloba. *Germen* dispermium. *Stylus* uncinatus. *Stigma* capitatum. *Legumen* ventricosum. *Semina* stropholata.

1. *D. floribunda*, floribus axillaribus geminis, foliis subulatis mucronatis. *Brown mss.*

Dillwynia floribunda. *Smith exot. bot. t. 26. Linn. soc. transact. 9, p. 262.*

Nat. of New South Wales. 2.

2. *D. ericifolia*, corymbis terminalibus sessilibus, foliis (4-lin.) subulatis punctato-scabris divaricatis subtortis: mucrone pungenti, ramulis pubescentibus. *Brown mss.*

Dillwynia ericifolia. *Smith exot. bot. t. 25. Linn. soc. transact. 9, p. 262.*

Pultenæa retorta. *Wendl. hort. herrenhus. 2, p. 13, t. 9.*

Nat. of New South Wales. 2.

3. *D. glaberrima*, corymbis terminalibus pedunculatis, foliis filiformibus erectis lævibus: mucrone innocuo recurvo. *Brown mss.*

Dillwynia glaberrima. *Smith in linn. soc. transact. 9, p. 263. Botan. magaz. 944.*

Nat. of New South Wales. The Right Hon. Sir Joseph Banks, Bart. K. B. 2.

EUTAXIA. *Brown mss.*

[iii. 16]

Cal. bilabiatus. *Cor. papilionacea, vexilli lamina paulo latior quam longa. Germen dispermum. Stylus uncinatus. Stigma capitatum. Legumen modice ventricosum. Semina strophololata. Folia opposita.*

1. *E. myrtifolia*, foliis lanceolatis lanceolato-obovatisve, pedunculis axillaribus geminis, alarum appendicibus brevissimis. *Brown mss.*

Dillwynia myrtifolia. *Smith in linn. soc. transact. 9, p. 263.*

Dillwynia obovata. *Labillard. nov. holl. 1, p. 110, t. 140. Botan. magaz. 1274.*

Nat. of the South-west coast of New Holland. 2.

foliis

SCLEROTHAMNUS. *Brown mss.*

Cal. 5-fidus, bilabiatus, basi bibracteatus. *Cor.* papilionacea: *carina* longitudine alarum. *Germen* dispermum, pedicellatum. *Stylus* adscendens, filiformis. *Stigma* simplex. *Legumen* ventricosum.

1. *S. microphyllus*, *Brown mss.*

Nat. of the South coast of New Holland. *Robert Brown.* h.

GASTROLOBIUM. *Brown mss.*

Cal. 5-fidus, bilabiatus, ebracteatus. *Cor.* papilionacea, petalis longitudine subæqualibus. *Germen* dispermum, pedicellatum. *Stylus* subulatus, adscendens. *Stigma* simplex. *Legumen* ventricosum. *Semina* strophiolata.

1. *G. bilobum*, foliis (uncialibus) subtus subsericeis re-tusis: lobulis rotundatis mucronulo longioribus, leguminis
iii. 17] pedicello tubum calycis æquante. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* h.

EUCHILUS. *Brown mss.*

Cal. profunde 5-fidus, bilabiatus, labio superiore maximo; basi bibracteatus. *Cor.* papilionacea, *carina* longitudine alarum. *Germen* dispermum, pedicellatum. *Stylus* subulatus, adscendens. *Stigma* simplex. *Legumen* compressum. *Strophiola* seminis lobis posticis integris.

1. *E. obcordatus*, *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* h.

PULTENÆA.

Smith in annals of bot. 1, p. 502. *Brown mss.*

Cal. 5-fidus, labiis proportionatis, bibracteatus (bracteis sæpius ipso tubo insidentibus). *Cor.* papilionacea. *Ger-*
men sessile, dispermum. *Stylus* subulatus, adscendens.
Stigma simplex. *Strophiola* seminis lobis posticis incis.

1. *P. daphnoides*, capitulis terminalibus, foliis obovato-oblongis planis glaberrimis lævibus (uncialibus) triplo longioribus quam latis: mucrone pungenti. *Brown mss.*

Pultenæa daphnoides. *Smith in linn. soc. transact.* 9, p. 247. *Willden. sp. pl.* 2, p. 507. *Andrews's reposit.* 98. *Wendl. hort. herrenhus.* 3, p. 7, t. 17.

Nat. of New South Wales. The Right Hon. Sir Joseph Banks, Bart. K. B. h

2. *P. obcordata*, capitulis terminalibus, foliis cuneato-obcordatis retusis planis glaberrimis lævibus (semuncialibus) vix duplo longioribus quam latis: mucrone pungenti. *Brown mss.*

Pultenæa obcordata. *Andrews's reposit.* 574.

Nat. of Van Diemen's Island. *Robert Brown.* h.

3. *P. scabra*, capitulis terminalibus paucifloris, foliis cuneiformibus truncatis setacco-mucronatis margine recurvis: supra scabris; subtus villosis, stipulis setaceis recurvis. *Brown mss.*

Nat. of New South Wales. *Robert Brown.* h.

4. *P. retusa*, capitulis terminalibus, foliis linearibus retusis muticis planis glabris, bracteis calyce paulo longioribus. *Brown mss.*

Pultenæa retusa. *Smith in linn. soc. transact.* 9, p. 247.

Nat. of New South Wales. h.

5. *P. linophylla*, bracteis capitulo 6—8-floro brevioribus foliis linearibus (semuncialibus) mucronulo marginibusque

recurvis: subtus sericeis, stipulis petiolo brevioribus.
Brown mss.

Pultenæa linophylla. *Smith in linn. soc. transact.* 9, p. 247. *Schrad. sert. hannov.* 3, p. 28, t. 18. *Willden. sp. pl.* 2, p. 505.

iii. 19] *Nat. of New South Wales.* The Right Hon. Sir *Joseph Banks*, Bart. K. B. h.

6. *P. stipularis*, capitulis multifloris: bracteis calyces subæquantibus, foliis planis linearibus acutis: junioribus ciliatis, stipulis bifidis planis imbricatis. *Brown mss.*

Pultenæa stipularis. *Smith in linn. soc. transact.* 9, p. 245. *Smith nov. holl.* 1, p. 35, t. 12. *Botan. magaz.* 435. *Willden. sp. pl.* 2, p. 506.

Nat. of New South Wales. h.

7. *P. vestita*, floribus axillaribus, foliis lineari-lanceolatis mucronatis glabris: marginibus inflexis, stipulis imbricatis ciliatis, calycibus bracteisque aristatis. *Brown mss.*

Nat. of the South coast of New Holland. *Robert Brown.*

h.

8. *P. villosa*, racemis foliatis, foliis lineari-oblongis: supra concaviusculis; subtus et calycibus ramulisque pilosis. *Brown mss.*

Pultenæa villosa. *Willden. sp. pl.* 2, p. 507. *Smith in linn. soc. transact.* 9, p. 248. *Botan. magaz.* 967.

Nat. of New South Wales. h.

9. *P. flexilis*, glaberrima, floribus axillaribus, foliis oblongo-linearibus mucronatis planis. *Brown mss.*

Pultenæa flexilis. *Smith in linn. soc. transact.* 9, p. 248.

iii. 20] *Nat. of New South Wales.* Colonel *William Paterson.*

h.

DAVIESIA.

Smith in linn. soc. transact. 4, p. 222. Brown mss.

Cal. angulatus, ebracteatus. *Cor.* papilionacea, *carina* vexillo brevior. *Germen* pedicellatum, dispermum. *Stylus* strictus. *Stigma* simplex. *Legumen* compressum, angulatum, elastice dehiscens. *Strophiola* seminis postice integra.

1. *D. ulicina*, ramis spinescentibus glabris patulis, foliis lanceolatis linearibusve margine lævibus: mucrone spinoso, pedunculis axillaribus unifloris: bracteis (8) imbricatis. *Brown mss.*

Daviesia ulicina. Smith in linn. soc. transact. 9, p. 256.

Daviesia ulicifolia. Andrews's reposit. 304.

Nat. of New South Wales. h.

2. *D. mimosoides*, ramis inermibus, foliis elongato-lanceolatis: mucrone brevissimo innocuo, corymbis axillaribus geminis solitariisve: bracteis adpressis, calycis labio superiore retuso. *Brown mss.*

Daviesia corymbosa. Andrews's reposit. 526. (excluso synonymo.)

Nat. of New South Wales. Robert Brown. h.

3. *D. latifolia*, ramis inermibus, foliis ellipticis ovalibusve venosis basi subattenuatis, racemis axillaribus multifloris. *Brown mss.*

Nat. of Van Diemen's Island. Robert Brown. h. [iii. 21]

MIRBELIA.

Smith in annals of bot. 1, p. 541. Brown mss.

Cal. 5-fidus, bilabiatus. *Cor.* papilionacea. *Legumen* biloculare, sutura utraque inflexa.

1. *M. reticulata*, foliis lanceolato-linearibus venosis, geminibus dispermis. *Brown mss.*

Mirbelia reticulata. *Smith in linn. soc. transact.* 9, p. 265. *Venten. malmais.* 119.

Pultenæa rubiæfolia. *Andrews's reposit.* 351.

Nat. of New South Wales. Mr. *David Burton.* 2.

2. *M. dilatata*, foliis cuneiformibus : apice dilatato trifido. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* 2.

CERCIS. *Gen. pl.* 696. *Brown mss.*

Cal. 5-dentatus. *Corolla* papilionacea. *Legumen* compressum, sutura seminifera alata. *Semina* obovata, embryone recto.

1. *C. Siliquastrum*, foliis orbiculatis cordatis. *Willden. sp. pl.* 2, p. 507. *Botan. magaz.* 1138. *Schmidt arb.* 1, p. 21, t. 20.

Nat. of the South of Europe and the Levant. 2.

iii. 2. *C. canadensis*, foliis cordatis acuminatis. *Willden. sp. pl.* 2, p. 508. *Schmidt arb.* 1, p. 22, t. 21.

Nat. of North America. 2.

DIADELPHIA DECANDRIA.

iv. 266] PLATYLOBIUM. (*Smith in linn. soc. transact.* 9, p. 302.) *Brown mss.*

Cal. bracteatus, bilabiatus : labio superiore bifido rotundato maximo. *Stam.* omnia connexa. *Legumen* pedicellatum, compresso-planum, dorso alatum, polyspermum.

1. *P. formosum*, foliis ovatis subcordatis, germine unidique villosa, bracteis sericeis, pedicello leguminis calyce brevior. *Brown mss.*

Platylobium formosum. *Willden. sp. pl.* 3, p. 921.

Smith new holl. 17, t. 6. *Botan. magaz.* 469. *Venten. malmais.* 31.

Nat. of New South Wales. h.

2. *P. parviflorum*, foliis lanceolato-ovatis, germine ciliato glabriusculo, bracteis glabris, pedicello leguminis calycem superante. *Brown mss.*

Platylobium parviflorum. *Willden. sp. pl.* 3, p. 921.

Nat. of New South Wales. h.

3. *P. triangulare*, foliis deltoideis subhastatisve: angulis spinosis, pedunculis basi apiceque bracteatis medio nudis, legumine calycem aliquoties superante. *Brown mss.*

Nat. of Van Diemen's Island. *Robert Brown.* h.

BOSSIÆA. (*Venten. cels.* 7.) *Brown mss.*

Cal. bilabiatus: labio superiore majore semibifido obtuso. *Stam.* omnia connexa. *Legumen* plano-^(iv. 267) compressum, pedicellatum, polyspermum, margine utroque incrassatum. *Semina* strophiolata.

1. *B. Scolopendrium*, ramis complanatis linearibus aphyllis: denticulis floriferis, caule erecto, carina nuda, bracteis superioribus persistentibus imbricatis pedunculum æquantibus, calyce glaberrimo. *Brown mss.*

Bossiæa scolopendria. *Smith in linn. soc. transact.* 9, p. 303.

Platylobium scolopendrium. *Andrews's reposit.* 191.

Platylobium scolopendrium. *Venten. malmais.* 55.

Nat. of New South Wales. h.

2. *B. rufa*, ramis complanatis linearibus aphyllis: denticulis floriferis, carina fimbriata, bracteis superioribus caducis ab inferioribus remotis, calycibus glaberrimis. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* h.

3. *B. heterophylla*, ramis foliosis compressis, foliis obo-

vatis linearibusque planis, legumine multiloculari: septis spongiosis. *Brown mss.*

Bossiaea heterophylla. *Smith in linn. soc. transact.* 9, p. 302. *Willden. sp. pl.* 3, p. 972. *Venten. cels.* 7.

Bossiaea lanceolata. *Botan. magaz.* 1144.

Platylobium lanceolatum. *Andrews's reposit.* 205.

Platylobium ovatum. *Andrews's reposit.* 276.

Nat. of New South Wales. h.

[iv. 268] 4. *B. linophylla*, ramis foliosis compressis, foliis linearibus: marginibus recurvis, legumine uniloculari. *Brown mss.*

Nat. of the south-west coast of New Holland. *Robert Brown.* h.

5. *B. prostrata*, ramis foliosis filiformibus, caule procumbente, foliis ovalibus glabris, stipulis petiolo brevioribus, legumine uniloculari. *Brown mss.*

Nat. of New South Wales. *Robert Brown.* h.

6. *B. cinerea*, ramis foliosis teretibus, caule erecto ramosissimo, foliis ovato-lanceolatis: supra scabris; subtus pubescentibus; marginibus recurvis. *Brown mss.*

Nat. of Van Diemen's Island. *Robert Brown.* h.

7. *B. microphylla*, ramis foliosis spinescentibus teretibus, foliis obcordato-cuneiformibus. *Brown mss.*

Bossiaea microphylla. *Smith in linn. soc. transact.* 9, p. 303.

Platylobium microphyllum. *Sims in botan. magaz.* 863.

Nat. of New South Wales. *Mr. George Caley.* h.

SCOTTIA. *Brown mss.*

Cal. bracteis imbricatus, 5-dentatus dentibus parum
[ii. 269] inæqualibus. *Vexillum* complicatum brevius Alis
æquantibus Carinam. *Stam.* omnia connexa. *Legumen*
pedicellatum, compressum; margine utroque incrassato.
Semina (3-4) strophiolata.

1. *SCOTTIA dentata*. *Brown mss.*

Nat. of the south-west coast of New Holland. *Robert Brown.* ♀.

TEMPLETONIA. *Brown mss.*

Cal. ebracteatus, 5-dentatus, dentibus parum inæqualibus. *Carina* oblonga. *Stam.* omnia connexa, antheris uniformibus. *Legumen* pedicellatum, plano-compressum, polyspermum. *Semina* strophiolata.

1. *TEMPLETONIA retusa*. *Brown mss.*

Rafnia retusa. *Venten. malmais.* 53.

Nat. of the south-west coast of New Holland. *Robert Brown.* ♀.

GOODIA. (*Salisb. paradis.* 41.) *Brown mss.*

Cal. bilabiatus labiis longitudine subæqualibus; superiore semibifido acuto. *Vexillum* explanatum majus. *Stam.* omnia connexa. *Legumen* pedicellatum, compressum, subdispermum. *Semina* strophiolata.

1. *G. lotifolia*, foliolis obovatis calycibusque glaberrimis, legumine varicoso. *Brown mss.*

Goodia lotifolia. *Salisb. paradis.* 41. *Botan. magaz.* 958.

Nat. of Van Diemen's Island. Mr. *David Nelson.* ♀.

2. *G. pubescens*, foliolis obovato-cuneatis calycibusque pubescentibus, legumine lævi. *Brown mss.* [iv. 270]

Goodia pubescens. *Sims. in botan. magaz.* 1310.

Nat. of Van Diemen's Island. *Robert Brown.* ♀.

HOVEA. *Brown mss.*

[iv. 275]

Cal. bilabiatus: labio superiore semibifido retuso. *Stam.* omnia connexa. *Carina* obtusa. *Legumen* sessile subrotundum ventricosum dispermum. *Semina* strophiolata.

1. *H. linearis*, foliis linearibus subtus pilosiusculis, leguminibus glabris. *Brown mss.*

Poiretia linearis. *Smith in linn. soc. transact.* 9, p. 304.
Nat. of New South Wales. h.

2. *H. longifolia*, foliis elongato-linearibus: subtus venosis leguminibusque tomentosis. *Brown mss.*

Nat. of New South Wales. *Robert Brown.* h.

iv. 299]

KENNEDIA. *Venten. malmais.* 104.

Vexillum recurvum a carina non reflexum. *Legumen* multiloculare, polyspermum. *Semina* strophiolata.

1. *K. rubicunda*, foliis ternatis: foliolis ovatis, pedunculis subtrifloris, leguminibus hirsutissimis. *Venten. malmais.* 104.

Glycine rubicunda. *Willden. sp. pl.* 3, p. 1065. *Curtis magaz.* 268. *Schneevoogt ic.* 28.

Nat. of New South Wales. The Right Hon. Sir *Joseph Banks*, Bart. K. B. h.

2. *K. coccinea*, foliis ternatis: foliolis obovatis, floribus capitatis, leguminibus glabriusculis. *Venten. malmais.* 105, excluso synonymo *Curtisii*.

Nat. of the South-West Coast of New Holland. *Robert Brown.* h.

3. *K. prostrata*, foliis ternatis: foliolis obovatis villosis, pedunculis 1-2-floris, carina alas oblongas superante, caule prostrato. *Brown mss.*

Glycine coccinea. *Willden. sp. pl.* 3, p. 1065. *Curtis magaz.* 270. h.

4. *K. monophylla*, foliis simplicibus glabris reticulatis: basi subcordata, floribus racemosis. *Brown mss.*

iv. 300] *Kennedia monophylla.* *Venten. malmais.* 106.

Glycine bimaculata. *Willden. sp. pl.* 3, p. 1067. *Curtis magaz.* 263. *Schneevoogt ic.* 29.

Nat. of New South Wales. The Right Hon. Sir Joseph Banks, Bart. K. B. h.

COLUTEA. (*Gen. pl.* 1196.) *Brown mss.* [iv. 325]

Cal. 5-dentatus. *Vexillum* bicallosum, explanatum, majus Carina obtusa. *Stigma* laterale sub apice uncinato Styli postice longitudinaliter barbati. *Legumen* inflatum scariosum. [iv. 326]

1. *C. arborescens*, foliolis ellipticis retusis, vexilli gibbis abbreviatis, caule fruticoso. *Willden. sp. pl.* 3, p. 1139. *Curtis magaz.* 81. *Schmidt arb. t.* 117.

Nat. of France and Italy. h.

2. *C. cruenta*, foliolis obovatis emarginatis glaucis, vexilli gibbis obtusis minimis, leguminibus apice apertis, caule fruticoso. *Willden. sp. pl.* 3, p. 1139.

Colutea aperta. Schmidt arb. t. 119.

Nat. of the Levant. h.

3. *C. Pocockii*, foliolis subrotundo-ellipticis obtusissimis mucronatis, vexilli gibbis elongatis adscendentibus, caule fruticoso. *Willden. sp. pl.* 3, p. 1140. *Schmidt arb. t.* 120.

Nat. of the Levant. h.

SWAINSONA. (*Salisb. parad.* 28.) *Brown mss.*

Loxidium. Venten dec. gen. nov.

Cal. 5-dentatus. *Vexillum* explanatum majus Carina obtusa. *Stigma* terminale. *Stylus* postice longitudinaliter barbatus; antice imberbis. *Legumen* turgidum (non vesicarium).

1. *S. galegifolia*, vexillo bicalloso, caule suffruticoso erecto, pedicello leguminis filamentis persistentibus longiore. *Brown mss.* [iv. 327]

Colutea galegifolia. *Sims in botan. magaz.* 792.

Vicia galegifolia. *Andrews's reposit.* 319.

Nat. of New South Wales. Colonel *William Paterson.*

h.

2. *S. coronillifolia*, vexillo bicalloso, caule suffruticoso erecto, pedicello leguminis filamentis persistentibus (parum) brevior. *Brown mss.*

Swainsona coronillæfolia. *Salisb. paradis.* 28.

Nat. of New South Wales. Colonel *William Paterson.*

h.

SUTHERLANDIA. *Brown mss.*

Cal. 5-dentatus. *Vexillum* ccallosum, lateribus replicatis, brevius *Carina* oblonga. *Stigma* terminale. *Stylus* postice barba longitudinali; antice transversa apicis. *Legumen* inflatum scariosum.

1. *SUTHERLANDIA frutescens.*

Colutea frutescens. *Willden. sp. pl.* 3, p. 1140. *Curtis magaz.* 181.

Nat. of the Cape of Good Hope. h.

LESSERTIA. (*Decand. astrag.*) *Brown mss.*

Cal. semi-5-fidus. *Vexillum* explanatum. *Carina* obtusa. iv. 328] *Stigma* capitatum. *Stylus* antice barba transversa apicis; postice imberbis. *Legumen* scariosum, evalve (compressum v. inflatum).

1. *L. annua*, calycibus bibracteatis nigro-pilosis foliolis linearibus supra glabris. *Brown mss.*

Lessertia annua. *Decand. astrag.* 38.

Colutea herbacea. *Willden. sp. pl.* 3, p. 1141.

Colutea africana annua foliolis parvis mucronatis, vesiculis compressis. *Commel. hort.* 2, p. 87, t. 44.

Nat. of the Cape of Good Hope. ☉.

2. *L. diffusa*, foliolis linearibus emarginatis utrinque cauleque diffuso hirsutis, calycibus ebracteatis nigro-pilosis. *Brown mss.*

Galega dubia. Jacqu. ic. 3, t. 576.

Nat. of the Cape of Good Hope. Mr. Fr. Masson. ☉.

3. *L. perennans*, calycibus ebracteatis, foliolis ovalibus: subtus sericeis; supra pubescentibus. *Brown mss.*

Lessertia perennans. Decand. astrag. 37.

Colutea perennans. Willden. sp. pl. 3, p. 1142. Jacqu. hort. vindob. 3, p. 5, t. 3. Murray in nov. comm. gotting. 5, p. 38, t. 7, f. 1.

Nat. of the Cape of Good Hope. ♀.

SESBANA.

(iv. 330)

Brown mss. Sesban. Poiret. in encycl. 7, p. 126.

Cal. 5-dentatus. Legumen elongatum (teres v. lineare), bivalve, septis transversis multiloculare.

1. *S. grandiflora*, racemis subtrifloris, foliolis oblongis emarginatis, calycis labiis distantibus; dentibus coherentibus, caule arboreo.

Coronilla grandiflora. Willden. sp. pl. 3, p. 1145.

Agaty. Rheed. mal. 1, p. 95, t. 51.

Nat. of the East Indies. ♂.

2. *S. ægyptiaca*, racemis multifloris, foliolis linear-oblongis obtusis mucronulatis, rachi foliorum lævi.

Sesban ægyptiacus. Poiret in encycl. 7, p. 128.

Coronilla Sesban. Willden. sp. pl. 3, p. 1147.

Sesban. Alpin. ægypt. 81, t. 82.

Nat. of Egypt. ♂.

3. *S. aculeata*, racemis paucifloris, foliolis linearibus obtusis mucronulatis, rachi foliorum aculeata.

Coronilla aculeata. Willden. sp. pl. 3, p. 1147.

Æschynomene bispinosa. *Jacqu. ic. 3, t. 564.*
Nat. of the East Indies. ☉.

iv. 336] **SMITHIA.** (*Gen. pl. 1760*). *Brown mss.*

Stamina divisa in duas phalanges æquales. *Legumen* articulatum, plicatum, inclusum Calyce bifido.

1. *S. sensitiva.* *Willden. sp. pl. 3, p. 1161.* *Salisb. paradis. 92.*

Nat. of the East Indies. *John Gerard Kænig, M.D.* ♂.

ÆSCHYNOMENE. (*Gen. pl. ed. 1, p. 350.* *Palis. fl. d'Oware 1, p. 88.*) *Brown mss.*

Stamina divisa in duas phalanges æquales. *Legumen* articulatum strictum exsertum. *Calyx* bipartitus, labiis dentatis.

1. *Æ. sensitiva*, caule fruticoso lævi, foliolis linearibus obtusis, racemis paucifloris, lomentis lævibus. *Willden. sp. pl. 3, p. 1162.*

Hedysarum arborescens, foliis mimosæ. *Plum. ic. 140, t. 149.*

Nat. of the West Indies. ♀.

iv. 337] 2. *Æ. aspera*, caule herbaceo inferne scabro, foliolis linearibus obtusis, racemis compositis hispidis, lomenti articulis medio scabris. *Willden. sp. pl. 3, p. 1163.*

Mimosa non spinosa major zeylanica. *Breyn. cent. 51, t. 52.*

Nat. of the East Indies. ☉.

3. *Æ. hispida*, caule herbaceo hispido, foliolis linearibus obtusis, racemis simplicibus, lomentis hispidis. *Willden. sp. pl. 3, p. 1163.*

Nat. of North America. ☉.

4. *Æ. americana*, caule herbaceo hispido, foliolis lineari-falcatis acuminatis, racemis simplicibus, lomenti articulis

subrotundis distinctis glabris. *Willden. sp. pl.* 3, p. 1163.

Æschynomene caule hispido, foliolis acuminatis, legumini articulis suborbiculatis. *Hebenstreit in nov. comm. petropol.* 8, p. 321, tab. 12.

Nat. of Jamaica. ☉.

5. *Æ. indica*, caule herbaceo lævi, leguminibus lævibus hinc torosis obtusis, foliolis obtusis. *Willden. sp. pl.* 3, p. 1164.

Neli Tali. *Rheed mal.* 9, p. 31, t. 18.

Nat. of the East Indies. ☉.

[iv. 338]

POLYGAMIA MONŒCIA.

ACACIA.

[v. 460]

3. *A. acicularis*, foliis tereti-subulatis mucronatis sparsis rigidis, stipulis deciduis, ramulis glabris, spicis globosis slitariis. *Brown mss.*

Nat. of New South Wales. Colonel *William Paterson.* ♀.

4. *A. sulcata*, foliis filiformibus undique sulcatis: mucronulo innocuo, capitulis subgeminis; bracteis basis pedunculi ovatis concavis persistentibus, leguminibus flexuosis. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* ♀.

12. *A. melanoxyton*, foliis lanceolato-oblongis nervosis subfalcatis, capitulis subracemosis, ramulis ultimis pedunculisque angulatis: furfure tenuissimo tectis, funiculo umbilicali colorato plicato semen subcingente. *Brown mss.*

Nat. of Van Diemen's Island. *Robert Brown.* ♀.

13. *A. Sophoræ*, foliis oblongis æquilateris nervosis, spicis geminis sessilibus, corollis tetrapetalis, leguminibus torosis, funiculo umbilicali plicato. *Brown mss.*

Mimosa Sophoræ. *Labillard. nov. holl.* 2, p. 87, t. 237.
Nat. of Van Diemen's Island. *h.*

14. *A. marginata*, foliis elongato-lanceolatis subfalcatis marginatis uninerviis: margine antico parum exciso unglanduloso, capitulis racemosis sub-4-floris. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* *h.*

v. 463] 17. *A. decipiens*, foliis triangularibus passimque trapezoideis: angulo exteriori spinoso; interiore glandulifero, stipulis setaceis caducis, ramulis glabris, capitulis solitariis 7—10-floris. *Brown mss.*

Mimosa decipiens. *König in annals of bot.* 1, p. 366, t. 8.

Adiantum truncatum. *Linn. syst. veg. ed.* 13, p. 790.

Nat. of the South-west coast of New Holland. *Archibald Menzies, Esq.* *h.*

18. *A. biflora*, foliis triangularibus: angulo exteriori spinoso; interiore glandulifero, stipulis setaceo-spinosis persistentibus, ramulis pubescentibus, capitulis bifloris. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* *h.*

19. *A. armata*, foliis dimidiato-oblongis glabris mucronulatis uninerviis: nervo parallelo approximato margini v. 464] interiori subtruncato, stipulis spinosis, capitulis globosis solitariis, ramis hirsutis. *Brown mss.*

Nat. of the South coast of New Holland. *Robert Brown.* *h.*

20. *A. alata*, caule bifariam alato, foliis decurrentibus uninerviis spinula terminatis: margine interiore dente unico glandulifero, stipulis spinosis, capitulis pedunculatis sub-solitariis. *Brown mss.*

Nat. of the South-west coast of New Holland. *Robert Brown.* *h.*

22. *A. pulchella*, foliis conjugato-pinnatis: glandula

pedicellata inter pinnas 5—7-jugas, stipulis spinosis folia subæquantibus, capitulis solitariis, ramis flexuosis. *Brown mss.*

Nat. of the South-west coast of New Holland. Robert Brown. h.

23. *A. ciliata*, inermis pilosa, foliis bipinnatis: partialibus bijugis: propriis 2—3-jugis, stipulis subsetaceis caducis, capitulis solitariis. *Brown mss.*

Nat. of the South-west coast of New Holland. Robert Brown. h.

24. *A. nigricans*, inermis glabra, foliis bipinnatis: partialibus bijugis: superioris propriis 5—7-jugis; inferioris 2—3-jugis, stipulis subulato-setaceis, capitulis solitariis. *Brown mss.*

Mimosa nigricans. Labillard. nov. holl. 2, p. 88, t. 238.

Nat. of the South-west coast of New Holland. h.

33. *A. pubescens*, inermis, foliis bipinnatis: partialibus suboctojugis; propriis subquindecimjugis, racemis axillari-bus solitariis: capitulis globosis pedicellatis, ramis hirsutis, petiolis eglandulosis.

Mimosa pubescens. Venten. malmais. 21. Botan. magaz. 1263.

Nat. of New South Wales.

[v. 468]

GENERA ET SPECIES QUÆDAM

PLANTARUM MYRTACEARUM,

QUÆ

IN HORTO KEWENSI COLUNTUR.

[*Extracted from 'Hortus Kewensis,' 2nd Edition, Vol. IV,*
pp. 410—419.]

LONDON.

1812.

POLYADELPHIA ICOSANDRIA.

[416]

MELALEUCA. (*Smith in linn. soc. transact.* 3, p. 273.)
Brown mss.

Staminum phalanges 5, petalis oppositæ, elongatæ: *Antheræ* incumbentes. *Caps.* 3-loc. polysperma, connata et inclusa calycis tubo incrassato basi adnato (ramo).

Folia alterna.

1. *M. Leucadendron*, foliis elongato-lanceolatis acuminatis falcatis aversis 3—5-nervibus, ramulis floriferis pendulis spicisque glaberrimis. *Brown mss.*

Melaleuca Leucadendron. *Willden. sp. pl.* 3, p. 1428.

Arbor alba Caju Puti. *Rumph. amb.* 2, p. 72, t. 16.

Nat. of the East Indies. ½.

2. *M. viridiflora*, foliis elliptico-lanceolatis æquilateris 5-nervibus, ramulis et rachi spicæ calycisque tubo pubescentibus, phalangium unguibus calycis laciniis brevioribus obsoletisve. *Brown mss.*

Melaleuca viridiflora. *Willden. sp. pl.* 3, p. 1429.

Metrosideros quinquenervia. *Cavanill. ic.* 4, p. 19, t. 333.

Nat. of New South Wales. *John White, M.D.* ½.

3. *M. paludosa*, foliis lineari-lanceolatis elongatis æquilateris strictis 3-nervibus: nervis lateralibus margini scabro approximatis, rachi spicæ cylindricæ calycisque tubo sericeis, phalangium unguibus brevissimis. *Brown mss.*

Nat. of the South-west Coast of New Holland. *Robert Brown.* ♀.

4. *M. globifera*, foliis oblongis 5-nerviis æquilatis basi attenuatis, capitulis sphaericis, capsulis connatis. *Brown mss.*

Nat. of the South Coast of New Holland. *Robert Brown.* ♀.

5. *M. diosmifolia*, foliis ovalibus oblongisve obsolete univerviis petiolatis planis confertis ramulisque glaberrimis, spicis oblongis glabris, phalangibus 3—5-andris. *Brown mss.*

Melaleuca diosmæfolia. *Andrews's reposit.* 476.

Nat. of the South-west Coast of New Holland. *Archibald Menzies, Esq.* ♀.

6. *M. stypheloides*, foliis ovatis acuminatis mucronatopungentibus striato-multinerviis sessilibus glabris, spicis pubescentibus, calycis laciniis acutis nervosis. *Brown mss.*

Melaleuca stypheloides. *Willden. sp. pl.* 3, p. 1430.

Nat. of New South Wales. *Mr. David Burton.* ♀.

7. *M. genistifolia*, foliis lineari-lanceolatis obsolete 1—3-⁴²nerviis, spicis laxis foliatis glabris, phalangibus polyandris: unguibus petala subæquantibus. *Brown mss.*

Melaleuca genistifolia. *Willden. sp. pl.* 3, p. 1431. *Smith exot. bot.* 1, p. 107, t. 55.

Nat. of New South Wales. *Mr. David Burton.* ♀.

8. *M. striata*, foliis lanceolato-linearibus acutis punctatis obsolete striatis rigidis subsessilibus, spicis ovalibus oblongisve, calycis tubo lanato, phalangibus subdecandris: unguibus petala bis superantibus. *Brown mss.*

Melaleuca striata. *Labillard. nov. holl.* 2, p. 26, t. 165.

Nat. of the South Coast of New Holland. *Robert Brown.* ♀.

9. *M. thymoides*, foliis lanceolatis (passim oblongis) tri-

nervibus petiolatis ramulisque glaberrimis, capitulis globosis ovalibusve, calycis laciniis acutis trinerviis, phalangibus subdecandris: unguibus petalis brevioribus. *Brown mss.*

Melaleuca thymoides. *Labillard. nov. holl. 2, p. 27, t. 167.*

Nat. of the South-west Coast of New Holland. *Robert Brown.* ♀.

10. *M. squamea*, foliis ovatis lanceolatis acuminatis trinerviis: novellis ramulisque villosis, capitulis globosis pubescentibus, phalangibus 5—6-andris: unguibus brevissimis. *Brown mss.*

Malaleuca squamea. *Labillard. nov. holl. 2, p. 28, t. 168.*

Nat. of Van Diemen's Island. ♀.

[413]

11. *M. nodosa*, foliis subulato-linearibus mucronatis rigidis uninerviis planis, capitulis globosis, calycis laciniis membranaceis glabris, phalangibus 3—6-andris: unguibus petalis brevioribus. *Brown mss.*

Malaleuca nodosa. *Willden. sp. pl. 3, p. 1431. Smith exot. bot. 1, p. 67, t. 35. Venten. malmais. 112.*

Nat. of New South Wales. The Right Hon. Sir *Joseph Banks*, Bart. K.B. ♀.

12. *M. ericifolia*, foliis lineari-subulatis enerviis muticis patentibus subrecurvisque, spicis ovalibus glabris, phalangibus 8—10-andris: unguibus petala vix æquantibus. *Brown mss.*

Malaleuca ericifolia. *Willden. sp. pl. 3, p. 1430. Smith exot. bot. 1, p. 65, t. 34.*

Nat. of New South Wales. *John White*, M.D. ♀.

13. *M. armillaris*, foliis lineari-subulatis mucronatis apice recurvis, spicis cylindricis glaberrimis, phalangibus polyandris: unguibus petala superantibus. *Brown mss.*

Melaleuca armillaris. *Willden. sp. pl. 3, p. 1431.*

Melaleuca ericæfolia. *Andrews's reposit. 175. Venten. malmais. 76.*

Nat. of New South Wales. The Right Hon. Sir *Joseph Banks*, Bart. K.B. h.

14. *M. uncinata*, foliis angulato filiformibus mucronatis erectis: apice uncinato recurvo, ramulis virgatis, capitulis ovalibus: rachi lanata, phalangibus 5—6-andris: unguibus petala superantibus. *Brown mss.*

Nat. of the South Coast of New Holland. *Robert Brown*. h.

15. *M. scabra*, foliis teretiusculis mucronulatis scabris confertis, capitulis globosis, phalangibus 4—6-andris: unguibus petala subæquantibus. *Brown mss.*

Nat. of the South Coast of New Holland. *Robert Brown*. h.

16. *M. pulchella*, foliis sparsis suboppositisque ovalibus obtusis obsolete 3-nervibus floribusque subsolitariis glabris, phalangibus polyandris intus basin usque ramosis. *Brown mss.*

Nat. of the South Coast of New Holland. *Robert Brown*. h.

Folia opposita v. verticillata.

17. *M. thymifolia*, foliis oppositis lanceolatis enervibus, spicis paucifloris, phalangibus polyandris: unguibus intus medium usque ramosis. *Brown mss.*

Melaleuca thymifolia. *Willden. sp. pl.* 3, p. 1432. *Smith exot. bot.* 1, p. 69, t. 36.

Melaleuca gnidiæfolia. *Venten. malmais.* 7.

Melaleuca coronata. *Andrews's reposit.* 278.

⁴¹⁵*Nat.* of New South Wales. The Right Hon. Sir *Joseph Banks*, Bart. K.B. h.

18. *M. decussata*, foliis oppositis decussatis ovali-lanceolatis trinervibus, spicis ovalibus glaberrimis, phalangibus polyandris: unguibus brevissimis. *Brown mss.*

Nat. of the South Coast of New Holland. Robert Brown.
h.

19. *M. fulgens*, foliis oppositis lanceolato-linearibus acutis uninerviis, spicis ovalibus glaberrimis, phalangibus multifidis: unguibus longitudine petalorum. *Brown mss.*

Nat. of the South Coast of New Holland. Robert Brown. *h.*

20. *M. linariifolia*, foliis oppositis lanceolato-linearibus acutis trinerviis, spicis oblongis glabris, phalangibus polyandris longitudinaliter pinnatis. *Brown mss.*

Melaleuca linariifolia. Willden. sp. pl. 3, p. 1482. Smith exot. bot. 1, p. 109, t. 56.

Nat. of New South Wales. Colonel William Paterson.
h.

21. *M. hypericifolia*, foliis oppositis elliptico-oblongis 3-nerviis: nervis lateralibus obsoletis margini recurvo approximatis, spicis glaberrimis, phalangibus polyandris: unguibus elongatis; filamentis radiantibus. [116]

Melaleuca hypericifolia. Willden. sp. pl. 3, p. 1433. Andrews's reposit. 200. Venten. cels. 10.

Nat. of New South Wales. Admiral Phillip. *h.*

22. *M. squarrosa*, foliis oppositis ovatis acutis 5—7-nerviis petiolatis, spicis oblongis ovalibusque: bracteis foliaceis, calycis laciniis enerviis obtusis, phalangium unguibus brevissimis.

Melaleuca squarrosa. Willden. sp. pl. p. 430. Labillard. nov. holl. 2, p. 28, t. 169.

Melaleuca myrtifolia. Venten. malmais. 47.

Nat. of New South Wales and Van Diemen's Island. Mr. David Nelson. *h.*

23. *M. calycina*, foliis oppositis ovato-lanceolatis 3—5-nerviis subsessilibus, glomerulis paucifloris, calycis laciniis acutis enerviis, phalangibus polyandris: unguibus petalo brevioribus. *Brown mss.*

Nat. of the South-west Coast of New Holland. Robert Brown. h.

24. *M. densa*, foliis ternis-obovatis trinerviis glabris, spicis oblongis ovalibusve. *Brown mss.*

Nat. of the South-west Coast of New Holland. Robert Brown. h.

471

TRISTANIA. *Brown mss.*

Staminum phalanges 5, petalis oppositæ iisdemque vix longiores. *Antheræ* incumbentes. *Caps.* 3-loc. polysperma (semiexserta v. inclusa et) connata calycis tubo turbinato pedicellato (in pedunculo communi).

1. *T. nereifolia*, foliis oppositis lanceolatis, phalangibus 3—5-andris. *Brown mss.*

Melaleuca nereifolia. Sims in botan. magaz. 1058.

Melaleuca salicifolia. Andrews's reposit. 485.

Nat. of New South Wales. Robert Brown. h.

2. *T. laurina*, foliis alternis cuneato-lanceolatis, ramulis calycibusque pubescentibus, capsulis semisuperis. *Brown mss.*

Melaleuca laurina. Willden. sp. pl. 3, p. 1429.

Nat. of New South Wales. Admiral Phillip. h.

3. *T. conferta*, foliis lanceolato-ellipticis acutis alternis: terminalibus confertis, calycis laciniis acutis foliaceis. *Brown mss.*

Nat. of New South Wales. Robert Brown. h.

CALOTHAMNUS. (*Labillard. nov. holl.* 2, p. 24.)

Brown mss.

Staminum phalanges (4-5) petalis oppositæ (aliquæ nunc connatæ v. steriles). *Antheræ* basi insertæ, integerrimæ. *Caps.* 3-loc. polysperma, connata et inclusa calycis tubo incrassato basi adnato (ramo).

1. *C. quadrifida*, floribus 4-fidis, phalangibus distinctis æqualibus 12—15-andris, foliis adultis fructibusque glabris. *Brown mss.*

Nat. of the South-west Coast of New Holland. *Robert Brown.* h.

2. *C. villosa*, floribus 5-fidis, phalangibus distinctis æqualibus polyandris, foliis adultis fructibusque villosis. *Brown mss.*

Nat. of the South-west Coast of New Holland. *Robert Brown.* h.

3. *C. gracilis*, floribus 5-fidis, phalangibus distinctis æqualibus 3-andris, foliis longissimis fructibusque exsertis glabris, caule ramoso. *Brown mss.*

Nat. of the South-west Coast of New Holland. *Robert Brown.* h.

BEAUFORTIA. *Brown mss.*

Staminum phalanges 5, petalis oppositæ. *Antheræ* basi insertæ: apice bifidæ lobis deciduis! *Capr.* 3-loc. monosperma, connata et inclusa calycis tubo incrassato basi adnato (ramo).

1. *B. decussata*, foliis oppositis decussatis ovatis ovalibusve multinerviis, phalangium unguibus longissimis; filamentis radiantibus. *Brown mss.*

Nat. of the South-west Coast of New Holland. *Robert Brown.* h.

2. *B. sparsa*, foliis sparsis ovalibus multinerviis. *Brown mss.*

Nat. of the South-west Coast of New Holland. *Robert Brown.* h.

GENERA ET SPECIES QUÆDAM

PLANTARUM COMPOSITARUM,

QUÆ

IN HORTO KEWENSI COLUNTUR.

[*Extracted from 'Hortus Kewensis,' 2nd Edition, Vols. IV and V,
var. pages.*]

LONDON.

—
1812-1813.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

COMPOSITÆ.

IXODIA. *Brown mss.*

[iv. 617]

Recept. paleaceum. *Pappus* nullus. *Cal.* imbricatus, squamis interioribus radiantibus coloratis.

1. IXODIA *Achillæoides*. *Brown mss.*

Nat. of the South Coast of New Holland. *Robert Brown.* ♀.

DONIA. *Brown mss.*

[v. 82]

Recept. nudum. *Pappus* setaceus, caducus. *Calyx* imbricatus, hemisphæricus.

1. D. *glutinosa*, foliis ovato-oblongis serratis calycibusque viscidis. *Brown mss.*

Doronicum glutinosum. *Willden. sp. pl.* 3, p. 2115.

Aster glutinosus. *Cavanill. ic.* 2, p. 52, t. 168.

Nat. of Mexico. ♀.

PODOLEPIS. (*Labill. nov. holl.* 2, p. 56.) *Brown mss.*

Recept. nudum. *Pappus* pilosus. *Calyx* imbricatus scariosus hemisphæricus: squamis unguiculatis.

1. P. *rugata*, squamis calycis rugosis obtusis, caule simplicis. *Brown mss.*

Podolepis rugata. *Labill. nov. holl.* 2, p. 57, t. 208.

Nat. of the South-west Coast of New Holland. ♀.

2. *P. acuminata*, squamis calycis æquatis ovatis acuminatis, caule subsimplici. *Brown mss.*

Scalia Jaceoides. Sims in Botan. magaz. 956.

Nat. of New South Wales. Robert Brown. 4.

v. 137]

CULLUMIA. *Brown mss.*

Recept. favosum. Semina glabra. Pappus nullus. Cal. monophyllus: foliolis imbricatis tectus.

1. *C. ciliaris*, foliis ovatis glabris imbricatis bifariam ciliato-spinosis: spina terminali reflexa. *Brown mss.*

Berckheya ciliaris. Willden. sp. pl. 3, p. 2273.

Carlina foliis imbricatis oblongis reticulatis et in aculeum aduncum desinentibus. Burm. afr. 151, t. 54, f. 1.

Nat. of the Cape of Good Hope. 2.

2. *C. setosa*, foliis ovato-lanceolatis glabris recurvatis ciliato-spinosis. *Brown mss.*

Berckheya setosa. Willden. sp. pl. 3, p. 2273. (exclus. syn. Commelini.)

Nat. of the Cape of Good Hope. 2.

v. 138] 3. *C. squarrosa*, foliis subulatis recurvatis patulisve spinuloso-ciliatis subtus sulcatis: adultis ramisque glabriusculis. *Brown mss.*

Berckheya squarrosa. Willden. sp. pl. 3, p. 2272.

Rohria squarrosa. Thunb. in act. soc. nat. scrut. hafn. 3, part. 1, pag. 100, t. 5.

Nat. of the Cape of Good Hope. 2.

BERCKHEYA. (*Gen. pl. 1329.*) *Brown mss.*

Recept. favosum. Semina villosa. Pappus paleaceus (nunc setoso-paleaceus ciliatus). Cal. monophyllus: foliolis imbricatis tectus.

1. *B. cynaroides*, foliis caulinis alternis amplexicaulibus spinuloso-ciliatis; radicalibus elongatis integris inermibus

subtus tomentosis, calycis foliolis ovatis integerrimis. *Brown mss.*

Berckheya cynaroides. *Willden. sp. pl.* 3, p. 2275.

Nat. of the Cape of Good Hope. ♀.

2. *B. obovata*, foliis alternis cuneato-lanceolatis spinoso-dentatis utrinque glabris, foliolis calycis dentato-spinosis. *Brown mss.*

Berckheya obovata. *Willden. sp. pl.* 3, p. 2269.

Basteria aculeata. *Houtt. nat. hist.* 6, p. 158, t. 34, f. 2.

Nat. of the Cape of Good Hope. ♀.

3. *B. incana*, foliis alternis ovatis spinoso-dentatis subtus caulibusque incanis, calycis foliolis dentato-spinosis [v. 139] subtus tomentosis. *Brown mss.*

Berckheya incana. *Willden. sp. pl.* 3, p. 2269.

Gorteria asteroides. *Jacqu. ic.* 3, t. 591.

Nat. of the Cape of Good Hope. ♀.

4. *B. cernua*, foliis alternis lanceolatis amplexicaulibus spinoso-dentatis ciliatis utrinque glabris, floribus cernuis, pappo setoso-ciliato. *Brown mss.*

Gorteria cernua. *Willden. sp. pl.* 3, p. 2268.

Gorteria araneosa. *Meerb. ic.* 1, t. 40.

Nat. of the Cape of Good Hope. ♂.

DIDELTA. (*Gen. pl.* 1351.) *Brown mss.*

Recept. favosum includens semina. *Pappus* multipartitus setoso-paleaceus dentatus. *Cal.* monophyllus: tectus foliolis: exterioribus maximis.

1. *D. carnosum*, foliis alternis.

Didelta carnosa. *Willden. sp. pl.* 3, p. 2262.

Didelta tetragonæfolia. *L'Herit. stirp. nov.* 55, t. 28.

Nat. of the Cape of Good Hope. Mr. Fr. Masson. ♀.

2. *D. spinosum*, foliis oppositis.

Didelta spinosa. Willden. *sp. pl.* 3, p. 2262. *Wendl. obs.* 33, t. 4, f. 32.

Nat. of the Cape of Good Hope. Mr. Fr. Masson. ♀.

v. 140] GORTERIA. (*Gen. pl.* 1328.) *Brown mss.*

Recept. scrobiculatum. *Pappus*: margo ciliatus. *Cal.* monophyllus: foliolis imbricatis tectus; fructus induratus connivens deciduus.

1. GORTERIA *personata*.

Gorteria personata. Willden. *sp. pl.* 3, p. 2265. *Jacqu. collect.* 4, p. 224, t. 21, f. 1.

Nat. of the Cape of Good Hope. ☉.

GAZANIA. (*Gærtn. sem.* 2, p. 451.) *Brown mss.*

Recept. epaleatum (nudum; v. alveolatum germinibus exsertis.) *Semina* villosissima. *Pappus* piloso-paleaceus. *Cal.* monophyllus: tubo foliolis imbricatis tecto v. nudo.

1. *G. rigens*, foliis spathulato-lanceolatis indivisis passim pinnatifidis subtus tomentosis, scapo unifloro calycibusque glabris, caule decumbente. *Brown mss.*

Gorteria rigens. Willden. *sp. pl.* 3, p. 2267. *Curtis magaz.* 90.

Nat. of the Cape of Good Hope. ♀.

2. *G. Pavonia*, foliis pinnatifidis supra pilosis subtus tomentosis: laciniis ovali-lanceolatis, scapo unifloro, caule decumbente. *Brown mss.*

Gazania Pavonia. *Andrews's reposit.* 523.

Nat. of the Cape of Good Hope. Mr. James Niven. ♂.

3. *G. subulata*, caule folioso decumbente unifloro, foliis subulato-linearibus margine revolutis subtus tomentosis. [v. 141] *Brown mss.*

Nat. of the Cape of Good Hope. Mr. Fr. Masson. ♀.

CRYPTOSTEMMA. *Brown mss.*

Recept. favosum. *Pappus* paleaceus lana implexa seminis tectus. *Cal.* imbricatus.

1. *C. calendulaceum*, ligulis indivisis, foliis pinnatifidis dentatis subtus tomentosis. *Brown mss.*

Arctotis calendulacea. *Willden. sp. pl.* 3, p. 2347. *Jacqu. hort. schænbr.* 2, p. 16, t. 157.

Nat. of the Cape of Good Hope. ☉.

2. *C. hypochondriacum*, ligulis (3-5-) partitis, foliis lyratis tomentosis. *Brown mss.*

Arctotis hypochondriaca. *Willden. sp. pl.* 3, p. 2348.

Nat. of the Cape of Good Hope. ☉.

3. *C. runcinatum*, ligulis (3-5-) partitis, foliis runcinatis dentatis subtus tomentosis. *Brown mss.*

Nat. of the Cape of Good Hope. Mr. Fr. Masson. ☉.

ARCTOTHECA. (*Wendl. hort. herrenhus.* 8.) *Brown mss.*

Recept. favosum. *Pappus* nullus. *Cal.* imbricatus.

1. *ARCTOTHECA repens*. *Willden. sp. pl.* 3, p. 2365. [v. 142]

Arctotis repens. *Jacqu. hort. schænbr.* 3, p. 31, t. 306.

Nat. of the Cape of Good Hope. ♀.

SPHENOGYNE. *Brown mss.*

Recept. paleis distinctis. *Pappus* paleaceus, simplex. *Stigmata* apice dilatato subtruncato. *Cal.* imbricatus: squamis intinis (omnibusve) apice dilatato scarioso.

1. *S. anthemoides*, herbacea glabra, foliis bipinnatifidis pinnatifidisve lineari-filiformibus, pappi foliolis albis: axi

subulato discolori. *Willden. sp. pl.* 3, p. 2361. *Botan. magaz.* 544.

Arctotis paradoxa. [*Hort. Kew.*] *Edit. prior.* 3, p. 273.

Nat. of the Cape of Good Hope. ☉.

2. *S. crithmifolia*, fruticosa glabra, foliis pinnatifidis lineari-filiformibus, foliolis exterioribus calycis subulatis. *Brown mss.*

Arctotis paleacea. *Willden. sp. pl.* 3, p. 2359.

Arctotis crithmoides. *Berg. cap.* 326.

Nat. of the Cape of Good Hope. ♀.

3. *S. scariosa*, fruticosa, foliis bipinnatifidis pinnatifidisve lineari-filiformibus glabris, calycis squamis omnibus scariosis obtusis nitidis: extimis sessilibus. *Brown mss.*

Arctotis scariosa. *Willden. sp. pl.* 3, p. 2360.

v. 148] *Nat. of the Cape of Good Hope.* Mr. *Fr. Masson.* ♀.

4. *S. abrotanifolia*, fruticosa, foliis 2-3-pinnatifidis calycibusque tomentosis. *Brown mss.*

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* ♀.

5. *S. dentata*, fruticosa, foliis pinnatifidis glabriusculis: laciniis 2-3-dentatis: dentibus piliferis, calycis foliolis exterioribus lanceolatis. *Brown mss.*

Arctotis dentata. *Willden. sp. pl.* 3, p. 2359.

Chrysanthemum foliorum pinnis brevissimis dentatis. *Burm. afr.* 175, t. 64.

Nat. of the Cape of Good Hope. ♀.

6. *S. odorata*, fruticosa, foliis planis glabris apice inciso-pinnatifidis, calycis foliolis exterioribus apice scariosis, seminibus glabris: pappo obsoleto. *Brown mss.*

Anthemis odorata. *Willden. sp. pl.* 3, p. 2186.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* ♀.

ARCTOTIS. (*Gen. pl.* 1340.) *Brown mss.* [v. 169]

Recept. setoso-alveolatum. *Semina* dorso semi-biloculari v. bisulco. *Pappus* paleaceus. *Cal.* imbricatus: squamis apice scariosis.

1. *A. acaulis*, caule brevissimo decumbente, foliis utrinque incanis ternato-lyratis dentatis, calycinis squamis exterioribus adpressis. *Willden. sp. pl.* 3, p. 2348.

Arctotis acaulis. *Trew, ehret.* 53, t. 93.

Nat. of the Cape of Good Hope. 4.

2. *A. tricolor*, foliis subtus tomentosis ovatis integerrimis lyratisve dentatis, scapo sulcato unifloro, squamis calycinis exterioribus reflexis. *Willden. sp. pl.* 3, p. 2349.

Arctotis tricolor. *Jacqu. hort. schænbr.* 2, p. 27, t. 159.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* 4.

3. *A. undulata*, foliis subtus tomentosis undulato-dentatis ovatis lyratisve, scapis unifloris striatis, calycinis squamis exterioribus reflexis. *Willden. sp. pl.* 3, p. 2349.

Arctotis undulata. *Jacqu. hort. schænbr.* 2, p. 17, t. 160.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* 4.

4. *A. grandiflora*, foliis pinnatifidis denticulatis arachnoideis triplinerviis.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* ♂.

Obs. Caulescens. *Radius* maximus. *Petala* straminea. subtus rubore tincta, supra prope basin lutescentia, ore nigro-purpureo.

5. *A. glaucophylla*, foliis canescentibus pinnatifidis repando-subdentatis, caule basi ramoso, calycinis squamis exterioribus reflexis. *Willden. sp. pl.* 3, p. 2350.

Arctotis glaucophylla. *Jacqu. hort. schænbr.* 2, p. 22, t. 170.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* 4.

6. *A. plantaginea*, foliis lanceolato-ovatis nervosis denticulatis amplexicaulibus. *Willden. sp. pl.* 3, p. 2350.

Nat. of the Cape of Good Hope. 4.

v. 171] 7. *A. argentea*, foliis lanceolato-linearibus integerrimis tomentosis. *Willden. sp. pl.* 3, p. 2351.

Nat. of the Cape of Good Hope. Mr. Fr. Masson. ♂.

8. *A. rosea*, caule procumbente, foliis spathulato-lanceolatis repando-dentatis incanis, pappo duplici. *Willden. sp. pl.* 3, p. 2351.

Arctotis rosea. *Jacqu. hort. schænbr.* 2, p. 18, t. 162.

Nat. of the Cape of Good Hope. Mr. Fr. Masson. 4.

9. *A. decumbens*, caule procumbente, foliis oblongo-lanceolatis inæqualiter dentatis canescentibus: subtus tomentosis trinerviis, calycinis squamis exterioribus adpressis. *Willden. sp. pl.* 3, p. 2351.

Arctotis decumbens. *Jacqu. hort. schænbr.* 3, p. 68, t. 381.

Nat. of the Cape of Good Hope. Mr. Fr. Masson. 4.

10. *A. angustifolia*, caule ramoso adscendente, foliis tomentosis spathulato-lanceolatis subdentatis trinerviis pubescentibus, calycinis squamis exterioribus reflexis. *Willden. sp. pl.* 3, p. 2352.

Nat. of the Cape of Good Hope. 2.

11. *A. flaccida*, caule ramoso herbaceo adscendente, foliis spathulato-lanceolatis integerrimis trinerviis pubescentibus, calycinis squamis omnibus reflexis. *Willden. sp. pl.* 3, p. 2352.

v. 172] *Arctotis flaccida.* *Jacqu. hort. schænbr.* 2, p. 19, t. 163.

Nat. of the Cape of Good Hope. Mr. Fr. Masson. ☉.

12. *A. decurrens*, caule fruticoso, foliis hirtis oblongis indivisis subdentatis. *Willden. sp. pl.* 3, p. 2353.

Arctotis decurrens. *Jacqu. hort. schænbr.* 2, p. 20, t. 165.

Nat. of the Cape of Good Hope. Mr. Fr. Masson. 2.

13. *A. reptans*, caule herbaceo adscendente, foliis hirtis subtus incanis: inferioribus lyrato-dentatis; superioribus lanceolatis dentatis. *Willden. sp. pl.* 3, p. 2354.

Arctotis reptans. *Jacqu. hort. schænbr.* 3, p. 69, t. 382.
Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* 4.

14. *A. auriculata*, caule fruticoso niveo-tomentoso, foliis lyratis amplexicaulibus tomentosis dentatis: lobo terminali rhombico. *Willden. sp. pl.* 3, p. 2354.

Arctotis auriculata. *Jacqu. hort. schænbr.* 2, p. 22, t. 169.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* 2.

15. *A. fastuosa*, caule erecto herbaceo, foliis hirtis oblongis dentatis, calycinis squamis exterioribus reflexis ciliatis. *Willden. sp. pl.* 3, p. 2354.

Arctotis fastuosa. *Jacqu. hort. schænbr.* 2, p. 21, t. 166.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* (v. 173) 3.

16. *A. spinulosa*, caule erecto herbaceo, foliis cano-hirtis viscosis oblongis amplexicaulibus mucronato-dentatis, calycinis squamis exterioribus patulis. *Willden. sp. pl.* 3, p. 2355.

Arctotis spinulosa. *Jacqu. hort. schænbr.* 2, p. 21, t. 167.

Nat. of the Cape of Good Hope. Mr. *Fr. Masson.* 2.

17. *A. aspera*, caule erecto fruticoso, foliis pinnatifidis scabris subtus tomentosis margine revolutis: laciniis lanceolatis obtusis mucronato-dentatis. *Willden. sp. pl.* 3, p. 2356.

a. *Anemonospermum africana*, foliis cardui benedicti, florum radiis intus sulphureis. *Commel. hort.* 2, p. 43, t. 22.

β. *Anemonospermum afra*, folio Jacobææ tenuiter laciniato, flore aurantio pulcherrimo. *Boerh. lugdb.* 1, p. 100, cum tab.

Nat. of the Cape of Good Hope. 2.

v. 180] ERIOCEPHALUS. (*Gen. pl.* 1344.) *Brown mss.*

Recept. paleaceum *Pappus* nullus. *Cal.* duplex: interior 1-phyllus; exterior 5-phyllus.

1. *E. frutescens*, foliis sericeis, floribus corymbosis.

Eriocephalus africanus. *Willden. sp. pl.* 3, p. 2384.
Botan. magaz. 833.

Nat. of the Cape of Good Hope. 2 .

2. *E. racemosus*, foliis sericeis, floribus racemosis.

Eriocephalus racemosus. *Willden. sp. pl.* 3, p. 2385.

Nat. of the Cape of Good Hope. 2 .

v. 181] CASSINIA. *Brown mss.*

Calyces 2-flori, 4-phylli. *Corollulæ* hermaphroditæ. *Pappus* paleaceus, penicillatus. *Recept. partiale* nudum.

v. 185] 1. CASSINIA aurea. *Brown mss.*

Nat. of the South Coast of New Holland. *Robert Brown.* 4 .

88.
: in-
2384.
5.
Pap-
obert

GENERA ET SPECIES

PLANTARUM ORCHIDEARUM,

QUE

IN HORTO KEWENSIS COLUNTUR.

[*Extracted from 'Hortus Kewensis,' 2nd Edition, Vol. V,*
pp. 188—222.]

LONDON.

1813.

GYNANDRIA.

[188

MONANDRIA.

1. *Anthera* adnata subterminalis persistens. *Pollinis* *massæ* e lobulis angulatis elasticæ cohærentibus; basi affixæ.

ORCHIS. *Brown prodr.* 312.

Cor. ringens. *Labellum* basi subtus calcaratum. *Glandulæ* (1—2) pedicellorum pollinis inclusæ cucullo unico.

1. *O. Morio*, labello trilobo: lobis crenatis obtusis: medio emarginato, petalis adscendentibus obtusis, cornu conico adscendente germine brevior. *Willden. sp. pl.* 4, p. 18. *Curtis lond. Engl. bot.* 2059.

Nat. of Britain. 4.

2. *O. mascula*, labello trilobo crenulato obtuso: lobo medio bifido, petalis acutis: exterioribus reflexis, cornu conico adscendente germinis longitudine. *Willden. sp. pl.* 4, p. 18. *Engl. bot.* 631.

Nat. of Britain. 4.

3. *O. ustulata*, labello tripartito: laciniis linearibus punctato-scabris: media bipartita, petalis erectis acutis, cornu uncinato germine triplo brevior, bracteis germen subæquantibus. *Willden. sp. pl.* 4, p. 20. *Engl. bot.* 18.

Nat. of England. 4.

4. *O. fusca*, labello tripartito punctato-scabro: laciniis lateralibus oblongis; intermedia ampla biloba crenata cum mucrone interjecto, petalis acutis conniventibus, cornu rectiusculo germine triplo brevior, bracteis germine quadruplo brevioribus. *Willden. sp. pl.* 4, p. 23. *Curtis lond.*

Orchis militaris. Engl. bot. 16.

Nat. of England. 4.

5. *O. militaris*, labello tripartito punctato-scabro: laciniis lateralibus linearibus; intermedia biloba obtusa cum mucrone interjecto, petalis acutis conniventibus, cornu recto germine duplo brevior, bracteis obsoletis. *Willden. sp. pl.* 4, p. 22. *Engl. bot.* 1873.

Nat. of England. 4.

6. *O. globosa*, labello tripartito: lacinia media emarginata, petalis apice mucronatis, cornu germine duplo brevior, spica dense ovata, foliis lanceolatis. *Willden. sp. pl.* 4, p. 14. *Jaqu. austr.* 3, p. 36, t. 265.

Nat. of Austria and Switzerland. 4.

7. *O. pyramidalis*, labello tripartito: superne basi bicorni; laciniis æqualibus integerrimis, cornu germen sub-¹⁹⁰¹æquante, spica densa oblonga. *Willden. sp. pl.* 4, p. 14. *Engl. bot.* 110.

Nat. of Britain. 2.

8. *O. hircina*, labello tripartito: laciniis lateralibus lineari-subulatis; media elongata germine triplo longiore lineari bifida, petalis conniventibus, cornu brevissimo conico didymo. *Willden. sp. pl.* 4, p. 28.

Satyrium hircinum. Eng. bot. 34.

Nat. of England. 2.

9. *O. latifolia*, labello leviter trilobo lateribus reflexo, petalis superioribus conniventibus; binis lateralibus reflexis, cornu conico germine brevior, bracteis flore longioribus. *Willden. sp. pl.* 4, p. 28. *Curtis lond. Engl. bot.* 2308.

Nat. of Britain. 4.

10. *O. maculata*, labello plano trilobo crenato, petalis superioribus conniventibus; lateralibus patentibus, cornu cylindraceo germine brevior, bracteis longitudine geminis. *Willden. sp. pl. 4, p. 31. Engl. bot. 632.*

Nat. of Britain. 4.

11. *O. spectabilis*, labello obovato indiviso crenato retuso, petalis rectis; lateralibus longioribus, cornu clavato germine brevior, bracteis flore longioribus, caule aphylo. *Willden. sp. pl. 4, p. 36.*

Nat. of North America. 4.

12. *O. papilionacea*, labello obovato indiviso dentato emarginato, petalis nervosis conniventibus, cornu subulato germine brevior, bracteis membranaceis coloratis germine longioribus. *Willden. sp. pl. 4, p. 24.*

Nat. of Spain and Naples. 4.

GYMNADENIA. *Brown mss.*

Cor. ringens. Labellum basi subtus calcaratum. Glandulæ pedicellorum pollinis nudæ approximatae.

1. *GYMNADENIA conopsea. Brown mss.*

Orchis conopsea. Willden. sp. pl. 4, p. 32. Engl. bot.

10.

Nat. of Britain. 4.

ACERAS. *Brown mss.*

Cor. ringens. Labellum ecalcaratum. Glandulæ pedicellorum pollinis cucullo communi inclusæ.

1. *A [anthropophora] labello germine longiore. Brown mss.*

Ophrys anthropophora. Willden. sp. pl. 4, p. 63.

Curtis lond. Engl. bot. 29.

Nat. of England. 4.

HERMINIUM. *Brown mss.*

Cor. subpatens. *Labellum* ecalcaratum. *Glandulæ* massarum pollinis nudæ distinctæ.

1. *H. Monorchis*, foliis radicalibus lanceolatis binis. *Brown mss.*

Ophrys Monorchis. *Willden. sp. pl.* 4, p. 61. *Engl. bot.* 71.

¹⁹²⁷ *Nat.* of England. 4

HABENARIA. *Brown prodr.* 312.

Cor. ringens. *Labellum* basi subtus calcaratum. *Glandulæ pollinis* nudæ distinctæ (loculis pedicellorum adnatis v. solutis distinctis).

1. *H. nigra*, cornu abbreviato didymo, labello ovato acuminato, spica densa ovata, foliis linearibus. *Brown mss.*

Orchis nigra. *Willden. sp. pl.* 4, p. 35.

Satyrium nigrum. *Fl. dan.* 998.

Nat. of the Alps of Switzerland, Austria, and Lapland. 4.

2. *H. viridis*, cornu abbreviato didymo, labello lineari tridentato: lateralibus acutis; medio brevissimo, bracteis flore sesquolongioribus. *Brown mss.*

Orchis viridis. *Willden. sp. pl.* 4, p. 33.

Satyrium viride. *Engl. bot.* 94.

Nat. of Britain. 4.

3. *H. bracteata*, cornu abbreviato didymo, labello lineari retuso-tridentato: lateralibus obtusis; medio obsoleto, bracteis flore duplo longioribus. *Brown mss.*

Orchis bracteata. *Willden. sp. pl.* 4, p. 34.

Orchis bractealis. *Salisb. paradis.* 110.

Nat. of North America. 4.

4. *H. albida*, cornu obtuso germine triplo brevior, ¹⁹³labello tripartito : laciniis acutis : media majore. *Brown mss.*

Orchis albida. *Willden. sp. pl. 4, p. 38.*

Satyrium albidum. *Engl. bot. 505.*

Nat. of Britain. 4.

5. *H. hyperborea*, cornu cylindrico germine brevior, labello integerrimo lineari-oblongo. *Brown mss.*

Orchis hyperborea. *Willden. sp. pl. 4, p. 37.*

Nat. of Iceland and Labrador. 4.

6. *H. bifolia*, cornu filiformi germine duplo longior, labello lineari integerrimo, foliis radicalibus binis oblongis basi attenuatis. *Brown mss.*

Orchis bifolia. *Willden. sp. pl. 4, p. 10. Curtis lond. Engl. bot. 22.*

Nat. of Britain. 4.

7. *H. herbiola*, cornu filiformi germine brevior, labello oblongo obtuso basi utrinque dentato ; palato unidentato, bracteis flore longioribus. *Brown mss.*

Nat. of North America. 4.

8. *H. fimbriata*, cornu filiformi germine longior, labello tripartito laciniis cuneiformibus fimbriatis. *Brown mss.*

Orchis fimbriata. *Willden. sp. pl. 4, p. 39.*

Nat. of Canada and Newfoundland. 4.

9. *H. cristata*, cornu filiformi germine brevior, labello ¹⁹⁴lanceolato pinuatim-fimbriato, petalis interioribus dentato-incisis. *Brown mss.*

Orchis cristata. *Willden. sp. pl. 4, p. 9.*

Nat. of North America. 4.

10. *H. ciliaris*, cornu filiformi germine longior, labello lanceolato pinnatim-fimbriato, petalis interioribus fimbriato-incisis. *Brown mss.*

Orchis ciliaris. *Willden. sp. pl. 4, p. 8. Andrews's reposit. 42.*

Nat. of North America. 4.

BARTHOLINA. *Brown mss.*

Cor. ringens: petalis interioribus inferne labello connatis. *Labellum* basi subtus calcaratum. *Pedicelli pollinis* elongati: loculis adnatis; glandulis distinctis lobulo exteriori semitectis.

1. BARTHOLINA *pectinata*. *Brown mss.*

Orchis pectinata. *Willden. sp. pl.* 4, p. 11.

Orchis burmanniana. *Swartz in Web. et Mohr archiv.*

1, p. 55, t. 3.

Nat. of the Cape of Good Hope. 4.

SERAPIAS. (*Swartz in act. stockholm.* 1800, p. 223.)
Brown mss.

Cor. ringens. *Labellum* ecalcaratum. *Columna* cuspidata. *Pedicelli pollinis* inserti glandula unica cucullo inclusa.

195] 1. *S. Lingua*, labello tripartito: laciniis lateralibus obtusis erectis conniventibus; media oblonga lanceolata acutiuscula glabriuscula dependente. *Willden. sp. pl.* 4, p. 70.

Orchis montana italica, flore ferrugineo, lingua oblonga. *Moris. hist.* 3, p. 495, s. 12, t. 14, f. 21.

Nat. of the South of Europe. 4.

2. *S. cordigera*, labello tripartito: laciniis lateralibus obtusis erectis conniventibus; media ovata acuminata disco pilosa dependente. *Willden. sp. pl.* 4, p. 71. *Andrews's reposit.* 475.

Nat. of Spain, Italy, and the Levant. 4.

OPHRYS. *Brown prodr.* 313.

Cor. subpatens. *Labellum* ecalcaratum. *Glandulæ pol-
linis* cucullis distinctis inclusæ.

1. *O. apifera*, labello trifido: lobo medio majore semitri-
fido: lacinia media longiore subulata deflexa. *Brown mss.*
Ophrys apifera. *Willden. sp. pl.* 4, p. 66. *Curtis. lond.*
Engl. bot. 383.

Nat. of England. 2.

2. *O. aranifera*, labello trilobo: lateralibus abbreviatis
obtusis; medio retuso. *Brown mss.*

Ophrys aranifera. *Willden. sp. pl.* 4, p. 66. *Engl.*
bot. 65.

Nat. of England. 2.

3. *O. muscifera*, labello trifido: lobo medio majore
bilobo, anthera obtusa. *Brown mss.*

Ophrys muscifera. *Smith fl. brit.* 3, p. 937. *Engl.*
bot. 64.

Ophrys myodes. *Willden. sp. pl.* 4, p. 64. *Andrews's*
reposit. 471.

Nat. of England. 2.

SATYRIUM. (*Swartz in act. stockholm.* 1800, p. 214.)

Brown mss.

Cor. ringens: petala 5 antica basi connata. *Labellum*
posticum, fornicatum, basi bicalcaratum v. bisaccatum. *An-
thera* resupinata. *Stigma* bilabiatum.

1. *S. cucullatum*, foliis radicalibus binis cordato-subro-
tundis concavis; caulinis remotis cucullatis subretusis, flori-
bus cernuis. *Willden. sp. pl.* 4, p. 54.

Orchis bicornis. *Andrews's reposit.* 315.

Nat. of the Cape of Good Hope. 2.

2. *S. carneum*, foliis radicalibus binis cordato-subrotundis; caulinis vaginæformibus approximatis, spica compacta, petalis extus carinatis.

Orchis carnea. [*Hort. Kew.*] *Edit. prior.*

Nat. of the Cape of Good Hope. Mr. Fr. Masson. ۛ.

DISA. *Swartz in act. stockholm. 1800, p. 208.*

Cor. ringens: galea basi unicalcarata v. saccata petala interiora columnæ accreta. *Labellum* ecalcaratum.

1973 1. *D. cornuta*, galea obtusa; calcare conico deflexo, petalis interioribus bidentatis, labello obvato velutino plano, spica laxa. *Willden. sp. pl. 4, p. 45.*

Nat. of the Cape of Good Hope. ۛ.

2. *D. spathulata*, galea erecta acuta, labello petiolato apice dilatato trifido, caule paucifloro, foliis linearibus. *Willden. sp. pl. 4, p. 52.*

Nat. of the Cape of Good Hope. ۛ.

PTERYGODIUM. (*Swartz in act. stockholm. 1800, p. 217.*) *Brown mss.*

Cor. subringens: petala exteriora antica horizontalia concava. *Labellum* columnæ adnatum, bilabiatum. *Antheræ* loculis distantibus. *Stigma* adnatum.

1. *P. volucris*, caule triphylo, labello triangulari sagittato. *Willden. sp. pl. 4, p. 57.*

Nat. of the Cape of Good Hope. ۛ.

II. *Anthera* stigmati parallela persistens. *Massæ pollinis* v. *farinaceæ* v. e corpusculis angulatis; apicibus stigmati affixæ.

GOODYERA. *Brown mss.*

Cor. ringens : petalis exterioribus anticis labello inferne gibboso superne indiviso suppositis. *Columna libera. Pollen angulatum.*

1. *G. repens*, foliis radicalibus ovatis, labello petalisque lanceolatis. *Brown mss.*

Neottia repens. Willden. sp. pl. 4, p. 75.

Satyrium repens. Engl. bot. 289.

Nat. of Scotland. 4.

1. *G. pubescens*, foliis radicalibus ovatis, labello ovato acuminato, petalis ovatis. *Brown mss.*

Neottia pubescens. Willden. sp. pl. 4, p. 76.

Nat. of North America. 4.

NEOTTIA. (*Swartz in act. stockholm. 1800, p. 224.*)
Brown prodr. 319.

Cor. ringens : petalis exterioribus anticis labello imberbi suppositis; interioribus conniventibus. *Columna aptera. Pollen farinaceum.*

1. *N. speciosa*, labello lanceolato indiviso, scapo bracteato, bracteis flore longioribus, foliis oblongis margine apicem versus undulatis. *Willden. sp. pl. 4, p. 72. Andrews's reposit. 3. Botan. mag. 1374.*

Nat. of the West Indies. 4.

2. *N. orchioides*, foliis radicalibus lato-lanceolatis, spica erecta labello basi cum petalis saccato; lamina acuminata. *Willden. sp. pl. 4, p. 75. Botan. mag. 1036.*

Nat. of Jamaica. 4.

3. *N. picta*, foliis radicalibus oblongo-lanceolatis, scapo

bracteato, petalis anticis decurrentibus labello semiinfero ecalcarato suppositis.

Neottia acaulis. *Smith exot. bot.* 2, p. 91, t. 105. (varietas exscapa.)

Nat. of Trinidad. Mr. *Alexander Anderson.* 2.

4. *N. elata*, labello obovato emarginato, scapo vaginato, bracteis flore brevioribus, foliis ovatis petiolatis margine planis. *Willden. sp. pl.* 4, p. 72. *Redouté liliac.* 164.

Neottia minor. *Andrews's reposit.* 376.

Nat. of Guiana and the West Indies. 2.

5. *N. cernua*, foliis lanceolatis trinerviis, caule vaginato, floribus recurvato-cernuis, labello oblongo integerrimo acuto. *Willden. sp. pl.* 4, p. 75.

Nat. of North America. 2.

6. *N. spiralis*, foliis radicalibus oblongis subpetiolatis, spica tortili floribus secundis, labio ovato. *Willden. sp. pl.* 4, p. 73.

Orchrys spiralis. *Curtis lond. Engl. bot.* 541.

Nat. of Britain. 2.

PONTHIEVA. *Brown mss.*

Cor. irregularis. *Labellum* posticum, cum petalis interioribus columnæ insertum. *Pollen* farinaceum.

2001 1. *P. glandulosa*, labello unguiculato acuminato, petalis interioribus dimidiato-ovatis.

Neottia glandulosa. *Sims in botan. magaz.* 842.

Nat. of the West Indies. Mr. *Alexander Anderson.* 2.

DIURIS. (*Smith. in linn. soc. transact.* 4, p. 222.) *Brown prodr.* 315.

Cor. irregularis: petala 2 exteriora antica linearia *labelle* trifido supposita; interiora unguiculata patula. *Columna*: lobis lateralibus petaloideis. *Pollen* farinaceum.

1. *D. aurea*, labelli lacinia intermedia basi intus bicarinata lateralibus duplo longiore, petalis integerrimis: interioribus ellipticis acutis, foliis linearibus canaliculatis scapo brevioribus. *Brown prodr.* 315.

Diuris aurea. *Smith exot. bot.* 1, p. 15, t. 9.

Nat. of New South Wales. 4.

THELYMITRA. *Brown prodr.* 314.

Cor. regularis. *Labellum* sessile petalis subconforme. *Columna* cucullata: lobis lateralibus penicillatis nudisve. *Pollen* farinaceum.

1. *T. ixioides*, corolla patenti, cuculli laciniis lateralibus penicillatis; intermedia trifida: lobulis lateralibus apice erosis medio brevioribus bifido dorso cristato. *Brown prodr.* 314.

Thelymitra ixioides. *Willden. sp. pl.* 4, p. 79. *Smith exot. bot.* 1, p. 55, t. 29. (absque maculis saturatoribus corollæ.)

Nat. of New South Wales. 4.

LISTERA. *Brown mss.*

Cor. irregularis. *Labellum* bilobum. *Columna* aptera. *Anthera* basi inserta. *Pollen* farinaceum.

1. *L. ovata*, caule bifolio, foliis ovatis oppositis, columna postice cucullo anthera incumbente aucta.

Ophrys ovata. *Curtis lond. Engl. bot.* 1548.

Epipactis ovata. *Willden. sp. pl.* 4, p. 87.

Nat. of Britain. 4.

2. *L. cordata*, caule bifolio, foliis cordatis oppositis, labello inferne bidentato; lobis apicis lineari-subulatis. *Brown mss.*

Ophrys cordata. *Engl. bot.* 358.

Epipactis cordata. *Willden. sp. pl.* 4, p. 88.

Nat. of Britain. 4.

III. *Anthera terminalis*, inserta, persistens. *Massæ pol-
linis* v. *pulveræ* v. e. corpusculis angulatis : basi v. infra
apicem affixæ.

EPIPACTIS. *Brown mss.* Epipactidis species. *Swartz
in act. stockholm.* 1800, p. 231.

Labellum inferne ventricosum ; superne v. indivisum ; v.
3-lobum lobo medio majore articulatum connexo. *Pollen*
farinaceum.

1. *E. latifolia*, foliis ovatis amplexicaulibus ; bracteis
inferioribus flore longioribus, floribus pendulis, labello
integerrimo acuminato petalis brevioribus, germinibus pubes-
centibus. *Willden. sp. pl.* 4, p. 83.

Serapias latifolia. *Engl. bot.* 269. *Select specim.* 3.
Nat. of Britain. 4.

2. *E. palustris*, foliis lanceolatis amplexicaulibus, brac-
teis flore brevioribus, floribus pendulis, labello crenato ob-
tuso petalis æquali, germinibus pubescentibus. *Willden.*
sp. pl. 4, p. 84.

Serapias palustris. *Engl. bot.* 270.
Nat. of Britain. 4.

3. *E. pallens*, foliis oblongo-lanceolatis sessilibus, bracteis
flore longioribus, floribus erectis, labello obtuso petalis bre-
vioribus, germinibus glabris. *Willden. sp. pl.* 4, p. 85.

Serapias grandiflora. *Engl. bot.* 271.
Nat. of Britain. 4.

4. *E. ensifolia*, foliis lanceolatis acuminatis subdistichis,
bracteis minutissimis subulatis, floribus erectis, labello
obtuso petalis duplo brevioribus, germinibus glabris. *Willden.*
sp. pl. 4, p. 85.

Serapias ensifolia. *Engl. bot.* 494.
Nat. of Britain. 4.

5. *E. rubra*, foliis lanceolatis, bracteis germine longiori-

bus, floribus erectis, labello acuto lineis elevatis undulatis, germinibus glabris. *Willden. sp. pl. 4, p. 86.*

Serapias rubra. Engl. bot. 437.

Nat. of Britain. 4.

[208]

POGONIA. *Brown mss. Pogoniæ species. Juss. gen. 65.*

Labellum sessile, cucullatum, intus cristatum. Petala 5 distincta eglandulosa. Pollen farinaceum.

1. *P. divaricata*, radice subpalmata, scapi folio bracteaque lineari-oblongis, petalis exterioribus lanceolato-linearibus.

Arethusa divaricata. Willden. sp. pl. 4, p. 31. Lamarek illustr. t. 729, f. 3.

Nat. of North America. 4.

CALADENIA. *Brown prodr. 323.*

Labellum subunguiculatum, disco glandulis seriatis. Petala extus glandulosa: posticum planiusculum; quatuor antica plana labello supposita. Pollen farinaceum.

1. *C. alba*, petalis acutiusculis, columna labelloque exvittatis, glandulis labelli biseriatis superioribus arcuatis capitulo concolori; lobo medio longitudinaliter fimbriato disco eglanduloso. *Brown prodr. 323.*

Nat. of New South Wales. Robert Brown. 4.

GLOSSODIA. *Brown prodr. 325.*

Appendix inter labellum eglandulosum et columnam. Petala 5 patula æqualia. Pollen farinaceum.

1. *G. major*, appendice semibifida: lobis patentibus acutis. *Brown prodr. 326.*

Nat. of New South Wales. Robert Brown. 4.

PTEROSTYLIS. *Brown prodr.* 326.

Labellum unguiculatum: lamina basi appendiculata v. gibbosa. *Cor.* ringens petalis anticis inferne connatis et ungui labelli accretis. *Columna* superne alata. *Pollen* farinaceum.

1. *P. obtusa*, caule folioso unifloro lamina obtusa apice-haud angustata, appendice penicillata. *Brown prodr.* 327.
Nat. of New South Wales. *Robert Brown.* 4.

CALEYA. (Caleana.) *Brown prodr.* 329.

Labellum unguiculatum posticum: lamina peltata cava, foramine exteriore. *Pollen* farinaceum

1. *C. major*, folio lanceolato-lineari plano, scapo medio unibracteato, lamina labelli lævi: utraque extremitate angustata semiovata. *Brown prodr.* 329.
Nat. of New South Wales. *Robert Brown.* 4.

CALOPOGON. *Brown mss.*

Labellum posticum, unguiculatum: lamina barbata. *Petala* 5 distincta. *Columna* libera. *Pollen* angulatum.

1. *CALOPOGON pulchellus.* *Brown mss.*
^{205]} *Cymbidium pulchellum.* *Willden. sp. pl.* 4, p. 105.
Limodorum tuberosum. *Curtis magaz.* 116.
Nat. of North America. 4.

ARETHUSA. *Brown mss.* *Arethusa* species, *Swartz in act. stockholm.* 1800, p. 230.

Labellum inferne cum columna connatum; superne cucullatum; intus cristatum. *Petala* 5 basi connata. *Pollen* angulatum.

1. *ARETHUSA bulbosa*.

Arethusa bulbosa. *Willden. sp. pl.* 4, p. 80.

Helleborine mariana monanthos, flore longo purpurascens liliaceo. *Pluk. mant.* 100, t. 348, f. 7.

Nat. of North America. 4 . .

IV. *Anthera terminalis mobilis decidua. Massæ pollinis demum cereaceæ.*

BLETIA. (*Ruiz et Pavon prodr.* 119.) *Brown mss.*

Labellum sessile, cucullatum; nunc basi calcaratum. *Petala* 5 distincta. *Columna* libera. *Massæ pollinis*, 8 v. 4 bilobæ.

1. *B. Tankervilliae*, labello calcarato indiviso: cornu abbreviato, foliis radicalibus ovato-lanceolatis. *Brown mss.*

Limodorum Tankervilliae. [*Hort. Kew.*] *Edit. prior.* 3, p. 302, t. 12. *Willden. sp. pl.* 4, p. 122. *Andrews's reposit. Redouté liliac.* 43. *Schneevooght ic.* 5.

Nat. of China. 4 .

2. *B. verecunda*, labello ecalcarato: costis disci ⁽²⁰⁶⁾ ramosis; lobo medio latiore quam longo; lateralibus superne angustioribus, scapo subramoso. *Brown mss.*

Cymbidium verecundum. *Willden. sp. pl.* 4, p. 105.

Limodorum altum. *Botan. magaz.* 930.

Nat. of the West Indies. 4 .

3. *B. florida*, labello ecalcarato: costis disci simplicibus; lobo medio subcuneiformi; lateralibus apice latioribus, scapo subramoso. *Brown mss.*

Cymbidium floridum. *Salisb. prodr.* 9.

Limodorum purpureum. *Redouté liliac.* 83.

Nat. of the West Indies. 4 .

4. *B. hyacinthina*, labello ecalcarato imberbi, massis pollinis quatuor bilobis, caule folioso, floribus racemosis. *Brown mss.*

Cymbidium hyacinthinum. *Smith exot. bot.* 1, p. 117, t. 60. *Botan. magaz.* 1492.

Nat. of China. Mr. *William Ker.* 4.

5. *B. capitata*, labello ecalcarato: prope basin intus callo adnato, caule folioso, floribus capitatis. *Brown mss.*

Nat. of the West Indies. 4.

207] GEODORUM. (*Jackson in Andrews's reposit.* 626.)

Brown mss.

Labellum cucullato-ventricosum (nunc basi calcaratum,) sessile, cum columna non articulatum. *Petala* conformia subsecunda. *Massæ pollinis* 2, postice lobulo auctæ.

1. *G. purpureum*, scapo foliis longiore, racemo pendulo: floribus alternis, labello ovato acuto picto. *Brown mss.*

Malaxis nutans. *Willden. sp. pl.* 4, p. 93.

Limodorum nutans. *Roxb. corom.* 1, p. 33, t. 40.

Nat. of the East Indies. *William Roxburgh, M.D.* 4.

2. *G. citrinum*, scapo foliis brevior, spica pendula: floribus congestis, labello basi subcalcarato; apice obtuso integerrimo. *Brown mss.*

Geodorum citrinum. *Andrews's reposit.* 626.

Nat. of the East Indies. *William Roxburgh, M.D.* 4.

3. *G. dilatatum*, scapo foliis brevior, spica pendula: floribus congestis, labello subcalcarato; apice dilatato crenulato. *Brown mss.*

Limodorum recurvum. *Willden. sp. pl.* 4, p. 124. *Roxb. corom.* 1, p. 33, t. 39.

Nat. of the East Indies. *William Roxburgh, M.D.* 4.

208] CALYPSO. (*Salisb. paradis.* 89.) *Brown mss.*

Labellum ventricosum prope apicem subtus calcaratum.

Petala adscendentia secunda. *Columna* petaloideo-dilatata.
Massæ pollinis 4.

1. *C. americana*, labello basi angustato subunguiculato :
calcare semibifido laminam superante dentibus acutis, pe-
dunculo germine longiore.

Calypso borealis. *Salisb. paradis.* 89. (exclusis, ni fallor,
synonymis.)

Nat. of North America. 4.

MALAXIS. (*Swartz in act. stockholm.* 1800, p. 233.)
Brown mss.

Labellum explanatum indivisum sessile (sæpius posticum).
Petala 5 angustiora, patula v. deflexa. *Massæ pollinis*
4 parallelæ, apicibus stigmati affixæ.

1. *M. paludosa*, foliis subquaternis apice scabris, scapo
pentagono, labello concavo acuto. *Willden. sp. pl.* 4, p. 91.
Engl. bot. 72.

Nat. of England. 4.

2. *M. Loeselii*, foliis binis ovato-lanceolatis, scapo trigono,
labello apice ovato recurvato. *Willden. sp. pl.* 4, p. 93.

Ophrys Loeselii. *Engl. bot.* 47.

Nat. of England. 4.

3. *M. lilifolia*, foliis binis ovato-lanceolatis, scapo tri-
quetro, petalis interioribus reflexis discoloribus, labello
concavo obovato apice acuto. *Willden. sp. pl.* 4, p. 92. [209

Ophrys lilifolia. *Andrews's reposit.* 65.

Nat. of North America 4.

CORALLORRHIZA. (*Hall. hist.* 2, p. 159.) *Brown mss.*

Labellum basi productum : calcare adnato v. libero.
Columna libera. *Massæ pollinis* 4 obliquæ (nec parallelæ).

1. *C. innata*, calcare abbreviato omnino adnato.
Cymbidium Corallorhizon. *Willden. sp. pl.* 4, p. 109.
Ophrys corallorhiza. *Engl. bot.* 1547.
Nat. of Scotland. 4.

ISOCHILUS. *Brown mss.*

Labellum petalis distinctis conniventibus subconforme.
Massæ pollinis 4, parallelæ.

1. *I. linearis*, spica terminali, foliis distichis linearibus
obtusis emarginatis, caule simplici.
Cymbidium lineare. *Willden. sp. pl.* 4, p. 97.
Epidendrum lineare. *Jacqu. amer.* 221, t. 131, f. 1.
Nat. of the West Indies. 4.

2. *I. prolifer*, floribus axillaribus, foliis distichis lanceo-
lato-oblongis, caule prolifero; bulbis axillaribus diphyllis.
²¹⁰ *Cymbidium* proliferum. *Willden. sp. pl.* 4, p. 95.
Nat. of the West Indies. 4.

ORNITHIDIUM. (*Salisb. in hort. soc. transact.* 1, p. 293, absque caractere.) *Brown mss.*

Labellum sessile cucullatum cum basi columnæ connatum.
Petala conniventia. *Massæ pollinis* 4, obliquæ, postice
sulcatæ.

1. *ORNITHIDIUM coccineum.*
Ornithidium coccineum. *Salisb. in hort. soc. transact.* 1,
p. 293.
Cymbidium coccineum. *Willden. sp. pl.* 4, p. 94. *Botan.*
magaz. 1437.
Nat. of the West Indies. 4.

109.

STELIS. (*Swartz in Schrader's journal* 1799, 2 band, p. 239.) *Brown mss.*

Labellum petalis interioribus nanis s abfornicatis conforme.
Petala 3 exteriora basi connata. *Massæ pollinis* 2.

forme.

1. *S. ophioglossoides*, caule unifolio, folio oblongo-lanceolato racemum subæquante, floribus trigonis. *Willden. sp. pl.* 4, p. 138.

earibus

Epidendrum ophioglossoides. Jacqu. amer. 225, t. 133, f. 2.

Nat. of the West Indies. 4.

1.

2. *S. micrantha*, caule unifolio elongato, folio lato-lanceolato racemo brevior, floribus hexagonis. *Willden.* [21] *sp. pl.* 4, p. 139.

lanceo-
yllis.

Stelis micrantha. Smith exot. bot. 2, p. 31, t. 75.

Nat. of Jamaica. 4.

5.

PLEUROTHALLIS. *Brown mss.*

. 293,

Labellum articulatum connexum cum basi simplici v. brevissime producta columnæ. *Petala* 2 antica exteriorum inferne connata. *Massæ pollinis* 2, exsulcæ.

atum.

ostice

1. *P. ruscifolia*, caule elongato unifolio, folio ovato-lanceolato, floribus in sinu folii aggregatis. *Brown mss.*

Dendrobium ruscifolium. Willden. sp. pl. 4, p. 135.

Epidendrum ruscifolium. Jacqu. amer. 226, t. 133, f. 3.

Nat. of the West Indies. 4.

act. 1,

Botan.

OCTOMERIA. *Brown mss.*

Labellum articulatum cum processu unguiformi, cujus lateribus petala antica adnata. *Massæ pollinis* 8.

1. *O. graminifolia*, caule elongato unifolio, folio lanceolato, pedunculis geminatis unifloris, radice repente.

Dendrobium graminifolium. Willden. *sp. pl.* 4, p. 135.
Helleborine graminea repens biflora. Plum. *ic.* 176, f. 1.
Nat. of the West Indies. 4.

212] *AERIDES.* (Swartz in Schrader's journal 1799,
 2 band, p. 233.) Brown mss.

Labellum calcaratum v. saccatum, insertum apice processus unguiformis, cujus lateribus petala antica exteriorum adnata. *Massæ pollinis* 2 postice bilobæ, processu communi stigmatis medio affixæ.

1. *A. odoratum*, calcare adscendente conico-subulato, labelli lobo medio lateralibus brevior, foliis retusis.
Aerides odoratum. Willden. *sp. pl.* 4, p. 131.
Nat. of China. 4.

DENDROBIUM. (Swartz in nov. act. upsal. 6, p. 82.)
 Brown mss.

Labellum ecalcaratum, articulatum cum apice processus unguiformis, cujus lateribus petala antica adnata, calcar æmulantia. *Massæ pollinis* 4, parallelæ.

1. *D. speciosum*, caulibus erectis apice 2-3-phyllis, foliis ovali-oblongis racemo terminali multifloro brevioribus, petalis angusto-oblongis, labello infra divisuram carinia unica; lobo intermedio ecarinato dilatato. *Brown prodr.* 332.

Dendrobium speciosum. Smith *exot. bot.* 1, p. 17, t. 10.
Nat. of New South Wales. 4.

2. *D. linguiforme*, caulibus repentibus, foliis ovalibus obtusis depressis carnosissimis racemo aliquoties brevioribus, 213] petalis elongato-linearibus acutis, labelli lobo medio undulato tricarinato. *Brown prodr.* 333.

Dendrobium linguiforme. Smith *exot. bot.* 1, p. 19, t. 11. Willden. *sp. pl.* 4, p. 138.
Nat. of New South Wales. 4.

3. *D. Barringtoniæ*, foliis subternis oblongis nervosis bulbo innatis, scapo subunifloro vaginato. *Willden. sp. pl. 4, p. 132.*

Epidendrum Barringtoniæ. Smith ic. pict. 25.
Nat. of the West Indies. 2.

CYMBIDIUM. *Brown mss.* Cymbidii species. *Swartz in nov. act. upsal. 6, p. 70.*

Labellum ecalcaratum concavum, cum basi (simplici nec producta) columnæ articulatum. *Petala* patentia distincta. *Massæ pollinis* 2, postice bilobæ.

* *Cymbidia vera.*

1. *C. aloifolium*, foliis radicalibus lato-linearibus canaliculatis carnosius apice retusis, scapis multifloris pendulis.

Cymbidium aloifolium. Willden. sp. pl. 4, p. 101.

Epidendrum aloifolium. Curtis magaz. 387. Redouté liliac. 144.

Nat. of the East Indies. 2.

2. *C. ensifolium*, foliis radicalibus ensiformibus nervosis, scapo tereti paucifloro, labello ovato sub-recurvato maculato. *Willden. sp. pl. 4, p. 110.*

Epidendrum ensifolium. Smith spicil. 22, t. 24, [214 Andrews's reposit. 344.]

Epidendrum sinense. Redouté liliac. 113.

Nat. of China and Japan. 2.

3. *C. sinense*, foliis radicalibus ensiformibus nervosis, scapo paucifloro, floribus secundis, petalis striatis: tribus exterioribus reflexis, labello oblongo obtuso reflexo. *Willden. sp. pl. 4, p. 111.*

Epidendrum sinense. Andrews's reposit. 216. Botan. magaz. 888.

Nat. of China. 2.

4. *C. præmorsum*, caulescens, foliis remotiusculis distichis

lato-linearibus canaliculatis apice præmorsis, spicis oppositifoliis. *Willden. sp. pl.* 4, p. 103.

Epidendrum præmorsum. *Roxb. corom.* 1, p. 34, t. 43.

Nat. of the East Indies. *William Roxburgh, M.D.* 4.

** *Cymbidium spurium.*

5. *C. tripterum*, acaule, foliis bulbo innatis radicalibus vaginatis, scapis multifloris, germine trialato. *Willden. sp. pl.* 4, p. 94.

Epidendrum tripterum. *Smith ic. pict.* 14.

Nat. of Jamaica. 4.

215]

BRASSIA. *Brown mss.*

Labellum explanatum indivisum. *Petala* patula distincta. *Columna* aptera. *Massæ pollinis* 2, postice bilobæ; medio affixæ processu communi stigmatis.

1. *BRASSIA maculata.* *Brown mss.*

Nat. of Jamaica. 4.

ONCIDIUM. (*Swartz in act. stockholm.* 1800, p. 239.)
Brown mss.

Labellum explanatum lobatum basi tuberculatum. *Petala* patentia (2 antica nunc connata). *Columna* alata. *Massæ pollinis* 2, postice bilobæ; medio affixæ processu communi stigmatis.

1. *O. altissimum*, petalis lanceolatis labello longioribus: anticis distinctis, scapo paniculato.

Oncidium altissimum. *Willden. sp. pl.* 4, p. 112.

Epidendrum altissimum. *Jacqu. amer.* 229, t. 141.

Nat. of the West Indies. 4.

2. *O. carthagenense*, petalis obovatis unguiculatis labello parum brevioribus anticis distinctis scapo paniculato.

Oncidium carthagenense. Willden. *sp. pl.* 4, p. 112.

Epidendrum undulatum. *Botan. magaz.* 777.

Nat. of the West Indies. 4.

3. *O. bifolium*, petalis obovatis undulatis : anticis connatis, labello petalis longiore : lobo medio dilatato-reniformi semibifido, scapo racemoso, bulbis diphyllis.

Oncidium bifolium. Sims in *Botan. magaz.* 1491.

Nat. of South America. 4.

4. *O. triquetrum*, petalis acutis : anticis connatis, labelli lobo medio subrotundo indiviso, scapo racemoso, foliis triquetris.

Cymbidium triquetrum. Willden. *sp. pl.* 4, p. 104.

Nat. of Jamaica. 4.

CYRTOPODIUM. *Brown mss.*

Labellum ungue geniculato ; lamina 3-partita. *Petala* 5 distincta. *Massæ pollinis* 2 postice bilobæ.

1. CYRTOPODIUM *Andersonii.* *Brown mss.*

Cymbidium Andersonii. Lambert in *Andrews's reposit.* 651.

Nat. of the West Indies. Mr. Alex. Anderson. 4.

BRASSAVOLA. *Brown mss.*

Labellum ungue simplici ; lamina indivisa. *Petala* distincta patula. *Massæ pollinis* 8 (vel plures!).

1. *B. cucullata*, caule unifloro, lamina labelli ciliata.

Cymbidium cucullatum. Willden. *sp. pl.* 4, p. 100.

Epidendrum cucullatum. *Botan. magaz.* 543.

Nat. of the West Indies. 4.

BROUGHTONIA. *Brown mss.*

Columna libera v. basi tantum connata labello unguiculato (nunc inferne producto in tubulum ovario adnatum). Massæ pollinis 4 parallelæ, septis completis persistentibus distinctæ, basi filo granulato elastico auctæ.

1. *B. sanguinea*, foliis geminis oblongis bulbo innatis, scapo diviso. *Brown mss.*

Dendrobium sanguineum. Willden. sp. pl. 4, p. 132.

Viscum radice bulbosa minus, delphinii flore rubro specioso. Sloan. jam. 1, p. 250, t. 121, f. 2.

Nat. of Jamaica. 4.

EPIDENDRUM. (*Swartz in nov. act. upsal. 6, p. 66.*)

Brown mss.

Columna cum ungue labelli longitudinaliter connata in tubum (quandoque decurrentem ovarium). Massæ pollinis 4 parallelæ, septis completis persistentibus distinctæ, basi filo granulato elastico auctæ.

1. *E. cochleatum*, foliis geminis oblongis bulbo innatis, scapo elongato, lamina labelli cordata obtusa. *Willden. sp. pl. 4, p. 114. Andrews's reposit. 13. Botan. magaz. 572.*

Nat. of the West Indies. 4.

2. *E. fragrans*, folio lanceolato bulbo innato, scapo^[218] abbreviato multifloro, lamina labelli cordata acuminata.

Epidendrum fragrans. Willden. sp. pl. 4, p. 114. Andrews's reposit. 645.

Epidendrum cochleatum. Curtis magaz. 152.

Nat. of Jamaica. 4.

3. *E. secundum*, caule simplici, foliis oblongis emarginatis, pedunculo terminali longissimo, spica laxa secunda, columna longitudine petalorum. *Willden. sp. pl. 4, p. 119.*

Epidendrum secundum. *Jacq. amer.* 224, t. 137.

Nat. of the West Indies. ۲.

4. *E. fuscatum*, caule simplici, foliis oblongis acuminate, pedunculo terminali elongato, spica globosa, columna petalis brevior. *Willden. sp. pl.* 4, p. 120. *Smith spicil.* 21, t. 23.

Nat. of the West Indies. ۲.

5. *E. elongatum*, caule simplici, foliis oblongis, pedunculo terminali elongato, spica laxa, lamina labelli dentato-ciliato. *Willden. sp. pl.* 4, p. 120. *Botan. magaz.* 611.

Nat. of the West Indies. ۲.

6. *E. umbellatum*, caule simplici, foliis oblongis subemarginatis, floribus in sinu folii terminalis confertis, lamina labelli triloba: lobo intermedio emarginato. *Willden. sp. pl.* 4, p. 117.

Nat. of Jamaica. ۲.

7. *E. nutans*, caule simplici, foliis ovato-lanceolatis amplexicaulibus, floribus suspicatis nutantibus, lamina labelli triloba: lobo intermedio tridentato. *Willden. sp. pl.* 4, p. 117.

Nat. of Jamaica. *Arthur Broughton, M.D.* ۲.

8. *E. conopseum*, caule simplici, floribus spicatis erectis, lamina labelli triloba: medio retuso, petalis interioribus angustioribus, foliis lanceolatis.

Nat. of Florida. *Mr. William Bartram.* ۲.

9. *E. ciliare*, caule simplici, foliis binis oblongis aveniis, lamina labelli tripartita: lacinia intermedia lineari. *Willden. sp. pl.* 4, p. 119. *Botan. magaz.* 463. *Redouté liliac.* 82.

Nat. of the West Indies. ۲.

VANILLA (*Swartz in nov. act. upsal.* 6, p. 66.) *Brown mss.*

Cor. cum germine articulata decidua! Labellum inferne

^{220]} columnna connatum. *Massæ pollinis granulosæ. Capsula* carnosa. *Semina* globosa aptera.

1. *V. aromatica*, foliis ovato-oblongis nervosis, petalis unilatis, labello acuto, capsulis cylindraceis longissimis.

Vanilla aromatica. Willden. sp. pl. 4, p. 121.

Vanilla flore viridi et albo, fructu nigricante. *Plum. ic. 183, t. 188.*

Nat. of South America. 7.

2. *V. planifolia*, foliis oblongo-lanceolatis planis obsolete striatis, labello retuso.

Vanilla planifolia. Andrews's reposit. 538.

Myrobroma fragrans. Salisb. paradis. 82.

Nat. of the West Indies. 4.

DIANDRIA.

CYPRIPEDIUM. (*Swartz in act. stockholm. 1800, p. 250.*) *Brown mss.*

Labellum ventricosum, inflatum (nunc saccatum). *Columnna* postice terminata lobo petaloideo (stamine sterili), antheras distinguente. *Petala 2 antica sæpius connata.*

1. *C. Calceolus*, caule folioso, lobo columnæ elliptico obtuso, labello petalis brevior compresso. *Willden. sp. pl. 4, p. 142. Engl. bot. 1. Fl. dan. 999. Redouté liliac. 19.*

Nat. of England. 4.

^{221]} 2. *C. parviflorum*, caule folioso, lobo columnæ triangulari acuto, labello petalis brevior compresso. *Willden. sp. pl. 4, p. 143.*

Cypripedium parviflorum. Botan. magaz. 911.

Nat. of North America. 4.

3. *C. pubescens*, caule folioso, lobo styli triangulari-ob-

longo-obtus, labello petalis brevior compresso. *Willden. sp. pl. 4, p. 143.*

Cypripedium pubescens. Willden. hort. berlin. 13.

Cypripedium flavescens. Redouté liliac. 20.

Nat. of North America. 4.

4. *C. spectabile*, caule folioso, lobo styli elliptico cordato obtuso, labello petalis obtusis longior antice fissus. *Willden. sp. pl. 4, p. 143.*

Cypripedium album. Curtis magaz. 216. Schnevooght ic. 2.

Nat. of North America. 4.

5. *C. humile*, scapo aphylo unifloro, foliis radicalibus geminis oblongis obtusis, lobo columnæ subrotundo-rhombeo acuminato deflexo, labello petalis lanceolatis longior antice fissus. *Willden. sp. pl. 4, p. 144.*

Cypripedium acaule. Curtis magaz. 192.

Nat. of North America. 4.

6. *C. arietinum*, corollis pentapetalis, labello saccato-calcarato, caule folioso.

Nat. of North America. 4.

GENERA ET SPECIES

PLANTARUM E VARIIS FAMILIIS,

QUÆ

IN HORTO KEWENSI COLUNTUR.

[*Extracted from 'Hortus Kewensis,' 2nd Edition, Vol. III, IV, & V.*]

LONDON.

—
1811—1813.

PLANTÆ VARLÆ HORTI KEWENSIS.

ANISOMELES. *Brown prodr.* 503.

[iii. 364]

Cal. 5-fidus. *Cor.* labium superius abbreviatum, integrum; inferius trifidum: lacinia media biloba. *Stam.* exserta, approximata: *Antheræ breviorum* biloculares, lobis parallelis; *longiorum* dimidiatæ v. dissimiles.

1. *A. ovata*, foliis ovatis subcordatis crenatis, verticillis multifloris, bracteis linearibus, calycibus pilosis: glandulis inconspicuis.

Nepeta indica. *Linn. sp. pl.* 799 (exclusis synonymis). *Willden. sp. pl.* 3, p. 57.

Ballota disticha. [*Hort. Kew.*] *Edit. prior.* 2, p. 304. *Linn. mant.* 83?

Marrubium odoratissimum *betonicae* folio. *Burm. zeyl.* 153, t. 71, f. 1.

Nat. of the East Indies. ☉.

PYCNANTHEMUM. *Michaux amer.* 2, p. 7. [iii. 376]

Brachystemum. *Michaux amer.* 2, p. 5.

Cal. 5-fidus. *Cor.* labium inferius lacinia media longiore *Stam.* distantia: antherarum loculis parallelis. *Involucrum* capituli multibracteatum.

1. *P. incanum*, staminibus exsertis, foliis oblongo-ovatis acutis subseratis tomentosis, capitulis compositis: lateralibus pedunculatis.

Pycnanthemum incanum. *Michaux amer.* 2, p. 7.

Clinopodium incanum. *Willden. sp. pl.* 3, p. [132].

Clinopodium menthæ folio incanum et odoratum. *Dill. elth.* 87, t. 74, f. 85.

Nat. of North America. 4.

iii. 377] 2. *P. aristatum*, staminibus exsertis, foliis lanceolatis, capitulis sessilibus, bracteis aristatis.

Pycnanthemum aristatum. *Michaux amer.* 2, p. 8, t. 33.

Nepeta virginica. *Willden. sp. pl.* 3, p. 56.

Nat. of North America. 4.

3. *P. virginicum*, staminibus inclusis, foliis lanceolatis linearibusve integerrimis, capitulis subcorymbosis.

Brachystemum virginicum. *Michaux amer.* 2, p. 6.

Thymus virginicus. *Willden. sp. pl.* 3, p. 145.

Satureja virginiana, floribus in summitate. *Herm. parad.* 218, cum tab.

Nat. of North America. 4.

CAPRARIA. *Gen. pl.* 1030.

iv. 46] 1. *C. biflora*, glabra, foliis alternis lanceolatis serratis, floribus geminis, calycis laciniis subulatis, corollæ fauce imberbi.

Capraria biflora. *Willden. sp. pl.* 3, p. 323. *Jacqu. amer.* 182, t. 115.

Nat. of South America. ½.

2. *C. cuneata*, pilosa, foliis alternis rhombeo-cuneiformibus inciso-serratis, floribus geminis, calycis laciniis linearibus: pilis capitatis, corollæ fauce imberbi.

Erinus frutescens. *Mill. dict. ed.* 8.

Nat. of South America. ½.

iv. 52] STEMODIA. *Gen. pl.* 1043.

Cal. 5-part. *Cor.* bilabiata. *Antheræ* lobis distantibus. *Caps.* bilocularis.

1. *S. parviflora*, procumbens ramosissima pubescens, foliis ternatis petiolatis ovatis crenatis.

Erinus verticillatus. *Mill. dict. ed. 8.*
Nat. of South America. ½.

MAZUS. *Lour. cochinch.* 385. *Brown prodr.* 439. [iv. 53]

Cal. 5-fid. campanulatus. *Cor.* ringens: labio superiore semibilobo; inferiore 3-fido, basi bigibboso. *Caps.* biloc. bivalvis: valvis integris, medio septiferis.

1. *M. rugosus*, racemo laxo caulem paucifolium superante, calycibus pubescentibus: fructiferis auctis.

Mazus rugosus. *Lour. cochinch.* 385.

Lindernia japonica. *Willden. sp. pl.* 3, p. 326.

Nat. of China and Japan. ☉.

CROSSANDRA. *Salish. paradis.* 12. *Brown prodr.* [iv. 54]
 475, in obs.

Cal. 5-part. inæqualis. *Cor.* unilabiata. *Stam.* inclusa: *Antheris* unilocularibus. *Caps.* biloc. bivalvis: dissepimento contrario. *Semina* retinaculis subtensa.

1. *CROSSANDRA undulæfolia.* *Salish. paradis.* 12.

Ruellia infundibuliformis. *Andrews's reposit.* 542.

Justicia infundibuliformis. *Willden. sp. pl.* 1, p. 99.

Nat. of the East Indies. ½.

APHELANDRA. *Brown prodr.* 475, in obs. [iv. 55]

Cal. 5-part. inæqualis. *Cor.* bilabiata. *Antheræ* uniloculares! *Caps.* biloc. bivalvis: dissepimento contrario. *Semina* retinaculis subtensa.

1. *A. cristata*, foliis ellipticis oblongisve acuminatis, spicis tetragonis, bracteis ovatis integerrimis, corollis glabris.

Justicia cristata. *Jacqu. hort. schænbr.* 3, p. 38, t. 320.

Justicia pulcherrima. *Willden. sp. pl.* 1, p. 86.

Justicia tetragona. *Willden. sp. pl.* 1, p. 85.

Justicia arborea. *Mill. dict. ed. 8.*

Ruellia cristata. *Andrews's reposit. 506.*

Nat. of the West Indies. ½.

BLECHUM. *Brown prodr. 478, in obs. Blechi species.*
Jussieu in annales du mus. 9, p. 269.

Cal. 5-part. *æqualis.* *Cor.* infundibuliformis. *Caps.* subbilocularis, bivalvis: dissepimenti contrarii segmentis demum liberis. *Semina* plura, retinaculis subtensa.

1. *B. Brownei*, foliis ovato-ellipticis subdentatis, spicis tetragonis, bracteis ovatis pubescentibus.

Blechum Brownei. *Jussieu in annales du mus. 9, p. 270.*

Ruellia Blechum. *Willden. sp. pl. 3, p. 362.*

Frunella elatior, flore albo. *Sloan. jam. 1, p. 173, t. 109, f. 1.*

Nat. of the West Indies. ¼.

iv. 61] **BONTIA.** *Brown prodr. 517, in obs.*

Cal. 5-part. *Cor.* ringens: labio superiore emarginato; inferiore trifido. *Stigma* bilobum. *Drupa* putamine biloculari: loculis bipartitis 4-spermis.

1. *BONTIA daphnoides.* *Willden. sp. pl. 3, p. 394.*

Bontia laureolæ facie. *Dill. elth. 57, t. 49, f. 57.*

Nat. of the West Indies. ½.

iv. 62] **VOLKAMERIA.** *Gen. pl. 1056.*

Cal. 5-fidus. *Cor.* tubo cylindrico; limbo 5-part. laciniis æqualibus subsecundis. *Stam.* exserta adscendentia. *Bacca* dipyrenea: ossiculis bilocularibus.

1. *VOLKAMERIA aculeata.*

Volkameria aculeata. *Willden. sp. pl. 3, p. 383.*

Clerodendrum fruticosum spinosum, foliis inferioribus confertis, superioribus oppositis; pedunculis tripartitis trifloris alaribus. *Brown jam.* 262, t. 20, f. 2.

Nat. of the West Indies. $\frac{1}{2}$.

Obs. Genus vix distinctum a *Clerodendro*.

CLERODENDRUM. *Brown prodr.* 510.

Cal. 5-fid. (nunc 5-dent.) *Cor.* tubo cylindrico; limbo [iv. 68] 5-partito patenti laciniis subæqualibus. *Stam.* juxta faucem inserta, exserta, adscendentia: antherarum loculis parallelis. *Bacca* pyrenis 4 monospermis.

1. *C. fragrans*, tomentosum, foliis subcordatis dentatis basi glandulosis, corymbis terminalibus densis hemisphæricis. *Venten. malmais.* 70 (floribus simplicibus.)

Volkmania japonica. *Jacqu. hort. schænbr.* 3, p. 48, t. 338 (floribus plenis).

Nat. of Japan and China. $\frac{1}{2}$.

2. *C. viscosum*, tomentosum, foliis cordatis dentatis, corymbo terminali paniculato, calycibus viscosis laxis dimidio tubi corollæ longioribus.

Clerodendrum viscosum. *Venten. malmais.* 25.

Peragu. *Rheed. mal.* 2, p. 41, t. 25.

Nat. of the East Indies. *William Roxburgh*, M.D. $\frac{1}{2}$.

3. *C. squamatum*, foliis cordatis denticulatis subtus squamatis, panicula terminali divaricata, pedunculis calycibusque coloratis glabris, staminibus corolla tota longioribus.

Clerodendrum squamatum. *Willden. sp. pl.* 3, p. 385.

Volkameria Kæmpferi. *Willden. sp. pl.* 3, p. 385.

Volkameria Kæmpferiana. *Jacqu. ic.* 3, t. 500.

Nat. of China and Japan. $\frac{1}{2}$.

4. *C. paniculatum*, foliis cordatis quinquelobis subdenticulatis glabris; summis sæpius indivisis, panicula [iv. 64] brachiata, corollæ tubo calycem multoties superante.

Clerodendrum paniculatum. Willden. *sp. pl.* 3, p. 388.

Clerodendrum pyramidale. Andrews's *reposit.* 628.

Nat. of Java and Pulo Pinang. h.

5. *C. tomentosum*, foliis ellipticis acutis integris, calycibusque tomentosis: fructiferis auctis incrassatis coloratis, corymbis congestis. Brown *prodr.* 510. Andrews's *reposit.* 607.

Volkameria tomentosa. Venten. *malmais.* 84?

Nat. of New South Wales. h.

6. *C. ligustrinum*, foliis elliptico-lanceolatis integerrimis, petiolis pedunculis calycibusque (5-fidis) hirsutis, corymbis axillaribus, corollæ tubo limbum vix æquante.

Volkameria ligustrina. Willden. *sp. pl.* 3, p. 383. Jacq. *collect. suppl.* 117, t. 5, f. 1.

Nat. of the Island of Mauritius. h.

7. *C. heterophyllum*, foliis lanceolatis lineari-lanceolatisve integerrimis glaberrimis, corymbis axillaribus terminalibusque, calycibus 5-dentatis pedunculisque glabris, corollæ tubo limbum vix æquante.

Volkameria heterophylla. Poiret in *encycl. botan.* 8, p. 687.

Volkameria angustifolia. Andrews's *reposit.* 554.

Nat. of the Island of Mauritius. h.

iv. 66] 8. *C. inerme*, foliis ovatis ovalibusve integerrimis calycibusque (5-dentatis) glabris, corymbis axillaribus, corollæ tubo limbum aliquoties superante. Brown *prodr.* 514.

Volkameria inermis. Willden. *sp. pl.* 3, p. 383. Jacq. *collect. suppl.* 117, t. 4, f. 1.

Nat. of the East Indies. h.

9. *C. Siphonanthus*, foliis verticillatis elongato-lanceolatis integris glabris, corymbis axillaribus paucifloris, corollis longissimis.

Siphonanthus indica. Willden. *sp. pl.* 1, p. 606.

388. *Siphonanthus angustifolia*. Willden. *sp. pl.* 1, p. 606.
Ovieda mitis. Willden. *sp. pl.* 3, p. 382. *Burm. ind.* 136,
t. 43, *f.* 1.
Nat. of the East Indies. William Roxburgh, M.D. $\frac{1}{2}$.

MELHANIA. *Forsk. descr.* 64.

[iv. 146]

Cal. duplex: exterior 3-phyllus (nunc caducus). *Cor.*
 5-petala. *Filamenta* 5 alterna sterilia. *Capsula* 5-locul.
polysperma: dissepimentis contrariis.

1. *M. Erythroxyton*; calyce exteriori caduco; interiore
 intus sericeo petalis dimidio brevioribus, foliis cordatis crenato-
 repandis: subtus reticulatis tomentosis.

Pentapetes Erythroxyton. [*Hort. Kew.*] *Edit. prior* 2,
p. 438.

Dombeya Erythroxyton. Willden. *sp. pl.* 3, p. 725.
Botan. Magaz. 1000.

Nat. of the Island of St. Helena. $\frac{1}{2}$.

2. *M. Melanoxyton*, calyce exteriori caduco; interiore
 intus glabro petalis parum brevioribus, foliis cordatis integer-
 rimis utrinque tomentosis lævibus.

*Alcea arbor populnea fronde tota argentea quinque-
 capsularis, seu Ebenus viridis.* *Pluk. mant.* 6, *t.* 333, *f.* 5.

Nat. of the Island of St. Helena. $\frac{1}{2}$.

ABROMA. *Gen. pl.* 1217.

[iv. 409]

Cal. 5-part. *Petala* 5 unguibus dilatatis saccatis. *Sta-
 minum urceolus* 10-fidus: laciniis 3-antheriferis; 5 alternis
 petaloideis. *Styli* 5. *Caps.* 3-loc. 5-alata, polysperma.

1. *A. augusta*, ramis tomentosis lævibus, foliis adultis
 subtus pube simplicissima, capsulæ alis apice truncatis:
 angulo exteriori acutiusculo.

Abroma augusta. Willden. *sp. pl.* 3, p. 1424.

Abroma Wheleri. Willden. *sp. pl.* 3, p. 1425.

Abroma fastuosum. *Jacqu. hort. vindob.* 3, p. 3, *t.* 1.

Nat. of the East Indies. $\frac{1}{2}$.

2. *A. fastuosa*, ramis muricatis, foliis adultis subtus pube simplici et stellari, capsulæ alis apice subtruncatis: angulo exteriore elongato-acuminato. *Brown mss.*

Abroma fastuosum. Salisb. paradis. 102.

Nat. of New South Wales and the Moluccas. 7.

v. 529]

TRICHOMANES. *Gen. pl. 1635.*

Sorus marginalis receptaculo columnari sæpius setiformi insertus. Indusium urceolato-campanulatum monophyllum erectum sorum includens.

1. *T. brevisetum*, frondibus 2—4-pinnatifidis glabris: pinnulis unifloris: laciniis linearibus margine integerrimis, involucris turbinato-oblongis apteris: apice simplici indiviso, columella subinclusa, stipite alata. *Brown mss.*

Hymenophyllum alatum. Willden. sp. pl. 5, p. 526. Engl. bot. 1417.

Nat. of Britain. 4.

EXTRACTS

FROM

DR. RICHARDSON'S

BOTANICAL APPENDIX

TO THE

"NARRATIVE OF A JOURNEY TO THE
SHORES OF THE POLAR SEA.

BY

CAPTAIN FRANKLIN."

AND

ADDENDA

BY

ROBERT BROWN, F.R.S.

[*Pp.* 729—768.]

LONDON.

1823.

PLANTS FROM THE
A P P E N D I X

TO

CAPTAIN FRANKLIN'S NARRATIVE.

[“In drawing up the list, imperfect as it is, I have received much assistance from able botanists. To Mr. Brown I am under the greatest obligations, not only for the liberal use of the Herbarium and Library, which, so happily for science, have been placed in his possession; but also for the friendly manner in which he aided my researches, and condescended to solve the doubts so frequently presenting themselves to one little versant in these pursuits. In addition to this general assistance, he kindly superintended the botanical drawings, and has enriched my catalogue with the lists of the Cyperoideæ (including the Carices), the Gramineæ, Junci, and Filices, and, with the accounts of the genera Eutoca, Heuchera, and Cryptogramma.”—Dr. RICHARDSON.]

12. ELEOCHARIS PALUSTRIS, *Roem. et Sch. Syst.* ii, [2 p. 151. Scirpus palustris, *Willd.* i, p. 291. *Pursh*, i,

p. 54. (W.)¹

13. SCIRPUS CÆSPITOSUS, *Willd.* i, p. 292. (W.B.)²

14. S. LACUSTRIS, *Willd.* i, p. 296. *Pursh*, i, p. 56. (W.)

15. S. MARITIMUS, *Willd.* i, p. 306. (W.)

¹ W. denotes the wooded country from latitude 54° to 64° north.—*Richardson.*

² B. denotes the barren grounds from lat. 64° to the Arctic Sea, in lat. 69°.—*Richardson.*

16. *S. SYLVATICUS*, Willd. i, p. 307. Pursh, i, p. 56. (W.)

17. *ERIOPHORUM VAGINATUM*, Willd. i, p. 312. (W.B.)

18. *E. ANGUSTIFOLIUM*, Willd. i, p. 313. Pursh, i, p. 58. (W.B.)

31 19. *E. STRICTUM*, spicis pedunculatis involucri brevioribus, squamis acutiusculis tenuissimè ciliatis, foliis strictis planis apice triquetris, culmo teretiusculo. Brown MS. (W.)

Obs. Dubia species *E. tenello* (Nutt.) proxima. Brown.

20. *ALOPECURUS ALPINUS*, Smith. Fl. Brit. iii, p. 1386. Brown, Suppl. Parry's Voyage, No. 54. (B.)

21. *A. ARISTULATUS*, Mich. Am. i, p. 43? *A. subaristatus*: Pers. Ench. i, p. 80. Pursh, i, p. 66. (W.)

22. *ACROSTIS LAXIFLOEA*, *Trichodium laxiflorum*, Mich. Am. i, p. 42, t. 8. Pursh, i, p. 61. (W.)

23. *STIPA CANADENSIS*, Poiret, Encyclop. Bot. vii, p. 452. Pursh, i, p. 72. *Stipa juncea*, Mich. Am. i, p. 54. *Oryzopsis parviflora*, Nuttall (fide Nutt. ipso). (W.)

24. *ORYZOPSIS ASPERIFOLIA*, Mich. Am. i, p. 51, t. 9. Pursh, i, p. 60. (W.)

25. *CALAMAGROSTIS CANADENSIS*, Nuttall, Gen. Amer. Pl. i, p. 46. *Arundo Canadensis*, Mich. i, p. 73. (W.)

26. *C. STRICTA*, *Arundo stricta*, Schrad. Germ. i, p. 215, t. 4, f. 5. Smith, Comp. Fl. Brit. p. 20. Eng. Bot. t. 2160. (W.)

27. *C. PURPURASCENS*, panicula spicata, glumis glabris, perianthii valvula inferiore scabra: apice 4-dentato; dorso aristata, rudimento plumoso villis baseos duplò longiore. Brown MS. (B.)

28. *AIRA AQUATICA*, Willd. i, p. 376. (W.)

29. *HIELOCHLOA FRAGRANS*, Roem. et Sch. Syst. ii, p. 515. Brown Suppl. Parry's Voy. No. 66. *Holcus alpinus*, Wahl. Fl. Lapp. p. 31, t. 2. (B.)

31. *TRisetum SUBSPICATUM*, Brown Suppl. Parry's Journ. No. 65. *Trisetum airoides*, Roem. et Sch. Syst. ii, p. 666. *AIRA subspicata*, Willd. i, p. 377. (W.B.)

32. *AVENA STRIATA*, Mich. Am. i, p. 73. Pursh, i, p. 86. (W.)

33. BECKMANNIA ERUCÆFORMIS, *Roem. et Sch. Syst.* ii, p. 695. *Cynosurus erucæformis*, *Willd.* i. p. 412. (W.)
34. POA CROCATATA, *Pursh*, i, p. 80. (W.)
35. P. ALPINA, *Willd.* i, p. 386. *Pursh*, i, p. 79. (W.)
36. FESTUCA OVINA, *Willd.* i, p. 419. (W.B.)
37. BROMUS PURGANS, *Willd.* i, p. 431. *Pursh*, i, p. 85. (W.B.)
38. HORDEUM JUBATUM, *Pursh*, i, p. 89. (W.)
39. ELYMUS CANADENSIS, *Willd.* i, p. 468. *Pursh*, i, p. 89. (W.)
40. E. MOLLIS, spica erecta villosa, locustis geminatis 4-5-floris brevè setigeris glumas setaceas superantibus. *Brown MS.* (W.)
111. LUZULA CAMPESTRIS, *Decand. Flor. Franc.* iii, p. 161. *Juncus campestris*, *Pursh*, i, p. 238. (B.)
112. L. MELANOCARPA, *Desvaux, Journ. de Botan.* i, p. 142, t. 5, f. 2. *Juncus melanocarpus*, *Pursh*, i, p. 238. (W.)
113. JUNCUS TRIGLUMIS, *Willd.* ii, p. 216. (B.)
114. J. CASTANEUS, *Smith, Fl. Brit.* i, p. 383. *Engl. Bot. t.* 900. (W.B.)
115. J. ECHINATUS, *Muhl. Gram. Am.* p. 207? (W.)
116. J. AFFINIS, foliis subulatis nodoso-articulatis, capitulis subpaniculatis paucifloris (3-5 floris), capsulis ovalibus calyce obtuso hexandro longioribus. *Brown MS.* (W.)
117. J. FILIFORMIS, *Pursh*, i, p. 236. (W.)
118. J. TENUIS, *Willd.* -ii, p. 214. *J. bicornis*, *Mich. Am.* i, p. 191. (W.)
119. J. GLAUCUS, *Willd.* ii, p. 206. (W.B.)
363. KOBRESIA SCIRPINA, *Willd.* iv, p. 205. (B.)
- CAREX. 1. *Spicis dioicis.*
364. C. DIOICA, *Willd.* iv, p. 207. (W.) [34]
365. C. SCIRPOIDEA, *Mich. Am.* ii, p. 171. *Pursh*, i, p. 34. C. Wormskioldiana, *Hornemann, Fl. Dan.* t. 1528. (W.)
2. *Spica androgyna simplici, arista stricta (vel exserta vel inclusa).* [35]

366. *C. FILIFOLIA*, *Nutt. Am. ii, p. 204.* (W.)

3. *Spica androgyna simplici mutica.*

367. *C. AFFINIS*, spica androgyna simplici superne mascula, stigmatibus tribus, squamis lanceolatis acutis maticis; infima aristata. *Brown MS.* (W.)

Obs. Proxima *C. polytrichoidi.* *Br.*

368. *C. ATTENUATA*, spica androgyna simplici: superne mascula densa: femineis paucioribus alternis, squamis omnibus obtusis. *Brown MS.* (B.)

4. *Spicis androgynis pedunculatis.*

369. *C. MEDIA*, spicis androgynis ternis brevissimè pedunculatis sessilibusve approximatis basi masculis, stigmatibus tribus, capsulis ovatis rostellatis glaberrimis squama ovata obtusiuscula longioribus. *Brown MS.*

Obs. Prope *C. bicolorem.* *Br.*

370. *C. FULIGINOSA*, *Stern. et Hoppe in Act. Soc. Bot. Ratisb. i, p. 159, t. 3.* (B.)

5. *Spicis androgynis sessilibus alternis.*

371. *C. SCOPARIA*, *Willd. iv, p. 230.* *Pursh, i, p. 37.* (W.)

372. *C. LOLIACEA*, *Willd. iv, p. 237.* (W.)

373. *C. REMOTA*, *Willd. iv, p. 239.* *Pursh, i, p. 37.* (W.)

6. *Spicis sexu distinctis, mascula solitaria, femineis subsessilibus scapo nudo bracteisque membranaceis vaginatis.*

374. *C. RICHARDSONII*, spica mascula pedunculata, femineis binis alternis subsessilibus exsertis multifloris, stigmatibus tribus, fructibus obtusis pubescentibus. *Brown MS.* (W.)

7. *Spicis sexu distinctis, mascula solitaria, femineis sessilibus s. incluse pedunculatis.*

375. *C. CONCINNA*, spica mascula sessili cylindracea, femineis ternis subsessilibus approximatis, bracteis semimembranaceis, stigmatibus 3, capsulis trigono-obovatis brevissimè rostellatis pubescentibus duplo ferè longioribus squamis obovatis, foliis caulinis superioribus abbreviatis. *Brown MS.* (W.B.)

Obs. Affinis *C. marginatæ*, quæ diversa spica mascula majori, femineis binis brevioribus, squamis obtusiusculis capsulam subæquantibus. *Br.*

376. *C. VARIA*, Willd. iv, p. 259. *Pursh*, i, p. 40. (W.)

377. *C. COMPACTA*, *Brown App. Ross's Voyage*. (B.)

Obs. Prope *C. saxatilem*. *Br.*

378. *C. LUPULINA*, Willd. iv, p. 266. *Pursh*, i, p. 41. (W.)

8. *Spicis sexu distinctis, mascula solitaria, femineis superioribus sessilibus vel inclusè pedunculatis, inferioribus pedunculatis.*

379. *C. MUTICA*, spica mascula squamis obtusis, femineis tribus distantibus subexserte pedunculatis erectis raris, stigmatibus binis, capsulis ovalibus muticis lævibus squama ovata mucronata longioribus, foliis bracteisque planis. [36 *Brown MS.* (W.)

380. *C. OLIGOCARPA*, Willd. iv, p. 279. *Pursh*, i, p. 41. (W.B.)

9. *Spicis sexu distinctis, mascula solitaria, femineis omnibus pedunculatis.*

381. *C. CAPILLARIS*, Willd. iv, p. 290. (W.)

382. *C. LIMOSA*, Willd. iv, p. 293. (W.)

383. *C. PODOCARPA*, spica mascula solitaria, femineis binis pendulis oblongis, stigmatibus tribus, fructibus ellipticis brevissimè rostellatis integris lævibus acheniisque pedicellatis, foliis caulinis inferioribus brevioribus lanceolatis. *Brown MS.* (B.)

10. *Spicis sexu distinctis masculis pluribus.*

384. *C. PELLITA*, Willd. iv, p. 302. *Pursh*, i, p. 44. (W.)

385. *C. AMPULLACEA*, Willd. iv, p. 308. (W.)

386. *C. ARISTATA*, spicis femineis ternis quaternisve cylindraceis distantibus brevè pedunculatis, stigmatibus 3, capsulis glaberrimis nervosis rostro longissimo alte bifido lævi: laciniis patentibus, squamis omnibus aristatis, foliis subtus vaginisque villosis. *Brown MS.* (W.)

Obs. Inter *C. bullatam* et *lacustrem*. *Br.*

424. *EQUISETUM ARVENSE*, Willd. v, p. 1. *Pursh*, ii, p. 651. (W.)

425. *E. SYLVATICUM*, Willd. v, p. 3. Pursh, ii, p. 651. (W.)

426. *E. UMBROSUM*, Willd. v, p. 4? (W.)

Obs. Caractere specifico quadrat. Vaginæ glaucae dentibus circiter 14 nigro-fuscis, marginibus hyalinis.

427. *E. PALUSTRE*, Willd. v, p. 5. Pursh, ii, p. 651. (B.)

428. *E. VARIEGATUM*, Smith, Comp. Fl. Brit. p. 154. Eng. Bot. t. 1987. (B.)

429. *E. SCIRPOIDES*, Willd. v, p. 7. Pursh, ii, p. 652. (W.)

430. *LYCOPodium COMPLANATUM*, Willd. v, p. 19. Pursh, ii, p. 652. (W.B.)

431. *L. SELAGO*, Willd. v, p. 49. Eng. Bot. t. 233. (W.B.)

432. *L. DENDROIDEUM*, Willd. v, p. 21. Pursh, ii, p. 653. (W.)

433. *L. ANNOTINUM*, Willd. v, p. 23. Pursh, ii, p. 653. (W.B.)

434. *L. ALPINUM*, Willd. v, p. 20. Eng. Bot. t. 234. (B.)

435. *POLYPODIUM VULGARE*, Willd. v, p. 172. Pursh, ii, p. 658. (W.)

436. *P. DRYOPTERIS*, Willd. v, p. 209. Eng. Bot. t. 616. (W.)

437. *WOODSIA ILVENSIS*, Pursh, ii, p. 660. *Nephrodium rufidulum*, Mich. Am. ii, p. 269, fide exempl. à D. Richard. (W.)

438. *W. GLABELIA*, frondibus (lanceolato-linearibus) pinnatis glaberrimis, pinnis triangularibus pinnatifidis: imis dilatatis: laciniis cuneiformibus, rachi nuda, stipite squamato. *Brown MS.* (W.)

439. *ATHYRIUM FRAGILE*, *Aspidium fragile*, Willd. v, p. 280. (W.)

440. *NEPHRODIUM FRAGRANS*, *Aspidium fragrans*, Willd. v, p. 253. (W.B.)

441. *PTERIS GRACILIS*, Willd. v, p. 376?? Pursh, ii, p. 668?? (W.)

Var.? *nana*, fronde unguiculari, pinnis omnibus indivisis.

442. CRYPTOGRAMMA ACROSTICHOIDES, frondibus bipinnatifidis, sterilium pinnulis ovalibus crenatis; fertilium demum explanatis, soris linearibus discum totum occupantibus. *Brown, ad finem Catalogi* [p. 526]. (W.)

ADDENDA TO CAPTAIN FRANKLIN'S NARRATIVE.

EUTOCA. *Brown.*

SYST. LINN. Pentandria Monogynia, post Hydrophyllum.

CHAR. GEN. *Calyx* 5-partitus, persistens. *Corolla* subcampanulata: membranulæ tubi decem, per paria filamentis alternantes. *Stamina* exserta. *Stylus* bifidus. *Capsula* polysperma, unilocularis, bivalvis, valvis indivisis medio placentiferis.

ORD. NAT. Hydrophylleæ, *Brown in* (Prodr. Fl. Nov. Holl. 492) *Bot. Regist.* 242, post *Phaceliam*.

CHAR. GEN. *Calyx* quinquepartitus, sinibus edentulis. *Capsula* polysperma.

Herbæ annuæ, pubescentes, eglandulosæ. Folia alterna, sæpius pinnatifida, nunc indivisa! Racemi terminales et quandoque e summis alis, secundi, ebracteati, novelli spiritaliter revoluti.

Obs. Ad hoc genus, a coordinatis (*Hydrophylo*, *Nemophila*, *Ellisia*, *Phacelia*,) diversum ovulis placentæ singulæ numerosis (unde nomen), pertinent *E. Menziesii*, (erecta, foliis linearibus integerrimis nonnullis quandoque incisis, ovulis placentæ singulæ viginti pluribus,) quæ forsitan *Hydrophyllum* lineare, *Pursh, Am.* 1, p. 134; et *E. parviflora* (diffusa, foliis pinnatifidis trifidisve; superioribus quandoque indivisis lanceolatis, ovulis placentæ singulæ 6—8,) quæ

Phacelia parviflora, *Pursh, Am. i, p. 140.* *Polemonium dubium*, *Linn.*

EUTOCA FRANKLINII, *Tab. 27.*¹

Eutoca erecta, foliis pinnatifidis bipinnatifidisve, ovulis placentæ singulæ viginti pluribus.

Hab. Abundantly amongst trees that have been destroyed by fire, on the banks of the river Mississippi.

Herba annua, erecta, pubescens, 6-10-uncialis. *Radix* descendens subramosa. *Caulis* simplicissimus teres cavus, crassitiè pennæ columbinæ, fuscus, pilis acutis tenuibus patulis pubescens. *Folia* radicalia conferta, caulina alterna; omnia petiolata exstipulata pinnatifida (in plantis prioribus bipinnatifida), circumscriptione lanceolata, laciniis lanceolatis integerrimis, inferioribus quandoque dentatis incisive, plana utrinque viridia pilisque caulinis similibus pubescentia. *Petioles* foliis breviores super canaliculati basi parum dilatata semiamplexanti. *Spicæ* racemosæ breves secundæ, ebracteatae, novellæ arcuato-recurvæ, approximatae, terminales et e summis alis.

Calyx monophyllus, æqualis, quinquepartitus sinibus edentulis, foliaceus, persistens: *laciniæ* lineares acutiusculæ planæ, trinerviæ nervis lateralibus margini approximatis medio venoso, hirsutæ pilis acutis strictis marginalibus crebrioribus longioribusque.

Corolla monopetala, hypogyna, regularis, subcampanulata, calyce sesquolongior, glabra limbo extus pube rara consperso, cærulea immaculata, siccatione sæpius albescens, decidua. *Tubus* intus nervis 15, ternatim cum staminibus alternantibus, lateralibus ternationis singulæ infra medium auctis membranula lineari imberbi apice subtruncato medium fere tubi attingenti, basi ad ortum filamenti respondentis producta et cum eodem fere confluenti, versus nervum medium conniventi. *Faux* nuda. *Limbus* tubo brevior, 5 partitus, æqualis, patens, lobis subrotundis integerrimis planis, venosis venis anastomosantibus, æsti-

¹ [See note at vol. i, p. 187.]

vatione imbricatis germina obtusa. *Stamina* 5, epipetala, æqualia, limbum corollæ paulò superantia. *Filamenta* prope basin tubi inserta cum limbi laciniis alternantia, filiformia, basi parum dilatata, pilosiuscula pilis longiusculis sparsis in ipsa dilatata basi crebrioribus, æstivatione induplicata. *Antheræ* ovali-oblongæ imberbes, incumbentes, basi semibifidæ lobis approximatis, loculis parallelo-contiguis medio longitudinaliter dehiscentibus. *Pollen* sphaericum simplex læve, in cumulo albicans.

Pistillum altitudine staminum. *Ovarium* liberum, sessile ovatum pilosum, basi auctum disco annulari adnato, uniloculare, placentis duabus parietalibus semiseptiformibus cavitatem fere bipartientibus, polyspermis lateribus ovuliferis ventre plano nudiusculo: *ovulis* plurimis (30 pluribus) in singula placenta confertim nec utrinque seriatim affixis. *Stylus* 1, filiformis pilosiusculus, semibifidus, laciniis filiformibus æqualibus modice patentibus glabris. *Stigmata* obtusa papulosa ipsis apicibus laciniarum styli paulo crassiora.

Capsula calyce persistenti parum aucto cincta eoque paulò brevior, ovata modice ventricosa, pilosa, unilocularis, bivalvis, valvis indivisis medio placentiferis, placentis adnatis longitudine fere loculi, lateribus seminiferis ventre nudiusculo. *Semina* numerosa in singula placenta viginti plura maturescentia, parva, circumscriptione ovalis, trigona angulis anguste alatis, areolata fusca, nucleo ovali.

TABULÆ 27 EXPLICATIO.

EUTOCA FRANKLINII, magnitudine naturali cum foliis duobus separatis, quorum alterum bipinnatifidum ad procerius, alterum ad nanum exemplar pertinet.

Sequentes omnes ad lentem auctæ.

- a. ejusdem. Calyx apertus cum Pistillo.
- b. ——— Corolla aperta exhibens paria quinque membranularum tubi et Stamina cum iisdem alternantia.
- c. ——— Anthera cum portione filamenti.
- d. ——— Ovarii sectio transversalis, exhibens placentas duas parietales polyspermas.

e. Capsula haud penitus matura.

f. Capsulæ sectio transversalis.

g. ——— Semen.

1. EUTOCÆ MENZIESII, Capsula matura aperta. 2. Ejusdem Capsulæ valvula altera, exhibens placentam seminibus denudatam. 3. Ejusdem Semen. 4. Idem testa remota. 5. Sectio longitudinalis Albuminis Embryonem ostendens.

HEUCHERA, Linn.

Pentandria Digynia. *Flores pentapetali, superi, capsulares, cum Vahlia.*

CHAR. GEN. *Caps. unilocularis, placentis 2 parietalibus adnatis polyspermis. Calyx 5-fidus (nunc inæqualis), æstivatione imbricata.*

ORD. NAT. Saxifragæ inter Tellimam et Vahlia.

CHAR. GEN. *Calyx 5-fidus, æstivatione imbricata. Petala indivisa. Stamina 5. Ovarium uniloculare, placentis duabus adnatis polyspermis. Styli 2. Capsula semi-infera, flore emarcido coronata, bircisiris, inter rostra dehiscens.*

Herbæ perennes, sæpius virentes. Folia radicalia conferta, elongato-petiolata, cordata, sublobata lobis serrato-incisis; stipulis lateralibus infra adnatis apicibus distinctis: caulina, dum adsunt, alterna, minora, brevius petiolata. Scapi paniculati, pedunculis trifidis basi unibracteatis, divisuris bibracteatis.

Obs. I. In ordine Saxifragearum locus Heucheræ est inter Mitellam gradifloram, Pursh. (Tellimam nob.) convenientem capsulæ unilocularis birostris dimidio supero flore persistente tecto, diversam staminibus decem, petalis laciniatis; et Vahlia, Thunb. (cujus certè species genuina est Oldenlandia pentandra, Retz. quæ Oldenlandia, Smith in Rees, Cyclop.) similem floribus pentandris, petalis indivisis, et capsulæ unilocularis dehiscentia, distinguendam placentis ab apice cavitatis pendulis solutis, æstivatione valvata calycis et habitu diversissimo.

Obs. II. A Saxifrageis transitus haud difficilis ad Ribes structura floris et ovarii admodum simile Heucheræ, diver-

sum stylo semibifido, fructu baccato, albumine densiore, et habitu. Attamen vel in eadem prorsus familia cum Ribe includenda, vel in proximo ordine (Escalloneæ) disponenda Escallonia *Mutis*. (Stereoxylon *Ruiz et Pavon*) et nonnulla genera inedita Novæ Hollandiæ, nec non Anopterus, *Labill.* (et *Prodr. Fl. Nov. Holl.* 457) cujus petala distincta, calyci inserta et ovarii capsulæque basis adhærens. Sed Anopteri cum Saxifrageis affinitas mediante *Heuchera* in fructificatione saltem obvia, obstante quidem habitu omnino cum Escalloneis et non Cunoniaceis nec Saxifrageis conveniente.

HEUCHERA RICHARDSONII, *Tab.* 29.¹

Heuchera calycis limbo inæquali obliquo.

Hab. On the rocky banks of rivers from lat. 54° to 64° north.

Desc. *Herba* acaulis, perennis. *Folia* radicalia conferta, elongato-petiolata, cordata, diametro sesqui-pollicari, semiseptemloba, lobo medio majore, omnibus inæqualiter serrato-incisis, super scabra subter lævia, secundum nervos venasque primarias pubescentia pilis brevissimis acutis, eglandulosa. *Petiolus* folio longior, triuncialis, pilosiusculus, scaber, antice canaliculatus. *Stipulæ* laterales infra adnatæ, apicibus lingulatis scariosis subciliatis. *Scapus* simplex, erectus, pedalis—sesquipedalis, gracilis, teres, striatus, pilosiusculus, dimidio inferiore pilis acutis divaricatis glandulisque intermixtis paucissimis, superiore pube brevissima glanduloso-capitata copiosa pilisque acutis rarioribus. *Panicula* laxè thyrsoidæa, angusta, ramis alternis bractea lineari apice bi-trifido subtensis, trifidis pericellis lateralibus 1-2-floris medio unifloro præcociore, divisis singulis bibracteatis, bracteolis lineari subulatis glanduloso-pubescentibus.

Calyx monophyllus, campanulatus, irregularis, persistens, extus glandulis subsessilibus parvis copiosis tectus: *limbus* obliquus 5-fidus, laciniis obtusis obovatis magnitudine

¹ [See note at vol. i, p. 187.]

subæqualibus sed ob faucis obliquitatem altitudine inæqualibus, duabus altioribus paribus impari demissiori, omnibus quinquenerviis nervis venosis, æstivatione imbricatis duabus altioribus reliquas equitantibus.

Petala 5, ipsis sinubus calycis inserta, unguiculata, indivisa, brevè ciliata, vix calycis lacinias superantia, persistentia, altiora lamina cuneato-obovata, demissiora sublanceolata, omnia trinervia nervis extus venosis, unguibus, angustis linearibus.

Stamina 5, summo tubo calycis inserta, ejusdem laciniiis opposita, petala vix superantia, persistentia. *Filamenta* subulata, glabra, laciniis calycis altitudine proportionata. *Antheræ* incumbentes, brevè ovaes, obtusæ, loculis parallelo-contiguis, longitudinaliter dehiscentibus. *Pollen* lateritium.

Pistillum: *Ovarium* semi-inferum, uniloculare, placentis duabus parietalibus longitudinaliter adnatis prominulis, cum stylis alternantibus, polyspermis: *ovulis* indefinite numerosis confertis superficiem ventralem lateraque placentæ operientibus. *Styli* duo, subulati, glabri, altitudine staminum. *Stigmata* obtusa imberbia, apicibus stylorum vix crassiora.

Capsula semi-infera, dimidio supero brevior calyce aucto persistente cum petalis vix emarcidis et filamentis sæpe antheris orbatis tecto, stylis rostrato, apice bilobo lobis intus dehiscentibus, unilocularis, placentis proportionatim auctis.

Semina numerosa, sessilia, ovalia, punctis elevatis acutis longitudinaliter seriatis aspera, altera extremitate obtusior. *Testa* crasso-membranacea, fusca, facile separabilis. *Membrana interna* albumini arcte adhærens absque chalaza rapheve manifesta. *Albumen* semini conforme, album, carnosum, molle, oleosum. *Embryo* minutus, subovatus albus ad albuminis extremitatem obtusior sitis. *Cotyledones* brevissimæ. *Radicula* processu capillari terminata, centrifuga.

TABULÆ 29 EXPLICATIO.

HEUCHERA RICHARDSONII, magnitudine naturali. i. basis dilatata petioli cum stipulis infra adnatis.

Figuræ sequentes ad lentem plus minus auctæ.

a. Flos plurimum auctus. b. Idem hinc longitudinaliter apertus. c. Ovarium longitudinaliter sectum, placentas parietales adnatas accurate, sed perperam stylos infra conatos, exhibens. d. idem transversè sectum. e. Capsula matura flore stylisque coronata, paulo tantum aucta. f. Ejusdem sectio transversalis. g. Semen plurimum auctum. h. Idem testa orbatum.

CRYPTOGRAMMA. *Brown.*

ORD. NAT. Filices. *Gyrata* (*Polypodiaceæ*) inter Onocleam et Lomariam.

CHAR. GEN. *Sori* lineares (v. subrotundi) venulis costæ (pinnulæ) obliquis insidentes. *Capsulæ* pedicellatæ, receptaculo communi elevato nullo. *Involucrum commune* (pinnulæ) marginale, continuum, disco venoso, margine scarioso libero sæpius induplicato; *partiale* nullum.

Filiculæ *glabellæ*. Frondibus *cæspitosis bi-tripinnatifidis*: *centralibus mutato-contractis fertilibus, exterioribus sterilibus*. *Involucris dorsum totum pinnulæ tegentibus*. *Capsulis breve pedicellatis, annulo incompleto*. *Sporulis obtuse trigonis, lævibus*.

Obs. Typus generis est Cryptogramma acrostichoides, sed character constructus pro receptione Pteridis crispæ auctor. quæ dubia quidem species, ob soros abbreviatos potius subrotundos quam lineares, venulas terminantes sinum ferè involucri occupantes et cito confluentes, tunc æmulantes sorum linearem continuum costæ parallelum Pteridis, cum cujus speciebus pinnulis angustatis involucri omnino tectis, habitu benè satis convenit. C. acrostichoides habitu et sororum dispositione et forma Grammitidibus nonnullis

præsertim *G. leptophyllæ* analoga, diversa frondibus fertilibus distinctis et præsentia involucri dorsum totum pinnulæ tegentis: His notis autem *Cryptogramma* convenit cum *Pteridi thalictroidi Swartz* (à nobis tredecem adhuc annis in *Prodr. Fl. Nov. Holl.* i. p. 154, uti genus distinctissimum indicata). In hoc genere, (*Teleozoma nob.*) Capsulæ sessiles, annulo completo latissimo, distinctæ in serie simplici insident venis costæ parallelis, utrinque vel solitariis marginalibus, vel binis altera superficialia. *Teleozoma* insuper insigne est sporis in ordine naturali forsân, in tribu quantum scio maximis obtuse trigonis pulchre striatis, puncto unico opaciore.

CRYPTOGRAMMA ACROSTICHOIDES.

Cryptogramma, frondibus bipinnatifidis, sterilium pinnulis ovalibus crenatis; fertilius demum explanatis, soris linearibus discum totum occupantibus.

Hab. In shady rocky woods, between lat. 56° and 60° north. (First found by Mr. Menzies at Nootka Sound.)

Desc. *Filiæ* 4-8-uncialis, glabra, lætè virens, frondibus cæspitosis, stipitatis, crasso-membranaceis opacis venis vix adversus lucem conspicuis.

Frondes exteriores cæspitis *steriles*, circumscriptione lanceolatæ bipinnatifidæ apice pinnatæ pinnis distinctis alternis, brevissimè petiolatis; pinnulis confluentibus, ovalibus obtusis, crenatis, inferioribus passim dentatis, omnium venis e costa vix prominula acutangulis distinctis simplicibus bifidisve intra marginem crenaturæ desinentibus. *Stipites* pallidi, subtus semiteretes anticè bisulci, infra medium conspersi squamulis lineari-lanceolatis acuminatis disco fusco-atro limbo pallido, supra glabri. *Rachis* epaleata, angusta, compressa, colore et fere substantia frondis. *Frondes* centrales *fertiles* stipite sterilium longiore alioqui simili insidentes bipinnatæ: pinnæ petiolatæ; *pinnulæ* omnes fructiferæ distinctæ inferiores brevissimè petiolatæ; ante explicationem involucri lineares, limbo utrinque recurvo efformante *involucrum* fornicatum discum

totum operiens pallide viride venosum margine scarioso
 avenio sæpius induplicato: demum explanatæ angusto-
 oblongæ involuero quasi nullo: costa subtus particulis
 parvis flavis conspersa, venis costæ obliquis, altè bifidis fere
 bipartitis, distinctis.

Sori ramulos omnes venarum a basi fere ad apicem occu-
 pantes, lineares, distincti sed ita approximati ut discus
 totus pinnulæ explanatæ capsulis maturis tectus est, et in
 hoc stadio filix species Grammitidis vel Acrostichi quasi
 evadit.

Capsulæ brevè pedicellatæ, turgide lenticulares annulo
 incompleto, altero lateri infra annuli terminationem trans-
 versim dehiscentes, *receptaculo* communi elevato nullo sed
 venæ parum incrassatæ insidentes.

Sporulæ triangulares, læves, absque striis manifestis
 vel puneto opaciori, in cumulo dilutè flavæ separatim
 hyalinæ.

NAMES OF AND NOTES ON

INDIAN PLANTS.

BY

ROBERT BROWN, Esq., D.C.L., F.R.S., V.P.L.S., &c.

[*Extracted from 'A Numerical List of Dried Specimens of Plants in the East India Company's Museum, collected under the superintendence of Dr. Wallich.'*]

LONDON.

1828—1849.

NAMES OF AND NOTES ON
INDIAN PLANTS.

- No.
184. For *Prionopteris* substitute *Matonia pectinata*,
Brown (correction at p. 23).
286. *Polypodium Horsfieldii*, R. Brown MSS.
287. *Polypodium Wallichii*, R. Brown MSS.
293. Duæ species sub nomine *Polypodium propinquum*,
Wall.; quar. altera *Polopod. melanopus* Brown
MSS. (correction at p. 83).
815. *Abelia triflora*, Brown.
819. *Knoxia teres*, Br. MSS.
820. *Knoxia mollis*, Brown.
821. *Knoxia brachycarpa*, Brown.
826. *Spermacoce longicaulis*, Brown.
829. *Spermacoce ramosissima*, Brown.
832. *Spermacoce lasiocarpa*, Brown. *Sp. stricta*, Roxb.
et Heyne vix Linn. (ex R. Br.).
833. *Spermacoce tenera*, Brown. *Sperm. ocymoides*,
Heyne haud Burm. (MSS. Br.).
836. *Spermacoce tubularis*, Brown. *Sp. hirta*, Heyne
haud Linnæi (MSS. Brownii).
838. *Hedyotis polycarpa*, Br.
840. *Hedyotis volubilis*, Brown. *H. scandens*, Wall.
apud Roxb. (MSS. Br.).
842. *Hedyotis cephalophora*, Brown.
844. *Hedyotis congesta*, Br. β *longifolia*, Br.
846. *Hedyotis* ? *macrocephala*, Br.
847. *Hedyotis vestita*, Br.
848. *Hed.* ? *glabra*, Br.

- No.
849. *Hedyotis costata*, Br.
852. *Hedyotis approximata*, Br.
853. *Hed. stylosa*, Br.
854. *Hedyotis articularis*, Br.
864. *Hedyotis?* *arguta*, Brown.
865. *Hedyotis?* *elongata*, Br.
867. *Hed. Heynii*, Br. *Oldenlandia herbacea*, Heyne et Roxb. haud Linn. (Brown).
868. *Hed. Burmanniana*, Br. *Oldenlandia biflora*, Roxb. haud Linn. (Brown).
869. *Hedyotis extensa*, Brown.
870. *Hedyotis linarifolia*, Brown.
873. *Hedyot. alsinifolia*, Brown.
874. *Hedyot. brachypoda*, Br.
875. *Hed. racemosa*, Lam. *H. alata*, Roxb. minimè Koenigii (MSS. Brown).
876. *Hed. tubularis*, Brown.
879. *Hedyotis biflora*, Br. (haud Lam.) *Oldenlandia*, Linn.
881. *Hed. scapigera*, Brown.
884. *Hed. puberula*, Brown.
885. *Hedyot. cymosa*, Brown.
886. *Hedyot. glabella*, Brown.
6195. *Hedyotis stipulata*, Brown.
1915. *Bamia Abelmoschus*, Brown.
2170. *Hemionitis hastata*, Brown in Herb. Banks.
2702. *Helicia robusta*, Brown. Rhopalæ, No. 1038—1041, sunt species *Heliciæ* fide illustr. Brown in suppl. primo Prodr. Nov. Holl. p. 32.
3538. *Scleria macrocarpa*, Wall. *Ptychocarya*, illustr. R. Brown (gen. nov.).
3651. *Deutzia staminea*, Brown.
3652. *Deutzia corymbosa*, Brown.
3719. *Marlea begonifolia*, Roxb. *Stylidium*, Lour. ex illustr. R. Brown.
3764. *Sporobolus coromandelianus*, Br.
3765. *Sporobolus diander*, Br.¹

¹ In a note on *Sporobolus* at the foot of p. 132 of his "List," Dr. Wallich

.No.

3796. *Hierochloa laxa*, Br.
 3807. *Michrochloa elongata*, Br.
 3815. *Eleusine brevifolia*, Br.
 4095. *Sonerila arguta*, Brown.
 4098. *Sonerila tenera*, Brown.
 4099. *Sonerila grandiflora*, Brown.
 4361. *Canscora diffusa*, Brown.
 4950. *Stauntonia latifolia*, Brown.
 4951. *Stauntonia angustifolia*, Brown.
 5225. *Podostemon Wallichii*, Brown.
 5432. *Crotalaria Trifolium*, Hb. Madr. Hæc et species
 sequentes (N. 5432—5437) ad *Cyrtolobum*, Brown;
 genus certè distinctum pertinent.
 6049. *Podocarpus Horsfieldii*, Brown.
 6098. *Nauclea Wallichiana*, Brown.
 6104. *Uncaria elliptica*, Brown.
 6112. *Uncaria ovata*, Brown.
 6119. *Ixora propinqua*, Brown.
 6134. *Ixora grandifolia*, Br.
 6135. *Ixora lucida*, Brown.
 6141. *Ixora opaca*, Br.
 6146. *Ixora diversifolia*, Brown.
 6147. *Ixora oblonga*, Brown.
 6149. *Ixora concinna*, Brown.
 6150. *Ixora densa*, Brown.
 6151. *Ixora coriacea*, Brown.
 6153. *Ixora elliptica*, Brown.
 6154. *Ixora nigricans*, Br.
 6157. *Ixora ? arguta*, Br.
 6165. *Ixora macrophylla*, Brown.
 6168. *Pavetta ? lucens*, Br.
 6171. *Pavetta nauciflora*, Br.
 6176. *Pavetta polyantha*, Br.

says: "Mr. Brown having undertaken the elaboration of the grasses, the specific names which are wanting will be supplied hereafter by that gentleman, who has had the goodness in the mean time to furnish this provisional list of the family." [As this intention was never carried out, those names only are quoted here which have Mr. Brown's authority specially attached to them.—
 EDIT.]

No.

6179. *Pavetta mollis*, Br.
6181. *Pavetta canescens*, Br.
6182. *Pavetta weberæfolia*, Br.
6225. *Myrioneura* (R. Brown) *nutans*, Wall.
6228. *Ophiorrhiza bracteolata*, Brown.
6232. *Ophiorrhiza discolor*, Brown.
6239. *Spermadictyon suaveolens*, Br.
6890. *Allmania nodiflora*, Brown (*Chamissoa nodiflora*, Mart.)
6891. *Allmania albida*, Br. (*Chamissoa*, Mart.)
6892. *Allmania esculenta*, Br.
6950. *Axyris Moorcroftiana*, Brown.
7129. *Marlea barbata*, Brown.
7130. *Pisonia morindifolia*, Brown.
7131. *Cyrtandra* ? *lanuginosa*, Brown.
7246. *Balanophora dioica*, Brown.
7369. *Eulophia virens*, R. Br.
7414. *Bucklandia populnea*, Brown MSS. (Fam. *Hamamelidacearum*, ex R. B.)
7415. *Bragantia Wallichii*, Brown MSS. (Fam. *Aristolochiear.* ex R. B.)
7417. *Indeterminatus frutex*, decumbens, fol. altern. simplicibus serratis, racemis amentaceis diclinibus (?) baccis monostylis. R. Brown.
7418. *Terminalia* ? (R. B.)
7419. *Myrtacea* (R. B.)
7421. *Arbor indeterminata*, fol. alternis coriaceis glabris elongatis acuminatis integerrimis exstipulatis (R. B.)
7422. *Arborea indeterminata macrophylla*, inflorescentia axillari (R. Br.)
7423. *Frutex indeterminatus alternifol. pilosus* (R. B.)
7424. *Frutex ramis virgatis*, fol. alternis integerrimis oblongis nitidis lævibus stipulatis acumine brevi obtuso (R. B.)
7425. *Urticea Trophidi accedens* (R. B.)
7426. *Uvariæ facie frutex oppositifolius* (R. B.)
7427. *Grewia* ? (R. B.)

No.

7428. *Uvaria* facie frutex sed stipulatus; folia alterna (R. B.)
7431. Frutex indeterminatus alternifolius, axillis nonnullis foliolis corrugatis inflorescentiam referentibus (R. B.)
7432. *Samydea* (R. B.)
7433. Indeterm. arborea fol. altern. simplicibus integerrimis impunctatis exstipulatis (R. B.)
7434. Indeterm. arborescens fol. alternis, integerrimis, magnis, pellucido-punctatis (R. B.)
7435. *Impatiens* habitu herba foliis immerse lineolatis (R. B.)
7436. *Oleina* quodammodo habitu frutex vel arbor (R. B.)
7437. Indet. arborescens fol. simplicibus alternis stipulatis impunctatis (R. B.)
7438. Indeterm. arborescens fol. simpl. alternis exstip. impunctatis (R. B.)
7439. Indeterm. frutex fol. altern. simpl. exstipulatis impunctatis pilosis (R. B.)
7440. *Chenopodiacearum* facie (R. B.)
7441. Indeterm. arborea fol. alternis simpl. longe petiolatis (R. B.)
7442. Indeterm. arborescens fol. altern. simpl. punctatis (R. B.)
7443. *Chaillieticearum* quodammodo facie arbor fol. simpl. altern. integerr. impunctatis pulchre reticulatis, cymis axillaribus (R. B.)
7444. Indet. arboresc. fol. simpl. subsessil. alternis argute dentatis, stipulis lateralibus spica axillari (R. B.)
7445. *Ammannia* facie planta fol. oppos. simpl. integerrimis (R. B.)
7446. Indet. arbor, fol. altern. simpl. elongato-petiolat., petiolo apice incrassato-tereti, stipulis subulatis, glomerulis axillaribus (R. B.)
7447. Indet. frutex fol. simpl. altern. subtus argenteis stipulatis, ramis tomentosis (R. B.)

No.

7448. *Celastrinea*? Arborea fol. alt. simplicib., stipulis minutis? flor. axillarib. subumbellat (R. B.)
7449. *Myrsinearum* familiæ arborea fol. simpl. suboppos. racemis laxis (R. B.)
7450. *Myrsinea*? Arborea priori similis (R. B.)
7451. Indeterm. arborea fol. simpl. altern. exstipulat. laurinis, fol. axillar. subcymos. parvis, duplici ordine trifidis, subvalvatis (R. B.)
7452. Indet. arborea, fol. altern. simpl. dentato-serratis impunctatis, gemmulis amentiformibus supra axillaribus (R. B.)
7453. *Samydearum* facie arbor, sed fol. impunctatis, stipulis falcatis deciduis (R. B.)
7454. *Elatines* facie plantula (R. B.)
7456. Indeterm. arborea, fol. simpl. alternis obovatis integerrimis coriaceis (R. B.)
7457. Indet. arborea fol. simpl. alternis (R. B.)
7458. *Samydea* ob fol. punctata, punctis passim linearibus (R. B.)
7459. Indeterm. arborea fol. pinnatis denticulatis, dentibus setigeris (R. B.)
7460. Indet. arborea fol. simpl. altern. coriac. integerrimis exstipulatis, gemmis axillaribus tomentosis (R. B.)
7461. *Maba*? (R. B.)
7462. Indet. arborescens resinifera, fol. oppositis simpl. integerr. oblongis acuminatis pellucido-punctatis, glomerulis axillaribus (R. B.)
7463. Indet. planta submersa litorea, conf. *Acorum maritimum*, König, MS. in Herb. Banks. (R. B.)
7464. *Cansjeræ* v. *Opiliæ* facie frutex alternifol. spicis axillaribus (R. B.)
7465. *Diplanthera*, Du Petit Thouars? Affinis *Cauliniæ ovali* (R. B.)
7466. Indeterm. frutex dichotomus, fol. simpl. altern., corymb. terminalib. laxiusculis (R. B.)
7467. *Rosacearum*? frutex icosandrus digynus (R. B.)
7468. Indeterm. frutex decumb. fol. altern. sim-

No.

- plicib. remotè dentatis, spicis terminal. dichot.
longiss. laxis (R. B.)
7469. *Aurantiaecus* frutex ramis aculeatis foliis ternatis
(R. B.)
7470. Indet. frutex fol. altern. simpl. subpunctatis,
petiolo ganglionoso, fructu pisiformi axillari
fasciculato, loculis momospermis (R. B.)
7471. *Samydearum* habitu frutex fol. altern. simpl. sti-
pulis deciduis, glomerulis axill. deciduis (R. B.)
7472. *Aurantiaecus* frutex inermis fol. ternatis in-
florescentiâ terminali floribus parvis 4-5-fidis iso-
stemonibus (R. B.)
7473. *Euonymus* (R. B.)
7474. *Viburnum* (R. B.)
7475. Indeterm. frutex, fol. altern. simpl. cymis late-
ralibus (R. B.)
7476. Indet. arborea fol. altern. simpl. serratis fructu
libero piperis formâ et magnit. monostylo baccato
oligospermo (R. B.)
7477. Indeterm. frutex fol. altern. simpl. dentat.
fructu subracemoso 1-stylo calyce persist. suffulto
(R. B.)
7480. Indet. frutex alternifol. fol. simpl. acuminat. inte-
gerr. exstipulatis, spicis axillaribus (R. B.)
7491. *Garcinioides* frutex (R. B.)
7495. *Cyrtandracearum* familiæ (R. B.)
7496. *Chrysobalaneus* frutex fol. alternis, racemis axil-
laribus, florib. icosandris, flor. lanatis (R. B.)
7497. *Sapindacea* ? arborescens macrophylla fol. altern.
pinnat. foliol. oblong. acuminat. spithamam
longis, petiolis partialibus basi incrassatis, racemis
abbreviatis, filamentis 5 lanatis (R. B.)
7498. *Hippocratiea* (R. Br.)
7499. *Brucea* ? Arbor fol. altern. pinnat. nunc ternatis,
foliol. lanceol. acuminatis marginatis, flor. cymosis,
ligno amaro (R. B.)
7501. *Urticeæ* habitu, arborescens (R. B.)
7502. *Rhamnaceus* ? frutex decumbens (R. B.)

No.

7503. *Elæodendri* facie arbor (R. B.)
7505. Indeterm. frutex decumbens, fol. oppositis subsessil. basi inæquali, stipulis interpetiolarib. amentis pisiformibus axillar. sessil. an merè foliaceis? (R. B.)
7506. Indeterm. frutex fol. altern. simpl. lanceol. obtusè acuminatis pellucido-punctatis, petiolis teretibus epidermide decidua rugosa, fructu semibivalvi monospermo (R. B.)
7508. Indeterm. arboresc. fol. altern. simplicib. petiolis apice incrassatis, pedunc. brevibus axillaribus multifloris (R. B.)
7509. Indet. frutex fol. altern. simpl. impunctatis exstipulatis (R. B.)
7510. *Olacinea*? fruticosa, fol. altern. simplicib. augustè lanceolat. (R. B.)
7511. Indeterm. fruticosa fol. altern. simplicibus stipulis foliaceis inciso-lobatis lateral. oppos. distinctis (R. B.)
7512. Indeterm. arborca fol. altern. subrotundis subtus tomento incanis, flor. glomeratis, perianthiis æstivatione valvatis (R. B.)

NOTES AND OBSERVATIONS

ON

INDIAN PLANTS.

BY

ROBERT BROWN, Esq., D.C.L., F.R.S., V.P.L.S., &c.

[*Extracted from 'Plantæ Asiaticæ Rariores,' by Nathaniel Wallich.
M. and Ph. D., Vols. I and III.*]

LONDON.

1830—1832.

Ar

dil
Ov
ter
cu
Ba

loc

ce
Ba

cr

te
na

ne
fo
A
te
un

NOTES AND OBSERVATIONS

ON

INDIAN PLANTS.

ABELIA. Brown apud Abel, in Itin. Chin. Append. (t. 14
B. p. 376.

CALYX foliaceus 5- vel 2-partitus. COROLIA subinfundibuliformis, 5-loba. STAMINA 4, didyma vel subæqualia. OVARIUM 3-loculare; loculis 2 polyspermis, abortientibus! tertio 1-spermo, fertili. PERICARPIUM 1-spermum, exsuscum, indehiscens, calycis limbo foliaceo aucto coronatum. *Brown MSS.*

Classis Linnæana, *Didynamia Angiosperma*. — *Brown loc. cit.*

Ordo naturalis, *Caprifoliaceæ veræ*, Brown; *Caprifoliacearum sectio* 1. Juss. Gen. p. 211, *Linnææ* prox. — *Brown.*

Habitus: FRUTICES. FOLIA opposita, petiolata, dentato-crenata vel integerrima. INFLORESCENTIA varia. — *Brown.*

ABELIA *triflora*. (Brown MSS.) Tab. 15.

FLORIBUS corymbi congesti ternatis: lateralibus 3-bracteatis; calycibus 5-partitis; foliis ovato-lanceolatis, acuminatis integerrimis. — *Brown.*

This new species of *Abelia* has in its inflorescence nearly the same relation to *Abelia Chinensis*, that *Caprifolium* has to *Xylosteum*; the central flower present in *A. triflora* and in *Caprifolium* being abstracted in *Xylosteum* and in *A. Chinensis*. In a third species (*Abelia uniflora*) sent from China by Mr. John Reeves, and for a

specimen of which I am indebted to the Horticultural Society, another modification of inflorescence occurs. The peduncles in this species have but one flower, which is considered as one of the lateral flowers, the ovarium being subtended by three bracts, placed exactly as in those of the corresponding flower of *Abelia triflora*.

Of the Natural Order *Caprifoliaceæ*, the only certain genera appear to me to be,—*Caprifolium*, *Xylosteum*, *Symphoricarpos*, *Linnaea*, *Abelia*, *Diervilla* (to which belongs *Weigelia* of Thunberg, who described the epigynous gland as the ovarium which he overlooked) *Triosteum*, and *Leyseria* of Dr. Wallich. I am not acquainted, however, with any other character distinguishing them from the genera, which have hitherto been associated with them, except the very minute one of the raphe, or adnate portion of the funiculus umbilicalis being on the outer instead of the inner side of the ovulum.—*Brown MSS.*

167

MATONIA. (Brown MSS.)

SORI dorsales, rotundi, e puncto confluentiæ venularum plurium orti. INDUSIUM orbiculatum, peltatum. *Capsula* sessiles, in serie simplici circa receptaculum dispositæ.—*Brown.*

Ordo Naturalis, FILICES, tribus *Polypodiaceæ*.—*Brown.*

Habitus: Filix pulcherrima, lævis, fronde subbipinnatâ. Pinnæ plures hinc superiores, pauciores inde secundæ, omnes pinnatifidæ, lobis integerrimis, singulis basi, nunc utroque latere, nunc inferiore tantum monosoris, raro bisoris.—*Brown l. c.*

MATONIA PECTINATA. *Brown MSS.*

INDUSIUM depresso-sphæroideum, capsulas omninò includens, hemisphærio superiore crassiusculo, inferiore tenuissimo, membranaceo, capsulis arctè applicito, et mox subevanido. RECEPTACULUM parvum, paulò elevatum. CAPSULÆ annulo obliquè verticali incompleto cinctæ. SEMINA angulata.—*Brown MSS.*

The beautiful ramification of the veins, and their union, from which the sorus originates in *Matonia*, is not altogether peculiar to it. Among the genera of *Polypodiaceæ* having an indusium, one remarkable example occurs in a genus as yet undescribed (*Hypoderris*), which with an indusium not materially different from that of *Woodsia*, has exactly the habit of *Aspidium trifoliatum*; while of those genera of *Polypodiaceæ* which are without an indusium, the same kind of vascularity is found in an extensive and very natural section of *Polypodium*, to which *Polypodium phymatodes* and the greater number of those species *soris saccatis* belong.—*Brown*.

COMETES. (Burm. Flor. Ind. p. 39). 117

CALYX 5-partitus. PETALA nulla. STAMINA imo calyci inserta, ferè hypogyna, antherifera 5, infrà cum totidem sterilibus membranaceis in utercolo connata. ANTHERÆ 2-loculares. OVARIUM 1-spermum, ovulo adscendente funiculo e basi cavitatis orto. STYLUS 1. STIGMATA 3. PERICARPIUM: UTRICULUS calyce persistente inclusus, appendicibus setaceo-ramosis, post anthesin auctis et expansis involueratis. SEMEN adscendens, chalazâ laterali. ALBUMEN unilaterale. EMBRYO periphericus rectus; RADICULA infera.—*Brown MSS.*

Ordo Naturalis: Cum *Pterantho* Forsk. parvulum tribum efformat, hinc *Illecebreis* proximum, inde ad *Amaranthaceas* veras per *Desmochætam*, *Digeram*, et *Saltiam* Nob. hodiè (quæ *Achyranthus papposa* Forsk.) minùs arcè tamen accedens.—*Brown*.

Habitus: HERBÆ (annuæ?) ramosæ oppositifoliæ. FOLIA integerrima, stipulis scariosis, subsetaccis, vel lateralibus liberis, vel basi cum petiolo connatis. PEDUNCULI e superioribus alis alterni, apice 3-flori. BRACTEÆ ternationis singulæ 6, quarum 2 oppositæ communes; reliquæ 4 per paria floribus lateralibus, intra communes pedicello brevissimo indentibus, pertinentes, subulatæ: singulæ appendice (ramo mutato) axillari, setaceo-diviso, ramulis sub anthesi fasciculatim approximatis, florecentiâ peractâ auctis

et patulis, subdivisionibus singulis bracteolâ subulatâ sub-
tensis.—*Brown*.

COMETES SURATTENSIS. Burm. Tab. 17.

FOLIIS cuneato-obovatis, ellipticisve; ramulis lævibus;
stipulis petiolaribus; fructus involucris ramulis fasciculatis,
imixtis deflexis.—*Brown MSS.*

The genus *Cometes*, proposed by the younger Burmannus (in *Flora Indica*, p. 39), was adopted in *Mantissa prima* by Linnæus, whose generic character agrees in most respects with the short description of Burmannus, from which it was no doubt chiefly formed: as, however, it differs in some points, he probably had seen and slightly examined the original specimen, which Burmannus may have taken with him to Upsal, as it is known he carried there for Linnæus's determination many of his rarer unpublished plants. But Linnæus, in describing the fruit ¹⁸⁷ of *Cometes* to be a "*capsula tricocca*," must have presumed on the affinity which he erroneously supposed the genus to have to *Dalechampia*.

Burmannus's specimen of *Cometes surattensis* I have seen in his Herbarium, now in the possession of Baron Delessert. It corresponds tolerably with the figure in *Flora Indica*, which, notwithstanding some differences, was probably made from it.

When engaged in drawing up the catalogue of Mr. Salt's Abyssinian plants, it occurred to me that the genus which I have in that catalogue named *Saltia*, was at least nearly related to *Cometes*; but I had at that time no means of verifying my conjecture. I afterwards, however, requested M. Decandolle to examine the specimen in M. Delessert's Herbarium, and the result of that examination is given in a note attached to the specimen, written by M. Delessert in 1816, in which it is stated, on M. Decandolle's authority, to be a species of his genus *Desmochæta*, or *Pupalia* of Jussieu.

In September of the same year I examined the specimen, and left attached to it the following note, which refers to

M. Decandolle's opinion, as well as to the Abyssinian genus, which I had published:—"Non *Desmochætæ* sed *Saltia* species, vid. Catal. Pl. Abyssin. in Itin. D. Salt. *Cometis* nomen restituendum."—R. B.

M. Guillemin, in the *Dictionnaire Classique d'Histoire Naturelle*, tom. 4, p. 356, states that M. De Jussieu, who examined, or at least saw, the specimen in M. Delessert's Herbarium, recognised it to belong to *Amaranthaceæ*, and that M. Decandolle regarded it as a species of *Desmochæta*. M. Guillemin himself, in adopting M. Decandolle's opinion proposes to apply to that genus the older name *Cometes*; and he adds that in a manuscript note in Burmannus's specimen I have proposed to do the same thing. But from that note, which I have already given verbatim, it appears that my proposing to restore the name *Cometes* referred to *Saltia*, and not to *Desmochæta*, to which it was evident to me *Cometes* did not belong.

In the Linnean Herbarium the specimen named *Cometes*, I believe in the writing of the younger Linnæus, proves to be a plant belonging to *Convolvulaceæ*, and it is probably a species of *Convolvulus* or *Ipomæa*.

Burmannus (in *Flora Indica*) has given the specific name of *Surattensis* to his *Cometes*, and that name Linnæus has adopted in his first *Mantissa*. In the twelfth edition of the *Systema Naturæ* (vol. ii, p. 127), published in the same year, but subsequently to the *Mantissa*, he changed the specific name to *alterniflora*, no doubt derived from the account of the inflorescence given both by Burmannus and himself. It is, however, not a very apt name for a plant whose flowers are always in threes, though the common peduncles are generally alternate. I have therefore recurred to the original name.

Sir James Smith, in a (pencil) note on the specimen in the Linnean Herbarium, though aware that the specimen is not really *Cometes*, supposes the specific name last given by Linnæus to have been suggested by it. This might have been the fact had that name been *alternifolia*, which, when he wrote the note, I have no doubt he believed it to be; but the actual name *alterniflora* could not well be

suggested by this plant, which has its flowers in a capitulum.
—*Brown's MSS.*

COMETES ABYSSINICA. (Brown MSS.) Tab. 18.

FOLIIS lineari-lanceolatis, mucronatis, pungentibus; stipulis liberis; ramulis pulvereo-pubescentibus, scabris; fructus involucri ramulis pinnatis, patulis.—*Brown MSS.*

Saltia abyssinica, Brown apud Salt, in Itin. Abyssin. Append. B. p. 376.

39]

ABELMOSCHUS.

CALYX 5-dentatus, spathaceus, deciduus, cinctus *involucro* (*calyce exteriori*) 5—10-partito. STIGMATA 5. CAPSULA 5-locularis, valvis medio septiferis, polysperma. SEMINA calva.—*Brown MSS.*

41]

SPHÆROPTERIS.

INVOLUCRA dorsalia, e medio venulæ orto, pedicellata, sphaerica, clausa, verticaliter dehiscentia, bivalvia. CAPSULÆ pedicellatæ, receptaculo communi convexo insidentes.—*Brown MSS.*

Habitus *Aspidii*, caudice nullo, frondibus decompositis, venulis subtus glandulosis, stipite rachique paleaceis.—Genus nimis forsitan affine *Diacalpi*, Blume Enum. Plant. Javan. p. 241 (fide speciminis Javanici a Domino Horsfield anno 1818 communicati), quæ similis habitu, venulis glandulosis, medio soriferis, involucro sphaerico, clauso, reticulato, areolis subrotundis, parietibus moniliformibus, nodulosisque; diversa involucro sessili, laceratim dehiscente, capsulis vix pedicellatis, et receptaculo obsoletiori insidentibus. Hoc ultimo caractere et reticulatione involucri a *Cyatheis* nonnullis (involucro e medio venæ orto) præsertim distinguenda; arctiore tamen affinitate nixa cum *Woodsia*, mediante specie mexicanâ (*W. mexicana* Nob.) quæ, ni fallor, *Physematium molle*, Kaulf. in Regensb. Bot. Zeit. 1829, 1 Band, p. 341, cui cel. auctor attribuit involucrum

"circumcirca clausum" quod verò in nostra planta apice divisum est, lobis subciliatis, arcuè conniventibus.—*Brown MSS.*

APOSTASIA. *Blume Bijdrag.* p. 423. (74)

PERIANTHIUM limbo sexpartito, regulari. STAMINA antherifera 2, quorum FILAMENTA foliolis lateralibus interiorum perianthii opposita, infra connata cum basi STYLI supra cylindracci, et vel nudi, vel hinc FILAMENTO TERTIO castrato, altiùs adnato, foliolo antico exteriorum opposito, appendiculati. ANTHERÆ biloculares, longitudinaliter dehiscentes. POLLEN e granulis simplicibus, solutis. STIGMA obtusum, obsolete bi-trilobum. CAPSULA trilocularis, polysperma. SEMINA ovata, testâ nucleo conformi.—*Brown MSS.*

This very remarkable genus, founded on *Apostasia odorata*, was first published in 1825 by Dr. Blume in the work referred to; but in 1821 a nearly related species was discovered in the valley of Noakote in Nipal, by the plant collectors of Dr. Wallich, who, in his manuscripts, which I have had the advantage of consulting, named it *Mesodactylis deflexa*, and at the same time had the drawing made which is here given.

I have followed these two distinguished botanists in regarding *Apostasia* as belonging to, or at least as most nearly related to, *Orchideæ*. It exhibits, however, very few of those characters generally considered essential to that family of plants.

In its antheræ, pollen, style, and stigma (all which parts are so remarkably modified in *Orchideæ*), *Apostasia* does not materially differ, either in form, structure, or economy from the more regular-flowered families of *Monocotyledones*; and in its trilocular ovarium it is distinguished from all other genera of the order to which it is here appended.

On the other hand, it agrees with *Orchideæ* in the structure, as far as I am able to ascertain, of its minute seeds, in the reduced number of stamina, and probably, with some genera of the family, in the order of their reduction;

in the filaments being at base connate with the lower part of the style, and in a great degree in habit. In endeavouring to estimate the importance of the several points of resemblance and difference here enumerated, with a view to decide on the degree of relationship *Apostasia* bears to *Orchideæ*, it is necessary to consider the relative position of the parts of the flower in that order, and also in *Scitamineæ*, the family most nearly allied to it.

The relation of stamina to the parts of the floral envelope in *Apostasia* is in the first place to be determined. The two antheriferous filaments, which I have more particularly examined in the unexpanded flowers of *Apostasia nuda*, appear to be opposite to the two lateral segments of the inner series of the perianthium; and the sterile filament in *Apostasia Wallichii*, and, no doubt, also in *A. odorata*, is opposite to the anterior segment of its outer series.

Several years since, I advanced the opinion, "That in a complete flower, whose parts are definite, the number of stamina, and also of pistilla, is equal to that of the calyx and corolla united in *Dycotyledones*, and of both series of the perianthium in *Monocotyledones*."¹

It may be further observed that, in cases of reduction of pistilla, it is generally found that the remaining carpella, when more than one, but inferior in number to that of one series of the floral envelope, correspond in position with parts of both series, and, with very few exceptions, whether distinct or confluent, are all equally developed. Stamina, on the other hand, in cases of equal reduction, generally belong to one of the series only, or, if corresponding with parts of both series, are usually in different states of development, as they are here described to be in two species of *Apostasia*.

This appearance of part of the inner series of stamina has not hitherto been expressly remarked in *Orchideæ*. It is not improbable, however, that the same relation to perianthium exists in the lateral antheriferous stamina of

¹ In 1826, in *Appendix to 'Denham and Clapperton's Travels,'* p. 237 (vol. i, p. 293).

Cypripedium,¹ as well as in the sterile petaloid processes similarly situated in the other genera, as in *Diuris*. And the third stamen of the inner series, still more altered in form, may be considered as present in certain New Holland genera, especially *Glossodia*, where this supposed stamen is placed within the labellum, but entirely distinct from it; in *Epiblema*, *Pterostylis*, and *Chiloglottis*, in which an analogous appendage similarly situated coheres in various degrees with that division of the perianthium; and perhaps it may be considered as indicated in all cases where the labellum is furnished with a process, however minute, arising from its axis.

If the view here taken of the position of the lateral filaments in *Cypripedium* and *Diuris* be adopted, it may be remarked that indications or rudiments of the two stamina necessary to complete the order in *Orchideæ*, of those, namely, corresponding with the lateral segments of the outer series of the perianthium, have not yet been observed in the regular structure of any plant of the order. They have, however, been occasionally met with in monstrous flowers of *Habenaria bifolia*, in more than one spike of which I have found the greater number of flowers triandrous, the three antheræ being equidistant, and placed exactly opposite to the three divisions of the outer series of the perianthium, the inner series of which remains in its ordinary state.

M. Achille Richard² has given an account of an 176 analogous monstrosity in *Orchis latifolia*. In this case of a triandrous *Orchis*, M. Richard having adopted the opinion, which I believe I was the first to advance,³ of the origin or nature of the auriculæ of the anthera of many genera of *Orchideæ*, considers the additional antheræ as formed by the perfect development of these auriculæ. This view, however, cannot be taken of the monstrosity of *Habenaria bifolia*, in which not only the auriculæ of the anterior or ordinary stamen are distinctly present, but two

¹ 'Prod. Fl. Nov. Holl.,' i, p. 309.

² 'Mém. de la Soc. d'Hist. Nat. de Paris,' i, p. 202.

³ 'Prod. Fl. Nov. Holl.,' i, pp. 309 & 311.

other similar processes, one on the anterior side of each of the additional antheræ, also exist; a fact which throws considerable doubt on the correctness of the view here referred to of the nature of these processes in *Orchideæ*, unless the same hypothesis could likewise be extended to all cases of trifid filaments, as those of *Allium* and *Deutzia*, to which the auriculæ in *Orchideæ* may be said to be analogous.

In *Scitamineæ*, the family most nearly akin to *Orchideæ*, the complete number of stamina may be considered very generally present. Only one, however, is antheriferous; and this perfect stamen, instead of corresponding, as in *Orchideæ*, with the anterior segment of the outer series of the perianthium, is placed within the posterior segment of the inner series, the two remaining barren stamina of the same series being the epigynous glands or filaments existing in all the genera of this order except *Costus*;¹ while the outer series of stamina, very differently modified, form the innermost or supplementary series of the perianthium.

This view of the origin of that series was many years ago communicated to me in conversation by the celebrated Correa de Serra; but was first, I believe, published in 1826 by Professor Lestiboudois, in a memoir² in which the correctness of the opinion held, namely, that *Scitamineæ* and *Cannæ* possess rudiments or modifications of six stamina, is remarkably contrasted with the erroneous views taken, or rather adopted, of the greater part of the structures adduced in support of it.

A more accurate account of the relative position of parts was given in 1828 by my ingenious friend Professor Von Martius. In confirmation of the opinion, I may remark that the cells of the ovarium, whose relation to the floral envelope appears to be very uniform in *Monocotyledones*,³ are in *Scitamineæ* opposite to the supposed petaliform stamina, and to the divisions of the outermost series of perianthium. I have formerly pointed out the difference in

¹ 'Prod. Fl. Nov. Holl.,' i, p. 305.

² 'Mémoire sur la *Canna indica* et sur les familles des *Balisiers* et des *Bananiers*.'

³ Appendix to 'Denham and Clapperton's Travels,' p. 243 (vol. i, p. 300).

position of the antheriferous stamen in *Scitamineæ* and that of *Canneæ* or *Maranteæ*, and have remarked that this difference is in some degree analogous to that existing between *Cypripedium* and the other genera of *Orchideæ*.¹

Apostasia in its trilocular ovary differs from all the genera of *Orchideæ*; but an analogous difference occurs in *Scitamineæ*, in which *Globba* is distinguished from every other genus in having its ovary unilocular, with three parietal placentæ. And in both these families it may be proved that the constituent parts of the compound ovary, whether unilocular or trilocular, agree in position or in their relation to the divisions of the perianthium.

Lastly, *Apostasia*, in the economy of impregnation, or the state of the pollen, and the manner of its application to the stigma, probably differs essentially from all *Orchideæ*, except perhaps *Cypripedium* and possibly *Vanilla*. But a similar difference, and in a degree still more striking, exists between *Apocineæ*, as I have formerly proposed to limit that order, and *Asclepiadeæ*, which can only be regarded as a subdivision of the same natural class.—*Brown MSS.*

APOSTASIA WALLICHII, Brown MSS.

ANTHERARUM lobis posticis inæqualibus, filamento tertio castrato.—*Brown MSS.*

APOSTASIA ODORATA, Blume, Bijdr., p. 423. [76]

ANTHERARUM lobis posticis æqualibus, filamento tertio castrato.—*Brown MSS.*

APOSTASIA NUDA, Brown MSS.

ANTHERARUM lobis posticis æqualibus, filamento tertio nullo.—*Brown MSS.*

¹ 'Gen. Remarks on Bot. of Terra Aust.,' p. 43 (vol. i, p. 49).

HUMBOLDTIA.

CALYX bibracteatus, tubo turbinato, limbo quadripartito, lacinia postica (e duabus conflata) binervi. Petala 5 vel (suppressione inferiorum) 3. Stamina 5, omnia antherifera, distincta, fauci calycis inserta, ejusdemque laciniis opposita. Ovarium pluriovulatum, stipite tubo calycis hinc adnato. Stylus subulatus. Stigma capitatum. Legumen (oblongum, compressum, *Vahl*).—*Brown MSS.*

Frutices (vix arbores) inermes. Folia abruptè pinnata, subtùs glandulis nonnullis adpressis instructa. Stipulae foliaceae, persistentes, peltatim adnatae, infra insertionem productae in lobum transversim dilatatum, subtùs foliorum instar glandulosum, (in sicco saltem) saepiùs reduplicatum stipulamque accessoriam exteriorem æmulantem. Racemi axillares, pedicellis bracteâ, præter duos calycinos, unicâ caducâ subtensis. Calycis limbus, æstivatione imbricatâ, deciduus, tubum persistentem superans. Filamenta æstivatione induplicata. Antherae versatiles.—*Brown.*

18] *H. laurifolia*, *Vahl*; pentapetala, foliis 3—5 jugis; stipularum lobo postico hinc productiore, acuto; ramulorum internodiis superioribus incrassato-fistulosis.—*Brown.*

RATZEBURGIA PULCHERRIMA, Kunth.

Ratzeburgia, Kunth, Revis. Gramin., p. 487.—Spica articulata, rachi flexuosâ. Locustae in singulo articulo 3. Duae sessiles, per glumas superiores parallelo-contiguæ, bifloræ. Glumæ valvula inferior cartilaginea, dorso depresso nervoso scrobiculato, apice membranaceo bilobo; superior chartacea planiuscula lævis. Flosculi hyalini mutici; superior hermaphroditus univalvis enervis. Squamulae hypogynae 3, quarum duae exteriores subcollaterales dilatato-cuneatae retusae bidentatae; tertia interior triloba (pro palea superiore habita a Dom. Kunth). Stamina 3. Ovarium imberbe. Styli ad basin distincti, interjecto umbone. Stigmata coarctato-plumosa. Flosculus inferior neuter univalvis ob-

soletè binervis marginibus inflexis. Locusta tertia pedicello libero insidens, neutra, univalvis.—*Brown MSS.*

Gramen perpulchrum, humile, stoloniferum, glabriusculum. Culmi adscendentes, monostachyi. Folia brevissima, rigidiuscula, plana, nunc conduplicata, obtusa, marginata, vaginâ compressâ, ligulâ imberbi, indivisâ.—*Brown.*

Obs. *Ratzeburgia* may be readily distinguished from all the other genera of *Panicæ*, as I have defined that extensive tribe, in having at each joint of its rachis three locustæ, of which two are sessile, fertile, and without aristæ, while the third, which is merely rudimentary, consisting of a single valve, is supported on a pedicel entirely distinct from the rachis.

In most of these characters, indeed, it agrees with *Mnesithea* of Kunth (*Rotböllia perforata* of Roxburgh), which differs, however, in having the pedicel of its third still more imperfect locusta, connate at both extremities with the rachis, and thus contributing to form the perforated joints so remarkable in this genus. In *Mnesithea* M. Kunth considers the upper valve of the glume as belonging to a third univalvular neuter flosculus—an opinion which seems to be founded solely on its membranaceous texture, and which, if adopted, would exclude this genus from *Panicæ*, as I have defined them, a definition which M. Kunth certainly does not adopt, as he has included in his tribe of *Rotbölliaceæ* several genera belonging to *Poaceæ*, namely, *Lepturus*, *Oropetium*, *Psilurus*, and *Nardus*.

According to the view here taken of the structure of the two genera, *Mnesithea* would have nearly the same relation to *Ratzeburgia* in its more important characters that *Rotböllia* has to *Ischænum Rotbölllioides*, 'Prodr. Flor. Nov. Holl.,' which I have in that work suggested might form a distinct genus. This genus has very recently been adopted by M. Brongniart, and named *Cælorachis*, with a character, however, which, as far as regards the structure of the pedicelled locusta, is so limited as to exclude both *Ischænum Rotbölllioides*, whose pedicelled locusta contains an hermaphrodite and a neuter flower, and *Rotböllia Cælorachis* of Forster, in which that locusta contains two male flowers

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

15 28
12 25
10 22
8 20

10 01

This last degree of development of the pedicelled locusta extends to other species, and such species differ from *Manisuris* merely in the form of the outer valve of the gluma of the sessile locusta.

Two other views, different from that here adopted, of the structure of *Ratzeburgia*, have been taken; the first by M. Kunth, who considers, and perhaps more correctly, the third squamula as the upper valve of the hermaphrodite flower. According to the second view, which is obviously suggested by the figure, the locusta would be regarded as containing one flower only.—*Brown*.

66]

AIKINIA BRUNONIS, Wall.

FOLIIS petiolatis, pedunculis solitariis pluribusve seriatis.—*Brown*.

Habitat in rupibus subumbrosis prope Cœpang in insula Timor; lecta florida et fructifera initio Aprilis anno 1803.—*Brown*.

RADIX annua, fibrosa, brevissima. CAULIS teres, succulentus, tenuissimè pubescens pilis brevibus acutis; 2—4 pollicaris, sæpius parùm ramosus, nunc simplicissimus. FOLIA petiolata, cordata, subovata, obtusa, repando-dentata, dentibus brevissimis, subsesquiuncialia, inferiora alterna, terminalia quandoque subopposita; omnia super viridia, pilis brevibus acutis sparsis scabriuscula; subter lætè viridia, glabra, pube brevissimâ nervum venasque primarias pinatas tantùm occupante. PETIOLI inferiores folia subæquant, pilosiusculi, semiteretes. SPICÆ unilaterales, circinatæ, racemosæ, demùm corymbosæ, pedunculatæ, rachi hinc (anticè) pedicellis confertis ebracteatis, florescentiâ peractâ sensim elongatis; indè (posticè) foliolio sessili, subrotundo, cucullato (*Commelinæ* ad instar) involucratâ. PEDUNCULI laterales solitarii, et terminales 2—4 collaterales, seriati inter folia duo subopposita, petiolata. CALYX liber, monophyllus, tubulosus, pilosiusculus, viridis, semiquinquefidus, dentibus æqualibus, erectis, acutis, persistens, post anthesin parùm auctus limboque patente

subcampanulatus. *Corollæ* magnitudine et figurâ ferè *Erini*; *tubus* calyce paullò longior, albus; *faux* imberbis, angusta; *limbus* quinquefidus, bilabiatus, patens, cœruleus, immaculatus, lobis obtusis, crenulatis, labio superiore bifido; inferiore tripartito, laciniis subæqualibus æstivatione superius equitante. STAMINA 4, epipetala, tubo inclusa. FILAMENTA omnia simplicia (ex icone *Baueriana bifida*), 2 superiora sub sinubus interlabialibus inserta, conniventia, tubi lateri superiori approximata, antherifera. ANTHERÆ apici obliquo filamenti basi adnatæ, pallidè flavæ, imberbes, apicibus mutuò coherentibus, bivalves, septo vix completo, contrario; pollen hyalinum, subglobosum, læve, exsulcum. FILAMENTA 2 inferiora absque antheris e lata basi subulata, apice parùm dilatato, subovato, acuto, plano. OVARIUM sessile, ovatum, glaberrimum, *squamis* hypogynis 2 oppositis, crassiusculis, truncatis, indivisis subtensum, uniloculare. PLACENTÆ 2 subrotundæ, indivisæ, compressæ, solutæ, polyspermæ, utrinque ovuliferæ, e pedicellis parietalibus adnatis angustis ortum ducentes. STYLUS filiformis, hyalinus, glaber, situ staminum iisque paullò longior. STIGMA peltatum, convexum, papulosum, indivisum. CAPSULA tubo calycis aucti limbo patente arcti cincta, obovata, apice exserto, stylo emarcido terminata, unilocularis, transversim dehiscens, marginibus laceris, operculo crassiusculè membranaceo, dimidio inferiore tenuissimè lacero. PLACENTÆ subrotundæ compressæ, liberæ, utrinque seminiferæ, e pedicellis adnatis filiformibus ortæ. SEMINA minuta, ovali-oblonga, castanea, striis spiralibus pulchrè insignita, basi subobliquè inserta *funiculo* elongato, semen ipsum superante, capillari, hyalino, per lentem pluriès augentem quasi e tubulo unico vel duplici annulari formato. TESTA crasso-membranacea, opaca, striis elevatis spiralibus hinc indè transversim connexis. MEMBRANA interna crasso-membranacea, tenax, albicans, areolis subquadratis tenuissimis. ALBUMEN vix ullum. EMBRYO dicotyledoneus rectus, orthotropus, albus. COTYLEDONES semiovatæ, obtusæ, vix longitudinæ *radiculæ* inferæ. — *Brown*.

Gratiola involucrata, Roxb. 'Flor. Ind.' [Ed. Wall.], vol. i,

p. 138, certainly belongs to this genus, and is perhaps not specifically different from the plant here described. But a second species, discovered in Java by Dr. Horsfield, in 1815, may be readily distinguished by the following character:—*AIKINIA* (*Horsfieldii*) folio unico sessili, caulem simplicissimum terminante, pedunculo solitario pluribusve seriatis.—*Brown*.

[EXTRACTS FROM]

PLANTÆ JAVANICÆ RARIORES.

DESCRIPTÆ ICONIBUSQUE ILLUSTRATÆ,

QUAS IN INSULA JAVA, ANNIS 1802—1818, LEGIT
ET INVESTIGAVIT

THOMAS HORSFIELD, M.D.;

E SICCIS

DESCRIPTIONES ET CHARACTERES PLURIMARUM ELABORAVIT

JOANNES J. BENNETT;

OBSERVATIONES STRUCTURAM ET AFFINITATES PRÆSERTIM
RESPICIENTES PASSIM ADJECIT

ROBERTUS BROWN.

LONDINI.

1838—1852.

sta
lee
th
Co
ta
Br
th
A
se
ge
di
an
th
ba
de

in
gr
th
no
po
R

fig
in
nu

PLANTÆ JAVANICÆ RARIORES.

IN the postscript to the 'Plantæ Javanicæ' Dr. Horsfield states—"On my arrival in England my zoological collections required my first attention, both with a view to their preservation and to their exhibition in the Honorable Company's Museum. It was therefore no less advantageous to myself than important to science, that Robert Brown, Esq., with a ready and disinterested zeal, undertook the examination and arrangement of my Herbarium. After much time and labour, every specimen of an extensive series of duplicates was examined, all the species of each genus were brought together, the entire Herbarium was distributed into families according to the natural method, and the basis of an accurate catalogue was formed. By this operation the number of species composing the Herbarium, which had previously been vaguely estimated, was determined to be 2196.

"While engaged in this arrangement Mr. Brown noted in each family those subjects which appeared to possess the greatest interest, either on account of their novelty, or of their peculiarity of structure; and from the subjects thus noted he finally made a selection, which it was proposed to publish under the title of '*Plantæ Javanicæ Rariores*.'

"It was at the same time determined that a series of figures for the illustration of all the subjects to be described in the work should be prepared. Although a considerable number of drawings had been made in Java by native

artists, these were found useful only in the representation of the general character and habit of the subjects. Mr. Brown therefore undertook the task of preparing such dissections as were necessary for the illustration of the generic characters, or other interesting points of structure, and of generally superintending the execution of the drawings; to this portion of the work Mr. Brown cheerfully devoted particular care and attention, and his experience in analysis of vegetable structure will be duly appreciated by Botanists.

“After the completion of the drawings and of the illustrative details, the subjects were put into the hands of the engraver, and Mr. Brown commenced the preparation of the text. In the prosecution of this task his public engagements, and other important scientific inquiries accumulating in a degree beyond what was expected when he commenced the work, it was agreed between Mr. Brown and myself that the original plan should be modified, and that the assistance of a coadjutor should be obtained.

“On this point I have the satisfaction to state that J. J. Bennett, Esq., Mr. Brown’s assistant in the Botanical Department of the British Museum, was found willing to prepare for the press such articles as were left unfinished by Mr. Brown, and likewise to co-operate generally in the preparation of the work. Mr. Brown has, agreeably to his original intention, contributed his remarks on the affinity and structure of the subjects described; he has also afforded many valuable suggestions in the progress of the work, and the whole has received his examination and revision.

“I embrace with pleasure and satisfaction the opportunity now afforded me of publicly expressing my great obligations to Mr. Brown. The examination and arrangement of my Herbarium, the laborious duties connected with the superintendence of the figures contained in this work, the preparation of the illustrative details, and the time devoted to the description of the subjects, are by no means the only marks of friendship which I have received

from that distinguished Botanist; who ever since his arrival in England has afforded to me his advice and assistance in my researches connected with Natural History, and on many other important occasions."

[The Extracts which follow are limited to Mr. Brown's direct contributions, but his invaluable suggestions enrich almost every page of the work.—EDIT.]

POLYPODIUM (DIPTERIS) HORSFELDII.

TAB. I.¹

CHAR. SUBGEN. (DIPTERIS) *Sori* subrotundi, sparsi (v. transversim subseriati), inter (frondis palmatæ) venas primarias dichotomas earumque divisiones, venulis divaricatissimis anastomosantibus insidentes. *Indusium* (verum) nullum.

Caudex repens teres. Frondes elongato-stipitata binatæ; partiales dimidiatæ palmato-lobatæ. Venulæ secundariæ tertiariæ et ultimæ divaricatissimæ, crebre anastomosantes, penultimæ latere soriferæ, ultimæ apice vix dilatato libero. Indusium spurium vel (in D. Horsfieldii) pili sorum cingentes et capsulis intermixti; vel (in D. Wallichii) materia pulposo-gummosa capsulas immaturas obtegens.

POLYPODIUM (DIPTERIS) *Horsfieldii*, frondis binatæ palmatæ subtus glaucæ lobis serratis, soris pilosis.

Dipteris conjugata. *Reinv. in Syllog. Pl. Nov. Soc. Bot. Ratisb., tom. ii, p. 3.*

Polypodium Dipteris. *Blume Enum. Pl. Jav., p. 135.*

DESCR. Filix 2—3-pedalis, vel elongatione stipitis usque orgyalis. Caudex repens crassitie pennæ olorinæ fibrillas copiosas nigricantes subramosas dimittens squamisque rigidis strigosis atris subulatis setaceo-acuminatis tectus, solidus, constans ligno quasi duplici, utroque e cellulis elongatis impunctatis utrinque acutis composito, cum strato interposito tubum completum constituyente, e vasis scalaris

¹ The Plates of "Plantæ Javanicæ Rariores," not being at present accessible, will not be given among the Illustrations. They are entirely systematic, and the reader is referred for them to the original work.—EDIT.

formibus in laminam transversim interrupte striatam vix definitæ latitudinis facilè, ut in plerisque Filicibus Lycopodineisque, solubilibus. Stipites elongatæ frondem ipsam sæpius superantes crassitie pennæ anatinæ, adultæ glabræ læves pallidè castaneæ teretiuseculæ anticè canaliculatæ; vasis scalariformibus tubum ferè completum anticè deficientem efformantibus. Frondes ad basin usque bipartitæ potiusve binatæ; partiales dimidiatæ palmato-lobatæ, lobis incisus indivisisque acuminatis serratis; novellæ utrinque tectæ vellere ferrugineo-castaneo e pilis articulatis laxis constante; adultæ glabræ pilis nonnullis secundum latera venarum primarium aliisque soros cingentibus relictis. Venæ primariæ dichotomæ validæ, subter prominentes teretes, super vix emersæ, divisionibus ultimis sæpe infra apicem acuminatum lobi conniventibus et unitis. Venulæ secundariæ et tertiariæ divaricatissimæ creberrimè anastomosantes areolas subquadratas sed amorphas efformantes, ultimæ liberæ apice vix dilatato. Epidermidis areolæ minutæ parietibus paulo flexuosis. Stomata in pagina tantum inferiore. Sori parvi subrotundi passim e confluentia duorum amorphi, lateribus venularum tertiarium sæpiusque ubi plures confluunt insidentes, novelli pilis articulatis copiosis semitecti, adulti pilis brevioribus capsulis interstinctis. Capsulæ brevè pedicellatæ annulo verticali incompleto lateribus areolatis. Spora lævia subovalia inæquilatera margine altero convexo altero rectiusculo.

OBS. I.—Dr. Horsfield, who discovered this remarkable species in 1814, has the following observations respecting it.

“I have four localities for this Fern. The most western is the mountain Karang, situated south of Bantam; next the mountain Gede, south of Batavia; proceeding westward, Ptarangan; and finally Mount Prahu; the latter lies south-west of Samarang. It is remarkable that I never noticed it in my numerous excursions through the eastern provinces; and I have concluded that it is not at all, or at least that it is very rarely, found eastward of the longitude of Samarang, which lies near the middle of the island, as far

as regards its extent from west to east. From its striking physiognomy it could not, I think, have escaped my notice.

"It grows in clusters with fronds of different height, the tallest often equalling five or six feet.

"Its range is above that at which rice can be cultivated in Java, which is about 5000 feet above the level of the ocean. It springs from the black vegetable mould, which very generally covers the mountains of Java."

OBS. II.—Before entering on the subject of the affinities of *Polypodium* (*Dipteris*) *Horsfieldii*, there are some parts of its structure deserving of notice.

In the first place, the existence of the complete circle of vasa scalariformia separating the ligneous or fibrous vessels of the caudex into an outer and inner portion, though not peculiar to *Dipteris*, seems to be of rare occurrence among Ferns. It exists, however, in the caudex of *Platyzoma* (though not in that of *Gleichenia*); and I have observed it also in some (probably it will be found in all) species of *Anemia*.

I have described *Polypodium* (*Dipteris*) *Horsfieldii* as having hairs surrounding the sori, and mixed with the capsules, while in the nearly related species *Polypod.* (*Dipteris*) *Wallichii*, the capsules, even when fully formed, are imbedded in a gum-like pulpy substance, the remains of which may always be found, even after the spora are discharged. In the dried specimens examined this pulpy substance had no appearance of organization; in the living plant, however, it will probably be found to be minutely cellular, as I have observed in the nascent sori of several species of *Polypodium*, in which the production of this cellular pulp precedes the formation of the capsules, and is generally evanescent soon after they become visible. The peculiarity in *Polypod.* (*Dipteris*) *Wallichii* probably, therefore, consists merely in the longer duration of this substance, in which perhaps the capsules of the greater number of *Polypodia* are formed.

3] OBS. III.—The Fern here described, and the nearly related species *Polypodium* (*Dipteris*) *Wallichii*, form a very

distinct section or subgenus of *Polypodium*, to which they must be referred, while the characters of that genus consist merely in its round sori, and absence of true indusium. But in adding two species, having so singular a habit, to a genus already too extensive, and including so many different forms, it must be evident that the present sources of generic distinction in this Natural Family are very insufficient for its satisfactory analysis.

The number of Ferns at present known, including unpublished species existing in collections, may be stated at upwards of 1800. The Family has been subdivided into five primary groups. These groups are strictly natural; they are not, however, of equal value as to the importance of their characters, and they are extremely unequal in extent; the great mass belonging to that called *Polypodiaceæ*, which has been divided into upwards of fifty genera, and contains about 1600 species. Of this number nearly one half, or 800, belong to five genera, namely, *Polypodium*, *Aspidium*, *Nephrodium*, *Asplenium*, and *Pteris*, as they are at present constituted, *Polypodium* alone consisting of about 300 species. The extent of *Polypodium*, therefore, renders its subdivision into natural sections or subgenera absolutely necessary.

For such subdivision, not in *Polypodium* only, but in other extensive genera of Ferns the most obvious, as well as the most advantageous source of character seems to be the modifications of vascular structure, or the various ramifications of the bundles of vessels, or veins of the frond, combined with the relation of the sori to their trunks or branches.

On this subject many important observations have been made, and subdivisions founded on them proposed by several writers, especially MM. Bory, Gaudichaud, Kaulfuss, Ad. Brongniart, and Blume. I may add, that I have at a still earlier period¹ introduced the ramification and relation of sori to veins into the definitions of several genera, in which these characters had not been before employed, and have more recently,² in treating of *Mutonia*, pointed out a natural and extensive group in *Polypodium*,

¹ [*Prod. Fl. Nov. Holl.*, i, p. 156-8.]

² [*Antè*, pp. 542-3.]

as being readily distinguishable by the sorus originating at the point of confluence of several veins.

With respect to DIPTERIS, the section to which our plant belongs, there is no difficulty in distinguishing it from all other groups of Polypodium, and particularly from that now alluded to, if the dichotomous ramification of the primary veins be admitted into its definition. And as that ramification may be said to be necessarily connected with the peculiar division of the frond, this section, so constituted appears to rest on characters at least as important as those of several groups at present generically distinguished from Polypodium, as Cyclophorus, Pleopeltis, Adenophorus, and even Grammitis, Selligera, and Meniscium.

If, however, the dichotomous primary veins are left out of consideration, no sufficient character remains to distinguish Dipteris from that section of Polypodium, including *P. quercifolium*, *diversifolium*, and several other species, and which M. Bory has established, chiefly from the presence of dissimilar sterile fronds, as a subgenus under the name of *Drynaria*. But the existence of these sterile fronds being neglected, *Drynaria* cannot be separated from that more extensive section comprehending *P. phymatodes*, *lycopodioides*, &c., and to which (including *Drynaria*) I have referred in my observations on *Matonia*, in Dr. Wallich's *Plantæ Asiaticæ Rariores*.¹

These three subdivisions of Polypodium agree in having their sori placed on the point of confluence, or perhaps sometimes of divarication, of several branches of the anastomosing veins; and Dipteris being distinguished by its dichotomous primary veins, the remaining two sections may form one subgenus, for which the name proposed by M. Bory may be adopted.

In many species of DRYNARIA so constituted, the principal vein of the sorus is manifestly that in which the tendency to produce capsules is generally the greatest in the natural order; namely, the lowest branch of the upper or inner side of the primary vein, or that branch which in the appendix to Captain Flinders's Voyage² is considered as

¹ [*Antè*, p. 543.]

² [Vol. i, p. 60.]

having a relation to two different trunks; in other species, in those particularly where the sori are most numerous, and not confined to that branch, it is often difficult to determine the most important vein of the sorus; and in *Dipteris* it is hardly practicable.

From *Drynaria*, and from those species of it especially in which the principal vein of the sorus is distinctly marked, the transition is easy to *Polypodium aureum*, *decumanum*, and a few other species having anastomosing veins, and in which the sori are placed on the apices of two, or more rarely three, connivent ultimate ramuli, included in an area formed by the anastomosing secondary veins. But these species, from the identity of habit, may be included in, or appended to, a more extensive group, whose anastomosing veins form areæ or meshes, in each of which only one sorus exists, and that terminating a single included branch. This section, which may be named *PHLEBODIUM*, and whose species have either pinnate, deeply pinnatifid, or more rarely simple fronds, appears to me strictly natural, though it includes several species having the spurious indusium of *Pleopeltis*, and at least one with an oval or even oblong sorus.

Next to *Phlebodium* in affinity, as well as in the arrangement of the sori, may be placed a group, most of whose species have simple fronds, and all of which are natives of America. In this group, which may be distinguished by the subgeneric name *CYRTOPHLEBIUM*, the primary parallel veins are connected by transverse arched branches, from the convex upper side of which generally three (and never more than three) upright parallel simple veins arise, terminating within the area included between the proximate transverse arched branches: of these simple tertiary ramuli the two lateral are soriferous, generally below the apex, the middle branch being always sterile. In one species at least this branch reaches, and is united with, the arched secondary vein above it; and in a few others it is entirely wanting. This arrangement in *Cyrtophlebium* produces two series of sori between each primary vein, except in the lowest area, or that next the

midrib, in which there is one only, and that terminating the single vein exactly as in *Phlebodium*, into which this group passes by species having the habit of *Cyrtophlebium*, but with fronds so narrow, that they are reduced to the lowest areolæ, and consequently agree in character with *Phlebodium*.

An arrangement of veins and of sori analogous to *Cyrtophlebium* exists in the real species of *CYCLOPHORUS* or *Niphobolus*, none of which are natives of America: in all these the secondary veins are straight and parallel, instead of being arched; they are also given off at an angle more or less acute from the primary parallel veins, which they connect; and the tertiary or ultimate branches originating only on the upper side of each secondary vein are parallel with each other, more than three in number, and all of them bearing terminating sori.

An extensive and strictly natural group may be next noticed, though it cannot be considered nearly akin either to any of the preceding sections or to the principal part of that which follows.

This group or subgenus, the *LASTREA* of M. Bory, whose fronds are either bipinnatifid or simply pinnate, is chiefly intratropical. Its character consists in the secondary veins of the pinnate, and the only veins of the segments of the bipinnatifid fronds being perfectly simple and parallel, with one known exception reaching the margin of the segment, or in the pinnate species uniting with the corresponding vein, and each bearing a lateral sorus, generally about the middle, in some cases near the base, and in a few others proceeding from the base itself.

The closest affinity of *Lastrea* is not to any group of *Polypodium*, but to that section of *Gymnogramma*, the division of whose fronds, and the disposition of veins, are exactly similar, and in which the sori form very short lines of like origin. As the only distinction therefore consists in a difference, generally very slight, in the form of the sorus, it appears to me (and Dr. Blume has made a similar remark) that these two tribes cannot be generically separated, especially as species belonging to both agree in

having pilose capsules; and it is at least certain that the section of *Gymnogramma* referred to should be removed from that genus, and if still distinguished as a genus or subgenus, might receive the name of *PLEUROGRAMMA*.

This section is also manifestly related to *MENISCIMUM*, which, though readily distinguished by its technical character, may be considered as a subgenus of *Polypodium*, nearly approaching in vascular structure to the pinnate species of *Lastrea*, but differing in the sori terminating the corresponding veins at their point of junction, and by their confluence forming a semilunar line. It is deserving of remark, that in some, and especially the original species of *Meniscium* in that part of the frond where the power of producing capsules is the weakest, the sorus is transferred from the point of union of veins to the upright branch arising from that point, and from semilunar becomes round or sublinear.

Lastrea may even be considered as approaching in affinity rather more nearly to that section of *Nephrodium*, which M. Gaudichaud has separated under the generic name of *Polystichum*, than to any subdivision of *Polypodium*; an approximation which appears to be confirmed by more than one fern, entirely agreeing in habit, in undivided veins and lateral fructification with this group of *Nephrodium*, but having a short linear sorus, with an *Indusium* of corresponding form, inserted by its longitudinal axis in the middle of the sorus. To this group the name of *MESOCHLÆNA* may be given; and though in general appearance it is abundantly different from *Didymochlæna*, it can only be distinguished from that genus, according to my view of the structure of its *indusium*, by its simple veins and lateral sori.

The most extensive but least natural section of *Polypodium* is that in which none of the veins anastomose, the sorus standing either on the apex, generally more or less dilated, of the branch, or distinctly below the apex, which is then scarcely or not at all dilated; the vein when simple, or its upper scriferous division when branched, terminating within the margin of the frond or pinna.

Polypodium vulgare is one of the best examples of this section, which, however, comprehends species with undivided, pinnatifid, pinnate, bipinnate, and even decompound fronds, and these as various in texture as in division. Many species are found within the tropics; but the section includes all the European Polypodia and most of those found in the higher latitudes of the southern hemisphere.

As this section includes many of the earliest described species of Polypodium, it would require, if retained entire, no subgeneric name. But in a complete analysis of the genus it ought to be divided. The strictly natural subdivisions, however, can hardly be characterised from modifications of vascular structure alone; and I have not yet been able to detect sufficient differences, either in the capsules or seeds, by which they may be distinguished.

That subgeneric or sectional characters may in several instances be obtained or assisted from the seeds of this Natural Order is not improbable, and in one case, namely CERATOPTERIS (or Teleozoma), including Parkeria in that genus, even the generic character appears chiefly to reside in the seeds, which in their unusual size and peculiar marking or striation entirely agree in all the species of the genus, while in the original species the annulus is nearly complete; and in Parkeria, differing from the rest of the genus in no other point whatever, the ring is reduced to a few faint striæ.

TAB. I. *Fig. 1.* A frond of *Polypodium (Dipteris) Horsfieldii*, of the natural size. *Fig. 2.* A portion of the under surface magnified, showing the arrangement of the veins and sori. *Fig. 3.* Another portion of the same, from which a sorus has been removed to show its insertion. *Fig. 4.* A sorus, more highly magnified. *Fig. 5.* A capsule before bursting. *Fig. 6.* A capsule after bursting. *Fig. 7.* Sporules.

ATAXIA HORSFIELDII.

[8

TAB. III.

ATAXIA, Br. in *Chlor. Melv.* p. 35. Kunth, *Agrost. Synopt.*, p. 39.

CHAR. GEN. *Gluma* locustam trifloram subæquans. *Flosculus inferior* masculus bivalvis; *intermedius* univalvis neuter: uterque dorso aristatus; *terminalis* hermaphroditus diandrus.

Gramen *tam habitu quam structurâ* inter *Anthoxanthum* et *Hierochloam medium*, *pariterque odoratum*. *Gluma inæquivalvis*. *Flosculi a basi brevissimâ persistenti racheos conjunctim solubiles*. Valvula superior hermaphroditi flosculi *uninervis*; staminibus *axibus valvularum oppositis*.

Obs. *Ataxia* along with *Hierochloe* and *Anthoxanthum* form a very natural and well characterised section, which belongs rather to the tribe *Avenaceæ* than to *Phalarideæ*; and these three genera are in reality so nearly related that they may perhaps be more properly considered as sections of one and the same genus; or at least *Hierochloe* and *Ataxia* might be united. It is deserving of notice that in all of them the upper valve of the hermaphrodite flosculus has a single nerve occupying its axis, and that one of the two stamina is placed opposite to this nerve. The co-existence of these two characters, both of which are remarkable deviations from the usual arrangement in *Gramineæ*, seems to invalidate the hypothesis respecting the composition of the inner valve of the flower in this family.¹ It might, however, be assumed that the median nerve in these genera is formed of two confluent cords, a view to a certain extent supported by the somewhat analogous structure in the corolla of *Compositæ*. It might also be assumed

¹ General Remarks, &c. in Flinders's Voyage, vol. ii, App. p. 580 [vol. i, p. 55].

that the stamen belongs to the inner or complementary series, which is rarely developed in triandrous genera. One remarkable apparent exception to the usual order of development of stamina occurs indeed in a genus of grasses found in Abyssinia by Dr. Rüppell, to whom I am indebted for the specimens I have examined. In this genus the locusta contains apparently a single flower, of which the gluma consists of two minute obtuse nerveless valves; the perianthium is formed of two valves nearly equal in size, form, texture, and nerves, which are three in number, the middle nerve of each valve ending in a seta; the stamina are three in number, but instead of being inserted as I have described those of triandrous grasses generally to be, they are placed within the upper or inner valve, the middle stamen being opposite to the median nerve; the embryo also is placed on the side of the inner valve: hypogynous squamulæ are entirely wanting. If the flower here described be really simple, it would present a still more formidable objection than *Ataxia* to the composition of the inner valve of the perianthium. But the arrangement of stamina, and direction of scutellum or embryo, suggest another hypothesis with respect to the Abyssinian genus; namely, that the flower is not simple, but made up of two flowers reduced to their outer valves. This latter view I am disposed to adopt, not only on considering the usual order of suppression of the parts of the floral envelope in grasses; but from the same degree of reduction actually existing in several *Paniceæ*, to which primary division of *Gramineæ* the Abyssinian genus would according to this view belong. It may be added that the genus referred to very remarkably agrees, both in habit and structure, with an unpublished genus discovered by Ehrenberg, likewise in Abyssinia (*Podopogon*, Ehrenb. MSS.), and which unquestionably belongs to this primary division of the order.

SCLERACHNE PUNCTATA.

[15]

TAB. IV.

SCLERACHNE.

CHAR. GEN. *Spicæ* androgynæ, fasciculatæ, singulæ involucro monophyllo foliaceo inclusæ; constantes locustæ inferiore (rarò duâbus) femineâ sessili, et alterâ pedicellatâ masculâ, utrâque biflorâ.

♂ *Gluma* bivalvis, subæquivalvis, nervosa, herbacea; *flosculo* utroque bivalvi, membranaceo.

♀ *Gluma* bivalvis; *valvula exterior* dimidio inferiore cartilagineo pedicellum locustæ masculæ amplectens, margine altero equitante, superiore compresso semiherbaceo nervoso; *interior* membranacea, acuminata. *Flosculus inferior* univalvis, neuter, glumæ interiori analogus; *superior* femineus, bivalvis, valvis angustatis acutis, stylo bifido, stigmatibus hispidulis.

Gramen *glabrum*; *culmo ramoso, geniculato*; *foliis planis*; *ligulâ brevissimâ, ciliatâ*. Affinitate proxima *Coici arundinaceæ*, Willd., quæ genus proprium (CHIONACHNE) efformat, a *Coice* diversum defectu veri involucris osseo-cartilaginei; in hac plantâ enim involucrum auctorum gluma inferior locustæ femineæ est, ut in *Sclerachne*, a quâ *Chionachne* distinguitur præsertim figurâ et texturâ uniformi glumæ inferioris locustæ femineæ, et insuper spicâ locustis masculis pluribus, nec unicâ, etiam habitu.

SCLERACHNE *punctata*, R. Br.

POLYTOCA BRACTEATA.

TAB. V.

POLYTOCA.

CHAR. GEN. *Spicæ terminales* masculæ; *axillares* androgynæ, infrâ femineæ; utriusque rachi ad articulos singulos bifloros solubili. *Locustæ masculæ* bifloræ, biglumes; *stamina* 3; *squamulæ hypogynæ* 2. *Locusta feminea* biflora, biglumis: *Glumæ valvula exterior* cartilaginea, rachin angustam amplexens, nervosa, alata; *interior* lævis, acuminata: *flosculus exterior* univalvis, neuter, *glumæ interiori* analogus; *interior femineus*, bivalvis, lodiculâ nullâ.

Gramen elatum: foliis planis, margine asperis; ligulâ brevi, ciliatâ. Articuli feminei *spicæ androgynæ locustæ neutre, univalvi, herbaceæ, articulum amplexente, stipati.* *Locustæ masculæ flosculus superior paulò præcocior.* *Stigmata longissima, colorata.* Affinitate hinc *Tripsaco*, inde *Chionackni*. An hujus generis, vel forsân generis distincti valde approximati, *Coix heteroclita*, Roxb. Fl. Ind. 3, p. 572.

POLYTOCA *bracteata*, R. Br.

HEXAMERIA DISTICHA.

[26]

TAB. VII.

HEXAMERIA.

CHILOGLOTTIS V. APPENDICULA, *Blume Bijdr.* p. 296-7, ad MALAXIDEAS pertinens.

CHAR. GEN. *Perianthium* erectiusculum, sepalis lateralibus exteriorum basi connatis et infra productis in saccum maximum calcar obtusum emulans. *Labellum* e basi calcaris ortum, unguiculatum; laminâ supernâ indivisâ, basi bisetâ. *Columna* brevissima. *Anthera* (filamento insidens) stigmati bilobo parallela, bilocularis. *Massæ pollinis* cereaceæ, in singulo loculo tres (!), quarum duæ inferiores collaterales, tertia superior, omnes apice acuto affixæ corpusculo septiformi loculum longitudinaliter bipartienti et cum eodem deciduæ.

Epiphyta *nana*; caule *ramoso*; foliis *distichis*, *mucronatis*, basi in *petiolum* brevem *semivaginantem* *attenuatis*; racemis *terminalibus*, *paucifloris*.

OBS. *Hexameria* proxima *Chiloglottidi* esse videtur, forsâ ab ipsâ *Chiloglottidi serpyllifolia*, Bl. haud diversa; et si in hac necnon in *Appendiculis* omnibus septa mobilia decidua loculos antheræ bipartientia et massas pollinis affigentia exstant hæc tria genera conjungi merentur. Numerus ternarius enim massarum, quamvis characterem insignem, et in *Hexameriâ distichâ* constantem, præbens, minoris certè valoris considerari debet.

HEXAMERIA *disticha*, R. BR.

8)

PHALÆNOPSIS AMABILIS, Bl.

TAB. VIII.

PHALÆNOPSIS, *Blume Bijdr.*, p. 294. *Lindl. Orchid.*,
p. 213.

VANDEÆ, *Lindl. Orchid.*

CHAR. GEN. *Perianthium* explanatum, 5-partitum. *Labellum* ecalcaratum, unguiculatum: *luminâ* tripartitâ: *lobo medio* hastato, apice laciniis duâbus elongatis subulatis, dente intermedio obsoleto. *Columna* libera, cum ungue labelli continua; apice obliquo trifido. *Anthera* bilocularis. *Massæ pollinis* cereaceæ, duæ (in singulo loculo singula), subglobosæ, singulæ ex apice depresso funiculum elasticum exserentes, ope cujus apici dilatato processûs e glandulâ stigmatis orti separatim affixæ.

Herba *arboribus innascens*; caule *abbreviato foliato haud dilatato*; foliis (2—4) *oblongis explanatis rigidis*; scapo *elongato*; racemo *laxo*; floribus *speciosis albis*; sepalis *lateralibus interiorum dilatatis*; labelli ungue *apice tuberculo aucto*, laminæ lobis *lateralibus unguiculatis*; columnæ laciniis *lateralibus deflexis*, *glandulam stigmatis ejusque processum tegentibus*.

LOXOTIS OBLIQUA.

[102]

TAB. XXIV.

CHAR. GEN. *Calyx* tubulosus, 5-fidus, æstivatione valvata. *Corollæ personatæ labium superius* abbreviatum, bilobum; *inferius* productum, semitrilobum (lobis lateralibus nanis). *Stamina* inclusa; *antherifera* duo, antheris reniformibus. *Stigma* capitatum (vix divisum). *Capsula* subovata, calyce inclusa, bivalvis; *placentis* utrinque seminiferis.

Herba *alternifolia*, annua, erecta, *glabriuscula*. Caule *succulento*. Foliis *integerrimis*, basi *hinc altè excisis*. Racemis *subsecundis*, *indivisis*, *pedicellis solitariis*, *unibracteatis*, *floribus deflexo-porrectis*, *cæruleis*.

Obs. Genus ad tribum Cyrtandracearum pertinens, proximum Glossantho, quæ, foliis et inflorescentia omnino conveniens, vix aliter differt nisi staminibus quatuor antheriferis antheris in coronulam cohærentibus, calyce basi inæquali angulisque tubi altius alatis, labio corollæ inferiore indiviso et disco hypogyno completo.

LOXOTIS OBLIQUA.

a. *Calycis lacinie acutiusculæ tubo breviores.*

Antonia vel Loxotis. *R. Br. in Wallich, Pl. Asiat. Rar.*, vol. iii, p. 65, in obs. sub Aikinia.

Loxotis obliqua. *Benth. Scrophul. Ind.*, p. 57.

Wulfenia obliqua. *Wallich, Tent. Flor. Nepal.*, p. 45, t. 35. *Don, Flor. Nepal. Prodr.*, p. 92.

Rhynchoglossum obliquum. *Blume, Bijdr. Stuk. 14*, p. 741?

β. *Calycis lacinie acutissimæ tubo longiores.*

Loxotis intermedia. *Benth. Scrophul. Ind.*, p. 57.

Wulfenia intermedia. *Wallich, Cat. n. 408?*

DESCR. Herba annua, erecta, glabriuscula, pube brevi articulata acuta rara conspersa. *Radix* fibrosa. *Caulis* teres, succulentus, crassitie pennæ anserinæ, 3—12-uncialis, modo simplicissimus sæpius parum ramosus, ramis ad ortum cum ipsa basi petiolorum confluentibus sæpe duobus ¹⁰⁸³ v. tribus seriatis, inferiore (cauli proximo) præcociore. *Folia* alterna, petiolata, exstipulata, integerrima, membranacea, lætè viridia, circumscriptione subovata cum acumine brevi, inæquilatera, basi hinc altè et ad nervum usque excisa inde rotundata, venis primariis vix eminentibus costata, venulis immersis, ultimis obsoletis, utriusque pube in adultis rarissima instructa, quandoque glaberrima, 2½ usque 4 pollices longa. *Petiol*i antice canaliculati 1—1½ unciales. *Racemi* terminales et axillares, simplicissimi, secundi, 2—4-unciales, rari, basiflori, pedicellis alternis recurvis unifloris unibracteatis. *Bracteolæ* vel subtendentes vel ipso pedicello insidentes, angustissimè lineares subfiliformes, erectæ, virides, pedicellis breviores. *Flores* deflexo-porrecti, cærulei. *Calyx* liber, tubulosus, 5-gonus, viridis, subbilinearis, æqualis, persistens; *tubi* lateribus planis, angulis in aciem alamve angustissimam exstantibus, minutè denticulatis, basi obtusa æquali; *limbi laciniis* semilanceolatis acutis sæpius acutiusculis tantum, æstivatione valvata alabastro acuto. *Corolla* monopetala, lætè cærulea, glaberrima, personata; *tubus* calyce longior subcylindræus; *fauz* clausa arcu elevato transverso tenuissimè pubescenti ad basin utriusque labii, inferioris stria verticali brevi in areas duas pro receptione antherarum distincto: *limbus* bilabiatus: *labium superius* brevius, bifidum, respectu axeos racemi adscendens, lobis obtusis sinu rotundato; *labium inferius* lingulatum, cæruleum basi macula flava insignitum, apice semitrifido lobulis obtusis, lateralibus nanis. *Stamina* antherifera duo inclusa; *filamentis* sinibus lateralibus labii inferioris suboppositis, compressis, adscendentibus, glabris: *antheris* reniformibus, imberbibus, conniventibus leviterque coherentibus, violaceis, bilocularibus, loculis basi divaricatis apice confluentibus, utriusque valvula interiore brevior et angustior: *pollen* brevè ovale, obtusè trigonum. *Filamenta sterilia* tria, quorum duo sinibus

interlabialibus respondentia, apicibus simplicibus acutiusculis: *tertium* nanum axi labii superioris oppositum. *Ovarium* liberum, sessile, ovatum, compressum, glabrum, viride, utrinque sulco longitudinali axi placentæ lateralis respondenti insculptum, uniloculare, polyspermum: *placentis* duabus parietalibus, lateralibus, e lamina angusta ortis, bilobis, lobis planis, utrinque et per totam ferè superficiem ovuliferis. *Stylus* filiformis glaber pallidus, situ et longitudine staminum antheriferorum, diu persistens. *Stigma* depresso-capitatum indivisum papulosum stylo manifestè crassius. *Vaginula* hypogyna incompleta, venoso-striata, glaberrima, subtruncata sæpè lobata. *Capsula* deflexa, calyce persistenti arcuè cincta ejusque longitudine, ovata, compressa, utrinque sulco longitudinali insculpta, unilocularis, bivalvis. *Placentæ* parietales duæ, axibus valvularum lamina angusta in lamellas duas fissili adnatæ, bilobæ, lobis expansis utrinque seminiferis. *Semina* undique versa, parva, numerosissima, elliptico-oblonga utrinque acuta, basi funiculo brevi crassiusculo pallido affixa; *testa* crassiusculo-membranacea; *membrana interna* obsoleta. *Embryo* exalbuminosus, aqueo-pallidus, cavitatem testæ replens, rectus; *cotyledones* breves; *radicula* teretiuscula umbilicum attingens.

I observed the plant here described and figured in the Island of Timor near Coepang, chiefly in shady places, but sometimes in more exposed situations, in April, 1803. The same species was several years after collected in many parts of Java by Dr. Horsfield, who states that it grows "in shaded situations at no great elevation above the ocean, near rivulets and ponds, rarely in forests; and that the native name is *Turu-pencheng*." If I am correct in referring *Rhynchoglossum obliquum* to our plant, it was observed also by Dr. Blume in mountainous situations of the same island, flowering in February.

By Dr. Wallich it was found in Nepal, Sylhet, and also in Martaban; for although he has considered the plant from the last-mentioned region as a different species, named *L. intermedia* in his list, and for which Mr. Benthams

proposed a specific difference, yet I find that the greater number of specimens collected in Nepal in 1819, and sent by Dr. Wallich to Sir Joseph Banks in the following year, have the deeply divided calyx with very acute and narrow segments characteristic of *L. intermedia*, while among the specimens from Martaban, in Dr. Wallich's Indian Herbarium, in the Museum of the Linnean Society, both states of calyx occur. But though I am not disposed to regard these differences in calyx as of specific importance, it is right to state that all the specimens which I have examined from Java and Timor, as well as those from Jurreepanee in the collection of Dr. Royle, agree in having the broader less acute and shorter segments of calyx, as represented in Mr. Bauer's figure, and also in that of Dr. Wallich.

With respect to the generic name *Loxotis* here adopted, it is that which I first gave in my manuscripts to the plant now described. This, however, I many years ago changed to *Antonia*, in compliance with the request of my lamented friend and fellow-traveller Mr. Ferdinand Bauer, to whom I was indebted for the figure here published. But as that name, by which it was introduced into a celebrated flower piece, painted in honour of the late Baron Jacquin at Vienna, and well known to the botanists of that capital, was never otherwise made public, and as *Antonia* of Pohl since published in his work on the "Plants of Brazil (vol. ii, p. 13, tab. 109)" is sufficiently established as a genus, I have been obliged to recur to my original name, under which indeed it has already appeared in Mr. Bentham's "Essay on Scrophularinæ Indicæ." The name *Loxotis*, however, may now be objected to from its too close resemblance in sound and identity of meaning, to *Loxonia*, another genus of the same family, more recently established by Dr. Jack; and the specific name *obliqua* is hardly less exceptionable, being merely a translation of that of the genus. This difficulty would be easily removed were it absolutely certain that *Rhinchoglossum* of Dr. Blume was identical with *Loxotis*; but from some of the characters ascribed to it I am not entirely satisfied that such is the case; and, indeed, as it is arranged by its author with *Rhinantheæ*, had I not re-

marked that it had been referred with a doubt to *Cyrtandraceæ* by Professor Lindley, in his enumeration of the genera belonging to that family, I should probably have overlooked it altogether, as I had previously done with respect to *Epithema* of Blume, which the author included in *Primulaceæ*, though unquestionably the same genus with my *Aikinia*, also belonging to *Cyrtandraceæ*.

With regard to the genus itself, it may be doubted whether *Loxotis* and *Glossanthus* ought to be generically distinguished merely or chiefly on account of the difference in the number of their antheriferous stamina, especially as they entirely agree in habit, in which there is something peculiar. It is not a little remarkable, that in some of the more minute and less important differences between them, the intermediate structure or connecting link should be found in a species sent by Dr. Schiede from Mexico (*Glossanthus Mexicana*, Br. ined.), and that this should be the only plant belonging to *Cyrtandraceæ* hitherto observed in any part of America.

TAB. XXIV. *Fig. 1. Loxotis obliqua*, natural size. *Fig. 2.* A front view of a flower, slightly magnified. *Fig. 3.* An opposite view of corolla only. *Fig. 4.* Corolla laid open, showing the antheriferous and the two lateral barren stamina (the minute rudiments of the 5th omitted). *Fig. 5.* Calyx, after the falling of corolla, with the persistent style and stigma. *Fig. 6.* An antheriferous stamen. *Fig. 7.* Pistillum separate and magnified (the hypogynous incomplete disk wanting). *Fig. 8.* Upper part of style with the slightly and unequally bilobed stigma. *Fig. 9.* Capsule with its persistent style, natural size. *Fig. 10.* The same magnified. *Fig. 11.* Capsule after bursting, showing the form of one of the parietal placentæ. *Fig. 12.* A placenta separate. *Fig. 13.* A transverse section of a capsule, showing the origin of the placenta and insertion of seeds on both surfaces. *Fig. 14.* A valve of the capsule with its placenta, from which the seeds have been removed. *Fig. 15.* A side view of the same. *Fig. 16.* A seed, with its testa. *Fig. 17.* A seed, deprived of its testa. *Fig. 18.* The embryo.

105]

LOXONIA ACUMINATA.

TAB. XXV.

LOXONIA, *Jack in Linn. Trans.*, xiv, p. 40.

CHAR. GEN. *Calyx* 5-partitus æqualis, æstivatione valvata. *Corolla* bilabiata; *fauce* aperta; *labio superiore* semibifido; *inferiore* trifido. *Stamina antherifera* 4, didynamia, exserta; *antheris* reniformibus, per paria approximatis. *Stigma* bilobum, obtusum. *Capsula* ovata, calyce persistente longiore cincta, bivalvis; *placentis* utrinque seminiferis.

Herbæ v. Suffrutices. *Folia opposita, inæqualia: majore inæquilatere, dentato; altero nano, integerrimo, stipuliformi. Racemi subcymosi, recurvi, sæpè bifidi, ex alis foliorum stipuliformium.*

Obs. Genus Cyrtandracearum a b. Jack conditum et ad eandem sectionem cum *Loxoti* referendum, sed affinitate arctius junctum cum *Stauranthera* in quâ folia opposita nec alterna, pari modo inæqualia, altero nempe nano stipuliformi; altero inæquilatere.

LOXONIA acuminata, pubescens, foliis lanceolato-oblongis acuminatis denticulatis; nanis reniformibus, racemis subbifidis folio brevioribus.

DESCR. Suffrutex parcè tenuissimèque pubescens. Caulis erectus vel adscendens, ramis teretibus parum flexuosis, pube subappressa. Folia opposita, quorum alterum lanceolato-oblongum acumdatum inæquilaterum, denticulis brevissimis, subsessile, ni fallor aversum, basi hinc (lateris inferioris) obtusâ rotundatâ, superioris et angustioris excisâ, costatum, venis primariis alternis arcuatis intra marginem deliquescentibus anastomosantibus, secundariis tertiariisque divaricatissimis, ultimis reticulantibus immersis, 5—8

pollices longum $2\frac{1}{2}$ —3 pollicis latum, membranaceum, viride, subtus pallidius, in cujus alâ rudimentum minutum gemmæ foliaceæ: alterum nanum stipuliforme sessile reniforme latius quam longum integerrimum, e cujus alâ pedunculus folio majore brevior, nano ipso multoties longior. *Racemus* subbifidus, ramis 3—5-floris, bracteis subulatis. *Calyx* 5-partitus: *laciniæ* lanceolatæ acuminatæ herbaceæ pilosæ pilis crebris articulatis acutis patulis, longitudine æquales, duæ laterales superiorum paulum angustiores, trinerviæ, æstivatione valvatâ alabastro ovato acuminato, acuminum apicibus distinctis. *Corolla* monopetala bilabiata, calyce parum longior, glabra; *tubo* ventricoso calyce brevior; *fauce* apertâ infernè barbatâ; *labio superiore* semibifido plano, lobis obtusis; *inferiore* profundè trifido, lobis obtusiusculis integerrimis, medio paulo longiore. *Stamina* *antherifera* quatuor didynama exserta limbo tamen breviora. *Filamenta* subulata conniventia glabra, duo inferiora longiora. *Antheræ* conformes per paria approximatae et leviter cohærentes, imberbes, reniformes, loculis divaricatis apice confluentibus medio anticè longitudinalitèr dehiscentibus. *Pollen* subglobosum. Rudimentum staminis quinti brevissimum absque antheræ simulacro. *Ovarium* liberum sessile ovatum glaberrimum disco annulari angustissimo imberbi basi cinctum, uniloculare, placentis duabus parietalibus bifidis lobis divaricatis planis utrinque ovuliferis. *Stylus* cylindraceus glaber longitudine circiter staminum. *Stigma* obtusum papulosum obsoletè bilobum stylo parum crassius. *Capsula* ovata cuspidata membranacea calyce persistenti brevior unilocularis bivalvis, cuspidè bipartibili; valvulæ medio placentiferæ; placenta singula utrinque dilatata in laminam rectam marginibus utrinque seminiferis. *Semina* numerosissima minuta obovata sessilia, sæpè angulata mutuâ pressione, castanea: testa membranacea tenax: membrana interna tenuissima albumini parco v. parciissimo adhærens. *Embryo* rectus aqueo-pallidus subovatus, cotyledonibus brevissimis.

I have considered the plant here figured as distinct from

Loxonia hirsuta of Jack, which, however, it appears from his description to resemble in so many points that it may actually belong to the same species, differing only somewhat in the form of the leaves and in being less pubescent. Dr. Jack did not find his plant in fruit, neither did he ascertain the dehiscence of the capsule in *L. discolor*, from which the character of the genus was formed. In both species he describes the ovary as bilocular, and the lobes of the placentæ as revolute.

Dr. Horsfield found this plant in 1818, when he accompanied Sir Stamford Raffles on his journey from Padang—one of the principal stations on the west coast of Sumatra—to the Menangaboo country, growing on the ranges of hills which extend parallel to the coast from N.W. to S.E., in shaded forests between 500 and 1000 feet above the level of the ocean. He did not observe it in Java.

TAB. XXV. *Fig. 1. Loxonia acuminata*, natural size. *Fig. 2.* A flower, magnified. *Fig. 3.* An anther, with a portion of the filament. *Fig. 4.* Style and stigma. *Fig. 5.* Capsule surrounded by the calyx. *Fig. 6.* Capsule after dehiscence, the calyx being removed. *Fig. 7.* One of the valves of the capsule. *Fig. 8.* Transverse section of the ripe capsule. *Fig. 9.* A seed. *Fig. 10.* The embryo.

CYTANDRACEÆ, to which *Loxonia* and *Loxotis* belong, was established in 1822¹ by the late Dr. Jack as a natural order, according to him most nearly allied to *Bignoniaceæ*, but differing sufficiently from that family in the structure of its fruit, especially in the placentation of its minute seeds.

The existence or absence of albumen in the ripe seed is not expressly stated in his character of the order, nor is it noticed in the description of any of the species he has referred to it. It may, however, be presumed that he

¹ 'Linn. Soc. Trans.,' vol. xiv. p. 23.

believed the whole family to agree in this respect with *Bignoniaceæ*, next to which he had placed it; and he had at least the proof of the absence of albumen in Roxburgh's figure of his *Incarvillea parasitica*, in which the structure of ovarium as well as of seeds is correctly given.

The genera referred by Dr. Jack to this new family are *Cyrtandra* of Forster, *Didymocarpus* of Wallich, and ¹⁰⁷ two other genera for the first time proposed, namely, *Æschynanthus*, to which Roxburgh's *Incarvillea parasitica* belongs, and *Loxonia*.

Nearly about the same time Mr. Don published his *Didymocarpeæ*,¹ a family consisting of two of Dr. Jack's *Cyrtandraceæ*, namely, *Didymocarpus* and *Æschynanthus* (his *Trichosporum*), and a new genus *Lysionotus*. He in like manner depends chiefly on placentation, which is not very clearly described; and he also introduces into his character the absence of albumen, the pendulous position of seeds, and the undivided stigma. From this family he excludes *Cyrtandra*, which has, he states, erroneously however, a copious albumen; while *Chirita* of Dr. (Buchanan) Hamilton, of which Dr. Jack's *Didymocarpus* includes at least one species, is doubtfully referred to *Scrophularinæ*, chiefly on account of its bilamellar stigma.

In 1826 Dr. Blume¹ refers *Cyrtandraceæ* to the natural order *Bignoniaceæ*, distinguishing it as a tribe from true *Bignoniaceæ* by its pendulous seeds, and subdividing it into two sections; the first, *Trichosporæ*, with capsular fruit, and seeds either winged or with some other form of appendage; the second, *Cyrtandraceæ*, having baccate fruit and seeds without appendage. To each of these sections he has added more than one new genus, but *Loxotis* and *Loxonia* are not included in either of them; though *Loxotis*, as I have already stated, is probably his *Rhynchoglossum*, referred by him to *Rhinantheæ*; and *Loxonia*, notwithstanding his account of the fruit, may possibly be his *Loxophyllum*, which he has placed in *Scrophularinæ*;

¹ 'Edinburgh Philosophical Journal,' vii. p. 83, and 'Prodromus Floræ Nepalensis,' p. 121.

² 'Bijdragen,' p. 759.

while his *Epithema*, my *Aikinia*, also belonging to *Cyrtandraceæ*, he refers to *Primulaceæ*.

It is somewhat remarkable that none of these writers should have adverted to the affinity of this new family to *Besleriaceæ* of Richard and De Jussieu, now generally named *Gesneriaceæ*. This affinity, however, did not escape Dr. Von Martius, who in his elaborate account of *Gesneriaceæ*, published in 1829,¹ considers *Cyrtandraceæ* as sufficiently distinct from that order in the absence of albumen and in having an inverted embryo: the latter character he states on the authority of Mr. Don, who, in employing the term "Embryo inversus," can only have intended to express its direction with respect to pericarpium; such at least is the real structure of those genera which he referred to his *Didymocarpeæ*, and it is certain that in the relation of embryo to hilum both families entirely agree.

Dr. Von Martius also notices the difference in the order of abortion of stamina between these two families, which is no doubt generally true, but admits in each of at least one exception; *Sarmienta* in *Gesneriaceæ*, agreeing with *Cyrtandraceæ* in having only its two anterior or lower stamina antheriferous: and in this latter family *Aikinia* or *Epithema*, which, as in the greater part of diandrous *Gesneriaceæ*, has its two posterior or upper stamina perfect.

There is indeed another, and that a very remarkable, distinction noticed in the position of the lobes of the stigma, which in *Gesneriaceæ*, according to Von Martius, are placed right and left in relation to the parts of the flower, and consequently opposite to the lateral parietal placentæ; while in *Cyrtandraceæ* the lips of the stigma—for so it is necessary to express the fact in this family—are anterior and posterior, and therefore alternate with the lateral placentæ; the latter being the ordinary relation in unilocular ovaria, where the placentæ and lobes, or rather lips, of stigma, correspond in number. This difference, however, even were it fully established, would hardly be available here as a technical distinction, several genera in each

¹ 'Nov. Gen. Pl. Bras.,' iii. p. 72.

family having an undivided stigma; unless in such cases the position of the confluent parts could be determined by that of the two vascular cords generally observable in the style, and continued into the axes of the lobes of a regularly bifid stigma, when belonging to an ovarium composed of two carpels. But even if this distinguishing character should be admitted to be general, it is certainly not without exception; and in the only cases that I have examined in *Gesneriaceæ*, where the lateral position of the lobes of stigma may be supposed to exist, the apparent position arises from the extreme breadth and manifest division of the lips, the two vascular cords of the style being still anterior and posterior.¹

The only point of difference remaining, therefore, is ^[108] the existence of albumen in *Gesneriaceæ* and its absence in *Cyrtandraceæ*. This character, however, is not absolutely constant, there being cases in *Cyrtandraceæ* where the ^[109] remains of albumen are visible in the apparently ripe seed; and in several *Gesneriaceæ* it exists so sparingly as to become a character of very little value,² especially as it is not here connected with other more important differences.

¹ Here follows a note "On the relative position of the Divisions of Stigma and Parietal Placentæ in the Compound Ovary of Plants," which, having been originally distributed also in a separate form, has been already given in vol. i. pp. 553—563.—*EDIT.*

² The late Correa de Serra, in a very ingenious essay published in 1811,¹ endeavoured to establish a test for ascertaining the importance of albumen in relation to the affinities of plants, namely, that where the albumen is of a texture very different from that of the embryo, which does not absorb it in germination, its constancy may be depended on; while in these cases where its texture is nearly similar to that of the embryo, which derives from it its earliest nourishment, its presence or absence becomes of little value. His hypothetical expression of this difference is, that in the latter case the embryo before germination converts part of a uniform substance into its own body, and in germinating derives nourishment from the remainder; in the former it selects what is suited for its nourishment, leaving a residuum which it does not afterwards act upon, and whose presence is therefore constant. Among the examples given of families in which this selection and residuum exist are *Gramineæ*, *Palmæ*, *Nyctagineæ*, *Caryophylleæ*, and *Euphorbiaceæ*.

Soon after the publication of this essay a paper was read before the Linnean Society of London, in which I endeavoured to prove that the test attempted to be established by Correa was liable to many exceptions, and that his hypo-

¹ 'Annales du Muséum,' xviii. p. 206.

110] In describing the genus *Aikinia* (*Epithema* of Dr. Blume) I regarded *Cyrtandraceæ*, or *Cyrtandreæ*, for the reasons now assigned, as a tribe merely of *Gesneriaceæ*,
 111] distinguishable from that portion of the order with hypogenous corolla, or *Beslerieæ*, by characters either of little importance or which required confirmation. For although, in addition to the characters referred to, *Cyrtandreæ* differ very remarkably in geographical distribution from the rest of the family, yet this difference is not entirely without exception, as I have already noticed in my account of *Loxotis*.

But whether these groups be considered as distinct families, or as tribes only, it will probably be admitted that in a natural classification *Cyrtandreæ* must stand next to *Beslerieæ*; while on the other hand they appear to be very nearly related to *Bignoniaceæ*, with which they are connected through *Incarvillea*, particularly with that section of it which in Dr. Royle's Illustrations I have described as a sub-genus, and named *Amphicome*. This sub-

theoretical expression of the facts was not applicable even to all the families he has cited in support of it. And I concluded that as a general rule the point most to be depended on in proving the importance of albumen in systematic botany was its relative quantity, especially when accompanied with a low degree of development of embryo; for where the albumen forms the great mass of the seed in any known portion of a natural family, it may in most cases, though not always, be safely inferred, not only to be present, but to exist in like proportion in the whole of that family. This rule, however, I regarded as merely empirical, founded on extensive experience, but not necessarily connected either with uniformity or even apparent importance of function; for while in some families in which its proportion to the whole body of the seed is the greatest, it constitutes the early nourishment of the embryo, in others, where it exists in equal quantity, it is either not at all or but slightly acted upon in germination. I stated also that there were cases in which this character was of reduced importance, existing only in certain tribes of one and the same great natural family, as in *Rubiaceæ*; nor are there wanting instances in which it is only of generic value.¹ And, lastly, I noticed that in several families, in which the constancy of the character was very general, exceptions occurred, dependent on the apparent necessity for an unusual development and increased energy in the embryo, connected either with the unfavorable circumstances in which it was destined to vegetate, as in plants growing in or exposed to the action of salt-water; or where great resistance, arising from the structure of the pericarpium, or even from the texture of the proper integuments of the seed itself, was to be overcome in germination.

¹ 'Linn. Soc. Trans.,' vol. x. p. 36 [vol. i. p. 23], et 'Prodr. Flor. Nov. Holl.,' vol. i. *passim*.

genus Professor Lindley has lately considered generically distinct from *Incarvillea*. But except those differences in the seeds and calyx, which I have regarded, and still consider, as only of sectional value, I find no other characters whatever in the flower or fruit to justify the separation; for in both species of *Amphicome*, as in the original *Incarvillea*, the ovarium as well as the capsule is certainly bilocular, and not unilocular, as it is described in the species figured in the 'Botanical Register' (for 1838, t. 19), and the two subgenera entirely agree in the peculiar structure of the anthera, the spur of each of whose loculi originates not on the back, but front of the cell, in the line of dehiscence, which it limits.

The following characters of *Gesneriaceæ*, and of the three tribes of which, according to my view, it consists, may serve to distinguish the family from the nearly related orders, and the tribes readily from each other.

GESNERIACEÆ, Richard et de Jussieu.

Calyx 5-divisus, æqualis (raro parùm inæqualis). *Corolla* monopetala, irregularis, limbo 5-lobo, æstivatione imbricata. *Stamina* antherifera 2 v. 4, cum v. absque quinti postici rudimento. *Ovarium* (liberum v. adnatum) uniloculare (nunc approximatione placentarum quasi biloculare); basi disco lobato v. indiviso cinctum; *placentis* 2 parietalibus lateralibus (sæpiùs bilamellosis) polyspermis; *ovulis* anatropis. *Pericarpium* capsulare v. baccatum. *Semina* parva (raphe nulla), albuminosa v. exalbuminosa; albumine carnosio, molli, copioso v. parco. *Embryo* rectus, axilis, orthotropus, dimidium albuminis dum adsit æquans v. superans.

Herbæ v. *Suffrutices* foliis *simplicibus*, *indivisis*, *exstipulatis*, *oppositis*, *verticillatis alternisve*, sæpiùs *serratis crenatisve nunc integerrimis*, in *plerisque pube simplici*, *acuta v. capitata*. *Inflorescentia varia*.

GESNERIÆÆ.

Calyx cum ovario plùs minùs connatus. *Pericarpium* capsulare. *Semina* albumine copioso.

BESLERIÆ.

Calyx liber. *Pericarpium* baccatum v. capsulare. *Semina* albuminosa.

CYRTANDRÆ.

Calyx liber. *Pericarpium* capsulare v. baccatum. *Semina* exalbuminosa v. albumine parco.

OBS. The following remarks relate to the modifications of the different parts of fructification in *Cyrtandrea* and their relative importance in characterising genera.

The CALYX admits of every degree of depth of division. Its segments are generally acute, always so when divided to the base; and wherever acute the æstivation appears to be valvular. In several cases where the calyx is tubular, particularly in the greater part of true *Didymocarpus*, and in one of the sections of *Æschynanthus*, the segments are rounded, and the æstivation necessarily overlapping. The tubular calyx, when accompanying an elongated capsule, is generally thrown off or separates transversely at the base, except in the few cases in which the capsule is pedicellated. The deeply divided calyx of the genera with elongated capsules is persistent; as it also is, whether deeply divided or tubular, in all those with short capsules and in the ¹¹³⁷ baccate genera. None of these modifications appear to be of generic value, though some of them form the principal characters of very natural sections of genera.

The tube of the COROLLA varies greatly in length, but its various proportions, either with respect to the limb or calyx, are seldom of generic importance; a difference of this kind, however, forms the only distinguishing character between *Bæa* and *Streptocarpus*. The limb varies considerably in form, direction, and proportion of lips; but the most important modification occurs in *Glossanthus* and *Loxotis*, in both of which the lateral lobes of the lower lip are obsolete; or, in the former genus, perhaps, altogether

wanting. This modification is necessarily accompanied by a different æstivation, which in all the other genera is quincuncial, the lateral lobes of the lower lip overlapping the upper, which covers the middle lobe of the lower. The spur of the tube is found only in one of the two species of *Stauroanthera*.

The number of antheriferous STAMINA, or the difference between the diandrous and didynamous plants of the tribe, is not always of much value; for in *Didymocarpus* there are several didynamous species which certainly do not form a natural section. *Stamina inclusa* and *exserta* generally mark distinct genera, but yet not in all cases. The difference between parallelism and divarication of the lobes of antheræ is always, I believe, of generic importance. The various degrees of confluence of the divaricate lobes, and the apparently peculiar dehiscence in some of its modifications, seem not to be of equal value.

The STIGMA exhibits various remarkable differences, some of which are considerable, though not always of equal value in the definition of genera. The most important of these, and which hitherto has been overlooked, is the abortion, or great reduction in size, of the upper lip, while the lower is proportionally dilated, and, in some cases, deeply divided. Thus in *Chirita*, in which the stigma is described as bilamellar, both lamellæ belong to the lower lip. And in many species of *Didymocarpus* the apparent obliquity of stigma arises from the abortion of the upper lip, and the lamellar expansion of the lower, which, however, is never divided as in *Chirita*. In several genera the lips are equal, and either lamellar or so short as to be hardly distinguishable; in other cases there is no trace of division. These different modifications, in most cases, mark the limits of genera.

As some of the most important characters of the tribe reside in the structure of OVARIIUM and PERICARPIUM, so the principal natural divisions are founded on modifications of the same organ. The ovarium may be in all cases described as properly unilocular: though, from the approximation and slight cohesion of the parallel portions of the

inflected parts, or, as they are commonly called, placentæ, it not unfrequently appears to be bilocular. But this cohesion only occurs when the production of ovula is confined to the upper or inner surface of the carpel, which is the case in the greater part of the genera with elongated capsules; for where both surfaces are ovuliferous, as in the baccate and most of the genera with included capsules, no such cohesion can take place. The general direction of the margins of the placenta of each component part of ovarium and pericarpium may at first appear a deviation from the ordinary structure, the general rule being that the margins only unite to form a complete cell, whereas the completion of the cell in that manner is incompatible with the direction of these margins, which in each carpel are turned from, not towards each other. This difference, however, is more apparent than real, and the structure in *Cyrtandrea* may be justly compared with that of such genera of other families as have the placenta of a multi-locular, or that of the single distinct, carpel projecting considerably into the cavity.

The great elongation of pericarpium in many of the genera having capsular fruit, is the more remarkable in *Cyrtandrea*, as there is no instance of similar elongation, or any approach to it, in either of the two other tribes of *Gesneriaceæ*. In this elongation of capsule, however, they approach to *Bignoniaceæ*, where it is both more general and exists in a still greater degree.

In most of the *Cyrtandrea* with elongated capsules, the valves, though membranaceous, are perfectly straight, but in a few others they are spirally twisted, though nearly of the same texture. The spiral torsion of the valves certainly does not depend on the length of the capsule merely, the greatest length being found conjoined with straight valves, as in *Eschynanthus*; nor is it the consequence of drying, for the twisting in all cases commences long before the ripening of the fruit. The mechanism explaining these differences is, however, in general obvious. In the twisted valve the endocarp consists of a stratum of vertically elongated fibres, with an extremely thin or hardly manifest

inner membrane, while in the valve of the straight capsule the vertically elongated stratum has a manifest inner covering, consisting of transversely elongated cells, which no doubt counteract the tendency to torsion of the longitudinal fibres. But the transversely elongated cells are found in the inflected or placentiferous portion of the elongated capsule, both in the twisted and straight-fruited genera; and their function here seems to be to determine the involution or revolution of the ovuliferous margin, which in these fruits is probably necessary for the protection of the seeds even after dehiscence.

The more remarkable differences in placentation are almost always important: thus, in many genera the ovula are produced on the inner surface of the margin only; in others both surfaces are equally productive: and in some, the production, instead of being confined to the marginal region, extends over the whole of the inflected and included portion of the carpel.

The SEEDS are generally pendulous, but in a few genera, as *Epithema* and *Loxocarpus*, erect; and in some others they vary in pericarpial direction, according to their different heights in the same placenta. They are always minute, generally oval, oblong, or nearly cylindrical, and inserted at or very near one extremity; in most cases sessile, or nearly so, but in a few furnished with a very long and extremely slender funiculus.

Although the ovulum is anatropous, there is no apparent raphe in the ripe seed. The capillary appendages existing in some cases at both extremities of the seed vary considerably in number and form in that genus where they are most remarkable, namely *Eschynanthus*, in which, as well as in *Agalmyla*, and probably in *Tromsdorffia*, they are mere appendages, performing no other function; but in *Lysionotus* the upper hair in the pendulous seed is in reality its funiculus or attenuated base.

The integument of the ripe seed is, in most cases, apparently simple; but in a few, especially *Eschynanthus*, the inner membrane is easily separable from the testa.

Before the complete ripening of the seed, the semifluid

remains of ALBUMEN are generally obvious; and even in the ripe seed, in several cases, slight traces of it are visible: in *Rhabdothermus* it is more abundant, and of firmer consistence.

CYRTANDREARUM SYNOPSIS GENERUM.

A. PERICARPIUM CAPSULARE.

† *Capsulæ* elongatæ. *Semina* utrinque appendiculata.

a. *Antheræ* exsertæ, inappendiculatæ, loculis linearibus parallelis. *Semina* pendula, extremitate superiore nuclei (nec appendicis) affixa.

ÆSCHYNANTHUS. *Stamina* antherifera 4. *Stigma* indivisum, dilatatum.

TROMSDORFFIA. *Stamina* antherifera 4. *Stigma* bilamellatum.

AGALMYLA. *Stamina* antherifera 2. *Stigma* bilamellatum.

β. *Antheræ* (2) inclusæ, dorso appendiculatæ. *Semina* pendula extremitate setæ superioris affixa.

LYSIONOTUS.

†† *Capsulæ* elongatæ. *Semina* inappendiculata, sessilia. *Antheræ* inclusæ, loculis divergentibus.

CHIRITA. *Stamina* antherifera 2. *Stigma* labio superiore obsoleto; inferiore bilamellato. *Capsula* valvis strictis (nec spiritaliter tortis).

DIDYMOCARPUS. *Stamina* antherifera 2—4. *Stigma* indivisum (sæpè ex abortione labii superioris obliquum). *Capsula* valvis strictis.

STREPTOCARPUS. *Stamina* antherifera 2. *Capsula* valvis spiritaliter tortis. *Corollæ* tubus calyce duplò multotiesve longior.

BÆA. *Stamina* antherifera 2. *Capsula* valvis spiralitèr t^{or}tis. *Corollæ* tubus calycem vix æquans.

+++ *Capsulæ* calyce longiores, hinc longitudinalitèr dehiscentes. *Semina* erecta, funiculis clongatis.

LOXOCARPUS.

++++ *Capsulæ* subovatæ, calyce inclusæ: *placentis* utrinque seminiferis.

EPITHEMA. *Capsula* circumscissa. *Stamina* duo superiora antherifera!

STAUANTHERA. *Capsula* circumscissa. *Stamina* antherifera 4. *Calyx* 5-fidus, sinibus plicatis!

LOXONIA. *Stamina* antherifera 4, exserta. *Capsula* bivalvis. *Calyx* 5-partitus.

GLOSSANTHUS. *Stamina* antherifera 4, inclusa. *Calyx* 5-fidus, tubo 5-gono. *Capsula* bivalvis.

LOXOTIS. *Stamina* antherifera 2, inclusa. *Calyx* 5-fidus, tubo angulato. *Capsula* bivalvis.

MONOPHYLLÆA. *Stamina* antherifera 4. *Calyx* 5-partitus, æstivatione imbricatâ. *Capsula* ruptilis?

PLATYSTEMMA. *Stamina* antherifera 4, exserta. *Calyx* 5-fidus, æstivatione valvatâ. *Corollæ* tubo brevissimo, limbo patenti. *Capsula*?

RHABDOTHAMNUS. *Stamina* antherifera 4. *Corolla* tubo campanulato. *Stigma* bilobum. *Capsula* 4-valvis. *Semina* albuminosa!

B. PERICARPIUM BACCATUM.

FIELDIA. *Stamina* antherifera 4, antherarum loculis parallelis. *Calyx* 5-partitus. *Corolla* tubulosa. *Stigma* bilobum. *Bacca* exsucca.

RHYNCHOTHECUM. *Stamina* antherifera 4, antheris bivalvibus, valvulâ interiore minore. *Calyx* 5-partitus.

CENTRONIA. *Stamina* antherifera 4, antheris basi calcaratis. *Calyx* spathaceus. *Bacca* siliquæformis.

CYRTANDRA. *Stamina* antherifera 2, inclusa, antherarum loculis parallelis. *Calyx* 5-fidus.

WHITIA. *Stamina* antherifera 2, semiexserta, antherarum loculis haud parallelis. *Calyx* 5-partitus.

ÆSCHYNANTHUS, Jack in Linn. Trans. 14, p. 42.

CHAR. GEN. *Calyx* 5-divisus (tubulosus v. partitus). *Corolla* bilabiata. *Stamina* antherifera 4, antherarum loculis parallelis. *Stigma* indivisum, dilatatum, sub-infundibuliforme. *Capsula* elongata, valvis strictis. *Semina* pendula, apice nucleî affixa, utrinque pilifera, pilo inferiore unico; superiore unico, duplici pluribusve.

Suffrutices in arborum cortice radicantes; foliis oppositis aequalibus, integerrimis, coriaceis venis obsoletis.

† *Calyx* tubulosus, ore 5-lobo obtuso. *Semina* utrinque monotricha, pilo superiore basi dilatata.

1. *Æschynanthus volubilis*, Jack, l. c. t. 2, f. 3.

2. *Æschynanthus radicans*, Jack, l. c. p. 43.

3. *Æschynanthus parvifolia*, calyce pilosiusculo sub-campanulato, foliis elliptico-lanceolatis glabris.

Loc. Banca, 1813. Dr. Horsfield.

†† *Calyx* tubulosus 5-dentatus acutus. *Semina* . . .

4. *Æschynanthus fulgens*, Wall. List, No. 797.

††† *Calyx* 5-fidus acutus. *Semina* extremitate superiore ditrichâ.

5. *Æschynanthus parasitica*, Wall. List, No. 796.

6. *Æschynanthus ramosissima*, Wall. List, No. 799, and Pl. Asiat. Rar., 1, p. 55, t. 71.

7. *Æschynanthus Griffithii*, calyce 5-partito glabro: laciniis lanceolatis, filamentis hirsutis, foliis lanceolatis.

Loc. Tavoy? D. Griffith.

++++ *Calyx* 5-partitus v. altè 5-fidus, acutus. *Semina* utrinque monotricha, pilo superiore basi simplici.

8. *Æschynanthus Horsfieldii*, glabra, calyce 5-partito : laciniis lineari-lanceolatis, seminum pilis subulatis, foliis ovato-lanceolatis sub-acuminatis.

Loc. Java, an. 1814. Dr. Horsfield.

9. *Æschynanthus bracteata*, Wall. List, No. 794.

10. *Æschynanthus acuminata*, Wall. List, No. 6397.

+++++ *Calyx* abbreviatus, cyathiformis, dentatus. *Semina* utrinque monotricha, pilo superiore basi dilatâtâ.

11. *Æschynanthus Wallichii*.

Æschynanthus radicans, Wall. List, No. 798, non Jack.

Loc. Singapore.

+++++ *Calyx* *Semina* extremitate superiore comosâ, pilis indefinitè numerosis.

12. *Æschynanthus longicaulis*, Wall. List, No. 888.

TROMSDORFFIA.

Tromsdorffia Sp. *Blume, Bijdr.*, p. 762.

CHAR. GEN. *Calyx* 5-fidus, acutus. *Corolla* tubulosa, bilabiata. *Stamina* antherifera 4; antheris exsertis, loculis parallelis. *Stigma* bilamellatum (labiis æqualibus). *Capsula* . . . *Semina* . . .

Herba radicans; foliis alternis dentatis.

Tromsdorffia? elongata, Blume, *Bijdr.*, p. 763.

AGALMYLA.

Agalmyla Sp., *Blume, Bijdr.*, p. 766.

CHAR. GEN. *Calyx* 5-partitus, acutus. *Corolla* tubulosa, bilabiata. *Stamina* antherifera 2; antheris exsertis, loculis parallelis. *Stigma* bilamellatum (labiis æqualibus). *Capsula*

elongata, valvis strictis. *Semina* pendula, apice nuclei affixa, utrinque monotricha.

Herba radicans; foliis alternis dentatis.

Agalmyla staminea, Blume, Bijdr., p. 767.

LYSIONOTUS, *Don, Prodr. Fl. Nepal.*, p. 124.

CHAR. GEN. *Calyx* 5-partitus, acutus. *Corolla* bilabiata, fauce bicallosa. *Stamina* antherifera 2, inclusa. *Antheræ* dorso appendiculatæ, imberbes. *Stigma* indivisum. *Capsula* elongata, valvis strictis. *Semina* utrinque in pilum producta, apice pili superioris affixa.

Herba; foliis subverticillatis, serratis. Inflorescentia cymosa.

Lysionotus serratus, *Don, Prodr. Fl. Nepal.*, p. 124.

L. ternifolia, *Wall. Pl. Asiat. Rar.*, 2, p. 20, t. 118.

CHIRITA, *Buchan. Hamilt. in Don, Prodr. Fl. Nepal.*,
p. 89.

CHAR. GEN. *Calyx* 5-fidus, æstivatione valvatâ. *Corolla* tubulosa, bilabiata. *Stamina* duo antherifera; *antheræ* (sæpius barbatae) loculis divergentibus. *Stigma*: labio superiore abortiente v. nano; inferiore bilamellato. *Capsula* elongata, valvis strictis. *Semina* inappendiculata, pendula.

Herbæ v. Suffrutices; foliis oppositis sæpius inæqualibus, altero in quibusdam nano v. abortiente; pedunculis axillaribus.

1. *Chirita urticifolia*, foliis inæqualibus ovatis acutis serratis elongato-petiolatis, pedunculis subsolitariis medio bibracteatis, calycis laciniis subulatis, antheris imberbibus!

Chirita urticifolia, *Buchanan Hamilton MSS. in Don, Nepal.* 90.

Chirita grandiflora, *Wall. Pl. Asiat. Rar.*, 1, p. 43, t. 50

2. *Chirita flava*, Wall. List, n. 801. (*Calosacme*.) [117]
Chirita pumila, Don, *Nepal.*, p. 90.
3. *Chirita acuminata*, Wall. List, n. 802. (*Calosacme*.)
Incarvillea oblongifolia, *Roxb. Ind.*, vol. 3, p. 113.
4. *Chirita dimidiata*, Wall. List, n. 803. (*Calosacme*.)
5. *Chirita bifolia*, Don, *Prodr. Fl. Nepal.*, p. 90, Royle, *Illustr.*, p. 294, t. 70, f. 2.
6. *Chirita macrophylla*, Wall. List, No. 805 (*Calosacme*),
 & *Pl. Asiat. Rar.*, 1, p. 56, t. 72.
7. *Chirita Horsfieldii*, foliis ovato-oblongis acutis serratis scabris inæqualibus, bracteis orbiculatis calycibusque coloratis, stigmatis lamellis latioribus quam longis.
Didymocarpus barbata, Jack, in *Linn. Soc. Trans.*, 14, p. 38?
Tromsdorffia speciosa, Blume, *Bijdragen*, p. 762??
Loc. Java, an. 1814. Dr. Horsfield.
8. *Chirita scaberrima*, foliis ovatis acutis crenatis scaberrimis, stigmatis labellis longioribus quam latis.
Loc. Java orientalis. Dr. Horsfield.
9. *Chirita cærulea*, annua, foliis ovatis acutiusculis obsolete dentatis subæqualibus lævibus, pedunculis apice diphyllis: foliolis reniformibus basi hinc (sæpiùs) connatis, pedicellis ebracteatis subseriatis.
Loc. Java, an. 1814. Dr. Horsfield.
10. *Chirita hamosa*, foliis oppositis subæqualibus oblongis subovatisve, pedunculo communi cum petiolo connato; pedicellis seriatis bifidis simplicibusve.
Didymocarpus? *hamosa*, Wall. List, n. 788.
Loc. Troglä in Martabania. D. Wallich.

DIDYMOCARPUS, Wall. in *Malay. Misc.* 1, No. 5, p. 1.

CHAR. GEN. *Calyx* 5-divisus (v. tubulosus 5-fidus, 5-partitus). *Corolla* tubulosa, bilabiata. *Stamina* anthe-

rifera 2 (raro 4) inclusa; antheris imberbibus, loculis divergentibus. *Stigma* indivisum (sæpè, ex abortione labii superioris, obliquum). *Capsula* valvis strictis. *Semina* inappendiculata, sessilia.

Herbæ *pubescentes, caulescentes vel acaules; foliis serratis crenatisve petiolatis, caulinis oppositis, verticillatis alternisve; inflorescentia subcymosa.*

CLAVIS SPECIERUM.

A. DIANDRÆ.

† *Calyx* infundibuliformis, coloratus, tubo lobos superante, in plerisque basi transversim secedens.

α. *Calycis* lobi rotundati. *Stigma* obliquum (unilabiatum). Bractæ perfoliatæ, coloratæ.

1. *Caulescentes.*

Didymocarpus aromatica, villosa, oblonga, acuminata, punduana.

2. *Acaules.*

Didymocarpus macrophylla, pedicellata.

β. *Calycis* lobi acuti (glanduloso-pilosi).

Didymocarpus subalternans, obtusa.

†† *Calyx* 5-partitus acutus persistens.

α. Placentæ latitudine valvularum, marginibus ovuliferis longitudinalitèr revolutis.

1. *Caulescentes.*

Didymocarpus crinita, serrata, racemosa, corniculata, cordata, corchorifolia, reptans.

2. *Acaules.*

Didymocarpus missionis, Zeylanica.

118] β. *Capsula* hinc dehiscens, inde diù cohærens. Placentæ valvis angustiores.

Didymocarpus Rottleriana.

B. DIDYNAMÆ.

α. Caulescentes.

Didymocarpus frutescens, elongata.

β. Acaulis.

Didymocarpus lanuginosa.

SPECIERUM DIAGNOSES.

A. Diandræ. Calyce tubuloso infundibuliformi.

α. Calycis lobi rotundati.

1. *Didymocarpus aromatica*, foliis ovatis obovatisve inciso-crenatis, caule apice foliato petiolisque pube appressâ.

Didymocarpus aromatica, Wall. *Pl. Asiat. Rar.* 2, p. 34, t. 141, exclus. fig. 4—7, ad *Didymocarpum macrophyllum* pertinentibus.

Didymocarpus primulifolia, Don, *Nepal.*, p., 123.

Loc. Nepal.

2. *Didymocarpus villosa*, foliis obovatis inciso-crenatis adultis subsericeis, caule apice foliato petiolisque villis patulis hirsutissimo.

Didymocarpus villosa, Don, *Nepal.*, p. 123.

Loc. Nepal.

3. *Didymocarpus oblonga*, foliis oblongo-lanceolatis inciso-serratis adultis cauleque apice foliato glabriusculis.

Didymocarpus oblonga, Wall. *in litt.* 1819, et *Plant. Asiat. Rar.* vol. 2, p. 34, tab. 140. Don, *Nepal.*, p. 123.

Didymocarpus verticillata, Wall. *List*, No. 783.

Loc. Nepal.

4. *Didymocarpus Punduana*, foliis ternis caulem glabriusculum terminantibus oblongis integris, pedunculis pedicellis et pube glandulosâ conspersis.

Didymocarpus Punduana, Wall. *List*, No. 777.

Loc. Montes Sylhet.

5. *Didymocarpus acuminata*, foliis caulem terminantibus approximatis latè ovatis acuminatis duplicato-crenatis, pedicello capsulæ calycem persistentem superante.

Loc. Chura-Poongi. *Wallich.*

6. *Didymocarpus pedicellata*, acaulis, ovariis glandulosis, capsulæ pedicello calycem persistentem superante, stylo brevissimo.

Didymocarpus macrophylla, *Royle, Illustr.* p. 294, t. 70, f. 1.

Loc. Kamoan. *Royle.*

7. *Didymocarpus macrophylla*, acaulis, ovariis glandulis sessilibus conspersis, capsulæ pedicello brevissimo.

Didymocarpus macrophylla, *Wall. List*, No. 784. *Don, Nepal.* p. 122.

Didymocarpus plicata, *Don, Nepal.* p. 122.

Loc. Nepal. *Wallich.*

β. Calycis lobi acuti.

8. *Didymocarpus subalternans*, caulescens, foliis oppositis suboppositisve obtusis elongato-petiolatis.

Didymocarpus subalternans, *Wall. List*, No. 782.

Didymocarpus aromatica, *Don, Nepal.* p. 123.

Loc. Nepal. *Wallich.*

9. *Didymocarpus obtusa*, scapis subradicalibus bifoliatis, foliis radicalibus elongato-petiolatis cordato-ovatis crenatis obtusis.

Didymocarpus obtusa, *Wall. List*, No. 786.

Didymocarpus aromatica, *Don, Nepal.* p. 122.

Loc. Nepal. *Wallich.*

B. Diandræ. Calyce 5-partito acuto. Capsula utrinque simul dehiscens.

10. *Didymocarpus crinita*, Jack, in Linn. Trans., 14, p. 33, t. 2, f. 2.

Loc. Java, an. 1814. *Horsfield.* Penang. *Jack, Wallich.*

11. *Didymocarpus serrata*, foliis oppositis æqualibus elliptico-lanceolatis serrulatis acutissimis : basi acutâ æquali, cymis bifidis, corollæ tubo limbum quadruplò superante.

Loc. Sumatra? *Horsfield.*

12. *Didymocarpus racemosa*, Jack, l. c., p. 34.

13. *D. corniculata*, Jack, l. c., p. 36.

14. *D. cordata*, Jack, Wall. List, No. 781.

15. *D. corchorifolia*, Wall. List, No. 792.

16. *D. reptans*, Jack, l. c. p. 35.

17. *Didymocarpus missionis*, caule brevissimo, pedunculis axillaribus scapiformibus, foliis cordato-ovatis.

Didymocarpus? *missionis*, Wall. List, No. 639.

18. *Didymocarpus Zeylanica*, sub-acaulis, scapis paucifloris, foliis orbiculato-ovatis altè cordatis dentatis petiolo longioribus, calycis laciniis linearibus acutis.

Loc. Zeylona.

C. Diandra. Calyce 5-partito acuto. Capsula hinc dehiscens inde diù cohærens.

19. *Didymocarpus Rottleriana*, Wall. List, No. 778.

D. Didynamæ.

20. *Didymocarpus frutescens*, Jack, in Linn. Soc. Trans., vol. xiv, p. 39. Wall. List, No. 780.

Loc. Penang.

21. *Didymocarpus elongata*, Jack, in Linn. Soc. Trans., vol. xiv, p. 37.

Loc. Sumatra. *Jack, Horsfield.*

22. *Didymocarpus lanuginosa*, Wall. List, No. 791.

STREPTOCARPUS, *Lindl. Bot. Regist.* 1173.

CHAR. GEN. *Calyx* 5-partitus. *Corolla* bilabiata : tubo calycem duplò multotièsvè superante. *Stamina* antherifera 2, inclusa : antherarum loculis divergentibus. *Stigma* bilobum : labiis æqualibus brevissimis.

Capsula elongata valvis (dextrorsum) spiralibus. *Semina* inappendiculata.

Herbæ pubescentes vel caulescentes, foliis oppositis subæqualibus, inflorescentiâ cymosâ ; vel caule brevissimo, folio altero nano, pedunculis seriatis.

† *Caule abbreviato ; foliis oppositis, altero nano ; pedunculis seriatis axillaribus.*

1. *Streptocarpus Rexii*, Lindl. in *Bot. Regist.*, t. 1173.

Didymocarpus Rexii, *Hooker in Bot. Mag.*, t. 3005.

Loc. Africa australis.

†† *Caulescentes, foliis oppositis subæqualibus petiolatis. Inflorescentia axillaris subcymosa.*

2. *Streptocarpus Helsingbergii*, foliis ovatis crenatis petiolo quadruplo longioribus, cymis paucifloris, corollæ tubo calycem 4—5-iès superante.

Loc. Madagascar. *Helsingberg & Bojer.*

3. *Streptocarpus Bojeri*, foliis ovatis acutis grossè et subduplicato-crenatis petiolo quadruplo longioribus, corollæ tubo calyce duplò longiore.

Loc. Madagascar. *Helsingberg & Bojer.*

4. *Streptocarpus Thompsonii*, foliis subovatis ovalibusque crenato-serratis petiolo paulò longioribus, corollæ tubo calyce duplò longiore.

Loc. Madagascar. *D. I. V. Thompson.*

5. *Streptocarpus paniculata*, foliis ovatis acutis crenato-serratis brevè petiolatis, cymis elongato-pedunculatis paniculatis.

Loc. Madagascar.

Bæa, Commerson, in *Lam. Enc. Méth.* 1, p. 401, et in ⁽¹²⁰⁾
Juss. Gen. Plant., p. 121.

CHAR. GEN. *Calyx* 5-partitus. *Corolla* bilabiata tubo calycem vix æquante. *Stamina* antherifera 2, antherarum loculis divergentibus. *Stigma* obtusum (subbilobum). *Capsula* elongata, valvis (dextrorsum) spiralibus. *Semina* inappendiculata.

Herbæ subacaules, foliis confertis; v. caulescentes, foliis oppositis.

1. *Bæa Commersonii*, caule abbreviato, foliis ovalibus ellipticisve obsolete crenato-serratis petiolatis pube appressâ subsericeis, pedunculis scapiformibus uni-pauciflorisve: pedicellis calycibusque pube appressâ eglandulosâ.

Bæa Magellanica, *Lam. Enc. Méth.* i, p. 401.

Loc. "Isles Praslin," Commerson.

2. *Bæa hygrometrica*, acaulis, foliis ovatis obovatisve crenatis utrinque lanatis basi cuneatis subsessilibus, scapis folio longioribus apice divisis, pedicellis calycibusque pube glandulosâ patulâ.

Dorcoceras hygrometrica, Bunge, in *Mém. Acad. Imper. Sc. Petersb. Div. Sav.* tom. ii. p. 128.

Loc. China Borealis. Bunge.

3. *Bæa? Wallichii*, acaulis, foliis obovatis crenatis crassis, scapis apice 2—4-floris.

Didymocarpus helicteroides, Wall. *List.* No. 789.

Loc. Toong Dong. Wallich.

Obs. Flores nondum visi, ideoque dubii generis. *Capsula Streptocarpi* et *Bæa*.

4. *Bæa? multiflora*, caule suffruticoso, foliis oppositis petiolatis oblongis ovalibusve crenatis, paniculis axillaribus pedunculatis lanatis.

Didymocarpus? multiflora, Wall. *List.* No. 793.

Loc. Sylhet in montibus Punduah.

Obs. Flores inexpansi solum a nobis visi: tubo brevi

corollæ ni fallor cum Bæa convenit sed habitus diversissimus.

LOXOCARPUS.

CHAR. GEN. *Calyx* 5-partitus. *Corolla* tubo brevi. *Stamina* antherifera duo, antherarum loculis divergentibus. *Stigma* indivisum. *Capsula* calyce (paulò) longior, hinc dehiscentis, placentis angustissimis. *Semina* erecta, funiculis capillaribus elongatis.

Herba annua, incana; foliis omnibus radicalibus petiolatis. Scapi apice subcorymbosi.

Loxocarpus incana.

Loxonia? alata. Wall. List, No. 809.

Loc. Penang.

EPITHEMA, Blume, Bijdrag. p. 737.

Aikinia, Br. in Wall. 1 viat. Rar. 3, p. 65, t. 288.

CHAR. GEN. *Calyx* tubulosus 5-fidus. *Corolla* bilabiata. *Stamina* duo superiora antherifera! *Stigma* indivisum. *Capsula* calyce cincta, circumscissa; placentis liberis, utrinque seminiferis, pedicellis parietalibus adnatis. *Semina* erecta, funiculis elongatis.

Herbæ annuæ, pilosiusculæ. Folia cordata, subdentata, inferiora, dum uno plura, alterna, terminalia subopposita. Spicæ unilaterales, circinales; hinc pedicellis subduplici serie confertis ebracteolatis, inde foliolo cucullato subrotundo involucreatæ.

STAUANTHERA, Bentham in Scrophul. Ind., p. 57.

CHAR. GEN. *Calyx* turbinato-campanulatus, 5-fidus, sinibus plicatis (quandoque in dentibus productis). *Corolla* subrotata, bilabiata, 5-loba; tubo nunc basi calcarato. *Stamina* antherifera 4; antheræ in coronulam conniventes, loculis divaricatis. *Stigma* *Capsula* calyce per-

sistente cincta, circumscissa; placentis utrinque seminiferis.

Herba *oppositifolia*, folio altero nano stipuliformi.

1. *Stauranthera grandifolia*, corollæ tubo basi calcarato, calycis sinubus in dentibus productis. (121)

Stauranthera grandifolia, Benth. *Scrophul. Ind.*, p. 57.

Glossanthus? *grandiflora*, Benth. in Wall. List, No. 6395.

Loc. Penang.

2. *Stauranthera ecalcarata*, corollæ tubo ecalcarato.

Miquelia cœrulea, Blume in *Bullct. des Sc. Phys. et Nat. en Neerlande*, an. 1836, p. 94.

Loc. Java. Dr. Horsfield.

GLOSSANTHUS, Klein in Wall. List, No. 6394.

CHAR. GEN. *Calyx* 5-fidus, tubo 5-gono angulis marginatis, laciniis parum inæqualibus; æstivatione valvata. *Corolla* personata; labium superius abbreviatum bilobum; inferius indivisum (lobis lateralibus abortientibus). *Stamina* antherifera 4 inclusa, antheris in coronulam cohærentibus. *Stigma* indivisum. *Ovarium* disco completo cinctum. *Capsula* calyce inclusa, bivalvis, placentis utrinque seminiferis.

Herbæ *alternifoliæ*, *glabriusculæ*; foliis integris basi altè excisis; racemis secundis.

1. *Glossanthus malabarica*, Klein in Wall. List, No. 6394. Benth *Scrophul. Ind.*, p. 57.

2. *Glossanthus Notoniana*.

Wulfenia Notoniana, Wall. *Tent. Flor. Nepal.*, p. 46; List, No. 409.

3. *Glossanthus Zeylanica*.

4. *Glossanthus Mexicana*, R. Br.

Klugia azurea, Schlecht. in Linn., 8, p. 248.

MONOPHYLLÆA.

CHAR. GEN. *Calyx* 5-partitus, laciniis ovatis; æstivatione imbricata. *Corolla* bilabiata: labio superiore bilobo; inferiore trilobo. *Stamina* antherifera 4, antherarum loculis divergentibus. *Ovarium* disco dimidiato basi instructum. *Stigma* indivisum? *Capsula* calycē tecta, ruptilis? semi-bilocularis.

Herba *glabra*. Folium *unicum caulem simplicissimum terminans, sessile, costatum, integrum; racemi corymbosi, subseriati, ex ipsâ basi folii.*

Monophyllæa Horsfieldii.

Loc. Sumatra. *Dr. Horsfield.*

PLATYSTEMMA, *Wall. Pl. Asiat.*, 2, p. 42, t. 151.

CHAR. GEN. *Calyx* altè 5-fidus; æstivatione valvatâ. *Corolla* tubo brevissimo, limbo bilabiato patenti: labio superiore bilobo; inferiore trifido. *Stamina* antherifera 4, exserta, antherarum loculis divergentibus. *Ovarium* disco annulari basi cinctum, placentis utrinque ovuliferis. *Stigma* indivisum. *Capsula* . . . ?

Herba *pubescens; caule simplicissimo terminato folio unico inciso-crenato, quandoque cum altero nano stipuliformi; racemo terminali unico paucifloro.*

RHABDOTHAMNUS, *Cunningh. in Ann. Nat. Hist.*, 1, p. 460.

CHAR. GEN. *Calyx* altè 5-fidus. *Corolla* tubo campanulato, limbo bilabiato. *Stamina* antherifera 4, exserta, antheris in coronulam cohærentibus, loculis divaricatis. *Stigma* . . . *Capsula* demùm 4-valvis, placentis utrinque seminiferis. *Semina* albuminosa.

Frutex ramosissimus oppositifolius.

Rhabdothamnus Solandri, *Cunningh.*, l. c.

Loc. Nova Zelandia. 1769, *J. Banks & Solander*, 1826, *Cunningham.*

122] *FIELDIA*, *Cunningh. in Field's Mem. N. S. Wales*,
p. 363.

CHAR. GEN. *Calyx* 5-partitus. *Corolla* tubulosa. *Stamina* antherifera 4, antherarum loculis parallelis. *Stigma* bilobum. *Bacca* subexsucca.

Suffrutex oppositifolius.

Fieldia australis, *Cunningh. l. c.* p. 364, *cum. tab.*

RHYNCHOTHECUM, *Blume Bijdr.*, p. 775.

Corysanthera, *Wall. List.*, No. 6411.

CHAR. GEN. *Calyx* 5-partitus. *Corolla* bilabiata, tubo brevi. *Stamina* antherifera 4, inclusa; *antheræ*, loculis confluentibus, bivalves, valvulâ interiore minore. *Bacca* calyce cincta.

Suffrutices oppositifolii, fasciculis paniculisve axillaribus.

CENTRONIA, *Blume Bijdr.*, p. 776.

CHAR. GEN. *Calyx* spathaceus, hinc fissus. *Corolla* infundibuliformis, limbo bilabiato patenti. *Stamina* antherifera 4 inclusa; *antheræ* uniloculares, dorso ad basin calcaratæ, liberæ. *Stigma* capitatum. *Bacca* siliquæformis; placentæ carnosæ, lobis revolutis seminiferis.

Herba carnosa in radicibus arborum parasitica aphylla; scapis squamatis.

Obs. Character ex D. Blume qui ad hanc familiam retulit plantam *Æginetiæ* forsan affiniorem.

Centronia mirabilis, *Blume, Bijdr.*, p. 777.

CYRTANDRA, *Forst. Gen.*, t. 3.

Getonia, *Banks et Soland. MSS.*

CHAR. GEN. *Calyx* 5-fidus. *Corolla* infundibuliformis, bilabiata. *Stamina* antherifera 4, 2, inclusa, antherarum

loculis parallelis. *Stigma* bilobum. *Bacca* placentis revolutis undique seminiferis.

WHITIA, *Blume, Bijdr.*, p. 774.

CHAR. GEN. *Calyx* 5-partitus. *Corolla* infundibuliformis, bilabiata. *Stamina* antherifera 2 semiexserta, antherarum loculis haud parallelis. *Stigma* infundibuliforme. *Bacca* siliquæformis.

Frutices (*Javanicæ*) *radicantes*; *foliis oppositis integerrimis inæqualibus inæquilateris*; *inflorescentiâ axillari fasciculatâ*.

Obs. Character a D. Blume.

PTEROCYMBIUM JAVANICUM.

[219]

TAB. XLV.

CHAR. GEN. *Flores* polygamo-monoici (ferè hermaphrodit). *Calyx* 5-fidus, pateris. *Antheræ* (10) simplici serie; loculis omnibus parallelis. *Ovaria* disperma. *Stigmata* filiformia, recurvata. *Pericarpia*: *Folliculi* foliacei, naviculares, longè ante maturitatem aperientes. *Semen* unicum, albuminosum. *Embryonis Radicula* hilo proxima.

Arbor, foliis lobatis indivisisve; paniculis subterminalibus.

Obs. Genus *Sterculiacearum* SCAPHIO proximum et forsàn nimis affine. *Scaphium* verò differt: *Antheris* (15—20) congestis; *stylis* arctè cohærentibus; *stigmatibus* coalitis, unicum lobato-capitatum efformantibus.

DESCR.—Arbor sylvestris, 50—60-pedalis; trunco modicè crasso, cortice lævi, rufescenti-fusco; comâ patenti (D. Horsfield). Folia alterna, 5—3-loba, passim indivisa, circumscriptiōne latè ovata, basi cordata 5-nervia, lobis ellipticis brevè acuminatis, sinubus obtusis; dum indivisa, paulò altius cordata, 3—5-nervia, brevissimè acuminata: omnium adulta supèr glabra præter nervos venasque primarias tenuissimè pubescentes, pube simplici: subtèr pube simili magis copiosâ, præsertim in nervis venisque ferè omnibus et insuper, rariore tamen et tenuiore, per totam paginam: novella utrinque pube stellari copiosâ tecta. Petioli teretiusculi, undique pubescentes, anticè sulco tenui exarati, basi apiceque pululū incrassati, quandoque folia subæquantes, sæpius dimidio et ultrà breviores. Stipulæ laterales, subulatæ, caducæ. Panicula subterminalis, e racemis alternis approximatis subcorymbosis. Pedicelli extra medium articulati, ipso articulo constricto, inter dilatationes duas manifestas. Bracteæ omnes deciduæ v. caducæ. Calyx monophyllus, ad medium quinquefidus, æqualis, turbinato-canpanulatus, coriaceus, glaber, laciniis

tomento minutissimo cinereo marginatis, immersè nervosus
 venisque anastomosantibus, latè semilanceolatis; æstivatione
 valvatâ, alabastro 5-gono, turbinato; tubo intùs basi cor-
 pusculis scobiformibus minutis densissimè tecto. Corolla
 nulla. Flores hermaphroditi vel polygamo-monoici. Her-
 maphrodito-*Masculi*: Columna genitalium (gynostemium)
 e basi parùm dilatatâ cylindracea; dimidio inferiore bre-
 viore, pube tenuissimâ simplici patenti; superiore glabro,
 apice parùm dilatato (staminibus occultato) in urceolum
 decem- v. octo-dentatum antheriferum. Antheræ 10 v. 8
 220] simplici serie dispositæ, incumbentes, supra medium
 affixæ; loculis appositis, contiguis, linearibus, longitudinaliter
 dehiscentibus. Pollen læve, subovale. Pistilla omnibus
 punctis ferè ut in *Femineis*. Hermaphrodito-*Feminei*:
 Columna quandoque brevissima. Antheræ ut in masculis
 numero et dispositione, polline forsân effecto. Ovaria 5
 (rarè 6) ipso apice columnæ insidentia, sessilia, arcuè
 approximata, distincta tamen, dorsi apice gibboso, ita ut
 stylus quasi lateralis evadit, disperma; ovulis collateralibus
 erectis anatropis. Styli 5—6, arcuè approximati, leviter
 cohærentes, breves. Stigmata filiformia, recurvata, intus
 longitudinaliter papulosa. Pericarpia 5—6, aliqua sæpè
 abortientia, follicularia, singula stipiti filiformi pubescenti
 insidentia; stipite communi 5-angulato, vix longitudine
 calycis diù persistentis haud mutati. Folliculi citò post
 fœcundationem et longè ante seminis maturitatem aperientes,
 foliacei, reticulato-venosi, navicularcs; dimidio inferiore
 duplò profundiore ibique carinâ in lobulum obtusum pro-
 ductâ, superiore planiusculo. Semen unicum maturescens,
 exsertum, basin folliculi occupans, erectum, basi insertum,
 sphæricum, læve, tenuissimè pubescens pube simplici. Inte-
 gumentum triplex; *externum* crustaceo-membranaceum,
 latere interiore, respectu axeos communis folliculorum,
 lineâ parùm elevatâ ab hilo ad apicem ductâ raphem indi-
 cante insignitum, superficie interiore præter raphem venis
 per alterum latus recurrentibus ad basin tendentibus, ramo
 crassiore raphi opposito; inter hocce integumentum (v.
 epidermidem testæ) et medium substantia adest parca
 subfloccosa: *medium* (endospermium) nucamentaceum,

apice extus areolâ circulari (chalazâ) insignitum: *intimum* (integumentum interius) membranaceum. Albumen nueleo conforme, bipartibile, amygdalinum, album. Embryo erectus, ferè longitudine seminis, albus. Cotyledones latitudine albuminis idque bipartientes, foliaceæ: Radicula hilo proxima, brevis, subovata.

Pterocymbium javanicum, according to Dr. Horsfield, grows scattered among other trees in the luxuriant forests which, at a small elevation above the sea, cover many of the plains in the central and eastern districts of Java.

The native name is *Wining* or *Kemoonoong*.

TAB. XLV. *Fig. 1.* A branch bearing leaves. *Fig. 2.* A portion of the panicle, bearing ripe fruits. *Fig. 3.* An unexpanded flower. *Fig. 4.* The same, slightly magnified, opened longitudinally, the sexual organs being at that period sessile and the anthers closed. *Fig. 5.* An anther seen in front. *Fig. 6.* The same, seen from behind. *Fig. 7.* The column thickened at top, which in *Fig. 8* is seen divided into ten teeth. *Fig. 9.* One of the pistilla, cut longitudinally to show the two erect collateral ovula. *Fig. 10.* The half of an expanded flower, in which the column is elongated. *Fig. 11.* The base of a carpel, with its single seed. *Fig. 12.* The seed, deprived of its outer integument. *Fig. 13.* The same, after the removal of the inner integument. *Figs. 14 and 15.* The embryo, with its bipartite albumen.

STERCULIACÆ, to which *Pterocymbium* evidently belongs, was first proposed as a distinct natural family by Ventenat in 1804;¹ his only character separating it from *Malvacæ* being the presence of albumen surrounding the embryo, and from *Tiliacæ* its monadelphous stamina.

In 1814,² in suggesting the formation of Natural Classes of plants and proposing *Malvacæ* as one of these classes, I referred *Sterculiæ* to it, but regarded the order as

¹ 'Hort. Malmais,' fol. 91.

² 'Appendix to Capt. Flinders's 'Voyage to Terra Australis,' vol. ii. p. 540 [vol. i. p. 11].

more limited than Ventenat appears to have done; considering it as sufficiently distinct from *Buttneriaceæ*, to great part of which, however, Ventenat's two principal characters of *Sterculiaceæ* equally belong.

221.] In endeavouring to establish the characters and affinities of *Pterocymbium*, it became necessary to examine those of the established genera most nearly related to it, and as this examination has led me to adopt considerable alterations in arrangement, and enabled me to make some additions to the number of species, I shall here give the characters of the genera and species of that tribe to which the name of *Sterculiaceæ* has been applied by DeCandolle and Endlicher, prefacing that account with a history of the tribe, and especially of the genus *Sterculia*, from its formation to the present time.

In the botanical history of *Sterculia*, it is not necessary to go farther back than 1747, when Linnæus first published his character of the genus,¹ founded on the specimens and unpublished figures of Hermannus, whose Ceylon herbarium, of which the Museum Zeylanicum is a catalogue, had been sent from Copenhagen for his inspection. The result of a careful examination of this herbarium was the publication in 1748 of the 'Flora Zeylanica.'

Hermann's herbarium (purchased by Sir Joseph Banks and now in the British Museum) contained flowering specimens of *Sterculia fœtida* and *Balanghas*, and the fruits of both species were figured by Hermann himself in the volume of drawings which accompanied the herbarium. From these materials, confirmed no doubt by the figures of the fruits in 'Hortus Malabaricus' and 'Herbarium Amboinense,' the Linnean genus was entirely established. The character given, however, is strictly applicable to *Sterculia fœtida* only; and Linnæus was probably induced to refer *Balanghas* to the same genus, either from not having distinctly seen the remarkable form of the flower in that species, or, which is more likely, disregarding that difference, was determined by the exact resemblance in its

¹ 'Nova Genera Plantarum, respondente Dasso,' p. 13.

fruit to that of *S. foetida*, both being figured in the volume of drawings already referred to. These were the only materials he appears ever to have examined, and his own herbarium contained absolutely nothing except a single leaf of *S. foetida*: the generic character consequently remains unchanged in all his subsequent works.

In 'Flora Zeylanica,' Linnæus correctly includes *Sterculia* in his Class *Monœcia*, notices the imperfect stamina in the female flower, and only overlooks the minute rudiments of ovaria in the male flower. He referred the genus to his Natural Order *Tricoccæ* (very nearly corresponding with the *Euphorbiacæ* of Jussieu), as appears first in 'Philosophia Botanica,' published in 1751, and afterwards in the sixth edition of his 'Genera Plantarum' in 1764. In this determination of its affinity he was followed by Bernard de Jussieu in 1759, by Adanson in 1763; and in a manuscript list of the arrangement of plants adopted in 1779 in the Paris Garden, I find *Sterculia* still placed in the same family. The generic character of *Sterculia*, as given by Linnæus, who does not notice, and had no opportunity of ascertaining the structure of the seed, might with very slight alteration stand for that of the whole tribe, *Heritieria* alone excepted.

In the order of time, the next work in which the same genus is described, though under a different name, is Aublet's 'History of the Plants of French Guiana,'¹ which appeared in 1775. He describes his genus *Ivira*, which all subsequent botanists have referred to *Sterculia*, as having hermaphrodite flowers, with ten stamina, and the capsules or follicles surrounded at the base with rigid filiform processes, formed as he states by the enlargement of the hairs which according to him exist in the flowering state. But from an examination of the specimens in his own herbarium (purchased by Sir Joseph Banks and now in the British Museum), as well as from others collected in the same country by the late celebrated Professor Richard, it appears that the flowers are unisexual; that the hairs of the

¹ p. 695, t. 279.

urceolus antherarum which Aublet describes, and which, as he states, by their enlargement form the mass of rigid processes surrounding the ripe fruit, have no existence; and that this singular appendage to the fruit is an accidental structure or monstrosity, probably of rare occurrence, of which I have found a single specimen among the loose fruits of his herbarium.

In 1775 also was published the 'Flora Ægyptiaco-Arabica' of Forskael, who describes a supposed new genus, *Culhamia*,¹ which is no doubt a species of *Sterculia*, certainly, however, not *platanifolia* as Vahl affirms, but ²²²¹ perhaps either *S. tomentosa* of the 'Flora Senegambiæ,' or the nearly related species *abyssinica*. *Culhamia* is described as having hermaphrodite flowers, and it is stated that the style, which is originally concealed by the stamina, only becomes visible after the expansion of the flower, so that here we have the same account of structure and gradual development of pistillum which was afterwards given by Cavanilles and adopted by de Jussieu.

In 1786, in the first volume of the 'Acta Acad. Patavin,'² Marsili proposed a new genus related to *Sterculia*, and formed of *St. platanifolia* of the 'Supplementum Plantarum' of the younger Linnæus.

In 1788 Cavanilles published his fifth dissertation on Monadelphous plants, in which *Sterculia*³ appears with its characters considerably modified, *Ivira* of Aublet being included in it, and some additional species described. He refers the genus to *Monadelphia*, states the flowers to be hermaphrodite, and accounts for their unisexual appearance by assuming the gradual and more tardy development of the female organ; he also describes the stigma as either bifid or with five rays, the supposed existence of the bifid stigma being founded on an error in the drawing of *S. platanifolia* by Mlle. Basseporte, from which his engraving is copied.

In 1789 the immortal work of A. L. de Jussieu appeared. *Sterculia* is there⁴ for the first time correctly

¹ p. 96.

² p. 106.

³ p. 284.

⁴ p. 278.

referred to the Natural Order *Malvaceæ* as then understood: the generic characters, however, are stated to be taken from Aublet and Cavanilles. The erroneous assertion of the latter that the flowers are hermaphrodite, and his account of the gradual development of pistillum are adopted, and finally Jussieu expresses a doubt whether the embryo is corrugated, which he describes it to be in all the other genera of the same section of *Malvaceæ*, and which he considers as one of the principal characters of that Natural Order.

In Schreber's edition of the 'Genera Plantarum'¹ of Linnæus, published the same year, *Sterculia* is referred to *Dodecandria*, and the few changes made in the character are apparently taken from Cavanilles, and among these is the *stigma bifidum*.

In 1789 also Dryander, in the first edition of 'Hortus Kewensis,' established, partly from the manuscripts of J. G. Kœnig, the genus *Heritiera*,² which he justly places next to *Sterculia*, and refers both to *Monœcia Monadelphica*.

Neither in Willdenow's edition of the 'Species Plantarum' of Linnæus, nor in Persoon's 'Synopsis' is any important alteration made in the character of the genus, which, following Schreber, is referred by Willdenow to *Dodecandria*, and by Persoon to *Monadelphia*.

In 1804 Ventenat, in his 'Jardin de la Malmaison,'³ gives a considerably altered, and in most respects improved character of *Sterculia*, of which with some other genera of *Malvaceæ*, and the first section of Jussieu's *Tiliaceæ*, he proposes to form a new Natural Order, *Sterculiaceæ*, his principal distinguishing character of which has been already mentioned. He refers the genus to *Monadelphia Dodecandria* of the Linnean system for no very sufficient reason, namely the existence of the rudiments of the other sex in both the male and female flowers. He is the first to notice the *albumen bipartibile* and *radicula embryonis umbilico opposita* or *contraria*, both of which he introduces into

¹ p. 324.² Vol. iii. p. 546.³ Vol. ii. fol. 91.

the character of the genus. Besides these improvements in the account of the genus, he has given the distinguishing characters of several new species.

In 1805 M. Palisot de Beauvois, in his 'Flore d'Oware et de Benin,'¹ in describing his *Sterculia acuminata*, introduces several alterations into his character of the genus, most of which belong only to the species he has there described, and some of which may be considered of generic importance, especially the remarkable structure of antheræ.

In 1806 Salisbury proposes a new genus, *Southwellia*, formed of all such species of *Sterculia* as have the segments of the calyx connivent with cohering apices; this being the only character distinguishing it from the rest of the genus.

In the same year Poiret, in the article *Sterculia* of the Botanical Dictionary of the 'Encyclopédie Méthodique,'² considers the flowers as hermaphrodite, and adopts the explanation given by Forskael, Cavanilles and de Jussieu. In his generic character there is no improvement; but he suggests the probable expediency of afterwards dividing the genus, when the structure of the various plants referred to it is better known.

In 1819 Sir James (then Dr.) Smith, in Rees's 'Cyclopedia,' in his account of *Sterculia*, takes no notice of the structure of seed, and is inclined to agree with Schreber and Willdenow in referring it to *Dodecandria*. If the genus should be hereafter subdivided, he seems more disposed to trust to differences in the styles and stigmata than to those very remarkable modifications of fruit, with some of which he was acquainted, believing them not to be supported by other characters, and in proof of this instancing *St. platanifolia* and *colorata*, so similar in fruit, and so unlike each other in the form of the flower.

In 1824 DeCandolle, in the first volume of his 'Prodromus,' forms a tribe which he calls *Sterculiæ*, consisting of *Sterculia* and *Heritiera*, referring it to *Buttneriaceæ*. The

¹ Vol. i. p. 40.

² Vol. vii. p. 423.

most remarkable circumstance in his character of *Sterculiæ*, is his describing the embryo as erect, which, if I rightly interpret his meaning, although applicable to a few species, is directly contrary to the more usual structure. *Sterculia* as a genus he distinguishes from *Heritiera* by the existence of albumen, which, however, is not universally present, and dehiscence of the follicles; the direction of embryo is not noticed in his characters of either of these two genera.

In 1827 M. Auguste de St. Hilaire, in the 'Plantes Usuelles des Brasiiliens,' gives, appended to his account of a supposed new species of *Sterculia*, an improved character of the genus, though in this character he describes the albumen as a coat of the seed; the "embryon antitrope" is considered as common to the whole genus. In the remarks that follow he supports Linnæus's account of his classification of the genus, in opposition to the observations of Jussieu which he refers to, but which he does not seem to be aware were adopted from Cavanilles, and perhaps also from Forskael, with whom a similar account originated.

In the same year he published a *Livraison* of his 'Flora Brasiiliæ Meridionalis,' in which work, as in the 'Plantes Usuelles,' owing to the state of his health, he was assisted by MM. Adrien de Jussieu and Cambessèdes. In the 'Flora Brasiiliæ'¹ an enlarged, and in many respects improved character is given of *Sterculia*, of which the principal difference from that in the 'Plantes Usuelles' is, his admitting the existence of albumen divided into two equal segments, which, he correctly states, frequently cohere with the corresponding cotyledons, the first distinct notice I believe of that remarkable economy: he states also, that where the seeds are ascendent, the radicle of the embryo points to the hilum, and where they are transverse, in the diametrically opposite direction. His character, therefore, of the genus is so framed as to include every species of *Sterculia* of DeCandolle, as far as the author was acquainted with their structure. It will hereafter appear, however, that neither is the existence of albumen universal, nor is

the direction of the embryo dependent on that of the seed, even in the carpella of all the species then published.

In 1831, in the 'Flore de Sénégambie,' the joint work of MM. Guillemain, Perrottet and Richard, a new point is introduced into the character of *Sterculia*,¹ namely an incomplete arillus, which, however, if it really exists in any case, is probably to be found in one species only, namely, *Sterculia cordifolia*: in all the other species which I have examined, there is either only a minute *caruncula umbilicalis* or *strophiola*, as in *St. fœtida*, or more generally no trace whatever of this appendage.

In 1832, in the 'Meletemata Botanica' of Schott and Endlicher, the Natural Order *Sterculiaceæ* is divided into three principal tribes—*Bombaceæ*, *Helicterææ*, and *Sterculiææ*. This last tribe, as in DeCandolle's 'Prodromus,' is limited to the genera *Sterculia* and *Heritiera*. But *Sterculia* is divided into twelve genera, chiefly from modifications of the flower, or from the texture and period of dehiscence of the folliculi, and in one case from the seed being winged; no modification of internal structure of seed being introduced into any of the characters.

In the same year, the third volume of Dr. Roxburgh's 'Flora Indica' was printed at Calcutta. In this valuable work such a generic character of *Sterculia*² is given as to comprehend all the Indian species, and indeed so constructed as to include all that are now known, except *Courtenia*, a new genus of the present essay; and even that would be excluded only from its generally having double the usual number of ovaria. Several new species are well described in the work, and the direction of embryo noticed in most of them; the only species in which the radicle is described as pointing to the umbilicus being his *Sterculia alata*.

In 1840 Professor Endlicher, in the 13th part of his 'Genera Plantarum,' modifies the arrangement of the Linnean genus *Sterculia* given in the 'Meletemata,' all the genera there established, except his *Pterygota* (the *St. alata*

¹ Vol. i. p. 79.

² p. 144.

of Roxburgh), being considered as only subgenera of *Sterculia*, of which the principal characters distinguishing it from *Pterygota* are, as he states, orthotropous ovula and embryo either antitropous or parallel to the umbilicus. It would seem that these distinctions were adopted chiefly from the facts stated in the 'Flora Indica' of Dr. Roxburgh, and in part also perhaps from the generic character of *Sterculia* given in the 'Flora Brasiliæ.'

In 1841 the same arrangement and characters are given in his 'Enchiridion.'

Before proceeding to the description and characters of *Sterculiæ*, I have to make a few preliminary observations, chiefly on the relative importance of the different organs in the formation of genera.

In the general description of the tribe, I have enclosed in brackets such exceptions to the more usual structure as I believe to be in most cases of generic value.

The ordinary structure follows separately; no notice being taken of the exceptions, which are numerous, and such parts of the general description being excluded as are of least importance, and it may be remarked that this character is but little different from that of the genus *Sterculia*.

I then give the few characters to which hitherto no exception has been found.

The direction of *Embryo* with relation to the insertion or umbilicus of the seed appears to be by far the most important character, or that which is best supported by other modifications of structure; and it is worthy of remark, that in this point the ordinary direction of the embryo in the tribe, namely the radicle seated at the opposite extremity or apex of the seed, is itself a deviation from the more usual structure of Phænogamous plants, and an exception not only to the other tribes of *Sterculiaceæ*, but to the whole of the Natural Class *Malvaceæ*, to which that order belongs; and it becomes still more remarkable in

regard to the state of the unimpregnated ovulum, which I have some reason to believe is not orthotropous as might be expected, and as it has been described, but apparently anatropous, and that perhaps in the whole tribe. As, however, my observations on this subject are entirely made from the macerated ovaria of dried specimens, the statement here made must be received as requiring confirmation from the examination of living plants, and of a greater number of species.¹

From this ordinary direction of embryo in the tribe, the deviations are of two kinds: the first, and no doubt the more important, is that in which the radicle is placed at a point close to the umbilicus, which is the most general structure in Phanogamous plants; but as it never points directly within the umbilicus, either in this or any other family, I have modified the expression generally employed in such cases. The second deviation is where the umbilicus is placed on or near the middle of the ripe seed with the radicle pointing to its lower extremity; in other words, where the embryo is parallel to the umbilicus. But this position of umbilicus of the ripe seed does not necessarily imply an exactly similar insertion in the unimpregnated ovulum; and in this tribe I am inclined to believe that in many cases the foramen of the ovulum is so close to the ^{225]} umbilicus as to appear anatropous, and that it ultimately becomes more distant from the unequal growth of the opposite extremities of the seed.

The exceptions to the ordinary structure in *Sterculiæ* which appear to be next in importance are the modifications in texture, and especially in the period of dehiscence of the seed-vessel or carpel, or even its nondehiscence, for in this respect the tribe admits of the two extremes. In the first, where the carpel opens long before the ripening of the seed, its texture is always foliaceous, and the embryo may

¹ The species of *Sterculia* with orthotropous embryo in which I have found this unexpected position of foramen in the unimpregnated ovulum, are *foetida*, *guttata*, *carthagenensis*, *nobilis*, and *angustifolia*; and in the ripe seeds of *tragantha*, *urens*, *villosa*, and *quadrifida*, an indication of a lateral foramen near the base is still visible, but which in *foetida* I have not been able to detect.

be either transverse, that is, having its radicle distant from the umbilicus, or approximated to it.

The want of *Albumen* in this tribe is also a character of considerable value, but its absence is not quite so frequent as has been supposed, there being some room for doubt where its union with the cotyledons is most intimate.

The deviations from ordinary arrangement of the *Antheræ* may be considered as next in value to those already enumerated; of these modifications, that of the greatest importance is where the antheræ are disposed in a simple series and equidistant; this occurs I believe only when they are ten in number, or in the very rare case where they are reduced to five; the mere number, where they are not equidistant nor closely approximated, seems to be of much less importance.

The degrees of development of stamina in the female flower can never be employed in the formation of genera: in several, perhaps in many cases, the antheræ contain pollen, but, on comparing it with that of the male flower, I am disposed to believe it to be imperfect or effete. It would seem at first sight, that, in the numerous cases where the style is deflected, the stigmata are so closely approximated to the antheræ, that impregnation by their pollen is at least probable; but by this deflection of style the stigmata are equally exposed to the influence of the antheræ of the male flowers, which are generally lateral, while the female is terminal, the exposure being nearly equal, whether the inflorescence is pendulous or erect.

The modifications of calyx in *Sterculiæ*, especially in the depth of division and direction of its segments, have hardly more than specific or sectional value; and even those processes which occur in several species, either in the segments or tube, seem to be of no greater importance. The most remarkable anomaly of all is the induplicate æstivation found in a single species belonging to New Holland and in *Sterculia villosa* of Roxburgh. This mode of æstivation readily passes into the valvular in the corolla of several families, and is therefore of no great weight.

It is singular that so great a difference as that between

simple and compound leaves should in this tribe be unconnected with other characters, and the very few plants in which compound leaves occur differ widely from each other in the form of their flowers or direction of their segments.

STERCULIÆ.

DESC. TRIBUS.—*Flores* diclines, monoici, *Masculi* cum rudimento, sæpiùs minuto et antheris occultato, pistilli: rarò stigmatibus ovulisque ferè ut in femineis. *Feminei* antheris manifestis sæpè donatis polline sed semper effecto. *Calyx* monophyllus, 5-fidus, nunc 5-partitus, rarò 4- v. 6-fidus, æqualis, coloratus; æstivatione valvatâ rarissimè induplicatâ; deciduus, quandoque marcescens. *Corolla* nulla. *Gynostemium*, columna genitalia sustinens, calyce brevius v. longius rarò abbreviatum ferè nullum, in cuius apice *Antheræ* subsessiles quindecim, nunc decem, quandoque viginti (rarò quinque) urceolum abbreviatum cyathiformem rarissimè tubulosum terminantes, in capitulum inordinatim congestæ, v. in fasciculos quinque cum ovariis alternantes polyadelphæ (nunc dum decem v. quinque, sæpiùs in serie simplici dispositæ); biloculares, oculis distinctis, parallelis (rarò divaricatis). *Pollen* simplex, læve, sphæroideum. *Ovaria* 5, rarò 4 v. 6 (rarissimè 10—12), leviter cohærentia, polysperma (nunc tetra- v. disperma, rarò monosperma). *Ovula* angulo interno ovarii, dum inde-
226] finita duplici rarò quadruplici serie, inserta. *Styli* tot quot ovaria, coaliti v. arcuè approximati, et tunc sæpè arcuè deflexi (nunc ad basin ferè distincti, quandoque nulli). *Stigmata* in unicum subcapitatum quinquelobum cohærentia, v. distincta subcylindræa indivisa. *Pericarpia*: *Folliculi* coriacei v. lignei (rarò submembranacei) hæud ante maturitatem seminum dehiscetes (in nonnullis in quibus ovula definita (2 v. 4) citò post fecundationem aperientes, foliacei) (rarò clausi, nucamentacei). *Semina* aptera rarissime apice alata, umbilico sæpè omninò nudo, quandoque strophiolâ parvâ, rarissimè si unquam? arillo carnosio incompleto? aucto. *Integumentum* triplex (in exalbuminosis simplex:

dum triplex *extimum*, quasi testæ lamina exterior, tenue, crustaceum, superficie interiore sæpiùs vasculosâ, inter hoc et medium substantia laxè cellulosa vel stiposa v. pulvorea sæpè obvia; *medium*, quasi testæ lamina interna, cartilagineo-nucamentaceum, evasculosum semper; *intimum* membrana-ceum, quandoque evanidum. *Albumen* amygdalino-carnosum, album, bipartibile! segmenta sæpiùs cotyledonibus cohærentia, in quibusdam cum iisdem conferruminata, et texturâ v. colore paullò diverso tantùm indicata quandoque nullum. *Embryo* dicotyledoneus, albus, longitudine albuminis, antitropus (nunc orthotropus v. transversus). Cotyledones in albuminosis latitudine albuminis et sæpiùs foliaceæ (in exalbuminosis semper carnosæ). *Radicula* v. hilo contraria et centrifuga (v. approximata et centripeta, v. transversa et infera). *Plumula* parva sed manifesta.—Arbores *intra tropica* sæpiùs provenientes, et tamen omnes *foliis deciduis*. Folia *alterna, petiolata, simplicia, indivisa* v. *lobata, rarò digitato-composita, foliolis cum petiolo articulatis: omnia pube sæpissimè stellatâ, in quibusdam nullâ, presertim in paginâ superiore, v. in adultis deciduâ*. Petiolus *teretiusculus, basi et apice paullò incrassatis*. Stipulæ *laterales, distinctæ, plerumque subulatæ et caducæ*. Inflorescentia sæpiùs *paniculata* v. *racemosa, pendula, quandoque fasciculis axillaribus* v. *rameis erectis: bracteolis caducis pedicellis medio v. juxta apicem articulatis*.

CHAR. ORDINARIUS TRIBUS. *Flores* polygamo-monoici. *Calyx* 5-fidus, æstivatione valvatâ. *Corolla* nulla. *Genitalia* stipitem terminantia. *Antheræ* 15, congestæ v. polyadelphæ, biloculares; loculis parallelis. *Ovaria* 5, pluriovulata. *Styli* cohærentes. *Stigma* divisum v. lobatum. *Folliculi* ligneo-coriacei, polyspermi. *Embryo* rectus, longitudine albuminis bipartibilis. *Cotyledones* albumini cohærentes. *Radicula* hilo contraria.

CHARACTERES UNIVERSALES. *Flores* unisexuales. *Calyx* coloratus, æqualis, æstivatione valvata rarò ejusdem modificatione induplicata. *Corolla* 0. *Stamina* hypogyna. *Antheræ* biloculares. *Pericarpia* distincta.

STERCULIA.

Sterculiæ pars auctorum ferè omnium.

CHAR. GEN. *Calyx* 5-fidus (rarò 4-fidus). *Antheræ* v. inordinatim congestæ v. polyadelphæ. *Ovaria* pluriovulata. *Styli* cohærentes (in unicum sæpiùs arcè deflexum). *Stigmata* in unicum 5-lobum coalita quandoque distincta. *Folliculi* lignei v. coriacei. *Semina* albuminosa; albumine bipartibili cum cotyledonibus plùs minùs arcè cohærenti. *Embryonis radícula* hilo contraria.—Arbores intra tropicos Asiæ (et insularum adjacentium) Africæ et Americæ provenientes. Folia indivisa, v. lobata, v. in paucis digitatim ²²⁷ composita. Inflorescentia v. paniculata v. racemosa, in plerisque nutans; floribus terminalibus sæpiùs præcocioribus et femineis, et in his stylus arcè deflexus.

* Folia composita, digitata: foliolis cum petiolo articulatis et separatim solubilibus.

a. Calycis lacinia patulæ.

STERCULIA (*fœtida*) foliolis 7—5 elliptico-lanceolatis acuminatis petiolatis vix quadruplo longioribus quam latis utrinque glabris.

S. fœtida, Linn., DeCand.

Loc. Nat. India orientalis, Insulæque Moluccanæ.

Obs. Varietatem in orâ septentrionali Novæ Hollandiæ, anno 1802, legi cujus venæ primariæ subflexuosæ et magis deliquescentes.

S. (polyphylla) foliolis 9—11 lincari-lanceolatis elongatis sessilibus sexies longioribus quam latis utrinque glabris.

Loc. Nat. E Sumatra, ad fretum Sunda, folium unicum reportavit D. G. Staunton (v. s.).

Obs. Figura Clompani majoris, Rumph. Amb. iii, t. 107, sat benè respondet, arboremque juvenilem biorgyalem, foliis

omnibus sumatrano similibus, in Hort. Reg. Paris. cultam anno 1843 vidi.

S. (mexicana) foliolis 7 cuneato-oblongis basi attenuatis petiolatis glaberrimis, floribus (masculis) paniculatis, calycibus intus barbatis.

Loc. Nat. Mexico ad Chiapas D. Linden v. s. in Herbb. Hort. Paris. et D. Delessert.

Obs. Feminei flores et pericarpia desiderantur ideoque genus incertum.

* *β. Calycis laciniæ conniventes, apicibus diù coherentibus.*

S. (versicolor) foliolis oblongo-ellipticis acuminatis subtùs tomentosis.

S. versicolor, *Wall., Pl. asiat. rar.* i, p. 48, tab. 59.

Loc. Nat. Ind. or. ad ripas Irrawaddi D. Wallich.

Obs. Fructus ignotus.

** *Folia simplicia, aliqua v. omnia lobata. Calycis laciniæ patentes.*

S. (urens) foliis 3—5-lobis subtùs tomentosis velutinis, paniculis compositis erectis, calycis laciniis basi lingułā deflexâ, floribus decandris, urceolo antherarum cylindræo, stylo erecto.

S. urens, *Roxb. Corom. pl.* i, p. 25, t. 24.

Loc. Nat. India orientalis, Roxburgh.

S. (villosa) foliis 5—7-lobis acuminatis subtùs tomentosis velutinis, paniculis compositis pendulis, calycis æstivatione induplicata! stylo deflexo.

S. villosa, *Roxb. Fl. ind.* iii, p. 153.

Loc. Nat. Ind. or., Roxburgh.

S. (alyssinica) foliis 3—5-lobis adultis supèr glabris subtèr pubescentibus, racemis (femineis) sub-simplicibus nutantibus, calycibus glabriusculis: laciniis intùs extra

medium villosis, columna genitalium glaberrimo calyce brevior, stylo recurvo, folliculis polyspermis intus setosis.

S. abyssinica, *R. B. in Salt, Trav. app.*, p. lxiv.

Loc. Nat. Abyssinia, D. Salt.

^{228]} *S. (tomentosa)* foliis cordatis plerisque trilobis adultis utrinque tomentosis velutinis, floribus racemosis pendulis, calycibus extus tomentosis, stipiti genitalium piloso.

S. tomentosa, *Guillem. Perrottet et Richard, Flor. senegamb.*, i, p. 81, tab. 16.

Loc. Nat. Africa æquinotialis.

Obs. *S. abyssinica* nimis affinis.

S. (Triphaca) foliis reniformibus acutis indivisis obsoletè trilobisve adultis supèr glaberrimis subtèr glabriusculis, ovariis 2—3, stylo recurvo, stigmatè lobato-capitato.

Triphaca africana, *Lour. Cochîn.*, 577 (et *DeCand. Prodr.*) fid. fragm. ab ipso Loureiro in Herb. Mus. Paris.

Loc. Nat. Africa orientali-australis contra Mozambique, Loureiro.

S. (carthaginensis) foliis 3—5-lobis obtusis acutiusculisve cordatis adultis supèr glabratibus subnitentibus subtèr velutino-tomentosis cinereis venulis immersis, calycibus extus furfuraceo-tomentosis intus glabris, stylo recurvo.

S. carthaginensis, *Cav. Diss.* vi, p. 353.

S. Helicteres, *Pers. Syn.* ii, p. 240. *DeCand. Prodr.* i, p. 483.

S. Chicka, *Aug. St. Hil. Pl. usuell. des Brasil.*, tab. 46, et *Fl. Brasil. merid.*, i, p. 278, fid. exempl. ab ipso Auct. in Herb. Mus. Paris.

Helicteres apetala, *Jacq. Amer.*, p. 238, ubi desc. opt. cum ic. flor. in tab. 181, fig. 97, fide floris unici feminei in Herbario ipsius Jacquin in Museo Britannico asservato.

Obs. Ab hac vix differt *Chichæa acerifolia*, Presl, Reliq. Hænck. ii, p. 141, quamvis ad plantam suam stylum erectum attribuit; et hujus loci forsitan *St. punctata*, DeCand. Prodr. i, p. 483, vix obstante foliorum lobis acuminatis figuræ ineditæ.

S. (striata) foliis 3—5-lobis acutiusculis obtusisve cordatis adultis supèr glabris non nitentibus subtèr pubescentibus rugosis venulis emersis, calycibus extùs pilosis.

S. striata, Aug. St. Hilaire et Naudin, in *Annal. des sc. nat.*, ser. 2, tom. xviii, bot. p. 213.

Loc. Nat. Brasilia, v. Martius (in cujus herb. *S. chicka*) et D. Gardner.

*** *Folia simplicia, omnia indivisa.*

a. *Calyx* 5-partitus, patens; laciniis medio intùs lingulâ (squamulâ abbreviatâ) auctis. *Antheræ* decem, sed subcongestæ (nec æquidistantes).

S. (Ivira) foliis ovato-oblongis acutis basi obtusâ (nec cordatâ) adultis supèr glabris subtèr petiolisque tomentosis.

Ivira pruriens, Aubl. *Guian.*, p. 695, t. 279. Character ab ipso exemplari Aubletii in Herb. Banks. Mus. Brit.

S. Ivira, Sw. *Prodr.* p. 98, *Flor. Ind.-occident.* ii, p. 1160.

Obs. Character specific. et descr. Swartzii præsertim e descriptione Aubletii deprompta est, sed partim a sequenti.

S. (caribæa) foliis ovato-oblongis acutis basi obtusâ adultis supèr petiolisque glaberrimis subtèr glabriusculis.

Loc. Nat. In Insulis Caribæis legit De Ponthieu.

S. (propinqua) foliis oblongo-ovatis acutis basi cordatâ subtèr tomentosis.

Loc. Nat. A Surinam misit D. Hostmann.

Obs. A *S. Ivira* foliis basi cordatâ præsertim et vix aliter differt.

S. (frondosa) foliis ad apicem ramuli confertis oblongis obtusissimis basi obtusiusculâ lævibus adultis supèr glaberrimis nitidis subtèr glabris venulis altè immersis, racemis axillaribus pedunculatis divisis.

Sterculia frondosa, *Richard in Act. soc. hist. nat. paris.*, p. 111.

Loc. Nat. Guiana Gallica, *D. Richard.*

S. (Surinamensis) foliis ad apicem rami confertis oblongis obtusissimis basi obtusiusculâ adultis utrinque pubescentibus venulis subtèr parùm emersis, racemis axillaribus pedunculatis divisis.

Loc. Nat. A Surinam misit *D. Hostmann*, v. s. in *Herb. D. Hooker.*

S. (rugosa) foliis oblongis obtusissimis venis venulisque subtèr prominentibus rugosis.

Loc. Nat. In Demerara detexit *A. Anderson*, Horti Botanici Ins. Sti. Vincentii tunc præfectus.

*** *β. Calyx patens, laciniis absque lingulâ.*

S. (guttata) foliis oblongo-ovatis acutis acuminatisve basi obtusâ subcordatâ adultis subtèr tomentosis, racemis axillaribus indivisis, floribus subternis subsessilibus, calycibus extùs furfuraceo-tomentosis intùs villosis.

Sterculia guttata, *Roxb. Fl. ind.* iii, p. 148.

α. folia oblongo-ovata acuta.

β. folia obovata acuminata, racemis folio multotiès brevioribus.

Loc. Nat. Ind. Or., *D. Roxburgh.*

S. (Madagascariensis) foliis oblongis altè cordatis obtusis adultis utrinque glaberrimis, racemis compositis, pedunculis calycibusque extùs pube stellari cinereis.

Loc. Nat. Madagascar, *D. Richard de Bourbon in Herb. Mus. Paris.*

S. (Roxburghii) foliis ovalibus oblongisve acuminatis basi obtusâ utrinque glabris, racemis axillaribus folio brevioribus, calycibus profundè 5-fidis: laciniis lanceatis imberbibus.

Sterculia Roxburghii, *Wall. List*, n. 1124.

Sterculia lanceæfolia, *Roxb. Fl. ind.* iii, p. 150.

Loc. Nat. Ind. Or., Sillet, *Dr. Wallich.*

S. (lanceolata) foliis elliptico-lanceolatis acutis basi acutiusculâ adultis utrinque glabris, paniculis axillaribus folio brevioribus, calycibus quinquepartitis patulis imberbis glabriusculis.

Sterculia lanceolata, *Cav. Diss.* v, p. 287, t. 143, f. 1.
Lindl. in Bot. regist., 1256.

Loc. Nat. China.

S. (parvifolia) foliis lanceolatis utrinque glaberrimis acutis subacuminatisve basi acutâ, racemis axillaribus simplicibus folio brevioribus.

Sterculia parvifolia, *Wall. List*, n. 1123.

[230]

Loc. Nat. Ins. Penang, *Dr. Wallich*.

S. (Javanica) foliis obovatis acumine brevissimo basi obtusâ subcordatâ subtèr tomentosis, paniculis cernuis, calycibus patulis utrinque tomentosis.

Sterculia cordata, *Blume Bijdr.*, p. 86?

Loc. Nat. Java, *Dr. Horsfield*.

S. (macrophylla) foliis altè cordatis obtusis indivisis subtèr tomentosis, paniculis lateralibus cernuis, calycibus 5-fidis patentibus.

Sterculia macrophylla, *Vent. Malm.* ii, n. 91 *in adnot.*
De Cand. Prodr. i, p. 483.

Loc. Nat. Java, *Dr. Horsfield*.

S. (comosa) foliis ovatis cordatis acuminatis subtèr tomentosis, paniculis decompositis, calycibus patentibus.

Sterculia comosa, *Wall. Pl. asiat. rar.* ii, p. 25, t. 127.

Loc. Nat. Amboina, *in Hort. Calc. cult.*, *Wallich*.

S. (longifolia) foliis oblongis acutiusculis glabris, racemis subsimplicibus, pedicellis subgeminis folio brevioribus, calycis laciniis patentibus intus barbatis tubum extus glabriusculum subæquantibus.

Sterculia longifolia, *Vent. Malm.* ii, n. 91 *in adnot.* fid. specim. in *Herb. Vent.*

Obs. Cfr. *Sterculiam lanceolatam*, *Blume, Bijdr.* quoad

folia et inflorescentiam similem sed calycis laciniis conniventibus et apice connexis diversam.

Loc. Nat. Java, Dr. Horsfield.

S. (lævis) foliis oblongo-lanceolatis acumine lineari obtuso basi acutiusculis utrinque glabris, racemis subsimplicibus folio brevioribus, calyce extus glabro: laciniis patentibus tubo duplò longioribus e latâ basi linearibus intus barbatis.

Sterculia lævis, Wall. List, n. 1138.

Sterculia coccinea, Jack in Mal. miscel. i, non Roxburgh.

Loc. Nat. Pulo Pinang.

S. (Spangleri) foliis lanceolato-oblongis acutis subacuminatisve utrinque glabris, racemis subcompositis, calycibus extus glabris: laciniis patentibus e latiore basi subulatis intus barbatis tubo duplò longioribus.

Obs. Proxima *Sterculiæ lævi* et *longifoliæ*; ab illâ foliorum latiorum et basi obtusiorum acumine brevior et ab apice folii sensim angustato nec subitò constricto et lineari; ab hac præsertim calycis laciniis tubo duplò longioribus diversa.

Loc. Nat. Java, D. Spangler in Herb. D. Hooker.

S. (coccinea) foliis lanceolatis subacuminatis basi obtusiusculâ adultis utrinque glabris, paniculis axillaribus cernuis, calyce extus glabro: laciniis e latâ basi filiformibus barbatis tubo abbreviato 4—5-plò longioribus.

^{231]} *β. folia cuneato-lanceolata, racemi subsimplices folio breviores.*

Sterculia coccinea, Roxb. Fl. ind. iii, p. 151. Wall. List, n. 1122.

Loc. Nat. a. Sillet, Roxb., Wall. β. Assam, D. Griffith.

Obs. Ad hanc sectionem forsàn referendæ *Helicteres undulata* et *paniculata*, *Lour. Cochîn.*, p. 531: et ad eandem primo intuitu pertinere videtur *Sterculia punduana*, *Wall. List, n. 2701*, quæ autem *Reevesiæ* species est, *R. Wallichii, nob.*

*** γ. *Calycis lacinia conniventes apicibus diu
coherentibus.*

S. (nobilis) foliis ovali-oblongis brevè acuminatis basi obtusâ utrinque glabris, paniculis pendulis calycibusque extûs pube rarâ conspersis, stigmatibus subrotundis stylo deflexo multoties brevioribus.

Sterculia monosperma, *Vent. Malm.* ii, n. 91.

Sterculia Balanghas, *Roxb. Fl. ind.* iii, p. 144, non *Linnæi*.

Southwellia nobilis, *Salisb. Parad. lond.*, t. 69, exclus. synn. *Linnæi*, *Cavanilles* et *Hort. Malab.*

Loc. Nat. China; in *Hort. Bot. Calc. et Europæis* culta.

S. (Balanghas) foliis oblongo-ovalibus obtusiusculis v. acumine brevissimo obtuso basi obtusâ (nec cordatâ) adultis supèr glabris subtèr pubescentibus, paniculis calycibusque extûs tomentosis: laciniis intûs densè barbatis, folliculis polyspermis extûs tomentosis intûs glabris.

Sterculia, *Linn. Flor. zeyl.*, n. 350, *fide specimenum Herb. Hermannii*, vol. ii, fol. 42.

Sterculia Balanghas, *Linn. Sp. pl.* ed. 1, p. 1007.

Cavalam, *Hort. malab.* i, p. 89, t. 49.

Loc. Nat. Zeylona et Peninsula Indiæ Orientalis.

Obs. Hujus varietas ut videtur foliis acutis acutiusculisve nec unquam constrictè acuminatis, petiolis adultis pubescentibus, ex *Herb. D. Wight*, et probabiliter illius *S. Balanghas*, *Illustr.*, tab. 30.

S. (angustifolia) foliis oblongo-lanceolatis attenuato-acuminatis basi obtusâ adultis supèr glabriusculis subtèr copiosè pubescentibus, paniculis nutantibus folia superantibus, calycibus extûs tomentosis: laciniis intûs modicè barbatis tubo longioribus, stigmatibus distinctis recurvis stylum æquantibus.

Sterculia angustifolia, *Roxb. Fl. ind.* iii, p. 148. *Wall. List*, n. 1133.

Loc. Nat. Nepalia.

S. (mollis) foliis elliptico-oblongis acutis acuminatisve basi obtusâ adultis supèr pube rarâ scabriusculis subtèr velutino-tomentosis, paniculis laxis nutantibus ramulis pedicellisque capillaribus pilis patulis, calycibus extûs tomentosis: laciniis intûs barbatis longitudine tubi hemisphærici, stigmatibus distinctis stylum æquantibus, folliculis polyspermis extûs velutinis intûs pilosiusculis.

Sterculia mollis, *Wall. List*, n. 1131, *fid. sp. Herb. Wallichiani in Musco Soc. Linn. Lond.*

Loc. Nat. Martabania, *Dr. Wallich.*

S. (rubiginosa) foliis cuneato-oblongis acutis v. constrictè acuminatis basi obtusâ adultis supèr glabris subtèr pubescentibus venoso-rugosis, stipulis longitudine petiolorum, ^{2392]} paniculis folia vix æquantibus, calycibus extûs tomentosis: laciniis intûs densè barbatis tubum turbinatum superantibus.

Sterculia rubiginosa, *Vent. Malm.* ii, fol. 91 *in adnot. fide specim. in Herb. D. de Jussieu et D. Smith a Thouin.*

Sterculia angustifolia, *Jack in Mal. miscell.*, vol. i.

Sterculia Jackiana, *Wall. List*, n. 1134, *sec. specimina in Herb. Wallich. Musei Soc. Linn.* c quibus character specif. desumptus.

Loc. Nat. Java, *Ventenat.* Pulo Pinang et Singapore, *Jack et Wallich.*

S. (cuneata) foliis cuneato-obovatis ellipticisve acutiusculis basi obtusâ adultis supèr glabris subtèr pubescentibus, petiolis stipulas subulatas vix æquantibus, racemis terminalibus subsimplicibus, calycis laciniis intûs strigoso-barbatis tubo cyathiformi longioribus.

Loc. Nat. Insulæ Philippinæ, *D. H. Cuming.*

S. (ferruginea) foliis oblongo-lanceolatis attenuato-acuminatis basi obtusiusculâ adultis supèr glabris subtèr pubescentibus venoso-rugosis, ramulis petiolis calycibusque extûs ferrugineo-tomentosis, racemis subcompositis, calycis laciniis tubo duplè longioribus, stigmatibus recurvis stylo arcè deflexo dimidio brevioribus.

Loc. Nat. Ins. Philippinæ, *D. H. Cuming.*

S. (stipularis) foliis cuneato-oblongis acuminatis membranaceis basi cordatâ adultis supèr glabriusculis subtèr petiolisque undique tomentosis, stipulis petiolos æquantibus, racemis compositis elongatis pendulis furfuraceo-pubescentibus, calycis laciniis intùs strigoso-barbatis tubo turbinato paullò longioribus.

Loc. Nat. Insulæ Philippinæ, *D. H. Cuming.*

S. (oblongata) foliis oblongis acutiusculis basi obtusissimâ adultis supèr glaberrimis subtèr glabriusculis, paniculâ foliis (spithameis—pedalibus) breviorè, calycis laciniis tubum urceolatum glabriusculum vix æquantibus subulatis pubescentibus demùm solutis.

Loc. Nat. Insulæ Philippinæ, *D. H. Cuming.*

S. (grandifolia) foliis cuneato-oblongis basi (apiceque?) obtusis adultis glabris, racemis simplicibus petiolo elongato brevioribus, calycis laciniis linearibus utrinque tomentosis: tubo brevissimo.

Loc. Nat. Java, *Spangler in Herb. D. Hooker.*

S. (cuspidata) foliis oblongo-ovatis acumine lineari basi obtusis utrinque glaberrimis, racemis simplicibus, calycibus extùs glabris: laciniis subulatis intùs pilosiusculis tubum turbinatum æquantibus.

Loc. Nat. Sumatra, *D. Raffles.*

S. (insularis) foliis ovali-oblongis acutis basi obtusâ utrinque glabris, floribus paniculatis, calycis tubo turbinato glabro laciniis intùs barbatus demùm distinctas et patentes æquante.

Loc. Nat. Insula Moluccana Honimoa, *Ch. Smith.*

S. (parviflora) foliis oblongo-ovatis acuminatis basi transversâ subcordatâ adultis subtèr tenuissimè pubescentibus, floribus 5-fidis paniculatis tomento adpresso, calycis tubo urceolato laciniis adpressé tomentosis longiore.

Sterculia parviflora, *Roxb. Fl. ind.* iii, p. 147. *Wall. List*, n. 1121.

Loc. Nat. Sillet, *Roxb. et Wallich.*

S. (quadrifida) foliis ovatis cordatis acutis obtusiusculisve adultis utrinque glabris, floribus subpaniculatis plerisque 4-fidis, calycibus extus laciniisque tubum urceolatum æquantibus intus tomentosus.

α. folia obtusiuscula, tomentum paniculæ et calycis laud adpressum.

β. folia acuta, tomentum paniculæ et calycis extus subadpressum.

Loc. Nat. Nova Hollandia; *α.* in orâ orientali, *β.* in orâ septentrionali, ann. 1803 legi.

S. (urceolata) foliis ovatis ellipticisve acutis basi obtusâ adultis glabriusculis, floribus paniculatis, calyce tomentoso: laciniis tubum ovatum subæquantibus intus barbatis pilis strictis.

Sterculia urceolata, *Smith in Rees's Cyclop.*: fid. exempl. in *Herb. Smith. in Mus. Soc. Linnean.* An hujus loci *S. Candollei*, *Wall. Pl. asiat. rar.* i, p. 4, quæ *Sterculia populifolia α*, *DeCand. Prodr.* i, p. 483?

Loc. Nat. Honimoa, Ins. Moluc., *Ch. Smith.*

S. (ceramica) foliis ovatis subcordatis acutiusculis adultis utrinque paniculisque glabris, calycibus glaberrimis alabastro obtuso.

Loc. Nat. Insula Moluccana Ceram, *Ch. Smith.*

Obs. An lacinia calycis apice connexæ?

S. (Tragacanthæ) foliis ovatis acutis basi obtusâ adultis super pube rarâ conspersis subtèr tomentosus, calycis laciniis tubum turbinatum æquantibus.

Sterculia Tragacanthæ, *Lindl. in Bot. regist.*, 1353.

Loc. Nat. Africa Æquinotialis prope Sierra Leone.

S. (obovata) foliis obovatis acumine brevissimo basi obtusâ adultis super glabriusculis subtèr pubescentibus, folliculis tomentosus.

Loc. Nat. Africa Æquinoctialis ad flumen Congo, Christian Smith.

TETRADIA.

CHAR. GEN. *Calyx* 4-part. (nunc 3-part.). *Antheræ* 8, simplici serie, loculis omnibus parallelis. *Ovaria* 4, ovulis numerosis. *Stigmata* recurva. *Arbor* (Javanica) *foliis indivisis, racemis axillaribus brevibus monoicis, floribus hermaphrodito-femineis columna nulla, aliquoties majoribus masculo cui columna manifesta gracilis et rudimenta minuta pistilli.*

Obs. Pericarpia et Semina ignota statio generis? incerta: propè *Brachychiton* posui ob ovula basi parum obliqua foramen hilo proximum indicante.

Tetradium, *Lour. cochin.*, p. 91, fide exemplaris ab auctore ad D. Banks missi *Fagaræ* species est.

TETRADIA *Horsfieldii*.

Loc. Nat. Java, Dr. Horsfield.

BRACHYCHITON.

[234]

Sterculiæ subgenera sequentia. *Brachychiton*, *Poecilodermis* et *Trichosiphon*, *Endl. Gen.*, p. 994.

CHAR. GEN. *Calyx* 5-fidus. *Antheræ* congestæ. *Styli* cohærentes. *Stigmata* distincta, v. in unicum peltatum coalita. *Folliculi* coriaceo-lignei, polyspermi. *Semina* albuminosa, pube stellari tecta, mutuò et fundo folliculi cohærentia. *Embryonis radícula* hilo proxima!—Arbores (Novæ Hollandiæ) *foliis lobatis indivisisve.*

* *Calycis æstivatio induplicata (tubus squamis numerosis inflexis).* *Sterculiæ* subgenus BRACHYCHITON, *Endl.*

BRACHYCHITON (*ramiflorum*) *foliis cordatis circumscriptione subrotundis trilobis obtusis utrinque folliculisque extus tomentosis.*

Loc. Nat. In orâ septentrionali Novæ Hollandiæ ann. 1802—3 legi.

** *Calycis æstivatio valvata.*

Br. (*incanum*) foliis altè 5-lobis tomentosis subtèr incanis : lobis acuminatis, folliculis extùs tomentosis.

Obs. Æstivatio ignota, an induplicata?

Loc. Nat. In orâ septentrionali Novæ Hollandiæ ann. 1819 legit b. A. Cunningham.

Br. (*platanoides*) foliis altè 5-lobis acutis glabris, racemis subsimplicibus, calyce infundibuliformi, stigmatè peltato, folliculis extùs glabris.

Trichosiphon, *Endl. Gen.*, p. 994.

Loc. Nat. In orâ orientali, intra tropicum, Novæ Hollandiæ ann. 1802 legi.

Br. (*populneum*) foliis ovatis acuminatis indivisis trilobisve basi acutâ v. obtusâ glaberrimis, racemis axillaribus subsimplicibus, calycibus campanulatis, folliculis elongato-stipitatis.

Poecilodermis, *Endl. Gen.*, p. 994.

Loc. Nat. In orâ orientali, extra tropicum, Novæ Hollandiæ ann. 1803—4 legi.

Br. (*diversifolium*) foliis ovatis elongato-acuminatis basi obtusâ v. cordatâ glaberrimis (quandoque oblongis linearibusve), folliculis extùs glabriusculis abbreviato-stipitatis.

Loc. Nat. In orâ septentrionali Novæ Hollandiæ ann. 1803 legi.

PTERYGOTA, *Endl. Gen.*, p. 995.

CHAR. GEN. *Calyx* 5-partitus. *Antheræ* congestæ v. polyadelphæ. *Stigmata* distincta. *Folliculi* lignei, polyspermi. *Semina* apice alata! albuminosa. *Radicula*

embryonis hilo proxima.—Arbor (Indiæ Orientalis) *foliis indivisis*.

PTERYGOTA alata.

Sterculia alata, *Roxb. Coromand.* iii, p. 84, t. 287.

Loc. Nat. Ind. Or., Silhet et Chittagong, D. Roxburgh.

HILDEGARDIA.

[235]

Sterculiæ subgenus *HILDEGARDIA*, *Endl. Gen.*, p. 994.

CHAR. GEN. *Calyx* 5-partitus. *Antheræ* congestæ. *Folliculi* membranacei, venosi, apice alati, tardiùs dehiscences. *Semina* albuminosa. *Radicula embryonis* hilo proxima.—Arbor (Indiæ Orientalis) *foliis cordatis indivisis*.

HILDEGARDIA populifolia.

Sterculia populifolia, *Roxb. Fl. ind.* iii, p. 148. *Wall. Pl. asiat. rar.* i, p. 3, t. 3.

Loc. Nat. Ind. Or., Roxburgh et Wallich.

FIRMIANA, Marsili in Act. Patav. i, p. 116, t. 1 et 2.

Sterculiæ subgenera *FIRMIANA* et *ERYTHROPSIS*, *Endl. Gen.*, p. 994.

CHAR. GEN. *Calyx* v. 5-partitus tubo brevissimo, v. tubulosus semiquinquefidus. *Antheræ* (15) congestæ v. polyadelphæ. *Styli* coaliti, *stigmata* lobato. *Pericarpia* longè ante maturitatem aperientia, membranacea, foliiformia. *Semina* 2—4, albuminosa. *Embryo* transversus; *Radicula* infera.—Arbores Asiatici (Chinensis et Indiæ Orientalis) *foliis lobatis*.

* *Calyx* 5-partitus; tubo brevissimo fasciculis quinque pilorum intus barbato. *Sterculiæ* subgenus *FIRMIANA*. *Endl. Gen.*

FIRMIANA platanifolia.

Sterculia platanifolia, *Linn. fil. Suppl. Plant.* p. 422.

Sterculia pyriformis, *Bunge in Act. Petrop., sav. etrang.* ii, p. 83, fid. exempl. e Mus. Acad. Petrop. in Herb. D. Hooker.

Sterculia tomentosa, *Thunb. Ic. pl. Japon. dec.* 4.
Loc. Nat. China et Japonia.

** *Calyx tubulosus quinquedentatus v. semiquinquefidus; tubo intus fasciculis 5 pilorum.* ERYTHROPSIS, *Lindl.* *Sterculiæ* subgenus ERYTHROPSIS, *Endl. Gen.*

F. colorata.

Sterculia colorata, *Roxb. Coromand.* i, p. 26, t. 25.

α. calyx extus furfuraceo-pubescent, pube stellari radiis abbreviatis rigidis: tubo dentibus ferè quater longiore foliis adultis subtus glabriusculis.

β. calyx extus furfuraceo-pubescent, pube stellari radiis abbreviatis rigidis: tubo dentibus quater et ultra longiore, foliis adultis subtus pube copiosa scabris.

St. fulgens, *Wall. List*, n. 1135.

Forsan distincta species.

γ. calyx extus pube stellari radiis laxis patentibus velutino: tubo dentibus vix ter longiore.

An species distincta?

Loc. Nat. Ind. Or. et Ins. adj., *Roxburgh et Wallich.*

236]

SCAPHIUM.

Sterculiæ subgenus SCAPHIUM, *Endl. Gen.*

CHAR. GEN. *Flores* monoici. *Calyx* 5-fidus. *Antheræ* (15) congestæ. *Ovaria* disperma. *Styli* coaliti. *Stigma* 5-lobum. *Pericarpia* longè ante maturitatem aperientia, foliacea, navicularia. *Semen* unicum, albuminosum? *Embryonis radícula* hilo proxima.—Arbor (Indiæ Orientalis) *foliis indivisis.*

SCAPHIUM *Wallichii.*

Sterculia scaphigera, *Wall. List*, n. 1130.

Loc. Nat. Martabania, *Wallich.*

PTEROCYMBIUM (*v. suprà*).

COURTENIA.

CHAR. GEN. *Calyx* 5- (raro 4-) fidus, patens. *Antheræ* (10) simplici serie æquidistantes; loculis omnibus parallelis. *Ovaria* 10! (nunc 11—12, rarò 5?). *Stigmata* 10 recurva (stylo brevissimo). *Folliculi* crassi, polyspermi. *Semina* exalbuminosa. *Radicula* hilo proxima.—Arbores (Africæ Æquinoctialis); foliis *altè lobatis*; floribus *sæpius congestis*.

COURTENIA *Afzelii*, caudice simplicissimo, foliis palmato 5—3-lobis cordatis: lobi medii dimidio inferiore angustiore, floribus caulinis congestis.

Loc. Nat. Africa Æquinoctialis prope Sierra Leone et Congo. In Hort. Reg. Kew. florentem vidi ann. 1808 et 1843.

C. triloba, foliis circumscriptione cuneatis basi obtusa palmato-trilobis: lobi medii dimidio inferiore haud angustiore, floribus axillaribus congestis.

Loc. Nat. Africa Æquinoct., Senegambia, *Heudelot in Mus. Paris. et Herb. D. Delessert.*

C. ? heterophylla, foliis palmato-trilobis indivisisve basi cuneata, racemis laxis, ovariis quinque.

Sterculia heterophylla, *Palis. Fl. d'Oware*, p. 67, t. 40.

Loc. Nat. Africa Æquinoct., Oware.

Obs. Fructus ignotus ideoque et ob ovaria tantum quinque dubii generis planta, sed habitus ferè Courteniae trilobae quacum etiam convenit antherarum dispositione et numero, defectu columnæ et stigmatum forma.

COLA.

Sterculiæ subgenus COLA, *Endl. Gen.*

CHAR. GEN. *Calyx* 5-fidus, patens. *Antheræ* (10) simplici serie; loculis divaricatissimis! *Ovaria* multiovulata. *Stigmata* sessilia, distincta, filiformia. *Folliculi* dehiscentes, polyspermi (quandoque monospermi). *Semina* exalbuminosa. *Radicula embryonis* hilo proxima.—Arbores *foliis indivisis glabris*.

^{237]} *Obs.* Ad hoc genus vix pertinet planta e Sierra Leone, ab Afzelio lecto, habitu quidem simili, stipiti genitalium nullo, stigmatibus distinctis sessilibus, floribus axillaribus, foliis cuneato-ellipticis basi attenuatis utrinque glaberrimis glabris, sed diversa antherarum loculis parallelis.

COLA *acuminata*, foliis elliptico-oblongis acuminatis basi acutiuscula adultis utrinque glabris, calycibus altè 5-fidis, genitalibus sessilibus.

Sterculia acuminata, *Palis. Flor. d' Oware*, i, p. 41, t. 24.

——— *grandiflora*, *Vent. Malm.* ii, p. 91 *in adnot.*, ramulus cum racemo hermaphrodito-femineo.

——— *nitida*, *Vent. Malm.*, l. c., ramulus floribus masculis.

——— *macrocarpa*, *Don, Dict.* i, p. 515.

——— *verticillata*, *Schumach. Plant. guin.*, p. 240.

——— *oblongifolia*, *DeCand. Prodr.* i, p. 482?

Lunania Bichy, *DeCand. Prodr.* ii, p. 92.

Loc. Nat. In Africâ Æquinoctiali ad Sierra Leone, Congo; a nigritis olim in Antillis introducta; culta etiam in Brasiliâ, Mexico?, et Insulâ Mauritiî.

C. *cordifolia*, foliis orbiculato-ovatis cordatis indivisis passimque semitrilobis adultis subtèr pubescentibus, calycibus urceolatis semi-5-fidis, genitalibus brevissimè stipitatis.

Stereulia cordifolia, *Cav. diss.* v, p. 286, t. 143, f. 2 (exclus. fruct. ad St. tomentosam sec. auctores flor. senegamb. pertinenti), *Guillem. Perrottet et Richard, Flor. senegamb.*, p. 79, t. 15.

Loc. Nat. Africa Æquinoct., Senegambia.

Obs. Ad Colam retuli præsertim ob antherarum loculos divaricatissimos, semina exalbuminosa et radiculam embryonis hilo proximam, attamen differt a *C. acuminata* floribus (multoties minoribus) suburecolatis 5-dentatis et, secundum D. Perrottet, seminibus arillatis, necnon habitu.

HERITIERA, *Dryand. in Hort. Kew.* ed. 1, iii, p. 546.

CHAR. GEN. *Calyx* 5-fidus. *Antheræ* 5. *Ovaria* uniovulata! *Pericarpia* folliculiformia, lignea, clausa, dorso alata. *Semen* exalbuminosum. *Radicula* hilo proxima.—*Arbores* (Asiaticæ) *littorææ*; foliis *indivisis subtus lepidotis squamulis incisis*; floribus *axillaribus subpaniculatis*.

HERITIERA littoralis, foliis elliptico-oblongis ovatisve obtusiusculis.

Heritiera littoralis, *Dryander in Ait. Kew.* ed. 1, vol. iii, p. 546.

H. fomes, *Buchanan in Sym. Ava. Willd. sp.* iv, p. 972. *DeCand. prodr.* i, p. 484.

H. minor, *Lam. dict.* iii, p. 229. *DeCand. prodr.* i, p. 484.

Loc. Nat. Littora Ind. Or. Insul. Molucc. Philipp Java et Nov. Holl.

H. attenuata, foliis lanceolatis acuminatis.

Heritiera attenuata, *Wall. List*, n. 1140.

Loc. Nat. Ora Martabanicæ et Tenasserim, *Wallich*.

GENUS DUBIÆ TRIBUS.

MICRANDRA.

CHAR. GEN. *Masc.*—*Calyx* turbinatus, 5-fidus, patens, ^{238]} æqualis, æstivatione valvatâ. *Cor.* 0. *Antheræ* 5 (6?) biloculares, loc. parallelis, duplici verticillo, adnatæ medio columnæ filiformis ultra productæ, indivisæ. *Fem.* ignota.

Arbor? floribus *paniculatis*; foliis *ternatis foliolis petiolatis, petiolis partialibus cum communi haud articulatis, nec apice teretibus v. incrassatis*, ideoque vix ad Sterculiaceas pertinens.

MICRANDRA ternata.

Loc. Nat. Brasilia? forsan Para? v. s. in Mus. Paris; ex Herb. Ulyssip.

Obs. Foliola elliptico-oblonga acuta, adulta super glaberrima nitida subtus pube tenui simplici conspersa venis primariis parùm eminentibus costata.

ACTINOPHORA FRAGRANS, Wallich.

(339)

TAB. XLVI.

ACTINOPHORA, Wallich.

CHAR. GEN. *Calyx* 5-partitus; fructûs auctus, patens, foliaceus. *Petala* angusta, diù persistentia. *Stamina* indefinita, distincta. *Ovarium* 4-loculare; loculis biovulatis. *Stigmata* 4 (rarò 3 v. 5), indivisa. *Pericarpium* evalue, ruptile, moniospermum. *Semen* erectum.

Arbor parva v. *Frutex subarborescens, multicaulis*. *Folia alternâ, stipulata*. *Inflorescentia axillaris*.

ACTINOPHORA fragrans, Wallich, List, n. 1163.

DESCR.—*Arbor* v. *Frutex multicaulis*, 10—12-pedalis; caules partiales inordinatè ramosi, ramis ramulisque patulis parumque flexuosis, ultimis pube brevissimâ tomentosus. *Folia* simplicia, alterna, petiolata, stipulata, cuneato-oblonga, acutiuscula, 3—3½ uncias longa, 1½ unciam lata, coriacea, trinervia, nervis venisque primariis subtus elevatis, extra medium dentato-repanda, adulta super glabra subter pube stellari et tomento simplici flexuoso tecta: *petioli* breves. *Stipulae* laterales, subulatæ, petiolo breviores. *Racemi* corymbosi, pauciflori, axillares, folio breviores. *Flores*, paulò ante expansionem visi, parvi. *Calyx* 5-partitus, æqualis, æstivatione valvatâ; laciniis coriaceis, ovatis, acutiusculis, planis, pube stellatâ utrinque tectis extûs copiosiore, post anthesin plurimùm auctis, trinerviis, reticulato-venosis, planis, subscarioso-membranaceis, pube rariore conspersis. *Petala* 5 foliolis calycis alternantia, mox ante expansionem calycis parva, lanceolata, basi parùm attenuata, post anthesin diù remanentia, elongata, subemarcida, lineari-lanceolata, basi valdè attenuata, filamenta emarcida superantia. *Stamina*

numerosa, simplici serie hypogyna, æquidistantia, distincta. *Filamenta* subulata, glabra, omnia antherifera. *Antheræ* oblongæ, obtusæ, paulò supra basin emarginatam insertæ; connectivo angusto; loculis parallelis longitudinaliter dehiscentibus. *Pollen* subglobosum, hispidulum. *Ovarium* ovatum, sessile, pube stellatâ copiosâ, 4-loculare; loculis dispermis; ovulis erectis, collateralibus, cuneato-obovatis. *Stigmata* 3—4 (quandoque 5), indivisa, semiteretia, papillosa. *Pericarpium* evalve, pergamenæo-crustaceum, subglobosum, calyce aucto patulo, filamentis emarcidis, et diù ²⁴⁰¹ petalis elongatis sed calyce brevioribus cinctum, ruptile, monospermum. *Semen* subsphæroideum, glabrum, irregulariter sulcis nonnullis longitudinalibus lobatum; umbilico nudo (nec stropholato) prope basin lateris interioris.

Dr. Wallich, with whom the name *Actinophora fragrans* originates, and with whose specimens the plant here described and figured entirely agrees, has given no other account than that it was introduced into the Calcutta Botanic Garden in 1825, from the Isle of France. There is, however, no sufficient reason to believe that *Actinophora* is indigenous either in that island or in Bourbon or in Madagascar, whereas it is certainly a native of Java, Dr. Horsfield having found it in several districts, chiefly in the medial and eastern parts of the island, in teak forests in a rich soil, or in low ridges extending to the sea-shore. He adds that the wood, which is very hard and durable, is used for various domestic and mechanical purposes, and that it is so ponderous as to be occasionally employed for anchors of small native praoes and canoes. Its Javanese name is *Walli-kookoon*.

With respect to the affinities of *Actinophora*, it certainly does not belong to *Buttneriaceæ* as I originally defined it, but this may equally be said of several genera at present included in that order, and which like *Actinophora* are more obviously referable to *Tiliaceæ*; at the same time, as I observed in proposing the separation of *Buttneriaceæ*, these two families gradually pass into each other.

From all the genera of *Tiliaceæ* at present described,

Actinophora is readily distinguishable; its more remarkable characters being the enlarged subfoliaceous spreading calyx, accompanying the crustaceous evalvular monospermous pericarpium.

TAB. XLVI. *Fig. 1.* A branch of *Actinophora fragrans*, of the natural size. *Fig. 2.* A flower just before expansion, with the calyx removed. *Fig. 3.* A stamen, magnified. *Fig. 4.* The ovarium, cut open longitudinally, crowned with the stigmata. *Fig. 5.* An ovule. *Fig. 6.* The fruit, with the enlarged persistent calyx, petals and stamina. *Fig. 7.* The enlarged calyx, seen from without. *Fig. 8.* The fruit, separated and slightly enlarged. *Fig. 9.* The same, cut open and with the seed removed. *Fig. 10.* The seed. *Fig. 11.* The convoluted cotyledons.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 272-4503

241]

SARCOSTIGMA HORSFIELDII.

TAB. XLVII.

SARCOSTIGMA, *Wight and Arnott in Edin. New Philos. Journ.* xiv, p. 299 (1833).

CHAR. GEN. *Flores* unisexuales (dioici). *Perianthium* duplex; utrumque calycinum 5-fidum (rarò 4-fidum); interius intra exterius sessile, æstivatione valvatâ. Masc. *Stamina* 5 (rarò 4), tubo brevissimo perianthii inserta, ejusdem laciniis alternantia, exserta. *Antheræ* versatiles. Fem. *Ovarium* uniloculare, biovulatum; ovulis ab apice cavitatis pendulis. *Stigma* sessile, depresso-capitatum, indivisum. *Pericarpium* drupaceum, perianthio utroque, inferiore indurato, subtensum. *Semen* . . .

Frutices volubiles, tenuè pubescentes, glabriusculi. Folia alterna, simplicia, integerrima, penninervia, reticulato-venosa, coriacea, petiolata, exstipulata. Spicæ e glomeris alternis paucifloris indivisæ.

SARCOSTIGMA *Horsfieldii*, foliis obovato-oblongis, spicarum rachibus perianthioque interiore extus pubescentibus.

Loc. Nat. In provinciâ orientali Blambangan, ubi anno 1806 et in pago Pagetan anno 1809 detexit D. *Horsfield*. Javanis *Kamaras*.

DESCR.—Frutex volubilis, ramosissimus; caule tereti (diametro sæpè bipollicari, D. *Horsfield*). Folia alterna, petiolata, exstipulata, integerrima, coriacea, glabra, impunctata, penninervia, reticulato-venosa, obovato-oblonga, extra medium paulò latiora, quandoque acutiuscula, nunc obtusissima et emarginata, 5—6 pollicaria, ultra duos pollices lata. Petioli teretes, semuiciales, juxta basin

arcuatim recurvi. Spicæ longæ, sæpiùs 2—3 aggregatæ quandoque solitariae, nunc ramæ absque cicutrice folii delapsi, nunc axillares, indivisæ sed e glomerulis alternis 3—4 floris formatæ; rachi tereti pedicellisque flosculorum, perianthio brevioribus, extùs pubescentibus. Flores parvi. Perianthium exterius abbreviatum, 5-lobum lobis rotundatis; interius æstivatione valvatâ altè 5-fidum; laciniis planis, lanceato-linearibus, acutis, intùs glabris, supra medium recurvis. Stamina 5: filamenta laciniis perianthii^[242] interioris alternantia, ejusdem tubo inserta, filiformia, lacinias limbi æquantia v. superantia. Antheræ versatiles, ovales; loculis parallelis, approximatis, longitudinaliter dehiscentibus. Rudimentum pistilli parvum, pubescens. Fem. perianthii, ut in Masc. interioris, laciniis apice recurvis. Ovarium ovatum, pubescens, uniloculare, bivulatum; ovulis collateralibus, ab apice cavitatis pendulis. Pericarpium drupaceum, oblongum, ipsâ basi perianthio interno indurato vix aucto subtensum; putamine crasso-pergameneo, rugoso (pulpâ succulentâ eduli, *D. Horsfield*). Semen

My remarks on the affinities of *Sarcostigma* are annexed to the account of the following genus, *Iodes*.

TAB. XLVII. *Fig. 1.* Flowering branch of a female plant of *Sarcostigma Horsfieldii*. *Fig. 2.* Part of a male spike; both of natural size. *Fig. 3.* A male flower, moderately magnified. *Fig. 4.* A male flower with quaternary division; probably a rare occurrence. *Fig. 5.* A stamen, to show the mode of dehiscence and insertion of filament. *Fig. 6.* The outer floral envelope or calyx. *Fig. 7.* A female flower, magnified in the same degree as the male; to show the expansion, recurvation, and external pubescence of the segments of the inner perianthium or corolla. *Fig. 8.* An ovary, slightly magnified and cut open to show the two pendulous ovula. *Fig. 9.* A drupe, nearly ripe, natural size.

IODES OVALIS, Bl.

243]

TAB. XLVIII.

IODES, *Blume, Bijdr.* p. 29.

CHAR. GEN. *Flores* unisexuales (dioici), 5-fidi. *Perianthium* duplex: utrumque calycinum; interius 5-partitum, majus, æstivatione valvatâ. Masc. *Stamina* 5, tubo abbreviato perianthii (corollæ?) inserta, ejusdem laciniis alternantia iisque breviora. *Antheræ* stantes. Fem. *Ovarium* uniloculare, biovulatum; ovulis ab apice cavitatis pendulis. *Stigma* sessile, depresso-capitatum, umbilicatum. *Drupa* exsucca, stigmate apiculata. *Semen* unicum: *Albumen* semini conforme; *Embryo* dimidio albumine longior; *Radicula* supera.

Frutex *volubilis, cirrhis axillaribus scandens; foliis oppositis, integerrimis, petiolatis, exstipulatis; floribus parvis, paniculatis, axillaribus.*

IODES *ovalis*, Blume, *Bijdr.* p. 30 (1825); Hassk. Hort. Bogor. p. 172, n. 798.

Loc. Nat. In plagis orientalibus primum 1806 et postea in variis locis detexit D. Horsfield. Javanis *Jungetan* et *Jagal-Kedang*.

DESCR. Frutex (quandoque subarborescens, D. Horsfield) pubescens, scandens, cirrhis axillaribus indivisis nunc oppositifloris. Folia opposita, petiolata, exstipulata, coriacea, penninervia, reticulato-venosa, ovata, acuta, basi obtusa, 4—5-uncialia, ultra 2 pollices lata; petioli semunciales, teretiusculi. Paniculæ axillares, compositæ, corymbosæ. Flores parvi, pubescentes (odorati D. Horsfield). Masc. *Stamina* 5, perianthio interiori (corollâ potiùs) aliquoties breviora, ejusdem laciniis alternantia; filamentis brevissimis; antheris basifixis, loculis approximatis parallelis longitudinaliter dehiscentibus. Rudimentum pistilli. Fem.

Floribus paulò post anthesin solùm visis, et tunc etiamque fructu maturo perianthio unico tantùm persistenti nec aucto nec indurato. Drupa exsueca, rugosa, magnitudine pisi majoris. Semen unicum maturescens; albumen carnosum, album, magnitudine seminis; Embryo respectu pericarpium inversus; Cotyledonibus subrotundis, planis, foliaceis; Radiculâ rectâ, brevissimâ, superâ.

Obs. I. Botanists are not agreed as to the light in which the two floral envelopes of *Sarcostigma* and *Iodes* are to be viewed. Both Klein, in describing the plant from which *Sarcostigma* was formed, and Blume, in his character of *Iodes*, have termed the outer envelope calyx and the inner corolla; while Drs. Wight and Arnott, by whom the genus *Sarcostigma* was established, have considered the outer envelope an involucre and the inner calyx. In support of this opinion the persistence and even induration of the inner envelope in *Sarcostigma* would no doubt be adduced; but they were also, it may be supposed, determined by referring their genus to *Hernandiaceæ*, there being in *Hernandia* an undoubted involucre, but that involucre containing several flowers. On the other hand, the alternation of division of the two envelopes, and their close approximation in both these genera, are characters of at least equal importance; to which may be added the many analogous points of structure between these genera and *Cardiopteris*, in which it will hardly be disputed that both calyx and corolla are present.

Obs. II. As to the affinities of *Sarcostigma* and *Iodes*, the former genus was referred by Wight and Arnott to *Hernandiaceæ*, a family proposed by Blume in his 'Bijdragen,' p. 550, and adopted by Endlicher, consisting of *Hernandia* and *Inocarpus*, two genera which do not appear to me very nearly related to each other, and from both of which *Sarcostigma* is readily distinguished by many important characters. In the following year Professor Arnott, in describing *Nansiatum* of Dr. Buchanan Hamilton, proposed to place it next in affinity to *Phytocrene* of Wallich (*Gyno-*

cephalum of Blume), and to form of these two genera the new family of *Phytocrenæ*. It is remarkable that he did not also include in this family *Sarcostigma*, which so obviously belongs to it: this has been recently done by M. Planchon, who also justly proposes to add *Iodes*, appended by Blume to *Menispermeæ*, as well as *Miquelia* or *Jenkinsia*, which he regards as distinct genera, but which I think ought to be united. He also includes *Pyrenacantha*, properly united with *Adelanthus* of Endlicher; but this genus I exclude, admitting, however, its near relationship.

Of PHYTOCRENEÆ so constituted I subjoin the distinguishing characters of the family, as well as the differential characters of the genera belonging to it.

PHYTOCRENEÆ, *Arnott, in Edinb. New Phil. Journ.* xvi, p. 314 (1834).

Flores diclines (dioici), inconspicui. *Perianthium* duplex (*Calyx* et *Corolla*) utrumque 4—5-fidum, calycinum; interius majus, æstivatione valvatâ, intra exterius in quibusdam pedicellatum. MASC. *Stamina* 4—5, laciniis perianthii interioris alternantia; *filamenta* nunc hypogyna et ipsâ basi coalita, nunc tubo perianthii inserta; *Antheræ*: loculis longitudinaliter dehiscentibus. Rudimentum *pistilli*. FEM. *Ovarium* liberum, uniloculare, biovulatum; ovulis ab apice cavitatis suspensis. *Stigma* sessile, indivisum v. bifidum. *Drupa* monosperma. *Albumen* semini conforme; *Embryo* magnitudine fere albuminis, cotyledonibus foliaceis.

Frutices volubiles v. scandentes; foliis alternis rarò oppositis, integerrimis v. lobatis, exstipulatis.

PHYTOCRENE, *Wallich, Pl. Asiat. rar.* vol. iii, p. 11, t. 216.

Flores utriusque sexûs 4-fidi, capitati. *Stamina* hypogyna. *Antheræ* versatiles. *Stigma* bilobum, obtusum.

Folia indivisa v. lobata.

SARCOSTIGMA, *Wight and Arnott, in Edinb. New Phil. Journ.* xiv, p. 299.

Flores spicati, 5-fidi (rarò 4-fidi). *Stamina* tubo perianthii inserta, ejusdem laciniis longiora. *Antheræ* versatiles. *Stigma* depresso-capitatum. *Drupa* pulposa.

Frutices volubiles; foliis alternis integerrimis.

IODES, *Blume.*

[245]

Flores 5-fidi, paniculati. *Stamina* tubo perianthii longioris inserta, ejusdem laciniis breviora. *Antheræ* stantes. *Stigma* depresso-capitatum. *Drupa* exsucca.

Frutex cirrhis axillaribus scandens; foliis oppositis, integerrimis.

NANSIATUM, *Buchanan Hamilton, in Edinb. New Phil. Journ.* xvi, p. 314.

Flores 5-fidi, spicati. *Stigmata* duo, acuta, recurva. *Pericarpium*

Suffrutex volubilis; foliis alternis, indivisis.

MIQUELIA, *Meisn. Gen.* 152. JENKINSIA, *Griffith, in Calc. Journ. Nat. Hist.* iv, p. 231, t. 12.

Flores 5-fidi. MASC. *Perianthium* interius intra exterius pedicellatum. FEM. *Perianthium* interius intra exterius sessile. *Stamina* sub ovarii rudimento inserta. *Stigma* depresso-capitatum, umbilicatum. *Drupa* exsucca.

Suffrutices volubiles; foliis alternis, integerrimis.

Phytocreneis affine genus,

PYRENACANTHA, *Hooker, Bot. Misc.* ii, p. 107, *Tabb. Suppl.* 9, 10.

Pyrenacantha Adelantho congener, a *Phytocreneis* diversum: *Perianthio* simplici; *Stigmate* radiatim multifido:

quadrat *Staminibus* cum calycis segmentis alternantibus ;
æstivatione valvatâ ; *Pericarpio* indehiscente.

TAB. XLVIII. *Fig.* 1. A flowering male branch of *Iodes ovalis*. *Fig.* 2. The outer perianthium. *Fig.* 3. The deeply divided inner perianthium or corolla. *Fig.* 4. An anthera burst longitudinally, with insertion at base. *Fig.* 5. Rudiment of ovarium in male flower. *Fig.* 6. Ovarium after flowering, subtended by one only of the perianthia. *Fig.* 7. The same laid open, to show the number and insertion of ovula. *Fig.* 8. A ripe drupe, natural size. *Fig.* 9. The seed. *Fig.* 10. The seed cut open lengthways, to show the relative proportion of albumen and embryo. *Fig.* 11. The embryo inverted in respect to pericarpium.

CARDIOPTERIS LOBATA, Wall. List, n. 8033. [246]

TAB. XLIX.

CARDIOPTERIS, Wallich.

CHAR. GEN. *Flores* hermaphroditi? (v. monoici). *Calyx* 5-partitus, æstivatione imbricatâ, persistens. *Corolla* monopetala, rotata, limbo æquali 5-fido, æstivatione imbricatâ, decidua. *Stamina* 5, tubo corollæ sub sinibus limbi inserta. *Antheræ* longitudinaliter dehiscentes. *Ovarium* liberum, uniloculare, biovulatum; *ovulis* pendulis. *Stigmata* duo: altero (vero) post anthesin aucto, emarginato, tardè deciduo: altero (effæcto) capitato, pedicellato, persistenti. *Samara* compressa, alata, monosperma. *Albumen* semini conforme. *Embryo* minutissimus, in apice (respectu pericarpium) albuminis.

Plantæ annuæ v. biennes, volubiles, glabræ, lacte albo scatentes (fid. D. D. Blume et Hasskarl); *Foliis alternis, petiolatis, cæstipulatis, v. lobatis v. integerrimis, cordatis, venosis; Cymis furcatis v. dichotomis v. paniculatis; floribus parvis, secundis, ebracteatis.*

CARDIOPTERIS lobata (Wall. List, n. 8033), foliis angulato-lobatis, cymis furcatis v. semel dichotomis.

Cardiopteris javanica, Blume, *Rumphia*, vol. iii. p. 206, tab. 177.

Peripterygium quinquelobum, Hassk. *Hort. Bogor.* p. 235.

An *Dioscorea sativa*? Kœnig, *MSS. in Mus. Brit.* vol. iii. p. 81; exclus. syn. *Rumph. Amb.* tom. v. p. 482, ad *Cardiopteridem moluccanam*, Blume, *Rumph.* vol. iii. p. 207 pertinente.

Loc. Nat. In plagis orientalibus et medianis Javæ; in planitiis haud multum elevatis supra mare anno 1806

primùm detexit *D. Horsfield*. Javanis orientalibus *Parianom*, medianis *Kanjar Kawang* vel *Rindengan* appellata.

DESCR.—Planta annua? volubilis, glaberrima, præter raches spicarum tenuissimè pubescentes pube acutâ simplici. Rami ramulique flexiles, striati, inanes. Folia alterna, remota, elongato-petiolata, exstipulata, flaccidè membranacea, angulato-lobata, altè cordata; lobo medio acuminato cum mucronulo brevi; reliquis obtusis, muticis; diametro 2—3-pollicari; basi nervis 5—7 lobis respondentibus, immersè reticulato-venosis. Cymæ axillares, pedunculatæ, semel v. bis furcatæ. Flores unilaterales, ebracteati, brevissimè pedicellati, parvi, caput aciculæ mediocris vix superantes. Calyx 5-partitus, æqualis, membranaceus, texturâ subcorollinâ, æstivatione imbricatâ; sepalis ovatis, obtusiusculis, marginibus tenuissimè ciliatis, subnerviis, axi parùm opaciorè. Corolla monopetala, calycem vix superans, subrotata, virescenti-alba, glaberrima; tubo perbrevis, limbo 5-partito æquali; laciniis obovatis, obtusis, concaviusculis, obsoletè immersè venosis, æstivatione imbricatis. Stamina 5, epipetala, æqualia, corollâ breviora, ejusdem sinubus inserta. Filamenta brevina, filiformia, glabra. Antheræ pro ratione flosculi majusculæ, didymæ, utrinque profundè emarginatæ; connectivo membranaceo angusto; loculis apposis, longitudinaliter dehiscentibus. Discus hypogynus nullus. Ovarium uniloculare, biovulatum; ovulis collateralibus, ab apice cavitatis pendulis, altero sæpè minore effæto. Stigmata duo: alterum (imperfectum) sub anthesi magis conspicuum, stylo manifesto insidens, capitatum, læve nec papillosum; alterum (perfectum) sub anthesi vix obvium, postea sensim valdè auctum, demùm emarginatum, superficie interiore imperfectum spectante papillosum. Samara obcordata, nitida, vix uncialis, ipsâ basi angustatâ calyce persistenti vix aucto conniventi subtensa; alâ marginatâ utrinque ipso nucleo subcylindraceo ter quater vè latiore pergamenâ; stigmatè vero plurimùm aucto emarginato, tardè deciduo, stigmatè effæto, capitato, pedicellato, persistenti, haud mutato, subemarcido, coronata.

Semen unicum, testâ? cum samaræ cavitare cohærens et omninò replens, striatum, subcylindraceum; integumentum præterea unicum, membranaceum, albumini arcè adhærens. Albumen semini conforme, aqueo-pallidum, dense carnosum. Embryo in apice (respectu pericarpîi) albuminis, minutissimus; radícula brevis, supera; cotyledon adhucdum indivisus, subglobosus, obtusissimus.

Obs. There are some points both in the botanical history and in the structure of the genus *Cardiopteris* which deserve to be specially noticed. It is probable that a short time before the publication of the 10th edition of his 'Systema Naturæ,' in 1759, Linnæus had particularly examined the figures of the 'Herbarium Amboinense,' for in that edition almost the only figures quoted of the different species of *Dioscorea* are those of that work, and under *Dioscorea sativa* he refers to *Olus sanguinis*, vol. v, p. 482, tab. 180, which is an undoubted and a tolerably good representation of *Cardiopteris moluccana* of Blume; and in proof of Linnæus having no doubt as to the correctness of his reference, he has in his own copy of the work written *Dioscorea sativa* under the figure quoted. This synonym is adopted by Willdenow in his edition of 'Species Plantarum.' J. G. Kœnig, in vol. iii, p. 81 of his manuscripts, formerly in Sir Joseph Banks's Library, now in the Banksian or Botanical Department of the British Museum, quotes the same figure of Rumphius (with a doubt, however, as to the correctness of Linnæus's reference to *Dioscorea*) for a plant which is no doubt *Cardiopteris lobata*, from his description in many respects very good, and with regard to stigma more nearly correct than some recent accounts. Kœnig considers his plant to be monoicous, as does Hasskarl (*supr. cit.*): I also entertained the same opinion, having never found flowers with completely developed pistillum in which the corolla and consequently stamina were present. This, however, might depend on the advanced state of the flower whose corolla had fallen soon after expansion. Dr. Blume, therefore, is perhaps more correct in considering the flowers as hermaphrodite, and I am

inclined to adopt his view. His statements respecting the structure of the pistillum are more liable to objection: he describes the ovulum as orthotropous, having the micropyle at its lower extremity, and the embryo consequently existing at the same point of the seed; my own observations, which may indeed require to be verified, placing the embryo at the upper extremity, or close to the insertion of the seed, the ovulum being consequently anatropous. The external structure of the pistillum is very singular. In an early stage of the flower, immediately before or even at the time of expansion, there are apparently two stigmata: of these the more obvious is capitate, undivided, fleshy, but not papillose, and is supported on a distinct style; the second is quite sessile, much shorter in this stage than the capitate branch, and having its upper or inner surface distinctly stigmatic or papillose. In the next stage, the latter, which I regard as the efficient stigma, gradually enlarges, becoming longer than the capitate organ, which in my opinion is an imperfect stigma; and as in this stage the ovarium though enlarged has not perceptibly increased in diameter, this capitate stigma has the appearance of being lateral. The perfect stigma, which continues to lengthen, its upper surface becoming more evidently hispid or papillose, not unfrequently remains crowning the samara even when ripe; but frequently also it is then deciduous, while the imperfect capitate stigma, which has undergone no change either in size or surface, more generally remains after the real stigma has fallen.

In one of the flowers of a specimen preserved in spirits by Dr. Wallich, and in which the corolla was wanting, and a slight enlargement of ovarium had taken place, I found only one ovulum, the lower extremity of which seemed in some degree to support Dr. Blume's account of the position of micropyle. In a second flower of the same specimen only one apparently perfect ovulum existed, but the funiculus or remains of a second was visible; in the perfect ovulum a more transparent point, which might possibly be micropyle, was lateral; and in a third flower, long after fœcundation, the samara being distinctly formed though not of its full

size, I was able to observe only one ovulum, suspended from a funiculus which was continued on one side into a raphe ending at the lower and wider extremity of the ovulum. These appearances leave the question of the original position of micropyle undetermined.

Dr. Blume considers the capitate body as the true stigma, and I have some reason to believe that this is also the opinion of an excellent and more recent observer. I have little doubt, however, that the emarginate or bidentate branch is the organ performing the function. In connection with and in support of this opinion, I regard the ovarium or pericarpium of *Cardiopteris* as composed of two carpella, the line of junction being in the axis of the samara. This I infer from the relative position of the supposed perfect and imperfect stigmata, the former presenting its papillose surface to the latter, and both being in the plane of the wings of the samara; and these wings, as well as I can judge, being anterior and posterior with relation to the axis of the unilateral spike.

With respect to the affinities of *Cardiopteris*, Dr. Wallich, with whom the genus originated, and who first hazarded a conjecture on the subject, was disposed to refer it to *Sapindaceæ*; Hasskarl (l. supra cit.) has placed it among *Euphorbiaceæ*; and Blume, who considers it as forming a separate family (*Cardiopterideæ*), has placed this family in proximity with *Verbenaceæ* and *Boraginæ*. None of these approximations appear to me satisfactory, and although I am aware of several important objections to the view, I am inclined to consider *Cardiopteris* as an isolated genus or family to be placed at no great distance from *Phytocreneæ*, chiefly through *Iodes*, under which genus I have given some account of that family.

TAB. XLIX. *Fig. 1.* A branch of *Cardiopteris lobata*, in flower and fruit, natural size. *Fig. 2.* An expanded flower, slightly magnified, which shows the relative proportion of calyx, corolla, and stamina, and the external appearance of pistillum, of which the capitate stigma has nearly attained its full size, while the true stigma is as yet

barely visible. *Fig. 3.* A corolla laid open, to show that it is monopetalous, that the stamina are inserted below the sinuses, and that its segments are slightly ciliated, somewhat more magnified than figure 2. *Fig. 4.* The corolla removed, to show more distinctly the external structure of pistillum. *Fig. 5.* A stamen. *Fig. 6.* The calyx surrounding the somewhat enlarged pistillum, on which, from the elongation of the true emarginate stigma, the capitate or spurious stigma has become apparently lateral. *Fig. 7.* A ripe seed-vessel, natural size, with its short pedicle covered by the withered calyx, and crowned by the two stigmata, of which the true or emarginate has greatly increased in size, especially length, while the capitate, not at all enlarged, is barely visible. *Fig. 8.* A seed slightly magnified. *Fig. 9.* The embryo separate, magnified more than fig. 8, and so placed as to indicate that it must have been situated at the upper extremity of the seed.

BENNETTIA JAVANICA.

[249]

TAB. L.

BENNETTIA.

CHAR. GEN. *Flores* unisexuales (dioici), regulares, 5-fidi (minimi). *Calyx* 5-partitus. *Petala* 5 concava, æstivatione marginibus inflexis mutuo applicitis valvata. Masc. *Stamina* 10, petalis sæpiùs cucullatis inclusa. *Antheræ* loculis longitudinaliter dehiscentibus. Fem. *Ovarium* uniloculare, biovulatum, ovulis pendulis. *Stigmata* 3—4, filiformia, uno alterove quandoque bipartito. *Drupa* latior quam longa, monosperma. *Semen* figurâ drupæ. *Albumen* magnitudine seminis. *Embryo* transversus, rectus, longitudine fere albuminis; radícula alteri margini approximata, recta, cotyledonibus foliaceis.

Arbusculæ; foliis *simplicibus*, *alternis*, *integerrimis*, stipulis *lateralibus*, sæpiùs *caducis*. Spicæ *ramulos graciles terminantes*, *solitariae*, *longissimæ*, *pendulæ*.

BENNETTIA *javanica*, masculi floris petalis altè cucullatis glabris: alabastro ultra medium 5-fido depresso lobis retusis, antheris rudimentoque pistilli imbertibus, drupis paulo tantum latioribus quam longis, foliis utrinque ramulisque glabris.

Loc. Nat. In plagis Javæ orientalibus, ubi primum detexit anno 1806 *D. Horsfield*; et anno 1814 regionibus medianis Insulæ observavit. Javanis orientalibus *Klimo Koncher*; medianis *Jirikan* appellata.

DESCR. *Arbuscula* glabra, ramosissima, ramulis virgatis. *Folia* alterna, integerrima, stipulata, brevè petiolata, impunctata, oblonga, super glabra, subter secundum nervum et venas saltem primarias tenuissimè pubescentia. *Stipulæ* laterales, angusto-subulatæ, petiolo paulo breviores, caducæ. *Spica* terminalis, solitaria, longissima (usque pedalis), indivisa, nutans; *mascula* e fasciculis 4—5-floris appropi-

mato-alternis, sessilibus, pedicellis unifloris basi unibracteatis superne ebracteatis; *feminea* floribus solitariis. *Flores masculi* parvi, caput aciculi mediocris subæquant. ²⁵⁰⁷ *Calyx* 5-partitus, æqualis, patens, laciniis planis acutiusculis pubescentibus, pube brevissimâ simplici acutâ. *Petala* 5, cum laciniis calycis alternantia, patentia, altè cucullata, obovata, glabra, apice cuculli retuso, calyce plus duplò longiora, brevissimè unguiculata, axi laminæ intùs incrassatâ prominulâ, æstivatione marginibus inflexis mutuò applicitis valvatâ, alabastro depresso 5-lobo. *Stamina* decem distincta. Filamenta brevissima, latiuscula, sepalis et petalis opposita. *Antheræ* biloculares, loculis connectivo lato distinctis longitudinaliter dehiscentibus; omnes cucullis petalorum semi-inclusæ, ita ut duæ petalo singulo oppositæ esse videantur (exterior interiorque), sed dum exterior ad filamentum petalo oppositum pertinet, interior e loculis respondentibus filamentorum duorum petalis alternantium formata est. *Pollen* simplex, sphæroideum. *Rudimentum pistilli* subovatum, apice depresso indiviso. *Feminei flores* paulo post fecundationem solum visi; petalis concavis haud cucullatis. *Ovarium* sphæroideum, 4—5-sulcum, pubescens, basi calyce parvo subtensum, stigmate triplici vel duplici singulo bifido v. laciniato coronatum, uniloculare, biovulatum, ovulis apici cavitatis affixis, collateralibus, latioribus quam longis, transversim oblongis, utroque angulo acutiusculo. *Drupa* paulò nec duplò longior quam lata, tenuissimè pubescens, pulpâ parcâ; putamine intùs venoso, basi excavato, depressione latâ longitudine totius baseos in cujus centro calyx persistens haud auctus. *Semen* unicum, apici cavitatis affixum, transversim ovale: integumento (testâ ?) simplici. *Albumen* semini conforme, carnosum. *Embryo* transversus, longitudine ferè albuminis. *Cotyledones* foliaceæ, quoad apicem cavitatis accumbentes, ovaes. *Radicula* recta cotyledonibus multoties brevior, margini alteri approximata.

Obs. *Bennettia* is dedicated to the principal author of the present work, John Joseph Bennett, Esq., my friend and colleague in the British Museum.

The generic characters of *Bennettia* are clear and important, but the affinities of the genus, or its place in a natural arrangement, though determinable, are not equally obvious. Many important points of agreement will readily present themselves between *Euphorbiaceæ* and *Bennettia*, but the general resemblance it bears to *Antidesmeæ* is still more striking. In their simple entire alternate leaves with lateral deciduous stipules and in their inconspicuous unisexual flowers they entirely agree, very nearly also in inflorescence, in the structure of ovary, in their monospermous drupaceous pericarpium, and lastly, in the presence and texture of albumen, and in the degree of development of embryo. The principal distinctions would therefore be reduced to the existence of petals in *Bennettia*, to its stamens being equal in number to the divisions or parts of both floral envelopes, to a remarkable and obvious difference in the structure and æstivation of antheræ, and to the singular character of its transverse embryo. The presence of petals may even be regarded as of more than ordinary importance, their usual form in the male flower being necessarily connected with the æstivation of stamens. According to this view, therefore, *Bennettia* may be said to bear the same relation to *Antidesma* (for *Antidesmeæ* contains at present no other well-established genus) that the polypetalous bear to the apetalous genera of *Euphorbiaceæ*. But according to a principle which I proposed for adoption in 1810, *Bennettia* ought not only to determine the place, but also give the name (*Bennettiaceæ*) to the family. The principle referred to is stated in the 'Prodromus Floræ Novæ-Hollandiæ,' p. 351, in a note relative to *Combretaceæ* (an order then first proposed and characterised), in the following terms:—"Hunc ordinem inter polypetalos posui, non solum propter petalorum in pluribus existentiam, sed quia vera natura partium affinitatesque ordinum, ex contemplatione generum in quibus structura magis evoluta quam ex iis in quibus aliqua pars suppressa, tutius erui queant." In 1814, in conformity with the same principle, I placed among polypetalæ *Euphorbiaceæ*, a family to which the same reasoning is still more

strikingly applicable. It is not my intention, however, to propose any change in this respect, for in both cases the names must be considered as established.

^{251]} The affinity between *Euphorbiaceæ* and *Antidesmeæ* is rendered more obvious by the addition to the latter of *Bennettia*. But the structure of ovarium and the monospermous drupaceous pericarpium readily distinguish them.

Iodes and *Sarcostigma* also agree with *Bennettia* in several important points, particularly in their unisexual minute flowers, ovarium with two pendulous ovula, monospermous drupa, and in most respects in the structure of seed. They differ in habit, being twining or scandent shrubs without stipules, in their monopetalous persistent inner perianthium or corolla, in æstivation and reduced number of stamina, in structure of antheræ, and in the embryo being inverted, not transverse.

Obs. II. Several species of *Bennettia* have been discovered in India. One of these, first observed by the late Dr. Jack, at Singapore, was referred by him, though doubtfully, to *Limonia*. Dr. Blume, who had observed a plant of the same genus in Java, for which he adopted the generic name here proposed, having liberally communicated specimens and drawings of this plant, it proves to be identical with *Bennettia javanica*; and lastly, Dr. Wallich has, in the concluding distribution of his great Indian Herbarium, noticed several plants of the same genus, to which he has also given the name *Bennettia*. These plants so closely resemble *Bennettia javanica* and each other, that they are chiefly distinguishable by minute, but, as I believe, constant differences in their male flowers, and in the form of their fruits. Of these supposed species the differential characters are here subjoined.

Bennettia Wallichii, masculi floris petalis concavis vix cucullatis glabris, antheris rudimentoque pistilli imberbibus, drupis transversim oblongis duplò latioribus quam longis varicosis, ramulis foliisque utrinque glaberrimis.

Doc. Nat. India Or. Tavoy. Wall. List, 8585 E.

B. phlebocarpa, drupis latè cordatis sinu clauso reticulato-venosis paulo latioribus quam longis, foliis oblongis subacuminatis utrinque glabris.

Loc. Nat. Singapore. *Wall. List*, 8585 D.

B. Finlaysoni, masc. petalis modicè cucullatis glabris apice deflexo acuto: axi intùs elevatâ, antheris imberbibus, rudimento pistilli apice barbato, foliis oblongis acuminatis basi inæquilaterali subtùs nervo venis primariis ramulisque pubescentibus, spicis masc. foliis longioribus.

Loc. Nat. Pulo-Penang, b. Finlayson. *Wall. List*, 8585 B.

B. affinis, masc. petalis cucullatis glabris: axi intùs elevatâ, antheris imberbibus, rudimento pistilli apice barbato, stipulis ramulis venisque primariis subtùs pubescentibus, spicis folia lanceolato-oblonga acuminata subsessilia vix superantibus.

Loc. Nat. Pulo-Penang, b. Finlayson. *Wall. List*, 8585 B.

B. pedicellata, masc. petalis modicè concavis obtusis: axi intùs parum elevatâ, antheris filamentisque barbatis.

Obs. Sequenti nimis forsàn affinis.

Loc. Nat. Pulo-Penang. *Wall. List*, 8585 C.

B. Jackiana, masc. petalis orbiculato-ovatis concaviusculis: axi intus haud elevatâ, antheris barbatis.

Loc. Nat. Pulo-Penang b. Jack, qui ad *Limoniam* dubitanter retulit. *Wall. List*, 8585 A.

TAB. L. *Fig. 1.* A flowering branch of the male plant of *Bennettia javanica*, natural size. *Fig. 2.* An expanded male flower, slightly magnified. *Fig. 3.* A male flower before expansion. *Fig. 4.* A petal of the same flower, somewhat magnified. *Fig. 5.* Back view of one of the longer stamina, but in which the loculi are too divergent. *Fig. 6.* A front view of one of the shorter stamina. *Fig. 7.* A male flower, the petals and stamina removed, to show the calyx and rudiment of pistillum. *Fig. 8.* A drupe, slightly reduced. *Fig. 9.* The same,

longitudinally divided, which shows the single seed. *Fig.* 10. A seed, longitudinally divided to show its insertion, with the direction of the embryo and the relative proportion of its radicle and cotyledons.

g.
n,
or-

CONTRIBUTIONS
TO THE
BOTANICAL MAGAZINE.

1811—1825.

ce
2
p
ce

g
n
fl
p
L

CONTRIBUTIONS
TO THE
BOTANICAL MAGAZINE.

Zieria Smithii, Bot. Repos., No. 606.
Zieria lanceolata, Brown in *Herb. Banks.*
Bot. Mag. 1395 (1811).

Dillwynia parvifolia, Brown in *Herb. Banks.*
Bot. Mag. 1527 (1813).

GOMEZA RECURVA.

Labellum ecalcaratum, indivisum, bicristatum, sessile, cum basi columnæ apteræ liberæ continuum. *Petala* 2 antica exteriorum connata, labello supposita. *Massæ pollinis* 2, hinc sulco obliquo bilobæ, apice connexæ processu communi stigmatis. *Brown MSS.*

Gomezia recurva, Brown MSS.

We are indebted to our friend, Robert Brown, Esq., for generic character and name, given to it in honour of Bernardinus Antonius Gomez, Physician to the Portuguese fleet, and author of *Botanico-Medical Observations* on the plants of Brazil, a very respectable work, written in the Latin and Portuguese languages.

Bot. Mag. 1748 (1815).

CYRTOPODIUM WOODFORDII.

Petala 5, distincta, secunda. *Labellum* trilobum, cum processu unguiformi baseos columnæ apteræ subarticulatum. *Massæ pollinis* 2, postice bilobæ. *Brown MSS.*

Cyrtopodium Woodfordii, labello ventricoso: lobis laminæ lateralibus intermedio verrucoso callis cristato brevioribus.

Cyrtopodium Andersonii, labello angusto unguiculato: lobis laminæ lateralibus divaricatis intermedio excavato longioribus.

Bot. Mag. 1814 (1816).

Anisomeles malabarica, tomentosa, subglandulosa, caule incano, foliis lanceolatis subtus rugosis, verticillis multifloris, calycibus lanatis dentibus subulatis tubum æquantibus. *Brown MSS.*

Bot. Mag. 2071 (1819).

Pultenæa tenuifolia, capitulis terminalibus subbifloris, fructibus lateralibus, foliis subulato-linearibus muticis supra concavis subtus convexis ramulisque pilosis. *Brown MSS.*

Native of the south coast of New Holland, and probably of Van Diemen's Island. It was observed on several parts of the former by Mr. Brown, to whom we are indebted for enabling us to settle the species, and to distinguish it from the *vestita* of the "Hortus Kewensis," under which name we received it.

Bot. Mag. 2086 (1819).

Pultenæa biloba, capitulis terminalibus paucifloris, foliis cuneiformibus apice dilatato-bilobis supra tuberculato-scabies subtus sericeis; mucrone brevi marginibusque recurvis. *Brown MSS.*

We are indebted to our friend Mr. Robert Brown for

his assistance in determining the species, by whom it was first discovered in New South Wales.

Bot. Mag. 2091 (1819).

CHLORANTHUS MONOSTACHYS.

MONANDRIA MONOGYNIA.

* *Flores Apetali.*

Anthera lateri ovarii insidens (indivisa, bilocularis; vel triloba, 4-locularis). *Ovarium* monospermum, ovulo pendulo. *Stigma* capitatum. *Drupa*.—Brown.

Chloranthus monostachys, antheris incurvis trilobis indivisive, spicis solitariis, floribus alternis. Br.

By the advice of Mr. Brown we have placed *Chloranthus* in the first class of the Linnean system, although, he observes, that only one known species is really and at the same time constantly monandrous (*C. monander*, Br.); for in *inconspicuus* and *elatio*r (Br.) the three-lobed, four-celled anther, hitherto regarded as an antheriferous petal, is according to him composed of three confluent anthers, of which the middle one only is perfect or two-celled, corresponding with the anther of *monander*, the lateral being dimidiate or one-celled; and in *monostachys* both simple and compound anthers are found on the same spike.

Bot. Mag. 2190 (1820).

RULINGIA PANNOSA.

PENTANDRIA PENTAGYNIA.

(Inter Maherniam et Commersoniam.)

Petala 5, e cucullata basi ligulata. *Stamina* sterilia 5,

indivisa (*Nectarium*, Linn.). *Ovarium* 5-loculare: loculis dispermis. *Capsula*: septis duplicatis dermum 5-partibilis. *Brown*.

Rulingia pannosa, capsulis echinatis exsertis, foliis dentato-serratis acutis planis supra scabris subtis tomentosis: inferioribus ovatis subcordatis passimque lobatis; superioribus oblongo-lanceolatis. *Br. prodr. fl. nov.-holl.* 2 *inedit*.

Our friend Mr. Brown, to whom we are indebted for the above generic and specific characters, places this genus in the Natural system in his order of *Buttneriaceæ*, which he has defined in the appendix to 'Flinders's Voyage,' vol. ii, p. 540.¹

It is nearly related to *Commersonia*, from which it differs in the number of sterile filaments, or divisions of the nectarium, in the cells of the ovarium being two-seeded, and in the capsule. Named in memory of John Philip Rüling, author of an essay on the 'Natural Orders,' in which he has published the ideas of Professor Büttner upon this subject. A greenhouse shrub, native of Port Jackson, in New Holland, where Mr. Brown discovered several other species of the same genus.

Bot. Mag. 2191 (1820).

SYMPHORIA RACEMOSA.

PENTANDRIA MONOGYNIA.

Germen 4-loculare: loculis 2 polyspermis, sterilibus; 2 monospermis, fertilibus. *Bacca* coronata. *Cor.* tubulosa, brevis, 5-fida, subæqualis. *Cal.* 5-dentatus. *Brown*. *Lonicæræ* sp. *Linn.*

Symphoria racemosa, racemo interrupto subterminali, corolla intus barbata.

Germen ovate, with two or three small bracts applied close to the base. Mr. Brown has observed that it is four-

¹ [*Antè* vol. i, p. 11.]

celled, 2 of the cells bearing each several ovula, but which are always abortive; the other two having each a single seed, which alone come to maturity. An analogous circumstance takes place, he observes, in the fruit of *Linnaea*, and of *Abelia*, two nearly related genera belonging to the same family, the *Caprifoliaceæ*, in both which the germen is three-celled, two of the cells having several imperfect ovula, and the other a single ovulum, which alone comes to maturity. *Vide* 'Characters and Descriptions of Three New Species of Plants, found in China by Clarke Abel, Esq. By Robert Brown, F.R.S.' [*Ante*, p. 324.]

Bot. Mag. 2211 (1822).

Dillwynia cinerascens, corymbis terminalibus sessilibus, foliis filiformibus erectis; mucrone innocuo brevissimo sub-recurvo, ramulis calycibusque sericeis. *Brown MSS.*

Mr. Brown possesses many species of this genus, which he divides into two sections; in one of which (*Xeropetalum*, Br.) these parts are marcescent. It is to the latter section that the present plant belongs. The three species recorded in the 'Hortus Kewensis' all belong to the first; and from this division the above generic character seems to have been more especially framed.

Bot. Mag. 2247 (1822).

JACARANDA OVALIFOLIA.

Cal. 5-dentatus. *Cor.* fauce campanulata: limbo bilabiato. *Filamentum* quintum sterile, longius, barbatum. *Caps.* ovata, bilocularis, bivalvis: dissepimento contrario, valvis pleniuseculis adnato. *Semina* alata.—Arbores. *Folia opposita, abrupte rariusve impari-pinnata, pinnis impari-pinnatis.* Flores *paniculati, violacei.* Calyx sæpius *cyathiformis nunc breve campanulatus vel subcylindricus dentibus obsoletis.* *Corollæ æstivatio imbricata, labio superiore bre-*

viore bilobo equitante inferius, cujus lobus medius major indivisus, lateralibus tectus. Stamina antherifera inclusa: Antheræ sæpius dimidiatæ, cum rudimento nano lobi alterius; nunc completæ, oculis æqualibus divaricatis apice solum connexis. Brown MSS.

Jacaranda ovalifolia, antheris dimidiatis, corollis extus sericeis, foliolis pubescentibus; lateralibus ovalibus cum mucrone; terminali lanceolato. *Brown MSS.*

Jacaranda ovalifolia is very nearly related both to *J. acutifolia* and *J. obtusifolia* of Humboldt and Bonpland (*Plant. Æquinoc. tabs. 17 and 18*), between which it may be placed. *J. acutifolia* differs from it chiefly in all the leaflets being lanceolate, and in having a smaller number of pinnæ. *J. obtusifolia* is still more distinct in its leaflets entirely wanting the mucro, which is both obvious and constant in our plant, and in having a smooth corolla. *J. Bahamensis*, Nob. (*J. caroliniana*, Persoon; *Bignonia cærulea*, Linn.), of which there is in the Banksian Herbarium a single imperfect specimen that may be supposed to be authentic, and *J. rhombifolia*, of Meyer (*Flor. Essequib.*, 213), which is probably not different from the plant found by the late Dr. Anderson, of St. Vincent, on the banks of the Essequibo, and cultivated in some of the gardens, under his name of *Bignonia filicifolia*, are easily distinguished from the three species already mentioned, by their rhomboidal leaflets, and from each other by differences in the surface of corolla, which is silky in *J. Bahamensis* and smooth in *J. rhombifolia*.

J. procera, Nob. (*Bignonia Copaia*, Aublet, *B. procera*, Willd.), is sufficiently different from all the others in the much greater size of its leaflets, which are frequently upwards of an inch in length; in the rachis of the pinnæ not being winged; and in the cylindrical calyx, of which the teeth are extremely minute.

In five of the above-mentioned species, I have ascertained that the antheræ are dimidiate, with a hardly visible rudiment of a second lobe; a structure which M. Meyer (l. c.) has expressed by "Antheræ simplices," and intro-

duced into his character of the genus.' But in an undescribed species lately found in Brazil (*J. tomentosa*, Nob.) they are perfect, consisting of two equal divaricate lobes, as in the greater part of the natural order. This species differs, indeed, from the rest of the genus in its leaves, having constantly a terminal pinna. It agrees with them, however, in all the other characters of the flower, and entirely in the structure of its fruit; it can therefore be regarded as forming only a section of a strictly natural and well defined genus, still depending on characters not materially different from those proposed for it by M. de Jussieu; the only doubtful species being the *Jacaranda* 2, of Piso, from which the generic name was adopted. *Brown.*

Bot. Mag. 2327 (1822).

BRACHYSTELMA TUBEROSA.

Asclepiadea. *Corolla* campanulata; sinubus angulatis. *Columna* inclusa. *Corona* 1-phylla, 5-fida: lobis antheris oppositis, dorso simplicibus. *Antheræ* absque membranâ apiculari. *Massæ pollinis* erectæ, basi insertæ. *Brown MSS.*

Brachystelma tuberosa. *Brown MSS.*

Stapelia tuberosa, Meerb. ic. t. 54, f. 1, Monente D. Aiton.

Mr. Brown, in his valuable treatise on the "Natural Orders of *Asclepiadææ* and *Apocineæ*," published in the first volume of the 'Transactions of the Wernerian Society,' has divided the genus *Stapelia*, but our present plant will not unite with any of the genera there established; we are therefore highly indebted to the friendship of this learned botanist, for enabling us to establish the genus of this very rare plant.

Meerburg's figure, above quoted, which was pointed out to Mr. Brown by Mr. Aiton, represents the flower much

larger than in our specimen, with the segments elongated to a very narrow point.

Bot. Mag. 2343 (1822).

Arthropodium cirratum, racemo diviso, bracteis foliaceis, pedicellis fasciculatis, petalis interioribus integerrimis, filamentorum dimidio barbato basi biappendiculato, foliis lanceolato-ensiformibus. *Brown MSS.*

Anthericum cirratum, foliis lanceolatis planiusculis, scapo paniculato, filamentis bicirratiss. *Forster Prodr. Fl. Austr.*, p. 24. Willd. sp. pl. 2, p. 146. *Anthericum latifolium*, *Banks and Sol. MSS.*

Another species of *Arthropodium*, the *paniculatum*, will be found at No. 1421 of this work; from which our present plant differs in the much greater width of the leaves and revolute bractes; greater size of its white flowers; and both from that and every known species in the greater length of the spur-like processes at the base of the bearded part of the filaments, of which Mr. Brown could only trace mere rudiments in any of the other species; in the entire margins of the inner petals; and in the thicker root, approaching to tuberous.

Bot. Mag. 2350 (1822).

AMMOBIUM ALATUM.

SYNGENESIA POLYGAMIA ÆQUALIS.

Sect. DISCOIDEI.

Receptaculum paleis distinctis. *Pappus*, margo dentatus. *Antheræ* basi bisetæ. *Involucrum* imbricatum, coloratum, radians.—Herba erecta, tomentosa. Folia integerrima: radicalia lanceolata, basi attenuata, caulina minora, decurrentia. Caulis alatus, ramis unifloris. Involucra hemisphærica, laminis albis, intimis patulis radium abbreviatum

efformantibus. Corollulæ uniformes, flavæ. Stigmata truncato-dilatata. Semina ancipitia. Receptaculum convexum. Brown.

Ammobium alatum, Brown MSS.

An undescribed genus belonging to the same tribe with *Gnaphalium*, discovered by Robert Brown, Esq., in 1804, growing plentifully, near the shores of Port Hunter (or the Coal River), in New South Wales, and named by him from its growing in sand.

Bot. Mag. 2459 (1824).

ENTELÆA ARBORESCENS.

POLYANDRIA MONOGYNIA.

Cal. 4—5-phyllus. *Petala* 4. *Stamina* indefinita, uniformia, *antheris* subrotundis, incumbentibus. *Stigma* denticulatum. *Capsula* sphæroidea, echinata, sex locularis, semi-sexvalvis, polysperma.—Arbor (*parva, ligno lævissimo spongioso*, SOLANDER) *pube stellari* (*in pagina superiore folii adulti parciore et subsimplici*) tomentosa. *Folia* cordata, angulata, duplicato-crenata, 5-nervia, stipulis persistentibus, parvis, foliaceis. *Umbellæ simplices, terminales, pedunculatæ, involucri polyphylo, brevi, pedicellis ebracteatis.*

Flores albi, raro 5-fidi; calyce deciduo, foliolis aristatis, gemma (alabastro) quadricorni; petalis flaccidis, apice sublaceris, staminibus corolla brevioribus, subtorulosis, omnibus antheriferis; stylo angulato, stigmate umbilicato, margine denticulato. Capsularum setæ longæ, rigidae, opacæ, pubescentes, spinula pellucida acuta recta terminatæ. Semina in singulo loculo biseriata, parva, obovata; albumine carnosio, semini conformi; embryone recto, albo, longitudine fere albuminis; cotyledonibus planis, cordatis.

ORD. NAT. Inter *Tiliaceas*, proximum genus *Sparmanniæ*, quacum habitu, inflorescentia, antheris, stigmate, capsula-

rum figura, earumque setis spinula pellucida terminatis convenit; diversum filamentis omnibus antheriferis (unde nomen) vix manifeste torulosis, capsulis basi indivisis nec longitudinaliter dehiscentibus, loculis polyspermis, nec oligospermis (secundum Thunberg in *Sparmannia* dispermis, cujus tamen ovarii loculi certe polyspermi sunt) calycis foliolis aristatis nec muticis. *R. Brown MSS.*

Entelæa arborescens, R. Brown MSS.

Bot. Mag. 2480 (1824).

Stylidium adnatum, foliis cuneato linearibus, spica subcomposita; partialibus paucifloris, labello inappendiculato, capsulis angusto-linearibus: loculo postico angustissimo vacuo adhærente. *R. Brown MSS.*

Stylidium adnatum, Br. Prodr. Fl. Nov.-Holl., p. 572.

β *spica pedunculata, caulibus partialibus indivisis, Br.*

"It is possible that the plant here figured," Mr. Brown observes, "may belong to a species different from *S. adnatum*, whose spike is generally sessile, and more subdivided; I am inclined, however, to unite them, and to add, as a third variety, *S. propinquum*, which was taken up from imperfect materials. In the section of the genus to which *S. adnatum* belongs, the best specific differences are found in modifications of the structure of the capsule, and by these alone the species at present known may be readily determined. Thus, in *S. breviscapum* the cells and valves of the capsule are in form and contents perfectly equal, as in the other sections of the genus. In *fasciculatum* the cells are dissimilar in form, but of nearly equal size, and both fertile. In *falcatum* the posterior cell, still more different in form, is also considerably reduced in width and entirely without ovula; nor is it, as I have stated in the specific character, fertile. Lastly, in *adnatum*, of which I am now disposed to consider *propinquum* a variety, the posterior cell is not only entirely destitute of ovula, but is reduced to a capillary tube, which continues to adhere to the fertile cell. From this series of species, in which the upper or posterior cell is gradually obliterated, we are led

to the true explanation of the structure of *Lysipoma*. This genus, belonging to *Lobeliaceæ*, is described by M. Kunth as having a single parietal placenta; it may, however, be regarded as having a septum so closely pressed to the upper side of the capsule as to have no manifest cavity, and consequently produces ovula on the one side only." *Brown MSS.*

Bot. Mag. 2598 (1825).

Bæckia camphorata, Brown MSS.

Bot. Mag. 2694 (1826).

CONTRIBUTIONS
TO THE
BOTANICAL REGISTER.

1816—1826.

ce
ap
as

in

b
p

p
S
d
(
(
(
2
e

CONTRIBUTIONS
TO THE
BOTANICAL REGISTER.

CRYPTARRHENA. Petala 5, distincta, patentia. *Labellum* ecalcaratum: *lamina* dilatata explanata. *Columna* libera, aptera. *Anthera* apice cucullato columnæ inclusa, porrecto-ascendens. *Massæ pollinis* quatuor. *Brown MSS.*

Cryptarrhena lunata, Brown MSS.

Bot. Reg. 153 (1816).

VESTIA.

On Mr. Brown's suggestion, we have placed the genus in the Order *Solaneæ*. It comes very close to *Lycium*.

Bot. Reg. 299 (1818.)

Callistemon rigidum, foliis linearibus (lanceolato-linearibusve) planis acutissimis mucronatis lævibus, ovariis pubescentibus, capsulis distinctis. *Brown MSS.*

Bot. Reg. 393 (1819).

CALYTRIX. *Cal.* superus, tubo cylindraceo, limbo 5-partito, laciniis aristatis persistentibus. *Pet.* 5, decidua. *Stam.* omnia (sæpius indefinita rarò decem) antherifera, decidua. *Ovarium* uniloculare, dispermum. *Pericarpium* (*Achenopsis*) monospermum, indehiscens, exsuccum. Frutices (*Novæ Hollandiæ*) ericoidei. *Folia parva, sæpiùs angulata, sparsa, imbricata, glanduloso-punctata, in plerisque petiolata et stipulis! setacis liberis minutis deciduis instructa.* Flores

(*albi vir purpurei*) *axillares, solitarii, bibracteali, bracteis membranaceis, carinatis, persistentibus, basi connatis*. Brown MSS.

C. glabra, icosandra; foliis petiolatis stipulatis adultis bracteisque glabris. Brown MSS.

The generic character now offered is from the pen of Mr. Brown, and has been framed with the accustomed skill and foresight of that learned naturalist with a view to the cluster of confining genera from the same regions, which will be found near it in the concluding volume of his 'Prodromus of the Flora of New Holland.' Achenium implying an indelhiscent originally one-seeded seed-vessel, while that of the present genus, though eventually one-seeded, having been found by Mr. Brown to originate in a germen with two ovula, one of which proves constantly abortive, he has thought it advisable to suggest *Achenopsis* as a more precise denomination for this sort of seed-vessel.

Glabra is the first of the genus that has presented itself in our European gardens, and is native within the colony of Port Jackson, where it was found by Mr. Brown, whose herbarium contains likewise four more of its congeners, of which that gentleman has favoured us with the following account:—Three, our plant being one, were observed by him in the colony above mentioned, as well as in Van Diemen's Island, and agree in having a petioled stipuled foliage and icosandrous flowers; a fourth he discovered on the south-west coast of New Holland, this had likewise a petioled stipuled foliage, but the flowers were decandrous; the fifth he observed on the north coast of the same continent (in the Bay of Carpentaria), in that the flowers were icosandrous as in the three first-mentioned species, but it differed from all the others in having a foliage without either petioles or stipules. Through these modifications the genus will be found to unite with its confining co-ordinates at different points. Bot. Reg. 393 (1819).

Melaleuca incana, foliis ternis lineari-lanceolatis utrinque ramulisque incano-pubescentibus, spicis ovalibus oblongisve. Brown MSS. Bot. Reg. 410 (1819).

ARTABOTRYS. *Cal. tripartitus. Petala sex. Stamina hypogyna. Ovaria distincta, disperma. Baccæ dispermæ (abortione quandoque monospermæ). Semina collateralia erecta exarillata albumine rimoso.*—Frutices decumbentes. *Folia alterna, integerrima, exstipulata. Pedunculi extra- alares, suboppositifolii pauciflori, pedicellis (1—2) laterali- bus, apice uncinati.* Brown MSS.

Obs. *Characteribus suprâ datis proximè accedit Kadsura, cujus baccæ dispermæ seminibus collateralibus pariter distinctæ receptaculo carnosio insidentes fide iconis et descriptionis Kämpferi (Amæn. Exot. 476); sed Kadsura viz Annonacea ob summam affinitatem cum Uvariâ heteroclitâ, Roxb. Flor. Ind. Ined. cui albumen indivisum nec processibus membranæ interioris rimosum auctoritate descriptionis operis citati.* Brown MSS.

A. odoratissimus, petalorum laminis planis lanceolatis, foliis oblongis acuminatis. Brown MSS.

Uvaria odoratissima. Roxburgh, Flor. Ind. Ined.

Unona uncinata, Dunal Anonacées, 105, t. 12 et 12a, DeCand. Syst. Nat. 1, 490.

Unona hamata. Dunal Anonacées, 107. DeCand. Syst. Nat. 1, 491.

Uvaria esculenta. Rottler in Nov. Act. Soc. Nat. Cur. berol. 4, 201.

Uvaria uncata. Loureiro cochîn. 349.

Anona uncinata. Lamarck. encyc. 2, 127.

Anona hexapetala. Linn. Suppl. 270. Hort. Kew. 2, 253. Ed. 2, 3, 335. Willd. sp. pl. 2, 1266.

The present plant, after various shiftings from one inappropriate group to another, according to the wavering views of different botanists, is now placed in a new genus constituted for its reception by Mr. Brown, to whom the generic name has been suggested by the curious grapple or tendril belonging to the peduncle, by which the growing fruit is conveniently suspended on the nearest support, during its advance to maturity, and the slender flexile branch relieved from the disproportionate burden, which would be otherwise laid on the ground.

The synonymy is also a valuable contribution from

Mr. Brown, and presents a critical view of the scientific history of the species.

We understand that three species of *ARTABOTRYS*, besides the present, are already known, one of which with curiously small flowers has been recently discovered by Dr. Horsfield during his residence in Java, and is in the rich Herbarium that gentleman has brought to this country.

Bot. Reg. 423 (1820).

MODECCA. Flores diclines (dioici v. monoici). *Cal.* 5-fidus. *Petala* 5, calyci inserta. *Squamæ* (*Nectarium*, Linn.), numero definitæ (5—10) raro nullæ. *MAS.* *Stamina* 5; *antheræ* stantes. *FEM.* *Caps.* (pedicellata) unilocularis 3-valvis polysperma. *Herbæ* (*Indiæ Orientalis*, *Novæ Hollandiæ*, et *Africa Equinoctialis*) scandentes cirrhis axillaribus simplicibus vel e divisuris pedunculorum. Folia lobata v. indivisa basi et subtus sæpius glandulosa. Pedunculi axillares apice divisi dichotomiis cirrhiferis. Brown MSS.

Bot. Reg. 433 (1820).

CRYPTOSTEGIA. Cor. infundibuliformis: *tubus* squamis 5, inclusis (bipartitis *subulatis*) laciniis limbi alternantibus. *Stamina* inclusa, imo tubo inserta; *filamentis* distinctis; *antheris* basi stigmate cohærentibus. *Glandulæ* 5, spathulatae, angulorum stigmatis pollen granulosum (simplex) colligentes. *Ovaria* 2. *Styli* 2. *Stigma* (commune) pentagonum. *Folliculi* *Semina* comosa.—Frutex volubilis, glaber. Folia opposita. Pedunculi terminales, 3-fidi. Flores speciosi. Corolla æstivatione contorta. Folliculi unguati, divaricatissimi. Brown MSS.

Cryptostegia grandiflora. Brown MSS.

CRYPTOSTEGIA is founded upon a single species, native of the Peninsula of India, where it was first observed by Dr. Roxburgh, and strangely mismatched by him with the species of the genus *Nerium*, belonging to the *Apocynæ*; while our plant belongs to the third section of Mr. Brown's *Asclepiadeæ*, distinguished from the other sections by granular solitary pollen-masses and filaments which are

distinct in whole or in part. *Cryptostegia* is, however, conceived by Mr. Brown to be the link by which the two orders connect through his genus *Cryptolepis*, also a native of India and a climber.

The name was suggested to Mr. Brown by the circumstance of the enclosure of the five-scaled crown within the tube of the corolla, and its not being exposed to view, as in other bordering genera.

Bot. Reg. 435 (1820).

BURCHELLIA. *Capitulum* involocratum. *Cor.* clavato-infundibuliformis: *limbo* 5-fido abbreviato fauceque imberbi; æstivatione mutuò imbricatâ contortâ. *Stamina* supra medium tubi inserta; *antheris* subsessilibus inclusis. *Stigma* clavatum. *Bacca* calyce altè 5-fido coronata, bilocularis, polysperma. *Brown MSS.*

Frutex ramosissimus pubescens, ramulis compressis. Folia opposita. Stipulæ interpetiolares, e dilatatâ basi subulatæ, indivisæ, caducæ. Capitulum terminale, extra involucreum monophyllum pluridentatum abbreviatum, pari unico foliorum minorum stipulis proportionatim latioribus subtensum. Ovaria supra receptaculum convexum villosum bracteolisque nonnullis minutissimis conspersum sessilia, distincta. Calyx: limbo foliaceo æquali, tubum aliquoties superante. Corolla coccinea, extûs pilis appressis, intûs glabra præter barbam annularem juxta basin tubi. Antheræ lineares. Discus epigynus carnosus, indivisus, imberbis. Stigma exsertum, utrinque sulco exaratum. Bacca turbinato-globosa, bilocularis septo completo. Placenta adnata. Semina angulata. Embryo axilis, dimidio albuminis cartilaginei longior. Brown MSS.

Burchellia capensis. Brown MSS.

By its original observer the species had been ranked under the head of *Lonicera*, but was afterwards referred to its true station in *Rubiaceæ*; though that learned botanist, in adopting an erroneous description, representing the seeds of the berry as solitary, has placed the plant in a wrong section of the order. Mr. Brown having proved it not to be consistently reducible to any established genus, has

formed a new one from it, and with that liberality and vigilance which he extends to every interest of science, availed himself of the appropriate occasion of honouring the merits of Mr. Burchell, the zealous and enterprising investigator of the regions to which our plant belongs.

Bot. Reg. 466 (1820).

CALOTIS. *Recept. epaleaceum. Achenia coronata paleis duabus oppositis aristisque 1-3 glochidatis. Involucrum subæquale, simplici vel duplici serie polyphyllum. Brown MSS.*

Herbæ ramosæ, pilosiusculæ. Folia alterne. Involucra ramos terminantia, solitaria, planiuscula, foliacea. Ligulæ fæminæ numerosæ, imbricatæ, cæruleæ, post anthesin spiritaliter revolutæ, et diu persistentes. Flosculi masculi quinquefidi, flavi, antheris basi muticis. Achænia (semina, Linn.) verticaliter compressa cuneata. Pappus persistens, e paleis lateralibus, dilatatis latioribus quam longis, auriculiformibus; et aristis longioribus sæpius duabus (anticâ et posticâ), extra medium vel apice solùm aculeolis reversis. Receptaculum scrobiculatum vel subalveolatum convexum. Brown MSS.

Calotis cuneifolia, foliis cuneatis apice inciso-dentatis. Brown MSS.

The character of *Calotis* was formed, but not published about fifteen years ago, from *C. dentæ*, a species first observed by Mr. Brown himself in New Holland, where it is not uncommon in the neighbourhood of Port Jackson. The present has been since found, during an expedition into the interior of the above country, growing on the banks of the River Lachlan in 1817, by Mr. Allan Cunningham, who is commended by Mr. Brown as "an indefatigable collector and acute observer."

Bot. Reg. 504 (1820).

VANDA. *Labellum calcaratum, cum basi simplici (breviusve productâ) columnæ apteræ continuum, trifidum, lobo medio carnoso. Petala patentia distincta. Massæ pollinis 2, obliquè bilobæ. Brown MSS.*

OBS. *Aerides paniculatum* (*Bot. Reg. v. 3, f. 220*) generis *pristini species impar nimis huic* (monente *D. Brown*) *aptè satis associari potest. Parasiticæ ambæ.*

Vanda Roxburghii, ovariis contortis, petalis oblongo-ovatis undatis, foliis obliquè tridentatis. *Brown MSS.*

The generic character now given has been formed by Mr. Brown so as to include *AERIDES paniculatum*, published provisionally by that generic name in our third volume (fol. 220); but now it will be seen that the species is not a very close congener of the one before us. The *CYMBIDEUM tessellatum* of Roxburgh is not included in the present genus.

Mr. Brown had some scruple in not ranging the group under *ANGRÆCUM* of M. Du Petit-Thouars (of which a species is given in 'Voy. de Bory de St. Vincent,' 1359, tab. 19), not being able, from want of sufficient detail in the description of the structure of the parts of that genus, to decide satisfactorily whether he ought to do so or not; and has at last been determined principally by the *label* in *ANGRÆCUM* being undivided, and probably membranous, while in *VANDA* it is three-lobed and fleshy. He expects that several species of Swartz's genus *LIMODORUM* will be found to rank under the present.

Bot. Reg. 506 (1820).

GRIFFINIA parviflora.

Semina secundum observationes Dom. Brown, obovata, ventricosa, nitentia, ochroleuca; umbilicus basilaris; apex chalazæ fuscæ insignitus; raphe tenuis, immersa; integumentum duplex, exterius membranaceum, nitens, per lentem pulchrè areolatum, ventre longitudinaliter incrassatum, præsertim in regione umbilici; membrana propria tenuissima, albumini adhærens, separabilis tamen, evasculosa, basi chalazæ inserta cæterum libera; albumen semini conforme, densè carnosum; embryo axilis, longitudine dimidii albuminis, cylindræus; radícula umbilico approximata, apice exserto. Brown MSS.

We are to thank Mr. Brown for the description of the seed of this genus, the nature of which had been misunderstood until the present subject was submitted to the test of his accurate and skillful investigation.

Bot. Reg. 511 (1821).

Hedychium elatum, foliis oblongo-lanceolatis glabris, spicis laxis; fasciculis ternis subtrifloris, laciniis interioribus limbi cuneato-linearibus, labelli laminâ bifidâ. *Brown MSS.*

Caulis 8—10-pedalis. Folia inferiora oblongo-lanceolata, omnia acumine subulato-filiformi citius marcescenti. Spica 6-uncialis, fasciculis distinctis sæpissimè ternis, patentibus, trifloris. Ovarium (germen) villosum. Perianthium exterius (calyx) spathaceum, scariosum, pubescens: interius (corolla) tubo cylindræo ochroleuco, limbo exteriori tripartito laciniis angusto-linearibus acutis, interiore laciniis lateralibus cuneato-linearibus, apice parùm dilatato obliquo: labellum ungue brevi lato-lineari: laminâ circumscriptione ovatâ, lobis dimidiato-ovatis obtusiusculis. Filamentum labello longius. Antheræ linearis basi bifidâ lobis poliniferis. Stigma cyathiforme indivisum ciliatum. Nectaria (stamina sterilia) 2, brevissima, teretiuscula, approximata. *Brown MSS.*

An unrecorded species, determined and described by Mr. Brown from native samples in the Banksian Museum, of which that gentleman is now, fortunately for science, in the well-earned possession. The samples formed part of the collection of Nepaulese plants transmitted by Dr. Wallich, the able superintendent of the Calcutta Botanic Garden, by whom, as was plain from the inscribed notice, they had been taken for *speciosum*, accurately described by himself in the 'Flora Indica' of Roxburgh; but on examination by Mr. Brown, proved to be essentially distinct from that species.

Bot. Reg. 526 (1821).

Anthemis apiifolia, foliis glaberrimis pinnatifidis: lobis cuneatis trifidis incisive, floribus solitariis, involucri foliolis linearibus apice scariosis. *Brown MSS.*

Involucrum (calyx) hæmisphæricum, foliolis subæquali-

bus, appressis, linearibus, apice brevi scarioso vix dilatato. Pappus: margo membranaceus brevissimus indivisus. Receptaculum convexiusculum, paleaceum, paleis linearibus acutis. Brown MSS.

With our present imperfect knowledge of this plant, it is perhaps necessary to refer it to *ANTHEMIS*, though in real affinity it will probably be found to approach more nearly to *CHRYSANTHEMUM Indicum*, of which the supposed double-flowered varieties are in like manner generally furnished with *paleæ*. As that plant, however, in its single state has always a naked receptacle, and some of its double-flowered varieties are constantly, while others are occasionally without *paleæ*, in the second edition of the 'Hortus Kewensis,' it was continued in *CHRYSANTHEMUM*; an arrangement which it would not perhaps be necessary to alter, were it even proved that two species have been confounded under the name of *CHRYSANTHEMUM Indicum*, both of them belonging decidedly to the same natural genus, and to this genus I am inclined to think *ANTHEMIS apifolia*, when completely known, may be referred. As to the narrow *margo* crowning the seed, on account of which our plant has been considered a *PYRETHRUM*, it does not appear to me to constitute a sufficient generic character, and it equally exists in *CHRYSANTHEMUM Indicum*, particularly in the single-flowered specimen of the Linnean Herbarium. *Brown MSS.*

For the above account we are indebted to the unwearied liberality of Mr. Brown, who has taken this opportunity to account for his retaining *CHRYSANTHEMUM Indicum* in that genus in the second edition of the 'Hortus Kewensis,' and for not following some of his predecessors in referring the species to *ANTHEMIS*.

Bot. Reg. 527 (1821).

ASTELMA. Receptaculum nudum (nec paleatum nec favosum). Pappus plumosus, sessilis; radiis basi connatis. Involucrum (calyx) imbricatum, squamis scariosis: intimis conniventibus. Brown MSS.

Astelma eximium, Brown MSS.

ASTELMA is now first detached from GNAPHALIUM by the above able hand; and founded upon the present species characterised as a genus by a naked *receptacle of the flower*. (neither chaffy nor honeycombed); a sessile feathered *seed-crown*, the radii of which are connected at the base; an imbricated *calyx* composed of scariose *leaflets*, the innermost of which are convergent, and whose florets are all both stamen-bearing and pistil-bearing. It is the nearest genus to ELICHRYSUM, from the genuine species of which, according to Mr. Brown, it only differs by the rayless calyx.

Bot. Reg. 532 (1821).

Dendrobium cucullatum, caulibus pendulis, foliis bifariis lanceolatis acuminatis, pedunculis oppositifoliis subbifloris, labello indiviso circumscriptione ovato basi cucullato. *Brown MSS.*

It is said to be in several collections, and to have been generally conceived to be *DENDROBIUM Pierardi* of the unedited part of Roxburgh's 'Flora Indica'; and which, according to Mr. Brown, to whom, as in so many instances, we are indebted for all we have to say on the subject, it is possible it may really be; but finding by both figure and description the flowering stems of that species represented as leafless, the lamina of the labellum of an obovate form, and leaves as emarginate, while in the present plant the flowering stems are in full leaf, its labellum of an ovate outline, and the leaves taper-pointed, he has deemed it safer to regard the present sample as belonging to a distinct species.

Bot. Reg. 548 (1821).

LISSOCHILUS. *Massæ pollinis* 2, obliquè bilobæ. *Label-lum* basi saccatum: laminâ sessili indivisâ convexâ, posticè excisâ et cum basi columnæ apteræ edentulæ connatâ. *Brown MSS.*

L. speciosus, petalis interioribus divaricatis duplò majoribus exterioribus reflexis, labello altè cordato. *Brown MSS.*

Herba 4-pedalis et ultrâ. Folia radicalia ensiformia

acutissima crassiuscula rigidula viridia nec glauca, marginibus carinâque lævibus: basibus dilatatis incrassatis in bulbum emersum connatis, scapo aliquoties breviora. Scapus vaginatus teres viridis solidus, fasciculis vasorum sparsis. Vaginæ distantes strictæ foliaceæ, in foliola lanceolata, appressa, acuta longitudine vix ipsius vaginæ productæ. Spica racemosa simplicissima erecta laxa sesquipedalis. Bracteæ foliaceæ lanceolatae, patulae sæpiùsve divaricatae, immersè nervosæ, marginibus lævibus, infimæ in scapum paulò decurrentes ovarium cum pedicello suo superantes, reliquæ sessiles, superiores ovario breviores. Flores speciosi, e torsione pedicelli baseosque ovarii verè resupinati, labello antico. Perianthii foliola 3 exteriora utrinque viridia calycina, distincta arcuè reflexa ovata acutiuscula immersè nervosa, postico plano, lateralia paulò latiora lavissimè concava; interiorum lateralia petaloidea, flava, immaculata, divaricatissima, oblongo-ovata, obtusiuscula, limbo parum reflexo, disco hinc convexiusculo; nervosa nervis lateralibus alternatim simplicibus bifidisque, ipsa basi alba. Labellum lateralibus interiorum vix brevius, basi subtus calcarato-saccatum cornu brevissimo compresso albo: lamina sessilis, circumscriptione ovato-oblonga, flava basi albicante venis purpureis, apice subretuso, basi altè cordata lobis posticis rotundatis dimidio inferiore liberis superiore cum basi columnæ cohærentibus suturis manifestis, lateribus subparallelè-reflexis plicatis plicis paulò tantum elevatis obtusis sulcis interstitialibus angustis: disco convexo axi gibbosiusculo. Columna PROPORTIONATA semiteres e basi albâ pallidè virens, aptera apice edentulo posticè rotundato. Anthera terminalis mobilis decidua, in expanso flore apicem anticè declivem columnæ operiens, didyma, dorso incrassato dilutè viridi apice retuso, bilocularis loculis approximatis septulo partiali incompleto contrario in latere exteriori instructis. Massæ pollinis in singulo loculo singulæ, in gemmâ floralis juniori minutissimè granulosa, in expanso flore cereaceæ ovata, post imprægnationem in apice obliquo columnæ (clinandrio) inversæ, posticè (respectu antheræ anticè) instructæ lobulo minore operiente majoris cavitatem illitam materia viscidâ elasticâ flavescenti processus cujus

massæ insertæ lingulæ communi brevi lato-lineari membranaceæ albæ e glandulâ subrotundâ apicis stigmati ortæ. Stigma anticum cavum subrotundum viride secernens. Brown MSS.

The plant here described is the only certain species of *Lissochilus*; *Cymbidium giganteum*, however, according to Thunberg's description, may also belong to the same genus. This genus, in affinity, comes nearest to those plants of South Africa at present referred to *Limodorum* and *Cymbidium*, namely, *Limodorum barbatum*, *triste*, *longicorne*, and perhaps also *hians* of Thunberg (not, however, *Satyrium hians*, Linn, which seems to be a *Disa*), *Cymbidium pedicellatum* and *aculeatum*. These, along with several Indian species, also referred by Swartz and Willdenow to *Limodorum*, especially *L. virens*, *carinatum* and perhaps *epidendroides*, form a genus (EULOPHUS) related on the one hand to *Dipodium* (vide *Prodr. Flor. Nov. Holl.*, p. 300), and essentially agreeing with *Lissochilus* in the structure of Anthera, but sufficiently different from it in its trilobed crested labellum, which is neither cordate nor connected at base with the column. In both these characters *Lissochilus* equally differs from *Angræcum* of M. Du Petit-Thouars, which has also an elongated spur, and a considerably different habit.

Among the plants at present referred to *Limodorum*, another very distinct genus may be noticed, consisting of *Limodorum veratrifolium*, and judging from Kæmpfer's figure, *L. striatum* also. This genus (CALANTHE) agrees with *Bletia* in having eight pollen-masses, but differs from it in the claw of the labellum being connected with the column. It is not unlikely that this may be the *Cyanorchis* of M. Du Petit-Thouars, who will probably hereafter publish it under a different name. Brown MSS.

Bot. Reg. 573 [578 by mistake in orig.] (1821).

OPHIOPOGON *spicatus*.

Liriope spicata, *Loureiro coch.* 200 (*monente D. Brown ab autopsyiã edocto exemplaris archetypi in Herb. Parisiano asservati*).

In *O. spicato* (*monente D. Brown*) semen (sæpè 2—3), *nudum baccatum de ovulo mox post factum conceptum auctu præcociori cætera illidente exque germinis vertice erumpente excretum: basi cum exuto germine cohærens*.

In *O. spicatus*, Mr. Brown has found the fruit to be of the same kind he has ascribed to *PELIOSANTHES Teta*, in the twelfth volume of the 'Linnean Transactions' [vol. 1, p. 362].

Bot. Reg. 593 (1822).

MACRADENIA. *Labellum* sessile, cucullato-concavum indivisum acuminatum. *Corollæ* petala distincta patula. *Columna* libera, lobis apicis conniventibus. *Massæ pollinis* 2, exsulcæ. *Brown MSS.*

Macradenia lutescens, *Brown MSS.* *Cor. sordidè lutea, intus purpureo-fusco marmorata. Labellum corollæ brevius, ecalcaratum, album, intus purpureo interruptè lineatum, obovatum, appendice lineari acutâ reflexâ. Columna longitudine ferè labelli, semiteres, suprâ sensim latior lobis lateralibus apicis in cucullum antheram amplexantem conniventibus. Anthera e basi paulò gibbosiore intusque biloculare producta in acumen lato-lineare concavum ipsâ basi aliquoties longius processum subulatum stigmatis, glandulam, ejusque appendicem, involvens. Massæ pollinis cuneiformes, modicè compressæ, post migrationem apicibus suis acutis affixæ disco apicis dilatati appendicis elongatæ proscollæ. Brown MSS.*

An undescribed species, now first introduced by Mr. Griffin, who received the plant, represented in the annexed plate, from Trinidad. It flowered last December in the hothouse at South Lambeth, and was communicated to Mr. Brown, who has made it the foundation of the present

genus, to be placed next to *ONCIDIUM*, from which it differs by a convolutely concave undivided taper-pointed *label*, a *corolla* with the *petals* all distinct, a *column* with the two top-lobes converging convolutely (not extended), and by two furrowless or unindented *pollen-masses* (furrowed or indented in *ONCIDIUM*).

Bot. Reg. 612 (1822).

CEROPEGIA. *Asclepiadea*. *Cor.* tubulosa basi ventricosa, limbi laciniis ligularibus conniventibus: æstivatione valvata. *Columna fructificationis* inclusa. *Corona staminea* 10-fida; laciniæ 5 antheris distinctis oppositæ longiores cum 5 reliquis (indivisis bifidisve) alternantes. *Massæ pollinis* erectæ. *Herbæ glabræ, sæpius volubiles, radice tuberosâ*. *Umbellæ interpetiolares, nunc paucifloræ*. Brown MSS.

C. africana, pedunculis 2—3-floris, foliis ovato-lanceolatis carnosis, limbo corollæ barbato, coronæ laciniis longioribus aversis falcato-recurvis: brevioribus indivisis. Brown MSS.

Radix: tuber pro ratione herbæ 3—6-uncialis magnum. *Caulis simplex, sæpius volubilis*. *Folia acuta, planiuscula, avenia, brevè petiolata, divaricata, apicibus subrecurvis, semuncialia*. *Spica rara e pedunculis alternis brevibus 2—3-floris, pedicellis brevibus minutè bracteolatis*. *Flores erecti, inferiore præcociore*. *Cal. 5-partitus, laciniis lineari-subulatis modicè patentibus, basi ventricosâ tubi corollæ ferè ter brevioribus*. *Corolla: tubus e basi obovato-globosâ obsolete pentagonâ viridi cylindraceus, fauce infundibuliformi striatâ; limbi laciniæ lineares intus marginibusque pilis coloratis patulis acutis barbatae, erectæ apicibus conniventibus læviterque cohærentibus: limbus inexpansus prismatico-pentagonus fauce pentagonâ angulis extantibus cum laciniis limbi alternantibus*. *Columna fructificationis basi ventricosâ tubi inclusa*. *Corona apici tubi staminei brevis inserta monophylla decemloba: lobi quinque cum staminibus alternantes abbreviati indivisi obtusi basi saccati: quinque reliqui quasi interiores antheris oppositi aliquoties longiores margine interiori truncato-canaliculato exteriori in aciem*

attenuato. Antherae distinctae ovatae loculis lateralibus longitudinaliter dehiscentibus appendice apicis carnosâ brevi acutâ stigmatibus incumbente. Massae pollinis minutae supra basin affixae subglobosae. Stigma muticum. Brown MSS.

A very curious unrecorded species, now first introduced from the Cape of Good Hope. The masterly description is by Mr. Brown, who has so far modified the character of the genus with respect to that defined by himself in his illustration of the Natural Order, in the 'Memoirs of the Wernerian Natural History Society,' as to fit it for the reception of the species before us; which he considers too naturally allied to the East Indian group he had originally allotted to CEROPEGIA to be separated.

Bot. Reg. 626 (1822).

SCHIZOPETALON. *Calyx* clausus. *Petala* pinnatifida! *Siliqua* torosa, seminibus uniseriatis. *Cotyledones* quatuor! separatim contortuplicatae! *Herba* annua alternifolia, *pubescenti-ramosa*. *Folia* sinuato-pinnatifida. *Racemus* foliaceo-bracteatus.

Calyx pubescens, foliolis albo-marginatis, lateralibus altius descendentibus. *Petala* alba, unguibus calycem paulò superantibus, laminis circumscriptione ovatis, pinnatifidis, laciniis linearibus siccitate (et forsân aestivatione) involutis. *Stamina* 6, filamentis subæqualibus, edentulis, antheris unifornibus, sagittato-linearibus, introrsis. *Glandulae* hypogynae quatuor lineares, erectae, petalis suboppositae, geminatim basibus dilatatis confluentibus filamenta lateralia stipantes. *Stylus* brevissimus. *Stigmata* papulosa, connato-approximata, in stylum decurrentia, basibus solutis, unicum quasi extensoriiforme efformantibus. *Siliqua* sessilis, bilocularis, angusto-linearis, pubescenti-ramosa brevi conspersa, valvis venosis. *Semina* sphaerico-lenticularia. *Embryo* viridis *Radicula* albicans, arcuata, semine paulò longior. *Cotyledones* verticillatae, aequales, elongatae, angustatae, semiteretes, separatim subspiraliter involuta.

In ordine *Cruciferarum* genus nulli cognito affine, et

¹ [See ante, p. 205.]

equidem ob numerum, figuram et vernationem cotyledonum, petala pinnatifida, stigmatis structuram et stamina subæqualia tribus distinctæ initium efformare videtur. *Brown MSS.*

For the elaborate character which we have given above, we are entirely indebted to Mr. Brown, who alone, in this country, was capable of defining the limits, and of fully appreciating the peculiarities of this truly singular plant.

Bot. Reg. 752 (1823).

CARMICHAELIA. *Calyx* cyathiformis 5-dentatus. *Ovarium* polyspermum. *Legumen* oligospermum (1—3-sp.), *replum* post lapsum valvularum persistente!

Frutex ramosissimus, sub statu florescentiæ sæpius aphyllus. Caulis ramique primarii teretes, teneriores v. plano-compressi, v. ancipites, stipulis minutis alternatim dentati. *Folia e dentibus ramorum fruticis junioris, ternata v. pinnata (foliolis 3—7 obcordatis). Racemi simplicissimi e denticulis ramorum: pedicelli basi bractea abbreviata, apice binis minutissimis. Flores parvi. Calycis dentes subæquales, brevissimi. Petala longitudine subæqualia. Vexillum lamina latiore quam longiore, basi absque callis auriculisve. Carina obtusa. Filamenta 1—9-fid. Antheræ uniformes, subovales. Ovarium lineare, 5—6-spermum. Stylus subulatus, adscendens. Stigma obtusum, imberbe. Semina subreniformia, sinu clauso umbilico nudo. Brown MSS.*

C. australis, Brown MSS.

Lotus arboreus, Forst. Prodr., No. 2, 278.

For the above character of this very remarkable genus we are indebted to the kindness of Mr. Brown, who, fortunately for science, is in possession of perfectly ripe fruit. The term *replum*, which is employed in Mr. Brown's generic character to designate the persistent circumscription of the legume, was used by Vitruvius for the frame of a door, and we believe has been already applied to the purposes of botanical description in the sense in which it is used here, which is certainly unexceptionable.

The garden plant has been compared by Mr. Brown with the specimens from Cook's 'Voyage,' in his own collec-

tion, and with an authentic specimen from George Forster's 'Herbarium,' as well as by ourselves, with others from some part of New Zealand; and no difference has been detected between them.

In explanation of the name applied to this plant, we make the following extract from Mr. Brown's communication upon the subject:

"I have named the genus in honour of my friend, Captain Dugald Carmichael, F.L.S., a very accurate botanist, whose interesting account of the Island of Tristan d'Acunha is published in the 12th volume of the 'Linnean Society's Transactions,' and to whom I am indebted for extensive collections, and many excellent descriptions of the plants of the Island of Mauritius and of the Cape of Good Hope."

Bot. Reg. 912 (1825).

Chorizema Henchmanni, foliis acicularibus pungentibus solitariis v. ternatim fasciculatis, calycibus villosis.

C. Henchmanni, *R. Brown, ined.*

This plant, which recedes very much in habit from the species of *Chorizema*, previously published, has been named by Mr. Brown in honour of Francis Henchman, Esq., a very successful importer of New Holland plants, for whom the present subject was collected by Mr. William Baxter.

Bot. Reg. 986 (1826).

NAMES, &c., OF PLANTS

CONTRIBUTED TO

VARIOUS PUBLICATIONS BETWEEN
1818 AND 1826;

AND

NAMES QUOTED

FROM

R. BROWN'S "HERBARIUM"

IN

MR. BENTHAM'S 'FLORA AUSTRALIENSIS,'
Vols. I—III, 1863—6.

NAMES, &c., OF PLANTS

CONTRIBUTED TO

VARIOUS PUBLICATIONS BETWEEN 1818 AND 1826.

I. Names and Descriptions of Plants contributed to De Candolle's 'Systema Naturale Regni Vegetabilis,' Paris, 1818—1821.

RANUNCULACEÆ.

- Clematis stenopetala*, R. Brown, *vol.* 1, *p.* 147.
— *aristata*, R. Brown, *ibid.*
Ranunculus inundatus, R. Brown, 1, *p.* 269.
— *collinus*, R. Brown, 1, *p.* 271.
— *Pumilio*, R. Brown, *ibid.*
— *plebeius*, R. Brown, 1, *p.* 288.
— *sessiliflorus*, R. Brown, 1, *p.* 302.

DILLENIACEÆ.

- Pachynema*, R. Brown, 1, *p.* 411.
— *complanatum*, R. Brown, 1, *pp.* 411, 412.
Hemistemma dealbatum, R. Brown, 1, *p.* 413.
— *Banksii*, R. Brown, 1, *p.* 414.
— *angustifolium*, R. Brown, *ibid.*
Pleurandra bracteata, R. Brown, 1, *p.* 415.
— *nitida*, R. Brown, 1, *p.* 415.
— *sericea*, R. Brown, *ibid.*
— *cinerea*, R. Brown, 1, *p.* 417.
— *furfuracea*, R. Brown, *ibid.*
— *parviflora*, R. Brown, 1, *p.* 418.

- Pleurandra scabra*, R. Brown, 1, p. 418.
 — *riparia*, R. Brown, 1, p. 419.
 — *pedunculata*, R. Brown, *ibid.*
 — *acerosa*, R. Brown, 1, p. 422.
 — *stricta*, R. Brown, *ibid.*
Candollea pedunculata, R. Brown, 1, p. 424.
 — *fasciculata*, R. Brown, *ibid.*
Hibbertia dentata, R. Brown, 1, p. 426.
 — *saligna*, R. Brown, 1, p. 427.
 — *virgata*, R. Brown, 1, p. 428.
 — *fasciculata*, R. Brown, *ibid.*
 — *linearis*, R. Brown, *ibid.*
 — *diffusa*, R. Brown, 1, p. 429.
 — *monogyna*, R. Brown, *ibid.*
 — *pedunculata*, R. Brown, 1, p. 430.
 — *serpyllifolia*, R. Brown, *ibid.*
 — *oblongata*, R. Brown, 1, 431.
 — *cistifolia*, R. Brown, *ibid.*
 — *tomentosa*, R. Brown, 1, p. 432.
 — *lepidota*, R. Brown, *ibid.*
Wormia alata, R. Brown, 1, p. 434.

MAGNOLIACEÆ.

- TASMANNIA, R. Brown, 1, p. 445.
Tasmannia aromatica, R. Brown, *ibid.*
 — *insipida*, R. Brown, *ibid.*

Hujus generis character sic a cl. Brown observatus fuit, nempe:

Flores diici vel polygami. Calyx 2-sepalus. Petala 2—5. ♂ Stamina ∞ , cum vel absque rudimento pistilli. ♀ vel ♂ Ovarium 1, 1-loculare; stigma latere interiore ovarii longitudinaliter adnatum! Bacca polysperma. (Brown, in litt.)

Obs. Cl. Brown autumat Tasmanniam cum Illicio et Wintera seu Drymide constituere Ordinem proprium, WINTEREÆ nomine ab eo insignitum.—*T. insipidam* nunc potius *dipetalam* vocat quia cortex ejus non est omnino aromate destitutus. Vol. 1, pp. 547, 548.

MAGNOLIA. Sect. II. GWILLIMIA.

Cl. Brown me monet in litteris specimina fructifera Magnoliarum Asiaticarum in Musæo Banksiano servata non essentialiter esse diversa a speciebus Americanis. *Vol. 1, p. 548.*

CRUCIFERÆ.

Draba Pumilio, Brown, ii, *p. 353.*

CHORISPORA, R. Brown.

Chorispermi nomen ob nimiam cum Corispermo similitudinem ipse mutavit Cl. Brown, ut me monuit in colloquio, ii, *p. 435.*

STENOPETALUM, R. Brown, ii, *p. 513.*

Stenopetalum lineare, R. Brown, *ibid.*

II. Names of Plants contributed to Decandolle's 'Prodromus Systematis Naturalis Regni Vegetabilis,' vol. i, Parisiis, 1824; vol. iii, 1828.

CAPPARIDÆ.

Capparis lasiantha, Brown, i, *p. 247.*

— *umbellata*, Brown, *ibid.*

VIOLARIÆ.

Hymenanthera angustifolia, Brown, i, *p. 315.*

— *dentata*, Brown, *ibid.*

DROSERACEÆ.

Drosera petiolaris, Brown, i, *p. 318.*

— *Banksii*, Brown, i, *p. 319.*

— *Menziesii*, Brown, *ibid.*

TREMANDREÆ.

TREMANDRA, Brown, i, *p. 344.*

— *stelligera*, Brown, *ibid.*

— *diffusa*, Brown, *ibid.*

PASSIFLOREÆ.

Modecca australis, Brown, iii, *p. 337.*

III. Names of species of *Hydrocotyle*, contributed to the 'Monographie du Genre *Hydrocotyle*, par M. Achille Richard, fils' (Extrait du tome 4 des 'Annales générales des Sciences Physiques,' Bruxelles, 1820).

- Hydrocotyle plebeia*, Brown MSS., p. 46.
 — *pulchella*, Brown MSS., p. 59.
 — *flaccida*, Brown MSS., p. 60.
 [H. heteromeria, A. Rich.]
 — *peduncularis*, Brown MSS., p. 62.
 — *intertexta*, Brown MSS., p. 63.
 — *hirta*, Brown MSS., p. 64.
 — *muscosa*, Brown MSS., p. 68.
 — *tripartita*, Brown MSS., p. 69.
 — *pusilla*, Brown MSS., p. 75.
 [H. alata, A. Rich.]

IV. Names of species of *Coniferae*, contributed to Mirbel's "Essai sur la Distribution Géographique des Conifères," in 'Mémoires du Muséum d'Histoire Naturelle,' vol. 13, pp. 74—76, Paris, 1825.

- Callitris* sp., R. Br., p. 74.
 — *glauca*, R. Br., *ibid.*
 — *verrucosa*, R. Br., *ibid.*
 — *calcarata*, R. Br., *ibid.*
 — *robusta*, R. Br., *ibid.*
 — *Ventenatii*, R. Br., *ibid.*
 — *propinqua*, R. Br., *ibid.*
 — *tuberculata*, R. Br., *ibid.*
 — *australis*, R. Br., *ibid.*
Podocarpus Nageia, R. Br., p. 75.
 — *cupressina*, R. Br., *ibid.*
 — *polystachya*, R. Br., *ibid.*
 — *ensifolia*, R. Br., *ibid.*

- Podocarpus elata*, R. Br., p. 75.
 — *spinulosa*, R. Br., *ibid.*
 — *spicata*, R. Br., *ibid.*
 — *thuyoides*, R. Br., *ibid.*
 — *alpina*, R. Br., *ibid.*
 — *latifolia*, R. Br., *ibid.*
 — *falcata*, R. Br., *ibid.*
 — *Antillarum*, R. Br., p. 76.

V. Names of species of *Coniferae*, contributed to Richard's
 'Commentatio Botanica de Coniferis et Cycadeis,'
 Stutgardiae, 1826.

- Callitris rhomboidea*, Brown, p. 47.
 — *fruticosa*, R. Brown MS., p. 49.
Cunninghamia (Brown, in litt. non Willd.), pp. 86
 and 150.

VI. Names referred to by various authors :

Diplachne Baueri, R. Brown, in Desfontaines' "Supplément au Mémoire sur le genre *Chamaelaucium*," in 'Mémoires du Muséum d'Histoire Naturelle,' tom. v, p. 272, Paris, 1819.

Lissochilus speciosus, Br. MSS.

"The name of this fine genus has been communicated to us by Mr. Brown, from his unpublished manuscripts."
 Lindley, 'Collectanea Botanica,' sub tab. 31 ; London. 1821.

HARRISONIA, *Brown*.

"Nomen generi impositum, quo specimina ejus in Herbario Mus. Paris. inscribuntur," Adrien de Jussieu, in Mémoire sur le groupe des Rutacées, in 'Mémoires

du Muséum d'Histoire Naturelle,' vol. xii, p. 517 (1825).

Cryptandra pyramidalis, R. Brown MSS., in 'Ad. Brongniart, 'Mémoire sur la Famille des Rhamnées,' in Ann. Sci. Nat., Ser. 1, 10, p. 373 (Paris, 1827).

Pycnospora, Brown MSS., Wight and Arn. 'Prodr. Fl. Pen. Ind. Orient.' 1, p. 197 (Lond. 1834).

Pycnospora hedysaroides, Brown, *ibid.*

Triphelia brunoides, R. Brown MSS., in Hügel, 'Enumeratio Plantarum Novæ Hollandiæ, Austro-Occidentalis,' p. 48 (Vindob. 1837).

Polyosma Cunninghamii, R. Br. MSS., in 'Plantæ Javanicæ Rariores,' p. 196 (1840).

VI. Names of Plants quoted from Robert Brown's Herbarium in Bentham's 'Flora Australiensis,' vols. 1—6, London, 1863—6.

DILLENACEÆ.

Hibbertia stricta, R. Br. i, p. 27.

— *vetulina*, R. Br., 30.

— *echiifolia*, R. Br., 31.

— *scabra*, R. Br., *ibid.*

MENISPERMEÆ.

Adelioides decumbeus, R. Br., 59.

CAPPARIDEÆ.

Capparis lucida, R. Br., 96.

POLYGALEÆ.

Comesperma aphyllum, R. Br., 143.

— *lanceolatum*, R. Br., 148.

MALVACEÆ.

- Sida micropetala*, R. Br., 201.
Hibiscus thespesioides, R. Br., 220.

STERCULIACEÆ.

- Lasiopetalum rufum*, R. Br., 263.

TILIACEÆ.

- Triumfetta glaucescens*, R. Br., 273.
— *denticulata*, R. Br., 274.
Corchorus Pumilio, R. Br., 277.

LINEÆ.

- Erythroxyton ellipticum*, R. Br., 284.

RUTACEÆ.

- Zieria canescens*, R. Br., 305.
— *involutrata*, R. Br., 306.
— *furfuracea*, R. Br., *ibid.*
Boronia affinis, R. Br., 311.
— *albiflora*, R. Br., 317.
Crowea dentata, R. Br., 330.
Eriostemon parvifolius, R. Br., 335.

LEGUMINOSÆ.

- Brachysema uniflorum*, R. Br., ii, p. 12.
Sphærolobium grandiflorum, R. Br., 66.
Daviesia concinna, R. Br., 75.
— *teretifolia*, R. Br., 82.
Pultenæa viscosa, R. Br., 127.
— *rigida*, R. Br., 130.
— *acerosa*, R. Br., 131.
— *hispidula*, R. Br., 133.

- Galega reticulata*, R. Br., 205.
— *crocea*, R. Br., *ibid.*
— *oblongata*, R. Br., *ibid.*
— *porrecta*, R. Br., 206.
— *astragaloides*, R. Br., 208.
— *juncea*, R. Br., *ibid.*

HALORAGÆ.

- Haloragis paniculata*, R. Br., ii, p. 481.
— *pusilla*, R. Br., *ibid.*
— *lanceolata*, R. Br., 482.
— *stricta*, R. Br., *ibid.*

COMBRETACEÆ.

- Terminalia volucris*, R. Br., 498.

MYRTACEÆ.

- Calythrix decandra*, R. Br., iii, p. 48.
— *la* z, R. Br., 52.
Bæckea Arbuscula, R. Br., 79.
— *pygmæa*, R. Br., 86.
Melaleuca lanceolata, R. Br., 143.
— *cylindrica*, R. Br., 146.
Eucalyptus hypericifolia, R. Br., 203.
— *Baxteri*, R. Br., 207.
— *grandifolia*, R. Br., 250.
— *perfoliata*, R. Br., 253.

LYTHRARIÆ.

- Entelia ammannioides*, R. Br., 296.
Ammannia triflora, R. Br., 297.

LORANTHACEÆ.

- Loranthus angustifolius*, R. Br., 390.

RUBIACEÆ.

- Gardenia suffruticosa*, R. Br., iii, p. 410.
— *fucata*, R. Br., 410.
Ixora triflora, R. Br., 416.
Canthium attenuatum, R. Br., 421.
Opercularia volubilis, R. Br., 435.
Spermacoce brachystema, R. Br., 439.
— *membranacea*, R. Br., 441.
— *purpureo-cærulea*, R. Br., 442.
— *suffruticosa*, R. Br., 443.

COMPOSITÆ.

- Aster argutus*, R. Br., 488.
Calotis pterosperma, R. Br., 505.
Blumea diffusa, R. Br., 525.
Pluchea dentex, R. Br., 529.

SHORT CONTRIBUTIONS

TO

VARIOUS WORKS,

AND

OCCASIONAL NOTICES,

BOTANICAL AND GEOLOGICAL.

de
fro
aie
ex
ale
of
be
“
‘T
Lo

tu
th
of
va
of
it
Ne
ca
co

SHORT CONTRIBUTIONS

TO

VARIOUS WORKS.

"In the preparation of the following notices, I have derived most essential assistance in almost every article, from the information of Mr. Robert Brown; without that aid the account would have wanted those elucidations and explanations which his profound knowledge and research alone could supply, and which has given interest to some of the subjects far beyond what I anticipated or hoped for, before I obtained his promise of co-operation." J. SABINE, "Account of the Edible Fruits of Sierra Leone," in the 'Transactions of the Horticultural Society,' vol. v, p. 441, London, 1824.

ANISOPHYLLÆA, *Brown MSS., ibid.*, p. 446.

Ficus Brassii, *Brown MSS., ibid.*, p. 448.

CARPODINUS, *Brown MSS., ibid.*, p. 455.

"The specimens [of *Hoya*] which I have had the opportunity of examining, afford evidence of the existence of three hitherto undescribed species. The first is a native of the more northern part of New South Wales, with leaves varying from elliptic to obovate; this is in the Herbarium of Mr. Robert Brown, and is named by him *Hoya australis*, it having been referred by him in his 'Prodromus Floræ Novæ Hollandiæ,' vol. i, p. 460, with a doubt, to *H. carnosa*. The second is also in the same collection, and comes from the Nicobar Islands; it has beautiful globose

umbels of flowers, which are very numerous in each umbel, and ovate-lanceolate leaves: Mr. Brown has named this *Hoya nicobarica*." TRAILL, *Accounts and Descriptions of several plants belonging to the genus Hoya, &c.*, in the 'Transactions of the Horticultural Society,' vol. vii, London, 1830.

MAYNA, *Ruddi*.

"Ihre stellung im Natürlichen System betreffend, reiht sich unsere Pflanze unstreitig zunächst an die *Flacourtiaceæ* und *Bixineæ*. Wir haben sie vorläufig mit Fragezeichen zu ersterer Familie in die Nähe von *Hydnocarpus* gestellt, müssen aber dabei einer mündlichen Äusserung R. Brown's gedenken, gemäss welcher sie mit *Hydnocarpus* und *Gynocardia*, Roxb., eine eigene Familie bilden dürfte, deren Aufstellung unser unsterblicher meister hoffentlich später selbst übernehmen wird. ZUCCARINI in *Fasciculus Secundus Plantarum Minus Cognitarum, in Abhandlungen der Königlich Bayerischen Akademie*, band ii, p. 368 (1837).

Prof. Euckland acknowledges assistance from Mr. Brown, in determining the nature of the fossils, for which he states that, at Mr. Brown's suggestion, he had established a new family with the designation *Cycadoideæ*. *Transactions of Geological Society of London*, 2nd series, vol. ii, p. 395. (*Read June 6th, 1828.*)

In a note following the Preface to Dr. Buckland's 'Bridgewater Treatise,' 2 vols., 8vo, 1836, the author says—

"The scientific reader will feel that much additional value has been added to the present work—from the botanical part having been submitted to Mr. Robert Brown."

It is probable that most of the observations on the structure of *Cycadeæ* and Cycadeous fossils, both in this work and in Prof. Buckland's Paper in the 'Geological Transactions,' were contributed by Mr. Brown; but the following are the only particulars for which he is specifically quoted:

CYCADITES.—M. Ad. Brongniart has referred these two fossil species to a new genus, by the name of *Mantellia nidiformis* and *Mantellia cylindrica*. In my paper just quoted, I applied to them the provisional name of *Cycadoidea megalophylla* and *Cycadoidea microphylla*; but Mr. Brown is of opinion that until sufficient reasons are assigned for separating them from the genus *Cycas* or *Zamia*, the provisional name of *Cycadites* is more appropriate, as expressing the present state of our knowledge upon this subject. Buckland's *Bridgewater Treatise*, vol. i, p. 496, note (1836).

In the vascular bundles within the fossil petioles, Mr. Brown has recognised the presence of spiral, or scalariform vessels (*Vasa scalariformia*), such as are found in the petioles of recent *Cycadeæ*; he has also detected similar vessels in the laminated circle within the trunk of the fossil buds next to be described. The existence of vessels with discs peculiar to recent *Cycadeæ* and *Coniferæ*, such as have been described in speaking of fossil *Coniferæ*, has not yet been ascertained. *Ibid.*, p. 499, note.

PODOCARYA.

This fossil was found by the late Mr. Page, of Bishport, near Bristol, in the lower regions of the Inferior Oolite formation on the east of Charmouth, Dorset, and is now in the Oxford Museum. The size of this fruit is that of a large orange; its surface is occupied by a stellated covering or Epicarpium composed of hexagonal tubercles, forming the summit of cells, which occupy the entire circumference of the fruit.

Within each cell is contained a single seed, resembling a small grain of rice, more or less compressed, and usually hexagonal. Where the Epicarpium is removed, the points of the seeds are seen, thickly studded over the surface of the fruit. The bases of the cells are separated from the receptacle, by a congeries of footstalks formed of a dense mass of fibres, resembling the fibres beneath the base of

the seeds of the modern *Pandanus*. As this position of the seeds upon footstalks, composed of long rigid fibres, at a distance from the receptacle, is a character that exists in no other family than the *Pandaneæ*, we are hereby enabled to connect our fossil fruit with this remarkable tribe of plants, as a new genus, *Podocarya*. I owe the suggestion of this name, and much of my information on this subject, to the kindness of my friend, Mr. Robert Brown.

The large spherical fruit of *Pandanus*, hanging on its parent tree, is represented at pl. lxxxiv, fig. 1. Fig. 11 is the summit of one of the many drupes into which this fruit is usually divided. Each cell, when not barren, contains a single, oblong, slender seed; the cells in each drupe vary from two to fourteen in number, and many of them are abortive. The seeds within each drupe of *Pandanus* are enclosed in a hard nut. These nuts are wanting in *Podocarya*, whose seeds are smaller than those of *Pandaneæ*, and not collected into drupes, but dispersed uniformly in single cells over the entire circumference of the fruit. The collection of the seeds into drupes, surrounded by a hard nut, in the fruit of *Pandanus*, forms the essential difference between this genus and our new genus *Podocarya*.

In the fruit of *Pandanus* the summit of each cell is covered with a hard cup or tubercle, irregularly hexagonal; and crowned at its apex with the remains of a withered stigma. We have a similar covering of hexagonal tubercles over the cells of *Podocarya*. The remains of a stigma appear also in the centre of those hexagons above the apex of each seed. Buckland's *Bridgewater Treatise*, vol. i pp. 504, 505 (1836).

In the title-page to the edition of Dr. Buckland's 'Bridgewater Treatise,' published in 1858, after the death both of the author and of Mr. Brown, Mr. Brown's name is placed as having made additions to it. That Mr. Brown suggested to Dr. Buckland many of the remarks contained in the botanical portion of his Treatise, as well as of those contained in his previous papers on the *Cycadoideæ*, in the

'Geological Society's Transactions,' there can be no doubt; but the only "additions" indicated in the notes to this, the third edition, appear to be the following:

Mr. Robert Brown has ascertained, by examination of a trunk of *Cycadites microphyllus*, from Portland, the existence of scalariform vessels without discs, in the mature trunk; a point in which, he informs me, these fossils agree with the American portion of the order *Cycadeæ*, though, in other respects, they bear a greater resemblance to the African and Australian species. Mr. Brown observes further, that the order *Cycadeæ* presents but one genus in America, namely, the *Zamia*, on which this genus was originally founded, and to which it has been recently restricted; and that the coincidence in the structure of the scalariform vessels in the trunk of this *Zamia* of the New World, with that of the fossil *Cycadites* of Europe, is very remarkable. Buckland's *Bridgewater Treatise*, new edition, 1858, vol. i, pp. 461, 462.

Mr. Robert Brown has noticed in the cellular tissue of a silicified trunk of *Cycadites*, portions of chalcedony bearing the form of extravasated gum within the trunks of recent *Cycadeæ*. He has also recognised spiral vessels in the laminated woody circle of a mature trunk of fossil *Cycadites*, and also in the laminated circle within a silicified bud of the same, near its origin. *Ibid.*, vol. ii, p. 102.

Mr. [now Sir] R. I. Murchison, in a paper on the "Fossils found at Eningen," says—

"Mr. R. Brown has discovered that among these plants, one is almost indistinguishable in the leaf from the *Acer villosum*, a species of maple brought from Nepaul by Dr. Wallich." *Geol. Trans.*, 2nd ser., vol. iii, p. 287 (1835).

Mr. Horner, in a note to his paper "On the Geology of the Environs of Bonn," says—

"I visited the lignite deposit at Friesdorf, in September, 1835, in company with Mr. Robert Brown, when he collected several specimens of the vegetable remains. He in-

forms me that all the wood he found is coniferous." *Geol. Trans.*, 2nd ser., vol. iv, p. 474 (1836).

Mr. Charles Stokes in his "Notice respecting a piece of Recent Wood partly petrified by Carbonate of Lime" (*Trans. Geol. Soc. of London*, 2nd series, vol. v, p. 207, 1840), acknowledges assistance obtained from Mr. Brown, and in a "Further Notice" appended to this paper (l. c., p. 213) he says—

"Since I communicated to the Geological Society the preceding notice on the partly petrified wood from the ancient Roman aqueduct of Eilsen, in the principality of Lippe Buckeberg, Mr. Robert Brown has shown to me a specimen from the same piece of wood, which was presented to him at Tharand, in the month of October last, by M. Cotta of that place, who discovered the wood in the aqueduct, and remarked its peculiar condition. Mr. Brown has pointed out to me, in the longitudinal section, that the petrified portions, in his specimen, are about two inches in length, and in the middle part, nearly a quarter of an inch in diameter, and terminate in a point at each end. The petrified portions are, in these instances, completely enclosed within and surrounded by the unchanged wood. See pl. xvi, fig. 3."

"Mr. Brown has observed another remarkable circumstance in the condition of these petrified portions. The change of the longitudinal fibres appears to be complete, but the medullary rays, of which the ends are seen in this section are still in their ligneous state, as shown in the magnified engraving, pl. xvi, fig. 4."

Dr. Fitton, in his paper "On the Strata below the Chalk," says—

"From the evidence afforded by thin transparent slices, both of the transverse and longitudinal sections which have been examined under the microscope by Mr. Brown, the fossil trunks of Portland are found to possess the characters uniformly belonging to coniferous wood; but it must be observed that these characters are not absolutely confined

to Coniferæ." *Geol. Trans.*, 2nd series, vol. iv, p. 222 (1836).

"A point which Mr. Brown considers as well deserving of remark is, that the only remains of vegetables hitherto found in these strata, under the circumstances above described, belong to two nearly related families, Coniferæ and Cycadeæ, which have lately been regarded as forming a distinct class, characterised not only by greater simplicity of the parts of fructification, but also by some peculiarities of the internal structure, and thence have been considered as intermediate between Phænogamous and Cryptogamous or Acotyledonous plants." *Ibid.* p. 225.

"A cone has recently been found on the shore of the Isle of Portland, not improbably derived from one of the beds of clay, or 'Dirt,' subordinate to the lower part of the Portland strata, the structure of which, according to Mr. Brown, approaches in some respects to that of *Araucaria*. I am indebted to the kindness of the Rev. David Williams, of Cross, near Bleadon, in Somersetshire, to whom it belongs, for an opportunity of submitting this beautiful specimen to the examination of Mr. Brown, who will, I hope, describe its structure in the 'Transactions of the Linnean Society.'" *Ibid.* p. 349.

Mr. [now Prof.] Morris, in his appendix to Mr. Prestwich's memoir "On the Geology of Coalbrook Dale," under the description of pl. xxxviii, says—

"*Stigmaria ficoides*. In the 'Fossil Flora,' t. 31—36, are figures and descriptions of *Stigmaria ficoides*, and at t. 156 is shown the structure of the same; and although we cannot add much new information to that previously given by Prof. Lindley, it has been thought advisable to have another section represented, with a view of showing what has hitherto not been well illustrated in the published figures of its structure. The internal cylinder in the specimen (fig. 3) is eccentric, and consists of wedge-like portions of vascular tissue, the rounded origin of which, internally, is well defined; these wedges are generally of equal or nearly equal size, but they occasionally become confluent

by the joining of two or more of them together. The form of the space necessarily left, or interstices between the sections where these are distinct, varies a little; in some cases being of nearly equal breadth throughout, and in others becoming narrower outwards, and appearing to terminate or contract about the middle of the vascular tissue, beyond which they again frequently widen outwards: these spaces often contain portions of oblique and smaller vascular cords, apparently arising at different depths in the vascular cylinder; the origin and connection of which with the cylinder is shown in the oblique section, where a single series of vessels is seen passing from it, surrounded by tissue of smaller diameter. Pl. xxxviii, fig. 3 a."

"In no specimen yet examined has the course of the oblique cords been absolutely ascertained, but there can scarcely be any doubt, as suggested by Mr. Brown (to whom we are also indebted for the above observations), that those vessels after arising from the cylinder passed to the tubercles of the surface, through the thick cellular tissue which once probably occupied the larger space in the original plant. The discovery of these smaller oblique vessels is an interesting feature in the anatomy of *Stigmara*; and they have also been pointed out by Mr. Brown as existing in *Anabathra*, and one of these is actually figured by Mr. Witham, in his work, ('On the Internal Structure of Fossil Vegetables,' 2nd edit., t. 8, f. 12), but considered by him (p. 41) as a section of a medullary ray. The analogous vessels existing in *Lepidodendron Harcourtii*, as figured by Mr. Witham ('Trans. of the Nat. Hist. Soc. of Newcastle, 1832'), appear to arise from the outer part of the vascular cylinder. A somewhat similar arrangement is also found in that division of *Lycopodiaceæ*, consisting of *Psilotum* and *Thesipteris*: in those genera the vascular cylinder, from which the oblique cords proceed, includes a central pith (Brongniart, 'Hist. des Végét. Foss.,' tom. ii, pp. 44, 45).

"Fig. 3 b. Shows that the vessels are much smaller at the internal rounded portions of the wedges.

"Fig. 3 c. Exhibits the oblique cords, consisting of

smaller vessels surrounding vessels of a larger diameter nearly equal to those forming the vertical tissue.

"Fig. 3a. Is an oblique section, which exhibits the connection of these vascular cords with the vertical tissue." *Trans. Geol. Soc.*, 2nd ser. vol. v (1840).

PETROPHILOIDES.

"Upon showing the fossil cones to Dr. R. Brown, he very kindly pointed out to me the affinity existing between them and the genera *Petrophila* and *Leucadendron*, and particularly with one species of the former genus—*Petrophila diversifolia*—described in his 'Prodromus Floræ Novæ Hollandiæ,' page 365."

Dr. Bowerbank's '*History of the Fossil Fruits and Cones of the London Clay*,' page 43 (1840).

Mr. [now Sir] C. Lyell, in a paper 'On the Boulder Formation and Freshwater Deposits of Eastern Norfolk,' says—

"Among the vegetable fossils the most common and best preserved are the seed-vessels of an aquatic plant which Mr. R. Brown refers to *Ceratophyllum demersum*, English Botany, 947."—*London and Edin. Phil. Mag.*, vol. 15, p. 355 (1840).

Lieut. Newbould says that Mr. R. Brown determined the specimens of fossil wood brought by him from Egypt "to be dicotyledonous, and not coniferous."—*Geol. Proc.* iii, p. 787. (*Read Jan. 29, 1842.*)

Dr. Mantell, in a paper 'On Fruits from the Cretaceous Rocks,' says, under *Carpolithes Smithiæ*—

"I am indebted to Dr. Robert Brown for the careful examination of this fossil, and he informed me that he knew of no fruit to which it bore any near affinity, but

suggested that the original was probably a succulent compound berry, the seeds appearing to have been imbedded in a pulpy substance, like the fruit of a mulberry, which is a spurious compound berry, formed by a partial union of the enlarged and fleshy calyces, each enclosing a dry membranous pericarp.—*Quart. Journ. Geol. Soc.*, vol. ii, p. 54 (1846).

The President exhibited portions of trunks of Winter's-bark trees from the Straits of Magellan, cut down in 1826 by Capt. P. P. King, R.N., offering inscriptions made through the bark by a midshipman who accompanied the Spanish expedition under Captain Cordoba in 1786, and by one of the companions of Captain Bougainville in 1767; the annual rings in the former case distinctly corresponding with the interval between 1786 and 1826. He also made some observations on the structure of the woody vessels of the genus.—*Proc. Linn. Soc.*, ii, p. 95 (June 18, 1850).

The President exhibited specimens of stems of *Kingia australis*, R. Br., and *Xanthorrhœa arborea*, R. Br., together with drawings of the former illustrative of its structure, especially of the siliceous covering of the vascular fasciculi of the persistent bases of the leaves; and in both genera, the means by which the stems are protected from the scorching fires of the natives.—*Proc. Linn. Soc.*, ii, p. 113—4 (Feb. 4, 1851).

The President exhibited specimens of the three known species of *Rafflesia* (*R. Arnoldi*, *R. Patma*, and *R. Cumingii*), particularly with the view of showing that they are all three diœcious.—*Proc. Linn. Soc.*, ii, p. 128 (April 15, 1851).

The President exhibited numerous specimens of recent and fossil *Cycadeæ*. Among these was a fine specimen of a new species (*Cycadites Saxbyanus*, R. Br.) found in the Isle of Wight by Mr. Saxby, of Bonchurch. The President remarked, that all the specimens of *Cycadites* hitherto found

in the Isle of Wight agreed in having an elliptical outline, unaccompanied with any inequality in the woody ellipsis, and also in having a bud in the axilla of each leaf; in these respects differing from the *Cycadites* of the Isle of Portland, and from all the recent species of *Cycadæa* with which we are acquainted, which have a circular outline and only scattered buds.—*Proc. Linn. Soc.*, ii, p. 130 (May 6, 1851).

INCARVILLEA.

Having, in examining the Himalayan species of *Incarvillea*, observed some appearances not mentioned in, or at variance with, the descriptions of that genus by Jussieu and Lamarek, I was induced to ask the opinion of the highest authority on the subject, and Mr. Brown has been good enough to favour me with his opinion that the Himalayan species are too closely allied to be separated into a genus distinct from the Chinese *Incarvillea*, but are yet sufficiently distinguished to constitute a sub-genus. He has been good enough to enrich my work with the character of this, and an amendment of the genus *Incarvillea*.

INCARVILLEA. Juss. Gen. 138.

“*Calyx* 5-dentatus, nunc 10-dentatus, [dentibus] alternis (accessoriis) membranaceis. *Corolla* fauce ventricosa; limbo 5-lobo inæquali. *Stamina* antherifera quatuor, didynama, inclusa. *Antheræ* lobis distinctis apice convergentibus, singulis infra apicem processu subulato munitis! *Stigma* bilamellosum. *Capsula* siliquæformis, bilocularis, bivalvis; septo contrario libero intra marginem seminifero. *Semina* suspensa (v. membranaceo-marginata v. comosa), radícula supéra.

“*Herbæ* alternifoliæ foliis pinnatis bipinnatifidisve; racemi terminalis longi pedicellis tribracteatis. Rudimentum filamenti quinti (primum in *Amphicome* detectum a D. Royle). Genus *Bignoniaceis* adjiciendum, in sectione tamen propria ob habitum et semina pendula, radícula supéra (v. *Prodr.*

Flor. Nov.-Holl. 1, p. 471) locandum et in duo subgenera dividendum. Horum alterum INCARVILLEA *Juss.* Chinense, distinguitur, foliis bipinnatifidis; calyce, præter normales, dentibus 5 accessoriis sinuum loco membranaceis emarginatis; seminibus obovatis margine lato membranaceo indiviso cinctis.

“Alterum, AMPHICOME (Cyrtandraceis quodammodo accedens) ex Indiæ septentrionalis montibus: foliis pinnatis foliolis incis; calyce 5-dentato sinubus simplicibus; seminibus fere subulatis utrinque acutis, basi et apice membrana capillaceo-lacera comosis.”

Illustrations of the Botany of the Himalayan Mountains, by J. Forbes Royle, M.D. Pp. 295, 296. London, 1839.

HEDYCHIUM.

SECT. D. *Brachychilum*, Brown MSS. *Labellum* nanum, retusum, sessile. *Stigma* bilabiatum, labio inferiore triplo longiore.

H. (Brachychilum) Horsfieldii, Brown MSS.; glabrum, foliis lanceolatis acuminatis, spica laxiuscula, bracteis ovatis 2—3-floris, tubo gracili, limbi interioris laciniis lateralibus ovalibus obtusiusculis exteriores lanceolatas acutas latitudine triplo excedentibus.

Patria, Mons Prahn Javæ, *Horsfield in Herb Banks.*

I am indebted to my illustrious friend, Mr. Brown, for the following details of the inflorescence of this most singular plant, which I have copied from the original manuscript, lent to me on purpose, and which was written in 1815, when Dr. Horsfield sent the specimen from Java to the late Sir Joseph Banks.

“Planta elegans, glabra, rhizomatosa, juxta specimen unicum in herbario citato circiter bipedalis. *Folia* punctis lineolisque longitudinalibus creberrimis notata. *Calyx* glanduloso-punctatus, tubo tertiâ parte brevior, ore hinc fisso inde acervulo pubescentiæ.

“*Limbi exterioris laciniæ* angusti-lanceolatae, concavius-

culæ, acutæ, tubo breviores, post expansionem reflexæ; tertia acumine brevi subulato; *interioris lacinia laterales* ovales, obtusiusculæ, longitudine limbi exterioris, ejusdem laciniiis duplo et ultra latiores, punctis minutis glandulosis creberrimis instructæ, immerse venosæ. *Labellum* minutum, vix octavam partem longitudinis lateralium emetiens, latius quam longum, retusum, punctatum. *Filamentum* intra laciniam acuminatam limbi exterioris insertum, cum basi labelli oppositi quasi continuum, breve, lineare, concaviusculum. *Anthera* lato-linearis, apice nuda, basi semibifida. *Stigma* bilabiatum, hispidum, labiis ciliatis, inferiore triplo longiore subovato."

Wallich, "Attempt to define the Species of *Hedyckium*," in 'Kew Journal of Botany,' vol. v, p. 376-7 (1853).

Al

Ac

A

A

A

A

A

A

A

INDEX OF SUBJECTS

TO

VOL II.

- Abel, Clarke, characters and descriptions of three new species of plants found in China by, 319—328; notice of his collection, 326
- Acicarpa, observations on the genus, with description of a new species, 307—309; proposed as the type of a new family, 309; further observations on, 312—318
- Actinophora, affinities of the genus, 646
- Albumen, its origin, 23; occasional existence in Proteaceæ, *ibid.*; and in Cyrtandraceæ, 587; its importance, 587 *note*; in Sterculiaceæ, 623
- Allium descendens, exceptional order of expansion of the flowers of, 280
- Anadenia, exceptional order of expansion of the flowers of certain species of, 280
- Angianthus, synonymy of the genus, 283
- Antheræ, their examination at an early period universally neglected, 6; great advantage of this examination, *ibid.*; their structure in Orchideæ and Asclepiadæ, *ibid.*; views of Linnaeus, Jussieu, and Richard on this subject in Asclepiadæ, *ibid.*; views of Jacquin, Koelreuter, Rottboell, Cavanilles, Smith, and Desfontaines, 7; views of the author, 7, 8; and in Orchideæ, 9; development of in Asclepiadæ, 197, 198; arrangement of in Sterculiaceæ, 623
- Antennaria, history and characters of the genus, 300—302
- Apocineæ, difficulty of circumscribing the order, 195; subdivision of it, *ibid.*; characters and descriptions of the genera, and species of the section of the order "seminibus comosis," 232—247
- Ascidia of Cephalotus and other pitcher-plants, observations on the, 357
- Asclepiadæ, views of various authors as to the structure of the Antheræ in, 6—8; examination of, 8; memoir on the order, 193, &c.; characters and descriptions of the order, genera, and species, 200—231; observations on the flower in its earlier stages, 196—198
- Asphodeleæ, order of reduction of stamina in, 278
- Baccharis, history and characters of the genus, 293-4
- Banks, Sir Joseph, his invaluable herbarium and library, 31, 199; important results of his voyage, 163
- Begonia, order of expansion of flowers, and position of female flowers in, 279 *note*
- Bennettia, observations on the structure and affinities of the genus, 663, 664; characters of the species of, 665
- Bentham's Flora Australiensis, names of plants quoted in, from Mr. Brown's herbarium, 708—711
- Bocconia, reductions of pistillum in, 272
- Boopidæ, proposed as a family, by M.

- Cassini, 312; observations on, 312—318; position of ovulum and radicle in, 313; other distinctive characters of the family, 314, &c.
- 'Botanical Magazine,' contributions to, 667, &c.
- 'Botanical Register,' contributions to, 681, &c.
- Bowerbank, Dr., notice of a fossil cone referred to *Petrophila*, in his 'History of the Fossil Fruits of the London Clay,' 723
- Brunonia*, its relations to *Goodenoviæ* and *Compositæ*, 267 *note*; observations on its affinities, 310, 311; nervures of its corolla, 310; structure of its ovary, 311; structure of the indusium of its stigma, 312
- Buckland, contributions to his papers on *Cycadites*, and to his 'Bridgewater Treatise,' 716—719
- Buttneriaceæ*, gradually pass into *Tiliaceæ*, 646
- Buxbaumia*, observations on the genus, 349—351
- Buxbaumia aphylla*, nature of its peristomium, 350, 351; inner membrane of its capsule, 351; furnished with perfect leaves, *ibid.*
- Cæsulia*, nature of capitulum of, 275-6
- Calca*, history, synonymy, and subdivision of the genus, 287—307
- Calyceræ*, proposed as a new family, 309; further observations on, 312—318: the name superseded by that of *Boopideæ*, 313
- Calytrix*, observations on the genus and on the species composing it, 684
- Calyx* and adhering tube of corolla, separability of, in some species of *Goodenia*, 267 *note*
- Campanulaceæ*, æstivation of corolla in, 269 *note*
- Capitulum of *compositæ*, order of expansion of florets in the, 273-7
- Caprifoliaceæ*, limitation of the family, with observations on the genera composing it, 324; further observations on the genera, 673
- Cardiopteris*, observations on the structure and affinities of the genus, 657—659
- Cassiniæ*, characters of the genus, with a synopsis of the species, 304—306
- Catesbæa*, arrangement of the nerves of the corolla in, 268
- Celtis*, order of expansion of flowers, and position of female flowers in, 279
- Cephalotæa*, establishment of the family, 357
- Cephalotus*, remarks on the structure and affinities of, 353—359; account of the ripe fruit of, 356; entitled to rank as a distinct family, 357; nature of its ascidia, *ibid.*
- Cestrum*, arrangement of nerves in the corolla of, 267
- Chalaza*, manifest in the seeds of all *Proteaceæ*, 22, 23; its nature and function, 23
- China, characters and descriptions of three new species of plants found in, 319—328
- Cleome*, species quæ in horto Kewensi coluntur, 413—415
- Compositæ*, observations on the natural family of plants called, 267—318: object of the memoir, 259; arrangement of nerves in the corolla of, 260—268; claim of M. Cassini to priority in regard to the arrangement, 261; refuted, 261-2; observations of Grew, Van Berkhey, Schmidel, Batsch, Schkuhr, Mirbel, and Cassini, 262—264; remarks on M. Cassini's statement, 264—267; nerves in the corolla of *Goodenoviæ*, 266; in *Ernodeæ*, *Datura* and *Cestrum*, 267; æstivation of the corolla, 268-9; exceptions to in *Chuquiraga* and *Corymbium*, 269; form of pollen, 269-70; disposition of the stigmata, 270; structure of ovary, 270-72; inflorescence and order of expansion of the florets, 272—281; on what dependent, 276: genera et species quædam quæ in horto Kewensi coluntur, 459—470
- Craspedia*, history and synonymy of the genus, 285
- Crucifæræ*, type of pistillum, and of obliterations in, 272; genera et species cruciferarum quæ in horto Kewensi coluntur, 367—413
- Cycadites*, observations on the struc-

- ture of, 716—719; Saxbyanus, account of, 724
- Cyrtandraeae, observations on, and monograph of the order, 584—610; history of the order, 584—586; affinity of, 386—588; relation of stigmata and placenta in, 386
- Cyrtandrae, modifications of the different parts of fructification in, 590—594
- Datura, arrangement of nerves in the corolla of, 267
- Dawsonia, nature and origin of its peristomium and columella, 331-2
- Decandolle's 'Systema Vegetabilium,' names, &c., of plants contributed by Mr. Brown to, 703; 'Prodromus,' names, &c., of plants, contributed to, 705
- Dipsacae, inflorescence of, and order of expansion of flowers in, 277-8; position of ovary in, 314-5
- Dryander, advice and assistance derived from him, 31
- Dryandra, observations on the genus, 179
- Echinops, nature of capitulum, 274-5
- Embryo, direction of, in the Sterculiaceae, 621-2
- Ernodea, arrangement of nerves in the corolla of, 267
- Euphorbia, its inflorescence, 279 and note, 280; relative position of sexes in, 280
- Ferns, number of species known to Linnæus, 251; to Willdenow, *ibid.*; known in, 1838, 565; vasa sealariformia, their arrangement in, 564; ramification of veins, and relation of veins to sori in, 565—570; seeds of, useful in distinguishing genera, 276
- Fitton, Dr., notice of fossil Coniferae and Cycadeae in his paper 'On the Strata below the Chalk,' 720-1
- Franklin, extracts from the botanical appendix to his 'Narrative of a Journey to the Shores of the Polar Sea,' and addenda, 511—527
- Fruit of Proteaceae, its nature, 21, 22; its varieties in the order, 22
- Globularia cordifolia, arrangement of the nerves of the corolla in, 315-6
- Goodenia, separability of calyx and of adhering tube of the corolla in some species of, 267
- Goodenovieae, arrangement of nerves in the corolla of, 266; its analogy to that of Compositae, 267 note; aestivation of corolla in, 269 note; type of pistillum and reductions of, 272; structure of the indusium of its stigma, 312
- Gramineae, nature of inflorescence of, 276-7; order of expansion of flowers in, 277; composition of inner valve of the flower in, 571-2
- Grevillea, exceptional order of expansion of the flowers in certain species of, 280
- Grindelia, history, characters, and extent of the genus, 282-3
- Gymnostomum, structure of various species referred to the genus, 342—5; great diversity in these species, *ibid.*
- Hedychium, account of a new subgenus and species of, 725-6
- Hernandiaceae, observations on the family, 651
- Hornier, Mr. Leonard, note on coniferous wood in his paper on the 'Geology of Bonn,' 719
- Horsfield, extracts from his 'Plantae Javanicae Rariores,' 557, &c.
- Hortus Kewensis, extracts from the second edition of, 367—510
- Humboldt, his observations on a remarkable difference in the distribution of the species of plants, 12
- Hymenostomum, establishment of the genus, 342
- Iucarvillea, observations on the genus, 725
- Indian plants, names of and notes on, extracted from Dr. Wallich's 'List,' 529—538; notes and observations on, extracted from Dr. Wallich's 'Plantae Asiaticae Rariores,' 539—556
- Inflorescence, types of, and order of expansion of flowers, 272—281; in Compositae, *ibid.*; in Umbelliferae, 273 note; in Gramineae, 276-7;

- in Dipsacæ, 277-8; in B. opidæ, 316
- Iôdes, affinities of the genus, 651-2
- Jacaranda, observations on the genus and on its species, 670-1
- Junceæ, order of reduction of stamina in, 278
- Kingia Australis*, notice of the structure of its stem, 724
- Lagasca, nature of capitulum of, 374-5
- Lambertia, inflorescence of, and order of reduction of the flowers of, 280-1
- Leguminosæ, genera et species quædam quæ in horto Kewensi coluntur, 417-447
- Leontopodium, establishment of the genus, 302
- Leptostomum, observations on the genus, 341-349; reasons for retaining it, 342-3
- Leucadendron, reasons for differing from Salisbury on the application of the Linnean name, 29-31; history of the genus, *ibid.*; observations on the separation of sexes in, 40-1
- Liliacæ, order of reduction of stamina in, 178
- Limnanthus, characters and description of, 361-364
- Limnanthes, establishment of the family, 363; characters, 364
- Linnean System of Botany, advantages derived from the, 5
- Lobeliaceæ, æstivation of corolla in, 269 *note*
- Lyellia, characters and description of, a new genus of mosses, &c., 329-356; characters of the genus, 332-334; structure of its columella, and mode of dispersion of its seeds, 335; use of the pores at the base of its capsule, *ibid.*; its position in the family, 336
- Mantell, Dr., note on *Carpolithes Smithie*, in his paper on 'Fruits from the Cretaceous Rocks,' 723
- Melampodium, history and synonymy of the genus, 284-5
- Metalasia, establishment of the genus, 302
- Meyera, history and synonymy of the genus, 283-4
- Mirbel, names of species of Coniferæ contributed to his 'Essai,' 706
- Morris, Prof., notice of *Stigmaria*, in his paper 'On the Geology of Coalbrook Dale,' 721-723
- Mosses, necessity of additional sources of distinction in the genera of, 345; nature of some of these sources, 346
- Murchison, Sir R. I., note on a fossil leaf, in his paper on Eningen fossils, 719
- Myrtacæ, genera et species quædam quæ in horto Kewensi coluntur, 449-487
- Nepenthes, its place in the natural series, 358; distinctions between *Cytinæ* and, *ibid.*; its spiral vessels, *ibid.*
- Newbould, Lieut., notice of fossil wood from Egypt, in his paper on the subject, 723
- Nivenia, reasons for changing Salisbury's generic name, 110
- Orchideæ, structure of anthera in, 6, 9; genera et species quæ in horto Kewensi coluntur, 471-499
- Ovarium, utility of ascertaining its internal condition before fecundation, 5; steadily kept in view by A. du Petit-Thouars, *ibid.*; its importance in Proteacæ, 19, 20; its structure in Compositæ, 270
- Ovula, importance of their insertion as well as number, 20; insertion, direction, and relation to ovarium, 313
- Ozothamnus, characters of and observations on the genus, 303-304
- Peristomium of mosses, observations on the, 346-349; of *Buxbaumia*, 350-1
- Petit-Thouars, Aubert du, almost the only botanist who has steadily kept in view the internal condition of the ovarium before fecundation, 5
- Petrophila, observations on the sexes in the genus, 55, 56
- Phytocrenæ, characters of the family

and of the genera composing it, 652-3
 Pistillum of all Phanogamous plants considered as formed on the same plan, of which a polyspermous legumen or folliculus, whose seeds are disposed in a double series, may be taken as the type, 270; a circular series of these pistilla disposed round an imaginary axis, and whose number corresponds with that of the parts of the calyx or corolla, enters into the author's notion of a flower complete in its parts, *ibid.*; deviations from this type and number, how accounted for, *ibid.*; the type and its reduction in the family of Proteaceæ, 272; in Goodenoviæ, *ibid.*; in Cruciferae, *ibid.*; order of development of stamina and pistilla, 281
 Pittosporum revolutum, arrangement of nerves of the corolla in, 268 *note*
 Plantæ Javanicæ Rariores, extracts from, 557, &c.
 Plantago, an exception to the ordinary rule of development of stamina and pistilla, 281
 Podocarya, a fossil fruit referrible to Pandanæ, observations on, 717-8
 Pollen, importance of its figure in fixing our notions of the limits of genera, 19; form of, in Compositæ, 269-70
 Polypodium, expediency of subdividing the genus, 252; divisions proposed, 565-570
 Polytrichoidæ, distinguishing characters of the group, 336-341; structure of peristomium in, 340
 Polytrichum, characters of the genus, 336; structure of the lamellæ in, *ibid.*; form and position of male flowers in, 337; double calyptra of, *ibid.*; teeth of peristomium in, *ibid.*; transverse membrane of tympanum in, *ibid.*; various forms of capsule in, 338; inner membrane of capsule in, *ibid.*; inner peristomium of, 339; columella of, 339-40; seeds of, 340.
 Portlandia, arrangement of the nerves of the corolla in, 268
 Poterium, order of expansion of flowers

and position of female flowers in its spike, 279
 Protea, reasons for differing from Salisbury in the application of the Linnean name, 24-29; history of the genus, 25-29
 Proteaceæ, on the natural order of, 3; remarks on its geographical distribution, 9; almost entirely confined to the southern hemisphere, 9; occurs in the greatest abundance and variety in Southern Africa and New Holland, 10; its distribution in New Holland, 10-12; stations and elevations at which its species are found, 13; pubescence of, 14; disposition of, 14, 15; form and division of leaves, 15; inflorescence of, *ibid.*; calyx of, why so considered, 15, 16; identity of calyx with the organ so called in Thymelæ, 16; better called perianthium, or perigonium, *ibid.*; invariably divided into four leaves or segments, *ibid.*; æstivation of, 16, 17; its colour useful in indicating genera, 17; scales surrounding the ovarium, their use, importance, and number, *ibid.*; varieties in structure, or apparent origin of stamina, 17, 18; deviations from the usual structure of antheræ in, 18; figure of the pollen in, 19; its importance in fixing the limits of genera, *ibid.*; internal structure of the ovarium of the greatest importance, 19, 20; number and insertion of ovula, 20; position of radicle, *ibid.*; form of style, *ibid.*; form and direction of stigma, 20, 21; nature of fruit, 21, 22; its varieties in the order, 22; chalaza always manifest, 22, 23; its nature and functions, 23; albumen, its origin, 23; occasional existence in Proteaceæ, *ibid.*; radicle always points towards the base of the fruit, 23, 24; plumula inconspicuous in the whole order, 24; cotyledons, when more than two of little importance, 24; genera formed before the publication of Salisbury's arrangement of the family, 24; reasons for differing from Salisbury in the application of the Linnean

- names *Protea* and *Leucadendron*, 24—31: characters and descriptions of the order, its genera and species, 32—192; type of the pistillum, and its reductions in, 272
- Radicula*, its direction in *Proteaceæ* uniformly pointing towards the base of the fruit, 20, 23, 24; relation of its direction to ovulum and to ovary, 313-4
- Rafflesia*, all the three known species shown to be diœcious, 724
- Restiaceæ*, order of reduction of stamina in, 278
- Richard, Achille, names of species of *Hydrocotyle*, contributed to his 'Monographie,' 706
- Richard, Louis Claude, names of species of *Coniferae*, contributed to his 'Commentatio,' 707
- Richardson, Dr., extracts from his botanical appendix to the narrative of a journey to the shores of the Polar Sea, by Captain Franklin, and addenda, 511—527
- Royle, Dr., observations on *Incarvillea*, from his 'Botany of the Himalayan Mountains,' 725
- Rubiaceæ*, difficulty of circumscribing the order, 195; restivation of corolla in, 269 *note*
- Sabine, Joseph, contributions to his 'Account of the Edible Fruits of Sierra Leone,' 715
- Sarcostigma*, affinities of the genus, 651-2
- Scabiosa*, inflorescence of certain species of, 278; order of expansion of flowers in, *ibid.*
- Seeds, non-existence of naked, 21; dispersion of, in *Polytrichoidæ*, how secured, 335, 340-1
- Sexes, in the separation of, in flowers, the female the more perfect, 279; and the earliest developed, *ibid.*; exceptions to, 279, 280
- Solander, Dr., his description of *Knightia* given as a specimen of the accuracy with which he described specimens of natural history, 163
- Soliva*, history, characters, and extent of the genus, 282
- Spike, order of expansion of flowers in a simple and in a compound, 273
- Spiral vessels of *Nepenthes*, 358-9
- Stamina* and *pistilla*, importance of attending to their state before the expansion of the flower, 5; order of reduction of *stamina* in various families, 278; order of development of *stamina* and *pistilla*, 281; origin of glandular disk in various families from imperfect *stamina*, 317-8; objections to this view, *ibid.*
- Sterculiaceæ*, history of the family, 613—621; observations on the relative importance of the different organs in the formation of genera in, 621—624; direction of embryo in, 621; monograph of, 611—644
- Stigma, in *Proteaceæ*, its form and direction, 20, 21; in *Asclepiadæ*, its development, 196—198; disposition of *stigmata* in *Compositæ*, 270; nature of the *indusium* in *Brunonia* and *Goodenovia*, 312
- Stokes, Mr. Charles, notice on partly petrified wood, in his paper on the subject, 720
- Style, its utility in the generic characters of *Proteaceæ*, 20; coherence of its base with the apex of the tube of the calyx in *Boopideæ*, 315
- Stylidæ*, position of *stamina* in, 312
- Tiliaceæ*, gradually pass into *Buttneriaceæ*, 646
- Traill, contributions to his 'Account of the genus *Hoya*,' 715-6
- Tridax*, history of the genus, 283
- Umbelliferae*, surface of the pericarpium in, 267 *note*; order of expansion of flowers in, 273 *note*
- Vestia* referred to *Solanæ*, 683
- Wallich, Dr., account of a new sub-genus and species of *Hedychium*,

<p>from his monograph of the genus,
725-6</p> <p>Winter's bark, notice of trunks of,
containing inscriptions, and of the
structure of their woody vessels,
724</p> <p>Woodsia, a new genus of ferns, obser-
vations on, 249—255; singularity of
structure of its involucre, 252-3;</p> | <p>characters of the genus and its spe-
cies, 253-4</p> <p>Xanthorrhoea arborea, notice of the
structure of its stem, 724</p> <p>Zuccarini, notice of the genus Mayna
and its affinities, in his 'Plantæ
minus Cognitæ,' 716</p> |
|--|--|

Ab

Ab
Ab

Ab
Ac

Ac
Ac
Ac
Ac
Ac

Ac
Ac
Ac

A

INDEX OF NAMES OF PLANTS

TO

BOTH VOLUMES.

Abelia, i, 448; ii, 324, 541, 673
 Chinensis, ii, 325, 541
 triflora, ii, 531, 541
 uniflora, ii, 542
Abelmoschus, ii, 546
Abroma, i, 11; ii, 509
 augusta, ii, 509
 fastuosa, ii, 509, 510
 Wheleri, ii, 509
Abrus precatorius, i, 111, 165, 168
Acacia, i, 23, 62, 64, 110, 339; ii, 445
 acicularis, ii, 445
 alata, ii, 446
 armata, ii, 446
 biflora, ii, 446
 ciliata, ii, 447
 decipiens, ii, 446
 fasciculata, i, 95
 læta, i, 95
 marginata, ii, 446
 melanoxydon, ii, 445
 nigricans, ii, 447
 nilotica, i, 288
 pubescens, ii, 447
 pulchella, ii, 446
 Sophoræ, ii, 445
 sulcata, ii, 445
Acaciæ, i, 311
Acanthaceæ, i, 134, 380
Acanthus tetragonus, i, 94
Acer villosum, ii, 719
Aceras, ii, 475
 anthropophora, ii, 475
Achenopsis, ii, 684
Achyranthus papposa, ii, 543
Acicarpa, ii, 309, 312
 lanata, ii, 309
 spathulata, ii, 307
 tribuloides, ii, 309, 312
Acioa, i, 115

Acorum maritimum, ii, 536
Acrostichum, ii, 250
 alcicorne, i, 147, 165
 alpinum, ii, 254
 aureum, i, 147, 165, 169
 calomelanos, i, 473
 hyperboreum, ii, 254
 ilvense, ii, 254
 Stemaria, i, 147, 165
 velleum, i, 303
Actinophora, ii, 645
 fragrans, ii, 645
Actinotus, i, 29
Adansonia, i, 153
 digitata, i, 288
Adelanthus, ii, 652
Adelioides decumbens, ii, 708
Adenanthera, i, 22
Adenanthos, ii, 16, 20, 21, 36, 125
 cuneata, ii, 125
 obovata, ii, 125
 sericca, ii, 125
 terminalis, ii, 125
Adenophorus, ii, 566
Adiantum, i, 146
 Africanum, i, 148 *note*
 capillus-veneris, i, 148 *note*
 tenerum, i, 148 *note*
 truncatum, i, 83; ii, 446
Ægiceras, i, 150 *note*, 364, 380
Ægopricon, i, 127
Ærides, ii, 492
 odoratum, ii, 492
 paniculatum, ii, 689
Æschynanthus, ii, 585, 590, 594, 596
 acuminata, ii, 597
 bracteata, ii, 597
 fulgens, ii, 596
 Griffithii, ii, 596
 Horsfieldii, ii, 597
Æschynanthus longicaulis, ii, 597

- Æschynanthus parasitica*, ii, 596
 parvifolia, ii, 596
 radicans, ii, 596, 597
 ramosissima, ii, 596
 volubilis, ii, 596
 Wallichii, ii, 597
Æschynomene, ii, 444
 Americana, ii, 444
 aspera, ii, 444
 bispinosa, ii, 444
 hispida, ii, 444
 indica, ii, 445
 sensitiva, ii, 444
Ætheilema, i, 134
Æthionema, i, 271 ; ii, 376
 monospermum, ii, 376
 saxatile, ii, 376
Agalmyla, ii, 593, 594, 597
 staminea, ii, 598
Agaricus alneus, i, 70
 campestris, i, 70
 muscarius, i, 70
Agastachys, ii, 17, 37, 132
 odorata, ii, 132
Agathis, i, 47, 458, 461
Agaty, ii, 443
Ageratum conyzoides, i, 165, 167
Agoul, i, 295
Agrostis, i, 221
 algida, i, 177
 laxiflora, ii, 514
 paradoxa, i, 177, 222
 virginica, i, 58, 164, 166
Aikinia, ii, 581, 606
 Brunonis, ii, 554
 Horsfieldii, ii, 556
Aira, i, 232
 aquatica, ii, 514
 spicata, i, 231
 subspicata, i, 231 ; ii, 514
Akee, i, 161
Alchornea, i, 127, 164
Alcina, ii, 285
Alfonzia oleifera, i, 141
Alhagi Maurorum, i, 295
Alisma Plantago, i, 68
Allamanda, i, 532
Allium, ii, 550
 descendens, ii, 250
Allmania albida, ii, 534
 esculenta, ii, 534
 nodiflora, ii, 534
Aloc, i, 64
Alopecurus, i, 55
 alpinus, i, 67, 177, 219 ; ii, 514
Alopecurus antarcticus, i, 219
 aristulatus, ii, 514
 ovatus, i, 219
 pratensis, i, 219
Alsino rubella, i, 199
 stricta, i, 201
Alsodeia, i, 123
Alsodina, i, 123
Alstonia, ii, 233, 245
 costata, ii, 246
 scholaris, ii, 245
 spectabilis, ii, 246
 venenata, ii, 246
Alyssum, i, 266, 267 *note*, 271 ; ii, 386
 alpestre, ii, 387
 austriacum, ii, 386
 calycinum, i, 267 *note* ; ii, 387
 campestre, i, 267 *note* ; ii, 387
 clypeatum, ii, 389
 deltoideum, ii, 389
 densiflorum, i, 267 *note*
 Dioscoridis, ii, 389
 fulvescens, i, 267 *note*
 halimifolium, ii, 388
 hirsutum, i, 267 *note*
 incanum, ii, 389
 lunarioides, ii, 388
 maritimum, i, 266 ; ii, 388
 micropetalum, i, 267 *note*
 minimum, i, 267 *note* ; ii, 388
 montanum, ii, 387
 mutabile, ii, 389
 rostratum, i, 267 *note*
 sativum, ii, 386
 saxatile, ii, 387
 sinuatum, ii, 390
 spinosum, ii, 388
 strictum, i, 267 *note*
 strigosum, i, 267 *note*
 umbellatum, i, 267 *note*
 utriculatum, ii, 389
Alyssinae, i, 263
Alyxia, i, 37
Amaranthaceae, i, 36 ; ii, 543, 545
Amaryllideae, i, 51
Amaryllis, i, 364
Ambora, i, 25
Ambun Ambun, i, 393
Ambrosia, i, 31
Amellus, ii, 295, 296
Ammannia, ii, 535
 triflora, ii, 710
Ammobium, ii, 676
 alatum, ii, 677

Amomum granum Paradisi, i, 155
Ampelopsis, i, 151
Amphicome, ii, 588, 725
Amygdaleæ, i, 115
Amyridæ, i, 112
Amyris Gileadensis, i, 94
 Kataf, i, 94
Anabathra, ii, 722
Anacardæ, i, 112
Anacardium, i, 43, 112
Anadenia, ii, 37, 139, 280
 illicifolia, ii, 140
 pulchella, ii, 140
 trifida, ii, 140
Anagallis, i, 297
 cærulea, i, 297
Anagyris, ii, 421
 fetida, i, 25; ii, 421
Anasser, i, 37, 132
Anastatica, ii, 376
 hieracantha, ii, 376
 Syriaca, ii, 371
Anchusa affinis, i, 94
Aneistrocarpus, i, 139 *note*
Andromeda tetragona, i, 177, 213
Andropogon, i, 55, 57, 146
 acicularis, i, 302
Angianthus, ii, 283
 tomentosus, ii, 283
Angophora, i, 311
Angræcum, ii, 689, 694
Anictangium aquaticum, ii, 335
 ciliatum, i, 69
 Humboldtii, ii, 335
 pulvinatum, ii, 344
Anisomeles, i, 380; ii, 503
 malabarica, ii, 670
 ovata, ii, 503
Anisophyllæa, ii, 715
Anodontium, i, 348
Anona, i, 164
 Asiatica, i, 105
 hexapetala, ii, 685
 microcarpa, i, 162 *note*
 mucosa, i, 105
 muricata, i, 105
 myristica, i, 162 *note*
 obtusiflora, i, 105
 Senegalensis, i, 105, 153, 155
 uncinata, ii, 685
Anonaceæ, i, 37, 74, 105, 382; ii, 685
Anopterus, ii, 523
Antennaria, ii, 301
 alpina, i, 212

Anthemis apiifolia, ii, 690
 odorata, ii, 466
Anthenantia, i, 57
Anthericum, i, 50
 cirratum, ii, 676
 latifolium, ii, 676
Antherylum, i, 120
Anthistiria, i, 57
 australis, i, 43, 308, 309
Anthobolus, i, 42
Anthoceros punctatus, i, 69
Anthoelcista, i, 133, 153
Anthospermum, i, 131
Anthoxanthum, i, 231; ii, 571
Antiaris, i, 78
 macrophylla, i, 78
 toxicaria, i, 79
Antidaphne, i, 413 *note*
Antidesma, i, 155; ii, 663
Antidesmea, ii, 663
Antirrhinum majus, i, 479
Antonia, ii, 577
Aotus, ii, 430
 ferruginea, ii, 430
 villosa, ii, 430
Aphanes arvensis, i, 68
Aphelandra, ii, 505
 cristata, ii, 505
Aphyteia, i, 388, 391, 396, 401, 461
 Hydnora, i, 416
 multiceps, i, 429
Aplodon, i, 210, 217
Apocineæ, i, 36, 37, 131, 133, 151, 295, 469; ii, 6, 22, 195, 551
Apocinum, ii, 233, 239
 androsæmifolium, ii, 239
 cannabinum, ii, 240
 cordatum, ii, 229
 frutescens, ii, 236
 hastatum, ii, 227
 hypericifolium, ii, 240
 laucolatum, ii, 229
 lineare, ii, 228
 minutum, ii, 227
 pubescens, ii, 240
 Sibiricum, ii, 240
 triflorum, ii, 228
 Venetum, ii, 240
Apodanthæ, i, 429
Apodanthes, i, 424, 430
 Berteri, i, 430
 Blanchetii, i, 430
 Calliandra, i, 431
 Casearia, i, 430
Apostasia, i, 503, 560; ii, 547

- Apostasia nuda*, ii, 548, 551
 odorata, ii, 547, 551
 Wallichii, ii, 548, 551
Appendicula, ii, 575
Aquilaria, i, 126
Aquilarinæ, i, 126
Aquilegia, ii, 318
Aquilicia, i, 151
Arabis, i, 195, 271 ; ii, 395
 alpina, ii, 395
 arenosa, ii, 396
 bellidifolia, ii, 395
 cærulea, ii, 395
 Canadensis, i, 268 ; ii, 397
 ciliata, ii, 396
 falcata, ii, 397
 hirsuta, ii, 397
 hispida, ii, 396
 nudicaulis, i, 197
 nutans, ii, 395
 pendula, i, 268 ; ii, 397
 pumila, ii, 395
 stricta, ii, 396
 thaliana, ii, 396
 turrita, i, 268 ; ii, 397
 verna, ii, 395
Arachis, i, 24
 hypogæa, i, 154, 159
Araliacæ, ii, 322
Araliæ, i, 29
Araucaria, i, 456
 excelsa, i, 48, 64 ; ii, 721
Arbor radulifera, i, 72
Arctotheca, ii, 465
 repens, ii, 465
Arctotis, ii, 298, 467
 acaulis, ii, 467
 angustifolia, ii, 468
 argentea, ii, 468
 aspera, ii, 469
 auriculata, ii, 469
 calendulacea, ii, 465
 crithmoides, ii, 466
 decumbens, ii, 468
 decurrens, ii, 468
 dentata, ii, 466
 fastuosa, ii, 469
 flaccida, ii, 468
 glaucophylla, ii, 467
 grandiflora, ii, 467
 hypochondriaca, ii, 465
 paleacea, ii, 466
 paradoxa, ii, 466
 plantaginea, ii, 465
 repens, ii, 465
Arctotis reptans, ii, 469
 rosea, ii, 468
 scariosa, ii, 466
 spinulosa, ii, 469
 tricolor, ii, 467
 undulata, ii, 467
Ardisia, i, 150 *nole*
Areca, i, 51
Arenaria marina, i, 66, 68, 310
 quadrivalvis, i, 199
 Rossii, i, 200
Arethusa, i, 507, 538 ; ii, 486
 bulbosa, ii, 487
 divaricata, ii, 485
Aristida ramosa, i, 93
Aristolochia, i, 384 ; ii, 358, 534
 cordiflora, i, 385
 grandiflora, i, 385
Arnica angustifolia, i, 210
 fulgens, i, 211
 montana, i, 210
 plantaginica, i, 211
Aroideæ, i, 169, 298, 365, 381, 383
Arrhenatherum, i, 232
Artabotrys, ii, 685
 odoratissimus, ii, 685
Arthrochloa, i, 232
Arthropodium cirratum, ii, 676
 paniculatum, ii, 676
Artocarpæ, i, 138
Arundo, i, 58
 Canadensis, ii, 514
 Phragmites, i, 43, 58, 68, 301
 stricta, ii, 514
Asarinæ, i, 384, 385, 392, 396, 401 ;
 ii, 358
Asclepiadæ, i, 36, 37, 133, 134, 295,
 466, 487, 513, 516, 545, 549 ;
 ii, 6, 195, 551
Asclepias, ii, 6, 216
 amœna, i, 524 ; ii, 216
 aphylla, ii, 226
 arborescens, ii, 217
 carnosa, ii, 209
 citrifolia, ii, 216
 cordata, ii, 226
 crispa, ii, 217
 curassavica, i, 519, 534 ; ii, 216
 Davurica, ii, 223
 decumbens, ii, 216
 elevata, ii, 216
 fruticosa, ii, 217
 gigantea, ii, 218
 grandiflora, ii, 217
 incarnata, ii, 216

- Asclepias laetifera*, ii, 213
laniflora, i, 94; ii, 218
linaria, ii, 216
Mexicana, ii, 216
nigra, ii, 224
nivea, ii, 216
parviflora, ii, 216
phytolaccoides, i, 528, 535, 549
procera, ii, 218
pubescens, ii, 217
pulchra, ii, 216
purpurascens, i, 524, 534, 535;
 ii, 216, 528
rubra, ii, 216
scandens, ii, 226
setosa, ii, 217
Sibirica, ii, 224
stipitacea, ii, 226
Syriaca, ii, 196, 216
tuberosa, ii, 216
undulata, ii, 217
variegata, ii, 216
verticillata, ii, 216
viminalis, ii, 226
Vincetoxicum, ii, 224
volubilis, ii, 209
vomitaria, ii, 220
Asparagæ, i, 50
Asperuicæ, i, 36
Asphodelæ, i, 50, 52, 298, 433; ii, 278
Aspidium, ii, 546
 fragile, ii, 518
 fragrans, ii, 518
 trifoliatum, ii, 543
Asplenium, i, 59, 146; ii, 250
 Filix-fœmina, i, 60
Astelia, i, 50, 64
Astelma, ii, 691
 eximium, ii, 691
Astephanus, ii, 228
Aster argutus, ii, 711
 glutinosus, ii, 461
Asteranthos, i, 387 *note*, 388 *note*
Astragalus alpinus, i, 209
Astrantia, ii, 273
Astranthus, i, 120
Ataxia, i, 231; ii, 571
 Horsfieldii, ii, 571
Atherospermæ, i, 25, 64
Athyrium fragile, ii, 518
Atlas tree, ii, 30
Atriplex Halimus, i, 66, 68
Attil, i, 283
Attyah, i, 112 *note*
Atylus, ii, 55, 58
Aubrietia, i, 271
Aulax, ii, 19, 21, 35, 39, 41
 pinifolia, ii, 39, 41
 umbellata, ii, 39, 40
Aurantiacæ, ii, 537
Avena, i, 231
 arundinacea, i, 301
 Forskalii, i, 301
 striata, ii, 514
Avenacæ, ii, 571
Averrhoa, i, 113
 Carambola, i, 114
Avicennia, i, 40, 364
Axyris Moorcroftiana, ii, 534
Azolla, i, 87
 pinnata, i, 87
Baccharis, ii, 293, 294
 nereifolia, ii, 294 *note*
 scoparia, ii, 306
Bac, ii, 590, 593, 605
 Commersonii, ii, 605
 hygrometrica, ii, 605
 multiflora, ii, 605
 Wallichii, ii, 605
Bæckia, i, 19
 arbuscula, ii, 710
 camphorata, ii, 679
 pygmæa, ii, 710
Bæobotrys, i, 151 *note*, 293
Balanophora, i, 389, 391
 dioica, i, 392 *note*; ii, 534
 fungosa, i, 392 *note*
Balanophoreæ, i, 412
Balanites, i, 44
 Ægyptiaca, i, 287
Balbisia, ii, 283
 elongata, ii, 283
Balfouria, ii, 233, 241
Ballota disticha, ii, 503
Bambusa, i, 56
Bamia Abelmoschus, ii, 532
Banana, i, 156
Banisteria, i, 164
 Leona, i, 105
Banksia, i, 116, 308, 310, 450, 455;
 ii, 11, 15, 17, 19, 38
 œmula, ii, 177
 asplenifolia, ii, 176
 attenuata, ii, 176
 australis, ii, 174
 coccinea, ii, 175
 collina, ii, 172
 compar, ii, 175
 conchifera, ii, 177

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

8
18
20
22
25
28
32
36
40
45
50

10
01

- Banksia dactyloides*, ii, 22, 156
 dentata, ii, 177
 depressa, ii, 173
 elatii, ii, 177
 ericifolia, ii, 172
 gibbosa, ii, 152
 glauca, ii, 174
 grandis, ii, 178
 ilicifolia, ii, 178
 insularis, ii, 174
 integrifolia, ii, 12, 174
 latifolia, ii, 176
 littoralis, ii, 173
 marcescens, ii, 176
 marginata, ii, 173
 microstachya, ii, 173
 musculiformis, ii, 190
 nivea, ii, 181
 nutans, ii, 172
 oblongifolia, ii, 175
 occidentalis, ii, 172
 oleaefolia, ii, 156, 174
 paludosa, ii, 175
 patula, ii, 173
 pinifolia, ii, 152
 præmorsa, ii, 176
 pulchella, ii, 171
 pyriformis, ii, 159
 quercifolia, ii, 178
 repens, ii, 178
 robur, ii, 176
 salicifolia, ii, 176
 serrata, ii, 177
 serratifolia, ii, 177
 speciosa, ii, 13, 178
 sphaerocarpa, ii, 171
 spicata, ii, 174
 spinulosa, ii, 172
 tenuifolia, ii, 152
 teretifolia, ii, 150
 verticillata, ii, 175
Baptisia, ii, 422
 alba, ii, 423
 australis, ii, 423
 perfoliata, ii, 423
Barbarea, ii, 398
 præcox, ii, 398
 tinctoria, ii, 423
 vulgaris, ii, 398
Barbula leucostoma, i, 238, 253
 ruralis, i, 239
 unguiculata, i, 69
Barleria brevispina, i, 94
 eranthemoides, i, 93
 grandiflora, i, 94
Barleria macracantha, i, 94
 mollis, i, 94
 parviflora, i, 94
Barraldeia, i, 119 *note*
 pomiformis, i, 69
Bartholinia, ii, 478
Bartramia, ii, 335
 pectinata, ii, 478
Bassia, i, 296
 rufescens, i, 294
Basteria aculeata, ii, 463
Bauera, i, 20
Bauhinia, i, 25
Beaufortia, i, 19, 311; ii, 457
 decussata, ii, 457
 sparsa, ii, 457
Beckmannia erucæformis, ii, 515
Begonia, ii, 279 *note*
Begoniaceæ, i, 149
Belis, i, 461
Bellenda, i, 131 *note*
Bellendena, ii, 17, 18, 23, 37
Belvisæ, i, 388 *note*
Belvisia, i, 387 and *note*, 388 *note*
Bennettia, ii, 661
 affinis, ii, 665
 Finlaysoni, ii, 665
 Jackiana, ii, 665
 javanica, ii, 661
 pedicellata, ii, 665
 phlebocarpa, ii, 665
 Wallichii, ii, 664
Bennettiaceæ, ii, 663
Berberidæ, i, 124 *note*, 381
Berberides, i, 65, 124, 361
Berberis, i, 124 *note*
Berckheya, ii, 462
 cernua, ii, 463
 ciliaris, ii, 462
 cynaroides, ii, 462
 incana, ii, 463
 obovata, ii, 463
 setosa, ii, 462
 squarrosa, ii, 462
Berteroia, i, 271
Besleriaceæ, ii, 586
Besleriæ, ii, 590
Bidens arborea, ii, 292 *note*
 hirsuta, ii, 291
 nivea, ii, 290, 295
 scandens, ii, 290
Bignonia cærulea, ii, 674
 Copaia, ii, 674
 discolor, i, 94
 ilicifolia, ii, 674

- Bignonia procera*, ii, 674
Bignoniaceæ, ii, 584
Billardiera, i, 13
Billottia, i, 311
 flexuosa, i, 311 *note*
 linearifolia, i, 311 *note*
 marginata, i, 311 *note*
Biscutella, i, 264; ii, 373
 apula, ii, 373
 auriculata, ii, 373
 coronopifolia, ii, 373
 lævigata, ii, 374
 sempervirens, ii, 374
Bixineæ, ii, 716
Blackwellia, i, 120
Bladhia, i, 150 *note*
Blechnum, i, 164; ii, 506
 Brownei, ii, 506
Blennodia, i, 316
 canescens, i, 316
Blepharis, i, 135
Bletia, ii, 694
 capitata, ii, 488
 florida, ii, 487
 hyacinthina, ii, 487
 Tankervilleæ, i, 491, 498, 512;
 ii, 487
 verecunda, ii, 487
Blighia sapida, i, 161 *note*
Blumea diffusa, ii, 711
Bocconia, i, 270 *note*; ii, 271
 cordata, i, 270 *note*; ii, 271
 frutescens, i, 270 *note*; ii, 271
Bœomyces paschalis, i, 252
 vermicularis, i, 252
Boerhavia mutabilis, i, 164, 167
Boletus igniarius, i, 70
Bombacæ, ii, 620
Bombax pentandrum, i, 109, 153, 166
Bonatea speciosa, i, 503, 537
Bontia, i, 40; ii, 506
 daphnoides, ii, 506
Boopideæ, ii, 312
Boopis, ii, 309, 312
 balsamitifolia, ii, 317
Boraginæ, i, 297; ii, 659
Boronia, i, 17
 affinis, ii, 709
 albiflora, ii, 709
Borreria aurantiaca, i, 250, 253
 flavicans, i, 250
Bossiaea, i, 25; ii, 437
 cincree, ii, 438
 heterophylla, ii, 437
 lanceolata, ii, 438
 linophylla, ii, 438
Bossiaea microphylla, ii, 438
 prostrata, ii, 438
 rufa, ii, 437
 Scolopendrium, ii, 437
Botria, i, 151
Botrychium Lunaria, i, 67
Brabeium stellatum, ii, 16, 18, 36, 41,
 137, 138
Brabyla, ii, 138
Bracheilema paniculatum, i, 95
Brachychilum, ii, 726
 Horsfieldii, ii, 726
Brachychiton, i, 315; ii, 637
 diversifolium, ii, 638
 incanum, ii, 638
 platanoides, ii, 638
 populneum, ii, 638
 ramiflorum, ii, 637
Brachylæna, ii, 294 *note*
Brachysema, ii, 427
 latifolium, ii, 427
 uniflorum, ii, 709
Brachystelma, ii, 675
 tuberosa, ii, 675
Brachystemum, ii, 503
 virginicum, ii, 504
Bragantia, i, 385
 Wallichii, ii, 534
Brassavola, ii, 495
 cucullata, ii, 495
Brassia, ii, 494
 maculata, ii, 494
Brassica, ii, 408, 494
 alpina, ii, 410
 arvensis, ii, 398
 campestris, ii, 410
 chinensis, ii, 409
 elongata, ii, 409
 Eruca, ii, 409
 Erucastrum, ii, 409
 monensis, ii, 409
 muralis, ii, 412
 Napus, ii, 408
 oleracea, ii, 408
 orientalis, ii, 404
 Rapa, ii, 408
Braya, i, 192, 193
Breweria evolvuloides, i, 94
Bridelia, i, 127
Bromelia Ananas, i, 154
Bromeliaceæ, i, 413 *note*
Bromus purgans, ii, 515
Brosimum, i, 79
Broughtonia, ii, 496
 sanguinea, ii, 496
Brucea, ii, 537

- Brugmannsia*, i, 402, 427, 558
 Zippelii, i, 427
Bruguiera, i, 20, 119, 364
Brunia, i, 64; ii, 322
 Levisanus, ii, 45
Bruniaceæ, ii, 322
Brunonia, i, 31; ii, 267 *note*, 269
 note, 310, 357
 australis, ii, 310
 sericea, ii, 310
Bryum, i, 178; ii, 335
 argenteum, i, 69
 cæspititium, i, 277
 calophyllum, i, 235
 capillaceum, i, 238
 capillare, i, 69
 macrocarpum, i, 353, 354; ii, 335
 tenuifolium, 8, 238
 rostratum, i, 235
Buchanania, i, 112
Buchnera orobanchoides, i, 94
Bucklandia populnea, ii, 534
Buddlea acuminata, i, 93
 foliata, i, 93
Bulbocodium, i, 298
Bunias, ii, 372
 ægyptiaca, ii, 372
 balearica, ii, 375
 Cakile, ii, 369
 Erucago, ii, 372
 myagroides, ii, 408
 orientalis, ii, 372
 syriaca, ii, 371
Burchellia, ii, 687
 Capensis, ii, 687
Burmannia, i, 53
Bursaria, i, 13, 132
Bursera, i, 112
Burtonia, ii, 428
 scabra, ii, 428
Butomæ, i, 379 *note*, 555
Butter and tallow tree, i, 161, 295
Buttneriaceæ, i, 11, 26, 63, 109; ii, 614, 646, 672
Buxbaumia, i, 348, 351; ii, 349
 aphylla, i, 351; ii, 350

Cabomba, i, 451
Cacalia abyssinica, i, 95
Cachio, i, 140 *note*
Cactus quadrangularis, i, 128
Cadaba dubia, i, 276
 farinosa, i, 276
Cænopteris, i, 59
Cæsalpineæ, i, 110, 116, 294

Cæaulia, ii, 274
 radicans, ii, 284
Cakile, i, 263; ii, 369
 maritima, ii, 369
 perennis, ii, 369
 rugosa, ii, 369
Caladenia, 485
 alba, ii, 485
Calamagrostis Canadensis, ii, 514
 purpurascens, ii, 514
 stricta, ii, 514
Calamus secundiflorus, i, 142
Calanchoë pubescens, i, 94
Calanthe, ii, 694
Caldasia, i, 389
Calea, ii, 287
 aculeata, ii, 303, 305, 306
 affinis, ii, 305
 Amellus, ii, 287, 290
 arcuata, ii, 306
 aspera, ii, 295, 306, 307
 aurea, ii, 305
 cordata, ii, 306
 corditolia, ii, 287, 306
 denticulata, ii, 305
 Jamaicensis, ii, 287, 295, 299
 lævis, ii, 305
 leptophylla, ii, 300, 303, 304
 lobata, ii, 298
 longifolia, ii, 305
 oppositifolia, ii, 287, 290
 pinifolia, ii, 300, 302
 quinquefaria, ii, 306
 scoparia, 293, 306
 spectabilis, ii, 304, 306
Caleacte, ii, 289
 pinnatifida, ii, 289
 urticifolia, ii, 289
Caleana, ii, 486
Calectasia, i, 52, 84, 436
 cyanea, i, 85
Caleya, ii, 486
 major, ii, 486
Calicium claviculare, i, 69
Calladium, i, 365
Callistachys elliptica, ii, 426
Callistemon, i, 19
 rigidum, ii, 683
Callitriche, i, 22
Callitris, i, 48, 308, 339
 australis, ii, 706
 calcarata, ii, 706
 fruticosa, ii, 707
 glauca, ii, 706
 propinqua, ii, 706
 rhombocidea, ii, 707

- Callitris robusta*, ii, 706
 tuberculata, ii, 706
 Ventenatii, ii, 706
 verrucosa, ii, 706
Calocephalus, ii, 286
Calopogon, ii, 486
 pulchellus, ii, 486
Calosacme, ii, 599
Calothamnus, i, 19, 311; ii, 456
 gracilis, ii, 457
 quadrifida, ii, 457
 villosa, ii, 457
Calothyrsus, ii, 147
Calotis, ii, 688
 cuneifolia, ii, 688
 dentex, ii, 688
 pterosperra, ii, 711
 radicans, i, 190
Calotropis, ii, 217
 gigantea, ii, 218
 procera, ii, 218
Caltha arctica, i, 190
 natans, i, 190
Calvaria, i, 296
Calycera, ii, 309
Calycereæ, ii, 309, 312
Calycomis, i, 20
Calymperes, ii, 343, 345
Calypso, ii, 488
 Americana, ii, 489
 borealis, ii, 489
Calystegia sepium, i, 68, 296
Calythrix, i, 311; ii, 683
 decandra, ii, 710
 glabra, ii, 684
 laricina, ii, 710
Camelina, ii, 386
 austriaca, ii, 386
 sativa, ii, 386
Cameraria, ii, 247
Campanula uniflora, i, 212
Campanulacæ, i, 31, 212; ii, 269
 note
Campulosus, i, 58
Candollea, i, 12
 fasciculata, ii, 704
 pedunculata, ii, 704
Canella, i, 381
Canna Indica, i, 158
Canneæ, i, 49; ii, 550
Canscora diffusa, ii, 533
Cantharellus lobatus, i, 253
Canthium attenuatum, ii, 711
 lucidum, i, 93
Capparidæ, i, 272, 282
Capparis, i, 273
 lasiantha, ii, 705
 lucida, ii, 708
 sodada, i, 279
 trifoliata, i, 277
 umbellata, ii, 705
Capraria, ii, 504
 biflora, ii, 504
Caprifoliacæ, ii, 324, 541, 673
Caprifolium, ii, 541
Capsella bursa-pastoris, i, 263, 271
Capsicum, i, 155, 156, 158
 frutescens, i, 153
Carallia, i, 20, 119
Caralluma, ii, 200, 208
 adscendens, ii, 209
Cardamine, ii, 392
 Africana, ii, 393
 amara, ii, 394
 articulata, i, 197
 asarifolia, ii, 393
 bellidifolia, i, 198
 bulbifera, ii, 393
 chelidonia, ii, 393
 enneaphylla, ii, 392
 græca, ii, 393
 hastulata, ii, 396
 hirsuta, ii, 394
 impatiens, ii, 394
 latifolia, ii, 394
 nivalis, ii, 397
 nudicaulis, i, 197
 pentaphylla, ii, 392
 pinnata, ii, 392
 pratensis, ii, 394
 resedifolia, ii, 394
 trifolia, ii, 393
Cardiopteridæ, ii, 659
Cardiopteris, ii, 655
 Javanica, ii, 655
 lobata, ii, 655
 moluccana, ii, 655, 657
Cardiospermum grandiflorum, i, 107,
 165, 172
Carduacæ, ii, 270
Carex, i, 53, 143, 145
 affinis, ii, 516
 ampullacea, ii, 517
 aristata, ii, 517
 attenuata, ii, 516
 bicolor, ii, 516
 bullata, ii, 517
 cæspitosa, i, 68, 255
 capillaris, ii, 517
 compacta, i, 178; ii, 517

- Carex concinna*, ii, 516
 concolor, i, 218, 255
 dioica, ii, 515
 filifolia, ii, 516
 fuliginosa, i, 218, 255; ii, 516
 lacustris, ii, 517
 limosa, ii, 517
 lioliacea, ii, 516
 lupulina, ii, 517
 marginata, ii, 517
 media, ii, 516
 misandra, i, 217, 255
 mutica, ii, 517
 oligocarpa, ii, 517
 pellita, ii, 517
 podocarpa, ii, 517
 polytrichoides, ii, 516
 pseudo-cyperus, i, 68
 pulla, i, 178
 remota, ii, 516
 Richardsonii, ii, 516
 saxatilis, ii, 517
 scirpoides, ii, 515
 scoparia, ii, 516
 varia, ii, 517
 Wormskioldiana, ii, 515
Cargillia australis, i, 162 *note*
Carica papaya, i, 154, 158
Carissa abyssinica, i, 94
 edulis, i, 94
 mitis, ii, 236
Cartonema, i, 53
Carmichaelia, ii, 698
 australis, ii, 698
Carphephorus, ii, 299 *note*
Carpodinus, ii, 715
Carpolithus Smithii, ii, 723
Cartodium, ii, 286
Caryophyllæ, i, 65, 198, 285; ii, 587
 note
Cassa, i, 111, 153
Cassava, i, 154, 155
Cassia, i, 24, 153, 339
 artemisiæfolia, i, 326
 canaliculata, i, 326
 circinata, i, 327
 eremophila, i, 326
 glaucescens, i, 326
 linearis, i, 327
 obovata, i, 294
 occidentalis, i, 111, 163, 165,
 167
 phyllodinea, i, 326
 platypoda, i, 326
 pubescens, i, 94
Cassia spectabilis, ii, 306
 Sturtii, i, 325
 zygophylla, i, 326
Cassinia, ii, 304, 470
 aculeata, ii, 306
 aurea, ii, 283, 470
 leptophylla, ii, 304
Cassipourea, i, 164
Casuarina, i, 45, 308, 339, 549
 equisetifolia, i, 45
 nodiflora, i, 45
 quadrivalvis, i, 45
Casuarinæ, i, 45
Cassuvie, i, 112
Cassytha, i, 149
 pubescens, i, 150, 165
Catabrosa, i, 220, 228
Catesbæa, ii, 268
Catharinea glabrata, i, 234
 lævigata, i, 234
Caulinia oceanica, i, 68
 ovalis, ii, 536
Caulophyllum, i, 124 *note*
 thalictroides, i, 361
Causjera, ii, 536
Ceanothus, i, 26
Cecropia, i, 138
Cedreleæ, i, 72, 149
Celastrinæ, i, 114
Celastrinea f., ii, 536
Celastrinea, i, 13, 27, 107
Celastrus glaucus, i, 94
 serrulatus, i, 94
Celosia albida, i, 165
 argentea, i, 165
 margaritacea, i, 165
Celtis, i, 138; ii, 279
 orientalis, i, 165
Cenarrhenes, ii, 37, 132
 nitida, ii, 132
Cenchrus, i, 57
 tripsacoides, i, 93
Cenomyce coccifera, i, 69
 cornuta, i, 69
 deformis, i, 69
 fimbriata, i, 178
 pyxidata, i, 69, 252
 rangiferina, i, 6, 9, 178
 vermicularis, i, 69, 252
Centotheca lappacea, i, 165
Centronia, ii, 595, 609
 mirabilis, ii, 609
Centrophorum, i, 302
Centropodia, i, 302
Cephalotæ, ii, 357

Cephalotus, i, 76 ; ii, 353
 follicularis, i, 76, 440
 Cerania vermicularis, i, 251
 Ceranthera dentata, i, 123
 Cerastium alpinum, i, 178, 198
 latifolium, i, 198
 Ceratopetalum, i, 20, 120
 Ceratophyllum, ii, 155
 compressum, ii, 152
 demersum, ii, 723
 Ceratopteris, ii, 570
 Cerbera, i, 133
 Cercis, i, 24 ; ii, 436
 siliquastrum, i, 25 ; ii, 436
 Ceropegia, i, 549 ; ii, 200, 205, 227, 696
 acuminata, ii, 205
 Africana, ii, 696
 biflora, ii, 205
 bulbosa, ii, 205
 Candelabrum, ii, 205
 juncea, ii, 205
 sagittata, ii, 228
 tenuiflora, ii, 228
 tuberosa, ii, 205
 Cestrum, ii, 267
 Ceterach alpinum, ii, 254
 Cetraria cucullata, i, 251
 islandica, i, 178, 251
 juniperina, i, 250
 nivalis, i, 178, 250
 odontella, i, 251
 Chailletea, i, 125
 Chailletia, i, 125, 164
 sessiliflora, i, 126
 Chailletiacæ, ii, 535
 Chamærops humilis, i, 52
 Chamissoa albida, ii, 534
 Chasme, ii, 40
 Cheilanthes leptophylla, i, 95
 Cheiranthus, ii, 404
 albus, ii, 405
 alpinus, ii, 403
 annuus, ii, 406
 Cheiri, i, 379 note ; ii, 404
 coccineus, ii, 405
 erysimoides, ii, 403
 Farsertia, ii, 388
 fenestralis, ii, 406
 glabrus, ii, 406
 helveticus, ii, 404
 incauus, ii, 405
 littoreus, ii, 407
 longifolius, ii, 404
 maritimus, ii, 407
 mutabilis, ii, 404

Cheiranthus odoratissimus, ii, 406
 parviflorus, ii, 406
 quadrangulus, ii, 403
 sinuatus, ii, 406
 tenuifolius, ii, 404
 tricuspidatus, ii, 406
 tristis, ii, 406
 Chenopodeæ, i, 138
 Chenopodiaceæ, ii, 535
 Chichæa aserifolia, ii, 628
 Chilli, i, 158
 Chiloglottis, ii, 549, 575
 serpyllifolia, ii, 575
 Chilurus, ii, 129
 Chionachne, ii, 573, 574
 Chirita, ii, 585, 591, 594, 598
 acuminata, ii, 599
 bifolia, ii, 599
 cærulea, ii, 599
 dimidiata, ii, 599
 flava, ii, 599
 grandiflora, ii, 598
 hamosa, ii, 599
 Horsfieldii, ii, 599
 macrophylla, ii, 599
 pumila, ii, 599
 scaberrima, ii, 599
 urticifolia, ii, 598
 Chlenaceæ, i, 11, 108, 109, 110
 Chloanthes, i, 39
 Chloranthæ, i, 215
 Chloranthus, ii, 671
 monostachys, ii, 671
 Chondrachne, i, 145
 Chorispermum, ii, 413, 705
 tenellum, ii, 413
 Chorispore, ii, 705
 Chorizandra, i, 145
 Chorizema, ii, 425
 Henchmanii, ii, 699
 ilicifolia, ii, 425
 ilicifolium, ii, 425
 nana, ii, 425
 rhombea, ii, 425
 trilobatum, ii, 426
 Chrysanthemum Indicum, ii, 691
 Chrysobalanæ, i, 115 ; ii, 537
 Chrysobalanus, i, 115
 Icaco, i, 115, 153, 155, 167
 Chrysocoma, ii, 293
 cinerea, ii, 302
 Chrysosplenium alternifolium, i, 205
 Chuquiraga, i, 81 ; ii, 269
 Chuncoa, i, 20
 Cichoraceæ, i, 30 ; ii, 263, 269

- Cinclidotus*, ii, 335
Cineraria congesta, i, 211, 255
 palustris, i, 211, 255
Cinerocephala, i, 30; ii, 298
Cissampelos, i, 381
 nymphaeifolia, i, 95
Cissus, i, 151
 angustifolia, i, 394 *note*
Cistinea, i, 65, 124, 282, 288, 453
 and note, 532
Cistus, i, 453 *note*
Citrus aurantium, i, 154
 medica, i, 154
Cladium Mariscus, i, 68
Cladonia subuliformis, i, 252
Clarekia pulchella, i, 466
Clavaria coralloides, i, 70
 pistillaris, i, 70
Clavija, i, 151 *note*
Clematis aristata, ii, 703
 stenopetala, ii, 703
Clome, ii, 413
 arabica, i, 273, 275; ii, 415
 arborea, i, 278
 Chelidonii, ii, 413
 deflexa, i, 276
 Erucago, ii, 414
 gigantea, ii, 414
 Houstoni, ii, 414
 monophylla, ii, 415
 ornithopodioides, ii, 414
 pentaphylla, i, 273; ii, 414
 procumbens, ii, 415
 Siliquaria, i, 94
 spinosa, ii, 414
 violacea, i, 276; ii, 414
 viridiflora, ii, 414
 viscosa, ii, 413
Clerodendron, i, 135; ii, 507
 fragrans, ii, 507
 incerne, ii, 508
 ligustrium, ii, 508
 myricoides, i, 94
 paniculatum, ii, 507
 pyramidale, ii, 507
 Siphonanthus, ii, 508
 squamatum, ii, 507
 tomentosum, ii, 508
 viscosum, ii, 507
Clanthus Dampieri, i, 320, 330, 339
 Oxleyi, i, 320, 321
Clidantha, i, 322
 psoraloides, i, 322
Cliffortia, i, 64
Clinopodium incanum, ii, 503
Clypeola, ii, 374
 Ionthlaspi, ii, 374
Cnestis, i, 112, 113, 294
Cochlearia, i, 271; ii, 383
 anglica, i, 178; ii, 383
 Armoracia, i, 379 *note*; ii, 384
 Coronopus, ii, 373
 danica, i, 178; ii, 383
 Draba, ii, 381
 fencstrata, i, 178, 191
 glastifolia, ii, 383
 officinalis, ii, 383
 saxatilis, ii, 384
Cocoa-nut, i, 158, 161
Cocoinae, i, 141
Cocos indica, i, 141
Codarium acutifolium, i, 161
 nitidum, i, 161 *note*
Codia, i, 20
Cœlorachis, ii, 553
Coffea, ii, 269 *note*
Coix arundinacea, ii, 573
 heteroclita, ii, 574
Cola, i, 153; ii, 642
 acuminata, ii, 642
Colchicum fasciculare, i, 300
 Ritchii, i, 298
Coleanthus, i, 220
Collema fasciculare, i, 70
 nigrum, i, 70
 tremelloides, i, 70
Colletia, i, 26
Collomia, i, 549
Colpodium, i, 210, 221
 compressum, i, 221
 latifolium, i, 221, 255
 Stevenii, i, 221
Coluria, i, 206
Colutea, ii, 441
 aperta, ii, 441
 arborescens, ii, 441
 cruenta, ii, 441
 frutescens, ii, 442
 galegifolia, ii, 442
 herbacca, ii, 442
 Novæ Hollandiæ, i, 320
 perenuans, ii, 443
 Pocockii, ii, 441
Combretaceæ, i, 19; ii, 663
Combretum molle, i, 94
 ovale, i, 94
Comesperma, i, 15
 aphyllum, ii, 708
 laucolatum, ii, 708
Cometes, ii, 543

- Cometes abyssinica*, ii, 546
 alterniflora, ii, 545
 Surattensis, ii, 544
Commelina, i, 298, 299 *note*; 513
 acuminata, i, 93
 hirsuta, i, 93
Commelineæ, i, 53
Commersonia, i, 11; ii, 672
Compositæ, i, 30, 62, 65, 81, 104, 128,
 210, 295, 308, 448; ii, 257, 571
Conchium, ii, 20, 149
 aciculare, ii, 152
 cornutum, ii, 152
 dactyloides, ii, 22, 157
 drupaceum, ii, 157
 ellipticum, ii, 157
 gibbosum, ii, 152
 longifolium, ii, 150
 nervosum, ii, 157
 oleifolium, ii, 155
 pugioniforme, ii, 150
 salicifolium, ii, 156
 salignum, ii, 156
 sphæroideum, ii, 152
 teretifolium, ii, 150
 trifurcatum, ii, 153
Conferva ebenea, i, 70
 ericetorum, i, 70
Coniferæ, i, 44, 46, 138, 312, 453,
 506 *note*, 509; ii, 720, 721
Connaracæ, i, 112
Conarus, i, 112, 294
Conocarpodendron, ii, 26, 40, 78, 380
 note, 413 *note*
Conocarpus, i, 164; ii, 40
Conohoria, i, 123
Conospermum, i, 41, 82, 381; ii, 11,
 15, 17, 18, 21, 37, 127
 cæruleum, ii, 129
 capitatum, ii, 129
 distichum, ii, 129
 ericifolium, ii, 128
 longifolium, ii, 128
 reticulatum, ii, 130
 taxifolium, ii, 128
 teuuiifolium, ii, 128
 teretifolium, ii, 129
Contortæ, i, 384; ii, 269
Convolvulacæ, i, 81, 104, 130, 168,
 296; ii, 545
Convolvulus, i, 135; ii, 545
 Batatas, i, 161
 Brasiliensis, i, 164
 cirrhosus, i, 94
 congestus, i, 94
Convolvulus pes-capræ, i, 164
 pilosus, i, 94
 repens, i, 296
Conyza, ii, 294, 298
 bifrons, ii, 294
 squarrosa, ii, 294
Corallorrhiza, ii, 489
 innata, ii, 490
Corchorus, i, 108
 gracilis, i, 94
 Pumilio, ii, 709
Cordia abyssinica, i, 94
 ovalis, i, 94
Cordylocarpus lavigatus, ii, 408
Coreopsis tripteris, ii, 263, 297
Corethrum, i, 42
Cornicularia bicolor, i, 178
 lanata, i, 70, 251
 ochroleuca, i, 251
 pubescens, i, 70
 spadicea, i, 70
Cornulaca monacantha, i, 297
Cornus, ii, 322
Coronilla aculeata, ii, 443
 grandiflora, ii, 443
 Sesban, ii, 443
Coronopus, ii, 372
 didyma, ii, 373
 Ruellii, ii, 373
Correa, i, 17
Corymbifera, i, 30; ii, 270
Corymbium, ii, 269
Corypha, i, 141, 161
 Taliera, i, 142
 umbraculifera, i, 142
Corysanthera, ii, 609
Corysanthes, i, 85
 fimbriata, i, 86
Costus, ii, 550
Coupea, i, 115
Courtenia, ii, 620, 641
 Afzelii, ii, 641
 heterophylla, ii, 641
 triloba, ii, 641
Coussapoa, i, 138
Crambe, ii, 369
 filiformis, ii, 371
 fruticosa, ii, 370
 hispanica, ii, 370
 maritima, ii, 370
 orientalis, ii, 370
 pinnatifida, ii, 370
 strigosa, ii, 371
 suecica, ii, 370
 tatarica, ii, 370

- Craspedia*, ii, 285
 uniflora, ii, 286
Crassula puberula, i, 94
Crassulacæ, ii, 318, 356
Crateva, i, 273
 acuminata, i, 278
 Adansonii, i, 277
 fragrans, i, 277
 gynandra, i, 278
 læta, i, 277
 Roxburghii, i, 277
 Tapia, i, 278
Cream fruit, i, 153, 155
Cremolobæ, i, 264
Cremolobus, i, 264
Crinum, i, 364
Crossandra, ii, 505
 undulæfolia, ii, 505
Crotalaria axillaris, i, 165
 Cunninghamii, i, 320
 farceta, i, 95
 trifoliastrum, ii, 533
 propinqua, i, 95
 Saltiana, i, 95
 Sturtii, i, 319, 325
Croton acuminatum, i, 95
Crowea dentata, ii, 709
Cruciferae, i, 65, 182 *note*, 186, 261,
 282, 294, 503, 559; ii, 271,
 367
Cryphiospermum, ii, 284
 repens, ii, 284
Cryptandra, i, 26
 pyramidalis, ii, 708
Cryptarrhena, ii, 683
 lunata, ii, 683
Cryptodon, i, 247
Cryptogramma, ii, 525
 acrostichoides, ii, 519, 525
Cryptolepis, ii, 233, 240, 687
Cryptostegia, ii, 686
 grandiflora, ii, 686
Cryptostemma, ii, 465
 calendulaceum, ii, 465
 hypochondriacum, ii, 465
 runcinatum, ii, 465
Cucifera, i, 141
Cucurbita Pepo, i, 154
Cucurbitacæ, i, 121, 384, 388, 444
Culhamia, ii, 616
Cullumia, ii, 462
 ciliaris, ii, 462
 setosa, ii, 462
 squarrosa, ii, 462
Cunninghamia, i, 458; ii, 707
Cunonia, i, 20
Cunoniacæ, i, 20, 21, 120; ii, 523
Cuphea, ii, 238
Cupressina, i, 458
Curatella, i, 12
Curculigo, i, 51
Cusparia, i, 16
Cyanorchis, ii, 694
Cyathæa, ii, 546
Cycadæ, i, 149
Cycadites, ii, 380 *note*, 453, 717
 microphyllus, ii, 719
 Saxhyanus, ii, 724
Cycadoidea, ii, 716, 717
 megalophylla, ii, 717
 microphylla, ii, 717
Cycas, i, 458; ii, 717
Cyclophorus, ii, 566, 568
Cyclopia, ii, 422
 genistoides, ii, 422
Cyclopterae, ii, 11, 20, 148
Cylindria, ii, 190
Cymbidium, i, 50; ii, 493
 aculcatum, ii, 694
 aloefolium, ii, 493
 Andersonii, ii, 495
 coccineum, ii, 490
 Corallorhizon, ii, 490
 cucullatum, ii, 495
 ensifolium, ii, 493
 floridum, ii, 487
 giganteum, ii, 694
 hyacinthinum, ii, 488
 lineare, ii, 490
 pedicellatum, ii, 694
 præmorsum, ii, 493
 proliferum, ii, 490
 pulchellum, ii, 486
 sinense, ii, 493
 tesselatum, ii, 689
 tripterum, ii, 494
 triquetrum, ii, 495
 verecundum, ii, 487
Cynanchum, ii, 6, 221
 acutum, ii, 221
 bicolor, ii, 226
 Capense, ii, 223
 Carolinense, ii, 215
 Chinense, ii, 221
 crassifolium, ii, 223
 crispiflorum, ii, 215
 erectum, ii, 212
 crubescens, ii, 222
 extensum, ii, 226
 floribundum, ii, 222

Cynanchum grandiflorum, ii, 215

hirtum, ii, 215

maritimum, ii, 215

medium, ii, 224

Mouspeliacum, ii, 221

nigrum, i, 215, 224, 529

obliquum, ii, 215

obtusifolium, ii, 223

parviflorum, ii, 227

pauciflorum, ii, 222

pedunculatum, ii, 222

pilosum, ii, 223

planiflorum, ii, 215

prostratum, ii, 215

pyrotechnicum, ii, 226

racemosum, ii, 215

roseum, ii, 223

rostratum, ii, 215

Sibiricum, ii, 224

suberosum, ii, 215

undulatum, ii, 215

viminale, ii, 226

Vincetoxicum, ii, 224

Cynodon Dactylon, i, 65, 68

Cynodontium capillaceum, i, 238

Cynomorium, i, 389, 391

cayancense, i, 389 note

coccineum, i, 412

jamaicense, i, 389 note

Cynontodium capillaceum, i, 69, 238

Cynosurus eruceformis, ii, 515

Cyperaceæ, i, 53, 54, 62, 85, 104,

142, 217, 298, 300

Cyperus, i, 143, 300

articulatus, i, 144, 162, 164, 166

densus, i, 93

involutus, i, 93

laxiflorus, i, 144

laxus, i, 93

ligularis, i, 144, 165

melanocephalus, i, 93

niloticus, i, 164

Papyrus, i, 143, 162, 166, 300

rotundus, i, 66, 68

scirpoides, i, 93

Cypha, i, 32, 34

Cypripedium, i, 48, 503; ii, 489, 549, 551

acaule, ii, 499

album, ii, 499

arietinum, ii, 499

Calceolus, ii, 498

flavescens, ii, 499

humile, ii, 499

parviflorum, ii, 498

Cypripedium pubescens, ii, 498

spectabile, i, 508; ii, 499

Cyrtandra, i, 558; ii, 585, 596, 609

C? lanuginosa, ii, 534

Cyrtandraceæ, i, 558; ii, 537, 577, 584

Cyrtandree, ii, 585, 590

Cyrtodon, i, 241

Cyrtolobus, ii, 533

Cyrtopodium, ii, 495, 567, 670

Andersonii, ii, 495, 670

Woodfordii, ii, 670

Cytineæ, i, 428; ii, 358

Cytinus, i, 385, 389, 391, 396, 401, 428

americanus, i, 429

dioicus, i, 429

Hypocistis, i, 428

Cytisus Cajan, i, 154, 159

Dacrydium, i, 44, 47, 455, 458, 460

Dahlia, ii, 321

Dalechampia, ii, 544

tripartita, i, 95

Dammara, i, 458

Dampiera, i, 315; ii, 271

Danais, i, 131

Danthonia, i, 301

Forskali, i, 301

Dasypogon, i, 52, 83, 436

bromeliifolius, i, 84

Datisca, i, 285

Datisceæ, i, 285

Datura, ii, 267

Daviesia, ii, 435

concinna, ii, 709

corymbosa, ii, 435

denudata, ii, 429

latifolia, ii, 435

mimosoides, ii, 435

teretifolia, ii, 709

ulicifolia, ii, 435

ulicina, ii, 435

Dawsonia, i, 348; ii, 331, 335, 340

polytrichoides, i, 349

Deidamia, i, 386, 387 note

Delabechia, i, 315

Dendrobium, i, 49; ii, 492

Barringtoniæ, ii, 493

cucullatum, ii, 692

graminifolium, ii, 492

linguiforme, ii, 492

moniliforme, i, 49

Pierardi, ii, 692

ruscifolium, ii, 491

sanguineum, ii, 496

speciosum, ii, 492

- Dentaria*, ii, 392
 bulbifera, ii, 393
 enncaphylla, ii, 392
 heptaphyllos, ii, 393
 pentaphylla, ii, 392
 pinnata, ii, 393
Deschampsia, i, 221, 228, 231
 brevifolia, i, 230
Desmochæta, ii, 543, 544
 lappacea, i, 165, 167
 staminea, ii, 532
Deutzia, ii, 550
 corymbosa, ii, 532
Dialalpe, ii, 546
Dianella, i, 50
Dianthus, i, 285
 abyssinicus, i, 94
Diaspasis, i, 32
Dichapetalum, i, 125
Dicksonia antarctica, i, 59
Dicoryphe, ii, 321
Dicranum, ii, 345
 falcifolium, ii, 335
 flexuosum, i, 69
 pulvinatum, i, 69
 purpureum, i, 69
 scoparium, i, 69, 178
Dictamnus, i, 16
Didelta, ii, 463
 caruosum, ii, 463
 tetragonæfolium, ii, 463
 spinosum, ii, 463
Didymocarpus, i, 532, 585, 590, 594, 599
 acuminata, ii, 602
 aromatica, ii, 601, 602
 barbata, ii, 599
 corchorifolia, ii, 603
 cordata, ii, 603
 corniculata, ii, 613
 crinita, ii, 602
 elongata, ii, 603
 frutescens, ii, 603
 hamosa, ii, 599
 helicteroides, ii, 605
 lanuginosa, ii, 603
 macrophylla, ii, 602
 missionis, ii, 603
 multiflora, ii, 605
 oblonga, ii, 601
 obtusa, ii, 602
 pedicellata, ii, 602
 plicata, ii, 602
 primulifolia, ii, 601
 Punduana, ii, 601
Didymocarpus racemosa, ii, 603
 reptans, ii, 603
 Rexii, ii, 604
 Rotleriana, ii, 603
 serrata, ii, 603
 subalternans, ii, 602
 verticillata, ii, 601
 villosa, ii, 601
 Zeylanica, ii, 603
Didymochlæna, ii, 569
Didymodon, ii, 348
 capillaceum, i, 238
 latifolium, ii, 348
 subulatum, i, 238
Diervilla, ii, 541
Digera, ii, 543
Dillenia, i, 12, 381
Dilleniaceæ, i, 12, 63, 64, 293
Dillwynia, ii, 430
 cinerascens, ii, 673
 crucifolia, ii, 431
 floribunda, ii, 430
 glaberrima, ii, 431
 myrtifolia, ii, 431
 obovata, ii, 431
 parvifolia, ii, 669
Diomedea, ii, 297 *note*
Dioscorea, i, 154, 160; ii, 655, 657
Diosmeæ, i, 16, 63, 72, 114, 285
Diphylleia, i, 124 *note*
Diplachne Baueri, ii, 707
Diplacrum, i, 54, 145
Diplanthera, ii, 531
Diplasia, i, 145
Diplazium, i, 60
Diplolæna, i, 17
Diplolepis, ii, 219
Diplomeris, i, 504
Dipodium, ii, 694
Dipsacæ, i, 65; ii, 277, 309
Dipsacus, ii, 277
Dipteris, ii, 562, 566
 conjugata, ii, 562
Disa, ii, 480, 694
 cornuta, ii, 480
 spatulata, ii, 480
Dischidia, ii, 213, 357
Ditassa, ii, 225
Diuris, ii, 482, 549
 aurea, ii, 483
Dobera glabra, i, 93
Dœmia, i, 295; ii, 225
Dombeya Erythroxylon, ii, 509
Donia, i, 177, 321; ii, 282, 461
 glutinosa, ii, 461

- Donia speciosa*, i, 320
Doom, i, 141
Doreoceras hygrometrica, ii, 605
Doronicum glutinosum, ii, 461
Doubah, i, 329
Doura, i, 289
Draba, i, 268, 271; ii, 384
 aizoides, ii, 385
 alpina, i, 190
 androsacea, i, 191
 ciliaris, ii, 385
 corymbosa, i, 178
 hirta, ii, 385
 incana, ii, 385
 lapponica, i, 191
 muralis, ii, 385
 muricella, i, 178
 nemorulis, ii, 385
 oblongata, i, 178
 pauciflora, i, 190, 253
 Pumilio, ii, 705
 pyrenaica, ii, 386
 rupestris, i, 178; ii, 384
 verna, ii, 384
Dracophyllum verticillatum, i, 38
Drapetes, i, 82, 83
Drepanophyllum, ii, 335
Drosera, i, 65
 Banksii, ii, 705
 petiolaris, ii, 705
 Menziesii, ii, 705
Droseraceæ, i, 282
Dryandra, i, 116, 450, 455; ii, 38, 179
 armata, ii, 180
 blechnifolia, ii, 182
 cuneata, ii, 180
 falcata, ii, 180
 floribunda, ii, 179
 formosa, ii, 180
 longifolia, ii, 181
 mucronulata, ii, 180
 nivea, ii, 181
 obtusata, ii, 181
 plumosa, ii, 181
 pteridifolia, ii, 182
 tenuifolia, ii, 182
Dryas integrifolia, i, 178, 206
 octopetala, i, 206
 pentapetala, i, 206
 tenella, i, 206
Drymis, ii, 704
Drynaria, ii, 566
Dufourea rugosa, i, 178
Dunalia acaulis, i, 94
Dupontia, i, 221, 228
 Fisheri, i, 229, 253
Dysodium, ii, 285
Ebenaceæ, i, 109
Echinops, ii, 274
 Ritro, ii, 263
Echites, ii, 232-4
 acuminata, ii, 234
 agglutinata, ii, 234
 annularis, ii, 234
 asperuginis, ii, 234
 biflora, ii, 234
 bispinosa, ii, 235
 circinalis, ii, 234
 corymbosa, ii, 235, 238
 costata, ii, 246
 Domingensis, ii, 234
 floribunda, ii, 235, 238
 glandulosa, ii, 235
 hirsuta, ii, 234
 laxa, ii, 234
 quinguangularis, ii, 234
 repens, ii, 234
 scholaris, ii, 246
 siphilitica, ii, 235
 spicata, ii, 235, 238
 subcrecta, ii, 234
 succulenta, ii, 235
 torulosa, ii, 234
 umbellata, ii, 234
Eclipta, ii, 297 *note*
 sessilis, ii, 253
Edwardsia, ii, 419
 grandiflora, ii, 419
 microphylla, ii, 419
Ehretia abyssinica, i, 94
 obovata, i, 94
Ehrlharta, i, 56
Elæagneæ, i, 65, 82
Elæis, i, 164
 guineensis, i, 140, 153, 154, 161
Elæodendrum, i, 107; ii, 538
Elate sylvestris, i, 142
Elatine, ii, 536
 Hydropiper, i, 65
Eleocharis capitata, i, 144, 164
 palustris, ii, 513
Eleusine brevifolia, ii, 533
 indica, i, 165
 E? stolonifer, i, 93
Elichrysum, ii, 692
Ellisia, ii, 519
Elymus canadensis, ii, 515
 mollis, ii, 515

- Emba*, i, 140 *note*
Embelia, i, 150 *note*
Embothrium, ii, 11, 13, 18, 19, 22,
 38, 271
 buxifolium, ii, 146
 chaparro, ii, 187
 coccineum, ii, 165
 cytisoides, ii, 142
 dentatum, ii, 170
 emarginatum, ii, 13, 166
 ferrugineum, ii, 13, 169
 genianthum, ii, 146
 grandiflorum, ii, 166
 herbaceum, ii, 163
 hirsutum, ii, 170
 lanceolatum, ii, 165
 lineare, ii, 143
 linearifolium, ii, 143
 monospermum, ii, 162
 myricoides, ii, 169
 obliquum, ii, 170
 pinnatum, ii, 162
 salignum, ii, 155
 sericeum, ii, 142; β , ii, 142; γ ,
 ii, 143
 silaifolium, ii, 168
 spathulatum, ii, 167
 speciosum, ii, 167
 speciosissimum, ii, 167
 strobilinum, ii, 187
 tinctorium, ii, 168, 169
 truncatum, ii, 137
 umbellatum, ii, 170
Empetrum nigrum, i, 178
Empleurum, i, 16
Encalypta, ii, 349
 vulgaris, i, 69
Endocarpum hepaticum, i, 69
Enhydra, ii, 284
Ensete, i, 157
Enteleæa, ii, 677
 arborescens, ii, 678
Entelia ammannioides, ii, 710
Epacridæ, i, 36, 38, 62, 63, 64, 308,
 339, 380
Ephedra, i, 46, 453, 455, 457
Epiblema, ii, 549
Epidendrum, i, 49, 515; ii, 496
 aloifolium, ii, 493
 altissimum, ii, 474
 autumnale, i, 50
 Barringtoniæ, ii, 493
 ciliare, ii, 497
 cochleatum, ii, 496
 conopseum, i, 49; ii, 497
Epidendrum cucullatum, ii, 495
 elongatum, ii, 497
 ensifolium, ii, 493
 fragrans, ii, 496
 fuscatum, ii, 497
 humile, i, 49 *note*
 lineare, ii, 490
 nutans, ii, 497
 ophioglossoides, ii, 491
 præcox, i, 49 *note*
 præmorsum, ii, 494
 ruscifolium, ii, 491
 secundum, ii, 496
 sinense, ii, 493
 tripterum, ii, 493
 umbellatum, ii, 497
 undulatum, ii, 495
Epimedium, i, 124 *note*
Epipactis, i, 492, 542
 cordata, ii, 483
 ensifolia, ii, 484
 latifolia, ii, 484
 ovata, ii, 483
 pallens, ii, 484
 palustris, i, 528; ii, 484
 rubra, ii, 484
Epistephium, i, 501
Epithema, ii, 581, 586, 595, 606
Equisetacæ, i, 65
Equisetum, i, 469
 arvense, ii, 517
 palustre, ii, 518
 scirpoides, ii, 518
 sylvaticum, ii, 518
 umbrosum, ii, 518
 variegatum, ii, 518
 virgatum, i, 474 *note*
Eragrostis ciliaris, i, 165
Erasma, ii, 322
Eremocharis, i, 322
Eremodendron Cunninghamii, i, 332
Eremophila, i, 332, 339
 alternifolia, i, 332
 arborescens, i, 332
 Cunninghamii, i, 332
 Mitchelli, i, 333
 oppositifolia, i, 332
 Sturtii, i, 333
Ericæ, i, 12, 65
Ericinæ, i, 213, 381
Erigeron, i, 329
Erinus frutescens, ii, 504
 verticillatus, ii, 505
Eriocaulon, i, 53
Eriocephalus, ii, 470

- Eriocephalus africanus*, ii, 470
frutescens, ii, 470
racemosus, ii, 470
Eriophorum angustifolium, i, 218 ; ii, 514
capitatum, i, 218
gracile, i, 218
polystachyon, i, 177, 218
strictum, ii, 514
tenellum, ii, 514
vaginatum, ii, 514
Eriostemon, i, 17
parvifolius, ii, 709
Eriostylis, ii, 145
Ernodea, ii, 267
Fedendrum, ii, 15, 29, 60
amplexicaule, ii, 77
formosum, ii, 65
tenax, ii, 72
turbiniiflorum, ii, 76
Erodium, i, 288
Eruca peregrina, ii, 390
Erucaria, ii, 408
aleppica, ii, 408
Erysimum, ii, 402
Alliaria, ii, 404
angustifolium, ii, 405
Barbarea, ii, 398
bicorne, ii, 404
cheiranthoides, ii, 402
diffusum, ii, 403
helveticum, ii, 403
hieracifolium, ii, 404
lanceolatum, ii, 403
odoratum, ii, 404
officinale, ii, 400
orientale, ii, 404
præcox, ii, 398
repandum, ii, 403
sessiliflorum, ii, 403
Erythraea compar, i, 94
Erythrina tomentosa, i, 95
Erythrophileum, i, 111, 290
Erythropsis, ii, 639
Erythrospermum, i, 121
Erythroxylon, i, 106
ellipticum, ii, 709
Escallonea, ii, 523
Escallonia, ii, 523
Ethulia conyzoides, i, 162
Eucalyptus, i, 18, 62, 64, 74, 75, 311, 339
Baxteri, ii, 710
calophylla, i, 311
ciliata, i, 240
globulus, i, 18, 75
Eucalyptus grandifolia, ii, 710
hypericifolia, ii, 710
perfoliata, ii, 710
Euchilus, ii, 432
obcordatus, ii, 432
Euclidium, ii, 371
syriacum, ii, 371
Eudema, i, 254
Eudesmia, i, 74
tetragona, i, 75
Eugenia, i, 364
Eulophia virens, ii, 534
Eulophus, ii, 694
Euodia, i, 16
Euonymus, i, 448, 537
Eupatorium, ii, 299 *note*
ferrugineum, ii, 302
rosmariuifolium, ii, 302
Euphorbia, i, 28, 62, 128 ; ii, 279
propinqua, i, 95
puicea, ii, 279 *note*
Euphorbiaceæ, i, 27, 28, 104, 127, 283, 559 ; ii, 587 *note*, 615, 659, 663
Eupomatia, i, 73, 352
laurina, i, 73
Eurya, ii, 326
Chinensis, ii, 327
Japonica, ii, 328
Euryale, i, 451
Euryspermum, ii, 30, 40
grandiflorum, ii, 47
salicifolium, ii, 49
Eustegia, ii, 226
Eustrephus, i, 50
Eutaxia, ii, 431
myrtifolia, ii, 431
Euthales, i, 33 ; ii, 266 *note*
Eutoca, ii, 519
Franklinii, ii, 520
parviflora, ii, 519
Menziesii, ii, 519, 522
Eutrema, i, 193
Edwardsii, i, 193, 254
Evernia prunastri, i, 69
Exocarpus, i, 42, 137, 308, 310
cupressiformis, i, 43, 308, 310
Fagara, i, 16, 37, 133
Fagonia arabica, i, 285
armata, i, 94
cretica, i, 285
Mysorensis, i, 286
Oudneyi, i, 285
Farsetia, i, 262, 264, 265, 269 *note* ; ii, 388

- Farsetia Ægyptiaca*, i, 269 *note*
chciranthoides, ii, 383
clypeata, ii, 389
deltoidea, ii, 389
incana, ii, 389
lunarioides, ii, 388
mutabilis, ii, 389
stylosa, i, 269 *note*
Festuca brevifolia, i, 225, 256
Halleri, i, 225
ovina, i, 225 ; ii, 515
vivipara, i, 225
Fibigia, i, 271
Ficus, i, 137
Brassii, ii, 715
religiosa, i, 137
Fieldia, ii, 609
australis, ii, 609
Filices, i, 59, 62, 104, 146
Fimbristylis, i, 143, 300
Firmiana, ii, 639
colorata, ii, 640
platanifolia, ii, 639
Fissidens exilis, i, 69 ; ii, 344, 348
Fissilia, i, 136
disparilis, i, 136
Flabellaria pinnata, i, 106
Flacourtiaceæ, ii, 716
Flagellaria, i, 59
indica, i, 165, 166, 169
Flindersia, i, 71
australis, i, 71
Floerkea, ii, 363
Forstera, i, 34 ; ii, 312
Fothergilla, ii, 321
Francoaceæ, ii, 357
Franklandia, i, 79 ; ii, 15, 16, 18, 19, 37 ; ii, 130
fucifolia, i, 80 ; ii, 131
Fresiera, ii, 327
Frostia, i, 430
Fructus musculiformis, ii, 190
Fuchsia, i, 21
Fucus articulatus, i, 70
corneus, i, 70
obtusius, i, 70
palmatus, i, 70
pinnatifidus, i, 70
plicatus, i, 70
rubens, i, 70
sinuosus, i, 70
Fuirena, i, 53
umbellata, i, 144, 164, 169
Fumariaceæ, i, 65, 272
Funaria hygrometrica, i, 69, 346
Funaria minor, ii, 334
Fungi, i, 253
Fusanus, i, 42
alternifolia, i, 93
Gærtnera, i, 37, 132
Gaguedi, i, 163 *note*
Galega astragaloides, ii, 709
crocea, ii, 710
juncea, ii, 709
oblongata, ii, 709
porrecta, ii, 709
reticulata, ii, 710
Galipea, i, 16
Galium, i, 295
Gangi, i, 155
Garcinioides, ii, 537
Gardenia, i, 131, 269 *note*
fucata, ii, 711
suffruticosa, ii, 711
Gastrolobium, ii, 432
bilobum, ii, 432
Gazania, ii, 464
Pavonia, ii, 464
rigens, ii, 464
subulata, ii, 464
Geissorhiza abyssinica, i, 93, 163
Geniostoma, i, 37, 132
Gentianæ, i, 36, 37, 133, 295
Geodorum, ii, 488
citrinum, ii, 488
dilatatum, ii, 488
purpureum, ii, 488
Geraniaceæ, i, 114, 288
Geranium compar, i, 95
Geropogon glabrum, ii, 270
Gesneriaceæ, ii, 586, 589
Gesneriæ, ii, 589
Getonia, ii, 609
Geum, i, 206
congesta, i, 206
dilatata, i, 206
glaciale, i, 206
humilis, i, 206
montanum, i, 206
Peckii, i, 206
radiatum, i, 206
reptans, i, 206
triflorum, i, 206
urbanum, i, 65
Gisekia, i, 139
pharuaceoides, i, 165, 172
Gleichenia, i, 59
Hermanni, i, 146, 164
Glins lotoides, i, 162, 165, 166

- Globba, ii, 551
 Globularia cordifolia, ii, 315
 Globularia, i, 65
 Gloriosa simplex, i, 170
 superba, i, 165, 169, 170
 Glossanthus, ii, 577, 590, 595, 607
 grandifolia, ii, 607
 Malabarica, ii, 607
 Mexicana, ii, 581, 607
 Notoniana, ii, 607
 Zeylanica, ii, 607
 Glossodia, ii, 485, 549
 major, ii, 485
 Glyceria, i, 56, 226
 fluitans, i, 68
 Glycine bimaculata, ii, 440
 coccinea, ii, 440
 rubicunda, ii, 440
 subterranea, i, 155, 160
 Glycyrrhiza, i, 323
 Glyphocarpa capensis, ii, 335
 Gnaphalium, ii, 294, 300
 alpinum, i, 212; ii, 302
 carpathicum, ii, 302
 dioicum, ii, 302
 Leontopodioides, ii, 302
 Leontopodium, ii, 302
 luteo-album, ii, 300
 margaritaceum, ii, 301 *note*, 302
 mucronatum, ii, 302
 muricatum, ii, 302
 plantagineum, ii, 302
 seriphioides, ii, 302
 sylvaticum, ii, 300
 uliginosum, ii, 300
 Gnaphaloideæ, i, 30
 Gnetum, i, 453, 455
 Gomeza, ii, 669
 recurva, ii, 669
 Gomphocarpos, ii, 216
 arborescens, ii, 217
 crispus, ii, 217
 fruticosus, i, 542; ii, 217
 setosus, ii, 217
 Gompholobium, ii, 427
 ellipticum, ii, 426
 fimbriatum, ii, 427
 latifolium, ii, 427
 maculatum, ii, 422
 marginatum, ii, 427
 polymorphum, ii, 427
 psoralæfolium, ii, 427
 scabrum, ii, 428
 spinosum, ii, 429
 tomentosum, ii, 428
 Gompholobium venustum, ii, 428
 Gonocarpus, i, 22
 Gonolobus, ii, 215
 Carolinensis, ii, 215
 crispiflorus, ii, 215
 grandiflorus, ii, 215
 hirtus, ii, 215
 maritimus, ii, 215
 niger, ii, 215
 obliquus, ii, 215
 planiflorus, ii, 215
 prostratus, ii, 215
 racemosus, ii, 215
 rostratus, ii, 215
 suberosus, ii, 215
 undulatus, ii, 215
 Goodenia, ii, 267 *note*
 beilidifolia, ii, 267 *note*
 cycloptera, i, 331
 decurrens, ii, 267 *note*
 Goodenovia, i, 31, 63, 81, 308; ii, 266, 271, 272, 310
 Goodenovia littoralis, i, 34, 64
 Goodia, ii, 439
 lotifolia, ii, 439
 pubescens, ii, 439
 Goodyera, ii, 481
 pubescens, ii, 481
 repens, ii, 481
 Gorteria, ii, 464
 araneosa, ii, 463
 asteroides, ii, 463
 cernua, ii, 463
 personata, ii, 464
 rigens, ii, 464
 Gossypium, i, 317
 Gouania, i, 26
 Gramineæ, i, 54, 62, 85, 104, 145, 169, 182 *note*, 219, 298, 300, 468, 559; ii, 276, 587 *note*
 Grammitis, ii, 525, 566
 Ceterach, i, 303
 leptophylla, ii, 526
 Grangea maderaspatana, i, 162, 165, 167
 Grangeria, i, 115
 Gratiola involucreta, ii, 555
 Grevillea, i, 41, 339; ii, 11, 18, 20, 22, 37, 140, 271, 280
 acuminata, ii, 145
 arenaria, ii, 144
 aspera, ii, 144
 asplenifolia, ii, 147
 australis, ii, 143
 Banksii, ii, 147

- Grevillea Baueri*, ii, 145
buxifolia, ii, 146
ceratophylla, ii, 148
Chrysodendron, ii, 16, 147
cinerea, ii, 145
concinna, ii, 144
Dryandri, ii, 147
dubia, ii, 142
gibbosa, ii, 149
Goodii, ii, 146
heliosperma, ii, 148
juncea, i, 335
juniperina, ii, 143
linearis, ii, 142
lineata, i, 335
longistyla, i, 335
lorea, i, 335; ii, 149
mimosoides, ii, 148
Mitchellii, i, 335
montana, ii, 144
mucronulata, ii, 145
neglecta, i, 335
occidentalia, ii, 145
parviflora, ii, 143
pauciflora, ii, 144
phylicoides, ii, 146
polystachya, ii, 148
pungens, ii, 147
puicea, ii, 142
refracta, ii, 148
riparia, ii, 143
sericea, ii, 142
sphacelata, ii, 146
striata, i, 335; ii, 148
stricta, ii, 143
Sturtii, i, 334
teuuiifolia, ii, 144
venusta, ii, 146
Grewia, i, 108; ii, 534
Griffithia Daviesii, ii, 334
parviflora, ii, 689
Grimmia, ii, 338, 344
apocarpa, i, 69
Daviesii, ii, 344
pulvinata, i, 69
Grindelia, ii, 282
Grossularinæ, i, 65
Ground nuts, i, 154
Guevina, ii, 18, 36, 138
Avellana, ii, 138
Guiera, i, 20
Guilandina, i, 294
Bonduc, i, 111, 165, 168
Bonducella, i, 165, 171
Guttifera, i, 149
Gwillimia, ii, 705
Gymnadenia, ii, 475
conopsea, ii, 475
Gymnanthera, ii, 231
Gymnema, ii, 213
lactifera, ii, 213
sylvestris, ii, 214
Gymnocarpus decandrum, i, 297
Gymnogonia, i, 275
Gymnogramma, ii, 568
Gymnostomum, ii, 341, 354
Bonplandii, ii, 344
fasciculare, ii, 341, 344
Griffithianum, ii, 341
julaceum, ii, 345
lapponicum, ii, 344
microstomum, ii, 341, 344
obtusifolium, i, 240, 253
pennatum, ii, 344
pulvinatum, ii, 344
pyrifforme, i, 69; ii, 339
Rottleri, ii, 344
viridissimum, ii, 344
Gymnostyles, ii, 281
anthemifolia, ii, 282
nasturtiifolia, ii, 282
Gynandropsis, i, 273, 275
Gynocardia, ii, 716
Gynoecephalum, ii, 651
Gyratæ, ii, 525
Gyrinops, i, 126
Gyromium proboscideum, i, 250
Gyrophora erosa, i, 178, 250
heteroidea β , i, 69
hirsuta, i, 178, 250
polyphylla, i, 69
proboscidea, i, 69
Habenaria, ii, 476
albida, ii, 477
bifolia, i, 491, 500; ii, 477, 549
bracteata, ii, 476
ciliaris, ii, 477
cristata, ii, 477
fimbriata, ii, 477
herbiola, ii, 477
hyperborea, ii, 477
nigra, ii, 476
viridis, i, 540 *note*; ii, 476
Hakea, i, 310, 339; ii, 11, 15, 18, 22, 37, 187
acicularis, ii, 152
amplexicaulis, ii, 155

Hakea arborescens, ii, 157

- attenuata*, ii, 154
- ceratophylla*, ii, 155
- cinerea*, ii, 156
- clavata*, ii, 15
- cycloptera*, ii, 153
- dactyloides*, ii, 156
- elliptica*, ii, 157
- epiglottis*, ii, 150
- flexilis*, ii, 151
- florida*, ii, 154
- gibbosa*, ii, 152
- glabra*, ii, 150
- ilicifolia*, ii, 154
- leucoptera*, ii, 151
- linearis*, ii, 154
- lissosperma*, ii, 151
- lorea*, i, 335
- marginata*, ii, 156
- microcarpa*, ii, 153
- nitida*, ii, 154
- nodosa*, ii, 151
- obliqua*, ii, 151
- oleifolia*, ii, 155
- prostrata*, ii, 155
- pubescens*, ii, 152
- pugioniformis*, ii, 150
- pyriformis*, ii, 159
- rugosa*, ii, 150
- ruscifolia*, ii, 156
- saligna*, ii, 155
- sericea*, ii, 152
- suaveolens*, ii, 153
- sulcata*, ii, 151
- trifurcata*, ii, 153
- undulata*, ii, 155
- varia*, ii, 153
- vittata*, ii, 152

Haloragæ, i, 21, 63

Haloragis, i, 21, 22

- lanceolata*, ii, 710
- paniculata*, ii, 710
- pusilla*, ii, 710
- stricta*, ii, 710

Hamamelidæ, i, 381; ii, 321, 334

Hamamelis, ii, 321

- Chineusis*, ii, 322
- Virginica*, ii, 323

Harrisonia, ii, 707

Hedwigia secunda, ii, 335

Hedychium elatum, ii, 690

- Horsfieldii*, ii, 725
- speciosum*, ii, 690

Hedyotis, i, 131, 295

- alata*, ii, 532

Hedyotis alsinifolia, ii, 532

- approximata*, ii, 532
- H. ? arguta*, ii, 532
- articularis*, ii, 532
- biflora*, ii, 532
- Burnianniana*, ii, 532
- cephalophora*, ii, 531
- congesta*, ii, 531
- costata*, ii, 532
- cymosa*, ii, 532
- H. ? elongata*, ii, 532
- brachypoda*, ii, 532
- extensa*, ii, 532
- H. ? glabra*, ii, 531, 532
- Heynii*, ii, 532
- linarifolia*, ii, 532
- H. ? macrocephala*, ii, 531
- polycarpa*, ii, 531
- puberula*, ii, 532
- racemosa*, ii, 532
- scandens*, ii, 531
- scapigera*, ii, 532
- stipulata*, ii, 532
- stylosa*, ii, 532
- tubularis*, ii, 532
- vestita*, ii, 531
- volubilis*, ii, 531

Hedysarum adscendens, i, 165

- pictum*, i, 165
- triflorum*, i, 111, 165
- vaginale*, i, 165

Helianthæ, i, 128; ii, 297

Helianthemum, i, 453 *note*

Helianthus, ii, 266

- multiflorus*, ii, 265

Helicia, ii, 532

- cochinchinensis*, ii, 161
- robusta*, ii, 532

Helicteræ, ii, 620

Helicteres apetalæ, ii, 628

- paniculata*, ii, 632
- undulata*, ii, 632

Heliophila, ii, 391

- amplexicaulis*, ii, 391
- arabioides*, ii, 391
- filiformis*, ii, 391
- fœniculacea*, ii, 392
- incana*, ii, 391
- integrifolia*, ii, 391
- pendula*, ii, 392
- pilosa*, ii, 391
- pinnata*, ii, 392
- platisiliqua*, ii, 391

Heliotropium cinereum, i, 94

- H. ? dubium*, i, 94

- Heliotropium ellipticum*, i, 94
 gracile, i, 94
 indicum, i, 165
Helosis guianensis, i, 124
Helvella membranacea, i, 253
Hemarthria, i, 58
Hemerocallidæ, i, 513
Hemidesmus, ii, 230
Hemigenia, i, 39
Hemionitis hastata, ii, 532
Hemistemma, i, 12
 angustifolium, ii, 703
 Banksii, ii, 703
 dealbatum, ii, 703
Hepaticæ, i, 66, 249
Heritiera, ii, 615, 643
 attenuata, ii, 643
 Fomes, ii, 643
 littoralis, ii, 643
 minor, ii, 643
Hermannia, i, 63
Hermanniaceæ, i, 109
Herminium, ii, 476
 Monorechis, ii, 476
Hermodactylum, i, 300
Hernandia, ii, 651
Hernandiaceæ, ii, 651
Hesperidæ, i, 72, 149
Hesperis africana, ii, 407
 inodora, ii, 408
 matronalis, ii, 408
 nitens, i, 272; ii, 407
 ramosissima, i, 272
 tristis, ii, 407
 verna, ii, 396
Heuchera, ii, 522
 Richardsonii, ii, 523
Hexameria, ii, 575
 disticha, ii, 575
Hibbertia, i, 12
 cistifolia, ii, 704
 dentata, ii, 704
 diffusa, ii, 704
 echinifolia, ii, 708
 fasciculata, ii, 704
 lepidota, ii, 704
 linearis, ii, 704
 monogyna, ii, 704
 oblongata, ii, 704
 pedunculata, ii, 704
 saligna, ii, 704
 scabra, ii, 708
 serpyllifolia, ii, 704
 stricta, ii, 708
 tomentosa, ii, 704
Hibbertia vetulina, ii, 708
 virgata, ii, 704
 volubilis, i, 449
Hibiscus, i, 109
 crianthus, i, 95
 parvifolius, i, 95
 surattensis, i, 165
 thespesioides, ii, 709
 tiliaceus, i, 165, 171
Hierochloë, i, 231; ii, 571
 alpina, i, 178, 232
 fragrans, ii, 514
 laxa, ii, 533
 pauciflora, i, 232
Hildegardia, ii, 639
 populifolia, ii, 639
Hingslia, ii, 284
Hippia minuta, ii, 281
 stolonifera, ii, 282
Hippocratea, i, 107
Hippocrateæ, ii, 537
Hippocrateæ, i, 27, 107
Hippuris, i, 21, 32
Hiræa pinnata, i, 106
Hirculus propinquus, i, 202
Hirtella, i, 115
Holarrhena, ii, 232, 236, 244
Holcus, i, 57, 232
 acicularis, i, 302
 alpinus, i, 178, 232; ii, 514
 Gryllus, i, 66, 68
 saccharatus, i, 160
 sorghum, i, 160
Holostemma, ii, 220
Homalinæ, i, 120, 384, 386
Homalium, i, 120
Hookeria, ii, 337
 lucens, i, 69
Hordeum jubatum, ii, 515
Hoslundia, i, 135
Iostea, ii, 215
Hovea, i, 25; ii, 439
 linearis, ii, 440
 longifolia, ii, 440
Hovenia, i, 26
Hoya, ii, 200, 208, 716
 australis, ii, 715
 carnosa, i, 527, 529; ii, 209, 715
 Nicobarica, ii, 716
 viridiflora, ii, 209
Hudsonia, i, 453 *note*
Huernia, ii, 200, 205
 campanulata, ii, 206
 guttata, ii, 206

- Huernia venusta*, ii, 206
Humboldtia, ii, 552
 laurifolia, ii, 552
Hutchinsia, i, 263; ii, 378
 alpina, i, 263; ii, 378
 petraea, i, 263; ii, 378
 rotundifolia, ii, 378
Hybanthus, i, 122
Hydnocarpus, ii, 716
Hydnora, i, 401, 427, 558
 africana, i, 402, 412, 416, 427
 americana, i, 428
 triceps, i, 412, 428
Hydnorea, i, 427
Hydrocotyle alata, ii, 706
 asiatica, i, 165
 flaccida, ii, 706
 heteromeria, ii, 706
 hirta, ii, 705
 intertexta, ii, 706
 muscosa, ii, 706
 peduncularis, ii, 706
 plebeia, ii, 706
 pulchella, ii, 706
 pusilla, ii, 706
 tripartita, ii, 706
 vulgaris, i, 68
Hydrolea, i, 135
Hydrolea, i, 135
Hydropeltideæ, i, 556
Hydropeltis, i, 54, 451
Hydrophyllææ, ii, 519
Hydrophyllum, ii, 519
 lineare, ii, 519
Hymenanthera, i, 125
 angustifolia, ii, 705
 dentata, ii, 705
Hymenopappus, ii, 265, 267 *note*, 268
 note
 scabiosæus, ii, 265
 tenuifolius, ii, 266
Hymenophyllum, i, 146; ii, 342
 alatum, ii, 510
 tunbridgense, i, 68
Hypælyptum argenteum, i, 144, 164
Hypælytrum, i, 144
Hyphæne, i, 141
 coriacea, i, 141
Hyppnum, ii, 345
 aduncum, i, 178, 235
 cordifolium, i, 234
 nitens, i, 234
 recognitum, i, 68
 rufescens, i, 235
 velutinum, i, 346
Hypoderris, ii, 543
Hypœlyptum, i, 54
Hypoestes, i, 134
 Forskalii, i, 93
Hypophyllocarpodendron, ii, 25, 86
Hypoxidææ, i, 51
Hypoxis, i, 51
 fascicularis, i, 298
Hyptis, i, 135, 164
 obtusifolia, i, 165, 167
Iberis, i, 265; ii, 379
 amara, ii, 380
 ciliata, ii, 379
 gibraltaria, ii, 379
 linifolia, ii, 380
 nudicaulis, ii, 378
 pinната, ii, 380
 rotundifolia, ii, 378
 saxatilis, ii, 379
 semperflorens, ii, 379
 sempervirens, ii, 379
 umbellata, ii, 379
 violacea, ii, 380
Ibetssonia, ii, 422
 genistoides, ii, 422
Ichnocarpus, ii, 232, 235
Ilex Japonica, i, 124 *note*
Illecebreææ, i, 138; ii, 543
Illicium, ii, 704
Impatiens, ii, 535
 tenella, i, 94
Imperata aurundinacea, i, 68
 cylindrica, i, 165
Incarvillea, ii, 725
 oblongifolia, ii, 599
 parasitica, ii, 585
Indigofera, i, 295
 albicans, i, 95
 diffusa, i, 95
 lateritia, i, 165
Incumba, i, 155
Inga biglobosa, i, 288
 Senegalensis, i, 289
Inocarpus, ii, 651
Inula, ii, 294
Iodes, ii, 650
 ovalis, ii, 650
Ionidium, i, 122 *note*
 aquatica, i, 296
 Clappertoni, i, 296
 pentaphylla, i, 164
 pes-caprææ, i, 164
Ipomœa, i, 135, 296; ii, 545
Ipomopsis, i, 532

- Iridæ*, i, 54, 56, 64, 502, 513, 559
Iris, i, 298, 559
Isachne, i, 57
Isatis, i, 270 *note*; ii, 271, 375
 aleppica, ii, 375
 lusitanica, ii, 375
 orientalis, ii, 375
 tinctoria, ii, 375
Ischæmum, i, 57
 rottboëlioides, ii, 553
Isocarpha, ii, 289
Isophilus, ii, 490
 linearis, ii, 490
 prolifer, ii, 490
Isolepis fluitans, i, 68
 setacea, i, 68
Isonema, ii, 232, 236
Isopogon, i, 310; ii, 11, 14, 16, 17, 35, 58, 185
 anemonifolius, ii, 59, 185
 anethifolius, ii, 58, 185
 attenuatus, ii, 60
 axillaris, ii, 60
 buxifolius, ii, 60
 ceratophyllus, ii, 59
 cuneatus, ii, 59
 formosus, ii, 58
 longifolius, i, 59
 polyccephalus, ii, 60
 teretifolius, ii, 58
 trilobus, ii, 59
Iva frutescens, ii, 266 *note*
Ivira, ii, 615, 616
 pruriens, ii, 629
Ixodia, ii, 461
 achillæoides, ii, 461
Ixora, ii, 269 *note*
 ?, *arguta*, ii, 533
 concinna, ii, 533
 coriacea, ii, 533
 densa, ii, 533
 diversifolia, ii, 533
 elliptica, ii, 533
 grandifolia, ii, 533
 lucida, ii, 533
 macrophylla, ii, 533
 nigricans, ii, 533
 oblonga, ii, 533
 opaca, ii, 533
 propinqua, ii, 533
 triflora, ii, 711
Jacaranda, ii, 673
 acutifolia, ii, 674
 bahamensis, ii, 674
Jacaranda caroliniana, ii, 674
 obtusifolia, ii, 674
 ovalifolia, ii, 674
 procera, ii, 674
 rhombifolia, ii, 674
 tomentosa, ii, 675
Jacksonia, ii, 428
 scoparia, ii, 429
 spinosa, ii, 429
Jacquinia, i, 151 *note*
Jamaica nutmeg, i, 161
Jambolifera, i, 16
Jasminum abyssinicum, i, 93
 lineare, i, 330, 339
 micanthum, i, 331
 Mitchellii, i, 330
Jasione, ii, 314
Jatropha Manihot, i, 154
Jeffersonia, i, 124 *note*
Jenkinsia, ii, 652, 653
Josephia, ii, 11, 19
Juncæ, i, 52, 53, 216, 209 *note*, 436;
 ii, 278
Juncus, i, 298
 affinis, ii, 515
 arcuatus, i, 217
 bicornis, ii, 515
 biglumis, i, 216
 campestris, i, 217; ii, 515
 castaneus, ii, 515
 echinatus, ii, 515
 effusus, i, 68
 filiformis, ii, 515
 glaucus, ii, 515
 maritimus, i, 68
 melanocarpus, ii, 515
 tenuis, ii, 515
 triglumis, ii, 515
Jungermannia, i, 178; ii, 351
 bicornis, i, 249
 bidentata, i, 69
 byssacea, i, 69
 complanata, i, 69
 furcata, i, 69
 minuta, i, 249
 punguis, i, 69
 tamarisci, i, 69
 tomentella, i, 69
 ventricosa, i, 249
Jussiaea arborea, ii, 505
 bivalvis, ii, 93
 cristata, ii, 505
 cynanchifolia, i, 93
 erecta, i, 165, 167, 171
 Forskaliæ, i, 93

Jussiea infundibuliformis, ii, 505
puleherrima, ii, 505
tetragona, ii, 505

Kadsura, ii, 685
Kallstroemia, i, 318
Kanabia, ii, 218
laniflora, i, 94; ii, 218

Kangaroo grass, i, 308
Kantuffa, i, 94
Kasheia, i, 303
Kennedia, i, 25; ii, 440
coccinea, ii, 440
monophylla, ii, 440
prostrata, ii, 440
rubicunda, ii, 440
speciosa, i, 321

Kerneria, i, 271
Kingia, i, 433, 436
australis, i, 308, 436; ii, 724

Kino gum, i, 290
Klugia azurea, ii, 607
Knightia, ii, 12, 38, 163
Kuoxia mollis, ii, 531
teres, ii, 531

Kobresia scirpina, ii, 515
Koniga, i, 266 *note*, 271
libyca, i, 263
maritima, i, 263

Kouka, i, 288
Krameria, i, 14
Krubut, i, 372, 393
Kyllinga, i, 54, 144

Labiata, i, 38, 135, 297, 380

Labioclea, i, 328
Lada Tchilli, i, 158
Lactuca virosa, ii, 263
Lagasca, ii, 274
Laguncularia, i, 20

Lambertia, ii, 11, 15, 18, 38, 157, 280
echinata, ii, 158
formosa, ii, 158, 280
inermis, ii, 158, 280
teretifolia, ii, 150
uniflora, ii, 158, 280

Langsdorfia, i, 412
Lantana, i, 136
polycephala, i, 94

Lappago, i, 57, 379 *note*
Lardizabalea, i, 555
Lasianthera, i, 151
Lasiodictyon, i, 11, 12
rufum, ii, 708
Lastrea, ii, 568

Lauradia, i, 124

Laurelia, i, 26, 64
Laurinae, i, 25, 149, 380
Laurus, i, 150

Lavandula, i, 297
multifida, i, 297
Lavenia erecta, i, 165, 167; ii, 290
Laxmannia, ii, 292 *note*

Lecanora atra, i, 69
brunnea, β *nebulosa*, i, 69
elegans, i, 250
decipiens, i, 69
 β *dendritica*, i, 69
fusco-atra, i, 69
gelida, i, 69
lentigera, i, 69
lepidosa, i, 69
microphylla, i, 69
parella, i, 69
subfusca, i, 69
sulphurea, i, 69
ventosa, i, 69

Lechea, i, 453 *note*

Lecidea confluens, i, 69
geographica, i, 69
lurida, i, 69
luteola, i, 69
parasema, i, 69

Leca, i, 151
sambucina, i, 165, 169

Leersia ciliata, i, 240

Legnotis, i, 119, 164

Leguminosae, i, 22, 62, 104, 110, 113,
 168, 288, 291, 307, 503

Leichhardtia, i, 329
australis, i, 329

Lemna, i, 448
minor, i, 68
trisulca, i, 68

Leontice, i, 124 *note*
altaica, i, 362
Leontopetalum, i, 362
thalictroides, i, 361

Leontodon lividus, i, 210
nana, i, 178

palustre, i, 210
Taraxacum, i, 178, 210

Leontopodium, ii, 302

Lepeyrodia, i, 145

Lepidium, ii, 380
alpinum, ii, 378
bidentatum, ii, 382
bonariense, ii, 381
campestre, ii, 382
Cardamines, ii, 381

- Lepidium didymum*, ii, 373
divaricatum, ii, 381
Draba, ii, 381
graminifolium, ii, 381
Iberis, ii, 381, 383
latifolium, ii, 380
lyratum, ii, 380
oleraceum, ii, 382
perfoliatum, ii, 382
petraeum, ii, 378
piscidium, ii, 382
procumbens, i, 271
rotundifolium, ii, 378
ruderales, ii, 381
salivum, ii, 382
spinosum, ii, 382
subulatum, ii, 382
virginicum, ii, 383
Lepidocarpodendron, ii, 25, 26, 29
Lepidodendron Harecourtii, ii, 722
Lepidosperma, i, 53
Lepraria botryoides, i, 70
flava, i, 70
incana, i, 70
Leptadenia, i, 214
Leptomeria, i, 42
Leptospermum, i, 19, 311 *note*
Leptostomum, i, 352; ii, 341, 345
erectum, i, 352
gracile, i, 353
inclians, i, 352
Menziesii, i, 353
Lepturus, i, 55; ii, 553
Lepyrodia, i, 54
Leskea, ii, 335
complanata, i, 68
rufescens, i, 235
Lessertia, ii, 442
annua, ii, 442
diffusa, ii, 443
perennans, ii, 443
Leucadendron, ii, 15, 17, 21, 22, 24,
29, 35, 40, 41, 60, 183, 184,
723
abietinum, ii, 52
acaulon, ii, 73
adscendens, ii, 49, 51, 184
æmulum, ii, 52
angustatum, ii, 43, 183
argenteum, ii, 41
buxifolium, ii, 44
cancellatum, ii, 40
cartilagineum, ii, 54
cinaroides, α , ii, 61
cinereum, ii, 46
Leucadendron comosum, ii, 52
concinnum, ii, 50
concolor, ii, 47
Conocarpodendron, ii, 51
corymbosum, ii, 46
crassifolium, ii, 54
cucullatum, ii, 88
cyanooides, ii, 97
decorum, ii, 46
decurrens, ii, 48
divaricatum, ii, 91
elongatum, ii, 109
crucifolium, ii, 54
floridum, ii, 51
fusculorum, ii, 183
glabrum, ii, 48, 50
Globularia, ii, 53
glomeratum, ii, 28, 106
grandiflorum, ii, 47, 82
hirtum, ii, 87
Hypophyllocarpodendron, ii, 84
imbricatum, ii, 44
Lepidocarpodendron, ii, 66, 68
Levisanus, ii, 45
linifolium, ii, 183
nanum, ii, 71
oleæfolium, ii, 25
ovale, ii, 48
phylicoides, ii, 103
pinifolium, ii, 28, 39
platyspermum, ii, 51
plumosum, ii, 42, 184
protocoides, ii, 27, 91
pubescens, ii, 53
racemosum, ii, 120
repens, ii, 69; β , ii, 75
retusum, ii, 43, 62, 184, 187
salignum, ii, 50
scabrum, ii, 53
Scolymocephalum, ii, 70
sericeum, ii, 53
Serruria, α , ii, 108; β , ii, 106
sessile, ii, 43
spathulatum, ii, 43
speciosum, ii, 28, 64
sphærocephalum, ii, 186
spicatum, ii, 112
squarrosus, ii, 47, 184
strictum, ii, 48
thymelæoides, ii, 90
tortum, ii, 45, 183
uliginosum, ii, 51
venosum, ii, 48
virgatum, ii, 49, 50
Leucadendros, ii, 26, 30, 31

- Leucadendros africana*, ii, 30
Leucadendrum, ii, 78
Leucus affinis, i, 94
 quinquedentata, i, 94
Leucodon, ii, 318
Leucolæna, i, 29
Leucophyta, ii, 286
Leucosin, i, 125
Leucospermum, ii, 20, 35, 78, 185, 271
 attenuatum, ii, 79
 luxifolium, ii, 82
 Conocarpum, ii, 81, 190
 orinitum, ii, 85
 diffusum, ii, 85
 ellipticum, ii, 80, 185
 grandiflorum, ii, 81, 82
 Hypophyllum, ii, 84
 lineare, ii, 79
 medium, ii, 80
 molle, ii, 85
 nutans, ii, 80
 oleæfolium, ii, 85
 patulum, ii, 82, 86
 puberum, ii, 82, 83
 spithulatum, ii, 83
 tomentosum, ii, 83
 Tottum, ii, 79
Levenhookia, i, 34
Leycesteria, ii, 542
Liatris, ii, 299 *note*
 spicata, ii, 270
Licania, i, 115
Lichen aphthosus, i, 251
 cucullatus, i, 251
 elegans, i, 250
 islandicus, i, 251
 lanatus, i, 251
 nivalis, i, 250
 normöricus, i, 251
 ochroleucus, i, 251
 odontellus, i, 251
 paschalis, i, 252
 ramulosus, i, 252
 vermicularis, i, 252
Lichenosæ, i, 66, 250
Liliacæ, i, 299, 439, 449, 450, 513;
 ii, 278
Lime, i, 154, 156
Limnanthæ, ii, 363
Limnanthus, ii, 363
 Douglasii, ii, 364
Limodorum, ii, 689
 altum, ii, 487
 barbatum, ii, 694
Limodorum carinatum, ii, 694
 epidendroides, ii, 694
 hians, ii, 694
 longicornis, ii, 694
 nutans, ii, 488
 purpureum, ii, 487
 striatum, ii, 694
 Tankervillei, ii, 487
 triste, ii, 694
 tuberosum, ii, 486
 veratrifolium, ii, 694
 virens, ii, 694
Limonia, ii, 664
Linaria gracilis, i, 94
 hastata, i, 94
 propinqua, i, 94
Linconia, ii, 322
Lindernia japonica, ii, 505
Linkia, ii, 133
 lævis, ii, 136
Linnæa, ii, 324, 541, 673
Linschotenia, i, 315
Lipocarypha, i, 144
 argentea, i, 164, 169
Lipotrichis, ii, 297 *note*
Lippia, i, 136
Liriope spicata, ii, 695
Lissochilus, ii, 692, 694
 speciosus, ii, 692, 707
Listera, ii, 483
 cordata, ii, 483
 ovata, ii, 483
Lithospermum? ambiguum, i, 94
Lobelia, i, 32
Lobeliacæ, i, 292; ii, 269 *note*, 310,
 678
Lobularia, i, 266
Loganæ, i, 37
Logania, i, 37, 132
Lolium, i, 55; ii, 277
 perenne, i, 468 *note*, 530 *note*
Lomaria, ii, 525
Lomatia, i, 41, 64; ii, 38, 168
 dentata, ii, 169
 ferruginea, ii, 168
 ilicifolia, ii, 169
 longifolia, ii, 169
 obliqua, ii, 170
 polymorpha, ii, 169
 silaifolia, ii, 168
 tinctoria, ii, 168
Lomentacæ, i, 23
Lonicera, ii, 687
Lonicercæ, ii, 324

- Loranthaceæ, i, 413 *note*, 501
 Loranthææ, i, 288
 Loranthus, i, 21, 120, 137, 288
 angustifolius, ii, 710
 calycinus, i, 94
 congestus, i, 94
 floribundus, i, 308
 lætus, i, 94
 Loropetalum, ii, 323
 Lotus arboreus, ii, 698
 Loxidium, ii, 442
 Loxocarpus, ii, 595, 606
 incana, ii, 606
 Loxonia, ii, 580, 582, 595
 acuminata, ii, 582
 discolor, ii, 582
 hirsuta, ii, 584
 Loxophyllum, ii, 585
 Loxotis, ii, 577, 590, 595
 intermedia, ii, 577
 obliqua, ii, 577
 Lunaria, ii, 390
 annua, ii, 390
 libyen, i, 262, 265
 parviflora, i, 262
 rediviva, ii, 390
 Luzula campestris, i, 68, 217; ii,
 515
 hyperborea, i, 216
 melanocarpa, ii, 515
 pilosa, i, 217
 Luzuriaga, i, 50
 Lychnis, i, 285
 apetala, i, 177, 198
 triflora, i, 178
 Lycoperdon pratense, i, 253
 Lycopodiaceæ, ii, 722
 Lycopodineæ, i, 147
 Lycopodium alpinum, ii, 518
 annotinum, ii, 518
 complanatum, ii, 518
 dendroideum, ii, 518
 Selago, i, 178; ii, 518
 Lycopus europæus, i, 65
 Lyellia, ii, 331, 345, 351
 crispa, ii, 333
 Lygodium, i, 146
 Lyonsia, ii, 232, 238
 Lysionotus, ii, 585, 593, 594, 598
 serratus, ii, 598
 ternifolia, ii, 598
 Lysipoma, ii, 679
 Lyssostylis, ii, 142
 Lythrum, ii, 238
 Lythrum Salicaria, i, 68
 Maba, i, 140; ii, 536
 Macleaya cordata, i, 270 *note*
 Macradenia, ii, 695
 lutescens, ii, 695
 Macromitrium, ii, 335
 Macropodium, ii, 397
 nivale, ii, 397
 Madia, ii, 266
 Mærua, i, 280 *note*
 angolensis, i, 281 *note*
 rigida, i, 280
 senegalensis, i, 280
 uniflora, i, 281 *note*
 Mæsa, i, 151 *note*
 Magnolia, ii, 705
 Magnoliaceæ, i, 12, 149
 Mahernia, i, 63
 Maize, i, 155, 160
 Malachra radiata, i, 109, 165
 Malaguetta Pepper, i, 155
 Malaxideæ, ii, 575
 Malaxis, ii, 489
 lilifolia, ii, 489
 Loeselii, ii, 489
 nutans, ii, 488
 paludosa, ii, 489
 Malcomia, i, 272; ii, 407
 littorea, ii, 407
 Malesherbia, i, 387 *note*
 Malpighiaceæ, i, 105
 Malvaceæ, i, 11, 26, 104, 109, 168,
 288; ii, 613
 Mammillaria, i, 549
 Mangifera, i, 112
 Mangrove, i, 363
 Manioc, i, 155, 160
 Manisuris, ii, 554
 Mantellia cylindrica, ii, 717
 nidiformis, ii, 717
 Marantææ, i, 49, 380; ii, 551
 Marchantia hemispherica, i, 69
 polymorpha, i, 69, 249
 Maregravia, ii, 357
 Marigold, ii, 262
 Marlea, ii, 322
 barbata, ii, 534
 begonifolia, ii, 532
 Marsdenia, ii, 210
 cinarescens, ii, 211
 clausa, ii, 211
 erecta, ii, 212
 rostrata, ii, 212

Marsdenia suaveolens, ii, 211
tinctoria, ii, 211
velutina, ii, 211
viridiflora, ii, 211
Marshallia, ii, 267 *note*
Marsilea quadrifolia, i, 68
Matelea, ii, 215
Mathiola, i, 272, 316; ii, 405
elliptica, i, 94
fenestralis, ii, 406
incana, ii, 405
odoratissima, ii, 406
parviflora, ii, 406
sinuata, ii, 406
tricuspidata, ii, 406
tristis, ii, 406
Matonia, ii, 542, 565
pectinata, ii, 531, 542
Mazus, ii, 505
rugosus, ii, 505
Meioneetes, i, 21
Meisarrhena tomentosa, i, 93
Melaleuca, i, 19, 311; ii, 451
armillaris, ii, 453
Cajeputi, i, 19
calycina, ii, 455
coronata, ii, 454
cylindrica, ii, 710
decussata, ii, 454
densa, ii, 456
diosmifolia, ii, 452
ericifolia, ii, 453
fulgens, ii, 455
genistifolia, ii, 452
globifera, ii, 452
gnidiæfolia, ii, 454
hypericifolia, ii, 455
incana, ii, 684
lanceolata, ii, 710
laurina, ii, 456
Leucadendron, i, 19; ii, 451
linariifolia, ii, 455
myrtifolia, ii, 455
nerifolia, ii, 456
nodosa, ii, 453
paludosa, ii, 451
pulchella, ii, 454
salicifolia, ii, 456
scabra, ii, 454
squamea, ii, 453
squarrosa, ii, 455
striata, ii, 452
stypelioides, ii, 452
thymifolia, ii, 454
thymoides, ii, 452

Melaleuca uncinata, ii, 454
viridiflora, ii, 451
Melampodium, ii, 284
americanum, ii, 285
australe, ii, 284
humile, ii, 284
longifolium, ii, 284
Melanthera, ii, 291, 296
aspera, ii, 306, 307
Melanthaceæ, i, 298, 299 *note*
Melanthera, ii, 296
Melanthium punctatum, i, 298
Melastomaceæ, i, 116
Melastoma decumbens, i, 116
octandra, i, 117
Melhania, ii, 509
Denhamii, i, 288
Erythroxylon, ii, 509
Melunoxylon, ii, 509
Meliaceæ, i, 72, 123, 124, 149
Melica, i, 56
Melicocœa, i, 108
Melicope, i, 16
Melocactus, i, 549
Memecylon, i, 118
Meniocus, i, 265, 268
Meniscium, ii, 566, 569
Menispermæ, i, 379; ii, 652
Menonvillea, i, 264
Merendera, i, 298
Merriana, i, 118
Mertensia, i, 138
dichotoma, i, 146 *note*, 164
Mesembryanthemum, i, 64, 556
æquilaterale, i, 43, 309
cordifolium, i, 556
crystallinum, i, 556
nodiflorum, i, 556
papulosum, i, 556
Mesochlœna, ii, 569
Mesodactylis deflexa, ii, 547
Mestotes, i, 125
Metalasia, ii, 302
Metaplexis, ii, 224
Metastelma, ii, 227, 229
Metrosideros, i, 19
quinquenervia, ii, 451
Meyera, ii, 283
Micadania, i, 295
Micraura, ii, 644
ternata, ii, 644
Micranthus, i, 134 *note*
Microchloa elongata, ii, 533
Microcorys, i, 39
Microloma, ii, 227, 229

- Microlooma lineare*, ii, 228
 sagittatum, ii, 228
Micropus, ii, 285
Microstemma, ii, 200, 208
Microtea, i, 139, 139 *note*
Mikania chenopodiifolia, i, 128
Millet, i, 160
Miltus, i, 139 *note*
Mimetes, ii, 20, 36, 86, 186
 capitulata, ii, 87
 cucullata, ii, 88, 190
 divaricata, ii, 91
 Hartogii, ii, 88, 190
 Hibbertii, ii, 89
 hirta, ii, 86, 191
 Massoni, ii, 89
 myrtifolia, ii, 90
 pauciflora, ii, 87
 purpurea, ii, 91
 thymelæoides, ii, 90
Mimosa, i, 294
 aspera, i, 110
 canescens, i, 110
 decipiens, ii, 446
 Habbas, i, 288
 nigricans, ii, 447
 pubescens, ii, 447
 Sophoræ, ii, 446
 stellata, i, 23
 taxifolia, i, 290 *note*
Mimoseæ, i, 22, 110, 288
Miquelia, ii, 652, 653
 cærulea, ii, 607
Mirbelia, ii, 435
 dilatata, ii, 436
 reticulata, ii, 435
Misodendron, i, 413 *note*
Mitella grandiflora, ii, 522
Mitrasacme, i, 37
Mnesithea, ii, 553
Mnium rostratum, i, 235
 turgidum, i, 178, 235
Modecca, i, 150, 387 *note*; ii, 686
 australis, ii, 705
Molina, ii, 294
Molucella integrifolia, i, 94
 repanda, i, 94
 scariosa, i, 94
Monacchine, i, 57
Monimia, i, 25
Moninnicæ, i, 25
Monkey Pepper, i, 161
Monneria, i, 16
Mounia, i, 14
Monocaryum, i, 300
Monodora Myristica, i, 162 *note*
Monogamia, ii, 276
Monophyllea, ii, 595, 608
 Horsfieldii, ii, 608
Moquileca, i, 115
Morcea, i, 559
Moriuda, i, 131
Moringa, i, 294
Moringæ, i, 294
Mosses, i, 469
Murucua, i, 386
Musa, i, 156, 157 *note*
 sapientum, i, 154, 156
 superba, i, 157 *note*
Musanga, i, 138, 153
Musci, i, 66
Myagrum, ii, 371
 arborescens, ii, 371
 hispanicum, ii, 412
 paniculatum, ii, 372
 perenne, ii, 369
 perfoliatum, ii, 371
 rugosum, ii, 369
 sativum, ii, 386
 saxatile, ii, 384
Myoporinæ, i, 339
Myoporum, i, 40, 63
Myrianthus, i, 138
Myrioneura nutans, ii, 534
Myriophyllum, i, 21, 22
Myristica, i, 381
Myrobroma fragrans, ii, 498
Myrospermum, i, 127 *note*
Myroxylon, i, 127 *note*
Myrsine, i, 150 *note*
Myrsinæ, i, 149, 293; ii, 536
Myrtaceæ, ii, 534
Myrtaceæ, i, 18, 20, 63, 75, 118, 307, 311
Mystropetalon, i, 414 *note*
Myzodendron, i, 413 *note*
 brachystachyum, i, 413 *note*
 punctulatum, i, 413 *note*
 quadriflorum, i, 413 *note*
Nama, i, 135
Nandina, i, 124 *note*
Nausiatum, ii, 651, 653
Napimoga, i, 120
Napolcona, i, 387, 388 *note*
Nardus, ii, 553
Nasturtium, ii, 399
 amphibium, i, 68; ii, 399
 officinale, ii, 399

- Nasturtium pyrenaicum*, ii, 399
 sagittatum, ii, 400
 sylvestre, ii, 399
 terrestre, ii, 399
Natter Wurtz, i, 214
Nauclea, i, 131
 Wallichiana, ii, 533
Neckera, ii, 335
 heteromalla, i, 69
 pennata, i, 69
Neli Tali, ii, 445
Nelsonia, i, 134
Nemophila, ii, 519
Neottæ, i, 538
Ncottia, ii, 481
 acaulis, ii, 481
 cernua, ii, 482
 clata, ii, 481
 glandulosa, ii, 482
 minor, ii, 482
 orchioides, ii, 481
 picta, i, 501; ii, 481
 pubescens, ii, 481
 repens, ii, 481
 speciosa, ii, 481
 spiralis, ii, 482
Nepenthes, i, 77, 386, 449; ii, 357
Nepeta azurea, i, 94
 indica, ii, 503
 virginica, ii, 504
Nephrodium, i, 61, 146; ii, 569
 fragrans, ii, 518
 rufidulum, ii, 518
Nerium, ii, 233, 242, 686
 antidysentericum, ii, 242, 244
 coronarium, ii, 242
 divaricatum, ii, 242
 obesum, ii, 242
 odorum, ii, 242
 Oleander, ii, 242, 243
 salicinum, ii, 242
 Zeylanicum, ii, 242, 243, 244
Nety, i, 289
Neurachne, i, 58
 alopescuroides, i, 337
 Mitchelliana, i, 337
 paradoxa, i, 336
Neurada prostrata, i, 288
Neurocarpæa, i, 131
 lanceolata, i, 94
Neurolæna, ii, 299
Nicandra, i, 133
Nicotiana, i, 158
 Australasica, i, 158
 undulata, i, 158
Niphobolus, ii, 568
Nisa, i, 120
Nitta, i, 289
Nivenia, ii, 20, 36, 110, 187
 capitata, ii, 115
 erithimifolia, ii, 113
 Lagopus, ii, 114
 marginata, ii, 111
 media, ii, 113
 mollissima, ii, 114, 187
 parvifolia, ii, 112
 sceptrum, ii, 111
 spathulata, ii, 111
 spicata, ii, 112
Notoceras, ii, 404
 canariensis, ii, 404
 mollis, ii, 274
Nuphar, i, 379 *note*, 451
Nürvala, i, 278
Nuxia congesta, i, 93
 dentata, i, 93
Nuytsia floribunda, i, 308
Nyctagineæ, ii, 587 *note*
Nyctagineæ, i, 81
Nymphæa, i, 74, 379 *note*, 451
 Lotus, i, 162, 165, 166
 pubescens, i, 165
Nymphæaceæ, i, 451, 555

Ochradenus, i, 281
Octadenia, i, 267
Octoblepharum, ii, 337, 339
 albidum, i, 165; ii, 337
 serratum, ii, 337
Octodiceras, ii, 334
Octomeria, ii, 491
 graminifolia, ii, 491
Ocymum, i, 135
 cinereum, i, 94
 monadelphum, i, 94
Ænothera, i, 467
Ogheglie, i, 155
Olacinae, i, 136
Olacinea ? ii, 538
Olax, i, 44, 136
 scandens, i, 136
 zeylanica, i, 136
Oldenlandia, ii, 532
 biflora, ii, 532
 herbacea, ii, 532
 pentandra, ii, 522
Oleina, ii, 535
Olmedia, i, 79
Omphalobium, i, 113
Onagrarice, i, 21, 467

- Oncidium*, i, 515; ii, 494, 696
 altissimum, ii, 494
 bifolium, ii, 494
 carthagense, ii, 494
 triquetrum, ii, 495
Oncoba, i, 108
Onobrychis simplicifolia, i, 95
Onoclea, ii, 525
Opercularia, i, 36, 131, 132
 volubilis, ii, 711
Ophiopogon spicatus, ii, 695
Ophiorrhiza bracteolata, ii, 534
 discolor, ii, 534
Ophiospermum, i, 126 *note*
Ophioxylum, i, 132
Ophrydæ, i, 538
Ophrys, i, 493, 538; ii, 479
 anthropophora, ii, 475
 apifera, i, 491, 502, 508, 540
 note, 547; ii, 479
 aranifera, ii, 479
 corallorhiza, ii, 490
 cordata, ii, 483
 lilifolia, ii, 489
 Loeselii, ii, 489
 Monorchia, ii, 476
 muscifera, ii, 479
 myodes, ii, 479
 ovata, ii, 483
 spiralis, ii, 482
Opilia, ii, 536
Orange, i, 154, 156
Orchidæ, i, 48, 53, 62, 380, 405, 469,
 487, 545, 557, 560; ii, 6, 547
Orchis, i, 493; ii, 473
 albida, ii, 477
 bicornis, ii, 479
 bifolia, ii, 477
 bractealis, ii, 476
 bracteata, ii, 476
 Burmanniana, ii, 478
 carnea, ii, 480
 ciliaris, ii, 477
 conopsea, ii, 475
 cristata, ii, 477
 fimbriata, ii, 477
 fusca, i, 547; ii, 474
 globosa, ii, 474
 hircina, ii, 474
 hyperborea, ii, 477
 latifolia, i, 500; ii, 474, 540
 mascula, ii, 473, 475, 491, 500,
 502
 militaris, ii, 474
 Morio, i, 500, 540, 547; ii, 473
Orchis nigra, ii, 476
 papilionacea, ii, 475
 pectinata, ii, 478
 pyramidalis, ii, 474
 spectabilis, ii, 475
 ustulata, i, 547; ii, 473
 viridis, ii, 476
Ordeal tree, i, 153
Oreobolus, i, 53
Oreocallis, ii, 38, 165
 grandiflora, ii, 166
Orites, ii, 38, 159
 diversifolia, ii, 160
 revoluta, ii, 160
Oritina acicularis, ii, 189
Ormosia, ii, 420
 dasycarpa, ii, 420
Ornithidium, ii, 490
 coccineum, ii, 490
Orobanchæ compacta, i, 296
Orobanchæ, i, 557
Oropetium, ii, 553
Orthodon, ii, 337, 338
Orthotrichum, ii, 344, 345, 348, 349
 cupulatum, i, 178
Oryzopsis asperifolia, ii, 514
 parviflora, ii, 514
Osbeckia, i, 117
Osmunda barbara, i, 64
Osmundacæ, i, 146
Othra, i, 150 *note*
Oudneya, i, 272 *note*
 africana, i, 272 *note*
Ovieda mitis, ii, 509
Ovvando, i, 159
Oxalidæ, i, 114
Oxalis, i, 64, 113
 corniculata, i, 65
 sensitiva, i, 114
Oxylobium, ii, 426
 arborescens, ii, 426
 cordifolium, ii, 426
 ellipticum, ii, 426
Oxypetalum, ii, 219
Oxyria, i, 177 *note*, 214
 reniformis, ii, 215
Oxystelma, i, 134, 295; ii, 218
 Bornouense, i, 295 *note*
 esculentum, i, 134, 165, 295;
 ii, 219
Oxytropis arctica, i, 209, 254
 uralensis, i, 210
Ozothamnus, ii, 303
Pachynema complanatum, ii, 703

Pæonia Moutan, ii, 318
Pagamea, i, 37, 132
Paliurus, i, 26
Palmæ, i, 51; ii, 587 *note*
Pancratium, i, 364
Pandanus, ii, 718
 Candelabrum, i, 153
Panicæ, i, 57, 145, 300; ii, 553, 572
Panicum, i, 55, 57, 146
 crus-galli, i, 68, 165
 ovale, i, 93
 turgidum, i, 303
Papaver nudicaule, i, 178, 190, 379
 note
 radicatum, i, 190
Papaveracæ, i, 190, 272, 559
Papaw, i, 154, 155, 156, 158
Papilionacæ, i, 23, 24, 209, 295, 339
Pappophorum, i, 55
Paranomus, ii, 110
Parinarium, i, 115
Parkeria, ii, 570
Parkia, i, 289
 africana, i, 290 *note*
Parmelia caperata, i, 69
 conspersa, i, 69
 olivacea, i, 69
 parietina, i, 69
 physodes, i, 69
 plumbca, i, 69
 stellaris, i, 69
Parnassia, i, 559
Paronychia sedifolia, i, 94
Paropsia, i, 121, 387 *note*
Parrya, i, 195, 268, 271, 272 *note*
 arctica, i, 196
 macrocarpa, i, 197
Parsonsia, ii, 232, 237, 238
Paspalum, i, 55, 57
Passalia, i, 123
Passifloreæ, i, 121, 149, 280, 282,
 384, 386, 392
Passura, i, 123
Patrisia, i, 125
Paullinia, i, 164
 japonica, i, 107
 pinnata, i, 107, 165, 172
Pavetta, ii, 269 *note*
 canescens, ii, 534
 congesta, i, 93
 lucens, ii, 533
 mollis, ii, 534
 nauchleiflora, ii, 533
 polyantha, ii, 533
 reflexa, i, 93

Pavetta webernefolia, ii, 534
Pavonia, i, 25, 64
Pedicularis arctica, i, 213, 253
 hirsuta, i, 178
 Langsdorffii, i, 253
 sudetica, i, 214, 253
Peganum, i, 16
Pelargonium, i, 64
 abyssiunicum, i, 95, 163
Peliosanthes Teta, i, 362; ii, 695
Peltaria, ii, 374
 alliacea, ii, 374
Peltidea aphthosa, i, 251
 canina, i, 69
Penæa, i, 64
Pennisetum dichotomum, i, 302
 glaucum, i, 68
 villosum, i, 93
Pentadactylon, ii, 133
Pentadynamis, i, 325
 incana, i, 325
Pentaloba sessilis, i, 123
Pentapetes Erythroxylon, ii, 509
Pentatropis cynanchoides, i, 94
Peperomia, i, 150
Pergularia, ii, 6, 210, 212, 237
 edulis, ii, 212
 glabra, ii, 212
 Japonica, ii, 212
 minor, ii, 212
 odoratissima, ii, 212
 purpurea, ii, 212
Periploca, i, 134; ii, 198, 231
 angustifolia, ii, 231
 capsularis, ii, 238
 emetica, ii, 230
 esculenta, i, 165; ii, 219
 græca, ii, 231
 indica, ii, 230
 lævigata, ii, 231
 Secamone, ii, 230
 sylvestris, ii, 214
 tunicata, ii, 222
Periplocæ, i, 469; ii, 230
Perripterygium quinquelobum, ii, 655
Perotis latifolia, i, 165
Persoonia, i, 116, 339; ii, 11, 17,
 18, 20, 22, 23, 24, 37, 133
 articulata, ii, 137
 elliptica, ii, 137
 falcata, ii, 136
 flexifolia, ii, 135
 furruginea, ii, 136
 graminea, ii, 137
 hirsuta, ii, 134

- Persoonia juniperina*, ii, 134
lanceolata, ii, 136
latifolia, ii, 136
laurina, ii, 136
linearis, ii, 134
longifolia, ii, 137
lucida, ii, 135
microcarpa, ii, 134
mollis, ii, 134
nutans, ii, 135
pinifolia, ii, 134
prostrata, ii, 136
salicina, ii, 136
scabra, ii, 135
spathulata, ii, 135
teretifolia, ii, 133
virgata, ii, 135
Petaloma, i, 118
Petalostemma chenopodii, i, 94
Petalostylis, i, 328
labicheoides, i, 328
Petiveria, i, 139
Petrobium, ii, 292 *note*, 314
Petrocallis, i, 268; ii, 386
pyrenaica, ii, 386
Petrophila, i, 310; ii, 11, 17, 35, 55,
723
acicularis, ii, 56
diversifolia, ii, 57, 723
filifolia, ii, 56, 57
pedunculata, ii, 57
pulchella, ii, 56
rigida, ii, 56
squamata, ii, 57
teretifolia, ii, 56
trifida, ii, 57
Peziza scutellata, i, 70
Phaca astragalina, i, 209
Phacelia, ii, 519
parviflora, ii, 520
Phalænopsis, ii, 576
amabilis, ii, 576
Phalaridæ, ii, 571
Phascom, i, 247
alternifolium, i, 348
bryoides, i, 247
curvicollum, i, 248
flexuosum, i, 248
muticum, i, 69
Phaseolus, i, 154
Phayloopsis, i, 13
Phebalium, i, 17
Phelpæa sanguinea, i, 429
Philydram, i, 48, 53; ii, 357
lanuginosum, i, 53
Philydram pygmæum, i, 53
Phippsia, i, 219
algida, i, 220
Phlebodium, ii, 567
Phleum alpinum, i, 67
Phragmites, i, 58
Phylca, i, 26
Phyllanthus, ii, 279 *note*
Phyllocladus, i, 460
Physcia islandica, i, 251
Physematium molle, ii, 546
Physiphora, i, 123
Phytocrene, ii, 651, 652
Phytocrenæ, ii, 652
Phytolacca, i, 139
abyssinica, i, 139
Phytolaccæ, i, 138
Piarranthus, ii, 200, 206
pulla, ii, 206
punctata, ii, 206
Picris hieracioides, i, 68
Pileanthus, i, 76
Pilocarpus, i, 16
Pilostyles, i, 414 *note*, 424, 430
Berterii, i, 430
Blanchetii, i, 430
Calliandræ, i, 431
Pimelea, ii, 16
Pine apples, i, 154, 155
Pinus, i, 47, 455
Piper Æthiopicum, i, 105, 161
caudatum, i, 150
Cubeba, i, 150
Piperacæ, i, 138, 149, 452
Piptocarpha, ii, 299 *note*
Pisonia morindifolia, ii, 534
Pistia Stratiotes, i, 164, 166, 169, 298
Pittosporæ, i, 13, 63
Pittosporum, i, 13
revolutum, ii, 268 *note*
Plagioloba, i, 272
Plagiopoda, ii, 146
Planta Aphyteja, i, 416
Plantago, ii, 281
Plantain, i, 154, 156
Platylobium, i, 25; ii, 436
formosum, ii, 436
lanceolatum, ii, 438
microphyllum, ii, 438
ovatum, ii, 438
parviflorum, ii, 437
Scolopendrium, ii, 437
triangulare, ii, 437
Platypetalum, i, 192, 193
dubium, i, 193, 253

- Platypetalum purpurascens*, i, 192, 254
Platystemma, ii, 595, 608
Platyzoma, i, 59
Pleopeltis, ii, 566, 567
Pleurandra, i, 12
 acerosa, ii, 704
 bracteata, ii, 703
 cinerea, ii, 703
 furfuracea, ii, 703
 nitida, ii, 703
 parviflora, ii, 703
 pedunculata, ii, 704
 riparia, ii, 704
 scabra, ii, 704
 sericea, ii, 703
 stricta, ii, 704
Pleurogramma, ii, 569
Pleuropogon, i, 226
 Sabinii, i, 226, 256
Pleurothallis, i, 515; ii, 491
 ruscifolia, ii, 491
Pluchea Dentex, ii, 711
Plumbaginæ, i, 297
Plumbago eglandulosa, i, 94
Plumeria, ii, 247
Poa, i, 221
 abbreviata, i, 223, 255
 alpina, ii, 515
 angustata, i, 223, 255
 arctica, i, 224
 ciliaris, i, 165
 crocata, ii, 515
 flexuosa, i, 225
 laxa, i, 177, 224, 225
Poaceæ, i, 53, 145, 228, 300; ii, 553
Podalyria, ii, 422, 423
 alba, ii, 423
 argentea, ii, 424
 aurea, ii, 421
 australis, ii, 423
 biflora, ii, 424
 bracteata, i, 111
 calyptata, ii, 424
 capensis, ii, 422
 cordata, ii, 424
 genistoides, ii, 422
 hirsuta, ii, 424
 lupinoides, ii, 421
 myrtillifolia, ii, 423
 sericea, ii, 424
 tinctoria, ii, 423
Podocarpus, i, 44, 47, 455, 460
 alpina, ii, 707
 antillarum, ii, 707
 asplenifolia, i, 47
 cupressina, ii, 706
 elata, ii, 707
 ensifolia, ii, 706
 falcata, ii, 707
 Horsfieldii, ii, 533
 latifolia, ii, 707
 Nageia, ii, 706
 polystachya, ii, 706
 spicata, ii, 707
 spinulosa, ii, 707
 thuyoides, ii, 707
Podocarya, ii, 717
Podocoma, i, 329
 acuminata, ii, 462
 cuneifolia, i, 329
Podolepis, ii, 461
 rugata, ii, 461
Podolobium, ii, 425
 trilobatum, ii, 425
Podophyllum, i, 124 *note*
Podopogon, ii, 572
Podostemon Wallichii, ii, 533
Podostigma, i, 520
Pœcilorodermis, ii, 637, 638
Pogonia, ii, 485
 divaricata, ii, 485
Pohlia, i, 236
 arctica, i, 236
 bryoides, i, 236
 purpurascens, i, 237
Poiretia linearis, ii, 440
Polanisia, i, 273
Polemoniaceæ, i, 135
Polemonidææ, i, 65
Polemonium dubium, ii, 520
Polygala abyssinica, i, 95
 linearis, i, 95
Polygalæ, i, 13, 16, 63, 125, 282,
 291, 380
Polygamia congregata, ii, 276
 segregata, ii, 276
Polygonæ, i, 138, 214
Polygonum sinuatum, i, 94
 viviparum, i, 214
Polyosma Cunninghamii, ii, 708
Polypodiaceæ, ii, 525
Polypodium, i, 166; ii, 205
 arvonicum, ii, 254
 aureum, ii, 567
 decumanum, ii, 567
 Dipteris, ii, 562
 diversifolium, ii, 566
 Dryopteris, ii, 518
 Horsfieldii, ii, 531, 562
 hyperboreum, ii, 252

- Polypodium ilvense*, ii, 252
 lycopodioides, ii, 566
 melanopus, ii, 531
 phymatodes, ii, 543, 566
 propinquum, ii, 531
 quercifolium, ii, 566
 spinulosum, i, 83
 vulgare, ii, 518, 570
 Wallichii, ii, 531, 562
Polystichum, ii, 569
Polytoca, ii, 574
 bracteata, ii, 574
Polytrichoides, ii, 336, 351
Polytrichum, i, 348; ii, 331, 345, 349, 351
 alpinum, i, 234; ii, 336
 angustatum, ii, 338
 brevifolium, i, 233
 commune, i, 233; ii, 339
 hyperboreum, i, 233
 juniperinum, i, 178
 lævigatum, i, 234; ii, 337, 340
 magellanicum, ii, 337
 microstomum, ii, 339
 piliferum, i, 233
 propinquum, i, 233
 septentrionale, i, 234
 subrotundum, ii, 338
 undulatum, ii, 336, 338, 340
 urnigerum, ii, 339
Pomaderris, i, 26
Pomax, i, 36, 132
Pombalia, i, 122
Ponthieva, ii, 482
 glandulosa, ii, 482
Portlandia, ii, 268
Portulaca oleracea, i, 68
Potalia, i, 133
Potamogeton crispum, i, 68
 gramineum, i, 66, 68
 natus, i, 68
 perfoliatum, i, 68
Potentilla anserina, i, 68
 Brauniana, i, 178
 frigida, i, 178, 209
 grœnlandica, i, 178, 209
 hirsuta, i, 209
 Jamesoniana, i, 209
 macrantha, i, 209
 nivca, i, 208
 pulchella, i, 178, 208
 sericea, i, 178, 208
 Vahlana, i, 209
 verna, i, 209
Poterium, ii, 279
Poupartia, i, 112
Prestonia tomentosa, ii, 241
Primulacæ, i, 151 *note*, 293; ii, 315, 581, 586
Prionopteris, ii, 531
Proserpinaca, i, 21
Prosopis, i, 22
Prostanthera, i, 39
Protea, ii, 10, 14, 15, 20, 24, 26, 29, 35, 40, 60, 185, 272
 abrotanifolia, ii, 103, 186
 abyssinica, i, 163; ii, 70
 acaulis, ii, 28, 72
 acerosa, ii, 78
 acuifolia, ii, 71
 adscendens, ii, 104
 alba, ii, 46
 alopeceuiroides, ii, 111
 amplexicaulis, ii, 77
 anemonifolia, ii, 59
 anethifolia, ii, 58
 angustata, ii, 73
 angustifolia, ii, 70
 arcuata, ii, 47; β , ii, 47
 argentea, ii, 12, 24, 25, 27, 30, 31, 42; β , ii, 50
 argentiflora, ii, 108
 aulacea, ii, 40
 barbata, ii, 64
 bracteata, ii, 40
 bruniades, ii, 46
 cæspitosa, ii, 76
 canaliculata, ii, 72
 cancellata, ii, 28, 30
 candicans, ii, 83, 187
 caudata, ii, 124, 125
 Chamelæa, ii, 48
 cinerea, ii, 45, 46
 coccinea, ii, 63
 comosa, ii, 52
 compacta, ii, 62
 concaua, ii, 187
 conica, ii, 49
 conifera, ii, 27, 28, 49; A, ii, 49; α , ii, 50
 conocarpa, ii, 26, 28, 30, 31, 81; A, ii, 80
 cordata, ii, 77
 cordifolia, ii, 77
 coronata, ii, 64, 65
 corymbosa, ii, 46
 criniflora, ii, 85
 crinita, ii, 85
 cristata, ii, 66

Protea cucullata, ii, 88 ; β , ii, 89
cyanoides, ii, 97, 102
cynaroides, ii, 61, 185
decumbens, ii, 100, 104
dichotoma, ii, 56, 186
divaricata, ii, 91, 185
elliptica, ii, 80
elongata, ii, 73
florida, ii, 104
formosa, ii, 64, 80
fucifolia, ii, 56, 183, 184
fusca, ii, 27, 45
glabra, ii, 184
glaucophylla, ii, 73
globosa, ii, 47
globularia, ii, 53
glomerata, ii, 28, 98, 101, 106,
 109
grandiflora, ii, 66, 69, 185
Gustaviana, ii, 112
heterophylla, ii, 85
hirta, ii, 30, 87
humiflora, ii, 77
humilis, ii, 77
hypophylla, ii, 31, 84
Hypophyllocarpodendron, ii, 84
imbricata, ii, 44, 118
incompta, ii, 67
incurva, ii, 52, 124
lacticolor, ii, 63
levis, ii, 74
Lagopus, ii, 113, 114
lanata, ii, 117
latifolia, ii, 61, 62
Laureola, ii, 47
Lepidocarpodendron, ii, 28 ; β ,
 ii, 64, 66
Lepidocarpon, ii, 65
Levisanus, ii, 28, 44, 45
linearis, ii, 79, 183
linifolia, ii, 183
longiflora, ii, 62, 68, 185
lorea, ii, 76
macrophylla, ii, 64
melaleuca, ii, 65
mellifera, ii, 13, 68
mucronifolia, ii, 70
myrtifolia, ii, 90
nana, ii, 71, 73
nectarina, ii, 158
neriifolia, ii, 66
obliqua, ii, 42 ; β , ii, 47 ; α , ii,
 47
ochroleuca, ii, 63
odorata, ii, 184

Protea odoratissima, ii, 71
pallens, ii, 28, 48, 49
parviflora, ii, 28, 40
patens, ii, 67
patula, ii, 106
pedunculata, ii, 98
pendula, ii, 71
phylicoides, ii, 99, 100
pinifolia, ii, 39
pinnata, ii, 96
plumosa, ii, 42
procumbens, ii, 104
prolifera, ii, 122
prostrata, ii, 187
pubera, ii, 82
pulchella, ii, 56, 66
purpurea, ii, 27, 91
racemosa, ii, 25, 121, 122
repens, ii, 75, 77
revoluta, ii, 73
rosacea, ii, 71
saligna, ii, 25, 27, 28, 50, 51
scabra, ii, 74
Sceptrum, ii, 111, 112
Scolopendrium, ii, 76
Scolymus, ii, 70
sericea, ii, 53
Serraria, ii, 26, 108
spatulata, ii, 111, 112
speciosa, ii, 28, 63, 66, 67
speciosa nigra, ii, 65
sphærocephala, ii, 97, 103, 186 ;
A, ii, 98
spicata, ii, 112, 113
strobilina, ii, 28, 47, 184
tenax, ii, 72
tenuifolia, ii, 74
teretifolia, ii, 52
thyrsoides, ii, 109
tomentosa, ii, 83
torta, ii, 45
totta, ii, 28, 79
tridactylites, ii, 59
tritermata, ii, 108, 186
turbiniflora, ii, 76
vestita, ii, 80
villosa, ii, 100, 186
villosiuscula, ii, 82
virgata, ii, 78
Proteaceæ, i, 21, 36, 40, 63, 80, 81,
 307, 310, 339, 449, 501 ; ii, 5,
 271, 317
Protææ, ii, 9
Præstonia, ii, 233, 241
Primulacæ, i, 297

- Prunella vulgaris*, i, 68
Psilotum, ii, 722
Psilurus, ii, 553
Psoralea, i, 322
Psychotria, i, 131
Ptelidium, i, 107
Pteranthus, ii, 543
Pteris, i, 146; ii, 250
 crispa, ii, 525
 esculenta, i, 309
 gracilis, ii, 518
 thalictroides, ii, 526
Pterocarpus, i, 106, 111, 290
 Ecastophyllum, i, 111, 165, 172
 erinacea, i, 290
 lunatus, i, 111, 165, 172
 senegalensis, i, 200
Pterocymbium, ii, 611, 641
 javanicum, ii, 611
Pterogonium declinatum, i, 334
Pterolobium lacerans, i, 94
Pterostylis, ii, 486, 549
 obtusa, ii, 486
Pterygodium, ii, 480
 Volucris, ii, 480
Pterygota, ii, 620, 638
 alata, ii, 639
Pterygandrum, ii, 345
Ptilotus latifolius, i, 336
Ptychocarpa, ii, 144
Ptychocarya, ii, 532
Ptychostomum pulchellum, i, 237
Pulicaria aromatica, i, 95
 involuta, i, 95
 viscida, i, 95
Pultenæa, ii, 433
 acerosa, ii, 709
 biloba, ii, 670
 daphnoides, ii, 433
 ericoides, ii, 430
 flexilis, ii, 434
 hispidula, ii, 709
 illicifolia, ii, 425
 linophylla, ii, 433
 nana, ii, 425
 obcordata, ii, 433
 retorta, ii, 431
 retusa, ii, 433
 rigida, ii, 709
 rubinefolia, ii, 436
 scabra, ii, 433
 stipularis, ii, 434
 tenuifolia, ii, 670
 vestita, ii, 434, 670
 villosa, ii, 434
Pultenæa viscosa, ii, 709
Pumpkins, i, 154
Pupalia, ii, 544
Pycnanthemum, ii, 503
 aristatum, ii, 504
 incanum, ii, 503
 virginicum, ii, 504
Pycnospora, ii, 708
 hedysaroides, ii, 708
Pyrenacantha, ii, 652, 653
Pyrethrum, ii, 691
Pyrola rotundifolia, i, 177

Quadria, ii, 138
 heterophylla, ii, 139
Quinchamalium, i, 137
Quirivelia Zeylanica, ii, 236

Racomitrium lanuginosum, i, 238
Rafflesia, i, 374, 558; ii, 358
 Arnoldi, i, 374, 401, 415, 424;
 ii, 724
 Cumingii, 405 *note*, 425; ii, 724
 Horsfieldii, i, 390 *note*, 425
 Manillana, i, 405 *note*, 425
 Patma, i, 402, 424; ii, 724
 Titan, i, 415
Rafflesia, i, 424
Rafflesiaceæ, i, 423
Rafnia perfoliata, ii, 423
 retusa, ii, 439
Ramalina fastigiata, i, 70
 fraxinea, i, 70
Ranunculaceæ, i, 65, 188, 318
Ranunculus affinis, i, 189, 253
 auricomus, i, 189, 253
 collinus, ii, 703
 frigidus, i, 188
 glacialis, i, 178
 hyperboreus, i, 189
 inundatus, ii, 703
 Laxmanni, i, 188
 nivalis, i, 188, 253
 plebeius, ii, 703
 Pumilio, ii, 703
 pygmæus, i, 188
 Sabinii, i, 189, 253
 sessiliflorus, ii, 703
 Smithii, i, 188
 sulphureus, i, 178, 188
 Willdenovii, i, 188
Raphanus, ii, 412
 erucoides, ii, 410
 maritimus, ii, 412
 Raphanistrum, ii, 412

Raphanus sativus, ii, 412
 tenellus, ii, 413
 Raphia vinifera, i, 142, 153, 154
 Rapinia herbacea, i, 171 *note*
 Rapistrum, ii, 372
 ægyptiacum, ii, 372
 paniculatum, ii, 372
 rugosum, ii, 319
 Ratzburgia, ii, 552
 pulchella, ii, 552
 Ravenala, i, 157 *note*
 Red Snow, i, 178
 Red Water Tree, i, 153
 Reevesia Wallichii, ii, 632
 Reimaria, i, 57
 Renanthera coccinea, i, 548
 Reseda, i, 363
 odorata, i, 284
 pedunculata, i, 94
 propinqua, i, 281
 suffruticulosa, i, 281
 undata, i, 281
 Resedaceæ, i, 65, 272, 273, 281
 Restiaceæ, i, 52, 53, 54, 63, 339; ii,
 278
 Rhabdotheramnus, ii, 594, 595, 608
 Solandri, ii, 608
 Rhamnaceæ, ii, 537
 Rhamnus, i, 12, 13, 26, 27, 114, 120
 Rhamnus, i, 26
 inebrians, i, 94
 Rhaphis, i, 302
 Rheum, i, 214
 digynum, i, 215
 Rhexia, i, 116
 Rhinanthaceæ, i, 14
 Rhinanthus, ii, 580
 Rhinoglossum obliquum, ii, 577
 Rhizophora, i, 20, 119, 364, 381
 setiformis, i, 70
 Rhizophoreæ, i, 20, 119
 Rhodolæna, i, 108
 Rhodoraceæ, i, 109
 Rhopala, ii, 13, 15, 18, 21, 22, 38,
 160, 532
 cochinchinensis, ii, 161
 dentata, ii, 162
 diversifolia, ii, 162
 media, ii, 161
 moluccana, ii, 161
 montana, ii, 160
 nitida, ii, 161
 peruviana, ii, 162
 serrata, ii, 162
 sessilifolia, ii, 163

Rhus, i, 112
 Rhynchothecum, ii, 609
 Rianza, i, 123
 Ribes, ii, 522
 Riccia fluitans, i, 69
 glauca, i, 69
 natans, i, 69
 Rice, i, 161
 Richea, ii, 286
 glauca, ii, 286
 Richcia, i, 119
 Ricinus, ii, 279
 Ricotia, i, 264; ii, 390
 ægyptiaca, ii, 390
 Rimorea, i, 123
 Ritcheia, i, 279
 Rivina, i, 139, 164
 octandra, i, 139 *note*
 Robergia, i, 112
 Robinia subdecandra, ii, 421
 Roccella, i, 252
 fuciformis, i, 69, 165
 Rohria, i, 126
 Rolandra, ii, 274
 Rondeletia febrifuga, i, 131
 Ropala nitida, ii, 161
 hameliæfolia, ii, 163
 Roridula, i, 94
 paradoxa, i, 94
 parviflora, i, 94
 Rosa abyssinica, i, 94
 Rosaceæ, ii, 536
 Rosaceæ, i, 65, 77, 109, 115, 120,
 288, 291; ii, 356
 Rottbölliacæ, ii, 553
 Rottböllia Cælorachis, ii, 553
 perforata, ii, 553
 Roupala, ii, 160
 montana, ii, 161
 pinnata, ii, 188
 sessilifolia, ii, 163
 Rourea, i, 112
 Rubiaceæ, i, 36, 37, 104, 130, 295;
 ii, 14, 195, 269 *note*, 314, 588
 note, 687
 Rubus compar, i, 94
 Ruellia balsamea, i, 135
 Blechum, ii, 506
 cristata, ii, 506
 infundibuliformis, ii, 505
 uliginosa, i, 135
 Ruizia, i, 25
 Rulingia, ii, 671, 672
 Rumex, i, 214
 digynus, i, 177, 215

- Rupala montana*, ii, 161
Ruta, i, 16
Rutacea, i, 285
Ryania, i, 121, 387 *note*

Saccharum, i, 57, 146
Sadoo, i, 94
Safu, i, 112, 154, 160
Sagonea palustris, i, 135
Sagus Palma-pinus, i, 142
Salacia, i, 107
Salicaria, i, 120
Salisburya, i, 460
Salix arctica, i, 178, 215
 curviflora, ii, 292
 hirsuta, ii, 292
 oleifolia, i, 379 *note*
 scandens, ii, 292
Salmca, ii, 291, 317
Salomonina, i, 14
Saltia, ii, 543, 544, 545
 abyssinica, i, 94; ii, 546
Salvia abyssinica, i, 93
Salvia, i, 380
Samolus, i, 65, 151 *note*, 293, 397
 ebracteatus, i, 297
 Valerandi, i, 68, 297
Samydeæ, i, 122, 126; ii, 535, 536, 537
Santalacæ, i, 20, 41, 44, 136, 413
 note, 501
Santalum, i, 42
Sautolina, ii, 26, 287, 291
Sapindacæ, i, 107; ii, 537, 659
Sapindus, i, 107
Sapotæ, i, 295
Sapria, i, 426
 Himalayana, i, 426
Sarcocephalus, i, 153, 155
Sarcobolus, ii, 214
Sarcophyte, i, 389
Sarcostemma, ii, 226
Sarcostigma, ii, 648, 651
 Horsfieldii, ii, 648
Sarmienta, ii, 586
Sarracenia, ii, 357
Satureja ovata, i, 94
 punctata, i, 94
Satyrion, i, 504; ii, 479
 albidum, ii, 477
 carneum, ii, 480
 cucullatum, ii, 479
 bians, ii, 694
 hirenium, ii, 474
 nigrum, ii, 476

Satyrion pustulatum, i, 497
 repens, ii, 481
 viridis, ii, 476
Saururus, i, 452
Savignya, i, 265 *note*, 271
 Egyptiaca, i, 262
Saxifraga, i, 201
 bulbifera, i, 205
 cæspitosa, i, 177, 205, 254
 cernua, i, 177, 205
 flagellaris, i, 177, 202
 foliolosa, i, 205
 Hirculus, i, 177, 202
 hyperborca, i, 204
 longiscapa, i, 205
 nivalis, i, 204
 oppositifolia, i, 177, 201
 petiolaris, i, 177, 204
 propinqua, i, 177, 202
 rivularis, i, 177, 204
 setigera, i, 177, 202
 stellaris, i, 205
 tricuspidata, i, 177, 203
 uniflora, i, 204, 254
 venosa, i, 204
Saxifragæ, i, 20, 65, 182 *note*, 201;
 ii, 522
Scabiosa, ii, 275, 278, 315
 atropurpurea, ii, 278
 succisa, ii, 278
Scævola, i, 34
 depauperata, i, 331
Scalia jaceoides, ii, 462
Scaphium, ii, 611
Schistostegia pennata, ii, 344
Schivereckia, i, 265
Schizopetalon, ii, 697
Schlotheimia, ii, 335
Schœnus nemorum, i, 54
Schwenckia, i, 164
 americana, i, 165, 167
Scitamineæ, i, 49, 53, 59, 380, 452,
 501; ii, 317, 548, 550
Scirpus, i, 300
 cæspitosus, ii, 513
 lacustris, i, 68; ii, 513
 maritimus, i, 68; ii, 513
 mucronatus, i, 68
 palustris, ii, 513
 sylvaticus, ii, 514
 triqueter, i, 68
Sclerachne, ii, 573
 punctata, ii, 573
Scleria macrocarpa, ii, 532
Sclerothamnus, ii, 432

Sclerothamnus microphyllus, ii, 432
 Scolopendrium, i, 60
 Scolymocephalus, ii, 30, 49
 Scoparia dulcis, i, 164, 167
 Scottia, i, 25; ii, 438
 dentata, ii, 439
 Scrophularinæ, i, 135, 213, 296, 532;
 ii, 585
 Secamone, ii, 229
 Securidaca, i, 14, 291
 Seetzenia africana, i, 286
 Seguieria, i, 139
 Selligera, ii, 566
 Semecarpus, i, 43, 112
 Sempervivum tectorum, i, 379 *note*
 Senna obovata, i, 294
 Senra, i, 317
 Sensitivæ, i, 114
 Serapias, ii, 478
 cordigera, ii, 478
 ensifolia, ii, 484
 grandiflora, ii, 484
 latifolia, ii, 484
 Lingua, ii, 478
 palustris, ii, 484
 rubra, ii, 485
 Sergilus, ii, 293
 Serpicula, i, 21
 Serraria, ii, 92
 Serratula tinctoria, ii, 301 *note*
 Serruria, ii, 14, 15, 17, 19, 36, 92,
 186
 acrocarpa, ii, 92, 186
 adscendens, ii, 104
 æmula, ii, 103
 Aitoni, ii, 94
 arenaria, ii, 96
 Bergii, ii, 186
 Burmanni, ii, 107
 candicans, ii, 107, 187
 ciliata, ii, 101
 compar, ii, 107
 congesta, ii, 101, 186
 erithimifolia, ii, 109
 cyanoides, ii, 96
 cygnea, ii, 92
 decipiens, ii, 106, 107
 decumbens, ii, 104
 diffusa, ii, 95
 elevata, ii, 94
 elongata, ii, 109
 flagellaris, ii, 105
 florida, ii, 104
 foeniculacea, ii, 101
 furcellata, ii, 97

Serruria glaberrima, ii, 92
 glomerata, ii, 106
 hirsuta, ii, 99, 186
 nitida, ii, 102
 Niveni, ii, 100
 pedunculata, ii, 98, 186
 phylicoides, ii, 103
 pinnata, ii, 95
 Roxburghii, ii, 107
 rubricaulis, ii, 105
 scariosa, ii, 97
 scoparia, ii, 98
 simplicifolia, ii, 95
 squarrosa, ii, 102
 Stilbe, ii, 99
 tritermata, ii, 108
 villosa, ii, 100
 Sesameæ, i, 295
 Sesamum pterospermum, i, 94, 295
 Sesban, ii, 443
 Sesbania, ii, 443
 aculeata, ii, 443
 ægyptiaca, ii, 443
 grandiflora, ii, 443
 Shea Tree, i, 295
 Shrubby Ilocus, i, 154, 160
 Sida, i, 109
 acuminata, i, 95
 gracilis, i, 95
 juncea, i, 165
 micropetala, ii, 709
 pannosa, i, 95
 periplorefolia, i, 165, 171
 Sideroxylum, i, 296
 Sieversia, i, 206
 humilis, i, 253
 Rossii, i, 207, 253
 Silene, i, 285
 acaulis, i, 177
 Siliquaria arabica, i, 275
 glandulosa, i, 94; ii, 415
 Silphium trifoliatum, ii, 263
 Simsia, i, 41; ii, 15, 17, 18, 37, 126
 anethifolia, ii, 127
 tenuifolia, ii, 127
 Sinapis, ii, 410
 alba, ii, 410
 Allionii, ii, 410
 arvensis, ii, 410
 chinensis, ii, 411
 erucoides, ii, 411
 frutescens, ii, 411
 hispanica, ii, 412
 incana, ii, 412
 juncea, ii, 411

- Sinapis lævigata*, ii, 411
millefolia, ii, 402
muralis, ii, 412
nigra, ii, 411
orientalis, ii, 410
pubescens, ii, 411
pyrenaica, ii, 401
tenuifolia, ii, 411
Siphonanthus angustifolia, ii, 509
indica, ii, 509
Siphonia, ii, 279
Sisymbrium, ii, 400
altissimum, ii, 401
amphibium, ii, 399
arenosum, ii, 396
asperum, ii, 401
austriacum, ii, 401
Columnæ, ii, 400
Irio, ii, 400
Loeselii, ii, 398
millefolium, ii, 402
molle, ii, 400
monense, ii, 409
murale, ii, 412
Nasturtium, i, 279; ii, 399
officinale, ii, 401
orientale, ii, 400
palustre, ii, 399
panuonicum, ii, 401
polyceratum, ii, 401
pyrenaicum, ii, 399
sagittatum, ii, 400
sinapoides, ii, 401
Sophia, ii, 402
supinum, ii, 401
strictissimum, ii, 402
sylvestre, ii, 399
tanacetifolium, ii, 402
tenuifolium, ii, 412
terrestre, ii, 399
Skitophyllum, ii, 334
Smcathmannia, i, 121, 150, 386
lævigata, i, 387
media, i, 387
pubescens, i, 387
Smithia, ii, 444
sensitiva, ii, 444
Sobreya, ii, 284
Sodada decidua, i, 279
Solanaceæ, ii, 267
Solanum, i, 381
cinereum, i, 94
uncinatum, i, 94
Solidago urticæfolia, ii, 289
Soliva, ii, 281
Soliva pedicellata, ii, 281
Sonchus oleraceus, i, 68
sibiricus, ii, 263
Sonerila arguta, ii, 533
grandiflora, ii, 533
tenera, ii, 533
Sophora, ii, 419
alopecuroides, ii, 420
australis, ii, 423
cærulea, ii, 423
cordata, ii, 424
flavescens, ii, 420
japonica, ii, 420
junca, ii, 429
lupinoides, ii, 421
microphylla, ii, 419
monosperma, ii, 420
occidentalis, ii, 420
sericea, ii, 424
tetraptera, ii, 419
tomentosa, ii, 420
Sorindeia, i, 112
Sorocephalus, ii, 36, 115
diversifolius, ii, 118
imberbis, ii, 116
imbricatus, ii, 118
lanatus, ii, 117
salsoloides, ii, 116
setaceus, ii, 116
spatilloides, ii, 117
tenuifolia, ii, 117
Southwellia, ii, 618
Sparmannia, i, 108
Spatalla, ii, 14, 15, 17, 19, 20, 21,
36, 115, 119
bracteata, ii, 121
brevifolia, ii, 125
caudata, ii, 124
iucurva, ii, 124
laxa, ii, 121
mollis, ii, 119
nivea, ii, 120
pedunculata, ii, 120
polystachya, ii, 123
prolifera, ii, 117, 122
propinqua, ii, 124
pyramidalis, ii, 123
ramulosa, ii, 120
sericea, ii, 122
Thunbergii, ii, 125
Spermaceæ, i, 131, 295
brachystema, ii, 711
lasiocarpa, ii, 531
longicaulis, ii, 531
membranacea, ii, 711

- Spermacoce purpureo-cerulea*, ii, 711
ramosissima, ii, 531
stricta, ii, 531
suffruticosa, ii, 711
tenera, ii, 531
tubularis, ii, 531
Spermadietyon suaveolens, ii, 534
Sphæria ophioglossoides, i, 70
Sphærolobium, ii, 429
grandiflorum, ii, 709
medium, ii, 430
minus, ii, 430
vimineum, ii, 429
Sphærophoror compressum, i, 70
coralloides, i, 70
Sphæropteris, ii, 540
Sphagnum capillifolium, i, 69
Sphæroclea zeylanica, i, 162, 165, 167, 170
Sphenogyne, ii, 465
abrotanifolia, ii, 466
anthemoides, ii, 465
erithmifolia, ii, 466
dentata, ii, 466
odorata, ii, 466
scariosa, ii, 466
Spilanthus, ii, 291, 317
arbores, ii, 292 *note*
atriplicifolius, ii, 290
tetraundrus, ii, 292 *note*
Splachnæ, ii, 337
Splachnum, i, 240; ii, 343, 347
angustatum, i, 243; ii, 337
arcticum, i, 243, 244, 253
exsertum, i, 245
Frælichianum, i, 241
longicollum, i, 241
luteum, ii, 337
magellanicum, i, 242
mnioides, i, 244, 246, 253
octoblepharum, i, 242
paradoxum, i, 245, 246
propinquum, i, 243, 244
rubrum, ii, 337
squamosum, ii, 337
tenue, i, 241
vasculosum, i, 243
Wormskioldii, i, 241, 243, 256
Sporobolus coromandelianus, ii, 532
diander, ii, 532
Staavia, ii, 327
Stachytarpheta cinerea, i, 93
indica, i, 136
Stackhousia, i, 27, 63
Stæhelia, ii, 285
Stapelia, i, 64; ii, 6, 200, 206, 675
adscendens, ii, 208
ambigua, ii, 207
articulata, ii, 207
Asterias, ii, 207
campanulata, ii, 206
chinensis, ii, 209
geminata, ii, 207
glandulifera, ii, 207
guttata, ii, 206
hirsuta, ii, 207
lepida, ii, 207
mixta, ii, 207
pedunculata, ii, 207
pulla, ii, 206
punctata, ii, 206
revoluta, ii, 207
sororia, ii, 207
stellaris, ii, 207
tuberosa, ii, 675
variegata, ii, 207
venusta, ii, 206
verrucosa, ii, 207
vetula, ii, 207
Statice Armeria, i, 297
Taxanthema, i, 297
Stauntonia, i, 379 *note*
angustifolia, ii, 533
latifolia, ii, 533
Staurothera, ii, 582, 591, 595, 606
cealearata, ii, 607
grandifolia, ii, 607
Stelis, i, 515; ii, 491
micrantha, ii, 491
ophioglossoides, ii, 491
Stellaria Edwardsii, i, 199, 254
nitida, i, 199, 254
Stellata, i, 36, 130
Stemodia, ii, 504
parviflora, ii, 504
Stenocarpus, i, 41; ii, 12, 38, 170
Forsteri, ii, 170
salignus, ii, 171
Stenochilus, i, 333, 339
curvipes, i, 334
glaber, i, 334
longifolius, i, 334
maculatus, i, 334
ochroleucus, i, 334
pubiflorus, i, 334
salicinus, i, 334
Stenopetalum, i, 192; ii, 705
lineare, ii, 705
Sterculia, ii, 614, 626

- Sterculia abyssinica*, i, 94; ii, 616, 627
acuminata, i, 153; ii, 618, 642
alata, ii, 620, 639
angustifolia, ii, 622 *note*, 633
Balanghas, ii, 614, 683
Candollei, ii, 636
caribæa, ii, 629
carthaginensis, ii, 622 *note*, 628
ceramica, ii, 636
Chickæ, ii, 628
coccinea, ii, 632
colorata, i, 363; ii, 618, 640
comosa, ii, 631
cordata, ii, 631
cordifolia, ii, 620, 642
cuaenta, ii, 634
cuspidata, ii, 635
ferruginea, ii, 634
fetida, ii, 614, 620, 622 *note*, 626
frondosa, ii, 629
fulgens, ii, 640
grandiflora, ii, 642
grandifolia, ii, 635
guttata, ii, 630
Helicteres, ii, 628
heterophylla, ii, 641
insularis, ii, 635
Ivira, ii, 629
Jackiana, ii, 634
Javanica, ii, 631
lævis, ii, 632
lanceæfolia, ii, 630
lanceolata, ii, 631
longifolia, ii, 631
macrocarpa, ii, 642
macrophylla, ii, 631
madagascariensis, ii, 630
mexicana, ii, 627
mollis, ii, 634
monosperma, ii, 633
nitida, ii, 642
nobilis, ii, 622 *note*, 633
oblongata, ii, 635
oblongifolia, ii, 642
obovata, ii, 636
parviflora, ii, 635
parvifolia, ii, 631
platanifolia, i, 363; ii, 616, 639
polyphylla, ii, 626
populifolia, ii, 636, 639
propinqua, ii, 629
punctata, ii, 628
Punduana, ii, 632
pyriformis, ii, 640
quadrifida, ii, 622 *note*, 636
- Sterculia Roxburghii*, ii, 630
rubiginosa, ii, 634
rugosa, ii, 630
scaphigera, ii, 640
Scaphium, ii, 640
Spangleri, ii, 632
stipularis, ii, 635
striata, 629
surinamensis, ii, 630
tomentosa, ii, 616, 628, 640, 643
Tragacanthæ, ii, 622 *note*, 636
Triphaca, ii, 628
urceolata, ii, 636
urens, ii, 622 *note*, 627
versicolor, 627
verticillata, ii, 642
villosa, ii, 622 *note*, 623, 627
Wallichii, ii, 640
- Sterculiaceæ*, i, 11, 384, 388; ii, 611
Sterculicæ, ii, 614, 620, 624
Stereocaulon paschale, i, 70, 252
Stereoxylon, ii, 523
Sticta crocata, i, 69
pulmonacea, i, 69
scrobiculata, i, 69
Stigmaria ficoides, ii, 721
Stipa, i, 56
canadensis, ii, 514
juncea, ii, 514
Strelitzia, i, 157 *note*
Streptocarpus, ii, 590, 594, 604
Bojeri, ii, 604
Helsingbergii, ii, 604
paniculata, ii, 604
Roxii, ii, 604
Thompsonii, ii, 604
Stroemia farinosa, i, 94, 276
longifolia, i, 94
rotundifolia, i, 94
Strophanthus, ii, 233, 243
Struchium americanum, i, 165
Sturtia, i, 317
gossypioidea, i, 317
Stylidæ, i, 34, 63, 339; ii, 269 *note*
312
Stylidium, i, 34; ii, 322, 532
adnatum, ii, 678
breviscapum, ii, 678
falcatum, ii, 678
fasciculatum, ii, 678
propinquum, ii, 678
Stypandra, i, 50
Stypheliæ, i, 38
Suag, i, 279
Subularia, i, 192, 254; ii, 384

- 630
 0
 628, 640, 643
 2 note, 636
 627
 623, 627
 388; ii, 611
 624
 70, 252
 4, 604
 6
 165
 , 269 note
 32
 84
- Subularia aquatica*, ii, 384
Succowia, ii, 375
 balearica, ii, 375
Sugar-cane, i, 154, 156
Sutherlandia, ii, 442
 frutescens, ii, 442
Symphionema, ii, 17, 37, 131
 montanum, ii, 131
 paludosum, ii, 13, 131
Symphoria, ii, 672
 racemosa, ii, 672
Symphoricarpus, ii, 324, 542
Symplocere, i, 388 note
Symplocos, ii, 245
Synaphea, i, 41, 82, 381; ii, 11, 15,
 17, 18, 21, 37, 129
 dilatata, i, 82; ii, 130
 favosa, ii, 130
 petiolaris, ii, 130
 polymorpha, ii, 130
 ruralis, i, 239
Syntrichia mucronifolia, i, 239
Systylium, i, 241, 247; ii, 337
Swainsona, ii, 441
 coronillifolia, ii, 442
 grandiflora, i, 323; ii, 441
 Greyana, i, 323, 324
 laxa, i, 324
Swartzia capillacea, i, 238

Tabernæmontana, ii, 242
Tamarinc, i, 154, 156
Tamarindus indica, i, 154
Tamariscinæ, i, 288
Tamarix gallica, i, 285
Tapura, i, 126
 guianensis, i, 126
Targionia hypophylla, i, 69
Tasmania, i, 64, 559; ii, 704
 aromatica, ii, 704
 dipetala, ii, 704
 insipida, ii, 704
Taxanthemum attenuatum, i, 94
Taxus, i, 43, 137
Tayloria, ii, 337
Tchilli, i, 158
Teesdalia, ii, 378
 nudicaulis, ii, 378
Teichostemma fruticosum, i, 95
Telanthra, i, 164
Teleozoma, ii, 526, 570
Tellima, ii, 522
Telopia, ii, 38, 166
 speciosissima, ii, 167
 truncata, ii, 167

Templetonia, ii, 439
 retusa, ii, 439
Terebintaceæ, i, 291
Terminalia, i, 20?; ii, 534
 cycloptera, i, 94
 volucris, ii, 710
Testudinaria, i, 408
Tetracera, i, 12
Tetradia, ii, 637
 Horsfieldii, ii, 637
Tetrameles, i, 285
Tetraphis, ii, 337
 pellucida, ii, 348
Tetratheca, i, 15, 381
Tetrathecacceæ, i, 15
Thamnea, ii, 322
Thelira, i, 115
Thelotrema lepadinum, i, 69
Thelymitra, ii, 483
 ixioides, ii, 483
Theophrasta, i, 151 note
Thermopsis, ii, 421
 lanceolata, ii, 421
Thesium, i, 137
Thlaspi, i, 263, 271; ii, 376
 alliaceum, ii, 377
 alpestre, ii, 377
 arvense, ii, 376
 Bursa-pastoris, ii, 377
 campestre, ii, 382
 ceratocarpon, ii, 377
 montanum, ii, 377
 perfoliatum, ii, 377
 saxatile, ii, 376
Thlaspidium, ii, 379
Thoa, i, 453
Thompsonia, i, 387 note
Thonningia sanguinea, i, 414 note
Thottea, i, 384
 grandiflora, i, 384 note
Thunbergia angulata, i, 94
Thymelææ, i, 82, 127; ii, 16, 31
Thymus virginicus, ii, 504
Thyrsine, i, 429
Ticorea, i, 16
Tiliacæ, i, 11, 108, 109; ii, 613, 646
Tillandsia usneoides, i, 413 note
Timmia cucullata, i, 235
 mcgapolitana, i, 235
Tmesipteris, ii, 722
Tobacco, i, 155, 156
Toluisera, i, 127 note
Tomex glabra, i, 93
Tonina, i, 53
Toresia, i, 231

- Tortula mucronifolia*, i, 239
 ruralis, i, 239
Tradescantia, i, 532
 virginica, i, 513, 530 *note*
Tremandra, i, 15; ii, 705
 diffusa, ii, 705
 stelligera, ii, 705
Tremandrea, i, 15, 63
Trematodon, ii, 348
Tremella cruenta, i, 178
Tribulopsis, i, 319
 angustifolia, i, 319
 pentandra, i, 319
 Solandri, i, 319
Tribulus Histrix, i, 318
 occidentalis, i, 318
 terrestris, i, 285
Trichodium, i, 55
 algidum, i, 220
 laxiflorum, ii, 514
Trichomanes, i, 146; ii, 510
 brevisetum, ii, 510
Trichosiphon, ii, 637, 638
Trichosporæ, ii, 585
Trichosporum, ii, 585
Trichostomum, ii, 335, 348
 canescens, i, 69
 capillaceum, i, 238
 lanuginosum, i, 178, 237
Tricocæ, ii, 615
Tridax, ii, 283
 procumbens, ii, 283
Triostcum, ii, 324, 542
Triphaca africana, ii, 628
Triphelia brunoides, ii, 708
Tripsacum, ii, 574
Triraphis molle, i, 302
 pumilio, i, 302
 pungens, i, 302
Trisetum, i, 230
 airoides, i, 231; ii, 514
 subspicatum, i, 231; ii, 514
Tristania, i, 19; ii, 456
 conferta, ii, 456
 laurina, ii, 456
 nereifolia, ii, 456
Tristemma hirtum, i, 118
 incompletum, i, 116
Triticum, ii, 277
Triumfetta denticulata, ii, 709
 glaucescens, ii, 709
Tropæolum majus, i, 379 *note*, 557
Tromsdorfia, ii, 593, 594, 597
 elongata, ii, 597
 speciosa, ii, 599
Tubercularia vulgaris, i, 70
Turritis, ii, 397
 alpina, ii, 396
 arvensis, ii, 398
 cærulea, ii, 395
 ciliata, ii, 396
 glabra, ii, 398
 hirsuta, ii, 397
 Loeselii, ii, 398
Tussilago corymbosa, i, 211, 253
 frigida, i, 212, 253
 odorata, ii, 270
Tylophora, ii, 210
Typha angustifolia, i, 66, 165

Ulva crispa, i, 178
 Lactuca, i, 70
 plumosa, i, 70
Umbelliferæ, i, 29
Umfar, i, 93
Uncaria elliptica, ii, 533
 ovata, ii, 533
Uniola latifolia, i, 226
Unona Æthiopica, i, 105, 161
 aromatica, i, 105
 hamata, ii, 685
 uncinata, ii, 685
Upas Antiar, i, 79
Urania, i, 157 *note*
Urceola, i, 133
Urena, i, 559
 Americana, i, 109, 165, 171
 glabra, i, 95
 mollis, i, 95
 reticulata, i, 165, 171
Urticea, ii, 534, 537
Urticæ, i, 79, 137
Usnea, i, 178
 florida, i, 70
 melaxantha, i, 252
 ochroleuca, i, 251
 sp. racelata, i, 252
Usteria, i, 37, 132
Utricularia, i, 65
Uvaria, ii, 534, 535
 esculenta, ii, 685
 heteroclita, ii, 685
 odoratissima, ii, 685
 uncata, ii, 685

Vahea, i, 133
Vahlia, ii, 522
Valerianæ, i, 65
Vallaris, ii, 232, 237
Valisneria spiralis, i, 68

Vanda, ii, 688
 Roxburghii, ii, 689
 Vandee, ii, 576
 Vandellia, i, 164
 Vanilla, ii, 497, 551
 aromatica, ii, 498
 planifolia, ii, 498
 Vella, ii, 375
 annua, ii, 375
 Pseudo-Cytisus, ii, 376
 Velleæ, i, 253
 Velleia, i, 33; ii, 266 *note*
 Velvet Tamarind, i, 161
 Ventenatia, i, 108
 Ventilago, i, 26
 Verbena officinalis, i, 68
 triphylla, i, 136
 Verbenaceæ, i, 38, 40, 135; ii, 659
 Verbesina gigantea, ii, 291
 Vernoniaceæ, i, 128
 Verrucaria nitida, i, 69
 Vesicaria, ii, 389
 sinuata, ii, 389
 ntriculata, ii, 389
 Vestia, ii, 683
 Viburnum, ii, 537
 Vicia galegifolia, ii, 442
 Vigna, i, 325
 Villa, i, 219, 221
 Viminaria, ii, 429
 denudata, ii, 429
 Vinca, ii, 247
 Vincetoxicum nigrum, i, 529
 Viniferæ, i, 151
 Viola, i, 122, 532
 tricolor, i, 469 *note*
 Violeæ, i, 122
 Virgilia, ii, 421
 aurea, ii, 421
 capensis, ii, 422
 intrusa, ii, 422
 Viscum, i, 21, 120, 137, 381
 album, i, 460
 Visnia, i, 164
 Vitellaria paradoxa, i, 296
 Vitex, i, 136
 Vitis, i, 151, 288, 390
 Vitmannia, i, 106
 Voacanga, i, 133
 Voanda, i, 159
 Voandzeia, i, 155
 Voandzou, i, 155
 Voitia, i, 247
 hyperborea, i, 248, 256
 nivalis, i, 249

Voitia vogesiana, i, 248
 Volkameria, ii, 506
 aculeata, ii, 506
 angustifolia, ii, 508
 heterophylla, ii, 508
 japonica, ii, 507
 Kæmpferi, ii, 507
 Kæmpferiana, ii, 507
 ligustrina, ii, 508
 tomentosa, ii, 508
 inermis, ii, 508
 Waltheria americana, i, 165, 171
 indica, i, 165, 161
 Wanzey, i, 94
 Wedelia, ii, 285, 297 *note*
 Weigelia, ii, 541
 Weinmannia, i, 20
 Weissia, ii, 348, 349
 affinis, ii, 344
 controversa, i, 69
 nuda, ii, 348
 radians, ii, 344
 splachnoides, i, 241; ii, 347
 Templetoni, ii, 343, 344
 trichodes, ii, 344
 Westringia, i, 39, 380
 Whittia, ii, 596, 610
 Wintera, i, 64; ii, 704
 Winteranæ, i, 413 *note*
 Wintereæ, ii, 704
 Winter's bark, ii, 724
 Woodsia, ii, 253, 543, 546
 glabella, ii, 518
 hyperborea, ii, 253
 ilvensis, ii, 253, 518
 mexicana, ii, 546
 Wormia, i, 12
 alata, ii, 704
 Wrightia, ii, 233, 242, 243
 antidysenterica, ii, 244
 pubescens, ii, 244
 tinctoria, ii, 244
 Zeylanica, ii, 244
 Wulfenia intermedia, ii, 577
 Notoniana, ii, 607
 obliqua, ii, 577
 Xanthium, i, 31
 Xanthorrhæa, i, 50, 308, 435
 arborea, ii, 724
 Xeropetalum, ii, 673
 Xerotes, i, 52, 436
 Ximenia, i, 155
 americana, i, 551

Xylomelum, ii, 15, 18, 21, 38; ii, 159

pyriforme, ii, 159

Xylophylla, i, 43

longifolia, i, 43

Xylosteum, ii, 541

Xyris, i, 53

Xysmalobium, ii, 217

grandiflorum, ii, 217

undulatum, ii, 517

Yam, i, 154, 160

Zaffo, i, 160 *note*

Zamia, i, 458, 506 *note*; ii, 717

spiralis, i, 308, 310

Zanthoxylon, i, 16

Zapania, i, 136

nodiflora, i, 66, 68

Zea, i, 506 *note*

Mays, i, 154, 469 *note*

Zieria, 17

canescens, ii, 709

furfuracea, ii, 709

involucrata, ii, 709

lanceolata, ii, 669

Smithii, ii, 669

Zizania aquatica, i, 469 *note*

Ziziphus, i, 26

Zostera marina, i, 68

Zygodon, ii, 344

Zygophyllæ, i, 144, 285

Zygophyllum album, i, 285

coccineum, i, 287

Fabago, i, 287

lanatum, i, 286

simplex, i, 285