IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 871-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1985

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- ☒ Coloured coverz/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- ☐ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☐ Bound with other material/
Relié avec d'autres documents
- ☐ Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distorsion le long de la marge intérieure
- ☐ Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- ☐ Additional comments:/
Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☒ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
				✓							

The copy filmed here has been reproduced thanks to the generosity of:

Seminary of Quebec
Library

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Séminaire de Québec
Bibliothèque

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

Horse Doctor.

TORONTO:
PUBLISHED BY ADAM MILLER
1873

THE
COMPLETE FARRIER
OR,
HORSE DOCTOR.

A GUIDE FOR THE

Treatment of Horses in all Diseases

TO WHICH

THAT NOBLE ANIMAL IS LIABLE.

BEING THE RESULT OF

Fifty Years Extensive Practice of the Author.

BY JOHN C. KNOWLSON.

DURING HIS LIFE AN ENGLISH FARRIER OF HIGH POPULARITY.

Twenty-Ninth American Edition.

Containing the Latest Discoveries in the Cure of Strains.

=====

TORONTO:

PUBLISHED BY ADAM MILLER,

AND FOR SALE BY ALL BOOKSELLERS.

1868.

INDEX TO CONTENTS.

Back Sinews, sprain in.....	46	Kidneys and Bladder, disorders of.....	29
Bladder, disorders of.....	29	Knees, sprain in.....	47
Blood Spavin.....	51	Lameness in the Stifle.....	47
Bone Spavin.....	50	Lameness in the Cup-Bone.....	48
Bots and Worms.....	11	Mallenders and Sallenders.....	52
Broken Wind.....	22	Mange, the.....	35
Bruises of the Withers.....	53	Moulten Grease.....	32
Coffin Joint, sprain in.....	46	Pasterns, sprain in.....	47
Cold, remedy for.....	3	Poll Evil.....	43
Colic, or Gripes.....	7	Plushes from Saddles.....	59
Colic, Flatulent or Windy.....	7	Remedy for a Cold.....	3
Colic, Billous or Inflammatory.....	9	Ring-Bone.....	52
Convulsions.....	16	Sallenders.....	52
Corb.....	52	Scouring.....	27
Cough and Asthma.....	5	Scratches.....	61
Crown Scab, the.....	62	Spavin, Bone.....	50
Cup-Bone, lameness in the.....	48	Spavin, Blood or Bog.....	51
Farcy, or Farcin.....	37	Sprain in the Back.....	44
Farcy, the Water.....	41	Sprain in the Shoulder.....	45
Feet, hurts in the.....	56	Sprain in the Coffin Joint.....	46
Fevers.....	17	Sprain in the Back Sinews.....	46
Fever, a Compound.....	19	Sprains in the Knees and Pasterns.....	47
Fistula, or Bruises of the Withers.....	58	Stag Evil.....	16
Founders.....	42	Stifle, lameness in the.....	47
Girth-Galls, Plushes, &c.....	59	Strangles, the.....	52
Glanders, the.....	54	Surfeit.....	33
Grease, the.....	60	Swellings and Impostumes.....	54
Gripes, the Dry.....	10	Twitter-Bone.....	57
Gripes, or Colic.....	7	Warbles, Girth-Galls, &c.....	59
Hidebound, the.....	35	Water Farcy, the.....	41
How to Manage a Horse on a Journey.....	62	Whirlbone, lameness in the.....	48
Hurts in the Feet.....	56	Windgalls.....	60
Impostumes.....	54	Wind, a Broken.....	22
Intestines, disorders of.....	27	Withers, bruises of.....	58
Jaundice, or Yellows.....	13	Worms and Bots.....	11
Journey, how to manage on a.....	62	Wounds.....	55
		Yellows, the.....	13

THE COMPLETE FARRIER,

OR,

HORSE DOCTOR.

Remedy for a Cold.

People not much accustomed to study the diseases of horses, look upon a cold as a trivial matter. But they should bear in mind that nearly the whole of the disorders to which this noble animal is liable, derive their origin from a Cold. The causes of Colds are various, but the most usual one is riding the horse until he is hot, and then suffering him to stand still exposed to the cold air; removing him from a hot stable to a cold one: if the horse has been high-fed, and clothed, the cold contracted in this manner often proves very violent; and this is the reason why horses often catch a severe cold on their first coming out of the dealer's hands: neglecting to rub him properly down, and to rub the sweat carefully off when he comes in from a journey; and I have known grievous disorders brought on by removing horses into a new stable before the walls and plastering were dry. Workmen are often in fault for not leaving air-holes above; as when a horse comes into a new stable, and gathers heat, it will cause the walls and plastering to sweat very much, especially if there are no air-holes left. Many a horse has lost his eyes, and some their lives, by being put into new stables before they were dry.

SYMPTOMS.—When a horse has caught a cold, a cough will follow, and he will be heavy and dull in proportion to the severity of the disease: his eyes will be watery; the kernels about his ears, and under his jaws, will swell, and a thin, mucous gleet will run from his nose. If the cold be violent, the horse will be feverish; his flanks will heave, and he will refuse his food. The owners should be very careful to observe these last symptoms, because when they appear, and are attended with a slimy mouth, cold ears and feet, moist eyes, and a great inward soreness, there is danger of a fever, and generally of a malignant kind. But when the horse coughs strongly, and snorts after it, eats scalded bran, and drinks warm water, moves briskly in his stall, dungs and

stales freely, and without pain, his skin feels kindly, and his coat does not stare, there is no danger, nor any occasion for medicine. You should, however, bleed him, keep him warm, give him some feeds of scalded bran, and let him drink warm water.

THE CURE.—If the horse feels hot and refuses his meat, it will be necessary to bleed him plentifully, and to give the following drink :

2 ounces of Juice of Liquorice.
2 ounces of Salt of Tartar.
2 drams of Saffron.
2 ounces of Honey.

Cut the juice small, dissolve all together in hot water, and give it nearly cold. This drink may be repeated as occasion requires, but let twenty-four hours elapse first. Or give

4 ounces of Aniseeds.
2 ounces of Liquorice Root.
1 ounce of Gum Scammony.
1 ounce of Nitre.

Boil these together in three pints of water for ten or twelve minutes ; strain the liquor through a cloth ; and add two ounces of honey to it when you give it to the horse.

The following ball, commonly called *The Cordial Ball*, is one of the best yet found out for coughs or colds. Be careful to get your drugs good, for this ball is of great worth in many diseases ; and few things will remove a cough or cold, or clear a horse's wind, sooner or better :

Take 4 ounces of Aniseeds Powder.
4 ounces of Fennigreek.
4 ounces of Liquorice Powder.
4 ounces of Elecampane Powder.
4 ounces of Flour of Brimstone.
6 ounces of Grains of Paradise, in fine powder.
4 ounces of Liquorice, cut small, and dissolved in White Wine.
1 ounce of Saffron, pounded small.
1 ounce of Oil of Aniseeds.
8 ounces of Olive Oil.
8 ounces of Honey

Bray them all well together till they come into paste, and if they should be too dry, add a little more olive oil and honey. The dose is about two ounces, and may be given three or four times a day, if needful. These balls, consisting of warm, opening ingredients, are of great use ; and given in small quantities, about the size of a pullet's egg, will encourage a free perspiration ; but in case of a Fever, they should be given with the greatest caution.

It will be of great use to put scalding-hot bran into the manger, that the horse may hold his head over it, and receive

the steam up his nostrils, which will cause a running from them, and relieve him very much. I have known asarabacca, dried and rubbed to powder, and blown up the nostrils, to cause a discharge; for when a horse has caught a violent cold, he is often troubled with a pain in his head, which a good discharge at the nose is very likely to cure. For the same purpose the horse should be warmly clothed, especially about the head, neck, and throat, as it has a tendency to promote a running at the nostrils.

By this simple method, with proper care, hot manes, and warm water, most colds may be cured; and as soon as the horse begins to feed heartily, and snorts after coughing, an hour's exercise every day will greatly hasten the cure. If the legs swell, and the horse be full of flesh, rowels are necessary.

A Cough, and Asthma.

Among all diseases to which this noble creature is subject, none has given more perplexity to Farriers than a settled Cough; indeed it too often defies all the attempts of art, and the horse frequently becomes *Asthmatical*, or *Broken-winded*.

CAUSES.—The causes are various. Sometimes it is owing to colds imperfectly cured; sometimes to pleurisies, or malignant fevers, which have left a taint upon the lungs or other vessels; sometimes to small eruptions in the glands, which cause the lungs to be much larger than they ought to be, and a quantity of phlegm, and mucilaginous juices, which stuff up the glands and branches of the windpipe; and sometimes to fleshy substances engendered in the large blood vessels; for all these things hinder a free respiration, and excite a cough. It is of the utmost importance to distinguish one kind of cough from another, and this makes the disorder so hard to cure: for it can not be cured till the seat of the complaint be found out.

If the cough be of long standing, attended with a loss of appetite, wasting of flesh, and weakness, it denotes a Consumption; and that the lungs are full of knotty, hard substances, called tubercles. When a cough proceeds from phlegm, and mucilaginous matter, stuffing up the vessels of the lungs, the flanks have a sudden, quick motion; the horse breathes quick, but not with his nostrils distended like one that is broken-winded; his cough is sometimes moist, and sometimes dry and husky; before he coughs he wheezes, and sometimes throws out of his nose or mouth large pieces

of white phlegm, especially after drinking, and this discharge generally gives very great relief.

CURE.—If the horse be full of flesh, take from him a moderate quantity of blood. The next day give him scalded bran, and in the evening the following ball :

- 1 ounce of Powder of Aniseeds.
- 1 ounce of Liquorice Powder.
- 1 dram of Calomel, 8 drams to an ounce.

Work them into a ball with Barbadoes tar. Give this ball the last thing at night, and be careful to keep the horse out of wet, and from cold water the next day. On the second morning give the following purge :

- 1 ounce of Barbadoes Aloes.
- 1 ounce of Castile Soap.
- $\frac{1}{2}$ ounce of Powdered Ginger.
- 1 dram of Oil of Aniseeds.

Bray them together in a mortar, with a little syrup of buckthorn to make them into a ball, which is to be given in the morning, and plenty of warm water, and walking exercise, till it be wrought off. It will not work the first day. In three days after, give six ounces of the Cordial ball in a little warm ale, fasting, and to fast two hours after. Repeat the Calomel ball, physic, and Cordial ball, six days after, in the same manner as before. Let the horse's hay be sweet, and his manger-meat scalded bran, with a spoonful of honey in each feed : let him have walking exercise in the open air, but be careful of wet, and of cold water.

When this course has been pursued two or three times, give two or three ounces of the Cordial ball every morning. The above method will remove most Coughs, but if it fail, try the following :

- 1 ounce of Gum Ammoniacum, in fine powder.
- $\frac{1}{2}$ ounce of Gum Galbanum, in powder.
- 2 drams of Saffron, brayed.
- 2 drams of Assafetida, in powder.

Work them up with honey, or Barbadoes tar, into one ball ; roll it in liquorice powder, and give it fasting, and to fast two hours after. This ball must be given every morning, for six or seven times, before it can have a fair trial. In the cure of this disease, the diet should be very moderate, the usual quantity of hay should be abridged, and sprinkled with water, and the usual allowance of grain and water divided into several portions ; for with these regulations in diet the disease will soon be cured ; and where it is incurable, the horse will be so far recovered as to be able to do a great deal of work.

this dis-

a mode-
scalded

this ball
orse out
e second

yrup of
given in
ng exe-
rst day.
all in a
Repeat
after, in
e sweet,
f honey
pen air,
e times,
orning.
it fail,

to one
and to
morn-
al. In
erate,
inkled
ter di-
ons in
ncura-
do a

It may not be improper here to add that some young horses are subject to coughs when cutting their teeth, and their eyes are also affected from the same cause. In these cases always bleed, and if the cough be obstinate, repeat it, and give warm mashes, which are often sufficient alone to remove the complaint. When young horses have a cough that is caused by worms, as is often the case, such medicines must be given as are proper to destroy those vermin. [See Worms.]

The Colic, or Gripes.

The Colic proceeds from various causes; therefore the method of cure varies; for otherwise the medicines intended to cure it may increase it, and perhaps render it fatal. We shall, therefore, divide this disorder into three different species, and endeavor to give such plain directions for managing each, as can not fail to prove very beneficial. The three species are these:—1. The Flatulent, or Windy Colic;
2. Bilious, or Inflammatory Colic;
3. The Dry Gripes.

The Flatulent, or Windy Colic.

SYMPTOMS.—The horse is very restless, lying down and starting up again. He strikes his belly with his hind feet, stamps with his fore feet, and refuses his meat. When the pain is violent, he has convulsive twitches; his eyes are turned up, and his limbs stretched out, as if dying; and his ears and feet alternately cold; he falls into profuse sweats, and then into cold damps; often tries to stale, and turns his head frequently to his flanks; he then falls down, rolls about, and often turns on his back. This last symptom proceeds from a stoppage of urine, which generally attends this species of colic, and may be increased by a load of dung pressing on the neck of the bladder.

CAUSES.—This disease often proceeds from catching cold by drinking cold water when hot, and the perspirable matter is by that means thrown upon the bowels, which causes them to distend violently, and sometimes brings on an inflammation in the small intestines, when the body begins to swell, and the cure is despaired of.

CURE.—The first thing to be done is to empty the straight gut with a small hand dipped in oil. This frequently gives room for the wind, before confined in the bowels, to discharge itself; and, by taking off the weight that pressed upon the neck of the bladder, the suppression of urine is removed.

upon which the horse immediately stales, and becomes much easier. If the horse be young, and full of blood, it will be proper to take a sufficient quantity of blood from the neck.

When these purgative operations have been performed, the following may be given, as it seldom fails to give relief :

- 4 ounces of Tincture of Senna, or Daffy's Elixir.
- 6 drams of Tincture of Opium.
- 1 dram of Oil of Juniper.
- 8 ounces of Juniper Berries, braised.

Put one quart of boiling water on the juniper berries, let them stand a few minutes, strain it off, put all together, and give them to the horse.

If he does not find relief soon after taking this dose, both by staling and breaking wind, it is doubtful whether he will receive any benefit from it ; so you must prepare the following clyster for him as soon as you can :

- Take 4 ounces of Camomile Flowers.
- 2 ounces each of Aniseeds, Fennel and Coriander.

Boil them in one quart of water, and add 2 ounces of Castile Soap, cut small, while the water is hot, that the soap may dissolve. Give it blood warm. During the fit the horse may be walked about, or trotted a little, but should by no means be harrassed, or driven about till he is jaded. If no better, give the following :

- 2 drams of Camphor.
- 1 dram of Pellitory of Spain.
- 2 ounces of Ginger Powder.
- 3 gills of Holland Gin.

If the horse sweat much at times, and then falls into cold sweats, give four ounces of mithridate, in three gills of Holland gin, and repeat the clyster. If the disorder continue three or four hours, give one ounce of tincture of opium, in three gills of Holland gin. When the horse begins to recover, he will lie quiet, without starting and trembling ; and if he continues in this quiet state an hour, you may conclude that the danger is over. Dress him down well, and give him a small quantity of warm water, if he will drink it : bed him down well, cover him to keep him warm, and then leave him to get a little rest. You must consider that the disorder has left a little soreness on him, both within and without ; therefore, make him a little gruel, with a pint of red wine in it ; and if any skin be knocked off about his eyes, or his huck-bones, rub it with the wash recommended for bruises.

Sometimes the Colic is received into the stomach, and does not act so violently, nor cause the horse's pains to be so strong. You may best judge of this by his motions. He

will draw his four feet together, lay himself down, stretch out his feet and head, throw his head back, and often put his nose to his chest; after standing a little, he will lie down again as before. When the colic is easier, he will lie for an hour or more together, with his feet stretched out and his head thrown back, or with his nose upon his ribs. This is caused by bad feed, or bad water, or both: sometimes by drinking hard water when hot, or by a change from soft grit water to limestone or iron water, or by the break of a storm. I have had five or six horses under my care in this disorder in one day, at the break of a frost, by drinking ice or snow water. Sour grains, sour grass, dry meal, dust, bad hay, and many other things cause this disorder. Give the following, which is almost a certain cure in two hours:

1 ounce of Spirits of Sweet Nitre.
 1 ounce of Spirits of Nitre.
 1 ounce of Tincture of Opium.
 1 ounce of Sweet Oil.

All to be given together in a gill of warm ale. Bed the horse well down, and leave him that he may get a little sleep, after which he will get up and feed. This is one of the best medicines that has yet been found out. It has saved hundreds of horses, and will save hundreds more if rightly applied.

The Bilious or Inflammatory Colic.

SYMPTOMS.—This kind of Colic, besides most of the symptoms of the former, is attended with a fever, great heat, panting, and dryness of the mouth. The horse also generally parts with a little loose dung, and a little scalding-hot water; which, when it appears blackish, or reddish, indicates an approaching mortification.

Take 3 ounces of Senna.

1 ounce of Salt of Tartar. Infuse them in a quart of boiling water an hour—strain, and add

2 ounces of Lenitive Electuary.

4 ounces of Glauber's Salt. Mix when hot.

If the disorder be not removed by the above medicine, but, on the contrary, the fever and inflammation continue to increase, attended with a discharge of flesh-colored matter, the event will probably be fatal; and the only medicine likely to prevent it, is a strong decoction of Jesuit's bark, a pint of which may be given every three hours, mixed with a gill of red port wine; or you may give one ounce of the powder of bark with the wine. Or, if these can not be got easily, give four ounces of tincture of rhubarb in three gills

of red port wine. Also give a clyster every two hours, made of two new-laid eggs, well broken, and two ounces of molasses, in one quart of milk. Give it warm.

If the horse recover, it will be proper to give him a gentle purge or two in a week after.

Take 1 ounce of Rhubarb, in Powder.
 $\frac{1}{2}$ ounce of Jalap, do.

Work them up into a ball with syrup of buckthorn, and give it to the horse, with warm water to work it off.

The Dry Gripes.

SYMPTOMS.—This disorder mostly proceeds from costiveness, and is discovered by the horse's frequent and fruitless attempts to dung, the blackness and hardness of the dung, the frequent motion of his tail, the high color of his urine, and his great uneasiness.

CURE.—The first thing to be done is to draw the dung out of the fundament, with a small hand, as far as you can reach, and then give the following :

4 ounces of Castor Oil.
 4 ounces of Tincture of Senna.
 $\frac{1}{2}$ ounce of Oil of Juniper.

Give them all together, and then the following clyster :

Boll in a quart of water a handful each of
 Camomile Flowers and
 Marshmallows. Strain off and add
 2 ounces of Linseed Oil, or Pale Oil.

If the horse do not mend, repeat both the drink and the clyster. During this disorder the horse must not have any dry food ; but boiled linseed and scalded bran, with warm water to drink. Gentle walking exercise is a great means to cause the physic to work ; but be careful of cold.

From the account that I have given of the different species of the Colic, the reader will be abundantly convinced how necessary it is to be acquainted with each, that he may be able to give proper medicines and to relieve the creature's excruciating pains. He should carefully avoid all hot, violent medicines, which always prove hurtful in every species of this disorder, and frequently fatal. Nor is it any wonder that horses treated in that manner should die, for such medicines stimulate the neck of the bladder, augment the heat of the blood, before much too great, and inflame the bowels, by which a mortification is brought on, and the horse is lost by the very means used for his recovery.

Sharp fits of the Gravel are sometimes taken for the Colic ; but should this happen, the drink recommended for the Colic will also be proper for the Gravel.

Worms and Bots.

Horses are subject to five sorts of worms, and perhaps to many more, but I shall only describe three, which are the most common. The worst sort to destroy are long, round worms, resembling earth-worms, but smaller at the tail ; they have a seam all the length of their bodies, and are very hard : these are called *Round Worms*. The next are small worms, about the size of a sewing needle ; they have reddish flat heads, having nine feet on each side, and are called *Ascarides* : these are also very troublesome to horses. The third sort are short, thick worms, called *Bots* : their seat is mostly at the stomach, but when horses get any food that they are fond of, they fill themselves so full, that they lose their hold, and come along with the dung to the fundament, and there catch hold and stick to the end gut, partly out of the horse ; this happens mostly in Spring, when they get the juice of fresh grass.

It is well known that horses which have many worms can never thrive, or carry much flesh. If the breeding of these vermin were prevented, it would add much to the strength of the horse ; and it might be done by giving him a decoction of bitter herbs, such as wormwood, in Spring. It may be boiled, or steeped in hot water, and given two or three times a week. Or a decoction of wormwood, buck-bean, gentian root, and camomile flowers, of each a large handful, boiled in a sufficient quantity of water, and given as above, will answer the end.

SYMPTOMS.—The symptoms which indicate worms are various, as the animals are different, and seated in different parts of the body. When the Bots are seated in the straight gut, they are never dangerous, but are often thrust out with the dung. They generally come in the months of May and June, and scarcely ever continue in a horse above a fortnight. But when they breed in the stomach, they often cause convulsions, and even death. The Bots that breed in the stomach are about the size of a large maggot, composed of circular rings, and have little, sharp, prickly feet along the sides of their bellies, by means of which they fasten themselves to the part from whence they derive their nourishment, to prevent their being loosed from such adhesion before they come to maturity ; and as they drain the coats of the stomach like

leeches, it is no wonder that they often throw the horse into convulsions, which terminate in death, unless the cause be removed. The violent agonies of the creature are the only indications of their existence. The other kinds of worms are more troublesome than dangerous, and are discovered by the following signs : There is a white fur on the end of the straight gut ; the horse is lean and jaded ; his coat is rough and staring ; and if you rub your hand backward on the hair, a white scurf will rise, as if he had been surfeited ; and though he eats with a remarkable appetite, he does not thrive. He often strikes his hind feet against his belly, and is sometimes griped, but without the violent pains that attend the colic, or stranguary ; for he never rolls or tumbles, but is uneasy, often laying himself down quietly on his belly for a little while, and then rising and beginning to feed. But the surest symptom is when the horse voids the worms with his dung.

The way these creatures get into the stomach is curious. While the horse is feeding, the bot-fly lights on his side in a place convenient to the animal's mouth. He then stings the horse and flies away. The horse naturally turns his head and throws his mouth to the place which has been stung—the eggs adhere to his lips, and in this way are taken into the stomach, where they hatch into worms, which instantly fasten themselves to the coats of the stomach. Keeping a horse well curried and brushed, and keeping him in stable during the fly-bot season, will often prevent the disease.

CURE.—Many medicines have been given to destroy these vermin, without knowledge or judgment, and even contrary to common reason. Some give coarse sugar for that purpose, but, in my opinion it will rather increase than destroy them ; although a few will fill themselves so full as to lose their hold, and come away with the dung.

Take 1 ounce of Socotrine Aloes.
 1 dram of Calomel, 8 drams to an ounce.
 1 dram of Oil of Aniseeds.
 2 drams of Powdered Ginger.
 $\frac{1}{2}$ ounce of Syrup of Buckthorn.

Beat all up together in a mortar till the Aloes are well broken, and the whole is brought into a paste ; which give in the morning, fasting, and to fast one hour after ; also give warm water, and walking exercise, till wrought off. It will not work the first day. Be careful that the horse be open in his body before you give the ball. In grass time you will have nothing more to do than to give it, and to put the horse

where he can get water. This dose is for a pretty strong horse, so you must add or diminish according to size. This dose must be repeated as need requires, but not within seven days. It will destroy most kind of worms; but the hard round worms require different treatment, as they are the worst of any to get rid of. To destroy them give the following :

1 dram of Calomel, 8 drams to an ounce.
6 drams of Jalap.
6 drams of Rhubarb, in powder.

Wrought up into a paste with conserve, or honey; and two days after give the following :

1 dram of Calomel, 8 drams to an ounce.
1 ounce of dried Foxglove Leaves, powdered.
 $\frac{1}{2}$ ounce of Worm Seed, powdered.
1 ounce of Jalap, in powder.

To be given in three gills of malt liquor from the mash-tub. If the above be given every week for three weeks together, you may be sure that most of the vermin will be expelled. If the medicines be given in the house, let the food be light and opening, and warm water for two days, with walking exercise.

I advise all who have horses troubled with worms, to give safin, dried and powdered, before they give the worm physic. If one ounce a day be given for a week before, in a mash of bran, it will be much better. The above ball is good for many disorders besides worms.

The Yellows, or Jaundice.

This disorder is very common in horses, and sometimes it is either unknown, or overlooked, till it gets much worse to cure. A young horse is easily cured, but an old one is bad to cure. Some people may say, *How can a horse have the Jaundice, when he has no gall-bladder?* I answer. *Though he has no gall-bladder in sight, he has a large vessel in the liver which answers the same end.* We may here observe, the handiwork of God in placing a horse's gall-bladder differently from those of other animals, when we consider that the horse is the swiftest and most laborious creature in the world; so that if the gall-bladder had been placed on the liver, it would always have been exposed to injuries.

SYMPTOMS.—The white of the eyes is yellow, also the inside of the mouth, the tongue, and the bars in the mouth, are of a dusky yellow; the horse is dull, and refuses all kinds of food; a slow fever is perceived, which increases with the

yellowness; the dung is often hard and dry, and of a pale yellow color; the urine is commonly of a dark, dirty brown, and when it has settled, sometimes looks like blood: the horse stales with pain and difficulty; and, if the disorder is not checked, becomes in a short time unable to stir about. When this disease gets strong hold of a horse before proper medicines are applied, it is often fatal, or it brings on some other disorder as bad, or worse than itself.

Cure.—Bleed plentifully, and as this disease is always attended by a costive habit of body, it will be proper to give a clyster or two before you give the physic. A clyster may be made of one ounce of camomile flowers, boiled in a quart of water with two ounces of Castile soap. Then give the following ball:

4 drams of Indian Rhubarb.
2 drams of Saffron.
6 drams of Scocotrine Aloes.
1 ounce of Castile Soap.

To be brayed in a mortar with a little syrup of buckthorn, and made into two balls; one to be given the last thing at night, and the other the first thing in the morning; and give plenty of warm water to work them off. If the disease is obstinate and will not yield to the above, give the following ball:

1 dram of Calomel, 8 drams to an ounce.
1 ounce of Barbadoes Aloes.
1 ounce of Venice Soap.
1 ounce of Turmeric.
1 dram of Oil of Juniper.

All to be brayed in a mortar with a little syrup of buckthorn, and made into two balls; one to be given at night, and the other in the morning, with plenty of warm water, and walking exercise till wrought off.

If the horse is better after the first medicine, repeat it, for it is much safer than the last; but if not, the last may be given, and repeated as need requires; but be careful not to give it in less than six days distance.

If, after giving one dose of the last ball, you give the following powders every day in celantine tea, if it can be got, or in warm ale, you will find them very useful in removing the complaint.

2 ounces of Salts of Tartar.
1 ounce of Æthiop's Mineral.
2 ounces of Turmeric.

These powders are proper to be given after either of the former doses.

The Staggers.

This disease is caused by the liver making blood so fast that the cavity of the heart is overloaded and the blood flies up the neck vein till the head is overloaded too; and if relief can not be obtained, the horse soon dies.

SYMPTOMS.—The most common are, drowsiness, watery and somewhat full and inflamed eyes, a disposition to reel, feebleness, a bad appetite, the head generally hanging down, or resting on the manger. There is little or no fever, and the dung and urine are very little altered. The horse soon begins to reel, and falls down, and sometimes is so outrageous as to bite every thing in his way.

CURE.—In the first place, bleed him well as soon as you possibly can, by striking the veins in several places at once, and taking away four or five quarts at one time; and, in order to raise up his head and shoulders, support them with plenty of straw. If he survive the first fit, cut several rowels, give him clysters at night and morning, made of barley-water and a little sweet oil and salt; and blow up his nostrils a little Cayenne pepper, or white hellebore. Also give him

4 drams of Bark.
16 grains of Turbeth Mineral.
 $\frac{1}{2}$ ounce of Camphor.

Give it in a little warm ale. When ale cannot be conveniently had, any other beverage to disguise the taste will answer. If the horse be outrageous, give him

1 ounce of Tincture of Opium.
1 gill of Syrup of Poppies.
1 ounce of Tincture Gualacum.

Be careful not to let him knock his head, for it will increase the disorder. If he gets through the first fit, give him two ounces of crocus metalorum every day, to thin his blood for fear of a relapse. It will be proper to give him the following ball once a month for some time after:

1 ounce of Rhubarb, in powder.
 $\frac{1}{2}$ ounce of Jalap.
1 dram of Calomel, 8 drams to the ounce.

To be made into a ball with syrup of buckthorn. This ball will be of great use in thinning the blood, and preventing a return of the disorder; for when a horse has had one fit of it, he is very likely to have another, if care be not taken to prevent it.

Convulsions, or the Stag Evil.

Of all disorders to which horses are subject, this is the worst, and is scarcely discoverable till the horse falls down raging mad. It seizes him all at once, without any previous warning. He raises his head, with his nose toward the rack, pricks up his ears, and cocks his tail. In this posture he continues, and those who do not understand the disorder never suppose that he ails any thing of consequence. But other symptoms soon convince them of their mistake; for his neck grows stiff, cramped, and almost immovable; his jaws are locked, and every tendon in his body becomes stiff. If he can get his mouth open he will bite any thing that comes in his way; and if he lives a few days in this condition, several knots will arise on the tendinous parts of it. Every muscle is so much cramped and extended, that the horse looks as if he were fastened to the place, with his legs stiff, wide, and staggering, and the skin drawn so tight over every part of his body, that it is almost impossible for him to move; and if you attempt to make him walk, he will be ready to fall at every step, unless he be well supported. At the same time his eyes are so fixed by the contraction of the muscles as to give him a dead look. He snorts and sneezes often, pants continually, and his shortness of breath increases till the distemper takes a favorable turn, or the horse falls down and dies.

CURE.—In the first place, bleed plentifully, unless the horse be old and low in flesh, or taken from some hard duty, and then you must not take so much blood. After bleeding give the following ball, if you can get it in, but the horse is very often jaw-locked till nothing can be got in but by clyster-pipe put between his fore and axle teeth.

½ ounce of Assafetida.

½ ounce of Gum Gualacum.

½ ounce of Gum Camphor.

Make them up into a ball with honey, and give one of these balls every twelve hours, for two days, if you can get them in; and if not, dissolve them in a little hot beer, and give them with the clyster-pipe. Be careful to powder the Gums. Then make an ointment or lotion of the following:

1 ounce of Oil of Spike.

1 ounce of Oil of Amber.

1 ounce of Oil of Bricks.

1 ounce of Spirits of Sal Ammoniac.

Shake them well together, and rub the jaws just below the

ears, where they lock in the upper chap ; also rub the small of the back well, where the cross bones join the back bone.

In this, as in most other disorders, the body should be kept gently open with laxative purges and emollient clysters. When the jaws are so locked that you can not get any thing in, do not open them by force, for that would increase the disorder instead of relieving it.

Sheep skins, newly taken off, and laid with the flesh side to the horse, will sweat him greatly, and by that means draw a quantity of water from the blood : indeed, there are few things that will relieve a locked jaw more : but if they be laid on the loins, they must not lie above three hours at a time before they are turned with the wool side to the horse. You may keep the skins on twenty-four hours if you change sides every two or three hours.

If the horse can not take either food or water in at his mouth, he must be supported by clysters, made of barley-water and milk, and given both at the mouth and the fundament. I once supported a large wagon-horse in this manner for twelve days, and he recovered.

Convulsions are caused by different things, but often by bots in the stomach ; for these destructive vermin suck up the juices that should feed the blood, and bring it into a thin, bad state ; indeed they are sure to destroy the horse by one means or other when there are a great number of them in the stomach.

When you suspect that these vermin are the cause of the disorder, give the ball recommended to destroy bots. If the horse get better the first time, be sure to guard against a relapse, for he will not be apt to get better a second time.

Tapping under the jaws, and at the breast, is sometimes of great service in this disorder, but I am of opinion that sweating with sheep skins will relieve much sooner. If the stiffness of the jaws continues after the convulsions have ceased, the following medicine should be given :

$\frac{1}{2}$ ounce of Matthews' Pill.
 $\frac{1}{2}$ ounce of Assafœtida.

Make them into a ball, and give it twice, (one day between the doses,) and it will give relief.

Fevers.

Horses are subject to few disorders which are not attended with more or less fever.

CAUSES.—Fevers are often brought on by sudden heats and colds ; by going out of warm stables into cold ones ; by be-

ing clothed, and then having the clothes stript off; and by being turned out to grass; for many people turn their horses out to grass in the morning, and let them lay out, which is quite wrong: for when they are turned out to grass, to be there night and day, it is best to turn them out at night, for then they will graze all night; but if you turn them out in the morning, they will fill themselves in the day-time, and lie still all night, which is the way to catch cold. Most fevers are brought on by colds, therefore be careful to keep your horses as much as possible from catching cold.

SYMPTOMS.—The horse is remarkably restless, ranging from one end of his rack to the other: his flanks work, his eyes appear red and inflamed, his tongue is parched, and his breath hot and of a strong smell. He often smells at the ground, he loses his appetite, and though he will take hay into his mouth, he does not chew it; his whole body is hotter than usual, but not parched: he dungs often, but little at a time, and it is generally hard, and in small pieces; his urine is high-colored, and he generally stales with pain and difficulty; he is always craving for water, but drinks very little at a time; and his pulse is much quicker than usual.

CURE.—Whenever a fever takes place, the first part of a cure is bleeding, and if the horse be strong and in good condition, the quantity should be two or three quarts. When this has been done, give him a pint at a time of the following infusion, three or four times a day:

4 ounce of Juice of Liquorice.
 4 ounces of Liquorice Root.
 2 ounces of Salt Petre.
 4 ounces of Salt of Tartar.
 8 ounces of Good Raisins.
 2 ounces of Aniseeds.
 2 Drachms of Saffron.

Boil all these together in six quarts of water, for ten minutes; let it stand till cold, and then strain it off. It is one of the best medicines for colds, coughs, hoarseness, or fevers, in either horse or man; and if it were more known, and more used, it would give greater relief in violent colds than any thing yet found out. It is kind in its operations, opening to the lungs, works gently by stool and urine, is free in its passage, and opening in its nature.

The horse should scarcely eat any thing but mashes made of linseed and bran, and given in small quantities. If he refuses them, let him have dry bran sprinkled with water, and put a little hay into his rack, as a small quantity of it will not hurt him, and a horse will often eat hay when he

t off; and by
n their horses
out, which is
o grass, to be
t at night, for
n them out in
day-time, and
e cold. Most
areful to keep
cold.

ranging from
work, his eyes
shed, and his
smells at the
will take hay
e body is hot-
n, but little at
d pieces; his
rith pain and
t drinks very
than usual.

first part of a
d in good con-
narts. When
ne of the fol-

for ten min-
off. It is one
ess, or fevers,
known, and
nt colds than
ations, open-
e, is free in

ashes made
ities. If he
with water,
quantity of it
ay when he

will not eat any thing else. His water should be rather warm, and given often, but in small quantities; and his clothing moderate, too much heat being pernicious in a fever. If he refuse his meat, do not let it lie before him, but take it away, and clean his rack and manger. If he be able to go about, a little walking exercise in the open air will be very proper, but you must be careful not to get him wet.

This method, with good nursing, will often be sufficient to restore the horse to health; but if he refuse his meat, more blood should be taken from him, and the drink continued; and if his dung be hard and knotty, a clyster should be given.

Take Marshmallows and

Camomile Flowers, a handful each.

Boil one-third away in three quarts water—strain, and add

4 ounces of Venice Treacle.

1 pint of Pale Rape Oil.

The above will make three clysters, to be given at four hours distance. If his pulse continue high and quick, give the following:

2 ounces of Nitre.

2 ounces of Cream of Tartar.

4 ounces of Glauber's Salt.

2 ounces of Lenitive Electuary.

Dissolve them in hot water, give one-half, and the other half the day following. If the horse be very open in his body, you need not give the above; but if dry, be sure to give it him. If he be very open, give him four drams of bark in a gill of red port. By pursuing this method, the horse will begin to recover, and will relish his hay, though his flanks will continue to heave for a fortnight. Nothing more will be requisite to complete the cure than walking him abroad in the air, and giving him plenty of clean litter to rest on in the stable.

There is another and much worse kind of fever to which horses are often subject, and which often proves fatal if not properly treated, viz.:

A Compound Fever.

SYMPTOMS.—The symptoms of this disease are—a slow fever, with great depression; and sometimes inward heat and outward cold, and at other times heat all over, but not excessive. The horse's eyes are moist and languid; his mouth is continually moist, so that he is not desirous of drinking, and when he does drink, a very little satisfies him; he eats very little, and moves his joints in a loose, feeble manner,

grating his teeth very disagreeably; his body is generally open, his dung soft and moist, and he stales irregularly, sometimes making little water, and at others a large quantity, which is of a pale color, and has very little sediment.

CURE.—In the first place, take from the horse a moderate quantity of blood. Let it not exceed three pints, but repeat the operation according to its strength, if there be any tendency to inflammation; after this the nitre drink already described may be given, with the following addition:

- 1 ounce of Snake Root.
- 3 drams of Saffron.
- 3 drams of Camphor, dissolved in Spirits of Wine.

The horse's diet should be scalded bran; and linseed, boiled, and wrought up with bran. Also give him the best hay by a handful at a time. It is often necessary to feed him by the hand, for sometimes he is not able to lift his head to the rack.

In this disease, drinking is absolutely necessary to thin the blood; and therefore if the horse refuses warm water, he should be indulged with such only as had the cold taken off. This may be done with a hot iron, or by letting it stand in the pail in a warm stable; and this will be better than forcing warm water on the horse's stomach. If this method does not prove sufficient, but the fever shall continue to increase, the following ball should be given immediately, as the danger augments every hour.

- 1 ounce of Camphor.
- $\frac{1}{2}$ ounce of Gum Myrrh.
- 1 ounce of Squills.
- 2 drams of Castor.

Make them up into two balls, and give one at night, and the other in the morning. If no better in a short time, give the following infusion:

- 1 ounce of Snake Root.
- 2 ounces of Gentian Root.
- 2 ounces of Lemon Peel.
- 2 drams of Saffron.

Boil these well together in three quarts of water, and give a pint once a day. If the above ball fails of success, give the following:

- 1 ounce of Camphor, dissolved in Spirits of Wine.
- 1 ounce of Sal Ammoniac.
- 1 pint of good Vinegar.

Put them all together, and stir them about till the fume subsides. This is for two doses to be taken at twelve hours' distance, diluted with water. There is not perhaps a more

dy is generally
les irregularly,
s a large quan-
tle sediment.

orse a moderate
ints, but repeat
re be any ten-
drink already
ddition :

drirts of Wine.

; and linseed,
e him the best
essary to feed
to lift his head

ecessary to thin
warm water, he
cold taken off.
ing it stand in
be better than
If this method
continue to in-
mmediately, as

at night, and
a short time,

ater, and give
success, give

Wine.

till the fume
twelve hours'
erhaps a more

powerful and effectual medicine known than camphor in all kinds of putrid fevers, it being active, attenuating, and particularly calculated to promote urine and perspiration, the two principal outlets by which relief is to be obtained ; and if this medicine were more often given than it is, it would be a greater credit to the farrier, and give greater relief to the horse.

If the horse be costive, clysters, or an opening drink, should be given ; and should he purge moderately, be careful not to suppress it ; but if it continue so long as to enfeeble the horse, give him a little red port wine and bark. Also observe to let the animal drink plentifully, for that will greatly promote the operation of the above named medicines, as both the disorder and the medicines will cause a thirst. If the horse can bear walking about, a little open air will be proper. but be careful to keep him well covered. Particular attention should also be paid to his staling, which if it flow in too great quantities, must be repressed by proper astringents, and by giving him lime-water ; and, on the other hand, if he stale so little as to occasion a fullness or swelling in his body and legs, give the following drink :

- 1 ounce of Nitre.
- 2 ounces of Castile Soap.
- 1 ounce of Venice Turpentine.
- 2 drams of Oil of Juniper

Make them into a ball with liquorice powder, and give them at twice twenty-four hours distance. These balls may be given as occasions may require, and are very proper to convey off the greasy, slimy matter from the passage of the urine, and to settle swelled legs.

These are the best methods of management, and will generally prove successful ; but sometimes art will fail, and the horse will discharge a greenish or a reddish gleet from his nostrils, and sneeze very frequently ; he will continue to lose his flesh, become hide-bound, refuse his meat, swell about his joints, and his eyes will appear fixed and dead ; a purging also ensues, and a dark-colored, foetid matter is discharged. When these symptoms appear, the case may be considered desperate, and all attempts to save the horse will be fruitless.

In this disorder you must take care not to let the horse eat too much, for his diet should be light, and in small quantities at once, and increased gradually as he may gain strength. When his skin feels kind, his ears and feet continue moderately warm, his eyes look lively, his nose remains clean and dry, his appetite mends, he lies down with

ease, and dungs and stales well, you may conclude that the danger is nearly over, and that nothing more is needful but care to complete the cure. On the contrary, by overfeeding, you will run the risk of bringing on a bad surfeit, and the horse may be, according to the old saying,—*killed with keeping*.

Sometimes the fever returns; so that every one who has a horse in a fever should be careful of cold for some time after, as his blood is left in a thin, bad state. His legs will probably be subject to swell; and if the swelling leave a dimple when you press your finger upon it, it is a sign of a dropsy; in which case it will be advisable to put two rowels on each side of his belly, and to give him half an ounce of the best yellow bark every day for some time. At other times a fever leaves a running at his nose; of a thin, yellow, glueish matter, and small swellings below his ears and chaps. When you find these symptoms, give one ounce of crocus metalorum every day in a mash of bran, and rub the swellings with mercurial ointment.

Care should be taken to keep the head and throat warmer than common, as the kernels about the latter are swelled; and also to promote a free perspiration, and to increase the running at the nose, which has the same effect in horses as spitting has in the human species; but never syringe the nose, as is often done, to promote the discharge, for it has an effect quite contrary, and lessens the quantity of matter instead of increasing it; and checking the discharge of matter at the nostrils often causes swellings at the glands, and other bad consequences. Let me once for all remind you that all such discharges are critical, and thrown by Nature to free herself from the load that oppresses her, and consequently should by all means be promoted.

A Broken Wind.

This disorder may sometimes be prevented, but can not be cured; and it has hitherto been as little understood as any to which a horse is subject. People have had various opinions respecting its cause, and why some horses are more subject to it than others; but of all the opinions hitherto delivered, that of Mr. Gibson seems the best founded. He thinks that it is frequently owing to the hasty or injudicious feeding of young horses for sale; by which means the growth of the lungs is rapidly increased, and all the contents of the chest so much enlarged, that in a few years the cavity of the chest is not sufficient to contain them when

conclude that the
re is needful but
by overfeeding,
surfeit, and the
ing,—*killed with*

ery one who has
d for some time
. His legs will
welling leave a
it is a sign of a
put two rowels
alf an ounce of
ime. At other
a thin, yellow,
his ears and
one ounce of
n, and rub the

throat warmer
r are swelled;
o increase the
ct in horses as
er syringe the
rge, for it has
tivity of matter
harge of mat-
he glands, and
ll remind you
vn by Nature
er, and conse-

but can not
nderstood as
had various
e horses are
ions hitherto
ounded. No
r injudicious
means the
all the con-
ew years the
them when

they are expanded to perform their proper functions. Nor is this opinion founded upon bare conjecture, for horses that have died broken-winded have been opened, and the lungs and other parts found too large for the chest. But although hasty feeding is often the cause of this disorder, yet it is not always, for a narrow chest may cause it. It has been observed that horses rising eight years old are most subject to it. The reason of this is, because a horse arrives at his full strength and maturity at that age. At six he generally finishes his growth in height; then he lets down his belly, and spreads, and all his parts gain their full size; so that the pressure on the lungs and the midriff is now increased.

Also, when the horse catches cold and gets a dry cough, the lungs are much larger than they ought to be, and at that time riding sharply is enough to force the lungs so hard against the midriff as to force a passage through it.*

A few years back some people pretended to cure the complaint by boring; but none were ever cured by it yet, nor ever will be. They made a hole above the fundament, to let out the wind that was forced through the midriff into the bowels; and this caused the horse to be continually discharging wind out of the place; so that the pretended cure was worse than the disease.

Dissections of horses that have died broken winded, have sufficiently proved the truth of the above observations; and that not only the lungs, together with the heart and its bag, were preternaturally large, but also the membrane which divides the chest; and that the midriff was remarkably thin. In some horses the disproportion has been so great that the heart and lungs have been almost twice their natural size, yet perfectly sound: and without any ulceration whatever, or the least defect in the windpipe or in its glands.

From these observations it abundantly appears, that the enormous size of the lungs, and other contents of the chest, by hindering the free action of the midriff, is the principal cause of this disorder; and as the lungs are found much more fleshy than usual, they must consequently have lost a great part of their spring and tone.

Therefore, as this disorder is caused by the largeness of the lungs, we may conclude that it is one of those diseases which can not be cured by art; and that the boastings of those who pretend to cure it are built on a sandy foundation. They may indeed relieve the complaint, but will never cure

* The Midriff, or Diaphragm, is that which is commonly called the Skirts, and separates the Chest (where the lungs lie) from the Bowels.

it, for an absolute cure is not in the power of any human being. All that I can do is to lay down some rules which have a great tendency to prevent this disorder, if pursued in time; and some remedies that will afford relief when it has taken place and render the horse capable of performing good service, notwithstanding his misfortune.

SYMPTOMS.—The first symptoms of a Broken Wind is an obstinate dry cough, which is neither attended with sickness nor loss of appetite; but on the contrary, with a disposition to foul feeding, eating the litter, and drinking large quantities of water.

PREVENTION.—When a horse is troubled with an obstinate dry cough, and eats his litter, it will be necessary to bleed him, and to give him the mercurial physic already prescribed, repeating it two or three times. Afterwards give the following balls for some time, which have been found of great service:

- 4 ounces of Gum Ammoniacum.
- 4 ounces of Galbanum.
- 4 ounces of Assafoetida.
- 4 ounces of Squilla.
- $\frac{1}{2}$ ounce of Saffron.
- 6 drams of Cinnabar of Antimony.

Make the whole up into balls with honey and a little liquorice powder, and give one about the size of a pullet's egg every other morning. This is a very good ball for a dry cough.

But it is not enough to give proper medicines; the horse's diet should also be carefully attended to at the same time, if we would hope for success. In order to this, the horse should eat very sparingly of hay, which, as well as his grain, should be wetted with chamber-ley, which is much better than water. Chamber-ley is best for this purpose, because of the volatile salts which it contains, as they are a means of removing the thirst. For the same reason, garlic is very efficacious in this disorder. Two or three cloves being given in each feed; or three ounces bruised, and boiled in a quart of milk and water, and given every morning for a fortnight, has been found very serviceable. So easy a remedy should never be neglected; for, by warming and stimulating the solids, and at the same time dissolving the tenacious juices which choke up the vessels of the lungs, it greatly relieves this complaint. Moderate exercise should never be omitted; and although broken-winded horses are not able to endure much labor the first summer, yet many have been found less oppressed the second, and scarcely perceptibly

of any human
ne rules which
r, if pursued in
ief when it has
performing good

en Wind is an
d with sickness
with a disposi-
drinking large

h an obstinate
ssary to bleed
dy prescribed,
give the fol-
found of great

a little liquo-
pullet's egg
ball for a dry

s; the horse's
the same time,
his, the horse
as his grain,
much better
dose, because
are a means
garlic is very
being given
ed in a quart
for a fortnight,
medy should
mulating the
acious juices
atly relieves
ver he omit-
not able to
have been
perceptibly

affected the third, being then able to perform a long journey, and to endure great fatigue. A horse kept constantly in the field, when not in work, will be able to do good service for many years.

It may not be improper to observe that those who hope to cure a broken-winded horse, or even one that is troubled with an obstinate cough, by putting him to grass, will find themselves wretchedly mistaken; for on being taken into the stable and fed with dry meat, he will be much worse than before; and some that had only a dry cough when they were put to grass, have returned broken-winded. Therefore, always remember that if you can not keep a horse of this description constantly abroad, it is best not to put him to grass at all, as, instead of curing, it will tend to augment the disorder.

In short, the grand secret of managing horses of this kind, consists in having particular regard to their diet and exercise. A moderate quantity of hay or grain, and water, should be given at a time, and the former constantly moistened, to prevent their wanting too much of the latter. They should have moderate exercise, but never any that is violent. By this method, and giving the following ball once every fortnight or three weeks, the horse will be able to do good service for many years:

6 drams of Socotrine Aloes.
2 drams of Myrrh.
2 drams of Galbanum.
2 drams of Ammoniacum.
2 ounces of Bayberries, in powder.

Make the whole into a ball with a little oil of amber, and a sufficient quantity of syrup of buckthorn. This ball operates so gently that there is no need for confinement, except a little the day following that on which it is given. The horse must have warm mash and warm water, and the utmost care must be taken to prevent his catching cold.

A Consumption.

SYMPTOMS.—A Consumption is a want of nourishment, and a waste of flesh. The horse's eyes look dull, his ears and feet are commonly hot, he coughs violently by fits, sneezes often, and groans at the same time; he gleans at the nose, and sometimes throws a yellowish matter, rather curdled, from his nose; his flanks have a quick motion, and he has little appetite to hay, though he will eat grain, but he grows hot after it.

CAUSES.—Damp stables are most likely to bring on this

disorder, though it may be brought on by many other things. In my time I have known many horses suffer much by damp stables. I knew a gentleman who had two valuable horses, and he built a brick stable for them, without any air-holes above their heads. He put the horses in as soon as the stable appeared dry, and their heat soon caused the walls of the place to sweat, and to run down with water, by which means both the horses were thrown into a Consumption and died. I mention this to caution others.

CURE.—The first, and indeed one of the principal things to be done, is to bleed in small quantities. A pint, or at most a pint and a half, is sufficient at once, and the operation is to be repeated whenever the breath is more than commonly oppressed. We are assured, by dissection, that in a Consumption both the glands of the lungs and the mesentery are swelled, and often indurated. The only medicines that can be depended upon, are mercurial purges and ponderous alteratives. I have already given you examples of the former, and the following is a formula of the latter. Mix

4 ounces of Crocus Metalorum.

1 ounce of Calomel pp.

1 pound of Gum Gualacum, finely powdered.

Give about an ounce every day in a mash of bran and linseed. Iceland liverwort, a handful boiled in a gallon of water, is much better to make mashes up with than water; for it is a great helper of the blood. But it is to be observed that nothing will answer so good an end as Spring grass; so that if the horse be afflicted with this disease in Spring time, turn him out to grass as soon as you can; and if the nights be cold, turn him out in the day-time, and take him in at nights. Salt marshes are the most proper places, when they can be met with. When a horse has had this disorder, he can never more bear cold and hard service as before.

A Scouring, and Other Disorders of the Intestines.

You should consider well what the Scouring proceeds from, whether it is caused by foul feeding, bad water, hard exercise, sudden heat or cold, an overflowing of bile, or a weakness of the intestines.

If it is brought on by foul feeding, or bad water, it should not be stopped, but rather be promoted; for it should be remembered that Nature by this means throws off the seeds of disease, and evacuates the morbid matter which would otherwise be retained to the great disadvantage, and per-

ny other things.
much by damp
valuable horses,
at any air-holes
as soon as the
caused the walls
water, by which
consumption and

ncipal things to
int, or at most
operation is to
an commonly
that in a Con-
mesentery are
icines that can
nd ponderous
mples of the
atter. Mix

ed.
of bran and
a gallon of
than water ;
is to be ob-
nd as Spring
is disease in
u caz ; and
ne, and take
oper places,
has had this
d service as

Intestines.

g proceeds
water, hard
f bile, or a

r, it should
should be
the seeds
ich would
, and per-

haps to the destruction, of the animal. The great difficulty therefore consists in knowing when these discharges are critical and salutary, and when detrimental and noxious ; for the former must not be checked, but the aid of medicine must be called in to put a stop to the latter.

For instance,—if a healthy horse, upon taking cold, or after hard riding, over feeding, or at the beginning of a slight fever, have a moderate purging, you must be careful not to stop it, but on the contrary to promote it, by an open diet, and plenty of warm gruel. But if this purging continue a long time, with smart gripings, and the inner skin of the bowels come away with the dung, and the horse lose both his flesh and his appetite at the same time, recourse must immediately be had to proper medicines ; among which the following are very effectual. I do not wish any one to give medicines upon merely hearing the names of the drugs, but to know first in what manner the drugs will operate.

Take 1 ounce of Rhubarb, in powder.
2 drams of Myrrh, do.
2 drams of Saffron.

Give altogether in warm ale, and warm water for two days after. This dose will not only work gently, but will be of great service to the horse, as it will bring away the slime which lodges in the small intestines, and correct the bile of the stomach, which is the cause of this disorder. When the disorder continues, and the horse's flesh keeps wasting away, recourse must be had to astringents. Tormentil Root dried, and pounded in a mortar, and put through a sieve, is one of the best astringents yet found out, though very little known. The dose is from an ounce to an ounce and a half. I believe that this valuable root has done more good in my time, in stopping looseness and bowel complaints, than any thing else. When the purging is attended with fever, a different method of practice is necessary.

Take $\frac{1}{2}$ ounce of Rhubarb, in powder.
1 ounce of Lenitive Electuary.
 $\frac{1}{2}$ ounce of Camphor.
1 ounce of Powdered Ginger.

To be given in a pint of old ale. This is a very proper medicine when the horse is troubled with a fever ; but if he have no fever upon him, give the following :

1 ounce of Tormentil Root, in powder.
 $\frac{1}{2}$ ounce of Japan Earth, do.

Give these in red wine, or if that be thought too expensive, in oak bark tea. Japan earth is a great healer of the bow-

els. Repeat this last medicine three or four times to allow it a fair trial ; giving the horse at the same time but little exercise, for he can not then bear much. Should this medicine fail, and the disorder increase instead of decreasing, which may be known by his flanks and belly being full and distended, and his appearing to suffer strong griping pains, give the following clyster :

$\frac{1}{2}$ ounce of Isinglass, dissolved in a quart of warm milk.
2 ounces of Mithridate.

Sometimes the flux is so violent as not to be overcome by the preceding medicine, when recourse must be had to the following. Boil a handful of oak bark in a quart of water, strain it off, and add

2 ounces of Tormentil Root, in powder.
2 ounces of Bole.

Give them all together. This should be repeated once a day, for two or three days.

The practitioner should carefully attend to the symptoms that accompany this disorder ; for if the discharge be attended with any acrid mucus, or slime, the griping pains being very severe, there is then a sure indication that the common lining of the bowels is wasted away ; and then it will be necessary frequently to inject the following clyster, warm, in order to prevent the fatal consequences which will otherwise soon ensue :

4 ounces of Starch, dissolved in a quart of water.
Half a pint of Sweet Oil.
Yolks of three Eggs, well broken.
A little Loaf Sugar.

This will do for twice, at four hours' distance.

In treating this disease, be exceedingly careful to follow my directions, and to get good drugs, for your success will depend upon this, and a cautious watching of the symptoms.

It is also necessary to observe that some horses, from having weak stomachs and bowels, throw out their aliment undigested, and their dung is habitually soft, and of a pale color ; they also feed sparingly, and are always low in flesh. This complaint, which often proves fatal at last, may be removed by the following medicines :

6 drams of Socotorine Aloes.
3 drams of Rhubarb, in powder.
1 dram of Myrrh.
1 dram of Saffron.

Make all up into a ball with syrup of ginger. After the above stomachic purge shall have been given two or three

times, a pint of the following infusion should be given every morning :

Take 2 ounces of Orris Root,
A handful each of
Gentian, Columbia Root,
Winter Bark, Aniseeds,
Orange Peel, Fennel Seeds,
and Camomile Flowers.

Boil all together in a gallon of strong ale, and when cold, clear it off, and add one pint of spirits of wine. If this be thought too strong, two quarts more of ale may be added. This is an excellent cordial both for healing and strengthening the stomach and bowels. These are the best methods of treating the above disorders, to which horses are often subject, and in which they are often lost for want of proper treatment.

But before we conclude this chapter, it is necessary to observe that the scourings which succeed long-continued sickness, such as the Farcy, Putrid Fevers, or an inflamed state of the blood, where bleeding and other proper evacuations have been neglected, too often terminate fatally ; especially when the creature discharges a foetid slime, and when the same matter gleans from his nose ; for in these cases the blood is dissolved, and the whole mass of the fluids is become putrid, and discharges itself by those drains.

Disorders of the Kidneys and Bladder.

Inflammation of the kidneys and bladder are sometimes brought on by other disorders, and often by sprains, hard exercise, catching cold when hot, or from the want of proper care.

SYMPTOMS OF A HURT IN THE KIDNEYS.—The horse is very weak in the back and loins ; he stales with difficulty, is faint, eats very little ; his eyes appear languid and of a dead color ; his urine is foul, thick, and often bloody, especially after a violent strain ; he can not move backwards without great pain, which may be seen at every attempt. It is sometimes hard to distinguish an inflammation from a sprain, or what is called, "tified in the back." There is no method but observing the gait of the horse. If he have got tified in the back, he will be hard put to it to keep from coming down behind, and indeed will often come down ; his eyes will look rather red, but his urine the same as before. A tife lies in the marrow, or pith of the back.

CURE.—The principal remedy for a hurt in the kidneys is bleeding, which should be done pretty plentifully, as by this

means an inflammation will be prevented; and if you have reason to think that the inflammation is already begun, from the creature's being feverish, and staling with great difficulty, the operation should be repeated; for unless the inflammation be prevented, or immediately removed, the consequence will be fatal. But, although bleeding is the principal, it is not the only remedy, for rowelling is of great service. Put a rowel on each side of his belly, and give the following balls twice a day, in a pint of the decoction of marshmallows, having an ounce of gum arabic, and an ounce of honey dissolved in it:

1 ounce of Salts of Prunella.
6 drams of Spermaceti.
2 ounces of Castile Soap.

Add as much honey as will make a ball; and if the urine be bloody, an ounce of Japan earth must be added. If the fever continue, you must repeat the bleeding, and give emollient clysters, and the cooling, opening drink, before recommended for Fevers, till it abate. These methods will often prove successful, but sometimes the disease is too obstinate to be overcome by them, and the urine still passes with pain and difficulty. Recourse must then be had to the following balls, and they must be repeated twice a day till the horse stales without pain, and his urine becomes clear, and without any purulent settlement:

1 ounce of Venice Turpentine.
1 ounce of Castile Soap.
6 drams of Nitric.
2 drams of Myrrh, in powder.

Make the whole into a ball with honey, and wash it down with a strong decoction of marshmallows.

These are the best methods of treatment in this disorder, and will in general prove successful. Sometimes indeed this malady is too strong for the power of medicine, and then the urine continues turbid, and daily becomes of a deeper color, with a fetid smell, a sure sign that the kidneys are ulcerated; which generally terminates in a consumption, and the creature becomes absolutely incurable.

In treating of the Colic I have recommended a method for removing the Stranguary, when it proceeds from wind, or from dung pressing upon the neck of the bladder; but sometimes it proceeds from an inflammation, and a retention of urine.

SYMPTOMS OF AN INFLAMMATION OF THE NECK OF THE BLADDER.—When a horse is seized with a Stranguary from the above cause, he will make frequent motions to stale,

standing wide and straddling, his bladder being full of urine, and his flanks distended; he will be uneasy, constantly shifting his hind feet, and often giving clicks in his motions; he also sometimes hangs his head, and then raises it suddenly.

CURE.—First bleed largely, and then give the following :

- 1 ounce of Spirits of Sweet Nitre.
- 2 ounces of Syrup of Marshmallows.
- 1 ounce of Venice Soap.

Cut the soap small, dissolve it in a gill of hot water, put the above to it, and give it to the horse. Repeat it every eight hours. Also dissolve an ounce of gum arabic and an ounce of nitre in a gallon of water, and let him drink plentifully of it. If he will drink it of his own accord, it is best; but if not, horn a little into him; for it will greatly tend to remove the cause of the disease, and consequently to terminate the effects.

There is a disease of the kidneys, viz. : a Diabetes, or profuse staling, which produces effects directly opposite. This disorder is seldom cured in old horses, as their fibres are become rigid, and unable to perform their office, a misfortune which all the power of medicine can not remove. But in young horses this disease is often cured, and the following method will generally be attended with success :

CURE OF A DIABETES.—In order to cure this threatening disorder, great care must be taken not to let the horse drink too much water, and never to give him any moist food. Attention to these particulars will go a great way towards a cure : and instead of giving him common water to drink, give him lime-water; to make which, take about three pounds of lime, unslacked, put it into a clean vessel, and for every pound of lime pour six quarts of water into it; let it stand three days, take the scum off the top, and give the horse the water to drink. Be careful not to stir the lime at the bottom of the vessel. This is a very clear, wholesome water, and very good for many disorders. In the mean time the following medicine should be given :

- 1 ounce of Peruvian Bark.
- 1 ounce of Japan Earth, in powder.
- 1 ounce of Irish Slate, do.

Give these in lime-water, and repeat as often as needful.

Some Farriers give strong alum possets two or three times a day; but this kind of treatment cannot be proper, for the alum takes so fast hold of the juices of the stomach, that if the horse were not ill, it would be enough to make him so.

The Molten Grease.

This is so common a disorder, that it ought to be well understood; but, alas! like many others, it is often mistaken for something else. This disease is a melting down of the fat of the horse's body, caused by violent exercise in very hot weather; or if the horse be full of flesh, it may happen in cold weather. Hard riding, or sudden colds, will bring on this disorder.

SYMPTOMS.—It is always attended with a fever, heat, restlessness, starting and trembling, inward sickness and shortness of breath. Also the horse's dung is extremely greasy, and he often falls into a scouring. His blood, when cold, is covered with a thick scum of fat, of a white or yellow color, but generally the latter. The congealed part, or sediment appears like a mixture of size and grease, and is so extremely slippery that it will not adhere to the fingers; and the small proportion of serum is also slippery and clammy. The horse soon loses his flesh and fat, the latter being probably dissolved into the blood. Such as have sufficient strength to sustain the first shock, commonly become hidebound, and their legs swell greatly; and in this state they continue till the blood and juices are rectified; and if that is not done effectually, the Farcy, or an obstinate Surfeit, is generally the consequence, which can not be removed but with the utmost difficulty.

CURE.—In the first place bleed pretty plentifully, and repeat the operation two or three days successively; but take care that you take only a small quantity at a time after the first bleeding, as otherwise the creature would be rendered too weak to support himself, and his blood too poor to be easily recruited. As soon as he has been bled the first time, let two or three rowels be put in, and the emollient clysters prescribed for Fevers be thrown up daily, to mitigate the fever, and to cleanse the intestines from the greasy matter. At the same time plenty of water-gruel should be given, and sometimes a small quantity of water, with a little nitre dissolved in it. The latter will be of great service, as it will prevent the blood from running into grumous concretions, and proving the source of innumerable disorders, if not causing a total stagnation, and consequently the death of the animal.

The horse must be treated in this manner till the fever be entirely gone, and he shall have recovered his appetite; and then it will be necessary to give him three or four purges, a week distant from each other, which will make

him stale and perspire plentifully, and at the same time bring down the swelling of his legs. The following is well calculated for the purpose :

6 drams of Socotrine Aloes.
4 ounces of Gum Gualacum, in powder.
2 drams of Ginger.
2 drams of Jalap.
2 drams of Oil of Juniper.

To be made into a ball with syrup of buckthorn. By pursuing this method the horse will soon be recovered, for this purge will mend his appetite, and increase his flesh. If it be too weak, add a dram more of aloes. It will bring down his swelled legs, and carry away all the superfluous matter that clogs the blood. When you give the physic, be careful to give plenty of warm water all the time.

A Surfeit.

Some people pronounce every ill-thriven horse surfeited, whether he is so or not. A Surfeit is nothing more than the effects of an ill cured disease, and therefore what is called a Surfeit in horses is very different to the disease which bears that name in the human body ; the latter being the beginning of a disease, and the former the remains of one.

SYMPTOMS.—The horse's coat will stare, look of a rusty color, and even appear dirty, although the greatest pains have been taken to keep him clean. His skin will be covered with scales and dandruff, which will appear like meal among the hair, and when cleaned off will be followed by a succession of the same matter, occasioned by the perspiration being obstructed. Some horses will be covered with a kind of scab, sometimes moist, attended with heat and inflammation, and the humor so very sharp, and causing so violent an itching, that the creature is incessantly rubbing himself, and by that means making himself raw in different parts of his body. Some horses have neither scales, dandruff, nor scab ; but look dull, sluggish, and lazy. Some are hidebound, and others afflicted with flying pains and temporary lameness. In short, the symptoms are very various, and almost as numerous as those of the scurvy itself.

CAUSES.—The causes are as various as the symptoms. Some horses are surfeited by high feeding and want of proper exercise ; which produce a bad digestion, and generate ill humors. Some are surfeited by unwholesome food ; some by hard riding ; some by drinking cold water when they are hot ; some by bad or improper physic, and some by standing

in stables through which the rain drops, or by lying wet. But as many also get surfeited by standing when hot at the doors of public houses, such bad practices should be guarded against.

CURE.—If we duly consider the nature of Surfeits, their cure will be much more easily performed. All allow that they arise from a bad state of the blood; but how is this to be remedied? By bleeding and purging. But this must be done in a very gentle manner. Take about a quart of blood, and the next day give the following ball:

- 1 ounce of Socotrine Aloes.
- 1 dram of Calomel, pp.
- 2 drams Oil of Aniseeds.

Add as much lenitive electuary as will make it into a ball. Give plenty of warm water to work it off. Repeat both bleeding and physic in eight days; and when the last dose is wrought off, give him six ounces of the cordial balls in a little warm ale. Also give a spoonful of the following powders every day in a mash:

- 4 ounces of Flour of Brimstone.
- 4 ounces of Crocus Metalorum.
- 4 ounces of Nitre, in powder.

Mix all up well together. By persevering in the above method you may cure the most inveterate Surfeit, and if any scabs or runnings appear in the skin, rub them with the following:

- 4 ounces of Sulphur Vivum.
- 2 ounces of White Copperas.
- 2 ounces of White Hellebore, in powder.

Mix these powders with churn-milk, rub the places affected well, and the grievance will soon disappear. Sometimes a Surfeit settles in the legs, and they swell much, and then break out and run very much. When this is the case, two taps put into the lower belly will be of great use. Staling balls should be given every third day, and plenty of warm water to work them off with. Make the staling balls as follows:

- 1 ounce of White Resin.
- 4 drams of Castile Soap.
- 2 drams of Oil of Juniper.
- 2 drams of Camphor.
- 4 drams of Saltpetre.

Bray all well together into a paste, in a mortar, and form it into a ball. The above is only for one dose, but you may make as many as you think proper at once, and keep them for use. Give them at night, and they will work off the next

by lying wet.
 when hot at the
 and be guarded

surfeits, their
 will allow that
 now is this to
 this must be
 part of blood,

into a ball.
 Repeat both
 the last dose
 al balls in a
 lowing pow-

a the above
 t, and if any
 m with the

ces affected
 sometimes a
 , and then
 e case, two
 e. Staling
 ty of warm
 g balls as

and form
 t you may
 eep them
 r the next

day. These balls are of great service in many of the disorders of horses, and are some of the best staling balls yet found out.

By following the foregoing directions, a cure will generally be performed. Sometimes in this disorder little knots break out, especially upon the hind parts of the horse, and these knots throw out a little matter. When this is the case, you must rub them with strong mercurial ointment. Sometimes these little tubes, or pustules, have living insects in them: but by rubbing them as I have just directed you, they will be destroyed and the cure completed.

The Hidebound.

This disorder is too often brought on by the horse being worked too hard, and badly kept; although this is not always the case. When the skin of a horse sticks so close to his ribs that it appears immovable, the horse is said to be hidebound. But this is not properly a disease, but rather a symptom, being often caused by previous disorders, such as fevers, convulsions, surfeits, worms, or disorders of the kidneys or lungs.

CURE.—As the hidebound may proceed from various causes, it is necessary to determine the cause, before such medicines can be applied as will remove it. If it owe its origin to hard labor and want of food, rest and plenty will soon remove it. If it be caused by worms, worm medicines must be applied; or if it be left by any imperfectly-cured disorder, the following drink must be given:

- 2 ounces of Aniseeds, in powder.
- 2 ounces of Ginger, in powder.
- 1 ounce of Grains of Paradise.
- 2 ounces of Mustard.
- 2 ounces of Turmeric.

All to be powdered, and to be given in warm ale, fasting, and to fast two hours after. Bleeding, tapping and physic are also necessary, when the hidebound is left by any disorder.

The Mange.

This disorder is more shameful than dangerous, for you can not go abroad with a scabbed horse without being hissed at, neither is it proper; for this disease is so infectious that every horse that may come near it will be in danger.

The Mange is too well known to need a long description, though some have been mistaken, and have taken a hot itching eruption for it.

SYMPTOMS.—At first it is confined to the skin, but by long continuance it vitiates and pollutes the blood. The skin is generally thick, and full of wrinkles, especially about the mane, the loins, and the tail; and the little hair remaining on those parts stands erect. The ears and eyebrows are commonly naked; and when the limbs are affected, they have the same appearance: but at the same time the horse is not raw, nor does the skin peel off as in a Surfeit.

CAUSES.—The Mange is generally taken by infection, for it is so very catching, that if a horse be put into a stable where one in the Mange has stood, before it be thoroughly cleansed, he will hardly fail being infected. But though infection is the general, it is not the only cause of the Mange. Low feeding, and running long abroad in cold, pinching weather, without sufficient provender, will cause a horse to have the Mange.

CURE.—When the horse has been infected by another, the disorder is not so obstinate as when caused by starvation, for the blood will not be in so bad a state. When you think a horse has got the Mange, apply the following where you think it needful, and it will cure it at the beginning without much trouble or expense:

- 4 ounces of Sulphur Vivum.
- 4 ounces of White Copperas.
- 4 ounces of White Hellebore Root, in powder.

Mix all together in two quarts of butter-milk, and rub the places well. By this method you can cure most scabs of short duration; but when once the Mange has got great hold, it will require sharper treatment. Make the following for one horse:

- 4 ounces of Sulphur Vivum.
- 2 ounces of White Hellebore Root, in powder.
- 2 ounces of Blue Stone Vitriol, in powder.
- $\frac{1}{2}$ ounce of Verdigris, in powder.
- 4 ounces of Flanders Oil of Bays.
- 3 gills of Whale Oil.

Mix all well together, and rub the horse well with it all over—in the sun, if in summer, but before a fire if in the winter. In summer you must also turn him out to grass after rubbing, but in winter keep him warm in the house. You must be careful to wash your saddles and bridles, cart gears, stands, mangers, racks, &c., well with quick-lime and chamber-ley; for if you do not clean all that the horse may have used, the infection will remain. You will find the above a certain cure, if managed rightly, for I have cured hundreds with it, and I do not remember one instance of its failure.

At the same time give freely of flour of sulphur and liver of antimony ; and if you have a number of horses infected, be sure to rub them all together.

Some people say that when a horse is rubbed for the scab he will infect others, but I am of opinion that he will not, neither do I remember an instance of it.

The Farcy, or Farcin.

There have been many opinions respecting this disease. Some authors reckon five kinds ; but although there are so many different branches, yet four of them have the same root. The Water Farcy is different from the others, and therefore I shall put it afterwards by itself. There is a scurvy which horses are subject to, and which is often called a Farcy ; but it is no such thing, for there are only the two kinds of Farcy, which I shall here treat upon. Horses are often said to have the Farcy when they have not, for sometimes when people do not know the proper name of a disorder, they call it the Farcy. The true Farcy is a disorder of the blood-vessels, and generally follows the course of the veins, and when inveterate, thickens their coats and integuments in such a manner that they become like so many cords.

SYMPTOMS.—At the beginning of this disorder a few small knobs or tumors, resembling grapes, are found on the veins, which are so painful to the touch that the creature shows evident marks of uneasiness on their being pressed with the finger. They are at first very hard like unripe grapes, but in a very little time they grow soft, and break and discharge a bloody matter, and become very foul and untoward ulcers. This disease appears in different places on different horses. Some show it first on the head ; some on the external jugular vein ; some on the plate vein, extending from thence downward, on the inside of the fore-leg, towards the knee, or upwards towards the brisket. In some it first appears about the pasterns, on the sides of the large veins, and on the insides of the thighs, extending towards the groin ; in others on the flanks, spreading by degrees towards the lower belly ; and some horses are nearly covered all over the body at once.

CURE.—When the Farcy attacks only one part of a horse, and that where the blood-vessels are small, it may be easily cured ; but when the plate vein is affected and turns corded, and especially when the crural veins, withinside the thigh, are in that condition, the cure is very difficult, and the creature is rarely fit for any thing but the lowest work after it.

Therefore those who depend upon some particular medicine, and flatter themselves with being able to cure every species of the Farcy with it, will find themselves wretchedly mistaken ; for different medicines are needful, according as the disease is superficial or inveterate. The former is easily cured, for sometimes moderate exercise is sufficient ; but the latter requires knowledge and experience ; and often baffles the most skillful, and defies the whole power of medicine.

From the above description of this disease it appears that it is of the inflammatory kind, and that the blood-vessels are affected. Copious bleedings are therefore absolutely necessary, especially if the horse be fat and full of blood. This evacuation always checks the progress of the Farcy in its beginning, but its good effects soon vanish, especially if the horse be low in flesh. After bleeding, mix the following :

4 ounces of Cream of Tartar.
 4 ounces of Liver of Antimony.
 4 ounces of Lenitive Electuary.
 4 ounces of Castile Soap.
 2 drams of Calomel, 8 drams to an ounce.

Make these into balls, and give two ounces a day for some time. While giving these balls, dissolve a little nitre in the water given to the horse to drink. These medicines will keep his body open, and allay the inflammatory heat of his blood, which is the principal cause of the disease ; and while they are given inwardly to remove the cause, let the tumors be rubbed twice a day with the following ointment :

4 ounces of Elder Ointment.
 4 ounces of Flanders Oil of Bays.
 2 ounces of White Vitriol.
 1 ounce of Red Precipitate.
 2 ounces of Sugar of Lead.

Beat all well together into an ointment, and keep it for use. This ointment will soon disperse the tumors, which will leave small bald spots on the skin, but the hair will grow again in time. If the tumors break, and run a thick, well digested matter, it is a sign that the disease is conquered, and the horse will soon be well ; but it will be necessary to give him two ounces of liver of antimony every day for a fortnight, and two ounces every other day for a fortnight after, in order to sweeten his blood, and disperse the small bunches that remain. This method never fails when the small veins only are affected ; and a short time will complete the cure.

But when the Farcy affects the large blood-vessels, the cure is far more difficult. Let the practitioner always attempt it at the beginning of the disease, as he then will have fewer

difficulties to encounter; for delay renders that almost impossible to be overcome which at first might have been easily conquered. Therefore, when the plate or crural veins are corded, lose no time, but bleed immediately on the opposite side, and apply to the distempered vein the following mixture, which is proper to dress the wounds with, but not before they are broken out:

1 dram of Corrosive Sublimate.
1 ounce of Spirits of Salt.

Powder the sublimate, and put it into a bottle, and put the spirits of salt upon it to dissolve it; then add two ounces of vinegar, by degrees. This is a very proper mixture to dress the ulcers with; but if it can not be easily got, take

6 ounces of Oil of Turpentine.
3 ounces of Oil of Vitriol.

Put the pot in water with the turpentine in it, and pour the oil of vitriol in, a little at a time, and keep stirring it till it shall have subsided. If the Farcy be situated in the loose and fleshy parts, such as the flanks and belly, the mixture should consist of equal parts of oil of turpentine and oil of vitriol; but when the seat of the disease is in the parts which are less fleshy, the proportions above are best calculated to perform a cure. The medicine must be used in the following manner. Rub the parts affected with a woollen cloth, and then apply some of the compound oil to every bad and tumor. Continue this method twice a day, and at the same time give cooling physic every other day. The balls and nitrous draughts before mentioned will answer the intention. By this treatment the tumors will be digested, and the cords dissolved; but it will be necessary to give liver of antimony to complete the cure; and to prevent a relapse; and also to dress the sores, when well digested, with a mixture of bees' wax and oil, which will heal them, and smooth the skin.

Sometimes the disease will not yield to this treatment, especially when situated near the flanks or the lower belly. In that case it will be necessary to bathe the parts with the compound oil, as far as the centre of the belly; and at the same time to give a course of antimonial medicines. The following composition is stronger than the last, and on that account is often used when the disease is obstinate:

4 ounces of Spirits of Wine.
2 ounces of Oil of Turpentine.
4 ounces of Oil of Vitriol.
2 ounces of Vinegar.

Mix all together, with the caution before directed. When

this method fails, and the disorder becomes inveterate, try the following, which is recommended by an eminent practitioner :

- ½ pint of Linseed Oil.
- 3 ounces of Oil of Turpentine.
- 3 ounces of Oil of Peter.
- 2 ounces of Oil of Bays.
- ½ ounce of Oil of Origanum.
- ½ ounce of Strong Aquafortis.
- 2 ounces of Barbadoes Tar.
- 2 drams of Tincture of Euphorbium.

Mix all together with caution, as before directed. This medicine must be rubbed on the tumors and corded veins once in two or three days, observing that if the mouths of the ulcers are choked up, or so thick as to confine the matter, to open a passage with a small hot iron ; and also to destroy the proud flesh, which may be kept down by touching it occasionally with oil of vitriol, aquafortis or butter of antimony.

In this disorder, these are the best ways of proceeding that have yet been discovered ; but it is to be considered as an obstinate one, and is sometimes very bad to cure. It has hitherto baffled many an able practitioner, and it is to be feared will baffle many more ; for when the blood has got into such a corrupted state, it bids defiance to medicines.

The ingenious Dr. Brackden recommends the strong mercurial ointment for rubbing the cords and tumors with before they break ; and in order to disperse them when they are broken, to dress the sores with a mixture composed of equal parts of Venice turpentine and quicksilver. If the mouth becomes sore by this means, a gentle purge should be given to prevent salivation. This is doubtless a very good method, and if care be taken, will often prove effectual. He also recommends the following alterative ball :

- 1 ounce of Liver of Antimony.
- 1 ounce of Bezoar Mineral.
- 4 ounces of Cordial Ball.

Beat all well together, and give about the size of a walnut every day for some time, fasting, and to fast two or three hours after.

I have given you the best prescriptions that I am able to give, and such as will not fail to cure if properly applied, if the horse be not incurable. I have been more particular in treating of this disease, because it is common among horses, and very often managed improperly by those who pretend to cure it. Such, therefore, as have valuable horses in this disease, would do well to be careful whom they em-

ploy, and assiduous in observing the methods they make use of to cure it, if they do not think proper to attempt a cure themselves. In my opinion they may do it better themselves by following these directions, than by trusting to farriers.

The Water Farcy.

This disease varies very much from the last, and would more properly be called a Dropsy than a Farcy. There are two kinds of this disorder, but they are nearly of the same nature. One of them is produced by indisposition terminating in the skin, as is often the case in epidemical colds; the other is a true dropsy, where the water is not confined to the belly and limbs, but is found in different parts of the body, and a great number of soft swellings appear. When you press the finger pretty hard upon any of the swelled parts, or under the horse's belly, or on any part of the body, you may take it for granted that he has got the Dropsy, or what is called the Water Farcy.

This disorder mostly proceeds from foul feeding, or a continuance of very wet weather in the end of summer.

CURE.—Whenever the swellings appear, make scarifications, that is, holes through the skin. This may be done by a short fleam; and if you have not one short enough, put a collar made of a piece of leather on it. If the swelling be under the belly, strike a good many holes in at a time, but be careful to avoid the veins. By this means a great quantity of water will run out. Taps in the brisket are also often of great service. I have fleamed horses four or five times before I could get the swelling to subside. When the water has subsided, the blood is left in a bad state, to remedy which, a gentle purge should be given two or three times, eight days distance, to recover the crisis of the blood, and brace up the relaxed fibres of the whole body. Lime water is very proper, with a little nitre in it, and let the horse's food be warm mashies of bran, with a little malt in it. His keep must be increased by degrees. The disorder mostly happens to young horses that have not been used to high keep. You must give one of the following balls every day, omitting the time when the physic is working:

2 ounces of Squills.
1 ounce of Camphor.
1 ounce of Castile Soap.
 $\frac{1}{2}$ ounce of Turpentine.
4 ounces of Yellow Resin.

Make these into a ball with honey, and give one ounce at a time. These balls will work the water off by urine. When

the horse has been treated in this manner till the water is evacuated, and he begins to recover, give him a pint of the following infusion every day for a fortnight, fasting, and let him fast one hour after each dose :

4 ounces of Gentian Root.
4 ounces of Black Hellebore.
2 ounces of Jesuit's Bark.
4 ounces of Camomile Flowers.
1 Handful of Centaury.

Boil all together in six quarts of water, for ten minutes, let it stand till cool, and then strain it through a cloth. This strengthening drink will brace the fibres, cause the fluids to circulate quicker, and complete the cure.

Founders.

The Chest Founder is produced by violent exercise on a full stomach, and drinking large quantities of cold branch water ; by the use of mouldy bran, corn, or oats, or by eating large quantities of green food, such as oats, wheat, peas, &c., while performing hard labor. The seat of the disease is in the lungs ; the heart and liver are also considerably enlarged, insomuch that there is not room for them to perform their office with ease. The liver, lungs, diaphragm, and surrounding parts, are all covered with large brown spots, and are much inflamed.

SYMPTOMS.—A horse that is chest foundered, will straddle or stake with his fore legs, showing an unwillingness to bring his feet together ; and if they are placed near each other, he will not permit them to remain so for a minute. Indeed, they are frequently twelve or eighteen inches apart, which is caused by a fullness and continual uneasiness about the chest : the cavity being too small to contain the lungs, &c., in their enlarged situation, with ease. The hind legs are free from the pained appearance of those before, and it is not difficult to distinguish it from a common founder, as it is wanting in all its symptoms, except the stiff and numbed appearance of the legs.

Large bleedings, and half an ounce of aloes given internally in a ball, have sometimes afforded momentary relief. A horse laboring under this disease, is worth but little more than his board, as he is unable to bear fatigue, or undergo severe service.

The Common Founder is caused by riding the horse until much heated and fatigued, and then allowing him to cool suddenly, by drinking freely of cold water, or standing in an exposed situation, or in a cold stable without covering ; or,

without sudden cooling, it may be produced by too bountiful feeding, and his swallowing his food greedily, while very warm and hungry. Under such unfavorable circumstances, the poor animal, after resting, instead of being refreshed, is stiff and sore; his rest, food, and drink, being more destructive to health than constant action and abstinence. On the contrary, had he been allowed to cool gradually, and fed sparingly, he would have escaped injury.

CURE.—Take from the neck vein a gallon of blood, and give as a drink, in a quart of strong sassafras tea, a table-spoonful of saltpetre, and a quarter of an ounce of assa-fœtida; withholding any drink for six hours; at the end of which, should he not be better, take half the quantity of blood, repeat the drink, offering him bran or oats scalded with sassafras tea, his drink being mixed with tea. His feet should be well cleaned, and filled with cow dung.

OTHER REMEDIES.—As soon as your horse is foundered, bleed him in the neck in proportion to the severity of the case. In extreme cases, bleed him as long as he can stand. Then draw his head up, and, with a spoon, lay salt back on his tongue, till he has swallowed one pint. Let him drink only moderately. Anoint his feet with spirits of turpentine, and he will soon be well. The founder pervades every part of the system. Bleeding arrests it in the blood, and the salt in the stomach and bowels. Attend to it immediately after the injury.

The Prairie Farmer recommends, as a simple and certain cure, to bleed the horse freely in the neck, as soon as the founder is discovered, and as soon as practicable, place him in water about up to his belly—the colder the better—and let him stand two-thirds of a day; or, if badly foundered, longer. This drives the founder from his feet and legs, and prevents its settling there.

Poll Evil.

This disease in the horse proceeds sometimes from striking the poll of his head against any hard substance, or from bruising the part with the halter, which causes it to swell. First abate the inflammation by bleeding, physic, and the application of cold lotions to the part. This will sometimes disperse the swelling. If it matters, hasten its formation with warm fomentations, poultices, or stimulating embrocations; then open the swelling so that *the whole of the matter should run out*, and continue to do so. This is done by a seton. Keep it clean with warm water. A piece of the skin

of old bacon rubbed on with a hot iron, is a good application. Poll Evil has been cured by the following mixture (apparently at least by it in some instances.) Take finely pulverized flint glass three spoonsful, put into urine, one pint in a bottle, and bury it in the ground for three or four days; after which take one spoonful or more of the mixture, well shaken up, and put that much into each ear, once a day for three, four or five days.

A Sprain in the Back.

All kinds of Sprains resemble each other. They are a relaxation of the tendinous fibres, from the muscular parts being overstretched. Sprains in the back are mostly caused by overweighing the horse, or by his losing his hind feet on the side of a hill; and sometimes by putting him back too quickly. There is much difference between a Sprain in the back, and what is called tified in the back. Sometimes a horse catches cold in his loins by having his clothes or his saddle taken off when he is hot, and being turned out of doors; and some horses are subject to Rheumatics, which make them lame in many places, and are generally taken for Sprains.

To cure a sprain in the back, first bleed pretty freely, and then give the following drink:

1 ounce of Tincture of Gualacum.
1 ounce of Balsam Capivi.
1 ounce of Oil of Juniper.

To be put into a quart of strong parsley-root tea: half to be given at night, half in the morning, and plenty of warm water to work them off.

Also, lay a sheep's skin with the flesh side to his loins for six or eight hours, if he can bear it; and then turn the wool side to him, and let it stay on a day; and when you take it off, sheet him well for fear of catching cold. Also, make a plaster of the following:

2 ounces of Oxyeroseum.
2 ounces of Paracellis.
2 ounces of Red Dominion.
2 ounces of Burgundy Pitch.

Melt all together, and lay them on pretty warm, and put a little wool, clipped short, on the charge while hot, and then pour on a little cold water to fasten the whole together. These are the best methods I am acquainted with, and if properly pursued, and the mixtures carefully made, there will be prompt relief.

A Sprain in the Shoulder.

When the shoulder of a horse is sprained, he does not put out the leg like the other, but in order to ease it sets the sound foot firmly on the ground. When trotted in hand he forms a kind of circle with his lame leg, instead of putting it forward, and when he stands in the stable that leg is advanced before the other.

There is what is called a Shoulder-slip, which is worse than a Sprain. When this happens, the horse can neither lift his leg nor put it forward. You may know this by the shoulder-blade standing higher than the other; but to discern that, you must make him stand on the lame leg, for the leg he stands on will always appear a little higher than the other. The flesh will also shortly waste away from the shoulder-blade, which is a sure sign of a Shoulder-slip. When this accident has taken place, put a tap in the lame side of the animal's breast, and blow the shoulder full of wind with a pipe. When you have rinded the skin to put the tap in, hold the skin fast to the pipe, and blow the part that you have rinded full, and let some one draw the wind up into the shoulder with the edge of his hand as far as the top of the shoulder-blade, and then put in the tap, or rowel, and stop the hole up well with tow and salve. Give warm water for three days, and then open the place, stir the tap round, and rub the shoulder all over with the following liniment:

2 ounces of Spirits of Wine.
2 ounces of Sweet Oil.
2 ounces of Spirits of Sal Ammoniac.

Shake them well together, rub the shoulder well with the mixture every third day for some time, and if the horse do not get better with three or four times rubbing, use the following:

2 ounces of Oil of Turpentine.
1 ounce of Oil of Origanum.
1 ounce of Oil of Bricks.

Shake these up together, rub all on at a time, and walk the horse about a little afterwards. When near the sea, swimming in the salt water is very proper, and I have known swimming in fresh water to be of great use, particularly in clear running streams.

A sprain in the shoulder point requires nearly the same treatment as a slip, but you need not blow it. When it is attended by inflammation, cooling mixtures, such as extract of lead and water, must be used. But when a swelling or an inflammation takes place, it is mostly caused by a hurt,

or by a stroke from another horse. If there be no swelling, rub the shoulder point well with the following mixture every third day :

1 ounce of Oil of Peter.
1 ounce of Oil of Amber.
1 ounce of Oil of Spike.
1 ounce of Oil of Bricks.

Shake these well together, and rub the shoulder point every other day. If the horse be not better, take

1 ounce of Oil of Turpentine.
1 ounce of Oil of Origanum.
1 ounce of Oil of Swallows.
1 ounce of Oil of Amber.

Shake these together, and rub the shoulder point well with them every third day ; and if the horse continue lame, recourse must be had to blistering.

A Sprain in the Coffin Joint.

This is often a grievous disease, and it is difficult to discover where the lameness is. It is often neglected till the joint grows stiff, and then the horse pitches upon his toe, and is afraid of bearing any weight upon his foot. If you press with your thumb in the hole in the horse's heel, and upon the cornet of his foot, you will soon discover whether the hurt is in the Coffin Joint.

When people can not tell the cause of a horse's lameness, they often say that he has got sprained in the coffin. In my opinion it is better not to doctor a horse than to apply stuff to you know not what. If people would have a little patience, most lamenesses would soon show themselves, especially a Sprain in the Coffin Joint, for it would raise a ring round the corner of the foot, not much unlike a Ring-bone, but closer to the foot.

The first thing to be done is to draw a little blood from the spurn vein, then mix an equal quantity of oil of bays, or oil of origanum, beat well together, and rub well all round, just above the hoof. Apply this for three or four days together, and if no better, you must have recourse to repeated blistering.

A Sprain in the Back Sinews.

This kind of Sprain is more frequent among horses than any other, and is so common that I need not describe it, but only inform you how to cure it.

If it be recent, bathe the leg with a little hot vinegar, or verjuice, with a little saltpetre dissolved in it, and put round

It a proper bandage ; or, curriers' shavings, wetted with a composition made of vinegar, spirits of wine, and a little tar, and laid on the swelling with a pretty tight bandage round them, will be of great use. Take it off once a day, and soak the shavings again, or get fresh. Injuries of this kind must not be expected to be removed immediately. Rest is absolutely necessary, and turning the horse out to grass would be of great service as soon as the swelling disappears, but not before. If these methods fail, the next thing is to blister ; for I have known blistering succeed when all the former have failed. The last thing to have recourse to, is firing ; and this must be done carefully, and by some experienced person.

Sprains of the Knees and Pasterns.

The knees are liable to many misfortunes besides sprains. The Speedy Cut is done by striking one foot against the other leg, just below the knee, and is frequently done by a horse that trots high. Sometimes it swells very much, and is taken for a sprain. Sometimes horses get kicked by others, or meet with some other accident which causes a swelled knee, which is sometimes bad to remove. A poultice, made as follows, and carefully applied to the part injured, will have a great tendency to remove the swelling :

Take 4 ounces of Tar.
4 ounces of Spirits of Wine.
3 ounces of Hogs' Lard.

Melt these together over a slow fire, and be careful not to set fire to them, and put in as much linseed-meal as will make them of a proper consistence. This is a very good poultice for many other kinds of swellings, and although but little known hitherto, I hope that it will be found of great service. If any substance be left which will not give way to this method, you must lay on a little blistering ointment.

Lameness in the Stifle.

The Stifle is the tenderest part of a horse, except the eye. How many horses have lost their lives by misfortunes in this part, and how many have been left lame by not being properly cured ! A horse that is lame in the stifle generally treads on his toe, and can not set his heel to the ground without great pain and difficulty. When you find this is the case, bathe the part well with warm vinegar ; and if a puny swelling appear, foment it well with a woollen cloth wrung out of hot vinegar, wormwood and rosemary, having

added half a pint of spirits of wine to a quart of the decoction. Let this operation be continued till the swelling is nearly gone.

Owners of horses should watch closely for this lameness. They cannot be too careful of the feet of their animals. You should examine the feet of your horses frequently, and with much minuteness.

When a horse has got a stroke and cut by the heel of his shoe, great care must be taken to keep out the cold air, and to keep him from drinking cold water, especially in winter time; for if the horse take cold, and an inflammation come on, there is reason to fear that death will follow. When you perceive that a horse has got a stroke and is cut through the skin, bathe it well with the following mixture :

2 ounces of Spirits of Wine.
1 ounce of Spirits of Sal Ammoniac.
1 ounce of Oil of Amber.

Shake all well together, bathe the place well with it, and lay on a diachylon plaster with gum, in order to keep out the cold. If the horse grow very lame, and the swelling much, foment with the following :

Take a handful each of Wormwood,
Elder Leaves, Camomile Flowers,
Juniper Berries, Marshmallow Leaves.
Boil in 2 gallons of Chamber-ley.

If the above can not all be got, take double the quantity of juniper berries and camomile flowers, and foment for some time, as before directed; and when you leave off fomenting, rub with the above mixture. Be careful to wrap the part up warm, and as soon as it comes to matter, dress it with basilicon ointment. If any brown lee appear, syringe tincture of Benjamin, or balm drops, into the wound. When a callous substance, or proud flesh appears, eat it off with red precipitate. These are the best ways of curing a horse that has got lamed in the stifle.

Lameness in the Cup-Bone, or Whirlebone.

A lameness in this part of the hip is discovered by the horse's dragging his leg after him, and dropping backwards on his heel when he trots. If the muscles of the hips only are injured, the lameness may be soon cured; but if the ligaments of the cup are affected, or relaxed, the cure is often very tedious; and when the cup is full of glueish liquor, the cure is doubtful. I have known the ligament which holds the hip bone in the cup to be broken, and then the hip bone

has come loose ; but this seldom happens, and when it does, nothing can be done to relieve it. When a horse is lame in the cup-bone, rub him well with the following mixture :

- 1 ounce of Oil of Amber.
- 1 ounce of Oil of Bricks.
- $\frac{1}{2}$ ounce of Oil of Origanum.
- 1 ounce of Oil of Turpentine.

Shake these well up, rub one half on the cup, and the other half the third day. I have known this mixture remove the complaint when of long standing ; but if it fail, you must try what blistering will do. The last thing to be tried is firing. I have fired horses that have been lame above a year, and cured them by it.

It is very easy to fire a horse in this part. Take the iron and make a circle, eight inches in diameter, like a cart-wheel, with scores about an inch asunder. Cut nearly through the skin ; and if you do cut through in some places, it will be no worse. Then lay on the following charge :

- 2 ounces of Oxycrosum.
- 2 ounces of Paracellia.
- 2 ounces of Red Dominion.
- 2 ounces of Burgundy Pitch.

Melt these all together, lay them on pretty hot, lay a little wool, clipped short, on before it cools, and then a little water on to fasten all together.

Sprains in the Fetlock, or anywhere in the limbs, may be treated in the same manner, and if the oils will not remove the substances, blisters will ; therefore I shall give you a recipe for a blister which is not only suitable for those complaints, but for others where a blister is wanted.

- Take $\frac{1}{2}$ ounce of Quicksilver.
- $\frac{1}{2}$ ounce of Venice Turpentine.
 - 2 ounces of Flanders Oil of Bays.
 - $\frac{1}{2}$ ounce of Gum Euphorbium.
 - 1 ounce of Spanish Flies.
 - 2 drams of Oil of Origanum.

Rub the Venice turpentine and quicksilver well in a mortar for some time, and then put in the oil of bays and rub for some time longer, till the quicksilver is quite killed. Powder the gum and flies well, and mix them all together. When you lay the blister on, rub it well against the hair to get it to the bottom ; lay it on with a broad pointed knife, and cover it with a cloth, or the horse will get his mouth to it. This blister will answer any purpose you want a blister for, but you must observe that there must be fifteen days between the times of laying it on.

The Bone-Spavin.

Although this is a common disorder among horses, yet it is little understood by either breeders or farriers. The Bone-spavin is a bony excrescence, or hard swelling, on the inside of the hock of a horse's leg, and sometimes owes its origin to kicks and blows, and sometimes to natural causes; but in the former case it is much more easily cured than in the latter; and those that grow spontaneously on colts, or young horses, are not so bad as those that appear in horses that have arrived at their full strength and maturity. In old horses they are generally incurable.

Sometimes the horse is very lame when the Spavin is first coming out, and when it has come out is better for some time, and then grows lamer again as the bone hardens. I would advise you to apply a blister as soon as you have any suspicion that a horse is likely to put out a spavin, and to continue blistering, every fortnight, for some time, by which means you may stop a Spavin in a young horse.

CURE.—Mild medicines should be used if the horse is young, as they will in a short time wear the tumor down by degrees, which is much better than trying to remove it at once by severer methods, which often have a very bad effect, and produce worse consequences than those they were intended to remove. But in full-grown horses other treatment is necessary. The following ointment is probably the best that can be used:

- 2 ounces of Cantharides.
- 4 ounces of Mercurial Ointment.
- 3 ounces of Tincture of Iodine.
- 4 ounces of Turpentine.
- 3 drams of Corrosive Sublimata.
- Mix with 2 pounds of Lard.

Cut off the hair from the part affected, and grease thoroughly with the ointment thus made, rubbing it in well with the naked hand. Let it remain two days, then grease the part with lard. In two days more wash it off with soap and water, and then apply the ointment again. Continue this till a cure is effected. By these means Bone-spavin may often be cured; but when they fail, recourse must be had to firing.

Before you fire a horse for the Bone-spavin, be careful to take the vein out of the way, for it generally lies over the Spavin, and you can not fire deep enough to come at the callos substance without its removal. In order to destroy the vein, cut a nick through the skin upon it, just below the Spavin, and another just above it, and put a crooked needle

under the vein, and tie both ends : then cut the vein across between the tyings, both above and below, and you may either draw the piece of the vein out or leave it in. Let the iron you fire with be pretty sharp ; cut four or five nicks upon the bone, and let the iron take hold of the superfluous bone, in order that it may waste away by mattering ; and when you have done, lay on some white pitch, pretty hot, and put a cloth round it to keep it on. In three days open the place and dress it with yellow basilicon.

The Blood-Spavin, or Bog-Spavin.

Many farriers and horse-dealers divide this disease into two heads, and give them different names ; but to my certain knowledge they are both one, for I have proved it in many ways. A Blood-spavin does not come by breeding from spavined mares, nor by being got by spavined horses, as the Bone-spavin does ; but you may safely breed out of a Blood-spavined mare, or have the foals got by a Blood-spavined horse. In my opinion Blood-spavins are generally brought on either by Sprains, or hard labor when the horse is young, and sometimes when he is full-grown.

The Blood-spavin, or Bog-spavin, is a dilation of the vein that runs along the inside of the hock, and forms a small soft swelling in the hollow part, which in time renders the creature lame, but seldom till the gelatinous matter becomes ropy, like melted glue in a bag, and is situate on the inside of the hough. Sometimes it goes through the back part of the joint, and then it is called a thorough-pin.

CURE.—Soon as you discover the vein puffed up, or forming a bag, lay on some blistering ointment, and in four days after bathe the swelling well with hot vinegar, with a little saltpetre dissolved in it. Also put a bandage round it to disperse the swelling as much as you can. If this method do not succeed, you must make two incisions in the skin lengthwise, as the vein runs, one just above, and the other just below the joint, and lay the vein bare : then put the end of a buck's horn under it, raise it up, and fasten it in both places with waxed thread : then cut the vein in two at both places, within the tyings, and if you think proper, draw the vein out. This method of proceeding will cure most Bog-spavins at the beginning. The Blood-spavin may also be cured by applying the same ointment used in Bone-spavin, only it should be applied once in six days instead of once in four. This spavin ointment is very powerful, and when properly used is the best ever invented.

A Corb.

This is a soft swelling that rises out of the joint on the back part of the hind-leg, just below the hock, and mostly lames the horse, besides being unpleasant to the eye. To cure it, strike a few holes into it with a pricker, made so as just to go through the skin, then rub well with oil of origanum, and blister as often as needful. If this will not answer, use the Spavin ointment once in six days.

A Ring Bone.

This is so well known that I need not describe it, but only point out the remedy; yet I must observe that a Sprain in the Coffin is sometimes taken for a Ring-bone, when it causes a rim to rise just above the foot. Ring-bones come out from the pastern, between the fetlock and the foot; but if the pastern is long, they are nearer the foot. They will generally yield to the same method of cure as a Corb, especially if just coming out, but if not, use the Spavin ointment in same manner as in Bone-spavin, only do not apply it so often. Once in six days is often enough.

Mallenders and Sallenders.

The first is on the fore-leg, at the bend of the knee, and the last on the hind-leg, at the bend of the hough. They crack and throw out a thin brown matter, and sometimes, a hard scurf, or scab, which prevents the horse from bending the limbs which are affected as he should do.

CURE.—They both proceed from the same cause, and consequently require the same treatment; which consists in washing the parts with old chamber-ley, or a warm lather of soap-suds, and afterwards applying strong mercurial ointment, spread on tow, to the cracks, once a day till the scabs fall off, when the cure will be completed; and then it will be necessary to give him a dose or two of physic. If the disorder will not yield to the mercurial ointment, make a strong mixture of vitriol water, and wash the cracks with it, and it will dry them up, and cause the scabs to fall off.

The Strangles.

Most horses have this disorder while young, but at seven years old they are out of danger. There are two kinds of this disorder. The common kind is a swelling under or between the jaw-bones. The other, which is called the bastard kind, is much the worst. Sometimes swellings appear

on the buttocks, break, and discharge matter for a few days, and then dry up, after which others appear in a fresh place in the same manner. I have known horses that have had this complaint eight or ten weeks. The common kind begins with a swelling between the jaw-bones, which sometimes extends to the muscles of the tongue, and is often attended with so much heat, pain and inflammation, that before the matter is formed the creature swallows with the utmost difficulty.

SYMPTOMS.—The Strangles is attended with great heat and fever, a painful cough, and great inclination to drink, without being able. Some horses lose their appetites entirely, and others eat but very little, occasioned by the pain resulting from the motion of the jaws in chewing and swallowing. When the horse runs much at the nose, it is not a good sign. Although this disease is very troublesome, it is not dangerous, except when the swelling turns upwards against the wind-pipe and gullet, and then there is danger of suffocation if it do not break soon.

CURE.—The Strangles is not properly a disease, but a discharge common to young horses, and therefore it follows that the discharge must be promoted in order to throw off the offensive matter. The best method of doing this is to keep the swelling always soft by soaking it with softening ointment, such as marshmallows, or elder ointment. I have known oil of swallows, with a little spirit of hartshorn in it, to be very useful in bringing the swelling forward and causing it to break. A cloth in the form of a cap, put on the horse's head, and stuffed with wool to keep the swelling warm, will be of great service. Some people apply a poultice, but there is no need of this if the above be properly used. Give plenty of warm water, with a little meal on it; for in this disorder a horse can not swallow dry food enough for its support. When they gather and break, see that the matter runs freely; if not, open them with the point of a sharp knife. It will be five or six days before they break.

When the swelling is broken, and the orifice of a proper size to discharge the matter, dress with the following ointment spread on tow :

Take 1 pound of Yellow Resin.
 1 pound of Burgundy Pitch.
 $\frac{1}{2}$ pound of Honey.
 $\frac{1}{2}$ pound of Common Turpentine.
 $1\frac{1}{2}$ pounds of Hog's Lard.
 4 ounces of Beeswax.
 1 ounce of Verdigris, finely powdered.

Melt the ingredients together, but do not put the Verdi-

gris in till nearly cold, and keep stirring all the time till cold, or the Verdigris will fall to the bottom. This is one of the best salves for wounds that has been found out, and especially for old ones.

The Bastard Strangles requires the same kind of treatment, but it is proper to give the horse some calomel physic also.

The Glanders.

This disease has baffled all who have tried to cure it, and probably will do so to the end of time; so I advise those who may have a glandered horse, to put him off as soon as they shall be certain that he is so. People often mistake other disorders for the Glanders. A violent cold sometimes causes a running at the nostrils, and kernels under the jaws, when the horse is free from the Glanders. Sometimes a running at the nostrils is caused by laying too much weight on a horse. I do not pretend to cure Glanders.

SYMPTOMS.—The matter discharged from the nostrils of a glandered horse is either white, yellow, greenish, or streaked or tinged with blood. When the disease has been of long standing, and the bones are fouled, the matter turns blackish and becomes very bad. The Glanders is always attended with a swelling of the kernels, or glands under the jaws, but in every other respect the horse is generally healthy and sound, till the disorder has continued some time, and the morbid matter has affected other parts. If a thin, limpid fluid be first discharged, and afterwards a whitish matter—if the gland under the jaw do not continue to swell, and the disorder shall have been recently contracted, a speedy cure may be effected by applying the following:

1 ounce of Reach Alum.
1 ounce of White Vitriol.

Powder these well, put them into a pint of warm vinegar, and syringe about an ounce up his nostrils every day. This may do good if the disorder be newly caught.

Swellings and Imposthumes.

When a swelling is in its first stage, bathe it well with verjuice or vinegar, with a little saltpetre dissolved in it; and if the swell still continue, mix

1 ounce of Extract of Lead.
1 ounce of Spirits of Wine.
1 ounce of Spirits of Sal Ammoniac.
3 ounces of Vinegar.

Rub the swelling well with it; but if it be very hot, add

four ounces of water instead of the vinegar. Should the swelling come forward and form matter, which you may feel by the pressure of your finger, let the matter out, but be sure to make the incision large enough that you may dress it with ease. When you have laid the part open, dress it with the green salve before recommended, on tow, for you can not have a more proper salve. The next time you dress it, that is the day after, make a wash of the following :

2 ounces of Spirits of Wine.
2 ounces of Spirits of Rosemary.
1 ounce of Roach Alum.
2 ounces of Water.

Mix these all together, and they will answer the purpose extremely well. When you have washed the wound with the above mixture, lay on a little green salve, on tow, and bind it on if you can, but if you can not, lay on a plaster to keep it on.

Some swellings, such as have been caused by bad barfen on the shoulders, or blows on the legs, will not submit to weak mixtures, nor come to matter in a reasonable time. Mix the following, and it will either take them off or bring them to matter :

2 ounces of Oil of Spike.
1 ounce of Oil of Amber.
 $\frac{1}{2}$ ounce of Oil of Origanum.
1 ounce of Oil of Turpentine.

Mix these well together, and rub the swelling well with them every other day. I have known this remove obstinate swellings.

Wounds.

Wounds are caused by accidents of various kinds. When the skin is much torn from the flesh, if you are at hand while the wound is quite fresh, take a square-pointed needle and a waxed thread, and sew it up. Mind to put the needle in straight, one side over against the other; draw the skin tight, and tie a knot; cut off the thread, and then take another stitch about an inch off, and so proceed. When some people sew up a wound they do it the same as they would sew up a piece of cloth, but that is quite wrong, for they should tie a knot at every stitch, and cut the thread off. But when you do not see the wound until the place is growing dead, and the skin is drawing up, then take off the loose skin; for if you keep it on it will curl up, and leave a blemish. Always keep the lips of a wound down.

When a wound is upon or near a joint, there is danger of

throwing out a joint-lee, of which there are three kinds. One, and mostly the first, is thin and brown, something like sweet wort; the second is rather thicker and tougher, something like melted glue; and the third, which is the worst, is like muddy water and snort mixed together. This last has deceived many people; for when the wound has thrown out this kind of lee, with little white slippery pieces, something like matter, it has often been taken for such. When you find any of these kinds of lee, get a bottle of Riga Balsam, and syringe the wound every day. If Riga Balsam can not be got, use Tincture of Benjamin. I have known fomentations be of great use, especially on the stifle joint. When a wound is of a dead color, and the lips rise, and the dirty lee flows profusely, the cure is to be despaired of. If proud flesh arise when a wound is in a fair way for healing, take

1 ounce of Basilicon.

2 drams of Red Precipitate.

Mix them well together, and lay them on the proud flesh. This ointment is also very proper to dress a wound with that appears dead, and does not discharge a proper matter. If the proud flesh do not submit to the above, lay on a little blue vitriol, in powder, or touch it with caustic or oil of vitriol; and should none of these make it give way, lay on a little corrosive sublimate. The salves, tinctures and mixtures already laid down, are sufficient to cure any curable wound.

Hurts in the Feet.

Horses are oftener hurt in the Feet than any where else, and those hurts are often received from the blacksmith. Every blacksmith should be a farrier, and every farrier a blacksmith, for these businesses should go hand in hand. Sometimes a horse is lamed by being shod into the quick. The foot should always be carefully examined on the first appearance of lameness. Sometimes a nail with a flaw in it will cause a great grievance, as one part will come out and the other will go into the foot. Blacksmiths should never use such nails except in a frost, when they can put them into the old holes. Sometimes they leave stumps or pieces of nails in the feet; and sometimes when they drive a nail they turn the point into the foot, and then draw it back and put in another, taking no further thought about it, and when the horse becomes lame they say he is gravelled.

When a horse has got lamed in the foot, be careful to cut it well out, and to damage the hoof as little as possible, and

dress the place with oil of turpentine, spirits of tar, and common tar. Lay no hot drying drugs on, unless proud flesh rise.

Sometimes corns on the heels cause a horse to be lame; cut them out, and dress the place with aquafortis. There is another fault which blacksmiths are guilty of, and which is a great hurt to a horse, that is, cutting or paring the heel down too much. The heel being best to come at, they clap the paring-knife there and cut them down, when there is no need to take any off them at all. The heels are the greatest support of the horse, and by paring them too thin, both corns and lameness in the back sinews are produced.

Before a blacksmith begins to shoe a valuable horse, especially a road-horse, he should examine how he stands and how he goes. If he go low, heavy shoes should be used; but if he go high, light ones. If he turn his toes out, he will cut with the heel, and if he turn them in, will cut with the spurn. If he have a thin, flat foot, he should have broad shoes; but if a hollow, dished foot, narrow ones. If the crust be thin, and the vein near, small nails should be used. Some horses are hoof-bound—that is, have straight heels which pinch the vein between the hoof and the coffin. When this is the case, thin the soles of the feet till the blood springs through, and then put on screw-shoes, and screw the feet out. Let the screw-shoes stay on a fortnight. You may screw the feet out more than half an inch.

The next thing to be considered is gravel rising from the bottom to the top of the foot, and breaking out at the corner. How many twitter-bones have been thus formed that might have been prevented! When the gravel comes up to the top of the foot, take away the sole at the bottom, and the hoof at the top, and mix equal parts of oil of turpentine and oil of origanum, and bathe the top part of the foot. This will prevent a twitter from forming.

A Twitter-Bone.

A Twitter-bone makes a horse very lame, and not fit for work. It keeps throwing a tough white matter out of what is called a pipe. In order to cure it, first find out where the pipe leads. This you may do with a piece of round lead, the size of a small quill-barrel. Have the following ready, viz.:

$\frac{1}{2}$ an ounce of Sublimate, in fine powder.

1 ounce of Spirits of Salt.

Add juice of one lemon when Sublimate is dissolved.

Mix all together, and syringe a squirt full into the pipe when you take the lead out; and when you draw out the

point of the syringe, put your thumb upon the place to prevent it coming back again, and then put on a pledget of green salve and tow. Do this every other day, for three or four times, and by this method most twitters may be removed. But if this method fail, the next thing to be done is to put a hot iron, the thickness of a small finger, where the pipe leads to, and to fill the hole with sublimate, and bind on it a pledget of green salve with a cloth. Let it be five days, and then lay on some more salve and tow, and in five days more the Twitter-bone and pipe will come out. The wound must be washed with the following mixture, and dressed with green salve and tow every other day, till nearly well :

1 ounce of Bole.
1 ounce of Oil of Origanum.
4 ounces of Oil of Turpentine.

Shake them up well together, and they will both help to heal the wound, and to bring down the substance. If any more Twitter-bones or pipes form, you must take them out in the same manner.

A Fistula, and Bruises of the Withers.

The Withers are much subject to bruises, caused by bad saddles, or such as are too wide in the front ; but, whatever be the cause, it is well known that by neglect they often terminate in a Fistula.

When the Withers are bruised and a swelling appears, lay a poultice of bran and vinegar over the part. People who have a horse which is crushed in the Withers, and pay no regard to it, are worse than savages, and are not fit to have such a useful animal.

CURE.—After a horse has been bruised in the Withers, the skin breaks and matter is discharged, and then the owner supposes that the abscess will subside, but he is often mistaken ; and the pipe which he perceives throwing out matter, is at the same time running forwards in the Withers, and forming a Fistula. When you find the disease proceeding in this manner, lay the pipe open with a sharp penknife all the length, for if you leave any unopened, it will form a Fistula. After you have laid the place open, dress it with the following mixture :

4 ounces of Potashes.
4 ounces of Honey.
2 ounces of White Vitriol.
1 pint of Vinegar.

Boil all together, wash the wound well with the mixture,

and lay on the green salve and tow. If the above be not dry enough, add two ounces more of white vitriol, and two ounces of bole. These will make a salve by themselves: and by these methods you may cure any disease of the Withers arising from external injuries.

But tumors often arise in the Withers from internal causes, such as the crisis of Fevers. When this happens, you must not attempt to stop it, nor use any thing to put it back, for by this means you would drive it more into the shoulder-blades, and make it worse to cure; but, on the contrary, do every thing you can to assist Nature in bringing it forward. You can not do better than lay on poultices twice a day till it breaks, for reason tells us that it is better to do so than to cut it; but when it is broken, open the orifice with a knife, that you may have more freedom in dressing it; but be careful in using the knife, that you do not catch hold of the ligament which turns along the neck to the Withers. Sometimes it runs to the other side of the neck, under the Withers. The cure is the same as before. Lay all the cavities open with the knife, and do not cut across if you can avoid it. Then take the following never-failing mixture to dress with:

4 ounces of Crude Sal Ammoniac.
2 ounces of Bole.
8 ounces of Honey.
2 ounces of Pearl Ashes.
2 ounces of White Vitriol.
1 pint of Vinegar.

Boil all together, and apply it to the wound every day at first, and afterwards every other day till well.

Warbles, Girth-Galls, and Plushes from Saddles.

These grievances are commonly known, and the following mixture will effect a cure, take off the soreness, and leave the skin kind. I may affirm that it is one of the best recipes yet found out for the purpose:

2 ounces of Extract of Lead.
 $\frac{1}{2}$ an ounce of White Copperas.
2 ounces of Spirits of Wine.
4 ounces of Soft Water.
1 ounce of Spirits of Sal Ammoniac.

Mix all together in a bottle, keep shaking it up, and rub the affected places well with it, and put your saddles and barfens on while the places are wet in order to prevent them from infecting other horses.

Sometimes horses have what are called Setfasts on their backs. Rub them with a little mercurial ointment, in order

to raise them and make them come off; and if you can not pull them out after using the ointment, cut them out with a sharp knife, and apply to the place the above mixture, or heal it up as a common wound. Most excrescences on horses are cured in this manner.

Windgalls.

These are mostly on the hind legs, near the fetlock, but I have known them above the fetlock, and on the arm. Windgalls are not only eye-sores, but lame many a horse. Many methods are tried to disperse them without effect. If you put your finger on one side, and your thumb on the other, and press with one of them, you will find the Windgall to go quite through the leg. The reason that they are mostly on the hind leg is, because the horse stands lower behind than before, and throws most weight on the hind legs.

CURE.—On the first appearance of a Windgall, bathe the place well with warm vinegar and spirits of wine, and put a pretty tight bandage round it. If this do not remove it, lay on blistering ointment till the cure shall be completed. But should this method also fail, which it seldom does, you must lay the Windgall open and dress it as a common wound. Before you use the knife, be careful to get the horse's body into a proper cool state by physic.

The Grease.

This disorder is mostly brought on by soft corn, hard usage, want of proper cleaning, or a depraved state of the blood and juices; therefore it is proper to divide it into two heads.

CURE, WHEN THE VESSELS ARE RELAXED.—On first observing the legs of a horse to swell after standing several hours, and to recover their proper dimensions with exercise, be careful to wash them clean with chamber-ley, soap-suds, or vinegar and water, every time he comes in, for this will prevent or remove the disorder. Horses that have round or fleshy legs are more subject to the Grease than those that have flat legs; but a flat-legged horse is more easily sprained. Nitre, sulphur, and liver of antimony, are proper both to prevent the Grease and to refine the blood. Mix equal parts of each, and give a meat-spoonful every day in his food.

CURE FOR THE GREASE FROM INTERNAL CAUSES.—If the horse be full of flesh, the cure must be begun by bleeding, rowels, and repeated purging; after which two ounces of the follow-

ing balls should be given every other day for some time, and they will work by urine the day following :

4 ounces of Yellow Resin.
2 ounces of Salt of Prunel.
1 ounce of Oil of Juniper.
2 ounces of Salt of Tartar.
8 ounces of Castile Soap.
1 ounce of Camphor.

Put these into a mortar with about two ounces of honey, or as much as will make them into balls, and they will carry off the offending humors, and free the blood from its noxious qualities. But at the same time that these internal remedies are taken, outward ones should not be omitted. The legs should be bathed with warm verjuice ; and, if very bad, a poultice of boiled turnips applied. If turnips can not be got, make a poultice of linseed meal and bran, with a little hog's lard in to prevent it from growing hard.

Sometimes horses are neglected when in the Grease till they have what are called grape-legs. These may be cured on their first appearance, when they are in the bud, by laying on caustic, or corrosive sublimate. When the swelling is abated, make the following into a salve to dress the sores with :

1 ounce of Blue Stone Vitriol, in powder.
2 ounces of White Lead, in powder.
4 ounces of Honey.

Mix these well together, and lay them on the sores with tow, to heal them ; but, should they continue foul, and not frame to heal, mix four ounces of green salve, and four ounces of Ægyptiacum ointment well together, and lay it on in the above manner. The mixture will both heal and dry up the sores.

Scratches.

This disease is a forerunner of the Grease, and is a hot oozing matter that breaks out like links, with a nauseous smell. It is very troublesome to a horse, causing his heels to look red and angry, and to be very sore ; and sometimes it is so violent as to render a horse lame, and make him unfit for use.

If you do not bleed and physic as soon as you find a heat in a horse's heels, you may soon have him laid in the Grease. Sometimes in slight touches of this complaint the heels are not hotter than usual. In that case, take a little flour of sulphur and spirits of wine, mix them together into an ointment, and fill the cracks well with it. When the disorder is

too hot to be healed by this ointment, rub with a little *Ægyptiacum* ointment, but be careful not to lay too much on, for it is a great drier.

Rat-tails at first resemble the Grease, and are attended with a hot scorbutic humor at the beginning, and often by neglect destroy the roots of the hair, and then there is no complete cure for them, but they may be relieved by rubbing with equal parts of strong mercurial ointment and Flanders oil of bays.

If a horse have got the Scurvy, or Scabs on his legs, this ointment will mostly take them off; but if not, mix well with hog's lard, one dram of sublimate, in fine powder. These two last mixtures will cure most Scabs, and mixed together they will cure the Itch.

The Crown Scab.

This disorder is similar to the last. It breaks out on the cornet of the foot, and is at first a hot running tumor, and afterwards a dry, scurvy scab. Let the horse be bled and physicked, and then rub him with either of the above ointments, and the cure will be effected.

How to Manage a Horse on a Journey.

In the first place, find out whether the horse is in health, and has been properly fed and exercised; for when a horse is flushed up, and has had no exercise, he is very unfit for a journey. Before you set out be careful to observe if his shoes are fast, and if they sit easy; also whether he cuts before or behind, or interferes, as it is called. If a horse cut with bad shoes, he will probably do it with new ones, notwithstanding what horse-dealers may say to the contrary.

If your horse's back inflame by the rubbing of the saddle, wash the part as soon as you perceive it, with salt and water, or vinegar and water, and have the stuffing of the saddle altered so as to remove the pressure from the part affected: but if the skin be broken before you perceive it, the injured part should be washed with a mixture, composed of equal parts of extract of lead, spirits of wine, and water. You must look well after his back, and walk sometimes, in order to ease him, especially when going down a hill.

It often happens, especially to young horses, that the legs swell, or become gourdy, as farriers call it, with traveling, and more frequently when a horse cuts. It is also observed that the hind legs swell oftener than the fore legs, because the fore parts stand highest in the stall, and consequently

the greatest stress lies on the hind legs. The best method of prevention is to wash the legs with warm water every time you bait, by which means the disease will be prevented, and much time, trouble and expense saved.

Nothing is of more consequence in traveling than to take care that your horse has water at proper times and in proper quantities. When a horse travels he perspires considerably, especially in hot weather, and should therefore be allowed to drink a little more than usual, as opportunity offers, which will refresh him greatly; but never suffer him to drink much at a time, for if you suffer him to drink his fill he will be dull and sluggish afterwards, besides the harm he may receive from drinking too much cold water when he is hot. Never allow your horse to be led to water, or to have his heels washed, after you have arrived at your inn, but let him have lukewarm water, or water moderately cool, when he has stood some time in the stable.

It is a general rule that when any extraneous body, or foreign matter, such as sand or gravel, is lodged in any part of the animal's body, it must be extracted as soon and as easily as possible. When gravelly matter has got into the quick at a nail-hole, or any other aperture, it ought to be removed as soon as possible, but with as little loss of substance as the nature of the case will admit of, for it is a folly to cut and pare away the hoof as some ignorant farriers do; because by that means they increase the evil instead of removing it, as it is a considerable time before the breach is repaired, and till that is done, the same part is likely to admit more gravel. So much, therefore, and no more, of the hoof should be taken away as is necessary, viz.: till the blackness or discoloration vanishes; then the wound should be dressed with the following balsam:

- $\frac{1}{2}$ an ounce of Gum Benjamin.
- $\frac{1}{2}$ an ounce of Balsam of Peru.
- 6 drams of Myrrh.
- 3 drams of Storax.
- 3 drams of Socotrine Aloes.
- 3 drams of Frankincense.
- 3 drams of Gum Guaiacum.

Powder the ingredients that will powder, and put them in a bottle with one quart of spirits of wine, and let them stand warm for eight or ten days, shaking the bottle up every day. This is an excellent balsam for green wounds, and no person that keeps valuable horses should be without it. Apply it warm to the wound, dipping a piece of tow or lint into it, and fastening it on to the part out of which the gravel or thorn has been taken, and renew it as it grows dry.

A horse is often seized with the Gripes on a journey, the best and surest cure for which is the following :

9 drams of Tincture of Opium.
1 ounce of Spirits of Sweet Nitre.
2 ounces of Olive Oil.

Mix the above in half a pint of mint-water, give it to the horse, and at the same time bed him well, that he may lie down to rest, and be careful not to give him cold water for a day or two after.

A horse that is subject to scouring or purging on the road, has what is called a washy constitution, and such horses seldom bear hardships well, because the aliment passes off before it is thoroughly digested, which is a sign that the fibres of the stomach and bowels are lax ; therefore such horses should not be chosen. The best method of curing this disorder is, to keep the horse chiefly on hard food, and to give him proper exercise, also adding a little of the following composition to each of his feeds :

Aleppo Gall, finely powdered.
Japan Earth, in powder.
Irish Slate, in powder.

Mix these all together, and put them into a bladder for use. Sprinkle a little water on the horse's oats, rub them a while, and then sprinkle about half a spoonful of the powder upon them, and if he have not a very squeamish stomach, he will eat them as well as usual. But remember to proportion the quantity of the powder to the degree of the disorder, or you may make him too costive, which will be as bad as the other extreme when the horse's constitution shall require his being laxative rather than costive.

When a horse purges upon the road, occasioned by foul feeding, or catching cold, give him the following :

4 ounces of Venice Treacle.
3 ounces of Armenian Bole.
1 ounce of Japan Earth.
1 ounce of Powdered Ginger.

Make these up into four balls, and give them night and morning.

When a horse has got a cough by catching cold, give

4 ounces of Cordial Balls.
1 ounce of Liquorice Powder.
1 ounce of Elecampane Bark.

Give all together in warm ale, fasting, and to fast two hours after.

THE END.

READING BOOK

ILLUSTRATED BY

Published by the Council of Public Education
for Ontario

No other or superior has been found to make these books, in every respect,
in type, illustration, paper and binding, the best reading books
that have ever been published in Canada.

PRICES ARE VERY LOW.

Orders solicited.

ADAM HILLMAN, Publisher