IMAGE EVALUATION TEST TARGET (MT-3)

**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14590
(716) 872-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1983

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

- ☐ Coloured covers/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- ☐ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☐ Bound with other material/
Relié avec d'autres documents
- ☒ Tight binding may cause shadows or distortion
along interior margin/
Lare liure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- ☐ Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- ☐ Additional comments:/
Commentaires supplémentaires:

L'Institut a microfilmé le meilleur exemplaire
qu'il lui a été possible de se procurer. Les détails
de cet exemplaire qui sont peut-être uniques du
point de vue bibliographique, qui peuvent modifier
une image reproduite, ou qui peuvent exiger une
modification dans la méthode normale de filmage
sont indiqués ci-dessous.

- ☐ Coloured pages/
Pages de couleur
- ☒ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☒ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
☐	☐	☐	☒	☐	☐
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:

Library,
Geological Survey of Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque,
Commission Géologique du Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

é
étalis
s du
modifier
r une
image

s

errata
to

pelure,
n à

32X

GEO

A

NORTH AMERICAN

GEOLOGY AND PALÆONTOLOGY

FOR THE USE OF

AMATEURS, STUDENTS, AND SCIENTISTS

BY

S. A. MILLER

CINCINNATI, OHIO

1889

Entered, according to Act of Congress, in the year 1889, by

S. A. MILLER,

In the Office of the Librarian of Congress, at Washington.

YBARRILL
Y3VRLB JACOBSON
ADAMAS 90

PRESS OF
WESTERN METHODIST BOOK CONCERN,
CINCINNATI, OHIO.

A c
to the p
any othe
state of
common
no speci
difficult
evidence
make the
names ar
of Natur
the name
than the
work an c
to define

Gene
the fossils
are printe
rules of
other reas
species to
found in
which it
reference v

An a
Palæozoic
is given, t
book and
of the typ
order, will
authors of

4355

PREFACE.

A GENERAL knowledge of Geology is probably of greater importance to the people of the United States than a like amount of information in any other department of natural science; but every one will admit the state of learning in this branch is not of a very high grade. There is a common complaint among well-informed people who have given Geology no special study that the language used is technical, the names long, difficult to understand, and not unfrequently bear upon their face the evidence of affectation, as if those coining the words had attempted to make them as obscure as possible. It is a fact, however, that technical names are absolutely essential to a correct understanding of every branch of Natural History; and when the system of nomenclature is once learned the names are readily understood, and much more easily remembered, than the arbitrary names of individual things possibly can be. In this work an effort has been made to popularize the rules of nomenclature, and to define the technical words in the text or in the Glossary.

Generic and specific names, which have been used by authors where the fossils are not known to occur in the Palæozoic rocks of North America, are printed in italics. Synonyms, names not described as required by the rules of nomenclature, preoccupied names, and those condemned for any other reason, are also printed in italics. When an author has referred his species to a genus to which it does not belong, the specific name will be found in italics under such generic name, and referred to the genus to which it belongs, and at the latter place the original erroneous generic reference will be found in parenthesis.

An attempt has been made to define all genera known from the Palæozoic rocks of North America; the name of the author of each genus is given, the date of coining the word, and an abbreviated reference to the book and page where published, and the etymology of the word and name of the type species. The names of all the species, arranged in alphabetical order, will be found under the genera to which they belong; and also the authors of them, the dates and places of publication, and very frequently

references to two places of publication, especially where, in the first instance, the species was defined without illustration, as has been too frequently done in society publications; the place above and beyond all others where no species should be described unless accompanied by proper illustrations.

An attempt has also been made to correct the misspelling of words so as to perfect the nomenclature, and we call special attention to the Index of Genera, where a few corrections are made that were overlooked in the text, and where the gender of each genus is indicated.

After the author had commenced the preparation of this work, which was several years ago, knowing the great expense attending the making of illustrations, he applied to several State Geologists and others for the privilege of taking electrotypes from the wood-cuts belonging to the State Governments and to the individuals; and he has now to express his acknowledgments to Alfred R. C. Selwyn, F. R. S., F. G. S., Director of the Geological Survey of Canada, who placed at his disposal all belonging to the Canadian Survey, and he availed himself of about one hundred and sixty of the original figures used by the late Prof. Billings; and also to express his obligations to the late Prof. A. H. Worthen, from whom he obtained nearly all those used in the Geological Survey of Illinois. After a very large number of figures had been made by the expensive process of wood-engraving, he learned of the much cheaper electrotype process, and engaged the services of the Kline Photo-engraving Company, of Cincinnati, and for the accuracy and faithfulness with which many figures have been reproduced he is indebted to the skill of the artists in that company.

CINCINNATI, November, 1889.

NO

§ 1.

investiga
upon the
and later
upon the
at the bo
the fact
sea, the p
darkness.
overlying
velopment
of the ear
and migh
branches o
synclinal t
position, a
rarely affo
except by
the dip of
or flexures
study of th

§ 2.

vestigation
occurred w
ages, from
the process
cavities an
characters
nor animal
to preserve
penetrate t
decays, we
operation.
and the up
but are dist

NORTH AMERICAN GEOLOGY.

CHAPTER I.

DEFINITIONS AND LAWS OF GEOLOGY.

§ 1. GEOLOGY is the science which comprehends the structure of the earth and investigates its history. It does not extend to the beginning, nor throw any light upon the astronomical theory that the world was, at one time, in a gaseous state, and later in a condition of fluidity. It commences at the most ancient rocks found upon the surface of the earth. These had their origin, in sedimentary deposition, at the bottom of an ocean. The world was then as large as it is now, and beyond the fact that these rocks were once merely sedimentary layers at the bed of a sea, the previous history of the earth is unknown, and all prior time is impenetrable darkness. Geology commences at the lowest discovered rocks, and investigates the overlying strata, the changes that have taken place, the lapse of time, and the development of organic life, to the present moment. If the strata of rocks on the surface of the earth were horizontal, the science would extend over a short period of time, and might be learned as rapidly as we progress in zoölogy, anatomy, or other branches of Natural History; but the rocks are inclined at various angles, and form synclinal troughs and anticlinal ridges, and expose, in the order of sedimentary deposition, at the maximum more than forty miles in thickness. Mountain regions rarely afford so good opportunities for the study of Geology as a country unbroken, except by the exposures in stone-quarries and the banks of streams. In some States the dip of the strata is quite uniform for a hundred miles or more, without any folds or flexures. It is in these areas the student will find the most inviting fields for the study of the science.

§ 2. The laws of the science have been ascertained, from observation and investigation of the changes now taking place, from a knowledge of those which have occurred within the historical period, from the evidence of change in more remote ages, from the study of the skeletons and harder parts of animals and plants, and the process of infiltration of mineral matter into these organisms, which fills up the cavities and produces petrifications, and from the study and determination of the characters of the petrifications found in the rocks of nearly all ages. Neither plants nor animals turn to stone; flesh can not petrify. When a body is sufficiently firm to preserve its form until water, holding lime or silica in chemical solution, can penetrate the cavities, saturate it, and deposit the stony matter as the organism decays, we have a fossil or petrification. The laws of nature are uniform in their operation. The diversified character of the rocks has resulted from general causes, and the uplifting and inclination of sediments did not occur in one period of time, but are distributed through and belong to all geological ages. We do not assume

the intensity of any forces exceeded, in times past, those which are now in activity.

FIG. 1. Anticlinal axis at *a*. Strata disturbed, folded, and denuded, and afterward unconformable strata deposited upon them, followed by conformable layers.

The changes which the earth has undergone within the scope of geological investigations were produced by the same laws, acting with the same degree of power, as those we may daily witness. This is true of aqueous and igneous action and of all organic and inorganic movements.

§ 3. An *anticlinal axis* is that line from which strata dip to either side. The ridge of a house-top, the slope of the roof representing the dip of the strata, will convey an idea of an anticlinal axis; but an upheaval may be in the form of a dome, or the arc of a circle, and, in such case, the strata incline in all directions from a given point, which is the anticlinal axis. A *synclinal axis* is the reverse of an anticlinal axis. Rocks are called *stratified* whether the planes of the beds are parallel to each other, or rest unconformably. *Conformable* strata have the planes of the beds parallel to each other, and *unconformable* strata have the planes of the strata of one bed resting upon the edges of the strata of another. This must necessarily mark an interval of time between the two which is not represented by a deposit. A *fault* is a dislocation of strata so that the continuity of the

FIG. 2. Section of the Jura Mountains, illustrating the folding of strata, anticlinal axis at *A* and *B*. Synclinal axis between *A* and *B* and between *B* and *C*. Strata unbroken at *A* and *B*, but broken at *C*, *a*, *b*, *c*, and *d*. Strata conformable, though disturbed and thrown into waves.

FIG. 3. Escarpment on the right, debris representing the slope from the escarpment. Outliers, lone rocks, or standing columns in the center and to the left of the illustration. The central figure is a form sometimes called a cheese rock.

mass is destroyed by one side of the fracture being elevated higher than the other. A *dyke* is a wall of rock between the two sides of a fault or fracture, interrupting the continuity of the beds on either side. Sometimes a dyke shows an overflow at the top. When strata terminate abruptly, they terminate in an *escarpment*. An *outlier* is a lone rock in place, or a hill detached by erosion from the surrounding mass of similar beds, of which it evidently once formed a part.

§ 4. The erosion of the earth never ceases. Decomposed and disintegrated substances are being constantly removed by rain and superficial waters to a lower level than they previously occupied. The erosion or denudation must be followed by the deposition of the materials. The deposit at one place can only progress at the rate with which it is transported from another. All strata consist of transported matter, and, as Lyell said, the evidence of the work of denudation is defective, because it is the tendency of every destroying cause to obliterate in great part the signs of its own agency. Stratified rocks, therefore, indicate only part of the erosion which the earth's surface has undergone,

FIG. 4. Strata inclined but conformable. Fault from *f* to *2* and dyke in the fault.

because t
again, an
forms thr
were form
see any p
§ 5.
are gener
union wit
siderable.
of disinteg
and soils,
of sand, d
and someti
Sand blow
from the d
the charac
to the unal
atmospheric
winds, blow
pile it in m
upon the de
exposed to
track at Ea
several tons
parts of the
§ 6. W
creases with
enter the m
and so it wi
burst from l
by expansion
and freezing
will, when b
winds, transp
are transport
in the tempe
boulders are
§ 7. Th
increases with
currents of t
in colder latit
rain. The ra
the disintegr
agencies, dow
the sea 6,368
basin of the r
of a foot thic

because the same materials in a multitude of cases have been broken up again and again, and re-stratified, presenting for our observation only the last of the many forms through which they have passed. The oldest rocks, as well as the most recent, were formed from the waste of older rocks than themselves, therefore we can never see any part of the primitive earth or original solidified matter.

§ 5. The atmospheric forces, in activity and disturbing the surface of the earth, are generally combined with the aqueous, as in frost; or the chemical, as in the union with carbonic acid; but the effects of air and wind are, by no means, inconsiderable. The surface of all exposed rocks and earthy materials bear the evidences of disintegration and denudation. The sun dries up the mud and cracks the earth and soils, while the winds sweep the dust from roads and barren places. Grains of sand, driven by the wind, will groove and polish the hardest rocks and minerals, and sometimes fairly dissolve and carry away limestone and more friable substances. Sand blowing is used in the arts for etching hard materials. All soils have resulted from the disintegration of rocks, and when not transported, the quality depends upon the character of the parent rock immediately below; and the penetration of the soil to the unaltered parent rock will reveal the different stages of the change effected by atmospheric agencies, aided more or less by the effects of frost and water. The winds, blowing inland from large bodies of water, carry sand from the beaches, and pile it in mounds and ridges, called sand-dunes; and the same effects are produced upon the deserts, and to a greater or less extent wherever light or loose materials are exposed to its action. A wind-storm blew a standing locomotive off the railroad track at East St. Louis, and other storms have been known to move bodies weighing several tons. The geological effects of the wind therefore are conspicuous in some parts of the world, while in others they are so slight as to be quite overlooked.

§ 6. Water is an active solvent of rocky substances, and the solvent power increases with heat and pressure. It is also a powerful mechanical agent. It will enter the minute openings in the hardest rocks, freeze, and chip up minute scales; and so it will enter larger cracks and orifices, freeze and break open large rocks, or burst from ledges immense masses. Ice, freezing at the margin of lakes and ponds, by expansion, crowds the loose rocks on the shores in the form of ridges of boulders, and freezing around the free rocks at the bottom in shallow water or near the shore, will, when broken up by partial thawing, and assisted by the force of waves and winds, transport such rocks to distant places. Mud, sand, gravel, and pieces of rock are transported down stream by all rivers, and the transportation is aided by the ice in the temperate and colder latitudes. On the shores of the St. Lawrence transported boulders are found weighing many tons.

§ 7. The capacity of the atmosphere to take up aqueous vapor in suspension, increases with the temperature, and when saturated the least interference with the currents of the air will precipitate rain. Hence there is more rain in warmer than in colder latitudes. Clouds drifting against mountains and high lands will discharge rain. The rain falls upon the ground, disintegrates earthy substances, and transports the disintegrated materials resulting from its own action, and from atmospheric agencies, down the valleys to the ocean. It is said the Ganges annually carries to the sea 6,368,000,000 cubic feet of sediment, which, being spread over the whole basin of the river, comprehending 400,000 square miles, would make a layer 1-1751 of a foot thick. The Ganges, therefore, erodes its basin one foot in 1,751 years.

The area of the Mississippi basin is 1,244,000 square miles, and the annual discharge of sediment by the river is estimated at 7,471,411,200 cubic feet, an amount sufficient to cover the whole basin 1-4640 of a foot. Therefore the Mississippi River removes from its basin a thickness of one foot in 4,640 years.

§ 8. The greater number of valleys in North America have been carved out by the streams flowing in them at substantially the same rate of excavation that is now in progress. All the valleys in Ohio, Indiana, and Illinois, have been excavated by the slow process of the action of rain and the rivers. The Mississippi and all its tributaries have excavated their own valleys, with the exception of a few in the mountain regions. Not only have the valleys been thus excavated, but much of the intervening land has been denuded of many feet of surface rocks. While the beds of the older streams sink extremely slow, if at all, the valleys are gradually widening by the wear and tear of rain and storm. This erosion has taken place since the close of Paleozoic time. The hills are usually terraced because the strata are of different degrees of hardness and durability, the softer and more easily disintegrated are gradually removed by atmospheric influences and the transporting power of rains and springs, leaving the harder and more solid standing out in more or less abrupt slopes and cliffs.

§ 9. The lower limit of perpetual snow under the equator is 16,000 feet above the sea, in the Swiss Alps, in latitude 46 N., it is 8,500 feet, and in the arctic and antarctic regions it reaches the level of the sea. The isothermal lines, around the earth, being affected by the distribution of the land and water surface and the ocean currents, do not follow the degrees of latitude; therefore, in ages past, when the land and water occupied different areas, and the ocean currents moved in other routes, the isothermal lines were correspondingly changed. Above the line of perpetual snow there is an augmentation from year to year, and below it, during the colder seasons, the snow falls many feet in thickness. An equilibrium is preserved by the melting of the snow in sunshine, by occasional rains to which it is subjected, and by the natural tendency to creep down the mountain side by the force of its own gravity. This movement gives rise to glaciers, which follow the depressions or ravines on the sides of the mountains to a considerable distance below the perpetual line of snow. They move very slowly, but transport sand, gravel, and masses of rock, and smooth, polish, and groove their rocky channels, because fragments of rock get interposed between the glacier and the rocks of the valley. The stones carried along on the ice are called the "moraines" of the glacier. There is always one line of blocks on each side, these are called the "lateral moraines." Where there are confluent glaciers the lateral moraines of the tributary glacier are carried into the larger stream of ice, and are called "medial moraines."

§ 10. The effects of glaciers upon the face of the earth are not important, notwithstanding so much has been said about them, and it is evident they have not been much more imposing in past geological ages than they are now. There are probably no evidences of glacial action upon the continent of North America where they do not now exist, except in a few places in the Rocky Mountain region, where they have departed on account of the drainage of adjacent lakes, and some indications in the New England Mountains where they are unknown now, either because that region is somewhat depressed, or because the Arctic Current does not hug the shore as far south as it did in the Pliocene or Post-pliocene period.

§ 11. where fi
the earth
in with c
and let
marine s
rock. C
winds.
surface, a
of the cu
intervene
bergs dri
mouth of
Hope, an
bearing a
known la
than glaci
is inconsp

§ 12. in chemica
this man
storms and
terials at c
where the
never ceas
realize the

§ 13. and depres
historical p
past geolog
neither per
the surface
experienced
phenomena
of sand an
water, and
that steam
the layers o
of temperat
the first 2,0
tinuing inc
rock, or at c
of heat, nor
earth, but,
caverns near
causes which

§ 14. A

§ 11. In high northern and southern latitudes glaciers descend into the sea, where fragments are broken off, which are called "icebergs." Icebergs bear all the earth and rocks they did when constituting part of a glacier, and they soon fall in with ocean currents, and are drifted great distances before they are dissolved, and let the "moraines" fall to the bottom of the sea. In this manner the submarine surface is strewn with foreign mud, sand, gravel, boulders, and fragments of rock. Coast-ice acts in the same manner when blown out into the sea by off-shore winds. The ice sinks into the ocean eight times as deep as it projects above the surface, and when in shallow water it impinges upon the submarine bottom, the force of the current or the winds may cause it to polish or groove the rocks, if fragments intervene, in the same manner that glaciers will polish or groove their valleys. Icebergs drift from Baffin's Bay to the latitude of the Azores, from Greenland to the mouth of the Gulf of St. Lawrence, from the antarctic regions to the Cape of Good Hope, and also to Chili, in South America. Darwin saw one in the southern seas bearing a rock visible twelve feet above the surface, 1,400 miles distant from any known land. Icebergs have a transporting power more than a thousand-fold greater than glaciers, and an eroding power but little inferior, and yet the action of icebergs is inconspicuous now, and has been, so far as we know, in all the ages gone by.

§ 12. A large part of the rain sinks into the ground, takes up mineral matter in chemical solution, flows out in springs, and transports its load to the ocean. In this manner many caves and caverns are excavated. The waves produced by storms and tides beat down the shores of large bodies of water, and deposit the materials at other places. The ocean currents have a drifting and denuding action where the water is shallow. The wear and tear of the earth by the action of water never ceases, and the more we contemplate the subject, the better able we are to realize the magnitude of the never-ending destruction.

§ 13. The violence of earthquakes, and the fires of the volcanoes, the elevations and depressions of land with respect to the sea, seem to have operated within the historical period on as grand a scale as we are warranted in believing they did in past geological ages. Earthquakes and volcanic fires are intimately connected, and neither penetrate the earth to any great depth. Earthquakes have been felt upon the surface of the earth when miners, at a depth of 1,000 feet or more, have not experienced the sensation. The transmission of the vibration is more distinct, and phenomena more apparent where the strata are hard rocks than where they consist of sand and gravel, or softer material. All volcanoes are near large bodies of water, and observation has shown that water gains access to the volcanic foci, and that steam is a powerful agent in all eruptions. The pressure or force of gravity of the layers of the surface of the earth develops the latent heat, so there is an increase of temperature at the rate of about one degree for every sixty feet penetrated for the first 2,000 or 3,000 feet. The deeper borings have not shown the regular continuing increase of the heat, nor is the increase uniform through different kinds of rock, or at different places. The better opinion seems to be that neither this increase of heat, nor the volcanic fires afford any evidence of the internal fluidity of the earth, but, on the contrary, the earth is probably solid, with exception of local caverns near the surface, and local masses of melted matter resulting from chemical causes which are in operation at no great depth.

§ 14. All are more or less familiar with the story of the buried cities of Hercu-

laneum and Pompeii, and the great eruptions of Vesuvius. In 1669 a current of lava flowed from Etna, having a width of 600 yards, and a depth of 40 feet when it reached the sea at the distance of fifteen miles. In 1783 Skapter Jokul, in Iceland, sent forth two currents of lava in opposite directions, one of which extended fifty miles, and the other forty-five. The extreme breadth of the one in Skapter valley was fifteen miles, and the other had a breadth of seven miles. The ordinary height of the current was 100 feet, but in narrow defiles it sometimes amounted to 600 feet. There is no evidence of a volcanic eruption on the continent of North America in past geological ages that surpassed this in volume.

About midnight, August 11, 1772, a luminous cloud appeared to envelop Papandayang, a volcano on the island of Java, and in a short time it actually fell in with a great noise. Immense quantities of volcanic substances were thrown out and distributed for many miles around. It is estimated the mountain for fifteen miles in length and six in breadth was swallowed up in the earth by this commotion. Forty villages were engulfed or destroyed, and 2,957 inhabitants perished. It seems in this instance the eruptions had formed a corresponding cavity beneath the surface, and when the weight above overcame the resistance, the volcano suddenly fell into the abyss beneath.

A volcano forced its way from beneath the sea into the atmosphere off St. Michael's, Azores, in 1811. It was first seen above the sea on June 13th. The appearances were exceedingly beautiful, the volcano shooting up columns of the blackest cinders to the height of between 700 and 800 feet above the surface of the water. When not ejecting ashes, an immense body of vapor or smoke revolved almost horizontally on the sea. The bursts were accompanied by explosions resembling a mixed discharge of cannon and musketry, and a great abundance of lightning. By the 4th of July an island was formed a mile in circumference and 300 feet high. In the center there was a crater full of hot water, which discharged itself through an opening facing St. Michael's. The island subsequently disappeared beneath the water.

Twelve islands constitute the Hawaiian Group, four of these are mere barren rocks; the remaining eight have an area of about 6,000 square miles. All of these islands are volcanic, and no other rocks than volcanic are found upon them save a few remnants of sea-beaches. They are all mountainous, and the deep sea surroundings have shown the islands are only the summits of gigantic mountain masses. Mauna Kea, on Hawaii, is 13,900 feet above the sea, and Mauna Loa 13,700 feet. If the ocean were driven away, it is said these mountain peaks would stand 30,000 feet above the foot of the mountain range. On Hawaii the volcanic forces are still in operation. On Maui they rested at a recent epoch, or within a few hundred years. On the other islands they have long been extinct, and the piles built up have been greatly eroded. On Hawaii there are two grand foci of volcanic eruption where the fires are now raging, Mauna Loa and Kilauea. Mauna Loa is the largest volcano in the world, and none approach it in the magnitude of its eruptions. A moderate eruption represents more material than Vesuvius has emitted since the days of Pompeii, and the flow of 1855 would have nearly built Vesuvius. On the whole, it appears there are as many active volcanoes, and some as vast and frightful in eruptive power as seem to have existed at any other single period in geological time.

§ 1
ing 4,6
of Chil
more.
1819 an
about fi
the delta
quakes i
every ce
are warr

§ 1
terventio
the Balti
coast of
area in C
of Fundy
feet of wa
the New
tion of sev
elevations

§ 17.
accompany
rise or fall
volcanic an
show the i
present on
they act by
to internal
intermittent
up until str
earth, whe
forever. V
heat contain
sidered as l
earth was a

Geyser
intermittent
The analog
erupting ho
supposed to
and by its m
plosive gas,
out the colu
is furnished
dition of th
counted for.
The me

§ 15. An earthquake in New Zealand in 1856 raised a tract of land comprising 4,600 square miles, from one to nine feet. In 1822, and again in 1835, the coast of Chili for several hundred miles was elevated from one to three or four feet or more. The estimated area raised in 1822 amounted to 100,000 square miles. In 1819 an earthquake at Cutch, in the delta of the Indus, raised an extent of country about fifty miles long and sixteen miles wide, ten feet, while a considerable tract in the delta of the Indus sank down. Such are a few of the effects produced by earthquakes in the present century; they are similar to those which have occurred in every century during the historical period, and are quite as extensive as any we are warranted in believing occurred in any of the earlier geological ages.

§ 16. It is said large tracts of land are elevated and depressed without the intervention of earthquakes. It is said there has been an elevation of land bordering the Baltic, during the historic period, of about three feet in a century. The whole coast of Scandinavia is said to be gradually rising at a very slow pace. A large area in Greenland is reported as slowly subsiding. At Fort Lawrence, in the Bay of Fundy, there is a pine and beach forest covered at high tide by about thirty feet of water. And it is claimed there is some evidence of subsidence on part of the New England Coast, where we have the most indubitable evidence of an elevation of several hundred feet since the beginning of the Post-pliocene period, but these elevations and depressions may have been accompanied with earthquakes.

§ 17. Earthquakes and volcanoes have a common origin, the former always accompany the eruption of the latter, and it is not likely any great areas of land rise or fall without the intervention of the same energies. The proximate cause of volcanic and earthquake phenomena is not fully known, and it is much easier to show the improbability of the many theories offered for their explanation than to present one free from objections. Volcanoes are intermittent in their eruptions; they act by spasms of activity, separated by intervals of repose. If they were vents to internal fluidity of the earth, the streams of flowing fire would be constant, not intermittent explosions. If they were vents to any great mass of melted matter pent up until strength enough were obtained to force a passage way to the surface of the earth, when the vents would open the reservoirs would exhaust themselves and close forever. Volcanoes are not to be attributed to the remains or residue of enormous heat contained in the globe, at some remote period of its physical evolution, or considered as lending any support to the nebular hypothesis, or the theory that the earth was at one time in a gaseous or fluid condition.

Geyser (from the Icelandic word *geysa*, to gush,) is a periodically eruptive or intermittent hot spring, from which the water is projected in a fountain-like column. The analogy between it and a volcano is so striking that it might be called a volcano erupting hot water instead of melted lava. In the case of a geyser, cold water is supposed to sink from the surface to heated rocks; it starts as a passive liquid, and by its molecular absorption of heat is converted in the depths into an elastic, explosive gas, which ejects it through another orifice to the surface. The gas forces out the column of water and escapes; then quiet ensues until a new supply of water is furnished. This accounts for the intermitting flows. Grant the local heated condition of the rocks below, and all the phenomena of the geysers may be accounted for.

The melted lavas of volcanoes bring up with them great quantities of the vapor

of water, having an enormous expansive power which is given off as steam at the moment of eruption. Lava is generally a sponge-like mass of myriads of visible vesicles formed by the sudden exclusion of the water-vapor in the act of solidification. There is abundant evidence of the participation of water and its constituent gases in volcanic phenomena. From the proximity of volcanoes to or occurrence in the sea, it has been supposed their active state is produced by the percolation of seawater to metallic bases of the earths, or alkalies, at various depths, which bases become inflamed and chemical action ensues, producing the eruption. The oxygen of the water is supposed to unite with the metallic base, the hydrogen to unite with sulphur, forming sulphureted hydrogen gas, and with the chlorine forming muriatic acid gas, etc. The gases evolved from volcanoes are muriatic acid gas, sulphur combined with oxygen or hydrogen, carbonic acid gas, nitrogen, and aqueous vapor. Electricity is a factor in all earthquakes and volcanic eruptions. Its action is manifest in the atmospheric disturbances, in the undulatory movement on the surface of the earth, and in the speed with which the earthquake wave travels. An earthquake moves in the direction of the wave at a rate frequently exceeding fifty miles in a minute, and when the movement is communicated to the waters of the ocean, the waves follow at a pace hundreds of times slower. Suppose a powerful current of electricity near the surface of the earth, to be broken, and suddenly restored, the shock may be supposed to resemble that of an earthquake. Fusion might result in consequence of such restoration. The crystallization of stratified rocks might break such electrical currents, if any exist in the earth, or it might disturb the equanimity of the electricity if it exists in a passive state, to the same extent as if it were a broken and restored current. In other words, subterranean electric currents, if once excited, may melt the rocks and produce the heat necessary, when assisted by the presence of a sufficient quantity of water, to produce volcanic eruptions. Such are some of the theories to account for the instigating or proximate causes of earthquakes and volcanoes.

The mouth of a volcano is called a *crater*, though the pit on Kilauea has been called a *caldera*. If steam alone escapes through a vent, it is called a *fumarole*; but if sulphurous vapors also escape, it is called a *solfatara*. When hot springs deposit lime, it is called *tufa*; but if the deposit is silicious, it is called *sinter* or *geyserite*. Lava consists of silica, alumina, lime, magnesia, soda, potash, and iron oxide. If the silica is in excess, it is *trachyte*, and belongs to what lithologists call the acidic group, from the large quantity of silicic acid it contains; but if there is a large proportion of soda or potash and lime or magnesia, and not more than 50 per cent of silica, it is a *basalt*, and belongs to the basic group, from the larger quantity of alkaline and earthy bases it contains. Trachyte is a grayish igneous rock, of rough fracture owing to the grains of glassy feldspar which mainly constitute it. Basalt may be light-colored crystalline or granitoid, or dark colored, compact, massive, like dolerite; but in addition to labradorite and pyroxene, it contains chrysolite in disseminated grains. When lava becomes glassy, it is called *obsidian*.

§ 18. The most important change taking place upon the earth is in constant operation at the bed of the ocean. Near the shore it is a littoral deposit; farther away it is a chalky deposit, consisting of foraminifera and shells, and in deeper water it is a red, silicious clay. The character of the deposit is dependent upon the depth of the ocean, except where washings from land affect it. The depth of the pure

globiger
depths t
consists
moderat
increases
the chem
of water
The simp
of the d
which ex
they do
induratio
arenaceou

§ 19.
modifying
changes,
spot, for
corals, is
operation,
past to de
and to ele
denude it

§ 20.
much of v
a large pa
time, been
form of ri
existing in
different d
with the s
great diver
separately
rocks. Cre
hence, mar
the strata u
the animal
tions of typ
approximat
equal depth
water and
farther from
tinuous ove
or limestone
mony to pr
been ascerta
characterize
order of sup

globigerina ooze, or chalky deposit, is limited to about 2,250 fathoms, and at greater depths the deposit gradually passes into fine pure clay, and below 2,500 fathoms it consists almost wholly of a silicate of the red oxide of iron and alumina. At moderate depths shells fall upon the bottom, in perfect condition; as the depth increases they become more and more brittle, and finally break up and disappear by the chemical action which affects them, until, having passed through 2,500 fathoms of water, nothing is left save an insoluble residue, which constitutes the red clay. The simple fact of the increasing depth of the ocean gives variety to the character of the deposits. But at the greatest depth to which the dredge has descended, which exceeds five miles, the silicious shells of Radiolarians exist as abundantly as they do in the shallower depths of the ocean. Such deposits, in the process of induration, become stratified and laminated, and form calcareous, argillaceous, and arenaceous or silicious rocks.

§ 19. Animals, secreting carbonate of lime, have played an important part in modifying the surface of the earth. The coral-making polyp has wrought great changes, because the reef-forming genera continue the accumulation, on the same spot, for centuries, and the influence of the Bryozoa, which produce only delicate corals, is everywhere conspicuously engraved. There are other agents, inferior in operation, affecting the surface of the earth, and all combined have served in times past to deposit in water all the rocks constituting the continent of North America, and to elevate the land above the seas and lakes, after such deposition, and again to denude it and present it to us with its mountains and valleys as they now exist.

§ 20. Every part of the surface of the earth has been covered with water, and much of what is now dry land has been several times inundated; and it is supposed a large part, if not the whole area covered by the oceans, has, at some period of time, been above the water line. The elevations and depressions have been in the form of ridges, with intervening basins, in different ages of the world; and basins, existing in the same age, have been filled with deposits of different kinds and in different degrees of rapidity,—some being filled with drifted materials, and others with the secretions of animal and vegetable organisms. Consequently there is a great diversity in the structure of the land of different continents, and they must be separately investigated. The most recent deposits may be made on the most ancient rocks. Cretaceous deposits may occur upon the Silurian, or Jurassic on the Devonian; hence, many difficulties are encountered in ascertaining the chronological order of the strata upon each continent; and this would be utterly impossible were it not for the animal and vegetable remains, which have followed the progress of time in evolutions of type and structure in different oceanic basins, so as to furnish the means of approximately parallelizing the strata. Different kinds of rocks are forming at unequal depths of the ocean, at the same time; conglomerates and sandstones in shallow water and near the shores; chalky, and slaty or shaly in deeper water, and silicious farther from land and at still greater depths. Strata of the same kind are not continuous over large areas; but change within short distances from sandstone to shale or limestone; hence, it is never safe to trust to the character of the rock for the testimony to prove its age. We must go to the fossils for the evidence, because it has been ascertained that species did not generally live beyond a geological period, and characterized different Groups of rocks, and thus become infallible guides to the order of superposition. No two periods are represented by like assemblages of fossil

forms, and this dissimilarity furnishes the facts upon which the Groups of rocks are distinguished from each other. Comparison of the fossils shows a progression in development along an ascending scale toward the higher and more enduring plants and animals, and the extinction of lower or less highly organized forms.

§ 21. *Sandstone* is a rock made of sand derived from a silicious rock. When pure it is used for making glass. Iron usually colors it red or yellowish, and often cements it into good building stone. When a little clay is intermixed it is called freestone, and if it contains gravel it is conglomerate, or if loosely cemented in the air and not under water a pudding-stone. When sandstone is subjected to heat and pressure it is metamorphosed and becomes quartzite.

Shale is a soft, fine-grained, aluminous rock, in layers. If it is pure it is clay shale; if it contains sand it is sandy shale; if bituminous matter, bituminous shale. When the shale is hardened it becomes slate. Slate rocks among the metamorphic series are called schists. The clay slate used in North Carolina for making slate-pencils is called pyrophyllite.

Limestone is ordinarily composed of lime and carbonic acid, with impurities of clay, sand, and iron. Hydraulic limestone contains clay and magnesia. Magnesian limestone is called dolomite, after Dolomieu, a mineralogist. Lithographic stone is a very even-grained, compact limestone, usually of buff or drab color. Chalk is a soft limestone, and marble is a hard crystalline limestone. Gypsum, alabaster, calcite, dogtooth spar and satin spar are names given to crystalline limestone.

§ 22. The general order of superposition of the rocks of North America has been ascertained, and they have been divided into Systems and Groups. Another division has been made, founded on the organisms that occur in the rocks, viz: Eozoic, Palæozoic, Mesozoic and Cænozoic. Some use the word Archæan instead of Eozoic. The Eozoic includes the Laurentian and Taconic Systems. The Palæozoic includes the Lower Silurian, Upper Silurian, Devonian, Subcarboniferous and Carboniferous Systems. The Mesozoic includes the Triassic, Jurassic and Cretaceous Systems. The Cænozoic is synonymous with the Tertiary System. These Systems may be very closely parallelized with the strata of Europe and other parts of the world. The words "System" and "formation" are in use with this nomenclature, as Devonian "System" or Devonian "formation," but more generally they are both omitted as unnecessary appendages to the names of the divisions.

The Taconic is introduced in many places with conglomerate layers resting unconformably upon the Laurentian; the Lower Silurian commences with the Potsdam sandstone, the Upper Silurian with the Medina sandstone, the Devonian with the Oriskany sandstone, the Subcarboniferous with the Waverly sandstone, and the Coal Measures with the Carboniferous Conglomerate. Each of these great divisions commences with drifted materials, and important changes of the fauna. They are each capable of subdivision into Groups, and they are not only convenient in the discussion of the science, but they are, to a certain extent, founded in nature.

§ 23. For the purpose of more definite classification these larger divisions are subdivided. Each subdivision is called a "Group," and it generally bears the name of the place where first studied and described; as, the Potsdam Group, so named because the strata were first studied and described at Potsdam, New York. This method is preferred to any other, because the geographical name, when combined with the word "Group," is sufficiently technical. It can not be used for any other

purpose
strata,
erates,
geologic
acters o
simple
not in c
To spea
of its
characte

§ 2
palæozoi
cant of f
different
Scalent,
tine, Un
break, M
noon, D
Unfortun
ture them
use of co
extensive
consequen
suggested

§ 25.
epoch, ar
cause exp
palæontol
When pro
name for
it at a p
Groups ha
Utica in t
or Hudson
black slate
though cal
becki, Lep
plainly dis
Utica had
Cincinnati
or both of
the Utica

§ 26.
been name
another loc
priority in
defined so

purpose; it can never mislead as to the mineral structure or relative position of the strata, and it indicates the typical locality of the exposure. Sandstones, conglomerates, limestones, and shales, occur in nearly every Group, and for this reason geological subdivisions can not be established upon the mineral or chemical characters of the rocks. The rocks which form these Groups are composed of a few simple minerals, which are repeated over and over again in the different layers, but not in chronological succession, nor in any other way indicative of age or position. To speak of a rock as a limestone, sandstone, shale, slate, or clay, conveys no idea of its geological age or place. It is merely the expression of a mineralogical character.

§ 24. Prof. Rogers conceived the idea of improving the nomenclature of the palæozoic rocks by dividing them into fifteen parts, and giving them names significant of their relative ages. This he did by using words suggesting metaphorically different parts of a day, as follows: Primal, Auroral, Matinal, Levant, Surgent, Scalent, Premeridian, Meridian, Post Meridian, Cadent, Vergent, Ponent, Vespertine, Umbral, and Seral, meaning respectively the formations of the Dawn, Day-break, Morning, Sunrise, Mounting Day, Climbing Day, Forenoon, Noon, Afternoon, Declining Day, Descending Day, Sunset, Evening, Dusk, and Nightfall. Unfortunately for his attempt to substitute another for the geographical nomenclature then quite well established and susceptible of indefinite expansion, without the use of conflicting terms or words that could mislead the student, there were several extensive Groups of rocks full of the remains of animal life, then unexplored, and consequently quite unknown to his system. For obvious reasons the nomenclature suggested by Mr. Rogers has not been adopted.

§ 25. The words series, strata, layer, deposit, zone, bed, horizon, period, age, epoch, and era are not technical names, but are used in geological descriptions, because expressive and convenient. Each Group must, in all cases, depend upon the palæontological characters, and can never rest upon the structure of the rocks. When properly defined, it is established, and no one has a right to substitute another name for it, nor to propose a name, simply because of inability to properly distinguish it at a particular locality. For example, the Trenton, Utica, and Hudson River Groups had been long established, when some one, being unable to distinguish the Utica in the vicinity of Cincinnati, and not knowing whether the rocks are Trenton or Hudson River, proposed to call the exposure the "Cincinnati Group." The black slate, which characterizes the Utica in New York, does not exist at Cincinnati, though calcareous slates and shales of the same age do, and they contain *Triarthrus becki*, *Leptobolus lepis*, and other characteristic fossils, while the Hudson River is plainly distinguishable above, and the Trenton as readily determined below. If the Utica had thinned out in its extension westward from New York before reaching Cincinnati, there would have been no excuse for calling the Trenton or Hudson River, or both of them together, by a new name, nor is there any excuse for so doing when the Utica is easily distinguished.

§ 26. Another kind of synonymy to be deplored exists where a Group has been named and thoroughly defined, and for some trivial reason, the geologists of another locality use another name for rocks of the same age without regard to priority in nomenclature. For example, the Calcareous Group was established and defined so as to include rocks other than Calcareous sandrock, and ten years after-

ward rocks of the same age on the Mississippi were called the "Lower Magnesian Limestone," and geologists of that locality persist in the use of the latter name, because they say the word Calcareous is not admissible from the lithological character of the rock. It is to be regretted that the name Calcareous has come down to this generation as the name of a Group of rocks, but it is as well established as the name of any other Group, and like the word Tertiary, which has no application to the rocks to which it is applied, is fastened upon the science, and so interlocked with it that it can not be eradicated even were it desirable so to do. With how much less reason should we encourage the use of another mineralogical name, having more limited application, in its stead!

§ 27. The rule is, the law of priority should be rigorously enforced where a Group has been named, and the fossils have been so described and illustrated that it may be identified by a palæontologist elsewhere than at the typical locality. Synonymy is always the result of ignorance, and much of it has come from those whose work has been absolutely worthless.

§ 28. Experience has shown the impracticability of making lesser subdivisions for the purpose of geological nomenclature, than Groups, especially in the present state of the science, though it is eminently fit and proper to speak of the marl-beds or sandstone layers in any Group, or of the *Glyptocrinus* or *Orthis* beds at any locality. Minute and careful definition and description of the characters of each and every part of a Group is one thing, and the suggestion of a geological subdivision, founded upon a marked peculiarity at one locality, which can not be distinguished at another, is quite a different thing. It must not be supposed none of the Groups will be subdivided, but proposing a name is not establishing a Group. The Coal Measures ought to be divided into Groups because of the great thickness of the fossiliferous rocks, and a temporary division in some localities is indicated by the use of the words Upper, Middle, and Lower Coal Measures, but great palæontological information must be acquired before any practicable subdivision can be made.

§ 29. The stratigraphical division of the rocks of North America into Groups bearing geographical names, with an approximate thickness in ascending order, is as follows:

Laurentian System, not divided into Groups,		40,000 feet.
Taconic.	{ Lower Taconic, not divided into Groups,	25,000 "
	{ Georgia Group, including St. John Group,	10,000 "
	{ Swanton Group,	10,000 "
Lower Silurian.	{ Potsdam Group,	4,000 "
	{ Calciferous Group,	2,000 "
	{ Quebec Group (very doubtful),	6,000 feet ?
	{ Chazy Group,	1,000 "
	{ Black River Group,	150 "
	{ Trenton Group,	1,100 "
	{ Utica Slate Group,	600 "
	{ Hudson River Group,	1,200 "
Upper Silurian.	{ Medina Group,	2,500 "
	{ Clinton Group,	500 "
	{ Niagara Group,	800 "
	{ Guelph Group,	1,200 "
	{ Onondaga Group,	1,200 "
	{ Lower Helderberg Group,	2,000 "
Carried forward,		102,050 feet.

Devonian.
Subcarboniferous.

Carboniferous.
Triassic
Jurassic

Cretaceous.
Tertiary.

§ 30.

and not fr
rocks of C
of the Ge
of these ro
the Potsda
rests uncon
and hence
consist of s
thicknesses
and quartz
phic, or pr
Granite
applied. I
usually giv
being separ
excess, it i

		<i>Brought forward,</i>	102,050 feet.	
Devonian.	{	Oriskany Group,	300	"
		Upper Helderberg Group,	900	"
		Hamilton Group,	1,400	"
		Portage Group,	1,400	"
		Chemung Group,	3,000	"
		Catskill Group,	7,500	"
Subcarboniferous.	{	Waverly Group,	500	"
		Burlington Group,	500	"
		Keokuk Group,	200	"
		Warsaw Group,	100	"
		St. Louis Group,	400	"
		Kaskaskia Group,	720	"
In Pennsylvania, where the Subcarboniferous can not be separated into Groups, there is a thickness of 5,000 feet, and in Nova Scotia 6,000 feet.				
Carboniferous.	{	Carboniferous Conglomerate, or Millstone Grit,	6,000	"
		Coal Measures,	10,000	"
		Permian Group,	1,000	"
Triassic not divided into Groups,		25,500	"	
Jurassic not divided into Groups,		10,000	"	
Cretaceous.	{	Dakota Group,	1,000	"
		Fort Benton Group,	900	"
		Niobrara Group,	500	"
		Fort Pierre Group,	1,200	"
		Fox Hills Group,	4,000	"
		Fort Union or Laramie Group,	7,000	"
Tertiary.	{	Eocene,	15,000	"
		Miocene,	3,000	"
		Pliocene,	3,000	"
		Post-pliocene,	1,000	"
Total,		211,150 feet.		

CHAPTER II.

LAURENTIAN SYSTEM.

§ 30. THE Laurentian System was so named from the Laurentian Mountains, and not from the St. Lawrence River. The name was applied to the metamorphic rocks of Canada as a scientific term, by Sir William Logan, in the Report of Progress of the Geological Survey of Canada for the years 1852-53. His special study of these rocks began as early as 1846. He applied the name to all rocks lower than the Potsdam; but Emmons had preceded him in defining the Taconic System, which rests unconformably upon the rocks that comprise nearly all which Logan studied; and hence the Laurentian is confined to the rocks below the Taconic. The rocks consist of sedimentary strata altered to a highly crystalline condition—great vertical thicknesses of gneiss and granitoid rocks, separated by masses of crystalline limestone and quartzite. Previous to this geographical name they were called azoic, metamorphic, or primary rocks.

Granite is a word derived from the granular texture of the rock to which it is applied. It is crystalline and composed of quartz, felspar, and mica. The felspar usually gives the predominating color. When the granite is stratified, the laminae being separated by thin scales of mica, it is *gneiss* or *granite schist*; if mica is in excess, it is *mica schist*; when hornblende displaces the mica, it is *syenite* (named

from Syene, in Egypt); and if it only partially displaces it, it is *syenitic granite*. Many of the granites and syenites are intrusive, while others, not distinguishable from these, take the place of sedimentation and pass into gneiss or mica schist. *Felspar* signifies rock-spar from the German word *fels*, a rock, though it is usually spelled *feldspar* from the German word *feld*, a field, and therefore made to signify field-spar. There are several species of felspar, dependent upon the potash, soda, or lime they contain. That which usually enters granite is *orthoclase*, or potash felspar, and is compact laminated, or compact crypto-crystalline, consisting of about the following substances: silica 64.6, alumina 18.5, and potash 16.9. When soda enters into the composition of the felspar, it becomes *albite*, and the granite is then disposed to undergo spontaneous disintegration, which sometimes takes place below direct atmospheric influences at great depths in the earth. The kaolin of the Chinese is derived from felspar from the disintegration of granitic rocks, and porcelain clay is often from the same source. Garnets are common in gneiss and mica schist. The most common mica, and that which generally enters into granite, gneiss, and related rocks, is called *muscovite*. Other species in the mica group are called phlogopite, biotite, lepidomelane, astrophyllite, lepidolite, and cryophyllite.

§ 31. Logan said of the Laurentian System: "Stretching on the north side of the St. Lawrence from Labrador to Lake Huron, this series occupies by far the larger portion of Canada, and its strata probably possess a great thickness. To determine the superposition of the various members of such an ancient series of rocks is a task which has never yet been accomplished in geology, and the difficulties attending it arise from the absence of fossils to characterize its different members. Bands of the crystalline limestone are easily distinguished from bands of the gneiss; but it is scarcely possible to know from local inspection whether any mass of limestone in one part is equivalent to a certain mass in another. They all resemble one another lithologically, and although masses dipping in the same direction are met with, running for considerable distances rudely parallel with one another, it is scarcely ever safe to take for granted that they are stratigraphically distinct. The dips avail but little in tracing out the structure; for in the numerous folds of the series the dips are frequently overturned, and the only reliable mode of pursuing the investigation and working out the physical structure, is patiently and continuously to follow the outcrop of each important mass in all its windings as far as it can be traced, until it becomes covered up by superior, unconformable strata; is cut off by a great dislocation, or disappears by thinning out."

§ 32. The surface area accupied by the Laurentian series in Canada and British America, exclusive of any exposure that may exist in the Cordillera or Rocky Mountains, is not less than 250,000 square miles. The northern limit is the Arctic Ocean; from here it may be traced south upon the western side of Hudson's Bay, and appearing upon its eastern side it spreads over the greater part of Labrador, and extends to the Gulf of St. Lawrence. The southern limit is the St. Lawrence from Labrador to Cape Tourmente, a distance of 600 miles, except a narrow border of Taconic on the Strait of Belle Isle; another at the mouth of the Mingan River; a third near the Seven Islands, and two on Murray Bay River, and the Gouffre. Extending westwardly it occurs 30 miles north of Montreal, and follows up the Ottawa River for a distance. It then strikes off to the Thousand Islands, and crosses over into New York, where it exposes an area of 10,000 square miles. From there it

extends
Lake Su
of Rainy
Wisconsin
Woods in
that divi
directions
covered w

§ 33.

of Michig
one of wh
granite, g
useful min
posures in
North Can
belt from
eastern sec
ward into
of gneiss,
southern b
the Virgin
and crosser
another sm
in Georgia
at other pla
Iron Mount
many expos
rection of
Nevada, Ut

§ 34.

as follows:

1. Ortho
2. Cryst
3. Ortho
4. Cryst
5. Ortho
6. Cryst
7. Ortho
8. Procto
9. Ortho
10. Anorth

This is
thickness in
by no means
be that the

extends north-westerly a short distance north of Lake Huron, and bordering upon Lake Superior, a great part of its length, it appears at Lake of the Woods, north of Rainy River, though an arm extends south of Lake Superior into Michigan and Wisconsin. The western boundary of this great area extends from Lake of the Woods in a sinuous northerly direction among the lakes, and following the highlands that divide the waters which flow into Hudson's Bay from those flowing in other directions, to the Arctic Ocean. There are some patches within this general outline covered with Taconic rocks, or those of Post-pliocene age.

§ 33. The arm of this great exposure, which appears in the Upper Peninsula of Michigan, has an area of about 1,839 square miles, consisting of several tracts, one of which touches Lake Superior west of Marquette. The rocks are chiefly granite, gneiss, syenite, and crystalline limestone, which thus far have afforded no useful minerals. The surface area in Wisconsin is somewhat greater. Other exposures in the United States are confined to irregular areas in the mountain regions. North Carolina exposes about 20,000 square miles, or nearly half the State. One belt from twenty to twenty-five miles wide, crosses the northern part of the sub-estern section of the State upon which the capital is situated. It extends northward into Virginia, and southward beyond Cape Fear River. It consists generally of gneiss, which passes into granite or mica schist. Another belt extends from the southern border of the State at Catawba River in a north-east direction, almost to the Virginia line near Roxboro, and reappears eight or ten miles to the eastward and crosses the northern border about midway of Granville County. There is another small area in the southern part of Orange County. Limited areas are found in Georgia, Virginia, Pennsylvania, New Jersey, Vermont, New Hampshire, and at other places in the Appalachian chain. A small area occurs in Missouri near Iron Mountain, and another in Arkansas. In the Rocky Mountain region there are many exposures, some of which are quite large. They generally trend in the direction of the mountain chain, and are found in Mexico, New Mexico, Arizona, Nevada, Utah, Colorado, Idaho, Nebraska, Wyoming, and Montana.

§ 34. A section taken by Logan in the region where he studied the rocks, is as follows:

1. Orthoclase gneiss, composing Trembling Mountain,	5,000 feet.
2. Crystalline limestone of Trembling Lake,	1,500 "
3. Orthoclase gneiss,	4,000 "
4. Crystalline limestone of Great Beaver and Green Lakes, with interstratified garnetiferous rock and hornblendic orthoclase gneiss,	2,500 "
5. Orthoclase gneiss, garnetiferous gneiss and quartzite below the Grenville limestone,	3,500 "
6. Crystalline limestone of Grenville, with interstratified gneiss,	750 "
7. Orthoclase gneiss,	1,580 "
8. Proctor's Lake limestone,	20 "
9. Orthoclase gneiss, passing gradually into Anorthosite between Proctor's Lake and Morin band,	3,400 "
10. Anorthosite above the Morin band,	10,000 "
Total,	32,250 feet.

This is about the thickness in New Hampshire, and not equal to the estimated thickness in Wisconsin. Enough is known, however, to show this section of Logan's by no means represents the total thickness in Canada. The better opinion seems to be that the Laurentian series has a thickness in Canada of more than 40,000 feet,

and probably more than eight miles, or quite as great as it is in Bohemia or any other European country.

§ 35. While these are the oldest rocks known, they were, in their unmetamorphosed condition, ordinary sediment in water derived from materials that preceded them. They were formed by the disintegration, denudation, and redeposition of older rocks, which in their turn preceded others, in how many cycles of change we have no means of knowing. Their upheaval above the surface of the sea was the beginning of the North American continent. The trend of the range in this upheaval is as nearly east and west as the later elevations of the Appalachian and Rocky Mountain chains are north and south. These rocks were until recently supposed to have preceded the existence of both vegetable and animal organisms, and were, therefore, called azoic, but in addition to the fossil *Eozoon canadense* there are other evidences of organic life, as follows:

1. The iron ore evidences organic life, because all the accumulations of iron now in progress are formed by the agency of organic matter. The peroxide of iron existing in the rocks is not soluble in water alone, but the addition of decomposing organic matter deoxidizes it, and carbonate of iron is formed, which is soluble and may be precipitated. Peroxide of iron being insoluble, the infiltrating waters which take up soda, lime, and magnesia from sediments, can not remove this metal unless they contain organic matter. The evidence of the reducing and dissolving action of organic matter is, in the great thickness of sediments, almost destitute of iron and in the extensive beds of iron ore.

2. The masses of limestone tend to prove the existence of organic matter, because limestone in process of formation is almost wholly composed of shells, corals, tests of foraminifera, and other animal secretions, and nearly all the unmetamorphosed limestones of past ages are largely composed of organic relics.

3. Graphite occurs in beds, imbedded masses, and in scales; in granite, gneiss, mica schist, and crystalline limestones; it results from the alteration by heat of coal in the Coal Measures, and is a common product of furnaces. Its presence is, therefore, an evidence of organic matter, because we know of no other source for its derivation, and are able to trace its origin to vegetable matter in rocks of a less remote date. It is inferred the carbon was collected by marine vegetation at that early period.

4. In the lowest non-metamorphosed rocks, and in the shales and limestones of the Taconic System, several classes of the animal subkingdom are represented, which indicates, if we judge by analogy with subsequent changes and progress of life, that the seas in much earlier times must have teemed with life. This is the only view consistent with the modern theory of evolution and the present state of knowledge concerning the development of animals and vegetables.

5. The *Eozoon canadense*, a fossil rhizopod, is found in the Grenville band of limestone near the middle of the series. The limestone is thus described: "The general character of the rock connected with the fossil produces the impression that it is a great foraminiferal reef, in which the pyroxene masses represent a more ancient portion, which, having died and become much broken up and worn into cavities and deep recesses, afforded a seat for a new growth of foraminifera, represented by the calcareo-serpentinous part. This in its turn became broken up, leaving, however, in some places, uninjured portions of the organic structure. The main difference

between
with the
more and
reef."

6. T
abundant
in potash,
in the sed

7. A
lifeless; b
from those
magnesium
rocks in th

§ 36.

silica, alun
transforma
and new co
acting unde
sea would
volcanic an
as active ag
sedimentary
accumulation
the crust of
is too chime

§ 37. S

continued e
mospheric a
land and in
furnished m
and will fore
rentian rock
The Taconic
glomerate co
unrevealed
and never c
formably up
gneiss, and
precious met
of later time

between this foraminiferal reef and more recent coral reefs seems to be, that while with the latter are usually associated many shells and other organic remains, in the more ancient one the only remains yet found are those of the animal which built the reef."

6. The relatively large amount of potash in the Laurentian series indicates an abundant marine vegetable life, because later fossil fucoidal layers frequently abound in potash, and living algæ secrete potash from the ocean in such form as to retain it in the sediments now accumulating, and in which they are buried.

7. And, negatively, we have no good reason to think the Laurentian Age was lifeless; beside, the actual elements composing the Laurentian rocks are not different from those in succeeding formations; indeed, oxygen, hydrogen, silica, aluminum, magnesium, calcium, potassium, sodium, iron, and carbon constitute .99 of all the rocks in the world.

§ 36. The change which sedimentary strata of sands and clays, composed of silica, alumina, and potash, underwent to form granite, gneiss, and mica schist; the transformation of sand into quartzite, and all other changes caused by crystallization and new combinations, are supposed to be owing to chemical and molecular forces, acting under the conditions of pressure, heat, and moisture. The pressure of a deep sea would develop a high degree of heat. The mountain ranges have undergone volcanic and earthquake upheavals which may have accompanied the metamorphism as active agencies. It would seem to be a law that mountain upheavals follow great sedimentary deposits, and the chemical action is most powerful under the grandest accumulations; but the idea that such accumulations bend the crust of the earth, or the crust of the earth contracts and wrinkles up mountain chains in the act of cooling, is too chimerical for consideration.

§ 37. Sedimentation ceased when the beds were forced above the ocean, but continued elsewhere. When the beds were elevated, the wear and wash from atmospheric and aqueous forces began, and deposits ensued upon the margin of the land and in the depths of the ocean. The denudation of the anticlinal heights has furnished many geological sections, but the older rocks remain hidden from view, and will forever remain unknown. There is absolute nonconformability of the Laurentian rocks with overlying Groups at every locality which has been examined. The Taconic is introduced by total nonconformability, and frequently with a conglomerate containing pebbles derived from the adjacent Laurentian. Here is an unrevealed chapter of geological history, one that has not been reached and read, and never can be unless some region is unearthed where the Taconic rests conformably upon the Laurentian. The Laurentian is the home of granite, marble, gneiss, and other valuable building rocks, and the best mica quarries; but the precious metals have been found only in the intrusive, altered, or sedimentary rocks of later times.

CHAPTER III.

TACONIC SYSTEM.

§ 38. In 1842, Ebenezer Emmons, in his Report on the Second Geological District of New York, described the rocks lying on the sides of the Taconic Mountains, parallel with the boundary line between New York and Vermont, under the name of the Taconic System. He found the belt on the western border of the mountains more than fifteen miles wide, and on the eastern side nearly twenty-five miles, making a total width of nearly forty miles. The rocks occur in Westchester, Columbia, Rensselaer, and Washington Counties, and stretching the whole length of Vermont, enter Canada, and extend beyond Quebec. He mentioned a typical locality in Berkshire, Massachusetts. The general character of the rocks was given as follows:

1. A coarse, granular limestone of various colors called Stockbridge limestone from the quarries at that place.
2. Granular quartz rock, generally fine-grained, in firm, tough crystalline masses of a brown color, but sometimes white, granular, and friable.
3. Magnesian slate.
4. Sparry limestone.
5. Taconic slate, which is extremely fine-grained and only slightly coherent.

He traced the rocks in a north and south course for 150 or 200 miles, and observed the fact that they underlie the Potsdam sandstone wherever it does not rest upon the gneissoid strata.

§ 39. In 1844 he published the "Taconic System," reviewed his former work, furnished numerous evidences in support of the existence of these rocks below the Potsdam and above the gneissoid rocks, or what are now known as Laurentian, and ascertained they had a thickness, as shown by a single section, of more than two miles. He said, taking one broad view of the whole system, it might be described as consisting of fine and coarse slates, with subordinate beds of chert, fine and coarse limestones, and gray, brown, and white sandstone; these admitting, however, of further divisions. The leading divisions recognized were:

1. Granular quartz, or brown sandstone, resting unconformably upon the older gneiss. It is the least regular in its continuation of any of the rocks of the Taconic System, and generally appears in insulated mountain masses, as at Oak Hill between Adams and Williamstown, Mass., at Monument Mountain, in the south part of Berkshire, in the east part of Bennington, Vt., and in Dutchess, Putnam, and Westchester Counties, New York.
2. Stockbridge limestone, generally known as Stockbridge marble, and occurring in New York, Vermont, Massachusetts, and Connecticut. Commencing at Sing Sing, it runs a northerly course through Westchester, Dutchess, and Columbia Counties, and extends into Connecticut. It passes up the valley of the Housatonic into the upper valleys of the Hoosic, and onward into Vermont, and is well represented at Williamstown, Massachusetts.
3. Magnesian slate, which composes the highest mountains in the Taconic ranges. The range of mountains composed of this slate, extending along the western

border
hundre
It cros
throug
4.
Eaton.
Renssel
Arling
5.
ated la
Countie
Vermont
above M
roofing-
ton Cou
Nereites
murchiso
6.
System,
§ 4
bridge li
stone Va
in the sl
Nereites
dinate to
with Dor
Upper H
pressions
his work
an appen
granite r
§ 41
and estab
others, m
his fossils
erent nan
knowledg
sustained
position t
him, but
Marcou, a
ability of
gan and
name, Hu
have been
Huronian
§ 42.

border of Massachusetts and through Vermont, often rising to the height of fifteen hundred feet, known as the Taconic range, furnished the name to this System. It crosses the Hudson about thirty miles above New York City, and passing south through New Jersey, enters Pennsylvania.

4. Sparry limestone, a name given to it many years before by Prof. Amos Eaton. It occupies a belt of country in the eastern part of Dutchess, Columbia, Rensselaer, and Washington Counties, and passing north strikes the west line of Arlington, Vermont.

5. Taconic slate, with its subordinate beds of roofing-slate and coarse brecciated layers, occupies almost the whole of Columbia, Rensselaer, and Washington Counties, and extends to the base of the Taconic range, which separates New York from Vermont and Massachusetts, and has an immense thickness. It crosses the Hudson above Newburg, and passes through Orange County into New Jersey. From the roofing-slate he defined *Diplograptus simplex*, and from the Taconic slate in Washington County *Bythotrephes flexuosa*, *B. rigida*, *Palæochorda marina*, *Nemapodia tenuissima*, *Nereites deweyi*, *N. gracilis*, *N. jacksoni*, *N. lanceolatus*, *N. loomisi*, *N. pugnus*, *Myrianites murchisoni*, and *M. sillimani*.

6. Black slate, forming, so far as he knew, the highest member of the Taconic System, and from which he defined *Elliptocephala asaphoides* and *Atops trilineatus*.

§ 40. He identified the Smithfield limestone in Rhode Island with the Stockbridge limestone, and an accompanying slate with the Magnesian Slate, and in Blackstone Valley found the brown sandstone and fine granular quartz. He recognized in the slates at Waterville, Maine, the Taconic Slate of New York, and found the *Nereites* at Kennebec. The fine roofing-slates on the Piscataqua he found subordinate to the Taconic Slate, in like manner as they exist in New York. And, jointly with Douglas Houghton, the Taconic System was found largely developed in the Upper Peninsula of Michigan; the slates of the formation with their fucoidal impressions and the granular quartz were both recognized. In 1846, he reproduced his work on the Taconic System in a book on the Agriculture of New York, with an appendix describing a conglomerate at the base, resting unconformably upon granite rocks.

§ 41. In this manner this geological subdivision was first determined, defined, and established, and it should have been recognized from that time forward. But others, much less informed, disputed the existence of the rocks, erroneously referred his fossils to more recent genera; and some, finding the same rocks, gave them different names, which added to the confusion, and seriously retarded the progress of knowledge respecting them. It may be later researches have not, in every respect, sustained his determinations, but Ford's work near Albany, New York, where the position taken by Emmons was most violently assaulted, has not only corroborated him, but has forever set the questions at rest in that locality. Wing, Dale, Marcou, and Dwight have sustained his assertions respecting the want of conformity of the Hudson River Slates with the Taconic. All the surveys of Michigan and Wisconsin have sustained him, though the geologists apply the later name, Huronian, to the Strata. His determinations of the rocks in North Carolina have been most fully confirmed by later geologists, though some use the word Huronian when referring to them.

§ 42. In 1849, Alexander Murray, an assistant on the Geological Survey of

Canada, in the Report of Progress for the year 1847, described the rocks on the north side of Lake Huron, and constituting many of the adjacent islands, under the name of "quartz rocks and sandstones, conglomerates, slates, and limestones," and correctly identified them as resting unconformably upon the older granite and syenitic gneiss, and succeeded unconformably by the Potsdam, but he did not call them by any geological name. If he had read Emmons's "Taconic System," it is difficult to conceive why he should have hesitated in referring the rocks to that System. In the Report of Progress of 1856, he redescribed the rocks, under the name of the "Huronian Series," which was adopted by the officers of the Canadian Survey, without once mentioning the Taconic System. From that time forward authors have generally used the name Huronian, and have almost annihilated the name Taconic. The word Taconic, however, has priority over Huronian. It is equally appropriate, and the definition of the fossils in the Upper Slates at once furnished the means of tracing it and determining it at different and distant places. The word "Huronian" is, therefore, a synonym for Taconic, and comprehended, as used originally by the Canadian Geologists, substantially the same series of rocks, though not ascending quite so high.

§ 43. A section of the so-called Huronian, but more properly called the Lower Taconic, between Missisquoi and St. Mary's Rivers, in ascending order, is as follows:

1. Gray quartzite,	500 feet.
2. Greenish, red-weathering chloritic and epidotic slates,	2,000 "
3. White quartzite, etc.,	1,000 "
4. Slate conglomerate,	1,280 "
5. Limestone,	300 "
6. Slate conglomerate, etc.,	3,000 "
7. Red quartzite, etc.,	2,300 "
8. Red jasper conglomerates, etc.,	2,150 "
9. White quartzite, etc.,	2,970 "
10. Yellowish chert, etc.,	400 "
11. White quartzite, etc.,	1,500 "
12. Yellowish chert, etc.,	200 "
13. White quartzite,	400 "
Total,	18,000 "

Another section adds to this one 4,000 feet, and even then the maximum thickness of the series in that locality has not been reached.

§ 44. Throughout the Huronian region, the whole series bears evidence of great disturbance, and is frequently cut with intrusive masses of greenstone, granite, or other igneous rocks. The more recent disturbances frequently bear metalliferous veins, which give to the country its value as a mineral region. Copper and iron are the chief minerals, and abound in nearly every section. Gold and silver sometimes occur. The Taconic of Michigan contains vast beds of iron ore. The ores are magnetic, red specular hematite and soft hematite resembling the brown hematite of other States. The magnetic and specular ores are the most prized, and usually contain from 60 to 70 per cent of iron, and hardly a trace of phosphorus or sulphur. (Phosphorus makes iron brittle when it is cold, and is therefore called cold-short, though it is malleable when hot, while sulphur makes it brittle when it is hot, and it is therefore called red-short.) The Lake Superior region is the chief locality of the world for native copper. It is so pure the aborigines manufactured it into implements. The copper-bearing rocks extend eastward

along the south shore of the lake for more than forty miles, then forming a narrow belt stretch in a north-east direction for about a hundred miles to the extremity of Keweenaw Point. The copper occurs in a rock called melaphyre, associated with beds of conglomerate, and appears to be interstratified with them. Sometimes bands of slate separate beds of melaphyre. The native copper exists in sheets, strings, and masses, and is sometimes associated with silver. In Ashland County, Wisconsin, the copper-bearing series has a thickness of more than four miles, though not very rich in the ore. The Taconic area in Minnesota is large. It extends across the northern border, and forming an elbow in the north-east extends diagonally through the State to the south-west corner. Here there is a hard, reddish, metamorphic sandstone, called the Sioux quartzite, interstratified with which is a layer of red indurated clay or pipestone, one foot thick, called Catlinite, largely used for the manufacture of pipes. The quarry is thirty miles north of the south-west corner of the State, and four miles east of the west line. The Sioux quartzite occurs in the north-west corner of Iowa.

§ 45. The geographical extent of these rocks in Canada is very great. They may be traced from near Lake Temiscaming 80 miles north-west of Lake Nipissing, south-westward to Lake Huron, and from thence westward on the north shore of the lake and the north shore of Lake Superior, and on beyond Lake of the Woods, a distance in all of about 800 miles. They pass beneath the lakes and expose a large area in the Upper Peninsula of Michigan at Marquette and Menominee, and a great thickness, extending from the lowest to the highest Taconic, as first ascertained by Houghton; thence they pass into Wisconsin, exposing a large area and quite as complete a representation of the series, while another arm extends from Duluth into Minnesota. The thickness in Michigan is about four miles; but in Wisconsin, including the copper-bearing series, which is three-fourths of igneous material, the thickness is much greater; and even excluding the igneous material the thickness exceeds four miles. The upper part of the Taconic System in Wisconsin, formerly called the "Copper-bearing series," has received the unattractive name of the Keweenawan formation, from the Keweenaw Point; but as it is part of the Taconic System the preferable name is the older one of the "Copper-bearing series." The rocks appear between Scoresby Bay and Cape Cresswell, in Lat. 82° 40' N., where Nares and Feilden called them Cape Rawson beds.

§ 46. In 1856 Emmons divided the System into Upper and Lower Taconic. The Canadian Geologists in 1863 placed his Upper Taconic in the Silurian System and called it "Lower Potsdam," which name therefore became a synonym. The only geographical names which have been used to subdivide the Upper Taconic into Groups, which seem in the present state of learning to be worthy of retention, are, in descending order, the Swanton Group, the Georgia Group, and the St. John Group—if in fact the latter is below the Georgia, and therefore not a synonym. Emmons placed the Stockbridge limestone in the Lower Taconic; but it would seem from the examinations made by others, that his division would have been more clearly marked if the Stockbridge limestone had been retained in the Upper Taconic. The Paradoxides beds at Braintree, Mass., in Newfoundland and New Brunswick, and wherever found on the continent, belong to the Upper Taconic. The same difficulty exists in the West, in separating the Upper Taconic from the overlying rocks of the Potsdam, that has led to so much discussion in the East; and the confusion is

increased by the addition of numerous synonyms—the ready weapon to which ignorance resorts.

§ 47. In 1863 G. F. Matthew named the rocks exposed at St. John, New Brunswick, the “St. John Group.” He described them as arenaceous, argillaceous, and carbonaceous shales, and clay slates; often sandy, with sandstone and quartzite, having a thickness of 4,500 feet, and having an exposure about 30 miles long and 4 miles wide. He collected *Paradoxides*, *Conocoryphe*, *Obolella*, *Orthis*, *Orthisina*, *Stenotheca*, *Hyolithes*, and *Lingula*. In 1865 he and Bailey and Hartt correlated these rocks with the slates of Vermont having *Elliptocephala asaphoides*, and the schistose beds at Braintree, Mass., holding *Paradoxides harlani*, and thus proved their “St. John Group” to be a synonym for Emmons’s “Black Slate,” in the Upper Taconic System. Furthermore, they identified the slates with some found in Newfoundland containing *Paradoxides* and *Conocoryphe*. Later they divided the Lower Taconic of New Brunswick, which they called Huronian, into the “Coldbrook Group,” the “Coastal Group,” and the “Kingston Group,” and estimated the thickness as exceeding 10,000 feet.

§ 48. The Vermont Geologists in 1861 called the Black Slate, Taconic Slate, and Roofing-slate of Emmons the “Georgia Group.” The name has priority over the “St. John Group,” and if the Taconic System is to be divided into Groups with geographical names, and these three divisions of Emmons are to be thrown together in one Group, then they must under the laws of nomenclature bear the name of the Georgia Group. The Black Slate has, however, been called the Swanton Group, and if this name should become desirable then the Upper Taconic would be divided into the Swanton Group and the Georgia Group, and their maximum thickness in Vermont exceeds two miles. This division is that adopted by Perry, who has shown the Potsdam sandstone rests directly upon the Swanton Group, or Black Slate, as originally asserted by Emmons, and that both the Swanton Group and the Georgia Group are fossiliferous.

§ 49. The Taconic rocks extend from Canada East and Maine to Georgia and Alabama, flanking almost continuously the ranges of mountains upon both the eastern and western slopes. Their thickness in New Hampshire is over four miles, and in Vermont the maximum must exceed five miles. The slate belts of York and Lancaster Counties, Pa., and the rocks containing the valuable ores of nickel and copper belong to this System. There are five extensive outcrops in North Carolina, and three or four subordinate ones. They rest unconformably upon the belts of the exposed Laurentian, and very much resemble in their character the subdivisions in Vermont and New York. The largest outcrop is from twenty to forty miles wide, and extends quite across the State. The maximum thickness exceeds five miles. There are large outcrops in Virginia, South Carolina, Georgia, and Tennessee, and limited outcrops in Alabama. Gold, silver, copper, lead, iron, and other valuable minerals, occur in these rocks not only in veins, fissures, and dykes, but in seams following the stratification and as part of the sedimentary materials. In Northern Georgia gold exists in seams, with milky quartz, following the stratification of hornblende schists, and constituting as truly sedimentary rocks as the schists themselves do. The seams are stratified within the slaty sediments, and are of the same age as the Taconic System. These seams are so constant they characterize the slates or schists in the Appalachian System. They are metalliferous,

and frequent also occurring partially in iron ore is of this very thin ore, containing and bearing numerous often exposed usually

§ 50. pass up in *coryphe*, *A. luthellus*, *A.* rocks, but closed the from this to live to velopen single spe separates determina formability, and the c of the dis sents the in this sys bers (if th Group bel author bel first order become ex very likely

§ 51. Keweenaw whole basin face area 18,000 square with an est feet for the last ha feet. The sive dykes. sunk beneath representin

and frequently auriferous, or cupriferous. The magnetic and specular iron ores also occur with the material of the slates as a deposit of the same age, and constituting part of the same system. This mineral wealth is so distributed it is practically inexhaustible. The Taconic appears in Missouri, Arkansas, and Texas. The iron ore district about Iron Mountain and Pilot Knob containing porphyry rocks is of this age, but the granite to the east is Laurentian. The ore is found in very thick veins in Iron and Shepard Mountain, and Pilot Knob. It is specular ore, containing between sixty and seventy-five per cent of iron, free from sulphur and bearing no more than a mere trace of phosphorus. The rocks appear in numerous places in the Rocky Mountain ranges from Mexico to British Columbia, often exposing great geographical areas and an immense thickness, and they are usually metalliferous.

§ 50. The genera, regarded as typical of the Taconic fauna, and which do not pass up into Silurian rocks, are *Paradoxides*, *Microdiscus*, *Atops*, *Elliptocephala*, *Conocoryphe*, *Anopolenus*, *Bathynotus*, *Solenopleura*, *Acrothele*, *Salterella*, *Scenella*, *Iphidea*, *Hypolithellus*, *Archæocyathus*, and *Ethmophyllum*. There are some others peculiar to these rocks, but they are either obscure or limited in their distribution. Some genera closed their existence in Silurian time, others reached the Devonian age, and some from this remote period, as *Orthis*, *Orthisina*, *Orthoceras*, and *Leperditia*, continued to live to the Carboniferous, though *Orthoceras* reached its most remarkable development in the Black River Group, and *Orthis* in the Hudson River. Not a single species belonging to the Upper Taconic system crossed over the line that separates it from the Potsdam Group of the Lower Silurian, so far as any reliable determination has thus far been made. This, supported as it is by a want of conformability, indicates a vast lapse of time between the deposit of the Upper Taconic and the commencement of the Potsdam period. The Taconic is composed in part of the disintegrated materials of prior Laurentian rocks, while the Potsdam represents the washings of the Laurentian and Taconic. The order Graptolida appeared in this system, and reached its maximum development of genera, species, and numbers (if the Point Levis beds referred by the Canadian Geologists to the Quebec Group belong to the Upper Taconic, as claimed by Marcou and others, and as the author believes), and became extinct in the Upper Silurian System. This is the first order of animal life to reach the highest stage of its existence, and the first to become extinct. It is referred to the class Hydrozoa, but if more was known of it, very likely it would form a distinct class.

§ 51. The Cupriferous series of the lake region, called also the Keweenaw, Keweenawian, Keweenawan, and Nipigon series, is supposed to underlie nearly the whole basin of Lake Superior, or an area of about 28,000 square miles, and a surface area upon the borders of the lakes and their immediate vicinity of about 18,000 square miles. This series has been divided into an upper and lower division, with an estimated maximum thickness of 15,000 feet for the upper division, and 35,000 feet for the lower, which rests upon the slates and quartzites of the Taconic System, the last having a variable thickness that reaches a maximum of at least 22,000 feet. The Cupriferous series consists of eruptive flows and detrital rocks, with massive dykes. The region was, in Taconic days, represented by a volcano, which has sunk beneath the waters of the lake. The flows were followed by detrital rocks, representing the intervals of time between them; but these detrital rocks are com-

posed largely of conglomerate layers and large-sized pebbles, indicating strong currents of water. The flows visible upon the borders of the lakes were forced through fissures by volcanic energies. The copper which occurs in the conglomerates, amygdaloids, epidote veins, and otherwise, is supposed to have been precipitated from water holding it in solution, or leached from detrital rocks where it was originally deposited in a sulphureted form. R. D. Irving, who has studied closely the copper-bearing rocks of this region, says the explorer for transverse veins should bear in mind that epidote, prehnite, and chlorite are favorite associates of copper, while laumontic veins, and those bearing a predominating quantity of calcite, are not so rich; that a wide vein in amygdaloidal or other soft rock will pinch to a mere seam within the massive and compact layers; and in sandstone and conglomerate deposits the valuable belts have been found where the conglomerate is overlaid with trap, or in sandstone very rich in basic detritus. Any of the conglomerate seams from Keweenaw Point to Minnesota may be cupriferous. All of the upper division of the series is noncupriferous, except the Nonesuch sandstone belt in the Porcupine Mountains; and all the belts and areas of acid rocks, such as the central area of the Porcupine Mountains, and the great spread of red rock in the Brulé Lake country in Minnesota, and all belts and areas of coarse-grained basic rocks, such as the great area of coarse gabbro in the Bad River region in Wisconsin, and the similar area which occupies the belt of country from Duluth to Brulé Lake, are also noncupriferous. The slates and quartzites of the Taconic System which lie below the Cupriferous series on the north shore of Lake Superior, have been called the Animikie Group. About three-fourths of the great thickness of the rocks is referred to volcanic overflows, and does not, therefore, belong to the geological column, the whole of which is the result of sedimentary deposition.

CHAPTER IV

SILURIAN SYSTEM.

§ 52. IN 1833, Sir R. I. Murchison, in a memoir read before the Geological Society of London, divided the fossiliferous rocks below the Devonian into six Groups. He founded this subdivision upon the fossils, and mentioned such species as were then defined. This was followed in the succeeding year by other memoirs, and in 1835 he concluded all these Groups might be placed in one System, and in honor of the ancient tribe of Silures, who inhabited Wales, he named it the Silurian. He placed three Groups in the Upper Silurian, and three in the Lower Silurian. Before this time no knowledge of the order of the strata had been ascertained, and hence he is entitled to the credit of the name. Subsequently palæontologists found his discoveries were world-wide in their application, and it was not long until the distinction between the Upper and the Lower Silurian had been observed in North America, and the base of the Lower Silurian had been recognized in the Potsdam sandstone, and the base of the Upper Silurian in the Medina sandstone.

§ 53. Some years later, the word Cambrian was applied in England to the rocks which belong to the Lower Silurian, and to inferior strata which are the equiva-

lent in part but in the was never breaks in g No careful strata, tho times we se

§ 54. the followi Trenton, Ut ing order, Helderberg

§ 55. I of New Yor and propose described qu gists in 186 is a granitic ure, are som veloped in th composition compact lay slaty sandsto Calciferous, gradual into 200 feet. T to Lake Cha

§ 56. It ranging from with magnes more diversit sometimes oc and fills up rests unconfo to have been and wind ma Lake Huron variations in again thins o Canada, the which the tra ripple-marked marks, and t as to show th dant in the

lent in part of the Taconic System. It was not used in the sense of a "Group," but in the higher sense of a "System," as these words were then understood. It was never well defined, and it crossed one of the grandest and most important breaks in geological time—that which separates the Taconic and Silurian Systems. No careful geologist or palæontologist uses the word in the nomenclature of American strata, though it occasionally occurs in incoherent geological papers, and sometimes we see such monsters in nomenclature as Cambro-Silurian and Siluro-Cambrian.

§ 54. The Lower Silurian in North America is divided, in ascending order, into the following Groups: viz., Potsdam, Calciferous, Quebec, Chazy, Black River, Trenton, Utica Slate, and Hudson River. The Upper Silurian is divided, in ascending order, into the Medina, Clinton, Niagara, Onondaga, Guelph, and Lower Helderberg.

POTSDAM GROUP.

§ 55. Prof. Ebenezer Emmons, in the Annual Report of the Geological Survey of New York for 1838, described the sandstone at Potsdam in St. Lawrence County, and proposed for it the designation "Potsdam Sandstone." It was subsequently described quite fully in the New York Reports, and finally the Canadian Geologists in 1863 called the rocks the Potsdam Group. The lowest portion at Potsdam is a granitic conglomerate, in which large masses of quartz, the size of a peck measure, are sometimes inclosed. These were water-worn and rounded before being enveloped in the deposit. The sandstone is quite variable in texture and color, but its composition is uniformly silicious. At some places it is an even-grained mass in compact layers, and at others it is traversed by joints. In some localities a dark, slaty sandstone, about ten feet in thickness, intervenes between the Potsdam and Calciferous, at others a coarse brecciated rock, and at others the passage is very gradual into the Calciferous sand-rock. The thickness in New York is from 100 to 200 feet. The exposure is narrow, but extends from near the Thousand Islands to Lake Champlain, and enters Vermont with a thickness of about fifty feet.

§ 56. It extends from New York into Canada, where it attains a thickness ranging from 300 to 700 feet, and at the summit the sandstone is interstratified with magnesian limestone that constitutes a passage to the Calciferous. There is more diversity in the rocks in Canada than in New York, and limestones and slate sometimes occur with the sandstone. It rests unconformably upon the Laurentian, and fills up inequalities where the Taconic System does not intervene, and it also rests unconformably upon the Taconic when it is present. The sandstone appears to have been deposited in shallow water along the margin of a sea. The tracks and wind marks support that view. In its extension westerly, by the way of Lake Huron and Lake Superior, across Wisconsin and into Minnesota, the same variations in thickness occur. Sometimes it attains a thickness of 3,000 feet, and again thins out to 40 or 50 feet. For several miles in distance near Beauharnois, Canada, the strata are marked by the tracks of *Protichnites*. The surfaces on which the tracks are impressed are sometimes smooth, and sometimes beautifully ripple-marked. On the latter the tracks have often beaten down the ripple-marks, and the sand of the ridge has been dragged into the furrow, in such a way as to show the direction in which the animal was progressing. Fucoids are abundant in the upper part of the Group, and *Scolithus* so common as to be quite

characteristic, and near St. Genevieve the rock is completely honey-combed with it to the depth of three feet.

§ 57. It is largely distributed in Northern Michigan, and striking into Wisconsin north of Green Bay gradually widens southerly as a surface rock, until it reaches the central part of the State, where it has a width of 100 miles. It then curves northwardly and enters Minnesota, forming the high hills on the Mississippi River. It is unconformable with the rocks below, and rests upon an exceedingly irregular surface, sometimes filling depressions in the quartzite or metamorphic rocks of several hundred feet. Its upper surface is uniform, and graduates into the Calciferous Group or the Lower Magnesian limestone, as the rocks in these States are called. The exposed area in Wisconsin is about 12,000 square miles, the thickness very irregular by reason of the great depressions and elevations at the base, and the maximum thickness is fully 1,000 feet. The rock is chiefly composed of cemented grains of silicious sand, but presents several varieties, as the calcareous, argillaceous, ferruginous, and green sand, and the waters issuing from it in places contain a small percentage of lime salts. In the argillaceous class the clayey material becomes so abundant as to render the rock shaly, and so impervious to water that valuable springs occur at its upper exposed surface. In the calcareous class the lime becomes so great in some layers that they are more properly limestones than sandstones, and so associated with magnesia that they become arenaceous dolomites. In the ferruginous class, at one extreme, the amount of iron oxide is barely sufficient to color or cement the mass, and at the other so great as to make an iron ore. In the green sand there are two classes, one in which the grains are colored by iron, and the other consisting of deep green grains of glauconite. The green sand is not restricted to the Potsdam in Wisconsin, for it also occurs in the Calciferous and St. Peter's Sandstone. It is almost identical with the Cretaceous green sand of New Jersey, and similar deposits in existing strata. The surface area in Michigan, Wisconsin, Iowa, and Minnesota is estimated at 25,000 square miles, which is about half the surface area on the continent; but it is generally believed to exist under many of the more recent deposits, and, therefore, to cover several hundred thousand square miles. Springs and streams of soft water are abundant where it forms the surface rock, and a good supply of soft water has been found wherever it has been penetrated with the drill; its existence, therefore, becomes a question of much economical interest where a supply of good water is desired from artesian boring. The drill has never reached it in Ohio, though a supply of good water is imperatively demanded in some parts of the State; and it is to be hoped an effort will be made to determine whether it exists below the Calciferous, which has been reached with the drill many times.

§ 58. It is exposed at numerous places in the Appalachian System from New York to Tennessee. In New Jersey it reaches a thickness of 3,000 feet, and if both the Chilhowee sandstone and Knox Group in Tennessee belong to it, it has a thickness of 9,000 feet, but probably 5,000 feet of this belongs to the Taconic. In the southern and south-eastern counties of Missouri it has a thickness of 700 feet. It appears in several counties in Northern Texas, along the margins of the Big Horn, Laramie, and Wind River ranges, at the Black Hills, and in other regions of the Rocky Mountain System from Mexico to British America. The erosion by water and weathering has left picturesque scenery in the sandstone at many places. The

" Pictured
" Chasm o
§ 59.

from the a
and schists
tains ripple
evidence sh
not have o
climate was
ence of lan
prevailed o
Dicellocypha
somewhat c
slowly depo
ity, indicati
is no doubt
time the sar
dam age ha

§ 60. T

lime and fin
contains calc
taining beds
defined by V
He united th
fucoidal laye
cious, magn
They pass fr
ties lined wi
sessed of a s
Middleville
contain a flu
structure of
structure, as
Counties.

At Chaz

1. Silice
2. Lime
3. Dull,
4. Cystic

15 feet.

"Pictured Rocks" of Lake Superior, the "Dalles" of the Wisconsin, and the "Chasm of the Au Sable" in New York are examples.

§ 59. Everywhere it is essentially an accumulation of sandstone and pebbles from the adjacent Laurentian gneisses, granites and syenites, and Taconic quartzites and schists, resulting from the disintegrating influences of air and water. It contains ripple-marks, wave-lines, mud cracks, animal tracks, and worm burrows, which evidence shallow seas and shore lines. The continent at the time of its deposit did not have one twentieth its present area. There is nothing known to indicate the climate was different then from what it is now, except so far as the relative difference of land and water surface would necessarily change it. Some species of fossils prevailed over great areas, as *Hyalithes primordialis*, *Lingulepis pinniformis*, *L. prima*, *Dicelloccephalus minnesotensis*, *D. osceola*, and *Ptychaspis minuta*, and therefore become somewhat characteristic of the Group. Though composed almost wholly of sand it was slowly deposited. The sandstone is frequently charged with fossils to its full capacity, indicating a formation almost as slow as marine limestone is now made. There is no doubt that Calcareous mud was forming in the depths of the ocean at the same time the sand was deposited nearer the shore, but no limestone group of the Potsdam age has been found, unless it exists in the Eureka district of Nevada.

CHAPTER V.

CALCIFEROUS GROUP.

§ 60. This name was first applied by Prof. Eaton to a gray rock consisting of lime and fine grains of sand, so intimately blended as to appear homogeneous. It contains calcite and a sparkling surface, but passes into a carbonate of lime, containing beds of magnesian limestone and a small amount of iron. The Group was defined by Vanuxem in 1842, in the Geology of the Third District of New York. He united the silicious layers above the Potsdam, the calciferous sand-rock, and the fucoidal layers in one Group. The rocks consist in general of three varieties—silicious, magnesian, and carbonate of lime, with intermediate grades of composition. They pass from compact to granular, and granular to porous, the latter having cavities lined with crystals of quartz, calcareous spar; or, instead of being lined, possessed of a single beautiful perfect crystal of limpid quartz, nearly filling the space. Middleville and Little Falls are noted localities for these crystals, some of which contain a fluid or anthracite, which enhances their value as cabinet specimens. The structure of the rock is often oolitic, passing into thick layers having a concretionary structure, as in agate. The typical localities are in Montgomery and Herkimer Counties.

At Chazy the following ascending section occurs:

1. Silico-calcareous beds, more or less interspersed with sparry masses, 30 to 35 feet; fossils rare and cherty.
2. Limestone, in which the plates of Cystideans abound, 20 feet.
3. Dull, gray, earthy mass, without fossils, and passing into oolitic beds, 10 feet.
4. Cystidean limestone, similar to the first though of a brighter red color, 15 feet.

5. Massive earthy and silicious limestone containing trilobites, 20 feet, followed by beds of similar character of greater thickness containing brachiopods.

6. Red Cystidean limestone, susceptible of a fine polish, 15 feet.

7. Drab-colored, thin-bedded, earthy magnesian beds, suitable for hydraulic lime, of considerable thickness; fossils rare, except fucoids. Toward the top of the rock it is blue and frequently cherty, oolitic, and concretionary, the upper masses from 20 to 30 feet thick.

§ 61. The Group is persistent, and surrounds the irregular dome of Laurentian rocks, which form the northern highlands of New York, in a belt, overlying the Potsdam. It is chiefly a hard calcareous sandstone or arenaceous limestone, resting upon the margin of the Potsdam sandstone, from Lake Ontario eastwardly to Vermont, and from New Jersey north, near the line of New York and Vermont, into Canada. It forms a narrow belt of surface exposure, with a variable thickness from 50 to 350 feet. Lake Champlain has cut a channel through it for twenty miles. In Canada, adjacent to New York and Vermont, it is, in the lower part, a dark, bluish-gray, crystalline, strongly coherent dolomite or magnesian limestone, and in the upper part a bluish-gray, calcareous argillite, but its characters are different in different localities. It is usually found as a narrow belt following the sinuosities of the Potsdam Sandstone, from west of Lake of the Woods to the Atlantic sea-board; but where the rocks have been disturbed by volcanic energies it may be absent or difficult of detection. The surface area of its distribution in Canada is several thousand square miles, and in its undisturbed condition the maximum thickness rarely exceeds 450 feet, though in Newfoundland, where it is a definitely stratified limestone, it has a thickness of more than 2,000 feet. In the region of the Mingan Islands, in the Gulf of St. Lawrence, the fossil casts and shells are in a good state of preservation.

§ 62. In Northern Michigan and on the Menominee and Escanaba it preserves its New York characters in a remarkable degree, although its thickness may not exceed 50 feet. The upper portions are highly calcareous, and on fresh fracture show the peculiar granular structure so characteristic in New York. It is thin-bedded, and contains small cavities lined with crystals of calc-spar, quartz, or hornblende. The surfaces of the layers are often covered with fucoidal impressions. From St. Mary's River westerly to the Wisconsin and the Mississippi there is a gradual augmentation in the thickness of the rocks and a material change in their composition. The Group enters Wisconsin from Michigan a few miles from Green Bay, and striking south-west upon the border of the Potsdam it forms a serrated margin from five to fifteen miles in width, until it reaches the streams that flow into the Mississippi in the south-western part of the State, where it is exposed upon some of the streams for a distance of 75 or 100 miles. It crosses the Mississippi and the north-eastern corner of Iowa, appearing in the bluffs and hills more conspicuous than the Potsdam, though not so thick, because it is a much firmer rock. It is a buff-colored dolomite, without uniformity of texture or stratification, and weathers into rough, bold, and often picturesque fronts along the valleys, and has a thickness of about 250 feet.

§ 63. From Iowa the area of exposure is a little west of north through Minnesota, reaching as far west as the second tier of counties from the Mississippi and following the bluffs with limited outcrops in Wisconsin to Lake Pepin, north of

which and Wisconsin. It is covered by the glacial drift, cropping out only at the mouth of the Wisconsin River. In Wisconsin, Iowa, and Minnesota, the surface is uneven, and the distances in the sea. These beds have elapsed before the present time, frequently called it the Laurentian, saying that in the Laurentian to a serviceable sand. The Laurentian talline formation is abundant, and the amount of the Laurentian face rocks to be weathered by weathering.

§ 64. The

Missouri, with an intermediate order, of the "Third Magnesian" earthy magnesian, with occasional cellular, silicious, Osage and it is often a few feet, though brownish sandiness, though the sandiness rarely exceeds of intercalated bedded, coarse chert-beds. It is frequently high. The craves, or dissections, times in many layers of limestone other places heavy spar (sulfate of lime) seldom ex-

which and east of the St. Croix it forms the surface of nearly two large counties of Wisconsin. It follows the Mississippi north of Minneapolis for several miles before it is covered with later formations. The conspicuous perpendicular walls of rock, cropping out from the hills and bluffs along the Mississippi from the St. Croix to the mouth of the Wisconsin, belong to this Group. Throughout the exposures in Wisconsin, Iowa, and Minnesota, it is conformable with the underlying Potsdam, and unconformable with the overlying rocks. The lower surface is plane, while the upper surface is undulating, and in some instances the undulations are said to swell in short distances into elliptical domes, rising 100 feet above their bases, like billows on the sea. These undulations are the work of denudation during the interval that elapsed before the deposition of superimposed strata. The Group in Wisconsin is frequently called the Lower Magnesian limestone, and some one in Minnesota has called it the Shakopee Group, because the stone has been quarried at a village bearing that Indian name. The Magnesian limestone is usually sufficiently pure to burn to a serviceable quicklime. The chief impurities are quartz, clay, iron, and green sand. The dolomite occurs in the earthy, granular, crystalline, and crypto-crystalline forms, and chert is irregularly distributed. Argillaceous material is not abundant, except in shaly bands, where it may constitute 20 per cent of the whole; and the amount of silica disseminated through the rock varies from 1 to 10 per cent. The difference in the composition and hardness of the layers causes the surface rocks to present great irregularities, which are much enhanced and exaggerated by weathering, and hence outliers have a rough and often grotesque exterior.

§ 64. The Group is displayed in grand proportions in the southern counties of Missouri, where it consists of an upper and lower division of magnesian limestone with an intermediate division of sandstone. These received the names, in descending order, of the "Second Magnesian limestone," the "Second Sandstone," and the "Third Magnesian limestone." The upper division is generally composed of beds of earthy magnesian limestone, interstratified with shale-beds and layers of white chert, with occasionally thin beds of white sandstone, and near the lower part thick, cellular, silico-magnesian limestone-beds. It constitutes many of the bluffs of the Osage and its tributaries, and also of the Missouri from Osage to Jefferson City. It is often a lead-bearing rock, as in Cole County. The thickness rarely exceeds 200 feet, though on the Meramec it is 300 feet. The middle division is usually a brownish sandstone, stratified in firm, regular beds from 2 inches to 3 feet in thickness, though sometimes friable. The surfaces are often ripple-marked. The thickness rarely exceeds 150 feet. The upper part often occurs in thin strata with beds of intercalated chert abounding in fossils. The third division is generally a thick-bedded, coarsely crystalline, bluish-gray magnesian limestone, with occasional thick chert-beds. It is the chief lead-bearing rock of South-east and Southern Missouri, and is frequently exposed along the streams in bold escarpments from 200 to 300 feet high. The ores of lead, zinc, copper, nickel, and cobalt, occur in fissures and caves, or disseminated in small masses in the limestone itself. The lead occurs sometimes in masses of galena accompanied with copper pyrites disseminated through layers of limestone, while the ores of nickel and cobalt occur in clay slate. At other places bands of red clay inclose calamine (silicate of zinc), galena, and heavy spar (sulphate of baryta). The maximum thickness is about 600 feet, though it seldom exceeds 300 feet. The maximum thickness of the three divisions is more

than 1,000 feet, but the Group at no single locality displays so great a thickness. From Missouri the Group extends southerly across Arkansas into San Saba, Llano, McCulloch, Menard, Mason, and Lampasas Counties, in Texas, where the maximum thickness is more than 400 feet. It is exposed in narrow belts in the Appalachian chain from New York and New Jersey to Tennessee and Georgia, but has not been very clearly distinguished in the mountain regions of the West.

§ 65. It is said this Group in some localities graduates into the Quebec; but on the other hand it is claimed the Quebec belongs to the Taconic System, and is below the Potsdam. It is certain many of the rocks referred to the Quebec Group belong to the Taconic, and some of them may belong to the Calciferous or the Chazy, or may form passage beds from one to the other. The oldest known Lamellibranchiata are found in this Group. Among the fossils having the greatest distribution, and which are most characteristic, we may mention *Ophileta complanata*, *O. uniaugularis*, *Holopea turgida*, *H. dilicula*, and *Orthoceras prinigenium*. *Pleurotomaria canadensis* and *Leptena barabuenensis* occur in this Group and in the Potsdam. *Pleurotomaria calcifera*, *P. postumia*, *Holopea dilicula*, *Helicotoma perstriata*, *Maclurea matutina*, *M. sordida*, *Eccyliomphalus canadensis*, *Camarella calcifera*, *Lingulella mantelli*, *L. irene*, *Amphion salteri*, *Bathyurus cordai*, *B. conicus*, and *Asaphus canalis* have been described from this Group and from the Quebec. These identifications may well be doubted, unless the rocks containing all these species really belong to the Calciferous.

CHAPTER VI.

QUEBEC GROUP.

§ 66. THE Quebec Group was first characterized and its position between the Calciferous and Chazy asserted, upon palæontological evidence, in 1862, by Prof. Billings. His position was supported by the Canadian Geology in 1863, and in Decade 2 of a later date. The limits of the Group are still a subject of discussion, and part of the rocks originally referred to it belong to the Upper Taconic; but another part of them may form passage beds from the Calciferous to the Chazy, and occupy a position which warrants the name of an independent Group. The name was derived from the city of Quebec, where it was subdivided into the Levis, Lauzon, and Sillery divisions. The Levis was named from Point Levis, where it is fossiliferous, and has a thickness of 6,145 feet; the Lauzon from Lauzon, where its thickness is 4,000 feet, and it is non-fossiliferous; and the Sillery from Sillery Cove, where it is 2,000 feet thick and almost barren of fossils. The Lauzon division is below the Levis, and, from fragments of fossils found in pebbles, it has been since ascertained that it belongs to the Upper Taconic. A great fault at the Island of New Orleans and another near the Falls of Montmorency, with lesser faults, are said to account for the erroneous reference of this division to the Quebec. It was supposed at one time that the Sillery and Lauzon were upper members of the Quebec Group; but both of them belong to the Upper Taconic.

§ 67. The Group has been traced from Vermont to Newfoundland, a distance of 1,000 miles, and the Levis division noted at several localities, where it consists of a variety of shales, with some sandstones and conglomerates, distinguished by

the general graptolitic shales, 1,000 feet of black slates, division. Some have undiscovered and therefore be an erroneous may be found the trilobites primordial examine the of the learned should be re undoubtedly East from w that which h Lower Silurian in the Calciferous the rocks re belong to th

§ 68. The Pogonip where the Canadian mountain beds of the terminated bed *Acrotreta gem* *C. unisulcatus*

§ 69. In animal kingdom, of the Upper Group, and of these forms distinguished Group is said and *Asaphus* mon to all th

§ 70. The lines of passage-beds occur conformability geographical System, and vast period of animals, and

the general black or dark color. In Newfoundland the Levis division consists of graptolitic shales, having a thickness of 4,000 feet, which are followed by about 1,000 feet of serpentines and diorites referred to the Lauzon division; and these by black slates and limestones, having a thickness of 4,000 feet, referred to the Sillery division. Serpentines, diorites, and slates sound like Upper Taconic, and it may be undiscovered faults have given rise to an erroneous determination of the order, and therefore the so-called Lauzon and Sillery may be below the Levis; or it may be an erroneous identification of the Lauzon and Sillery; and yet the true solution may be found in the fact that all three divisions belong to the Upper Taconic, for the trilobites described by Billings, from these rocks in Newfoundland, have a primordial or Taconic aspect. The author has never had an opportunity to examine the rocks of the Quebec Group, but an examination of the present state of the learning respecting it, makes it very doubtful whether or not the name should be retained. If the Group belongs to the Taconic System, as most of it undoubtedly does, possibly the name should be retained. If that part of it in the East from which Calciferous fossils have been obtained, constitutes all of it except that which belongs to the Taconic, then probably the name should be stricken from Lower Silurian nomenclature, and the part containing such fossils should be included in the Calciferous Group, in which event the Chazy Group would include some of the rocks referred to the Quebec in the Western mountains, and the rest would belong to the Upper Taconic.

§ 68. The Quebec Group has been recognized in the Wahsatch Range, in Utah, at Pogonip Mountain, Nevada, and other places in the Western mountain chains, where the Calciferous and Chazy have not been distinguished from it. In the Pogonip mountain-beds the following species are said to pass from clearly distinguished beds of the Potsdam Group up three or four thousand feet into as certainly determined beds of the Quebec Group, viz.: *Lingulepis maera*, *L. minuta*, *L. manticula*, *Aerotreta gemma*, *Agnostus communis*, *A. bidens*, *A. neon*, *Crepicephalus haquei*, and *C. unisulcatus*.

§ 69. In this Group we find the first illustration of an important branch of the animal kingdom reaching its highest stage of development, and subsequently declining, and finally becoming extinct. The first known *Graptolites* appear in slates of the Upper Taconic System, and reach the climax of evolution in the Quebec Group, and become extinct in the Upper Silurian era. The development of these forms seems to have been wonderful. About thirty genera have been distinguished in America, and to these have been referred about 170 species. The Group is said to be connected specifically with higher Groups by *Maclurea atlantica* and *Asaphus canalis*, that occur in the Chazy, and by *Leptena sericea*, which is common to all the Groups in some of its varietal forms as high as the Clinton.

§ 70. This Group is said to graduate up into the Chazy without lithological lines of separation, and without an abrupt break in the chain of fossils. Clear passage-beds occur where the Groups are well developed, and even where there is non-conformability some fossil species are said to be common to the two Groups. The geographical surface distribution is confined to limited areas east of the Appalachian System, and to small exposures among the Western chains; but it must represent a vast period of time, as evidenced by the great development and evolution of its animals, and by the erosion of the Calciferous where it does not exist.

§ 71. Bitumen, or mineral pitch, is a product resulting from the distillation of vegetable and animal matter within the earth. It has a pitch-like odor, and burns with a bright flame without any ash, and varies from liquid naphtha to solid asphaltum. Naphtha is a nearly colorless fluid, having a pungent smell, that issues from the rocks in Persia. Its specific gravity is about 7-10, and by exposure it loses its transparency and odor, and acquires a yellowish or brown color, becomes thicker and heavier, and approaches petroleum. Petroleum is so called from exuding as an oil from the rocks. Its specific gravity is 87-100, and by exposure to the air and the application of heat it may be converted into asphaltum. Asphaltum was so named from a lake in Judea, where it rises in a liquid form to the surface of the water and then hardens. Its specific gravity varies from 1.07 to 1.65. It is quite brittle and electric, though coal is not. Bituminous matter occurs in the limestones and dolomites of the Quebec Group, and the odor may be detected in many places by striking or heating the rocks. A black, combustible, coal-like matter is found with crystals of bitter spar and quartz, sometimes coating the crystals or the walls of cavities, and at other times in the form of buttons or drops, evidently having been introduced in a liquid state and subsequently hardened. It fills veins and fissures in limestones, shales, and sandstones, and even in the trap-rocks which traverse these. It is very pulverulent, brittle, of a shining black color, and yields from ten to twenty per cent of volatile matter. It approaches anthracite in its characters. The volatile matter is a hydrocarbon gas. It has resulted from the slow alteration of liquid bitumen in the fissures of the strata. The bitumen was derived from marine vegetation or marine animals, which underwent a special mineralization, producing the bituminous matter instead of coal. It is due to chemical reactions, by which it retained a greater proportion of hydrogen in its combination than would have been retained if it had been converted into coal.

CHAPTER VII.

CHAZY GROUP.

§ 72. THE Chazy Group was first defined in the Report of the Second District of New York in 1842, by Prof. Emmons, under the name of the Chazy limestone. The name was derived from the town of Chazy, where it has a thickness of 130 feet, reposes unconformably upon the Calciferous, and is succeeded by the Birdseye limestone. It is a dark, irregular, thick-bedded limestone, containing many rough, flinty, or cherty masses, and extends as a belt into Vermont, where it exposes more surface area than any other Group of the Lower Silurian, and has a maximum thickness of 300 feet. It was called the "Chazy Formation" in the Geology of Canada for 1863, because shales and sandstones are there associated with the limestone. It occupies a narrow area about the Ottawa and Montreal, and extends to the Mingan Islands and Newfoundland, its thickness not exceeding 300 feet. The western extension of the belt appears in cliffs on the coast of Lake Winnipeg, in the region of Lakes Huron and Superior, in Michigan, Wisconsin, Iowa, and Minnesota. In the lake region it consists of arenaceous and arenaceous

calcareous b
lowed by be

§ 73. I
the river of
ally m le u
occupies par
forms the
at the Fall
erous, fills u
the lower p
the Calcifer
yellow, frial
ferruginous
posed to be
surface ther
yellow or re
larly. The
or painted r
the Calcifer
ally ripple-n
layers, but t
absence of f

§ 74. P
are derived i
particles of c
talline miner
tion, and oc
ready taken
about the cry
in its own cry
reduced to e
grains. By t
of the waves
and finally d
worn into sor
The angular
crystals, but
inal crystalli
quartz, whic
sandstone its
they have b
questionable,
and fragmen
their charact

§ 75. It
banks, stretch
north-eastern

calcareous beds, sometimes difficult to distinguish from the Calciferous layers, followed by beds of argillo-calcareous composition.

§ 73. In 1852 David Dale Owen called it the "St. Peter's Sandstone," after the river of that name, now sometimes called the Minnesota River, where it is usually made up of grains of limpid and colorless quartz, remarkable for whiteness. It occupies part of the slope between the first and second terrace at Prairie du Chien, forms the base of the bluffs at the St. Peter's, and the lower nineteen feet at the Falls of St. Anthony. It rests upon the billowy surface of the Calciferous, fills up the depressions, and is followed conformably by the Trenton. In the lower part there is some shaly material and conglomerate matter washed from the Calciferous and older rocks, but above this it is a remarkably uniform, white or yellow, friable quartzose sandstone, substantially free from silt and calcareous or ferruginous cement. There are oblique and discordant lines of stratification, supposed to be due to the shifting of the waves during deposition, and near the upper surface there is more or less argillaceous material. In some localities it is tinged yellow or red by the oxides of iron, and cemented in streaks, and weathers irregularly. The outliers and standing rocks are brightly colored, and are called pictured or painted rocks. The thickness will exceed 200 feet where filling a depression in the Calciferous; but the average thickness does not exceed 100 feet. Occasionally ripple-marks, fucoidal impressions, and tubes of *Scolithus* occur in the harder layers, but the only fossil yet described from this region is *Lingulepis morsei*. The absence of fossils is due to want of preservation.

§ 74. Prof. T. C. Chamberlin says the constituent grains of sand in this Group are derived in the main from granitoid and schistose rocks, which are composed of particles of quartz intermixed with a variety of softer and more decomposable crystalline minerals. In the metamorphism the quartz was usually last in crystallization, and occupied the angular interstitial spaces between the crystals that had already taken shape, and hence while crystalline in internal structure it molded itself about the crystals of the previously formed minerals. It was thus angular, but not in its own crystalline form. Upon decomposition the associated minerals were mainly reduced to earths and clays, while the undecomposable quartz remained in angular grains. By the action of streams in carrying these down to the sea, and by the agency of the waves in distributing them, the grains were sifted, assorted, rolled, rounded, and finally deposited in the forms in which we now find them. The majority are worn into somewhat spherical grains; others less acted upon remain quite angular. The angularity, however, is not what is characteristic of freely forming quartz crystals, but is due to the circumstances under which it was formed. In the original crystalline rock occasional cracks and cavities occurred filled with secondary quartz, which in such a situation assumed its own crystalline form; and in the sandstone itself secondary crystals might have been formed after deposition, just as they have been in adjacent limestone-beds where their secondary origin is unquestionable, and the degradation of the rock inclosing these would furnish points and fragments of true crystals of quartz, which might not be so far worn as to lose their characteristic form.

§ 75. It occupies a narrow area fringing the Calciferous, or exposed in river banks, stretching in an irregular course from the Lower Menominee River on the north-eastern border of Wisconsin to the mouth of the Wisconsin River. It occurs

in North-eastern Iowa and the eastern part of Minnesota, where its dip is westerly. It occurs in Illinois, at Oregon, on Rock River, and at La Salle, on the Illinois, caused by a local uplift. The unevenness of the Calcareous bed, as proven by artesian boring, is greater near the margin or shore-line of the oceanic deposit than elsewhere. It is known, by artesian boring, in Minnesota more than 100 miles from the Mississippi, and in Illinois more than 100 miles from its exposure in Wisconsin. In some places the sand mingles with the calcareous materials and forms passage beds to the Trenton, and at other places the transition, while conformable, is abrupt. In Missouri the upper part of the Group received the name of the "First Magnesian Limestone," and the lower part the "First Sandstone" and the "Saccharoidal Sandstone." The latter presents very few characters not found in the exposures in Illinois and Wisconsin; while the former is limited in its distribution, and indicates local changes in the deposition of the upper part of the Group. It is usually a gray or buff crystalline, cherty, magnesian limestone, filled with silicious patches, breaking readily with the hammer, and extremely variable in thickness. In New Jersey it consists of a fine, even-grained limestone, sometimes a pure dolomite, except near the base, where there are sandy and calcareous layers. It occurs in long, narrow belts, in a north-east and south-west direction, corresponding to synclinal and anticlinal axes. From this State and from Pennsylvania it is exposed in numerous places within the Appalachian System as far south as Alabama, and may generally be detected by the presence of *Maclurea magna*. In Tennessee the lower part is an argillaceous limestone, varying in thickness from 50 to 600 feet; and if the marble of Knox County is referred to it, its upper part will have a thickness of more than 400 feet. It occurs in the Wahsatch Range in Utah, in the White Pine district of Nevada, in the Wind River Mountains of Wyoming, and in numerous other localities in the great system of mountain ranges of the West, where it also bears the name of the Quebec Group. It has been identified in the Arctic regions, on King William's Island, North Devon, and Depot Bay, in Bellot's Strait, where it is a dolomitic limestone. It graduates into the Black River wherever the latter is separable from the Trenton, and especially where the Birdseye limestone is present. Numerous fossil species connect it intimately with the overlying rocks, many of which occur as high as the Hudson River, viz.: *Strophomena alternata*, *S. incrassata*, *Orthis perveta*, *Leperditia canadensis*, *L. louckana*, *L. amygdalina*, *Orinoceras multicameratum*, *O. bilineatum*, and *Modiolopsis nasuta*. The most characteristic fossil is *Maclurea magna*.

§ 77
logical
posures
cliff ext
compose
well-cha
stone ap
Group o
been so
eye lime
ers, and
organism
and wher
sistent in
about 30
Cephalop
feet in le
The thick
§ 77.
thickness
received t
exceeds 2
greater th
a belt up
on the St
been iden
Lake Win
St. Mary's
It has bee
westwardl
as a local
merely be
logical rea
many spec
are confor
But the st
ness of lim
menced its
succeeding
quently, it
a home in
The *Cyrtoc*

CHAPTER VIII.

BLACK RIVER GROUP.

§ 76. THE Black River Group was defined by Lardner Vanuxem, in the Geological Report for the Third District of New York in 1842, and named from its exposures on Black River. The name "Black River limestone" was applied to the cliff extending from Boonville through Lewis into Jefferson County, the cliff being composed of the Birdseye limestone of the Mohawk and the rocks upon which the well-characterized Trenton limestone is placed. We find the name Birdseye limestone applied to rocks in the report of 1838, but not in the sense of the name of a Group of rocks, as the term Black River was used in 1842, and if the name had been so used it would necessarily give way to the geographical name. The Birdseye limestone was distinguished on the Mohawk by its light dove-color, thick layers, and the presence of crystalline particles representing *Phytopsis tubulosa* or other organisms, which caused the rock to break readily or possess a kind of brittleness, and when broken to clearly show the crystalline spots. This character is not persistent in geographical distribution, and the greatest thickness of the rocks is only about 30 feet. The Black River limestone is distinguished by the abundance of Cephalopoda, and especially by remarkably large *Orthoceras*, some of which are 10 feet in length and a foot in diameter; beside, it has quite an extensive distribution. The thickness on Black River is about 50 feet.

§ 77. From New York it extends into Vermont, where about 12 or 14 feet in thickness becomes a black, finely granular mass, susceptible of a high polish, and has received the name of the Black Marble of Isle La Motte. In Vermont it rarely exceeds 20 feet in thickness; but it outcrops in Pennis Valley, Pennsylvania, with greater thickness than it possesses in New York. It crosses into Canada, and forms a belt upon the margin of the Chazy, but rarely attains any great thickness, though on the St. Lawrence, 90 miles below Quebec, it has a thickness of 130 feet. It has been identified by the presence of gigantic *Orthoceras* on the north-west side of Lake Winnipeg; and its existence has been noted in the Lake Superior region, on St. Mary's, Escanaba, and Menominee Rivers, and on St. Joseph and Sugar Islands. It has been identified at various places in the Appalachian System, but it thins out westwardly and has a limited area of surface distribution. By some it is regarded as a local and peculiar phase of the lower part of the Trenton, or as constituting merely beds of passage from the Chazy to the Trenton, but there are paleontological reasons for retaining the name as a geological subdivision. It contains many species unknown in the Trenton, though others pass up, as the two Groups are conformable, and both represent the deeper oceanic deposits of limestone. But the strongest reason for holding to the geological separation of so small a thickness of limestone from other Groups is that the family *Orthoceratidae*, which commenced its existence in the Upper Taconic, increased in genera and species in succeeding ages until it reached its maximum development in this Group. Subsequently, it diminished in number of species and size of specimens, though it found a home in every Group, until it became extinct in the latest Carboniferous epoch. The *Cyrtoceratidae* and *Endoceratidae* were highly developed, and the *Gomphoceratidae*,

Phragmoceratidæ, and *Gyroceratidæ* here first developed their essential characters. In the Birdseye limestone at Montmorency, Canada, petroleum exudes in drops from fossil corals, supposed to have its origin either in the marine animals or fu-
coid vegetation.

CHAPTER IX.

TRENTON GROUP.

§ 78. THE Trenton Group was named from Trenton, Oneida County, New York. The limestone at the Falls, where it is more than 100 feet thick, was called the Trenton limestone long prior to the use of the words in a geological sense. In 1838 Vanuxem referred to the Trenton limestone, but it was not until 1842 that he and Prof. Emmons so described the Group as to establish it. At Trenton Falls there are two kinds of stone—one a dark, fine-grained limestone, in thin layers, separated by black shale, and abounding in fossils; the other a gray, coarse-grained limestone, in thick layers, forming the top of the mass, and much less fossiliferous. The Group has quite an extensive surface distribution in belts upon the margin of the older rocks in New York, and varies somewhat in its characters, but seems at all times to be a limestone, with the exception of shaly partings. It is 400 feet thick at Chazy, the greatest exposed thickness, and from here it thins toward the east.

§ 79. It enters Vermont from New York in three narrow outcrops, consisting of black layers and seams of limestone and occasional argillaceous matter, with a maximum thickness of about 400 feet. It enters New Jersey, and crosses the counties of Warren and Sussex, with a maximum thickness of about 200 feet. It is frequently exposed in the broken-up hills and mountains of Pennsylvania, showing a thickness from 300 to 700 feet. The exposures continue to occur southerly in the Appalachian Mountains in crossing Virginia, North Carolina, and Tennessee, where, in the eastern part of the latter State, there is a thickness of 1,100 feet, and in the middle part of about 500 feet. It is exposed by an ancient uplift in the central part of Kentucky over several counties, forming a large part of what is called the Blue-grass Region, and reaches as far north as the Ohio River. The thickness is about 700 feet.

§ 80. It has an extensive geographical distribution in Canada. The Montreal and Ottawa sections have each a thickness of 600 feet. The sections in Western Canada, on the Trent River and at Collingwood, have a thickness of 750 feet, but it thins westerly, and in following the outcrops around Lakes Huron and Michigan the exposures rarely exceed 50 feet in thickness. In passing south of Lake Superior it crosses Sulphur, St. Joseph's, and Great Encampment Islands, and thence stretches west and south-west near Little Bay de Noquet and Green Bay, and enters Wisconsin near the mouth of the Menominee River. From here the exposure extends south-west across the State, displaying a large area in the south-western part, and, entering the State of Illinois, occupies more or less of the surface in four or five of the north-western counties. From here the exposures bear north-west and north, occupying several counties in North-eastern Iowa, with a continuing belt across

Minnesota,
Iowa, the
called the
which is
divisions
upper or
feet. It
disappear
ginous ool
the east a
from west
posit from
Northern
British Am
curs in fis
supposed t
Galena lin
Illinois.
consin and

§ 81.

crops occur
mountain
Somerset,
ville, Ky.,
most any p
upon the s
is a regular
fathoms.
with little
of nearly a
in the ocean
Trilobites,
on the dec
Lamellibranch

§ 82.

vision of th
chisonia maj
characterist
ing its age
Kentucky,
and Kentuc
Minnesota;
Blastoidocrin
Leperditia f
tucky; and
The genus
sil in every

Minnesota to St. Paul and the Falls of St. Anthony. In Wisconsin, Illinois, and Iowa, the Group is separable into two divisions, the lower one being a blue limestone called the Trenton limestone, and the upper a dolomite called the Galena limestone, which is the lead and zinc bearing rock of that region. The thickness of these divisions is variable, but where most persistent, as in South-western Wisconsin, the upper or Galena has a thickness of 250 feet, and the lower or blue limestone 120 feet. It thins northward through Minnesota, and the Galena division gradually disappears. The layers of limestone become thin and shaly, with sometimes ferruginous oolitic layers. The Galena appears as a lenticular mass or as thinning toward the east and north, and was apparently derived, so far as detrital matter occurs, from western sources, though a great part, like that of other limestones, was a deposit from the harder parts of animal organisms. Erratics and boulders found in Northern Dakota indicate the existence of Galena limestone in the mountains of British America. The Galena contains about 86.6 lead and 13.4 sulphur, and occurs in fissures and crevices in the limestone, and not in true veins. The ore is supposed to have been precipitated from an aqueous solution. It was called the Galena limestone from the lead or galena, and from its typical exposure at Galena, Illinois. The lead area is about 4,000 square miles, two-thirds of which is in Wisconsin and the rest about equally divided between Illinois and Iowa.

§ 81. It forms some large surface exposures in Southern Missouri, where outcrops occur 400 feet in thickness. Numerous outcrops occur among the Western mountain ranges and in the Arctic regions, on King William's Island, at North Somerset, Boothia, and other places. It was found by the artesian boring at Louisville, Ky., at Columbus, Ohio, and it is expected it will be found by boring at almost any place upon the continent, save where the rocks of older date are exposed upon the surface. It was not formed upon the margin of an island or continent, but is a regular sea deposit of general distribution where the depth did not exceed 2,000 fathoms. The materials are marine, the mass being remains of organic secretions, with little detrital matter. The fauna was abundant and embraced representatives of nearly all the great subdivisions of invertebrate life that now have an existence in the ocean, and several orders and classes, as the Graptolites, Cystideans, and Trilobites, which have become extinct. The Graptolites and Trilobites were then on the decline, while Crinoids, Cystideans, Brachiopods, Corals, Gasteropods, and Lamellibranchs were on the increase.

§ 82. *Receptaculites oweni* is peculiar to and characteristic of the Galena division of this Group, and it is usually accompanied with *Lingula quadrata*, *Murchisonia major*, *Fusispira elongata*, and other characteristic species. The species most characteristic of the Trenton Group, and which may be relied upon as determining its age wherever they occur, are *Orthis tricenaria*, found in New York, Canada, Kentucky, Missouri, and Nevada; *Orthis pectinella*, found in New York, Canada, and Kentucky; *Cyrtolites compressus*, found in New York, Canada, Wisconsin, and Minnesota; *Hyboerinus tumidus*, *H. conicus*, *Amygdalocystites florealis*, *A. radiatus*, *Blastoidocrinus carcharidens*, found at Ottawa, Canada, and High Bridge, Kentucky; *Leperditia fabulites* and *Conularia quadrata*, found in New York, Canada, and Kentucky; and *Orthis borealis*, found in Canada, Wisconsin, Minnesota, and Kentucky. The genus *Amygdalocystites* has a wide geographical distribution, though a rare fossil in every locality, and, so far as known, is confined to this Group. Other char-

acteristic species are *Bythotrephix succulens*, *Monticulipora lycoperdon*, *Schizocrinus nodosus*, *Stictopora elegantula*, *Orthis bellarugosa*, *O. æquivalvis*, *Trochonema umbilicatum*, *Subulites elongatus*, and *Helicotoma planulata*.

§ 83. There are numerous species which continued to live until the Hudson River age, and are therefore common to three Groups, as *Strophomena alternata*, *S. rhomboidalis*, *Leptena sericea*, *Zygospira modesta*, *Rhynchonella capax*, *Calymene callicephala*, *Asaphus gigas*, and *Ceraurus pleurexanthemus*. Such species are usually quite variable in form and size, and seem to have changed to suit the conditions of their habitat, and also, in accordance with the theory of evolution, to have reached the climax of development, and subsequently gradually declined. *Strophomena rhomboidalis* occurs in Trenton, Utica Slate, Hudson River, Clinton, Niagara, Lower Helderberg, Upper Helderberg, Hamilton, Chemung, Waverly, Burlington, and Keokuk Groups. Its vertical range exceeds that of any other species in any of the rocks of the known world, and its geographical distribution is common to every continent where strata of these ages have been studied and described. The varietal forms have been called *S. tenuistriata* from the Lower Silurian, *S. depressa* from the Upper Silurian, and *S. rhomboidalis* from the Devonian and Subcarboniferous. The Lower Silurian specimens are usually smaller, and have fewer concentric wrinkles over the visceral region, than those from the Upper Silurian and Devonian, while the length of the front and lateral margins from the geniculation is usually greater in the Upper Silurian than it is in the Lower Silurian, Devonian, or Subcarboniferous specimens; but these differences are not so constant as to form inflexible characters, and hence it is that many of the learned and better palæontologists have classed them all together under the first and oldest specific name. The various forms which *Strophomena alternata* assume in the same Group of rocks are wonderful; the radiating striæ differ in size and number; the hinge line is sometimes longer and at other times shorter than the greatest width of the shell. The shells are sometimes much longer than wide, and at other times as much shorter. The lateral sides are sometimes straight, and at other times rounded. Some shells are nearly flat, others are deeply concave on the dorsal side and highly convex on the ventral. Age in some specimens appears to have materially thickened the shells, and preserved strong imbricating lines of growth, while in other cases we have much larger shells that are very thin and destitute of imbrications. Like differences may be distinguished in other species having great vertical distribution, as in *Rhynchonella capax* and *Zygospira modesta*.

§ 84. The rocks of this Group are composed almost entirely of remains of the hard parts of animals that swarmed in the seas of that age. Some shells are preserved in good condition, but generally the comminuted fragments are held together by lime cement, forming the limestone strata, leaving well-preserved specimens to be found only in the shaly partings. It is common to find that one animal has grown upon another, as a *Lichenocrinus* upon a brachiopod, and a bryozoan upon the former, under such circumstances as to show the shell was at the bottom of the ocean during the growth of the *Lichenocrinus*, and that the latter must have ceased to grow before the bryozoan attached. From this we infer the clearness of the water, for otherwise mud would have intervened; and we also infer a slow deposition of materials, for the lives of two animals transpired before the deposit was

sufficient to
temperatur
§ 85.

with it; an
formable wi
in New Yor
rocks, and
up into cal
the line of
changing fa

§ 86. I
tion of orga
rocks from
from the T
per minute.
discharges h
surprising w
a considerab
offers succes
trating water
hydrogen ga
Niagara Fall
well-known
collieries, an
with an unp
rocks charged
the Christian
occurs in the
at Pakenham
petroleum oc
has been fou
absence of p
and vegetable
It has been a
Group, but th

sufficient to cover a thin shell. There is no evidence of any difference between the temperature of the water then and now, nor between the climate then and now.

§ 85. Wherever the Black River limestone exists, the Trenton is conformable with it; and where the Black River is not distinguished, the Trenton is usually conformable with the Chazy. The Trenton is conformable with the Utica Slate above, in New York and Canada; but there is an abrupt change in the character of the rocks, and a marked difference in the fauna, while in Kentucky it graduates up into calcareous shales of the age of the Utica Slate by imperceptible grades, so the line of separation can not be determined, except as based upon a slowly changing fauna.

§ 86. Light carbureted hydrogen gas is often the product of the transformation of organic matter at ordinary temperatures, and is abundant in the palæozoic rocks from the Chazy to the Permian. A spring at Caledonia, Canada, issuing from the Trenton Group, evolves 300 cubic inches of carbureted hydrogen gas per minute. It is saline water. Another discharges somewhat less, and another discharges large quantities of sulphureted hydrogen gas. This is not considered surprising when it is remembered the Chazy Group in the Ottawa Valley includes a considerable thickness of shales and argillaceous limestones, and the Quebec Group offers successions of limestones and shales, whose slow decomposition from infiltrating waters will furnish such gases. In higher strata, however, the carbureted hydrogen gas escapes in much greater quantities, as at the burning spring near Niagara Falls, and in the region of the oil-wells. Carbureted hydrogen gas is the well-known "fire-damp" of the coal-mines. It collects in ill-ventilated galleries of collieries, and when sufficiently mixed with the atmosphere, if it comes in contact with an unprotected flame, it explodes with great violence. It exudes from all rocks charged with petroleum or naphtha, and was known and used for fuel before the Christian era on the Caspian Sea, where it is evidently inexhaustible. Petroleum occurs in the cavities of fossils, *Orthoceras* sometimes holding several ounces of it, at Pakenham and Lancaster, Canada. While both carbureted hydrogen and petroleum occur in the rocks of the Quebec and all succeeding Groups, yet none has been found of commercial value as low as the Trenton. The reasons are, absence of porous strata and cavities for its collection, and because the animal and vegetable matter was not collected in sufficient quantity at any single locality. It has been asserted the gas in Western Ohio and Northern Indiana is from this Group, but the author thinks all the evidence is against such conclusion.

CHAPTER X.

UTICA SLATE GROUP.

§ 87. This Group was named the Utica Slate from Utica, New York, and quite fully defined as a geological subdivision in 1842 by both Vanuxem and Emmons in their respective reports. It is in typical localities a dark-colored slate or shaly mass, highly charged with carbon, and agreeing in its composition with the dark layers that separate the limestone strata in the Trenton Group. The surface exposure forms a belt resting upon the Trenton, extending from New Jersey across New York into Vermont, passing under Lake Champlain and entering Canada. The greatest thickness in New York is about 600 feet, and in Vermont about 100 feet. It exposes considerable surface in Canada, never exceeding 500 feet in thickness, and extends from Lake Huron, where it thins out, to the eastern shores of the continent, appearing on the Saguenay, in Newfoundland, and the Island of Anticosti. It is very fossiliferous, and everywhere characterized by the presence of *Triarthrus becki*; and in the vicinity of Ottawa *Triarthrus spinosus* is abundant, and the Scotch fossil, *Siphonotreta scotica*, occurs. It is often interstratified with thin bands of limestone.

§ 88. It is exposed in numerous places in the Appalachian System, and attains a thickness in Huntingdon County, Pennsylvania, of more than 1,000 feet. It thins out westerly, and loses its character as a black slate before reaching the Ohio River, where it is composed of blue calcareous shales and marls with interstratified thin limestones, apparently forming beds of passage from the Trenton to the Hudson River without any want of conformability. The change in its lithological characters would have prevented forever its identification in the banks of the Ohio, had it not been for the tell-tale fossils. The abundance of *Triarthrus becki* and *Leptobolus lepis* and associate fossils settled the question of its identity. It is unknown farther west, but exists in the Arctic regions as a more or less calcareous slate. The fossils of the greatest geographical distribution, and by which it may generally be recognized, are *Triarthrus becki*, *Leptobolus lepis*, *Asaphus canadensis*, *Lingula probye*, and *Graptolithus quadrimucronatus*. The rocks are composed in part of mechanical sediment, derived from sources east of the Appalachian System, and not almost wholly of shells and the harder parts of animals, as the Trenton is below and the Hudson River above. It thins westerly, and as the mechanical sediment disappears the marine deposits form continuous passage beds from the Trenton to the Hudson River. The strongest reason for its retention as a geological subdivision is found in the fauna with which it abounds; for at many localities, *e. g.*, Cincinnati, Ohio, and Jefferson County, New York, it can only be separated from the Hudson River Group by an arbitrary line; and at other localities, *e. g.*, Deerfield, New York, and in Kentucky, the Trenton Group is so blended with it that the line of demarcation is wholly obscured. The Galena limestone of Northern Illinois, Eastern Iowa, and South-western Wisconsin occupies substantially the same geological position, though its affinities are more closely allied with the Trenton, while the relations of this Group are nearer the Hudson River; beside, none of the characteristic fossils of this Group are found in the Galena, and none of the

characterist
from this C
evolve larg
gas has bee
shales of th
distilled the
shales of Co

§ 89. T
River in Ne
1842. At t
thick-bedded
regular order
mentions, as
Cone," which
thickness in

§ 90. T
Appalachian
of 1,200 feet
surface rock
Ohio, to near
western part
limestone, an
Blue limesto
consin, and n
240 feet. In
it appears in
ada, extendin
the Great La
ain ranges bo
1,100 feet, bu
Great Lakes
about 2,000

§ 91. Th
upon the olde
to find it alm
It is the equi
England and
Trenton and
able depth, n
the sandstone
and derived fr

characteristic fossils of the Galena occur in this Group. A petroleum spring rises from this Group on the Grand Manitoulin Island, and saline springs at Varennes evolve large volumes of carbureted hydrogen gas. At one of these springs the gas has been collected in a holder, and employed in lighting a house. The black shales of this Group contain variable amounts of combustible matter, and when distilled they give, beside inflammable gases, portions of oily matter, which in the shales of Collingwood are equal to four or five per cent.

CHAPTER XI.

HUDSON RIVER GROUP.

§ 89. THE Hudson River Group was named from an exposure near Hudson River in New York, and first defined in the geological report by Vanuxem in 1842. At the typical locality it consists of shales, shaly sandstones, slates, and thick-bedded grits, stratified and conformable, alternating many times without any regular order of alternation. It was called the Lorraine Shales by Emmons, who mentions, as occurring at one place in New York, that structure called "Cone within Cone," which is so common in the Devonian and later formations. Its maximum thickness in New York is about 800 feet.

§ 90. The Group is largely exposed in Pennsylvania and other States in the Appalachian System, as far south as Tennessee, and has a thickness in some places of 1,200 feet. In the latter State it has been called the Nashville Group. It is the surface rock of many counties in Kentucky, extending from above Maysville on the Ohio, to near Louisville. In the south-eastern part of Indiana and the south-western part of Ohio, it consists of alternating layers of blue calcareous clay and limestone, and has a thickness of about 800 feet. It has been called in this section the Blue limestone. It occurs in the northern part of Illinois, southern part of Wisconsin, and north-eastern part of Iowa. Its thickness in these States does not exceed 240 feet. In the south-eastern part of Missouri its thickness is about 250 feet, and it appears in Texas and New Mexico. It has a wide geographical range in Canada, extending from the Island of Anticosti and the eastern border west, by way of the Great Lakes, to the Red River of the north, and again appearing in the mountain ranges bordering the Pacific. In the vicinity of Toronto its thickness is about 1,100 feet, but it is much thinner in its western extension, and in the region of the Great Lakes rarely exceeds 100 feet. Its greatest thickness in Eastern Canada is about 2,000 feet.

§ 91. This Group is persistent and of almost universal distribution, except upon the older rocks that were dry land before its deposition. We would expect to find it almost anywhere on the continent by boring through more recent deposits. It is the equivalent, to some extent, of the Caradoc sandstone, or Bala Group, of England and Wales, and is represented in different European exposures. Like the Trenton and all earlier Groups, it is a marine deposit made in water of considerable depth, not a littoral or shore-line deposit as the Potsdam Group was, though the sandstone occurring in many of the northern exposures was evidently mechanical and derived from land at no great distance to the north.

§ 92. The seas swarmed with animal life and fucoid organisms, and the rocks are composed almost wholly of their remains. It is literally a graveyard of invertebrate life. The Brachiopoda and Bryozoa reached in this age the stage of their greatest varietal development, and possibly the highest state of their existence.

§ 93. As the exposure in Ohio, Indiana, and Kentucky is very large and quite characteristic of it in other places, it may be fit and proper to further define it. To go from the Ohio River, at Cincinnati, west 51 miles to Osgood, Indiana, or north to Dayton, or north-east to Xenia, Ohio, one will pass across the upturned edges of this Group, and reach the Niagara. The rocks dip westerly and northerly at the rate of about ten feet in a mile. The hills at Cincinnati expose about 400 feet in thickness, constituting the lower half of the Group; and the upper half, or about 400 feet, occurs between the top of these hills and the bordering Niagara Group, about 50 miles distant to the north and west. The area of its exposure in Ohio is all of Hamilton, Butler, Warren, Clermont, and Brown Counties, and part of eight counties that border upon these. The exposure in Indiana is about half as great, reaching as far north as Richmond, and bordering the Ohio nearly half-way from Madison to Jeffersonville. The exposure in Kentucky is greater than in Ohio, for it surrounds the Trenton Group in that State. Throughout the whole area it is composed of alternate layers of calcareous clay and limestone of varying thickness. In some places calcareous clay is 6 or 8 feet thick, without a layer of stone. At other places one layer of stone, 4, 6, 8, or 10 inches in thickness, follows another, with intervening layers of calcareous clay of much less thickness, for 40 or 50 feet. It is rare to find a layer of limestone more than a foot in thickness. All the layers are broken into small, irregular pieces of suitable size for cellar and other light stone-work, for which they are used. The blue calcareous clay exposed to the action of the weather for a few years loses its color and becomes of a dull gray hue. The sulphuret of iron occurs in the blue rocks, but instead of this we find iron oxide and sulphate of lime in the gray. The silicious matter prevails over the carbonate of lime in the layers of calcareous clay, while the carbonate of lime is much in excess of the silicious matter in the stone, due, in part at least, to the fact that the stones are a mass of more or less comminuted shells, corals, and crinoids. There is nothing in the general character and appearance of the rocks and calcareous clays to indicate the changes which the fossils undergo; that is, the changes are not to be attributed to surrounding conditions without the aid of that law of animal evolution which the science of palæontology teaches us has taken place in all past geological ages.

§ 94. Some fossils, as *Calymene callicephala*, *Asaphus megistus*, *A. gigas*, *Beyrichia chambersi*, *Leptæna sericea*, *Bellerophon bilobatus*, *Zygospira modesta*, *Strophomena alternata*, and *Orthis testudinaria*, pass from the extreme lower part to the extreme upper part of the Group; and all of them save *Beyrichia chambersi* are known from lower rocks, and *Leptæna sericea* occurs in higher ones. *Streptorhynchus hallianum* has a limited range in the lower part, *S. planoconvexum* and *S. sinuatum* a limited range below the middle of the Group, *S. nutans* and *S. sulcatum* in the middle of the upper half of the Group, and *S. subtentum* and *S. filitextum* in the upper part. *Lichenocrinus crateriformis*, *L. dyeri*, and *L. pattersoni* are confined to the lower half, and *L. tuberculatus* and *L. affinis* to the upper part. *Acidaspis crossotus* occurs in the lower part, *A. anchoralis* and *A. cincinnatiensis* in the middle part, while *A. onealli* occurs in the

upper part.

lulu, *Tetradium*

Such are a f

completely pr

Crinoids, as a

particular rar

istic of the c

alis. *O. subqu*

Glyptocrinus a

§ 95. W

the greatest b

of the Potsdam

nearer the line

Silurian, than

Hudson River

unconformable

almost comple

unconformably

Clinton Group

upon it. On

by rocks appar

cal break. O

at once below

by 45 species t

s at any other

probably the st

§ 96. Ther

ability and pal

represented by

been described

which had beco

which had come

to have an exi

ver been disc

and had existed

ain, by warmth

ion. If the lar

course none wil

or fresh-water a

upper part. *Rhynchonella capax*, *R. dentata*, *Streptelasma corniculum*, *Favistella stellata*, *Tetradium fibratum*, *Cypriocardites haynesi*, etc., are confined to the upper part. Such are a few illustrations of the changing fauna at different elevations. To completely present the subject would require the enumeration of all the species. Crinoids, as a rule, are limited vertically, and hence each species is sought in its particular range. Species having a wide geographical distribution, and characteristic of the Group are *Aulopora arachnoidea*, *Stomatopora inflata*, *Orthis occidentalis*, *O. subquadrata*, *O. retrorsa*, *Pterinea demissa*, *P. insueta*, *Cyclonema bilix*, and *Glyptocrinus decadactylus*.

§ 95. With this Group the Lower Silurian closes, because at its top we have the greatest break stratigraphically and palæontologically that occurs from the base of the Potsdam to the top of the Lower Helderberg, and because it approaches nearer the line of division established by Murchison, between his Lower and Upper Silurian, than any other line, if, indeed, it is not identical with it. Wherever the Hudson River has been examined on the continent, the superimposed rocks are unconformable with it, no passage-beds are found, and the palæontological break is almost complete. In the Western States the Niagara Group succeeds it, and rests unconformably upon it. In the Eastern States it is succeeded by the Medina and Clinton Groups before the Niagara is reached, but the Medina rests unconformably upon it. On the Island of Anticosti it has a thickness of 950 feet, and is followed by rocks apparently conformable with it, although there is an abrupt palæontological break. Of 121 species known to Prof. Billings from Anticosti, 80 disappear at once below the dividing line, and 41 only appear above it, where they are joined by 45 species that are not found below. This palæontological break is less than it is at any other known place on the continent; but it is so great as to show that probably the strata are not strictly conformable.

§ 96. There is an important period of time indicated by this want of conformity and palæontological change. Vast ages must have intervened, which are not represented by any known rocks on the continent. More than 400 genera have been described as existing previous to this time, more than three-fourths of which had become extinct. Or, in other words, less than one-fourth of the genera which had come into existence prior to the close of the Lower Silurian Age continued to have an existence afterward. No evidence of the existence of land-plants has ever been discovered in Lower Silurian rocks. We are convinced, however, that land had existed above water for ages; that it was necessarily refreshed by sun and rain, by warmth and air, and that it may have sustained some kind of land vegetation. If the land vegetation did not possess hard parts capable of preservation, of course none will ever be found. Neither has any evidence of the existence of land or fresh-water animals of this era ever been discovered.

CHAPTER XII.

UPPER SILURIAN.

§ 97. ALL the rocks of the Upper Silurian System are marine; but land-plants, or such as may have existed in marshes, and received support from sunlight and air, have been found within them. No remains of land or fresh-water animals, or marine vertebrates, have been discovered in North America. There is no radical difference in the general character of the Lower Silurian and Upper Silurian fossils, because vertebrates had not made their appearance, and the same orders of invertebrates were represented in each era; but the separation into two Systems is very convenient, because both are introduced with sandstone Groups, and the Trenton in the Lower Silurian, and Niagara in the Upper Silurian, are alike extensive in geographical distribution, and some analogy may be traced between the upper Groups in each System. On the whole, the calling of one System Lower Silurian, and the other Upper Silurian, was a happy hit in nomenclature as well as correct in science.

MEDINA GROUP.

§ 98. THIS Group took its name from Medina, New York. The rocks were described by Vanuxem in 1842, under the names Oneida Conglomerate, Gray Sandstone of Oswego, and Medina Sandstone. At the typical localities they are conglomerate, and gray and red sandstone. The conglomerate is hard and gritty, and composed of quartz pebbles and sand so firmly cemented as to be used for millstones. The sandstone is argillaceous, thinly laminated, and of red, gray, and mottled colors. Where it is not fragile, but firmly cemented, it makes a good building stone, and has been largely used for paving streets, as it readily breaks into stones of regulation size. The Group borders Lake Ontario on the south, and extends in an east and west line of exposure about three-fourths the length of the State, and entering Canada at the Niagara River, continues to Lake Huron. In Oneida and Oswego Counties the thickness is from 500 to 600 feet; at the west end of Lake Ontario 614 feet, and at Lake Huron 100 feet. It thins so rapidly that few, if any, traces have been discovered west of this lake. A small surface area in New Jersey has a thickness of 900 feet, and a larger one in Pennsylvania has a thickness of 2,500 feet. It occurs in patches among the broken ranges of the Appalachian System in Maryland and other States, as far south as Tennessee; but is unknown in the Western States.

§ 99. The conglomerate is 500 feet thick in the Shawangunk Mountains, and 700 feet in the Kittatinny Valley in Pennsylvania. It graduates into the gray sandstone, and then into the red sandstone, so they can scarcely be distinguished except by color; and the gray sandstone in like manner graduates into the conglomerate by enlarging and increasing the number of its pebbles; so there is no reason, stratigraphical or palæontological, for subdividing the Group, as was done in early work on the New York Survey. It always rests unconformably upon the Hudson River Group, and bears the internal evidence of having been derived from land immediately north and east, and of having been deposited in shallow water, subject

to waves and conglomerate ous, though definition, has been deposited about shells, In all these from the Low § 100. In fairly well pre and species *Lingulella cune* the whole ext brine is frequ ured hydroge at Gasport it

§ 101. The defined by Va Survey of that the typical lo sandstone, ofte at other places limestones, co narrow belt of westward south width in Wayn Huron, appear probably enter and rapidly thi The two upper Clinton Group fossils peculiar extension into as far as Antic of the divisions Hudson River It occurs in the crossing Penns thins out before borders of the from the mecha

to waves and currents which transported the materials only short distances. The conglomerate indicates a shore-line and rapid deposition, and is almost non-fossiliferous, though a few fragments of fucoids and shells, generally too imperfect for definition, have been found in it. The sandstone, too, bears the evidence of having been deposited near the land in shallow water, not only in wave-lines, rill-marks about shells, and ripple-marked slabs, but in mud-cracks produced by sun-drying. In all these respects it compares with the Potsdam, which separates the Taconic from the Lower Silurian.

§ 100. In the more argillaceous part of the sandstone, fossils are sometimes fairly well preserved. The characteristic fossils are *Arthropycus harlani*, both genus and species being confined to this Group, and having a wide distribution, and *Lingulella cuneata*, a strongly marked species. Saline springs are common throughout the whole extent of these rocks, and brine is universally found by boring. The brine is frequently impure from the presence of muriatic lime and iron. Carbonated hydrogen gas rises in many places on the Erie Canal east of Lockport, and at Gasport it was collected and used for illuminating purposes a half century ago.

CHAPTER XIII.

CLINTON GROUP.

§ 101. This Group was named from the town of Clinton, in New York, and defined by Vanuxem in 1842; and re-defined by Hall in 1843 in the Geological Survey of that State. The rocks have no uniformity in color or composition. At the typical locality there is green and black-blue shale; green, gray, and red sandstone, often laminated; calcareous sandstone and red fossiliferous iron ore beds; at other places, it consists of shaly sandstones and shales of various colors, impure limestones, conglomerates, and oolitic iron ore, with concretions. It occupies a narrow belt of country in New York, commencing near Canajoharie, and stretching westward south of Lake Ontario, resting on the Medina Group, with the greatest width in Wayne County, and, entering Canada at Hamilton, extends west to Lake Huron, appearing on Drummond, Manitoulin, Cockburn, and other islands, and probably enters the Peninsula of Michigan with a thickness of less than 50 feet, and rapidly thins out. The maximum thickness in New York is about 400 feet. The two upper bands of limestone included by the New York geologists in the Clinton Group, are now generally classed with the Niagara, as they possess no fossils peculiar to the Clinton, and the shales which separate them thin out in their extension into Canada. In its easterly extension from New York, outcrops occur as far as Anticosti Island and Newfoundland. On Anticosti it is described as one of the divisions of the Anticosti Group, which there includes the rocks from the Hudson River to the Niagara, and has a maximum thickness of about 500 feet. It occurs in the Appalachian chain as far south as Georgia and Tennessee, and in crossing Pennsylvania develops a thickness of more than 2,000 feet. The Group thins out before reaching the Western States, and is unknown except upon the borders of the Appalachian and Laurentian elevations. It appears to have resulted from the mechanical deposition of materials derived from land lying north and east

of it, and to represent a border-land and shallow water deposit, that extended only a short distance from the primitive source of its materials.

§ 102. In Western New York the dividing line between the Medina and Clinton is sharply defined, and the materials of which each are composed are quite distinct; but in the central part they graduate into each other, the Clinton being largely composed of sandstone. There is strong resemblance between the marine vegetation which abounds in the two periods. Westerly the Clinton is more calcareous and more fossiliferous, and graduates up into the Niagara in its lithological and fossil characteristics. The Medina, Clinton, and Niagara are clearly defined in some localities; but in others the Medina graduates into the Clinton, and in others the Clinton blends with the Niagara. There is no want of conformability between them where best developed, and the lines of separation show only a changed condition or altered circumstances under which the deposition was continued from one Group to the other. *Pentamerus oblongus*, *Spirifera radiata*, *Meristella cylindrica*, and *Lingulella lamellata* are among the species accredited both to the Clinton and Niagara, and which show the intimate relation between the Groups. The Clinton abounds in fucoids, tracks, and trails, the former being more abundant than in any earlier Group. The fossils having the greater distribution and being most characteristic are *Ichnophycus tridactylus*, *Graptolithus clintonensis*, *Helopora fragilis*, *Athyris naviformis*, *Leptocelia hemispherica*, *Triplesia congesta*, *Cyclonema cancellatum*, and *Cornulites distans*. The iron ore beds are frequently thick enough to be valuable, and are worked successfully. They are sometimes very fossiliferous, and the quantity of iron is decisive proof of the vegetative character of the fucoids of that age, and the absence of land-plants among the fossils is almost conclusive against their existence at that period.

CHAPTER XIV.

NIAGARA GROUP.

§ 103. This Group was named from its development at Niagara Falls, where the rock over which the water is precipitated belongs to it. It was defined by Vanuxem in 1842, and by Hall in 1843. It is the most persistent in its geographical distribution of any Upper Silurian Group; indeed, wherever the Upper-Silurian is found it is present, except with the exposed belts of the lower Groups, and not unfrequently it constitutes the whole formation. It generally consists of limestone and shales, but sometimes becomes arenaceous, argillaceous, or highly ferruginous. In New York it exposes an east and west belt almost the entire length of the State, a short distance south of Lake Ontario, with a maximum thickness of 300 feet. Near Niagara Falls there are 165 feet of limestone (directly at the falls 85 feet) overlying 80 feet of shale. In its western extension it crosses the Niagara River into Canada, appears at Lake Huron, on Manitoulin and Drummond Islands, occupies the southern part of the northern peninsula of Michigan, spreads over the south-eastern part of Wisconsin and the northern part of Illinois. Keeping south of the Lower Silurian area in the north-western part of Illinois, it enters Iowa below Dubuque, and presents a surface exposure 160 miles in length by 40 or 50 in

breadth. In
sometimes too
Chicago and
Joliet. It so
where it indic
ties near its b
Iron Ridge, in
640 feet, in W
eastern Canad
mum thicknes
lachian System
shales, it has
Tennessee and
a porous mag
tion, and now
forms a sub-ci
great Lower S
South-eastern
and whitish-ye
ing, with a ma
there is iron-st
stone is used f
Paul, and othe
rounds the Lo
and frequently
in the Arctic
its exposures o
from Capes H
equivalent of t
Russia, German
ring in the upp
occurring at t
Baltic Sea. It
to the Devonian
nearly all the
§ 104. It is
shore-line, and
is generally lim
ing the entire s
It is so thorough
culty in recogni
plants occur—A
pression of unce
have been a ma
mud, which pre
plant in Americ
in the Clinton.

breadth. In Wisconsin, Illinois, and Iowa it is principally a magnesian limestone, sometimes too porous or friable for building purposes, but suitable for lime, as at Chicago and Racine; at other places having a good reputation for buildings, as at Joliet. It sometimes occurs more or less saturated with petroleum, as at Chicago, where it indicates the presence of shales immediately below it, and in some localities near its base it contains beds of hematite in small lenticular concretions, as at Iron Ridge, in Dodge County, Wisconsin. The maximum thickness in Illinois is 640 feet, in Wisconsin 800 feet, and in Illinois and Iowa 600 feet. It occurs in south-eastern Canada, in New Brunswick, Newfoundland, and Anticosti, where its maximum thickness is 800 feet. It occurs in nearly all the States to which the Appalachian System extends. In crossing Pennsylvania, where it consists mostly of shales, it has a maximum thickness of 1,600 feet. It occupies extensive areas in Tennessee and Alabama; and in the latter State that part of it which was originally a porous magnesian limestone, subsequently became infiltrated with iron in solution, and now constitutes the celebrated fossiliferous iron ore of Alabama. It forms a sub-circular belt of exposures from 5 to 60 miles in width surrounding the great Lower Silurian area in the middle part of Kentucky, South-western Ohio, and South-eastern Indiana, where it consists of hard, blue and gray limestone, yellowish and whitish-yellow magnesian limestone, and shales, variously alternating and combining, with a maximum thickness of about 600 feet. In some places near the base there is iron-stained chert. At Cedarville, near the top, the porous magnesian limestone is used for the manufacture of lime, and the harder limestone at Dayton, St. Paul, and other places is used for building and other economic purposes. It surrounds the Lower Silurian and Taconic uplift in the southern part of Missouri, and frequently occurs in the Rocky Mountain ranges. It outcrops far to the north, in the Arctic regions north of British America. Fossils have been described from its exposures on Beechy, Cornwallis, Griffiths, Seal, Napoleon, and Offley Islands, from Capes Hilgard, Hotham, Louis, and other points. It is substantially the equivalent of the Wenlock in England, and has its representative in Scandinavia, Russia, Germany, and other European countries. Several species of fossils occurring in the upper part of the Group at Waldron, Indiana, are identical with those occurring at the equally celebrated locality on the Island of Gottland, in the Baltic Sea. It is so constantly present where the rocks from the Lower Silurian to the Devonian are exposed, that it is regarded as a universal Group underlying nearly all the more recent rocks on this continent.

§ 104. It is a deep-sea deposit, as distinguished from all mechanical, littoral, shore-line, and marsh deposits, and, like most other undisturbed marine sediments, is generally limestone. The ocean must have swarmed with invertebrate life during the entire age, as the rocks are almost wholly constituted of their harder parts. It is so thoroughly characterized by its fossils that a paleontologist has little difficulty in recognizing it wherever it exists. It is in this Group the earliest land-plants occur—*Psilophyton* and *Glyptodendron*. The latter was founded upon an impression of uncertain value in a magnesian limestone. *Psilophyton* is supposed to have been a marsh-plant that drifted in the ocean and became imbedded in the mud, which preserved its characters. *Psilophyton princeps* is the oldest fossil land-plant in America. Fucoids are scarce; in striking contrast with their abundance in the Clinton. Sponges were more numerous than in any preceding age. Coral-

reefs were formed, which may now be traced for many miles; single masses were several feet in diameter, and the beauty of their structure is not surpassed by any of the corals which now abound in the ocean. Some of the species, too, were almost world-wide in their distribution, as *Halysites catenulatus*, *Heliolites pyriformis*, and *Favosites forbesi*. It is famous, too, for its Echinoderms. The Cystideans commenced their existence in Taconic times, as evidenced by the plate called *Ecocystites*, and reached the climax of their evolution and development in this Group, and almost suddenly disappeared from the face of the earth, a few small species only being found in the Lower Helderberg and Lower Devonian, where the entire order became extinct. Cystideans were marine animals, related to the Crinoidea. Some were sessile; others possessed a column and roots, by which they attached to other objects; and others were free, and possessed a flexible column tapering to a point, which could be used for attaching purposes. The head was globular, oval, pyriform, conical, cylindrical, or of any other shape, but always covered with an external skeleton composed of polygonal calcareous plates, which are sometimes very richly ornamented. The fracture of the plates presents the same crystalline structure as crinoidal plates do. In some species the number of plates and order of arrangement remained constant throughout the life of the animal, the size of the animal increasing by the growth of the original plates, which enlarged throughout, instead of by addition to the edges. In other species the plates are not limited in number, and have no order of arrangement; they increase in size, or new plates are introduced, so as to destroy uniformity in different specimens in the same species. In other species the dorsal side has a definite number of plates and regular order of arrangement without any increase, while the ventral side has no order of arrangement of the plates, and they increase in number to cover the increased growth of the animal. There are usually two principal apertures, and often many smaller ones through which the most important functions of the animal economy were exercised. One of these is called the mouth, and is found on the side near the base or near the apex. It is a curious fact that so important an organ as the mouth occurs almost anywhere on the body of a Cystidean, but, of course, always occupying the same position in each species. Another aperture, called the ambulacral orifice, occurs near the center of the upper part of the body, and between the bases of the arms, when the species possessed such organs. The other apertures are called calycine pores and pectinated rhombs. The calycine pores served in some manner to introduce water into the interior of the animal, but they bear little resemblance to each other in different species, and one can form no adequate idea of the system of circulation. Pectinated rhombs differ in number and position in different species, and sometimes do not occur at all. Their function, too, is an absolute mystery, except they furnished another medium of communication from the exterior to the interior of the body. The Blastoida commenced existence in this Group by the appearance of *Stephanocrinus*, and became extinct in Carboniferous times. The order Myelodactyloidea, another Echinoderm of very uncertain affinity seems to have been confined to this age. The development of the Crinoidea was wonderful, no less than 15 genera making their first appearance, eight of which are unknown in later rocks. The Graptolitidae here became extinct.

§ 105. *Holocystites* occurs in Ohio, Indiana, Illinois, New York, and other States. Twenty-five species have been defined, and none are known from higher

or lower rocks.
lyptocrinus
for the same
O. hybrida
distribution,
§ 106.

In some parts
exudes from
these rocks
spring" on
rose from be
its way in co
plication of
which seldom
and only obt
height of sev
the shale, wh
through 85 f
limestone at
sian boring fa
from surface
mercial impor
which have p
sea. There n
substances in

§ 107. Tr
fined by Logan
and having a
larly distinguish
fossils closely
It may have b
north-western
in Canada, but
daga Group, a
few fossils foun
the Guelph.
grandis are fou
dentalis, *Murchi*
oides, and *Din*

or lower rocks, and it may therefore be considered a characteristic genus. *Eucalyptocrinus* has a wider geographical distribution, and is more abundant, and for the same reason may be called characteristic. *Orthis elegantula*, *O. flabellum*, *O. hybrida*, *Calymene blumenbachi*, and *Ilkenus barriensis* have almost world-wide distribution, and are characteristic of rocks of this age.

§ 106. The dolomites of this Group in Canada are more or less bituminous. In some parts of Western New York they contain so much solid bitumen that it exudes from the rocks when heated. The escape of carbureted hydrogen from these rocks is of common occurrence. Lyell described in 1841 a "burning spring" on the river just above Niagara Falls, where the light hydro-carbon gas rose from beneath the water out of the limestone rock. The invisible gas makes its way in countless bubbles through the clear, transparent water, and on the application of a lighted candle it plays about with a lambent, flickering flame, which seldom touches the water, the gas being at first too pure to be inflammable, and only obtaining sufficient oxygen after mingling with the atmosphere at the height of several inches above the surface of the river. This gas had its origin in the shale, which forms the lower part of the Falls, and has found its way up through 85 feet or more of quite compact limestone. Petroleum occurs in Niagara limestone at Chicago, which had its origin in some shaly strata beneath, but artesian boring failed to discover it in commercial quantities. Where gas or oil escapes from surface limestone there is little prospect of finding accumulations of commercial importance by artesian boring, because so much has escaped in the ages which have passed away since the elevation of the limestone above the water of the sea. There must be an impervious covering of clay or stone to retain such volatile substances in valuable quantities.

CHAPTER XV.

GUELPH GROUP.

§ 107. THIS Group was named from the town of Guelph in Canada, and defined by Logan in 1863. It appears as a lenticular mass, resting upon the Niagara, and having a maximum thickness of 160 feet. It is a limestone dolomite, particularly distinguished for having no fossil Echinoderms, while it is rich in other fossils closely allied to those in the Niagara, some of the species being identical. It may have been a brackish water-deposit in an arm of the sea. It occurs in the north-western part of Ohio with all the fossils and characteristics pertaining to it in Canada, but is unknown elsewhere. It is doubtless of the same age as the Onondaga Group, and probably should not bear a distinct name, as among the very few fossils found in the latter, *Murchisonia boydi* and *Cyclonema sulcatum* occur in the Guelph. *Megalomus canadensis*, the most common species, and *Trimerella grandis* are found in the Niagara. The characteristic fossils are *Pentamerus occidentalis*, *Murchisonia bivittata*, *M. longispira*, *Subulites ventricosus*, *Pleurotomaria solaroides*, and *Dinobolus gallensis*.

CHAPTER XVI.

ONONDAGA GROUP.

§ 108. This Group was named the Onondaga Salt Group, by the New York Geologists, from Onondaga County, New York, in 1839, and re-defined by Vanuxemi in 1842, and by Hall in 1843. The Canadian Geologists very properly dropped the word "salt" from the name. It consists, on Oneida Creek and Cayuga Lake, in the lower part, of clayey deposits and red shale, showing green spots, followed by gypseous shales and impure limestones, which at the commencement alternate with the red shale, and this is followed by the gypseous deposit, which embraces the great lenticular masses quarried for plaster, and this by a magnesian rock having groups of needle-form cavities caused by the crystallization of sulphate of magnesia, and the upper member is the Waterlime. It rests upon the Niagara from the western line of New York, east to the middle part of Herkimer County, where the Niagara thins out; it then rests upon the Clinton until it disappears, and then upon older rocks until it reaches the Hudson River. It is therefore unconformable with the underlying rocks in middle and Eastern New York. The red shale loses its color west of the Genesee, becomes a bluish green, and gradually thins out, showing the unconformability in Western New York. The passage from the Niagara to the Onondaga is abrupt, offering no gradation in character of products or in continuation of fossil species. The great mass of gypseous deposits consists of yellowish or drab, and brownish colored argillaceous, and calcareous shale and slate, or of hard and compact slate, which weathers as if hacked by an instrument. The dark color of the gypsum, and brownish color of other rocks, is due to carbonaceous matter. An important member is called the vermicularimerock, which is gray or blue, and perforated with holes and cells, once filled with soluble saline material, which subsequently dissolved, leaving the cavities, some of which are hopper-shaped, and were produced by common salt, as no other common soluble mineral presents similar ones. The sulphate of magnesia cavities are lined with carbon, showing the liquid that held the salt in solution, contained bituminous matter, the salt ejecting its particles in the act of assuming form, as occurs in the purification of acetic acid when obtained from the distillation of wood. This Group is celebrated for its salines, and formerly furnished nearly all the salt consumed in New York; for this reason it has been called the Salina and Saliferous Group. Sulphate of Strontian and sulphurets of lead and zinc occur in small quantities. Sulphuric acid escapes with the water from the earth in many localities, giving rise to acid springs, and sometimes destroying the water in wells for culinary purposes, as the sulphuric acid becomes strong enough to coagulate milk.

§ 109. The Group attains its greatest thickness at about 1,000 feet in Wayne County, and gradually diminishes westerly, so that on Grand River, Canada, it does not exceed 300 feet, which belongs chiefly to the upper portions, from the summit to a little below the gypsum-beds. The beds of gypsum are never continuous for long distances, but appear as detached lenticular or dome-like masses; the strata above them being arched over and often broken, while those below constitute an even, undisturbed floor. The Group is continued through Lake Huron to

the Straits of
land. The
broken up in
southern lin
Island, and
be seen on
County, and
not been acc
rable, the th
varying from
Iowa, and is
composition o
it does not ap

§ 110. I
extremely ran
part of it and
quetra were e
fera, *Atrypa*,

§ 111.
used for mak
while the Can
berg. It has
Geologists, it
and Pennsylv
nized by its
brownish lime
small *Orthocer*
Waterlime, wh
New York.
Onondaga, is

§ 112. Th
quarries usual
odor when str
gas supplied b
salt manufact
City, rock-salt
surface, which
afterward pur
been formed a
York, Michiga

the Straits of Mackinac, where it forms the island and the points of the main land. The thickness on the peninsula of Michigan does not exceed 50 feet. It is broken up in a ridge extending west from the west end of Lake Erie near the southern line of Michigan, where it is much thicker, and again at Put-in Bay Island, and at Sandusky and other places in Ottawa County, Ohio, and may be seen on the western and south-west anticlinals, which pass through Wood County, and as far south as Delaware and Pike. The thickness in Ohio has not been accurately ascertained, but including the Waterlime, which is not separable, the thickness is several hundred feet. It has been identified in Missouri, varying from 10 to 75 feet in thickness. It does not occur in Wisconsin or Iowa, and is unknown south of Pennsylvania in the Appalachian system. The composition of the rocks indicates shallow water; but as there is no conglomerate, it does not appear as a shore deposit.

§ 110. It is not very fossiliferous at any locality, and generally fossils are extremely rare. In addition to the two species mentioned as common to the lower part of it and the Guelph, *Orthoceras sublaeve*, *Euomphalus sulcatus*, and *Avicula triquetra* were early described from Wayne County; but the indistinct forms of *Spirifer*, *Atrypa*, and *Cornulites* remain without specific names.

§ 111. The Waterlime takes its name from the earthy, drab-colored limestone used for making hydraulic cement, and is regarded by some as a distinct Group, while the Canadian Geologists regard it as the lower member of the Lower Helderberg. It has its characteristic minerals and fossils; but, following the New York Geologists, it is here treated as the upper member of the Onondaga. In New York and Pennsylvania its thickness is from 30 to 300 feet, and is well-defined and recognized by its mineral nature, its fossils and position. In Eastern New York a brownish limestone, often mottled, containing corals, fragments of crinoids, and small *Orthoceras* forms the base of it. All the species of *Pterygotus* belong to the Waterlime, while *Eurypterus remipes* and *Pterinea rugosa* are characteristic of it in New York. The species which has the greatest geographical distribution in the Onondaga, is that peculiar form called *Pleurodictyum problematicum*.

§ 112. The whole Group contains more or less carbonaceous matter, and the quarries usually smell of petroleum, and the limestone generally gives up the odor when struck with a hammer. This Group is the source of a large part of the gas supplied by the gas-wells of Ohio and Indiana. It is the chief source of the salt manufactured in New York and in Michigan. On the St. Clair River, at Marine City, rock-salt occurs in a mass, extending from 1,633 feet to 1,748 feet below the surface, which is mined by forcing fresh water down into it to take up the salt, and afterward pumping the brine and evaporating it. Thick masses of rock-salt have been formed at various other places in this Group within the salt districts of New York, Michigan, and Ontario.

CHAPTER XVII.

LOWER HELDERBERG GROUP.

§ 113. This Group was named from the Helderberg Mountains, and defined by Hall in 1859, in the third volume of the Paleontology of New York. The lower member is a thin-bedded, often thinly laminated, dark-blue limestone, resting on the Waterlime-beds called Tentaculite limestone. The second member is a thin limestone full of *Stromatopora*, followed by a dark-gray concretionary limestone, in irregular layers, charged with *Pentamerus galeatus* and other fossils, which has a maximum thickness in Otsego County of 80 feet, and is called the *Pentamerus* limestone. The third is a blue, drab-weathering, calcareous shale and blue limestone, full of *Spirifera macropleura* and other fossils, having a maximum thickness in Albany County of 70 feet, called the Delthyris or Catskill Shaly limestone, from Catskill Creek, near Madison, Greene County. The fourth member is a light-gray limestone, full of broken Encrinites, having a thickness of 25 feet. And above this there is a bluish-gray limestone, charged with Brachiopoda, called the Upper *Pentamerus* limestone. These local subdivisions are not recognized at any distance from the Helderberg Mountains, nor does the Group occur in Western New York or Western Canada. Strata of this age occur in two or three small outliers in the great basin near Montreal, at the distance of 200 miles from the nearest exposure of the Group in New York. The most important of these is on the Island of St. Helen's, opposite Montreal. The Group, however, is quite largely developed in the Eastern Provinces, where it includes part of the Gaspé limestones. It is exposed on both sides of the Hudson River, and forms the outlier known as Becrafts Mountain, and appears in Maine and New Hampshire. Its maximum thickness in New York is about 400 feet, and nearly as much in Maine, while at Gaspé it is 2,000 feet. It extends southwardly to Tennessee, having a thickness in Pennsylvania of 1,400 feet, in Virginia 1,000 feet, in New Jersey 150 feet, and in Tennessee 100 feet. It has been identified at Cape Frazier in latitude 80°.

§ 114. This is an important Group on the eastern part of the continent, but does not occur west of the Appalachian system, which is in striking contrast with the Onondaga, that spreads out westerly from New York instead of southerly. It abounds in limestone strata, and the evidences of marine life, the latter apparently succeeding that of the Niagara age, by gradual change and development. Crinoids, Corals, Bryozoans, Brachiopods, Gasteropods, Lamellibranchs, and Crustaceans were abundant, but we have no evidence that a vertebrate land or fresh-water animal had yet made its appearance on this continent. The evidence of swamp or air vegetation is on the increase, and here we discover the genus *Annularia*, which subsequently became so abundant in the Coal Measures. The characteristic fossils are: *Tentaculites gyraanthus*, *Spirifera macropleura*, *S. vanuxemi*, *Eatonia singularis*, *E. medialis*, *Pentamerus galeatus*, *P. pseudogaleatus*, *Streptorynchus radiatum*, *Strophonella punctulifera*, *Meristella laevis*, *Rhynchonella semiplicata*, *R. ventricosa*, *Strophodonta varistriata*, *Avicula naviformis*, *A. manticula*, *Beyrichia granulata*, and *B. notata*.

§ 115. Petroleum springs occur on the St. John's River and on Silver Brook,

in the Gaspé which has a peculiar odor is from the fossil good well has § 116. stratigraphic System as the overlying, represented

§ 117. T England. Its fossils in ascending order Group, Port

§ 118. It principally of at all places, been regularly in New York rine deposits. within the sea gan to assume we now behold libbranches, and was a long and organisms that genera and species presented a strong alence of Gym the two Systems The masses and rocks older than carbonaceous strata of rocks, and, continent. Ve different States exposed sandstone

in the Gaspé series, and in cavities of an amygdaloidal greenstone at Tar Point, which has hardened in some instances to the consistency of pitch, and from its peculiar odor the name Tar Point was given to the locality. The source of this oil is from the fossiliferous rocks or shales beneath, and exudes from an anticlinal. No good well has, however, been discovered by boring in these rocks.

§ 116. With this Group the Upper Silurian closes, because we have another stratigraphical and palæontological chasm, and have arrived at the top of the System as established by Murchison. The absolute want of conformability, with the overlying rocks, is everywhere apparent, and an age of time is therefore unrepresented in the geological column.

CHAPTER XVIII.

DEVONIAN SYSTEM.

§ 117. The Devonian was named in 1837, by Murchison, from Devonshire, in England. It has greater thickness, and is capable of more subdivisions based upon its fossils in this country than in any other part of the world. It is subdivided in ascending order as follows: Oriskany Group, Upper Helderberg Group, Hamilton Group, Portage Group, Chemung Group, and Catskill Group.

§ 118. It commences with a sandstone formation, after which it consists principally of limestone and shales. It is unconformable with the Upper Silurian at all places, except possibly Gaspé, Canada, where the sediment seems to have been regularly deposited from one age to the other. Its greatest development is in New York and Pennsylvania, where mechanical detritus accompanies the marine deposits. During this era land-plants became abundant, and fish swarmed within the seas, while the Archipelago, which had existed in the Silurian era, began to assume somewhat the outlines of a continent, though by no means such as we now behold. Corals, Crinoids, Brachiopods, Gasteropods, Cephalopods, Lamellibranchs, and Crustaceans were abundant, while Cystideans became extinct. It was a long and glorious era, marked by more progress in animal and vegetable organisms than characterized earlier ages. The plants increased in number of genera and species from the Lower to the Upper Devonian, until the flora presented a strong resemblance to that of the Subcarboniferous, especially in the prevalence of Gymnosperms and Cryptogams, though very few species are identical in the two Systems. It is everywhere unconformable with the Subcarboniferous. The masses and dykes of intrusive granite in Nova Scotia, which penetrate all the rocks older than the Subcarboniferous, belong to the close of the Devonian. The carbonaceous shales of this System exceed in thickness those of any other System of rocks, and, as a result, they are the chief oil and gas producing rocks on the continent. Very valuable iron ores and manganese ores occur in this System in different States. In Virginia huge masses of manganese are found imbedded in exposed sandstone ledges, where the supply seems to be practically inexhaustible.

CHAPTER XIX.

ORISKANY GROUP.

§ 119. This Group was defined as the Oriskany sandstone by Vanuxem, in 1839, and named from the white sandstone occurring at the Falls of the Oriskany, in Oneida County, N. Y., where it is about 20 feet in thickness. It forms a narrow belt of rough sandstone from the Hudson to Cayuga Lake, charged with peculiar fossils, and varying from a few inches to 30 feet in thickness. It stretches south in the Appalachian region through Pennsylvania, Maryland, and Virginia, and has a thickness in Pennsylvania of 300 feet. It appears in New Jersey with a thickness of 130 feet. In Maine there is a large exposure between Parlin Pond and Aroostook, and it exists at Gaspé and in Nova Scotia. It is known in Canada at but few places, one of the principal exposures being at North Cayuga, and covering only 230 acres. In Southern Illinois it is underlaid with silicious limestone, called the Clear Creek limestone, which constitutes incomplete passage-beds from the Upper Silurian. It is also known in Missouri.

§ 120. It appears as a belt deposited upon the shores of the islands which then existed, and to mark their outlines in a greater or less degree. Like other arenaceous deposits, it indicates the presence of land and shallow water. It abounds in the casts of Brachiopods and Gasteropods in New York, Maryland, and Virginia, and in some places Crinoids occur. The characteristic species are *Spirifera arenosa*, *S. arrecta*, *S. pyxidata*, *Rensselaeria ovoides*, *Orthis proximus*, *O. muscosa*, *Strophodontia magniventra*, *S. magnifica*, *Cyrtina rostrata*, *Eatonia peculiaris*, *Leptocælia flabellites*, and *Platystoma ventricosum*. In some places in Virginia the shells are silicified and quite free from adhering matter, and the exterior markings and internal structure are well preserved, even the internal coils of Brachiopoda are beautifully represented. Near Cumberland, Md., a few elegant crinoids have been found, and one Cystidean, *Anomalocystites disparilis*, which is the latest known representative of that order, except *Strobilocystites calvini*.

§ 121. The Brachiopods are Devonian in their character rather than Silurian, and there is graduation to the succeeding rocks through the Cauda-galli grit, which is a dark, gritty slate, bearing few fossils. The rocks are not such as to have preserved land-plants very well; but they should have preserved fish-teeth if any then existed, but no trace of them has been discovered.

§ 122.

divided into
erous limes
great geog
erous forma
ogical word
gritty slate
grit, which
in the east
thin out an
bony plates
subcrystallin
which bears
suggested th
these two d
chert, or ho
or protophy
pods, and o
is about 300

§ 123.

sula of Cana
into Michiga
western Ohio
the Scioto, r
State. It c
Ohio River
resting on th
serving the
exceeding a
galli grit has
ing a total t
is one of the

§ 124.

I
corals, and
specimens be
earlier period
or the outer
of layers, 3
and singularly
in length con
tubercles. C
drifted land-p

CHAPTER XX.

UPPER HELDERBERG GROUP.

§ 122. This Group was named from the Helderberg Mountains, where it was divided into the Cauda-galli grit, Schoharie grit, Onondaga limestone, and Corniferous limestone. The Corniferous limestone being the only one which has any great geographical distribution, the Canadian Geologists in 1863 used "Corniferous formation" instead of Upper Helderberg; but as Corniferous is a mineralogical word, Upper Helderberg is to be preferred. The Cauda-galli grit is a dark gritty slate covered with *Taonurus cauda-galli*, and graduates into the Schoharie grit, which is an arenaceous limestone weathering to a brownish color. These occur in the eastern counties of New York, Albany, Greene, and Schoharie, but soon thin out and are not found west as far as the center of the State. The scales and bony plates of fish are first found in the Schoharie grit. The Onondaga is a gray subcrystalline, coralline limestone. It is followed by the Corniferous limestone, which bears dark-colored, cherty beds, that break with a horny fracture, which suggested the name Corniferous; but the cherty beds occur in various places in these two divisions, and there is no real line of separation between them. The chert, or hornstone, is largely composed of microscopic, silicious forms of plants or protophytes, spiculae of sponges, fragments of the dental apparatus of Gastropods, and other organisms. The aggregate thickness of the Group in New York is about 300 feet.

§ 123. From New York the Group extends in a belt west across the peninsula of Canada to Mackinac Island, where it is 250 feet thick, and from thence into Michigan where its thickness is 354 feet. It appears at Sandusky and Northwestern Ohio, at Columbus, and on the Ohio a few miles below the mouth of the Scioto, resting upon the Waterlime Group, which has great thickness in this State. It crosses into Northern Indiana, and striking south-westerly, crosses the Ohio River at Louisville. It appears in Illinois, Iowa, Missouri, and Tennessee, resting on the Oriskany, or the Waterlime, or the Niagara, and everywhere preserving the character of the great coral-reef period of the Devonian, but never exceeding a thickness of about 300 feet. In New Jersey, however, the Cauda-galli grit has a thickness of 400 feet, and the Corniferous limestone 500 feet, making a total thickness of 900 feet. It occurs in the western mountain ranges, and is one of the most persistent and generally distributed Groups.

§ 124. It is a marine limestone, distinguished for the remarkable abundance of corals, and coral reefs, the variety in form, number, and size of species, some specimens being several feet in diameter, and larger than any belonging to any earlier period. It is distinguished also for its fish remains, which consist of teeth, or the outer bony covering, sometimes so abundant as to constitute the major part of layers, 3 or 4 inches, or even more, in thickness. Some were very large and singularly constructed. The *Macropetalichthys sullivanii* had a head 15 inches in length composed of hard, bony plates, covered with a thick skin dotted with tubercles. Cephalopods are abundant and quite characteristic, and in a few places drifted land-plants have been found, but they are not of general occurrence. The

most characteristic species among the invertebrates, and those by which the rocks may be readily identified are *Cyathophyllum rugosum*, *Favosites goldfussi*, *Syringopora maclurii*, *Phillipsastrea verneuili*, *Nucleocrinus verneuili*, *Spirifera acuminata*, *S. gregaria*, *Pentamerus knighti*, *P. aratus*, *Stricklandinia elongata*, *Paracyclas occidentalis*, *Conocardium subtrigonale*, *Platyceras dumosum*, *Tentaculites scalariformis*, and *Dalmanites selenurus*. In the vicinity of Davenport, Iowa, it furnishes an abundance of durable and massive building material and contains cavernous openings, as if worn out by the action of water, and filled up subsequently with material derived from higher rocks, and especially those of the Hamilton Group. The quarries at Columbus, Ohio, and North Vernon, Indiana, are in this Group. The strata in the vicinity of the Straits of Mackinac have been eroded and excavated so as to produce the Island of Mackinac, and large masses of the materials have been transported and distributed over Southern Michigan and Ohio.

§ 125. The limestones of this Group in Canada are usually bituminous, and petroleum frequently fills the cells of corals and other fossils. The corals often prevail in distinct bands, some of which will be saturated with the oil, while others will not. Petroleum springs rise from this Group at Tilsonburg, and other places along an anticlinal which runs through the Western Peninsula. The oil being lighter than water, and permeating the strata, naturally rises to the highest part of the anticlinal between the impervious layers of rock, and escapes to the surface. In other localities the bitumen is solid, and takes the form of asphaltum or mineral pitch, as at Kincardine, where slaty beds contain from 10 to 15 per cent of bitumen soluble in benzole. No good well, however, has been discovered in Canada by boring in these rocks, though it has been contended the oil at Enniskillen and on the Thames has its source here. Where the oil has been found in this Group, it has had its source in the Waterlime or in the shales below.

CHAPTER XXI.

HAMILTON GROUP.

§ 126. This Group was named from Hamilton, Madison County, New York, and defined by Vanuxem in 1842, though he did not include within it the Marcellus Shale, Tully Limestone, and Genesee Slate. The divisions made for it in New York are Marcellus Shale, Ludlowville Shale, Enderinal Limestone, Moscow Shale, Tully Limestone, and Genesee Slate. The rocks are not susceptible of this division, except locally, and they all belong to a single Group. The Marcellus Shale was named from Marcellus, where it is an argillaceous slaty rock, bearing much carbonaceous matter, and sometimes small pieces of coal, and has a thickness of about 200 feet. It contains layers of impure limestone, and abounds in fossils. In many places it contains so much bitumen as to give out flame when thrown into the fire, which led the early settlers to explore it throughout its whole extent for coal, only, of course, to suffer disappointment. It is not separable from the Ludlowville Shale by any well-defined characters. The Ludlowville Shales were named from the town of that name, and separated from the Moscow Shale by a layer of limestone 3 or 4 feet thick, called the Enderinal limestone; but such

division is from 300 to 400 feet thick. The rock has a thick, fissile mass, Group in N.

§ 127. of variable thickness in feet in the excavated face afford the best stones in the State. The wave-lines, Fucoids and good size, and as to show more in the extent and the Gro Wisconsin, Milwaukee, the manufacture Helderberg and Falls of the Iowa, and also on the Mackinac thickness in the Silurian chain ment. In the remains, while

§ 128. its invertebrates *Lepidodendron* shales. The species are distributed usually be distributed *Tropidoleptus limitare*, *L. recurva*, *Pterin sulcomarginata*

§ 129. The known to the upon the surface sinking through considerable s

division is scarcely worthy of recognition. The three have a thickness varying from 300 to 900 feet, extend from Lake Erie to the Hudson, and abound in fossils. The Tully limestone was named from Tully, where it is burnt for lime, and has a thickness of 14 to 20 feet. The Genesee slate, named from the opening of the gorge of the Genesee River at Mount Morris, where it is a black, argillaceous fissile mass, attains a thickness of 150 feet and closes the era of the Hamilton Group in New York.

§ 127. The Group extends from the Hudson to Lake Erie, occupying a belt of variable width in the central part of the State, and attaining a maximum thickness in the eastern part of 1,200 to 1,400 feet, and diminishing to about 300 feet in the western part. The valleys of Seneca and Cayuga Lakes are excavated for more than half their length in these rocks, and the banks and ravines afford the best facilities for examination. It is an olive shale, with slates and sandstones in the eastern, and calcareous shale and limestone in the western part of the State. The bedded rocks are remarkable for the abundance of ripple-marks, and wave-lines, and the shales abound in carbonaceous material, due to vegetation. Fucoids and marine plants are common, and coniferous trees and ferns grew to a good size, and drifted into the ocean, where they were imbedded and preserved, so as to show much of their form and structure. The New York subdivisions are lost in the extension across the peninsula of Canada from Lake Erie to Lake Huron, and the Group becomes a limestone in Michigan. It occurs at only one place in Wisconsin, which consists of a strip about 10 miles long and 5 or 6 wide, near Milwaukee, where it is an impure limestone, quite fossiliferous, and largely mined for the manufacture of hydraulic cement. It occurs in Ohio, resting on the Upper Helderberg as far south as Columbus, and the upper part of the limestone at the Falls of the Ohio, is referred to it. It occurs at Davenport and New Buffalo, in Iowa, and also in Illinois and Missouri. It appears among the western mountains, on the Mackenzie River, in Alaska, and in the Arctic regions. It has greater thickness in Pennsylvania, New Jersey, Virginia, and other States in the Appalachian chain, than it has in the West, and contains much more mechanical sediment. In the East it is a mud rock supplied with drift materials and marine remains, while more westerly it is exclusively a marine calcareous rock.

§ 128. It is of quite general distribution and usually readily determined by its invertebrate fossils, which exceed in number almost all earlier Groups. *Lepidodendron*, which became so common in the Coal Measures, is found in the shales. The remains of fish are much like those of the Upper Helderberg, though species are distinct. The characteristic fossils, and those by which the Group may usually be determined, are *Helicophyllum halli*, *Spirifera pennata*, *S. grandifera*, *Tropidoleptus carinatus*, *Rhynchonella venustula*, *Athyris spiriferoides*, *Leiorhynchus limitare*, *L. quadricostatum*, *Orthonota undulata*, *Cypriocardella bellistriata*, *Cimitaria recurva*, *Pterinea flabellum*, *Modiomorpha concentrica*, *Bellerophon patulus*, *Neurotomaria sulcomarginata*, *Styliola fissurella*, *Homalonotus dekayi*, and *Phacops bufo*.

§ 129. The oil-springs of Enniskillen: and of the Thames, in Canada, were known to the Indians and to the settlers from an early period. The oil floated upon the surface of the waters, and formed by its drying beds of tarry bitumen. On sinking through the clay from 40 to 60 feet, a bed of gravel is reached, from which considerable supplies of petroleum are obtained. Such are called surface-wells,

and are less productive than the deeper ones. Below the gravel thin limestones, shales, and clays occur for a distance of about 230 feet before the Upper Helderberg limestones are reached. One of these wells, when sunk to a depth of 200 feet below the surface, yielded, when first opened, 2,000 barrels of oil in twenty-four hours. In some of the wells bored in this vicinity, both oil and water flowed to the surface, and in some of the deeper ones the water is saline. Wells bored into the Upper Helderberg limestone sometimes reached small quantities of oil, but no valuable wells have thus far been discovered in Canada by boring below the Hamilton Group. The flowing wells soon become intermittent, and within a year cease to flow altogether; they continue, however, to furnish oil by pumping for a limited period, and then appear to be exhausted. The petroleum differs in volatility; the less volatile contains paraffine in solution, and is suited for lubricating machinery, while the more volatile is best suited for light. The alliaceous odor of some of the unrefined oil is due to the presence of a little sulphureted hydrogen. Petroleum is modified on exposure to the air by volatilization and oxidation, and eventually assumes a solid form. Thus near Oil Creek, in Enniskillen, the thickened oil formed two layers, called gum-beds, of a viscid, tarry consistence, covering two or three acres with a thickness from a few inches to two feet. In sinking a well, a bed of this asphaltum, from 2 to 4 inches thick, was met with at a depth of 10 feet, upon a layer of gravel. It contained the remains of leaves and insects, which were imbedded in it during its slow accumulation and solidification. In boring the oil-wells there is always a greater or less disengagement of inflammable carbureted hydrogen-gas, and sometimes it is liberated with explosive violence. The strata almost everywhere in that region hold in a condensed state portions of light carbureted hydrogen, which is discharged wherever a natural fissure or an artificial boring furnishes a vent. The shale on Sulphur Island, at the mouth of Thunder Bay in Lake Huron, is so highly charged with bituminous matter that it has been set on fire and burned for months. The bitumen burns out and leaves the shale with a reddened appearance.

CHAPTER XXII.

PORTAGE GROUP.

§ 130. This Group was named from Portage, New York, and defined by Hall in 1843. It consists of variable shales and sandstones, forming in New York an east and west band, resting upon the Hamilton Group, and dipping south about 25 feet in a mile. The sandstones produce falls in the streams, beautiful cascades, and grand and striking scenery. The highest perpendicular fall of water and deepest canons and gorges in the State exist in this Group. It thickens westerly and thins easterly, and does not extend to the extreme eastern part of the State. Sandstones greatly predominate in the eastern part, while shales increase westerly, until the whole Group becomes a mass of black, bituminous shale. The thickness on the Genesee is 1,000 feet, on Lake Erie 1,400 feet. A considerable part of Lake Erie is excavated out of this Group, which shows a belt on the south side extending nearly to Sandusky; and from here it bends southerly across Ohio, leaving Columbus to the west, and, reaching the Ohio River below the mouth of the Scioto,

it crosses in
ranges. It
sula, and
the Huron
and enters
reaching the
toward Ten
Ohio for 18
Indiana, and
in Indiana
been recogn
to the Appa

§ 131.
throughout
with those
he spent in
greatest abu
which are of
lavia occur in
istic. Fish
teeth strong
a carnivorous
mian. It wa
of any other
seas or arms
large geograph
local in their
limestone, son
monster *Dini*

§ 132. T
posed to be th
New York, H
rocks until th
supply. Ten
shale yields
and in those
vania belong
of this Group
The gas at F
1820. Lyell
with little var

it crosses into Kentucky, and is soon broken up in the spurs of the mountain ranges. It crosses Lake Erie, and occupies a small part of the Canadian peninsula, and enters the southern peninsula of Michigan, where Winchell called it the Huron Group. From Michigan it crosses the north-western corner of Ohio, and enters Indiana, forming a belt across that State by way of Indianapolis, and, reaching the Ohio River at New Albany, crosses into Kentucky, and extends far toward Tennessee. It was called the Black Shales in the Geological Survey of Ohio for 1838, and in that of Indiana for 1839, and in later surveys of Kentucky, Indiana, and Tennessee. The thickness in Ohio is from 200 to 1,000 feet or more, in Indiana from 100 to 200, and in Tennessee from 10 to 150 feet. It has never been recognized west of these States, and is therefore classed as a Group belonging to the Appalachian mountain system.

§ 131. Fucoids, wave-lines, and ripple-marks are numerous, and occur throughout its distribution. The paucity of fossils in this Group, when compared with those above and below it, is one of its striking characters. Whole days may be spent in some parts of it without finding a shell, though fucoids are in the greatest abundance. Land-plants occur in profusion in New Brunswick, some of which are of gigantic size. *Goniatites complanatus*, *Panenukka speciosa*, and *Spirifera levis* occur in New York and in Ohio, and may therefore be considered characteristic. Fish of large size, covered with thick heavy plates, and having jaws and teeth strong enough to crush a body the size of a man, occur in it. *Cladodus*, a carnivorous fish, became abundant in this period, and flourished until the Permian. It was world-wide in its distribution, and its vertical range exceeds that of any other genus of fishes. The Group seems to have been deposited in internal seas or arms of the ocean, and is the last Group of the Devonian System, having a large geographical distribution, for the Chemung and Catskill are comparatively local in their extension. In Ohio there are large concretionary balls of impure limestone, some of them several feet in diameter, and it was in one of these the monster *Dinichthys* was discovered.

§ 132. The Group is distinguished as the great seat of petroleum, and is supposed to be the source from which the chief supply in this country is derived. In New York, Pennsylvania, and Ohio the wells are bored through the overlying rocks until the Portage is reached, or the saturated sands that overlie it furnish the supply. Ten per cent of the shales is bituminous and carbonaceous matter. The shale yields oil by distillation, and gas and oil springs abound in its sandstones, and in those which overlie it. The great oil-sands in the oil regions of Pennsylvania belong to the Chemung, and have doubtless been fed as well from the shales of this Group as from those of the Chemung, which furnish the same products. The gas at Fredonia, New York, in this Group, was used for lighting houses in 1820. Lyell described it in his travels in 1841, and it has been in constant use, with little variation in the supply, ever since.

CHAPTER XXIII.

CHEMUNG GROUP.

§ 133. This Group was named from the exposure at the Chemung Upper Narrows, at Chemung, New York, and defined by Vanuxem and Hall in 1842 and 1843. The shale and sandstone at Ithaca, having a thickness at Hector's Falls of 400 feet, was called the Ithaca Group, but it is only part of the Chemung. The Chemung consists of a highly fossiliferous series of shales and thin-bedded sandstones and impure limestones, and an infinite variety formed from admixture of these. Except in a few localities there is no marked line between it and the Portage below. The two are distinguished by their fossils. The shales vary in color from a deep black to olive-green, with every grade of intermixture; the sandstones are gray, olive, or green, and almost the whole series weathers to a brownish olive. The Group forms an east and west belt across the southern part of New York, having a thickness in the eastern part of 2,000 feet, dipping southerly at 25 feet or more to the mile, and thinning westwardly, so as not to be determined a short distance from where it crosses the line of Ohio. It is unknown farther west. In its extension from Eastern New York into Pennsylvania the thickness increases until it exceeds 3,000 feet. It occurs at New Brunswick and at Gaspe, Canada, but has not been satisfactorily determined at many other places, though it probably occurs in many other regions of the Appalachian system. The rocks which have been called Chemung in Ohio, Indiana, Illinois, Missouri, Iowa, and Michigan belong to the Waverly, except the thin, tapering belt in North-eastern Ohio, already mentioned.

§ 134. The alternations and interlaminations of shales and sandstones show deposition under similar circumstances to those under which the Portage was deposited. The source of the materials was to the east or south-east of New York, as evidenced by the thinning of the deposits and diminution of sandy strata toward the west. The land-plants occur in Eastern New York, and disappear westerly, proving the land existed in that direction. The marine and land plants are abundant in the sandstones, while marine shells increase with the decline of the sandstones and augmentation of the shales westerly, though fucoids continue in abundance wherever the Group exists. The plants foreshadow the approaching Carboniferous System by the presence of *Archæopteris*, *Cyclopteris*, *Sigillaria*, *Lepidodendron*, and *Trigonocarpus*. The fauna has more of a Carboniferous aspect than any which preceded it, and there is a diminution of the types which characterized the earlier Devonian. The species having the greater distribution and most characteristic are *Lepidodendron chemungense*, *Archæopteris lucæ*, *Asterophyllites parvulus*, *Orthis impressa*, *Orthis tioga*, *Streptorhynchus chemungense*, *S. pectinaceus*, *Strophodontia caryata*, *S. mucronata*, *Chonetes muricata*, *Productella hirsuta*, *Spirifer disjuncta*, *S. mesacostalis*, *Atrypa dumosa*, *A. hystrix*, *Aviculopecten duplicatus*, *A. rugistriatus*, *Leptodesmus longispinus*, *L. spinigerum*, *Leiopteria chemungensis*, *Pterinopecten dispandus*, *P. crenicostatus*, *P. suborbicularis*, *Pterinea consimilis*, *Crenipecten crenulatus*, *Mytilarca chemungensis*, and *Phacops nupera*.

§ 135. Springs, evolving carbureted hydrogen-gas, or gas accompanied with petroleum, are common throughout nearly all that part of New York and Pennsyl-

vania cover
minous odo
and gas pr
the oil had
rived from
taining earl
possible the
making the
both formed
organisms,
evidence als
for the shale
the greater
are porous
roleum, whi
are bored un
the Chemung
as well as fr
with anticlin
supported wi
much less ha
Wells are as
rock is struck

§ 136. Th
quite fully def
conglomerates,
portion is bric
shales are gra
eastern New Y
feet, and dips
feet, and soon
New York, and
It is conformal
lithology, and
been described
erally very po
the rocks belon
though these ar
Old Red Sands
is cemented and
and strata uneq

vania covered with the Chemung. The rocks in nearly all localities emit a bituminous odor on percussion, and petroleum often exudes from the crevices. The oil and gas products are the same in the Chemung as in the Portage. The gas and the oil had the same origin. They are both hydrocarbons. They were both derived from vegetable and animal organisms. Wherever shales are found containing carbonaceous matter, evidence of these products may be obtained. It is possible the gas was first produced, and from it the petroleum has been derived, making the latter a secondary product; but the evidence seems to prove they were both formed at the same period of time, and during the decomposition of the organisms, and before the mud had indurated or hardened into rock. And the evidence also seems to prove they were derived almost wholly from marine plants, for the shales bearing the greater number of fucoids are those to which we ascribe the greater supplies of hydrocarbons. The sandstones which overlie these shales are porous and capable of holding from one-eighth to one-tenth their bulk of petroleum, which is sufficient to account for the flowing wells of Pennsylvania which are bored until they penetrate the sandstone. Many of the wells penetrate only the Chemung sandstone, though the oil is derived from the shales of the Portage as well as from the Chemung. The supposed connection of petroleum and gas with anticlinal axes, or synclinal ones, has not been verified by observation, nor supported with reason, neither are they dependent upon faults or crevices, and much less has the depth of the well any connection with the level of the sea. Wells are as valuable when bored below the sea level as they are when the proper rock is struck above that horizon.

CHAPTER XXIV.

CATSKILL GROUP.

§ 136. This Group was named by Emmons from the Catskill Mountains, and quite fully defined by Vanuxem in 1842. It consists of sandstones, shales, slates, conglomerates, and impure limestones. The prevailing color of the arenaceous portion is brick-red, though all of it is more or less colored with iron, and the shales are gray, olive-red, or green. It exists only in a few counties in South-eastern New York, in the Catskill Mountains, where it has a thickness of 3,000 feet, and dips rapidly toward Pennsylvania, where it reaches a thickness of 7,500 feet, and soon disappears. It does not extend west of the Genesee Valley in New York, and is wholly unknown on any part of the continent west of that State. It is conformable with the Chemung, and is distinguished only by the change in lithology, and by the fossils. No Corals, Crinoids, Brachiopods, or Trilobites have been described from it, and only a few Lamellibranchs. The land-plants are generally very poorly preserved. The fish remains are relied upon to really prove the rocks belong to the Devonian rather than to the Subcarboniferous age, and though these are rare and poorly preserved, they show it is the equivalent of the Old Red Sandstone of England, and therefore Devonian. In some places the sand is cemented and forms a grindstone grit, and there are hard concretionary masses, and strata unequally hardened, that weather into picturesque rocks. The Group is

§ 139. In Nova Scotia the lower part is called Lower Coal Measures, and the upper part Lower Carboniferous Marine Formation, or more generally the whole is called Lower Carboniferous, even where its thickness is 6,000 feet. It consists of sandstones, shales, conglomerates, and limestones, with beds of gypsum. The limestones bear Brachiopods specifically identical with those of corresponding age in the Illinois basin. In Pennsylvania and in Nova Scotia thin seams of coal occur in the strata, which is not the case farther west. On Cape Breton the thickness is 4,600 feet. In the Rocky Mountain region there is a thickness of 4,000 to 7,000 feet or more, and the several Groups may be determined at different places. The System has been divided in the west into the Lodore Group, Tonto Group, Red Wall Group, Lower Aubrey Group, and Upper Aubrey Group. Prof. Dawson found no paleontological or stratigraphical reason for regarding the Subcarboniferous as a System distinct from the Carboniferous, but as it is generally capable of subdivision into Groups, is always unconformable with the Devonian, begins with a sandstone, and is followed by a conglomerate or sandstone unconformable with it, there is good reason for retaining the name, though if the lines were not better defined elsewhere than in Nova Scotia, we might join Prof. Dawson in discarding it.

§ 140. There are some fossils in this System almost world-wide in distribution, and belonging alike to all the Groups into which it has been subdivided; viz., *Spirifer striata*, *Athyris lamellosa*, *A. planosulcata*, *Orthis michelini*, *O. resupinata*, and *Productus semireticulatus*. There are some that occur in the rocks of this age in each of the Coal-basins on this continent; as, *Athyris subtilita* and *Productus cora*. It is in this System at Hillsborough, New Brunswick, the bituminous mineral Albertite is so abundant. The rocks are thin-bedded shales, composed of fine, indurated clay, with much bituminous matter, and are full of fossil fishes in a good state of preservation. The shales have been disturbed and contorted, and contain the vein of asphaltic mineral called Albertite. The theory of its creation is as follows: The argillaceous mud which formed the indurated shales, was charged with finely comminuted vegetable matter, which in its decomposition furnished the petroleum that at some later age escaped into a vein or fissure in the rocks, and by losing its more volatile parts and partial oxidation, it hardened into the coaly or asphaltic substance. No extra heat for such transformation was necessarily required. Springs yielding petroleum flow from these rocks in various places. Peroxide of manganese, used in bleaching and in gas manufacture, occurs in limestone near the base of the System, and wad or black manganese ore is abundant at different places. Alum frequently occurs from the spontaneous weathering of pyritous shales, and is sometimes manufactured from them. Saline springs are not uncommon; indeed, they are numerous from the commencement of the Upper Silurian rocks to the close of this System, and occur occasionally both above and below such range. The conglomerate on the Stewiacke, Musquodoboit, and St. Mary's Rivers, is auriferous. It was formed from auriferous quartz-veins, derived from the Taconic System, and gold occurs in it exactly as in modern auriferous gravels, being found in the lower part of the conglomerate, and in the hollows and crevices of the underlying unconformable rocks. The rocks of the age of this System in Europe are commonly known as the Mountain Limestone.

CHAPTER XXVI.

WAVERLY GROUP.

§ 141. This Group was named in 1838, by Mr. C. Briggs, an assistant geologist on the Ohio Survey, from Waverly, Ohio, where it consists of a fine-grained sandstone, about 300 feet in thickness, superimposed upon a black argillaceous slate 200 or 300 feet thick, and is followed by from 40 to 80 feet of conglomerate. He identified the rocks at Portsmouth, Piketown, and Chillicothe. Mr. J. W. Foster, another assistant, followed them through Licking and Fairfield Counties. In 1839 David Dale Owen, after having examined the rocks in Ohio, found them in Indiana, Illinois, and Kentucky, and described the freestone knobs displayed back of New Albany as the Waverly Sandstone series, and referred them to the base of his Subcarboniferous System. Owen established this Group as a geological subdivision by a fair definition. Owen, Norwood, Pratten, and other Western geologists recognized the Group from that time forward. In 1841 Hubbard recognized the Group in the geological survey of Michigan. Hall and some Eastern geologists erroneously asserted the rocks were of Devonian age. In 1861, Meek and Worthen, having ascertained, upon paleontological evidence, the limestones at Rockford, Indiana, at Choteau, Missouri, and at Kinderhook, in Pike County, Illinois, belong to the base of the Subcarboniferous rocks, proposed to call them the Kinderhook Group. They understood they were making a synonym, but supposed they were including less in their Group than is included in the Waverly. In the same year Alexander Winchell described the Marshall Group of Michigan, and afterward thoroughly defined it, and proved its identity with the Waverly Group, the Kinderhook, the Yellow sandstone series of Iowa, and Choteau limestone, Vermicular sandstone and shale, and Lithographic limestone of Missouri.

§ 142. The Group in Ohio forms a belt from 10 to 20 miles in width, commencing near the mouth of the Scioto, and bearing north and north-east toward Cleveland, but widening as it approaches Lake Erie, until its width exceeds 40 miles. It rests upon the Portage Group, and has been called in its northern extension the Cuyahoga shale, Berea Grit, Bedford and Cleveland Shales. It crosses the Ohio from the Scioto, and entering Kentucky is soon broken up among the mountain ranges. In Indiana it forms a belt extending from New Albany north, by way of Rockford, and south across the Ohio River, by way of Danville and Knob Lick, Kentucky. The fossiliferous, greenish, mottled limestone at Rockford, so famous for its *Goniatites* is at the base of the Group. The maximum thickness in Indiana is 500 feet, in Kentucky 200 feet. In Michigan, at Marshall, Hillsdale, and other places, it consists of reddish, yellowish, and greenish sandstones, having a thickness of 160 feet, and the Napoleon sandstone, 123 feet in thickness. It furnishes large quantities of salt and gypsum. The brine is obtained by boring and pumping, and very large salt-works are established on the Lower Saginaw River. Salt has been largely manufactured from brine obtained from the rocks in Ohio. The celebrated Ohio freestone, so much used for building purposes, is from this Group.

§ 143. In Missouri, the Lithographic limestone has a thickness of 55 feet; is a fine-grained, compact limestone, breaking with a free, conchoidal fracture, and is

especially ch
thickness of
burrows. T
Choteau Spr
At Burlingto
and sandston
it disappears.
sists of grit-s
oolitic limesto
other places

§ 144. .
able in the sp
fish remains.
ductella conce
luris, Syringo
Centronella a
dianense, Gon

§ 145. The
Burlington lim
geologist seems
science, though
The limestone
of crinoidal rex
increases in sili
beds, which, w
Group. Hall
and others refe
dips below the
fine exposure
Mississippi, fro
and at Sedalia.
fossils received
Valley Mining
the great West.

§ 146. The
tained were it n
and this resulte
western localitie
ties. The detrit
the Appalachian
and other States

especially characterized by *Pentremites roemeri*. The Vermicular sandstone has a thickness of 75 feet, and is ramified with irregular perforations resembling worm-burrows. The Choteau limestone has a thickness of 100 feet, and was named from Choteau Springs, in Cooper County. It has an extensive geographical distribution. At Burlington, Iowa, the Group has a thickness of 77 feet, and consists of shales and sandstones, capped by a four-foot bed of oolitic rock. It thins northerly until it disappears. It has a thickness in Illinois of 200 feet, and at Kinderhook it consists of grit-stones, sandy and argillaceous shales, with thin beds of fine-grained and oolitic limestone. It has been identified in the Wahsatch Range, in Utah, and at other places in the great West.

§ 144. The fauna, on the whole, has assumed a Carboniferous aspect, noticeable in the species which pass to higher Groups, and more strongly in the genera of fish remains. Fossils having a wide distribution and characteristic species are *Productella concentrica*, *Productus cooperensis*, *Spirifera carteri*, *S. extenuata*, *S. peculiaris*, *Syringothyris halli*, *Athyris hannibalensis*, *Rhynchonella hubbardi*, *R. missouriensis*, *Centronella allii*, *Bellerophon cyrtolites*, *Grammysia hannibalensis*, *Orthoceras indianense*, *Goniatites oweni*, *G. marshallensis*, and *Phillipsia doris*.

CHAPTER XXVII.

BURLINGTON GROUP.

§ 145. This Group was named from Burlington, Iowa, where it was called the Burlington limestone before it was described as a geological subdivision. No single geologist seems to have established the Group, or to have introduced the name to science, though the first full definition is in the geological survey of Iowa for 1858. The limestone at Burlington is subcrystalline, often friable, and largely composed of crinoidal remains, has a thickness of 100 feet, and thins out northwardly. It increases in silicious matter toward the top, until the limestone merges into silicious beds, which, without evidence of unconformability, separate it from the Keokuk Group. Hall referred these cherty layers to the Keokuk, but White, Wachsmuth, and others refer them to the Burlington. In its southern extension, the Group dips below the bed of the Mississippi, and rises again at Quincy, and furnishes a fine exposure at Hannibal, Missouri. It exists in nearly every county on the Mississippi, from St. Louis to Iowa, and west from St. Charles to Howard County, and at Sedalia. The thickness varies from 100 to 500 feet. From a collection of fossils received from Prof. Cope, the author identified the Group in the Lake Valley Mining District of New Mexico; and it doubtless exists at other places in the great West.

§ 146. The separation of the Burlington from the Keokuk could not be maintained were it not for the great change in the specific characters, of the Crinoids, and this resulted probably from the deeper, or clearer, or less disturbed water in the western localities during the Burlington period, than existed in the eastern localities. The detrital material may have prevented the recognition of the Group in the Appalachian system, and rocks of the same age in Ohio, Kentucky, Indiana, and other States may be referred to the Waverly or the Keokuk. In no other

period did the harder parts of Crinoids so completely form the limestone, and hence it is pre-eminently the age of Crinoids. As the *Graptolida* reached the height of development in the Quebec or Upper Taconic, the *Orthoceras* in the Black River, and the *Cystidea* in the Niagara; so did the *Crinoidea* in the Burlington. The bed of the ocean was covered with a dense growth of Crinoids, one generation after another, while the superincumbent water swarmed with fish and invertebrate life. About 400 species of Crinoids, or one-fourth of all known, are from this Group. Among those having the greater distribution and being most characteristic are *Dorycerinus missouriensis*, *D. parvus*, *D. unicornis*, *Batoerinus christyi*, *B. pyriformis*, *B. rotundus*, *Actinocrinus proboscidealis*, *Platycrinus planus*, *Amphocrinus divergens*, *Belemnoocrinus typus*, *Strotoerinus regalis*, *Steganoocrinus concinnus*, and *Physetoerinus ventricosus*.

CHAPTER XXVIII.

KEOKUK GROUP.

§ 147. This Group was named from Keokuk, Iowa, where it was extensively quarried, and known as the Keokuk limestone, before it was known as a geological subdivision. It was first defined by Owen in 1852, and afterward by Hall in 1858. As defined by Hall, it consisted of fifty feet of fossiliferous limestone capped by 40 feet of shale, abounding in geodes of quartz, called the geode bed. Others refer the chert layers, which separate it from the Burlington, to this Group. It rapidly thins out to the north, but maintains its thickness southerly to the mouth of the Illinois River, and appears in the south-western part of Missouri, with a thickness of 200 feet, where it is a lead-bearing rock. It crops out in Indiana, 40 or 50 miles north-west of Crawfordsville, and extends southerly, crossing into Kentucky a short distance below New Albany. The thickness does not much exceed 100 feet. It is celebrated at Crawfordsville for the abundance and perfection of the Crinoids; entire specimens—roots, column, head, arms, and pinnules—have been collected. It is well displayed in Southern Kentucky, at King's Mountain tunnel, and in Tennessee, where the thickness is 200 feet. It occurs in Richland County, Ohio, and at other places on the western border of the Appalachian coal basin, but has not been described on the eastern border. It has been identified at numerous places in the western mountain ranges.

§ 148. Ores of lead and zinc occur in South-western Missouri in pockets and fissures associated with limestone and chert, and some of the mines are very rich and have been largely worked. In New Mexico and south of there, in Mexico, silver and lead occur in veins and fissures, some of the mines being very valuable. Some of the fossils having an extensive distribution, and being characteristic, are *Dorycerinus mississippiensis*, *Cyathocrinus multibrachiatus*, *Barycerinus hoveyi*, *Forbesioerinus wortheni*, *Platycrinus hemisphericus*, *Agaricoerinus americanus*, *A. wortheni*, *Actinocrinus lowei*, *A. pernodosus*, *Batoerinus biturbinatus*, *B. indianensis*, *Gonioeroidocrinus tuberosus*, *Cyathocrinus subtumidus*, *Palaacis compressus*, *Amplexus fragilis*, *Productus vittatus*, *Orthis keokuk*, *Spirifera keokuk*, *S. suborbicularis*, *Platyceras fissurellum*, *P. equilaterale*, and *Lithophaga illinoensis*.

§ 149. The more fully developed magnesian, and able with the a member of been described tances from Alton, Illinois County, Missouri, probably be re having great c *crinus simplex*, *mus*, *Productus* *nori*, *R. mutata* *sublaevis*, *Natic* *tomaria subglob*

§ 150. The ical Survey of splendid quarries thin layers of a forms bluffs bel Mississippi, but soon are 175 feet high Missouri and Io band of red ch crosses Kentucky palachian coal-f Cincinnati South limestones, more geodes, and havi in Kentucky or T cavernous, and a face, funnel-sha Wyandotte Cave 240 feet high, and

CHAPTER XXIX.

WARSAW GROUP.

§ 149. THIS Group was named from Warsaw, Illinois, by Hall, in 1856, and more fully defined in 1858. At the typical locality, near Warsaw, it consists of magnesian, arenaceous, and shaly limestones, abounding in Bryozoa. It is conformable with the Keokuk, only a few feet in thickness, and generally considered as a member of the Keokuk. I have retained it, because so many small fossils have been described from it, which have been the means of identifying it, at great distances from the typical locality. It occurs below the limestone of the cliffs at Alton, Illinois; at Bloomington and Spergen Hill, Indiana; and in St. Genevieve County, Missouri, where it attains its maximum thickness of 100 feet. It should probably be regarded as a mere member of the Keokuk Group. Some of the fossils having great distribution, and therefore characteristic, are *Endothyra baileyi*, *Dichocrinus simplex*, *Alloprosallocrinus conicus*, *Batocrinus icosidactylus*, *Pentremites koninckanus*, *Productus biseriatus*, *Spiriferina norwoodana*, *Athyris hirsuta*, *Rhynchonella grosvenori*, *R. mutata*, *Terebratula turgida*, *T. formosa*, *Cypriocardinia indianensis*, *Bellerophon sublaevis*, *Naticopsis carleyana*, *Holopea prontana*, *Cyclonema leavenworthianum*, *Pleuronotaria subglobosa*, and *Spirorbis annulatus*.

CHAPTER XXX.

ST. LOUIS GROUP.

§ 150. THIS Group was named and described by Dr. Shumard in the Geological Survey of Missouri, in 1855. In St. Louis County it is celebrated for its splendid quarries, and consists of hard crystalline limestone, sometimes cherty, with thin layers of argillaceous shales, and has a maximum thickness of 250 feet. It forms bluffs below St. Louis as far as Carondelet, where it dips beneath the Mississippi, but soon rises again, and forms bluffs as far as the Meramec, some of which are 175 feet high. It is exposed in the western part of Illinois and eastern part of Missouri and Iowa, thinning out a short distance north of Keokuk. It forms a band of red clay, chert, and limestone bordering the Indiana coal-fields, and crosses Kentucky and Tennessee, south, by way of Clarksville. It borders the Appalachian coal-field in Southern Kentucky, and may be seen at Burnside, on the Cincinnati Southern Railroad and in Eastern Kentucky. In Indiana it consists of limestones, more or less argillaceous, with beds of red clay, sometimes containing geodes, and having a thickness of 200 to 300 feet. It does not lose its thickness in Kentucky or Tennessee, but becomes more cherty and silicious. It is everywhere cavernous, and abounds in sunken rivers, lost or subterranean streams, and in surface, funnel-shaped sink-holes. The Mammoth Cave of Kentucky, and the Wyandotte Cave of Indiana, which has been explored 23 miles, and has a room 240 feet high, are in this Group.

§ 151. These underground avenues have resulted from percolating water, without the intervention of earthquakes or other extraordinary agency. Surface water from ordinary rain-storms, finding its way through the ground as it does, to supply common springs, will take up carbonate of lime in chemical solution in limestone countries, and by so doing the fissures through which it passes will be enlarged. In massive limestones with thin, shaly partings, the constant action forges of percolating water, aided by disengaged carbonic-acid gas, will enlarge the fissures into rivulets, which will culminate in a subterranean river, finding an outlet in some open stream at a lower level. Such is the process by which the sink-holes, caverns, and subterranean streams in this Group of rocks have been formed. Slight projections on the walls record the different stages of the streams as they were slowly cutting their way to greater depths in the limestone. At the bottom of caverns where little or no water is now flowing, rounded pebbles that have played their part in grinding out the channels occur, as well as sand and clay.

§ 152. When water, holding bicarbonate of lime in solution, slowly drops from the ceiling of a cavern, exposed to the air long enough to allow one equivalent of carbonic-acid gas to escape, the lime is crystallized. If the deposit takes place from above downward, in the form of an icicle, it constitutes stalactite; but if it forms on the floor, from below upward, it is stalagmite. These two sometimes meet and form columns. If the solution which forms the stalactites is free from oxide of iron and other impurities, they will be translucent or milk-white. The presence of iron gives them a dirty yellow, red, or brown color. The chambers in which gypsum occurs are dry, and when rosettes of alabaster or translucent lime are formed the caverns must be dry, as they will not form in a damp atmosphere.

§ 153. The fossils having the greatest distribution, and which are most characteristic of this Group are *Lithostrotion canadense*, *L. proliferum*, *Productus ovatus*, *P. marginicinctus*, *Melonites multiporus*, *Myalina st ludovici*, *Temnocheilus coxatum*, and *Solenocleilus collectum*. Ores of lead and zinc occur in pockets and fissures in Livingston, Crittenden, and Caldwell Counties, Kentucky, and at Rosiclare, Illinois. The ores are associated with fluor spar and calc spar. The principal gangue with which the lead is associated in Hardin County, Illinois, is fluor spar, and it is thoroughly disseminated through it. The fluor spar is used for the manufacture of hydro-fluoric acid, and as a flux for smelting ores, where sulphuret of zinc is associated with galena. Lead occurs associated with different minerals and in many Groups of rocks, but never appears to have had an igneous origin.

§ 154.

Kentucky, In 1856 Hn In 1866 Pr the name in 1855; but Chester used has priority ceous, and bedded sands by a mass of Chester is 1 Tennessee 72 forms a belt s and south-we der of the Mi limestones and

§ 155. T

Aeroerinus sha godoni, *P. su Talarocrinus c Spirifera incre spectabile.*

§ 156. T

Measures, and formably upon basins. It is the occasional thickness is ab Ohio 200 feet, feet, and in N pebbles are we time on the b cemented into Coal Measures everywhere of does not under are free from i

CHAPTER XXXI.

KASKASKIA GROUP.

§ 154. DR. GEO. G. AND B. F. SHUMARD were acquainted with this Group, in Kentucky, Indiana, Illinois, Missouri, and Arkansas, in 1852, but did not name it. In 1856 Hall named it, from Kaskaskia, Illinois, and more fully defined it in 1858. In 1866 Prof. Worthen called it the Chester Group, because he had proposed the name in 1853, and had so informed Prof. Hall while acting as his assistant in 1855; but the latter published the information, and instead of using the name Chester used Kaskaskia. Chester is the shortest and best name, but Kaskaskia has priority of publication. At the typical locality it consists of a compact, arenaceous, and coarse-textured limestone, with shaly partings, in the lower part, heavy-bedded sandstone and limestone, with shaly partings, in the central part, followed by a mass of green shale, succeeded by heavy-bedded limestone. The thickness at Chester is 198 feet, at Huntsville, Alabama, 635 feet, on the southern line of Tennessee 720 feet, at the northern line 400 feet, and in Indiana 300 feet. It forms a belt surrounding the Illinois and Indiana Coal-basin, exists upon the western and south-western border of the Appalachian Coal-basin, and upon the eastern border of the Missouri and Arkansas Coal-basin. It consists everywhere of fossiliferous limestones and sandstones, and is followed by rocks unconformable with it.

§ 155. The fossils having the greatest distribution and most characteristic are *Acrocrinus shumardi*, *Agassizocrinus conicus*, *Hydreionocrinus depressus*, *Pentremites godoni*, *P. sulcatus*, *P. cervinus*, *P. obesus*, *P. pyriformis*, *Pterotoocrinus capitalis*, *Taluroocrinus cornigerus*, *Zeacrinus maniformis*, *Athyris sublamellosa*, *A. subquadrata*, *Spirifera increbescens*, *Spiriferina spinosa*, *Euomphalus planidorsatus*, and *Temnocheilus spectabile*.

CHAPTER XXXII.

CARBONIFEROUS SYSTEM.

§ 156. This system is divided into the Carboniferous Conglomerate, Coal Measures, and Permian Group. The Carboniferous Conglomerate rests unconformably upon the Subcarboniferous rocks, and forms a belt around all the coal-basins. It is a massive sandstone or conglomerate, almost nonfossiliferous, except the occasional presence of *Stigmara*, *Calanites*, and *Lepidodendron*. In Indiana the thickness is about 200 feet, in Illinois about 300 feet, in Kentucky 500 feet, in Ohio 200 feet, in Michigan 100 feet, in Pennsylvania 1,500 feet, in Virginia 1,000 feet, and in Nova Scotia, where it is called the Millstone grit, 6,000 feet. The pebbles are well rounded, showing the fragments of rock were rolled for a long time on the beaches by the action of the winds and waves, before they were cemented into rock. A similar conglomerate separates the Subcarboniferous and Coal Measures in Europe, where it is called the Millstone Grit. It bears the marks everywhere of a shore-line deposit that surrounded the basins of internal seas. It does not underlie the whole of the Coal Measures—the central parts of the basins are free from it, as is shown by artesian boring.

CHAPTER XXXIII.

COAL MEASURES.

§ 157. THE name "Coal Measures" originated among the miners of England before Geology became a science. It is familiarly used in the earliest text-books on Geology, as a scientific term, which was understood without a definition. It is applied to part of the Carboniferous System, and not to Cretaceous or Tertiary Coal regions. The Coal Measures consist of beds of sandstone, shale, slate, limestone, clay, and coal, which are variable in their geographical distribution. The area covered in North America is estimated at about 210,000 square miles, nearly all of which is included in five fields, four of which are in the United States and one in Nova Scotia. Canada and British America are destitute of this important deposit, as well as many States in the Union, among which are Maine, New Hampshire, Vermont, Connecticut, New York, New Jersey, Delaware, South Carolina, Florida, Mississippi, Louisiana, Minnesota, and Wisconsin.

§ 158. The Coal Measures of Nova Scotia rest upon Subcarboniferous rocks, and are divided into the Millstone Grit, Middle Coal Formation, and Upper Coal Formation. A section of the Millstone Grit is as follows: 1. Reddish shales and red and gray sandstones, having a thickness of 2,082 feet, containing no coal, and poor in fossils, except a few drifted trunks of trees. 2. Sandstones, red shales, and a few dark-colored shales, with nine small or rudimentary coal-beds, with a total thickness of 3,240 feet. The underclays abound in *Sigillaria*, and some strata are quite fossiliferous, containing plants, crustaceans, and fish. 3. Red and gray sandstones, red and chocolate shales, arenaceous conglomerates, and thin beds of concretionary limestones, having a thickness of 700 feet, making a total thickness of 6,000 feet. The Middle Coal Formation includes the productive coal-beds, and contains no marine limestones or conglomerates. It consists of shales and sandstones, and has a thickness of 4,000 feet. The Upper Coal Formation consists of shales, sandstones, conglomerates, limestone, and coal, and has a thickness of 3,000 feet. On Cape Breton, the last two divisions have a thickness of 10,000 feet, making the maximum thickness of the Measures, 16,000 feet. From Nova Scotia the Measures dip south-west, and reappear in the form of a subtriangular basin in New Brunswick. The area in Nova Scotia and New Brunswick is 18,000 square miles. The coal is all bituminous. There are 72 seams and numerous dark bands containing more or less carbonaceous material. A coal-bed at Pictou is 37½ feet thick, and another 22½ feet. A large part of the coal-basin is beneath the waters of the Atlantic and the Gulf of St. Lawrence.

§ 159. The first coal-field in the United States is the Appalachian, which extends over important parts of Pennsylvania, Virginia, West Virginia, Maryland, Ohio, Kentucky, Tennessee, and Alabama. Its length is 875 miles, and width from 30 to 200 miles. The anthracite region is in the north-eastern part of Pennsylvania, and does not cover 500 square miles. The coal-beds form synclinals, anticlinals, or stand highly tilted on their edges, but are never horizontal. All the other parts of this great area, estimated at 60,000 square miles, produce only bituminous coal, and the beds may be horizontal or possessed of a slight dip, to which all the strata

are subjected
is 120 feet,
feet. The
The best se
River Coal-
§ 160.

the western
of 47,000 sq
the coal-bed
are ten seam
are from 2 s

§ 161. '
of Iowa, Mi
80,000 squar
and eastern
lowest Group
There are 22
Ten of them
excellent.

§ 162. T
miles, with a
one bed from
thinnest near
seams in clos
fern-leaves an
and Massachu
but possessing
of the rocks a
anthracite, an

§ 163. T
much in thick
the Group in
feet; Ohio, 2
Indiana, 1,000
an upper and
than one cong
localities, and
stones, and co
and open seas
have accompan
bearing *Stigma*
of the vegetati
lucous sedime
the coal vegeta
potash. This
peat, the alkali
waters were fre

are subjected. The aggregate thickness of the coal-beds in the Pottsville district is 120 feet, in the Wilkesbarre district 62 feet, and in the Pittsburg district 25½ feet. The thickest vein at Wilkesbarre is 29½ feet, and at Pittsburg 8 feet. The best seam in Ohio is from 6 to 12 feet in thickness, and is called the Hocking River Coal-bed.

§ 160. The second coal-field in importance covers nearly two-thirds of Illinois, the western part of Indiana, and the western part of Kentucky, and has an area of 47,000 square miles. The coal is bituminous, and the aggregate thickness of the coal-beds is about 40 feet. Indiana is celebrated for her block coal. There are ten seams of coal in a vertical thickness of 600 feet in Illinois, and six of them are from 2 and one-half to 6 feet each in thickness.

§ 161. The third coal-field in importance is the larger one, and occupies parts of Iowa, Missouri, Kansas, Nebraska, Arkansas, and Texas, and has an area of 80,000 square miles. The coal is all bituminous. The western part of Missouri and eastern part of Kansas bear coal in abundance. The Coal Measures are the lowest Group of rocks exposed in Kansas, and have a thickness of 2,000 feet. There are 22 seams of coal, varying in thickness from a few inches to seven feet. Ten of them are more than a foot each in thickness. The coal in Arkansas is excellent.

§ 162. The fourth coal-field is in Michigan, and occupies about 6,700 square miles, with a thickness of about 125 feet. The coal is bituminous, and consists of one bed from 3 to 5 feet in thickness throughout the whole shallow basin, being thinnest near the border. Toward the central axis of the basin there are 2 or 3 thin seams in close proximity to the main seam. The shales are well stocked with fern-leaves and other terrestrial vegetation. There is a small area in Rhode Island and Massachusetts of about 1,000 square miles, having a thickness of 6,500 feet, but possessing no valuable coal-seam. The basin has suffered by the metamorphism of the rocks and plication of the strata. The coal-seams have been changed to anthracite, and are often somewhat wedge-shaped or of irregular thickness.

§ 163. The Coal Measures were deposited in basins, and must necessarily vary much in thickness, the Group in Nova Scotia being thicker than elsewhere, and the Group in Michigan thinner. The maximum thickness in Pennsylvania is 8,000 feet; Ohio, 2,500 feet; Tennessee, 2,500 feet; Western Kentucky, 3,500 feet; Indiana, 1,000 feet, and Missouri, 2,000 feet. The Group is frequently separated into an upper and lower series by the intervention of a conglomerate, and sometimes more than one conglomerate exists in the Group. Marine vegetation abounds at some localities, and land or marsh plants are distributed throughout the shales, sandstones, and coal. Coal was formed from plants which grew in swamps, marshes, and open seas, and, where valuable, it is quite free from sediment, such as would have accompanied much disturbance of the water. The beds usually rest on clay, bearing *Stigmara* and stumps of trees, and are followed by rocks bearing the leaves of the vegetation of that era. The clay beneath the coal-beds is usually an argillaceous sediment, almost devoid of alkalies, and represents the ancient soil in which the coal vegetation flourished, and apparently deprived it of the greater part of its potash. This clay is usually excellent fire-clay. From the coal, as from modern peat, the alkalies were almost entirely removed by the action of water. The waters were fresh, brackish, and salt at different times and at different places. The

marshes were subject to overflows, as shown by the remains of fish and beds of sand and shale, while land-shells, air-breathing reptiles, and trees show the presence of land. The bark of the trees was the 'most durable part, and it is not unusual in sandstone to find only a cast of the tree, covered with a thin film of coal, retaining the original markings of the bark. Some blocks of coal are composed of thin layers formed from the bark of trees and nothing else. Beds vary in purity, from coal with less than one per cent of earthy matter to dark-colored shales, with only a trace of coal.

§ 164. When bituminous coal has lost part of its hydrocarbon gas, it is semi-bituminous, as at Blossburg and Broad Top Mountain coal-fields in Pennsylvania; but if the bitumen is all driven off, it is converted into anthracite. At gas-works bituminous coal is put in a retort, and by the application of heat the gas is driven off, leaving a residue of coke; but if the gas is driven off under great pressure, the residuum is anthracite. When coal melts and runs together in the fire, forming a crust which must be broken to give vent to the draft, it is coking coal. Splint-coal or block-coal does not melt and run together, and is therefore dry-burning coal. Cannel-coal burns with a bright flame like that of a candle, from which circumstance it derived its name. Cannel was the pronunciation of candle in Scotland and England, where this coal received its name. Coal containing sulphur is unfit for smelting iron ores in a blast-furnace, and is not suitable for the manufacture of illuminating gas.

§ 165. Bituminous shales frequently contain iron ore disseminated through them, either as a carbonate or sesquioxide, and sometimes forming black-bands. The same layer of shale which constitutes black-band ore at one place will have the ore gathered in balls, arranged in rows, at another place. By chemical affinity the disseminated particles were brought together, and formed into balls or discs; and hence the iron exists in all stages, from fine distribution through the shales to layers of kidney ores, with whitened shales intervening. The iron ores of the Coal Measures are generally hardened mud, charged with iron, or clay-iron stone, and rarely yield more than 40 per cent of iron, and they are not of much value except as they exist around the margin of the Appalachian coal-field in the Lower Coal Measures. No good iron-mines are found in the other coal-basins. The greater part of iron manufactured from these ores has been obtained in Pennsylvania.

§ 166. The first trace of reptiles observed in the Carboniferous System consisted of foot-prints, found in 1841, in the Lower Coal Measures of Horton Bluff, in Nova Scotia. This was followed in 1844 by the discovery of reptilian bones at Saarbruck, and in 1851 to 1853, bones in Nova Scotia, and the land-snail, *Pupa vetusta*. Since that time the discoveries have been numerous. There is no reason to suppose the atmosphere was charged then with any more carbonic acid than it is now; on the contrary, the air-breathing animals prove it was not. The life of plants and animals is controlled by oxygen, and the adaptation of organs is in accordance with its properties. If there was less oxygen in the atmosphere, the membranous reptile lung could not supply the demands of its system, and analogy proves these animals could not have existed in the coal period with a less proportion of oxygen than is required now.

§ 167. The coal-beds and the vegetation of the coal period are usually suffi-

cient to de-
fossils of w
liferum, Sp
les mesolob
isma subcon
Crenipecten
ulata, P. sp
nata, Nautil

§ 168. Ural Mount
this country
Norwood and
lupe Mounta
thickness of
shales, congl
New Mexico
marks, and so
It is conforma
ures, having
mains in Illin
unconformabl
species are *P*
Pleurophorus
§ 169. T
voted. All t
visions of the
Group has no
38,000 feet.
other States th
in Nova Scotia
western ranges

cient to determine the age of the rocks, but some of the invertebrate characteristic fossils of wide geographical distribution are: *Fusulina cylindrica*, *Lophophyllum proliferum*, *Spirifera camerata*, *Productus rogersi*, *P. nebraskensis*, *P. longispinus*, *Chonetes mesolobus*, *Athyris subtilita*, *Spiriferina kentuckiensis*, *Macrodon carbonarius*, *Allorisma subcuneatum*, *Aviculopecten rectilaterarius*, *Pernopecten aviculatus*, *Pinna peracuta*, *Crenipecten retiferus*, *Myalina subquadrata*, *Bellerophon carbonarius*, *Pleurotomaria tabulata*, *P. sphaerulata*, *Macrochilina gracilis*, *M. primigenia*, *M. kansasensis*, *M. variata*, *Nautilus missouriensis*, *Phillipsia missouriensis*, and *P. sangamonensis*.

CHAPTER XXXIV.

PERMIAN GROUP.

§ 168. This Group was described by Murchison in 1845, in Russia and the Ural Mountains, and named from Perm, in Russia. It was first ascertained in this country by Swallow in 1858, in Kansas, where it has a thickness of 320 feet. Norwood announced its existence in Illinois, and Shumard described it in the Guadalupe Mountains of New Mexico, where it consists of white limestone, having a thickness of 1,000 feet. In Kansas it consists of magnesian limestone, marls, shales, conglomerates, and gypsum; the magnesian character increases southerly to New Mexico. Fossils are abundant on the Cottonwood, with sun-cracks and ripple marks, and sometimes small piles of fossils and fragments appear, as if washed together. It is conformable with the Coal Measures. In Pennsylvania the Upper Barren Measures, having a thickness of 1,000 feet, are referred to it. It is claimed the reptilian remains in Illinois and Texas have shown its existence in those States. It is always unconformable with the rocks above, in this country and elsewhere. Characteristic species are *Pseudomonotis hawni*, *Myalina permiana*, *Bakevella parva*, *Monotis halli*, and *Pleurophorus subcuneatus*.

§ 169. This Group closes the Palæozoic series, to which this work is chiefly devoted. All the Groups exist in New York and Pennsylvania, except the subdivisions of the Subcarboniferous can not be distinguished, and the doubtful Quebec Group has no existence there. The maximum thickness in these States is about 38,000 feet. Some of the Groups in the Lower Silurian have greater thickness in other States than they have in these two, and the Coal Measures are much thicker in Nova Scotia than they are in Pennsylvania. The whole Palæozoic series in the western ranges of mountains has an estimated thickness of about 40,000 feet.

CHAPTER XXXV.

TRIASSIC SYSTEM.

§ 170. The Mesozoic era is divided into three grand ages—Triassic, Jurassic, and Cretaceous. The name Triassic was applied to the rocks in Germany, in allusion to a threefold division which they present in that country; but no such division exists in America. Indeed, notwithstanding the vast thickness of the rocks, they have thus far baffled all attempts to divide them into Groups, and, on account of the similarity of the rocks with the Jurassic, and the barrenness of fossils in the eastern exposures, these Systems have not been satisfactorily defined and separated. On the eastern part of the continent they fill synclinal troughs, and have been very much disturbed by intrusive rocks and volcanic action. They generally rest on Laurentian or Taconic strata, and, of course, the bed is always unconformable. But on the western part of the continent they are frequently undisturbed, and spread over great areas of country, resting on unconformable rocks. The Triassic in the Connecticut Valley extends from Northfield, in the northern part of Massachusetts, across the latter State and Connecticut to New Haven, on Long Island Sound, a distance of 105 miles. It fills a synclinal trough, and has its greatest width at the mouth of the Farmington River, which is about 20 miles. The rocks consist of red sandstones, conglomerates, shales, and occasionally impure limestone. The maximum thickness is about 20,000 feet, but the upper 8,500 feet is referred to the Jurassic, leaving 11,500 feet for the Triassic. A great many reptilian tracks, some fish and a few land-plants and fucoids, have been described from these rocks. Much excellent building-stone has been quarried from the sandstone. About 15 miles west of the exposure, on Long Island Sound, there is another exhibit, about 6 or 7 miles long and 2 miles wide.

§ 171. A long trough and great exposure begins at Stony Point, on the Hudson, and extends across New Jersey, Pennsylvania, and Maryland to Culpeper County, Virginia. It has a length of about 350 miles, and, though frequently narrowing to a breadth of 4 or 5 miles, expands in New Jersey to a width of about 36 miles. The general character of the rocks is like those in the Connecticut Valley, and the total thickness on the Delaware River is 27,000 feet, part of which is probably Jurassic. Another range crosses the Potomac near Washington City, and extends 25 or 30 miles beyond Richmond, and another exists 25 miles west of this one. There is a valuable coal-field in this System in Virginia, which is about 26 miles long and 4 to 12 wide. The James River flows through the middle of it, about 15 miles from the northern extremity, while the Appomattox traverses it near its southern border, and on its eastern side it is distant from Richmond about 13 miles. A great many fossil plants have been described from this locality. There are two basins in North Carolina. One begins at Lakeville, and extends about 30 miles south-west to Germantown, being from 4 to 6 miles wide; and the other commences in Granville County, six miles south of Oxford, and extends south-west about 120 miles, reaching 6 miles into South Carolina. Its width is generally about 6 miles, but at the widest part 18 miles. The thickness in some places exceeds 25,000 feet; the area is about 1,000 square miles, nearly one-third

of which consists of
lanceous iron
these rocks,
discovered in
sides of Col
south side of
miles along
rocks of this

§ 172.

and all vari
system from
square miles.
from nearly
extensive are
and bear into
without any
rocks earlier
few genera are

§ 173. I

the Shinarump
of the Shinar
coloring is str
terraced butte
red, approxim
great cliff-form
homogeneous
vening; the sh
off vertically.
and sloping to
been called the
imum thickness
columns in the
weathered out

of which contains coal-beds. Very valuable beds of coal and beds of good argillaceous iron ore are distributed through it. Many fossils have been described from these rocks, and among them *Dromatherium sylvestre*, the earliest fossil mammal yet discovered in America. The rocks occur in Nova Scotia, on the north and south sides of Cobequid Bay, from Moose River to the mouth of North River, and on the south side of the Bay of Fundy. Prince Edward's Island, which stretches for 125 miles along the northern coast of Nova Scotia and New Brunswick, consists of rocks of this age.

§ 172. The red beds of the Triassic, consisting of every texture of sandstone and all varieties of red, are distributed almost throughout the Rocky Mountain system from Mexico to the Arctic regions, covering hundreds of thousands of square miles. Fossils have been collected and described from every territory and from nearly every mountain range throughout this vast extent of country. Over extensive areas of country the Triassic rocks are more than a mile in thickness, and bear internal evidence of having been deposited in the depths of the ocean without any mechanical sediment. Not a single species of any organism found in rocks earlier or later than the Triassic have ever been found within it, and very few genera are common to it and rocks of earlier or more recent date.

§ 173. In Colorado and Utah the lower part of the Triassic has been called the Shinarump Group, and the upper part the Vermilion Cliff Group. The rocks of the Shinarump are persistent in their characters for hundreds of miles, and the coloring is strong and deep. They weather into striking architectural forms and terraced buttes. The rocks of the Vermilion Cliff Group are colored a brilliant red, approximating vermilion, or sometimes inclining to orange, and constitute the great cliff-forming series of the West. The Group consists of massive layers of homogeneous sandstone, from 100 to 300 feet in thickness, with shaly layers intervening; the shales disintegrate, and thereby the sandrock is undermined and breaks off vertically. This process, in time, has presented a series of perpendicular walls and sloping taluses. In the West Humboldt Range of Mountains the lower part has been called the Kaipato Group, and the upper part the Star Peak Group. The maximum thickness in this region has been estimated at 16,000 feet. The fantastic columns in the "Garden of the Gods" and in Pleasant Park, Colorado, have been weathered out of the sandstones of this System.

CHAPTER XXXVI.

JURASSIC SYSTEM.

§ 174. The Jurassic System was named from the Jura Mountains, of Switzerland. No *Trigonia*, *Belemnites*, *Ammonites*, or specially characteristic fossils of the Jurassic, have been found on the Atlantic side of the continent, notwithstanding the upper part of the rocks described in the last chapter may be Jurassic. The Jurassic fossils, however, occur in the Rocky Mountain Ranges from Mexico to the Arctic regions. The rocks exist in every State and Territory throughout that vast extent of country, varying in thickness from a few hundred feet to 10,000 feet. They follow the Triassic, and generally rest upon it. Fossils have been described from California, Arizona, New Mexico, Idaho, Colorado, Nevada, Montana, Dakota, British Columbia, Cook's Inlet, Alaska, Point Wilkie on Prince Patrick's Land, and the islands north of Grinnell Land. In some parts of its grand geographical distribution it is composed of sandstones and clays, resembling, in appearance, the Triassic; but in others it consists of limestones, sandstones, shales, and clays, indicating shallow water, and bearing no resemblance to the Triassic. The limestones are frequently fossiliferous, and show the progress animal life had made in the ocean, and vegetation had made on the land. Of 50 genera of vertebrates described from the Jurassic, none of them are Paleozoic, and only two have been doubtfully identified in the Cretaceous. *Ammonites*, *Ceratites*, and *Belemnites* made their first appearance in the Jurassic, and became extinct in the Cretaceous. The genus *Spirifera*, so abundant in the Devonian and Carboniferous, became extinct in the Jurassic. Several genera of mammalian remains have been defined from the Jurassic, but they are all peculiar to it. No single species of plant or animal is common to the Jurassic and any other formation. Ten genera of Carboniferous plants have been identified in the Jurassic, and four genera occurring in the Jurassic have been identified in the Cretaceous. There is a general progress among the invertebrates toward succeeding ages, but the evolution of the vertebrates is very much more marked. There is almost universal unconformability with the overlying Cretaceous, and hence there is an era of time not represented by the rocks. It has been called the Reptilian age, because of the gigantic saurians which then infested the seas. Some of the rocks belonging to this System in California, and, especially about Mariposa, are said to be gold-bearing, but minerals are generally very scarce.

§ 175.

to the rocks of the Cretaceous system. The Tertiary York to Florida with the exception of Mississippi, West of the country is covered by regions, exposed and swept away, and includes, of course, it is found in the British American rock, or is over American cor fairly presumed its present distribution but in the west.

§ 176. M

and the great ton Group, the Dakota Group with alternating silicified wood trees, with considerable thickness in the San Juan Group of Alaska contains beds of

§ 177. Th

Missouri, where with seams of limestone and other fossils

§ 178. Th

consists of marl fish, and has a thin, but rarely

§ 179. Th

it consists of clay sulphuret of iron

CHAPTER XXXVII.

CRETACEOUS SYSTEM.

§ 175. THE name Cretaceous is from the Latin *Creta*, chalk, and was applied to the rocks in Europe long before its use as a geological term. The existence of the Cretaceous on this continent was first ascertained, in 1827, by Morton and Vanuxem. The Cretaceous is found either exposed upon the surface, or covered by the Tertiary, forming a border of variable width on the Atlantic Coast, from New York to Florida. In like manner it occurs everywhere south of the 33d parallel, with the exception of limited areas in the mountain regions. It covers nearly all Mississippi, extends into Tennessee and Arkansas, and reaches Southern Illinois. West of the 97th Meridian, from the 33d parallel to the Arctic Ocean, the whole country is covered with this formation, with the exception of the areas in the mountain regions, exposing older rocks and inconsiderable extensions of land, where it has been swept away, and an area of some magnitude north and west of Hudson's Bay. This includes, of course, the whole extent covered by the Tertiary and more recent deposits. It is found east of the 97th Meridian, extending into Iowa, Minnesota, and some parts of British America. Or, approximately stated, the Cretaceous forms the surface-rock, or is overlaid with the Tertiary and recent strata over nearly half the North American continent, and from the extensive denudation it has suffered, we may fairly presume, at the commencement of the deposit, the land surface was not half its present dimension. In the east and south the formation is exclusively marine, but in the west the marine is succeeded by a brackish-water deposit.

§ 176. Meek and Hayden divided the marine Cretaceous of Kansas, Nebraska, and the great West, in 1861, in ascending order, into the Dakota Group, Fort Benton Group, Niobrara Group, Fort Pierre Group, and Fox Hills Group. The Dakota Group was named from Dakota County, where it consists of sandstones, with alternations of various colored clays, and beds, and seams of impure lignite, silicified wood, and great numbers of leaves of the higher types of dicotyledonous trees, with casts of *Pharella dakotensis*, *Acineta siouzensis*, and *Cyrena arenaria*. The thickness in that locality is 400 feet, in North-western Colorado 600 feet, and in the San Juan region 1,000 feet. It is the supposed equivalent of the Eutaw Group of Alabama and Mississippi, which has a thickness of about 400 feet and contains beds of lignite.

§ 177. The Fort Benton Group was named from Fort Benton on the Upper Missouri, where it consists of dark-gray, laminated clays, sometimes alternating with seams of limestone. It abounds in *Inoceramus*, *Ammonites*, *Scaphites*, *Nautilus*, and other fossils, and has a thickness of 800 feet.

§ 178. The Niobrara Group was named from Niobrara, in Nebraska, where it consists of marls and limestones, and abounds in *Inoceramus*, *Ostrea*, and remains of fish, and has a thickness of 200 feet. It has an extensive geographical distribution, but rarely exceeds 500 feet in thickness.

§ 179. The Fort Pierre Group was named from Fort Pierre, in Dakota, where it consists of clays containing carbonaceous matter, seams of gypsum, and masses of sulphuret of iron, and abounds in the shells of Cephalopods, Lamellibranchs,

remains of fish and saurians, and has a thickness of 700 feet. In Northern Colorado it is 800 feet thick, and in Alabama and Mississippi it is known as the Rotten limestone, and reaches a thickness of 1,200 feet.

§ 180. The Fox Hills Group was named from Fox Hills, in Dakota, where it consists of gray, ferruginous, and yellowish sandstones, and arenaceous clays, abounding with shells of Cephalopods, Lamellibranchs, Gasteropods, remains of fish and saurians, and has a thickness of 500 feet. East of the Colorado Range its thickness is 1,500 feet, in the valley of Bitter Creek 3,000 feet, and in that of the North Platte 4,000 feet. It is the same as the Ripley Group of North Carolina, Alabama, and Mississippi, which has a thickness of about 400 feet.

§ 181. The thickness of the marine Cretaceous in New Jersey is about 700 feet. It is valued in that State for its fertile marl, and beds of kaolin in its lower part. In Louisiana its thickness is more than 1,000 feet, in the Uintah Mountain region 7,000 feet, and in New Mexico and British America more than a mile at many places. The cañon of San Carlos, on the Rio Grande, exposes a clear perpendicular height above the river level of 1,500 feet of Cretaceous strata. The Cretaceous is the Coal-bearing formation at Vancouver's Island and other points on the Pacific Coast.

§ 182. There is in the West, superimposed upon the marine Cretaceous strata, rocks which were deposited in brackish water, and form transition-beds from the strictly marine condition of the Cretaceous to the epoch of numerous fresh-water lakes, which were scattered all over the country west of the Mississippi, and north in British America to the Arctic regions. These rocks were named in 1861, by Meek and Hayden, the Fort Union Group. They consist of beds of clay and sand, with numerous seams and local deposits of lignite and beds of coal. The passage from the marine to the brackish-water deposits, and from the latter to the fresh-water deposits, is without abrupt change in the sediment, and with complete conformability. There is no evidence of any important physical or climatic change, beyond the gradual filling up of the basins of the sea and the recession of the salt and brackish water, appearance of fresh-water lakes, and their gradual disappearance. The Fort Union Group has been called the Judith River Group, the Bitter Creek Group, the Bear River Group, the Laramie Group, and by divers other names. It has a thickness, in Bitter Creek Valley, Wyoming, of 6,000 feet, and in Bear River Valley, in Utah, of 7,000 feet. Its geographical distribution extends for a thousand miles in length, and a maximum width of 500 miles or more, with a varying thickness from 100 feet or less, to 7,000 feet or more. It abounds in plants belonging to Eocene genera, which connect the Cretaceous and Tertiary flora by insensible degrees, while the Dinosaurian remains demonstrate its Cretaceous age.

§ 183. Before the discovery of this Group, absolute nonconformability was supposed to exist between Cretaceous and Tertiary rocks, and this is the case where marine Tertiary follows the marine Cretaceous, wherever known in the world. But here, where the marine Cretaceous is as recent as elsewhere, and the continuance of the period is represented by brackish-water deposits, and then fresh-water deposits in lakes cut off from the ocean, the rocks are conformable, and the vegetable and animal kingdoms show the slow progress of advancing ages. About one-third of the genera of plants belonging to that period have become extinct, but the living plants, *Corylus americana*, *C. rostrata*, *Davallia tenuifolia*, and *Onoclea sensibilis*, have

been identified with the Cretaceous era with the important genera of mammals hitherto considered as chasm di-
era. No such fossils are not furnished, encouraged to imperceptible Group.

§ 184. The living, the Primary and so interwoven notwithstanding which it relates to conchological the living species of geological time rocks generally impressions in the continuous strata order of super- species. This per cent of living things; those com- plic less recent Pliocene, which identical with the Pliocene. Instead of a contrary course is Certain species be, and from the This subdivision species and the nomenclature.

been identified from the Fort Union Group, thus specifically uniting the Cretaceous era with the present time. It is possible, too much confidence in this identification may lead to error, and better specimens may show specific distinctions; but it is an important fact, they so closely resemble the living forms as to be mistaken for them, and show how closely the living are connected with the ancient dead. Among the Cretaceous genera of invertebrates, about one-third survive; three genera of reptiles, *Crocodylus*, *Trionyx*, and *Emys* survive; but no genus of birds or mammals has come down from that age to the present. There is no great break or chasm discoverable in vegetable or animal life in passing back to the Cretaceous era. No sudden physical change has taken place over which some deposit may not furnish a connecting bridge. No evidence of any great climatic change is furnished, either in the animal or vegetable world, but on every hand we are encouraged to look at uniformity in the organisms, subject only to a constant, almost imperceptible evolution. Seams of productive coal occur at different places in this Group.

CHAPTER XXXVIII.

TERTIARY SYSTEM.

§ 184. THE organic remains of the Tertiary are so completely blended with the living, that no Quaternary age or period can be distinguished. The words Primary and Secondary have become quite obsolete in Geology, while Tertiary is so interwoven with the science as to be permanently fastened to the nomenclature, notwithstanding its definition, as the third age, has no application to the period to which it relates. The subdivision of the Tertiary, with reference to the survival of conchological species into Eocene, Miocene, Pliocene, and Post-pliocene, brings us to the living species as gradually as the species change within any of the subdivisions of geological time, or within any division of the strata into Groups. The Tertiary rocks generally consist of marls, clays, sands, or other friable material, filling depressions in the underlying rocks, and, though widely distributed, seldom form hard, continuous strata. This condition of the rocks made it difficult to determine the order of superposition, until a comparison of the shells had been made with living species. This comparison led to the naming of the rocks containing about 3 or 4 per cent of living species, the Eocene, which signifies the dawn of the present state of things; those containing 15 to 20 per cent of living species, the Miocene, which implies less recent; and those containing 90 to 95 per cent of living species, the Pliocene, which means more recent; and those having all the imbedded fossil shells identical with living species, though containing extinct mammalian remains, Post-pliocene. Instead of determining the rocks by the per cent of living species, the contrary course is now adopted, and the age is determined by the extinct species. Certain species are regarded as types of Eocene age, or Miocene, as the case may be, and from the presence of these the rocks are referred to the proper Group. This subdivision of the Tertiary, with reference to the survival of conchological species and the division into geographical Groups, have made a double system of nomenclature.

§ 185. The marine Eocene, commencing in New Jersey with a thickness of 37 feet, and exposing only a narrow surface area, crosses Maryland by way of Fort Washington; Virginia, by way of Fredericksburg, Richmond, and Petersburg; North Carolina, by way of Newbern and Wilmington; South Carolina, by way of Charleston and Shell Bluff, on the Savannah River; Georgia, by way of Milledgeville; Alabama, by way of Claiborne; and Mississippi, by way of Jackson and Vicksburg. In South Carolina it consists of loose sand, clay, gravel, sandstone, limestone, and marl, covers a large area, and has a thickness of 1,100 feet. It is divided into the Buhrstone Group, Santee beds, and Ashley and Cooper beds. It is exposed in Florida, and reaches up into Tennessee, where it is called the Porter's Creek Group. Conrad subdivided it in Alabama and Mississippi, where it has a thickness of about 900 feet, into the Claiborne Group, Jackson Group, St. Stephen's Group, and Vicksburg Group. It crosses Louisiana, appears in Arkansas, and offers numerous exposures in Texas, Mexico, and California. It is extremely fossiliferous at many places, and nowhere conformable with the underlying rocks.

§ 186. The gradual elevation of the western ranges of mountains through Cretaceous and Tertiary time, the formation of bays and arms of the sea, and lakes which have drained themselves in continuing succession, have linked the Tertiary with the Cretaceous, and bound the Eocene, Miocene, Pliocene, and Post-pliocene with the present, almost as one connected age. In these lake regions the Eocene is divided into the Wahsatch Group, Green River Group, Bridger Group, and Brown's Park Group, and there are numerous synonyms for each one of them. The Wahsatch is characterized by its brick-red color, and has a thickness of 8,000 feet; the Green River Group is quite fossiliferous, and has a thickness of 7,500 feet; the Bridger Group rests conformably on the Green River, consists of Bad Land sandstones, limestones, shells, and marls, and has a thickness of 2,000 feet; and the Brown's Park Group has a thickness of 2,500 feet. The combined thickness of the Eocene in the Western Territories is therefore 20,000 feet.

§ 187. The marine Miocene beginning at Martha's Vineyard, though it may exist as far north as Maine, crosses New Jersey through Cumberland County, and forms a border upon the east and south of the Eocene exposure a large part of the way to the Mississippi River, and west across Louisiana, Texas, and Mexico. It is not conformable with the Eocene, and in some parts does not intervene between it and later deposits. It has its greatest thickness in California, where it exceeds 3,000 feet. The Coast Range of mountains is composed in large part of strata of this age, and hence its elevation has been since the Miocene period. It is highly fossiliferous, and the shells generally belong to living genera, and many of the species still survive in the waters bordering the adjacent coast, thus indicating no material change in the climate since that period. The Miocene lake deposits, like the Eocene, cover great extensions of Territory and reach an enormous thickness. In Nebraska it has been divided into the Wind River Group, which has a thickness of 2,000 feet, and the White River Group, which has a thickness of 1,000 feet. On the divide between the Arkansas and South Platte, where the thickness is from 1,500 to 2,000 feet, it is called the Monument Creek Group, and in Oregon it is called the Truckee Group.

§ 188. The marine Pliocene strata are found in Maryland, superimposed upon the Miocene, in South Carolina, upon the Eocene, and generally forming a narrow border at the east of these outcrops on the Atlantic coast, and a wider border on

the south adjacent coast. The climate and in California evidence, so this continent now, no more. Post-pliocene an arbitrary deposits in 400 feet and thickness of called the Ni Group and thickness. § 189. T by drift, sand it is confined those of special thickness is 5 thickness may been formed, a thickness. TH have a thickness able with the In South Carolina other mammal. At some time of his work is f are associated v survived the d § 190. Du Bay was submerged clays occurring too, are striated fragments of rock of Canada south was submerged, are shown at M above the bay; part of Canada. species in the G surface of the rock the St. Lawrence New England St they appear on n or fresh-water fac of the clays with

the south adjoining the Gulf Coast. Fossil shells of species now living on the adjacent coast, abound at every point, and demonstrate beyond reasonable doubt the climate and the waters on the eastern and southern coast of the United States, and in California, were then the same that now prevails. There is no palæontological evidence, so far known, that the Pliocene climate was different from the present on this continent, and as the outlines of the continent were then nearly as they are now, no material difference can be inferred. The Pliocene graduates into the Post-pliocene, so that separation of the strata frequently becomes impracticable, and an arbitrary approximating line for separation is assumed. The Pliocene lake deposits in Nebraska, are called the Loup Fork Group, and have a thickness of 400 feet and cover a great extent of territory, and in North-western Kansas have a thickness of 500 feet. In Wyoming they have a thickness of 1,500 feet, and are called the Niobrara Pliocene. In Bear River Valley they are called the Salt Lake Group and the Cache Valley Group, and the thickness is from 500 to 1,500 feet.

§ 189. The Post-pliocene is represented by marine deposits on the coast, and by drift, sand, and gravel, in the middle part of the continent. In South Carolina it is confined to a belt along the coast 8 or 9 miles wide, and the fossil shells are those of species inhabiting the coast. In Los Angeles Valley, in California, the thickness is 500 feet; but where depressions upon the coast have been filled the thickness may be 1,000 or 1,500 feet, and so at the mouths of rivers where a delta has been formed, as at the mouth of the Mississippi, the Post-pliocene becomes of very great thickness. There are some Lake deposits of this age in the great West, which have a thickness of 500 feet or more. The marine Post-pliocene is usually conformable with the Pliocene, and graduates into the present deposits without disturbance. In South Carolina the bones of horses, hogs, dogs, rabbits, beavers, tapirs, and other mammals occur in the layers of blue mud and sand throughout the period. At some time during this age, man made his appearance on this continent, for none of his work is found preceding it, nor preceding the drift; but his stone implements are associated with the remains of the mastodon and mammoth, and such animals as survived the drift period in such condition as to show they lived at the same time.

§ 190. During the Post-pliocene era, a portion of the country about Hudson's Bay was submerged by the ocean, as shown by the fossiliferous marine sands and clays occurring at 300 or 400 feet above the present level of the ocean. The rocks, too, are striated in all directions, as if done by icebergs or shore-ice holding angular fragments of rock. The New England States and New Brunswick, and that portion of Canada south of the St. Lawrence River and east of the vicinity of Montreal, was submerged, with the exception of the mountain elevations. Several beaches are shown at Murray Bay 90 miles below Quebec, varying from 30 to 326 feet above the bay; like beaches occur at Montreal and at various other places in this part of Canada. All these deposits abound in marine fossils belonging to living species in the Gulf of St. Lawrence and on the near coast of the Atlantic. The surface of the rocks below these deposits is polished and striated in the direction of the St. Lawrence Valley. Like phenomena occur over New Brunswick and the New England States, and extending as far south as the mouth of the Hudson; but they appear on no other part of the continent. These deposits contain no terrestrial or fresh-water fauna, and, so far as the marine life is concerned, connect the lowest of the clays with the present time by an unbroken chain of animal existence.

§ 191. South of the Laurentian Mountains the surface of the rocks beneath the boulder clay is striated in the direction of the valleys, but there is no connection between these and those occurring north of the mountains in the Hudson's Bay region. The force which produced the scratches did not cross the mountains nor exist upon them. Prof. Dawson has proven the bodies which produced them came from the Atlantic Ocean, and following up the St. Lawrence drifted to the south, at various angles, some floating over New Brunswick, and others over Maine, and others through Lake Champlain, and re-entering the Atlantic Ocean by the Hudson River, while others were driven beyond Montreal into the mouth of the Ottawa River. In New Brunswick the striæ are related to the contour of the surface of the land, and conform to the direction of the river valleys. A south-easterly course prevails in the western part of Charlotte County, and a south-western course in the valleys east and north-east of St. John. A map of Maine showing the course of the rivers will show the course of the striæ. The appearance of the surface geology of this State early suggested the fact that a great rush of waters poured over it from a northerly source, and transported by its power the surface *débris* which had accumulated in earlier ages by subaerial forces, and large masses of rock from parent ledges, and deposited them in regions more or less distant from the several sources; and as they passed along they striated and grooved the rocks against which they impinged, or over which they rubbed in the traveled course. The striæ conform to the valleys as a rule, and therefore have their courses in all directions, though some are found deflected at right angles to their original course. The Katahdin Mountains formed an obstruction around which the striating agency operated, but it did not cross the summit. The striæ occur on the north side of the mountains, but not upon the south side. In Vermont, New Hampshire, Massachusetts, and Connecticut, beneath the drift, sand, gravel, boulders, and clay, the surface of the rocks is grooved and furrowed in a general southern direction, though varying with the contour and course of the valleys. At the Island of New York the current swept from the north-west to the south-east, and the furrows are most strongly marked on the north-western slopes of the hills, and least on the south-eastern. In many instances they are very distinct on the western and north-western slopes, extending to the highest point of the rocks; but no traces exist on the eastern and south-eastern slopes, although both slopes are equally exposed. The striæ are most numerous in the middle part of the island, somewhat less in the western, and least in the eastern, showing the current was deflected southward in the middle part of the island. Throughout all this area south of the Gulf of St. Lawrence and the St. Lawrence Valley, we have, in the striæ and furrows and in the distribution of clay, boulders, gravel, sand, and fossils, the evidence of an overflow of the whole country, except the higher hills and mountains, the overflow resulting from subsidence of the coast, and the evidence that the Arctic current, instead of leaving the coast on approaching the mouth of the gulf, as it does now, flowed into the gulf and across the depressed New England area, transporting its fields of ice, which grounded upon the northern slopes of hills and mountains, and rubbed the rocks in the valleys and plains wherever the surface soil and subaerial accumulations were swept off by the grinding weight of a mass, driven by a current through water too shallow to float it. In the Gaspé Peninsula, ocean-terraces and stratified clay, containing marine testacea, occur at the height of 600 feet above the sea. In the Champlain region of Ver-

mont, and
Ottawa V.
high as 50
on the coast
and sand a
200 fathom
feet, becau
the clay, to
a covering
which then
which have
§ 192.
spreads out
of the States
boulders, an
which now fl
restrial anim
which show
were stationa
of sand and
than it does
have overflow
These terraces
Michigan, Hu
Superior durin
Ontario. Her
large as all of
were rolled in
and terraces, w
spread over W
Illinois, and
Mississippi River on
far as the Ohio
soon the gravel
and reach to th
nuded of their
This is especiall
and furrows app
inks when occu
boulders, that w
directions, and
been made by o
§ 193. Com
hills are from 60
south over all the
Georgia, Alabam
Mexico, there is

mont, and the triangular area of 9,000 square miles extending from there to the Ottawa Valley, the marine fossiliferous clays and sand occur at all elevations, as high as 500 feet. They form a coating for New Brunswick, and a continuous belt on the coast of Maine 150 feet above the ocean. The marine species in these clays and sand are such as live at moderate depths, or varying from the littoral zone to 200 fathoms. The submergence must therefore have been much more than 600 feet, because the shells and bones must have had some depth of water, as well as the clay, to protect them, in order to produce the fossilization, and they received a covering of drift materials sufficient to protect them from the ocean currents, which then swept over that region, and the disintegrating and denuding agencies which have prevailed during the long train of centuries that have since elapsed.

§ 192. The fresh-water drift surrounds the great central lakes of the continent, spreads out over a large country in British America, and overspreads part of each of the States in the Valley of the Mississippi. This drift consists of clays, gravel, bowlders, and sand, containing no marine organisms, but bearing land vegetation which now flourishes in the same latitude, and fresh-water shells and the bones of terrestrial animals of the Post-pliocene age. There are beaches surrounding the lakes which show the lakes have occupied much higher levels than they now do, and were stationary for a time at each of these beaches. The terraces and lake deposits of sand and clay in Wisconsin show that Lake Superior stood 600 feet higher than it does now, at one time, in the Post-pliocene age, at which time it could have overflowed nearly the whole country south of it to the Gulf of Mexico. These terraces and lake deposits occur at different elevations surrounding Lakes Michigan, Huron, Erie, and Ontario, showing they were elevated as high as Lake Superior during this period. They have been noticed 750 feet higher than Lake Ontario. Here was then one grand central Post-pliocene lake, several times as large as all of them combined are now. Upon the shores of this lake angular rocks were rolled into bowlders and beaten down to gravel and sand, that formed beaches and terraces, which were subsequently swept south by the overflowing lake, and spread over Western Ohio, Western Kentucky, nearly all of Michigan, Indiana, Illinois, and Mississippi, and the eastern part of the States bordering the Mississippi River on the west. Large bowlders are spread over these States south as far as the Ohio River, though they gradually diminish in size in that direction, and soon the gravel disappears, and only the finer materials are spread over Mississippi and reach to the Gulf. Beneath these clays and sands, where the rocks were denuded of their subaerial *débris*, the surface is frequently scratched and furrowed. This is especially the case where the higher lands were overflowed. The scratches and furrows appear to have been made by shore-ice on the margin of the lake or lakes when occupying different elevations, and by ice carrying angular rocks and bowlders, that were driven against the shores or shallow places. They bear in all directions, and frequently cross each other, which proves they could not have been made by one body, or by any number of bodies moving in the same direction.

§ 193. Commencing in the lower tier of counties in New York, where the hills are from 600 to 800 feet above the level of the narrow valleys, and extending south over all the highlands of Pennsylvania, Virginia, West Virginia, the Carolinas, Georgia, Alabama, Eastern Kentucky, and Tennessee, and south to the Gulf of Mexico, there is an absolutely driftless area, and the surface rocks are free from

scratches and furrows. It was dry land, and much of it high and mountainous, when the marine clays and sands were strewn over the territory adjacent to the Gulf of St. Lawrence and the New England States, and dry land during the period of the drift of the central part of the continent, and for geological ages antecedent thereto. The precipitous ledges and profound valleys of denudation, the overhanging rocks and castellated outliers, furnish incontestable evidence of the ordinary eroding agencies through a period of time commencing anterior to the Tertiary epoch. There are extensive driftless areas in Eastern and Southern Ohio free from scratches and furrows on the surface rocks, and from drift, sand, gravel, and boulders, and they are characterized by outliers, monument rocks, sharp ridges, and rugged scenery. The drift materials extend from the lakes to the sources of the rivers that flow into the Ohio, and over more or less of the land intervening between the head-waters; but below this they occur only in the valleys of the larger rivers. Wherever the valley was large enough to carry off the flow of water from the north, the adjacent land was not overflowed, and the height of the water in the valley is marked by river terraces. In Eastern Ohio, only those rivers having their sources in the central and northern part of the State have river terraces, as the Scioto, Hocking, and Muskingum, while the smaller tributaries, such as Raccoon, Shade, and little Muskingum, have not a vestige of drift, or scratch, or furrow, from their sources to the Ohio. The Ohio River Valley was large enough to carry off the water that flowed across Ohio and Indiana, and hence no drift crossed the valley until it reached the western part of Kentucky. Throughout the drift area of Ohio, Indiana, and Illinois, it is common in excavations below the drift to find an ancient soil of vegetable mold resting upon stratified rocks in place. Beech, sycamore, hickory, and cedar have been found where they grew prior to the drift; but beneath the ancient soil no striated or furrowed rock has ever been discovered.

§ 194. There is a driftless area in the south-western part of Wisconsin, covering about 13,000 square miles, or nearly one-fourth of the State, and which extends into Northern Illinois, North-eastern Iowa, and Eastern Minnesota. There is no drift, sand, clay, or gravel, and, as in all cases where these do not occur, there are no scratches or furrows on the surface of the rocks. This area was not overflowed by the lake, and is a region of narrow, ramifying valleys, narrow, steep-sided, dividing ridges, whose directions are toward every point of the compass, and whose perfectly coinciding horizontal strata prove conclusively their subaerial erosion. The ravines are all in direct proportion to the relative sizes of the streams in them. North and east of this driftless area, from 25 to 75 miles, there is a scantiness of drift and numerous outliers, attesting the ordinary effects of erosion. The "Stand Rock," in the dells of the Wisconsin, the isolated ridges and peaks in the central part of the State, rising from 100 to 300 feet abruptly from the low ground around them, and composed of horizontally stratified sandstone, or of sandstones capped with limestone, prove the regular erosion for ages, and are quite inconsistent with any single mechanical eroding power that must have operated upon the whole country alike. In Dakota County, Minnesota, there is an outlier of the St. Peter's sandstone known as "Lone Rock," whose summit is 100 feet higher than the surrounding country, and from which many other outliers are in view; and yet in the valleys the drift prevails and boulders abound. In Wabasha County, the "Twin Mounds," and in Olmsted County the "Sugar Loaf Mound" and the "Lone Mound," attest in like manner

the continent
the south
rise as high
inconsistent
or across it

§ 195.

British Col
ain regions,
of scratched
with boulders
of scratched
ders, as occur
country was
produced up
rivers that, i
borders of th
various place
the existence
of a heavy l
that border u
as evidence o
River Eocene
White River
sandstone tow
Montana, Dal
Garden of the
lone mountain
Mexico to Ala
ever moved so

§ 196. In

the continent;
On account of
if one had eve
such dimension
have been such
Had there been
ing their place
trary, no such
present flora an
and passing bac
The scratches a
exist upon the
overflowed by w
outliers that a g
The scratches a
glacial period;
elevations of lan

the continuing erosion since Silurian times. The two lonely towers in the valley of the south branch of Root River, in Fillmore County, known as "Eagle Rocks," rise as high as the rocky walls of the valley, and evidence subaerial erosion, but are inconsistent with the idea that any large body of ice ever passed down the valley or across it.

§ 195. There is no drift in California, nor on the Pacific Coast as far north as British Columbia and Alaska. There are no indications of it in the Rocky Mountain regions, or upon the great plains of the West. There are no such exhibitions of scratched and grooved rocks succeeded by fossiliferous marine clays and sands, with bowlders, as occur in the New England States and St. Lawrence region, nor of scratched rocks and ancient soils succeeded by clay, sand, and gravel, with bowlders, as occur in the central part of the continent; but, on the contrary, the whole country west of the Mississippi Valley is absolutely driftless, except as to local drift produced upon the shores of Tertiary lakes, and more or less distributed by the rivers that, in the course of time, cut out the cañons which drained them. On the borders of the ancient lakes and rivers there are terraces, marking shore-lines at various places from Mexico to Alaska, but they are standing monuments to disprove the existence of a continental ice-sheet; for no one can conceive of the movement of a heavy body of ice across a valley without disturbing the graveled terraces that border upon both sides at different elevations. The natural towers that stand as evidence of erosion from the Wahsatch times to the present; from the Green River Eocene to the present; from the Bridger Eocene to the present; from the White River Miocene to the present; the columnar masses, irregular pyramids, sandstone towers, and turreted outliers of the Bad Lands of Colorado, Wyoming, Montana, Dakota, and British Columbia; the monuments on Monument Creek; the Garden of the Gods; the buttes in all the mountain chains; the transverse ridges, lone mountains, and exalted peaks; and the whole array of cañons from Texas and Mexico to Alaska,—all alike tell us, in language unmistakable, that no glacial sheet ever moved south upon the western plains or mountain ranges.

§ 196. Indeed, there is no evidence a glacial sheet ever existed on any part of the continent; none that gives any warrant to the hypothesis of a glacial period. On account of the valleys, hills, and mountains, no glacial sheet could move; and if one had ever existed, the waters flowing from it would have cut out channels of such dimensions they could have been not only traced, but their dimensions would have been such they could not be mistaken for any of the valleys now existing. Had there been a glacial period, northern plants and shells would be found occupying their places as far south as Florida, Louisiana, and Texas. But, on the contrary, no such flora or fauna is found farther south than it now exists, while the present flora and fauna occur in the same latitude throughout the Post-pliocene age, and passing back through earlier ages, unmolested by any visible climatic changes. The scratches and furrows so often cited as evidence of the glacial period do not exist upon the mountains, but occur only in the valleys and lower lands that were overflowed by water; and in these valleys there are now standing lone rocks and outliers that a glacier moving in the valleys would necessarily have swept away. The scratches and furrows are readily accounted for without the hypothesis of a glacial period; and on account of their position on the northern side of the higher elevations of land and not upon the southern, and their universal course up the

valleys from the lakes without regard to the direction of the valleys, they can not be accounted for as glacial phenomena, for they are wholly inconsistent with it. The glacial epoch is a theoretical blunder, not supported by scientific facts or intelligent reasoning, and contrary to all geographical, geological, and palæontological information. There is no such geological period, and no gap into which it can possibly be injected.

CHAPTER XXXIX.

NOMENCLATURE.

THE rules of nomenclature are, with few exceptions, firmly established. They have resulted from years of experience and reflection, and tend to secure fixity and convenience in the designation of animals and plants. *Each animal and each plant has a name consisting of two words—the first generic, and the second specific. This is called the binomial system, or Linnæan method of nomenclature. The genera are arranged in families, the families in orders, the orders in classes, and the classes in subkingdoms. These divisions are sometimes further separated into sections or intermediate groups, often distinguished by the prefixes *sub* and *super*.

Linnæus first consistently applied the binomial system of nomenclature to all classes of organisms in 1758, in the 10th edition of *Systema Naturæ*; but he applied it to botany in *Species Plantarum*, published in 1753. It had been used intermittingly by earlier authors. Naturalists have generally adopted 1753 as the starting-point for the binomial system in botany, and 1758 for zoology, or, without reason, the 12th edition of *Systema Naturæ*, published in 1766. It can make no difference in palæontology which is regarded as the starting-point, for the last precedes the science. The names in the binomial system assume the Latin form by taking a Latin termination.

DENOMINATION OF HIGHER GROUPS THAN GENERA.

The names of groups higher than genera are usually taken from some of the principal characters. They are expressed by single words of Greek or Latin origin, in which a certain harmony of form and termination is preserved for groups of similar nature; as, *Phanerogamæ*, *Cryptogamæ*; *Cephalopoda*, *Gasteropoda*.

Compounds of Greek and Latin words are not allowable. In cryptogamic botany, ancient names of families, such as *Musci* and *Filices*, have been employed as names of classes or sub-classes. Botanical cohorts or sub-cohorts are designated by the name of one of their principal families, with the termination *ales*.

The families in botany are designated by the name of one of their principal genera, with the termination *aceæ*, as *Rosa*, *Rosaceæ*; *Ranunculus*, *Ranunculaceæ*. To which there are the following exceptions: 1. When the genus from which the

*NOTE.—See Report of the 12th Meeting of the British Association for the advancement of Science, held at Manchester in June, 1842, Reprinted Cin. Quar. Jour. Sci., Vol. I, p. 351; Report of the British Association at Birmingham, in 1865, and Report of the Committee (W. H. Dall) on Zoological Nomenclature, to section B. of the American Association for the Advancement of Science, at the Nashville Meeting in 1877. The authorities are quite fully cited in the latter report.

name of the
the termin
Tamarix,
is derived
family form
terocarpeæ,
under their
Compositæ,
but applied
a family nar
Botanic
in them, wit
tribes which
The nar
the earliest
families by a
not *Strixidæ*;
but in *ine* it
Names o
because their
therefore allo
And when a
another famil
the remaining

The rule
Latinized by
the following

α
 β
 γ
 δ
 ϵ
 ζ
 η
 η fi
 θ , θ
 ι
 κ
 λ
 μ
 ν
 ξ
 $\omicron$, ω
 π

name of the family is taken ends in Latin with *ix* or *is* (genitive *icis*, *idis*, or *iacis*), the termination *iceæ*, *ideæ*, or *ineæ* is permitted; as, *Salix*, *Salicineæ*; *Berberis*, *Berberideæ*; *Tamarix*, *Tamariscineæ*. 2. When the genus from whence the name of the family is derived has a name of inconvenient length, and there is not a tribal name in the family formed from the same generic name, the termination *æ* is admitted; as, *Dipterocarpeæ*, from *Dipterocarpus*. 3. For some very large families universally known under their exceptional names, the ancient designation is preserved; as, *Cruciferaæ*, *Compositæ*, and *Gramineæ*. 4. An old generic name no longer preserving that rank, but applied only to a section, or even a species, may be maintained as the base of a family name; as, *Hippocastaneæ*, from *Aesculus hippocastanum*.

Botanical sub-families are formed from the name of one of the genera contained in them, with the termination *æ* or *ineæ*, and also the names of tribes and sub-tribes which take the termination *æ*; as, *Roseæ*, from *Rosa*.

The names of zoological families are formed by adding the termination *idæ* to the earliest known, or most characteristic genus contained in them; and of sub-families by adding the termination *inæ*; as, *Terbebratula*, *Terebratulidæ*; *Strix*, *Strigidæ*, not *Strixidæ*; *Buceros*, *Bucerotidæ*, not *Bucerosidæ* or *Buceridæ*. The *i* in *idæ* is short; but in *inæ* it is long.

Names of higher rank than genera are not rigidly subject to the law of priority, because their limits fluctuate with the advancement of science, and changes are therefore allowable when newly discovered facts have made the name erroneous. And when a genus from whose name a family name has been taken, is removed to another family, the family name may be dropped, and a new one may be coined for the remaining genera.

ORTHOGRAPHY.

The rules of Latin orthography must be adhered to. Greek names are Latinized by substituting for the Greek letters their Latin equivalents, according to the following table:

α	=	a;	(βῆτα)	Beta.
β	=	b;	(βραχίων)	Brachium.
γ	=	g;	(γλῶσσα)	Glossa.
δ	=	d;	(διψάς)	Dipsas.
ϵ	=	e;	(δάλειος)	Hyalea, not Hyalæa.
ζ	=	z;	(ζίζυφον)	Zizyphus, Zizyphinus.
η	=	e;	(πειρήνη)	Pirena, not Pirina.
η final	=	a;	(πειρήνη)	Pirena, not Pirene.
θ, θ	=	th;	(τηθύς)	Tethys; (θέτις) Thetis.
ι	=	i;	(βαλιός)	Balia, not Balea.
κ	=	c;	(ιπποκρήνη)	Hippocrena, not Hippochrenes.
λ	=	l;	(φυλλίς)	Phyllis.
μ	=	m;	(μέλας)	Melas.
ν	=	n;	(πειρήνη)	Pirena.
ξ	=	x;	(ξένος)	Xenus, Xenophora.
$\omicron, \omega$	=	o;	(φορός)	Phorus; (πῶμα) Poma.
π	=	p;	(ποταμός)	Potamus.

Resolution Test Chart Labels:

- 1.0
- 1.1
- 1.25
- 1.4
- 1.6
- 1.8
- 2.0
- 2.2
- 2.5
- 2.8
- 3.2
- 3.6
- 4.0

Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

1.5 1.8 2.0 2.2 2.5 2.8 3.2 3.6

10
01

ρ	=	r;	(πτερόν)	Pterum.
ρρ	=	rrh;	(φυλλίρ-ροη)	Phyllirrhoa, not Phyllirhoe.
σ, ς	=	s;	(γλωσσός)	Glossus.
τ	=	t;	(πτερόν)	Pterum.
υ	=	y;	(ὕβος)	Hybolithus, not Hibolites.
φ	=	ph;	(φορός)	Phorus.
χ	=	ch;	(κοχλίας)	Cochlias.
ψ	=	ps;	(ψάμμος)	Psammus.
αι	=	æ;	(λιμναῖος)	Limnæa, not Limnea.
αυ	=	au;	(γλαυκός)	Glaucus.
ει	=	e;	(τεῖνω)	Exotenobranchia.
ῥι	=	i;	(χέιλος)	Chilostoma, not Cheilostoma.
ευ	=	eu;	(εὐρος)	Eurus.
αι, οι	=	oe;	(δῖς, οἰκέω)	Dioeca, not Dioica.
ον final	=	um;	(ἐφίππιον)	Ephippium, not Ephippion.
ος final	=	us;	(εὐομφαλός)	Euomphalus, not Euomphalos.
ου	=	u;	(λουτήριον)	Luterium, not Lotorium.
γγ	=	ng;	(ἀγγαρεία)	Angaria.
γχ	=	nch;	(ἀγχιστρον)	Anchistoma, not Angistoma.
γκ	=	nc;	(ἀγκιστρον)	Ancistrodon, not Agkistrodon.
ρ	=	rh;	(ῥέα)	Rhea.
ερμαία	=	h;	(ερμαία)	Hermæa, not Ermæa.

It follows therefore, that *Bythotrephix* must, according to the laws of etymology, be spelt *Bythotrephix*; *Xenophasia*, instead of *Zenophasia*; *Pæocephala*, instead of *Poiocephala*. In Latinizing modern words where the rules of classic usage do not apply, the etymology must be preserved, even though it includes letters and combinations unknown in Latin; thus, *woodwardi*, instead of *vudvardi*; *knighti*, instead of *enichti*; *bullocki*, instead of *bullocci*; *escheholtzi*, instead of *essolzi*; *nebrascensis*, instead of *nebrascensia*. But words of barbarous origin should be rendered as classical in appearance as is consistent with the preservation of their original sound; as, *toecus*, instead of *tockus*; *ansure*, instead of *ansuree*; *argunda*, instead of *argyondat*.

In Latinizing proper names and converting them into specific ones, they assume a distinctive character, which they did not before possess. The rule is to use the termination *us*, genitive *i*, when the name ends with a consonant; as, *Miller*, *milleri*. But when it ends in a vowel, *ius*, genitive *ii*; as, *Moore*, *moorii*. This rule is often violated, but it would be much better strictly to adhere to it.

PRIORITY.

It is of the highest importance that we retain the first defined and illustrated names of genera and species. The British Association said:

"It being admitted on all hands that words are only the conventional signs of ideas, it is evident that language can only attain its end effectually by being permanently established and generally recognized. This consideration ought, it would seem, to have checked those who are continually attempting to subvert the established language of Zoology, by substituting terms of their own coinage. But, forgetting the true value of language, they persist in confounding the name of a species or

group w
of expre
some new
unknown
were to
cases wh
of the n
and shou
would no
before us
we agree
stamped
Zoology,
by the pe
hence it
of elegan
principle
injustice
owe our fi
mission of
selves int

"The
a species,
synonyms.

"As
science is
complete,
of genera
It thus bec
definitions
justice to
and it is n
its nomencl
being made

"A g
subsequent
constituent

"Wh
retained for
acters as at
species as
When they
first species
was regarde
serve to poi
the original
the old genu

group with its *definition*; and because the former always falls short of the fullness of expression found in the latter, they cancel it without hesitation, and introduce some new term which appears to them more characteristic, but which is utterly unknown to the science, and is therefore devoid of all authority. If those persons were to object to such names of men as *Long*, *Little*, *Armstrong*, *Golightly*, etc., in cases where they fail to apply to the individuals who bear them, or should complain of the names of *Gough*, *Lawrence*, or *Harvey*, that they were devoid of meaning, and should hence propose to change them for more characteristic appellations, they would not act more unphilosophically or inconsiderately than they do in the case before us; for, in truth, it matters not, in the least, by what conventional sound we agree to designate an individual object, provided the sign to be employed be stamped with such an authority as will suffice to make it pass current. Now, in Zoology, no one person can subsequently claim an authority equal to that possessed by the person who is the first to define a new genus or describe a new species; and hence it is that the name originally given, even though it may be inferior in point of elegance or expressiveness to those subsequently proposed, ought as a general principle to be permanently retained. To this consideration we ought to add, the injustice of erasing the name originally selected by the person to whose labors we owe our first knowledge of the object; and we should reflect how much the permission of such a practice opens a door to obscure pretenders for dragging themselves into notice at the expense of original observers."

"The name originally given by the founder of a group, or the describer of a species, should be permanently retained to the exclusion of all subsequent synonyms."

"As the number of known species which form the ground-work of zoological science is always increasing, and our knowledge of their structure becomes more complete, fresh generalizations continually occur to the naturalist, and the number of genera and other groups requiring appellations is ever becoming more extensive. It thus becomes necessary to subdivide the contents of old groups, and to make their definitions continually more restricted. In carrying out this process, it is an act of justice to the original author that his generic name should never be lost sight of, and it is no less essential to the welfare of the science, that all which is sound in its nomenclature should remain unaltered amid the additions which are continually being made to it."

"A generic name, when once established, should never be canceled in any subsequent subdivision of the group, but retained in a restricted sense for one of the constituent portions."

"When a genus is subdivided into other genera, the original name should be retained for that portion of it which exhibits in the greatest degree its essential characters as at first defined. Authors frequently indicate this by selecting some one species as a fixed point of reference, which they term the 'type of the genus.' When they omit doing so, it may still in many cases be correctly inferred that the first species mentioned on their list, if found accurately to agree with their definition, was regarded by them as the type. A specific name or its synonyms will also often serve to point out the particular species, which by implication must be regarded as the original type of a genus. In such cases we are justified in restoring the name of the old genus to its typical signification, even when later authors have done otherwise."

"The generic name should always be retained for that portion of the original genus which was considered typical by the author."

"Example.—The genus *Picumnus* was established by Temminck, and included two groups, one with four toes, the other with three, the former of which was regarded by the author as typical. Swainson, however, in raising these groups at a later period to the rank of genera, gave a new name, *Asthenurus*, to the former group, and retained *Picumnus* for the latter. In this case we have no choice but to restore the name *Picumnus* Tem., to its correct sense, canceling the name *Asthenurus* Sw., and imposing a new name on the three-toed group which Swainson had called *Picumnus*."

"When no type is indicated, then the original name is to be kept for that subsequent subdivision which first received it."

"When the evidence as to the original type of a genus is not perfectly clear and indisputable, then the person who first subdivides the genus may affix the original name to any portion of it at his discretion, and no later author has a right to transfer that name to any part of the original genus."

"When an author infringes the law of priority by giving a new name to a genus, which has already been properly defined and named, the only penalty which can be attached to this act of negligence or injustice, is to expel the name so introduced from the pale of science."

"When two authors define and name the same genus, both making it exactly of the same extent, the later name should be canceled *in toto*, and not retained in a modified sense."

"No special rule is required for the cases in which the later of two generic names is so defined as to be *less extensive* in signification than the earlier; for if the later includes the type of the earlier genus, it would be canceled by the operation of the rule that the generic name should always be retained for that portion of the original genus which was considered typical by the author."

"If the later name be so defined as to be equal in extent to two or more previously published genera, it must be canceled, *in toto*."

"A genus compounded of two or more previously proposed genera, whose characters are now deemed insufficient, should retain the name of one of them. If these original generic names differ in date, the oldest one should be the one adopted."

The committee on zoological nomenclature, appointed by the American Association for the Advancement of Science, said:

"A change in the diagnostic characters, or a revision which carries with it the exclusion of certain elements of a group, or the inclusion of new elements, does not authorize the change of the name or names of a group."

"When a group or genus is divided into two or more groups, the original name must be preserved and given to one of the principal divisions. The division including the typical species of the primitive genus, if any type had been specified, or the oldest, best known, or most characteristic of the species originally included when the primitive genus was first described by its author, is the portion for which the original name is to be preserved. If there is no section specially so distinguished, that which retains the larger number of species should retain the old name, but the latter can not be applied to a restricted group containing none of the

species r
scribed, c

The
its type
to be we
it under
violate it
genus *Re*
ing that
Rhynchot
not been
Rhynchos
Rhynchot
a genus,
nated cha
and reliab

When
as the typ
species m
genus, du
species as

In di
typified b
genus, the
founded o
may have
divided a
which oth

In ca
oldest nar
date, the
to be equ
When it i
name mus

A gen
bear the s

Public
acters in a
tributed an
an underst
nition or i
mand in
recognized.
so minute,
those conv

species referred to the primitive group by its author at the time when it was described, or when he enumerated the species contained in it."

The rule that a subsequent author can not revise a genus and substitute as its type a species different from that relied upon by the founder of the genus seems to be well settled in England and America. The instances of strictly adhering to it under circumstances where it would have seemed to accommodate the author to violate it, are numerous. For instance, Professor Hall, mistaking the type of the genus *Retzia*, proposed and defined the genus *Rhynchospira*; afterward ascertaining that *Rhynchospira* was a synonym for *Retzia*, he abandoned it and proposed *Rhynchotreta* for the form which he had originally mistaken for *Retzia*. Had it not been for this rule he might have abandoned *Retzia evax* as the type of his genus *Rhynchospira*, and substituted *Rhynchonella cuneata*, which became the type of *Rhynchotreta*. If you can substitute another than the original species as the type of a genus, I can substitute another, and so we destroy all fixity in the type and designated characters, throw the science into confusion, and seriously impair the value and reliability of generic characters.

When an author has specified no type, the first species defined is to be taken as the type, or if the genus is to be divided, no type having yet been selected, a species may be chosen from among those originally specified as belonging to the genus, due regard being paid to the necessity of retaining as many of the original species as possible in the division which is to retain the old name.

In dividing a genus of which there are already synonyms, if the synonyms are typified by the same species or group of species selected as types of the primitive genus, they should not be again used. When, however, the so-called synonyms are founded on species belonging to different sections of the genus, although the names may have been considered coextensive in their application, and the genus is to be divided accordingly, the so-called synonyms become the proper designations for which other names can not be applied.

In case of the consolidation of two or more groups of the same nature, the oldest name must be retained for the whole. If both, or all, are of the same date, the reviser may select the one to be retained. If a name be so defined as to be equal in extent to two or more previously described, it must be canceled. When it is necessary to divide a species, the form which received the old specific name must retain it.

A generic name must have a single meaning, and therefore two genera can not bear the same name, even though belonging to distinct subkingdoms.

AS TO PUBLICATION.

Publication consists of the insertion of a distinct exposition of essential characters in a printed book which is kept for sale, or which has been generally distributed among those conversant with the subject. Where figures are necessary to an understanding of the character of the organism, they must accompany the definition or it will be invalid. The tendency of the science of palæontology is to demand in all cases both definition and illustration before the publication is to be recognized. There are many species whose characters are so complicated and parts so minute, that an exposition of the essential ones, so they may be understood by those conversant with the fossils in the class, can only be made by illustration

accompanied by proper definition; the science therefore demands the rule shall be co-extensive with its necessities, and good authors refuse to recognize names unless the publication is such that their meaning may be readily comprehended.

A communication in a public assembly or learned society, or the reading of a paper containing new names at such meeting, printing of the names in a catalogue, labeling the fossils in a collection, printing the names and description in a newspaper, either one or all these attempts to introduce the names, does not constitute a publication within the rule, and hence give the names no place in science. Nor does the printing of the names with brief definitions in an obscure pamphlet, or even in the Journal of a learned society, where the definition will not enable an ordinary paleontologist to identify or distinguish the species at another locality than the typical one, give them any right to claim recognition. Occasional pamphlets independently issued, and insufficiently advertised and distributed, or very small editions that can not reach the students of the science generally, are not publications within the rule.

The date borne by a publication will be presumed to be accurate, though this presumption is only *prima facie*, and may always be contested, and the true date shown, from which time alone do names have any validity.

A species is not to be considered as named unless both generic and specific names are simultaneously applied to it.

Where a genus or species is announced in a publication, and subsequently described in another publication, the latter only is entitled to recognition. It is essential in establishing a genus that some species be referred to it.

NAMES TO BE REJECTED, CHANGED, OR MODIFIED.

A generic name should be rejected when it has been previously applied to another valid genus of organisms, even if it has received general currency. It should also be rejected when it expresses a positively false character in the genus, and is therefore liable to propagate error, and especially is this the case where the definition is so erroneous as not to entitle it to recognition; but where the name has received general circulation, and the error is not such as to seriously mislead, the name is retained; as, *Athyris* and *Atrypa*. So a specific name should be rejected when it is already applied to another species or subdivision in the same genus, or when a geographical name of a country entirely removed from the habitat of the species is used.

A name should be rejected when it is formed of two words belonging to different languages, as *en* put before a Latin name, *sub* before a Greek name, *oides*, *opsis* suffixed to a Latin name; or when it is identical if properly spelled, according to its true derivation, with a prior valid name, as *Platystoma* of Conrad, being preoccupied, can not be retained simply because he misspelled it *Platystoma*.

A name should always be rejected when it outrages decency.

It is inelegant and tautological to derive a generic name from the specific name of its typical species. For example, *Corvus pyrrhocrax*, Linn., was afterward advanced to a genus under the name of *Pyrrhocrax*. The name therefore became *Pyrrhocrax pyrrhocrax*. The rule is now to reject all such generic names, except those which, from long usage, have imbedded themselves into science; none of which, however, can claim a place in palæontology.

Wh
species o
synonym
name is
to anothe
and the
of the la
Wh
other one

The
guishing
it is more
Greek and
faulty con
der is giv
of a natu

Whe
real ortho
first syllab
dictionarie
botanical o
after the C
Glyptocrin
only becau
tionaries, h
the fixity
named afte
sistent wit
the name t

Name
is prefixed
cause the r
has no plac
be rejected

The n
name can n
binomial, a
eminently f
collected.
have been c
trades and p
generally in
mythologica
is conceded.
centennialis

When a species is transferred from one genus to another in which there is a species of the same name, the older specific name is retained, and the oldest tenable synonym is adopted for the other form, if there be one; and if not, a new specific name is proposed. But if the form bearing the prior specific name is transferred to another genus, the original specific name of the later species must be restored, and the new specific name must fall into synonymy. This is the necessary result of the law of priority.

When a name is published, the author has no more control over it than any other one. He has the same rights, no more and no less, than other naturalists.

SELECTION OF NAMES AND MODIFICATION.

The best names are derived from Latin and Greek, and express some distinguishing characteristic of the object to which they are applied. In palæontology it is more consistent with practice and uniformity to derive the generic names from Greek and the specific names from Latin; and if the name as proposed exhibits a faulty construction, any naturalist is authorized to correct it. When a wrong gender is given to a species by its termination, not agreeing with a genus, it is the duty of a naturalist to correct it.

When a name derived from a person has not been written according to the real orthography of his name, it may be changed, provided it does not involve the first syllable and thereby disturb the arrangement of indices, tables, catalogues, and dictionaries, in alphabetical order, or interfere with long-established usage. The botanical congress at London, in 1866, refused to change the name *Cinchona*, named after the Countess Chincon, because of established usage. In 1866 Hall described *Glyptocrinus nealli* in honor of O'Neill; but the name must stand as described, not only because its change would interfere with indices, tables, catalogues, and dictionaries, but Hall had the right to construct the specific name *nealli* as he did, and the fixity of nomenclature will not allow another to change it. *Scalaria turtoni*, named after Miss Turton, may be changed to *S. turtonæ*; and *Viviparus* being inconsistent with itself may be changed to *Vivipara*, because the change is in the end of the name to conform to the rules of grammar.

Names of persons are Latinized and not adopted in Greek form, but where *en* is prefixed or *oides* or *opsis* suffixed, one may not be authorized to change it, because the name is not of Latin origin, though it is in very bad taste. Buffoonery has no place in science; hence Latin puns on names, as *faba* after Mr. Bean, should be rejected in all cases as a poor joke.

The name of a person must have the termination Latinized, but the specific name can not be composed of the Christian and surname, because it would not be binomial, and can not be made to conform to the rules. Geographical names are eminently fit and suitable when they indicate the locality from which the type was collected. Barbarous names are not in good taste in Palæontology, though they have been defended in other departments of Natural History. Names expressive of trades and professions are not in good taste. Mythological and historical names are generally in bad taste for specific names, though they have been largely used; but mythological names for genera have usually been defended. The right to use both is conceded. Names expressive of something else than a character of the fossil, as *centennialis* for a *Hyolithes* are in very bad taste, and sometimes even absurd. Com-

parative names are often appropriate; but those expressive of size, as *maximus*, *minor*, and *minimus*, are too frequently rendered inaccurate by after discoveries, and are therefore objectionable.

Both generic and specific names derived from persons engaged in palæontological pursuits are very appropriate. Names of harsh and inelegant pronunciation ought to be avoided, as also words of too great length or having more than five syllables.

Generic names may be compounded from other genera to express the position of the genus as intermediate to or allied with two other genera, care being taken not to adopt such as are of too great length, and not to corrupt them in trying to render them shorter. *Aviculopecten* and *Aviculopinna* are examples of the appropriate use of compound words, notwithstanding their length, while *Tellinomya* is more fanciful than real, and yet not to be discarded.

In compounding words all the radical or essential parts of the constituent members must be retained, and no change made except in the variable terminations. Words coined at random, or without any derivation or meaning, will not be recognized.

The names of genera are in all cases essentially substantive, and hence adjective terms can not be employed for them without doing violence to grammar; for instance, *Anomaloides* proposed as a generic name must be disregarded. The same may be said of names in the genitive case, which are wholly inadmissible, without reformation.

FORMATION OF NAMES.

The generic name always begins with a capital letter, the specific name with a small initial letter, even when derived from person or place. The generic name is a noun, while the specific name has the force of an adjective. The specific name is in no instance a proper noun, but all species are equal, and should therefore be written alike. It is a violation of a plain rule of grammar to write a specific name with a capital letter; beside, there is an advantage in obeying the rule, for by so doing the eye at a glance distinguishes specific from generic names.

The generic name retains the gender which belongs to it in the language from which it is taken. Where no change is made in the termination of the last word in a generic name, the gender of that word determines the gender of the genus. Thus *cerus*, *nemus*, *stoma*, and *desma* are in the Greek of the neuter gender, and consequently all genera ending with these words, such as *Orthocerus*, *Loxonema*, *Phragmostoma*, *Lyrodesmus*, are neuter.

In defining a new genus the etymology should be given, and a species should be selected as the type. There is no excuse for neglecting these rules, except that the author is incapable of giving the etymology of his proposed generic name, and is not sufficiently confident of his definition to dare venture to rely upon one of his species as the type.

When a generic name is derived from the name of a person, it is stripped of all titles and preliminary particles, reduced to the genitive case, and the letter *a* is appended, thus taking on a feminine form. The following examples illustrate the method, viz.: *Names*, Brun, Bruni, Bruno, Brunus, Bruna, Brune, Bruny. Generic form, Brunia, Bruniia, Brunoia, Brunusia, Brunæa, Brunia, Brunyia. Y at the end of a word of one syllable is treated as a consonant, as Quoy, Quoyia; Gay, Gayia; and mute *e* final becomes *i*, or is dropped entirely, as Pouse, Pousia.

Even
if an ad
the term
of a com
of a per
The

—ad
in words
be annex

—an
be added
confined
mination
euz (lescu
dins, claus
tion must
geinitzanus

—atu
the name
only; as,
tus; lobus,
tion some
from a lin
the nomen

—form
confined to
i; as, later
first given
necting vo
more awk
squama, squ

—ensi
fore be cor
been traver
illustration
morsii, bein
ensis, Card
terms are fi
become loca
In using th
If the name
word; as, e
dropped, an
nesolensis;
3d. If the r
ensis; Misso
chouteauensi

Every specific name agrees in gender with the genus to which it belongs, and if an adjective, its termination must show it. If the specific name is a substantive, the termination is not necessarily changed. The rule is not to change the ending of a common noun or mythological name, but to make an adjective, and the name of a person or place, indicate the gender of the genus to which it belongs.

The following rules govern the use of these terminations:

—*alis*. This Latin termination, implying resemblance, is seldom used, except in words already compounded in Greek and Latin; and when otherwise, it must be annexed to the stem of the word, as *rectilateralis*, *quadrilateralis*.

—*anus*. This Latin termination implies resemblance or association, and may be added to proper names, personal or local; though in science its use is almost confined to the former. If the word is capable of taking a classic form, the termination should be simply annexed to the stem as *Linneus*, *linneanus*; *Lesqueret* (*lescuria*), *lescurianus*; in conformity with classic usage; *pagus*, *paganus*; *Claudius*, *claudianus*; *Neapolis*, *neapolitanus*. In other cases, the addition of this termination must follow the same rule as those for *ensis*, as *America*, *americanus*; *Geinitz*, *geinitzianus*; *Meek*, *meekanus*; *Erie*, *erianus*; *Italy* (*ia*), *italianus*.

—*atus*. This Latin termination strictly implies the possession of the thing to the name of which it is added. It is therefore affixed to the stem of common names only; as, *costa*, *costatus*; *galea*, *galeatus*; *fornix*, *fornicatus*; *sinus*, *sinuatus*; *stria*, *striatus*; *lobus*, *lobatus*; *rostrum*, *rostratus*. It is worthy of remark here that this termination sometimes loses its *at*, to shorten the word. The practice is not commendable from a linguistic stand-point, but some of the terms so made have become fixed in the nomenclature; as, *Orthis biloba*.

—*formis*. This Latin termination implies resemblance of shape, and should be confined to Latin words, to the stem of which it is joined by the connecting vowel *i*; as, *laterna*, *laterniformis*; *pistillum*, *pistilliformis*. In forming terms, such as the first given above from Latin words ending in *a*, the error of using *a* as the connecting vowel should be avoided; being inconsistent with classic usage, as well as more awkward and lengthy, thus we have from *terra*, *terricola*; *gemma*, *gemmafer*; *squama*, *squamiger*; *tuba*, *tubiformis*; etc.

—*ensis*. This is a Latin termination, expressive of locality, and can not therefore be correctly employed, except as an affix to the name of a place. This rule has been traversed in few real, but in many apparent instances. *Lingula morsensis* is an illustration of the former. In accordance with law, this has been changed to *L. morsii*, being given in honor of Mr. Morse. *Zygospira cincinnatiensis*, *Pupa vermilionensis*, *Cardium napoleonense*, *Athyris hannibalensis* are apparent exceptions; but these terms are formed from words which, though originally personal or trivial, have now become local names, and consequently no valid objection can be raised against them. In using this termination the following rules have been generally followed: 1st. If the name of the place ends in a consonant, the termination is annexed to the word; as, *Clinton clintonensis*. 2d. If the name ends in *a* or *e*, these letters are dropped, and the termination then annexed; as, *Canada*, *canadensis*; *Minnesota*, *minnesotensis*; *Iowa*, *iowensis*; *Indiana*, *indianensis*; *Lasalle*, *lasallensis*; *Erie*, *erienensis*. 3d. If the name ends in *i*, *o*, or *u*, that vowel is retained; as, *Mississippi*, *mississippiensis*; *Missouri*, *missouriensis*; *Chicago*, *chicagoensis*; *Colorado*, *coloradoensis*; *Chouteau*, *chouteauensis*. 4th. If the name ends in *y*, that letter becomes *i* upon the addition

of the termination; as, *Kentucky, kentuckiensis; Alleghany, alleghaniensis*; in accordance with classic usage, as *Sicily, sicilensis*.

—*i*. The termination *i* is to be considered a mere indication of the Latin genitive case, and custom rather than correctness has, in some sense, legalized its addition to any name. In practice, however, it is almost restricted to proper names. Thus we have *knighti, littoni, flemingi, ivesi*.

—*icus*. This Greek termination implies resemblance, and may be added to common names under the same rules as those given for —*ensis*, except that, in forming the word, a vowel is suppressed if it would precede the termination; thus, *Macedon, macedonicus; Italy (ia), italicus*. It is little used, except as an affix to the name of a river or country; as, *euphraticus, anglicus*, or in such words as *ellipticus*.

—*eus*. This Latin termination has been occasionally employed; but as it implies "made of," it is evidently seldom, if ever, admissible in palæontology. The term *eboraceus*, from *eboracum*, the Latin name for York, is a misnomer and should have been *eboracensis*.

—*inus*. This termination is applied to both common and proper names. Latin usage restricted its application more than modern scientific practice has done, and applied it mainly to proper names, local terms, and living beings; as, *caninus, alpinus*; but did not sanction such words as *rugatinus, sulcatinus, secalinus, taxinus*, and *velutinus*. The termination is used subject to the same laws as —*ensis*.

—*ites*. This termination expresses the fossil nature of the specimen. It is a contraction of the Greek word *lithos*, a stone. In most instances it coalesces with the last vowel of the root. This and long usage in many words, such as *Ammonites, Belemnites, Pyrites*, have completely established the long *i*, while the gender is determined by that of the Greek word to be masculine. All specific terms in the genus must, therefore, be of this gender.

—*oides*. This Greek termination, signifying "like," should be added only to the stems of words of Greek origin. No connecting vowel is necessary. Thus we have *dactylos, dactyloides; discos, discoides*. The Latin form —*oideus* obeys the same laws, except the Greek termination is alike in all genders, while the Latin is inflected as Latin adjectives of similar termination.

Compound terms. In forming compound terms care should be taken to connect them rightly. If an adjective of three terminations, or a noun of the second Latin declension, composes the first part of the word, either *i* or *o* may be employed as a connecting vowel, the choice being largely determined by the ear. Thus *sulcomarginatus* is better than *sulcimarginatus*, and *crassicaulis* than *crassocaulis*. If the adjective has but one or two terminations, or the noun be of the first, third, or fourth Latin declension, the connecting vowel *i* should always be employed; as, *tenuistriatus, pinniformis, ilicifolius, retiformis, cornifer*. The connecting vowel *o* is admissible by Greek usage in all declensions; as, *Ulodendron, Cycloconcha, Syringodendron, Alethopteris, Dictyonema, Dictyopteris*, except that where the first part of the word is an adjective ending in —*ys*, it is shorter, and at the same time consonant with classic usage to employ no connecting vowel at all; thus, *pachyderma, euryteines, Platystoma*, etc., are better than *pachyderma, euryteines, Platystoma*, etc.

P

By the
world. But
added the v
all which h
and the "A

THE PR

Equisetaceae,
called Sargassum,
the leathery
casts, showing
readily disintegrate
the sand, so
exposed to a
isms for more
gelatinous matter
the living Fungi
representative

The fossil
Families. The
Asterophycus
phycus, Chonetes
cophycus, Dystactos
Licorophycus,
Taenidites, Trilobites

The Fungi
riage). They
as to escape de
in other cases
Palaeozoic rock
zomorphia to it

The vascular
iferous era, the

NORTH AMERICAN PALÆOZOIC FOSSILS.

By the little words plants and animals we include all the organisms in the world. But science, demanding technical words and controlling characteristics, has added the word "Kingdom" to these common names; and hence all organisms and all which have existed in the past are divided between the "Vegetable Kingdom" and the "Animal Kingdom."

VEGETABLE KINGDOM.

THE Palæozoic Fossil plants are divided into seven classes; viz., Fucoides, Fungi, Equisetaceæ, Filicaceæ, Lycopodiaceæ, Cordaitæ, and Coniferæ. The Fucoides are also called Sargassites and Thalassophytes. They are supposed to have some affinity with the leathery marine vegetation called Fucus or the Sargassum. The fossils are merely casts, showing, as a rule, no structure whatever. Lesquereux says marine vegetation readily disintegrates and passes into a gelatinous, half-fluid matter, which penetrates the sand, so that the lowest strata of the great heaps thrown up by the waves and exposed to atmospheric action, do not generally preserve traces of their organisms for more than a year. The fossil forms may have been harder, and contained less gelatinous matter in their cells, and probably had only a remote resemblance to the living Fucus or Sargassum, though there can be no reasonable doubt they are representatives of extinct marine cryptogamous plants.

The fossils referred to this Class have never been distributed into Orders and Families. The genera are as follows: Archæophyton, Arthraria, Arthrophycus, Asterophycus, Astropolithon (Graptolite?), Blastophycus, Bythotrephes, Calamophycus, Chondrites, Conostichus, Cruziana, Dactylophycus, Dendrophycus, Discophycus, Dystactophycus, Eophyton, Heliophycus, Hippodophycus, Ichnophycus, Licrophycus, Palæophycus, Phytopsis, Protostigma, Rusophycus, Sphenothallus, Taonurus, Trichophycus.

The Fungi are cellular cryptogamous plants (*kruptos*, hidden; *gamos*, marriage). They are flowerless plants, in which the fructifying organs are so minute as to escape detection without a microscope. The spores are sometimes naked, and in other cases inclosed in a theca. The evidence of the existence of this Class in Palæozoic rocks is extremely meager, though Lesquereux refers a species of Rhizomorpha to it.

The vascular cryptogamous plants flourished to such an extent in the Carboniferous era, that it has been called the "Age of Acrogens," and the "Age of Coal-

plants." The Classes and Orders have been named as Latin adjectives in the feminine plural, to agree with *plantæ* (plants), which is said to be always understood. Thus from *Equisetum*, by prolonging the termination into *aceæ*, we have *Equisetacæ*; from *Filices*, *Filicacæ*, etc.

The *Equisetacæ* are either cellular or vascular flowerless plants, producing spores instead of seeds. The Palæozoic fossils are all referred to one Order, the *Calamariæ*. The genera are as follows: *Anarthrocanna*, *Annularia*, *Arthrostigma*, *Asterophyllites*, *Bechera*, *Bornia*, *Calamites*, *Calamodendron*, *Calamostachys*, *Equisetites*, *Macrostachya*, *Nematophyllum*, *Sphenophyllum*, *Volkmannia*.

The *Filicacæ*, or ferns, are too common among existing plants to have escaped the notice of any one. The Palæozoic ferns are divided into Orders as follows:

1. ORDER, NEUROPTERIDEÆ.

Cyclopteris, *Dictyopteris*, *Lesleya*, *Neuropteris*, *Odontopteris*.

UNCERTAIN RELATION TO THE ORDER.

Baiera, *Cardiopteris*, *Danaëites*, *Idiophyllum*, *Megalopteris*, *Neriopteris*, *Orthogoniopteris*, *Tæniopteris*.

2. ORDER, ALETHOPTERIDEÆ.

Alethopteris, *Callipteridium*, *Callipteris*, *Lescuropteris*, *Protoblechnum*.

3. ORDER, PSEUDOPECOPTERIDEÆ.

Pseudopecopteris.

4. ORDER, PECOPTERIDEÆ.

Beinertia, *Cymoglossa*, *Lonchopteris*, *Oligocarpia*, *Pecopteris*, *Phyllopteris*.

5. ORDER, SPHENOPTERIDEÆ.

Eremopteris, *Hymenophyllites*, *Sphenopteris*.

6. ORDER, ADIANTITES.

Aneimites, *Archæopteris*, *Triphyllopteris*.

FERNS OF UNCERTAIN AFFINITY.

Asteropteris, *Crematopteris*, *Pachypteris*, *Rhacophyllum*.

SEPARATE FRUCTIFICATION OF FERNS.

Sorocladus.

RACHIS OF FERNS.

Rhachiopteris.

RHIZOMA OF FERNS.

Stigmarioides.

STEMS OR TRUNKS OF FERNS.

Caulopteris, *Megaphytum*, *Psaronius*, *Stemmatopteris*.

The
bogs, or
a few spe
and open
Pine," are
Carbonife
or more in
of Carbon

"The
sometimes
row, linea
midrib.
forms, whi
of the part
tially from
been estab
"The

or ovate sp
base of leav
the outside
small ones
distinct only
lizing pollen
ular seeds,
The clau

Acanthophyt
Knorria, Lep
lum, Lepido
angites, Spor

Didymophyllu
Stigmaria, Sy

The Corda
ally by fragme
and flowers.
intermediate b

The existing Lycopodiaceæ inhabit the deep shade of the forests, the surface of bogs, or the slopes of mountains, where there is a high degree of humidity, except a few species, which have the power of closing the leaves under the heat of the sun and opening them to receive the rain or fog. Some of them, like the "Ground Pine," are evergreens, and none of them grow beyond a few feet in length. Many Carboniferous plants of this Class, however, were grand and stately trees, two feet or more in diameter, and fifty feet or more in length. Lesquereux says, in speaking of Carboniferous plants:

"The leaves of the Lycopodiaceæ are generally in a spiral order, modified sometimes in their relative disposition, even in the same species. They are narrow, linear-lanceolate, of various length, according to species, all with a strong midrib. Their point of attachment upon the stems is marked by scars of divers forms, which greatly vary in size, according to the age of the fragments, or rather of the part of the tree from which the fragments of bark are derived. It is essentially from the characters of these leaf-scars that species of the *Lepidodendræ* have been established."

"The fructifications, rarely found attached to their support, are in cylindrical or ovate spikes, sessile or pedicellate, composed of sporanges attached to the anterior base of leaves or blades of various forms, which, curved upward and imbricated, cover the outside of the cones. The sporanges contain organisms of two kinds, either very small ones (microspores), which are like powder, or agglutinated globules of matter, distinct only with microscopes of great power. They may represent the male fertilizing pollen. Or, and more generally, they contain macrospores, large, true globular seeds, angular on one side, and rounded on the other."

The class may be divided into three orders, as follows:

1. ORDER, LEPIDODENDRÆ.

Acanthophyton (?), *Cyclostigma*, *Dechenia*, *Diplostegium*, *Glyptodendron*, *Halonia*, *Knorria*, *Lepidocystis*, *Lepidodendron*, *Lepidop. æum*, *Lepidophlojos*, *Lepidophyllum*, *Lepidostrobus*, *Leptophloeum*, *Lycopodites*, *Plumalina*, *Psilophyton*, *Sporangites*, *Sporocystis*, *Ulodendron*.

2. ORDER, TÆNIOPHYLLÆ.

Tæniophyllum.

3. ORDER, SIGILLARIÆ.

Didymophyllum, *Pinnularia*, *Sigillaria*, *Sigillarioides*, *Sigillariostrobus*, *Spirangium*, *Stigmara*, *Syringodendron*.

4. ORDER, NOEGGERATHIÆ.

Noeggerathia, *Whittleseya*.

The *Cordaites*, an extinct class, are represented in the Coal Measures, generally by fragments of ribbon-like leaves, and most rarely by stems bearing leaves and flowers. They belong to the *Gymnosperms*, and occupy a position somewhat intermediate between the *Noeggerathie* and *Coniferæ*. The genera are as follows:

Antholithes, *Asterocarpus*, *Cardiocarpon*, *Carpolithes*, *Cordiaanthus*, *Cordaicarpus*, *Cordaistrobus*, *Cordaites*, *Desmiophyllum*, *Dicranophyllum*, *Lepidoxylon*, *Rhabdocarpus*, *Trigonocarpum*.

FRUIT OF UNCERTAIN AFFINITY.

Gulielmites.

The *Coniferæ* are exogenous evergreen trees and shrubs, with branching trunks containing a resinous juice. They have a strobile cone or solitary seed. Three Palæozoic genera have been referred to the *Coniferæ*: viz., *Dadoxylon*, *Saportea*, and *Walchia*, but there must be doubt about the reference of *Dadoxylon* to this Class.

WOOD OF UNCERTAIN AFFINITY.

Celluloxylon, *Nematoxylon*, *Ormoxyton*, *Prototaxites*, *Sternbergia*, *Syringoxylon*.

ACANTHOPHYTON, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 324. [Ety. *akartha*, thorn; *phyton*, plant.] Cylindrical branches, ramifying in alternate manner, striated, with scattered tubercles, on which are borne short spines. Type *A. spinosum*.

spinosum, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 324, Chemung Gr.

ALETHOPTERIS, Sternberg, 1825, Vers. Darst. Flora der Vorwelt, p. 21. [Ety. *alethos*, true; *ptēris*, fern.] Fronds polypinnate; pinnules coriaceous, simple, mostly entire, enlarged at the base, cuneate or free, borders reflexed; midrib distinct, immersed into the epidermis, marked by a groove on the upper surface; prominent on the lower; lateral veins simple or forking once, open, often in right angle to the rachis; fructifications marginal. Type *A. lonchitica*.

acuta, see *Pecopteris acuta*.

ambigua, Lesquereux, 1880, Coal Flora of Pa., p. 182, Coal Meas.

aquilina, Schlotheim, 1820, (Filicites *aquilinus*.) Petrefaktenkunde, p. 405, and Coal Flora of Pa., p. 181, Coal Meas.

bunburyi, Andrews, 1875, Ohio Pal., vol. 2, p. 421, Coal Meas.

coxana, Lesquereux, 1861, Geo. Sur. Ky., vol. 4, p. 433, Coal Meas.

crassa, Lesquereux, 1884, Coal Flora of Pa., p. 748, Coal Meas.

crenulata, Brongniart, as identified by Lesquereux, in Geo. Sur. Ill., vol. 2, p. 439, is *Pseudopecopteris subcrenulata*.

cristata, see *Pecopteris cristata*.

discrepans, Dawson, 1862, Jour. Geo. Soc., vol. 18, p. 222, Devonian.

distans, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 865, is a variety of *A. lonchitica*.

emarginata, see *Pecopteris emarginata*.

erosa, see *Pecopteris erosa*.

evansi, Lesquereux, 1884, Coal Flora of Pa., p. 834, Coal Meas.

falcata, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 396, Coal Meas.

gibsoni, Lesquereux, 1880, Coal Flora of Pa., p. 183, Coal Meas.

grandifolia, Newberry, 1873, Ohio Pal., vol. 1, p. 384, Coal Meas.

grandis, Dawson, 1863, Can. Nat. & Geol., vol. 8, and Acad. Geol. p. 484, Coal Meas.

halleri, see *Pecopteris halli*.

helenæ, Lesquereux, 1880, Coal Flora of Pa., p. 179, Coal Meas.

heterophylla, Lindley & Hutton, 1833, (*Pecopteris heterophylla*.) Foss. Flora, vol. 1, p. 113, Coal Meas.

holdeni, see *Protoblechnum holdeni*.

hymenophylloides, see *Pseudopecopteris hymenophylloides*.

inflata, see *Callipteridium inflatum*.

ingens, Dawson, 1868, Acad. Geol. p. 553, Devonian.

lævis, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 865, Coal Meas.

lanceolata, see *Pecopteris lanceolata*.

lonchitica, Schlotheim, 1820, (Filicites *lonchiticus*.) Nachtrage zur Petrefaktenkunde, p. 411, and Coal Flora of Pa., p. 177, Coal Meas.

longifolia, see *Pecopteris longifolia*.

macrophylla, see *Danæites Macrophyllum*.

massilionis, see *Callipteridium massilionum*.

FIG. 5.—*Alethopteris lonchitica*.

maxim
p. 42
mazona
nana
muricat
nervosa,
obscura,
vol.
rugos
owen, s
pectina
Ill., v
pennsylv
Sur. I
perleyi,
Devon
pluckene
eneti.
preciosa,
pteroides,
robusta,
Pa., p.
rugosa, s
serlii, Br
Hist. o
Flora o
serrula, s
serrulata,
sheferi, s
solida, see
spinulosa,
losa.
stellata, se
tenioptero
oides.
urophylla
urophyll
Meas.
virginiana
or Up. C
Permian
ANARTHROCA
Voy. [c
canna, s
more or
ribs flat
of formi
perryana,
Soc. vol.
Dev. an
Catskill
ANEIMITES, D
Soc. vol.
a genus.
or attach
flabellate
closely re
acadicus, D
Soc., vol.
Low. Co
bockshii,
bockshii,
Foss. pla
Can. p. 4
obtusus, Le
obtus, a)
and Foss.
of Can., p
validus, Dav

maxima, Andrews, 1875, Ohio Pal., vol. 2, p. 421, Coal Meas.

mazonana, see *Pseudoplectopteris mazonana*.

muricata, see *Pseudoplectopteris muricata*.

nervosa, see *Pseudoplectopteris nervosa*.

obscura, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 865, syn. for *Callipteridium rugosum*.

oweni, see *Callipteridium oweni*.

pectinata, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 469, Coal Meas.

pennsylvanica, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 864, Coal Meas.

perleyi, Hart, 1868, Acad. Geol. p. 554, Devonian.

pluckeneti, see *Pseudoplectopteris pluckeneti*.

preciosa, see *Pecopteris preciosa*.

pteroides, see *Pecopteris pteroides*.

robusta, Lesquereux, 1884, Coal Flora of Pa., p. 835, Coal Meas.

rugosa, see *Callipteridium rugosum*.

serlii, Brongniart, 1828, (*Pecopteris serlii*.) Hist. d. Veg. Foss. p. 292, and Coal Flora of Pa., p. 176, Coal Meas.

serrula, see *Pecopteris serrula*.

serrulata, see *Pecopteris serrulata*.

sheaferi, see *Pseudoplectopteris sheaferi*.

solida, see *Pecopteris solida*.

spinulosa, see *Pseudoplectopteris spinulosa*.

stellata, see *Pecopteris stellata*.

teniopteroides, see *Pecopteris teniopteroides*.

urophylla, Brongniart, 1828, (*Pecopteris urophylla*.) Hist. d. Veg. Foss. Coal Meas.

virginiana, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 88, Coal Meas. or Permian.

ANARTHROCANNA, Göppert, 1845, in Tchih. Voy. [Ety. an, without; arthron joint; canna, a plant.] Cylindrical stems, more or less swelling at the nodes, with ribs flattened and continuous instead of forming joints as in *Calamites*.

perryana, Dawson, 1863, Quar. Jour. Geo. Soc. vol. 19, p. 461, and Foss. plants of Dev. and Up. Sil. formations, p. 27, Catskill Gr.

ANEIMITES, Dawson, 1861, Quar. Jour. Geo. Soc. vol. 17, p. 5. [Ety. from *Anemia*, a genus.] Pinnules clustered, petiolate or attached by a narrow base, with flabellate venation. Type *A. acadicus*, closely related to *Cyclopteris*.

acadicus, Dawson, 1861, Quar. Jour. Geo. Soc., vol. 17, p. 5, and vol. 21, p. 153, Low. Coal Meas.

bockshii, Göppert, 1836, (*Adiantites bockshii*.) Syst. Filic. Foss. p. 384, and Foss. plants of Dev. and Up. Sil. of Can. p. 46, Chemung Gr.

obtusus, Lesquereux, 1858, (*Noeggerathia obtusa*.) Geo. Sur. Pa., vol. 2, p. 854, and Foss. plants of Dev. and Up. Sil. of Can., p. 46, Catskill Gr.

validus, Dawson, 1862, (*Cyclopteris valida*.)

Quar. Jour. Geo. Soc., vol. 18, p. 319, and Foss. plants of Dev. and Up. Sil. of Can., p. 46, Ham. Gr.

FIG. 6.—*Aneimites obtusus*.

ANNULARIA, Sternberg, 1820, Essai d'un exposé Geognostico-botanique d. l. Flore. du monde primitif. 2d Cahier, p. 36. [Ety. annulus, a ring.] Stem articulate, striate, with a strong diaphragm traversing it at the articulations; branches opposite, nearly in right angles from the articulations; leaves verticillate, lanceolate, spatulate, or lingulate, abruptly or gradually acuminate, or obtuse, even emarginate at the apex; fructifications in long cylindrical spikes, with close articulations, and narrowly lanceolate bracts, bearing round sporanges in the axils of the leaves, or double, oval ones, pedicellate and attached in the middle of the internodes. Type *A. spinulosa*. This name is preoccupied in the subkingdom Mollusca, and Wood in 1860 proposed to substitute *Trochophyllum*; but *Trochophyllum* was preoccupied for a genus of corals in 1851, by Edwards and Haine.

acuminata, see *Sporangites acuminatus*.

antiqua, Dawson, 1861, Can. Nat. and Geol. vol. 6, p. 170, Devonian.

calamitoidea, Schimper, 1869, Pal. Veget., vol. 1, p. 349, and Coal Flora of Pa., p. 48, Coal Meas.

clavata, Lesquereux, 1880, (*Trochophyllum clavatum*.) Coal Flora of Pa., p. 65, Coal Meas.

cuspidata, Lesquereux, 1884, Coal Flora of Pa., p. 725, Subcarboniferous.

dawsoni, Schimper, 1869, Palæontologie Vegetale, vol. 1, p. 350, and Coal Flora of Pa., p. 51, Devonian. Proposed for *Asterophyllites latifolius*, of Dawson, because that name was preoccupied; but I have retained *Asterophyllites latifolius* because it is doubtful whether it is an *Annularia*.

ersi.) Quar. Jour. Geo. Soc., vol. 19, p. 463, and Coal Flora of Pa., pp. 307, 776, Catskill Gr.
sphenophyllifolia Lesquereux, 1884, Coal Flora of Pa., p. 775, Catskill Gr.
stricta, Andrews, 1875, Ohio Pal. vol. 2, p. 418, Coal Meas.

FIG. 10.—*Archæopteris stricta*.

ramosum var. *germanum*, Miller and Dyer, 1878, Cont. to Pal. No. 2, p. 4, Hud. Riv. Gr. Probably inorganic.

ARTHRARIA, Billings, 1874,

Pal. Foss., vol. 2, p. 66.

[Ety. *arthron*, a joint.]

Cylindrical stems with an

expansion at each end in

the form of a dumb-bell.

Type A. *antiquata*.

antiquata, Billings, 1874,

Pal. Foss., vol. 2, p. 66,

Upper Taconic.

biclavata, S. A. Miller, 1875, Cin. Quar.

Jour. Sci., vol. 2, p. 354, Hud. Riv. Gr.

ARTHROPHYCUS, Hall, 1852, Pal. N. Y., vol.

2, p. 4. [Ety. *arthron*, joint; *phykos*, sea-

plant.] Simple or branching, rounded

or subangular, flexuous, transversely

ridged or furrowed. Type A. *harlani*.

FIG. 12.—*Arthropycus harlani*.

harlani, Conrad, 1833, (Fucoides *harlani*.)

Ann. Rep. N. Y., p. 113, and Pal. N. Y.,

vol. 2, p. 5, Medina sandstone.

montalto, Simpson, 1888, Dict. Foss.,

found in Pa. Medina (?) Gr.

ARTHROSTIGMA, Dawson, 1871, Foss. Plants

Canada, p. 41. [Ety. *arthron*, joint;

Aristophycus, Miller and Dyer, 1878,

Cont. to Pal.

No. 2, p. 3.

Probably inorganic, and, if

fucoidal, too irregular and too

little known to be retained as a

genus.

ramosum, Miller and Dyer, 1878,

Cont. to Pal.

No. 2, p. 4, Hud.

Riv. Gr. Probably inorganic.

Probably inorganic.

FIG. 11.
Artharia antiquata.

stigma, a dot or puncture.] Stems elongated, cylindrical, bifurcating, and giving off lateral branches; irregularly furrowed longitudinally, with circular leaf scars arranged in whorls, and bearing linear rigid leaves with circular bases, structure apparently cellular, with a slender vascular axis. Type A. *gracile*.

gracile, Dawson, 1871, Foss. Plants Can., p. 41, Devonian.

Artisia, Sternberg, syn. for *Sternbergia*.

transversa, see *Sternbergia transversa*.

Asolanus, Wood, 1860, syn. for *Sigillaria*.

camptotenia, syn. for *Sigillaria monostigma*.

manephleus, a doubtful species of *Sigillaria*.

ornithicoides, see *Sigillaria ornithicoides*.

Asplenites, Göppert, 1836, Systema Filicum

Fossilium. [Ety. *Asplenium*, a genus of

ferns.]

elegans, see *Eremopteris elegans*.

ruber, Lesquereux, 1858, Geo. Sur. Pa.,

vol. 2, p. 864, Coal Meas. This species

does not seem to be recognized by

Lesquereux in his later work.

ASTEROCARPUS, Göppert, 1836, Syst. Fil.

Foss., p. 188. [Ety. *aster*, star; *karpos*,

fruit.] Fructification on lanceolate pin-

nules, marked by large star-like sori.

Type A. *sternbergi*.

grandis, Lesquereux, 1870, Geo. Sur. Ill.,

vol. 4, p. 469, Coal Meas.

sternbergi, Göppert, 1836, Syst. Filic.

Foss., p. 188, Coal Meas.

FIG. 13.—*Asterophycus simplex*.

ASTEROPHYCUS, Lesquereux, 1876, 7th Ann.

Rep. Geol. Sur. Ind., p. 139. [Ety.

aster, star; *phykos*, a sea-weed.] Stem

short, cylindrical; frond expanded and

divided star-like from the top of the

central axis; segments flattened or in-

flated. Type A. *coxi*.

coxi, Lesquereux, 1876, 7th Ann. Rep.

Geol. Sur. Ind., p. 139, Low. and Up.

Coal Meas.

simplex, Lesquereux, 1880, Coal Flora of

Pa., p. 13, Coal Meas.

ASTEROPHYLLITES, Brongniart, 1822, Mem.

du Mus. t. 8, p. 203. [Ety. *aster*, star;

phyllon, leaf; *lithos*, stone.] Stems articulate; branches opposite; central axis hollow or solid; leaves verticillate, free to the base, linear, acuminate, simple nerved; fructifications in elongated ears, bearing round sporanges in the axils of the leaves. Type *A. equisetiformis*.
acicularis, Dawson, 1862, Quar. Jour. Geol. Soc., vol. 18, p. 310, Devonian.
anthracinus, Heer, 1877, Fl. Foss. Helv., vol. 4, p. 50, and Coal Flora of Pa., p. 36, Coal Meas.
apertus, see *Macrostachya aperta*.
brardi, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 159, Coal Meas.
crassicaulis, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 851, Coal Meas.
curtus, see *Bechera curta*.
equisetiformis, Schlotheim, 1804, (Casuarinites *equisetiformis*.) Beitrag Zur Flora der Vorwelt, tab. 1, fig. 1, and Coal Flora of Pa., p. 35, Coal Meas.
erectifolius, Andrews, 1875, Ohio Pal., vol. 2, p. 425, Coal Meas.
fasciculatus, Lesquereux, 1880, Coal Flora of Pa., p. 41, Coal Meas.

FIG. 14.—*Asterophyllites foliosus*.

foliosus, Lindley & Hutton, 1833, Foss. Flora, vol. 1, p. 77, and Coal Flora of Pa., p. 38, Coal Meas.
gracilis, Lesquereux, 1860, Geo. Sur. Ark., vol. 2, p. 310, Coal Meas.
grandis, see *Bechera grandis*.
lanceolatus, see *Macrostachya lanceolata*.
latifolius, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 311, Devonian. The same form was called by Schimper *Annularia dawsoni*.
laxus, Dawson, 1868, Acad. Geol., p. 539, Devonian.
lentus, Dawson, 1871, Foss. Plants Can., p. 29, Devonian.
longifolius, Sternberg, 1823, (Bruckmannia *longifolia*.) Vers. Darst. Flora der Vorwelt fasc. 4, p. 58, Coal Meas.
minutus, Andrews, 1875, Ohio Pal., vol. 2, p. 424, Coal Meas.
ovalis, see *Calamostachys ovalis*.
parvulus, Dawson, 1861, Can. Nat. and Geo., vol. 6, p. 168, and Acad. Geol. p. 539, Chemung Gr.
radiatus, see *Annularia radiata*.

rigidus, Sternberg, 1824, (Bruckmannia *rigida*.) Vers. Darst. Flora der Vorwelt, p. 29, and Coal Flora of Pa., p. 37, Coal Meas.
scutigerus, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 311, Devonian.
stachioides, Wood, 1860, (Lepidostrobus *stachioides*.) Proc. Acad. Nat. Sci. Phil., vol. 12, p. 240, Coal Meas.
sublaevis, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 851, Coal Meas.
trinervis, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol. p. 479, Coal Meas.
tuberculatus, see *Annularia tuberculata*.

ASTEROPTERIS, Dawson, 1881, Quar. Jour. Geo. Soc., vol. 37, p. 299. [Ety. *aster*, star; *pteria*, fern.] Stems of ferns having the axial portion composed of vertical radiating plates of scalariform tissue imbedded in parenchyma, and having the outer cylinder composed of elongated cells traversed by leaf-bundles similar to those of *Zygopteris*. Type *A. novoboracensis*.
novoboracensis, Dawson, 1881, Quar. Jour. Geo. Soc., vol. 37, p. 299, Portage Gr.
ASTROPOLITHON, Dawson, 1888, Geo. Hist. Plants, p. 31. A peculiar impression, supposed by Prof. Dawson to be tucoidal. Judging from the illustration, I would refer it to the *Graptolida*. Type *A. hindii*.
hindii, Dawson, 1888, Geo. Hist. plants, p. 31, Utaconic.
BAIERA, Fr. Braun, 1840, Die Petrefakten d. Naturalien Samml. [Ety. proper name.] Leaves petiolate, flabelliform, dichotomous, many parted; nerves in each lacinia, several, dichotomous, and proceeding parallel with each other; leaf substance leathery. Type *B. teniata*.
virginiana, Fontaine & White, 1880, Perm or Up. Carb. Flora, p. 103, Coal Meas. or Permian.

FIG. 16.—*Baiera virginiana*. FIG. 15.—*Astropolithon hindii*.

BECHERA,
 der Vo
 Like
 leaves
 joints
 stems,
 grandis,
 Flora
 Coal F
 tenuis, F
 series,
 BEINERTIA,
 p. 273.
 guished
 flexuous
 ing, an
 tative
 gymno
 goeperti
 and Ac
Bergeria ma
 ginatur
Bergeria
 rhombi
BLASTOPHY
 Soc. Na
 bud; p
 with a
 junction

FIG. 17.—*Blastophyllum diadematum*.

ing, thi
 thick; l
 lanceola
 inornata, l
 natus), (c
 p. 310, G
 radiata, B
 diatus),
 Coal Flora

FIG. 18.—*Baiera teniata*.

transitionis
 transition
 116, and
 p. 309, H

BECHERA, Sternberg, 1824, Vers. Darst. Flora der Vorwelt, p. 30. [Ety. proper name.] Like Asterophyllites in its verticillate leaves, but distinguished by its tumid joints and deeply and widely furrowed stems. Type *B. grandis*.

grandis, Sternberg, 1824, Vers. Darst. Flora der Vorwelt, fasc. 4, p. 30, and Coal Flora of Pa., p. 41. Coal Meas.

tenuis, Bunbury, 1846, Am. Jour. Sci., 2d series, vol. 2, p. 232, Coal Meas.

BEINERTIA, Göppert, 1836, Syst. Filic. Foss. p. 273. [Ety. proper name.] Distinguished from Pecopteris by the treble flexuous nerves; sometimes anastomosing, and may have its actual representative in the Gymnogramme. Type *B. gymnogrammoides*.

göpperti, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol. p. 485, Coal Meas.

Bergeria marginata, see *Lepidodendron marginatum*.

Bergeria rhombica, see *Lepidodendron rhombicum*.

BLASTOPHYCUS, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 24. [Ety. *blastos*, bud; *phukos*, sea-weed.] Plant bilobate with a button-like protuberance at the junction. Type *B. diadematum*.

diadematum, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 24, Utica Slate Gr.

BORNIA, F. A. Roemer, 1854, Palæontographica, vol. 3. [Ety. proper name.] Stems cylindrical, articulate and furrowed as in *Calamites*; articulations scarcely contracted; ribs cut square or obtuse at the articulations, continuous, not alternating, thinly striate; cortical cylinder thick; leaves verticillate, free, linear-lanceolate. Type *B. radiata*.

inornata, Dawson, 1862, (Calamites *inornatus*), Quar. Jour. Geo. Soc., vol. 18, p. 310, Genesee Slate.

radiata, Brongniart, 1828, (Calamites *radiatus*), Hist. d. Veg. Foss., p. 122, and Coal Flora of Pa., p. 30, Subconglomerate.

FIG. 18.—*Bornia transitionis*.

transitionis, Göppert, 1852, (Calamites *transitionis*), Foss. Fl. d. Uebergsg., p. 116, and Quar. Jour. Geo. Soc., vol. 18, p. 309, Ham. Gr.

Bothrodendron punctatum, see *Ulodendron punctatum*.

Brachyphyllum obtusum, see *Lepidocystis obtusum*.

Bruckmannia longifolia, see *Asterophyllites longifolius*.

rigida, see *Asterophyllites rigidus*.

tuberculata, see *Annularia tuberculata*.

BYTHOTREPHIS, Hall, 1847, Pal. N. Y., vol. 1, p. 8. [Ety. *buthos*, depth of the sea; *trephos*, to grow.] Stems subcylindrical or compressed; branches numerous, divaricating, sometimes leaf-like. Type *B. antiquata*.

antiquata, Hall, 1847, Pal. N. Y., vol. 1, p. 8, Calcif. Gr.

asteroides, Fitch, 1849, Trans. Ag. Soc., and Emmons Am. Geol., p. 101, Upper Taconic.

cæspitosa, Hall, 1850, 3d Rep. N. Y. St., Mus. Nat. Hist., p. 178, Trenton Gr.

flexuosa, Emmons, 1844, (Fucoides *flexuosa*), Taconic system, p. 69, Upper Taconic.

FIG. 19.—*Bythotrephes ramulosa*, showing the ends, and branches as they appear on a nodule.

gracilis, Hall, 1843, Geo. Rep., 4th Dist., N. Y., p. 69, and Pal. N. Y., vol. 1, p. 62, Trenton to Clinton Gr.

gracilis var. *crassa*, Hall, 1852, Pal. N. Y., vol. 2, p. 19, Clinton Gr.

gracilis var. *intermedia*, Hall, 1852, Pal. N. Y., vol. 2, p. 19, Trenton to Clinton Gr.

granti, Dawson, 1888, Geo. Hist. of Plants, p. 37, Clinton Gr.

gregaria, Ringueberg, 1888, Proc. Acad. Nat. Sci. Phil., p. 131, Niagara Gr.

impudica, Hall, 1852, Pal. N. Y., vol. 2, p. 20, Clinton Gr.

lesquereuxi, Grote & Pitt, 1876, Bull. Buff. Soc. Nat. Hist., vol. 3, p. 88, Waterlime Gr.

palmeta, Hall, 1852, Pal. N. Y., vol. 2, p. 20, Clinton Gr.

ramosa, Hall, 1852, Pal. N. Y., vol. 2, p. 21, Clinton Gr.
ramulosa, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 235, Utica Slate Gr.
rigida, Emmons, 1844, (Fucoides rigidus,) Taconic System, p. 69, Upper Taconic.
subnodosa, Hall, 1847, Pal. N. Y., vol. 1, p. 262, Hnd. Riv. Gr.
succulens, Hall, 1847, Pal. N. Y., vol. 1, p. 62, Trenton Gr.
tenuis, Hall, 1852, Pal. N. Y., vol. 2, p. 18, Trenton Gr. The Trenton form of *B. gracilis*.

CALAMITES, Guettard, 1751, Mem. Ac. Sci., Paris. [Ety. *calamus*, a reed.] Plants arborescent; trunks cylindrical, articulate; articulations variable in distance, rapidly closer toward the narrowed obconical base; surface narrowly ribbed lengthwise; ribs equal, simple, parallel, contracted or rounded at the articulations; branches nearly at right angles, verticillate like the leaves, which are lanceolate, acuminate, simple nerved. Type: *C. suckovi*.

approximatus, Sternberg, 1820, Essai d'un exposé Geognostico-botanique d. l. Fl. d. Monde primitif 2d Cahier, p. 3, and Coal Flora of Pa. p. 26, Coal Meas.

betriatus, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 850, Coal Meas. This name was preoccupied by Sternberg.

canniformis, Schlotheim, 1820, Petrefactenkunde, p. 398, and Coal Flora of Pa., p. 24, Coal Meas.

cisti, Brongniart, 1828, Hist. d. Veg. Foss. p. 129, and Coal Flora of Pa., p. 27, Coal Meas.

cruciatus, Brongniart, 1828, Hist. d. Veg. Foss. t. 1, p. 128, Coal Meas.

disjunctus, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 850, Coal Meas.

dubius, Artis, 1825, Antedil. Phytology, pl. 13, and Coal Flora of Pa., p. 27, Coal Meas.

gigas, Brongniart, 1828, Hist. d. Veg. Foss., 1, p. 136, and Coal Flora of Pa., p. 25, Coal Meas.

gracilis, Lesquereux, 1861, Geo. Sur. Ky., vol. 4, p. 436, Coal Meas.

inornatus, see *Bornia inornata*.

major, Weiss, 1872, Fossil Flora d. jungsten Steinkolen formation, p. 119, and Coal Flora of Pa., p. 21, Coal Meas.

nodosus, Sternberg, 1820, Essai d'un Exp. Geog. Bot. d. l. Fl. d. Monde primitif 2d Cahier, p. 36, Coal Meas.

nova-scoticus, Dawson, 1863, Can. Nat. & Geol., vol. 8, and Acad. Geol. p. 479, Coal Meas.

pachyderma, Brongniart, 1828, Hist. d. Veg. Foss., 1, p. 132, and Coal Flora of Pa., p. 28, Coal Meas.

radiatus, see *Bornia radiata*.

ramifer, Stur, 1875, Culm Flora d. Mährisch-Schlesischen Dachschiefers, p. 82, and Coal Flora of Pa., pp. 23, 703, Coal Meas.

ramosus, Artis, 1825, Antedil. Phytology, pl. 2, and Coal Flora of Pa., pp. 22, 702, Coal Meas.

suckovi, Brongniart, 1828, Hist. d. Veg. Foss., t. 1, p. 124, and Coal Flora of Pa., p. 20, Coal Meas.
transitionis, see *Bornia transitionis*.

undulatus, Brongniart, 1828, Hist. d. Veg. Foss. 1, p. 127, Coal Meas.

voltzi, Brongniart, 1828, Hist. d. Veg. Foss. 1, p. 135, and Acad. Geol. p. 194, Coal Meas.

Calamocladus, Schimper, 1869, Pal. Veget., vol. 1, p. 423. Not clearly distinguished from *Asterophyllites* and founded upon *A. longifolius* as the type, and including *A. equisetiformis*, *A. foliosus*, *A. rigidus*, and *Bechera grandis*.

CALAMODENDRON, Brongniart, 1828, Hist. d. Veg. Foss. vol. 1, p. 133. [Ety. *calamus*, reed; *dendron* tree.] Central cylinder striate lengthwise and articulate, surrounded by a thick, woody cylinder or bark, with outside surface smooth. The structure is allied to *Sigillaria*, but the appearance is like *Calamites*. Type *C. approximatum*.

antiquum, Dawson, 1871, Foss. Plants Canada, p. 24, Devonian.

approximatum, Brongniart, 1828, Hist. d. Veg. Foss., vol. 1, p. 133, Coal Meas.

obscurum, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol. p. 476, Coal Meas.

tenuistriatum, Dawson, 1871, Foss. Plants Canada, p. 25, Devonian.

CALAMOPHYCUS, Lesquereux, 1877, Proc. Am. Phil. Soc. p. 165. [Ety. *calamus*, reed; *phukos*, sea-plant.] Fronds simple, elongated, gradually tapering to a point; cavity divided by transverse membranes, either passing through the whole diameter, or connected in the middle to vertical subdivisions. Type *C. septum*.

septum, Lesquereux, 1877, Proc. Am. Phil. Soc. p. 165, Low. Held. Gr.

CALANOSTACHYS, Schimper, 1869, Traité de Paléontologie Végétale, vol. 1, p. 328. [Ety. *calamus*, reed; *stachys*, plant.] Spikes doubtfully considered as fructifications of *Asterophyllites*. Type *C. typicus*.

brevifolius, Lesquereux, 1884, Coal Flora of Pa., p. 718, Coal Meas.

lanceolatus, Lesquereux, 1884, Coal Flora of Pa., p. 715, Subconglomerate.

FIG. 20.—*Calamites* Suckovi.

FIG. 21.—*Calamodendron* approximatum.

ovalis, L.
 ovalis,
 Flora
 praelongu
 CALLIPTER
 jungste
 from
 large,
 to the
 decurre
 the m
 at the
 solved
 obliqu
 ders, d
 to the
 aldrichi,
 Pa., p.
 journa
 dourna
 and Co
 dawson
 Perm.
 Meas. c
 grandifo
 Perm. c
 Meas. o

FIG. 22.

grandini,
 grandini
 and Coal
 inflatum,
 inflata,
 Coal Me
 inequale,
 Pa., p. 16
 mansfieldi,
 of Pa., p.
 massillone
 opteris m
 2, p. 438,
 membran
 Flora of
 neuropter
 Flora of
 oblongifoli
 Perm. or
 Meas. or
 odontopter
 Perm. or
 Meas. or

ovalis, Lesquereux, 1858, (*Asterophyllites ovalis*), Geo. of Pa., p. 851, and Coal Flora of Pa., p. 717, Coal Meas.

prælongus, see *Volkmania prælonga*.

CALLIPTERIDIUM, Weiss, 1872, Foss. Flora d. jungsten Steinkohlen formation. [Ety. from the genus *Callipteris*.] Fronds large, polypinnate; pinnules attached to the rachis by the whole base, often decurrent, and the lower descending to the main rachis, connate or disjointed at the base; primary nerve strong, dissolved below the apex; lateral veins oblique, curved in passing to the borders, dichotomous, the basilar attached to the rachis. Type *C. sullivanti*.

aldrichi, Lesquereux, 1880, Coal Flora of Pa., p. 171, Coal Meas.

dournaisi, Brongniart, 1828, (*Pecopteris dournaisii*), Hist. d. Veg. Foss., p. 282, and Coal Flora of Pa., p. 747, Coal Meas.

dawsonianum, Fontaine & White, 1880, Perm. or Up. Carb. Flora., p. 56, Coal Meas. or Permian.

grandifolium, Fontaine & White, 1880, Perm. or Up. Carb. Flora., p. 58, Coal Meas. or Permian.

FIG. 22.—*Callipteridium sullivanti*.

grandini, Brongniart, 1828, (*Pecopteris grandini*), Hist. d. Veg. Foss., p. 286, and Coal Flora of Pa., p. 748, Coal Meas.

inflatum, Lesquereux, 1870, (*Alethopteris inflata*), Geo. Sur. Ill., vol. 4, p. 393, Coal Meas.

inequale, Lesquereux, 1880, Coal Flora of Pa., p. 168, Coal Meas.

mansfieldi, Lesquereux, 1880, Coal Flora of Pa., p. 166, Coal Meas.

massilloneum, Lesquereux, 1866, (*Alethopteris massillonis*), Geo. Sur. Ill., vol. 2, p. 438, Low. Coal Meas.

membranaceum, Lesquereux, 1880, Coal Flora of Pa., p. 172, Coal Meas.

neuropteroides, Lesquereux, 1880, Coal Flora of Pa., p. 166, Coal Meas.

oblongifolium, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 56, Coal Meas. or Permian.

odontopteroides, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 59, Coal Meas. or Permian.

oweni, Lesquereux, 1860, (*Alethopteris oweni*), Geo. Ref. of Arkansas, vol. 2, p. 309, Coal Meas.

pardeeii, Lesquereux, 1880, Coal Flora of Pa., p. 169, Coal Meas.

rigidum, Lesquereux, 1884, Coal Flora of Pa., p. 746, Coal Meas.

rugosum, Lesquereux, 1858, (*Alethopteris rugosa*), Catal. Potts, Ass'n, p. 11, and Coal Flora of Pa., p. 169, Coal Meas.

sinuatum, Brongniart, 1828, (*Pecopteris sinuata*), Hist. d. Veg. Foss., p. 296, and Coal Flora of Pa., p. 745, Coal Meas.

sullivanti, Lesquereux, 1854, (*Callipteris sullivanti*), Bost. Jour. Nat. Hist., vol. 6, p. 423, and Geo. Sur. Pa., vol. 2, p. 866, Coal Meas.

unitum, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 60, Coal Meas.

CALLIPTERIS, Brongniart, 1828, Hist. d. Veg. Foss., p. 249. [Ety. *kallos*, beautiful; *pterus*, fern.] Fronds polypinnate; pinnules sessile and sometimes occurring on the principal rachis, thick; parenchyma dense, nerves immersed, showing creases in the leaf-substance, simple or forking once. Type *C. conferta*.

conferta, Sternberg, 1824, (*Sphenopteris conferta*), Vers. Darst. Flor. d. Vorwelt and Perm. or Up. Carb. Flora of Pa., p. 54, Coal Meas. or Permian.

sullivanti, see *Callipteridium sullivanti*.

CARDIOCARFON, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 87. [Ety. *kardia*, heart; *karpos*, fruit.] Seeds of various forms, composed of a compressed, generally cordiform or oval nucleus, surrounded by a flattened, fibrous border, or a membranaceous wing. Type *C. majum*.

affine, Lesquereux, 1860, Geo. of Ark., vol. 2, p. 311, Coal Meas.

annulare, Sternberg, 1824, (*Carpolithes annularis*), Vers. Darst. Flor. d. Vorwelt and Coal Flora of Pa., p. 814, Subconglomerate.

annulatum, Newberry, 1873, Ohio Pal., vol. 1, p. 374, Coal Meas.

apiculatum, Göppert & Berger, 1848, De fructibus et seminibus, p. 23, and Coal Flora of Pa., p. 571, Subconglomerate.

baileyi, Dawson, 1868, Acad. Geol., p. 554, Devonian.

bicornutum, Lesquereux, 1870, (*Ptilocarpus bicornutus*), Geo. Sur. Ill., vol. 4, p. 443, Coal Meas.

bicuspidatum, Sternberg, 1820, (*Carpolithes bicuspidatus*), Flora der Vorwelt, and Coal Flora of Pa., p. 573, Coal Meas.

bisectum, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 491, Coal Meas.

circulare, Lesquereux, 1884, Coal Flora of Pa., p. 812, Coal Meas.

conglobatum, Lesquereux, 1884, Coal Flora of Pa., p. 810, Coal Meas.

congruens, Grand Eury, 1877, Flore Carbonifère, p. 236, and Coal Flora of Pa., p. 573, Coal Meas.

cornutum, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 324, Devonian.
 crampi, Hartt, 1868, Acad. Geol., p. 554, Devonian.
 crassum, Lesquereux, 1884, Coal Flora of Pa., p. 812, Coal Meas.
 dilatatum, Lesquereux, 1884, Coal Flora of Pa., p. 806, Subcarboniferous.
 diminutivum, Lesquereux, 1880, Coal Flora of Pa., p. 570, Coal Meas.
 diplotesta, Lesquereux, 1884, Coal Flora of Pa., p. 812, Coal Meas.
 divergens, Lesquereux, 1884, Coal Flora of Pa., p. 811, Coal Meas.
 ellipticum, Sternberg, 1820, (Carpolithes ellipticus), Flor. d. Vorw., p. 40, and Coal Flora of Pa., p. 814, Coal Meas.
 elongatum, Newberry, 1873, Ohio Pal., vol. 1, p. 324, Coal Meas.
 fasciculatum, Lesquereux, 1880, Coal Flora of Pa., p. 570, Coal Meas.
 fruticosum, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 491, Coal Meas.
 harveyi, Lesquereux, 1884, 13th Rep. Ind. Geol., p. 102, and Coal Flora of Pa., p. 808, Coal Meas.
 ingens, Lesquereux, 1860, Geo. of Ark., vol. 2, p. 311, Coal Meas.
 late-alatum, Lesquereux, 1880, Coal Flora of Pa., p. 568, Coal Meas.
 lator, Lesquereux, 1884, Coal Meas. of Pa., p. 811, Coal Meas.
 latum, Newberry, 1873, Ohio Pal., vol. 1, p. 372, Coal Meas.
 lescurianum, n. sp. Coal Meas. Proposed instead of *C. ovale* Lesquereux, in Coal Flora of Pa., p. 810, which name was preoccupied.
 longicollis, Lesquereux, 1884, Coal Flora of Pa., p. 808, Coal Meas.
 mamillatum, see *Rhabdocarpus mamillatus*.
 marginatum, Artis, 1828, Antedil. Phytol., pl. 22, Coal Meas.
 minus, Newberry, 1873, Ohio Pal., vol. 1, p. 372, Coal Meas.
 newberryi, Andrews, 1875, Ohio Pal., vol. 2, p. 425, Coal Meas.
 obliquum, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 324, Devonian.
 orbiculare, Newberry, 1853, Ann. of Sci., vol. 1, p. 374, Coal Meas.
 ovale, Dawson, 1871, Foss. Plants Can., p. 60, Devonian.
 ovale, Lesquereux, 1884, Coal Flora of Pa., p. 810, Coal Meas. The name was preoccupied. See *C. lescurianum*.
 ovatum, Grand'Eury, 1877, Flore Carbonifere, p. 236, Coal Meas.
 pachyteta, Lesquereux, 1880, Coal Flora of Pa., p. 565, Coal Meas.
 patens, Lesquereux, 1884, Coal Flora of Pa., p. 807, Coal Meas.
 picatum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 876, Coal Meas.
 punctatum, Gœppert, 1836, Syst. Filic. Foss., p. 24, and Coal Flora of Pa., 597, Coal Meas.
 pusillum, Lesquereux, 1884, Coal Flora of Pa., p. 815, Coal Meas.

regulare, Sternberg, 1820, (Carpolithes regularis), Flor. d. Vorw., and Coal Flora of Pa., p. 572, Coal Meas.
 retusum, Sternberg, 1820, (Carpolithes retusum), Flora der Vorwelt, and Ohio Pal., vol. 1, p. 374, Coal Meas.

samariforme, Newberry, 1873, Ohio Pal., vol. 1, p. 375, Coal Meas.

simplex, Lesquereux, 1880, Coal Flora of Pa., p. 569, Coal Meas.

speciosus, Lesquereux, 1884, Coal

Flora of Pa., p. 807, Coal Meas.
 tenellum, Dawson, 1873, Rep. Foss. Plants, p. 28, Subcarboniferous.

trevortoni, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 876, Coal Meas.

zonulatum, Lesquereux, 1880, Coal Flora of Pa., p. 568, Coal Meas.

CARDIOPTERIS, Schimper, 1869, Traité de Paléontologie Végétale, vol. 1, p. 457. [Ety. *kardia*, heart; *pteris*, fern.] Leaves simple, pinnate; stipe striated, rounded, base spoon-like, dilated; pinnae perpendicular, opposite close, imbricated, cordato-ovate, leathery, margins reflexed; primary nerves numerous, equal, dichotomous. Type *C. polymorpha*.

eriana, Dawson, 1881, Quar. Jour. Geo. Soc. Lond., vol. 37, p. 305, Devonian.

CARPOLITHES, Schlotheim, 1820, Petrefactenkunde. [Ety. *karpos*, fruit; *lithos*, stone.] Seeds of uncertain relation: not referable by their characters to other genera.
 acuminatus, Sternberg, 1821, Flor. d. Vorw. and Coal Flora of Pa., p. 596, Coal Meas.

bicarpus, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 98, Coal Meas. or Permian.

bicuspidatus, see *Cardiocarpon bicuspidatum*.

bifidus, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 877, Coal Meas.

bullatus, see *Lepidocystis bullatus*.

butlerianus, Lesquereux, 1884, Coal Flora of Pa., p. 824, Coal Meas.

cerasiformis, Sternberg, Flor. d. Vorw., vol. 2, p. 208, and Coal Flora of Pa., p. 824, Coal Meas.

cistula, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 451, Coal Meas.

clavatus, see *Rhabdocarpus clavatus*.

compactus, Dawson, 1871, Foss. Plants Canada, p. 63, Devonian.

conicus, Lesquereux, 1884, Coal Flora of Pa., p. 824, Coal Meas.

corticosus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 462, Coal Meas.

FIG. 23.—*Cardiocarpon samariforme*.

CAS.—CH
disjunct
 Pa.,
 pon
 fascicul
 Ill., v
 'ragaria
 vol. 1
 granula
 Vorw
 Coal
 jackson
 lator, I
 p. 826
 lunatus
 Soc.,
 marginat
 or Up
 was p
 C. wh
 minimu
 welt a
 Meas.
 multistri
 striat
 perpusil
 of Pa.
 persicari
 Ill., vo
 platimarg
 margin
 retusum, s
 siliqua,
 Soc. L
 spicatus,
 Soc. L
 transect
 of Pa.,
 trilocular
 umbonat
 Flora
 venosus, s
 vesiculari
 whitanu
 Propos
 taine d
 Carb.
 preocc
 Casuarinites
 equiset
 Caulerpites,
 d. Ve
 stem;
 mose, d
 form of
 Type C
 marginat
 CAULOPTER
 Flora,
 pteris,
 disk ei
 mains d
 to the
 impress
 shape,
 without
 vascula
 been, in
 of the
 Type C

- disjunctus*, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 877, syn. for *Trigonocarpon dawesi*.
- fasciculatus*, Lesquereux, 1860, Geo. Sur. Ill., vol. 2, p. 457, Coal Meas.
- fragarioides*, Newberry, 1873, Ohio Pal., vol. 1, p. 370, Coal Meas.
- granularis*, Sternberg, 1820, Flora der Vorwelt, and Coal Flora of Pa. p. 825, Coal Meas.
- jacksonensis*, see *Rhabdocarpus jacksonensis*.
- laticorpus*, Lesquereux, 1884, Coal Flora of Pa., p. 826, Coal Meas.
- lunatus*, Dawson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 464, Devonian.
- marginatus*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 98. The name was preoccupied by Artis in 1825. See *C. whitianus*.
- minimus*, Sternberg, 1820, Flora der Vorwelt and Coal Flora of Pa., p. 825, Coal Meas.
- multistriatus*, see *Rhabdocarpus multistriatus*.
- perpusillus*, Lesquereux, 1884, Coal Flora of Pa., p. 825, Coal Meas.
- persicaria*, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 462, Coal Meas.
- platinmarginatus*, see *Rhabdocarpus platinmarginatus*.
- retusus*, see *Cardiocarpon retusum*.
- siliqua*, Dawson, 1863, Quar. Jour. Geo. Soc. Lond., vol. 19, p. 465, Devonian.
- spicatus*, Dawson, 1863, Quar. Jour. Geo. Soc. Lond., vol. 19, p. 461, Devonian.
- transsectus*, Lesquereux, 1884, Coal Flora of Pa., p. 826, Coal Meas.
- trilocularis*, see *Trigonocarpum triloculare*.
- umbonatus*, Sternberg, 1820, Vers. Darst. Flora der Vorwelt, Coal Meas.
- venosus*, see *Rhabdocarpus venosus*.
- vesicularis*, see *Lepidocystis vesicularis*.
- whitianus* n. sp. Coal Meas. or Perm. Proposed instead of *C. marginatus*, Fontaine & White, 1880, in Perm. or Up. Carb. Flora, p. 98, which name was preoccupied.
- Casuarinites equisetiformis*, see *Asterophyllites equisetiformis*.
- Caulerpites*, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 21. [Ety. *kaulos*, stem; *erpo*, creep.] Stem simple or ramose, covered with short branches, in form of leaves doubled or imbricated. Type *C. lycopodioides*.
- marginatus*, see *Taonurus marginatus*.
- CAULOPTERIS**, Lindley & Hutton, 1833, Foss. Flora, vol. 1, p. 121. [Ety. *kaulos*, stem; *pteria*, a fern.] Scars with the inside disk either marked by linear bands, remains of vessels passing from the trunk to the base of the rachis, or covered by impressions of rootlets obliterating its shape, or merely ovate or elliptical, without traces of horseshoe-shaped vascular lines. These lines may have been, in some cases, effaced by abrasion of the surface or covered by rootlets. Type *C. primæva*.

acanthophora, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 458, Coal Meas. It may be a synonym for

Ulodendron punctatum.

antiqua, Newberry, 1871, Quar. Jour. Geo. Soc., vol. 27, p. 271, Up. Held. Gr.

cisti, Brongniart, 1828, (Sigillaria cisti,) Hist. Veg. Foss. p. 418, and Coal Flora of Pa., p. 345, Coal Meas.

elliptica, Fontaine &

White, 1880, Perm. or Up. Carb. Flora, p. 95, Coal Meas. or Permian.

giffordi, Lesquereux, 1880, Coal Flora of Pa., p. 343, Coal Meas.

gigantea, see *Stemmatopteris gigantea*.

gigantea, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 95, Coal Meas. or Permian.

insignis, see *Stemmatopteris insignis*.

intermedia, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 459, Coal Meas.

lacoii, Lesquereux, 1880, Coal Flora of Pa., p. 344, Coal Meas.

lockwoodi, Dawson, 1871, Quar. Jour. Geo. Soc., vol. 27, p. 270, Chemung Gr.

mansfieldi, Lesquereux, 1880, Coal Flora of Pa., p. 346, Coal Meas.

obtecta, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 457, Coal Meas.

peregrina, Newberry, 1871, Quar. Jour. Geo. Soc., vol. 27, p. 272, Up. Held. Gr.

punctata, see *Stemmatopteris punctata*.

wortheni, see *Stemmatopteris wortheni*.

CELLULOXYLON, Dawson, 1881, Lond. Quar. Jour. Geo. Soc., vol. 37, p. 302. [Ety.

cellula, a small apartment; *xylo*, wood.] Trunk showing in cross section, large and somewhat unequal cells disposed in narrow concentric bands, between wider bands of fine fibrous tissue; no medullary rays; longitudinal section shows either cells superimposed in vertical rows, or a sort of banded prosenchymatous tissue. The structure appears to have been of exogenous growth. Type *C. primævum*.

primævum, Dawson, 1881, Quar. Jour. Geo. Soc., vol. 37, p. 302, Ham. Gr. Penhallow says this is an *Alga*, and belongs to the genus *Nematophycus*.

Chloephycus, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 3. Probably inorganic.

plumosum, Miller & Dyer, Cont. to Pal., No. 2, p. 3, Hud. Riv. Gr. Probably inorganic.

FIG. 24.—*Caulopteris primæva*.

CHONDRITES, Sternberg, 1833, Vers. Darst. Flora der Vorwelt, p. 25. [Ety. from its resemblance to *Chondrus crispus*, or Irish moss.] Fronds cartilaginous, filiform or robust stems, dichotomous, branched; rounded or compressed. Type *C. antiquus*.

antiquus, Brongniart, 1828, (*Fucoides antiquus*.) Hist. d. Veg. Foss., vol. 1, p. 63, Devonian.

colletti, see *Taonurus colletti*.

targioni, Brongniart, 1828, (*Fucoides targioni*.) Hist. d. Veg. Foss., t. 1, p. 56, Coal Meas.

CONOSTICHUS, Lesquereux, 1876, 7th Ann. Rep. Geo. Sur. Ind., p. 142. [Ety. *konos*, cone; *stichos*, row.] Stipe cylindrical, continuous; frond enlarging from the base upward in the shape of a plate, or of a cup, or increasing by successive superposed layers or concentric laminae; top concave, cup-shaped. Type *C. ornatus*.

broadheadi, Lesquereux, 1880, Coal Flora of Pa., p. 15, Coal Meas.

FIG. 25.—*Conostichus ornatus*.

ornatus, Lesquereux, 1876, 7th Ann. Rep. Geo. Sur. Ind., p. 142, Coal Meas. *prolifer*, Lesquereux, 1880, Coal Flora of Pa., p. 16, Coal Meas.

CORDAIAANTHUS, Grand'Eury, 1877, Flore Carbonifere, p. 228. [Ety. *Cordaites*, a genus; *anthos*, flower.] Flowers and fruits of Cordaites, found isolated or in fragments where their relation to stem-bearing leaves is unknown. A provisional name only. Type *C. gemmifer*.

bracteatus, Lesquereux, 1870, (Schultz's *bracteata*.) Geo. Sur. Ill., vol. 4, p. 427, Coal Meas.

dichotomus, Lesquereux, 1880, Coal Flora of Pa., p. 546, Coal Meas.

ebracteatus, Lesquereux, 1884, Coal Flora of Pa., p. 844, Coal Meas.

flexuosus, Lesquereux, 1884, Coal Flora of Pa., p. 802, Coal Meas.

gemmifer, Grand'Eury, 1877, Flore Carbonifere, p. 228, and Coal Flora of Pa., p. 545, Coal Meas.

ovatus, Lesquereux, 1880, Coal Flora of Pa., p. 545, Coal Meas.

rugosus, Lesquereux, 1884, Coal Flora of Pa., p. 803, Coal Meas.

scaber, Lesquereux, 1884, Coal Flora of Pa., p. 844, Coal Meas.

simplex, Lesquereux, 1880, Coal Flora of Pa., p. 538, Coal Meas.

spicatus, Lesquereux, 1884, Coal Flora of Pa., p. 802, Coal Meas.

CORDAIOCARPUS, Grand'Eury, 1877, Flore Carbonifere, p. 236. [Ety. *Cordaites*, a genus; *karpos*, fruit.] Seeds of variable size and shape. Type *C. ovatus*.

apiculatus, Lesquereux, 1880, Coal Flora of Pa., p. 551, Coal Meas.

cinctus, Lesquereux, 1884, Coal Flora of Pa., p. 804, Coal Meas.

costatus, Lesquereux, 1880, (Cordaites *costatus*.) Coal Flora of Pa., p. 540, Coal Meas.

FIG. 26.

guthieri, Grand'Eury, 1877, Flore Carbonifere, p. 236, Coal Meas.

lineatus, Lesquereux, 1884, Coal Flora of Pa., p. 805, Coal Meas.

ovatus, Grand'Eury, 1877, Flore Carbonifere, p. 236, and Coal Flora of Pa., p. 550, Coal Meas.

stabilis, Lesquereux, 1884, Coal Flora of Pa., p. 805, Coal Meas.

CORDAISTROBUS, Lesquereux, 1880, Coal Flora of Pa., p. 551. [Ety. *Cordaites*, a genus; *strobilus*, cone.] Strobile cylindrical, tapering to a blunt acumen, covered by transversely rhomboidal scars placed in spiral, bearing narrow, linear leaves, with the characters, form, and nervation of leaves of Cordaites.

Type *C. grandeuryi*.

grandeuryi, Lesquereux, 1880, Coal Flora of Pa., p. 552, Coal Meas.

CORDAITES, Unger, 1850, Gen. et sp., p. 277. [Ety. proper name.] Trunks of large size, irregularly branching, formed of a large medullar canal or pith; marked on the outer surface by transverse, narrow, parallel, simple ribs, rarely joined by divisions, covered by double or triple layers of wood and bark, converted by fossilization into thin layers of coal; leaves in spiral order, more or less distant, ribbon-like, of various length and width, linear, or more generally gradually enlarging upward, obtuse, entire or undulate, and split at the apex; borders curving to the sessile, or semi-embracing, somewhat inflated base; surface marked lengthwise by primary and secondary parallel simple nerves, generally more distant in the middle of the leaves, and slightly inflated toward the base; flowers in racemes from the axils of the leaves; fruits generally oval, sessile, of various size. Type *C. borassifolius*.

angustifolius, Dawson, 1861, Can. Nat., vol. 6, p. 10, Ham. Gr.

angustifolius, Lesquereux, see *C. diversifolius*.

borassifolius, Sternberg, 1820, (Flabellaria *borassifolia*.) Essai d. Exp. Geogn.-botan. d. l. Flora d. monde primitif, 2d Cabier, p. 36, and Coal Flora of Pa., p. 532, Coal Meas.

commu
Phil.
p. 534
costatus

FIG. 27.—C

Coal F
diversif
of Pa.
Instead
which
flexuosu
Soc., v
gracilis,
Phil. S
p. 539,
grandifol
Phil. S
Pa., p.
lacoel, L
Pa., p.
lingulatu
bonifer
p. 533,
mansfield
Phil. S
p. 537,
radiatus,
Pa., p.
robbl, D
p. 8, H
serpens, l
Pa., p.
simplex, l
and Acc
validus,
Phil. So
p. 523, C

CREMATOPTER

Paleont
[Ety. *Crema*
Rachis
oblong,
auricula
pennsylv
Flora of

CRUZIANA,

l'Amer.
proper
kled fu
Type C.
linnaeoni
Foss., p
Merid.,

communis, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 320, and Coal Flora of Pa., p. 534, Coal Meas.

costatus, Lesquereux, 1878, Proc. Am.

FIG. 27.—*Cordaites costatus*.

crassinervis, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 97, Coal Meas. or Perm.

crassus, Lesquereux, 1880, Coal Flora of Pa., p. 530, Coal Meas.

diversifolius, Lesquereux, 1880, Coal Flora of Pa., p. 535, Coal Meas. Proposed instead of *C. angustifolius* Lesquereux, which was preoccupied.

flexuosus, Dawson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 462, Catskill Gr.

gracilis, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 322, and Coal Flora of Pa., p. 539, Coal Meas.

grandifolius, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 318, and Coal Flora of Pa., p. 530, Coal Meas.

lanceolatus, Lesquereux, 1880, Coal Flora of Pa., p. 535, Coal Meas.

lingulatus, Grand'Eury, 1877, Flore Carbonifère, p. 218, and Coal Flora of Pa., p. 533, Coal Meas.

mansfieldi, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 321, and Coal Flora of Pa., p. 537, Coal Meas.

radiatus, Lesquereux, 1880, Coal Flora of Pa., p. 540, Coal Meas.

robbei, Dawson, 1861, Can. Nat., vol. 6, p. 8, Ham. Gr.

serpens, Lesquereux, 1880, Coal Flora of Pa., p. 324, Coal Meas.

simplex, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 490, Coal Meas.

validus, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 317, and Coal Flora of Pa., p. 523, Coal Meas.

CREMATOPTERIS, Schimper, 1869, Traité de Paléontologie Végétale, vol. 1, p. 596. [Ety. *krematos*, hanging; *pterus*, fern.]

Rachis thick; pinnules sessile, ovate-oblong, contracted at the base, and subauriculate. Type *C. typica*.

pennsylvanica, Lesquereux, 1880, Coal Flora of Pa., p. 307, Coal Meas.

CRUZIANA, D'Orbigny, 1842, Voy. dans l'Amer. Merid. t. 3, pt. 2, p. 30. [Ety. proper name.] A transversely wrinkled fucoid, much like *Rusophycus*. Type *C. rugosa*.

linnarsoni, White, 1874, Rep. Invert. Foss., p. 5, and Geo. Sur. W. 100th Merid., vol. 4, p. 32, Upper Taconic.

rustica, White, 1874, Rep. Invert. Foss., p. 5, and Geo. Sur. W. 100th Merid., vol. 4, p. 32, Up. Taconic.

similis, Billings, 1874, Pal. Foss., vol. 2, p. 68, Up. Taconic.

CYCLOPTERIS, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 51. [Ety. *kuklos*, circle; *pterus*, fern.] Leaflets orbicular or reniform, large, veins numerous, and not positively referable to other genera. Type *C. orbicularis*.

acadica, see *Anelmities acadicus*.

alleghaniensis, Meek, 1876, Desc. Foss. Plants Va. Syn. for *Archæopteris rogersi*.

antiqua, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 481, Coal Meas.

bockshii, see *Anelmities bockshii*.

browni, see *Rhacophyllum browni*.

crispata, Germ. & Kaulf, 1831, (Filicites crispus,) Nova. Acta. Acad., vol. 15, p. 229, Coal Meas.

elegans, Lesquereux, 1858, Zost. Jour. Nat. Hist., vol. 6, p. 416, and Geo. Sur. Pa., vol. 2, p. 856, Coal Meas.

FIG. 28.—*Cyclopteris elegans*.

fimbriata, see *Neuropteris fimbriata*.

flabellata, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 52, Coal Meas.

germari, see *Neuropteris germari*.

hallana, see *Archæopteris hallana*.

hispida, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 481, Coal Meas.

hirsuta, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 856, Coal Meas.

incerta, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 320, Ham. Gr.

jacksoni, see *Archæopteris jacksoni*.

laciniata, see *Neuropteris laciniata*.

lescuriana, see *Triphyllopteris lescuriana*.

oblata, Lindley & Hutton, 1837, Foss. Flora, vol. 3, pl. 217, Coal Meas.

obliqua, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 52, Coal Meas.

orbicularis, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 52, Coal Meas.

problematica, Dawson, 1871, Foss. Plants Dev. and Up. Sil., p. 47, Devonian.

rogersi, see *Archæopteris rogersi*.

trichomanoides, see *Neuropteris trichomanoides*.

undans, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 855, Coal Meas.

valida, see *Aneimites validus*.
varia, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 319, Devonian.
virginiana, see *Pseudopteris virginiana*.
wilsoni, Wood, 1860, Proc. Acad. Nat. Sci., p. 519, Coal Meas.

CYCLOSTIGMA, Houghton, 1860, Ann. and Mag. Nat. Hist., 3d ser., vol. 5, p. 444. [Ety. *kuklos*, circle; *stigma*, a dot or puncture.] Steins arborescent, surface tuberculate, rugose lengthwise; tubercles in regular spiral order, small, subglobose, more generally conical, acute, topped with a vascular terminal and prominent point, or more rarely flattened at the top into small, round areoles, with the vascular point in the middle; decorticated surface smooth or obscurely striate lengthwise by the series of tubercles, which are oval, elevated or prominent, and gradually effaced downward or decurring, preserving the impressions of the central vascular scars. Type *C. kiltorkense*.
affine, Dawson, 1881, Quar. Jour. Geo. Soc., vol. 37, p. 301, Chemung Gr.
densifolium, Dawson, 1871, Foss. Plants Can., p. 43, Devonian.

kiltorkense, Houghton, 1860, Ann. and Mag. Nat. Hist., 3d ser., vol. 5, p. 444, Subcarboniferous.

CYMOGLOSSA, Schimper, 1869, Traité de Paléontologie Végétale, vol. 1, p. 553. [Ety. *kumo*, wavy; *glossa*, tongue.] Frond pinnate, or bipinnate; pinnae oblong, sessile, alternate, margin lobed; nerves simple or branching, reaching the margin and leaving triangular spaces without nerves. Type *C. goeppertana*.

breviloba, Fontaine & White, 1880, Perm. and Up. Carb. Flora, p. 86, Coal Meas. or Permian.

formosa, Fontaine & White, 1880, Perm. and Up. Carb. Flora, p. 86, Coal Meas. or Permian.

lobata, Fontaine & White, 1880, Perm. and Up. Carb. Flora, p. 87, Coal Meas. or Permian.

FIG. 29.—
Dactylo-
phycus
tridigi-
tatur.

obtusifolia, Fontaine & White, 1880, Perm. and Up. Carb. Flora, p. 85, Coal Meas. or Permian.

DACTYLOPHYCUS, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 1. [Ety. *dactylos*, finger; *phycus*, sea-plant.] Stem divided at one or both ends into three or more short subequal branches, and closely related to *Ichnophycus*. Type *D. tridigitatus*.

quadrupartitum, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 2, Utica Slate Gr.
tridigitatum, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 1, Utica Slate Gr.

DADOXYLON, Endlicher, 1840, Syn. Con. [Sig. pine or torch-wood.] Branching trunks, with distinct zones of growth and a pith of Sternbergia type; wood-cells, with rows of areoles with oval pores; medullary rays with series of cells.

acadianum, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 473, Coal Meas.

annulatum, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol. p. 473, Coal Meas.

antiquum, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol. p. 473, Coal Meas.

clarkii, Dawson, 1882, Foss. Plants Erian and Up. Sil. Formations, pt. 2, p. 124, Genesee shales.

halli, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 306, Ham. Gr.

materialium, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 473, Up. Coal Meas.

newberryi, Dawson, 1871, Foss. Plants Can., p. 14, Portage Gr.

ouangondianum, Dawson, 1861, Can. Nat., vol. 6, and Acad. Geol. p. 534, Up. Devonian.

DANEITES, Göppert, 1836, Syst. Filic. Foss., p. 380. [Ety. from the genus *Danea*.] Fronds pinnate; secondary veins coming out in right angles from the primary straight nerve, simple or dichotomous; sporanges, on the lower side of the lamina, placed in rows from the medial nerve to near the borders along the lateral veins, oval or linear exannulate. Type *D. asplenoides*.

asplenoides, var. *major*, Bunbury, 1846, Quar. Jour. Geo. Soc., vol. 2, p. 85, Coal Meas.

emersoni, Lesquereux, 1880, Coal Flora of Pa., p. 157, Coal Meas.

macrophyllus, Newberry, 1873, (Alethopteris macrophylla.) Ohio Pal., vol. 1, p. 383, Low. Coal Meas.

DECHENIA, Göppert, 1841, Die Gattungen der fossilen Pflanzen, p. 43. [Ety. proper name.] Stems arborescent;

leaf-scars in continuous spiral lines; bolsters oblong, rounded, marked by obscure concentric striae, on the middle

FIG. 30.—Dadoxylon. a, bark; b, woody zone or fiber (pleurochyma); c, medulla or pith; d, cast of hollow pith or sternbergia.

FIG. 31.—
Dactylo-
phycus
macrophyllus.

of w
ably
idea.
striata,
Pa.,
DENROPI
Flora
dron,
comp
ments
spread
rhizon
thick;
large,
when
branch
one sl
sions
Type
desori, L
Pa., vo
DESMITOP
Flora
band;
leaves
larger
or join
late at
leaves
promin
buried
thick, i
ting of
gracile.
gracile, L
Pa., p. 5
DICRANOPHY
Carboni
pointed
leaves n
various
filaments
with a f
intermed
mersed i
dichotomu
of Pa.,
dimorphum
Am. Phil
Pa., p. 55
Dictyoites, see
Protozoa.
becki see Di
DICRYOPTERIS,
Schwarz
pteris, fer
nules co
at the ba
oblong, o
veins fle
and inter
less disti
of polygo
niarti.
cordata, Ro
Steinkohle
in Paleoe

of which were attached leaves, probably cylindrical. Type *D. euphorbioides*.

striata, Lesquereux, 1880, Coal Flora of Pa., p. 431, Coal Meas.

DENDROPHYCUS, Lesquereux, 1884, Coal Flora of Pa., vol. 3, p. 699. [Ety. *dendron*, tree; *phukos*, sea-weed.] Root composed of tubulose flattened filaments, irregularly branching and widely spreading from the base of the rhizoma; rhizoma cylindrical, simple, long and thick; fronds at first top-shaped, very large, tree-like, and many times divided when opened; primary and secondary branches thick and somewhat flat on one side, dichotomous; ultimate divisions cylindrical, narrow and pointed. Type *D. desori*.

desori, Lesquereux 1884, Coal Flora of Pa., vol. 3, p. 699, Devonian.

DESMIOPHYLLUM, Lesquereux, 1880, Coal Flora of Pa., p. 556. [Ety. *desmos*, band; *phyllon*, leaf.] Stems slender; leaves narrow, sublinear, gradually enlarged from the base, single and sparse or joined 3 or 4 together and fasciculate at the base; surface of stem and leaves irregularly ribbed lengthwise by prominent large bundles of nerves buried under the epidermis, which is thick, irregularly granulose, by splitting of the coaly layer. Type *D. gracile*.

gracile, Lesquereux, 1880, Coal Flora of Pa., p. 557, Coal Meas.

DICRANOPHYLLUM, Grand'Eury, 1877, Flore Carbonifère, pl. 30. [Ety. *dikranos*, two-pointed; *phyllon*, leaf.] Stems slender, leaves narrow, linear, subcoriaceous, of various length, forking, or dividing in filaments in the upper part, marked with a few thick primary nerves, and intermediate nervilles, more or less immersed into the epidermis.

dichotomum, Lesquereux, 1880, Coal Flora of Pa., p. 553, Coal Meas.

dimorphum, Lesquereux, 1879, Proc. Am. Phil. Soc., p. 329, and Coal Flora of Pa., p. 554, Coal Meas.

Dictyolites, see *Dictyophyton* in the class Protozoa.

becki see *Dictyophyton becki*.

DICTYOPTERIS, Gutbier, 1835, Verst. Zwisch. Schwarzk., p. 63. [Ety. *dictyon*, net; *pterus*, fern.] Frond bipinnate; pinules cordate, truncate or rounded at the base, sessile or short pedicelled, oblong, obtuse or lanceolate, entire; veins flexuous, connected by flexures and intersections, forming a more or less distinct and close reticulation of polygonal meshes. Type *D. bronniarti*.

cordata, Roemer, in Pflanzen d. prod. Steinkohleng. am Harz und Piesberg in Paleontographica, vol. 9, p. 186,

and Coal Flora of Pa., p. 833, Coal Meas.

neuropteroidea, Gutbier, 1852, Verst. Stein Sachs., p. 23, and Coal Flora of Pa., p. 833, Coal Meas.

obliqua, Bunbury, 1847, Quar. Jour. Geo. Soc., vol. 3, p. 427, and Coal Flora of Pa., p. 146, Coal Meas.

rubella, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 388, Coal Meas.

scheuchzleri, Hoffmann, in Roem. Pflanz. d. Kohlengeb. am Harz in Paleont. IX, p. 186, and Coal Flora of Pa., p. 832, Coal Meas.

DIDYMOPHYLLUM, Göppert, 1841, Gatt. der Foss. Pflanzen, p. 35. [Ety. *didymos*, double; *phyllon*, a leaf.] Trunk arborescent, cylindrical; leaves double, united at the base, disposed in spiral order, appressed; areoles prominent, reniform, each resembling a pair of small areoles attached to each other. Type *D. schottini*.

oweni, Lesquereux, 1870, (Sigillaria *oweni*.) Geo. Sur. Ill., vol. 4, p. 498, Coal Meas.

reniforme, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 309, Ham. Gr.

Diplazites emarginatus, see *Pecopteris emarginata*.

DIPLOSTEGIUM, Corda, 1845, Beiträge zur Flora der Vorwelt, p. 112. [Ety. *diplos*, double; *stega*, a covering; but spelled by Corda *Diplotegium*.] Thick trunks of trees longitudinally furrowed; bark thick, and imbricated in short cylindrical overlaps. Type *D. brownianum*.

FIG. 83
Diplotegium retusum.

brownianum, Corda, 1845, Beiträge zur Flora der Vorwelt, p. 112, Coal Meas.

retusum, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 490, Coal Meas.

truncatum, Lesquereux, 1860, Geo. Sur. Ark., vol. 2. Syn. for *Knorria imbricata*.

DISCOPHYCUS, Walcott, 1879, Trans. Alb. Inst., vol. 10, p. 19. [Ety. *diskos*, disk; *phukos*, sea-plant.] Frond discoid, slightly convex, and substance coriaceous. Type *D. typicale*.

typicale, Walcott, 1879, Trans. Alb. Inst., vol. 10, p. 19, Utica slate.

DYSTACTOPHYCUS, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 2. [Ety. *dustaktos*, hard to arrange; *phukos*, sea-plant.] Frond mammiiform, expanded and concentrically wrinkled. Type *D. mammillanum*.

FIG. 32.—*Dictyopteris obliqua*.

mammillanum, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 2, Hud. Riv. Gr.

FIG. 34.—*Dystactophycus mammillanum*.

EOPHYTON, Torell, 1868, Bidr. t. Sparagm. geogn. och. paleont., p. 36. [Ety. *eos*, dawn; *phyton*, a plant.] Slender, cylindrical, reed-like fucoids, longitudinally striated. Type *E. linnæanum*.

explanatum, Dawson, 1870, Can. Nat. and Geol., Low. Arenig rocks.

3. jukesii, Billings, 1874, Pal. Foss., vol. 2, p. 65, Up. Taconic.

linnæanum (?), Torell, 1868, Bidr. t. Sparagm. geogn. och. paleont., p. 36, Up. Taconic.

EQUISETITES, Sternberg, 1833, Vers. Darst. Flora der Vorwelt, vol. 2, p. 43. [Ety. *equus*, a horse; *seta*, a hair or bristle; in allusion to the resemblance to a horse-tail.] Arborescent; stems articulate; articulations surrounded with costate sheaths, dentate on the border. Type *E. gigantea*.

FIG. 35.—*Equisetites curtus*.

taine & White, 1880, Perm. or Up. Carb. Flora, p. 33, Coal Meas. or Permian.

gracilis, Lesquereux, 1884, Coal Flora of Pa., p. 729, Coal Meas.

macrodonatus, Wood, 1860, Proc. Acad. Nat. Sci. Not satisfactorily defined.

occidentalis, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 425, Coal Meas.

stellifolius, Harlan, 1835, (Equisetum stellifolium.) Trans. Geo. Soc. Pa., vol. 1, p. 261, Coal Meas. Syn. for *Annularia longifolia*?

striatus, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 34, Coal Meas. or Permian.

wrightianus, see *Echinocaris wrightiana*.

Equisetum, see *Equisetites*.

columnaris, Brongniart, 1828, (Equisetum columnare.) Hist. Veg. Foss., t. 1, p. 115, Coal Meas.

curtus, Dawson, 1863, Syn. Carb. Flora in Can. Nat., vol. 8, and Acad. Geol., p. 443, Coal Meas.

elongatus, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 33, Coal Meas. or Permian.

gracilis, Lesquereux, 1884, Coal Flora of Pa., p. 729, Coal Meas.

macrodonatus, Wood, 1860, Proc. Acad. Nat. Sci. Not satisfactorily defined.

occidentalis, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 425, Coal Meas.

stellifolius, Harlan, 1835, (Equisetum stellifolium.) Trans. Geo. Soc. Pa., vol. 1, p. 261, Coal Meas. Syn. for *Annularia longifolia*?

striatus, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 34, Coal Meas. or Permian.

wrightianus, see *Echinocaris wrightiana*.

Equisetum, see *Equisetites*.

columnare, see *Equisetites columnaris*.

stellifolium, see *Equisetites stellifolius*.

EREMOPTERIS, Schimper, 1869, Traité de Palæontologie Vegetale, vol. 1, p. 416. [Ety. *eremos*, isolated; *pterus*, fern.]

Upper part of fronds dichotomous; pinnae open or oblique, irregularly pinnatifid; laciniae long, obovate or wedge-form, the lower ones deeply cut; the lateral veins enter the lobes in acute angles of divergence from the midrib, and passing up to the borders are flabellate, dichotomous, parallel, and close. Type *E. artemisiifolia*.

artemisiifolia, Sternberg, 1824, (Sphenopteris artemisiifolia.) Vers. Darst. Flora der Vorwelt, p. 44, and Coal Flora of Pa., p. 293, Coal Meas.

cheathamii, Lesquereux, 1884, Coal Flora of Pa., p. 770, Coal Meas.

crenulata, Lesquereux, 1876, Geo. Rep. of Alabama, p. 75, and Coal Flora of Pa., p. 292, Coal Meas.

dissocata, Lesquereux, 1876, Geo. Rep. of Alabama, p. 75, and Coal Flora of Pa., p. 293, Coal Meas.

elegans, Ettingshausen, 1852, (Asplenites elegans.) Die Steinkohlen flora v. Stradonitz in Bohmen, p. 15, and Coal Flora of Pa., p. 294, Coal Meas.

flexuosa, Lesquereux, 1876, Geo. Rep. of Alabama, p. 75, and Coal Flora of Pa., p. 293, Coal Meas.

marginata, Andrews, 1875, Ohio Pal., vol. 2, p. 422, Coal Meas.

microphylla, Lesquereux, 1880, Coal Flora of Pa., p. 296, Coal Meas.

missouriensis, Lesquereux, 1880, Coal Flora of Pa., p. 295, Coal Meas.

Ficoidites scabrousus, Hildreth, 1837, Am. Jour. Sci. and Arts, vol. 31, p. 30, Low. Coal Meas. Not recognized, but probably a *Sigillaria*.

Filicites, Schlotheim, 1820, Nachtr. zur Pctref. It was used for all fossil ferns, and hence is not of generic value.

acuminatus, see *Neuropteris acuminata*.

aquilinus, see *Alethopteris aquilina*.

arborescens, see *Pecopteris arborescens*.

crispus, see *Cyclopteris crispa*.

gracilis, see *Plumalina gracilis*.

lonchiticus, see *Alethopteris lonchitica*.

miltoni, see *Pecopteris miltoni*.

penniformis, see *Pecopteris penniformis*.

plukenetii, see *Pseudopecopteris plukenetii*.

plumosa, see *Pecopteris plumosa*.

oreopteridis, see *Pecopteris oreopteridis*.

trifoliata, see *Pseudopecopteris trifoliata*.

Flabellaria borassifolia, see *Cordaites borassifolia*.

Fucoides, Brongniart, 1822, in Mem. d. Hist.

FIG. 36.—*Eremopteris marginata*.

Nat. a
It was
sites of
more
allegan
auriform
bilobatus
caudagal
demissa,
losa.
dentatus,
tus pr
filiciform
flexuosus,
gracilis,
graphica
mined
harlani,
heterophy
be deta
retort, see
rigidus, s
secalinus,
plex.
serra, Bro
noides
simplex, s
velum, see
verticalis,
Galium sp
sphenop
GLYPTODEND
Sci. and
[Ety. *gly*
Stem th
areoles,
depressed

FIG. 37.

eatonense,
and Arts
ara Gr.
Goniopteris ne
berryana
oblonga, see
Gordia martin
GUELDMITES,
Rothlieg.
19. [Ety.
kind of f
orbicularis,
Perm. an
Meas. or
permianus,
d. Rothlieg.
19, Permian

Nat. and Hist. d. Veg. Foss., t. 1., p. 50. It was used to comprehend the *Sargassites* or *Thalassophytes*, and hence is of more than generic value.

alleghaniensis, see *Arthropycus harlani*.

auriformis, Hall, 1843. Not organic.

bilobatus, see *Rusophycus bilobatus*.

caudagalli, see *Taonurus caudagalli*.

demissa, Conrad probably *phytopsis tubulosa*.

dentatus, Brongniart probably *Diplograptus pristiniiformis*.

filiciformis, see *Rhacophyllum filiciforme*.

flexuosus, see *Bythotrephis flexuosa*.

gracilis, see *Bythotrephis gracilis*.

graphica. Not defined so as to be determined.

harlani, see *Arthropycus harlani*.

heterophyllus, Hall. Not defined so as to be determined.

retort, see *Taonurus retortus*.

rigidus, see *Bythotrephis rigida*.

secalinus, Hall syn. for *Diplograptus simplex*.

serra, Brongniart, see *Graptolithus bryonoides*.

simplex, see *Diplograptus simplex*.

velum, see *Taonurus velum*.

verticalis, see *Scolithus verticalis*.

Gulium sphenophylloides, see *Annularia sphenophylloides*.

GLYPTODENDRON, Claypole, 1878, Am. Jour. Sci. and Arts, 3d ser., vol. 15, p. 302.

[Ety. *glyptos* sculptured; *dendron*, tree.]

Stem thick, covered with rhomboidal areoles, the lower portions of which are depressed. Type *G. eatonense*.

FIG. 37.—*Glyptodendron eatonense*.

eatonense, Claypole, 1878, Am. Jour. Sci. and Arts, 3d. ser., vol. 15, p. 302, Niagara Gr.

Goniopteris newberryana, see *Pecopteris newberryana*.

oblonga, see *Pecopteris oblonga*.

Gordia marina, see *Palaeochorda marina*.

GULIELMITES, Geinitz, 1858, Leithpflanzen d.

Rothleig. u. d. Zechstein, Sachsen, p.

19. [Ety. from the genus *Gulielma*.] A

kind of fruit, of uncertain affinity.

orbicularis, Fontaine & White, 1880, Perm. and Up. Carb. Flora, p. 99, Coal Meas. or Permian.

permanus, Geinitz, 1858, Leithpflanzen d. Rothleig. u. d. Zechstein, Sachsen, p.

19, Permian.

HALONIA, Lindley & Hutton, 1835, Foss.

Flora, vol. 2, p. 11. [Ety. from its close

affinity with *Halonia*.] Stems of me-

diu size, dichotomous; cortex tuber-

culate; spaces intermediate to the tu-

bercles marked with rhomboidal scars;

decorticated surface, covered with punc-

tiform round or oval papillæ, obtuse or

perforated in the center, placed in

spiral order. Type *H. tortuosa*.

flexuosa, Goldenberg, 1855, (*Ulodendron*

flexuosum,) Flora Sarræpontana fossilis,

vol. 1, pl. 2, fig. 10, and Coal Flora of

Pa., p. 416, Coal Meas.

mansfieldi.

Lesquereux,

1880, Coal

Flora of Pa.,

p. 414, Coal

Meas.

pulchella, Les-

quereux,

1860, Geo.

Sur. Ark.,

vol. 2, p. 311,

Coal Meas.

secreta, Les-

quereux,

1880, Coal

Flora of Pa.,

p. 417, Coal

Meas.

tortuosa, Lind-

ley & Hut-

ton, 1835, Foss.

Flora, vol. 2, p. 11, Coal

Meas.

tuberculata, Brongniart, 1838, Hist. d.

Veg. Foss., vol. 2. pl. 28, and Coal Flora

of Pa., p. 411, Coal Meas.

Harlania, syn. for *Arthropycus*.

halli, syn. for *Arthropycus harlani*.

FIG. 38.—*Halonia flexuosa*.

FIG. 39.—*Helophycus stelliforme*.

HELOPHYCUS, Miller & Dyer, 1878, Cont. to

Pal. No. 2, p. 2. [Ety. *helios*, the sun;

phukos, sea-plant.] Star-like frond, having five rays; transversely wrinkled. Type H. stelliforme.

stelliforme, Miller & Dyer, 1878, Cont. to Pal. No. 2, p. 2, Hud. Riv. Gr.

HYPONOPHYLLITES, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 203. [Ety. *hippodon*, horse-foot; *phukos*, sea-plant.] Founded upon cavities in sandstone, having a form similar to that which a putty ball will assume, when pressed between thumb and finger, leaving a rounded rim on three sides of the disc, the compressed margin being truncate. Type H. cowlesii.

cowlesii, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 203, Chemung Gr.

HYMENOPHYLLITES, Gœppert, 1836, Syst. Filic. Foss. [Ety. from the genus *Hymenophyllum*.] Frond membranaceous, many times regularly pinnately divided or irregularly cut, lobed with pinnatifid or dichotomous divisions, decurring on a broad common rachis, which is sometimes indistinct; veins pinnate, percurrent, solitary in each division. Type H. gerardii.

FIG. 40.
Hymenophyllum curtilobus.

adnascens, see *Rhacophyllum adnascens*.

alatus, see *Sphenopteris alata*.

arborescens, see *Rhacophyllum arborescens*.

ballantini, see *Sphenopteris ballantini*.

capillaris, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 863, Coal Meas.

clarki, see *Rhacophyllum clarki*.
curtilobus, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 321, and Acad. Geol., p. 552; Devonian.

delicatulus, Brongniart, 1828, Hist. d. veg. Foss., p. 185, Coal Meas.

flexicaulis, see *Sphenopteris flexicaulis*.

furcata, see *Sphenopteris furcata*.

gerardii, Gœppert, 1836, Syst. Filic. Foss. Devonian.

giganteus, see *Rhacophyllum lactuca*.

gutbieranus, Unger, 1850, Gen. et. sp., p. 132, Coal Meas.

hildrethi, see *Sphenopteris hildrethi*.

inflatus, see *Rhacophyllum inflatum*.

lactuca, see *Rhacophyllum lactuca*.

molle, see *Rhacophyllum molle*.

myriophyllus, Brongniart, 1828, (Sphenopteris myriophylla.) Hist. d. Veg. Foss., p. 184, Coal Meas.

obtusilobus, Gœppert, 1836, Syst. Filic. Foss., Devonian.

pentadactylus, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 485, Coal Meas.

pinnatifidus, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 436, Coal Meas.

schlotheimi, Brongniart, 1828, (Sphenopteris schlotheimi.) Hist. d. Veg. Foss., p. 193, Coal Meas.

spinosa, see *Sphenopteris spinosa*.

splendens, see *Sphenopteris splendens*.

strongi, see *Rhacophyllum strongi*.

subfurcatus, Dawson, 1868, Acad. Geol., p. 55, Devonian.

tenuifolius, Brongniart, 1828, (Sphenopteris tenuifolia.) Hist. d. Veg. Foss., p. 190, Coal Meas.

thalliformis, see *Rhacophyllum thalliforme*.
trichomanoides, see *Sphenopteris trichomanoides*.

tridactylites, see *Sphenopteris tridactylites*.

ICHNOPHYLLITES, Hall, 1852, Pal. N. Y., vol. 2, p. 56.

[Ety. *ichnos*, a footprint; *phukos*, a seaweed.] Tridactyle impressions somewhat resembling a foot-track, the middle stem being the longer. Type I. *ichnophycus tridactylum*.

FIG. 41.—*Ichnophycus tridactylum*.

tridactylum, Hall, 1852, Pal. N. Y., vol. 2, p. 26, Clinton Gr.

INDIOPHYLLUM, Lesquereux, 1880, Coal Flora of Pa., p. 159. [Ety. *idios*, peculiar; *phyllon*, leaf.] Leaves small, round, or broadly obovate; medial nerve thick, gradually narrowed and effacing in joining the borders; lateral secondary veins sub-opposite, thick, passing in an inside curve toward the borders, gradually effaced in the reticulation; venules more or less continuous; sometimes crossing each other in contrary directions, and forming, by intersections, quadrate or rhomboidal meshes. Type I. *rotundifolium*.

rotundifolium, Lesquereux, 1880, Coal Flora of Pa., p. 160, Coal Meas.

KNORRIA, Sternberg, 1825, Essai d. Exp. Geogn.-botan. d. l. Flor. du Monde primitif fasc. 4, p. 37. [Ety. proper name.]

Trunks covered with elongated semiconical or truncate tubercles placed in spiral order more or less imbricated, leaving, after falling off, round convex marks, with a single, vascular scar in the middle; leaves long, linear, more or less inflated at the base, with a flat medial nerve. Type K. *imbricata*.

compacta, Lesquereux, 1884, Coal Flora of Pa., p. 839, Coal Meas.

imbricata, Sternberg, 1825, Flor. d. monde primitif fasc. 4, p. 37, Kaskaskia Gr.

FIG. 42.
Knorrria imbricata.

selloni,
primi
taxina,
Flora
LEPIDOCY
Pa., p.
der.]
in rig
rachis
upon
spora
unkno
natus.
angulari
Pa., p.
bullatus
bullat
Coal M
fraxinifo
(Carp
et Sem
lineatus,
Pa., p.
obtusul
lum o
876, Co
pectinatu
of Pa.,
quadrang
Flora d
vesiculari
vesicul
Coal M
LEPIDODEN
expose
flore du
25. [E
Surface
points a
(bolster
bark of
variable
tion, oft
(inside
dotted l
bearing
gin two
bundles
on each
which, l
less di
wrinkles
Type L.
aculeatum
Geogn.-b
itif, 1st
Pa., p. 3
acuminatu
Ueberga
iferous.
andrewsi,
Pa., p. 3
altivolare, s
binerve, B
Soc., vol.
borda, W
p. 239. C
brittsi, Les
Pa., p. 36

- selloni, Sternberg, 1825, Flor. d. monde primitif fasc. 4, p. 37-50, Coal Meas.
 taxina, Lindley & Hutton, 1833-5, Foss. Flora, vol. 2, p. 37, Coal Meas.
- LEPIDOCYSTIS, Lesquereux, 1880, Coal Flora of Pa., p. 454. [Ety. *lepis*, scale; *kustia*, bladder.] Spore cases long, naked, attached in right angle and opposite to a broad rachis; or short, placed in spiral order upon long, flexuous axes; or isolated sporanges, detached from strobiles of unknown character. Type *L. pectinatus*.
- angularis, Lesquereux, 1880, Coal Flora of Pa., p. 456, Coal Meas.
 bullatus, Lesquereux, 1870, (Carpolithes bullatus,) Geo. Sur. Ill., vol. 4, p. 463, Coal Meas.
 fraxiniformis, Gœppert & Berger, 1848, (Carpolithes fraxiniformis,) De Fruct. et Sem., p. 26, Coal Meas.
 lineatus, Lesquereux, 1880, Coal Flora of Pa., p. 454, Coal Meas.
 obtusus, Lesquereux, 1858, (Brachyphyllum obtusum,) Geo. of Pa., vol. 2, p. 876, Coal Meas.
 pectinatus, Lesquereux, 1880, Coal Flora of Pa., p. 454, Coal Meas.
 quadrangularis, Lesquereux, 1880, Coal Flora of Pa., p. 455, Coal Meas.
 vesicularis, Lesquereux, 1870, (Carpolithes vesicularis,) Geo. Sur. Ill., vol. 4, p. 462, Coal Meas.
- LEPIDODENDRON, Sternberg, 1820, Essai d'un expose Geognostico-botanique de la flore du monde primitif, 1st Cahier, p. 25. [Ety. *lepis*, scale; *dendron*, tree.] Surface of the stem marked by scars, points of leaf attachments; leaf scars (bolsters) rhomboidal, oblong, upon the bark of large trees or small branches, variable in size according to their position, often disfigured; central cicatrices (inside scars) rhomboidal, transversely dotted by three points (vascular scars) bearing, generally, under the lower margin two oval small tubercles, scars of bundles of vessels (appendages) placed on each side of a medial line (cauda), which, like the appendages, is more or less distinct, sometimes deep and wrinkled across, sometimes obsolete. Type *L. dichotomum*.
- aculeatum, Sternberg, 1820, Essai d. Exp. Geogn.-botan. d. l. flor. d. monde primitif, 1st Cahier, p. 25, and Coal Flora of Pa., p. 371, Coal Meas.
 acuminatum, Gœppert, 1852, Foss. Fl. d. Uebergangsgebirge, p. 185, Subcarboniferous.
 andrewsi, Lesquereux, 1880, Coal Flora of Pa., p. 380, Coal Meas.
 alveolare, see *Sigillaria alveolaris*.
 binerve, Bunbury, 1847, Quar. Jour. Geo. Soc., vol. 3, p. 431, Coal Meas.
 bordie, Wood, 1860, Proc. Acad. Nat. Sci., p. 239, Coal Meas.
 brittsi, Lesquereux, 1880, Coal Flora of Pa., p. 368, Coal Meas.
- carinatum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 875, Coal Meas.
 chemungense, Hall, 1843, (Sigillaria chemungensis,) Geo. Rep. 4th Dist. N. Y., p. 275, Chemung Gr.
 chilalloeum, Syn. for *L. distans*.
 clypeatum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 875, Coal Meas.
 conicum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 874, Coal Meas.
 corrugatum, Dawson, 1860, Quar. Jour. Geo. Soc., vol. 15, p. 313, and Acad. Geol., p. 253, Waverly Gr.
 costatum, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 453, Kaskaskia Gr.
 crenatum, Sternberg, 1820, Flor. d. monde primitif, 1st Cahier, p. 25, and Coal Flora of Pa., p. 394, Coal Meas.
 cruciatum, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 432, Coal Meas.
 cuspidatum, Lesquereux, 1880, Coal Flora of Pa., p. 388, Coal Meas.
 cyclostigma, Lesquereux, 1880, Coal Flora of Pa., p. 394, Coal Meas.
 decurtatum, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 487, Coal Meas.
 dichotomum, Sternberg, 1820, Flor. d. monde primitif, 1st Cahier, p. 25, and Coal Flora of Pa., p. 384, Coal Meas.
 dikrocheilum, Wood, 1860, Proc. Acad. Nat. Sci., p. 239, Coal Meas.
 dilatatum, Lindley & Hutton, 1831, Foss. Flora, vol. 1, p. 27, Coal Meas.
 diplostegiodes, Lesquereux, 1860, Geo. Sur. Ark., vol. 2, p. 311, Coal Meas.
- distans, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 874, Coal Meas.
- drepanaspis, Wood, 1860, Proc. Acad. Nat. Sci., Phil., vol. 12, p. 240, Coal Meas.
- dubium, Wood, syn. for *L. rimosum*.
- elegans, Sternberg, 1824, (Lycopodiolithes elegans,) Vers. Darst. Flor. d. Vorwelt 4 fasc., p. 8, Coal Meas.
- forulatum, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 431, Coal Meas.
- gaspanum Dawson, 1859, Quar. Jour. Geo. Soc., vol. 15, p. 484, and Acad. Geol., p. 541, Catskill Gr. Probably the same as Vanuxem's *Sigillaria simplicitas*.
- giganteum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 874, Coal Meas.
- gracile, Lindley & Hutton, 1831, Foss. Flora, vol. 1, p. 30, Coal Meas.
- greeni, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 43, Coal Meas.
- harcourtii, Witham, 1832, Trans. Nat. Hist. Soc., New, upon Tyne, p. 51, Coal Meas.
- ichthyolepis, Wood, 1860, (Lepidophlois ichthyolepis,) Proc. Acad. Nat. Sci. Phil., p. 240, Coal Meas.

FIG. 43.—*Leptodendron distans*.

ingens, Wood, syn. for *L. aculeatum*.
lanceolatum, Lesquereux, 1880, Coal Flora of Pa., p. 369, Coal Meas.
latifolium, Lesquereux, 1880, Coal Flora of Pa., p. 370, Coal Meas.
lesquereuxi, Wood, syn. for *L. clypeatum*.
longifolium, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 85, and Coal Flora of Pa., p. 373, Coal Meas.
magnum, Wood, 1860, Proc. Acad. Nat. Sci., Phil., p. 239, Coal Meas.
mammillatum, Lesquereux, syn. for *L. veltheimianum*.
marginatum, Presl, 1826, (*Bergeria marginata*), in Sternberg Flor. d. Vorw., p. 134, and Coal Flora of Pa., p. 784, Coal Meas.
mekiston, Wood, syn. for *L. modulatum*.
mielcki, Goppert, 1836, Syst. Filic. Foss., p. 465, and Coal Flora of Pa., p. 395, Coal Meas.
modulatum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 874, Coal Meas.
morrisanum, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 430, Coal Meas.
obovatum Sternberg, 1820, Flor. d. monde primitif, 1st Cahier, p. 25, Coal Meas.
obscurum, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 453, Kaskaskia Gr.
obtusum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 875, Coal Meas.
oculatum, Lesquereux, syn. for *L. distans*.
ovum, Wood, syn. for *L. vestitum*.
personatum, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 488, Coal Meas.
pictoense, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol. p. 487, Coal Meas.
plicatum, Dawson, 1863, Can. Nat. and Geo. vol. 8, and Acad. Geol., p. 488, Coal Meas.
plumarium, Lindley & Hutton, 1835, Foss. Flora, vol. 3, p. 151, Coal Meas.
politum, syn. for *L. modulatum*.
primævum, Rogers, 1858, Geo. Sur. Pa., vol. 2, p. 675, Ham. Gr.
quadrangulatum, Schlotheim, 1820, (Pal-macites quadrangulatus,) Petrefacten-kunde, p. 395, and Coal Flora of Pa., p. 383, Coal Meas.
quadrilaterale, Lesquereux, 1880, Coal Flora of Pa., p. 389, Coal Meas.
radiato-plicatum, Dawson, 1873, Rep. Foss. Plants, p. 32, Subcarboniferous.
radicans, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 454, Coal Meas.
rectangulum, Wood, 1860, Proc. Acad. Nat. Sci. Phil., vol. 12, p. 519, Coal Meas.
rhombicum, Presl, 1833, (*Bergeria rhombica*), in Sternberg's Flor. d. Vorw., vol. 2, p. 184, Coal Meas.
rigens, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 429, Coal Meas.
rigidum, Lesquereux, 1884, Coal Flora of Pa., p. 839, Coal Meas.
rimosum, Sternberg, 1820, Flor. d. monde primitif, 1st Cahier, p. 25, and Coal Flora of Pa., p. 392, Coal Meas.

rugosum, syn. for *L. dichotomum*.
rushvillense, Andrews, 1875, Ohio Pal., vol. 2, p. 423, Coal Meas.
salebrosum, Wood, 1860, Proc. Acad. Nat. Sci., Phil., p. 520, Coal Meas.
scobiniforme, Meek, 1876, Bull. Phil. Soc. Wash., p. 13, Waverly Gr. Probably a syn. for *L. corrugatum*.

scutatum, Lesquereux, 1880, Coal Flora of Pa., p. 369, Coal Meas.

selaginoides, Sternberg, 1820, Flor. d. monde primitif, 2d Cahier, p. 35, Coal Meas.

sigillarioides, Lesquereux, 1858, founded upon a decorticated specimen of *L. vestitum* or *L. latifolium*.

simplex, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 454, Coal Meas.

squamiferum, Lesquereux, 1880, Coal Flora of Pa., p. 376, Coal Meas.

sternbergi, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 85, Coal Meas.

tetragonum, Sternberg, 1821, Flor. d. monde primitif, 2d Cahier, p. 35, Coal Meas.

tijoui, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 431, Coal Meas.

tumidum, Bunbury, 1847, (Lepidophloios tumidum,) Quar. Jour. Geo. Soc., vol. 3, p. 432, Coal Meas.

turbinatum, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 453, Kaskaskia Gr.

undulatum, Sternberg, 1820, Flor. d. monde primitif, 1st Cahier, p. 25, Coal Meas.

ursum, Wood, 1860, Proc. Acad. Nat. Sci., p. 240, Coal Meas.

veltheimianum, Sternberg, 1823, Vers. Darst. Flora der Vorwelt, vol. 1, p. 12, Kaskaskia Gr.

venustum, Wood, syn. for *L. obtusum*.
vestitum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 874, Coal Meas.

wortheni, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 452, Coal Meas.

LEPIDOPHLOIOS, Sternberg, 1823, Vers. Darst. Flora der Vorwelt. [Ety. *lepis*, scale; *phloios*, the bark.] Stems arborescent, erect, with four ranked branches disposed in spiral order; leaves coriaceous, linear, long, narrow, with a thick medial

FIG. 44.—*Lepidodendron sternbergi*.

ner
or
upp
cula
rho
thre
lose
nus.
acadia
Geo
Coal
antiqu
Can
auricu
Ill.,
crassic
der
dilat
Pa.,
ichthy
Flor
ichthyo
lepis
irregul
Ark.
laricin
larici
Cahie
lesquer
vol. 2
macrole
sarrac
Coal

FIG. 45.—*macrolepis*.

platystig
Geo.,
Coal
prominu
Geo.,
Coal
protuber
Ill., vo
sigillari
of Pa.
tetragon
Geo.,
Coal
tumidus,
tumid
LEPIDOPHY
Hist.
scale;
either
giopho
Type I

nerve, bearing at base thick, suberect or recurved bolsters, inflated in the upper part and dotted with small vascular points; leaf-scars transversely rhomboidal, marked horizontally by three vascular scars, minutely papillose under the cortex. Type *L. laricinus*.

acadianus, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol. p. 489, Coal Meas.

antiquus, Dawson, 1871, Foss. Plants Canada, p. 36, Devonian.

auriculatus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 439, Coal Meas.

crassicaulis, Corda, 1845, Beitrage zur Flora der Vorwelt, p. 18, Coal Meas.

dilatatus, Lesquereux, 1884, Coal Flora of Pa., p. 781, Coal Meas.

ichthyoderma, Lesquereux, 1880, Coal Flora of Pa., p. 426, Coal Meas.

ichthyolepis, see *Lepidodendron ichthyolepis*.

irregularis, Lesquereux, 1860, Geo. Sur. Ark. vol. 2, p. 311, Coal Meas.

laricinus, Sternberg, 1820, (*Lepidodendron laricinum*,) Flor. d. monde primitif, 1st Cahier, p. 25, Coal Meas.

lesquereuxi, Andrews, 1875, Ohio Pal., vol. 2, p. 423, Coal Meas.

macrolepidotus, Goldenberg, 1862, Flora sarræpontana fossilis, vol. 3, p. 37, Coal Meas.

FIG. 45.—*Lepidophloeos macrolepidotus*.

obcordatus, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 457, Coal Meas.

parvus, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 490, Coal Meas.

platystigma, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 490, Coal Meas.

prominulus, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 489, Coal Meas.

protuberans, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 440, Coal Meas.

sigillarioides, Lesquereux, 1880, Coal Flora of Pa., p. 425, Coal Meas.

tetragonum, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 490, Coal Meas.

tumidus, Lesquereux, see *Lepidodendron tumidum*.

LEPIDOPHYLLUM, Brongniart, 1828, Prodr. d. Hist. Veg. Foss., p. 87. [Ety. *lepis*, scale; *phyllon*, leaf.] Blades or bracts, either joined to sporanges, or sporangio-phores of *Lepidostrobus*, or isolated. Type *L. majum*.

acuminatum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 875. The name was pre-occupied by Gutbier in 1843, but as it is a *Lepidostrobus* the name may be retained.

affine, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 875, Coal Meas.

auriculatum, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 457, Coal Meas.

brevifolium, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 876, Coal Meas.

campbellianum, Lesquereux, 1884, Coal Flora of Pa., p. 786, Coal Meas.

coriaceum, Lesquereux, 1884, Coal Flora of Pa., p. 787, Coal Meas.

cultriforme, Lesquereux, 1884, Coal Flora of Pa., p. 785, Coal Meas.

elegans, Lesquereux, 1884, Coal Flora of Pa., p. 787, Coal Meas.

fallax, Lesquereux, 1884, Coal Flora of Pa., p. 786, Coal Meas.

foliaceum, see *Lepidostrobus foliaceus*.

gracile, Lesquereux, 1884, Coal Flora of Pa., p. 786, Coal Meas.

hastatum, see *Lepidostrobus hastatus*.

intermedium, Lindley & Hutton, 1831, Foss. Flora, vol. 1, p. 125, Coal Meas.

lanceolatum, see *Lepidostrobus lanceolatus*.

linearifolium, Lesquereux, 1880, Coal Flora of Pa., p. 452, Coal Meas.

majum, Brongniart, 1828, Prodr. d'une Hist. Veg. Foss., p. 87, and Coal Flora of Pa., p. 449, Coal Meas.

mansfieldi, Lesquereux, 1880, Coal Flora of Pa., p. 449, Coal Meas.

minutum, Lesquereux, 1884, Coal Flora of Pa., p. 787, Coal Meas.

morrisanum, Lesquereux, 1880, Coal Flora of Pa., p. 448, Coal Meas.

obtusum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 875, Coal Meas.

plicatum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 876, Coal Meas.

rostellatum, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 443, Coal Meas.

striatum, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 443, Coal Meas.

trinerve, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 87, and Lindley & Hutton's Foss. Flora, vol. 2, p. 195, Coal Meas.

truncatum, see *Lepidostrobus truncatus*.

tumidum, Lesquereux, 1880, Coal Flora of Pa., p. 448, Coal Meas.

LEPIDOSTROBUS, Brongniart, 1828, Prodr. d. Hist. Veg. Foss., p. 87. [Ety. *lepis*, scale; *strobilus*, cone.] Strobiles cylindrical or ovate, oblong, conical; composed of sporanges (spore-cases) subcylindrical or clavate, emarginate at the apex, supported in the middle lengthwise by bracts formed of a pedicel, attached like the sporanges in right angle to the axis, linear or oblanceolate, either simple, not longer than the sporanges, or prolonged into lanceolate, obtuse or acuminate laminae, curved upward on the outside of the strobiles and imbricated on their sides, or merely inflated

at the outer end, and covering the apex of the sporanges by a rhomboidal small shield; spores, triquetre on one side, half globular on the other, like those of the Lycopods, homomorphous or dimorphous. Type *L. ornatus*.

acuminatus, Lesquereux, 1858, (*Lepidophyllum acuminatum*), Geo. Pa., vol. 2, p. 875, Coal Meas.

aldrichi, Lesquereux, 1880, Coal Flora of Pa., p. 441, Coal Meas.

butleri, Lesquereux, 1884, Coal Flora of Pa., p. 840, Coal Meas.

connivens, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 442, Coal Meas.

foliaceus, Lesquereux, 1870, (*Lepidophyllum foliaceum*), Geo. Sur. Ill., vol. 4, p. 444, Coal Meas.

globosus, Dawson, 1861, Can. Nat. and Geo., vol. 6, p. 174, Devonian.

goldenbergi, Schimper, 1872, *Traité de Paléontologie Végétale*, vol. 2, p. 61, Coal Meas.

hastatus, Lesquereux, 1858, (*Lepidophyllum hastatum*), Geo. Sur. Pa., vol. 2, p. 876, Coal Meas.

incertus, Lesquereux, 1880, Coal Flora of Pa., p. 442, Coal Meas.

lacoel, Lesquereux, 1880, Coal Flora of Pa., p. 439, Coal Meas.

FIG. 46.—*Lepidostrobus hastatus*.

lanceolatus, Brongniart, 1828, (*Lepidophyllum lanceolatum*), Prodr. d. Hist. d. Vég. Foss., p. 87, and Coal Flora of Pa., p. 436, Coal Meas.

lancifolius, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 442, Coal Meas.

latus, Lesquereux, 1884, Coal Flora of Pa., p. 841, Coal Meas.

longifolius, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol. p. 489, Coal Meas.

manfieldi, Lesquereux, 1880, Coal Flora of Pa., p. 444, Coal Meas.

mirabilis, Newberry, 1873, (*Polysporia mirabilis*), Ohio Pal., vol. 1, p. 362, Low. Coal Meas.

oblongifolius, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 441, Coal Meas.

FIG. 47.—*Lepidostrobus ornatus*. Cone $\frac{1}{2}$ size.

ornatus, Parkinson, 1811, Organic Remains, vol. 1, pl. 9, fig. 1, and Coal Flora of Pa., p. 440, Coal Meas.

ovatifolius, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 441, Coal Meas.

pinaster, Lindley & Hutton, 1837, Foss. Flora, vol. 3, p. 129, Coal Meas.

prælongus, Lesquereux, 1880, Coal Flora of Pa., p. 433, Coal Meas.

princeps, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 455, Coal Meas.

quadratus, Lesquereux, 1880, Coal Flora of Pa., p. 444, Coal Meas.

richardsoni, Dawson, 1861, Can. Nat. and Geo., vol. 6, p. 174, Devonian.

salisburyi, Lesquereux, 1880, Coal Flora of Pa., p. 443, Coal Meas.

spectabilis, Lesquereux, 1880, Coal Flora of Pa., p. 435, Coal Meas.

squamosus, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 489, Coal Meas.

stachoides, see *Asterophyllites stachoides*, trigonolepis, Bunbury, 1847, Quar. Jour. Geo. Soc., vol. 3, p. 432, Coal Meas.

truncatus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 442, Coal Meas.

variabilis, Lindley & Hutton, 1833, Foss. Flora, vol. 1, p. 31, and Coal Flora of Pa., p. 434, Coal Meas.

LEPIDOXYLON, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 334, and Coal Flora of Pa., p. 557. [Ety. *lepis*, scale; *xylon*, wood.]

Stems large, tapering to a point; bark thin, covered with leafy scales; leaves variable, sublinear, narrowed or enlarged to the point of attachment, forking upward in two or more laciniae; nervation distinct with the glass; primary nerves parallel, buried in the epidermis, inflated or half round; intermediate veinlets thin, visible on the decorticated face. Type *L. anomalum*. *anomalum*, Lesquereux, 1880, Coal Flora of Pa., p. 557, Coal Meas.

LEPTOPHYLLUM, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 316. [Ety. *leptos*, slender; *phyllois*, the bark of a tree.]

Stem covered with continuous rhombic areoles, each with a single small scar a little above its center, and above this a very slight furrow; decorticated stems, with spiral punctiform scars in slight depressions; bark thin, pith cylinder very large, with transverse markings of the character of *Sternbergia*. Type *L. rhombicum*.

rhombicum, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 316, Devonian.

LESCUROPTERIS, Schimper, 1869, *Paläontologie Végétale*, vol. 1, p. 465. [Ety. proper name; *pteris*, a fern.] Fronds large, bi-

FIG. 48.—*Lescuropteris adiantites*.

tripinnate; rachis broad; foliate; pinnae pinnatifid, close, oblique; divisions ovate, acute, inclined outside, connate

to the
prima
lower
from
from
forkin
adiant
teris a
vol. 6
163, C
moorii,
moori
Meas.
LESLEYA,
Pa., p
ne sin
gradua
traver
der th
equal,
L. gra
grandis,
Pa., p
microph
of Pa.,

FIG.
LICHOPHYCU
1, p. 99
sea-wee
elongate
ing from
ing sing
angle.
flabellum,
Soc. Na
Riv. Gr

to the middle, decurrent to the rachis, primary nerve thin, dichotomous; lower pairs of lateral veins emerging from the rachis, the other alternately from the midrib, forking twice, the upper forking once or simple. Type *L. moorii*. *adiantites*, Lesquereux, 1854, (*Neuropteris adiantites*,) Bost. Jour. Nat. Hist., vol. 6, p. 419, and Coal Flora of Pa., p. 163, Coal Meas.

moorii, Lesquereux, 1858, (*Neuropteris moorii*,) Geo. Sur. Pa., vol. 2, p. 860, Coal Meas.

LESLEYA, Lesquereux, 1880, Coal Flora of Pa., p. 142. [Ety. proper name.] Pinnæ simple, very entire, sublanceolate, gradually narrowing toward the base, traversed by a thick costa effaced under the apex; veins oblique, curved, equal, repeatedly dichotomous. Type *L. grandis*.

grandis, Lesquereux, 1880, Coal Flora of Pa., p. 143, Coal Meas.

microphylla, Lesquereux, 1884, Coal Flora of Pa., p. 831, Coal Meas.

FIG. 49.—*Lycophyus ottawense*.

LICOPHYCUS, Billings, 1862, Pal. Foss., vol. 1, p. 99. [Ety. *likros*, a fan; *phykos*, sea-weed.] Composed of numerous, elongated, subcylindrical stems, radiating from a common root, and remaining single, or branching at an acute angle. Type *L. ottawense*.

flabellum, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 25, Hud. Riv. Gr.

formosum, Billings, 1866, Catal. Sil. Foss.

Antic., p. 72, Hud. Riv. Gr.

hiltonense, Billings, 1862, Pal. Foss., vol.

1, p. 101, Black Riv. and Trenton Gr.

hudsonicum, Billings, 1862, Pal. Foss., vol.

1, p. 101, Hud. Riv. Gr.

minor, Billings, 1862, Pal. Foss., vol. 1,

p. 100, Trenton Gr.

ottawense, Billings, 1862, Pal. Foss., vol.

1, p. 99, Trenton Gr.

robustum, Billings, 1866, Catal. Sil. Foss.

Antic., p. 72, Hud. Riv. Gr.

vagans, Billings, 1866, Catal. Sil. Foss.

Antic., p. 72, Hud. Riv. Gr.

Lithodictyon becki, Conrad. Not properly defined; but see *Dictyophyton becki*.

LONCHOPTERIS, Brongniart, 1828, Prodr. Hist.

Veg. Foss., p. 59. [Ety. *lonche*, spear;

pterus, fern.] Pinnate or bipinnate;

pinnules contiguous at the base, nearly

at right angles to petiole, oblong-elon-

gate, obtuse, middle-sized veins reticu-

lated with finer ones. Type *L. bricei*.

tenuis, Dawson, 1863, Can. Nat. and Geol.,

vol. 8, and Acad. Geol., p. 483, Coal

Meas.

Lycopodiolithes elegans, see *Lepidoden-*

dron elegans.

LYCOPIDITES, Brongniart, 1822, Mem. du

Mus. d'Hist. Nat. de Paris, and Lycopodiolithis of Schlottheim and

Sternberg. [Ety. from *Lycopodium*,

the club moss.] Plants herbaceous;

leaves of the same or of two differ-

ent forms upon the same branches,

distichous or in spiral order; fruc-

tifications in small cylindrical spikes.

Type *L. pinniformis*.

annulariifolius, Lesquereux, 1870,

Geo. Sur. Ill., vol. 4, p. 426, Coal

Meas.

arborescens, Lesquereux, 1884, Coal

Flora of Pa., p. 778, Coal Meas.

asterophyllitifolius, Lesquereux, 1866,

Geo. Sur. Ill., vol. 2, p. 447, Coal

Meas.

cavifolius, Lesquereux, 1861, Geo. Sur.

Ky., vol. 4, p. 437, Coal Meas.

comosus, Dawson, 1863, Quar. Jour.

Geo. Soc., vol. 19, p. 462, Devo-

nian.

flexifolius, Lesquereux, 1884, Coal

Flora of Pa., p. 779, Coal Meas.

lacoeci, Lesquereux,

1884, Coal Flora of

Pa., p. 780, Coal

Meas.

matthewi, Dawson,

1861, Can. Nat. and

Geo., vol. 6, p. 171, and

Acad. Geol., p. 540, De-

vonian.

meeki, Lesquereux, 1870,

Geo. Sur. Ill., vol. 4, p.

426, Coal Meas.

ortoni, Lesquereux, 1880,

Coal Flora of Pa., p. 357, Coal Meas.

pendulus, Lesquereux, 1880, Coal Flora

of Pa., p. 357, Coal Meas.

4.

5.

6.

7.

8.

9.

10.

FIG. 50.—*Lycopodites matthewi*, Dawson, 1861, Can. Nat. and Geo., vol. 6, p. 171, and Acad. Geol., p. 540, Devonian. *a*, branch and leaves; *b*, *c*, *d*, leaves.

- plumula*, see *Plumalina plumula*.
richardsoni, Dawson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 461, Devonian.
simplex, Lesquereux, 1884, Coal Flora of Pa., p. 779, Coal Meas.
strictus, Lesquereux, 1880, Coal Flora of Pa., p. 360, Coal Meas.
uncinatus, Lesquereux, 1886, (Selaginites *uncinatus*.) Geo. Sur. Ill., vol. 2, p. 446, Coal Meas.
vannuxemi, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 314, Syn. for *Plumalina plumula*.
MACROSTACHYA, Schimper, 1869, *Traité de Paléontologie Végétale*, vol. 1, p. 332. [Ety. *makros*, long; *stachys*, a plant.] Plants arborescent, articulate; articulations close; cortex thin, smooth or striate; impressions of the internal surface plano-costate; furrows narrow, alternating at the articulations; leaves appressed, linear, carinate or marked with a medial nerve, acuminate, finely truncate; leaf scars marked upon the articulations by transversely oval rings, like the links of a chain; scars of branches verticillate, large, round, umbonate, with a stigmaroid central mamilla; spikes very large, cylindrical; bracts lanceolate, costate in the middle, imbricate, scarcely longer than the internodes. Type *M. infundibuliformis*.
aperta, Lesquereux, 1858, (Asterophyllites *apertus*.) Geo. Sur. Pa., vol. 2, p. 852, Coal Meas.
communis, Lesquereux, 1884, Coal Flora of Pa., p. 828, Coal Meas.
infundibuliformis, Brongniart, 1828, (Equisetum *infundibuliforme*.) Hist. Vég. Foss., t. 1, p. 119, Coal Meas.
lanceolata, Lesquereux, 1858, (Asterophyllites *lanceolatus*.) Geo. Sur. Pa., vol. 2, p. 852, Coal Meas.
minor, Lesquereux, 1884, Coal Flora of Pa., p. 829, Coal Meas.
MEGALOPTERIS, Dawson, 1871, Foss. Plants Dev. and Up., Sil. Formations, p. 51. [Ety. *megale*, great; *pterus*, fern.] Fronds very large, pinnate, ultimate pinnæ oblique, sublinear or lanceolate, entire, the lower side broadly decurrent on the rachis, which thus becomes alate, the upper narrowed in a curve, confluent; midrib thick, canalliculate on the upper surface, half cylindrical on the lower, gradually narrowed, but distinct to the apex of the leaves; veins open, emerging from the rachis in a more open angle of divergence, curving upward in reaching the borders, close dichotomous. Type *M. dawsoni*.
abbreviata, Lesquereux, 1880, Coal Flora of Pa., p. 151, Coal Meas.
dawsoni, Hart, 1868, (Neuropteris *dawsoni*.) Acad. Geol., p. 550, Devonian.
dentata, Lesquereux, 1884, Coal Flora of Pa., p. 833, Coal Meas.
fasciculata, Lesquereux, 1880, Coal Flora of Pa., p. 150, Coal Meas.

- hartti*, Andrews, 1875, Ohio Pal., vol. 2, p. 416, Coal Meas.
lata, Andrews, 1875, Ohio Pal., vol. 2, p. 417, Coal Meas.
marginata, Lesquereux, 1880, Coal Flora of Pa., p. 152, Coal Meas.
minima, Andrews, 1875, Ohio Pal., vol. 2, p. 416, Coal Meas.
ovata, Andrews, 1875, Ohio Pal., vol. 2, p. 417, Coal Meas.
rectinervis, Lesquereux, 1884, Coal Flora of Pa., p. 744, Coal Meas.
serrata, Lesquereux, 1884, Coal Flora of Pa., p. 834, Coal Meas.

FIG. 51.—*Megalopteris southwelli*.

- southwelli*, Lesquereux, 1880, Coal Flora of Pa., p. 148, Coal Meas.
MEGAPHYTON, Artis, 1828, Antedil. Phytol., p. 20. [Ety. *megas*, great; *phyton*, a plant.] Scars large, round-quadrate in outline, mostly contiguous, placed in opposite biserial rows; internal disks convex, with central or vascular impressions in the form of a horseshoe, or a medial band dividing the disks into two lobes, joined in the middle. Type *M. frondosum*.
goldenbergi, Weiss, 1860, Zeitsch. d. deutsch. Geol. Gesellsch. XII, p. 510, Coal Meas.
grandeuryi, Lesquereux, 1880, Coal Flora of Pa., p. 350, Coal Meas.
humile, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol. p. 486, Coal Meas.
maclayi, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 458, Coal Meas.
magnificum, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 486, Coal Meas.

FIG. 52.—*Megaphyton protuberans*.

protub.
 Ill.,
Nematoph
Micr
logani,
 NEMATOP
Pern
nema
 ered
 epide
 leave
 and t
 out,
 unite
 band
 angust
 Perm
 Meas
 NEMATOP
 Geo.
 a thr
 Penh
 belon
 Nema
 with
 comp
 with

FIG. 53.—*Nematopteris lanceolata*. Single pin-nule.

base,
 margin
 lanceolat
 1, p. 3

protuberans, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 458, Kaskaskia Gr.
Nematophycus, Carruthers, 1872, Month. Micro. Jour. Syn. for Prototaxites.
logani, see Prototaxites logani.

NEMATOPHYLLUM, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 35. [Ety. *nema*, thread; *phyllon*, leaf.] Stem covered with a thick, very finely striate epidermis, internodes remote, swollen; leaves verticillate, numerous, very long and thread-like, of equal width throughout, finely striate, without nerves, united at the base in a narrow annular band. Type *N. angustum*.

angustum, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 35, Coal Meas. or Permian.

NEMATOTRYLON, Dawson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 466. [Ety. *nema*, a thread; *trylon*, wood.] Carruthers, Penhallow, and others say this genus belongs to the Alge, and is a syn. for *Nematophycus*. Fragments of wood, with a smooth bark and a tissue wholly composed of elongated cylindrical cells, with irregular pores or markings; no pith, medullary rays or rings of growth. Type *N. crassum*.

crassum, Dawson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 466, Devonian.

tenuis, Dawson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 467, Devonian.

Nephropteris, Brongniart, 1828, Tab. des gener.

elegans, see Cyclopteris elegans.

fimbriata, see Neuropteris fimbriata.

germari, see Cyclopteris germari.

hirsuta, see Cyclopteris hirsuta.

laciniata, see Cyclopteris laciniata.

orbicularis, see Cyclopteris orbicularis.

trichomanoides, see Cyclopteris trichomanoides.

undans, see Cyclopteris undans.

NEUROPTERIS, Newberry, 1873, Ohio Pal., vol. 1, p. 378.

[Ety. *nerion*, the oleander; *pteris*, a fern.] Frond pinnate or bipinnate; rachis strong, punctate; pinnules lanceolate, simple, entire; medial nerve strong, extending from base to summit; secondary nerves given off at an acute angle, numerous simple or forked at

base, parallel, equal; fructifications marginal. Type *N. lanceolata*.

lanceolata, Newberry, 1873, Ohio Pal., vol. 1, p. 381, Coal Meas.

NEUROPTERIS, Brongniart, 1822, Mem. du Mus. d'Hist. Nat. de Paris, t. 8, p. 203, and Prodr. d. Hist. d. Veg. Foss., p. 52.

[Ety. *neuron*, nerve; *pteris*, fern.] Fronds simple, bi, tri-pinnate; pinnules varying from round to ovate, obtuse, or obtusely acuminate, mostly entire, rounded, cordate, or auricled at the base, attached to the rachis by the middle; sessile, or rarely short pedicled; veins either from the base of the pinnules or from a costa, diverging fan-like and arched backward, in passing toward the borders, many times dichotomous; costa generally dissolved at or below the middle; basilar veins simple or in fascicles. Type *N. acuminata*.

acuminata, Schlotheim, 1820, (Filicites acuminatus,) Petrefactenkunde, p. 412, and Coal Flora of Pa., p. 123, Coal Meas.

acutifolia, Brongniart, 1828, Hist. d. Veg. Foss., p. 229, Coal Meas.

adiantites, see Lescuropteris adiantites.

agassizi, Lesquereux, 1880, Coal Flora of Pa., p. 117, Coal Meas.

angustifolia, Brongniart, 1828, Hist. d. Veg. Foss., p. 231, and Coal Flora of Pa., p. 89, Coal Meas.

anomala, Lesquereux, 1880, Coal Flora of Pa., p. 118, Coal Meas.

aspera, Lesquereux, 1880, Coal Flora of Pa., p. 121, Coal Meas.

attenuata, Lindley & Hutton, 1837, Foss. Flora, vol. 3, p. 65, Coal Meas.

auriculata, Brongniart, 1828, Hist. d. Veg. Foss., p. 236, Coal Meas.

biformis, Lesquereux, 1880, Coal Flora of Pa., p. 121, Coal Meas.

blissi, Lesquereux, 1884, Coal Flora of Pa., p. 737, Coal Meas.

callosa, Lesquereux, 1880, Coal Flora of Pa., p. 115, Coal Meas.

capitata, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 363, Coal Meas.

carri, Lesquereux, 1884, Coal Flora of Pa., p. 731, Coal Meas.

cisti, Brongniart, 1828, Hist. d. Veg. Foss., p. 238, Coal Meas.

clarksoni, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 857, Coal Meas.

collinsi, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 382, Coal Meas.

cordata, Brongniart, 1828, Hist. Veg. Foss., p. 229, and Coal Flora of Pa., p. 91, Coal Meas.

cordato-ovata, see Pseudocopteris cordato-ovata.

coriacea, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 387, Coal Meas.

crassa, Dawson, 1868, Acad. Geol., p. 551, Devonian.

crenulata, Brongniart, 1828, Hist. Veg. Foss., p. 234, and Coal Flora of Pa., p. 116, Coal Meas.

cyclopteroides, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol. p. 482, Coal Meas.

dawsoni, see Megalopteris dawsoni.

FIG. 53.—*Neuropteris lanceolata*. Single pinnule.

decipiens, Lesquereux, 1880, Coal Flora of Pa., p. 93, Coal Meas.
 delicatula, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 858, Coal Meas.
 dentata, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 859, Coal Meas.
 desori, Lesquereux, 1854, Bost. Jour. Nat. Hist., vol. 6, p. 418, and Geo. Sur. Pa., vol. 2, p. 859, Coal Meas.
 dictyopteroides, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 49, Coal Meas. or Permian.
 dilatata, Lindley & Hutton, 1835, (Cyclopteris dilatata,) Foss. Flora, vol. 2, p. 29, Coal Meas.
 elrodi, Lesquereux, 1880, Coal Flora of Pa., p. 107, Coal Meas.
 eveni, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 430, Coal Meas.
 fasciculata, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 381, Coal Meas.
 fimbriata, Lesquereux, 1854, (Cyclopteris fimbriata,) Jour. Bost. Soc. Nat. Hist., p. 416, and Coal Flora of Pa., p. 81, Coal Meas.
 fissa, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 857, Coal Meas.
 flexuosa, Sternberg, 1825, Vers. Darst. Flora der Vorwelt, p. 16, Coal Meas.
 germari, Goepfert, 1836, (Adiantites germari,) Systema Filicum Fossilium, p. 218, and Coal Flora of Pa., p. 113, Coal Meas.
 gibbosa, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 858, Coal Meas.
 gigantea, Sternberg, 1825, Vers. Darst. Flora der Vorwelt, p. 16, Coal Meas.
 grangeri, Brongniart, 1828, Hist. Veg. Foss., p. 237, and Coal Flora of Pa., p. 105, Coal Meas.
 griffithi, Lesquereux, 1884, Coal Flora of Pa., p. 737, Coal Meas.
 heterophylla, Brongniart, 1822, (Filicites heterophylla) Mem. du Mus. d'Hist. Nat. de Paris, t. 8, p. 203, Coal Meas.

FIG. 54.—Neuopteris hirsuta.

hirsuta, Lesquereux, 1854, Bost. Jour. Nat. Hist., vol. 6, p. 417, and Coal Flora of Pa., p. 88, Coal Meas.
 inflata, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 431, Coal Meas.
 ingens, Lindley & Hutton, 1835, Foss. Flora, vol. 2, p. 29, Coal Meas.
 lacerata, syn. for Neuropteris fimbriata.
 laciniata, Lesquereux, 1858, (Cyclopteris laciniata,) Geo. Sur. Pa., vol. 2, p. 855, Coal Meas.
 loshi, Brongniart, 1828, Hist. d. Veg. Foss., p. 242, and Coal Flora of Pa., p. 98, Coal Meas.
 microphylla, Brongniart, 1828, Hist. Veg. Foss., p. 245, Coal Meas.

minor, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 859, Coal Meas.
 missouriensis, Lesquereux, 1880, Coal Flora of Pa., p. 104, Coal Meas.
 moori, see Lescuropteris moorii.
 oblongifolia, Lesquereux, 1884, Coal Flora of Pa., p. 732, Coal Meas.
 obscura, Lesquereux, 1880, Coal Flora of Pa., p. 108, Coal Meas.
 odontopteroides, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 50, Coal Meas. or Permian.
 pachyderma, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 430, Coal Meas.
 perelegans, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 482, Coal Meas.
 platynervis, Fontaine & White, 1880, Perm. or Up. Carb. Flora, pl. 8, fig. 2, Coal Meas. or Permian.
 plicata, Sternberg, 1825, Vers. Darst. Flora der Vorwelt, p. 74, and Coal Flora of Pa., p. 96, Coal Meas.
 polymorpha, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 320, Devonian.
 rarineris, Bunbury, 1847, Quar. Jour. Geo. Soc., vol. 3, p. 425, and Coal Flora of Pa., p. 109, Coal Meas.
 reniformis, Brongniart, 1828, (Cyclopteris reniformis,) Hist. d. Veg. Foss., p. 216, and Coal Flora of Pa., p. 77, Coal Meas.
 retorquata, Dawson, 1871, Foss. Plants Canada, p. 50, Devonian.
 rogersi, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 856, Coal Meas.
 rotundifolia, Brongniart, 1828, Hist. Veg. Foss., p. 238, and Coal Flora of Pa., p. 97, Coal Meas.
 selwyni, Dawson, 1871, Foss. Plants Canada, p. 50, Devonian.
 serrulata, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 320, Devonian.
 smilacifolia, Sternberg, 1824, Vers. Darst. Flora der Vorwelt, vol. 2, p. 29, Coal Meas.
 smithi, Lesquereux, 1876, Geo. Rep. of Alabama, p. 76, and Coal Flora of Pa., p. 106, Coal Meas.
 soreti, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 53, and Hist. d. Veg. Foss., t. 1, p. 244, Coal Meas.
 speciosa, Lesquereux syn. for N. rogersi.
 subfalcata, Lesquereux, 1880, Coal Flora of Pa., p. 102, Coal Meas.
 tenuifolia, Sternberg, 1825, Vers. Darst. Flora der Vorwelt, p. 17, and Coal Flora of Pa., p. 100, Coal Meas.
 tenuinervis, see Odontopteris tenuinervis.
 trichomanoides, Brongniart, 1828, (Cyclopteris trichomanoides,) Hist. d. Veg. Foss., p. 217, and Coal Flora of Pa., p. 79, Coal Meas.
 undans, Lesquereux, 1854, Bost. Jour. Nat. Hist., vol. 6, p. 418, and Geo. Sur. Pa., vol. 2, p. 859, Coal Meas.
 verbenifolia, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 431, Coal Meas.

vermi
Ky.
villier
Veg
NEUGER
exp
Flor
37.
a stu
attac
base
and
beiner
Pflar
bockach
bockach
dispar.
Geol
Coal

F
Neogger
d'Hist
odous,
large,
subalt
gener
rachis
decur
arate
middl
towar
ually
leaflet
tached
veins
rarely
dichot
curve,
O. bra
abbrevi
of Pa.
aequalis,
vol. 2,
affinis,
Pa., p.
alata, L
Foss.,
131, C
alpina, S
pina, S
and C
Meas.

vermicularis, Lesquereux, 1861, Geo. Sur. Ky., vol. 4, p. 434, Coal Meas.
villiersi, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 53, Coal Meas.

NEGGERATHIA, Sternberg, 1828, Essai d'un expose Geognostico-botanique de la Flore du monde primitif, 2d Cahier, p. 37. [Ety. proper name.] Branch with a slender rachis bearing pinnate leaves attached to the stem by a semi-twisted base, dilated upward, veins flabellate and dichotomous. Type *N. foliosa*.

belnertiana, Gœppert, 1842, Gatt. d. Foss. Pflanzen, Coal Meas.

bockschii, see *Aneimites bockschii*.

bockschiana, syn. for *Aneimites bockschii*. dispar, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 480, Coal Meas.

FIG. 55.
Neggerathia dispar.

d'Hist. Nat. de Paris, t. 8, p. 203. [Ety. *odous*, tooth; *pteria*, fern.] Fronds large, bipinnate; pinnæ opposite or subalternate; pinnules of various forms, generally oblong, obtuse, joined to the rachis by their whole base, sometimes decurrent, either disjointed and separate to the base or connate to the middle, generally becoming confluent toward the top of the pinnæ, and gradually effaced in passing to a terminal leaflet; lower pinnules sometimes attached to the main rachis and difform; veins emerging from the rachis, more rarely from a midrib; veinlets thin, dichotomous, diverging straight or in curve, in passing to the borders. Type *O. brardi*.

abbreviata, Lesquereux, 1880, Coal Flora of Pa., p. 138, Coal Meas.

æqualis, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 434, Coal Meas.

affinis, Lesquereux, 1884, Coal Flora of Pa., p. 742, Coal Meas.

alata, Lesquereux, 1858, Catal. Pottsville Foss., p. 6, and Coal Flora of Pa., p. 131, Coal Meas.

alpina, Sternberg, 1825, (*Neuropteris alpina*.) Flora d. Vorwelt, vol. 2, p. 76, and Coal Flora of Pa., p. 126, Coal Meas.

flabellata, Lindley & Hutton, 1832, Foss. Flora, vol. 1, p. 80, Coal Meas.

gilboensis, Dawson, 1871, Quar. Jour. Geo. Soc., vol. 27, p. 273, Chemung Gr.

minor, see *Archæopteris minor*.

obliqua, see *Archæopteris obliqua*.

obtus, see *Aneimites obtusus*.

ODONTOPTERIS, Brongniart, 1822, Mem. du Mus.

antiqua, Dawson, 1863, Can. Nat. and Geo., Coal Meas.

bradleyi, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 390, Coal Meas.

brardi, Brongniart, 1822, Mem. du Mus. d'Hist. Nat. de Paris, t. 8, p. 205, tab. 2, fig. 5, and Coal Flora of Pa., p. 132, Coal Meas.

britannica, Guthrie, 1842, Abdrucke u. Verst. d. Zwick. Schwarzk. u. sel. Umgeb. Zwick., p. 68, and Coal Flora of Pa., p. 830, Coal Meas.

cornuta, Lesquereux, 1880, Coal Flora of Pa., p. 128, Coal Meas.

crenulata, of Brongniart, as indentified by Lesquereux in Geo. Sur. Pa., vol. 2, p. 860, is *O. subcrenulata*.

dawsoniana, n. sp. Devonian. Proposed instead of *O. squamosa*, in Quar. Jour. Geo. Soc. Lond., vol. 37, p. 305, which was preoccupied.

deformata, Lesquereux, 1880, Coal Flora of Pa., p. 141, Coal Meas.

densifolia, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 54, Coal Meas. or Permian.

dilatata, Lesquereux, 1884, Coal Flora of Pa., p. 831, Coal Meas.

dubia, Lesquereux, 1858, Geo. Sur. Penn., vol. 2, p. 860, Coal Meas.

gracillima, Newberry, 1873, Ohio Pal., vol. 1, p. 382, Coal Meas.

heterophylla, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 433, Coal Meas.

intermedia, Lesquereux, 1860, Geo. Sur. Ark., vol. 2, p. 313, Coal Meas.

leacurei, Wood, 1860, Trans. Am. Phil. Soc., vol. 13, p. 348, Coal Meas.

monstruosa, Lesquereux, 1884, Coal Flora of Pa., p. 741, Coal Meas.

nervosa, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 52, Coal Meas. or Permian.

neuropteroides, Newberry, 1873, Ohio Pal., vol. 1, p. 381. The name was preoccupied by R. Emer, and the species has been named *O. newberryi*.

newberryi, Lesquereux, 1880, Coal Flora of Pa., p. 127, Low. Coal Meas.

obtusiloba var. *rarinervis*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 52, Coal Meas. or Permian.

pachyderma, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 53, Coal Meas. or Permian.

patens, Lesquereux, 1884, Coal Flora of Pa., p. 740, Coal Meas.

FIG. 56.
Odontopteris gracillima.

reichiana, Lesquereux, 1884, Coal Flora of Pa., p. 831, Coal Meas.
rotundifolia, Wood, 1866, Trans. Am. Phil. Soc., vol. 13, p. 348, Coal Meas.
schlotheimi, Brongniart, 1828, Hist. d. Veg. Foss., p. 256, and Coal Flora of Pa., p. 136, Coal Meas.

FIG. 57.—*Odontopteris schlotheimi*.

subcrenolata, Lesquereux, 1880, Coal Flora of Pa., p. 137, Coal Meas.
subcuneata, Bunbury, 1847, Quar. Geo. Jour., vol. 3, p. 427, and Coal Flora of Pa., p. 134, Coal Meas.

tenuinervis, Lesquereux, 1858, (*Neuropteris tenuinervis*), Geo. Sur. Pa., vol. 2, p. 859, Coal Meas.

wortheni, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 432, Coal Meas.

OLIGOCARPIA, Göppert, 1841-48, Die Gattungen der fossilen Pflanzen, p. 3. [*Ety. oligos*, few; *karpus*, fruit.] Fronds bipinnate or tripinnatifid; primary pinnules oblong-lanceolate; secondary divisions, open, linear, pinnately divided in oblong or half round lobes or leaflets, connate at the base, crenulate; primary and secondary veins nearly of the same size, thin but distinct; lateral veins curved to the borders, simple or forked. Type *O. gutbieri*.

alabamensis, Lesquereux, 1875, Geo. Rep. Ala., p. 76, and Coal Flora of Pa., p. 266, Coal Meas.

flagellaris, Lesquereux, 1858, (*Sphenopteris flagellaris*), Geo. Sur. Pa., vol. 2, p. 862, Coal Meas.

gutbieri, Göppert, 1841-48, Die Gattungen der fossilen Pflanzen, p. 3, Coal Meas.

ORMOXYLON, Dawson, 1871, Foss. Plants Canada, p. 14. [*Ety. ormos*, a chain, a cord; *xylon*, wood.] Woody stems, with cells of the character of those of *Dadoxylon*, very thick walled, with three rows of hexagonal areoles, having oval pores and medullary rays of one row of cells. Pith cavity composed of a series of spherical chambers, separated by thick, transverse cellular partitions. Type *O. erianum*.

erianum, Dawson, 1871, Foss. Plants Canada, p. 14, Portage Gr.

sphenopteroides, Lesquereux, 1880, Coal Flora of Pa., p. 139, Coal Meas.

squamosa, Lesquereux, 1854, Bost. Jour. Nat. Hist., vol. 6, p. 419, and Geo. Sur. Pa., vol. 2, p. 860, Coal Meas.

squamosa, Dawson, 1881, Quar. Jour. Geo. Soc. Lond., vol. 37, p. 306, Devonian. The name was preoccupied. See *O. dawsoniana*.

ORTHOGONIOPTERIS, Andrews, 1875, Ohio Pal., vol. 2, p. 418. [*Ety. orthogoniopteris*, rectangular-fern.] Frond simply pinnate; pinnules alternate, lanceolate or oblong-linear, rounded and tapering to an acute point, enlarged and decurrent on the lower side to an auricle rounded in the upper part in joining the lamina a little above its point of attachment to the rachis; medial nerve thick, ascending to the apex; nervules fine and numerous, uniform, at right angle to the midrib, decurring to it at the point of attachment, forking once near the base. Type *O. clara*.

clara, Andrews, 1875, Ohio Pal., vol. 2, p. 419, Coal Meas.
gilberti, Andrews, 1875, Ohio Pal., vol. 2, p. 420, Coal Meas.

Pachyphyllum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2. [*Ety. pachys*, thick; *phylon*, a leaf.] This name was preoccupied in the class Polypi. See *Rhacophyllum*.

affine, see *Rhacophyllum affine*.

fimbriatum, see *Rhacophyllum fimbriatum*.

hirsutum, see *Rhacophyllum hirsutum*.

laceratum, see *Rhacophyllum laceratum*.

lactuca, see *Rhacophyllum lactuca*.

PACHYPTERIS, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 49. [*Ety. pachys*, thick; *pterus*, fern.] Frond pinnate or bipinnate, bearing opposite coriaceous pinnules, with a medial nerve or without nervation, narrowed toward the base, not joined to the rachis. Type *P. lanceolata*.

gracillima, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 419, Coal Meas.

PALÆOPHYCUS, Hall, 1847, Pal. N. Y., vol. 1, p. 7. [*Ety. palaios*, ancient; *phykos*, seaweed.] Stems simple or dichotomous, branches cylindrical or slightly flattened, obtuse, surface smooth or dotted. Type *P. tubularis*.

articulatum, Winchell, 1864, Am. Jour. Sci. and Arts, 2d series, vol. 37, p. 231, Potsdam Gr.

beauharnoiseense, Billings, 1862, Pal. Foss., vol. 1, p. 98, Calif. Gr.

beverleyense, Billings, 1862, Pal. Foss., vol. 1, p. 97, Potsdam Gr.

congregatum, Billings, 1861, Pal. Foss., vol. 1, p. 3, Potsdam Gr.

divaricatum, Lesquereux, 1876, 7th Ann. Rep. Geol. Sur. Ind., p. 138, Coal Meas.

funiculus, Billings, 1862, Pal. Foss., vol. 1, p. 98, Calif. Gr.

FIG. 58. *Orthogoniopteris clara*, part of a pinnule.

gracillima
Geo.

FIG. 58.
Palæophycus gracillima

occidentalis
Black

p. 33
pinnules

Geo.
vol. 1

rugosus
63, T.

simplex
63, T.

striatum
22, C.

tortuosus
p. 6,

tubularis
7, Cal.

virgatus
263, H.

Palæopteris

by Geo.

acadiana,
hartii, se

Palæoxylon

d. Veg.

cient;

Type

have b

Ameri

ferred

appendic

latum.

corrugato

prendeli

Palæacites

naggarathi

rathi.

Pecopteris

Foss. i

Paris,

pterus,

pinnæ

hering

often r

decurr

ous, or

rived

pinnule

P. long

tioned

Foss.,

Coal F

pennifi

abbreviat

Veg. F

Pa., p.

gracile, Lesquereux, 1876, 7th Ann. Rep. Geol. Sur. Ind., p. 137, Coal Meas.

incipiens, Billings, 1861, Pal. Foss., vol. 1, p. 2, Potsdam Gr. informe, Winchell, 1864, Am. Jour. Sci. and Arts, vol. 37, p. 232, Potsdam Gr. irregular, Hall, 1847, Pal. N. Y., vol. 1, p. 8, Calcif. Gr. milleri, Lesquereux, 1876, 7th Ann. Rep. Geol. Sur. Ind., p. 136, Coal Meas.

obscurum, Billings, 1862, Pal. Foss., vol. 1, p. 98, Trenton Gr. occidentale, Whitfield, 1877, Rep. Pal. Black Hills, p. 7, and Geol. Black Hills, p. 332, Potsdam Gr.

plumosum, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 50, and Geo. Wis., vol. 4, p. 169, Potsdam Gr.

rugosum, Hall, 1847, Pal. N. Y., vol. 1, p. 63, Trenton Gr.

simplex, Hall, 1847, Pal. N. Y., vol. 1, p. 63, Trenton Gr.

striatum, Hall, 1852, Pal. N. Y., vol. 2, p. 22, Clinton Gr.

tortuosum, Hall, 1852, Pal. N. Y., vol. 2, p. 6, Medina sandstone.

tubulare, Hall, 1847, Pal. N. Y., vol. 1, p. 7, Calcif. Gr.

virgatum, Hall, 1847, Pal. N. Y., vol. 1, p. 263, Hud. Riv. Gr.

Paleopteris, Schimper, being preoccupied by Geinitz, see *Archæopteris*.

acádica, see *Aneimites acádicus*.

hartii, see *Archæopteris hartii*.

Paleoxyris, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 137. [Ety. *palaïos*, ancient; *xyris*, plant.] An inflorescence. Type *P. regularis*. The fossils which have been referred to this genus in the American paleozoic rocks are now referred to *Spirangium*.

appendiculata, see *Spirangium appendiculatum*.

corrugata, see *Spirangium corrugatum*.

prendeli, see *Spirangium prendeli*.

Palmucites oculatus, see *Sigillaria oculata*.

naggethi, see *Trigonocarpum naggethi*.

PECOPTERIS, Brongniart, 1822, Class d. Veg. Foss. in Mem. du Mus. d'Hist. Nat. d. Paris, tom. 8, p. 203. [Ety. *peko*, comb; *ptēris*, fern.] Fronds. bi, tripinnate; pinne long, pinnatifid; pinnules adhering to the rachis by the whole base, often more or less deeply connate, not decurring; borders generally contiguous, or nearly so; secondary veins derived from the medial nerve of the pinnules, simple, bi or trifurcate. Type, *P. longifolia* is the first species mentioned in the Prodr. d. Hist. d. Veg. Foss., and the first mentioned in the Coal Flora of Pa. is *P. unita*, while *P. penniformis* is a representative species. *abbreviata*, Brongniart, 1828, Hist. d. Veg. Foss., p. 337, and Coal Flora of Pa., p. 248, Coal Meas.

acuta, Brongniart, 1828, Hist. d. Veg. Foss., p. 350, and Coal Flora of Pa., p. 241, Coal Meas.

æqualis, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 58, Coal Meas.

alata, see *Sphenopteris alata*.

alata, Schimper, 1869, Pal. Veg., t. 1, p. 531, syn. for *Pseudopecopteris decurrens*.

angustipinna, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 76, Coal Meas. or Permian.

angustissima, Sternberg, 1820, Vers. Darst. Flor. d. Vorw., p. 18, and Coal Flora of Pa., p. 257, Coal Meas.

aquilina, see *Alethopteris aquilina*.

arborescens, Schlotheim, 1820, (Filicites *arborescens*.) Petrefaktenkunde, p. 404, and Coal Flora of Pa., p. 230, Coal Meas.

arguta, Sternberg, 1820, Vers. Darst. Flor. d. Vorw., p. 19, and Coal Flora of Pa., p. 227, Coal Meas.

aspera, Brongniart, 1828, Hist. d. Veg. Foss., p. 330, and Coal Flora of Pa., p. 242, Coal Meas.

aspidioides, Brongniart, 1828, Hist. d. Veg. Foss., p. 311, and Coal Flora of Pa., p. 756, Coal Meas.

asplenioides, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 72, Coal Meas. or Permian.

bucklandi, Brongniart, 1828, Hist. d. Veg. Foss., p. 319, and Coal Flora of Pa., p. 244, Coal Meas.

callosa, see *Pseudopecopteris callosa*.

candolliana, Brongniart, 1828, Hist. d. Veg. Foss., p. 305, Coal Flora of Pa. p. 243, Coal Meas.

carri, Lesquereux, 1884, Coal Flora of Pa., p. 758, Coal Meas.

chærophylloides, see *Sphenopteris chærophylloides*.

cisti, Brongniart, 1828, Hist. d. Veg. Foss., p. 330, and Coal Flora of Pa., p. 243, Coal Meas.

clarki, Lesquereux, 1880, Coal Flora of Pa., p. 261, Coal Meas.

clintoni, Lesquereux, 1880, Coal Flora of Pa., p. 251, Coal Meas.

concinna, Lesquereux, 1854, Bost. Jour. Soc. Nat. Hist., vol. 6, p. 424, and Gen. Sur. Pa., vol. 2, p. 867, Coal Meas. But the name was preoccupied by Presl in 1833.

crenulata, Brongniart, not American. The form sometimes referred to it is *Pseudopecopteris subcrenulata*.

cristata, Gutbier, 1843, Gaea von Sachsen, p. 80, and Coal Flor. of Pa., p. 256, Coal Meas.

cristata, see *Sphenopteris cristata*.

cyathea, Schlotheim, 1820, (Filicites *cyathea*.) Petrefaktenkunde, p. 403, Coal Meas.

decurrens, see *Pseudopecopteris decurrens*. *decurrens*, Dawson, 1862. The name being preoccupied, it was changed to *P. discrepans*.

- densifolia*, Dawson, 1874, Foss. Plants of Canada, p. 56, Devonian.
- dentata*, Brongniart, 1828, Hist. d. Veg. Foss., p. 346, and Coal Flora of Pa., p. 240, Coal Meas.
- distans*, Lesquereux, 1854, Bost. Jour. Soc. Nat. Hist., vol. 6, p. 423, and Geo. Sur. Pa., vol. 2, p. 866, Coal Meas. The name was preoccupied by Rost in 1839.
- dournaisi*, see *Callipteridium dournaisi*.
- dubia*, Sternberg, 1820, Tent. Flor. Primord., p. 19, and Gutbier in Gaea von Sachsen, Coal Meas.
- elegans*, Göppert, 1836, (Polypodites elegans,) Syst. Filic. Foss., p. 344, and Coal Flora of Pa., p. 228, Coal Meas.
- elliptica*, Bunbury, 1846, Quar. Jour. Geo. Soc., vol. 2, p. 82, and Coal Flora of Pa., p. 245, Coal Meas.
- elliptica*, Fontaine & White, 1880, (Goniopteris elliptica,) Perm. or Up. Carb. Flora, p. 83, Coal Meas. or Permian. The name was preoccupied.
- emarginata*, Göppert, 1836, (Diplazites emarginatus,) Syst. Filic. Foss., p. 274, and Coal Flora of Pa., p. 225, Coal Meas.
- erosa*, Gutbier, 1843, Gaea von Sachsen, p. 81, and Coal Flora of Pa., p. 255, Coal Meas.
- flavicans*, Presl, 1833, in Sternberg, Vers. Darst. Flor. d. Vorw., vol. 2, p. 127. Probably not American.
- georgiana*, Lesquereux, 1884, Coal Flora of Pa., p. 759, Coal Meas.
- germari*, Weiss, 1869, (Cyatheites germari,) Foss. Flora d. Jungsten Steink. Form., Up. Carb. Meas. or Permian.
- germari* var. *crassinervis*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 70, Coal Meas. or Permian.
- germari* var. *cuspidata*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 70, Coal Meas. or Permian.
- goniopteroides*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 80, Coal Meas. or Permian.
- halli*, Lesquereux, 1870, (Alethopteris halli,) Geo. Sur. Ill., vol. 4, p. 394, Coal Meas.
- heerana*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 77, Coal Meas. or Permian.
- hemiteloides*, Brongniart, 1828, Hist. d. Veg. Foss., p. 314, Coal Meas.
- heterophylla*, see *Alethopteris heterophylla*.
- imbricata*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 72, Coal Meas. or Permian.
- inclinata*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 80, Coal Meas. or Permian.
- incompleta*, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 868, Coal Meas.
- ingens*, Dawson, 1862, Quar. Jour. Geo. Soc., Lond., vol. 18, p. 322, Devonian.
- lanceolata*, Lesquereux, 1870, (Alethopteris lanceolata,) Geo. Sur. Ill., vol. 4, p. 398, Coal Meas.
- lanceolata*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 79, Coal Meas. or Permian. The name was preoccupied; beside, it is probably a syn. for *P. unita*.
- latifolia*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 79, Coal Meas. or Permian.
- lepidorachis*, Brongniart, 1828, Hist. d. Veg. Foss., p. 313, Coal Meas.
- lescuriana*, n. sp. Coal Meas. Proposed instead of *P. obsoleta*, Lesquereux, 1884, Coal Flora of Pa., p. 758, which name was preoccupied.
- longifolia*, Brongniart, 1828, Hist. d. Veg. Foss., p. 273, and Coal Flora of Pa., p. 226, Coal Meas.
- loschi*, Brongniart, 1828, Hist. d. Veg. Foss., p. 355, Coal Meas.
- lyratifolia*, Göppert, 1841, (Sphenopteris lyratifolia,) Die Gattungen d. Foss. Pflanzen, p. 71, and Coal Flora of Pa., p. 259, Coal Meas.
- mantelli*, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 57, Coal Meas.
- marginata*, see *Alethopteris marginata*.
- merianopteroides*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 78, Coal Meas. or Permian.
- microphylla*, Brongniart, 1828, Hist. d. Veg. Foss., p. 340, and Coal Flora of Pa., p. 263, Coal Meas.
- milleri*, Harlan, 1835, Trans. Geo. Soc. Pa., Coal Meas.
- miltoni*, Artis, 1825, (Filicites miltoni,) Anted. Phytol. pl. 4, and Coal Flora of Pa., p. 247, Coal Meas.
- muricata*, see *Pseudopecopteris muricata*.
- murrayana*, Brongniart, as identified by Lesquereux in Geo. Sur. Ill., vol. 2, p. 443, see *Sphenopteris pseudo-murrayana*.
- nervosa*, see *Pseudopecopteris nervosa*.
- newberryana*, Fontaine & White, 1880, (Goniopteris newberryana,) Perm. or Up. Carb. Flora, p. 84, Coal Meas. or Permian.
- newberryi*, see *Pseudopecopteris newberryi*.
- nodosa*, Göppert, 1836, (Aspidites nodosus,) Systema Filicum Fossilium, p. 372, and Coal Flora of Pa., p. 233, Coal Meas.
- notata*, Lesquereux, 1854, Bost. Jour. Soc. Nat. Hist., vol. 6, p. 424, and Geo. Sur. Pa., vol. 2, p. 866, Coal Meas.
- oblonga*, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 83, Coal Meas. or Permian.
- obsoleta*, Harlan, 1835, Trans. Geo. Soc. Pa., Coal Meas.
- obsoleta*, Lesquereux, 1884, Coal Flora of Pa., p. 758. The name was preoccupied. See *P. lescuriana*.
- oreopteroides*, Schlotheim, 1820, (Filicites oreopteridius,) Petrefaktenkunde, p. 407, and Coal Flora of Pa., p. 238, Coal Meas.
- ornata*, Lesquereux, 1884, Coal Flora of Pa., p. 760, Coal Meas.

obscura, see *Alethopteris obscura*.

ovata, Veg.
ovoide
Flora
mian
pachyl
Perm
Meas
pennif
penn
Mem
tom:
239, C
platyne
Perm
Meas.
platytr
Veg.
p. 232
pluckene
cneti.
plumosa
Anted
polymor
Hist. d
preciosa
Devon
pteroides
pteridif
and C
Meas.
pusilla, se
quadratif
Flora o
rarinervi
Perm.
Meas. o
rigida, D
vol. 8,
Meas.
robusta, I
Pa., p.
rotundifo
Perm. o
Meas. o
rotundilob
Perm. o
Meas. o
schimper
Perm. o
Meas. o
schimper
of Pa., p
sertii, see
serpillifoli
of Pa., p
serrula, L
serrula,
Coal Me
serrulata,
Devonia
sheaferti, see
sillimanii, se
sinuata, see
solida, Les
solida,
Coal Me
squamosa,
Ill., vol.

ovata, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss. p. 58, Coal Meas.
 ovoides, Fontaine & White, 1880, Coal Flora of Pa., p. 79, Coal Meas. or Permian.
 pachypteroides, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 76, Coal Meas. or Permian.
 penniformis, Brongniart, 1822, (Filicites pennaeformis,) Class des Veg. Foss., in Mem. du Mus. d'Hist. Nat. de Paris, tom. 8, p. 203, and Coal Flora of Pa., p. 239, Coal Meas.
 platynervis, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 73, Coal Meas. or Permian.
 platyrachis, Brongniart, 1828, Hist. d. Veg. Foss. p. 312, and Coal Flora of Pa., p. 232, Coal Meas.
pluckeneti, see *Pseudopteris pluckenetii*.
 plumosa, Artis, 1825, (Filicites plumosus,) Anted. Phytol. pl. 17, Coal Meas.
 polymorpha, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 56, Coal Meas.
 preciosus, Hartt, 1868, Acad. Geol., p. 553, Devonian.
 pteroides, Schlotheim, 1820, (Filicites pteridius,) Petrefaktenkunde, p. 406, and Coal Flora of Pa., p. 249, Coal Meas.
pusilla, see *Pseudopteris pusilla*.
 quadratifolia, Lesquereux, 1880, Coal Flora of Pa., p. 234, Coal Meas.
 rarinervis, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 71, Coal Meas. or Permian.
 rigida, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 485, Coal Meas.
 robusta, Lesquereux, 1880, Coal Flora of Pa., p. 229, Coal Meas.
 rotundifolia, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 73, Coal Meas. or Permian.
 rotundiloba, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 74, Coal Meas. or Permian.
 schimperana, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 75, Coal Meas. or Permian.
 schimperi, Lesquereux, 1884, Coal Flora of Pa., p. 835, Coal Meas.
serlii, see *Alethopteris serlii*.
 serpillifolia, Lesquereux, 1880, Coal Flora of Pa., p. 237, Coal Meas.
 serrula, Lesquereux, 1858, (Alethopteris serrula,) Geo. Sur. Ill., vol. 2, p. 865, Coal Meas.
 serrulata, Hart, 1868, Acad. Geol., p. 553, Devonian.
sheaferi, see *Pseudopteris sheaferi*.
sillimani, see *Pseudopteris sillimani*.
sinuata, see *Callipteridium sinuatum*.
 solida, Lesquereux, 1870, (Alethopteris solida,) Geo. Sur. Ill., vol. 4, p. 397, Coal Meas.
 squamosa, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 400, Coal Meas.

stellata, Lesquereux, 1866, (Alethopteris stellata,) Geo. Sur. Ill., vol. 2, p. 440, Low. Coal Meas.
 strongi, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 399, Coal Meas.
 subfalcata, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 70, Coal Meas. or Permian.
 taniopteroides, Bunbury, 1847, Quar. Jour. Geo. Soc., vol. 3, p. 428, Coal Meas.
 tenuinervis, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 77, Coal Meas. or Permian.
 tenuis, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 57, Coal Meas.
 unita, Brongniart, 1828, Hist. d. Veg. Foss., p. 342, Coal Meas.
urophylla, see *Alethopteris urophylla*.
 vellutina, Lesquereux, 1854, Bost. Jour. Soc. Nat. Hist., vol. 6, p. 423, and Geo. Sur. Pa., vol. 2, p. 866, Coal Meas.
 venulosa, Lesquereux, 1880, Coal Flora of Pa., p. 230, Coal Meas.
 vestita, Lesquereux, 1880, Coal Flora of Pa., p. 252, Coal Meas.
 villosa, Brongniart, 1828, Hist. d. Veg. Foss., p. 316, and Coal Flora of Pa., p. 253, Coal Meas.

PHYLLOPTERIS, Brongniart, 1849, Table d. Gen. d. Veget. Foss., pp. 22, 103. [Ety. *phyllon*, leaf; *pteris*, fern.] Pinnate,

FIG. 61.—Phyllopteris antiqua.

pinnules oblong or lanceolate, pointed, attached by the middle of the base; midrib strong, extending to the point, giving off oblique nerves, which have obliquely pinnate nervules not anastomosing. Type *P. phillipsi*.
antiqua, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 484, Coal Meas.
Physophycus, Schimper, 1869. Syn. for *Taonurus*.
marginatus, see *Taonurus marginatus*.
Phytolithus, Martin, 1809, Petrificata Derbiensis. [Ety. *phyton*, plant; *lithos*, stone.] Applied indiscriminately to fossil wood.
cancellatus, syn. for *Lepidodendron cancellatum*.
notatus, see *Sigillaria notata*.
tessellatus, see *Sigillaria tessellata*.
transversus, see *Sternbergia transversa*.

FIG. 60. Pecopteris unita.

PHYTOPHYS, Hall, 1847, Pal. N. Y., vol. 1, p. 38. [Ety. *phylon*, plant; *opsis*, resemblance.]

FIG. 62.—*Phytopsis tubulosa*.

Stems cylindrical or subcylindrical, straight or flexuous, erect or procumbent, branched; branches diverging and anastomosing; structure cellular, consisting apparently of thin laminae, with transverse divisions, or having a reticulated structure. This structure is too obscure for satisfactory determination. Type *P. tubulosa*.

cellulosa, Hall, 1847, Pal. N. Y., vol. 1, p. 39, Birdseye Gr. *tubulosa*, Hall, 1847, Pal. N. Y., vol. 1, p. 38, Birdseye Gr. PINNULARIA, Lindley & Hutton, 1835, Foss. Flora, vol. 2, p. 81. [Ety. *pinna*, a feather.]

Roots or rootlets divided in filaments of variable length and thickness, and generally possessing few definable characters. Type *P. capillacea*. *calamitarum*, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 878, Coal Meas. *capillacea*, Lindley & Hutton, 1835, Foss. Flora, vol. 2, p. 81, Coal Meas. *confervoides*, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 878, Coal Meas. *crassa*, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 480, Coal Meas. *dispalans*, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 312, Devonian. *elongata*, Dawson, 1871, Foss. Plants Can., p. 33, Devonian. *ficoides*, Lesquereux, 1868, Geo. Sur. Pa., vol. 2, p. 878, Coal Meas. *horizontalis*, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 878, Coal Meas. *nodosa*, Dawson, 1871, Foss. Plants Can., p. 33, Devonian. *palmatifida*, Lesquereux, 1860, (Rhizolites palmatifidus.) Geo. Sur. Ark., vol. 2, p. 313, Coal Meas. *pinnata*, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 878, Coal Meas.

FIG. 63.—*Pinnularia ramosissima*.

ramosissima, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 480, Coal Meas.

PLUMALINA, Hall, 1858, Can. Nat. and Geo., vol. 3, p. 175. [Ety. *pluma*, a small feather.] Simple fronds, with linear pinnules diverging from each side, in the same plane, and more or less ascending. It is a peculiar plant, described, originally, as a Graptolite, to which opinion Prof. Hall still adheres. On the other hand, Prof. Dawson claims the characters prove it is a vegetable, and in this he is supported by the fact that all Graptolites had become extinct, as shown by their absence in several groups of rocks before the appearance of this form. Type *P. plumaria*.

densa, Hall, 1879, 30th Rep. N. Y. St. Mus. Nat. Hist., pl. 4, fig. 6, Ham. Gr. *gracilis*, Shumard, 1855, (Filicites gracilis.) Geo. Rep. Mo., p. 208, Waverly Gr. in Lithographic limestone. *linearis*, Lesquereux, 1880, (Trochophyllum lineare.) Coal Flora of Pa., p. 64, Waverly Gr. *plumaria*, Hall, 1843, (Filicites?) Geo. Rep. p. 273, and 4th Dist. N. Y., Can. Nat. and Geo., vol. 3, p. 175, Chemung Gr.

FIG. 64.

plumula, Dawson, 1873, (Lycopodites plumula.) Rep. Foss. plants Low. Carb. and Millstone Grit p. 24, Subcarboniferous. *Polyporites*, Lindley & Hutton, 1833, Foss. Flora, vol. 1, p. 181. This genus was founded upon a fish-scale, and the form referred to it in Geo. Sur. Pa., vol. 2, p. 847, is quite a different thing.

Protoblechnum holdeni.

FIG. 65.

Polysporia, Newberry, syn. for *Lepidostrobus mirabilis*, see *Lepidostrobus mirabilis*.

PROTOBLECHNUM, Lesquereux, 1880, Coal Flora of Pa., p. 188. [Ety. *protos*, first; *blechnum*, a small fern frond.]

Ble
nat
pin
acu
the
gen
late
ders
holde
hold
Mea
PROTOST
Phil
stigm
rhon
with
Not
sigill
sigillar
Phil
PROTOST
Geo.
first;
the
istic
bran
of gr
pleur
cylind
series
pores
specie
with
P. log
logani,
Soc.,
the o
Amer
Carru
and ha
PSARONIUS
ziehur
Stems
adven
super
trunks
woody
branch
sels ei
plicate
tissue.
erianus,
p. 58, l
textilis, l
p. 59, l
PSEUDOPECE
Flora
false;
rachis
ing br
cate an
pinnate
forked;
the bas
tuse or
right an
borderi
veins ob
lowest

- Blechnum*, a genus.] Fronds large, pinnate; rachis thick, scaly toward the base; pinnae long, narrow linear-lanceolate, acuminate, entire, enlarged at base on the lower side to a decurring auricle, generally free; medial nerve percurrent; lateral veins open, curving to the borders, forking twice. Type *P. holdeni*. holdeni, Andrews, 1875, (*Alethopteris holdeni*), Ohio Pal., vol. 2, 420, Coal Meas.
- PROTOSTIGMA**, Lesquereux, 1877, Proc. Am. Phil. Soc., p. 169. [Ety. *protos*, first; *stigma*, a brand or dot.] Stems with rhomboidal scars as in *Sigillaria*, but without vascular scars in the middle. Not a land-plant, but a fucoid. Type *P. sigillarioides*.
- sigillarioides*, Lesquereux, 1877, Proc. Am. Phil. Soc., p. 169, Hud. Riv. Gr.
- PROTAXITES** Dawson, 1859, Quar. Jour., Geo. Soc., vol. 15, p. 484. [Ety. *protos*, first; *taxis*, yew-tree; so named from the spirally marked cells characteristic of the genus *Taxites*.] Woody and branching trunks, with concentric rings of growth and medullary rays; cells of pleurenychyma not in regular lines, cylindrical, thick-walled, with a double series of spiral fibers; discs or bordered pores few, circular and indistinct. The specimens found are usually silicified, with the bark in a coaly state. Type *P. logani*.
- logani*, Dawson, 1859, Quar. Jour. Geo. Soc., vol. 15, p. 484, Devonian. This is the oldest known exogenous tree in America, according to Dawson, but Carruthers says it is a huge sea-weed and has named it *Nematophycus logani*.
- PSARONIVUS**, Cotta, 1832, Dendrol in Beziehung, p. 27. [Ety. *psaros*, speckled.] Stems of tree-ferns, covered below by adventive roots, increasing by their superposition the conical base of the trunks; cortex thick, parenchymatous; woody cylinder, subdivided into branches composed of fascicles of vessels either half cylindrical or diversely plicate, immersed in cellular medullary tissue. Type *P. helmintholithus*.
- helmintholithus*, Dawson, 1871, Foss. Plants Can., p. 58, Ham. Gr.
- textilis*, Dawson, 1871, Foss. Plants Can., p. 59, Ham. Gr.
- PSEUDOPECOPTERIS**, Lesquereux, 1880, Coal Flora of Pa., p. 189. [Ety. *pseudo*, false; *Pecopteris*, a genus.] Primary rachis forking near the base in diverging branches of equal size, or divaricate and dichotomous; branches polypinnate, ultimate divisions sometimes forked; pinnules connate or separated to the base, of various shape, oblong-obtuse or ovate-lanceolate, oblique or in right angle, decurring to the rachis and bordering it by a narrow wing; lateral veins oblique, generally forking once, the lowest pair twice. Type *P. mazonana*.
- abbreviata*, Lesquereux, 1854, (*Sphenopteris abbreviata*), Bost. Jour. Soc. Nat. Hist., vol. 6, p. 419, and Geo. Sur. Pa., vol. 2, p. 861, Coal Meas.
- acuta*, Brongniart, 1828, (*Sphenopteris acuta*), Hist. d. Veg. Foss., p. 207, and Coal Flora of Pa., p. 215, Coal Meas.
- anceps*, Lesquereux, 1880, Coal Flora of Pa., p. 207, Coal Meas.
- andreaana*, Roehl, 1868, (*Sphenopteris andreaana*), Fossile Flora der Steinkohlenformation Westphalens, p. 62, and Coal Flora of Pa., p. 754, Coal Meas.
- callosa*, Lesquereux, 1866, (*Pecopteris callosa*), Geo. Sur. Ill., vol. 2, p. 442, Low. Coal Meas.
- cordato-ovata*, Weiss, 1869, (*Neuropteris cordato-ovata*), Foss. Flor. d. jungst. Steink. form., p. 28, and Coal Flora of Pa., p. 205, Coal Meas.
- decipiens*, Lesquereux, 1854, (*Sphenopteris decipiens*), Bost. Jour. Soc. Nat. Hist., vol. 6, p. 420, and Geo. Sur. Pa., vol. 2, p. 862, Coal Meas.
- decurrens*, Lesquereux, 1854, (*Pecopteris decurrens*), Bost. Jour. Soc. Nat. Hist., vol. 6, p. 424, and Geo. Sur. Pa., vol. 2, p. 867, Coal Meas.
- denudata*, Lesquereux, 1880, Coal Flora of Pa., p. 212, Coal Meas.
- dimorpha*, Lesquereux, 1880, Coal Flora of Pa., p. 201, Coal Meas.
- glandulosa*, Lesquereux, 1854, (*Sphenopteris glandulosa*), Bost. Jour. Soc. Nat. Hist., vol. 6, p. 420, and Geo. Sur. Pa., vol. 2, p. 862, Coal Meas.
- hispida*, Lesquereux, 1884, Coal Flora of Pa., p. 755, Coal Meas.
- hymenophylloides*, Lesquereux, 1870, (*Alethopteris hymenophylloides*), Geo. Sur. Ill., vol. 4, p. 393, Coal Meas.
- irregularis*, Sternberg, 1833, (*Sphenopteris irregularis*), Vers. Geog. Darst. Flor. d. Vorw., vol. 2, p. 68, Coal Meas.
- latifolia*, Brongniart, 1828, (*Sphenopteris latifolia*), Hist. d. Veg. Foss., p. 205, and Coal Flora of Pa., p. 215, Coal Meas.
- macilentia*, Lindley & Hutton, 1835, Foss. Flora, vol. 2, pl. 151, and Coal Flora of Pa., p. 219, Coal Meas.
- mazonana*, Lesquereux, 1870, (*Alethopteris mazonana*), Geo. Sur. Ill., vol. 4, p. 391, Low. Coal Meas.
- muricata*, Brongniart, 1828, (*Pecopteris muricata*), Hist. d. Veg. Foss., p. 352, and Coal Flora of Pa., p. 203, Coal Meas.
- nervosa*, Brongniart, 1828, (*Pseudopecopteris nervosa*), Hist. d. Veg. Foss., p. 207, and Coal Flora of Pa., p. 187, Coal Meas.
- newberryi*, Lesquereux, 1854, (*Sphenopteris newberryi*), Bost. Jour. Soc. Nat.

FIG. 66.

Pseudopecopteris mazonana.

Hist., vol. 6, p. 420, and Geo. Sur. Pa., vol. 2, p. 862, Coal Meas.
nummularia, Gutbier, 1842, Abdrucke u. Verst. d. Zwick. Schwarzk. u. Seiner Umgebungen, p. 43, and Coal Flora of Pa., p. 752, Coal Meas.
obtusiloba, Brongniart, 1828, (Sphenopteris *obtusiloba*.) Hist. d. Veg. Foss., p. 204, and Coal Flora of Pa., p. 753, Coal Meas.
pluckeneti, Schlottheim, 1820, (Filicites *pluckeneti*.) Petrefaktenkunde, p. 410, and Coal Flora of Pa., p. 199, Coal Meas.
polyphylla, Lindley & Hutton, 1835, (Sphenopteris *polyphylla*.) Foss. Flora, vol. 2, pl. 147, and Coal Flora of Pa., p. 218, Coal Meas.
pumilla, Lesquereux, 1854, (Pecopteris *pumilla*.) Bost. Jour. Soc. Nat. Hist., vol. 6, p. 424, and Geo. Sur. Pa., vol. 2, p. 866, Coal Meas.
sheaferi, Lesquereux, 1858, (Pecopteris *sheaferi*.) Catal. Potts. Ass'n, p. 11, and Coal Flora of Pa., p. 194, Coal Meas.
sillimani, Brongniart, 1828, (Pecopteris *sillimani*.) Hist. d. Veg. Foss., p. 353, and Coal Flora of Pa., p. 206, Coal Meas.
speciosa, Lesquereux, 1880, Coal Flora of Pa., p. 216, Coal Meas.
spinulosa, Lesquereux, 1870, (Alethopteris *spinulosa*.) Geo. Sur. Ill., vol. 4, p. 396, Coal Meas.
subcrenulata, Lesquereux, 1880, Coal Flora of Pa., p. 193, Coal Meas.
subnervosa, Roemer, F. A., 1860, Paleontographica, vol. 9, p. 192, and Coal Flora of Pa., p. 198, Coal Meas.
trifoliata, Artis, 1825, (Filicites *trifoliatus*.) Anted. Phytol., pl. 2, and Coal Flora of Pa., p. 217, Coal Meas.
virginiana, Meek, 1875, (Cyclopteris *virginiana*.) Bull. Phil. Soc. of Washington, p. 18, and Coal Flora of Pa., p. 217, Waverly Gr.
PSILOPHYTON, Dawson, 1859, Quar. Jour. Geo. Soc., vol. 15, p. 478. [Ety. *psilon*, smooth; *phyton*, stem.] Stems dichotomous; young branches carinate; rhizomes cylindrical, villous or scaly; marked with round scars, points of attachment of cylindrical rootlets; leaves in spiral order, small or rudimentary, acicular, squarrose, open; fructifications in small, naked sporanges, spindle-shaped or clavate, axillary, or in pairs at the extremity of the branches. Type *P. princeps*.
cornutum, Lesquereux, 1877, Proc. Am. Phil. Soc., p. 165, Low. Held. Gr.
elegans, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 315, Devonian.
glabrum, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 315, Devonian.
gracillimum, see *Dendrograptus gracillimus*.
princeps, Dawson, 1859, Quar. Jour. Geo. Soc., vol. 15, p. 479, Upper Silurian and

Devonian. This is the oldest known plant in America. It is supposed to have grown in a marsh.

FIG. 67.—*Psilophyton princeps*.

princeps var. *ornatum*, Dawson, 1871, Foss. Plants, p. 38, Devonian.
robustum, Dawson, 1859, Quar. Jour. Geo. Soc., vol. 15, p. 479, Devonian.
Ptilocarpus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 493, Syn. for *Cardiocarpon*.
bicornutus, see *Cardiocarpon bicornutum*.
Ptilophyton, Dawson, 1878, Scottish Devonian Plants in Can. Nat., vol. 8. This is founded upon *Lycopodites vanuxemi* as the type, which is the same as *Plumalina plumula*, and falls therefore as a synonym.
gracile, see *Plumalina gracilis*.
lineare, Lesquereux, see *Plumalina linearis*.
plumula, see *Lycopodites plumula*.
vanuxemi, syn. for *Plumalina plumula*.
RHABDOCARPUS, Goeppert & Berger, 1848, De Fruct. et Sem., p. 20. [Ety. *rhabdos*, stria; *karpus*, fruit.] Seeds ovate or oblong, costate or striate, acute or acuminate, surrounded by a putamen sometimes deficient. Type *R. tunicatus*.
abnormalis, Lesquereux, 1884, Coal Flora of Pa., p. 818, Coal Meas.
acuminatus, Newberry, 1873, Ohio Pal., vol. 1, p. 378, Coal Meas.
amygdaliformis, Goeppert & Berger, 1848, De Fruct. et Sem., p. 21, Coal Meas.
apiculatus, Newberry, 1873, Ohio Pal., vol. 1, p. 377, Coal Meas.
arcuatus, Lesquereux, 1861, Geo. Sur. Ky., vol. 4, p. 434, Coal Meas.
beinertianus, Goeppert & Berger, 1848, De Fruct. et Sem., p. 20, and Coal Flora of Pa., p. 844, Coal Meas.
bockshianus, Goeppert & Berger, 1848, De Fruct. et Sem., p. 21, and Coal Flora of Pa., p. 844, Coal Meas.
carinatus, Newberry, 1873, Ohio Pal., vol. 1, p. 376, Coal Meas.
clavatus, Sternberg, 1820, (Carpolithes *clavatus*.) Vers. Darst. Flora der Vor-

FIG. 68.—*Rhabdocarpus carinatus*.

- welt, and Coal Flora of Pa., p. 581, Coal Meas.
- cornutus, Lesquereux, 1880, Coal Flora of Pa., p. 583, Coal Meas.
- costatus, Newberry, 1873, Ohio Pal., vol. 1, p. 378, Coal Meas., syn. ? for *R. acuminatus*.
- danai, Foster, 1854, Ann. of Sci., vol. 1, p. 129, and Ohio Pal., vol. 1, p. 376, Coal Meas.
- emarginatus, Lesquereux, 1884, Coal Flora of Pa., p. 818, Coal Meas.
- howardi, Lesquereux, 1880, Coal Flora of Pa., p. 575, Coal Meas.
- inflatus, Lesquereux, 1884, Coal Flora of Pa., p. 815, Coal Meas.
- insignis, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 478, Coal Meas.
- insignis*, Lesquereux, 1880, Coal Flora of Pa., p. 575. The name being preoccupied, it has been called *R. lescurianus*.
- jacksonensis, Lesquereux, 1866, (Carpolithes jacksonensis.) Geo. Sur. Ill., vol. 2, p. 461, Low. Coal Meas.
- kevis, Newberry, 1873, Ohio Pal., vol. 1, p. 377, Coal Meas.
- laticostatus, Lesquereux, 1884, Coal Flora of Pa., p. 815, Coal Meas.
- lescurianus, S. A. Miller, 1883, 2d. Ed. Am. Pal. Foss., p. 256, Coal Meas. Proposed instead of *R. insignis*, Lesquereux, which was preoccupied.
- mammillatus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 461, Coal Meas.
- minutus, Lesquereux, 1860, Geo. Sur. Ark., vol. 2, p. 313, Coal Meas.
- multistriatus, Presl, 1833, (Carpolithes multistriatus.) in Sternberg's Flor. d. Vorw., vol. 2, p. 208, and Coal Flora of Pa., p. 578, Coal Meas.
- oblongus, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 98, Coal Meas. or Permian.
- pachytesta, Lesquereux, 1884, Coal Flora of Pa., p. 816, Coal Meas.
- platimarginatus, Lesquereux, 1860, (Carpolithes platimarginatus.) Geo. Sur. Ark., vol. 2, p. 312, Low. Coal Meas.
- subglobosus, Lesquereux, 1884, Coal Flora of Pa., p. 817, Coal Meas.
- tenax, Lesquereux, 1884, Coal Flora of Pa., p. 818, Coal Meas.
- venosus, Sternberg, as identified by Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 870, Coal Meas. Not noticed in Coal Flora of Pa., and probably not American.
- RHACHOPTERIS**, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 323. [Ety. *rachis*, a stalk; *pteris*, fern.] Detached leaf-stalks of ferns; stipes half an inch wide or less; unevenly striate, giving off opposite branches, which are abruptly broken off at short distances from the stipe. Type *R. pinnata*.
- affinis, Lesquereux, 1870, (Stigmarioides affinis.) Geo. Sur. Ill., vol. 4, p. 455, Coal Meas.
- cyclopteroides, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 323, Catskill Gr.
- gigantea, Dawson, 1871, Foss. Plants Can., p. 57, Ham. Gr.
- palmata, Dawson, 1871, Foss. Plants of Canada, p. 57, Ham. Gr.
- pinnata, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 323, Catskill Gr.
- punctata, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 323, Catskill Gr.
- selago, (Lesquereux, 1870, (Stigmarioides selago.) Geo. Sur. Ill., vol. 4, p. 456, Coal Meas.
- squamosa, Lesquereux, 1884, Coal Flora of Pa., p. 838, Coal Meas.
- striata, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 323, Chemung Gr.
- tenuistriata, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 323, Ham. Gr.
- RHACOPHYLLUM**, Schimper, 1869, Paleontologie Vegetale, vol. 1, p. 684. [Ety. *rakos*, rugged; *phylon*, leaf.] Fronds either flabelliform, many times subdivided or pinnate, irregularly pinnatifid, bipinnatifid; rachis flat, often much dilated, scarcely thicker than the foliaceous lamina, which is very variable in the size and the mode of its divisions; veins numerous, more or less indistinct, following the rachis in parallel bundles, dichotomous in the foliaceous divisions. Type, *R. flabellatum*.
- adnascens, Lindley & Hutton, 1835, (Schizopteris adnascens.) Foss. Flora, vol. 2, p. 57, and Coal Flora of Pa., p. 321, Coal Meas.
- affine, Lesquereux, 1858, (Pachyphyllum affine.) Geo. Sur. Pa., vol. 2, p. 863, Coal Meas.
- arborescens, Lesquereux, 1870, (Hymenophyllites arborescens.) Geo. Sur. Ill., vol. 4, p. 415, Coal Meas.
- browni, Dawson, 1861, (Cyclopteris browni.) Quar. Jour. Geo. Soc., vol. 17, p. 32, Portage Gr.
- clarki, Lesquereux, 1866, (Hymenophyllites clarki.) Geo. Sur. Ill., vol. 2, p. 438, Coal Meas.
- corallinum, Lesquereux, 1880, Coal Flora of Pa., p. 317, Coal Meas. Misspelled *corratum* in the text.
- cornutum, Lesquereux, 1880, Coal Flora of Pa., p. 317, Coal Meas.
- expansum, Lesquereux, 1880, Coal Flora of Pa., p. 313, Coal Meas.
- filiciforme, Gutbier, 1842, (Fucoides filiciformis.) Abdr. u. Verst. d. Zwick. Schwarzk. u. sein. Umg., p. 11, and Coal Flora of Pa., p. 316, Coal Meas.
- filiforme, Gutbier, 1842, (Fucoides filiformis.) Abdr. u. Verst. d. Zwick. Schwarzk. u. sein. Umg., p. 12, and Coal Flora of Pa., p. 838, Coal Meas.
- fimbriatum, Lesquereux, 1858, (Pachyphyllum fimbriatum.) Geo. Sur. Pa., vol. 2, p. 863, Coal Meas.
- flabellatum, Sternberg, 1833, (Aphlebia flabellata.) Flor. d. Vorw., vol. 2, p. 112, and Coal Flora of Pa., p. 311, Coal Meas.

fucoides, Lesquereux, 1880, Coal Flora of Pa., p. 325, Coal Meas.
 hamulosum, Lesquereux, 1880, Coal Flora of Pa., p. 321, Coal Meas.
 hirsutum, Lesquereux, 1858, (Pachyphyllum hirsutum,) Geo. Sur. Pa., vol. 2, p. 863, Coal Meas.
 inflatum, Lesquereux, 1870, (Hymenophyllites inflatus,) Geo. Sur. Ill., vol. 4, p. 414, Coal Meas.
 irregulare, Germar, 1844, (Aphlebia irregularis,) Verst. d. Steink. v. Wettin u. Löbejun, p. 57, and Coal Flora of Pa., p. 326, Coal Meas.
 laceratum, Lesquereux, 1858, (Pachyphyllum laceratum,) Geo. Sur. Pa., vol. 2, p. 863, Coal Meas.
 laciniatum, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 94, Coal Meas. or Permian.

FIG. 69.—*Rhacophyllum lactuca*.

lactuca, Sternberg, 1833, (Schizopteris lactuca,) Flor. d. Vorw. vol. 2, p. 112, and Coal Flora of Pa., p. 315, Coal Meas.
 membranaceum, Lesquereux, 1880, Coal Flora of Pa., p. 312, Coal Meas.
 molle, Lesquereux, 1870, (Hymenophyllites mollis,) Geo. Sur. Ill., vol. 4, p. 418, Coal Meas.
 scolopendrites, Lesquereux, 1858, (Scolopendrites dentatus,) Geo. Sur. Pa., vol. 2, p. 868, Coal Meas.
 spinosum, Lesquereux, 1880, Coal Flora of Pa., p. 320, Coal Meas.
 strongi, Lesquereux, 1870, (Hymenophyllites strongi,) Geo. Sur. Ill., vol. 4, p. 417, Coal Meas.
 thalliforme, Lesquereux, 1870, (Hymenophyllites thalliformis,) Geo. Sur. Ill., vol. 4, p. 417, Coal Meas.
 trichodeum, Lesquereux, 1880, Coal Flora of Pa., p. 322, Coal Meas.
 truncatum, Lesquereux, 1880, Coal Flora of Pa., p. 311, Coal Meas.
Rhizolites, F. Braun, 1847, in Flora, etc. [Ety. *rhiza*, root; *lithos*, stone.]
palmatifidus, see *Pinnularia palmatifidus*.
Rhizomopteris, Schimper, 1869, Traité de Paléontologie Végétale, vol. 1, p. 699. [Sig. the rhizomas of ferns.] This genus, as the name indicates, comprehends the

rhizomas of ferns. Type, *R. lycopodioides*. Some of the species of *Lycodites* as *L. uncinatus* have been referred to it.

RHIZOMORPHA, Roth, as identified by Lesquereux, Coal Flora of Pa., p. 3. [Ety. *rhiza*, root; *morpha*, form.] Fungous filaments of hard substance, disposed in branches abnormally divided, and often anastomosing; generally living under the decaying bark of trees.

sigillariae, Lesquereux, 1877, Proc. Am. Phil. Soc., p. 174, and Coal Flora of Pa., p. 3, Coal Meas.

Rotularia longifolia, see *Sphenophyllum longifolium*.

RUSOPHYCUS, Hall, 1852, Pal. N. Y., vol. 2, p. 23. [Ety. *rusos*, rugose; *phykos*, seaplant.] Simple or branched stems, transversely wrinkled, and often possessing a central longitudinal depression. Type *R. clavatum*.

asperum, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 25, Utica State.

bilobatum, Vanuxem, 1842, (Fucoides bilobatus,) Geo. Rep. N. Y., p. 79, Hud. Riv. and Clinton Gr.

clavatum, Hall, 1852, Pal. N. Y., vol. 2, p. 23, Clinton Gr.
grevillense, Billings, 1862, Pal. Foss., vol. 1, p. 101, Chazy Gr.

pudicum, Hall, 1852, Pal. N. Y., vol. 2, p. 24, Hud. Riv. and Clinton Gr.

subangulatum, Hall, 1852, Pal. N. Y., vol. 2, p. 23, Clinton Gr.

Sagenaria veltheimiana, see *Lepidodendron veltheimianum*.

SAPORTEA, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 99. [Ety. proper name.] Leaves simple, subreniform, flabellate or suborbicular, cuneate, bordered at the base with a woody rim, terminal margin incised; petiole long, slender, and grooved on the upper surface; nerves parting flabellately from the summit of the petiole and the woody basal margin, all passing into the laminae; leaf substance thin. Type *S. grandifolia*.

grandifolia, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 101, Coal Meas. or Permian.

salisburyoides, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 102, Coal Meas. or Permian.

Schizopteris, Brongniart, 1828, Prodr. d. Hist. d. Vég. Foss., p. 63. [Ety. *schizo*, I cleave; *pteris*, fern.] Frond lacinate,

FIG. 70.—*Rusophycus bilobatum*.

SCH.—SCH.
 or c
 ions
 mar
 bran
 the
 adnasc
 luctuca
 Schutzia
 161.
 singl
 alter
 biles
 pour
 point
 S. an
 bractea
 Scolopend
 Pa.,
 done
 dentat
 drites
 Selaginites
 Veg.
 leave
 some
 leavin
 paten
 synon
 cavifol
 crassus
 vol. 2
 ifolius
 formosus
 p. 176
 a crus
 uncinatu
 SIGILLARIA
 Foss.
 Paris,
 seal;
 leaves
 large,
 apex,
 series,
 spiral
 or less
 shape,
 nate,
 boidal
 simple
 others
 shape.
 carinat
 nerve.
 acumina
 land A
 of Pa.
 alternan
 vol. 2,
 alveolari
 dron
 Geogn
 monde
 Mens.
 angusta,
 Foss.;
 approxim
 Perm.
 Meas.

- or cut in linear erect or curved divisions, sometimes enlarged at the top, marked with thin parallel veins without branching, being split in fascicles with the divisions. Type *S. anomala*.
adnascens, see *Rhacophyllum adnascens*.
lactuca, see *Rhacophyllum lactuca*.
Schultzia, Göppert, 1848, Permian Flora, p. 161. [Ety. proper name.] Stems either single or branching, bearing on short alternate pedicels small cones or strobiles of an ovate, truncate form, a compound of imbricate, broadly linear pointed scales, united at the base. Type *S. anomala*.
bracteata, see *Cordianthus bracteatus*.
Scolopendrites, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 868. This name is abandoned.
dentatus, see *Rhacophyllum scolopendrites*.
Selaginites Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 84. Stems dichotomous; leaves small, numerous, imbricated, sometimes enlarged at the base, scarcely leaving any visible scars. Type *S. patens*. The genus is regarded as synonymous with *Lycopodites*.
cavifolius, see *Lycopodites cavifolius*.
crassus, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 446, syn. for *Lycopodites cavifolius*.
formosus, Dawson, 1861, Can. Nat., vol. 6, p. 176. Not a plant, but a fragment of a crustacean.
uncinatus, see *Lycopodites uncinatus*.
SIGILLARIA, Brongniart, 1822, Class. des Veg. Foss. in Mem. du Mus. d'Hist. Nat. de Paris, tom. 8, p. 203. [Ety. *sigillum*, a seal; from the seal-like scars of fallen leaves stamped upon the bark.] Trunks large, simple or dichotomous near the apex, marked by leaf-scars in vertical series, separated by furrows or placed in spiral order, either contiguous or more or less distant, very variable in size and shape, round, oval, truncate, or emarginate, hexagonal, transversely rhomboidal, with three vascular scars, one simple, medial, punctiform, the two others lateral of semi-lunar or linear shape. Leaves linear, long, triplicate, carinate, or plane, with a distinct medial nerve. Type *S. punctata*.
acuminata, Newberry, 1874, Proc. Cleveland Acad. Sci., p. 164, and Coal Flora of Pa., p. 496, Coal Meas.
alternans, Sternberg, 1833, Flor. der Vorw., vol. 2, p. 50, Coal Meas.
alveolaris, Sternberg, 1820, (Lepidodendron alveolare,) Essai d'un exposé Géognostico-botanique de la Flore du monde primitif, 1st Cahier, p. 25, Coal Meas.
angusta, Brongniart, 1828, Hist. d. Veg. Foss.; Coal Meas.
approximata, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 96, Coal Meas.

- attenuata*, Lesquereux, 1858, Catal. Potts. Foss., p. 17, and Coal Flora of Pa., p. 488, Coal Meas.
biercei, syn. for *S. ichthyolepis*.
brardi, Brongniart, 1822, Class. des Veg. Foss. tab. 1, fig. 5, and Coal Flora of Pa., p. 477, Coal Meas.
bretonensis, Dawson, 1865, Quar. Jour. Geo. Soc., vol. 20, p. 148, and Acad. Geol., p. 475, Coal Meas.
brochanti, Brongniart, 1828, Hist. d. Veg. Foss., p. 442, and Coal Flora of Pa., p. 842, Coal Meas.
brongniarti, Geinitz, 1855, Die Verst. d. Steink. form. Sachsen, p. 47, Coal Meas.
browni, Dawson, 1861, Quar. Jour. Geo. Soc., vol. 17, and Acad. Geol., p. 180, Coal Meas.

FIG. 71.—*Sigillaria brardi*.

- catenoides*, Dawson, 1865, Quar. Jour. Geo. Soc., vol. 20, p. 147, and Acad. Geol., p. 474, Coal Meas.
catenulata, Lindley & Hutton, 1831, Foss. Flora, vol. 1, p. 163, Coal Meas.
chemungensis, see *Lepidodendron chemungense*.
cisti, see *Caulopteris cisti*.
corrugata, Lesquereux, 1861, Geo. Sur. Ky., vol. 4, p. 437: redefined 1870, Geo. Sur. Ill., vol. 4, p. 445, Coal Meas.
cortei, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 64, and Coal Flora of Pa., p. 495, Coal Meas.
cuspidata, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 65, and Coal Flora of Pa., p. 486, Coal Meas.
cymatoides, Wood, 1860, Proc. Acad. Nat. Sci., vol. 12, p. 520, Coal Meas.
defranci, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 66, Coal Meas.
dentata, Newberry, 1874, Proc. Cleveland Acad. Sci., p. 165, Coal Meas.
dilatata, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 871, Coal Meas.
discoidea, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 873, Coal Meas.
dournaisi, Brongniart, 1828, Hist. d. Veg. Foss., p. 441, Coal Meas.
dubia, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 872, syn. for *S. cortei*.
elegans, Sternberg, 1826, (Favularia elegans,) Tent. flor. primord., p. 14, Coal Meas.
elliptica, Brongniart, 1828, Hist. d. Veg. Foss., p. 447, and Coal Flora of Pa., p. 494, Coal Meas.
elongata, Brongniart, 1822, Ann. des Sci. Nat., tom. 4, p. 23, Coal Meas.
eminens, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol. p. 475, Coal Meas.

- fissa*, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 871, Coal Meas.
flexuosa, Lindley & Hutton, 1837, Foss. Flora, vol. 3, p. 147, Coal Meas.
grandeyrii, Lesquereux, 1884, Coal Flora of Pa., p. 795, Coal Meas.
hexagona, Schlotheim, 1820, (Palmacites hexagonus,) Petrefaktenkunde, p. 394, and Coal Flora of Pa., p. 483, Coal Meas.
ichthyolepis, Sternberg, 1833, Flora d. Vorw., vol. 2, p. 38, and Coal Flora of Pa., p. 482, Coal Meas.
intermedia, Brongniart, 1828, Hist. d. Veg. Foss., p. 474, Coal Meas.
knorrii, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 65, Coal Meas.
lacoii, Lesquereux, 1880, Coal Flora of Pa., p. 490, Coal Meas.
lavigata, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 64, and Coal Flora of Pa., p. 500, Coal Meas.
leioderma, Brongniart, 1828, Hist. d. Veg. Foss., p. 422, and Coal Flora of Pa., p. 476, Coal Meas.
lepidodendrifolia, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 426, and Coal Flora of Pa., p. 477, Coal Meas.
leptoderma, Lesquereux, 1880, Coal Flora of Pa., p. 489, Coal Meas.
lescurii, Schimper, 1869, Trait. de Paléontologie Vegetale, vol. 2, p. 85, Coal Meas.
leveretti, Lesquereux, 1884, Coal Flora of Pa., p. 800, Coal Meas.
lorenzi, Lesquereux, 1880, Coal Flora of Pa., p. 473, Coal Meas.
lorwayana, Dawson, 1873, Rep. on Foss. Plants, p. 43, Subcarboniferous.
mammillaris, Brongniart, 1828, Hist. d. Veg. Foss., p. 451, and Coal Flora of Pa., p. 483, Coal Meas.
marginata, Lesquereux, 1880, Coal Flora of Pa., p. 498, Coal Meas.
marineria, Hildreth, 1837, Am. Jour. Sci. and Arts, vol. 31, p. 30, Low. Coal Meas.
massiliensis, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 446, Coal Meas.
menardi, Brongniart, 1828, Hist. d. Veg. Foss., p. 430, and Coal Flora of Pa., p. 479, Coal Meas.
monostigma, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 449, Coal Meas.
notata, Steinhaur, 1818, (Phytolithus notatus,) Trans. Am. Phil. Assoc., vol. 1, p. 294, and Coal Flora of Pa., p. 486, Coal Meas.
obliqua, Brongniart, 1828, Hist. Veg. Foss., p. 429, and Coal Flora of Pa., p. 470, Coal Meas.
obovata, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 872, Coal Meas.
oculata, Schlotheim, 1820, (Palmacites oculatus,) Petrefaktenkunde, p. 394, Coal Meas.
orbicularis, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 65, Coal Meas.
organum, Sternberg, 1820, (Syringodendron organum,) Flor. der Vorw., p. 23, and Lindley & Hutton, 1831, Foss. Flora, Vol. 1, p. 199, Coal Meas.
ornithicnoides, Wood, 1860, Proc. Acad. Nat. Sci., vol. 12, p. 238, and Trans. Am. Phil. Soc., vol. 13, p. 348, Coal Meas.
ovalis, Lesquereux, 1880, Coal Flora of Pa., p. 495, Coal Meas.
oweni, see Didymophyllum oweni.
pachyderma, see Syringodendron pachyderma.
pulpebra, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 307, and Acad. Geol. p. 536, Devonian.
perplexa, Wood, 1866, Proc. Acad. Nat. Sci. Phil. vol. 12, p. 237, Coal Meas.
pittstonana, Lesquereux, 1880, Coal Flora of Pa., p. 493, Coal Meas.
planicosta, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 474, Coal Meas.
polita, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 872, Coal Meas.
pulchra, Newberry, 1874, Proc. Cleveland Acad. Sci. p. 165, Coal Meas.
pyriformis, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 65, and Coal Flora of Pa., p. 799, Coal Meas.
reniformis, Brongniart, 1822, Ann. des Sci. Nat., t. 4, p. 32, and Coal Flora of Pa., p. 501, Coal Meas.
reticulata, Lesquereux, 1860, Geo. Sur. Ark., vol. 2, p. 310, Coal Meas.
rugosa, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 64, and Coal Flora of Pa., p. 497, Coal Meas.
saulli, Brongniart, 1823, Hist. Veg. Foss., vol. 1, p. 456, and Coal Flora of Pa., p. 842, Coal Meas.
schimperii, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 871, Coal Meas.
schlotheimiana, Brongniart, 1828, Hist. Veg. Foss., p. 469, Coal Meas. American Sp. (?)
sculpta, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 871, Coal Meas. Syn. for S. obliqua?
scutellata, Brongniart, 1822, Class. des Veg. Foss., tab. 1, fig. 4, Coal Meas.
semina, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 463, Coal Meas.
serlii, Brongniart, 1828, Hist. d. Veg. Foss., p. 433, and Coal Flora of Pa., p. 480, Coal Meas.
sillimani, Brongniart, 1828, Hist. Veg. Foss., p. 459, and Coal Flora of Pa., p. 493, Coal Meas.
simplicitas, Vanuxem, 1843, Geo. Rep. 3d Dist. N. Y., p. 190, Catskill Gr.
solanus, Wood, 1860, Proc. Acad. Nat. Sci., Coal Meas. [*Solanus* in text; *solenotus* on plate; *solena* in Trans. Am. Phil. Soc., vol. 13.]
spinulosa, Germ., 1844, Vers. v. Wettin, etc., p. 58, Coal Meas.
stellata, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 871, Coal Meas.
striata, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Quar. Jour. Geo. Soc., vol. 15, p. 147, Coal Meas.
sydenensis, Dawson, 1863, Can. Nat. and

Geol.
 Coal
 tessellat
 tessellat
 vol. 1
 481, C
 vanuxem
 Flora
 and C
 venosa,
 Foss.,
 842, C
 voltzi, B
 Veg. F
 p. 492
 williams
 Pa., p.
 yardleyi
 Foss.,
 491, C
 SIGILLARIO
 Ill., v
 sembla
 Fragn
 leaves
 boidal
 radicans,
 vol. 4,
 stellaris,
 SIGILLARIO
 Flora c
 Sigillar
 tached
 axis, su
 ophore
 up and
 laurenci
 of Pa.,
 Solenoula, V
 vol. 12,
 oulos, e
 Syringo
 psilophleu
 Sci., p.
 SOROCLADUS
 of Pa.,
 of the
 frond;
 name
 not wel

- Geol., vol. 8, and Acad. Geol., p. 475, Coal Meas.
- tessellata, Steinhaur, 1818, (Phytolithus tessellatus.) Trans. Am. Phil. Assoc., vol. 1, p. 295, and Coal Flora of Pa., p. 481, Coal Meas.
- vanuxemi, Gœppert, 1852, Die fossile Flora des Uebergangsgebirges, p. 546, and Coal Flora of Pa., p. 505, Coal Meas.
- venosa, Brongniart, 1828, Hist. d. Veg. Foss., p. 424, and Coal Flora of Pa., p. 842, Coal Meas.
- voltzi, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 65, and Coal Flora of Pa., p. 492, Coal Meas.
- williamsi, Lesquereux, 1880, Coal Flora of Pa., p. 488, Coal Meas.
- yardleyi, Lesquereux, 1858, Catal. Potts. Foss., p. 17, and Coal Flora of Pa., p. 491, Coal Meas.
- SIGILLARIOIDES, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 449. [Ety. from its resemblance to the genus *Sigillaria*.] Fragments of roots bearing stigmaroid leaves attached to sigillarioid rhomboidal scars. Type *S. radicans*.
- radicans, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 449, Coal Meas.
- stellaris, see *Stigmaria stellaris*.
- SIGILLARIOSTROBUS, Lesquereux, 1884, Coal Flora of Pa., p. 794. [Ety. the genus, *Sigillaria*; *strobis*, cone.] Sporangies attached in horizontal rows to a vertical axis, supported by persistent sporangio-phores, with lanceolate scales, turned up and imbricate. Type *S. laurencianus*.
- laurencianus, Lesquereux, 1884, Coal Flora of Pa., p. 794, Coal Meas.
- Solenoula*, Wood, 1860, Proc. Acad. Nat. Sci., vol. 12, p. 238. [Ety. *solen*, a channel; *oulos*, entire.] Probably a decorticated *Syringodendron*. Type *S. psilophleus*.
- psilophleus*, Wood, 1860, Proc. Acad. Nat. Sci., p. 238, Coal Meas.
- SOROCLADUS, Lesquereux, 1880, Coal Flora of Pa., p. 327. [Ety. *soros*, a heap: one of the fruit dots on the back of the frond; *klado*, I break in pieces.] A name proposed for fruiting fragments not well understood. Type *S. stellatus*.
- ophioglossoides, Lesquereux, 1880, Coal Flora of Pa., p. 329, Coal Meas.
- sagittatus, Lesquereux, 1870, (Staphylopteris sagittatus.) Geo. Sur. Ill., vol. 4, p. 407, Coal Meas.
- stellatus, Lesquereux, 1860, (Staphylopteris stellata.) Geo. Sur. Ark., vol. 2, p. 309, Coal Meas.
- wortheni, Lesquereux, 1870, (Staphylopteris wortheni.) Geo. Sur. Ill., vol. 4, p. 405, Coal Meas.
- SPHENOPHYLLUM, Brongniart, 1828, Prodr. d. Hist. Veg. Foss., p. 68. [Ety. *sphen*, a wedge; *phyllon*, a leaf.] It was called *Sphenophyllites* by Brongniart in 1822. Plant herbaceous; stems articulate, inflated at the articulations, pinnately, bipinnately divided; leaves verticillate, sessile, wedge-form, with lateral borders entire, crenulate, dentate, or laciniate-lobate at the upper margin; medial nerve none; veins straight dichotomous; fructifications in cylindrical spikes, with bracts curved upward in a sharp flexure from near the base; sporanges globular in the axils of the bracts. Type *S. schlotheimi*.
- angustifolium, Germar, 1844, Verst. d. Steink. v. Wett., u. Löbejün, and Coal Flora of Pa., p. 726, Coal Meas.
- antiquum, Dawson, 1861, Can. Nat., vol. 6, p. 170, Devonian.
- bifurcatum, Lesquereux, 1860, Geo. Sur. Ark., vol. 2, p. 309, Coal Meas.
- brevifolium, Newberry, not defined.
- cornutum, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 421, Coal Meas.
- densifolium, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 37, Coal Meas. or Permian. Syn. (?) for *S. angustifolium*.
- emarginatum, Brongniart, 1828, Prodr. d. Hist. Veg. Foss., p. 68, and Coal Flora of Pa., p. 53, Coal Meas.
- erosum, Lindley & Hutton, 1833, Foss. Flora, vol. 1, p. 43, and Coal Flora of Pa., p. 55, Coal Meas.
- filiculme, Lesquereux, 1858, Geo. Rep. Pa., vol. 2, p. 853, Coal Meas.
- fontainianum, S. A. Miller, 1883, 2d. Ed. Am. Pal. Foss., p. 258, Up. Coal Meas. Proposed instead of *S. latifolium*, in Perm. or Up. Carb. Flora, p. 36, which was preoccupied.
- latifolium, Wood, 1866, Trans. Am. Phil. Soc., vol. 13, p. 347, Coal Meas.
- latifolium, Fontaine & White, 1880. The name was preoccupied. See *S. fontainianum*.
- longifolium, Germar, 1831, (Rotularia longifolia.) Isis, p. 426, and Coal Flora of Pa., p. 53, Coal Meas.
- oblongifolium, Germar, 1844, Verst. d. Steink. v. Wett., u. Löbejün, p. 12, and Coal Flora of Pa., p. 57, Coal Meas.
- primævum, Lesquereux, 1877, Proc. Am. Phil. Soc., p. 167, Hud. Riv. Gr. I think this is not a plant.

FIG. 72.—*Sorocladus asteroides*.

asteroides, Lesquereux, 1870, (Staphylopteris asteroides.) Geo. Sur. Ill., vol. 4, p. 406, Coal Meas.

saxifragifolium, Sternberg, 1825, (Rotularia saxifragifolia,) Vera. Darst. Flora der Vorwelt, and Coal Flora of Pa., p. 726, Coal Meas.

FIG. 78.
Sphenophyllum
schlotheimi.

schlotheimi, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 68, and Coal Flora of Pa., p. 52, Coal Meas.

tenerimum, Stur, 1877, Culm. Flora, p. 108, and Coal Flora of Pa., p. 728, Coal Meas.

trifoliatum, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 853, Coal Meas.

SPHENOPTERIS, Brongniart, 1822, Mem. du Mus. d'Hist. Nat. de Paris, tom. 8, p. 203. [Ety. *sphen*, wedge; *pteris*, fern.] Fronds bi, tri, polypinnate; divisions open or in right angles; pinnules narrowed at base, often decurring or cuneiform, pinnately lobed; lobes rarely entire, crenulate, dentate, or lacinate; primary nerve slender, alternately dichotomous, simple, branches entering the base of each lobe to pass by branchlets into the subdivisions of the lamina. Type *S. elegans*.

abbreviata, see *Pseudopteris abbreviata*.

acrocarpa, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 40, Coal Meas. or Permian.

acuta, See *Pseudopteris acuta*.

adiantoides, Lindley & Hutton, 1835, Foss. Flora, vol. 2, p. 91, Coal Meas.

alata, Brongniart, 1828, (*Pecopteris alata*), Hist. d. Veg. Foss., p. 361, Coal Meas.

alabamensis, see *Oligocarpia alabamensis*.

artemesiifolia see *Eromopteris artemesiifolia*.

auriculata, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 42, Coal Meas. or Permian.

ballantini, Andrews, 1875, (*Hymenophyllites ballantini*), Ohio Pal., vol. 2, p. 422, Coal Meas.

brittsi, Lesquereux, 1880, Coal Flora of Pa., p. 277, Coal Meas.

canadensis, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol. p. 243, Coal Meas.

chacrophylloides, Brongniart, 1828, (*Pecopteris chacrophylloides*), Hist. d. Veg. Foss., p. 357, and Coal Flora of Pa., p. 270, Coal Meas.

communis, Lesquereux, 1884, Coal Flora of Pa., p. 762, Coal Meas.

coriacea, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 41, Coal Meas. or Permian.

crenata, Lindley & Hutton, 1835, Foss. Flora, vol. 2, pl. C., and Coal Flora of Pa., p. 835, Coal Meas.

cristata, Brongniart, 1828, (*Pecopteris cristata*), Hist. d. Veg. Foss., p. 356,

and Coal Flora of Pa., p. 273, Coal Meas.

davallana, Gœppert, 1841, Gatt. d. Foss. Pflanzen, Coal Meas.

decipiens, see *Pseudopteris decipiens*, *delicatula*, see *Hymenophyllites delicatulus*.

dentata, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 42, Coal Meas. or Permian.

dilatata, as identified by Lesquereux, Syn. for *Pseudopteris decipiens*.

dissecta, Brongniart, 1828, Hist. d. Veg. Foss., p. 183, and Coal Flora of Pa., p. 836, Coal Meas.

divaricata, Gœppert, 1836, (*Chellanthus divaricatus*), Syst. Filic. Foss., p. 238, and Coal Flora of Pa., p. 767, Coal Meas.

dubuissoni, Brongniart, 1828, Hist. d. Veg. Foss., p. 195, and Coal Flora of Pa., p. 275, Coal Meas.

elegans, Brongniart, 1822, Class. d. Veg. Foss. pl. 2, fig. 2, and Coal Flora of Pa., p. 287, Coal Meas.

fascicularis, Roemer, 1866, Beitr. in Palæont., vol. 9, p. 179, and Coal Flora of Pa., p. 837, Coal Meas.

flaccida, Crepin, 1874, Bull. Acad. Roy. of Belgium, p. 7, and Coal Flora of Pa., p. 291, Coal Meas.

FIG. 79.—*Sphenopteris crenata*.

flagellari, see *Oligocarpia flagellaris*.

flexicaulis, Lesquereux, 1860, (*Hymenophyllites flexicaulis*), Geo. Sur. Ark., vol. 2, p. 309, Coal Meas.

foliosa, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 44, Coal Meas. or Permian.

fuiformis, Lesquereux, 1884, Am. Naturalist, vol. 18, p. 921, Carboniferous.

furcata, Brongniart, 1828, Hist. d. Veg. Foss., p. 179, and Coal Flora of Pa., p. 282, Coal Meas.

gersdorffii, see *Hymenophyllites gersdorffii*, *glandulosa*, see *Pseudopteris glandulosa*.

goniopteroides, Lesquereux, 1880, Coal Flora of Pa., p. 269, Coal Meas.

gracilis, Brongniart, 1828, Hist. d. Veg. Foss., p. 197, and Coal Flora of Pa., p. 276, Coal Meas.

gravenhorsti, Brongniart, 1828, Hist. Veg. d. Foss., p. 191, and Coal Flora of Pa., p. 274, Coal Meas.

hartti, I
Soc., v
harveyi,
Pa., p.
hastata,
or Up.
or Per
hildrethi
phyllit
vol. 2,
hitchcocki
Geol. S
hoeningh
d. Foss
p. 288,
hymenop
Prodr.
and C
Meas.
inequiliat
Flora o
intermedia
Pa., vol
in 1852
median
irregulari
ularis.
larischii,
ischii.)
Flora o
latifolia, s
lator, D
and Ac
lara, Hull
Chemu
cupied
hallana
lescuriana
or Up.
or Perm
lesquereu
Pa., vol
linearis, S
d. Vorw
lyratifolia,
macilenta,
lenta.
marginata
Geo. So
mediana,
Pa., p. 2
microcarp
of Pa., p
microloba
Foss., p
minutisecc
Perm. o
Meas. o
mixta, Sch
tologie
of Pa., p
munda, D
vol. 8,
Meas.
myriophyll
ophyllu
newberryi,
berryi.

Coal
 Foss.
 ptiens.
 delica-
 Perm.
 Meas.
 ereux.
 ens.
 . Veg.
 Pa., p.
 anthes
 p. 238,
 Coal
 ist. d.
 ora of
 . Veg.
 of Pa.,
 n Pal-
 ora of
 Roy. of
 Pa., p.
 hartti, Dawson, 1862, Quar. Jour. Geol. Soc., vol. 18, p. 321, Devonian.
 harveyi, Lesquereux, 1884, Coal Flora of Pa., p. 766, Coal Meas.
 hastata, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 46, Coal Meas. or Permian.
 hildrethi, Lesquereux, 1858, (Hymenophyllites hildrethi,) Geo. Sur. of Pa., vol. 2, p. 863, Coal Meas.
 hitchcockiana, Dawson, 1862, Quar. Jour. Geol. Soc., vol. 18, p. 321, Devonian.
 hoenlghausi, Brongniart, 1828, Hist. Veg. d. Foss., p. 199, and Coal Flora of Pa., p. 288, Coal Meas.
 hymenophyllioides, Brongniart, 1828, Prodr. d. Hist. d. Veg. Foss., p. 51, and Coal Flora of Pa., p. 764, Coal Meas.
 inequilateralis, Lesquereux, 1884, Coal Flora of Pa., p. 765, Coal Meas.
 intermedia, Lesquereux, 1858, Geo. Sur. Pa., vol. 2. The name was preoccupied in 1852 by Ettingshausen. It is now *S. mediana*.
 irregularis, see *Pseudopteris irregularia*.
 larischii, Stur, 1877, (Calymmotheca larischii,) Culm Flora, p. 168, and Coal Flora of Pa., p. 288, Coal Meas.
 latifolia, see *Pseudopteris latifolia*.
 lator, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 483, Coal Meas.
 laca, Hall, 1843, Geo. Rep. 4th Dist. N. Y., Chenung Gr. This name was preoccupied by Sternberg. See *Archæopteris hallana*.
 lescuriana, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 44, Coal Meas. or Permian.
 lesquereuxi, Newberry, 1858, Geo. Sur. Pa., vol. 2, p. 862, Coal Meas.
 linearis, Sternberg, 1820, Vers. Darst. Flor. d. Vorw., p. 15, Low. Coal Meas.
 lyratifolia, see *Pecopteris lyratifolia*.
 macilenta, see *Pseudopteris macilenta*.
 marginata, Dawson, 1862, Quar. Jour. Geol. Soc., vol. 18, p. 321, Devonian.
 mediana, Lesquereux, 1880, Coal Flora of Pa., p. 271, Coal Meas.
 microcarpa, Lesquereux, 1880, Coal Flora of Pa., p. 280, Coal Meas.
 microloba, Gœppert, 1836, Syst. Filic. Foss., p. 238, Coal Meas.
 minutisecta, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 43, Coal Meas. or Permian.
 mixta, Schimper, 1869, Traité de Paléontologie Veg. tale, p. 382, and Coal Flora of Pa., p. 276, Coal Meas.
 munda, Dawson, 1863, Can. Nat. and Geo., vol. 8, and Acad. Geol., p. 483, Coal Meas.
 myriophylla, see *Hymenophyllites myriophyllus*.
 newberryi, see *Pseudopteris newberryi*.

obovata, Lindley & Hutton, 1835, Foss. Flora, vol. 2, p. 75, and Coal Flora of Pa., p. 769, Coal Meas.
 obtusiloba, see *Pseudopteris obtusiloba*.
 pachynervis, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 46, Coal Meas. or Permian.
 paupercula, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 435, Coal Meas.
 pilosa, see *Callipteris pilosa*.
 pilenta, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 862, Coal Meas.
 polyphylla, see *Pseudopteris polyphylla*.
 pseudomurrayana, Lesquereux, 1880, Coal Flora of Pa., p. 271, Coal Meas.
 pterota, Wood, 1866, Trans. Am. Phil. Soc., vol. 13, p. 348, Coal Meas.
 quercifolia, Gœppert, 1836, Syst. Filic. Foss., p. 252, and Coal Flora of Pa., p. 286, Coal Meas.
 recurva, Dawson, 1863, Quar. Jour. Geol. Soc., vol. 19, p. 464, Devonian.
 rigida, Brongniart, 1828, Hist. Veg. Foss., p. 201, Coal Meas.
 royi, Lesquereux, 1884, Coal Flora of Pa., p. 768, Coal Meas.
 scaberrima, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 408, Coal Meas.
 schlotheimi, see *Hymenophyllites schlotheimi*.
 solida, Lesquereux, 1884, Coal Flora of Pa., p. 769, Coal Meas.
 spinosa, Gœppert, 1841, Gatt. Foss. Pflanzen, p. 70, and Coal Flora of Pa., p. 281, Coal Meas.
 splendens, Dawson, 1871, Foss. Plants Canada, p. 53, Devonian.
 splendens, Lesquereux, 1870, (Hymenophyllites splendens,) Geo. Sur. Ill., vol. 4, p. 413, Coal Meas.
 squamosa, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 862, Coal Meas.
 subalata, Weiss, 1869, Foss. Flora d. jungst. Steink. form., p. 57, and Coal Flora of Pa., p. 272, Coal Meas.
 tenella, Brongniart, 1828, Hist. Veg. Foss., p. 186, and Coal Flora of Pa., p. 836, Coal Meas.
 tenuifolia, see *Hymenophyllites tenuifolius*.
 tracyana, Lesquereux, 1884, Coal Flora of Pa., p. 766, Coal Meas.
 trichomanoides, Brongniart, 1828, Hist. d. Veg. Foss., p. 182, and Coal Flora of Pa., p. 286, Coal Meas.
 tridactylites, Brongniart, 1828, Hist. d. Veg. Foss. p. 181, and Coal Flora of Pa., p. 284, Coal Meas.
 trifoliata, see *Pseudopteris trifoliata*.
 SPHENOTHALLUS, Hall, 1847, Pal. N. Y., vol. 1, p. 261. [Ety. *sphen*, a wedge; *thallos*, a branch or frond.] Stem with diverging wedge-formed leaves, thickened, and sometimes subcoriaceous. Type *S. angustifolius*.

angustifolius, Hall, 1847, Pal. N. Y., vol. 1, p. 261, Hud. Riv. Gr.

FIG. 75.—*Sphenothallus angustifolius*.

latifolius, Hall, 1847, Pal. N. Y., vol. 1, p. 262, Hud. Riv. Gr.

SPHRANGIUM, Schimper, 1874, *Traité de Paléontologie Végétale*, vol. 2, p. 514. [Ety. *speira*, that which is twisted; from the coiled marking around the pod.] Oblong or spindle-shaped bodies formed of narrow linear leaves, interwoven or twisted in spiral, with the ends united into a pedicel, which joins them horizontally or in umbels. Type *S. carbonarium*.

appendiculatum, Lesquereux, 1870, (*Palæoxyris appendiculata*.) Geo. Sur. Ill., vol. 4, p. 465, Coal Meas.

corrugatum, Lesquereux, 1870, (*Palæoxyris corrugata*.) Geo. Sur. Ill., vol. 4, p. 460, Coal Meas.

intermedium, Lesquereux, 1880, Coal Flora of Pa., p. 521, Coal Meas.

multiplicatum, Lesquereux, 1880, Coal Flora of Pa., p. 520, Coal Meas.

prendeli, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 464, Coal Meas.

Spirophyton, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist. p. 78. [Ety. *speira*, a coil; *phyton*, a plant.] Syn. for *Taonurus*.

cauda-galli, see *Taonurus caudagalli*.

crassum, see *Taonurus crassum*.

typus, see *Taonurus typus*.

velum, see *Taonurus velum*.

SPORANGITES, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Proc. Geo. Soc. Lond., vol. 15, p. 164. [Sig. seed-vessel.] Spores and spore-cases of *Lepidodendron*, *Calamites*, and similar plants, which can not be otherwise referred. Type *S. papillatus*.

acuminatus, Dawson, 1861, (*Annularia acuminata*.) Can. Nat., vol. 6, and Acad. Geol., p. 540, Portage Gr.

bilobatus, Dawson, 1883, Proc. Am. Ass. Ad. Sci., vol. 32, p. 260, Marcellus Shale.

glaber, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 491, Coal Meas.

huronensis, Dawson, 1871, Am. Jour. Sci. and Arts, p. 257, Ham. Gr.

papillatus, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 491, Coal Meas.

SPOROCYSTIS, Lesquereux, 1880, Coal Flora of Pa., p. 458. [Ety. *spora*, seed; *kystis*, bladder.] Agglomerations of macrospores grouped together or cohering or agglutinate by the borders, more generally without cases, and therefore of uncertain reference. Type *S. planus*.

planus, Lesquereux, 1880, Coal Flora of Pa., p. 458, Coal Meas.

Staphylopteris, 1838, Presl, in Sternb. Vort. Darst. Flora der Vorwelt. [Ety. *staphyle*, bunch of grapes; *pteris*, fern.] Not an American paleozoic genus.

asteroides, see *Sorocladus asteroides*.

sagittata, see *Sorocladus sagittatus*.

stellata, see *Sorocladus stellatus*.

wortheni, see *Sorocladus wortheni*.

STEMMATOPTERIS, Corda, 1845, Beiträge zur Flora der Vorwelt, p. 76. [Ety. *stemma*, a wreath; *pteris*, fern.] Trunks erect, cylindrical; scars large, disciform, oval, round, or ovate, not contiguous, disposed in quincuncial or spiral order; outside borders or rings flat; internal disk formed by impressions of fascicles of vascular tissues, shaped like a horseshoe, the horns curving inward in the upper part of the scars, either short and hooked, or descending below the middle of the scars, and there united. Type *S. peltigera*.

anceps, Lesquereux, 1884, Coal Flora of Pa., p. 838, Coal Meas.

angustata, Lesquereux, 1880, Coal Flora of Pa., p. 330, Coal Meas.

cyclostigma, Lesquereux, 1880, Coal Flora of Pa., p. 341, Coal Meas.

emarginata, Lesquereux, 1880, Coal Flora of Pa., p. 337, Coal Meas.

gigantea, Lesquereux, 1858, (*Caulopteris gigantea*.) Geo. of Pa., vol. 2, p. 869, Coal Meas.

hirsuta, Lesquereux, 1880, Coal Flora of Pa., p. 337, Coal Meas.

insignis, Lesquereux, 1870, (*Caulopteris insignis*.) Geo. Sur. Ill., vol. 4, p. 459, Coal Meas.

FIG. 77.—*Stemmatopteris mimica*, leaf scar.

(*Caulopteris punctata*.) Geo. Sur. Pa., vol. 2, p. 869, Coal Meas.

schimper, Lesquereux, 1880, Coal Flora of Pa., p. 338, Coal Meas.

FIG. 76.—*Sporangites papillatus*.

Sphenothallus of Pa., worthenii, Coal M. STERNBERG Phytoz. The plant other in sand perfect which in the transverse transverse vol. 1, var. angustifolius, Geo. Sur.

FIG.

var. approx. Jour. G. Meas. var. distans Geo. Sur. var. obscura Geo. Sur. SIGMATHIA, J. Foss. in Paris, to dot or roots, g. distantly scarcely length, pith, a trical, co. disposed linear w. ruption, round sc. trical rim mammil. punctif. ficoides. amena, L. Pa., p. 5. anabathra, der Vorw. areolata, D. ada, p. 2. costata, L. vol. 2, p. elliptica, L. vol. 4, p. eveni, see S.

squamosa, Lesquereux, 1880, Coal Flora of Pa., p. 330, Coal Meas.
wortheni, Lesquereux, 1866, (Caulopteris *wortheni*.) Geo. Sur. Ill., vol. 2, p. 459, Coal Meas.

STERNBERGIA, Artis, 1825, Antediluvian Phytology, p. 8. [Ety. proper name.] The piths of *Dadoxylon*, *Sigillaria*, and other plants usually preserved as casts in sandstone, retaining more or less perfectly the transverse partitions into which the pith *Sternbergs* were divided in the process of growth. Type *S. transversa*.
transversa, Steinhaur, 1818, (Phytolithus *transversus*.) Trans. Am. Phil. Ass'n., vol. 1, p. 295, Coal Meas.
var. angularis, Dawson, 1865, Quar. Jour. Geo. Soc., vol. 22, p. 165, Coal Meas.

FIG. 78.—*Sternbergia angularis*, pith of *Dadoxylon*.

var. approximata, Dawson, 1865, Quar. Jour. Geo. Soc., vol. 22, p. 165, Coal Meas.
var. distans, Dawson, 1865, Quar. Jour. Geo. Soc., vol. 22, p. 165, Coal Meas.
var. obscura, Dawson, 1865, Quar. Jour. Geo. Soc., vol. 22, p. 165, Coal Meas.
STIGMARIA, Brongniart, 1822, Class. d. Veg. Foss. in Mem. du Mus. d'Hist. Nat. d. Paris, tom. 8, p. 203. [Ety. *stigma*, a dot or puncture.] Floating stems or roots, generally growing horizontally, distantly dichotomous; branches long, scarcely variable in size in their whole length, subcylindrical or compressed; pith, a woody cylinder, often eccentric, composed of fascicles of vessels disposed star-like; leaves long, tubulose, linear when flattened, leaving after disruption, on the surface of the stems, round scars composed of two concentric rings, with a central umbonate mammilla, pitted in the middle by a punctiform vascular scar. Type *S. ficoides*.
amena, Lesquereux, 1880, Coal Flora of Pa., p. 516, Coal Meas.
anabathra, Corda, 1845, Beiträge zur Flora der Vorwelt, p. 34, Coal Meas.
areolata, Dawson, 1871, Foss. Plants Canada, p. 23, Devonian.
costata, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 870, Coal Meas.
elliptica, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 451, Coal Meas.
eveni, see *Stigmarioides eveni*.

exigua, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 308, Chemung Gr.
ficoides, Brongniart, 1822, Mem. du Mus. d'Hist. Nat. de Paris, tom. 8, p. 203, Coal Meas.
ficoides var. a, b, c, d, e, f, g, h, i, k, l, Dawson, 1865, Quar. Jour. Geo. Soc., vol. 22, p. 148, Coal Meas.
ficoides var. reticulata, Goeppert, 1841, Gatt. d. Foss. Pflanzen, p. 13, Coal Meas.
ficoides var. stellata, Goeppert, 1841, Gatt. d. Foss. Pflanzen, p. 13, Coal Meas.
ficoides var. undulata, Goeppert, 1841, Gatt. d. Foss. Pflanzen, p. 13, Coal Meas.
irregularis, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 870, Coal Meas. [Ety. from the irregularity of the scars.]
minor, Goeppert, 1841, Gatt. d. Foss. Pflanzen, p. 13, Coal Meas.
minuta, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 871, Coal Meas.
minutissima, Dawson, 1871, Foss. Plants Can., p. 23, Devonian.
perlata, Dawson, 1871, Foss. Plants Canada, page 22, Devonian.
pusilla, Dawson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 460, Devonian.
radicans, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 870, Coal Meas.
stellaris, Lesquereux, 1870, (Sigillarioides *stellaris*.) Geo. Sur. Ill., vol. 4, p. 450, Coal Meas.
umbonata, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 870, Coal Meas.

FIG. 79.—*Stigmaria ficoides*, $\frac{1}{4}$ diam.

STIGMARIOIDES, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 452. [Ety. from its resemblance to *Stigmaria*.] Fragments of rhizomas, with surface marked by small round impressions, irregularly disposed and without central vascular points, base of detached radicles or filaments. Type *S. eveni*.
affinis, see *Rachiopteris affinis*.
eveni, Lesquereux, 1866, (Stigmaria *eveni*.) Geo. Sur. Ill., vol. 4, p. 448, Coal Meas.
linearis, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 455, Coal Meas.
rugosus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 470, Coal Meas.
selago see *Rachiopteris selago*.

- truncatus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 453, Coal Meas.
 tuberosus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 453, Coal Meas.
 villosus, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 454, Coal Meas.
Strobilus caryophyllus, Hildreth, 1837, Am. Jour. Sci. and Arts, vol. 31, p. 32, Coal Meas. Possibly a *Stigmaria*.
 SYRINGODENDRON, Sternberg, 1820, Essai d'un exposé Geognostico-botanique de la Flore du monde primitif, 1st Cahier, p. 26. [Ety. *syrix*, a pipe; *dendron*, tree.] Cortex costate; vascular scars united in one; resembles decorticated stems of *Sigillaria*. Type *S. pes capreoli*.
bistriatum, Wood, 1860, Proc. Acad. Nat. Sci., vol. 12, p. 521, Coal Meas.
brongniarti, Geinitz, 1855, (*Sigillaria brongniarti*.) Verst. d. Steink. form. in Sachsen, p. 47, and Coal Flora of Pa., p. 504, Coal Meas.
cyclostigma, Brongniart, 1828, Hist. d. Veg. Foss., p. 480, and Coal Flora of Pa., p. 505, Coal Meas.
gracile, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 308, Waverly Gr.
kirtlandium, Hildreth, 1837, Am. Jour. Sci. & Arts, vol. 31, p. 29, Coal Meas.
magnificum, Wood, 1866, Trans. Am. Phil. Soc., vol. 13, p. 352, Coal Meas.
organum, see *Sigillaria organum*.
pachyderma, Brongniart, 1828, (*Sigillaria pachyderma*.) Prodr. d. Hist. d. Veg. Foss., p. 65, and Coal Flora of Pa., p. 503, Coal Meas.
pescapreoli, Sternberg, 1820, Essai d'un exposé Geognostico-botanique de la Flore du monde primitif, 1st Cahier, p. 26, Coal Meas.
porteri, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 448, Coal Meas.
 SYRINGOXYLON, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 305. [Ety. *syrix*, a pipe; *xylon*, wood.] Woody tissue close, thick-walled; ducts many times the diameter of the wood-cells, thin walled, with transverse pores in several series; medullary rays of two or more series of muriform cells; growth rings, distinct. Type *S. mirabile*.
mirabile, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 305, Ham. Gr.
 TÆNIOPHYLLUM, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 330. [Ety. *tainia*, ribbon; *phyllon*, leaf.] Stems large, leaves crowded, fistular, flat by compression, thick, exactly linear, decurring at the base; surface smooth, opaque, or shining. Type *T. decurrens*.
brevifolium, Lesquereux, 1880, Coal Flora of Pa., p. 788, Coal Meas.
contextum, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 332, Coal Meas.
decurrens, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 331, and Coal Flora of Pa., p. 464, Coal Meas.
deflexum, Lesquereux, 1878, Proc. Am. Phil. Soc., p. 331, Coal Meas.
 TÆNIOPTERIS, Brongniart, 128, Prodr. Hist. d. Veg. Foss., p. 61. [Ety. *tainia*, ribbon; *pteria*, fern.] Fronds simple, large, linear; medial nerve canaliculate, strong; veins open, or in right angle, thin, forking a little above the base or more generally simple, parallel, sometimes joined to a marginal nerve. Type *T. vittata*.
lescuriana, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 91, Coal Meas. or Permian.
newberryana, Fontaine & White, 1880, Perm. or Up. Carb. Flora, p. 91, Coal Meas. or Permian.
smithi, Lesquereux, 1875, Geo. Rep. Ala., p. 78, and Coal Flora of Pa., p. 153, Coal Meas.

FIG. 80.—Tæniopteris smithi.

- truncata*, Lesquereux, 1884, Coal Flora of Pa., p. 743, Coal Meas.
 TAONURUS, Fisher-Ooster, 1858, Foss. Fucoiden d. Schweizer Alpen, p. 41. [Ety. *taon*, peacock; *oura*, tail.] Frond membranaceous, derived from utricles attached to a lateral or central axis, erected or twisted in spiral, flattened in various ways, ribbed; ribs or striae curved, scythe-shaped, converging to the borders, which are either free, naked or attached on one side or all around to the axis or its branches. Type *T. caudagalli*.

FIG. 81.—Taonurus caudagalli.

- archimedes*, Ringueberg, 1884, (Spiral-ton archimedes.) Proc. Acad. Nat. Sci. Phil., p. 144, Medina Gr.
caudagalli, Vanuxem, 1842, (Fucoides

candag
p. 128,
colletti,
colletti,
Coal M
crassus,
sum.) 1
p. 83, V
marginat
pites m
vol. 13,
retortus,
Geo. R
Portage
typus, H
16 Rep.
Ham. o
veilm, V
Geo. Re
TRICHTOMAN
Foss. [
manes.]
Americ
nules at
are of d
filicula, D
Soc., vo
TRICHOPTERYC
Cin. Soc
trichos, h
branchi
by the f
T. lanos

FIG. 82.—Trichophyllum lanosum.

Foss., p
karpos, f
at the b
six cost
promine
disappe
small, r
cavity.
adansii, L
Pa., p. 8
anpullifor
Flora of
avellanum
Geo., vo
Coal Me
bertholleti
of Sci., v
369, Coa
carbonariu
Nat. Sci.
dawsii, Li
Flora, v
Pa., p. 5

- candagalli, Geo. Rep. 3d Dist. N. Y., p. 128, Devonian.
- colletti, Lesquereux, 1870, (Chondrites colletti,) Geo. Sur. Ill., vol. 4, p. 379, Coal Meas.
- crassus, Hall, 1863, (Spirophyton crassum,) 16 Rep. N. Y. St. Mus. Nat. Hist., p. 83, Waverly Gr.
- marginatus, Lesquereux, 1866, (Cauler-pites marginatus,) Trans. Am. Phil. Soc., vol. 13, p. 314, Subcarb.
- retortus, Vanuxem, 1842, (Retort fucoid,) Geo. Rep. 3d Dist., N. Y., p. 176, Portage Gr.
- typus, Hall, 1863, (Spirophyton typus,) 16 Rep. N. Y. St. Mus. Nat. Hist., p. 80, Ham. or Chemung Gr.
- velum, Vanuxem, 1842, (Fucoides velum,) Geo. Rep. 3d Dist. N. Y., p. 176, Ham. Gr.
- TRICHOMANITES, Gœppert, 1836, Syst. Filic. Foss. [Ety. from the plant *Trichomanes*.] This genus is only known in America by fragments of slender pin-nules attached to long petioles, which are of doubtful generic affinity.
- filiola, Dawson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 464, Devonian.
- TRICHOPHYCUS, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 24. [Ety. *trichos*, hair; *phukos*, sea-weed.] Simple branching stems having markings as if by the folding down of filaments. Type *T. lanosum*.
- lanosum, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 24, Hud. Riv. Gr.
- sulcatum, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 4, Hud. Riv. Gr.
- venosum, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 112, Hud. Riv. Gr.

FIG. 82.—*Trichophycus lanosum*.

- TRIGONOCARPUM, Brongniart, 1828, Prodr. d. Hist. Veg. Foss., p. 137. [Ety. *trigon*, triangle; *karpos*, fruit.] Fruits ovoid, compressed at the base point of insertion, three or six costate, the ribs more distinct and prominent toward the base, sometimes disappearing above; apex pitted by a small, round or triquetre mammillate cavity. Type *T. parkinsoni*.
- adamsi, Lesquereux, 1884, Coal Flora of Pa., p. 820, Coal Meas.
- anpulliforme, Lesquereux, 1884, Coal Flora of Pa., p. 823, Coal Meas.
- avellanum, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 478, Coal Meas.
- bertholletiforme, Foster, 1853, Ann. of Sci., vol. 1, and Ohio Pal., vol. 1, p. 369, Coal Meas.
- carbonarium, King, 1854, Proc. Acad. Nat. Sci., vol. 7, p. 66, Coal Meas.
- dawsi, Lindley & Hutton, 1837, Foss. Flora, vol. 3, p. 321, and Coal Flora of Pa., p. 586, Coal Meas.

- giffordi, Lesquereux, 1880, Coal Flora of Pa., p. 592, Coal Meas.
- grande, Lesquereux, 1884, Coal Flora of Pa., p. 821, Coal Meas.
- hildrethi, Lesquereux, 1858, Geo. Sur. Pa., vol. 2, p. 877, Coal Meas.
- hildrethi, Dawson, syn. (?) for *Trigonocarpum triloculare*.
- hookeri, Dawson, 1861, Quar. Jour. Geol. Soc., vol. 17, p. 525, Coal Meas.
- intermedium, Dawson, 1863, Can. Nat. vol. 8, and Acad. Geol., p. 478, Coal Meas.
- juglans, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 460, Low. Coal Meas.
- kansaseanum, Lesquereux, 1884, Coal Flora of Pa., p. 822, Coal Meas.
- magnum, Newberry, 1873, Ohio Pal., vol. 1, p. 369, Coal Meas.
- mentzelianum, Gœppert & Berger, 1848, De Fruct. et Sem., p. 19, and Coal Flora of Pa., p. 590, Coal Meas.
- minus, Dawson, 1863, Can. Nat. and Geol., vol. 8, and Acad. Geol., p. 478, Coal Meas.
- multicarinarum, Newberry, 1873, Ohio Pal., vol. 1, p. 478, Carb. Conglomerate.
- multistriatum, Lesquereux, 1884, Coal Flora of Pa., p. 823, Coal Meas.
- næggerathi, Sternberg, 1820, (Palmacites næggerathi,) Flor. d. Vorw., p. 55, and Coal Flora of Pa., p. 584, Coal Meas.
- oblongum, Lindley & Hutton, 1837, Foss. Flora, vol. 3, p. 193, Coal Meas.
- oliviforme, Lindley & Hutton, 1837, Foss. Flora, vol. 3, p. 222, and Coal Flora of Pa., p. 590, Coal Meas.
- ornatum, Newberry, 1873, Ohio Pal., vol. 1, p. 368, Carb. Conglomerate.
- parkinsoni, Brongniart, 1828, Prodr. Hist. Veg. Foss., p. 137, and Coal Flora of Pa., p. 589, Coal Meas.
- perantiquum, Dawson, 1871, Foss. Plants Canada, p. 62, Devonian.
- perpusillum, Lesquereux, 1884, Coal Flora of Pa., p. oliviforme, 820, Coal Meas.
- racemosum, Dawson, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 324, Devonian.
- rostellatum, Lesquereux, 1866, Geo. Sur. Ill., vol. 2, p. 460, Up. Coal Meas.
- rotundum, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 478, Coal Meas.
- saffordi, Lesquereux, 1880, Coal Flora of Pa., p. 587, Coal Meas.
- schultzeanum, Gœppert & Berger, 1848, De Fruct., etc., p. 19, and Coal Flora of Pa., p. 819, Coal Meas.
- sigillaria, Dawson, 1863, Can. Nat., vol. 8, and Acad. Geol., p. 477, Coal Meas.
- tricuspidatum, Newberry, 1873, Ohio Pal., vol. 1, p. 368, Coal Meas.
- triloculare, Hildreth, 1837, (Carpolithes trilocularis,) Am. Jour. Sci., vol. 31, p. 29, Conglomerate and Low. Coal Meas.

FIG. 83.—*Trigonocarpum perpusillum*.

woodruffi, Moss, 1852, Proc. Acad. Nat. Sci., vol. 5, Coal Meas.

TRIPLHYLLOPTERIS, Schimper, 1874, Traité de Paléontologie Végétale, vol. 2, p. 40. [Ety. *tria*, three; *phyllon*, a leaf; *pteris*, a fern.] Lower pinnules subopposite, tripartite or trifoliate, upper ones simple, all narrowed or contracted to a flat, slightly decurring pedicel; veins all equal, simple or dichotomous, diverging fan-like. Type *T. lescuriana*.

FIG. 81.—*Triphylopteris cheathamii*.

because the latter was preoccupied as a generic name in the subkingdom Mollusca; but *Trochophyllum* was preoccupied for a genus of fossil corals by Edwards & Haine, in 1851.

clavatum, see *Annularia clavata*.

lineare, see *Plumalina linearis*.

ULODENDRON, Rhode, 1823, Beiträge z. Pflanz. d. Vorwelt. [Ety. *ula*, wood; *dendron*, tree.] Arboresecent; rarely branching; bearing in two opposite rows round or oval scars, im-

pressions of the base of strobiles, marked with concentric scales and a central mamillilla; leaves short lanceolate, leaf scars disposed in spiral, small, rhomboidal or sub-rhomboidal; fructifications in long, cylindrical strobiles. Type *U. majus*.

commutatum, Schimper, 1874, Pal. Veg., vol. 2, p. 40, Coal Meas.

ellipticum, Sternberg, 1838, Vers. Darst. Flora der Vorwelt, vol. 2, p. 186, and Coal Flora of Pa., p. 405, Coal Meas.

FIG. 85.—*Ulodendron elongatum*.

elongatum, Lesquereux, 1870, Geo. Sur. Ill., vol. 4, p. 437, Coal Meas.

flexuosum, see *Halonnia flexuosa*.

lindleyanum, Presl, 1833, in Sternberg, Vers. Darst. Flora der Vorwelt, p. 185, Coal Meas. *majus*, Rhode, 1823, Beitr. z. Pflanz. d. Vorw., pl. 3, fig. 1, and Coal Flora of Pa., p. 401, Coal Meas.

minus, Lindley & Hutton, 1831, Foss. Flora, vol. 1, p. 6, Coal Meas. *punctatum*, Lindley & Hutton, 1833, (Bothrodendron *punctatum*), Foss. Flora, vol. 2, p. 80, and Coal Flora of Pa., p. 405, Coal Meas.

VOLKMANNIA, Sternberg, 1823, Tent. Flor. Primord., p. 30.

FIG. 87.—*Walchia pinniformis*.

fertilis, Lesquereux, 1884, Coal Flora of Pa., p. 420, Coal Meas.

FIG. 88.—*Whitlieseyia elegans*.

prælonga, Lesquereux, 1880, (Calamotachys *prælongus*), Coal Flora of Pa., p. 59, Coal Meas.

FIG. 86.—*Volkmaninia fertilis*.

WALCHIA, Flora name. Branches of seed form. *gracilis*, p. 474, robusta, ward I.

THE A
Protista, C
Vertebrata.

The P
Porifera.

The R
generally m
naked eye.
mass, consta
processes, b
occur in ma
greatest dep
and have tin
in the sands
shell or inve
an extensile
organic bodi
mold), and
character it
fixed organs
Rhizopoda, s

"If the
living substa
as is necessar

WALCHIA, Sternberg, 1825, Vers. Darst. Flora der Vorwelt, p. 22. [Ety. proper name.] Arborescent, like Araucaria; branches with feathery foliage; strobiles oblong, cylindrical, or elongated, with ovate scales, sharp or lanceolate; seed minute, ovate. Type *W. pinniformis*.
gracilis, Dawson, 1863, (Araucarites *gracilis*.) Can. Nat., vol. 8, and Acad. Geol. p. 474, Coal Meas.
robusta, Dawson, 1871, Rep. on Prince Edward Island, p. 43, Coal Meas.

WHITTLESEYA, Newberry, 1874, Proc. Cleveland Acad. Sci., p. 43. [Ety. proper name.] Frond simple or pinnate, nerves fasciculate, confluent to the base, not dichotomous. Type *W. elegans*.
elegans, Newberry, 1874, Proc. Cleveland Acad. Sci., p. 43, Coal Meas.
integrifolia, Lesquereux, 1880, Coal Flora of Pa., p. 524, Coal Meas.
microphylla, Lesquereux, 1884, Coal Flora of Pa., p. 843, Coal Meas.
undulata, Lesquereux, 1880, Coal Flora of Pa., p. 525, Coal Meas.

ANIMAL KINGDOM.

THE Animal Kingdom is divided into seven Subkingdoms, viz.: Protozoa or Protista, Cœlenterata, Echinodermata, Molluscoidea, Mollusca, Articulata, and Vertebrata.

SUBKINGDOM PROTOZOA.

(*protos*, first; *zoon*, animal).

The Palæozoic Protozoa are included in two Classes, viz.: Rhizopoda and Porifera.

CLASS RHIZOPODA (*riza*, root; *pous*, foot).

The Rhizopoda are the simplest and lowest forms of animal life. They are generally microscopic, though some of them are more or less conspicuous to the naked eye. They abound in fresh-water ponds, where each consists of a shapeless mass, constantly changing its form, and shooting out and withdrawing finger-like processes, but visible only under the magnifying power of a microscope. They occur in marshes, ponds, lakes, and seas, and wherever dampness exists, from the greatest depths to the snow-line of the mountains. The greater portion are marine, and have tiny shells that enter into the composition of the ocean mud, and abound in the sands of every ocean shore. The simplest kinds are not provided with a shell or investing membrane, but consist of a fluid, viscid, albuminoid jelly, having an extensile and contractile power, which is regarded as the elementary basis of organic bodies in general. This jelly is called protoplasm (*protos*, first; *plasso*, I mold), and resembles in motive power the flesh of higher animals, from which character it is called sarcode (*sarx*, flesh; *eidos*, form). The protoplasm has no fixed organs of any kind, internal or external. Dr. Carpenter, speaking of the Rhizopoda, says:

"If the views which I have expressed as to the nature and relations of their living substance be correct, that substance does not present any such differentiation as is necessary to constitute what is commonly understood as 'organization' even

of the lowest degree and simplest kind; so that the physiologist has here a case in which those vital operations which he is accustomed to see carried on by an elaborate apparatus are performed without any special instruments whatever—a little particle of apparently homogeneous jelly changing itself into a greater variety of form than the fabled Proteus, laying hold of its food without members, swallowing it without a mouth, digesting it without a stomach, appropriating its nutritious material without absorbent vessels or a circulating system, moving from place to place without muscles, feeling (if it has any power to do so) without nerves, propagating itself without genital apparatus; and not only this, but in many instances forming shelly coverings of a symmetry and complexity not surpassed by those of any testaceous animals."

The fresh-water, shapeless, gelatinous mass is called the *Amœba*, and it shows a voracious disposition by seizing upon minute substances and appropriating them to the nutrition of its own jelly. This it does by surrounding and inclosing the food supply, which is retained until it is dissolved or the desired part appropriated.

The lowest Order of Rhizopoda has received the name of Monera (*moneres*, simple), of which Prof. Haeckel says:

"In a state of rest most Monera appear as small globules of slime, invisible, or barely visible, to the naked eye, and at most about the size of a pin-head. When the Moner moves, there are produced on the surface of the little slime-ball finger-like processes or very fine radiating threads, the so-called false feet, or pseudopods. The latter are simple continuous processes of the structureless, albumen-like mass of which the body consists. We are unable to perceive different parts in it, and we can obtain direct proof of the absolute simplicity of the semi-fluid mass of albumen; for, with the aid of the microscope, we can follow the Moner as it receives its nourishment. When minute bodies suitable for food, as, for instance, small particles of decayed organic bodies or microscopic plants and infusoria, accidentally come into contact with the Moner, they remain hanging to the sticky surface of the semi-fluid mass of slime, and here produce an irritation, which is followed by a strong afflux from the slimy mass of the body, and they become finally completely enclosed by it, or they are drawn into the body of the Moner by displacement of the several albuminous particles, and there digested, being absorbed by simple diffusion (endosmosis).

"Just as simple as is the nourishment is the mode of reproduction of these primitive beings, which one can not positively call animal or plant. All Monera propagate themselves only in an asexual manner by self-division. When such a speck—for example, a *Protamœba* or a *Protophagus*—has attained a certain size by the assimilation of foreign albuminous matter, it falls into two pieces; there is formed a constriction around the middle, which finally leads to the separation of the two halves. Each half becomes rounded, and then appears as an independent individual, which commences anew the simple play of the vital phenomena of nutrition and propagation. In other Monera (*Vampyrella*) the body, in the process of propagation, instead of two, falls into four equal parts; and in others again (*Protomonas*, *Protomyxa*, *Myxastrum*), at once into a large number of small globules of slime, each of which again, by simple growth, becomes like the parent body."

The marine Rhizopoda are usually furnished with a horny shell, and live, singly or socially, in shells having a series of chambers. The Sub-class Monothalamia

(*monos*, single shells) are those in which the single shells are composed of several processes by which the animal is attached to the calcareous skeleton. The communication between the individual animals is effected by the cells, and only a single shell is pierced by a single thread.

The shells are single and simple, and have a central death the chief part of the thickness, composed of the Archipelago, 2,000 feet deep.

The C. families of bearing an and swim incessant run a few are any of the *Nonionina* t. lantic City. In an ounce were 280,000. Ocean is so ally called shells of F. made of the mulite limes is the *Fusuli* of the Palæozoic For.

FAMILY
FAMILY
FAMILY
FAMILY
FAMILY
FAMILY

(*monos*, one; *thalamos*, chamber), includes those Rhizopoda which are inclosed in a single shell, and have a minute opening for the extrusion of the filamentous processes by which motion is effected. The Sub-class Polythalamia includes those having calcareous shells, consisting of a series of distinct chambers, which sometimes communicate with each other, and at other times appear to be completely closed up. Each chamber is supposed to contain an independent animal, though the individual animals may be so connected, through the openings communicating between the cells, as to constitute a common mass. In some genera each chamber presents only a single external opening, but in most genera the substance of the shell is pierced by minute pores, like a sieve, through which delicate filaments are protruded.

The Order Radiolaria (*radiolus*, a little ray,) includes many beautiful forms, living and swimming in vast multitudes near the surface of the ocean. Most of them have a complex silicious skeleton of great beauty of form and symmetry, and after death the skeletons sink to the bottom of the ocean, where they often furnish the chief part of the mud. On the island of Barbadoes, Tertiary strata 1,100 feet in thickness, consisting of marls, tripoli, and ferruginous sandstone, are largely composed of the silicious skeletons of Radiolaria. The Nicobar Islands of the Indian Archipelago, consisting of clays, marls, and arenaceous marls, to the extent of 2,000 feet in thickness of Tertiary age, are largely composed of the remains of this Order.

The Order Foraminifera (*foramen*, an aperture; *fero*, I bear,) includes all the families of Palæozoic Rhizopoda noticed in this work. They are marine shell-bearing animals, living at the bottom of oceans and seas, attached, free, or pelagic, and swimming on the surface of the water, from whence their dead shells form an incessant rain to the bottom of the ocean. They are generally microscopic, though a few are several inches in diameter. Some extinct genera are much larger than any of the living forms. Prof. Leidy obtained 18,700 shells of a single species of *Nonionina* from an ounce of mud scraped from the surface, between tides, at Atlantic City. In another sample, from Cape May, he obtained 38,400 shells; and in an ounce from the bathing beach at Newport, Rhode Island, he estimated there were 280,000 shells of several genera and species. The sediment of the Atlantic Ocean is so largely constituted of one kind of foraminiferous shell, that it is generally called Globigerina ooze. Common chalk is almost wholly composed of the shells of Foraminifera. The building stone of the city of Paris is almost wholly made of the shells of Foraminifera belonging to the Sub-order Miliola. The Nummulite limestone of different countries is composed of foraminiferous shells, and so is the Fusulina limestone of Carboniferous age. The microscopic genera and species of the Palæozoic rocks have not been much studied. The classification of the Palæozoic Foraminifera, so far as they have been investigated, is as follows:

FAMILY CALCISPHERIDÆ.—Calcisphæra.

FAMILY EOOZONIDÆ.—Eozoon.

FAMILY FUSULINIDÆ.—Fusulina, Loftusia, Moellerina.

FAMILY GLOBIGERINIDÆ.—Calcarina.

FAMILY LITUOLIDÆ.—Endothyra, Nodosinella, Valvulina.

FAMILY AFFINITY, UNCERTAIN.—Rhabdaria.

CLASS PORIFERA (*poros*, canal; *phero*, I bear).

The Porifera include the Sponges, and are not to be regarded as any more highly organized than the Rhizopoda. A sponge consists of a congeries of horny filaments, interlaced in every direction so as to form an intricate network of intercommunicating cells. Imbedded in these filaments, in the majority of sponges, are a number of minute needle-shaped, or forked, or radiated silicious, or calcareous particles of various forms, called spiculæ. The spiculæ may be acicular and pointed at both ends, or have a small knob at one end, while the opposite end is pointed; or one end may be a fork, with two or three prongs. The horny filaments, with their contained spiculæ, constitute the skeleton which supports the living sponge. The living sponge consists of a mere coating of gelatinous matter spread over all the filaments, of the consistence of the white of an egg, which runs freely away from the skeleton or framework of the sponge when taken out of the water. Under the microscope this gelatinous matter is found to consist of an aggregation of sarcode cells, and each cell appears to possess an independent existence; and even when detached from its fellows it has the power to move by the extension of its substance in various directions. In a living sponge there is an infinite number of minute holes, and a lesser number of larger openings. The water is imbibed through the smaller pores, and thrown out from the larger ones. The circulation results from the action of cilia, in much the same way motion is effected by the Rhizopoda.

Sponges attach themselves to all kinds of objects, whether fixed or floating. Some cover rocks and shells with a spongy incrustation; others hang from floating sea-weeds, and others shoot up branched stems, or a massive, globular framework. The *Cliona* is a boring sponge, that imbeds itself in shells or other calcareous substances. Sponges of the same species assume very different forms. In fact, there are no animals in which the variations are as great in a single species. They attain their greatest development in tropical seas, but occur in the most northern latitudes.

The genera and species of living sponges are largely founded upon the framework and spiculæ, and of course the same characters are sought in fossil sponges for the purpose of classification. Among the Palæozoic sponges, form is of much more importance than it is among living sponges, as we may believe, because we find so many specimens of the same form and size in a given species, not only at one locality, but at distant places, even hundreds or thousands of miles apart, in the same Group of rocks; as, for instance, *Astylospongia præmorsa*, on the Island of Gottland, in the Baltic Sea, and in Tennessee and Indiana. When Silurian sponges are silicified, the surface is generally very poorly preserved, and the spiculæ perfectly preserved; but calcareous and unsilicified specimens of the same species will show a well-preserved exterior and no spiculæ. It is therefore impossible to determine whether the sponge in its living state had calcareous or silicified spiculæ. In the fossilization of sponges and other bodies, and even long after fossilization has taken place, silica will be taken up, and lime will be deposited in its place in some waters; while in other waters lime will be taken up, and silica will be deposited in its stead. An original calcareous sponge, when converted into a silicious fossil, will preserve the spiculæ; but if a sponge bears silicious spiculæ, and is converted

ANT.]

into a cal
possible e
were conv
termination
specimens
The s

FAMIL

FAMIL

FAMIL

FAMIL

t

T

FAMIL

FAMIL

FAMIL

t

F

FAMIL

FAMIL

FAMIL

FAMIL

FAMIL

FAMIL

sp

FAMIL

FAMIL

st

ri

FAMIL

to

ANTHASPIDE

Geo. Sp
anthos, f
minutiv
supporte
stem; in
numerou
face for
through
the cha
radiating
tical seri
of spicul
face; sp
bifurcati
gle, and
ranged a
angular,
surface s
mal laye

into a calcareous fossil, the spiculæ will disappear in the coarser lime materials. A possible exception might exist if, in the process of change, the interior of the sponge were converted into calcespar. The spiculæ, therefore, are of importance in the determination of genera and species among Palæozoic sponges, only when silicified specimens can be obtained.

The arrangement of the Palæozoic sponges into families is as follows:

FAMILY ANTHASPIDELLIDÆ.—Anthaspidella, Climacospongia, Edriospongia, Streptosolen, Zittellella.

FAMILY ARCHÆOCYATHIDÆ.—Archæocyathus, Ethmophyllum.

FAMILY ASTRÆOSPONGIDÆ.—Astræospongia.

FAMILY ASTYLOSPONGIDÆ.—Astylospongia, Aulocopina, Calathium, Conopterium, Cyathospongia, Eospongia, Palæomanon, Palæospongia, Trachyum, Trichospongia.

FAMILY BEATRICIDÆ.—Beatricea.

FAMILY BRACHIOSPONGIDÆ.—Brachiospongia, Chirosporgia.

FAMILY DICTYOSPONGIDÆ.—Cleodictya, Cyathophycus, Dictyophyton, Ectenodictya, Lyriodictya, Phragmodictya, Physospongia, Protospongia, Rauffella, Rhombodictyon, Thamnodictya, Uphantænia.

FAMILY DYSTACTOSPONGIDÆ.—Dystactospongia, Heterospongia, Saccospongia.

FAMILY LEPTONITIDÆ.—Leptonitus.

FAMILY MICROSPONGIDÆ.—Hindia, Microspongia.

FAMILY PALÆACIDÆ.—Palæacis.

FAMILY PASCEOLIDÆ.—Pasceolus.

FAMILY PATTERSONIIDÆ.—Pattersonia.

FAMILY PHARETRONES.—Batospongia, Camarocladia, Cylindrocœlia, Streptospongia.

FAMILY RECEPTACULITIDÆ.—Cerionites, Receptaculites.

FAMILY STROMATOPORIDÆ.—Caunopora, Cœnostroma, Cryptozoon, Dictyostroma, Megastroma, Strephochetus, Stromatocerium, Stromatopora, Syringostroma.

FAMILY AFFINITY UNCERTAIN.—Astroconia, Fungispongia, Lepidolites, Leptomitius.

ANTHASPIDELLA, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 256. [Ety. *anthos*, flower; *aspis*, shield; *ellus*, diminutive.] Saucer or funnel-shaped, supported by a short, subcylindrical stem; inosculating, radiating channels numerous, and those on the upper surface form radical canals that pass through the sponge-wall, and open into the channels of the lower surface; radiating canals closely arranged in vertical series, separated by vertical sheets of spicules; oscula on the upper surface; spicules bifid at each end, the bifurcations directed nearly at right angle, and slightly curving, and so arranged as to leave minute canals of triangular, quadrate, or polygonal form; surface sometimes covered with a dermal layer. Type A. *mammulata*.

fenestrata, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 264. Trenton Gr. *firma*, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 263, Trenton Gr. *florifera*, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 259, Trenton Gr. *grandis*, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 262, Trenton Gr. *magnifica*, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 265, Trenton Gr. *mammulata*, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 258, Trenton Gr. *obliqua*, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 265, Trenton Gr. *parvistellata*, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 260, Trenton Gr. *scutula*, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 261, Trenton Gr.

Archæocyathellus, Ford, 1873, Am. Jour. Sci. and Arts, 3d ser., vol. 6, p. 135, syn. for *Ethmophyllum*.

ARCHÆOCYATHUS, Billings, 1861, Pal. Foss., vol. 1, p. 3, and 354. [Ety. *arche*, beginning; *cyathus*, cup.] An elongated, cylindrical, sponge-like body; large end open; central cavity lined by an endotheca and external surface by an epitheca; intervening space being filled with poriferous and cellular tissue; walls perforated. Type *A. atlanticus*.

FIG. 89.—*Archæocyathus atlanticus*.
a, reduced; b, transverse section.

18.

atlanticus, Billings, 1861, Pal. Foss., vol. 1, p. 5, Up. Taconic.

billingsi, Walcott, 1886, Bull. U. S. Geo. Sur., No. 30, p. 74, Up. Taconic.

minganensis, see *Ethmophyllum minganensis*, though Hinde has made it the type of a new genus, *Archæoscyphia*.

profundus, see *Ethmophyllum profundum*.
rensselericus, see *Ethmophyllum rensselericum*.

ASTRÆOSPONGIA, Roemer, 1860, Sil. Fauna d. West Tenn., p. 13. [Ety. *aster*, star; *spongia*, sponge.] Globular or disk-like, free sponge composed of regular star-shaped spicules, without order, no epitheca or canals. Type *A. meniscus*.

hamiltonensis, Meek & Worthen, 1866, Proc. Chi. Acad. Sci., vol. 1, p. 12, Ham. Gr.

meniscus, Roemer, 1848, (*Blumenbachium meniscus*), Leonh. and Bronn's Jahrb., p. 683, Niagara Gr.

FIG. 90.—*Astræospongia meniscus*.

ASTROCONIA, Sollas, 1881, Quar. Jour. Geo. Soc. Lond., vol. 37, p. 254. [Ety. *aster*, star; *konis*, dust.] Founded upon the appearance of various spiculæ in a grayish silicious dolomite. Characters not distinct. Type *A. granti*. The name was preoccupied by Edwards & Haime in 1848.

granti, Sollas, 1881, Quar. Jour. Geo. Soc. Lond., vol. 37, p. 254, Niagara Gr.

ASTYLOSPONGIA, Roemer, 1860, Sil. Fauna d.

West Tenn., p. 7. [Ety. *astylos*, without a pillar; *spongia*, sponge.] Globular or disk-like, free sponge; inner texture formed of small, regular, star-shaped spicules, connected by their rays; canals running from the center to the surface crossed by concentric canals. Type *A. præmorsa*.

bursa, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 105, Niagara Gr.

christiana, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 344, Niagara Gr.

imbricato-articulata, Roemer, 1848, (*Siphonia imbricato-articulata*), Leonh. and Bronn's Jahrb., p. 685, and Sil. Fauna d. West Tenn., p. 12, Niagara Gr.

inciso-lobata, Roemer, 1848, (*Spongia inciso-lobata*), Leonh. and Bronn's Jahrb., p. 685, and Sil. Fauna d. West Tenn., p. 11, Niagara Gr.

inornata, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 70, syn. for *Hindia fibrosa*.

parvula, Billings, 1861, Pal. Foss., vol. 1, p. 20, Trenton Gr.

perryi, Billings, 1861, Geo. Vermont, p. 29, 957, Black Riv. Gr.

præmorsa, Goldfuss, 1826, (*Siphonia præmorsa*), Petref. Germ., p. 17, and Sil. Fauna d. West Tenn., p. 8, Niagara Gr.

FIG. 91.—*Astylospongia præmorsa*. Vertical section, showing cup.

stellatim-sulcata, Roemer, 1848, (*Spongia stellatim-sulcata*), Leonh. and Bronn's Jahrb., p. 686, and Sil. Fauna West Tenn., p. 11, Niagara Gr.

AULOCOPINA, Billings, 1875, Can. Nat. and Geol., vol. 7, p. 230. [Ety. *aulokopeo*, cut into pipes.] Elongate, ovate, or pyriform; upper face concave, with an osculum in the center, from which ridges radiate over the surface and descend to the base; the osculum is the opening of a central cavity, from which smaller branching canals radiate. Type *A. granti*.

granti, Billings, 1875, Can. Nat. and Geol., vol. 7, p. 231, Niagara Gr.

BATOSPONGIA, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, p. 246. [Ety. *batos*, prickly bush; *spongia*, sponge.] Subhemispherical or subglobose, consisting of small, inosculating, subcylindrical or flattened

bran-
base
whic
work
bran
bifid,
spica
spicata,
Ill., v
BEATRICE
Geo.
name
Hyatt
to be
and h
genus
a hor
family
long,
B. noc
12. nodulos
Geo.
Hud.
13. undulata
Geo. S
BELEMNI
Sur. Ill
a dart
elonga
upwar
base;
other
cicular
fascicula
Ill., vo
Blumenbach
sectiles
meniscus,
BRACHIOSP
Sci. an
[Ety. b
A shor
throwi
are cl
skeletal
ing a n
specim
face th
digitata

FIG. 92.—Brac

digitata,
Geo. of
hoveyi, Ma
Sci., p.

branches, which arise from a reticulated base; base covered with a dermal layer, which exhibits on its inner side a network of substellate or irregularly branched spicule fiber; spicules acerate, bifid, trifid, or four-rayed. Type B. spicata.

spicata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 248, Coal Meas.

BEATRICEA, Billings, 1875, Rep. of Progr. Geo. Sur. Can., p. 343. [Ety. proper name.] This genus was supposed by Hyatt (Am. Jour. Sci. and Arts, 1865,) to belong to the class Cephalopoda, and he proposed a new order for the genus, to-wit: Ceriolites, from *kerion*, a honey-comb; *lithos*, a stone; and a family Ceriolidae. They are, however, long, cylindrical spongeoid bodies. Type B. nodulosa.

22. nodulosa, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 344, Trenton and Hud. Riv. Gr.

23. undulata, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 344, Trenton Gr.

BELEMNOSPONGIA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 248. [Ety. *belemnos*, a dart; *spongia*, sponge.] Composed of elongate acerate spicules, which radiate upward and outward from a pointed base; spicules large, and joined to each other by short processes. Type B. fascicularis.

fascicularis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 248, Burlington Gr.

Blumenbachium, König, 1820, Icones fossiles, sectiles.

meniscus, see *Astræospongia meniscus*.

BRACHIOSPONGIA, Marsh, 1867, Am. Jour. Sci. and Arts, 2d ser., vol. 44, p. 88. [Ety. *brachium*, arm; *spongia*, sponge.] A short vase or hollow central nucleus, throwing out large, hollow arms, which are closed at the distal extremities; skeleton comparatively thin and bearing a network of spicules; all observed specimens are silicious, and outer surface therefore destroyed. Type B. digitata.

FIG. 92.—*Brachiospongia digitata*. $\frac{1}{4}$ diam., showing large gastral cavity.

digitata, Owen, 1857, (*Scyphia digitata*), Geo. of Ky., vol. 2, p. 111, Trenton Gr. *hoveyi*, Marsh, 1874, Trans. Kansas Acad. Sci., p. 344, syn. for B. digitata, but

founded on a specimen having twelve arms.

lyoni, Marsh, 1867, Am. Jour. Sci. and Arts, 2d ser., vol. 44, p. 88, syn. for B. digitata, but founded on a specimen having eleven arms.

roemerana, Marsh, 1867, Am. Jour. Sci. and Arts, 2d ser., vol. 44, p. 88, syn. for B. digitata.

CALATHIUM, Billings, 1865, Pal. Foss., vol. 1, p. 208. [Ety. *kalathos*, a small wicker basket.] Cylindro-turbinate in form, perforated by small canals arranged in longitudinal and transverse rows; apertures round, oval, or quadrangular; cup deep. Type C. formosum.

affine, Billings, 1865, Pal. Foss., vol. 1, p. 209, Quebec Gr.

anstedii, Billings, 1865, Pal. Foss., vol. 1, p. 201 and 337, Quebec Gr.

canadense, Billings, 1865, Pal. Foss., vol. 1, p. 377, Chazy Gr.

fittoni, Billings, 1865, Pal. Foss., vol. 1, p. 211, Quebec Gr.

formosum, Billings, 1865, Pal. Foss., vol. 1, p. 209, Quebec Gr.

infelix, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 274, Trenton Gr.

FIG. 94.—*Calathium formosum*.

pannosum, Billings, 1865, Pal. Foss., vol. 1, p. 335, Quebec Gr.

paradoxicum, Billings, 1865, Pal. Foss., vol. 1, p. 358, Calcif. Gr. Hinde, in 1889, Quar. Jour. Geo. Soc., p. 144, made this species the type of a new genus, *Nipterella*.

CALCARINA, D'Orbigny, 1826, Tableau Methodique de la Classe des Cephalopodes, in Annales des Sciences Naturelles, tome 7. [Ety. *calcis*, limestone.] Free, convoluted, depressed, spire-coiled, supplemental growths of the interior shell, aperture slit in the terminal chamber close to the penultimate convolution. A living genus in tropical seas.

ambigua, Brady, 1878, Monograph of Carboniferous and Permian foraminifera, p. 141, Carboniferous.

CALCISPHERA, Williamson, 1880, Mem. Org. of the plants of the Coal Meas., pt. 10. [Ety. *calcis*, limestone; *sphæra*, sphere.] A minute globular test, having an aperture; wall composed of minute calcareous grains. Type C. robusta.

24.

25.

26.

27.

28.

29.

30.

robusta, Williamson, 1880, Mem. Org. of the Coal Meas., pt. 10, Up. Held Gr.

FIG. 95.—*Culestheria robusta*. *a*, natural size; *b*, magnified, showing sculpture; *c*, showing aperture; *d*, magnified, showing aperture; *e*, section of wall magnified.

CAMAROCLADIA, Ulrich & Everett, (in press.) Geo. Sur. Ill., vol. 8, p. 280. [Ety. *kamara*, arching chamber; *klados*, twig.] Small, subcylindrical branching stems; interior canals irregular, separated by thin, cribose walls; spicules three-rayed. Type *C. dichotoma*.

dichotoma, Ulrich & Everett, (in press.) Geo. Sur. Ill., vol. 8, p. 281, Trenton Gr.

CAUNOPORA, Phillips, 1841, Pal. Foss., Cornwall and Devon. and W. Somerset, p. 18. [Ety. *chaunos*, loose; *poros*, perforation.] Amorphous, composed of concentric or nearly plain masses, perforated by flexuous or vermiform small tubuli, and by larger, straight, subparallel or radiating open tubes, persistent through the mass. Type *C. placenta*.

hudsonica, Dawson, 1879, Quar. Jour. Geo. Soc., vol. 35, p. 52, Niagara Gr. *incrustans*, Hall & Whitfield, 1873, (Stromatopora incrustans,) 23d Rep. N. Y. St. Mus. Nat. Hist., p. 227, Chemung Gr.

mirabilis, Spencer, 1884, Bull. No. 1, Univ. St. Mo., p. 47, Niagara Gr. *plannata*, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 228, Chemung Gr.

walkeri, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 46, Niagara Gr.

FIG. 96.—*Caunopora walkeri*, vertical and horizontal section enlarged.

CERIONITES, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 346. [Ety. *kerion*, honey-comb; *lithos*, stone.] Founded upon casts apparently holding an intermediate position between *Pasceolus* and *Receptaculites*. The pits are hexagonal and upon the convex side, perforated in the center by a minute circular opening, while those upon the

under side are imperforate. Type *C. dactyloides*. *dactyloides*, Owen, 1844, (Lunulites dactyloides,) Rep. on Min. Lands, p. 69, Niagara Gr.

CHIROSPONGIA, n. gen. FIG. 97.—*Cerionites dactyloides*.

[Ety. *cheir*, hand; *spongia*, sponge.]

General form hand-like, or somewhat like a compressed goblet; composed of internal filamentous or fibrous substance, which is covered with a thin, lobed, vesicular parenchyma; it was firmly fastened by an expanded base to a solid rock or the sea-bottom; above the base it is a flattened oboconoidal cup, with a deep sulcus down the middle of each side, bringing the sides nearly together; on each side of the sulcus the interior of the sponge is hollow, showing a large gastral cavity; the whole skeleton is openly vesicular or porous. The type species is silicified, and does not show the surface markings, but a calcareous specimen, supposed to belong to the same genus, is finely papillated. No microscopic sections have been made to ascertain the character of the spicules, but doubtless both parenchyma and fibrous substance bear spicules similar to those of *Brachiospongia*. In the surface lobes and filaments it resembles *Pattersonia*, but is distinguished by its vesicular and porous substance and coarser filaments. In its large gastral cavity, thin skeleton, and vesicular parenchyma, it resembles *Brachiospongia*. Type *C. wenti*.

FIG. 98.—*Chirosporgia wenti*.

faberi, n. sp. This species is founded upon a calcareous fragment of the parenchyma, about one-third of which is shown in the figure. It is thin, and

FIG. 99.—*Chirosporgia wenti*.

CLEODICTYA, Mus. N. closed u expanded globose a row of peripher above, a or slight compose rayed sp rods. T.

belonged to the side of a large, hollow specimen. The semi-elliptical lobes are nearly equal in size, and regularly distributed in rows over the surface. The surface is reticulated with fine papillae, presenting to the naked eye the appearance of a bryozoan. The illustration shows rhomboidal depressions instead of papillae. Found associated with Pattersonia and fragments of sponge filaments near the middle of the Hud. Riv. Gr., about 350 feet above low-water mark at Cincinnati. Collector, Charles Faber.

went, n. sp. This species is founded upon a silicified specimen having the characters above ascribed to the genus, and being well illustrated in the figure. The fibrous substance shown in the sulcus formed the basal attachment, as the parenchymatous surface tissue does not appear at the bottom. The lobes are large, somewhat semi-elliptical in outline, of unequal size, and irregularly disposed, but not pendent as in Pattersonia. The substance of the filaments and parenchyma, as shown, where broken off and weathered at the top and bottom of the specimen, is openly vesiculose or irregularly porous, resembling to the naked eye somewhat the appearance of Alveolites goldfussi. The species is named in honor of Mr. C. E. Went, of Frankfort, Ky., who found it in the Trenton Group near that city.

FIG. 99.—*Chirosporgia faberi*; reticulated depressions should indicate papillae.

CLEODICTYA, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 467. [Ety. *kleo*, closed up; *dictyon*, net.] Frond rapidly expanding from the base to a subglobose or hemispherical form, bearing a row of large, rounded nodes on the periphery; tube abruptly contracted above, and extending in a cylindrical or slightly expanded form. Substance composed of regular lattice-work of six-rayed spicules and bundles of acicular rods. Type C. gloriosa.

gloriosa, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 479, Keokuk Gr. mohri, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 479, Keokuk Gr. *Cnemidium*, Goldfuss, 1826, Petref. Germ., p. 15. [Ety. *knemidos*, armor for the legs, a sort of boot.] Type C. lamellosum. *trentonensis*, see *Palaeospongia trentonensis*.

CENOSTROMA, Winchell, 1867, Proc. Am. Ass. Ad. Sci., p. 91. [Ety. *koinos*, shared in common; *stroma*, layer.] Distinguished from *Stromatopora* by the absence of central, simple, radiating tubes, which in this genus is represented by a group of more or less divergent ascending tubuli, so that the surface of the last layer presents eminences, not with a single large pore at the summit, but with several small pores diverging from their sides. Type C. monticuliferum.

botryoideum, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 50, Niagara Gr.

constellatum, Hall, 1852, (*Stromatopora constellata*) Pal. N. Y., vol. 2, p. 324, Coralline limestone, Niagara Gr.

galtense, Dawson, 1879, Quar. Jour. Geo. Soc., vol. 35, p. 52, Guelph Gr.

monticuliferum, Winchell, 1866, (*Stromatopora monticulifera*) Rep. Low. Penin. Mich., p. 91, Ham. Gr.

pustuliferum, Winchell, 1866, (*Stromatopora pustulifera*) Rep. Low. Penin. Mich., p. 90, Ham. Gr.

ristigouchense, Spencer, 1884, Bull. No. 1, Univ. St. Mo., p. 49, Low. Held. Gr. solidulum, Hall & Whitfield, 1873, (*Stromatopora solidula*) 23d Rep. N. Y. St. Mus. Nat. Hist., p. 227, Chemung Gr.

CONOPTERUM, Winchell, 1865, Proc. Acad. Nat. Sci., p. 110. [Ety. *konos*, cone; *pteron*, cnp.] Cells crowded, inseparable, rapidly enlarging, walls marked by vertical striae, and a few pores communicate between the cells; epitheca exterior. Type C. effusum.

effusum, Winchell, 1865, Proc. Acad. Nat. Sci., p. 111, Waverly Gr. or Lithographic limestone.

Coscinopora infundibuliformis, see *Receptaculites infundibuliformis*.

Coscinopora sulcata, Owen, 1844, see *Receptaculites oweni*.

CRYPTOZOON, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 95. [Ety. *kryptos*, hidden; *zoon*, animal.] Composed of irregular, concentric laminae, resembling *Stromatopora*, substance traversed by minute canals, which branch and anastomose irregularly. Type C. proliferum. minnesotense, Winchell, 1886, 14th Ann. Rep. Geo. Minn., p. 313, Calciferous Gr. proliferum, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 95, Calciferous

FIG. 100.—*Cenostroma constellatum*, horizontal section.

- CYATHOPHYCUS**, Walcott, 1879, Trans. Alb. Inst., vol. 10, p. 18. [Ety. *kuathos*, cup; *phukos*, sea-weed.] Hollow, cyathiform, with a reticulated structure. Type C. reticulatum.
- reticulatum, Walcott, 1879, Trans. Alb. Inst., vol. 10, p. 18, Utica Slate.
- subsphericum, Walcott, 1879, Trans. Alb. Inst., vol. 10, p. 19, Utica Slate.
- CYATHOSPONGIA**, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 15. [Ety. *kuathos*, cup; *spongia*, sponge.] Body solid, turbinate, cyathiform; structure similar to *Astylospongia*. Type C. excrescens.
- excrescens, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 15, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 419, Niagara Gr.
- CYLINDROCELLA**, Ulrich, 1889, Am. Geo., vol. 3, p. 245. [Ety. *kulindros*, cylinder; *koiolos*, belly.] Cylindrical; central cloaca; walls thick, radiating canals. Type C. endoceroidea.
- covingtonensis, Ulrich, 1889, Am. Geo., vol. 3, p. 247, Hud. Riv. Gr.
- endoceroidea, Ulrich, 1889, Am. Geo., vol. 3, p. 246, Trenton Gr.
- minnesotensis, Ulrich, 1889, Am. Geo., vol. 3, p. 248, Trenton Gr.
- minor, Ulrich, 1889, Am. Geo., vol. 3, p. 248, Trenton Gr.
- Dentalina*, D'Orbigny, 1826, Ann. Des. Sci. Nat., t. 7, p. 89. [Ety. *dentale*, tooth; *inus*, implying resemblance.]
- priscilla*, see *Nodosinella priscilla*.
- DICTYOPHYTON**, Hall, 1863, 16th Rep. N. Y. St. Mus., p. 87. [Ety. *dictyon*, net; *phyton*, plant.] Turbinate or infundibuliform, with nodose or coralline protuberances or hollow stems externally, and marked by minute rectangular spaces, and consisting of a reticulate envelope. Type D. filitextile.
- abacus, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 474, Waverly Gr.
- annulatum, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 90, Chemung Gr.
- baculum, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 471, Chemung Gr.
- becki, Conrad, 1837, (Lithodictyon becki,) Ann. Rep. N. Y., p. 167, and Pal. N. Y., vol. 2, p. 6, Medina Sandstone.
- catilliforme*, see *Phragmodictya catilliformis*.
- cinctum, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 472, Chemung Gr.
- conradi, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 89, Chemung Gr.
- cylindricum, Whitfield, 1881, Bull., No. 1, Am. Mus. Nat. Hist., p. 19, Keokuk Gr.
- fenestratum, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 90, Chemung Gr.
- filitextile, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 88, Chemung Gr.
- hamiltonense, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 468, Ham. Gr.
- irregular, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 470, Chemung Gr.

newberryi, see *Thamnodictya newberryi*.

nodosum, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 91, Chemung Gr.

parallelum, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 471, Chemung Gr.

patulum, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 469, Chemung Gr.

prismaticum, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 469, Chemung Gr.

ramosum, Lesquereux, 1884, Coal Flora of Pa., p. 827, Up. Chemung Gr.

redfieldi, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 88, Waverly Gr.

rude, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 90, Chemung Gr.

sacculus, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 473, Waverly Gr.

telum, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 470, Chemung Gr.

tenu, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 474, Waverly Gr.

tuberosum, Conrad, 1842, (Hydnoceras tuberosum,) Jour. Acad. Nat. Sci. Phil., vol. 8, p. 267, Chemung Gr.

DICTYOSTROMA, Nicholson, 1875, Ohio Pal.,

vol. 2, p. 254. [Ety. *dictyon*, net; *stroma*, layer.] Allied to *Stromatopora*, but the upper surface of each lamina is developed into conical points, which support the lamina above instead of pillars. The laminae have horizontal canals, and are probably minutely perforate. Type D. undulatum.

reticulatum, Spencer, 1884, Bull. Mus. Univ. St. Mo., p. 51, Niagara Gr.

undulatum, Nicholson, 1875, Ohio Pal., vol. 2, p. 254, Niagara Gr.

FIG. 101.—*Dictyostroma undulatum*.

FIG. 102.

Dystactospongia insolens.

DYSTACTOSPONGIA, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 42. [Ety. *dystaktos*, hard to arrange; *spongia*,

ECT.—ET
spong
tache
work
insol
insolen
Nat.
minima
p. 242
minor,
Sur. I
rudis,
Sur. I
Ectenodictya
St. M
stretc
late f
planar
ing ra
E. im
burling
Y. St.
Gr.
excentric
St. M
Gr.
expansa
Mus. I
implexa
Mus. I
EDRIOSPON
Geo.
edri
ive, l
sides
canals
ones;
ules;
Type I
basalis,
Sur. I
ENDOTHYR
Polyte
p. 279.
Free,
merou
forate,
mani.
baileyi, I

FIG. 103.
Endothyra
baileyi,
magnified.
roemeri,
p. 19, C
varians, I
19, Cha
Eozoon, Da
2d ser
zoon, a
patches
at the

sponge.] Massive, hemispherical, attached with a strong radiating framework. Structure vesicular. Type *D. insolens*.

insolens, S. A. Miller, 1882, Jour. Clin. Soc. Nat. Hist., vol. 5, p. 43, Hud. Riv. Gr. minima, Ulrich, 1880, Am. Geol., vol. 3, p. 243, Hud. Riv. Gr.

minor, Ulrich & Everett, (in press.) Geo. Sur. Ill., vol. 8, p. 278, Trenton Gr.

rudis, Ulrich & Everett, (in press.) Geo. Sur. Ill., vol. 8, p. 279, Trenton Gr.

ECTENODICTYA, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 406. [Ety. *ektense*, stretched out; *dictyon*, net.] A reticulate frond irregularly expanded or explanate; reticulation irregular presenting radiating and concentric striae. Type *E. implexa*.

burlingtonensis, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 476, Waverly Gr.

excentrica, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 476, Keokuk Gr.

expansa, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 475, Waverly Gr.

implexa, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 475, Waverly Gr.

ENDOSPONGIA, Ulrich & Everett, (in press.) Geo. Sur. Ill., vol. 8, p. 271. [Ety. *edrion*, a seat; *spongia*, sponge.] Massive, lobate, attached by a broad base; sides irregularly dented; radiating canals, connected by tortuous, vertical ones; minute canals formed by spicules; sides covered with a dermal layer. Type *E. basalis*.

basalis, Ulrich & Everett, (in press.) Geo. Sur. Ill., vol. 8, p. 272, Trenton Gr.

ENDOTHYRA, Phillips, 1845, Proc. Geol. and Polytech. Soc. W. Riding Yorks., vol. 2, p. 279. [Ety. *endos*, within; *thura*, door.] Free, spiral, rotaliform, segments numerous, texture subarenaceous, imperforate, aperture simple. Type *E. bowmani*.

baileyi, Hall, 1858, (*Rotalia baileyi*), Trans. Alb. Ins., vol. 4, p. 34, and 1882, Bull. Mus. Nat. Hist., p. 42, Warsaw Gr.

FIG. 103.
Endothyra baileyi, magnified.

Eospongia, Billings, 1861, Pal. Foss., vol. 1, p. 18. [Ety. *eos*, dawn; *spongia*, sponge.] Subglobular, pyriform or subhemispherical, not free, pores radiating irregularly from the central axis; cup of variable depth. Type *E. roemerii*.

roemerii, Billings, 1861, Pal. Foss., vol. 1, p. 19, Chazy Gr.

varians, Billings, 1861, Pal. Foss., vol. 1, p. 19, Chazy Gr.

Eozoon, Dawson, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 54. [Ety. *eos*, dawn; *zoon*, animal.] Massive, in large sessile patches or irregular cylinders, growing at the surface, by the addition of suc-

cessive laminae, internally, the chambers are flattened, irregular, with numerous rounded extensions, and separated by walls of variable thickness, penetrated by septal orifices irregularly disposed; thicker parts of the walls with fine branching tubuli; the appearance to the naked eye is something like *Stromatopora*. Type *E. canadense*.

canadense, Dawson, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 54, Laurentian. The most ancient organism.

FIG. 104.—*Eozoon canadense*.

ETHMOPHYLLUM, Meek, 1868, Am. Jour. Sci. and Arts, 2d ser., vol. 45, p. 62. [Ety. *ethmos*, sieve; *phyllon*, plant.] Body simple, elongate, turbinat, cup-shaped, clavate or cylindro-conical, curved or straight, corrugated, lobed, or ribbed, penetrated by round or oval pores, in vertical or horizontal rows; vertical septa numerous, originating at the outer wall, and extending to the inner one, poriferous; inner wall with or without vesicular tissue, extending into the central cup; series of septa and walls sometimes repeated; spiculae branching. Type *E. whitneyi*. *gracile*, Meek, syn. for *E. whitneyi*.

FIG. 105.—*Ethmophyllum profundum*. Longitudinal and transverse section of a fragment.

minganense, Billings, 1859, (*Petraia minganensis*), Can. Nat. and Geo., vol. 4, p. 346, and Pal. Foss., vol. 1, p. 354. Calciferous Gr. Hinde, in 1889, Quar. Jour. Geo. Soc., p. 142, proposed this species as the type of a new genus *Archaeoscyphia*.

32.

profundum, Billings, 1861, (Archæocyathus profundus,) Pal. Foss., vol. 1, p. 4, Up. Taconic.

FIG. 106.—Ethmophyllum profundum. Base of attachment.

rarium, Ford, 1878, (Protocyathus rarus,) Am. Jour. Sci. and Arts, 3d ser., vol. 15, p. 124, Up. Taconic. rensselearicum, Ford, 1873, (Archæocyathus rensselearicus,) Am. Jour. Sci. and Arts, 3d ser., vol. 5, p. 211, Up. Taconic. whitneyi, Meek, 1868, Am. Jour. Sci. and Arts, 2d ser., vol. 45, p. 62, Up. Taconic.

FUNGISPONGIA, Ringueberg, 1884, Proc. Acad. Nat. Sci., p. 147. [Ety. fungus, a mushroom; spongia, a sponge.] Definition very poor. Type *F. irregularis*, Ringueberg, 1884, Proc. Acad. Nat. Sci., p. 147, Clinton Gr. Very poorly defined.

FUSULINA, Fischer, 1837, Oryct. du Gouv. de Moscou., p. 126. [Ety. fusus, spiralis, little.] Snell fusiform, symmetrically involute, surface furrowed coincident with the septa within; aperture a narrow slit in the middle part, foramina passing through the walls; septa widening toward the extremities. Type *F. cylindrica*.

cylindrica, Fischer, 1837, Oryct. du Gouv. de Moscou., p. 126, Coal Meas.

FIG. 107.—Fusulina cylindrica. Natural size, magnified, and transverse section.

cylindrica var. ventricosa, see *F. ventricosa*. depressa, Fischer, 1837, Oryct. du Gouv. de Moscou., p. 127, Coal Meas.

elongata, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 297. Permian Gr. gracilis, Meek, 1864, Pal. of California, vol. 1, p. 4, Coal Meas.

hyperborea, Salter, 1855, Belcher's Last Arctic Voyage, vol. 2, p. 380, Carboniferous.

robusta, Meek, 1864, Pal. California, vol. 1, p. 3, Coal Meas.

ventricosa, Meek & Hayden, 1864, Pal. Upper Mo., p. 14, Coal Meas.

HETEROSPONGIA, Ulrich, 1889, Am. Geol., vol. 3, p. 239. [Ety. heteros irregular; spongia, sponge.] Sublobate, compressed branches, covered with mouths of tortuous canals; skeleton composed of loosely interwoven spicule fibers. Type *H. subramosa*.

aspera, Ulrich, 1889, Am. Geol., vol. 3, p. 241, Hud. Riv. Gr.

knottii, Ulrich, 1889, Am. Geol., vol. 3, p. 241, Hud. Riv. Gr.

subramosa, Ulrich, 1889, Am. Geol., vol. 3, p. 240, Hud. Riv. Gr.

HINDIA, Duncan, 1879, Ann. and Mag. Nat. Hist., 5th ser., vol. 4, p. 91. [Ety. proper name.] Free, spheroidal, without involution of texture; small central space occupied by spicules which form a series of bifurcating, long, straight canals, that open at the surface; spicules more or less in shape of a stemmed tripod, with four limbs, and swollen or fringed at the ends. Type *H. fibrosa*. This may be a synonym for *Microspongia*; but as the latter is calcareous, and the spicules have not been determined, both generic names are retained.

fibrosa, Roemer, 1860, (Calamopora fibrosa,) Sil. Fauna W. Tenn., p. 20, Niagara Gr. inaequalis, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 275, Trenton Gr.

spheriodalis, Duncan, 1879, Ann. and Mag. Nat. Hist., 5th ser., vol. 4, p. 91, syn. for *H. fibrosa*.

parva, see *Microspongia parva*.

HYSTRISPONGIA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 245. [Ety. hystrix, porcupine; spongia, sponge.] Subglobular or ovoid; spicules arranged radiately from the base, most of them biacerate and taper each way to pointed ends, some trifid at one end, and others four-rayed. Type *H. carbonaria*.

carbonaria, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 245, Coal Meas.

Ischadites tessellatus, see *Receptaculites tessellatus*.

LASTIOCLADIA, Hinde, 1884. [Ety. lasios, shaggy; klados, twig.] Skeleton composed of elongate, slender, straight, acerate spicules, pointed at both ends. Type *L. compressa*.

hindii, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 249, Keokuk Gr.

LEPIDOLITES, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 20. [Ety. lepis, scale; lithos, stone.] Subspherical or subcylindrical bodies, hollow within and consisting of exteriorly imbricating scales. Type *L. dickhauti*. The name was preoccupied in mineralogy. *dickhauti*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 21, Hud. Riv. Gr. *elongatus*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 22, Hud. Riv. Gr. This is not distinct from *L. dickhauti*.

LEPTOMITUS, Walcott, 1886, Bull. U. S. Geo. Sur. No. 30, p. 89. [Ety. leptos, fine; mitos, thread.] Elongate bodies, formed of fine, thread-like, longitudinal lines, apparently imbedded in a delicate membrane, slowly expanding from a narrow base. Type *L. zitteli*.

zitteli, Walcott, 1886, Bull. U. S. Geo. Sur. No. 30, p. 89, Georgia Gr.

Leptopteris

p. 238

Obcor

thin,

L. ma

mammif

3, p. 2

LEPTUSIA,

Roy.

Small

tical

contin

with

Type

columb

Geo.

Lunulites

tyloid

LYRODICT

St. M

lyre;

late f

ules,

bands

altern

minge

rominge

St. M

MEGASTRO

path

great;

Strom

memb

ing in

other

Type

laminos

path M

Microspon

Cin. S

micros

no ex

openin

gregari

FIG. 108.

spongia

MELLERIN

Pal. E

sisting

chamf

inner

outer

ing, s

where

greeni

greeni,

35, U

NODOSINE

Carb.

[Ety.

Free,

constr

Leptopteron, Ulrich, 1889, Am. Geol., vol. 3, p. 239. [Ety. *leptos*, thin; *poterion*, cup.] Obconical, annulated free sponge; wall thin, outer surface reticulated. Type *L. mammiferum*. Not well defined.

mammiferum, Ulrich, 1889, Am. Geol., vol. 3, p. 239, Hud. Riv. Gr.

Leptostia, Carpenter & Brady, 1869, Trans. Roy. Soc., p. 742. [Ety. proper name.] Small foraminifer, with oval or elliptical test, consisting, primarily, of a continuous lamina coiled upon itself, with interspaces divided into chambers. Type *L. persica*.

columbiana, Dawson, 1879, Quar. Jour. Geo. Soc., vol. 35, p. 74, Coal Meas.

Lunulites ? *ductyloides*, see *Cerionites ductyloides*.

LYRODICTYA, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 466. [Ety. *lyra*, lyre; *dictyon*, net.] Cyathiform, reticulate fronds composed of stellate spicules, with broad, strong, longitudinal bands of acicular spicules, showing an alternating bifurcation. Type *L. romingeri*.

romingeri, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 476, Keokuk Gr.

MEGASTROMA, Dawson, 1883, Report on Redpath Mus. No. 2, p. 12. [Ety. *megas*, great; *stroma*, layer.] Somewhat like *Stromatopora*; layers consisting of two membranes, beset with spicules, pointing inwards like two brushes facing each other; membranes porous or reticulate. Type *M. laminosum*.

laminosum, Dawson, 1883, Rep. on Redpath Mus. No. 2, p. 12, Subcarboniferous.

MICROSPIONGIA, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 37. [Ety. *micros*, small; *spongia*, sponge.] Free, no epitheca; compact, without large openings; structure radiate. Type *M. gregaria*.

FIG. 108.—*Microspongia gregaria*.

gregaria, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 37, Hud. Riv. Gr.

parva, Ulrich, 1889, (Hindia *parva*), Am. Geol., vol. 3, p. 244, Trenton Gr.

MELLERINA, Ulrich, 1886, Cont. to Am. Pal., p. 34. [Ety. proper name.] Consisting of two suborbicular, thin-walled chambers, outer one with spiral ridges, inner one smooth; at the ends of the outer chamber there is a round opening, surrounded by an elevated border, where the ridges terminate. Type *M. greenii*.

greenii, Ulrich, 1886, Cont. Am. Pal., p. 35, Up. Held. Gr.

NODOSINELLA, Brady, 1876, Monograph Carb. and Perm. Foraminifera, p. 102. [Ety. *nodus*, knot; *ellus*, diminutive.] Free, straight, or arcuate, not spiral; constricted at intervals, test imperforate, texture finely arenaceous, aperture simple or compound. Type *N. digitata*.

priscilla, Dawson, 1868, (Dentalina *priscilla*), Acadian Geology, p. 285, Carboniferous.

Nullipora, Lamarck, 1801, Système des Animaux sans Vert. [Ety. *nullus*, no; *poros*, pore.] Not American Paleozoic.

? *obtexta*, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 33, Burlington Gr.

Orbiculites ? *reticulata*, see *Receptaculites reticulatus*.

PALÆACIS, Edwards & Haime, 1860, Hist. Nat. des Coralliaires, vol. 3, p. 171. [Ety. *palaos*, ancient; *akis*, barb.] Skeleton cuneate or turbinate, adherent, cups 1 to 12, cell-like, margins crenulate, separated by depressions; substance pierced by microscopic tubuli. Type *P. cuneiformis*.

compressus, Meek & Worthen, 1860, (Sphenopoterium *compressum*), Proc. Acad. Nat. Sci. Phil., p. 448, and Geo. Sur. Ill., vol. 2, p. 234, Keokuk Gr.

cuneatus, Meek & Worthen, 1860, (Sphenopoterium *cuneatum*), Proc. Acad. Nat. Sci., p. 448, syn. for *P. cuneiformis*.

cuneiformis, M. Edwards, 1860, Hist. Nat. d. Coralliaires, tome 3, p. 171, Warsaw Gr.

enormis, Meek & Worthen, 1860, (Sphenopoterium *enorme*), Proc. Acad. Nat. Sci., p. 448, and Geo. Sur. Ill., vol. 2, p. 146, Kinderhook Gr.

enormis, var. *depressus*, Meek & Worthen, 1866, (Sphenopoterium *enorme* var. *depressum*), Geo. Sur. Ill., vol. 2, p. 146, Kinderhook Gr.

obtusum, Meek & Worthen, 1860, (Sphenopoterium *obtusum*), Proc. Acad. Nat. Sci., p. 448, and Geo. Sur. Ill., vol. 2, p. 233, Keokuk Gr.

FIG. 111.—*Palæomanon cratera*.

large, dispersed openings, with inter-

FIG. 109.—*Nodosinella priscilla*. Natural size and enlarged.

FIG. 110. *Palæacis cuneiformis*.

PALÆOMANON, Roemer, 1860, Sil. Fauna West Tenn., p. 12. [Ety. *palaos*, ancient; *Manon*, a genus of sponges.] Cylindrical or irregular, cup-shaped, free, upper surface displaying

FIG. 112.—*Palaeospongia trentonensis*. View of calice.

vening space minutely porous. Type *P. cratera*.

cratera, Roemer, 1848, (*Siphonia cratera*.) Leonh. und Bronn's Jahrb., p. 685, Niagara Gr.

roemerl, Walcott, 1885, Monog. U. S. Geo. Sur., vol. 8, p. 99, Devonian.

PALÆOSPONGIA, D'Orbigny, 1850, Prodr. d. Palcont., t. 1, p. 26. [Ety. *palaia*, ancient; *spongia*, sponge.] Cyathiform, irregular, surface reticulated irregularly, by concentric and transverse lines. Type *P. cyathiformis*.

cyathiformis, Hall, 1847, (— *cyathiformis*.) Pal. N. Y., vol. 1, p. 72, Trenton Gr.

FIG. 113.—*Palaeospongia trentonensis*. Side view.

trentonensis, Worthen, 1875, (*Cnemidium trentonense*.) Geo. Sur. Ill., vol. 6, p. 491, Trenton Gr.

PASCEOLUS, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 342. [Ety. *pasceolus*, leather moneybag.] Sub-globular bodies marked on the cast as if by polygonal plates, and with a scar or depression for an attaching stem. Type *P. globosus*.

claudii, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 6, Hud. Riv. Gr.

darwini, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 5, Hud. Riv. Gr.

globosus, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 343, Trenton Gr.

gregarius, Billings, 1866, Catal. Sil. Foss. Antic., p. 72, Anticosti Gr.

halli, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 342, Anticosti Gr.

intermedius, Billings, 1866, Catal. Sil. Foss. Antic., p. 72, Anticosti Gr.

gregarius, Billings, 1866, Catal. Sil. Foss. Antic., p. 72, Anticosti Gr.

FIG. 115.—*Pasceolus darwini*. Upper surface. FIG. 116.—*Pasceolus darwini*. Under surface.

halli, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 342, Anticosti Gr.

intermedius, Billings, 1866, Catal. Sil. Foss. Antic., p. 72, Anticosti Gr.

FIG. 117.—*Pasceolus halli*. FIG. 114.—*Pasceolus claudii*.

PATTERSON

Cin. S.

[Ety. p.

no larg

pansio

bundle

and in

into th

spicule

aurita,

aurita,

Trento

difficilis,

Soc. N.

Gr.

tuberosa,

tubero

Trento

FIG. 118.—E

PHRAGMODI

Y. St.

phragm

Cylindr

a conc

expand

of a ret

rayed

roda.

catillifor

ton cat

Nat. H

lineata, I

Mus. N

patellifor

St. Mu

Gr.

PHYSOSPONG

N. Y. S

physa, b

cylindr

surface

tudinal

spicule

conce

bullate

terzona

P. daw

alternata,

Mus. N

colletti, I

Mus. N

dawsoni,

dawson

Hist., p

- PATTERNSONIA, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 43. [Ety. proper name.] Solid, amorphous, no large openings; lobed, pendent expansions on the upper surface, and bundles of fine filaments at the base and in the interior, which do not merge into the parenchyma of the sponge; spicules unknown. Type *P. difficilis*. aurita, Beecher, 1889, (Strobilospongia aurita,) Mem. Pea. Mus., vol. 2, p. 28, Trenton Gr.
- difficilis, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 43, Hud. Riv. Gr.
- tuberosa, Beecher, 1889, (Strobilospongia tuberosa,) Mem. Pea. Mus., vol. 2, p. 28, Trenton Gr.

FIG. 118.—Pattersonia difficilis. Fragment of upper surface.

- PHRAGMODICTYA, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 466. [Ety. phragmos, a partition; dictyon, a net.] Cylindrical or cup-shaped fronds, with a concave diaphragm near the broadly expanded base. Substance composed of a reticulate tissue of six and three rayed spicules and long cylindrical rods. Type *P. catilliformis*.
- catilliformis, Whitfield, 1881, (Dictyophyton catilliformis,) Bull. No. 1, Am. Mus. Nat. Hist., p. 18, Keokuk Gr.
- lineata, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 478, Keokuk Gr.
- patelliformis, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 478, Keokuk Gr.
- PHYSOSPONGIA, Hall, 1884, Abstr. 35th Rep. N. Y. St. Mus. Nat. Hist., p. 467. [Ety. physa, bladder; spongia, sponge.] Frond cylindrical, expanding from the base; surface divided into from 8 to 24 longitudinal areas by bands of tubular spicules, and into regular quadrules by concentric bands of spicules; surface bullate; spicules anchor-shaped. Interzonate tissue finely reticulated. Type *P. dawsoni*.
- alternata, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 481, Keokuk Gr.
- colletti, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 480, Keokuk Gr.
- dawsoni, Whitfield, 1881, (Uphantænia dawsoni,) Bull. No. 1, Am. Mus. Nat. Hist., p. 16, Keokuk Gr.

Protocyathus, Ford, 1878, Am. Jour. Sci. and Arts, 3d ser., vol. 15, p. 124, syn. for *Ethmophyllum*.

PROTOSPONGIA, Salter, 1864, Quar. Jour. Geo. Soc., vol. 20, p. 273. [Ety. *protos*, first; *spongia*, sponge.] Skeleton loose, reticulate formed of cruciform spiculae in one plane. Type *P. fenestrata*.

fenestrata, Salter, 1864, Quar. Jour. Geo. Soc., vol. 20, p. 238, and Mon. U. S. Geo. Sur., vol. 8, p. 11, Up. Taconic.

rarus, see *Ethmophyllum rarum*.

RAUFFELLA, Ulrich, 1889, Am. Geol., vol. 3, p. 235. [Ety. proper name.] Hollow cylindrical stems or radially arranged leaves; wall thin, composed of two layers of spicule tissue, inner one porous, outer one composed of large spicules appearing as threads interwoven. Type *R. filosa*.

filosa, Ulrich, 1889, Am. Geol., vol. 3, p. 237, Trenton Gr.

palmipes, Ulrich, 1889, Am. Geol., vol. 3, p. 238, Trenton Gr.

RECEPTACULITES, DeFrance, 1827, Dict. Sci. Nat., tome 45, p. 5. [Ety. *receptaculum*, receptacle; *lithos*, stone.] Subglobular, discoid, or infundibuliform; composed of cylindrical columns, connected at their upper and lower ends by transverse stolons. Type *R. neptunei*.

arcticus, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 576, Lower Silurian.

bursiformis, Hall, 1883, Rep. St. Geol., pl. 23, fig. 12-14, Schoharie Grit.

californicus, Billings, 1865, Pal. Foss., vol. 1, p. 351, Calif. Gr.

canadensis, Billings, 1863, (Ischadites canadensis,) Geo. of Can., p. 309, Anticosti Gr.

circularis, Emmons, 1856, Am. Geol., p. 230, Hud. Riv. Gr.

dactyloides, see *Cerionites dactyloides*.

devonicus, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 198, Up. Held. Gr.

eatoni, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 68-226, Schoharie Grit.

elegantulus, Billings, 1865, Pal. Foss., vol. 1, p. 360, Calif. Gr.

ellipticus, Walcott, 1885, Monog. U. S. Geo. Sur., vol. 8, p. 67, Chazy Gr.

elongatus, Walcott, 1885, Monog. U. S. Geo. Sur., vol. 8, p. 66, Chazy Gr.

formosus, Meek & Worthen, 1870, Proc. Acad. Nat. Sci., p. 23, and Geo. Sur. Ill., vol. 6, p. 500, Niagara Gr.

fungosus, Hall, 1861, Geo. Rep. Wis., p. 15, Galena Gr.

globularis, Hall, 1861, Supp. Geo. Sur. Wis., p. 16, and Geo. Sur. Ill., vol. 3, p. 301, Galena Gr.

hemisphericus, Hall, 1861, Geo. Rep. Wis., p. 16, and Geo. Wis., vol. 4, p. 269, Niagara Gr.

infundibuliformis, Eaton, 1832, (Coscinopora infundibuliformis,) Geo. Text Book, p. 44, Low. Held. Gr.

37.

38.

39.

- infundibulum, Hall, 1861, Geo. Rep. Wis., p. 16, Niagara Gr.
 40. insularis, Billings, 1866, Catal. Sil. Foss. Antic., p. 29, Anticosti Gr.
 iowensis, Owen, 1852, (Selenoides iowensis,) Geo. Sur. Wis., Iowa, and Minn., p. 587, Trenton Gr.
 41. jonesi, Billings, 1865, Pal. Foss., vol. 1, p. 389, Low. Held. Gr.

FIG. 119.—Receptaculites occidentalis, showing the tubes.

- mammillaris, Walcott, 1885, Monog. U. S. Geo. Sur., vol. 8, p. 65, Chazy Gr.
 monticulatus, Hall, 1883, Rep. St. Geol., pl. 23, fig. 3-11, Low. Held. Gr.

FIG. 120.—Receptaculites occidentalis, showing the endorhynch, the pores at the angles of the plates, and the deeply concave nucleus.

- reticulatus, Owen, 1844, (Orbituloides reticulata,) Rep. on Minn. Lands, p. 70, Niagara Gr.

FIG. 121.—Receptaculites occidentalis, showing the nucleus and the ectorrhynch.

- sacculus, Hall, 1879, Desc. New Species Foss. from Waldron, Ind., p. 1, and 11th Rep. Geo., and Nat. Hist. Ind., p. 222, Niagara Gr.
 squamifer, Hall, 1859, (Dictyocrinus squamifer,) Pal. N. Y., vol. 3, p. 135, Low. Held. Gr.
 subtrubrinatus, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 224, Niagara Gr.
 sulcatus, Owen, 1844. This name was pre-occupied by Goldfuss, and the species is now named *R. oweni*.

- tessellatus, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., vol. 1, p. 85, Niagara Gr.

RHABDARIA, Billings, Pal. Foss., vol. 1, p. 357. [Ety. *rhabdos*, rod.] Small, cylindrical bodies, with a rough surface and a perforation in the center. Type *R. fragilis*.

- fragilis, Billings, 1865, Pal. Foss., vol. 1, p. 357, Calciferous Gr.

- furcata, Billings, 1865, Pal. Foss., vol. 1, p. 358, Calciferous Gr.

RHOMBODICTYON, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 347. [Ety. *rhombos*, rhomb; *dictyon*, net.] Globular, discoid, or cyathiform, composed of two or more sets of rods crossing each other at various angles, but not dividing, and leaving rhombic spaces filled with another substance. Type *R. reniforme*.

- discum, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 348, Utica Slate.

- reniforme, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 347, Utica Slate.

- reniforme var. rhombiforme, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 348, Utica Slate.

Rotalia, Lamarck, 1804, Ann. Mus. [Ety. *rota*, wheel.] Not Palaeozoic.

baileyi, see *Endothyra baileyi*.

Saccamina, Sars, 1868, Vidensk.-Selsk. Forhandl., p. 248. [Ety. diminutive of *sakkos*, a bag.] Not American Palaeozoic.

eriana, Dawson, 1881, Can. Nat., vol. 10, syn. for *Calcisphera robusta*.

SACCOSPONGIA, Ulrich, 1879, Am. Geo., vol. 34, p. 242. [Ety. *sakkos*, bag; *spongia*, sponge.] Subcylindrical, with a central cloacal cavity extending through it; walls porous, traversed with tortuous branching canals intercommunicating with each other. Type *S. rudis*.

- danvillensis, Ulrich, 1889, Am. Geol., vol. 3, p. 243, Trenton Gr.

- rudis, Ulrich, 1889, Am. Geol., vol. 3, p. 242, Trenton Gr.

Scyphia, Oken, 1815. Not American Palaeozoic.

digitata, see *Brachiospongia digitata*.

stellata, Troost, 1840, not properly defined.

Selenoides, Owen, 1852, syn. for *Receptaculites*.

iowensis, see *Receptaculites iowensis*.

Siphonia, Parkinson, 1823, Organ. Rem. Not American Palaeozoic.

cratera, see *Palaecis cratera*.

imbricato-articulata, see *Astylospongia imbricato-articulata*.

praemorsa, see *Astylospongia praemorsa*.

Sphenopterium, Meek & Worthen, 1860, syn. for *Palaecis*.

compressum, see *Palaecis compressus*.

cuneatum, see *Palaecis cuneiformis*.

enormis, see *Palaecis enormis*.

enorme var. depressum, see *Palaecis enorme var. depressum*.

obtusum, see *Palaecis obtusum*.

Spongia, ozoic.

FIG. 122.—richmond several sponges.

twine-careous layers twinning latus.

atratus, Sci. ar.

Black brainerd.

Jour. vol. 32.

ocellatus, Sci. an.

Chazy richmond.

1882, monder.

Nat. H. Riv. G.

STREPTOSOLIA (in pres. p. 273).

solen, a pedunc.

having tendent.

surrounding canals.

are very obconic.

Geo. Sur.

STREPTOSPONGIA, p. 244.

sponge, twinning.

tortuous labyrinth.

labyrinthine, p. 24.

Strobilospongia Mus., v.

aurita, see *tuberosa*, s.

STROMATOCYBE, l, p. 48.

honey-composed of layers.

the pores.

richmondensis.

rugosum, 48, Bird.

Spongia, Linnaeus. Not American Palaeozoic.

FIG. 122.—*Strephochetus richmondensis*, showing several specimens on a slab.

inciso-lobata, see *Astylospongia incisolobata*.
stellatim-sulcata, see *Astylospongia stellatim-sulcata*.

STREPHOCHETUS, Seely, 1885, Am. Jour. Sci. and Arts, 3d ser., vol. 30, p. 355. [Ety. *strepho*, I

twine; *ochetos*, canal.] A free calcareous sponge, showing concentric layers composed of minute twining canals. Type *S. ocellatus*.

atratus, Seely, 1885, Am. Jour. Sci. and Arts, vol. 32, p. 32, Black Riv. Gr.

brainerdi, Seely, 1885, Am. Jour. Sci. and Arts, 3d ser., vol. 32, p. 32, Chazy Gr.

ocellatus, Seely, 1885, Am. Jour. Sci. and Arts, vol. 30, p. 357, Chazy Gr.

richmondensis, S. A. Miller, 1882, (*Stromatocentrum richmondense*), Jour. Cin. Soc. Nat. Hist., vol. 5, p. 41, Hud. Riv. Gr.

STREPTOSOLEN, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 273. [Ety. *streptos*, twisted; *solen*, a channel.] Obconical, pedunculate; central oscula having thin walled tubes extending to the base; oscula surrounded with radiating canals, between which there are vertical ones. Type *S. obconicus*.

obconicus, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 274, Trenton Gr.

STREPTOSPONGIA, Ulrich, 1889, Am. Geo., vol. 3, p. 244. [Ety. *streptos*, twisted; *spongia*, sponge.] Massive, composed of intertwining vertical lamellae, separated by tortuous linear interspaces. Type *S. labyrinthica*. Poorly defined.

labyrinthica, Ulrich, 1889, Am. Geo., vol. 3, p. 244, Hud. Riv. Gr.

Strobilospongia, Beecher, 1889, Mem. Pea. Mus., vol. 2, p. 14, syn. for *Pattersonia aurita*, see *Pattersonia aurita*.

tuberosa, see *Pattersonia tuberosa*.

STROMATOCERIUM, Hall, 1847, Pal. N. Y., vol. 1, p. 48. [Ety. *stroma*, layer; *kerion*, honey-comb.] Hemispherical, composed of numerous concentric vesicular layers, more or less wrinkled, without the pores that characterize *Stromatopora*. Type *S. rugosum*.

richmondense, see *Strephochetus richmondensis*.

rugosum, Hall, 1847, Pal. N. Y., vol. 1, p. 48, Birdseye and Black Riv. Gr.

STROMATOPORA, Goldfuss, 1826, Petref. Germ., p. 22. [Ety. *stroma*, stratum; *poros*, pore.] Dimorphous masses or extended sheets composed of delicate calcareous laminae, in successive layers, separated by minute, vertical pillars, dividing the interval into minute subquadrangular cavities; the whole is perforated by canals irregularly disposed and possessed of exhalant apertures. Type *S. concentrica*. *caespitosa*, Winchell, 1866, Rep. Low. Penin. Mich., p. 91, Ham. Gr. *compacta*, Billings, 1862, Pal. Foss., vol. 1, p. 55, Black Riv. Gr. There is some doubt about the reference of this species to this genus. Possibly it is a bryozoan. *concentrica*, Goldfuss, 1826, Germ. Petref.,

FIG. 123.—*Stromatocentrum rugosum*.

p. 22, and Pal. N. Y., vol. 2, p. 136, Niagara Gr.

constellata, see *Cœnostroma constellatum*. *erratica*, Hall, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 226, Up. Held. Gr.

expansa,

Hall &

Whitfield,

1873, 23d

Rep. N. Y.

St. Mus.

Nat. Hist.,

p. 226, Che-

mung Gr.

granulata,

Nicholson,

1873, Ann.

and Mag.

Nat. Hist.,

4th ser.,

vol. 12, p.

92, Cornif-

hindii, Nicholson, 1874, Ann. and Mag. Nat. Hist., 4th ser., vol. 13, and Pal. Prov. of Ont., p. 13, Niagara Gr.

FIG. 124.—*Stromatopora hindii*.

- incrustans*, see *Cænopora incrustans*.
mammillata, Nicholson, 1873, Ann. and Mag. Nat. Hist., 4th ser., vol. 12, p. 92, Corniferous Gr.
monticulifera, see *Cænostroma monticuliferum*.
nodulata, Nicholson, 1875, Ohio Pal., vol. 2, p. 249, Corniferous Gr.
nulliporoides, Nicholson, 1875, Pal. Prov. Ont., p. 78, Ham. Gr.
nux, Winchell, 1866, Rep. Low. Penin. Mich., p. 91, Ham. Gr.
ostiolata, Nicholson, 1873, Ann. and Mag. Nat. Hist., 4th ser., vol. 12, p. 90, Guelph Gr.
perforata, Nicholson, 1874, Ann. and Mag. Nat. Hist., 4th ser., vol. 13, and Pal. Prov. of Ont., p. 15, Corniferous Gr.
ponderosa, Nicholson, 1875, Ohio Pal., vol. 2, p. 246, Corniferous Gr.
pustulifera, see *Cænostroma pustuliferum*.
pustulosa, Safford. Not defined.
solidula, see *Cænostroma solidulum*.
subcylindrica, James, 1885, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 20, Hud. Riv. Gr. Poorly defined. Not a Stromatopora. None have been found in Lower Silurian rocks.
substriatella, Nicholson, 1875, Ohio Pal., vol. 2, p. 248, Corniferous Gr.
tuberculata, Nicholson, 1873, Ann. and Mag. Nat. Hist., 4th ser., vol. 12, p. 90, Corniferous Gr.
verrucosa, Troost, 1840, 5th Geo. Rep. Tenn., p. 66, Devonian? Not recognized.
STROTOSPONGIA, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 276. [Ety. *strotos*, twisted; *spongia*, sponge.] Funnel-shaped, composed of thin, intricately intertwined vertical leaves, arranged radiately around oscula; cloacal depressions, having apertures of vertical tubes in them; sponge-wall traversed by intertwined canals, having perforated thin walls; spicules minute, three-rayed. Type *S. maculosa*.
maculosa, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 277, Trenton Gr.
Syringophylum, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 250. The name was preoccupied by Edwards & Haime.

FIG. 125.—*Syringostroma columnare*.

SYRINGOSTROMA, Nicholson, 1875, Ohio Pal., vol. 2, p. 251. [Ety. *syrinx*, pipe; *stroma*, layer.] Massive, composed of con-

centric laminae, and vertical pillars firmly amalgamated. It is intimately related to Stromatopora. Type *S. columnare*.

columnare, Nicholson, 1875, Ohio Pal., vol. 2, p. 253, Corniferous Gr.

densum, Nicholson, 1875, Ohio Pal., vol. 2, p. 251, Corniferous Gr.

Textularia palæotrochus, see *Valvulina palæotrochus*.

THAMNODICTYA, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist. p. 466. [Ety. *thamnos*, shrub; *dictyon*, net.] Fronds tubular below, rapidly expanding and cyathiform or infundibuliform above, with twelve strong, longitudinal ridges dividing the surface into twelve areas. Substance reticulate. Type *T. newberryi*.

newberryi, Hall, 1863, (Dictyophyton newberryi), 16th Rep. N. Y. St. Mus. Nat. Hist., p. 87, Waterbury Gr.

TRACHYUM, Billings, 1865, Pal. Foss., vol. 1, p. 211. [Ety. *trachus*, rough, rugged.] Turbinate or cylindrical, with a cup on the upper surface. It has a close texture, without large canals. Type *T. cyathiforme*.

FIG. 126.—*Thamnodictya newberryi*. FIG. 127.—*Trachyum cyathiforme*.

cyathiforme, Billings, 1865, Pal. Foss., vol. 1, p. 211, Quebec Gr.

rugosum, Billings, 1865, Pal. Foss., vol. 1, p. 212, Quebec Gr.

TRICHOSPONGIA, Billings, 1865, Pal. Foss., vol. 1, p. 357. [Ety. *trichias*, to show hairs; *spongia*, sponge.] Large, rudely hemispheric, minutely fibrous, and full of elongate cylindrical or acerate spicules, just visible to the naked eye. There are also numerous irregular branching canals. Type *T. sericea*.

sericea, Billings, 1865, Pal. Foss., vol. 1, p. 257, Calceiferous Gr.

UPHANTENIA, Vauuxem, 1842, Geo. Rep. 3d Dist. N. Y., p. 183. [Ety. *uphantos*, woven; *tainia*, ribbon.] Composed of ligulate radiating and concentric bands, the reticulations being produced by the substance of the frond, and not by superficial striæ. For many years it was supposed to represent a marine plant. Type *U. chemungensis*.

chemungensis,
 3d Dist. N. Y.,
dawsoni,
 VALVULINA,
 d. l.
valva,
 blance,
 choid,
 cylind,
 somet,
 series.
 on the
 valvul,
 bulloide,
 Perm.
 erous,
 decurren,
 Scotlan,
 palæotro,
 ria pal,
 bonifer,
 plicata,
 land, p

SU

THE C

Classes: vi
 the palæozo
 animal) are
 They are a
 locomotive
 mouth, whi
 gastric cavi
 ding, dividi
 The sk
 secretions ta
 disk or ston
 tions or sep
 an expressio
 sentially a sk
 of the polyp
 made by the
 addition of s
 vesicular ma
 fill up the po
 ward, solid h
 Wherev
 the tissues b

- chemungensis, Vanuxem, 1842, Geo. Rep. 3d Dist. N. Y., p. 183, Chemung Gr. *dawsoni*, see *Physospongia dawsoni*.
- VALVULINA, D'Orbigny, 1826, Tabl. Method. d. l. Classe d. Cephulopodes. [Ety. *valva*, door; *inus*, implying resemblance.] Free or adherent, spirally trochoid, turbinoid, planoconvex or sub-cylindrical, chambers spirally arranged, sometimes terminating in a rectilinear series. Aperture in the umbilical angle, on the inferior surface, protected by a valvular tongue. Type *V. triangularis*. bulboides, Brady, 1876, Monog. Carb. and Perm. Foraminifera, p. 89, Carboniferous.
- decurrens, Brady, 1873, Mem. Geo. Sur. Scotland, pp. 63-95, Carboniferous.
- palæotrochus, Ehrenberg, 1854, (Textularia palæotrochus.) Mikogeologie, Carboniferous.
- plicata, Brady, 1873, Mem. Geo. Sur., Scotland, pp. 66-95, Carboniferous.
- randis, Brady, 1876, Monog. Carb. and Perm. Foraminifera, p. 90, Carboniferous.
- ZITTELELLA, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 267. [Ety. proper name.] Pedunculate, attached, variable in shape; upper surface with a shallow, central depression, with thin walled, vertical tubes extending to the base; radiating, inosculating canals, separated by spicular tissue, giving the appearance of vertical fissures. Type *Z. typicalis*. Ulrich & Everett refer *Palæospongia trentonensis* to this genus, so probably this genus is a synonym for *Palæospongia*.
- inosculata, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 271, Trenton Gr.
- lobata, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 270, Trenton Gr.
- typicalis, Ulrich & Everett, (in press,) Geo. Sur. Ill., vol. 8, p. 268, Trenton Gr. They have also made the varieties pistilliformis, subrotunda, and turbinata.

SUBKINGDOM CŒLEENTERATA.

THE Cœlenterata (*koilos*, hollow; *entera*, intestines) are divided into three Classes: viz., Anthozoa, Hydrozoa, and Ctenophora; the first two of which include the palæozoic fossils of this Subkingdom. The Anthozoa (*anthos*, flower; *zoon*, animal) are more generally known by the name Polypi (*polys*, many; *pous*, foot). They are all aquatic, usually cylindrical, organized for sedentary life, have no locomotive organs, and are provided with a circle of retractile tentaculæ around the mouth, which is destitute of any masticating apparatus, and they have a central gastric cavity. There are no special organs of sense, and they increase by budding, dividing, and by means of ova.

The skeleton which the polyps secrete is technically called the corallum. The secretions take place at the sides and lower part of the polyp, but not in the disk or stomach. Each septum is secreted between a pair of radiating, fleshy partitions or septa of the polyp, and hence the radiate structure of ordinary corals is an expression of the internal radiate structure of the polyp. The corallum is essentially a skeleton of carbonate of lime, the open spaces in which show the structure of the polyp animal. The bottom of the calyx, or calycle, in the corallum may be made by the meeting of the septa, or by the twisting of them together, with the addition of a point or columella at the center; or the bottom may be a porous or vesicular mass; or it may be solid, because the coral secretions of the polyp may fill up the pores, or because there are formed periodically, as the polyp grows upward, solid horizontal plates across the bottom, called tabulæ.

Wherever a tabula cuts off the connection of the polyp with the coral below, the tissues below the tabula dry and wither, and we have dead coral below the

tabula, and the living polyp above. In this way massive corals are formed ; the secretions take place at the top, and the animal cuts itself off from the coral skeleton below. Prof. Dana says :

“It is not more surprising, nor a matter of more difficult comprehension, that a polyp should form structures of stone (carbonate of lime) called coral, than that the quadruped should form its bones, or the mollusk its shell. The processes are similar, and so the result. In each case it is a simple animal secretion; a secretion of stony matter from the aliment which the animal receives, produced by the parts of the animal fitted for this secreting process; and in each, carbonate of lime is a constituent or one of the constituents of the secretion.”

Ordinary corals of the present seas have a hardness a little greater than common marble, or about equal to aragonite, and give a ringing sound when struck with a hammer.

The Anthozoa are divided into three Subclasses, two of which, Zoantharia (*zoon*, animal; *anthos*, flower) and Alcyonaria (*alkuoneion*, a zoophyte, like the kingfisher's nest), occur in palæozoic rocks. The Zoantharia are divided into seven orders, four of which are said to be palæozoic: viz., Perforata, Tabulata, Rugosa, and Tubulosa. To the Perforata the genera Protarea and Pleurodictyum have been very doubtfully referred, and the Auloporidae have been classed with the Tubulosa. All other palæozoic corals are referred to the Tabulata and Rugosa. The family Favositidae is typical of the Tabulata, and the family Cyathophyllidae of the Rugosa. Authors are not in accord respecting the ordinal relations of all the families, and hence we will simply arrange them alphabetically.

CLASS ANTHOZOA.

SUBCLASS ZOANTHARIA.

FAMILY AULOPORIDÆ.—Aulopora, Romingeria.

FAMILY CHETETIDÆ.—Chetetes, Dania, Ptychonema.

FAMILY COLUMNARIIDÆ.—*Calapœcia*, *Columnaria*, *Favistella*.

FAMILY CYATHOPHYLLIDÆ.—*Acervularia*, *Acrophyllum*, *Amplexus*, *Anisophyllum*, *Arachnophyllum*, *Astræophyllum*, *Aulacophyllum*, *Aulophyllum*, *Axophyllum*, *Baryphyllum*, *Blothrophyllum*, *Bucanophyllum*, *Campophyllum*, *Chonophyllum*, *Clisiophyllum*, *Coleophyllum*, *Craspedophyllum*, *Crepidophyllum*, *Cyathophyllum*, *Diphyphyllum*, *Duncanella*, *Elasmophyllum*, *Eridophyllum*, *Hadrophyllum*, *Hallia*, *Heliophyllum*, *Heteroprentis*, *Lithostrotion*, *Lophophyllum*, *Omphyma*, *Pachyphyllum*, *Palæophyllum*, *Phillipsastrea*, *Ptychophyllum*, *Pycnostylus*, *Streptelasma*, *Strombodes*, *Stylastrea*, *Trochophyllum*, *Zaphrentis*.

FAMILY CYCLOLITIDÆ.—Combophyllum, Discophyllum, Microcylus, Paleocylus.

FAMILY CYSTIPHYLLIDÆ.—*Cystiphorolites*, *Cystiphyllum*, *Cystostylus*.

FAMILY FAVOSITIDÆ.—Alveolites, Chonostegites, Cladopora, Cenites, Dendropora, Emmonsia, Favosites, Leptopora, Lunatipora, Michelinia, Pleurodictyum, Pachypora, Sphærolites, Striatopora, Syringolites, Trachypora, Vermipora.

FAMILY HALYSITIDÆ.—Halysites.

FAMILY PORITIDÆ.—Protarea.

FAMILY SYRINGOPORIDÆ.—Cannapora, Syringopora, Thecostegites.

FAMILY TETRADIDÆ.—Tetradium.

FAMILY THECIDÆ.—Thecia.

SUBCLASS ALCYONARIA.

FAMILY BOLBOPORITIDÆ.—Bolboporites.

FAMILY HELIOPORIDÆ.—Heliolites, Lyellia, Plasmopora.

FAMILY MONTICULIPORIDÆ.—Dekayella, Dekayia, Diplotrypa, Monotrypa, Monotrypella, Monticulipora, Nebulipora, Nyctopora, Prasopora.

FAMILY STELLIPORIDÆ.—Stellipora.

CLASS HYDROZOA.

This class is represented in paleozoic rocks by carbonaceous horny skeletons, called Graptolites. They are usually flattened, forming a thin film between shaly or slaty layers, and generally, in whatever rocks they occur, they are more or less compressed. Specimens are found in clay nodules and in calcareous clay beds, at Cincinnati and vicinity, which are cylindrical branching bodies, or have subquadrate stipes, covered with a thin, carbonaceous coating. The interior of one species is divided by longitudinal partitions of thin, carbonaceous films, into three departments, one of which is only about half the capacity of either of the other two. The denticulated edges on flattened films become projecting cells on more perfect specimens. The projecting cells may be subcircular or angular, and lead directly to the interior. When the interior substance is absent, and the cells are pressed together, instead of being pressed into the stipe, there is presented a diagrammatic side view of the cells, which furnishes the usual saw or denticulated aspect, but which gives a very imperfect, and frequently a very erroneous, idea of the form of the animal. This was the first Order of organisms to reach a high state of development, and the first to become extinct.

ORDER GRAPTOLIDA.

FAMILY CALLOGRAPTIDÆ.—Acanthograptus, Callograptus, Cyclograptus, Dendrograptus.

FAMILY DICTYONEMIDÆ.—Calyptograptus, Dictyonema, Rhizograptus.

FAMILY GRAPTOLITIDÆ.—Cladograptus, Climacograptus, Clonograptus, Dicranograptus, Didymograptus, Diplograptus, Graptolithus.

FAMILY GLOSSOGRAPTIDÆ.—Glossograptus, Retiograptus.

FAMILY NEMAGRAPTIDÆ.—Nemagraptus.

FAMILY MONOGRAPTIDÆ.—Monograptus.

FAMILY INOCAULIDÆ.—Inocaulus.

FAMILY MEGALOGRAPTIDÆ.—Megalograptus.

FAMILY OLDHAMIDÆ.—Oldhamia.

FAMILY PHYLLOGRAPTIDÆ.—Phyllograptus.

FAMILY PTILOGRAPTIDÆ.—Ptilograptus.

FAMILY RASTRITIDÆ.—Rastrites.

FAMILY RETIOLITIDÆ.—RETIOLITES.

FAMILY STAUROGRAPTIDÆ.—Staurograptus.

FAMILY THAMNOGRAPTIDÆ.—Bythograptus, Thamnograptus.

FAMILY UNCERTAIN.—Dawsonia.

ACANTHOGRAPTUS, Spencer, 1878, Can. Nat., vol. 8, p. 462.

FIG. 128.—*Acanthograptus pulcher*.

[Ety. *akantha*, spine; *grapho*, I write.] Shrub-like; one side spinous. Stronger and more bushy than *Dendrograptus*. Type A. *granti*.

granti, Spencer, 1878,

Can. Nat., vol. 8, p. 463, and Bull. No. 1, Mus. Univ., St. Mo., p. 31, Niagara Gr. *pulcher*, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 32, Niagara Gr.

ACERVULARIA, Schweigger, 1820, Handb. der Naturg., p. 418. [Ety. *acervus*, a heap; considered as a body.] Compound, massive, cells presenting two separated walls, as in *Aulophyllum*; septa well developed between the walls, but much less in the central area; no columella; tabulæ little developed; increasing by gemmation. Type A. *baltica*.

FIG. 129.—*Acervularia clintonensis*.

adjunctiva, White, 1880, Proc. U. S. Nat. Mus., vol. 2, p. 255, and Cont. to Pal. No. 6, p. 120, Carboniferous.

clintonensis, Nicholson, 1875, Ohio Pal., vol. 2, p. 227, Niagara Gr.

FIG. 130.—*Acervularia davidsoni*.

davidsoni, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 418, Up. Held. and Ham. Gr.

inequalis, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 233, Chemung Gr.

pentagona, Goldfuss, 1826, (Cyathophyllum *pentagonum*,) Petref. Germ., p. 60, Devonian.

profunda, Hall, 1858, Geo. Sur. Iowa, p. 477, Ham. Gr.

ACROPHYLLUM, Thomson & Nicholson, 1876, Ann. and Mag. Nat. Hist. 4th ser., vol. 17, p. 455. [Ety. *akros*, summit; *phyllon*, leaf.] Corallum simple, turbinate, or sub-cylindrical, straight, or curved; septa numerous, well-developed, coalescing, and curving as they reach the tabulæ, forming prominent, tortuous ridges on the central, elevated portion, and becoming complicated with the tabulæ to form the conspicuous, central prominence, which often forms a central axis; fossette reaches from the base of the elevation to the margin of the calyx; exterior usually constricted. Type A. *oneidaense*.

oneidaense, Billings, 1859, (Clisiophyllum *oneidaense*, Can. Jour., p. 128, Up. Held. Gr. *agarcia*, Lamarck, 1801, Syst. des Anim. sans Vert. Not Palæozoic. *swinderniana*, see *Thecia swinderniana*.

ALVEOLITES, Lamarck, 1801, Syst. des Anim. sans Vert., p. 375. [Ety. *alveus*, cavity; *lithos*, stone.] Dendroid, massive, or incrusting; corallites short, prismatic, or cylindrical; walls united; tabulæ complete; mural pores large, usually near the angles of the tubes, few in number; calices oblique, lower lip most prominent; septa absent, or forming tooth-like projections. Type A. *eschroidea*.

arctica, Woodward, 1879, Lond. Geo. Mag. n. s., vol. 5, Devonian.

billingsi, Nicholson, 1874, Geo. Mag. n. s., vol. 1, p. 55, Up. Held. Gr.

confertus, Nicholson, 1874, Geo. Mag. n. s., vol. 1, p. 54, Up. Held. Gr.

FIG. 131.—*Acrophyllum oneidaense*.

cryptod.
4, p. 1
distant,
vol. 1,
dubia, s
explana
13, fig
Low.

FIG. 132.

Not de
hemisph
Pulmon
to be r
irregular
Geo. S

vol. 4,
labechi,
Foss. d

50. labiosus,
4, p. 11

megastom
Penin.

multilam
40th Pa

niagarens
Can. Jo

niagarensi
ramulosu

s., vol.

repens, F

Amoen
reticulat

rockford

Rep. N

Chemu

41. roemerii,
p. 255,

selwyni,
vol. 1,

42. squamosu
5, p. 25

strigillatu
Peninsu

subramos
als, p. 4

undusus,
2d ed.,

the spec
1876, in
preoccu

cryptodens, Billings, 1850, Can. Jour., vol. 4, p. 115, Up. Held. Gr.
 distans, Nicholson, 1874, Geo. Mag. n. s., vol. 1, p. 54, Up. Held. Gr.
 dubia, see Favosites dubius.
 explanatus, Hall, 1883, Rep. St. Geol., pl. 13, fig. 16, and Pal. N. Y., vol. 6, p. 11, Low. Held. Gr.

FIG. 132.—*Alveolites goldfussi*.

Not defined so as to be recognized.

- hemisphericus*, D'Orbigny, 1850, Prodr. d. Paléont., t. 1, p. 49. Not defined so as to be recognized.
 irregularis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 72, and Geo. Wis., vol. 4, p. 251, Hud. Riv. Gr.
 labechi, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 257, Anticosti Gr.
 50. labiosus, Billings, 1850, Can. Jour., vol. 4, p. 114, Up. Held. Gr.
 megastoma, Winchell, 1866, Rep. Low. Penin. Mich., p. 89, Ham. Gr.
 multilamella, Meek, 1877, U. S. Geo. Sur. 40th Parallel, vol. 4, p. 25, Devonian.
 niagarensis, Nicholson & Hinde, 1874, Can. Jour., vol. 14, p. 150, Niagara Gr.
 niagarensis, Rominger, see *A. undosus*.
 ramulosus, Nicholson, 1874, Geo. Mag. n. s., vol. 1, p. 55, Up. Held. Gr.
 repens, Fought, 1749, (Millepora repens,) Amaen. Acad., vol. 1, p. 99, Niagara Gr.
 reticulata, see Favosites reticulatus.
 rockfordensis, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 229, Chemung Gr.
 51. roemeri, Billings, 1860, Can. Jour., vol. 5, p. 255, Ham. Gr.
 selwyni, Nicholson, 1874, Geo. Mag. n. s., vol. 1, p. 15, Up. Held. Gr.
 52. squamosus, Billings, 1860, Can. Jour., vol. 5, p. 257, Up. Held. Gr.
 strigillatus, Winchell, 1866, Rep. Low. Peninsula Mich., p. 89, Ham. Gr.
 subramosus, Rominger, 1876, Foss. Corals, p. 43, Ham. Gr.
 undosus, S. A. Miller, 1883, Am. Pal. Foss., 2d ed., p. 262, Niagara Gr. Proposed for the species described by Rominger in 1876, in Foss. Corals, p. 40, under the preoccupied name of *A. niagarensis*.

exsul, see Calopora exsul.
fischeri, see Pachypora fischeri.
frondosus, see Pachypora frondosa.
goldfussi, Billings, 1860, Can. Jour., vol. 5, p. 255, Ham. Gr.
granulosus, James, 1875, Catal. Cin. Foss., p. 2.

vallorum, Meek, 1868, Trans. Chi. Acad. Sci., p. 86, Devonian.
 AMPLEXUS, Sowerby, 1814, Mineral Conchology, vol. 1, p. 165. [Ety. *amplexus*, encircling.] Resembles *Zaphrentis*, except the septa do not extend to the center, they leave the upper surface of the tabulae exposed in that part; septal fossula highly developed in the upper portion of the corallum; tabulae well developed; surface usually constricted. Type *A. coralloides*.
 annulatus, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 80, and Geo. Wis., vol. 4, p. 314, Niagara Gr.
 53. cingulatus, Billings, 1862, Pal. Foss., vol. 1, p. 106, Mid. Sil.
 coralloides, Sowerby, 1814, Min. Conch., vol. 1, p. 165, Warsaw Gr.
 54. exilis, Billings, 1875, Can. Nat. and Geol., vol. 7, p. 232, Up. Held. Gr.
 fieldeni, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 589, Niagara Gr.
 fenestratus, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 80, and Geo. Wis., vol. 4, p. 278, Niagara Gr.
 fragilis, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 116, Keokuk Gr.
 hamiltoniae, Hall, 1876, Illust. Dev. Foss., pl. 19, Ham. Gr.
 intermittens, Hall, 1876, Illust. Dev. Foss., pl. 32, Ham. Gr.
 junctus, Hall, 1882, Foss. Corals Niagara and Up. Held. Groups, p. 11, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 415, Niagara Gr.

FIG. 133.—*Amplexus yandelli*

Sutherland's Jour., vol. 2, p. ccxxx, Niagara Gr.
 shumardi, Edwards & Haime, 1851, (Cya-

55. laxatus, Billings, (?) Can. Nat. and Geol., vol. (?) p. (?) Up. Held. Gr.
 56. mirabilis, Billings, 1875, Can. Nat. and Geol., vol. 7, p. 232, Coal Meas.
 phragmoceras, Salter, 1852, (Calophyllum phragmoceras)

- thophyllum shumardi,) Pol. Foss. Terr. Pal., p. 370, Niagara Gr.
 uniformis, Hall, 1882, Foss. Corals Niagara & Up. Held. Grs., p. 11, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 415, Niagara Gr.
 yandelli, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 344. Up. Held. Gr. zaphrentiformis, White, 1876, Geo. of Uinta Mountains, p. 107, and Cont. to Pul. No. 6, p. 120, Low. Aubrey Gr.
 ANISOPHYLLUM, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 351. [Ety. *anisos*, unequal; *phyllon*, leaf.] Distinguished from Zaphrentis by the great development of three primary septa, one of which faces the septal fossula; this fossula extends to the center of the visceral chamber, and there ceases to be distinct from the bottom of the calyces. Type A. agassizi.
 agassizi, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 351, Low. Held. Gr. bilamellatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 9, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 413, Niagara Gr.
 trifurcatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 9, and 12th Rep. Ind. Geol. & Nat. Hist., p. 273, Niagara Gr.

FIG. 134.—*Anisophyllum unilargum*.

- unilargum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 8, and 12th Rep. Ind. Geol. & Nat. Hist., p. 272, Niagara Gr.
Anthophyllum, Schweigger, 1820, Handb. der Naturg., p. 417, Not a Palaeozoic genus.
denticulatum, Goldfuss, 1826, Petref. Germ., p. 46, Niagara Gr. Not determined.
expansum, Owen, 1840, Rep. on Mineral Lands, p. 69. Not defined so as to be recognized.
 ARACHNOPHYLLUM, Dana, 1848, Zoophytes U. S. Expl. Exped., vol. 8, p. 360. [Ety. *arachne*, spider; *phyllon*, leaf.] Massive, encrusting, having obtusely defined polygonal scars, with a depressed, flattened center, in which the septa meet; septa thin, perforated; buds marginal, structure vesicular, arranged in transverse undulations, corresponding to the form of the cells; no defining walls to the center or between the stars; center marked by a few vertical striae, resulting from the twisted edges of the septa. Type A. baltica. (Acervularia baltica of authors.)
 richardsoni, Salter, 1852, Sutherland's Jour., vol. 2, p. cccxxii, Up. Sil.
 Astraea, Lamarck, 1816, Hist. Nat. d. Animaux Vert., vol. 2, p. 257. Not a Palaeozoic genus.
gigas, see *Phillipsastrea gigas*.
hennahi, see *Smithia hennahi*.
helianthoides, see *Heliophyllum halli*.
mammillaris, see *Strombodes mammillaris*.

- mammillaris*, see *Lithostrotion mammillare*.
rugosa, see *Cyathophyllum rugosum*.
tessellata, Troost. Not defined.
 ASTREOPHYLLUM, Nicholson & Hinde, 1874, Can. Jour., vol. 14, p. 152. [Ety. *aster*, star; *phyllon*, leaf.] Corallum aggregate; corallites cylindrical and united by numerous mural expansions, which form complete floors; septa meeting in the center, forming a columella; costal radii prolonged over the successive exothecal floors; tabulae rudimentary or absent (?). Type A. gracile.

FIG. 135.—*Astreophyllum gracile*, greatly enlarged, showing calyces, confluent mural expansions, and costal radii.

- gracile*, Nicholson & Hinde, 1874, Can. Jour., vol. 14, p. 153 and Pal. Ontario, p. 57, Niagara Gr.
Astrocerium, Hall, 1852, Pal. N. Y., vol. 2, p. 120. [Ety. *aster*, star; *kerion*, honeycomb.] It was supposed to be distinguished from Favosites by the presence of twelve or more slender spiniform rays, but it is a synonym. Type A. venustum.
constrictum, see Favosites constrictus.
parasiticum, see Favosites parasiticus.
pyriforme, see Favosites pyriformis.
venustum, see Favosites venustus.
 AULACOPHYLLUM, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxxvii. [Ety. *aulos*, furrow; *phyllon*, leaf.] Resembles Halia, though the septal fossula is not replaced by a primary septum, but forms a narrow groove at the bottom where the adjoining septa meet. Type A. sulcatum.
 bilaterale, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 25, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 429, Up. Held. Gr.
 convergens, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 22, and 12th Rep. Ind. Geo., p. 281, Up. Held. Gr.
 cruciforme, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 24, and 12th Rep. Ind. Geo., p. 283, Up. Held. Gr.
 pinnatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 23, and 12th Rep. Ind. Geo., p. 284, Up. Held. Gr.
 poculum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 25 and 12th Rep. Ind. Geo., p. 283, Up. Held. Gr.

praeceptum
 agara
 12th R
 praterifo
 agara
 12th R

FIG. 13

princeps,
 and Up.
 Ind. Ge.
 reflexum,
 and Up.
 Ind. Ge.
 sulcatum,
 cuta,) P.
 Rep. Ind.
 tripinnat
 agara a
 12th Rep
 trisculcat
 agara a
 12th Re
 Gr.
 AULOPHYLLUM
 Brit. Foss.
 pipe; *ph*
 septa w
 aents d
 visceral
 central s
 nal and
 not well
 richardson
 Sci., p. 8
 AULOPORA, G
 82. [Et
 Creeping
 gemmati
 pet-shap
 nicating
 springs;
 dimentar

praeceptum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 24, and 12th Rep. Ind. Geo., p. 280, Up. Held. Gr.
 prateriforme, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 23, and 12th Rep. Ind. Geo., p. 282, Up. Held. Gr.

FIG. 136.—*Aulacophyllum princeps*.

princeps, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 23, and 12th Rep. Ind. Geo., p. 281, Up. Held. Gr.
 reflexum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 24, and 12th Rep. Ind. Geo., p. 284, Up. Held. Gr.
 sulcatum, D'Orbigny, 1850, (Caninia sulcata,) Prodr. d. Pal. t. 1, p. 105, and 12th Rep. Ind. Geo., p. 279, Up. Held. Gr.
 tripinnatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 25, and 12th Rep. Ind. Geo., p. 285, Up. Held. Gr.
 trisulcatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 25, and 12th Rep. Ind. Geo., p. 285, Up. Held. Gr.

AULOPHYLLUM, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxx. [Ety. *aulos*, pipe; *phyllon*, leaf.] Corallum simple; septa well-developed; mural investments double, the interior dividing the visceral chamber into two parts—one central and columnar, the other external and annular; no columella; tabulae not well developed. Type A. proliferum.
 richardsoni, Meek, 1868, Trans. Chi. Acad. Sci., p. 81, Devonian.

AULOPORA, Goldfuss, 1826, Petref. Germ., p. 82. [Ety. *aulos*, pipe; *poros*, pore.] Creeping, increasing by latero-basal gemmation; corallites pyriform, trumpet-shaped, the cavity of each communicating with the one from which it springs; no pores; septa absent or rudimentary. Type A. serpens.

annectans, Clarke, 1885, Bull. 16. U. S. Geo. Sur., p. 63, Genesee shales.
 aperta, Winchell, 1866, Rep. Low. Penin. Mich., p. 91, Ham. Gr.
 arachnoidea, Hall, 1847, Pal. N. Y., vol. 1, p. 76, Trenton and Hud. Riv. Gr.
 canadensis, see *Hederella canadensis*.
 conferta, Winchell, 1866, Rep. Low. Penin. Mich., p. 91, and Rominger's Foss. Corals, p. 88, Ham. Gr.
 cornulites, Hall, 1883, Rep. St. Geo., pl. 2, figs. 21 and 22, Low. Held. Gr.
 cornuta, see *Romingeria cornuta*.
 cyclopora, Winchell, 1866, Rep. Low. Penin. Mich., p. 92, Ham. Gr.
 elongata, Hall, 1887, Pal. N. Y., vol. 6, p. 5, Low. Held. Gr.
 erecta, Rominger, 1876, Foss. Corals, p. 88, Ham. Gr.

filiformis, see *Hederella filiformis*.
 lowensis, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 235, Chemung Gr.

precus, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 107, Niagara Gr.
 repens, Walch, et Knorr, 1775, (Milleporites repens,) Sammlung von Merkw., vol. 3, p. 179, and Sil. Fauna W. Tenn., p. 28, Niagara Gr.

saxivada, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 235, Chemung Gr.

schoharie, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 110, Low. Held. Gr.

serpens, Goldfuss, 1826, Germ. Petref., p. 82, and Rominger's Foss. Corals, p. 87, Ham. Gr.

serpuloidea, Winchell, 1866, Rep. Low. Penin. Mich., p. 91, Ham. Gr.

subtenuis, Hall, 1883, Rep. St. Geo., pl. 2, fig. 9-20, Low. Held. Gr.

tubiformis, Goldfuss, 1826, Germ. Petref., p. 82, and Murch. Sil. Syst., Up. Held. and Ham. Gr.

tubula, Hall, 1883, Rep. St. Geo., pl. 2, fig. 7-8, Low. Held. Gr.

umbellifera, see *Romingeria umbellifera*.
 vancevii, Hall, 1883, 12th Rep. Ind. Geo., p. 255, Niagara Gr.

Axinura, Castlenau, syn. for *Lithostrotion*.

canadense, see *Lithostrotion canadense*.

AXOPHYLLUM, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxxii. [Ety. *axon*, axis; *phyllon*, leaf.] Corallum simple, trochoid, and in structure resembling *Lithostrotion*. Type A. expansum.

FIG. 138.—*Axophyllum rudé*.

FIG. 137.—*Aulopora serpens*.

infundibulum, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 525, Coal Meas.
rude, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 115, Coal Meas.

BARYPHYLLUM, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxvi. [Ety. *barys*, heavy; *phyllon*, leaf.] Corallum short; calice superficial; slight septal fossula corresponding to one of the branches of a cross, the other three of which are primary septa; younger septa inclined toward the primary ones. Type *B. verneuianum*.

arenarium, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 409, Onondaga Gr.

fungulus, White, 1878, Proc. Acad. Nat. Sci. Phil., p. 29, Niagara Gr.

verneuianum, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 352, Niagara Gr.

BLOTHROPHYLLUM, Billings, 1859, Can. Jour., vol. 4, p. 130. [Ety. *blothros*, tall-growing; *phyllon*, leaf.] Corallum simple, turbinate, or cylindrical, having the central region occupied by flat, transverse diaphragms; an intermediate area, with strong radiating septa, and an outer area, in which there are imperfect diaphragms, projecting upward, and having on their upper surface rudimentary septa; a thin, complete epitheca, and a septal fossette. Type *B. decorticatum*.

approximatum, Nicholson, 1873, Can. Nat. and Geo., vol. 7, p. 140, Up. Held. Gr.

caespitosum, Rominger, 1876, Foss. Corals, p. 114, Niagara Gr.

decorticatum, Billings, 1859, Can. Jour., vol. 4, p. 130, Up. Held. Gr.

multicalicatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 44, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 448, Up. Held. Gr.

papulosum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 44, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 448, Up. Held. Gr.

promissum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 45, and 12th Rep. Ind. Geo., p. 304, Up. Held. Gr.

sinuosum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 45, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 448, Up. Held. Gr.

FIG. 139. — Blotrophylum promissum.

BOLBOPORITES, Pander, 1830, Beitr. zur Geognosie Russlands, p. 106. [Ety. *bolbos*, bulb; *poros*, pore.] Small, globular, showing basal attachment; structure dense. The type of the genus is said to be neither a coral nor bryozoan, but to belong to the Echinodermata. The form which Billings referred to the genus is probably a coral.

americanus, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 429, Chazy Gr.

BUCANOPHYLLUM, Ulrich, 1886, Cont. to Am. Pal., p. 31. [Ety. *bukane*, trumpet; *phyllon*, leaf.] Corallum trumpet-shaped, consisting of a long, slender, cylindrical stem, with the upper end abruptly dilated into a cup, which becomes oblique in older specimens; interior of cup with numerous septal striae, which become obsolete at the bottom. Type *B. gracile*.

gracile, Ulrich, 1886, Cont. to Am. Pal., p. 31, Up. Held. Gr.

BYTHOGRAPTUS, Hall, 1861, Geo. Rep. Wis., p. 18. [Ety. *buthos*, in the deep; *graphos*, I write.] Frond consisting of a central stipe, with closely arranged lateral branches, flexuous or recurved; cellularous on one side; substance corneous brown or black. Type *B. laxus*.

laxus, Hall, 1861, Geo. Rep. Wis., p. 19, Trenton Gr.

Calamopora, Goldfuss, syn. for *Favosites basaltica*, see *Favosites basalticus*.

cellulata, Castelnau, 1843. Not recognized.

cristata, see *Favosites cristatus*.

cumberlandica, see *Favosites cumberlandicus*.

favosa, see *Favosites favosus*.

fibrosa, see *Monticulipora fibrosa*.

fibrosa, Roemer, see *Hindia fibrosa*.

forbesi var. *discoidea*, see *Favosites forbesi* var. *discoideus*.

goldfussi, see *Favosites goldfussi*.

gothlandica, see *Favosites gothlandicus*.

heliolitiformis, see *Favosites heliolitiformis*.

hemispherica, see *Favosites hemisphericus*.

infundibuliformis, Goldfuss, identified by D'Archiac and Verneuil. Not an American species.

mackrothi, see *Chetetes mackrothi*.

maxima, see *Favosites maximus*.

minuta, Castelnau. Not recognized.

minutissima, Castelnau. Not recognized.

radians, Castelnau. Not recognized.

tumida, see *Chetetes tumidus*.

verneuili, Castelnau, syn. for *Monticulipora fibrosa*.

winchelli, see *Favosites winchelli*.

CALAPGECIA, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 425. [Ety. *kalos*,

FIG. 140. — Bolboporites americanus. a, view of base; b, c, d, side views.

beautiful hemispherical sites, covered with a thin; the sp. Type anticosti Foss. camden Geo., 2

FIG. 141. — Calceola eribrita.

CALCEOLA, sans slipper subtriseptal americanis. attenuat. Sci. Ph. referre phyllu cornicul. Sci. Ph. C. tenu coxi, Lyc. Proc. Nat. Sc. p. 44, Gr. for C. seensis plicata, 1840, Rep. N. Y. St. Mus. Nat. Hist., p. 207, Held. pusilla, and U. N. Y. agara

FIG. 143. — Calceola eribrita.

Lindst genus CALLOGRAI Decade grapho

beautiful; *poikilos*, spotted.] Composite, hemispherical or subspherical, corallites slender, tubular, perforated as in *Favosites*, outside striated by imperfectly developed costae; septa about 24; tabulae thin; when corallites are not in contact the space is filled with vesicular tissue. Type *C. canadensis*.

59. *anticostiensis*, Billings, 1866, Catal. Sil. Foss. Antic., p. 32, Hud. Riv. Gr.

60. *canadensis*, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 426, Black Riv. Gr.

cribriformis, Nicholson, 1874, (Columnopora *cribriformis*,) Geo. Mag., vol. 1, p. 253, and Pal. Ohio, vol. 2, p. 186, Hud. Riv. Gr.

huronensis, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 426, Hud. Riv. Gr.

FIG. 141.—*Calappaecia cribriformis*.

CALCEOLA, Lamarck, 1801, Syst. des Animaux Vert., p. 139. [Ety. *calceola*, a slipper.] Corallum simple, operculated, subtriangular, pyramidal; calice deep; septa narrow; structure dense. Type *C. sandalina*.

americana, Safford, syn. for *C. tennesseensis*.

attenuata, Lyon, 1879, Proc. Acad. Nat. Sci. Phil., p. 45, Niagara Gr. Lindstrom referred this species to his genus *Rhizophyllum*.

corniculatum, Lyon, 1879, Proc. Acad. Nat. Sci. Phil., p. 43, Niagara Gr. Syn. (?) for *C. tennesseensis*.

coxi, Lyon, 1879, Proc. Acad. Nat. Sci. Phil., p. 44, Niagara Gr. Syn. (?) for *C. tennesseensis*.

plicata, Conrad, 1840, Ann. Rep. N. Y., p. 207, Low. Held. Gr.

pusilla, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 15, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 419, Niagara Gr.

FIG. 142.—*Calceola sandalina*, showing deep calice.

FIG. 143.—*Calceola sandalina*, operculum.

Lindstrom referred this species to his genus *Rhizophyllum*.

CALYPTOGRAPTUS, Hall, 1865, Can. Org. Rem. Decade 2, p. 133. [Ety. *kallos*, beautiful; *grapho*, I write.] Flabellate fronds, with

numerous slender, bifurcating branches proceeding from a strong stem; branches and divisions celliferous on one side, striate on the other; sometimes distantly and irregularly united by transverse dissepiments. Type *C. elegans*.

elegans, Hall, 1865, Can. Org. Rem. Decade 2, p. 134, Quebec Gr. or Up. Taconic.

granti, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 21, Niagara Gr.

minutus, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 22, Niagara Gr.

multiculus, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 22, Niagara Gr.

FIG. 144.—*Callograptus niagarensis*.

niagarensis, Spencer, 1878, Can. Nat., vol. 8, and Bull. No. 1, Mus. Univ. St. Mo., p. 21, Niagara Gr.

salteri, Hall, 1865, Can. Org. Rem. Decade 2, p. 135, Quebec Gr., or Up. Taconic.

Calophyllum, Dana, 1846, Am. Jour. Sci., p. 183, syn. for *Amplexus*.

phragmoceras, see *Amplexus phragmoceras*.

CALYPTOGRAPTUS, Spencer, 1878, Can. Nat., vol. 8, p. 459. [Ety. *kalyptos*, covered; *grapho*, I write.] Cyathiform, bifurcating branches, not connecting laterally; resembles *Dicyonema*. Type *C. cyathiformis*.

FIG. 145.—*Calyptograptus cyathiformis*.

cyathiformis, Spencer, 1878, Can. Nat., vol. 8, p. 459, Niagara Gr.

subretiformis, Spencer, 1878, Can. Nat., vol. 8, p. 460, Niagara Gr.

CAMPOPHYLLUM, Edwards & Haime, 1850, British Foss. Corals, p. lxxviii. [Ety. *kampto*, I bend; *phyllon*, leaf.] Simple, tall, protected by an epitheca; septa well developed; tabulæ very large and smooth toward the center; interseptal area vesicular. Type *C. flexuosum*.

minum, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 232, Chemung Gr.

texanum, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 388, Permian.

torquillum, Owen, 1852, (Cynathophyllum torquillum,)

Geo. Rep. Wis., Iowa, and Minn., pl. 4, fig. 2, Coal Meas.

Caninia, Michelin, syn. for *Zaphrentis*.

bilateralis, see *Zaphrentis* *bilateralis*.

punctata, D'Orbigny, 1850, Prodr. d. Paléont., t. 1, p. 105. Not defined so as to be recognized.

FIG. 147.—*Campophyllum torquillum*.

sulcata, D'Orbigny, 1850, Prodr. d. Paléont., t. 1, p. 105. Not defined so as to be recognized.

FIG. 148.—*Campophyllum torquillum*. Transverse section.

CANNAPORA, Hall, 1852, Pal. N. Y., vol. 2, p. 43. [Ety. *kanna*, reed; *poros*, pore.] Massive, tubular, united externally by

FIG. 146.—*Campophyllum flexuosum*. Vertical section showing smooth central tabulæ.

tabulæ; distinguished from *Syringopora* by the regular transverse external tabulæ and by the internal structure of the corallites. Type *C. junciformis*.

annulata, Nicholson & Hinde, 1874, Can. Jour., p. 154, and Pal. Prov. of Ontario, p. 58, Niagara Gr.

junciformis, Hall, 1852, Pal. N. Y., vol. 2, p. 43, Clinton Gr.

Caryophyllia, Lamarck, 1816. Not Palæozoic.

cornicula, see *Zaphrentis* *cornicula*.

gigantea, see *Zaphrentis* *gigantea*.

pulmonea, see *Zaphrentis* *pulmonea*.

Catenipora, Lamarck, 1816, syn. for *Halysites* *micelini*, Castelnau, syn. for *Halysites* *cateulatus*.

CHETETES, Fischer, 1837, Oryct. du Gouv. Moscou, p. 159. [Ety. *chaite*, hair.]

Corallum conglomerate; corallites very long, basaltiform; calyces polygonal; tabulæ not connected or on the same plane in different corallites; walls analagated, imperforate; growth fissiparous. Type *C. radians*.

abruptus, see *Monotrypella* *abrupta*.

aequidistans, Hall, 1881, Bryozoans Up. Held Gr., p. 4, Up. Held. Gr.

approximatus, Nicholson, syn. for *Monticulipora* *dalii*.

arbusculus, see *Monotrypella* *arbuscula*.

arcticus, Houghton, 1857, Jour. Roy. Dub. Soc., vol. 1, Silurian.

atritus, Nicholson, syn. for *Dekayia* *aspera*.

barrandii, see *Monticulipora* *barrandii*.

briareus, see *Monotrypella* *briareus*.

calicula, see *Aspidopora* *caliculus*.

carbonarius, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 526, Coal Meas.

cincinnatiensis, see *Monticulipora* *cincinnatiensis*.

clathratulus, James & Nicholson, syn. for *Monticulipora* *pavonia*.

clavacoides, see *Leptopora* *clavacoides*.

colliculatus, Hall, 1883, Rep. St. Geo., pl. 8, fig. 1-4, and Pal. N. Y., vol. 6, p. 11, Low. Held. Gr.

columnaris, see *Tetradium* *columnare*.

compressus, see *Peronopora* *compressa*.

consimilis, see *Monotrypella* *consimilis*.

corticans, Nicholson, syn. for *Spatiopora* *tuberculata*.

corticosa, see *Trematopora* *corticosa*.

crassus, Lonsdale, 1845, (Stenopora *crassa*,) Russ. and Ural Mts., vol. 1, p. 631, Coal Meas.

crebrirama, Hall, 1881, Bryozoans Up. Held. Gr., p. 4, Up. Held. Gr.

dalet, see *Monticulipora* *dalii*.

decipiens, see *Monticulipora* *decipiens*.

delicatulus, see *Monticulipora* *delicatula*.

discoideus, see *Amplexopora* *discoidea*.

egenus, Hall, 1881, Bryozoans Up. Held. Gr., p. 4, Up. Held. Gr.

elegans, see *Disotrypa* *elegans*.

exilis, Dawson, 1868, Acad. Geo., p. 287, (Stenopora *exilis*,) Subcarb.

expansus, Ringuéberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 20. Not properly defined.

fibrosus se
filiosa, se
fletcheri,
in Ohio
lipora t
frondosus,
fruticosus
Ham. C
fruticosus,
arbuscu
furcatus,
pl. 37, i
fusiformi
Geo. S
vol. 4,
Chetete
gracilis, s
granulifer
lifera.
hamiltoni
Penin.
helderbergi
gie.
humilis,
pl. 37,
internasc
Held. C
irregularis
jamesi, see
lycopodon
don.
mackrothi
mackrothi
Americ
mammula
uiata.
microscop
Penin.
millepora
Mon. d
Coal M
montiform
formis.
monticula
8, fig. 5-
Low. H
muscatine
Nat. Sc
newberryi,
nodulosus,
onealli, see
ortoni, see
pavonia, s
petechialis
petropolit
p. 105.
pulchellus,
fied in
ticulipo
quadrang
gularis.
quadratus,
ramosus, s
rhombicus,
pella q
rugosus, s
rugosus, E
Montic
sigillaria

fibrosus see *Monticulipora fibrosa*.
filiosa, see *Monticulipora filiosa*.
fletcheri, Edwards & Haime, as identified in Ohio Pal., vol. 2, p. 197, is *Monticulipora ulrichi*.
frondosus, see *Monticulipora frondosa*.
fruticosus, Hall, 1876, *Illust. Foss.*, pl. 38, Ham. Gr.
furcatus, Hall, 1883, see *Monotrypella arbuscula*.
furcatus, Hall, 1876, *Illust. Devon. Foss.*, pl. 37, Ham. Gr.
fusiformis, Whitfield, 1878, *Ann. Rep. Geo. Sur. Wis.*, p. 70, and *Geo. Wis.*, vol. 4, p. 248, *Hud. Riv. Gr.* Not a *Chetetes*; probably a Bryozoan.
gracilis, see *Batostomella gracilis*.
granuliferus, see *Monotrypella granulifera*.
hamiltonensis, Winchell, 1866, *Rep. Low. Penin. Mich.*, p. 89, Ham. Gr.
helderbergiae, see *Ptychonema helderbergiae*.
humilis, Hall, 1876, *Illust. Devon. Foss.*, pl. 37, Up. Held. Gr.
internascens, Hall, 1881, *Bryozoans Up. Held. Gr.*, p. 4, Up. Held. Gr.
irregularis, see *Monticulipora irregularis*.
jamesi, see *Batostoma jamesi*.
lycoperdon, see *Monticulipora lycoperdon*.
mackrothi, Geinitz, 1846, (*Calamopora mackrothi*.) Grund, p. 586, Permian, American (?)
mammulatus, see *Monticulipora mammulata*.
microscopica, Winchell, 1866, *Rep. Low. Penin. Mich.*, p. 90, Ham. Gr.
milleporaceus, Edwards & Haime, 1851, *Mon. d. Pol. Foss. d. Terr. Pal.*, p. 272, Coal Meas.
moniliformis, see *Monticulipora moniliformis*.
monticulatus, Hall, 1883, *Rep. St. Geo.*, pl. 8, fig. 5-7, and *Pal. N. Y.*, vol. 6, p. 12, Low. Held. Gr.
muscatinensis, White, 1876, *Proc. Acad. Nat. Sci. Phil.*, p. 27, Devonian.
newberryi, see *Prasopora newberryi*.
nodulosus, see *Callopora nodulosa*.
onealli, see *Callopora onealli*.
ortoni, see *Atactoporella ortoni*.
pavonia, see *Ptilodictya pavonia*.
petechialis, see *Petigopora petechialis*.
petropoliatus, Pander, 1830, *Russ. reiche*, p. 105. Not an American species.
pulchellus, Edwards & Haime, as identified in Ohio Pal., vol. 2, p. 195, is *Monticulipora andrewsi*.
quadrangularis, see *Palescharcha quadrangularis*.
quadratus, see *Monotrypella quadrata*.
ramosus, see *Monticulipora ramosa*.
rhombicus, Nicholson, syn. for *Monotrypella quadrata*.
rugosus, see *Monticulipora rugosa*.
rugosus, Edwards & Haime, is a variety of *Monticulipora ramosa*.
sigillarioides, see *Callopora sigillarioides*.

sphaericus, see *Favosites sphaericus*.
spinigerus, Lonsdale, 1845, (*Stenopora spinigera*.) *Geo. Russ. and Urul Mts.*, vol. 1, p. 631, Coal Meas.
subglobosus, see *Monticulipora subglobosa*.
subpulchellus, see *Monticulipora subpulchella*.
tabulatus, see *Ptychonema tabulatum*.
tuberculatus, see *Spatiopora tuberculata*.
tumidus, Phillips, 1836, (*Calamopora tumida*.) *Geo. Yorkshire*, p. 200, Subcarb.
undulatus, see *Monticulipora undulata*.
venustus, see *Monticulipora venusta*.

CHONOPHYLLUM, FIG. 149.—*Chetetes tumidus*. Edwards & Haime, 1850, *Brit. Foss. corals*, p. lxxix. [Ety. *chonos*, funnel; *phyllon*, leaf.] Corallum simple, constituted, principally, by a series of infundibuliform tabulae, superposed and invaginated, upon the surface of which, equally developed septal radii extend from center to circumference; no walls or columella. Type C. perfoliatum.

belli, Billings, 1865, *Can. Nat. and Geo.*, vol. 2, p. 431, Clinton Gr.
capax, Hall, 1882, *Foss. Corals Niagara & Up. Held. Grs.*, p. 6, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 410, Niagara Gr.
ellipticum, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 233, Chemung Gr.
magnificum, Billings, 1860, *Can. Jour.*, vol. 5, p. 264, Up. Held. Gr.

FIG. 150.—*Chonophyllum niagarense*.

niagarense, Hall, 1852, (*Conophyllum niagarense*.) *Pal. N. Y.*, vol. 2, p. 114, Niagara Gr.
ponderosum, Rominger, 1876, *Foss. Corals*, p. 117, Ham. Gr.
sedaliense, White, 1880, 12th Rep. U. S. Geo. Sur. Terr., p. 157, Choteau limestone.
vadum, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 410, Niagara Gr.
validum, Hall, 1882, *Foss. Corals Niagara and Up. Held. Grs.*, p. 6, and 12th Rep. Geo. Ind., p. 272, Niagara Gr.

CHONOSTROITES, Edwards & Haime, 1851, *Pol. Foss. d. Terr. Pal.*, p. 299. [Ety. *konos*, cone; *steges*, covering.] Subhemispheric; corallites cylindrical, annulated, connected at the expansions, imperforate at the constrictions; mural pores where the corallites are contiguous; tabulae

62.

63.

numerous; septa consisting of short spines; growth by gemmation. Type *C. clappi*.

FIG. 151.—*Chonostegites ordinatus*.

clappi, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 299, Up. Held. Gr. *ordinatus*, Billings, 1859, (Halmephylum ordinatum,) Can. Jour.,

vol. 4, p. 139, Up. Held. Gr.

CLADOGRAPTUS, Geinitz, 1852, (*Cladograpus*,) Verst. Grauw. Sachs. and Emmons, Am. Geo., p. 107. [Ety. *klados*, twig; *grapho*, I write.] Serrations, or cells, arranged on the outer sides of branching stipes; no axis.

dissimularis, Emmons, 1856, Am. Geo., p. 107, Upper Taconic.

inequalis, Emmons, 1856, Am. Geo., p. 107, Upper Taconic.

CLADOPORA, Hall, 1852, Pal. N. Y., vol. 2, p. 137. [Ety. *klados*, twig; *poros*, pore.] Ramose or reticulate; branches cylindrical or compressed; terminations terete; corallites radiating from the axis, and opening upon the surface in rounded or subangular expanded mouths; tabulae and septal crests usually obsolete, sometimes present; corallites connected by mural pores. Type *C. seriata*.

alpenensis, Rominger, 1876, Foss. Corals, p. 51, Ham. Gr.

aspera, Rominger, 1876, Foss. Corals, p. 56, Up. Held. Gr.

caespitosa, Hall, 1852, Pal. N. Y., vol. 2, p. 138, Niagara Gr.

canadensis, Rominger, 1876, Foss. Corals, p. 49, syn. for *Pachypora frondosa*.

cervicornis, Hall, 1852, Pal. N. Y., vol. 2, p. 139, Niagara Gr.

dichotoma, Hall, 1858, Geo. Sur. Iowa, p. 478, Ham. Gr.

expatiata, Rominger, 1876, Foss. Corals, p. 57, Up. Held. Gr.

fibrosa, Hall, 1852, Pal. N. Y., vol. 2, p. 139, Niagara Gr.

imbricata, Rominger, 1876, Foss. Corals, p. 56, Up. Held. Gr.

64. *labiosa*, Billings, 1859, Can. Jour., vol. 4, p. 133, Up. Held. Gr.

laqueata, Rominger, 1876, Foss. Corals, p. 46, Niagara Gr.

lichenoides, Winchell & Marcy, 1865, Bost. Soc. Nat. Hist., vol. 1, p. 84, Niagara Gr.

lichenoides, Rominger, 1876, see *C. winchellana*.

macrophora, Hall, 1852, Pal. N. Y., vol. 2, p. 140, Niagara Gr.

magna, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 230, Up. Held. Gr.

multiopora, Hall, 1852, Pal. N. Y., vol. 2, p. 140, Niagara Gr.

palmata, Hall, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 231, Up. Held. Gr.

pinguis, Rominger, 1876, Foss. Corals, p. 53, Up. Held. Gr.

prolifera, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 230, Up. Held. Gr.

pulchra, Rominger, 1876, Foss. Corals, p. 54, Up. Held. Gr.

reticulata, Hall, 1852, Pal. N. Y., vol. 2, p. 141, Niagara Gr.

rimosa, Rominger, 1876, Foss. Corals, p. 53, Up. Held. Gr.

robusta, Rominger, 1876, Foss. Corals, p. 55, Up. Held. and Ham. Gr.

sarmentosa, Hall, 1876, Desc. New Spec. Foss., p. 3, and 11th Geo. Sur. Ind., p. 230, Niagara Gr.

seriata, Hall, 1852, Pal. N. Y., vol. 2, p. 137, Niagara Gr.

turgida, Rominger, 1876, Foss. Corals, p. 49, Up. Held. Gr.

verticillata, Winchell & Marcy, 1865, Bost. Soc. Nat. Hist., vol. 1, p. 84, Niagara Gr.

winchellana, S. A. Miller, 1883, 2d Ed. Am. Pal. Foss., p. 265, Up. Held. Gr.

Proposed for the species described by Rominger under the preoccupied name of *C. lichenoides*, in Foss. Corals, p. 47.

CLIMACOGRAPTUS, Hall, 1865,

Can. Org. Rem. Decade 2, p. 111. [Ety. *klimar*, ladder; *grapho*, I write.]

Simple stipes, with subparallel margins, having a range of cells on each side;

axis subquadrate; apertures transversely oval or subquadrate; denticles on the upper side of the apertures. Type *C. bicornis*.

antennarius, Hall, 1863, (Graptolithus antennarius,) Geo. of Can., p. 955, and Can. Org. Rem. Decade 2, p. 112, Quebec Gr.

bicornis, Hall, 1847, (Graptolithus bicornis,) Pal. N. Y., vol. 1, p. 268, Hud. Riv. Gr.

commersi, Walcott, 1886, Bull. U. S. Geo. Sur. No. 30, p. 93, Upper Taconic.

parvus, Hall, 1865, Can. Org. Rem. Decade 2, p. 57, Hud. Riv. Gr. Not defined.

typicalis, Hall, 1865, Can. Org. Rem. Decade 2, p. 57, Hud. Riv. Gr.

FIG. 153.—*Climacograptus bicornis*.

FIG. 152.—*Cladopora reticulata*.

CLISTOPHYLLA
vol. 8, p.
lent.] S

FIG. 154.—*Clistophylla* ger

Pol. Foss.
Held. Gr.

gabbii, Meek,
1, p. 8, C.

oneidaense,
pluridiale,
Ont., p. 1.

tumulus,
Arctic V.

CLONOGRAPTUS
Nat. Hist.

twig; gr.

numerous
cylindrical
small der.

C. rigidus
flexilis, Ha.

Geo. Sur.

Rem. Dec.

rigidus, H.

idus, G.

Can. Org.

Quebec C.

GENITES, Eich.

vol. 1, p.

gether.]

or ramoso

CLISTOPHYLLUM, Dana, 1848, Explor. Exped., vol. 8, p. 361. [Ety. *klision*, tent; *phyllon*, leaf.] Simple, branched or aggregate,

FIG. 154.—*Clistophyllum conigerum*.

with vertical radiating lamellae or septa; central area vesicular and forming a conical boss or columella, exterior to which the vesicular plates incline outward and upward; calyces deep.

Type *C. danianum*. austini, Salter, 1852, (Strophodonta austini,) Sutherland's Jour., vol. 2, p. cccxx, Devonian.

conigerum, Rominger, 1876, (Zaphrentis conigera,) Foss. Corals, p. 40, Up. Held. Gr. danianum, Edwards & Haime, 1854,

Pol. Foss. d. Terr. Pal., p. 412, Low. Held. Gr.

gabbii, Meek, 1864, Pal. California, vol. 1, p. 8, Carboniferous.

oneidaense, see *Acrophyllum oneidaense*. pluridiale, Nicholson, 1874, Pal. Prov. Ont., p. 21, Up. Held. Gr.

tumulus, Salter, 1855, Belcher's Last Arctic Voyage, vol. 2, p. 383, Carb.

CLONOGRAFTUS, Hall, 1873, Ann. and Mag. Nat. Hist., 4th ser., vol. 13. [Ety. *klon*, twig; *grapho*, I write.] Composed of numerous slender, regular branching, cylindrical stipes; cells small, forming small denticulations on one side. Type *C. rigidus*.

flexilis, Hall, 1858, (Graptolithus flexilis,) Geo. Sur. Can., p. 119, and Can. Org. Rem. Decade 2, p. 103, Quebec Gr.

rigidus, Hall, 1857, (Graptolithus rigidus,) Geo. Sur. Can., p. 121, and Can. Org. Rem. Decade 2, p. 105, Quebec Gr.

CENTES, Eichwald, 1829, Zoologia specialis, vol. 1, p. 186. [Ety. *koios*, living together.] Corallum incrusting, massive, or ramose; corallites vertical or oblique

to the surface, remote, imbedded in a coenenchyma; calices irregular, prominent, triangular, quincuncially arranged; lower margin most prominent; interstices increasing by age, and reducing the cavity of the cell-tubes; no septa; tabulae distinct; mural pores large and few. Type *C. clathrata*.

crassus, Rominger, 1876, (Limaria crassa,) Foss. Corals, p. 45, Niagara Gr.

falcatus, Prout, 1859, (Limaria falcata,) Trans. St. Louis Acad. Sci., vol. 1, p. 445, Up. Held. Gr.

fruticosus, Steining, 1834, (Limaria fruticosa,) Bull. Soc. Geo. France, vol. 1, p. 339, and Pal. N. Y., vol. 2, p. 143, Niagara Gr.

laminatus, Hall, 1852, (Limaria laminata,) Pal. N. Y., vol. 2, p. 143, Niagara Gr.

FIG. 155.—*Coenites lunatus*.

lunatus, Nicholson & Hinde, 1874, Can. Jour., p. 149, and Pal. Prov. Ont., p. 55, Niagara Gr.

ramulosus, Hall, 1852, (Limaria ramulosa,) Pal. N. Y., vol. 2, p. 142, Niagara Gr.

COLEOPHYLLUM, Hall, 1883, 12th Rep. Geo. Sur. Ind., p. 317.

[Ety. *koios*, sheath; *phyllon*, leaf.] Corallum simple; substance composed of closely arranged, invaginated tabulae, more or less oblique to the axis;

rays obscure; calices oblique. Type *C. romingeri*.

pyriforme, Hall, 1883, 12th Rep. Geo. Sur. Ind., p. 318, Up. Held. Gr.

romingeri, Hall, 1883, 12th Rep. Geo. Sur. Ind., p. 317, Up. Held. Gr.

COLUMNARIA, Goldfuss, 1826, Germ. Petref., p. 72. [Ety. *columnarius*, formed of columns.] Aggregate, corallites polygonal, longitudinally sulcated, but readily separable; no mural pores; tabulae numerous; septa rudimentary; increase by fission. Type *C. alveolata*.

alveolata, Goldfuss, 1826, Germ. Petref., p. 72, and Pal. N. Y., vol. 1, p. 47, Black Riv. Gr.

FIG. 156.—*Coleophyllum romingeri*.

65. blainvilli, Billings, 1858, Can. Nat. and Geo., vol. 3, and Rep. of Progr. Geo. Sur. Can., p. 168, Hud. Riv. Gr.

FIG. 157.—*Columnaria alveolata*.

66. carterensis, Safford, 1869, Geo. of Tenn., p. 285, Trenton Gr.
divergens, Troost, 1840, 5th Geo. Rep. Tenn., p. 72, Devonian.
67. erratica, Billings, 1858, Can. Nat. and Geo., vol. 3, and Rep. of Progr. Geo. Sur. Can., p. 167, Trenton Gr.
goldfussi, Billings, 1858, Can. Nat. and Geo., vol. 3, and Rep. of Progr. Geo. Sur. Can., p. 168, Hud. Riv. Gr.
halti, Nicholson, 1879, Tabulate corals, syn. for *C. alveolata*.
herzeri, Rominger, 1876, syn. for *Favistella stellata*.
68. incerta, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 128, Chazy Gr.
inaequalis, Hall, 1852, Pal. N. Y., vol. 2, p. 223, Coralline limestone.
intermedia, Eaton, 1832, Geo. Text-book, p. 41. Not recognized.
mammillaris, Castelnau. Not recognized.
multiradiata, Castelnau, 1843, Syst. Sil., p. 44. Not recognized. Probably same as *Favistella stellata*.
69. parva, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 428, Chazy Gr.
70. rigida, Billings, 1858, Can. Nat. and Geo., vol. 3, and Rep. of Progr. Geo. Sur. Can., p. 167, Hud. Riv. Gr.
sutherlandi, Salter, 1852, Sutherland's Jour., vol. 2, p. cxxxii, Devonian.
troosti, Castelnau, 1843, Syst. Sil., p. 44, syn. for *Lonsdaleia papillata*.
Columnopora, Nicholson, 1874, London Geo. Mag. N. S., vol. 1, p. 253, and Ohio Pal., vol. 2, p. 186, syn. for *Calapocia*.
cribriformis, see *Calapocia cribriformis*.
COMBOPHYLLUM, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxvii. [Ety. *kom-bos*, strip of cloth; *phyllon*, leaf.] Coralium, in form like *Cyclolites*; single septal fossula; septa exsert and regularly radiate. Type *C. osismorum*.
multiradiatum, Meek, 1868, Trans. Chi. Acad. Sci., p. 84, Devonian.
Conophyllum, Hall, 1852, Pal. N. Y., vol. 2, syn. for *Chonophyllum*.
niagarensis, see *Chonophyllum niagarensis*.
Constellaria, Dana, 1848, syn. for *Stellipora*.

constellata, syn. for *Stellipora antheloidea*.
fischeri, see *Stellipora fischeri*.
floridensis, see *Stellipora florida*.
polystomella, see *Stellipora polystomella*.

CRASPEDOPHYLLUM, Dybowski, 1873, Beschreibung neuen aus Nordamerika, Stammenden, Devonischen art der *Zoantharia rugosa*, p. 153. [Ety. *kraspedos*, an edge; *phyllon*, leaf.] Probably a syn. for *Heliophyllum*. Type *C. americanum*.

americanum, Dybowski, 1873, Besch. n. a. Nord. Stamm. Dev. a. d. Zoanth. *rugosa*, p. 153, Up. Held. Gr.

CREPIDOPHYLLUM, Nicholson & Thompson, 1877, Proc. Roy. Soc. Edinburgh, vol. 9, p. 149. [Ety. *krepsis*, horseshoe; *phyllon*, leaf.] Distinguished from *Heliophyllum*, by the central part of the tabulate area being shut off from the rest of the visceral chamber by a secondary investment, in the form of a central pipe, which is crossed, by tabulae; this pipe is sometimes open or horseshoe-shaped. Type *C. archiaci*.

archiaci, Billings, 1860, (Diphyphyllum *archiaci*.) Can. Jour., vol. 5, p. 260, Ham. Gr.

subcespitosum, Nicholson, 1874, (Heliophyllum *subcespitosum*.) Lond. Geo. Mag. n. ser., vol. 1, p. 58, Ham. Gr.

CYATHAXONIA, Michelin, 1846, Icon. Zooph., p. 258. [Ety. *kyathos*, cup; *axones*, a tablet made to turn on its axis.] Simple; calice deep;

columella styliform, strong and prominent; septa extending to the columella; the place of one of them occupied, by a deep depression or septal fossula. Type *C. cornu*.

columellata, Hall, 1882, Foss. Corals Niagara and Up. Held. Gr., and 35th Rep. N. Y. Mus. Nat. Hist., p. 415, Niagara Gr.

cynodon, Rafinesque & Clifford, 1820, (Turbinolia *cynodon*.) Monog. d. Turbinolides in Ann. d. Phys. d. Brux., t. 5, p. 234, Waverly Gr.

distorta, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 526, Coal Meas.

herzeri, Hall, 1882, Foss. Corals Niagara and Up. Held. Gr., p. 11, and 12th Rep. Ind. Geo., p. 275, Niagara Gr.

profunda, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 323, Carboniferous.

FIG. 158.
Crepidophyllum subcespitosum

prolifer
wiscon

FIG.
Cyathaxo

being
exteri
Type
agglomer
ognize
ammonio
nized.

72. antioch

vol. 1,
arboresce

48. N
arctium

Sci., p
articulat

articul
p. 97,
arctifossa

agars,
12th H
Held C

allas, Ca
Not rec

billingsi,
287. S

bullatum,
and Up

N. Y. f
Held. C

bullatur
Niagara

35th Re
412, Ni

cespitosu
p. 60, U

calyculare
Lauds, p

canalicula
Niagara

35th Re
Up. Hel

ceratites, G

coalitum,
p. 108, U

coherens,

prolifera, see *Lophophyllum proliferum*.
wisconsinensis, Whitfield, 1878, Ann.

FIG. 159.
Cyathaxonia herzeri.

being filled with vesicular dissepiments; exterior wall provided with an epitheca. Type *C. caespitosum*.

agglomeratum, Castelnau, 1843. Not recognized.

ammonis, Castelnau, 1843. Not recognized.

72. *anticostiense*, Billings, 1862, Pal. Foss., vol. 1, p. 109, Anticosti Gr., Div. 4.

arborescens, Castelnau, 1843, Syst. Sil., p. 48. Not recognized.

arcticum, Meek, 1868, Trans. Chi. Acad. Sci., p. 79, Devonian.

articulatum, Wahlenberg, (Madrepores articulatus.) Nov. Act. Upsal., vol. 8, p. 97, Up. Sil.

arctifossa, Hall, 1882, Foss. Corals Niagara and Up. Held. Gr., p. 40, and 12th Rep. Ind. Geo., p. 297, Up. Held. Gr.

atlas, Castelnau, 1843, Syst. Sil., p. 47. Not recognized.

billingsi, Dawson, 1868, Acad. Geol., p. 287. Subcarboniferous.

bullatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 41, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 445, Up. Held. Gr.

bullulatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 12, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 412, Niagara Gr.

caespitosum, Goldfuss, 1826, Petref. Germ., p. 60, Up. Held. Gr.

calyculare, Owen, 1840, Rep. on Mineral Lands, p. 69, Devonian.

canaliculatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 39, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 443, Up. Held. Gr.

ceratites, Goldfuss. Not American.

coalitum, Rominger, 1876, Foss. Corals, p. 108, Up. Held. Gr.

coherens, Hall, 1882, Foss. Corals Ni-

agara and Up. Held. Grs., p. 41, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 445, Up. Held. Gr.

conatum, Hall, 1876, Illust. Dev. Foss., pl. 31, Ham. Gr.

concentricum, Hall, 1882, Foss. Corals Niagara and Up. Held.

Gr., p. 42, and 12th Rep. Geo.

Sur. Ind., p. 316, Up. Held. Gr.

conicum, Castelnau, 1843, Syst. Sil., p. 48. Not recognized.

corinthium, Owen, 1840, Rep. on Minn. Lands, p. 69, Devonian.

cratatum, Rominger, 1876, Foss. Corals, p. 108, Ham. Gr.

depressum, Hall, 1882, Foss. Corals Niagara and Up. Held. Gr., p. 40, and 12th Rep. Ind. Geol., p. 298, Up. Held. Gr.

dianthus, Goldfuss, 1826, Germ. Petref., p. 54, Onondaga Gr.

dilatatum, Castelnau, 1843, Syst. Sil., p. 48. Not recognized.

distinctum, Castelnau, 1843, Syst. Sil., p. 49. Not recognized.

d'orbignyi, Castelnau, 1843, Syst. Sil., p. 49. Not recognized.

eriphyle, Billings, 1862, Pal. Foss., vol. 1, p. 111, Anticosti Gr., Div. 4.

euryone, Billings, 1862, Pal. Foss., vol. 1, p. 110, Anticosti Gr., Div. 4.

excentricum, Goldfuss. Not American.

exfoliatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 39, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 443, Up. Held. Gr.

flexuosum, Owen, syn. for *Campophyllum torquium*.

galerum, Hall, 1876, Illust. Dev. Foss., pl. 32, Ham. Gr.

geniculatum, Rominger, 1876, Foss. Corals, p. 103, Ham. Gr.

gigas, Yandell & Shumard, syn. for *Zaphrentis gigantea*.

goldfussi, Castelnau, 1843, Syst. Sil., p. 47. Not recognized.

goliath, Castelnau, 1843, Syst. Sil., p. 47. Not recognized.

gracile, Troost, 5th Rep. Tenn., Subcarb. Not recognized.

gradatum, Hall, 1876, Illust. Dev. Foss., pl. 31, Ham. Gr.

helianthoides, Goldfuss, see *Heliophyllum halli*.

houghtoni, Rominger, 1876, Foss. Corals, p. 104, Ham. Gr.

impositum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 40, and

FIG. 160.—*Cyathophyllum caespitosum*. *a*, vertical section. *b*, transverse section. *c*, vertical section.

cratatum, Rominger, 1876, Foss. Corals, p. 108, Ham. Gr.

depressum, Hall, 1882, Foss. Corals Niagara and Up. Held. Gr., p. 40, and 12th Rep. Ind. Geol., p. 298, Up. Held. Gr.

dianthus, Goldfuss, 1826, Germ. Petref., p. 54, Onondaga Gr.

dilatatum, Castelnau, 1843, Syst. Sil., p. 48. Not recognized.

distinctum, Castelnau, 1843, Syst. Sil., p. 49. Not recognized.

d'orbignyi, Castelnau, 1843, Syst. Sil., p. 49. Not recognized.

eriphyle, Billings, 1862, Pal. Foss., vol. 1, p. 111, Anticosti Gr., Div. 4.

euryone, Billings, 1862, Pal. Foss., vol. 1, p. 110, Anticosti Gr., Div. 4.

excentricum, Goldfuss. Not American.

exfoliatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 39, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 443, Up. Held. Gr.

flexuosum, Owen, syn. for *Campophyllum torquium*.

galerum, Hall, 1876, Illust. Dev. Foss., pl. 32, Ham. Gr.

geniculatum, Rominger, 1876, Foss. Corals, p. 103, Ham. Gr.

gigas, Yandell & Shumard, syn. for *Zaphrentis gigantea*.

goldfussi, Castelnau, 1843, Syst. Sil., p. 47. Not recognized.

goliath, Castelnau, 1843, Syst. Sil., p. 47. Not recognized.

gracile, Troost, 5th Rep. Tenn., Subcarb. Not recognized.

gradatum, Hall, 1876, Illust. Dev. Foss., pl. 31, Ham. Gr.

helianthoides, Goldfuss, see *Heliophyllum halli*.

houghtoni, Rominger, 1876, Foss. Corals, p. 104, Ham. Gr.

impositum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 40, and

- 12th Rep. Geo. Sur. Ind., p. 299, Up. Held Gr.
75. *interruptum*, Billings, 1862, Pal. Foss., vol. 1, p. 109, Mid. Sil.
- intertrium*, Hall, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 416, Niagara Gr.
- intervesicula*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 38, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 442, Up. Held. Gr.
- juvenc*, Rominger, 1876, Foss. Corals, p. 101, Up. Held. Gr.
- lescuri*, Edwards & Haime, 1851, Pol. Foss. J. Terr. Pal., p. 371, Onondaga Gr.
- melchiorii*, Castelnau, 1843, Syst. Sil., p. 48. Not recognized.
- nanum*, Hall, 1876, Illust. Dev. Foss., pl. 22, Ham. Gr.
- nepos*, Hall, 1876, Illust. Dev. Foss., pl. 22, Ham. Gr.
- nevadense*, Meek, 1877, U. S. Geo. Sur. 40th Parallel, vol. 4, p. 60, Carboniferous.
76. *nympheale*, Billings, 1862, Pal. Foss., vol. 1, p. 111, Mid. Sil.
- palmeri*, Meek, 1877, U. S. Geo. Sur. 40th Parallel, vol. 4, p. 33, Devonian.
- palum*, Hall, 1876, Illust. Dev. Foss., pl. 31, Ham. Gr.
- panicum*, Winchell, 1866, Rep. Low Penin. Mich., p. 90, Ham. Gr.
- partitum*, Winchell, 1866, Rep. Low. Penin. Mich., p. 90, Ham. Gr.
77. *pasithea*, Billings, 1862, Pal. Foss., vol. 1, p. 112, Mid. Sil.
78. *pelagicum*, Billings, 1862, Pal. Foss., vol. 1, p. 108, Anticosti Gr., Div. 2.
79. *pennanti*, Billings, 1862, Pal. Foss., vol. 1, p. 107, Mid. Sil.
- perforosulatum*, Hall, 1882, Foss. Corals, Niagara and Up. Held. Gr., p. 42 and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 446, Up. Held. Gr.
- perlamellosum*, Hall, 1876, Illust. Devon. Foss., pl. 39, Up. Held. Gr.
- perplicatum*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 42, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 447, Up. Held. Gr.
- piethorni*, Salter, 1852, (Strephodes pithorni,) Sutherland's Jour., vol. 2, p. cccxxx, Devonian.
- plicatum*, Castelnau, 1843, Syst. Sil., p. 48. Not recognized.
- plicatum*, Goldfuss, 1826, Germ. Petref. Not American.
- profundum*, see *Streptelasma profundum*.
- pusillum*, Conrad, 1848. Not properly defined.
- quadrigenum*, Goldfuss. Not American.
- radicula*, Rominger, 1876, Foss. Corals, p. 109, Niagara Gr.
- robustum*, Hall, 1876, Illust. Devon. Foss., pl. 22, Ham. Gr.
- robustum*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 43, Up. Held. Gr. The name was preoccupied.
- rollini*, Castelnau, 1843, Syst. Sil., p. 49. Not recognized.

rugosum, Hall, 1843, (Astrea rugosa,) Geo. Sur. 4th Dist. N. Y., p. 159, Up. Held. Gr.

scalenum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 42, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 446, Up. Held. Gr.

scyphus, Rominger, 1876, Foss. Corals, p. 103, Ham. Gr.

septatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 41, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 445, Up. Held. Gr.

shumardi, see *Amplexus shumardi*.

solitarium, Billings, 1866, Catal. Sil. Foss. Antic., p. 93, Clinton and Niagara Grs.

striatum, Castelnau, Syst. Sil., p. 48. Not recognized.

subcaespitosum, Meek, 1872, 6th Rep. Hayden's Geo. Sur. Terr., p. 470, and U. S. Geo. 40 Parallel, vol. 4, p. 60, Sub-carboniferous.

torquum, see *Campophyllum torquum*.

turbatum, Goldfuss. Not American.

undulatum et multiplicatum, Owen, 1840, Rep. on Min. Lands. Not binomial.

validum, Hall, 1876, Illust. Devon. Foss., pl. 39, Up. Held. Gr.

vanuxemi, Hall, 1859, figured without specific name in 1843, Geo. Rep. 4th Dist. N. Y., Tab. 49, fig. 3, 3a, Ham. Gr.

vermiculare, Owen, syn. for *Campophyllum torquum*.

vesiculatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 41, and 12th Geo. Sur. Ind., p. 297, Up. Held. Gr.

vesiculosum, see *Cystiphyllum vesiculosum*.

vicinum, Castelnau, 1843, Syst. Sil., p. 48. Not recognized.

wahlenbergi, Billings, 1862, Pal. Foss., vol. 1, p. 108, Anticosti Gr., Div. 3.

zenkeri, Billings, 1860, Can. Jour., vol. 5, p. 262, Up. Held. Gr.

Cyathopora iowensis, Owen, see *Striatopora iowensis*. There is no genus *Cyathopora*, and if Dr. Owen did not intend to refer his species to *Cyathophora*, then he failed to establish a genus, by neglecting to define it.

CYCLOGRAFTUS. Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 42. [Ety. kuklos, disk; grapho, I write.] A circular disk, with stipes radiating from the radicle to the margin and in a free manner beyond. Type *C. rotadentatus*.

rotadentatus, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 42, Niagara Gr.

FIG. 102.—Cyclograftus rotadentatus.

FIG. 101.—Cyathophyllum rugosum.

Cyclolites, Lamarek, 1801, Syst. Anim. sans Vert., p. 360. Not a Paleozoic genus. *rotuloides*, see *Palæocyclus rotuloides*.

CYSTIPHYLITES. Recently proposed genus but the reference mislaid. [Ety. *kustis*, a small cavity; *phoros*, bearing; *lithos*, stone.] Corallum compound, formed of superimposed series of cups, which in vertical sections appear as layers of unequal, vesiculate plates, resembling *Cystiphyllum*; layers radiated, margins of cells broad, expanded, and confluent. Type *C. major*. Proposed instead of *Vesicularia*, Rominger, which was preoccupied.

major, Rominger, 1876, (*Vesicularia major*), Foss. Corals, p. 135, Niagara Gr. minor, Rominger, 1876, (*Vesicularia minor*), Foss. Corals, p. 136, Niagara Gr. *variolosus*, Rominger, 1876, (*Vesicularia variolosa*), Foss. Corals, p. 136, Niagara Gr. *CYSTIPHYLLUM*, Lonsdale, 1839, Murch. Sil. Syst., p. 691. [Ety. *kustis*, cavity; *phyllo-*, leaf.] Simple, turbinate, or cylindrical, rarely aggregate; interior filled with vesicular tissue; septa rudimentary or absent. Type *C. siluriense*.

aggregatum, Billings, 1859, Can. Jour., vol. 3, p. 136, Ham. Gr.

americanum, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 464, Ham. Gr.

americanum var. *arcticum*, Meek, 1868, Trans. Chi. Acad. Sci., p. 80, Ham. Gr.

bifurcatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 55, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 460, Up. Held. Gr.

bipartitum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 55, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 459, Up. Held. Gr.

conifolius, Hall, 1876, Illust. Dev. Foss., pl. 50, Ham. Gr.

corrugatum, Hall, 1876, Illust. Devon. Foss., pl. 29, Ham. Gr.

crateriforme, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 57, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 461, Up. Held. Gr.

cylindricum, Lonsdale. Not American.

fruticosum, Nicholson, 1875, Geo. Mag., vol. 2, N. S., p. 32, Corniferous Gr.

grande, Billings, 1859, Can. Jour., vol. 4, p. 138, Corniferous Gr.

granilineatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 14, and 12th Rep. Ind. Geo. p. 274, Niagara Gr.

huronense, Billings, 1860, Catal. Sil. Foss. Antic., p. 92, Clinton and Niagara Grs.

infundibulum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 58, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 462, Up. Held. Gr.

latiradius, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 57, and 12th Rep. Geo. Sur. Ind., p. 304, Up. Held. Gr.

maritimum, Billings, 1862, Pal. Foss., vol. 1, p. 112, Mid. Sil.

mundulum, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 234, Chemung Gr.

muricatum, Hall, 1882 Foss. Corals, Niagara and Up. Held. Grs., p. 56, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 460, Up. Held. Gr.

nanum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 56, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 460, Up. Held. Gr.

obliquum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 58, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 462, Up. Held. Gr.

 

FIG. 163. — *Cystiphyllum ohioense*.

Rep. Geo. Sur. Ind., p. 262, Up. Held. Gr.

quadrangulare, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 56, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 460, Up. Held. Gr.

scalatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 59, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 463, Up. Held. Gr.

senecaense, Billings, 1859, Can. Jour., vol. 4, p. 137, Devonian.

squamosum, Nicholson, 1875, Geo. Mag. N. S., vol. 2, p. 31, Corniferous Gr.

striatura, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 59, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 463, Up. Held. Gr.

sulcatum, Billings, 1859, Can. Nat. and Geo., vol. 3, p. 136, Corniferous Gr.

superbum, Nicholson, 1875, Geo. Mag. vol. 2, N. S., p. 33, Ham. Gr.

supraplanum, Hall, 1882, Foss. Corals Ni-

FIG. 164. — *Cystiphyllum vesiculosum*.

superbum, Nicholson, 1875, Geo. Mag. vol. 2, N. S., p. 33, Ham. Gr.

 

- agars and Up. Held. Grs., p. 57, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 461, Up. Held. Gr.
- tenuiradius, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 56, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 460, Up. Held. Gr.
- varians, Hull, 1876, Illust. Devon. Foss., pl. 29, Ham. Gr.
- vesiculosum, Goldfuss, 1826, (Cyathophyllum vesiculosum,) Germ. Petref., p. 58, Devonian.
- CYSTOSTYLUS**, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis. and Geo. Wis., vol. 4, p. 273. [Ety. *kustis*, cavity; *stylos*, stalk.] Aggregate, cylindrical, corallites in contact or united by transverse filaments; increase by bifurcation, structure cystose as in Cystiphyllum; formed by imperfect transverse plates arranged in circular, funnel-formed order; septa and tabulae obsolete. Type *C. typicus*.
- infundibulum, Whitfield, 1878, (Syringopora infundibulum,) Ann. Rep. Geo. Sur. Wis., p. 79, and Geo. Wis., vol. 4, p. 274, Niagara Gr.
- typicus, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis. and Geo. Wis., vol. 4, p. 274, Niagara Gr.
- DANIA**, Edwards & Haime, 1849, Comp. Rend., t. 29, p. 261. [Ety. proper name.] Corallum having most of the characters of Chetetes, but with the tabulae connected through the corallites so as to divide the mass into parallel strata. Type *D. huronica*.
- huronica, Edwards & Haime, 1849, Comp. Rend., t. 29, p. 261, Up. Sil.
- DAWSONIA**, Nicholson, 1873, Ann. Mag. Nat. Hist., 4th ser., vol. 12. [Ety. proper name.] Supposed to be the ovarian vesicles of Graptolites. Type *D. campanulata*.
- acuminata, Nicholson, 1873, Ann. Mag. Nat. Hist., 4th ser., vol. 12, Quebec Gr.
- campanulata, Nicholson, 1873, Ann. Mag. Nat. Hist., vol. 12, Quebec Gr.
- rotunda, Nicholson, 1873, Ann. Mag. Nat. Hist., vol. 12, Quebec Gr.
- tenuistriata, Nicholson, 1873, Ann. Mag. Nat. Hist., vol. 12, Quebec Gr.
- DEKAYELLA**, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 155. [Ety. diminutive of Dekayia.] Ramose, interstitial cells; spiniform tubuli of two kinds, larger ones arranged as in Dekayia, others more numerous; diaphragms in both sets of tubes. Type *D. obscura*.
- obscura, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 89, Hud. Riv. Gr.
- robusta, Foord, 1884, Ann. and Mag. Nat. Hist., 5th ser., vol. 14, p. 341, Hud. Riv. Gr.
- DEKAYIA**, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 277. [Ety. proper name.] Distinguished from Monticulipora by having little protuberances on the surface between the angles of the corallites. Type *D. aspera*.
- appressa, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 152, Hud. Riv. Gr.
- aspera, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 278, Hud. Riv. Gr.
- atrita, syn. for *D. aspera*.
- multispi- Fig. 165.—*Dekayia aspera*, natural size, and magnified.
- nosa, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 154, Hud. Riv. Gr.
- paupera, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 153, Hud. Riv. Gr.
- pelliculata, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 150, Hud. Riv. Gr.
- trentonensis, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 151, Trenton Gr.
- DENDROGRAPTUS**, Hall, 1865, Can. Org. Rem., Decade 2, p. 126. [Ety. *dendron*, tree; *grapho*, I write.] Simple or aggregate; foot-stalk strong, sometimes with a root-like bulb; ramified above into slightly divergent branches, celluliferous on one side. Type *D. hallanus*.
- compactus, Walcott, 1879, Utica Slate and related formations, p. 21, Utica Slate.
- dawsoni, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 18, Niagara Gr.
- diffusus, Hall, 1865, Can. Org. Rem., Decade 2, p. 132, Quebec Gr.
- divergens, Hall, 1865, Can. Org. Rem., Decade 2, p. 129, Quebec Gr.
- dubius, n. sp. Proposed instead of *D. simplex*, Spencer, in Bull. No. 1, Mus. Univ. St. Mo., p. 17, which was preoccupied. Niagara Gr.
- erectus, Hall, 1865, Can. Org. Rem., Decade 2, p. 130, Quebec Gr.
- flexuosus, Hall, 1865, Can. Org. Rem., Decade 2, p. 127, Quebec Gr.
- frondosus, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 18, Niagara Gr.
- fruticosus, Hall, 1865, Can. Org. Rem., Decade 2, p. 131, Quebec Gr.
- gracilis, Hall, 1865, Can. Org. Rem., Decade 2, p. 132, Quebec Gr.
- gracillimus, Lesquereux, 1877, Proc. Am. Phil. Soc. p. 164, (Psilophyton gracillimum,) Hud. Riv. Gr.
- hallanus, Prout, 1851, (Graptolithus hallanus,) Am. Jour. Sci. 3d ser., vol. 11, p. 187, Potsdam sandstone.
- novellus, Hall, 1879, Desc. New Spec. Foss., p. 2, and 11th Rep. Geo. Sur. Ind., p. 225, Niagara Gr.

FIG. 166.—*Dendrograptus hallanus*.

FIG. 167.—*Dicranograptus* ramosus net; nemo of flabellations, con branches,

praegracilis, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 19, Niagara Gr.
primordialis, Matthew, 1885, Trans. Roy. Soc. Can., p. 31, St. John Gr.
ramosus, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 17, Niagara Gr.
simplex, Walcott, 1879, Utica Slate and related formations, p. 20, Utica Slate.
simpler, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 17. The name was preoccupied. See *D. dubius*.

spinus, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 19, Niagara Gr.
striatus, Hall, 1865, Can. Org. Rem., Decade 2, p. 129, Quebec Gr.
tenuiramosus, Walcott, 1879, Utica Slate and related formations, p. 21, Utica Slate.

DENDROPORA, Michelin, 1846, Icon. Zooph., p. 187. [Ety. *dendron*, tree; *poros*, pore.] Corallum arborescent, with very delicate, smooth branches; calices distant, and surrounded by a narrow, obtuse margin; septa small, but distinct. Type *D. explicita*.

alternans, Rominger, 1876, Foss. Corals, p. 64, Ham. Gr.

neglecta, Rominger, 1876, Foss. Corals, p. 63, Up. Held. Gr.

ornata, see *Trachypora ornata*.

proboscoidalis, Rominger, 1876, Foss. Corals, p. 65, Ham. Gr.

reticulata, Rominger, 1876, Foss. Corals, p. 65, Ham. Gr.

Dichograptus, syn. for *Graptolithus*.

DICRANOGRAPTUS, Hall, 1865, Can. Org. Rem., Decade 2, p. 46. [Ety. *dikranos*, two pointed; *grapho*, I write.] The lower part of the stipe has a row of cells on each side, but above, the stipe bifurcates, and has cells only on the outer side of each bifurcation. Type *D. ramosus*.

divaricatus, Hall, 1859, (*Graptolithus divaricatus*), Pal. N. Y., vol. 3, p. 513, Hud. Riv. Gr.

furcatus, Hall, 1847, (*Graptolithus furcatus*), Pal. N. Y., vol. 1, p. 273, Utica Slate.

ramosus, Hall, 1847, (*Graptolithus ramosus*), Pal. N. Y., vol. 1, p. 270, Utica Slate.

sextans, Hall, 1847, (*Graptolithus sextans*), Pal. N. Y., vol. 1, p. 273, Utica Slate.

DICTYONEMA, Hall, 1852, Pal. N. Y., vol. 2, p. 174.

[Ety. *dictyon*, net; *nema*, thread.] Fronds consisting of flabelliform or funnel-shaped expansions, composed of slender, radiating branches, which frequently bifurcate

as they recede from the base; branches united laterally by transverse dissepiments; exterior striated; interior surface cellulariferous or serrate. Type *D. retiforme*.

expansum, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 25, Niagara Gr.
fenestratum, Hall, 1851, in Foster and Whitney's Rep. on Lake Superior Land Dist., p. 223, Up. Held. Gr.
gracile, Hall, 1852, Pal. N. Y., vol. 2, p. 175, Niagara Gr.

grande, Nicholson, 1873, Ann. Mag. Nat. Hist., 4th ser., vol. 12, Quebec Gr.

irregulare, Hall, 1865, Can. Org. Rem., Decade 2, p. 136, Quebec Gr.

murrayi, Hall, 1865, Can. Org. Rem., Decade 2, p. 138, Quebec Gr.

neenah, Hall, 1861, Geo. Rep. Wis., p. 17, Trenton Gr.

pergracile, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 181, and Acad. Geo., p. 563, Niagara Gr.

quadrangulare, Hall, 1865, Can. Org. Rem., Decade 2, p. 138, Quebec Gr.

FIG. 168.—*Dictyonema retiforme*.

retiforme, Hall, 1843, (*Gorgonia retiformis*), Geo. Rep. 4th Dist. N. Y., p. 115, and Pal. N. Y., vol. 2, p. 174, Niagara Gr.

robustum, Hall, 1865, Can. Org. Rem., Decade 2, p. 137, Quebec Gr.

splendens, Billings, 1874, Pal. Foss., vol. 2, p. 12, Gaspé limestone No. 1, Up. Sil.

tenellum, Spencer, 1878, Can. Nat., vol. 8, and Bull. No. 1, Mus. Univ. St. Mo., p. 26, Niagara Gr.

websteri, Dawson, 1860, Can. Nat. and Geo., vol. 5, and Acad. Geo., p. 563, Niagara Gr.

DIDYNOGRAPTUS, McCoy, 1851, Brit. Pal. Foss., p. 3-9. [Ety. *didymos*, double; *grapho*, I write.] Consisting of forked stipes, straight or curved; one cellulariferous side. Type *D. murchisoni*.

FIG. 167.—*Dicanograptus ramosus*.

128 125
122 120

10

caduceus, Salter, 1853, (*Graptolithus caduceus*.) Quar. Jour. Geo. Soc., vol. 9, p. 87, Quebec Gr.

FIG. 169.—*Didymograptus geminus*.

geminus, Hisinger, 1840, (*Prionotus geminus*.) Leth Suecia, Supp. 2, p. 5, pl. 38, Quebec Gr.

DIPHYPHYLLUM, Lonsdale, 1845, Russ. and Ural Mts., vol. 1, p. 624. [Ety. *diphya*, division; *phyllon*, leaf.] Corallum simple, composite, increasing by lateral gemmation; corallites tall, cylindrical, connected by epithecal or radiciform expansions, with each other; central area occupied by tabulae; circumscribed by an inner wall; exterior vesicular zone occupied by septa, which are confined between the outer and inner mural investment; no columella. Type *D. conicum*.

adnatum, Hall, 1882, Foss. Corals Niagara and Up. Held Grs., p. 54, and 12th Rep. Geo. Sur. Ind., p. 303, Up. Held. Gr.

apertum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 54, and 12th Rep. Geo. Sur. Ind., p. 303, Up. Held. Gr.

archiaci, see *Crepidophyllum archiaci*. *arundinaceum*, Billings, 1859, Can. Jour., vol. 4, p. 134, Corniferous limestone.

breve, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 55, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 459, Up. Held. Gr.

caespitosum, Hall, 1852, (*Diplophyllum caespitosum*.) Pal. N. Y., vol. 2, p. 116, Niagara Gr.

coralliferum, Hall, 1852, (*Diplophyllum coralliferum*.) Pal. N. Y., vol. 2, p. 322, Coralline limestone.

cylindraceum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 54, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 458, Up. Held. Gr.

fasciculum, Meek, 1877, U. S. Geo. Sur. 40th Parallel, vol. 4, p. 29, Devonian.

gigas, Rominger, 1876, Foss. Corals, p. 125, Niagara Gr.

gracile, McCoy, 1854, Brit. Pal. Foss., p. 88, Up. Held. Gr.

huronium, Rominger, 1876, Foss. Corals, p. 121, Niagara Gr.

rectiseptatum, Rominger, 1876, Foss. Corals, p. 124, Ham. Gr.

rugosum, Edwards & Haime, 1851, (*Eridophyllum rugosum*.) Pol. Foss. des Terr. Pal., p. 424, Up. Held. Gr.

simcoense, Billings, 1859, (*Eridophyllum simcoense*.) Can. Jour., vol. 4, p. 131, Up. Held. Gr.

stramineum, Billings, 1859, Can. Jour., vol. 4, p. 135, Corniferous Gr.

strictum, Edwards & Haime, 1851, (*Eri-*

dophyllum strictum.) Poll. Foss. des Terr. Pal., p. 424, Up. Held. Gr.

tumidulum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 55, and 12th Rep. Geo. Sur. Ind., p. 303, Up. Held. Gr.

FIG. 170.—*Diphyphyllum stramineum*.

vennori, Billings, 1865, (*Eridophyllum vennori*.) Can. Nat. and Geo., 2d ser., vol. 2, p. 431, Clinton Gr.

verneuianum, Edwards & Haime, 1850, (*Eridophyllum verneuianum*.) Brit. Foss. Corals, p. lxxi, and Pol. Foss. des Terr. Pal., p. 424, Up. Held. Gr.

DIPLOGRAPTUS, McCoy, 1854, (*Diplograptus*.) Brit. Pal. Rocks, p. 3. [Ety. *diploos*, duplex; *grapho*, I write.] Stipes simple, flattened, or quadrangular; cellules, in single series, on the two sides of a double central axis; cellules oblique, opening toward the apex; cell denticles prominent, often mucronate. Type *D. foliaceus*.

amplexicaulis, Hall, 1847, (*Graptolithus amplexicaulis*.) Pal. N. Y., vol. 1, p. 79, Trenton Gr.

angustifolius, Hall, 1859, (*Graptolithus angustifolius*.) Pal. N. Y., vol. 3, p. 515, Hud. Riv. Gr.

ciliatus, Emmons, 1856, Am. Geo., p. 105, Up. Taconic.

dissimilaris, Emmons, 1856, Am. Geo., p. 105, Up. Taconic.

foliaceus, (?) Murch, 1839, (*Graptolites foliaceus*.) Murch. Sil. Syst., p. 695, Hud. Riv. Gr.

foliosus,
Up. T

FIG. 171.
Diplograptus foliosus.

inutilis,
cade 2
laciniat
236, U
marcidu
cidus,
Riv. G
mucrona
mucro
Hud. I
obliquus
106, U
peosta, I
Geo. R
pristinif
pristin
and Ca
Quebec
pristis, (Y., vol
putillus,
cade 2,
rugosus,
Up. Ta
rectangul
Rocks,
secalinus,
Pal. N.
simplex
simplex,
plex.)
N. Y.,
spinulosu
spinulo
Hud. R
whitfieldi
fieldi,
Riv. Gr
Diplophyllum
p. 115,
caespitosum
coralliferum
liferum.
DIPLOTRYPA
Corals,
trypa, l
the la
tabulae
clusters
lites an
pletely
numero
infida, Ul
Minn.,
milleri, U
Hist., vo

foliosus, Emmons, 1856, Am. Geo., p. 105, Up. Taconic.

FIG. 171.
Diplo-
graptus
folium.

folium, Hisinger, 1837, (Prionotus folium.) Leth. Suec., p. 113, Hud. Riv. Gr.

hudsonicus, Nicholson, 1875, Pal. Proc. Ont., p. 38, Hud. Riv. Gr. hypniformis, White, 1874, (Graptolithus hypniformis.) Rep. Invert. Foss., p. 12, and Geo. Sur. W. 100th Mer., vol. 4, p. 63, Trenton Gr.

inutilis, Hall, 1865, Can. Org. Rem., Decade 2, p. 111, Quebec Gr.

laciniatus, Emmons, 1856, Am. Geo., p. 236, Up. Taconic.

marcidus, Hall, 1859, (Graptolithus marcidus.) Pal. N. Y., vol. 3, p. 514, Hud. Riv. Gr.

mucronatus, Hall, 1847, (Graptolithus mucronatus.) Pal. N. Y., vol. 1, p. 263, Hud. Riv. Gr.

obliquus, Emmons, 1856, Am. Geo., p. 106, Up. Taconic.

peosta, Hall, 1861, (Graptolithus peosta.) Geo. Rep. Wis., p. 17, Trenton Gr.

pristiniformis, Hall, 1858, (Graptolithus pristiniformis.) Geo. Sur. Can., p. 133, and Can. Org. Rem., Decade 2, p. 110, Quebec Gr.

pristis, (?) Hisinger, 1837, (Prionotus pristis.) Leth. Suec., p. 114, and Pal. N. Y., vol. 1, p. 265, Hud. Riv. Gr.

putillus, Hall, 1865, Can. Org. Rem., Decade 2, p. 44, Hud. Riv. Gr.

rugosus, Emmons, 1856, Am. Geo., p. 105, Up. Taconic.

rectangularis, McCoy, 1851, Brit. Pal. Rocks, p. 3, Low Sil.

secalinus, Hall, 1847, (Fucoides secalinus.) Pal. N. Y., vol. 1, p. 267, syn. for D. simplex.

simplex, Emmons, 1844, (Fucoides simplex.) Taconic system, p. 27, and Pal. N. Y., vol. 1, p. 267, Up. Taconic.

spinulosus, Hall, 1859, (Graptolithus spinulosus.) Pal. N. Y., vol. 3, p. 517, Hud. Riv. Gr.

whitfieldi, Hall, 1859, (Graptolithus whitfieldi.) Pal. N. Y., vol. 3, p. 516, Hud. Riv. Gr.

Diphyphyllum, Hall, 1852, Pal. N. Y., vol. 2, p. 115, syn. for Diphyphyllum.

caespitosum, see Diphyphyllum caespitosum.

coralliferum, see Diphyphyllum coralliferum.

DIPLOTRYPA, Nicholson, 1879, Pal. Tab. Corals, p. 292. [Ety. *diploos*, double; *trypa*, hole.] Corallites of two kinds, the larger thin walled, polygonal, tabulae remote; often aggregated in clusters, (monticules); smaller corallites angular, thin walled, and completely isolating the larger ones; tabulae numerous. Type D. petropolitana.

infida, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 88, Trenton Gr.

milleri, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 245, Niagara Gr.

regularis, Foord, 1883, Micropaleontology, p. 13, Trenton Gr.

DISCOPHYLLUM, Hall, 1847, Pal. N. Y., vol. 1, p. 277. [Ety. *diskos*, disk; *phyllon*, leaf.] Discoid flattened, rays numerous, proceeding from the center and terminating in a thickened border. Type D. peltatum.

peltatum, Hall, 1847, Pal. N. Y., vol. 1, p. 277, Up. Taconic.

DUNCANELLA, Nicholson, 1874, Ann. Mag. Nat. Hist., 4th ser., vol. 13, p. 333. [Ety.

proper name.] Corallum simple, obconical; calyces deep, circular; rays strong, exsert; epitheca striated vertically; closely allied to Streptelasma. Type D. borealis.

borealis, Nicholson, 1874, Ann. Mag. Nat. Hist., 4th ser., vol. 13, p. 333, Niagara Gr.

ELASMOPHYLLUM, Hall, 1882, Foss. Corals Niagara and Up. Held Grs., p. 38. [Ety.

elasma, lamellae; *phyllon*, leaf.] Simple, turbinate, lamellae extending to the center, twisted or not; interlamellar cysts continuing to the center; no tabulae. Type E. attenuatum.

attenuatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 38, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 442, Up. Held. Gr.

EMMONSIA, Edwards & Haime, 1851, Monographie des Polyp., Foss. des Terr. Palaeoz., p. 246. [Ety. proper name.] Distinguished from Favosites by the compound character of the diaphragms, but generally regarded as a synonym. Type E. hemispherica.

hemispherica, Troost, 1840, (Calamopora hemispherica.) 5th Rep. Geo. Tenn., p. 72, Up. Held. Gr.

hemispherica, Yandell & Shumard, 1847, (Favosites hemisphericus.) Contrib. to Geo. of Ky., p. 7. Same species described by Troost.

Eridiphyllum, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxxi, syn. for Diphyphyllum.

rugosum, see Diphyphyllum rugosum.

simcoense, see Diphyphyllum simcoense.

strictum, see Diphyphyllum strictum.

vennori, see Diphyphyllum vennori.

verneuilanum, see Diphyphyllum verneuilanum.

Favastrea, DeBlainville, 1830, Man. d. Actinol., p. 374. Not an American palaeozoic genus.

striata, D'Orbigny, 1850, Prodr. d. Paléont., t. 1, p. 48. Not defined so as to be recognized.

Faviphyllum, as used by Hall, 1852, Stans. Exped. to Great Salt Lake, p. 407. Not defined, and founded upon a silicified, indeterminate fragment.

FIG. 172.
Duncanella
borealis
magnified.

FAVISTELLA, Hall, 1847, Pal. N. Y., vol. 1, p. 275. [Ety. *favus*, honey-comb; *stellata*, star.] Massive, hemispherical, corallites polygonal, increasing by lateral development; walls not separable as in Favosites, nor perforated by pores; tabulæ close; septa of alternately larger and smaller size, the larger reaching the center; twelve or more in each corallite. Type *F. stellata*.
calicina, Nicholson, 1874, Rep. Brit. Ass'n. and Pal. Tab. Corals, p. 197, Hud. Riv. Gr.
favosidea, Hall, 1852, Pal. N. Y., vol. 2, p. 41, Clinton Gr.
franklini, Salter, 1852, Sutherland's Jour., vol. 2, p. cccxxi, Up. Sil.
reticulata, Salter, 1852, Sutherland's Jour., vol. 2, p. cccxxi, Up. Sil.

FIG. 173.—Favistella stellata.

stellata, Hall, 1847, Pal. N. Y., vol. 1, p. 275, Hud. Riv. Gr.
FAVOSITES, Lamarck, 1812, Cours. de Zool. du Mus. d'Hist. Nat. and Hist. des Animaux Vert., vol. 2, p. 204. [Ety. *favus*, honey-comb.] Massive or branched, composed of numerous more or less polygonal corallites; tabulæ present; septa absent or rudimentary; walls perforated by one or more rows of mural pores, connecting the corallites. Type *F. alveolatus*.
alpenensis, Winchell, 1866, Rep. Low. Penin. Mich., p. 88, Ham. Gr.
aeolaria, DeBlainville. Not American.
arbuscula, Hall, 1876, Illust. Devon. Foss., pl. 38, Ham. Gr.
argus, Hall, 1876, Illust. Dev. Foss., pl. 13, Ham. Gr.
asper, D'Orbigny, 1849, Prodr. de Paléont., t. 1, p. 49, Clinton Gr.
basalticus, Goldfuss, 1826, Germ. Petref., p. 78. (Calamopora basaltica.) Devonian.
billingsi, Rominger, 1876, Foss. Corals, p. 29, Ham. Gr.

FIG. 174.—Favosites canadensis.

canadensis, Billings, 1858, (Fistulipora canadensis.) Can. Nat. and Geol., vol. 4, p. 98, Up. Held. Gr.
capax, Billings, 1866, Catal. Sil. Foss. Antic., p. 6, Hud. Riv. Gr.
cervicornis, DeBlainville, 1830, (Alveolites cervicornis.) Dict., vol. 60, p. 369, Devonian.
chapmani, Nicholson, 1874, Pal. Prov. Ont., p. 52, Up. Held. Gr.

clausus, Rominger, 1876, Foss. Corals, p. 37, Up. Held. and Ham. Gr.
conicus, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 112, and Pal. N. Y., vol. 6, p. 9, Low. Held. Gr.
constrictus, Hall, 1852, (Astrocerium constrictum.) Pal. N. Y., vol. 2, p. 123, Niagara Gr.
cristatus, Edwards & Haime, 1851, Pol. Foss. Terr. Palæoz., p. 242, Niagara Gr.
cumberlandicus, Troost, 1840, (Calamopora cumberlandica.) 5th Geo. Rep. Tenn., p. 70, Kaskaskia Gr.
digitatus, Rominger, 1876, Foss. Corals, p. 39, Ham. Gr.
divergens, Winchell, 1862, Proc. Acad. Nat. Sci., p. 112, and Geo. Sur. W. 100th Merid., vol. 4, p. 79, Subcarb.
dubius, L. DeBlainville, 1830, (Alveolites dubius.) Dict., vol. 60, p. 370, Corniferous Gr.
dumosus, Winchell, 1866, Rep. Low. Penin. Mich., p. 89, Ham. Gr.
emmonsii, Rominger, 1876, Foss. Corals, p. 27, Up. Held. Gr. Syn. (?) for *F. heliophoriformis*.
emmonsii, Hall, 1876, Illust. Dev. Foss., pl. 9. The name was preoccupied.
epidermatus, Rominger, 1862, Am. Jour. Sci. and Arts, vol. 34, p. 396, Corniferous Gr.
epidermatus var. *biloculi*, Hall, 1876, Illust. Dev. Foss., pl. 7, Up. Held. Gr.
epidermatus var. *corticatus*, Hall, 1876, Illust. Dev. Foss., pl. 10, Up. Held. Gr.
excretus, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., syn. for *F. spinigerus*.
explanatus, Hall, 1876, Illust. Dev. Foss., pl. 14, Ham. Gr.
favosus, Goldfuss, 1826, Germ. Petref., p. 77, and Pal. N. Y., vol. 2, p. 126, (Calamopora favosa.) Niagara Gr.
flabelliformis, Troost, 1843. Not satisfactorily defined.
forbesi, Edwards & Haime, 1854, Brit. Foss. Corals, p. 258, Niagara Gr.

FIG. 176.—Favosites goldfussi.

ensis, Nicholson, 1870, syn. for *F. forbesi* var. *occidentalis*.

FIG. 175.—Favosites forbesi var. occidentalis.

forbesi var. *discoideus*, Roemer, 1860, (Calamopora forbesi var. *discoidea*.) Sil. Fauna W. Tenn., p. 19, Niagara Gr.
forbesi var. *occidentalis*, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 109, Niagara Gr.

forbesi var. *waldroni*, Rominger, 1870, syn. for *F. forbesi* var. *occidentalis*.

goldfussi
goldfussi
ont.
preocc
gothland
sans V
Ham.

FIG. 177.—Favosites gothlandicus.

hemisph
hemisph
72, Up
hemisph
hemisph
Illust.
hemisph
Illust.
hisingeri
Foss.
agara C
hispidus
23, Nia
infundib
iac & V
intertext
p. 38, I
invagin
vol. 2,
limitaris
36, Cor
lycoperd
mamilla
nized.
maneus,
Sci., p.
maximus
ima.) 5
vonian
minimus
Mus. N
niagarens
p. 125,
niagarens
nitellus,
Mich.,
obliquus
24, Nia
occidens
Sur. W
p. 313,
parasitic
siticum
agara G
by Phil

goldfussi, Castelnau, 1843, (Calamopora goldfussi,) Syst. Sil., p. 47, Up. Sil. goldfussi, D'Orbigny, 1850, Prodr. de Paléont., p. 107, Devonian. The name was preoccupied.
gothlandicus, Lamarck, 1816, Hist. An. sans Vert., vol. 2, p. 206, Up. Held. and Ham. Grs.

FIG. 177.—Favosites gothlandicus.

hamiltonensis, Rominger, 1876, Foss. Corals, syn. for F. dumosus.
hamiltoniae, Hall, 1876, Illust. Dev. Foss., pl. 34, Ham. Gr.
helderbergiae, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 111, and Pal. N. Y., vol. 6, p. 8, Low. Held. Gr.
heliolitiformis, Rominger, 1862, (Calamopora heliolitiformis,) Am. Jour. Sci., vol. 34, 2d series, p. 397, Devonian.

hemisphericus, Troost, 1840, (Calamopora hemispherica,) 5th Geo. Rep. Tenn., p. 72, Up. Held. Gr. Same as Emmonsia hemispherica.

hemisphericus var. distortus, Hall, 1876, Illust. Dev. Foss., pl. 5, Up. Held. Gr.

hemisphericus var. rectus, Hall, 1876, Illust. Dev. Foss., pl. 2C, Up. Held. Gr.
hisingeri, Edwards & Haime, 1851, Pol. Foss. des Terr. Paléoz., p. 240, Niagara Gr.

hispidus, Rominger, 1876, Foss. Corals, p. 23, Niagara Gr.

infundibuliformis, as identified by D'Archiac & Verneuil. Not American.

intertextus, Rominger, 1876, Foss. Corals, p. 38, Ham. Gr.

invaginatus, Nicholson, 1875, Ohio Pal., vol. 2, p. 232, Corniferous Gr.

limitaris, Rominger, 1876, Foss. Corals, p. 36, Corniferous Gr.

lycoperdon, see Monticulipora lycoperdon.
mamillaris, Castelnau, 1843. Not recognized.

manus, Winchell, 1865, Proc. Acad. Nat. Sci., p. 112, Kinderhook Gr.

maximus, Troost, 1840, (Calamopora maxima,) 5th Rep. Geo. Tenn., p. 73, Devonian.

minimus, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 113, Low. Held. Gr.

niagarensis, Hall, 1852, Pal. N. Y., vol. 2, p. 125, Niagara Gr.

niagarensis var. spinigerus, see F. spinigerus.

nitellus, Winchell, 1866, Rep. Low. Pen. Mich., p. 89, Ham. Gr.

obliquus, Rominger, 1876, Foss. Corals, p. 24, Niagara Gr.

occidens, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 78, and Geo. Wis., vol. 4, p. 313, Niagara Gr.

parasiticus, Hall, 1852, (Astrocerium parasiticum,) Pal. N. Y., vol. 2, p. 122, Niagara Gr. This name was preoccupied by Phillips in his Geol. of Yorkshire.

placenta, Rominger, 1876, Foss. Corals, p. 34, Ham. Gr.

pleurodictyoides, Nicholson, 1875, Ohio Pal., vol. 2, p. 231, Corniferous Gr.

polymorphus, Goldfuss, 1826, Germ. Petref., p. 79, Corniferous Gr.

præficus, Billings, 1865, Can. Nat. and Geol., 2d ser., vol. 2, p. 429, Hud. Riv. Gr.

proximus, Hall, 1883, Rep. St. Geol., pl. 7, fig. 13-15, and Pal. N. Y., vol. 6, p. 10, Low. Held. Gr.

pyriformis, Hall, 1852, (Astrocerium pyriforme,) Pal. N. Y., vol. 2, p. 123, Niagara Gr.

radiatus, Rominger, 1876, Foss. Corals, p. 33, Ham. Gr.

radiciformis, Rominger, 1876, Foss. Corals, p. 34, Devonian.

reticulatus, DeBlainville, 1840, (Alveolites reticulatus,) Dict., vol. 60, p. 369, Niagara Gr.

sphericus, Hall, 1874, (Chetetes sphericus,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 111, and Pal. N. Y., vol. 6, p. 9, Low. Held. Gr.

spinigerus, Hall, 1876, (F. niagarensis var. spinigerus,) 28th Rep. N. Y. St. Mus. Nat. Hist., p. 108, Niagara Gr.

spongilla, Rominger, 1876, Foss. Corals, p. 24, syn. for F. spinigerus.

striatus, Say, 1818, Am. Jour. Sci., vol. 1, p. 384, Niagara Gr.

troosti, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 238, Devonian.

tuberosus, Rominger, 1876, Foss. Corals, p. 31, Corniferous Gr.

turbinatus, Billings, 1859, Can. Jour., vol. 4, p. 109, Up. Held. & Ham. Gr.

venustus, Hall, 1852, (Astrocerium venustum,) Pal. N. Y., vol. 2, p. 120, Niagara Gr.

verneuilii, Castelnau, 1843, syn. for Monticulipora fibrosa.

whitfieldi, White, 1874, Rep. Invert. Foss., syn. for F. divergens.

winchelli, Rominger, 1862, (Calamopora winchelli,) Am. Jour. Sci., vol. 34, 2d ser., p. 397, Devonian.

Favositopora, Kent, 1870, Ann. and Mag. Nat. Hist., 3d ser., vol. 6, p. 384.

palæozoica, Kent, 1870, Ann. and Mag. Nat. Hist., 3d ser., vol. 6, p. 384. Not recognized.

Filicites gracilis, see Plumalina gracilis.

Geoporites americanus, D'Orbigny, 1850. Not defined so as to be recognized.

Glossograptus, Emmons, (Glossograptus,) 1856, Am. Geo., p. 108. [Ety. glosse, tongue; grapho, I write.] Stipe free; thin, membranaceous, ligulate, extremities rounded, axis distinct. Type G. ciliatus.

ciliatus, Emmons, 1856, Am. Geo., pt. 2, p. 108, Up. Taconic.

setaceus, Emmons, 1856, Am. Geo., pt. 2, p. 236, Up. Taconic.

GRAPTOLITHUS, Linnæus, 1736, Syst. Nat., 1st Ed., but it was not until 1767, in the 12th Ed., that any species were defined. [Ety. *grapho*, I write; *lithos*, stone.] Stipes elongated, slender, flattened, or quadrangular; they may be simple or bifurcating; the cells enter the central canal and open their mouths upward, so as to form denticles on the margins when compressed. Type *G. scalaris*.

abnormis, Hall, 1858, Geo. Sur. Can., p. 117, and Can. Org. Rem., Decade 2, p. 106, Quebec Gr.

alatus, Hall, 1858, Geo. Sur. Can., p. 127, and Can. Org. Rem., Decade 2, p. 93, Quebec Gr.

amplexicaulis, see *Diplograptus amplexicaulis*.

angustifolius, see *Diplograptus angustifolius*.

annectans, Walcott, 1879, Utica Slate and related formations, p. 20, Utica Slate.

antennarius, see *Climacograptus antennarius*.

approximatus, Nicholson, 1873, (*Tetragraptus approximatus*), Ann. and Mag. Nat. Hist., 4th ser., vol. 12, Quebec Gr.

arcuatus, Hall, 1865, Can. Org. Rem., Decade 2, p. 79, Quebec Gr.

bicornis, see *Climacograptus bicornis*.

bifidus, Hall, 1858, Can. Nat. and Geo., vol. 3, p. 73, Quebec Gr.

bigsbyi, Hall, 1865, Can. Org. Rem., Decade 2, p. 86, Quebec Gr.

bryonoides, Hall, 1858, Geo. Sur. Can., p. 126, and Can. Org. Rem., Decade 2, p. 84, Quebec Gr.

caduceus, see *Didymograptus caduceus*.

clintonensis, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 74, and Pal. N. Y., vol. 2, p. 39, Clinton Gr.

constrictus, Hall, 1865, Can. Org. Rem., Decade 2, p. 76, Quebec Gr.

crucifer, Hall, 1858, Geo. Sur. Can., p. 125, and Can. Org. Rem., Decade 2, p. 92, Quebec Gr.

dentatus, Emmons, 1842, Geo. Rep. N. Y., p. 279, Utica Slate.

denticulatus, Hall, 1858, Geo. Sur. Can., p. 132, and Can. Org. Rem., Decade 2, p. 88, Quebec Gr.

divaricatus, Hall, 1859, Pal. N. Y., vol. 3, p. 513, Hud. Riv. Gr. See *Dicranograptus divaricatus*.

divergens, Hall, 1859, Pal. N. Y., vol. 3, p. 509, Hud. Riv. Gr.

ensiformis, see *Ratiolites ensiformis*.

extensus, Hall, 1858, Geo. Sur. Can., p. 132, and Can. Org. Rem., Decade 2, p. 80, Quebec Gr.

extenuatus, Hall, 1865, Can. Org. Rem., Decade 2, p. 75, Quebec Gr.

flaccidus, Hall, 1865, Can. Org. Rem., Decade 2, p. 143, Utica Slate.

flexilis, see *Clonograptus flexilis*.

foliaceus, see *Diplograptus foliaceus*.

folium, see *Diplograptus folium*.

fruticosus, Hall, 1858, Geo. Sur. Can., p. 128 and Can. Org. Rem., Decade 2, p. 100, Quebec Gr.

furcatus, see *Dicranograptus furcatus*.

gracilis, Hall, 1847, Pal. N. Y., vol. 1, p. 274, Utica Slate.

hallanus, see *Dendrograptus hallanus*.

headi, Hall, 1858, Geo. Sur. Can., p. 127, and Can. Org. Rem., Decade 2, p. 94, Quebec Gr.

hypniformis, see *Diplograptus hypniformis*.

indentus, Hall, 1858, Geo. Sur. Can., p. 128, and Can. Org. Rem., Decade 2, p. 74, Quebec Gr.

lævis, Hall, 1847, Pal. N. Y., vol. 1, p. 274, Utica Slate.

logani, 1858, Geo. Sur. Can., p. 115, and Can. Org. Rem., Decade 2, p. 100, Quebec Gr.

marcidus, see *Diplograptus marcidus*.

milesi, Hall, 1861, Geo. Sur. Vermont, vol. 1, p. 372, Quebec Gr.

mucronatus, see *Diplograptus mucronatus*.

multifasciatus, Hall, 1859, Pal. N. Y., vol. 3, p. 508, and Can. Org. Rem., Decade 2, p. 10, Hud. Riv. Gr.

nitidus, Hall, 1858, Geo. Sur. Can., p. 129, and Can. Org. Rem., Decade 2, p. 69, Quebec Gr.

octobrachiatus, Hall, 1858, Geo. Sur. Can., p. 122, and Can. Org. Rem., Decade 2, p. 96, Quebec Gr.

oceanarius, Hall, 1858, Geo. Sur. Can., p. 124, and Can. Org. Rem., Decade 2, p. 95, Quebec Gr.

patulus, Hall, 1858, Geo. Sur. Can., p. 131, and Can. Org. Rem., Decade 2, p. 71, Quebec Gr.

pennatulus, Hall, 1865, Can. Org. Rem., Decade 2, p. 82, Quebec Gr.

peosta, see *Diplograptus peosta*.

pristis, see *Diplograptus pristis*.

putillus, see *Diplograptus putillus*.

pristiniformis, see *Diplograptus pristini-formis*.

quadrirachiatus, Hall, 1858, Geo. Sur. Can., p. 125, and Can. Org. Rem., Decade 2, p. 91, Quebec Gr.

quadrimumeronatus, Hall, 1865, Can. Org. Rem., Decade 2, p. 144, Utica Slate.

ramosus, see *Dicranograptus ramosus*.

ramulus, Hall, 1865, Can. Org. Rem., Decade 2, p. 108, Quebec Gr.

ramulus, White. The name was preoccupied. See *G. whittianus*.

richardsoni, Hall, 1865, Can. Org. Rem., Decade 2, p. 107, Quebec Gr.

rigidus, see *Clonograptus rigidus*.

scalaris, Linnæus, as identified by Hall in Pal. N. Y., vol. 1, p. 271, Utica Slate.

FIG. 178.—*Graptolithus logani*.

secalinus,
serratus,
p. 274,
sagittari-
identifi-
1, p. 2
sextans,
similis,
ade 2,
spinulosus,
subtenuis,
244, H
tentaculat-
latus.
tenius, F
T
Portloe
renosus, s
whitfieldi,
whittianus
Foss.,
instead
which
Sur. W
HADROPHY-
Brit. I
hadros,

FIG. 179.
Hadrophylum gians.

and Co-
lington
orbigny,
Foss. C
Haimnophyl-
vol. 4,
ites.
ordinatum,
HALLIA, E
d. Pol.
[Ety. p
turbinat-
ter; one
place o
neighbo
as to as
no colum
divergens,
agara a
35th Re
412, Nia
divisa, H
and Up.
N. Y. S
agara G
insignis,
d. Pol.
Held. G
pluma, H
and Up.
N. Y. S
agara G

secalinus, see *Diplograptus secalinus*.
serratus, Hall, 1847, Pal. N. Y., vol. 1, p. 274, Utica Slate.

sagittarius, Linnaeus, 1767, Syst. Nat., as identified by Hall in Pal. N. Y., vol. 1, p. 272, Utica Slate.

sextans, see *Dicranograptus sextans*.

similis, Hall, 1865, Can. Org. Rem., Decade 2, p. 78, Quebec Gr.

spinulosus, see *Diplograptus spinulosus*.
subtenuis, Hall, 1877, Am. Pal. Foss., p. 244, Hud. Riv. Gr.

tentaculatus, see *Retiograptus tentaculatus*.

tenuis, Hall, 1847, Pal. N. Y., vol. 1, p. 272. The name was preoccupied by Portlock in 1843. See *G. subtenuis*.

venosus, see *Retiolites venosus*.

whitfieldi, see *Diplograptus whitfieldi*.

whitianus, S. A. Miller, 1883, Am. Pal. Foss., p. 269, Hud. Riv. Gr. Proposed instead of *G. ramulus*, White, 1874, which was preoccupied. See Geo. Sur. W. 100th Mer., vol. 4, p. 62.

HADROPHYLLUM, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxxvii. [Ety. *hadros*, mighty; *phyllon*, leaf.] Coral-

FIG. 179.
Hadrophylum glans.

lim short; calice superficial; one very large septal fossula and three small ones representing a cross; radiate arrangement of the septa somewhat irregular. Type *H. orbigny*.

glans, White, 1862, (Zaphrentis *glans*.) Proc. Bost. Soc. Nat. Hist., vol. 9, p. 32.

and Cont. to Pal., No. 8, p. 156, Burlington Gr.

orbigny, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxxvii, Up. Held Gr.

Haimophyllum, Billings, 1859, Can. Jour., vol. 4, p. 139, syn. for *Chonostegites*.

ordinatum, see *Chonostegites ordinatum*.

HALLIA, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 353.

[Ety. proper name.] Corallum tall, turbinate; septa extending to the center; one large septum occupying the place of the septal fossula, and the neighboring septa directed toward it, so as to assume a pinnate arrangement; no columella. Type *H. insignis*.

divergens, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 8, and 35th Rep. N. Y. Mus. Nat. Hist., p. 412, Niagara Gr.

divisa, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 8, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 412, Niagara Gr.

insignis, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 353, Up. Held. Gr.

pluma, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 8, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 412, Niagara Gr.

scitula, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 7, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 411, Niagara Gr.

scitula, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 7, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 411, Niagara Gr.

HALYSITES, Fischer, 1813, Zoognosia, vol. 1, p. 387. [Ety. *halyson*, a small chain.] Corallites long, arranged in single series, united laterally in the form of elliptical expansions, presenting a chain-like arrangement; epitheca thick; septa usually absent or rudimentary, but, in perfect specimens, extending to the center of the visceral chamber; tabulae horizontal. Type *H. catenulatus*.

agglomeratus, Hall, 1843, (Catenipora *agglomerata*.) Geo. Rep. 4th Dist. N. Y., Tab. Foss. No. 22, fig. 2, and Pal. N. Y., vol. 2, p. 129, Niagara Gr.

catenulatus, Linnaeus, 1767, (Tubipora *catenulata*.) Syst. Nat., 12th Ed., p. 1270, Niagara Gr.

catenulatus var. *fieldeni*, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 582, Up. Sil.

catenulatus var. *harti*, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 583, Up. Sil.

catenulatus var. *microporus*, Whitfield, 1882, Geo. Wis., vol. 4, p. 272, Niagara Gr.

compactus, Rominger, 1876, Foss. Corals, syn. for *H. agglomeratus*.

escharoides, Lamarck, 1816, (Catenipora *escharoides*.) Hist. des Anim. sans Vert., vol. 2, p. 207, Niagara Gr.

gracilis, Hall, 1851, (Catenipora *gracilis*.) Geo. Lake Sup. Land Dist., vol. 2, p. 212, Hud. Riv. Gr.

labyrinthicus, Goldfuss, 1826, (Catenipora *labyrinthica*.) Petref. Germ., p. 71, Niagara Gr.

meandrina, Troost, 1840, (Catenipora *meandrina*.) 5th Geo. Rep. Tenn., Niagara Gr. The definition is too meagre for identification.

parryi, König, 1824, (Catenipora *parryi*.) Supp. to App. of Capt. Parry's Voyage for the Discovery of a North-west Passage, Up. Sil.

sexto-attenuatus, Owen, 1862, Geo. Sur. Ind., p. 362, Niagara Gr.

Harmodites rugosus, D'Orbigny, 1850, Prodr. de Paléont., t. 1, p. 50. Not defined so as to be recognized.

HELIOLITES, Guettard, 1770, Mem. 3, p. 454. [Ety. *helios*, sun; *lithos*, stone.] Corallum spheroidal, hemispherical or ramose; corallites of larger and smaller size, the

FIG. 180.—Halysites catenulatus.

Porpora
affinis
(Kier.)
1903.

larger ones cylindrical, with twelve infoldings of the wall or septa, not reaching the center, the smaller ones polygonal, investing the larger ones; walls amalgamated; tabulae numerous; no columella. Type *H. interstinctus*.

affinis, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 427, Hud. Riv. and Mid. Sil.

elegans, Hall, 1852, Pal. N. Y., vol. 2, p. 130, Niagara Gr.

exiguus, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 428, Mid. Sil.

FIG. 181.—*Heliolites interstinctus*.

interstinctus, Linnaeus, 1767, (*Madrepora interstincta*,) Syst. Nat., 12th Ed., p. 1276, Niagara Gr.

macrostylus, Hall, 1852, Pal. N. Y., vol. 2, p. 135, Niagara Gr.

megastoma, McCoy, 1846, Sil. Foss. of Ireland, p. 62, Niagara Gr.

pyriformis, Guettard, 1770, Mem. 3, p. 454, and Pal. N. Y., vol. 2, p. 133, Niagara Gr.

sparvus, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 428, Mid. Sil.

speciosus, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 426, Mid. Sil.

spiniporus, Hall, 1852, Pal. N. Y., vol. 2, p. 131, Niagara Gr.

subtubulatus, McCoy, as identified by Rominger, 1876, Foss. Corals, p. 13, Niagara Gr.

tenuis, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 428, Mid. Sil.

HELIOPHYLLUM, Hall, 1848, in Dana. Zooph., p. 356. [Ety. *helios*, sun; *phyllon*, leaf.]

Corallum simple; septa well developed and producing lateral lamellar prolongations, which extend from the wall toward the center of the visceral chamber, so as to represent ascending arches and to constitute irregular central tabulae, and which are united toward the circumference by means of vertical dissepiments. Type *H. halli*.

acuminatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 46, and 12th Rep. Ind. Geo., p. 310, Up. Held. Gr.

aequale, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 47, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 451, Up. Held. Gr.

aequum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 51, and 12th Rep. Ind. Geo., p. 314, Up. Held. Gr.

alternatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 45, and 12th Rep. Ind. Geo., p. 306, Up. Held. Gr.

annulatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 48, and 12th Rep. Ind. Geo., p. 307, Up. Held. Gr.

arachne, Hall, 1876, Illust. Dev. Foss., pl. 24, Ham. Gr.

campaniforme, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 53, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 457, Up. Held. Gr.

canadense, Billings, 1859, Can. Jour., vol. 4, p. 125, Up. Held. Gr.

cancellatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 53, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 457, Up. Held. Gr.

cayugaense, Billings, 1859, Can. Jour., vol. 4, p. 124, Up. Held. Gr.

colborneense, Nicholson, 1875, Can. Nat. and Geo., vol. 7, p. 143, Up. Held. Gr.

colligatum, Billings, 1859, Can. Jour., vol. 4, p. 126, Up. Held. Gr.

compactum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 48, and 12th Rep. Ind. Geo., p. 308, Up. Held. Gr.

confluens, Hall, 1876, Illust. Dev. Foss., pl. 26 and 27, Ham. Gr.

degener, Hall, 1876, Illust. Dev. Foss., pl. 25, Ham. Gr.

dentatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 48, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 452, Up. Held. Gr.

denticulatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 52, and 12th Rep. Ind. Geo., p. 313, Up. Held. Gr.

dentilineatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 13, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 417, Niagara Gr.

distans, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 50, and 12th Rep. Ind. Geo., p. 308, Up. Held. Gr.

eriense, Billings, 1859, Can. Jour., vol. 4, p. 124, Corniferous Gr.

exiguum, Billings, 1860, Can. Jour., vol. 5, p. 261, Corniferous Gr.

fasciculatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 48, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 452, Up. Held. Gr.

secundum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 49, and 12th Rep. Geo. Ind., p. 309, Up. Held. Gr.

fissuratum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 53, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 457, Up. Held. Gr.

gemmatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 49, and 12th Rep. Geo. Ind., p. 310, Up. Held. Gr.

gemmiferum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 13, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 417, Niagara Gr.

Porpora
speciosa (Kier.)
1903.

Rev. des
mittelsil.

Heliolites,

Christiania,

Videnskabs

Selskabet

Skripten

Math. natur

klasse

No. 10.

halli, Foss.
Corals

FIG. 182.—

Mus.
incrassat

agara
12th R

infundib

Ind., p

invagina

agara
12th R

irregular

pl. 24,

latericre

Niagar
12th R

lineolatu

35th R

454, U

mitella

Up. He

St. Mus

netteleroth

agara a

Rep. G

pocillatu

agara a

35th R

454, U

pravum,

and Up

Geo. In

proliferu

Ont. Cæ

halli, Edwards & Haime, 1850, Brit. Foss. Corals, p. 235, Ham. Gr.

FIG. 182.—*Heliophyllum halli*.

halli var. obconicum, Hall, 1876, Illust. Dev. Foss., pl. 25, Ham. Gr.

halli var. reflexum, Hall, 1876, Illust. Dev. Foss., pl. 23, Ham. Gr.

imbricatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 46, and 35th Rep. N. Y. St.

Mus. Nat. Hist., p. 450, Up. Held. Gr. incrassatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 46, and 12th Rep. Geo. Ind., p. 309, Up. Held. Gr. infundibulum, Hall, 1883, 12th Rep. Geo. Ind., p. 305, Up. Held. Gr. invaginatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 47, and 12th Rep. Geo. Ind., p. 306, Up. Held. Gr. irregulare, Hall, 1876, Illust. Dev. Foss., pl. 24, Ham. Gr.

laterescens, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 49, and 12th Rep. Geo. Ind., p. 314, Up. Held. Gr. lineolatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 50, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 454, Up. Held. Gr.

mitella, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 14, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 418, Niagara Gr. netterothi, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 51, and 12th Rep. Geo. Ind., p. 312, Up. Held. Gr. pocillatum, Hall, 1884, Foss. Corals Niagara and Up. Held. Grs., p. 50, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 454, Up. Held. Gr.

pravum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 13, and 12th Rep. Geo. Ind., p. 274, Niagara Gr.

proliferum, Nicholson, 1874, Rep. Pal. Ont. Can., p. 27, Up. Held. Gr.

proliferum, Hall, 1876, Illust. Dev. Foss., pl. 28, is probably a syn. for *H. proliferum*, Nicholson.

puteatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 14, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 418, Niagara Gr.

scyphulus, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 51, and 12th Rep. Geo. Ind., p. 306, Up. Held. Gr. sordidum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 52, and 12th Rep. Geo. Ind., p. 311, Up. Held. Gr.

subcapitosum, see *Crepidophyllum subcapitosum*.

tenuimurale, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 51, and 12th Rep. Geo. Ind., p. 307, Up. Held. Gr.

tenuiseptatum, Billings, 1850, Can. Jour., vol. 4, p. 126, Ham. Gr.

venatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 46, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 450, Up. Held. Gr.

verticale, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 47, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 451, Up. Held. Gr.

HETEROPHRENTIS, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 235. [Ety. *heteros*, irregular; *phren*, midriff or lamella.] Corallum simple, turbinate, calice large, septal fossette well-defined, bottom smooth or with a pseudo-columnella, septa, below the calice, sharp edged; often with their inner edges twisted together, usually rounded on approaching the margin; apparently only a single transverse diaphragm, which forms the floor of the cup. Type *H. spatiosa*. *compta*, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 236, Corniferous Gr. *excellens*, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 236, Corniferous Gr. *prolifera*, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 236, Corniferous Gr. *spatiosa*, Billings, 1858, (*Zaphrentis spatiosa*), Can. Nat. and Geo., vol. 3, p. 430, Onondaga and Corniferous Gr.

Heterotrypa, Nicholson, 1879, Pal. Tab. Cor., p. 291. Proposed as a subgenus of *Monticulipora*, making *M. mammulata* the type which is the type of *Monticulipora*. This is a violation of the elementary principles of nomenclature.

Houghtonia, syn. for *Calapæcia*.

huronica, see *Calapæcia huronica*.

INOCAULIS, Hall, 1852, Pal. N. Y., vol. 2, p. 176. [Ety. *inos*, small sprouts; *kaulos*, stem.] Expanded, bifurcating, fenestrate, and usually indicated by simple black rays connected by small cross bars. Type *I. plumulosus*.

FIG. 183.—*Inocaulis plumulosus*.

anastomica, Ringueberg, 1888, Proc. Acad. Nat. Sci. Phil., p. 131, Niagara Gr.

- arbuscula, Ulrich, 1870, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 28, Hud. Riv. Gr. bellus, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 122, Niagara Gr. cervicornis, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 37, Niagara Gr. diffusus, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 36, Niagara Gr. divaricatus, Hall, 1879, Desc. New sp. Foss., p. 2, and 11th Rep. Geo. Ind., p. 225, Niagara Gr. phycoides, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 38, Niagara Gr. plumulosus, Hall, 1851, Pal. N. Y., vol. 2, p. 176, Niagara Gr. problematis, Spencer, 1878, Can. Nat., vol. 8, and Bull. No. 1, Mus. Univ. St. Mo., p. 36, Niagara Gr. ramulosus, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 38, Niagara Gr. walkeri, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 35, Niagara Gr. *Lamellopora*, Owen, 1840, Rep. or Minn. Lands, p. 70. Very poorly defined, but a syn. for *Stromatopora*. *infundibularia*, Owen, 1840, Rep. on Minn. Lands, p. 70. A species of *Stromatopora*, poorly defined. *LEPTOPORA*, Winchell, 1863, Proc. Acad. Nat. Sci. Phil., p. 2. [Ety. *leptos*, shallow; *poros*, cell.] Discoidal, cells shallow; walls vertically striated; interior vesicular; cups elevated in the center, and displaying radial septa. Type *H. typus*. *typus*, Winchell, 1863, Proc. Acad. Nat. Sci. Phil., p. 3, Marshall Gr. *winchelli*, White, 1879, Bull. U. S. Sur., vol. 5, p. 211, and Cont. to Pal. No. 6, p. 121, Carboniferous. *Limaria*, Steininger, 1834, Bull. Soc. Geo. France, vol. 1, p. 339. The name was preoccupied by Link in 1807, and by Rafinesque in 1815. See *Cœnites*.

crassa, see *Cœnites crassus*.

falcata, see *Cœnites falcatus*.

fruticosa, see *Cœnites fruticosus*.

laminata, see *Cœnites laminatus*.

ramulosa, see *Cœnites ramulosus*.

LINDSTROMIA, Nicholson & Thompson, 1877, Proc. Roy. Soc. Edinb., vol. 9, p. 149. [Ety. proper name.] Type *L. columnaris*. *columnaris*, Nicholson & Thompson, 1877, Proc. Roy. Soc. Edinb., vol. 9, p. 149, Devonian.

Linipora rotunda, Troost, not defined.

LITHOSTROTION, Lhwyd, 1869, Lithophyl. Britann. Ichnographia, Epistola 5, tab. xxiii.

[Ety. *lithos*, stone; *strotion*, little rafter.] *Cœlum* composite, astreiform; corallites loosely approximated, and circu-

FIG. 184.—*Lithostrotion basaltiforme*.

lar or intimately united, and polygonal; septa numerous; calices unequal; structure as in *Clisiophyllum*. Type *L. basaltiforme*.

californiense, Meek, 1864, Pal. Californin., vol. 1, p. 6, Carb.

canadense, Castelnau, 1843, (*Axi-nura canadensis*), Syst. Sil., p. 49, St. Louis Gr.

harmodites, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 440, Carboniferous.

juncum, Fleming, 1828, (*Caryophylla juncum*), Brit. Anim., p. 509, Subcarb.

mammillare, Castelnau, 1843, (*As-trea mammillaris*), Syst. Sil., p. 50, syn. for *L. canadense*.

microstylum, White, 1880, 12th Rep. U. S. Geo. Sur. Terr., p. 158, Kinderhook or Waverly Gr.

pictoense, Billings, 1868, Acad. Geo., p. 285, Carb.

proliferum, Hall, 1858, Geo. Rep. Iowa, p. 668, St. Louis Gr.

stokesi, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 440, Carboniferous.

whitneyi, Meek, 1875, Wheeler's Sur. W. 100 Mer., vol. 4, p. 103 Coal Meas.

LONSDALIA, McCoy, 1849, Ann. & Mag. Nat. Hist. 2d ser., vol. 3, p. 10. [Ety. proper name.] Corallum aggregate; corallites circular, not laterally united; septa and tabulae numerous; visceral chamber separated into two zones, the outer one composed of curved vesicular plates extending upward and outward; walls rugose and striated; reproduction by circular germs arising from the outer zone. Type *L. duplicata*.

papillata, Fischer, 1837, (*Cyathophyllum papillatum*), Oryct. de Moscon., p. 155, Carboniferous. American. (?)

LOPHOPHYLLUM, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxxvi. [Ety. *lophos*, ridge; *phyllon*, leaf.] Corallum resembling *Zaphrentis*, excepting that a cres-

cent

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

of the

and b

with

prima

Type

calceola,

calceo

expansu

1876, I

Sci. P

Cont.

p. 157,

centif

occup

of t

is in

one o

a su

placed

centiform columella occupies the center of the callee, and is in continuity by one of its ends with a small septum placed in the middle of the septal fossula, and by the other end with the opposite primary septum. Type *L. konineki*.

calceola, see *Zaphrentis calceola*.

expansum, White, 1876, Proc. Acad. Nat. Sci. Phil., p. 27, and Cont. to. Pal., No. 6, p. 157, Keokuk Gr.

proliferum,

McChes-

ney, 1860,

(*Cyathax-*

onia pro-

liferum),

New Pal.

Foss., p.

75, and Pal. E. Neb., p. 144,

Coal Mens.

FIG. 187.—*L. LUNATIPORA*, Winchell, 1860, phophyllum Rep. Low. Penin. Mich. proliferum. p. 89. [Ety. *lunatus*, cres-

cent-formed; *poros*, pore.] Massive or with corallites consolidated; corallites long, curving outward from an imaginary axis; walls double; tabulae present; no mural pores. Type *L. michiganensis*.

michiganensis, Winchell, 1860, Rep. Low. Penin. Mich., p. 89, Ham. Gr.

LYELLIA, Edwards & Haime, 1851, Mon. Pol. Foss. Terr. Pal., p. 226 [Ety. proper name.] Corallum massive; corallites cylindrical; walls thick, costulated, free toward their terminations, and united by vesicular coenenchyma; septa 12; tabulae irregular. Type *L. americana*.

FIG. 188.—*Lyellia americana*.

americana, Edwards & Haime, 1851, Mon.

Pol. Foss. Terr. Pal., p. 226, Up. Heid. Gr.

decipiens, Rominger, 1876, Foss. Corals,

p. 17, Niagara Gr.

glabra, Owen, 1840, (*Sarcinula glabra*),

Rep. on Minn. Lands, p. 70, Niagara Gr.

FIG. 189.—*Megalograptus welchii*. Cylindrical part of the body depressed, showing cells.

papillata, Rominger, 1876, Foss. Corals,

p. 16, Ni-

agara Gr.

parvutuba,

Rominger,

1876, Pal.

Foss. Cor-

als, p. 17,

Niagara

Gr.

Madrepora

repens,

Troost.

Not satis-

factorily

defined.

MEGALOGRA-

TUS, S. A.

Miller,

1874, Cin.

Quar. Jour.

Sci., vol. 1,

p. 343.

[Ety. *me-*

gale, large;

grapho, I

write.]

Very large

cylindri-

cal, bear-

ing fronds

with spi-

nous pro-

cesses, and

covered

with cellu-

lar open-

ings. Type

M. welchii.

welchii, S.

A. Miller, 1874, Cin. Quar. Jour. Sci.,

vol. 1, p. 343, Hud. Riv. Gr.

FIG. 190.—*Megalograptus welchii*. Frond, showing cells and spinous processes.

MICHELINIA, DeKoninck, 1842, Descries Anim. Foss. Belg., p. 29. [Ety. proper name.] Corallum composite, forming hemispherical, depressed, or pyriform masses of prismatic or subcylindrical corallites; mural pores; tabulae; tubes having striae or ridges; epitheca concentrically wrinkled, with root-like prolongations. Type *M. favosa*.

FIG. 191.—*Megalograptus weichti*. Frond, showing cells and spinous processes.

convexa, D'Orbigny, 1850, Prodr. de Paléont., t. 1, p. 107, Onondaga and Corniferous Grs.

FIG. 192.—*Michelinita eugeneae*.

dividua, Hall, 1876, Illust. Dev. Foss., pl. 18, Ham. Gr.

eugeneae, White, 1884, 13th, Rep. Geo. Ind., p. 119, Coal Meas.

expansa, White, 1880, 12th Rep. U. S. Geo. Sur. Terr., p. 158, Waverly Gr.

favositoides, Billings, 1858, Rep. of Progr. Can. Geo. Sur., p. 175, Up. Held. Gr.

insignis, Rominger, 1876, Foss. Corals, p. 75, Up. Held. and Ham. Gr.

intermittens, Billings, 1859, Can. Nat. and Geo. Sur., vol. 4, p. 113, Corniferous Gr.

lenticularis, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 113, Low. Held. Gr.

placenta, White, 1880, 12th Rep. U. S. Geo. Sur. Terr., p. 157, Waverly Gr.

stylopore, Eaton, 1832, (*Astrea stylopore*.) Geo. Text book, p. 40, and Illust. Dev. Foss., pl. 18, Ham. Gr.

trochiscus, Rominger, 1876, Pal. Foss., p. 76, syn. for *Plenrodictyum americanum*.

MICROCYCLUS, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 420. [Ety. *mikros*, small; *kuklos*, circle.] Corallum free or with a minute central point of attachment, discoidal; no columella; calice shallow, with a single fossula; septa short, radiating regularly, or those nearest the fosse converging a little

FIG. 193.—*Michelinita eugeneae*. Another specimen, with larger corallites.

toward its sides; epitheca well developed. Type *M. discus*.

discus, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 420, Ham. Gr.

Millepora repens, see *Alveolites repens*.

MONOGRAPTUS, Emmons, 1856, (Monograptus.)

Am. Geo., p. 106. [Ety. *monos*, one; *grapho*, I write.] Serrations confined to one edge of the stipe; axis none. Type *M. elegans*.

convolutus var. *coppingeri*, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 577, Silurian.

elegans, Emmons, 1856, Am. Geo., p. 106, Up. Taconic.

rectus, Emmons, 1856, Am. Geo., p. 107, Up. Taconic.

MONOTRYPA, Nicholson, 1879, Pal. Tab. Corals, p. 320. [Ety. *monos*, one; *trypa*, hole.] Corallites of two kinds; the larger aggregated into clusters (monticules); the smaller occupying the space between the monticules; both larger and smaller thin-walled, polygonal tabulae remote. Type *M. undulata*. This was proposed as a subgenus for Monticulipora; upon microscopical examination and upon such a state of facts, I prefer, at present, to leave the species under the genus Monticulipora.

(?) spinulosa, Hall, 1887, Pal. N. Y., vol. 1, p. 67, Low. Held. Gr.

MONOTRYPELLA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 153. [Ety. *monotrypa*; and *illus*, diminutive.] Ramose, smooth or tuberculated, cells of one kind only; walls thin, in the axial region, and thicker toward the periphery; diaphragms straight; no spiniform tubuli. Type *M. aequalis*.

abrupta, Hall, 1879, (*Chetetes abruptus*.) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 148, Low. Held. Gr.

aequalis, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 247, Hud. Riv. Gr.

arbuscula, Hall, 1887, Pal. N. Y., vol. 6, p. 12, Low. Held. Gr.

briareus, Nicholson, 1875, (*Chetetes briareus*.) Ohio Pal., vol. 2, p. 202, Utica Slate.

consimilis, Hall, 1876, (*Chetetes consimilis*.) 28th Rep. N. Y. St. Mus. Nat. Hist., p. 110, Niagara Gr.

densa, Hall, 1874, (*Trematopora densa*.) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 105, Low. Held. Gr.

FIG. 195.—*Monotrypella quadrala*, natural size and magnified.

atopora densa, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 105, Low. Held. Gr.

FIG. 194.—*Micrograptus elegans*.

quadra-
ratus
3, an-
the
Riv.
subgen-
Nat.
Monitric-

Palaeo-
hillo-
form
two
ous;
the
often
nume-
mura-
Linda-
genus
son, R.
it wi-
belon-
adheren-

heren-
427, C.
andrew-
Affin.

Ulrich
barrand-
randi,
Pal. o-
bulbosa-
bosa.)
2, p. 4
calceolu-
Soc.

Riv. C.
cincinna-
cincin-
p. 2, s.
Monti-
compress-
pressa
5, p. 2
consimil-
Nat. 1
Gr.

dalii, Ed-
dalii,
Hud. 1
lopore-
dawsoni,
Affin.

decipien-
cipiens
3, Hud-

Fig. 196.—*Monotrypella*

dychiei, J.
Hist., v

- quadrata*, Rominger, 1886, (*Chetetes quadratus*.) Proc. Acad. Nat. Sci. Phil., p. 3, and Ohio Pal., vol. 2, p. 201, under the name of *Chetetes rhombicus*, Hud. Riv. Gr.
- subquadrata*, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 249, Hud. Riv. Gr.
- MONTICULIPORA*, D'Orbigny, 1850, Prodr. de Paléont., t. 1, p. 25. [Ety. *monticulus*, hillock; *poros*, pore.] Corallum of every form and shape; corallites usually of two kinds, one minute; tabulae numerous; walls separable, thickened toward the mouths of the tubes; corallites often aggregated, upon the surface, in numerous monticules; no septa; no mural pores; increase by gemmation. Lindstrom, Ulrich, and others, class this genus with the Bryozoa, while Nicholson, Edwards & Haime, and others, class it with the Polypt, where it seems to belong. Type *M. mammulata*.
- adherens*, Billings, 1859, (*Stenopora adherens*.) Can. Nat. and Geo., vol. 4, p. 427, Chazy Gr.
- andrewsi*, Nicholson, 1881, Struct. and Affin. of Montic., p. 128, Hud. Riv. Gr. Ulrich refers it to *Callopora*.
- barrandi*, Nicholson, 1874, (*Chetetes barrandi*.) Quar. Jour. Geo. Soc., vol. 30, and Pal. of Ontario, p. 60, Ham. Gr.
- bulbosa*, Billings, 1865, (*Stenopora bulbosa*.) Can. Nat. and Geo., 2d ser., vol. 2, p. 429, Mid. Sil.
- calceolus*, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 26, Hud. Riv. Gr.
- cincinnatiensis*, James, 1875, (*Chetetes cincinnatiensis*.) Int. Catal. Cin. Foss., p. 2, and Nicholson, Struct. and Affin. Montic., p. 226, Hud. Riv. Gr.
- compressa*, Ulrich, 1882, (*Peronopora compressa*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 244, Hud. Riv. Gr.
- consimilis*, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 238, Hud. Riv. Gr.
- dalii*, Edwards & Haime, 1851, (*Chetetes dalii*.) Pol. Foss. d. Terr. Pal., p. 266, Hud. Riv. Gr. Ulrich refers it to *Callopora*.
- dawsoni*, Nicholson, 1881, Struct. and Affin. Montic., p. 141, Hud. Riv. Gr.
- decepiens*, Rominger, 1866, (*Chetetes decepiens*.) Proc. Acad. Nat. Sci. Phil., p. 3, Hud. Riv. Gr.

FIG. 196.—*Monticulipora delicatula*.

dychei, James, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 235, Hud. Riv. Gr.

delicatula, Nicholson, 1874, (*Chetetes delicatulus*.) Quar. Jour. Geo. Soc., vol. 30, and Ohio Pal., vol. 2, p. 199, Hud. Riv. Gr. Probably a Bryozoum and not a *Monticulipora*.

- fibrosa*, Goldfuss, 1826, (*Calamapora fibrosa*.) Germ. Petref., p. 82, Hud. Riv. and Clinton Grs.
- filiosa*, D'Orbigny, 1850, Prodr. d. Pal., t. 1, p. 25, and Edwards & Haime, Pol. Foss. d. Terr. Pal., p. 266, Hud. Riv. Gr.
- frondosa*, D'Orbigny, 1850, Prodr. d. Pal., t. 1, p. 25, and Ohio Pal., vol. 2, p. 208, Hud. Riv. Gr.
- gracilis*, see *Batostomella gracilis*.
- grandis*, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 78, Trenton Gr.
- implicata*, see *Batostoma implicata*.
- irregularis*, Ulrich, 1879, (*Chetetes irregularis*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 129, Hud. Riv. Gr.
- lavis*, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 236, Hud. Riv. Gr.

FIG. 197.—*Monticulipora lycoperdon*.

- lycoperdon*, Say, 1847, (*Favosites lycoperdon*.) Hall, Pal. N. Y., vol. 1, p. 64, Trenton Gr.
- mammulata*, D'Orbigny, 1850, Prodr. de Paléont., t. 1, p. 25, and Ohio Pal., vol. 2, p. 207, Hud. Riv. Gr.
- molesta*, Nicholson, 1881, Struct. and Affin. of Montic., p. 224, Hud. Riv. Gr.
- moniliformis*, Nicholson, 1874, (*Chetetes moniliformis*.) Geo. Mag., vol. 1, p. 57, and Pal. of Ont., p. 60, Ham. Gr.
- monticula*, White, 1876, Proc. Acad. Nat. Sci. Phil., p. 27, Devonian.
- multituberculata*, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 71, and Geo. Wis., vol. 4, p. 250, Hud. Riv. Gr.
- parasitica*, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 238, Hud. Riv. Gr.
- patulus*, Billings, 1859, (*Stenopora patula*.) Can. Nat. and Geo., vol. 4, p. 427, Chazy Gr.
- pavonia*, see *Ptilodictya pavonia*.
- petasiformis*, Nicholson, 1881, Struct. and Affin. of Montic., p. 190, Hud. Riv. Gr.
- punctata*, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 71, and Geo. Wis., vol. 4, p. 249, Hud. Riv. Gr.

FIG. 198.—*Monticulipora ramosa*, natural size and magnified.

ramosa, D'Orbigny, 1850, Prodr. d. Pal., t. 1, p. 25, and Edwards & Haime, Pol. Foss. de Terr. Pal., p. 266, and Ohio Pal.,

- vol. 2, under the name of *Chetetes delii*, Hud. Riv. Gr. Ulrich refers it to *Calopora*.
- rectangularis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 70, and Geo. Wis., vol. 4, p. 249, Hud. Riv. Gr.
- rugosa, Hall, 1847, (*Chetetes rugosus*), Pal. N. Y., vol. 1, p. 67, Trenton Gr.
- rugosa, Edwards & Haime, 1851, Pol. Foss. de Terr. Pal., is merely a variety or form of *M. ramosa*, and associated with it in the Hud. Riv. Gr.
- selwyni, see *Prasopora selwyni*.
- selwyni var. *hospitatis*, see *Prasopora selwyni* var. *hospitatis*.
- solitaria, Ulrich, 1883, (*Heterotrypa solitaria*), Jour. Cin. Soc. Nat. Hist., vol. 6, p. 88, Hud. Riv. Gr.
- subglobosa, Ulrich, 1879, (*Chetetes subglobosus*), Jour. Cin. Soc. Nat. Hist., vol. 2, p. 129, Hud. Riv. Gr.
- subpulchella, Nicholson, 1875, (*Chetetes subpulchellus*), Ohio Pal., vol. 2, p. 196, Hud. Riv. Gr.
- trentonensis, Nicholson, 1881, Struct. and Affin. Montic., p. 149, Trenton Gr.
- tuberculata, see *Spatiopora tuberculata*.
- ulrichi, Nicholson, 1881, Struct. and Affin. Montic., p. 131, Hud. Riv. Gr.
- undulata, Nicholson, 1875, (*Chetetes undulatus*), Pal. of Ont., p. 10, and Struct. and Affin. Montic., p. 170, Trenton and Hud. Riv. Gr.
- uniformis, Ulrich, 1882, (*Peronopora uniformis*), Jour. Cin. Soc. Nat. Hist., vol. 5, p. 244, Hud. Riv. Gr.
- vaupelli, Ulrich, 1883, (*Heterotrypa vaupelli*), Jour. Cin. Soc. Nat. Hist., vol. 6, p. 85, Hud. Riv. Gr.
- venusta, Ulrich, 1878, (*Chetetes venustus*), Jour. Cin. Soc. Nat. Hist., vol. 1, p. 93, Utica Slate.
- westoni, Foord, 1883, Cont. to Micro-Pal., p. 7, Trenton Gr.
- wetherbyi, Ulrich, 1882, Jour. Cin. Soc. Nat. History, vol. 5, p. 239, Trenton Gr.
- whiteavesi, Nicholson, 1879, Pal. Tab. Corals, p. 316, and Struct. and Affin. of Montic., p. 160, Trenton Gr.
- NEBULIPORA, McCoy, 1850, Ann. and Mag. Nat. Hist., 2d ser., vol. 6, p. 284. [Ety. *nebula*, thick mist; *poros*, pore.] Incrusting or forming lenticular masses, with a concentrically wrinkled epitheca below, composed of small prismatic corallites perpendicular to the upper surface, with clusters of rather larger size, all in contact; tabulae at regular distances; no septa. Type *N. explanata*.
- papillata, McCoy, 1850, Ann. and Mag. Nat. Hist., vol. 6, p. 284, Hudson Riv. Gr.
- NEMAORAPTUS, Emmons, (*Nemagrapsus*), 1856, Am. Geo., pt. 2, p. 109. The termination *grapsus* is preferred because *grapsus* is used in the nomenclature of crustacea. [Ety. *nema*, thread; *grapho*, I write.] Axis elongated and thread-like, simple or com-
- pound branches, round at the base, and flattened at the extremities; cells arranged on the flattened part of the axis instead of the margin. Type *N. elegans*.
- capillaris, Emmons, 1856, Am. Geo., pt. 2, p. 109, Up. Taconic.
- elegans, Emmons, 1856, Am. Geo., pt. 2, p. 109, Up. Taconic.
- NYCTOPORA, Nicholson, 1879, Pal. Tab. Corals, p. 182. [Ety. *nyktos*, night; *poros*, pore.] Corallum composite, massive; corallites polygonal, in contact; walls thin, amalgamated; mural pores numerous, small; septa, in the form of marginal vertical ridges; 10 to 15 in each corallite; tabulae numerous, complete, horizontal. Type *N. billingsi*.
- billingsi, Nicholson, 1879, Pal. Tab. Corals, p. 184, Trenton Gr.
- OLDHAMIA, Forbes, 1850, Dub. Geo. Jour. [Ety. proper name.] Strong stems, with branches arranged in whorls; substance corneous; cellules undetermined. Type *O. antiqua*.
- antiqua, Forbes, 1850, Dublin Geo. Jour., Potad. Gr.
- fruticosa, Hall, 1865, Can. Org. Rem. Decade 2, p. 50, Trenton Gr.
- OMPHYMA, Rafinesque, 1820, Ann. des Sci. Phys. d. Bruxelles, vol. 5, p. 234. [Ety. *omphax*, precious stone.] Simple, turbinate, wall with rudimentary epitheca, producing radicleform appendages; septa numerous, equally developed and divided into four groups by an equal number of shallow fossulae; tabulae smooth toward the center. Type *O. turbinata*.
- congregata, Billings, 1866, Catal. Sil. Foss. Antic., p. 93, Clinton and Niagara Gr.
- drummondii, Billings, 1866, Catal. Sil. Foss. Antic., p. 93, Clinton and Niagara Gr.
- stokesi, Edwards & Haime, 1851, (*Ptychophyllum stokesi*), Polyp. Foss. Pal., p. 407, and Geo. Wis., vol. 4, p. 279, Niagara Gr.
- verrucosa, Rafinesque & Clifford, 1820, Monog. d. Turbinolides in Ann. d. Phys. d. Brux., t. 5, p. 235, Niagara Gr.
- PACHYPHYLLUM, Edwards & Haime, 1850, Brit. Foss. Corals, p. lxviii. [Ety. *pachys*, thick; *phyllon*, leaf.] Corallum, composite, increasing by lateral gemination; corallites united by the development of the costae and exotheca; tabulae abundant. Type *P. bouchardi*.

FIG. 200. — Omphyma turbinata.

FIG. 199. — Oldhamia antiqua.

solitari
Rep.
Cher

FIG.
woodm
mani
Chen
PACHYPO
K. V
[Ety.
droid
polyg
thick
centr
annul
obso
tabul
few,
P. lar
fischeri
Can.
Held.

FIG. 202. — P. froud.

circula
cemen
kirbyi,
85, De
rotuloid
(Cyclo
loides,
vol. 2,
ton Gi
PALEOPHY
lings,
Prog.
palaos

solitary, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 232, Chemung Gr.

FIG. 201.—Pachyphyllum woodmani.

woodmani, White, 1870, (Smithia woodmani,) Geo. Sur. Iowa, vol. 1, p. 188, Chemung Gr.

PACHYFORA, Lindstrom, 1873, Ofversigt af K. Vetensk. Akad. Forhandl., p. 14. [Ety. *pachys*, thick; *poros*, pore.] Dendroid or frondescens; corallites polygonal or subcylindrical walls thickened toward their mouths, by concentric layers of sclerenchyma; calices annular, oblique, or semilunar; septa obsolete or mere spiniform projections; tabulæ complete, remote; mural pores few, irregular, and often large. Type *P. lamellicornis*.

fischeri, Billings, 1860, (Alveolites fischeri,) Can. Jour. n. s., vol. 5, p. 256, Up. Held. Gr.

FIG. 202.—Pachypora frondosa.

frondosa, Nicholson, 1874, (Alveolites frondosus,) Geo. Mag., vol. 1, p. 15, and Rep. Pal. Ontario, p. 57, Ham. Gr. ornata, Roringer, 1876, (Dendropora ornata,) Pal. Foss. Corals, p. 62, Ham. Gr.

PALÆOCYCLUS, Edwards & Haime, 1849, Comptus rendus, t. 29, p. 71. [Ety. *palaos*, ancient; *kuklos*, circle.] Corallum circular; fossula deep, broad, circular; septa thick, not numerous or cemented together. Type *P. porpita*.

kirbyi, Meek, 1868, Trans. Chi. Sci., p. 85, Devonian. rotuloides, Hall, 1852, (Cyclolites rotuloides,) Pal. N. Y., vol. 2, p. 42, Clinton Gr.

PALÆOPHYLLUM, Billings, 1858, Rep. of Prog. Geo. Sur. Can., p. 168. [Ety. *palaos*, ancient; *phyllon*, leaf.] Fascicu-

FIG. 203.—Palæocyclus rotuloides.

late or aggregate; corallites surrounded by a thick wall; septa extending the whole length; tabulæ absent or rudimentary; increase by lateral budding. Distinguished from Streptelasma by forming aggregate masses. Type *P. rugosum*.

FIG. 204.—Palæophyllum divaricans.

divaricans, Nicholson, 1875, Pal. Ohio, vol. 2, p. 220, Hud. Riv. Gr.

rugosum, Billings, 1858, Rep. of Prog. Can. Geo. Sur., p. 168, Trenton Gr.

Palaetochia, Emmons, 1856, Geo. Rep. Midland counties of North Carolina. Two species were mentioned, *P. major* and *P. minor*, both of which are supposed to be concretions, and therefore inorganic.

Peronopora, Nicholson, syn. for Monticulipora.

PETRAIA, Munster, 1839, Beitrage zur Petrefaktenkunde, vol. 1, p. 42. [Ety. *petraios*, that grows among rocks.]

Simple, turbinate; septa of one or two sizes, the larger extending from the walls to the center, where they are more or less twisted; no tabulæ or connecting vesicular plates. Type *P. decussata*. Streptelasma is by some regarded as a synonym, by others as a subgenus, and by others as quite distinct. The forms in this country which have been referred to Petraia are all, probably, Streptelasma, and for that reason I have so referred them.

angulata, see Streptelasma angulatum.

aperta, see Streptelasma apertum.

fanningana, see Streptelasma fanninganum.

forresteri, Honeyman, 1868, Acadian Geology, p. 594. A catalogue name.

latusecula, see Streptelasma latusecula.

logani, see Streptelasma logani.

minganensis, see Archeocyathus minganensis.

ottawensis, see Streptelasma ottawense.

pulchella, see Streptelasma pulchellum.

pygmaea, see Streptelasma pygmaea.

rustica, see Streptelasma rusticum.

selecta, see Streptelasma selectum.

wayneensis, see Streptelasma wayneuse.

PHILLIPASTREA, D'Orbigny, 1849, Note Sur. des Polypters Fossiles, p. 12. [Ety. proper name; *aster*, star.] Composite, resembling Strombodes, but differing in the septa of neighboring corallites

being confluent, and consequently the calices are not definitely circumscribed; no exterior walls; interior mural investment well characterized; center of tabulæ presenting a columellar tubercle. Type *P. hennahi*.

affinis, Billings, 1874, Pal. Foss., vol. 2, p. 11, Gaspé limestone No. 8, Devonian. *gigas*, Owen, 1840, (*Astræa gigas*.) Rep. on Mineral lands, p. 70, Devonian. *hennahi*, Lonsdale, 1840, (*Astræa hennahi*.) Geo. Trans., vol. 5, p. 697, Devonian.

johanni, Hall & Whitfield, 1873, (*Smithia johanni*.) 23d. Rep. N. Y. St. Mus. Nat. Hist., p. 234, Chemung Gr.

mammillaris, see *Strombodes mammillaris*. *multiradiata*, Hall & Whitfield, 1873, (*Smithia multiradiata*.) 23d. Rep. N. Y. St. Mus. Nat. Hist., p. 234, Chemung Gr.

FIG. 205.—*Phillipsastrea verneuili*.

verneuili, Edwards & Haime, 1851, Polyiers Foss. des Terr. Pal., p. 447, Ham. Gr.

verrilli, Meek, 1868, (*Smithia verrilli*.) Trans. Chi. Acad. Sci., p. 83, Devonian. *yandelli*, Rominger, 1876, Foss. Corals, p. 130, Up. Held. Gr. Not well defined.

PHYLLOGRAPTUS, Hall, 1858, Rep. of Progr. Can. Geo. Sur., p. 135. [Ety. *phylon*, leaf; *grapho*, I write.]

FIG. 206.—*Phyllograptus typus*.

Frond consisting of simple or compound foliiform stipes, which are celluliferous on the two opposite sides, the margins having a mucronate extension from each cellule; supported on a slender radicle, or combined in groups. Type *P. typus*.

angustifolius, Hall, 1858, Rep. of Progr. Can. Geo. Sur., p. 139, and Dec. 2. Org. Rem., p. 125, Quebec Gr.

anna, Hall, 1865, Can. Org. Rem., Decade 2, p. 124, Quebec Gr.

dubius, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 15, Niagara Gr.

illicifolius, Hall, 1858, Rep. of Progr. Can. Geo. Sur., p. 139, and Dec. 2, Org. Rem., p. 121, Quebec Gr.

loringi, White, 1874, Rep. Invertebrate Foss., p. 9, and Geo. Sur. W. 100th Mer., vol. 4, p. 51, Quebec Gr.

similis, Hall, 1853, Can. Nat. and Geo., vol. 4, syn. for *Graptolithus bigsbyi*.

typus, Hall, 1858, Rep. of Progr. Can. Geo. Sur., p. 137, and Dec. 2, Org. Rem., p. 118, Quebec Gr.

PLASMOPORA, Edwards & Haime, 1849, Comptes rend., t. 29, p. 262. [Ety. *plasma*, cast; *poros*, pore.] Free, sub-hemispheric; basal epitheca, con-

FIG. 207.—*Plasmopora foliis*. Natural size, and magnified.

foliis, Edwards & Haime, 1851, Mon. Pol. Foss. de Terr. Pal., p. 220, Niagara Gr.

PLEURODICTYUM, Goldfuss, 1826, Petref. Germ., vol. 1, p. 209. [Ety. *pleura*, side; *dictyon*, net. Corallum discoidal, upper surface convex; corallites diverging from the center of the base, polygonal or subcylindrical; walls thick; mural pores irregular; tabulæ not numerous, but sometimes inosculating; septa rudimentary, in the form of marginal ridges. There is usually a vermiform body at the central part of the base. Type *P. problematicum*.

americanum, Roemer, 1876, Lethæ Palæozoica, pl. 33, figs. 2a and 2b. Ham. Gr.

FIG. 208.—*Pleurodictyum problematicum*. Under side, showing serpulæ like body.

problematicum, Goldfuss, 1826, Petref. Germ., vol. 1, p. 113, Onondaga Gr. *Polydylasma*, Hall, 1852, Pal. N. Y., vol. 2, syn. for *Zaphrentis*.

turbinatum, see *Zaphrentis turbinata*.

Porites, Lamarck, 1816, Hist. des Anim. sans Vert., t. 2, p. 267. Not an American Palæozoic genus.

astræa
era
sa
Pa
pyrif
Ve
vetust
PRASOP
An
20,
por
cor
atir
lar
gle
cul
lar
ves
cen
an
ran
and
ple
affin
tol
con
Sur
cont
Sur
new
new
Huc
nodo
His
ocul
pal
selw
selw
p. 2
selw
(Mo
Stru
Huc
simul
Sur.
Prionotu
folium
pristia,
PROTAR
Foss

- astræiformis*, Owen, 1840, Rep. on Mineral lands, Devonian. This may be the same species subsequently described as *Pachyphyllum woodmani*.
pyriformis, as identified by d'Archiac & Verneuil, not American.
vetustus, see *Protarea vetusta*.
PRASOPORA, Nicholson & Etheridge, 1877, Ann. and Mag. Nat. Hist., 4th ser., vol. 20, p. 382 [Ety. *prason*, sea-plant; *poros*, pore.] Corallum compound, concavo-convex or hemispheric; corallites radiating from a wrinkled basal epitheca; larger and smaller corallites intermingled throughout the colony; no monticules; corallites thin-walled, prismatic; large ones with an exterior zone of vesicular tabulae surrounding a vacant central tube, which may be crossed by an occasional tabula; smaller ones arranged in a zone around the larger ones, and crossed by numerous, close-set, complete, horizontal tabulae. Type *P. grave*.
affinis, Foord, 1883, Cont. to Micropalaontology, p. 12, Trenton Gr.
conoidea, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 87, Trenton Gr.
contigua, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 87, Trenton Gr.
newberryi, Nicholson, 1875, (Chetetes newberryi,) Ohio Pal., vol. 2, p. 212, Hud. Riv. Gr.
nodosa, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 245, Hud. Riv. Gr.
oculata, Foord, 1883, Cont. to Micropalaontology, p. 11, Trenton Gr.
selwyni, Nicholson, 1881, (Monticulipora selwyni,) Struct. and Affin. of Montic., p. 206, Trenton Gr.
selwyni var. *hospitalis*, Nicholson, 1881, (Monticulipora selwyni var. *hospitalis*,) Struct. and Affin. of Montic., p. 206, Hud. Riv. Gr.
simulatrix, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 85, Trenton Gr.
Prionotus, Nilsson, 1835, Leth. Suec.
folium, see *Diplograptus folium*.
pristis, see *Diplograptus pristis*.
PROTAREA, Edwards & Haime, 1851, Pol. Foss. des Terr. Pal., p. 208. [Ety.

FIG. 208.—*Protarea vetusta*, on *Strophomena alternata*.

protos, first; *araios*, porous.] Thin, incrusting; calices equal, hexagonal,

shallow; septa 12, extending but slightly into the visceral chamber; walls thick. Type *P. vetusta*.

vetusta, Hall, 1847, (Porites vetustus,) Pal. N. Y., vol. 1, p. 71, Trenton & Hud. Riv. Grs.

verneuili, Edwards & Haime, 1851, Pol. Foss. des Terr. Pal., p. 209, Silurian. (?)

PROTOGRAPTUS, Matthew, 1885, Trans. Roy. Soc. Can., p. 31. [Ety. *protos*, first; *grapho*, I write.] Stipes thin, flat, elongate, dichotomously branched; having a central axis, and being alate on each side; pores arranged along the axis of the stipe; axis and margin of the stipe connected by delicate nervules. Type *P. alatus*.

alatus, Matthew, 1885, Trans. Roy. Soc. Can., p. 22, St. John Gr.

PTILOGRAPTUS, Hall, 1865, Can. Org. Rem., Decade 2, p. 139. (Ety. *ptilon*, feather; *grapho*, I write.) Plant-like, rooted, simple or branching; branches plumose, pinnules alternate on opposite sides; celluliferous on one face only; branches cylindrical or flattened. Type *P. plumosus*.

foliaceus, Spencer, 1878, Can. Nat., vol. 8, and Bull. No. 1, Mus. Univ. St. Mo., p. 41, Niagara Gr.

geinitzanus, Hall, 1865, Can. Org. Rem., Decade 2, p. 140, Quebec Gr.

plumosus, Hall, 1865, Can. Org. Rem., Decade 2, p. 140, Quebec Gr.

PRYCHONEMA, Hall, 1887, Pal. N. Y., vol. 6, p. xiv. [Ety. *ptyche*, wrinkle; *nema*, thread.] Massive or ramose, composed of thin-walled, strongly corrugated cells, which are apparently without diaphragms. Type *P. tabulatum*.

helderbergiae, Hall, 1874, (Chetetes helderbergiae,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 110, Low. Held. Gr.

tabulatum, Hall, 1876, (Chetetes tabulatum,) Illus. Dev. Foss., pl. 37, and figs. 16-19, and Pal. N. Y., vol. 6, p. 14, Up. Held. Gr.

PRYCOPHYLLUM, Lonsdale, 1839, Sil. Syst., p. 691, and E. & H. Brit. Foss. Corals, p. lxxix. [Ety. *ptyche*, ridge; *phyllon*, leaf.] Corallum simple, having infundibuliform tabulae superposed and invaginated; septa strongly twisted toward the center of the tabulae so as to constitute a spurious columella. Type *P. stokesi*.

canadense, Billings, 1862, Pal. Foss., vol. 1, p. 107, Mid. Sil. Anticosti Gr., Division 4.

floriforme, Hall, 1832, Foss. Corals Niagara and Up. Held. Grs., p. 5, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 409, Niagara Gr.

FIG. 210.—*Ptilograptus foliaceus*.

- bulbosus, Spencer, 1878, Can. Nat., vol. 8, p. 460, and Bull. No. 1, Mus. Univ. St. Mo., p. 30, Niagara Gr.
- ROMINOERIA, Nicholson, 1879, Tab. Corals, p. 114. [Ety. proper name.] Corallum lax, spreading; corallites cylindrical, annulated, multiplying by lateral gemmation, and typically producing new tubes, in umbellate whorls or verticils, at short intervals; where the walls are in contact with the visceral chambers they are connected by mural pores; tabulae complete, remote; septa represented by vertical rows of spinules. It resembles Aulopora, but is only attached basally, and is therefore free throughout the greater part of its extent. Type R. umbellifera.
- cornuta, Billings, 1859, (Aulopora cornuta,) Can. Jour., vol. 4, p. 119, Up. Held. and Ham. Grs.
- niagarensis, Rominger, 1876, (Quenstedtia niagarensis,) Foss. Corals, p. 72, Niagara Gr.
- umbellifera, Billings, 1859, (Aulopora umbellifera,) Can. Jour., vol. 4, p. 119, Up. Held. Gr.
- Sarcinula, Lamarck, 1816, Hist. des Animaux Vert., t. 2, p. 222. Not an American Palaeozoic genus.
- glabra, Owen, 1840, Rep. on Mineral Lands. See Lyellia glabra.
- obsoleta, Hall, 1857, Geo. Lake Sup. Land Dist., vol. 2, Hud. Riv. Gr. Not recognized.
- ramosa, Eaton, 1832, Geo. Text Book, p. 41. Not properly defined.
- Smithia, Edwards & Haime, 1851, Pol. Foss. des Terr. Pal. The name was preoccupied for a genus in botany, and is a syn. for Phillipsastrea.
- johanni, see Phillipsastrea johanni.
- multiradiata, see Phillipsastrea multiradiata.
- woodmani, see Pachyphyllum woodmani.
- verrilli, see Phillipsastrea verrilli.
- SPHÆROLITES, Hinde, 1875, Proc. Geo. Soc. Lond., vol. 31, p. 514. [Ety. from the spheroidal form.] Type S. nicholsoni.
- nicholsoni, Hinde, 1875, Proc. Geo. Soc. Lond., vol. 31, p. 514, Low. Held. Gr.

FIG. 216.—Staurograptus dichotomus.

dichotomus, Emmons, 1856, Am. Geo., p. 109, Up. Taconic.

- STELLIPORA, Hall, 1847, Pal. N. Y., vol. 1, p. 79. [Ety. stella, star; poros, pore.] Corallum dendroid or incrusting; corallites dimorphic; apertures subcircular; no septa; tabulae abundant; surface covered with conspicuous star-shaped elevations and depressions. Type S. antheloidea.
- antheloidea, Hall, 1847, Pal. N. Y., vol. 1, p. 79, Trenton and Hud. Riv. Grs.
- fischeri, Ulrich, 1883, (Constellaria fischeri,) Jour. Cin. Soc. Nat. Hist., vol. 6, p. 270, Hud. Riv. Gr.
- florida, Ulrich, 1882, (Constellaria florida,) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 257, Hud. Riv. Gr.
- limitaris, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 126, Hud. Riv. Gr. Syn. (?) for S. polystomella.

FIG. 217.—Stellipora polystomella, natural size and magnified star.

- polystomella, Nicholson, 1873, (Constellaria polystomella,) Ohio Pal., vol. 2, p. 215, Hud. Riv. Gr.
- STENOPORA, Lonsdale, 1844, App. to Darwin's Volcanic Islands, p. 161, and Geo. Russ. and Ural Mts., vol. 1, p. 631. [Ety. stenos, narrow; poros, pore.] Corallum very similar to Chetetes, but having small styliform processes at the angles of the calices, as understood by Edwards & Haime. Nicholson defines the genus, and restricts it to specimens from Australia and Van Diemen's Land, which, as in the type, have constricted corallites and minute mural pores. Type S. ovata.
- bulbosa, see Monticulipora bulbosa.
- adherens, see Monticulipora adherens.
- crassa, see Chetetes crassus.
- fibrosa, see Monticulipora fibrosa.
- exilis, Dawson, 1868, Acad. Geo., p. 287, Subcarboniferous.
- huronensis, see Tetradium huronense.
- libana, Safford, 1869, Geo. of Tenn. Not defined.
- patula, see Monticulipora patula.
- spinigera, see Chetetes spinigerus.

FIG. 218.—Stenopora exilis.

Strephodes, McCoy, 1849, syn. for *Cyathophyllum*.
austini, see *Clisiophyllum austini*.
pickthorni, see *Cyathophyllum pickthorni*.

STREPTELASMA, Hall, 1847, Pal. N. Y., vol. 1, p. 17. [Ety. *streptos*, twisted; *elasma*, lamella.] Turbinate, gradually or abruptly expanding; cup deep; lamellæ or septa longitudinal, spirally twisted toward the center; no tabulæ or fessette. Type *S. expansum*.

aequidistans, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 20, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 424, Up. Held. Gr.

ampliatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 19, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 423, Up. Held. Gr.

angulatum, Billings, 1862, (*Petraia angulata*.) Pal. Foss., vol. 1, p. 103, Hud. Riv. Gr.

apertum, Billings, 1862, (*Petraia aperta*.) Pal. Foss., vol. 1, p. 102, Black Riv. Gr.

calyculus, Hall, 1852, Pal. N. Y., vol. 2, p. 111, Niagara Gr.

coarctatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs.; p. 21, and 12th Rep. Geo. Sur. Ind., p. 275, Up. Held. Gr.

conspicuum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 19, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 423, Up. Held. Gr.

conulus, Rominger, 1876, Foss. Corals, p. 144, Niagara Gr.

corniculum, Hall, 1847, Pal. N. Y., vol. 1, p. 69, Trenton and Hud. Riv. Grs.

crassum, Hall, 1847, Pal. N. Y., vol. 1, p. 70, Trenton Gr.

crateriforme, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 20, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 424, Up. Held. Gr.

dissimile, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 17, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 421, Up. Held. Gr.

FIG. 219.—*Streptelasma inflatum*, transverse section.

expansum, Hall, 1847, Pal. N. Y., vol. 1, p. 17, Chazy Gr.

extans, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 5, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 409, Niagara Gr.

fanningana, Safford, 1869, (*Petraia fanningana*.) Geo. Tenn., p. 320, Low. Held. Gr.

fossula, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 19, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 423, Up. Held. Gr.

inflatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 18, and 12th Rep. Geo. Ind., p. 276, Up. Held. Gr.

involutum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 20, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 424, Up. Held. Gr.

lamellatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 17, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 421, Up. Held. Gr.

laterarium, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 18, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 422, Corniferous limestone.

latuscula, Billings, 1862, (*Petraia latuscula*.) Pal. Foss., vol. 1, p. 104, Mid. Sil. Anticosti, Div. 4.

limitare, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 5, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 409, Niagara Gr.

logani, Nicholson, 1875, (*Petraia logani*.) Can. Nat., vol. 7, p. 143, Up. Held. Gr.

mammiferum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 21, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 425, Up. Held. Gr.

minimum, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 106, syn. for *Duncanella borealis*.

multilamellosum, Hall, 1847, Pal. N. Y., vol. 1, p. 70, Trenton Gr.

ottawensis, Billings, 1865, (*Petraia ottawensis*.) Can. Nat. and Geo., 2d ser., vol. 2, Trenton Gr.

papillatum, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 21, and 12th Rep. Geo. Ind., p. 276, Up. Held. Gr.

parvulum, Hall, 1847, Pal. N. Y., vol. 1, p. 71, Trenton Gr.

patulum, Rominger, 1876, Foss. Corals, p. 143, Niagara Gr.

profundum, Conrad, 1843, Proc. Acad. Nat. Sci. Phil., p. 335, (*Cyathophyllum profundum*.) and Hall, 1847, Pal. N. Y., vol. 1, p. 49, Birdseye, Black Riv. and Trenton Grs.

pulchellum, Billings, 1865, (*Petraia pulchella*.) Can. Nat. and Geo., 2d ser., vol. 2, p. 424, Mid. Sil.

pygmaeum, Billings, 1862, (*Petraia pygmaea*.) Pal. Foss., vol. 1, p. 103, Mid. Sil. Anticosti, Div. 4.

radicans, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 106, Niagara Gr.

rectum,
Geo.

FIG. 220.—*Streptelasma rectum*.

1876,
strictum,
Mus. L.

tenuis,
and U.

Rep. C.

ungula,
19, H.

waynem
nensis
Held.

STRIATOPOR
p. 156

pore.]

angula

the su

like o

nally s

bands

mural

carbona

Nat. F

caverno

p. 60,

flexuosa

156, N

formosa

p. 254

hurone

p. 58,

iowensis

sis.) R

p. 69,

issa, H

Nat. F

limbata

Geo. T

Illust.

Ham.

linnean

Jour.

(Gr.

missouri

Worth

Ill., v

Held.

rugosa

Iowa,

iowen

STROMBO

Natur

rectum, Hall, 1843, (*Strombodes rectus*),
Geo. Rep. 4th Dist. N. Y., p. 200, and
Illust. Dev. Foss., pl. 19,
Ham. Gr.

FIG. 220. *Streptelasma rectum*.

rusticum, Billings, 1858,
(*Petraia rustica*), Rep.
of Progr. Geo. Sur. Can.,
p. 168, Hud. Riv. Gr.
selectum, Billings, 1865,
(*Petraia selecta*), Can.
Nat. and Geo., 2d ser.,
vol. 2, p. 429, Mid. Sil.
simplex, Hall, 1882, Foss.

Corals Niagara and Up.
Held. Grs., p. 18, and
12th Rep. Geo. Ind.,
p. 277, Up. Held. Gr.

spongiaxis, Rominger,

1876, Foss. Corals, p. 144, Niagara Gr.
strictum, Hall, 1874, 26th Rep. N. Y. St.
Mus. Nat. Hist., p. 114, Low. Held. Gr.
tenuis, Hall, 1882, Foss. Corals Niagara
and Up. Held. Grs., p. 17, and 12th
Rep. Geo. Ind., p. 278, Up. Held. Gr.
ungula, Hall, 1876, Illust. Dev. Foss., pl.
19, Ham. Gr.
waynensis, Safford, 1869, (*Petraia waynensis*),
Geo. of Tenn., p. 314, Low. Held. Gr.

STRIATOPORA, Hall, 1852, Pal. N. Y., vol. 2,
p. 156. [Ety. *striatus*, striated; *poros*,
pore.] Ramose; corallites thick-walled,
angular, conical; cells opening upon
the surface in expanded, angular, cup-
like depressions, which are longitudi-
nally striated, and between the striae the
bands may bear spinules; tabulae and
mural pores common. Type *S. flexuosa*.
carbonaria, White, 1862, Proc. Bost. Soc.
Nat. Hist., vol. 9, Burlington Gr.

cavernosa, Rominger, 1876, Foss. Corals,
p. 60, Corniferus Gr.

flexuosa, Hall, 1852, Pal. N. Y., vol. 2, p.
156, Niagara Gr.

formosa, Billings, 1860, Can. Jour., vol. 5,
p. 254, Up. Held. Gr.

huronensis, Rominger, 1876, Foss. Corals,
p. 58, Niagara Gr.

iowensis, Owen, 1840, (*Cyathopora iowensis*),
Rep. on Min. Lands of Iowa, etc.,
p. 69, Ham. Gr.

issa, Hall, 1874, 26th Rep. N. Y. St. Mus.
Nat. Hist., p. 114, Low. Held. Gr.

limbata, Eaton, 1832, (*Madrepora limbata*),
Geo. Text Book, p. 30, and
Illust. Dev. Foss., pl. 33,
Ham. Gr.

linneana, Billings, 1860, Can.
Jour., vol. 5, p. 253, Ham.
Gr.

missouriensis, Meek &
Worthen, 1863, Geo. Sur.
Ill., vol. 3, p. 369, Low.
Held. Gr.

rugosa, Hall, 1858, Geo. of
Iowa, p. 479, syn. for *S.*
iowensis.

STROMBODES, Schweigger, 1820, Handb. der
Naturg., p. 418. [Ety. *strombos*, twisting.]

FIG. 221.—*Strombodes linneana*.

Composite, increasing by calicular gem-
mation; corallites constituted prin-
cipally by a series of superposed, in-
vaginates, infundibuliform tabulae,
united by ascending trabeculae, so as to
form a columnar mass; calices pen-
tagonal, well circumscribed, and com-
pletely covered with the septal radii;
outer walls not well developed, and inner
mural investment rudimentary. Type
S. pentagonus.

alpenensis, Rominger, 1876, Foss. Corals,
p. 133, Ham. Gr. Is this a syn. for *S.*
mammillaris?

diffuens, Edwards & Haime, 1851, Pol.
Fos. des Terr. Pal., p. 431, Auticosti Gr.
distortus, Hall, 1843, Geo. Rep. 4th Dist.
N. Y., p. 209, Ham. Gr. Too imper-
fectly described for recognition. Prob-
ably a *Heliophyllum*.

eximius, Billings, 1866, Catal. Foss.
Antic., p. 93, Clinton and Niagara Grs.

gracilis, Billings, 1862, Pal. Foss., vol. 1,
p. 113, Mid. Sil.

heliophyllum, (?) *Heliophyllum halli*.

mammillaris, Owen, 1840, (*Astrea mam-
miliaris*), Rep. on Min. Lands, p. 70, and
Rominger, in Pal. Foss., p. 133, Ni-
agara Gr.

FIG. 222.—*Strombodes pentagonus*.

pentagonus, Goldfuss, 1826, Germ. Petref.
p. 62, Niagara Gr.

pygmaeus, Rominger, 1876, Foss. Corals,
p. 132, Niagara Gr.

(?) *rectus*, see *Streptelasma rectum*.

separatus, Ulrich, 1886, Cont. to Am. Pal.,
p. 32, Niagara Gr.

simplex, see *Zaphrentis simplex*.

striatus, D'Orbigny, 1850, Prodr. de
Paléont., p. 48 Niagara Gr.

STYLASTREA, Lonsdale, 1845, Geo. and Pal.
of Russia, and Ural Mts., p. 621. [Ety.
stylōs, pillar; *aster*, star.] Composite,
large; corallites prismatic; easily sepa-
rable; walls thick, striated longitudi-
nally, and wrinkled transversely; within
the walls there is a narrow, vesicular,
perithecal zone, and within it a lamelli-
ferous area; septa numerous, not reach-
ing the center; tabulae abundant. Type
S. inconferta.

anna, Whitfield, 1882, Ann. N. Y. Acad.
Sci., vol. 2, p. 199, Up. Held. Gr.

SYRINGOLITES, Hinde, 1879, Geo. Mag., vol.
6, p. 244. [Ety. *syrix*, pipe; *lithos*,
stone.] Composite, large with basal
epithea; corallites polygonal, thin-

- walled, with mural pores, and a cylindrical tube in the center of each corallite. Type *S. huronensis*.
- huronensis*, Hinde, 1879, Geo. Mag., vol. 6, p. 246, Niagara Gr.
- SYRINGOPORA*, Goldfuss, 1826, Germ. Petref., p. 75. [Ety. *syrix*, pipe; *poros*, pore.] Corallum aggregating, at first creeping after the manner of *Aulopora*, then sending up numerous vertical, cylindrical corallites, usually flexuous, subparallel, and connected laterally by more or less transverse processes; septa rudimentary; tabule close set, infundibuliform; epitheca well developed. Type *S. reticulata*.
- alectiformis*, Winchell, 1866, Rep. Low. Penin. Mich., p. 90, Ham. Gr.
- annulata*, Rominger, 1876, Foss. Corals, p. 81, Niagara Gr.
- aulopora*, Salter, 1855, Belcher's Last of the Arctic Voyages, vol. 2, p. 385, Carboniferous.
- cleviana*, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 295, Corniferous Gr.
- compacta*, Billings, 1858, Can. Nat. and Geo., vol. 3, p. 422, Up. Sil.
- crassata*, Winchell, 1866, Rep. Low. Penin. Mich., p. 90, Ham. Gr.
- dalmani*, Billings, 1858, Can. Nat. and Geo., vol. 3, p. 423, Up. Sil.
- debilis*, Billings, 1858, Can. Nat. and Geo., vol. 3, p. 423, Up. Sil.
- eiegans*, Billings, 1858, Can. Nat. and Geo., vol. 3, p. 425, Corniferous Gr.
- fenesrata*, Winchell, 1866, Rep. Low. Penin. Mich., p. 90, Ham. Gr.
- fibrata*, Rominger, 1876, Foss. Corals, p. 82, Niagara Gr.
- harveyi*, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 32, Waverly or Kinderhook Gr.
- hisingeri*, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 116, Corniferous Gr.
- infundibulum*, see *Cystostylus infundibulum*.
- intermedia*, Nicholson, 1874, Rep. Pal. Prov. Ont. Can., p. 126, Ham. Gr.
- laxata*, Billings, 1859, Can. Jour., vol. 4, p. 118, Corniferous Gr.
- maclurii*, Billings, 1860, Can. Jour., vol. 5, p. 258, Corniferous Gr.
- multattenuata*, McChesney, 1860, New Pal. Foss., p. 75, and Pal. E. Neb., p. 144, Coal Meas.
- multicaulis*, Hall, 1852, Pal. N. Y., vol. 2, p. 119, Niagara Gr.
- nobilis*, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 118, Up. Held. Gr.
- parallela*, Etheridge, 1873, Quar. Jour. Geo. Soc., vol. 34, p. 583, Up. Sil.
- perelegans*, Billings, 1859, Can. Jour., vol. 4, p. 117, Up. Held. Gr.
- retiformis*, Billings, 1858, Can. Nat. and Geo., vol. 3, p. 424, Up. Sil.

FIG. 223.—*Syringopora maclurii*.

- reticulata*, Goldfuss, 1826, Petref. Germ., p. 76, Devonian.
- tabulata*, Edwards & Haime, 1851, Pol. Foss. des Terr. Pal., p. 288, Up. Held. Gr.
- tenella*, Rominger, 1876, Foss. Corals, p. 81, Niagara Gr.
- tubiporoides*, Yandell & Shumard, 1847, Cont. to Geo. of Ky., p. 8, Corniferous Gr.
- tubiporoides*, Billings, see *S. maclurii*.
- verneuli*, Edwards & Haime, 1851, Polyp. Foss. de Terr. Pal., p. 289, Corniferous Gr.
- verticillata*, Goldfuss, 1826, Petref. Germ., p. 76, Niagara Gr.
- TETRADIUM*, Dana, 1848, Wilkes. Expl. Exped. Zooph., vol. 8, p. 701. [Ety. *tetra*, four.] Aggregate, massive, subhemispheric; corallites long, prismatic, in close contact; septa few, not reaching the center of the visceral chamber (typically four); tabule numerous, complete; calices generally petaloid; no mural pores; increase by fission. Type *T. fibratum*.

FIG. 224.—*Tetradium fibratum*.

- columnare*, Hall, 1847, (Chetetes columnaria), Pal. N. Y., vol. 1, p. 68, Trenton Gr.
- fibratum*, Safford, 1856, Am. Jour. Sci., vol. 22, p. 237, Hud. Riv. Gr.
- fibratum* var. *aperatum*, Safford, 1856, Am. Jour. Sci., vol. 22, p. 237, Hud. Riv. Gr.
- fibratum* var. *minus*, Safford, 1856, Am. Jour. Sci., vol. 22, p. 238, Hud. Riv. Gr.
- huronense*, Billings, 1865, (Stenopora huronensis.) Pal. Foss., vol. 1, p. 185, Hud. Riv. Gr.
- peachi* var. *canadense*, Foord, 1883, Cont. to Micro. Pal., p. 24, Trenton Gr.
- Tetraraptus*, Salter, 1863, Quar. Jour. Geo. Soc., vol. 19. [Ety. *tetra*, four; *grapho*, I write.] This genus is not regarded with much favor. *Graptolithus bryonoides* is made the typical species. *G. quadribrachiatum* is also placed in it.

FIG. 225.—*Tetradium fibratum*. Corallites scattered through the rock.

approximate.
matus.
THAMNOGRA,
3, p. 519
I write.
or flexu
widely
long, sh
manner
and bra
longitud
the axis
anna, Ha
ade 2, p
bartonens
8, and H
p. 39, N
capillaris,

FIG. 226.—*Tetradium* type.

TRECEIA, Edw
rend, t.
Corallum
compact,
duced by
together
developed
shallow,
Type T.

major, Ro
67, Niag
minor, Ro
68, Niag
ramosa, R
69, Up.
swinderna
swinderna
Niagara
THECOSTEGIT
Comptes
theke, sh
lites cyl
short m

approximatus, see *Graptolithus approximatus*.

THAMNOGRAPTUS, Hall, 1856, Pal. N. Y., vol. 3, p. 519. [Ety. *thamnus*, shrub; *grapho*, I write.] Fronds consisting of straight or flexuous stipes, with alternating or widely diverging branches; branches long, simple, or ramose, in the same manner as the stipe; the main stipe and branches are marked by a central longitudinal, depressed line, indicating the axis. Type *T. typus*.

typus, Hall, 1856, Can. Org. Rem., Decade 2, p. 141, Quebec Gr.

hartonensis, Spencer, 1878, Can. Nat., vol. 8, and Bull. No. 1, Mus. St. Univ. Mo., p. 39, Niagara Gr.

capillaris, Hall, 1859, Pal. N. Y., vol. 3, p. 520, Hud. Riv. Gr.

multiformis, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 40, Niagara Gr.

typus, Hall, 1859, Pal. N. Y., vol. 3, p. 519, Hud. Riv. Gr.

FIG. 226.—*Thamnograptus typus*.

THECIA, Edwards & Haime, 1849, Comptes rend., t. 29, p. 263. [Ety. *theke*, sheath.] Corallum massive, with an abundant, compact, spurious coenenchyma, produced by the septa becoming cemented together laterally; septal system highly developed; tabulæ numerous; calices shallow, with a small deep fossula. Type *T. swindernana*.

FIG. 227.—*Thecia major*.

major, Rominger, 1876, Foss. Corals, p. 67, Niagara Gr.

minor, Rominger, 1876, Foss. Corals, p. 68, Niagara Gr.

ramosa, Rominger, 1876, Foss. Corals, p. 69, Up. Held. Gr.

swindernana, Goldfuss, 1829, (Agaricia *swindernana*.) Petref. Germ., p. 109, Niagara Gr.

THECOSTEGITES, Edwards & Haime, 1849, Comptes rend., t. 29, p. 261. [Ety. *theke*, sheath; *steges*, covering.] Corallites cylindrical, short and united by short mural expansions situated at

various heights; tabulæ horizontal.

Type *T. bouchardi*.

bouchardi, Michelin, 1845, (Harmodites *bouchardi*.) Icon. Zooph., p. 185. This species was described from France, and is probably not American.

FIG. 228.—*Thecostegites hemisphericus*, natural size and magnified.

hemisphericus, Roemer, 1860, Sil. Fauna W. Tenn., p. 25, Niagara Gr.

TRACHYPORA, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 305. [Ety. *trachys*, rough; *poros*, pore.] Dendroid; calices slightly salient; no septa; coenenchyma abundant, solid, and surface marked by strong, irregular, vermicular, subechinulated striae. Type *T. davidsoni*.

austini, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 81, Coal Meas.

elegantula, Billings, 1860, Can. Jour., vol. 5, p. 254, Ham. Gr.

ornata, Rominger, 1876, (Dendropora *ornata*.) Foss. Corals, p. 62, Ham. Gr.

FIG. 229.—*Trachypora elegantula*. Portion of two corallites—a longitudinal section and a corallite enlarged.

TROCHOPHYLLUM, Edwards & Haime, 1851, Mon. d. Pol. Foss. de Terr. Pal., p. 356. [Ety. *trochos*, wheel; *phyllon*, leaf.] Simple, trochoid; calice shallow; septa thick, not denticulate, extending almost to the center of the visceral chamber, where a small tabula is visible; fossula rudimentary and occupied by a small septum. Type *T. verneuilanum*.

verneuilanum, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 357, Subcarboniferous.

TUBIPORA, Linnæus, 1758, Syst. Nat., 10th Ed., p. 789. Not American Palæozoic.

lamellosa, Owen, 1840, Rep. on Min. Lands, p. 78. Not defined. Probably a *Syringopora*.

VERMIPORA, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 109. [Ety. *vermis*, worm; *poros*, pore.] Ramose; corallites cylindrical, close, increasing by lateral gemmation, and projecting at the surface; tabulæ remote; no mural pores connecting corallites. Type *V. serpuloides*.

- fasciculata*, Rominger, 1876, Foss. Corals, p. 70, Ham. Gr.
niagarensis, Rominger, 1876, Foss. Corals, p. 70, Niagara Gr.
robusta, Hall, 1883, Rep. St. Geo., pl. 2, figs. 32, 33, Low. Held. Gr.
serpuloidea, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 110, Low. Held. Gr.
tortuosa, Hall, 1833, Rep. St. Geo., pl. 2, fig. 23, Low. Held. Gr.
Vesicularia, Rominger, 1876, Foss. Corals, p. 135. This name was preoccupied among the Bryozoa. See *Cystiphorolites*.
major, see *Cystiphorolites major*.
minor, see *Cystiphorolites minor*.
variolosa, see *Cystiphorolites variolosus*.
ZAPHIRENTIS, Rafinesque, 1820, Ann. des Sci. Phys. Brux., vol. 5, p. 234. [Ety. *za*, very; *phrentis*, diaphragm.] Simple, turbinate; lamellae simple, alternate, extending from the epitheca to the center of the visceral chamber; tabulae well developed, extending from wall to wall, and deflected downward around the periphery; no columella; calice deep, with a single strongly developed fossula occupying the place of one of the lamellae. Type *Z. phrygia*.
acuta, White & Whitfield, 1882, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 306, Waverly or Choteau Gr.
affinis, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 430, Hud. Riv. Gr.
ampla, Hall, 1876, Illust. Dev. Foss., pl. 21, Ham. Gr.
annulata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 33, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 437, Up. Held. Gr.
bellistriata, Billings, 1865, Can. Nat. and Geo., 2d ser., vol. 2, p. 430, Hud. Riv. Gr.
bigsbyi, Billings, 1866, Catal. Sil. Foss. Antic., p. 92, Clinton and Niagara Grs.
bilateralis, Hall, 1852, (Caninia *bilateralis*), Pal. N. Y., vol. 2, p. 41, Clinton and Niagara Grs.
calcariformis, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 33, and 12th Geo. Ind., p. 293, Up. Held. Gr.
calceola, White & Whitfield, 1882, (Lophophyllum *calceola*), Proc. Bost. Soc. Nat. Hist., vol. 8, p. 305, and 1880, Cont. to Pal. No. 6, p. 156, Waverly or Choteau Gr.
canadensis, Billings, 1862, Pal. Foss., vol. 1, p. 105, Hud. Riv. Gr.
cannonensis, Winchell, 1860, Geo. of Tenn., p. 442, Waverly or Kinderhook Gr.
carniata, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 75, Keokuk Gr.
cassedayi, M. Edwards, 1860, Hist. d. Corallaires, t. 3, Warsaw Gr.
celator, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 107, Niagara Gr.
centralis, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 328, Up. Held. Gr.
chesterensis, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 73, Kaskaskia Gr.
cinctosa, Billings, 1866, Catal. Sil. Foss. Antic., p. 92, Clinton and Niagara Grs.
cingulosa, Billings, 1874, Pal. Foss., vol. 2, p. 10, Gaspe limestone No. 8, Devonian.
clappi, syn. for *Z. gigantea*.
cliffordiana, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 329, Sub-carboniferous.
colletti, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 28, and 12th Rep. Geo. Ind., p. 315, Up. Held. Gr.
complanata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 26, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 430, Up. Held. Gr.
compressa, M. Edwards, 1860, Hist. d. Corallaires, t. 3, Warsaw Gr.
compressa, see *Z. davisana*.
concava, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 35, and 12th Rep. Geo. Ind., p. 291, Up. Held. Gr.
conigera, see *Clisiophyllum conigerum*.
constricta, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 33, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 437, Up. Held. Gr.
contorta, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 37, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 441, Up. Held. Gr.
convoluta, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 37, and 12th Rep. Geo. Ind., p. 294, Up. Held. Gr.
cornicula, Lesueur, 1820, (Caryophyllia *cornicula*), Mem. du Mus., vol. 6, p. 297, Up. Held. Gr.
corrugata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 27, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 431, Schoharie Gr.
corticata, Billings, 1874, Pal. Foss., vol. 2, p. 9, Low. Devonian.
cristulata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 10, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 414, Niagara Gr.
cruciformis, Hall, 1883, 12th Rep. Geo. Ind., p. 315, Up. Held. Gr.
curvata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 35, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 439, Up. Held. Gr.
cyathiformis, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 27, and 12th Rep. Geo. Ind., p. 290, Up. Held. Gr.
cylindraceas, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 78, Kaskaskia Gr.
cystica, Winchell, 1860, Rep. Low. Penin. Mich., p. 90, Ham. Gr.
dalei, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 329, Warsaw Gr.

davisana,
 instead
 1876, F
 was pre
 deforma
 p. 290,
denticulata
 America
 desori, Ed
 Pol. Fo
 Held. G
 duplicata,
 and Up
 Rep. G
 edwardsi,
 2, p. 235
 egeria, Bil
 vol. 7, p
 elegans, H
 and Up
 Rep. Ge
 elliptica,
 Nat. Ill
 to Pal.,
 eriphyll
 Geo. vo
 excentric
 U. S. G
 Sur. W.
 Mens.
 fastigata,
 and Up
 N. Y. St
 Held. G
 fenestrata,
 Geo., vo
 foliata, H
 and Up
 Rep. Ge
 frequentat
 agara an
 35th Re
 435, Up
 fusiformis,
 and Up
 Rep. Ge
 genitiva,
 Geo., vo
 gibsoni, W
 p. 117, G
 gigantea,
 vol. 8, U
 glans, see
 gravis, Ha
 Up. He
 N. Y. St
 Held. G
 gregaria, I
 149, Nia
 halli, Edw
 Pol. Fos
 haysi, Me
 Arts, 2
 Held. G
 hecuba, B
 vol. 7, p
 herzeri, H
 and Up
 Rep. Ge

1, Mon.
328, Up.
Geo. Sur.
H. Foss.
ara Grs.
ss., vol.
No. 8,

51, Mon.
29, Sub-

Niagara
nd 12th
d. Gr.
rals Ni-
26, and
Hist., p.

Hist. d.

Niagara
nd 12th
d. Gr.
rum.
rals Ni-
33, and
Hist., p.

Niagara
nd 35th
p. 441,

rals Ni-
37, and
94, Up.

ophyllia
pl. 6, p.

rals Ni-
27, and
Hist., p.

p., vol. 2,

rals Ni-
10, and
Hist., p.

p. Geo.

Niagara
nd 35th
p. 439,

Corals
27, and
90, Up.

s.) (Geo.
ia Gr.
r. Penin.

n. d. Pol.
saw Gr.

davisana, n. sp. Up. Held. Gr. Proposed instead of *Z. compressa* of Rominger, 1876, Foss. Corals, p. 151, pl. 53, which was preoccupied.
deformis, Hall, 1883, 12th Rep. Geo. Ind., p. 290, Up. Held. Gr.
denticulata, Eichwald, 1857. Probably not American.
desori, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 333, Low. Held. Gr.
duplicata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 32, and 12th Rep. Geo. Ind., p. 293, Up. Held. Gr.
edwardsi, Nicholson, 1875, Ohio Pal., vol. 2, p. 235, Up. Held. Gr.
egeria, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 234, Up. Held. Gr.
elegans, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 37, and 12th Rep. Geo. Ind., p. 287, Up. Held. Gr.
elliptica, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 31, and 1880, Cont. to Pal., No. 6, p. 155, Burlington Gr.
eriphyle, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 233, Up. Held. Gr.
excentrica, Meek, 1873, Haydens, 6th Rep. U. S. Geo. Sur. Terr., p. 495, and Geo. Sur. W. 100th Mer., vol. 4, p. 101, Coal Meas.
fastigata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 30, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 434, Up. Held. Gr.
fenestrata, Nicholson, 1875, Can. Nat. and Geo., vol. 7, p. 138, Up. Held. Gr.
foliata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 34, and 12th Rep. Geo. Ind., p. 286, Up. Held. Gr.
frequentata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 31, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 435, Up. Held. Gr.
fusiformis, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 29, and 12th Rep. Geo. Ind., p. 296, Up. Held. Gr.
genitiva, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 235, Up. Held. Gr.
gibsoni, White, 1884, 13th Rep. Geo. Ind., p. 117, Coal Meas.
giganta, Lesueur, 1820, Mem. du. Mus., vol. 6, Up. Held. Gr.
glans, see *Hadrophryllum glans*.
gravis, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 36, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 440, Up. Held. Gr.
gregaria, Rominger, 1876, Foss. Corals, p. 149, Niagara Gr.
halli, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 341, Ham. Gr.
haysi, Meek, 1865, Am. Jour. Sci. and Arts, 2d ser., vol. 40, p. 32, Low. Held. Gr.
hecuba, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 234, Up. Held. Gr.
herzeri, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 35, and 12th Rep. Geo. Ind., p. 292, Up. Held. Gr.

ida, Winchell, 1865, Proc. Acad. Nat. Sci. Phil., p. 117, Waverly or Kinderhook Gr.
illinoisensis, Worthen, (in press.) Geo. Sur. Ill. vol. 8, p. 77, Keokuk Gr.
inequalis, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., syn. for *Palaeophyllum divaricans*.
inclinata, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 34, syn. for *Cynthophyllum angustatum*.
incondita, Billings, 1874, Pal. Foss., vol. 2, p. 7, Devonian.
invenusta, Billings, 1875, Can. Nat. vol. 7, p. 233, Up. Held. Gr.
irregularis, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 34, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 438, Up. Held. Gr.
knappi, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 34, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 438, Up. Held. Gr.
lanceolata, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 76, Warsaw Gr.
latisinus, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 10, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 414, Niagara Gr.
macfarlani, Meek, 1868, Trans. Cl. Acad. Sci., p. 83, Devonian.
marcoui, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 337, Niagara Gr.
minas, Dawson, 1868, Acad. Geo., p. 286, Subcarboniferous.
multilamella, Hall, 1852, Stans. Ex. to Gt. Salt Lake, p. 408, Coal Meas.
multilamellata, Nicholson, 1875, Ohio Pal., vol. 2, p. 236. The name was preoccupied, and the definition is very imperfect.
nitida, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 31, and 12th Rep. Geo. Ind., p. 288, Up. Held. Gr.
nodulosa, Rominger, 1876, Foss. Corals, p. 148, Corniferous Gr.
offeyensis, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 588, Up. Sil.
ovalis, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 29, and 12th Rep. Geo. Ind., p. 294, Up. Held. Gr.
ovibus, Salter, 1855, Belcher's Last of the Arctic Voyages, vol. 2, p. 382, Carboniferous.
parasitica, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 79, Kinderhook Gr.
patens, Billings, 1865, Can. Nat. and Geo. 2d. ser., vol. 2, p. 450, Mid. Sil.
pellaensis, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 74, St. Louis Gr.
planima, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 29, and 12th Rep. Geo. Ind., p. 292, Up. Held. Gr.
ponderosa, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 27, and 12th Rep. Geo. Ind., p. 288, Up. Held. Gr.

- pressula*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 10, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 414, Niagara Gr.
- profunda*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 31, and 12th Rep. Geo. Ind., p. 287, Up. Held. Gr.
- prolixa*, Billings, 1858, Can. Jour. N. S., vol. 4, p. 121, Up. Held. and Ham. Grs.
- prona*, M. Edwards, 1860, Hist. d. Corallaires, t. 3, Warsaw Gr.
- pulmonea*, Lesueur, 1820, (Caryophyllia pulmonea,) Mem. du Mus., vol. 6, Carboniferous.
- racinensis*, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis. and Geo. Wis., vol. 4, p. 277, Niagara Gr.
- raffinesquii*, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 329, Up. Held. Gr.
- recta*, Meek, 1868, Trans. Chi. Acad. Sci., p. 82, Devonian.
- reversa*, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 78, Warsaw Gr.
- rigida*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 9, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 413, Niagara Gr.
- roemeri*, Edwards & Haime, 1851, Mon. d. Pol. Foss. d. Terr. Pal., p. 341, Delthyris Shale, Low. Held. Gr.
- rugatula*, Billings, 1874, Pal. Foss., vol. 2, p. 8, Gaspe limestone No. 1, Up. Sil.
- sentosa*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 32, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 436, Up. Held. Gr.
- simplex*, Hall, 1843, (Strombodes simplex,) Geo. Rep. 4th Dist. N. Y., p. 200, and Illust. Dev. Foss., pl. 21, Ham. Gr.
- solida*, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 231, Chemung Gr.
- spatiosa*, see *Heterophrentis spatiosa*.
- spergenensis*, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 77, Warsaw Gr.
- spinulifera*, Hall, 1858, Geo. Sur. Iowa, p. 650, Warsaw Gr.
- spinulosa*, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 334, Kaskaskia Gr.
- spiassa*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 30, and 12th Rep. Geo. Ind., p. 289, Corniferous limestone.
- stansburyi*, Hall, 1852, Stans. Ex. to Gt. Salt Lake, p. 408, Coal Meas.
- stokesi*, Edwards & Haime, 1851, Pol. Foss. d. Terr. Pal., p. 330, Niagara Gr.
- subcompressa*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 28, and 12th Rep. Geo. Ind., p. 286, Up. Held. Gr.
- subrecta*, Billings, 1875, Can. Nat. and Geo., vol. 7, p. 235, Up. Held. Gr.
- subvada*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 11, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 415, Niagara Gr.
- subvesicularis*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 10, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 414, Niagara Gr.
- tabulata*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 27, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 431, Up. Held. Gr.
- terebrata*, Hall, 1883, 12th Rep. Geo. Ind., p. 316, Up. Held. Gr.
- torta*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 30, and 12th Rep. Geo. Ind., p. 285, Up. Held. Gr.
- transversa*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 36, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 440, Up. Held. Gr.
- transversensis*, Winchell, 1866, Rep. Low. Penin. Mich., p. 90, Ham. Gr.
- trisutura*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 30, and 12th Rep. Geo. Ind., p. 289, Up. Held. Gr.
- turbinata*, Hall, 1852, (Polydasma turbinatum,) Pal. N. Y., vol. 2, p. 112, Niagara Gr.
- ulrichi*, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 76, Warsaw Gr.
- umbonata*, Rominger, 1876, Foss. Corals, p. 146, Ham. Gr.
- undata*, Hall, 1883, 12th Rep. Geo. Ind., p. 291, Up. Held. Gr.
- ungula*, Rominger, 1876, Foss. Corals, p. 151, Up. Held. Gr.
- varsoviensis*, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 78, Keokuk Gr.
- venusta*, Hall, 1882, Foss. Corals Niagara and Up. Held. Grs., p. 38, and 35th Rep. N. Y. St. Mus. Nat. Hist., p. 442, Up. Held. Gr.

FIG. 230.—*Zaphrentis wortheni*.

wortheni, Nicholson, 1875, Ohio Pal., vol. 2, p. 235, Corniferous Gr.

SUI

THIS

Crinoidea,

The w

lished a bo

family nar

Paleozoic co

Cystoidea,

The Steller

Echinida by

The fo

composed o

are the caly

and Lichen

Cyclocystoi

loidea were

were sessile

roots for at

of them att

but some d

Blastoidea,

Prof.

characters

tinguished

1. The

squamous in

2. The

posteriorly,

probably, in

3. The

ing a dome

the dome o

Witho

can not alw

Heterocrini

Heterocrini

Ohioerinus

beyond the

We be

upon the co

ably no fa

SUBKINGDOM ECHINODERMATA.

THIS Subkingdom is represented, in the Palæozoic rocks, by the Classes Crinoidea, Stellerida, and Echinida.

The word "Crinoidea" was first used in 1821, by J. S. Miller, who published a book entitled "A Natural History of the Crinoidea." He used it as a family name, but later investigations raised it to the rank of a Class. The Palæozoic Orders, into which the Class is divided, are Palæocrinoidea, Blastoida, Cystoida, Lichenocrinoidea, Agelacrinoidea, Cyclocystoida and Myelodactyloidea. The Stellerida are represented by the Orders Asteroidea and Ophiuroidea, and the Echinida by the Order Perischoechinida.

The fossils consist of plates, variously arranged and connected, all of which are composed of peculiar crystalline lime. The principal parts of the Palæocrinoidea are the calyx or body, arms, pinnules, column, and base or root. The Agelacrinoidea and Lichenocrinoidea were attached, by one side, to some foreign substance. The Cyclocystoida were free or attached in like manner. Whether the Myelodactyloidea were free or attached to other bodies is unknown. Some of the Cystoida were sessile, others possessed columns tapering to a point, and others had bases or roots for attachment. The Blastoida possessed columns, but whether or not any of them attached by bases or roots is unknown. All Palæocrinoidea had columns, but some did not have bases or roots. The Orders bearing pinnules are the Blastoida, Palæocrinoidea, and part of the Cystoida.

Prof. Wachsmuth has claimed the construction of the vault affords good characters for the separation of the Palæocrinoidea into families, and has distinguished three plans upon which the summit is constructed, viz.:

1. The summit composed of a more or less pliable, sometimes perhaps squamous integument, yielding to motion, in the body and arms.
2. The summit composed of solid plates, with a porous ventral sac, located posteriorly, on the disk, and closed at the top. Anal opening rarely observed, but, probably, lateral.
3. The summit composed of heavy immovable plates, closely joining and forming a dome arching the entire oral side. Anal opening directly through the wall of the dome or at the extremity of a tube, the so-called proboscis.

Without underestimating his work, a single illustration will show that families can not always be distinguished by the construction of the vault; for in the family Heterocrinidae, there is no resemblance between the vaults of Ectenocrinus, Heterocrinus, Iocrinus, and Ohioocrinus. Ectenocrinus has no tube or proboscis, Ohioocrinus has a large spiral tube, and Iocrinus has a long cylindrical one, extending beyond the ends of the arms and flowing pinnules.

We believe the separation of the Palæocrinoidea into families must be based upon the construction of the calyx and vault, but chiefly upon the former. Probably no family should be made to include genera, some of which have subradials

and others do not. It is therefore of the first importance to ascertain whether the crinoid has one or two circles of plates below the radials. Those having only one circle have been called monocyclic, and those having two circles dicyclic. The circle at the base is composed of what we call the "basal plates," and the second circle, whenever it has an existence, is composed of "subradial plates." In this we follow Billings, Meek, Agassiz, and most other standard authors. Carpenter and Wachsmuth call the "subradials" the "basals" in all cases where they occur, and the lower plates "underbasals;" but where there are no "subradials," they follow the well-established nomenclature in calling the first circle of plates "basals."

The presence or absence of regular interradians, it seems, should always be regarded as of family importance.

The number of basal plates should also be regarded as of family importance. If not in all cases, then in connection with the general structure of the calyx and vault the families will be sufficiently well-defined. Those characters upon which genera are founded, when combined, in certain associations will form families; and under this head several important families have been created.

Generic characters, as a matter of course, are to a certain extent included in the family characters; but the form and construction of the column is of generic importance. The general form of the calyx and vault, and the number of primary radials, and the construction of the azygous area, are always of generic importance. Beside, certain combinations and associations of what are usually regarded as specific characters have been made the basis for establishing genera.

Wachsmuth, speaking from experience, says in young crinoids the basals are the most perfectly developed parts; they attain nearly their full size in young individuals, greater in proportion than the subradials and radials, which are comparatively early developed, and at a time when the interradian and anal plates have scarcely made their appearance. The latter develop the slowest, and in some genera increase continually, both in size and number, during the growth of the individual. Abnormal growths, or sudden modifications of specific characters, almost always take place in the interradian and azygous areas, the azygous rays and dome. His experience is corroborated by others, and the author never saw a small specimen that did not have its basals or first circle of plates as distinctly marked as they occur in large specimens of the same species.

The columns of crinoids very frequently show injuries received by the animal in its life-time. The column is sometimes much swollen on one side and depressed on the other; sometimes a parasite that attached to the column is found imbedded or enveloped in the crinoid column. The animal could also repair its arms and other parts of its body by secretions of lime in the same way.

Some of the Cystoidea may be arranged into families, upon characters similar to those upon which families are founded in the Palaeocrinoidea; but, generally, this is not the case. Some of the Cystoidea possessed an ambulacral opening and two other orifices, the purposes of which may not be fully understood; in others, the two openings referred to are absent. In addition to these, many bear openings called "pectinated rhombs," and all have pores passing through the plates. These pores passed to organs called "hydrospires," which were largely developed within the calyx of the Cystoidea and Blastoidea. The communication, through the test, with the outside water is supposed to show the hydrospires belonged to the respira-

tory system. The rhombs consist of orifices at the fissures and have five spires situated some have

In the or forked radials is or interradians and the test occupied wise by the marked by ambulacra of the hydrospires the gap between hydrospires externally anal inter The plates down the

The connect with and radial also had a Lichenocr Internally supported been common have main evidence and effluents sponges, but of this order three orders in rocks late

The C pentagonal face of the along the is a more of aeozoic order seas. In plates from

tory system. The number and position of the larger orifices and the pectinated rhombs constitute the principal basis for family classification. The Blastoida have orifices at the summit of the calyx which are important in classification. Some have fissures at the summit, others have slits along each side of the ambulacra, and others have five pairs surrounding an oral center. These openings connect with hydrospires situated beneath the ambulacra. These orifices are of family importance, and some have regarded the number of hydrospires as of generic importance.

In the nomenclature of the Blastoida the calyx consists of the basals, radials or forked plates, and orals or deltoid plates. The suture between the basals and radials is the basi-radial suture. The ridge at the median line of an oral is an oral or interradiar ridge. In the forked plates the lower part is the body of the radial, and the two prongs are the limbs. Between the limbs is the radial sinus, which is occupied by the ambulacrum, consisting of a lancet-piece, which is excavated lengthwise by the food-groove or ambulacrum, and against it rest side plates or pore pieces, marked by pinnule pits or sockets, and there are also side plates. Beneath the ambulacra there are interradiar systems of lamellar tubes or hydrospires. The openings of these tubes on the ventral surface of the calyx, as in *Codaster*, are called hydrospire slits; if they are concentrated beneath the ambulacra, as in *Codonites*, the gap between the edge of the lancet-plate and the sides of the radial sinus is the hydrospire cleft, which leads downward into the hydrospire canal. The canals open externally by spiracles, sometimes called ovarian openings. The spiracles of the anal interradius may be confluent with the anal opening to form the anal spiracle. The plates covering the mouth and peristome, and which are sometimes continued down the ambulacra covering the food-grooves, are the summit plates or the vault.

The Cyclocystoidea have tubes radiating from the center of the disk, which connect with a circular tube in the rim. It is evident there was both a circular and radiate system of circulation in this order of animals. The Myelodactyloidea also had a compound internal system of both circular and radiate circulation. The Lichenocrinoidea attached by a base that appears to have been a single solid plate. Internally there are numerous thin, upright septa radiating from the center, which supported the very small external plates, and the sarcode between which must have been connected with the tube in the column to have given support to it, and to have maintained it in an upright position. The column tapered to a point, and no evidence has been found of any external opening of these animals. The affluent and effluent openings that abound in all other Echinoderms, and even among the sponges, have thus far never been discovered in the Lichenocrinoidea. The notice of this order in Wachsmuth's Palæocrinoidea seems to be wholly erroneous. The three orders—Cyclocystoidea, Myelodactyloidea and Lichenocrinoidea—are unknown in rocks later than the Upper Silurian.

The Class Stellerida is composed of animals with a flattened and more or less pentagonal body and central disk. The mouth opens in the center of the lower surface of the disk; the skin is coriaceous, the whole body more or less flexible, and along the lower surface of each arm or prolonged ray from the central disk, there is a more or less distinct furrow from which the ambulacra are protruded. The Palæozoic orders, Asteroidea and Ophiuroidea, are exceedingly abundant in all existing seas. In the common starfish the arms are mere prolongations of the disk, and the plates from which the ambulacra are exerted are in deep furrows along the lower

surface of the arms. The mouth is in the center of the disk, and the ramifications of the stomach extend a greater or less distance into the arm-furrows. In the Ophiuroidea [*Ophis*, snake; *oura*, tail] there are usually five simple curving or flowing arms with undefined furrows and furnished with cirri, which give them a ragged and tangled exterior.

The class Echinida is composed of animals having a complete exterior calcareous shell of closely-fitting plates, which prevents all flexion of the body. The animal has no arms, but the holes, through which the sucking feet are protruded, are arranged upon five rows of plates running from the center of the top of the shell to the angles of the mouth at the bottom; or, when they are confined to the dorsal surface, they form a distinct five-rayed star surrounding the apex of the shell. A striking character in this class is the manner in which spines are articulated upon tubercles on the surface of the shell; the base of the spines being hollowed for the reception of the convex surface of the tubercle, and, being sustained in place by a ligament, the spines are movable, and serve economical purposes. The Palæozoic order Perischoechinida is extinct, but some of them had an internal masticatory apparatus that will compare with any that exists in the living representatives.

CLASS CRINOIDEA.

ORDER PALÆOCRINOIDEA.

FAMILY ACROCRINIDÆ.—Acrocrinus.

FAMILY ACTINOCRINIDÆ.—Actinocrinus, Agaricocrinus, Allopriosalocrinus, Amphorocrinus, Batocrinus, Dorycrinus, Eretmocrinus, Gennæocrinus, Megistocrinus, Melocrinus, Physetocrinus, Saccocrinus, Siphonocrinus, (?) Steganocrinus, Stereocrinus, Strotocrinus, Teleocrinus.

FAMILY AGASSIZOCRINIDÆ.—Agassizocrinus.

FAMILY ALLAGECRINIDÆ.—Allagecrinus.

FAMILY ANCYROCRINIDÆ.—Ancyrocrinus.

FAMILY ARTHRACANTHIDÆ.—Arthracantha.

FAMILY BELEMNOCRINIDÆ.—Belemnocrinus.

FAMILY CALCEOCRINIDÆ.—Calceocrinus, Deltacrinus.

FAMILY CAMAROCRINIDÆ.—Camarocrinus.

FAMILY CATILLOCRINIDÆ.—Catillocrinus.

FAMILY CUPRESSOCRINIDÆ.—Aspidocrinus.

FAMILY CYATHOCRINIDÆ.—Ampheristocrinus, Arachnocrinus, Atelestocrinus, Barycrinus, Carabocrinus, Cyathocrinus, Erisocrinus, Eupachyrcinus, Euspirocrinus, Menocrinus, Palæocrinus, Vasocrinus.

FAMILY DICHOCRINIDÆ.—Cotyledonocrinus, Dichocrinus, Pterotocrinus, Talarocrinus.

FAMILY DIMEROCRINIDÆ.—Coronocrinus, Cytoocrinus.

FAMILY EDRIOCRINIDÆ.—Edriocrinus.

FAMILY EUCALYPTOCRINIDÆ.—Eucalyptocrinus.

FAMILY GASTEROCOMIDÆ.—Myrtillocrinus.

FAMILY GAUROCRINIDÆ.—Gaurocrinus, Retiocrinus, Rhaphanocrinus, Thyasnocrinus.

FAMILY GLYPTASTERIDÆ.—Glyptaster, Lampterocrinus.

- FAMILY GLYPTOCRINIDÆ.—Archæocrinus, Compsoocrinus, Glyptocrinus, Pycno-
crinus, Schizocrinus.
- FAMILY HAPLOCINIDÆ.—Coccoocrinus, Haplocrinus.
- FAMILY HETEROCRINIDÆ.—Ectenocrinus, Heterocrinus, Iocrinus, Ohioocrinus.
- FAMILY HYBOCRINIDÆ.—Anomalocrinus, Hyboocrinus.
- FAMILY ICHTHYOCRINIDÆ.—Cleioocrinus, Ichthyocrinus, Lecanocrinus, Mespilo-
crinus, Nipterocrinus, Onychocrinus, Taxocrinus.
- FAMILY MELOCINIDÆ.—Alloocrinus, Dolatocrinus, Macrostylocrinus, Maria-
crinus, Technocrinus.
- FAMILY PISOCRINIDÆ.—Pisocrinus.
- FAMILY PLATYCRINIDÆ.—Cordylocrinus, Eucladocrinus, Marsupiocrinus, Platy-
crinus.
- FAMILY POTERIOCRINIDÆ.—Bursacrinus, Coeliocrinus, Dendrocrinus, Graphio-
crinus, Homocrinus, Hydreionocrinus, Meroocrinus, Ottawacrinus, Poter-
iocrinus, Stemmatoocrinus, Zeacrinus.
- FAMILY RHODOCRINIDÆ.—Goniasteroidocrinus, Hadrocrinus, Lyriocrinus,
Rhodocrinus.
- FAMILY SYNATHOCINIDÆ.—Synathocrinus.
- FAMILY TAXOCRINIDÆ.—Cupulocrinus, Forbesiocrinus, Taxocrinus.
- FAMILY XENOCRINIDÆ.—Xenocrinus.
- FAMILY AFFINITY UNCERTAIN.—Brachiocrinus, Closterocrinus, Cystocrinus.

ORDER CYSTOIDEA.

- FAMILY AMYGDALOCYSTIDÆ.—Amygdalocystites, Palæocystites.
- FAMILY ANOMALOCYSTIDÆ.—Anomalocystites.
- FAMILY CARYOCRINIDÆ.—Caryocrinus.
- FAMILY COMAROCYSTIDÆ.—Comarocystites.
- FAMILY ECHINOCYSTIDÆ.—Echinocystites.
- FAMILY EOCTIDÆ.—Eocystites.
- FAMILY GOMPHOCYSTIDÆ.—Gomphocystites, Hemicosmites.
- FAMILY HOLOCYSTIDÆ.—Allocystites, Crinocystites, Holocystites.
- FAMILY HYBOCTIDÆ.—Hybocystites.
- FAMILY LEPADOCRINIDÆ.—Apiocystites, Callocystites, Glyptocystites, Lepado-
crinus, Pleurocystites, Sphaerocystites, Strobilocystites.
- FAMILY PLATYCISTIDÆ.—Platycystites.
- FAMILY UNCERTAIN.—Heterocystites, Lysocystites, Malocystites, Porocrinus.

ORDER BLASTOIDEA.

- FAMILY BLASTOIDOCRINIDÆ.—Blastoidocrinus.
- FAMILY CODASTERIDÆ.—Codaster, Heteroschisma.
- FAMILY CODONITIDÆ.—Codonites.
- FAMILY ELEUTHEROCRINIDÆ.—Eleutheroocrinus.
- FAMILY GRANATOCRINIDÆ.—Granatocrinus, Schizoblastus.
- FAMILY NUCLEOCRINIDÆ.—Nucleocrinus.
- FAMILY PENTREMITIDÆ.—Pentremites, Pentremitidea.
- FAMILY STEPHANOCRINIDÆ.—Stephanocrinus.
- FAMILY TROOSTOCRINIDÆ.—Troostocrinus, Tricælocrinus.

ORDER AGELACRINOIDEA.

FAMILY AGELACRINIDÆ.—Agelacrinus, Echinodiscus, Edrioaster, Lepidodiscus.

FAMILY HEMICYSTIDÆ.—Hemicystites.

ORDER MYELODACTYLOIDEA.

FAMILY MYELODACTYLIDÆ.—Myelodactylus.

ORDER CYCLOCYSTOIDEA.

FAMILY CYCLOCYSTOIDIDÆ.—Cyclocystoides.

ORDER LICHENOCRINOIDEA.

FAMILY LICHENOCRINIDÆ.—Lichenocrinus.

CLASS STELLERIDA.

ORDER ASTEROIDEA.

FAMILY ONYCHASTERIDÆ.—Onychaster.

FAMILY PALÆASTERIDÆ.—Cholaster, Compsaster, Palæaster, Palæasterina, Petraster, Schoenaster, Stenaster, Tremataster.

ORDER OPHIUROIDEA.

FAMILY PROTASTERIDÆ.—Eugaster, Palæocoma, Protaster, Tæniaster.

CLASS ECHINIDA.

ORDER PERISCHOECHINIDA.

FAMILY ARCHÆOCIDARIDÆ.—Archæocidaris, Eocidaris, Lepidocidaris, Perichodonus, Pholidocidaris.

FAMILY LEPIDECHINIDÆ.—Hybochinus, Lepidechinus.

FAMILY PALÆCHINIDÆ.—Lepidesthes, Melonites, Oligoporus, Palæchinus.

ACROCRINUS, Yandell, 1855, Am. Jour. Sci. and Arts, 2d ser., vol. 20, p. 135. [Ety. *akros*, extreme, from the great number of plates covering the body; *krinon*, lily.] Body goblet or urn-shaped, consisting of many series of plates; two basals, the suture from the anterior to the posterior side, followed by a series of small plates, and these again by another and another, so that the plates reach the 5th to 10th series before the arms become free; the size of the plates increase as they approach the arms; arms 20, long, composed of two series of plates bearing pinnules; column round. Type A. shumardi.

shumardi, Yandell, 1855, Am. Jour. Sci. and Arts, 2d ser., vol. 20, p. 135, Kas-kaskia G.

urniformi Hall, 1858, Geo. Rep. Iowa, p. 690, Kas-kaskia Gr.

wortheni, Wachsmuth, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 4, and Geo. Sur. Ill., vol. 7, p. 343, Coal Meas.

ACTINOCRINUS, Miller, 1821, Nat. Hist. Crinoidea, p. 95. [Ety. *aktin*, ray; *krinon*, lily.] Body turbinate, plates sculptured; basals 3; primary radials 3x5; secondary 1x10, axillary; succeeding radials having a single series to each division, one axillary, the other simple; arms 20 to 50 or more; pinnules; regular interradianals, one in the first series, two in the second, and one or two in the third; azygous interradianals, one in line with the first primary radials, and of the same size, two in the second series, and one, two, or three in succeeding series; vault variable, plates nodose; tube or proboscis large, subcentral; column long. Type A. triacontadactylus.

abnormis,
agilops, se
aqualis, s
equibrach
iatius.
equibrach
crinus
agassizi, T
althea, see
amplus, se
andrewsia
araneolus,
arnoldi,
press.)
Kinder
asterias,
ney, 18
New. I
Syn. f
crinus
sus.
asteriscus,
crinus
biturbinate
Batocrin
turbina
brevicornis
gisticrin
cornis.
brevia, see
crinus b
brontes, F
Sup. to
Iowa, p
Geo. S
vol. 5,
Warsaw
celatus, F
Geo. St
p. 585,
Sur. Ill
p. 341,
ton (Gr.
calyculo
Eretmoc
yculoide
calyculus,
crinus c
calypso, s
neocrin
lypso.
cassedayi,
carica, see
caroli, see
cauliculus,
chloris, Ha
syn. for
christyi, St
christyi.
christyi, Ha
clarus, Ha
p. 2, and
Burlingt
clavigerus,
dio, see Er
clivus, see
clelia, see
clypeatus,
concavus, s

abnormis, see *Megistocrinus abnormis*.
ægilops, see *Teleocrinus ægilops*.
æqualis, see *Batocrinus æqualis*.
æquibrachiatus, see *Batocrinus æquibrachiatus*.
æquibrachiatus var. *alatus*, syn. for *Batocrinus æquibrachiatus*.
agassizi, Troost, 1850, Catal. Not defined.
althea, see *Teleocrinus althea*.
amplus, see *Saccocrinus amplus*.
andrewsanus, see *Batocrinus andrewsanus*.
araneolus, see *Steganocrinus araneolus*.
arnoldi, Wachsmuth & Springer, (in press,) Geo. Sur. Ill., vol. 8, p. 168, Kinderhook Gr.

asterias, McChesney, 1860, Desc. New. Pal. Foss. Syn. for *Batocrinus verrucosus*.

asteriscus, see *Batocrinus asteriscus*.
biturbinatus, see *Batocrinus biturbinatus*.

brevicornis, see *Megistocrinus brevicornis*.

brevis, see *Agaricocrinus brevis*.

brontes, Hall, 1860, Sup. to Geo. Sur. Iowa, p. 47, and Geo. Sur. Ill., vol. 5, p. 341, Warsaw Gr.

cælatus, Hall, 1858, Geo. Sur. Iowa, p. 585, and Geo. Sur. Ill., vol. 5, p. 341, Burlington Gr.

calyculoides, see *Eretmocrinus calyculoides*.

calyculus, see *Batocrinus calyculus*.

calypso, see *Gennæocrinus calypso*.

cassedayi, see *Gennæocrinus cassedayi*.

carica, see *Eretmocrinus carica*.

caroli, see *Batocrinus caroli*.

cauliculus, see *Gennæocrinus cauliculus*.

chloris, Hall, 1861, Desc. New Crinoidea, syn. for *A. tenuisculptus*.

christyi, Shumard, 1855, see *Batocrinus christyi*.

christyi, Hall, see *Saccocrinus christyi*.

clarus, Hall, 1861, Desc. New Crinoidea, p. 2, and Geo. Sur. Ill., vol. 5, p. 341, Burlington Gr.

clavigerus, see *Batocrinus clavigerus*.

clio, see *Eretmocrinus clio*.

clivus, see *Teleocrinus clivus*.

cllelia, see *Eretmocrinus cllelia*.

clypeatus, see *Batocrinus clypeatus*.

concaus, see *Dorycrinus concaus*.

concinus, see *Steganocrinus concinns*.

copei, see *Physetocrinus copei*.

corbulis, see *Eretmocrinus corbulis*.

coreyi, Lyon & Casseday, 1859, Am. Jour. Sci. and Arts, 2d ser., vol. 29, p. 78, Keokuk Gr.

corniculum, Hall, 1858, Geo. Rep. Iowa, p. 566, Burlington Gr. Wachsmuth says it is a syn. for *Agaricocrinus brevis*.

cornigerus, Hall, see *Dorycrinus cornigerus*.

cornigerus, Lyon & Casseday, see *Gennæocrinus cornigerus*.

cornutus, Troost, 1850, Catal. Not defined.

FIG. 231.—*Acrocrinus wortheni*; diagram.

coronatus, see *Eretmocrinus coronatus*.

dalyanus, S. A. Miller, 1881, Jour. Clin. Soc. Nat. Hist., vol. 4, p. 309, Burlington Gr.

daphne, Hall, 1864, 17th Rep. N. Y. St. Mus. Nat. Hist., p. 52, and Ohio Pal., vol. 2, p. 162, Waverly Gr.

decornis, see *Dorycrinus decornis*.

delicatus, Meek & Worthen; the young of *Teleocrinus umbrosus*.

desideratus, Hall, syn. for *Dorycrinus missouriensis*.

discoideus, see *Batocrinus discoideus*.

divaricatus, Hall, syn. for *Dorycrinus cornigerus*.

divergens, see *Amphoraocrinus divergens*.

dodecadactylus, see *Batocrinus dodecadactylus*.

doris, see *Batocrinus doris*.
erodus, see *Teleocrinus erodus*.
eryx, Hall, 1861, Desc. New Crinoidea, p. 12, Burlington Gr.
eucharis, see *Gennæocrinus eucharis*.
evansi, see *Megistocrinus evansi*.
excerptus, Hall, 1861, Desc. New Cri-

Foss., p. 24, and Trans. Chi. Acad. Sci., p. 17, Burlington Gr.
icosidactylus, see *Batocrinus icosidactylus*.
indianensis, see *Batocrinus indianensis*.
inflatus, see *Amphoracrinus inflatus*.
infrequens, Hall, 1861, Desc. New Crinoidea, p. 14, Burlington Gr.

FIG. 232.—*Actinocrinus arnoldi*.

noidea, p. 3, and Geo. Sur. Ill., vol. 5, p. 341, Burlington Gr.
fibula, Troost, 1850, Catal. Not defined.
facellus, see *Agaricocrinus facellus*.
fosteri, McChesney, 1860, Desc. New Pal. Foss., p. 19, and Trans. Chi. Acad. Sci., p. 14, Burlington Gr.
formosus, see *Batocrinus formosus*.
gemmaformis, see *Eretmocrinus gemmaformis*.
gibbosus, Troost, 1850. Not defined.
glans, Hall, 1860, Sup. to Geo. Sur. Iowa, p. 16, Burlington Gr.
glyptus, see *Strotocrinus glyptus*.
gouldi, see *Dorycrinus gouldi*.
hageri, see *Batocrinus hageri*.
helice, see *Agaricocrinus helice*.
helice var. *eris*, see *Agaricocrinus eris*.
humboldtii, Troost. Not defined.
hurduanus, McChesney, 1860, New Pal.

inornatus, see *Batocrinus inornatus*.
insculptus, see *Teleocrinus insculptus*.
irregularis, see *Batocrinus irregularis*.
jugosus, Hall, 1860, Supp. Geo. Sur. Iowa, p. 49, Keokuk Gr.
kentuckiensis, Shumard,

syn. for *Gennæocrinus cornigerus*.
konincki, see *Eretmocrinus konincki*.
lagena, Hall, 1861, Desc. New Crinoidea, p. 13, Burlington Gr.
lagunculus, see *Batocrinus lagunculus*.
laura, see *Batocrinus laura*.
lepidus, see *Batocrinus lepidus*.
leucosia, see *Eretmocrinus leucosia*.
limbrachistatus, Hall, 1861, Desc. New Crinoidea, p. 2, and Bost. Jour. Nat. Hist., p. 268, Burlington Gr.
liratus, see *Teleocrinus liratus*.
lobatus, Hall, 1860, Supp. Geo. Sur. Iowa, p. 51, Keokuk Gr.
locellus, Hall, 1861, Desc. New Crinoidea, p. 15, Burlington Gr.
longirostris, see *Batocrinus longirostris*.
longus, Meek & Worthen, Proc. Acad. Nat. Sci. Phil., p. 156, and Geo. Sur. Ill., vol. 5, p. 345, Burlington Gr.
lowii, Hall, 1858, Geo. Sur. Iowa, p. 611, Keokuk Gr.
lucina, Hall, 1861, Desc. New Crinoidea, p. 11, Burlington Gr.
matuta, see *Eretmocrinus matuta*.

matuta v.
attenu
meeki, p.
minor, p.
 573, B.
mississippiensis
mississippiensis
missouriensis
moniliformis
 Ameri
mortoni,
multibrachia
 Iowa,
multibrachia
 Desc. I.
multicornis
mundulus
multiradialis
 Louis
 Iowa,
nashville
nashville
nashville
novobrachia
 (in pre
 Kinder
nysa, see
oblatus, s
oppyramis
amidal
olla, McC
olliculus,

FIG. 233.—*noerinus tissimus*.

papillatus
parvus, se
pendens, s
penicillus
 Acad.
 Sur. Ill.
pentagonus
pentaspina
pernodos
 p. 608,
perumbros
pistilliform
pistillus, s
planobas
basalis
planodiscu
plumosus,
 N. Y.,
 180, Olf
 and th
 minatio
pocillum, t

ad. Sci.,
actylus.
nsis.
is.
ew Cri.

matuta var. *attenuatus*, see *Eretmocrinus attenuatus*.

meeki, see *Macrostylocrinus meeki*.

minor, Hall, 1858, Geo. Rep. Iowa, p. 573, Burlington Gr.

mississippiensis, see *Dorycrinus mississippiensis*.

mississippiensis var. *spiniger*, see *Dorycrinus mississippiensis* var. *spiniger*.

missouriensis, see *Dorycrinus missouriensis*.

moniliformis, Miller, cited by Troost. Not American.

mortoni, Troost, 1850. Not defined.

multibrachiatus, Hall, 1858, Geo. Rep. Iowa, p. 580, Burlington Gr.

multibrachiatus var. *echinatus*, Hall, 1861, Desc. New Crinoidea, p. 10, Warsaw Gr.

multicornis, see *Centrocrinus multicornis*.

mundulus, see *Batocrinus mundulus*.

multiradiatus, Shumard, 1857, Trans. St. Louis Acad. Sci., p. 75, and Geo. Rep. Iowa, p. 579, Burlington Gr.

nashvillae, see *Batocrinus nashvillae*.

nashvillae var. *subtractus*, see *Batocrinus nashvillae* var. *subtractus*.

novobrachiatus, Wachsmuth & Springer, (in press), Geo. Sur. Ill., vol. 8, p. 165, Kinderhook Gr.

nyssa, see *Gennæocrinus nyssa*.

oblatulus, see *Batocrinus oblatulus*.

obpyramidalis, see *Melocrinus obpyramidalis*.

olla, McCoy, 1849. Not American.

olliculus, syn. for *Megistocrinus whittii*.

opusculum, Hall, 1861, Bost. Jour. Nat. Hist., p. 264, Burlington Gr.

ornatissimus, Wachsmuth & Springer, (in press), Geo. Sur. Ill., vol. 8, p. 163, Kinderhook Gr.

ornatus, see *Physetocrinus ornatus*.

ovatus, Hall, 1861, Desc. New Crinoidea, p. 19, Burlington Gr.

papillatus, see *Batocrinus papillatus*.

parvus, see *Dorycrinus parvus*.

pendens, see *Dorycrinus pendens*.

penicillus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 155, and Geo. Sur. Ill., vol. 5, p. 342, Burlington Gr.

pentagonus, see *Steganoecrinus pentagonus*.

pentaspinus, see *Centrocrinus pentaspinus*.

pernodosus, Hall, 1858, Geo. Rep. Iowa, p. 608, Keokuk Gr.

perumbrosus, see *Strotocrinus perumbrosus*.

pistilliformis, see *Batocrinus pistilliformis*.

pistillus, see *Batocrinus pistillus*.

planobasalis, see *Amphoracrinus planobasalis*.

planodiscus, see *Batocrinus planodiscus*.

plumosus, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 72, and Pal. N. Y., vol. 2, p. 180, Clinton Gr. Not an *Actinocrinus*, and the fragments too poor for determination.

pocillum, see *Gennæocrinus pocillum*.

polydactylus, see *Mariacrinus polydactylus*.

præcursor, see *Dorycrinus præcursor*.

proboscivalis, Hall, 1858, Geo. Rep. Iowa, p. 584, Burlington Gr.

pyriformis, see *Batocrinus pyriformis*.

pyriformis, var. *rudis*, Meek & Worthen, see *Batocrinus pistilliformis*.

pyramidatus, see *Agaricocrinus pyramidatus*.

quadrifidus, see *Amphoracrinus quadrifidus*.

quaternarius, Hall, 1860, Supp. Geo. Rep. Iowa, p. 22, Burlington Gr.

quaternarius var. *spiniferus*, Hall, 1861, Desc. New Crinoidea, p. 11, Burlington Gr.

quinquelobus, see *Dorycrinus quinquelobus*.

ramulosus, see *Eretmocrinus ramulosus*.

regalis, see *Strotocrinus regalis*.

remibrachiatus, see *Eretmocrinus remibrachiatus*.

reticulatus, see *Physetocrinus reticulatus*.

rotundus, see *Batocrinus rotundus*.

rudis, see *Teleocrinus rudis*.

rusticus, Hall, 1861, Desc. New Crinoidea, p. 2, syn. for *A. scitulus*.

FIG. 234.—*Actinocrinus proboscivalis*.

Batocri-
ps.
Teleio-
ptus.
Batocri-
ria.
ria, 1860,
ur. Iowa,
k Gr.
humard,
igerus.
aincki.
y Crino-
unculus.

FIG. 233.—*Actinocrinus ornatissimus*.

FIG. 235.—*Actinocrinus scitulus*. Diagram x 2.

scitulus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 386, and Geo. Sur. Ill., vol. 2, p. 202, Burlington Gr.

sculptus, see *Steganoecrinus sculptus*.

securis, Hall, 1861, Desc. New Crinoidea, p. 14, Burlington Gr.

semiradiatus, see *Saccocrinus semiradiatus*.

senarius, see *Physetocrinus senarius*.

sexarmatus, Hall, 1860, Supp. Geo. Rep. Iowa, p. 21, Burlington Gr.

sillimani, Meek & Worthen, syn. for *A. scitulus*.

similis, see *Batocrinus similis*.

sinuosus, see *Batocrinus sinuosus*.

spicuosus, Meek & Worthen, syn. for *Strotocrinus regalis*.

spinobrachiatus, see *Amphoracrinus spinobrachiatus*.

osia.
esc. New
ur. Nat.
eo. Sur.
inoidea,
stris.
Acad.
eo. Sur.
Gr.
p. 611,
inoidea,

spinotentaculus, Hall, 1860, Supp. Geo. Rep. Iowa, p. 86, Burlington Gr.

spinulosus, see Dorycrinus spinulosus.

steropes, see Batocrinus steropes.

subaculeatus, see Dorycrinus subaculeatus.

subaequalis, see Batocrinus subaequalis.

subturbinatus, see Dorycrinus subturbinatus.

subumbrosus, Hall, syn. for Teleiocrinus liratus.

subventricosus, see Phytocrinus subventricosus.

superlatus, see Megistocrinus superlatus.

symmetricus, see Dorycrinus symmetricus.

tenuidiscus, Hall, 1861, Desc. New Crinoidea, p. 14, Burlington Gr.

tenuiradiatus, Hall, 1847, see Palaeocystites tenuiradiatus.

tenuiradiatus, Hall, 1861, see Teleiocrinus tenuiradiatus.

tenuisculptus, McChesney, 1860, Desc. New Pal. Foss., p. 15, and Trans. Chi.

thalia, Hall, 1861, Desc. New Crinoidea, p. 13, Burlington Gr.

themis, Hall, 1861, Desc. New Crinoidea, p. 11, Burlington Gr.

thetis, Hall, 1861, Desc. New Crinoidea, p. 11, Burlington Gr.

thous, Hall, syn. for A. reticulatus.

tholus, syn. for A. glans.

tricornis, see Dorycrinus tricornis.

trinodus, see Dorycrinus trinodus.

turbinatus, see Batocrinus turbinatus.

turbinatus var. elegans, see Batocrinus turbinatus var. elegans.

umbrosus, see Teleiocrinus umbrosus.

unicarinatus, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 48, Keokuk Gr.

unicornis, see Dorycrinus unicornis.

unispinus, see Dorycrinus unispinus.

urna, Troost, 1850. Not defined.

urniformis, McChesney, 1860, New Pal. Foss., p. 23, syn. for Eretmocrinus konincki.

validus, Meek & Worthen, 1860, syn. for Steganocrinus concinnus.

ventricosus, see Phytocrinus ventricosus.

ventricosus var. cancellatus, see Phytocrinus ventricosus var. cancellatus.

ventricosus var. internodus, see Phytocrinus ventricosus var. internodus.

verneuili, see Melocrinus verneuili.

verneuilianus, see Eretmocrinus verneuilianus.

verrucosus, Hall, 1858, Geo. Rep. Iowa, p. 578, Burlington Gr.

viaticus, White, 1874, Rep. Invert. Foss., p. 16, and Geo. Sur. W. 100th Merid., vol. 4, p. 82, Subcarboniferous.

viminalis, see Amphoracrinus viminalis.

wachsmuthi, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 17, syn. for A. scitulus.

wachsmuthi, White, 1880, see Batocrinus wachsmuthi.

whitfieldi, see Saccocrinus whitfieldi.

whitii, see Megistocrinus whitii.

yandelli, see Batocrinus yandelli.

AGARICOCRINUS, Troost, 1850, Catal. in Proc. Am. Ass'n, and Hall in Geo. Sur. Iowa, p. 560. [Ety. *Agaricus*, mushroom; *crinus*, lily.] The form of the calyx is that of an inverted basin or mushroom; plates smooth; dome composed of large nodose plates and smaller convex ones, the central plate being the largest in the body; basals 3, small; primary radials 3x5; secondary radials 1 or 2x10, which are succeeded by shorter arm-plates; regular interradials 3; azygous plates 4 to 7, or more; aperture, at the upper part, directed laterally; arms long, constructed of two rows of plates bearing pinnule; columns round. Type A. americanus.

americanus, Roemer, 1854, (Amphoracrinus americanus.) Bronn's Leth. Geog., vol. 2, p. 250, and Geo. Sur. Iowa, p. 617, Keokuk Gr.

bellitrema, Hall, 1861, Bost. Jour. Nat. Hist., p. 281, Burlington Gr. Wachsmuth says it is a syn. for A. ornorema.

brevis, Hall, 1858, (Actinocrinus brevis.) Geo. Sur. Iowa, p. 567, Burlington Gr.

bullatus, Hall, 1858, Geo. Sur. Iowa, p. 562, Burlington Gr. Wachsmuth says it is a syn. for A. americanus.

convexus, Hall, 1860, (A. pentagonus var. convexus.) Supp. to Geo. Sur. Iowa, p. 58, Burlington Gr.

Fig. 236.—Agaricocrinus crassus, azygous view of calyx.

corrugatus, Hall, 1861, Desc. New Spec. Crin., p. 4, and Bost. Jour. Nat. Hist., p. 283, Burlington Gr. Wachsmuth says it is a syn. for A. pyramidalis founded upon a mature specimen.

crassus, Wetherby, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 178, Keokuk Gr.

elegans, Wetherby, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 179, Keokuk Gr.

eris, Hall, 1864, (Actinocrinus helice var. eris.) 17th Rep. N. Y. St. Mus. Nat. Hist., p. 53,

Fig. 237.—Agaricocrinus elegans, view of the vault.

and
verly

Fig. 238.—
elegans,

New
Nat.
geomet
Sur.
gracilis
Acad.
ton G
helice,
tinocr
17th
Mus.
53, a
vol. 2
verly
inflatus
Desc.
p. 4,
Nat.
Burlin
macadar
(in p
ill.,
Keok
nodos
Worth
Phil.,
p. 38
says i
nodulos
ill., v
ornotre
oidea,
pentago
Iowa,
pentagon
planoco
Crino
Hist.,
pyrami
pyram
Burlin
springer
and l
kuk C
stellatus
564, B
tuberosus
Geo.
americ
whitfield
p. 621,
worthen
619, K

and Ohio Pal., vol. 2, p. 164, Waverly Gr.

FIG. 238.—Agaricocrinus elegans, basal view.

New Spec. Crin., p. 2, and Bost. Jour. Nat. Hist., p. 272, Burlington Gr. geometricus, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 56, Burlington Gr. gracilis, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 135, Burlington Gr.

helice, Hall, 1864, (Actinocrinus helice,) 17th Rep. N. Y. St. Mus. Nat. Hist., p. 53, and Ohio Pal., vol. 2, p. 163, Waverly Gr.

inflatus, Hall, 1861, Desc. New Crinoidea, p. 4, and Bost. Jour. Nat. Hist., p. 284, Burlington Gr.

macadamisi, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 94, Keokuk Gr.

nodosus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 167, and Geo. Sur. Ill., vol. 5, p. 387, Burlington Gr. Wachsmuth says it is a syn. for A. americanus.

nodulosus, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 94, Keokuk Gr.

ornotremis, Hall, 1861, Desc. New Crinoidea, p. 3, Burlington Gr.

pentagonus, Hall, 1860, Supp. Geo. Rep. Iowa, p. 57, Burlington Gr.

pentagonus var. convexus, see A. convexus.

planoconvexus, Hall, 1861, Desc. New Crinoidea, p. 3, and Bost. Jour. Nat. Hist., p. 280, Burlington Gr.

pyramidatus, Hall, 1858, (Actinocrinus pyramidatus,) Geo. Rep. Iowa, p. 565, Burlington Gr.

springeri, White, 1882, 11th Rep. Geo. and Nat. Hist. Indiana, p. 363, Keokuk Gr.

stellatus, Hall, 1858, Geo. Rep. Iowa, p. 564, Burlington Gr.

tuberosus, Troost, 1850, Catal. Hall, 1858, Geo. Rep. Iowa, p. 617, syn. for A. americanus.

whitfieldi, Hall, 1858, Geo. Rep. Iowa, p. 621, Keokuk Gr.

wortheni, Hall, 1858, Geo. Rep. Iowa, p. 619, Keokuk Gr.

excavatus, Hall, 1861, (Actinocrinus excavatus,) Desc. New Spec. Crin., p. 3, and Bost. Jour. Nat. Hist., p. 282, Burlington Gr. Wachsmuth says it is a syn. for A. americanus.

fiscellus, Hall, 1861, (Actinocrinus fiscellus,) Desc.

FIG. 239.—Agaricocrinus helice.

Agassizocrinus, Troost, 1850 Mss., Shumard, 1853, Marcy's Rep. Red Riv., and Hall, 1858, Geo. Rep. Iowa, p. 684. [Ety. proper name; *krinon*, lily.] Calyx conical or semielliptical; not ornamental; basals 5, usually ankylosed, very small inner cavity; subradials 5, thick, usually ankylosed; radials 2x5; arms 10; azygous plates 3 or 4; column evidenced by a small cylindrical tube extending from a minute cicatrix at the center of the basals to the interior of the cup. Type A. dactyliformis.

carbonarius, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 566, Up. Coal Meas.

chesterensis, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 558, Kaskaskia Gr.

conicus, Owen & Shumard, 1851, Jour. Acad. Nat. Sci. Phil., 2d. ser., vol. 2, p. 93, and Geo. Sur. Ill., vol. 5, p. 557, Kaskaskia Gr.

constrictus, Hall, 1858, Geo. Rep. Iowa, p. 687, Kaskaskia Gr.

dactyliformis, Troost, 1850, described by Shumard, 1853, Marcy's Rep. Red. Riv., p. 199, Kaskaskia Gr.

gibbosus, Hall, 1858, Geo. Rep. Iowa, p. 686, Kaskaskia Gr.

globosus, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 557, Kaskaskia Gr.

gracilis, Troost, 1850. Not defined.

hemisphericus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 37, and Geo. Sur. Ill., vol. 7, p. 316, Kaskaskia Gr.

occidentalis, Owen & Shumard, 1852, (Poterocrinus occidentalis,) Jour. Acad. Nat. Sci. Phil., vol. 2, p. 92, Kaskaskia Gr.

papillatus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 36, and Geo. Sur. Ill., vol. 7, p. 315, Kaskaskia Gr.

pentagonus, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 556, Kaskaskia Gr.

tumidus, Owen & Shumard, 1852, (Poterocrinus tumidus,) Jour. Acad. Nat. Sci. Phil., vol. 2, p. 90, Kaskaskia Gr.

Agelacrinus, Vanuxem, 1842, (Agelacrinus,) Geo. Rep. 3d Dist. N. Y., p. 158. [Ety. *agele*, herd; *krinon*, lily.] A thin, circular, parasitic disk; upper face more or less convex, and composed of thin imbricating plates; ambulacra consist-

FIG. 240.—Agassizocrinus dactyliformis.

ing of a double series of alternating plates, forming convex ridges, constituting part of the upper face, and bearing two or more rows of ambulacral pores; ovarian or anal aperture is situated within the azygous interambulacral area, surrounded by cuneiform plates. Type *A. hamiltonensis*. Billings, Chapman, 1860, Can. Jour., vol. 5, p. 358, Trenton Gr.

FIG. 241.—*Agelacrinus cincinnatiensis*.

- cincinnatiensis*, Roemer, 1851, Verh. Naturh. Rhein. Westph., vol. 8, p. 372, and Ohio Pal. vol. 1, p. 55, Hud. Riv. Gr.
- dicksoni*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 294, and Can. Org. Rem., Dec. 3, p. 84, Trenton Gr.
- hamiltonensis*, Vanuxem, 1842, Geo. Rep. 3d Dist. N. Y., p. 158, Ham. Gr.
- holbrookii*, James, 1887, Jour. Cin. Soc. Nat. Hist., vol. 10, p. 25, Hud. Riv. Gr.
- kaskaskiensis*, see *Echinodiscus kaskaskiensis*.
- pileus*, Hall, 1866, Adv. sheets, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 214, and Ohio Pal., vol. 1, p. 56, Hud. Riv. Gr.
- septembranchiatus*, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 27, Hud. Riv. Gr.
- squamosus*, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 357, and Geo. Sur. Ill., vol. 5, p. 573, Keokuk Gr.
- stellatus*, see *Hemicystites stellatus*.
- vorticellatus*, Hall, 1866, Adv. sheets, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 215, and Ohio Pal., vol. 1, p. 57, Hud. Riv. Gr.
- ALLAGECRINUS**, Etheridge & Carpenter, 1881, Ann. and Mag. Nat. Hist., p. 231. [Ety. *allage*, change; *krinon*, lily.] Calyx minute, pyriform, without ornamentation; basals 5, anchylosed; radials 1 x 5; arms 10; interradials none; column round. Type *A. austini*.
- carpenteri*, Wachsmuth, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 40, and Geo. Sur. Ill., vol. 7, p. 341, Kaskaskia Gr.
- ALLOCRINUS**, Wachsmuth & Springer, (in press.) Geo. Sur. Ill., vol. 8, p. 206. [Ety. *allos*, another; *krinon*, lily.] Calyx small; arms stout; basals 3, small; primary radials 3 x 5, first large, others smaller; secondary radials 2 or 3, rounded, quadrangular; arms composed of transverse plates; interradials two or more deeply impressed; column small; canal pentagonal. Type *A. typus*.
- typus*, Wachsmuth & Springer, (in press.) Geo. Sur. Ill., vol. 8, p. 207, Niagara Gr.
- ALLOCYSTITES**, n. gen. [Ety. *allos*, another; *kustis*, bladder.] Small, irregularly sub-elliptical, tapering below to a small column; plates polygonal, without definite order of arrangement and of very

unequal size; all the plates poriferous; mouth near the margin of the summit; the plates which form it cover part of the body, and on approaching the orifice curve up so as to form part of the opening. The collector says when found it projected an eighth of an inch, and the plates forming the projection were accidentally broken off. The ambulacral opening is upon the extreme height of the summit, and projects above the body, where it is covered by minute plates forming a pentagonal star. Type *A. hammelli*.

hammelli, n. sp., Niagara Gr. In addition to the characters above ascribed to the genus, the ranges of plates, if in regular series, would form about six series; the first series are anchylosed so that two plates only can be distinguished; in the second range there are seven plates; above this the plates are extremely variable in form and size, no two of them being alike; only four plates are distinguished as forming the mouth, but there is no reasonable doubt there are five, and that one is narrow, and situated between the mouth and ambulacral orifice, as is usual in this family of Cystidians. The projecting mouth-plates and elevated ambulacral opening specially characterize this genus and species. The specific name is in honor of Mr. J. F. Hammell, of Madison, Indiana, who collected it in Jefferson County.

FIG. 242.—*Allocystites hammelli*. Side and summit view.

- ALLOPROSALLOCRINUS**, Casseday & Lyon, 1860, Proc. Am. Acad. Arts and Sci., vol. 5, p. 29. [Ety. *allopriosallos*, inclining first to one side and then to another; *krinon*, lily.] Turbinate; basals 3; primary radials 3 x 5; secondary radials 2 x 10; regular interradials 1; azygous plates 3; vault elevated, bearing a central tube or proboscis; arms 11 to 13; distinguished from *Agaricocrinus* by general form and fewer interradials. Type *A. conicus*.
- conicus*, Casseday & Lyon, 1860, Proc. Am. Acad. Arts and Sci., vol. 5, p. 29, Warsaw Gr.
- euconus*, see *Batocrinus euconus*.
- depressus*, Casseday & Lyon, 1860, Proc. Am. Acad. Arts and Sci., vol. 5, p. 31, Warsaw Gr.

AMPHERIS
Spec.
and M
amph
Turbi
area,
als 5;
trix o
for th
regula

FIG. 243.—A

typus, H
p. 11,
Indian
AMPHORAC
Geo. S
Sur. I
cup; A
dome
excent
3; pri
dials I
compo
regula
radial
smalle
gilbert
american
bellatrem
divergen
vergen
and G
ington
excavatus
inflatus,
jerseyen
Sur. II

FIG. 243.—*planobas planobas*. Burlington syn.

AMPHERISTOCRINUS, Hall, 1879, Desc. New Spec. Foss., p. 11, and 11th Rep. Geo. and Nat. Hist. Indiana, p. 278. [Ety. *ampheristos*, disputed; *krinon*, lily.] Turbinate, base attenuate; azygous area, large; plates 6; basals 3; subradials 5; radials 1x5, with a narrow cinctrix on the middle of the top of each for the attachment of the arms; no regular interradials. Type *A. typus*.

FIG. 243.—*Ampheristocrinus typus*. Basal and side view of calyx.

typus, Hall, 1879, Desc. New Spec. Foss., p. 11, and 11th Rep. Geo. and Nat. Hist. Indiana, p. 278, Niagara Gr.

AMPHORACRINUS, Austin, 1848, Quar. Jour. Geo. Soc. Lond., vol. 4, p. 292, and Geo. Sur. Ill., vol. 5, p. 386. [Ety. *amphora*, cup; *krinon*, lily.] Body short, lobed, dome elevated, with tube or proboscis excentric on the azygous side; basals 3; primary radials 3x5; secondary radials 1x10; arms numerous, variable, composed of a double series of plates; regular interradials 3; azygous interradials, 3 or 4 large ones and a few smaller ones; column round. Type *A. gilbertsoni*.

americanus, see *Agaricocrinus americanus*. *bellatrema*, see *Agaricocrinus bellatrema*.

divergens, Hall, 1860, (Actinocrinus *divergens*.) Supp. Geo. Rep. Iowa, p. 36, and Geo. Sur. Ill., vol. 5, p. 388, Burlington Gr.

excavatus, see *Agaricocrinus excavatus*.

inflatus, see *Agaricocrinus inflatus*.

jerseyensis, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 96, Kinderhook Gr.

FIG. 244.—*Amphoracrinus viminalis*.

planobasalis, Hall, 1858, (Actinocrinus *planobasalis*.) Geo. Rep. Iowa, p. 19, Burlington Gr. Wachsmuth says it is a syn. for *A. divergens*.

quadrilobus, White, 1832, (Actinocrinus *quadrilobus*.) Proc. Bost. Soc. Nat. Hist., vol. 9, p. 15, Burlington Gr. Wachsmuth says it is a syn. for *A. divergens*.

spinobrachiatus, Hall, 1860, (Actinocrinus *spinobrachiatus*.) Supp. Geo. Rep. Iowa, p. 6, and Geo. Sur. Ill., vol. 5, p. 389, Burlington Gr.

viminalis, Hall, 1864, (Actinocrinus *viminalis*.) 17th Rep. N. Y. St. Mus. Nat. Hist., p. 54, and Ohio Pal., vol. 2, p. 165, Waverly Gr.

AMYGDALOCYSTITES, Billings, 1854, Can. Jour., vol. 2, p. 270, and Can. Org. Rem., Decade 3, p. 63. [Ety. *amygdalos*, almond; *cystis*, bladder.] Body flattened, ovate, covered with nonporiferous plates arranged without order; ambulacral opening at the apex, mouth near by; arms recumbent, composed of a double series of plates; column round. Type *A. florealis*.

florealis, Billings, 1854, Can. Jour., vol. 2, p. 270, and Can. Org. Rem., Decade 3, p. 63, Trenton Gr.

florealis var. *lævis*, W. R. Billings, 1883, Trans. No. 4, Ottawa Field Nat. Club, p. 52, Trenton Gr.

huntingtoni, Wetherby, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 177, Trenton Gr.

radiatus, Billings, 1854, Can. Jour., vol. 2, p. 271, and Can. Org. Rem., Decade 3, p. 65, Trenton Gr.

tenuistriatus, Billings, 1854, Can. Jour., vol. 2, p. 271, and Can. Org. Rem., Decade 3, p. 64, Trenton Gr.

FIG. 245.—*Amygdalocystites huntingtoni*.

ANCYROCINUS, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 89. [Ety. *ankura*, grapple; *krinon*, lily.] A bulb with lateral ascending processes and a central column. But little is known of this genus. Type *A. bulbosus*.

bulbosus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 90, Ham. Gr.

spinosus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 90, Up. Heid. Gr.

ANOMALOCINUS, Meek & Worthen, 1868, Geo. of Ill., vol. 3, p. 327. [Ety. *anomalos*, irregular; *krinon*, lily.] Calyx depressed, irregularly saucer-shaped; basals 5; sometimes a 6th intercalated one; radials 1x3 and 2x2, the last ones truncated in the central part for the free arms, and curving over on the vault on either side, thus widely separating the arms; arms irregular, frequently bifurcating, composed of a single series of plates, round on the exterior; pinnules strong; vault convex

and supposed to possess a tube or proboscis; column large and longitudinally from 5 to 20 partite. Type A. incurvus.

caponiformis, Lyon, 1869, (Ataxocrinus caponiformis,) Trans. Am. Phil. Soc., vol. 13, p. 464, and Jour. Clin. Soc. Nat. Hist., vol. 2, p. 109, Hud. Riv. Gr.

FIG. 246.—Anomalocrinus incurvus. Diagram.

incurvus, Meek & Worthen, 1865, (Heterocrinus incurvus,) Proc. Acad. Nat. Sci. Phil., p. 148, and Geo. Sur. Ill., vol. 3, p. 327, Hud. Riv. Gr.

ANOMALOCYSTITES, Hall, 1850, Pal. N. Y., vol. 3, p. 132. [Ety. *anomalos*, irregular; *kustis*, bladder.] Somewhat semielliptical, sides unequal, vertical outline oval or ovoid; first series of plates 3 on the convex and 2 on the flat or concave side; second series 4 or 5 on the convex side and 2 on the concave side; third series 4 on the convex and 1 on the other; succeeding series have smaller plates and the apex is unknown; column large at the body and very rapidly tapering; no pores or pectinated rhombs. Type A. cornutus. Wetherby supposed this to be a Crustacean and gave it the name of Euoploura.

FIG. 247.—Anomalocystites balanoides. Convex and flattened sides.

balanoides, Meek, 1872, Am. Jour. Sci., 3d ser., vol. 3, p. 423, and Ohio Pal., vol. 1, p. 41, Hud. Riv. Gr.
cornutus, Hall, 1859, Pal. N. Y., vol. 3, p. 133, Low. Held. Gr.
disparilis, Hall, 1859, Pal. N. Y., vol. 3, p. 145, Oriskany sandstone.
huxleyi, Billings, 1858, (Ateleocystites huxleyi,) Can. Org. Rem., Decade 3, p. 72, Trenton Gr.

Anomaloides, Ulrich, 1878, Jour. Clin. Soc. Nat. Hist., vol. 1, p. 92. A word constructed of adjectives making it meaningless, contrary to the rules of nomenclature, and the attempt to found a genus was made on a fossil fragment not understood.

reticulatus, Ulrich, 1878, Jour. Clin. Soc. Nat. Hist., vol. 1, p. 92, Hud. Riv. Gr. A fossil fragment not understood.

ARIOCYSTITES, Forbes, 1848, Mem. Geo. Sur. Great Brit., vol. 2, p. 502. [Ety. *apion*, pear; *kustis*, bladder.] Body ovoid, or oblong oval, angular and covered by four series of plates; first series has 4 plates; second series 5; third series 5 or 6; fourth series 5 or more; arms 4, recumbent and filling shallow grooves at the angles of the body, column rapidly tapering; ovarian aperture near the summit, on the anterior side; all the plates bearing calycine pores; a pectinated rhomb upon each side in the 3d and 4th series, and one on the anterior side in the 2d series, but these may be variable. Type A. pentremitoides.

canadensis, Billings, 1866, Catal. Sil. Foss. Antic., p. 90, Niagara Gr.

elegans, Hall, 1852, Pal. N. Y., vol. 2, p. 243, Niagara Gr.

huronensis, Billings, 1866, Catal. Sil. Foss. Antic., p. 91, Niagara Gr.

imago, Hall, 1867, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 358, Niagara Gr.

tecumsethi, Billings, 1866, Catal. Sil. Foss. Antic., p. 91, Niagara Gr.

ARACHNOCRINUS, Meek & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 177. [Ety. *arachne*, spider; *krinos*, lily.] Calyx small, resembling Cyathocrinus, but more depressed; basals 5; subradials 5; radials 3x5; arms long, robust, spreading, furrow deep; no pinnules; azygous plate supporting a lateral tube; column round. Type A. bulbosus.

bulbosus, Hall, 1860, (Cyathocrinus bulbosus,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

extensus, Wachsmuth & Springer, 1879, Revis. Paleocrinoidea, p. 93, Ham. Gr.

knappi, Wachsmuth & Springer, 1879, Revis. Paleocrinoidea, p. 93, Ham. Gr.

pisiformis, Roemer, 1860, (Poterocrinus pisiformis,) Sil. Fauna W. Tenn., p. 54, Niagara Gr.

Wachsmuth says it is a Lecanocrinus.

ARCHÆOCIDARIS, McCoy, 1844, Carb. Foss. Ireland, p. 173. [Ety. *archaios*, ancient; *cidaris*, turban.] Spherical; ambulacra narrow, each composed of two ranges of plates, with two pores in each plate; interambulacral plates large, thin, each with a large, perforated, central tubercle, surrounded, at its base, by a smooth ring, and rounded, at the base, for the artic-

FIG. 248. Arachnocrinus pisiformis.

ARCHÆOCIDARIS, McCoy, 1844, Carb. Foss. Ireland, p. 173. [Ety. *archaios*, ancient; *cidaris*, turban.] Spherical; ambulacra narrow, each composed of two ranges of plates, with two pores in each plate; interambulacral plates large, thin, each with a large, perforated, central tubercle, surrounded, at its base, by a smooth ring, and rounded, at the base, for the artic-

ula
wh
for
mo
cat
aut
orn
acule
Aca

FIG. 249. *Cidaris* *longispina*.

No.
edgare
Sur.
Mea
gracill
Ex.
illinois
Sur.
keokul
699,
longisp
Ex.
megast
Loui
Mea
mucron
Acad
Sur.
newbe
Loui
kask
norwo
701,
ornata,
p. 11
shuma
p. 69
spinocl
Geo.
triplex
Foss.
triserr
151,
trudire
p. 17
4, p.
verneul
Loui
occup
acule
worthe
700, s

ulation of a primary spine, and the whole surrounded by smaller tubercles for the articulation of secondary spines; mouth surrounded by numerous imbricating plates; jaws strong with mesial suture; primary spines large, variously ornamented. Type *A. uril*.

aculeata, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 223, Permian Gr.

agnassizi, Hall, 1858, Geo. Rep. Iowa, p. 698, Burlington Gr. *biangulata*, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 224, Coal Meas.

cratis, White, 1876, Geol. of Uinta Mountains, p. 109, and Cont. to Pal., No. 6, p. 130, Lower Aubrey Gr.

dininni, White, 1880, Proc. U. S. Nat. Mus., vol. 2, p. 260, and Cont. to Pal., No. 6, p. 131, Up. Coal Meas.

edgarensis, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 337, Up. Coal Meas.

gracilis, Newberry, 1861, Ives Col. Ex. Ex., p. 117, Up. Carb.

illinoisensis, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 338, St. Louis Gr.

keokuk, Hall, 1858, Geo. Rep. Iowa, p. 699, Keokuk Gr.

longispina, Newberry, 1861, Ives Col. Ex. Ex., p. 116, Up. Carb.

megastylus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 225, Up. Coal Meas.

mucronata, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 395, and Geo. Sur. Ill., vol. 2, p. 295, Kaskaskia Gr.

newberryi, Hambach, 1884, Trans. St. Louis Acad. Sci., vol. 4, p. 548, Kaskaskia Gr.

norwoodi, Hall, 1858, Geo. Rep. Iowa, p. 701, Kaskaskia Gr.

ornata, Newberry, 1861, Ives Col. Ex. Ex., p. 116, Up. Carb.

shumardana, Hall, 1858, Geo. Rep. Iowa, p. 699, Warsaw Gr.

spinoclavata, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 337, Coal Meas.

triplax, White, 1882, Rep. Carb. Invert. Foss. New Mex., p. xxii, Coal Meas.

triserrata, Meek, 1872, Pal. E. Neb., p. 151, Up. Coal Meas.

trudifera, White, 1874, Rep. Invert. Foss., p. 17, and Geo. Sur. W. 100th Mer., vol. 4, p. 104, Carb.

vernuihiana, Swallow, 1858, Trans. St. Louis Acad. Sci. This name was pre-occupied by King. The species is *A. aculeata*.

wortheni, Hall, 1858, Geo. Rep. Iowa, p. 700, St. Louis Gr.

FIG. 249.—*Archæodictya agassizi*, showing spines.

Archæocrinus, Wachsmuth & Springer, 1881, Proc. Acad. Nat. Sci. and Rev. Palæocrinolidea, p. 180. [Ety. *archaios*, ancient; *krinon*, lily.] Basals 5; sub-radials 5; primary radials 3 x 5; secondary radials 3 or 4 x 10; median line of radial plates keeled as in *Glyptocrinus*; interradial areas wide; arms composed of a double series of plates; column round. Type *A. lacunosus*.

desideratus, Billings, 1885, Trans. Ottawa Field Nat. Club, p. 248, Trenton Gr.

lacunosus, Billings, 1857, (*Glyptocrinus lacunosus*.) Rep. of Prog. Geo. Sur. Can., p. 261, and Org. Rem., Decade 4, p. 61, Trenton Gr.

marginatus, Billings, 1857, (*Glyptocrinus marginatus*.) Rep. of Prog. Geo. Sur. Can., p. 260, and Org. Rem., Decade 4, p. 59, Trenton Gr.

microbasalis, Billings, 1857, (*Rhodocrinus microbasalis*.) Rep. of Progr. Geo. Sur. Can., p. 264, and Org. Rem., Decade 4, p. 63, Trenton Gr.

pyriformis, Billings, 1857, (*Rhodocrinus pyriformis*.) Rep. of Prog. Geo. Sur. Can., p. 262, and Org. Rem., Decade 4, p. 61, Trenton Gr.

FIG. 250.—*Archæocrinus sculptus*.

sculptus, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 83 and 117, Trenton Gr.

ARTHROCANTHA, Williams, 1883, Proc. Am. Phil. Soc., p. 84. [Ety. *arthron*, joint; *akantha*, spine.] Calyx bowl-shaped;

plates of body and arms covered with spine-bearing tubercles; basals 3; primary radials 3 x 5, the lower one large, the others small; an azygous interradial as large as the primary radials rests upon the basals, and is followed by numerous small plates; regular interradials small; arms 10, bearing pinules; column round. Type *A. ithacensis*.

carpenteri, Hinde, 1885, (*Hystericrinus carpenteri*.) Ann. and Mag. Nat. Hist., p. 162, Ham. Gr. Probably a syn. for *A. punctobrachiata*.

ithacensis, Williams, 1883, Proc. Am. Phil. Soc., p. 83, Ham. Gr.

punctobrachiata, Williams, 1883, Proc. Am. Phil. Soc., p. 83, Ham. Gr.

ASPIDOCRINUS, Hall, 1859, Pal. N. Y., vol. 3, p. 122. [Ety. *aspis*, shield; *krinon*, lily.] Calyx broadly circular, depressed, hemispheric or scutelliform;

upper margin plain or plicate exteriorly; articulating edges irregular; point for attachment of column small. Type *A. scutelliformis*.

- callosus, Hall, 1859, Pal. N. Y., vol. 3, p. 123, Low. Held. Gr.
 digitatus, Hall, 1859, Pal. N. Y., vol. 3, p. 123, Low. Held. Gr.
 scutelliformis, Hall, 1859, Pal. N. Y., vol. 3, p. 122, Low. Held. Gr.
Asterias, Lamarck, 1815, Hist. Nat. Anim. sans Vert. Not Palæozoic.
anthonyi, see *Palæaster jamesi*.
antiqua, see *Palæaster antiqua*.
antiquata, see *Palæaster antiquata*.
matutina, see *Palæaster matutina*.
Asterocrinus, Lyon, 1857, Geo. Sur. Ky., vol. 3. This name was preoccupied by Münster. See *Pterotocrinus*.
capitatus, see *Pterotocrinus capitalis*.
coronarius, see *Pterotocrinus coronarius*.
Astrios, Troost, 1850, Catalogue. Not defined.
tennesseæ, Troost, 1850. Not defined.

FIG. 251.—*Berycrinus wachsmuthi*. Diagram.

- Astrocrinites*, Conrad in Catalogue Ann. Geo. Rep., 1840-'41. This name was proposed, but not defined; moreover it was preoccupied.
pachydactylus, see *Mariocrinus pachydactylus*.
Astylocrinus, Roemer, 1854, Leth. Geo., p. 229, syn. for *Agassizocrinus*.
levis, syn. for *Agassizocrinus dactyli-formis*.
Ataxocrinus, Lyon, 1869, syn. for *Anomalocrinus*.
caponiformis, see *Anomalocrinus caponiformis*.
Ateleocystites, Briggs, 1858, Can. Org. Rem., Decade 3, p. 72, syn. for *Anomalocystites*.
huxleyi, see *Anomalocystites huxleyi*.
ATELESTOCRINUS, Wachsmuth & Springer, 1886, Rev. Pal., pt. 3, p. 221. [Ety. *ateleitos*, incomplete; *krinon*, lily.] Calyx elongate, bell-shaped, sides concave, restricted along the suture between

- basals and subradials; basals 5; subradials 5; long, narrow, irregular; 3 hexagonal, 2 heptagonal; azygous radial, non-arm bearing, the other four supporting each from 2 to 5 brachials; the eight arms give off branching armlets; azygous plate large, resting upon two subradials; column pentangular Type A. *delicatus*.
delicatus, Wachsmuth & Springer, 1886, Rev. Pal., pt. 3, p. 223, Burlington Gr.
robustus, Wachsmuth & Springer, 1886, Rev. Pal., pt. 3, p. 223, Burlington Gr. base of Keokuk Gr.
Balanocrinus, Troost, 1850. This name was preoccupied. See *Lampterocrinus*.
inflatus, see *Lampterocrinus inflatus*.
BARYCRINUS, Wachsmuth, 1868, Proc. Acad. Nat. Sci., p. 338. [Ety. *barys*, heavy; *krinon*, lily.] Distinguished from *Cyathocrinus* by being more robust, having thicker plates, and a shallower cup; there are usually two azygous plates, while in *Cyathocrinus* there is never more than one; there are never more than two brachials, and these are shorter and wider than in *Cyathocrinus*; the arms are shorter, heavier, and have narrower grooves; the column is stouter; subpentagonal and longitudinally five partite, with a highly organized central canal. Type B. *angulatus*.
angulatus, Meek & Worthen, 1860, (*Cyathocrinus angulatus*), Proc. Acad. Nat. Sci. Phil., p. 391, and Geo. Sur. Ill., vol. 2, p. 234, Keokuk Gr.
bullatus, Hall, 1858, (*Cyathocrinus bullatus*), Geo. Sur. Iowa, p. 624, Keokuk Gr.
cornutus, Owen & Shumard, 1850, (*Cyathocrinus cornutus*), Jour. Acad. Nat. Sci., 2d ser., vol. 2, p. 63, and Geo. Sur. Wis., Iowa, and Minn., p. 591, Burlington Gr.
crassibrachiatus, Hall, 1860, (*Cyathocrinus crassibrachiatus*), Sup. to Geo. Sur. Iowa, p. 60, Keokuk Gr.
geometricus, Meek & Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 485, Keokuk Gr.
herculeus, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 341, and Geo. Sur. Ill., vol. 5, p. 485, Keokuk Gr.
hoveyi, Hall, 1861, (*Cyathocrinus hoveyi*), Desc. New Crin., p. 5, and Geo. Sur. Ill., vol. 5, p. 486, Keokuk Gr.

kello
 log
 Ke
 magi
 may
 Ke
 magn
 Aca
 Sur
 mam
 vol.
 penta
 Ill.
 protu
 pro
 Keo
 syn
 rhomb
 (Po
 Aca
 and
 595,
 sculpt
 scul
 Bur
 solidu
 Desc
 Nat.
 Gr.
 spect
 Aca
 vol.
 spuriu
 rius,
 okuk
 stellat
 latus
 okuk
 striatu
 vol.
 thoma
 Supp
 tumidu
 dus),
 wach
 (Cya
 Nat.
 Ill.,
 BATOCRIN
 Zeits
 and C
 batos,
 bitur
 mary
 2 x 10
 inter
 12 or
 to cu
 radial
 vated
 tral;
 colum
 tinocr
 radial
 numb
 areas
 radial
 of pla
 ning.

- kelloggi, White, 1862, (*Cyathocrinus kelloggi*), Proc. Bost. Soc. Nat. Hist., p. 8, Keokuk Gr.
- magister, Hall, 1858, (*Cyathocrinus magister*), Geo. Sur. Iowa, p. 628, Keokuk Gr.
- magnificus, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 340, and Geo. Sur. Ill., vol. 5, p. 483, Keokuk Gr.
- mammatus, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 486, Keokuk Gr.
- pentagonus, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 487, Keokuk Gr.
- protuberans, Hall, 1858, (*Cyathocrinus protuberans*), Geo. Sur. Iowa, p. 626, Keokuk Gr. Wachsmuth says it is a syn. for *B. bullatus*.
- rhombiferus, Owen & Shumard, 1850, (*Poteriocrinus rhombiferus*) Jour. Acad. Nat. Sci. Phil., 2d ser., vol. 2, and Geo. Wis., Iowa, and Minn., p. 595, Burlington Gr.
- sculptilis, Hall, 1860, (*Cyathocrinus sculptilis*), Supp. Geo. Sur. Iowa, p. 59, Burlington Gr.
- solidus, Hall, 1861, (*Cyathocrinus solidus*), Desc. New Crin., p. 5, and Bost. Jour. Nat. Hist., vol. 7, p. 293, Burlington Gr.
- spectabilis, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil. and Geo. Sur. Ill., vol. 5, p. 530, St. Louis Gr.
- spurius, Hall, 1858, (*Cyathocrinus spurius*), Geo. Sur. Iowa, p. 625, Keokuk Gr.
- stellatus, Hall, 1858, (*Cyathocrinus stellatus*), Geo. Sur. Iowa, p. 623, Keokuk Gr.
- striatus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 515, Keokuk Gr.
- thomae, Hall, 1860, (*Cyathocrinus thomae*), Supp. Geo. Sur. Iowa, p. 61, Warsaw Gr.
- tumidus, Hall, 1858, (*Cyathocrinus tumidus*), Geo. Sur. Iowa, p. 624, Keokuk Gr.
- wachsmuthi, Meek & Worthen, 1861, (*Cyathocrinus wachsmuthi*), Proc. Acad. Nat. Sci. Phil., p. 136, and Geo. Sur. Ill., vol. 3, p. 482, Burlington Gr.
- BATOCRINUS**, Casseday, 1854, Deutsche Zeitschr., d. Geol. Gesellsch., vol. 6, p. 237, and Geo. Sur. Ill., vol. 2, p. 150. [*Ety. batos*, prickly bush; *krinon*, lily.] Calyx biturbinate or globose; basals 3; primary radials 3x5; secondary radials 2x10; tertiary radials 2x2x10; regular interradials 1 to 5; azygous plates 6 to 12 or more; tertiary radials meet so as to cut off the connection of the interradials with the dome plates; vault elevated; tube or proboscis nearly central; arms 18 to 26 or 36 to 40; pinnules; column round, distinguished from *Actinocrinus* by the quadrangular second radial instead of hexagonal; by the number of plates in the interradial areas; by the number of secondary radials; and by having a double series of plates in each arm from the beginning. Type *B. icosidactylus*.
- aequalis*, Hall, 1858, (*Actinocrinus aequalis*), Geo. Rep. Iowa, p. 592, Burlington Gr.
- aequibrachiatus*, McChesney, 1860, (*Actinocrinus aequibrachiatus*), New Pal. Foss., p. 25, and Trans. Chi. Acad. Sci., p. 18, Burlington Gr.
- aequibrachiatus*, var. *alatus*, Hall, 1861, (*Actinocrinus aequibrachiatus* var. *alatus*), Bost. Jour. Nat. Hist., vol. 7, p. 263. Wachsmuth says it is a syn. for *B. aequibrachiatus*.
- andrewsanus*, McCheaney, 1859, (*Actinocrinus andrewsanus*), New Pal. Foss., p. 27, and Trans. Chi. Acad. Sci., p. 20, Burlington Gr.
- asteriscus*, Meek & Worthen, 1860, (*Actinocrinus asteriscus*), Proc. Acad. Nat. Sci. Phil., p. 385, and Geo. Sur. Ill., vol. 2, p. 207, Burlington Gr.
- biturbinatus*, Hall, 1858, (*Actinocrinus biturbinatus*), Geo. Sur. Iowa, p. 616, Keokuk Gr.
- calyculus*, Hall, 1860, (*Actinocrinus calyculus*), Supp. Geo. Sur. Iowa, p. 55, Warsaw Gr.
- calyculus* var. *hardinensis*, Meek & Worthen, 1866, (*Actinocrinus calyculus* var. *hardinensis*), Proc. Acad. Nat. Sci. Phil., p. 253, Warsaw Gr.
- caroli*, Hall, 1860, (*Actinocrinus caroli*), Supp. Geo. Rep. Iowa, p. 54, Warsaw Gr.
- cassedayanus*, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 353, and Geo. Sur. Ill., vol. 5, p. 370, Burlington Gr.
- christyi*, Shumard, 1855, (*Actinocrinus christyi*), Geo. Sur. Mo., p. 191, Burlington Gr.
- clavigerus*, Hall, 1860, (*Actinocrinus clavigerus*), Supp. Geo. Sur. Iowa, p. 44, Burlington Gr. Wachsmuth says it is a syn. for *B. similis*.
- clypeatus*, Hall, 1860, (*Actinocrinus clypeatus*), Supp. Geo. Sur. Iowa, p. 12, and Geo. Sur. Ill., vol. 2, p. 150, Burlington Gr.
- discoideus*, Hall, 1858, (*Actinocrinus discoideus*), Geo. Rep. Iowa, p. 594, Burlington Gr.
- dodecadactylus*, Meek & Worthen, 1861, (*Actinocrinus dodecadactylus*), Proc. Acad. Nat. Sci. Phil., p. 13, and Geo. Sur. Ill., vol. 2, p. 205, Burlington Gr.
- doris*, Hall, 1861, (*Actinocrinus doris*), Desc. New Crinoidea, p. 15, Burlington Gr. Wachsmuth says it is a syn. for *B. aequalis*.
- euconus*, Meek & Worthen, 1860, (*Alloprosallocrinus euconus*), Proc. Acad. Nat. Sci. Phil., p. 164, Warsaw Gr.

FIG. 252. - *Batocrinus dodecadactylus*. Diagram.

formosus, Hall, 1860, (*Actinocrinus formosus*), Supp. to Geo. Sur. Iowa, p. 30, Burlington Gr. Wachsmuth says it is a syn. for *B. discoideus*.

hageri, McChesney, 1860, (*Actinocrinus hageri*), New Pal. Foss., p. 28, and Trans. Chi. Acad. Sci., p. 21, Burlington Gr.

icosidactylus, Casseday, 1854, (*Actinocrinus icosidactylus*), Zeitsch. Deutsch. Geol. Gesellsch., vol. 6, p. 238, Warsaw Gr.

indianensis, Casseday & Lyon, 1859, Am. Jour. Sci. and Arts, 2d ser., vol. 29, p. 75, Keokuk Gr.

inornatus, Hall, 1860, (*Actinocrinus inornatus*), Supp. to Geo. Sur. Iowa, p. 34, Burlington Gr. Wachsmuth says it is a syn. for *B. clypeatus*.

irregularis, Casseday, 1854, Zeitsch. Deutsch. Geol. Gesellsch., vol. 6, p. 238, Warsaw Gr.

lagunculus, Hall, 1860, (*Actinocrinus lagunculus*), Supp. to Geo. Sur. Iowa, p. 41, Warsaw Gr.

laura, Hall, 1861, (*Actinocrinus laura*), Desc. New Crinoidea, p. 15, Burlington Gr.

lepidus, Hall, 1860, (*Actinocrinus lepidus*), Supp. to Geo. Sur. Iowa, p. 32, Burlington Gr.

longirostris, Hall, 1858, (*Actinocrinus longirostris*), Geo. Sur. Iowa, p. 589, Burlington Gr.

lovii, Wachsmuth & Springer, 1881, Proc. Acad. Nat. Sci., p. 342, Burlington Gr.

machridii, Wachsmuth & Springer, (in press), Geo. Sur. Ill., vol. 8, p. 172, Kinderhook Gr.

montgomeryensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 35, and Geo. Sur. Ill., vol. 8, p. 83, Keokuk Gr.

mundulus, Hall, 1860, (*Actinocrinus mundulus*), Supp. to Geo. Sur. Iowa, p. 39, Warsaw Gr.

nashvillae, Troost, Hall, 1858, (*Actinocrinus nashvillae*), Geo. Sur. Iowa, p. 609, Keokuk Gr.

nashvillae var. subtractus, White, 1863, (*Actinocrinus nashvillae* var. subtractus), Proc. Bost. Soc. Nat. Hist., vol. 9, p. 16, Keokuk Gr.

neglectus, see *Eretmocrinus neglectus*.

obolatus, Hall, 1860, (*Actinocrinus obolatus*), Supp. to Geo. Sur. Iowa, p. 38, Burlington Gr. Wachsmuth says it is a syn. for *B. rotundus*.

papillatus, Hall, 1860, (*Actinocrinus papillatus*), Supp. to Geo. Sur. Iowa, p. 29, Burlington Gr. Wachsmuth says it is a syn. for *B. clypeatus*.

pistilliformis, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 153, and Geo. Sur. Ill., vol. 2, p. 151 Waverly or Kinderhook Gr.

pistillum, Meek & Worthen, 1865, (*Actinocrinus pistillum*), Proc. Acad. Nat. Sci. Phil., p. 152, and Geo. Sur. Ill., vol. 3, p. 472, Burlington Gr.

planodiscus, Hall, 1860, (*Actinocrinus planodiscus*), Supp. to Geo. Sur. Iowa, p. 45, Warsaw Gr.

pyriformis, Shumard, 1855, (*Actinocrinus pyriformis*), Geo. Sur. Mo., p. 192, Burlington Gr.

quasillus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 352, and Geo. Sur. Ill., vol. 5, p. 369, Burlington Gr.

rotundus, Yandell & Shumard, 1855, (*Actinocrinus rotundus*), Geo. Sur. Mo., p. 191, Burlington Gr.

similis, Hall, 1860, (*Actinocrinus similis*), Supp. to Geo. Sur. Iowa, p. 40, Keokuk Gr.

sinuosus, Hall, 1860, (*Actinocrinus sinuosus*), Supp. to Geo. Sur. Iowa, p. 26 Burlington Gr.

steropes, Hall, 1860, (*Actinocrinus steropes*), Supp. to Geo. Sur. Iowa, p. 43, Keokuk Gr.

subaequalis, McChesney, 1860, (*Actinocrinus subaequalis*), New Pal. Foss., p. 17, and Trans. Chi. Acad. Sci., p. 13, Burlington Gr. Wachsmuth says it is a syn. for *B. discoideus*.

subconicus, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 26, and Geo. Sur. Ill., vol. 8, p. 84, Keokuk Gr.

trochiscus, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 254, and Geo. Sur. Ill., vol. 5, p. 372, Burlington Gr.

turbinatus, Hall, 1858, (*Actinocrinus turbinatus*), Geo. Rep. Iowa, p. 587, Burlington Gr.

turbinatus var. elegans, (*Actinocrinus turbinatus* var. elegans), Geo. Rep. Iowa, p. 588, Burlington Gr.

unionensis, Worthen, (in press), Geo. Sur. Ill., vol. 8, p. 84, St. Louis Gr.

wachsmuthi, White, 1860, (*Actinocrinus wachsmuthi*), 12th Rep. U. S. Geo. Sur. Terr., p. 162, and 2d Rep. Ind. Geo. Sur., p. 510 Keokuk Gr.

whitii, Wachsmuth & Springer, 1881, Proc. Acad. Nat. Sci., Phil., p. 343, Keokuk Gr.

yandelli, Shumard, 1857, (*Actinocrinus yandelli*), Trans. St. Louis Acad. Sci., vol. 1, p. 76, and Geo. Sur. Ill., vol. 5, p. 341, Keokuk Gr.

FIG. 253.—*Batoerinus machridii*.

FIG. 254.—*Batoerinus rotundus*.

FIG. 255.—*Batoerinus whitii*.

FIG. 256.—

BELEMNOCRINUS, White, 1862, Proc. Bost. Soc.

Nat. Hist., vol. 9, p. 14.

[Ety. *belemnion*, dart, javelin; *krinon*, lily.]

Basals 5, large, long, narrow and of irregular shape; radials 5; arms 10, with one or two syzygies in each joint; pinnules long, arm-like, and bifurcating; azygous plate rests between radials, and upon one of the basals, and supports a large ventral sac composed of numerous hexagonal plates; column pentagonal, with or without lateral cirrhi.

Type B. typus.

florifer, Wachsmuth & Springer, 1877, Am. Jour. Sci. and Arts, 3d ser., vol. 13, p. 256, Burlington Gr.

pourtalesi, Wachsmuth & Springer, 1877, Am. Jour. Sci. and Arts, 3d ser., vol. 13, p. 258, Burlington Gr.

typus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 14, Burlington Gr.

whitii, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 251, and Geo. Sur. Ill., vol. 3, p. 463, Burlington Gr.

FIG. 255.—Belemnocrinus typus.

FIG. 256.—Belemnocrinus whitii. Diagram x 2.

BLASTOIDOCRINUS, Billings, 1859, Can.

Org. Rem., Decade 4, p. 18. [Ety.

blastos, a bud; *eidos*, form; *krinon*, lily.]

The general form is like a *Pentremites*; basals do not rest upon the top of the column, but have their inner edges turned upward, and the column passes on into the visceral

cavity; the deltoids occupy the whole space between the pseudambulacra; the orifices unknown; column round.

Type B. *carcharidens*.

carcharidens, Billings, 1859, Geo. Sur. of Can., Decade 4, p. 18, Chazy Gr.

BRACHIOCRINUS, Hall,

1859, Pal. N. Y.,

vol. 3, p. 118. [Ety.

brachium, an arm;*krinon*, lily.]

Founded upon arms rounded at the base, composed of single articulating plates having thickened, node-like joints, and bearing pinnules. Type B. *nodosarius*.

nodosarius, Hall, 1859, Pal. N. Y., vol. 3, p. 118, Low. Held. Gr.

BURSACRINUS, Meek & Worthen, 1861, Proc.

Acad. Nat. Sci. Phil., p. 136. [Ety.

bursa, purse; *krinon*, lily.] Calyx some-what like *Graphiocrinus*, but arms

widely different; basals 5; subradials 5;

radials 2 x 5; regular interradians 0;

azygous interradians 1; arms wide, flat,

jointing below, in compact series, and

bifurcating above. Type B. *wachsmuthi*.

confirmatus, White, 1862, Proc. Bost. Soc.

Nat. Hist., vol. 9, p. 11, Burlington Gr.

wachsmuthi, Meek & Worthen, 1861, Proc.

Acad. Nat. Sci. Phil., p. 137, and Geo.

Sur. Ill., vol. 3, p. 479, Burlington Gr.

FIG. 257.—Blastoidocrinus carcharidens. d, deltoid plates extending the whole length of the pseudambulacra.

FIG. 258.—Bursacrinus wachsmuthi. Diagram.

Cacabocrinus, Troost, 1850. Never described. The fossils referred to it belong to *Dolatocrinus*.

Calathocrinus, Hall, 1861. The name was preoccupied by Von Meyer in 1848. See *Teliocrinus*.

CALCEOCRINUS, Hall, 1852, Pal. N. Y., vol. 2, p. 352, and 13th Rep. N. Y. St. Mus. Nat. Hist., p. 122. [Ety. *calceus*, shoe; *krinon*, lily.] Base, a single subtriangular or semioval plate, composed of four anchylosed pieces, with cicatrix, for columnar attachment at lower angle; body, above the base, consisting of 5 or 7 plates, of which two are much the larger; a central, elongated plate separates the two large lateral radial plates, and bears an arm; lateral radial plates, each, support brachials that bear bifurcating arms; azygous side arched and composed of 4 or more plates, after which a free arm arises. Type C. *chrysalis*.

articulosus, Billings, 1859, (*Heterocrinus articulosus*.) Can. Org. Rem., Decade 4, p. 51, Trenton Gr.

barrandii, Walcott, 1883, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 212, Trenton Gr.

barrisi, see *Deltacrinus barrisi*.

bradleyi, see *Deltacrinus bradleyi*.

chrysalis, Hall, 1860, (*Cheirocrinus chrysalis*.) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Niagara Gr.

clarus, see *Deltacrinus clarus*.

dactylus, see *Deltacrinus dactylus*.

furcillatus, Billings, 1887, Trans. Ottawa Field Nat. Club, vol. 3, p. 51, Trenton Gr.

inaequalis, Billings, 1859, (*Heterocrinus inaequalis*.) Can. Org. Rem., Decade 4, p. 51, Trenton Gr.

lamellosus, Hall, 1860, (*Cheirocrinus lamellosus*.) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Burlington Gr. Not well defined.

nodosus, see *Deltacrinus nodosus*.

perplexus, Shumard, 1866, (*Cheirocrinus perplexus*.) Trans. St. Louis Acad. Sci., vol. 2, p. 358, Keokuk Gr.

punctatus, Ulrich, 1886, (*Cremacrinus punctatus*.) 14th Rep. Geo. Sur. Minn., p. 107, Trenton Gr.

radicula, Ringueberg, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 120, Niagara Gr.

FIG. 259.—*Calceocrinus rugosus*.

robustus, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 92, Keokuk Gr.

rugosus, Billings, 1887, Trans. Ottawa Field Nat. Club, vol. 3, p. 53, Trenton Gr.

stigmatus, see *Deltacrinus stigmatus*.

tunicatus, Hall, 1860, (*Cheirocrinus tunicatus*.) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 124, Keokuk Gr.

ventricosus, Hall, 1860, (*Cheirocrinus ventricosus*.) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Burlington Gr.

wachsmuthi, see *Deltacrinus wachsmuthi*.

CALLOCYSTITES, Hall, 1852, Pal. N. Y., vol. 2, p. 238. [Ety. *kallos*, beautiful; *kyttis*, bladder.] Ovoid; 1st series of plates 4; 2d series 8; 3d series about the same number; small plates at the apex; arms recumbent, resting in a small shallow groove; pectinated rhombs in three pairs; oral, ovarian and anal apertures. Type C. *jewetti*.

FIG. 260.—*Callocystites jewetti*, Hall, 1852, Pal. N. Y., vol. 2, p. 239, Niagara Gr.

triplectinatus, Ringueberg, 1886, Bull. Bnf. Soc. Nat. Sci., vol. 5, p. 12, Niagara Gr.

CAMAROCINUS, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 205. [Ety. *kamara*, arching chambers; *krinon*, lily.] Body large, unsymmetrical, externally lobed, chambered within and bearing no arms; wall of the dome composed of two layers, the infolding of the inner one forming the partition dividing the chambers; subcircular area in the basal portion composed of spreading, radicleform, bifurcating rays, composed of plates resembling those of a crinoid column, and connected by irregular polygonal plates; ambulacral openings between bifurcations near the outer rim of the area; column cylindrical, internal canal five-rayed. Type C. *stellatus*.

clarki, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 209, Low. Held. Gr.

saffordi, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 208, Low. Held. Gr.

stellatus, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 207, Low. Held. Gr.

Campanulites, Troost, 1850. Not defined.

tesellatus, Troost, 1850. Not defined.

Canistocrinus, Wachsmuth & Springer, 1885, Palaeocrinidae, vol. 1, pt. 3, p. 94. Founded upon *Glyptocrinus richardsoni* and *G. patersoni*, two widely different species. The generic characters are not satisfactorily pointed out.

CARABOCCRINUS, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 275, and Can. Org. Rem., Decade 4, p. 30. [Ety. *karabos*,

FIG. 261.—*Carabocrinus radiatus*. Diagram.

a crab; *krinon*, lily.] Calyx globular or ovoid; basals 5; subradials 5; primary radials 5; arms five, and frequently

divi
inte
bas
groc
mar
C. r
rallat
Geo.
Org.
ton
tuberc
Rem
vancor
Rem
CARYOCR
Sci.,
. *krino*
1st s
series
or le
plater
gous
vanit
conic
or an
ous, a
from
pectin
Type
globosus,
granulit
hexagonu
insculptu
loricatus,
vol. 4,
meconoid

FIG. 262.—*Caryocrinus*
ocystite
CATILICRIN
scribed
Louis A
catillus,
Calyx
bottom;
secondar
arms nu
summit
Type C.
bradleyi,
Acad. N
Sur. Ill.,
tennessee
described

dividing; regular interradials 0; azygous interradials 3, the first one resting on a basal plate; five calycinal, ambulacral grooves on the dome; opening in the margin over the azygous plates. Type *C. radiatus*.

radiatus, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 276, and Can. Org. Rem., Decade 4, p. 31, Trenton Gr.

tuberculatus, Billings, 1859, Can. Org. Rem., Decade 4, p. 33, Hud. Riv. Gr. *vancortlandti*, Billings, 1859, Can. Org. Rem., Decade 4, p. 32, Trenton Gr.

CARYOCYRINUS, Say, 1825, Jour. Acad. Nat. Sci., vol. 4, p. 289. [Ety. *karyon*, a nut; *krinon*, lily.] Body ovoid or subglobose; 1st series of plates 4; 2d series 6; 3d series 6, which bear 9 to 13 arms more or less; vault covered by polygonal plates of moderate size; upon the azygous side, near the outer edge of the vault, 6 triangular plates, forming a conical elevation, represent the mouth or anal orifice; calycine pores numerous, and also in double rows radiating from the center of the body plates; no pectinated rhombs; column round. Type *C. ornatus*.

globosus, Troost, 1850. Not defined.

granulatus, Troost, 1850. Not defined.

hexagonus, Troost. Not defined.

insculptus, Troost. Not defined.

loricatus, Say, 1825, Jour. Acad. Nat. Sci., vol. 4, syn. for *C. ornatus*.

meconoides, Troost. Not defined.

ornatus, Say, 1825, Jour. Acad. Nat. Sci., vol. 4, p. 289, and Pal. N. Y., vol. 2, p. 216, Clinton and Niagara Gr.

Caryocystites, Von Buch, as cited by Hall in 1861, in Geo. Rep. Wis. See Holo-

alternatus, see Holo-

cylindricus, see Holo-

ocystites cylindricus.

CATILICRINUS, Troost, 1850, Cat. Foss. described by Shumard, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 357. [Ety. *catillus*, a small bowl; *krinon*, lily.] Calyx hemispherical concave at the bottom; basals 5; primary radials 1 x 5; secondary radials 1 x 5, very irregular; arms numerous, rising directly from the summit of the radials; column round. Type *C. tennesseæ*.

bradleyi, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 342, and Geo. Sur. Ill., vol. 5, p. 504, Keokuk Gr. *tennesseæ*, Troost, 1850, Catalogue, but described by Shumard, in 1866, in Trans.

St. Louis Acad. Nat. Sci., vol. 2, p. 358, Warsaw Gr.

wachsmuthi, Meek & Worthen, 1866, (Synbathocrinus *wachsmuthi*.) Proc. Acad. Nat. Sci. Phil., p. 251, and Geo. Sur. Ill., vol. 3, p. 465, Burlington Gr.

Centrocrinus, Wachsmuth & Springer, 1881, Proc. Acad. Nat. Sci. Proposed as a subgenus under *Actinocrinus*, to include *A. multicornis* and *A. pentaspinus*, but the name was preoccupied by Austin in 1843.

tennesseensis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 95, Niagara Gr.

Ceriocrinus, White, 1880, proposed as a subgenus of *Erisocrinus*, but the name was preoccupied.

Cheirocrinus, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 122. This name was preoccupied by Eichwald in 1856, and is a syn. for *Calceocrinus*.

chrysalis, see *Calceocrinus chrysalis*.

clarus, see *Deltacrinus clarus*.

dactylus, see *Deltacrinus dactylus*.

lamellosus, see *Calceocrinus lamellosus*.

nodosus, see *Deltacrinus nodosus*.

perplexus, see *Calceocrinus perplexus*.

stigmatus, see *Deltacrinus stigmatus*.

tunicatus, see *Deltacrinus tunicatus*.

ventricosus, see *Calceocrinus ventricosus*.

CHOLASTER, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 328. Ety. *cholos*, defective; *aster*, star.] Body truncated pentagonal; central area circular, large, deep; rays distant, small, short, truncated; centro-dorsal plate large, surrounded by five plates in the position of radials. Type *C. peculiaris*.

peculiaris, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 328, Kaskaskia Gr.

CLEIOCRINUS, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 276, and Can. Org. Rem., Decade 4, p. 52. [Ety. *kleio*, I close; *krinon*, lily.] Calyx conical or pyriform; basals 5; primary radials 3 x 5; secondary radials 4 x 10; tertiary radials numerous; azygous interradials forming a single series from the base to the top of the calyx; regular interradials none; arms numerous and compact. Type *C. regius*.

grandis, Billings, 1860, Can. Org. Rem., Decade 4, p. 54, Trenton Gr.

libanus, Safford, 1869, Geo. of Tenn. Not defined.

FIG. 263.—Catillocrinus wachsmuthi.

FIG. 292.—Caryocrinus ornatus.

magnificus, Billings, 1859, Can. Org. Rem., Decade 4, p. 54, Trenton Gr.
regius, Billings, 1857, Rep. of Prog. Geo. Sur. Can., p. 277, and Can. Org. Rem., Decade 4, p. 53, Trenton Gr.

FIG. 264.—*Cleioocrinus regius*. Diagram.

CLOSTEROCRINUS, Hall, 1852, Pal. N. Y., vol. 2, p. 79. [Ety. *kloster*, a spindle; *krinon*, lily.] Body obconic; basals 3; sub-radials 1x5; number of radials unknown; azygous interradials present; arms composed of a single series of plates; column round. Type *C. elongatus*.

elongatus, Hall, 1852, Pal. N. Y., vol. 2, p. 179, Clinton Gr.

COCCOOCRINUS, Muller, 1855, Verhand. Naturhist. Vereins Rhein und Westph., Jahr. 12, p. 20. [Ety. *kokkos*, a berry; *krinon*, lily.] Basals 3; radials 2x5; interradials 1; column round; distinguished from *Haploocrinus* by the characters of the first radials and the oral plates, and from *Platycrinus* by the character of the vault which is composed of five oral plates resting upon the five interradials, and by the character of the column. Type *C. rosaceus*.

FIG. 265.—*Coccoocrinus bacca*.

bacca, Roemer, 1860, Sil. Fauna West Tenn., p. 57, Niagara Gr.

CODASTER, McCoy, 1849, Ann. & Mag. Nat. Hist., 2d ser., vol. 3, p. 250. [Ety. *kodon*, a bell; *aster*, star.] Calyx inverted conical; summit broad; basals 3, one

tetragonal and two pentagonal, each having its inner apex notched to form part of the round columnar canal; radials 1x5, large, equal, reaching to the truncated summit, to which, from their mesial gibbosity, they give a pentagonal outline; deltoid plates on the summit; mouth central, and from it five prominent ambulacra diverge, one to each angle, each being on a thick tapering ridge, divided by a mesial sulcus; from the re-entering angles of these interradial ridges four other ridges extend to the middle of the four straight sides, the fifth space having no ridge, but, instead, a large ovate opening; hydrospire slits in four interradial areas, but no hydrospire canals, and no pores. Type *C. acutus*.

alternatus, Lyon, 1857, Geo. Sur. Ky., vol. 3, p. 493. A misprint for *C. attenuatus*.
americanus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 239, Up. Held. Gr. Syn. (?) for *C. pyramidatus*.

attenuatus, Lyon, 1857, Geo. Sur. Ky., vol. 3, p. 493–498, Up. Held Gr.

canadensis, Billings. Not defined.

gratiosus, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 257, Keokuk Gr. The hydrospire slits are visible on the casts but are very fine.

hindii, Etheridge & Carpenter, 1882, Ann. and Mag. Nat. Hist., p. 235, Ham. Gr. *C. canadensis*. (?)

kentuckiensis, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 239, Burlington Gr.

pentalobus, see *Stephanocrinus pentalobus*.

FIG. 267.—*Codaster pulchellus*. Summit and side views.

pulchellus, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 35, Niagara Gr. Possibly a *Stephanocrinus*.
pyramidatus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 238, Up. Held. Gr.

FIG. 266.—*Codaster gratiosus*. Summit and side view of cast, the latter showing an aperture at the summit.

FIG. 269.—*Codites form's*.

whitii, Hall, 1861, Desc. New Crinoidea, p. 10, and Bost. Jour. Nat. Hist., vol. 7, p. 237, Burlington Gr.

CODONITES, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 84, and Geo. Sur. Ill., vol. 5, p. 463. [Ety. *kodon*, bell; *lithos*, stone.] Calyx truncate-obpyramidal, elongate below, which distinguishes it, in form, from *Pentremites* and allied genera; deltoid plates constricted in the middle; anal opening large, remote from the center; ambulacra narrow, without marginal pores; side plates large, their apposed edges having pinnule sockets; ten spiracles parallel or subparallel to the ambulacra; the slits are equally developed in all the interradiar areas, while they are absent in the azygous interradius of *Codaster*. Type *C. stelliformis*.

campanulatus, Hambach, 1864, Trans. St. Louis Acad. Sci., vol. 4, p. 548, Burlington Gr.

FIG. 268.—*Codonites conicus*.

conicus, Wachsmuth & Springer, (in press,) (*Orophocrinus conicus*), Geo. Sur. Ill., vol. 8, p. 201, Waverly or Kinderhook Gr.

gracilis, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. and Geo. Sur. Ill., vol. 5, p. 467, Burlington Gr.

FIG. 269.—*Codonites fusi-formis*.

fusi-formis, Wachsmuth & Springer, (in press,) (*Orophocrinus fusi-formis*), Geo. Sur. Ill., vol. 8, p. 203, Waverly or Kinderhook Gr.

stelliformis, Owen & Shumard, 1850. (*Pentremites stelliformis*), Jour. Acad. Nat. Sci. Phil., 2d ser., vol. 2, p. 67, Burlington Gr.

CELOCRINUS, White, 1863, Jour. Bost. Soc. Nat. Hist., vol. 7, p. 499. [Ety. *koilia*, belly; *krinon*, lily.] Distinguished from *Hydreionocrinus* and *Zeacrinus* by its balloon-shaped ventral sac or proboscis, and from the former, also, by the less robust body and comparatively longer arms. Type *C. dilatatus*.

cariniferus, Worthen, 1873, (*Zeacrinus wortheni*), Geo. Sur. Ill., vol. 5, p. 535, St. Louis Gr.

dilatatus, Hall, 1861, (*Poteriocrinus dilatatus*), Desc. New Crinoidea, p. 6, and Bost. Jour. Nat. Hist., p. 300, Burlington Gr.

lyra, Meek & Worthen, 1869, (*Zeacrinus lyra*), Proc. Acad. Nat. Sci., p. 152, and Geo. Sur. Ill., vol. 5, p. 432, Burlington Gr.

subspinosus, White, 1863, Jour. Bost. Soc. Nat. Hist., vol. 7, p. 501, Burlington Gr.

ventricosus, Hall, 1861, (*Poteriocrinus ventricosus*), Desc. New Crinoidea, p. 6, and Bost. Jour. Nat. Hist., p. 301, Burlington Gr.

Celocrinus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., and Geo. Sur. Ill., vol. 2, p. 214, syn. for *Dorycrinus*.

concavus, see *Dorycrinus concavus*.

COMAROCYSTITES, Billings, 1854, Can. Jour., vol. 2, p. 269, and Can. Org. Rem.,

FIG. 270.—*Comarocystites shumardi*. Diagram of part of it.

Decade 3, p. 61. [Ety. *komaron*, strawberry; *kustis*, bladder.] Body ovate; 1st series of plates 3, above which there are from 5 to 11 series, in irregular order; mouth or valvular orifice near the summit; arms free, grooved, bearing pinnules; ambulacral orifice at the apex; column round; all the plates poriferous. Type *C. punctatus*.

- obconicus, Meek & Worthen, 1865, (C. shumardi var. obconicus,) Proc. Acad. Nat. Sci. Phil., p. 144, and Geo. Sur. Ill., vol. 3, p. 294, Trenton Gr.
- punctatus, Billings, 1864, Can. Jour., vol. 2, p. 270, and Can. Org. Rem., Decade 3, p. 61, Trenton Gr.
- shumardi, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 143, and Geo. Sur. Ill., vol. 3, p. 292, Trenton Gr.
- COMPSASTER, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 327. [Ety. *kompos*, elegant; *aster*, star.] Central disk small; rays large, long, fusiform; grooves deep, bordered by numerous adambulacral plates; several rows of disk-plates upon each side of the ambulacral furrows. Type C. formosus.
- formosus, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 327, Kaskaskia Gr.
- COMPSOCRINUS, S. A. Miller, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 233. [Ety. *kompos*, elegant; *krinon*, lily.] Basals 4; primary radials, 3x5; secondary radials 2 or more by 10; tertiary radials more or less numerous; median line of radials keeled; interradials numerous; column four-sided. Type C. harrisi.

FIG. 271.—*Compsocrinus harrisi*. Mag. 2 diam.

resembling *Platycrinus*; basals 3; unequal; radials 3x5; arms single or branching; pinnules long; interradials between the upper edges of the first radials; followed by 3 or 4 more; column cylindrical. Type C. comtus.

FIG. 272.—*Compsocrinus harrisi*. Diagram of basal part and a specimen flattened, natural size.

parvus, Hall, 1861, (*Platycrinus parvus*), Pal. N. Y., vol. 3, p. 114, Low Held. Gr.

harrisi, S. A. Miller, 1881, (*Glyptocrinus harrisi*), Jour. Cin. Soc. Nat. Hist., vol. 4, p. 74, Hud. Riv. Gr.

Conocrinus, Troost. Not defined.

CORDYLOCRINUS, Angelin, 1878, Icon. Crin. Suec., p. 3. [Ety. *kordyle*, a cudgel; *krinon*, lily.] Body

plumosus, Hall, 1861, (*Platycrinus plumosus*), Pal. N. Y., vol. 3, p. 113, Low Held. Gr.

ramulosus, Hall, 1861, (*Platycrinus ramulosus*), Pal. N. Y., vol. 3, p. 115, Low Held. Gr.

CORONOCRINUS, Hall, 1859, Pal. N. Y., vol. 3, p. 124. [Ety. *korone*, a crown; *krinon*, lily.] Founded upon the fragment of the upper part of the calyx showing great breadth, probably hemispheric form, and as many as 40 arm openings in the circumference. Wachsmuth says it is a syn. for *Dolatocrinus*, but as that genus is not known, in rocks, so low as this is found, there is great doubt about the synonymy. Type C. polydactylus.

polydactylus, Hall, 1859, Pal. N. Y., vol. 3, p. 124, Low Held. Gr.

COTYLEDONOCRINUS, Casseday & Lyon, 1860, Proc. Am. Acad. Arts and Sci., vol. 5, p. 26. [Ety. *kotyledon*, any cup-shaped cavity; *krinon*, lily.] Basals 2; radials 3x5; secondary radials 2x10; arms 10; interradials 3x5. Distinguished from *Dichocrinus* by having no azygous plate in line with the first radials, and believed, by Wachsmuth, to have been founded upon an abnormal *Dichocrinus*. Type C. pentalobus.

pentalobus, Casseday & Lyon, 1860, Proc. Am. Acad. Arts and Sci., vol. 5, p. 26, Kaskaskia Gr.

Cremacrinus, Ulrich, syn. for *Calceocrinus punctatus*, see *Calceocrinus punctatus*.

CRINOCYSTITES, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 361. [Ety. *krinon*, lily; *kystis*, bladder.] Elongate, swelling in the upper third of the azygous side, and contracting below the arms; covered by five or more ranges of irregularly disposed plates; central and submarginal apertures. Type C. chrysalis.

chrysalis, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 362. Niagara Gr.

(?) *rectus*, Hall, 1864, see *Rhodocrinus* (?) *rectus*.

Crinosoma antiqua, Castelnau, 1843, Syst. Sil. Probably a fucoid.

Cromyocrinus, Trautschold, 1867, syn. for *Eupachyrcrinus*.

gracilis, see *Eupachyrcrinus gracilis*.

Crumenacrinites, Troost, 1850. Not defined.

ovalis, Troost, 1850. Not defined.

Cryptoblastus, Etheridge & Carpenter, 1886, Catalogue of Blastoidea, p. 229. This genus is founded upon *Granatocrinus melo*, and distinguished from *Granatocrinus* by a slight difference, in the hydrospires. They also referred to it *G. pismum*, and two other species, about which they had very little information.

Ctenocrinus, Bronn, 1840, Leonh. und Bronn. Jahrb., syn. for *Melocrinus*. *bainbridgensis*, see *Melocrinus bainbridgensis*.

breviro-
tus,
Cupellae
Cupuloc-
Pal.
Sev-
l.
Z-n
lute
arm
eroc
a sy
hetero
hete
Tren
CYATHO
Crin
gobl
shap
1 x 5
with
part
irreg
bran
inter
whic
prob
plan
amulu
p. 1
Hist
angula

FIG. 273.

arbores
Acad.
Sur. I
barrisi
Desc.
Nat. I
barydac
1878,
lingto
bulbosus
bullatus
congloba

breviradiatus, see *Melocrinus breviradiatus*.

Capellacrinus, Troost, 1850. Not defined.

Cupulocrinus, D'Orbigny, 1850, Prodr. d. Pal., t. 1, p. 23. Proposed instead of *Scyphocrinus*, Hall, Pal. N. Y., vol. 1, p. 85, that was preoccupied, by Zanker. Basals 5; radials 4 x 5, regular interradials 3; azygous interradials 4; arms 10; column round. Type *C. heterocostalis*. Wachsmuth regards it as a syn. for *Taxocrinus*.

heterocostalis, Hall, 1847. (*Scyphocrinus heterocostalis*.) Pal. N. Y., vol. 1, p. 85, Trenton Gr.

CYATHOCRINUS, Miller, 1821, Nat. Hist. Crinoidea, p. 85. [Ety. *cyathos*, cup or goblet; *crinus*, lily.] Calyx saucer-shaped; basals 5; subradials 5; radials 1 x 5; as large or larger than the basals, with articulating facet occupying only part of the width of a plate; brachials irregular in number; arms long, branching; column round; no regular interradials; azygous interradial 1, which is followed, in the ventral sac or proboscis, by other plates. Type *C. planus*.

æmulus, Hall, 1879, Desc. new spec. foss., p. 10, and 11th Rep. Geo. and Nat. Hist., Indiana, p. 266, Niagara Gr.

angulatus, see *Barycrinus angulatus*.

FIG. 273.—*Cyathocrinus arboreus*. Diagram.

arboreus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 160, and Geo. Sur. Ill., vol. 3, p. 520, Keokuk Gr.

barrisi, Hall, 1861, (*Poteriocrinus barrisi*.) Desc. New Crin., p. 5, and Bost. Jour. Nat. Hist., p. 303, Burlington Gr.

barydactylus, Wachsmuth & Springer, 1878, Proc. Acad. Nat. Sci., p. 257, Burlington Gr.

bulbosus, see *Arachnocrinus bulbosus*.

bullatus, see *Barycrinus bullatus*.

conglobatus, Troost. Not defined.

cora, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 366, Niagara Gr.

cornutus, see *Barycrinus cornutus*.

corrugatus, Troost. Not defined.

crassibrachiatus, see *Barycrinus crassibrachiatus*.

crassus, see *Eupachycrinus crassus*.

crateriformis, Troost. Not defined.

crawfordavillensis, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 79, Keokuk Gr.

decadactylus, Lyon & Casseday, 1860, Am. Jour. Sci. and Arts, vol. 29, p. 73, Keokuk Gr.

depressus, Troost, see *Zeacrinus depressus*.

divaricatus, Hall, 1858, Geo. Sur. Iowa, p. 554, Burlington Gr.

enormis, Meek & Worthen, 1865, (*Poteriocrinus enormis*.) Proc. Acad. Nat. Sci. Phil., p. 152, and Geo. Sur. Ill., vol. 3, p. 481, Burlington Gr.

FIG. 274.—*Cyathocrinus farleyi*. Diagram.

farleyi, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 252, and Geo. Sur. Ill., vol. 3, p. 517, Keokuk Gr.

fasciatus, see *Macrostylocrinus fasciatus*.

floralis, see *Zeacrinus floralis*.

fragilis, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 237, and Geo. Sur. Ill., vol. 5, p. 401, Burlington Gr.

globosus, Troost. Not defined.

gilesi, Wachsmuth & Springer, 1878, Proc. Acad. Nat. Sci., p. 259, Burlington Gr.

granuliferus, Shumard, 1854, Red Riv. Expl. Louisiana, p. 199, Kaskaskia Gr.

hamiltonensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 32, syn. for *C. parvibrachiatus*.

harrisi, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 255, Keokuk Gr. As suggested at the time of describing this species, it may become the type of a new genus.

harrodi, Wachsmuth & Springer, 1879, Proc. Acad. Nat. Sci., p. 87, Keokuk Gr.

hexadactylus, Lyon & Casseday, 1860, Am. Jour. Sci., vol. 29, p. 74, syn. for *Vasocrinus lyoni*.

The name was essentially incorrect and definition wrong.

hoveyi, see *Barycrinus hoveyi*.

inæquidactylus, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 219, Kaskaskia Gr.

incipiens, Hall, 1861, Desc. New Crinoidea, p. 5, and Bost. Jour. Nat. Hist., vol. 7, p. 296, Burlington Gr.

inflatus, Troost. Not defined.

inflexus, see *Eriocrinus inflexus*.

inspiratus, Lyon, 1860, Trans. Am. Phil. Soc., vol. 13, p. 457, Keokuk Gr.

intermedius, Hall, 1858, Geo. Rep. Iowa, p. 627, Keokuk Gr.

iowensis, Owen & Shumard, 1850, Jour. Acad. Nat. Sci., 2d ser., vol. 2, p. 63, and Geo. Sur. Wis., Iowa, and Minn., p. 591, Burlington Gr.

kelloggi, see *Barycrinus kelloggi*.

læviculus, Lyon, 1861, Proc. Acad. Nat. Sci. Phil., p. 409, Up. Held. Gr.

latus, Hall, 1861, Desc. New Crinoidea, p. 5, syn. for *Barycrinus sculptilis*.

lamellosus, White, 1863, Jour. Bost. Soc. Nat. Hist., vol. 7, p. 504, Burlington Gr.

lyoni, see *Vasocrinus lyoni*.

macroleptus, see *Vasocrinus macroleptus*.

magister, see *Barycrinus magister*.

magnoliiformis, see *Zeacrinus magnoliiformis*.

malvaceus, Hall, 1858, Geo. Sur. Iowa, p. 554, Burlington Gr.

Wachsmuth says it is founded on a depressed specimen of *C. iowensis*.

maniformis, see *Zeacrinus maniformis*.

marshallensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 33, and Geo. Sur. Ill., vol. 7, p. 310, Waverly or Kinderhook Gr.

multibrachiatus, Lyon & Casseday, 1859, Am. Jour. Sci., vol. 28, p. 245, Keokuk Gr.

nucleus, Hall, 1876, (Dendrocrinus nucleus,) 28th Rep. N. Y. St. Mus. Nat. Hist., p. 136, Niagara Gr.

ornatissimus, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 247, Portage Gr.

parvibrachiatus, Hall, 1861, Desc. New Crinoidea, p. 6, and Bost. Jour. Nat. Hist., vol. 7, p. 294, Keokuk Gr.

pentelobus, see *Eupachyrinus pentelobus*.

planus, Troost. Not defined.

polyxo, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 199, and 28th Rep. N. Y. St. Mus. Nat. Hist., p. 135, Niagara Gr.

poterium, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 24, and Geo. Sur. Ill., vol. 5, p. 489, Keokuk Gr.

protuberans, see *Barycrinus protuberans*.

pusillus, see *Lecanocrinus pusillus*.

pyriformis, Murchison as identified by Hall. See *Ichthyocrinus lævis*.

FIG. 275.—*Cyathocrinus multibrachiatus*.

quinelobus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 150, and Geo. Sur. Ill., vol. 3, p. 519, Keokuk Gr.

rarus, Lyon, 1869, Trans. Am. Phil. Soc., vol. 13, p. 453, Up. Held. Gr.

rigidus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 8, Burlington Gr.

robustus, Troost. Not defined.

roemeri, Troost. Not defined.

rotundatus, Hall, 1858, Geo. Rep. Iowa, p. 555, Burlington Gr.

saffordi, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 371, and Geo. Sur. Ill., vol. 2, p. 338, Keokuk Gr.

sangamonensis, see *Eupachyrinus sangamonensis*.

scitulus, Meek & Worthen, syn. for *Barycrinus scitulus*.

sculptilis, see *Barycrinus sculptilis*. Not defined.

sculptus, Troost. Not defined.

solidus, see *Barycrinus solidus*.

pentelobus, see *Eupachyrinus pentelobus*.

planus, Troost. Not defined.

polyxo, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 199, and 28th Rep. N. Y. St. Mus. Nat. Hist., p. 135, Niagara Gr.

poterium, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 24, and Geo. Sur. Ill., vol. 5, p. 489, Keokuk Gr.

protuberans, see *Barycrinus protuberans*.

pusillus, see *Lecanocrinus pusillus*.

pyriformis, Murchison as identified by Hall. See *Ichthyocrinus lævis*.

FIG. 276.—*Cyathocrinus quinelobus*.

quinelobus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 150, and Geo. Sur. Ill., vol. 3, p. 519, Keokuk Gr.

rarus, Lyon, 1869, Trans. Am. Phil. Soc., vol. 13, p. 453, Up. Held. Gr.

rigidus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 8, Burlington Gr.

robustus, Troost. Not defined.

roemeri, Troost. Not defined.

rotundatus, Hall, 1858, Geo. Rep. Iowa, p. 555, Burlington Gr.

saffordi, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 371, and Geo. Sur. Ill., vol. 2, p. 338, Keokuk Gr.

sangamonensis, see *Eupachyrinus sangamonensis*.

scitulus, Meek & Worthen, syn. for *Barycrinus scitulus*.

sculptilis, see *Barycrinus sculptilis*. Not defined.

sculptus, Troost. Not defined.

solidus, see *Barycrinus solidus*.

FIG. 277.—*Cyathocrinus saffordi*. Diagram.

saffordi, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 371, and Geo. Sur. Ill., vol. 2, p. 338, Keokuk Gr.

sangamonensis, see *Eupachyrinus sangamonensis*.

scitulus, Meek & Worthen, syn. for *Barycrinus scitulus*.

sculptilis, see *Barycrinus sculptilis*. Not defined.

sculptus, Troost. Not defined.

solidus, see *Barycrinus solidus*.

somern

Acad

spurius

stellatus

stiliati

Mus

No. 6

subtun

Acad

Sur.

tennesse

tenuib

Phil.

Held

tenuide

Proc.

Geo.

ton C

thomæ,

tiaræfor

tumidus

vanho

Soc.

agara

riminal

p. 5,

wachsm

waldron

to Pal

muth

wankon

Mus.

worther

Sci. P

Cyclaster,

This

riost

bigaby,

Cyclocyst

Can. C

kuklos

form. J

disk,

cylind

the ir

ment

chann

with t

which

suppor

elevati

if for t

Type

antecept

St. M

ton G

FIG.

bellulus,

Soc. N

Riv. G

- somersi, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 226, Coal Meas.
- spurius*, see *Barycrinus spurius*.
- stellatus*, see *Barycrinus stellatus*.
- stillativus*, White, 188², Proc. U. S. Nat. Mus., vol. 2, p. 258, and Cont. to Pal. No. 6, p. 125, Up. Coal Meas.
- subtumidus*, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 151, and Geo. Sur. Ill., vol. 5, p. 487, Keokuk Gr.
- tennesseensis*, Troost. Not defined.
- tenuibrachiatus*, Lyon, 1869, Trans. Am. Phil. Soc., vol. 13, p. 460, Up. Held. Gr.
- tenuidactylus*, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 238, and Geo. Sur. Ill., vol. 5, p. 403, Burlington Gr.
- thomæ*, see *Barycrinus thomæ*.
- tiariformis*, see *Ichthyocrinus tiariformis*.
- tumidus*, see *Barycrinus tumidus*.
- vanhornii*, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 261, Niagara Gr.
- viminialis*, Hall, 1861, Desc. New Crin., p. 5, syn. for *C. lowensis*.
- wachsmuthi*, see *Barycrinus wachsmuthi*.
- waldronensis*, Miller & Dyer, 1878, Cont. to Pal. No. 2, p. 6, Niagara Gr. Wachsmuth refers it to *Macrostylocrinus*.
- waukoma*, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 367, Niagara Gr.
- wortheni*, Lyon, 1861, Proc. Acad. Nat. Sci. Phil., p. 410, Up. Held. Gr.
- Cyclaster*, Billings, 1857, Rep. of Progr. This name was preoccupied. See *Edrioaster*.
- bigsbyi*, see *Edrioaster bigsbyi*.
- Cyclocystoides*, Billings & Salter, 1858, Can. Org. Rem., Decade 3, p. 86. [Ety. *kuklos*, circle; *kustis*, bladder; *eidos*, form.] Body consisting of a circular disk, surrounded by a series of short, cylindrical, perforated, porous plates; the interior is covered by an integument of small plates, with radiating channels, which bifurcate and connect with the channel in the marginal series, which makes a complete circle; mouth supposed to be central; mammillary elevations on the exterior of the rim as if for the attachment of small spines. Type *C. halli*.
- anteceptus*, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 219, Trenton Gr.
- bellulus*, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 34, Hud. Riv. Gr.
- halli*, Billings, 1858, Can. Org. Rem., Decade 3, p. 86, Trenton Gr.
- huronensis*, Billings, 1865, Pal. Foss., vol. 1, p. 393, Hud. Riv. Gr.
- magnus*, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 32, and vol. 4, p. 70, Hud. Riv. Gr.
- minus*, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 33, Hud. Riv. Gr.
- mundulus*, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 34, Hud. Riv. Gr.
- nitidus*, Faber, 1886, Jour. Cin. Soc. Nat. Hist., vol. 9, p. 17, Hud. Riv. Gr.
- parvus*, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 33, Hud. Riv. Gr.
- salteri*, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 218, Trenton Gr.
- CYSTOCRINUS*, Roemer, 1860, Sil. Fauna West Tenn., p. 56. [Ety. *kustis*, bladder; *krinon*, lily.] A cylindrical body, the interior of which looks like a crinoid column, but the external part consists of a compact mass of tubes connecting with the central canal. Wachsmuth has called it a detached column, but it is anomalous, and I retain the genus. Type *C. tennesseensis*.
- tennesseensis*, Roemer, 1860, Sil. Fauna West Tenn., p. 56, Niagara Gr.
- Cytocrinus*, Roemer, 1860, Sil. Fauna West Tenn., p. 46, syn. for *Melocrinus*.
- levis*, see *Melocrinus levis*.
- Dæmonocrinites*, Troost. Not defined.
- Decadactylocrinites*, Owen. Not defined.
- Decadoocrinus*, Wachsmuth & Springer, 1879, Proc. Acad. Nat. Sci. Phil. and Revis. Palæocrinoidea, pt. 1, p. 119. It was described as a subgenus of *Poteriocrinus*, but it hardly arises to that dignity. Their type is *Scaphiocrinus scalaris*.
- DELTAOCRINUS*, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 109. [Ety. *delta*, (Greek letter; *krinon*, lily.) Basal piece triangular, composed of anchylosed plates; four plates form the dorsal side above the base; lower central plate triangular and separated from the upper triangular plate by the union of the two lateral radial plates. Distinguished from *Calceocrinus*, which has a long plate on the dorsal side between the lateral radials instead of the two triangular plates separated, as above described by the union of the two radials. Type *D. clarus*.

FIG. 278.—Cyclocystoides magnus.

FIG. 279.—Cyclocrinus tennesseensis.

barrisi, Worthen, 1875, (Calceocrinus *barrisi*, Geo. Sur. Ill., vol. 6, p. 510, Ham. Gr. *bradleyi*, Meek & Worthen, 1869, (Calceocrinus *bradleyi*,) Proc. Acad. Nat. Sci. Phil., p. 73, and Geo. Sur. Ill., vol. 5, p. 502, Keokuk Gr.

FIG. 280.—*Deltacrinus barrisi*.

clarus, Hall, 1862, (Cheirocrinus *clarus*,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 88, Niagara Gr. *dactylus*, Hall, 1860, (Cheirocrinus *dactylus*,) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Burlington Gr. *nodosus*, Hall, 1860, (Cheirocrinus *nodosus*,) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 124, Keokuk Gr. *stigmatus*, Hall, 1863, (Cheirocrinus *stigmatus*,) Trans. Alb. Inst., vol. 4, p. 225, Niagara Gr. *tunicatus*, Hall, 1860, (Cheirocrinus *tunicatus*,) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 124, Keokuk Gr. *wachsmuthi*, Meek & Worthen, 1869, (Calceocrinus *wachsmuthi*,) Proc. Acad. Nat. Sci. Phil., p. 74, and Geo. Sur. Ill., vol. 5, p. 444, Burlington Gr.

FIG. 281.—*Deltacrinus stigmatus*. Dorsal view of calyx enlarged 2 diam.

DENDROCRINUS, Hall, 1852, Pal. N. Y., vol. 2, p. 193. [Ety. *dendron*, tree; *krinos*, lily.] Calyx obconoidal; basals 5; subradials 5; radials 1x5, and an additional one caused by a division of the plate on the left side of the large azygous one; regular interradials 0; azygous interradial 1; large and long proboscis or ventral sac rises from the azygous interradial; arms long, branching; ambula-

FIG. 282.—*Dendrocrinus*. Diagram.

crural furrow deep; pinnules wanting; column round or pentagonal; without base or roots for attachment. Type *D. longidactylus*.

acutidactylus, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 266, and Can. Org. Rem., Decade 4, p. 37, Trenton Gr. *ancilla*, Hall, 1879, Desc. New Spec. Foss., p. 9, and 11th Rep. Geo. and Nat. Hist. Indiana, p. 271, Niagara Gr. *alternatus*, Hall, 1847, (Poteriocrinus *alternatus*,) Pal. N. Y., vol. 1, p. 83, Trenton Gr.

angulatus, see *Palaeocrinus angulatus*. *angustatus*, Meek & Worthen, 1870, (Homocrinus *angustatus*,) Proc. Acad. Nat. Sci. Phil., p. 30, and Geo. Sur. Ill., vol. 6, p. 492, Hud. Riv. Gr.

caduceus, Hall, 1860, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 208, Hud. Riv. Gr. *cassii*, Meek, 1871, Am. Jour. Sci. and Arts, 3d ser., vol. 2, p. 295, and Ohio Pal., vol. 1, p. 28, Hud. Riv. Gr. *cincinnatiensis*, Meek, 1872, Proc. Acad. Nat. Sci. Phil., p. 312, and Ohio Pal., vol. 1, p. 20, Hud. Riv. Gr. *conjugans*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 268, and Can. Org. Rem., Decade 4, p. 41, Trenton Gr. *celsus*, Ringueberg, 1888, Proc. Acad. Nat. Sci. Phil., p. 132, Niagara Gr. *curtus*, see *Merocrinus curtus*. *cylindricus*, Billings, 1859, Can. Org. Rem., Decade 4, p. 44, Trenton Gr. *dyeri*, Meek, 1872, Proc. Acad. Nat. Sci. Phil., p. 310, and Ohio, Pal., vol. 1, p. 24, Hud. Riv. Gr. *erraticus*, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 316, Hud. Riv. Gr. *gracilis*, Hall, 1847, (Poteriocrinus *gracilis*,) Pal. N. Y., vol. 1, p. 84, Trenton Gr. *gregarius*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 265, and Can. Org. Rem., Decade 4, p. 38, Trenton Gr. *humilis*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 265, and Can. Org. Rem., Decade 4, p. 39, Trenton Gr. *jewetti*, Billings, 1859, Can. Org. Rem., Decade 4, p. 43, Trenton Gr. *latibrachiatulus*, Billings, 1857, Rep. Progr. Can. Geo. Sur., p. 270, and Can. Org. Rem., Decade 4, p. 39, Hud. Riv. Gr. *longidactylus*, Hall, 1852, Pal. N. Y., vol. 2, p. 193, Niagara Gr. *modestus*, Safford. Not defined. *navigiolium*, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 235, Utica Slate Gr. *nucleus*, see *Cyathocrinus nucleus*.

FIG. 283.—*Dendrocrinus jewetti*.

FIG. 284.—*Dendrocrinus oswegoensis*. Diagram.

oswegoensis, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 333, Hud. Riv. Gr.

- polydactylus, Shumard, 1857, (Homo-
crinus polydactylus.) Trans. St. Louis
Acad. Sci., vol. 1, p. 78, and Ohio Pal.,
vol. 1, p. 22, Hud. Riv. Gr.
- posticus, Hall, 1872, (Pterotocrinus postic-
us.) 24th Rep. N. Y. St. Mus. Nat.
Hist., p. 209, and Ohio Pal., vol. 1, p.
22, Hud. Riv. Gr.
- proboscidiatus, Billings, 1857, Rep. of
Progr. Can. Geo. Sur., p. 267, and Can.
Org. Rem., Decade 4, p. 38, Tren-
ton Gr.
- retractilis, Walcott, 1883, 35th Rep. N. Y.
St. Mus. Nat. Hist., p. 211, Tren-
ton Gr.
- rusticus, Billings, 1857, Rep. of Progr. Geo.
Sur. Can., p. 270, and Can. Org. Rem.,
Decade 4, p. 41, Trenton Gr.
- similis, Billings, 1857, Rep. of Progr. Geo.
Sur. Can., p. 267, and Can. Org. Rem.,
Decade 4, p. 40, Trenton Gr.
- tener, Billings, 1860, Catal. Sil. Foss.
Antic., p. 9, Hud. Riv. Gr.
- Dicnocrinus, Munster, 1839, Beitrag. Zur.
Petrif., vol. 1, p. 2. [Ety. *dicha*,
in two parts; *krinon*, lily.] Calyx deep,
cup-shaped; plates delicate; basals 2;
primary radials 5, large, resting two
upon each basal, and the other in a
notch at one end of the basal suture,
opposite which there is a large azygous
plate in line with the first radials; suc-
ceeding radials 1 to 3, in each ray, the
last supporting arms; arms 10, long,
bifurcating and bearing stout pinnules;
interradials 4 or 5, small, situate
above the first radials; vault slightly
elevated, with a small opening upon
the azygous side; column round. Type
D. radiatus.
- angustus, White, 1862, Proc. Bost. Soc.
Nat. Hist., vol. 9, p. 19, Burling-
ton Gr.
- chesterensis, see Pterotocrinus chesterensis.
- constrictus, Meek & Worthen, 1860,
Proc. Acad. Nat. Sci. Phil., p. 381, and
Geo. Sur. Ill., vol. 2, p. 263, War-
saw Gr.
- conus, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 381, and Geo.
Sur. Ill., vol. 2, p. 169, Burlington Gr.
- cornigerus, see Talarocrinus cornigerus.
- coraxus, Worthen, 1882, Bull. No. 1, Ill.
St. Mus. Nat. Hist., p. 35, and Geo. Sur.
Ill., vol. 7, p. 313, Keokuk Gr.
- crassitestus, White, 1862, Proc. Bost. Soc.
Nat. Hist., vol. 9, p. 19, Burlington Gr.
- crassus, see Pterotocrinus crassus.
- dichotomus, Hall, 1860, Supp. to Geo.
Sur. Iowa, p. 85, Warsaw Gr.
- elegans, Casseday & Lyon, see Talarocrinus
elegans.
- expansus, Meek & Worthen, 1868. The
name was preoccupied by DeKoninck
& LeHon, but the name is probably a
synonym for *D. polydactylus*.
- ficus, Casseday & Lyon, 1860, Proc. Am.
Acad. Arts and Sci., vol. 5, p. 24, and
Geo. Sur. Ill., vol. 5, p. 502, Keokuk Gr.

inornatus, Wachsmuth & Springer, (in
press.) Geo. Sur. Ill.,
vol. 8, p. 190, Waverly
or Kinderhook Gr.

hamiltonensis, Worthen,
1882, Bull. No. 1, Ill.
St. Mus. Nat. Hist., p.
35, and Geo. Sur. Ill.,
vol. 7, p. 313, Keokuk
Gr.

lachrymosus, Hall, 1860,
Supp. to Geo. Sur.
Iowa, p. 84, Burlington
Gr. Wachsmuth says
it is a syn. for *Platy-*

crinus subspinulosus.

levis, Hall, 1860, Supp.
Geo. Sur. Iowa, p. 83,
Burlington Gr.

lineatus, Meek & Worth-
en, 1869, Proc. Acad.
Nat. Sci. Phil., p. 69,
and Geo. Sur. Ill., vol. 5, p. 440, Burl-
ington Gr.

liratus, Hall, 1861, Desc. New Crinoidea,
p. 5, and Jour. Bost. Nat. Hist., vol. 7,
p. 200, Burlington Gr.

ornatus, Wachsmuth & Springer, 1881,
Proc. Acad. Nat. Sci. Phil. and Revis.
Paleocrin., p. 84, Keokuk Gr. This
name was proposed instead of *D. sculptus*,
Casseday & Lyon, because the latter
was preoccupied.

ovatus, Owen & Shumard, 1850, Jour.
Acad. Nat. Sci., 2d ser., vol. 2, p. 61,
and Geo. Sur. Iowa, Wis., and Minn., p.
590, Burlington Gr.

pisum, Meek & Worthen, 1869, Proc.
Acad. Nat. Sci. Phil., p. 69, and Geo.
Sur. Ill., vol. 5, p. 441, Burlington Gr.

plicatus, Hall, 1861, Desc. New Crinoidea,
p. 4, and Jour. Bost. Soc. Nat. Hist.,
vol. 7, p. 288, Burlington Gr.

poecilum, Hall, 1861, Desc. New Crinoidea,
p. 5, and Jour. Bost. Soc. Nat. Hist.,
vol. 7, p. 291, Burlington Gr.

polydactylus, Casseday & Lyon, 1860,
Proc. Am. Acad. Arts and Sci., vol. 5,
p. 20, Keokuk Gr.

protuberans, Hall, see Pterotocrinus pro-
tuberans.

scitulus, Hall, 1861, Desc. New Crinoidea,
p. 4, and Jour. Bost. Soc. Nat. Hist.,
vol. 7, p. 289, Burlington Gr.

sculptus, Casseday & Lyon, 1860, Proc.
Am. Acad. Arts and Sci., vol. 5, p. 25.
The name was preoccupied by DeKoninck
& LeHon in 1853. See *D. or-*
natus.

sexlobatus, see Talarocrinus sexlobatus.

simplex, Shumard, 1857, Trans. St. Louis
Acad. Sci., p. 74, and Geo. Sur. Iowa,
p. 654, Warsaw Gr.

striatus, Owen & Shumard, 1850, Jour.
Acad. Nat. Sci., 2d ser., vol. 2, and Geo.
Sur. Iowa, Wis., and Minn., p. 590,
Burlington Gr.

symmetricus, see Talarocrinus symmet-
ricus.

pendro-
wettli.

ur. Cin.
5, Utica

Diagram.

68, Geo.
v. Gr.

Dictyocrinus, Conrad, 1841, (*Dictyocrinites*), Ann. Rep. N. Y. and Pal. N. Y., vol. 3, p. 135, syn. for *Receptaculites*.
squamifer, Hall, see *Receptaculites squamifer*.

DOLATOCRINUS, Lyon, 1857, Geo. Sur. Ky., vol. 3, p. 482. [Ety. *dolatus*, hewn or tooled; *krinon*, lily.] Body subaphe-roidal; calyx basin-shaped; vault hemispherical, depressed in the interradial areas; basals anchylosed and probably numbering 5; radials 3x5; secondary radials 2x10; sometimes tertiary radials 2x20; arms 20 to 40, bifurcating and bearing pinnules; interradials 5 or more, the first one large; apertures sub-central; column round. Type *D. lacus canadensis*, Whiteaves, 1887, Cont. to Can. Pal. vol. 1, p. 99, Ham. Gr.
glyptus, Hall, 1862, (*Cacabocrinus glyptus*), 15th Rep. N. Y. St. Mus. Nat. Hist., p. 140, Ham. Gr.

FIG. 236.—*Dolatocrinus lacus*. Side view.

FIG. 237.—*Dolatocrinus lacus*. Ventral view.

glyptus var. *intermedius*, Hall, 1862, (*Cacabocrinus glyptus* var. *intermedius*), 15th Rep. N. Y. St. Mus. Nat. Hist., p. 141, Ham. Gr.

FIG. 238.—*Dolatocrinus lacus*. Diagram of a ray.

lacus, Lyon, 1857, Geo. Sur. Ky., vol. 3, p. 482, Up. Held. Gr.

lamellosus, Hall, 1862, (*Cacabocrinus lamellosus*), 15th Rep. N. Y. St. Mus. Nat. Hist., p. 141, Up. Held. Gr.

liratus, Hall, 1862, (*Cacabocrinus liratus*), 15th Rep. N. Y. St. Mus. Nat. Hist., p. 139, Ham. Gr.

liratus var. *multilira*, Hall, 1862, (*Cacabocrinus liratus* var. *multilira*), 15th Rep. N. Y. St. Mus. Nat.

Hist., p. 139, Ham. Gr.
marshi, Lyon, 1869, Trans. Am. Phil. Soc., vol. 13, p. 461, Up. Held. Gr.
ornatus, Meek, 1871, Proc. Acad. Nat. Sci., p. 57, Up. Held. Gr.
speciosus, Hall, 1862, (*Cacabocrinus speciosus*), 15th Rep. N. Y. St. Mus. Nat. Hist., p. 137, Up. Held. Gr.
triadactylus, Barris, 1885, Proc. Dav. Acad. Sci., vol. 4, p. 100, Ham. Gr.
troosti, Hall, 1862, (*Cacabocrinus troosti*), 15th Rep. N. Y. St. Mus. Nat. Hist., p. 138, Ham. Gr.

Donacirinites, Troost. Not defined.

simplex, Troost. Not defined.

DORYCRINUS, Roemer, 1854, Archiv. f. Naturgesch. Jahrg. 19, p. 207. [Ety. *dory*, spear; *krinon*, lily.] Body turbinate or subglobose, truncate at the base, depressed in the interradial spaces so as to make it pentalobate; dome convex and usually bearing from 1 to 6 spines; basals 3; primary radials 3x5; secondary radials 2x2; or where there are tertiary radials, there are only 1x2 secondaries; arms 24 to 40; interradials 2 or 3, in two series; azygous area very different from the interradial areas, and having several more plates and an aperture near the top directed laterally; readily distinguished from *Batocrinus* and *Eretmocrinus* by the lobed form of the body, by the azygous area and lateral opening, and by the shortness of the arms. Type *D. mississippiensis*.

canaliculatus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci., p. 166, and Geo. Sur. Ill., vol. 5, p. 381, Burlington Gr.

concaus, Meek & Worthen, 1861, (*Actinocrinus concaus*), Proc. Acad. Nat. Sci. Phil., p. 131, and Geo. Sur. Ill., vol. 2, p. 215, Low. Burlington Gr.

cornigerus, Hall, 1858, (*Actinocrinus cornigerus*), Geo. Rep. Iowa, p. 576, Burlington Gr.

gouldi, Hall, 1858, (*Actinocrinus gouldi*), Geo. Rep. Iowa, p. 613, Keokuk Gr.

FIG. 239.—*Dorycrinus concaus*. Diagram.

FIG. 240.—*Dorycrinus immaturus*. Posterior and anterior views.

immaturus, Wachsmuth & Springer, (in press.) Geo. Sur. Ill., vol. 8, p. 175, Waverly or Kinderhook Gr.
kelloggi, Worthen, 1875, Geo. Sur. Ill. vol. 6, p. 513, Keokuk Gr.

lineatu
Nat.
ton
missis
fur
kuk
missis
Supp
kuk
missou
nus
190,
parvib
pres
derh

FIG. 291.—I

parvus,
vus,
lingto
penden
dene
Burli
præcar
carsoi
Hist.,
quinque
quinq
p. 15,

FIG. 292.—I

quinque
Worth
346, a
Burlin
radiatus
press,
derhod
roemeri,
Acad.
Geo. S
ton Gr
spinulos
spinul
52, Ke

lineatus, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 310, Burlington Gr.
 mississippiensis, Roemer, 1853, Archiv. fur Nat. Jahr. 19, p. 207, Keokuk Gr.
 mississippiensis, var. spiniger, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 53, Keokuk Gr.
 missouriensis, Shumard, 1858, (Actinocrinus missouriensis,) Geo. Rep. Mo., p. 190, Burlington Gr.
 parvibasis, Wachsmuth & Springer, (in press,) Geo. Sur. Ill., vol. 8, p. 177, Kinderhook Gr.

FIG. 291.—*Dorycrinus parvibasis*. Anterior, posterior, and ventral views.

parvus, Shumard, 1855, (Actinocrinus parvus,) Geo. Sur. Mo., p. 193, Upper Burlington Gr.
 pendens, Hall, 1860, (Actinocrinus pendens,) Supp. to Geo. Sur. Iowa, p. 31, Burlington Gr.
 precursor, Hall, 1862, (Actinocrinus precursor,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 131, Ham. Gr.
 quinquelobus, Hall, 1860, (Actinocrinus quinquelobus,) Supp. to Geo. Rep. Iowa, p. 15, Burlington Gr.

FIG. 292.—*Dorycrinus radiatus*. Posterior and anterior views.

quinquelobus var. intermedius, Meek & Worthen, 1868, Proc. Acad. Nat. Sci., p. 346, and Geo. Sur. Ill., vol. 5, p. 385, Burlington Gr.
 radiatus, Wachsmuth & Springer, (in press,) Geo. Sur. Ill., vol. 8, p. 176, Kinderhook Gr.
 roemer, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 346, and Geo. Sur. Ill., vol. 5, p. 383, Burlington Gr.
 spinulosus, Hall, 1860, (Actinocrinus spinulosus,) Supp. Geo. Sur. Iowa, p. 52, Keokuk Gr.

subaculeatus, Hall, 1858, (Actinocrinus subaculeatus,) Geo. Rep. Iowa, p. 570, Burlington Gr.

subturbatus, Meek & Worthen, 1860, (Actinocrinus subturbatus,) Proc. Acad. Nat. Sci. Phil., p. 388, and Geo. Sur. Ill., vol. 2, p. 212, Burlington Gr.

FIG. 293.—*Dorycrinus subturbatus*. Diagram.

symmetricus, Hall, 1858, (Actinocrinus symmetricus,) Geo. Sur. Iowa, p. 574, Burlington Gr.
 trinodus, Hall, 1858, (Actinocrinus trinodus,) Geo. Sur. Iowa, p. 575, Burlington Gr.

FIG. 294.—*Dorycrinus unicornis*.

unicornis, Owen & Shumard, 1850, (Actinocrinus unicornis,) Jour. Acad. Nat. Sci. Phil., vol. 2, new ser., p. 67, and Geo. Sur. Ill., vol. 5, p. 380, Burlington Gr.

unispinus, Hall, 1861, (Actinocrinus unispinus,) Desc. New Crinoidea, p. 2, and Bost. Jour. Nat. Hist., vol. 7, p. 270, Burlington Gr.

Echinocystites, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 360. The name was preoccupied by Wyville Thompson. See *Lysocystites*.

nodosus, see *Lysocystites nodosus*.

Echinodiscus, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 335. [Éty. *echinos*, sea urchin; *diskos*, quait.] Body discoid; depressed convex, larger plates in the center of the interradial areas, none imbricating; narrow elongate plates form the border and pass to the under side forming a non-sessile rim; ambulacra 5, connected near the center, and composed of numerous interlocking plates; mouth central or subcentral. Type *E. optatus*.

kaskaskiensis, Hall, 1858, (Agelacrinus kaskaskiensis,) Geo. Sur. Iowa, p. 696, Kaskaskia Gr.

optatus, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 336, Kaskaskia Gr.

Echino-encrinites, Meyer, 1826, Karst. Archiv. Nat., vol. 7. [Éty. *echinos*, sea urchin; *krinon*, lily.]

anatiformis, see *Glyptocystites anatiformis*.

feneistratus, Troost. Not defined.

Echinus drydenensis, see *Eocidaris drydenensis*.

gyracanthus, see *Tentaculites gyracanthus*.

ECTENOCRINUS, n. gen. [Ety. *ekteino*, I stretch out; *krinon*, lily.] General form very elongate; calyx small, subcylindrical, moderately expanding; basals 5, unequal; radials irregular, four plates in three series, before the bifurcation of the free arms, and three in each of the other two series; arms 10, long; pinnules strong; azygous plates 3, following each other, but not in a direct line; vault unknown; column very long, round, tripartite, and attaching by an expanded base. Type *E. simplex*. This genus is founded upon *Heterocrinus simplex*, Hall, as the type, because the genus *Heterocrinus* was founded upon *H. heterodactylus*, as

FIG. 205.
Ectenocrinus grandis.

the type, which is quite widely removed from *H. simplex*.

canadensis, Billings, 1859, (*Heterocrinus canadensis*,) Can. Org. Rem., Decade 4, p. 48, Trenton Gr.

grandis, Meek, 1873, (*Heterocrinus simplex* var. *grandis*,) Pal. Ohio, vol. 1, pl. 1, fig. 206.—*Ectenocrinus* fig. 7, Hud. simplex. Diagram.

simplex, Hall, 1847, (*Heterocrinus simplex*,) Pal. N. Y., vol. 1, p. 280, Trenton and Hud. Riv. Gr.

EDRIOASTER, Billings, 1858, Can. Org. Rem., Decade 3, p. 82. [Ety. *edrion*, seat; *aster*, star.] A substitute for *Cyclaster*, proposed in 1857, the latter name having been preoccupied. Body sessile, discoid; plates numerous, irregular, polygonal; ambulacral grooves 5, tapering, composed of two series of oblong ossicles, with four rows of ambulacral pores in each; mouth large, formed of five oral and five internal ossicles. Type *E. bigsbyi*.

bigsbyi, Billings, 1857, (*Cyclaster bigsbyi*,) Rep. of Progr. Geo. Sur. Can., p. 293, and Can. Org. Rem., Decade 3, p. 82, Trenton Gr.

EDRIOCRINUS, Hall, 1859, Pal. N. Y., vol. 3, p. 119. [Ety. *edrion*, seat; *krinon*, lily.] Body obconic; base solid, without column; radials 5, resting in depressions, in the base; azygous plates 2, one large, resting in a basal depres-

sion, the other smaller and succeeding the first; arms composed of transversely linear plates and bifurcating. Type *E. pocilliformis*.

pocilliformis, Hall, 1859, Pal. N. Y., vol. 3, p. 121, Low. Held. Gr.

pyriformis, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 88, Up. Held. Gr.

sacculus, Hall, 1859, Pal. N. Y., vol. 3, p. 143, Oriskany sandstone.

Elæocrinus, Roemer, 1852, syn. for *Nucleocrinus*.

kirkwoodensis, see *Nucleocrinus kirkwoodensis*.

verneuili, see *Nucleocrinus verneuili*.

ELEUTHEROCRINUS, Shumard & Yandell, 1856, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 73. [Ety. *eleutheros*, free; *krinon*, lily.] Calyx subelliptical, resembling *Nucleocrinus* in form but depressed on the azygous side; truncated at the summit and bulged on one side; subtriangular at the base and prolonged on one of its sides; basals 3, one small, two irregular and much elongated; radials 1x5, four-forked, occupying nearly the length of the calyx, one short and not forked; interradials 1x5; pseudambulacral areas 5, four linear, extending nearly the entire length of the calyx, one short, subtriangular, situated on the summit plane; apertures 8 (?). Type *E. casedayi*.

casedayi, Shumard & Yandell, 1856, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 74, Up. Held. Gr.

whitfieldi, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., App. C, p. 123, Ham. Gr.

ECODIARIS, Desor, 1858, Synopsis des Echinodermes Fossiles. [Ety. *eos*, dawn; *diarion*, day.] Plates hexagonal; one large tubercle on each plate, smooth at the base and perforated at the summit; distinguished from *Archæodiaris* by the absence of a second ring. Type *E.*

drydenensis, Vanuxem, 1842, (*Echinus drydenensis*,) Geo. Rep. 3d Dist. N. Y., p. 184, and 20th Rep. N. Y. St. Mus. Nat. Hist., p. 343, Chemung Gr.

hallanus, Geinitz, 1866, Carb. and Dyas. in Neb., p. 61, and Pal. E. Neb., p. 152, Up. Coal Meas.

squamosus, see *Lepidocrinus squamosus*.

EOCYSTITES, Billings, 1868, Acad. Geol., p. 643. [Ety. *eos*, dawn; *kustis*, bladder.] Plates numerous, varying in size, form, and ornamentation, usually radiately sculptured. Type *E. primævus*.

longidactylus, Walcott, 1886, Bull. U. S. Geo. Sur., No. 30, p. 94, Upper Taconic.

primævus, Billings, 1868, Acad. Geol., p. 643, Up. Taconic, St. John's Gr.

ERETMOCRINUS, Lyon & Caseday, 1859, Am. Jour. Sci. and Arts, vol. 28, p. 241. [Ety. *eretmos*, oar; *krinon*, lily.] Body bitur-

FIG. 207.
Eocystites primævus.

binat
the c
wide
openi
pound
portio
intern
boscis
the i
Type
adultus
Proc.
Keok
attenua
tuta v
p. 14,
calyculo
calycu
p. 17,
carica,
Desc.
clio, Ha
New C
Nat.
ton G
clælia, H
Desc.
Jour.
ton Gr
corbulis,
bulis, H
Jour.
ton Gr
coronatu
natus, H
Burlin
gemmafo
gemmi
p. 23, I

FIG. 208.
Eretmocrinus kontinckii.

seday,
vol. 28,
matuta, H
Desc. N
ton Gr
matuta var
neglectus
crinus
Sci., p.
p. 377,
originariu
Proc. A
ramulosu
ulosus,
kuk Gr.

binate or subglobose, vault exceeding the calyx in size; basals 3, forming a wide rim; primary radials 3x5; arm openings 12 to 22; arms simple or compound, long, flattened in the upper portions; interradials 1 to 3; azygous interradials, 8 or more; tube or proboscis excentric and extending beyond the infolding arms; column round. Type *E. magnificus*.

adultus, Wachsmuth & Springer, 1881, Proc. Acad. Nat. Sci. Phil., p. 349, Keokuk Gr.

attenuatus, Hall, 1861, (*Actinocrinus matuta* var. *attenuatus*.) Desc. New Crin., p. 14, Burlington Gr.

calyculoides, Hall, 1860, (*Actinocrinus calyculoides*.) Supp. to Geo. Sur. Iowa, p. 17, Burlington Gr.

carica, Hall, 1861, (*Actinocrinus carica*.) Desc. New Crin., p. 10, Burlington Gr.

clio, Hall, 1861, (*Actinocrinus clio*.) Desc. New Crinoidea, p. 1, and Bost. Jour. Nat. Hist., vol. 7, p. 262, Burlington Gr.

clœlia, Hall, 1861, (*Actinocrinus clœlia*.) Desc. New Crinoidea, p. 1, and Bost. Jour. Nat. Hist., vol. 7, p. 266, Burlington Gr.

corbulis, Hall, 1861, (*Actinocrinus corbulis*.) Desc. New Crin., p. 1, and Bost. Jour. Nat. Hist., vol. 7, p. 265, Burlington Gr.

coronatus, Hall, 1860, (*Actinocrinus coronatus*.) Supp. Geo. Sur. Iowa, p. 28, Burlington Gr.

gemmiformis, Hall, 1860, (*Actinocrinus gemmiformis*.) Supp. Geo. Sur. Iowa, p. 23, Burlington Gr.

FIG. 298.
Eretmocrinus konineki.

intermedius, Wachsmuth & Springer, 1881, Proc. Acad. Nat. Sci., p. 348, Keokuk Gr.

konineki, Shumard, 1855, (*Actinocrinus konineki*.) Geo. Sur. Mo., p. 194, Burlington Gr.

leucosia, Hall, 1861, (*Actinocrinus leucosia*.) Desc. New Crin., p. 1, and Bost. Jour. Nat. Hist., vol. 7, p. 261, Burlington Gr.

magnificus, Lyon & Caseday, 1859, Am. Jour. Sci. and Arts, vol. 28, p. 241, Keokuk Gr.

matuta, Hall, 1861, (*Actinocrinus matuta*.) Desc. New Crinoidea, p. 14, Burlington Gr.

matuta var. *attenuata*, see *E. attenuatus*.
neglectus, Meek & Worthen, 1869, (*Batocrinus neglectus*.) Proc. Acad. Nat. Sci., p. 355, and Geo. Sur. Ill., vol. 5, p. 377, Burlington Gr.

originarius, Wachsmuth & Springer, 1881, Proc. Acad. Nat. Sci., p. 348, Keokuk Gr.

ramulosus, Hall, 1858, (*Actinocrinus ramulosus*.) Geo. Sur. Iowa, p. 615, Keokuk Gr.

remibrachiatus, Hall, 1861, (*Actinocrinus remibrachiatus*.) Desc. New Crinoidea, p. 11, Burlington Gr.

varsouviensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 30, and Geo. Sur. Ill., vol. 7, p. 306, Warsaw Gr.
verneuianus, Shumard, 1855, (*Actinocrinus verneuianus*.) Geo. Sur. Mo., p. 193, Burlington Gr.

FIG. 299.
Eretmocrinus verneuianus.

ERISOCRINUS, Meek & Worthen, 1865, Am. Jour. Sci. and Arts, vol. 89, p. 174. [Ety. *eris*, contention; *krinon*, lily.] Calyx saucer-shaped; basals 5, small; subradials 5, large; radials 2x5, large; no interradials; arms 10; column round. Type *E. typus*.

antiquus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 71, and Geo. Sur. Ill., vol. 5, p. 447, Burlington Gr.

cognatus, Wachsmuth & Springer, 1887, Note to p. 255, Revis. Palæocrinoidea, syn. for *E. planus*.

conoides, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 150, and Geo. Sur. Ill., vol. 2, p. 318, Up. Coal Meas.

inflexus, Geinitz, 1866, (*Cyathocrinus inflexus*.) Carb. und Dyas, in Neb., p. 62, and White's Cont. to Pal., No. 6, p. 128, Coal Meas.

nebraskensis, Meek & Worthen, 1865, Am. Jour. Sci., vol. 89, p. 174, Up. Coal Meas. Regarded a variety of *E. typus*.
planus, White, 1880, Proc. U. S. Nat. Mus., vol. 2, p. 257, and Cont. to Pal., No. 6, p. 127, Coal Meas.

FIG. 301.—*Erisocrinus typus*. Two side views, basal view, and top view of calyx.

typus, Meek & Worthen, 1865, Am. Jour. Sci. and Arts, vol. 89, p. 174, and Geo. Sur. Ill., vol. 2, p. 319, Up. Coal Meas.
tuberculatus, see *Eupachycrinus tuberculatus*.

whitii, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 72, and Geo. Sur. Ill., vol. 5, p. 448, Burlington Gr.
EUCALYPTOCRINUS, Goldfuss, 1826, Petref. Germ., p. 212. [Ety. *eu*, well; *kalyptos*, covered; *krinon*, lily.] Body turbinate

or bowl-shaped from base to arms, and with arms and interbrachial plates subovate or subelliptical; basals 4, concealed in basal cavity and developed in the interior; primary radials 3 x 5, the first large; secondary radials 2 x 10; interradials 3, one very large; arms 20, composed of a double series of plates, which fill the interbrachial spaces; interbrachial plates solid, extending from the interradial plates as high as the arms reach, and uniting at the summit; proboscis or tube extending to the top, and sometimes far beyond; column round; attaching by branching roots. Type *E. rosaceus*.

armosus, see *Siphonocrinus armosus*.

cælatus, Hall, 1843, (*Hypanthocrinites cælatus*), *Geo. Rep.* 4th Dist. N. Y., p. 113, and 28th Rep. N. Y. St. Mus. Nat. Hist., p. 142, Niagara Gr.

chicagoensis, Winchell & Marcy, 1865, *Mem. Bos. Soc. Nat. Hist.*, p. 90, Niagara Gr.

conicus, Troost. Not defined.

constrictus, Hall, 1879, *Trans. Alb. Inst.*, vol. 10 (Abstract, p. 10), and 11th Rep. Geol. and Nat. Hist. Ind., p. 273, Niagara Gr.

cornutus, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 363, Niagara Gr.

cornutus var. *excavatus*, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 364, Niagara Gr.

FIG. 302.—*Eucalyptocrinus crassus*.

crassus, Hall, 1863, *Trans. Alb. Inst.*, vol. 4, p. 197, and 28th Rep. N. Y. St. Mus. Nat. Hist., p. 141, Niagara Gr.

decorus, Phillips, 1839, (*Hypanthocrinites decorus*), *Murch. Sil. Syst.*, p. 672, and *Pal. N. Y.*, vol. 2, p. 207, Niagara Gr.

depressus, S. A. Miller, 1880, *Jour. Cin. Soc. Nat. Hist.*, vol. 3, p. 232, Niagara Gr.

egani, S. A. Miller, 1880, *Jour. Cin. Soc. Nat. Hist.*, vol. 3, p. 140, Niagara Gr.

extensus, Troost. Not defined.

gibbosus, Troost. Not defined.

goldfussi, Troost. Not defined.

inconspicuous, Ringueberg, 1884, *Proc. Acad. Nat. Sci.*, p. 148. Not properly defined.

lævis, Troost. Not defined.

magnus, Worthen, 1875, *Geo. Sur.* III, vol. 6, p. 501, Niagara Gr.

nashvillæ, Troost. Not defined.

obconicus, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 365, Niagara Gr.

ornatus, Hall, 1861, *Rep. of Progr. Geo. Sur. of Wis.*, p. 20, Niagara Gr.

ovalis, Troost, as figured by Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 143, Niagara Gr.

papulosus, Hall, 1862, *Pal. N. Y.*, vol. 2, p. 211, Niagara Gr.

phillipsi, Troost. Not defined.

proboscidalis, S. A. Miller, 1882, *Jour. Cin. Soc. Nat. Hist.*, vol. 5, p. 224, Niagara Gr.

ramifer, Roemer, 1860, *Sil. Fauna West Tenn.*, p. 51, Niagara Gr.

rotundus, S. A. Miller, 1882, *Jour. Cin. Soc. Nat. Hist.*, vol. 5, p. 82, Niagara Gr.

splendidus, Troost, *Catal. Hall & Whitfield*, 1875, *Ohio Pal.*, vol. 2, p. 128, Niagara Gr.

tennesseæ, Troost. Not defined.

tuberculatus, Miller & Dyer, 1878, *Jour. Cin. Soc. Nat. Hist.*, vol. 1, p. 36, Niagara Gr.

turbinatus, S. A. Miller, 1882, *Jour. Cin. Soc. Nat. Hist.*, vol. 5, p. 82, Niagara Gr.

EUCALYPTOCRINUS, Meek, 1871, *U. S. Geo. Sur. Terr.*, p. 373. [Ety. *eu*, very; *kados*, branch; *krinon*, lily.] Calyx like *Platycrinus*, and distinguished by having the radial series extended in the form of tubular free rays, which bear arms, alternately, on either side, throughout their length; arms composed of a double series of interlocking plates. Type *E. montanensis*.

millebrachiatus, Wachsmuth & Springer, 1878, *Proc. Acad. Nat. Sci. Phil.*, p. 245, Burlington and Keokuk Gr.

montanensis, Meek, 1871, *Hayden's Rep. U. S. Geo. Sur. Terr.*, p. 373, Subcarboniferous.

pleurovinnus, White, 1862, (*Platycrinus pleurovinnus*), *Proc. Bost. Soc. Nat. Hist.*, vol. 9, p. 17, Up. Burlington Gr.

EUGASTER, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 332. [Ety. *euge*, pre-eminent, remarkable; *aster*, star.] A central, alated disk, with five long, slender flexuous rays; disk composed on the ventral side of small polygonal

FIG. 303.—*Eucalyptocrinus tuberculatus*.

plates series bulacra cral pl two ro and dis logani. concinnu Soc. N ara Gr. logani, H Mus. N EUPACHYCR Proc. A [Ety. e lily.] plates h defined dials 2 third r the sec bearing 5 or 10, of plate round. alis. asperatus, Ill. St. Geo. Sur kaskia basseti, W vol. 8, p boydi, M Acad. N Geo. Sur kaskia craigi, W S. p. 527 crassus, M crinus c Phil. p. p. 314, L

Fig. 304.—*Eup* fayettensis, vol. 5, p. 6 formosus, Kaskaskia germanus, S Soc. Nat. 1 kia Gr.

plates; rays consisting of a double series of alternating, subquadrate, ambulacral ossicles, with curved ambulacral plates; oral plates 10; pores large, two rows in each ray; adambulacral and disk-plates spine-bearing. Type E. logani.

concinus, Ringueberg, 1886, Bull. Buff. Soc. Nat. Sci., vol. 5, p. 8, Niagara Gr.

logani, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 333, Ham. Gr.

EUPACHYCRINUS, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 159. [Ety. *eu*, very; *pachys*, thick; *krinon*, lily.] Calyx saucer or bowl shaped; plates heavy, tumid; sutures strongly defined; basals 5; subradials 5; radials 2 x 5, and sometimes there are third radials in some of the rays; the second radials are often spine-bearing; azygous interradials 3; arms 5 or 10, composed of a double series of plates, bearing pinnules; column round. Type E. quatuordecembrachialis.

asperatus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 34, and Geo. Sur. Ill., vol. 7, p. 311, Kaskaskia Gr.

bassetti, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 528, Coal Meas.

boydi, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 30, and Geo. Sur. Ill., vol. 5, p. 554, Kaskaskia Gr.

craigi, Worthen, 1875, Geo. Sur. Ill., vol. 3, p. 527, Coal Meas.

crassus, Meek & Worthen, 1870, (Cyathocrinus crassus,) Proc. Acad. Nat. Sci. Phil., p. 392, and Geo. Sur. Ill., vol. 2, p. 314, Low. Coal Meas.

FIG. 304.—*Eupachyrcrinus crassus*. Diagram.

fayettensis, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 565, Up. Coal Meas.

formosus, Worthen, 1873, (Zeacrinus formosus,) Geo. Sur. Ill., vol. 5, p. 549, Kaskaskia Gr.

germanus, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 40, Kaskaskia Gr.

gracilis, Wetherby, 1880, (Cromyocrinus gracilis,) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 248, Kaskaskia Gr.

hemisphericus, Shumard, 1858, (Poteriocrinus hemisphericus,) Trans. St. Louis, Acad. Sci., vol. 1, p. 221, and Geo. Sur. Ill., vol. 5, p. 561, Coal Meas.

FIG. 305.—*Eupachyrcrinus sangamonensis*. Diagram.

monroensis, Worthen, 1882, Bull. No. 1, St. Mus. Nat. Hist., p. 30, Kaskaskia Gr.

orbicularis, Hall, 1861, (Scaphiocrinus orbicularis,) Bost. Jour. Nat. Hist., p. 311, Keokuk Gr.

peralobus, Hall, 1858, (Cyathocrinus peralobus,) Geo. Sur. Iowa, p. 687, Kaskaskia Gr.

platybasis, White, 1876, Geo. Uinta Mountains, p. 108, and Cont. to Pal., No. 6, p. 124, Low. Aubrey Gr.

quatuordecembrachialis, Lyon, 1857, (Graphiocrinus quatuordecembrachialis,) Geo. Sur. Ky., vol. 3, p. 477, Kaskaskia Gr.

sanctiludovici, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 98, St. Louis Gr.

sangamonensis, Meek & Worthen, 1861, (Cyathocrinus sangamonensis,) Proc. Acad. Nat. Sci. Phil., p. 392, and Geo. Sur. Ill., vol. 2, p. 310, Up. Coal Meas.

spartarius, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 38, Kaskaskia Gr.

subtumidus, Worthen, 1867, (Zeacrinus subtumidus,) Geo. Sur. Ill., vol. 5, p. 548, Kaskaskia Gr.

tuberculatus, Meek & Worthen, 1865, (Erisocrinus tuberculatus,) Proc. Acad. Nat. Sci. Phil., p. 150, and Geo. Sur. Ill., vol. 2, p. 319, Coal Meas.

verrucosus, White & St. John, 1869, Trans. Chi. Acad. Sci., p. 117, Coal Meas.

FIG. 306.—Euspirocrinus obconicus.

covered with large plates; arms bifurcate. Type E. spiralis.

obconicus, W. R. Billings, 1885, Ottawa Field Nat. Club, vol. 2, p. 248, Trenton Gr.

FIG. 307.—Euspirocrinus obconicus. Diagram.

FORBESOCRINUS, DeKoninck & LeHon, 1854, Resch. Crin. Carb. Belg., p. 118. [Ety. proper name; *krinon*, lily.] Calyx

large, plates heavy; basals 3; subradials 5; primary radials 3 or 4 x 5; secondary radials 2 to 4 x 10; tertiary radials 2 to 4 x 20; arms 50 to 60, long and sometimes dividing; regular interradials 10 to 20 or more; azygous interradials numerous; interaxillaries 10 to 20 or more. Type F. nobilis.

agassizi, Hall, 1858 and 1860, Geo. Sur. of Iowa, p. 631, Burlington Gr.

agassizi var. giganteus, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil. p. 131, and Geo. Sur. Ill., vol. 3, p. 495, Burlington Gr.

asteriformis, see Onychocrinus asteriformis.

cestriensis, Hall, 1860, Supp. to Geo. Iowa, p. 68, Kaskaskia Gr.

communis, see Taxocrinus communis.

giddingsi, see Taxocrinus giddingsi.

juvenis, see Taxocrinus juvenis.

kelloggi, see Taxocrinus kelloggi.

lobatus, see Taxocrinus lobatus.

lobatus var. tardus, see Taxocrinus lobatus var. tardus.

meeki, see Taxocrinus meeki.

monroensis, see Onychocrinus monroensis.

multibrachiatus, see Taxocrinus multibrachiatus.

norwoodi, see Onychocrinus norwoodi.

nuntius, see Taxocrinus nuntius.

parvus, Wetherby, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 138, Kaskaskia Gr.

FIG. 308.—Forbesocrinus wortheni.

pratteni, see Melocrinus pratteni.

ramulosus, Lyon & Casseday, see Onychocrinus ramulosus.

ramulosus, Hall, see Taxocrinus ramulosus.

saffordi, see Taxocrinus saffordi.

semiovatus, see Taxocrinus semiovatus.

shumardanus, see Taxocrinus shumardanus.

spiniger, see Taxocrinus spiniger.

subramulosus, Shumard, 1866, syn. for Taxocrinus ramulosus.

thiemii, see Taxocrinus thiemii.

whitfieldi, see Taxocrinus whitfieldi.

wortheni, Hall, 1858, Geo. Rep. Iowa, p. 632, Keokuk Gr.

FIG. 309.—Gaurocrinus neulli. Diagram.

GAUROCRINUS, S. A. Miller, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 228. [Ety. *gauros*, haughty; proud; *krinon*, lily.]

Calyx
depr
ary
prim
post
10 to
ous
by a
plate
whic
the
beari
angula
crinu
Hist.
cognatu

FIG. 310.—G
neu

GENNAEOCRINUS
1882, P
gennaei
Body
vault
3 x 5;
radials
numer
the se
Actino
Type G
calypso, P
15th R
133, H
cassedayi
sedayi,
410, U
cauliculus
caulicu
Nat. Hi
cornigeru
(Actino
Sci., vol
encharis,
charis.)
Hist., p
kentuckiens
crinus
Acad.
nigerus.

Calyx having strong radial ridges and depressed interradial and intersecondary radial areas; basals 5; subradials 5; primary radials 3x5, or the left posterior ray only 2; secondary radials 10 to 16x10; interradial plates numerous and small; azygous area supported by a ridge up the middle series of plates; vault covered by small plates, which are continued as a covering over the arm furrows; arms 20 or more, bearing pinnules. Type *G. nealli*.

angularis, Miller & Dyer, 1878, (Glyptocrinus *angularis*,) Jour. Cin. Soc. Nat. Hist., vol. 1, p. 28, Hud. Riv. Gr.

cognatus, S. A. Miller, 1881, (Glyptocrinus *cognatus*,) Jour. Cin. Soc. Nat. Hist., vol. 4, p. 75, Hud. Riv. Gr.

magnificus, S. A. Miller, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 230, Hud. Riv. Gr.

nealli, Hall, 1866, (Glyptocrinus *nealli*,) Adv. Sheets 24th Rep. N. Y. St. Mus. Nat. Hist., p. 206, and Ohio Pal., vol. 1, p. 34, Hud. Riv. Gr.

splendens, S. A. Miller, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 230, Trenton Gr.

FIG. 310.—*Gaurocrinus nealli*.

GENNÆOCRINUS, Wachsmuth & Springer, 1882, Proc. Acad. Nat. Sci., p. 334. [Ety. *gennaios*, of noble birth; *krinon*, lily.] Body wider than high, lobed, striated; vault low; basals 3; primary radials 3x5; secondary radials 1x10; interradials 5 to 7; azygous interradials more numerous, and having three plates in the second row instead of two, as in *Actinocrinus*; interaxillaries 1 to 3. Type *G. cornigerus*.

calypso, Hall, 1862, (Actinocrinus *calypso*,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 133, Ham. Gr.

cassedayi, Lyon, 1861, (Actinocrinus *cassedayi*,) Proc. Acad. Nat. Sci. Phil., p. 410, Up. Held. Gr.

cauliculus, Hall, 1862, (Actinocrinus *cauliculus*,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 132, Ham. Gr.

cornigerus, Lyon & Casseday, 1859, (Actinocrinus *cornigerus*,) Am. Jour. Sci., vol. 28, p. 238, Ham. Gr.

eucharis, Hall, 1862, (Actinocrinus *eucharis*,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 130, Ham. Gr.

kentuckiensis, Shumard, 1860, (Actinocrinus *kentuckiensis*,) Trans. St. Louis Acad. Sci., p. 345, syn. for *G. cornigerus*.

nyssa, Hall, 1862, (Actinocrinus *nyssa*,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 129, Ham. Gr.

pocillum, Hall, 1862, (Actinocrinus *pocillum*,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 134, Ham. Gr.

GLYPTASTER, Hall, 1852, Pal. N. Y., vol. 2, p. 187. [Ety. *glyptos*, sculptured; *aster*, star.] Calyx obconical, depressed between the arm bases, radial portion ridged; basals 5; subradials 5; primary radials 3x5; secondary radials 2 or more by 10; interradials 6 or more; azygous interradials more numerous; arms 10, composed of double series of plates. Type *G. brachiatus*.

brachiatus, Hall, 1852, Pal. N. Y., vol. 2, p. 187, Niagara Gr.

FIG. 311.—*Glyptaster egani*. Natural size and enlarged.

egani, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 261, Niagara Gr.

inornatus, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 205, and 28th Rep. N. Y. St. Mus. Nat. Hist., p. 134, Niagara Gr.

occidentalis, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 204, and 28th Rep. N. Y. St. Mus. Nat. Hist., p. 134, Niagara Gr.

occidentalis var. *crebescens*, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 133, Niagara Gr.

pentangularis, Hall, 1867, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 369, Niagara Gr.

GLYPTOCRINUS, Hall, 1847, Pal. N. Y., vol. 1, p. 280. [Ety. *glyptos*, sculptured;

FIG. 313.—*Glyptocrinus decadactylus*. Part of vault mag. 6 diam. showing excurrent opening.

krinon, lily.] Calyx obconoidal, interradial areas flattened or depressed; surface sculptured and having radial ridges; basals 5; primary radials 3x5;

secondary radials 1 or more by 10; tertiary radials usually present; arms 10 to 20 or more, bearing pinnules; first interradial resting upon the first

FIG. 314.—Glyptocrinus decadactylus. Diagram.

primary radials, and followed by succeeding ranges of two or more, which graduate into the vault; vault slightly convex, with sunken interradial areas; plates becoming smaller as they approach the inner face of the arms, and becoming a somewhat granular continuous cover over the ambulacral furrows; excurrent opening subcentral on the upper face of the vault; column round, without base or roots for attachment.

Type *G. decadactylus*.

angularis, see *Gaurocrinus angularis*.

FIG. 318.—Glyptocrinus forni-shelli.

A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 103, Hud. Riv. Gr.

argutus, Walcott, 1883, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 207, Trenton Gr.

armosus, see *Siphonocrinus armosus*.

baeri, see *Xenocrinus baeri*.

carleyi, see *Mariocrinus carleyi*.

cognatus, see *Gaurocrinus cognatus*.

decadactylus, Hall, 1847, Pal. N. Y., vol. 1, p. 281, Hud. Riv. Gr.

dryeri, Meek, 1872, Proc. Acad. Nat. Sci. Phil., p. 314, and Ohio Pal., vol. 1, p. 32, Hud. Riv. Gr.

dryeri var. *sublaevis*, S.

FIG. 315.—Glyptocrinus decadactylus.

fimbriatus, Shumard, 1855, Geo. Sur. Mo., p. 194, Trenton Gr.

fornshelli, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 348, Hud. Riv. Gr.

gracilis, Wetherby, syn. for *Gaurocrinus angularis*.

harrisi, see *Compsocrinus harrisi*.

lacunosus, see *Archæocrinus lacunosus*.

libanus, Safford, 1869, Geo. of Tenn.

Not defined.

marginatus, see *Archæocrinus marginatus*.

miamiensis, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 34, Hud. Riv. Gr.

nealli, see *Gaurocrinus nealli*.

nobilis, see *Siphonocrinus nobilis*.

ornatus, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 260, and Can. Org. Rem., Decade 4, p. 60, Trenton Gr.

parvus, Hall, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 207, Hud. Riv. Gr.

pattersoni, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 80, Utica Slate Gr.

plumosus, Hall, 1843, (Actinocrinus plumosus,) Geo. Rep. 4th Dist. N. Y., p. 72, and Pal. N. Y., vol. 2, p. 180, Clinton Gr. Founded upon fragments too poor for even generic determination.

priscus, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 257, and Can. Org. Rem., Decade 4, p. 56, Black Riv. and Trenton Grs.

quinquepartitus, Billings, 1859, Can. Org. Rem., Decade 4, pl. 8, fig. 4a, 4b, Trenton Gr.

ramulosus, Billings, 1856, Can. Nat. Geo., vol. 1, and Can. Org. Rem., Decade 4, p. 57, Trenton Gr.

richardsoni, Wetherby, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 245, Hud. Riv. Gr.

sculptus, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 37, Hud. Riv. Gr.

shafferi, see *Pycnocrinus shafferi*.

shafferi var. *germanus*, see *Pycnocrinus germanus*.

siphonatus, Hall, 1861, syn. for *Siphonocrinus armosus*.

subglobosus, Meek, 1873, (G. *dryeri* var. *subglobosus*,) Pal. Ohio, vol. 1, p. 34, Hud. Riv. Gr.

subnodosus, see *Rhaphanocrinus subnodosus*.

GLYPTOCYSTITES, Billings, 1854, Can. Jour., vol. 2, p. 215, and Can. Org. Rem., Decade 3, p. 53. [Ety. *glyptos*, sculptured; *kustis*, bladder.] Body elongate, cylindrical; four series of plates, 4 in the basal and 5 in each succeeding series; mouth in one of the plates of the second series; ambulacral orifice at the center of the summit where it receives the five ambulacral grooves; arms recumbent upon the apex of the fossil,

FIG. 318.

below, ranged upper sessile, ral orif of graa clavus, H Nat. H glans, Ha Nat. H

and grooves beset with small plates; 10 to 13 pectinated rhombs; column short, tapering to a point. Type *G. multiporus*. *anatifomis*, Hall, 1847, (*Echinoecrinites anatifomis*), Pal. N. Y., vol. 1, p. 80, Trenton Gr.

forbesi, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 283, and Can. Org. Rem., Decade 3, p. 59, Chazy Gr.

logani, Billings, 1857, Rep. of Progress, Geo. Sur. Can., p. 282, and Can. Org. Rem., Decade 3, p. 57, Trenton Gr.

logani var. *gracilis*, Billings, 1858, Can. Org. Rem., Decade 3, p. 59, Trenton Gr. *multiporus*, Billings, 1854, Can. Jour., vol. 2, p. 215, and Can. Org. Rem., Decade 3, p. 54, Trenton Gr.

GOMPHOCYSTITES, Hall, 1860, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 351, [Ety. *gomphos*, nail or rudder; *kustis*, bladder.] Elongate pyriform, narrow

indianensis n. sp., Niagara Gr. Upper part elliptical in outline and regularly convex; five ambulacral grooves curve spirally outward from an ambulacral orifice within the groove near the mouth, and extend below the summit; mouth round, situate between two of the ambulacral grooves; each ambulacral groove has a suture in the bottom of it, but there is no other evidence of the subdivision of the top into plates; even the mouth appears as a hole through a solid test; whole surface tuberculated, and each tubercle pierced with a pair of pores. Collected by J. F. Hammell in Jefferson County, Indiana.

tenax, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 352, Niagara Gr.

GONIASTEROIDOCRINUS, Lyon & Casseday, 1859, Am. Jour. Sci., vol. 28, 2d series, p. 233. [Ety. like the recent genus *Goniaster*; *krinon*, lily.] Body short, cylindrical, or subglobose; basals 5; subradials 5, often protuberant; primary radials 3 x 5; secondary radials 2 to 4 x 10; arms numerous, delicate, pend-

FIG. 319.—*Gomphocystites indianensis*. Summit view.

FIG. 318.—*Goniasteroidocrinus fascellus*. Diagram 2 diam.

below, inflated above; plates spirally arranged; ambulacral orifice central on the upper surface; mouth excentric; arms sessile, and curving from the ambulacral orifice outward to or below the point of greatest diameter. Type *G. glans*. *clavus*, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 353, Niagara Gr. *glans*, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 352, Niagara Gr.

ent, supporting pinnules; interradials 10 to 18 in each area; vault depressed and extending in five or six pseudo-brachial appendages star-like, which bifurcate, then spread, curve, and terminate each in a point; these brachial appendages separate the interradial areas from the dome; excurrent orifice sublateral, not protruding. Type *S. tuberosus*. There are some who use *Ollacrinus* as the

FIG. 320.—Diagrammatic view of *Gilbertocrinus bursa* to show it is distinct from *Goniasteroidocrinus*.

generic name, but it was not defined or established; there are others who use *Gilbertocrinus*, but it, probably, is a distinct genus, and, so far, not known in America.

fiscellus, Meek & Worthen, 1861, (*Trematocrinus fiscellus*.) Proc. Acad. Nat. Sci. Phil., p. 383, and Geo. Sur. Ill., vol. 2, p. 222, Burlington Gr.

FIG. 321.—*Gonlasteroideocrinus tuberosus*. A is the vault; B, the opening; C, side view of vault; D, under side of false arms; E, enlargement of base of arms.

FIG. 322.—Three views of *Gilbertocrinus calcaratus*, to show the genus is distinct from *Gonlasteroideocrinus*.

obovatus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci., Phil., p. 76, and Geo. Sur. Ill., vol. 5, p. 391, Burlington Gr.
papillatus, Hall, 1860, (*Trematocrinus papillatus*.) Supp. to Geo. Rep. Iowa, p. 76, Burlington Gr.
reticulatus, Hall, 1861, (*Trematocrinus reticulatus*.) Desc. New Crinoidea, p. 9, and Bost. Jour. Nat. Hist. vol. 7, p. 325, Burlington Gr.

robustus, Hall, 1860, (*Trematocrinus robustus*.) Supp. to Geo. Rep. Iowa, p. 77, Keokuk Gr.

spinigerus, Hall, 1862, (*Trematocrinus spinigerus*.) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 128, Ham. Gr.

tenulradiatus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 75, and Geo. Sur. Ill., vol. 5, p. 389, Burlington Gr.

tuberculosis, Hall, 1860, (*Trematocrinus tuberculosis*.) Supp. to Geo. Rep. Iowa, p. 75, Burlington Gr.

tuberosus, Lyon & Casseday, 1859, Am. Jour. Sci., vol. 28, 2d ser., p. 233, Kaskaskia Gr.

typus, Hall, 1860, (*Trematocrinus typus*.) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 73, Burlington Gr.

GRANATOCRINUS, Troost, 1850, Cat. Foss. in Am. Jour. Sci., vol. 8, p. 420, and described by Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 146. [Ety. *granatus*, granular; *krinos*, lily.] Calyx subglobose oval or elliptic; the proportions of the plates giving a very different outline to the calyx from that of a *Pentremites*; summit depressed convex; base flattened or concave; ambulacra areas like those in *Pentremites*, but narrower, and extending nearly or quite the entire length; basals 3, sunken so as not to be visible in a side view; radials and deltoids similar to those of *Pentremites* and proportionally as variable; slender, thread-like arms, or pinnules, as in *Pentremites*; ambulacra and lancet-plates in narrow sinuses; anal opening as in *Pentremites*; central opening and spiracles often closed by small plates; ten narrow hydrospiral canals open externally by either five or ten apertures. Type *G. norwoodi*.

cidariformis, Troost. Not defined.
cornutus, Meek & Worthen, 1861, (*Pentremites cornutus*.) Proc. Acad. Nat. Sci. Phil., p. 141, and Geo. Sur. Ill., vol. 2, p. 276, St. Louis Gr.

curtus, Shumard, 1855, (*Pentremites curtus*.) Geo. Rep. Mo., p. 187, Warsaw Gr.

glaber, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 91, and Geo. Sur. Ill., vol. 5, p. 537, St. Louis Gr.

granulatus, Roemer, 1852, (*Pentatrematites granulatus*.) Monog. Blast., p. 43, Warsaw Gr.

granulosus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 165, and Geo. Sur. Ill., vol. 5, p. 508, Keokuk Gr.

leda, Hall, 1862, (*Pentremites leda*.) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 149, Ham. Gr.

lotobla
Foss.
Mer.

melo,
mites
Phil.,
Gr.
speci
tobla
genus
ters,
melo var
jectus
melonia
missour
Louis
neglect
Acad.
Sur. I
norwood
tremi
Sci. P

FIG. 324.—
projecto

Phil.,
496, B
roemer,
meri,
sayi, see
shumard
Acad.

Sur. II
GRAPHIOCH
Rech.

[Ety.
krinos,
this g
radials
muth,
by the
1 x 5, u
1 x 5;
heavy,
pinnul
small,
strong
G. ene
carbonar
phiocri

lotoblastus, White, 1874, Rep. Invert. Foss., p. 15, and Geo. Sur. W. 100th Mer., vol. 4, p. 80, Subcarb.

FIG. 323.—*Granatocrinus melo*.

melo, Owen & Shumard, 1850, (Pentremites *melo*.) Jour. Acad. Nat. Sci. Phil., 2d ser., vol. 2, p. 65, Burlington Gr. Etheridge & Carpenter made this species the type of a new genus, *Cryptoblastus*, and referred to the same genus *G. pisum*. The generic characters, however, are not apparent.

melo var. projectus, see *Granatocrinus projectus*.

melonoides, see *Schizoblastus melonoides*. *missouriensis*, Shumard, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 375, Waverly Gr. *neglectus*, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 60, and Geo. Sur. Ill., vol. 5, p. 471, Burlington Gr. *norwoodi*, Owen & Shumard, 1850, (Pentremites *norwoodi*.) Jour. Acad. Nat. Sci. Phil., 2d ser., vol. 2, p. 64, Burlington Gr.

FIG. 324.—*Granatocrinus projectus*. Diagram.

pisum, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 89, and Geo. Sur. Ill., vol. 5, p. 470, Burlington Gr. *projectus*, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 42, and Geo. Sur. Ill., vol. 3, p. 496, Burlington Gr.

roemeri, Shumard, 1855, (Pentremites *roemeri*.) Geo. Rep. Mo., p. 186, Waverly Gr. *sayi*, see *Schizoblastus sayi*.

shumardi, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 257, and Geo. Sur. Ill., vol. 3, p. 498, Burlington Gr.

GRAPHIOCRINUS, DeKoninck & LeHon, 1853, Rech. Crin. Carb. Belg., p. 115. [Ety. *graphion*, writing instrument; *krinon*, lily.] The authors described this genus as having only basals and radials, but as re-defined, by Wachsmuth, there are five basals concealed by the column; subradials 5; radials 1 x 5, upper margins straight; brachials 1 x 5; sutures gaping; arms 10, long, heavy, short joints, parallel sutures; pinnules long; azygous interradial 1, small, but extending above the radials; strong ventral sac or proboscis. Type *G. encrinoides*.

carbonarius, Meek & Worthen, 1861, (Scaphiocrinus *carbonarius*.) Proc. Acad.

Nat. Sci. Phil., p. 140, and Geo. Sur. Ill., vol. 5, p. 562, Coal Meas.

dactylus, Hall, 1860, Supp. to Geo. Rep. Iowa, p. 80, and Geo. Sur. Ill., vol. 5, p. 559, St. Louis Gr.

longicirifer, Wachsmuth & Springer, (in press,) Geo. Sur., Ill., vol. 8, p. 193, Kinderhook Gr.

macadamsi, Worthen, 1873, (Scaphiocrinus *macadamsi*.) Geo. Sur. Ill., vol. 5, p. 495, Keokuk Gr. *quatuordecembrachialis*, see *Eupachyrinus decembrachialis*.

rudis, Meek & Worthen, 1869, (Scaphiocrinus *rudis*.) Proc. Acad. Nat. Sci. Phil., p. 39, and Geo. Sur. Ill., vol. 5, p. 412, Burlington Gr.

simplex, Hall, 1858, (Scaphiocrinus *simplex*.) Geo. Sur. Iowa, p. 551, Burlington Gr.

spinobrachiatus, Hall, 1861, (Scaphiocrinus *spinobrachiatus*.) New Pal. Crin., p. 8, and Bost. Jour. Nat. Hist., p. 306, Burlington Gr.

striatus, Meek & Worthen, 1869, (Scaphiocrinus *striatus*.) Proc. Acad. Nat. Sci. Phil., p. 142, and Geo. Sur. Ill., vol. 5, p. 418, Burlington Gr.

FIG. 327.—*Graphiocrinus wachsmuthi*. Diagram, 2 diam.

tortuosus, Hall, 1861, (Scaphiocrinus *tortuosus*.) Desc. New Crin., p. 7, and Bost. Jour. Nat. Hist., p. 309, Burlington Gr.

FIG. 325.—*Graphiocrinus longicirifer*.

FIG. 326.—*Graphiocrinus rudis*.

- wachsmuthi, Meek & Worthen, 1861, (Scaphiocrinus wachsmuthi.) Proc. Acad. Nat. Sci. Phil., p. 141, and Geo. Sur. Ill., vol. 3, p. 488, Burlington Gr.
- HADROCRINUS**, Lyon, 1869, Trans. Am. Phil. Soc., vol. 13, p. 445. [Ety. *adros*, full grown; *krinon*, lily.] Calyx broad, low vasiform, dome hemispherical; basals 3, hidden by the column; primary radials 2 x 5; secondary, tertiary, and higher orders of radials, having 2 in each series; arm-openings numerous, and not separated by Interradials; Interradials 3 or 4; column round. Type *H. plenissimus*.
- discus*, Lyon, 1869, Trans. Am. Phil. Soc., vol. 13, p. 448, Up. Held. Gr.
- pentagonus*, Lyon, 1869, Trans. Phil. Soc., vol. 13, p. 440, Up. Held. Gr.
- plenissimus*, Lyon, 1869, Trans. Phil. Soc., vol. 13, p. 445, Up. Held. Gr.
- Halysicrinus*, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 110, syn. for *Deltacrinus*.
- HAPLOCORINUS**, Steininger, 1834, Bul. Soc. Geol. France, t. 8, 1st series, p. 232. [Ety. *haploos*, simple; *krinon*, lily.] Calyx small subtrubinate; basals 5; radials 2 x 3 plus 1 x 2, protruding at the center of the superior face for the attachment of arms; dome convex, composed of 5 plates, having sutures from the center of the arm-openings toward the central part of the dome. Type *H. sphaeroideus*.
- clio*, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., App. C., p. 115, Marcellus shale.
- granulatus*, Troost, Not defined.
- hemisphericus*, Troost, Not defined.
- maximus*, Troost, Not defined.
- ovalis*, Troost, Not defined.
- HEMICRINUS**, Von Buch, 1840, Monatsber. d. Berlin Akad., p. 129, and Geol. Russia, vol. 2, p. 31. [Ety. *hemi*, half; *kosmos*, sphere.] Body having four series of plates; basals 4; second series 6; third series 9; ovarian orifice between second and third series; mouth central. Type *H. malum*.
- subglobosus*, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 359, Niagara Gr.
- HEMICYSTITES**, Hall, 1852, Pal. N. Y., vol. 2, p. 245. [Ety. *hemi*, half; *kystis*, bladder.] Parasitic, circular, more or less convex on the upper surface and sometimes sac-like in form; composed of numerous imbricating plates; ambulacra 5, straight, radiating from the center and composed each of a double series of alternating plates, forming part of the upper surface; aperture excentric. Type *H. parasiticus*.
- altus*, syn. for *H. granulatus*.
- granulatus*, Hall, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., pl. 6, fig. 164, Hud. Riv. Gr.
- parasiticus*, Hall, 1852, Pal. N. Y., vol. 2, p. 246, Niagara Gr.
- stellatus*, Hall, 1866, Adv. Sheets 24th Rep. N. Y. St. Mus. Nat. Hist., p. 215, Hud. Riv. Gr.
- HETEROCRINUS**, Hall, 1847, Pal. N. Y., vol. 1, p. 278. [Ety. *heteros*, irregular; *krinon*, lily.] Calyx, small, slightly expanded; basals 5; radials irregular, two or three of the rays having two plates each, and the others only one; four radials supported on the basals; the other is smaller and rests on the azygous plate, and supports the ventral sac on one side and the brachials on the other; brachials, generally, four to each ray, the last one axillary, and supporting two arms, which sometimes branch at irregular distances; pinnules strong; azygous plate pentagonal; column pentagonal, pentapartite; attaching base small. Type *H. heterodactylus*.
- articulosus*, see *Calceocrinus articulatus*.
- bellevillensis*, W. R. Billings, Trans. No. 4, Ottawa Field Naturalists Club, p. 49, Trenton Gr.
- canadensis*, see *Ectenocrinus canadensis*.
- constrictus*, see *Ohioocrinus constrictus*.
- constrictus* var. *compactus*, see *Ohioocrinus compactus*.
- crassus*, see *Iocrinus crassus*.
- exilis*, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 213, Trenton and Hud. Riv. Gr.
- exiguus*, Meek, syn. for *H. exilis*.
- gracilis*, Hall, 1847, Pal. N. Y., vol. 1, p. 280, Hud. Riv. Gr. Not properly defined.
- geniculatus*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 16, Utica Slate Gr.
- heterodactylus*, Hall, 1847, Pal. N. Y., vol. 1, p. 279, Hud. Riv. Gr.
- inaequalis*, see *Calceocrinus inaequalis*.
- incurvus*, see *Anomalocrinus incurvus*.
- isodactylus*, syn. for *Ohioocrinus compactus*.
- juvenis*, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 212, Hud. Riv. Gr.
- laxus*, see *Ohioocrinus laxus*.
- milleri*, Wetherby, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 153, Trenton Gr.
- obovatus*, see *Ohioocrinus obovatus*.
- pentagonus*, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, Hud. Riv. Gr.
- polyxo*, syn. for *Iocrinus subcrassus*.
- simplex*, see *Ectenocrinus simplex*.
- simplex* var. *grandis*, see *Ectenocrinus grandis*.
- subcrassus*, see *Iocrinus subcrassus*.
- tenuis*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 273, and Can. Org. Rem., Decade 4, p. 50, Trenton Gr.
- vaupelii*, syn. for *H. constrictus*.

FIG. 328.—Hemicystites stellatus.

HETEROCRINUS
2, p.
blad
secon
num
unde
armat
220, 1
HETEROCRINUS
III,
lar;
from

FIG. 330.—schlam Side view

gracile,
vol. 7

FIG. 331.—see

Holocystites
St. M
entire
drical
or att
nume
larger
ambu
centr
ing b
proces
abnorm
Mus.
alternat
natus,
p. 23,
Hist.,
baculus,
Nat. I

HETEROCYSTITES, Hall, 1852, Pal. N. Y., vol. 2, p. 229. [Ety. *heteros*, irregular; *kustis*, bladder.] Basals 4, irregular in size; second series 10, large; higher plates numerous, but exact order and number undetermined. Type *H. armatus*.
armatus, Hall, 1852, Pal. N. Y., vol. 2, p. 229, Niagara Gr.

HETEROSCHISMA, Wachsmuth, 1883, Geo. Sur. Ill., vol. 7, p. 352. [Ety. *heteros*, irregular; *schisma*, slit.] It is distinguished from *Codaster* by the sunken hydrospiral areas and exposure of the orals, only, immediately contiguous to the mouth; the limbs are extended interradially, into pyramidal ridges, which the hydrospires enter obliquely. Type *H. gracile*.

alternatum var. *elongatum*, Wachsmuth, 1883, Geo. Sur. Ill., vol. 7, p. 354. Founded upon a magnified view of *Codaster attenuatus*.
gracile, Wachsmuth, 1883, Geo. Sur. Ill., vol. 7, p. 354, Ham. Gr.

FIG. 330.—*Heteroschisma gracile*. Side view, 3 diam.

FIG. 331.—*Heteroschisma gracile*. Horizontal section of hydrospires, 5 diam.

Holocystites, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 353. [Ety. *holos*, entire; *kustis*, bladder.] Body cylindrical, subovate or globose, free, sessile, or attaching by roots, and covered by numerous ranges or irregular series of larger and smaller poriferous plates; ambulacral opening central or subcentral; mouth excentric; smaller opening between these; arms mere spinous processes. Type *H. cylindricus*.
abnormis, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 355, Niagara Gr.
alternatus, Hall, 1861, (Caryocystites *alternatus*), Rep. of Progress Geo. Sur. Wis., p. 23, and 20th Rep. N. Y. St. Mus. Nat. Hist., p. 355, Niagara Gr.
baculus, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 105, Niagara Gr.

brauni, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 130, Niagara Gr.
canneus n. sp.

Niagara Gr.

Body long, irregularly subcylindrical; summit prolonged on the flattened side in the direction of the ambulacral orifice; plates long, polygonal, of irregular size; eight ranges may be counted in our specimen, and one or two have been broken from the lower end; the ambulacral orifice is surrounded by six plates; below this there is a range of eight plates, three of which reach the mouth, and one of which bears the anal orifice; there are no arms, ambulacral spines, or cicatrices; there are eight plates in the next range, two of which join the mouth; the mouth in this genus is generally upon the flattened side of the specimen and opposite the posterior bulge, but not so in this species, for the ambulacral area is prolonged on the flattened side, and the bulge is opposite thereto, while the mouth is on the side of the summit between the bulge and the flattened side;

FIG. 332.—*Holocystites canneus*.

FIG. 333.—*Holocystites canneus*. Summit view.

all the plates are very poriferous, the pores penetrating the plates in clusters of from two to seven instead of by pairs as is usual in this genus; the flattened

side, covered by numerous pits and a thickening of the plates; these pits do not pass through the plates, though they cover a series of plates, sutures and all; such pits have been found on different species, and it is probable they represent a disease of the test, as they seem to destroy the pores and ankylose the sutures. Collected by J. F. Hammell, of Madison, in Jefferson County, Indiana. *cylindricus*, Hall, 1861, (*Caryocystites cylindricus*,) Ann. Rep. Geo. Wis., p. 23, and 20th Rep. N. Y. St. Mus. Nat. Hist., p. 354, Niagara Gr.

dyeri, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 108, Niagara Gr. *elegans*, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 136, Niagara Gr. *faberi*, n. sp. Niagara Gr. Body somewhat

FIG. 334.—*Holocystites faberi*.

obovate; large axial canal passes down into the column; seven plates in the first range; two small intercalated plates on the posterior side between the first and second ranges; seven plates in the second range; above the second range the plates are polygonal, of all sizes, and not disposed in ranges; if they were in ranges there would be about seven below the summit; ambulacral orifice on the posterior side of the summit, elliptical, surrounded by six plates, four of which are protuberant or swollen at the orifice, but no arms ever attached, nor are there cicatrices for spines; mouth pentagonal, on the anterior side of the summit, separated from the ambulacral orifice by two plates, one of which bears the anal orifice; on the posterior side, below the summit, there are three circular pits, which do not seem to have been of any economical use; entire surface poriferous in pairs, which open through small tubercles. Collected by Charles Faber, of Cincinnati, (in whose honor I have given the specific name,) in Jefferson County, Indiana.

globosus, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 133, Niagara Gr.

FIG. 335.—*Holocystites faberi*.

hammelli n. sp. Niagara Gr. Body subglobose, flattened on the anterior side; sessile, no perforation for a column; basals seven; between the first and second series there are five plates inserted on the posterior side; ten plates in the second series; ten in the third; nine in the fourth, and nine in the fifth, which series reaches the mouth; above these there are eight plates in the series which bore the ambulacral spines, and some smaller plates surrounding the ambulacral orifice on the summit; four cicatrices for ambulacral spines; mouth at the margin of the summit; anal opening in the plate adjoining the mouth and between it and the ambulacral orifice; all the plates poriferous in pairs; on the anterior side below the mouth there is one plate in the third series, and one in the fourth series, each bearing a prominent tubercle, with a circle of pores passing through it, giving it a radiate appearance on top; this character may not be of specific importance, but the tubercles are different from those observed on other specimens. Collected by J. F. Hammell, in Jefferson County, Indiana, in whose honor I have given the specific name.

FIG. 336.—*Holocystites hammelli*. Anterior side.

FIG. 337.—*Holocystites hammelli*. Summit view; mouth on the lower side.

jolietensis, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 223, Niagara Gr. *ornatus*, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 132, Niagara Gr. *ovatus*, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 357, Niagara Gr. *perlongus*, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 132, Niagara Gr. *plenus*, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 135, Niagara Gr. *pustulosus*, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 134, Niagara Gr. *rotundus*, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 107, Niagara Gr.

scutellat.
Mus.
sphaeric
Bost.
agara
recogn
subglobo
globos
species
first r
ranges
plates,
the th
ambul
summi
rounde
have c
orifice
orifice

FIG. 338.—*subglobosus*.

Miller,
vol. 2,
tumidus,
Nat. H

FIG. 339.

turbinatu
Soc. Na
ventricos
Soc. Na
wetherby
Soc. Na
winchelli
Mus. N
HOMOCRINUS
p. 185.
Basals
azygous
ventral

scutellatus, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 357, Niagara Gr. sphericus, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., vol. 1, p. 111, Niagara Gr. Not defined so as to be recognized.

subglobosus n. sp. Niagara Gr. Body globose; no axial canal, hence the species was sessile; seven plates in the first range; above this there are four ranges below the summit of fourteen plates, each with an additional plate in the third range on the posterior side; ambulacral orifice in the center of the summit, elongated transversely, surrounded by six plates, four of which have cicatrices for attaching spines; anal orifice near a cicatrix and near the oral orifice; the whole surface is poriferous

in pairs which open on the surface in ornamental sculptured figures, somewhat like the Greek letter ω , (Omega). Collected by Charles Faber, in Jefferson County, Indiana.

subrotundus, S. A.

Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 107, Niagara Gr.

tumidus, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 104, Niagara Gr.

FIG. 338.—*Holocystites subglobosus*.

FIG. 339.—*Holocystites turbidus*.

turbidus, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 259, Niagara Gr. ventricosus, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 108, Niagara Gr. wetherbyi, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 131, Niagara Gr. winchelli, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 356, Niagara Gr.

HOMOCRINUS, Hall, 1852, Pal. N. Y., vol. 2, p. 185. [Ety. *homos*, like; *krinon*, lily.] Basals 5; subradials 5; radials 1x5; azygous interradials 2; proboscis or ventral sac long and large; arms bifur-

cating; pinnules wanting; column round. Distinguished from *Dendrocrinus* by the proportionally larger basals, and absence of one radial, and from *Poteroicrinus* by the arrangement of the azygous plates and absence of pinnules. Type *H. parvus*.

angustatus, see *Dendrocrinus angustatus*.

crassus, Whiteaves, 1887, Cont. to Can. Pal., vol. 1, p. 95, Ham. Gr.

cylindricus, Hall, 1852, Pal. N. Y., vol. 2, p. 186, Niagara Gr.

parvus, Hall, 1852, Pal. N. Y., vol. 2, p. 185, Niagara Gr.

polydactylus, see *Dendrocrinus polydactylus*.

proboscoidialis, Hall, 1859, Pal. N. Y., vol. 3, p. 38, Oriskany sandstone.

scoparius, Hall,

1859, Pal.

N. Y., vol. 3,

p. 102, Low.

Held. Gr.

HYBOCRINUS,

Worthen &

Miller, 1883,

Geo. Sur. Ill.,

vol. 7, p. 331.

[Ety. *hubos*,

humpbacked;

echinos, sea-

urchin.] Test

flexible, sub-

spheroidal,

five ambula-

cral areas,

with numer-

ous ranges of

interlocking

plates imbric-

ating up-

ward, and

each perforated

with a pair of

pores; inter-

ambulacral

areas narrower;

plates

imbricate down-

ward; surface

granular;

jaws strong. Type *H. spectabilis*.

spectabilis, Worthen & Miller, 1883, Geo.

Sur. Ill., vol. 7, p. 332, Kaskaskia Gr.

HYBOCRINUS, Billings, 1857, Rep. of Progr.

Geo. Sur. Can., p. 274, and Can. Org.

Rem., Decade 4, p. 23. [Ety. *hubos*,

humpbacked; *krinon*, lily.] Calyx pro-

tuberant on the

azygous side;

basals 5; radials

1x5; azygous

interradials 2;

arms 5; no pin-

nules; column

round. Type *H.*

conicus.

conicus, Billings, 1857, Rep. of Progr. Geo.

Sur. Can., p. 274, and Can. Org. Rem.,

Decade 4, p. 29, Trenton Gr.

pristinus, Billings, 1859, Can. Org. Rem.,

Decade 4, p. 23, Chazy Gr.

tumidus, Billings, 1857, Rep. of Progr.

Geo. Sur. Can., p. 275, and Can. Org.

Rem., Decade 4, p. 28, Trenton Gr.

FIG. 340.—*Homocrinus scoparius*.

FIG. 341.—*Hyboocrinus conicus*. Diagram.

HYBOCYSTITES, Wetherby, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 150. [Ety. *hubos*, humpbacked; *kustis*, bladder.] Calyx like Hybocrinus, and order of

FIG. 342.—Hybocystites problematicus.

arrangement of the two series of plates is the same, but distinguished by having three arm-like projections, and two or more recumbent arms with ambulacral opening central; from the peristome, the five ambulacra diverge; three are directed to the arm-like projections, and are supposed to pass

over the top of them and extend downward upon the exterior; valvular opening between the upper azygous plate and the mouth. Type H. problematicus. problematicus, Wetherby, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 150, Trenton Gr.

HYDREIONOCRINUS, DeKoninck, 1858, Bull. Acad. Royale Belgique, vol. 8, pt. 2, p. 13. [Ety. *hydreion*, water-bucket; *krinon*, lily.] Calyx short, rounded below; basals 5; subradials 5; radials 1 x 5; brachials 1 x 4 + 2 x 1; arms as in Zeacrinus, and pinnules short; distinguished from Zeacrinus by the ventral sac, which extends beyond the arms and covers them like a roof; the upper plates are convex or spinous; respiratory pores in the sutures of the cylindrical part of the sac, which is covered, by the arms. Type H. woodanus.

acanthophorus, Meek & Worthen, 1870, (Zeacrinus acanthophorus,) Proc. Acad. Nat. Sci. Phil., p. 28, and Geo. Sur. Ill., vol. 5, p. 563, Up. Coal Meas.

armiger, Meek & Worthen, 1870, (Zeacrinus armiger,) Proc. Acad. Nat. Sci. Phil., p. 27, and Geo. Sur. Ill., vol. 5, p. 547, Kaskaskia Gr.

depressus, Troost, as defined by Hall, 1858, (Zeacrinus depressus,) Geo. Sur. Iowa, p. 546, Kaskaskia Gr.

discus, Meek & Worthen, 1860, (Zeacrinus discus,) Proc. Acad. Nat. Sci. Phil., p. 39, and Geo. Sur. Ill., vol. 2, p. 312, Up. Coal Meas.

mucrospinus, McChesney, 1859, (Zeacrinus mucrospinus,) Desc. New Pal. Foss., p. 10, and Trans. Chi. Acad. Sci., p. 7, and Geo. Sur. Ill., vol. 5, p. 563, Coal Meas.

verrucosus, see Eupachycrinus verrucosus.

wetherbyi, Wachsmuth & Springer, 1886, Revis. Palaeocrinoidea, pt. 3, p. 245, Kaskaskia Gr.

HYPANTHOCRINITES, Phillips, 1839, Murch. Sil. Syst. [Ety. *upo*, under; *anthos*, flower; *krinon*, lily.]

caelatus, see Eucalyptocrinus caelatus.

decorus, see Eucalyptocrinus decorus.

Hystericinus, Hinde, 1885, Ann. and Mag. Nat. Hist., p. 158, syn. for Arthracantha carpenteri, see Arthracantha carpenteri.

ICHTHYOCRINUS, Conrad, 1842, Jour. Acad. Nat. Sci. Phil., vol. 8, p. 279. [Ety. *ichthys*, fish; *krinon*, lily.] General form, including incumbent arms, ovoid or pear-shaped; calyx cup-shaped; basals 3; subradials 1 x 5; primary radials 3 or 4 x 5, short and increasing in width, upward; secondary and tertiary radials similar in form to the primaries; arms 40 to 60 or more. Type I. laevis.

FIG. 343.—Ichthyocrinus corbis.

burlingtonensis, Hall, 1858, Geo. Sur. Iowa, p. 557, Burlington Gr.

clintonensis, Hall, 1852, Pal. N. Y., vol. 2, p. 181, Clinton Gr.

corbis, Winchell & Marcy, 1865, Meis. Bost. Soc. Nat. Hist., vol. 1, p. 89, and Jour. Cin. Soc. Nat. Hist., vol. 4, p. 175, Niagara Gr.

laevis, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 279, and Pal. N. Y., vol. 2, p. 195, Niagara Gr.

nobilis, Wachsmuth & Springer, 1878, Proc. Acad. Nat. Sci., p. 254, Upper Burlington and Keokuk Gr.

FIG. 344.—Ichthyocrinus laevis.

subangularis, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 207, and 11th Rep. Geo. Ind., p. 268, Niagara Gr.

tiariformis, Troost, as defined by Hall, 1858, (Cyathocrinus tiariformis,) Geo. Sur. Iowa, p. 558, Subcarboniferous.

Icosidactylocrinites. Not defined.

IOCRINUS, Hall, 1866, Advance sheets, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 212. [Ety. *io*, in triumph; *krinon*, lily.] Calyx pentagonal, pyramidal sides deeply concave; basals 5, indented down the middle; radials 3 to 6 x 5; arms long, frequently bifurcating, but bearing no pinnules; ventral sac very long, extending beyond the arms, subcylindrical longi-

FIG. 345.—Iocrinus. Diagram; b, basals; r, radials; a, azygous plate; t, plates of tube.

tudina
transv
onal.
crassus,
rocin
Phil.,

FIG. 346.—Iocrinus suberectus.

radiale
interr
vault;
ous; v
suber
unkno
inflatus,
flatus,
22, an
Hist.,
parvus,
p. 9,
Ind., p
sculptus,
tennese
Roe
1860,
Fauna
Tenn.,
Niagar
LECANOC
Hall,
Pal. N.
vol. 2,
[Ety.
basin;
non,
Body
arms s
bose;
heavy
sals 3;
dials
primar
dials 1
3 x 5
ondary
dials 1
2; arm
round.

d Mag.
scantha.
nteri.
r. Acad.
[Ety.
General
s, ovoid
ed; ba-
radials
ing, in
tertiary
maries;
Type

1858,
57, Bur-

Pal. N.
nton Gr.
ey, 1865,
t. Hist.,
r. Clin.
75, Ni-

tudinally, five partite, and corrugated transversely; column sharply pentagonal. Type I. subcrassus.

crassus, Meek & Worthen, 1865, (Heterocrinus crassus.) Proc. Acad. Nat. Sci. Phil., p. 147, and Geo. Sur. Ill., vol. 3, p. 325, Hud. Riv. Gr.

FIG. 346.—Iocrinus subcrassus.

polyro, syn. for I. subcrassus.

subcrassus, Meek & Worthen, 1865, (Heterocrinus, subcrassus.) Proc. Acad. Nat. Sci. Phil., p. 148, and Geo. Sur. Ill., vol. 3, p. 325, Hud. Riv. Gr.

trentonensis, Walcott, 1884, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 210, Trenton Gr.

LAMPTEROCRINUS, Roemer, 1860, Sil. Fauna West Tenn., p. 37. [Ety. lampter, lamp; krinon, lily. Calyx urn-shaped, contracted between the arm-bases, and bulged out on the azygous side; basals 5; subradials 5; primary

radials 3 x 5; secondary radials, 1 x 10; interradials 8 or 10, graduating into the vault; azygous interradials more numerous; vault unsymmetrical, and bearing a subcentral proboscis or ventral sac; arms unknown. Type L. tennesseensis.

inflatus, Hall, 1861, (Balanocrinus inflatus.) Rep. of Progr. Sur. of Wis., p. 22, and 20th Rep. N. Y. St. Mus. Nat. Hist., p. 328, Niagara Gr.

parvus, Hall, 1879, Desc. New Spec. Foss., p. 9, and 11th Rep. Geo. Nat. Hist. Ind., p. 272, Niagara Gr.

sculptus, syn. for L. tennesseensis.

tennesseensis, Roemer, 1860, Sil. Fauna West Tenn., p. 37, Niagara Gr.

LECANOCRINUS, Hall, 1852,

Pal. N. Y.,

vol. 2, p. 199.

[Ety. lekane,

basin; krite-

non, lily.]

Body and

arms subglo-

biose; plates

heavy; ba-

asals 3; subra-

dials 1 x 5;

primary ra-

dials 2 or

3 x 5; sec-

ondary ra-

dials 1 to 3 x 10; azygous interradials

2; arms as in Ichthyocrinus; column

round. Type L. macropetalus.

FIG. 347.—Lampteroocrinus tennesseensis.

caliculus, Hall, 1852, Pal. N. Y., vol. 2, p. 203, Niagara Gr.

elegans, see Taxocrinus elegans.

excavatus, Ringueberg, 1886, Bull. Buf.

Soc. Nat. Sci., vol. 5, p. 11, Niagara Gr.

incisus, Ringueberg, Bull. Buf. Soc.

Nat. Sci., vol. 5, p. 10, Niagara Gr.

levis, see Taxocrinus levis.

macropetalus, Hall,

1852, Pal. N. Y.,

vol. 2, p. 199, Ni-

agara Gr.

nitidus, Ringue-

berg, 1886, Bull.

Buf. Soc. Nat.

Sci., vol. 5, p. 9,

Niagara Gr.

ornatus, Hall, 1852,

Pal. N. Y., vol.

2, p. 201, Ni-

agara Gr.

pusillus, Hall, 1863,

(Cyathocrinus

pusillus.) Trans. Alb. Inst., vol. 4, p. 200,

and 11th Rep. Geo. and Nat. Hist. Ind.,

p. 267, Niagara Gr.

pusillus, Winchell & Marcy, syn. for L.

pusillus.

puteolus, Ringueberg, 1886, Bull. Buf.

Soc. Nat. Sci., vol. 5, p. 11, Niagara Gr.

simplex, Hall, 1852, Pal. N. Y., vol. 2, p.

202, Niagara Gr.

solidus, Ringueberg, 1886, Bull. Buf. Soc.

Nat. Sci., vol. 5, p. 8, Niagara Gr.

LECYNOCRINUS, White, 1880, Proc. U. S.

Nat. Mus., vol. 2, p. 257. This name

was preoccupied by Muller in 1858,

and by Zittel in 1879. See Menocrinus.

adamsi, see Menocrinus adamsi.

olliculiformis, see Menocrinus olliculiformis.

LEPADOCRINUS, Conrad,

1840, (Lepocrinites.)

Ann. Rep. N. Y., p.

207. [Ety. from the

resemblance to the

Lepas or Barnacle An-

atifa; krinon, lily.]

Body oblong or ovoid,

consisting of four

series of plates; first

series 4; second series

5; third series 4;

fourth series 5; pec-

tinated rhombs 3 to 5;

arms 3 or 4, recum-

bent, and consisting

of a double series of

interlocking plates,

resting, in shallow

grooves; plates porif-

erous, column taper-

ing. Type L. geb-

hardi.

gebhardi, Conrad, 1840, Fig. 349.—Lepado-

(Lepocrinites geb-

hardi.) Ann. Rep. N. Y., p. 207, and

Pal. N. Y., vol. 3, p. 127, Low. Held.

Gr.

FIG. 348.—Lecanocrinus macropetalus.

FIG. 349.—Lepado-

crinus gebhardi.

hardi.) Ann. Rep. N. Y., p. 207, and

Pal. N. Y., vol. 3, p. 127, Low. Held.

Gr.

moorii, Meek, 1871, (Lepocrinites moorii.) Am. Jour. Sci., 3d series, vol. 2, p. 296, and Ohio Pal., vol. 1, p. 39, Hud. Riv. Gr.

LEPIDOCINUS, Hall, 1861, Desc. New Spec. Crinoidea, p. 18. [Ety. *lepis*, scale; *echinus*, sea-urchin.] Subspheroidal; ambulacral area having a double row of plates imbricating downward, with two pores in each plate, near the outer end; interambulacral areas wide, and having numerous ranges of plates, imbricating from below upward, and from the center outward. Type *L. imbricatus*.

imbricatus, Hall, 1861, Desc. New Crinoidea, p. 18, Burlington Gr.

rarispinus, Hall, 1867, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 340, Waverly Gr.

LEPIDESTHES, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 522. [Ety. *lepis*, scale; *esthes*, garment.] Subspheroidal;

FIG. 350.—*Lepidesthes coreyi*. Diagram 3 diam.; (a.) ambulacral; (b.) section of them; position of interambulacral.

ambulacral areas wide, having numerous plates, and imbricating from above downward, and having two pores in each plate, nearly central; interambulacral areas narrow, plates imbricating from below upward, as well as outward from the middle; jaws well developed; surface granular. Type *L. coreyi*.

colletti, White, 1878, Proc. Acad. Nat. Sci. Phil., p. 33, and Cont. to Pal., No. 8, p. 163, Keokuk Gr.

coreyi, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 525. Keokuk Gr.

formosus, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 41, Keokuk Gr.

LEPIDOCIDARIS, Meek & Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 478. Ety. *lepis*, scale; *Cidaris*, a genus.] Body large, globose, eight or more rows of imbricating plates in the middle of each interambulacral area, but only two reach the oral apertures; plates hexagonal or pentagonal; tubercles for the support of primary spines smooth and in the center of each plate; pustules near the outer edge of the plates for the secondary spines; ambulacra narrow; plates slightly imbricating in the opposite direction from the interambulacral series, and each pierced by two pores; jaws strong. Type *L. squamosus*.

squamosus, Meek & Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 478, Burlington Gr.

LEPIDODISCUS, Meek & Worthen, 1875, Geo. Sur. Ill., vol. 5, p. 573. [Ety. *lepis*, scale; *diskos*, quoin.] A proposed subgenus for Agelacrinites, founded upon *A. squamosus*.

Lepocrinites, Conrad, 1840. The correct orthography seems to be *Lepadocrinites*.

moorei, Meek, see *Lepadocrinites moorii*.

lichenocrinites, Hall, 1866, Adv. sheets 24th Rep. N. Y. St. Mus. Nat. Hist., p. 216. [Ety. *lichen*, tree-moss; *krinon*, lily.] Body parasitic, discoid, more or less crateriform, from the center of which arises a long tapering column, each ring of which is composed of small interlocking plates; upper surface of body covered with polygonal plates, which are supported in the interior by numerous radiating lamellae. Type *L. dyeri*.

affinis, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 229, Hud. Riv. Gr.

crateriformis, Hall, 1866, Adv. sheets 24th Rep. N. Y. St. Mus. Nat. Hist., p. 217, Hud. Riv. Gr.

dubius, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 234, Utica Slate Gr.

dyeri, Hall, 1866, Adv. sheets, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 216, Hud. Riv. Gr.

St. Mus. Nat. Hist., p. 216, Hud. Riv. Gr.

FIG. 351.—*Lichenocrinites tuberculatus*; one specimen shows the radiating lamellae.

pattersoni, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 118, Utica Slate Gr.

tuberculatus, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 346, Hud. Riv. Gr.

LYRIOCINUS, Hall, 1852, Pal. N. Y., vol. 2, p. 197. [Ety. *lyrion*, small lyre; *krinon*, lily.] Calyx hemispherical; basals 5; subradials 5; primary radials 3x5; secondary radials 2x10; interradial areas 4 or 5, which are in closed above by the secondary radials;

FIG. 352.—*Lyrrocinus dactylus*. Side view

azygo
intern
depre
small
10, co
lockin

FIG. 353.—*Lyrrocinus dactylus*.

198,
Hist.

FIG. 354.—

sculptili
Mus.
sculptus,
Archu
nus s
tus.

Lvsocys

n.
[Ety.
settin
kustis,
der.]
subs
roidal
pose
four r
of p
basal
probab
succ
by
range
plates
and
pla
mouth
tral;
rian
turela
Type
dosus.
gener

azygous area similar to the regular interradial areas; vault almost flat, depressed interradially, composed of small plates; opening excentric; arms 10, composed of a double series of interlocking plates. Type *L. dactylus*.

FIG. 353.—*Lyrioerinus dactylus*. Basal view.

198, and 11th Rep. Geo. and Nat. Hist. Ind., p. 269, Niagara Gr.

FIG. 354.—*Lyrioerinus melissa*. Basal view.

sculptilis, Hall, 1864, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 368, Niagara Gr.

sculptus, see *Archaeocrinus sculptus*.

LYROCYSTITES, n. gen.

[Ety. *lysis*, setting free; *kustis*, bladder.] Body subspheroidal, composed of four ranges of plates; basal plates probably 4, succeeded by two ranges of 5 plates each and dome plates; mouth central; ovarian aperture lateral. Type *L. nodosus*. This generic name is proposed instead of

FIG. 355.—*Macrostylocrinus fusibrachiatus*.

Echinocystites, Hall, which was preoccupied. The genus is known only from casts.

nodosus, Hall, 1864, (*Echinocystites nodosus*, 20th Rep. N. Y. Mus. Nat. Hist., p. 360, Niagara Gr.

MACROSTYLOCINUS, Hall, 1852, Pal. N. Y., vol. 2, p. 203. [Ety. *makros*, long; *stylos*, an arm; *krinon*, lily.] Calyx urn-shaped; basals 3; primary radials 3x5; secondary radials 1 or more by 10; regular interradials 3; azygous interradials 4; arms 10. Type *M. ornatus*.

fasciatus, Hall, 1876, (*Cyathocrinus fasciatus*), 28th Rep. N. Y. St. Mus. Nat. Hist., p. 130, Niagara Gr. Probably a syn. for *M. meeki*.

fusibrachiatus, Ringueberg, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 119, Niagara Gr.

meeki, Lyon, 1861, (*Actinocrinus meeki*), Proc. Acad. Nat. Sci. Phil., p. 411, Niagara Gr.

ornatus, Hall, 1852, Pal. N. Y., vol. 2, p. 204, Niagara Gr.

striatus, Hall, 1868, Trans.

Alb. Inst., vol. 4, p. 207,

and 20th Rep. N. Y. St.

Mus. Nat. Hist., p. 327,

Niagara Gr.

striatus var. *granulosus*,

Hall, 1879, 28th Rep.

N. Y. St. Mus. Nat. Hist.,

p. 129 Niagara Gr.

MALOCYSTITES, Billings, 1858, Can. Org. Rem.,

Decade 3, p. 66. [Ety. *malum*, apple;

kustis, bladder.] Body ovate or globular;

plates nonporiferous and in very

irregular series; first series 3; second

series 10 or 12, and in all the series 40

or 50; mouth apical; ambulacral orifice

near the upper part; arms recumbent,

8 or more. Type *M. murchisoni*.

barrandi, Billings, 1858, Can. Org. Rem.,

Decade 3, p. 67, Chazy Gr.

murchisoni, Billings, 1858,

Can. Org. Rem., Decade 3, p. 66, Chazy Gr.

MARIACRINUS, Hall, 1859,

Pal. N. Y., vol. 3, p.

104. [Ety. *Maria*, proper

name; *krinon*, lily.]

Body obconoidal, interradial areas depressed,

surface ornamented; basals 4;

primary radials 3x5;

secondary radials 3x10; tertiary radials 1 or more by 20; interradials 3 to 10, the first one supported by the first radials; azygous area large and plates numerous; vault inflated, plates small; arms composed of a double series of interlocking plates, and not unfrequently bearing armlets consisting also of a double series of interlocking plates; column round. Type *M. nobilissimus*.

FIG. 357.—*Malocystites murchisoni*.

FIG. 356.—*Macrostylocrinus striatus*.

carleyi, Hall, 1862, (Glyptocrinus *carleyi*.) Trans. Alb. Inst., vol. 4, p. 203, and 11th Rep. Geo. and Nat. Hist. Ind., p. 261, Niagara Gr. *macropetalus*, Hall, 1859, Pal. N. Y., vol. 3, p. 111, Low. Held. Gr.

FIG. 358.—*Marlaecrinus carleyi*.

pachydactylus, Conrad, 1841, (Astrocrinites *pachydactylus*.) Ann. Rep. N. Y., p. 34, and Pal. N. Y., vol. 3, p. 107, Low. Held. Gr. Syn. (?) for *M. polydactylus*. *paucidactylus*, Hall, 1859, Pal. N. Y., vol. 3, p. 109, Low. Held. Gr. *plumosus*, Hall, 1859, Pal. N. Y., vol. 3, p. 110, Low. Held. Gr. *polydactylus*, Bonny, 1837, (Actinocrinus *polydactylus*.) Am. Jour., vol. 31, syn. for *M. pachydactylus*? *ramosus*, Hall, 1859, Pal. N. Y., vol. 3, p. 147, Low. Held. Gr. *stoloniferus*, Hall, 1859, Pal. N. Y., vol. 3, p. 112, Low. Held. Gr. *warreni*, Ringueberg, 1888, Proc. Acad. Nat. Sci. Phil., p. 133, Niagara Gr. **MARSUPIOCRINUS**, Phillips, 1839, Murch. Sil. Syst., p. 672. [Ety. *marsupos*, bag; *krinon*, lily.] Basals 3; primary radials 2×5 ; secondary radials 2×5 ; arms 20; distinguished from *Platycrinus* by the higher order of radials, by the round column, instead of elliptic, and by having a larger canal. Type *M. cælatus*. *dactylus*, see *Lyriocrinus dactylus*. *tennesseensis*, Roemer, 1860, (Platycrinus *tennesseensis*.) Sil. Fauna West Tenn., p. 35, Niagara Gr. *tentaculatus*, Hall, 1861, (Platycrinus *tentaculatus*.) Pal. N. Y., vol. 3, p. 116, Low. Held. Gr.

MEGISTOCRINUS, Owen & Shumard, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 594. [Ety. *megistos*, very great; *krinon*, lily.] Body basin-shaped; basals 3; primary radials 3×5 ; secondary radials 1×10 ; tertiary radials 1 or more $\times 20$; inter-radials numerous; first azygous plate like the first radials, and resting on the basals, succeeded by three plates, and these by numerous smaller ones; arms in double series of short plates, bifurcating and bearing pinnules; vault convex; orifice excentric or lateral; column round. Type *M. evansi*. *abnormis*, Lyon, 1857, (Actinocrinus *abnormis*.) Geo. Sur. Ky., vol. 3, p. 479, Up. Held. Gr. *concavus*, Wachsmuth, 1885, Proc. Dav. Acad. Sci. vol. 4, p. 96, Ham. Gr. *brevicornis*, Hall, 1858, (Actinocrinus *brevicornis*.) Geo. Sur. Iowa, p. 571, Burlington Gr.

crassus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 17, Burlington Gr. *depressus*, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 134, Hamilton Gr. *evansi*, Owen & Shumard, 1850, (Actinocrinus *evansi*.) Jour. Acad. Nat. Sci. vol. 1, pt. 2, p. 68, and Geo. Sur. Wis., Iowa, and Minn., p. 594, Burlington Gr. *farnsworthi*, White, 1876, Proc. Acad. Nat. Sci., p. 29, Ham. Gr. *infelix*, see *Saccocrinus infelix*. *knappi*, Lyon & Casseday, 1857, Proc. Acad. Nat. Sci. Phil., p. 412, Up. Held. Gr. *latus*, Hall, 1858, Geo. Sur. Iowa, vol. 1, pt. 2, p. 480, Ham. Gr. *marcouanus*, see *Saccocrinus marcouanus*. *necis*, see *Saccocrinus necis*. *nobilis*, Wachsmuth & Springer, (in press.) Geo. Sur. Ill., vol. 8, p. 139, Waverly or Kinderhook Gr.

nodosus, Barris, 1879, Proc. Dav. Acad. Nat. Sci., vol. 2, p. 285, Up. Held. Gr. *nodosus* var. *multidecoratus*, Barris, 1885, Proc. Dav. Acad. Nat. Sci., vol. 4, p. 98, Ham. Gr.

ontario, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 136, Ham. Gr.

parvirostris, syn. for *M. plenus*. *parvus*, Wachsmuth & Springer, (in press.) Geo. Sur. Ill., vol. 8, p. 171, Kinderhook Gr.

pileatus, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 114, Up. Held. Gr.

plenus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 16, Burlington Gr.

rugosus, Lyon & Casseday, 1859, Am. Jour. Sci., vol. 28, p. 243, Up. Held. Gr.

spinulosus, Lyon, 1861, Jour. Acad. Nat. Sci. Phil., p. 413, Up. Held. Gr.

whitii, Hall, 1861, Jour. Bost. Soc. Nat. Hist., vol. 7, p. 271, Burlington Gr.

FIG. 359.—*Megistocrinus pileatus*. View of vault.

FIG. 360.—*Megistocrinus pileatus*. View from below.

to 12;
vault
Type
bainbui
Hall &
field,
(Cteno
bainbui
sis.) Oh
vol. 2,
Portag
brevirad
Hall &
field,
(Cteno
brevirad
tus, Oh
vol. 2,
Ham.
clarkii,
Sci., p.
lævis Ro
Sil. F
ara Gr
nodosus
19, De

obconicu
vol. 4,
obpyram
(Actin
Bost. S
agara C
pratteni,
pratten
Traus.
saw Gr

stewarti, Safford, 1860, Geo. of Tenn., p. 346, St. Louis Gr.

FIG. 366.—Melonites multiplorus. Diagram, showing pores in ambulacral area, 2 diam.

MENOCRINUS, n. gen. *Ety. menos*, strength of body; *krinon*, lily.] Calyx somewhat globular; basals 3; subradials 1 x 5; radials 1 x 5; azygous and interradials 0; which distinguishes the genus from *Platycrinus*. Type *M. olliculiformis*. This generic name is proposed as a substitute for *Lecythiocrinus*, White, 1880, because that name was preoccupied by Muller in 1858, and by Zittel in 1870.

adamsi, Worthen, 1882, (*Lecythiocrinus adamsi*), Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 37, and Geo. Sur. Ill., vol. 7, p. 317, Coal Meas.

olliculiformis, White, 1880, (*Lecythiocrinus olliculiformis*), Proc. U. S. Nat. Mus., vol. 2, p. 257, and Cont. to Pal., No. 6, p. 124, Up. Coal Meas.

MEROCRINUS, Walcott, 1883, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 208. [*Ety. meros*, hip-joint; *krinon*, lily.] Basals 5, pentangular, low, broad; subradials 5, hexagonal, short, broad; radials pentagonal, four support upon the upper truncate face of each a row of six or seven brachials, and the azygous plate from the same level, the fifth radial; right posterior radial like the azygous plate, but having an angular upper side, giving off, on one side, the ventral tube, and on the other a row of brachials; arms long, bifurcating, without pinnules. Type *M. typus*.

FIG. 367.—Meroocrinus curtus.

curtus, Ulrich, 1879, (*Dendrocrinus curtus*), Cin. Soc. Nat. Hist., vol. 2, p. 18, Utica Slate Gr.

corroboratus, Walcott, 1883, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 210, Trenton Gr.

typus, Walcott, 1883, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 209, Trenton Gr.

MESPILOCRINUS, DeKoninck & LeHon, 1854, Rech. Crin. Terr. Carb. Belg., p. 111. [*Ety. mespilum*, medlar; *krinon*, lily.] Body small, with arms globular or pyriform; basals 3; subradials 5;

radials 3 x 5; arms 10, which divide once, taper rapidly, infold and incline

obliquely from left to right; azygous plate 1, resting upon the larger basal; column round. Type *M. forbesianus*.

konincki, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 69, Burlington Gr.

scitulus, Hall, 1861, Desc. New Crinoidea, p. 9, Burlington Gr.

MYELODACTYLUS, Hall, 1852, Pal. N. Y., vol. 2, p. 191. [*Ety. myelos*, the inside pith; *dactylus*, finger.]

Body consisting of a coil rolled, in the same plane, with finger-like processes, from each coil, overlapping the next inner one; coil and processes perforated so as to form connecting channels. Type *M. convolutus*, *brachiatulus*, Hall, 1852, Pal. N. Y., vol. 2, p. 232, Niagara Gr.

FIG. 368.—Myelodactylus bridgportensis.

bridgportensis, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 141, Niagara Gr.

convolutus, Hall, 1852, Pal. N. Y., vol. 2, p. 192, Niagara Gr.

MYRTILOCRINUS, Sandberger, 1856, Verster Rhein. Schi. Syst. in Nassau. [*Ety. myrtilus*, myrtle; *krinon*, lily.]

Body subglobose or ovoid; basals 5; subradials 5; radials 1 x 5; dome consisting of 5 plates alternating with the radials; arms 5; columnar canal, quadrangular. Type *M. elongatus*.

americanus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., App. C., p. 114, Up. Held. Gr.

Nematocrinus, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 251, syn. for *Catillocrinus*.

NIPTEROCRINUS, Wachsmuth, 1868, Proc. Acad. Nat. Sci. Phil., p. 341. [*Ety. nipter*, washing vessel; *krinon*, lily.]

Calyx basin-shaped; basals 3, nearly hidden by the column; subradials 1 x 5; radials 3 or 4 x 5, the first one very large; arms bifurcating; column round. Type *N. wachsmuthi*.

arboresus, Worthen, 1863, Geo. Sur. Ill., vol. 5, p. 436, Burlington Gr.

wachsmuthi, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 341, and Geo. Sur. Ill., vol. 5, p. 435, Burlington Gr.

NUCLEOCRINUS, Conrad, 1842, Jour. Acad. Nat. Sci. Phil., vol. 8, p. 280. [*Ety. nucleus*, a little nut; *krinon*, lily.]

Calyx ellipsoidal; basals 3, small, hidden within the columnar cavity; radials 5, forming a small cup, deeply scalloped

for re
elong
jectin
deltoid
three-
interc
reache
and d
row c
exten
calyx
long
nume
tubes
an am
which
large,
azygou
and N
angular
laris.)
Ham.
canadens
and G
conradi,
Mus. 1
Held.
elegans,
Sci. Ph
hall's, syn
kirkwood
nus ki
Acad. t
lucina, F
Mus. 1

FIG. 369.—Nematocrinus. Crinodrosipres, 2

verneuilli,
neuilli.)
the Ge
14, and
vol. 3,
Held. G
verneuilli v
eridge d
simply a
very co
other s
without
acters.
OHIOCRINUS,
Palaeocr
proper
calyx a

zygous
basal;
forbi-

o Geo.
Gr.
w Cri-

N. Y.,
myelos,
inger.]
lled, in
ger-like
over-
oil and
m con-
olutes.
vol. 2,

msis.

0, Jour.
141, Ni-
vol. 2,

Verst.
Nassau.
n, lily.]
asals 5;
me con-
with the
l, quad-
p. N. Y.
p. 114,
66, Proc.
syn. for

8, Proc.
[Ety.
n, lily.]
nearly
brachials
first one
column

Sur. Ill.,

p, 1868,
341, and
Burling-

. Acad.
[Ety.
Calyx
hidden
radials 5,
calloped

for receiving the bases of the narrow, elongate ambulacra, and having projecting lips forming a quinquepod; deltoid plates 6, elongate, forming three-fourths of the calyx; a narrow intercalated plate, on the azygous side, reaches from the aperture to the radial, and divides the deltoid into two narrow curving plates; sinuses narrow, extending the entire length of the calyx; ambulacra narrow; lancet plates long and very narrow; side plates numerous; hydrospires pendent, two tubes or elongated sacs on each side of an ambulacrum; spiracles in five pairs, which notch the deltoid plates; mouth large, covered normally with plates; azygous opening large. Types *N. elegans* and *N. verneuili*.

angularis, Lyon, 1857, (*Olivanites angularis*.) Geo. Sur. Ky., vol. 3, p. 492, Ham. Gr.

canadensis, Montgomery, 1881, Can. Nat. and Geol. Vol. 10, p. 83, Ham. Gr.

conradi, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., App. C., p. 121, Up. Held. Gr.

elegans, Conrad, 1842, Jour. Acad. Nat. Sci. Phil., p. 280, Ham. Gr.

halli, syn. for *Nucleocrinus elegans*.

kirkwoodensis, Shumard, 1863, (*Eleacrinus kirkwoodensis*.) Trans. St. Louis Acad. Sci., vol. 2, p. 113, St. Louis Gr.

lucina, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., App. C., p. 120, Ham. Gr.

meloniformis, Barris, 1883, (*Eleacrinus meloniformis*.) Geo. Sur. Ill., vol. 7, p. 361, Ham. Gr.

obovatus, Barris, 1883, (*Eleacrinus obovatus*.) Geo. Sur. Ill., vol. 7, p. 358, Ham. Gr.

coreyi, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 34, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

danæ, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 474, Keokuk Gr.

arms 10, bearing bifurcating pinnules; ventral tube large, having a spiral form somewhat like the cast of a *Murchisonia*, and covered with hexagonal plates; column pentagonal and pentapartite. Type *O. laxus*.

compactus, Meek, 1873, (*Heterocrinus constrictus* var. *compactus*.) Ohio Pal., vol. 1, pl. 11, Hud. Riv. Gr.

constrictus, Hall, 1866, (*Heterocrinus constrictus*.) 24th Rep. N. Y. St. Mus. Nat. Hist., p. 210, Hud. Riv. Gr.

laxus, Hall, 1866, (*Heterocrinus laxus*.) 24th Rep. N. Y. St. Mus. Nat. Hist., p. 211, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

oëhanus, Ulrich, 1882, (*Heterocrinus oëhanus*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 175, Hud. Riv. Gr.

 

FIG. 371.
Oliocrinus compactus.

FIG. 372.
Oliocrinus constrictus.

FIG. 369.—*Nucleocrinus obovatus*. Cross section of hydrospires, 2 diam.

FIG. 370.—*Nucleocrinus verneuili*.

FIG. 373.—*Oligoporus danæ*; 2 diam.

danæ, Meek & Worthen, 1860, (*Melonites danæ*.) Proc. Acad. Nat. Sci. Phil., p. 397, and Geo. Sur. Ill., vol. 2, p. 249, Keokuk Gr.

nobilis, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 358, and Geo. Sur. Ill., vol. 5, p. 476, Burlington Gr.

parvus, Hambach, 1884, Trans. St. Louis Acad. Sci., vol. 4, p. 548, Keokuk Gr.

Olivanites, syn. for *Nucleocrinus*.

angularis, see *Nucleocrinus angularis*.

verneuili, see *Nucleocrinus verneuili*.

Ollaocrinus, Cumberland, 1826, Appendix to *Reliquiæ Conservata*. Figured without description, and subsequently declared by DeKoninck & LeHon to be a *Rhodoocrinus*. Wachsmuth & Springer claim priority for this name over *Goniasteroidocrinus*, without good reason, however, as shown by Meek in Ill. Geo. Sur., vol. 2, p. 217.

ONYCHASTER, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 526. [Ety. *onyx*, claw; *aster*, star.] A small sub discoid body, with five long, slender, rounded, flexible rays; dorsal side of disk composed of an outer circle of five pairs of plates each, pierced with an ovarian pore, and two inner circles of five pairs each, non poriferous, and surrounding a central anal opening; outside the pore plates, each pair is followed by two or three pairs of interlocking transverse plates, connecting with the dorsal side of the rays; farther there are lanceolate plates, furrowed and having pores between the inner ends. Type *O. flexilis*.

barrisi, Hall, 1861, (*Protaster barrisi*.) Desc. New Crinoidea, p. 18, and Geo. Sur. Ill., vol. 5, p. 476, Burlington Gr. *flexilis*, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 526, Keokuk Gr.

ONYCHOCRINUS, Lyon & Casseday, 1859, Am. Jour. Sci., 2d series, vol. 29, p. 77. [Ety. *onyx*, claw; *krinon*, lily.] Calyx low, arms

FIG. 374.—*Onychaster flexilis*.

like the talons of a fowl; basals 3; subradials 5; radials 4 to 7 x 5; arms short, branching; interradials 3 to 20; vault depressed; column large. Type *O. exculptus*.

asteriformis, Hall, 1861, (*Forbesiocrinus asteriformis*.) Desc. New Crin., p. 9, and Geo. Sur. Ill., vol. 2, p. 243, Keokuk Gr.

distensus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 31, and Geo. Sur. Ill., vol. 7, p. 307, Kaskaskia Gr.

FIG. 375.—*Onychocrinus diversus*. Diagram.

diversus, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 256, and Geo. Sur. Ill., vol. 3, p. 492, Burlington Gr.

FIG. 376.—*Onychocrinus exculptus*.

exculptus, Lyon & Casseday, 1859, Am. Jour. Sci. and Arts, vol. 29, p. 78, Keokuk Gr.

magnus
vol. 6
monroe
(Forb
Acad.
Sur. I

FIG. 3

norwoodi,
iocrinus
Sci. Pl
vol. 2,
ramulosus
(Forbe
Sci. an
Orophocrinus
in 186
sench.
mites
The de
was m
foreign
no circ
fossil o
scure,
not no
in 1878
been,
scribed
Codoni
their g
noia.
nition o
Orophoc
Codoni

FIG. 378.—

ical, ba
1 x 5; a

magnus, Worthen, 1875, *Geo. Sur. Ill.*, vol. 6, p. 520, St. Louis Gr.
monroensis, Meek & Worthen, 1861, (*Forbesiocrinus monroensis*.) *Proc. Acad. Nat. Sci. Phil.*, p. 130, and *Geo. Sur. Ill.*, vol. 2, p. 244, Keokuk Gr.

FIG. 377.—*Onychocrinus exculptus*. Diagram.

norwoodi, Meek & Worthen, 1860, (*Forbesiocrinus norwoodi*.) *Proc. Acad. Nat. Sci. Phil.*, p. 389, and *Geo. Sur. Ill.*, vol. 2, p. 245, syn for *O. exculptus*.

ramulosus, Lyon & Casseday, 1859, (*Forbesiocrinus ramulosus*.) *Am. Jour. Sci. and Arts*, vol. 28, p. 235, Keokuk Gr.

6, *Proc.*
 and *Geo.*
 gton Gr.

Orophocrinus was proposed by von Seebach, in 1864, in *Nachr. k. Gesellsch. Wissensch. Göttingen*, p. 110, for *Pentremites stelliformis*, Owen & Shumard. The definition was very imperfect, and was made in a foreign language, in a foreign country, and in a journal having no circulation in America, where the fossil occurs. The definition was so obscure, its application to the species was not noticed until Ludwig discovered it in 1878, and probably never would have been, had Meek & Worthen not described the genus, under the name of *Codonites*, in 1869, and illustrated it in their great work on the Geology of Illinois. Neither the publication or definition of von Seebach is such as to allow *Orophocrinus* to stand in preference to *Codonites*.

FIG. 378.—*Ottawacrinus typus*.

OTTAWACRINUS, W. R. Billings, 1887, *Ottawa Nat. Club*, vol. 1, p. 49. [Ety. propername; *krinon*, lily.] Calyx obcon-

ical, basals 5; subradials 1 x 5; radials 1 x 5; arms 5; azygous plate rests on a

basal as in *Dendrocrinus*, and from which it is distinguished only by the arrangement of the plates on the azygous side. Type *O. typus*.

typus, W. R. Billings, 1887, *Ottawa Nat. Club*, vol. 1, p. 49, Trenton Gr.

PACHYCRINUS, Billings, 1850, *Can. Org. Rem.*, Decade 4, p. 22. [Ety. *pachys*, thick; *krinon*, lily.] Calyx saucer-shaped; basals 1 x 5; radials 1 x 5. Type *P. crassibasalis*.

crassibasalis, Billings, 1850, *Can. Org. Rem.*, Decade 4, p. 22, Chazy Gr.

Pachyocrinus, Wachsmuth & Springer, 1879, *Proc. Acad. Nat. Sci. Phil.*, p. 115. Proposed for a division of *Poteriocrinus* of less than generic importance, but later the same authors referred their type to *Woodocrinus*.

PALAEASTER, Hall, 1852, *Pal. N. Y.*, vol. 2, p. 247. [Ety. *palaos*, ancient; *aster*, star.] Stellate, disk small; two ranges of plates in each ambulacral groove, and two on either side, adambulacral and marginal; four ranges of pores in each groove; oral plates

in pairs at the base of the rays; dorsal plates polygonal, sometimes spinous, madreporic tubercle. Type *P. niagarensis*.

antiqua, Locke, 1846, (*Asterias antiqua*.) *Proc. Acad. Nat. Sci. Phil.*, vol. 3, p. 38, Hud. Riv. Gr. Too poorly defined for determination.

antiquus, Troost, 1835, (*Asterias antiqua*.) *Trans. Geo. Soc. Penn.*, vol. 1, p. 232, Hud. Riv. Gr.

clarkanus, S. A. Miller, 1880, *Jour. Cin. Soc. Nat. Hist.*, vol. 3, p. 236, Hud. Riv. Gr.

clarki, S. A. Miller, *Jour. Cin. Soc. Nat. Hist.*, vol. 1, p. 102, see *Palaeaster clarkanus*.

crawfordsvillensis, S. A. Miller, 1880, *Jour. Cin. Soc. Nat. Hist.*, vol. 2, p. 256, Keokuk Gr.

dubius, Miller & Dyer, 1878, *Cont. to Pal.*, No. 2, p. 256, Utica Slate Gr.

dyeri, Meek, 1872, *Am. Jour. Sci.*, 3d series, vol. 3, p. 257, and *Ohio Pal.*, vol. 1, p. 58, Hud. Riv. Gr.

eucharis, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 330, Ham. Gr.

FIG. 379.—*Palaeaster crawfordsvillensis*, showing madreporic tubercle.

exculptus, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 60, Hud. Riv. Gr.

finli, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 19, Utica Slate Gr.

granti, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 53, Clinton Gr.

granulosus, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 327, Hud. Riv. Gr.

harrisi, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 117, Hud. Riv. Gr.

incomptus, Meek, 1872, Am. Jour. Sci., 3d series, vol. 3, p. 275, and Ohio Pal., vol. 1, p. 64, Hud. Riv. Gr.

jamesi, Dana, 1863, (*Palæsterina* (?) *jamesi*.) Am. Jour. Sci., 2d series, vol. 35, p. 295, Hud. Riv. Gr.

longibrachiatus, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 102, Hud. Riv. Gr.

magnificus, S. A. Miller, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 16, Hud. Riv. Gr.

matutinus, Hall, 1847, (*Asterias matutina*.) Pal. N. Y., vol. 1, p. 91, Trenton Gr.

miamiensis, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 143, Hud. Riv. Gr.

parviusculus, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 69, and Acad. Geol., p. 594, Mid. Sil.

niagarensis, Hall, 1852, Pal. N. Y., vol. 2, p. 247, Niagara Gr.

pulchellus, see *Stenaster pulchellus*.

shafteri, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 320, Hud. Riv. Gr.

FIG. 380.—*Palæaster simplex*.

Meek & Worthen, 1861, (*Petraster wilberianus*.) Proc. Acad. Nat. Sci. Phil., p. 142, Hud. Riv. Gr.

PALÆASTERINA, McCoy, 1851, Brit. Pal. Foss., p. 59, but first defined, by Salter, 1857, Ann. Mag. Nat. Hist. [Ety. *palaios*, ancient; *aster*, star; *inus*, resemblance.] Pentagonal, depressed, with plated disk that fills up the angles, leaving the rays but slightly produced; ambulacra shallow, bordered by subquadrate plates. Type P.

approximata, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 30, Hud. Riv. Gr.

fimbriata, see *Schaenaster fimbriatus*.

jamesi, see *Palæaster jamesi*.

rigida, see *Petraster rigida*.

rugosa, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 291, and Can. Org. Rem., Dec. 3, p. 77, Hud. Riv. Gr.

FIG. 381.—*Palæasterina speciosa*.

speciosa, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 30, Hud. Riv. Gr.

stellata, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 290, and Can. Org. Rem., Decade 3, p. 76, Trenton Gr.

PALÆCHINUS, McCoy, 1844, Carb. Foss. Ireland, p. 172. [Ety. *palaios*, ancient; *echinus*, sea-urchin.] Large, oval or spheroidal; plates spinous; 5 to 8 ranges of plates in the interambulacral areas; 2 ranges in the ambulacral areas, each plate perforated at the outer end by two pores. Type P. *koenigi*.

burlingtonensis, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 396, and Geo. Sur. Ill., vol. 2, p. 230, Burlington Gr.

FIG. 382.—*Palæchinus burlingtonensis*, 2 diam.

gracilis, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 82, and Geo. Sur. Ill., vol. 5, p. 473, Burlington Gr.

PALÆOCOMA, Salter, 1857, Ann. and Mag. Nat. Hist., 2d series, vol. 17. [Ety. *palaios*, ancient; *coma*, hair.] Disk small, plates spinous, rays shallow, and having ambulacral, adambulacral, and marginal plates, the latter bearing spines, inclined toward the extremity of the ray. Type P. *marstoni*.

cylindria,
princeps,
ceps,
Hist.,
spinosa,
PALÆOCOMA
Decad
krinon,
basal
radial
radiat
of the

FIG. 383.—*aeorinus* atus.

Can.
Decad
Trent
striatus,
1859,
Rem.,
p. 25,
sulcatus,
Not d
PALÆOCOMA
Rem.,
ancient
or py
porifer
tenuir
chapma
Decad
dawsoni
Decad
pulcher,
vol. 4,
tenuirac
tenuir
Chapm
Parascri
Proc.
[Ety.
A di
than
as the
P. sal
Cyath
Pentacrin
Rep.
merel
Pentagon
crinoid
PENTREMIT
2, p.
board
or py
lobate
line;
basals
or for
the g

cylindrica, see *Teniaster cylindricus*.

princeps, Hall, 1868, (Ptilonaster princeps), 20th Rep. N. Y. St. Mus. Nat. Hist., p. 334, Chemung Gr.

spinosa, see *Teniaster spinosus*.

PALÆOCRINUS, Billings, 1859, Can. Org. Rem., Decade 4, p. 24. [Ety. *palaios*, ancient; *krinon*, lily.] Calyx oval or pyriform; basals 5; radials 1 x 5; azygous inter-radials 1 to 3; calycinal ambulacra 5, radiating from the center to the bases of the arms. Type *P. striatus*.

FIG. 383. — Palæocrinus striatus.

angulatus, Billings, 1857, (Dendrocrinus angulatus,) Rep. of Prog. Geo. Sur. Can., p. 209, and Can. Org. Rem., Decade 4, p. 24, Trenton Gr.

pulchellus, Billings, 1859, Can. Org. Rem., Decade 4, p. 45, Trenton Gr.

rhombiferus, Billings, 1859,

Can. Org. Rem., Decade 4, p. 45, Trenton Gr.

striatus, Billings, 1859, Can. Org. Rem., Decade 4,

p. 25, Chazy Gr.

sulcatus, Safford,

Not defined.

PALÆOCYSTITES, Billings, 1858, Can. Org. Rem., Decade 4, p. 68. [Ety. *palaios*, ancient; *kystis*, bladder.] Body oval or pyriform; plates numerous and poriferous at the margins. Type *P. tenuiradiatus*.

chapmani, Billings, 1858, Can. Org. Rem., Decade 3, p. 71, Chazy Gr.

dawsoni, Billings, 1858, Can. Org. Rem., Decade 3, p. 70, Chazy Gr.

pulcher, Billings, 1858, Can. Nat. Geo., vol. 4, p. 49, Chazy Gr.

tenuiradiatus, Hall, 1847, (Actinocrinus tenuiradiatus,) Pal. N. Y., vol. 1, p. 18, Chazy Gr.

PARISOCRINUS, Wachsmonth & Springer, 1879, Proc. Acad. Nat. Sci. Phil., p. 115.

[Ety. *parios*, resembling; *krinon*, lily.] A division of *Poteroocrinus* of less than generic value, with *P. perplexus* as the type, and including *P. nereus*, *P. salignoides*, *P. tenuibrachiatus*, and *Cyathocrinus intermedius*.

Pentacrinites hamptoni, Emmons, 1842, Geo. Rep. N. Y., Trenton Gr. This is merely the plate of a crinoid column.

Pentagonites, proposed by Rafinesque for a crinoid column.

PENTREMITES, Say, 1820, Am. Jour. Sci., vol. 2, p. 36. [Ety. *pente*, five; *remos*, a board or plate.] Calyx globose, ovoid, or pyriform; base never distinctly trilobate, nor excavated in the middle line; section more or less triangular; basals 3, forming a small cup; radials or fork plates 1 x 5, long, forming the greater part of the calyx; limbs

long, with flat or concave sides and truncated above; sinus, subpetaloid; deltoid plates 1 x 5, small; ambulacra subpetaloid, resting in the sinuses or forks of the radials; lanceolate plates resting below on under lanceolate plates, and forming about a third the width of the ambulacra; side plates numerous and abutting the lanceolate plates; hydrosphere or ambulacral pores partially excavated out of the sides of the sinuses; pinnales attached between the pores; hydrospheres from 3 to 9, pendent, but partially contained within the substance of the radials near their distal ends; spiracles or apertures single or double, partially excavated in the deltoid plates; posterior spiracles confluent with the azygous opening; peristome covered by minute polygonal plates; column round. Type *P. godoni*.

abbreviatus, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, p. 155, Kaskaskia Gr.

angularis, Lyon, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 631, Kaskaskia Gr.

basilaris, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, p. 145, Kaskaskia Gr.

bipyramidalis, Hall, 1858, see *Troostocrinus bipyramidalis*.

bradleyi, Meek, 1873, 6th Rep. Geo. Sur. Terr., p. 470. Not satisfactorily defined.

broadheadi, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, p. 145, Kaskaskia Gr.

burlingtonensis, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., and Geo. Sur. Ill., vol. 5, p. 461, Burlington Gr.

calyce, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 122, Ham. Gr.

calycinus, Lyon, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 628, Kaskaskia Gr.

cervinus, Hall, 1858, Geo. Sur. Iowa, p. 690, Kaskaskia Gr.

cherokeeus, Troost, 1850, Catal. Proc. Am. Assoc. Ad. Sci. and Geo. Sur. Iowa, p. 691, Kaskaskia Gr.

chesterensis, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, p. 145, Kaskaskia Gr.

clavatus, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, Kaskaskia Gr.

conoideus, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 5, and Geo. Sur. Iowa, p. 655, Warsaw Gr.

cornutus, see *Granatocrinus cornutus*.

curtus, see *Granatocrinus curtus*.

decussatus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 243, Keokuk Gr.

elegans, Lyon, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 632, Kaskaskia Gr.

elongatus, Shumard, 1855, Geo. Rep. Mo., p. 187, Burlington Gr.

florealis, Schlothheim, 1820, syn. for *P. godoni*.

FIG. 384. — Palæocrinus striatus. Diagram.

gemmiformis, Hambach, 1884, Trans. St. Louis Acad. Sci., vol. 4, p. 548, Kaskaskia Gr.

globosus, Say, as identified by Troost, 1850, probably *Pentremites sulcatus*.

FIG. 375.—*Pentremites godoni*.

godoni, DeFrance, 1818, Dict. Sci. Nat., t. 14, p. 467, Kaskaskia Gr.

granulatus, see *Granatocrinus granulatus*.

grosvenori, Shumard, see *Troostocrinus grosvenori*.

hemisphericus, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, p. 145, Kaskaskia Gr.

kentuckiensis, see *Codaster kentuckiensis*.
koninckanus, Hall, 1858, Trans. A.S. Inst., vol. 4, p. 4, and Geo. Sur. Iowa, p. 656, Warsaw Gr.

laterniformis, Owen & Shumard, 1850, Jour. Acad. Nat. Sci., 2d series, vol. 2, p. 66, Kaskaskia Gr.

leda, Hall, see *Granatocrinus leda*.

lineatus, see *Troostocrinus lineatus*.

longicostalis, Hall, 1890, Supp. to Geo. Iowa, p. 85, Warsaw Gr. Not satisfactorily defined.

lycorias, Hall, 1863, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Ham. Gr.

maia, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 122, Ham. Gr.

melo, see *Granatocrinus melo*.

missouriensis, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 81, Kaskaskia Gr.

nodosus, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, p. 145, Kaskaskia Gr.

norwoodi, see *Granatocrinus norwoodi*.

obesus, Lyon, 1857, Geo. Sur. Ky., vol. 3, p. 469, Kaskaskia Gr.

obliquatus, see *Tricelocrinus obliquatus*.
ovalis, Owen. Not defined.

potteri, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, p. 156, Burlington Gr.

pyriformis, Say, 1825, Jour. Acad. Nat. Sci. Phil., vol. 4, p. 294, Kaskaskia Gr.

reinwardti, Troost, see *Troostocrinus reinwardti*.

robustus, Lyon, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 629, Kaskaskia Gr.

roemeri, see *Granatocrinus roemeri*.

sampsoni, Hambach, 1884, Trans. St. Louis Acad. Sci., vol. 4, p. 548, Choteau or Waverly Gr.

sayi, see *Schizoblastus sayi*.

sirius, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 20, Burlington Gr.

spinous, Hambach, 1880, Trans. St. Louis Acad. Sci., vol. 4, p. 145, Kaskaskia Gr.
stelliformis, see *Orophocrinus stelliformis*.
subconoides, Meek, 1873, Hayden's Geo. Sur. Terr., p. 471, Subcarb.

FIG. 387.—*Pentremites pyriformis*. One shows the deltoid plates extended to the summit.

subcylindricus, see *Troostocrinus subcylindricus*.

subtruncatus, see *Troostocrinus subtruncatus*.

sulcatus, Roemer, 1852, Monog. Blastoid., p. 354, Kaskaskia Gr.

symmetricus, Hall, 1858, Geo. Rep. Iowa, p. 694, Kaskaskia Gr.

tennessee, Troost. Not defined.

troosti, Shumard, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 386, Kaskaskia Gr.

truncatus, Conrad, 1843, Proc. Acad. Nat. Sci. Phil., vol. 1, p. 334, Warsaw Gr.

varsouviensis, see *Tricelocrinus varsouviensis*.

verneuili, see *Nucleocrinus verneuili*.

whitii, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 122, Ham. Gr.

woodmani, see *Tricelocrinus woodmani*.

wortheni, see *Tricelocrinus wortheni*.

PENTREMITEA D'ORBIGNY, 1849, Prodr. d. Paléont., t. 1, p. 102. [Ety. from *Pentremites*.] Number and disposition of plates as in *Pentremites*, but the deltoids are inconspicuous, confined to the summit, rarely visible in a side view; spiracles large. It is also closely connected with *Troostocrinus*, and is of doubtful generic value. Type *P. schultzi*.

americana, Barris, 1883, Geo. Sur. Ill., vol. 7, p. 363, Ham. Gr.

filosa, Whiteaves, 1887, Cont. to Can. Pal., vol. 1, p. 104, Ham. Gr.

PERISCHODOMUS, McCoy, 1849, Ann. Nat. Hist., vol. 3, p. 251. [Ety. *perischo*, inclosing; *domus*, house.] Spheroidal, subpentagonal; ambulacra narrow, two rows of small plates imbricating downward, each pierced by one pair of simple pores; interambulacra wide, with five rows of

FIG. 388.—*Pentremitea americana*; 2 diam.

plates
the c
the r
supp
forat
round
plates
open
biseri
illinois
Geo.
ka-ki
Pereichoc
Mag.
some
crinn
PETRASTE
Decad
aster,
ter,
adam
plates
rigid

bellulus
p. 393
rigidus,
idus,
291, a
80, T
wilberia
Philocrinus
favori
pelvis, M
Sci.
crinn
PHOLINOC
Proc.
pholia
ambu
upwa
rows
ties;
lower
show
surro
areas
plates
plate
larger
Type
irregula
Acad
Sur. I

plates imbricating upward, and from the center outward; primary spines on the rows adjoining the ambulacra, the supporting tubercle being small, perforated, but not crenulated, and surrounded by a double ring; ovarian plates having 6 pores; mouth and anal openings small, central. Type *P. biserialis*.

illinoensis, Worthen & Miller, 1883, *Geo. Sur. Ill.*, vol. 7, p. 333, *Kaskaskia Gr.*

Pereichocrinus, Austin, 1843, *Ann. and Mag. Nat. Hist.*, vol. 11, p. 203. Not defined so as to be recognized, though some authors use it instead of *Saccocrinus*.

PETRASTER, Billings, 1858, *Can. Org. Rem.*, Decade 3, p. 79. [Ety. *petros*, stone; *aster*, star.] Closely related to *Palæaster*, and having both marginal and adambulacral plates, with a few disk-plates, on the ventral side. Type *P. rigidus*.

FIG. 389.—*Petraster bellulus*.

bellulus, Billings, 1865, *Pal. Foss.*, vol. 1, p. 393, *Niagara Gr.*

rigidus, Billings, 1857, (*Palæasterina rigidus*.) *Rep. of Progr. Geo. Sur. Can.*, p. 291, and *Can. Org. Rem.*, Decade 3, p. 80, *Trenton Gr.*

wilberianus, see *Palæaster wilberianus*.

Philocrinus, Koninck, 1863. [Ety. *philos*, favorite; *krinon*, lily.]

pelvis, Meek & Worthen, 1865, *Am. Jour. Sci.*, 2d series, vol. 39, syn. for *Erisocrinus typus*.

PHOLIDOCIPARIS, Meek & Worthen, 1869, *Proc. Acad. Nat. Sci. Phil.*, p. 77. [Ety. *pholidos*, scale; *kideris*, turban.] Interambulacra thin, irregular, imbricating upward and laterally; five or more rows; only two reaching the extremities; marginal rows and those on the lower side having primary tubercles, showing a pit in the top, and being surrounded by two rings; ambulacral areas wide, with six or more rows of plates imbricating downward, each plate pierced by two pores, and the larger ones having additional pores. Type *P. irregularis*.

irregularis, Meek & Worthen, 1869, *Proc. Acad. Nat. Sci. Phil.*, p. 78, and *Geo. Sur. Ill.*, vol. 5, p. 512, *Keokuk Gr.*

PHYSETOCRINUS, Meek & Worthen, 1869, *Proc. Acad. Nat. Sci. Phil.*, p. 158.

[Ety. *physetos*, inflated; *krinon*, lily.] Distinguished from *Actinoocrinus*, in the form and construction of the vault, which has no proboscis, and has pores along the radial portions of the dome; and, also, in having no hook-like projections along the pinnules. Type *P. ventricosus*.

asper, Meek & Worthen, 1869, *Proc. Acad. Nat. Sci. Phil.*, p. 161, and *Geo. Sur. Ill.*, vol. 5, p. 351, *Burlington Gr.*

copei, S. A. Miller, 1881, (*Actinoocrinus copei*.) *Jour. Cin. Soc. Nat. Hist.*, vol. 4, p. 310, *Burlington Gr.*

dilatatus, Meek & Worthen, 1869, (*Strotoocrinus dilatatus*.) *Proc. Acad. Nat. Sci. Phil.*, p. 162, and *Geo. Sur. Ill.*, vol. 5, p. 363, *Burlington Gr.*

ornatus, Hall, 1858, (*Actinoocrinus ornatus*.) *Geo. Sur. Iowa*, p. 583, *Burlington Gr.*

reticulatus, Hall, 1861, (*Actinoocrinus reticulatus*.) *Desc. New Crin.*, p. 3, *Burlington Gr.*

subventricosus, McChesney, 1860, (*Actinoocrinus subventricosus*.) *Desc. New Crin. Pal. Foss.*, p. 21, and *Trans. Chi. Acad. Sci.*, p. 16, *Burlington Gr.*

ventricosus, Hall, 1858, (*Actinoocrinus ventricosus*.) *Geo. Sur. Iowa*, p. 595, *Burlington Gr.*

ventricosus var. *cancellatus*, Hall, 1861, (*Actinoocrinus ventricosus* var. *cancellatus*.) *Bost. Jour. Nat. Hist.*, vol. 7, p. 279, *Burlington Gr.*

ventricosus var. *internodus*, Hall, 1861, (*Actinoocrinus ventricosus* var. *internodus*.) *Bost. Jour. Nat. Hist.*, vol. 7, p. 278, *Burlington Gr.*

PISOCRINUS, DeKoninck, 1858, *Bull. Acad. Roy. Belgique*, 2me ser., tome 3, p. 24. [Ety. *pisos*, pea; *krinon*, lily.] Calyx round globular; basals 5, forming a triangle; these are followed by three large plates, forming nearly the entire calyx; one of these supports two small plates, and a small plate is supported, in a notch, between the other two larger plates; arms 5; column round. Type *P. pilula*.

gemmiformis, S. A. Miller, 1879, *Jour. Cin. Soc. Nat. Hist.*, vol. 2, p. 113, *Niagara Gr.*

globosus, Ringueberg, 1884, (*Triacrinus globosus*.) *Proc. Acad. Nat. Sci. Phil.*, p. 146, *Clinton Gr.*

FIG. 390.—*Physetocrinus copei*.

FIG. 391.—*Pisocrinus gemmiformis*. Natural size and magnified.

Sci. Phil., p. 146, Clinton Gr.

- pyriformis, Ringueberg, (Triacrinus pyriformis,) Proc. Acad. Nat. Sci. Phil., p. 145, Clinton Gr.
- PLATYCRINUS**, Miller, 1821, Nat. Hist. Crinoidea, p. 73. [Ety. *platys*, flat; *krinos*, lily.] Calyx bowl-shaped; basals 3; primary radials 2x5; regular interradials 1x4; azygous interradials, 1 large and 3 small; dome elevated; arms 10 to 35, bearing pinnules; column large and twisted. Type *P. laevis*.
- aequalis*, Hall, 1861, Desc. New Crin., p. 117, and Geo. Sur. Ill., vol. 5, p. 456, Burlington Gr.
- americanus*, Owen & Shumard, 1850, Jour. Acad. Nat. Sci. Phil., 2d ser., vol. 2, p. 89, Burlington Gr.
- andixoni*, Troost. Not defined.

FIG. 392.—*Platyacrinus asper*. Diagram.

- asper*, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 129, and Geo. Sur. Ill., vol. 3, p. 468, Burlington Gr.
- bedfordensis*, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 161, Erie Shales.
- bloomfieldensis*, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 257, Keokuk Gr.
- bonoensis*, White, 1878, Proc. Acad. Nat. Sci. Phil., p. 30, and Cont. to Pal., No. 6, p. 160, Keokuk Gr.
- brevinodus*, Hall, 1861, Desc. New Crinoidea, p. 4, and Bost. Jour. Nat. Hist., vol. 7, p. 286, Keokuk Gr.
- burlingtonensis*, Owen & Shumard, 1850, Jour. Acad. Nat. Sci., 2d ser., vol. 2, p. 60, Burlington Gr.
- calyculus*, Hall, 1861, Desc. New Crin., p. 16, Burlington Gr.
- canaliculatus*, Hall, 1858, Geo. Sur. Iowa, vol. 1, pt. 2, p. 539, Burlington Gr.
- cavus*, Hall, 1858, Geo. Sur. Iowa, p. 527, Burlington Gr.
- clytis*, Hall, 1861, Desc. New Crin., p. 4, and Bost. Jour. Nat. Hist., vol. 7, p. 285, Burlington Gr.
- contritus*, Hall, 1863, 17th Rep. N. Y. St. Mus. Nat. Hist., p. 54, and Ohio Pal., vol. 2, p. 166, Waverly Gr.
- corporeculus*, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 12, Niagara Gr.
- Not properly defined.

- corrugatus*, Owen & Shumard, 1850, Jour. Acad. Nat. Sci., vol. 2, p. 59, Burlington Gr.
- depressus*, Owen. Not defined.
- discoideus*, Owen & Shumard, 1850, Jour. Acad. Nat. Sci., 2d ser., vol. 2, p. 58, Burlington Gr.
- eboracensis*, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 119, Ham. Gr.
- elegans*, Hall, 1861, Desc. New Crin., p. 4, and Bost. Jour. Soc. Nat. Hist., vol. 7, p. 285, Burlington Gr.
- eminulus*, Hall, 1861, Desc. New Crin., p. 17, Burlington Gr.
- eriensis*, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 119, Ham. Gr.
- excavatus*, Hall, 1861, Desc. New Crin., p. 4, and Bost. Jour. Nat. Hist., vol. 7, p. 286, Burlington Gr.
- exsertus*, Hall, 1858, Geo. Sur. Iowa, p. 539, Burlington Gr.
- faberi*, n. sp. Calyx bowl-shaped: sub-cylindrical above; attaching point for column projecting below; base marked by three keels, corresponding with the sutures between the basal plates, and along which the sutures may be distinguished; radials large, width a little greater than height; cicatrix for attachment of arms in the center of the upper face of each radial, and occupying about one-third of the width of the plate; surface marked by a row of tubercles radiating on each

FIG. 393.—*Platyacrinus faberi*, x 2. FIG. 394.—*Platyacrinus hemisphericus*. Diagram.

plate from the angle of union of basal and radial plates, and also a few scattering tubercles; collected in Scott County, West Va., in the St. Louis or Kaskaskia Gr.

georgii,
Iowa,
glyptus,
16, Bu
graphia
Mus.
vol. 2
balli, S
Acad.
Ill., v
haydeni
Terr.,
p. 122

FIG. 395.—*Platyacrinus hemisphericus*.

Trans.
Up. H
lodensis
Pal., v
monroei
Ill. St.
Sur. I
montane
en-is.
multibr
1861.
135, W
nioten
Meek
Wort
1865, H
Acad.
Sci. P
p. 162,
Geo.
Ill., v
p. 513,
knk G
nodobrac
tus, I
1858,
Sur. I
p. 5
Burlin
nodulosu
541, Bu

georgii, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 82, Warsaw Gr.
glyptus, Hall, 1861, Desc. New Crin., p. 16, Burlington Gr.
graphiens, Hall, 1863, 17th Rep. N. Y. St. Mus. Nat. Hist., p. 55, and Ohio Pal., vol. 2, p. 166, Waverly Gr.
halli, Shumard, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 388, and Geo. Sur. Ill., vol. 5, p. 454, Burlington Gr.
haydeni, Meek, 1872, Hayden's Geo. Sur. Terr., p. 469, and Cont. to Pal., No. 6, p. 122, Subcarboniferous.

FIG. 395.—*Platyocrinus hemisphericus*.

hemisphericus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 16, and Geo. Sur. Ill., vol. 3, p. 511, Keokuk Gr.

huntsvillae, Troost. Not defined.

incomptus, White, 1863, Jour. Bost. Soc. Nat. Hist., vol. 7, p. 503, and Geo. Sur. Ill., vol. 5, p. 459, Burlington Gr.

inornatus, syn. for *P. burlingtonensis*.

insculptus, Troost. Not defined.

leai, Lyon, 1869, Trans. Am. Phil. Soc., vol. 13, p. 459, Up. Held Gr.

lodensis, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 168, Waverly Gr.
monroensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 30, and Geo. Sur. Ill., vol. 7, p. 306, St. Louis Gr.
montanensis, see *Eucladocrinus montanensis*.

multibrachiatus, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 135, Warsaw Gr.

niotensis,

Meek &

Worthen,

1865, Proc.

Acad. Nat.

Sci. Phil.,

p. 162, and

Geo. Sur.

Ill., vol. 3,

p. 513, Keo-

kuk Gr.

nodobrachi-

atus, Hall,

1853, Geo.

Sur. Iowa,

Fig. 396.—*Platyocrinus niotensis*.

p. 542,

Burlington Gr.

nodulosus, Hall, 1858, Geo. Sur. Iowa, p.

541, Burlington Gr.

Diagram.

nucleiformis, Hall, 1858, Geo. Sur. Iowa, p. 540, Burlington Gr.

olla, Hall, 1861, Desc. New Crin. The name was preoccupied. See *P. halli*.

ornigranulus, McChesney, 1860, Desc. New Pal. Foss., p. 5, and Trans. Chi. Acad. Sci., p. 3, Burlington Gr.

oweni, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 120, Burlington Gr.

parvinodus, Hall, 1861, Desc. New Crinoidea, p. 17, Burlington Gr.

parvulus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 163, and Geo. Sur. Ill., vol. 5, p. 555, Kaskaskia Gr.

parrus, see *Cordylocrinus parvus*.

penicillus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 380, and Geo. Sur. Ill., vol. 2, p. 266, St. Louis Gr.

perasper, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 161, Burlington Gr.

piliciformis, Hall, 1858, Geo. Sur. Iowa, p. 529, Burlington Gr.

planus, Owen & Shumard, 1850, Jour. Acad. Nat. Sci. Phil., 2d ser., vol. 2, p. 57, Burlington Gr.

plenus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 380, and Geo. Sur. Ill., vol. 2, p. 267, St. Louis Gr.

pleurovimineus, see *Eucladocrinus pleurovimineus*.

plumosus, see *Cordylocrinus plumosus*.

pocilliformis, Hall, 1858, Geo. Sur. Iowa, p. 528, Burlington Gr.

poculum, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 311, Burlington Gr.

polydactylus, Troost. Not defined.

praematurus, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 124, Niagara Gr.

praenuntius, Wachsmuth & Springer, 1878, Proc. Acad. Nat. Sci. Phil., p. 249, Burlington Gr.

prattenanus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 379, and Geo. Sur. Ill., vol. 2, p. 264, St. Louis Gr.

pratteni, Worthen, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 69, Burlington Gr.

pumilus, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 82, Warsaw Gr.

quinquenodus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 18, Burlington Gr.

ramulosus, see *Cordylocrinus ramulosus*.

regalis, Hall, 1861, Desc. New Crinoidea, p. 16, Burlington Gr.

richfieldensis, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 167, Waverly Gr.

saffordi, Troost, 1850, Hall, 1858, Geo. Sur. Iowa, p. 634, Keokuk Gr.

sarae, Hall, 1858, Geo. Sur. Iowa, p. 673, St. Louis Gr.

scobina, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 129, and Geo. Sur. Ill., vol. 3, p. 466, Burlington Gr.

sculptus, Hall, 1858, Geo. Sur. Iowa, p. 536, Burlington Gr.

shumardanus, Hall, 1858, Geo. Sur. Iowa, vol. 1, pt. 2, p. 532, Burlington Gr.
siluricus, Hall, 1879, Desc. New Spec. Foss., p. 9, and 11th Rep. Geo. Sur. Ind., p. 256, Niagara Gr.

FIG. 397.—*Platyterinus scobina*. Diagram.

striobrachiatum, Hall, 1861, Desc. New Crinoidea, p. 4, and Bost. Jour. Nat. Hist., vol. 7, p. 287, Burlington Gr.
subspinulosus, Hall, 1858, Geo. Sur. Iowa, p. 536, Burlington Gr.
subspinulosus, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 81, Burlington Gr.
symmetricus, Wachsmuth & Springer, (in press,) Geo. Sur. Ill., vol. 8, p. 186, Waverly or Kinderhook Gr.
tennesseensis, see Marsupiocrinus tennesseensis.
tentaculatus, see Marsupiocrinus tentaculatus.
tenuibrachiatum, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 16, and Geo. Sur. Ill., vol. 5, p. 450, Burlington Gr.
truncatulus, Hall, 1858, Geo. Sur. Iowa, p. 538, Burlington Gr.
truncatus, Hall, 1858, Geo. Sur. Iowa, p. 537, Burlington Gr.
tuberosus, Hall, 1858, Geo. Sur. Iowa, p. 534, Burlington Gr.
verrucosus, White, 1863, Jour. Bost. Soc. Nat. Hist., vol. 7, p. 502, Burlington Gr.
vexabilis, White, 1875, U. S. Sur. W. 100th Meridian, vol. 4, p. 81, Sub. Carb.
wortheni, Hall, 1858, Geo. Sur. Iowa, p. 530, Burlington Gr.
yandelli, Owen & Shumard, 1850, Jour. Acad. Nat. Sci., 2d ser., vol. 2, p. 58, Burlington Gr.

PLATYCYSTITES, n. gen. [Ety. *platys*, flat; *kustis*, bladder.] General form compressed elliptical, or like the kernel of a peach-seed, with a narrow rim on the border; axial canal passes down into the column; three plates in the first range, one of them bending around the rim and the other two having the dividing suture in the rim itself; there are five large plates in the second range, and above these there are three large plates, on the azygous side, with six or more smaller ones on the border; the whole surface is granular and every plate full of minute pores. Type *P. faberi*.

faberi, n. sp. Basal plates hexagonal, longer than wide, one of them bending around the bordered rim, and the other two uniting at the middle of the rim; a large subcentral hexagonal plate in the second range on the azygous side rests upon the upper side of the plate, which is separated by a suture in the rim from an adjoining basal plate, but does not reach the other basal; this large hexagonal plate joins two plates in the second range with its under sloping sides; three large plates rest upon the three upper faces of this large hexagonal plate; the one upon the superior face is octagonal, resting between the other two large plates and having five smaller ones joining its upper faces. The specimen is worn at the upper edge so as to destroy the orifices. It was received by Charles Faber among a lot of fossils from the Kaskaskia Group in the southern part of West Virginia, but as no cystideans have ever been found above the Lower Devonian, and as the specimen is worn as if it had been drifted, the probability is that it belongs to the Silurian rocks.

FIG. 398. *Platyterinus faberi*.

PLEUROCYSTITES, Billings, 1854, Can. Jour., vol. 2, p. 250. [Ety. *pleuron*, side; *kustis*, bladder.] Body oval, flat; dorsal side with large plates, ventral with smaller ones; two free arms: mouth at the base on the left side; small orifice near the apex; three pectinated rhombs, two in the upper half and one in the lower; column short. Type *P. squamosus*.

FIG. 399.—*Pleurocystites squamosus*.

252, and Can. Org. Rem., Decade 3, p. 49, Trenton Gr.

anticostiensis, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 288, and Can. Org. Rem., Decade 3, p. 52, Hud. Riv. Gr.

elegans, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 287, and Can. Org. Rem., Decade 3, p. 51, Trenton Gr.

exornatus, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 287, Trenton Gr.

flitextus, Billings, 1854, Can. Jour., vol. 2, p. 252, and Can. Org. Rem., Decade 3, p. 48, Trenton Gr.

robustus, Billings, 1854, Can. Jour., vol. 2, p. 252, and Can. Org. Rem., Decade 3, p. 49, Trenton Gr.

squamosus, vol. 2, p. 252, and Can. Org. Rem., Decade 3, p. 49, Trenton Gr.

FIG. 400.—I

Sur. C
Decade

FIG. 401

crassus, Acad. Sur. Ill. pentagon & W. 1865, Acad. Phil., and C. Ill., v. 332, Tr. smithi 1881, T. tawa F. uralists No. 2, POTERICRINOidea krinon, sub-rhombic n. ray of azygon succed. part of in a lora or bra Type I aequalis, Iowa, alternatus

squamosus, Billings, 1854, Can. Jour., vol. 2, p. 251, and Can. Org. Rem., Decade 3, p. 49, Trenton Gr.

POROCRINUS, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 279. [Ety. from the poriferous areas similar to pectinated rhombs.] Calyx conical; basals 5; sub-radials 1 x 5; radials 1 x 5; azygous interradials 2; pectinated rhombs numerous. Type P. conicus.

FIG. 400.—Porocrinus conicus. Diagram.

Sur. Can., p. 279, and Can. Org. Rem., Decade 4, p. 34, Trenton Gr.

FIG. 401.—Porocrinus crassus. Diagram.

crassus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 115, and Geo. Sur. Ill., vol. 3, p. 330, Hud. Riv. Gr.

pentagonus, Meek

& Worthen,

1865, Proc.

Acad. Nat. Sci.

Phil., p. 146,

and Geo. Sur.

Ill., vol. 3, p.

332, Trenton Gr.

smithi, Grant,

1881, Trans. Ot-

tawa Field Nat-

uralists' Club,

No. 2, p. 42, Trenton Gr.

POTERICRINUS, Miller, 1821, Nat. Hist. Crinoidea, p. 68. [Ety. *poterion*, goblet; *krinos*, lily.] Calyx obconical; basals 5; sub-radials 5; radials 1 x 5, with a variable number of smaller ones, the azygous ray often having more than the others; azygous plates 3 or 4, within the calyx, succeeded by smaller ones that form part of the ventral sac; vault produced in a long sac or proboscis; arms simple or branching and bearing pinnules. Type P. crassus.

aequalis, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 63, Burlington Gr.

alternatus, see Dendrocrinus alternatus.

anomalos, Wetlieryby, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 158, Kaskaskia Gr.

arachniformis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 13, and Geo. Sur. Ill., vol. 7, p. 281, Keokuk Gr.

asper, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 11, and Geo. Sur. Ill., vol. 7, p. 278, Keokuk Gr.

asperatus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 12, and Geo. Sur. Ill., vol. 7, p. 280, Keokuk Gr.

barrisi, see Cyathocrinus barrisi.

bayensis, see Scaphiocrinus bayensis.

bisselli, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 546, Kaskaskia Gr.

briareus, see Scaphiocrinus briareus.

buffaloensis, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 89, Ham. Gr.

burketi, see Scaphiocrinus burketi.

bursiformis, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 10, Burlington Gr.

caduceus, see Dendrocrinus caduceus.

calyculus, Hall, 1858, Geo. Sur. Iowa, p. 553, Burlington Gr.

calyx, Hall, 1879, Desc. New Spec. Foss., p. 10, and 11th Rep. Geo. Sur. Ind., p. 266, Niagara Gr.

carbonarius, see Graphiocrinus carbonarius.

FIG. 403.—Poterocrinus carinatus. Diagram.

carinatus, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 139, and Geo. Sur. Ill., vol. 3, p. 486, Burlington Gr.

clarkii, Williams, 1882, Proc. Acad. Nat. Sci. Phil., p. 21, Chemung Gr.

claytonensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 18, and Geo. Sur. Ill., vol. 7, p. 288, Warsaw Gr.

clytis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 16, and Geo. Sur. Ill., vol. 7, p. 294, St. Louis Gr.

columbiensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 22, and Geo. Sur. Ill., vol. 7, p. 293, Kaskaskia Gr.

concinus, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 26, and Geo. Sur. Ill., vol. 5, p. 490, Keokuk Gr.

coreyi, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.

cornellanus, Williams, 1882, Proc. Acad. Nat. Sci. Phil., p. 18, Chemung Gr.

corycia, Hall, 1863, 17th Rep. N. Y. St. Mus. Nat. Hist., p. 57, and Ohio Pal., vol. 2, p. 173, Waverly Gr.
cozanus, see Scaphiocrinus cozanus.
crineus, Hall, 1863, 17th Rep. N. Y. St. Mus. Nat. Hist., p. 56, and Ohio Pal., vol. 2, p. 172, Waverly Gr.
cultidactylus, see Scaphiocrinus cultidactylus.

FIG. 404.—*Poterioocrinus indianensis*. Diagram.

cylindricus, Lyon, 1860, Trans. Am. Phil. Soc., vol. 13, p. 458, Up. Held. Gr.
davisianus, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 226, Up. Held. Gr.
decadactylus, see Scaphiocrinus decadactylus.
diffusus, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 121, Ham. Gr.
dilatatus, see *Caeliocrinus dilatatus*.
divaricatus, Hall, 1860, Supp. to Geo. Sur. Iowa., p. 65, Warsaw Gr.
elsahensis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 88, Kinderhook Gr.
enormis, see *Cyathocrinus enormis*.
floralis, see *Zeacrinus floralis*.
fountainensis, Worthen, 1882, Bull. No. 1, Ill. Mus. Nat. Hist., p. 17, and Geo. Sur. Ill., vol. 7, p. 286, St. Louis Gr.
fusiformis, Hall, 1861, Desc. New Crinoidea, p. 6, and Bost. Jour. Nat. Hist., vol. 7, p. 302, Burlington Gr.
gracilis, see *Dendrocrinus gracilis*.
gregarius, Williams, 1882, Proc. Acad. Nat. Sci. Phil., p. 22, Chenung Gr.
hamiltonensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 7, and Geo. Sur. Ill., vol. 7, p. 273, Keokuk Gr.
hardinensis, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 533, St. Louis Gr.
hemisphericus, see *Eupachycrinus hemisphericus*.

loveyi, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
missouriensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 19, and Geo. Sur. Ill., vol. 7, p. 289, Warsaw Gr.
indentus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 122, Ham. Gr.
indianensis, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 155, and Geo. Sur. Ill., vol. 3, p. 515, Keokuk Gr.
iowensis, see Scaphiocrinus iowensis.
jesupi, Whitfield, 1881, Bull. No. 1, Am. Nat. Hist., p. 7, syn. for *P. swallowi*.
kaskaskiensis, see Scaphiocrinus kaskaskiensis.
keokuk, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 64, Keokuk Gr.
lasallensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 526, Coal Meas.
latidactylus, see Scaphiocrinus latidactylus.
lepidus, Hall, 1861, Desc. New Crin., p. 6, and Bost. Jour. Nat. Hist., vol. 7, p. 304, Burlington Gr.
longidactylus, Shumard, 1855, The name was preoccupied. See *P. missouriensis*.
macoupinensis, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 561, Up. Coal Meas.
mammiformis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 91, Warsaw Gr.
maniformis, see *Zeacrinus maniformis*.
meekanus, Shumard, 1855, Geo. Rep. Mo., p. 188, Burlington Gr.
milleri, Wetherby, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 330, Kaskaskia Gr.
missouriensis, Shumard, 1857, Trans. St. Louis Acad. Sci., p. 80, and Geo. Sur. Iowa, p. 669, St. Louis Gr.
montanensis, see Scaphiocrinus montanensis.
municipalis, Troost. Not defined.
nassa, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 120, Ham. Gr.
nauvoensis, see Scaphiocrinus nauvoensis.
nerens, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 121, Ham. Gr.
nettletothanus, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 227, Up. Held. Gr.
nodobasalis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 89, St. Louis Gr.

FIG. 405.—*Poterioocrinus missouriensis*.

norwoodensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
nycteus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
obuncus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
occidentalis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
sizocrius, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
occidentalis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
occidentalis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
okuwensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
oreales, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
otterensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
peculiaris, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
penicillatus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
perplexus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
pleias, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
pleias, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
popensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
posticus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
proboscis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
propinquus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
rhombifer, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
richfieldensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
rowleyi, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
rugosus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
salignoides, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
sculptus, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.
similis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.

FIG. 406.—*Poterioocrinus similis*, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 516, Keokuk Gr.

norwoodi, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 165, Kaskaskia Gr.
 nycteus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 120, Ham. Gr.
 obuncus, White, 1862, Proc. Bost. Soc. Nat. Hist., p. 10, Burlington Gr.
occidentalis, Owen & Shumard, see Agassizocrinus *occidentalis*.
occidentalis, Worthen, see Scaphiocrinus *occidentalis*.
okawensis, see Scaphiocrinus *okawensis*.
orestes, see Scaphiocrinus *orestes*.
otterensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 14, and Geo. Sur. Ill., vol. 7, p. 283, Keokuk Gr.
peculiaris, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 25, and Geo. Sur. Ill., vol. 7, p. 298, Kaskaskia Gr.
penicilliformis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 9, and Geo. Sur. Ill., vol. 7, p. 276, Keokuk Gr.
perplexus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 138, and Geo. Sur. Ill., vol. 5, p. 405, Burlington Gr.
pisiformis, see Arachniocrinus *pisiformis*.
pleias, Hall, 1863, 17th Rep. N. Y. St. Mus. Nat. Hist., p. 57, and Ohio Pal., vol. 2, p. 173, Waverly Gr.
popensis, see Scaphiocrinus *popensis*.
posticus, see Dendrocrinus *posticus*.
proboscidioides, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 518, St. Louis Gr.
propinquus, see Scaphiocrinus *propinquus*.
rhombiferus, see Barycrinus *rhombiferus*.
richfieldensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 15, and Geo. Sur. Ill., vol. 7, p. 285, Kinderhook Gr.
rowleyi, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 90, Kaskaskia Gr.
rugosus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 223, Coal Meas.
saliginoides, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 10, Burlington Gr.
salteri, see Scaphiocrinus *salteri*.
sculptus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 21, and Geo. Sur. Ill., vol. 7, p. 292, Kaskaskia Gr.

FIG. 406.—*Poteriocrinus subimpressus*.

similis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 23, and Geo. Sur. Ill., vol. 7, p. 295, Keokuk Gr.

simplex, Lyon, 1869, Trans. Am. Phil. Soc., vol. 13, p. 458, Up. Held. Gr.
solidus, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 140, Burlington Gr.
spinobrachiatus, Scaphiocrinus *brachiatus*.
spinuliferus, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 27, and Geo. Sur. Ill., vol. 8, p. 86, Kaskaskia Gr.
spinuliferus, Worthen, (in press,) see Zeacrinus *spinuliferus*.
spinosus, see Zeacrinus *spinosus*.
subimpressus, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 13, and Geo. Sur. Ill., vol. 3, p. 485, Burlington Gr.
subramulosus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 14, and Geo. Sur. Ill., vol. 7, p. 284, Keokuk Gr.
subtumidus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 159, Kaskaskia Gr.
swalovi, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 394, and Geo. Sur. Ill., vol. 2, p. 183, Burlington Gr.
talboti, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 7, and Geo. Sur. Ill., vol. 7, p. 287, St. Louis Gr.
tentaculatus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 10, and Geo. Sur. Ill., vol. 7, p. 277, Keokuk Gr.
tenuibrachiatus, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 138, and Geo. Sur. Ill., vol. 3, p. 484, Burlington Gr.
tenuidactylus, Meek & Worthen, see Scaphiocrinus *tenuidactylus*.
tenuidactylus, FIG. 407.—*Poteriocrinus tenuibrachiatus*. Diagram.

tenuidactylus, FIG. 407.—*Poteriocrinus tenuibrachiatus*. Diagram.
 Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 6, and Geo. Sur. Ill., vol. 7, p. 271, Keokuk Gr.
 Wachsmuth says this is a Scaphiocrinus, and he has proposed to call it Scaphiocrinus *obscurus*.

ulrichi, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 87, Keokuk Gr.

tumidus, see Agassizocrinus *tumidus*.

validus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 18, and Geo. Sur. Ill., vol. 7, p. 287, Warsaw Gr.

vanhornei, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 517, St. Louis Gr.

varsoviensis, see Scaphiocrinus *varsoviensis*.

ventricosus, see Coliocrinus *ventricosus*.

venustus, see Scaphiocrinus *venustus*.

FIG. 408.—*Poteriocrinus ulrichi*.

- verticillus, Hall, 1862, 15th R-p. N. Y. St. Mus. Nat. Hist., p. 94, Ham. Gr.
wachsmuthi, Meek & Worthen, see *Graphiocrinus wachsmuthi*.
wachsmuthi, Wetherby, 1880, (*Seyta-
 loerinus wachsmuthi*.) Jour. Cin. Soc.
 Nat. Hist., vol. 3, p. 155, Kaskaskia Gr.
wetherbyi, S. A. Miller, 1879, Jour. Cin.
 Soc. Nat. Hist., vol. 2, p. 36, Kas-
 kaskia Gr.
zethus, Williams, 1882, Proc. Acad. Nat.
 Sci., p. 27, Chemung Gr.
PROTASTER, Forbes, 1849, Mem. Geo. Sur.
 (Great Britain, Decade 1. [Ety. *protos*,
 first; *aster*, star.] Disk circular, com-
 posed of squamiform plates; rays
 flexuous, composed of two series of
 ambulacral plates, bordered by spinous
 adambulacral ones; oral plates five.
 Type *P. miltoni*.
barrisi, see *Onychaster barrisi*.

FIG. 400.—*Protaster flexuosus*. forbesi, Hall,
 1859, Pal.

- N. Y., vol. 3, p. 134, Low. H.-ld. Gr.
granuliferus, Meek, 1872, Am. Jour. Sci.,
 3d ser., vol. 3, p. 274, and Ohio Pal.,
 vol. 1, p. 68, Hud. Riv. Gr.
gregarius, Meek & Worthen, 1869, Proc.
 Acad. Nat. Sci. Phil., p. 169, and Geo.
 Sur. Ill., vol. 5, p. 509, Keokuk Gr.
miamiensis, S. A. Miller, 1882, Jour. Cin.
 Soc. Nat. Hist., vol. 5, p. 116, Hud.
 Riv. Gr.
stellifer, Ringueberg, 1886, Bull. Buff. Soc.
 Nat. Sci., vol. 5, p. 7, Niagara Gr.
Protasterina, syn. for *Protaster*.
fimbriata, syn. for *Protaster flexuosus*.
PTEROTOCRINUS, Lyon & Casseday, 1860, Am.
 Jour. Sci., vol. 29, p. 68. [Ety. *pteros*,
 feathered; *krinos*, lily.] Calyx saucer-
 shaped, wider than high; vault high,
 with five wing-like processes that char-
 acterize this genus; basals 2; radials 1
 or 2 x 5, the second being small; second-
 ary radials 1 x 10; tertiary radials 2 or
 3 x 20; azygous interradial 1; arms
 20, reaching only to the vault; column
 round. Type *P.*
capitalis.
acutus, Wetherby,
 1879, Jour. Cin.
 Soc. Nat. Hist., vol.
 2, p. 134, Kaskas-
 kia Gr.
bifurcatus, Wether-
 by, 1879, Jour.
 Cin. Soc. Nat. Hist., vol. 2, p. 136,
 Kaskaskia Gr.

FIG. 410.—*Pterotoerinus
 chesterensis*. Dia-
 gram.

- capitalis*, Lyon, 1857, (*Astrocrinus capi-
 talis*.) Geo. Sur. Ky., vol. 3, p. 472, Kas-
 kaskia Gr.
chesterensis, Meek & Worthen, 1860,
 (*Actinocrinus chesterensis*.) Proc. Acad.
 Nat. Sci. Phil., p. 383, and Geo. Sur.
 Ill., vol. 2, p. 292, Kaskaskia Gr.
coronarius, Lyon, 1857, (*Astrocrinus cor-
 onarius*.) Geo. Sur. Ky., vol. 3, p. 476,
 Kaskaskia Gr.

FIG. 411.—*Pterotoerinus crassus*. Diagram.

- crassus*, Meek & Worthen, 1860, (*Dicho-
 crinus crassus*.) Proc. Acad. Nat. Sci.
 Phil., p. 382, and Geo. Sur. Ill., vol. 2,
 p. 290, Kaskaskia Gr.
depressus, Lyon & Casseday, 1860, Am.
 Jour. Sci., vol. 29, p. 68, and Geo. Sur.
 Ill., vol. 5, p. 559, Kaskaskia Gr.
protuberans, Hall, 1858, (*Dichocrinus
 protuberans*.) Geo. Sur. Iowa, p. 689,
 Kaskaskia Gr.
pyramidalis, Lyon & Casseday, 1860,
 Am. Jour. Sci., vol. 29, p. 69, Kaskas-
 kia Gr.
rugosus, Lyon & Casseday, 1860, Am.
 Jour. Sci., vol. 29, p. 71, Kaskas-
 kia Gr.
sexlobatus, see *Talarocrinus sexlobatus*.
spatulatus, Wetherby, 1879, Jour. Cin.
 Soc. Nat. Hist., vol. 2, p. 137, Kaskas-
 kia Gr.
Pti onaster, Hall, 1868, syn. for *Palaeocoma*.
princeps, see *Palaeocoma princeps*.
Ptychocrinus, Wachsmuth & Springer, 1886,
 Revis. Palaeocrinoidea, pt. 3, p. 99, syn.
 for *Gaurocrinus*.
PYCNOCRINUS, S. A. Miller, 1883, Jour. Cin.
 Soc. Nat. Hist., vol. 6, p. 231. [Ety.
puknos, dense; *krinos*, lily.] Only
 small, cup-shaped; basals
 5; radials 3 x 5; regular
 interradials 3; arms 10,
 sometimes dividing after
 becoming free. Type *P.*
shafferi.
germanus, S. A. Miller, 1880,
 (*Glyptocrinus shafferi* var.
germanus.) Jour. Cin. Soc. Nat. Hist.,
 vol. 3, p. 233,
 Hud. Riv. Gr.
shafferi, S. A. Miller, 1875, (*Glyptocrinus
 shafferi*.) Cin. Quar. Jour. Sci., vol. 2,

FIG. 412.—*Pycnocrinus
 germanus*.

FIG. 413.—*Pycnocrinus
 shafferi*.

- Pycnogonich-
 Retinocrinus*
 Decade
 lily.] C
 very p
 radials
 second
 interra
 large c
 extend
 column
 fimbriatu
 Decade
 gracilis, V
 angulan

FIG. 416.—*Retinocrinus
 stellatus*.

- 1883, G
 Rep. N.
 Trenton
RUONOCRINUS
noidea,
krinos,
 wider th
 arm bas
 5; prim
 radials 1
 rated, a
 interlocl
 wide, p
 orifice e
 umn rou
 asperatus,
 Decade 4
 barrisi, H
 and Jour
 p. 322, B

p. 277, and Jour. Cin. Soc. Nat. Hist., vol. 3, p. 233, Hud. Riv. Gr.

FIG. 413.—Pycnoerinus shafferli.

FIG. 414.—Pycnoerinus shafferli. Enlarged $2\frac{1}{4}$ diam.

FIG. 415.—Pycnoerinus shafferli. Column coiled around a column of Glyptocrinus.

Pygorthynchus gouldi. Not recognized.

RETIOCRINUS, Billings, 1858, Can. Org. Rem., Decade 4, p. 63. [Ety. *retium*, net; *krinon*, lily.] Calyx basin-shaped; radial ridges very prominent; basals 5, large; subradials 5, large; primary radials 4×5 ; secondary radials 4 to 6×10 ; plates in interradial areas, numerous, with a large central row in the azygous area extending up the side of a ventral tube; column round. Type *R. stellaris*.

fimbriatus, Billings, 1859, Can. Org. Rem., Decade 4, p. 65, Hud. Riv. Gr.

gracilis, Wetherby, syn. for *Glaurocrinus angularis*.

FIG. 416.—Retiocrinus stellaris.

stellaris, Billings, 1859, Can. Org. Rem., Decade 4, p. 64, Trenton Gr.

RHAPHIANOCRINUS, Wachsmuth & Springer, 1885, Revis. Paleocrinoidea, pt. 3, p. 98. [Ety. *rhaphe*, ridge; *krinon*, lily.] Calyx short; basals 5, small; subradials 5; primary radials 3×5 ; interradial plates numerous; column round. Type *R. subnodosus*.

subnodosus, Walcott, 1883, Glyptocrinus subnodosus, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 208, Trenton Gr.

RHODOCRINUS, Miller, 1821, Nat. Hist., Crinoidea, p. 106. [Ety. *rhodon*, rose; *krinon*, lily.] Body subglobose, often wider than high, constricted near the arm bases; basals 5, small; subradials 5; primary radials 3 to 4×5 ; secondary radials 1 to 3×10 ; arms widely separated, and composed of two rows of interlocking plates; interradial areas wide, plates large; vault depressed; orifice excentric and protruding; column round. Type *R. verus*.

asperatus, Billings, 1859, Can. Org. Rem., Decade 4, p. 27, Chazy Gr.

barrisi, Hall, 1861, Desc. New Crin., p. 9, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 322, Burlington Gr.

barrisi var. *divergens*, Hall, 1861, Desc. New Crin., p. 9, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 322, Burlington Gr.

coxanus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 29, and Geo. Sur. Ill., vol. 7, p. 305, Keokuk Gr.

FIG. 417.—Rhodocrinus. Diagram.

gracilis, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 127, Ham. Gr.

halli, Lyon, 1861, Proc.

Acad. Nat. Sci. Phil.,

p. 412, Low. Held. Gr.

kirbyi, Wachsmuth &

Springer, (in press,)

(Geo. Sur. Ill., vol. 8,

p. 180, Kinderhook

Gr.

melissa, see *Lyriocrinus*

melissa.

microbasalis, see Arch-

aeocrinus *microba-*

salis.

nanus, Meek & Wor-

then, 1866, Proc. Acad. Nat. Sci. Phil.,

FIG. 418.—Rhodocrinus nanus.

FIG. 419.—Rhodocrinus nanus. Diagram.

p. 254, and Geo. Sur. Ill., vol. 3, p. 476, Burlington Gr.

nodulosus, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 126, Ham. Gr.
pyriformis, see *Archaeocrinus pyriformis*.
rectus, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 368, Niagara Gr.
spinosus, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 127, Ham. Gr.
varsoviensis, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 80, Warsaw Gr.
vesperalis, White, 1880, Proc. U. S. Nat.

wortheni, Hall, 1858, Geo. Sdr. Iowa, p. 556, Burlington Gr.
Saccoocrinus, Hall, 1852, Pal. N. Y., vol. 2, p. 205. [Ety. *sakkos*, bag; *krinon*, lily.] Calyx large, urn-shaped; basals 3; primary radials 3x5; secondary radials 1 to 4x10; tertiary radials, in some species; regular interradianals 10 to 17; vault depressed, opening subcentral; arms 10 to 30; column round. Type *S. speciosus*.

FIG. 420.—*Saccoocrinus amplius*. Diagram.

Mus., vol. 2, p. 259, and Cont. to Pal. No. 6, p. 129, Up. Coal Meas.
wachsmuthi, Hall, 1861, Desc. New Crin., p. 18, Burlington Gr.

FIG. 421.—*Rhodoocrinus watersianus*.

watersianus, *wachsmuth* & Springer, (in press,) (Geo. Sur. Ill., vol. 3, p. 184, Kinderhook, Gr.)

whitii, Hall, 1861, Desc. New Crin., p. 9, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 324, Burlington Gr.

whitii var. *burlingtonensis*, Hall, 1861, Desc. New Crin., p. 9, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 325, Burlington Gr.

amplius, Meek & Worthen, 1861, (Actinocrinus *amplius*.) Proc. Acad. Nat. Sci. Phil., p. 133, and Geo. Sur. Ill., vol. 3, p. 470, Burlington Gr.

christyi, Hall, 1863, (Actinocrinus, *christyi*.) Trans. Alb. Inst., vol. 4, p. 196, and 28th Rep. N. Y. Mus. Nat. Hist., p. 127, Niagara Gr.

eganii, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 173, Niagara Gr.

infelix, Winchell & Marcy, 1865, (Megistocrinus *infelix*.) Mem. Bost. Soc. Nat. Hist., p. 110, Niagara Gr.

FIG. 422.—*Saccoocrinus christyi*.

marco
 (Me
 Soc.
 neclis,
 crin
 Hist
 ornatu
 vol.
 pyrifo
 Soc.
 agar
 semira
 St. M
 specio
 205,
 tennes
 field
 agar
 urnifor
 Soc.
 agar
 whitfie
 crinu
 SCAPHIO
 p. 55
 lily.]
 radia
 diala
 arms
 proje
 Type
 Sprin
 and
 tomu
 abnorm
 vol. 6
 agina,
 Nat.
 aqualis
 and
 kuk G
 bayensi
 Acad.
 Geo.
 kia G
 briareus
 briare
 Hist.,
 279, K
 burketi,
 burke
 Hist.,
 270, K
 carbonar
 arius.
 carinatu
 8, and
 7, p. 3
 clio, Me
 Nat. S
 vol. 5,
 coreyi,
 Acad.
 Geo.
 kuk G
 coxanus,
 Hist.,
 269, K

marconianus, Winchell & Marcy, 1865, (Megistocrinus marconianus,) Mem. Bost. Soc. Nat. Hist., p. 87, Niagara Gr.
 necla, Winchell & Marcy, 1865, (Megistocrinus necla,) Mem. Bost. Soc. Nat. Hist., p. 110, Niagara Gr.
 ornatus, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 126, Niagara Gr.
 pyriformis, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 81, Niagara Gr.
 semiradiatus, Hall, 1867, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 370, Niagara Gr.
 speciosus, Hall, 1852, Pal. N. Y., vol. 2, p. 205, Niagara Gr.
 tennesseensis, Troost, Ms., Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 125, Niagara Gr.
 urniformis, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 170, Niagara Gr.
 whitfieldi, Hall, 1867, synonym for Saccocrinus christyi.

SCAPHIOCRINUS, Hall, 1858, Geo. Sur. Iowa, p. 550. [Ety. *scaphion*, skiff; *krinon*, lily.] Calyx obconoidal; basals 5; subradials 5; radials 2 x 5; regular interradials 0; zygous interradials 1 to 6; arms 10, simple or bifurcating, plates projecting laterally; sutures gaping. Type S. simplex. Wachsmuth & Springer refer the type to Graphiocrinus and substitute, as the type S. dichotomus.

abnormis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 519, St. Louis Gr.

regina, Hall, 1863, 17th Rep. N. Y. Mus. Nat. Hist., p. 57, Waverly Gr.

aqualis, Hall, 1861, Desc. New Crin., p. 8, and Geo. Sur. Ill., vol. 5, p. 494, Keokuk Gr.

bayensis, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 157, and Geo. Sur. Ill., vol. 5, p. 550, Kaskaskia Gr.

briareus, Worthen, 1882, (Poteriocrinus briareus,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 12, and Geo. Sur. Ill., vol. 7, p. 279, Keokuk Gr.

burketi, Worthen, 1882, (Poteriocrinus burketi,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 5, and Geo. Sur. Ill., vol. 7, p. 270, Keokuk Gr.

carbonarius, see Graphiocrinus carbonarius.

carinatus, Hall, 1861, Desc. New Crin., p. 8, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 310, Burlington Gr.

elio, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 144, and Geo. Sur. Ill., vol. 5, p. 408, Burlington Gr.

coreyi, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 148, and Geo. Sur. Ill., vol. 5, p. 494, Keokuk Gr.

coxanus, Worthen, 1882, (Poteriocrinus coxanus,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 43, and Geo. Sur. Ill., vol. 7, p. 269, Keokuk Gr.

cultidactylus, Hall, 1860, (Poteriocrinus cultidactylus,) Supp. to Geo. Sur. Iowa, p. 62, and Geo. Sur. Ill., vol. 7, p. 301, Burlington Gr.

dactyliformis, Hall, 1858, Geo. Sur. Iowa, p. 670, St. Louis Gr.

decabrachiatas, Hall, 1858, Geo. Sur. Iowa, p. 679, St. Louis Gr.

decadactylus, Meek & Worthen, 1860, (Poteriocrinus decadactylus,) Proc. Acad. Nat. Sci. Phil., p. 394, and Geo. Sur. Ill., vol. 2, p. 238, Keokuk Gr.

delicatus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 144, and Geo. Sur. Ill., vol. 5, p. 407, Burlington Gr.

depressus, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 27, and Geo. Sur. Ill., vol. 5, p. 492, Keokuk Gr.

dichotomus, Hall, 1858, Geo. Sur. Iowa, p. 553, Burlington Gr.

divaricatus, Hall, 1860, Supp. to Geo. Sur. Iowa, p. 65, Burlington Gr.

doris, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 312, Burlington Gr.

elegantulus, Wachsmuth & Springer, 1886, Revis. Palaeocrinoidea pt. 3, p. 237. Proposed instead of Poteriocrinus asper, Worthen, but the latter name was not pre-occupied.

fiscellus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 146, and Geo. Sur. Ill., vol. 5, p. 424, Burlington Gr.

gibsoni, White, 1878, Proc. Acad. Nat. Sci. Phil., p. 31, and Cont. to Pal., No. 8, p. 161, Keokuk Gr.

globosus, Wachsmuth & Springer, (in press) Geo. Sur. Ill., vol. 8, p. 196, Kinderhook Gr.

gurleyi, White, 1878, Proc. Acad. Nat. Sci. Phil., p. 32, and Cont. to Pal., No. 8, p. 162, Keokuk Gr.

halli, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

halleri, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 308, Burlington Gr.

FIG. 423.—Scaphiocrinus decadactylus. Diagram.

FIG. 424.—Scaphiocrinus elegantulus.

FIG. 425.—Scaphiocrinus globosus. Diagram.

Actino-

crinus,

vol. 3.

Saccocrinus

christyi.

(Megistocrinus)

Soc. Nat. Hist.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

28
32
36
40
45
50
56
63
71
80
90
100
112
125
140
160
180
200
224
256
288
320
360
400
450
500
560
630
710
800
900
1000

1.0
1.1
1.2
1.5
2.0
2.5
3.2
4.0
5.0
6.3
8.0
10.0
12.5
16.0
20.0
25.0
32.0
40.0
50.0
63.0
80.0
100.0
125.0
160.0
200.0
250.0
320.0
400.0
500.0
630.0
800.0
1000.0

- huntsvillae, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 534, St. Louis Gr.
- internodius, Hall, 1858, Geo. Sur. Iowa, p. 679, St. Louis Gr.
- iowensis, Worthen, 1882, (Poteriocrinus iowensis,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 6, and Geo. Sur. Ill., vol. 7, p. 272, Keokuk Gr.
- juvenis, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 146, and Geo. Sur. Ill., vol. 5, p. 417, Burlington Gr.
- kaskaskiensis, Worthen, 1882, (Poteriocrinus kaskaskiensis,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 27, and Geo. Sur. Ill., vol. 7, p. 300, Kaskaskia Gr.
- latidactylus, Worthen, 1882, (Poteriocrinus latidactylus,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 8, and Geo. Sur. Ill., vol. 7, p. 275, Keokuk Gr.
- liliformis, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 138, Burlington Gr.
- liriope, Hall, 1863, 17th Rep. N. Y. Mus. Nat. Hist., p. 58, Waverly Gr.
- longidactylus, McCheaney, 1860, New Pal. Foss., p. 7, Kaskaskia Gr.
- macadamsi, see Graphiocrinus macadamsi.
- macrodactylus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 140, and Geo. Sur. Ill., vol. 5, p. 415, Burlington Gr.
- montanensis, Meek, 1872, (Poteriocrinus montanensis,) Hayden's Geo. Sur. Terr., p. 469, and Cont. to Pal., No. 6, p. 128, Sub-arconiferous.
- nanus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 141, and Geo. Sur. Ill., vol. 5, p. 423, Burlington Gr.
- navoocensis, Worthen, 1882, (Poteriocrinus navoocensis,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 13, and Geo. Sur. Ill., vol. 7, p. 282, Keokuk Gr.
- nodobrachiatus, Hall, 1861, Desc. New Crin., p. 8, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 314, Keokuk Gr.
- notabilis, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 148, and Geo. Sur. Ill., vol. 5, p. 410, Burlington Gr.
- obscurus, Wachsmuth & Springer, 1886, Revis. Paleocrinoidea, pt. 3, p. 236, Keokuk Gr.
- occidentalis, Worthen, 1882, (Poteriocrinus occidentalis,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 10, and Geo. Sur. Ill., vol. 7, p. 278, Keokuk Gr.
- okawensis, Worthen, 1882, (Poteriocrinus okawensis,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 24, and Geo. Sur. Ill., vol. 7, p. 296, Kaskaskia Gr.
- orbicularis, see Eupachycrinus orbicularis.
- orestes, Worthen, 1882, (Poteriocrinus orestes,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 7, and Geo. Sur. Ill., vol. 7, p. 273, Keokuk Gr.
- penicillus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 142, and Geo. Sur. Ill., vol. 5, p. 414, Burlington Gr.
- popensis, Worthen, 1882, (Poteriocrinus popensis,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 23, and Geo. Sur. Ill., vol. 7, p. 296, Kaskaskia Gr.
- propinquus, Worthen, 1882, (Poteriocrinus propinquus,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 28, and Geo. Sur. Ill., vol. 7, p. 299, Kaskaskia Gr.
- ramulosus, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 307, Burlington Gr.
- randolphensis, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 551, Kaskaskia Gr.
- robustus, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 315, Keokuk Gr.
- rudis, see Graphiocrinus rudis.
- rusticellus, White, 1863, Proc. Bost. Soc. Nat. Hist., vol. 7, p. 505, Burlington Gr.
- salteri, Worthen, 1882, (Poteriocrinus salteri,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 13, and Geo. Sur. Ill., vol. 7, p. 291, Kaskaskia Gr.
- scalaris, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 145, and Geo. Sur. Ill., vol. 5, p. 421, Burlington Gr.
- scoparius, Hall, 1858, Geo. Sur. Iowa, p. 680, Kaskaskia Gr.
- simplex, see Graphiocrinus simplex.
- spinifer, Wetherby, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 157, Kaskaskia Gr.
- spinobrachiatus, see Graphiocrinus spinobrachiatus.
- spinobrachiatus, Worthen, 1882, (Poteriocrinus spinobrachiatus,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 20, and Geo. Sur. Ill., vol. 7, p. 290, Kaskaskia Gr.
- striatus, see Graphiocrinus striatus.
- subcarinatus, Hall, 1863, 17th Rep. N. Y. Mus. Nat. Hist., p. 58, Waverly Gr.

FIG. 426.—Scaphiocrinus tenuidactylus. Diagram.

- subtortuosus, Hall, 1863, 17th Rep. N. Y. Mus. Nat. Hist., p. 59, Waverly Gr.
- tenuidactylus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 156, and Geo. Sur. Ill., vol. 3, p. 490, Burlington Gr.

FIG. 427.
nus

striat
316
and
SCHÖEN
Acc
Ill.

tethys, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 143, and Geo. Sur. Ill.; vol. 5, p. 419, Burlington Gr.

tortuosus, see *Graphiocrinus tortuosus*.
unicus, Hall, 1861, Desc. New Crin., p. 8, and Geo. Sur. Ill., vol. 5, p. 493, Keokuk Gr.

varsoviensis, Worthen, 1882, (*Poteriocrinus varsoviensis*,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 20, and Geo. Sur. Ill., vol. 7, p. 290, Warsaw Gr.

venustus, Worthen, 1882, (*Poteriocrinus venustus*,) Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 24, and Geo. Sur. Ill., vol. 7, p. 297, Kaskaskia Gr.

wachsmuthi, see *Graphiocrinus wachsmuthi*.

whitti, Hall, 1861, Desc. New Crin., p. 7, and Jour. Bost. Soc. Nat. Hist., vol. 7, p. 306, Burlington Gr.

SCHIZOBLASTUS, Etheridge & Carpenter, 1882, Ann. & Mag. Nat. Hist., vol. 9, p. 243. [Ety. *schiza*, cleft; *blastos*, bud.] Calyx in form like *Granatocrinus*; basals confined to the base, sometimes visible, in a side view; deltoids always visible in a side view; ambulacra narrow and sublinear, extending the height of the calyx; lancet-plates nearly concealed by the side plates; latter from 20 to 80 in number; 1 to 4 hydropore folds on each side of an ambulacrum; spiracles minute linear slits between the lancet-plate and the deltoid ridges; surface ornamented with striae. Type *S. sayi*.

melonoides, Meek & Worthen, 1869, (*Granatocrinus melonoides*,) Proc. Acad. Nat. Sci. Phil., p. 88, and Geo. Sur. Ill., vol. 5, p. 468, Burlington Gr.

sayi, Shumard, 1855, (*Pentremites sayi*,) Geo. Rep. Mo., p. 185, Burlington Gr.

FIG. 427.—*Schizocrinus nodosus*.

SCHIZOCRINUS, Hall, 1847, Pal. N. Y., vol. 1, p. 81. [Ety. *schiza*, cleft; *krinon*, lily.] Basals 5; primary radials 3 x 5; secondary radials 2 x 10; interradials 5 or more; arms short, branching, bearing pinnules; column round. Type *S. nodosus*.

nodosus, Hall, 1847, Pal. N. Y., vol. 1, p. 81, Trenton Gr.

striatus, Hall, 1847, Pal. N. Y., vol. 1, p. 316, Trenton Gr. Probably belongs to another genus.

SCHENASTER, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 449, and Geo. Sur. Ill., vol. 2, p. 277. [Ety. *schenos*, rope;

aster, star.] Pentagonal disk, with angles produced into rays; margins between rays concave and spinous; plates alternating on dorsal side of rays, and on ventral side of disk imbricating inward and laterally toward the ambulacra; furrows wide, deep, bordered with a single row of adambulacra, which become the marginal plates of the free rays. Type *S. fimbriatus*.

fimbriatus, Meek & Worthen, 1860, (*Palaeasterina fimbriata*,) Proc. Acad. Nat. Sci. Phil., p. 449, and Geo. Sur. Ill., vol. 2, p. 278, St. Louis Gr.

wachsmuthi, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 259, and Geo. Sur. Ill., vol. 3, p. 499, Burlington Gr.

Scyphocrinus, Hall, 1847, Pal. N. Y., vol. 1. Preoccupied, by Zenker, in 1839. See *Cupulocrinus*.

heterocostalis, see *Cupulocrinus heterocostalis*.

Scytalocrinus, Wachsmuth & Springer, 1879, Proc. Acad. Nat. Sci. Phil. A division of *Poteriocrinus* of less than generic importance, with *P. robustus* as the type.

wachsmuthi, see *Poteriocrinus wachsmuthi*.

SIPHONOCRINUS, S. A. Miller, 1888, Am. Geol., vol. 1, p. 263. [Ety. *siphon*, bent tube; *krinon*, lily.] Basals 3 (?) small.

FIG. 428.—*Siphonocrinus nobilis*; lateral view of an internal cast.

Wachsmuth says there are 5; primary radials 3 x 5; first interradials nearly as large as primary radials, and succeeded by two smaller ones, and

these by three or more; first azygous plate as large as the primaries; it rests upon the basals and is succeeded by three plates; the following ranges have more plates and cover an expanded azygous side; vault very large, high, and bears a proboscis either projected upward or recumbent; surface of plates ornamented. Type *S. nobilis*.

armosus, McChesney, 1861. (*Eucalyptocrinus armosus*.) New. Pal. Foss., p. 95, and 20th Rep. N. Y. St. Mus. Nat. Hist., p. 373, Niagara Gr.

nobilis, Hall, 1861. (*Glyptocrinus nobilis*.) Geo. Sur. Wis., p. 21, and 20th Rep. N. Y. St. Mus. Nat. Hist., p. 328, Niagara Gr.

Sphaerocrinus, Meek & Worthen, 1866. The name was preoccupied, by Roemer. See *Caelocrinus*.

SPHEROCYSTITES, Hall, 1863, Pal. N. Y., vol. 3, p. 130. [Ety. *sphaira*, sphere; *kusta*, bladder.] Spheroidal, wider than high; arms, in two principal pairs, with numerous bifurcations; brachial sulci obliquely lobed; mouth apical; opening subapical; ovarian opening on the summit; basal plates 4, others unknown. Type *S. multifasciatus*.

multifasciatus, Hall, 1859, Pal. N. Y., vol. 3, p. 130, Low. Held. Gr.

Squamaster, Ringueberg, 1886, Bull. Buf. Soc. Nat. Hist., vol. 5, p. 5.

STEGANOCRINUS, Meek & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 195. [Ety. *steganos*, covered; *krinon*, lily.] General form like *Actinocrinus*: basals 3; primary

concinnus, Shumard, 1855, (*Actinocrinus concinnus*.) Geo. Sur. Mo., p. 189, Burlington Gr.

FIG. 430.—*Steganocrinus pentagonus*. Vault and part of the rays; *a* and *d* showing structure of the rays.

pentagonus, Hall, 1858, (*Actinocrinus pentagonus*.) Geo. Sur. Iowa, p. 577, Burlington Gr.

sculptus, Hall, 1858, (*Actinocrinus sculptus*.) Geo. Sur. Iowa, p. 582, Burlington Gr.

STEMMATOCRINUS, Trautschold, 1867, Crin. d. jungeren Bergkalkes b. Moskau, p. 28. [Ety. *stemma*, wreath; *krinon*, lily.] Calyx low, cup-shaped; basals 5, anchylosed; subradials 5; radials twice as wide as high; brachials 1 x 5; arms heavy. Type *S. cornuus*. This is closely related to *Erismocrinus* and *Eupachyrcrinus*.

trautscholdi, Wachs-muth & Springer, 1886, Revis. Palaeocrinoidea, pt. 3, p. 256, Keokuk Gr.

STENASTER, Billings, 1858, Can. Org. Rem., Decade 3, p. 77. [Ety. *stenos*, narrow; *aster*, star.] Disk small, rays extended, flexible; dorsal side covered with small

FIG. 431.—*Steganocrinus sculptus*. Diagram of a ray and transverse sections.

FIG. 429.—*Steganocrinus concinnus*.

radials 3 x 5; secondary radials 1 x 2 x 5, in each ray; regular interradials 3 to 6 x 4; azygous interradials 3 to 10 or more; vault elevated, with long subcentral tube; arms bifurcating; column round. Type *S. pentagonus*.

araneolus, Meek & Worthen, 1860. (*Actinocrinus araneolus*.) Proc. Acad. Nat. Sci. Phil., p. 387, and Geo. Sur. Ill., vol. 2, p. 198, Burlington Gr.

FIG. 433.—*anocrinus latus*.

cies
but i
pentad
1879,
pen
Desc
Foss.
11th
Sur.
Niag
pulche

plates; adambulacrals square or oblong; orals 10. Type *S. salteri*.

FIG. 432.—*Stenaster grandis*.

grandis, Meek, 1872, Am. Jour. Sci., 3d ser., vol. 3, p. 258, and Ohio Pal., vol. 1, p. 66, Hud. Riv. Gr.

huxleyi, Billings, 1865, Pal. Foss., vol. 1, p. 213, Quebec Gr.

pulchellus, Billings, 1857, (Palæaster *pulchellus*), Geo. Sur. Can., p. 292, and Can. Org. Rem., Decade 3, p. 79, Trenton Gr.

salteri, Billings, 1858, Can. Org. Rem., Decade 3, p. 78, Trenton Gr.

Stenocrinus, Wachsmuth & Springer, 1885, Palæocrinoides, pt. 3, p. 207, syn. for *Heterocrinus*.

STEPHANOCRINUS, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 278. [Ety. *stephanos*, coronet; *krinon*, lily.] A blastoid, with 3 basals, 5 fork pieces or radials, and 5 orals; aperture subcentral; ambulacral appendages, but thus far the hydrospires are unknown. Type *S. angulatus*. Some authors refer this genus to the Palæocrinoides.

FIG. 433.—*Stephanocrinus angulatus*.

angulatus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 279, and Pal. N. Y., vol. 2, p. 212, Niagara Gr.

gemmaformis, Hall, 1852, Pal. N. Y., vol. 2, p. 215, Niagara Gr.

osgoodensis, S. A. Miller, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 116, Niagara Gr. Wachsmuth has said this species was described from internal casts, but it was not.

pentalobus, Hall, 1879, (Codaster *pentalobus*), Desc. New Spec. Foss., p. 13, and 11th Rep. Geo. Sur. Ind., p. 280, Niagara Gr.

pulchellus, Miller & Dyer, 1878, (Codaster

FIG. 434.—*Stephanocrinus pulchellus*, instead of Codaster *pulchellus*, as shown by fig. 288.

pulchellus, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 35, Niagara Gr.

STEROCRINUS, Barris, 1879, Proc. Dav. Acad. Sci., vol. 2, p. 282. [Ety. *stereos*, firm; *krinon*, lily. Distinguished from *Dolatocrinus* by having 2x5 instead of 3x5 primary radials; one large interradial succeeded by a smaller one, and this by smaller ones, within the depressions, between the arm bases. Type *S. triangulatus*.

triangulatus, Barris, 1879, Proc. Dav. Acad. Sci., vol. 2, p. 283, Up. Held. Gr.

FIG. 435.—*Strotocrinus perumbrosus*. Under side of canopy with calyx broken away.

triangulatus var. *liberatus*, Barris, 1879, Proc. Dav. Acad. Sci., vol. 2, p. 284, Up. Held. Gr.

STROBILOCYSTITES, White, 1876, Proc. Acad. Nat. Sci. Phil., p. 28. [Ety. *strobilos*, pine cone; *kystis*, bladder.] Subspherical; 3 pectinated rhombs, two above the middle and one below; ovarian aperture below the summit; 4 principal arm grooves extending below the middle, and 4 secondary grooves. Type *S. calvini*.

calvini, White, 1876, Proc. Acad. Nat. Sci. Phil., p. 28, Devonian.

STROTOCRINUS, Meek & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 188. [Ety. *strotos*, spread; *krinon*, lily.] Calyx bowl-shaped, with vault spreading beyond like a canopy; basals 3; primary radials 3x5; secondary radials 1 or 2x10, succeeded by tertiary and other divisions, which, with the interaxillaries and interbranchials, unite to form the under side of a greatly expanded hor-

ocrinus
9, Bur-

fault and
structure of

ocrinus
p. 577,

izontal disk, completely isolating the azygous and interradial areas, from the vault, and supporting the free, ascend-

FIG. 486.—*Strotocrinus regalis*.

ing arms around its margin; interradials 9 or 10 or more; azygous interradials 9 to 13 or more, the first one resting on the basals; vault depressed, opening subcentral; arms 30 to 72 or more; column round. Type *S. perumbrosus*.
asperrimus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 160, and Geo. Sur. Ill., vol. 5, p. 349, Burlington Gr.

FIG. 487.—*Strotocrinus regalis*. Diagram, $\frac{1}{4}$ diam.

bloomfieldensis, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 258, and vol. 4, p. 78, Up. Burlington or Keokuk Gr.
dilatatus, see *Physetocrinus dilatatus*.

ectypus, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 159, and Geo. Sur. Ill., vol. 5, p. 353, Burlington Gr.

glyptus, Hall, 1860, (Actinocrinus *glyptus*,) Supp. to Geo. Sur. Iowa, p. 2, Burlington Gr.

perumbrosus, Hall, 1860, (Actinocrinus *perumbrosus*,) Supp. to Geo. Sur. Iowa, p. 7, Burlington Gr.

regalis, Hall, 1860, (Actinocrinus *regalis*,) Supp. to Geo. Sur. Iowa, p. 8, and Geo. Sur.

Ill., vol. 2, p. 192, Burlington Gr. *umbrosus*, Hall, 1858, (Actinocrinus *umbrosus*,) Geo. Sur. Iowa, p. 590, Burlington Gr.

SYNBATHOCRINUS, Phillips, 1836, Geol. Yorkshire, pt. 2, p. 206. [Ety. *syn*, together; *bathos*, depth; *krinon*, lily.] Calyx small; arms large and of great length; basals 3; radials 2x5; azygous plates 1

2 or more; arms simple and composed of plates in single series. Type *S. conicus*.

brevis, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 68, and Geo. Sur. Ill., vol. 5, p. 439, Burlington Gr.

dentatus, Owen & Shumard, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 597, Burlington Gr. *granulatus*, Troost. Not defined.

granuliferus Wetherby, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 250, Waverly Gr. *matutinus*, Hall, 1858, Geo. Sur. Iowa, p. 483, Ham. Gr.

FIG. 488.—*Synbathocrinus granuliferus*. Anterior and posterior views.

SVR.—TA
oweni,
Nat.
papilla
p. 18
robust
Acad
Ill.,
swallo
672,
tennes
tennes
Wes
unchar
worth
500,
SYRINOC
Rem
pipe
on t
all
genu
parado
Deca
TENIART
Deca
aster,
rays
lacra
amb
the r
Org
TALAROCH
1881,
259.
Calyx
press
nus h
open
of a
radia
corni
cornig
Sci., v
elegans,
crinus
and S
ovatus,
St. M
Ill., v

oweni, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 111, Waverly Gr.
 papillatus, Hall, 1861, Desc. New Crin., p. 18, Burlington Gr.
 robustus, Shumard, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 397, and Geo. Sur. Ill., vol. 6, p. 514, Keokuk Gr.
 swallowi, Hall, 1858, Geo. Sur. Iowa, p. 672, St. Louis Gr.
 tennesseus, Troost. Not defined.
 tennesseensis, Roemer, 1860, Sil. Fauna West Tenn., p. 55, Niagara Gr.
 wachamuthi, see *Catillocrinus wachamuthi*.
 wortheni, Hall, 1858, Geo. Sur. Iowa, p. 560, Burlington Gr.

SYNGOCRINUS, Billings, 1859, Can. Org. Rem., Decade 4, p. 65. [Ety. *syninx*, pipe; *krinon*, lily.] Founded, possibly, on the fragment of a ventral sac; at all events, not a well-characterized genus. Type *S. paradoxicus*.
paradoxicus, Billings, 1859, Can. Org. Rem., Decade 4, p. 65, Trenton Gr.

TENIASTER, Billings, 1858, Can. Org. Rem., Decade 3, p. 80. [Ety. *tania*, ribbon; *aster*, star.] No disk or marginal plates; rays long, flexible, spinous; two rows of ambulacral plates elongated; two rows of ambulacral pores; ossicles contracted in the middle. Type *T. spinosus*.

FIG. 439.—*Teniaster spinosus*.

Org. Rem., Decade 3, p. 80, Trenton Gr.

TALAROCRINUS, Wachamuth & Springer, 1881, Proc. Acad. Nat. Sci. Phil., p. 259. [Ety. *talaros*, basket; *krinon*, lily.] Calyx sub-conical; suture lines impressed; distinguished from *Dichocrinus* by its higher vault and having the opening through it and not at the end of a tube, and in having the secondary radials form part of the calyx. Type *T. cornigerus*.

cornigerus, Shumard, 1857, (*Dichocrinus cornigerus*.) Trans. St. Louis Acad. Sci., vol. 1, p. 72, Kaskaskia Gr.

elegans, Lyon & Casseday, 1860, (*Dichocrinus elegans*.) Proc. Am. Acad. Arts and Sci., vol. 5, p. 22, St. Louis Gr.

ovatus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 36, and Geo. Sur. Ill., vol. 7, p. 314, Kaskaskia Gr.

sexlobatus, Shumard, 1857, (*Dichocrinus sexlobatus*.) Trans. St. Louis Acad. Sci., vol. 1, p. 73, Kaskaskia Gr.

symmetricus, Lyon & Casseday, 1860, (*Dichocrinus symmetricus*.) Proc. Am. Acad. Arts and Sci., vol. 5, p. 22, Kaskaskia Gr.

TAXOCRINUS, Phillips, 1843, Morris Cat. Brit. Foss., p. 90. [Ety. *taxus*, yew-tree; *krinon*, lily.] Calyx short, cup-shaped; basals 3, small, unequal; subradials 5, one larger than the others; primary radials 3 or 4 by 5; secondary radials 3 to six by 10; tertiary radials supporting arms; interradials 0 to 9; azygous interradials 2 to 5; arms dividing once or twice. Type *T. egertonii*.

communis, Hall, 1863, (*Forbesocrinus communis*.) 17th Rep. N. Y. St. Mus. Nat. Hist., p. 55, and Ohio Pal., vol. 2, p. 169, Waverly Gr.

curtus, Williams, 1882, Proc. Acad. Nat. Sci. Phil., p. 30, Chemung Gr.

elegans, Billings, 1857, (*Lecanocrinus elegans*.) Geo. Sur. Can., p. 278, and Can. Org. Rem., Decade 4, p. 47, Trenton Gr.

fletcheri, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 31, and Geo. Sur. Ill., vol. 7, p. 308, Keokuk Gr.

giddingei, Hall, 1858, (*Forbesocrinus giddingei*.) Geo. Sur. Iowa, p. 633, Keokuk Gr.

FIG. 440.—*Taxocrinus* (?) *elegans*.

FIG. 441.—*Taxocrinus gracilis*. Diagram.

gracilis, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 142, and Geo. Sur. Ill., vol. 3, p. 421, Ham. Gr.
intermedius, Wachamuth & Springer, (in press.) Geo. Sur.

Ill., vol. 8, p. 199 Kinderhook Gr.
interscapularis, Hall, 1858, Geo. Sur. Iowa, p. 482, Ham. Gr.

ithacensis, Williams, 1882, Proc. Acad. Nat. Sci. Phil., p. 28, Chemung Gr.

juvenis, Hall, 1861, (*Forbesocrinus juvenis*.) Bost. Jour. Nat. Hist., vol. 7, p. 319, Burlington Gr.

kelloggi, Hall, 1863, (*Forbesocrinus kelloggi*.) 17th Rep. N. Y. St. Mus. Nat. Hist., p. 56, and Ohio Pal., vol. 2, p. 171, Waverly Gr.

levis, Billings, 1857, Geo. Sur. Can., p. 278, and Can. Org. Rem., Decade 4, p. 47, Trenton Gr.

lobatus, Hall, 1862, (Forbesocrinus lobatus,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 124, Ham Gr.

lobatus var. tardus, Hall, 1863, (Forbesocrinus lobatus var. tardus,) 17th Rep. N. Y. St. Mus. Nat. Hist., p. 56 and Ohio Pal., vol. 2, p. 171, Waverly Gr.

meeki, Hall, 1858, (Forbesocrinus meeki,) Geo. Sur. Iowa, p. 631, Keokuk Gr.

multibrachiatus, Lyon & Casseday, 1858, (Forbesocrinus multibrachiatus,) Am. Jour. Sci. and Arts, vol. 28, p. 235, Keokuk Gr.

multibrachiatus var. colletti, White, 1881, 2d Ann. Rep. Bureau of Statistics of Indiana, p. 506, Keokuk Gr.

nuntius, Hall, 1862, (Forbesocrinus nuntius,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 124, Ham Gr.

ramulosus, Hall, 1860, (Forbesocrinus ramulosus,) Supp. Geo. Sur. Iowa, p. 67, Burlington Gr.

robustus, Wachsmuth, (in press,) Geo. Sur. Ill., vol. 8, Kinderhook Gr.

semiovatus, Meek & Worthen, 1860, (Forbesocrinus semiovatus,) Proc. Acad. Nat. Sci. Phil., p. 389, and Geo. Sur. Ill., vol. 2, p. 272, St. Louis Gr.

shumardanus, Hall, 1858, (Forbesocrinus shumardanus,) Geo. Sur. Iowa, p. 671, St. Louis Gr.

thiemii, Hall, 1861, (Forbesocrinus thiemii,) Desc. New Crin., p. 8, and Geo. Sur. Ill., vol. 5, p. 389, Burlington Gr.

whitfieldi, Hall, 1858, (Forbesocrinus whitfieldi,) Geo. Sur. Iowa, p. 632, Kaskaskia Gr.

TECHNOCRINUS, Hall, 1859, Pal. N. Y., vol. 3, p. 139. [Ety. *technē*, art; *krinon*, lily.] Basals 4, one larger than the others; primary radials 3x5; secondary radials 1x10; tertiary radials 2x20; interradials 3x5; arms simple, bearing pinnules; column round. Type *T. andrewsi*.

andrewsi, Hall, 1859, Pal. N. Y., vol. 3, p. 141, Oriskany sandstone.

sculptus, Hall, 1859, Pal. N. Y., vol. 3, p. 143, Oriskany sandstone.

spinulosus, Hall, 1859, Pal. N. Y., vol. 3, p. 140, Oriskany sandstone.

striatus, Hall, 1859, Pal. N. Y., vol. 3, p. 142, Oriskany sandstone.

TALIOCRINUS, Wachsmuth & Springer, 1881, Proc. Acad. Nat. Sci. Phil., p. 320. [Ety. *teleios*, perfect; *krinon*, lily.] Distinguished from *Strotocrinus*, with which it has generally been classed, by having a long ventral tube, instead of a simple opening through the vault. Type *T. umbrosus*.

agilops, Hall, 1860, (Actinocrinus agilops,) Supp. to Geo. Sur. Iowa, p. 5, Up. Burlington Gr.

althaea, Hall, 1861, (Actinocrinus althaea,) Desc. New Crin., p. 13, Up. Burlington Gr.

clivus, Hall, 1861, (Actinocrinus clivus,) Bost. Jour. Nat. Hist., vol. 7, p. 274, Up. Burlington Gr.

erodus, Hall, 1861, (Actinocrinus erodus,) Desc. New Crin., p. 12, Up. Burlington Gr.

insculptus, Hall, 1861, (Actinocrinus insculptus,) Desc. New Crin., p. 12, Up. Burlington Gr.

liratus, Hall, 1860, (Actinocrinus liratus,) Supp. to Geo. Sur. Iowa, p. 1, and Geo. Sur. Ill., vol. 5, p. 355, Burlington Gr.

rudis, Hall, 1860, (Actinocrinus rudis,) Supp. to Geo. Sur. Iowa, p. 33, Burlington Gr.

tenuiradiatus, Hall, 1861, (Actinocrinus tenuiradiatus,) Desc. New Crin., p. 12, Burlington Gr.

umbrosus, Hall, 1858, (Actinocrinus umbrosus,) Geo. Sur. Iowa, p. 590, Up. Burlington Gr.

THYSANOCRINUS, Hall, 1852, Pal. N. Y., vol. 2, p. 188. [Ety. *thysanos*, fringed; *krinon*, lily.] Calyx small, subglobose; basals 5; subradials 5; primary radials 3x5; secondary radials 2 or more x10; regular interradials 3; azygous area wide, lower plates large, smaller above; arms composed of a double series of plates, with pinnules; column round. Type *T. liliiformis*.

aculeatus, Hall, 1852, Pal. N. Y., vol. 2, p. 190, Niagara Gr.

canaliculatus, Hall, 1852, Pal. N. Y., vol. 2, p. 189, Niagara Gr.

immaturus, Hall, 1852, Pal. N. Y., vol. 2, p. 191, Niagara Gr.

liliiformis, Hall, 1852, Pal. N. Y., vol. 2, p. 188, Niagara Gr.

microbasalis, see Archæocrinus microbasalis.

pyriformis, see Archæocrinus pyriformis.

TREMATASTER, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 330. [Ety. *trema*, opening; *aster*, star.] Central part discoid; rays long, flexuous, a double series of ambulacral plates, with tapering ends directed toward the apices of the rays, upon each side of which there is a series of curved adambulacral plates, which form the margin of the rays; pores large between the con-

FIG. 442.—*Taxocrinus robustus*.

TRE.—XEN.
tracti
and th
adam
each
ficilla
difficil
Sur.
kia G
Trematocri
nus.
fascellu
lus.
papillatu
reticulatu
robustus,
spinigru
tubercula
typus, see
Triacrinus
Nat. &
preocc
syn. fo
globose,
pyriform
TRICLOCR
Proc.
Geo. S
three;
Calyx
base x
along
contra
deltoid
deeply
spires
spiracle
large;
mani.
meekanu
Catal.
saw G
obliquatu
tites of
Jahrg.
varsouvi
Ill., vo
woodma
tremite
nus w
Phil., p
p. 506,
TROOSTOCRI
Leuis
proper
tinguis
slender
bulacr
angula
ture; s
imal s
open i
Type T
bipyram
bipyram
Keokuk
grosvener
grosvener
vol. 1,

vol. 3, p.
ger, 1881,
p. 320.
ly.] Dis-
lased, by
instead of
the vault.

inus segi-
wa, p. 5,

s althea,)
Burling-

inus clivo-
vol. 7, p.

s erodus,)
Burling-

crinus in-
p. 12, Up.

s liratus,)
p. 1, and
Burling-

s rudis,)
33, Bur-

crinus
in., p. 12,

inus um-
590, Up.

T. Y., vol.
anged; kri-
globose;

ry radials
or more

azygous
e, smaller
double se-
column

, vol. 2, p.

N. Y., vol.

Y., vol. 2,
vol. 2, p.

s micro-

yriformia.

1883, Geo.

ty. trema,
l part dis-

a double
with taper-

apices of
which there
ambulacral
gin of the
the con-

tracting sides of the ambulacral plates, and the concave sides of the curving adambulacral; four plates border on each pore; orals 10. Type *T. difficilis*.

difficilis, Worthen & Miller, 1883, Geo. Sur. Ill., vol. 7, p. 330, Kaskaskia Gr.

Trematocrinus, syn for *Goniasteroidocrinus*.

fascellus, see *Goniasteroidocrinus fascellus*.

papillatus, see *G. papillatus*.

reticulatus, see *G. reticulatus*.

robustus, see *G. robustus*.

spinigerus, see *G. spinigerus*.

tuberculatus, see *G. tuberculatus*.

typus, see *G. typus*.

Triacrinus, Ringueberg, 1887, Proc. Acad. Nat. Sci. Phil., p. 144. The name was preoccupied; beside it is probably a syn. for *Pisocrinus*.

globosus, see *Pisocrinus globosus*.

pyriformis, see *Pisocrinus pyriformis*.

TRICELOCRINUS, Meek & Worthen, 1868, Proc. Acad. Nat. Sci. Phil., p. 356, and Geo. Sur. Ill., vol. 5, p. 507. [Ety. *treis*, three; *koilos*, hollow; *krinon*, lily.] Calyx subpyramidal, or subfusiform; base short, trihedral, and excavated along the interbasal sutures; summit contracted; radials long and narrow; deltoids small; ambulacra narrow, deeply situated in the sinuces; hydrospires small, three (?) on a side; spiracles and mouth small; anus large; column circular. Type *T. woodmani*.

meekanus, Etheridge & Carpenter, 1886, Catal. of Blastoides, p. 208, Warsaw Gr.

obliquatus, Roemer, 1851, (Pentatremites *obliquatus*.) Archiv f. Naturgesch., Jahrg. xvii, p. 367, St. Louis Gr.

varsouviensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 521, St. Louis Gr.

woodmani, Meek & Worthen, 1868, (Pentremites, *Troostocrinus*) *Tricelocrinus woodmani*.) Proc. Acad. Nat. Sci. Phil., p. 356, and Geo. Sur. Ill., vol. 5, p. 506, Warsaw Gr.

TROOSTOCRINUS, Shumard, 1865, Trans. St. Louis Acad. Sci., vol. 2, p. 384. [Ety. proper name; *krinon*, lily.] Distinguished from *Pentremites* by the slender, subfusiform shape, linear ambulacra, lancet plates concealed, triangular base, and simple summit structure; spiracles at the sides of the proximal side plates; hydrospiracular canals open into linear spiracular apertures. Type *T. reinwardti*.

bipyramidalis, Hall, 1858, (Pentremites *bipyramidalis*.) Geo. Sur. Iowa, p. 607, Keokuk Gr.

grosvenori, Shumard, 1858, (Pentremites *grosvenori*.) Trans. St. Louis Acad. Sci., vol. 1, p. 240, Warsaw Gr.

lineatus, Shumard, 1858, (Pentremites *lineatus*.) Trans. St. Louis Acad. Sci., vol. 1, p. 241, Burlington Gr. This is made the type of the genus *Metablaustus* by Etheridge & Carpenter, to which they also refer *T. wortheni* and *Tricelocrinus varsouviensis*.

reinwardti, Troost, 1835, (Pentremites *reinwardti*.) Trans. Geo. Soc. Pa., vol. 1, p. 224, Niagara Gr.

subcylindricus, Hall & Whitfield, 1875, (Pentremites *subcylindricus*.) Ohio Pal., vol. 2, p. 129, Niagara Gr.

subtruncatus, Hall, 1858, (Pentremites *subtruncatus*.) Geo. Sur. Iowa, p. 485, Ham. Gr.

wortheni, Hall, 1858, (Pentremites *wortheni*.) Geol. Sur. Iowa, p. 606, Keokuk Gr.

VASOCRINUS, Lyon, 1857, Geo. Sur. Ky., vol. 3, p. 485. [Ety. *vas*, vessel; *crinon*, lily.] Calyx low, vase-shaped; basals 5; subradials, 5; primary radials, 1 x 5; secondary radials 2 x 5; arms, 10 or more; azygous interradians 2 or more, first one large; ventral sac. Type *V. valens*. *lyoni*, Hall, 1861, (Cyathocrinus *lyoni*.) Desc. New Crin., p. 3, and Bost. Jour. Nat. Hist., vol. 7, p. 298, Keokuk Gr. *macroleurus*, Hall, 1861. (Cyathocrinus *macroleurus*.) Desc. New Crin., p. 5, and Bost. Jour. Nat. Hist., vol. 7, p. 295, Burlington Gr. *sculptus*, Lyon, 1857, Geo. Sur. Ky., vol. 3, p. 486, Ham. Gr. *valens*, Lyon, 1857, Geo. Sur. Ky., vol. 3, p. 485, Ham. Gr.

XENOCRINUS, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 71 and 176. [Ety. *xenos*, strange; *krinon*, lily.] Ba-

FIG. 443.
TROOSTOCRINUS
wortheni.

FIG. 444.—*Xenocrinus penicillus*. Azygous and opposite side views and end of column.

sals 4; primary radials 3 x 5; secondary radials 4 to 6 x 10; interradian areas excavated and filled with numerous

plates; azygous area having a central vertical series of plates which continue up the ventral sac; column square. Type X. *penicillus*.

baeri, Meek, 1872, (*Glyptocrinus baeri*.) Am. Jour. Sci. and Arts, 3d ser., vol. 3, p. 260, and Ohio Pal., vol. 1, p. 37, Hud. Riv. Gr.

penicillus, S. A. Miller, 1881, Jour. Clin. Soc. Nat. Hist., vol. 4, p. 72, Hud. Riv. Gr.

ZEACRINUS, Troost, Catal. Foss. 1850, and described by Hall, 1858, Geo. Sur. Iowa, p. 541. [Ety. *zea*, Indian corn; *krinon*, lily.] Calyx low, basin-shaped; basals 5, hidden by the column; subradials 5; radials 2 x 5, with from 1 to 6 additional in the azygous ray; azygous interradials 4 to 7; arms 10 to 40, with pinnales; ventral sac subpyramidal, covered with small plates; column round. Type Z. *magnoliformis*.

acanthophorus, see *Hydreionocrinus acanthophorus*.

arboreus, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 534, St. Louis Gr.

armiger, see *Hydreionocrinus armiger*.

asper, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 150, and Geo. Sur. Ill., vol. 5, p. 430, Burlington Gr.

bifurcatus, McCheaney, 1860, New Pal. Foss., p. 10, and Trans. Chi. Acad. Sci., vol. 1, p. 71, Kaskaskia Gr.

cariniferus, see *Cellocrinus cariniferus*. *compactilis*, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 536, Kaskaskia Gr.

coxanus, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 27, and Geo. Sur. Ill., vol. 7, p. 302, Keokuk Gr.

crassus, see *Eupachycrinus crassus*.

crateriformis, Troost. Not defined.

depressus, see *Hydreionocrinus depressus*.

discus, see *Hydreionocrinus discus*.

elegans, Hall,

1858, Geo. Sur.

Iowa, p. 547,

Burlington Gr.

florealis, Yandell

& Shumard,

1847, (*Cyathocrinus flore-*

alis.) Cont. to

Geo. Ky., p.

24, Kaskaskia

Gr.

formosus, see *Eup-*

pachycrinus

formosus.

intermedius,

Hall, 1858,

Geo. Sur. Iowa,

p. 681, Kas-

kaskia Gr.

keokuk, Wor-

then, 1882, Bull. No. 1, Ill. St. Mus. Nat.

Hist., p. 28, and Geo. Sur. Ill., vol. 7, p.

30, Keokuk Gr.

lyra, see *Cellocrinus lyra*.

magnoliformis, Owen & Norwood, 1846, (*Cyathocrinus magnoliformis*.) Research Pot. Carb. Rocks Ky., and Geo. Sur. Iowa, p. 684, Kaskaskia Gr.

maniformis, Yandell & Shumard, 1847, (*Poteroocrinus maniformis*.) Cont. to Geo. Ky., p. 24, Kaskaskia Gr.

merope, Hall, 1863, 17th Rep. N. Y. St. Mus. Nat. Hist., p. 60, and Ohio Pal., vol. 2, p. 178, Waverly Gr.

mooril, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 227, Coal Meas.

mucrospinus, see *Hydreionocrinus mucro-*

spinus.

nodosus, Wachsmuth & Springer, 1886,

Revis. Palaeocrinoloides, pt. 3, p. 243,

Keokuk Gr.

ovalis, Lyon & Casseday, 1858, Am. Jour.

Sci., 2d ser., vol. 29, p. 71, Kaskaskia Gr.

paternus, Hall, 1863, 17th Rep. N. Y. St.

Mus. Nat. Hist., p. 59, Waverly Gr.

perangulatus, White, 1862, Proc. Bost. Soc.

Nat. Hist., vol. 9, p. 11, Burlington Gr.

pikensis, Worthen, 1882, Bull., No. 1, Ill.

St. Mus. Nat. Hist., p. 29, and Geo. Sur.

Ill., vol. 7, p. 304, Burlington Gr.

planobrachiatus, Meek & Worthen, 1860,

Proc. Acad. Nat. Sci. Phil., p. 391, and

Geo. Sur. Ill., vol. 2, p. 240, Keokuk Gr.

ramosus, Hall, 1858, Geo. Sur. Iowa, p.

548, Burlington Gr.

sacculus, White, 1862, Proc. Bost. Soc. Nat.

Hist., vol. 9, p. 12, Burlington Gr.

FIG. 446.—*Zeacrinus spinuliferus*.

sacculus var. *concinus*, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 12, Burlington Gr.

scobina, Meek &

Worthen, 1869,

Proc. Acad. Nat.

Sci. Phil., p. 149,

and Geo. Sur.

Ill., vol. 5, p.

426, Burlington

Gr.

scoparius, Hall,

1861, Desc. New

Crin., p. 6, and

Jour. Bost. Soc.

Nat. Hist., vol. 7,

p. 305, Burling-

ton Gr.

serratus, Meek &

Worthen, 1861,

Proc. Acad. Nat. Sci. Phil., p. 151, and

Geo. Sur. Ill., vol. 5, p. 428, Burling-

ton Gr.

FIG. 447.—*Zeacrinus troostianus*. Diagram.

ood, 1846,
als.) Re-
and Geo.
Gr.
ard, 1847,
Cont. to
r.
N. Y. St.
Ohio Pal.,

J. Y. Acad.
us mucro-

ger, 1886,
3, p. 243,

Am. Jour.
skaskia Gr.
N. Y. St.
ly Gr.

Bost. Soc.
ington Gr.
No. 1, Ill.
Geo. Sur.
Gr.

then, 1860,
p. 391, and
Keokuk Gr.
r. Iowa, p.

t. Soc. Nat.
a Gr.

erus.

1862, Proc.
p. 12, Bur-

Zenerinus
Diagram.

p. 151, and
3, Burling-

spinosus, Owen & Shumard, 1852, (Pote-
riocrinus spinosus,) Jour. Acad. Nat.
Sci. Phil., vol. 2, p. 91, Kaskaskia Gr.
spinuliferus, Worthen, (in press.) (Pote-
riocrinus spinuliferus,) Geo. Sur. Ill.,
vol. 8, p. 90, Kaskaskia Gr.
stimpsoni, Lyon, 1869, Trans. Am. Phil.
Soc., vol. 13, p. 465, Subcarb.

subtumidus, see Eupachyrcinus subtu-
midus.

troostanus, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 390, and
Geo. Sur. Ill., vol. 2, p. 186, Burling-
ton Gr.

wortheni, Hall, 1858, Geo. Sur. Iowa, p.
683, Kaskaskia Gr.

SUBKINGDOM MOLLUSCOIDA.

CLASS BRYOZOA.

THE Bryozoa are small animals that grow in clusters, forming branched or moss-
like compound structures. Each animal lives in a separate cell, called a zoecium,
into which it can retract itself, though some connection exists between the animals.
The Flustra or Sea-mats, abundant on the shores of the ocean, and the moss-like en-
crustations so common on marine shells, are examples. All known Palaeozoic Bry-
ozoa were marine, and lived in calcareous cells, forming a mass that is often difficult
to distinguish from the true corals.

This calcareous mass or skeleton is called the bryozoum or zoarium. It is found
encrusting other objects, or standing on a foot-stalk, with basal attachment, and, in
other cases, apparently free. There are rarely any such calcareous partitions in the
cell-tubes as abound in the true corals, and the method of reproduction was exclu-
sively gemmiparous, while the true corals were increased by both gemmiparous and
fissiparous reproduction.

The animal consists of a bent tube or alimentary canal, having an oesophagus,
stomach, and intestine. The two orifices of the canal are situated close together,
but the anal opening is beyond the ring of ciliated tentacles that surround the
mouth. Thus constituted, the alimentary canal is inclosed in a sac having two
openings corresponding to the two extremities of the canal. Generally the upper
side of this sac is flexible, and admits of being invaginated, so that when the ani-
mal retracts itself into its cell the inverted portion forms a sheath around the ten-
tacles. Ova may be developed in a receptacle attached to the zoecium, called the
oecium, or in an inflation of the surface of the zoarium, called a gonocyst. The
gonocyst is a modified zoecium. The term oecia is also applied to these struc-
tures. Many Bryozoa have appendicular organs called avicularia and vibracula.
The avicularia may be pedunculate, and sway to and fro, or fixed and firmly at-
tached to the zoecium. The vibracula are flexible, bristle-like structures, set in the
excavated summit of a knob-like elevation or blunt spine.

Some naturalists refer the Monticuliporidae and Stelliporidae to the Bryozoa,
and probably the latter should be so classed on as good grounds as the Fistu-
liporidae are referred to the Bryozoa. The Palaeozoic Bryozoa are referred to an
order called the Gymnolamata, which are supposed to have had a complete ring of

tentacles around the mouth. This order has been divided into five suborders, viz.: Chilostomata, Cryptostomata, Trepotomata, Cyclostomata, and Otenostomata. The families which we recognize are as follows:

FAMILY ACANTHOCLADIDÆ.—Acanthocladia, Diplopore, Glauconome, Ichthyorachis, Ptilopora, Ramipora, Septopora, Synocladia.

FAMILY AMPLEXOPORIDÆ.—Amplexopora, Atactopora, Discotrypa, Leptotrypa, Petalotrypa.

FAMILY ARTHROSTYLIDÆ.—Arthroclema, Arthrostylus, Helopora, Nematopora, Nematoporella, Sceptropora.

FAMILY ASCODICTYONIDÆ.—Ascodictyon, Rhopalonaria.

FAMILY BATOSTOMELLIDÆ.—Anisotrypa, Batostoma (?), Batostomella, Leiolema, Peronopora.

FAMILY BOTRYLLOPORIDÆ.—Botryllopora.

FAMILY BYTHOPORIDÆ.—Bythopora.

FAMILY CERAMOPORIDÆ.—Aspidopora, Ceramella, Ceramopora, Ceramoporella, Chiloporella, Crepipora, Eridopora, Glossotrypa, Idiotrypa, Lichenalia, Lichenotrypa, Odontotrypa, Petigopora, Phractopora, Pileotrypa, Sagenella, Selenopora, Spatiopora.

FAMILY CRISINELLIDÆ.—Crisinella.

FAMILY ENALLOPORIDÆ.—Diploclema, Enallopora, Protocrisina.

FAMILY FENESTELLIDÆ.—Archimedes, Clathropora, Coscinella, Coscinium, Coscinotrypa, Evactinopora, Fenestella, Fenestralia, Fenestrapora, Helicopora, Hemitrypa, Isotrypa, Loculipora, Lyropora, Phyllopora, Polypora, Ptilopora, Ptiloporella, Ptiloporina, Reptaria, Reteporina, Senuicoscium, Semiopora, Teetulipora, Unitrypa.

FAMILY FISTULIPORIDÆ.—Actinotrypa, Buscopora, Callopora, Calloporella, Callotrypa, Chilotrypa, Cœlocaulis, Eridopora, Favicella, Fistulipora, Lichenotrypa, Pinacotrypa, Selenopora, Strotopora.

FAMILY HELIOTRYPIDÆ.—Heliotrypa.

FAMILY LABECHIDÆ.—Labechia.

FAMILY PALESCHARIDÆ.—Paleschara.

FAMILY PHACELOPORIDÆ.—Phacelopora.

FAMILY PTILODICTYONIDÆ.—Coscinella, Cyclopora, Cycloporella, Escharopora, Graptodictya, Heterodictya, Phænopora, Proutella, Ptilodictya, Ptilotrypa, Streblotrypa, Worthenopora.

FAMILY RHABDOMESONTIDÆ.—Acanthoclema, Anisotrypa, Bactropora, Cœlocorus, Nemataxis, Rhombopora, Tropicopora.

FAMILY RHINOPORIDÆ.—Rhinopora.

FAMILY SPHRAGIOPORIDÆ.—Sphragiopora.

FAMILY STICTOPORIDÆ.—Acrogenia, Arthropora, Cystodictya, Dichotrypa, Dicranopora, Eurodictya, Euspilopora, Goniortrypa, Heliotrypa, Intrapora, Pachydictya, Phractopora, Phyllodictya, Prismopora, Rhinidictya, Scalaripora, Stictopora, Stictoporella, Stictoporina, Stictotrypa, Sulcopora, Tænidictya, Tæniopora, Thamnotrypa.

FAMILY SUBRETIPORIDÆ.—Chainodictyon, Drymotrypa, Subretopora.

FAMILY THAMNISCIDÆ.—Crisinella, Diplopore, Thamniscus.

FAM

FAM

FAM

ACANTHO

Nat. H

akanth

symm

more c

furcati

caslon

bilater

one sid

and ar

tudina

ridges.

american

Acad.

fruticosa

Ill., vo

ACANTHO

6, p. 7

twig.]

a cent

longitu

a bran

ridges

longitu

form n

alternatu

ternata

(Gr., an

Held.

bispinul

spinul

882, a

Ham.

confluen

fluens,

Cuyah

divergen

73, Up

ovatum,

73, Up

scutulatu

scutul

180, a

Ham.

sulcatum

192, H

triseriale

rialia.)

vol. 6,

ACROGENIA

[Ety.

Fron

ing fr

each

divisio

FAMILY THEONOIDÆ.—Scenellopora.

FAMILY TREMATOPORIDÆ.—Acanthoclema, Amplexopora, Atactopora, Atactoporella, Bactropora, Chilotrypa, Diamesopora, Homotrypa, Homotrypella, Nemataxis, Nicholsonella, Orthopora, Trematella, Trematopora, Trepidopora.

FAMILY TUBULIPORIDÆ.—Berenicea, Clonopora, Cystopora, Hederella, Hermodia, Stomatopora.

ACANTHOCLADIA, King, 1849, Ann. and Mag. Nat. Hist., 2d ser., vol. 3, p. 389. [Ety. *akantha*, spine; *klados*, branch.] Stem symmetrically and bilaterally branched, more or less on one plane; rarely bifurcating; branches short, simple, occasionally elongated and becoming bilaterally branched; celluliferous on one side only; cell apertures circular and arranged in three or more longitudinal series, separated by dividing ridges. Type A. anceps.

americana, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 180, Permian Gr. fruticosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 65, Up. Coal Meas.

ACANTHOCLEMA, Hall, 1887, Pal. N. Y., vol. 6, p. 72. [Ety. *akantha*, spine; *klema*, twig.] Ramose, solid, cells arising from a central axis; cell apertures oval, in longitudinal parallel rows, about ten on a branch, separated by longitudinal ridges; between the apertures, in the longitudinal direction, there are spiniform nodes. Type A. alternatum.

alternatum, Hall, 1881, (Trematopora alternata,) Bryozoa of the Up. Held. Gr., and Pal. N. Y., vol. 6, p. 72, Up. Held. Gr.

bispinulatum, Hall, 1881, (Callopora bispinulata,) Trans. Alb. Inst., vol. 10, p. 882, and Pal. N. Y., vol. 6, p. 182, Ham. Gr.

confluens, Ulrich, 1888, (Rhombopora confluens,) Bull. Denison Univ., p. 91, Cuyahoga Shales.

divergens, Hall, 1887, Pal. N. Y., vol. 6, p. 73, Up. Held. Gr.

ovatum, Hall, 1887, Pal. N. Y., vol. 6, p. 73, Up. Held. Gr.

scutellatum, Hall, 1881, (Trematopora scutellata,) Trans. Alb. Inst., vol. 10, p. 180, and Pal. N. Y., vol. 6, p. 190, Ham. Gr.

sulcatum, Hall, 1887, Pal. N. Y., vol. 6, p. 192, Ham. Gr.

triseriale, Hall, 1883, (Stictopora triseriale,) Rep. St. Geol. and Pal. N. Y., vol. 6, p. 74, Up. Held. Gr.

ACROGENIA, Hall, 1884, Rep. St. Geol. p. 51. [Ety. *akros*, sharp; *genea*, growth.] Frond ramose; two branches proceeding from the truncate termination of each preceding one; base of each division obconical, terete above and

strongly striated, gradually becoming flattened and celluliferous; margins noncelluliferous; apertures in rows separated by ridges, central range of apertures the smaller. Type A. prolifera.

prolifera, Hall, 1884, Rep. St. Geol., p. 52, and Pal. N. Y., vol. 6, p. 267, Ham. Gr.

ACTINOTRYPA, Ulrich, Geo.

Sur. Ill., vol. FIG. 44—Acrogenia prolifera. 8, p. 386, (in press.) [Ety. *aktin*, a ray; *trypa*, an opening.] Like Dichotrypa. Cell apertures showing the projecting ends of from eight to ten vertical septal-like ridges, that extend down on the inner side of the tubular vestibule nearly or quite to the primitive apertures. Type A. peculiaris.

peculiaris, Rominger, 1866, (Fistulipora peculiaris,) Proc. Acad. Nat. Sci. Phil., p. 10, Keokuk Gr.

Alecto, Lamouroux, 1821, Exposit. Method. It was preoccupied by Leach in the class Echinodermata, when Lamouroux used it, and hence Stomatopora is used in its place.

auloporoides, see Stomatopora auloporoides.

canadensis, see Hederella canadensis.

confusa, see Stomatopora confusa.

frondosa, see Stomatopora frondosa.

inflata, see Stomatopora inflata.

nexilis, see Stomatopora nexilis.

AMPLEXOPORA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 154. [Ety. *amplexus*, an encircling; *poros*, pore.] Ramose; cells of one kind only; walls thin in the axial part of the branches, but thicker in the peripheral region; acanthopores numerous. Type A. cingulata.

affinis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 36, Hud. Riv. Gr.

canadensis, Foord, 1883, Cont. to Micro-paleontology, p. 17, Trenton Gr.

cingulata, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 254, Hud. Riv. Gr.

FIG. 449.—*Amplexopora cingulata*. Tangential section x 50, showing wall structure, and vertical section x 50.

discoidea, Nicholson, 1875, (Chetetes discoideus,) Ohio Pal., vol. 2, p. 206, Hud. Riv. Gr.

FIG. 450.—*Amplexopora robusta*, funnel-shaped diaphragm, supposed to be a modified cystiphragm.

pustulosa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 36, Hud. Riv. Gr.
robusta, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 82, Hud. Riv. Gr.
septosa, Ulrich, 1879, (Atactopora septosa,) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 125, Hud. Riv. Gr.
superba, Foord, 1883, Cont. to Micropalaontology, p. 16, Trenton Gr.

winchelli, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 91, Trenton Gr.

ANISOTRYPA, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 275. [Ety. *anisos*, unequal; *trypa*, perforation.] Ramose, hollow, inner side lined with an epitheca; walls of tubes thin in the interior, and thickened exteriorly; no interstitial cells or spiniform tubuli. Type A. symmetrica.

fistulosa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 72, St. Louis Gr.

ramulosa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 72, St. Louis Gr.

solida, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 72, Kaskaskia Gr.

symmetrica, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 76, Kaskaskia Gr.

ARCHIMEDES, LeSueur, 1842, (Retepora archimedes,) Am. Jour. Sci., vol. 43, p. 19. [Ety. from its resemblance to the machine for raising water, consisting of a tube rolled in a spiral form around a cylinder, invented by Archimedes. Distinguished from Fenestella by its axis and mode of growth; the flabelliform expansion acquiring a solid central axis, around which it revolves in an ascending spiral form, spreading equally in all directions. Type A. wortheni.]

communis, Ulrich, (in press,) Geol. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

compactus, Ulrich, (in press,) Geol. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

distans, Ulrich, (in press,) Geol. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

grandis, Ulrich, (in press,) Geol. Sur. Ill., vol. 8, pl. 63, Keokuk Gr.

intermedius, Ulrich, (in press,) Geol. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

invaginatus, Ulrich, (in press,) Geol. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

FIG. 452.—*Archimedes wortheni*.

owenianus, Hall, 1857, Proc. Am. Ass'n Ad. Sci., vol. 10, p. 176, Keokuk Gr.

perminimus, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

proutianus, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

reversus, Hall, 1858, Geo. Rep. Iowa, p. 652, Warsaw Gr.

sublaxus, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

swallowianus, Hall, 1857, Proc. Am. Ass'n Ad. Sci., vol. 10, p. 176, Kaskaskia Gr.

terebriiformis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 63, Kaskaskia Gr.

wortheni, Hall, 1857, Proc. Am. Ass'n Ad. Sci., vol. 10, p. 176, and Geo. Sur. Iowa, p. 651, Warsaw Gr.

Archimediopora, D'Orb., 1850, Prod. de Pal., t. 1, p. 102, syn. for Archimedes.

Archimediopora archimedes was too tautological, and by common consent Archimedes has become the generic name.

archimedes, see Archimedes.

FIG. 451.—*Archimedes reversus*.

laxus, Hall, 1857, Proc. Am. Ass'n Ad. Sci., vol. 10, p. 176, Kaskaskia Gr.

meekianus, Hall, 1857, Proc. Am. Ass'n Ad. Sci., vol. 10, p. 176, Kaskaskia Gr.

negligens, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 63, Keokuk Gr.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

round, shafferi, section.

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

ARTHRODONTES, 1888, vol. 1, p. 101, arthrodon, subcylindrical, each side of the central axis, curved inwards, (curtus, tum,) p. 101,

Geol. Sur.
Gr.

chirmedes
as.

Geol. Sur.
Gr.

s, Hall,
1857, Proc.
m. Ass'n
d. Sci.,
vol. 10, p.
16, Kas-

askia Gr.
k and s,
all, 1857.

Proc. Am.
Ass'n Ad.
Sci., vol. 10,

176, Kas-

askia Gr.
ligens, Ul-

rich, (in
press), Geo.

Sur. Ill.,
vol. 8, pl.

3, Keokuk
Gr.

m. Ass'n
nk Gr.

Geo. Sur.
Gr.

Geo. Sur.
Gr.

Iowa, p.

Geo. Sur.
Gr.

Am. Ass'n
Kaskas-

Geo. Sur.
Gr.

m. Ass'n
Geo. Sur.

d. de Pal.,
chirmedes.

was too
n consent

the generic

ARTHROCLEMA, Billings, 1862, Pal. Foss., vol. 1, p. 54. [Ety. *arthron*, joint; *klessa*, twig.] Cylindrical jointed stem, with long, slender-jointed branches; pores oval. Type *A. pulchellum*.

angulare, Ulrich, (in press), Geo. Sur. Ill., vol. 8, pl. 29, Hud. Riv. Gr.

billingsi, Ulrich, (in press), Geo. Sur. Ill., vol. 8, Trenton Gr.

pulchellum, Billings, 1862, Pal. Foss., vol. 1, p. 54, Trenton Gr. (See p. 329.)

spiniforme, see *Helopora spiniformis*.

Arthronema, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 160. The name was preoccupied. See *Arthrostylus*.

FIG. 453.—*Arthropora shafferi*.

curtum, see *Arthrostylus curtus*.

tenuis, see *Arthrostylus tenuis*.

FIG. 454.—*Arthropora shafferi*. Magnified section.

rounded by interstitial pits. Type *A. shafferi*.

shafferi, Meek, 1872, (Stictopora *shafferi*.) Proc. Acad.

Nat. Sci., p. 317, and Ohio Pal., vol. 1, p. 69, Hud. Riv.

Gr.

simplex, Ulrich, 1886, 14th Rep. Geo. Sur.

Minn., p. 65, Trenton Gr.

ARTHROSTYLUS, Ulrich, 1888, Am. Geol.

vol. 1, p. 230. [Ety. *arthron*, joint;

stylos, pillar.] Ramose, composed of

subcylindrical segments, swollen at

each end, celluliferous on one side,

suolated on the other; cells between elevated lines. Type *A. tenuis*.

curtus, Ulrich, 1882, (*Arthronema curtum*.) Jour. Cin. Soc. Nat. Hist., vol. 5,

p. 161, Hud. Riv. Gr.

ARTHROPORA, Ulrich, 1882, Jour. Cin.

Soc. Nat. Hist., vol. 5, p. 152.

[Ety. *arthron*, a joint; *poros*, perforation.] Zoarium like Stictopora, but in

short, jointed, branching segments; cell apertures subcircu-

lar, and sur-

rounded by interstitial pits. Type *A. shafferi*.

shafferi, Meek, 1872, (Stictopora *shafferi*.) Proc. Acad.

Nat. Sci., p. 317, and Ohio Pal., vol. 1, p. 69, Hud. Riv.

Gr.

simplex, Ulrich, 1886, 14th Rep. Geo. Sur.

Minn., p. 65, Trenton Gr.

ARTHROSTYLUS, Ulrich, 1888, Am. Geol.

vol. 1, p. 230. [Ety. *arthron*, joint;

stylos, pillar.] Ramose, composed of

subcylindrical segments, swollen at

each end, celluliferous on one side,

suolated on the other; cells between elevated lines. Type *A. tenuis*.

curtus, Ulrich, 1882, (*Arthronema curtum*.) Jour. Cin. Soc. Nat. Hist., vol. 5,

p. 161, Hud. Riv. Gr.

tenuis, Ulrich, 1882, (*Arthronema tenue*.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 160, Trenton Gr.

Ascidictrvon, Nicholson, 1877, Ann. and Mag. Nat. Hist., 4th ser., vol. 19, p. 463. [Ety. *askos*, leather bottle; *dictyon*, net.] Organism composite, parasitic, composed of numerous calcareous cells; minutely perforated. Type *A. tusiforme*. It is probably a sponge.

tusiforme, Nicholson, 1877, Ann. and Mag. Nat. Hist., 4th ser., vol. 19, p. 463, Ham. Gr.

stellatum, Nicholson, 1877, Ann. and Mag. Nat. Hist., 4th ser., vol. 19, p. 464, Ham. Gr.

ASPIDOPORA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 155. [Ety. *aspis*, shield;

poros, perforation.] Thin, free expansions; concentrically wrinkled and striated epitheca on the lower side;

cells gradually increasing in size toward the center of the convex expansion;

interstitial cells numerous; diaphragms cross both kinds of tubes; spiniform tubuli present. Type *A. areolata*.

areolata, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 164, Utica Slate.

calculus, James, 1875, (Chetetes *calculus*.) Int. Catal. Cin. Foss., p. 1, and Nicholson Struct. and Affin. Montic., p. 165, Utica Slate.

parasitica, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 90, Trenton Gr.

ATACTOPORA, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 119. [Ety. *atactos*, without regularity; *poros*, pore.] Incrusting; surface with monticules or maculae; cell apertures petaloid, surrounded by rows of blunt spines; interstitial cells in clusters; tube walls infected; diaphragms present. Type *A. hirsuta*.

hirsuta, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 120, Hud. Riv. Gr.

maculata, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 121, Hud. Riv. Gr.

multigranosa, see *Atactoporella multigranosa*.

mundula, see *Atactoporella mundula*.

septosa, see *Amplexopora septosa*.

? *subramosa*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 124, Hud. Riv. Gr.

tenella, see *Atactoporella tenella*.

ATACTOPORELLA, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 247. [Ety. diminutive of *Atactopora*.] Incrusting; surface with monticules or maculae; cell apertures petaloid; interstitial cells numerous; spiniform tubuli and diaphragms. Type *A. typicalis*.

multigranosa, Ulrich, 1879, (*Atactopora multigranosa*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 122, Hud. Riv. Gr.

mundula, Ulrich, 1879, (*Atactopora mundula*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 123, Hud. Riv. Gr.

septosa, see *Amplexopora septosa*.

? *subramosa*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 124, Hud. Riv. Gr.

tenella, see *Atactoporella tenella*.

ATACTOPORELLA, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 247. [Ety. diminutive of *Atactopora*.] Incrusting; surface with monticules or maculae; cell apertures petaloid; interstitial cells numerous; spiniform tubuli and diaphragms. Type *A. typicalis*.

multigranosa, Ulrich, 1879, (*Atactopora multigranosa*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 122, Hud. Riv. Gr.

mundula, Ulrich, 1879, (*Atactopora mundula*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 123, Hud. Riv. Gr.

septosa, see *Amplexopora septosa*.

? *subramosa*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 124, Hud. Riv. Gr.

tenella, see *Atactoporella tenella*.

ATACTOPORELLA, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 247. [Ety. diminutive of *Atactopora*.] Incrusting; surface with monticules or maculae; cell apertures petaloid; interstitial cells numerous; spiniform tubuli and diaphragms. Type *A. typicalis*.

multigranosa, Ulrich, 1879, (*Atactopora multigranosa*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 122, Hud. Riv. Gr.

mundula, Ulrich, 1879, (*Atactopora mundula*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 123, Hud. Riv. Gr.

septosa, see *Amplexopora septosa*.

? *subramosa*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 124, Hud. Riv. Gr.

tenella, see *Atactoporella tenella*.

ATACTOPORELLA, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 247. [Ety. diminutive of *Atactopora*.] Incrusting; surface with monticules or maculae; cell apertures petaloid; interstitial cells numerous; spiniform tubuli and diaphragms. Type *A. typicalis*.

multigranosa, Ulrich, 1879, (*Atactopora multigranosa*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 122, Hud. Riv. Gr.

mundula, Ulrich, 1879, (*Atactopora mundula*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 123, Hud. Riv. Gr.

newportensis, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 250, Utica Slate.

FIG. 456.—*Atactopora newportensis*.

ortoni, Nicholson, 1874, (Chetetes *ortoni*.) Quar. Jour. Geo. Soc., vol. 30, p. 513, and Ohio Pal., vol. 2, p. 211, Hud. Riv. Gr.

schucherti, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 251, Hud. Riv. Gr.

tenella, Ulrich, 1879, (*Atactopora tenella*.) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 123, Hud. Riv. Gr.

typicalis, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 248, Utica Slate.

BACTROPORA, Hall, 1887, Pal. N. Y., vol. 6, p. 193. [Ety. *baktron*, staff; *poros*, pore.] Ramose, solid; base tapering, striated; cells tubular, curved oblique from the center; septa thin, apertures oval, distant near the base, closer above; interspaces granulose. Type *B. granistriata*. *curvata*, Hall, 1887, Pal. N. Y., vol. 6, p. 194, Ham. Gr.

granistriata, Hall, 1881, (Trematopora *granistriata*.) Trans. Alb. Inst., vol. 10, p. 182, and Pal. N. Y., vol. 6, p. 193, Ham. Gr.

simplex, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 70, Keokuk Gr.

BATOSTOMA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 154. [Ety. *batos*, prickly bush; *stoma*, mouth.] Ramose, base expanded; cell apertures ovate or circular, surrounded by a ring-wall; interstitial tubes numerous; spiniform tubuli abundant. Type *B. impicatum*.

fertile, Ulrich, 14th Rep. Geo. Sur. Minn., p. 92 Trenton Gr.

imperfectum, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 35, Hud. Riv. Gr.

impicatum, Nicholson, 1881, (Monticulipora *impicata*.) Struct. and Affin. of Montic., p. 147, Hud. Riv. Gr.

irrasum, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 94, Trenton Gr.

FIG. 457.—*Batostoma jamesi*. Natural size and magnified.

jamesi, Nicholson, 1874, (Chetetes *jamesi*.) Quar. Jour. Geo. Soc., vol. 30, p. 506, and Ohio Pal., vol. 2, p. 200 Hud. Riv. Gr.

manitobense, Ulrich, (in press.) Micropalaontology, p. 7, Hud. Riv. Gr. (?)

ottawense, Foord, 1883, Cont. to Micropalaontology, p. 18, Trenton Gr.

rugosum, Whitfield, 1882, (Fistulipora *rugosa*.) Geo. Wis., vol. 4, p. 255, Hud. Riv. Gr.

variabile, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 35, Hud. Riv. Gr.

BATOSTOMELLA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 154. [Ety. diminutive of *Batostoma*.] Ramose, smooth, cell apertures small; interstitial cells and spiniform tubuli; walls of tubes in the peripheral region thick. Type *B. gracilis*.

abrupta, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 75, Kaskaskia Gr.

FIG. 458.—*Batostomella gracilis*. Natural size and enlarged.

gracilis, Nicholson, 1874, (Chetetes *gracilis*.) Quar. Jour. Geo. Soc., vol. 30, p. 504, and Ohio Pal., vol. 2, p. 198, Hud. Riv. Gr.

interstincta, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 75, St. Louis Gr.

nitidula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 75, Kaskaskia Gr.

obliqua, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 46, Ham. Gr.

spinulosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 75, Kaskaskia Gr.

simulatrix, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 35, Hud. Riv. Gr.

BERENICEA, Lamoureux, 1821, Exp. Meth. des genres. d. pol., 80. [Ety. mythological name.] Incrusting, composed of a very thin, calcareous, foliaceous base, bearing numerous ovate, distinctly separated cells, not piled; aperture round near the broad anterior end; cells disposed in an obscurely radiated arrangement. Type *B. diluviana*.

insueta, Dawson, 1883, Rep. on Redpath, Mus. No. 2, p. 12, Subcarboniferous.

minnesotensis, Ulrich, 1886, 14th Rep. Geo. Sur. of Minn., p. 58, Trenton Gr.

primitiva, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 157, Hud. Riv. Gr.

vesiculosa, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 158, Utica Slate or lower part Hud. Riv. Gr.

BOTRYLLOPORA, Nicholson, 1874, Geo. Mag. Lond. n. s., vol. 1, p. 160. [Ety. *botryllos*, cluster; *poros*, pore.] Incrusting, forming systems of small circular disks, the upper surfaces of which are marked with radiating, cell-bearing ridges; non-

pori
disk
occu
B. so

FIG. 459.—*natural ray to sl*

socialis
n. s.,
BUCOPOR
p. 22
perfo
incru
a co
zoec
apert
or pe
tooth
natio
stitial
with
dentata,

FIG. 460.—*B*
nata. Ta
tion, sh
ture or lu

31, ar
Held.
BYTHOPOR
Pal. N
sea; p
small,

Fr
cell ap
rated
frutico

poriferous space in the center of each disk, round which the radiating ridges occupy a slightly elevated zone. Type *B. socialis*.

FIG. 459.—*Botryllopora socialis*. a, (Group on coral natural size; b, enlarged specimen; c, enlarged ray to show pores.

socialis, Nicholson, 1874, Geo. Mag. Lond. n. s., vol. 1, p. 160, Ham. Gr.

BUSCOPORA, Ulrich, 1886, Cont. to Am. Pal., p. 22. [Ety. *Busk*, proper name; *poros*, perforation.] Zoarium thin, lamellate; incrusting or free; under surface, with a concentrically wrinkled epitheca; zoecia tubular, short, with subcircular apertures and a faintly elevated border or peristome; posterior margin, with a tooth-like process divided at its termination; accessory cells present; interstitial spaces vesiculose; zoecial tubes, with diaphragms. Type *B. lunata*.

dentata, Ulrich, syn. for *B. lunata*.

FIG. 460.—*Buscopora lunata*. Tangential section, showing aperture or lunarium.

lunata, Rominger, 1868, (*Fistulipora lunata*.) Proc. Acad. Nat. Sci. Phil., p. 7, and Pal. N. Y., vol. 6, p. 77, Up. Held. Gr.

lunata var. *tubulata*, Hall, 1887, (*Lichenalia lunata* var. *tubulata*.) Rep. St. Geol. for 1885, pl. 31, and Pal. N. Y., vol. 6, p. 78, Up. Held. Gr.

BYTHOPORA, Miller & Dyer, 1878, Cont. to Pal. No. 2, p. 6. [Ety. *bythos*, depths of sea; *poros*, pore.] Dendroid, branches small, sometimes anastomosing, smooth;

FIG. 461.—*Bythopora fruticosa*.

cell apertures longer than wide, separated by impressed lines. Type *B. fruticosa*.

arctipora, Nicholson, 1875, (*Ptilodictya arctipora*.) Ann. and Mag., ser. 4, vol. 15, p. 180, Utica Slate.

delicatula, (?) instead of *Monticulipora delicatula*.

fruticosa, Miller & Dyer, 1878, Cont. to Pal. No. 2, p. 6, Hud. Riv. Gr.

herricki, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 99, Trenton Gr.

nashvillensis, S. A. Miller, 1880, Jour.

Cin. Soc. Nat.

Hist., vol. 3, p.

143, Trenton Gr.

striata, Ulrich, (in

pressa,) *Micropora*

laeontology, p. 10,

Hud. Riv. Gr.

CALLIOPORA, Hall,

1852, Pal. N. Y.,

vol. 2, p. 144.

[Ety. *kallos*, beau-

tiful; *poros*, pore.]

Ramose, smooth, or tuberculated; cell

tubes cylindrical; interstitial cells num-

erous; diaphragms numerous, no spini-

form tubuli; intercellular space occupied

by septate tubuli. Type *C. elegantula*.

aculeolata, see *Cœlocaulis aculeolata*.

? *aspera*, Hall, 1852, Pal. N. Y., vol. 2, p.

147, Niagara Gr.

bipunctata, Hall, 1884, Rep. St. Geol., p.

15, syn. for *Streblotrypa hamiltonensis*.

bispinulata, see *Orthopora bispinulata*.

cellulosa, Hall, 1883, Rep. St. Geol., pl. 12,

fig. 7-9, Low. Held. Gr.

cervicornis, Hall, 1879, Desc. New Spec.

Foss., p. 3, and 11th Rep. Ind. Geo.

Sur., p. 238, Niagara Gr.

cincinnatiensis, Ulrich, syn. for *Leioclema*

occidens.

diversa, Hall, 1879, Desc. New Spec. Foss.,

p. 4, and 11th Rep.

Ind. Geo. Sur., p.

239, Niagara Gr.

elegantula, Hall, 1852,

Pal. N. Y., vol. 2, p.

144, Niagara Gr.

? *exsul*, Hall, 1876, (*Al-*

veolites exsul.) 28th

Rep. N. Y. St. Mus.

Nat. Hist., p. 115,

Niagara Gr.

fistulosa, Hall, 1883, Rep. St. Geol., pl. 12,

fig. 1-6, Low. Held. Gr.

florida, Hall, 1852, Pal. N. Y., vol. 2, p.

146, Niagara Gr.

geniculata, Hall, 1887, Pal. N. Y., vol. 6,

p. 75, Up. Held. Gr.

hemispherica, see *Fistulipora hemispherica*.

heteropora, see *Callotrypa heteropora*.

hyale, see *Cœlocaulis hyale*.

incrassata, see *Fistulipora incrassata*.

incontroverta, Ulrich, 1886, 14th Rep.

Geo. Sur. Minn., p. 96, Trenton Gr.

internodata, see *Callotrypa internodata*.

irregularis, see *Cœlocaulis irregularis*.

laminata, Hall, 1852, Pal. N. Y., vol. 2, p.

146, Niagara Gr.

FIG. 462.—*Bythopora nashvillensis*. Magnified.

FIG. 463.—*Calliopora exsul*.

macropora, see *Callotrypa macropora*.
macropora var. *signata*, see *Callotrypa macropora* var. *signata*.
magnopora, Foerste, 1887, Bull. Denison University, p. 173, Niagara Gr.
minutissima, see *Leiclema minutissimum*.
missouriensis, Rominger, syn. for *Leiclema punctatum*.
multiseriata, see *Callotrypa multiseriata*.
nodulosa, Nicholson, 1874, (Chetetes nodulosus,) Quar. Jour. Geo. Soc., vol. 30, p. 506, and Ohio Pal., vol. 2, p. 200, Hud. Riv. Gr.
nummiformis, Hall, 1852, Pal. N. Y., vol. 2, p. 148, Niagara Gr.
oculifera, see *Callotrypa oculifera*.
ohioensis, Foerste, 1887, Bull. Denison Univ., p. 174, Niagara Gr.
onealli, James, 1875, (Chetetes onealli,) Int. Catal. Cin. Foss., p. 2, Hud. Riv. Gr.
oppleta, Hall, 1887, Pal. N. Y., vol. 6, p. 21, Low. Held. Gr.

FIG. 464.—*Callotrypa sigillarioides*. Natural size and magnified.

1875, (Chetetes sigillarioides,) Ohio Pal., vol. 2, p. 203, Hud. Riv. Gr.
singularis, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 115, Niagara Gr.
subnodosa, Ulrich, (in press), Geo. Sur. Ill., vol. 8, pl. 33, Hud. Riv. Gr.
subplana, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 253, Hud. Riv. Gr.
undulata, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 95, Trenton Gr.
unispina, see *Callotrypa unispina*.
venusta, see *Cælocaulis venusta*.
CALLOPORILLA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 154. [Ety. diminutive of *Callotrypa*.] Thin expansions, epitheca below; tubes with thick walls containing interstitial cells or angular

parasitica, see *Fistulipora parasitica*.
perelegans, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 102, Low. Held. Gr.
ponderosa, see *Fistulipora ponderosa*.
punctata, see *Leiclema punctatum*.
punctillata, Winchell, 1866, Rep. Low. Penin. Mich., p. 88, Ham. Gr.
sigillarioides, Nicholson,

FIG. 465.—*Callotrypa subnodosa*. Tangential section x50, showing amalgamation of walls.

mesopores; diaphragms and spiniform tubuli. Type *C. harrisi*.
harrisi, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 91, Hud. Riv. Gr.
nodulosa, Ulrich, (in press), Geo. Sur. Ill., vol. 8, pl. 33, Hud. Riv. Gr.
CALLOTRYPA, Hall, 1887, Pal. N. Y., vol. 6, p. 24. [Ety. *kallos*, beautiful; *trypa*, foramen.] Distinguished from *Callotrypa* by having a solid intercellular space, or one occupied with minute tubuli without septa. Type *C. macropora*.
heteropora, Hall, 1874, (Callotrypa heteropora,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 102, Low. Held. Gr.
internodata, Hall, 1881, (Callotrypa internodata,) Trans. Alb. Inst., vol. 10, p. 182, and Pal. N. Y., vol. 6, p. 189, Ham. Gr.
macropora, Hall, 1874, (Callotrypa macropora,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 101, Low. Held. Gr.
macropora var. *signata*, Hall, 1874, (Trematopora signata,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 104, Low. Held. Gr.
multiseriata, Hall, 1881, (Callotrypa multiseriata,) Bryozoans of the Up. Held. Gr., p. 7, and Pal. N. Y., vol. 6, p. 75, Up. Held. Gr.
oculifera, Hall, 1879, (Callotrypa oculifera,) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 155, Low. Held. Gr.
paucipora, Hall, 1887, Pal. N. Y., vol. 6, pl. 23, fig. 21, Low. Held. Gr.
striata, Hall, 1887, Pal. N. Y., vol. 6, p. 26, Low. Held. Gr.
unispina, Hall, 1874, (Callotrypa unispina,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 102, Low. Held. Gr.
Carinopora, Nicholson, 1874, Ann. and Mag. Nat. Hist., 4th ser., vol. 13, and Pal. Prov. Ont., p. 109, synonym for *Fenestella*.
hindi, Nicholson, 1874, Ann. and Mag. Nat. Hist., 4th ser., vol. 13, and Pal. Prov. Ont., p. 111. Not a good species.
CERAMELLA, Hall, 1887, Pal. N. Y., vol. 6, p. 19. [Ety. *keramis*, imbricated.] Thin, growing from a spreading base, celluliferous on both sides, tubes oblique; peristomes elevated; maculae sterile, depressed. Type *C. scidacea*.
scidacea, Hall, 1887, Pal. N. Y., vol. 6, p. 240, Ham. Gr.
CERAMOPORA, Hall, 1852, Pal. N. Y., vol. 2, p. 168. [Ety. *keramis*, imbricated like roof tile; *poros*, pore.] Discoidal, free or attached by the center of the base to foreign bodies; under surface with one or more layers of small, irregular, intercommunicating cells, which do not form tubes; cells large, oblique, imbricating, arranged in a radial manner around the depressed center, communicating with each other and the mesopores by means of remote perforations in their walls; mesopores irregular, short, numerous at the center of the colony, decreasing in number toward the margin. Type *C. imbricata*.

FIG. 466.—

ohioensis, 2, p. 2
orbicularis, Soc. N
erly d
parvicell
Mus.
rariopora
Foss.
Nat. I
CERAMOPORA
Soc. N
Sur. I
Ceram
one o
short,
lique,
by me
distincta
Ill., v
granulos
Ill., v
stellata,
vol. 8,
Ceripora,
[Ety.]
Not a
hamilton
ensis.

agellus, Hall, 1867, 28th Rep. N. Y. Mus. Nat. Hist., p. 120, Niagara Gr.
 beani, James, 1885, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 23, Hud. Riv. Gr.
 confluenta, Hall, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 119, Niagara Gr.
 explanata, Hall, 1879, Desc. New Spec. Foss., p. 5, and 11th Rep. Ind. Geo. and Nat. Hist., p. 245, Niagara Gr.
 foliacea, Hall, 1852, Pal. N. Y., vol. 2, p. 170, Niagara Gr.
 huronensis, Nicholson, 1875, Geo. Mag. n. s., vol. 2, p. 37, Ham. Gr.
 imbricata, Hall, 1852, Pal. N. Y., vol. 2, p. 169, Niagara Gr.
 incrustans, Hall, 1852, Pal. N. Y., vol. 2, p. 169, Niagara Gr.
 labecula, Hall, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 119, Niagara Gr.
 labeculoidea, Hall, 1883, Rep. St. Geol., pl. 16, fig. 1-2, and Pal. N. Y., vol. 6, p. 33, Low. Held. Gr.
 maculata, Hall, 1874, 26th Rep. N. Y. Mus. Nat. Hist., p. 108, Low. Held. Gr.
 maxima, Hall, 1874, 26th Rep. N. Y. Mus. Nat. Hist., p. 109, Low. Held. Gr.
 nicholsoni, James, 1875, Int. to Catal. Cin. Foss., p. 3, Hud. Riv. Gr.
 nothus, Hall, 1879, Desc. New Spec. Foss., p. 6, and 11th Rep. Ind. Geo. and Nat. Hist., p. 244, Niagara Gr.

FIG. 466.—*Ceratopora ohioensis*. Fragment natural size and magnified.

ohioensis, Nicholson, 1875, Ohio Pal., vol. 2, p. 265, Hud. Riv. Gr.
 orbiculata, Ringuenberg, 1886, Bull. Buf. Soc. Nat. Hist., vol. 5, p. 19. Not properly defined.
 parvicella, Hall, 1879, 32d Rep. N. Y. St. Mus. Nat. Hist., p. 158, Low. Held. Gr.
 raripora, Hall, 1879, Desc. New Spec. Foss., p. 6, and 11th Rep. Ind. Geo. and Nat. Hist., p. 244, Niagara Gr.
 CERATOPORELLA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 156, and Geo. Sur. Ill., vol. 8, (in press.) [Ety. from *Ceratopora*.] Incrusting, consisting of one or more thin layers; zoecial tubes short, apertures rounded, direct or oblique, and more or less nearly isolated by mesopores. Type *C. distincta*.
distincta, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 39, Hud. Riv. Gr.
granulosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 41, Hud. Riv. Gr.
stellata, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 41, Hud. Riv. Gr.
Ceratopora, Goldfuss, 1826, Germ. Petref. [Ety. *kerion*, honey-comb; *poros*, pore.] Not a Palaeozoic genus.
hamiltonensis, see *Streblotrypa hamiltonensis*.

CHAINODICTYON, Foerste, 1887, Bull. Denison Univ., p. 81. Zoaria flabellate, consisting of narrow inosculating branches, poriferous on one side only, the other with concentric or lunate plications. Fenestrules elliptical; zoecia subtubular in two to four alternating series, their apertures rounded and placed at the bottom of sloping areas. Type *C. laxum*.

laxum, Foerste, 1887, Bull. Denison Univ., vol. 2, p. 81, Low. Coal Meas.
laxum var. *minor*, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 62, Low. Coal Meas.
 CHILOPORELLA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 157. [Ety. *cheilos*, edge or lip; *poros*, pore; *ella*, dim.] Flabellate fronda or compressed branches, from a greatly expanded heavy crust; zoecial tubes long, very thin-walled, large, and of irregular shape in the axial region; walls much thickened near the surface; apertures ovate, the lunarium conspicuously elevated; mesopores numerous; diaphragms few, generally absent. Type *C. flabellata*.

flabellata, Ulrich, 1879, (Fistulipora flabellata,) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 28, Hud. Riv. Gr.
 CHILOTRYPA, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 49. [Ety. *cheilos*, edge; *trupa*, opening.] Ramose, small central tube to which the zoecia are attached; interstitial spaces vesiculose; diaphragms wanting, or few. Type *C. hispida*.

hispida, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 50, Kaskaskia Gr.
ostiolata, Hall, 1852, (Trematopora ostiolata,) Pal. N. Y., vol. 2, p. 152, Niagara Gr.

CLATHROPORA,

Hall, 1852, Pal. N. Y., vol. 2, p. 159. [Ety. *clathrum*, lattice; *poros*, pore.] Reticulate, uniformly poriferous on both sides of the bifoliate fronds; apertures more or less quadrangular, regularly arranged in parallel series or obliquely in quincunx order. Type *C. calcicornis*.

calcicornis, Hall, 1852, Pal. N. Y., vol. 2, p. 159, Niagara Gr.

FIG. 467.—*Clathropora frondosa*.

- carinata*, Hall, syn. for *Coscinotrypa cribriformis*.
- clintonensis*, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 113, Niagara Gr.
- flabellata*, Hall, 1851, Foster & Whitney's Rep., vol. 2, p. 207, Trenton Gr.
- frondosa*, Hall, 1852, Pal. N. Y., vol. 2, p. 160, Niagara Gr.
- gracilis*, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 54, Niagara Gr.
- intermedia*, Nicholson & Hinde, 1874, Can. Jour., p. 156, Niagara Gr.
- intertexta*, Nicholson, 1874, Geo. Mag. Lond. n. s., vol. 1, p. 125, Corniferous Gr.
- striatura*, see *Coscinium striatum*.
- CLOPOPORE**, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 20. [Ety. *klonos*, confusion; *poros*, pore.] Consisting of an aggregation of elongate, cylindrical, tubular cells, which at intervals become free and turn abruptly outward in an umbelliform expansion, or in alternation; cell apertures expanded or narrowly trumpet-shaped. Type C. *semireducta*.
- fasciculata*, Hall, 1887, Pal. N. Y. vol. 6, p. 28t, Up. Held. Gr.
- incurva*, Hall, 1881, Bryozoans of Up. Held. Gr., p. 20, Up. Held. Gr.
- semireducta*, Hall, 1881, Bryozoans of Up. Held. Gr., p. 20, Up. Held. Gr.
- CALLOCAULIS**, Hall, 1887, Pal. N. Y., vol. 6, p. 23. [Ety. *kallos*, hollow; *kaulos*, stem.] Ramose, structure like *Callopora*, but growing as hollow stems, the thin expansion lined with a striated epitheca. Type C. *venusta*.
- aculeolata*, Hall, 1881, (Callopora *aculeolata*.) Bryozoans of Up. Held. Gr., p. 7, and Pal. N. Y., vol. 6, p. 76, Up. Held. Gr.
- hyale*, Hall, 1874, (Callopora *hyale*.) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 100, and Pal. N. Y., vol. 6, p. 76, Up. Held. Gr.
- irregularis*, Hall, 1881, (Callopora *irregularis*.) Bryozoans of the Up. Held. Gr., p. 7, and Pal. N. Y., vol. 6, p. 76, Up. Held. Gr.
- mediopora*, Hall, 1887, Pal. N. Y., vol. 6, p. 23, Low. Held. Gr.
- venusta*, Hall, 1874, (Callopora *venusta*.) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 101, and Pal. N. Y., vol. 6, p. 23, Low. Held. Gr.
- CALLOCONUS**, Ulrich, Geo. Sur. Ill., vol. 8, p. 402. [Ety. *kallos*, hollow; *konos*, cone.] Zoaria simple, hollow, expanding gradually from the striated and sub-acute basal extremity, substance thin; external characters of zoecia as in *Rhombopora*; primitive portion short; hemisepta well developed. Type C. *rhombicus*.
- granosus*, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 72, Kaskaskia Gr.
- rhombicus*, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 72, St. Louis Gr.
- COSCINELLA**, Hall, 1887, Pal. N. Y., vol. 6, p. 19. [Ety. diminutive of *Coscinium*.] Distinguished from *Coscinium* by the presence of minute, angular pits between the cell apertures and around the margins of the fenestrules. Type C. *elegantula*.
- cosciniformis*, Nicholson, 1875, (Ptilodictya *cosciniformis*.) Geo. Mag., vol. 2, p. 35, and Pal. Prov. Ont., p. 80, Ham. Gr.
- elegantula*, Hall, 1887, Pal. N. Y., vol. 7, p. 239, Ham. Gr.
- COSCINIUM**, Keyserling, 1846, Geognost. beabacht., p. 192. [Ety. *koskinion*, a little sieve.] Lobed, leaf-like expansions, cells on each side, quincuncially arranged; perforated as in *Adeona cribriformis*; intercellular spaces wide, and permeated with capillary tubuli, which fill up with age; the dividing plate has a cancellous structure on either side, from the outer cellules of which the large oblique cells, terminating on the free surface, take their rise. Type C. *cyclops*.
- asterium*, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 574, Keokuk Gr.
- cribriforme*, see *Coscinotrypa cribriformis*.
- cyclops*, Keyserling, 1846, Geognost. beabacht., p. 192, Up. Held. Gr.
- elegans*, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 572, St. Louis Gr.
- escharoides*, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 574, (erroneously written *escharenæ*.) Keokuk Gr.
- keyserlingi*, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 269, Warsaw Gr.
- latum*, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 76, Burlington Gr.
- melchlini*, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 573, St. Louis Gr.
- plumosum*, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 572, St. Louis Gr.
- saganella*, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 573, St. Louis Gr.
- striatum*, Hall, 1887, Pal. N. Y., vol. 6, p. 238, Ham. Gr.
- striatum*, Hall, 1887, Pal. N. Y., vol. 6, p. 88, Up. Held. Gr.
- tuberculatum*, Prout, 1860, Trans. St. Louis Acad., vol. 1, p. 573, Keokuk Gr.
- wortheni*, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 571, Keokuk Gr.
- COSCINOTRYPA**, Hall, 1887, Pal. N. Y., vol. 6, p. 19. [Ety. *koskinion*, a little sieve; *trupa*, door.] Explanate, celluliferous on both sides, with fenestrules at varying distances; surface plicated; cells tubular, arising from a mesotheca; apertures trilobate, denticulated; intercellular tissue vesiculose. Type C. *cribriformis*.
- carinata*, Hall, syn. for C. *cribriformis*.
- cribriformis*, Prout, 1858, (Coscinium *cribriforme*.) Trans. St. Louis Acad. Sci., vol. 1, p. 267, Up. Held. Gr.
- Crateripora*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 29, C. *erecta*, C. *lineata*, and C. *lineata* var. *expansa*, rep-

FIG. 388.—
lans.
tion, sh

Crisina
bicu
CRISINEL
pl. 2
solid
obli
ter
ston
with
Type
scrib
lata,
p. 20
Held
Cryptop
Nat.
Prov
cast
mirabi
Cycloro
Acad
cle;
or in
pose
than
of a
work
mark
less
the
inter
disoid

resent the basal articulating sockets of Ptilodictya and Arthropoda, and are not entitled to rank as species.

CREPIPORA, Ulrich, (in press.) Geo. Sur. Ill., vol. 8. [Ety. *crepis*, horseshoe; *poros*, pore.] Incrusting, lamellate or massive, with a wrinkled epitheca on the lower side, in one case forming regular hollow branches; surface exhibiting, at subregular intervals, maculae of mesopores, appearing as minutely porous or subsolid elevations or depressions; zoecia very little oblique, the apertures varying from rhomboidal to subpyriform; lunarium well marked in perfect examples; best shown in tangential sections; mesopores usually restricted to the maculae; diaphragms present. Type C. simulans.

epidermata, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Group.

hemispherica, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

impressa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

simulans, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 39 and 40, Hud. Riv. Gr.

solida, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

stolidi, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

subulosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

ultrix, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

variosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

viridula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 40, Hud. Riv. Gr.

expatiata, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 68, Keokuk Gr.

fungia, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 577, Keokuk Gr.

jamesi, Prout, syn. for Ptilodictya pavonia.

polymorpha, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 578, Kaskaskia Gr.

CYCLOPORA, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, p. 404. [Ety. dim. of Cyclopora.] Thin discoidal expansion; zoecia subtubular, with a succession of superior hemisepta in the vestibular portion; irregular mesopores abundant; acanthopores of large size, numerous. Type C. spinifera.

perversa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 69, Keokuk Gr.

spinifera, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 69, Keokuk Gr.

CYSTODICTYA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist. vol. 5, p. 152. [Ety. *kusta*, a bladder; *dictyon*, net.] Zoarium like Stictopora, but with wide interstitial spaces occupied with vesicular tissue. Type C. ocellata.

americana, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 76, Keokuk Gr.

angusta, Ulrich, 1888, Bull. Denison Univ., p. 81, Waverly Gr.

hamiltonensis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 43, Ham Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

lineata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist. vol. 7, p. 37, Keokuk Gr.

FIG. 468.—*Crepipora simulans*. Tangential section, showing lunarium.

Crisina scrobiculata, see *Crisinella scrobiculata*.

CRISINELLA, Hall, 1883, Rep. St. Geol. Def., pl. 26. [Ety. from *Crisina*.] Ramose, solid, celluliferous on one side; cells in oblique, ascending rows from the center to the margin of the branch; peristomes prominent; interapertural spaces, with polygonal pits or mesopores. Type C. scrobiculata.

scrobiculata, Hall, 1881, (Crisina scrobiculata,) Bryozoans of the Up. Held. Gr., p. 20, and Pal. N. Y., vol. 6, p. 103, Up. Held. Gr.

Cryptopora, Nicholson, 1874, Ann. and Mag. Nat. Hist., 4th ser., vol. 13, and Pal. Prov. Ont., p. 102. Founded upon a cast from the under side of a Fenestella.

mirabilis, Nicholson. Not a species.

CYCLOPORA, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 574. [Ety. *kuklos*, circle; *poros*, pore.] Discoidal, frondescent or incrusting; plates sometimes superposed with subprismatic cells longer than broad, having their sides formed of a minutely porous interstitial network, developed from an epitheca marked by transverse bands more or less concentric, separating the bases of the cells; cells shallow and expanded; interstitial cells. Type C. fungia.

discoidea, see Proutella discoidea.

FIG. 469.—*Cystodycta ocellata*. Natural size and 18 diam.

lineata var. **major**, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 76, St. Louis Gr.

lineata, var. **stolidi**, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 76, St. Louis Gr.

nitida, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 76, Keokuk Gr.

ocellata, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 170, Keokuk Gr.

pustulosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 76, Keokuk Gr.

simulans, Ulrich, 1888, Bull. Denison Univ., p. 81, Waverly Gr.

zigzag, Ulrich, 1888, Bull. Denison Univ., p. 81, Cuyahoga Shales.

FIG. 470.—*Cystodycta ocellata*. Tangential section showing lunarium.

CYSTOPORA, Hall, 1881, Bryozoans of Up. Held. Gr., p. 19. [Ety. *kustis*, bladder; *poros*, pore.] Simple or branching subcylindrical stipes; cells arising from the axis; circular and subcylindrical below, enlarged above the middle and becoming ampullate, turning abruptly outward below the apertures, which are extremely contracted; cell-tubes exposed more than half their length. Type C. geniculata.

geniculata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 20, and Pal. N. Y., vol. 6, p. 103, Up. Held. Gr.

DIAMESOPORA, Hall, 1852, Pal. N. Y., vol. 2, p. 158, and vol. 6, p. 19. [Ety. *diamesos*, the part between; *poros*, opening.] Ramose, hollow, epitheca on inner surface; intercellular space solid; surface like Trematopora. Type D. dichotoma.

camerata, Hall, 1883, (Trematopora camerata.) Rep. St. Geol. and Pal. N. Y., vol. 6, p. 72, Up. Held. Gr.

communis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 39 and 41, Utica Slate.

constricta, Hall, 1874, (Trematopora constricta.) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 104, Low. Held. Gr.

dichotoma, Hall, 1852, Pal. N. Y., vol. 2, p. 158, Low. Held. Gr.

dispersa, Hall, 1879, (Trematopora dispersa.) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 150, Low. Held. Gr.

vaupeli, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 39 and 41, Utica Slate.

DICHOITYPA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 386. [Ety. *dicha*, double; *trupa*, opening.] Consisting of large bifoliate expansions; the surface with solid maculae; zoecial and minute structure as in Cystodictya. Type D. foliata.

elegans, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 76, St. Louis Gr.

expatiata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, St. Louis Gr.

flabellum, Rominger, 1866, (Fistulipora flabellum,) Proc. Acad. Nat. Sci., Phil., p. 9, St. Louis Gr.

foliata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 42, Ham. Gr.

grandis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 42, Niagara Gr.

intermedia, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 76, St. Louis Gr.

lyroides, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 77, St. Louis Gr.

DICRANOPORA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 166. [Ety. *di-* *kranos*, two-pointed; *poros*, pore.] Zoarium like Stictopora, but distinguished by being composed of ligulate joints, the edges being subparallel to near the upper end, when they diverge and bear two segments; cell-mouths between raised longitudinal lines; no interstitial cells. Type D. internodia.

emacrerata, Nicholson, 1875, (Ptilodictya emacrerata,) Pal. Ohio, vol. 2, p. 261, Hud. Riv. Gr.

fragilis, Billings, 1866, (Ptilodictya fragilis,) Catal. Sil. Foss. Antic., p. 9, Hud. Riv. Gr.

FIG. 471.—*Dicanopora internodia*. Natural size and magnified.

internodia, Miller & Dyer, 1878, (Ptilodictya internodia,) Cont. to Pal., No. 2, p. 7, Hud. Riv. Gr.

lata, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 166, Hud. Riv. Gr.

nitidula, Billings, 1866, (Ptilodictya nitidula,) Catal. Sil. Foss. Antic., p. 9, Hud. Riv. Gr.

trentonensis, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 167, Trenton Gr.

DIPLOCLEMA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 368. [Ety. *diploos*, double; *klema*, twig.] Ramose, ovate in cross section; zoecia tubular, long, apparently moniliform proximally; separated internally by an axial lamina, from which they gradually diverge to open on the two sides of the compressed branches; apertures prominent, isolated, somewhat constricted and circular; external wall thin. Type D. trentonense.

trentonense, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 53, Trenton Gr.

DIPLOPORA, Young & Young, 1875, Proc. Nat. Hist. Soc. Glasgow. [Ety. *diploos*, double; *poros*, pore.] Very slender straight stems, throwing off a few lateral branches of equal dimensions; obverse or poriferous side, with two ranges of zoecia apertures, and moderately developed medium keel; reverse striated. Type D. marginalis.

bifurcata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 62, Kaskaskia Gr.

biserialis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 62, Low. Coal Meas.

DISCOTRYPA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 155. [Ety. *diskos*, quoit; *trupa*, opening.] Free, thin, circular expansions; cells rhomboidal or hexagonal; low monticules, with clusters of large cells present; no interstitial cells or spiniform tubuli. Type D. elegans.

?devonica, Ulrich, 1886, Cont. to Am. Pal., p. 25, Up. Held. Gr.

elegans, Ulrich, 1879, (Chetetes elegans,) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 130, Hud. Riv. Gr.

ENALLOPORA, D'Orbigny, 1850, Prodr. d. Paléont., t. 1, p. 22. [Ety. *enallós*, changed; *poros*, pore.] Small bifurcating branches, without connecting bars; cell-mouths

(fragilis,
9, Hud.

tural size

3, (Ptilo-
l., No. 2,

Soc. Nat.
Gr.

ctya niti-
ic., p. 9,

Cin. Soc.
ton Gr.

Geo. Sur.
double;

in cross
appar-

separated
ina, from

s to open
impressed

isolated,
cular; ex-

ntonense.
Geo. Sur.

75, Proc.

ly. diploos,
slender

a few lat-
tions; ob-

with two
d moder-

reverse

Geo. Sur.
ir.

Geo. Sur.
Meas.

Cin. Soc.
ty. diskos,

thin, cir-
boidal or

with clus-
intersti-

Type D.

Am. Pal.,

elegans.)

.2, p. 130,

dr. d. Pa-

changed;

branches,
ell-mouths

prominent on each side, opening later-
ally and alternately. Type *E. peran-*
tiqua.

cinctosa, Ulrich, 1882, (Mitoclema *cinc-*
tosa), Jour. Cin. Soc. Nat. Hist., vol. 5,
p. 159, Trenton Gr.

perantiqua, Hall, 1847, (Gorgonia *peran-*
tiqua), Pal. N. Y., vol. 1, p. 76, Tren-
ton Gr.

ERIDOPORA, Ulrich, 1882, Jour. Cin. Soc.
Nat. Hist., vol. 5, p. 137. [Ety. *eridos*,
in dispute; *poros*, pore.] Zoarium thin,
incrusting; cell-mouths oblique, ovate,
or subtriangular, one side more promi-
nent than the other, surrounded by
angular interstitial cells, which do not
form tubes, and may be either open or
closed; intertubular spaces vesicular.
Type *E. macrostoma*. Should this genus
prove to be founded upon reliable
characters, then many of the parasitic
species now placed with *Fistulipora*
will be referred to it.

macrostoma, Ulrich, 1882, Jour. Cin. Soc.
Nat. Hist., vol. 5, p. 137, Kaskaskia Gr.
minima, Ulrich, 1886, Cont. to Am. Pal.,
p. 21, Up. Held. Gr.

punctifera, Ulrich, 1882, Jour. Cin. Nat.
Hist., vol. 5, p. 138, Kaskaskia Gr.
Eschara, Lamarck, 1801, Syst. An. sans
Vert. [Ety. *eschara*, scar.] Not Amer-
ican Palaeozoic.

? concentrica, Prout, Trans. St. Louis Acad.
Sci., vol. 1, p. 234, Coal Meas. Not
recognized.

ovatifera, Troost, 1840, 5th Geo. Rep.
Tenn. Low. Sil. Not recognized.

reticulata, Troost, 1840, 5th Geo. Rep.
Tenn. Low. Sil. Not recognized.

? tuberculata, Prout, 1858, Trans. St. Louis
Acad. Sci., vol. 1, p. 234, Coal Meas.
Not recognized.

ESCHAROPORA, Hall, 1847, Pal. N. Y., vol. 1,
p. 72. [Ety. *eschara*, scar; *poros*, pore.]
Cylindrical, solid, tapering above, ex-
panded and root-like below; cells oval,
inclosed in a rhomboid, by elevated
oblique lines; tubes radiating from an
imaginary axis. Type *E. recta*.

angusta, Hall, 1879, Desc. New Spec.
Foss., p. 6, and 11th Rep. Ind. Geo.
and Nat. Hist., p. 245, Niagara Gr.

lirata, see *Ptilodictya lirata*.

nebulosa, see *Ptilodictya nebulosa*.

recta, Hall, 1847, Pal. N. Y., vol. 1, p. 73,
Trenton Gr.

recta var. *nodosa*, Hall, 1847, Pal. N. Y.,
vol. 1, p. 73, Trenton Gr.

tenuis, see *Phanopora tenuis*.

EURYDICTYA, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, p. 389. [Ety. *eury*, broad;
dictyon, a net.] Broad, simple, or irregu-
larly divided, bifoliate expansions, with-
out nonporiferous parallel margins;
surface with more or less conspicuous,
small, solid macule or monticules;
zoecial structure very much as in *Sul-*
copora, the differences being of small
importance, and due to zoarial habit.

Type *E. montifera*. Syn. (?) for *Phae-*
nopora.

calhounensis, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, pl. 30, Trenton Gr.

montifera, Ulrich, (in press.) Geo. Sur. Ill.,
vol. 8, pl. 30, Hud. Riv. Gr.

sterlingensis, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, pl. 30, Hud. Riv. Gr.

EUSPILOPORA, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, p. 389. [Ety. *euspi*, full of
dots; *poros*, pore.] Small, bifoliate, lo-
bate or irregularly dividing branches;

cell apertures subcircular, arranged be-
tween longitudinal spinous ridges at the
center of the stipe; at intervals several
short oblique rows of cells extend out-
ward from the central rows to near the
margins of the frond; these alternate
with concave nonporiferous but finely
granular spaces, which do not extend
out as far as the celluliferous lobes,
and which cause the edges of the frond
to be serrate; internally a vertical row
of small vesicles behind the vestibul-
ar portion of the zoecia; all the re-
maining interspaces traversed by nu-
merous minute tubuli. Type *E. serrata*.
Syn. (?) for *Stictopora*.

barrisi, Ulrich, (in press.) Geol. Sur. Ill.,
vol. 8, pl. 43, Ham. Gr.

serrata, Ulrich, (in press.) Geo. Sur. Ill.,
vol. 8, pl. 43, Ham. Gr.

EVACTINOPORA, Meek & Worthen, 1865,
Proc. Acad. Nat. Sci. Phil., p. 165.

[Ety. *evactinos*, with beautiful rays;
poros, pore.] Free, consisting of four or
more vertical leaves which radiate from
an imaginary axis; rays thin, cellulif-
erous on both sides; inter-
stitial spaces occupied
by vesicular cells, filled
with sclerenchyma,
which is traversed by
canals. Type *E. radiata*.

grandis, Meek & Worthen,
1868, Geo. Sur. Ill., vol. 3, p. 503,
Burlington Gr.

FIG. 473.—*Evactinopora radiata*. Pores 2 diam.

FIG. 472.—*Evactinopora grandis*. Pores 2 diam.

quiqueradiata, Ulrich,
(in press.) Geo. Sur. Ill.,
vol. 8, pl. 73, Burling-
ton Gr.

radiata, Meek & Worthen,
1865, Proc. Acad. Nat.
Sci. Phil., p. 65, and
Geo. Sur. Ill., vol. 3, p.
502, Burlington Gr.

sexradiata, Meek & Worthen, 1868, Geo.
Sur. Ill., vol. 3, p. 502, Burlington Gr.

FAVICELLA, Hall, 1887, Pal. N. Y., vol. 6, p.

19. [Ety. *favus*, honey-comb; *ellus*,
diminutive.] Free or incrusting, thin
expansion; apertures inclosed in poly-
gonal vestibular areas, similar to *Selen-*
opora; intercellular surface occupied
by minute mesopores; structure vesic-
ulose. Type *F. inclusa*.

inclusa, Hall, 1881, (Thallostigma *inclusa*),
Trans. Alb. Inst., vol. 10, p. 188, and
Pal. N. Y., vol. 6, p. 234, Ham. Gr.

FENESTELLA, Lonsdale, 1839, Murch. Sil. Syst. [Ety. *fenestella*, little window.] Zoarium, flattened or infundibuliform, composed of rays radiating from a base and uniting laterally by diasegments, so as to form a net-work, the meshes of which are usually oblong; inner surface of rays rounded and striated, and without cells; cells on the outer side of the rays in two rows, one on each side of a median ridge; diasegments without cells. Type *F. antiqua*.
acaulis, see *Unitrypa acaulis*.
acmea, Hall, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 124, Niagara Gr.
aculeata, see *Polypora aculeata*.
acuticosta, Roemer, 1860, Sil. Fauna West. Tenn., p. 30, Niagara Gr.
adnata, see *Polypora adnata*.
adornata, Hall, 1887, Pal. N. Y., vol. vi, p. 66, Low. Held. Gr.
adraste, Hall, 1883, Rep. St. Geol., pl. 20, fig. 20-22, Low. Held. Gr.
aqualis, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 31, and Pal. N. Y., vol. 6, p. 112, Up. Held. Gr.
aeyle, Hall, 1883, Rep. St. Geol., pl. 19, fig. 11-13, and Pal. N. Y., vol. 6, p. 46, Low. Held. Gr.
albida, Hall, 1887, 6th Ann. Rep. Geo. N. Y., p. 48, Waverly Gr.
albida var. *richfieldensis*, Ulrich, 1888, Bull. Denison Univ., p. 66, Waverly Gr.
althaea, Hall, 1883, Rep. St. Geol., pl. 19, fig. 17-19, and Pal. N. Y., vol. 6, p. 48, Low. Held. Gr.
ambigua, see *Loculipora ambigua*.
angulata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 28, Up. Held. Gr.
angustata, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 60, Ham. Gr.
anonyma, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 34, Up. Held. Gr.
aperta, Hall, 1887, 6th Ann. Rep. St. Geol. N. Y., p. 58, Waverly Gr.
arctica, Salter, 1855, Belcher's Last Arctic Voyage, vol. 2, p. 385, Carboniferous.
arta, see *Polypora arta*.
aspectans, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 59, Ham. Gr.
assita, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 64, Ham. Gr.
banyana, Prout, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 450, Warsaw Gr.
belliatriata, Hall, 1879, Desc. New Spec. Foss., p. 7, and 11th Rep. Ind. Geo. and Nat. Hist., p. 252, Niagara Gr.
bicornis, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 55, Clinton Gr.
bifurca, Ulrich, 1886, Cont. to Am. Pal., p. 6, Up. Held. Gr.
bifurcata, Prout, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 411, Ham. Gr.
bigeneria, Ulrich, 1886, Cont. to Am. Pal., p. 11, Up. Held. Gr.
biimbricata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 31, and Pal. N. Y., vol. 6, p. 122, Up. Held. Gr.

biseriata, see *Hemitrypa biserialis*.
biseriata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 25, and Pal. N. Y., vol. 6, p. 113, Up. Held. Gr.
biserrulata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 30, and Pal. N. Y., vol. 6, p. 128, Up. Held. Gr.
brevillina, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 70, Ham. Gr.
breviusculata, see *Polypora brevisculata*.
burlingtonensis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 49, Burlington Gr.
cavernosa, Ulrich, 1888, Bull. Denison Univ., p. 69, Waverly Gr.
celsipora, see *Polypora celsipora*.
celsipora var. *minima*, see *Polypora celsipora* var. *minima*.
celsipora var. *minor*, see *Polypora celsipora* var. *minor*.
castriensis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 51, Kaskaskia Gr.
cinctuta, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 62, Ham. Gr.
cingulata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 52, Keokuk Gr.
clathrata, Hall, 1887, Pal. N. Y., vol. 6, p. 117, Up. Held. Gr.
cleia, Hall, 1883, Rep. St. Geol., pl. 20, fig. 14-15, Low. Held. Gr.
compacta, see *Polypora compacta*.
compressa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 50, Keokuk Gr.
compressa, var. *nododorsalis*, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 50, Keokuk Gr.
compressa, see *Polypora compressa*.
conferta, Hall, 1870, Desc. New Spec. Foss., p. 7, and 11th Rep. Ind. Geo., and Nat. Hist., p. 252, Niagara Gr.
confertipora, Hall, 1887, Pal. N. Y., vol. 6, p. 108, Up. Held. Gr.
conjunctiva, see *Isotrypa conjunctiva*.
coronis, Hall, 1883, Rep. St. Geol., pl. 21, fig. 10-13, Low. Held. Gr.
corticata, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 231, Coal Meas.
crebripora, Hall, 1874, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 25, Low. Held. Gr.
cribrosa, Hall, 1852, Pal. N. Y., vol. 2, p. 166, Niagara Gr.
cribrosa, see *Hemitrypa cribrosa*.
cultellata, see *Polypora cultellata*.
cultrata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 29, and Pal. N. Y., vol. 6, p. 119, Up. Held. Gr.
curvata, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 69, Ham. Gr.
curvijunctura, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 29, and Pal. N. Y., vol. 6, p. 107, Up. Held. Gr.
cylindracea, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 24, Up. Held. Gr.
davidsoni, Nicholson, 1875, Geo. Mag., vol. 2, n. s., p. 36, Ham. Gr.

delic
Sc

FIG. 474
leuta.

elega
164
elega
Ro
elega
pre
tise
eleva
Ill.
elong
emac
Mu
erect
Up
6, p
eudor
exigu
vol
exim
Mid
exorn
Mu
fastig
favos
filifor
1, n
per
Un
filistr
Ill.
filite
Mid
fistula
fiabell
p.
Am
fiabell
flexu
vol
foliat
p. 6
funicu
vol
gracil
granif
Up
vol
granif

FIG. 174.—*Fenestella delicata*. Part of a frond.

delicata, Meek, 1871, Proc. Acad. Nat. Sci. Phil., vol. 23, p. 159, and Ohio Pal., vol. 1, p. 273, Waverly Gr. depressa, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 30, and Pal. N. Y., vol. 6, p. 111, Up. Held. Gr.
dilatata, Prout., 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 411, Ham. Gr. diaspanda, Hall, 1887, Pal. N. Y., vol. 6, p. 114, Up. Held. Gr.
distans, see Polypora distans.
elegans, Hall, 1852, Pal. N. Y., vol. 2, p. 164, Niagara Gr.
elegantissima, Eichwald, 1860, Lethæa Rossica, p. 364, Up. Coal Meas.
elegantissima, Hall, 1881. This name was preoccupied, but see Unitrypa elegantissima.
elevatipora, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 51, Kaskaskia Gr.
elongata, see Polypora elongata.
emaciatata, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 68, Ham. Gr.
erectipora, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 33, and Pal. N. Y., vol. 6, p. 118, Up. Held. Gr.
eudora, see Polypora eudora.
exigua, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 51, Warsaw Gr.
eximia, Winchell, 1866, Rep. Low. Penin. Mich., p. 92, Ham. Gr.
exornata, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 67, Ham. Gr.
fastigata, see Unitrypa fastigata.
favosa, see Hemitrypa favosa.
filiformis, Nicholson, 1874, Geo. Mag., vol. 1, n. s., p. 199, Up. Held. Gr. The superficial network of some species of Unitrypa.
filistriata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 49, Burlington Gr.
filitexta, Winchell, 1866, Rep. Low. Penin. Mich., p. 92, Ham. Gr.
fistulata, see Polypora fistulata.
fiabellata, Phillips, 1836, Geo. York, pt. 2, p. 198, Coal Meas., or Permian. Not American. (?)
fiabelliformis, see Polypora fiabelliformis.
flexuosa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 51, Kaskaskia Gr.
foliata, Ulrich, 1888, Bull. Denison Univ., p. 67, Waverly Gr.
funicula, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 51, Keokuk Gr.
gracilis, see Subretepora gracilis.
granifera, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 33, and Pal. N. Y., vol. 6, p. 125, Up. Held. Gr.
granilinea, see Polypora granilinea.

granulosa, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 68, and Geo. Sur. Wis., vol. 4, p. 252, Hud. Riv. Gr.
hemitrypa, Prout, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 444, Warsaw Gr.
herrickana, Ulrich, 1888, Bull. Denison Univ., p. 63, Waverly Gr.
heutia, Hall, 1883, Rep. St. Geol., pl. 20, fig. 12-13, Low. Held. Gr.
hexagonalis, see Polypora hexagonalis.
hexagonalis var. *foraminulosa*, see Polypora hexagonalis var. *foraminulosa*.
idalia, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 95, Low. Held. Gr.
idothea, see Polypora idothea.
inequalis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 52, Coal Meas.
inflexa, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 64, Ham. Gr.
intermedia, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 231, Coal Meas.
interrupta, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 32, and Pal. N. Y., vol. 6, p. 123, Up. Held. Gr.
junceus, Hall, 1883, Rep. St. Geol., pl. 20, fig. 16-18, Low. Held. Gr.
laevistriata, see Polypora laevistriata.
largissima, see Polypora largissima.
lata, see Unitrypa lata.
latijunctura, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 31, and Pal. N. Y., vol. 6, p. 128, Up. Held. Gr.
latitruncata, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 58, Ham. Gr.
laevinodeata, see Polypora laevinodeata.
limbata, Foerste, 1887, Bull. Denison Univ., vol. 2, p. 83, Low. Coal Meas.
lineanoda, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 22, Up. Held. Gr.
lunulata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 31, and Pal. N. Y., vol. 6, p. 121, Up. Held. Gr.
lyelli, Dawson, 1868, Acad. Geol., p. 288, Subcarboniferous.
magnifica, Nicholson, 1874, Geo. Mag., vol. 1, n. s., p. 197, Up. Held. Gr.
marcida, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 57, Ham. Gr.
marginalis, Nicholson, 1874, Geo. Mag., vol. 1, n. s., p. 197, Up. Held. Gr.
marginata, McCoy, 1862, Carb. Foss. of Ireland, p. 206, Up. Coal Meas. Not American. (?)
meekana, Ulrich, 1888, Bull. Denison Univ., p. 64, Waverly Gr.
microtrema, D'Orbigny, 1850, Prodr. d. Paléont. Not properly defined.
mimica, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 52, Low. Coal Meas.
modesta, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 52, Low. Coal Meas.
multiplex, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 57, Ham. Gr.
multiportata var. *lodiensis*, Meek, 1875, Ohio Pal., vol. 2, p. 274, Waverly Gr.
multispinosa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 50, Keokuk Gr.
mutabilis, see Polypora mutabilis.
nerata, see Ptiloporella nervata.

nervia, see *Unitrypa nervia*.
nervia var. *constricta*, see *Unitrypa nervia* var. *constricta*.
nexa, see *Polypora nexa*.
nodosa, Prout, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 410, Ham. Gr.
norwoodana, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 233, Coal Meas.
oxfordensis, Ulrich, syn. for *F. granulosa*.
papillata, Hall, syn. for *Polypora paxillata*.
parallela, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 26, and Pal., N. Y., vol. 6, p. 107, Up. Held. Gr.
peculiaris, Hall, 1883, Rep. St. Geol., pl. 33, fig. 19-21, Up. Held. Gr.
parvulipora, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Niagara Gr.
patellifera, Ulrich, 1886, Cont. to Pal., p. 8, Up. Held. Gr.
paxillata, see *Polypora paxillata*.
perangulata, see *Polypora perangulata*.
perelegans, Meek, 1872, Pal. E. Nebraska, p. 153, Coal Meas.
perforata, see *Loculipora perforata*.
permarginata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 30, and Pal. N. Y., vol. 6, p. 127, Up. Held. Gr.
perminuta, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 52, Low. Coal Meas.
pernodosa, see *Unitrypa pernodosa*.
perplexa, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 33, and Pal. N. Y., vol. 6, p. 130, Up. Held. Gr.
pertenuis, Hall, 1879, Desc. New Spec. Foss., p. 6, and 11th Rep. Ind. Geo. and Nat. Hist., p. 251, Niagara Gr.
pertenuis, Hall, 1881. The name was preoccupied, see *F. proutana*.
perundata, see *Polypora perundata*.
perundulata, see *Reteporina perundulata*.
philia, Hall, 1883, Rep. St. Geol., pl. 20, fig. 9-11, Low. Held. Gr.
planiramosa, Hall, 1883, Rep. St. Geol., pl. 18, fig. 14-18, syn. for *Polypora compressa*.
planiramosa, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 65, Ham. Gr.
plebeia, McCoy, 1862, syn. Carb. Foss. Ireland, p. 203, Up. Coal Meas.
plumosa, see *Hemitrypa plumosa*.
popeana, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 229, Permian Gr.
porosa, see *Polypora porosa*.
precursor, see *Unitrypa precursor*.
prisca, Lonsdale, 1839, Murch. Sil. Syst., p. 178, Clinton Gr.
proceritas, Hall, 1887, Pal. N. Y., vol. 6, p. 115, Up. Held. Gr.
prolixa, Hall, 1879, Desc. New Spec. Foss., p. 8, and 11th Rep. Ind. Geo. and Nat. Hist., p. 253, Niagara Gr.
propria, see *Polypora propria*.
proutana, S. A. Miller, 1882, 2d Ed. Am. Pal. Foss., p. 291, Up. Held. Gr. Proposed instead of *F. pertenuis*, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 29, which was preoccupied.

pulchella, Ulrich, 1886, Cont. to Am. Pal., p. 9, Up. Held. Gr.
puncto-striata, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 68, Niagara Gr.
quadrangula, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 68, Ham. Gr.
quadrangularis, see *Polypora quadrangularis*.
quadrula, Hall, 1883, Rep. St. Geol., pl. 21, fig. 19-22, Low. Held. Gr.
regalis, Ulrich, 1888, Bull. Denison Univ., p. 70, and Geo. Sur. Ill., vol. 8, pl. 50, Keokuk and Waverly Grs.
remota, Foerste, 1887, Bull. Denison Univ., vol. 2, p. 84 and 87, Low. Coal Meas.
rhombifera, see *Reteporina rhombifera*.
rigida, see *Polypora rigida*.
robusta, see *Polypora robusta*.
rudis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 49, Keokuk Gr.
scalaris, see *Unitrypa scalaris*.
sculptilis, Ulrich, 1886, Cont. to Am. Pal., p. 10, Up. Held. Gr.
semitrunda, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 32, and Pal. N. Y., vol. 6, p. 125, Up. Held. Gr.
separata, see *Polypora separata*.
serrata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 28, and Pal. N. Y., vol. 6, p. 110, Up. Held. Gr.
serratura, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 50, Warsaw, St. Louis, and Kaskaskia Grs.
sevillensis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 52, Low. Coal Meas.
shumardji, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 232, Up. Coal Meas.
singularitas, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 29, and Pal. N. Y., vol. 6, p. 114, Up. Held. Gr.
sinuosa, Hall, 1887, Pal. N. Y., vol. 6, p. 116, Up. Held. Gr.
spio, Hall, 1887, Pal. N. Y., vol. 6, p. 47, Low. Held. Gr.
stellata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 29, and Pal. N. Y., vol. 6, p. 109, Up. Held. Gr.
stipata, see *Unitrypa stipata*.
striata, see *Reteporina striata*.
striatopora, see *Polypora striatopora*.
subflexuosa, Ulrich, 1888, Bull. Denison Univ., p. 68, Waverly Gr.
submutans, see *Polypora submutans*.
subretiformis, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 233, Coal Meas.
substriata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 35, Up. Held. Gr.
subtortilis, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 71, Ham. Gr.
sylvia, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 90, Low. Held. Gr.
tantulus, Hall, 1879, Desc. New Spec. Foss., p. 8, and 11th Rep. Ind. Geo. and Nat. Hist., p. 253, Niagara Gr.
tegulata, see *Unitrypa tegulata*.
tenax, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 51, Keokuk, St. Louis and Kaskaskia Grs.

tenella, Hall, 1887, Pal. N. Y., vol. 6, p. 105, Up. Held. Gr.

unicarpa, Hall, 1882, Pal. N. Y., vol. 2, p. 105, Niagara Gr.

tenula, Hall, 1882, Pal. N. Y., vol. 2, p. 51, Clinton Gr.

thyene, Hall, 1883, Rep. St. Geol., pl. 21, fig. 1-5, Low. Held. Gr.

toria, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 30, Up. Held. Gr.

trituberculata, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 228, Coal Meas.

tuberculata, Hall, 1887, Pal. N. Y., vol. 6, p. 110, Up. Held. Gr.

variabilis, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 231, Coal Meas.

varlopora, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 28, Up. Held. Gr.

vera, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 44, Ham. Gr.

verrucosa, Hall, 1883, Rep. St. Geol., pl. 33, fig. 11, and Pal. N. Y., vol. 6, p. 110, Up. Held. Gr.

virgosa, Eichwald, 1860, Lethaea Rossica, p. 358, Up. Coal Meas. Probably not American.

wortheni, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 52, Low. Coal Meas.

FENESTRALIA, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 235. [Ety. from genus *Fenestella*.]

Zoarium like *Fenestella*, from which it is distinguished by having two rows of cells on each side of the median ridge. Type *F. studovici*.

studovici, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 235, St. Louis Gr.

studovici var. compacta, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 50, St. Louis Gr.

FENESTRIPORA, Hall, 1885, Rep. St. Geol., p. 36. [Ety. *fenestra*, opening; *poros*, pore.]

Forms of *Fenestellidae* having the branches connected by diasepiments,

two ranges of cell apertures, separated by a carina bearing pores; noncelluliferous side with conspicuous pores. Type *F. biporifera*.

biporifera, Hall, 1885, Rep. St. Geol., pl. 2, fig. 17, Up. Held. Gr.

infraporosa, Ulrich, 1888, Cont. to Amer. Pal., p. 14, Up. Held. Gr.

occidentalis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 44 and 54, Ham. Gr.

FISTULIPORA, McCoy, 1849, Ann. and Mag. Nat. Hist., 2d ser., vol. 3, p. 130. [Ety. *fistula*, pipe; *poros*, pore.]

Incrusting or massive; corallites long, cylindrical, thick-walled, not in contact; tabulae

numerous; cells circular, smooth-edged; intervals between corallites filled with vesicular plates, tabulated. Type *F. minor*.

acervulosa, Rominger, 1866, Proc. Acad. Nat. Sci., p. 7, Ham. Gr.

astriata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 47 and 48, Ham. Gr.

canadensis, see *Favosites canadensis*.

carbonaria, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 45, Up. Coal Meas.

clausa, see *Meekopora clausa*.

collina, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 47 and 48, Ham. Gr.

communis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 47 and 48, Ham. Gr.

compressa, Rominger, 1866, Proc. Acad. Nat. Sci., p. 10, Keokuk Gr.

confertipora, Hall, 1881, (Thallostigma confertipora,) Trans. Alb. Inst., vol. 10, p. 184, and Pal. N. Y., vol. 6, p. 211, Ham. Gr.

constricta, Hall, 1881, (Lichenalia constricta,) Trans. Alb. Inst., vol. 10, p. 183, and Pal. N. Y., vol. 6, p. 227, Ham. Gr.

corrugata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 47 and 48, Ham. Gr.

crassa, Rominger, 1866, Proc. Acad. Nat. Sci., p. 8, Ham. Gr.

decipiens, Hall, 1881, (Thallostigma decipiens,) Trans. Alb. Inst., vol. 10, p. 187, and Pal. N. Y., vol. 6, p. 232, Ham. Gr.

densa, Hall, 1881, (Thallostigma densa,) Trans. Alb. Inst., vol. 10, p. 186, and Pal. N. Y., vol. 6, p. 231, Ham. Gr.

digitata, Hall, 1881, (Thallostigma digitata,) Trans. Alb. Inst., vol. 10, p. 185, and Pal. N. Y., vol. 6, p. 229, Ham. Gr.

elegans, see *Pinacotrypa elegans*.

erianis, Rominger 1866, Proc. Acad. Nat. Sci., p. 8, Ham. Gr.

excellens, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 46, Kaskaskia Gr.

flabellata, see *Chiloporella flabellata*.

flabellum, see *Dichotrypa flabellum*.

foordi, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 47 and 48, Ham. Gr.

halli, Rominger, 1866, Proc. Acad. Nat. Sci., p. 6, Niagara Gr.

helios, Rominger, 1866, Proc. Acad. Nat. Sci., p. 7, Corniferous Gr.

hemispherica, Hall, 1881, (Callopora hemispherica,) Trans. Alb. Inst., vol. 10, p. 183, and Pal. N. Y., vol. 6, p. 226, Ham. Gr.

FIG. 470.—Fenestripora foordi. Tangential section, showing lunulum.

FIG. 475.—Fenestripora biporifera. Magnified nonporiferous side.

FIG. 477.—Fenestripora foordi. Opercular cover x 50.

incrassata, Nicholson, 1874, (Callopora incrassata,) Geo. Mag. Lond. n. s., vol. 1, p. 13, and Rep. Pal. Ont., p. 61, Ham. Gr. intercellata, Hall, 1881, (Thallostigma intercellata,) Trans. Alb. Inst., vol. 10, p. 13, and Pal. N. Y., vol. 6, p. 87, Up. Held. Gr.

involvens, Hall, 1887, Pal. N. Y., vol. 6, p. 221, Ham. Gr.

labiosa, Winchell, 1866, Rep. Low. Penin. Mich., p. 88, Ham. Gr.

lamellata, Hall, 1881, (Thallostigma lamellata,) Trans. Alb. Inst., vol. 10, p. 13, and Pal. N. Y., vol. 6, p. 87, Up. Held. Gr.

lens, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 69, and Geo. Wis., vol. 4, p. 256, Hud. Riv. Gr.

longimacula, Hall, 1881, (Thallostigma longimacula,) Trans. Alb. Inst., vol. 10, p. 185, and Pal. N. Y., vol. 6, p. 209, Ham. Gr.

lunata, see Buscopora lunata.

micropora, Hall, 1884, Thallostigma micropora, Rep. St. Geol., p. 26, Ham. Gr.

minuta, Rominger, 1866, Proc. Acad. Nat. Sci., p. 7, Ham. Gr.

monticulata, Ulrich, Geo. Sur. Ill., vol. 8, pl. 47 and 48, Ham. Gr.

multiculatea, Hall, 1884, (Thallostigma multiculatea,) Rep. St. Geol., p. 23, Ham. Gr.

neglecta, Rominger, 1866, Proc. Acad. Nat. Sci., p. 6, syn. for Lichenalia concentrica.

FIG. 478.—*Flustra* (?) *tuberculata*. Sections x 50.

nodulifera, Meek, 1872, Pal. E. Neb., p. 143, Up. Coal Meas.

normalis, Ulrich, 1886, Cont. to Am. Pal., p. 20, Up. Held. Gr.

occidens, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 229, Chemung Gr.

oweni, James, 1885, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 21, Hud. Riv. Gr. Poorly defined.

parasitica, Hall, 1879, (Callopora parasitica,) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 157, Low. Held. Gr.

peculiaris, see Actinotrypa peculiaris.

prolifera, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 45, St. Louis Gr.

proporoides, Nicholson, 1879, Pal. Tab. Corals, p. 310, Ham. Gr.

ponderosa, Hall, 1874, (Callopora ponderosa,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 103, Low. Held. Gr.

rugosa, see Batostoma rugosa.

saffordii, Winchell, 1866, Rep. Low. Penin. Mich., p. 88, Ham. Gr.

scrobiculata, Hall, 1884, (Thallostigma scrobiculata,) Rep. St. Geol., p. 20, Ham. Gr.

segregata, Hall, 1884, (Thallostigma segregata,) Rep. St. Geol., p. 27, Ham. Gr.

serrulata, Hall, 1884, (Thallostigma serrulata,) Rep. St. Geol., p. 22, Ham. Gr.

solidissima, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 69, and Geo. Wis., vol. 4, p. 255, Hud. Riv. Gr.

spergenensis, Rominger, 1866, Proc. Acad. Nat. Sci., p. 9, Warsaw Gr.

spherioidea, Hall, 1884, (Thallostigma spherioidea,) Rep. St. Geol., p. 31, Ham. Gr.

spinulifera, Rominger, 1866, Proc. Acad. Nat. Sci., p. 8, Ham. Gr.

stellifera, Rominger, 1866, Proc. Acad. Nat. Sci., p. 7, Ham. Gr.

subtilis, Hall, 1884, (Thallostigma subtilis,) Rep. St. Geol., p. 30, Ham. Gr.

sulcata, Rominger, 1866, Proc. Acad. Nat. Sci., p. 7, Ham. Gr.

triangularis, Hall, 1884, (Thallostigma triangularis,) Rep. St. Geol., p. 32, Ham. Gr.

trifaria, Hall, 1887, Pal. N. Y., vol. 6, p. 222, Ham. Gr.

trifolia, Rominger, 1866, Proc. Acad. Nat. Sci., p. 9, Keokuk Gr.

triloba, Hall, 1887, Pal. N. Y., vol. 6, p. 29, Low. Held. Gr.

umbilicata, Hall, 1884, (Thallostigma umbilicata,) Rep. St. Geol., p. 23, Ham. Gr.

unilinea, Hall, 1887, Pal. N. Y., vol. 6, p. 217, Ham. Gr.

utriculus, Rominger, 1866, Proc. Acad. Nat. Sci., p. 8, Ham. Gr.

variopora, Hall, 1884, (Thallostigma variopora,) Rep. St. Geol., p. 18, Ham. Gr.

Flustra, Linnaeus, 1745, Amoenitates academicae. Not Palaeozoic.

carbascoides, Eaton, 1832, Geo. Text Book, p. 44. Not recognized.

spatulata, see Worthenopora spatulata.

tuberculata, Prout, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 447, Warsaw Gr. Not a flustra.

FIG. 479.—*Flustra* (?) *tuberculata*. Aperture x 50.

GLAUC
vo
in
my
ste
of
of
ate

FIG.

flexu
vol.
intern
terr
Cuy
minor
Bul
Sha
nereic
p. 1
vol.
nodat
Hel
p. 1
simul
sim
Cuy
sinuos
Hel
p. 1
suban
sub
76,
tenuir
79,
tenuis
Up.
vol.
triline
157,
vinii,
Bul
whitii
Bul
young
you
Sha
Glossor
p. x
open
ing

(GLAUCOME, Goldfuss, 1826, Germ. Petref., vol. 1, p. 100, as emended by Lonsdale in Murch. Sil. Syst., p. 677. [Ety. mythological name.] Narrow central stem, with lateral branches; two rows of cells separated by a keel on the face of each branch, and opposite side striated. Type *G. disticha*.

FIG. 480.—*Glaucome nereidis*.

bellula, Ulrich, (Pinnatopora bellula,) Geo. Sur. Ill., vol. 8, pl. 66, Low. Coal Meas. carinata, Hall, 1884, Rep. St. Geol., p. 60, Ham. Gr. curvata, Ulrich, 1888, (Pinnatopora curvata,) Bull. Denison Univ., p. 76, Cuyahoga Shales. flexuosa, Ulrich, (in press), (Geo. Sur. Ill., vol. 8, pl. 66, Keokuk Gr. intermedia, Ulrich, 1888, (Pinnatopora intermedia,) Bull. Denison Univ., p. 74, Cuyahoga Shales. minor, Ulrich, 1888, (Pinnatopora minor,) Bull. Denison Univ., p. 77, Cuyahoga Shales. nereidis, White, 1874, Rep. Invert. Foss., p. 18, and Geo. Sur. W. 100th Mer., vol. 4, p. 105, Carboniferous. nodata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 18, and Pal. N. Y., vol. 6, p. 102, Up. Held. Gr. simulatrix, Ulrich, 1888, (Pinnatopora simulatrix,) Bull. Denison Univ., p. 75, Cuyahoga Shales. sinuosa, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 18, and Pal. N. Y., vol. 6, p. 101, Up. Held. Gr. subangulata, Ulrich, 1888, (Pinnatopora subangulata,) Bull. Denison Univ., p. 76, Cuyahoga Shales. tenuiramosa, Ulrich, 1888, (Pinnatopora tenuiramosa,) Bull. Denison Univ., p. 79, Cuyahoga Shales. tenuistriata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 19, and Pal. N. Y., vol. 6, p. 102, Up. Held. Gr. trilineata, Meek, 1872, Pal. E. Neb., p. 157, Coal Meas. vinii, Ulrich, 1888, (Pinnatopora vinii,) Bull. Univ., p. 77, Cuyahoga Shales. whitii, Foerste, 1887, (Pinnatopora whitii,) Bull. Univ., p. 78, Low. Coal Meas. youngi, Ulrich, 1888, (Pinnatopora youngi,) Bull. Univ., p. 78, Cuyahoga Shales. GLAUCOTRYPA, Hall, 1887, Pal. N. Y., vol. 6, p. xvii. [Ety. glosse, the tongue; trupa, opening.] Zoarium tubular; cells arising from the epitheca lining the cylind-

rical frond, intersected by narrow projections from the cell walls, extending partially across the cell tube; apertures paliform; intercellular structure vesiculose. Type *G. paliformis*. paliformis, Hall, 1881, (Lichenalia paliformis,) Trans. Alb. Inst., vol. 10, p. 11, and Pal. N. Y., vol. 6, p. 85, Up. Held. Gr. Glyptotrypa, Ulrich, syn. (?) for Coscinium.

GONOTRYPA,

Ulrich,

(in press),

Micropal-

aeontolog-

gy, p. 14.

[Ety. gon-

nia, an-

gle; trupa,

opening.]

Bifoliate,

jointed,

segments

small,

each face

with a

central

ridge;

cells in

longitudinal

rows; apertures

oval, directed

obliquely outward.

Type

(*G. bilateralis*.

Syn. (?) for

Dicranopora.

bilateralis, Ulrich, (in press), Micropal-

aeontology, p. 15, Hud. Riv. Gr. (?)

Gorgonio, Linnaeus, 1745, *Amoenitates Acad.*

[Ety. mythological name.] Not Amer-

ican Paleozoic.

anticorum, Castelnau, 1843, *Syst. Sil.*, p. 50.

Not recognized.

(?) *aspera*, see *Subretrepora aspera*.

dubia, Goldfuss, 1826, *Petref. Germ. Per-*

mian. Not recognized.

ehrenbergi, see *Phyllopora ehrenbergi*.

infundibuliformis, Eaton, 1832, (Geo. Text

Book, p. 43. Not recognized.

perantiqua, see *Enallopora perantiqua*.

retiformis, see *Dictyonema retiforme*.

siluriana, Castelnau, 1843, *Syst. Sil.*, p.

50. Not recognized.

FIG. 481.—*Goniotrypa bilateralis*. 1, Transverse section; 2, transverse section lower down; 3, tangential section of the upper part.

bilateralis, Ulrich, (in press), Micropalaeontology, p. 15, Hud. Riv. Gr. (?) *Gorgonio*, Linnaeus, 1745, *Amoenitates Acad.* [Ety. mythological name.] Not American Paleozoic. *anticorum*, Castelnau, 1843, *Syst. Sil.*, p. 50. Not recognized. (?) *aspera*, see *Subretrepora aspera*. *dubia*, Goldfuss, 1826, *Petref. Germ. Permian*. Not recognized. *ehrenbergi*, see *Phyllopora ehrenbergi*. *infundibuliformis*, Eaton, 1832, (Geo. Text Book, p. 43. Not recognized. *perantiqua*, see *Enallopora perantiqua*. *retiformis*, see *Dictyonema retiforme*. *siluriana*, Castelnau, 1843, *Syst. Sil.*, p. 50. Not recognized.

FIG. 482.—*Graptodictya nitida*. Natural size and magnified.

GRAPTODICTYA, Ulrich, 1882, *Jour. Cin. Soc. Nat. Hist.*, vol. 5, p. 165. [Ety. grapho, I write; dictyon, net.] Zoarium pointed below, branching above, cell apertures circular, and separated by interstitial pits or sulci; distinguished from *Ptilodictya* by the circular cells and surrounding pits. Type *G. perelegans*.

nitida, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 166, Hud. Riv. Gr.
 perelegans, Ulrich, 1878, (Ptilodictya perelagans,) Jour. Cin. Soc. Nat. Hist., vol. 1, p. 94, Hud. Riv. Gr.

FIG. 483.—*Hederella canadensis*. Magnified.

HEDERELLA, Hall, 1884, Rep. St. Geol., p. 53. [Ety. *hedera*, ivy.] Bryozoom parasitic, procumbent, attached the entire length; main axis tubular, from which proceed lateral tubular cells, giving it the general appearance of Stomatopora. Type *H. canadensis*.

canadensis, Nicholson, 1873, (Alecto (?) *canadensis*.) Can. Nat. and Geol., vol. 7, p. 144, and Pal. Prov. Ont., p. 124, Up. Held. and Ham. Gr.
cirrhusa, Hall, 1884, Rep. St. Geol., p. 53, Ham. Gr.
conferta, Hall, 1884, (Ptilionella *conferta*.) Rep. St. Geol., p. 56, Ham. Gr.
filiformis, Billings, 1858, (Aulopora *filiformis*.) Can. Jour., vol. 4, p. 119, Ham. Gr.
magna, Hall, 1884, Rep. St. Geol., p. 55, Ham. Gr.

HELICOPORA, Claypole, 1883, Quar. Jour. Geo. Soc., p. 30. [Ety. *helix*, spiral; *poros*, pore.] Bryozoom expanded, fenestrate, and spiral; formed of slender, bifurcating rays, poriferous on one face, connected by nonporiferous bars, forming an open net-work; cells arranged in two rows along the rays, one row on each side of a median keel; axis none, or consisting only of the thickened inner border of the bryozoom, not straight, but forming a spiral, rounded, nonporiferous, or slightly poriferous, inner margin. Type *H. latispinalis*. Regarded by some as a synonym for *Fenestella*, and distinguished only by the spiral form.

archimedisformis, Claypole, syn. for *Archimedes laxus*.
latispinalis, Claypole, 1883, Quar. Jour. Geo. Sci., p. 32, Niagara Gr.
ulrichi, Claypole, 1883, Quar. Jour. Geo. Soc., p. 33, Up. Held. Gr.

FIG. 484.—*Heliotrypa bifolia*. Tangential section x 50.

HELIOTRYPA, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 277. [Ety. *helios*, sun;

trypa, opening.] Bifoliate, interstitial cells developed from the prostrate por-

tion of the zoecia; intercommunication by means of radially arranged tubuli. Type *H. bifolia*.

bifolia, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 278, Kaskaskia Gr.
 HELOPORA, Hall, 1852, Pal. N. Y., vol. 2, p. 44. [Ety. *helos*, nail; *poros*, pore.] Zoarium jointed; segments small, simple, cylindrical, often swollen at the extremities; cells oval or subangular, and arranged between longitudinal elevated lines or in quincunx. Type *H. fragilis*.
armata, Billings, 1866, Catal. Sil. Foss. Antic., p. 38, Anticosti Gr.
bellula, Billings, 1866, Catal. Sil. Foss. Antic., p. 38, Anticosti Gr.
circe, Billings, 1866, Catal. Sil. Foss. Antic., p. 39, Anticosti Gr.

FIG. 485.—a-l, *Helopora fragilis*; f-h, *Helopora lindstromi*, sections x 50.

conceva, Billings, 1866, Catal. Sil. Foss. Antic., p. 37, Anticosti Gr.
divaricata, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 59, Trenton Gr.
formosa, Billings, 1865, Catal. Sil. Foss. Antic., p. 37, Anticosti Gr.
fragilis, Hall, 1852, Pal. N. Y., vol. 2, p. 44, Clinton Gr.
fragilis var. *acadiensis*, Hall, 1860, Can. Nat. and Geo., vol. 5, Anticosti Gr.
irregularis, Billings, 1866, Catal. Sil. Foss. Antic., p. 39, Anticosti Gr.
imbricata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 29, Hud. Riv. Gr.

FIG. 486.—*Helopora fragilis*. Natural size and magnified.

lineata, Billings, 1866, Catal. Sil. Foss. Antic., p. 36, Anticosti Gr.
lineopora, Billings, 1866, Catal. Sil. Foss. Antic., p. 38, Anticosti Gr.
nodosa, Billings, 1866, Catal. Sil. Foss. Antic., p. 38, Anticosti Gr.
spiniformis, Ulrich, 1882, (Arthroclema spiniforme.) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 161, Trenton Gr.
striatopora, Billings, 1866, Catal. Sil. Foss. Antic., p. 39, Anticosti Gr.
strigosa, Billings, 1866, Catal. Sil. Foss. Antic., p. 37, Anticosti Gr.
tenuis, see *Arthrostylus tenuis*.
varipora, Billings, 1866, Catal. Sil. Foss. Antic., p. 40, Anticosti Gr.

FIG. 487. — Hemitrypa biordo.

are elevated, widened at the summit, and connected by scale, which meet midway and coalesce, forming pseudocarinæ. Type *H. oculata*.

aspera, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 57, Keokuk Gr.
biordo, Hall, 1887, Pal. N. Y., vol. 6, p. 149, Up. Held. Gr.
biserialis, Hall, 1879, 32d Rep. N. Y. St. Mus. Nat. Hist., p. 174, Low. Held. Gr.
biserialis var. *exilis*, Hall, 1887, Pal. N. Y., vol. 6, p. 57, Low. Held. Gr.
columellata, Hall, 1887, Pal. N. Y., vol. 6, p. 146, Up. Held. Gr.
cribrosa, Hall, 1881, (Fenestella cribrosa.) Trans. Alb. Inst., vol. 10, p. 35, Up. Held. Gr.
dubia, syn. for *Loculipora ambigua*.
favosa, Hall, 1881, Trans. Alb. Inst., vol. 10, p. 35, and Pal. N. Y., vol. 6, p. 148, Up. Held. Gr.
nodosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 57, Keokuk Gr.
prima, Hall, syn. for *Unitrypa nervia*.
pateriformis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 57, Keokuk Gr.
perstriata, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 57, Keokuk Gr.
plumosa, Prout, 1858, (Fenestella plumosa.) Trans. St. Louis Acad. Sci., vol. 1, p. 236, Keokuk and Warsaw Gr.
proutana, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 57, proposed instead of *Fenestella hemitrypa* of Prout, Keokuk and Warsaw Grs.
proutana var. *nodulosa*, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 57, Keokuk Gr.
proutana var. *vermifera*, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 57, Warsaw Gr.
tenera, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 44, Hamilton Gr.

ulrichi, Foerste, 1887, Bull. Denison Univ., vol. 2, p. 152, Clinton Gr.

HERNODIA, Hall, 1884, Rep. St. Geol., p. 58.

[Ety. *hernodes*, like a young sprout.] Bryozoon parasitic, procumbent, increasing by gemination like *Aulopora*; budding lateral, and for some distance in contact and frequently coalescing with the parent cells. Type *H. humifusa*.

humifusa, Hall, 1884, Rep. St. Geol., p. 58, Ham. Gr.

HETERODICTYA, Nicholson, 1875, Geo. Mag., vol. 2, n. s., p. 33, and Pal. Prov. Ont., p. 79. [Ety. *heteros*, irregular; *dictyon*, net.] The correct orthography is *Heterodictyon*. Flattened, two-edged frond, with subparallel sides, consisting of two series of cells upon opposite sides of a central membrane; cells are in longitudinal rows; tabulæ present. Type *H. gigantea*.

gigantea, Nicholson, 1875, Geo. Mag., vol. 2, p. 34, and Pal. Prov. Ont., p. 79, Subcarboniferous.

FIG. 488. — Heterodictya gigantea. Magnified.

Hippothoa, Lamouroux, 1821.

Expos. method. Not Palæozoic.

inflata, see *Stomatopora inflata*.

HOMOTRYPA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 240. [Ety. *homos*, similar; *trupa*, perforation.] Ramose or subfrondescent; surface smooth or bearing monticules; cells, circular, ovate or polygonal, thin-walled; groups of larger-sized cells; mesopores absent or restricted to the macule; spiniform tubuli, diaphragms and cystiphragms present. Type *H. curvata*.

arbuscula, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 38, Birdseye Gr.

curvata, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 241, Hud. Riv. Gr.

FIG. 489.—Homotrypa obliqua. Natural size and magnified.

exilis, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 80, Trenton Gr.

flabellaris, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 32, Hud. Riv. Gr.

gelasinosa, Ulrich, (in press), Geo. Sur. Ill., vol. 8, pl. 32, Hud. Riv. Gr.
 insignis, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 82, Trenton Gr.
 minnesotensis, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 79, Trenton Gr.
 obliqua, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 243, Hud. Riv. Gr.
 subramosa, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 81, Trenton Gr.

HOMOTRYPELLA, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 83. [Ety. *homotrypa*, a genus; *ellus*, diminutive.] Zoarium ramose; monticules wanting; interstitial cells present; zoecia small, moderately thick walls, and cystoid diaphragms straight; spiniform tubuli numerous. Type H. instabilis.

contexta, Ulrich, (in press), Geo. Sur. Ill., vol. 8, pl. 32, Hud. Riv. Gr.

granulifera, Ulrich, 1879, (Chetetes granuliferus,) Jour. Cin. Soc. Nat. Hist., vol. 2, p. 128, Trenton Gr.

instabilis, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 83, Trenton Gr.

HORNERA, Lamouroux, 1821, Expos. Method. des genres de L'Ordre des Pol. [Ety. proper name.] Not American Palæozoic.

dichotoma, see *Subretopora dichotoma*.

ICHTHYORACHIS, McCoy, 1844, Carb. Foss. Ireland, p. 205. [Ety. *ichthys*, fish; *rachis*, backbone.] Bryozoom plumose,

consisting of a rachis, with short lateral branches or pinnules; celluliferous on one side; cell apertures in two ranges on the branches, and in three or more on the main stem. Type I. nowenhami.

neris, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 98, Low. Held. Gr.

IDIOTRYPA, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 272. [Ety. *idios*, peculiar; *trypa*, opening.] Parasitic, interstitial cells angular, both cells with diaphragms; spiniform tubuli present. Type I. parasitica.

parasitica, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 273, Niagara Gr.

INTRAPORA, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 16. [Ety. *intra*, within; *poros*, pore.] Resembling *Stictopora*, branches broad; intercellular spaces regularly punctured or pitted, as if by minute cell apertures; cells with rounded mouths and short prostrate portion; intercellular space vesiculose. Type I. puteolata.

puteolata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 16, and Pal. N. Y., vol. 6, p. 97, Up. Held. Gr.

INTRICARIA, DeFrance, 1823, Dictionnaire des Sciences Naturelles. Not a Palæozoic genus.

clathrata, see *Subretopora clathrata*.

reticulata, see *Subretopora reticulata*.

ISOTRYPA, Hall, 1885, Rep. St. Geol., p. 37. [Ety. *isos*, equal; *trypa*, perforation.] Fenestelloid, having the branches con-

nected by dissepiments, and with two ranges of cell apertures, separated by carinae, elevated and much thickened above, connected by distinct lateral processes; the reverse face has on or near the dissepiments conspicuous pores larger than the cell apertures. Type I. conjunctiva.

FIG. 490.—*Isotrypa conjunctiva*. Celluliferous side and noncelluliferous side, with pores on the dissepiments.

bifaria, syn. for I. conjunctiva.

conjunctiva, Hall, 1881, (Fenestella conjunctiva,) Trans. Alb. Inst., vol. 10, p. 143, and Pal. N. Y., vol. 6, p. 143, Up. Held. Gr.

consimilis, Hall, 1885, Rep. St. Geol., pl. 2, fig. 14, Up. Held. Gr.

LABECIHA, Edwards & Haime, 1851, Pol. Foss. des Terr. Pal., p. 297. [Ety. proper name.] Bryozoom laminar, incrusting, or attached by part of the base, and having the remainder covered by an epitheca; surface covered with rounded or elongated, solid, tubercles, separated by an imperforate calcareous membrane; internally it consists of vertical columns extending from the epitheca below, and terminating above in the surface tubercles, the interspaces between the columns consisting of lenticular vesicles, the uppermost layer of which gives rise to the seemingly imperforate membrane between the tubercles. Type L. conferta. Probably this genus belongs to the Protozoa, and is related to the sponges.

montifera, Ulrich, 1886, Cont. to Am. Pal., p. 33, Hud. Riv. Gr.

LEOCLEMA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 141. [Ety. *leios*, smooth; *klerma*, twig.] Ramose, lamellate, or parasitic; surface even; cell-mouths small, rounded, surrounded by interstitial cells; tubes thin-walled; diaphragms remote; acanthopores abundant. Type L. punctatum.

araneum, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 75, Kaskaskia Gr.

foliatum, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 301, Warsaw Gr.

gracill
 Ill.,
 minut
 pora
 tario

FIG. 491.—*Leptotrypa*. Cross-section of zoecia, showing acanthopores, magnified 50 times.

puncta
 tata,
 and V
 subglo
 Ill., v
 wachm
 Ill., v
 wilming
 Geo.
 Riv. C

LEPTOTRYPA

Nat. I
 thin;
 crust
 surfac
 bnli;
 Type
 clavaco
 coidea
 Nicho
 p. 182

clavis, U
 Hist.,
 cortex,
 Hist.,
 hexagor
 Ill., v
 minima
 Hist.,
 ornata,
 Hist.,
 semipila
 Ill., v
 stidham
 Ill., v

LICHENALIA

p. 171.
 Memb
 circula
 ous on
 an ep
 from t
 triloba

gracillimum, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 75, Keokuk Gr.
minutissimum, Nicholson, 1875, (Callopora minutissima.) Pal. Prov. of Ontario, p. 77, Hamilton Gr.

tural space smooth: intercellular space vesiculose. Type *L. concentrica*.
alternata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 8, and Pal. N. Y., vol. 6, p. 80, Up. Held. Gr.

FIG. 491.—*Leloclema foliatum*. a, Vertical section x28, showing entire thickness of zoarium, tabulation of zooecia and mesopores, and structure of the acanthopores; b, horizontal section x28, showing distribution of acanthopores, mesopores, and zooecia; c, small portion of wall x50; d, acanthopore x50, showing its structure.

punctatum, Hall, 1858, Callopora punctata, Geo. Sur. Iowa, p. 653, Keokuk and Warsaw Grs.
subglobosum, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 75, Kinderhook Gr.
wachsmuthi, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 75, Kinderhook Gr.
wilmingtonense, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 34, Hud. Riv. Gr.

LEPTOTRYPA, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 158. [Ety. *leptos*, thin; *trupa*, perforation.] Thin, incrusting; cells polygonal, thin-walled; surface, with monticules; spiniform tubuli; no diaphragms or rudimentary. Type *L. minima*.

clavacoidea, James, 1875, (Chetetes clavacoidea.) Int. Catal. Cin. Foss., p. 2, and Nicholson on Struct. and Affin. Montic, p. 182, Hud. Riv. Gr.

clavis, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 161, Utica Slate.

cortex, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 162, Utica Slate.

hexagonalis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 36, Trenton Gr.

minima, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 159, Hud. Riv. Gr.

ornata, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 160, Hud. Riv. Gr.

semipilaris, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 36, Hud. Riv. Gr.

stidhami, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 36, Hud. Riv. Gr.

LICHENALLA, Hall, 1852, Pal. N. Y., vol. 2, p. 171. [Sig. from resemblance to a lichen.]

Membranous expansions, growing in circular or flabellate forms, celluliferous on one side, the other covered with an epitheca; cells septate, arising from the epitheca; apertures circular or trilobate, often denticulate; intersper-

Geol., p. 37, Ham. Gr.
colliculata, Hall, 1884, Rep. St. Geol., p. 36, Ham. Gr.
complexata, see *Selenopora complexata*.
concentrica, Hall, 1852, Pal. N. Y., vol. 2, p. 171, Niagara Gr.

FIG. 492.—*Lichenalla concentrica*. Under surface and upper surface magnified.

concentrica var. *maculata*, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 117, Niagara Gr.

concentrica var. *parvula*, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 117, Niagara Gr.

confusa, Hall, 1887, Pal. N. Y., vol. 6, p. 204, Ham. Gr.

constricta, see *Fistulipora constricta*.

conulata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 9, and Pal. N. Y., vol. 6, p. 81, Up. Held. Gr.

cornuta, Hall, 1887, Pal. N. Y., vol. 6, p. 203, Ham. Gr.

crassa, Hall, 1879, (Trematopora crassa,) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 152, Low Held. Gr.

crustacea, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 8, Up. Held. Gr.

cultellata, Hall, 1884, Rep. St. Geol., p. 35, Ham. Gr.

denticulata, see *Pileotrypa denticulata*.

alveata, see *Odontotrypa alveata*.

bistriata, Hall, 1881,

Bryozoans of the Up. Held. Gr., p.

8, and Pal. N. Y., vol. 6, p. 79, Up.

Held. Gr.

bullata, Hall, 1887,

Pal. N. Y., vol. 6, p. 205, Ham. Gr.

carinata, Hall, 1881,

Bryozoans of the Up. Held. Gr., p.

9, Up. Held. Gr.

circincta, see *Selenopora circincta*.

clivulata, see *Pileotrypa clivulata*.

clypeiformis, Hall,

1884, Rep. St.

- dissimilis, Hall, 1883, Rep. St. Geol., pl. 15, fig. 10-13, Low. Held. Gr.
 distans, Hall, 1883, Rep. St. Geol., pl. 15, fig. 8-9, Low. Held. Gr.
 foliacea, Hall, 1884, Rep. St. Geol., p. 35, Ham. Gr.
 geometrica, Hall, 1887, Pal. N. Y., vol. 6, p. 79, Up. Held. Gr.
granifera, see *Pileotrypa granifera*.
 imbricella, Hall, 1884, Rep. St. Geol., p. 35, Ham. Gr.
longispina, see *Lichenotrypa longispina*.
lunata, see *Puscopora lunata*.
 maculosa, Hall, 1884, (Trematopora maculosa,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 106, Low. Held. Gr.
 operculata, Hall, 1887, Pal. N. Y., vol. 6, p. 205, Ham. Gr.
 ovata, Hall, 1887, Pal. N. Y., vol. 6, p. 80, Up. Held. Gr.
paliformis, see *Glossotrypa paliformis*.
 permarginata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 10, and Rep. St. Geol., 1883, pl. 24, fig. 20, Up. Held. Gr.
 pustulosa, Hall, 1887, Pal. N. Y., vol. 6, p. 206, Ham. Gr.
pyriformis, see *Pileotrypa pyriformis*.
 radiata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 10, Up. Held. Gr.
 ramosa, Hall, 1887, Pal. N. Y., vol. 6, p. 199, Ham. Gr.
 serialis, Hall, 1887, Pal. N. Y., vol. 6, p. 32, Low. Held. Gr.
 stellata, Hall, 1884, Rep. St. Geol., p. 33, Ham. Gr.
 subcava, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 8, and Rep. St. Geol., 1883, pl. 24, fig. 23-25, Up. Held. Gr.
 substellata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 7, and Rep. St. Geol., 1883, pl. 24, fig. 26, Up. Held. Gr.
 subtrigona, Hall, 1887, Pal. N. Y., vol. 6, p. 196, Ham. Gr.
 tessellata, Hall, 1887, Pal. N. Y., vol. 6, p. 207, Ham. Gr.
 torta, Hall, 1883, Rep. St. Geol., pl. 15, fig. 1-7, Low. Held. Gr.
 tortuosa, Hall, 1883, Rep. St. Geol., pl. 13, fig. 17-18, Low. Held. Gr.
 vesiculata, Hall, 1887, Pal. N. Y., vol. 6, p. 197, Ham. Gr.
- LICHENOTRYPA, Ulrich, 1886, Cont. to Am. Pal., p. 23. [Ety. *lichen*, a tree-moss; *trupa*, perforation.] Zoarium thin, incrusting, in early growth like *Fistulipora*, with short, tubular zoecia, wide, concave interspaces, subcircular apertures, posterior margin elevated; in later growth peristomes of adjacent cells unite by thin, irregular walls, which traverse the interstitial spaces, and form an irregular net-work, with spine-like elevations; interstitial cells present. Type *L. cavernosa*. Syn. (?) for *Lichenalia*.
cavernosa, Ulrich, 1886, Cont. to Am. Pal., p. 24, Up. Held. Gr.
longispina, Hall, 1881, (*Lichenalia longispina*), Trans. Alb. Inst., vol. 10, p. 11,

and Pal. N. Y., vol. 6, p. 287, Up. Held. Gr.

LOCULIPORA, Hall, 1887, Pal. N. Y., vol. 6, p. xxiii. [Ety. *loculus*, cell; *poros*, pore.] Fenestelloid; branches connected by dissepiments; cell apertures in two ranges, surrounding the fenestrules; branches and dissepiments carinated; carinae elevated and much thickened above, having the appearance of the branches and dissepiments of the noncelluliferous face of the frond. Type *L. perforata*.

ambigua, Hall, 1876, (*Fenestella ambigua*), 28th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Niagara Gr.

circumstata, Hall, 1887, Pal. N. Y., vol. 6, p. 144, Up. Held. Gr.

perforata, Hall, 1884, (*Fenestella perforata*), 36th Rep. N. Y. St. Mus. Nat. Hist., p. 65, Ham. Gr.

LYROPORA, Hall, 1857, Proc. Am. Ass. Ad. Sci., vol. 10, p. 179. [Ety. *lyra*, lute; *poros*, pore.] Zoarium consisting of a reticulated expansion, margined by two strong diverging supports which curve outward and upward; the rays of the expansion carry from two to five rows of cells; but there are none in the dissepiments; fenestrules small, ovate. Type *L. lyra*.

cinctura, Hall, 1885, Rep. St. Geol., pl. 1, Ham. Gr.

divergens, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 58, Kaskaskia Gr.

lyra, Hall, 1857, Proc. Am. Ass. Ad. Sci., vol. 10, p. 179, Kaskaskia Gr.

ovalis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 58, Kaskaskia Gr.

quincuncialis, Hall, 1857, Proc. Am. Ass. Ad. Sci., vol. 10, p. 179, Kaskaskia Gr.

ranosculum, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 58, Kaskaskia Gr.

retrorsa, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 504, Burlington Gr.

subquadrans, Hall, 1857, Proc. Am. Ass. Ad. Sci., vol. 10, p. 179, Kaskaskia Gr.

MEKOPORA, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, p. 383. [Ety. proper name.] Bifoliate, sometimes branching; the median laminae thin, flexuose; cells arranged with their oblique apertures directed toward the distal margin of the expansion; lunarium moderate or obsolete; zoecial tubes oblique, the anterior walls thinnest and flexuose; diaphragms numerous, often recurved; oecum a large oval cell, showing as a convex space with a small apical perforation. Type *M. eximia*.

(?) *aperta*, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 76, Keokuk Gr.

FIG. 493.—LYROPORA CINCTURA. Noncelluliferous side.

FIG. 494.—TA CLICULAR

CINCTURA

NEMAT.

74.

Ra

fro

pan

face

of c

the

pea

fibros

74.

simpl

193.

NEMAT.

Ill.

poro

uou

zooe

a ra

min

subc

arra

one

pres

alterna

Ill.

delica

Ill.

quadra

Ill.

retrors

vol.

NICHOLSON

Ill.

Irreg

bran

tubu

ture

gran

occu

poro

zooec

meso

regio

buli

with

the t

meso

diaph

cumula

vol. 8

287, Up.

, vol. 6,
1; *poros*,
nes con-
apertures
the fene-
ments car-
and much
appear-
sements
the frond.

(ambigua.)
Hist., p.

. Y., vol.

lla perfo-
Mus. Nat.

Ass. Ad.
lyra, lute;
sting of a
ed by two
ich curve
ays of the
five rows
n the dis-
all, ovate.

porpora cine-
cellulifer.

p. Sur. Ill.,

Proc. Am.
9, Kaskas-

Geo. Sur.
Gr.

1868, Geo.
ington Gr.
2. Am. Ass.

Kaskasia Gr.
Geo. Sur.
proper name.]
ching; the
ous; cells
e apertures
margin of
moderate or
que, the an-
xuous; dia-
recurved;
owing as a
ypical perfo-

Geo. Sur.
Gr.

approximata, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, pl. 77, Kaskaskia Gr.

FIG. 494.—Meekopo-
ra clausa. Aper-
tural cover $\times 50$.

clausa, Ulrich, 1884,
(Fistulipora? clausa.)
Jour. Cin. Soc. Nat.
Hist., vol. 7, p. 47,
Kaskaskia Gr.
eximia, Ulrich, (in
press.) Geo. Sur. Ill.,
vol. 8, pl. 77, Kaskas-
kia Gr.

Mitoclema, Ulrich, 1882,
Jour. Cin. Soc. Nat.
Hist., vol. 5, p. 150.
Syn. for Enallopora.

cinctosa, see Enallopora cinctosa.

NEMATAXIS, Hall, 1887, Pal. N. Y., vol. 6, p.
74. [Ety. *nema*, thread; *axon*, axis.]
Ramoso, solid, bifurcating, cells arising
from a filiform axis, apertures oval, in
parallel rows, separated by ridges; sur-
face marked with monticules, destitute
of cell apertures, and extending across
the branch, give it an annulated ap-
pearance. Type N. fibrosus.
fibrosus, Hall, 1887, Pal. N. Y., vol. 6, p.
74, Up. Held. Gr.
simplex, Hall, 1887, Pal. N. Y., vol. 6, p.
193, Ham. Gr.

NEMATOPORA, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, p. 401. [Ety. *nema*, thread;
poros, pore.] Slender, ramoso, contin-
uous above the pointed basal extremity;
zoecia subtubular, short, arranged in
a radial manner around one or two
minute axial tubes; apertures ovate or
subcircular, with peristome, generally
arranged between longitudinal ridges;
one or two diaphragms occasionally
present. Type N. quadrata.

alternata, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, pl. 29, Galena Gr.

delicatula, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, pl. 29, Galena Gr.

quadrata, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, pl. 29, Trenton Gr.

retorsa, Ulrich, (in press.) Geo. Sur. Ill.,
vol. 8, pl. 29, Galena Gr.

NICHOLSONELLA, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, p. 374. [Ety. proper name.]
Irregularly intertwining, flattened
branches, sometimes laminated; zoecia
tubular, with diaphragms in the "ma-
ture" region; apertures circular, with a
granose peristome; interspaces wide,
occupied by numerous angular meso-
pores, that more or less isolate the
zoecia; walls of both the zoecia and
mesopores thin, and in the mature
region traversed longitudinally by tu-
buli; the interzoecial spaces are filled
with a calcareous deposit, into which
the tubuli continue, but in which the
mesopore walls become unrecognizable;
mesopores with thick and numerous
diaphragms. Type N. ponderosa.

cumulata, Ulrich, (in press.) Geo. Sur. Ill.,
vol. 8, pl. 33, Hud. Riv. Gr.

ponderosa, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, pl. 34, Trenton Gr.

ODONTOTRYPA, Hall, 1887, Pal. N. Y., vol.
6, p. xvii. [Ety. *odontos*, tooth; *trypa*,
opening.] Distinguished from Lichen-
alia, by the oblique trilobate, closely
arranged cell apertures, with strongly
elevated, denticulated margins, forming
a crescentic projection over the aper-
ture. Type O. alveata.

alveata, Hall, 1881, (Lichenalia alveata.)
Trans. Alb. Inst., vol. 10, p. 10, and Pal.
N. Y., vol. 6, p. 85, Up. Held. Gr.

ORTHOPORA, Hall, 1887, Pal. N. Y., vol. 6,
p. xiv. [Ety. *orthos*, straight; *poros*,
pore.] Zoarium ramoso, solid; cell
apertures arranged in parallel, longitu-
dinal rows; intercellular space solid, or
occupied near the surface by minute
tubuli; no septa. Type O. regularis.

bispinulata, Hall, 1884, (Callopora bispin-
ulata,) Rep. St. Geol., p. 14, Ham. Gr.
ornata, Hall, 1887, Pal. N. Y., vol. 6, p.
184, Ham. Gr.

regularis, Hall, 1874, (Trematopora regu-
laris,) 26th Rep. N. Y. St. Mus. Nat.
Hist., p. 105, Up. Held. Gr.

reticulata, Hall, 1887, Pal. N. Y., vol. 6,
p. 179, Ham. Gr.

rhombifera, Hall, 1874, (Trematopora
rhombifera,) 26th Rep. N. Y. St. Mus.
Nat. Hist., p. 106, Up. Held. Gr.

scutulata, Hall, 1881, (Trematopora scutu-
lata,) Trans. Alb. Inst., vol. 10, p. 6,
and Pal. N. Y., vol. 6, p. 70, Up. Held. Gr.

PACHYDICTYA, Ulrich, 1882, Jour. Cin. Soc.
Nat. Hist., vol. 5, p. 152. [Ety. *pachys*,
thick; *dictyon*, net.] Zoarium com-
posed of large, thick, branching fronds;
cells ovate, separated by interstitial
tubes; diaphragms in both sets of
tubes; median epithelial plates perfo-
rated by minute foramina. Type P.
robusta.

conciliatrix, Ulrich, 1886, 14th Rep. Geo.
Sur. Minn., p. 76, Trenton Gr.

everetti, Ulrich, (in press.) Geo. Sur. Ill.,
vol. 8, pl. 33, Trenton Gr.

fimbriata, Ulrich, 1886, 14th Rep. Geo.
Sur. Minn., p. 75, Trenton Gr.

firma, Ulrich, (in press.) Geo. Sur. Ill.,
vol. 8, pl. 31, Hud. Riv. Gr.

foliata, Ulrich, 1886, 14th Rep. Geo. Sur.
Minn., p. 73, Trenton Gr.

gigantea, Ulrich, (in press.) Geo. Sur. Ill.,
vol. 8, pl. 31, Hud. Riv. Gr.

occidentalis, Ulrich, 1886, 14th Rep. Geo.
Sur. Minn., p. 75, Trenton Gr.

robusta, Ulrich, 1882, Jour. Cin. Soc. Nat.
Hist., vol. 5, p. 173, Trenton Gr.

splendens, Ulrich, (in press.) Geo. Sur.
Ill., vol. 8, pl. 31 and 32, Hud. Riv. Gr.

PALESCHARA, Hall, 1874, 26th Rep. N. Y. St.
Mus. Nat. Hist., p. 107. [Sig. ancient
Eschara.] Incrusting expansions; cells
polygonal, oblong, separated by thin
solid walls; maculae present; no spini-
form tubuli or diaphragms. Type P.
incrustans.

amplectans, Hall, 1884, Rep. St. Geol., p. 7, Ham. Gr.

aspera, Hall, 1876, syn. for *P. maculata*.

bifoliata, syn. for *Ptilodictya nebulosa*.

bilateralis, Hall, 1883, Rep. St. Geol., pl. 16, fig. 22-25, Low. Held. Gr.

concentrica, Hall, 1887, Pal. N. Y., vol. 6, p. 67, Low. Held. Gr.

foliata, syn. for *Ptilodictya nebulosa*.

incrassata, Hall, 1879, 28th Rep. N. Y.

St. Mus. Nat. Hist., p. 121, Niagara Gr.

incrustans, Hall, 1874, 26th Rep. N. Y. St.

Mus. Nat. Hist., p. 107, Low. Held. Gr.

intercella, Hall, 1884, Rep. St. Geol., p. 5,

Ham. Gr.

maculata, Hall, 1876, 28th Rep. N. Y. St.

Mus. Nat. Hist., p. 121, Niagara Gr.

offula, Hall, 1876, 28th

Rep. N. Y. St. Mus. Nat.

Hist., p. 120, Niagara Gr.

pertenuis, Hall, 1884, Rep.

St. Geol., p. 7, Ham. Gr.

radiata, Hall, 1883, Rep.

St. Geol., p. 16, fig. 13-14,

Low. Held. Gr.

reticulata, Hall, 1884, Rep. St. Geol., p. 6,

Ham. Gr.

sphaerion, Hall, 1876, 28th Rep. N. Y. St.

Mus. Nat. Hist., p. 121, Niagara Gr.

tenuis, Hall, 1887, Pal. N. Y., vol. 6, p.

36, Low. Held. Gr.

variocella, Hall, 1884, Rep. St. Geol., p. 6,

Ham. Gr.

PETALOTRYPA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 377. [Ety. *petalos*, spread out; *trupa*, an opening.] Bifoliate, consisting of irregular, compressed branches or simple fronds, celluliferous on both sides; zoecial tubes prismatic, arising from a strongly flexuous mesial line; apertures subcircular or polygonal; mesopore-like interspaces, that do not differ in their tabulation from the zoecia, may occur; very small acanthopores (?) occupy many of the angles of junction. Type *P. compressa*.

compressa, Ulrich,

(in press,) Geo.

Sur. Ill., vol. 8, pl.

46, Ham. Gr.

delicata, Ulrich, (in

press,) Geo. Sur.

Ill., vol. 8, pl. 46,

Ham. Gr.

PETIGOPORA, Ulrich,

1882, Journal Cin.

Soc. Nat. Hist.,

vol. 5, p. 155.

[Ety. *petigo*, scab;

poros, pore.] Small

patches adhering to foreign objects, with a narrow nonporiferous band or germinating membrane along the outer margin; no interstitial cells; spiniform tubuli present. Type *P. gregaria*.

asperula, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 157, Hud. Riv. Gr.

FIG. 495.—Paleosephara offula.

gregaria, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 155, Hud. Riv. Gr.

petechialis, Nicholson, 1875. (Chetetes petechialis,) Ohio Pal., vol. 2, p. 213,

Hud. Riv. Gr.

FIG. 496.—Petigopora petechialis on a Monticulipora; also, specimen greatly enlarged.

PHACLOPORA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 368. [Ety. *phakelos*, bundle;

poros, pore.] Zoarium articulated; seg-

ments short, obconical, consisting of

two or more equal, conical zoecia, with

slightly contracted circular apertures.

Type *P. pertenuis*.

constricta, Ulrich, (in press,) Geo. Sur.

Ill., vol. 8, pl. 29, Trenton Gr.

pertenuis, Ulrich, (in press,) Geo. Sur.

Ill., vol. 8, pl. 29, Galena Gr.

PHÆNOPORA, Hall, 1852, Pal. N. Y., vol. 2,

p. 46. [Ety. *phaino*, to open or make a

window; *poros*, pore.] Zoarium form-

ing thin, broad, or ensiform expansions,

celluliferous on both sides; cellules

oval and arranged between elevated

lines; maculae often developed; distin-

guished from *Ptilodictya* and *Stictopora*

by the absence of a nonporiferous, stri-

ated edge. Type *P. explanata*.

FIG. 497.—Phænopora constellata. Sections x 50.

constellata, Hall, 1852, Pal. N. Y., vol. 2, p. 47, Clinton Gr.

ensiformis, Hall, 1852, Pal. N. Y., vol. 2, p. 48, Clinton Gr.

excellens, Billings, 1866, (*Ptilodictya excellens*,) Cat. Sil. Foss. Antic., p. 34,

Anticosti Gr.

expansa, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 114, Niagara Gr.

FIG. 498.—Phylloporina expansa.

explanata, Hall, 1852, Pal. N. Y., vol. 2, p. 46, Clinton Gr.

multiplora, Hall, 1851, Geo. Lake Supp. Land Dist., vol. 2, p. 206, Trenton Gr.

tenuis, Hall, 1874, (Escharopora tenuis,) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 99, Low. Held. Gr.

PHYLLOPORINA, Hall, 1881, Trans. Alb. Inst., vol. 10, p. 12. [Ety. phylloporos, inclosed; poros, pore.] Zoarium explanate, free or incrusting, frequently contorted, celluliferous on one or both faces; surface elevated at irregular intervals into prominent crests; cells tubular, without septa; intercellular structure vesiculate near the base, septate above. Type P. cristata.

cristata, Hall, 1881, Trans. Alb. Inst., vol. 10, p. 12, and Pal. N. Y., vol. 6, p. 99, Up. Held. Gr.

cristata var. lineata, Hall, 1887, Pal. N. Y., vol. 6, p. 99, Up. Held. Gr.

PHYLLODICTYA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 153. [Ety. phyllon, leaf; dictyon, net.] Zoarium forming simple, leaf-like expansions, sometimes branched; cell apertures small, oblique, with the lower margin lipped; interstitial spaces minutely granular or punctate. Type P. frondosa.

FIG. 499.—Phyllodictya frondosa.

frondosa, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 174, Trenton Gr.

PHYLLOPORINA, King, 1849, Ann. and Mag. Nat. Hist., 2d ser., vol. 3, p. 389. [Ety. phyllon, leaf; poros, perforation.] Zoarium like Fenestella, but having cellules on the whole of the under surface of the rays in two or more ranges. Type P. ehrenbergi.

aspera, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 46, Up. Held. Gr.

corticosa, see Subretepora corticosa, ehrenbergi, Geinitz, 1846, (Gorgonia ehrenbergi,) Grundriss, p. 585, Permian Gr. Very doubtfully identified in America.

superba, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 46 and 55, Ham. Gr. variolata, see Subretepora variolata.

Phylloporina, Ulrich, syn. for Subretepora.

PILEOTRYPA, Hall, 1887, Pal. N. Y., vol. 6, p. xvi. [Ety. pileos, cap; trupa, opening.] Distinguished from Lichenalia by having the posterior portions of the peristomes strongly elevated and arched, with distinct denticulations in the aperture, which, in the course of growth, form two longitudinal striations along the interior of the cell wall. Type P. denticulata.

clivulata, Hall, 1881, (Lichenalia clivulata,) Trans. Alb. Inst., vol. 10, p. 9, and Pal. N. Y., vol. 6, p. 83, Up. Held. Gr. denticulata, Hall, 1881, (Lichenalia denticulata,) Trans. Alb. Inst., vol. 10, p. 8, and Pal. N. Y., vol. 6, p. 84, Up. Held. Gr.

granifera, Hall, 1881, (Lichenalia granifera,) Trans. Alb. Inst., vol. 10, p. 11, and Pal. N. Y., vol. 6, p. 84, Up. Held. Gr.

pyriformis, Hall, 1881, (Lichenalia pyriformis,) Trans. Alb. Inst., vol. 10, p. 12, and Pal. N. Y., vol. 6, p. 82, Up. Held. Gr.

PINACOTRYPA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 384. [Ety. pinax, plank; trupa, opening.] Thin, contorted expansions, with a wrinkled epitheca below; zoecia with subcircular apertures, a well-developed granose peristome, thin walls, and, so far as observed, no lunarium; interspaces wide, occupied by a single series of very large angular mesopores, which never present the appearance of vesicular tissue; diaphragma horizontal, few in the zoecial tubes, numerous in the mesopores. Type P. elegans.

elegans, Rominger, 1866, (Fistulipora elegans,) Proc. Acad. Nat. Sci. Phil., p. 9, Ham. Gr.

Pinnatopora, syn. for Glaucanome.

curvata, see Glaucanome curvata.

intermedia, see Glaucanome intermedia.

minor, see Glaucanome minor.

simulatrix, see Glaucanome simulatrix.

subangulata, see Glaucanome subangulata.

tenuiramosa, see Glaucanome tenuiramosa.

vinei, see Glaucanome vinei.

whitii, see Glaucanome whitii.

youngi, see Glaucanome youngi.

POLYPORE, McCoy, 1845, Carb. Foss. Ireland, p. 206. [Ety. polys, many; poros, pore.] Zoarium like that of Fenestella, from which it is distinguished by having no median ridge on the celluliferous side of the rays, and in having from three to ten rows of cell openings. Type P. dendroides.

aculeata, Hall, 1881, (Fenestella aculeata,) Trans. Alb. Inst., vol. 10, p. 21, and Pal. N. Y., vol. 6, p. 157, Up. Held. Gr. adnata, Hall, 1881, (Fenestella adnata,) Trans. Alb. Inst., vol. 10, p. 25, and Pal. N. Y., vol. 6, p. 152, Up. Held. Gr.

albionensis, Spencer, 1884, Bull. No. 1, Univ. St. Mo., p. 55, Niagara Gr.

approximata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 61, Kaskaskia Gr.

arkonensis, S. A. Miller, 1883, 2d ed. Am. Pal. Foss., p. 202, Ham. Gr. Proposed instead of *P. tuberculata*, Nicholson, in Geo. Mag. for April, 1874, and Rep. Pal. Prov. Ont., p. 100, figs. 37, a, b, c. Found at Arkona, township of Bosanquet, Canada.

arta, Hall, 1879, (Fenestella arta,) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 163, Low. Held. Gr.

biarmica, Keyserling, 1846, Geognost. Beobacht., p. 191. Geinitz referred a form from the Coal Meas., and Prout referred one from the Kaskaskia Gr. to it. Probably not an American species.

FIG. 500.—*Polypora biserialis*. Aperture having the central perforation of the cover closed, x 50.

biseriata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 60, Warsaw and St. Louis Gr.

blandida, Ulrich, 1880, Contr. to Amer. Pal., p. 18, Up. Held. Gr.

brevisulcata, Hall, 1881, (Fenestella brevisulcata,) Trans. Alb. Inst., vol. 10, p. 26, and Pal.

N. Y., vol. 6, p. 168, Up. Held. Gr.
burlingtonensis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 59, Burlington Gr.
carinella, Hall, 1887, Pal. N. Y., vol. 6, p. 153, Up. Held. Gr.

celsipora, Hall, 1881, (Fenestella celsipora,) Trans. Alb. Inst., vol. 10, p. 24, and Pal. N. Y., vol. 6, p. 150, Up. Held. Gr.

celsipora var. minima, Hall, 1881, (Fenestella celsipora var. minima,) Trans. Alb. Inst., vol. 10, p. 24, and Pal. N. Y., vol. 6, p. 151, Up. Held. Gr.

celsipora var. minor, Hall, 1881, (Fenestella celsipora var. minor,) Trans. Alb. Inst., vol. 10, p. 24, and Pal. N. Y., vol. 6, p. 151, Up. Held. Gr.

cestriensis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 60, Kaskaskia Gr.

compacta, Hall, 1879, (Fenestella compacta,) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 163, Low. Held. Gr.

complanata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 60, Kaskaskia Gr.

compressa, Hall, 1879, (Fenestella compressa,) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 164, Low. Held. Gr.

corticosa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 61, Kaskaskia Gr.

crebescens, Hall, 1887, Pal. N. Y., vol. 6, p. 170, Up. Held. Gr.

crassa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 61, Up. Coal Meas.

cultellata, Hall, 1881, (Fenestella cultellata,) Trans. Alb. Inst., vol. 10, p. 21, and Pal. N. Y., vol. 6, p. 160, Up. Held. Gr.

distans, Hall, 1881, (Fenestella distans,) Trans. Alb. Inst., vol. 10, p. 24, and Pal.

N. Y., vol. 6, p. 161, Up. Held. Gr.

elegans, Hall, 1874, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 97, Low. Held. Gr.

elongata, Hall, 1882, Rep. St. Geol. and Pal. N. Y., vol. 6, p. 153, Up. Held. Gr.

eudora, Hall, 1887, Pal. N. Y., vol. 6, p. 58, Low. Held. Gr.

fistulata, Hall, 1884, (Fenestella fistulata,) 36th Rep. N. Y. St. Mus. Nat. Hist., p. 59, Ham. Gr.

flabelliformis, Hall, 1881, (Fenestella flabelliformis,) Trans. Alb. Inst., vol. 10, p. 23, and Pal. N. Y., vol. 6, p. 161, Up. Held. Gr.

gracilis, Prout, 1860, Trans. St. Louis Acad. Sci., p. 580, Warsaw Gr.

grandis, Toula, 1875, N. Jahrbuch, p. 230, Carboniferous.

granilinea, Hall, 1881, (Fenestella granilinea,) Trans. Alb. Inst., vol. 10, p. 27, and Pal. N. Y., vol. 6, p. 154, Up. Held. Gr.

hallana, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 580, Warsaw Gr.

hamiltonensis, Prout, 1860, Geo. Sur. Ill., vol. 2, p. 423, Ham. Gr.

hexagonalis, Hall, 1881, (Fenestella hexagonalis,) Trans. Alb. Inst., vol. 10, p. 27, and Pal. N. Y., vol. 6, p. 164, Up. Held. Gr.

hexagonalis var. foraminulosa, Hall, 1881, (Fenestella hexagonalis var. foraminulosa,) Trans. Alb. Inst., vol. 10, p. 27, and Pal. N. Y., vol. 6, p. 165, Up. Held. Gr.

idothea, Hall, 1879, (Fenestella idothea,) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 97, Low. Held. Gr.

imbricata, Prout, 1860, Trans. St. Louis Acad. Sci., vol. 2, p. 412, Devonian.

impressa, Ulrich, 1888, Bull. Denison Univ., vol. 4, p. 72, Cuyahoga Shale.

incepta, Hall, 1882, Pal. N. Y., vol. 2, p. 167, Niagara Gr.

intermedia, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 272, Up. Held. Gr.

levinodata, Hall, 1881, (Fenestella levinodata,) Trans. Alb. Inst., vol. 10, p. 28, and Pal. N. Y., vol. 6, p. 169, Up. Held. Gr.

lavistriata, Hall, 1883, Rep. St. Geol. and Pal. N. Y., vol. 6, p. 159, Up. Held. Gr.

largissima, Hall, 1881, (Fenestella largissima,) Trans. Alb. Inst., vol. 10, p. 22, and Pal. N. Y., vol. 6, p. 166, Up. Held. Gr.

libea, Hall, 1874, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 62, Low. Held. Gr.

maccoyana, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 59, Keokuk Gr.

megastoma, DeKoninck, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 5, Carboniferous.

mexicana, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 270, Permian Gr.

mutabilis, Hall, 1881, (Fenestella mutabilis,) Trans. Alb. Inst., vol. 10, p. 25, and Pal. N. Y., vol. 6, p. 166, Up. Held. Gr.

nexa, Hall, 1881, (Fenestella nexa,) Trans. Alb. Inst., vol. 10, p. 25, and Pal. N. Y., vol. 6, p. 165, Up. Held. Gr.

FIG. 501.—*Polypora* shum tared.

(Fenestella Inst. 6, p.

- olocarinata*, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 61, Coal Meas.
obliqua, Hall, 1887, Pal. N. Y., vol. 6, p. 64, Low. Held. Gr.
papillata, McCoy, 1862, Carb. Foss. of Ireland, p. 226, Up. Coal Meas.
paxillata, Hall, 1879, (Fenestella paxillata,) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 164, Low. Held. Gr.
perangulata, Hall, 1881, (Fenestella perangulata,) Trans. Alb. Inst., vol. 10, p. 23, and Pal. N. Y., vol. 6, p. 162, Up. Held. Gr.
perundata, Hall, 1881, (Fenestella perundata,) Trans. Alb. Inst., vol. 10, p. 27, and Pal. N. Y., vol. 6, p. 163, Up. Held. Gr.
porosa, Hall, 1881, (Fenestella porosa,) Trans. Alb. Inst., vol. 10, p. 26, and Pal. N. Y., vol. 6, p. 163, Up. Held. Gr.
propria, Hall, 1881, (Fenestella propria,) Trans. Alb. Inst., vol. 10, p. 22, and Pal. N. Y., vol. 6, p. 157, Up. Held. Gr.
(?) psyche, Billings, 1874, Pal. Foss., vol. 2, p. 11, Gaspe limestone No. 8, Devonian.
pulchella, Nicholson, 1874, Geo. Mag. Lond. n. s., vol. 1, p. 161, Corniferous Gr.
quadrangularis, Hall, 1881, (Fenestella quadrangularis,) Trans. Alb. Inst., vol. 10, p. 21, and Pal. N. Y., vol. 6, p. 158, Up. Held. Gr.
radialis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 60, Keokuk Gr.
retrorsa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 59, Keokuk Gr.
rigida, Prout, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 412, Up. Held. Gr.
rigida, Hall, 1881, (Fenestella rigida,) Trans. Alb. Inst., vol. 10, p. 22, and Pal. N. Y., vol. 6, p. 155, Up. Held. Gr.
robusta, Hall, 1881, (Fenestella robusta,) Trans. Alb. Inst., vol. 10, p. 22, and Pal. N. Y., vol. 6, p. 156, Up. Held. Gr.
rustica, Hall, 1887, Pal. N. Y., vol. 6, p. 169, Up. Held. Gr.
separata, Hall, 1882, Rep. St. Geol. and Pal. N. Y., vol. 6, p. 166, Up. Held. Gr.
shumardi, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 271, Up. Held. Gr.
simulatrix, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 59, Keokuk Gr.
spinulifera, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 61, Kaaskaskia Gr. and Coal Meas.
stragula, White, 1874, Rep. Invert. Foss., p. 19, and Geo. Sur. W. 100th Mer. vol. 4, p. 108, Coal Meas.
striatopora, Hall, 1881, (Fenestella striatopora,) Trans. Alb. Inst., vol. 10, p. 23, and Pal. N. Y., vol. 6, p. 168, Up. Held. Gr.

FIG. 501.—*Polypora shumardi*. Enlarged.

- stricta*, Hall, 1887, Pal. N. Y., vol. 6, p. 59, Low. Held. Gr.
submarginata, Meek, 1872, Pal. E. Neb., p. 154, Coal Meas.
submutans, Hall, 1881, (Fenestella submutans,) Trans. Alb. Inst., vol. 10, p. 21, and Pal. N. Y., vol. 6, p. 167, Up. Held. Gr.
tenella, Nicholson, 1874, Geo. Mag. Lond. n. s., vol. 1, p. 162, Corniferous Gr.
transversa, Ulrich, 1880, Cont. to Am. Pal., p. 18, Up. Held. Gr.
tuberculata, Prout, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 449, Kaskaskia Gr.
tuberculata, Nicholson, see *P. arkonensis*.
varsoviensis, Prout, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 237, Warsaw Gr.
varsoviensis var. *spininodata*, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 60, Warsaw Gr.
whitii, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 62, Coal Meas.
whitii var. *eximia*, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 62, Coal Meas.
PAMMOPORA, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 17. [Ety. *primos*, the hole made by a cylindrical saw; *poros*, pore.] Ramose, branches triangular, dichotomous, each side celluliferous; tubes radiate from the center to each angle, margins noncelluliferous; intersittal spaces smooth, vesicular. Type *P. triquetra*.
dilatata, Hall, 1884, Rep. St. Geol., p. 50, Ham. Gr.
lata, Hall, 1887, Pal. N. Y., vol. 6, p. 266, Ham. Gr.
minima, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 78, Coal Meas.
paucirama, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 17, and Rep. St. Geol., 1883, pl. 25, fig. 11, Up. Held. Gr.
serrata, Meek, 1875, (Ptilodictya serrata,) Pal. Ohio, vol. 2, p. 327, Coal Meas.
serrulata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 41, Kaskaskia Gr. Perhaps the same as *P. serrata*.
sparsipora, Hall, 1881, (Thallostigma sparsipora,) Trans. Alb. Inst., vol. 10, p. 13, and Pal. N. Y., vol. 6, p. 288, Up. Held. Gr.
triquetra, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 17, and Rep. St. Geol., 1883, pl. 25, fig. 8-10, Up. Held. Gr.
PROTOCRISINA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 369. [Ety. *protos*, first; *Crisina*, a genus.] Ramoso, celluliferous on one side only; cells subcylindrical, with prominent circular apertures; reverse finely grano-striate; small pores, apparently communicating with the interior of the zoecia, are rather irregularly distributed over both sides of the branches; axis thin, cruciform in transverse section; external walls thick. Type *P. exigua*.
exigua, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 29 and 53, Trenton and Hud. Riv. Gr.

PROUTELLA, Ulrich, (in press,) *Geo. Sur. Ill.*, vol. 8, p. 403. [Ety. proper name.] Discoid, thin, free, lower surface convex and lined with a concentrically wrinkled epitheca; primary zoea sub-tubular, the succeeding ones shorter, all rather thin-walled; aperture broad-elliptical, surrounded by a narrow, sloping area, hexagonal in outline; when perfect, with a depressed delicate calcareous plate, that closes a little less than two-thirds of the opening, the orifice left being subtriangular in form, without thickened margins, and situated at the anterior side; with age, a second, third, and more layers of zoea are developed directly over the first, so that they gradually form a zoecial tube seemingly having the cavity intersected by incomplete diaphragms; these appear to spring from the posterior wall, and extend about one-half the distance across. Type *Cyclopore discoidea*. Syn.?

discoidea, Prout, 1860, (*Cyclopore discoidea*), *Trans. St. Louis Acad. Sci.*, vol. 1, p. 575, Keokuk Gr. *Pteropora duogeneria*, Hall, syn. for *Tenipora exigua*.

Ptilonella, Hall, 1884, *Rep. St. Geol.*, p. 56, syn. for *Reptaria*. *nodata*, see *Reptaria nodata*. *penniformis*, see *Reptaria penniformis*.

Ptilodictya, Lonsdale, 1839, *Murch. Sil. Syst.*, p. 676. [Ety. *ptilon*, feather; *dictyon*, net.] The correct orthography is *Ptilodictyon*. Zoarium pointed below, articulating into a spreading base, above a leaf-like expansion, which is sometimes lobed at the distal extremity, celluliferous on both faces, divided by a mesial lamina; margin without cells; apertures circular or subquadrate; no intercellular tissue. Type *P. lanceolata*.

FIG. 502.—*Ptilodictya maculata*. Vertical section x 85, showing spinous process.

acuminata, James, 1876, *Int. Catal. Cin. Foss.*, p. 3, Hud. Riv. Gr. Not well defined.

alcyone, see *Pachydictya alcyone*. *arctipora*, see *Bythopora arctipora*.

arguta, see *Stictopora arguta*. *bipunctata*, Van Cleave, 1883, 12th Rep. Ind. Geol. and Nat. Hist., p. 206, Niagara Gr.

briareus, Ulrich, 1882, *Jour. Cin. Soc. Nat. Hist.*, vol. 5, p. 164, Trenton Gr. *canadensis*, Billings, 1866, *Catal. Sil. Foss. Antic.*, p. 9, Hud. Riv. Gr.

carbonaria, see *Stictopora carbonaria*. *cosciniformis*, see *Coscinella cosciniformis*. *dictyota*, Meek, 1873, Hayden's 6th Rep. Geo. Sur. Terr., p. 465, Subcarboniferous.

emacerrata, see *Dicranopora emacerrata*. *excellens*, see *Phenopora excellens*. *explicans*, Safford, 1869, *Geo. of Tenn.* Not defined.

falciformis, Nicholson, 1875, Ohio Pal., vol. 2, p. 259, Hud. Riv. Gr. *fenestelliformis*, Nicholson, 1875, Ohio Pal., vol. 2, p. 263, Hud. Riv. Gr.

FIG. 503.—*Ptilodictya magnifica*. Sections x 50.

flagellum, Nicholson, 1875, Ohio Pal., vol. 2, p. 262, Hud. Riv. Gr.

fragilis, see *Dicranopora fragilis*. *gladiola*, Billings, 1866, *Catal. Sil. Foss. Antic.*, p. 10, Anticosti Gr.

hilli, James, 1882, (as figured by Ulrich), *Jour. Cin. Soc. Nat. Hist.*, vol. 5, pl. 7, Trenton Gr.

internodia, see *Dicranopora internodia*. *libana*, Safford, 1869, *Geo. of Tenn.* p. 286, Trenton Gr.

lirata, Hall, 1874, (*Escharopora lirata*), 26th Rep. N. Y. St. Mus. Nat. Hist., p. 100, Low. Held. Gr.

maculata, Ulrich, 1882, *Jour. Cin. Soc. Nat. Hist.*, vol. 5, p. 163, Hud. Riv. Gr. *magnifica*, S. A. Miller, 1878, *Jour. Cin. Soc. Nat. Hist.*, vol. 1, p. 100, Hud. Riv. Gr.

meeki, Nicholson, 1874, *Geo. Mag. n. s.*, vol. 1, p. 123, Corniferous and Ham. Gr. *multiramis*, Safford. Not defined.

nebulosa, Hall, 1874, (*Escharopora nebulosa*), 26th Rep. N. Y. St. Mus. Nat. Hist., p. 99, Low. Held. Gr.

nitidula, see *Dicranopora nitidula*. *nodosa*, James, 1882, (as figured by Ulrich), *Jour. Cin. Soc. Nat. Hist.*, vol. 5, pl. 7, Hud. Riv. Gr. The name was preoccupied. See *P. variabilis*.

PTIL.
obl.
(
par
2
FIG.
pav
E
per
plu
r
p
plu
2
pun
J
ram
H
ret
p
rue
scu
le
serr
sub
S
sul
A
sup
A
sym
tar
12
FIG. 5
var
verse
show
port
layer
and
char
med
Betw
ther
med
In
pa
ro

12th Rep.
p. 200, Ni-

a. Soc. Nat.
Gr.

i. Sil. Foss.

onaria.
einiformis,
s 6th Rep.
boniferous,
acerata.
ons.
Tenn. Not

io Pal., vol.

1875, Ohio
Gr.

ous x 50.

io Pal., vol.

is.

. Sil. Foss.,

by Ulrich,)

vol. 5, pl. 7,

ernodia.

Tenn. p. 286,

ora lirata,)

at. Hist., p.

r Cin. Soc.

nd. Riv. Gr.

our Cin. Soc.

nd. Riv. Gr.

Mag. n. s.,

nd Ham. Gr.

ed.

opora nebu-

Mus. Nat.

ula.

red by Ul-

Hist., vol. 5,

e name was

ilia.

obliqua, Ringueberg, 1884, (Stictopora
obliqua,) Proc. Acad. Nat. Sci., p. 140,
Clinton Gr. Not well defined.
parallela, Hall, 1887, Pal. N. Y., vol. 6, p.
270, Ham. Gr.

FIG. 504.—*Ptilodictya pavonia*. Natural size,
and magnified.

pavonia, D'Orbigny, 1850, Prodr. de
Paleont., t. 1, p. 22, Hud. Riv. Gr.
perlegans, see *Graptodictya perlegans*.
plumaria, James, 1882, (as figured by Ul-
rich,) Jour. Cin. Soc. Nat. Hist., vol. 5,
pl. 7, Hud. Riv. Gr.
plumea, Hall, 1887, Pal. N. Y., vol. 6, p.
271, Ham. Gr.
punctata, Nicholson & Hinde, 1874, Can.
Jour., p. 7, Clinton Gr.
ramosa, Ulrich, 1882, Jour. Cin. Soc. Nat.
Hist., vol. 5, p. 164, Trenton Gr.
retiformis, Hall, 1887, Pal. N. Y., vol. 6,
p. 272, Ham. Gr.
rustica, see *Stictopora rustica*.
scutulata, Hall, 1884, (Stictopora scutu-
lata,) Rep. St. Geol., p. 47, Ham. Gr.
serrata, see *Prismopora serrata*.
subrecta, Ulrich, 1886, 14th Rep. Geo.
Sur. Minn., p. 63, Trenton Gr.
sulcata, Billings, 1886, Catal. Sil. Foss.
Antic., p. 35, Anticoeti Gr.
superba, Billings, 1886, Catal. Sil. Foss.
Antic., p. 35, Anticoeti Gr.
symmetra, Safford. Not defined.
tarda, Billings, 1874, Pal. Foss., vol. 2, p.
13, Gaspé Limestone No. 8, Devonian.

FIG. 505.—*Ptilodictya*
variabilis. Trans-
verse section x 50,
showing the basal
portion of the two
layers of zoecia,
and the duplex
character of the
median lamina.
Between the plates
there is no series of
median tubuli.

Ireland, p. 200. [Ety. *ptilon*, plume;
poros, pore.] Flabelliform attached by
roots, from which a strong midrib

arises, giving origin on each side to
thin, equidistant rays, connected by
regular dissepiments; external face of
the rays carinate and bearing two rows
of pores. Type *P. flustriformis*.
acuta, Ulrich, (in press,) Geo. Sur. Ill.,
vol. 8, pl. 65, Burlington and Keo-
kuk Gr.
cylindracea, Ulrich, (in press,) Geo. Sur.
Ill., vol. 8, pl. 66, Keokuk Gr.
infrequens, Hall, 1887, Pal. N. Y., vol. 6,
p. 284, Ham. Gr.
nodosa, Hall, 1884, Rep. St. Geol., p. 59,
Ham. Gr.
paupera, Ulrich, (in press,) Geo. Sur. Ill.,
vol. 8, pl. 66, Keokuk Gr.
pronti, Hall, 1858, Geo. Rep. Iowa, p. 653,
Warsaw Gr.
atriata, Hall, 1884, Rep. St. Geol., p. 58,
Ham. Gr.
valida, Ulrich, (in press,) Geo. Sur. Ill.,
vol. 8, pl. 65 and 66, Keokuk Gr.

FIG. 506.—*Ptilodictya variabilis*. Vertical section
x 35, showing hemisepta.

PTILOPORELLA, Hall, 1887, Pal. N. Y., vol.
6, p. xxiv. [Ety. from the genus *Ptilo-*
pora.] Bryozoom growing in the
same manner as *Ptilopora*, but with
only two ranges of cell apertures on
the branches. Type *P. laticrescens*.
inaequalis, Hall, 1887, Pal. N. Y., vol. 6, p.
171, Up. Held. Gr.
laticrescens, Hall, 1887, Pal. N. Y., vol. 6,
p. 171, Up. Held. Gr.
nervata, Nicholson, 1875, (Fenestella
nervata,) Ohio Pal., vol. 2, p. 264, Nia-
gara Gr.
PTILOPORINA, Hall, 1887, Pal. N. Y., vol. 6,
p. xxiv. [Ety. from the genus *Ptilo-*
pora.] Resembling *Fenestella*, some
branches larger than others; ordinary
branches originate laterally from one
or both sides of the primary branches,
not bifurcating as in ordinary forms of
Fenestella; cell apertures in three or
more ranges. Type *P. conica*.
conica, Hall, 1887, Pal. N. Y., vol. 6, p.
172, Up. Held. Gr.
disparilis, Hall, 1887, Pal. N. Y., vol. 6,
p. 173, Up. Held. Gr.
pinnata, Hall, 1887, Pal. N. Y., vol. 6, p.
172, Up. Held. Gr.

sinistralis, Hall, 1887, Pal. N. Y., vol. 6, p. 174, Up. Held. Gr.

PTILOTRYPA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 393. [Ety. *ptilon*, feather; *trupa*, an opening.] Bifoliate, forming large ramose expansions. Zoecial tubes and apertures very oblique; at the upper extremity of the acutely ovate aperture there is a small cell which is best seen in tangential sections; surface with irregular, longitudinally channeled spots. Type *T. obliquata*.

obliquata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 30, Hud. Riv. Gr.

RAMPORA, Toulou, 1875, Permo-Carbon-Fossilien von der West Küste von Spitzbergen, p. 6. [Ety. *ramus*, branch; *poros*, pore.] Staff in cross section, rounded, rhombic, with keel on both sides; branches in pairs, one on each side, and these connected in like manner by rays, upward and downward; pores on one side, on each side of the keel. Type *R. hochstetteri*.

hochstetteri, Toulou, 1875, Permo-Carbon-Fossilien von der West Küste von Spitzbergen, p. 6, Carboniferous.

REPTARIA, Rolle, 1851, Leonhard & Bronn, Neues Jahrb., p. 810. [Ety. *repto*, to creep.] Zoarium parasitic, procumbent, attached its entire length; consisting of a rachis, from which proceed laterally, at regular intervals, cylindrical cell tubes, and at irregular distances tubes which have the same manner of growth as the primary rachis; cell-tubes turn abruptly outward at their distal extremities, and open in an aperture parallel with the axis of the branch. Type *R. stolonifera*.

nodata, Hall, 1884, (*Ptilionella nodata*), Rep. St. Geol., p. 57, Ham. Gr.

penniformis, Hall, 1884, (*Ptilionella penniformis*), Rep. St. Geol., p. 56, Ham. Gr.

stolonifera, Rolle, 1851, Leonhard & Bronn, Neues Jahrb., p. 810, Ham. Gr.

Retepora, Lamarck, 1801, Syst. An. sans. Vert. [Ety. *rete*, net; *poros*, pore.] Not a Palæozoic genus.

angulata, see *Subretepora angulata*.

antiqua, as identified by d'Archiac & Verneuil. Not American.

archimedes, see *Archimedes*.

asperato-striata, see *Subretepora asperato-striata*.

clintoni, Vanuxem, 1842, Geo. Rep. 3d Dist. N. Y. Not recognized.

diffusa, see *Thamniscus diffusus*.

fenestrata, see *Subretepora fenestrata*.

foliacea, Hall, 1847. This name Prof.

Hall says may be erased from the list.

gracilis, see *Subretepora gracilis*.

hamiltonensis, see *Reteporina hamiltonensis*.

incepta, see *Subretepora incepta*.

phillipsi, see *Reteporina phillipsi*.

prisca, see *Reteporina prisca*.

trentonensis, see *Subretepora trentonensis*.

RETEPORINA, D'Orbigny, 1850, Prodr. J. Paléont., t. 1, p. 101. [Ety. from *Retepora*.]

Resembling *Retepora*, but having on the greater part of the branches only two ranges of cell apertures; branches connected by anastomosis or by dissepiments so short as to be essentially wanting. Type *R. prisca*.

coalescens, Hall, 1887, Pal. N. Y., vol. 6, p. 120, Up. Held. Gr.

hamiltonensis, Prout, 1866, (*Retepora hamiltonensis*), Trans. St. Louis, Acad. Sci., vol. 2, p. 412, Ham. Gr.

perundulata, Hall, 1884, (*Fenestella perundulata*), 36th Rep. N. Y. St. Mus.

Nat. Hist., p. 63, Ham. Gr.

phillipsi, Nicholson, 1874, (*Retepora phillipsi*), Geo. Mag. n. s., vol. 1, p. 163, Corniferous Gr.

prisca, (Goldfuss, 1831, (*Retepora prisca*), Petref. Germ., vol. 1, p. 103, Ham. Gr.

rhombifera, Hall, 1881, (*Fenestella rhombifera*), Trans. Alb. Inst., vol. 10, p. 32, and Pal. N. Y., vol. 6, p. 120, Up.

Held. Gr.

striata, Hall, 1884, (*Fenestella striata*), 36th Rep. N. Y. St. Mus. Nat. Hist., p. 72, Ham. Gr.

RHINIDICTYA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 152. [Ety. *rhine*, file; *dictyon*, net.] Zoarium narrow,

branching at long intervals; cells surrounded by a close series of small spiniform tubuli; otherwise like *Stictopora*. Type *R. nicholsoni*. Syn. for *Sulcopora* probably.

FIG. 507.—*Rhinidictya nicholsoni*. Natural size and magnified 18 diam.

granulosa, Hall, 1887, Pal. N. Y., vol. vi, p. 40, Low. Held. Gr.

FIG. 508.—*Rhinopora verrucosa*. Natural size and enlarged.

nicholsoni, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 170, Trenton Gr.

RHINOPORA, Hall, 1852, Pal. N. Y., vol. 2, p. 48. [Ety. *rhine*, file; *poros*, pore.] Expanded or subcylindrical and hollow; celluliferous on two sides; cells arranged in quincunx order, roundish or oval, and raised in little pustules over the surface. Type *R. verrucosa*.
curvata, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 19, Niagara Gr.
frondosa, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 112, Niagara Gr.

FIG. 500.—*Rhinopora frondosa*.

St. Mo., p. 54, Clinton Gr.
verrucosa, Hall, 1852, Pal. N. Y., vol. 2, p. 48, Clinton Gr.

RHOMBOPORA, Meek, 1872, Pal. Eastern Nebraska, p. 141. [Ety. *rhombos*, rhomb; *poros*, pore.] Ramose, tubular, cells short; septa none; corallites radiating obliquely outward and upward on all sides from an imaginary axis; mouths rhombic or rhombic oval, and arranged in longitudinal and oblique spiral rows; interspaces thick, with minute pores visible in microscopic sections. Type *R. lepidodendroidea*.

armata, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 31, Kaskaskia Gr.

(?) *asperima*, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 70, Keokuk Gr.

attenuata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 70, Keokuk or Warsaw Gr.

confluens, see *Acanthoclema confluens*.

crassa, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 25, Up. Coal Meas.

decipiens, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 71, St. Louis Gr.

dichotoma, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 70, Burlington Gr.

elegantula, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 33, Kaskaskia Gr.

exigua, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 70, Burlington Gr.

gracilis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 70, Burlington Gr.

incrassata, Ulrich, 1888, Bull. Denison Univ., p. 89, Cuyahoga shales.

lepidodendroidea, Meek, 1872, Pal. Eastern Nebraska, p. 141, Up. Coal Meas.

tuberculosa, Hall, 1852, Pal. N. Y., vol. 2, p. 170, Niagara Gr.

tubulosa, Hall, 1852, Pal. N. Y., vol. 2, p. 49, Clinton Gr.

venosa, Spencer, 1884, Bull. No. 1, Mus. Univ.

FIG. 510.—*Rhombopora lepidodendroidea*.

ohioensis, Ulrich, 1888, Bull. Denison Univ., p. 90, Cuyahoga Shales.

persimilis, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 30, Kaskaskia Gr.

pulehella, Ulrich, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 31, Kaskaskia Gr.

simulatrix, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 71, St. Louis Gr.

(?) *spiralis*, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 71, Keokuk Gr.

subannulata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 45, Ham. Gr.

sulcifera, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 45, Ham. Gr.

tabulata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 70, Kaskaskia Gr.

tenuirama, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 70, Kaskaskia Gr.

transversalis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 71, Keokuk Gr.

varia, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 71, Keokuk Gr.

wortheni, Ulrich, 1884, Cin. Soc. Nat. Hist., vol. 7, p. 32, Kaskaskia Gr.

RHOPALONARIA, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 26. [Ety. *rhopalon*, a club.] Cells slender, fusiform, in single anastomosing series; cell mouths near the middle of the cells. Type *R. venosa*.

pertenuis, see *Stomatopora pertenuis*.

FIG. 511.—*Rhopalonaria venosa*.

venosa, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 26, Hud. Riv. Gr.

SAGENELLA, Hall, 1852, Pal. N. Y., vol. 2, p. 172. [Ety. *sagenella*, a little drag-net.] Membranous net incrusting other bodies; cells in parallel or diverging series, more or less oblong-quadrangular, and separated by a thin lamina. Type *S. membranacea*.

ambigua, Walcott, 1879, Utica Slate and related formations, p. 22, Utica Slate.

elegans, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 118, Niagara Gr.

membranacea, Hall, 1852, Pal. N. Y., vol. 2, p. 172, Niagara Gr.

FIG. 512.—*Sagenella elegans*. Magnified.

SCALARIPORA, Hall, 1881, Bryozoans of Up. Held. Gr., p. 17. [Ety., *scalare*, ladder; *poros*, pore.] Irregular groups of triangular branches, more or less concave, traversed transversely by sharp, elevated laminae at regular distances; cel-

luliferous on each face; cells radiating from the center to each angle of the branch; margins and summit of lamina noncelluliferous. Type *S. scalariformis*.

approximata, Ulrich, (in press,) *Geo. Sur. Ill.*, vol. 8, pl. 43, Ham. Gr.

scalariformis, Hall, 1881, *Bryozoans of Up. Held. Gr.*, p. 18, and *Pal. N. Y.*, vol. 6, p. 100, *Up. Held. Gr.*

separata, Ulrich, (in press,) *Geo. Sur. Ill.*, vol. 8, pl. 43, Ham. Gr.

subconcaua, Hall, 1881, *Bryozoans of Up. Held. Gr.*, p. 18, and *Pal. N. Y.*, vol. 6, p. 100, *Up. Held. Gr.*

SCENELLOPORA, Ulrich, 1882, *Jour. Cin. Soc. Nat. Hist.*, vol. 5, p. 150. [Ety. *scene*, tent; *ellus*, diminutive; *poros*, pore.] Zoarium broad, obconical; cell apertures on ridges, which radiate from the subsolid and depressed center of the upper surface. Type *S. radiata*.

radiata, Ulrich, 1882, *Jour. Cin. Soc. Nat. Hist.*, vol. 5, p. 158, *Trenton Gr.*

SCEPTROPORA, Ulrich, 1888, *Am. Geo.*, vol. 1, p. 228. [Ety. *skeptron*, staff; *poros*,

FIG. 513.—*Sceptropora facula*, x 18. a, Segment; b, vertical section; c, transverse section; d, transverse section of expanded part.

pore.] Zoarium articulated; segments short, numerous, club-shaped, lower half striated, noncelluliferous; upper half expanded, celluliferous, and having one or more articulating sockets; zoecia subcylindrical, radially arranged about a central axis; apertures subovate. Type *S. facula*.

facula, Ulrich, 1888, *Am. Geo.*, vol. 1, p. 228, *Hud. Riv. Gr.*

SELENOPORA, Hall, 1887, *Pal. N. Y.*, vol. 6, p. xvii. [Ety. *selen*, moon; *poros*, pore.] Zoarium explanate, incrusting; apertures subcircular, with an elevated denticulate peristome, and situated within polygonal vestibular areas formed by connecting walls, which traverse the surface; interior structure as in *Lichenalia*. Type *S. circincta*.

circincta, Hall, 1881, (*Lichenalia circincta*), *Trans. Alb. Inst.*, vol. 10, p. 11, and *Pal. N. Y.*, vol. 6, p. 86, *Up. Held. Gr.*

complexa, Hall, 1881, (*Lichenalia complexa*), *Trans. Alb. Inst.*, vol. 10, p. 11, and *Pal. N. Y.*, vol. 6, p. 87, *Up. Held. Gr.*

SEMISCOSCIUM, Prout, 1859, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 443. [Sig. somewhat like *Coscinum*.] Leaf-like expansion, somewhat penniform, without a shaft; sole formed of longitudinal and horizontal parallel ridges, surmounted by a cellular tissue, divided perpendicularly by thin, longitudinal septa, corresponding to the ridges, and supporting parallel lines of tortuous tubes alternately approximating and receding from each other; covered by a dense, strong crust, divided into a net-work of rays and dissepiments bounding rhomboidal or ovate fenestrules, giving passage to oblique cells; the tortuous tubes give place to quincuncial, oval openings in the fenestrules; each tortuous tube has a line of cells on each side. Type *S. rhomboideum*.

eriense, Prout, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 579, *Up. Held. Gr.*

obliquatum, Ulrich, 1886, *Cont. to Am. Pal.* p. 13, *Up. Held. Gr.*

planodorsatum, Ulrich, (in press,) *Geo. Sur. Ill.*, vol. 8, pl. 45, *Up. Held. Gr.*

rhomboideum, Prout, 1859, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 443, *Up. Held. Gr.*

rhombicum, Ulrich, (in press,) *Geo. Sur. Ill.*, vol. 8, pl. 45-54, *Ham. Gr.*

tuberculatum, Prout, 1860, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 579, *Up. Held. Gr.*

SEMIOFORA, Hall, 1884, *Rep. St. Geol.*, p. 51.

[Ety. *semi*, half; *poros*, pore.] Bryo-

zoum ramose; branches infrequent, bifurcating or trifurcating; margins parallel; celluliferous on both sides; cells arising from a mesial epitheca; apertures in longitudinal parallel rows, separated by ridges; two minute pits on the transverse space between adjacent apertures; apertures near the margin larger and more oblique than the others; margin striated; noncelluliferous. Type *S. bistigmata*.

bistigmata, Hall, 1884, *Rep. St. Geol.*, p. 57, *Ham. Gr.*

SEPTOPORA, Prout, 1859, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 448. [Ety. *septum*, partition; *poros*, pore.] Zoarium like *Fenestella*, but distinguished by the dissepiments, which have from one to four rows of cells. Type *S. cestransia*.

cestransia, Prout, 1859, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 448, *Kaskaskia Gr.*

decipiens, Ulrich, (in press,) *Geo. Sur. Ill.*, vol. 8, pl. 66, *Kaskaskia Gr.*

delicatula, Ulrich, (in press,) *Geo. Sur. Ill.*, vol. 8, pl. 64, *Low Coal Meas.*

robusta, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 64, Up. Coal. Meas.
 subquadrans, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 64, Kaskaskia Gr.
 SPATIOPORE, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., p. 155. [Ety. *spatium*, spread out; *poros*, pore.] Thin, incrusting; surface smooth or tuberculated; cells shallow; interstitial cells and spiniform tubuli. Type *S. aspera*.
 areolata, Foord, 1883, Cont. to Micropalaeontology, p. 21, Trenton Gr.
 aspera, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 166, Hud. Riv. Gr.
 lineata, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 167, Hud. Riv. Gr.
 maculosa, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 167, Hud. Riv. Gr.
 montifera, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 168, Hud. Riv. Gr.

FIG. 514.—*Spatiopora tuberculata* on an *Orthoceras*.

tuberculata, Edwards & Haime, 1851, (Chetetes tuberculatus.) Pol. Foss. d. Terr. Pal., p. 268, Hud. Riv. Gr.
 SPURAGIOPORA, (in press.) Ulrich, Geo. Sur. Ill., vol. 8, p. 303. Parasitic, forming very small subhemispheric patches on foreign bodies; cells with circular apertures and slight peristome, arranged in a subradial manner, in single or double rows. Type *S. parasitica*.
 parasitica, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 65, Kaskaskia Gr. and Coal Meas.
 STICTOPORA, Hall, 1847, Pal. N. Y., vol. 1, p. 73. [Ety. *stictos*, punctured; *poros*, pore.] Zoarium attached to foreign objects by an expanded base, ramose, branches thin, furnishing an acutely elliptical transverse section, and composed of two layers of cells, separated by epithelial laminae; cell apertures oval or circular, surrounded by peristome, separated by raised longitudinal lines; no interstitial cells; margins nonporiferous and striated. Type *S. elegantula*.
 acuta, Hall, 1847, Pal. N. Y., vol. 1, p. 74, Trenton Gr.
 alcyone, Billings, 1865, (Ptilodictya alcyone.) Catal. Sil. Foss. Antic., p. 36, Anticosti Gr.
 alternata, Hall, 1887, Pal. N. Y., vol. 6, pl. xxiii, A. Low. Held. Gr.
 angularis, Hall, 1887, Pal. N. Y., vol. 6, p. 252, Ham. Gr.
 arguta, Billings, 1865, (Ptilodictya arguta.) Catal. Sil. Foss. Antic., p. 36, Anticosti Gr.

basalis, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 169, Trenton Gr.
 bifurcata, VanCleve, 1883, 12th Rep. Ind. Geo. and Nat. Hist., p. 267, Niagara Gr.
 bifurcata, Hall, see *S. bristolensis*.
 bristolensis, n. sp. Ham. Gr. Proposed instead of *S. bifurcata*, Hall, 1887, Pal. N. Y., vol. 6, p. 254, which name was preoccupied.
 carbonaria, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 160, and Ohio Pal., vol. 2, p. 328, Coal. Meas.
 compressa, VanCleve, 1883, 12th Rep. Ind. Geol., and Nat. Hist., p. 267, Niagara Gr.
 crassa, Hall, 1852, Pal. N. Y., vol. 2, p. 45, Clinton Gr.
 crescens, Hall, 1887, Pal. N. Y., vol. 6, p. 91, Up. Held. Gr.
 crenulata, Hall, 1884, Rep. St. Geol., p. 44, Ham. Gr.
 dichotoma, Hall, syn. for *FIG. 515.—Stictopora carbonaria*.
 divergens, Hall, 1887, Pal. N. Y., vol. 6, p. 257, Ham. Gr.
 elegantula, Hall, 1847, Pal. N. Y., vol. 1, p. 75, Trenton Gr.
 fenestrata, see *Sulcopora fenestrata*.
 fidelis, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 68, Trenton Gr.
 fragilis, see *Dicranopora fragilis*.
 fruticosa, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 14, and Rep. St. Geol., pl. 25, fig. 12, 13, Up. Held. Gr.
 gilberti, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 63, and Ohio Pal., vol. 1, p. 94, Up. Held. Gr.
 glomerata, Hall, 1847, Pal. N. Y., vol. 1, p. 17, Chazy Gr.
 granatula, Hall, 1887, Pal. N. Y., vol. 6, p. 38, Low. Held. Gr.
 granifera, Hall, 1884, Rep. St. Geol., p. 45, Ham. Gr.
 graminifolia, Ringueberg, 1884, Proc. Acad. Nat. Sci., p. 147, Niagara Gr. Very poorly defined.
 incisurata, Hall, 1884, Rep. St. Geol., p. 38, Ham. Gr.
 incrassata, Hall, 1884, Rep. St. Geol., p. 47, Ham. Gr.
 indenta, Hall, syn. for *S. incisurata*.
 interstriata, Hall, 1884, Rep. St. Geol., p. 45, Ham. Gr.
 invertis, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 15, and Rep. St. Geol., pl. 25, fig. 24-26, Up. Held. Gr.
 labyrinthica, Hall, 1847, Pal. N. Y., vol. 1, p. 50, Birdseye Gr.
 lichenoides, Meek, 1873, Ohio Pal., vol. 1, p. 194, Up. Held. Gr.

FIG. 516.—*Stictopora gilberti*. Tangential section, showing lunarium.

t. Louis
 g. some-
 expansion
 thout a
 nal and
 mounted
 pendic-
 ata, cor-
 support-
 s tubes
 receding
 a dense,
 work of
 g rhom-
 ing pas-
 tubes
 openings
 ous tube
 e. Type
 lead Sci.,
 to Am.
 ss.) Geo.
 d. Gr.
 a, Prout,
 ns. St.
 d. Sci.,
 443, Up.
 Ulrich, (in
 Sur. Ill.,
 l. 45-54,
 a, Prout,
 St. Louis
 vol. 1, p.
 eld Gr.
 Hall, 1884,
 Pol., p. 51.
 i, half;
 e.] Bryo-
 quent, bi-
 rgins par-
 des; cells
 ca; aper-
 rows, sep-
 e pits on
 adjacent
 e margin
 than the
 cellulifer-
 Geol., p.
 St. Louis
 y. septum,
 rium like
 by the dis-
 ne to four
 ensia.
 St. Louis
 Kaskas-
 Geo. Sur.
 Gr.
 Geo. Sur.
 Meas.

limata, Hall, 1887, Pal. N. Y., vol. 6, p. 250, Ham. Gr.
 linearis, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 15, and Rep. St. Geol., 1883, pl. 25, fig. 4-5, Up. Held. Gr.
 lobata, Hall, 1887, Pal. N. Y., vol. 6, p. 256, Ham. Gr.
 magna, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 112, Niagara Gr.
 multifida, VanCleve, 1883, 12th Rep. Ind. Geol. and Nat. Hist., p. 268, Niagara Gr.
 multipora, Hall, syn. for *S. incisurata*.

FIG. 517.—*Stictopora mutabilis*. Deep tangential section $\times 50$, showing the primitive portion of the zoecia and the median tubuli in their walls; *d*, tangential section $\times 50$; *e*, transverse section $\times 50$, showing median tubuli; *f*, transverse section $\times 50$; *g*, vertical section $\times 50$; *h*, vertical section $\times 50$.

mutabilis, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 66, Trenton Gr.
nitidula, see *Dicranopora nitidula*.
obliqua, syn. for *S. incisurata*.
obliqua, Ringueberg, see *Ptilodictya obliqua*.
obsoleta, Hall, 1887, Pal. N. Y., vol. 6, p. 37, Low. Held. Gr.

FIG. 518.—*Stictopora ovatifora*.

orbipora, Hall, 1879, Desc. New Spec. Foss., p. 5, and 11th Rep. Ind. Geo. and Nat. Hist., p. 248, Niagara Gr.

ovata, Hall, 1887, Pal. N. Y., vol. 6, p. 248, Ham. Gr.
ovatifora, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 14, and Rep. St. Geol., pl. 25, fig. 23, 23a, Up. Held. Gr.
palmipes, Hall, 1884, Rep. St. Geol., p. 41, Ham. Gr.
papillosa, Hall, 1883, Rep. St. Geol., pl. 13, fig. 12-13, Low. Held. Gr.
paupera, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 69, Trenton Gr.
perarcta, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 15, and Pal. N. Y., vol. 6, p. 46, Up. Held. Gr.
permarginata, Hall, 1884, Rep. St. Geol., p. 46, Ham. Gr.
punctipora, Hall, 1852, Pal. N. Y., vol. 2, p. 157, Niagara Gr.
ramosa, Hall, 1847, Pal. N. Y., vol. 1, p. 51, Birdseye Gr.
rarpipora, Hall, 1852, Pal. N. Y., vol. 2, p. 46, Clinton Gr.
recta, Hall, 1887, Pal. N. Y., vol. 6, p. 253, Ham. Gr.
rectilinea, Hall, 1887, Pal. N. Y., vol. 6, p. 245, Ham. Gr.
rectilatera, Hall, syn. for *S. linearis*.
recubans, Hall, 1884, Pal. N. Y., vol. 6, p. 260, Ham. Gr.
rhomboidea, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 15, and Pal. N. Y., vol. 6, p. 95, Up. Held. Gr.

rigida, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 14, and Rep. St. Geol., 1883, pl. 25, fig. 15-16, Up. Held. Gr.

rustica, Billings, 1865, (Ptilodictya rustica.) Catal. Sil. Foss. Antic., p. 36, Anticosti Gr.

scitula, Hall, 1887, Pal. N. Y., vol. 6, pl. lxi, Niagara Gr.

scutolata, see *Ptilodictya scutolata*.

semistriata, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 14, and Rep. St. Geol., 1883, pl. 25, fig. 17-20, Up. Held. Gr.

serrata, see *Prismopora serrata*.

shafferi, see *Arthropora shafferi*.

similis, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 122, Niagara Gr.

sinuosa, Hall, 1884, Rep. St. Geol., p. 42, Ham. Gr.

FIG. 519.—*Stictopora punctipora*. Section $\times 50$.

stria
 24
 subco
 subri
 Ha
 sulca
 Mi
 tener
 Ca
 cos
 trillin
 243
 triser
 tumu
 246
 vanc
 and
 variat
 Ac
 vermi
 p. 9
 Stictop
 Nat
 tive
 Stic
 the
 ture
 angula
 Sur.
 ? basal
 vol.

cribros
 Minn
 frondif
 Sur.

Fig.
 interstr
 Nat. H
 ? undula
 Ill., vo

striata, Hall, 1887, Pal. N. Y., vol. 6, p. 246, Ham. Gr.

subcarinata, see *Tenipora subcarinata*.

subrigida, Hall, 1884, Rep. St. Geol., p. 43, Ham. Gr.

sulcata, Winchell, 1866, Rep. Low. Penin. Mich., p. 92, Ham. Gr.

tenera, Billings, 1865, (Ptilodictya tenera,) Catal. Sil. Foss. Antic., p. 36, Anticosti Gr.

trilineata, Hall, 1887, Pal. N. Y., vol. 6, p. 243, Ham. Gr.

triseriale, see *Acanthoclema triseriale*. tumulosa, Hall, 1887, Pal. N. Y., vol. 6, p. 246, Ham. Gr.

vancelei, Hall, 1883, 12th Rep. Ind. Geol. and Nat. Hist., p. 268, Niagara Gr.

variabilis, Prout, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 413, Up. Held. Gr.

vermicula, Hall, 1887, Pal. N. Y., vol. 6, p. 93, Up. Held. Gr.

STICTOPORELLA, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 152. [Ety. diminutive of *Stictopora*.] Distinguished from *Stictopora* by interstitial pits between the longer diameters of the cell apertures. Type *S. interstincta*.

angularis, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 71, Trenton Gr.

? basalis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 68, 69, and 75, Keokuk Gr.

FIG. 520.—*Stictoporella interstincta*. Natural size and magnified 18 diam.

cribrosa, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 69, Trenton Gr.

frondifera, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 72, Trenton Gr.

FIG. 521.—*Stictoporella interstincta*.

interstincta, Ulrich, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 169, Utica Slate Gr.

? undulata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 69, Kaskaskia Gr.

STICTOPORINA, Hall, 1887, Pal. N. Y., vol. 6, p. xx. [Ety. diminutive of *Stictopora*.]

Zoarium obtusely pointed at the base, enlarging above and becoming flattened; bifurcations, few; cells tubular arising from a mesotheca; apertures oval; interspersed space elevated, angular, inclosing the apertures in rhomboidal or polygonal areas. Type *S. claviformis*. claviformis, Hall, 1881, (Trematopora claviformis,) Trans. Alb. Inst., vol. 10, p. 181, and Pal. N. Y., vol. 6, p. 269, Ham. Gr.

Stictotrypa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8. Syn. (?) for *Stictopora*.

STOMATOPORA, Bronn, 1825, System d. urwelt. Pflanzenthier. [Ety. *stoma*, mouth; *poros*, perforation.] Zoarium adnate; cells in single branching series, mouths elevated, and at the end of the tubular cells. Type *S. dichotoma*.

alternata, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 235, Chemung Gr.

auloporoides, Nicholson, 1875, (Alecto auloporoides,) Ohio Pal., vol. 2, p. 267, Hud. Riv. Gr.

confusa, Nicholson, 1875, (Alecto confusa,) Ohio Pal., vol. 2, p. 267, Hud. Riv. Gr.

frondosa, Nicholson, 1875, (Alecto frondosa,) Ohio Pal., vol. 2, p. 266, Hud. Riv. Gr.

inflata, Hall, 1847, (Alecto inflata,) Pal. N. Y., vol. 1, p. 77, Trenton and Hud. Riv. Grs.

nexilis, James, 1875, (Alecto nexilis,) Int. to Catal. Cin. Foss., p. 3, Hud. Riv. Gr.

pertenuis, Ulrich, 1886, (Rhopalonia pertenuis,) 14th Ann. Rep. Geol. Sur. Minn., p. 59, Trenton Gr.

proutana, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 39, Hud. Riv. Gr.

STREBLOTRYPA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 403. [Ety. *streblon*, turned about; *trupa*, an opening.] Ramose, slender, solid; cells radiating from an imaginary axis, their primitive portion long, tubular; or from a linear axis when they are somewhat shorter; inferior hemisepta best developed, situated rather far down; apertures regularly elliptical, or somewhat truncated at the posterior margin, surrounded by a slight peristome and, within this, sometimes a narrow sloping area; arranged usually in rather regular longitudinal series; back of the aperture, occupying the depressed front of the cell, there are from one to twelve or more small pits, which, when numerous, are arranged in two or three rows; small acanthopores occasionally present. Type *S. nicklesi*.

FIG. 522.—*Stomatopora inflata*. Natural size and magnified.

- denticulata, Ulrich, 1888, (in press,) Bull. Denison Univ., vol. 4, p. 85, Cuyahoga shale.
- distincta, Ulrich, Geo. Sur. Ill., vol. 8, pl. 71, Kaskaskia Gr.
- hamiltonensis, Nicholson, 1874, (Ceriopora hamiltonensis,) Geo. Mag., vol. 1, p. 161, Ham. Gr.
- hertzeri, Ulrich, 1888, Bull. Denison Univ., vol. 4, p. 85, Cuyahoga shale.
- major, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 71, Keokuk Gr.
- multiporata, Ulrich, 1888, Bull. Denison Univ., vol. 4, p. 87, Waverly Gr.
- nicklesi, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 71, Kaskaskia Gr.
- obliqua, Ulrich, 1888, Bull. Denison Univ., vol. 4, p. 85, Cuyahoga shale.
- radialis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 72, Keokuk Gr.
- regularis, Ulrich, 1888, Bull. Denison Univ., vol. 4, p. 88, Waverly Gr.
- striata, Ulrich, 1888, Bull. Denison Univ., vol. 4, p. 87, Waverly Gr.
- subspinosa, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 71, Kaskaskia Gr.
- STROTOPORA, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, p. 383. [Ety. *strotos*, spread; *poros*, pore.] Ramose, branches large, irregular, solid or hollow; large, abruptly spreading cells, which are supposed to represent oecia, are distributed among the ordinary zoecia; when well-preserved they appear on the zoarial surface as strongly convex nodes, about 0.5 mm. in diameter, with an opening on one side. In all other respects like *Fistulipora*. Type *S. foveolata*.
- dermata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 77, Keokuk Gr.
- foveolata, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 77, Keokuk Gr.
- perminuta, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 47, Up. Held. Gr.
- SUBRETEPORA, D'Orbigny, 1850, Prodr. d. Paléont., t. 1, p. 22. [Ety. from *Retepora*.] Filiform, cylindrical branches, irregularly anastomosing; cells in a single row on the upper side of the branches (Ulrich says from 2 to 8 rows); apertures circular or oval. Type *S. reticulata*.

FIG. 523.—Subretepora angulata.

- angulata, Hall, 1852, (Retepora angulata,) Pal. N. Y., vol. 2, p. 163, Niagara Gr.
- aspera, Hall, 1847, (Gorgonia (?) aspera,) Pal. N. Y., vol. 1, p. 16, Chazy Gr.
- asperatostrata, Hall, 1852, (Retepora asperatostrata,) Pal. N. Y., vol. 2, p. 161, Niagara Gr.
- clathrata, Miller & Dyer, 1878, (Intricaria clathrata,) Cont. to Pal., No. 2, p. 7, Hud. Riv. Gr.
- corticosa, Ulrich, 1886, (Phyllopora corticosa,) 14th Rep. Geo. Sur. Minn., p. 61, Trenton Gr.

- dawsoni, Ulrich, (in press,) (Phylloporina dawsoni,) Geo. Sur. Ill., vol. 8, pl. 54, Trenton Gr.
- dichotoma, Hall, 1852, (Hornera dichotoma,) Pal. N. Y., vol. 2, p. 163, Niagara Gr.
- fenestrata, Hall, 1850, (Retepora fenestrata,) 3d Rep. N. Y. St. Mus. Nat. Hist., p. 178, Trenton Gr.
- gracilis, Hall, 1847, (Retepora gracilis,) Pal. N. Y., vol. 1, p. 15, Chazy Gr.
- incepta, Hall, 1847, (Retepora incepta,) Pal. N. Y., vol. 1, p. 15, Chazy Gr.
- reticulata, Hall, 1847, (Intricaria reticulata,) Pal. N. Y., vol. 1, p. 77, Trenton Gr.
- trentonensis, Nicholson, 1875, (Retepora trentonensis,) Geo. Mag., vol. 2, p. 37, Trenton Gr.
- variolata, Ulrich, 1882, (Phyllopora variolata,) Jour. Cin. Soc. Nat. Hist., vol. 5, p. 160, Hud. Riv. Gr.
- SULCOPORA, D'Orbigny, 1850, Prodr. d. Paléont., t. 1, p. 22. [Ety. *sulcus*, furrow; *poros*, pore.] Distinguished from *Stictopora* by the obtuse extremities of the branches, and by the perpendicular rows of apertures separated by elevated ridges and cross bars. Type *S. fenestrata*.

FIG. 525.—Sulcopora fenestrata. Natural size and magnified.

- fenestrata, Hall, 1847, (Stictopora fenestrata,) Pal. N. Y., vol. 1, p. 16, Chazy Gr.

- SYNOCLADIA, King, 1849, Ann. and Mag. Nat. Hist., 2d ser., vol. 3, p. 389. [Ety. *syn*, together; *klados*, young branch.] Cup-shaped,

with a central root-like base; reticulated, composed of rounded, narrow, often branched interstices, bearing on the inner face from 3 to 5 alternating, longitudinal rows of prominent edged cells, separated by narrow keels, studded with vesicles; dissepiments thin, spur-shaped, extending upward, and meeting those from the adjoining interstice, and bearing two rows of cells. Type *S. virgulacea*.

FIG. 524.—Subretepora reticulata. Natural size and magnified.

bise
A
rect
A
TENIO
III
ne
pa
na
str
of
ap
rou
rid
Ty
Fig. 526.
dely
Tangen
tion x
ing a t
lining c
tral reg
walls, a
often p
among
dietyon
subred
vol.
TENIOPO
Long
ribb
expa
on
Ptilo
long
from
prom
exigua
n. s.
occide
Ill.,
42, H
penni
Nic
187-
Mag,
s., v
123,
subca
Hall,
(Sti
subc
Trans
Inst.,
p. 1
Pal.
vol. 6
Ham.
Thallistig

ylloporina
8, pl. 54,
ora dich-
163, Ni-
ora fenes-
Mus. Nat.

biserialis, Swallow, 1858, Trans. St. Louis Acad. Sci., p. 179, Up. Coal Meas.
rectistyla, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 220, Kaskaskia Gr.
TENTODICTYA, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, [Ety. *tainia*, ribbon; *dictyon*, net.] Zoaria growing from a basal expansion into dichotomously divided narrow branches or broad fronds; cell structure very much as in some species of *Ptilodictya* (P. pavonia, D'Orb.); apertures elliptical or subcircular, surrounded by a sloping area; interspaces ridge-like; both "hemisepta" present. Type T. ramulosa.

FIG. 521.—*Tentodictya cingulata*. Tangential section x 50, showing a transverse lining of the central region of the walls, a character often present among the *Ptilodictyonidae*.

cingulata, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 67, Keokuk Gr.

frondosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 70, Keokuk Gr.

interpolata, Ulrich, 1888, Bull. Denison Univ., vol. 4, p. 80, Cuyahoga Shale.

ramulosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 67, Keokuk Gr.

ramulosa var. burlingtonensis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 67, Burlington Gr.

subrecta, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 67, St. Louis Gr.

TENIOPORA, Nicholson, 1874, Geo. Mag. Lond. n. s., vol. 1, p. 120. [Ety. *tainia*, ribbon; *poros*, pore.] Flattened linear expansion; dichotomous; cellulariferous on both sides. Distinguished from *Ptilodictya* and *Stictopora* by a central, longitudinal keel, which divides the frond into two lateral halves, and by prominent cell-mouths. Type T. exigua.

exigua, Nicholson, 1874, Geo. Mag. Lond. n. s., vol. 1, p. 122, Ham. Gr.

occidentalis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 42, Ham. Gr.

penniformis, Nicholson, 1874, Geo. Mag. Lond. n. s., vol. 1, p. 123, Ham. Gr.

subcarinata, Hall, 1881, (Stictopora subcarinata,) Trans. Alb. Inst., vol. 10, p. 191, and Pal. N. Y., vol. 6, p. 261, Ham. Gr.

FIG. 527.—*Teniopora penniformis*. a, Natural size; b, transverse section enlarged; c, fragment enlarged.

Thallistigma, Hall, syn for *Fistulipora*.

confertipora, see *Fistulipora confertipora*.
decipiens, see *Fistulipora decipiens*.

densa, see *Fistulipora densa*.

digitata, see *Fistulipora digitata*.

inclusa, see *Favicella inclusa*.

intercellatum, see *Fistulipora intercellata*.

lamellatum, see *Fistulipora lamellata*.

longimacula, see *Fistulipora longimacula*.

micropora, see *Fistulipora micropora*.

multaculeata, see *Fistulipora multaculeata*.

plana, see *Fistulipora plana*.

scrobiculata, see *Fistulipora scrobiculata*.

segregata, see *Fistulipora segregata*.

serrulata, see *Fistulipora serrulata*.

sparsipora, see *Prismopora sparsipora*.

spheroidea, see *Fistulipora spheroidea*.

subtilis, see *Fistulipora subtilis*.

triangula, see *Fistulipora triangularis*.

umbilicata, see *Fistulipora umbilicata*.

variopora, see *Fistulipora variopora*.

THAMNISCUS, King, 1849, Ann. and Mag. Nat. Hist., 2d ser., vol. 3, p. 389. [Ety. *thamniskos*, little shrub.] Stems frequently bifurcating more or less on one plane; cellulariferous on the side overlooking the imaginary axis; cells imbricated and arranged in quincunx; gemmuliferous vesicles overlying the cell apertures. Type T. dubius.

cisseis, Hall, 1883, Rep. St. Geol., pl. 22, fig. 23-30, Low. Held. Gr.

diffusus, Hall, 1852, (Retepora diffusa,) Pal. N. Y., vol. 2, p. 160, Niagara Gr.

fruticella, Hall, 1883, Rep. St. Geol. pl. 22, fig. 33, Low. Held. Gr.

divaricans, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 62, Keokuk Gr.

furcillatus, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 62, Kaskaskia Gr.

multiramus, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 19, and Rep. St. Geol. 1883, pl. 26, fig. 1-5, Up. Held. Gr.

nanus, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 19, Up. Held. Gr.

niagarensis, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 126, Niagara Gr.

nysa, Hall, 1883, Rep. St. Geol., pl. 22, fig. 47-48, Lower Held. Gr.

FIG. 528.—*Thamniscus niagarensis*.

octonarius, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 62, Up. Coal Meas.

pauciramus, Hall, 1884, Rep. St. Geol., p. 60, Ham. Gr.

ramulosus, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 62, Kaskaskia Gr.

ramulosus var. sevellensis, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 62, Low. Coal Meas.

oretopora natural size

rod. d. Pa-
cus, furrow;
d from Stic-
mities of the
pendicular
by elevated
pe S. fenes-

Hall., 1847,
pora fenes-
al. N. Y.,
16, Chazy

King, 1849,
d Mag. Nat.
ser., vol. 3, p.
ty. syn. to-
clados, young
Cup-shaped,
reticulated,
arrow, often
gon the inner
longitudinal
ills, separated
with vesicles;
shaped, ex-
eeting those
ce, and bear-
ype S. virgu-

- variolata*, Hall, 1883, Rep. St. Geol., pl. 22, fig. 34-46, Low. Held. Gr.
sculptilis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 62, Keokuk Gr.
Thamnopora, Hall. This name was preoccupied. See *Thamnotrypa*.
divaricata, see *Thamnotrypa divaricata*.
THAMNOTRYPA, Hall, 1887, Pal. N. Y., vol. 6, p. 101. [Ety. *thamnos*, bush; *trupa*, perforation.] Narrow, branching stipe, celluliferous on both sides; the divisions are not by bifurcation, as in *Stictopora*, but by lateral and abrupt divergence from the main stipe. Type *T. divaricata*.
divaricata, Hall, 1881, Trans. Alb. Inst., vol. 10, p. 16, and Pal. N. Y., vol. 6, p. 101, Up. Held. Gr.
TREMATELLA, Hall, 1886, Rep. St. Geol. and Pal. N. Y., vol. 6, p. xiv. [Ety. *trema*, hole; *ellus*, diminutive.] Ramose, solid; cells tubular, in contact below, diverging near the surface, intersected by septa; interapertural surface marked by pseudo-pores. Type *T. annulata*.
annulata, Hall, 1881, (Trematopora annulata,) Bryozoans of the Up. Held. Gr., p. 5, and Pal. N. Y., vol. 6, p. 69, Up. Held. Gr.
arborea, Hall, 1881, (Trematopora arborea,) Bryozoans of the Up. Held. Gr., p. 5, and Pal. N. Y., vol. 6, p. 69, Up. Held. Gr.
glomerata, Hall, 1887, Pal. N. Y., vol. 6, p. 70, Up. Held. Gr.
nodosa, Hall, 1887, Pal. N. Y., vol. 6, p. 176, Ham. Gr.
perspinulata, Hall, 1881, (Trematopora perspinulata,) Trans. Alb. Inst., vol. 10, p. 181, and Pal. N. Y., vol. 6, p. 175, Ham. Gr.
TREMATOPORA, Hall, 1852, Pal. N. Y., vol. 2, p. 149. [Ety. *trema*, hole; *poros*, pore; Ramose, branches solid, tuberculated or smooth; interstitial cells, spiniform in tubuli, and diaphragms present. Type *T. tuberculosa*.
alternata, see *Acanthoclema alternatum*.
americana, S. A. Miller, Jour. Cin. Soc. Nat. Hist., p. 312, Burlington Gr.
annulifera, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 67, and Geo. Wis., vol. 4, p. 254, Hud. Riv. Gr.
annulata, see *Trematella annulata*.
annulata var. *pronaspina*, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 6, Up. Held. Gr.
arborea, see *Trematella arborea*.
aspera, Hall, 1852, Pal. N. Y., vol. 2, p. 154, Niagara Gr.
calloporoides, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 38, Galena Gr.
camerata, see *Diamesopora camerata*.
canaliculata, Hall, 1883, Rep. St. Geol. pl. 11, fig. 12, Low. Held. Gr.
carinata, Hall, 1887, Pal. N. Y., vol. 6, p. 179, Ham. Gr.
claviformis, see *Stictopora claviformis*.
coalescens, Hall, 1852, Pal. N. Y., vol. 2, p. 150, Niagara Gr.
constricta see *Diamesopora constricta*.
corticosa, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 105, and Pal. N. Y., vol. 6, p. 15, Low. Held. Gr.
crassa, see *Lichenalia crassa*.
crebripora, Hall, 1879, Desc. New Spec. Foss., p. 3, and 11th Rep. Ind., Geol. and Nat. Hist., p. 236, Niagara Gr.
debilis, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 34, Galena Gr.
densa, Hall, 1874, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 105, Low. Held. Gr.
dispersa, see *Diamesopora dispersa*.
echinata, Hall, 1876, 26th Rep. N. Y. St. Mus. Nat. Hist., p. 112, Niagara Gr.
elongata, Hall, 1887, Pal. N. Y., vol. 6, p. 183, Ham. Gr.
fragilis, Winchell, 1863, Proc. Acad. Nat. Sci., p. 3, Waverly Gr.
granifera, Hall, 1887, Pal. N. Y., vol. 6, p. 186, Ham. Gr.
granistriata, see *Bactropora granistriata*.
granulata, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., vol. 4, p. 253, Hud. Riv. Gr.
granulifera, Hall, 1852, Pal. N. Y., vol. 2, p. 154, Niagara Gr. The same species is marked "n. sp." in 28th Rep. N. Y. St. Mus. Nat. Hist., probably by mistake.
halli, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 261, Niagara Gr.
hexagona, Hall, 1887, Pal. N. Y., vol. 6, p. 178, Ham. Gr.
immersa, Hall, 1887, Pal. N. Y., vol. 6, p. 185, Ham. Gr.
infrequens, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 111, Niagara Gr.
interplana, Hall, 1887, Pal. N. Y., vol. 6, p. 186, Ham. Gr.
lineata, Hall, 1887, Pal. N. Y., vol. 6, p. 181, Ham. Gr.
macropora, Hall, 1879, Desc. New Spec. Foss., p. 4, and 11th Rep. Ind. Geo. and Nat. Hist., p. 236, Niag. Gr.
maculosa, see *Lichenalia maculosa*.
minuta, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 113, Niagara Gr.
nitida, Ulrich, (in press,) Geo. Sur. Ill., vol. 8, pl. 34, Hud. Riv. Gr.
nodosa, Hall, 1887, Pal. N. Y., vol. 6, pl. xxiii, Low. Held. Gr.
orbipora, Hall, 1884, Rep. St. Geol., p. 12, Ham. Gr.
ornata, Ulrich, 1886, 14th Rep. Geo. Sur. Minn., p. 98, Trenton Gr.
osculum, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 110, Niagara Gr.
ostiolata, see *Chilotrypa ostiolata*.

FIG. 529. — Trematopora infrequens. Much enlarged.

ovati
fig.
paral
fig.
persp
p.
polyg
Ha
pond
Mu
punct
151,
primi
Sur

rectilin
Up.
regular
rhombif
scutulat
scutulat
signata
nata.
solida,
153,
sparsa,
155,
spicula
Foss.
spinulo
155,
spinulos
Mus.
cupie
striata,
153,
subimb
Foss.
234,
subquad
11, H
superba
Antic

stricta.
N. Y. St.
Pal. N. Y.,

New Spec.,
Ind., Geol.
a Gr.
o. Sur. Ill.,
N. Y. St.
Held. Gr.
ersa.
N. Y. St.
ara Gr.
., vol. 6, p.
Acad. Nat.
Z., vol. 6, p.
anistriata.
n. Rep. Geo.
nd. Riv. Gr.

Trematopora
s. Much en-

Y., vol. 6, p.

Rep. N. Y.
111, Niag-

N. Y., vol. 6,

Y., vol. 6, p.

New Spec.
Ind Geo. and

ulosa.

ep. N. Y. St.
agara Gr.

Geo. Sur. Ill.,

r.

Y., vol. 6,

Geol., p. 12,

ep. Geo. Sur.

ep. N. Y. St.

agara Gr.

olata.

FIG. 530.—*Aethroclema pulchellum*. a, Magnified view. (See page 293.)

rectilinea, Hall, 1881, Bryozoans of the Up. Held. Gr., p. 6, Up. Held. Gr.
regularis, see *Orthopora regularis*.
rhombifera, see *Orthopora rhombifera*.
scutulata, see *Acanthoclema scutulatum*.
scutulata, see *Orthopora scutulata*.
signata, see *Callotrypa macropora* var. *signata*.
solida, Hall, 1852, Pal. N. Y., vol. 2, p. 153, Niagara Gr.
sparsa, Hall, 1852, Pal. N. Y., vol. 2, p. 155, Niagara Gr.
spiculata, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 245, Niagara Gr.
spinulosa, Hall, 1852, Pal. N. Y., vol. 2, p. 155, Niagara Gr.
spinulosa, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist. The name was preoccupied. See *T. spiculata*.
striata, Hall, 1852, Pal. N. Y., vol. 2, p. 153, Niagara Gr.
subimbricata, Hall, 1879, Desc. New Spec. Foss., p. 4, and 11th Rep. Ind. Geo., p. 234, Niagara Gr.
subquadrata, Hall, 1884, Rep. St. Geol., p. 11, Ham. Gr.
superba, Billings, 1866, Catal. Sil. Foss. Antic., p. 93, Clinton and Niagara Grs.

tortallinea, Hall, 1884, Rep. St. Geol., p. 10, Ham. Gr.
transversa, Hall, 1884, Rep. St. Geol., p. 8, Ham. Gr.
tuberculosa, Hall, 1852, Pal. N. Y., vol. 2, p. 149, Niagara Gr.
tubulosa, Hall, 1852, Pal. N. Y., vol. 2, p. 151, Niagara Gr.
varia, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 111, Niagara Gr.
variolata, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 113, Niagara Gr.
vesiculosa, Winchell, 1863, Proc. Acad. Nat. Sci., p. 3, Burlington Gr.
whitfieldi, Ulrich, 1883, Jour. Cin. Soc. Nat. Hist., vol. 6, p. 262, Niagara Gr.
Trematopora, Hall, 1887, Pal. N. Y., vol. 6, p. 71. [Ety. *tropis*, keel; *poros*, pore.] Ramose, solid, cells in irregular longitudinal rows, separated by sinuous ridges; peristomes thin, slightly elevated. Type *T. nana*.
nana, Hall, 1887, Pal. N. Y., vol. 6, p. 71, Up. Held. Gr.
Tuberculopora, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 21. Not properly defined.
inflata, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 21. Not properly defined.
UNITRYPA, Hall, 1885, Rep. St. Geol., p. 36. [Ety. *unus*, one; *trupa*, perforation.] Form like *Fenestella*, having the branches connected by dissepiments; cell apertures in two ranges, separated by carinae, which are elevated, widened at the summit, and connected by thin, lateral processes or scales more or less numerous. Type *U. lata*.
acaulis, Hall, 1881, (Fenestella *acaulis*.) Bryozoans of Up. Held. Gr., p. 33, and Pal. N. Y., vol. 6, p. 131, Up. Held. Gr.
acaulis var. *inclinis*, Hall, 1887, Pal. N. Y., vol. 6, p. 132, Up. Held. Gr.
acclivis, Hall, 1887, Pal. N. Y., vol. 6, p. 138, Up. Held. Gr.
biserialis, Hall, 1882, (Fenestella *biserialis*.) Rep. St. Geol. and Pal. N. Y., vol. 6, p. 57, Low. Held. Gr.
conferta, Ulrich, 1886, Cont. to Am. Pal., p. 17, Up. Held. Gr.
consimilis, Hall, 1887, Pal. N. Y., vol. 6, p. 142, Up. Held. Gr.
elegantissima, Hall, 1881, (Fenestella *elegantissima*.) Trans. Alb. Inst., vol. 10, p. 36, and Pal. N. Y., vol. 6, p. 140, Up. Held. Gr.
fastigata, Hall, 1881, (Fenestella *fastigata*.) Trans. Alb. Inst., vol. 10, p. 36, and Pal. N. Y., vol. 6, p. 141, Up. Held. Gr.
facticius, Hall, 1887, Pal. N. Y., vol. 6, p. 137, Up. Held. Gr.

FIG. 531.—*Unitrypa lata*

lata, Hall, 1881, (*Fenestella lata*.) Trans. Alb. Inst., vol. 10, p. 34, and Pal. N. Y., vol. 6, p. 136, Up. Held. Gr.
nana, Hall, 1887, Pal. N. Y., vol. 6, p. 133, Up. Held. Gr.
nervia, Hall, 1874, (*Fenestella nervia*.) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 93, Low. Held. Gr.
nervia var. *constricta*, Hall, 1879, (*Fenestella nervia* var. *constricta*.) 32d Rep. N. Y. St. Mus. Nat. Hist., p. 174, Low. Held. Gr.
præcursor, Hall, 1874, (*Fenestella præcursor*.) 26th Rep. N. Y. St. Mus. Nat. Hist., p. 94, Low. Held. Gr.
pernodosa, Hall, 1881, (*Fenestella pernodosa*.) Trans. Alb. Inst., vol. 10, p. 35, Pal. N. Y., vol. 6, p. 139, Up. Held. Gr.
projecta, Hall, 1887, Pal. N. Y., vol. 7, p. 132, Up. Held. Gr.
retrorsa, Ulrich, 1886, Cont. to Am. Pal., p. 15, Up. Held. Gr.
scalaris, Hall, 1884, (*Fenestella scalaris*.) 36th Rep. N. Y. St. Mus. Nat. Hist., p. 66, Ham. Gr.
spatiosa, Hall, syn. for *U. lata*.

stipata, Hall, 1881, (*Fenestella stipata*.) Trans. Alb. Inst., vol. 10, p. 34, and Pal. N. Y., vol. 6, p. 134, Up. Held. Gr.
tegulata, Hall, 1881, (*Fenestella tegulata*.) Trans. Alb. Inst., vol. 10, p. 34, and Pal. N. Y., vol. 6, p. 135, Up. Held. Gr.
transversa, Hall, 1887, Pal. N. Y., vol. 7, p. 132, Up. Held. Gr.

WORTHENOPORA, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, p. 403. [Ety. proper name.] Bifoliate, branching or palmate; cells regularly arranged, sub-tubular or elongate rhomboidal, with the aperture semi-elliptical; on the surface the line of junction between the cells is marked by an elevated ridge; the truncated posterior margin of the aperture is raised into a less strong transverse bar; the elongate triangular depressed front appears perfectly plane. Type *W. spinosa*.

spatulata, Prout, 1859, (*Flustra spatulata*.) Trans. St. Louis Acad. Sci., vol. 1, p. 446, Warsaw Gr.
spinosa, Ulrich, (in press.) Geo. Sur. Ill., vol. 8, pl. 68, Keokuk and Warsaw Grs.

SUBKINGDOM MOLLUSCA.

CLASS BRACHIOPODA.

[Ety. *brachium*, arm; *pous*, foot.]

THE Brachiopoda are all marine animals, having a bivalve shell and a pair of long, ciliated, and usually spiral arms, with which they produce a current of water that carries the food to the mouth, which is close to the middle of the base of the shell. The valves of the shell, instead of being placed on each side of the animal, as in the Lamellibranchiata, are placed above and below it; so they are dorsal and ventral valves, instead of right and left valves. The ventral valve is generally larger than the dorsal, and projects beyond it at the beak. The beak is generally perforated, for the passage of a muscular peduncle, for the attachment of the animal; but in the Lingulidæ, the peduncle projects from the interior of the shell, between the umbones. When there is no peduncle, the shell attaches by the beak, or by the whole surface of the ventral valve. The dorsal valve is always free and imperforate. There is generally a pair of teeth in the ventral valve, developed from the hinge margin, that lock in corresponding cavities in the dorsal valve. Some genera have no teeth or hinge.

The shells of the living Rhychonellidæ and of many fossil genera consist of flattened prisms, parallel with each other, and directed obliquely to the surface of the shells, the interior of which is imbricated by their outcrop. The substance of

the shell is traversed by small canals from one surface to the other, through which little cœcal processes of the outer layer of the mantle pass, and are covered externally by a thickening of the epidermis.

They have no special branchial apparatus. The respiratory function is performed by the mantle, which is traversed by numerous blood-vessels. The arms are frequently supported upon a calcareous framework on the interior of the dorsal valve, as shown in the illustration of *Waldheimia australis*. The valves are opened by cardinal muscles, which originate on each side of the center of the ventral valve, and converge toward the hinge margin of the dorsal valve, behind the dental sockets, where there is usually a prominent cardinal process. The valves are closed by adductor muscles, of which there are four in *Crania* and *Discina*. In many fossil genera there are spiral processes, or loops, upon which are founded family distinctions.

Shells are sometimes silicified, and become so transparent that they show the coils when held up to the light. Sometimes the coils are preserved in empty shells; and when shells are found wholly filled with spar, both valves may be removed, and the sparry matrix scraped away on either side until the spirals may be clearly seen by holding the specimen up to the light.

The class was divided by King into two orders—the *Clistenterata* and *Tretenterata*—which correspond with the *Arthropomata* and *Lyropomata* of other authors. These divisions include the families as follows:

ORDER ARTHROPOMATA.

Athyridæ, *Atrypidæ*, *Orthidæ*, *Nucleospiridæ*, *Pentameridæ*, *Porambonitidæ*, *Productidæ*, *Rhynchonellidæ*, *Spiriferidæ*, *Strophomenidæ*, *Terebratulidæ*, *Triplesiidæ*.

ORDER LYOPOMATA.

Craniidæ, *Discinidæ*, *Lingulidæ*, *Obolidæ*, *Pholidopidæ*, *Siphonotretidæ*, *Trimerellidæ*.

FAMILY ATHYRIDÆ.—*Acambona*, *Athyris*, *Eumetria*, *Merista*, *Meristella*, *Whitfieldia*.

FAMILY ATRYPIDÆ.—*Anazyga*, *Atrypa*, *Cœlospira*, *Glassia*, *Koninekia*, *Zygospira*.

FAMILY CRANIIDÆ.—*Crania*, *Pseudocrania*.

FAMILY DISCINIDÆ.—*Discina*, *Orbiculoidea*, *Schizocrania*, *Schizobolus*, *Trematis*.

FAMILY LINGULIDÆ.—*Dignomia*, *Lingula*, *Lingulella*, *Lingulasma*, *Lingulepis*.

FAMILY NUCLEOSPIRIDÆ.—*Hindella*, *Meristina*, *Nucleospira*, *Retzia*, *Trematospira*.

FAMILY OBOLIDÆ.—*Dicellomus*, *Elkania*, *Leptobolus*, *Linnarsonia*, *Obolella*, *Obolus*.

FAMILY ORTHIDÆ.—*Meekella*, *Orthis*, *Orthisina*, *Skenidium*, *Vitulina*.

FAMILY PENTAMERIDÆ.—*Amphigenia*, *Anastrophia*, *Gypidula*, *Pentamerella*, *Pentamerus*, *Stenoschisma*.

FAMILY PHOLIDOPIDÆ.—*Pholidops*.

FAMILY PORAMBONITIDÆ.—*Porambonites*.

FAMILY PRODUCTIDÆ.—*Aulosteges*, *Chonetes*, *Productella*, *Productus*, *Strophalosia*.

FAMILY RHYNCHONELLIDÆ.—Camarella, Camarophoria, Eatonia, Eichwaldia, Leiorhynchus, Rhynchonella, Rhynchotreta, Stenoschisma.

FAMILY SIPHONOTRETIDÆ.—Acrothele, Acrotreta, Iphidæ, Kutorgina, Schizambon, Siphonotreta.

FAMILY SPIRIFERIDÆ.—Ambocælia, Cyrtia, Cyrtina, Martinia, Spirifera, Spiriferina, Syntrielasma, Syringothyris, Trigonotreta.

FAMILY STROPHOMENIDÆ.—Leptæna, Streptorhynchus, Strophodonta, Strophomena, Strophonella.

FAMILY TEREBRATULIDÆ.—Centronella, Cryptonella, Leptocælia, Rensseleria, Terebratula, Tropidoleptus, Vitulina, Waldheimia.

FAMILY TRIMERELLIDÆ.—Dinobolus, Lingulops, Monomerella, Trimerella.

FAMILY TRIPLESIDÆ.—Triplesia.

ACAMBONA, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 27. [Ety. *ake*, point; *ambon*, umbo.] Syn. for Eumetria.

prima, see Eumetria *prima*.

ACROTHELE, Linnarsson, 1876, Bihang till K. Vet. Akad. Handl., p. 20, Swed. Acad. Sci. on the Brachiopoda of the Paradoxides beds. [Ety. *akros*, pointed; *thele*, nipple; from the apex of the valve.] Shell thin, corneous, subcircular in outline, depressed, concentrically marked, and sometimes radiated; apex of ventral valve teat-like; subcentral or near the posterior margin, perforated; dorsal valve slightly convex, posterior margin slightly reflexed, and internally a low median longitudinal septum represented by an impression in the cast.

Type A. coriacea.

FIG. 532.—Acrothele subsidua. Interior of dorsal valve enlarged.

dichotoma, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 14, Up. Taconic. matthewi, Hartt, 1868, (Lingula matthewi), Acad. Geol., p. 644, St. John Gr. subsidua, White, 1874, (Acrotreta subsidua),

Rep. Invert. Foss., p. 6, and Geo. Sur. W. 100th Mer., vol. 4, p. 34, Up. Taconic.

Acrotreta, a

FIG. 533.—Acrotreta gemma. Small outline figures natural size; a and c, dorsal valves; b, ventral valve; d, area of ventral valve, which shows a central groove; e, area of another specimen having no groove; e, side view.

1, p. 133. [Ety. *akros*, the top or summit; *tretos*, perforated.] Shell triangu-

lar, larger valve conical, false area flat, bent back at right angles to the margin of the valve, longitudinally grooved along the center, and perforated at its extremity by a small circular aperture, the lines of growth encircle the shell and pass uninterruptedly over the false area; the smaller valve flat, operculiform, smooth, marked by concentric lines of growth; valves unarticulated. Type A. subconica.

attenuata, 1873, 6th Rep. Hayden's Geo. Sur. Terr., p. 463, Up. Taconic.

baileyi, Matthew, 1885, Trans. Soc. Can., p. 36, St. John Gr.

gemma, Billings, 1865, Pal. Foss., vol. 1, p. 216, Quebec Gr.

guelielma, Matthew, 1885, Trans. Roy. Soc. Can., p. 37, St. John Gr.

pyxidicula, White, 1874, Rep. Invert. Foss., p. 9, and Geo. Sur. W. 100th Mer., vol. 4, p. 53, Potsdam Gr.

subsidua, see Acrothele subsidua.

Egilops, Hall, 1850, 3d Rep. N. Y. St. Mus. Nat. Hist., p. 179. The name was pre-occupied for a genus in botany; beside it was founded on the cast of a Lamellibranch.

subcarinata. Name not to be retained.

AMBOCÆLIA, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 71. [Ety. *ambon*, umbo; *koilos*, the belly.] Distinguished from Orthis, Spirifera, etc., by the interior markings in the ventral valve, the thickened margins of the fissure are produced in short, strong teeth, but there is scarcely any extension of the dental plates; in the dorsal valve the bases of the crura continue attached to the inner surface of the valve for more than one-third of its length before becoming free; there is a lateral projection from these crural bases bounding the teeth sockets; the cardinal process is elongate, lying between the crura, and is bifurcated at the outer extremity as in Cyrtina; the muscular impressions are below the middle of the valve, often near the front and quadruple; the dorsal valve being concave, flat or de-

chwaldia,
a, Schiz-
ra, Spirif-
ta, Stro-
nsscheria,
erella.

area flat,
the margin
grooved
at its
aperture,
the shell
er the false
operculi-
concentric
articulated.

on's Geo.

. Soc.

oss, vol. 1.

Roy. Soc.

p. Invert.

W. 100th

Gr.

aa.

. St. Mus.

e was pre-

ny; beside

f a Lamel-

retained.

. N. Y. St.

ty. ambon.

tinguished

by the in-

tral valve,

fissure are

teeth, but

ion of the

valve the

attached to

for more

ore becom-

ection from

g the teeth

ss is elong-

ura, and is

mity as in

essions are

live, often

e; the dor-

lat or de-

pressed convex, the spires lie in the ventral valve. Type *A. umbonata*.
limbriata, Clappole, 1883, Proc. Am. Phil. Soc., p. 232, Portage Gr.

gemmula, syn. for *Spirifera planconvexa*.

gregaria, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 81, and Pal. N. Y., vol. 4, p. 261, Chemung Gr.

minuta, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 26, Waverly Gr.

nucleus, syn. for *Ambocella umbonata*.

præumbona, Hall, 1857, (Orthis præumbona,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 167, and Pal. N. Y., vol. 4, p. 262, Ham. Gr.

subumbona, see *Spirifera*, sub-umbona.

umbonata, Conrad, 1842, Fig. 534—*Ambocella umbonata*, Jour. Acad. Nat. Sci., vol. 8, p. 264, and Pal. N. Y., vol. 4, p. 259, Marcellus Shale and Ham. Gr.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

umbonata var. *gregaria*, see *A. gregaria*.

and Pal. N. Y., vol. 4, p. 383, Schokarie grit and Up. Held. Gr.

FIG. 536.—*Amphigenia elongata*.

elongata var. *undulata*, Hall, 1867, Pal. N. Y., vol. 4, p. 384, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

elongata var. *subtrigonalis*, Hall, 1857, (Meganteris subtrigonalis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 123, Up. Held. Gr.

FIG. 537.—*Amphigenia elongata*.

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

ANASTROPHIA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *ana*, with; *strophe*, a turning round; the relation of the valves is the reverse of that of *Pentamerus*.] Gib-

FIG. 538.—*Anastrophia internascens*.

internascens, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 168, Niagara Gr.

- interplicata, Hall, 1852, (*Atrypa interplicata*.) Pal. N. Y., vol. 2, p. 275, Niagara Gr.
- reversa, Billings, 1857, (*Pentamerus reversus*.) Rep. of Prog. Geo. Sur. Can., p. 215, Mid. Sil.
- verneuli, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 104, and Pal. N. Y., vol. 3, p. 260, Low. Held. Gr.
- ANAZYGA, Davidson, 1883, Supp. to Brit. Brachiopoda, vol. 5, pt. 1, p. 128. [Ety. *ana*, upward; *zygos*, a connecting band.] Small, longitudinally oval and striated; position of spiral cones as in *Zygospira*; about four coils in each spiral cone; stems attach to the hinge plate of the dorsal valve, extend parallel for a short distance, and then, bending at right angles, form two large curves facing the lateral parts of the valve before reaching their furthest extension in front, they give off a circular band or loop, which is directed upward toward the beak, and is exterior to the spiral cones on their dorsal side. Type *A. recurvirostra*.
- recurvirostra, Hall, 1847, (*Atrypa recurvirostra*.) Pal. N. Y., vol. 1, p. 140, Trenton Gr.
- Anomia, Linnaeus, 1767, Syst. Nat., 12th Ed. [Ety. *anomis*, unequal.] Not Palaeozoic.
- biloba*, see *Orthis biloba*.
- pecten*, see *Strophomena pecten*.
- reticularis*, see *Atrypa reticularis*.
- Anomiles, Wahlenberg, 1821, Act., Upsal.
- exporrectus*, see *Cyrtia exporrecta*.
- glaber*, see *Spirifera glabra*.
- punctatus*, see *Productus punctatus*.
- resupinatus*, see *Orthis resupinatus*.
- reticularis*, see *Atrypa reticularis*.
- rhomboidalis*, see *Strophomena rhomboidalis*.
- scabriculus*, see *Productus scabriculus*.
- semireticulatus*, see *Productus semireticulatus*.
- ATHYRIS, McCoy, 1844, Carb. Foss. Ireland, pp. 128 and 146. [Ety. *a*, without; *thuris*, a small door; in allusion to the absence of a deltidium or door. But the name is erroneous.] Nearly orbicular or ovate, both valves convex; no cardinal area, foramen, or hinge-line; spiral appendages attached to the hinge plate of the dorsal valve, very large, nearly filling the shell; a strong mesial septum in rostral part of dorsal valve; dental lamellae moderate; pallial and ovarian impressions thick, numerous, dichotomous; tissue of shell fibrous. Type *A. spiriferoides*.
- americana, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 89, Kaskaskia Gr.
- angelica, Hall, 1861, 14th Rep. N. Y. St. Mus. Nat. Hist., p. 99, and Pal. N. Y., vol. 4, p. 292, Chemung Gr.
- argentea, Shepard, 1838, Am. Jour. Sci. and Arts, vol. 34, p. 152, Up. Coal Meas.
- biloba, Winchell, 1865, (*Spirigera biloba*.) Proc. Acad. Nat. Sci., p. 118, Kinderhook Gr.
- blancha, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 115, Low. Held. Gr.
- caputserpentis, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 90, Up. Coal Meas.
- charitonensis, Swallow, 1860, (*Spirigera charitonensis*.) Trans. St. Louis Acad. Sci., vol. 1, p. 651, Coal Meas.
- chloe, Billings, 1860, Can. Jour., vol. 5, p. 282, Ham. Gr.
- clara, Billings, 1860, Can. Jour., vol. 5, p. 274, Up. Held. Gr.
- claytoni, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 256, Waverly Gr.
- clintonensis, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 89, Kaskaskia Gr.
- clusia, Billings, 1860, Can. Jour., vol. 5, p. 279, Up. Held. Gr.
- cora, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 94, and Pal. N. Y., vol. 4, p. 291, Ham. and Chemung Grs.
- corpulenta, Winchell, 1863, (*Spirigera corpulenta*.) Proc. Acad. Nat. Sci., p. 6, Waverly Gr.
- crassiscardinalis, White, 1860, Bost. Jour. Nat. Hist., vol. 7, p. 229, Waverly Gr.
- eborea, Winchell, 1866, (*Spirigera eborea*.) Rep. Low. Peninsula Mich., p. 94, Ham. Gr.
- euzona, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 91, Kaskaskia Gr.
- differens, McChesney, 1860, New Pal. Foss., p. 47, syn. for *A. subtilita*.
- formosa, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 91, Kaskaskia Gr.
- fultonensis, Swallow, 1860, (*Spirigera fultonensis*.) Trans. St. Louis Acad. Sci., vol. 1, p. 650, Ham. Gr.
- hannibalensis, Swallow, 1860, (*Spirigera hannibalensis*.) Trans. St. Louis Acad. Sci., vol. 1, p. 649, Waverly or Kinderhook Gr.
- hawni, Swallow, 1860, (*Spirigera hawni*.) Trans. St. Louis Acad. Sci., vol. 1, p. 652, Coal Meas.
- headi, see *Zygospira headi*.
- headi var. *anticostiensis*, see *Zygospira headi* var. *anticostiensis*.
- headi var. *borealis*, see *Zygospira headi* var. *borealis*.
- harpalyce, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 116, Low. Held. Gr.
- hirsuta, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 8, and Bull. Am. Mus. Nat. Hist., p. 49, Warsaw Gr.
- incrassata, Hall, 1858, Geo. Rep. Iowa, p. 600, Burlington Gr.
- intervarica, McChesney, 1860, Pal. Foss., p. 78, Burlington Gr. Not recognized.
- jacksoni, Swallow, 1860, (*Spirigera jacksoni*.) Trans. St. Louis Acad. Sci., vol. 1, p. 651, Coal Meas.
- julia, see *Meristella julia*.

era biloba,
8, Kinder-

Port. Soc.
Held. Gr.
Trans. St.
Up. Coal

(Spirigera
ouis Acad.
s.,
ur., vol. 5,
r., vol. 5,

1877, U. S.
l. 4, p. 256,

Trans. St.
9, Kaskas-
., vol. 5, p.

Y. St. Mus.
Y., vol. 4,
Gr.
(Spirigera
Sci., p. 6,

Bost. Jour.
verly Gr.
era eborea,
h., p. 94,

St. Louis
kaskia Gr.
New Pal.
illita.

St. Louis
1, Kaskas-

spirigera ful-
Acad. Sci.,

(Spirigera
ouis Acad.
or Kinder-

ra hawnii,
1, vol. 1, p.

Zygospira

pira beadi

e. Port. Soc.
Held. Gr.
Inst., vol.
Nat. Hist.,

ep. Iowa, p.

Pal. Foss.,
recognized.
igera jack-
Sci., vol. 1,

junia, Billings, 1866, Catal. Sil. Foss. |
Antic., p. 46, Anticosti Gr.

lamellosa, Leveille, 1835, (Spirifer
lamellosus,) Mem. Geol. Soc. France,
vol. 2, p. 39, Waverly Gr.

lara, Billings, 1866, Catal. Sil. Foss.
Antic., p. 47, Anticosti Gr.

maconensis, Swallow, 1860, (Spirigera
maconensis,) Trans. St. Louis Acad.
Sci. vol. 1, p. 651, Coal Meas.

maia, see Spirigera maia.

minima, Swallow, 1860, (Spirigera min-
ima,) Trans. St. Louis Acad. Sci.,
vol. 1, p. 649, Ham. Gr.

missouriensis, Swallow, 1860, (Spirigera
missouriensis,) Trans. St. Louis Acad.
Sci., vol. 1, p. 650, Coal Meas.

missouriensis, Winchell, 1865, (Spirigera
missouriensis,) Proc. Acad. Nat. Sci.,
p. 117, Lithographic limestone. This
name was preoccupied.

monticola, White, 1874, (Spirigera mon-
ticola,) Rep. Invert. Foss., p. 16, and
Geo. Sur. W. 100th Mer., vol. 4, p. 91,
Subcarboniferous.

naviformis, Hall, 1843, (Atrypa navifor-
mis,) Geo. 4th Dist. N. Y., p. 71, and
Pal. N. Y., vol. 2, p. 76, Clinton Gr.

obmaxima, McChesney, 1860, Desc., New
Pal. Foss., p. 80, and Geo. Sur. W. 100th
Mer., vol. 4, p. 92, Waverly Gr.

obvia, McChesney, 1860, Pal. Foss., p. 81,
Kaskaskia Gr. Not recognized.

ohioensis, Winchell, 1865, Proc. Acad.
Nat. Sci., p. 118, Waverly Gr.

orbicularis, McChesney, 1860, New Pal.
Foss., Coal Meas. Not recognized.

papilioniformis, McChesney, 1867, Trans.
Chi. Acad. Sci., vol. 1, Kaskaskia Gr.

parvirostris, Meek and Worthen, 1860,
Proc. Acad. Nat. Sci. Phil., p. 451,
Keokuk Gr. Referred later to A.

planosulcata.

pectinifera, Swallow, 1863, Trans. St.
Louis Acad. Sci., vol. 2, p. 88, Keo-
kuk Gr.

perinflata, McChesney, 1860, Desc. New
Pal. Foss., p. 81, Keokuk Gr. Not
recognized.

persinuata, Meek, 1877, U. S. Geo. Sur.,
40th parallel, p. 81, Carboniferous.

planosulcata, Phillips, 1836, Geo. York.,
vol. 2, p. 220, Keokuk Gr.

plattensis, Swallow, 1863, Trans. St. Louis
Acad. Sci., vol. 2, p. 87, Up. Coal Meas.

polita, Hall, 1843, (Atrypa polita,) Geo.
4th Dist. N. Y., pl. 65, fig. 5, and Pal.
N. Y., vol. 4, p. 293, Chemung Gr.

prinstana, see Mexistella prinstana.

prouti, Swallow, 1860, (Spirigera proutii,
Trans. St. Louis Acad. Sci., vol. 1, p.
649, Kinderhook or Waverly Gr.

reflexa, Swallow, 1863, Trans. St. Louis
Acad. Sci., vol. 2, p. 88, Warsaw Gr.

singletoni, Swallow, 1863, Trans. St. Louis
Acad. Sci., vol. 2, p. 87, Low. Coal
Meas.

solitaria, Billings, 1866, Catal. Sil. Foss.
Antic., p. 48, Anticosti Gr.

spiriferoides, Eaton, 1831, (Terebratula
spiriferoides,) Am. Jour. Sci., vol. 21, p.

FIG. 539.—Athyris spiriferoides. Dorsal and ventral view.

137, and Pal. N. Y., vol. 4, p. 285, Cor-
nif. and Ham. Gr.

squamosa, Worthen, 1884,
Bull. No. 2, Ill. St. Mus.

Nat. Hist., p. 24, and Geo.
Sur. Ill., vol. 8, p. 103, St.
Louis Gr.

sublamellosa, Hall, 1858,
Geo. Rep. Iowa, p. 702,
Kaskaskia Gr.

subquadrata, Hall, 1858,
Geo. Rep. Iowa, p. 703,
Kaskaskia Gr.

subtilita, Hall, 1852, Stans-
bury's Exped. to Great
Salt Lake, p. 409, Coal
Meas.

trinuclea, Hall, 1858, (Terebratula trin-
clea,) Trans. Alb. Inst., vol. 4, p. 7, and
Geo. Sur. Iowa, p. 659, Warsaw Gr.

tumida, Dalman, 1827, (Atrypa tumida.)
The fossil usually referred to this species
is Whitfieldia maria, which Davidson
regarded as a synonym for W. tumida.

tumidula, Billings, 1866, Catal. Sil. Foss.
Antic., p. 47, Anticosti Gr.

turgida, Shaler, 1865, Bulletin No. 4, M. C.
Z., Anticosti Gr. Not defined so as to
be recognized.

ultravarica, McChesney, 1861, Desc. New
Pal. Foss., p. 79, Keokuk Gr. Not
recognized.

umbonata, see Hindella umbonata.

vittata, Hall, 1860, 13th Rep. N. Y. St.
Mus. Nat. Hist., p. 89, and Pal. N. Y.,
vol. 4, p. 289, Cornif. and Ham. Grs.

ATRYPA, Dalman, 1827, Vet. Acad. Handl.,
p. 102. [Ety. a, without; trypa, a hole
or perforation. It was supposed the
shells had no foramen in the beak. The
name is erroneous.] Suborbicular,
transverse or elongated; articulating by
teeth and sockets; beak of the ventral
valve produced and incurved, the apex
truncated by a small, round perforation,
sometimes separated from the hinge-
line by a deltidium; valve more or less
convex with or without a defined sinus;
a strong tooth on each side at the base
of the broad fissure is somewhat bilobed
at the summit, with a crenulated groove
on the back; from the base of the teeth
a curving ridge extends forward and
partially incloses a broad, muscular

FIG. 540.
Athyris spirif-
eroides. Side
view.

scar; dorsal valve convex, with or without a mesial fold; hinge plate divided in the middle with a tooth-like plate on each side, the crura originating outside of these close to the dental sockets, and outside of the latter, close to the shell margins, there is a crenulated fold, which occupied the groove at the base of the tooth; the spires originating from the crura form two hollow cones, directed into the cavity of the dorsal valve, their adjacent sides being flattened and apices brought close together near the center of the bottom of the cavity; the processes at the base of the crura are directed into the cavity of the dorsal valve, and unite to form a loop; surface smooth, striate, or costate; structure fibrous. Type *A. reticularis*.

acutiplicata, see *Leptocœlia acutiplicata*.
acutirostra, see *Rhynchonella acutirostra*.
æquiradiata, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 266, Low. Held. Gr.
æquiradiata, see *Rhynchonella æquiradiata*.

affinis, syn. for *Atrypa reticularis*.

altilis, see *Rhynchonella altilis*.

ambigua, see *Camarella ambigua*.

aprinis, see *Rhynchonella aprinis*.

arata, see *Pentamerella arata*.

aspera, Schlotheim, 1813, (*Terebratula aspera*.) Petrefaktenkunde, p. 263, Ham. and Chemung Grs.

aspera var. *occidentalis*, Hall, 1858, Geo. Rep. Iowa, vol. 1, pt. 2, p. 515, Ham. Gr.

bidens, see *Rhynchonella bidens*.

bisulcata, see *Camarella bisulcata*.

borealis, Schlotheim, as identified by d'Archiac & Verneuil. Not American.

brevirostris, as identified by Hall, Pal. N. Y., vol. 2, p. 278. See *Pentamerus brevis* and *Anastrophia verneuli*.

camura, see *Trematospira camura*.

capax, see *Rhynchonella capax*.

cassidea, as identified by d'Archiac & Verneuil. Not American.

chemungensis, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 265, Chemung Gr.

circulus, see *Camarella circulus*.

concinna, see *Nucleospira concinna*.

comis, see *Pentamerus comis*.

concentrica, syn. for *Athyris spiriferoides*.

congesta, see *Tripllesia congesta*.

congregata, see *Stenochisma congregatum*.

contracta, see *Stenochisma contracta*.

corallifera, see *Eichwaldia corallifera*.
crassirostra, Hall, 1852, Pal. N. Y., vol. 2, p. 269, Niagara Gr.

crenulata, see *Terebratula crenulata*.

cuboides, as identified by Hall and others. See *Rhynchonella venustula*.

cuneata, see *Rhynchonella cuneata*.

cuspidata, see *Tripllesia cuspidata*.

cylindrica, see *Meristella cylindrica*.

deflecta, Hall, 1847, Pal. N. Y., vol. 1, p. 140, Trenton Gr.

dentata, see *Rhynchonella dentata*.

disparilis, see *Calospira disparilis*.

dubia, see *Rhynchonella dubia*.

dumosa, Hall, 1843, Geo. Rep., 4th Dist. N. Y., p. 272, Chemung Gr.

duplicata, see *Stenochisma duplicatum*.

elongata, syn. for *Rensseleria ovoides*.

emacerata, see *Rhynchonella emacerata*.

exigua, Hall, 1847, Pal. N. Y., vol. 1, p. 141, Trenton Gr.

extima, see *Stenochisma eximium*.

extans, see *Tripllesia extans*.

flabella, syn. for *Leptocœlia hemispherica*.

flabellites, see *Leptocœlia flabellites*.

galeata, see *Pentamerus galeatus*.

gibbosa, Hall, 1852, Pal. N. Y., vol. 2, p. 79, Clinton Gr.

globuliformis, see *Leiorhynchus globuliformis*.

hemiplicata, see *Camarella hemiplicata*.

hemispherica, see *Leptocœlia hemispherica*.

hirsuta, see *Trematospira hirsuta*.

hystrix, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 272, and Pal. N. Y., vol. 4, p. 326, Chemung Gr.

impressa, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 122, and Pal. N. Y., vol. 4, p. 315, Schoharie Grit.

FIG. 541.—*Atrypa reticularis*. a, Dorsal valve; b, ventral valve; c, anterior view.

impressa, Shaler. The name was preoccupied.

increbescens, syn. for *Rhynchonella capax*.

inflata, Conrad, 1843, Geo. Rep. 3d Dist. N. Y. Not defined.

intermedia, Hall, 1852, Pal. N. Y., vol. 2, p. 77, Clinton Gr.

interplicata, see *Anastrophia interplicata*.

lavis, see *Meristella lavis*.

lamellata, see *Rhynchonella lamellata*.

laticosta, Phillips, 1841, (*Terebratula laticosta*.) Pal. Foss., Chemung Gr. This species is not clearly identified in America.

lentiformis, syn. for *Atrypa reticularis*.

limitaris, see *Leiorhynchus limitare*.

mansoni, Salter, 1852, (*Rhynchonella mansoni*.) Sutherland's Jour., vol. 2, p. cccxi, Devonian.

marginalis, (?) Dalman, 1827, (*Terebratula marginalis*.) Vet. Acad. Handl., p. 143, Niagara Gr.

medialis, see *Eatonia medialis*.

mesacostalis, see *Leiorhynchus mesacostalis*.

modesta, see *Zygospira modesta*.

nasuta, see *Meristella nasuta*.

naviformis, see *Athyris naviformis*.

neglecta, see *Rhynchonella neglecta*.
nitida, see *Meristina nitida*.
nitida var. *oblata*, see *Meristina nitida* var. *oblata*.
nodostriata, Hall, 1852, Pal. N. Y., vol. 2, p. 272, Niagara Gr.
nucleolata, Hall, 1852, Pal. N. Y., vol. 2, p. 328, Coralline limestone.
nucleus, see *Triplesia nucleus*.
rustella, Castelnau, 1843, Syst. Sil., p. 39. Not recognized.
oblata, Hall, 1852, Pal. N. Y., vol. 2, p. 9, Medina Gr.

FIG. 542.—*Atrypa reticularis*. Interior of ventral valve; a, impression of adductor muscle; c, cardinal muscle; p, pedicle muscle; o, ovarian sinus; d, deltidium.

obtusiplicata, see *Rhynchonella obtusiplicata*.
octocostata, see *Pentamerella arata*.
peculiaris, see *Eatonia peculiaris*.
phoca, Salter, 1852, (*Rhynchonella phoca*,) Sutherland's Jour., vol. 2, p. cccxvi, Devonian.

planoconvexa, see *Leptocoelia planoconvexa*.
plebeia, Conrad, 1843, Geo. Rep. 3d Dist. N. Y., Ham. Gr. Preoccupied name.

pleiopleura, see *Rhynchonella pleiopleura*.
plena, see *Rhynchonella plena*.

plicata, see *Rhynchonella plicata*.

plicatula, (?) Linnæus, as identified by Hall, in Pal. N. Y., vol. 2, p. 279. May be stricken from the list as an erroneous identification.

plicatula, see *Rhynchonella plicatula*.

plicifera, see *Rhynchonella plicifera*.

polita, see *Athyris polita*.

prisca, syn. for *Atrypa reticularis*.

pseudomarginalis, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 84, and Pal. N. Y., vol. 4, p. 327, Up. Held. Gr.

quadrucostata, see *Leiorhynchus quadrucostatus*.

quadrucostata, Hall, 1852, see *Rhynchonella quadrucostata*.

rectiplicata, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 265, Low. Held. Gr.

recurvirostra, see *Anazyga recurvirostra*.

reticularis, Linnæus, 1767, (Anomia reticularis,) Syst. Nat., ed. 12, p. 1132, and Pal. N. Y., vol. 2, p. 72. It

FIG. 543.—*Atrypa reticularis*. Interior of dorsal valve, showing spirals; p, hinge plate.

occurs, with its varieties, in all the Groups of the Upper Silurian and Devonian formations, except the Oriskany sandstone. Some of its varieties or synonyms are, *Atrypa affinis*, *A. lentiformis*, *A. prisca*, *A. tribulis*, *Hipparionyx consimilis*, etc.

robusta, see *Rhynchonella robusta*.

rostrata, see *Meristella rostrata*.

rugosa, see *Rhynchonella rugosa*.

scitula, see *Meristella scitula*.

semiplicata, see *Rhynchonella semiplicata*.

singularis, see *Eatonia singularis*.

sordida, see *Rhynchonella sordida*.

spinosa, Hall, 1843, Geo. 4th Dist. N. Y., p. 200, Cornif., Ham., Tully, and Chemung Grs. Equal to *Atrypa aspera* var. *occidentalis*.

subcuboides, D'Orbigny, see *Rhynchonella venustula*.

subtrigonalis, see *Rhynchonella subtrigonalis*.

sulcata, see *Merista sulcata*.

tenuilineata, Hall, 1843, Geo. 4th Dist. N. Y., p. 272, Chemung Gr.

tribulis, syn. for *Atrypa reticularis*.

tumida, see *Athyris tumida*.

unguiformis, syn. for *Orthis proximus*.

unisculcata, see *Meristella unisculcata*.

AULOSTEGES, Helmerston, 1847, Bull. de la Classe Physi. Math. Acad. Sci. St. Petersburg, vol. 6, p. 135. [Ety. *aulos*, tube; *steges*, chamber.] Shell subpentagonal; ventral valve most convex,

FIG. 544.—*Aulosteges wangenheimi*. a, Triangular hinge area; d, convex pseudodeltidium; j, cardinal process; a, adductor impression.

beak produced, twisted, area triangular, interrupted by a pseudodeltidium not reaching the hinge-line, which is straight and toothless; dorsal valve convex at the umbo, depressed or concave laterally; cardinal edge more or less developed; surface of valves with short tubular spires; in the interior of the dorsal valve a trifid cardinal process is made to fill the uncovered portion of the fissure, and serve as the point of attachment to the cardinal muscle; under this process a longitudinal mesial ridge extends nearly to the margin, and on either side are elongated, ramified adductor scars; the reniform impressions, after dividing the above named muscle, extend by an outward oblique curve to near the margin, when, turning backward and inward, termi-

nate some distance from their origin; two brachial elevations under the adductor move toward the center of the valve. Type *A. wangenheimi*.
guadalupensis, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 292, Permian Gr.
spondyliiformis, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 118, Up. Coal. Meas.
Billingsia, Ford, 1885. The name being preoccupied, see *Elkania*.
Brachymerus, Shaler. The name was preoccupied for a genus of Coleoptera. See *Anastrophia*.
Brachyprium, Shaler, syn. for *Strophomena geniculatum*, see *Strophomena geniculata*.
leda, see *Strophomena leda*.
ventricosum, see *Strophomena ventricosa*.
CAMARELLA, Billings, August, 1858, Can. Nat. and Geol., vol. 4, p. 301. [Ety. *kamara*, arching chamber; *ellus*, diminutive.] Shell ovate or subcircular, beaks small, hinge-line short; mesial fold and sinus becoming obsolete in the middle part of the shell, below which the radiating striae are more or less numerous, while above concentric striae occur. Type *C. volborthi*.

FIG. 545.—*Camarella hemiplicata*. Dorsal, ventral, and side views.

ambigua, Hall, 1847, (*Atrypa ambigua*), Pal. N. Y., vol. 1, p. 143, Trenton Gr.
antiquata, Billings, 1861, Pal. Foss., vol. 1, p. 10, Georgia Gr.
bisulcata, Emmons, 1842, (*Orthis bisulcata*), Geo. Rep. N. Y., p. 395, and Pal. N. Y., vol. 1, p. 139, Trenton Gr.
breviplicata, Billings, 1865, Pal. Foss., vol. 1, p. 304, Quebec Gr.
calcifera, Billings, 1861, Can. Nat. and Geo., vol. 6, p. 318, Calcif. Gr.
circulus, Hall, 1847, (*Atrypa circulus*), Pal. N. Y., vol. 1, p. 142, Trenton Gr.
congesta, see *Tripllesia congesta*.
costata, Billings, 1865, Pal. Foss., vol. 1, p. 305, Quebec Gr.
cuspidata, see *Tripllesia cuspidata*.
extans, see *Tripllesia extans*.
hemiplicata, Hall, 1847, (*Atrypa hemiplicata*), Pal. N. Y., vol. 1, p. 144, Trenton Gr.
lenticularis, Billings, 1866, Catal. Sil. Foss. Antic., p. 45, Anticosti Gr.
longirostra, Billings, 1858, Can. Nat. and Geo., vol. 4, p. 302, Chazy Gr.
nucleus, see *Tripllesia nucleus*.
ops, Billings, 1862, Pal. Foss., vol. 1, p. 148, Mid. Sil.
ortoni, see *Tripllesia ortonii*.
panderi, Billings, 1858, Can. Nat. and Geo., vol. 4, p. 301, Black Riv. Gr.

parva, Billings, 1865, Pal. Foss., vol. 1, p. 219, Quebec Gr.
polita, Billings, 1865, Pal. Foss., vol. 1, p. 305, Quebec Gr.
primordialis, see *Tripllesia primordialis*.
reversa, see *Anastrophia reversa*.
varians, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 445, Chazy Gr.
volborthi, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 301, Black Riv. Gr.
waldronensis, see *Tripllesia waldronensis*.
Camarium, Hall, 1859, Pal. N. Y., vol. 3, p. 486, syn. for *Merista*.
elongatum, see *Merista elongata*.
typum, see *Merista typum*.

FIG. 546.—*Camarella volborthi*. Dorsal, ventral, and side views.

CAMAROPHORIA, King, 1844, Ann. and Mag. Nat. Hist., vol. 14, p. 313. [Ety. *kamara*, an arched chamber; *phoreo*, I carry.] Subtrigonal, convex longitudinally; mesial fold and sinus; beak acute, more or less incurved, small fissure beneath; no area or deltidium; plicated, impunctate, articulating by teeth and sockets; dental plates in the ventral valve, conjoined at their dorsal margins, forming a trough-shaped process affixed to a low, medio-longitudinal plate; the space between the sockets in the dorsal valve is occupied by a small, cardinal, muscular protuberance, on either side of which two slender processes curve upward; from beneath the cardinal process a vertical mesial septum, a third or more of the length of the valve, supporting along its upper edge a spatula-shaped process, dilated toward its free extremity, and projected with a curve to near the center of the shell. Type *C. schlotheimi*.
bisulcata, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 296, Permian Gr.
eucharis, Hall, 1867, Pal. N. Y., vol. 4, p. 368, Corniferous Gr.

FIG. 547.—*Camarophoria giffordii*. a, Dorsal view; b, ventral valve; c, profile view.

giffordii, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 39, and Geo. Sur. Ill., vol. 7, p. 318, Middle Coal Meas.
globulina, Phillips, 1844, as identified by Geinitz, is *Rhynchonella uta*.
occidentalis, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 313, Burlington Gr.

, vol. 1, p.

, vol. 1, p.

ordialis.

arella vol-
t, ventral,v. Gr.
onensis.
vol. 3, p.

and Mag.
ty, kamara,
I carry.
itudinally;
eak acute,
fissure be-
; plicated,
teeth and
the ventral
dorsal mar-
ed process
ngitudinal
he sockets
upied by a
tubulance,
two slender
m beneath
ical mesial
the length
g its upper
ess, dilated
d projected
nter of the

s. St. Louis
man Gr.
vol. 4, p.

sal view; b.

No. 1, Ill.
d Geo. Sur.
l Meas.
entified by

Jour. Cin.
3, Burling-

schlothelmi, Von Buch, 1834, (Terebratulites schlothelmi,) Mem. de la Soc. Geol., vol. 3, p. 138, Permian Gr.
subtrigona, Meek & Worthen, 1860, (Rhynchonella subtrigona,) Proc. Acad. Nat. Sci. Phil., p. 451, and Geo. Sur. Ill., vol. 2, p. 251, Keokuk Gr.
swallowana, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 394, Permian Gr.

wortheni, Hall, 1858, (Rhynchonella wortheni,) Trans. Alb. Inst., vol. 4, p. 11, and Bull. Am. Mus. Nat. Hist., p. 54, Warsaw Gr.

CENTRONELLA, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 131. [Ety. a little point.] General form like Terebratula; dorsal valve with a loop consisting of two ribbon-like lamellæ, which extend about half the length of the shell, at first curving outward and then approaching until their lower extremities meet at an acute angle; here they unite and are reflected backward toward the beak in a thin, flat, vertical plate; near their origin each bears upon the ventral side a single triangular crural process. Type C. glansfagea.

v allii, Winchell, 1865, Proc. Acad. Nat. Sci., p. 123, Waverly or Marshall Gr.

alveata, Hall, 1857, (Rhynchonella alveata,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 124, Onondaga Gr.

anna, Hartt, 1868, Acad. Geol., p. 300, Subcarb.

billingsana, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 352, Niagara Gr.

crassicaudinalis, Whitfield, 1882, Bull. Ann. Mus. Nat. Hist., No. 3, p. 55, Warsaw Gr.

flora, Winchell, 1879, Proc. Am. Phil. Soc., vol. 12, p. 254, Marshall Gr.

glansfagea, Hall, 1857, (Rhynchonella glansfagea,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 125, and Pal. N. Y., vol. 4, p. 399, Schoharie grit, Cornif. Gr. and Oriskany sandstone.

glaucia, Hall, 1867, Pal. N. Y., vol. 4, p. 403, Ham. Gr.

hecate, Billings, 1861, Can. Jour. vol. 6, p. 272, Up. Held. Gr.

impressa, Hall, 1861, 14th Rep. N. Y. St. Mus. Nat. Hist., p. 102, and Pal. N. Y., vol. 4, p. 402, Ham. Gr. Prof. Billings said this is a syn. for C. hecate.

julia, Winchell, 1862, Proc. Acad. Nat. Sci., vol. 14, p. 405, and Pal. N. Y., vol. 4, p. 419, Marshall Gr.

ovata, Hall, 1867, Pal. N. Y., vol. 4, p. 419, Up. Held. Gr.

Charionella, Billings, 1861, Can. Jour. Ind. Sci., and Art., p. 148, syn. for Meristella.

circe, see Meristella circe.

doris, see Meristella doris.

(?) hyale, see Meristella hyale.

CHONRTES, Fischer, 1837, Oryct. Moscou, p. 131. [Ety. chone, a little cup.] Shell thin, semi-cylindrical, transverse section semi-oval, ventral valve convex, dorsal concave hinge-line straight; external margin of the area of ventral valve bearing a row of tubular spines, foramen distinct but partially closed by a pseudo-deltidium; dorsal valve with a cardinal process, simple at the base, but bifid or grooved at the extremity; valves articulated by teeth, surface radiately striated, often spinous, interior pustulose or papillose. Type C. arcinulatus.

acutiradiatus, Hall, 1843, (Strophomena acutiradiata,) Geo. Rep. 4th Dist. N. Y., p. 171, and Pal. N. Y., vol. 4, p. 120, Up. Held. Gr.

antiope, Billings, 1874, Pal. Foss., vol. 2, p. 19, Low Devonian.

arcuatus, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 116, and Pal. N. Y., vol. 4, p. 119, Up. Held. Gr.

armatus, DeKoninck, the specimens referred to this species belong to C. pusillus.

canadensis, Billings, 1874, Pal. Foss., vol. 2, p. 17, Lower Devonian.

carinatus, Conrad, 1842, (Strophomena carinata,) Jour. Acad. Nat. Sci., vol. 8, p. 257, and Pal. N. Y., vol. 4, p. 133, Ham. Gr.

complanatus, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 56, and Pal. N. Y., vol. 3, p. 418, Oriskany sandstone.

cornutus, Hall, 1843, (Strophomena cornuta,) Geo. Rep. 4th Dist. N. Y., and Pal. N. Y., vol. 2, p. 64, Clinton Gr.

dawsoni, Billings, 1874, Pal. Foss., vol. 2, p. 18, Low Devonian.

deflectus, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 149, and Pal. N. Y., vol. 4, p. 126, Ham. Gr.

emmetensis, Winchell, 1866, Rep. Low. Penin. Mich., p. 92, Ham. Gr.

filistriatus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 127, Devonian.

fischeri, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 25, Kinderhook Gr.

flemingi, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 26, Permian Gr.

geinitzianus, N. Sp., Up. Coal Meas. Proposed instead of C. glabra of Geinitz in Carb. und Dyas in Neb., p. 60, tab. 4, fig. 15 to 18, which name was preoccupied.

geniculatus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 29, Waverly or Marshall Gr.

gibbosa, syn. for C. deflectus.

glaber, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 117, Up. Held. Gr.

glabra, Geinitz, 1866, Carb. und Dyas. The name was preoccupied. See C. geinitzianus.

granuliferus, Owen, 1852, Geo. Rep. Wis., Iowa, and Minn., p. 583, Coal Meas.

FIG. 518.—Centronella hecate, a. Showing loop; b, c, and d, different views.

hecate, Billings, 1861, Can. Jour. vol. 6, p. 272, Up. Held. Gr.

impressa, Hall, 1861, 14th Rep. N. Y. St. Mus. Nat. Hist., p. 102, and Pal. N. Y., vol. 4, p. 402, Ham. Gr.

- hemisphericus, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 116, and Pal. N. Y., vol. 4, p. 118, Schoharie grit and Cornif. Gr.
- illinoisensis, Worthen, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 571, and Geo. Sur. Ill., vol. 3, p. 505, Kaskaskia Gr.
- iowensis, Owen, 1852, Geo. Rep. Iowa, Wis. and Minn., p. 584, Carb.
- koninckanus, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, 2d ser., p. 30, Devonian.
- levis, Keyes, 1888, Proc. Acad. Nat. Sci. Phil., pl. xii, figs. 3a, 3b, Coal Meas.
- laticosta, syn. for *C. mucronatus*.
- lepidus, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 148, and Pal. N. Y., vol. 4, p. 132, Marcellus shale and Ham. Gr.
- lineatus, Conrad, 1839, (*Strophomena lineata*), Ann. Geo. Rep. N. Y., p. 64, and Pal. N. Y., vol. 4, p. 121, Up. Held. Gr.
- littoni, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 25, Ham. Gr.
- loganensis, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 253, Waverly Gr.
- logani, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci. vol. 3, p. 30, Burlington Gr.
- logani var. aurora, Hall, 1867, Pal. N. Y., vol. 4, p. 137, Tully limestone and Ham. Gr.
- maclurii, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 28, Ham. Gr.
- macrostriatus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 126, Devonian.
- martini, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 29, Ham. Gr.
- melonicus, Billings, 1874, Pal. Foss., vol. 2, p. 15, Gaspe limestone No. 8, Devonian.
- mesolobus, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 27, Coal Meas.
- michiganensis, Stevens, 1858, Am. Jour. Sci., vol. 25, p. 262, Marshall Gr.
- millepunctatus, Meek & Worthen, 1870, Proc. Acad. Nat. Sci., p. 35, and Geo. Sur. Ill., vol. 5, p. 566, Coal Meas.
- minimus, Hall. Being preoccupied by Sowerby. See *C. undulatus*.
- mucronatus, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 180, and Pal. N. Y., vol. 4, p. 124, Corniferous and Ham. Grs.
- mucronata, Meek & Hayden, 1858, Proc. Acad. Nat. Sci., p. 262, Coal Meas. This name was preoccupied; moreover it is a syn. for *C. granuliferus*.
- multicosta, Winchell, 1863, Proc. Acad. Nat. Sci., p. 5, Marshall Gr.
- muricatus, Hall, 1867, Pal. N. Y., vol. 4, p. 143, Chemung Gr.
- novascoticus, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 144, Niagara Gr.
- ornatus, Shumard, 1855, Geo. of Mo., p. 202, Waverly or Kinderhook Gr.
- parvus, Shumard, 1855, Geo. of Mo., p. 201, Coal Meas.
- permianus, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 390, Permian Gr.
- planumbonus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 450, and Geo. Sur. Ill., vol. 2, p. 253, Keokuk Gr.
- platynotus, White, 1874, Rep. Invert. Foss., p. 19, and Geo. Sur. W. 100 Mer., vol. 4, p. 121, Subcarboniferous.
- pulchellus, Winchell, 1862, Proc. Acad. Nat. Sci., p. 410, Marshall Gr.
- pusillus, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 149, and Pal. N. Y., vol. 4, p. 128, Ham. Gr.
- reversus, Whitfield, 1882, Desc. New Spec. Foss., from Ohio, p. 213, Marcellus shale.
- scitulus, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 147, and Pal. N. Y., vol. 4, p. 130, Ham. Gr.
- setigerus, Hall, 1843, (*Strophomena setigera*), Geo. Rep. 4th Dist. N. Y., p. 180, and Pal. N. Y., vol. 4, p. 129, Ham. and Chemung Grs.
- shumardanus, DeKoninck, 1847, Recherches sur les Anim. Foss., p. 192, Waverly Gr.
- smithi, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 24, Coal Meas.
- striatellus, Dalman, 1827, (*Orthis striatella*), Kongl. Svenska Ak. Handl., p. 111, Up. Sil.
- syrtalis, syn. for *C. carinata*.
- tenuistriatus, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 144, Up. Sil.
- tuomeyi, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, 2d ser., p. 28, Ham. Gr.
- undulatus, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 155, Niagara Gr.
- variolatus, DeKoninck, 1847, Monogr. du genre Chonetes, p. 206, Coal Meas.
- verneuillanus, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 26, Coal Meas.
- verneuillanus var. utahensis, Meek, 1876, Simpson's Rep. on Gt. Basin of Utah, p. 348, Carboniferous.
- yandellanus, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 118, and Pal. N. Y., vol. 4, p. 123, Corniferous Gr.
- CELOSPIRA, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 146. [Ety. *koilos*, hollow; *speira*, spire.] Ovate or suborbicular, concavo-convex, surface finely plicated, usually undefined mesial fold and sinus, beak small, foramen triangular; internal spires forming two flattened coils connected by a strong loop. Type *C. concava*.

FIG. 549.—*Chonetes mesolobus*. Ventral valve.

FIG. 551.

107
nif

FIG. 551.—*Chonetes parvulus*. Ventral valve.

disj
CRANIA

Geol
vol.
par
rate
by
obje
less
or
radi
not
of t
is a
liga
four
app
post
and
reco
bur

FIG. 552.—*Chonetes aurora*. Posterior muscle scar.

acadie
vol.
anna,
Univ
aurora
Mus.

concava, Hall, 1857, (*Leptocœlia concava*),
10th Rep. N. Y. St. Mus. Nat. Hist., p.

FIG. 550.—*Crœlospira concava*. Magnified view of spirals.

107, and Pal. N. Y., vol. 3, p. 245, Corniferous Gr.

FIG. 551.—*Crœlospira disparilis*. Dorsal and ventral views.

dichotoma, Hall, 1859, (*Leptocœlia dichotoma*), Pal. N. Y., vol. 3, p. 452, Oriskany sandstone.

disparilis, Hall, 1852, (*Atrypa*

disparilis), Pal. N. Y., vol. 2, p. 277, Niagara Gr.

CRANIA, Retzius, 1781, *Schriften der Berliner Gesellschaft Naturforschende Freund*, vol. 2, p. 72. [Ety. *kranton*, the upper part of a skull.] Shell circular, subquadrate, transverse, or elongated, attached by its ventral valve to some foreign object; upper or dorsal valve more or less convex or conical; apex central or subcentral; surface smooth, spiny, radiated, or concentrically lined, and not unfrequently having the markings of the object to which the lower valve is attached; no articulating hinge or ligament, but valves held in place by four muscles; anterior adductor scars approximate and close to the center; posterior pair near the cardinal edge, and widely separated; structure calcareous and tubular. Type C. brattenburgensis.

FIG. 552.—*Crania anomala*, 2 diam. a, Anterior adductors; d, posterior adductors; c, protractor sliding muscles; e, cardinal muscle; r, o, retractor sliding muscles.

acadiensis, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 144, Up. Sil.

anna, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 57, Niagara Gr.

aurora, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 30, Schoharie Grit.

bella, Billings, 1874, Pal. Foss., vol. 2, p. 15, passage beds between Up. Sil. and Devonian.

bordeni, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 187, Up. Held. Gr.

carbonaria, Whitfield, 1882, Desc. New Spec. Foss., from Ohio, p. 229, Coal Meas.

corrugata, Hall, 1843, (*Orbicula corrugata*), Geo. Rep. N. Y., p. 109, Niagara Gr.

crenistriata, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 78, and Pal. N. Y., vol. 4, p. 28, Ham. Gr.

deformata, Hall, 1847, (*Orbicula deformata*), Pal. N. Y., vol. 1, p. 23, Chazy Gr. Is it a *Crania*?

dentata, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 16, Niagara Gr.

dyeri, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 13, Hud. Riv. Gr.

eccentrica, Emmons, 1856, (*Orbicula eccentrica*), Am. Geol., p. 112, Up. Taconic.

famelica, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 236, Chemung Gr.

gracilis, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 17, Niagara Gr.

granulosa, Winchell, 1880, 8th Rep. Geo. Sur. Minn., p. 63, Trenton Gr.

gregaria, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 31, Ham. Gr.

hamiltoniae, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 77, and Pal. N. Y., vol. 4, p. 27, Ham. Gr.

laelia, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 220, Hud. Riv. Gr.

leoni, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 78, and Pal. N. Y., vol. 4, p. 30, Chemung Gr.

modesta, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 118, Up. Coal Meas.

multipunctata, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 13, Hud. Riv. Gr.

pannosa, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 17, Niagara Gr.

parallela, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 98, Hud. Riv. Gr.

percarinata, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 98, Hud. Riv. Gr.

permiana, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 395, Permian Gr.

prima, Owen, 1852, (*Orbicula prima*), Geo. Sur. Iowa, Wis., and Minn., p. 583, Potsdam Gr.

radicans, Winchell, 1866, Rep. Low. Peninsula Mich., p. 92, Ham. Gr.

FIG. 553.—*Crania dyeri*.

FIG. 554.—*Crania multipunctata*.

- reposita, White, 1866, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 8, Ham. Gr.
 reticularis, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 280, Hud. Riv. Gr.
 rowleyi, Gurley, 1883, New Carb. Foss. Kinderhook Gr. Not defined and published as required by the rules of nomenclature.
 scabiosa, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 220, Hud. Riv. Gr.
 setifera, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 209, Niagara Gr.
 setigera, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 220, Trenton Gr.
 sheldoni, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 8, Ham. Gr.
 siluriana, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 208, Niagara Gr.
 socialis, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 99, Hud. Riv. Gr.
 spinigera, Hall, 1879, Desc. New Spec. Foss., p. 13, and 11th Rep. Geo. and Nat. Hist. Ind., p. 283, Niagara Gr.
 trentonensis, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 219, Trenton Gr.
 truncata, Emmons, 1856, (Orbicula truncata,) Am. Geol., p. 200, Trenton Gr.
 CRYPTONELLA, Hall, 1861, 14th Rep. N. Y. St. Mus. Nat. Hist., p. 102. [Sig. a little cavity.] Equilateral, inequivalve, elongate oval or ovoid; valves unequally convex, no mesial fold or sinus; ventral valve with beak extended or incurved, perforate; foramen terminal; punctate smooth or with concentric striae; articulating by teeth and sockets; dental lamellae of the ventral valve extending downward into the cavity of the shell; crura extend in a long recurved loop, with long processes into the ventral valve, between which and the apex they are united by a transverse band. Type C. rectirostra.
 calvini, Hall & Whitfield, 1870, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 239, Chemung Gr.
 circula, Walcott, 1885, Monogr. U. S. Geo. Sur., p. 163, Devonian.
 eudora, Hall, 1867, Pal. N. Y., vol. 4, p. 398, Chemung Gr.
 iphis, Hall, 1867, Pal. N. Y., vol. 4, p. 396, Up. Held. Gr.
 lens, Hall, 1860, (Terebratula lens,) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 89, Up. Held. Gr.
 lincklaeni, Hall, 1860, (Terebratula lincklaeni,) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 88, Ham. Gr.
 pinonensis, Walcott, 1885, Monogr. U. S. Geo. Sur., p. 163, Devonian.
 planirostra, Hall, 1860, (Terebratula planirostra,) 13th Rep. N. Y. St. Mus. Nat.

FIG. 556. — *Cryptonella lincklaeni*. Dorsal and profile views.

FIG. 555. — *Crania reticularis*.

- Hist., p. 89, and Pal. N. Y., vol. 4, p. 395, Ham. Gr.
 rectirostra, Hall, 1860, (Terebratula rectirostra,) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 88, and Pal. N. Y., vol. 4, p. 394, Ham. Gr.
 CYRTIA, Dalman, 1827, Kongl. Vet. Acad. Handl., p. 93. [Ety. *kyrtia*, a fishing basket.] Shell somewhat trigonal, valves convex, hinge-line nearly as long as the width of the shell, articulating by teeth and sockets; ventral valve deep, more or less pyramidal, beak straight or slightly recurved, area wide and triangular, fissure covered by a convex pseudodeltidium, generally perforated close to the beak by a circular foramen, a longitudinal depression in the deltidium sometimes shows, at the extremity a circular aperture for the passage of pedicle muscular fibers; dorsal valve less convex; a mesial longitudinal septum, in the ventral valve, extends from the fissure to near the margin, to the sides of which the dental plates converge, and are united after having formed the fissure walls. Type C. exporrecta.
 acutirostra, see *Cyrtina acutirostra*.
 biplicata, see *Cyrtina biplicata*.
 curvilineata, see *Cyrtina curvilineata*.
 dalmani, see *Cyrtina dalmani*.
 exporrecta, Wahlenberg, 1821, Nova. Acta. Regie. Soc. Sci., vol. 8, p. 64, and 24th Rep. N. Y. St. Mus. Nat. Hist., p. 183, Niagara Gr.
 exporrecta var. arrecta, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 183, Niagara Gr.
 hamiltonensis, see *Cyrtina hamiltonensis*.
 missouriensis, see *Cyrtina missouriensis*.
 myrtea, Billings, 1862, Pal. Foss., vol. 1, p. 165, Mid Sil.
 occidentalis, see *Cyrtina occidentalis*.
 rostrata, see *Cyrtina rostrata*.
 triquetra, see *Cyrtina triquetra*.
 umbonata, see *Cyrtina umbonata*.
 CYRTINA, Davidson, 1853, Monog. Brit. Carb. Brach., p. 66. [Ety. the diminutive of *Cyrtia* is *Cyrtidium*, but the author said he preferred bad (Greek to a long name.) Spirifer-like shells; valves very unequal, ventral being extremely elevated, with high area and narrow fissure, closed by a pseudodeltidium; dental plates converge from the inner margins of the fissure, and, uniting, form a septum to the bottom of the internal cavity, thus dividing it into two parts; shell punctate. Type C. heteroclyta.
 acutirostra, Shumard, 1855, (Cyrtia acutirostra,) Geo. Rep. Mo., p. 204, Waverly or Chateau Gr.
 affinis, Billings, 1874, Pal. Foss., vol. 2, p. 49, Gaspe No. 8, Devonian.

FIG. 557. — *Cyrtina exporrecta*.

billin
Sci.
biplic
10th
165,
crassa
267,
curvil
linea
vol.
270,
dalma
10th
64,
david
Geo.
euphe
Geo.
hamilt
tone
Hist
268,
Ham
hamilt
N. Y.
missou
sour
vol.
occide
denta
vol.
panda,
p. 10
pyrami
mida
ara C
rostrata
Rep.
Orisk
triquet
Geo.
Ham
umbon
Geo.
Ham
Delthyris,
acantho
acuminat
nata.
acuminat
costal
acutiro
arenosa,
audacul
bivalve
bilobata,
brachy
chemung
congesta,
cuspidat
decmpla

billingsi, Meek, 1868, Trans. Chi. Acad. Sci., p. 97, Ham. Gr.

biplicata, Hall, 1857, (Cyrtia biplicata,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 165, Schoharie grit and Cornif. Gr.

crassa, Hall, 1867, Pal. N. Y., vol. 4, p. 267, Up. Held. Gr.

curvilineata, White, 1865, (Cyrtia curvilineata,) Proc. Bost. Soc. Nat. Hist., vol. 9, p. 25, and Pal. N. Y., vol. 4, p. 270, Ham. Gr.

dalmani, Hall, 1857, (Cyrtia dalmani,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 64, Low. Held. Gr.

davidsoni, Walcott, 1885, Monogr. U. S. Geol. Sur., vol. 8, p. 146, Devonian.

euphemis, Billings, 1863, Can. Nat. and Geol., vol. 8, p. 19, Corniferous Gr.

FIG. 558.—*Cyrtina hamiltonensis*. Dorsal, ventral, and side views.

hamiltonensis, Hall, 1857, (Cyrtia hamiltonensis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 166, and Pal. N. Y., vol. 4, p. 268, Schoharie grit, Cornif. and Ham. Grs.

hamiltonensis var. *recta*, Hall, 1867, Pal. N. Y., vol. 4, p. 270, Ham. Gr.

missouriensis, Swallow, 1860, (Cyrtia missouriensis,) Trans. St. Louis Acad. Sci., vol. 1, p. 647, Ham. Gr.

occidentalis, Swallow, 1860, (Cyrtia occidentalis,) Trans. St. Louis Acad. Sci., vol. 1, p. 648, Ham. Gr.

panda, Meek, 1868, Trans. Chi. Acad. Sci., p. 100, Ham. Gr.

pyramidalis, Hall, 1852, (Spirifer pyramidalis,) Pal. N. Y., vol. 2, p. 266, Niagara Gr.

rostrata, Hall, 1857, (Cyrtia rostrata,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 64, Oriskany sandstone.

triquetra, Hall, 1858, (Cyrtia triquetra,) Geo. Rep. Iowa, vol. 1, pt. 2, p. 513, Ham. Gr.

umbonata, Hall, 1858, (Cyrtia umbonata,) Geo. Rep. Iowa, vol. 1, pt. 2, p. 512, Ham. Gr.

Delthyris, Dalman, 1827, syn. for *Spirifera*.

acanthoptera, syn. for *Spirifera* *disjuncta*.

acuminata, Conrad, see *Spirifera* *acuminata*.

acuminata, Hall, syn. for *Spirifera* *mesacostalis*.

acutilirata, see *Orthis* *acutilirata*.

arenosa, see *Spirifera* *arenosa*.

audacula, see *Spirifera* *audacula*.

bialveata, see *Spirifera* *bialveata*.

bilobata, see *Orthis* *bilobata*.

brachynota, see *Spirifera* *brachynota*.

chemungensis, syn. for *Spirifera* *disjuncta*.

congesta, see *Spirifera* *congesta*.

cuspidata, syn. for *Spirifera* *disjuncta*.

decomplicata, see *Spirifera* *decomplicata*.

deltoides, syn. for *Orthis* *lynx*.

disjuncta, see *Spirifera* *disjuncta*.

duodenaria, see *Spirifera* *duodenaria*.

dupplicata, see *Spirifera* *dupplicata*.

euruteines, see *Spirifera* *euruteines*.

expansa, see *Pterotheca* *expansa*.

fimbriata, see *Spirifera* *fimbriata*.

granulifera, see *Spirifera* *granulifera*.

granulosa, see *Spirifera* *granulosa*.

inermis, see *Spirifera* *disjuncta*.

laevis, see *Spirifera* *laevis*.

macronota, see *Spirifera* *macronota*.

macropleura, see *Spirifera* *macropleura*.

medialis, see *Spirifera* *medialis*.

mesacostalis, see *Spirifera* *mesacostalis*.

mesastrialis, see *Spirifera* *mesastrialis*.

microptera, syn. for *Orthis* *lynx*.

mucronata, see *Spirifera* *mucronata*.

niagarensis, see *Spirifera* *niagarensis*.

pachyptera, see *Spirifera* *pachyptera*.

perlata, see *Spirifera* *disjuncta*.

prolata, see *Spirifera* *prolata*.

prora, see *Spirifera* *prora*.

radiata, see *Spirifera* *radiata*.

ruricosta, see *Spirifera* *ruricosta*.

rugatina, see *Spirifera* *rugatina*.

sculptilis, see *Spirifera* *sculptilis*.

staminea, see *Spirifera* *staminea*.

triloba, see *Spirifera* *triloba*.

undulata, see *Spirifera* *undulata*.

varica, see *Orthis* *varica*.

ziczac, see *Spirifera* *ziczac*.

Dicellomus, Hall, 1873, 23d Rep. N. Y. St.

Mus. Nat. Hist., p. 246. A generic name proposed for the reception of *Obolella* *crassa* and *O. polita*, without distinguishing the generic characters.

Dicraniscus, Meek, syn. for *Triplisia*.

ortoni, see *Triplisia* *ortoni*.

DIGNOMIA, Hall, 1873, 23d Rep. N. Y. St.

Mus. Nat. Hist., p. 245. [Ety. *di*, from;

dia, twice; *gnoma*, a sign.] Lingula-like shells having a longitudinal septum in one or both valves. Type *D. alveata*.

alveata, Hall, 1873, 23d Rep. N. Y. St.

Mus. Nat. Hist., p. 245, Ham. Gr.

DINOBOLUS, Hall, March, 1871, 23d Rep.

N. Y. St. Mus. Nat. Hist., p. 247. [Ety.

di, twice; *Obolus*, a genus of shells.]

Shell subcircular, valves thick; umbo

of the ventral valve slightly prominent;

area wider than long; platform sinu-

ated, widely V-shaped; crescent prom-

inently marked in crown and sides;

hinge moderately thick, edge rounded,

with a pair of subcardinal scars in front

of the cardinal facet; umbo of the

brachial valve tumid; platform trilobed;

outer margins raised; antemedian por-

tion rounded, projecting, and terminat-

ing in a median plate; crescent marked

linear scar on the hinge; arching for-

ward in front of the cardinal facet; an

indentation on the inner border of its

sides near the hinge, another further

forward; outer border a fine line; sub-

cardinal scar in the umbonal cavity;

rhomboidal, postmedian scar in front of

the latter. Type *D. conradi*.

FIG. 557.—*Cyrtia eximiosa*.

eximiosa, Hall, 1857, (Cyrtia eximiosa,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 166, and Pal. N. Y., vol. 4, p. 268, Schoharie grit, Cornif. and Ham. Grs.

hamiltonensis, Hall, 1857, (Cyrtia hamiltonensis,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 166, and Pal. N. Y., vol. 4, p. 268, Schoharie grit, Cornif. and Ham. Grs.

missouriensis, Swallow, 1860, (Cyrtia missouriensis,) Trans. St. Louis Acad. Sci., vol. 1, p. 647, Ham. Gr.

occidentalis, Swallow, 1860, (Cyrtia occidentalis,) Trans. St. Louis Acad. Sci., vol. 1, p. 648, Ham. Gr.

panda, Meek, 1868, Trans. Chi. Acad. Sci., p. 100, Ham. Gr.

pyramidalis, Hall, 1852, (Spirifer pyramidalis,) Pal. N. Y., vol. 2, p. 266, Niagara Gr.

rostrata, Hall, 1857, (Cyrtia rostrata,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 64, Oriskany sandstone.

triquetra, Hall, 1858, (Cyrtia triquetra,) Geo. Rep. Iowa, vol. 1, pt. 2, p. 513, Ham. Gr.

umbonata, Hall, 1858, (Cyrtia umbonata,) Geo. Rep. Iowa, vol. 1, pt. 2, p. 512, Ham. Gr.

Delthyris, Dalman, 1827, syn. for *Spirifera*.

acanthoptera, syn. for *Spirifera* *disjuncta*.

acuminata, Conrad, see *Spirifera* *acuminata*.

acuminata, Hall, syn. for *Spirifera* *mesacostalis*.

acutilirata, see *Orthis* *acutilirata*.

arenosa, see *Spirifera* *arenosa*.

audacula, see *Spirifera* *audacula*.

bialveata, see *Spirifera* *bialveata*.

bilobata, see *Orthis* *bilobata*.

brachynota, see *Spirifera* *brachynota*.

chemungensis, syn. for *Spirifera* *disjuncta*.

congesta, see *Spirifera* *congesta*.

cuspidata, syn. for *Spirifera* *disjuncta*.

decomplicata, see *Spirifera* *decomplicata*.

deltoides, syn. for *Orthis* *lynx*.

disjuncta, see *Spirifera* *disjuncta*.

duodenaria, see *Spirifera* *duodenaria*.

dupplicata, see *Spirifera* *dupplicata*.

euruteines, see *Spirifera* *euruteines*.

expansa, see *Pterotheca* *expansa*.

fimbriata, see *Spirifera* *fimbriata*.

granulifera, see *Spirifera* *granulifera*.

granulosa, see *Spirifera* *granulosa*.

inermis, see *Spirifera* *disjuncta*.

laevis, see *Spirifera* *laevis*.

macronota, see *Spirifera* *macronota*.

macropleura, see *Spirifera* *macropleura*.

medialis, see *Spirifera* *medialis*.

mesacostalis, see *Spirifera* *mesacostalis*.

mesastrialis, see *Spirifera* *mesastrialis*.

microptera, syn. for *Orthis* *lynx*.

mucronata, see *Spirifera* *mucronata*.

niagarensis, see *Spirifera* *niagarensis*.

pachyptera, see *Spirifera* *pachyptera*.

perlata, see *Spirifera* *disjuncta*.

prolata, see *Spirifera* *prolata*.

prora, see *Spirifera* *prora*.

radiata, see *Spirifera* *radiata*.

ruricosta, see *Spirifera* *ruricosta*.

rugatina, see *Spirifera* *rugatina*.

sculptilis, see *Spirifera* *sculptilis*.

staminea, see *Spirifera* *staminea*.

triloba, see *Spirifera* *triloba*.

undulata, see *Spirifera* *undulata*.

varica, see *Orthis* *varica*.

ziczac, see *Spirifera* *ziczac*.

canadensis, Billings, 1857, (Obolus canadensis,) Rep. of Progr. Geo. Sur. of Can., p. 189, and Can. Nat., vol. 6, p. 222, Black Riv. Gr.
 conradi, Hall, 1868, (Obolus conradi,) 20th Rep. N. Y. St. Mus. Nat. Hist., p. 368, Niagara Gr.
 galtensis, see Trimerella galtensis.
 magnificus, Billings, 1872, (Obolus magnificus,) Canadian Naturalist, vol. 6, p. 330, Black Riv. Gr.
 parvus, Whitfield, 1882, Geo. Wis., vol. 4, p. 347, Galena Gr.
 DISCINA, Lamarck, 1819, Hist. Nat. Anim. sans Vert., vol. 6, p. 236. [Ety. *discus*, a flat, round plate; the termination *inus*, implying resemblance.] Circular, longitudinally or transversely oval; dorsal valve conical, with apex inclined toward the posterior margin; ventral valve opercular, flat, or partly convex, perforated by a narrow, oval, longitudinal slit, reaching to near the posterior margin, and placed in the

FIG. 559.—*Discina ostreoides*.

middle of an oval depressed disk; surface smooth, striated from the apex to the margin, or having concentric lines of growth produced in foliaceous expansions; structure horny, and perforated by minute tubuli. Type *D. ostreoides*.

acadica, see *Stenotheca acadica*.

allegahania, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 77, and Pal. N. Y., vol. 4, p. 25, Chemung Gr.

ampla, Hall, 1867, Pal. N. Y., vol. 4, p. 17, Oriskany sandstone. Proposed instead of *D. grandis* of Hall.

✓ capax, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 30, Waverly or Marshall Gr.

capuliformis, McChesney, syn. for *D. nitida*.

circe, Billings, 1862, Pal. Foss., vol. 1, p. 51, Trenton Gr. See remarks on, *D. lamellosa*.

✗ clara, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 56, Niagara Gr.

✓ connata, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 214, Devonian.

conradi, Hall, 1859, Pal. N. Y., vol. 3, p. 161, Low. Held. Gr.

FIG. 560.—*Discina circe*.

convexa, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 221, Coal Meas.

discus, Hall, 1859, Pal. N. Y., vol. 3, p. 159, Low. Held. Gr.

doria, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 26, Ham. Gr.

elmira, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 29, Chemung Gr.

gallaheri, Winchell, 1865, Proc. Acad. Nat. Sci., p. 112, Marshall Gr.

grandis, Vanuxem, 1842, Geo. Rep. 3d Dist. N. Y., p. 152, and Pal. N. Y., vol. 4, p. 17, Cornif. and Ham. Gr.

grandis, Hall, 1859, Pal. N. Y., vol. 3. The name was preoccupied. See *D. ampla*.

humilis, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 25, Marcellus Slate and Ham. Gr.

inutilis, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 130, Potsdam Gr.

lamellosa, Hall, 1847, (Orbicula lamellosa.) The name was preoccupied by Broderick in 1833. Billings has described it as *D. circe*.

lodensis, Vanuxem, 1842, (Orbicula lodensis,) Geo. Rep. 3d Dist. N. Y., p. 168, and Pal. N. Y., vol. 4, p. 22, Genesee Slate.

manhattanensis, Meek and Hayden, 1859, Proc. Acad. Nat. Sci., p. 25, Coal Meas.

FIG. 561.—*Discina ostreoides*, 2 diam. u, Umbo; f, foramen; d, disk; a, anterior adductors; p, posterior adductors; c, c, protractor sliding muscles; r, retractor muscles.

marginalis, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 70, and Geo. Wis., vol. 4, p. 325, Ham. Gr.

media, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 27, Ham. and Chemung Grs.

meekana, Whitfield, 1882, Desc. New Spec. Foss. from Ohio, p. 228, Coal Meas.

microscopica, Shumard, 1861, Am. Jour. Sci. and Arts, vol. 32, p. 213, Potsdam Gr.

minuta, Hall, 1843, (Orbicula minuta,) Geo. Rep. 4th Dist. N. Y., p. 180, and Pal. N. Y., vol. 4, p. 16, Marcellus Shale.

missouriensis, Shumard, 1858, Trans. St. Louis Acad. Sci., Coal Meas. Syn. for *D. nitida*.

Acc. to Hall, Pal. N. Y., Vol. 8. Brachiopoda, Expl. of Pl. IV E, R. is *Orbiculoides* (*Schizotreta*?) *tenuilamellata*, Hall.

St. Louis
Meas.
vol. 3, p.
Y. St. Mus.
N. Y. St.
ung. Gr.
roc. Acad.
r.
b. Rep. 3d
N. Y. vol.
ir.
Y., vol. 3.
d. See D.
N. Y. St.
ce. us Slate
p. N. Y. St.
dam Gr.
lamellosa,
by Broder-
described it

Orbicula lo-
t. N. Y., p.
p. 22, Gen-
ayden, 1859,
Coal Meas.

men; d, disk;
rotractor slid-

Ann. Rep.
1 Geo. Wis.,

p. N. Y. St.
m. and Che-

Desc. New
8, Coal Meas.
1, Am. Jour.
p. 213, Pots-

ula minuta,
t. 180, and Pal.

lus Shale.
3, Trans. St.

bas. Syn. for

2. IV E, Ric

neglecta, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 29, Chemung Gr.
newberryi, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 30, Waverly Gr.
nitida, Phillips, 1836, (Orbicula nitida,) Geo. of York., vol. 2, p. 221, and Geo. Sur. Ill., vol. 5, p. 572, Coal Meas.
patellaris, Winchell, 1863, Proc. Acad. Nat. Sci., p. 4, Waverly or Marshall Gr.
pelopea, Billings, 1862, Pal. Foss., vol. 1, p. 52, Trenton Gr.

pleurites, Meek, 1875, Ohio Pal., vol. 2, p. 278, Waverly Gr.

randalli, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 25, Ham. Gr.

saffordi, Winchell, 1869, Geo. of Tenn., and, in 1870, Proc. Am. Phil. Soc., p. 248, Marshall Gr.

seneca, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 26, Ham. Gr.

sublamellosa, Ulrich, 1878, Jour. Clin. Soc. Nat. Hist., p. 97. Probably the east of a Trematis.

subtrigonalis, McChesney, 1865, Desc. New Pal. Foss., Coal Meas. Not recognized.

tennilamellata, Hall, 1852, (Orbicula tennilamellata,) Pal. N. Y., vol. 2, p. 250, Niagara Gr.

tennilamellata var. subplana, Hall, 1860, Can. Nat. and Geol., vol. 5, p. 144, Up. Sil.

tennilineata, Meek & Hayden, 1859, Proc. Acad. Nat. Sci., p. 25, Coal Meas.

tennistriata, Ulrich, 1878, Jour. Clin. Soc. Nat. Hist., p. 96. Probably the east of a Trematis.

trigonalis, syn. for D. subtrigonalis.

truncata, see Schizobolus truncatus.

tullia, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 28, Tully limestone.

vanuxemi, Hall, 1859, Pal. N. Y., vol. 3, p. 162, Water-lime or Low. Held. Gr.

varsoviensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 23, and Geo. Sur. Ill., vol. 8, p. 102, Keokuk Gr.

EATONIA, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 90, and 12th Rep., p. 35. [Ety. proper name.] Oval, ovoid, subcircular, elongate, or transverse;

valves very unequally convex; mesial fold and sinus; beak of ventral valve small, perforate, closely incurved over the umbo of the dorsal valve; two teeth in the ventral, with corresponding sockets in the dorsal valve; a prominent bifurcating cardinal process and four crural processes in the dorsal valve distinguish this genus. Type E. medialis.

eminens, Hall, 1857, N. Y. St. Mus. Nat. Hist., p. 92, and Pal. N. Y., vol. 3, p. 242, Low. Held. Gr.

medialis, Vanuxem, 1842, (Atrypa mediaialis,) Geo. Rep. 3d Dist. N. Y., p. 121, and Pal. N. Y., vol. 3, p. 241, Low. Held. Gr.

peculiaris, Conrad, 1841, (Atrypa peculiaris,) Ann. Rep. N. Y., p. 58, and

Pal. N. Y., vol. 3, p. 244, Oriskany and Low. Held. Gr.

pumila, Hall, 1859, Pal. N. Y., vol. 3, p. 437, Oriskany sandstone.

singularis, Vanuxem, 1842, (Atrypa singularis,) Geo. Rep. 3d Dist. N. Y., p. 120, and Pal. N. Y., vol. 3, p. 243, Low. Held. Gr.

sinuata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 91, and Pal. N. Y., vol. 3, Oriskany sandstone.

whitfieldi, Hall, 1859, Pal. N. Y., vol. 3, p. 437, Oriskany sandstone.

EICHWALDIA, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 190. [Ety. proper name.] Ovate or subtrigonal, with or without mesial fold and sinus; ventral valve obscurely perforate on the umbo; apex acute and entire; space beneath occupied by an imperforate concave plate; interior of the rostral cavity containing a transverse septum; dorsal valve with a slender cardinal process and a very elevated medio-longitudinal septum; valves articulated in a narrow groove in the dorsal valve; surface of the shell reticulate, solid, and fibrous beneath. Type E. subtrigonalis.

anticostiensis, Billings, 1866, Catal. Sil. Foss. Antic., p. 10, Ind. Riv. Gr.

concinna, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

corallifera, Hall, 1852, (Atrypa corallifera,) Pal. N. Y., vol. 2, p. 281, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

FIG. 503.—Eatonia singularis.

FIG. 504.—Eichwaldia reticulata.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

regarded this shell as identical with E. capewelli, which was described in 1848, in Bull. Soc. Geol. France, vol. 3.

gibbosa, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 319, Niagara Gr.

reticulata, Hall, 1863, (Rhynchonella?) reticulata, Trans. Alb. Inst., vol. 4, p. 217, Niagara Gr.

Davidson said a syn. for E. capewelli.

subtrigonalis, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 192, Black Riv. Gr.

FIG. 502.—Eatonia medialis. Anterior view.

FIG. 505.—Eichwaldia subtrigonalis. Dorsal, ventral, side, front, and apex views.

ELKANIA, Ford, 1886, Am. Jour. Sci. and Arts, 3d ser., vol. 32, p. 325. [Ety. proper name.] Shell thin, calcareous, inarticulate, longitudinally ovate or subcircular, convex; ventral valve, with solid beak and minute-grooved area; muscular scars, six in each valve; beneath the rostrum a spoon-shaped pit separates the scars. Type *E. desiderata*.

desiderata, Billings, 1862, (*Obolella desiderata*.) Pal. Foss., vol. 1, p. 69, Up. Taconic.

EUMETRIA, Hall, 1864, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 59. Shell longitudinally suboval; striated, without mesial fold and sinus; structure punctate; beak of the ventral valve incurved; hinge area contracted; foramen large; internal spire as in *Athyris*; dorsal valve in the form of a pectinoid shell, with diverging lamellæ, which extend beneath the cardinal area of the ventral valve on either side of the center; processes extending into the cavity of the dorsal valve, gradually converge, and are united by a transverse concave septum. Type *E. vera*.

prima, White, 1862, (*Acambona prima*.) Proc. Bost. Soc. Nat. Hist., vol. 9, p. 27, Burlington Gr.

vera, Hall, 1858, (*Retzia vera*.) Geo. Sur. Iowa, p. 704, Kaskaskia Gr.

FIG. 566.—*Eumetria verneuillana*.

vera var. *costata*, Hall, 1858, (*Retzia vera* var. *costata*.) Geo. Sur. Iowa, p. 704, Kaskaskia Gr.

verneuillana, Hall, 1858, (*Retzia verneuillana*.) Trans. Alb. Inst., vol. 4, p. 19, and Geo. Sur. Iowa, p. 657, Warsaw Gr.

GLASSIA, Davidson, 1881, Lond. Geo. Mag., vol. 8, p. 11. [Ety. proper name.] Shell ovate; spiral coils in the dorsal valve for the support of the brachial appendages connected by a loop as in *Atrypa*; lamellæ converge downward like the letter V, with the extremities turned slightly upward before uniting; principal coils face the lateral margins; ends of the spirals meet in the center of the shell; spirals consist of four or five compressed coils. Type *G. obovata*.

headi, Meek, 1873, (*Zygospira headi*.) Ohio Pal., vol. 1, p. 127, Hud. Riv. Gr.

Goniocelia, Hall, syn. for *Pentagonia*.

GYPIDULA, Hall, 1867, Pal. N. Y., vol. 4, p. 373. [Ety. *gyps*, vulture; in allusion to the strongly incurved beak.] Short, gibbous or ventricose, ventral valve much the larger, with or without mesial fold; a large fissure, and elongate, much incurved, trough-shaped pit; dorsal valve depressed in front; area on both valves, that of the ventral striated as in *Spirifera*; lamellæ of dorsal valve separate and diverging. Type *G. occidentalis*.

levisculæ, Hall, 1867, Pal. N. Y., vol. 4, p. 381, Devonian.

munda, Calvin, 1878, Bull. U. S. Geo. Sur., vol. 4, No. 3, p. 730, Low. Devonian.

obsolescens, see *Pentamerella obsolescens*, *occidentalis*, Hall, 1858, (*Pentamerus occidentalis*.) Geo. Rep. Iowa, vol. 1, pt. 2, p. 514, Ham. Gr.

unguliformis, Ulrich, 1880, Cont. to Am. Pal., p. 28, Niagara Gr.

Hemipronites, Pander, 1830. This name, not having been defined, has been superseded by *Streptorhynchus*, if the two names refer to the same form.

americanus, see *Streptorhynchus americanus*.

HINDELLA, Davidson, 1882, Monogr. Brit. Foss., Brachiopoda, vol. 5, p. 130. [Ety. proper name.] Shell elongate, ovate; about six coils in each spiral; apices directed laterally; stems attached to the hinge plate, and extending into the interior, they are abruptly bent backward, and then form a broad, rounded curve, facing the bottom of the dorsal valve; when they reach the front they give off a semicircular loop, having a spike-like process at the top, directed toward the beak. Type *H. umbonata*.

umbonata, Billings, 1865, (*Athyris umbonata*.) Pal. Foss., vol. 1, p. 144, Mid. Sil., Anticosti Div. 1.

Hipparionyx, Vanuxem, 1842, Geo. 3d Dist. N. Y., p. 124, syn. for *Orthis*. The genus was founded on a cast.

constrictilis, syn. for *Atrypa reticularis*.

proximus, see *Orthis proximus*.

similaris, Vanuxem, 1842, Geo. Rep. 3d Dist. N. Y., Oriskany sandstone. Not defined.

IPHIDEA, Billings, 1874, Pal. Foss., vol. 2, p. 76. [Ety. proper name.] Ventral valve

conical, elevated at the beak, hinge-line nearly straight, posterior angles rounded, sides and front nearly uniformly rounded; posterior side with a large false area and a convex pseudodeltidium; dorsal valve semicircular, moderately convex, most elevated at the beak; surface concentrically marked. Type *I. bella*.

bella, Billings, 1872, Can. Nat., vol. 6, p. 477, and Pal. Foss., vol. 2, p. 76, Up. Taconic.

sculptilis, see *Kutorgina sculptilis*.

KONINKIA, Suess, 1853. MS. published by Woodward, 1854, in Manual of Mollusca, p. 231. [Ety. proper name.] Shell circular, inequivalve, compressed; ventral valve convex, with a slight longitudinal depression; beak incurved, with auricular expansions; dorsal valve concave; surface smooth; no area or deltidium; valves inarticulated; mesial ridge in dorsal valve; oral appendages supported by a

FIG. 567.—*Ipheidea bella*.

Y., vol. 4,

U. S. Geo.
Low. Devon.obsolescens,
stamens or-
a, vol. 1, pt.

Cont. to Am.

This name,
t, has been
us, if the two
rm.
echus ameri-Monogr. Brit.
p. 130. [Ety.
ngate, ovate;
spiral; apices
attached to the
into the in-
nt backward,
unded earve.
dorsal valve;
nt they give
ving a spike-
ected toward
nata.Athyris uni-
p. 144, Mid.Geo. 3d Dist.
Orthis. The
ast.

eticularis.

is.
Geo. Rep. 3d
date. Not

oss., vol. 2, p.

Ventral valve

e
ys
s
t

FIG. 567.—Iph-

iden bella.

x
valve semi-
vex, most ele-
concentrically

nat., vol. 6, p.

2, p. 76. Up.

lptilis.

published by

annual of Mol-

proper name.]

valve, com-

convex, with

depression;

icular expan-

cave; surface

idium; valves

edge in dorsal

supported by a

spiral, calcified lamella. Type *K. leon-*
hardi.americana, Swallow, 1863, Trans. St. Louis
Acad. Sci., vol. 2, p. 94, Kaokaakia Gr.
Kutadina, Billings, 1861, Pal. Foss., vol. 1,p. 3. [Ety. proper name.] Shell more
or less subquadrate in out-
line; hinge-line straight;
sides slightly convex, an-
terior angles rounded,
front slightly convex; sur-
face with concentric ridges
terminating on the car-
dinal edges, and the course
conforming to the margin
of the shell, and sometimes FIG. 568.—*Ku-*
tordina pan-
nulla. En-
larged 6
diam.with lines radiating from the beak to
the margin; ventral valve tumid, most
convex about the middle, beak slightly
depressed; cardinal edges straight or
slightly concave and diverging from the
beak at an obtuse angle; dorsal valve
less convex, most elevated at the beak,
and along the middle there is a shallow
concavity extending to the front mar-
gin. Type *K. cingulata*.cingulata, Billings, 1861, Pal. Foss., vol. 1,
p. 8, Up. Taconic.labradorica, Billings, 1861, (Obolus lubra-
doricus,) Pal. Foss., vol. 1, p. 6, Up.
Taconic.latourensis, Matthew, 1885, Trans. Roy.
Soc. Can., p. 42, St. John Gr.minutissima, Hall & Whitfield, 1877, U. S.
Geo. Expl. 40th parallel, syn. for *K.*
sculptilis.pannulla, White, 1874, (Trematis pannulus,)
Rep. Invert. Foss., p. 6, and Geo. Sur. W.
100th Mer., vol. 4, p. 36, Up. Taconic.prospectensis, Walcott, 1885, Monogr.
U. S. Geo. Sur. Terr., vol. 8, p. 19, Up.
Taconic.pterinoides, Matthew, 1885, Trans. Roy.
Soc. Can., p. 43, St. John Gr.sculptilis, Meek, 1873, (Iphidesculptilis,)
6th Ann. Rep. U. S. Geo. Sur. Terr., p.
479, and Monogr. U. S. Geo. Sur. Terr.,
vol. 8, p. 20, Potsdam Gr.stissingensis, Dwight, 1889, Am. Jour.
Sci. and Arts, 3d ser., vol. 38, p. 145,
Up. Taconic.whitfieldi, Walcott, 1885, Monogr. U. S.
Geo. Sur. Terr., vol. 8, p. 18, Up. Ta-
conic.LEPTORHYNCHUS, Hall, 1860, 13th Rep. N. Y.
St. Mus. Nat. Hist., p. 75. [Ety. *leios*,
smooth; *rhynchos*, beak.] Ovate, cir-
cular or transverse, valves unequally
convex; mesial fold and sinus, which
are plicated; articulating by teeth and
sockets; apex of ventral valve perfor-
ate, two diverging lamellæ extend
into and join the sides or bottom of the
rostral cavity; muscular impressions oc-
cupy a narrow triangular cavity below
the dental lamellæ; median septum in
the dorsal valve extending half the
length of the shell; hinge plates, nar-
row, strong processes, embraced by thecurving teeth of the opposite valve; sub-
stance fibrous. Type *L. quadricostatum*.
dubium, Hall, 1867, Pal. N. Y., vol. 4,
p. 364, Marcellus Shale.globuliforme, Vanuxem, 1842, (Atrypa
globuliformis,) Geo. 3d Dist. N. Y., p.
182, and Pal. N. Y., vol. 4, p. 364, Che-
mung Gr.hecate, Clarke, 1885, Bull. U. S. Geo. Sur.
No. 16, p. 31, Genesee Shales.huronense, Nicholson, 1874, Geo. Mag.
Lond., n. s., vol. 1, p. 120, Ham. Gr.iris, Hall, 1867, Pal. N. Y., vol. 4, p. 360,
Chemung Gr.kelloggi, Hall, 1867, Pal. N. Y., vol. 4, p.
361, Chemung Gr.laura, Billings, May, 1860, (Rhynchonella
laura,) Can. Jour., vol. 5, p. 273, Ham. Gr.limiture, Vanuxem, 1842,
(Orthis limitaris,) Geo.3d Dist. N. Y., p. 146,
and Pal. N. Y., vol. 4, p.356, (Atrypa limitaris,)
4th Dist. N. Y., Mar-
cellus Shale.mesacostale, Hall, 1843,
(Atrypa mesacostalis,) Geo. 4th Dist. N. Y., p.

64, and Pal. N. Y., vol.

4, p. 362, Chemung Gr.

multicosta, Hall, 1860, 13th Rep. N. Y.
St. Mus. Nat. Hist., p. 85, and Pal. N. Y.,
vol. 4, p. 358, Ham. Gr.mysia, Hall, 1867, Pal. N. Y., vol. 4, p.
357, Marcellus Shale.nevadense, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 157, Devonian.newberryi, Hall & Whitfield, 1873, 23d
Rep. N. Y. St. Mus. Nat. Hist., p. 240,
Waverly Gr.quadricostatum, Vanuxem, 1842, (Orthis
quadricostata,) Geo. 3d Dist. N. Y., p.168, and Pal. N. Y., vol. 4, p. 357, Gen-
esee Slate.sesquiplicatum, Winchell, 1866, Rep. Low.
Penin. Mich., p. 95, Ham. Gr.sinuatum, Hall, 1867, Pal. N. Y., vol. 4, p.
362, Chemung Gr.LEPTENA, Dalman, 1827, Kongl. Vet. Acad.
Handl., p. 93. [Ety. *leptos*, thin.] Shellthin, semicircular, transversely elon-
gated, smooth or finely striated; hinge-
line straight, ventral valve convex, fis-
sure partly covered by a deltidium; beak
inconspicuous, sometimes per-
forated; cardinal area narrow; muscular
scars small, not marginal; adductor
scars close to a mesial ridge, while the
cardinal scars are on either side; vas-
cular impressions radiating; dorsal
valve concave; socket ridges large, car-
dinal process small, multifid, connate
with their bases; adductor impressions
large, produced, elongated, and bor-
dered by ridges; area on both valves.Type *L. transversalis*.*alternata*, see *Strophomena alternata*.*alternistriata*, see *Strophomena alterni-*
striata. FIG. 569.—*Leio-*
rhynchus
quadricosta-
tum.

anuloga, see *Strophomena anuloga*.
aspera, James, syn. for *L. sericea*.
barabuenensis, Winchell, 1864, (*Orthis barabuenensis*,) Am. Jour. Sci. and Arts, 2d ser., vol. 37, p. 229, and Geo. Wis., vol. 4, p. 171, Potsdam Gr.

bipartita, see *Strophomena bipartita*.
camerata, see *Strophomena camerata*.
concava, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 47, and Pal. N. Y., vol. 3, p. 197, Low. Held. Gr.

decipiens, Billings, 1862, Pal. Foss., vol. 1, p. 74, Quebec Gr.

deflecta, see *Streptorhynchus deflectum*.

delloidea, see *Strophomena delloidea*.

depressa, see *Strophomena depressa*.

fasciata, see *Strophomena fasciata*.

filitexta, see *Streptorhynchus filitextum*.

fragaria, syn. for *Productella subaculeata*.

incrassata, see *Strophomena incrassata*.

indenta, see *Strophodonta indenta*.

laticosta, syn. for *Tropidoleptus carinatus*.

melita, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 208, Potsdam Gr.

membranacea, see *Productella hirsuta*.
mesacosta, Shumard, 1855, Geo. Rep. Mo., p. 205, Trenton Gr.

nasuta, see *Strophomena nasuta*.

nucleata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 47, and Pal. N. Y., vol. 3, p. 419, Oriskany sandstone.

obscura, see *Strophomena obscura*.

orthideia, see *Strophomena orthideia*.

planconvexa, see *Streptorhynchus planconvexum*.

planumbona, see *Streptorhynchus planumbonum*.

plicatella, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 15, Utica Slate Gr.

plicifera, see *Strophomena plicifera*.

profunda, see *Strophodonta profunda*.

prolongata, Foerste, 1885, Bull. Sci. Lab. Denison Univ., p. 79, Niagara Gr.

punctulifera, see *Strophonella punctulifera*.

quadrilatera, syn. for *Strophomena rhomboidalis*.

recta, see *Streptorhynchus rectum*.

rugosa, see *Strophomena rugosa*.

semiovalis, syn. for *L. sericea*.

sericea, Sowerby, 1839, Murch. Sil. Syst., p. 636, and Pal. N. Y., vol. 1, p. 110, Trenton to Clinton Gr.

FIG. 570.—*Leptæna sericea*. Dorsal view, and interior of dorsal valve.

sordida, Billings, 1862, Pal. Foss., vol. 1, p. 73, Quebec Gr.

subquadrata, Hall, 1883, Rep. St. Geol., pl. 46, fig. 32, 33, Low. Held. Gr.

subtenta, see *Streptorhynchus subtentum*.

tenuilineata, see *Strophomena tenuilineata*.

tenuistriata, see *Strophomena tenuistriata*.

transversalis, Wahlenberg, 1821, (*Anomites transversalis*,) Act. Soc. Upsal., vol. 8, p. 64, and Pal. N. Y., vol. 2, p. 256, Anticosti and Clinton Gr.

trilobata, see *Strophomena trilobata*.

vicina, Castelnau, 1843, Syst. Sil., p. 39. Not recognized.

LEPTONOLUS, Hall, 1871, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 226. [*Ety. leptos*, minute; *Obolus*, a genus.] Shell small, ovate, fragile, semiphosphatic, concentrically lined; ventral valve with an area and pedicel groove, muscular scar elevated, subquadrate; dorsal valve with trifid muscular impressions. Type *L. lepis*.

insignis, Hall, 1871, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 227, Utica Slate.

lepis, Hall, 1871, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 226, Utica Slate.

occidentalis, Hall, 1871, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 227, Utica Slate.

LEPTOCHELIA, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 107, and 12th Rep., p. 32. [*Ety. leptos*, minute; *cheilia*, belly; in allusion to the shallow visceral cavity.] Shell inequivalve, variable in form, plicated, usually mesial fold and sinus, substance lamellose or fibrous; ventral valve convex, beak extended, and more or less incurved; foramen terminal, the lower side formed by two deltoid pieces; two strong teeth, denticulated; muscular impressions marking a flabelliform area with a thin median septum, adductor imprints small; dorsal valve flat, concave, or depressed convex; on each side of a strong cardinal process are the deep, oblique, dental fosses, from the inner margins of which the crural processes proceed, supported below by thickened plates, extending obliquely on the border of the muscular impression toward the middle of the shell; muscular impression divided by a low median septum; the crura, in their extension, are united, in a flattened disk, which terminates in an acute point; on the center of the cardinal side a slender process extends downward, and near the junction of the crura two slender processes extend into the cavity of the ventral valve. Type *L. flabellites*.

acutiplicata, Conrad, 1841, (*Atrypa acutiplicata*,) Ann. Rep. N. Y., p. 54, and Pal. N. Y., vol. 4, p. 365, Up. Held. Gr.

concava, see *Caelospira concava*.

dichotoma, see *Caelospira dichotoma*.

disparilis, see *Caelospira disparilis*.

ambriata, Hall, 1859, Pal. N. Y., vol. 3, p. 451, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

coelia flabellites, Conrad, 1841, (*Atrypa coelia flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

flabellites, Conrad, 1841, (*Atrypa flabellites*,) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 3, p. 449, Oriskany sandstone.

FIG. 571.—*Leptæna flabellites*.

821, (Ano-
soc. Upsal.
, vol. 2, p.
n Gr.

obata.
Sil., p. 39.

p. N. Y. St.
Ety. *leptos*,
Shell small,
tic, concent-
re with an
muscular scar
oral valve
sions. Type

p. N. Y. St.
ca Slate.
Y. St. Mus.
te.

Rep. N. Y.
Utica Slate.

ep. N. Y. St.
d 12th Rep.,

ute; *koilin*,
shallow vis-
ivalve, vari-
ually mesial

lamellose or
ex, beak ex-
s incurved;

side formed
two strong
ular impres-
ner area with
actor imprints

ceave, or de-
le of a strong
eep, oblique,
ner margins

sses proceed,
ened plates,
he border of

toward the
ular impres-
ian septum;

n, are united,
terminates in
center of the

ccess extends
junction of
sses extend
entral valve.

Atrypa acuti-
p. 54, and
p. Held. Gr.

va.
otoma.

571. — *Lepto-*
flabellites.

p. 449, Oris-

hemispherica, Sowerby, 1839, (*Atrypa hemispherica*.) Murch. Sil. Syst., p. 639, and Pal. N. Y., vol. 2, p. 74, Clinton Gr.

imbricata, see *Trematospira imbricata*.
internedia, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 144, Up. Sil.

planoconvexa, Hall, 1852, (*Atrypa planoconvexa*.) Pal. N. Y., vol. 2, p. 75, Clinton Gr.

propria, Hall, syn. for *L. flabellites*.
LINGULA, Bruguiere, 1792, Encyc. Meth., tab. 250. [Ety. *lingula*, a little tongue.]

Shell oblong or ovoid, depressed, thin, gaping at each end, rounded or subtruncate in front, pointed at the beaks, consisting of alternate fibrous, corneous, and tubular testaceous, phosphatic laminæ; valves convex, held together by the action of muscles, beak of ventral valve more pointed and prominent than the other; surface smooth or concentrically lined; peduncle long, thick, cylindrical, fleshy, and flexible; there are twelve muscular impressions in the dorsal, and thirteen in the ventral, valve. Type *L. anatina*. No Palæozoic

shell is positively known to agree with this genus in its muscular impressions, and probably none belong to it. Many referred to it belong to *Lingulella*, others to *Lingulepis*, and others, may be, to undefined genera. The external appearance, however, resembles *Lingula*, and for want of material to distinguish internal characters, they are left, provisionally, where the authors of the species left them.

acuminata, Conrad, 1839, Ann. Rep. N. Y., p. 64, Calcif. Gr.

FIG. 572.—*Lingula acuminata*. Various forms; a, b, c, and e are ventral valves; d, dorsal; and f and g are young shells.

acutangula, Roemer, 1852, Kreid. von Texas, p. 90, Silurian.

acutirostra, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 77, and Pal. N. Y., vol. 2, p. 56, Clinton Gr.

æqualis, Hall, 1847, Pal. N. Y., vol. 1, p. 95, Trenton Gr.

albapinensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 108, Devonian.

alveata, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 23, and Pal. N. Y., vol. 4, p. 12, Ham. Gr.

ampla, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 583, Potsdam Gr.

antiqua, Emmons, 1842, Geo. Rep. N. Y., p. 238, and Pal. N. Y., vol. 1, p. 3, Potsdam Gr.

antiquata, Emmons, 1856, Am. Geol., p. 202, Potsdam Gr.

artemis, Billings, 1874, Pal. Foss., vol. 2, p. 14, passage beds between Up. Sil. and Devonian.

FIG. 573.—*Lingula anatina*. aa, Anterior adductors; a, posterior adductor; pp, external protractors; pp, central protractors; rr, anterior retractors; r, posterior retractors; c, capsule of pedicel; n, visceral sheath; o, oesophagus; s, stomach; l, liver; v, vent; h, auricles, etc.

attenuata, Sowerby. The fossil referred by Hall to this species is described by Billings under the name of *L. daphne*.

aurora, see *Lingulella aurora*.

belli, Billings, 1859, Can. Nat. Geo., vol. 4, p. 431, Chazy Gr.

bicarinata, Ringueberg, 1884, Proc. Acad. Nat. Sci., p. 149, Niagara Gr. Not defined so as to be recognized.

billingsana, Whiteaves, 1878, Am. Jour. Sci. and Arts, 3d ser., vol. 16, p. 226, St. John's Gr.

bisulcata, Ulrich, 1889, Am. Geol., vol. 3, p. 380, Utica Slate.

briseis, Billings, 1862, Pal. Foss., vol. 1, p. 48, Trenton Gr.

cabmet, N. H. Winchell, 1885, 13th Ann. Rep. Geo. Sur. Minn., p. 65, Taconic. Probably an *Obolella*.

canadensis, Billings, 1862, Pal. Foss., vol. 1, p. 114, Hud. Riv. Gr.

carbonaria, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 215, Coal Meas.

centrilineata, Hall, 1859, Pal. N. Y., vol. 3, p. 155, Low. Held. Gr.

ceryx, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 19, and Pal. N. Y., vol. 4, p. 5, Schoharie grit.

clintoni, Vanuxem, 1842, Geo. Rep. N. Y., p. 79, and Pal. N. Y., vol. 2, p. 54, Clinton Gr.

cobourgensis, Billings, 1862, Pal. Foss., vol. 1, p. 50, Trenton Gr.

- complanata, Williams, 1882, Proc. A. A. A. S., vol. 30, p. 188, Chemung Gr.
 concentrica, Conrad, 1839, Ann. Rep. N. Y., p. 64, and Geo. Rep. 3d Dist. N. Y., p. 108, Genesee Slate.
 covingtonensis, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 67, Utica Slate.
 crassu, Hall, 1847, Pal. N. Y., vol. 1, p. 98, Trenton Gr.
crawfordsvillensis, Gurley, 1883, New Carb. Foss., p. 2, Keokuk Gr. The publication is not such as to entitle it to recognition.
cuneata, see *Lingulella cuneata*.
 curta, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 266, and Pal. N. Y., vol. 1, p. 97, Utica Slate.
 cuyahoga, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 24, and Pal. N. Y., vol. 4, p. 15, Waverly Gr.
 cyane, Billings, 1865, Pal. Foss., vol. 1, p. 216, Quebec Gr.
 daphne, Billings, 1862, Pal. Foss., vol. 1, p. 50, Trenton Gr. See *L. attenuata*.
 dawsoni, Mathew, 1884, Bull. U. S. Geo. Sur., vol. 2, p. 283, St. John Gr.
 delia, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 22, and Pal. N. Y., vol. 4, p. 12, Ham. Gr.
 densa, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 22, and Pal. N. Y., vol. 4, p. 11, Ham. Gr.
 desiderata, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 19, and Pal. N. Y., vol. 1, p. 6, Up. Held. Gr.
 elderi, Whitfield, 1880, Am. Jour. Sci. and Arts, 3d ser., vol. 19, p. 472, and Geo. Wis., vol. 4, p. 345, Trenton Gr.
elegantula, syn. for *Lingula quadrata*.
elliptica, Hall, 1843, Geo. Rep. 4th Dist. N. Y. The name was preoccupied by Phillips in 1836. See *L. subelliptica*.
elliptica, Emmons, 1856, Am. Geol. The name was preoccupied.
 elongata, Hall, 1847, Pal. N. Y., vol. 1, p. 97, Trenton Gr.
 exilis, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 77, and Pal. N. Y., vol. 4, p. 7, Marcellus Shale.
 eva, Billings, 1861, Can. Nat. Geo., vol. 6, p. 150, Black Riv. Gr.
 forbesi, Billings, 1862, Pal. Foss., vol. 1, p. 115, Hud. Riv. and Mid. Sil. Grs.
 gibbosa, Hall, 1879, Desc. New Spec. Foss., p. 13, and 11th Rep. Geo. and Nat. Hist. Ind., p. 284, Niagara Gr.
 halli, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 8, Burlington Gr.
 hurlbuti, Winchell, 1880, Geo. Sur. Minn., 8th Rep., p. 62, Galena Gr.
 huronensis, Billings, 1859, Can. Nat. Geo., vol. 4, p. 433, Chazy and Black Riv. Grs.
 ingens, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 56, Niagara Gr.
 insularis, Billings, 1866, Catal. Sil. Foss. Antic., p. 40, Anticosti Gr.
 iole, Billings, 1865, Pal. Foss., vol. 1, p. 215, Quebec Gr.
iowensis, see *Lingulella iowensis*.
 irene, Billings, 1862, Pal. Foss., vol. 1, p. 71, Quebec Gr.
 iris, Billings, 1865, Pal. Foss., vol. 1, p. 301, Quebec Gr.
 kingstonensis, Billings, 1862, Pal. Foss., vol. 1, p. 48, Black Riv. Gr.
 lamellata, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 108, and Pal. N. Y., vol. 2, p. 249, Clinton and Niagara Grs.
 leana, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 20, and Pal. N. Y., vol. 4, p. 9, Ham. Gr.
 ligea, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 76, and Pal. N. Y., vol. 4, p. 7, Ham. Gr.
 ligea var. Hall, 1867, Pal. N. Y., vol. 4, p. 8, Portage Gr.
 ligea var. nevadensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 107, Devonian.
 lonensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 108, Devonian.
 lucretia, Billings, 1874, Pal. Foss., vol. 2, p. 14, passage beds between Up. Sil. and Devonian.
 lyelli, Billings, 1859, Can. Nat. Geo., vol. 4, p. 348, Calcif. and Chazy Grs.
 maida, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 20, and Pal. N. Y., vol. 4, p. 9, Ham. Gr.
 manni, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 20, and Pal. N. Y., vol. 4, p. 6, Up. Held. Gr.
 mantelli, Billings, 1859, Can. Nat. Geo., vol. 4, p. 349, Calcif. Gr.
 manticula, White, 1864, Rep. Invert. Foss., p. 9, and Geo. Sur. W. 100th Mer., vol. 4, p. 52, Up. Taconic.
matthewi, see *Acrothele matthewi*.
 melie, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 24, and Pal. N. Y., vol. 4, p. 14, Waverly Gr.
 membranacea, Winchell, 1863, Proc. Acad. Nat. Sci. Phil., vol. 15, p. 3, Marshall Gr.
 minuta, Meek, 1868, Trans. Chi. Acad. Sci., p. 87, Devonian.
 mosia, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 126, Potsdam Gr.
 murrayi, Billings, 1874, Pal. Foss., vol. 2, p. 66, Up. Taconic.
 mytiloides, Sowerby, 1812, Min. Couch., p. 55, tab. 19, Coal Meas.
 nebraskensis, Meek, 1872, (L. scotica var. nebraskensis), Pal. E. Neb., p. 158, Coal Meas.
 norwoodi, James, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 10, Utica Slate Gr.
 nuda, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 22, Ham. Gr.
 nymph, Billings, 1865, Pal. Foss., vol. 1, p. 214, Quebec Gr.
 oblata, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 77, and Pal. N. Y., vol. 2, p. 54, Clinton Gr.

FIG. 574.—*Lingulella norwoodi*.

- oblonga*, Conrad, 1839, Ann. Rep. N. Y. The name was preoccupied, and afterward it was called *L. clintoni*.
- obtusa*, Hall, 1847, Pal. N. Y., vol. 1, p. 98, Trenton Gr.
- ovata*, McCoy, 1844, Syn. Sil. Foss. Ireland, p. 24. Not clearly identified in America.
- paliformis*, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 76, and Pal. N. Y., vol. 4, p. 8, Ham. Gr.
- papillosa*, Emmons, 1856, Am. Geol., p. 209, Trenton Gr.
- perlata*, Hall, 1859, Pal. N. Y., vol. 3, p. 156, Low. Held. Gr.
- perovata*, Hall, 1852, Pal. N. Y., vol. 2, p. 55, Clinton Gr.
- perplexa*, Hall, 1877, 1st ed. Am. Pal. Foss., p. 244. Proposed instead of *L. elliptica*, which was preoccupied, but D'Orbigny had previously proposed *L. subelliptica*.
- perryi*, Billings, 1861, Pal. Foss., vol. 1, p. 20, Black Riv. Gr.
- philomela*, Billings, 1862, Pal. Foss., vol. 1, p. 49, Trenton Gr.
- pinniformis*, see *Lingulepis pinniformis*.
- polita*, see *Obolella polita*.
- prima*, see *Lingulepis prima*.
- prima*, Emmons, 1856, Am. Geol. This name was preoccupied.
- proctori*, Ulrich, 1889, Am. Geol., vol. 3, p. 377, Trenton Gr.
- progne*, Billings, 1862, Pal. Foss., vol. 1, p. 47, Utica Slate and Trenton Gr.
- punctata*, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 21, and Pal. N. Y., vol. 4, p. 10, Ham. Gr.
- quadrata*, Eichwald, 1829, (Crania quadrata.) Zool. Specialis, vol. 1, p. 273, and Pal. N. Y., vol. 1, p. 96, Trenton Gr.
- quebecensis*, Billings, 1862, Pal. Foss., vol. 1, p. 72, Quebec Gr.
- rectilatera*, Hall, 1859, Pal. N. Y., vol. 3, p. 156, Low. Held. Gr.
- rectilateralis*, Emmons, 1842, Geo. Rep. N. Y., p. 399, Utica Slate.
- riciniformis*, Hall, 1847, Pal. N. Y., vol. 1, p. 95, Trenton Gr.
- scotica*, Davidson, 1860, Monogr. Scot. Carb. Brach., p. 62, Waverly Gr.
- scotica* var. *nebraskensis*, see *L. nebraskensis*.
- spathata*, Hall, 1859, Pal. N. Y., vol. 3, p. 157, Low. Held. Gr.
- spatiosa*, Hall, 1859, Pal. N. Y., vol. 3, p. 158, Low. Held. Gr.
- spatulata*, Vanuxem, 1842, Geo. Rep. 3d Dist. N. Y., p. 168, and Pal. N. Y., vol. 4, p. 13, Genesee slate.
- stoneana*, Whitfield, 1882, Geo. Wis., vol. 4, p. 344, Potsdam Gr.
- striata*, Emmons, 1856, Am. Geol., p. 112, Up. Taconic.
- subelliptica*, D'Orbigny, 1850, Prodr. d. Paleont., t. 1, p. 34, Clinton Gr. Proposed instead of *L. elliptica*, Hall, in 1843, Geo. Rep. 4th Dist. N. Y., p. 77.

- suboblonga*, D'Orbigny, syn. for *L. clintoni*.
- subspatulata*, Meek & Worthen, 1878, Geo. Sur. Ill., vol. 3, p. 437, Ham. Gr.
- thedfordensis*, Whiteaves, 1887, Cont. to Can. Pal., vol. 1, p. 111, Har. Gr.
- trentonensis*, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 266, Trenton Gr.
- triquetra*, Clarke, 1885, Ill. U. S. Geo. Sur., vol. 16, p. 62, Portage Gr.
- umbonata*, Cox, 1857, Geo. Sur. Ky., vol. 3, p. 576, Coal Meas.

- vanhorni*, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 9, Hud. Riv. Gr.
- varsoviensis*, Worthen, 1884, Bull. No. 2 Ill. St. Mus. Nat. Hist., p. 24, and Geo. Sur. Ill., vol. 8, p. 104, Warsaw Gr.
- whitii*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 109, Devonian.
- whitfieldi*, Ulrich, 1889, Am. Geol., vol. 3, p. 381, Utica Slate.
- winona*, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 126, Potsdam Gr.
- LINGULELASMA*, Ulrich, 1889, Am. Geol., vol. 3, p. 383. [Ety. *Lingula*, a genus; *clasma*, plate.] Form and composition like *Lingula*; pedicle valve with slightly projecting beak, faintly arched deltidium, no area, small socket on each side of the deltidial borders, and subtriangular scar opposite their anterior ends subtriangular, trilobed platform from base of detidium to middle of valve, with central part produced below in a low median ridge; two muscular scars on the lower lateral sides of the platform; brachial valve, with transverse ridge and swollen ends for sockets on the opposite valve; platform concave, elevated in front, and prolonged in a median plate, subcardinal, umbolateral and postmedian scars. Type *L. schucherti*.

FIG. 576.—*Lingulelasma schucherti*. a, Posterior transverse ridge; b, subcardinal scars; u, umbolateral scars; p, postmedian scars; l, lateral scars; m, median scars; a, anterior scars; t, transverse scars; s, septum; va, ventral valve.

- schucherti*, Ulrich, 1889, Am. Geol., vol. 3, p. 389, Hud. Riv. Gr.

FIG. 574.—*Lingula norwoodi*.

LINGULELLA, Salter, 1861, Mem. Geo. North Wales, and Geo. Sur. Gt. Brit., vol. 3, p. 333. [Ety. diminutive of Lingula.]

FIG. 577.—*Lingulella cincinnatiensis*.

General form like *Lingula*; nearly equivalve, ventral valve pointed with a pedicle groove; muscular scars as in obolus, but anterior retractors more linear, and sliding muscles small and not quite as much external. Type *L. davisii*.

?affinis, Billings, 1874, Pal. Foss., vol. 2, p. 67, Up. Taconic.

aurora, Hall, 1861, (*Lingula aurora*), (Geo. Rep. Wis., p. 24, Potsdam Gr.

cuculata, Hall, 1847, (*Orbicula cuculata*), (Pal. N. Y., vol. 1, p. 290, Georgia Gr.

cincinnatiensis, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 67, Illud. Riv. Gr.

cuneata, Conrad, 1839, (*lingula cuneata*), (Geo. Rep. N. Y., p. 64, and Pal. N. Y., vol. 2, p. 8, Clinton Gr.

dawsoni, Matthew, 1885, Trans. Roy. Soc. Can., p. 33, St. John Gr.

ella, Hall & Whitfield, 1877, (*Lingulepis ella*), (Geo. Expl. 40th Parallel, vol. 4, p. 232, Up. Taconic.

granvillensis, Walcott, 1887, Am. Jour. Sci. and Arts., 3d ser., vol. 34, p. 187, Up. Taconic.

inflata, Matthew, 1885, Trans. Roy. Soc. Can., p. 33, St. John Gr.

iowensis, Owen, 1840, (*Lingula iowensis*), Rep. Min. Lands, p. 70, Galena Gr.

lamborni, Meek, 1871, Proc. Acad. Nat. Sci., p. 185, Calciferous or Potsdam Gr.

linguloides, Matthew, 1885, Trans. Roy. Soc. Can., p. 34, St. John Gr.

?spissa, Billings, 1874, Pal. Foss., vol. 2, p. 67, Up. Taconic.

LINGULEPIS, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 126. [Ety. *lingula*, little tongue; *lepis*, scale.] Linguloid, inequivalve, equilateral, ovate or spatulate, corneous, phosphatic; visceral impressions in dorsal valve flabelliform, in ventral valve tripartite, the lateral divisions the larger. Type *L. pinniformis*.

cuneolus, Whitfield, 1877, Prelim. Rep. Pal. Black Hills, p. 8, and Geol. Black Hills, p. 336, Potsdam Gr.

dakotensis, Meek & Hayden, 1864, Pal. Up. Mo., p. 3, and Geol. Black Hills, p. 337, Potsdam Gr.

ella, See *Lingulella ella*.

mæra, Hall & Whitfield, 1877, U. S. Expl. 40th Parallel, vol. 4, p. 206, Potsdam Gr.

FIG. 578.—*Lingulella cuneata*.

minima, Whitfield, 1884, Bull. Am. Mus. Nat. Hist., vol. 1, p. 139, Up. Taconic.

minuta, Hall & Whitfield, 1877, U. S. Expl. 40th Parallel, vol. 4, p. 206, Potsdam Gr.

morsii, N. H. Winchell, 1876, (*Lingula morsensis*), Geol. Fillmore Co., Minn., p. 31, St. Peters sandstone.

perattenuata, Whitfield, 1877, Prelim. Rep. Pal. Black Hills, p. 9, and Geol. Black Hills, p. 337, Potsdam Gr.

pinniformis, Owen, 1852, (*Lingula pinniformis*), (Geo. Rep. Iowa, Wis., and Minn., p. 583, Potsdam Gr.

prima, Conrad, 1847, (*Lingula prima*), (Pal. N. Y., vol. 1, p. 3, Potsdam Gr.

FIG. 579.—*Lingulella pinniformis*.

LINGULOPS, Hall, 1871, 23d

Rep. N. Y. St. Mus. Nat. Hist., p. 245.

[Ety. *Lingula*, a genus; *opsis*, appearance.] In external appearance like *Lingula* or *Lingulella*; the ventral valve presents a small area, with a narrow pedicle groove and a large lobed muscular impression, which, in the east, extends as a narrow groove toward the base of the shell; the ramifications of the vascular lines originate at nearly the same point as in existing *Lingula*, but do not extend so far backward toward the beak. Type *L. whitfieldi*.

whitfieldi, Hall, 1871, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 245, Low. Sil.

LINNAARSSONIA, Walcott, 1885, Am. Jour. Sci. and Arts., 3d ser., vol. 29, p. 114. [Ety. proper name.] Ovate or subcircular; inarticulate; apex of ventral valve perforated by a minute foramen; no area; cardinal edge thin; two scars in the interior, on each side of the foramen, close to the posterior margin; dorsal valve, with no area; two scars in the interior, close to the posterior margin, separated by a ridge that extends forward between two small divaricator scars. Type *L. transversa*.

taconica, Walcott, 1887, Am. Jour. Sci. and Arts., 3d ser., vol. 34, p. 189, Up. Taconic.

transversa, Hartt, 1868, (*Obolella transversa*), Acad. Geol., p. 644, St. John Gr.

MARTINIA, McCoy, 1844, syn. Carb. Foss., Ireland, p. 128. [Ety. proper name.] General characters the same as *Spirifera*, for which it is usually regarded as a synonym. It is distinguished by its smooth surface without radiating ribs, and by having smaller spiral appendages. Type *M. decora*.

athyroides, Winchell, 1866, Rep. Low. Penin. Mich., p. 94, Ham. Gr.

planiconvexa, see *Spirifera planiconvexa*. subumbonata, Hall, 1867, (*Spirifera subumbonata*), (Pal. N. Y., vol. 4, p. 234, Ham. Gr. and Tully limestone.

Am. Mus.
Faonic.
77, U. S.
206, Pots-

(Lingula
o., Minn.,

Prelim.
and Geol.
Gr.

—Lingule-
niformis.

st., p. 245.
ts, appear-
ance like
ventral valve
a narrow
obed mus-
toward the east.
fications of
at nearly
g Lingula,
ckward to-
nitfieldi.

b. N. Y. St.
Sil.

a. Jour. Sci.
114. [Ety.
subcircular;
valve per-
no area;
rs in the in-
amen, close
rsal valve,
he interior,
, separated
rd between
Type L.

Jour. Sci.
p. 189, Up.

ella trans-
644, St.

Carb. Foss.,
per name.]
as Spirifer,
arded as a
shed by its
iating ribs,
ral append-

Rep. Low.

inoconvexa.
pirifer sub-

4, p. 234,
one.

MEEKELLA, White & St. John, 1868, Trans. Chi. Acad. Sci., vol. 1, p. 120. [Ety. proper name.] Globose, rather longer than wide, plications large, hinge-line shorter than greatest breadth of the valves; dorsal valve most prominent on the umbo, beak incurved, no mesial sinus; cardinal process long, curving backward in front of the pseudodeltidium, and having upon each side a wing like expansion, which is curved up at its outer edge forming an elongated dental fossette; ventral valve more convex, cardinal area high, no median septum; two broad dental lamellae, continuous from the cardinal teeth to the beak, pass directly in front of the sutures between the cardinal area and the pseudodeltidium, and thence, slightly diverging, extend forward along the bottom of the valve about half-way to the front, the anterior margins of the lamellae arching backward and upward to the dental processes; a cross section shows three chambers opening anteriorly into the shell. Type *M. striato-costata*. *striato-costata*, Cox, 1857, (Plicatula *striato-costata*.) Geo. Rep. Ky., vol. 3, p. 568, Coal Meas.

FIG. 580.—Meekella striatocostata. Dorsal and ventral view.

Meganteris aequiradiata, see *Rensseleria aequiradiata*.

cumberlandia, see *Rensseleria cumberlandia*.

elliptica, see *Rensseleria elliptica*.

elongata, see *Amphigenia elongata*.

lævis, see *Rensseleria lævis*.

mutabilis, see *Rensseleria mutabilis*.

ovalis, see *Rensseleria ovalis*.

ovoides, see *Rensseleria ovoides*.

subtrigonalis, see *Amphigenia elongata* var. *subtrigonalis*.

suessana, see *Rensseleria suessana*.

MERISTA, Suess, 1851, Jahrb. Geol. Reichs. Aust., vol. 2, p. 150. [Ety. *meros*, a part.] General form like *Athyris*, usually mesial fold and sinus poorly defined; the principal stems forming the spirals attach to the hinge plate, incline forward toward the interior of the shell, then abruptly bend backward and make a curve facing the bottom of the dorsal valve, and, after converging to about half their length, again diverge toward the front and form the

first spiral coil; there are 10 or 12 whorls in each spiral; the genus is distinguished by a shoe-lifter process under the beak of the ventral valve, consisting of two roof-shaped plates, fixed by their lateral margins to the medio-longitudinal region of the valve, and with their narrow end fitting under the extremity of the beak. Type *M. herculea*.

arcuata, see *Meristella arcuata*.

FIG. 581.—Merista bella. Dorsal and anterior view.

bella, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 92, and Pal. N. Y., vol. 3, p. 248, Low. Held. Gr.

bisulcata, Vanuxem, 1843, (Atrypa *bisulcata*.) Geo. Rep. 3d Dist. N. Y., p. 112, and Pal. N. Y., vol. 3, p. 253, Low. Held. Gr.

elongata, Hall, 1859, (Camarium *elongatum*.) Pal. N. Y., vol. 3, p. 488, Low. Held. Gr.

houghtoni, Winchell, 1862, Proc. Acad. Nat. Sci., vol. 6, 2d ser., p. 407, Portage Gr.

lævis, see *Meristella lævis*.

lata, Hall, 1859, Pal. N. Y., vol. 3, p. 431, Oriskany sandstone.

lens, Winchell, 1866, Rep. Low. Penin. Mich., p. 94, Ham. Gr.

meeki, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 97, and Pal. N. Y., vol. 3, p. 252, Low. Held. Gr.

princeps, see *Meristella princeps*.

subquadrata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 93, and Pal. N. Y., vol. 3, p. 249, Low. Held. Gr.

sulcata, Vanuxem, 1842, (Atrypa *sulcata*.) Geo. Rep. N. Y., p. 112, Waterlime Gr.

typus, Hall, 1859, (Camarium *typus*.) Pal. N. Y., vol. 3, p. 487, Low. Held. Gr.

MERISTELLA, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 74. [Ety. diminutive of *Merista*.] Shells variable in form, ovoid or transverse; valves unequally convex, with or without a median fold and sinus; beak of ventral valve imperforate, incurved over the beak of the other valve; no arca; valves articulating by teeth and sockets; surface smooth or with fine concentric striae; interior of dorsal valve having a longitudinal septum and the upper part of the ventral valve a deep subtriangular muscular impression, which unites with the rostral cavity; spires are continued from their origin obliquely backward into the cavity of the ventral valve, and then,

recurving upon themselves, are reunited laterally. Type *M. lævis*.

FIG. 582. — *Meristella circe*. Showing remains of spiral appendages in the dorsal valve.

cylindrica, Hall, 1852, (*Atrypa cylindrica*), Pal. N. Y., vol. 2, p. 76, Clinton and Niagara Gr.

doris, 1860, Hall, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 84, and Pal. N. Y., vol. 4, p. 303, Schoharie grit and Corniferous Gr.

elissa, syn. for *Meristella nasuta*. *haskinsi*, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 84, and Pal. N. Y., vol. 4, p. 306, Ham. Gr.

(?) *hyale*, Billings, 1862, (*Charionella* (?) *hyale*), Pal. Foss., vol. 1, p. 160, Guelph Gr.

julia, Billings, 1862, (*Athyris julia*), Pal. Foss., vol. 1, p. 146, Mid. Sil.

lævis, Vanuxem, 1843, (*Atrypa lævis*), Geo. Rep. 3d Dist. N. Y., p. 120, and Pal. N. Y., vol. 3, p. 247, Low Held. Gr. *lenta*, Hall, 1867, Pal. N. Y., vol. 4, p. 420, Oriskany sandstone.

maria, see *Whitfieldia maria*.

meta, Hall, 1867, Pal. N. Y., vol. 4, p. 308, Ham. Gr.

nasuta, Conrad, 1840, (*Atrypa nasuta*), Ann. Rep. N. Y., p. 18, and Pal. N. Y., vol. 4, p. 299, Schoharie grit, Up. Held. and Ham. Gr.

princeps, Hall, 1857, (*Merista princeps*), 10th Rep. N. Y. St. Mus. Nat. Hist., p. 95, and Pal. N. Y., vol. 3, p. 252, Low Held. Gr.

prinstana, Billings, 1862, (*Athyris prinstana*), Pal. Foss., vol. 1, p. 145, Mid. Sil. *rectirostra*, Hall, 1879, Desc. New Spec. Foss., p. 15, and 11th Rep. Geo. and Nat. Hist. Ind., p. 301, Niagara Gr.

rostrata, Hall, 1843, (*Atrypa rostrata*), Geo. Rep. 4th Dist. N. Y., p. 202, and Pal. N. Y., vol. 4, p. 307, Ham. Gr. and Tully limestone.

scitula, Hall, 1843, (*Atrypa scitula*), Geo. 4th Dist. N. Y., p. 171, and Pal. N. Y., vol. 4, p. 302, Corniferous Gr. Hall regards *M. circe* as a syn. for this species.

themselves, are reunited laterally. Type *M. lævis*.

arcuata, Hall, 1857, (*Merista arcuata*), 10th Rep. N. Y. St. Mus. Nat. Hist., p. 95, and Pal. N. Y., vol. 3, p. 249, Low Held. Gr.

barrisi, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 84, and Pal. N. Y., vol. 4, p. 304, Marcellus shale and Ham. Gr.

circe, Billings, 1861, (*Charionella circe*), Can. Jour., vol. 6, p. 273, Up. Held. Gr.

cylindrica, Hall, 1852, (*Atrypa cylindrica*), Pal. N. Y., vol. 2, p. 76, Clinton and Niagara Gr.

doris, 1860, Hall, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 84, and Pal. N. Y., vol. 4, p. 303, Schoharie grit and Corniferous Gr.

elissa, syn. for *Meristella nasuta*. *haskinsi*, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 84, and Pal. N. Y., vol. 4, p. 306, Ham. Gr.

(?) *hyale*, Billings, 1862, (*Charionella* (?) *hyale*), Pal. Foss., vol. 1, p. 160, Guelph Gr.

julia, Billings, 1862, (*Athyris julia*), Pal. Foss., vol. 1, p. 146, Mid. Sil.

lævis, Vanuxem, 1843, (*Atrypa lævis*), Geo. Rep. 3d Dist. N. Y., p. 120, and Pal. N. Y., vol. 3, p. 247, Low Held. Gr. *lenta*, Hall, 1867, Pal. N. Y., vol. 4, p. 420, Oriskany sandstone.

maria, see *Whitfieldia maria*.

meta, Hall, 1867, Pal. N. Y., vol. 4, p. 308, Ham. Gr.

nasuta, Conrad, 1840, (*Atrypa nasuta*), Ann. Rep. N. Y., p. 18, and Pal. N. Y., vol. 4, p. 299, Schoharie grit, Up. Held. and Ham. Gr.

princeps, Hall, 1857, (*Merista princeps*), 10th Rep. N. Y. St. Mus. Nat. Hist., p. 95, and Pal. N. Y., vol. 3, p. 252, Low Held. Gr.

prinstana, Billings, 1862, (*Athyris prinstana*), Pal. Foss., vol. 1, p. 145, Mid. Sil. *rectirostra*, Hall, 1879, Desc. New Spec. Foss., p. 15, and 11th Rep. Geo. and Nat. Hist. Ind., p. 301, Niagara Gr.

rostrata, Hall, 1843, (*Atrypa rostrata*), Geo. Rep. 4th Dist. N. Y., p. 202, and Pal. N. Y., vol. 4, p. 307, Ham. Gr. and Tully limestone.

scitula, Hall, 1843, (*Atrypa scitula*), Geo. 4th Dist. N. Y., p. 171, and Pal. N. Y., vol. 4, p. 302, Corniferous Gr. Hall regards *M. circe* as a syn. for this species.

FIG. 583. — *Meristella rostrata*. Side view.

unisulcata, Conrad, 1841, (*Atrypa unisulcata*), Ann. Rep. N. Y., p. 56, and Pal. N. Y., vol. 4, p. 309, Up. Held. and Ham. Gr.

MERISTINA, Hall, 1867, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 186, and Pal. N. Y., vol. 4, p. 299. [Ety. *Merista*, a genus; *inus*, implying resemblance.] General form like *Meristella*; apex perforated; lamellæ of the spires united by a simple loop; spirals oval, and each contains about nine convolutions; the two principal siems attach to the hinge plate and extend into the interior between the spirals, where they bend backward and give forth converging lamellæ, which cross between the spirals to the ventral side, where they unite in an angular point. Type *M. nitida*.

nitida, Hall, 1843, (*Atrypa nitida*), Geo. Rep. 4th Dist. N. Y., pl. 14, and Pal. N. Y., vol. 2, p. 268, Niagara Gr.

nitida var. *oblata*, Hall, 1852, (*Atrypa nitida* var. *oblata*), Pal. N. Y., vol. 2, p. 269, Niagara Gr.

nitida var. *oblata*, Hall, 1852, (*Atrypa nitida* var. *oblata*), Pal. N. Y., vol. 2, p. 269, Niagara Gr.

MONOMERELLA, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 220. [Ety. *monos*, one; *meros*, a part; *ella*, diminutive termination.] Shell thick, circular or transversely oval in its marginal outline; umbo of the pedicle; valve large, projecting, double-chambered; area and deltidium large; hinge thick, elevated, ledge-shaped, concave in the middle portion; cardinal facet a wall-like space behind the ledge or flat of the hinge; cardinal buttress strong, lamelliform; platform flat, slightly elevated, widest, highest, and obtusely angulated in front; umbo of the brachial valve rounded; hinge moderately thick; platform trilobed; usually with a thin margin. Type *M. prisca*.

newberryi, Hall & Whitfield, 1875, Ohio, Pal., vol. 2, p. 131, Niagara Gr.

orbicularis, Billings, 1871, Can. Nat., vol. 6, p. 220, Guelph Gr.

ovata, Whiteaves, 1884, Pal. Foss., vol. 3, p. 5, Guelph Gr.

ovata var. *lata*, Whiteaves, 1884, Pal. Foss., vol. 3, p. 6, Guelph Gr.

prisca, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 220, Guelph Gr.

prisca, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 220, Guelph Gr.

prisca, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 220, Guelph Gr.

prisca, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 220, Guelph Gr.

prisca, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 220, Guelph Gr.

FIG. 584. — *Meristina nitida*. Dorsal view.

FIG. 585. — *Monomerella prisca*.

Nucleo-

p. 2

spin-

vers-

and

era

trac-

tien-

smo-

erec-

ven-

tenc-

ben-

bas-

from

spat-

upw-

beak-

the

supp-

conf-

N. v

Fig. 586.

cleopatra

elma.

pisiform

1852,

3, p.

rotunda

Spec.

Held

ventric

p. 220

was f

Rep.,

Cholella

p. 7.

small

lar or

conve

area,

usuall

pedun

pa uni-
56, and
eld, and

N. Y. St.
l. N. Y.,
genus;
General
rforated;
by a sim-
ach com-
ns; the
he hinge
erior be-
y bend
nverging
mespirals
unite in
tida.

Meristina
Dorsal

Nat. and
nos, one;
termina-
or trans-
outline;
large, pro-
area and
elevated,
e middle
wall-like
at of the
strong, la-
ghtly ele-
obtusely
o of the
ge moder-
l; usually
M. prisca.
wberryi,
Hall &
Whitfield,
1875, Ohio,
Pal., vol.
2, p. 131,
Niagara
Gr.

ocularis,
Billings,
1871, Can.
Nat., vol.
6, p. 220,
Guelph Gr.
Foss., vol.

1884, Pal.
and Geol.,

NUCLEOSPIRA, Hall, 1859, Pal. N. Y., vol. 3, p. 219. [Ety. *nucleus*, kernel; *spira*, spire.] Shell spheroidal, or transversely elliptical, more or less gibbous, and furnished with spires as in Spirifera; hinge-line short, cardinal extremities rounded, valves subequal, articulating by teeth and sockets; surface smooth, structure punctate and covered with minute hair-like spines; ventral valve having the beak extended, with a triangular depression beneath, on each side of which at the base is a strong tooth, a narrow septum from beak to base; dorsal valve with spatulate cardinal process, which bends upward into the cavity of the opposite beak; from the sides of this process the brachial processes originate, which support the spires; muscular imprints confined to a narrow oval space. Type *N. ventricosa*.

Fig. 586.—*Nucleospira concinna*.

barrisi, White, 1860, Bost. Jour. Nat. Hist., vol. 7, p. 227, Kinderhook Gr.
concentrica, Hall, 1859, Pal. N. Y., vol. 3, p. 223, Low. Held. Gr.
concinna, Hall, 1843, (*Atrypa concinna*,) Geo. 4th Dist. N. Y., p. 200, and Pal. N. Y., vol. 4, p. 279, Hamilton Gr.
elegans, Hall, 1859, Pal. N. Y., vol. 3, p. 222, Low. Held. Gr.
pisiformis, Hall, 1859, (*Orthis pisum*, 1852, Pal. N. Y., vol. 2,) Pal. N. Y., vol. 3, p. 218, Niagara Gr.
rotundata, Whitfield, 1882, Desc. New Spec. Foss. from Ohio, p. 194, Low. Held. Gr.

Fig. 587.—Spiral coils of *Nucleospira pisiformis*.

ventricosa, Hall, 1859, Pal. N. Y., vol. 3, p. 220, Low. Held. Gr. This species was first described in 1856, in 9th Reg. Rep., as *Spirifera ventricosa*.
OBOLELLA, Billings, 1861, Pal. Foss., vol. 1, p. 7. [Ety. diminutive of *obolus*, a small Greek coin.] Shell ovate, circular or subquadrate, convex or plano-convex; ventral valve with a false area, which is sometimes minute and usually grooved for the passage of the peduncle; dorsal valve with or without

an area; muscular impressions in the ventral valve four, one pair in front of the beak near the middle or in the upper half of the shell, and the others situated one on each side near the cardinal edge; shell calcareous; surface concentrically striated. Type *O. chromatica*.

ambigus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 67, Chazy Gr.
chromatica, Billings, 1861, Pal. Foss., vol. 1, p. 7, Up. Taconic.
cingulata, Billings, 1861, Pal. Foss., vol. 1, p. 8, Up. Taconic.
circe, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 219, Up. Taconic.
crassa, Hall, 1847, (*Orbicula crassa*,) Pal. N. Y., vol. 1, p. 290, Up. Taconic Gr.

Fig. 588.—*Obolus chromatica*. c, Showing muscular impressions; d, side view.

desiderata, see *Elkania desiderata*.
discoidea, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 205, Potsdam Gr.
gemma, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 218, Up. Taconic.
ida, Billings, 1862, Pal. Foss., vol. 1, p. 71, Quebec Gr.
misera, Billings, 1874, Pal. Foss., vol. 2, p. 69, Up. Taconic.
nana, Meek & Hayden, 1861, Proc. Acad. Nat. Sci. Phil., p. 435, and Pal. Up. Mo., p. 4, Potsdam Gr.
nitida, Ford, 1873, Am. Jour. Sci. and Arts, 3d ser., vol. 5, p. 213, Up. Taconic.
polita, Hall, 1861, Geo. Rep. Wis., p. 24, and Geol. Black Hills, p. 339, (*Lingula polita*,) Potsdam Gr.
pretiosa, Billings, 1862, Pal. Foss., vol. 1, p. 68, Quebec Gr.
transversa, see *Linnarssonsonia transversa*.
Obolellina, Billings, Dec., 1871, Can. Nat., vol. 6, p. 220, syn. for *Dinobolus*.
canadensis, see *Dinobolus canadensis*.
galtensis, see *Dinobolus galtensis*.
magnifica, see *Dinobolus magnificus*.

OBOLOUS, Eichwald, 1829, Zoologiu Specialis, vol. 1, p. 274. [Ety. *obolus*, a small coin.] Shell orbicular, equilateral, transverse or elongated, depressed; valves not articulated; larger valve most convex, beak obtuse or pointed, wide flattened cardinal edge or false area, over which the concentric lines of surface growth pass uninterruptedly; cardinal edge grooved longitudinally by a semicylindrical furrow; smaller valve shorter, slightly convex, without prominent beak; hinge-line an arch; cardinal edge flattened, horizontally striated; surface smooth or having minute undulating wrinkles; interior of larger valve with a mesial ridge, on each side of which are two oval muscular scars, one pair near the cardinal angles, the other toward the center of the valve beyond the mesial ridge; structure calcareo-concorneous. Type *O. apollinis*.

canadensis, see *Dinobolus canadensis*.
conradi, see *Dinobolus conradi*.

FIG. 589.—*Obolus apollinis*. b, Dorsal valve; a, interior of ventral valve.

galtensis, see *Trimerella galtensis*.

labradoricus, see *Kutorgina labradorica*.

(?) *murrayi*, Billings, 1865, *Pal. Foss.*, vol. 1, p.

362, Quebec Gr. or Up. Taconic.

pectinoides, Whitfield, 1875, *Ludlow's Rep. Black Hills of Dakota*, p. 103, Up. Taconic.

Orbicula, Cuvier, 1808, *Tabl. Elem. d'Hist. Nat.*, p. 435, syn. for *Crania*.

cyclata, see *Lingulella cyclata*.

cancellata, see *Trematis cancellata*.

corrugata, see *Crania corrugata*.

crassa, see *Obolella crassa*.

deformata, see *Crania deformata*.

eccentrica, see *Crania eccentrica*.

filosa, see *Schizocrania filosa*.

grandis, see *Discina grandis*.

lamellosa, see *Discina lamellosa*.

lodensis, see *Discina lodensis*.

minuta, see *Discina minuta*.

nitida, see *Discina nitida*.

prima, see *Crania prima*.

squamiformis, see *Pholidops squamiformis*.

subtruncata, see *Pholidops subtruncatus*.

tenuilamellata, see *Discina tenuilamellata*.

terminalis, see *Trematis terminalis*.

truncata, see *Crania truncata*.

ORBICULOIDEA, D'Orbigny, 1847, *Comptes rendus de l'Académie des Sciences*, and *Prodr. de Paléont.*, t. 1, p. 44. [Ety. *Orbicula*, a genus; *oides*, like.] Suborbicular, patelliform, longitudinally or transversely oval, upper valve convex, with vertex near the posterior margin; lower valve conical or concave; no pedicle disk; a narrow oval or circular aperture, more or less confined in its shape, is situated in a furrow or depression. Type *O. elliptica*.

conica, Dwight, 1880, *Am. Jour. Sci. and Arts*, 3d ser., vol. 19, p. 452, Trenton Gr.

ORTHIS, Dalman, 1827, *Kongl. Vet. Acad. Handl.*, p. 93. [Ety. *orthos*, straight, in allusion to the straight hinge-line.] Shell variable in form, hinge-line straight; valves convex or plano-convex, plicated, with or without mesial fold and sinus; cardinal area notched in the center; ventral valve with two prominent diverging teeth, muscular impression saucer-shaped, divided by a median septum on which the central adductor attached; divaricator and pedicle impressions, lateral, fan-like; dorsal valve with a tooth-like, cardinal process between two curved brachial processes; adductor impression quadruple; vascular impressions numerous, spreading; no coiled spiral arms. Type *O. zonata* and *O. callactis*.

acuminata, Billings, 1859, *Can. Nat. Geo.*, vol. 4, p. 440, Chazy Gr.

acutilirata, Conrad, 1842, (*Delthyris acutilirata*), *Jour. Acad. Nat. Sci.*, vol. 8, p. 260, Hud. Riv. Gr.

acutitiloba, Ringueberg, 1888, *Proc. Acad. Nat. Sci. Phil.*, p. 134, Niagara Gr.

aequivalvis, Hall, 1847, *Pal. N. Y.*, vol. 1, p. 129, Trenton Gr.

aequivalva, Shaler. The name was preoccupied.

aequivalvis, Hall, see *Orthis cryna*.

alata, Shaler. The name was preoccupied. *alsus*, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 33, Schoharie gr.

alternans, Castelnau, 1843, *Syst. Sil.*, p. 38. Not recognized.

amona, Winchell, 1880, *Geo. Sur. Minn.* 8th Rep., p. 65, Hud. Riv. Gr.

anticostiensis, syn. for *Orthis porcata*.

apicalis, Billings, 1865, *Pal. Foss.*, vol. 1, p. 301, Quebec Gr.

armanda, Billings, 1865, *Pal. Foss.*, vol. 1, p. 303, Quebec Gr.

assimilis, Hall, 1859, *Pal. N. Y.*, vol. 3, p. 175, Low. Held. Gr.

aurelia, Billings, 1874, *Pal. Foss.*, vol. 2, p. 34, Gaspé limestone No. 8, Devonian.

barabuensis, see *Leptæna barabuensis*.

battis, Billings, 1865, *Pal. Foss.*, vol. 1, p. 185, Quebec Gr.

bellarugosa, Conrad, 1843, *Proc. Acad. Nat. Sci. Phil.*, vol. 1, p. 333, and *Pal. N. Y.*, vol. 1, p. 118, Trenton Gr.

bellula, Meek, 1873, *Ohio Pal.*, vol. 1, p. 103, Hud. Riv. Gr.

biforata, Schlotheim, 1820, (*Terebratulites biforatus*), *Petrefact.*, p. 265, Trenton and Hud. Riv. Grs.

billingsi, Hartt, 1868, *Acad. Geol.*, p. 644, St. John Gr.

biloba, Linnaeus, 1767, (*Anomia biloba*), *Linne. Syst.*, ed. 12, p. 1154, Niagara Gr.

bilobata, Conrad, 1838, (*Delthyris bilobata*), *Ann. Rep. N. Y.* The name was preoccupied by Sowerby.

bisulcata, see *Camarella bisulcata*.

borealis, Billings, 1859, *Can. Nat. Geo.*, vol. 4, p. 436, Chazy and Trenton Grs.

carbonaria, Swallow, 1858, syn. for *Orthis pectosi*.

carinata, Hall, 1843, *Geo. Rep.* 4th Dist. N. Y., p. 267, and *Pal. N. Y.*, vol. 4, p. 58, Portage and Chemung Grs.

carleyi, Hall, syn. for *Orthis retrorsa*.

centrilineata, Hall, 1847, *Pal. N. Y.*, vol. 1, p. 289, Hud. Riv. Gr.

centrosa, n. sp., Hud. Riv. Gr. Proposed instead of *O. crassa*, in *Cin. Quar. Jour. Sci.*, vol. 1, p. 20, and *Ohio Pal.*, vol. 1, p. 117, pl. 10, fig. 3.

charlottæ, Winchell, 1880, *Geo. Sur. Minn.* 8th Rep., p. 67, Hud. Riv. Gr.

cincinnatiensis, S. A. Miller, 1883, 2d Ed. *Am. Pal. Foss.*, p. 296, Hud. Riv. Gr.

cincinnatiensis, S. A. Miller, 1883, 2d Ed. *Am. Pal. Foss.*, p. 296, Hud. Riv. Gr.

cincinnatiensis, S. A. Miller, 1883, 2d Ed. *Am. Pal. Foss.*, p. 296, Hud. Riv. Gr.

costata, Hall, 1845, *Am. Jour. Sci. and Arts*, vol. 48, p. 294.

circ

80

circ

N

80

clar

A

cleo

M

sto

clyti

M

colo

Lo

da

colo

ple

conc

173

conra

No

conri

8th

coop

Low

sew

corin

p

costa

20

costa

eur

cin

crassa

vol

by

crisp

p

cumb

3

p

cune

low

cycl

Nat

p

cycl

cypha

esta

dalya

Nat

david

Fr

dayto

Den

deflect

deform

Mus

3

p

delicat

1

p

dentat

tatu

Tren

desmo

Rep

Prop

dichoto

discus

165

Nat. Geo.,
 hyrisaenti-
 ., vol. 8, p.
 Proc. Acad.
 ra Gr.
 Y., vol. 1.
 e was pre-
 na.
 reoccupied.
 N. Y. St.
 harie grit.
 yst. Sil., p.
 Sur. Minn.
 Gr.
 orcata.
 Foss., vol. 1.
 Foss., vol. 1.
 Y., vol. 3, p.
 ss., vol. 2, p.
 Devonian.
 buensis.
 ss., vol. 1, p.
 Proc. Acad.
 33, and Pal.
 on Gr.
 ul., vol. 1, p.
 erebratulites
 265, Trenton
 eol., p. 644.
 mia biloba.)
 1154, Niag-
 ris bilobata.)
 me was pre-
 cata.
 . Nat. Geo.
 Trenton Grs.
 n. for Orthis
 ep. 4th Dist.
 Y., vol. 4, p.
 Grs.
 retrorsa.
 . N. Y., vol.
 r. Proposed
 Quar. Jour.
 o Pal., vol. 1.
 o. Sur. Minn.
 Gr.
 1883, 2d Ed.
 ud. Riv. Gr.
 ed instead of
 5, Am. Jour.
 4.

circularis, Winchell, 1880, Geo. Sur. Min.
 8th Rep., p. 66, Hud. Riv. Gr.
 circulus, Hall, 1843, Geo. Rep. 4th Dist.
 N. Y., p. 71, and Pal. N. Y., vol. 2, p.
 86, Clinton Gr.
 clarkensis, Swallow, 1863, Trans. St. Louis
 Acad. Sci., vol. 2, p. 81, Keokuk Gr.
 cleobis, Hall, 1863, 16th Rep. N. Y. St.
 Mus. Nat. Hist., p. 35, Onondaga lime-
 stone, and Up. Held. Gr.
 clytie, Hall, 1861, 14th Rep. N. Y. St.
 Mus. Nat. Hist., p. 90, Trenton Gr.
 coloradoensis, Shumard, 1860, Trans. St.
 Louis Acad. Sci., vol. 1, p. 627, Pots-
 dam Gr.
 coloradoensis, Meek, 1870, see *O. desmo-*
pleura.
 concinna, Hall, 1850, Pal. N. Y., vol. 3, p.
 172, Low. Held. Gr.
 conradi, Castelnau, 1843, Syst. Sil. p. 37.
 Not recognized.
 conradi, Winchell, 1880, Geo. Sur. Minn.
 8th Rep., p. 68, Hud. River. Gr.
 cooperensis, Swallow, 1863, Trans. St.
 Louis Acad. Sci., vol. 2, p. 82, Wur-
 saw Gr.
 corinna, Billings, 1865, Pal. Foss., vol. 1,
 p. 302, Quebec Gr.
 costalis, Hall, 1847, Pal. N. Y., vol. 1, p.
 20, Chazy Gr.
 costata, Hall, 1845. This name was preoc-
 cupied by Sowerby in 1839. See *O.*
cincinnatiensis.
 crassa, James, 1874, Cin. Quar. Jour. Sci.,
 vol. 1, p. 20. The name was preoccupied
 by Lindstrom in 1860. See *O. centrosa*.
 crispata, Emmons, 1842, Geo. Rep. N. Y.,
 p. 404, Trenton Gr.
 cumberlandia, Hall, 1859, Pal. N. Y., vol.
 3, p. 481, Oriskany sandstone.
 cuneata, Owen, 1852, Geo. Sur. Wis.,
 Iowa, and Minn., p. 585, Devonian.
 cycas, Hall, 1860, 13th Rep. N. Y. St. Mus.
 Nat. Hist., p. 78, and Pal. N. Y., vol. 4,
 p. 52, Ham. Gr.
 cycus, James, syn. for *Orthis multisecta*.
 cypha, James. Not characterized so as to
 establish a species.
 dalyana, S. A. Miller, 1881, Jour. Cin. Soc.
 Nat. Hist., vol. 4, p. 313, Burlington Gr.
 davidsoni, Verneuil, 1840, Bull. Geol. Soc.
 France, vol. 5, p. 341, Up. Sil.
 daytonensis, Foerste, 1885, Bull. Sci., Lab.
 Denison Univ., p. 87, Niagara Gr.
 deflecta, see *Streptorhynchus deflectum*.
 deformis, Hall, 1857, 10th Rep. N. Y. St.
 Mus. Nat. Hist., p. 44, Pal. N. Y., vol.
 3, p. 174, Low. Held. Gr.
 delicatula, Billings, 1865, Pal. Foss., vol.
 1, p. 217, Quebec Gr.
 dentata, Pander, 1830, (Porambonites den-
 tatus,) Biétr. Geogn. Russl., p. 100,
 Trent. and Hud. Riv. Gr.
 desmopleura, Meek, 1872, Hayden's Geo.
 Rep. of Wyoming, p. 295, Silurian.
 Proposed instead of *O. coloradoensis*.
 dichotoma, syn. for *Orthis fissicosta*.
 discus, Hall, 1859, Pal. N. Y., vol. 3, p.
 165, Low. Held. Gr.

disparilis, Conrad, 1843, Proc. Acad. Nat.
 Sci., vol. 1, p. 333, and Pal. N. Y., vol. 1,
 p. 119, Black Riv. and Trenton Gr.
 dubia, Hall, 1858, Trans. Alb. Inst., vol.
 4, p. 12, and Bull. Am. Mus. Nat. Hist.,
 p. 45, Warsaw Gr.
 eboracensis, n. sp., Up. Held. Gr. Pro-
 posed instead of *O. lenticularis* of
 Vanuxem in Rep. 3d Dist. N. Y., p.
 147, which was preoccupied. It was re-
 described by Hall in Pal. N. Y., vol. 4,
 p. 35.
 electra, Billings, 1862, Pal. Foss. vol. 1, p.
 79, Quebec Gr.
 elegantula, Dalman, 1827, Kongl. Vet.
 Acad. Handl., p. 117, and Pal. N. Y.,
 vol. 2, p. 57, and 252, Clinton and Ni-
 agara Gr.

FIG. 500.—*Orthis elegantula*. Dorsal
 and ventral views.

ella, Hall, 1861, 13th Rep. N. Y. St. Mus.
 Nat. Hist., p. 121, Hud. Riv. Gr.
 emacerata, Hall, 1860, 13th Rep. N. Y.
 St. Mus. Nat. Hist., p. 121, Hud. Riv. Gr.
 emarginata, see *Orthis oblata* var. *emargi-*
nata.
 eminens, Hall, 1857, 10th Rep. N. Y. St.
 Mus. Nat. Hist., p. 42, and Pal. N. Y.,
 vol. 3, p. 167, Low. Held. Gr.
 erratica, Hall, 1847, Pal. N. Y., vol. 1, p.
 288, Hud. Riv. Gr.
 eryna, Hall, 1863, (Corrigenda eryna.)
 16th Rep. N. Y. St. Mus. Nat. Hist., p.
 35, and Pal. N. Y., vol. 4, p. 42, Cor-
 nif. Gr. Named instead of *O. aequal-*
vis in 10th Rep., p. 102.
 eudocia, Billings, 1862, Pal. Foss., vol.
 1, p. 83, Quebec Gr.
 eurekensis, Walcott, 1885, Monogr. U. S.
 Geo. Sur., vol. 8, p. 22, Up. Taconic.
 euryone, Billings, 1862, Pal. Foss., vol. 1,
 p. 78, Quebec Gr.
 evadne, Billings, 1862, Pal. Foss., vol. 1,
 p. 81, Quebec Gr.
 fasciata, Hall, 1852, Pal. N. Y., vol. 2, p.
 255, Niagara Gr.
 fausta, Foerste, 1885, Bull. Sci. Lab. Deni-
 son Univ., p. 85, Niagara Gr.
 fissicosta, Hall, 1847, Pal. N. Y., vol. 1,
 p. 121, Hud. Riv. Gr.
 fissiplica, Roemer, 1860, Sil. Fauna West
 Tenn., p. 64, Niagara Gr.
 flabellum, Sowerby, 1839, in Murch. Sil.
 Syst., p. 639, and Pal. N. Y., vol. 2, p.
 254, Niagara Gr.
 flava, Winchell, 1865, Proc. Acad. Nat.
 Sci., p. 117, Marshall Gr.
 gemmicula, Billings, 1862, Pal. Foss., vol.
 1, p. 75, Quebec Gr.
 gibbosa, Billings, 1857, Rep. of Progr.
 Geo. Sur. Can., p. 296, Black Riv. Gr.
 hamburgensis, Walcott, 1885, Monogr.
 U. S. Geo. Sur., vol. 8, p. 73, Chazy Gr.

- highlandensis, Walcott, 1886, Bull. U. S. Geo. Sur., No. 30, p. 119, Upper Taconic.
- hipparionyx*, syn. for *O. proximus*.
- hippolyte*, Billings, 1862, Pal. Foss., vol. 1, p. 81, Quebec Gr.
- huronensis*, Castelnaud, 1843, Syst. Sil., p. 37. Not recognized.
- hybrida*, Sowerby, 1830, Murch. Sil. Syst., p. 630, Niagara Gr.
- ldonea*, Hall, 1867, Pal. N. Y., vol. 4, p. 52, Ham. Gr.
- imperator*, Billings, 1850, Can. Nat. Geo., vol. 4, p. 435, Chazy Gr.
- impressa*, Hall, 1843, Geo. Rep. 4th Dist., N. Y., p. 268, and Pal. N. Y., vol. 4, p. 60, Chemung Gr.
- inaequalis*, Hall, 1858, Geo. of Iowa, p. 490, Ham. Gr.
- infera*, Calvin, 1878, Bull. U. S. Geo. Sur. Terr., vol. 4, No. 3, p. 728, Low. Devonian.
- insculpta*, Hall, 1847, Pal. N. Y., vol. 1, p. 125, Hud. Riv. Gr.
- insignis*, see *Skenidium insignis*.
- interlineata*, Sowerby, see *Orthis tioga*.
- interstitialis*, Phillips, 1841, Pal. Foss., Devonian. This species is probably foreign to America.
- iowensis*, Hall, 1858, Geo. of Iowa, p. 488, Ham. Gr.
- iowensis* var. *furnarius*, Hall, 1858, p. 489, Geo. of Iowa, Ham. Gr.
- iphigenia*, Billings, 1862, Pal. Foss., vol. 1, p. 133, Trenton Gr.
- jamesi*, Hall, 1861, 14th Rep. N. Y. St. Mus. Nat. Hist., p. 89, Hud. Riv. Gr.
- kankakensis*, McChesney, 1860, Desc. New Pal. Foss., p. 77, Hud. Riv. Gr.
- kassabae*, Winchell, 1880, Geo. Sur. Minn. 8th Rep., p. 65, Hud. Riv. Gr.
- kennicotti*, McChesney, syn. for *O. retrorsa*.
- keokuk*, Hall, 1858, Geo. Rep. Iowa, p. 640, Keokuk Gr. This species was referred to *Orthis umbraculum* of DeKoninck by Owen.
- lasallensis*, McChesney, 1860, New Pal. Foss., p. 32, syn. for *Streptorhynchus crassum*.
- laticosta*, Meek, 1873, Pal. Ohio, vol. 1, p. 116, Hud. Riv. Gr.
- laurentina*, Billings, 1857, Rep. of Geo. Sur. Can., p. 297, Mid. Sil., Anticosti Gr., Div. 1.
- lenticularis*, Vanuxem, 1842, Geo. Rep. 3d Dist. N. Y., p. 139. The name was preoccupied by Wahlenberg in 1821. See *O. eboracensis*.
- leonensis*, Hall, 1867, Pal. N. Y., vol. 4, p. 62, Chemung Gr.
- lepida*, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 78, and Pal. N. Y., vol. 4, p. 46, Ham. Gr.
- lepis*, as identified by d'Archiac & Verneuil. Not American.
- leptenoides*, Emmons, 1842, Geo. Rep. N. Y., p. 398, Trenton Gr.
- leucosia*, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 80, and Pal. N. Y., vol. 4, p. 48, Ham. Gr.
- limitaria*, see *Leiorhynchus limitaria*.
- livia*, Billings, 1860, Can. Jour. Ind., Sci. and Art, vol. 5, p. 267, Up. Held. Gr.
- lonensis*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 74, Trenton Gr.
- lucia*, Billings, 1874, Pal. Foss., vol. 2, p. 35, Gaspe Limestone No. 8, Devonian.
- lynx*, Eichwald, 1830, (*Terebratula lynx*.) Nat. Skizze von Podol., p. 202, and Pal. N. Y., vol. 1, p. 133, Trenton and Hud. Riv. Grs.
- maria*, Billings, 1862, Pal. Foss., vol. 1, p. 137, Anticosti Gr., Div. 1, Mid. Sil.
- macfarlanii*, Meek, 1848, Trans. Chi. Acad. Sci., vol. 1, p. 88, Ham. Gr.
- macleodi*, Whitfield, 1889, Bull. Am. Mus. Nat. Hist., vol. 2, p. 43, Calciferous Gr.
- media*, Shaler, 1865, Bull. No. 4, M. C. Z., p. 65, Anticosti Gr. This is probably only a variety of *O. elegantula*.
- media*, Winchell, 1880, Geo. Sur. Minn. 8th Rep., p. 64, Hud. Riv. Gr. The name was preoccupied.
- meeki*, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 20, Hud. Riv. Gr. A variety of *O. testudinaria*.
- merope*, Billings, 1862, Pal. Foss., vol. 1, p. 130, Trenton Gr.
- melchiori*, (*Terebratula melchiori*.) L'Eveille, 1835, Mem. Soc. Geol. France, vol. 2, p. 39, Subcarboniferous.
- melchiori* var. *burlingtonensis*, Hall, 1858, Geo. Rep. Iowa, p. 596, Burlington Gr.
- minna*, Billings, 1865, Pal. Foss., vol. 1, p. 303, Quebec Gr.
- minneapolis*, Winchell, 1880, Geo. Sur. Minn., 8th Rep., p. 63, Hud. Riv. Gr.
- missouriensis*, Shumard, 1855, Geo. Rep. Mo., p. 205, Up. Sil.
- missouriensis*, Swallow, 1860, Trans. St. Louis Acad. Sci. This name was preoccupied.
- mitis*, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 34, Schoharie grit.
- morrowensis*, James, not defined so as to be recognized.
- multisecta*, Meek, 1873, Ohio Pal., vol. 1, p. 112, Hud. Riv. Gr.
- multistriata*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 45, and Pal. N. Y., vol. 3, p. 176, Low. Held. Gr.
- musculosa*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 43, Oriskany sandstone.
- mycale*, Billings, 1862, Pal. Foss., vol. 1, p. 82, Quebec Gr.
- nisis*, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 181, Niagara Gr.
- nucleus*, Hall, syn. for *Ambocelia umbonata*.
- oblata*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 41, and Pal. N. Y., vol. 3, p. 162, Low. Held. Gr.

FIG. 591.—*Orthis lynx*. Small Trenton specimen.

oblate var. *emarginata*, Hall, 1859, Pal. N. Y., vol. 3, p. 164, Low. Held. Gr.
occusus, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 111, Waverly Gr.
occidentalis, Hall, 1847, Pal. N. Y., vol. 1, p. 127, Trenton to Hud. Riv. Gr.
orbicularis, Sowerby, 1839, Murch. Sil. Sys., p. 611, Up. Sil.
orthambonites, Eichwald, 1840, Sil. Syst. in Esthl., p. 150, Quebec Gr.
pecosi, Marcou, 1858, Geo. N. America, p. 48, Coal Meas. This species was subsequently described by Swallow under the name of *Orthis carbonaria*.
pecten, as identified by d'Archiac & Verneuil. Not American.
pectinella, Emmons, 1842, Geo. Sur. 2d Dist. N. Y., p. 394, and Pal. N. Y., vol. 1, p. 123, Trenton Gr.
pectinella var. *semiovalis*, Hall, 1847, Pal. N. Y., vol. 1, p. 124, Trenton Gr. Not distinguishable from the type species.
peduncularis, Hall, 1859, Pal. N. Y., vol. 3, p. 174, Low. Held. Gr.
peloris, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 32, Schoharie grit.
penelope, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 79, and Pal. N. Y., vol. 4, p. 50, Ham. Gr.
pepina, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 135, Potsdam Gr.
perelegans, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 44, and Pal. N. Y., vol. 3, p. 171, Low. Held. Gr.
perversa, see *Streptorhynchus perversum*.
perveta, Conrad, 1843, Proc. Acad. Nat. Sci., vol. 1, p. 333, and Pal. N. Y., vol. 1, p. 120, Black Riv. and Trenton Grs.
pigra, Billings, 1859, Can. Nat. Geo., vol. 4, p. 442, Chazy Gr.
pisum, as identified by Hall, see *Nucleospira pisiformis*.
planoconvexa, Hall, 1859, Pal. N. Y., vol. 3, p. 168, Low. Held. Gr.
platys, Billings, 1859, Can. Nat. Geo., vol. 4, p. 438, Chazy Gr.
plicata, Vanuxem, see *Spirifera vanuxemi*.
plicatella, Hall, 1847, Pal. N. Y., vol. 1, p. 122, Trenton and Hud. Riv. Grs.
pogonipensis, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 232, Chazy Gr.
porcata, McCoy, 1844, Sil. Foss. of Ireland, p. 32, Trenton, [Hud. Riv., and Mid. Sil.
porcia, Billings, 1859, Can. Nat. Geo., vol. 4, p. 439, Chazy Gr.
præumbona, see *Ambocoelia præumbona*.
pratteni, McChesney, 1860, New Pal. Foss., Coal Meas. Not recognized.
prava, Hall, 1858, Geo. of Iowa, p. 490, Ham. Gr.
propinqua, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 110, and Pal. N. Y., vol. 4, p. 43, Up. Held. Gr.
proximus, Vanuxem, 1842, (Hipparionyx proximus,) Geo. Rep. 3d Dist. N. Y., p. 124, and Pal. N. Y., vol. 3, p. 407, Oriskany sandstone.

punctostriata, Hall, 1852, Pal. N. Y., vol. 2, p. 254, Niagara Gr.
pyramidalis, see *Skenidium pyramidalis*.
quacoensis, Matthew, 1885, Trans. Roy. Soc. Can., p. 43, St. John Gr.
quadriradiata, see *Leiorhynchus quadriradiatus*.
remnicha, Winchell, 1886, 14th Ann. Rep. Geo. Minn., p. 317, Potsdam Gr.
resupinata, Martin, 1809, Petref. Derb., tab. 49, figs. 13 and 14, Subcarb.
resupinoides, Cox, 1857, Geo. Sur. Ky., vol. 3, p. 570, Coal Meas.
retrorsa, Salter, 1858, Mem. Geo. Sur. of Gt. Brit., vol. 2, p. 373, Trenton and Hud. Riv. Grs.
rhynchonelliformis, Shaler, 1865, Bull. No. 4, M. C. Z., p. 66, Anticosti Gr.
richmondi, McChesney, 1860, New Pal. Foss., p. 32, syn. for *Streptorhynchus crassum*.
robusta, Hall, 1858, Geo. Rep. Iowa, p. 713, syn. for *Streptorhynchus crassum*.
rugiplicata, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 182, Niagara Gr.
ruida, Billings, 1866, Catal. Sil. Foss. Antic., p. 42, Anticosti Gr.
saltemensis, Walcott, 1887, Am. Jour. Sci. and Arts, 3d ser., vol. 34, p. 190, Up. Taconic.
sandbergi, Winchell, 1886, 14th Ann. Rep. Geo. Minn., p. 318, Potsdam Gr.
schohariensis, Castelnau, 1843, Syst. Sil., p. 36. Not recognized.
scovillii, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 40, Hud. Riv. Gr.
sectostriata, Ulrich, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 15, syn. for *O. ella*.
semele, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 34, Onondaga and Up. Held. Grs.
sinuata, Hall, 1847, Pal. N. Y., vol. 1, p. 128, Hud. Riv. Gr.
sola, Billings, 1866, Catal. Sil. Foss. Antic., p. 12, Hud. Riv. Gr.
solitaria, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 80, and Pal. N. Y., vol. 4, p. 45, Ham. Gr.
stonensis, Safford, 1869, Geo. of Tenn., p. 286, Trenton and Nashville Grs.
striatella, see *Chonetes striatellus*.
striatula, Emmons, 1842, Geo. Rep. N. Y. This name was preoccupied by Schlottheim.
strophomenoides, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 46, and Pal. N. Y., vol. 3, p. 177, Low. Held. Gr.
subcarinata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 42, and Pal. N. Y., vol. 3, p. 169, Low. Held. Gr.
subequata, Conrad, 1843, Proc. Acad. Nat. Sci., vol. 1, p. 333, and Pal. N. Y., vol. 1, p. 118, Chazy to Trenton Gr.
subelliptica, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 202, Waverly or Kinderhook Gr.
subjugata, syn. for *Orthis occidentalis*.

subnotosa, Hall, 1879, Desc. New Spec. Foss., p. 14, and 11th Rep. Geo. and Nat. Hist. Ind., p. 286, Niagara Gr.

suborbicularis, Hall, 1858, Geo. of Iowa, p. 480, Ham. Gr.

subquadrata, Hall, 1847, Pal. N. Y., vol. 1, p. 120, Trenton to Hud. Riv. Gr.

subumbona, see *Martinia subumbonata*.

swallowi, Hall, 1858, Geo. Rep. Iowa, p. 597, Burlington Gr.

tenuidena, Hall, 1852, Pal. N. Y., vol. 2, p. 58, Clinton Gr.

tenuistriata, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 244, Portage Gr. The name was preoccupied by Sowerby.

testudinaria, Dalman, 1827, Vet. Acad. Hand., p. 116, and Pal. N. Y., vol. 1, p. 117, Trenton and Hud. Riv. Grs.

thiemii, White, 1860, Jour. Bost. Soc. Nat. Hist., vol. 7, p. 231, and Cont. to Pal. No. 8, p. 164, Kinderhook Gr.

tioa, Hall, 1867, Pal. N. Y., vol. 4, p. 59, (O. interlineata, Sow.) Geo. Rep. 4th Dist. N. Y., Portage and Chemung Grs.

tricenaria, Conrad, 1843, Proc. Acad. Nat. Sci., vol. 1, p. 333, and Pal. N. Y., vol. 1, p. 121, Trenton Gr.

trinucleus, Hall, 1852, Pal. N. Y., vol. 2, p. 58, Clinton Gr.

triplicatella, Meek, 1873, Ohio Pal., vol. 1, p. 109, Hud. Riv. Gr.

tritonina, Billings, 1862, Pal. Foss., vol. 1, p. 76, Quebec Gr.

tubulostriata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 42, and Pal. N. Y., vol. 3, p. 166, Low. Held. Gr.

tulliensis, Vanuxem, 1843, Geo. Rep. 3d Dist. N. Y., p. 163, and Pal. N. Y., vol. 4, p. 55, Tully limestone.

uberis, Billings, 1866, Catal. Sil. Foss. Antic., p. 42, Anticosti Gr.

umbonata, see *Ambocelia umbonata*.

umbraculum, DeKoninck, see *Orthis keokuk* and *Streptorhynchus umbraculum*.

unguiculus, Phillips, as identified by Hall in 1843, see *Ambocelia gregaria*.

unguiformis, Castlenau, 1843, Syst. Sil., p. 37, syn. for *Orthis hippurionyx*.

vanuxemi, Hall, 1857, 10 Rep. N. Y. St. Mus. Nat. Hist., p. 135, and Pal. N. Y., vol. 4, p. 47, Ham. Gr.

vanuxemi, Winchell, 1862, Proc. Acad. Nat. Sci., vol. 6, 2d ser., p. 409, Portage Gr. The name was preoccupied.

varica, Conrad, 1842, (Delthyris varica,) Jour. Acad. Nat. Sci., vol. 8, p. 262, and Pal. N. Y., vol. 3, p. 179, Low. Held. Gr.

ORTHISINA, D'Orbigny, 1850, Prodr. d. Pal., vol. 1, p. 16. [Ety. *Orthis*, a genus; *inus*, implying resemblance to.] External characters of *Orthis*, but the triangular pit in the cardinal area of the ventral valve is closed by a cicatrix with an oval perforation near the apex; interior

of ventral valve with two broad, dental lamellae bordering the cardinal pit, and converging to a mesial line at the surface of the shell; interior of dorsal valve with a trifid rostral tooth, from which a small mesial septum extends toward the margin; lateral cardinal teeth as in *Orthis*. Type *O. verneuili*.

alternata, see *Streptorhynchus perversum*.

arctostriata, see *Streptorhynchus arctostriatum*.

crassa, see *Streptorhynchus crassum*.

diversa, Shaler, syn. for *Orthisina verneuili*.

festinata, Billings, 1861, Pal. Foss., vol. 1, p. 10, Georgia Gr.

grandæva, Billings, 1859, Can. Nat. Geo., vol. 4, p. 349, Calif. Gr.

missouriensis, Swallow, 1858, syn. for *Meekella striatorostata*.

occidentalis, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 82, Up. Coal Meas.

orientalis, Whitfield, 1884, Bull. Am. Mus. Nat. Hist., vol. 1, p. 139, Georgia Gr.

shumardana, Swallow, 1858, Trans. St. Louis Acad. Sci., p. 183, Permian Gr.

transversa, Walcott, 1886, Bull. U.S. Geo. Sur., No. 30, p. 121, Up. Taconic.

verneuili Eichwald, 1842, (Orthis verneuili,) Urwelt Russie, vol. 2, p. 51, Trenton and Anticosti Gr.

Pentagonia, Cozzens, 1846, Ann. N. Y. Lyceum, vol. 4, p. 158. [Ety. *pente*, five; *gonia*, an angle.] This genus seems to have been founded upon Conrad's *Atrypa unisulcata*, which is now referred to *Meristella*. The genus is not recognized by authors.

peersi, Cozzens, 1846, Ann. N. Y. Lyceum, vol. 4, p. 158, syn. for *Meristella unisulcata*.

FIG. 504.—*Pentamerella arata*. Dorsal view.

PENTAMERELLA, Hall, 1867, Pal. N. Y., vol. 4, p. 375. [Ety. diminutive of *Pentamerella*.]

merus.] Ventral valve gibbous, beak incurved, fissure triangular, area narrow, mesial sinus; in the interior an elongate, spoon-shaped pit, the upper part supported on a central septum; dorsal valve convex, mesial fold; crura conjoined at their bases, making a V-shaped pit, which is attached to the valve in its upper part, and continues sessile for about half the length of the shell; surface plicated. Type *P. arata*.

FIG. 505.—*Pentamerella arata*. Side view.

arata, Conrad, 1841, (*Atrypa arata* and *Atrypa octo-costata*.) Ann. Rep. N. Y., p. 55, and Pal. N. Y., vol. 4, p. 375, Schöhariegrit and Up. Held. Gr. *compressa*, Ringueberg, 1880, Bull. Nat. Soc. Nat. Sci., vol. 5, p. 15, Niagara Gr. *dubia*, Hall, 1860, (Spirifer dubius.) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 90, Ham. Gr. *micula*, Hall, 1867, Pal. N. Y., vol. 4, p. 378, Ham. Gr. *obsolescens*, Hall, 1867, Pal. N. Y., vol. 4, p. 379, Devonian. *papilionensis*, Hall, 1858, (*Pentamerus papilionensis*.) Geo. Rep. Iowa, vol. 1, pt. 2, p. 514, Ham. Gr.

PENTAMERUS, Sowerby, 1812, Min. Conch., vol. 1, p. 73. [Ety. *penta*, five; *meros*, apartments.] Shell globose, ovate, receiving valve largest; generally destitute of mesial fold and sinus, but when present, the fold is in the receiving, and the sinus in the entering valve; no hinge-line; area large, undefined, and having a deep, triangular pit in the center, under the beak of the dorsal valve, and into which the beak of the entering valve is strongly incurved; internally the receiving valve has one large bipartite central septum, the walls of which suddenly divaricate as they approach the entering valve, forming the walls of the external triangular opening, and inclosing between them a triangular chamber much smaller than the two lateral ones; in the entering valve the two corresponding plates are subparallel, and separate from their origin, being so curved that internal casts show one of their edges, like the diverging cardinal teeth of *Orthis*, and the inner edges form the long, subparallel slits, the middle one of the three resulting chambers being much the narrower. Type *P. knighti*.

aratus, see *Pentamerella arata*.

arcuosus, McChesney, 1861, New Pal. Foss., p. 87, Niagara Gr. Not recognized.

barrandi, Billings, 1857, Rep. of Progr., Geo. Sur. Can., p. 290, Mid Sil.

beumonti, Castelnau, 1843, Syst. Sil., p. 38. Not recognized.

bisinuatus, McChesney, 1859, New Pal. Foss., p. 85, and Trans. Chi. Acad. Sci., vol. 1, p. 30, Niagara Gr.

borealis, Meek, 1868, Trans. Chi. Acad. Sci., p. 95, Ham. Gr. This name was preoccupied by Eichwald in 1840.

brevirostris, Sowerby, 1839, (Terebratula brevirostris.) Murch. Sil. Syst., p. 631, and Pal. N. Y., vol. 2, p. 278, Niagara Gr.

chicagoensis, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 94, Niagara Gr.

comis, Owen, 1852, (*Atrypa comis*.) Geo. Sur. Wis., Iowa and Minn., p. 583, Ham. Gr.

copplingeri, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 593, U. p. Silurian.

crassiradiatus, McChesney, 1861, New Pal. Foss., p. 87, Niagara Gr. Not recognized.

deshayesi, Castelnau, 1843, Syst. Sil., p. 38. Not recognized. Probably syn. for *Amphigenia elongata*.

elongatus, see *Amphigenia elongata*.

fornicatus, Hall, 1852, Pal. N. Y., vol. 2, p. 81, Clinton Gr.

galeatiformis, Meek & Worthen, syn. for *P. galeatus*.

galeatus, Dalman, 1827, (*Atrypa galeatus*.) Vet. Acad. Handl., p. 130, and Pal. N. Y., vol. 3, p. 257, Low. Held. Gr.

intralineatus, Winchell, 1866, Rep. Low. Penin. Mich., p. 94, Ham. Gr.

knappi, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 184, Niagara Gr.

knighti, Sowerby, 1812, Min. Conch., vol. 1, p. 73, Devonian. *laqueatus*, Conrad, 1855, Proc. Acad. Nat. Sci., p. 441, Niagara Gr.

lenticularis, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 295, Kinderhook Gr.

littoni, Hall, 1859, Pal. N. Y., vol. 3, p. 262, Low. Held. and Niagara Gr.

lotis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 161, Devonian.

multicostatus, Hall, 1861, Rep. of Progr. Wis. Sur., p. 1, Niagara Gr.

nucleus, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 200, Niagara Gr.

FIG. 506.—*Pentamerus knighti*.

Dorsal view.

l. N. Y., vol. 1, p. 51, of *Pentamerus*.

nysius, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 184, Niagara Gr. There are two varieties, one having coarse and the other finer radii. These are designated *P. nysius* var. *crassicostus* and *P. nysius* var. *tenuicostus*.

oblongus, Sowerby, 1839, Murch. Sil. Syst., p. 641, and Pal. N. Y., vol. 2, p. 79, Clinton and Niagara Gr.

oblongus var. *cylindricus*, Hall & Whitfield, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 183, Niagara Gr.

occidentalis, Hall, 1852, Pal. N. Y., vol. 2, p. 341, Guelph Gr.

occidentalis, see *Gypidula occidentalis*.

ovalis, Hall, 1852, Pal. N. Y., vol. 2, p. 103, Clinton Gr.

papilionensis, see *Pentamerella papilionensis*.

pergibbosus, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 139, Niagara Gr.

pesovis, Whitfield, 1882, Desc. New Spec. Foss., from Ohio, p. 195, Low. Held. Gr.

pseudogaleatus, Hall, 1857, 10th. Rep. N. Y. St. Mus. Nat. Hist., p. 106, and Pal. N. Y., vol. 3, p. 259, Low. Held. Gr.

reversus, see *Anastrophia reversa*.

salinensis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 652, Devonian.

similior, Winchell & Marcy, 1865, (Spirifer similior,) Mem.

Bost. Soc. Nat. Hist., p. 93, Niagara Gr.

subglobosus, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 429, Ham. Gr.

trisinuatus, McCheyney, 1861, Desc. New Pal. Foss., p. 86, Niagara Gr.

ventricosus, Hall, 1861, Rep. Progr. Wis. Sur., p. 2, and 20th Rep. N. Y. St. Mus. Nat. Hist., p. 374, Niagara Gr.

verneuili, see *Anastrophia verneuili*.

PHOLIDOPS, Hall, 1859, Pal. N. Y., vol. 3, p. 489. [Ety. *pholis*, *pholidos*, a scale.]

Small, thin, subelliptical, inequivalve; apex excentric, foramen in front of the apex of the ventral valve; surface marked by concentric lamellae of growth; dorsal valve marked with bilobed muscular impressions. Type *P. squamiformis*.

arenaria, Hall, 1867, Pal. N. Y., vol. 4, p. 413, Oriskany sandstone.

areolata, Hall, 1863, 16th Rep. N. Y. Mus. Nat. Hist., p. 31, Schoharie grit.

bellula, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 113, Devonian.

cincinnatiensis, Hall, 1872, 24th Rep. N. Y. St. Mus. Nat. Hist., pl. 7, fig. 10, Hud. Riv. Gr.

hamiltoniae, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 92, and Pal. N. Y., vol. 4, p. 32, Ham. Gr.

linguloides, Hall, 1867, Pal. N. Y., vol. 4, p. 414, Ham. Gr.

oblata, Hall, 1867, Pal. N. Y., vol. 4, p. 414, Ham. Gr.

ovalis, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 209, Niagara Gr.

ovata, Hall, 1859, Pal. N. Y., vol. 3, p. 490, Low. Held. Gr.

quadrangularis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 114, Devonian.

squamiformis, Hall, 1843, (Orbicula squamiformis,) Geo. Rep. 4th Dist. N. Y., p. 108, and Pal. N. Y., vol. 2, p. 250, Niagara Gr.

subtruncata, Hall, 1847, (Orbicula subtruncata,) Pal. N. Y., vol. 1, p. 290, Hud. Riv. Gr.

terminalis, Hall, 1859, Pal. N. Y., vol. 3, p. 490, Oriskany sandstone.

trentonensis, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 221, Trenton Gr.

Platystrophia, King, syn. for *Orthis*.

regularis, syn. for *Orthis lynx*.

Plectambonites areca, syn. for *Leptæna transversalis*.

glabra, syn. for *Leptæna sericea*.

tenes, a, syn. for *Leptæna transversalis*.

Plicatula, Lamarck, 1809. Not Palæozoic.

striatocostata, see *Meekella striatocostata*.

PORAMBONITES, Pander, 1830, Beiträge zur Geog. des Russischen Reiches, p. 99.

linguloides, Hall, 1867, Pal. N. Y., vol. 4, p. 414, Ham. Gr.

oblata, Hall, 1867, Pal. N. Y., vol. 4, p. 414, Ham. Gr.

ovalis, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 209, Niagara Gr.

ovata, Hall, 1859, Pal. N. Y., vol. 3, p. 490, Low. Held. Gr.

quadrangularis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 114, Devonian.

squamiformis, Hall, 1843, (Orbicula squamiformis,) Geo. Rep. 4th Dist. N. Y., p. 108, and Pal. N. Y., vol. 2, p. 250, Niagara Gr.

subtruncata, Hall, 1847, (Orbicula subtruncata,) Pal. N. Y., vol. 1, p. 290, Hud. Riv. Gr.

terminalis, Hall, 1859, Pal. N. Y., vol. 3, p. 490, Oriskany sandstone.

trentonensis, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 221, Trenton Gr.

Platystrophia, King, syn. for *Orthis*.

regularis, syn. for *Orthis lynx*.

Plectambonites areca, syn. for *Leptæna transversalis*.

glabra, syn. for *Leptæna sericea*.

tenes, a, syn. for *Leptæna transversalis*.

Plicatula, Lamarck, 1809. Not Palæozoic.

striatocostata, see *Meekella striatocostata*.

PORAMBONITES, Pander, 1830, Beiträge zur Geog. des Russischen Reiches, p. 99.

FIG. 598.—*Porambonites ottawensis*. a, b, c, d, Different views; e, interior of ventral valve; f, interior of dorsal valve; g, showing oval arims.

FIG. 597.—*Pholidops cincinnatiensis*.

[Ety. *poros*, opening; *ambon*, umbone.]

Subglobose, depressed, dorsal valve the larger, beaks obtuse, subequal, separated by a small cardinal area in each valve; foramen in each valve small, triangular, reaching the hinge-line; two long, slightly diverging dental lamellae in each valve, those of the ventral valve closer together; surface coarsely punctured in lines. Type *P. aequirostris*.

dentatus, see *Orthis dentatus*.

obscurus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 234, Quebec Gr.

ottawensis, Billings, 1862, Pal. Foss., vol. 1, p. 140, Black Riv. Gr.

PRODUCTELLA, Hall, 1867, Pal. N. Y., vol. 4, p. 153. [Sig. diminutive of *Productus*.]

Shells having the general form of *Productus*, but with a narrow area on each valve, a foramen or callosity on the ventral area, small teeth, and more or less distinct tooth-sockets. Type *P. subaculeata*.

arctirostrata, Hall, 1857, (Productus arctirostratus,) 10th Rep. N. Y. St. Mus.

hirsut., 10th Rep. N. Y. St. Mus.

hirsut., 175, N. Y. St. Mus.

hystri., p. 17, N. Y. St. Mus.

lachry., 8, p. 4, N. Y. St. Mus.

lachry., 8, p. 4, N. Y. St. Mus.

lachry., 8, p. 4, N. Y. St. Mus.

lachry., 8, p. 4, N. Y. St. Mus.

lachry., 8, p. 4, N. Y. St. Mus.

lachry., 8, p. 4, N. Y. St. Mus.

lachry., 8, p. 4, N. Y. St. Mus.

Y., vol. 4,
vol. 4, p.
Inst., vol.
vol. 3, p.

Monogr.
Devonian.
cula squa-
st. N. Y.,
2, p. 250,
Y., vol. 3,
ep. N. Y.
enton Gr.
s.

ena trans-

rsalis.
Paleozoic.
atocostata.
itragr zur
es, p. 90.

ews; e, in-
flowing oval

umbone.]
valve the
ual, sepa-
ea in each
ve small,
hinge-line;
dental la-
f the ven-
; surface
Type P.

377, U. S.
4, p. 234,

Foss., vol.

Y., vol. 4,
Productus.]
rm of Pro-
rea on each
ity on the
id more or
Type P.

uctus arcti-
St. Mus.

Nat. Hist., p. 177, and Pal. N. Y., vol. 4, p. 182, Chemung Gr.
bivalveata, Hall, 1867, Pal. N. Y., vol. 4, p. 183, Chemung Gr.
boydii, Hall, 1857, (Productus boydii,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 179, and Pal. N. Y., vol. 4, p. 169, Chemung Gr.
concentrica, Hall, 1857, (Productus concentrica,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 180, Kinderhook Gr.
costatula, Hall, 1867, Pal. N. Y., vol. 4, p. 180, Chemung Gr.
costatula var. strigata, Hall, 1867, Pal. N. Y., vol. 4, p. 181, Chemung Gr.
dissimilis, see P. hallana.
dumosa, Hall, 1861, (Productus dumosus,) 14th Rep. N. Y. St. Mus. Nat. Hist., p. 99, and Pal. N. Y., vol. 4, p. 162, Ham. Gr.
eriensis, Nicholson, 1874, Geo. Mag., n. s., vol. 1, p. 118, Cornif. Gr.
exanthemata, Hall, 1857, (Productus exanthematus,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 174, and Pal. N. Y., vol. 4, p. 163, Ham. Gr.
hallana, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 130, Ham. Gr. Proposed instead of P. dissimilis of Hall, which was preoccupied by DeKoninck.

FIG. 599.—Productella hirsuta.

hirsuta, Hall, 1857, (Productus hirsutus,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 175, Chemung Gr.
hirsuta var. rectispina, Hall, 1867, Pal. N. Y., vol. 4, p. 168, Chemung Gr.
hystricula, Hall, 1867, Pal. N. Y., vol. 4, p. 178, Chemung Gr.
lachrymosa, Conrad, 1842, (Strophomena lachrymosa,) Jour. Acad. Nat. Sci., vol. 8, p. 256, and Pal. N. Y., vol. 4, p. 174, Chemung Gr.
lachrymosa var. lima, Conrad, 1842, (Strophomena lima,) Jour. Acad. Nat. Sci., vol. 8, p. 256, and Pal. N. Y., vol. 4, p. 174, Chemung Gr.
lachrymosa var. stigmata, Hall, 1867, Pal. N. Y., vol. 4, p. 174, Chemung Gr.
navicella, Hall, 1857, (Productus navicella,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 172, and Pal. N. Y., vol. 4, p. 156, Cornif. and Ham. Grs.
newberryi, Hall, 1857, (Productus newberryi,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 180, Chemung Gr.
onusta, Hall, 1867, Pal. N. Y., vol. 4, p. 184, Chemung Gr.
pyxidata, Hall, 1858, (Productus pyxidatus,) Geo. of Iowa, p. 498, Ham. Gr.

rarisipina, Hall, 1857, (Productus rarisipinus,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 178, and Pal. N. Y., vol. 4, p. 170, Chemung Gr.
shumardana, Hall, 1858, (Productus shumardanus,) Geo. Rep. of Iowa, vol. 1, pt. 2, p. 499, and Pal. N. Y., vol. 4, p. 157, Up. Held. Gr., Marcellus shale, Ham. and Chemung Grs.
speciosa, Hall, 1857, (Productus speciosus,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 176, and Pal. N. Y., vol. 4, p. 175, Chemung Gr.
spinulicosta, Hall, 1857, (Productus spinulicostus,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 173, and Pal. N. Y., vol. 4, p. 160, Marcellus shales and Ham. Gr.
striatula, Hall, 1867, Pal. N. Y., vol. 4, p. 177, Chemung Gr.
subaculeata, Murchison, 1840, (Productus subaculeatus,) Bul. Soc. Geo. de France, vol. 11, p. 255, and Pal. N. Y., vol. 4, p. 154, Waverly Gr.
subalata, Hall, 1857, (Productus subalata,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 174, and Pal. N. Y., vol. 4, p. 165, Ham. Gr.
truncata, Hall, 1857, (Productus truncatus,) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 171, and Pal. N. Y., vol. 4, p. 160, Marcellus shales and Ham. Gr.
tullia, Hall, 1867, Pal. N. Y., vol. 4, p. 164, Ham. Gr.

FIG. 600.—Productella subalata. Ventral valve.

PRODUCTUS, Sowerby, 1812, Min. Conch., vol. 1, p. 153. [Ety. productus, produced—so named from one valve of the shell being prolonged beyond the other, and often to a great extent.] Shell inequivalve, transverse, or elongated with auricular expansions; ventral valve convex, geniculated, or perpendicularly incurved; hinge-line straight; area narrow, or the cardinal edge thickened; beak incurved; in the interior a narrow mesial ridge separates two elongated, ramified, muscular adductor scars; under and outside these are two deep, longitudinally subquadrate impressions for cardinal muscles, widely separated by a crest, and lower down toward the center of the shell two deep concave subspiral depressions for spiral or labial appendages; dorsal valve concave, following the other valve; cardinal process for the attachment of muscles prominent, trifid, and below it a mesial ridge, upon each side of which are the ramified adductor scars; outside and in front of these are two reniform impressions; a prominence on each side the mesial ridge indicates the origin of spiral arms; surface of shell striated, more or less concentrically

- wrinkled, and bearing tubular spines. Types *P. longispinus* and *P. semireticulatus*.
- aequicostatus*, Shumard, 1855, Geo. Rep. Mo., p. 201, Coal Meas.
- alternatus*, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., 2d series, vol. 3, p. 20, Keokuk Gr.
- altonensis*, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., 2d series, vol. 3, p. 7, Kaskaskia Gr.
- americanus*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 91, Up. Coal Meas.
- arctirostratus*, see *Productella arctirostrata*.
- arcuatus*, Hall, 1858, Geo. Rep. Iowa, p. 518, Kinderhook Gr.
- asper*, McChesney, syn. for *P. nebrascensis*.
- auriculatus*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 92, Coal Meas.
- biseriatus*, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 12, and Bull. Am. Mus. Nat. Hist., p. 46, Warsaw Gr.
- boonensis*, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 217, Coal Meas.
- boydi*, see *Productella boydi*.
- calhounianus*, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 215, Coal Meas. Prof. Meek regarded this name as a synonym for *P. semireticulatus*.
- callawayensis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 640, Ham. Gr.
- cancrini*, as identified by Geinitz, is *P. pertenuis* of Meek.
- capaci*, D'Orbigny, 1843, as identified by early authors, is referred to *P. longispinus*.
- cestriensis*, Worthen, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 570, Kaskaskia Gr.
- clavus*, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., 2d series, vol. 3, p. 10, Coal Meas.
- comoides*, as identified by d'Archiac & Vernel. Not American.
- concentricus*, see *Productella concentrica*.
- confragosus*, Conrad, 1835, Trans. Geo. Soc. Penn., vol. 1, p. 2, p. 267, Coal Meas. This species is not recognized.
- cooperensis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 640, Waverly or Choteau Gr.
- cora*, D'Orbigny, 1842, Paléont. d. l'Am. Merid., p. 48, Coal Meas.
- cora* var. *mogoyoni*, Marcou, 1858, Geo. N. Amer., p. 45, Subcarboniferous.
- coriformis*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 94, Kaskaskia Gr.
- costatoides*, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 217, Up. Coal Meas.
- costatus*, Sowerby, 1827, Min. Conch., vol. 6, p. 115, Coal Meas. It is doubtful whether this species has been identified in America.
- curtirostratus*, Winchell, 1865, Proc. Acad. Nat. Sci., p. 114, Marshall Gr.
- delawari*, Marcou, 1858, Geol. N. Amer., p. 45, Subcarb.
- depressus*, Sowerby, 1825, see *Strophomena depressa*.
- depressus*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 93, Keokuk Gr.
- dissimilis*, see *Productella dissimilis*.
- dolorosus*, Winchell, 1865, Proc. Acad. Nat. Sci., p. 114, Marshall Gr.
- dumosus*, see *Productella dumosa*.
- duplicostatus*, Winchell, 1865, Proc. Acad. Nat. Sci., p. 113, Marshall Gr.
- elegans*, Norwood & Pratten, 1854. This name was preoccupied, and the fossil is now named *P. cestriensis*.
- exanthematus*, see *Productella exanthemata*.
- fasciculatus*, McChesney, 1860, New Pal. Foss., Coal Meas. Not recognized.
- fentonensis*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 93, Keokuk Gr.
- flemingi*, Sowerby, 1812, Min. Conch., vol. 1, p. 155, Subcarb.
- flemingi* var. *burlingtonensis*, Hall, 1858, Geo. Rep. Iowa, p. 598, Burlington Gr.
- gracilis*, Winchell, 1865, Proc. Acad. Nat. Sci., p. 112, Cuyahoga shale.
- gradatus*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 93, Keokuk Gr.
- hepar*, Morton, 1836, Am. Jour. Sci. and Arts, vol. 29, p. 149, Coal Meas. Not recognized.
- hildrethianus*, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., 2d ser., vol. 3, p. 18, Coal Meas.
- hirsutiformis*, Walcott, 1895, Monogr. U. S. Geo. Sur., vol. 8, p. 133, Up. Devonian.
- hirsutus*, see *Productella hirsuta*.
- horridus*, as identified by Geinitz, 1866, Prof. Meek regarded the fossil as *P. longispinus*.
- incurvatus*, Shepard, 1838, Am. Jour. Sci., vol. 34, p. 144. Not recognized. Probably a *Streptorhynchus* or *Strophodonta*.
- indianensis*, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 13, and Bull. Am. Mus. Nat. Hist., vol. 4, p. 47, Warsaw Gr.
- inflatus*, syn. for *P. semireticulatus*.
- ivesi*, Newberry, 1861, Ives's *U. S. Ex. Exped.*, p. 122, Mid. Carb.
- lasallensis*, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 569, Up. Coal Meas.
- laevicostus*, White, 1860, Bost. Jour. Nat. Hist., vol. 7, p. 230, Kinderhook Gr.
- latissimus*, Sowerby, 1822, Min. Conch., vol. 4, p. 32, Carb.

FIG. 601.—*Productetus longispinus*. Dorsal and ventral views.

longispinus, Sowerby, 1812, Min. Conch., vol. 1, p. 154, Coal Meas.

- Proc. Acad. Nat. Sci. Phil.*, p. 142, and *Geo. Sur. Ill.*, vol. 3, p. 528, Keokuk Gr.
- magnicostatus*, Swallow, 1860, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 641, Coal Meas.
- magnus*, Meek & Worthen, 1861, *Proc. Acad. Nat. Sci. Phil.*, p. 142, and *Geo. Sur. Ill.*, vol. 3, p. 528, Keokuk Gr.
- marginicinctus*, Prout, 1857, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 43, St. Louis Gr.
- mesialis*, Hall, 1858, *Geo. Rep. Iowa*, p. 636, Keokuk Gr.
- mexicanus*, Shumard, 1858, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 291, Permian Gr.
- morbillianus*, Winchell, 1865, *Proc. Acad. Nat. Sci.*, p. 113, Burlington Gr.
- multistriatus*, Meek, 1860, *Proc. Acad. Nat. Sci.*, vol. 12, p. 309, and Simpson's *Rep. Gt. Basin of Utah*, p. 350, Coal Meas.
- muricatus*, Norwood & Pratten, 1854, *Jour. Acad. Nat. Sci. Phil.*, vol. 3, p. 14, Coal Meas. Prof. Meek regarded this as a syn. for *P. longispinus*.
- nanus*, Meek & Worthen, 1860, *Proc. Acad. Nat. Sci.*, p. 450, and *Geo. Sur. Ill.*, vol. 2, p. 320, Coal Meas.
- navicella*, see *Productella navicella*.
- nebraskensis*, Owen, 1852, *Geo. Rep. Wis., Iowa, and Minn.*, p. 584, Coal Meas.
- nevadensis*, Meek, 1877, *U. S. Geo. Sur.* 40th parallel, p. 64, Carboniferous.
- nodosus*, Newberry, 1861, *Ives' Col. Ex. Exped.*, p. 124, Carb.
- norwoodi*, Swallow, 1858, *Trans. St. Louis Acad. Sci.*, p. 182, Permian Gr.
- occidentalis*, Newberry, 1861, *Ives' Col. Ex. Exped.*, p. 122, Up. Carb.
- orbignyianus*, DeKoninck, 1847, *Mon. du genre Productus*, p. 152, Up. Coal Meas.
- ovatus*, Hall, 1858, *Geo. Rep. Iowa*, p. 674, St. Louis Gr.
- parvulus*, Winchell, 1863, *Proc. Acad. Nat. Sci.*, p. 4, Marshall Gr.
- parvus*, Meek & Worthen, 1860, *Proc. Acad. Nat. Sci. Phil.*, p. 450, and *Geo. Sur. Ill.*, vol. 2, p. 297, Kaskaskia Gr.
- pectenoides*, Shepard, 1838, *Am. Jour. Sci.*, vol. 34, p. 150. Not recognized. Probably a *Streptorhynchus*.
- pertenuis*, Meek, 1872, *Pal. E. Neb.*, p. 164, Coal Meas.
- phillipsi*, Norwood & Pratten, 1854, *Jour. Acad. Nat. Sci.*, vol. 3, 2d series, p. 8, Subcarb.
- pileiformis*, syn. for *Productus cora*.
- pileolus*, Shumard, 1858, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 291, Permian Gr.
- poecilum*, Morton, 1836, *Am. Jour. Sci. and Arts*, vol. 29, p. 149, Coal Meas. Not recognized.
- popii*, Shumard, 1858, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 290, Permian Gr.
- portlockanus*, Norwood & Pratten, 1854, *Jour. Acad. Nat. Sci.*, 2d series, vol. 3, p. 15, Coal Meas.
- prattenanus*, Norwood, 1854, *Jour. Acad. Nat. Sci. Phil.*, 2d series, vol. 3, p. 17, Coal Meas.
- punctatus*, Martin, 1809, *Petrif. Derb.*, pl. 37, fig. 6, Low. Carb. and Coal Meas.
- pyxidatus*, see *Productella pyxidata*.
- pyxidiformis*, DeKoninck, 1847, *Monographie du genre Productus*, p. 220, Subcarboniferous.
- rarispinus*, see *Productella rarispinus*.
- rogersi*, Norwood & Pratten, 1854, *Jour. Acad. Nat. Sci.*, vol. 3, p. 9, Coal Meas. Prof. Meek regarded this as a synonym for *P. nebraskensis*.
- scabriculus*, (*Conchylolithus Anomites scabriculus*), Martin, 1809, *Petrif. Derb.*, p. 8, tab. 36, fig. 5, Carb.
- scitulus*, Meek & Worthen, 1860, *Proc. Acad. Nat. Sci.*, p. 451, and *Geo. Sur. Ill.*, vol. 2, p. 280, St. Louis Gr.
- semipunctatus*, Shepard, 1838, *Am. Jour. Sci.*, vol. 34, p. 153, Coal Meas.
- semipunctatus*, Hildreth, 1838, syn. for *P. punctatus*.
- semireticulatus*, Martin, 1809, (*Conchylolithus Anomites semireticulatus*), *Petrif. Derb.*, p. 7, Keokuk Gr.
- semistriatus*, Meek, 1860, *Proc. Acad. Nat. Sci.*, vol. 12, p. 309, and Simpson's *Rep. Gt. Basin of Utah*, p. 349, Coal Meas.
- setigerus*, Hall, 1858, *Geo. Rep. Iowa*, p. 638, Keokuk Gr.
- setigerus* var. *Keokuk*, Hall, 1858, *Geo. Rep. Iowa*, p. 639, Keokuk Gr.
- shumardanus*, see *Productella shumardana*.
- speciosus*, see *Productella speciosa*.
- spinulicostus*, see *Productella spinulicosta*.
- spinulosus*, Sowerby, 1812, *Min. Conch.*, vol. 1, p. 155, Carb.
- splendens*, Norwood & Pratten, 1854, *Jour. Acad. Nat. Sci. Phil.*, vol. 3, p. 11, Coal Meas. Prof. Meek regarded this as a synonym for *P. longispinus*.
- subaculeatus*, see *Productella subaculeata*.
- subalatus*, see *Productella subalata*.
- subhorridus*, Meek, 1877, *U. S. Geo. Sur.* 40th parallel, p. 75, Carboniferous.
- sulcatus*, Castelnau, 1843, *Syst. Sil.*, p. 39, Not recognized.
- symmetricus*, McChesney, 1860, *Desc. New Pal. Foss.*, p. 35, and *Pal. E. Neb.*, p. 167, Coal Meas.
- tennicostus*, Hall, 1858, *Geo. Rep. Iowa*, p. 675, St. Louis Gr.
- tenuistriatus*, Verneuil, 1845, *Geol. Russia and Ural Mountains*, vol. 2, p. 260, Carb.

FIG. 602.—*Productus semireticulatus*.

Dorsal and

Min. Conch.,

truncatus, see *Productella truncata*.
tubulospinus, McChesney. Syn. for. *P. semipunctatus*.

viminalis, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 29, Burlington Gr.

vittatus, Hall, 1858, Geo. Rep. Iowa, p. 639, Keokuk Gr.

wabashensis, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 13, Coal Meas.

wilberianus, McChesney, syn. for *P. nebraskensis*.

wortheni, Hall, 1858, Geo. Rep. Iowa, p. 635, Keokuk Gr.

PSEUDOCRANIA, McCoy, 1851, Ann. and Mag. Nat. Hist., 2d series, vol. 8, p. 387. [Ety. *pseudo*, false; *Crania*, a genus.] Shell slightly inequivalve, free; each valve depressed, subconical; dorsal valve with or without a small cardinal area; internally, margin broad, flat, smooth, or minutely striated concentrically; anterior pair of muscular impressions much larger than the posterior pair; pallial impressions numerous, linear, not interrupted along the middle. Type *P. divaricata*.

anomala, Winchell, 1866, Rep. Low. Pen. Mich., p. 92, Ham. Gr.

RENSELERIA, Hall, 1859, Pal. N. Y., vol. 3, p. 454. [Ety. proper name.] Inequivalve, oval, ovoid, or suborbicular, elongated, rarely transverse, sometimes subtrigonal, gibbous or ventricose; no mesial fold or sinus; beak prominent, incurved, foramen terminal; articulation by two widely separated teeth and sockets; surface striated; structure punctate. Type *R. ovoidea*.

aequiradiata, Conrad, 1842, (Atrypa aequiradiata.) Jour. Acad. Nat. Sci., vol. 8, p. 266, and Pal. N. Y., vol. 3, p. 255, Low. Held. Gr.

condoni, McChesney, 1861, New Pal. Foss., p. 85, Oriskany sandstone.

cumberlandia, Hall, 1857, (Meganteris cumberlandia.) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 101, and Pal. N. Y., vol. 3, p. 464, Oriskany sandstone.

elliptica, Hall, 1857, (Meganteris elliptica.) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 98, Low. Held. Gr.

elongata, see *Amphigenia elongata*.

intermedia, Hall, 1859, Pal. N. Y., vol. 3, p. 463, Oriskany sandstone.

johanni, Hall, 1867, Pal. N. Y., vol. 4, p. 385, Up. Held. Gr.

laevis, Hall, 1857, (Meganteris laevis.) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 99, Low. Held. Gr.

laevis, Meek, 1868, Trans. Chi. Acad. Sci., p. 108. This name was preoccupied.

marylandica, Hall, 1859, Pal. N. Y., vol. 3, p. 461, Oriskany sandstone.

mutabilis, Hall, 1857, (Meganteris mutabilis.) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 97, Low. Held. Gr.

ovalis, Hall, 1857, (Meganteris ovalis.) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 101, and Pal. N. Y., vol. 3, p. 458, Oriskany sandstone.

ovoidea, Eaton, 1832, (Terebratula ovoidea.) Geo. Textbook, p. 45, and Pal. N. Y., vol. 3, p. 456, Oriskany sandstone.

suessana, Hall, 1857, (Meganteris suessana.)

10th Rep. N. Y. St. Mus. Nat. Hist., p. 100, and Pal. N. Y., vol. 3, p. 459, Oriskany sandstone.

portlandica, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 115, Low. Held. Gr.

RETZIA, King, 1850, Monograph of Permian Foss., p. 137. [Ety. proper name.] Longitudinally oval, ribbed, with large punctures; foramen in ventral valve; area triangular; fissure closed. Type *R. adrieni*.

FIG. 604.
Retzia evax.

altirostris, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 28, Marshall Gr.

compressa, Meek, 1864, Pal. California, vol. 1, p. 14, Coal Meas.

deweyi, Hall, 1857, (Waldheimia deweyi.) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 89, Low. Held. Gr.

dubia, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 113, Low. Held. Gr.

electra, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 114, Low. Held. Gr.

eugenia, Billings, 1861, Can. Jour., vol. 6, p. 147, Ham. Gr.

evax, Hall, 1863, (Rhynchospira evax.) Trans. Alb. Inst., vol. 4, p. 213, and Rep. Geol. and Nat. Hist. Ind., Niagara Gr.

formosa, Hall, 1857, (Waldheimia formosa.) 10th Rep. N. Y. St. Mus. Nat. Hist., p. 88, Low. Held. Gr.

hippolyte, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 112, Low. Held. Gr.

lepida, Hall, 1860, (Rhynchospira lepida.) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 83, and Pal. N. Y., vol. 4, p. 275, Ham. Gr.

marcyi, Shumard, 1854, (Terebratula marcyi.) Marcy's Exp. Red Riv., p. 177, Kaskaskia Gr.

maria, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 112, Low. Held. Gr.

FIG. 603.—Rensseleria ovoidea.

meekian
Louisian
mormo

FIG. 605.—mormo

osagens

Acad

Chote

papillat

Louisian

mormo

popana

Acad

Chote

punctilij

mormo

polyple

Nat.

age G

sexplic

Bost.

derhc

sinuata

uata,

Up. S

subglo

subglo

Up. F

subglobo

moni

vera, sec

vera, var

costat

verneulo

wooster

vol. 5

No. 6

Rhynch

Imp.

chos,

trigon

mesia

beak

promi

beak,

somet

a delt

pieces

suppo

tend

socket

two, s

the h

separ

scars

muscl

shape

plates

loxia.

abrupta

Mus.

vol. 3

meekana, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 295, Permian Gr.

mormoni, Marcou, 1858, (Terebratula mormonii,) Geo. N. Amer., p. 51, Coal Meas. This species was subsequently, though in the same year, described by Shumard under the name *R. punctilifera*.

osagensis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 653, Waverly or Choteau Gr.

papillata, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 294, Permian Gr.

popana, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 654, Waverly or Choteau Gr.

punctilifera, Shumard, 1858, syn. for *Retzia mormoni*.

polypleura, Winchell, 1862, Proc. Acad. Nat. Sci., 2d ser., vol. 6, p. 406, Portage Gr.

semplicata, White & Whitfield, 1862, Proc. Post. Soc. Nat. Hist., vol. 8, p. 294, Kinderhook Gr.

sinuata, Hall, 1860, (Rhynchospira sinuata,) Can. Nat. and Geol., vol. 5, Up. Sil.

subglobosa, Hall, 1867, (Rhynchospira subglobosa,) Pal. N. Y., vol. 4, p. 421, Up. Held. Gr.

subglobosa, McChesney, syn. for *Retzia mormoni*.

vera, see *Eumetria vera*.

vera var. costata, see *Eumetria vera* var. costata.

verneuilana, see *Eumetria verneuilana*.

woosteri, White, 1879, Bull. U. S. Sur., vol. 5, No. 2, p. 215, and Cont. to Pal., No. 6, p. 134, Coal Meas.

RHYNCHONELLA, Fischer, 1809, Mem. Soc. Imp. Mosc., vol. 2, p. 35. [Ety. *rhynchos*, beak; *ella*, little.] Shell oval or trigonal, subglobose, with or without mesial fold and sinus; surface plicated; beak of ventral valve acute, entire, prominent, curved; foramen under the beak, by the incurving of which it is sometimes closed, partly surrounded by a deltidium, which is composed of two pieces; two teeth in the ventral valve, supported by dental plates, which extend to the bottom of the valve; two sockets in the dorsal valve; apophyses two, short, flattened, curved, attached to the hinge plate; adductor scars four, separated by a mesial ridge; pedicle scars on the cardinal plates; pedicle muscles of the ventral valve in a saucer-shaped cavity at the base of the dental plates; shell impunctate. Type *R. loxia*.

abrupta, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 68, and Pal. N. Y., vol. 3, p. 228, Low. Held. Gr.

acadiensis, Davidson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 172, Low Carb.

acinus, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 215, Niagara Gr.

acutiplicata, Hall, 1857, N. Y. St. Mus. Nat. Hist., p. 73, and Pal. N. Y., vol. 3, p. 232, Low Held. Gr.

acutirostris, Hall, 1847, (Atrypa acutirostra,) Pal. N. Y., vol. 1, p. 21, Chazy Gr.

aequalis, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 66, and Pal. N. Y., vol. 3, p. 224, Low. Held. Gr.

aequiradiata, Hall, 1852, (Atrypa aequiradiata,) Pal. N. Y., vol. 2, p. 70, Clinton Gr.

ainslaei, Winchell, 1860, 14th Ann. Rep. Geo. Minn., p. 315, Trenton Gr.

algeri, McChesney, 1860, New Pal. Foss. Carb. Not recognized.

altis, Hall, 1847, (Atrypa altis,) Pal. N. Y., vol. 1, p. 23, Chazy Gr.

altiplicata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 72, and Pal. N. Y., vol. 3, p. 231, Low Held. Gr.

alveata, see *Centronella alveata*.

ambigua, Calvin, 1878, Bull. U. S. Geo. Sur., vol. 4, No. 3, p. 729, Low. Devonian.

angulata, Linnaeus, as identified by Geinitz, syn. for *Syntrielasma hemiplicatum*.

anticostiensis, Billings, 1862, Pal. Foss., vol. 1, p. 142, Hud. Riv. Gr.

aprinis, DeVerneuil, 1845, (Terebratula aprinis,) Geo. Russia and Ural Mts., vol. 2, p. 90, and Pal. N. Y., vol. 2, p. 280, Niagara Gr.

arctirostrata, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 84, Kaaskasia Gr.

argentea, Billings, 1866, Catal. Sil. Foss. Antic., p. 43, Anticosti Gr.

argenturica, White, 1874, Rep. Invert. Foss., p. 14, and Geo. Sur. W. 100th Mer., vol. 4, p. 75, Hud. Riv. Gr.

aspasia, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 111, Low. Held. Gr.

barquensis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 408, Marshall Gr.

barrandi, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 82, and Pal. N. Y., vol. 3, p. 442, Oriskany sandstone.

bialveata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 73, and Pal. N. Y., vol. 3, p. 233, Low. Held. Gr.

bidens, Hall, 1852, (Atrypa bidens,) Pal. N. Y., vol. 2, p. 69, Clinton Gr.

bidentata, Hisinger, 1826, (Terebratula bidentata,) Vet. Acad. Handl., p. 343, and Pal. N. Y., vol. 2, p. 276, Niagara Gr.

billingsi, see *Stenoschisma billingsi*.

boonensis, Shumard, 1855, Geo. Rep. Mo., p. 205, Burlington Gr.

brevirostris, see *Pentamerus brevirostris*.

campbellana, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 79, and Pal. N. Y., vol. 3, p. 239, Low. Held. Gr.

FIG. 605.—*Retzia mormoni*.

camerifera, Winchell, 1862, Proc. Acad. Nat. Sci., p. 408, Marshall Gr.
capax, Conrad, 1842, (Atrypa capax,) Jour. Acad. Nat. Sci., vol. 8, p. 264, Hud. Riv. Gr.

FIG. 606.—Rhynchonella capax.

caput-testudinis, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 23, Burlington Gr.

carica, see Stenoschisma carica.

carbonaria, McChesney, 1860, New Pal. Foss., Coal Meas. Not recognized.

carolina, see Stenoschisma carolina.

castanea, Meek, 1868, Trans. Chi. Acad. Sci., p. 93, Devonian.

congregata, see Stenoschisma congregatum.

contracta, see Stenoschisma contractum.

cooperensis, Shumard, 1855, Geo. Rep. Mo., p. 204, Waverly or Choteau Gr.

corinthia, Billings, 1865, Pal. Foss., vol. 1, p. 220, Quebec Gr.

cuboides, Sowerby, (Atrypa cuboides,) see R. venustula.

cuneata, see Rhynchotretra cuneata var. Americana.

dawsoniana, Davidson, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 172, Subcarboniferous.

dentata, Hall, 1847, (Atrypa dentata,) Pal. N. Y., vol. 1, p. 148, Hud. Riv. Gr.

dotis, see Stenoschisma dotis.

dryope, Billings, 1874, Pal. Foss., vol. 2, p. 37, Gaspe limestone No. 8, Devonian.

dubia, Hall, 1847, (Atrypa dubia,) Pal. N. Y., vol. 1, p. 21, Chazy Gr.

duplicata, syn. for Stenoschisma contractum.

eatoniformis, McChesney, 1860, New Pal. Foss., syn. for R. rockymontana.

emacerata, Hall, 1852, (Atrypa emacerata,) Pal. N. Y., vol. 2, p. 71, Clinton Gr.

eminens, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 78, and Pal. N. Y., vol. 3, p. 237, Low. Held. Gr.

emmonsii, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 247, Devonian.

endlichi, Meek, 1876, U. S. Geo. Sur. of Colorado, p. 47, and White's Cont. to Pal. No. 6, p. 133, Up. Devonian.

eurekensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 223, Subcarboniferous.

eva, Billings, 1866, Catal. Sil. Foss. Antic., p. 44, Anticosti Gr.

evangelina, Hartt, 1868, Acad. Geol., p. 299, Subcarboniferous.

excellens, Billings, 1874, Pal. Foss., vol. 2, p. 36, Gaspe limestone No. 8, Devonian.

eximia, see Stenoschisma eximium.

explanata, McChesney, 1860, Desc. New Pal. Foss., Kaskaskia Gr. Not recognized.

fitchiana, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 85, and Pal. N. Y., vol. 3, p. 441, Oriskany sandstone.

formosa, see Stenoschisma formosum.

fringilla, Billings, 1862, Pal. Foss., vol. 1, p. 141, Anticosti Gr., Div. 1, Mid. Sil.

glacialis, Billings, 1862, Pal. Foss., vol. 1, p. 143, Anticosti Gr., Div. 1, Mid. Sil.

glansfugea, see Centronella glansfugea.

greenana, Ulrich, 1886, Cont. to Am. Pal., p. 26, Waverly Gr.

grosvenori, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 10, and Bull. Am. Mus. Nat. Hist., p. 53, Warsaw Gr.

guadalupae, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 295, Permian Gr.

heteropsis, Winchell, 1865, Proc. Acad. Nat. Sci., p. 121, Marshall Gr.

horsfordi, see Stenoschisma horsfordi.

hubbardi, Winchell, 1862, Proc. Acad. Nat. Sci., p. 407, Marshall Gr.

huronensis, Winchell, 1862, Proc. Acad. Nat. Sci., 2d ser., vol. 6, p. 409, Portage Gr.

hydraulica, Whitfield, 1882, Desc. New Spec. Foss. from Ohio, p. 194, Low. Held. Gr.

ida, Hartt, 1868, Acad. Geol., p. 298, Subcarboniferous.

illinoisensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 24, and Geo. Sur. Ill., vol. 8, p. 104, Coal Meas.

increbescens, syn. for Rhynchonella capax.

indentata, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 393, Permian Gr.

indianensis, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 215, Niagara Gr.

inaequiplicata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 126, Up. Held. Gr.

intermedia, Barris, 1879, Proc. Davenport Acad. Sci., vol. 2, p. 285, Up. Held. Gr.

inutilis, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 74, and Pal. N. Y., vol. 3, p. 235, Low. Held. Gr.

janea, Billings, 1866, Catal. Sil. Foss. Antic., p. 43, Anticosti Gr.

lacunosa, Not an American species.

lamellata, Hall, 1852, (Atrypa lamellata,) Pal. N. Y., vol. 2, p. 329, Coralline Limestone.

laura, see Leiorhynchus laura.

macra, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 11, and Bull. Am. Mus. Nat. Hist., p. 52, Warsaw Gr.

mainensis, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 110, Low. Held. Gr.

marsh
Nat.
medea,
p. 27
metalli
p. 20,
4, p.
mica,
Antic
microp
Nat.
misson
Mo.,
multist
St. M
vol. 3
mutabili
Mus.
vol. 3
mutata
4, p.
saw
neenah
Sur.
p. 26
neglect
Pal.
neglect
Jour.
and
2, p.
nobilis
Mus.
vol. 3
nucleol
St. M
N. Y
nucula
nuc
Up.
nutrix,
Anti
oblata,
Mus.
vol. 3
obsoles
St.
verly
obtus
plic
agars
occiden
Geo.
opposit
Bost.
derh
orbicul
orienta
vol. 3
osagens
etion
ottum
Nat.
Pal.,
parvini
phor
perlman
Geo.
Wis.

l. Foss., vol.
e No. 8, De-

mium.

Desc. New
askaskia Gr.
ed.

1857, 10th
t. Mus. Nat.
d Pal. N. Y.,
41, Oriskany

Stenoschisma

gs, 1862, Pal.
p. 141, Anti-

Foss., vol. 1,
1, Mid. Sil.
ansfagen.
to Am. Pal.,

s. Alb. Inst.,
m. Mus. Nat.

, Trans. St.
p. 295, Per-

Proc. Acad.
Gr.

orsfordi.
oc. Acad. Nat.

Proc. Acad.
9, Portage Gr.
Desc. New
p. 194, Low.

, p. 298, Sub-

Bull. No. 2, Ill.
and Geo. Sur.

ns.
onella capax.

ans. St. Louis
ermian Gr.

ns. Alb. Inst.,

10th Rep.
p. 126, Up.

oc. Davenport
Up. Held. Gr.

rep. N. Y. St.
d Pal. N. Y.,
Gr.

al. Sil. Foss.
r.

species.
pa lamellata.)

oralline Lime-

ra.
b. Inst., vol. 4,
Nat. Hist., p.

Proc. Port.
p. 110, Low.

marshallensis, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 408, Marshall Gr.

medea, Billings, 1860, Can. Jour., vol. 5,
p. 271, Corniferous Limestone.

metallica, White, 1874, Rep. Invert. Foss.,
p. 20, and Geo. Sur. W. 100th Mer., vol.
4, p. 129, Carb.

mica, Billings, 1866, Catal. Sil. Foss.
Antic., p. 44, Anticosti Gr.

micropleura, Winchell, 1865, Proc. Acad.
Nat. Sci., p. 122, Marshall Gr.

missouriensis, Shumard, 1855, Geo. of
Mo., p. 204, Waverly or Choteau Gr.

multistriata, Hall, 1857, 10th Rep. N. Y.
St. Mus. Nat. Hist., p. 85, and Pal. N. Y.,
vol. 3, p. 440, Oriskany sandstone.

mutabilis, Hall, 1857, 10th Rep. N. Y. St.
Mus. Nat. Hist., p. 66, and Pal. N. Y.,
vol. 3, p. 225, Low. Held. Gr.

mutata, Hall, 1858, Trans. Alb. Inst., vol.
4, p. 10, and Geo. Sur. Iowa, p. 658, War-
saw Gr.

neenah, Whitfield, 1880, Ann. Rep. Geo.
Sur. Wis., p. 62, and Geo. Wis., vol. 4,
p. 265, Trenton Gr.

neglecta, Hall, 1852, (Atrypa neglecta,) Pal.
N. Y., vol. 2, p. 274, Niagara Gr.

neglecta var. scobina, Meek, 1872, Am.
Jour. Sci. and Arts, 3d ser., vol. 4, p. 277,
and Ohio Pal., vol. 1, p. 179, and vol.
2, p. 116, Niagara Gr.

nobilis, Hall, 1857, 10th Rep. N. Y. St.
Mus. Nat. Hist., p. 80, and Pal. N. Y.,
vol. 3, p. 240, Low. Held. Gr.

nucleolata, Hall, 1857, 10th Rep. N. Y.
St. Mus. Nat. Hist., p. 68, and Pal.
N. Y., vol. 3, p. 227, Low. Held. Gr.

nucula, Sowerby, 1839. (Terebratula
nucula.) Murch. Sil. Syst., p. 611,
Up. Sil.

nutrix, Billings, 1866, Catal. Sil. Foss.
Antic., p. 43, Anticosti Gr.

oblata, Hall, 1857, 10th Rep. N. Y. St.
Mus. Nat. Hist., p. 86, and Pal. N. Y.,
vol. 3, p. 439, Oriskany sandstone.

obsolescens, Hall, 1860, 13th Rep. N. Y.
St. Mus. Nat. Hist., p. 111, Wa-
verly Gr.

obtusiplicata, Hall, 1852, (Atrypa obtusi-
plicata,) Pal. N. Y., vol. 2, p. 273, Ni-
agara Gr.

occidens, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 152, Devonian.

opposita, White & Whitfield, 1862, Proc.
Bost. Soc. Nat. Hist., vol. 8, p. 294, Kin-
derhook Gr.

orbicularis, see Stenoschisma orbiculare.

orientalis, Billings, 1859, Can. Nat. Geo.,
vol. 4, p. 443, Chazy Gr.

osagensis, Swallow, 1858, syn. for Rhy-
nchonella uta.

ottumwa, White, 1862, Proc. Bost. Soc.
Nat. Hist., vol. 9, p. 23, and Cont. to
Pal., No. 8, p. 165, St. Louis Gr.

parvini, McChesney, syn. for Camero-
phoria subtrigona.

perlamellosa, Whitfield, 1878, Ann. Rep.
Geo. Sur. Wis., p. 73, and Geo. Sur.
Wis., vol. 4, p. 265, Hud. Riv. Gr.

perrostellata, Swallow, 1863, Trans. St.
Louis Acad. Sci., vol. 2, p. 85, Kas-
kaskia Gr.

persinnata, Winchell, 1865, Proc. Acad.
Nat. Sci., p. 121, Marshall Gr.

phoca, see Atrypa phoca.

pisum, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 135, Niagara Gr.

planoconvexa, Hall, 1857, 10th Rep. N. Y.
St. Mus. Nat. Hist., p. 75, and Pal.
N. Y., vol. 3, p. 235, Low. Held. Gr.

pleiopleura, Conrad, 1841, (Atrypa pleio-
pleura,) Ann. Rep. N. Y., p. 53, and
Pal. N. Y., vol. 3, p. 440, Oriskany
sandstone.

plena, Hall, 1847, (Atrypa plena,) Pal.
N. Y., vol. 1, p. 21, Chazy Gr.

plicata, Hall, 1852, (Atrypa plicata,) Pal.
N. Y., vol. 2, p. 10, Medina Gr.

plicatula, Hall, 1843, (Atrypa plicatula,) Geo.
Rep. 4th Dist. N. Y., p. 71, and
Pal. N. Y., vol. 2, p. 74, Clinton Gr.

plicifera, Hall, 1847, (Atrypa plicifera,) Pal.
N. Y., vol. 1, p. 22, Chazy Gr.

principalis, Hall, 1857, 10th Rep. N. Y.
St. Mus. Nat. Hist., p. 84, and Pal. N.
Y., vol. 3, p. 443, Oriskany sandstone.

prolifera, see Stenoschisma prolificum.

pugnus, Martin, 1809, (Conchiliolithus
Anonites pugnus,) Petrif. Derb., pl.
22, figs. 4 and 5, Subcarboniferous.

pustulosa, White, 1860, Bost. Jour. Nat.
Hist., vol. 7, p. 236, Burlington Gr.

pyramidata, Hall, 1857, 10th Rep. N. Y.
St. Mus. Nat. Hist., p. 70, and Pal. N. Y.,
vol. 3, p. 229, Low. Held. Gr.

pyrrha, Billings, 1866, Catal. Sil. Foss.
Antic., p. 44, Anticosti Gr.

quadriscopata, Hall, 1852, (Atrypa quadri-
costata,) Pal. N. Y., vol. 2, p. 68, Clin-
ton Gr.

ramsayi, Hall, 1859, Pal. N. Y., vol. 3, p.
446, Oriskany sandstone.

raricosta, Whitfield, 1882, Desc. New Spec.
Foss., from Ohio, p. 201, Up. Held. Gr.

recurvirostra, see Anazyga recurvirostra.

reticulata, see Eichwaldia reticulata.

ricinula, Hall, 1858, Trans. Alb. Inst., vol.
4, p. 9, and Bull. Am. Mus. Nat. Hist.,
p. 53, Warsaw Gr.

ringens, Swallow, 1860, Trans. St. Louis
Acad. Sci., vol. 1, p. 653, Burlington Gr.

robusta, Hall, 1852, Pal. N. Y., vol. 2, p.
71, (Atrypa robusta,) Clinton Gr.

rockymontana, Marcou, 1858, (Terebratula
rockymontana,) Geo. North America,
p. 50, Coal Meas.

royana, see Stenoschisma royanum.

ridleyana, Safford, 1869, Geo. of Tenn.
Not defined.

rudis, Hall, 1857, 10th Rep. N. Y. St. Mus.
Nat. Hist., p. 75, and Pal. N. Y., vol. 3,
p. 235, Low. Held. Gr.

rugosa, Hall, 1852, (Atrypa rugosa,) Pal.
N. Y., vol. 2, p. 271, Niagara Gr.

saffordi, Hall, 1860, Can. Nat. and Geo.,
vol. 5, p. 144, Low. Held. Gr.

sagerana, Winchell, 1862, Proc. Acad. Nat.
Sci., p. 407, Marshall Gr.

sappho, see *Stenoschisma sappho*.
semiplicata, Conrad, 1841, (*Atrypa semiplicata*.) Ann. Rep. N. Y., p. 56, and Pal. N. Y., vol. 3, p. 224, Low. Held. Gr. septata, Hall, 1859, Pal. N. Y., vol. 3, p. 443, Oriskany sandstone.
sordida, Hall, 1847, (*Atrypa sordida*.) Pal. N. Y., vol. 1, p. 148, Trenton Gr.
speciosa, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 81, and Pal. N. Y., vol. 3, p. 444, Oriskany sandstone.
stephani, see *Stenoschisma stephani*.
stricklandi, Sowerby, 1839, (*Terebratula stricklandi*.) Murch. Sil. Syst., p. 631, Niagara Gr.
subcircularis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 408, Marshall Gr.
subcuboides. Not an American species.
subcuneata, Hall, 1856, Trans. Alb. Inst., vol. 4, p. 11, and Geo. Sur. Iowa, p. 658, Warsaw Gr.
subtrigona, see *Camarophoria subtrigona*.
subtrigonalis, Hall, 1847, (*Atrypa subtrigonalis*.) Pal. N. Y., vol. 1, p. 145, Trenton Gr.
sulcoplicata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 76, and Pal. N. Y., vol. 3, p. 236, Low. Held. Gr.
tennesseensis, Roemer, 1860, Sil. Fauna West Tenn., p. 72, Niagara Gr.
tethys, see *Stenoschisma tethys*.
tetraptix, Winchell, 1865, Proc. Acad. Nat. Sci., p. 120, Kinderhook Gr.
texana, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 393, Permian Gr.
thalia, see *Stenoschisma billingsi*.
thera, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 223, Subcarboniferous.
transversa, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 74, and Pal. N. Y., vol. 3, p. 234, Low. Held. Gr.
tuta, S. A. Miller, 1881, Jour. Cin. Soc., Nat. Hist., vol. 4, p. 315, Burlington Gr.
unica, Winchell, 1865, Proc. Acad. Nat. Sci., p. 122, Marshall Gr.
unisulcata, see *Meristella unisulcata*.
uta, Marcou, 1858, (*Terebratula uta*.) Geo. N. Amer., p. 58, Coal Meas. This was subsequently described by Swallow as *R. osagensis*.
vellicata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 71, and Pal. N. Y., vol. 3, p. 230, Low. Held. Gr.
ventricosa, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 78, and Pal. N. Y., vol. 3, p. 238, Low. Held. Gr.
venustula, Hall, 1867, Pal. N. Y., vol. 4, p. 346, Tully limestone. This was identified by Vanuxem, 1842, Geo. 3d Dist. N. Y., as *Atrypa cuboides* of Sowerby.
vicina, Billings, 1866, Catal. Sil. Foss. Antic., p. 44, Anticosti Gr.
warrenensis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 653, Ham. Gr.
wasatchensis, White, 1874, Rep. Invert. Foss., p. 19, and Geo. Sur. W. 100th Mer., vol. 4, p. 130, Carb.

whitiana, S. A. Miller, 1883, 2d Ed. Am. Pal. Foss., p. 297, Niagara Gr., from Waldron, Indiana. Proposed instead of *R. whitii*, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 216, and also in 28th Rep. N. Y. St. Mus. Nat. Hist., p. 164, pl. 26, figs. 23-33, and again in the 11th Ann. Rep. Geol. and Nat. Hist. of Indiana, p. 307, pl. 26, figs. 23-33.
whitii, Winchell, 1862, Proc. Acad. Nat. Sci., p. 407, Marshall Gr.
whitii, Hall, see *R. whitiana*.
wilsoni, Sowerby, 1816, (*Terebratula wilsoni*.) Min. Conch., vol. 2, p. 38, Niagara Gr.
wortheni, see *Camarophoria wortheni*.
Rhynchospira, Hall, 1859, Pal. N. Y., vol. 3, syn. for *Retzia*.
deweyi, see *Retzia deweyi*.
evax, see *Retzia evax*.
formosa, see *Retzia formosa*.
lepida, see *Retzia lepida*.
nobilis, see *Trematospira nobilis*.
rectirostra, see *Trematospira rectirostra*.
subglobosa, see *Retzia subglobosa*.
sinuata, see *Retzia sinuata*.
RHYNCHOTRETA, Hall, 1879, 28th, Rep. N. Y. St. Mus. Nat. Hist., p. 166. [*Ety. rhynchos*, beak; *treto*, with a hole in it.] Distinguished from *Rhynchonella* by the straight, produced, perforated beak of the ventral valve and divided deltidium, and by the crura which rise near the dorsal beak, curve into the ventral cavity, and re-

FIG. 607.—*Rhynchotretra cuneata* var. *americana*

curve to the dorsal side. Type *R. cuneata*.

cuneata var. *americana*, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 167, Niagara Gr.

FIG. 608.—*Rhynchotretra quadriplicata*

quadriplicata, S. A. Miller, 1875, (*Trematospira quadriplicata*.) Cin. Quar. Jour. Sci., vol. 2, p. 60, Trenton Gr.
Rhynobolus, Hall, 1871, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 247, syn. for *Trimerella*.
galtensis, Hall, see *Trimerella galtensis*.
SCHIZAMBON, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 69. [*Ety. schiza*, a cleft; *ambon*, the boss of a shield.] Inequivalve, ovate; valves inarticulate; no area or deltidium; foramen oblong; structure calcareo-corneous; two scars in each valve. Type *S. typicalis*.
typicalis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 70, Chazy Gr.
SCHIZOBOLUS, Ulrich, 1886, Cont. to Am. Pal., p. 25. [*Ety. schiza*, a cleft; *obolus*, a genus.] Ventral valve with apex at the terminus of a notch in the posterior margin; two pair of adductor

scars with pairs septa truncal 16th 28, a esee
 SCHIZOCR 1875, 73.
 Cran paracr artic uppe ated Type filosa, Pal. Uti
 SIPHONOT ural siphon Shell tral v thick hing back with cylin sage valve form cepti valve dinal
 FIG. 610
 treta
 a, Vent interior
 slend base.
 scotica
 ser.,
 SKENIDIU
 Mus.
 little
 by it
 dina
 tum
 and
 extre
 devoni
 Geo.
 hali, S
 defin
 insign
 N. Y
 pyram
 alis,)
 ara C
 SPIRIFER
 2, p.
 p. 51

2d Ed. Am.
a Gr., from
posed instead
Trans. Alb.
also in 28th
Hist., p. 164,
in the 11th
Hist. of In-
t-33.

Acad. Nat.

abratura wil-
p. 38, Niag-

ortheni.

N. Y., vol. 3.

dia.

ectirostra.

osa.

8th, Rep. N.

at. Hist., p.

mches, beak;

hole in it.]

from Rhyn-

the straight,

orated beak

l valve and

um, and by

ch rise near

curve into

ity, and re-

e. Type R.

scars separated by a ridge; dorsal valve with truncated posterior margin; two pairs of muscular scars separated by a septum. Type *S. truncatus*.

truncatus, Hall, 1862, (*Discina truncata*), 16th Rep. N. Y. St. Mus. Nat. Hist., p. 28, and Pal. N. Y., vol. 4, p. 23. Genesee slate to Chemung Gr.

SCHIZOCRANIA, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 73. [Ety. *schiza*, a cleft; *Crania*, a genus.] Shell

parasitic inequivalve, inarticulated, surface of the upper valve finely striated; interior with six muscular scars. Type *S. filosa*.

filosa, Hall, 1847, (*Orbicula* (?) *filosa*), Pal. N. Y., vol. 1, p. 99, Hud. Riv. and Utica Slate.

SIPHONOTRETA, DeVerneuil, 1845, Russia and Ural Mountains, vol. 2, p. 286. [Ety. *siphon*, siphon; *tretos*, with a hole in it.]

Shell oblong oval, unarticulated; ventral valve most convex with a straight, thick, perforated, conical beak near the hinge-line; foramen opening on the back of the beak, and communicating with the interior of the shell by a cylindrical tube or siphon for the passage of the muscle of attachment; dorsal valve slightly convex, the hinge-line forming an arch which merges imperceptibly into the lateral margins; each valve has a wide, crescent-shaped cardinal edge, covered by horizontal lines

of growth; structure calcareo-corneous, with a distinctly punctured structure arranged in tubular layers; surface smooth, with numerous lines of growth and slender hollow spines dilated at the base. Type *S. unguiculata*.

scotica, Davidson, 1877, Geol. Mag., new ser., vol. 4, p. 13, Utica slate.

SKENIDION, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 70. [Ety. *skenidion*, a little tent.] Distinguished from *Orthis* by its large triangular area; the cardinal process extends as a median septum through the length of the shell, and may be simple or divided at the extremity. Type *S. insigne*.

devonicum, Walcott, 1885, Monogr. U. S. Geo. Surv., vol. 8, p. 116, Devonian.

halli, Safford, 1869, Geo. of Tenn. Not defined.

insigne, Hall, 1859, (*Orthis insignis*), Pal. N. Y., vol. 3, p. 173, Low. Held. Gr.

pyramidale, Hall, 1852, (*Orthis pyramidalis*), Pal. N. Y., vol. 2, p. 251, Niagara Gr.

SPHIFERA, Sowerby, 1815, Min. Conch., vol. 2, p. 42, and Linnæan Trans., vol. 12, p. 514. [Ety. *spira*, spire; *fero*, to bear.]

Triangular semicircular, transversely elongate, subglobose or otherwise variable in form, with or without mesial fold and sinus; structure impunctate; surface smooth, striated or plicated; cardinal line straight, area in each valve; hinge articulated by short teeth and sockets; area of the ventral valve larger than the other, and divided by a triangular foramen more or less closed by a false deltidium; area of the dorsal valve divided in the middle by a fissure occupied by the cardinal muscular process; beak of ventral valve more prominent than that of the other; in the interior of the dorsal valve the spiral supports of the labial arms are attached by their crura to the hinge plates, some distance from which they are nearly or quite connected by a small process extending inward from each; the cardinal muscles seem to have been attached to the cardinal process, under and in front of which four scars of the adductor muscles occur; on each side of a mesial ridge in the ventral valve occur the scars of the adductors, and outside of these the scars of the cardinal muscles. Type *S. striata*.

acanthoptera, Conrad, 1842, (*Delthyris acanthoptera*), Jour. Acad. Nat. Sci., vol. 8, p. 264, Chemung Gr.

acuminata, Conrad, 1839, (*Delthyris acuminata*), Ann. Rep. N. Y., p. 65, and Pal. N. Y., vol. 4, p. 198, Up. Held. and Ham. Grs.

acuticostata, DeKoninck, 1843, Desc. Ann. Foss. Terr. Carb. Belg., p. 265, Subcarboniferous.

agelaia, Meek, 1873, Hayden's Geo. Sur. Terr., p. 470, and White's Cont. to Pal. No. 6, p. 135, Subcarboniferous.

alata, Castelnau, 1843, Syst. Sil., p. 42. Not recognized.

albapinensis, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 255, Waverly Gr.

aldrichi, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 634, Devonian.

alta, Hall, 1867, Pal. N. Y., vol. 4, p. 248, Chemung Gr.

amara, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 642, Waverly or Choteau Gr.

angusta, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 164, and Pal. N. Y., vol. 4, p. 230, Ham. Gr.

annæ, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 641, Ham. Gr.

annctans, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 216, Subcarboniferous.

arata, syn. for *S. granulifera*.

archiaci, see *S. disjuncta*.

arctica, Houghton, 1857, Jour. Roy. Soc. Dub., vol. 1, p. 183, Devonian.

arctisegmenta, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 131, and Pal. N. Y., vol. 4, p. 208, Up. Held. Gr.

FIG. 609. — *Schizocrania filosa*.

FIG. 610. — *Siphonotreta unguiculata*. a, Ventral valve; b, interior of same.

Sphenotreta plicata.

- arenosa*, Conrad, 1839, (*Delthyris arenosa*.) Ann. Rep. N. Y., p. 65, and Pal. N. Y., vol. 3, p. 425, Oriskany sandstone.
- argentaria*, Meek, 1877, U. S. Geo. Sur. 40th Parallel, p. 42, Devonian.
- arrecta*, Hull, 1859, Pal. N. Y., vol. 3, p. 422, Oriskany sandstone.
- aspera*, Hall, 1858, Geo. Rep. Iowa, p. 508, Ham. Gr.
- asperata*, Ringuenberg, 1886, Bull. Nat. Sci., vol. 5, p. 16, Niagara Gr.
- atwaterana*, S. A. Miller, 1878, Proc. Davenport Acad. Sci., vol. 2, p. 221, Ham. Gr. Proposed instead of *S. penata*, Owen, which was preoccupied.
- audacula*, Conrad, (*Delthyris audacula*.) 1842, Jour. Acad. Nat. Sci., vol. 8, p. 262, Ham. Gr.
- belphegor*, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 30, Genesee shales.
- biulveata*, Conrad, (*Delthyris bialveata*.) 1842, Jour. Acad. Nat. Sci., vol. 8, p. 261, Niagara Gr. Probably a syn. for *S. radiata*.
- bicostata*, Vanuxem, 1842, (*Orthis bicostatus*.) Geol. Rep. 3d Dist. N. Y., p. 91, and Pal. N. Y., vol. 2, p. 263, Niagara Gr.
- bicostata* var. *petila*, Hall, 1879, Desc. New Spec. Foss., p. 15, and 11th Rep. Geo. and Nat. Hist. Ind., p. 297, Niagara Gr.
- bidorsalis*, Winchell, 1866, Rep. Low. Penin. Mich., p. 93, Ham. Gr.
- bifurcata*, Hull, 1858, Trans. Alb. Inst., vol. 4, p. 8, and Bull. Am. Mus. Nat. Hist., p. 47, Warsaw Gr.
- billingsana*, n. sp., Upper Devonian Gaspe limestone, No. 8. Proposed instead of *S. superba*, Billings, 1874, Pal. Foss., vol. 2, p. 45, which name was preoccupied.
- biloba*, Linneus, 1768, (*Anomia biloba*.) Syst. Nat., p. 115, and Pal. N. Y., vol. 2, p. 200, Niagara Gr.
- bimesialis*, Hall, 1858, Geo. Rep. Iowa, vol. 1, pt. 2, p. 507, Ham. Gr.
- biplicata*, Hall, 1858, Geo. Rep. Iowa, vol. 1, pt. 2, p. 519, Kinderhook Gr.
- boonensis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 640, Low Coal Meas.
- brachynota*, Hall, 1843, (*Delthyris brachynota*.) Geo. 4th Dist. N. Y., p. 71, Clinton Gr. Not well defined.
- calcarata*, syn. for *S. disjuncta*.
- camerata*, Morton, 1836, Am. Jour. Sci., vol. 29, p. 150, Coal Meas.
- camerata* var. *kansasensis*, Swallow, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 409, Coal Meas.
- camerata* var. *percrassa*, Swallow, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 409, Coal Meas. This name was preoccupied as a species.
- capax*, Hall, 1858, Geo. Rep. Iowa, vol. 1, pt. 2, p. 520, syn. for *S. parryana*.
- carteri*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 170, Waverly Gr.
- cedarensis*, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 585, Ham. Gr.
- centronota*, Winchell, 1865, Proc. Acad. Nat. Sci., p. 118, and Geo. Sur. W. 100th Mer., vol. 4, p. 87, Cuyahoga Shale.
- clara*, Swallow, 1853, Trans. St. Louis Acad. Sci., vol. 2, p. 86, Kaskaskia Gr.
- clavata*, McChesney, 1861, Desc. New Pul. Foss., p. 84, Burlington Gr. Not recognized.
- clintoni*, syn. for *S. granulifera*.
- clio*, syn. for *S. ziczac*.
- compacta*, Meek, 1868, Trans. Chi. Acad. Sci., p. 102, Ham. Gr.
- concinna*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 60, Low. Held. Gr.
- congesta*, syn. for *S. granulifera*.
- conradana*, S. A. Miller, 1883, 2d Ed. Am. Pal. Foss., p. 298, Oriskany, Up. Held. and Ham. Grs. Proposed instead of *S. fimbriata* of Conrad in Jour. Acad. Nat. Sci., vol. 8, p. 263, and Pal. N. Y., vol. 4, p. 214, which was preoccupied.
- consobrina*, D'Orbigny, 1850, Prodr. d. Puléont., t. 1, p. 98, Ham. Gr. Proposed instead of *S. ziczac*, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 200, which was preoccupied by Roemer.
- consors*, Winchell, 1866, Rep. Low. Peninsula Mich., p. 93, Ham. Gr.
- cooperensis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 643, Waverly or Choteau Gr.
- corticosa*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 160, Ham. Gr.
- costalis*, Castelnau, 1843, Syst. Sil., p. 41. Not recognized.
- crenistriata*, see *Streptorhynchus crenistriatum*.
- crispa*, Hisinger, 1826, (*Terebratula crispa*.) Act. Acad. Sci., Holm., t. 7, fig. 4, and Pal. N. Y., vol. 2, p. 262, Niagara Gr.
- crispa* var. *simplex*, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 157, Niagara Gr.
- cumberlandiae*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 63, and Pal. N. Y., vol. 3, p. 421, Oriskany sandstone.
- cuspidatiformis*, n. sp., Keokuk Gr. Proposed instead of *S. subcuspidata*, Hall, 1858, Geo. Sur. Iowa, p. 646, pl. 20, fig. 5 a, b, which name was preoccupied.
- cycloptera*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 58, and Pal. N. Y., vol. 3, p. 199, Low. Held. Gr.
- cyrtiniformis*, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 238, Chemung Gr.
- decemplicata*, Hall, 1843, (*Delthyris decemplicata*.) Geo. Rep. 4th Dist. N. Y., p. 106, Niagara Gr.
- desiderata*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 217, Subcarb.
- disjuncta*, Sowerby, 1840, Trans. Geo. Soc., 2d ser., vol. 5, p. 704, and Pal. N. Y., vol. 4, p. 243, Chemung Gr.
- disparilis*, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 134, and Pal. N. Y., vol. 4, p. 204, Up. Held. Gr.

distan
divar
Mu
vol
dubia
duod
der
and
gri
duplic
plig
p. 3
eaton
engel
Sci.
398
eudon
Sur
eurnt
teir
and
eacute
expor
expor
rect
extens
exten
520,
fascige
Spir
fastiga
and
fastiga
Acad
preo
tona
filitos
insu
fimbri
and
fimbria
occu
fischeri
Not
forbes
Acad
ton
formos
Nat.
4, p.
fornac
Mus
fornax
Nat.
frankl
Sci.
fulton
Ill.,
gaspen
2, p.
gibbos
Sur.
gigante
glabra
Petr
carbo
glabra
1861
Geo.

distans, syn. for *S. disjuncta*.
divaricata, Hall, 1857, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 133, and Pal. N. Y., vol. 4, p. 213, Cornif. and Ham. Grs.
dubia, see *Pentamerella dubia*.
duodenaria, Hall, 1843, (*Delthyris duodenaria*), Geol. 4th Dist. N. Y., p. 171, and Pal. N. Y., vol. 4, p. 189, Schoharie grit and Cornif. Gr.
duplicata, Conrad, 1842, (*Delthyris duplicata*), Jour. Acad. Nat. Sci., vol. 8, p. 261, Ham. Gr.
eutoni, see *S. medialis* var. *eutoni*.
engelmanni, Meek, 1860, Proc. Acad. Nat. Sci., p. 308, and Geo. Sur. Ill., vol. 3, p. 398, Oriskany sandstone.
eudora, Hall, 1861, Rep. of Prog. Wis. Sur., p. 25, Niagara Gr.
eurteines, Owen, 1844, (*Delthyris eurteines*, Report on Min. Lands, p. 74, and Pal. N. Y., vol. 4, p. 209, Up. Held. Gr. *eurteines* var. *forncacula* see *S. forncacula*.
exporrecta, see *Cyrtia exporrecta*.
exporrecta var. *arrecta*, see *Cyrtia exporrecta* var. *arrecta*.
extensa, syn. for *S. disjuncta*.
extenuata, Hall, 1858, Geo. Rep. Iowa, p. 520, Kinderhook Gr.
fasciger, Keyserling in Owen's report, see *Spirifera camerata*.
fastigata, Morton, 1836, Am. Jour. Sci. and Arts, vol. 29, p. 149, Coal Meas.
fastigata, Meek & Worthen, 1870, Proc. Acad. Nat. Sci., p. 36. The name was preoccupied by Morton. See *S. mortonana*.
filicosta, Winchell, 1866, Rep. Low. Peninsula Mich., p. 94, Ham. Gr.
fimbriata, Morton, 1836, Am. Jour. Sci. and Arts, vol. 29, p. 149, Coal Meas.
fimbriata, Conrad. The name was preoccupied. See *S. conradana*.
fischeri, Castelnau, 1843, Syst. Sil., p. 42. Not recognized.
forbesi, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., vol. 3, p. 73, Burlington Gr.
formosa, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 154, and Pal. N. Y., vol. 4, p. 220, Ham. Gr.
forncacula, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 154, Ham. Gr.
fornax, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 155, Ham. Gr.
franklini, Meek, 1868, Trans. Chi. Acad. Sci., p. 107, Ham. Gr.
fultonensis, Worthen, 1873, Geol. Rep. Ill., vol. 5, p. 572, Low. Coal Meas.
gaspensis, Billings, 1874, Pal. Foss., vol. 2, p. 74, Devonian.
gibbosa, Hall, 1861, Rep. of Prog. Wis. Sur., p. 25, Niagara Gr.
gigantea, syn. for *S. disjuncta*.
glabra, Martin, 1809, (*Anomites glabra*), Petrif. Derb., tab. 28, figs. 9 and 10, Subcarboniferous.
glabra var. *contracta*, Meek & Worthen, 1861, Proc. Acad. Nat. Sci., p. 143, and Geo. Sur. Ill., vol. 2, p. 298, Kaskaskia Gr.

glabra var. *nevadensis*, Walcott, 1885, Monog. U. S. Geo. Sur., vol. 8, p. 139, Up. Devonian.
glanscerasi, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 8, Ham. Gr.
grandeva, syn. for *S. disjuncta*.
granulifera, Hall, 1843, (*Delthyris granulifera*), Geol. 4th Dist. N. Y., p. 207, and Pal. N. Y., vol. 4, p. 223, Ham. Gr.
granulosa, Conrad, 1839, (*Delthyris granulosa*), Ann. Rep. N. Y., p. 65, Low. Held. Gr.

FIG. 611.—*Spirifera gregaria*.

gregaria, Clapp, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 127, and Pal. N. Y., vol. 4, p. 195, Up. Held. Gr.
grieri, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 127, and Pal. N. Y., vol. 4, p. 194, Schoharie grit and Up. Held. Gr.
grimesi, Hall, 1858, Geo. Rep. of Iowa, p. 604, Burlington Gr.
guadalupensis, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 391, Permian Gr.
hannibalensis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 647, Waverly or Choteau Gr.
hemicycla, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 399, Oriskany sandstone.
hemiplicata, see *Syntrielasma hemiplicatum*.
heterochilus, syn. for *S. granulifera*.
hirtus, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 293, Kinderhook Gr.
hungerfordi, Hall, 1858, Geo. Rep. Iowa, vol. 1, pt. 2, p. 501, Ham. Gr.
huronensis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 405, Portage Gr.
huronensis, Castelnau, 1843, Syst. Sil., p. 41. Not recognized.
imbrex, Hall, 1858, Geo. Rep. Iowa, p. 601, Burlington Gr.
inequivalvis, Castelnau, 1843, Syst. Sil., p. 40. Not recognized.
incerta, Hall, 1858, Geo. Rep. Iowa, p. 602, Burlington Gr.
inconstans, syn. for *Spirifera racinensis*.
increbescens, Hall, 1858, Geo. Rep. Iowa, p. 706, Kaskaskia Gr.
increbescens var. *americana*, Swallow, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 410, Kaskaskia Gr.
increbescens var. *transversalis*, Hall, 1858, Geol. Rep. Iowa, p. 708, Kaskaskia Gr.
inequicostata, Owen, 1852, Geo. Rep. Wis., Iowa, and Min., p. 586, Carb.
inornata, syn. for *S. disjuncta*.
insolita, Winchell, 1862, Proc. Acad. Nat. Sci., p. 405, Portage Gr.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

- intermedia*, Hall, 1859, Pal. N. Y., vol. 3, p. 424, Oriskany sandstone. This name was preoccupied by Brongniart in 1829.
- inutilis*, Hall, 1858, Geo. Rep. Iowa, vol. 1, pt. 2, p. 505, Ham. Gr.
- iowensis*, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 585, Ham. Gr.
- keilloggi*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 86, Keokuk Gr.
- kennicotti*, Meek, 1868, Trans. Chi. Acad. Sci., p. 101, Ham. Gr.
- kentuckensis*, see *Spiriferina kentuckiensis*.
- kentuckensis* var. *propatula*, see *Spiriferina kentuckiensis* var. *propatula*.
- keokuk*, Hall, 1858, Geo. Rep. Iowa, p. 642, Keokuk Gr.
- keokuk* var. *shelbyensis*, Swallow, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 410, Keokuk Gr.
- laevigata*, Swallow, 1853, Trans. St. Louis Acad. Sci., vol. 2, p. 86, Keokuk Gr.
- laevis*, Hall, 1843, (*Delthyris laevis*), Geol. 4th Dist. N. Y., p. 345, and Pal. N. Y., vol. 4, p. 239, Portage Gr.
- lamellosa*, see *Athyris lamellosa*.
- laminosus*, McCoy, as identified by Geinitz, is *Spiriferina kentuckiensis*.
- lateralis*, Hall, 1858, Geo. Rep. Iowa, p. 661, Warsaw Gr.
- laticus*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 86, Waverly or Choteau Gr.
- leidy*, Norwood & Pratten, 1855, Jour. Acad. Nat. Sci., 2d series, vol. 3, p. 72, Kaskaskia Gr.
- leidy* var. *chesterensis*, Swallow, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 409, Kaskaskia Gr.
- leidy* var. *merrimacensis*, Swallow, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 410, Warsaw Gr.
- ligus*, Owen, 1852, Rep. Geo. Sur. Wis., Iowa, and Minn., p. 585, Ham. Gr.
- lineatoides*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 645, Burlington Gr.
- lineata*, Martin, 1809, (*Conchiliolithus Anomites lineatus*), Petrif. Derb., tab. 36, fig. 3, and 13th Rep. Geo. Sur. Ind., p. 133, Coal Meas.
- lineata* var. *striato-lineata*, Swallow, 1866, Trans. St. Louis Acad. Sci., vol. 2, p. 408, Coal Meas.
- littori*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 646, St. Louis Gr.
- logani*, Hall, 1858, Geo. Rep. Iowa, p. 647, Keokuk Gr.
- lonsdalei*, syn. for *S. disjuncta*.
- macra*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 134, and Pal. N. Y., vol. 4, p. 190, Schoharie grit and Up. Held. Gr.
- macra*, Meek. This name was preoccupied. See *S. strigosa*.
- macronota*, Hall, 1843, (*Delthyris macronota*), Geo. 4th Dist. N. Y., p. 206, and Pal. N. Y., vol. 4, p. 231, Ham. Gr.
- macropleura*, Conrad, 1840, (*Delthyris macropleura*), Anz. Rep. N. Y., p. 217, and Pal. N. Y., vol. 3, p. 202, Low. Held. Gr.
- macropleura*, Castelnau, 1843, Syst. Sil., p. 41. The name was preoccupied.
- macroptera*, as identified by d'Archiac & Verneuil, is *S. pennata*.
- macrothyris*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 132, and Pal. N. Y., vol. 4, p. 202, Up. Held. Gr.
- maia*, Billings, 1860, (*Athyris maia*), Can. Jour. Ind. Sci. and Arts, vol. 5, p. 276, Up. Held. Gr.
- manni*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 128, and Pal. N. Y., vol. 4, p. 211, Up. Held. Gr.
- marcyi*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 158, and Pal. N. Y., vol. 4, p. 220, Ham. Gr.
- marionensis*, Shumard, 1855, Geo. Rep. Mo., p. 203, Waverly or Choteau Gr.
- medialis*, Hall, 1843, (*Delthyris medialis*), Geo. 4th Dist. N. Y., p. 208, and Pal. N. Y., vol. 4, p. 207, Ham. Gr.
- medialis* var. *eatoni*, Hall, 1857, (*Spirifer eatoni*), 10th Rep. N. Y. Mus. Nat. Hist., p. 157, and Pal. N. Y., vol. 4, p. 229, Ham. Gr.
- meeki*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 645, Burlington Gr.
- meristoides*, Meek, 1868, Trans. Chi. Acad. Sci., p. 106, Ham. Gr.
- mesacostalis*, Hall, 1843, (*Delthyris mesacostalis* and *D. acuminata*), Geo. 4th Dist. N. Y., p. 269, and Pal. N. Y., vol. 4, p. 240, Chemung Gr.
- mesastrialis*, Hall, 1843, (*Delthyris mesastrialis*), Geo. 4th Dist. N. Y., p. 269, and Pal. N. Y., vol. 4, p. 242, Ham. and Chemung Gr.
- meta*, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 380, Niagara Gr.
- meusebachianus*, syn. for *Spirifer camera*.
- mexicana*, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 292, Permian Gr.
- missouriensis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 643, Waverly or Choteau Gr.
- modesta*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 61, and Pal. N. Y., vol. 3, p. 203, Low Held. Gr.
- mortonana*, S. A. Miller, 1883, 2d Ed., Am. p. 298, Pal. Foss. Keokuk Gr. Proposed instead of *S. fastigata* of Meek and Worthen, 1870, in Proc. Acad. Nat. Sci., p. 36, and afterward in Geo. Sur. Ill., vol. 6, p. 521, pl. 30, fig. 3, from Crawfordsville, Indiana.
- mucronata*, Conrad, syn. for *S. pennata*.
- multicostata*, Castelnau, 1843, Syst. Sil., p. 42. Not recognized.
- multigranosa*, Worthen, (in press), Geo. Sur. Ill., vol. 8, p. 105, Coal Meas.
- multistriata*, see *Trematospira multistriata*.
- murchisoni*, Castelnau, 1843, Syst. Sil., p. 41. Not recognized.
- mysticensis*, Meek, 1873, Hayden's Geo. Sur. Terr. 6th Rep., p. 466. Not satisfactorily defined.
- neglecta*, Hall, 1858, Geo. Rep. Iowa, p. 642, Keokuk Gr.

Syst. Sil., p.
upied.

d'Archiac &

Rep. N. Y.
d Pal. N. Y.,

r.
maia.) Can.
ol. 5, p. 276,

N. Y. Mus.
N. Y., vol. 4,

N. Y. Mus.
N. Y., vol.

Geo. Rep.
Choteau Gr.

ris medialis.)
08, and Pal.

Gr.
857, (Spirifer

Y. Mus. Nat.
Y., vol. 4, p.

ms., St. Louis
urlington Gr.

na. Chi. Acad.

Delthyris mes-
ta.) Geo. 4th

il. N. Y., vol.

Delthyris mes-
N. Y., p. 269,

42, Ham. and

N. Y. Mus.
Gr.

pirifera cam-

rana. St. Louis
Permian Gr.

30, Trans. St.
643, Waverly

ep. N. Y. Mus.
N. Y., vol. 3,

3, 2d Ed., Am.
Gr. Proposed

Meek and Wor-
Nat. Sci., p.

Sur. Ill., vol.
m Crawford-

S. pennata.
Syst. Sil., p.

press.) Geo.
al Meas.

a multistriata.
Syst. Sil., p. 41.

ayden's Geo.
66. Not satis-

Rep. Iowa, p.

newberryi, Hall, 1883, Rep. St. Geol. pl.
56, fig. 9, 10, Waverly Gr.

niagarensis, Conrad, 1842, Jour. Acad.
Nat. Sci., vol. 8, p. 261, and Pal. N. Y.,
vol. 2, p. 264, Niagara Gr.

niagarensis var. oligoptycha, Roemer,
1860, Sil. Fauna West Tenn., p. 68,
Niagara Gr.

nictavensis, Dawson, 1868, Acad. Geol., p.
499, Devonian.

norwoodana, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 7, and Bull. Am. Mus. Nat.
Hist., p. 48, Warsaw Gr.

norwoodi, Meek, 1860, Proc. Acad. Nat.
Sci., vol. 12, p. 308, Devonian.

novamexicana, S. A. Miller, 1881, Jour.
Cin. Soc. Nat. Hist., vol. 4, p. 314, Bur-
lington Gr.

nymphia, Billings, 1863, Proc. Port. Soc.
Nat. Hist., vol. 1, p. 116, Low.
Held. Gr.

octocostata, Hall, 1857, 10th Rep. N. Y.
Mus. Nat. Hist., p. 62, and Pal. N. Y.,
vol. 3, p. 205, Low. Held. Gr.

opima, Hall, 1858, Geo. Rep. Iowa, p. 711,
syn. for S. rockymontana.

orestes, Hall & Whitfield, 1873, 23d Rep.
N. Y. Mus. Nat. Hist., p. 237, Che-
mung Gr.

oregonensis, Shumard, 1863, Trans. St.
Louis Acad. Sci., vol. 2, p. 108, Coal
Meas.

osagensis, Swallow, 1860, Trans. St. Louis
Acad. Sci., vol. 1, p. 641, Waverly or
Choteau Gr.

oweni, Hall, 1857, 10th Rep. N. Y. Mus.
Nat. Hist., p. 129, Up. Held. Gr.

pachyptera, Goldfuss, as identified by Con-
rad in 1839, (Delthyris pachyptera).
Not American.

parryana, Hall, 1858, Geo. Rep. Iowa, vol.
1, pt. 2, p. 509, Ham. Gr.

peculiaris, Shumard, 1855, Geo. Rep. Mo.,
p. 202, Waverly or Choteau Gr.

pennata, Atwater, 1820, (Terebratula
pennata.) Am. Jour. Sci. and Arts, vol.
2, p. 242, Ham. Gr.

FIG. 612.—*Spirifer pennata*.

pennata, Owen. The name was preoccu-
pied, see *S. atwaterana*.

percrassa, McCoy, 1855, Brit. Pal. Rocks,
p. 194, Sil. Not satisfactorily identified
in America.

perextensa, Meek & Worthen, 1868, Geo.
Sur. Ill., vol. 3, p. 414, Ham. Gr.

(?) *perforata*, see *Trematospira perforata*.

perlamellosa, Hall, 1857, 10th Rep. N. Y.
Mus. Nat. Hist., p. 57, and Pal. N. Y.,
vol. 3, p. 200, Low. Held. Gr.

perplexa, McChesney, 1860, New Pal.
Foss., syn. for *S. lineata*.

pertenuis, Hall, 1857, 10th Rep. N. Y.
Mus. Nat. Hist., p. 163, Ham. Gr.

pharovicina, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 405, Portage Gr.

pinonensis, Meek, 1870, Proc. Acad. Nat.
Sci., p. 60, and Expl. 40th Parallel, vol.
4, p. 45, Up. Held. Gr.

planoconvexa, Shumard, 1855, Geo. Rep.
Mo., p. 202, Coal Meas.

plena, Hall, 1858, Geo. Rep. Iowa, p. 603,
Burlington Gr.

plicata, Vanuxem, 1843, see *S. vanuxemi*.

pluto, Clarke, 1885, Bull. U. S. Geo. Sur.,
No. 16, p. 31, Genesee shales.

praematura, Hall, 1867, Pal. N. Y., vol. 4,
p. 250, Chemung Gr.

prolata, Vanuxem, 1842, (Delthyris pro-
lata.) Geo. Rep. N. Y., p. 181, Che-
mung Gr.

propinqua, Hall, 1858, Geo. Rep. Iowa, p.
647, Keokuk Gr.

prora, Conrad, 1842, (Delthyris prora.)
Jour. Acad. Nat. Sci., vol. 8, p. 263,
Ham. Gr.

protena, syn. for *S. disjuncta*.

pseudolineata, Hall, 1858, Geo. Rep. Iowa,
p. 645, Keokuk Gr.

pulchra, Meek, 1860, Proc. Acad. Nat. Sci.,
p. 310, and Simpson's Gt. Basin of
Utah, p. 352, Coal Meas.

pyramidalis, see *Cyrtina pyramidalis*.

pyxidata, Hall, 1859, Pal. N. Y., vol. 3, p.
428, Oriskany sandstone.

racinensis, McChesney, 1860, Pal. Foss.,
p. 84, Niagara Gr.

radiata, Sowerby, 1839, Murch. Sil. Syst.,
p. 637, and Pal. N. Y., vol. 2, pp. 66, 265,
Niagara Gr.

rariocosta, Conrad, 1842, (Delthyris rari-
costa.) Jour. Acad. Nat. Sci., vol. 8, p.
262, and Pal. N. Y., vol. 4, p. 192, Scho-
harie grit and Up. Held. Gr.

resupinata, as identified by d'Archiac &
Verneuil. Not American.

richardsoni, Meek, 1868, Trans. Chi. Acad.
Sci., p. 104, Ham. Gr.

rockymontana, Marcou, 1858, Geo. N.
Amer., p. 50, Coal Meas.

rostellata, Hall, 1858, Geo. Rep. Iowa, p.
641, Keokuk Gr.

rostellum, Hall & Whitfield, 1872, 24th
Rep. N. Y. Mus. Nat. Hist., p. 182, Ni-
agara Gr.

rostrata, Morton, 1836, Am. Jour. Sci. and
Arts, vol. 29, p. 149, Coal Meas.

rugicosta, Hall, 1860, Can. Nat. Geo., vol.
5, p. 144, Up. Sil.

rugatina, Conrad, 1842, (Delthyris ruga-
tina.) Jour. Acad. Nat. Sci., vol. 8, p.
261, Niagara Gr.

saffordi, Hall, 1859, Pal. N. Y., vol. 3, p.
203, Low. Held. Gr.

scobina, Meek, 1860, Proc. Acad. Nat.
Sci., p. 310, and Simpson's Gt. Basin of
Utah, p. 351, Coal Meas.

sculptilis, Hall, 1843, (Delthyris sculpti-
lis.) Geo. Rep. 4th Dist. N. Y., p. 202,
and Pal. N. Y., vol. 4, p. 221,
Ham. Gr.

segmenta, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 131, and Pal. N. Y., vol. 4, p. 207, Up. Held. Gr.
semiplicata, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 111, Kinderhook Gr.
setigera, Hall, 1858, Geo. Rep. Iowa, p. 705, Kaskaskia Gr.
sheppardi, Castelnau, 1843, Syst. Sil., p. 42. Not recognized, but probably a variety of *Orthia lynx*.
sillana, Winchell, 1865, Proc. Acad. Nat. Sci., p. 119, Cuyaboga shale.
similior, see *Pentamerus similior*.
solidirostris, White, 1860, Bost. Jour. Nat. Hist., vol. 7, p. 232, Kinderhook Gr.
sowerbyi, Castelnau, 1843, Syst. Sil., p. 43. Not recognized.
spinosa, see *Spiriferina spinosa*.
staminea, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 105, Niagara Gr.
striatiformis, Meek, 1875, Ohio Pal., vol. 2, p. 289, Waverly Gr.

FIG. 613.—*Spirifer striata*. Interior of ventral valve.

North America, p. 49, Subcarboniferous.

substriatulus, as identified by d'Archiac & Verneuil. Not American.

FIG. 614.—*Spirifer striata*. Interior of dorsal valve.

strigosa, Meek, 1860, Proc. Acad. Nat. Sci., p. 309, and Simpson's Rep. Gt. Basin of Utah, p. 347, Devonian. Proposed instead of *S. macra*, Meek, which was preoccupied.
subaequalis, Hall, 1858, Geo. Rep. Iowa, p. 663, Warsaw Gr.
subattenuata, Hall, 1858, Geo. Rep., Iowa, index, p. 3, Ham. Gr.
subcardiformis, Hall, 1858, Geo. Rep. Iowa, p. 660, Warsaw Gr.
subcuspidata, Hall, 1858, Geo. Rep. Iowa, p. 646, Keokuk Gr. Preoccupied by Schnur in 1831. See *S. cuspidatiformis*.
subdecussata, Whiteaves, 1887, Cont. to Can. Pal., vol. 1, p. 114, Ham. Gr.
subelliptica, McChesney, 1860, New Pal. Foss. Not recognized, Coal Meas.
sublineata, Meek, 1868, Trans. Chi. Acad. Sci., p. 103, Ham. Gr.
submucronata, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 62, and Pal. N. Y., vol. 3, p. 419, Oriskany sandstone.

submucronata, Hall, 1858, Geo. Rep. Iowa, vol. 1, pt. 2, Ham. Gr. This name was preoccupied. See *S. subattenuata*.
suborbicularis, Hall, 1858, Geo. Rep. Iowa, p. 644, Keokuk Gr.
subrotundata, Hall, 1858, Geo. Rep. Iowa, vol. 1, pt. 2, p. 521, Kinderhook Gr.
subsulcata, Hall, 1860, Can. Nat. and Geol., vol. 5, Up. Sil. This name was preoccupied by Dalman in 1828.
subumbonata, see *Martinia subumbonata*.
subundifera, Meek & Worthen, 1868 Geo. Sur. Ill., vol. 3, p. 434, Ham. Gr.
subvaricosa, Hall & Whitfield, 1873, 23d Rep. N. Y. Mus. Nat. Hist., p. 237, Up. Held. Gr.
subventricosa, McChesney, syn. for *S. rockymontana*.
sulcata, Hisinger, 1831, (*Delthyris sulcata*) Anteckn. Physik. Och. Geognosi., p. 119, Pal. N. Y., vol. 2, p. 261, Niagara Gr.
sulcifera, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 293, Permian Gr.
superba, Billings, 1874, Pal. Foss., vol. 2, p. 45, Devonian. The name was preoccupied by Eichwald in 1842. See *S. billingsana*.
taneyensis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 645, Kinderhook Gr.
temeraria, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 314, Burlington Gr.
tenuicostata, Hall, 1858, Geo. Rep. Iowa, p. 662, Warsaw Gr.
tenuimarginata, Hall, 1858, Geo. Rep. Iowa, p. 641, Keokuk Gr.
tenuis, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 162, Ham. Gr.
tenuistriata, Hall, 1859, Pal. N. Y., vol. 3, p. 204, Low. Held. Gr.
tenuistriata, Shaler, 1865. The name was preoccupied.
texana, Meek, 1871, Proc. Acad. Nat. Sci., p. 179, Coal Meas.
texta, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 169, Waverly Gr.
translata, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 85, Kaskaskia Gr.
transversa, McChesney, 1860, New Pal. Foss., Kaskaskia Gr. Not recognized.
tribulis, Hall, 1859, Pal. N. Y., vol. 3, p. 420, Oriskany sandstone.
triplicata, Hall, syn. for *Spirifer camerata*.
troosti, Castelnau, 1843, Syst. Sil., p. 41. Not recognized.
tullia, Hall, 1867, Pal. N. Y., vol. 4, p. 218, Ham. Gr.
undulata, Vanuxem, 1843, (*Delthyris undulatus*) Geo. 3d Dist. N. Y., p. 132, Onondaga Gr. The name was preoccupied.
unica, Hall, 1867, Pal. N. Y., vol. 4, p. 203, Cornif. Gr.
utahensis, Meek, 1860, syn. for *S. norwoodii*.

o. Rep. Iowa,
his name was
tenuata.

Geo. Rep.

o. Rep. Iowa,
rhook Gr.
n. Nat. and
is name was
n 1828.

ubumbonata.
en, 1868 Geo.
am. Gr.
ld, 1873, 23d
t., p. 237, Up.

syn. for S.

lthyris sulca-
h. Geognosi.,
p. 261, Niag-

ans. St. Louis
Permian Gr.
Foss., vol. 2,
ame was pre-
1842. See S.

, Trans. St.
p. 645. Kinder-

1, Jour. Cin.
314, Burling-

o. Rep. Iowa,

, Geo. Rep.

o. N. i. Mus.

Gr.

N. Y., vol. 3,

the name was

cad. Nat. Sci.,

. N. Y. Mus.

ly Gr.

ans. St. Louis

85, Kaskas-

60, New Pal.

ot recognized.

. Y., vol. 3, p.

Spirifer cam-

st. Sil., p. 41.

, vol. 4, p. 218,

(Delthyris un-

ne was preoc-

Y., vol. 4, p.

. for S. nor-

vanuxemi, Hall, 1859, Pal. N. Y., vol. 3,
p. 198, Low. Held. Gr., described as
Orthis plicata by Vanuxem in the Geo.
Rep. 3d Dist. N. Y., but that name was
preoccupied.

varicosa, Hall, 1857, 10th Rep. N. Y. Mus.
Nat. Hist., p. 130, and Pal. N. Y., vol.
4, p. 205, Up. Held. Gr.

ventricosa, see Nucleospira ventricosa.

venusta, syn. for Spirifer divaricata.

vernonensis, Swallow, 1860, Trans. St.
Louis Acad. Sci., vol. 1, p. 644, Wa-
verly or Choteau Gr.

verneuili, syn. for S. disjuncta.

? waldronensis, see Triplesia waldronensis.

waverlyensis, Winchell, 1870, Proc. Am.

Phil. Soc., vol. 12, p. 251, Marshall Gr.

whitneyi, Hall, 1858, Geo. Rep. Iowa, p.

502, Ham. and Chemung Gr.

wortheni, Hall, 1857, 10th Rep. N. Y.

Mus. Nat. Hist., p. 156, Ham. Gr.

ziczac, Hall, 1843. The name was preoc-

cupied by Roemer. See S. consobrina.

SPIRIFERINA, D'Orbigny, 1847, Consid. Zool.

et Geol. Sur. les Brachiopodes, Com-

ptes rendus des Sciences de l'Académie

des Sciences. [Ety. *Spirifera*, a genus;

inus, implying resemblance.] Shell

transverse, valves unequally convex;

with or without mesial fold and sinus;

smooth or costated; beak straight or

recurved; area large, and interrupted

by a pseudodeltidium, notched near

the cardinal edge; structure punctate;

surface spinous; tooth on each side of

the fissure, supported by vertical,

shelly plates, the space intervening oc-

cupied by the cardinal muscles; me-

sial septum wide at the base, and taper-

ing to an acute blade; dorsal valve

with dental sockets and shelly lamellæ,

for the support of serrated arms in the

form of two large spiral, horizontal

cones. Type *S. rostrata*.

billingsi, Shumard, 1858, Trans. St. Louis

Acad. Sci., vol. 1, p. 294, Permian Gr.

binacuta, Winchell, 1865, Proc. Acad.

Nat. Sci., p. 120, Burlington Gr.

clarksvillensis, Winchell, 1865, Proc. Acad.

Nat. Sci., p. 119, Marshall Gr.

kentuckiensis, Shumard, 1855, (*Spirifera*

kentuckiensis,) Geo. Rep. Mo., p. 203,

Coal Meas.

FIG. 615.—*Spiriferina kentuckiensis*.

kentuckiensis var. propatula, Swallow,

1866, (*Spirifera kentuckiensis* var. pro-

patula,) Trans. St. Louis Acad. Sci.,

vol. 2, p. 409, Coal Meas.

spinosa, Norwood & Pratten. (*Spirifera*

spinosa,) 1855, Jour. Acad. Nat. Sci.,

vol. 3, 2d series, p. 71, Kaskaskia Gr.

spinosa var. campestris, White, 1874, Rep.

Invert. Foss., p. 21, and Geo. Sur. W.

100th Mer., vol. 4, p. 139, Carb.

subtexta, White, 1862, Proc. Bost. Soc.

Nat. Hist., vol. 9, p. 8, Burlington Gr.

Spirigera, D'Orbigny in Comptes Rendus, t.

25, p. 268, syn. for *Athyris*.

americana, see *A. americana*.

biloba, see *A. biloba*.

caput-serpentis, see *A. caput-serpentis*.

charitonensis, see *A. charitonensis*.

clintonensis, see *A. clintonensis*.

concentrica, syn. for *A. spiriferoides*.

corpulenta, see *A. corpulenta*.

eborea, see *A. eborea*.

euzona, see *A. euzona*.

formosa, see *A. formosa*.

fultonensis, see *A. fultonensis*.

hannibalensis, see *A. hannibalensis*.

hawni, see *A. hawni*.

jacksoni, see *A. jacksoni*.

maconensis, see *A. maconensis*.

minima, see *A. minima*.

missouriensis, see *A. missouriensis*.

monticola, see *A. monticola*.

obmaxima, see *A. obmaxima*.

ohioensis, see *A. ohioensis*.

pectinifera, see *A. pectinifera*.

plattensis, see *A. plattensis*.

prouti, see *A. prouti*.

reflexa, see *A. reflexa*.

singletoni, see *A. singletoni*.

spiriferoides, see *A. spiriferoides*.

STENOSCHISMA, Conrad, 1839, Ann. Rep.

N. Y., p. 59. [Ety. *stenos*, narrow;

schisma, fissure.] Written *Stenocisma*

by Conrad. Subtriangular, ovoid, or

subglobose, hinge-line short; beak of

ventral valve extended, attenuate, more

or less arcuate, and appressed upon the

opposite valve; mesial fold and sinus;

surface plicated, valves articulated by

teeth and sockets; median septum in

dorsal valve, on each side of which the

crura are supported. Type *S. formo-*

sum. Conrad mentioned *Terebratula*

schlotheimi as the type which is now

the type of *Camarophoria*; but, as Hall

shows, Conrad was mistaken in identi-

fying what is now known as *S. formo-*

sum, with the European *Camarophoria*

schlotheimi.

billingsi, Hall, 1867, Pal. N. Y., vol. 4, p.

336, Cornif. Gr. The same that Bill-

ings called *Rbynchonella thalia*, Can.

Jour. 1860, but the name was preoc-

cupied.

carica, Hall, 1867, Pal. N. Y., vol. 4, p. 344, ✓

Ham. Gr.

carolina, Hall, 1867, Pal. N. Y., vol. 4, p.

337, Cornif. Gr.

congregatum, Conrad, 1841, (*Atrypa con-*

gregata,) Ann. Rep. N. Y., p. 55, and

Pal. N. Y., vol. 4, p. 341, Ham. Gr.

contractum, Hall, 1843, (*Atrypa con-*

tracta,) Geo. 4th Dist. N. Y., pl. 66, fig.

3a, and Pal. N. Y., vol. 4, p. 351, Che-

mung Gr.

contractum var. saxatile, Hall, 1867, Pal.

N. Y., vol. 4, p. 417, Chemung Gr.

dotis, Hall, 1867, Pal. N. Y., vol. 4, p. 344,

Ham. Gr.

ame was pre-

Geol., pl. 41,

nn. Rep. Geo.

. Wis., vol. 4,

13, (Stropho-

r. Acad. Nat.

al. N. Y., vol.

—Streptorhynchus sulcatum. Dorsal

Strophomena
Sci. Phil., p.
p. 113, Tren-

a. Quar. Jour.
Riv. Gr. A

ena filitexta,
Trenton and

Desc. New
p. 200, Up.

um. Exterior
valve.

4, Cin. Quar.
Hud. Riv. Gr.

, syn. for S.
num.

umbraculum, Whit-
1882, Desc.

Spec. Foss.,
Ohio, p. 193,

Held. Gr.
, White &

st. Soc. Nat.
rhook Gr.

st. Soc. Nat.
ing Gr.

U. S. Geo.
Gr.

ipronites nu-
p. 77, Hud.

for Meekella

pandora, Billings, 1860, Can. Jour., vol. 5,
p. 266, and Pal. N. Y., vol. 4, p. 68,
Schlotheim grit and Cornif. Gr.

pectinaceum, Hall, 1843, (Strophomena
pectinacea and S. bifurcata,) Geo. Rep.
4th Dist. N. Y., p. 266, and Pal. N. Y.
vol. 4, p. 73, Chemung Gr.

perversum, Hall, 1857, (Orthis perversa,) 10th
Rep. N. Y. Mus. Nat. Hist., p. 137,
and Pal. N. Y., vol. 4, p. 72, (Orthisina
alternata, 1860, 13th Rep.,) Cornif. and
Ham. Gr.

planoconvexum, Hall, 1847, (Leptæna
planoconvexa,) Pal. N. Y., vol. 1, p.
114, Hud. Riv. Gr.

planumbonum, Hall, 1847, (Leptæna
planumbona,) Pal. N. Y., vol. 1, p. 112,
Trenton and Hud. Riv. Gr.

primordiale, Whitfield, 1886, Bull. Am. Mus.
Nat. Hist., vol. 1, p. 301, Birdseye Gr.
pyramidale, Newberry, syn. for Meekella
striatocostata.

radium, Vanuxem, 1843, (Strophomena
radiata,) Geo. Rep. 3d Dist. N. Y., p.
122, and Pal. N. Y., vol. 3, p. 193, Low.
Held. Gr.

rectum, Conrad, 1843, (Strophomena
recta,) Proc. Acad. Nat. Sci., vol. 1, p.
332, and Pal. N. Y., vol. 1, p. 113, Black
Riv. and Trenton Grs.

sinuatum, Emmons, 1855, Am. Geol., p.
199, Hud. Riv. Gr.

subplanum, Conrad, 1842, Jour. Acad.
Nat. Sci., vol. 8, p. 258, and Pal. N. Y.,
vol. 2, p. 259, (Strophomena subplana,) 1
Niagara Gr.

subtentum, Conrad, 1847, (Strophomena
subtenta,) Pal. N. Y., vol. 1, p. 115,
Trenton and Hud. Riv. Gr.

FIG. 620.—Streptorhynchus sulcatum.
Interior of dorsal
valve.

sulcatum, Verneuil,
1848, (Leptæna sul-
cata,) Bull. Geol. Soc.
France, vol. 5, p.
350, and Ohio Pal.,
vol. 1, p. 85, Hud.
Riv. Gr.

tenue, Hall, 1863,
Trans. Alb. Inst.,
vol. 4, p. 210, Niag-
ara Gr.

thalia, Billings, 1860, (Strophomena
thalia,) Can. Nat. and Geol., vol. 5, p.
39, Trenton Gr.

umbraculum, Schlotheim, 1820, (Tere-
bratulites umbraculum,) Petrefakten-
kunde, p. 256, Devonian to the Per-
mian Gr.

vetustum, James, 1874, Cin. Quar. Jour.
Sci., vol. 1, p. 241, Hud. Riv. Gr. One
of the forms of S. subtentum.

woolworthanum, Hall, 1857, (Stropho-
mena woolworthiana,) 10th Rep. N. Y.
Mus. Nat. Hist., p. 43, and Pal. N. Y.,
vol. 3, p. 192, Low. Held. Gr.

Stricklandia, Billings, 1859, Can. Nat. Geo.,
vol. 4. This name having been pre-
viously applied to a genus of fossil
plants, the author abandoned it and
proposed Stricklandinia.

STRICKLANDINIA, Billings, 1863, Can. Nat.
and Geo., vol. 8, p. 370. [Ety. proper
name.] Large, elongate-oval, trans-
versely subcircular, sometimes com-
pressed, valves subequal; short mesial
septum in the interior of the ventral
valve, supporting a small triangular
chamber, beneath the beak, as in Pen-
tamerus; in the dorsal valve no longi-
tudinal septum, spires, or loop; two
short, rudimental plates, bearing pro-
cesses. Type S. gaspensis.

anticostiensis, Billings, 1863, Can. Nat.
Geo., vol. 8, p. 370, Anticosti Gr.

(?) arachne, Billings, 1862, Pal. Foss., vol.
1, p. 85, Quebec Gr.

(?) arethusa, Billings, 1862, Pal. Foss., vol.
1, p. 85, Quebec Gr.

brevis, Billings, 1859, Can. Nat. Geo., vol.
4, p. 135, Mid. Sil.

canadensis, Billings, 1859, Can. Nat. Geo.,
vol. 4, p. 135, Clinton Gr.

castellana, White, 1876, Proc. Acad. Nat.
Sci., p. 30, Niagara Gr.

davidsoni, Billings, 1868, Lond. Geo.
Mag., vol. 5, p. 59, Up. Sil.

deformis, Meek & Worthen, 1870, Proc.
Acad. Nat. Sci., Phil., p. 37, and Geo.
Sur. Ill., vol. 6, p. 502, Niagara Gr.

elongata, see Amphigenia elongata.

elongata var. curta, see Amphigenia curta.
gaspensis, Billings, 1859, Can. Nat. Geo.,
vol. 4, p. 134, Mid. Sil.

melissa, Billings, 1874, Pal. Foss., vol. 2,
p. 89, Mid. Sil.

multilirata, Whitfield, 1878, Ann. Rep.
Geo. Sur. Wis., p. 81, and Geo. Wis.,
vol. 4, p. 315, Niagara Gr.

salteri, Billings, 1874, Pal. Foss., vol. 2, p.
87, Anticosti Gr., Mid. Sil.

STROPHALOSIA, King, 1844, Ann. and Mag.
Nat. Hist., vol. 14, p. 313. [Ety. strophe,
a bending; alos, a disk.] Having the
general form and muscular impressions
of Leptæna, with the tubuliferous or
spinous surface of Productus; pos-
sessed of a well-developed condyloid
hinge, area, and deltidium. Type S.
excavata.

horrescens, Geinitz, 1866, Carb. und Dyas
in Neb., p. 49. Prof. Meek regarded this
name as a syn. for Productus nebras-
kensis.

numularis, Winchell, 1863, Proc. Acad.
Nat. Sci., p. 4, Marshall Gr.

STROPHODONTA, Hall, 1852, Pal. N. Y., vol.
2, p. 63. [Ety. strophos, bent; odous,
tooth.] General form and characters as
in Strophomena, one valve convex and
the other concave, and following nearly
the same curve as the convex one, leav-
ing only a thin space for the animal,
and the surface radiated; distinguished,
however, by a crenulated hinge-line;
the absence of a foramen in the area of
the ventral valve; dental lamellæ ab-
sent, or nearly so; the divaricator mus-
cular impressions spreading, flabelli-
form, without limitation, by an elevated

- ridge; cardinal process in the dorsal valve bifurcated from its origin, and directed backward beneath the area of the ventral valve; on the lower side of the ventral area a bilobed process is embraced by the divisions of the cardinal process of the dorsal valve. Type *S. demissa*.
- ✓ *aequicostata*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 639, Ham. Gr.
- ✓ *altidorsata*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 637, Ham. Gr.
- alveata*, Hall, 1863, 10th Rep. N. Y. Mus. Nat. Hist., p. 36, and Pal. N. Y., vol. 4, p. 81, Schoharie grit.
- ampla*, see *Strophonella ampla*.
- arcuata*, Hall, 1858, Geo. of Iowa, p. 492, Ham. Gr.
- becki*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 52, and Pal. N. Y., vol. 3, p. 191, Low. Held. Gr.
- boonensis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 638, Ham. Gr.
- cælata*, see *Strophonella cælata*.
- callawayensis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 638, Ham. Gr.
- callosa*, Hall, 1863, 16th Rep. N. Y. Mus. Nat. Hist., p. 36, and Pal. N. Y., vol. 4, p. 82, Schoharie grit.
- ✓ *calvini*, S. A. Miller, 1883, 2d Ed. Am. Pal. Foss., p. 298, Upper Helderberg Gr. Proposed instead of *S. quadrata*, Calvin, 1878, in Bull. U. S. Geo. Sur. Terr., vol. 4, No. 3, p. 728, which was preoccupied.
- canace*, Hall & Whitfield, 1873, 23d Rep. N. Y. Mus. Nat. Hist., p. 236, Chemung Gr.
- cavumbona*, see *Strophonella cavumbona*.
- cayuta*, Hall, 1867, Pal. N. Y., vol. 4, p. 110, Chemung Gr.
- cincta*, Winchell, 1866, Rep. Low. Penin. Mich., p. 93, Ham. Gr.
- concava*, Hall, 1857, (*Strophomena concava*), 10th Rep. N. Y. Mus. Nat. Hist., p. 115, and Pal. N. Y., vol. 4, p. 96, Cornif. and Ham. Grs.
- costata*, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 585, Devonian.
- crebristriata*, Conrad, 1842, (*Strophomena crebristriata*), Jour. Acad. Nat. Sci., vol. 8, p. 254, and Pal. N. Y., vol. 4, p. 86, Schoharie grit.
- cymbiformis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 635, Ham. Gr.
- demissa*, Conrad, 1842, (*Strophomena demissa*), Jour. Acad. Nat. Sci., vol. 8, p. 258, and Pal. N. Y., vol. 4, p. 81, Schoharie grit, Cornif., Ham., and Chemung Grs.
- erratica*, Winchell, 1866, Rep. Low. Peninsular Mich., p. 93, Ham. Gr.
- feildeni*, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 693, Up. Sil.
- fragilis*, Hall, syn. for *Strophodonta perplana*.
- geniculata*, Hall, 1859, Pal. N. Y., vol. 3, p. 483, Low. Held. Gr.
- headleyana*, Hall, 1857, N. Y. Mus. Nat. Hist., p. 49, and Pal. N. Y., vol. 3, p. 185, Low. Held. Gr.
- hemispherica*, Hall, 1857, (*Strophomena hemispherica*), 10th Rep. N. Y. Mus. Nat. Hist., p. 113, and Pal. N. Y., vol. 4, p. 90, Schoharie grit and Cornif. Gr.
- hybrida*, Hall & Whitfield, 1873, 23d Rep. N. Y. Mus. Nat. Hist., p. 239, Chemung Gr.
- imitata*, Winchell, 1866, Rep. Low. Penin. Mich., p. 93, Ham. Gr.
- inequigradiata*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 113, and Pal. N. Y., vol. 4, p. 87, Schoharie grit and Cornif. Gr.
- inequistriata*, Conrad, 1842, (*Strophomena inequistriata*), Jour. Acad. Nat. Sci., vol. 8, p. 254, and Pal. N. Y., vol. 4, p. 93, Cornif. and Ham. Grs., Moscow shales.
- indentata*, Conrad, 1838, (*Leptæna indentata*), Ann. Rep., N. Y., p. 117, Low. Held. Gr. Not properly defined.
- inflexa*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 637, Ham. Gr.
- intermedia*, Hall, 1859, Pal. N. Y., vol. 3, p. 482, Oriskany sandstone.
- iowensis*, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 585, Devonian.
- junia*, Hall, 1867, Pal. N. Y., vol. 4, p. 108, Cornif., Ham. and Tully Grs. (Changed from *textilis*, in the corrigenda and index.)
- kemperri*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 636, Ham. Gr.
- leavenworthana*, see *Strophonella leavenworthana*.
- lepida*, Hall, syn. for *S. naerea*.
- lincklaeni*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 55, and Pal. N. Y., vol. 3, p. 415, Oriskany sandstone.
- magnifica*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 54, and Pal. N. Y., vol. 3, p. 414, Oriskany sandstone.
- magniventra*, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 54, and Pal. N. Y., vol. 3, p. 411, Oriskany sandstone.
- mucronata*, Conrad, 1842, (*Strophomena mucronata*), Jour. Acad. Nat. Sci., vol. 8, p. 257, and Pal. N. Y., vol. 4, p. 111, Chemung Gr.
- naerea*, Hall, 1857, (*Strophomena naerea*), 10th Rep. N. Y. Mus. Nat. Hist., p. 144, and Pal. N. Y., vol. 4, p. 104, Cornif. and Ham. Grs.
- navalis*, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 635, Ham. Gr.
- parva*, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 584, Ham. Gr.

FIG. 621.—*Strophodonta inequistriata*.

N. Y., vol. 3,
Y. Mus. Nat.
Y., vol. 3, p.

Strophomena
N. Y. Mus.
N. Y., vol.
t and Cor.

37373, 23d Rep.
p. 239, Che-

Low. Penin.

h Rep. N. Y.
d Pal. N. Y.,
rit and Cor.

Strophodonta
alstriata.

ena indenta,
ow Held. Gr.

ms. St. Louis
am. Gr.

N. Y., vol. 3,

Sur. Wis.,
evonian.

7, vol. 4, p.
Tully Gr.

n the corri-

ms. St. Louis
am. Gr.

ella leaven-

ea.

Rep. N. Y.
d Pal. N. Y.,

ndstone.

Rep. N. Y.
d Pal. N. Y.,

ndstone.

h Rep. N. Y.
d Pal. N. Y.,

ndstone.

Strophomena
at. Sci., vol.

ol. 4, p. 111,

Strophomena
at. Sci., vol.

ol. 4, p. 111,

Strophomena
at. Sci., vol.

ol. 4, p. 111,

Strophomena
at. Sci., vol.

ol. 4, p. 111,

parva, Hall, 1863, 16th Rep. N. Y. Mus.
Nat. Hist., p. 37, Schoharie grit. This
name was preoccupied.

patersoni, Hall, 1857, (*Strophomena* *patersoni*), 10th Rep. N. Y. Mus. Nat. Hist.,
p. 114, and Pal. N. Y., vol. 4, p. 89, Scho-
harie grit and Cornif. Gr.

perplana, Conrad, 1842, (*Strophomena*
perplana), Jour. Acad. Nat. Sci., vol. 8,
p. 257, and Pal. N. Y., vol. 4, p. 92,
Onondaga, Schoharie, Cornif., Ham.,
and Chemung. Grs.

perplana var. *nervosa*, Hall, 1843, (*Stro-*
phomena nervosa), Geo. Rep. 4th Dist.
N. Y., p. 286, and Pal. N. Y., vol. 4, p.
113, Chemung Gr.

planulata, Hall, 1859, Pal. N. Y., vol. 3,
p. 184, Low. Held. Gr.

plicata, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 90, Ham. Gr.

prisca, Hall, 1852, Pal. N. Y., vol. 2, p. 63,
Clinton Gr.

profunda, Hall, 1852, (*Leptaena profunda*),
Pal. N. Y., vol. 2, p. 61, Clinton and
Niagara Grs.

punctulifera, see *Strophonella punctu-*
lifera.

quadrata, Swallow, 1860, Trans. St. Louis
Acad. Sci., vol. 1, p. 639, Ham. Gr.

quadrata, Calvin, 1878, Bull. U. S. Geo.
Sur. Terr., vol. 4, No. 3, p. 728. The
name was preoccupied. See S. calvini.

reversa, see *Strophonella reversa*.

semifasciata, see *Strophonella semifasciata*.

striata, Hall, 1843, (*Strophomena striata*),
Geo. Rep. 4th Dist. N. Y., p. 104, Ni-
agara Gr.

subcymbiformis, Swallow, 1860, Trans.
St. Louis Acad. Sci., vol. 1, p. 636,
Ham. Gr.

subdmissa, Hall, 1857, 10th Rep. N. Y.
Mus. Nat. Hist., p. 145, Ham. Gr.

textilis, Hall, 1852, Pal. N. Y., vol. 2, p.
327, Coralline Limestone.

variabilis, Calvin, 1878, Bull. U. S. Geo.
Sur., vol. 4, No. 3, p. 727, Up.
Held. Gr.

varistriata, Conrad, 1842, (*Strophomena*
varistriata), Jour. Acad. Nat. Sci., vol.
8, p. 255, and Pal. N. Y., vol. 3, p. 180,
Low. Held. Gr.

varistriata var. *arata*, Hall, 1859, Pal.
N. Y., vol. 3, p. 183, Low. Held. Gr.

vascularia, Hall, 1859, Pal. N. Y., vol. 3,
p. 412, Oriskany sandstone.

STROPHOMENA, Rafinesque, 1825, Manuel de
Malacologie of Blainville, p. 513. [Ety.
strophos, bent; *mene*, a crescent.] Shell
somewhat semicircular, or somewhat
semioval, though variable in outline;
thin; one valve convex, the other con-
cave, with a thin space between them for
the animal; surface with radiating striae;
hinge-line straight, longer or shorter
than the width of the shell below;
anterior end of the shell usually de-
flected or bent over toward the dorsal
valve; ventral valve convex, flat on the
umbo; beak small, and minutely per-

forated; cardinal area nearly cut in two
parts by an angular notch, which is
closed, or nearly closed, by the bifid
cardinal process of the dorsal valve;
two divergent teeth, two adductor scars,
and two cardinal muscular impressions;
dorsal valve having a linear area, two
cardinal processes close together at the
middle of the hinge-line, directed for-
ward; sockets for the reception of the
teeth of the ventral valve; two mus-
cular scars in front of the cardinal pro-
cesses. Type *S. rugosa*, which is gener-
ally regarded as synonymous with *S.*
rhomboidalis.

acutiradiata, see *Chonetes acutiradiatus*.

alternata, Conrad, 1838, (*Leptaena alter-*
nata), Ann. Rep. N. Y., p. 115, and
Pal. N. Y., vol. 1, pp. 102, 286, Trenton
and Hud. Riv. Grs.

alterniradiata, Shaler, 1865, Bulletin No.
4, M. C. Z., Anticosti Gr. Not defined
so as to be recognized.

alternistriata, Hall, 1847, Pal. N. Y., vol.
1, p. 109, Trenton and Hud. Riv. Grs.

alternata var.

loxorhytis,

Meeke, 1873,

Ohio Pal., vol.

1, p. 91, Hud.

Riv. Gr.

ampla, see *Stro-*

phonella

ampla.

analoga, Phillips,

1836, Geol.

Yorkshire, vol. 2, pl. 7, fig. 10, Sub-

carb.

anticostiensis, syn. for *Strophomena alter-*

nata.

antiquata, see *Streptorhynchus anti-*

quatam.

arctostriata, see *Streptorhynchus arcto-*

striatum.

arcuata, Shaler, 1865. This name was pre-

occupied.

arethusa, Billings, 1862, Pal. Foss., vol. 1,

p. 132, Hud. Riv. Gr.

aurora, Billings, 1865, Pal. Foss., vol. 1,

p. 218, Quebec. Gr.

bifurcata, syn. for *Streptorhynchus pecti-*

naceum.

bipartita, Hall, 1852, (*Leptaena bipartita*),

Pal. N. Y., vol. 2, p. 326, Coralline Lime-

stone.

blainvillii, Billings, 1874, Pal. Foss., vol.

2, p. 28, Up. Sil.

camerata, Conrad, 1842, Jour. Acad. Nat.

Sci., vol. 8, p. 254, and Pal. N. Y., vol.

1, p. 106, Trenton Gr.

carinata, Conrad, 1838, see *Chonetes*

carinatus.

carinata, Conrad, 1842, see *Chonetes*

carinatus.

ceres, Billings, 1860, Can. Nat. and Geo.,

vol. 5, p. 54, Hud. Riv. Gr. and

Mid. Sil.

chemungensis, Conrad, 1842, Jour. Acad.

Nat. Sci., vol. 8, p. 257, Chemung Gr.

FIG. 622.—*Strophomena*
alternata.

- concava*, see *Strophodonta concava*.
conradi, Hall, 1859, Pal. N. Y., vol. 3, p. 194, Low. Held. Gr.
convexa, Owen, 1840, Rep. on Mineral Lands, p. 70, Calcif. Gr.
cornuta, see *Chonetes cornutus*.
corrugata, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 3, p. 256, and Pal. N. Y., vol. 2, p. 59, Clinton Gr.
crebristriata, see *Strophodonta crebri-striata*.
crenistrata, syn. for *Strophodonta perplana*.
declivis, James, syn. for *Strophomena alternata*.
deflecta, see *Streptorhynchus deflectum*.
deltoides, Conrad, 1839, Ann. Rep. N. Y., p. 64, and Pal. N. Y., vol. 4, p. 106, Trenton Gr.
delthyris, syn. for *Strophodonta perplana*.
demissa, see *Strophodonta demissa*.
depressa, Sowerby, 1825, (Producta depressa,) Min. Conchology, vol. 5, p. 86, and Pal. N. Y., vol. 2, p. 257, Up. Sil. Generally regarded as a synonym for *S. rhomboidalis*.
depressa var. *ventricosa*, see *Strophomena rugosa* var. *ventricosa*.
✓ *donneti*, Salter, 1852, Sutherland's Jour., vol. 2, App., p. 218, Devonian.
elegantula, Hall, 1843, Geo. Rep., 4th Dist. N. Y., p. 73, Clinton Gr.
elongata, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 259, Low. Held. Gr.
elliptica, Conrad, 1839, Ann. Rep. N. Y., p. 64, Niagara Gr.
fasciata, Hall, 1847, (Leptæna fasciata,) Pal. N. Y., vol. 1, p. 20, Chazy Gr.
filixeta, see *Streptorhynchus filixetum*.
fluctuosa, Billings, 1860, Can. Nat. Geo., vol. 5, p. 57, Trenton and Hud. Riv. Grs.
fontinalis, White, 1874, Rep. Invert. Foss., p. 10, and Geo. Sur. W. 100th Mer., vol. 4, p. 54, Quebec Gr.
fracta, Meek, 1873, (S. alternata var. fracta,) Pal. Ohio, vol. 1, p. 91, Hud. Riv. Gr.
fragilis, syn. for *Strophodonta perplana*.
galatea, Billings, 1874, Pal. Foss., vol. 2, p. 20, Gaspé limestone No. 8, Devonian.
geniculata, Shaler, (Brachyprion geniculatum.) The name was preoccupied.
gibbosa, Conrad, 1841, Ann. Geo. Rep. N. Y., p. 25, Onondaga Gr.
hecuba, Billings, 1860, Can. Nat. Geo., vol. 5, p. 60, Hud. Riv. Gr.
hemispherica, see *Strophodonta hemispherica*.
imbecilis, Billings, 1865, Pal. Foss., vol. 1, p. 219, Quebec Gr.
imbrex, Pander, 1845, in Russia and Ural Mountains, Hud. Riv. Gr. The identification very doubtful in America.
impressa, syn. for *Strophodonta varistriata*.
incrassata, Hall, 1847, (Leptæna incrassata,) Pal. N., vol. 1, p. 19, Chazy to Hud. Riv. Gr.
inequigradiata, see *Strophodonta inequigradiata*.
inequistriata, see *Strophodonta inequistriata*.
interstitialis, Phillips, in Geo. 4th Dist. N. Y., see *Strophodonta cayuta*.
irene, Billings, 1874, Pal. Foss., vol. 2, p. 27, Devonian.
ithacensis, Vanuxem, 1842, Geo. Rep. N. Y., p. 174, Portage Gr.
julia, Billings, 1862, Pal. Foss., vol. 1, p. 127, Anticosti Gr., Div. 4, Mid. Sil.
kingi, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 72, and Geo. Wis., vol. 4, p. 261, Hud. Riv. Gr.
lachrymosa, see *Productella lachrymosa*.
levis, Emmons, 1842, Geo. Rep. N. Y., p. 385, Birdseye Gr.
leda, Billings, 1860, Can. Nat. Geo., vol. 5, p. 55, Mid. Sil.
lepida, syn. for *Strophodonta naerea*.
lima, see *Productella lachrymosa* var. *lima*.
lineata, see *Chonetes lineatus*.
macra, syn. for *Strophodonta semifasciata*.
magniventra, see *Strophodonta magniventra*.
membranacea, of Phillips, as identified by Vanuxem, 1842, Geo. 3d Dist. N. Y., see *Productella hirsuta*.
modesta, Conrad, 1839, Ann. Rep. N. Y., p. 64, Niagara Gr.
mucronata, see *Strophodonta mucronata*.
naerea, see *Strophodonta naerea*.
nasuta, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 260, Trenton and Hud. Riv. Grs.
nasula, Conrad, 1846, Proc. Acad. Nat. Sci., vol. 3, p. 23. Not defined so as to be recognized.
nemea, Hall & Whitfield, 1877, U. S. Geo. Sur. 40th parallel, vol. 4, p. 233, Quebec Gr.
nervosa, see *Strophodonta perplana* var. *nervosa*.
niagarensis, Winchell & Marcy, syn. for *Strophodonta profunda*.
nitens, Billings, 1860, Can. Nat. Geo., vol. 5, p. 53, Hud. Riv. Gr.
nutans, see *Streptorhynchus nutans*.
obscura, Hall, 1852, (Leptæna obscura,) Pal. N. Y., vol. 2, p. 62, Clinton Gr.
orthididea, Hall, 1852, (Leptæna orthididea,) Pal. N. Y., vol. 1, p. 62, Clinton Gr.
patenta, Hall, 1852, (Leptæna patenta,) Pal. N. Y., vol. 2, p. 60, Clinton Gr.
patersoni, see *Strophodonta patersoni*.
pecten, Linnæus, 1758, (Anomia pecten,) Syst. Nat., Niagara Gr. Not American.
pectinacea, see *Streptorhynchus pectinaceum*.
perplana, see *Strophodonta perplana*.
philomela, Billings, 1860, Can. Nat. Geo., vol. 5, p. 56, Mid. Sil.
planoconvexa, see *Streptorhynchus planoconvexum*.
planumbona, see *Streptorhynchus planumbonum*.

plicata, syn. for *Streptorhynchus strigentum*.

plicifera, Hall, 1847, (*Leptæna plicifera*.) Pal. N. Y., vol. 1, p. 19, Chazy Gr.

pluriatriata, syn. for *Strophodonta periplana*.

profunda, see *Strophodonta profunda*.

punctulifera, see *Strophonella punctulifera*.

pustulosa, syn. for *Productella truncata*.

radiata, see *Streptorhynchus radiatum*.

recta, see *Streptorhynchus rectum*.

rectilateria, syn. for *Strophodonta varistriata*.

reticulata, Shaler, 1865, Bulletin No. 4, M. C. Z., Anticosti Gr. Not defined so as to be recognized.

rhomboidalis, Wilckins, 1769, (*Conchites rhomboidalis*.) Nachricht von Seltenen Verst., p. 77. This species ranges from the Trenton Gr. to the Keokuk, regarding *S. tenuistriata*, *S. depressa*, and *S. rugosa* as varieties only. The type, however, is the Devonian form.

FIG. 623.—*Strophomena rugosa*.

rugosa, Dalmann, 1827, (*Leptæna rugosa*.) Vet. Acad. Handlinger, p. 106, and Pal. N. Y., vol. 3, p. 195, Niagara and Low.

Held. Gr. This form is supposed to be the type of Rafinesque's genus *Strophomena*. The species is usually regarded as merely a variety of *S. rhomboidalis*. *rugosa* var. *ventricosa*, Hall, 1857, (*S. depressa* var. *ventricosa*.) 10th Rep. N. Y. Mus. Nat. Hist., p. 53, and Pal. N. Y., vol. 3, p. 417, Oriskany sandstone.

setigera, see *Chonetes setigerus*.

semiovalis, Conrad, syn. for *Leptæna sericea*.

semiovalis, Shaler. The name had been twice preoccupied.

squamula, James, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 335, Hud. Riv. Gr.

striata, see *Strophodonta striata*.

subdemissa, syn. for *Strophodonta demissa*.

subplana, see *Streptorhynchus subplanum*.

subtenta, see *Streptorhynchus subtentum*.

syrtalis, syn. for *Chonetes carinatus*.

tenuilineata, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 259, and Pal. N. Y., vol. 1, p. 115, Trenton Gr.

tenuistriata, Sowerby, 1839, (*Leptæna tenuistriata*.) Murch. Sil. Syst., p. 636, and Pal. N. Y., vol. 1, p. 108, Low. Sil.

textilis, see *Strophodonta junia*.

thalia, see *Streptorhynchus thalia*.

transversalis, see *Leptæna transversalis*.

trilobata, Owen, 1852, (*Leptæna trilobata*.) Geo. Sur. Wis., Iowa, and Minn., p. 584, Trenton Gr.

tullia, Billings, 1874, Pal. Foss., vol. 2, p. 29, Low. Devonian.

undulata, syn. for *S. rhomboidalis*.

undulosa, Conrad, 1841, Ann. Rep. N. Y., p. 54, Low. Held. Gr.

unicostata, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 335, Hud. Riv. Gr.

varistriata, see *Strophodonta varistriata*.

ventricosa, Shaler, (*Brachyprion ventricosum*.) The name was preoccupied.

wisconsinensis, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 61, and Geo. Wis., vol. 4, p. 263, Hud. Riv. Gr.

woolworthana, see *Streptorhynchus woolworthanum*.

STROPHONELLA, Hall, 1879, 28th Rep. N. Y., Mus. Nat. Hist., p. 153. [Ety. diminutive of *strophos*, twisted.] Distinguished from *Strophodonta* by the resupinate character, the strong and more restricted muscular impression of the ventral valve and strong median septum of the dorsal valve; and from *Streptorhynchus* by the cardinal process, the crenulations on the inner margins of the cardinal area, and the solid area, with sometimes a triangular deltidium.

Type *S. semifasciata*.

ampla, Hall, 1857, (*Strophomena ampla*.) 10th Rep. N. Y., Mus. Nat. Hist., p. 111, and Pal. N. Y., vol. 4, p. 93, Up. Held. Gr.

cælata, Hall, 1867, (*Strophodonta cælata*.) Pal. N. Y., vol. 4, p. 112, Chemung Gr.

cavumbona, Hall, 1857, (*Strophodonta cavumbona*.) 10th Rep. N. Y. Mus. Nat. Hist., p. 51, and Pal. N. Y., vol. 3, p. 187, Low. Held. Gr.

leavenworthana, Hall, 1857, (*Strophodonta leavenworthana*.) 10th Rep. N. Y. Mus. Nat. Hist., p. 53, and Pal. N. Y., vol. 3, p. 189, Low. Held. Gr.

punctulifera, Conrad, 1838, (*Leptæna punctulifera*.) Ann. Rep. N. Y., p. 117, and Pal. N. Y., vol. 3, p. 188, Low. Held. Gr.

reversa, Hall, 1858, (*Strophodonta reversa*.) Geo. Rep. Iowa, p. 494, Ham. Gr.

semifasciata, Hall, 1863, (*Strophodonta semifasciata*.) Trans. Alb. Inst., vol. 4, p. 210, Niagara Gr.

SYNTRIELASMA, Meek & Worthen, 1865, Proc. Acad. Nat. Sci., p. 277. [Ety. *syn*, together; *treis*, three; *elasma*, plate.]

Shell thin, gibbous; valves articulated by teeth and sockets; hinge-line short; area higher in the ventral valve than

FIG. 624.—*Strophonella punctulifera*.

reversa, Hall, 1858, (*Strophodonta reversa*.) Geo. Rep. Iowa, p. 494, Ham. Gr.

semifasciata, Hall, 1863, (*Strophodonta semifasciata*.) Trans. Alb. Inst., vol. 4, p. 210, Niagara Gr.

SYNTRIELASMA, Meek & Worthen, 1865, Proc. Acad. Nat. Sci., p. 277. [Ety. *syn*, together; *treis*, three; *elasma*, plate.]

Shell thin, gibbous; valves articulated by teeth and sockets; hinge-line short; area higher in the ventral valve than

in the dorsal; beaks incurved, subequal; surface radiated, forming interlocking angular projections at their terminations; shell structure punctate. Type *S. hemiplicatum*.

FIG. 625.—*Syntrielasma hemiplicatum*. *l*, Dental lamina; *m*, mesial septum.

hemiplicatum, Hall, 1852, (*Spirifera hemiplicata*,) Stan.'s Ex. to Great Salt Lake, p. 409, Coal Meas.

SYRINGOTHYRIS, Winchell, 1863, Proc. Acad. Nat. Sci., p. 6. [Ety. *syrix*, a tube; *thyris*, a window.] General aspect like *Spirifera*; shell substance fibrous and impunctate; beak extremely elevated; area of ventral valve large, with a narrow triangular fissure closed toward the apex by an external, convex pseudo-deltidium, beneath which, and diverging from it, is another transverse plate, connecting the vertical dental lamellæ, arched above, and beneath giving off a couple of median parallel lamellæ, which are incurved so as to nearly join their inferior edges, thus forming a slit-bearing tube, which projects into the interior of the shell. Type *S. typus*.

halli, Winchell, 1863, Proc. Acad. Nat. Sci., p. 8, Marshall Gr.

typus, Winchell, 1863, Proc. Acad. Nat. Sci. Phil., p. 7, Marshall Gr.

FIG. 626.—*Terebratula maxillata*.

TEREBRATULA, Lihwyd, 1699, Lith. Brit. Ich. [Ety. diminutive of *terebratus*, perforated.] Shell variable in form, oval, elongated or transverse; smooth or plicated; valves unequally convex; hinge-line curved; beak

short, truncated by a foramen; deltidium in one or two pieces; loop short; confined to the posterior portion of the shell, not more than one-third the length of the valve, simply attached to a hinge plate; two ribbon-shaped lamellæ are united by a transverse lamella bent upward in the middle; the cirrated arms are supported by the crura, and project in front of the loop; no median septum in the socket valve. Type *T. vitrea* and *T. maxillata*.

acuminatissima, Castelnau, 1843, Syst. Sil., p. 40. Not recognized.

affinis, syn. for *Atrypa reticularis*.

aprinis, see *Rhynchonella aprinis*.

arcuata, Swallow, 1863, Trans. St. Louis Acad. Sci. The name was preoccupied by Roemer in 1840. See *T. shumardana*.

argentea, see *Athyris argentea*.

aspera, see *Atrypa aspera*.

bidentata, see *Rhynchonella bidentata*.

bisacula, McChesney, 1860, New Pal. Foss., p. 82, Kaskaskia Gr. Not recognized.

borealis, Castelnau, 1843, Syst. Sil., p. 40. Not recognized.

bovidens, Morton, 1836, Am. Jour. Sci., vol. 20, p. 150, Coal Meas.

brevirostris, see *Rhynchonella brevirostris*.

brevilobata, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 84, Kaskaskia Gr.

burlingtonensis, White, 1860, Bost. Jour. Nat. Hist., p. 228, and Geo. Sur. W. 100th Mer., vol. 4, p. 93, Kinderhook Gr. *concentrica*, syn. for *Athyris spiriferoides*. *cooperensis*, n. sp., Keokuk Gr. Proposed instead of *T. parva*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 83, which name was preoccupied.

FIG. 627.—*Terebratula bovidens*.

crenulata, Sowerby, 1840, (*Atrypa crenulata*,) Geo. Trans., 2d series, vol. 5, p. 704, Devonian.

cuneata, see *Rhynchonella cuneata*.

elia, Hall, 1867, Pal. N. Y., vol. 4, p. 390, Up. Held Gr.

formosa, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 6, and Bull. Am. Mus. Nat. Hist., p. 56, Warsaw Gr.

geniculosa, syn. for *Terebratula bovidens*.

gracilis, Swallow, 1863, Trans. St. Louis Acad. Sci. The name was preoccupied by Von Buch in 1834. See *T. swallowana*.

harmonia, Hall, 1867, Pal. N. Y., vol. 4, p. 388, Up. Held Gr.

inornata, McChesney, 1860, New Pal. Foss. Carb.

insperata, Phillips, 1841, Pal. Foss., Devonian.

interplicata, see *Anastrophia interplicata*.

jucunda, Hall, 1867, Pal. N. Y., vol. 4, p. 390, Up. Held Gr.

lacunosa. Not American.

lapillus, Morton, 1836, Am. Jour. Sci. and Arts, vol. 29, p. 149, Coal Meas.

lens, see *Cryptonella lens*.

laticosta, see *Atrypa laticosta*.

lincklaeni, see *Cryptonella lincklaeni*.

lynx, see *Orthis lynx*.

marcyi, see *Retzia marcyi*.

43, Syst. Sil.,

laris.
rinia.na. St. Louis
preoccupied
Shumardana.
a.

identata.

w Pal. Foss.,
recognized.

t. Sil., p. 40,

n. Jour. Sci.,

brevirostris.

Trans. St.
84, Kaskas-

, Bost. Jour.

eo. Sur. W.

nderhook Gr.

spiriferoides.

r. Proposed

allow, 1863,

vol. 2, p. 83,

ed.

idens.

trypa crenu-

vol. 5, p.

neata.

ol. 4, p. 390,

b. Inst., vol.

i. Nat. Hist.,

la bovidens.

as. St. Louis

preoccupied

See T. swal-

Y., vol. 4, p.

, New. Pal.

l. Foss., De-

interplicata.

Y., vol. 4, p.

our. Sci. and

Meas.

cklaeni.

marginalis, see *Atrypa marginalis*.*micbelini*, see *Orthis micbelini*.*millepunctata*, syn. for *T. bovidens*.*mormoni*, see *Retzia mormoni*.*navicella*, Hall, 1867, Pal. N. Y., vol. 4, p.

391, Ham. Gr.

nuciformis, Morton, 1836, Am. Jour. Sci.

and Arts, vol. 29, p. 149, Coal Meas.

nucula, see *Rhynchonella nucula*.*ontario*, Hall, 1867, Pal. N. Y., vol. 4, p.

418, Ham. Gr.

ovoides, see *Rensselaeria ovoides*.*parva*, Swallow, 1863, Trans. St. Louis

Acad. Sci., p. 83. The name was pre-

occupied by d'Archiac in 1846. See

Terebratulina cooperensis.*pennata*, see *Spirifera pennata*.*perinflata*, Shumard, 1859, Trans. St. Louis

Acad. Sci., vol. 1, p. 392, Permian Gr.

planirostra, see *Cryptonella planirostra*.*rectirostra*, see *Cryptonella rectirostra*.*reticularis*, see *Atrypa reticularis*.*rockymontana*, see *Rhynchonella rocky-**montana*.*roemingeri*, Hall, 1863, 16th Rep. N. Y.

Mus. Nat. Hist., p. 48, and Pal. N. Y.,

vol. 4, p. 389, Ham. Gr.

rowleyi, Worthen, 1884, Bull. No. 2, Ill.

St. Mus. Nat. Hist., p. 23, Burling-

ton Gr.

sacculus, Martin, 1809, Petrif. Derb., Low.

Carb.

schlotheimi, see *Camerophoria schlotheimi*.*shumardana*, S. A. Miller, 1883, 2d Ed.

Am. Pal. Foss., p. 299, Kaskaskia Gr.

Proposed instead of *T. arcuata*, Swallow,

1863, Trans. St. Louis Acad. Sci., vol. 2,

p. 83, which was preoccupied.

simulator, Hall, 1867, Pal. N. Y., vol. 4, p.

391, Ham. Gr.

spiriferoides, see *Athyris spiriferoides*.*stricklandi*, see *Rhynchonella stricklandi*.*subretziaforma*, McChesney, 1860, Pal.

Foss., p. 82, Kaskaskia Gr. Not recog-

nized.

subtilita, see *Athyris subtilita*.*sullivanti*, Hall, 1867, Pal. N. Y., vol. 4,

p. 387, Up. Held. Gr.

swallovana, S. A. Miller, 1883, 2d Ed. Am.

Pal. Foss., p. 299, Kaskaskia Gr. Pro-

posed instead of *T. gracilis*, Swallow,

1863, Trans. St. Louis Acad. Sci., vol. 2,

p. 83, which was preoccupied.

traversensis, Winchell, 1866, Rep. Low.

Penin. Mich., p. 95, Ham. Gr.

trinuclea, see *Athyris trinuclea*.*turgida*, Hall, 1858, Trans. Alb. Inst., vol.

4, p. 6, and Bull. Am. Mus. Nat. Hist.,

p. 54, Warsaw Gr.

uta, see *Rhynchonella uta*.*utah*, Hall & Whitfield, 1877, U. S. Geo.

Expl. 40th parallel, vol. 4, p. 258, Wa-

verly Gr.

valenciennae, Castelnau, 1843, Syst. Sil., p.

39. Not recognized.

wilsoni, see *Rhynchonella wilsoni*.*Terebratulites*, Schlotheim, syn. for *Spir-**ifera*.*biforatus*, see *Orthis biforata*.*Trematis*, Sharpe, 1848, Quar. Jour. Geo.Soc., vol. 13, p. 66. [Ety. *trema*, an

opening.] Shell suborbicular or trans-

versely oval, lenticular; valves un-

equally convex; umbo of the upper or

dorsal valve submarginal, slightly pro-

jecting; lower or ventral valve with a

subcentral umbo, behind which a nar-

row, oblong, oval slit reaches to near

the posterior margin, and afforded pas-

sage to the pedicle fibers of attachment;

shell punctate in the outer layers, and

fibrous and of a pearly luster within.

Type *T. terminalis*.*cyclata*, see *Obolella cyclata*.*cancellata*, Sowerby, 1825, (Orbicula can-

cellata.) Zool. Jour., vol. 2, Trenton Gr.

crassa, see *Obolella crassa*.*crassipuncta*, Ulrich, 1889, Am. Geo., vol.

4, p. 22, Hud. Riv. Gr.

dyeri, S. A. Miller, 1874,

Cin. Quar. Jour. Sci., vol.

1, p. 347, Hud. Riv. Gr.

filosa, see *Schizocrania filosa*.*fragilis*, Ulrich, 1889, Am.

Geo., vol. 4, p. 21, Utica

Slate.

huronensis, Billings, 1862, Fig. 628.—Tre-Pal. Foss., vol. 1, p. 53, matls *dyeri*.

Black Riv. Gr.

montrealensis, Billings, 1862, Pal. Foss.,

vol. 1, p. 52, Trenton Gr.

millepunctata, Hall, 1866,

24th Rep. N. Y. Mus.

Nat. Hist., p. 221,

Hud. Riv. Gr.

oblata, Ulrich, 1889, Am.

Geo., vol. 4, p. 23,

Hud. Riv. Gr.

ottawensis, Billings,

1862, Pal. Foss., vol. 1,

p. 53, Trenton Gr.

pannulus, see *Kutorgina pannulus*.*puncto-riata*, Hall & Whitfield, 1873, 23d

Rep. N. Y. Mus. Nat. Hist., p. 243,

Trenton and Hud. Riv. Gr.

(?) *pustulosa*, Hall, 1866, 24th Rep. N. Y.

Mus. Nat. Hist., p. 222, Hud. Riv. Gr.

quincuncialis, Miller & Dyer, 1878, Cont.

to Pal. No. 2, p. 8, Hud. Riv. Gr.

rudis, Hall & Whitfield, 1873, 23d Rep.

N. Y. Mus. Nat. Hist., p. 243, Tren-

ton Gr.

terminalis, Emmons, 1842, (Orbicula ter-

minalis.) Geo. Rep. N. Y., p. 395, and

Pal. N. Y., vol. 1, p. 100, Trenton Gr.

umbonata, Ulrich, 1889, Am. Geo., vol. 4,

p. 23, Hud. Riv. Gr.

Trematospira, Hall, 1859, 12th Rep. N. Y.Mus. Nat. Hist., p. 27. [Ety. *trema*,foramen; *spira*, a spire; in allusion

to the perforation in the beak of the

ventral valve.] Transverse, elliptical

or subrhomboidal, inequivalve; mesial

fold and sinus; surface plicated; internal

spires, as in *Spirifera*; hinge-line

short; cardinal angles rounded; valves

articulated by teeth and sockets; beak

of ventral valve produced or incurved,

FIG. 629.—Trematis millepunctata.

and truncated by a small, round perforation, separated from the hinge-line by a deltidium; deep, triangular pit beneath the beak of the ventral valve, which is filled by the closely incurved beak of the dorsal valve; structure punctate. Type *T. costata* and *T. multistriata*.

FIG. 630.—*Trematospira acadiæ*.

- acadiæ*, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 144, Up. Sil.
camura, Hall, 1852, (*Atrypa camura*), Pal. N. Y., vol. 2, p. 273, Low. Held. Gr.
costata, Hall, 1859, Pal. N. Y., vol. 3, p. 210, Low. Held. Gr.
deweyi, see *Rhynchospira deweyi*.
formosa, see *Rhynchospira formosa*.
gibbosa, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 82, and Pal. N. Y., vol. 4, p. 272, Ham. Gr.
globosa, Hall, 1857, 10th Rep. N. Y. Mus. Nat. Hist., p. 87, and Pal. N. Y., vol. 3, p. 215, (Waldheimiaglobosa), Low. Held. Gr.
granulifera, Meek, 1872, Proc. Acad. Nat. Sci. Phil., p. 318, and Ohio Pal., vol. 1, p. 128, Hud. Riv. Gr.
hirsuta, Hall, 1857, (*Atrypa hirsuta*), 10th Rep. N. Y. Mus. Nat. Hist., p. 168, and Pal. N. Y., vol. 4, p. 274, Up. Held. and Ham. Gr.
imbricata, Hall, 1857, (*Leptocoelia imbricata*), 10th Rep. N. Y. Mus. Nat. Hist., p. 108, and Pal. N. Y., vol. 3, p. 246, Low. Held. Gr.
infrequens, Walcott, 1885, Mongr. U. S. Geo. Sur., vol. 8, p. 151, Lower Devonian.
mathewsoni, McChesney, 1861, New Pal. Foss., p. 71, Niagara Gr.
multistriata, Hall, 1857, (*Spirifer multistriatus*), 10th Rep. N. Y. Mus. Nat. Hist., p. 59, Low. Held. Gr.
liniuscula, Winchell, 1866, Rep. Low. Peninsula Mich., p. 94, Ham. Gr.
 (?) *nobilis*, Hall, 1860, (*Rhynchospira nobilis*), 13th Rep. N. Y. Mus. Nat. Hist., p. 83, Ham. Gr.
perforata, Hall, 1857, (*Spirifer perforata*), 10th Rep. N. Y. Mus. Nat. Hist., p. 60, Low. Held. Gr.
 (?) *quadruplicata*, see *Rhynchotretra quadruplicata*.
rectirostris, Hall, 1856, (*Waldheimia rectirostra*), 10th Rep. N. Y. Mus. Nat. Hist., p. 49, and Pal. N. Y., vol. 3, p. 217, Low. Held. Gr.
simplex, Hall, 1856, Pal. N. Y., vol. 3, p. 211, Low. Held. Gr.
Trigonotretra, König, 1825, Icon. Foss. Sect., p. 3. [Ety. *trigonos*, a triangle; *tretos*, perforated.] Syn. for *Spirifera*. Meek, concluding that *S. cuspidatus* mentioned by Sowerby in Minn. Conch., vol. 2, p. 42, should be considered as the type of *Spirifera*, proposed to retain *Trigonotretra* for shells of the type of *S. striata*, contrary to the views of most authors. See Pal. Up. Mo., p. 18.

TRIMERELLA Billings, 1862, Pal. Foss., vol. 1, p. 153. [Ety. *treis*, three; *meros*, part; *ella*, diminutive.] Large subovate or subcircular; valves convex; beaks solid and transversely striated;

FIG. 631.—*Trimerella grandis*. Cast of dorsal valve.

shell thick, and surface concentrically striated; area of pedicle valve longer than wide; deltidium large; hinge thick, elevated, rudely or slightly dentary; cardinal facet large; crescent well defined; platforms elevated and doubly vaulted, occasionally solid and slightly raised; median plate in both valves, longest in the brachial one. Type *T. grandis*.

FIG. 632.—*Trimerella grandis*. Cast of ventral valve.

acuminata, Billings, 1862, Pal. Foss., vol. 1, p. 167, Guelph Gr.
billingsi, Dall, 1871, Am. Jour. Conch., vol. 7, p. 82, Guelph Gr.

Foss., vol.
ree; *peros*,
large subo-
es convex;
ly striated;

st of dorsal

ncentrically
valve longer
rge; hinge
lightly den-
e; crescent
levated and
ly solid and
ate in both
achial one.

t of ventral

. Foss., vol.
ur. Conch.,

dalli, Davidson & King, 1872, Brighton Meeting Brit. Assoc. and Quar. Jour. Geo. Soc., 1874, p. 154, Guelph Gr.

FIG. 633.—*Trimerella galtensis*.

galtensis, Billings, 1862, (Obolus galtensis,) Pal. Foss., vol. 1, p. 168, Guelph Gr. grandis, Billings, 1862, Pal. Foss., vol. 1, p. 166, Guelph Gr. minor, Dall, syn. for T. galtensis.

ohioensis, Meek, 1871, Am. Jour. Sci., 2d series, vol. 1, p. 315, and Ohio Pal., vol. 1, p. 183, Niagara Gr.

TRIPLESIA, Hall, 1858, 12th Rep. N. Y.

FIG. 634.—*Triplesia extans*.

Mus. Nat. Hist., p. 44. [Ety. *triplosos*, thrice; in allusion to the trilobate character of the shell.] Shell transverse or elongate, trilobate or subtrilobate; ventral valve deeply sinuous and dorsal, having a corresponding fold; hinge-line straight; area small; foramen triangular; surface concentrically striated; ventral valve with a strong tooth on each side of the fissure; muscular impressions small; dorsal valve with a prominent bifurcating cardinal process, on each side of which there is a brachial process directed obliquely inward and laterally. Type T. extans.

congesta, Conrad, 1842, (*Atrypa congesta*), Jour. Acad. Nat. Sci., vol. 8, p. 265, and Pal. N. Y., vol. 2, p. 67, Clinton Gr. cuspidata, Hall, 1847, (*Atrypa cuspidata*), Pal. N. Y., vol. 1, p. 138, Trenton Gr.

FIG. 635.—*Triplesia ortonii*.

extans, Emmons, 1842, (*Atrypa extans*), Geo. Rep. 2d Dist. N. Y., p. 395, and Pal. N. Y., vol. 1, p. 137, Trenton Gr. lateralis, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 303, Birdseye Gr.

nucleus, Hall, 1847, (*Atrypa nucleus*), Pal. N. Y., vol. 1, p. 138, Trenton Gr.

ortonii, Meek, 1872, (*Dicraniscus ortonii*), Am. Jour. Sci. and Arts, 3d ser., vol. 4, p. 280, and Ohio Pal., vol. 1, p. 178, Niagara Gr.

primordialis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 51, and Geo. Wis., vol. 4, p. 172, Potsdam Gr.

putillus, Hall, syn. for T. waldronensis.

radiata, Whitfield, 1889, Bull. Am. Mus.

Nat. Hist., vol. 2, p. 43, Calciferous Gr.

waldronensis, Miller & Dyer, 1878, (Spirifera ? waldronensis,) Jour. Cin. Soc.

Nat. Hist., vol. 1, p. 37, Niagara Gr.

TROPIDOLEPTUS, Hall, 1857, proposed in 10th Rep., but described in 1859 in 12th Rep. N. Y. Mus. Nat. Hist., p. 31. [Ety. *tropis*, the keel or bottom of a ship; *leptos*, slender.] General form of Strophomena; surface plicated; structure punctate; ventral valve convex, with an area and wide fissure beneath the beak; dental lamellae distinct from the margin of the fissure, crenulate; dorsal valve concave, with crenulate dental fossets; a strong, cardinal process, with diverging lobes in the interior, which support slender crura that converge to and unite with the median crest. Type T. carinatus.

FIG. 636.—*Tropidoleptus carinatus*.

carinatus, Conrad, 1839, (*Strophomena carinata*), Ann. Geo. Rep. N. Y., p. 64, and Pal. N. Y., vol. 4, p. 407, Ham. Gr. occidentis, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 91, and Pal. N. Y., vol. 4, p. 408, Ham. Gr.

VITULINA, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 72. [Ety. mythological name.] Externally it is like Leptocoelia, but distinguished by its strong dental lamellae and processes. Type V. pustulosa.

pustulosa, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 72, and Pal. N. Y., vol. 4, p. 410, Tully limestone.

WALDHEIMIA, King, 1849, Monograph of Permian Fossils, p. 145. [Ety. proper name.] Shell circular, subquadrate, transverse or

elongated; valves unequally convex, smooth, or plicated; beak truncated, perforated; deltidium in one or two pieces; loop long, formed of lamellae attached by the crura to the hinge

plate; one tooth on each side of the deltidium, supported by plates, and fitting in the sockets of the dorsal valve; structure punctate. Type *W. australis*.

FIG. 637.—*Waidhelmia australis*. Dorsal valve: *f*, cardinal process; *t*, dental sockets; *p*, hinge plate; *s*, septum; *e*, crura of the loop; *l*, reflected portion of the loop; *m*, quadruple adductor impression. Ventral valve: *f*, foramen; *d*, deltidium; *t*, teeth; *a*, single adductor impression; *r*, cardinal muscle; *z*, accessory muscles; *p*, pedicle muscles; *v*, position of the vent; *z*, attachment of pedicle sheath.

compacta, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

deweyi, see *Retzia deweyi*.

formosa, see *Retzia formosa*.

globosa, see *Trematospira globosa*.

rectirostra, see *Trematospira rectirostra*.

FIG. 638.—*Waidhelmia australis*.

WHITFIELDIA, Davidson, 1881, Lond. Geo. Mag., vol. 8, p. 289. [Ety. proper name.] While *Meristella* is distinguished from *Merista* by having

no shoe-lifter process; *Whitfieldia* is distinguished from both by the absence of those peculiar ring-shaped processes attached to the loop, and has instead only a short, bifurcating process, where in *Merista* and *Meristella* the rings are formed. These internal differences in the spirals seem to distinguish the genera. Type *W. tumida*.

maria, Hall, 1863, (*Meristella maria*.) Trans. Alb. Inst., vol. 4, p. 212, Niagara Gr.

FIG. 639.—*Whitfieldia maria*. Internal casts.

ZYGOSPIRA, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 126. [Ety. *zygos*, yoke; *spira*, spire.] General form of *Atrypa* with internal spires having a broad loop passing from the outer limbs of the spiral band entirely across from side to side, near to or above the center, and close to the inner side of the dorsal valve. Type *Z. modesta*.

concentrica, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 14, Hud. Riv. Gr.

headi, Billings, 1862, (*Athyris headi*.) Pal. Foss., vol. 1, p. 147, Hud. Riv. Gr.

headi var. *anticostiensis*, Billings, 1862, (*Athyris headi* var. *anticostiensis*.) Pal. Foss., vol. 1, p. 147, Hud. Riv. Gr.

headi var. *borealis*, Billings, 1862, (*Athyris headi* var. *borealis*.) Pal. Foss., vol. 1, p. 147, Hud. Riv. Gr.

headi, Meek, see *Glossia headi*.

minima, Hall, 1879, Desc. New Spec. Foss., p. 14, and 11th Rep. Geo. and Nat. Hist. Ind., p. 305, Niagara Gr.

modesta, Say, 1847, (*Atrypa modesta*.) Pal. N. Y., vol. 1, p. 141, Trenton and Hud. Riv. Gr.

modesta var. *cincinnatiensis*, Meek, 1872, Pal. Ohio, vol. 1, p. 126, Hud. Riv. Gr.

pauper, Billings, 1866, Catal. Sil. Foss. Antic., p. 46, Anticosti Gr.

subconca, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 380, Low. Held. Gr.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

tumida, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 119, Up. Coal Meas.

 

FIG. 640.—*Zygospira modesta*.

maria,) Trans.
Niagara Gr.

terrestrial casts.

p. N. Y. Mus.
zygos, yoke;
n of Atrypa
ing a broad
ster limbs of
across from
ve the center,
o of the dorsal

ur. Cin. Soc.
ud Riv. Gr.
n heads,) Pal.
Riv. Gr.
Billings, 1862,
stiensis,) Pal.
Riv. Gr.
1862, (Athyris
Foss., vol. 1,

adi.
New Spec.
ep. Geo. and
Niagara Gr.
pa modesta,
Trenton and

FIG. 640.—Zygospira modesta.

vol. 3, p. 380,

CLASS PTEROPODA.

THE Class Pteropoda consists wholly of marine animals of small size, furnished with a pair of fins at the sides of the head, by means of which they swim in the open sea. The living forms are divided into two orders: the Gymnosomata and Thecosomata. The Gymnosomata have no shells, and occur in such prodigious numbers that they furnish food for whales and many sea-birds. The Thecosomata have either straight or coiled shells, some of which are glassy in their texture and very beautiful. It may well be doubted whether or not any of the Palaeozoic fossils belong to this order. They are referred to the following families:

FAMILY ASPIDELLIDÆ.—Aspidella.

FAMILY CLATHROCELIIDÆ.—Clathrocellia.

FAMILY CONULARIDÆ.—Conularia.

FAMILY HYOLITHIDÆ.—Coleolus, Coleoprion, Diplothea, Hyolithellus, Hyolithes, Pharetrella, Stenotheca.

FAMILY MATTHEVIDÆ.—Matthevia.

FAMILY PTEROTHECIDÆ.—Pterotheca,

FAMILY SCENELLIDÆ.—Scenella.

FAMILY TENTACULITIDÆ.—Styliola, Tentaculites.

ASPIDELLA, Billings, 1872, Am. Jour. Sci., 3d ser., vol. 3, and Pal. Foss., vol. 2, p. 76. [Ety. *aspidella*, little shield.] Small, ovate, bordered by a narrow ring within which it is concave; in the middle there is a ridge, from which grooves radiate to the border. Type A. terranovica.

FIG. 641.—Aspidella terranovica.

terravovica, Billings, 1872, Am. Jour. Sci., 3d ser., vol. 3, and Pal. Foss., vol. 2, p. 77, Taconic.

Camerotheca, Matthew, 1885, Can. Rec. Sci., vol. 1, p. 149, syn. for Hyolithes gracilis, see Hyolithes gracilis.

CLATHROCELLIA, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 203. [Ety. *clathro*, latticed; *cellia*, b-ly.] An oblique conical tube, expanding more rapidly on one side than the other; interior crossed by unsymmetrical, arching, septal lines and longitudinal ones, which give it a cancellated aspect; shell thin, translucent, lamellose. Type C. eborica.

eborica, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 204, Ham. Gr.

Cloderna, Hall, syn. for Pterotheca attenuata, see Pterotheca attenuata. expansa, see Pterotheca expansa.

COLEOLUS, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 184. [Ety. *coleos*, sheath.] Tubuliform, elongate-conical, straight or slightly curved, annulated, sometimes obliquely, sometimes longitudinally striated, interior smooth. Type C. tenuicinctus.

FIG. 642.—Coleolus acicula.

acicula, Hall, 1843, (Orthoceras acicula,) Geo. Sur. 4th Dist. N. Y., p. 243, and Pal. N. Y., vol. 5, pt. 2, p. 187, Genesee Slate.

aciculatus, Hall, 1860, (Dentalium aciculatum,) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 107, Marcellus Shale and Portage Grs.

crenatocinctus, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 188, Up. Held. Gr.

gracilis, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 190, Chemung Gr.

herzeri, Hall, 1888, Pal. N. Y., vol. 7, p. 7, Waverly Gr.

lævis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 199, Devonian.

mohri, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 189, Up. Held. Gr.

spinulus, Hall, 1879, Desc. New Spec. Foss., p. 18, and 11th Rep. Geo. and Nat. Hist. Ind., p. 322, Niagara Gr.

- tenuicinctus, Hall, 1876, (Coleoprion tenuicinctum,) Illust. Devon. Foss., pl. 27, Ham. Gr.
- COLEOPRION, Sandberger, 1847, Leonhardt & Bronn, Jahrbuch, vol. 1, p. 25. [Ety. *koloos*, sheath; *prion*, saw.] Tubuliform, appearing as an elongate cone, encircled by oblique annulations, which are interrupted along a longitudinal line, and attenuate at their extremities; internal walls smooth. Type *C. gracilis*.
- minutum, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 85, Trenton Gr.
- tenue, Hall, 1879, Pal. of N. Y., vol. 5, pt. 2, p. 184, Ham. Gr.
- tenuicinctum*, see *Coleolus tenuicinctus*.
- CONULARIA, Miller, 1821, in Sowerby's Minn. Conch., vol. 3, p. 107. [Ety. *conulus*, little cone.] Elongate pyramidal; transverse section varying from quadrangular to octagonal; angles indented by longitudinal grooves; septum near the apex; surface reticulated and ornamented. Type *C. quadrisulcata*.
- asperata, Billings, 1866, Catal. Sil. Foss. Antic., p. 21, Hud. Riv. Gr.
- bifurca, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 18. Not properly defined.
- byblis, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 22, Waverly Gr.
- cayuga, Hall, 1876, Illust. Devonian Foss., pl. 28, and Pal. N. Y., vol. 5, pt. 2, p. 211, Ham. Gr.
- chesterensis, Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 325, Kaskaskia Gr.
- congregata, Hall, 1876, Illust. Devonian Foss., pl. 28, and Pal. N. Y., vol. 5, pt. 2, p. 214, Portage Gr.
- continens, Hall, 1876, Illust. Devonian Foss., pl. 28, and Pal. N. Y., vol. 5, pt. 2, p. 212, Marcellus Shale.
- continens var. rudis, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, pl. 28, Ham. Gr.
- crawfordsvillensis, Owen, 1862, Geo. Sur. Ind., p. 362, Keokuk Gr.
- crebristriata, Hall, 1876, Illust. Devonian Foss., pl. 29, and Pal. N. Y., vol. 5, pt. 2, p. 210, Ham. Gr.
- crustula, White, 1880, 12th Rep. U. S. Geo. Sur. Terr., p. 170, Coal Meas.
- elegantula, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 85, and Ohio Pal., vol. 1, p. 288, Up. Held. Gr.
- formosa, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 38, Hud. Riv. Gr.
- gattingeri, Safford, 1869, Geo. of Tenn., p. 289, Trenton Gr.
- gracilis, Hall, 1847, Pal. N. Y., vol. 1, p. 224, Trenton Gr.
- granulata, Hall, 1847, Pal. N. Y., vol. 1, p. 223, Trenton Gr.
- hudsoni, Emmons, 1856, Am. Geo., vol. 1, p. 208, Hud. Riv. Gr.
- huntana, Hall, 1859, Pal. N. Y., vol. 3, p. 348, Low. Held. Gr.
- indentata, Conrad, 1854, Proc. Acad. Nat. Sci., vol. 7, p. 31, Trenton Gr.
- infrequens, Hall, 1879, Desc. New Spec. Foss., p. 17, and 11th Rep. Geo. and Nat. Hist. Ind., p. 321, Niagara Gr.
- laqueata, Conrad, 1841, Ann. Rep. N. Y., p. 57, Niagara Gr.
- lata, Hall, 1859, Pal. N. Y., vol. 3, p. 479, Oriskany sandstone.
- longa, Hall, 1852, Pal. N. Y., vol. 2, p. 295, Niagara Gr.
- magnifica, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 58, Niagara Gr.
- marionensis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 656, Ham. Gr.
- miconema, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 84, and Ohio Pal., vol. 2, p. 316, Waverly Gr.
- missouriensis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 657, and Geo. Sur. Ill., vol. 5, p. 541, St. Louis Gr.
- molaris, White, 1876, Proc. Acad. Nat. Sci., p. 33, Devonian.
- multicostata, Meek & Worthen, 1865, Proc. Fro. 343.—*Conularia microneuma*.
- Waverly Gr.
- multipuncta, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 18. Not properly defined.
- newberryi, Winchell, 1865, Proc. Acad. Nat. Sci., p. 130, Waverly Gr.
- niagarensis, Hall, 1852, Pal. N. Y., vol. 2, p. 294, Niagara Gr.
- osagensis, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 98, Kaskaskia Gr.
- papillata, Hall, 1847, Pal. N. Y., vol. 1, p. 223, Trenton Gr.
- planocostata, Dawson, 1868, Acad. Geol., p. 307, Carb.
- pyramidalis, Hall, 1859, Pal. N. Y., vol. 3, p. 347, Low. Held. Gr.
- quadrata, Walcott, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 93, Trenton Gr.
- quadrisulcata, (?) Miller, 1821, Min. Conch., vol. 3, p. 107, Niagara Gr.
- rugosa, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 59, Niagara Gr.
- splendida, Billings, 1866, Catal. Sil. Foss. Antic., p. 21, Hud. Riv. Gr.
- subcarbonaria, Meek & Worthen, 1865, Proc. Acad. Nat. Sci., p. 253, and Geo. Sur. Ill., vol. 5, p. 520, Keokuk Gr.
- subulata, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 32, and Bull. Arc. Mus. Nat. Hist., p. 91, Warsaw Gr.
- transversa, Ringueberg, 1883, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 19, Niagara Gr.

FIG. 644.—*Conularia quadrisulcata*.

New Spec.
p. Geo. and
gara Gr.
Rep. N. Y.,

ol. 3, p. 479,

vol. 2, p.

Bull. No. 1,
58, Niag-

Trans. St.
56, Ham. Gr.

Fig. 643.—Conular
microneurite.

36, Bull. Buf.
3. Not prop-

chell, 1865,
Sci., p. 130,

, 1852, Pal.
p. 294, Niag-

y, 1863, Trans.
Sci., vol. 2,

ia Gr.
1847, Pal. N. Y.,

Trenton Gr.
awson, 1868,

307, Carb.
11, 1859, Pal.

p. 347, Low.

1876, 28th
Hist., p. 93,

1821, Min.
gara Gr.

No. 1, Mus.
gara Gr.

atal. Sil. Foss.
Gr.

Vorthen, 1865,
253, and Geo.

Keokuk Gr.
ns. Alb. Inst.,

Am. Mus. Nat.

trentonensis, Hall, 1847, Pal. N. Y., vol. 1, p. 222, Trenton and Hud. Riv. Grs. triplicata, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 657, Ham. Gr. undulata, Conrad, 1841, Ann. Rep. N. Y., p. 57, and Pal. N. Y., vol. 5, pt. 2, p. 208, Ham. Gr. verneuilana, Emmons, 1846, Am. Quar. Jour. Agr. and Sci., vol. 4, p. 330. Subcarboniferous. victa, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 22, Burlington Gr. whitii, Meek & Worthen, 1865, Proc. Acad. Nat. Sci., p. 253, Waverly Gr. wilkinsi, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 59, Niagara Gr.

DIPLOTHECA, Matthew, 1885, Am. Jour. Sci. and Arts, 3d ser., vol. 30, p. 293. [Ety. *diploos*, double; *Theca*, a genus.] Slender, conical, section triangular; internal septa dividing it in segments; body cavity separated from one side by a thin partition, supported by delicate transverse septa; distinguished from Hyolithes by more rapidly expanding, and by a firmer, rounder side, where it has the support of the lateral septa. Type D. acadica.

acadica, Matthew, 1885, Am. Jour. Sci. and Arts, 3d ser., vol. 30, p. 294, St. John Gr.

hyattana, Matthew, 1885, Am. Jour. Sci. and Arts, 3d ser., vol. 30, p. 294, St. John Gr.

hyattana var. caudata, Matthew, 1885, Am. Jour. Sci. and Arts, 3d ser., vol. 30, p. 294, St. John Gr.

HYOLITHELLUS, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 240, and Am. Jour. Sci. and Arts, 3d ser., vol. 3, p. 360. [Ety. diminutive of Hyolithes.] Distinguished from Hyolithes by its long, slender form and structure of the operculum. Type H. micans.

FIG. 645.—Hyolithes micans. Terminal portion enlarged.

the base to an acute extremity; dorsal side usually more convex than the ventral, and often longitudinally sinuate; aperture usually oblique and extended on the ventral side; surface smooth, or having arching or transverse striae. Type H. acutus.

acilis, Hall, 1876, Illust. Devonian Foss., pl. 27, and Pal. N. Y., vol. 5, pt. 2, p. 197, Ham. Gr.

aculeatus, Hall, 1860, (Theca aculeata,) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 107, Kinderhook Gr.

americanus, Billings, 1871, (Theca triangularis,) Hall, Can. Nat. and Geol., vol. 6, p. 213, Up. Taconic.

baconi, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 77, and Geo. Wis., vol. 4, p. 225, Trenton Gr.

carbonaria, Walcott, Monogr. U. S. Geo. Sur., vol. 8, p. 264, Subcarboniferous.

centennialis, Barrett, 1877, Ann. Lyc. Nat. Hist., vol. 11, p. 299, Low. Held. Gr.

communis, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 213, Up. Taconic.

danicus, Matthew, 1884, Bull. U. S. Geo. Sur., vol. 2, p. 283, St. John Gr.

emmonsii, Ford, 1873, Am. Jour. Sci., 3d ser., vol. 5, p. 214, Up. Taconic.

excellens, Billings, 1874, Pal. Foss., vol. 2, p. 70, Up. Taconic.

gibbosus, Hall & Whitfield, 1873, 23d Rep. N. Y. St. Mus. Nat. Hist., p. 242, Potsdam Gr.

gracilis, Matthew, 1885, (Camerotherca gracilis,) Can. Rec. Sci., vol. 1, p. 149, St. John Gr.

gregarius, Meek & Hayden, 1861, Proc. Acad. Nat. Sci. Phil., p. 436, and Pal. Up. Mo., p. 5, Potsdam Gr.

heros, Hall, 1888, Pal. N. Y., vol. 7, p. 7, Low. Held. Gr.

impar, Ford, 1872, Am. Jour. Sci., 3d ser., vol. 3, p. 419, Up. Taconic.

ligea, Hall, 1863, (Theca ligea,) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 62, and Pal. N. Y., vol. 5, pt. 2, p. 195, Up. Held. Gr.

micans, see Hyolithes micans.

micmac, Matthew, 1884, Full. U. S. Geo. Sur., vol. 2, p. 283, St. John Gr.

neapolis, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 56, Portage Gr.

parviusculus, Hall, 1862, (Theca parviuscula,) Geo. Rep. Wis., p. 425, Hud. Riv. Gr.

primordialis, Hall, 1861, (Theca primordialis,) Geo. Rep. Wis., p. 48, and Geo. Wis., vol. 4, p. 175, Potsdam Gr.

princeps, Billings, 1871, Can. Nat. and Geol., vol. 6, p. 213, and Am. Jour. Sci. and Arts, 3d ser., vol. 3, p. 355, Up. Taconic.

principalis, Hall, 1876, Illust. Devonian Foss., pl. 27, and Pal. N. Y., vol. 5, pt. 2, p. 196, Schoharie grit.

shaleri, Walcott, 1885, Bull. U. S. Geo. Sur., p. 283, Up. Taconic.

singulus, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 202, Ham. Gr.

FIG. 646.—Hyolithes primordialis.

striatus, Hall, 1876, *Illust. Devonian Foss.*, pl. 27, and *Pal. N. Y.*, vol. 5, pt. 2, p. 199, *Ham. Gr.*
 subimbricatus, Ringueberg, 1888, *Proc. Acad. Nat. Sci. Phil.*, p. 135, *Niagara Gr.*
 triliratus, Hall, 1879, *Pal. N. Y.*, vol. 5, pt. 2, p. 201, *Ham. Gr.*
 vanuxemi, Walcott, 1885, *Monogr. U. S. Geo. Sur.*, vol. 8, p. 85, *Chazy Gr.*

MATTHEVIA, Walcott, 1885, *Am. Jour. Sci. and Arts*, 3d ser., vol. 30, p. 17. [Ety. proper name.] Shell conical; aperture sinuous, transverse section, ovate, elliptical or rounded subquadrate; two elongate interior chambers, diverging from the apex, open into the terminal chamber, and are crossed by a single

FIG. 647.—Matthewia variabilis.

imperforate septum; surface papillose; operculum calcareous, nucleus excentric, lines of growth concentric. Type M. variabilis.

variabilis, Walcott, 1885, *Am. Jour. Sci. and Arts*, 3d ser., vol. 30, p. 18, *Calceiferous Gr.*

PHARETRELLA, Hall, 1888, *Pal. N. Y.*, vol. 7, p. 7. Shell large, elongate, Hyolithes-like in outline; apex acute; surface ornamented with transverse, undulating striae. Type P. tenebrosa.

tenebrosa, Hall, 1888, *Pal. N. Y.*, vol. 7, p. 7, *Genesee Slate*.

PTEROTHECA, Salter, 1852, *Rep. Brit. Ass'n*, p. 61. [Ety. pteron, wing; Theca, a genus.] Shells arcuate, somewhat calyptraeform, subtriangular, or oval; apex marginal and incurved on the same plane, carinate upon the back, abruptly and broadly expanding, with the anterior margin sinuate; interior concave, shallow; a concave, shelly partition covers the posterior half of the cavity. Type P. transversa.

anatiformis, Hall, 1847, (*Tellinomya anatiformis*), *Pal. N. Y.*, vol. 1, p. 154, *Trenton Gr.*

attenuata, Hall, 1861, (*Cleioderma attenuata*), 14th Rep. N. Y. St. Mus. Nat. Hist., p. 98, *Trenton Gr.*

canaliculata, Hall, 1861, (*Cleioderma canaliculata*), 14th Rep. N. Y. St. Mus. Nat. Hist., p. 97, *Trenton Gr.*

expansa, Emmons, 1842, (*Delthyris expansus*), *Geo. Rep. N. Y.*, p. 397, *Black Riv. and Trenton Grs.*

saffordi, Hall, 1861, (*Cleioderma saffordi*), 14th Rep. N. Y. St. Mus. Nat. Hist., p. 96, *Trenton Gr.*

transversa, Salter, 1852, *Rep. Brit. Ass'n*, p. 61, *Hud. Riv. Gr.*

undulata, Hall, 1861, (*Cleioderma undulata*), 14th Rep. N. Y. St. Mus. Nat. Hist., p. 97, *Trenton Gr.*

Pugiunculus aculeatus, see *Hyolithes aculeatus*.

SCENELLA, Billings, 1872, *Can. Nat. and Geol.*, vol. 6, p. 479, and *Pal. Foss.*, vol. 2, p. 77. [Ety. scene, tent; ella, diminutive.] Shell small, depressed, conical; apex central, an obscure carina extending from the apex to the margin; apex slightly incurved opposite the carina; aperture nearly circular; surface finely reticulated. Type S. reticulata.

conica, Whiteaves, 1884, *Pal. Foss.*, vol. 3, p. 32, *Guelph Gr.*

conula, Walcott, 1885, *Fig. 648.—Scenella conica. Side view.*

Sur., vol. 8, p. 15, *Up. Taconic*.
 reticulata, Billings, 1872, *Can. Nat. and Geol.*, vol. 6, p. 479, and *Pal. Foss.*, vol. 2, p. 77, *Up. Taconic*.

retusa, Ford, 1873, *Am. Jour. Sci. and Arts*, 3d series, vol. 5, p. 213, *Up. Taconic*.

varians, Walcott, 1886, *Bull. U. S. Geo. Sur.*, vol. 30, p. 127, *Up. Taconic*.

STENOITHECA, Hicks, 1872, *Quar. Jour. Geo. Soc.*, vol. 28, p. 180. [Ety. stenos, narrow; Theca, genus.] Shell small, curved; lines of growth strongly marked transversely. Type S. cornucopia.

acadica, Hartt, 1868, (*Discina acadica*), *Acad. Geol.*, p. 644, *St. John Gr.*

concentrica, Matthew, 1885, *Trans. Roy. Soc. Can.*, p. 57, *St. John Gr.*

elongata, Walcott, 1885, *Monogr. U. S. Geo. Sur.*, vol. 8, p. 23, *Up. Taconic*.

hicksana, Matthew, 1885, *Trans. Roy. Soc. Can.*, p. 56, *St. John Gr.*

nasuta, Matthew, 1885, *Trans. Roy. Soc. Can.*, p. 58, and *Can. Nat. and Geol.*, vol. 6, p. 479, *St. John Gr.*

pauper, Billings, 1872, *Pal. Foss.*, vol. 2, p. 77, *Up. Taconic*.

radiata, Matthew, 1885, *Trans. Roy. Soc. Can.*, p. 57, *St. John Gr.*

rugosa, Hall, 1847, (*Metoptoma rugosa*), *Pal. N. Y.*, vol. 1, p. 306, *Up. Taconic*.

triangularis, Matthew, 1885, *Trans. Roy. Soc. Can.*, p. 58, *St. John Gr.*

STYLIOLA, Lesueur, 1826. [Ety. stylos, pillar.] Small, conical, without annulations which distinguishes it from *Tentaculites*.

fissurella, Hall, 1843, (*Tentaculites fissurellus*), *Geo. 4th Dist. N. Y.*, p. 180, and *Pal. N. Y.*, vol. 5, pt. 2, p. 178, *Marcellus Shale and Genesee Slate*.

fissurella var. intermittens, Hall, 1879, *Pal. N. Y.*, vol. 5, pt. 2, p. 181, *Genesee Slate*.

fissurella var. obsolescens, Hall, 1879, *Pal. N. Y.*, vol. 5, pt. 2, p. 180, *Ham. Gr.*

fissurella var. strigata, Hall, 1879, *Pal. N. Y.*, vol. 5, pt. 2, p. 180, *Marcellus Shale*.

FIG. 649.—Stenotheca rugosa.

olithes acu-
n. Nat. and
Foss., vol.
lla, dimin-
ed, conical;
ina extend-
argin; apex
the carina;
urface finely
ata.

Scenella con-
side view.

onic.

n. Nat. and
l. Foss., vol.

ir. Sci. and
p. 213, Up.

U. S. Geo.
conic.

Jour. Geo.
stenos, nar-
all, curved;
arked trans-

ia.
na acadica,
n Gr.

Tra's. Roy.
tr.

noogr. U. S.
Taconic.

ns. Roy. Soc.

ns. Roy. Soc.
nd Geo., vol.

Foss., vol. 2,

ns. Roy. Soc.

57, St. John

ll, 1847, (Me-

rugosa, Pal.

ol. 1, p. 306,

onic.

is, Matthew,

n., p. 58, St.

stolos, pillar.]

annulations

from Tentac-

olithes fissurel-

p. 180, and

178, Marcel-

te.

Hall, 1879,

p. 181, Gen-

obtusa, Hall, 1879, Pal. N. Y., vol. 5, pt.
2, p. 182, Ham. Gr.
spica, Hall, 1888, Pal. N. Y., vol. 7, p. 7,
Ham. Gr.

TENTACULITES, Schlotheim, 1820, Petrefakten.,
p. 377. [Ety. *tentaculum*, feeler; *lithos*,
stone.] Straight, elongate, attenuately
elevated rings closely arranged near the
apex, and more distant and stronger
with the increasing size of the shell
and distance from the apex; surface
marked by fine transverse striae, rarely
by longitudinal striae. Type *T. scalaris*.
acula, Hall, 1888, Pal. N. Y., vol. 7, p. 6,
Low. Held. Gr.

arenosus, Hall, 1876, Illust. Devon. Foss.,
pl. 26, and Pal. N. Y., vol. 5, pt. 2, p.
166, Oriskany sandstone.

attenuatus, Hall, 1876, Illust. Devonian
Foss., pl. 26, and Pal. N. Y., vol. 5, pt.
2, p. 170, Ham. Gr.

bellulus, Hall, 1876, Illust. Devonian
Foss., pl. 26, and Pal. N. Y., vol. 5, pt.
2, p. 169, Ham. Gr.

dexitheia, Hall, 1888, Pal. N. Y., vol. 7, p.
6, Schoharie grit.

distans, see *Cornulites distans*.

elongatus, Hall, 1859, Pal. N. Y., vol. 3,
p. 136, Low. Held. Gr.

fissurella, see *Styliola fissurella*.

flexuosa, see *Conchicolites flexuosus*.

gracilistriatus, Hall, 1879, Pal. N. Y., vol.
5, pt. 2, p. 173, Marcellus Shale.

hoyti, White, 1876, Proc. Acad. Nat. Sci.,
p. 34, Devonian.

gyracanthus, Eaton, 1832, (*Echinus gy-*
racanthus.) Geo. Text-book, p. 128, Low.
Held. Gr.

incurvus, Shumard, 1856, Geo. Rep. Mo.,
p. 195, Trenton Gr.

irregularis, Hall, 1859, Pal. N. Y., vol. 3,
syn. for *T. gyracanthus*.

minutus, Hall, 1843, Geo. Rep. 4th Dist.
N. Y., p. 72, and Pal. N. Y., vol. 2, p.
183, Clinton Gr.

neglectus, Nicholson & Hinde, 1874, Can.
Jour., p. 9, Clinton Gr.

niagarensis,

Hall, 1852,

Pal. N. Y.,

vol. 2, p.

352, Niag-

ara Gr.

niagarensis var. *cumberlandiae*, Hall, 1888,

Pal. N. Y., vol. 7, p. 5, Niagara Gr.

oswegoensis, Meek & Worthen, 1865,

Proc. Acad. Nat. Sci. Phil., p. 254, and

Geo. Sur. Ill., vol. 3, p. 342, Hud. Riv. Gr.

richmondensis, S. A. Miller, 1874, Cin.

Quar. Jour. Sci., vol. 1, p. 234, Hud.

Riv. Gr.

scalariformis, Hall, 1876, Illust. Devonian

Foss., pl. 26, and Pal. N. Y., vol. 5, pt.

2, p. 167, Up. Held. Gr.

scalaris, Schlotheim, 1820, Petref. Not an

American species.

sicula, Hall, 1876, Illust. Devonian Foss.,

pl. 26, Up. Held. Gr.

spicula, Hall, 1876, Illust. Devonian

Foss., pl. 26, and Pal. N. Y., vol. 5, pt.

2, p. 172, Chemung Gr.

sterlingensis, Meek & Worthen, 1865,

Proc. Acad. Nat. Sci. Phil., p. 255,

and Geo. Sur. Ill., vol. 3, p. 342, Hud.

Riv. Gr.

subtilis, Winchell, 1866, Rep. Low. Pen-

insula Mich., p. 92, Ham. Gr.

tenuistriatus, Meek & Worthen, 1865,

Proc. Acad. Nat. Sci. Phil., p. 255, and

Geo. Sur. Ill., vol. 3, p. 343, Hud.

Riv. Gr.

Theca, Sowerby, 1845, syn. for *Hyolithes*.

aculeata, see *Hyolithes aculeatus*.

gregaria, see *Hyolithes gregarius*.

ligea, see *Hyolithes ligea*.

parviuscula, see *Hyolithes parviusculus*.

primordialis, see *Hyolithes primordialis*.

triangularis, Hall, 1847, Pal. N. Y., vol. 1,

p. 313. This name was preoccupied by

Portlock in 1843. Billings described it

as *Hyolithes americanus*.

FIG. 650.—Tentaculites rich-

mondensis.

CLASS GASTEROPODA.

[Ety. *gaster*, under side of body; *pous*, foot.]

THE locomotive organ in the Gasteropoda consists of a broad, muscular under-surface, or foot, upon which the animal creeps with a gliding motion. The expansion and contraction of the muscles may be seen when a snail is moving over glass. This form of the foot is the most important characteristic of the Gasteropoda. The head is distinct, and usually furnished with tentacles and eyes. The mouth is on the lower surface, and is often furnished with one or two teeth, or jaws, in the upper part, and a ribbon-like tongue, with minute silicious teeth on its upper surface, which are used with the upper teeth in separating the food. The teeth on the tongue are called the lingual teeth.

The body is generally much larger on one side than on the other, which produces a spiral shell in the growth of the animal, because the shell is secreted at the edge of the mantle. The shell nearly always consists of one piece (univalve), forming a conical tube, twisted spirally; but the tube is not perfect, because the inner wall of each whorl is formed of the preceding whorl with only a thin coating of shelly matter. Sometimes the tube is rolled in a plane, producing a discoid shell; and there are all grades of form, from the discoid to the upright. The right side of the animal is usually the larger, and the shell produced is dextral; but in some species and in some abnormal specimens of dextral species, the spire is turned in the opposite direction, and the shell is therefore called sinistral. The winding of the tube in the spiral shells as the animal grows, produces a central axis, which is called the columella. It extends from the apex to the base, and forms the inner margin of the aperture. The columella is usually hollow, and terminates at the base of the shell with a small opening, called the umbilicus. The margins of the aperture are called the lips. When the columella forms the inner lip, it is called the columellar lip. The outer lip forms the convexity of the shell. Sometimes the lips are continuous, and sometimes the outer lip is more or less deeply notched; and both lips may be furnished with teeth or denticulated edges. The last whorl of the shell is called the body whorl, from its receiving the body of the animal, and the remaining whorls constitute the spire. The line which separates the whorls is the suture. Many Gasteropoda have a calcareous plate attached to the hinder part of the foot, which closes the aperture when the animal retracts itself within the shell; this covering is called an operculum.

The Gasteropoda are divided into two subclasses: the Heteropoda and Gasteropoda proper. The Heteropoda, also called the Nucleobranchiata, are all inhabitants of the ocean, and usually have a shell covering only the essential organs of the body. They swim rapidly near the surface of the water with the back downward, and when the foot is present it is used to attach the animal to floating sea-weeds.

The Gasteropoda proper are divided into two orders: one breathing air, the Pulmonifera; and the other water, the Branchifera. The Pulmonifera include the

land snails and their allies; the Branchifera are furnished with gills, and include nearly all Paleozoic shells of this Class.

FAMILY BELLEROPHONTIDÆ.—Bellerophon, Bucanella, Bucania, Phragmotoma, Porcellia, Tremanotus.

FAMILY BULIMORPHIDÆ.—Bulimorpha.

FAMILY CALYPTRÆIDÆ.—Capulus, Conchopeltis, Metoptoma, Platyceras.

FAMILY CHITONIDÆ.—Chiton.

FAMILY CLISOSPIRIDÆ.—Billingsia, Clisospira.

FAMILY CODONCHILIDÆ.—Codonochilus.

FAMILY CYCLONEMIDÆ.—Cyclonema, Eunema, Holopea, Holopella, Platyschisma, Orthonema, Palæacmaea, Trochonema.

FAMILY CYCLORIDÆ.—Cyclora.

FAMILY CYRTOLITIDÆ.—Carinaropsis, Conchopeltis, Cyrtolites, Cyrtionella, Microceras.

FAMILY DENTALIIDÆ.—Dentalium.

FAMILY EUOMPHALIDÆ.—Calaurops, Ecyclylimphalus, Euomphalus, Omphalotrochus, Ophileta, Pleuronotus, Straparollina, Straparollus.

FAMILY FUSISPIRIDÆ.—Fusispira.

FAMILY HELICIDÆ.—Anthracopupa, Dawsonella, Pupa, Streptaxis, Strophites, Zonites.

FAMILY LITTORINIDÆ.—Xenophora.

FAMILY MACLURIDÆ.—Maclurea.

FAMILY NATACOSIDÆ.—Callonema, Isonema, Naticopsis, Trachydomia.

FAMILY PATELIDÆ.—Lepetopsis, Tryblidium.

FAMILY PLATYSTOMIDÆ.—Orthostoma, Platystoma, Scævogyra, Strophostylus.

FAMILY PLEUROTOMARIIDÆ.—Helicotoma, Lophospira, Microdoma, Murchisonia, Pleurotomaria, Raphistoma, Scalites.

FAMILY PSEUDOPHORIDÆ.—Pseudophorus.

FAMILY PYRAMIDELLIDÆ.—Loxonema, Macrochilina, Soleniscus, Zaptychius.

FAMILY ROTELLIDÆ.—Anomphalus, Rotella.

FAMILY SUBULITIDÆ.—Polyphemopsis, Subulites.

FAMILY TROCHIDÆ.—Eotrochus, Palæotrochus.

FAMILY TURRITELLIDÆ.—Aclisina, Turritella.

Aclis, Loven, 1846, Index, Mollusc. litora Scandin. occid. habit., p. 16. Not an American Paleozoic genus.

minuta, see *Aclisina minuta*.

robusta, see *Aclisina robusta*.

stevensoni, see *Aclisina stevensoni*.

swallowana, see *Aclisina swallowana*.

ACLISINA, DeKoninck, 1881, Faune du Calcaire Carbonifère de la Belgique Ann. d. Mus. Roy. d'Hist. Nat., t. 6, p. 86. [Ety. diminutive of *Aclis*.] An elongated, banded, conical, spiral shell; distinguished from *Murchisonia* by its oval aperture, and from *Loxonema* by its spiral bands. Type *A. striatula*.

minuta, Stevens, 1858, (*Aclis minuta*),

Am. Jour. Sci., vol. 25, p. 259, Coal Meas.

robusta, Stevens, 1858, (*Aclis robusta*),

Am. Jour. Sci., vol. 25, p.

259, and Geo. Sur. Ill., vol.

5, p. 526, Coal Meas.

stevensoni, White, 1882, (*Aclis*

stevensoni), Rep. Invert.

Foss. New. Mex., p. xxxv,

Coal Meas.

swallowana, Geinitz, 1866,

(*Turbonilla swallowana*),

Carb. und Dyas in Neb., p.

5, Coal Meas.

Acroculia, Phillips, 1841, Pal.

Foss. Cornwall, Devon, and

W. Somerset, p. 93, syn.

Aclisina

for *Platyceras*.

angulata, see *Platyceras angulatum*.

FIG. 651.

Aclisina

swallowana.

- erecta*, see *Platyceras erectum*.
ovalis, see *Platyceras ovale*.
niagarensis, see *Platyceras niagarense*.
trigonalis, see *Platyceras trigonale*.
Ampullaria, Lamarck, 1801, Syst. An. sans Vert. [Ety. *ampulla*, a flask.] Not a Palaeozoic genus.
helicoidea, see *Soleniscus helicoides*.
powelli, Walcott, 1883, Science, vol. 2, p. 808, and Monogr. U. S. Geo. Sur., vol. 8, p. 261, Subcarboniferous.
ANOMPHALUS, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 268. [Ety. *anomphalos*, without an umbilicus.] A helicoid shell of three or more volutions, and having an aperture transversely suboval. Type *A. rotulus*.
meeki, see *Dawsonella meeki*.
rotulus, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 268, and Geo. Sur. Ill., vol. 5, p. 597, Coal Meas.
ANTHRACOPUPA, Whitfield, 1881, Am. Jour. Sci. and Arts, 3d ser., vol. 21, p. 126. [Ety. *anthrax*, coal; *Pupa*, a genus.] Shell minute, pupiform, few volutions, last unsymmetrical; axis imperforate; aperture large, nearly vertical; peristome thickened, united above by a thin callus, on which may occur one or more palatal teeth; other tooth-like projections on the inner margin of lip; circular notch, as in *Pupina*, on inner edge of outer limb, near body whorl; surface vertically lined. Type *A. ohioensis*.
ohioensis, Whitfield, 1881, Am. Jour. Sci. and Arts, 3d ser., vol. 21, p. 126, Coal Meas.
BELLEROPHON, Montfort, 1808, Conch. Syst., vol. 1, p. 50. [Ety. mythological name.] Shell thick, symmetrical, globose, involute; sinus in the middle of the outer lip, from which a band extends backward along the outer surface of the volution; inner lip thickened, expanded on the inrolled spire. Type *B. vasulites*.
acutillra, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 56, Ham. Gr.
acutus, Sowerby, 1839, Murch. Sil. Syst., p. 643, Low. Silurian.
allegoricus, White, 1874, Rep. Invert. Foss., p. 10, and Geo. Sur. W. 100th Mer., vol. 4, p. 55, Quebec Gr.
alternodosus, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 225, Kaskaskia Gr.
angustata, see *Bucania angustata*.
antiquatus, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 52, and Geo. Wis., vol. 4, p. 176, Potsdam Gr.
apertus, Sowerby, 1825, Min. Conch., vol. 5, p. 108, Subcarboniferous.
argo, Billings, 1860, Can. Nat. and Geol., vol. 5, p. 167, Black Riv. and Trenton Gr.
auriculatus, Hall, 1852, Pal. N. Y., vol. 2, p. 334, Coralline limestone.
barquensis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 425, Marshall Gr.
bidorsatus, see *Bucania bidorsata*.

bilabiatus, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 304, Kinderhook Gr.

bilobatus, Sowerby, 1839, Murch. Sil. Syst., p. 643, and Pal. N. Y., vol. 1, p. 184, Black Riv. to Mid. Sil.

FIG. 652.—*Bellerophon bilobatus*.

- bilobatus* var. *acutus*, Hall, 1847, Pal. N. Y., vol. 1, p. 185, Trenton Gr.
bilobatus var. *corrugatus*, Hall, 1847, Pal. N. Y., vol. 1, p. 185, Trenton Gr.
blaneyanus, syn. for *B. carbonarius*.
bowmani, White, 1876, Proc. Acad. Nat. Sci., p. 32, Devonian.
brevilineatus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 269, and Pal. N. Y., vol. 5, p. 2, p. 107, Ham. Gr.
canadensis, Billings, 1866, Catal. Sil. Foss. Antic., p. 18, Hud. Riv. Gr.
cancellatus, Hall, 1847, Pal. N. Y., vol. 1, p. 307, Hud. Riv. Gr.
cancellatus, Hall, 1858, Trans. Alb. Inst., vol. 4. The name was preoccupied.
 See *B. textilis*.
carbonarius, Cox, 1857, Geo. Rep. Ky., vol. 3, p. 562, Coal Meas.
carbonarius var. *subpapillosus*, White, 1876, Geo. Uinta Mountains, p. 92, Up. Aubrey Gr.
carinatus, Sowerby, 1839, Murch. Sil. Syst., p. 634, Devonian.
cassinensis, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 318, Birdseye Gr.
charon, Billings, 1860, Can. Nat. and Geol., vol. 5, p. 169, Black Riv. and Trenton Grs.
combsi, Wolcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 193, Devonian.
convolutus, Eaton, 1832, Geo. Text-book, p. 28, Up. Sil.
crassus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 458, and Geo. Sur. Ill., vol. 2, p. 385, Coal Meas.
crenistris, Hall, 1876, Illust. Devonian Foss., pl. 25, and Pal. N. Y., vol. 5, pt. 2, p. 116, Ham. Gr.
curvilineatus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 269, Onondaga, Schoharie and Up. Held. Gr.
cyrtolites, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 107, Kinderhook Gr.
declivis, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 269, Trenton Gr.
disculus, Billings, 1860, Can. Nat. and Geol., vol. 5, p. 168, Black Riv. and Trenton Gr.
ellipticus, McChesney, 1860, Desc. New Pal. Foss., p. 58, Coal Meas.
expansus, see *Bucania expansa*.
explanatus, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 109, Chemung Gr.

i, 1862, Proc.
p. 304, Kin-

Bellerophon
tutus.

Y., vol. 1, p.

ll, 1847, Pal.

on Gr.

arius.

. Acad. Nat.

Jour. Acad.

l Pal. N. Y.,

r.

tal. Sil. Foss.

N. Y., vol. 1,

. Alb. Inst.,

preoccupied.

. Rep. Ky.,

osus, White,

is, p. 92, Up.

Murch. Sil.

Bull. Am.

. 318, Birds-

at. and Geol.

. and Tren-

r. U. S. Geo.

an.

. Text-book,

1860, Proc.

and Geo. Sur.

as.

t. Devonian

r., vol. 5, pt.

Jour. Acad.

Onondaga,

Gr.

Rep. N. Y.

17, Kinder-

. Acad. Nat.

Gr.

n. Nat. and

ck Riv. and

, Desc. New

s.

a.

N. Y., vol. 5,

ficellostriatus, Foerste, 1885, Bull. Sci.,
Lab. Denison Univ., p. 99, Niagara Gr.
fraternus, Billings, 1866, Catal. Sil. Foss.,
Antic., p. 19, Hud. Riv. Gr.

galericulatus, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 426, Marshall Gr.

gibsoni, White, 1882, 11th Rep. Geol. and
Nat. Hist. Indiana, p. 360, St. Louis Gr.
giganteus, Worthen, 1884, Bull. No. 2,
Ill. St. Mus. Nat. Hist., p. 8, and Geo.

Sur. Ill., vol. 8, p. 143, Low. Coal Meas.
globosus, Stevens, 1858, Am. Jour. Sci.,
vol. 25, p. 258, Coal Meas.

harrodi, Gurley, 1883, New Carb. Foss.,
p. 5. Publication not such as to establish
a species.

helena, Hall, 1879, Pal. N. Y., vol. 5, pt.
2, p. 114, Ham. Gr.

hindus, Sowerby, Min. Conch. Not
American.

hyalina, Hall, 1879, Pal. N. Y., vol. 5, pt.
2, p. 99, Up. Held. Gr.

incisus, Clarke, 1885, Bull. U. S. Geo.
Sur., vol. 16, p. 53, Portage Gr.

inspeciosus, White, 1882, Rep. Invert.
Foss. New Mex., p. xxx, Coal Meas.

interlineatus, Portlock, 1843, Geo. of Lon-
donderry, p. 402, Coal Meas. Probably
not American.

kansasensis, Shumard, 1858, Trans. St.
Louis Acad. Sci., vol. 1, p. 204, Coal
Meas.

leda, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 58, and Pal. N. Y., vol. 5,
pt. 2, p. 110, Ham. Gr.

lineolatus, Hall, 1860, 13th Rep. N. Y.
Mus. Nat. Hist., p. 107, Waverly or
Kinderhook Gr.

lindsleyi, Safford, 1869, Geo. of Tenn., p.
289, Nashville Gr.

lyra, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 59, and Pal. N. Y., vol. 5,
pt. 2, p. 113, Ham. Gr.

macer, Billings, 1865, Pal. Foss., vol. 1, p.
347, Calciferous Gr.

maria, Hall, 1876, Illust. Devonian Foss.,
pl. 22, and Pal. N. Y., vol. 5, pt. 2, p.
119, Chemung Gr.

FIG. 633.—*Bellerophon mohri*.

majusculus, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 256, Subcarb.

marcouanus, Geinitz, 1866, Carb. und
Dyas in Neb., p. 7, and Pal. E. Neb.,
p. 226, Coal Meas.

mekkanus, Swallow, 1858, Trans. St.
Louis Acad. Sci., vol. 1, p. 204, Coal
Meas.

michiganensis, Winchell, 1862, Proc.
Acad. Nat. Sci., p. 427, Marshall Gr.

miser, Billings, 1866, Catal. Sil. Foss.
Antic., p. 20, Hud. Riv. Gr.

missouriensis, Swallow, 1863, Trans. St.
Louis Acad. Sci., vol. 2, p. 100, Kas-
kaskia Gr.

mohri, S. A. Miller, 1874, Cin. Quar. Jour.
Sci., vol. 1, p. 300, Hud. Riv. Gr.

montfortanus, Norwood & Pratten, 1855,
Jour. Acad. Nat. Sci., vol. 3, p. 74, Coal
Meas.

FIG. 634.—*Bellerophon palmurus*.

morrowensis, Miller & Dyer, 1878, Cont-
to Pal. No. 2, p. 8, Hud. Riv. Gr.

nactus, Hall, 1879, Pal. N. Y., vol. 5, pt.
2, p. 121, Chemung Gr.

nashvillensis, Troost, 1840, 5th Geo. Rep.
Tenn., p. 54, Trenton Gr.

natator, Hall, 1862, (Phragmostoma na-
tator,) 15th Rep. N. Y. Mus. Nat. Hist.,
p. 60, and Pal. N. Y., vol. 5, pt. 2, p.
108, Ham. Gr.

nautiloides, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 427, Marshall Gr.

nemens, Hall & Whitfield, 1876, Illust.
Devonian Foss., pl. 22, and U. S. Geo.

40th Parallel, p. 250, Chemung Gr.

newberryi, Meek, 1871, Proc. Acad. Nat.
Sci., p. 77, and Ohio Pal., vol. 1, p. 222,
Up. Held. Gr.

nodocarinatus, Hall, 1858, Geo. Rep. Iowa,
p. 723, Coal Meas.

obsoletus, Hall, 1876, Illust. Devonian
Foss., pl. 22, Chemung Gr.

otsego, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 60, and Pal. N. Y., vol. 5,
pt. 2, p. 104, Ham. Gr.

palmurus, Billings, 1865, Pal. Foss., vol.
1, p. 311, Quebec Gr.

pannus, White, 1862, Proc. Bost. Soc.
Nat. Hist., vol. 9, p. 21, Marshall Gr.

patersoni, Hall, 1862, Geo. Rep. Wis., p.
55, Hud. Riv. Gr.

patulus, Hall, 1843, Geo. Rep. 4th Dist.
N. Y., p. 196, and Pal. N. Y., vol. 5, pt.
2, p. 100, Ham. Gr.

pelops, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 56, and Pal. N. Y., vol. 5,
pt. 2, p. 95, Schoharie and Up. Held. Gr.

pelops var. exponens, Hall, 1879, Pal.
N. Y., vol. 5, pt. 2, p. 96, Up. Held. Gr.

percarinatus, Conrad, 1842, Jour. Acad.
Nat. Sci., vol. 8, p. 268, Coal Meas.

perelegans, White & Whitfield, 1862, Proc.
Bost. Soc. Nat. Hist., vol. 9, p. 304,
Kinderhook Gr.

perforatus, Winchell & Marcy, 1866, syn-
for *Tremanotus chicagoensis*.

- ✓ perplexus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 193, Devonian.
- perlatius, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 270, Coal Meas.
- platystoma, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 312, Galena Gr.
- ✓ plenus, Billings, 1874, Pal. Foss., vol. 2, p. 62, Gaspe limestone No. 8, Devonian.
- profundus, Emmons, Geo. Rep., 2d Dist. N. Y., p. 303, Trenton Gr.
- ✓ propinquus, Meek, 1871, Proc. Acad. Nat. Sci., p. 78, and Ohio Pal., vol. 1, p. 226, Up. Held. Gr.
- ✓ punctifrons, see Bucania punctifrons.
- ✓ repertus, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 128, Ham. Gr.
- ✓ rotalina, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 115, Ham. Gr.
- ✓ rudis, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 57, and Pal. N. Y., vol. 5, pt. 2, p. 103, Ham. Gr.
- rugosiusculus, Winchell, 1862, Proc. Acad. Nat. Sci., p. 425, Marshall Gr.
- rugosus, Emmons, 1856, Am. Geol., p. 160, Hud. Riv. Gr.
- scriptiferus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 21, Marshall Gr.
- scissile, Conrad, 1844, Proc. Acad. Nat. Sci., vol. 2, p. 175, Kaskaskia Gr. Very poorly defined.
- solitarius, Billings, 1866, Catal. Sil. Foss. Antic., p. 20, Hud. Riv. Gr.
- stamineus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 209, Marshall Gr.
- stevensanus, McChesney, 1860, Desc. New Pal. Foss., p. 61, Coal Meas.
- sublaevis, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 32, and Geo. Sur. Iowa, p. 666, Warsaw Gr.
- subpapillosus, White, 1879, Bull. U. S. Geo. Sur. Ter., vol. 5, p. 218, and Cont. to Pal. No. 6, p. 138, Carboniferous.
- sulcatinus, see Bucania sulcatina.
- textiliformis, Gurley, 1883, New Carb. Foss., p. 6. Publication not valid.
- textilis, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 243, Warsaw Gr. Proposed instead of *B. cancellatus*, Hall, 1858, which was preoccupied.
- ✓ thalia, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 60, and Pal. N. Y., vol. 5, pt. 2, p. 105, Ham. Gr.
- tricarinatus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 204, Coal Meas.
- tricarinata, Hall, 1876, Illust. Devonian Foss. The name was preoccupied. See *B. triliratus*.
- ✓ triliratus, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 243, and Pal. N. Y., vol. 5, pt. 2, p. 117, Chemung Gr. Proposed instead of *B. tricarinatus*, Hall, 1876, which was preoccupied.
- troosti, D'Orbigny, 1840, Cephal., p. 206, and Geo. of Tenn., p. 289, Trenton Gr.
- tuber, Hall, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 177, Niagara Gr.
- urii, Fleming, 1828, British Animals, p. 338, Devonian. American species. (?)
- vinculatus, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 304, Kinderhook Gr.
- vittatus, syn. for *B. carbonarius*.
- volutus, Eaton, 1832, Geol. Text-book, p. 28, Up. Sil.
- whittleseyi, Winchell, 1865, Proc. Acad. Nat. Sci., p. 130, Cuyahoga shale.
- wisconsinensis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 76, and Geo. Wis., vol. 4, p. 223, Trenton Gr.
- Billingsia, Walcott, 1888, Bull. No. 30, U. S. Geo. Sur., p. 61. [Ety. proper name.] Syn. (?) for *Clisospira*. Turbinate, whorls, subcircular. Type *B. saratogensis*. Preoccupied.
- saratogensis, Walcott, 1888, Bull. No. 30, U. S. Geo. Sur., p. 61, Up. Taconic. FIG. 653.—1111. Ingsa saratogensis.
- BUCANELLA, Meek, 1870, Proc. Am. Phil. Soc., vol. 11, p. 426. [Ety. diminutive of *Bucania*.] Type *B. nana*.
- nana, Meek, 1870, Proc. Am. Phil. Soc., vol. 11, p. 426, Silurian.
- BUCANIA, Hall, 1847, Pal. N. Y., vol. 1, p. 32. [Ety. *bukane*, trumpet.] Convolute, spire equally concave on either side; volutions in the same plane, all visible; outer one ventricose, inner one usually angulated on the edge, concave on the ventral side; aperture rounded oval, somewhat compressed on the inner side by contact with the next volution, laterally and dorsally abruptly expanded. Type *B. sulcatina*.
- angustata, Hall, 1852, Pal. N. Y., vol. 2, p. 349, Niagara and Guelph Gr.
- bellipuncta, Hall, 1852, Pal. N. Y., vol. 2, p. 93, Clinton Gr.
- bidorsata, Hall, 1847, (Bellerophon bidorsatus), Pal. N. Y., vol. 1, p. 186, Trenton Gr.
- 2 buelli, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 76, and Geo. Wis., vol. 4, p. 24, Trenton Gr.
- chicagoensis, see *Tremanotus chicagoensis*.
- costata, James, 1872, (Cyrtolites costatus), Am. Jour. Sci., 3d ser., vol. 3, p. 26, and Ohio Pal., vol. 1, p. 150, Hud. Riv. Gr.
- crassolaris, McChesney, 1861, New Pal. Foss., p. 91, Niagara Gr.
- devonica, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 195, Up. Held. Gr.
- euomphaloides, Owen, 1862, Geo. Sur. Ind., p. 362. Not very satisfactorily defined.
- exigua, Foerste, 1825, Bull. Sci. Lab. Denison Univ., p. 99. Not properly defined.
- expansa, Hall, 1847, Pal. N. Y., vol. 1, p. 186, Trenton Gr.
- intexta, Hall, 1847, Pal. N. Y., vol. 1, p. 317, Trenton Gr.
- lirata, Hall, 1862, Geo. Rep. Wis., p. 55, Trenton Gr.

eld, 1862, Proc.
8, p. 304, Kin-

rius.
Text-book, p.

, Proc. Acad.
ga shale.

1878, Ann. Rep.
and Geo. Wis.,

Fig. 653.—Bulimella sulcatina.

p. 426. [Ety.
Type B. nana.
m. Phil. Soc.,

Y., vol. 1, p. 32.
] Convolute,
ne, on either side;
one, all visible,
er one usually
concave on the
rounded oval,
on the inner
next volution,
abruptly ex-
na.

N. Y., vol. 2, p.
Gr.

. N. Y., vol. 2,
erophon bidor-
, p. 186, Tren-

m. Rep. Geo.
so. Wis., vol. 4.

otus chicago-

blites costatus.)
vol. 3, p. 26,
p. 150, Hud.

1861, New Pal.

eld, 1872, 24th
st., p. 195, Up.

3602, Geo. Sur.
y satisfactorily

Sci. Lab. Deni-
properly defined.

N. Y., vol. 1, p.

. Y., vol. 1, p.

ep. Wis., p. 55,

pervoluta, McChesney, 1801, New Pal.
Foss., p. 91, Niagara Gr.
profunda, Conrad, 1841, (Euomphalus pro-
fundus,) Ann. Rep. N. Y., p. 37, and
Pal. N. Y., vol. 3, p. 341, Up. Held. Gr.

FIG. 658.—Bucania sulcatina.

punctifrons, Emmons, 1842, (Bellerophon
punctifrons,) Geo. Rep. 2d Dist. N. Y.,
p. 392, and Pal. N. Y., vol. 1, p. 187,
Black River and Trenton Grs.

rotundata, Hall, 1847, Pal. N. Y., vol. 1,
p. 33, Chazy Gr.

stigmata, Hall, 1852, Pal. N. Y., vol. 2, p.
92, Clinton Gr.

sulcatina, Emmons, 1842, (Bellerophon
sulcatinus,) Geo. Rep. 2d Dist. N. Y.,
p. 312, Pal. N. Y., vol. 1, p. 32, Chazy,
Black Riv., and Trenton Grs.

trilobata, Conrad, 1839, (Planorbis tri-
lobatus,) Ann. Rep. N. Y., p. 65, and
Pal. N. Y., vol. 2, pp. 13 and 93, Me-
dina sandstone and Clinton Gr.

tripla, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 55, Calcifer-
ous Gr.

Bulimella, Hall, 1858, Trans. Alb. Inst., vol.
4. This name was preoccupied by
Pfeiffer in 1852. See Bulimorpha.

bulimiformis, see Bulimorpha bulimiformis.

canaliculata, see Bulimorpha canaliculata.

elongata, see Bulimorpha elongata.

BULIMORPHA, Whitfield, 1882, Bull. Am.
Mus. Nat. Hist., No. 3, p. 74. [Ety.

Bulimus, a genus; morphe, form.] Fu-
siform, volutions convex; columella

bent, truncated at the base, separated
from the outer lip by a notch,
as in Achatina; outer lip

slightly notched near the
upper end; surface smooth.
Type B. bulimiformis.

bulimiformis, Hall, 1858,
(Bulimella bulimiformis,) Trans. Alb. Inst., vol. 4, p.

29, and Bull. Am. Mus.
Nat. Hist., p. 74, Warsaw Gr.

canaliculata, Hall, 1858,
Bulimella canaliculata,) Trans. Alb. Inst., vol. 4, p.

29, and Bull. Am. Mus. Nat. Hist., p.
74, Warsaw Gr.

29, and Bull. Am. Mus. Nat. Hist., p.
74, Warsaw Gr.

29, and Bull. Am. Mus. Nat. Hist., p.
74, Warsaw Gr.

elongata, Hall, 1858, Bulimella elongata,) Trans. Alb. Inst., vol. 4, p. 30, and Bull. Am. Mus. Nat. Hist., p. 75, Warsaw Gr. CALLONEMA, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 50. [Ety. kallos, beautiful;

nema, thread.] Subglobose, turbinate or ovoid-contal; volutions rounded or sub-angular above and below; outer lip thin; columnar lip thickened, spreading over the volution above and extended below; axis umbilicate; surface marked by striae extending backward from the sutures over the volutions. Type C. bellatulum.

bellatulum, Hall, 1861, (Loxonema bellatulum,) 14th Rep. N. Y. Mus. Nat. Hist., p. 104, and Pal. N. Y., vol. 5, pt. 2, p. 51, Up. Held. Gr.

imitator, Hall & Whitfield, 1872, (Pleuronomaria imitator,) 24th Rep. N. Y. Mus. Nat. Hist., p. 195, Ham. Gr.

lichas, Hall, 1861, (Platy-
stoma lichas,) 14th Rep. N. Y. Mus. Nat. Hist., p. 106, Up. Held. Gr.

occidentale, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 180, Devonian.

CALAUROPS, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 314. [Ety. kalaurops, a shepherd's crook.] Univalve, discoidal, convolute, inner volutions closely

coiled, outer one disunited and projected in a straight line. Type C. lituiformis. It seems to be distinguished from Ecyliomphalus only by having the last whorl straightened, which may or may not be of generic importance.

lituiformis, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 315, Chazy Gr.

CAPULUS, Montfort, 1810, Conch. Syst., vol. 2, p. 55. [Ety. capulus, a head-piece or cap.] Shell wide, cap-shaped, apex obliquely inclined backward and in-rolled toward the left side; aperture broad, oval, edge irregularly sinuated; muscular scar horseshoe-shaped, open in front. Type C. hungaricus. The horseshoe-shaped, muscular impression has never been observed in any American Palaeozoic fossil, and hence the species referred to this genus do not belong to it. Those named have been so poorly defined, their generic relations can not be determined, and they may as well be struck from the list of names.

acutirostris, see Platyceras acutirostrum.

auriformis, Hall, 1847, Pal. N. Y., vol. 1, p. 31, Chazy Gr.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

parvus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

triplicatus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 205, Coal Meas.

CARINAROPSIS, Hall, 1847, Pal. N. Y., vol. 1, p. 183.

[Ety. from its resemblance to *Carinaria*.] Shell subconical, patelliform; apex incurved or convolute, subcentral; aperture oval, expanded, narrowed posteriorly. Type *C. carinata*.

carinata, Hall, 1847, Pal. N. Y., vol. 1, p. 183, Trenton Gr.

orbiculata, Hall, 1847, Pal. N. Y., vol. 1, p. 306, Hud. Riv. Gr.

patelliformis, Hall, 1847, Pal. N. Y., vol. 1, p. 183, Trenton and Hud. Riv. Grs.

Chemnitzia, D'Orbigny, 1837, Mollusques, Echinodermes, Foraminiferes et Polypiers, etc. Slender, elongated, many whorled, plaited; apex sinistral; aperture simple, ovate; peristome incomplete; operculum horny; subspiral. Type *C. elegantissima*. Not an American Palaeozoic genus.

attenuata, see *Loxonema attenuatum*.

parva, see *Loxonema parvum*.

swallowana, see *Loxonema swallowanum*.

tenuilineata, see *Loxonema tenuilineatum*.

Chiton, Linnaeus, 1758, Syst. Nat., ed. 10, p. 667. [Ety. *chiton*, a coat of mail.] Shell composed of eight transverse imbricating plates, lodged in a coriaceous mantle, which forms an expanded margin round the body. Type *C. squamosus*. Not an American Palaeozoic genus.

canadensis, see *Metopoma canadense*.

carbonarius, Stevens, 1859, Am. Jour. Sci., vol. 25, p. 264, and Geo. Sur. Ill., vol. 5, p. 608. Probably a crustacean, Coal Meas.

parvus, Stevens, 1859, Am. Jour. Sci., vol. 25, p. 264, Coal Meas.

CLISOSPIRA, Billings, 1865, Pal. Foss., vol. 1, p. 186 and 420.

[Ety. *kleio*, to lock; *spira*, whorl.] Shell conical; aperture widely expanded all round in a plane at a right angle to the longitudinal axis of the corical spire; suture in the spire, but obsolete below. Type *C. curiosa*.

curiosa, Billings, 1865, Pal. Foss., vol. 1, pp. 186 and 420, Up. Taconic.

lirata, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 308, Birdseye Gr.

occidentalis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 75, and Geo. Wis., vol. 4, p. 222, Trenton Gr.

CODONOCILUS, Whiteaves, 1884, Pal. Foss., vol. 3, p. 17. [Ety. *kodon*, a trumpet; *cheilos*, lip.] Tur-

reted, subfusiform; volutions numerous, compressed, closely inrolled; outer half of body whorl produced obliquely outward and downward; lip thin, expanded; aperture nearly circular. Type *C. striatum*.

striatum, Whiteaves, 1884, Pal. Foss., vol. 3, p. 17, Guelph Gr.

CONCHOPELTIS, Walcott, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 93. [Ety. *conche*, shell; *pelte*, shield.] Patelliform, more or less conical, apex central or subcentral, vertically striated, older specimens lined concentrically. Type *C. alternata*.

alternata, Walcott, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 93, Trenton Gr.

minnesotensis, Walcott, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 94, Trenton Gr.

CYCLONEMA, Hall, 1852, Pal. N. Y., vol. 2, p. 89. [Ety. *kuklos*, circle; *nema*, thread.] Turbinate, thin, whorls ventricose, striae concentric and crossed by oblique lines of growth; no umbilicus; mouth rounded and with an im-

perfect peritreme; inner lip thin, closely reflected, and a little concave. Type *C. bilix*.

bellulum, Billings, 1866, Catal. Sil. Foss. Antic., p. 55, Anticosti Gr.

bilix, Conrad, 1842, (Pleurotomaria bilix,) Jour. Acad. Nat. Sci., vol. 8, p. 271, and Pal. N. Y., vol. 1, p. 305, Trenton and Hud. Riv. Grs.

bilix var. *conicum*, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 320, Hud. Riv. Gr.

bilix var. *fluctuatum*, James, 1874, (Cyclonema fluctuata,) Cin. Quar. Jour. Sci., vol. 1, p. 152, Hud. Riv. Gr.

cancellatum, Hall, 1843, (Littorina cancellata,) Geo. Rep. 4th Dist. N. Y., p. 72, and Pal. N. Y., vol. 2, p. 90, Clinton Gr.

cincinnatiense, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 230, Utica Slate Gr.

commune, Billings, 1866, Catal. Sil. Foss. Antic., p. 55, Anticosti Gr.

concinnum, Hall, 1876, Illust. Devonian Foss., pl. 12, and Pal. N. Y., vol. 5, pt. 2, p. 38, Chemung Gr.

crenistris, Hall, 1876, Illust. Devonian Foss., pl. 12, Schoharie grit.

crenulatum, Meek, 1871, Proc. Acad. Nat. Sci., p. 79, and Ohio Pal., vol. 1, p. 213, Up. Held. Gr.

decorum, Billings, 1866, Catal. Sil. Foss. Antic., p. 56, Anticosti Gr.

FIG. 659.—*Carinaropsis patelliformis*.

FIG. 662.—*Codonocylus striatum*.

FIG. 660.—*Chiton squamosus*.

FIG. 663.—*Cyclonema bilix*.

FIG. 661.—*Clisospira curiosa*.

78, Ann. Rep.
and Geo. Wis.,

84, Pal. Foss.,

on, a

Tur-

ose-

body

uely

lip

ture

C.

FIG. 662.—
Cyclonema
hallanum.

Pal. Foss.,

76, 28th Rep.

p. 93. [Ety.

] Patelliform,

hex central or

triated, older

rically. Type

8th Rep. N. Y.

Trenton Gr.

876, 28th Rep.

94, Trenton Gr.

N. Y., vol. 2,

nema, thread.]

is ventricose,

assed by oblique

bilicus; mouth

l with an im-

itreme; inner

osely reflected,

concave. Type

Billings, 1866,

Foss. Antic., p.

Gr.

1842, (Pleuro-

x.) Jour. Acad.

p. 8, p. 271, and

vol. 1, p. 305,

Grs.

Miller, 1874,

1, p. 320, Hud.

s, 1874, (Cyclo-

nar. Jour. Sci.,

Gr.

ittorina cancel-

t. N. Y., p. 72,

90, Clinton Gr.

er, 1882, Jour.

5, p. 230, Uta

atal. Sil. Foss.

r.

ust. Devonian

Y., vol. 5, pt.

ust. Devonian

grit.

roc. Acad. Nat.

, vol. 1, p. 213,

atal. Sil. Foss.

Gr.

doris, Hall, 1862, (Pleurotomaria doris.)

15th Rep. N. Y. Mus. Nat. Hist., p. 43,

and Pal. N. Y., vol. 5, pt. 2, p. 34,

Up. Held. Gr.

elevatum, Hall, 1863, 20th Rep. N. Y. Mus.

Nat. Hist., p. 391, Niagara Gr.

FIG. 664.—Cyclonema hageri.

hageri, Billings, 1862, Pal. Foss., vol. 1, p.

29, Trenton Gr.

hallanum, Salter, 1859, Can. Org. Rem.,

Decade 1, p. 26, Black Riv. Gr.

hamiltoniae, Hall, 1862, 15th Rep. N. Y.

Mus. Nat. Hist., p. 47, and Pal. N. Y.,

vol. 5, pt. 2, p. 37, Ham. Gr.

humile, Billings, 1866, Catal. Sil. Foss.

Antic., p. 56, Anticosti Gr.

leavenworthianum, Hall,

1858, (Pleurotomaria

leavenworthianum.)

Trans. Alb. Inst., vol.

4, p. 24, and Bull. Am.

Mus. Nat. Hist., p. 75,

Warsaw Gr.

liratum, Hall, 1862, 15th

Rep. N. Y. Mus. Nat.

Hist., p. 47, and Pal.

N. Y., vol. 5, pt. 2, p.

35, Ham. Gr.

mediocre, Billings, 1866,

Catal. Sil. Foss. Antic., p. 56, Anticosti Gr.

montrealense, Billings, 1862, Pal. Foss.,

vol. 1, p. 30, Trenton Gr.

multiliratum, Hall, 1862, 15th Rep. N. Y.

Mus. Nat. Hist., p. 48, and Pal. N. Y.,

vol. 5, pt. 2, p. 36, Ham. Gr.

obsolescens, Hall, 1877, 1st Ed. Am. Pal.

Foss., p. 243, and Pal. N. Y., vol. 5, pt.

2, p. 38, Chemung Gr. Proposed in-

stead of *C. obsoleta*, Hall, 1876, which

was preoccupied.

obsoletum, Hall, 1852, Pal. N. Y., vol. 2,

p. 90, Clinton Gr.

obsoleta, Hall. The name was preoc-

cupied. See *C. obsolescens*.

percarinatum, Hall, 1847, (Pleurotomaria

percarinata.) Pal. N. Y., vol. 1, p. 177,

Trenton and Hud. Riv. Grs.

percingulatum, Billings, 1857, Rep. of

Progr. Geo. Sur. Can., p. 304, Clinton

and Niagara Grs.

phedra, Billings, 1865, Pal. Foss., vol. 1,

p. 188, Quebec Gr.

pyramidatum, James, 1874, Cin. Quar.

Jour. Sci., vol. 1, p

carinatus, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 311, Utica Slate. compressus, Conrad, 1838, (Phragmolites compressus,) Ann. Rep. N. Y., p. 119, and Pal. N. Y., vol. 1, p. 188, Black Riv. and Trenton Grs.

conradi, Hall, 1862, Geo. Rep. Wis., p. 55, Trenton Gr.

costatus, see Bucania costata.

cristatus, Safford, 1860, Geo. of Tenn., p. 289, Nashville Gr.

desideratus, Billings, 1866, Catal. Sil. Foss. Antic., p. 21, Hud. Riv. Gr.

dyeri, Hall, 1871, 24th Rep. N. Y. Mus. Nat. Hist., p. 230, Hud. Riv. Gr.

FIG. 668.—Cyrtolites elegans.

elegans, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 310, Hud. Riv. Gr.

expansus, Hall, 1859, Pal. N. Y., vol. 3, p. 479, Oriskany sandstone.

filosus, Emmons, 1842, Geo. Rep. 2d Dist. N. Y.,

p. 372, and Pal. N. Y., vol. 1, p. 190, Trenton Gr.

gillanus, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 123, Coal Meas.

imbricatus, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 340, Hud. Riv. Gr.

magnus, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 103, Hud. Riv. Gr.

mitella, see Cyrtionella mitella.

nitidulus, Ulrich, 1878,

Jour. Cin. Soc. Nat. Hist., vol. 2, p. 12, Utica Slate Gr.

ornatus, Conrad, 1838, Ann. Rep. N. Y., p. 118, and Pal. N. Y., vol. 1,

p. 308, Hud. Riv. Gr.

pannosus, Billings, 1866, Catal. Sil. Foss. Antic.,

p. 20, Hud. Riv. Gr.

pileolus, see Cyrtionella lites ornatus.

sinuatus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 237,

Quebec Gr.

sinuosus, Hall, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 178, Niagara Gr.

trentonensis, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 270, and Pal. N. Y.,

vol. 1, p. 189, Trenton Gr.

CYRTONELLA, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 123. [Ety. diminutive of Cyrtolites.] Shells ovoid, trumpet-shaped;

FIG. 670.—Cyrtionella pileolus.

by the rapid expansion. Type C.

mitella, Hall, 1862, (Cyrtolites mitella,) 15th Rep. N. Y. Mus. Nat. Hist., p. 61, and Pal. N. Y., vol. 5, pt. 2, p. 123, Ham. Gr.

pileolus, Hall, 1862, (Cyrtolites pileolus,) 15th Rep. N. Y. Mus. Nat. Hist., p. 61, and Pal. N. Y., vol. 5, pt. 2, p. 125, Ham. Gr.

DAWSONELLA, Bradley, 1874, Am. Jour. Sci., 3d series, vol. 7, p. 151. [Ety. proper name.] Helicoid, having a thin plate

attached to the columella, covering half or more than half of the aperture of the shell as in Navicella. Type D. meeki.

meeki, Bradley, 1872, (Anomphalus meeki,) Am. Jour. Sci., 3d series, vol. 4, p. 88, Coal Meas.

DENTALIUM, Linnaeus, 1740, Syst. Nat., 2d Ed., p. 64. [Ety. dens, tooth.] Shell

elongate, terete, or angular, smooth, costate, or striate; aperture circular; lip simple, entire; margin of the posterior opening without a fissure. Type D. elephantinum.

aciculatum, see Coleolus aciculatus.

acutiusculatum, Gurley, 1883, New Carb. Foss., p. 7. Publication not valid.

annulostriatum, Meek & Worthen, 1870, Proc. Acad. Nat. Sci., p. 45, and Geo. Sur. Ill., vol. 5, p. 589, Coal Meas.

barquense, Winchell, 1862, Proc. Acad. Nat. Sci., p. 425, Marshall Gr.

canna, White, 1874, Rep. In-

vert. Foss., p. 23, and Geo. Sur. W. 100th Mer., vol. 4,

p. 156, Carb.

grandævum, Winchell, 1863, Proc. Acad. Nat. Sci., p. 18, Marshall Gr.

illinoisense, Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 325, and Geo. Sur. Ill., vol. 8,

p. 145, Kaskaskia Gr.

martini, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 203, Up. Held. Gr.

meekianum, Geinitz, 1866, Carb. und Dyas in Neb., p. 13, and Geo. Sur. Ill., vol. 5,

p. 590, Coal Meas.

missouriense, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 99, Kaskaskia Gr.

obsoletum, Hall. Preoccupied by Schlot-

heim in 1832. See D. sublaeve.

primarium, Hall, 1858, Geo. Rep. Iowa, p. 666, Warsaw Gr.

sublaeve, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 244, Coal Meas. Proposed instead of D. obsoletum, Hall, 1858, Geo. Sur. Iowa, which was preoccupied.

venustum, Meek & Worthen, 1861, Proc. Acad. Nat. Sci., p. 145, and Geo. Sur. Ill., vol. 2, p. 284, St. Louis Gr.

Discolites, Emmons, syn. for Cyclora.

minutus, see Cyclora minuta.

ECCOLIOMPHALUS, Portlock, 1843, Geol. Rep. Lond., p. 411. [Ety. eculiophalus, un-

rolled umbilicus.] Shell discoid, a few

FIG. 671.—Dentalium elephantinum.

olites mitella,
at. Hist., p. 61,
pt. 2, p. 123,
olites pilcolus,
at. Hist., p. 61,
pt. 2, p. 125,
Am. Jour. Sci.,
[Ety. proper
g a thin plate
a, covering half
aperture of the
type D. meeki.
(Anomphalus
3d series, vol.

Syst. Nat., 2d
tooth.] Shell
gular, smooth,
rture circular;
gin of the pos-
fissure. Type

cicu-
883,
Pub-

Vor-
Nat.
Sur.
leas.
862,
425,

In FIG. 671.—
Geo. Denta-
lum ele-
phanti-
um.

863,
8, Marshall Gr.
Geo. Sur. Ill.,
Sur. Ill., vol. 8,

Ann. N. Y.
Up. Held. Gr.
Carb. und Dyas
Sur. Ill., vol. 5,

863, Trans. St.
p. 99, Kaskas-

ied by Schlott-
bleve.

eo. Rep. Iowa,

Ed. Am. Pal.
Proposed in-
fall, 1858, Geo.
reoccupied.
ien, 1861, Proc.
and Geo. Sur.
uis Gr.
Cyclora.

a.
843, Geol. Rep.
liomphalus,
discoid, a few

tapering, widely disconnected whorls;
upper surface usually flattened in one
plane, or slightly elevated; lower sur-
face of whorls round; no chambers.
Type *E. bucklandi*.
atlanticus, Billings, 1865, Pal. Foss., vol.
1, p. 250, Quebec Gr.
canadensis, Billings, 1861, Can. Nat. and
Geol., vol. 6, p. 320, Quebec Gr.
circinatus, Whiteaves, 1884, Pal. Foss.,
vol. 3, p. 35, Guelph Gr.
comes, Hall, 1876, Illust. Devon. Foss.,
pl. 16, Ham. Gr.
devonicus, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 187, Devonian.

FIG. 672.—*Eocyclomphalus distans*.

distans, Billings, 1865, Pal. Foss., vol. 1,
p. 249, Quebec Gr.
eboracensis, Hall, 1861, (Euomphalus
eboracensis,) 15th Rep. N. Y. Mus. Nat.
Hist., p. 55, and Pal. N. Y., vol. 5, pt.
2, p. 61, Ham. Gr.
gyroceras, Roemer, 1852, (Euomphalus
gyroceras,) Kreid. von Texas, p. 91,
Silurian.
intortus, Billings, 1861, Can. Nat. and
Geol., vol. 6, p. 321, Quebec Gr.
laxus, Hall, 1861, (Euomphalus laxus,) 15th
Rep. N. Y. Mus. Nat. Hist., p. 54,
and Pal. N. Y., vol. 5, pt. 2, p. 60, Up.
Held. Gr.
paradoxus, Winchell, 1863, (Phanerotinus
paradoxus,) Proc. Acad. Nat. Sci., p. 21,
and Pal. N. Y., vol. 5, pt. 2, p. 60, Mar-
shall Gr.
priscus, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 46, Calciferous Gr.
spiralis, Billings, 1861, Can. Nat. and
Geol., vol. 6, p. 321, Quebec Gr.
superbus, Billings, 1865, Pal. Foss., vol.
1, p. 250, Quebec Gr.
undulatus, Hall, 1861, Geo. Rep. Wis., p.
37, Trenton Gr.
volutatus, Whitfield, 1886, Bull. Am. Mus.
Nat. Hist., vol. 1, p. 314, Birdseye Gr.
EOTROCHUS, Whitfield, 1882, Bull. Am. Mus.
Nat. Hist., p. 77. [Ety. eos, dawn;

Trochus, a genus.] Conical above, flat
or concave beneath, and broadly and
deeply umbilicated; aperture very ob-
lique, and the outer angle of volutions
strongly carinated; surface ornamenta-
tion unlike on the upper and lower
parts. Type *E. concavus*.
concavus, Hall, 1858,
(Pleurotomaria con-
cava,) Trans. Alb.
Inst., vol. 4, p. 24,
and Bull. Am. Mus.
Nat. Hist., p. 78,
Warsaw Gr.

Eulima, Risso, 1826, His-
toire Naturelle des
Principales, p. 123. Not an American
Paleozoic genus.

peracuta, see Polyphemopsis peracuta.

EUNEMA, Salter, 1859, Can. Org. Rem., Dec-
ade 1, p. 24. [Ety. eu, beautiful; nema,
line.] Turbinate, thin; few angular
whorls, strong concentric ridges, crossed
by sinuate or oblique lines of growth;
inner lip not reflected; peritreme
simple; mouth rather effuse below;
no umbilicus. Type *E. strigillatum*.

FIG. 674.—*Eunema*
cerithioides.

cerithioides, Salter, 1859,
Can. Org. Rem., Dec-
ade 1, p. 30, Black
Riv. Gr.
erigone, Billings, 1862,
Pal. Foss., vol. 1, p.
35, Black Riv. Gr.
pagoda, Salter, 1859,
Can. Org. Rem., Dec-
ade 1, p. 30, Black
Riv. Gr.

priscum, Billings, 1859, Can. Nat. and
Geo. vol. 4, p. 360, Calciferous Gr.
salteri, see Orthonema salteri.
strigillatum, Salter, 1859, Can. Org. Rem.,
Decade 1, p. 29, Black Riv. Gr.
trilineatum, Hall, 1867, 20th Rep. N. Y.
Mus. Nat. Hist., p. 397, Niagara Gr.

EUOMPHALUS, Sowerby, 1812, Min. Conch.,
vol. 1, p. 97. [Ety. eu, wide; omphalos,
umbilicus.] Shell
discoid, spire flat-
tened; whorls num-
erous, angu-
lated; umbilicus
very wide, expos-
ing volutions;
mouth nearly cir-
cular; peritreme
entire, not in-
dented by the
preceding whorl;
surface usually
sculptured. Type
E. pentangulatus.

FIG. 675.—*Euomphalus*
pentangulatus.

ammon, White & Whitfield, 1862, Proc.
Bost. Soc. Nat. Hist., vol. 8, p. 301, Kin-
derhook Gr.
boonensis, Swallow, 1863, Trans. St. Louis
Acad. Sci., vol. 2, p. 99, Burlington Gr.
calciferus, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 47, Calciferous Gr.

- catilloides, Conrad, 1842, (Inachus catilloides,) Jour. Acad. Nat. Sci., vol. 8, p. 273, Coal Meas.
- circumliratus, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 308, Birds-eye Gr.
- clymenioides, see Straparollus clymenioides.
- comes, Hall, syn. for Phanerotinus laxus.
- conradi, syn. for Pleuronotus decewi.
- cyclostomus, see Straparollus cyclostomus.
- decewi, see Pleuronotus decewi.
- decollatus, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 244, Low. Held. Gr. Proposed instead of E. disjunctus, Hall, 1859, Pal. N. Y., vol. 3, p. 340.
- depressus, Hall, 1843. Preoccupied by Goldfuss in 1832. See Straparollus hecale.
- disjunctus, Hall. Preoccupied by Goldfuss. See E. decollatus.
- eboracensis, see Eecyliomphalus eboracensis.
- exortivus, Dawson, 1868, Acad. Geol., p. 308, Carboniferous.
- expansus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 273, Niagara Gr.
- gyroceras, see Eecyliomphalus gyroceras.
- hecale, see Straparollus hecale.
- hecale var. corpulens, see Straparollus hecale var. corpulens.
- hemispherica, see platystoma hemisphericum.
- inops, see Straparollus inops.
- latus, Hall, 1858, Geo. Rep. Iowa, p. 605, Burlington Gr.
- laxus, see Eecyliomphalus laxus.
- lens, see Straparollus lens.
- luxus, White, 1875, Expl. W. 100th Meridian, vol. 4, p. 94, Subcarboniferous.
- macrolineatus, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 82, and Geo. Sur. Wis., vol. 4, p. 294, Niagara Gr.
- minnesotensis, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 581, Calciferous Gr.
- minutissimus, Castelnau, 1843, Syst. Sil., p. 35. Not recognized.
- obtusus, Hall, 1858, Geo. Rep. Iowa, p. 523, Kinderhook Gr.
- ophirensis, see Straparollus ophirensis.
- pernodosus, Meek & Worthen, 1870, (Straparollus pernodosus,) Proc. Acad. Nat. Sci., p. 45, and Geo. Sur. Ill., vol. 5, p. 604, Coal Meas.
- perspectivus, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 98, Kaskaskia Gr.
- pervetus, Conrad, 1843, (Inachus pervetus,) Proc. Acad. Nat. Sci., vol. 1, p. 334, Trenton Gr.
- planidorsatus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 462, and Geo. Sur. Ill., vol. 2, p. 302, Kaskaskia Gr.
- planispira, see Straparollus planispiratus.
- planodiscus, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 109, and Pal. N. Y., vol. 5, pt. 2, p. 57, F. m. Gr.
- polygyratus, Roemer, 1852, Kreidl. von Texas, p. 91, Silurian.
- profundus, see Bucania profunda.
- quadrivolvus, see Straparollus quadrivolvus.
- roberti, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 22, Burlington Gr.
- rotuliformis, Meek, 1870, Proc. Acad. Nat. Sci., p. 61, Calciferous Gr.
- rotundus, see Pleurotomaria rotunda.
- rudis, see Straparollus rudis.
- rugilineatus, see Cyclonema rugilineatum.
- rugosus, Hall, 1858, Geo. Sur. Iowa, p. 722. Preoccupied by Sowerby in 1812. See E. subrugosus.
- sanctisabae, see Straparollus sanctisabae.
- sinuatus, see Straparollus sinuatus.
- spergenensis, see Straparollus spergenensis.
- spergenensis var. planorbiformis, see Straparollus spergenensis var. planorbiformis.
- spirorbis, see Straparollus spirorbis.
- springvalensis, White, 1876, Proc. Acad. Nat. Sci., p. 32, and Cont. to Pal., No. 8, p. 167, Kinderhook Gr.
- strongi, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 66, and Geo. Wis., vol. 4, p. 200, Lower Magnesian Gr.
- subplanus, see Straparollus subplanus.
- subquadratus, Meek & Worthen, 1870, (Straparollus subquadratus,) Proc. Acad. Nat. Sci., p. 46, and Geo. Sur. Ill., vol. 5, p. 605, Up. Coal Meas.
- subrugosus, Meek & Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 607, Coal Meas. Proposed instead of E. rugosus, Hall, which was preoccupied.
- sulcatus, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 138, Onondaga Gr.
- tioga, Hall, 1876, Illust. Devonian Foss., pl. 15, and Pal. N. Y., vol. 5, pt. 2, p. 56, Chemung Gr.
- triliratus, Conrad, 1843, Proc. Acad. Nat. Sci., p. 333, Trenton Gr.
- trochiscus, see Raphistoma trochiscum.
- umbilicatus, see Straparollus umbilicatus.
- uniangulatus, see Ophileta uniangulata.
- utahensis, see Straparollus utahensis.
- vaticinus, Hall, 1863, 16th Rep. N. Y. Mus. Nat. Hist., p. 136, Potsdam Gr.
- verneuili, Castelnau, 1843, Syst. Sil., p. 34. Not recognized.
- whitneyi, see Omphalotrochus whitneyi.
- FUSISPIRA, Hall, 1871, 24th Rep. N. Y. Mus. Nat. Hist., p. 229. [Ety. *fusus*, spindle; *spira*, spire.] Fusiform, imperforate; spire elevated, with rounded volutions; aperture elongate-ovate or elliptical, produced below, forming a subrimate canal; columella slightly twisted, without folds; peristome sharp. Type F. ventricosa.
- compacta, Hall & Whitfield, 1877, U. S. Expl. 40th Parallel, vol. 4, p. 236, Quebec Gr.
- elongata, Hall, 1871, 24th Rep. N. Y. Mus. Nat. Hist., p. 229, Trenton Gr.
- subfusiformis, Hall, 1847, (Murchisonia subfusiforme,) Pal. N. Y., vol. 1, p. 180, Trenton and Hud. Riv. Grs.

FIG. 676.—*Fusispira ventricosa*.

terebiformis, Hall, 1871, 24th Rep. N. Y. Mus. Nat. Hist., p. 230, Hud. Riv. Gr.
ventricosa, Hall, 1871, 24th Rep. N. Y. Mus. Nat. Hist., p. 229, Trenton Gr.
vittata, Hall, 1847, (Murchisonia *vittata*,) Pal. N. Y., vol. 1, p. 181, Trenton Gr.
Fusus, Bruguiere, 1789, Encyc. Meth. This genus is unknown in the Palaeozoic rocks.
inhabilis, syn.

for *Macrochilina primigenia*.

HELICOTOMA, Salter, 1859, Can. Org. Rem., Decade 1, p. 13. [Ety. *Helix*, genus of shells; *tome*, notch.] Depressed discoid, spire nearly flat, whorls obtusely angular externally, rounded below; umbilicus broad; form helicoid. Type H. *planulata*.

declivis, Safford, 1869, Geo. of Tenn. Not defined.

eucharis, Billings, 1865, Pal. Foss., vol. 1, p. 249, Quebec Gr.

gorgonea, Billings, 1865, Pal. Foss., vol. 1, p. 248, Quebec Gr.

larvata, Salter, 1859, Can. Org. Rem., Decade 1, p. 15, Black Riv. and Trenton Gr.

misera, Billings, 1865, Pal. Foss., vol. 1, p. 309, Quebec Gr.

muricata, Salter, 1859, (H. *planulata* var. *muricata*,) Can. Org. Rem., Decade 1, p. 14, Black Riv. and Trenton Gr.

naresi, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 602, Up. Sil.

perstriata, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 356, Calciferous Gr.

planulata, Salter, 1859, Can. Org. Rem., Decade 1, p. 14, Black Riv. and Trenton Gr.

proserpina, Billings, 1865, Pal. Foss., vol. 1, p. 247, Quebec Gr.

serotina, Nicholson, 1874, Rep. Pal. Ont., p. 120, Up. Held. Gr.

spinosa, Salter, 1859, Can. Org. Rem., Decade 1, p. 15, Black Riv. Gr.

tennesseensis, Safford, 1869, Geo. of Tenn. Not defined.

tritonia, Billings, 1865, Pal. Foss., vol. 1, p. 247, Quebec Gr.

HOLOPEA, Hall, 1847, Pal. N. Y., vol. 1, p. 169. [Ety. *holos*, entire; *ope*, aperture.] Shell conical, ventricose, more or less oblique, or nearly direct; aperture round, ovate; margin entire; surface marked by fine curved striae or cancellated; distinguished from *Cyclonema* by the presence of an umbilicus. Type H. *symmetrica* and H. *obliqua*.

antiqua, Vanuxem, 1843, (Littorina *antiqua*,) Geo. Rep. 3d Dist. N. Y., p. 112, and Pal. N. Y., vol. 3, p. 294, Low. Held. Gr.

antiqua var. *pervetusta*, Hall, 1859, Pal. N. Y., vol. 3, p. 295, Low. Held. Gr.

cassina, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., p. 310, Birdseye Gr.

chicagoensis, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 99, Niagara Gr.

conica, Winchell, 1863, Proc. Acad. Nat. Sci., p. 21, Marshall Gr.

danai, Hall, 1859, Pal. N. Y., vol. 3, p. 295, Low. Held. Gr.

dilucula, Hall, 1847, (Turbo *dilucula*,) Pal. N. Y., vol. 1, p. 12, Calciferous Gr.

(?) *elongata*, Hall, 1859, Pal. N. Y., vol. 3, p. 295, Low. Held. Gr.

eriansis, Nicholson, 1874, Rep. Pal. Ont., p. 120, Up. Held. Gr.

gracia, Billings, 1862, Pal. Foss., vol. 1, p. 159, Guelph Gr.

guelphensis, Billings, 1862, Pal. Foss., vol. 1, p. 159, Guelph Gr.

harmonia, Billings, 1862, Pal. Foss., vol. 1, p. 158, Guelph Gr.

lavinia, Billings, 1862, Pal. Foss., vol. 1, p. 28, Trenton Gr.

leiosoma, Billings, 1865, Pal. Foss., vol. 1, p. 187, Quebec Gr.

magniventra, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 83, and Geo. Wis., vol. 4, p. 316, Niagara Gr.

nana, Meek, 1871, Proc. Acad. Nat. Sci., p. 172, syn. for *Cyclora minuta*.

neris, Billings, 1862, Pal. Foss., vol. 1, p. 27, Trenton and Black Riv. Grs.

newtonensis, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 224, Kaskaskia Gr.

niagarensis, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 99, Niagara Gr.

obesa, Whitfield, 1882, Geo. Wis., vol. 4, p. 348, Low. Magnesian Gr.

obliqua, Hall, 1847, Pal. N. Y., vol. 1, p. 170, Trenton and Hud. Riv. Gr.

obscura, Hall, 1847, (Turbo *obscura*,) Pal. N. Y., vol. 1, p. 12, Calciferous Gr.

occidentalis, Nicholson, 1875, Quar. Jour. Geo. Soc. Lond., vol. 31, p. 550, Guelph Gr.

ophelia, Billings, 1865, Pal. Foss., vol. 1, p. 222, Quebec Gr.

FIG. 678.
Holopea dilucula. FIG. 677.—*Helicotoma eucharis*.

- ovalis, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 351, Calciferous Gr.
 paludiformis, Hall, 1847, Pal. N. Y., vol. 1, p. 171, Trenton Gr.
 prosperina, Billings, 1862, Pal. Foss., vol. 1, p. 28, Calciferous and Chazy Grs.
 proutana, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 30, and Bull. Am. Mus. Nat. Hist., p. 72, Warsaw Gr.
 pyrene, Billings, 1862, Pal. Foss., vol. 1, p. 27, Black Riv. Gr.
 reversa, Hall, 1860, Can. Nat. and Geol., vol. 5, p. 154, Up. Silurian.
 subconica, Hall, 1859, Pal. N. Y., vol. 3, p. 294, Low. Held. Gr.
 subconica, Winchell, 1863, Proc. Acad. Nat. Sci., p. 21. This name was preoccupied.
 sweeti, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis. and Geo. Wis., vol. 4, p. 174, Potsdam Gr.
 symmetrica, Hall, 1847, Pal. N. Y., vol. 1, p. 170, Black Riv. Gr.
 turgida, Hall, 1847, (Pleurotomaria turgida), Pal. N. Y., vol. 1, p. 12, Calciferous Gr.
 ventricosa, Hall, 1847, Pal. N. Y., vol. 1, p. 171, Trenton Gr.
 HOLOPELLA, McCoy, 1855, Brit. Pal. Foss., p. 303. [Ety. diminutive of *Holopea*.] Shell spiral, elongate, slender, of numerous gradually increasing whorls, generally crossed by slightly arched striae; mouth circular, with the peristome entire; base rounded, with or without a minute umbilicus. Type *H. cancellata*.
 mira, Winchell, 1863, Proc. Acad. Nat. Sci., p. 22, Marshall Gr.
Inachus catilloides, see *Euomphalus catilloides*.
pervetus, see *Euomphalus pervetus*.
pervetus, see *Pleurotomaria pervetusta*.
undatus, see *Lituites undatus*.
 ISONEMA, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 251. [Ety. *isos*, equal; *nema*, thread.] Depressed turbinate, subglobose, obtusely angular around the middle of the body whorl; aperture subrhombic; outer lip thin, entire; inner lip a little flattened in the umbilical region; surface ornamented with transverse, very regular lines on the upper side of the volutions. Type *I. depressum*.
 bellatulum, see *Callonema bellatulum*.
 depressum, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 251, and Geo. Sur. Ill., vol. 3, p. 443, Ham. Gr.
 humile, Meek, 1871, Proc. Acad. Nat. Sci., p. 79, and Ohio Pal., vol. 1, p. 214, Up. Held. Gr.
 lichas, see *Callonema lichas*.
 LEPETOPSIS, Whitfield, 1882, Bull. Am. Mus. Nat. Hist., No. 3, p. 67. [Ety. *Lepeta*, a

FIG. 679.—*Isonema depressum*.

genus; *opsis*, resemblance.] Shell patelliform, more or less regularly round or oval, apex subcentral, posterior to the middle, directed backward, the nucleus dextrally coiled; muscular imprint horseshoe-shaped, open in front, consisting of an irregular narrow band, which expands more or less at the anterior extremities; surface with six radiating lines, two anterior, two posterior, and two lateral.

Type *L. levettii*.

chesterensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 25, and Geo. Sur. Ill., vol. 8, p. 140, Kaskaskia Gr.

levettii, White, 1882, (Patella levettii), 11th Rep. Geo. of Indiana, p. 359, Warsaw Gr.

Littorina, Ferussac, 1821, Tab. Syst. An. Mollusques, etc.

antiqua, see *Holopea antiqua*.

cancellata, see *Cyclonema cancellatum*.

wheeleri, see *Naticopsis wheeleri*.

LOPHOSPIRA, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 312. [Ety. *lophos*, the keel; *spira*, a whorl.] Spire elevated, strongly keeled, and axis minutely perforate, when whorls are not disconnected. The types are *Murchisonia milleri*, Hall, and *M. helictæres*; but as the generic characters are not very satisfactorily or clearly defined, I leave the species with *Murchisonia*.

cassina, see *Murchisonia cassina*.

calcifera, see *Murchisonia calcifera*.

LOXONEMA, Phillips, 1841, Pal. Foss., Cornwall, etc., p. 98. [Ety. *loxos*, oblique; *nema*, thread.] Shell elongate, many whorled; aperture simple, attenuate above, effuse below; lines of growth sigmoidal; no umbilicus. Type *L. sinuosum*.

aculeatum, Billings, 1866, Catal. Sil. Foss. Antic., p. 55, Anticosti Gr.

acutulum, Dawson, 1868, Acad. Geol., p. 309, Carboniferous.

approximatum, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 191, Devonian.

attenuatum, Stevens, 1858, (Chemnitzia attenuata), Am. Jour. Sci. and Arts, 2d ser., vol. 25, p. 259, Coal Meas.

attenuatum, Hall, 1859, Pal. N. Y., vol. 3, p. 296. The name was preoccupied.

See *L. emaceratum*.

attenuatum var. *semicostatum*, see *L. semicostatum*.

bellatulum, see *Isonema bellatulum*.

bellona, Hall, 1876, Illust. Devonian Foss., pl. 14, and Pal. N. Y., vol. 5, pt. 2, p. 46, Ham. Gr.

bellum, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 258, Subcarboniferous.

boydi, see *Murchisonia boydi*.

breviculum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 132, Ham. Gr.

FIG. 680.—*Leptopsis levettii*.

Shell particularly round posterior to forward, the muscular impression in front, narrow band, as at the anterior with six radiating, two posterior.

FIG. 680.—*Lopeopsis levettii*.

b. Syst. An.

a. acellatum.

cleri.

ull. Am. Mus.

[Ety. *lophos*,

] Spire ele-

nd axis mi-

horls are not

are Murchi-

M. helictares;

acters are not

ly defined, I

rchisonia.

isina.

leiferia.

l. Foss., Corn-

oxos, oblique;

ngate, many

ble, attenuate

nes of growth

Type L. sinu-

atal. Sil. Foss.

cad. Geol., p.

1885, Monogr.

191, Devonian.

(Chemnitzia

and Arts, 2d

Meas.

N. Y., vol. 3,

preoccupied.

, see L. semi-

latulum.

evonian Foss.,

5, pt. 2, p. 46,

ogr. U. S. Geo.

arboniferous.

di.

N. Y., vol. 5,

cara, Dawson, 1883, Rep. on Redpath Museum, No. 2, p. 11, Subcarboniferous. carinatum, see *Macrochilina carinata*.

cerithiiforme, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 465, and Geo. Sur. Ill., vol. 2, p. 379, Up. Coal Meas.

coaptum, Hall, 1876, Illust. Devonian Foss., pl. 13, and Pal. N. Y., vol. 5, pt. 2, p. 44, Ham. Gr.

compactum, Hall, 1859, Pal. N. Y., vol. 3, p. 297, Low. Held. Gr.

cotteranum, Billings, 1861, Can. Jour., vol. 6, p. 360, Corniferous limestone.

crassum, Webster, 1888, Am. Nat., p. 446. Not defined so as to be recognized.

danvillense, Stevens, 1858, Am. Jour. Sci., vol. 25, p. 259, Coal Meas.

delphicola, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 52, and Pal. N. Y., vol. 5, pt. 2, p. 47, Ham. Gr.

emaceratum, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 244, Low. Held. Gr. Proposed instead of *L. attenuatum*, Hall, 1859, in Pal. N. Y., vol. 3, p. 296, which was preoccupied.

eurekense, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 190, Devonian.

fasciatum, King, 1850, Permian Foss., p. 209, Permian Gr.

fitchi, Hall, 1859, Pal. N. Y., vol. 3, p. 296, Low. Held. Gr.

gigantea, Webster, 1888, Am. Nat., p. 445. Not properly defined.

halli, Norwood & Pratten, 1855, Jour. Acad. Nat. Sci., 2d series, vol. 3, p. 77, Coal Meas.

hamiltoniae, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 33, and Pal. N. Y., vol. 5, pt. 2, p. 45, Ham. Gr.

hydraulicum, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 193, and Pal. N. Y., vol. 5, pt. 2, p. 44, Ham. Gr.

inornata, see *Polyphemopsis inornata*.

kanii, Meek, 1865, Am. Jour. Sci. and Arts, 2d ser., vol. 40, p. 33, Low. Held. Gr.

leviusculum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 131, Ham. Gr.

laxum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 49, Chemung Gr.

leda, Hall, 1868, 20th Rep. N. Y. Mus. Nat. Hist., p. 398, Niagara Gr.

magnum, Whitfield, 1878, Ann. R. p. Geo. Sur. Wis., p. 83, and Geo. Wis., vol. 4, p. 317, Niagara Gr.

macclintochi, Haughton, 1857, Jour. Roy. Dub. Soc., vol. 1, Devonian.

minutum, Stevens, 1858, Am. Jour. Sci., 2d series, vol. 25, p. 260, Coal Meas.

moloch, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 30, Genesee shales.

multicostatum, Meek & Worthen, 1861, Proc. Acad. Nat. Sci., p. 123, and Geo. Sur. Ill., vol. 2, p. 378, Coal Meas.

murrayanum, Salter, 1859, Can. Org. Rem., Decade 1, p. 31, Black Riv. Gr.

newberryi, see *Soleniscus newberryi*.

nerile, Sowerby. Not an American species.

nitidula, see *Polyphemopsis nitidula*.

nobile, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 190, Devonian.

nodosum, Stevens, 1858, Am. Jour. Sci., 2d ser., vol. 25, p. 260, Coal Meas.

noe, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 55, Portage Gr.

obtusum, Hall, 1859, Pal. N. Y., vol. 3, p. 297, Low. Held. Gr.

oligospiratum, Winchell, 1863, Proc. Acad. Nat. Sci., p. 22, Marshall Gr.

owenense, Webster, 1888, Am. Nat., p. 446. Not defined so as to be recognized.

parvum, Cox, 1857, (Chemnitzia parva,) Geo. Sur. Ky., vol. 3, p. 567, Coal Meas.

parvulum, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 204, Up. Held. Gr.

peoriense, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 7, and Geo. Sur. Ill., vol. 8, p. 139, Coal Meas.

pexatum, Hall, 1861, 14th Rep. N. Y. Mus. Nat. Hist., p. 104, and Pal. N. Y., vol. 5, pt. 2, p. 42, Up. Held. Gr.

pexatum var. obsoletum, Hall, 1876, Illust. Devonian Foss., pl. 13, and Pal. N. Y., vol. 5, pt. 2, p. 43, Up. Held. Gr.

planogyratum, Hall, 1839, Pal. N. Y., vol. 3, p. 298, Low. Held. Gr.

plicatum, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 231, Coal Meas.

politum, Stevens, 1858, Am. Jour. Sci., 2d series, vol. 25, p. 260, Coal Meas.

postrenum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 132, Chemung Gr.

quadricarinatum, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 7, and Geo. Sur. Ill., vol. 8, p. 140, Coal Meas.

rectistriatum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 130, Ham. Gr.

regulare, Cox, 1857, Geo. Sur. Ky., vol. 3, p. 566, Coal Meas.

robustum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 52, and Pal. N. Y., vol. 5, pt. 2, p. 40, Schoharie grit.

rossi, Haughton, 1857, Jour. Roy. Soc. Dub., vol. 1, Devonian.

rugosum, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 465, and Geo. Sur. Ill., vol. 2, p. 378, Up. Coal Meas.

scitulum, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 464, and Geo. Sur. Ill., vol. 2, p. 372, Low. Coal Meas.

semicostatum, Meek, 1871, (L. attenuatum var. semicostatum,) Proc. Acad. Nat. Sci., p. 174, and Geo. Sur. Ill., vol. 5, p. 596, Coal Meas.

sicula, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 43, Up. Held. Gr.

solidum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 51, and Pal. N. Y., vol. 5, pt. 2, p. 41, Schoharie grit.

styliola, Hall, 1876, Illust. Devon. Foss., pl. 14, and Pal. N. Y., vol. 5, pt. 2, p. 48, Chemung Gr.

- subattenuatum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 52, and Pal. N. Y., vol. 5, pt. 2, p. 40, Scholharie grit.
- subulata*, see Murchisonia subulata.
- swallowanum, Shumard, 1859, (Chemnitzia swallowana,) Trans. St. Louis Acad. Sci., vol. 1, p. 399, Permian.
- tennicarinatum, Stevens, 1858, Am. Jour. Sci., 2d series, vol. 25, p. 260, Coal Meas.
- tenuilineatum, Shumard, 1855, (Chemnitzia tenuilineata,) Geo. Rep. Mo., p. 207, Waverly Gr. or Choteau limestone.
- terobra*, Hall, 1876, Illust. Devon. Foss., pl. 14, and Pal. N. Y., vol. 5, pt. 2, p. 48, Chemung Gr.
- teres*, Hall, 1876, Illust., Devonian Foss., pl. 13, and Pal. N. Y., vol. 5, pt. 2, p. 42, Corniferous Gr.
- turritiforme, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 109, Kinderhook Gr.
- vincta*, see Murchisonia vincta.
- yandellianum, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 28, and Bull. Am. Mus. Nat. Hist., p. 77, Warsaw Gr.
- MACLUREA, LeSueur, 1818, (Maclurites,) Jour. Acad. Nat. Sci., vol. 1, p. 312. [Ety. proper name.] Discoidal, few whorled, reversed, upper surface convex, deeply perforate, outer side spirally grooved; operculum sinistrally subspiral, solid, with two internal projections for the attachment of muscles. Type *M. magna*.

FIG. 682.—*Loxonema yandellianum*.

FIG. 683.—*Maclurea crenulata*.

- acuminata*, Billings, 1865, Pal. Foss., vol. 1, p. 240, Quebec Gr.
- affinis*, Billings, 1865, Pal. Foss., vol. 1, p. 238, Quebec Gr.
- annulata*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 81, Chazy Gr.
- atlantica*, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 459, Chazy Gr.
- bigshbyi*, Hall, 1861, Geo. Rep. Wis., p. 37, and Geo. Wis., vol. 4, p. 222, Trenton Gr.
- carinata*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 82, Trenton Gr.
- crenulata*, Billings, 1865, Pal. Foss., vol. 1, p. 236, Quebec Gr.
- cuneata*, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 75, and Geo. Wis., vol. 4, p. 246, Trenton Gr.

- emmonsi*, Billings, 1865, Pal. Foss., vol. 1, p. 242, Quebec Gr.
- labiata*, see *Raphistoma labiata*.
- logani*, Salter, 1851, Rep. British Assoc., p. 63, Black Riv. Gr.
- magna*, LeSueur, 1818, Jour. Acad. Nat. Sci., vol. 1, p. 312, and Pal. N. Y., vol. 1, p. 26, Chazy Gr.
- matutina*, Hall, 1847, Pal. N. Y., vol. 1, p. 10, Calciferous Gr.
- minima*, Hall & Whitfield, 1877, U. S. Geo. Expl., 40th parallel, vol. 4, p. 235, Chazy Gr.
- oceanica*, Billings, 1865, Pal. Foss., vol. 1, p. 237, Quebec Gr.
- ponderosa*, Billings, 1865, Pal. Foss., vol. 1, p. 239, Quebec Gr.
- psyche*, Billings, 1865, Pal. Foss., vol. 1, p. 244, Quebec Gr.
- rotundata*, Billings, 1865, Pal. Foss., vol. 1, p. 245, Quebec Gr.
- speciosa*, Billings, 1865, Pal. Foss., vol. 1, p. 240, Quebec Gr.
- sordida*, Hall, 1847, Pal. N. Y., vol. 1, p. 10, Calciferous Gr.
- striata*, see *Scalites striatus*.
- striata*, Troost, 1840. Not defined.
- subannulata*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 82, Trenton Gr.
- subrotunda*, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 75, and Geo. Wis., vol. 4, p. 246, Trenton Gr.
- sylpha*, Billings, 1865, Pal. Foss., vol. 1, p. 244, Quebec Gr.
- transitionis*, Billings, 1865, Pal. Foss., vol. 1, p. 241, Quebec Gr.

wadsworthi, Whitfield, 1884, Bull. Am. Mus. Nat. Hist., vol. 1, p. 139, Up. Taconic.

Macrocheilus, Phillips, 1841, Pal. Foss., Cornwall, etc., p. 103. This name was preoccupied by Hope, in 1838, for a genus of Coleopterous insects. Bayle has

- proposed *Macrochilina*, to which all the species are referred.
- altonense*, see *Macrochilina worthenanus*.
- altonense*, see *Macrochilina altonensis*.
- anguliferum*, see *Macrochilina angulifera*.
- attenuatum*, Hall, syn. for *Soleniscus fusiformis*.
- cooperense*, see *Macrochilina cooperensis*.
- fusiforme*, see *Soleniscus fusiformis*.
- gracile*, see *Macrochilina gracilis*.
- hallanum*, see *Soleniscus hallanum*.
- hamiltoniae*, see *Macrochilina hamiltoniae*.
- hebe*, see *Macrochilina hebe*.
- hildrethi*, see *Macrochilina hildrethi*.
- humile*, see *Macrochilina humilis*.
- inhabile*, syn. for *Macrochilina primitivus*.
- intercalare*, see *Macrochilina intercalaris*.

- kansasense*, see *Macrochilina kansasensis*.
klipparti, see *Soleniscus klipparti*.
macrostomum, see *Macrochilina macrostomus*.
mediale, see *Macrochilina medialis*.
missouriense, see *Macrochilina missouriensis*.
newberryi, see *Soleniscus newberryi*.
paludinaformis, see *Soleniscus paludini-formis*.
pingue, see *Macrochilina pinguis*.
ponderosum, see *Macrochilina ponderosus*.
primævum, see *Macrochilina primævus*.
primigenium, see *Macrochilina primigenius*.
priscum, see *Macrochilina prisca*.
pulchellum, syn. for *Macrochilina intercalaris*.
spiratum, see *Macrochilina spirata*.
subcorpulentum, see *Macrochilina subcorpulenta*.
terranovicum, see *Macrochilina terranovica*.
texanum, see *Soleniscus texanus*.
ventricosum, see *Soleniscus ventricosus*.
MACROCHILINA, Bayle, 1880, *Journal de Conchyliologie*, 3me. ser., t. 19. Proposed instead of *Macrocheilus* of Phillips, which was preoccupied by Ifope. [Ety. diminutive of *Macrocheilus*.] Subglobose, elongate; apex pointed; whorls convex, smooth, last one large; aperture subovate; columella imperforate; outer lip thin, without notch or sinus; inner lip thin above and thickened below. Type *M. acuta*.
altonensis, Worthen, 1873, (*Macrocheilus altonense*), *Geo. Sur. Ill.*, vol. 5, p. 593, Coal Meas.
angulifera, White, 1874, (*Macrocheilus anguliferum*), *Rep. Invertebrate Foss.*, p. 22, and *Geo. Sur. 100th Mer.*, vol. 4, p. 160, Carboniferous.
carinata, Stevens, 1858, (*Loxonema carinatum*), *Am. Jour. Sci.*, vol. 25, p. 259, Coal Meas.
cooperensis, Swallow, 1863, (*Macrocheilus cooperense*), *Trans. St. Louis Acad. Sci.*, vol. 2, p. 100, Kaskaskia Gr.
gracilis, Cox, 1857, (*Macrocheilus gracile*), *Geo. Sur. Ky.*, vol. 3, p. 570, Coal Meas.
hamiltonia, Hall, 1862, (*Macrocheilus hamiltonia*), 15th Rep. N. Y. Mus. Nat. Hist., p. 49 and Pal. N. Y., vol. 5, pt. 2, p. 33, Ham. Gr.
hebe, Hall, 1862, (*Macrocheilus hebe*), 15th Rep. N. Y. Mus. Nat. Hist., p. 43, and Pal. N. Y., vol. 5, pt. 2, p. 32, Ham. Gr.
hildrethi, Conrad, 1842, (*Plectostylus hildrethi*), *Jour. Acad. Nat. Sci.*, vol. 8, p. 275, Coal Meas.
humilis, Keyes, 1888, (*Macrocheilus humile*), *Proc. Acad. Nat. Sci. Phil.* pl. xii, fig. 1, Coal Meas.
intercalaris, Meek & Worthen, 1860, (*Macrocheilus intercalare*), *Proc. Acad. Nat. Sci.*, p. 467, and *Geo. Sur. Ill.*, vol. 2, p. 371, Up. Coal Meas.

- kansasensis*, Swallow, 1858, (*Macrocheilus kansasense*), *Trans. St. Louis Acad. Sci.*, vol. 1, p. 201, Coal Meas.
littonana, Hall, 1858, (*Natica littonana*), *Trans. Alb. Inst.*, vol. 4, p. 30, and *Bull. Am. Mus. Nat. Hist.*, p. 72, Warsaw Gr.
macrostoma, Hall, 1862, (*Macrocheilus macrostomum*), 15th Rep. N. Y. Mus. Nat. Hist., p. 49, and Pal. N. Y., vol. 5, pt. 2, p. 33, Ham. Gr.
medialis, Meek & Worthen, 1860, (*Macrocheilus mediale*), *Proc. Acad. Nat. Sci.*, p. 466, and *Geo. Sur. Ill.*, vol. 2, p. 370, Up. Coal Meas.
missouriensis, Swallow, 1858, (*Macrocheilus missouriense*), *Trans. St. Louis Acad. Sci.*, vol. 1, p. 201, Coal Meas.
pinguis, Winchell, 1863, (*Macrocheilus pingue*), *Proc. Acad. Nat. Sci.*, p. 21, Marshall Gr.
ponderosa, Swallow, 1858, (*Macrocheilus ponderosum*), *Trans. St. Louis Acad. Sci.*, vol. 1, p. 202, Coal Meas.
primæva, Hall, 1876, (*Macrocheilus primævum*), *Illust. Devonian Foss.*, pl. 12, and Pal. N. Y., vol. 5, pt. 2, p. 35, Schoharie grit.
primigenia, Conrad, 1835, (*Stylifer primigenia*), *Trans. Geo. Soc. Penn.*, vol. 1, p. 267, Coal Meas.
prisca, Whitfield, 1882, (*Macrocheilus priscum*), *Ann. N. Y. Acad. Sci.*, vol. 2, p. 204, Up. Held. Gr.
spirata, McCoy, 1850, (*Macrocheilus spiratum*), *Brit. Pal. Rocks*, p. 549, Coal Meas.
subcorpulenta, Whitfield, 1882, (*Macrocheilus subcorpulentum*), *Ann. N. Y. Acad. Sci.*, vol. 2, p. 224, Kaskaskia Gr.
terranovica, Dawson, 1883, (*Macrocheilus terranovicum*), *Rep. on Redpath Museum*, No. 2, p. 14, Carboniferous.
worthenianus, n. sp. St. Louis Gr. Proposed instead of *Macrocheilus altonense* in *Geo. Sur. Ill.*, vol. 8, p. 143, which name was preoccupied.
METOPTOMA, Phillips, 1836, *Geo. of Yorkshire*, pt. 2, p. 223. [Ety. *metopon*, front; *tome*, incision.] Patelliform, truncated under the apex, at the posterior side; horseshoe-shaped muscular scar, with the open end directed from the truncated side. Type *M. oblonga*.
alceste, Billings, 1862, *Pal. Foss.*, vol. 1, p. 153, Hud. Riv. Gr.
alta, Whitfield, 1889, *Bull. Am. Mus. Nat. Hist.*, vol. 2, p. 44, Calciferous Gr.
analoga, Walcott, 1885, *Monogr. U. S. Geo. Sur.*, vol. 8, p. 84, Trenton Gr.
angusta, Billings, 1862, *Pal. Foss.*, vol. 1, p. 88, Quebec Gr.
anomala, Billings, 1862, *Pal. Foss.*, vol. 1, p. 89, Quebec Gr.

FIG. 684. — *Macrocheilus littonana*.

barabuensis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 80, and Geo. Wis., vol. 4, p. 195, Low. Magnesian Gr.
 billingsi, Walcott, 1883, 35th Rep. N. Y. Mus. Nat. Hist., p. 212, Trenton Gr.

FIG. 685. — *Metoptoma canadensis*. a, Upper side; b, side view; c, under side.

canadensis, Billings, 1865, Pal. Foss., vol. 1, p. 394, (*Chiton canadensis*), Black Riv. Gr.
 cornutiformis, Walcott, 1879, Desc. New Spec. Foss., p. 1, Calciferous Gr.
 devonica, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 195, Devonian.
 dubia, Hall, 1847, Pal. N. Y., vol. 1, p. 23, Chazy Gr.
 erato, see *Tryblidium erato*.
 estella, Billings, 1862, Pal. Foss., vol. 1, p. 153, Hud. Riv. Gr.
 eubule, see *Tryblidium eubule*.
 hyrie, see *Tryblidium hyrie*.
 instabilis, Billings, 1865, Pal. Foss., vol. 1, p. 251, Quebec Gr.
 melissa, Billings, 1862, Pal. Foss., vol. 1, p. 86, Quebec Gr.
 montrealensis, Billings, 1865, Pal. Foss., vol. 1, p. 394, Chazy Gr.
 niobe, see *Tryblidium niobe*.
 nycteis, see *Tryblidium nycteis*.
 orithyia, Billings, 1862, Pal. Foss., vol. 1, p. 38, Calcif. Gr.
 orphyne, Billings, 1862, Pal. Foss., vol. 1, p. 88, Quebec Gr.
 peroccidens, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 260, Subcarb.
 perovallis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 74, and Geo. Wis., vol. 4, p. 211, Trenton Gr.
 phillipsi, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 83, Trenton Gr.
 quebecensis, Billings, 1865, Pal. Foss., vol. 1, p. 308, Quebec Gr.
 recurva, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 61, and Geo. Wis., vol. 4, p. 196, Low. Mag. Gr.
 retrorsa, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 54, and Geo. Wis., vol. 4, p. 197, Low. Mag. Gr.
 rugosa, see *Stenotheca rugosa*.
 similis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 61, and Geo. Wis., vol. 4, p. 196, Low. Mag. Gr.
 simplex, see *Tryblidium simplex*.
 superba, Billings, 1865, Pal. Foss., vol. 1, p. 172, Black Riv. Gr.
 trentonensis, Billings, 1862, Pal. Foss., vol. 1, p. 40, Trenton Gr.
 undata, Winchell, 1865, Proc. Acad. Nat. Sci. Phil., p. 131, Kinderhook Gr.
 umbella, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 267, and Geo. Sur. Ill., vol. 3, p. 506, Burlington Gr.

venilia, Billings, 1862, Pal. Foss., vol. 1, p. 88, Quebec Gr.
 MICROCERAS, Hall, 1845, Am. Jour. Sci., vol. 48, p. 294. [Ety. *mikros*, small; *keras*, horn.] General form like *Cyrtolites*, but distinguished by its minute size, smooth surface, and less angular dorsal margin. Type *M. inornatum*.
 inornatum, Hall, 1845, Am. Jour. Sci., vol. 48, p. 294, and Ohio Pal., vol. 1, p. 147, Hud. Riv. Gr.
 minutissimum, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 13, Hud. Riv. Gr.
 MICRODOMA, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 269. [Ety. *mikros*, small; *domus*, house.] Shell small, subtrochiform; volutions seven or more, flattened on a line with the slope of the shell; suture deep; aperture oblique; surface ornamented with nodular ridges. Type *M. conica*.
 conica, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 269, and Geo. Sur. Ill., vol. 5, p. 598, Low. Coal Meas.
 MURCHISONIA, D'Archiac & Verneuil, 1841, Bull. Soc. Geo. Fr., vol. 12, p. 154, and Phillips Pal. Foss. Cornwall, etc., p. 101. [Ety. proper name.] Shell elongated, many whorled; whorls variously sculptured and zoned; outer lip deeply notched; aperture slightly channeled in front. Type *M. bilineata*.
 abbreviata, Hall, 1847, Pal. N. Y., vol. 1, p. 32. The name was preoccupied by DeKoninck in 1841. See *M. subabbreviata*.
 aciculata, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 154, Up. Silurian.
 acrea, Billings, 1865, Pal. Foss., vol. 1, p. 232, Quebec Gr.
 ada, Billings, 1865, Pal. Foss., vol. 1, p. 346, Calciferous Gr.
 adelina, Billings, 1865, Pal. Foss., vol. 1, p. 232, Quebec Gr.
 agilis, Billings, 1865, Pal. Foss., vol. 1, p. 235, Quebec Gr.
 alexandra, Billings, 1865, Pal. Foss., vol. 1, p. 172, Black Riv. Gr.
 angulata, Phillips, 1836, (*Rostellaria angulata*), Geo. of Yorkshire, p. 230, Devonian. Very doubtfully identified in America.
 angustata, Hall, 1847, Pal. N. Y., vol. 1, p. 41, Birdseye Gr.
 anna, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 358, Calciferous Gr.
 archimedeia, McChesney, 1861, Desc. New Pal. Foss., p. 89, Coal Meas.
 arenaria, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 359, Calciferous Gr.
 arisaigensis, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 154, Silurian.

FIG. 686. *Murchisonia bilineata*.

Foss., vol. 1,

our. Sci., vol.
small; *keras*,
e *Cyrtolites*,
minute size,
angular dorsal
num.our. Sci., vol.
vol. 1, p. 117,
p. 13, Hud.n, 1866, Proc.
Ety. *mitros*,
all small, sub-
n or more,
e slope of the
ure oblique;
ith nodular1866, Proc.
269, and Geo.
Low. Coalerneuil, 1841,
2, p. 154, and
ll, etc., p. 101.
ell elongated,
iously sculp-
lip deeply
channeledY., vol. 1, p.
occupied by
M. subabbre- FIG. 686.
Murchisonia
bilineata.Riv. Gr.
stellaria an-
e, p. 230, De-
identified in

Y., vol. 1, p.

at. and Geol.,

Gr.

Bl, Desc. New

as.

n. Nat. and

erous Gr.

n. Nat. and

an.

artemesia, Billings, 1865, Pal. Foss., vol.
1, p. 345, Calciferous Gr.aspera, Billings, 1859, Can. Nat. and Geo.,
vol. 4, p. 458, Chazy Gr.attenuata, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 27, and Bull. Am. Mus. Nat.

Hist., p. 88, Warsaw Gr.

angustina, Billings, 1865, Pal. Foss., vol.
1, p. 234, Quebec Gr.bellicineta, Hall, 1847, Pal. N. Y., vol. 1,
p. 179, Trenton and Hud. Riv. Grs.bicineta, Hall, 1847, Pal. N. Y., vol. 1, p.
177. Preoccupied by McCoy in 1844.

See M. milleri.

bilitata, Hall, 1859, Pal. N. Y., vol. 3, p.
299, Low. Held. Gr.billingsana, n. s., Guelph Gr. Proposed
instead of M. hercynia in Pal. Foss., vol.

1, p. 157, which was preoccupied.

bivittata, Hall, 1852, Pal. N. Y., vol. 2, p.
345, Guelph Gr.bowdeni, Safford, 1869, Geo. of Tenn., p.
288, Nashville Gr.boydi, Hall, 1843, (Loxonema boydi,) Geo.
Rep. 4th Dist. N. Y., p. 138, and Pal.

N. Y., vol. 2, p. 346, Guelph Gr.

boylii, Nicholson, 1875, Quar. Jour. Geo.
Soc. Lond., vol. 31, p. 547, Guelph Gr.calcifera, Whitfield, 1889, (Lophospira
calcifera,) Bull. Am. Mus. Nat. Hist.,

vol. 2, p. 55, Calciferous Gr.

carinifera, Shumard, 1863, Trans. St. Louis
Acad. Sci., vol. 2, p. 106, Calcifer-

ous Gr.

cassandra, Billings, 1865, Pal. Foss., vol. 1,
p. 189, Quebec Gr.cassina, Whitfield, 1886, (Lophospira cas-
sina,) Bull. Am. Mus. Nat. Hist., vol. 1,

p. 312, Birdseye Gr.

catharina, Billings, 1865, Pal. Foss., vol.
1, p. 231, Quebec Gr.chamberlini, Whitfield, 1878, Ann. Rep.
Geo. Sur. Wis., p. 84, and Geo. Wis.,

vol. 4, p. 317, Niagara Gr.

confusa, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 54, Calcifer-

ous Gr.

cicelia, Billings, 1865, Pal. Foss., vol. 1, p.
233, Quebec Gr.conoidea, Hall, 1852, Pal. N. Y., vol. 2, p.
13, Medina Gr.conradi, Hall, 1867, 20th Rep. N. Y. Mus.
Nat. Hist., p. 396, Niagara Gr.constricta, Whiteaves, 1884, Pal. Foss.,
vol. 3, p. 25, Guelph Gr.copii, White, 1882, Rep. Invert. Foss.
New Mex., p. xxx, Coal Meas.decurta, Hall, 1877, 1st Ed. of Am. Pal.
Foss., p. 244, syn. for M. subab-

breviata.

desiderata, Hall, 1862, 15th Rep. N. Y.
Mus. Nat. Hist., p. 50, and Pal. N. Y.

vol. 5, pt. 2, p. 89, Up. Held. Gr.

V egressa, Billings, 1874, Pal. Foss., vol. 2, p.

58, Up. Held. Gr.

elegantula, see Pleurotomaria elegantula.

estella, Billings, 1862, Pal. Foss., vol. 1,

p. 157, Guelph Gr.

extenuata, Hall, 1859, Pal. N. Y., vol. 3,
p. 298, Low. Held. Gr.funata, Billings, 1866, Catal.
Sil. Foss. Antic., p. 55, An-

ticosti Gr.

gigantea, Billings, 1857, Rep.
of Progr. Geo. Sur. Can.,

p. 298, Mid. Sil.

gracilens, Whitfield, 1889,
Bull. Am. Mus. Nat. Hist.,

vol. 2, p. 53, Calcifer-

ous Gr.

gracilis, Hall, 1847, Pal. N. Y.,
vol. 1, p. 181, Trenton and

Hud. Riv. Gr.

gypsea, Dawson, 1868, Acad.
Geol., p. 310, Carbonif-

erous.

hebe, Billings, 1874, Pal. Foss., vol. 2, p. 57, Gaspe

limestone No. 8, Devonian.

helicteres, Salter, 1859, Can. Org. Rem.,
Decade 1, p. 21, Black Riv. and Tren-

ton Grs.

hercyna, Billings, 1862, Pal. Foss., vol. 1,
p. 158. The name was preoccupied by

Roemer in 1843. See M. billingsana.

hermione, Billings, 1862, Pal. Foss., vol. 1,
p. 33, Chazy or Black Riv. Gr.hespelerensis, Whiteaves, 1884, Pal. Foss.,
vol. 3, p. 24, Guelph Gr.hyale, Billings, 1862, Pal. Foss., vol. 1, p.
33, Chazy or Black Riv. Gr.infrequens, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 457, Chazy Gr.inornata, Meek & Worthen, 1866, Proc.
Acad. Nat. Sci., p. 274, and Geo. Sur.

Ill., vol. 5, p. 599, Coal Meas.

insculpta, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 26, and Bull. Am. Mus. Nat.

Hist. p. 85, Warsaw Gr.

intercedens, Hall, 1879, Pal. N. Y., vol. 5,
pt. 2, p. 92, Up. Held. Gr.jessica, Billings, 1865, Pal. Foss., vol. 1, p.
189, Quebec Gr.kansasensis, Swallow, 1858, Trans. St.
Louis Acad. Sci., vol. 1, p. 195, Coal

Meas.

keokuk, Worthen, (in press,) Geo. Sur.
Ill., vol. 8, p. 141, Keokuk Gr.laphami, Hall, 1861, Rep. of Progr. Wis.,
p. 36, Niagara Gr.latifasciata, Etheridge, 1878, Quar. Jour.
Geo. Soc., vol. 34, p. 600, Up. Sil.lasallensis, Worthen, (in press,) Geo. Sur.
Ill., vol. 8, p. 141, Up. Coal Meas.leda, Hall, 1861, 14th Rep. N. Y. Mus. Nat.
Hist., p. 103, and Pal. N. Y., vol. 5, pt.

2, p. 91, Up. Held. Gr.

limitaris, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 108, Kinderhook Gr.

linearis, Billings, 1859, Can. Nat. and Geo.,

vol. 4, p. 359, Calciferous Gr.

logani, Hall, 1852, Pal. N. Y., vol. 2, p.

346, Guelph Gr.

longispira, Hall, 1852, Pal. N. Y., vol. 2, p.

345, Guelph Gr.

macrospira, Hall, 1852, Pal. N. Y., vol. 2,

p. 346, Guelph Gr.

FIG. 687.—
Murchisonia
gracilla.

- maia*, Hall, 1861, 14th Rep. N. Y. Mus. Nat. Hist., p. 103, and Pal. N. Y., vol. 5, pt. 2, p. 91, Up. Held. Gr.
major, Hall, 1851, Geo. Lake Sup. Land Dist., vol. 2, p. 209, Trenton Gr.
marcouana, Geinitz, 1866, Carb. und Dyas in Neb., p. 11, Coal Meas.
melaniformis, Shumard, 1855, Geo. Rep. Mo., p. 208, Calciferous Gr.
micula, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 244, and Pal. N. Y., vol. 5, pt. 2, p. 93, Ham. Gr. Proposed instead of *M. turricula*, Hall, 1862, which was preoccupied.
milleri, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 244, Trenton and Hud. Riv. Grs. Proposed instead of *M. bicincta*, Hall, 1847, Pal. N. Y., vol. 1, p. 177, which was preoccupied.
minima, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 203, Middle Coal Meas.
minuta, Hall, 1850, Pal. N. Y., vol. 3, p. 208, Low. Held. Gr.
missisquoi, Billings, 1865, Pal. Foss., vol. 1, p. 307, Quebec Gr.
modesta, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 299, Hud. Riv. Gr.
mucro, Winchell, 1866, Rep. Low. Peninsula Mich., p. 96, Ham. Gr.
multigruma, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 104, Hud. Riv. Gr.
multivolvis, Billings, 1857, Rep. Progr. Geo. Sur. Can., p. 299, Hud. Riv. Gr.
mylitta, Billings, 1862, Pal. Foss., vol. 1, p. 157, Guelph Gr.
nebraskensis, Geinitz, 1866, Carb. und Dyas in Neb., p. 12, and Pal. E. Neb., p. 234, Coal Meas.
neglecta, Winchell, 1863, Proc. Acad. Nat. Sci., p. 20, Marshall Gr.
obelisca, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 317, Birdseye Gr.
obsoleta, Meek, 1871, Proc. Acad. Nat. Sci., p. 175, Coal Meas.
obtusa, Hall, 1852, Pal. N. Y., vol. 2, p. 333, Coralline limestone.
ozarkensis, Shumard, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 196, Calciferous Gr.
papillosa, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 301, Mid. Sil.
perangulata, Hall, 1847, Pal. N. Y., vol. 1, p. 41, Black Riv. and Trenton Grs.
perversa, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 195, Up. Coal Meas.
petilla, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 186, Niagara Gr.
placida, Billings, 1865, Pal. Foss., vol. 1, p. 235, Quebec Gr.
procris, Billings, 1862, Pal. Foss., vol. 1, p. 34, Black Riv. Gr.
prava, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 316, Birdseye Gr.
prolixa, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 303, Kinderhook Gr.
quadrineta, Winchell, 1863, Proc. Acad. Nat. Sci., p. 19, Marshall Gr.
rugosa, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 299, Hud. Riv. Gr.
serrulata, Salter, 1859, Can. Org. Rem., Decade 1, p. 20, Black Riv. and Trenton Grs.
shumardana, Winchell, 1863, Proc. Acad. Nat. Sci. Phil., p. 20, Marshall Gr.
simulatrix, Billings, 1865, Pal. Foss., vol. 1, p. 232, Quebec Gr.
soluta, Whiteaves, 1884, Pal. Foss., vol. 3, p. 28, Guelph Gr.
sorocula, Billings, 1865, Pal. Foss., vol. 1, p. 233, Quebec Gr.
subabbreviata, D'Orbigny, 1850, Prodr. d. Paleont. t. 1, p. 8, Chazy Gr. Proposed instead of *M. abbreviata*, Hall, 1847, Pal. N. Y., vol. 1, p. 32, which was preoccupied.
subfusiformis, see *Fusispira subfusiformis*.
subtæniata, see *Orthonema subtæniatum*.
subulata, Conrad, 1842, (*Loxonema subulatum*), Jour. Acad. Nat. Sci., vol. 8, p. 273, and Pal. N. Y., vol. 2, p. 91, Clinton Gr.
summerensis, Safford, 1869, Geo. of Tenn., p. 288, Nashville Gr.
sylvia, Billings, 1865, Pal. Foss., vol. 1, p. 190, Quebec Gr.
terebra, White, 1879, Bull. U. S. Geo. Sur. Terr., vol. 5, No. 2, p. 219, and Cont. to Pal., No. 6, p. 139, Carboniferous.
terebialis, Hall, 1852, Pal. N. Y., vol. 2, p. 334, Coralline limestone.
terebriiformis, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 28, and Bull. Am. Mus. Nat. Hist., p. 86, Warsaw Gr.
teretiformis, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 298, Hud. Riv. Gr.
texana, Shumard, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 626, Coal Meas.
tricarinata, Hall, 1847, Pal. N. Y., vol. 1, p. 178, Trenton Gr.
tringulata, Dawson, 1868, Acad. Geol., p. 310, Carboniferous.
tropidophora, Whiteaves, 1884, Pal. Foss., vol. 3, p. 29, Guelph Gr.
turricula, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 301, Mid. Sil.
turricula, Hall. The name was preoccupied. See *M. micula*.
turritella, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 27, and Bull. Am. Mus. Nat. Hist., p. 88, Warsaw Gr.
turritiformis, Hall, 1852, Pal. N. Y., vol. 2, p. 347, Guelph Gr.
uniangulata, Hall, 1847, Pal. N. Y., vol. 1, p. 179, Trenton and Hud. Riv. Grs.
uniangulata var. *abbreviata*, Hall, 1847, Pal. N. Y., vol. 1, p. 304, Hud. Riv. Gr.
varians, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 300, Hud. Riv. Gr.
varicosa, Hall, 1847, Pal. N. Y., vol. 1, p. 42, Birdseye Gr.
ventricosa, Hall, 1847, Pal. N. Y., vol. 1, p. 41, Black Riv. and Trenton Grs.
vermicula, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 27, and Bull. Am. Mus. Nat. Hist., p. 87, Warsaw Gr.

63, Proc. Acad. Nat. Sci., vol. 1, p. 32, Calciferous Gr.
 of Progr. Geo. Surv. Gr.
 Org. Rem., Dec. and Trenton Grs.
 63, Proc. Acad. Nat. Sci., vol. 1, p. 156, C. elph Gr.
 Pal. Foss., vol. 3, p. 225, Niagara Gr.
 xanthippe, Billings, 1862, Pal. Foss., vol. 1, p. 155, Guelph Gr.
Natica, Adanson, 1757, Histoire Naturelle du Senegal, p. 172. [Ety. *nato*, to swim with a fluctuating motion.] This genus is unknown in Paleozoic rocks.
altonensis, see *Naticopsis altonensis*.
carleyana, see *Naticopsis carleyana*.
chesterensis, see *Naticopsis chesterensis*.
littonana, see *Macrochilina littonana*.
magister, syn. for *Naticopsis ventricosa*.
shumardi, see *Naticopsis shumardi*.
ventricosa, see *Naticopsis ventricosa*.
Naticopsis, McCoy, 1844, Synop. Carb. Foss. Ireland, p. 33. [Ety. from resemblance to the genus *Natica*.] Subglobose, solid, imperforate; whorls few, convex, rapidly expanding; last one large; spire short; aperture subovate, straighter on the inner side, rounded below; columella callous, flattened, longitudinal impression for the operculum; lip sharp, entire; surface smooth in part or marked with oblique striae. Type *N. phillipsi*.
acquistriata, Meek, 1871, Proc. Acad. Nat. Sci., p. 76, and Ohio Pal., vol. 1, p. 216, Up. Held Gr.
altonensis, McChesney, 1865, (Natica altonensis,) Desc. New. Pal. Foss., and Geo. Sur. Ill., vol. 5, p. 595, Coal Meas.
carleyana, Hall, 1858, (Natica carleyana,) Trans. Alb. Inst., vol. 4, p. 31, and Bull. Am. Mus. Nat. Hist., p. 71, Warsaw Gr.
chesterensis, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 100, Kaskaskia Gr.
comperta, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, pl. 29, Up. Held Gr.
cretacea, Hall & Whitfield, 1873, 23d Rep. N. Y. Mus. Nat. Hist., p. 240, Up. Held Gr.
depressa, Winchell, 1863, Proc. Acad. Nat. Sci., p. 22, Marshall Gr.
dispassa, Dawson, 1868, Acad. Geol., p. 309, Carbonif.
gigantea, Hall & Whitfield, 1873, 23d Rep. N. Y. Mus. Nat. Hist., p. 238, Chemung Gr.
hollidayi, see *Trachydromia hollidayi*.
howi, Hartt, 1868, Acad. Geol., p. 309, Carboniferous.
humilis, see *Isonema humile*.
lavis, Meek, 1871, Proc. Acad. Nat. Sci., p. 76, and Ohio Pal., vol. 1, p. 215, Up. Held Gr.

littonana var. *genevievensis*, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 268, Kaskaskia Gr.
maddisonensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 9, and Geo. Sur. Ill., vol. 8, p. 144, St. Louis Gr.
magister, syn. for *N. ventricosa*.
monilifera, White, 1880, 12th Rep. U. S. Geo. Sur. Terr., p. 168, Up. Coal Meas.
nana, Meek & Worthen, 1860, (Platystoma nana,) Proc. Acad. Nat. Sci., p. 463, and Geo. Sur. Ill., vol. 2, p. 365, Up. Coal Meas.
nodosa, see *Trachydromia nodosa*.
ortoni, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 230, Coal Meas.
pricii, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 202, Up. Coal Meas.
remex, White, 1876, Geo. Uinta Mountains, p. 109, and Cont. to Pal., No. 6, p. 139, Low. Aubrey Gr.
shumardi, McChesney, 1860, (Natica shumardi,) Desc. New. Pal. Foss., p. 62, Coal Meas.
subovata, Worthen, 1873, Geo. Sur. Ill., vol. 5, p. 595, Coal Meas.
ventricosa, Norwood & Pratten, 1854, (Natica ventricosa,) Jour. Acad. Nat. Sci., 2d ser., vol. 3, p. 76, Coal Meas.
wheeleri, Swallow, 1860, (Littorina wheeleri,) Trans. St. Louis Acad. Sci., vol. 1, p. 658, and Geo. Sur. Ill., vol. 5, p. 595, Coal Meas.
ziczac, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 223, Kaskaskia Gr.
OPHIALOTROCHUS, Meek, 1864, Geo. California, vol. 1, p. 15. [Ety. *ophthalmus*, umbilicus; *Trochus*, a genus.] Distinguished from *Euomphalus* by having a more prominent spire, in having its whorls flattened or broadly concave around the outer side, and flattened with an outward slope; it is a more ponderous shell, with a more oblique outline to its lip, in consequence of which it projects much farther forward on the upper than on the lower side of the aperture. Type *O. whitneyi*.
whitneyi, Meek, 1864, Pal. of California, vol. 1, p. 15, Carboniferous.
OPHILETA, Vanuxem, 1842, Geo. Rep. N. Y., p. 36. [Ety. *ophis*, snake.] Discoidal spire sunk above; umbilicus perfectly open, exposing the whorls on one plane; whorls slender, numerous, truncate, and biangular exteriorly; aperture having a sinus below and a notch above. Type *O. complanata*.
abditata, Billings, 1865, Pal. Foss., vol. 1, p. 189, Quebec Gr.
 (?) *bella*, Billings, 1865, Pal. Foss., vol. 1, p. 310, Quebec Gr.
compacta, Salter, 1859, Can. Org. Rem., Decade 1, p. 16, syn. for *O. complanata*.

FIG. 388.—*Naticopsis lavis*.

complanata, Vanuxem, 1842, Geo. Rep. N. Y., p. 36, and Pal. N. Y., vol. 1, p. 11, Calciferous Gr.
 complanata var. nana, Meek, 1870, Hayden's U. S. Geo. Sur. Terr., p. 295, and Geo. 4th Parallel, vol. 4, p. 17, Calciferous Gr.
 disjuncta, Billings, 1865, Pal. Foss., vol. 1, p. 344, Calciferous Gr.
 levata, Vanuxem, 1842, Geo. Rep. N. Y., p. 36, Calciferous Gr.

FIG. 689.—*Ophileta nerine*.

nerine, Billings, 1865, Pal. Foss., vol. 1, p. 245, Quebec Gr.
 ottawensis, Billings, 1860, Can. Nat. and Geol., vol. 5, p. 167, Trenton Gr.
 owenana, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 313, Galena Gr.
 primordialialis, Winchell, 1864, (Straparolus primordialialis,) Am. Jour. Sci. and Arts, 2d ser., vol. 37, p. 228, and Geo. Wis., vol. 4, p. 173, Potsdam Gr.
 profunda, Billings, 1865, Pal. Foss., vol. 1, p. 188, Quebec Gr.
 uninangula a, Hall, 1847, (Euomphalus uninangulatus,) Pal. N. Y., vol. 1, p. 9, Calcifer. Gr.

ORMATHICHNUS, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 222. [Ety. *ormathos*, string of beads; *ichnos*, track.] Supposed to be the trail of a Gasteropod, and consisting of a continuous beaded track or trail. Type *O. moniliformis*.

moniliformis, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 222, Utica Slate Gr.

ORTHONEMA, Meek & Worthen, 1861, Proc. Acad. Nat. Sci., Phil., p. 146. [Ety. *orthos*, straight; *nema*, thread.] elongate, many whorled, ornamented with revolving carinae, crossed by nearly straight lines of growth; body whorl angular, not much enlarged or produced below; aperture angular above, slightly effuse below; peristome incomplete; outer lip simple, nearly straight; axis imperforate. Type *O. salteri*.

carbonarium, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 7, and Geo. Sur. Ill., vol. 8, p. 145, Coal Meas.

FIG. 690.
Orthonema newberryi.

conicum, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 270, and Geo. Sur. Ill., vol. 5, p. 590, Coal Meas.
 newberryi, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 81, and Ohio Pal., vol. 1, p. 217, Up. Held. Gr.
 salteri, Meek & Worthen, 1860, (Eunema (?) salteri,) Proc. Acad. Nat. Sci. Phil., p.

464, and Geo. Sur. Ill., vol. 2, p. 381, Low. Coal Meas.

subteniatum, Geinitz, 1866, (Murchisonia subteniatum,) Carb. and Dyas in Neb., p. 12, and Pal. E. Neb., p. 228, Coal Meas.
Orthonychia, Hall, 1843, syn. for *Platyceras*.
 ORTHOSTOMA, Conrad, 1838, Ann. Rep. N. Y., p. 119. [Ety. *orthos*, straight; *stoma*, mouth.] Shell spiral, spire plain, convex, terminal volution, ending in a straight tube. Type *O. commune*.

FIG. 691.—*Orthostoma commune*.

commune, Conrad, 1838, Ann. Rep. N. Y., p. 119, figured in 1841, Ann. Rep. pl. 2, fig. 16, Birdseye Gr.

PALEACMÆA, Hall & Whitfield, 1873, 23d Rep. N. Y. Mus. Nat. Hist., p. 242. [Ety. *palaos*, ancient; *Acmea*, an existing genus of shells.] Conical, more or less elevated; apex subcentral, erect or slightly curved; peristome entire, not sinuate; surface marked concentrically. Type *P. typica*.

FIG. 692.—*Palæotrochus kearneyi*.

irvingi, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 51, and Geo. Wis., vol. 4, p. 173, Potsdam Gr.
 typica, Hall & Whitfield, 1873, 23d Rep. N. Y. Mus. Nat. Hist., p. 242, Potsdam Gr.

PALEOTROCHUS, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 133. [Ety. *palaos*, ancient; *Trochus*, a genus.] Conical, trochiform; spire elevated; volutions moderately

, vol. 2, p. 381.

66, (Murchisonia
Dyas in Neb.
p. 228, Concl. Meas.
h. for Platyceras.
Ann. Rep. N. Y.,
straight; stoma,
spire plain, con-
n, ending in a
t. commune.

commune.

Ann. Rep. N. Y.,
Ann. Rep. pl.

field, 1873, 23d
Hist., p. 242.
Acmæa, an exist-
Conical, more or
central, erect or
ome entire, not
d concentrically.

keyi.

Ann. Rep. Geo.
eo. Wis., vol. 4,

1873, 23d Rep.
p. 242, Pots.

Pal. N. Y., vol.
alataios, ancient;
al, trochiform;
ns moderately

convex; aperture transverse. Type *P. kearneyi*.

kearneyi, Hall, 1862, (Pleuronomaria *kearneyi*,) 14th Rep. N. Y. Mus. Nat. Hist., p. 105, Up. Held. Gr.

precursor, Clarke, 1885, Bull. U. S. Geo. Sur. No. 16, p. 55, Portage Gr.

Patella, Linnaeus, 1758, Syst. Nat. 10th Ed. [Ety. *patella*, dish.] Not a Palaeozoic genus.

levetii, see *Lepetopsis levetti*.

Phanerolites, Sowerby, 1844, Min. Conch., vol. 7, p. 29. [Ety. *phaneros*, aperture; *litos*, extended. Syn. for *Eccyliomphalus*.

paradoxus, see *Eccyliomphalus paradoxus*.

Phragmolites, syn. for *Cyrtolites*.

compressus, see *Cyrtolites compressus*.

PHRAGMOSTOMA, Hall, 1861, 14th Rep. N. Y. Mus. Nat. Hist., p. 94. [Ety. *phragmos*, a partition; *stoma*, the mouth; from the septum within the aperture, which distinguishes this genus from *Carinaropsis* and *Bellerophon*.] Type *P. cymbula*.

FIG. 603.—*Phragmostoma cymbula*. View of aperture; *L*, lamina; *S*, septum.

cunulae, Hall, 1861, 14th Rep. N. Y. Mus. Nat. Hist., p. 94, Hud. Riv. Gr.

cymbula, Hall, 1861, 14th Rep. N. Y. Mus. Nat. Hist., p. 94, Hud. Riv. Gr.

natator, see *Bellerophon natator*.

Physa, Dapernaud, 1801, Hist. Nat. d. Moll. Not a Palaeozoic genus.

prisca, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 262, Subcarboniferous.

Pileopsis conoides, *P. naticoides*, *P. rotundata*, and *P. spiralis*, Castelnau, 1843, Syst. Sil. Not recognized.

tubifer, syn. for *Platyceras dumosum*.

vetustus, Sowerby. Not American.

Planorbis, Guettard, 1756, Mem. Acad. Sci. Paris. Not a Palaeozoic genus.

trilobatus, see *Bucania trilobata*.

PLATYCERAS, Conrad, 1840, Ann. Rep. N. Y., p. 205. [Ety. *platys*, broad; *keras*, horn.] Depressed subglobose, subovoid, or obliquely subconical; spire small; volutions none, or very few, without columella; aperture more or less expanded, often campanulate, and sometimes with lip reflexed; peristome entire or sinuous; surface striated, cancellated, lamellose, or spiniferous. Type *P. dumosum*.

acutirostre, Hall, 1858, (Capulus *acutirostris*,) Trans. Alb. Inst., vol. 4, p. 31, and Geo. Sur. Iowa, p. 665, Warsaw Gr.

agreste, Hall, 1859, Pal. N. Y., vol. 3, p. 338, Low. Held. Gr.

ammon, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 37, and Pal. N. Y., vol. 5, pt. 2, p. 20, Up. Held. Gr.

angulatum, Hall, 1852, (Acroculia *angulata*,) Pal. N. Y., vol. 2, p. 289, Clinton and Niagara Grs.

aretistoma, Ulrich, 1886, Cont. to Am. Pal., p. 30, Up. Held. Gr.

arcuatum, Hall, 1859, Pal. N. Y., vol. 3, p. 336, Low. Held. Gr.

argo, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 39, and Pal. N. Y., vol. 5, pt. 2, p. 19, Up. Held. Gr.

attenuatum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 30, and Pal. N. Y., vol. 5, pt. 2, p. 6, Ham. Gr.

attenuatum, see *P. dumosum* var. *attenuatum*.

auriculatum, Hall, 1876, Illust. Devonian Foss., pl. 3, Ham. Gr.

billingsi, Hall, 1859, Pal. N. Y., vol. 3, p. 315, Low. Held. Gr.

biseriale, Hall, 1860, Supp. to Geo. Iowa, vol. 1, pt. 2, p. 90, Burlington Gr.

bisinuatum, Hall, 1859, Pal. N. Y., vol. 3, p. 318, Low. Held. Gr.

bisulcatum, Hall, 1859, Pal. N. Y., vol. 3, p. 327, Low. Held. Gr.

bivolve, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 302, Kinderhook Gr.

bucculentum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 33, and Pal. N. Y., vol. 5, pt. 2, p. 10, Ham. Gr.

calantica, Hall, 1859, Pal. N. Y., vol. 3, p. 328, Low. Held. Gr.

caliosum, Hall, 1859, Pal. N. Y., vol. 3, p. 478, Oriskany sandstone.

campanulatum, Winchell & Marey, 1855, Mem. Bost. Soc. Nat. Hist., p. 99, Niagara Gr.

capax, Keyes, 1888, Proc. Am. Phil. Soc., (author's copy, p. 13,) Burlington Gr.

capulus, Hall, 1860, Supp. Geo. Iowa, p. 91, Burlington Gr.

carinatum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 33, and Pal. N. Y., vol. 5, pt. 2, p. 5, Up. Held. Gr.

chesterense, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 265, Kaskaskia Gr.

cirriforme, Conrad, 1841, Ann. Rep. N. Y. Not clearly defined.

clavatum, Hall, 1859, Pal. N. Y., vol. 3, p. 337, Low. Held. Gr.

concavum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 30, and Pal. N. Y., vol. 5, pt. 2, p. 3, Up. Held. Gr.

conicum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 31, and Pal. N. Y., vol. 5, pt. 2, p. 3, Ham. Gr.

conradi, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 182, Devonian.

cornuforme, Winchell, 1863, Proc. Acad. Nat. Sci., p. 18, Marshall Gr.

crassum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 36, and Pal. N. Y., vol. 5, pt. 2, p. 18, Up. Held. Gr.

curvirostrum, Hall, 1859, Pal. N. Y., vol. 3, p. 338, Low. Held. Gr.

cymbium, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 35, and Pal. N. Y., vol. 5, pt. 2, p. 12, Up. Held. Gr.

cyrtolites, McChesney, 1859, Pal. Foss., p. 71, Coal Meas.

- ✓ dentalium, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 29, and Pal. N. Y., vol. 5, pt. 2, p. 2, Up. Held. Gr.
- dilatatum, Hall, 1859, Pal. N. Y., vol. 3, p. 322, Low. Held. Gr.
- ✓ dumosum, Conrad, 1840, Ann. Rep. N. Y., p. 205, and Pal. N. Y., vol. 5, pt. 2, p. 14, Up. Held. Gr.

FIG. 694.—*Platyoceras dumosum*.

- ✓ dumosum var. attenuatum, Meek, 1871, Proc. Acad. Nat. Sci., p. 75, and Ohio Pal., vol. 1, p. 212, Up. Held. Gr.
- ✓ dumosum var. rarispinum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 38, and Pal. N. Y., vol. 5, pt. 2, p. 16, Up. Held. Gr.
- ✓ echinatum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 38, and Pal. N. Y., vol. 5, pt. 2, p. 13, Ham. Gr.
- elongatum, Hall, 1859, Pal. N. Y., vol. 3, p. 335, Low. Held. Gr.
- equilaterale, Hall, 1860, Supp. to vol. 1, pt. 2, Iowa Rep., p. 89, and Geo. Sur. Ill., vol. 5, p. 518, Keokuk Gr.
- ✓ erectum, Hall, 1843, (*Acroculia erecta*) Geo. 4th Dist. N. Y., p. 174, and Pal. N. Y., vol. 5, pt. 2, p. 5, Cornif. & Ham. Grs.
- expansum, see *Strophostylus expansus*.
- fissurellum, Hall, 1860, Supp. to Geo. Rep. Iowa, vol. 1, pt. 2, p. 90, and Geo. Sur. Ill., vol. 5, p. 519, Keokuk Gr.
- ✓ fluctuosum, Ulrich, 1886, Cont. to Am. Pal., p. 31, Up. Held. Gr.
- formosum, Keyes, 1888, Proc. Am. Phil. Soc., (author's copy, p. 14,) Kinderhook Gr.
- ✓ fornicatum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 35, and Pal. N. Y., vol. 5, pt. 2, p. 11, Up. Held. Gr.
- ✓ fornicatum var. contractum, Hall, 1876, Illust. Devonian Foss., pl. 5, Up. Held. Gr.
- ✓ gebhardi, Conrad, 1840, Ann. Rep. N. Y., p. 206, and Pal. N. Y., vol. 3, p. 312, Low. Held. and Oriskany Grs.
- gibbosum, Hall, 1859, Pal. N. Y., vol. 3, p. 322, Low. Held. Gr.
- haliotoides, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 264, and Geo. Sur. Ill., vol. 3, p. 458, Kinderhook Gr.
- herzeri, Winchell, 1870, Proc. Am. Phil. Soc., p. 256, Marshall Gr.

incile, Hall, 1859, Pal. N. Y., vol. 3, p. 332, Low. Held. Gr.

infundibulum, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 266, and Geo. Sur. Ill., vol. 5, p. 517, Keokuk Gr.

intermedium, Hall, 1859, Pal. N. Y., vol. 3, p. 321, Low. Held. Gr.

laciniosum, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 14, Niagara Gr.

laevigatum, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 263, Kaskaskia Gr.

lamellosum, Hall, 1859, Pal. N. Y., vol. 3, p. 330, Low. Held. Gr.

latum, Keyes, 1888, Proc. Am. Phil. Soc., (author's copy, p. 14,) Burlington Gr.

lodiense, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 170, and Ohio Pal., vol. 2, p. 313, Waverly Gr.

magnificum, Hall, 1859, Pal. N. Y., vol. 3, p. 476, Oriskany sandstone.

membranaceum, Ringueberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 15, Niagara Gr.

minutissimum, Walcott, 1879, Desc. New Spec. Foss., p. 1, Calciferous Gr.

multisinuatum, Hall, 1859, Pal. N. Y., vol. 3, p. 319, Low. Held. Gr.

multispinosum, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 73, and Ohio Pal., vol. 1, p. 210, Cornif. Gr.

naticoides, Etheridge, 1878, Quar. Jour. Geo. Soc., vol. 34, p. 603, Up. Sil.

nebraskense, Meek, 1872, Pal. E. Neb., p. 227, Coal Meas.

newberryi, Hall, 1859, Pal. N. Y., vol. 3, p. 333, Low. Held. Gr.

niagarensis, Hall, 1852, (*Acroculia niagarensis*) Pal. N. Y., vol. 2, p. 288, Niagara Gr.

nodosum, Conrad, 1841, Ann. Rep. N. Y., p. 56, and Pal. N. Y., vol. 3, p. 473, Oriskany sandstone.

obesum, Hall, 1859, Pal. N. Y., vol. 3, p. 329, Low. Held. Gr.

obliquum, Keyes, 1888, Proc. Am. Phil. Soc., (author's copy, p. 13,) Burlington Gr.

occidens, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 254, Subcarboniferous.

ovale, Stevens, 1858, (*Acroculia ovalis*) Am. Jour. Sci., vol. 25, p. 261, Subcarboniferous.

pabulocrinus, Owen, 1862, (*Pileopsis pabulocrinus*) Geo. Sur. Indiana, p. 364, Keokuk Gr.

parallum, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 302, Kinderhook Gr.

patulum, Hall, 1859, Pal. N. Y., vol. 3, p. 477, Oriskany sandstone.

pentalobus, Hall, 1859, Pal. N. Y., vol. 3, p. 319, Low. Held. Gr.

perlatum, Hall, 1859, Pal. N. Y., vol. 3, p. 328, Low. Held. Gr.

perplexum, Hall, 1876, Illust. Devonian Foss., pl. 2, Up. Held. Gr.

N. Y., vol. 3, p. 322.
 & Worthen, 1866,
 Phil., p. 26, and
 5, p. 517. Keo-
 59, Pal. N. Y., vol.
 . Gr.
 5, 1886, Bull. Buf.
 5, p. 14. Niag-
 orthen, 1866, Proc.
 53, Kaskaskia Gr.
 Pal. N. Y., vol. 3,
 r.
 oc. Am. Phil. Soc.,
) Burlington Gr.
 Proc. Acad. Nat.
 Ohio Pal., vol. 2,
 Pal. N. Y., vol. 3,
 istone.
 eberg, 1886, Bull.
 ol. 5, p. 15, Niag-
 , 1879, Desc. New
 ciferous Gr.
 1859, Pal. N. Y.,
 eid. Gr.
 1871, Proc. Acad.
 3, and Ohio Pal.,
 Gr.
 1878, Quar. Jour.
 303, Up. Sil.
 2, Pal. E. Neb., p.
 Pal. N. Y., vol. 3,
 r.
 (Acroculia niag-
 ol. 2, p. 288, Ni-
 Ann. Rep. N. Y.,
 , vol. 3, p. 473,
 . N. Y., vol. 3, p.
 Proc. Am. Phil.
 p. 13,) Burlington
 5, Monogr. U. S.
 254, Subcarbon-
 Acroculia ovalis,
 5, p. 261, Subcar-
 32, (Pileopsis pa-
 , Indiana. p. 364,
 tfield, 1862, Proc.
 , vol. 8, p. 302,
 . N. Y., vol. 3, p.
 ne.
 Pal. N. Y., vol. 3,
 i. N. Y., vol. 3, p.
 Illust. Devonian
 Gr.

perplicatum, Hall, 1859, Pal. N. Y., vol. 3,
 p. 325, Low. Held. Gr.
 pileiforme, Hall, 1859, Pal. N. Y., vol. 3,
 p. 327, Low. Held. Gr.
 piso, Walcott, 1885, Monogr. U. S. Geo.
 Sur., vol. 8, p. 254, Subcarboniferous.
 platystoma, Hall, 1859, Pal. N. Y., vol. 3,
 p. 326, Low. Held. Gr.
 platystoma var. alveatum, Hall, 1859, Pal.
 N. Y., vol. 3, p. 326, Low. Held. Gr.
 plicatile, Hall, 1859, Pal. N. Y., vol. 3, p.
 325, Low. Held. Gr.
 plicatum, Conrad, 1840, (Calceola plicata,)
 Ann. Rep. N. Y., p. 207, and Pal. N. Y.,
 vol. 3, p. 334, Low. Held. Gr.
 princeps, Billings, 1871, Can. Nat. and
 Geol., vol. 6, p. 220, Georgia Gr.
 primordiale, Hall, 1863, 16th Rep. N. Y.
 Mus. Nat. Hist., p. 136, Potsdam Gr.
 proclive, Ringueberg, 1886, Bull. Buf. Soc.
 Nat. Sci., vol. 5, p. 14, Niagara Gr.
 pyramidatum, Hall, 1859, Pal. N. Y., vol.
 3, p. 336, Low. Held. Gr.
 quincyense, McChesney, 1861, New Pal.
 Foss., p. 90, and Geo. Sur. Ill., vol. 3,
 p. 510, Burlington Gr.
 quinquesinuatum, Ulrich, 1886, Cont. to
 Am. Pal., p. 29, Up. Held. Gr.
 reflexum, Hall, 1859, Pal. N. Y., vol. 3, p.
 477, Oriskany sandstone.
 retrorsum, Hall, 1859, Pal. N. Y., vol. 3,
 p. 320, Low. Held. Gr.
 retrorsum var. abnorme, Hall, 1859, Pal.
 N. Y., vol. 3, p. 321, Low. Held. Gr.

FIG. 605.—*Platyceeras re-
 versum*.

reversum, Hall,
 1860, Supp. to
 Geo. Rep. Iowa,
 vol. 1, pt. 2, p.
 91, Burlington
 Gr.
 rictum, Hall, 1862,
 15th Rep. N. Y.
 Mus. Nat. Hist.,
 p. 35, and Pal.
 N. Y., vol. 5, pt.
 2, p. 13, Ham.
 and Up. Held.
 Grs.
 robustum, Hall,
 1859, Pal. N. Y., vol. 3, p. 313, Low.
 Held. Gr.
 senex, Winchell & Marcy, 1865, (Porcel-
 lia senex.) Mem. Bost. Soc. Nat. Hist.,
 p. 111, Niagara Gr.
 serratum, Ulrich, 1886, Cont. to Am. Pal.,
 p. 30, Up. Held. Gr.
 sinuatum, Hall, 1859, Pal. N. Y., vol. 3,
 p. 314, Low. Held. Gr.
 spinigerum, Worthen, 1873, Geo. Sur. Ill.,
 vol. 5, p. 594, Coal Meas.
 spirale, Hall, 1859, Pal. N. Y., vol. 3, p.
 331, Low. Held. Gr.
 squalodens, Whitfield, 1882, Ann. N. Y.
 Acad. Sci., vol. 2, p. 202, Up. Held. Gr.
 subnodosum, Hall, 1859, Pal. N. Y., vol.
 3, p. 474, Oriskany sandstone.
 subplicatum, Meek & Worthen, 1866, Proc.
 Acad. Nat. Sci. Phil., p. 265, and Geo.
 Sur. Ill., vol. 3, p. 457, Kinderhook Gr.

subrectum, Hall, 1859, 12th Rep. N. Y.,
 Mus. Nat. Hist., p. 18, and Pal. N. Y.,
 vol. 5, pt. 2, p. 1, Up. Held. Gr.
 subrectum, Hall, 1860, Supp. to Iowa Rep.
 The name being preoccupied, Meek &
 Worthen proposed *P. infundibulum*.
 subsinuatum, Worthen, 1882, Bull. No. 1,
 Ill. St. Mus. Nat. Hist., p. 38, Low.
 Held. Gr. Proposed instead of *P. sub-*
undatum, M. & W., which was preoc-
 cupied.
 subundatum, Conrad, 1841, Ann. Rep.
 N. Y., p. 56, Up. Held. Gr.
 subundatum, Meek & Worthen, 1868, Geo.
 Sur. Ill., vol. 3, p. 457. The name was
 preoccupied. See *P. subsinuatum*.
 sulcatum, Conrad, 1841, Ann. Rep. N. Y.,
 p. 56, Oriskany sandstone.
 sulcopicatum, Hall, 1859, Pal. N. Y., vol.
 3, p. 324, Low. Held. Gr.
 symmetricum, Hall, 1862, 15th Rep. N. Y.
 Mus. Nat. Hist., p. 34, and Pal. N. Y.,
 vol. 5, pt. 2, p. 9, Ham. and Up.
 Held. Grs.
 tenniliratum, Hall, 1859, Pal. N. Y., vol.
 3, p. 317, Low. Held. Gr.
 thetiforme, Walcott, 1885, Monogr. U. S.
 Geo. Sur., vol. 8, p. 184, Devonian.
 thetis, Hall, 1862, 15th Rep. N. Y. Mus.
 Nat. Hist., p. 32, and Pal. N. Y., vol. 5,
 pt. 2, p. 8, Up. Held. and Ham. Grs.
 thetis var. subspinosum, Hall, 1876,
 Illust. Devonian Foss., pl. 3, Ham. Gr.
 tortum, Meek, 1871, Proc. Acad. Nat. Sci.,
 p. 171, and Ohio Pal., vol. 2, p. 345,
 Coal Meas.
 tortuosum, Hall, 1859, Pal. N. Y., vol. 3,
 p. 472, Oriskany sandstone.
 tribulosum, White, 1880, 12th Rep. U. S.
 Geo. Sur. Terr., p. 168, Burling-
 ton Gr.
 trigonale, Stevens, 1858, (Acroculia tri-
 gonalis.) Am. Jour. Sci. and Arts, vol.
 25, p. 260, Carboniferous.
 trilobatum, Hall, 1859, Pal. N. Y., vol. 3,
 p. 316, Low. Held. Gr.
 tubiforme, Hall, 1859, Pal. N. Y., vol. 3,
 p. 332, Low. Held. Gr.
 uncum, Meek & Worthen, 1866, Proc.
 Acad. Nat. Sci., p. 264, and Geo. Sur.
 Ill., vol. 5, p. 516, Keokuk Gr.
 undatum, Hall, 1876, Illust. Devonian
 Foss., pl. 7, and Pal. N. Y., vol. 5, pt.
 2, p. 17, Up. Held. Gr.
 undulatum, Walcott, 1885, Monogr. U. S.
 Geo. Sur., vol. 8, p. 184, Devonian.
 undulostriatum, Hall, 1859, Pal. N. Y.,
 vol. 3, p. 336, Low. Held. Gr.
 unguiforme, Hall, 1859, Pal. N. Y., vol. 3,
 p. 322, Low. Held. Gr.
 uniserialle, Nicholson, 1874, Rep. Pal.
 Ont., p. 116, Up. Held. Gr.
 unisulcatum, Hall, 1859, Pal. N. Y., vol.
 3, p. 316, Low. Held. Gr.
 ventricosum, Conrad, 1840, Ann. Rep.
 N. Y., p. 206, and Pal. N. Y., vol. 3, p.
 311, Low. Held. Gr.
 vomerium, Winchell, 1863, Proc. Acad.
 Nat. Sci., p. 19, Marshall Gr.

PLATYSCHISMA, McCoy, 1844, Syn. Carb. Foss. Ireland, p. 38. [Ety. *platys*, wide; *schisma*, slit.] Obtusely conical, ventricose; spire short, obtuse, few whorls; aperture very oblique, large, lunate, deeply indented by the preceding whorl, rounded anteaally, narrow retrally, with a very wide, shallow sinus in the middle part of the outer lip not forming a definite band, sometimes obsolete; no trace of inner lip; pillar thin, a little reflected; surface smooth or only marked by the retrally waved lines of growth; umbilicus small, round, open. Type *P. helicites*.

✓ **ambiguum**, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 188, Devonian.

FIG. 696.—*Platyschisma dubium*.

dubium, Dawson, 1868, Acad. Geol., p. 309, Carboniferous.

✓ **mccoyi**, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 188, Devonian.

PLATYSTOMA, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 275. [Ety. *platys*, broad; *stoma*, mouth.] Spire short; aperture large, dilated; labrum joining the body whorl. This name was preoccupied for a shell by Klein in 1753, for an insect by Meigen in 1803, and for a fish by Agassiz in 1829. The genus has been named *Platyserina*. Type *P. ventricosum*.

✓ **affine**, Billings, 1874, Pal. Foss., vol. 2, p. 60, Gaspé limestone, No. 8, Devonian.

✓ **aplatum**, Hall, 1876, Illust. Devonian Foss., pl. 11, and Pal. N. Y., vol. 5, pt. 2, p. 26, Schoharie grit.

arenosum, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 276, and Pal. N. Y., vol. 3, p. 302, Low. Held. Gr.

✓ **belial**, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 30, Genesee shales.

✓ **defiguratum**, Hall, 1876, Illust. Devonian Foss., pl. 9, and Pal. N. Y., vol. 5, pt. 2, p. 24, Ham. Gr.

depressum, Hall, 1859, Pal. N. Y., vol. 3, p. 301, Low. Held. Gr.

✓ **euomphaloides**, Hall, 1876, Illust. Devonian Foss., pl. 9, and Pal. N. Y., vol. 5, pt. 2, p. 25, Ham. Gr.

hemisphericum, Hall, 1843, (Euomphalus hemisphericus.) Geo. Rep. 4th Dist. N. Y., p. 109, and Pal. N. Y., vol. 2, p. 288, Niagara Gr.

grayvillense, Worthen, 1882, Bull. No. 11, St. Mus. Nat. Hist., p. 38, Coal Meas. Proposed instead of *P. tumidum*, M. & W., which was preoccupied.

inornatum, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 255, Subcarboniferous.

lichas, see *Callonema lichas*.

✓ **lineatum**, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 276, and Pal. N. Y., vol. 5, pt. 2, p. 21, Up. Held and Ham. Grs.

✓ **lineatum var. amplum**, Hall, 1876, Illust. Devonian Foss., pl. 9, and Pal. Foss. N. Y., vol. 5, pt. 2, p. 23, Ham. Gr.

lineatum var. callosum, Hall, 1876, Illust. Devonian Foss., pl. 9, and Pal. Foss. N. Y., vol. 5, pt. 2, p. 23, Ham. Gr.

lineatum var. sinuosum, Hall, 1876, Illust. Devonian Foss., pl. 11, and Pal. Foss. N. Y., vol. 5, pt. 2, p. 24, Ham. Gr.

minutissimum, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 55, Portage Gr.

nana, see *Naticopsis nana*.

niagarensis, Hall, 1852, Pal. N. Y., vol. 2, p. 287, Niagara Gr.

peoriense, McChesney, 1860, Desc. New Pal. Foss., Up. Coal Meas.

plebeium, Hall, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 175, Niagara Gr.

pleurotoma, Hall, 1876, Illust. Devonian Foss., pl. 10, and Pal. N. Y., vol. 5, pt. 2, p. 30, Up. Held. Gr.

pleuratum, Whiteaves, 1887, Cont. to Can. Pal., vol. 1, p. 118, Ham. Gr.

shumardi, Verneuil, 1846, (Turbo shumardi) Bull. d. l. Soc. Geol. d. France, and Pal. N. Y., vol. 5, pt. 2, p. 135, Ham. Gr.

strophium, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 41, and Pal. N. Y., vol. 5, pt. 2, p. 25, Cornif. Gr.

subangulatum, Hall, 1859, Pal. N. Y., vol. 3, p. 301, Low. Held. Gr.

trigonostoma, Meek, 1871, Proc. Acad. Nat. Sci., p. 169, and Ohio Pal., vol. 1, p. 185, Niagara Gr.

tumidum, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 463, Up. Coal Meas.

turbinatum, Hall, 1861, 14th Rep. N. Y. Mus. Nat. Hist., p. 106, and Pal. N. Y., vol. 5, pt. 2, p. 27, Up. Held. Gr.

turbinatum var. cochleatum, Hall, 1876, Illust. Devonian Foss., pl. 10, and Pal. N. Y., vol. 5, pt. 2, p. 28, Up. Held. Gr.

unisulcatum, Conrad, 1842, (Pleurotomaria unisulcata.) Jour. Acad. Nat. Sci., vol. 8, p. 271, and Pal. N. Y., vol. 5, pt. 2, p. 27, Up. Held. Gr.

ventricosum, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 275, and Pal. N. Y., vol. 3, p. 300, Low. Held. Gr.

Plectostylus, Conrad, 1842, Jour. Acad. Nat. Sci., p. 275. The name was preoccupied.

hildrethi, see *Macrochilina hildrethi*.

FIG. 697.—*Platystoma niagarensis*.

FIG. 698.—*Platystoma trigonostoma*.

trigonostoma, Meek, 1871, Proc. Acad. Nat. Sci., p. 169, and Ohio Pal., vol. 1, p. 185, Niagara Gr.

tumidum, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 463, Up. Coal Meas.

turbinatum, Hall, 1861, 14th Rep. N. Y. Mus. Nat. Hist., p. 106, and Pal. N. Y., vol. 5, pt. 2, p. 27, Up. Held. Gr.

turbinatum var. cochleatum, Hall, 1876, Illust. Devonian Foss., pl. 10, and Pal. N. Y., vol. 5, pt. 2, p. 28, Up. Held. Gr.

unisulcatum, Conrad, 1842, (Pleurotomaria unisulcata.) Jour. Acad. Nat. Sci., vol. 8, p. 271, and Pal. N. Y., vol. 5, pt. 2, p. 27, Up. Held. Gr.

ventricosum, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 275, and Pal. N. Y., vol. 3, p. 300, Low. Held. Gr.

Plectostylus, Conrad, 1842, Jour. Acad. Nat. Sci., p. 275. The name was preoccupied.

hildrethi, see *Macrochilina hildrethi*.

Hall, 1876, Illust.
9, and Pal. Foss.
23, Ham. Gr.
Hall, 1876, Illust.
1, and Pal. Foss.
24, Ham. Gr.
1885, Bull. U. S.
S. Portage Gr.
na.

—*Platystoma niagarensis*.

yonian Foss., pl.
ol. 5, pt. 2, p. 30,

37, Cont. to Can.
n. Gr.

46, (Turbo shu-
Geol. d. France,
5, pt. 2, p. 135,

15th Rep. N. Y.
and Pal. N. Y.
if. Gr.

, Pal. N. Y., vol.
r.

onostoma, Meek,
1871, Proc. Acad.
Nat. Sci., p. 169,
nd Ohio Pal., vol.
, p. 185, Niagara
Gr.

oidum, Meek &
Worthen, 1860,
roc. Acad. Nat.
feas.

4th Rep. N. Y.
and Pal. N. Y.,
Held. Gr.

um, Hall, 1876,
, pl. 10, and
2, p. 28, Up.

2, (Pleurotoma-
Acad. Nat. Sci.,
N. Y., vol. 5, pt.

2, Jour. Acad.
and Pal. N. Y.,
i. Gr.

Jour. Acad. Nat.
ne was preoc-

hildrethi.

PLEURONOTUS, Hall, 1879, Pal. N. Y., vol.
5, pt. 2, p. 138. [Ety. *pleura*, side;
notos, back.] Distinguished from *Euom-*
phalus by the broadly expanded aper-
ture, sinuate on the upper margin,
making a deep retral angle, which
meets a peripheral band. Type *P.*
decewii.

Fig. 699.—*Pleuronotus decewii*.

decewii, Billings, 1861, (*Euomphalus*
decewii.) Can. Jour., p. 358, Up.
Held. Gr.

PLEUROTOMARIA, DeFrance, 1824, Tableau d.
Corps. Organises Fossiles, p. 114, and
Dict. Sci. Nat., t. 41, p. 381. [Ety. *pleura*,
side; *tome*, cut or notch.] Shell trochi-
form, more or less conical, pearly within,
variable in thickness, with or without
an umbilicus; volutions angular, flat-

tened, or rounded;
surface ornamented
with striae, nodes,
granulations, or car-
inae; aperture sub-
quadrate, semioval,
suborbicular, or sub-
rhombic; inner lip
thin; fissure of outer
lip narrow and
deep; revolving
band corresponding
in depth with the sinus. Type *P. an-*
glica.

abrupta, Billings, 1859, Can. Nat. and Geo.,
vol. 4, p. 354, Calciferous Gr.

acadica, Dawson, 1883, Rep. on Redpath
Mus., p. 11, Subcarboniferous.

adamsi, Worthen, 1884, Bull. No. 2, Ill.
St. Mus. Nat. Hist., p. 5, and Geo. Sur.

Ill., vol. 8, p. 137, Coal Meas.

adjutor, Hall, 1879, Pal. N. Y., vol. 5, pt.
2, p. 80, Up. Held Gr.

advena, Winchell, 1864, Am. Jour. Sci.
and Arts, 2d series, vol. 37, p. 228, Pots-

dam Gr.

agarista, Billings, 1865, Pal. Foss., vol. 1,
p. 230, Quebec Gr.

Fig. 700.—*Pleuroto-*
maria anglica.

agave, Billings, 1865, Pal. Foss., vol. 1, p.
170, Trenton Gr.

ambigua, Hall, 1847, Pal. N. Y., vol. 1, p.
176, Trenton Gr.

americana, Billings, 1860, Can. Nat. and
Geo., vol. 5, p. 164, Trenton Gr.

amphitrite, Billings, 1862, Pal. Foss., vol.
1, p. 32, Chazy or Black P. v. Gr.

angulata, Conrad, 1843, Proc. Acad.
Nat. Sci. Phil. This name was

preoccupied by Sowerby.

antiquata, Hall, 1847, Pal. N. Y., vol.
1, p. 31, Chazy Gr.

aperta, see *Raphistoma apertum*.

apicalis, Hall, 1876, Illust. Devonian
Foss., pl. 20, and Pal. N. Y., vol.

5, pt. 2, p. 88, Chemung Gr.

arabella, Billings, 1865, Pal. Foss.,
vol. 1, p. 343, Calciferous Gr.

arachne, Billings, 1862, Pal. Foss.,
vol. 1, p. 31, Black Riv. Gr.

arata, Hall, 1862, 15th Rep. N. Y.
Mus. Nat. Hist., p. 42, and Pal.

N. Y., vol. 5, pt. 2, p. 64, Scho-
harie grit.

arata var. *clausa*, Hall, 1879, Pal.
N. Y., vol. 5, pt. 2, p. 65, Up.

Held. Gr.

axion, Hall, 1867, 20th Rep. N. Y.
Mus. Nat. Hist., p. 394, Niag-

ara Gr.

beckwithana, McChesney, 1860, Desc.
New Pal. Foss., p. 61, Coal Meas.

beekmanensis, Whitfield, 1889, Bull. Am.
Mus. Nat. Hist., vol. 2, p. 53, Calcifer-

ous Gr.

biangulata, Hall, 1847, Pal. N. Y., vol. 1,
p. 31, Chazy Gr.

bicarinata, McChesney, 1860. Preoccu-
pied. See *P. turbiniformis*.

bilix, see *Cyclonema bilix*.

bispiralis, Hall, 1852, Pal. N. Y., vol. 2,
p. 348, Guelph Gr.

bonbarborensis, Cox, 1857, Geo. Sur. Ky.,
vol. 3, p. 567, Coal Meas.

brazoensis, Shumard, 1860, Trans. St.
Louis Acad. Sci., vol. 1, p. 624, and

Geo. Sur. Ill., vol. 2, p. 354, Low. Coal
Meas.

broadheadi, White, 1880, 12th Rep. U. S.
Geo. Sur. Terr., p. 169, Coal Meas.

calcifera, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 352, Calciferous Gr.

calphurnia, Billings, 1865, Pal. Foss., vol.
1, p. 230, Up. Taconic, Quebec Gr.

calyx, Billings, 1859, Can. Nat. and Geo.,
vol. 4, p. 454, Chazy Gr.

canadensis, Billings, 1865, Pal. Foss., vol.
1, p. 342, Calciferous Gr.

capillaria, Conrad, 1842, Jour. Acad. Nat.
Sci., vol. 8, p. 271, and Pal. N. Y., vol.

5, pt. 2, p. 77, Ham. Gr.

carbonaria, Norwood & Pratten, 1854,
Jour. Acad. Nat. Sci., 2d series, vol. 3,
p. 75, Coal Meas.

cassii, Meek & Worthen, 1868, Geo. Sur.
Ill., vol. 3, p. 359, Niagara Gr.

cavumbilicata, Winchell, 1866, Rep. Low.
Penin. Mich., p. 96, Ham. Gr.

- chesterensis, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 460, and Geo. Sur. Ill., vol. 2, p. 303, Kaskaskia Gr.
- chesterensis*, Swallow, 1863, Trans. St. Louis Acad. Sci. The name was preoccupied, but it is probably a synonym.
- circe*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 303, Hud. Riv. Gr.
- clipeiformis*, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 57, Niagara Gr.
- conca*, see *Eotrochus concavus*.
- coniformis*, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 38, Coal Meas. Proposed instead of *P. conoides*, M. & W.
- conoides*, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 271. Preoccupied by Deshayes in 1831. See *P. coniformis*.
- conulus*, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 26, and Bull. Am. Mus. Nat. Hist., p. 84, Warsaw Gr.
- cooperensis*, n. s. Kaskaskia Gr. Proposed instead of *P. trochiformis*, Swallow, Trans. St. Louis Acad. Sci., vol. 2, p. 99, that was preoccupied.
- coronula*, Hall, syn. for *P. sphaerulata*.
- coxana*, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 272, and Geo. Sur. Ill., vol. 5, p. 600, Coal Meas.
- coxana*, Worthen, 1884. The name was preoccupied. See *P. iowensis*.
- crevieri*, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 456, Chazy Gr.
- cryptata*, Billings, 1866, Catal. Sil. Foss. Antic., p. 54, Anticosti Gr.
- cyclonemoides*, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 360, Niagara Gr.
- cyclostoma*, Whiteaves, 1884, Pal. Foss., vol. 3, p. 23, Guelph Gr.
- deiopea*, Billings, 1862, Pal. Foss., vol. 1, p. 155, Guelph Gr.
- delia*, Billings, 1874, Pal. Foss., vol. 2, p. 61, Gaspé limestone No. 8, Devonian.
- delicatula*, Hall, 1876, Illust. Devonian Foss., pl. 19, and Pal. N. Y., vol. 5, pt. 2, p. 70, Up. Held. Gr.
- delphinuloides*, Goldfuss, as identified by d'Archiac & Verneuil. Not American.
- depauperata*, Hall, 1862, Geo. Rep. Wis., p. 55, Hud. Riv. Gr.
- depressa*, Cox, 1857, Geo. Sur. Ky., vol. 3, Coal Meas. The name was preoccupied by Passy in 1832, by Phillips in 1836, and by DeKoninck in 1841. See *P. kentuckiensis*.
- disjuncta*, Hall, 1876, Illust. Devonian Foss., pl. 20, and Pal. N. Y., vol. 5, pt. 2, p. 84, Ham. Gr.
- dispersa*, Dawson, 1868, Acad. Geol., p. 310, Carboniferous.
- occens*, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 452, Chazy Gr.
- doris*, see *Cyclonema doris*.
- dryope*, Billings, 1865, Pal. Foss., vol. 1, p. 170, Black Riv. Gr.
- durhamensis*, Whiteaves, 1884, Pal. Foss., vol. 3, p. 24, Guelph Gr.
- elegantula*, Hall, 1868, (Murchisonia elegantula,) Trans. Alb. Inst., vol. 4, p. 27, and Bull. Am. Mus. Nat. Hist., p. 84, Warsaw Gr.
- ella*, Hall, 1876, Illust. Devonian Foss., pl. 19, and Pal. N. Y., vol. 5, pt. 2, p. 72, Ham. Gr.
- elora*, Billings, 1862, Pal. Foss., vol. 1, p. 154, Guelph Gr.
- emmetensis*, Winchell, 1866, Rep. Low. Penin. Mich., p. 96, Ham. Gr.
- estella*, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 195, Ham. Gr.
- etna*, Billings, 1865, Pal. Foss., vol. 1, p. 226, Quebec Gr.
- eugenia*, Billings, 1862, Pal. Foss., vol. 1, p. 30, Black Riv. Gr.
- euomphaloides*, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 18, Ham. Gr.
- exigua*, Winchell, 1862, Proc. Acad. Nat. Sci., p. 424, Marshall Gr.
- filitexta*, Hall, 1876, Illust. Devonian Foss., pl. 19, and Pal. N. Y., vol. 5, pt. 2, p. 73, Ham. Gr.
- galtensis*, Billings, 1862, Pal. Foss., vol. 1, p. 154, Guelph Gr.
- giffordi*, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 5, and Geo. Sur. Ill., vol. 8, p. 135, Coal Meas.
- glandula*, Shumard, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 626, Coal Meas.
- gonopleura*, Winchell & Marcy, 1865, Bost. Soc. Nat. Hist., p. 98, Niagara Gr.
- granulostriata*, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 459, and Geo. Sur. Ill., vol. 2, p. 356, Low. Coal Meas.
- grayvillensis*, Norwood & Pratten, 1854, Jour. Acad. Nat. Sci., 2d series, vol. 3, p. 75, Coal Meas.
- gregaria*, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 355, Calciferous Gr.
- gurleyi*, Meek, 1871, Proc. Acad. Nat. Sci., p. 177, Coal Meas.
- halii*, see *Trochonema halii*.
- hallana*, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 399, Permian Gr.
- halli*, see *Raphistoma halli*.
- harpys*, Billings, 1865, Pal. Foss., vol. 1, p. 227, Quebec Gr.
- haydenana*, Geinitz, 1866, Carb. und Dyas in Neb., p. 11, and Pal. E. Neb., p. 231, Coal Meas.
- hebe*, Hall, 1861, 14th Rep. N. Y. Mus. Nat. Hist., p. 105, and Pal. N. Y., vol. 5, pt. 2, p. 68, Up. Held. Gr.
- helena*, Billings, 1860, Can. Nat. and Geol., vol. 5, p. 165, Hud. Riv. Gr.
- hickmanensis*, Winchell, 1869, Geo. of Tenn. and Proc. Am. Phil. Soc. vol., 12, p. 257, Waverly Gr.
- hortensia*, Billings, 1865, Pal. Foss., vol. 1, p. 227, Quebec Gr.
- hoyi*, Hall, 1861, Rep. of Progr. Wis. Sur., p. 35, Niagara Gr.
- humerosa*, Meek & Hayden, 1858, Proc. Acad. Nat. Sci. Phil., p. 262, and Pal. Up. Mo., p. 46, Coal Meas.

1884, Pal. Foss.,
Churichsonia ele-
 mentaria, vol. 4, p. 27,
 Nat. Hist., p. 84,
 Devonian Foss., pl.
 5, pt. 2, p. 72,
 Foss., vol. 1, p.
 366, Rep. Low.
 n. Gr.
 1872, 24th Rep.
 p. 195, Ham. Gr.
 Foss., vol. 1, p.
 al. Foss., vol. 1,
 1862, 15th Rep.
 p. 18, Ham. Gr.
 Proc. Acad. Nat.
 Hist., Devonian
 N. Y., vol. 5, pt.
 al. Foss., vol. 1,
 Bull. No. 2, Ill.
 and Geo. Sur.
 Meas.
 Trans. St. Louis
 Coal Meas.
 Arcy, 1865, Bost.
 Niagara Gr.
 Worthen, 1860,
 459, and Geo.
 low. Coal Meas.
 Pratt, 1854,
 series, vol. 3,
 Nat. and Geo.,
 Gr.
 Acad. Nat. Sci.,
 Trans. St. Louis
 Permian Gr.
 Foss., vol. 1,
 Carb. und Dyas
 E. Neb., p. 231,
 p. N. Y. Mus.
 Pal. N. Y., vol.
 Gr.
 Nat. and Geo.,
 Gr.
 1869, Geo. of
 il. Soc. vol. 12,
 Pal. Foss., vol.
 Progr. Wis. Sur.,
 1858, Proc.
 262, and Pal.
 ss.

humilis, Hall, 1858, Trans. Alb. Inst., vol.
 4, p. 21, and Bull. Am. Mus. Nat. Hist.,
 p. 82, Warsaw Gr.
humilis, Winchell, 1862. This name was
 preoccupied.
huronensis, Winchell, 1862, Proc. Acad.
 Nat. Sci. Phil., vol. 6, 2d ser., p. 425,
 Portage Gr.
hyale, Billings, 1865, Pal. Foss., vol. 1, p.
 228, Quebec Gr.
idia, Hall, 1861, Rep. of Progr. Wis. Sur.,
 p. 35, Niagara Gr.
ignobilis, Dawson, 1868, Acad. Geol., p.
 310, Carboniferous.
illinoisensis, Worthen, 1884, Bull. No. 2,
 Ill. St. Mus. Nat. Hist., p. 4, and Geo.
 Sur. Ill., vol. 8, p. 135, Coal Meas.
imitator, see *Callonema imitator*.
immatura, Billings, 1859, Can. Nat. and
 Geo., vol. 4, p. 454, Chazy Gr.
indenta, Hall, 1847, Pal. N. Y., vol. 1, p.
 176, Trenton Gr.
inexpectans, Hall & Whitfield, 1875, Ohio
 Pal., vol. 2, p. 117, Clinton Gr.
inornata, Meek, 1872, Pal. E. Neb., p. 232,
 Coal Meas.
insolita, Hall, 1876, Illust. Devonian
 Foss., pl. 20, and Pal. N. Y., vol. 5, pt.
 2, p. 81, Ham. Gr.
iowensis, Worthen, (in press.) Geo. Sur.
 Ill., vol. 8, p. 138, Keokuk Gr.
isaacsi, Hall & Whitfield, 1873, 23d Rep.
 N. Y. Mus. Nat. Hist., p. 238, Che-
 mung Gr.
itys, Hall, 1876, Illust. Devonian Foss.,
 pl. 20, and Pal. N. Y., vol. 5, pt. 2, p.
 76, Ham. Gr.
itys var. *tenuispira*, Hall, 1879, Pal. N. Y.,
 vol. 5, pt. 2, p. 87, Ham. Gr.
kearneyi, see *Palaeotrochus kearneyi*.
kentuckiensis, n. s. Coal Meas. Proposed
 instead of *P. depressa* in Geo. Sur. Ky.,
 vol. 3, p. 569, which was preoccupied.
labrossa, Hall, 1859, Pal. N. Y., vol. 3, p.
 339, Low. Held. Gr.
laphami, Whitfield, 1878, Ann. Rep. Geo.
 Sur. Wis., p. 84, and Geo. Wis., vol. 4,
 p. 296, Niagara Gr.
lapicida, see *Raphistoma lapicidum*.
laurentina, Billings, 1859, Can. Nat. and
 Geo., vol. 4, p. 354, Calcif. Gr.
leavenworthana, see *Cyclonema leaven-*
worthanum.
lenticularis, see *Raphistoma lenticulare*.
lineata, Hall, 1843, (Turbo lineatus.) Geo.
 Rep. 4th Dist. N. Y. Preoccupied.
 See *P. itys*.
litorea, Hall, 1852, Pal. N. Y., vol. 2, p.
 12, Medina sandstone.
lonensis, Walcott, 1885, Monogr. U. S.
 Geo. Sur., vol. 8, p. 80, Trenton Gr.
lucina, Hall, 1862, 15th Rep. N. Y. Mus.
 Nat. Hist., p. 42, and Pal. N. Y., vol. 5,
 pt. 2, p. 67, Up. Held. and Ham. Grs.
lucina var. *perfasciata*, Hall, 1876, Illust.
 Devonian Foss., pl. 20, and Pal. N. Y.,
 vol. 5, pt. 2, p. 83, Ham. Gr.
lydia, Billings, 1874, Pal. Foss., vol. 2, p.
 62, Gaspe limestone, No. 8, Devonian.

marcouana, Geinitz, 1866, Carb. und Dyas
 in Neb., p. 10, and Pal. E. Neb., p. 233,
 Coal Meas.
meekana, Hall, 1858, Trans. Alb. Inst.,
 vol. 4, p. 22, and Bull. Am. Mus. Nat.
 Hist., p. 82, Warsaw Gr.
meta, Meek & Worthen, 1865, Proc. Acad.
 Nat. Sci., p. 252, Keokuk Gr.
micula, Hall, 1862, Geo. Rep. Wis., p. 55,
 Hud. Riv. Gr.
misera, Billings, 1859, Can. Nat. and Geo.,
 vol. 4, p. 354, Calcif. Gr.
mississippiensis, White & Whitfield, 1862,
 Proc. Bost. Soc. Nat. Hist., vol. 8, p.
 302, Kinderhook Gr.
missisquoi, Billings, 1865, Pal. Foss., vol.
 1, p. 191, Quebec Gr.
missouriensis, Swallow, 1860, (Trochus
missouriensis.) Trans. St. Louis Acad.
 Sci., vol. 1, p. 657, Coal Meas.
mitigata, Hall, 1860, 13th Rep. N. Y. Mus.
 Nat. Hist., p. 108, Kinderhook Gr.
modesta, Keyes, 1888, Proc. Acad. Nat.
 Sci. Phil., pl. xii, figs. 2a, 2b, Coal Meas.
mohawkensis, n. sp. Birdseye limestone.
 Proposed instead of *P. nodulosa*, in
 Pal. N. Y., vol. 1, p. 44, which was pre-
 occupied.
montezuma, Worthen, 1883, Geo. Sur. Ill.,
 vol. 7, p. 324, Burlington Gr.
muralis, Owen, 1852, Geo. Sur. Wis.,
 Iowa, and Minn., p. 581, Trenton Gr.
nasoni, Hall, 1861, Geo. Rep. Wis., p. 34,
 and Geo. Wis., vol. 4, p. 215, Trenton Gr.
nauvoocensis, Worthen, 1884, Bull. No. 2,
 Ill. St. Mus. Nat. Hist., p. 5, and Geo.
 Sur. Ill., vol. 8, p. 137, Keokuk Gr.
nevadensis, Walcott, 1885, Monogr. U. S.
 Geo. Sur., vol. 8, p. 259, Subcar-
 boniferous.
newportensis, White, 1880, 12th Rep. U. S.
 Geo. Sur. Terr., p. 169, Coal Meas.
niota, Hall, 1861, Geo. Rep. Wis., p. 33,
 Trenton Gr.
nitela, Hall, 1879, Pal. N. Y., vol. 5, pt. 2,
 p. 85, Up. Held. Gr.
nodomarginata, McChesney, 1860, Desc.
 New. Pal. Foss., p. 70, and Trans. Chi.
 Acad. Sci., p. 47, Ham. Gr.
nodulosa, Hall, 1847, Pal. N. Y., vol. 1, p.
 44. The name was preoccupied by
 Sandberger in 1842, and by King in
 1844. See *P. mohawkensis*.
nodulostriata, Hall, 1858, Trans. Alb. Inst.,
 vol. 4, p. 21, and Bull. Am. Mus. Nat.
 Hist., p. 80, Warsaw Gr.
normani, Billings, 1865, Pal. Foss., vol. 1,
 p. 228, Quebec Gr.
nuculolata, Hall, 1847, Pal. N. Y., vol. 1,
 p. 42, Birdseye Gr.
numeria, Billings, 1865, Pal. Foss., vol. 1,
 p. 229, Quebec Gr.
obsoleta, Hall, 1847, Pal. N. Y., vol. 1, p.
 44, Birdseye Gr.
obtusispira, Shumard, 1859, Trans. St.
 Louis Acad. Sci., vol. 1, p. 401, Coal
 Meas.
occidens, Hall, 1867, 20th Rep. N. Y. Mus.
 Nat. Hist., pp. 342, 364, Niagara Gr.

- ✓ *parvispira*, Winchell, 1862, Rep. Low. Peninsula Mich., p. 96, Ham. Gr.
pauper, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 457, Chazy Gr.
pauper, syn. for *Trochonema hallii*.
percarinata, see *Cyclonema percarinatum*.
perhumerosa, Meek, 1872, Pal. E. Neb., p. 232, Coal Meas.
perizomata, White, 1882, Rep. Invert. Foss. New Mex., p. xxxi, Coal Meas.
perlata, Hall, 1852, Pal. N. Y., vol. 2, p. 349, Guelph Gr.
perornata, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 401, Coal Meas.
pervetusta, Conrad, 1838, (*Cyclostoma pervetusta*.) Ann. Rep. N. Y., p. 65, Medina sandstone.
piasensis, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 22, and Bull. Am. Mus. Nat. Hist., p. 83, Warsaw Gr.
 ✓ *planidorsalis*, Hall, 1876, Illust. Devonian Foss., pl. 20, and Pal. N. Y., vol. 5, pt. 2, p. 82, Ham. Gr.
 ✓ *plena*, Hall, 1876, Illust. Devonian Foss., pl. 17, and Pal. N. Y., vol. 5, pt. 2, p. 66, Ham. Gr.
postumia, Billings, 1862, Pal. Foss., vol. 1, p. 91, Quebec Gr.
 ✓ *poulsoni*, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 270, Onondaga Gr.
pratteni, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 459, and Geo. Sur. Ill., vol. 2, p. 357, Low. Coal Meas.
 ✓ *princessa*, Billings, 1874, Pal. Foss. vol. 2, p. 59, Up. Held. Gr.
progne, Billings, 1860, Can. Nat. and Geol., vol. 5, p. 163, Black Riv. and Trenton Grs.
proutana, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 401, Coal Meas.
quadracarinata, Hall, 1847, Pal. N. Y., vol. 1, p. 43, Birdseye Gr.
 ✓ *quadrilix*, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 86, Up. Held. Gr.
quebecensis, Billings, 1865, Pal. Foss., vol. 1, p. 190, Quebec Gr.
quinesulcata, Winchell, 1865, Proc. Acad. Nat. Sci., p. 131, Marshall Gr.
racinensis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 84, and Geo. Wis., vol. 4, p. 296, Niagara Gr.
ramsayi, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 351, Calciferous Gr.
rota, Winchell, 1863, Proc. Acad. Nat. Sci., p. 19, Marshall Gr.
rotalia, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 46, and Pal. N. Y., vol. 5, pt. 2, p. 71, Ham. Gr.
rotuloides, see *Raphistoma rotuloides*.
rotunda, Hall, 1843, (*Euomphalus* (?) *rotundus*.) Geo. Rep. 4th Dist. N. Y., p. 172, and Illust. Devon. Foss., pl. 18, Corniferous Gr.
rotundata, Hall, see *P. subglobosa*.
rotundispira, Billings, 1865, Pal. Foss., vol. 1, p. 191, Quebec Gr.
rugulata, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 108, and Pal. N. Y., vol. 5, pt. 2, p. 75, Ham. Gr.
scitula, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 461, and Geo. Sur. Ill., vol. 2, p. 353, Low. Coal Meas.
selecta, Billings, 1865, Pal. Foss., vol. 1, p. 224, Quebec Gr.
semele, Hall, 1861, Geo. Rep. Wis., p. 36, Hud. Riv. Gr.
shumardi, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 462, and Geo. Sur. Ill., vol. 2, p. 260, Keokuk Gr.
sigaretoides, Winchell & Marcy, 1865, Bost. Soc. Nat. Hist., vol. 1, p. 98, Niagara Gr.
sinistrorsa, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 203, Coal Meas.
solarioides, Hall, 1852, Pal. N. Y., vol. 2, p. 348, Guelph Gr.
speciosa, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 459, and Geo. Sur. Ill., vol. 2, p. 352, Low. Coal Meas.
sphaerulata, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 272, Coal Meas.
spironema, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 272, and Geo. Sur. Ill., vol. 5, p. 601, Coal Meas.
sponsa, Billings, 1865, Pal. Foss., vol. 1, p. 226, Quebec Gr.
stella, Winchell, 1862, Proc. Acad. Nat. Sci., p. 424, Marshall Gr.
subangulata, see *Cyclonema subangulatum*.
subconica, Hall, 1847, Pal. N. Y., vol. 1, p. 174, Black Riv., Trenton, and Hud. Riv. Grs.
subconstricta, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 458, and Geo. Sur. Ill., vol. 2, p. 351, Low. Coal Meas.
subdecussata, Geinitz, 1866, Carb. and Dyas in Neb., p. 10, and Pal. E. Neb., p. 233, Coal Meas.
subdepressa, Hall, 1852, Pal. N. Y., vol. 2, p. 333, Coralline limestone.
subglobosa, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 245, Warsaw Gr. Proposed instead of *P. rotundata*, Hall, 1858, which was preoccupied.
subscalaris, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 460, and Geo. Sur. Ill., vol. 2, p. 360, Low. Coal Meas.
subsinuata, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 460, and Geo. Sur. Ill., vol. 2, p. 358, Low. Coal Meas.

FIG. 701.—*Placrotomaria ramsayi*.

- ✓ *regulata*, Hall, 1860, 13th Rep., p. 108, Ham. Gr.
riddelli, Shumard, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 625, Coal Meas.

c. Acad. Nat.
 ep. N. Y. Mus.
 l. N. Y., vol. 5,
 rotuloides.
 aphalus (?) ro-
 Dist. N. Y., p.
 Foss., pl. 18,
 lobosa.
 Pal. Foss., vol.
 ep. N. Y. Mus.
 al. N. Y., vol.
 n, 1860, Proc.
 461, and Geo.
 low. Coal Meas.
 l. Foss., vol. 1,
 ep. Wis., p. 36,
 en, 1860, Proc.
 462, and Geo.
 Keokuk Gr.
 Marcy, 1865,
 pl. 1, p. 98, Ni-
 Trans. St. Louis
 3, Coal Meas.
 l. N. Y., vol. 2,
 en, 1860, Proc.
 459, and Geo.
 low. Coal Meas.
 Jour. Acad.
 Coal Meas.
 en, 1866, Proc.
 272, and Geo.
 Coal Meas.
 l. Foss., vol. 1,
 r. Acad. Nat.
 subangulatum.
 l. N. Y., vol. 1,
 aton, and Hud.
 Worthen, 1860,
 phil., p. 458, and
 351, Low. Coal
 66, Carb. and
 l. Pal. E. Neb.,
 ul. N. Y., vol. 2,
 one.
 t Ed. Am. Pal.
 Gr. Proposed
 ta, Hall, 1858,
 en, 1860, Proc.
 460, and Geo.
 low. Coal Meas.
 en, 1860, Proc.
 460, and Geo.
 58, Low. Coal.

subtilstriata, see *Raphistoma subtilstri-
 atum*.
subturbinata, Meek & Hayden, 1858, Proc.
 Acad. Nat. Sci. Phil., p. 264, and Pal.
 Up. Mo., p. 47, Coal Meas.
sulcomarginata, Conrad, 1842, Jour. Acad.
 Nat. Sci., vol. 8, p. 272, and Pal. N. Y.,
 vol. 5, pt. 2, p. 60, Ham. Gr.
supracingulata, Billings, 1857, Rep. of
 Progr., Geo. Sur. Can., p. 302, Tren-
 ton Gr.
swallowana, Hall, 1858, Trans. Alb. Inst.,
 vol. 4, p. 24, and Bull. Am. Mus. Nat.
 Hist., p. 80, Warsaw Gr.
sybillina, Billings, 1866, Catal. Sil. Foss.
 Antic., p. 54, Anticosti Gr.
tabulata, Conrad, 1835, (Turbo *tabulata*.)
 Trans. Geo. Soc. Penn., vol. 1, p. 267,
 Coal Meas.
taggarti, Meek, 1874, 7th Rep. Hayden's
 U. S. Geo. Sur. Terr., p. 271, and Cont.
 to Pal., No. 6, p. 140, Coal Meas.
tectoria, Winchell, 1863, Proc. Acad. Nat.
 Sci., p. 19, Marshall Gr.
tenuicincta, Meek & Worthen, 1860, Proc.
 Acad. Nat. Sci., p. 459, and Geo. Sur.
 Ill., vol. 2, p. 355, Up. Coal Meas.
tenuimarginata, Hall, syn. for *Eotrochus*
concavus.
tenuistriata, Shumard, 1860, Trans. St.
 Louis Acad. Sci., vol. 1, p. 625, Coal
 Meas.
textiliger, Meek, 1871, Proc. Acad. Nat.
 Sci., p. 176, and Ohio Pal., vol. 2, p. 314,
 Waverly Gr.
thalia, see *Cyclonema thalia*.
trilineata, Hall, 1858, Trans. Alb. Inst.,
 vol. 4, p. 25, and Bull. Am. Mus. Nat.
 Hist., p. 80, Warsaw Gr.
trilix, Hall, 1862, 15th Rep. N. Y. Mus.
 Nat. Hist., p. 45, and Pal. N. Y., vol. 5,
 pt. 2, p. 79, Ham. Gr.
trochiformis, Swallow, 1863. The name
 was preoccupied by Portlock in 1843.
 See P. Cooperensis.
tropidophora, Meek, 1872, Am. Jour. Sci.
 and Arts, 3d series, vol. 4, p. 278, and
 Ohio Pal., vol. 1, p. 154, Hud.
 Riv. Gr.
turbiniformis, Meek
 & Worthen, 1860,
 Proc. Acad. Nat.
 Sci., p. 461, and Geo.
 Sur. Ill., vol. 2, p.
 359, Up. Coal Meas.
turgida, see *Holopea*
turgida.
umbilicata, see *Trocho-*
nema umbilicatum.
uniusculata, Conrad,
 1842, Jour. Phil.
 Acad. Sci., vol. 8, p. 271, Up. Held. Gr.
vadosa, Hall, 1860, 13th Rep. N. Y.
 Mus. Nat. Hist., p. 108, Kinder-
 hook Gr.
vagrans, Billings, 1862, Pal. Foss., vol. 1,
 p. 90, Quebec Gr.
valeria, Billings, 1865, Pal. Foss., vol. 1, p.
 169, and vol. 3, p. 23, Guelph Gr.

FIG. 702.—*Pleuro-
 tomaria turbinif-
 ormis*.

valvatiformis, Meek & Worthen, 1866,
 Proc. Acad. Nat. Sci. Phil., p. 273,
 and Geo. Sur. Ill., vol. 5, p. 602, Coal
 Meas.
viola, Billings, 1865, Pal. Foss., vol. 1, p.
 169, Guelph Gr.
virgo, Billings, 1865, Pal. Foss., vol. 1, p.
 224, Quebec Gr.
virguncula, Billings, 1865, Pal. Foss., vol.
 1, p. 225, Quebec Gr.
vitruvia, Billings, 1865, Pal. Foss., vol. 1,
 p. 171, Black Riv. Gr.
volumina, Billings, 1874, Pal. Foss., vol.
 2, p. 61, Gaspe limestone No. 8, De-
 vonian.
whitii, Winchell, 1862, Proc. Acad. Nat.
 Sci., p. 423, Marshall Gr.
wortheni, Hall, 1858, Trans. Alb. Inst.,
 vol. 4, p. 23, and Geo. Sur. Iowa, p. 664,
 Warsaw Gr.
POLYPHEMUS, Portlock, 1843, Geol. Lon-
 donderry, p. 415. [Ety. *Polypemus*, a
 genus of shells; *opsis*, appearance.]
 Subfusiform; spire elongated; whorls
 flattened, last one produced below and
 forming half the length of the shell;
 outer lip thin, nearly straight; inner
 lip wanting; columella without folds,
 slightly twisted and truncated at the
 connection with the outer lip; aper-
 ture narrow, subovate, effuse or slightly
 notched at the base of the columella;
 surface smooth, or only with obscure
 lines of growth. Type *P. elongata*.
chrysalis, Meek & Worthen, 1866, Proc.
 Acad. Nat. Sci. Phil., p. 267, and Geo.
 Sur. Ill., vol. 5, p. 596, Coal Meas.
keokuk, Worthen, (in press.) Geo. Sur.
 Ill., vol. 8, p. 144, Keokuk Gr.
inornata, Meek & Worthen, 1860, (*Loxo-*
nema inornatum), Proc. Acad. Nat. Sci.
 Phil., p. 463, and Geo. Sur. Ill., vol. 2,
 p. 374, Up. Coal Meas.
louisville, Hall & Whitfield, 1872, 24th
 Rep. N. Y. Mus. Nat. Hist., p. 193, Up.
 Held. Gr.
melanoides, Whitfield, 1882, Ann. N. Y.
 Acad. Sci., vol. 2, p. 225, Kaskaskia Gr.
nitidula, Meek &
 Worthen, 1860,
 (*Loxonema nitid-*
ula), Proc. Acad.
 Nat. Sci. Phil.,
 p. 465, and Geo.
 Sur. Ill., vol. 2, p.
 374, Up. Coal
 Meas.
peracuta, Meek &
 Worthen, 1860,
 (*Eulima* (?) *per-*
acuta), Proc. Acad.
 Nat. Sci. Phil., p.
 466, and Geo. Sur.
 Ill., vol. 2, p. 375,
 Up. Coal Meas.
teretiformis, Hall, 1877, 1st Ed. Am. Pal.
 Foss., p. 245, Warsaw Gr. Proposed in-
 stead of *P. elongata*, Hall, 1858, which
 was preoccupied.

FIG. 703.—*Polyphem-
 us nitidula*.

PORCELLIA, Leveille, 1835, Mem. Soc. Geol. France, vol. 2, p. 39. [Ety. proper name.] Discoid, depressed; whorls very slightly embracing, exposed in a very wide umbilicus, slightly deeper on one side than the other, from a trifling obliquity of the first one or two turns; a narrow band extends along the middle of the exterior, ending in a narrow slit in the lip; surface often nodular and ornamented with rough striae. Type *P. puzosi*.

FIG. 704.—*Porcellia puzosi*.

crassinoda, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 303, Kinderhook Gr.
hertzeri, Hall, 1876, Illust. Devonian Foss., pl. 16, and Pal. N. Y., vol. 5, pt. 2, p. 126, Up. Held Gr.
nais, Hall, 1862, (*Gyroceras nais*), 15th Rep. N. Y. Mus. Nat. Hist., p. 68, and Pal. N. Y., vol. 5, pt. 2, p. 127, Chemung Gr.
nodosa, Hall, 1860, Supp. to vol. 1, pt. 2, Iowa Geo. Sur., p. 92, Kinderhook Gr.
obliquinodus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 21, Marshall Gr.
peoriensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 6, and Geo. Sur. Ill., vol. 8, p. 138, Coal Meas.
rectinoda, Winchell, 1863, Proc. Acad. Nat. Sci., p. 18, Marshall Gr.
rotatoria, Hall, see *Goniatites plebeiformis*.
scioto, Hall & Whitfield, 1873, 23d Rep. N. Y. Mus. Nat. Hist., p. 240, Up. Held Gr.
senex, see *Platyceras senex*.
PSEUDOPHORUS, Meek, 1873, Ohio Pal., vol. 1, p. 221. [Ety. *pseudes*, false; *Phorus*, a genus. Shell depressed, subtrochiform; umbilicus broad, shallow, eccentric; volutions two or three; suture obscure; aperture transversely rhombic, three times as wide as high, acutely angular at the outer and inner extremities; upper side of lip oblique and extended forward; surface bearing lines of growth directed obliquely backward. Type *P. antiquus*.

FIG. 705.—*Pseudophorus antiquus*.

antiquus, Meek, 1871, (*Trochita antiqua*), Proc. Acad. Nat. Sci. Phil., p. 82, and Ohio Pal., vol. 1, p. 221, Up. Held Gr.

PUPA, Humphrey, 1797, Museum Calounianum, and Lamarck Syst. Anim. sans Vert., p. 88. [Ety. *Pupa*, chrysalis shell.] Shell rimate or perforate, cylindrical or oblong; aperture rounded, often toothed, margins distant, mostly united by a callous lamina. Type *P. uva*.

FIG. 706.—*Pupa uva*.

bigshyl, Dawson, 1880, Am. Jour. Sci. and Arts, 3d ser., vol. 20, p. 410, Coal Meas.
vermillionensis, Bradley, 1872, Am. Jour. Sci., 3d series, vol. 4, p. 87, Coal Meas.
vetusta, Dawson, 1860, Quar. Jour. Geo. Soc., vol. 16, p. 268, and Acad. Geol., p. 383, Coal Meas.
vetusta var. *tenuistriata*, Dawson, 1880, Am. Jour. Sci. and Arts, 3d ser., vol. 20, p. 406, Coal Meas.
RAPHISTOMA, Hall, 1847, Pal. N. Y., vol. 1, p. 28. [Ety. *raphe*, seam or suture; *stoma*, mouth.] Depressed, often discoid; spine flat or nearly so; sutures close; whorls acute-angular externally and often with an angular edge to the moderate umbilicus. Type *R. striatum*.
acutum, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 235, Chazy Gr.
affinis, Foerste, 1885, Bull. Sci. Lab. Denison Univ., p. 95. Not properly defined.
angulatum, Emmons, 1856, (*Straparollus angulatus*), Am. Geol. p. 157, Calciferous Gr.
apertum, Salter, 1859, Can. Org. Rem., Decade 1, p. 12, Black Riv. and Trenton Gr.
compressum, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 309, Birdseye Gr.

FIG. 707.—*Raphistoma halli*.

halli, S. A. Miller, 1874, (*Pleurotomaria halli*), Cin. Quer. Jour. Sci., vol. 1, p. 318, Hud. Riv. Gr.
labiatum, Emmons, 1842, (*Maclurea labiatum* and Birdseye Gr.
lenticulare, Emmons, 1842, (*Pleurotomaria lenticularis*), Geo. Rep. N. Y., p. 392, and Pal. N. Y., vol. 1, p. 172, Trenton and Hud. Riv. Gr.
niagarensis, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 82, and Geo. Wis., vol. 4, p. 295, Niagara Gr.
planistria, Hall, 1847, Pal. N. Y., vol. 1, p. 30, Chazy Gr.
planistria var. *parvum*, Hall, 1847, Pal. N. Y., vol. 1, p. 30, Chazy Gr.

FIG. 708.—*Raphistoma lenticulare*.

eum Calomni-
t. Anim. sans
rysalis shell.]
rate,
per-
phed,
ited
e P.

Am.
ser.,
a.
872,
vol. 4, p. 87,
Acad. Jour. Geol.
Acad. Geol., p.

Dawson, 1880,
3d ser., vol.

N. Y., vol. 1, p.
suture; stoma,
discoid; spire
close; whorls
and often with
oderate umbili-

1, 1877, U. S.
vol. 4, p. 235,

Sci. Lab. Deni-
properly defined.
(Straparollus
157, Calcifer-

a. Org. Rem.,
div. and Tren-

86, Bull. Am.
p. 309, Birds-

A. Miller, 1874,
tomaria halli,
ser. Jour. Sci.,
p. 318, Illud.

Emmons,
Maclurea labi-
312, Calcifer-

n. Org. Rem.,
div. and Tren-

p. 708.—Raphisto-
ma lenticulare.

r. Wis., p. 82,
4, p. 295, Ni-

N. Y., vol. 1,

hall, 1847, Pal.
ty Gr.

prævium, Whitfield, 1889, Bull. Am.
Mus. Nat. Hist., vol. 2, p. 52, Calcifer-
ous Gr.
rotuliforme, Meek, 1870, Proc. Acad. Nat.
Sci., p. 61, and U. S. Geo. Sur. 40th
Parallel, vol. 4, p. 18, Calciferous Gr.
rotuloides, Hall, 1847, (Pleurotomaria ro-
tuloides,) Pal. N. Y., vol. 1, p. 173, Tren-
ton Gr.
stamineum, Hall, 1847, Pal. N. Y., vol.
1, p. 29, Chazy Gr.
striatum, Emmons, 1842, (Maclurea stri-
ata,) Geo. Rep. N. Y., p. 312, and Pal.
N. Y., vol. 1, p. 28, Chazy Gr.

FIG. 709.—Raphistoma striatum.

subplanum, Shumard, 1863, Trans. St.
Louis Acad. Sci., vol. 2, p. 106, Calcifer-
ous Gr.
subtilstriatum, Hall, 1847, (Pleurotomaria
subtilstriata,) Pal. N. Y., vol. 1, p. 172,
Trenton Gr.
trochiscum, Meek, 1870, (Euomphalus
trochiscus,) Proc. Acad. Nat. Sci., p. 61,
and Geo. Sur. W. 100th Mer., vol. 4, p.
77, Calciferous or Trenton Gr.

ROTELLA, Lamarck, 1822, Hist. Nat. Anim. sans
Vert., vol. 7, p. 6. [Ety.
diminutive of *rota*, a
wheel.] Lenticular, pol-
ished; spire depressed;
base callous, lingual
teeth 13; uncini, num-
erous, subequal. Type *R. vestiaria*.
verruculifera, White, 1882, Rep. Invert.
Foss., New Mexico, p. xxxi, Coal
Meas.

FIG. 710.—Rotella
vestiaria.

SCÆVOGYRA, Whitfield, 1878, Ann. Rep.
Geo. Sur. Wis., p. 61, and Geo. Wis.,
vol. 4, p. 198. [Ety. *scævus*, toward
the left; *gurus*, circle.] Sinistral, spire
elevated, volutions rounded; umbilicus
open, broad, no callus; peristome en-
tire, uniting with the volution on the
inner side and spreading externally.
Type *S. swezeyi*.
elevata, Whitfield, 1878, Ann. Rep. Geo.
Sur. Wis., p. 62, and Geo. Wis., vol. 4,
p. 199, Low. Mag. Gr.
obliqua, Whitfield, 1878, Ann. Rep. Geo.
Sur. Wis., p. 63, and Geo. Wis., vol. 4, p.
199, Low. Mag. Gr.
swezeyi, Whitfield, 1878, Ann. Rep. Geo.
Sur. Wis., p. 62, and Geo. Wis., vol. 4,
p. 198, Low. Mag. Gr.

SCALITES, Emmons, 1842, Geo. Rep. N. Y.,
p. 312. [Ety. *scala*, staircase.] Tur-
binate, whorls flat above, turritid, pro-

duced below; no umbilicus; form
elongate. Type *S. angulatus*.

angulatus, Emmons,
1842, Geo. Rep.
N. Y., p. 312, and
Pal. N. Y., vol. 1,
p. 27, Chazy Gr.

SCOLIOSTOMA, Braun,
1838, Neues Jahr.
Min. Geo. Geol. Pe-
tref., p. 298. [Ety.
skolios, curved;
stoma, mouth.]
Small, upper part
pupiform; aperture
extended, curved
outward. Type *S.*
dannenbergi.

americana, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 195, Low. De-
vonian.

Solarium, Lamarck, 1801, Syst. An. sans Vert.
Not Palæozoic.

leui, one of Troost's catalogue names.

SOLENISCUS, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 467. [Ety.
soleniskos, little channel or gutter.] Fus-
iform, pointed; whorls nearly flat, last
one contracted and produced below
into a straight canal; surface smooth
or obscurely marked by lines of growth;
aperture narrow; outer lip thin, en-
tire; inner lip thickened and bearing
a more or less distinct revolving fold;
columella straight, inperforate. Type
S. typicus.

brevis, White, 1882, Rep. Invert. Foss.,
New Mex., p. xxvii, Coal Meas.

fusiformis, Hall, 1858, (Macrocheilus fus-
iforme,) Geol. of Iowa, p. 718, Coal
Meas.

hallanus, Geinitz, 1866, (Macrocheilus
hallanum,) Carb. and Dynas in Neb. p.
6, Coal Meas.

helicoides, Sowerby,
1829, (Ampullaria
helicoides), Min.
Conch., vol. 6, p. 40,
Coal Meas.

klipparti, Meek, 1872,
(Macrocheilus klip-
parti,) Proc. Acad.
Nat. Sci., vol. 24, p.
328, and Ohio Pal.,
vol. 2, p. 346, Low.
Coal Meas.

newberryi, Stevens,
1858, (Loxonema
newberryi), Am.
Jour. Sci. and Arts,
2d ser., vol. 25, p.
259, and Geo. Sur.
Ill., vol. 5, p. 594,
Coal Meas.

paludiformis, Hall,
1858, (Macrocheilus
paludiformis), Geo.

of Iowa, p. 719, Coal Meas.
planus, White, syn. for *S. newberryi*.

FIG. 711.—Scalites an-
gulatus.

FIG. 712.—Soleniscus
klipparti.

texanus, Shumard, 1859, (Macrocheilus texanum,) Trans. St. Louis Acad. Sci., vol. 1, p. 402, Coal Meas.

FIG. 713.—*Soleniscus typicus*.

typicus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 467, and Geo. Sur. Ill., vol. 2, p. 384, Up. Coal Meas.

ventricosus, Hall, 1858, (Macrocheilus ventricosum,) Geo. Sur. Iowa, p. 718, Coal Meas.

STRAPAROLLINA, Billings, 1865, Pal. Foss., vol. 1, p. 223. [Ety. from the resemblance to shells of the genus *Straparollus*.] Shell turbinate, with round or obscurely angulated whorls; aperture nearly circular, sometimes with a notch in the inner lower angle of the lip. Type *S. pelagica*.

asperostriata, Billings, 1860, (Straparollus asperostriatus,) Can. Nat. and Geol., vol. 5, p. 162, Black Riv. Gr.

circe, Billings, 1860, (Straparollus circe,) Can. Nat. and Geol., vol. 5, p. 161, Black Riv. Gr.

eurydice, Billings, 1860, (Straparollus eurydice,) Can. Nat. and Geol., vol. 5, p. 162, Black Riv. Gr.

pelagica, Billings, 1865, Pal. Foss., vol. 1, p. 223, Quebec Gr.

remota, Billings, 1874, Pal. Foss., vol. 2, p. 70, Up. arollina pelagica. Taconic.

STRAPAROLLUS, Montfort, 1810, Conch. Syst., vol. 2, p. 174. [Ety. *strabos*, turned about.] Discoid, depressed conic, smooth or transversely striated; whorls rounded; umbilicus wide, exposing the whorls; mouth indented by the penultimate whorl; peritreme simple, thin, most so on the left side. Type *S. dionysii*.

ammon, White and Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 307, Marshall Gr.

angulatus, see *Raphistoma angulatum*.

asperostriatus, see *Straparollina asperostriata*.

barrisi, Winchell, 1863, Proc. Acad. Nat. Sci. Phil., p. 20, Marshall Gr.

✓ canadensis, Billings, 1861, Can. Jour., vol. 6, p. 359, Up. Held. Gr.

✓ circe, see *Straparollina circe*.

✓ clymenioides, Hall, 1862, (Euomphalus clymenioides,) 15th Rep. N. Y. Mus. Nat. Hist., p. 54, and Pal. N. Y., vol. 5, pt. 2, p. 62, Up. Held. Gr.

cornudanus, Shumard, 1859, Trans. St. Louis Acad. Sci., vol. 1, p. 400, Coal Meas.

✓ crenulatus, Whiteaves, 1884, Pal. Foss., vol. 3, p. 21, Guelph Gr.

✓ cyclostomus, Hall, 1858, (Euomphalus cyclostomus,) Geo. Sur. Iowa, p. 516, Ham. Gr.

daphne, Billings, 1862, Pal. Foss., vol. 1, p. 160, Guelph Gr.

eurydice, see *Straparollina eurydice*.

hecale, Hall, 1876, (Euomphalus hecale,) Illust. Devon. Foss., pl. 16, Chemung Gr.

hecale var. corpulens, Hall, 1876, (Euomphalus hecale var. corpulens,) Illust. Dev. Foss., pl. 27, Chemung Gr.

hippolyta, Billings, 1862, Pal. Foss., vol. 1, p. 160, Guelph Gr.

inops, Hall, 1876, (Euomphalus inops,) Illust. Devonian Foss., pl. 16, Up. Held. Gr.

labiatus, see *Raphistoma labiatum*.

lens, Hall, 1860, (Euomphalus lens) 13th Rep. N. Y. Mus. Nat. Hist., p. 109, Kinderhook Gr. FIG. 715.—*Straparollus hippolyta*.

macromphalus, Winchell, 1863, Proc. Acad. Nat. Sci. Phil., p. 20, Marshall Gr. *magnificus*, Shumard, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 110, Carboniferous. Too poorly defined for recognition.

minnesotensis, see *Euomphalus minnesotensis*.

mopsus, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 390, Niagara Gr.

newarkensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 187, Devonian.

niagarensis, Hall & Winchell, 1875, Ohio Pal., vol. 2, p. 144, N. Gr.

ophirensis, Hall & Winchell, 1877, (Euomphalus ophirensis,) U. S. Geo. 40th Parallel, vol. 4, p. 261, Waverly Gr.

pernodorus, see *Euomphalus pernodorus*.

planispira, Hall, 1858, (Euomphalus planispira,) Trans. Alb. Inst., vol. 4, p. 20, and Bull. Am. Mus. Nat. Hist., p. 70, Warsaw Gr.

primordialis, see *Ophileta primordialis*.

quadrivolvus, Hall, 1858, (Euomphalus quadrivolvus,) Trans. Alb. Inst., vol. 4, p. 19, and Bull. Am. Mus. Nat. Hist., p. 71, Warsaw Gr.

rudis, Hall, 1876, (Euomphalus rudis,) Illust. Dev. Foss., pl. 16, and Pal. N. Y., vol. 5, pt. 2, p. 58, Ham. Gr.

sanctisabae, Roemer, 1852, (Euomphalus sanctisabae,) Kreid. von Texas, p. 91, Silurian.

similis, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 145, and Geo. Sur. Ill., vol. 2, p. 285, St. Louis Gr.

similis var. planus, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 146, and Geo. Sur. Ill., vol. 2, p. 286, St. Louis Gr.

sinuatus, Hall, 1859, (Euomphalus sinuatus,) Pal. N. Y., vol. 3, p. 340, Low. Held. Gr.

✓ spergenensis, Hall, 1858, (Euomphalus spergenensis,) Trans. Alb. Inst., vol. 4, p. 19, and Bull. Am. Mus. Nat. Hist., p. 69, Warsaw Gr.

✓ spergenensis var. planorbiformis, Hall, 1858, (Euomphalus spergenensis var.

Foss., vol. 1,
 eurydice.
 phalus hecale,
 pl. 16, Che.
 1876, (Euom-
 phalus), Illust.
 ing Gr.
 pal. Foss., vol.
 phalus inops,
 pl. 16, Up.

715.—*Strophostylus*
hippolytus

1863, Proc.
 Marshall Gr.
 Trans. St.
 p. 110, Carbon-
 iferous for recog-

phalus minneso-

pp. N. Y. Mus.
 a Gr.

Monogr. U. S.
 Devonian.

ld, 1875, Ohio
 Gr.

1877, (Euom-
 phalus), 40th

S. Geo. 40th
 averly Gr.

s pernodosus.
 phalus plan-

, vol. 4, p. 20,
 t. Hist., p. 70,

rimordialis.

(Euomphalus
 Inst., vol. 4,

s. Nat. Hist.,

phalus rudis,
 and Pal. N. Y.,

ir.

(Euomphalus
 Texas, p. 91,

1861, Proc.

145, and Geo.

Louis Gr.

Worthen, 1861,
 l., p. 146, and

St. Louis Gr.

phalus sin-

p. 340, Low.

(Euomphalus
 Inst., vol. 4,

Nat. Hist., p.

planorbiformis, Trans. Alb. Inst., vol.
 4, p. 20, and Bull. Am. Mus. Nat. Hist.,
 p. 70, Warsaw Gr.

spirorbis, Hall, 1859, (Euomphalus spiror-
 bis,) 13th Rep. N. Y. Mus. Nat. Hist.,
 p. 109, Kinderhook Gr.

subplanus, Hall, 1852, (Euomphalus sub-
 planus,) Stans. Ex. to Gt. Salt Lake,
 p. 414, Coal Meas.

subquadratus, see Euomphalus subquad-
 ratus.

subrugosus, see Euomphalus subrugosus.

subumbilicatus, Worthen, (In press)
 Geo. Sur. Ill., vol. 8, p. 142, Kaskas-
 kia Gr.

umbilicatus, Meek & Worthen, 1860, (Eu-
 omphalus umbilicatus,) Proc. Acad.
 Nat. Sci. Phil., p. 492, and Geo. Sur. Ill.,
 vol. 2, p. 362, Coal Meas.

utahensis, Hall & Whitfield, 1877, U. S.
 Geo. Expl., 40th parallel, vol. 4, p. 259,
 Waverly Gr.

valvatiiformis, Shumard, 1863, Trans. St.
 Louis Acad. Sci., vol. 2, p. 105, Calcifer-
 ous Gr.

varsoviensis, Worthen (In press) Geo.
 Sur. Ill., vol. 8, p. 142, Keokuk Gr.

whitneyi, see Omphalotrochus whitneyi.

STREPTAXIS, Gray, 1837, Mag. Nat. Hist., p.
 484, [Ety. *streptos*, twisted; *axis*, axis.]

Shell ovate or oblong; when young, sub-
 hemispherical, deeply umbilicated, with
 rapidly enlarging whorls; at length the
 penultimate whorl is bent toward the

right and dorsal side of the axis and the
 umbiliculus become compressed and
 often nearly closed; the mouth lunate;
 the edge slightly thickened and re-

flexed, and often with a single tooth on
 the outer side of the inner or hinder
 lip. Type *S. corboides*. Not a Palae-
 ozic genus.

whitfieldi, Meek, 1871, Proc. Acad. Nat.
 Sci. Phil., p. 173, and Geo. Sur. Ill., vol.
 5, p. 596, Coal Meas.

STROPHITES, Dawson, 1880, Am. Jour. Sci.
 and Arts, 3d ser., vol. 20, p. 413. [Ety.

from the genus *Strophia*.] Shell resem-
 bling the modern *Strophia*, conical;
 apex obtuse; whorls four or more;
 surface covered with sharp vertical

ridges, separated by spaces three times
 as wide. Type *S. grandævus*.

grandævus, Dawson, 1880, Am. Jour. Sci.
 and Arts, 3d ser., vol. 20, p. 413, De-
 vonian.

STROPHOSTYLUS, Hall, 1859, Pal. N. Y., vol.
 3, p. 303. [Ety. *strophe*, turning round;
stylos, column.] Subglobose or ovoid

globose; spire small with a large ven-
 tricose body whorl; outer lip thin, not
 reflected; columella twisted or spirally
 grooved within, not reflected; no um-
 bilicus; aperture somewhat round,

ovate or transversely broad oval. Type
S. elegans.

andrewsi, Hall, 1859, Pal. N. Y., vol. 3,
 p. 472, Oriskany sandstone.

cancellatus, Meek & Worthen, 1868, Geo.
 Sur. Ill., vol. 3, p. 404, Oriskany sand-
 stone.

FIG. 716.—*Strophostylus cancellatus*. B. Surface
 markings enlarged.

cyclostomus, Hall, 1863, Trans. Alb. Inst.,
 vol. 4, p. 218, Niagara Gr.

cyclostomus var. *disjunctus*. Hall, 1879,
 28th Rep. N. Y. Mus. Nat. Hist., p. 177,
 Niagara Gr.

depressus, Hall, 1859, Pal. N. Y., vol. 3,
 p. 306, Low. Held. Gr.

elegans, Hall, 1859, Pal. N. Y., vol. 3, p.
 304, Low. Held. Gr.

expansus, Conrad, 1841, (Platyceras ex-
 pansum,) Ann. Rep. N. Y., p. 55, and
 Pal. N. Y., vol. 3, p. 470, Oriskany sand-
 stone.

fitchi, Hall, 1859, Pal. N. Y., vol. 3, p. 306,
 Low. Held. Gr.

globosus, Hall, 1859, Pal. N. Y., vol. 3, p.
 305, Low. Held. Gr.

matheri, Hall, 1859, Pal. N. Y., vol. 3, p.
 471, Oriskany sandstone.

obliquus, Nicholson, 1874, Rep. Pal. Ont.,
 p. 119, Up. Held. Gr.

obtusus, Hall, 1859, Pal. N. Y., vol. 3, p.
 305, Low. Held. Gr.

ovatus, Nicholson, 1874, Rep. Pal. Ont.,
 p. 118, Up. Held. Gr.

rotundatus, Hall, 1859, Pal. N. Y., vol. 3,
 p. 307, Low. Held. Gr.

subglobosus, Nicholson, 1874, Rep. Pal.
 Ont., p. 118, Up. Held. Gr.

transversus, Hall, 1859, Pal. N. Y., vol. 3,
 p. 470, Oriskany sandstone.

unicus, Hall, 1862, 15th Rep. N. Y. Mus.
 Nat. Hist., p. 41, and Pal. N. Y., vol. 5,
 pt. 2, p. 30, Schoharie grit.

varians, Hall, 1876, Illust. Devonian
 Foss., pl. 11, and Pal. N. Y., vol. 5, pt.
 2, p. 31, Up. Held. Gr.

Stylifer, Broderip, 1829, in Sowerby, Gen.
 Shells.

primigenia, see Macrochilina primigenia.

SUBULITES, Conrad, 1847, Pal. N. Y., vol. 1,
 p. 182. [Ety. *subula*, an awl.] Sub-

ulate, volutions wide, suture oblique;
 aperture very elongate, narrow, pointed
 above, but wider below. Type *S. elon-*
gatus.

abbreviatus, Hall, 1850, 3d Rep. N. Y.
 Mus. Nat. Hist., p. 180, Trenton Gr.

brevis, Winchell & Marcy, 1865, Mem.
 Bost. Soc. Nat. Hist., vol. 1, p. 100, Ni-
 agara Gr.

calciferus, Billings, 1859, Can. Nat. & Geo.,
 vol. 4, p. 360, Calciferous Gr.

compactus, Whiteaves, 1884, Pal. Foss. vol. 3, p. 16, Guelph Gr.

FIG. 717.—*Subulites calciferus*.

daphne, Billings, 1865, Pal. Foss., vol. 1, p. 223, Quebec Gr.

FIG. 718.—*Subulites elongatus*.

elongatus, Emmons, 1842, Geo. Rep. N. Y., p. 392, and Pal. N. Y., vol. 1, p. 182, Trenton Gr.

gracilis, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 116, Niagara Gr. *inflatus*, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 47, and Geo. Sur. Ill., vol. 6, p. 495, Galena Gr.

notatus, Billings, 1866, Catal. Sil. Foss. Antic., p. 54, Anticosti Gr.

obesus, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 318, Birdseye Gr.

parvulus, Billings, 1862, Pal. Foss., vol. 1, p. 36, Black Riv. Gr.

psyche, Billings, 1865, Pal. Foss., vol. 1, p. 188, Quebec Gr.

richardsoni, Billings, 1857, Rep. of Progr., Geo. Sur. Can., p. 306, Hnd. Riv. Gr.

terebriiformis, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 141, Niagara Gr.

ventricosus, Hall, 1852, Pal. N. Y., vol. 2, p. 347, Niagara and Guelph Gr.

TRACHYDOMIA, Meek & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 364. [Ety. *trachys*, rough; *dema*, house.] In

form like *Naticopsis*, but distinguished by having the surface ornamented with regularly disposed nodes. Type *T. nodosum*.

hollidayi, Meek & Worthen, 1860, (Naticopsis *hollidayi*.) Proc. Acad. Nat. Sci. Phil., p. 463, and Geo. Sur. Ill., vol. 2, p. 367, Low. Coal Meas.

nodosum, Meek & Worthen, 1860, (Naticopsis *nodosum*.) Proc. Acad. Nat. Sci. Phil., p. 463, and Geo. Sur. Ill., vol. 2, p. 366, Low. Coal Meas.

nodulosum, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 8, and Geo. Sur. Ill., vol. 8, p. 146, Coal Meas.

TREMANOTUS, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 347. [Ety. *trema*, hole; *notos*, back.] Distinguished from *Bucania* by a single row of hollow spines upon the back of the last whorl. In casts the spines are usually broken off, and hence Carpenter argues they never had spines. Type *T. chicagoensis*.

alpheus, Hall, 1864, 10th Rep. N. Y. St. Mus. Nat. Hist. Syn. for *T. chicagoensis*.

chicagoensis, McChesney, 1860, (*Bucania chicagoensis*.) New Pal. Foss., p. 69, Niagara Gr.

trigonostoma, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 146, Niagara Gr.

Trochita, Schumacher, 1817, Essai N. Syst., p. 184. [Ety. *trochus*, wheel.] Not a Palaeozoic genus.

antiqua, see *Pseudophorus antiquus*. *carbonaria*, Meek, 1866, Proc. Acad. Nat. Sci., p. 270, Kaskankia Gr. Not recognized.

TROCHONEMA, Salter, 1859, Can. Org. Rem., Decade 1, p. 27. [Ety. *trochus*, a wheel; *nema*, a thread.] Turbinate, thin, of few angular whorls; strong concentric ridges, crossed by oblique lines of growth; umbilicus wide, open; inner lip thin, scarcely reflected; peritreme complete. Type *T. umbilicatum*.

beloitense, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 74, and Geo. Wis., vol. 4, p. 212, Trenton Gr.

beachi, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 74, and Geo. Wis., vol. 4, p. 213, Trenton Gr.

emacreratum, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 193, Ham. Gr.

exile, Whitfield, 1880, Bull. Am. Mus. Nat. Hist., vol. 2, p. 57, Calciferous Gr.

fatua, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 394, Niagara Gr.

halli, Hall, 1861, (Pleuronomaria *halei*.) Geo. Sur. Wis., p. 34, Niagara Gr.

inornatum, Whiteaves, 1884, Pal. Foss., vol. 3, p. 19, Guelph Gr.

meekianum, n. sp. Up. Held. Gr. at Marblehead, Ohio. Proposed instead of *T. tricarinatum*, Meek, 1871, Proc. Acad. Nat. Sci., p. 82, and Ohio Pal., vol. 1, p. 218, which was preoccupied.

t distinguished
namented with
les. Type T.

en, 1860, (Nati-
Acad. Nat. Sci.
Sur. Ill., vol. 2,
en, 1860, (Nati-
Acad. Nat. Sci.
Sur. Ill., vol. 2,

4, Bull. No. 2,
p. 8, and Geo.
Coal Meas.
th Rep. N. Y.
7. [Ety. *trema*,
ingnished from
row of hollow
the last whorl.
usually broken
er argues they
pe T. chicago-

Rep. N. Y. St.
for T. chicago-

1860, (Bucania
Foss., p. 69, Ni-

Whitfield, 1875,
Niagara Gr.
Esai N. Syst.,
[Ety.] Not a

antiquus.
oc. Acad. Nat.
Gr. Not rec-

an. Org. Rem.,
chus, a wheel;
uate, thin, of
strong concen-
blique lines of
, open; inner
ed; peritreme
ilicatum.

8, Ann. Rep.
nd Geo. Wis.,
r.
m. Rep. Geo.
o. Wis., vol. 4,

field, 1872, 24th
Hist., p. 193,

ll. Am. Mus.
Calciferous Gr.
p, N. Y. Mus.
a Gr.
maria hnei.)
agara Gr.
4, Pal. Foss.,

Held. Gr. at
posed instead
k, 1871, Proc.
nd Ohio Pal.
oreoccupied.

nana, Foerste, 1885, Bull. Sci. Lab. Denison
Univ., p. 94. Not properly defined.
pauper, Hall, syn. for P. hali.
pauper var. ohioense, Hall & Whitfield,
1875, Ohio Pal., vol. 2, p. 144, Niagara Gr.
rectilatera, Hall & Whitfield, 1872, 24th
Rep. N. Y. Mus. Nat. Hist., p. 193, Up.
Held. Gr.
tricarinarum, Billings, 1859, Can. Nat. &
Geo., vol. 4, p. 356, Calciferous Gr.
tricarinata, see Trochonema meckanum.

FIG. 719.—Trochonema umbilicatum.

umbilicatum, Hall,
1847, (Pleuroto-
maria umbilica-
ta,) Pal. N. Y., vol.
1, p. 43, Chazy to
Ind. Riv. Gr.
yandellianum, Hall
& Whitfield, 1872,
24th Rep. N. Y.
Mus. Nat. Hist.,
p. 194, Up. Held.
Gr.
Trochus, Adanson,
1757, Voy. Sene-
gal. [Ety. *trochus*, a hoop.] Not a
Palaeozoic genus.
huronensis, Castelnau, 1843, Syst. Sil., p.
35. Not recognized.
missouriensis, see Pleurotomaria missouri-
ensis.

TRYBLIDIUM, Lindstrom, 1880, Fragmenta
Silurica, p. 15. [Ety. *trublion*, a cup.]
Patelliform, obovate, acuminate anteri-
orly, enlarged posteriorly; muscular
scars in six disconnected pairs arranged
in an oblong circle open toward the
front. Type T. reticulatum.

acutum, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 45, Calciferous Gr.
canadense, Whiteaves, 1884, Pal. Foss.,
vol. 3, p. 31, Guelph Gr.
conicum, Whitfield, 1886, Bull. Am. Mus.
Nat. Hist., vol. 1, p. 306, Birdseye Gr.
erato, Billings, 1862, (Metoptoma erato,)
Pal. Foss., vol. 1, p. 39, Black Riv. Gr.
eubule, Billings, 1862, (Metoptoma eu-
bule,) Pal. Foss., vol. 1, p. 38, Calcifer-
ous and Black Riv. Gr.

FIG. 720.—Tryblidium nycteis.

hyrie, Billings, 1862, (Metoptoma hyrie,)
Pal. Foss., vol. 1, p. 87, Quebec Gr.
niobe, Billings, 1862, (Metoptoma niobe,)
Pal. Foss., vol. 1, p. 37, Calciferous Gr.
nycteis, Billings, 1862, (Metoptoma nycteis,
Pal. Foss., vol. 1, p. 38, Calciferous Gr.

ovale, Whitfield, 1886, Bull. Am. Mus.
Nat. Hist., vol. 1, p. 305, Birdseye Gr.
ovatum, Whitfield, 1886, Bull. Am. Mus.
Nat. Hist., vol. 1, p. 305, Birdseye Gr.
pileolum, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 46, Calciferous Gr.
simplex, Billings, 1865, (Metoptoma sim-
plex,) Pal. Foss., vol. 1, p. 346, Calcifer-
ous Gr.

TURBO, Klein, 1753, Tent. Meth. Ostr. [Ety.
turbo, top.] Shell thick, ovate; body
whorl rounded, ventricose; spire small,
of several convex whorls, pointed; sur-
face spirally grooved or nodulated;
aperture large, nearly circular, slightly

FIG. 721.—Turbo marmoratus.

produced and
broadly
rounded in
front, more
or less mod-
ified by the
preceding
whorl; outer
and inner
lips thin; op-
erculum
thick, shelly,
rugged with-
out, flattened
and spirally
sulcated

within. Type T. marmoratus. Not an
American Palaeozoic genus. The spe-
cies left here is, for want of material,
to refer them where they belong.
bicaratus, Troost, 1840. Not defined.
dilucula, see Holoepa dilucula.

guadalupensis, Shumard, 1859, Trans St.
Louis Acad. Sci., vol. 1, p. 398, Per-
mian Gr.

huronensis, Castelnau, 1843. Not recog-
nized.
lineatus, see Pleurotomaria lineata.

obesus, Shumard, 1858, Trans. St. Louis
Acad. Sci., vol. 1, p. 202, Up. Coal
Meas.

(?) obscura, see Holoepa
obscura.

shumardi, see Platys-
toma shumardi.

tabulata, see Pleuroto-
maria tabulata.

tennesseensis, see Cyclo-
nema tennesseense.

texanus, Shumard,
1859, Trans. St. Louis
Acad. Sci., vol. 1, p.
400, Coal Meas.

Turbonilla, Leuch, 1826,
Risso Eur. Merid.

4. [Ety. diminutive
of Turbo, a genus.]
Not a Palaeozoic
genus.

swallowana, see Aclis-
ina swallowana.

TURRITELLA, Lamarck, 1801, Syst. An. sans
Vert., p. 39. Not a Palaeozoic genus.

Type T. imbricata.

FIG. 722.—Turritella imbricata.

schohariensis, Castelnau, 1843, Syst. Sil., p.

35. Not recognized.
stevensana, Meek &
Worthen, 1866, Geo.
Sur. Ill., vol. 2,
p. 382, Up. Coal
Meas.

Xenophora, Fischer, 1806,
Museum Demidovianum,
p. 213. Not an
American Palæozoic
genus.

antiqua, see *Pseudophorus antiquus*.

ZAPTYPHIUS, Walcott,
1884, Monogr. U. S.
Geo. Sur., vol. 8, p.
263. Shell minute,
elongate; aperture
large, oblong, nearly
vertical; outer lip
thin; collumellar lip
reflected, plicated;
surface marked by
slightly oblique
vertical striæ.

Type *Z. carbonarius*.

FIG. 723.—*Zaptypius carbonarius*.

carbonarius, Walcott, 1884, Monogr. U. S.
Geo. Sur., vol. 8, p. 263, Subcarbonif-
erous.

FIG. 724.—*Zonites priscus*.

ZONITES, Montfort, 1810, Conch. Syst.,
vol. 2, p. 282. [Ety. *zone*, belt.] A
coiled shell, closely resembling a *Helix*,
having an open umbilicus; the sur-
face is transversely sculptured. Type
Z. algireus.

priscus, Carpenter, 1867, Quar. Jour. Geo.
Soc., vol. 23, p. 331, and Acadian Geol.,
p. 385, Coal Meas.

CLASS CEPHALOPODA.

[Ety. *kephale*, head; *pous*, foot.]

THE animals of this class are all marine, and they reach a higher state of animal development than any other marine group among the Mollusca. Some of them have a rudimentary, cartilaginous, cephalic skeleton, which indicates superiority over other marine Mollusca. The locomotive organs consist of arms surrounding the head, furnished with sucking cups that take a firm hold on other objects. Many have fins, and all can propel themselves by the forcible expulsion of water from the respiratory chamber. They swim rapidly, creep on the bottom of the sea, and are very predatory in their habits. The body is short, thick, and symmetrical, with branchiæ on both sides.

The Palæozoic fossils of this Class belong to the Order Tetrabranchiata (four-gilled), which is represented in tropical seas by the *Nautilus*. The shells are straight, as in the family *Orthoceratidæ*; curved, as in *Cyrtoceratidæ*; discoid, as in the *Gyroceratidæ* and *Trocholitidæ*; spiral, as in the *Trochoceratidæ*; involute, as in the *Nautilidæ*; or involute and having lobed sutures, as in the *Goniatitidæ*. Internally the shell is divided into numerous chambers by partitions, or septa, the animal inhabiting the last chamber, and retaining connection through the preceding chambers by a tube, or siphuncle, but having no connection with the interior of the several chambers after having cut itself off by the secretion of the shelly septa. The outlines of the septa are called sutures, and in *Goniatites* the elevations of the folded sutures are called saddles, and the intervening depressions lobes.

184, Monogr. U. S.
263, Subcarbonif.

priscus.

Conch. Syst.,
zone, belt.] A
sembling a Helix,
bilicus; the sur-
culptured. Type

Quar. Jour. Geol.
Acadian Geol.,

gher state of an-
lusca. Some of
indicates superi-
arms surround-
on other objects.
pulsion of water
ttom of the sea,
nd symmetrical,

ranchiata (four-
The shells are
æ; discoid, as in
æ; involute, as
oniatitidæ. In-
s, or septa, the
ugh the preced-
on of the shelly
atites the eleva-
pressions lobes.

Each septum began to form at the circumference of the shell, and slowly approached the siphuncle as the animal moved forward in the body chamber. The siphuncle, being a point of muscular attachment, was not vacated by the animal between any two septa until the anterior one had been firmly closed by attachment to the siphuncle, forming a chamber of support.

The fossil shells are very thin in proportion to their size. They are not porous, like those of the Brachiopoda; nor horny, like the Crustacea; nor of the same composition as the Gasteropoda or Lamellibranchiata. Generally the exterior shell is destroyed, even when the associated shells of other classes are well preserved. Sometimes the shell appears as if it had melted and run together, or run down upon the siphuncle. Such molecular change will occur in one part of a specimen while another part is unchanged. The general form of the shell is of family importance. The shape of the siphuncle and the external markings are of generic importance. We recognize the following families:

FAMILY ASCOCERATIDÆ.—Ascoceras.

FAMILY CYROCERATIDÆ.—Cyrtoceras, Cyrtocerina, Oncoceras.

FAMILY DISCOSORIDÆ.—Discosorus.

FAMILY ENDOCERATIDÆ.—Cameroceras, Colpoceras, Endoceras.

FAMILY GOMPHOCERATIDÆ.—Gomphoceras.

FAMILY GONIATITIDÆ.—Goniatites.

FAMILY GYROCERATIDÆ.—Gyroceras.

FAMILY LITUITIDÆ.—Lituites.

FAMILY NAUTILIDÆ.—Discites, Nautilus, Pteronutilus, Solenochilus, Temnochilus, Trematodiscus.

FAMILY ORTHOCERATIDÆ.—Actinoceras, Bactrites, Gonioceras, Huronia, Orthoceras, Orthoceras, Trematoceras.

FAMILY PHRAGMOCERATIDÆ.—Phragmoceras, Streptoceras.

FAMILY PILOCERATIDÆ.—Piloceras.

FAMILY TROCHOCERATIDÆ.—Trochoceras.

FAMILY TROCHOLITIDÆ.—Trocholites.

FAMILY UNCERTAIN.—Petalichnus, Sericichnus, Teratichnus, Trachomatichnus.

ACTINOCERAS, Bronn, 1837, Lethaea Geognostica, p. 97. [Ety. *aktin*, ray; *keras*, horn.] Exterior like Orthoceras; siphuncle very large, inflated between the chambers, and connected with a slender central tube by radiating plates. Type *A. bigsbyi*, *A. richardsoni*, and *A. lyoni*. The genus was established before the species were defined.

beaudanti, Castelnau, 1843, Système Silurien, p. 31. Not recognized.

beaudanti, Castelnau, 1843, Système Silurien, p. 32. Not recognized.

bigsbyi, Stokes, 1840, Trans. Geo. Soc., 2d series, vol. 5, p. 707, Chazy Gr.

blainvilliei, Castelnau, 1843, Système Silurien, p. 31. Not recognized.

cordieri, Castelnau, 1843, Système Silurien, p. 31. Not recognized.

deshayesi, Castelnau, 1843, Système Silurien, p. 32. Not recognized.

dufresnoyi, Castelnau, 1843, Système Silurien, p. 32. Not recognized.

inops, Dawson, 1868,

Acad.

Geol.

p.

314, Carb.

lyoni, Stokes,

1840,

Trans.

Geol. Soc.,

vol. 5, p.

707, Black

Riv. Gr.

richardsoni,

Stokes,

1840,

Trans.

Geol. Soc.,

2d series, vol. 5, p. 708, Black Riv. Gr.

simmsi, Stokes, 1840, Trans. Geo. Soc., 2d

series, vol. 5, p. 708, Sil.

FIG. 725.—*Actinoceras richardsoni*.

Ammonites bellicosus, Morton, 1836, Am. Jour. Sci. and Arts, vol. 29, Coal Meas. Not recognized.

colubrellus, see *Goniatites colubrellus*.

hildrethi, see *Goniatites hildrethi*.

ASCO CERAS, Barrande, 1855, Bull. de la Soc. Geol. de France, vol. 12, 2d ser., p. 157.

[Ety. *askos*, leather bottle; *keras*, horn.] Chambers behind the living one short and rapidly tapering; living chamber long and constricted near the aperture; aperture somewhat T-shaped. Type A. bohemicum.

anticostiense, Billings, 1866, Catal. Sil. Foss. Antic., p. 60, and Pal. Foss., vol. 1, p. 164, fig. 148b, Anticosti Gr.

canadense, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 310, Hud. Riv. Gr. This species is made the type of the genus Billingsites by Hyatt.

FIG. 726.—*Ascoceras canadense*.

newberryi, Billings, 1862, Pal. Foss., vol. 1, p. 163, Hud. Riv. and Anticosti Grs. *southwelli*, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 151, Niagara Gr. *townsendi*, Whiteaves, 1884, Pal. Foss., vol. 3, p. 41, Guelph Gr.

BACTRITES, Sandberger, 1841, Leonh. u. Bronn's Jahrb., p. 240. [Ety. *baktron*, staff.] Shell long, straight, gradually tapering, many-chambered; sutures curve abruptly backward over the siphuncle, forming "the dorsal lobe" similar to that of a *Goniatites*. Type B. carinatus.

✓ *clavus*, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 316, Ham. Gr.

CAMEROCERAS, Conrad, 1842, Jour. Acad. Nat. Sci. Phil., vol. 8, p. 267. [Ety. *kamara*, chamber; *keras*, horn.] Shell straight, and in form like *Endoceras*; siphuncle

marginal, and obliquely annulated at the junction of the septa. Type C. trentonense.

trentonense, Conrad, 1842, Jour. Acad. Nat. Sci. Phil., vol. 8, p. 267, and Pal. N. Y., vol. 1, p. 221, Trenton Gr.

COLPO CERAS, Hall, 1850,

3d Rep. N. Y. Mus. Nat. Hist., p. 181.

[Ety. *kolpos*, furrow;

keras, horn.] Dis-

tinguished from

Orthoceras by the

oblique septa,

arched upon the

dorsal side, and

bending down in a

deep sinus on the

ventral side, and

strongly arching

toward the mouth.

Type C. virgatum.

arcuatum, James, a

poorly defined si-

phuncle of an *En-*

doceras.

clarkii, Wetherby,

1881, Jour. Cin.

Soc. Nat. Hist., vol.

4, p. 77, Tren-

ton Gr.

virgatum, Hall, 1850,

3d Rep. N. Y. Mus. Nat. Hist., p. 182,

Birdseye and Black Riv. Grs.

Chymenia, Munster, 1839. [Ety. mythological name.]

complanata, see *Goniatites complanatus*.

erato, see *Goniatites erato*.

Conulites, Schlottheim, 1820, Petrefaktenkunde, etc. [Ety. *konos*, cone, *lithos*, stone.]

capricornulus, Troost, 1840, 5th Geo. Rep. Tenn. Not satisfactorily defined.

Conotubularia, Troost, syn. for *Orthoceras*.

brongniarti, see *Orthoceras brongniarti*.

cuvieri, see *Orthoceras cuvieri*.

defranci, see *Orthoceras defranci*.

goldfussi, see *Orthoceras goldfussi*.

Conulites, Cozzens, 1848. Not satisfactorily defined.

angulosum, Cozzens, 1848. Not satisfactorily defined. It may be a plant.

Cryptoceras, D'Orbigny, 1850. [Ety. *kryptos*, concealed; *keras*, horn.] This name

was preoccupied by Latreille for a genus of insects, and had been previously used by Barrande for a genus of Cephalopods.

capax, see *Solenochilus capax*.

CYRTO CERAS, Goldfuss, 1832, in De la Beche's

Handbuch der Geognosie bearbeitet

von v. Deschen, p. 536. [Ety. *kurtos*,

curved; *keras*, horn.] Shell long,

conical, gently curved, aperture some-

times contracted; siphuncle straight or

expanded between the septa, and vari-

able in position, but usually at the

outer edge.

absens, see *Gomphoceras absens*.

FIG. 727.—*Colpoceras clarkii*.

ely annulated at septa. Type C. 2, Jour. Acad. Nat. 7, and Pal. N. Y. Gr.

FIG. 727.—*Colpoceras clarkii*.

Nat. Hist., p. 182, v. 5, Grs. b. [Ety. mytho-
s complanatus.

o.
20, Petrefakten-
mos, cone, lithol.

40, 5th Geo. Rep.
ily defined.
for Orthoceras.
is brongniartii.
vieri.
defrancii.
goldfussi.
Not satisfactorily

18. Not satisfac-
y be a plant.
50. [Ety. *kryptos*,
rn.] This name
Latreille for a ge-
had been previ-
le for a genus of

apax.
7, in De la Beche's
nosie bearbeitet
36. [Ety. *kurtos*,
.] Shell long,
aperture some-
uncle straight or
septata, and vari-
usually at the

absens.

acinacellum, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 327, Birds-eye Gr.

amulum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 371, Up. Held. Gr.

alethes, Billings, 1865, Pal. Foss., vol. 1, p. 193, Quebec Gr.

alternatum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 365, Marcellus Shale. Proposed instead of *C. undulatum* of Hall.

ammon, Billings, 1861, Can. Jour., vol. 6, p. 361, Corniferous limestone.

amentum, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 105, Hud. Riv. Gr.

ampliocorne, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 358, syn for *C. hercules*.

annulatum, Hall, 1847. This name was preoccupied by Goldfuss in 1832, see *C. subannulatum*.

arcticameratum, Hall, 1852, Pal. N. Y., vol. 2, p. 349, Guelph Gr.

arcuatum, Hall, 1847, Pal. N. Y., vol. 1, p. 196. The name was preoccupied by Steininger in 1830, see *C. subarcuatum*.

aristides, Billings, 1865, Pal. Foss., vol. 1, p. 316, Quebec Gr.

ashmanni, n. sp. Shell small, gently curved; section subelliptical, becoming subcircular near the point, the dorsal side a little less convex, than the ventral; siphuncle near the dorsal side; surface longitudinally furrowed and finely sculptured transversely, the furrows and transverse lines most distinct on the ventral side; there are

FIG. 728.—*Cyrtoceras ashmanni*. The two section views are natural size.

eight chambers in the specimen figured, which is enlarged one-half diameter; body chamber unknown. Collected by Mr. George Ashmann, among the minute fossils at Spergen Hill, Indiana, Warsaw Gr.,

and is in the collection of Charles Faber. beekmanense, Whitfield, 1889, Bull. Am. Mus. Nat. Hist., vol. 2, p. 57, Calciferous Gr.

bannisteri, see *Trochoceras bannisteri*.

belus, Billings, 1861, Can. Jour., vol. 6, p. 361, Corniferous Gr.

boycei, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 326, Birds-eye Gr.

billingsi, Salter, 1859, Can. Org. Rem., Decade 1, p. 33, Chazy or Black Riv. Grs.

bondi, Safford, 1869, Geo. of Tenn., p. 290, Nashville Gr.

brevicorne, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 407, Niagara Gr.

camurun, Hall, 1847, Pal. N. Y., vol. 1, p. 196, Trenton Gr.

cancellatum, Hall, 1852, Pal. N. Y., vol. 2, p. 290. The name was preoccupied by Roemer in 1844. See *C. subcancellatum*.

carrollense, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 496, Galena Gr.

cessator, Hall & Whitfield, 1877, U. S. Expl. Exped. 40th parallel, vol. 4, p. 278, Coal Meas.

citum, Hall, 1879, Pal. N. Y., vol. 5, p. 372, Up. Held. Gr.

clavatum, see *Gomphoceras clavatum*.

clitus, Billings, 1866, Catal. Sil. Foss. Antic., p. 85, Niagara Gr.

confertissimum, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 327, Birds-eye Gr.

conicum, Owen, 1840, Rep. on Min. Lands, p. 70, Up. Magnesian Gr.

conoidale, Wetherby, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 78, Hud. Riv. Gr.

constrictostriatum, Hall, 1847, Pal. N. Y., vol. 1, p. 195, Trenton Gr.

corniculum, Hall, 1862, Geo. Rep. Wis.

The name was preoccupied by Bur-
rande in 1848, and again by Eichwald
in 1860, see *C. tenuistriatum*.

corydon, Billings, 1866, Catal. Sil. Foss. Antic., p. 85, Niagara Gr.

cretaceum, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 209, Up. Held. Gr.

curtum, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 463, and Geo. Sur. Ill., vol. 2, p. 388, Up. Coal Meas.

Was this name preoccupied by Eich-
wald?

dactyloides, Dwight, 1884, Am. Jour. Sci. and Arts, 3d ser., vol. 27, p. 255, Cal-
ciferous Gr.

dardanus, Hall, 1861, Rep. of Progr. Geo. Sur. of Wis., p. 43, Niagara Gr.

densum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 363, Ham. Gr.

dictys, Billings, 1865, Pal. Foss., vol. 1, p. 192, Quebec Gr.

dictyum, White, 1876, Proc. Acad. Nat. Sci., p. 33, Devonian.

dilatatum, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 468, and Geo. Sur. Ill., vol. 2, p. 389, Up. Coal Meas.

dorsatum, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 197, Permian Gr.

eugenium, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 70, and Pal. N. Y., vol. 5, pt. 2, p. 369, Schoharie grit.

eugium, Hall, 1861, Rep. of Progr. Wis., p. 40, Chazy and Black Riv. Grs.

exiguum, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 172, Trenton Gr. This is made the type of the genus *Climoceras* by Hyatt.

faberi, James, 1886, Jour. Cin. Soc. Nat. Hist., vol. 8, p. 246, Hud. Riv. Gr.

falx, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 314, and Can. Org. Rem., Decade 1, p. 32, Black Riv. and Tren-
ton Grs.

filosum, Emmons, 1842, Nat. Hist. N. Y., vol. 4, p. 392, Trenton Gr.

formosum, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 362, Ham. Gr.

fosteri, Hall, 1861, Rep. of Progr. Geo. Sur. Wis., p. 41, Niagara Gr.

fragile, Billings, 1866, Catal. Sil. Foss. Antic., p. 59, Anticosti Gr.

gibbosum, Hall, 1876, Illust. Devonian Foss., syn. for *Gomphoceras oviforme*.
giganteum, McChesney, Jan. 1860, New Pal. Foss., Niagara Gr. In 1861 McChesney referred this species to the genus *Lituites*, and proposed for it the name *Lituites cancellatus*. Prof. Hall, in the meantime, described it as *Lituites occidentalis*. It is now referred to the genus *Nautilus*, and as both the earlier names were preoccupied, McChesney's name *cancellatus* has precedence.
hallanum, D'Orbigny, 1850, Prodrôme de Pal., tome 1, p. 1, Trenton Gr. Proposed instead of *C. lamellosum*, Hall, 1847, which was preoccupied. Hyatt founded his genus *Zitteloceras* on this species.

kirbyi, Whitfield, 1889, Bull. Am. Mus. Nat. Hist., vol. 2, p. 57, Calciferous Gr.
lamellosum, Hall, 1847, Pal. N. Y., vol. 1, p. 193. The name was preoccupied by d'Archiac & Verneuil in 1842. See *C. hallanum*.
laterale, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 407, Niagara Gr.
ligarius, Billings, 1865, Pal. Foss., vol. 1, p. 176, Hud. Riv. Gr.
liratum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 72, Ham. Gr.
loculosum, Hall, 1861, Rep. of Progr. Wis., p. 42, Trenton Gr.
lucillus, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 406, Niagara Gr.
lysander, Billings, 1862, Pal. Foss., vol. 1, p. 161, Hud. Riv. Gr.

FIG. 720.—*Cyrtoceras magister*.

hector, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 364, Up. Chemung Gr.
hercules, Winchell & Marcy, 1865, (*Lituities hercules*.) M.-m. Bost. Soc. Nat. Hist., p. 102, Niagara Gr.
hertzeri, see *Gomphoceras hertzeri*.
huronense, Billings, 1865, Pal. Foss., vol. 1, p. 176, Black Riv. or Trenton Grs.
infundibulum, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 66, and Geo. Wis., vol. 4, p. 300, Niagara Gr.
irregularis, Wetherby, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 79, Hud. Riv. Gr.
isidorus, Billings, 1865, Pal. Foss., vol. 1, p. 175, Black Riv. or Trenton Gr.
janus, see *Streptoceras janus*.
jason, see *Gyroceras jason*.
juvenale, Billings, 1865, Pal. Foss., vol. 1, pp. 177, 420, Trenton Gr.

macrostomum, Hall 1847, Pal. N. Y., vol. 1, p. 194, Black Riv. and Trenton Grs.
magister, S. A. Miller, 1875, Cin. Quar. Jour. of Sci., vol. 2, pp. 132, 284, Hud. Riv. Gr.
marginale, Conrad, 1843, Proc. Acad. Nat. Sci., p. 334. The name was preoccupied by Phillips in 1841, and the species is poorly defined.
markwi, Castelnau, 1843, Système Silurien, p. 30, Trenton Gr. Not recognized.
massiense, Safford, 1869, Geo. of Tenn., p. 290, Nashville Gr.
matheri, see *Gyroceras matheri*.
maccoyi, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 467 Chazy Gr.
maximum, see *Nautilus maximus*,
mercurius, see *Cyrtoceras mercurius*.

Bull. Am. Mus.
p. 57, Calceifer.

al. N. Y., vol. 1,
preoccupied by
in 1842. See C.

Rep. N. Y. Mus.
Niagara Gr.
Pal. Foss., vol. 1

Rep. N. Y. Mus.
Gr.
p. of Progr. Wis.,

Rep. N. Y. Mus.
Niagara Gr.
Pal. Foss., vol. 1,

7. Pal. N. Y.,
Riv. and Tren-

1875, Cin. Quar.
p. 132, 284, Hud.

Proc. Acad. Nat.
e was preoccu-
and the species

ystème Silurien,
t recognized.
Geo. of Tenn.,

atheri.
Can. Nat. and
y Gr.

aximus,
mercurius.

metellus, Billings, 1865, Pal. Foss., vol. 1,
p. 191, Quebec Gr.

metula, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 72, and Illust. Devon. Foss.,
pl. 46, Up. Held. Gr.

microscopicum, Dwight, 1884, Am. Jour.
Sci. and Arts, 3d ser., vol. 27, p. 256,
Calceiferous Gr.

missisquoi, Billings, 1865, Pal. Foss., vol.
1, p. 314, (Orthoceras missisquoi,) Que-
bec Gr.

morsum, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 71, and Pal. N. Y., vol. 5,
pt. 2, p. 367, Up. Held. Gr.

multicameratum, Hall, 1847, Pal. N. Y.,
vol. 1, p. 195, Black Riv. and Trenton Gr.
myrice, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 149, Niagara Gr.

neleus, Hall, 1861, Rep. of Progr. Wis., p.
40, Chazy and Black Riv. Grs.

nevadense, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 203, Devonian.

obscurum, S. A. Miller, changed to magis-
ter because obscurum was preoccupied.

ohioense, Meek, 1871, Proc. Acad. Nat.
Sci. Phil., p. 86, and Ohio Pal., vol. 1,
p. 229, Up. Held. Gr.

olens, Hall, 1877, syn. for Trochoceras
orion.

opinum, Keyes, 1888, Proc. Acad. Nat.
Sci. Phil., pl. xii., fig. 5, Ham. Gr.

orcas, see Oncoceras orcas.

orestes, Billings, 1865, Pal. Foss., vol. 1,
p. 177, Niagara Gr.

orion, see Trochoceras orion.

orodes, Billings, 1862, Pal. Foss., vol. 1,
p. 162, Guelph Gr.

planidorsatum, Whitfield, 1880, Ann. Rep.
Geo. Sur. Wis., p. 57, and Geo. Wis.,
vol. 4, p. 231, Trenton Gr.

postumius, Billings, 1865, Pal. Foss., vol.
1, p. 178, Hud. Riv. Gr.

pusillum, Hall, 1867, 20th Rep. N. Y.
Mus. Nat. Hist., p. 407, Niagara Gr.

raei, Whitfield, 1889, Bull. Am. Mus. Nat.
Hist., vol. 2, p. 58, Calceiferous Gr.

rectum, Whitfield, 1880, Ann. Rep. Geo.
Sur. Wis., p. 85, and Geo. Wis., vol. 4,
p. 319, Niagara Gr.

regulare, Billings, 1857, Rep. of Progr.
Geo. Sur. Can., p. 314, Black Riv. and
Trenton Grs.

reversum, Spencer, 1884, Bull. No. 1, Mus.
Univ. St. Mo., p. 60, Niagara Gr.

rigidum, Hall, 1867, 20th Rep. N. Y.
Mus. Nat. Hist., p. 408, Niagara Gr.

rockfordense, Winchell, 1865, Proc.
Acad. Nat. Sci., p. 132, Kinder-
hook Gr.

septoris, see Gomphoceras septore.

simplex, Billings, 1857, Rep. of
Progr. Geo. Sur. Can., p. 313,
Black Riv. and Trenton Grs.

sinuatum, Billings, 1857, Rep. of
Progr. Geo. Sur. Can., p. 314,
Black Riv. Gr.

spinosum, see Gyroceras spinosum.

stonense, Safford, 1869, Geo. of Tenn., p.
290, Trenton Gr.

subannulatum, D'Orbigny, 1850, Prodr.
de Pal., t. 1, p. 1, Black Riv. and Tren-
ton Grs. Proposed instead of C. unnu-
lulum, Hall, 1847, which was preoccu-
pied.

subarcuatum, D'Orbigny, 1850, Prodr. de
Pal., t. 1, p. 2, Trenton Gr. Proposed
instead of C. arcuatum, Hall, 1847,
which was preoccupied.

subcancellatum, Hall, 1877, 1st Ed. Am.
Pal. Foss., p. 243, Niagara Gr. Proposed
instead of C. cancellatum, Hall, 1852,
which was preoccupied.

subcompressum, Beecher, 1888, Pal. N. Y.,
vol. 7, p. 35, Clinton Gr.

subrectum, Hall, 1859, Pal. N. Y., vol. 3,
p. 342, Low. Held. Gr.

subturbatum, Billings, 1857, Rep. of
Progr. Geo. Sur. Can., p. 312, Chazy and
Black Riv. Grs.

surgens, Barrande, 1870, Syst. Sil. de
Boh., vol. 2, p. viii, pl. 431, Quebec Gr.

syphax, Billings, 1865, Pal. Foss., vol. 1,
p. 194, Quebec Gr. This species is the
type of Eremoceras, by Hyatt.

tenuiseptum, Faber, 1886, Jour. Cin. Soc.
Nat. Hist., vol. 9, p. 18, Hud. Riv. Gr.

tenuistriatum, Hall, 1877, 1st Ed. Am. Pal.
Foss., p. 243. Proposed instead of C.
corniculum, Hall, 1862, which was pre-
occupied. Trenton Gr.

tessellatum, DeKoninck. Not American.

transversum, see Gyroceras transversum.

trentonense, Emmons, 1842, (Orthoceras
trentonensis,) Geo. Rep. N. Y., p. 396,
Trenton Gr.

trivolve, see Gyroceras trivolve.

typicum, see Cyrtoceras typica.

undulatum, Hall, 1876, see C. alternatum.

undulatum, Vanuxem, see Gyroceras un-
dulatum.

unicorne, Winchell,
1863, Proc. Acad.
Nat. Sci., p. 23, Mar-
shall Gr.

vallandighami, S. A.
Miller, 1874, Cin.
Quar. Jour. Sci., vol. 1, p. 232, Hud. Riv.
Gr.

vassarinum, Dwight, 1884, Am. Jour. Sci.
and Arts., 3d ser., vol. 27, p. 254, Cal-
ceiferous Gr.

FIG. 730.—Cyrtoceras
vallandighami.

FIG. 731.—Cyrtoceras ventricosum.

ventricosum, S. A. Miller, 1875, Cin. Quar.
Jour. Sci., vol. 2, p. 131, Hud. Riv. Gr.

whitneyi, Hall 1861, Rep. of Progr. Wis., p. 39, Hud. Riv. Gr.

CYRTOCERINA, Billings, 1865, Pal. Foss., vol. 1, p. 178. [Ety. from the termination *inus*, signifying resemblance to *Cyrtoceras*.] In form like a short, rapidly tapering *Cyrtoceras*, and having a large siphuncle on the concave side.

FIG. 732.—*Cyrtoceras* typical. a, Dorsal view, showing cavity of siphuncle; b, outline, side view.

Type C. typical. *mercurius*, Billings, 1865, Pal. Foss., vol. 1, p. 194, Quebec Gr.

typica, Billings, 1865, Pal. Foss., vol. 1, p. 178, Black Riv. Gr. *Diploceras*, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 267,

[Ety. *diploos*, double; *keras*, horn.] Founded upon the fragment of an *Endoceras*, and very poorly defined.

vanuxemi, see *Endoceras vanuxemi*.

DISCITES, DeHaan, 1825, Mongr. Ammon., etc., p. 31. [Ety. *diskos*, quoit.] Discoid; umbilicus wide; whorls quadrangular, sides flattened, and dorsum gibbous; longitudinally striated and sometimes lined transversely; siphon above the center; living chamber from one-fourth to three-fourths of a whorl in length; aperture with deep ventral sinus. Type D. *costellatus*.

✓ *ammonis*, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 425, Up. Held. Gr.

disciformis, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 261, and Geo. Sur. Ill., vol. 5, p. 522, Keokuk Gr.

hartti, Dawson, 1868, (*Gyroceras hartti*) Acadian Geol., p. 311, Subcarboniferous. Made the type of Hyatt's genus, *Stroboceras*.

highlandensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 531, Coal Meas.

✓ *inopinatus*, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 426, Up. Held. Gr.

FIG. 733.—*Discites marcellensis*.

✓ *marcellensis*, Vanuxem, 1842, (*Goniatites marcellensis*) Geo. Sur. 3d Dist. N. Y., p. 146, Marcellus Shale. The type of Hyatt's genus *Centroceras*.

ornatus, syn. for *D. marcellensis*.

toddanus, Gurley, 1883, New Carb. Foss., p. 7. Publication invalid.

tuberculatus, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 581, Subcarb.

DISCOSORUS, Hall, 1852, Pal. N. Y., vol. 2, p. 99. [Ety. *diskos*, quoit; *soros*, heap or pile.] Composed of a series of disks, gradually diminishing in size from the body chamber; outer edes rounded; joining surfaces flat. Type D. *conoides*.

conoides, Hall, 1852, Pal. N. Y., vol. 2, p. 99, Clinton and Niagara Gr.

ENDOCERAS, Hall, 1847, Pal. N. Y., vol. 1, p. 58. [Ety. *endos*, within; *keras*, horn.]

An elongated conical shell, resembling an *Orthoceras*, and possessed of one or more smooth siphuncles, which do not expand in passing through the chambers. Type E. *annulatum*.

angusticameratum, Hall, 1847, Pal. N. Y., vol. 1, p. 218, Trenton Gr.

annulatum, Hall, 1847, Pal. N. Y., vol. 1, p. 207, Trenton Gr.

approxinatum, Hall, 1847, Pal. N. Y., vol. 1, p. 219, Trenton Gr.

arctiventrum, Hall, 1847, Pal. N. Y., vol. 1, p. 217, Trenton Gr.

atlanticum, Barrande, 1870, Syst. Sil. de Boh., vol. 2, p. viii, pl. 430, Quebec Gr.

bristolense, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 85, Hud. Riv. Gr.

distans, Hall, 1847, Pal. N. Y., vol. 1, p. 220, Trenton Gr.

duplicatum, Hall, 1847, Pal. N. Y., vol. 1, p. 219, Trenton Gr.

egani, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 84, Hud. Riv. Gr.

FIG. 734.—*Endoceras longistemma*.

gemelliparum, Hall, 1847, Pal. N. Y., vol. 1, p. 60, Black Riv. Gr.

inequabile, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 86, Hud. Riv. Gr.

Geo. Sur. Wis.
Sil. Subcarb.
l. N. Y., vol. 2, p.
t; soros, heap or
series of disks,
in size from the
edges rounded;
Type D. con-

al. N. Y., vol. 2,
gara Gr.

. N. Y., vol. 1, p.
n; kera, horn.]
shell, resembling
assessed of one or
es, which do not
ough the cham-
um.

1847, Pal. N. Y.,
Gr.

al. N. Y., vol. 1,

7, Pal. N. Y., vol.

. Pal. N. Y., vol.

370, Syst. Sil. de
430, Quebec Gr.
82, Jour. Cin. Soc.
Hud. Riv. Gr.
N. Y., vol. 1, p.

al. N. Y., vol. 1,

Jour. Cin. Soc.
Hud. Riv. Gr.

ngissimum.

Pal. N. Y., vol.

1882, Jour. Cin.
5, p. 86, Hud.

insulare, Barrande, 1870, Syst. Sil. de Boh.,
vol. 2, p. viii, pl. 430-431, Quebec Gr.
laiventrum, Hall, 1860, 3d Rep. N. Y.
Mus. Nat. Hist., p. 181, Trenton Gr.
longissimum, Hall, 1847, Pal. N. Y., vol.
1, p. 59, Black Riv. and Trenton Gr.
magniventrum, Hall, 1847, Pal. N. Y., vol.
1, p. 218, Trenton Gr.

marcoui, Barrande, 1869, Syst. Sil. de Boh.,
2d ser., 4me, Quebec Gr.

multitubulatum, Hall, 1847, Pal. N. Y.,
vol. 1, p. 59, Black Riv. and Trenton Grs.

The type of Hyatt's genus *Vaginoceras*.

proteiforme, Hall, 1847, Pal. N. Y., vol.
1, p. 208, Hud. Riv. and Trenton Grs.

proteiforme var. elongatum, Hall, 1847,
Pal. N. Y., vol. 1, p. 216, Trenton Gr.

proteiforme var. lineolatum, Hall, 1847,
Pal. N. Y., vol. 1, p. 211, Trenton Gr.

proteiforme var. strangulatum, Hall, 1847,
Pal. N. Y., vol. 1, p. 212, Trenton Gr.

proteiforme var. tenuistriatum, Hall, 1847,
Pal. N. Y., vol. 1, p. 209, Trenton Gr.

proteiforme var. tenuitextum, Hall, 1847,
Pal. N. Y., vol. 1, p. 210, Trenton Gr.

rapax, Billings, 1860, (*Orthoceras rapax*),
Can. Nat. and Geol., vol. 5, p. 176,
Black Riv. Gr.

rottermundi, Barrande, 1866, (*Orthoceras*
rottermundi), Syst. Sil. de Boh., 2d ser.,
2me, p. xiii, pl. 230, Trenton Gr.

subannulatum, Whitfield, 1880, Ann. Rep.
Geo. Sur. Wis., p. 56, and Geo. Wis.,
vol. 4, p. 230, Trenton Gr.

subcentrale, Hall, 1847, Pal. N. Y., vol.
1, p. 59, Black Riv. Gr.

vanuxemi, Conrad, 1842, (*Diploceras van-*
uxemi), Jour. Acad. Sci., vol. 8, p. 267,
Trenton Gr.

Endolobus, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci. Phil., p. 259. [Ety. *endos*,
within; *lobos*, lobe.] Prof. Meek said
later that this genus is not distinct from
Temnochilus, and if distinct it would
probably be a synonym for Montfort's
genus *Bisiphites*.

peramplus, see *Temnochilus peramplus*.

spectabilis, see *Temnochilus spectabile*.

Glossoceras, Barrande, 1865, Cephalopods of
Bohemia, vol. 2, p. 372. [Ety. *glosse*,
tongue; *keras*, horn.] Having a slender
annulated whorl, and an obscurely

Y-shaped

aperture.

It is not
known as
an American
genus.

desideratum,
Billings,
1866, Catal.
Sil. Foss.

Antic., p.
60. Not
defined so

astoberec-

ognized.

GOMPHOCERAS, Sowerby, 1839, Murch. Sil.

Syst. p. 620. [Ety. *gomphos*, club; *keras*,

horn.] Shell fusiform or globular with

a tapering apex; aperture contracted

in the middle; siphuncle moniliform,

subcentral. Type *G. pyriforme*.

abruptum, Hall,
1879, Pal. N. Y.,
vol. 5, pt. 2, p.
339, Ham. Gr.

absens, Hall,
1876, (*Cyrtoceras*
absens),
Illust. Devon.
Foss., pl. 47,
and Pal. N. Y.,
vol. 5, pt. 2, p.
324, Up. Held.
Gr.

ajax, Hall, 1879,
Pal. N. Y., vol.
5, pt. 2, p. 350,
Portage Gr.

amphora, Whit-
field, 1882, Ann.
N. Y. Acad.
Sci., vol. 2, p.
207, Up. Held.
Gr.

beta, Hall, 1862,
15th Rep. N. Y. Mus. Nat. Hist., p. 72,
and Pal. N. Y., vol. 5, pt. 2, p. 326, Up.
Held. Gr.

breviposticum, Whitfield, 1882, Geo. Wis.,
vol. 4, p. 339, Ham. Gr.

cammarus, Hall, 1879, Pal. N. Y., vol. 5,
p. 333, Up. Held. Gr.

cassinense, Whitfield, 1886, Bull. Am.
Mus. Nat. Hist., vol. 1, p. 322, Birds-
eye Gr.

cincinnatiense, S. A. Miller, 1884, Jour.
Cin. Soc. Nat. Hist., vol. 7, p. 19, Hud.
Riv. Gr.

clavatum, Hall, 1876, (*Cyrtoceras clava-*
tum), Illust. Devon. Foss., pl. 47, Up.
Held. Gr.

conradi, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, Ham. Gr.

crenatum, Beecher, 1883, Pal. N. Y., vol.
7, p. 33, Up. Held. Gr.

cruciferum, Hall, 1879, Pal. N. Y., vol. 5,
p. 328, Schoharie grit.

eos, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 100, Hud. Riv. Gr.

eximium, Hall, 1861, 14th Rep. N. Y.
Mus. Nat. Hist., p. 109, and Pal.
N. Y., vol. 5, pt. 2, p. 299, Up.
Held. Gr.

faberi, S. A. Miller, 1884, Jour. Cin. Soc.
Nat. Hist., vol. 7, p. 19, Hud. Riv. Gr.

fax, Hall, 1879, Pal. N. Y., vol. 5, p. 321,
Schoharie grit.

fischeri, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, and Pal. N. Y., vol.
5, pt. 2, p. 336, Ham. Gr.

tusiforme, Whitfield, 1882, Geo. Wis., vol.
4, p. 338, Ham. Gr.

hertzeri, Hall & Whitfield, 1875, (*Cyrtoceras*
hertzeri), Ohio Pal., vol. 2, p. 150,
Niagara Gr.

horn.] Shell fusiform or globular with

a tapering apex; aperture contracted

in the middle; siphuncle moniliform,

subcentral. Type *G. pyriforme*.

abruptum, Hall,
1879, Pal. N. Y.,
vol. 5, pt. 2, p.
339, Ham. Gr.

absens, Hall,
1876, (*Cyrtoceras*
absens),
Illust. Devon.
Foss., pl. 47,
and Pal. N. Y.,
vol. 5, pt. 2, p.
324, Up. Held.
Gr.

ajax, Hall, 1879,
Pal. N. Y., vol.
5, pt. 2, p. 350,
Portage Gr.

amphora, Whit-
field, 1882, Ann.
N. Y. Acad.
Sci., vol. 2, p.
207, Up. Held.
Gr.

beta, Hall, 1862,
15th Rep. N. Y. Mus. Nat. Hist., p. 72,
and Pal. N. Y., vol. 5, pt. 2, p. 326, Up.
Held. Gr.

breviposticum, Whitfield, 1882, Geo. Wis.,
vol. 4, p. 339, Ham. Gr.

cammarus, Hall, 1879, Pal. N. Y., vol. 5,
p. 333, Up. Held. Gr.

cassinense, Whitfield, 1886, Bull. Am.
Mus. Nat. Hist., vol. 1, p. 322, Birds-
eye Gr.

cincinnatiense, S. A. Miller, 1884, Jour.
Cin. Soc. Nat. Hist., vol. 7, p. 19, Hud.
Riv. Gr.

clavatum, Hall, 1876, (*Cyrtoceras clava-*
tum), Illust. Devon. Foss., pl. 47, Up.
Held. Gr.

conradi, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, Ham. Gr.

crenatum, Beecher, 1883, Pal. N. Y., vol.
7, p. 33, Up. Held. Gr.

cruciferum, Hall, 1879, Pal. N. Y., vol. 5,
p. 328, Schoharie grit.

eos, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 100, Hud. Riv. Gr.

eximium, Hall, 1861, 14th Rep. N. Y.
Mus. Nat. Hist., p. 109, and Pal.
N. Y., vol. 5, pt. 2, p. 299, Up.
Held. Gr.

faberi, S. A. Miller, 1884, Jour. Cin. Soc.
Nat. Hist., vol. 7, p. 19, Hud. Riv. Gr.

fax, Hall, 1879, Pal. N. Y., vol. 5, p. 321,
Schoharie grit.

fischeri, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, and Pal. N. Y., vol.
5, pt. 2, p. 336, Ham. Gr.

tusiforme, Whitfield, 1882, Geo. Wis., vol.
4, p. 338, Ham. Gr.

hertzeri, Hall & Whitfield, 1875, (*Cyrtoceras*
hertzeri), Ohio Pal., vol. 2, p. 150,
Niagara Gr.

horn.] Shell fusiform or globular with

a tapering apex; aperture contracted

in the middle; siphuncle moniliform,

subcentral. Type *G. pyriforme*.

abruptum, Hall,
1879, Pal. N. Y.,
vol. 5, pt. 2, p.
339, Ham. Gr.

absens, Hall,
1876, (*Cyrtoceras*
absens),
Illust. Devon.
Foss., pl. 47,
and Pal. N. Y.,
vol. 5, pt. 2, p.
324, Up. Held.
Gr.

ajax, Hall, 1879,
Pal. N. Y., vol.
5, pt. 2, p. 350,
Portage Gr.

amphora, Whit-
field, 1882, Ann.
N. Y. Acad.
Sci., vol. 2, p.
207, Up. Held.
Gr.

beta, Hall, 1862,
15th Rep. N. Y. Mus. Nat. Hist., p. 72,
and Pal. N. Y., vol. 5, pt. 2, p. 326, Up.
Held. Gr.

breviposticum, Whitfield, 1882, Geo. Wis.,
vol. 4, p. 339, Ham. Gr.

cammarus, Hall, 1879, Pal. N. Y., vol. 5,
p. 333, Up. Held. Gr.

cassinense, Whitfield, 1886, Bull. Am.
Mus. Nat. Hist., vol. 1, p. 322, Birds-
eye Gr.

cincinnatiense, S. A. Miller, 1884, Jour.
Cin. Soc. Nat. Hist., vol. 7, p. 19, Hud.
Riv. Gr.

clavatum, Hall, 1876, (*Cyrtoceras clava-*
tum), Illust. Devon. Foss., pl. 47, Up.
Held. Gr.

conradi, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, Ham. Gr.

crenatum, Beecher, 1883, Pal. N. Y., vol.
7, p. 33, Up. Held. Gr.

cruciferum, Hall, 1879, Pal. N. Y., vol. 5,
p. 328, Schoharie grit.

eos, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 100, Hud. Riv. Gr.

eximium, Hall, 1861, 14th Rep. N. Y.
Mus. Nat. Hist., p. 109, and Pal.
N. Y., vol. 5, pt. 2, p. 299, Up.
Held. Gr.

faberi, S. A. Miller, 1884, Jour. Cin. Soc.
Nat. Hist., vol. 7, p. 19, Hud. Riv. Gr.

fax, Hall, 1879, Pal. N. Y., vol. 5, p. 321,
Schoharie grit.

fischeri, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, and Pal. N. Y., vol.
5, pt. 2, p. 336, Ham. Gr.

tusiforme, Whitfield, 1882, Geo. Wis., vol.
4, p. 338, Ham. Gr.

hertzeri, Hall & Whitfield, 1875, (*Cyrtoceras*
hertzeri), Ohio Pal., vol. 2, p. 150,
Niagara Gr.

horn.] Shell fusiform or globular with

a tapering apex; aperture contracted

in the middle; siphuncle moniliform,

subcentral. Type *G. pyriforme*.

abruptum, Hall,
1879, Pal. N. Y.,
vol. 5, pt. 2, p.
339, Ham. Gr.

absens, Hall,
1876, (*Cyrtoceras*
absens),
Illust. Devon.
Foss., pl. 47,
and Pal. N. Y.,
vol. 5, pt. 2, p.
324, Up. Held.
Gr.

ajax, Hall, 1879,
Pal. N. Y., vol.
5, pt. 2, p. 350,
Portage Gr.

amphora, Whit-
field, 1882, Ann.
N. Y. Acad.
Sci., vol. 2, p.
207, Up. Held.
Gr.

beta, Hall, 1862,
15th Rep. N. Y. Mus. Nat. Hist., p. 72,
and Pal. N. Y., vol. 5, pt. 2, p. 326, Up.
Held. Gr.

breviposticum, Whitfield, 1882, Geo. Wis.,
vol. 4, p. 339, Ham. Gr.

cammarus, Hall, 1879, Pal. N. Y., vol. 5,
p. 333, Up. Held. Gr.

cassinense, Whitfield, 1886, Bull. Am.
Mus. Nat. Hist., vol. 1, p. 322, Birds-
eye Gr.

cincinnatiense, S. A. Miller, 1884, Jour.
Cin. Soc. Nat. Hist., vol. 7, p. 19, Hud.
Riv. Gr.

clavatum, Hall, 1876, (*Cyrtoceras clava-*
tum), Illust. Devon. Foss., pl. 47, Up.
Held. Gr.

conradi, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, Ham. Gr.

crenatum, Beecher, 1883, Pal. N. Y., vol.
7, p. 33, Up. Held. Gr.

cruciferum, Hall, 1879, Pal. N. Y., vol. 5,
p. 328, Schoharie grit.

eos, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 100, Hud. Riv. Gr.

eximium, Hall, 1861, 14th Rep. N. Y.
Mus. Nat. Hist., p. 109, and Pal.
N. Y., vol. 5, pt. 2, p. 299, Up.
Held. Gr.

faberi, S. A. Miller, 1884, Jour. Cin. Soc.
Nat. Hist., vol. 7, p. 19, Hud. Riv. Gr.

fax, Hall, 1879, Pal. N. Y., vol. 5, p. 321,
Schoharie grit.

fischeri, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, and Pal. N. Y., vol.
5, pt. 2, p. 336, Ham. Gr.

tusiforme, Whitfield, 1882, Geo. Wis., vol.
4, p. 338, Ham. Gr.

hertzeri, Hall & Whitfield, 1875, (*Cyrtoceras*
hertzeri), Ohio Pal., vol. 2, p. 150,
Niagara Gr.

horn.] Shell fusiform or globular with

a tapering apex; aperture contracted

in the middle; siphuncle moniliform,

subcentral. Type *G. pyriforme*.

abruptum, Hall,
1879, Pal. N. Y.,
vol. 5, pt. 2, p.
339, Ham. Gr.

absens, Hall,
1876, (*Cyrtoceras*
absens),
Illust. Devon.
Foss., pl. 47,
and Pal. N. Y.,
vol. 5, pt. 2, p.
324, Up. Held.
Gr.

ajax, Hall, 1879,
Pal. N. Y., vol.
5, pt. 2, p. 350,
Portage Gr.

amphora, Whit-
field, 1882, Ann.
N. Y. Acad.
Sci., vol. 2, p.
207, Up. Held.
Gr.

beta, Hall, 1862,
15th Rep. N. Y. Mus. Nat. Hist., p. 72,
and Pal. N. Y., vol. 5, pt. 2, p. 326, Up.
Held. Gr.

breviposticum, Whitfield, 1882, Geo. Wis.,
vol. 4, p. 339, Ham. Gr.

cammarus, Hall, 1879, Pal. N. Y., vol. 5,
p. 333, Up. Held. Gr.

cassinense, Whitfield, 1886, Bull. Am.
Mus. Nat. Hist., vol. 1, p. 322, Birds-
eye Gr.

cincinnatiense, S. A. Miller, 1884, Jour.
Cin. Soc. Nat. Hist., vol. 7, p. 19, Hud.
Riv. Gr.

clavatum, Hall, 1876, (*Cyrtoceras clava-*
tum), Illust. Devon. Foss., pl. 47, Up.
Held. Gr.

conradi, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 106, Ham. Gr.

crenatum, Beecher, 1883, Pal. N. Y., vol.
7, p. 33, Up. Held. Gr.

cruciferum, Hall, 1879, Pal. N. Y., vol. 5,
p. 328, Schoharie grit.

eos, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 100, Hud. Riv. Gr.

eximium, Hall, 1861, 14th Rep. N. Y.
Mus. Nat. Hist., p. 109, and Pal.
N. Y., vol. 5, pt. 2, p. 299, Up.
Held. Gr.

faberi, S. A. Miller, 1884, Jour. Cin. Soc.
Nat. Hist., vol. 7, p. 19, Hud. Riv. Gr.

fax, Hall, 1879

- ✓ *gomphus*, Hall, 1879, Pal. N. Y., vol. 5, p. 334, Up. Held. Gr.
- ✓ *hyatti*, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 208, Up. Held. Gr.
- ✓ *illæus*, Hall, 1879, Pal. N. Y., vol. 5, p. 332, Scholharie grit.
- ✓ *impar*, Hall, 1879, Pal. N. Y., vol. 5, p. 332, Up. Held. Gr.
- ✓ *lunatum*, Hall, 1879, Pal. N. Y., vol. 5, p. 341, Ham. Gr.
- ✓ *manes*, Hall, 1879, Pal. N. Y., vol. 5, p. 330, Genesee Slate.
- marcyæ*, Winchell & Marcy, 1885, Mem. Bost. Soc. Nat. Hist. Syn. for G. *scrinium*.
- minimum*, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 321, Birdseye Gr.
- ✓ *minum*, Beecher, 1888, Pal. N. Y., vol. 7, p. 34, Ham. Gr.
- ✓ *mitra*, Hall, 1879, Pal. N. Y., vol. 5, p. 330, Up. Held. Gr.
- nasutum*, Beecher, 1888, Pal. N. Y., vol. 7, p. 34, Chemung Gr.
- obsum*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 311, Utica Gr.
- ✓ *omicron*, Winchell, 1866, Rep. Low. Peninsula Mich., p. 97, Ham. Gr.
- ✓ *oviforme*, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 105, Ham. Gr.
- ✓ *pingue*, Hall, 1879, Pal. N. Y., vol. 5, p. 346, Ham. Gr.
- ✓ *planum*, Hall, 1879, Pal. N. Y., vol. 5, p. 352, Ham. Gr.
- ✓ *plenum*, Beecher, 1888, Pal. N. Y., vol. 7, p. 33, Up. Held. Gr.
- ✓ *poculum*, Hall, 1879, Pal. N. Y., vol. 5, p. 340, Ham. Gr.
- ✓ *potens*, Hall, 1879, Pal. N. Y., vol. 5, p. 351, Waverly Gr.
- powersi*, James. Not recognized.
- ✓ *raphanus*, Hall, 1879, Pal. N. Y., vol. 5, p. 347, Ham. Gr.
- ✓ *rude*, Hall, 1879, Pal. N. Y., vol. 5, p. 327, Ham. Gr.
- ✓ *sacculus*, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 258, and Geo. Sur. Ill., vol. 3, p. 445, Ham. Gr.
- ✓ *sciotoense*, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 208, Up. Held. Gr.
- scrinium*, Hall, 1864, 20th Rep. N. Y. Mus. Nat. Hist., p. 410, Niagara Gr.
- septore*, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 410, Niagara Gr.
- ✓ *solidum*, Hall, 1879, Pal. N. Y., vol. 5, p. 338, Marcellus Shale.
- subgracile*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 311, Up. Sil.
- ✓ *suboviforme*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 202, Devonian.
- ✓ *tumidum*, Hall, 1879, Pal. N. Y., vol. 5, p. 351, Chemung Gr.
- ✓ *turbiniforme*, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 258, and Geo. Sur. Ill., vol. 3, p. 444, Ham. Gr.
- GONIATITES**, DeHaan, 1825, Monographie Ammoniteorum et Goniatiteorum, p. 159. [Ety. *gonia*, an angle; *lithos*, stone. This name, it seems, should be spelled Goniolites.] Discoid; whorls embracing, sometimes closing the umbilicus; septa marked by zigzag lines or sutures; when the septa are folded the elevations are called *saddles*; body chamber long, sometimes constituting a whorl, but never expanding but slightly; siphonicle ventral. Type *Goniatites sphericus*.
- allii*, Winchell, 1862, Am. Jour. Sci., 2d series, vol. 33, p. 363, Marshall Gr.
- amplexus*, Beecher, 1888, Pal. N. Y., vol. 7, p. 39, Tully limestone.
- andrewsi*, Winchell, 1870, Proc. Am. Phil. Soc., vol. 12, p. 259, Marshall Gr.
- astarte*, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 29, Marcellus Shale.
- bicostatus*, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 245, and Pal. N. Y., vol. 5, pt. 2, p. 450, Portage Gr.
- canadensis*, Castelnau, 1843, Syst. Sil. p. 34. Probably a syn. for *Bellerophon bilobatus*.
- chemungensis*, Vanuxem, 1842, Geo. Rep. 3d Dist. N. Y., p. 182, and Pal. N. Y., vol. 5, pt. 2, p. 467, Chemung Gr.
- chemungensis* var. *aquicostatus*, Hall, 1875, 27th Rep. N. Y. Mus. Nat. Hist., p. 135, Chemung Gr.
- choctawensis*, Shumard, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 109, Coal Meas.
- colubrellus*, Morton, 1836, (Ammonites *colubrellus*), Am. Jour. Sci. and Arts, vol. 29, p. 154, Waverly Gr.
- compactus*, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 154, and Geo. Sur. Ill., vol. 5, p. 611, Coal Meas.
- complanatus*, Hall, 1843, (Clymenia (?) *complanatus*.) Geo. Rep. 4th Dist. N. Y., p. 244, and Pal. N. Y., vol. 5, p. 455, Portage Gr.
- complanatus* var. *perlatus*, Hall, 1875, 27th Rep. N. Y. Mus. Nat. Hist., p. 132, Chemung Gr.
- desideratus*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 203, Devonian.
- discoideus*, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 97, and Illust. Devon. Foss., pl. 71, Ham. Gr. The type of Hyatt's genus *Parodiceras*.
- discoideus* var. *ohioensis*, Hall, 1874, 27th Rep. N. Y., p. 200, Up. Held. Gr.
- entogonus*, Gabb, 1861, Proc. Acad. Nat. Sci., p. 372, Carboniferous.

FIG. 737.—*Goniatites sphericus*.

5, Monographia
atiteorum, p. 159,
hos, stone. This

latites sphericus.

when the septa
ons are culled
ong, sometimes
t never expand-
in cle ventral.

us.
Jour. Sci., 2d
Marshall Gr.

Pal. N. Y., vol.
p.

0, Proc. Am.
9, Marshall Gr.

U. S. Geo. Sur.,
Shale.

p. Rep. 4th Dist.
N. Y., vol. 5, pt.

13, Syst. Sil. p.
or Bellerophon

1842, Geo. Rep.
and Pal. N. Y.,

emung Gr.
costatus, Hall,

ius. Nat. Hist.,

1863, Trans. St.
2, p. 109, Coal

3, (Ammonites
Sci. and Arts,
Gr.

nen, 1865, Proc.
154, and Geo.

Coal Meas.
(Clymenia (?)

4th Dist. N. Y.,
vol. 5, p. 455.

us, Hall, 1875,
at. Hist., p. 132.

Monogr. U. S.
Devonian.

th Rep. N. Y.
7, and Illust.

am. Gr. The
urodiceras.

p, Hall, 1874,
Up. Held. Gr.

oc. Acad. Nat.
us.

FIG. 738.—Goniattites globulosus.

erato, Hall, 1862, (Clymenia erato,) 15th
Rep. N. Y. Mus. Nat. Hist., p. 64, and
Illust. Devon. Foss., pl. 70, Ham. Gr.
expansus, Vanuxem. The name was pre-
occupied by Von Buch in 1838. See
G. vanuxemi.

FIG. 739.—Goniattites globulosus. Outline.

goniolobus, Meek, 1877, U. S. Geo. Sur.
40th Parallel, vol. 4, p. 98, Carboniferous.
hathawayanus, McChesney, 1860, Desc.
New Pal. Foss., p. 66, Coal Meas.

globulosus, Meek
& Worthen,
1860, Proc.
Acad. Nat. Sci.
Phil., p. 471,
and Geo. Sur.
Ill., vol. 2, p.
390, Up. Coal
Meas.

globulosus var.
excelsus, Meek,
1875, Bull. U. S.
Geo. Sur. Terr.,
vol. 1, No. 6,
p. 445, Coal
Meas.

hildrethi, Morton, 1836, (Ammonites hil-
drethi,) Am. Jour. Sci. and Arts, vol.
29, p. 149, Waverly Gr.

holmesi, Swallow, 1860, Trans. St. Louis
Acad. Sci., vol. 1, p. 659, Waverly or
Choteau Gr.

houghtoni, Winchell, 1862, Am. Jour.
Sci., 2d ser., vol. 33, p. 363, Marshall Gr.

hyas, Hall, 1860, syn. for G. lyoni.

iowensis, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 471, and Geo.
Sur. Ill., vol. 2, p. 392, Coal Meas.

Type of Hyatt's genus Paralegoceras.
ixion, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 125, Kinderhook Gr. This
species is founded on the form which
has been identified with the European
species G. rotatorius.

kentuckiensis, n. sp. Shell very globose
and wide or broadly rounded on the
dorsal side; outer volution embracing
the inner ones; umbilicus small and
disclosing none of the inner volutions,
though the body chamber is broken
from our specimens; suture having a
sharp, dorso-lateral lobe and an equally

deep, bifid, central dorsal lobe; dorsal saddle subangular. This species is readily distinguished by its deep, globose form and sharply bifid lobe on the dorsal side. Collected by Charles Faber at Crab Orchard, Kentucky, in the St. Louis Group.

FIG. 740.—*Goniatites kentuckiensis*. The saddles are generally more angular than they appear in the figure.

kingi, Hall & Whitfield, 1877, U. S. Geo. Expl. Exped., 40th parallel, vol. 4, p. 279, Coal Meas.
lutheri, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 50, Chemung Gr.
lyoni, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., vol. 12, p. 471, and Geo. Sur. Ill., vol. 2, p. 165, Kinderhook Gr.
marcellensis, see *Discites marcellensis*.
marshallensis, Winchell, 1862, Am. Jour. Sci., 2d ser., vol. 33, p. 362, Marshall Gr.
minimus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 200, Coal Meas.
mithrax, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 98, and Pal. N. Y., vol. 5, pt. 2, p. 433, Up. Held. Gr.
monroensis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 150, St. Louis Gr.
morganeusis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 652, Waverly or Choteau Gr.
nodifer, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 21, Marcellus Shale.
nolinensis, Cox, 1857, Geo. Sur. Ky., vol. 3, p. 574, Coal Meas.
nundaia, Hall, 1875, syn. for *G. sinuosus*.
ohioensis, Winchell, 1870, Proc. Am. Phil. Soc., vol. 12, p. 259, Marshall Gr.
opimus, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 305, Kinderhook Gr.
orbicella, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 99, Ham. Gr.
osagensis, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 659, Waverly or Choteau Gr.
oweni, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 100, Kinderhook Gr.
oweni var. *parallelus*, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 101, Kinderhook Gr. Type of Hyatt's genus *Munsteroceras*.
parvus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 199, Coal Meas.
patersoni, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 99, Portage Gr.

peracutus, Hall, 1876, Illust. Devonian Foss., pl. 69, and Pal. N. Y., vol. 5, pt. 2, p. 403, Portage Gr.
plunorbiformis, Shumard, 1855, Geo. Sur. Mo., p. 208, Coal Meas.
plebeiformis, Hall, 1870, Pal. N. Y., vol. 5, p. 448, Marcellus Shale.
politus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 199, Coal Meas.
propinquus, Winchell, 1862, Am. Jour. Sci. and Arts, 2d series, vol. 33, p. 365, Marshall Gr.
punctatus, Conrad, 1838, Ann. Rep. N. Y., p. 117, Ham. Gr. Not properly defined.
pygmaeus, Winchell, 1862, Am. Jour. Sci. and Arts, 2d series, vol. 33, p. 366, Marshall Gr.
romingeri, Winchell, 1862, Proc. Acad. Nat. Sci., p. 427, Marshall Gr.
rotatorius, DeKoninck, 1843, Desc. des Anim. Foss. du Terr. Carb. See *G. icion*.
shumardanus, Winchell, 1865, Am. Jour. Sci. and Arts, 2d series, vol. 33, p. 363, Marshall Gr.
simulator, Hall, 1875, 27th Rep. N. Y. Mus. Nat. Hist., p. 133, Chemung Gr. Type of Hyatt's genus *Manticoceras*.
sinuosus, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 243, and Pal. N. Y., vol. 5, pt. 2, p. 460, Portage Gr. Type of Hyatt's genus *Gephuroceras*.

subcircularis n. sp. Shell small, circular; outer volution embracing the inner ones; umbilicus does not expose any of the inner volutions; four furrows or constrictions radiate from the umbilicus and divide the shell into four subequal parts, but become obsolete on the dorsal side, and in this respect resemble *Goniatites divinus* of DeKoninck; dorsal side round; body chamber unknown; suture lobed; entire surface longitudinally striated. Collected by Charles Faber, at Crab Orchard, Kentucky, in the St. Louis Group.

FIG. 741.—*Goniatites subcircularis*. Lateral and dorsal views magnified 2 diam.

sulciferus, Winchell. Not defined.
texasus, Shumard, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 109, Coal Meas.
uniangularis, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 3, p. 268, Ham. Gr. Type of Hyatt's genus *Tornoceras*.
unilobatus, Hall, 1875, 27th Rep. N. Y. Mus. Nat. Hist., p. 133, and Illust. Devon. Foss., pl. 71, Ham. Gr.
vanuxemi, Hall, 1879, Pal. N. Y., vol. 5, p. 434, Marcellus Shale. Proposed instead of *G. expansus*, of Vanuxem, which was preoccupied by Von Buch.
whitii, Winchell, 1862, Proc. Acad. Nat. Sci., vol. 6, p. 428, Portage Gr.

Illust. Devonian
N. Y., vol. 5, pt.

1855, Geo. Sur.

Pal. N. Y., vol.
le.

Trans. St. Louis
Coal Meas.

1862, Am. Jour.
s, vol. 33, p. 365,

Ann. Rep. N. Y.,
properly defined,
Am. Jour. Sci.
33, p. 366, Mar-

2, Proc. Acad.
all Gr.

1843, Desc. des
rb. See *G. lion*,
1865, Am. Jour.
vol. 33, p. 363,

15th Rep. N. Y.,
3, Chemung Gr.
Manticoceras.
Rep. 4th Dist.
N. Y., vol. 5, pt.
Type of Hyatt's

Nautilites subtr.
lateral and dorsal
enlarged 2 diam.

as radiate from
the shell into
t become obso-
and in this res-
divisus of De-
round; body
ture lobed; en-
ly striated. Col-
lateral, at Crab
the St. Louis

defined.

Trans. St. Louis
Coal Meas.

1842, Jour. Acad.
Ham. Gr. Type

ceras.

15th Rep. N. Y.
33, and Illust.

am. Gr.

l. N. Y., vol. 5,
Proposed in-
of Vanuxem,

by Von Buch.

Proc. Acad. Nat.
ge Gr.

GONIOCERAS, Hall, 1847, Pal. N. Y., vol. 1,
p. 54. [Ety. *gonia*, angle; *keras*, horn.]
Somewhat in the form of an Orthoceras,
but more or less flattened and subfusiform;
transverse section in the form of a
depressed ellipse with projecting
angles; siphuncle ventral, septa curve
over the ventral side, as shown in the
illustration. Type *G. anceps*.

FIG. 742.—*Gonioceras anceps*.

anceps, Hall, 1847, Pal. N. Y., vol. 1, p.
54, Black Riv. Gr.

occidentale, Hall, 1861, Rep. of Progr. Wis.,
p. 47, Trenton Gr.

GYROCERAS, DeKoninck, 1844, Desc. An.
Foss. Belg., p. 530. [Ety. *gyros*, circle;
keras, horn.] Not Gyrocera-
tites of Meyer, 1820. Discoid, rolled in one
plane; volutions in contact or open,
but not embracing; transverse section
circular, elliptical, scutiform, or poly-
gonal; body chamber large and some-
times straight or tangent to the spiral;
opening hollowed out on the exterior
border like the Nautilus; septa arched
and frequently project, curving back-
ward; siphon slender, cylindrical, and
usually subcentral toward the convex
border, but sometimes found within
the concave border; surface tubercu-
lous, having imbricated excrescences or
ringed with projecting fringes from the
septa. Type *G. paradoxicum*.

abruptum, Hall, 1879, Desc. New Spec.
Foss., p. 19, and 11th Rep. Geo. and
Nat. Hist., Ind. p. 325, Niagara Gr.

americanum, Billings, 1857, Rep. of Progr.
Can. Geo. Sur., p. 309, Up. Sil.

baeri, Meek & Worthen, 1865, (Trochoce-
ras *baeri*.) Proc. Acad. Nat. Sci., p. 263,
and Ohio Pal., vol. 1, p. 157, Hud.
Riv. Gr.

bannisteri, Winchell & Marcy, 1865, Mem.
Bost. Soc. Nat. Hist., vol. 1, p. 102, Ni-
agara Gr.

burlingtonense, see *Nautilus burlingtonensis*.
columbiense, Whitfield, 1882, Ann. N. Y.

Acad. Sci., vol. 2, p. 210, Up. Held. Gr.

constrictum, Meek & Worthen, 1868, Geo.
Sur. Ill., vol. 3, p. 446, Ham. Gr.

cornutum, Owen, 1840, Rep. on Min.
Lands, p. 69, Devonian.

cyclops, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 68, and Illust. Devon.

Foss., pl. 53, Up. Held. Gr.

duplicostatum, Whitfield, 1878, Ann. Rep.
Geo. Sur. Wis., p. 78, and Geo. Wis.,
vol. 4, p. 235, Trenton Gr.

elrodi, White, 1882, 11th Ann. Rep. Geol.
and Nat. Hist. Indiana, p. 356, Niagara Gr.

eryx, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 67, Ham. Gr.

expansum, Saeman, Dunker & Von Meyer,
1853, Palaeontographica, vol. 4. See
Nautilus buccinum.

gracile, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 105, Kinderhook Gr.

Probably a syn. for *Trematodiscus*
digonus.

hartii, see *Discites hartii*.

inelegans, Meek, 1871, Proc. Acad. Nat.
Sci. Phil., p. 80, and Ohio Pal., vol. 1,
p. 232, Up. Held. Gr.

jason, Hall, 1862, (Cyrtoceras *jason*.) 15th
Rep. N. Y. Mus. Nat. Hist., p. 71, Up.
Held. Gr. Type of Hyatt's genus *Ru-*

toceras.

laciniosum, Hall, 1879, Pal. N. Y., vol. 5,
p. 376, Up. Held. Gr.

liratum, see *Nautilus liratus*.

logani, Meek, 1868, Trans. Chi. Acad. Sci.,
p. 110, Devonian.

magnificum, see *Lituites magnificus*.

matheri, Conrad, 1840, Ann. Rep. N. Y.,
p. 206, and Pal. N. Y., vol. 5, pt. 2, p.
377, (Cyrtoceras *matheri*.) Up. Held. Gr.

nais, see *Porcellia nais*.

nereus, Hall, 1862, 15th Rep. N. Y. Mus.
Nat. Hist., p. 67, and Pal. N. Y., vol. 5,
pt. 2, p. 373, Up. Held. Gr.

numa, Billings, 1875, Can. Nat. and Geol.,
vol. 7, p. 238, Up. Held. Gr.

ohioense, Meek, 1871, Proc. Acad. Nat.
Sci. Phil., p. 87, and Ohio Pal., vol. 1,
p. 230, Up. Held. Gr.

puncinodum, Hall, 1876, Illust. Devonian
Foss., pl. 55, and Pal. N. Y., vol. 5, pt.
2, p. 380, Up. Held. Gr.

pratti, Barris, 1879, Proc. Dav. Acad. Sci.,
vol. 2, p. 287, Up. Held. Gr.

rhombolineure, Owen, 1862, Geo. Sur.
Indiana, p. 362, Silurian.

rockfordense, Meek & Worthen, 1866,
(*Nautilus* (*Cryptoceras*) *rockfordensis*.)
Proc. Acad. Nat. Sci. Phil., p. 275, and
Geo. Sur. Ill., vol. 3, p. 459, Kinder-

hook Gr.

seminodosum, Whitfield, 1882, Ann. N. Y.
Acad. Sci., vol. 2, p. 211, Up. Held. Gr.

spinosum, Conrad, 1840, (Phragmoceras
spinosum.) Ann. Rep. N. Y., p. 206,
and Pal. N. Y., vol. 5, pt. 2, p. 382,
Schoharie grit.

stebos, Beecher, 1888, Pal. N. Y., vol. 7, p.
36, Waverly Gr.

subliratum, see *Nautilus subliratus*.

transversum, Hall, 1860, (Cyrtoceras
transversum.) 13th Rep. N. Y. Mus.
Nat. Hist., p. 104, and Pal. N. Y., vol.
5, pt. 2, p. 384, Ham. Gr.

trivolve, Conrad, 1840, (Cyrtoceras *tri-*
volvius.) Ann. Rep. N. Y., p. 206, and Pal.
N. Y., vol. 5, pt. 2, p. 374, Up. Held. Gr.

undulatum, Vanuxem, 1842, (Cyrtoceras
undulatum.) Geo. Rep. N. Y., p. 139, and
Pal. N. Y., vol. 5, pt. 2, p. 378, Up. Held.

Gr. Type of Hyatt's genus *Halloceras*.

vagrans, Billings, 1857, Rep. of Progr. Can. Geo. Sur., p. 308, Black Riv. Gr. validum, Hall, 1876, Illust. Devonian Foss., pl. 51, t. 1 Pal. N. Y., vol. 5, pt. 2, p. 385, Schoharie grit.

FIG. 743.—*Gyroceras undulatum*.

Hortholus americanus, see *Lituites americanus*.

HURONIA, Stokes, 1823, Geo. Trans., n. s., vol. 1, p. 203. When this genus was proposed, the author thought he was describing a coral. Prof. Billings said the name was proposed for the siphuncle of an *Orthoceras*, and is, therefore, merely a synonym. This seems to be the correct opinion, though the shells to which the peculiar siphuncles belong are unknown. Type *H. bigsbyi*. *annulata*, Hall, 1851, Lake Superior Land Dist. by Foster & Whitney, p. 221, Niagara Gr.

bigsbyi, Stokes, 1823, Trans. Geo. Soc., vol. 1, p. 195, Clinton Gr.

minuens, Barrande, 1869, Syst. Sil. de Boh., 2d series, vol. 4, pl. ix, p. 435, Clinton Gr.

obliqua, Stokes, 1823, Trans. Geo. Soc., 2d series, vol. 1, p. 203, Clinton Gr.

portlocki, Stokes, 1840, Trans. Geo. Soc., 2d series, vol. 5, p. 710, Clinton Gr.

sphaericialis, Stokes, 1840, Trans. Geo. Soc., 2d series, vol. 5, p. 710, Clinton Gr.

stokesi, Castelnau, 1843, Syst. Sil., p. 33, Schoharie grit. Not recognized.

turbinata, Stokes, 1823, Trans. Geo. Soc., 2d series, vol. 1, p. 203, Clinton Gr.

vertebralis, Stokes, 1840, Trans. Geo. Soc., 2d series, vol. 5, p. 710,

Niagara and Clinton Grs. See *Orthoceras canadense*.

FIG. 744.—*Huronia vertebralis*.

Hydnoceras, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8. See *Dictyophyton*, a sponge, *tuberosum*, see *Dictyophyton tuberosum*.

LITUITES, Mont-

fort, 1808,

Conch. Syst.,

vol. 1, p. 279,

[*Ety. lituus*,

trumpet.]

Shell spiral

in the begin-

ning; last

chambers pro-

duced straight;

whorls free or

open in one plane;

septa simple;

siphuncle central;

section circular. Type

L. lituus.

americanus, D'Orbigny, 1850, (*Hortholus americanus*.) Prodr. d. Paléont., t. 1, p. 1, Black Riv. Gr.

apollo, Billings, 1862, Pal. Foss., vol. 1, p. 25, Calciferous Gr.

bickmoreanus, Whitfield, 1885, Bull. Am. Mus. Nat. Hist., vol. 1, p. 191, Niagara Gr.

cancellatus, McChesney, 1861, New Pal. Foss., Niagara Gr. See *L. occidentalis* and *Nautilus cancellatus* and *N. occidentalis*. If this species, as Prof. Hall suggests, is a true *Nautilus*, McChesney's name has precedence.

capax, see *Nautilus capax*.

complanatus, Shumard, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 107, Calciferous Gr.

convolvans, Schlotheim, 1813, in Jahrbuch, as identified by Hall, Pal. N. Y., vol. 1, p. 53. See *L. americanus*.

eatonii, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 331, Birdseye Gr.

eatonii var. *cassinensis*, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 332, Birdseye Gr.

farnsworthii, Billings, 1861, Pal. Foss., vol. 1, p. 21, Calciferous Gr.

grafftonensis, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil. p. 51, and Geo. Sur. Ill., vol. 6, p. 507, Niagara Gr.

hercules, Winchell & Marey, 1865, Mem. Bost. Soc. Nat. Hist., Niagara Gr. Syn. for *Cyrtoceras ampliacorne*. See 20th Rep. N. Y. Mus. Nat. Hist.

imperator, Billings, 1861, Pal. Foss., vol. 1, p. 23, Calciferous Gr.

magnificum, Billings, 1857, (*Gyroceras magnificum*.) Rep. of Progr. Geo. Sur. Can., p. 307, Hud. Riv. Gr. Type of Hyatt's genus *Aspidoceras*.

marshi, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 404, Niagara Gr.

internistriatus, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 332, Birdseye Gr.

multicostatus, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 67, and Geo. Wis., vol. 4, p. 303, Niagara Gr.

murchisoni, Troost. Not defined so as to be recognized.

FIG. 745.—*Lituites glauco-teus*. Quarter size.

Jour. Acad. Nat.
phyton, a sponge,
on tuberosum.

—Lituities gigan-
Quarter size.

septa simple; si-
circular. Type

1850, (Hortholus
Paléont., t. 1, p.

Foss., vol. 1, p.

1885, Bull. Am.
1, p. 191, Niag-

1861, New Pal.
e L. occidentalis
us and N. occi-
des, as Prof. Hall
utilus, McChes-
ence.

1863, Trans. St.
2, p. 107, Calcif-

13, in Jahrbuch,
al. N. Y., vol. 1,
us.

Bull. Am. Mus.
31, Birdseye Gr.
Whitfield, 1886,
Hist., vol. 1, p.

, Pal. Foss., vol.

then, 1870, Proc.
p. 51, and Geo.

Niagara Gr.

rey, 1865, Mem.

orne. See 20th
Hist.

Pal. Foss., vol.

1857, (Gyroceras
Progr. Geo. Sur.

v. Gr. Type of
ras.

Rep. N. Y. Mus.
ara Gr.

1886, Bull. Am.
1, p. 332, Birds-

1880, Ann. Rep.
and Geo. Wis.,

r.
defined so as to

niagarensis, Spencer, 1884, Bull. No. 1,
Mus. Univ. St. Mo., p. 60, Niagara Gr.
occidentalis, Hall, 1861, Rep. of Progr. Geo.
Sur. Wis., Niagara Gr. This species is
now referred by Prof. Hall to the genus
Nautilus, see 20th Rep. N. Y. St. Mus.
Nat. Hist., p. 400. It was first described
by McChesney, Jan. 1860, as *Cyrtoceras*
giganteum, but that name being preoc-
cupied, in 1861 he proposed *Lituities*
cancellatus. If it is a *Nautilus*, the
word *occidentalis* being preoccupied,
McChesney's name *cancellatus* has pre-
cedence.

ortoni, Meek, 1873, Ohio Pal., vol. 1, p.
186, Niagara Gr.

palinurus, Billings, 1862, Pal. Foss., vol. 1,
p. 25, Calciferous Gr.

pluto, Billings, 1865, Pal. Foss., vol. 1, p.
259, Quebec Gr.

robertsoni, Hall, 1861, Rep. of Progr. Wis.,
p. 38, Chazy and Black Riv. Grs.

seelyi, Whitfield, 1886, Bull. Am. Mus.
Nat. Hist., vol. 1, p. 330, Birdseye Gr.

undatus, Emmons, 1842, (Inachus unda-
tus.) Geo. Rep. N. Y., p. 394, and Pal.

N. Y., vol. 1, p. 52, Black Riv. and
Trenton Grs. It is not a *Lituities*.

undatus var. *occidentalis*, Hall, 1861, Rep.
of Progr. Wis., p. 38, Black Riv. and
Trenton Grs.

Melia cancellatus, Emmons, 1856, Am. Geol.
Not defined so as to be recognized.

cincinnati, D'Orbigny, 1850, Prodr. d.
Paléont., t. 1, p. 4. Not defined so as to
be recognized.

NAUTILUS, Bréynius, 1732, Dissert. Polyth.,
p. 11. [E.g. *Nautilus*, sailor or naviga-
tor.] Shell subglobose, compressed;
volutions coiled in the same plane, con-
tiguous; umbilicus open or closed;
septa simple, arched or waved on the
lateral margins; siphuncle central or
subcentral; lip sinuous on the dorsal
and ventro-lateral margins; surface
smooth, striate, costate, or bearing nodes.
Type *N. pompilius*.

acereus, Hall, 1879, Pal. N. Y., vol. 5, pt.
2, p. 417, Ham. Gr.

avonensis, see *Solenochilus avonense*.

avus, Barrande, 1869, Syst. Sil. de Boh.,
vol. 4, p. viii, pl. 435, Quebec Gr.

barrandi, Hall, 1876, see *N. Magister*.

biserialis, Hall, 1860, Sapp. to vol. 1, pt.
2, Iowa Geo. Sur., p. 92, Coal Meas.

bucinum, Hall, 1876, Illust. Devonian
Foss., pl. 60, and Pal. N. Y., vol. 5, pt.
2, p. 412, Ham. Gr. Type of Hyatt's

genus *Nephriticeras*.

calciferus, Billings, 1865, Pal. Foss., vol.
1, p. 258, Calciferous Gr.

cancellatus, McChesney, 1861, (*Lituities*
cancellatus.) New Pal. Foss., p. 96, Ni-
agara Gr.

canaliculatus, Cox, 1857, Geo. Sur. Ky.,
vol. 3, p. 575, Coal Meas. Type of Hy-
att's genus *Solenoceras*.

capax, Hall 1860, (*Lituities capax*.) Rep. of
Progr. Geo. Sur. Wis., p. 3, Niagara Gr.

capax, Meek & Worthen, 1865. This was
preoccupied and must yield unless it
can be retained in the subgenus *Solenochi-
lus*.

cavus, Hall, 1870, Pal. N. Y., vol. 5, p. 416,
Ham. Gr.

chAMPLAINENSIS, Whitfield, 1886, Bull. Am.
Mus. Nat. Hist., vol. 1, p. 329, Birds-
eye Gr.

chesterensis, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 469, and Geo.

Sur. Ill., vol. 2, p. 306, Kaskaskia Gr.

clarkanus, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 32, and Bull. Am. Mus. Nat.

Hist., p. 92, Warsaw Gr.

collectus, see *Solenochilus collectum*.

cornulum, Hall, 1876, Illust. Devonian
Foss., pl. 60, and Pal. N. Y., vol. 5, pt.
2, p. 414, Ham. Gr.

coXanus, see *Temnochilus coxanum*.

dANVILLENSIS, White, 1878, Proc. Acad.
Nat. Sci., p. 36, and Cont to Pal., No. 8,
p. 170, Coal Meas.

decoratus, Cox, 1857, Geo. Sur. Ky., vol.
3, p. 572, Coal Meas.

desertus, Billings, 1865, Pal. Foss., vol. 1,
p. 258, Quebec Gr.

digonus, see *Trematodiscus digonus*.

disciformis, see *Discites disciformis*.

discoidalis, see *Trematodiscus discoidalis*.

divisus, White & St. John, 1868, Trans.
Chi. Acad. Sci., p. 124, Up. Coal Meas.

The name was preoccupied by Meyer
in 1831.

eccentricus, Meek & Hayden, 1858, Trans.
Alb. Inst., vol. 4, p. 83, and Pal. Up.

Mo., p. 65, Permian Gr.

FIG. 746.—*Nautilus faberi*. Lateral and front
views. Magnified 2 diam.

faberi, n. sp. Shell small, smooth, all vo-
lutions embraced in the outer one,
leaving only a small round umbilicus;
aperture semielliptical above the inter-
ior volution. The species is founded
on a single specimen having a piece
chipped from the dorsal side of the last
volution, and a small piece broken from
the inner volution, but nowhere dis-
closing the septa. It was found in the
Coal Measures, on Elk Horn Creek,
Kentucky, and belongs to Mr. Charles
Faber's collection.

ferox, Billings, 1865, Pal. Foss., vol. 1, p.
351, Calciferous Gr.

ferratus, Cox, 1857, Geo. Sur. Ky., vol. 3, p. 574, Coal Meas.

FIG. 747.—*Nautilus forbesianus*.

forbesianus, McChesney, 1860, Desc. New Pal. Foss., p. 63, and Trans. Chi. Acad. Sci., p. 50, Coal Meas.
gipini, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 658, Coal Meas.
globatus, Sowerby, 1825, Min. Conch., vol. 5, p. 129, Kaskaskia Gr.
hercules, Billings, 1857, Rep. of Progr. Can. Geo. Sur., p. 306, Hud. Riv. Gr.
highlandensis, see *Discites highlandensis*.
hyatti, Beecher, 1888, Pal. N. Y., vol. 7, p. 37, Ham. Gr.
illinoisensis, McChesney, 1860, Desc. New Pal. Foss., p. 64, Coal Meas.
ingentior, Winchell, 1862, Am. Jour. Sci., 2d series, vol. 33, p. 361, Marshall Gr.
insolens, Billings, 1865, Pal. Foss., vol. 1, p. 258, Quebec Gr.
jason, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 464, Chazy Gr. Type of Hyatt's genus *Plectoceras*.
kelloggii, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 328, Birdseye Gr.
lasallensis, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 261, and Geo. Sur. Ill., vol. 5, p. 610, Up. Coal Meas.
latus, see *Temnochilus latus*.
lawsi, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 658, Ham. Gr.
leidy, see *Solenochilus leidy*.
liratus, Hall, 1860, (*Gyroceras liratum*), 13th Rep. N. Y. Mus. Nat. Hist., p. 104, Marcellus Shale.
liratus var. *juvenis*, Hall, 1879, Pal. N. Y., vol. 5, Ham. Gr.
magister, Hall, 1879, Pal. N. Y., vol. 5, p. 422, Ham. Gr. Proposed instead of *N. barrandi*, Hall, which was preoccupied.
marcellensis, see *Discites marcellensis*.
maximus, Conrad, 1838, (*Cyrtoceras maximus*) Ann. Rep. N. Y., p. 117, and Pal. N. Y., vol. 5, pt. 2, p. 418, Ham. Gr.
meekianus, see *Trematodiscus meekianus*.
missouriensis, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 198, Coal Meas.
montgomeryensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 4, and Geo. Sur. Ill., vol. 8, p. 148, Up. Coal Meas.
natator, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 466, Chazy Gr. Type of Hyatt's genus *Barrandoceras*.

niotensis, see *Temnochilus niotense*.
nodocarinatus, McChesney syn. for *N. occidentalis*.
nodoso-dorsatus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 198, Coal Meas.
occidentalis, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 196, and Pal. E. Neb., p. 234, Permian Gr.
occidentalis, Hall, 1860, 20th Rep. N. Y. Mus. Nat. Hist., p. 400, Niagara Gr. This name being preoccupied, McChesney's name *cancellatus*, has precedence. See *Lituities cancellatus*.
oceanus, Hall, 1879, Desc. New Spec. Foss., p. 19, and 11th Rep. Geo. and Nat. Hist. Ind., p. 325, Niagara Gr.
oriens, Hall, 1876, Illust. Devonian Foss., pl. 61, and Pal. N. Y., vol. 5, pt. 2, p. 420, Marcellus Shale.
ornatus, Hall, 1860, syn. for *N. marcellensis*.
ortoni, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 231, Coal Meas.
parallelus, Beecher, 1888, Pal. N. Y., vol. 7, p. 38, Coal Meas.
pauper, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 226, Kaskaskia Gr.
permanianus, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 196, Permian Gr.
planidorsalis, see *Trematodiscus planidorsalis*.
planorbiformis, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 469, and Geo. Sur. Ill., vol. 2, p. 386, Coal Meas.
planovolvis, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 198, Coal Meas.
pomponius, Billings, 1862, Pal. Foss., vol. 1, p. 26, Calciferous Gr.
ponderosus, White, 1872, Pal. of E. Neb., p. 236, Coal Meas. Type of Hyatt's genus *Titanoceras*.
quadrangularis, McChesney, 1860, Desc. New Pal. Foss., p. 65, and Trans. Chi. Acad. Sci., vol. 1, p. 57, Coal Meas. Type of Hyatt's genus *Tainoceras*.
rockfordensis, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 275, Kinderhook Gr. Probably a *Gyroceras*. See Ill. Geo. Sur., vol. 3.
sangamonensis, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 470, and Geo. Sur. Ill., vol. 2, p. 386, Coal Meas. Type of Hyatt's genus *Metaceras*.
seebachianus, see *Pteronutilus seebachianus*.
spectabilis, see *Temnochilus spectabile*.
springeri, see *Solenochilus springeri*.
striatulus, see *Trematodiscus striatulus*.
subglobosus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 469, syn. for *N. globatus*, see Geo. Sur. Ill., vol. 3, p. 305.
subliratus, Hall, 1876, (*Gyroceras subliratum*), Illust. Devon. Foss., pl. 58, and Pal. N. Y., vol. 5, pt. 2, p. 409, Ham. Gr.
subquadrangularis, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 232, Coal Meas.

is niotense.
syn. for N. oc.

rd, 1858, Trans.
l. 1, p. 198, Coal

1858, Trans. St.
p. 196, and Pal.
in Gr.

20th Rep. N. Y.
1900, Niagara Gr.
occupied, McChes-
has precedence.

Desc. New Spec.
Rep. Geo. and
Niagara Gr.
Devonian Foss.,
vol. 5, pt. 2, p.

for N. marcel-

nn. N. Y. Acad.
Meas.

Pal. N. Y., vol.

Ann. N. Y. Acad.
askia Gr.

1858, Trans. St.
1, p. 196, Per-

atodiscus plani-

Worthen, 1860,
hil., p. 469, and
386, Coal Meas.
1858, Trans. St.
p. 198, Coal Meas.
Pal. Foss., vol.

Pal. of E. Neb.,
of Hyatt's ge-

hey, 1860, Desc.
and Trans. Chi.

57, Coal Meas.
tainoceras.

Worthen, 1860,
hil., p. 275, Kin-
y a Gyroceras.

3.

Worthen, 1860,
hil., p. 470, and
386, Coal Meas.

Metacoceras.

utilus seebuch-

is spectabile.

springeri.

us striatulus.

hen, 1860, Proc.

p. 469, syn. for

Sur. Ill., vol. 3,

roceras sublira-

ss., pl. 58, and

p. 409, Ham. Gr.

eld, 1882, Ann.

232, Coal Meas.

subsulcatus, Phillips, 1836, Geo. York. Not
clearly identified in this country.

sulcatus, see *Trematodiscus sulcatus*.

trigonus, see *Trematodiscus trigonus*.

trisulcatus, see *Trematodiscus trisulcatus*.

tyrans, Billings, 1859, Can. Nat. and Geo.,

vol. 4, p. 465, Chazy Gr.

versutus, Billings, 1865, Pal. Foss., vol. 1,

p. 259, Quebec Gr. Type of Hyatt's

genus *Litoceras*.

winslowi, see *Temnochilus winslowi*.

Nelimenia incognita, Castelnau, 1843, Syst.

Sil., p. 33. Probably a fragment of

Phragmoceras or *Oncoceras*.

ONCOCERAS, Hall, 1847, Pal. N. Y., vol. 1, p.

190. [Ety. *onkos*, swelling; *keras*, horn.]

Curved, aperture constricted; lower

part of the body chamber, and upper

part of septate portion ventricose; ab-

solutely contracted to-

ward the apex; siphun-

cle dorsal; septa plane,

nearly flat, slightly ele-

vated on the dorsal

margin. Type *O. con-*

strictum.

abruptum, Hall, 1861, Rep.

of Progr. Wis., p. 44,

Trenton Gr.

alceus, Hall, 1861, Rep. of

Progr. Wis., p. 46, Chazy

and Black Riv. Grs.

amator, Billings, 1866,

Catal. Sil. Foss. Antic.,

p. 59, Clinton Gr.

brevicurvatum, Whitfield, 1880, Ar. Rep.

Geo. Sur. Wis., p. 59, and Geo. Wis.,

vol. 4, p. 234, Trenton Gr.

constrictum, Hall, 1847, Pal. N. Y., vol. 1,

p. 197, Black Riv. and Trenton Grs.

dilatatum, Hall, 1860, 13th Rep. N. Y.

Mus. Nat. Hist., p. 105, Ham. Gr.

expansum, Hall, 1852, Pal. N. Y., vol. 2,

p. 337, Coralline limestone.

futile, Billings, 1866, Catal. Sil. Foss.

Antic., p. 59, Clinton Gr.

gibbosum, Hall, 1852, Pal. N. Y., vol. 2,

p. 13, Medina sandstone.

lycus, Hall, 1861, Rep. of Progr. Wis., p.

45, Chazy and Black Riv. Gr.

mummiforme, Whitfield, 1880, Ann. Rep.

Geo. Sur. Wis., p. 58, and Geo. Wis.,

vol. 4, p. 232, Trenton Gr.

orcas, Hall, 1861, (Cyrtoceus *orcas*.) Rep.

of Progr. Geo. Sur. of Wis., p. 42, Niag-

ara Gr.

ovoides, Hall, 1859, Pal. N. Y., vol. 3, p.

342, Low. Held. Gr.

pandion, Hall, 1861, Rep. of Progr. Wis.,

p. 45, and Geo. Wis., vol. 4, p. 233,

Chazy and Black Riv. Grs.

pettiti, Billings, 1866, Catal. Sil. Foss.

Antic., p. 36, Niagara Gr.

plebeium, Hall, 1861, Geo. Rep. Wis., p.

44, Trenton Gr.

subrectum, Hall, 1852, Pal. N. Y., vol. 2,

p. 94, Clinton Gr.

teucer, Billings, 1866, Catal. Sil. Foss.

Antic., p. 86, Niagara Gr.

thales, Billings, 1866, Catal. Sil. Foss.

Antic., p. 87, Niagara Gr.

vasiforme, Dwight, 1884, Am. Jour. Sci.

and Arts, 3d ser., vol. 27, p. 257, Calcif-

erous Gr.

ORMOCERAS, Stokes, 1840, Trav. S. Geo. Soc.,

2d ser., vol. 5, p. 799. [Ety. *ormos*, a

chain or necklace; *keras*, horn; from

the appearance of the siphuncle.] Ex-

ternally like *Orthoceras*, and dis-

tinguished only by having the siphun-

cle constricted within each chamber in-

stead of at the place of union with the

septa. Types *O. backi*, and *O. bayfieldi*.

backi, Stokes, 1840, Trans. Geo. Soc., 2d

ser., vol. 5, p. 709, Clinton Gr.

bayfieldi, Stokes, 1840, Trans. Geo. Soc.,

2d ser., vol. 5, p. 709, Clinton Gr.

crebrisep-

tum, Hall,

1847, Pal.

N. Y., vol.

1, p. 313,

Ind. Riv.

Gr.

gracile, Hall,

1847, Pal.

N. Y., vol.

1, p. 58,

Black Riv.

Gr.

remotisep-

tum, Hall,

1850, 3d

Rep. N. Y. Mus. Nat. Hist., p. 181,

Trenton Gr.

tenuifilum, Hall, 1847, Pal. N. Y., vol. 1,

p. 55, Black Riv. and Trenton Gr.

tenuifilum var. *distans*, Hall, 1847, Pal.

N. Y., vol. 1, p. 58, Black Riv. Gr.

vertebratum, Hall, 1852, Pal. N. Y., vol.

2, p. 94, Clinton Gr.

whitii, Stokes, 1840, Trans. Geo. Soc., 2d

ser., vol. 5, p. 709, Clinton Gr.

ORTHOCERAS, Breynius, 1732, Dissertatio

physica de Polythalamis. [Ety. *orthos*,

straight; *keras*, horn.] Shell conical,

straight, or nearly so; body chamber

large, behind which the shell is com-

posed of numerous chambers separated

by convex, transverse septa, with simple

edges, at right angles to the longer axis

of the shell; siphuncle central, sub-

central or eccentric, cylindrical or di-

lated in the chambers; surface smooth

or transversely, or longitudinally stri-

ated, or furrowed. Typical *O. breynii*,

O. annulatum, and *O. striatum*.

abnorme, Hall, 1867, 20th Rep. N. Y.

Mus. Nat. Hist., p. 415, Niagara Gr.

abruptum, Hall, 1852, Pal. N. Y., vol. 2,

p. 97, Clinton Gr.

acicula, Hall, see *Coleolus acicula*.

aciculoides, Clarke, 1885, Bull. U. S. Geo.

Sur., No. 16, p. 51, Chemung Gr.

aculeatum, Swallow, 1858, Trans. St. Louis

Acad. Sci., vol. 1, p. 200, Coal Meas.

agea, Hall, 1862, 15th Rep. N. Y. Mus.

Nat. Hist., p. 80, Ham. Gr.

FIG. 748.—*Oncoceras constrictum*.

FIG. 749.—*Ormoceras bayfieldi*.

- aequale*, Emmons, 1842, Geo. Rep. N. Y., p. 404, Hud. Riv. Gr.
- alienum*, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 414, Niagara Gr.
- alumettense*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 331, Chazy and Black Riv. Grs.
- amplicameratum*, Hall, 1847, Pal. N. Y., vol. 1, p. 205, Black Riv. and Trenton Grs.
- amycus*, Hall, 1879, Desc. New Spec. Foss., p. 18, and 11th Rep. Geo. Sur. Ind., p. 324, Niagara Gr.
- anax*, Billings, 1875, Can. Nat. and Geol., vol. 7, p. 238, Up. Held. Gr.
- anguis*, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 312, Chemung Gr.
- angulatum*, (?) Wahlenberg, 1821, Nova Acta. Soc. Sci. Upsal., p. 90, Niagara Gr. See remarks on this species by Prof. Hall in 20th Rep. N. Y. Mus. Nat. Hist., p. 413.
- anellus*, Conrad, 1843, Proc. Acad. Nat. Sci. Phil., vol. 1, p. 334, and Pal. N. Y., vol. 1, p. 202, Black Riv. and Trenton Grs.
- annulato-costatum*, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 147. This name was preoccupied by Boll in 1857. See *O. randolphense*.
- annulatum*, Sowerby, 1818, Min. Conch., vol. 2, p. 77, Clinton and Niagara Grs.
- anterior*, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 463, Chazy Gr.
- anticostiense*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 316, Hud. Riv. Gr.
- aptum*, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 282, Marcellus Shale.
- arcuatellum*, Sandberger. Is not an American species.
- arcuoliratum*, Hall, 1847, Pal. N. Y., vol. 1, p. 198, Black Riv. and Trenton Grs.
- arenosum*, Hall, 1859, Pal. N. Y., vol. 3, p. 480, Oriskany sandstone.
- asmodeus*, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 30, Genesee Shale.
- atreus*, Hall, 1879, Pal. N. Y., vol. 5, p. 305, Portage Gr.
- atticus*, Billings, 1865, Pal. Foss., vol. 1, p. 312, Quebec Gr.
- aulax*, Hall, 1879, Pal. N. Y., vol. 5, p. 293, Ham. Gr.
- autolytus*, Billings, 1862, Pal. Foss., vol. 1, p. 91, Quebec Gr.
- baculum*, Meek, 1860, Proc. Acad. Nat. Sci., p. 310, Subcarboniferous.
- baculum*, Hall, 1862. The name was preoccupied. See *O. stylus*.
- balteatum*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 318, Hud. Riv. Gr.
- barquianum*, Winchell, 1862, Am. Jour. Sci., 2d ser., vol. 33, Marshall Gr.
- bartonense*, Spencer, 1884, Bull. No. 1, Mus. Univ. St. Mo., p. 60, Niagara Gr.
- bebryx*, Hall, 1876, Illust. Devonian Foss., pl. 39, and Pal. N. Y., vol. 5, pt. 2, p. 275, Ham. Gr.
- bebryx* var. *cayuga*, Hall, 1879, Pal. N. Y., vol. 5, p. 276, Chemung Gr.
- becki*, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 362, Calciferous Gr.
- bellatulum*, Billings, 1866, Catal. Sil. Foss. Antic., p. 58, Clinton Gr.
- beloitense*, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 97, and Geo. Wis., vol. 4, p. 226, Trenton Gr.
- bilineatum*, Hall, 1847, Pal. N. Y., vol. 1, p. 199, Chazy, Black Riv., Trenton, and Hud. Riv. Grs.
- bilineatum* var. *a*, Hall, 1847, Pal. N. Y., vol. 1, p. 200, Trenton Gr.
- bipartitum*, Hall, 1879, Pal. N. Y., vol. 5, p. 313, Up. Chemung Gr.
- brainerdi*, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 319, Birdseye Gr.
- brongniarti*, Troost, 1838, (Conotubularia *brongniarti*.) Mem. Soc. de France, 3, p. 89, Low. Sil.
- brontes*, Billings, 1866, Catal. Sil. Foss. Antic., p. 83, Niagara Gr.
- bucklandi*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 330, Up. Sil.

FIG. 750.—*Orthoceras byrnesi*.

- bullatum*, (?) Sowerby, 1839, Murch. Sil. Syst., p. 705, Trenton Gr.
- byrnesi*, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 126, and Jour. Cin. Soc. Nat. Hist., vol. 4, p. 319, Hud. Riv. Gr.
- cadmus*, Billings, 1866, Catal. Sil. Foss. Antic., p. 83, Niagara Gr.
- cælamen*, Hall, 1879, Pal. N. Y., vol. 5, p. 298, Ham. Gr.
- cameolare*, McChesney, 1861, New Pal. Foss., p. 93, Niagara Gr.
- canadense*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 321, Mid. Sil. Prof. Billings proposed this name as a substitute for *Huronina vertebralis* for the reason that *Huronina* is a syn. for *Orthoceras*, and there is one *O. vertebralis*.
- cancellatum*, Hall, 1852, Pal. N. Y., vol. 2. The name was preoccupied by Eichwald in 1842. See *O. subcancellatum*.
- capitolinum*, Safford, 1869, Geo. of Tenn., p. 290, Trenton Gr.
- carleyi*, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 98, Hud. Riv. Gr.
- carltonense*, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 85, and Geo. Wis., vol. 4, p. 318, Niagara Gr.
- carnosum*, Hall, 1879, Pal. N. Y., vol. 5, p. 258, Scholarie grit.
- cataline*, Billings, 1865, Pal. Foss., vol. 1, p. 315, Quebec Gr.

1879, Pal. N. Y.,
ing Gr.
i. Nat. and Geol.,
us Gr.
6, Catal. Sil. Foss.
Gr.
878, Ann. Rep.
and Geo. Wis.,
Gr.
Pal. N. Y., vol. 1,
iv., Trenton, and
1847, Pal. N. Y.,
Gr.
Pal. N. Y., vol. 5,
Gr.
1886, Bull. Am.
1, p. 319, Birds-
, (Conotubularia
Geo. de France,
Catal. Sil. Foss.
Gr.
Rep. of Progr.
Up. Sil.

byrnesi.

1839, Murch. Sil.
r.
1875, Cin. Quar.
and Jour. Cin.
4, p. 319, Hud.
Catal. Sil. Foss.
Gr.
N. Y., vol. 5, p.
1861, New Pal.
r.
Rep. of Progr.
Mid. Sil. Prof.
name as a sub-
tribe for the
is a syn. for
is one O. verte-
al. N. Y., vol. 2,
upied by Eich-
abancellatun.
Geo. of Tenn.,
1875, Ohio Pal.,
Gr.
1878, Ann. Rep.
and Geo. Wis.,
Gr.
il. N. Y., vol. 5,
al. Foss., vol. 1,

catro, Billings, 1865, Pal. Foss., vol. 1, p.
314, Quebec Gr.
catulus, Billings, 1865, Pal. Foss., vol. 1,
p. 313, Quebec Gr.
chemungense, Swallow, 1860, Trans. St.
Louis Acad. Sci., vol. 1, p. 660, Waverly
or Choteau Gr.
chesterense, Swallow, 1863, Trans. St.
Louis Acad. Sci., vol. 2, p. 98, Kaskas-
kia Gr.
chouteauense, Swallow, 1860, (O. che-
mungense var. chouteauense.) Trans. St.
Louis Acad. Sci., vol. 1, p. 660, Waverly
or Choteau Gr.
cincinnatiense, S. A. Miller, 1875, Cin.
Quar. Jour. Sci., vol. 2, p. 127, and
Jour. Cin. Soc. Nat. Hist., vol. 4, p. 319,
Hud. Riv. Gr.
cingulum, Hall, 1879, Pal. N. Y., vol. 5,
p. 240, Schoharie grit.
clathratum, Hall, 1847, Pal. N. Y., vol. 1,
p. 201, Trenton Gr.
clavatum, Hall, 1852, Pal. N. Y., vol. 2, p.
104, Clinton Gr.
clavatum, Hall, 1859. The name was ap-
propriated. See O. desideratum.
clinocameratum, Winchell, 1862, Am.
Jour. Sci., 2d ser., vol. 33, p. 356, Mar-
shall Gr.
clintoni, Hall, 1877, 1st Ed. Am. Pal.
Foss., p. 244, Chazy Gr. Proposed in-
stead of O. subarcuatum, Hall, 1847,
which was preoccupied.
clonei, Barrande, 1869, Sys. Sil. de Boh.,
4me ser., p. viii, pl. 432 to 434, Que-
bec Gr.
cochleatum, Hall, 1879, Pal. N. Y., vol. 5.
The name was preoccupied by Schlot-
heim in 1813. See O. warrenense.
collatum, Hall, 1879, Pal. N. Y., vol. 5, p.
252, Schoharie grit.
colon, White, 1874, Rep. Invert. Foss., p.
10, and Geo. Sur. W. 100th Mer., vol. 4,
p. 56, Quebec Gr.
columnare, Hall, 1860, Rep. Progr. Geo.
Sur. Wis. The name was preoccupied
by Mark in 1857. See O. orus.
conicum, Castelnau, 1843, Syst. Sil., p. 29.
The name was preoccupied by His-
inger.
consortale, Beecher, 1888, Pal. N. Y., vol.
7, p. 29, Chemung Gr.
constrictum, Vanuxem, 1842, Geo. Rep.
3d Dist. N. Y., p. 152, and Pal. N. Y.,
vol. 5, pt. 2, p. 288, Ham. Gr.
constrictum, Conrad, 1838. Not defined so
as to be recognized.
constrictum, see Oncoceras constrictum.
coralliferum, Hall, 1847, Pal. N. Y., vol.
1, p. 312, Utica and Hud. Riv. Grs.
cornuoryx, Whitfield, 1886, Bull. Am.
Mus. Nat. Hist., vol. 1, p. 320, Birds-
eye Gr.
cornum, Billings, 1857, Rep. of Progr.
Geo. Sur. Can., p. 329, Chazy Gr.
crebescens, Hall, 1867, 20th Rep. N. Y.
Mus. Nat. Hist., p. 354, Niagara Gr.
crebristatum, Meek & Worthen, 1865.
Proc. Acad. Nat. Sci. Phil., p. 255, and

Geo. Sur. Ill., vol. 6, p. 503, Niag-
ara Gr.
creon, Hall, 1879, Pal. N. Y., vol. 5, p.
260, Schoharie grit.
cribrosum, Geinitz, 1866, Carb. und Dyas
in Neb., p. 4, and Pal. E. Neb., p. 234,
Coul. Meas.
crocus, Billings, 1866, Catal. Sil. Foss.
Antic., p. 22, Hud. Riv. Gr. Proposed
instead of O. perannulatum, which was
preoccupied.
crotalum, Hall, 1862, 15th Rep. N. Y.
Mus. Nat. Hist., p. 78, Ham. Gr. Type
of Hyatt's genus Spyroceras.
cuvieri, Troost, 1838, (Conotubularia cu-
vieri,) Mem. Soc. Geo. de France, t. 3,
p. 88, Low. Sil.
dagon, Beecher, 1888, Pal. N. Y., vol. 7,
p. 28, Up. Held. Gr.
darwini, Billings, 1868, Pal. Foss., vol. 1,
p. 161, Guelph Gr.
dawsonanum, S. A. Miller, 1883, 2d Ed.
Am. Pal. Foss., p. 307, Carboniferous.
Proposed instead of O. perstrictum,
Dawson, in Acadian Geology, p. 312,
fig. 129, as the name was preoccupied
by Barrande.
decrescens, Billings, 1857, Rep. of Progr.
Geo. Sur. Can., p. 337, Black Riv. and
Trenton Gr.
defranchi, Troost, 1838, (Conotubularia de-
franchi,) Mem. Soc. Geo. de France, t. 3,
p. 90, Low. Sil.
demon, Hall, 1879, Pal. N. Y., vol. 5, p.
311, Chemung Gr.
deparcum, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 363, Calciferous Gr.
desideratum, Hall, 1877, 1st Ed. Am. Pal.
Foss., p. 244, Low. Held. Gr. Proposed
instead of O. clavatum, Hall, 1859, which
was preoccupied.
diffidens, Billings, 1865, Pal. Foss., vol. 1,
p. 174, Chazy Gr.
directum, Beecher, 1888, Pal. N. Y., vol. 7,
p. 27, Up. Held. Gr.
dolatum, Dawson, 1868, Acad. Geol. p.
311, Carboniferous.
drummondii, Billings, 1865, Pal. Foss., vol.
1, p. 173, Black Riv. Gr.
duramen, Beecher, 1888, Pal. N. Y., vol.
7, p. 25, Schoharie grit.
duseri, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 97, Hud. Riv. Gr.

FIG. 751.—Orthoceras dyeri.

dyeri, S. A. Miller, 1875, Cin. Quar. Jour.
Sci., vol. 2, p. 125, and Jour. Cin. Soc.
Nat. Hist., vol. 3, p. 236, Hud. Riv. Gr.

edax, Billings, 1865, Pal. Foss., vol. 1, p. 340, Calcif. Gr.
 elegantulum, Dawson, 1860, Can. Nat. and Geo., vol. 5, p. 155, and Acad. Geol., p. 607, Up. Sil.
 emaceratum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 170, Ham. Gr.
 epigrus, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 33, and Bull. Am. Mus. Nat. Hist., p. 91, Warsaw Gr.
 eriense, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 274, and Pal. N. Y., vol. 5, p. 274, Ham. Gr. Proposed instead of *O. robustum*, which was preoccupied.
 eurekaense, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 265, Subcarboniferous.
 exile, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 78, and Pal. N. Y., vol. 5, pt. 2, p. 290, Ham. Gr.
 exornatum, Dawson, 1860, Can. Nat. and Geo., vol. 5, p. 198, Up. Sil.
 expansum, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 468, and Geo. Sur. Ill., vol. 2, p. 286, St. Louis Gr.
 explorator, Billings, 1865, Pal. Foss., vol. 1, p. 253, Quebec Gr.
 expositum, Beecher, 1888, Pal. N. Y., vol. 7, p. 29, Chemung Gr.
 ferum, Billings, 1866, Catal. Sil. Foss. Antic., p. 22, Hud. Riv. and Anticosti Gr.
 filiforme, Castelnau, 1843, Syst. Sil., p. 30, Niagara Gr. Not recognized.
 filosum, Clarke, 1885, Bull. U. S. Geo. Sur., No. 16, p. 52, Chemung Gr.
 flavus, Billings, 1865, Pal. Foss., vol. 1, p. 255, Quebec Gr.
 fluctum, Hall, 1879, Pal. N. Y., vol. 5, p. 239, Schoharie grit.
 foliatum, syn. for *Cyrtoceras eugenium*.
 formosum, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 317, Trenton, Hud. Riv., and Anticosti Grs.
 fosteri, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 127, and Jour. Cin. Soc. Nat. Hist., vol. 4, p. 319, Hud. Riv. Gr.
 foxense, Safford, 1869, Geo. of Tenn. Not defined.
 fulgidum, Hall, 1879, Pal. N. Y., vol. 5, p. 310, Chemung Gr.
 fulgur, Billings, 1866, Catal. Sil. Foss. Antic., p. 22, Hud. Riv. Gr. Proposed instead of *O. propinquum*, which was preoccupied.
 fulvum, Billings, 1865, Pal. Foss., vol. 1, p. 348, Calcif. Gr.
 fusiforme, Hall, 1847, Pal. N. Y., vol. 1, p. 60, Black Riv. and Trenton Grs.
 fustis, Hall, 1879, Pal. N. Y., vol. 5, p. 281, Marcellus Shale.
 glaucus, Billings, 1865, Pal. Foss., vol. 1, p. 350, Calciferous Gr.
 goldfussi, Troost, 1838, (Conotubularia goldfussi.) Mem. Soc. Geo. de France, t. 3, p. 90, Low. Sil.
 gracilium, Winchell, 1862, Proc. Acad. Nat. Sci., p. 429, Portage Gr.

gregarium, Hall, 1861, Rep. of Progr. Wis. Preoccupied by Sowerby in 1839, Murch. Sil. Syst. See *O. sociale*.
 griffithi, Haughton, 1857, Jour. Roy. Dub. Soc., vol. 1, Devonian.?
 hesitans, Billings, 1865, Pal. Foss., vol. 1, p. 254, Quebec Gr.
 hageri, Hall, 1861, Geol. of Vermont, p. 718, Calciferous Gr.

FIG. 752.—*Orthoceras hallanum*.

hallanum, S. A. Miller, 1877, 1st Ed. Am. Pal. Foss., p. 245, Hud. Riv. Gr. Proposed instead of *O. halli*, in Cin. Quar. Jour. Sci., vol. 2, p. 128, which was preoccupied by Barrande.
 halli, see *O. hallanum*.
 harperi, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 128, and Jour. Cin. Soc. Nat. Hist., vol. 4, p. 319, Hud. Riv. Gr.
 hartmanum, S. A. Miller, 1883, 2d Ed. Am. Pal. Foss., p. 307, Carboniferous. Proposed instead of *O. laqueatum*, Hartt, in Acadian Geol., p. 312, fig. 128, which was preoccupied.
 hastatum, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 333, Black Riv. and Trenton Grs. Type of Hyatt's genus *Tripteroceas*.
 helderbergiae, Hall, 1859, Pal. N. Y., vol. 3, p. 345, Low. Held. Gr.
 henrietta, Dwight, 1884, Am. Jour. Sci. and Arts, 3d ser., vol. 27, p. 256, Calciferous Gr.
 herculaneum, Verneuil, 1846, Bull. de la Soc. Geol. de France, vol. 4, Low. Sil.
 hercules, Castelnau, 1843, Syst. Sil., p. 29, Up. Sil. Not recognized.
 heterocinctum, Winchell, 1863, Proc. Acad. Nat. Sci., p. 23, Kinderhook Gr.
 hindei, James. Founded on fragments of different species, most of them *O. transversum*.
 hoyi, McChesney, 1861, New Pal. Foss., p. 92, Niagara Gr.
 huronense, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 337, Trenton Gr.
 hyas, Hall, 1862, syn. for *O. thoas*.
 icarus, Beecher, 1888, Pal. N. Y., vol. 7, p. 31, Kinderhook Gr.
 idmon, Hall, 1879, Pal. N. Y., vol. 5, p. 302, Ham. Gr.
 illinoisense, Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 323, and Geo. Sur. Ill., vol. 8, p. 148, Kaskaskia Gr.

ep. of Progr. Wis.
oy in 1839, Murch.
ale.
Jour. Roy. Dub.
?
Pal. Foss., vol. 1,
d. of Vermont, p.

hallanum.

1877, 1st Ed. Am.
d. Riv. Gr. Pro-
alli, in Cin. Quar.
8, which was pre-

1875, Cin. Quar.
8, and Jour. Cin.
4, p. 319, Hud.

1883, 2d Ed. Am.
boniferous. Pro-
aqueatum, Hartt,
2, fig. 128, which

, Rep. of Progr.
3, Black Riv. and
of Hyatt's genus

Pal. N. Y., vol.
Gr.
Am. Jour. Sci.
27, p. 256, Cal-

1846, Bull. de la
vol. 4, Low. Sil.
Syst. Sil., p. 29,
ed.

ill, 1863, Proc.
Kinderhook Gr.
on fragments of
of them O. trans-

New Pal. Foss.,
7, Rep. of Progr.
Trenton Gr.
O. thoas.

al. N. Y., vol. 7,
N. Y., vol. 5, p.

83, Geo. Sur. Ill.,
Sur. Ill., vol. 8,

imbricatum, Sowerby, 1839, Murch. Sil.
Syst., p. 620, and Pal. N. Y., vol. 2, p.
291, Niagara Gr.

inceptum, Foerste, 1885, Bull. Sci. Lab.
Denison Univ., p. 117. Not properly
defined.

inlagator, Billings, 1865, Pal. Foss., vol.
1, p. 349, Calciferous Gr.

indianense, Hall, 1860, 13th Rep. N. Y.
Mus. Nat. Hist., p. 107, Kinder-
hook Gr.

infelix, Billings, 1866, Catal. Sil. Foss.
Antic., p. 57, Clinton Gr.

inoptatum, Hall, 1879, Pal. N. Y. vol. 5,
p. 267, Up. Held. Gr.

lowense n. sp. Devonian. Proposed in-
stead of O. undulatum in Rep. on Min.
Lands, p. 69, pl. 12, fig. 6, which name
was preoccupied.

irregulare, McChesney, 1861, New Pal.
Foss., Niagara Gr. The name was pre-
occupied. See O. woodworthi.

isogramma, Meek, 1871, Proc. Acad. Nat.
Sci., p. 172, Coal Meas.

jaculum, Hall, 1879, Pal. N. Y., vol. 5, p.
266, Up. Held. Gr.

jamesi, Hall & Whitfield, 1875, Ohio Pal.
vol. 2, p. 118, Clinton Gr.

jolietense, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci. Phil., p. 256, and Geo.
Sur. Ill., vol. 6, p. 505, Niagara Gr.

juncum, Hall, 1847, Pal. N. Y., vol. 1, p.
204, Trenton Gr.

kickapooense, Swallow, 1858, Trans. Acad.
Sci. St. Louis, vol. 1, p. 197, Up. Per-
mian Gr.

kingi, Meek, 1877, U. S. Geo. Sur. 40th
Parallel, vol. 4, p. 47, Devonian.

knoxense, McChesney, 1860, New Pal.
Foss., p. 69, Coal Meas.

lave, Hall, 1843, Geo. Rep. 4th Dist. N. Y.
The name was preoccupied by Fleming
in 1825. See O. sublave.

lamarecki, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 362, Calciferous Gr.

lamellosum, Hall, 1847, Pal. N. Y., vol. 1,
p. 312, Hud. Riv. Gr.

laphami, McChesney, 1861, New Pal.
Foss., p. 91, Niagara Gr.

laqueatum, Hall, 1847, Pal. N. Y., vol. 1,
p. 13, Calciferous to Trenton Gr.

laqueatum var. a, Hall, 1847, Pal. N. Y.,
vol. 1, p. 206, Trenton Gr.

laqueatum, Hartt, 1868, Acad. Geol. The
name was preoccupied. See O. Hartt-
anum.

lasallense, Worthen, 1883, Geo. Sur. Ill.,
vol. 7, p. 324, and Geo. Sur. Ill., vol. 8,
p. 149, Coal Meas.

lathropanum, Winchell, 1862, Am. Jour.
Sci. and Arts, 2d ser., vol. 33, p. 357,
Marshall Gr.

latiannulatum, Hall, 1847, Pal. N. Y., vol.
1, p. 204, Trenton Gr.

leander, Hall, 1879, Pal. N. Y., vol. 5, p.
309, Chemung Gr.

lima, Hall, 1879, Pal. N. Y., vol. 5, p. 303,
Ham. Gr.

lineolatum, McChesney, 1861, New Pal.
Foss., p. 93, Niagara Gr. The name
was preoccupied by Phillips in 1841.

lintonum, Hall, 1879, Pal. N. Y., vol. 5, p.
277, Ham. Gr.

longicameratum, Hall, 1859, Pal. N. Y.,
vol. 3, p. 343, Low. Held. Gr.

loxias, Hall, 1867, 20th Rep. N. Y. Mus.
Nat. Hist., p. 416, Low. Sil.

luxum, Hall, 1876, Illust. Devonian Foss.
pl. 35, and Pal. N. Y., vol. 5, pt. 2, p.
244, Schoharie grit.

lyelli, Billings, 1857, Rep. of Progr. Geo.
Sur. Can., p. 320, Hud. Riv. Gr.

magnisulcatum, Billings, 1857, Rep. of
Progr. Geo. Sur. Can., p. 330, Hud.
Riv. Gr.

marcellense, Vanuxem, 1842, Geo. Rep.
N. Y., p. 147, and Pal. N. Y., vol. 5, pt.
2, p. 278, Hum. Gr.

marginale, Owen, 1840, Rep. on Min.
Lands, p. 70, Up. Magnesian Gr.

maro, Billings, 1859, Can. Nat. and Geol.,
vol. 4, p. 461, Chazy Gr.

marshallense, Winchell, 1862, Am. Jour.
Sci., 2d series, vol. 33, p. 356, Mar-
shall Gr.

masculum, Hall, 1879, Pal. N. Y., vol. 5,
p. 238, Schoharie grit.

medium, Hall, 1879, Pal. N. Y., vol. 5, p.
254, Schoharie grit.

medon, Billings, 1866, Catal. Sil. Foss.
Antic., p. 57, Clinton Gr.

medullare, Hall, 1860, Rep. of Progr. Geo.
Sur. Wis., p. 4, Niagara Gr.

meeki, S. A. Mil-
ler, 1875, Cin.
Quar. Jour.
Sci., vol. 2, p.
126, Hud. Riv.
Gr.

menelaus, Bill-
ings, 1862, Pal.
Foss., vol. 1, p.
26, Black Riv.
Gr.

mephisto, Clarke, 1885, Bull. U. S. Geo.
Sur., No. 16, p. 29, Genesee Shale.

michiganense, S. A. Miller, 1883, 2d Ed.
Am. Pal. Foss., p. 303, Marshall Gr. in
the southern part of Michigan. Pro-
posed instead of O. multicinctum,

Winchell, Proc. Acad. Nat. Sci., Phil.,
Sept., 1862, p. 421.

minganense, Billings, 1857, Rep. of Progr.
Geo. Sur. Can., p. 319, Chazy and Black
Riv. Grs.

missisquoi, see Cyrtoceras missisquoi.

FIG. 753.—Orthoceras meeki.

mephisto, Clarke, 1885, Bull. U. S. Geo.
Sur., No. 16, p. 29, Genesee Shale.

michiganense, S. A. Miller, 1883, 2d Ed.
Am. Pal. Foss., p. 303, Marshall Gr. in
the southern part of Michigan. Pro-
posed instead of O. multicinctum,

Winchell, Proc. Acad. Nat. Sci., Phil.,
Sept., 1862, p. 421.

minganense, Billings, 1857, Rep. of Progr.
Geo. Sur. Can., p. 319, Chazy and Black
Riv. Grs.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

missisquoi, see Cyrtoceras missisquoi.

FIG. 574.—Orthoceras mohri.

mobri, S. A. Miller, 1875, Cin. Quar. Jour.
Sci., vol. 2, p. 124, Hud. Riv. Gr.

- molestum*, Hall, 1876, *Illust. Devonian Foss.*, pl. 35, and *Pal. N. Y.*, vol. 5, pt. 2, p. 265, Up. Held. Gr.
moniliforme, Hall, 1847, *Pal. N. Y.*, vol. 1, p. 35, Chazy Gr.
moniliforme, Swallow, 1858, *Trans. St. Louis Acad. Sci.*, vol. 1. The name was preoccupied. See *O. swallovanum*.
montrealense, Billings, 1859, *Can. Nat. and Geo.*, vol. 4, p. 383, Calciferous Gr.
multicameratum, Emmons, 1842, *Geo. Rep. N. Y.*, p. 382, and *Pal. N. Y.*, vol. 1, p. 45, Birdseye Gr.
multicinctum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 76, and *Pal. N. Y.*, vol. 5, pt. 2, p. 263, Schoharie grit.
multicinctum, Winchell, 1862. The name was preoccupied. See *O. michiganense*.
multilineatum, Emmons, 1842, *Geo. Rep. N. Y.*, p. 397, Trenton Gr.
multiseptum, Hall, 1852, *Pal. N. Y.*, vol. 2, p. 14, Medina Gr.
murrayi, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 332, Black Riv. and Trenton Grs.
niagarensis, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 416, Niagara Gr.
nobile, Meek & Worthen, 1865, *Proc. Acad. Nat. Sci.*, p. 256, Kaskaskia Gr.
notocostum, McClesney, 1861, *New Pal. Foss.*, p. 94, Niagara Gr.
novamexicanum, Marcou, 1858, *Geol. North America*, p. 44, Subcarboniferous.
nummularium, (?) 1839, Murch. Sil. Sys., p. 632, Up. Sil.
nuntium, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 79, and *Pal. N. Y.*, vol. 5, pt. 2, p. 269, Ham. Gr.
oberon, Billings, 1866, *Catal. Sil. Foss. Antic.*, p. 82, Niagara Gr.
occidentale, Swallow, 1858, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 201, Coal Meas., Permian Gr.
occidentale, Winchell, 1862. This name was preoccupied. See *O. vinchellanum*.
oedipus, Hall, 1879, *Pal. N. Y.*, vol. 5, p. 294, Ham. Gr.
okawense, Worthen, 1883, *Geo. Sur. Ill.*, vol. 7, p. 324, and *Geo. Sur. Ill.*, vol. 8, p. 149, Kaskaskia Gr.
olorus, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 245, Trenton Gr. Proposed instead of *O. vertebrale*, Hall, 1847, which was preoccupied.
ommaneyi, Salter, 1852, in Sutherland's Jour., vol. 2, Devonian.
oneidense, Walcott, 1879, *Trans. Alb. Inst.*, vol. x, p. 22, Utica Slate Gr.
ontario, Clarke, 1885, *Bull. U. S. Geo. Sur.*, No. 16, p. 51, Chemung Gr.
oppletum, Hall, 1879, *Pal. N. Y.*, vol. 5, p. 248, Schoharie grit.
ordinatum, Billings, 1865, *Pal. Foss.*, vol. 1, p. 350, Calciferous Gr.
ortoni, Meek, 1872, *Proc. Acad. Nat. Sci. Phil.*, p. 330, and *Ohio Pal.*, vol. 1, p. 155, Hud. Riv. Gr.
orus, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 245, Niagara Gr. Proposed instead of *O. columnare*, Hall, 1860, which was preoccupied.
ottawense, Billings, 1857, Rep. of Progr. Geo. Fur. Can., p. 331, Black Riv. and Trenton Grs.
ozarkense, Shumard, 1863, *Trans. St. Louis Acad. Sci.*, vol. 2, p. 107, Calciferous Gr.
pacator, Hall, 1879, *Pal. N. Y.*, vol. 5, p. 307, Portage Gr.
palmatum, Hall, 1879, *Pal. N. Y.*, vol. 5, p. 312, Chemung Gr.
pauciseptum, Hall, 1859, *Pal. N. Y.*, vol. 3, p. 346, Low. Held. Gr.
pelops, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 73, and *Pal. N. Y.*, vol. 5, pt. 2, p. 233, Schoharie grit.
pelops var. *ohioense*, Hall, 1876, *Illust. Devonian Foss.*, pl. 36, and *Pal. N. Y.*, vol. 5, pt. 2, p. 236, Up. Held. Gr.
perannulatum, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 319. This name was preoccupied by Portlock in 1843. See *O. crocus*.
perelegans, Salter, 1848, *Mem. Geo. Sur. Gr. Brit.*, vol. 2, p. 354, Ham. Gr.
perparvum, Billings, 1862, *Pal. Foss.*, vol. 1, p. 27, Black Riv. Gr.
persens, Billings, 1865, *Pal. Foss.*, vol. 1, p. 313, Quebec Gr.
persiphonatum, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 329, Mid. Sil. If the genus *Huron* is valid, this species will belong to it.
perstriatum, Hall, 1859, *Pal. N. Y.*, vol. 3, p. 346, Low. Held. Gr.
perstrictum, Dawson, 1865. The name was preoccupied by Barrande. See *O. dawsonianum*.
pertextum, Hall, 1879, *Pal. N. Y.*, vol. 5, p. 314, Chemung Gr.
pertinax, Billings, 1860, *Can. Nat. and Geo.*, vol. 5, p. 75, Black Riv. Gr.
pervicax, Hall, 1879, *Pal. N. Y.*, vol. 5, p. 257, Schoharie grit.
pileolum, Billings, 1866, *Catal. Sil. Foss. Antic.*, p. 58, Medina Gr.
piscator, Billings, 1865, *Pal. Foss.*, vol. 1, p. 251, Quebec Gr.
piso, Billings, 1862, *Pal. Foss.*, vol. 1, p. 168, Hud. Riv. Gr.
planoconvexum, Hall, 1861, Rep. of Progr. Wis., p. 47, and *Geo. Wis.*, vol. 4, p. 228, Black Riv. and Trenton Grs.
pravum, Hall, 1879, *Pal. N. Y.*, vol. 5, p. 255, Schoharie grit.
pressum, Rogers, 1868, Bigsby, *Thesaurus Siluricus*, p. 180. Not defined.
prismus, Billings, 1865, *Pal. Foss.*, vol. 1, p. 253, Quebec Gr.
primigenium, Vanuxem, 1842, *Geo. Rep. N. Y.*, p. 36, and *Pal. N. Y.*, vol. 1, p. 13, Calciferous Gr.
procerum, Hall, 1876, *Illust. Devonian Foss.*, pl. 35, and *Pal. N. Y.*, vol. 5, pt. 2, p. 249, Schoharie grit.

Am. Pal. Foss., p.
posed instead of
1860, which was

, Rep. of Progr.
, Black Riv. and

1803, Trans. St.
2, p. 107, Calcif.

N. Y., vol. 5, p.

Pal. N. Y., vol. 5,

Pal. N. Y., vol.

Rep. N. Y. Mus.

Pal. N. Y., vol. 5,

grit.

all, 1876, Illust.

, and Pal. N. Y.,

Up. Held. Gr.

1857, Rep. of

, p. 319, This

l by Portlock in

Mem. Geo. Sur.

, Ham. Gr.

2, Pal. Foss., vol.

Pal. Foss., vol. 1,

, 1857, Rep. of

, p. 329, Mid. Sil.

ia is valid, this

t.

Pal. N. Y., vol.

Gr.

The name was

de. See O. daw-

al. N. Y., vol. 5,

Can. Nat. and

ck Riv. Gr.

N. Y., vol. 5, p.

Catal. Sil. Foss.

gr.

Pal. Foss., vol. 1.

Foss., vol. 1, p.

61, Rep. of Progr.

Wis., vol. 4, p.

profundum, Hall, 1862, 15th Rep. N. Y.

Mus. Nat. Hist., p. 76, Up. Held. Gr.

propinquum, Billings, 1857, Rep. of Progr.

(Geo. Sur. Can.), p. 320. Preoccupied.

See O. fulgur.

punctostratum, Hall, 1860, Can. Nat. and

(Geo., vol. 5, p. 154, Up. Sil.

pastulosum, Winchell, 1866, Rep. Low.

Peninsula Mich., p. 97, Ham. Gr.

pylades, Billings, 1866, Catal. Sil. Foss.

Antic., p. 84, Niagara Gr.

pythion, Billings, 1857, Rep. of Progr. Geo.

Sur. Can., p. 335, Trenton Gr.

rundolphense, Worthen, 1882, Bull. No. 1,

Ill. St. Mus. Nat. Hist., p. 38, Kaskaskia

Gr. Proposed instead of O. annulato-

costatum, Meek & Worthen, which was

preoccupied.

rapax, see Endoceras rapax.

raptor, Billings, 1866, Catal. Sil. Foss.

Antic., p. 57, Medina Gr.

recedens, Barriande, 1869, Sys. Sil. de

Boh., 4me ser., p. viii, pl. 433, Quebec Gr.

rectianulatum, Hall, 1847, Pal. N. Y.,

vol. 1, p. 34, Chazy and Birdseye Grs.

recticameratum, Hall, 1847, Pal. N. Y.,

vol. 1, p. 46, Birdseye Gr.

rectum, Worthen, 1875, Geo. Sur. Ill.,

vol. 6, p. 504, Niagara Gr.

remus, Billings, 1866, Catal. Sil. Foss.

Antic., p. 85, Niagara Gr.

repens, Billings, 1865, Pal. Foss., vol. 1,

p. 312, Quebec Gr.

reticulatum, Phillips, 1836, Geol. York.

Chemung Gr. Not clearly identified

in this country.

rigidum, Hall, 1859, Pal. N. Y., vol. 3, p.

344, Low. Held. Gr.

robustum, Winchell, 1862, Am. Jour.

Sci., 2d ser., vol. 33, p. 356, Mar-

shall Gr.

robustum, Hall, 1876. The name was pre-

occupied. See O. erienne.

rotulatum, Billings, 1857, Rep. of Progr.

Geo. Sur. Can., p. 334, Niagara Gr.

rude, Hall, 1859, Pal. N. Y., vol. 3, p. 346,

Low. Held. Gr.

rudens, Beecher, 1888, Pal. N. Y., vol. 7,

p. 28, Ham. Gr.

rudicula, Hall, 1876, Illust. Devonian

Foss., pl. 37, and Pal. N. Y., vol. 5, pt.

2, p. 268, Up. Held. Gr.

rushiense, McChesney, 1860, New. Pal.

Foss., p. 68, and Geo. Sur. Ill., vol. 5,

p. 612, Coal Meas.

sayi, Billings, 1865, Pal. Foss., vol. 1, p.

315, Quebec Gr.

scammoni, McChesney, 1861, New Pal.

Foss., p. 92, Niagara Gr.

sceptrum, Beecher, 1888, Pal. N. Y., vol.

7, p. 26, Up. Held. Gr.

scintilla, Hall, 1879, Pal. N. Y., vol. 5, p.

293, Ham. Gr.

sedgwicki, Billings, 1857, Rep. of Progr.

Geo. Sur. Can., p. 320, Hud. Riv. Gr.

selwyni, Billings, 1862, Pal. Foss., vol. 1,

p. 161, Guelph Gr.

servile, Billings, 1865, Pal. Foss., vol. 1, p.

252, Quebec Gr.

shumardi, Billings, 1859, Can. Nat. and

(Geo., vol. 4, p. 460, Chazy Gr.

steinus, Hall, 1879, Pal. N. Y., vol. 5, p.

301, Marcellus Shale.

sieboldi, Billings, 1866, Catal. Sil. Foss.

Antic., p. 23, Hud. Riv. and Anti-

costi Grs.

simpsoni, Billings, 1859, Rep. of Progr.

Assiniboine and Saskatchewan Ex.

Exp., p. 186, Silurian.

simulator, Hall, 1876, 28th Rep. N. Y.

Mus. Nat. Hist., p. 179, Niagara Gr.

sirpus, Hall, 1879, Pal. N. Y., vol. 5, p.

269, Up. Held. Gr.

sociale, Hall, 1877, 1st Ed. Am. Pal.

Foss., p. 245, Hud. Riv. Gr. Proposed

instead of O. gregarium, Hall, 1861,

which was preoccupied.

sordidum, Billings, 1859, Can. Nat. and

(Geo., vol. 4, p. 363, Calciferous Gr.

spissiscriptum, Dwight, 1884, Am. Jour.

Sci. and Arts, 3d. ser., vol. 27, p. 256,

Calciferous Gr.

spissum, Hall, 1879, Pal. N. Y., vol. 5, p.

287, Ham. Gr.

stebos, Clarke, 1885, Bull. U. S. Geo.

Sur., No. 16, p. 29, Genesee Shale.

striatum, (?) Sowerby, 1812, Min. Conch.,

vol. 1, p. 129, Devonian.

striatilineatum, McChesney, 1861, New

Pal. Foss., p. 94, Niagara Gr.

strigatum, Hall, 1847, Pal. N. Y., vol. 1,

p. 205, Trenton Gr.

strix, Hall & Whitfield, 1875, Ohio Pal.,

vol. 2, p. 149, Niagara Gr.

stylus, Hall, 1877, 1st Ed. Am. Pal. Foss.,

p. 245, and Pal. N. Y., vol. 5, pt. 2, p.

253, Schoharie grit. Proposed instead

of O. baculum, Hall, 1862, which was

preoccupied.

subarcuatum, Hall, 1847, Pal. N. Y., vol.

1, p. 34. Preoccupied by Portlock in 1843.

See O. clintoni.

subbaculum, Meek & Worthen, 1865, Proc.

Acad. Nat. Sci., p. 256, Niagara Gr.

subcancellatum, Hall, 1877, 1st Ed. Am.

Pal. Foss., p. 245, Niagara Gr. Pro-

posed instead of O. cancellatum, Hall,

1852, which was preoccupied.

sublaeve, D'Orbigny, 1850, Prodrome de

Pal., t. 1, p. 28, Onondaga Gr. Pro-

posed instead of O. laeve, Hall, 1843,

which was preoccupied.

subtextile, Hall, 1859, Pal. N. Y., vol. 3,

p. 344, Low. Held. Gr.

subulatum, Hall, 1843, 4th Dist. (Geo.

Rep. N. Y., p. 180, and Pal. N. Y., vol.

5, pt. 2, p. 283, Marcellus Shale.

swallowanum, S. A. Miller, 1883, 2d Ed.

Am. Pal. Foss., p. 308, Coal Measures

in the Valley of Verdigris in Kansas.

Proposed instead of O. moniliforme,

Swallow, in Trans. St. Louis Acad. Sci.,

vol. 1, p. 200, which was preoccupied

by Hall.

tantalus, Hall, 1879, Pal. N. Y., vol. 5, p.

241, Schoharie grit.

telamon, Hall, 1879, Pal. N. Y., vol. 5, p.

291, Ham. Gr.

tener, Hall, 1879, Pal. N. Y., vol. 5, p. 285, Ham. Gr.
 tenerum, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 174, Black Riv. Gr.
 tenui-annulatum, Hall, 1859, Pal. N. Y., vol. 3, p. 345, Low. Held. Gr.
 tenuiseptum, Hall, 1847, Pal. N. Y., vol. 1, p. 35, Chazy Gr.
 teretiforme, Hall, 1847, Pal. N. Y., vol. 1, p. 198, Trenton Gr.
 tersum, Hall, 1879, Pal. N. Y., vol. 5, p. 286, Ham. Gr.
 tetricum, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 45, and Pal. N. Y., vol. 5, pt. 2, p. 251, Schoharie grit.
 textile, Hall, 1847, Pal. N. Y., vol. 1, p. 199, Trenton Gr.
 textum, Hall, 1879, Pal. N. Y., vol. 5, p. 285, Ham. Gr.
 thestor, Hall, 1879, Pal. N. Y., vol. 5, p. 302, Marcellus shales.
 thoms, Hall, 1862, 15th Rep. N. Y. Mus. Nat. Hist., p. 75, and Pal. N. Y., vol. 5, pt. 2, p. 61, Schoharie grit.
 thyestes, Hall, 1879, Pal. N. Y., vol. 5, p. 306, Portage Gr.
 tityrus, Billings, 1865, Pal. Foss., vol. 1, p. 316, Quebec Gr.
 transversum, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 129, Hud. Riv. Gr.

FIG. 755.—*Orthoceras transversum*.

trentonense, see *Cyrtoceras trentonense*.
 turbidum, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 100, Hud. Riv. Gr.
 typus, Saemann, as identified by Hall, 1876, Illust. Devonian Foss., is *O. marcellense*.
 undulatum, Owen, 1840, Rep. on Min. Lands, Niagara Gr. The name was preoccupied by Sowerby in 1812. See *O. iowense*.
 undulostriatum, Hall, 1847, Pal. N. Y., vol. 1, p. 202, Trenton Gr.
 unionense, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 505, Niagara Gr.
 varro, Billings, 1866, Catal. Sil. Foss. Antic., p. 84, Niagara Gr.
 varum, Hall, 1879, Pal. N. Y., vol. 5, p. 259, Schoharie grit.
 vastator, Hall, 1879, Pal. N. Y., vol. 5, p. 243, Schoharie grit. Correct in the index, but printed *O. obliquum*, on page 243.
 velox, Billings, 1865, Pal. Foss., vol. 1, p. 173, Chazy Gr.
 vertebrate, Hall, 1847, Pal. N. Y., vol. 1, p. 201. Preoccupied by Schlotheim in 1820, and by Eichwald in 1840. See *O. olorus*.
 veterator, Billings, 1865, Pal. Foss., vol. 1, p. 350, Calciferous Gr.
 viator, Hall, 1879, Pal. N. Y., vol. 5, p. 270, Up. Held. Gr.

vinchellanum, S. A. Miller, 1883, 2d Ed. Am. Pal. Foss., p. 308, Marshall Gr. in Southern Michigan. Proposed instead of *O. occidentale*, Winchell, 1862, Am. Jour. Sci. and Arts, 2d ser., vol. 33, p. 356, which was preoccupied by Swallow.
 vindobonense, Dawson, 1868, Acad. Geol., p. 311, Carboniferous.
 virgatum, Sowerby, 1839, Murch. Sil. Sys., p. 632, and Pal. N. Y., vol. 2, p. 291, Niagara Gr.
 virgulatum, Hall, 1852, Pal. N. Y., vol. 2, p. 90, Clinton and Niagara Grs.
 vittatum, Sandberger. Not American.
 vulgatum, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 337, Trenton Gr.
 warrenense n. sp., Chemung Gr. Proposed instead of *O. cochleatum*, Hall, Pal. N. Y., vol. 5, p. 308, pl. cxlii, fig. 10, which name was preoccupied.
 wauwatosense, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 66, and Geo. Wis., vol. 4, p. 297, Niagara Gr.
 whitii, Winchell, 1863, Proc. Acad. Nat. Sci., p. 22, Kinderhook Gr.
 winchelli, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 257, and Geo. Sur. Ill., vol. 6, p. 512, Ham. Gr.
 woodworthi, McChesney, 1865, New Pal. Foss., p. 53, Niagara Gr. Proposed instead of *O. irregulare*, which was preoccupied.
 xerxes, Billings, 1865, Pal. Foss., vol. 1, p. 316, Quebec Gr.
 xiphias, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 318, Trenton Gr.
 zeus, Hall, 1879, Pal. N. Y., vol. 5, pt. 2, p. 235, Schoharie grit.
 PETALICINUS, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 221. [Ety. *petalos*, spread out; *ichnos*, track.] A wide trail composed of numerous transversely elongated depressions arranged without order. Type *P. multipartitus*.
 multipartitus, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 222, Utica Slate Gr.
 PHRAGMOCERAS, Broderip, 1839, Murch. Sil. Syst., p. 621. [Ety. *phragmos*, partition; *keras*, horn.] Shell short, arched, compressed, more or less conical; sides of the mouth lapping toward each other; septa simple, crossed by the sigmoidal lines of growth; siphuncle on the internal edge, dilate between the septa. Type *P. arcuatum*.
 byronense, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 506, Niagara Gr.

FIG. 756.—*Phragmoceras then*. 1875, hector. Cast of interior. Geo. Sur. Ill., vol. 6, p. 506, Niagara Gr.

ler, 1883, 2d Ed.
Marshall Gr. in
proposed instead
chell, 1862. Am.
l ser., vol. 33, p.
pied by Swallow.
1868, Acad. Geol.,

Murch. Sil. Sys.,
., vol. 2, p. 291,

Pal. N. Y., vol. 2,
gara Grs.

of American.

, R. p. of Progr.

Trenton Gr.

lung Gr. Pro-

ochleatum, Hall,

08, pl. cxiii, fig.

reoccupied.

, 1880, Ann. Rep.

, and Geo. Wis.,

Gr.

Proc. Acad. Nat.

Gr.

chen, 1866, Proc.

p. 257, and Geo.

Ham. Gr.

, 1865, New Pal.

Gr. Proposed in

which was preoc-

l. Foss., vol. 1, p.

Rep. of Progr.

Trenton Gr.

Y., vol. 5, pt. 2, p.

1880, Jour. Cin.

p. 221. [Ety. *pe-*

track.] A wide

numerous trans-

sections arranged

P. multipartitus.

ler, 1880, Jour.

l. 2, p. 222, Utica

1839, Murch. Sil.

ragmos, partition;

ort, arched, com-

ellipticum, Hall & Whitfield, 1875, Ohio
Pal., vol. 2, p. 152, Niagara Gr.
expansum, Winchell, 1863, Proc. Acad.
Nat. Sci., p. 23, Kinderhook Gr.
hector, Billings, 1862, Pal. Foss., vol. 1, p.
163, Guelph Gr.
hoi, Whitfield, 1878, Ann. Rep. Geo. Sur.
Wis., p. 86, and Geo. Wis., vol. 4, p. 300,
Niagara Gr.

FIG. 757.—*Phragmoceras hector*. Side view.

hoi var. compressum, Whitfield, 1878,
Ann. Rep. Geo. Sur. Wis., p. 82, and
Geo. Wis., vol. 4, p. 301, Niagara Gr.
labiatum, Whitfield, 1878, Ann. Rep. Geo.
Sur. Wis., p. 86, and Geo. Wis., vol. 4,
p. 302, Niagara Gr.
nestor, Hall, 1867, 20th Rep. N. Y. Mus.
Nat. Hist., p. 405, Niagara Gr.
nestor var. canadense, Whiteaves, 1884,
Pal. Foss., vol. 3, p. 39, Guelph Gr.
parvum, Hall & Whitfield, 1875, Ohio
Pal., vol. 2, p. 151, Niagara Gr.
prematurum, Billings, 1866, Can. Nat.
and Geo., vol. 5, p. 173, Black Riv. and
Trenton Grs. Type of Hyatt's genus
Melinoceras.

FIG. 758.—*Pilo-*
ceras wortheni.

amplum, Dawson, 1881, Can. Nat., vol. 10,
p. 1, Calciferous Gr.

canadense, Billings, 1860, Can. Nat. and
Geol., vol. 5, p. 171, Calciferous Gr.
explanator, Whitfield, 1880, Bull. Am.
Mus. Nat. Hist., vol. 1, p. 323, Birds-
eye Gr.

gracile, Billings, 1865, Pal. Foss., vol. 1,
p. 257, Quebec Gr.

triton, Billings, 1865, Pal. Foss., vol. 1, p.
257, Quebec Gr.

wortheni, Billings, 1865, Pal. Foss., vol. 1,
p. 256, Quebec Gr.

Polycronites haani, Troost, 1840, 5th Geo.
Rep. Tenn., Devonian. Not clearly
defined, but probably a *Gyroceras*.

PTERONAUTILUS, Meek, 1864, Pal. of Up. Mo.,
p. 64. [Ety. *pteron*, wing; *nautilus*, a
genus.] Shell with the involute body
portion comparatively very small, and
globular in form, scarcely umbilicate;
outer chamber very large, and deflected
from the involute body, its inner or
ventral side being widely open, and the
lateral margins greatly dilated, so as to
form a very large, wing-like expansion
on each side. Type *P. seebachianus*.

seebachianus, Geinitz, (*Nautilus seebach-*
ianus.) Carb. und Dyas, p. 43, Per-
mian Gr.

SERICHNITES, Billings, 1866, Catal. Sil. Foss.
Antic., p. 73. The author supposed the
tracks might have been made by a spe-

cies of *Cephalopoda*. They consist of
two parallel rows of semicircular or
subquadrate pits; each pit is about one-
half inch in diameter, and separated
from the succeeding one by about one-
fourth of an inch. Type *S. abruptus*.
abruptus, Billings, 1866, Catal. Sil. Foss.
Antic., p. 73, Hud. Riv. Gr.

Sidemina infundibuliforme, Castelnau, 1843,
Syst. Sil., p. 33. Probably the fragment
of an *Endoceras*.

SOLENOCHILUS, Meek & Worthen, 1870, Proc.
Acad. Nat. Sci. Phil., vol. 20, p. 47.

[Ety. *solen*, a channel; *cheilos*, a lip.]
Nautiloid in form, with small siphuncle
in contact, or nearly in contact, with
the outer shell; margins of the lip near
the umbilicus, terminating in spout-
like auricles. Type *S. collectum*.

FIG. 759.—*Solenochilus avonense*.

avonense, Dawson, 1868, (*Nautilus avon-*
ensis.) Acad. Geol., p. 311, Carbonif-
erous.

- capax, Meek & Worthen, 1865, (Cryptoceras capax.) Proc. Acad. Nat. Sci. Phil., p. 262, and Geo. Sur. Ill., vol. 6, p. 532, Coal Meas.
- collectum, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 48, and Geo. Sur. Ill., vol. 5, p. 544, St. Louis Gr.
- indianense, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 150, St. Louis Gr.
- leidy, Meek & Worthen, 1865, (Nautilus leidy,) Proc. Acad. Nat. Sci. Phil., p. 262, and Geo. Sur. Ill., vol. 5, p. 524, Keokuk Gr.
- springeri, White & St. John, 1868, (Nautilus springeri,) Trans. Chi. Acad. Sci., vol. 1, p. 124, Up. Coal Meas.
- Spirula*, Lamarck, 1801, Syst. An. sans Vert.
- mortoni*, Troost, 1840, 5th Geo. Rep. Tenn., Niagara Gr. Not clearly defined.
- STREPTOCERAS**, Billings, 1866, Catal. Sil. Foss. Antic., p. 88. [Ety. *streptos*, twisted; *keras*, horn.] Having the general form of *Oncoceras*, but with a trilobed aperture resembling *Phragmoceras*. Type *S. janus*.
- heros, Billings, 1866, Catal. Sil. Foss. Antic., p. 89, Niagara Gr.
- janus, Billings, 1866, Catal. Sil. Foss. Antic., p. 88, Niagara Gr.

FIG. 760.—Aperture of *Streptoceras janus*.

- TEMNOCHILUS**, McCoy, 1844, Synop. Carb. Foss. Ireland, p. 20. [Ety. *temno*, I divide; *cheilos*, lip.] Nautiloid in form, and characterized by a broad, deep, open umbilicus, showing all the volutions, with the outer side of the volutions broad or flattened, and the middle of each lateral margin prominently angular; the angle being sometimes nodose, while the transverse diameter of the volutions is always greater than the dorso-ventral; siphuncle between the middle and the outer side of the whorls. Type *T. biangulatus*.
- coxanum, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 50, and Geo. Sur. Ill., vol. 5, p. 543, St. Louis Gr.
- latum, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 49, and Geo. Sur. Ill., vol. 5, p. 608, Coal Meas.

- niotense, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 200, and Geo. Sur. Ill., vol. 5, p. 523, Keokuk Gr.
- Type of Hyatt's genus *Edaphoceras*.

FIG. 761.—*Temnochilus coxanum*.

- peramplum, Meek & Worthen, 1865, (Endolobus peramplum,) Proc. Acad. Nat. Sci. Phil., p. 259, Kaskaskia Gr.
- scottense, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 151, Warsaw Gr.
- spectabile, Meek & Worthen, 1860, (Nautilus spectabilis,) Proc. Acad. Nat. Sci. Phil., p. 469, and Geo. Sur. Ill., vol. 2, p. 308, Kaskaskia Gr.
- winslowi, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 50, and Geo. Sur. Ill., vol. 5, p. 609, Coal Meas.
- TERATICHNUS**, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 221. [Ety. *teras*, a wonder; *ichnos*, track.] A track supposed to have been made by a cephalopod, and consisting of numerous elongated, more or less bifurcated impressions. Type *T. confertus*.
- confertus, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 221, Utica Slate Gr.
- TRACHONATICHNUS**, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 219. [Ety. *trachoma*, that which is made rough; *ichnos*, track.] A track supposed to have been made by a cephalopod and consisting of numerous simple or compound impressions arranged in two series. Type *T. numerosus*.
- cincinnatiensis, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 220, Utica Slate Gr.
- numerosus, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 219, Utica Slate Gr.
- permultus, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 220, Utica Slate Gr.
- TREMATOCERAS**, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 205. [Ety. *trema*,

hen, 1865, Proc.
s., 200, and Geo.
23, Keokuk Gr.
Edaphoceras.

coxanum.

Worthen, 1865,
s.), Proc. Acad.
259, Kaskas-

press.), Geo. Sur.
Law Gr.

hen, 1860, (Nau-
Acad. Nat. Sci.
Sur. Ill., vol. 2.

hen, 1870, Proc.
p. 50, and Geo.
Coal Meas.

1880, Jour. Cin.
Sci., p. 221. [Ety.
track.] A track
made by a
sting of numer-
ous bifurcated
confertus.

1880, Jour. Cin.
Sci., p. 221, Utica

ller, 1880, Jour.
vol. 2, p. 219,
which is made

A track sup-
side by a cephal-
numerous sim-
pansions arranged
numerous.

ller, 1880, Jour.
vol. 2, p. 220,

r, 1880, Jour.
vol. 2, p. 219,

1880, Jour. Cin.
Sci., p. 220, Utica

1882, Ann. N. Y.
Sci., p. 220, Utica

hole; *keras*, horn.] Shell straight, ob-
conical, like *Orthoceras* as to tube,
septa, and siphuncle; characterized by
a line of elongated, raised tubercles
along one side of the shell, which at one
stage of growth formed perforations,
which were closed as the animal ex-
tended the shell. Type *T. ohioense*.

ohioense, Whitfield, 1882, Ann. N. Y.
Acad. Sci., vol. 2, p. 206, Up. Held. Gr.
Trematodiscus, Meek & Worthen, 1861,
Proc. Acad. Nat. Sci. Phil., p. 147. [Ety.
trema, hole; *diskos*, quoit.] Discoid,
wide, shallow, umbilicus, perforated in
the middle, showing all the whorls;
whorls slender, merely in contact pos-
sessed of revolving angles, grooves, or
striae; siphuncle central or subcentral
on the dorsal side. Type *T. stygialis*.
The name having been used in 1860 by
Haeckel for *Radiolaria*, Hyatt proposed
Trematoceras.

altidorsalis, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 429, Marshall Gr.

digonus, Meek & Worthen, 1860, (Nau-
tilus *digonus*.) Proc. Acad. Nat. Sci., p.
470, and Geo. Sur. Ill., vol. 2, p. 163,
Kinderhook Gr.

discoidalis, Winchell, 1862, Am. Jour.
Sci., vol. 33, 2d series, p. 360, Mar-
shall Gr.

konincki, Wetherby, 1881, Jour. Cin. Soc.
Nat. Hist., vol. 4, p. 79, Waverly Gr.

FIG. 762.—*Trematodiscus konincki*.

meekanus, Winchell, 1862, Am. Jour.
Sci., 2d series, vol. 33, p. 360, Mar-
shall Gr.

planidorsalis, Winchell, 1862, Am. Jour.
Sci., 2d series, vol. 33, p. 358, Mar-
shall Gr.

rockymontanus, S. A. Miller, 1881, Jour.
Cin. Soc. Nat. Hist., vol. 4, p. 312, Bur-
lington Gr.

striatulus, Winchell, 1862, Proc. Acad.
Nat. Sci., 2d series, vol. 33, p. 358, Mar-
shall Gr.

strigatus, Winchell, 1862, Proc. Acad. Nat.
Sci., p. 426, Marshall Gr.

sulcatus, Meek & Worthen, 1866, Proc.
Acad. Nat. Sci. Phil., p. 274, Kaskas-
kia Gr.

trigonus, Winchell, 1862, Am. Jour. Sci.,
2d series, vol. 33, p. 358, Marshall Gr.

trisulcatus, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 470, and Geo.
Sur. Ill., vol. 2, p. 162, Kinderhook Gr.

Trochoceras, Hall, 1852, Pal. N. Y., vol. 2,
p. 335. [Ety. *trochos*, hoop; *keras*, horn.]
This name was proposed by Barrande
at about the same time. Turbinate or
trochiform, spire elevated, more or less
ventricose; umbilicated; aperture
rounded or round oval; volutions above
the outer one septate; siphuncle sub-
marginal or dorsal. Type *T. gebhardi*.
uneas, Hall, 1870, Rev. Ed. 20th Rep.
N. Y. Mus. Nat. Hist. Expl., pl. 25,
Niagara Gr.

baeri, see *Gyroceras baeri*.

barrandii, Hall, 1879, Pal. N. Y., vol. 5,
p. 398, Schoharie grit.

biton, Hall, 1879, Pal. N. Y., vol. 5, p. 395,
Schoharie grit.

clio, Hall, 1861, 14th Rep. N. Y. Mus.
Nat. Hist., p. 108, Schoharie grit. Type
of Hyatt's genus *Sphyraceras*.

costatum, Hall, 1861, Geo. Rep. of Wis.,
Niagara Gr.

desplainense, McChesney, 1860, New Pal.
Foss., p. 68, Niagara Gr.

discoldeum, Hall, 1862, 15th Rep. N. Y.
Mus. Nat. Hist., p. 64, and Illust. De-
von. Foss., pl. 59, Schoharie grit.

eugenium, Hall, 1861, 14th Rep. N. Y.
Mus. Nat. Hist., p. 108, Schoharie grit.
Type of Hyatt's genus *Nedoceras*.

expansum, Hall, 1879, Pal. N. Y., vol. 5,
p. 402, Schoharie grit.

gebhardi, Hall, 1852, Pal. N. Y., vol. 2,
p. 335, Coralline Gr.

incipiens, Barrande, 1869, Syst. Sil. de
Boh., 4me ser., Quebec Gr.

notum, Hall, 1867, 20th Rep. N. Y. Mus.
Nat. Hist., p. 403, Niagara Gr.

obliquatum, Hall, 1876, Illust. Devonian
Foss., pl. 48, Up. Held. Gr.

orion, Hall, 1876, (Cyrtoceras *orion*.) Il-
lust. Devonian Foss., pl. 48, Up. Held. Gr.

pandion, Hall,
1876, Illust. De-
vonian Foss.,
pl. 48, and Pal.
N. Y., vol. 5, pt.
2, p. 400, Scho-
harie grit.

pandion, Hall,
1879, Pal. N. Y.,
vol. 5, p. 403,
Schoharie grit.

turbinatum, Hall,
1852, Pal. N. Y.,
vol. 2, p. 336,
Coralline Gr.

waldronense,
Hall, 1876, 28th
Rep. N. Y. Mus. Nat. Hist., p. 179, Ni-
agara Gr.

FIG. 763.—*Trochoceras waldronense*.

TROCHOLITES, Conrad, 1838, Ann. Geo. Rep. N. Y., p. 119. [Ety. *trochos*, hoop; *lithos*, stone.]

FIG. 764.—Trocholites ammonius.

Discoidal volutions in the same plane, about four, rounded, slightly concave on the ventral side, gradually enlarging toward the aperture; septa direct; outer chamber large; siphuncle ventral. Type *T. ammonius*. ammonius, Conrad, 1838, Ann. Geo. Rep. N. Y., p. 119, and Pal. N. Y., vol. 1, p. 192, Trenton, Utica, and Hud. Riv. Grs. circularis, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 9, Hud. Riv. Gr.

minusculus, Miller & Dyer, 1878, Cont. to Pal. No. 2, p. 9, Utica Slate Gr.

FIG. 765.—Trocholites circularis.

planorbiformis, Conrad, 1842, Jour. Acad. Nat. Sci. Phil., vol. 8, p. 274, and Pal. N. Y., vol. 1, p. 310, Utica and Hud. Riv. Grs.

CLASS LAMELLIBRANCHIATA.

[Ety. *lamella*, a thin plate; *branchiæ*, gills.]

THE Lamellibranchiata, Blainville, or Conchifera, Lamarck, have bivalve shells, abundant in the rivers of North America, in every ocean, and were common in all geological ages, back to early Silurian time. The river shells are known by the common name of mussels, and nearly all belong to three genera, *Unio*, *Anodonta*, and *Margaritana*. All known Palæozoic shells of this class inhabited salt water. The animals have a bilobed mantle, the sides of which secrete a calcareous shell having two valves, which are attached by some kind of a hinge. The hinge frequently has teeth on one valve that fit in cavities on the other. The valves being on each side of the animal, one is a right valve and the other a left valve. In most genera the valves are equal, and the animals lived in an erect position, resting on the edge of the shell opposite to the hinge, and, when moving, plowed a furrow in the sand or mud by the extension of a tongue-like foot. In some genera one valve is much larger than the other, and the shell lies on the larger valve, and adheres to some foreign object, as is the case with the common oyster; in other instances the locomotion is by suddenly opening and closing the valves, which causes the shell to dart through the water, first in one direction and then another, as the *Pecten* does. Some genera have a byssus by which they are attached to submarine bodies. Each valve commences to grow at the apex or beak, which is also called the *umbo*. The umbones are almost always directed toward the ante-

1878, Cont. to
ate Gr.

rcularis.

42, Jour. Acad.
p. 274, and Pal.
tica and Hnd.

A.

have bivalve
were common
are known by
ra, Unio, Ano-
inhabited salt
te a calcareous
ge. The hinge
r. The valves
er a left valve.
erect position,
moving, plowed
oot. In some
on the larger
common oyster;
ng the valves,
n and then an-
ey are attached
or beak, which
ward the ante-

rior side of the shell, and sometimes project as far as the anterior margin. The length of a shell is the distance from the anterior to the posterior side; the width is measured from the hinge or dorsal side to the base; the thickness is measured through the center of the two valves. The surface of the shells is generally marked with ribs, radiating from the umbones, or concentric lines marking the growth of the shell from the umbones. A depression, anterior to the beak, is called a lunule, and when a depression exists posterior to the beak, it is called an escutcheon. Many shells have an external hinge ligament behind the umbones; some have a ligament between the umbones. When the valves are connected internally by a single muscle, the contraction of which brings the valves together, they belong to the Order Monomyaria; if there are two equally developed contracting mussels, they belong to the Dimyaria; or if there are two muscles, one large and functionally active, the other small, they belong to the Heteromyaria. These contracting muscles are called the adductors, and their places of attachment are indicated by scars. The border of the mantle makes an impression, which is called the pallial line, and if there is a sinus in the posterior part of the pallial line, it shows the animal had a retractile siphon, which, in burrowing shells, is often of great length. The Class has also been divided into two Orders, based on the presence or absence of a siphon, to wit: Asiphonida, Asiphonata, or Asiphonophora, and Siphonida, Siphonata, or Siphonophora. Each Order is spelled three different ways by different authors. Shells having a siphon are always gaping at the posterior or anterior side or at both.

It will be observed from the foregoing, the essential characters upon which Palæozoic shells are classified are the following: Equality or inequality of the valves; the presence or absence of an external ligament; the number of muscular scars; the character of the hinge and its dentition; the presence or absence of a pallial sinus; the position of the umbones; the radiate or concentric surface markings; whether the valves fit each other or are gaping at one or both ends; and the presence or absence of a byssal sinus.

ORDER ASIPHONIDA.

Ambonychiidæ, Amphiceliidæ, Anodontopsidæ, Arcidæ, Aviculidæ, Aviculopectenidæ, Cytherodontidæ, Modiomorphidæ, Mytilidæ, Nuculidæ, Nyassidæ, Orthotidæ, Ostreidæ, Palæoconchidæ, Pinnidæ, Prothyridæ, Pteriniidæ, Technophoridæ, Trioniidæ, Unionidæ.

ORDER SIPHONIDA.

Cardiidæ, Cardiomorphidæ, Conocardiidæ, Cyprinidæ, Eopteriidæ, Grammysiidæ, Lucinidæ, Myacidæ, Palæanatinidæ, Pholadellidæ, Sanguinolitidæ, Solenidæ, Spirodomidæ, Tellinidæ.

FAMILY AMBONYCHIIDÆ.—Ambonychia, Angellum, Anomalodonta, Byssopteria.

FAMILY AMPHICELIIDÆ.—Amphicælia.

FAMILY ANODONTOPSIDÆ.—Anodontopsis, Cycloconcha.

FAMILY ARCIDÆ.—Carbonarca, Clinopistha, Macrodon, Megalomus, Ptychodesma.

FAMILY AVICULIDÆ.—Actinodesma, Avicula, Aviculopinna, Bakevella, Ectenodesma, Glyptodesma, Inoceramus, Liopteria, Leptodesma, Limoptera, Monopteria, Monotis, Palæopinna, Posidonomya, Pseudomonotis, Pteronitella, Pteronites.

FAMILY AVICULOPECTENIDÆ.—Aviculopecten, Crenipecten, Euchondria, Lyrripecten, Pernopecten, Pterinopecten, Streblopteria.

FAMILY CARDIIDÆ.—Cardiola, Cardiopsis, Cardium, Dexiobia, Glyptocardia, Lunulicardium, Palæocardia, Panenka, Paracardium, Pararca.

FAMILY CARDIOMORPHIDÆ.—Cardiomorpha, Edmondia, Euthydesma, Protomya.

FAMILY CHÆNOCARDIIDÆ.—Chænocardia.

FAMILY CONOCARDIIDÆ.—Conocardium.

FAMILY CYPRINIDÆ.—Astartella, Cardinia, Clidophorus, Cypricardia, Cypricardites, Matheria, Pleurophorus, Vanuxemia.

FAMILY CYTHERODONTIDÆ.—Cytherodon, Lyrodesma, Schizodus.

FAMILY EOPTERIDÆ.—Eopteria, Euchasma.

FAMILY GRAMMYSIDÆ.—Allorisma, Chænomya, Cuneamya, Grammysia, Lepetodomus, Sedgwickia.

FAMILY LUCINIDÆ.—Paraelyas.

FAMILY MODIOMORPHIDÆ.—Amnigenia, Cypricardella, Elymella, Goniophora, Glossites, Modiomorpha.

FAMILY MYACIDÆ.—Anthracomya.

FAMILY MYTILIDÆ.—Anthracoptera, Gosselettia, Lithophaga, Megambonia, Modiella, Modiolopsis, Myalina, Mytilarca, Mytilops, Plethomytilus, Pyanomya.

FAMILY NUCULIDÆ.—Nucula, Nuculana, Nuculites, Palæoneilo, Pyrenomorus, Solenomya, Tellinomya, Yoldia.

FAMILY NYASSIDÆ.—Nyassa.

FAMILY ORTHONOTIDÆ.—Orthodesma, Orthonota, Orthonotella, Palæosolen, Sphenolium.

FAMILY OSTREIDÆ.—Ostrea, Placunopsis.

FAMILY PALÆANATINIDÆ.—Ilionia, Palæanatina, Prorhynchus.

FAMILY PALÆOCONCHIDÆ.—Palæoconcha.

FAMILY PHOLADELLIDÆ.—Cimitaria, Pholadella, Phthonia.

FAMILY PINNIDÆ.—Pinna.

FAMILY PROTHYRIDÆ.—Prothyris.

FAMILY PTERINIDÆ.—Actinopteria, Pterinea, Ptychopteria, Veicumnia.

FAMILY SANGUINOLITIDÆ.—Cypricardinia, Promacrus, Spathella, Sphenotus, Sanguinolites.

FAMILY SOLENIDÆ.—Solenopsis.

FAMILY SPIRODOMIDÆ.—Spirodomus.

FAMILY TECHNOPHORIDÆ.—Technophorus.

FAMILY TELLINIDÆ.—Tellinopsis.

FAMILY TRIGONIDÆ.—Dolabra, ? Ischyrinia, ?

FAMILY UNIONIDÆ.—Anthracosia, Priscunaia.

Actinodesma subrectans, see *Glyptodesma subrectum*.

ACTINOPTERIA, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, explanation of plate 17, fig. 6. [Ety. *aktin*, a ray; *Pteria*, a genus.] In the text published in 1884, pt. 1, p. 107, he wrote the word *Actinopteria*, which indicates he derived the name from the genus *Pteria*. Distinguished from *Pterinea* by strong cardinal and lateral teeth, and no striations on the ligamental area; right valve convex. The first species mentioned on page 3, where

the genus is defined, is *A. decussata*, but the first one mentioned on page 107 of the text is *A. eximia*. No type is designated.

auriculata, Hall,

1884, Pal. N. Y., vol. 5, pt. 1, p. 121, Chemung Gr.

boydi, Conrad,

1842, (*Avicula boydi*), Jour.

Acad. Nat. Sci.

Phil., vol. 8, p.

237, and Pal.

N. Y., vol. 5, p.

113, Ham. Gr.

decussata, Hall,

1843, (*Avicula*

decussata),

Geo. 4th Dist.

N. Y., p. 203, and Pal. N. Y., vol. 5, p.

111, Ham. Gr.

delta, Hall, 1883, Pal. N. Y., vol. 5, p. 121,

Chemung Gr.

doris, Hall, 1884, Pal. N. Y., vol. 5, p. 109,

Marcellus Shale.

epsilon, Hall, 1883, Pal. N. Y., vol. 5, pt. 1,

p. 122, Chemung Gr.

eta, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p.

124, Chemung Gr.

eximia, Hall, 1883, Pal. N. Y., vol. 5, pt.

1, p. 107, Up. Held. Gr.

iota, Hall, 1884, Pal. N. Y., vol. 5, pt. 1,

p. 127, Chemung Gr.

kappa, Hall, 1884, Pal. N. Y., vol. 5, pt. 1,

p. 128, Chemung Gr.

muricata, Hall, 1843, (*Avicula muricata*),

Geo. Sur. 4th Dist. N. Y., p. 181, and Pal.

N. Y., vol. 5, p. 108, Marcellus Shale.

perobliqua, Conrad, 1842, (*Avicula perobliqua*), Jour.

Acad. Nat. Sci. Phil., vol.

8, p. 235, and Pal. N. Y., vol. 5, p. 116,

Ham. Gr.

perstrialis, Hall, 1883, Pal. N. Y., vol. 5,

pt. 1, p. 118, Chemung Gr.

pleuroptera, Conrad, 1842, (*Avicula pleuroptera*), Jour.

Acad. Nat. Sci., vol. 8,

p. 242, Ham. Gr.

pusilla, Hall, 1884, Pal. N. Y., vol. 5, pt.

1, p. 117, Ham. Gr.

subdecussata, Hall, 1884, Pal. N. Y., vol.

5, pt. 1, p. 110, Ham. Gr.

FIG. 766.—*Actinopteria boydi*. Right valve.

FIG. 767.—*Actinopteria boydi*. Left valve.

tenuistriata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 120, Chemung Gr.

theta, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 125, Chemung Gr.

zeta, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 123, Chemung Gr.

ALLOSISMA, King, 1844, Ann. Mag. Nat. Hist.,

vol. 14, p. 315. [Ety. *allos*, variable; *ereisma*,

support, expressive of the variable nature

of the cartilage support or fulcrum.]

Equivalve, inequilateral, elongate, thin;

anterior side short; posterior long and

gaping at the extremity; beaks anterior,

depressed; surface concentrically

ridged or undulated; hinge edentulous;

ligament external; dorsal margin inflected,

forming a lanceolate depression

along the cardinal border behind the

beaks; anterior adductor scar occupying

a low position; pallial line faintly

marked. Type *A. sulcatum*.

altirostratum, see *Sedgwickia altirostrata*.

andrewsi, Whitfield, 1882, Ann. N. Y.

Acad. Sci., vol. 2, p. 222, Kaskaskia Gr.

antiquum, Swallow, 1863, Trans. St. Louis

Acad. Sci., vol. 2, p. 95, Kaskaskia Gr.

capax, Newberry, 1861, Ives' Col. Ex.

Exped., p. 120, Coal Meas.

clavatum, McChesney, 1860, New Pal.

Foss., p. 56, Kaskaskia Gr.

cooperi, see *Chenomya cooperi*.

costatum, Meek & Worthen, 1869, Proc.

Acad. Nat. Sci. Phil., p. 171, and Geo.

Sur. Ill., vol. 5, p. 585, Coal Meas.

cuneatum, Swallow, 1858, Trans. St. Louis

Acad. Sci., vol. 1, p. 210, Mid. Coal

Meas.

curtum, Swallow, 1858, Trans. St. Louis

Acad. Sci., vol. 1, p. 194, Permian Gr.

elegans, King, as identified by Geinitz.

See *A. geinitzi*.

elongatum, Morton, 1836, (*Pholadomya*

elongata), Am. Jour. Sci. and Arts, vol.

29, p. 153, Coal Meas.

elongatum, Worthen, see *A. worthenianum*.

ensiforme, Swallow, 1860, Trans. St. Louis

Acad. Sci., vol. 1, p. 656, Coal Meas.

geinitzi, Meek, 1867, Am. Jour. Sci., vol.

44, 2d ser., p. 170, and Geo. Sur. Ill.,

vol. 5, p. 586, Coal Meas.

gilberti, White, 1879, Bull. U. S. Geo.

Sur., vol. 5, No. 2, p. 217, and Cont. to

Pal., No. 6, p. 137, Carboniferous.

granosum, Shumard, 1858, (*Leptodomus*

granosus), Trans. St. Louis Acad. Sci.,

vol. 1, p. 207, and Pal. E. Neb. p. 220,

Coal Meas.

hannibalense, see *Grammysia hannibal-*

ensis.

hybridum, Meek & Worthen, 1865, (*Chæ-*

nomys hybridus), Proc. Acad. Nat. Sci.

Phil., p. 250, and Geo. Sur. Ill., vol. 3,

p. 538, Keokuk Gr.

illinoisense, Worthen, 1884, Bull. No. 2,

Ill. St. Mus. Nat. Hist., p. 11, and Geo.

Sur. Ill., vol. 8, p. 132, Keokuk Gr.

lanceolatum, Swallow, 1858, Trans. St.

Louis Acad. Sci., vol. 1, p. 194, Permian Gr.

latum, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 210, Mid. Coal Meas.

leavenworthense, see *Chænomya leavenworthensis*.

FIG. 768.—*Allorisma subcuneatum*.

marionense, White, 1876, Proc. Acad. Nat. Sci., p. 31, and Cont. to Pal., No. 8, p. 167, St. Louis Gr.

maxvillense, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 222, Kaskaskia Gr. minnehaha, see *Chænomya minnehaha*.

pleuropistha, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 70, and Ohio Pal., vol. 2, p. 309, Waverly Gr.

reflexum, Meek, 1872, Pal. E. Neb., p. 217, Coal Meas.

sinuatum, McChesney, 1860, New Pal. Foss., p. 56, Chester Gr.

FIG. 769.—*Allorisma subcuneatum*.

Geo. Sur. Ill., vol. 8, p. 133, which was preoccupied.

AMBONYCHIA, Hall, 1847, Pal. N. Y., vol. 1, p. 163. [Ety. *ambon*, the boss of a shield; *onyx*, a claw or talon.] Equivalve, inequilateral, subulate poste-

riorly, abrupt or curving down anteriorly; umbones high; beak incurved, cardinal line oblique; sinuate on the anterior side for the passage of the byssus; muscular impression large; cardinal tooth below the beak anteriorly; two or three remote lateral teeth, elongated and ranging parallel with the cardinal line posteriorly; surface radiately furrowed and concentrically lined. Type *A. bellistriata*.

acutirostra, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 383, Niagara Gr.

alata, see *Anomalodonta alata*. amygdalina, see *Cypicardites amygdalinus*.

aphaca, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 383, Niagara Gr.

attenuata, Hall, 1861, Geo. Rep. Wis., p. 33, and Geo. Wis., vol. 4, p. 206, Trenton Gr.

bellistriata, Hall, 1847, Pal. N. Y., vol. 1, p. 163, Trenton Gr.

cancellosa, Hall, 1861, Geo. Rep. Wis., p. 31. Mistake for *A. lamellosa*.

carinata, Goldfuss, 1826, (Pterinea carinata,) Germ. Petref., p. 136, and Pal. N. Y., vol. 1, p. 292, 294, Trenton and Hud. Riv. Gr.

cassii, Meek & Worthen, 1866, Proc. Chi. Acad. Nat. Sci., p. 22, Hud. Riv. Gr.

costata, Meek, 1873, Ohio Pal., vol. 1, p. 130, Hud. Riv. Gr.

erecta, Hall, 1861, Geo. Rep. Wis., p. 32, Trenton Gr.

illinoisensis, Worthen, 1875, Geo. Rep. Ill., vol. 6, p. 495, Hud. Riv. Gr.

intermedia, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 306, Galena Gr.

jamesi, Meek, 1872, (Megambonia jamesi,) Proc. Acad. Nat. Sci. Phil., p. 321, and Ohio Pal., vol. 1, p. 136, Hud. Riv. Gr.

lamellosa, Hall, 1861, Geo. Rep. Wis., p. 31, and Geo. Wis., vol. 4, p. 205, Trenton Gr.

maxima, Safford, 1869, Geo. of Tenn. Not defined.

mytiloides, Hall, 1847, Pal. N. Y., vol. 1, p. 315, Chazy Gr.

neglecta, see *Amphicoelia neglecta*.

nitida, Billings, 1866, Catal. Sil. Foss. Antic., p. 50, Anticosti Gr.

obtusa, see *Cypicardites obtusus*.

orbicularis, Emmons, 1842, (Pterinea orbicularis,) Geo. Rep. N. Y., p. 397, and Pal. N. Y., vol. 1, p. 164, Trenton Gr.

FIG. 770.—*Ambonychia bellistriata*. *h*, Byssal sinus; *t*, cardinal teeth; *u*, lateral teeth.

ing down ante-
beak incurved,
sinuate on the
passage of the
pression large;
below the beak
or three remote
elongated and
el with the car-
erriorly; surface
sowed and con-
d. Type A. bel-

1867, 20th Rep.
t. Hist., p. 383,

odonta alata.
Cypricardites

, 20th Rep. N. Y.
Niagara Gr.
eo. Rep. Wis.,
vol. 4, p. 206,

70.—*Ambonychia*
striata. *h.* Bassal
as; *l.* cardinal
th; *u.* lateral
h.

p. 22, Hud.

Pal., vol. 1, p.

ep. Wis., p. 32,

875, Geo. Rep.
Riv. Gr.

hen, 1868, Geo.
Palena Gr.

mbonia jamesi,
hil., p. 321, and

Hud. Riv. Gr.
p. Rep. Wis., p.

4, p. 205, Tren-

o. of Tenn. Not

. N. Y., vol. 1,

neglecta.
atal. Sil. Foss.

Gr.
obtusus.

2, (Pterinea or-
N. Y., p. 397,

p. 164, Tren-

planistriata, Hall, 1861, Geo. Rep. Wis.
p. 32, Trenton Gr.

FIG. 71.—*Ambonychia*
radiata.

Soc. Nat. Hist., vol. 1, p. 104, Hud.
Riv. Gr.

robusta, S. A. Miller, 1880, Jour. Cin. Soc.
Nat. Hist., vol. 3, p. 315, Hud. Riv. Gr.

striacosta, see *Pterinea striacosta*.

superba, Billings, 1866, Catal. Sil. Foss.
Antic., p. 50, Anticosti Gr.

sumana, Safford, 1869, Geo. of Tenn. Not
defined.

undata, Emmons, 1842, (*Pterinea undata*),
Geo. Rep. N. Y., p. 395, and Pal. N. Y.,
vol. 1, p. 165, Black Riv. and Trenton
Grs. Possibly belonging to an unde-
fined genus.

radiata, Hall, 1847,
Pal. N. Y., vol. 1,
p. 292, Trenton,
Hud. Riv. Grs.,
and Mid. Sil. Prob-
ably a syn. for *A.*
carinata.

rauchi, McChesney,
1860, New Pal.
Foss., p. 89, Hud.
Riv. Gr. Not rec-
ognized.

retrorsa, S. A. Miller,
1878, Jour. Cin.

large triangular cartilage pit beneath
the beaks, and smaller pit just anterior.
Type *A. leidyi*.

FIG. 73.—*Amphicella costata*.

costata, Hall & Whitfield, 1875, Ohio Pal.,
vol. 2, p. 140, Niagara Gr.

leidyi, Hall, 1867, 20th Rep. N. Y. Mus.
Nat. Hist., p. 387, Niagara Gr.

FIG. 72.—*Amnigenia catskillensis*.

AMNIGENIA, Hall, 1883, Pal. N. Y., vol. 5.
[Ety. *amnias*, a river; *gigno*, to bear.]
Like *Anodonta* in form and external
characters; anterior muscular impres-
sions large and prominent; posterior
ones large and shallow. Type *A. cats-*
killensis.

catskillensis, Vanuxem, 1842, (*Cypricar-*
dites catskillensis.) Geo. Rep. 3d Dist.
N. Y., p. 186, and Pal. N. Y., vol. 5, p.
516, Catskill Gr.

AMPHICELLA, Hall, 1868, 20th Rep. N. Y.
Mus. Nat. Hist., p. 386. [Ety. *amphi*,
both; *koilos*, hollow.] Equivalve, ine-
quilateral, subrhomboidal; umbones
gibbous; beaks elevated and incurved;
external ligamental area flattened;

neglecta, McChesney, 1861. (*Ambony-*
chia neglecta.) Pal. Foss., p. 88,
and Geo. Sur. Ill., vol. 3, p. 358, Niag-
ara Gr.

Amphidesma deloefeldi, Castelnau, 1843, Syst.
Sil., p. 44. Not recognized.

Anatina, Lamarck, 1809, Phil. Zool. [Ety.
pertaining to the duck, or like the
duck's bill.] Oblong, ventricose, atten-
uated, and gaping posteriorly; umbones
fissured; spoon-shaped cardinal process
in each valve. Type *A. rostrata*. Not
a Paleozoic genus.

leda, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 110, Ham. Gr. Not prop-
erly defined.

sinuata, see *Ilionia sinuata*.

ANGELLUM, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 105. [Ety.

FIG. 774.—*Angellum cuneatum*.

ANODONTOPSIS, McCoy, 1851, Ann. and Mag. Nat. Hist., 2d series, vol. 7, p. 54. [Ety.

from the resemblance to the shells of the genus *Anodonta*.] Equivalve, inequilateral, compressed; rotundato-quadrato or subtrigonal; posterior side wide, round, or obliquely subtruncate; anterior end slightly contracted in front of the beak; beaks small, prominent nearer the anterior than posterior end; hinge-line shorter than the length of the shell, with a posterior long, slender, lateral tooth extending just below it (double in the right valve), and another similar but shorter one in front of the beaks; anterior and posterior muscular impressions ovate; slight clavicular ridge between the beak and the adductor impressions; pallial impression entire; surface smooth or concentrically lined. Type *A. angustifrons*. Part of the generic definition is from *A. milleri*, as the interior of the type is not known.

amygdaliformis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 180, Devonian. *concinna*, Whiteaves, 1884, Pal. Foss., vol. 3, p. 12, Guelph Gr.

FIG. 775.—*Anodontopsis milleri*.

(?) *milleri*, Meek, 1871, Am. Jour. Sci., 3d series, vol. 2, p. 297, and Ohio Pal., vol. 1, p. 140, Hud. Riv. Gr.

unionoides, see *Modiolopsis unionoides ventricosa*, Billings, 1874, Pal. Foss., vol. 2, p. 55, Gaspe limestone No. 8, Devonian.

ANOMALODONTA, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 16. [Ety. *anomalos*, irregular; *odontos*, tooth.] Equivalve, inequilateral, alate posteriorly, abrupt anteriorly; umbones high; beak incurved; deeply sinuate for the byssus; cardinal ridge beneath the umbone sloping posteriorly; cartilage grooves extending from the cardinal

ridge to the termination of the posterior wing, and also from the cardinal

FIG. 776.—*Anomalodonta gigantea*. Hinge-line of right valve.

ridge to the byssal sinus; anterior muscular scar below the byssal sinus; surface radiately furrowed and concentrically lined. Type *A. gigantea*.

alata, Meek, 1872, (Amboynychia *alata*.) Proc. Acad. Nat. Sci. Phil., p. 319, and Ohio Pal., vol. 1, p. 131, Hud. Riv. Gr.

FIG. 777.—*Anomalodonta gigantea*. Left valve, showing hinge-line and muscular impression.

gigantea, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 17, Hud. Riv. Gr.

ANTHRACOMYA, Salter, 1861, Mem. Geo. Sur. Gr. Brit. Iron Ores, pt. 3, p. 229. [Ety. *anthrax*, coal; *Mya*, a genus.] Equivalve, inequilateral, mytiliform; ligament external; beak anterior; hinge-line straight; no teeth; surface concentrically marked; shell composed of an internal, lamellar, and subnacreous layer, a thin layer of vertical prismatic shell, and an epidermis; structure similar to the *Unionidae*. Type *A. elongata*.

angulata, Dawson, 1860, (*Naiadites angulatus*.) *Acadian Geology*, p. 205, *Coal Meas.*

FIG. 778.—*Anomalodonta gigantea*, external surface.

arenacea, Dawson, 1860, (*Naiadites arenacea*.) *Acadian Geology*, p. 205, *Coal Meas.*

carbonaria, Dawson, 1860, (*Naiadites carbonarius*.) *Acadian Geology*, p. 204, *Coal Meas.*

elongata, Dawson, 1860, (*Naiadites elongata*.) *Acadian Geology*, p. 204, *Coal Meas.*

FIG. 779.—*Anthracomya elongata*.

levis, Dawson, 1860, (*Naiadites levis*.) *Acadian Geology*, p. 204, *Coal Meas.*

obtusata, Dawson, 1860, (*Naiadites obtusatus*.) *Acadian Geology*, p. 205, *Coal Meas.*

ovalis, Dawson, 1860, (*Naiadites ovalis*.) *Acadian Geology*, p. 205, *Coal Meas.*

ANTHRACOPTERA, Salter, 1862, *Mem. Geo. Sur. Country Around Wigan*, p. 37. [Ety. *anthrax*, coal; *pteron*, a wing.] Shells small, aviculoid; height greater than width; valves subequal, wing short, hinge straight; surface concentrically marked.

carbonaria, see *Anthracomya carbonaria*. (?) *fragilis*, Meek & Worthen, 1866, *Proc. Chi. Acad. Sci.*, p. 18, *Keokuk Gr.*

levis, see *Anthracomya levis*. *polita*, White, 1880, 12th Rep. U. S. Geo. Sur. Terr., p. 166, *Coal Meas.*

ANTHRACOSTIA, King, 1844, *Azn. and Mag. Nat. Hist.*, p. 313. [Ety. *anthrax*, coal.] Equivalve, inequilateral; tooth in each valve below the umbo; crown of tooth of right valve excavated anteriorly and ridged posteriorly; crown of tooth of left valve ridged anteriorly and sloped posteriorly; furrow in hinge-plate, between umbone and tooth; scars of anterior pedal muscles above the anterior adductor impressions. Type A. *beanana*.

bradorica, Dawson, 1868, *Acad. Geol.*, p. 314, *Carb.*

Arca, Linne, 1758. This genus is unknown in the Palaeozoic rocks.

carbonaria, Cox. See Fig. 780.—*Anthracostia bradorica*. *Macrondon carbonarius*.

cuspidata, Swallow, 1858, *Trans. St. Louis Acad. Sci.*, vol. 1, p. 209, *Up. Coal Meas.* Founded on a cast. Genus unknown.

modesta, Winchell, 1863, *Proc. Acad. Nat. Sci.*, p. 15, *Marshall Gr.* Not an *Arca*.

striata, Schlottheim, as identified by Geinitz, is *Macrondon tenuistriatus*.

punctifera, Dawson, 1868, *Acad. Geol.*, *Carb.* The name was preoccupied by Deshayes in his work, 1824-1836.

Astarte, Sowerby, 1818, *Min. Conch.*, vol. 2, p. 85. Not a Palaeozoic genus.

mortonensis, see *Edmondia mortonensis*. *nebraskensis*, see *Edmondia nebraskensis*. *subtextilis*, see *Euthydesma subtextile*.

ASTARTELLA, Hall, 1858, *Geo. Rep. Iowa*, p. 715. [Ety. diminutive of *Astarte*.] Shell thick, smooth, or concentrically furrowed; lunule impressed, ligament external; hinge teeth, two in each valve; anterior tooth in right valve large and strong, with a longitudinal pit in the summit. Type A. *vera*.

concentrica, McChesney, 1860, (*Edmondia concentrica*.) *Descr. New Pal. Foss.*, p. 55, *Coal Meas.*

gurleyi, White, 1878, *Proc. Acad. Nat. Sci.*, p. 35, and *Cont. to Pal.*, No. 8, p. 166, *Coal Meas.*

newberryi, Meek, 1875, *Ohio Pal.*, vol. 2, p. 340, *Coal Meas.*

varica, McChesney, 1860, *Descr. New Pal. Foss.*, p. 55, *Coal Meas.*

vera, Hall, 1858, *Geo. Rep. Iowa*, p. 715, *Coal Meas.*

AVICULA, Klein, 1753, *Ostrac.* [Ety. *avicula*, a little bird.] Very inequivalve, inequilateral, obliquely oval; hinge produced posteriorly into a flattened defined wing; the inferior or right valve flattened, notched for the passage of the byssus; anterior muscular impression very small and faintly marked; adductor large, superficial, a little behind the middle; cartilage external, linear, simple, placed on a narrow marginal facet, extending from the beak toward the cardinal angle; hinge edentulous, or with two small cardinal teeth beneath the beak in one valve, and one in the other, and a long, slender, posterior bifid lateral tooth in each; substance corneo-calcareous, lamellar without, pearly within. Type A. *hirundo*. Not a Palaeozoic genus. Species are left here for want of better material to determine their generic relations.

acanthoptera, Hall, 1843, *Geo. Rep. 4th Dist. N. Y.*, p. 263, *Chemung Gr.*

acosta, Cox, 1857, Geo. Sur. Ky., vol. 3, p. 572, Coal Meas. The correct etymology would make this word *incosta*.
aequilatera, see *Aviculopecten aequilaterus*.
aequiradiata, Hall, 1859, Pal. N. Y., vol. 3, p. 285, Low. Held. Gr.

FIG. 781.—*Avicula hirundo*.

aesopus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 238, Ham. Gr.
angustirostra, Conrad, 1842, Jour. Acad. Nat. Sci., p. 236, Ham. Gr.
antiqua, see *Bakevellia antiqua*.
arenaria, Not American.
aviformis, see *Pterinea aviformis*.
bella, see *Aviculopecten bellus*.
bellula, Hall, 1859, Pal. N. Y., vol. 3, p. 289, Low. Held. Gr.
boydi, see *Actinopteria boydi*.
cancellata, see *Pterinea cancellata*.
chemungensis, see *Liopteria chemungensis*.
chemungensis, see *Pterinea chemungensis*.
circulus, see *Entolium circulus*.
communis, Hall, 1859, Pal. N. Y., vol. 3, p. 286, Low. Held. Gr.
cooperensis, see *Pterinea cooperensis*.
corrugata, see *Pterinea corrugata*.
cruciformis, see *Glyptodesma cruciforme*.
dannoniensis, Sowerby, as identified in the early N. Y. Reports. See *Liopteria chemungensis*.
decussata, see *Actinopteria decussata*.
demissa, see *Pterinea demissa*.
desquamata, Hall, 1847. The dorsal valve of *Obolella crassa*.
elliptica, see *Pterinea elliptica*.
emacerata, Conrad, 1842, Jour. Acad. Nat. Sci., p. 241, and Pal. N. Y., vol. 2, p. 83 and 282, Clinton and Niagara Grs.
erecta, see *Glyptodesma erectum*.
ferruginea, Conrad, 1848, Proc. Acad. Nat. Sci., vol. 3, p. 23, Up. Sil.
flabella, see *Pterinea flabellum*.
fragilis, see *Lunulicardium fragile*.
gebhardi, Conrad, 1841, Ann. Rep. N. Y., p. 54, Oriskany sandstone.
hermione, Billings, 1862, Pal. Foss., vol. 1, p. 40, Trenton Gr.
honeymani, see *Pterinea honeymani*.
insueta, see *Pterinea insueta*.
laevis, see *Liopteria laevis*.
leptonota, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 76, syn. for *A. emacerata*.
limiformis, Hall, 1852, Pal. N. Y., vol. 2, p. 332, Coralline limestone.
longa, Geinitz, 1866, (*Gervillia longa*), Carb. und Dyas in Neb., p. 32, and Pal. E. Neb., p. 199, Coal Meas.
longispina, see *Leptodesma longispinum*.
magna, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 98, Kaskaskia Gr.
manticula, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 241, and Pal. N. Y., vol. 3, p. 284, Low. Held. Gr.

morganensis, Meek & Worthen, 1866, (*Pterinea morganensis*), Proc. Acad. Nat. Sci., p. 259, and Geo. Sur. Ill., vol. 5, p. 576, Coal Meas.
multilineata, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 241, Chemung Gr.
muricata, see *Actinopteria muricata*.
naviformis, Conrad, 1842, Jour. Acad. Nat. Sci., p. 240, and Pal. N. Y., vol. 3, p. 279, Low. Held. Gr.
obliquata, Hall, 1859, Pal. N. Y., vol. 3, p. 285, Low. Held. Gr.
oblonga, see *Aviculopecten oblongus*.
obscura, Hall, 1859, Pal. N. Y., vol. 3, p. 280, Low. Held. Gr.
orbicularis, Stevens, 1858, Am. Jour. Sci., vol. 25, 2d ser., Coal Meas. The name was preoccupied by Sowerby in 1839.
orbiculata, Hall, 1843, see *Lyriopecten orbiculatus*.
orbiculata, Hall, 1852, Pal. N. Y., vol. 2, p. 284, Niagara Gr.
parilis, see *Aviculopecten parilis*.
pauceiradiata, Hall, 1859, Pal. N. Y., vol. 3, p. 287, Low. Held. Gr.
pectiniformis, see *Aviculopecten pectiniformis*.
perobliqua, see *Actinopteria perobliqua*.
pinniformis, Geinitz, 1848, (Solon pinniformis), Versteinerungen d. deutsch Zechsteingebirg, p. 8, and Carb. und Dyas in Neb., p. 31, Coal Meas.
pleuroptera, see *Actinopteria pleuroptera*.
protexta, see *Leptodesma protextum*.
quadrula, syn. for *Actinopteria boydi*.
rectilateraria, see *Aviculopecten rectilateraria*.
recticosta, Hall, 1859, Pal. N. Y., vol. 3, p. 466, Oriskany sandstone.
rhomboidea, Hall, 1852, Pal. N. Y., vol. 2, p. 84, Clinton Gr.
rugosa, see *Pterinea rugosa*.
schohariae, Hall, 1859, Pal. N. Y., vol. 3, p. 283, Low. Held. Gr.
securiformis, Hall, 1852, Pal. N. Y., vol. 2, p. 331, Coralline limestone.
securiformis, Hall, 1859, Pal. N. Y., vol. 3, p. 290. This name was preoccupied.
semielliptica, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 210, Up. Coal Meas.
shawneensis, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 211, Up. Coal Meas.
shumardi, see *Entolium shumardi*.
signata, see *Aviculopecten signatus*.
speciosa, see *Panenka speciosa*.
spinigera, see *Leptodesma spinigerum*.
spinnulifera, Hall, 1859, Pal. N. Y., vol. 3, p. 282, Low. Held. Gr.
subaequilatera, Hall, 1859, Pal. N. Y., vol. 3, p. 281, Low. Held. Gr.
subfalcata, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 242, Ham. Gr.
subplana, Hall, 1852, Pal. N. Y., vol. 2, p. 283, Niagara Gr.
subquadrans, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 236, Devonian.
subrecta, see *Aviculopecten subrectus*.

Worthen, 1866, Proc. Acad. Nat. Sci. Ill., vol. 5, p. 2, Jour. Acad. Chemung Gr. muricata. Jour. Acad. N. Y., vol. 3, p. 1, N. Y., vol. 3, oblongus. N. Y., vol. 3, p. 1, Am. Jour. Sci., 1865. The name verby in 1839. Lyriopecten or. N. Y., vol. 2, parilis. Pal. N. Y., vol. 1, Pecten pectini- a perobliqua. (Solon pinnæ- n d. deutsch Carb. und Dyas us. ia plenroptera. protextum. teria boydi. pecten rectila- N. Y., vol. 3, ne. N. Y., vol. 2, N. Y., vol. 3, al. N. Y., vol. stone. N. Y., vol. 3, preoccupied. 1858, Trans. St. 210, Up. Coal 1858, Trans. St. 211, Up. Coal umardi. signatus. losa. pinigerum. N. Y., vol. 3, Pal. N. Y., vol. ur. Acad. Nat. Gr. N. Y., vol. 2, Jour. Acad. Devonian. subrectus.

aurugosa, D'Orbigny, 1850, Prodr. d. Paléont. t. 1, p. 33. Syn. for *Pterinea rugosa*. *tenuilamellata*, Hall, 1859, Pal. N. Y., vol. 3, p. 281, Low. Held. Gr. *textilis*, Hall, 1859, Pal. N. Y., vol. 3, p. 288, Low. Held. Gr. *textilis* var. *arenaria*, Hall, 1859, Pal. N. Y., vol. 3, p. 465, Oriskany Gr. *trentonensis*, see *Pterinea trentonensis*. *tricolata*, see *Lyriopecten tricolatus*. *trilobata*, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 240, Ham. Gr. *triplistriata*, Stevens, 1858, Am. Jour. Sci., vol. 25, p. 265, Coal Meas. *triquetra*, Hall, 1843, Geo. Rep. N. Y., p. 137, Onondaga Gr. *tuberculata*, Conrad, 1838, Ann. Rep. N. Y., p. 117, Corniferous Gr. *umbonata*, Hall, 1859, Pal. N. Y., vol. 3, p. 284, Low. Held. Gr. *undata*, Hall, 1852, Pal. N. Y., vol. 2, p. 283, Niagara Gr. *undosa*, Ringuenberg, 1886, Bull. Buf. Soc. Nat. Sci., vol. 5, p. 18, Niagara Gr. *welchi*, James, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 239, Hud. Riv. Gr. *whitii*, Winchell, 1863, Proc. Acad. Nat. Sci., p. 8, Marshall Gr.

AVICULOPECTEN, McCoy, 1851, Ann. Mag. Nat. Hist., 2d ser., vol. 7, p. 171. [Ety. from the genera *Avicula* and *Pecten*.] Inequivalve, inequilateral; straight or slightly extended obliquely toward the posterior side; anterior ear flattened, smaller than the posterior, sharply and deeply defined, with a notch in the right valve between it and the body of the shell for the passage of the byssus; posterior ear pointed, extending about as far as the margin of the shell, defined or not; ligament confined to a narrow facet along the hinge margin, or having a wider cardinal area with cartilage furrows; no medial cartilage pit; muscular impression and pallial scar as in *Pecten*. Type *A. docens*.

aeadicus, Hartt, 1868, Acad. Geol., p. 307, Carb. *acutialatus*, Swallow, 1858, (*Avicula acutialata*.) Trans. St. Louis Acad. Sci., p. 185, Permian Gr. *aquilatralia*, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 19, Chemung Gr. *aquilaterus*, Hall, 1843, (*Avicula aequilata*.) Geo. Rep. 4th Dist. N. Y., p. 181, Up. Held. Gr. and Marcellus Shale. *affinis*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 229, Subcarboniferous. *amplus*, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 454, and Geo. Sur. Ill., vol. 2, p. 257, Keokuk Gr. *armigerus*, Conrad, 1835, (*Pecten armigerus*.) Trans. Geo. Soc. Penn., p. 268, Coal Meas. *bellus*, Conrad, 1841, (*Avicula bella*.) Ann. Rep. N. Y., p. 54, and Pal. N. Y., vol. 5, pt. 1, p. 35, Ham. Gr.

burlingtonensis, Meek & Worthen 1860, Proc. Acad. Nat. Sci., p. 453, and Geo. Sur. Ill., vol. 2, p. 231, Burlington Gr. *cancellatus*, Hall, 1843, (*Pecten cancellatus*.) Geo. Rep. 4th Dist. N. Y., p. 264, and Pal. N. Y., vol. 5, pt. 1, p. 18, Chemung Gr. *carboniferus*, Stevens, 1858, (*Pecten carboniferus*.) Am. Jour. Sci. and Arts, vol. 25, p. 261, and Pal. E. Neb., p. 193, Coal Meas. *caroli*, Winchell, 1863, Proc. Acad. Nat. Sci., p. 9, and Pal. N. Y., vol. 5, pt. 1, p. 29, Waverly Gr. *catactus*, Meek, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 93, Carboniferous. *celsus*, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 23, Chemung Gr. *chesterensis*, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 20, and Geo. Sur. Ill., vol. 8, p. 115, Kaskaskia Gr. *cleon*, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 6, Up. Held. Gr. *clevelandicus*, Swallow, 1858, (*Pecten clevelandicus*.) Trans. St. Louis Acad. Sci., vol. 1, p. 184, Permian Gr. *colletti*, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 21, and Geo. Sur. Ill., vol. 8, p. 119, Keokuk Gr. *coloradoensis*, Newberry, 1861, Ives' Col. Ex. Exped., p. 129, Coal Meas. *convexus*, Hall, 1843, (*Pecten convexus*.) Geo. Rep. 4th Dist. N. Y., p. 265, and Pal. N. Y., vol. 5, pt. 1, p. 28, Chemung Gr. *cora*, Dawson, 1868, Acad. Geol., p. 307, Carb. *coreyanus*, White, 1874, Rep. Invert. Foss., p. 21, and Geo. Sur. W. 100th Mer., vol. 4, p. 147, Coal Meas. *coxanus*, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 453, and Geo. Sur. Ill., vol. 2, p. 326, Low. Coal Meas. *crassicostratus*, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 183, Up. Held. Gr. *crenistriatus*, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 60, and Ohio Pal., vol. 2, p. 295, Waverly Gr. *curticardinalis*, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 273, Coal Meas. *debertanus*, Dawson, 1868, Acad. Geol., p. 307, Carboniferous. *dolabriformis*, Hall, 1843, (*Pecten* (?) *dolabriformis*.) Geo. Rep. 4th Dist. N. Y., p. 265, and Pal. N. Y., vol. 5, pt. 1, p. 26, Chemung Gr. *duplicatus*, Hall, 1843, (*Pecten duplicatus*.) Geo. Rep. 4th Dist. N. Y., p. 264, and Pal. N. Y., vol. 5, pt. 1, p. 17, Chemung Gr. *edwardsi*, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 22, and Geo. Sur. Ill., vol. 8, p. 119, Keokuk Gr. *ellipticus*, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 25, Chemung Gr.

elsahensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 19, and Geo. Sur. Ill., vol. 8, p. 115, Kinderhook Gr.
eurekensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 227, Subcarboniferous.
exacutus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 8, Ham. Gr.
fasciculatus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 11, Ham. Gr.
formio, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 9, Ham. Gr.
glaber, see *Pernopecten glaber*.
gradocostatus, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 31, Marshall Gr.
haguei, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 226, Subcarboniferous.
halli, Swallow, 1860, (Avicula halli,) Trans. St. Louis Acad. Sci., vol. 1, p. 656, Coal Meas.

FIG. 782.—*Avicutopecten princeps*.

hardinensis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 117, St. Louis Gr.
hertzeri, Meek, 1871, Proc. Acad. Nat. Sci., p. 61, and Ohio Pal., vol. 2, p. 330, Coal Meas.
idas, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 13, Ham. Gr.
ignotus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 33, Up. Held. Gr.
incultus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 30, Up. Chemung Gr.
indianensis, Meek & Worthen, 1866, Proc. Chi. Acad. Sci., vol. 1, p. 14, Keokuk Gr.
insignis, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 34, Ham. Gr.
intercostalis, Winchell, 1866, Rep. Low. Peninsula Mich., p. 95, Ham. Gr.
internatus, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 454, and Geo. Sur. Ill., vol. 2, p. 329, Low. Coal Meas.

invalidus, Hall, 1883, (Pterinopecten invalidus,) Pal. N. Y., vol. 5, pt. 1, p. 31, Marcellus Shale.

lowensis, S. A. Miller, 1883, 2d Ed. Am. Pal. Foss., p. 310, Marshall or Kinderhook Gr., at Burlington, Iowa. Proposed instead of *A. occidentalis* of Winchell, in 1863, in Proc. Acad. Nat. Sci., Phil., p. 9, which was preoccupied by Shumard.

itys, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 20, Chemung Gr.

konincki, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 453, and Geo. Sur. Ill., vol. 2, p. 328, Low. Coal Meas.
latus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 14, Ham. Gr.

limaformis, see *Pernopecten limaformis*.

lyelli, Dawson, 1868, Acad. Geol., p. 305, Carb.

lyelli var. *alternans*, Dawson, 1883, Rep. on Redpath Mus., p. 12, Carboniferous.

maccoyi, Meek & Hayden, 1865, Pal. Up. Mo., p. 50, Permian Gr.

macwhorteri, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 118, Kinderhook Gr.

mazonensis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 117, Coal Meas.

menardi, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 120, Coal Meas.

missouriensis, Shumard, 1855, (Pecten missouriensis,) Geo. Rep. Mo., p. 207, St. Louis Gr.

monroensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 21, and Geo. Sur. Ill., vol. 8, p. 114, St. Louis Gr.

mucronatus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 38, Ham. Gr.

newarkensis, Winchell, 1870, Notices and Desc. Foss. from Marshall Gr., Proc. Acad. Nat. Sci., p. 255, Marshall Gr.

niotensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 19, and Geo. Sur. Ill., vol. 8, p. 113, Keokuk Gr.

nodocostatus, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 296, Kinderhook Gr.

oblongus, Meek & Worthen, 1860, (Avicula oblonga,) Proc. Acad. Nat. Sci. Phil., p. 454, and Geo. Sur. Ill., vol. 2, p. 258, Keokuk Gr.

occidaneus, Meek, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 96, Carboniferous.

occidentalis, Shumard, 1855, (Pecten occidentalis,) Geo. Rep. Mo., p. 207, Carboniferous and Permian.

occidentalis, Winchell, 1863, Proc. Acad. Nat. Sci., p. 9. This name was preoccupied.

riopecten in-
5, pt. 1, p. 31,
3, 2d Ed. Am.
Hall or Kinder-
Iowa. Pro-
entalis of Win-
Acad. Nat. Sci.,
oreoccupied by
Z., vol. 5, pt. 1,
en, 1860, Proc.
453, and Geo.
ow. Coal Meas.
Y., vol. 5, pt. 1,

limiformis.
Geol., p. 305,
son, 1883, Rep.
h Mus., p. 12,
ous.
ek & Hayden,
Up. Mo., p. 50,
r.
Worthen, (in
p. Sur. Ill., vol.
Kinderhook Gr.
Worthen, (in
p. Sur. Ill., vol.
Coal Meas.
Worthen, (in
p. Sur. Ill., vol.
Coal Meas.
s, Shumard,
en missourien-
Rep. Mo., p. 207,
r.
Worthen, 1884,
2, Ill. St. Mus.
p. 21, and Geo.
ol. 8, p. 114, St.

Hall, 1883, Pal.
5, pt. 1, p. 38,
70, Notices and
Hall Gr., Proc.
Marshall Gr.
Bull. No. 2, Ill.
p. 19, and Geo.
Keokuk Gr.
Whitfield, 1862,
lisc., vol. 8, p.

en, 1860, (Avic-
Acad. Nat. Sci.
Sur. Ill., vol. 2.

. S. Geo. Expl.
96, Carbonif-

55, (Pecten oc-
c., p. 207, Car-
3, Proc. Acad.
me was proc-

orbiculatus, see *Lyriopecten orbiculatus*.
orestes, Worthen, 1884, Bull. No. 2, Ill.
St. Mus. Nat. Hist., p. 18, and Geo. Sur.
Ill., vol. 8, p. 112, Keokuk Gr.
ornatus, Hall, syn. for *A. fasciculatus*.
ornatus, Hall, 1883, Pal. N. Y., vol. 5, pt.
1, p. 37, Ham. Gr.
oweni, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 452, and Geo.
Sur. Ill., vol. 2, p. 256, Keokuk Gr.
parilis, Conrad, 1842, (*Avicula parilis*,)
Jour. Acad. Nat. Sci. Phil., vol. 8, p. 239,
and Ohio Pal., vol. 1, p. 197, Cornif. Gr.
parvulus, Hall & Whitfield, 1877, U. S.
Geo. Expl. 40th parallel, vol. 4, p. 274,
Coal Meas.

plenus, Hall, 1883, Pal. N. Y., vol. 5, pt.
1, p. 21, Chemung Gr.

FIG. 783.—*Avienlopecten princeps*. Cardinal
part showing ligamental area.

princeps, Conrad, 1838, (*Monotis prin-
ceps*,) Ann. Rep. N. Y., p. 117, and Pal.
N. Y., vol. 5, pt. 1, p. 1, Ham. Gr.

FIG. 784.—*Avienlopecten varsoviensis*.

patulus, Hall, 1883, Pal. N. Y., vol. 5, pt.
1, p. 24, Up. Chemung Gr.
pecteniformis, Conrad, 1842, (*Avicula
pecteniformis*,) Jour. Acad. Nat. Sci.,
vol. 8, p. 240, and Pal. N. Y., vol. 5, pt.
1, p. 4, Up. Held. Gr. and Marcellus
Shale.
pellucidus, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 455, and Geo.
Sur. Ill., vol. 2, p. 327, Low. Coal Meas.
perocidens, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 227, Subcarbonif-
erous.
phorceus, Hall, 1883, Pal. N. Y., vol. 5, pt.
1, p. 10, Ham. Gr.
pintoensis, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 228, Subcarbonif-
erous.

providencensis, Cox, 1857, (*Pecten provi-
dencensis*,) Geo. Sur. Ky., vol. 3, p. 566,
Coal Meas.
rectilaterarius, Cox, 1857, (*Avicula recti-
lateraria*,) Geo. Sur. Ky., vol. 3, p. 571,
Coal Meas.
repletus, Hall, syn. for *A. fasciculatus*.
reticulatus, Dawson, 1868, Acad. Geol., p.
306, Carboniferous.
ringens, Swallow, 1858, (*Pecten ringens*,)
Trans. St. Louis Acad. Sci., p. 184, Per-
mian Gr.
rugistriatus, Hall, 1843, (*Lima rugestri-
ata*,) Geo. Rep. 4th Dist. N. Y., p. 264,
and Pal. N. Y., vol. 5, pt. 1, p. 15, Chem-
mung Gr.
sanduskiensis, Meek, 1871, Proc. Acad.
Nat. Sci. Phil., p. 161, Up. Held. Gr.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

28
25
22
20
18

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

- scabridus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 7, Ham. Gr.
 signatus, Hall, 1843, (Avicula signata,) Geo. Rep. 4th Dist. N. Y., p. 265, and Pal. N. Y., vol. 5, pt. 1, p. 29, Chemung Gr.
 simplex, Dawson, 1868, Acad. Geol., p. 306, Carboniferous.
 spinuliferus, Meek & Worthen, 1870, Proc. Acad. Nat. Sci., p. 39, and Geo. Sur. Ill., vol. 8, p. 116, Keokuk Gr.
 squama, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 27, Chemung Gr.
 striatus, Hall, 1843, (Pecten striatus,) Geo. Rep. 4th Dist. N. Y., p. 264, and Pal. N. Y., vol. 5, pt. 1, p. 22, Chemung Gr.
 subcancellatus, Hall, 1883, syn. for A. cancellatus.
 subrectus, Hall, 1852, (Avicula subrecta,) Pal. N. Y., vol. 2, p. 331, Coralline limestone.
 talboti, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 21, St. Louis Gr.
 tenuicostus, Winchell, 1863, Proc. Acad. Nat. Sci., p. 10, Marshall Gr.
 tenuis, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 39, Up. Chemung Gr.
 terminalis, Hall, 1883, (Pterinopecten terminalis,) Pal. N. Y., vol. 5, pt. 1, p. 32, Up. Held. Gr.
 unionensis, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 511, Corniferous Gr.
 utahensis, Meek, 1860, (Pecten utahensis,) Proc. Acad. Nat. Sci., p. 310, Coal Meas.
 varsoviensis, Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 321, Keokuk Gr.
 weberensis, Hall & Whitfield, 1877, U. S. Geo. Sur., 40th parallel, vol. 4, p. 273, Coal Meas.
 whitii, Meek, 1872, Pal. E. Neb., p. 195, Coal Meas.
 williamsi, Meek, 1871, Proc. Acad. Nat. Sci., p. 178, Choteau limestone.
 winchelli, see Crenipecten Winchelli.
 AVICULOPINNA, Meek, 1867, Am. Jour. Sci., vol. 44, 2d ser., p. 282. [Ety. the genera *Avicula* and *Pinna*.] Compressed, slender, elongated, subtrigonal, or nearly in the form of a *Pinna*; beaks nearly obsolete, extremely oblique, and slightly behind the anterior extremity. Type *A. americana*.
 americana, Meek, 1867, Am. Jour. Sci., vol. 44, 2d ser., p. 232, and Pal. E. Neb., p. 197, Coal Meas.

FIG. 785.—*Aviculopinna americana*.

- illinoisensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 13, and Geo. Sur. Ill., vol. 8, p. 123, Coal Meas.
Azinas, Sowerby, 1821, Min. Conch., vol. 3. [Ety. *axine*, battle-axe.] This genus is unknown in Palaeozoic rocks.

- ovatus, see Schizodus ovatus.
securis, Shumard, 1859, Trans. St. Louis Acad. Sci., Permian Gr. Not recognized.
 BAKEVELLIA, King, 1849, Perm. Foss., p. 166. [Ety. proper name.] Shell aviculiform, subequivalve; valves sinuous, gaping in front for the passage of the byssus; umbones depressed, oblique; surface with concentric striae; hinge with linear anterior and posterior lateral teeth parallel to the cardinal margin; muscular scars as in *Pteria*; cardinal area in both valves; two to five cartilage furrows in each valve. Type *B. antiqua*.
antiqua, Munster, 1826, (Avicula antiqua,) Goldfuss Germ. Petref. Not American.
 illinoisensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 14, and Geo. Sur. Ill., vol. 8, p. 126, Up. Coal Meas.
 parva, Meek & Hayden, 1858, Trans. Alb. Inst., vol. 4, p. 78, and Pal. Up. Mo., p. 57, Permian Gr.
 (?) pulchra, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 189, Permian Gr.
 sulcata, Geinitz, 1866, (Gervillia sulcata,) Carb. und Dyas in Neb., p. 33, Coal Meas.
 BYSSOPTERIA, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 4. [Ety. *byssos*, byssus; *Pteria*, a genus.] Shell erect, equivalve, alate posteriorly, truncate, with a nasute projection in front; surface radiately furrowed and concentrically lined. Type *B. radiata*.

FIG. 786.
Bakevella
parva. FIG. 787.—*Byssopteria radiata*.

- radiata*, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 252, Up. Chemung Gr.
 CARBONARCA, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 39. [Ety.

19.
rans. St. Louis
Not recognized.
m. Foss., p. 166.
nell aviculiform,
uous, gaping in
of the byssus;
blique; surface
inge with linear
lateral teeth par-
argin; muscular
nal area in both
ilage furrows in
antiqua.

vicula antiqua,
Not American.
1884, Bull. No. 2,
p. 14, and Geo.
Up. Coal Meas.

358,
p. 57,
FIG. 786.
Bakevillia
parva.

Sci.,
Gr.
ervillia sulcata,
p. 33, Coal Meas.
al. N. Y., vol. 5,
p. byssus; Pteria,
equivalve, alate
with a nasute pro-
cess radiately fur-
row lined. Type

la radiata.

N. Y., vol. 5, pt.
g Gr.
rthen, 1870, Proc.
l., p. 39. [Ety.

carbo, coal; Arca, a genus.] Inequi-
valve, inequilateral, very convex, trans-
versely oblong or oval; umbones gib-
bous, prominent, strongly incurved,
with subangular posterior slopes;
valves closed all around with smooth
margins; ligament external; cardinal
margin arched; two anterior oblique
teeth, and behind these minute crenu-
lations, as in Arca. Type C. gibbosa.
gibbosa, Meek & Worthen, 1870, Proc.
Acad. Nat. Sci. Phil., p. 40, and Geo.
Sur. Ill., vol. 6, p. 531, Coal Meas.

FIG. 788.—Cardinia
listeri.

teeth obscure, lateral, remote, promi-
nent; adductor impressions deep; pal-
lial line simple. Type C. listeri.

aequimarginalis, see Edmondia aequimar-
ginalis.

antigonesensis, Dawson, 1868, Acad. Geo.,
p. 304, Carb.

complanata, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 413, Portage Gr.

concentrica, see Sanguinolites concentricus.

cordata, Swallow, 1858, Trans. St. Louis
Acad. Sci., vol. 1, p. 191, Permian Gr.

(?) fragilis, Cox, 1857, Geo. Sur. Ky., vol.
3, p. 570, Coal Meas.

occidentalis, Swallow, 1860, Trans. St.
Louis Acad. Sci., p. 655, Waverly or
Choteau Gr.

subangulata, Swallow, 1858, Trans. St.
Louis Acad. Sci., vol. 1, p. 192, Per-
mian Gr.

subangulata, Dawson, 1868, Acad. Geol., p.
304. This name was preoccupied.

CARDIOLA, Broderip,
1844, Trans.

Geo. Soc. [Ety.
kardia, the
heart.] Oblique-

ly oval or sub-
circular, tumid,
equivalve, in-

equilateral;
beaks large,
prominent,

obliquely incurved anteriorly; ends
subequal, rounded; ventral margin
convex; hinge-line shorter than the
shell, with a flattened cardinal area,
widest between the beaks, extending
its whole length; surface radiately
ribbed. Type C. interrupta.

equilatera, see Panenka equilatera.

dichotoma, see Panenka dichotoma.

doris, see Paracardium doris.

elevata, see Panenka ventricosa.

erecta, see Pararca erecta.

ilicostata, Walcott, 1885, Monogr. U. S. Geo.
Sur., vol. 8, p. 251, Subcarboniferous.

hero, see Panenka hero.

lincklani, see Panenka lincklani.

radians, see Panenka radians.

robusta, see Panenka robusta.

salteri, Haughton, 1857, Jour. Roy. Soc.
Dub., vol. 1, Devonian.

sao, see Pararca sao.

speciosa, Hall, 1883, Pal. N. Y., vol. 5, pl.
70, fig. 2-9, and pl. 80, fig. 10, Genesee
Slate.

transversa, see Pararca transversa.

CARDIOMORPHA, DeKoninck, 1844, Anim.
Foss. Carb. Belg., p. 101. [Ety. kardia,
heart; morphe, form.] Shell very thin,
equivalve, inequilateral, margins closed,
oblique, tumid; beaks tumid, produced,
spirally inrolled to the anterior side;
no hinge teeth; hinge margin inflected
nearly at right angles to form a hollow
lunette, running from the beak
nearly to the cardinal angle; two ad-
ductor impressions in each valve;
pallial scar simple, very faintly marked;
a shallow anterior depression beneath
the beaks, but the margin sharp and
prominent. Type C. elongata.

FIG. 790.—Cardiomorpha cordata.

archiacana, DeKoninck, 1843, Desc. An.
Foss. Belg., p. 104, Carboniferous.

bellatula, Hall, 1870, Prelim. Notice Lam.
Shells, p. 92, and Pal. N. Y., vol. 5, pl.
63, figs. 1-3, Ham. Gr.

capuloides, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 416, Marshall Gr.

concentrica, Hall, 1883, Pal. N. Y., vol. 5,
pl. 63, fig. 4, syn. for C. zonata.

cordata, Hall, 1883, Pal. N. Y., vol. 5, pl.
62, figs. 10-19, Ham. Gr.

donaciformis, Hall, 1883, Pal. N. Y., vol.
5, pl. 63, fig. 6, Ham. Gr.

eriopia, Hall, 1870, Prelim. Notice Lam.
Shells, p. 92, and Pal. N. Y., vol. 5, pl.
63, figs. 7-8, Ham. Gr.

julia, Winchell, 1862, Proc. Acad. Nat.
Sci., p. 416, Marshall Gr.

kansasensis, Swallow, 1858, Trans. St.
Louis Acad. Sci., vol. 1, p. 191, Per-
mian Gr.

missouriensis, Shumard, 1858, Trans. St.
Louis Acad. Sci., vol. 2, p. 207, and Geo.
Sur. Ill., vol. 5, p. 588, Coal Meas.

modiolaris, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 416, Marshall Gr.

(?) obliquata, Meek, 1872, Proc. Acad.
Nat. Sci. Phil., p. 327, and Ohio Pal.,
vol. 1, p. 146, Had. Riv. Gr.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 790.—Cardiomorpha cordata.

FIG. 789.—Cardiola in-
terrupta.

- oblonga*, see *Protomya oblonga*.
ovata, see *Dexiobia ovata*.
parvirostris, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 31, syn. for *Dexiobia ovata*.
pellensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 16, and Geo. Sur. Ill., vol. 8, p. 126, St. Louis Gr.
radiata, see *Cardiopsis radiata*.
rhomboidea, Hall, see *Cardiomorpha subrhomboidea*.
rhomboidea, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 191, Permian Gr.
rotunda, Hall, 1883, Pal. N. Y., vol. 5, pl. 63, figs. 17-20, refer figs. 18 and 19 to *Paracyclas rotunda*, fig. 17 to *Schizodus degener*, and fig. 20 to *S. patulus*.
subglobosa, Meek, 1875, Ohio Pal., vol. 2, p. 304, Waverly Gr.
suborbicularis, Hall, 1843, (Ungulina suborbicularis,) Geo. Rep. 4th Dist. N. Y., p. 244, and Pal. N. Y., vol. 5, pl. 63, figs. 9-10, Portage Gr.
subrhomboidea, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 186, (proposed instead of *Cypricardites rhomboidea*, in Geo. Rep. Iowa, p. 523, which was preoccupied,) Kinderhook Gr.
textilis, Hall, 1883, Pal. N. Y., vol. 5, pl. 63, figs. 11-15, Chemung Gr.
triangulata, Swallow, 1860, Trans. St. Louis Acad. Sci., vol. 1, p. 655, Waverly or Choteau Gr.
trigonalis, Winchell, 1863, Proc. Acad. Nat. Sci., p. 15, Marshall Gr.
undulata, Hall, 1883, Pal. N. Y., vol. 5, pl. 63, fig. 16, Portage Gr.
 (?) *vetusta*, see *Cypricardites vetustus*.
vindobonensis, Hartt, 1868, Acad. Geol., p. 304, Carboniferous.
zonata, Hall, 1883, Pal. N. Y., vol. 5, pl. 63, fig. 5, Ham. Gr.
- CARDIOPSIS**, Meek & Worthen, 1861, Proc. Acad. Nat. Sci. Phil., p. 144. [Ety. *kardia*, the heart; *opsis*, appearance.] Equivalve, somewhat inequilateral, oblique, ovate or cordiform, entirely closed; beaks elevated, incurved, directed anteriorly; cardinal margin short; rounding into the posterior border; two anterior teeth in each valve; surface radiated. Type C. *radiata*.
crassicostata, Hall & Whitfield, 1873, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 188, Schoharie grit and Corniferous limestone.
crenistriata, see *Pterinea crenistriata*.
jejuna, Winchell, 1862, Proc. Acad. Nat. Sci., p. 417, Marshall Gr.
megambonata, Winchell, 1862, Proc. Acad. Nat. Sci., p. 417, Marshall Gr.
parvirostris, White, 1862, Proc. Bost. Soc. Nat. Hist., vol. 9, p. 31, syn. for *Dexiobia ovata*.
radiata, Meek & Worthen, 1860, (*Cardiomorpha radiata*,) Proc. Acad. Nat. Sci. Phil., p. 458, and Geo. Sur. Ill., vol. 2, p. 157, Kinderhook Gr.

- Cardium*, Linnaeus, 1758, Syst. Nat., 10th Ed. [Ety. *kardia*, the heart.] Not a Palaeozoic genus.
iowensis, see *Cypricardites iowensis*.
lexingtonensis, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 207, Mid. Coal Meas.
nautiloides, Castelnau, 1843, Syst. Sil. Seneca Lake, N. Y. Not recognized.
vetustum, see *Præcardium vetustum*.

CHÆNOCARDIA

- CHÆNOCARDIA**, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 170. [Ety. *chaino*, to gape; *kardia*, the heart.] Ovate, ventricose, gaping anteriorly, edgetruncated, hinge-line short, beaks small, incurved; surface concentrically marked. Type C. *ovata*.

FIG. 791.—Chænocardia ovata.

- ovata*, Meek & Worthen, 1869, Proc. Acad. Nat. Sci. Phil., p. 170, and Geo. Sur. Ill., vol. 5, p. 586, Coal Meas.
CHENOMYA, Meek, 1864, Pal. of Up. Mo., p. 42. [Ety. *chaino*, to open or gape; *Mya*, a genus of shells.] Shell thin, equivalent, longitudinally oblong, subcylindrical; anterior side rounded, closed; posterior side long, truncated, gaping at the extremity; surface granulose and concentrically marked; cardinal margin inflected as in *Allorisma*; ligament external; hinge edentulous; posterior muscular impressions near the posterior extremity of the dorsal margin; scars of the anterior adductor and pedal muscles connected; pallial line with a broad shallow sinus. Type C. *leavenworthensis*.

FIG. 792.—Chenomya maria. Right valve.

- cooperi*, Meek & Hayden, 1858, (Panopaea cooperi,) Trans. Alb. Inst., vol. 4, p. 83, and Pal. Up. Mo., p. 44, Coal Meas.
hybrida, see *Allorisma hybridum*.

Nat., 10th Ed.
Not a Palaeo-

owensia.
3, Trans. St.
p. 207, Mid.

3, Syst. Sil.
recognized.
tustum.

occardia ovata.
ncurved; sur-
ked. Type C.

, 1869, Proc.
170, and Geo.
al Meas.

of Up. Mo., p.
open or gape;
Shell thin,

oblong, sub-
side rounded,
ng, truncated.

surface gran-
marked; cardi-
in Allorisma;

edentulous;
pressions near
of the dorsal

erior adductor
nected; pallial
v sinus. Type

Right valve.

858, (Panopaea
t., vol. 4, p. 83,
Coal Meas.
ridum.

leavenworthensis, Meek & Hayden, 1858,
(Allorisma leavenworthense,) Proc.
Acad. Nat. Sci. Phil., p. 263, and Pal.
Up. Mo., p. 43, Coal Meas.

maria, Worthen, 1882, Bull. No. 1. Ill. St.
Mus. Nat. Hist., p. 39, and Geo. Sur.
Ill., vol. 7, p. 319, Up. Coal Meas.

FIG. 793.—Chenomya
maria. Dorsal
view.

face concentrically lined. Type C. re-
curva.

angulata, Hall, 1885, Pal. N. Y., vol. 5, p.
468, Chemung Gr.

corrugata, Conrad, 1842, (Cypricardites
corrugatus,) Jour. Acad. Nat. Sci., vol. 8,
p. 244, and Pal. N. Y., vol. 5, pl. 77, figs.
1-4, Ham. Gr.

FIG. 794.—Cimitaria recurva.

elongata, Conrad, 1841, (Cypricardites
elongatus,) Ann. Rep. N. Y., p. 51, and
Pal. N. Y., vol. 5, pl. 77, figs. 5-8,
Ham. Gr.

recurva, Conrad, 1842, (Cypricardites re-
curvus,) Jour. Acad. Nat. Sci., vol. 8, p.
245, and Pal. N. Y., vol. 5, pl. 77, figs.
9-16, Ham. Gr.

CLIDOPHORUS, Hall, 1847, Pal. N. Y., vol. 1,
p. 300. [Ety. *kleidos*, a clavicle; *phoros*,
bearing.] Equivalve, inequilateral;
hinge without teeth or crenulations;
cast marked by an oblique linear de-

pression extending from the anterior
cardinal margin toward the base, indi-
cating the existence of a clavicle as in
Solecortus; surface concentrically lined.
Type C. planulatus.

chicagoensis, S. A. Miller, 1880, Jour. Cin.
Soc. Nat. Hist., vol. 3, p. 314, Niag-
ara Gr.

concentricus, Hall, 1860, Can. Nat. and
Geo., vol. 5, p. 149, Low. Sil.
concentricus, Dawson, 1868. The name was
preoccupied.

cuneatus, Hall, 1860, Can. Nat. and Geo.,
vol. 5, p. 148, Low. Sil.

ellipticus, Ulrich, 1879, Jour. Cin. Soc.
Nat. Hist., vol. 2, p. 25, Hud. Riv. Gr.

elongatus, Hall, 1860, Can. Nat. and Geo.,
vol. 5, p. 150, Low. Sil.

erectus, Hall, 1860, Can. Nat. and Geo.,
vol. 5, p. 149, and Acad. Geol., p. 600,
Up. Sil.

erectus, Dawson, 1868. The name was
preoccupied.

faberi, n. sp. Shell
small, smooth,
subelliptical in
outline, length
greater than
height; anterior
end narrower
than the poste-

FIG. 795.—Clidophorus fa-
beri. Mag. 5 diam.

rior; basal mar-
gin a semi-
elliptic curve; beaks prominent, and
but little in advance of the middle;
umbonal slope rounded, and tapering
to the postero-basal margin of the
shell; cardinal line gently curving,
reaching the highest point posterior to
the middle of the shell; pallial line simple
and well defined; furrow deep, and ex-
tending from immediately in front of
the beaks to the pallial line.

Distinguished from C. fabula
by the more prominent
beaks and higher arch in
the cardinal line posterior to
the beaks and other minor
particulars; beside it is gen-
erally a larger shell, though
variable in size. Collected
in the upper part of the
Hud. Riv. Gr., near Ver-
sailles, Indiana, and in Butler
County, Ohio.

fabula, Hall, 1845, (Nucula fab-
ula,) Am. Jour. Sci.
and Arts, vol. 48,
p. 295, and Ohio
Pal., vol. 1, p. 138,
Hud. Riv. Gr.

FIG. 796.—Clidophorus
maccheneyanus, syn.
for Modiopsis
recta.

major, Ulrich, 1879,
Jour. Cin. Soc. Nat. Hist., vol. 2, p. 25,
Hud. Riv. Gr.

neglectus, Hall, 1862, Geo. Rep. Wis., p.
55, Hud. Riv. Gr.

nuculiformis, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 150, Up. Sil.

planulatus, Conrad, 1841, (Nuculites planulatus,) Ann. Rep. N. Y., p. 48, and Pal. N. Y., vol. 1, p. 300, Hud. Riv. Gr.

semiradiatus, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 150, Arisaig series of Up. Sil.

solenoides, see Solenopsis solenoides.

subovatus, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 151, Arisaig series of Up. Sil.

CLINOPISTHA, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 43. [Ety. *kline*, I lean; *opisthe*, backward.] Shell short, gibbous, subquadrate, beaks posterior, and muscular impressions immediately behind the beaks; muscular impres-

FIG. 797.—*Clinopecten antiqua*.

sus near the margins of the valves; ligament external. Type *C. lævis*.

insularis, Walcott, 1885, (Dystactella insularis,) Monogr. U. S. Geo. Sur., vol. 8, p. 172, Devonian.

antiqua, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 67, and Ohio

Pal., vol. 1, p. 208, Corniferous Gr.

lævis, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 44, and Geo. Sur. Ill., vol. 5, p. 584, Coal Meas.

radiata, Hall, 1858, (Edmondia radiata,) Geo. Rep. Iowa, p. 716, Coal Meas.

subnasuta, Hall & Whitfield, 1872, (Tellinomya subnasuta,) 24th Rep. N. Y. Mus. Nat. Hist., p. 192, and Pal. N. Y., vol. 5, p. 512, Ham. Gr.

telliniformis, Hall, 1883, (Dystactella telliniformis,) Pal. N. Y., vol. 5, p. 513, Up. Held. Gr.

CONOCARDIUM, Bronn, 1835, Leth. Geo., vol. 1, p. 92. [Ety. *konos*, a cone; *kardia*, the heart.] Equivalve, very inequilateral, hemifusiform; beaks prominent, incurved close to the anterior end, which is broad, flattened, more or less truncate nearly at right angles to the straight hinge-line, which is prolonged as an abruptly contracted, slender, tubular wing from the dorsal part of the anterior face; body of the shell diminishing conoidally from the edge of the anterior face toward the posterior end, which is attenuated, roundly and widely gaping; substance of the shell very thick, of a minute quadrangular cellular tissue, with strong internal ribs radiating from the beak, and often smaller external ones, strongest anteriorly. Type *C. hibernicum*.

acadianum, Hartt, 1868, Acad. Geol., p. 304, Carb.

æquilaterale, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 18, and Bull. Am. Mus. Nat. Hist., p. 62, Warsaw Gr.

altum, Keyes, 1888, Proc. Acad. Nat. Sci. Phil., pl. xii, figs. 4a, 4b, Ham. Gr.

antiquum, Owen, 1852, (Pleurorhynchus antiqua,) Geo. Wis., Iowa, and Minn., pl. 2, fig. 19, Silurian.

attenuatum, Conrad, 1842, (Pleurorhynchus attenuatus,) Jour. Acad. Nat. Sci., vol. 8, p. 252, Up. Held. Gr.

bifarium, Winchell, 1856, Rep. Low. Peninsula Mich., p. 95, Ham. Gr.

blumenbachium, see Euchasma blumenbachii.

bovipedale, Winchell, 1862, Proc. Acad. Nat. Sci., p. 419, Marshall Gr.

carinatum, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 14, and Bull. Am. Mus. Nat. Hist., p. 59, Warsaw Gr.

catostomum, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 13, and Bull. Am. Mus. Nat. Hist., p. 58, Warsaw Gr.

concinnum, Hall, 1883, Pal. N. Y., vol. 5, pl. 68, figs. 26-27, Ham. Gr.

crassifrons, Conrad, 1842, (Pleurorhynchus crassifrons,) Jour. Acad. Nat. Sci., vol. 8, p. 252, Ham. Gr.

cuneatum, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 14, and Bull. Am. Mus. Nat. Hist., p. 60, Warsaw Gr.

cuneus, Conrad, 1840, (Pleurorhynchus cuneus,) Ann. Rep. N. Y., p. 206, and Pal. N. Y., vol. 5, pl. 67, figs. 21-32, Up. Held. Gr.

denticulatum, Hall, 1883, Pal. N. Y., vol. 5, pl. 68, figs. 24-25, Ham. Gr.

eboraceum, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 91, and Pal. N. Y., vol. 5, p. 412, Ham. Gr.

elegantulum, Billings, 1866, Catal. Sil. Foss., Antic., p. 53, Anticosti Gr.

emmetense, Winchell, 1866, Rep. Low. Peninsula Mich., p. 95, Ham. Gr.

immaturum, Billings, 1862, Pal. Foss., vol. 1, p. 41, Black Riv. Gr.

inceptum, Hall, 1859, Pal. N. Y., vol. 3, p. 491, Low. Held. Gr.

liratum, Hall, 1883, Pal. N. Y., vol. 5, pl. 68, figs. 28-29, Chemung Gr.

meekianum, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 15, and Bull. Am. Mus. Nat. Hist., p. 61, Warsaw Gr.

napoleonense, Winchell, 1862, Proc. Acad. Nat. Sci., p. 419, Marshall Gr.

nasutum, Hall, 1883, Pal. N. Y., vol. 5, pl. 67, figs. 12-20, Schoharie grit.

nevadense, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 177, Devonian.

niagarensis, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 97, Niagara Gr.

normale, Hall, 1883, Pal. N. Y., vol. 5, pl. 68, figs. 17-19, Ham. Gr.

obliquum, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 249, and Geo. Sur. Ill., vol. 6, p. 529, Coal Meas.

FIG. 798.—*Clinopecten radiata*.

FIG. 799.—*Concocardium subtrigonale*. Side view.

ohloense, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 65, and Ohio Pal., vol. 1, p. 203, Corniferous Gr.
ornatum, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 111, Niagara Gr.
parrishi, Worthen, (in press.) Geo. Sur. Ill., vol. 8, p. 112, Up. Coal Meas.
prattenanum, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 15, and Bull. Am. Mus. Nat. Hist., p. 61, Warsaw Gr.
pulchellum, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 5, p. 299, Kinderhook Gr.
reluquum, Hall, 1883, Pal. N. Y., vol. 5, pl. 68, fig. 33, Chemung Gr.
rugosum, Hall, 1883, Pal. N. Y., vol. 5, pl. 68, fig. 32, Ham. Gr.
subtrigonale, D'Orbigny, 1850, Prodr. Paleont., t. 1, p. 80, Up. Held. Gr. Proposed instead of *C. trigonale*, Hall, 1843, Geo. Rep. 4th Dist., N. Y., p. 171, which was preoccupied.

FIG. 801.—*Concocardium subtrigonale*. a, Shows part of the alation; b, points to the posterior hiatus.

tegulum, Hall, 1883, Pal. N. Y., vol. 5, pl. 68, figs. 30-31, Niagara Gr.
trigonale, Phillips, 1836, (Pleurorhynchus trigonale), Geol. Yorkshire, p. 211, Devonian.
trigonale, Hall, see *C. subtrigonale*.
ventricosum, Hall, 1860, 13th Rep. N. Y. Mus. Nat. Hist., p. 91, Ham. Gr.
vomer, Conrad, 1842, (Pleurorhynchus vomer), Jour. Acad. Nat. Sci., vol. 8, p. 253, Devonian.
CRENIPECTEN, Hall, 1883, Pal. N. Y., vol. 5, p. 3. (Plates and Explanations.) [Ety. *crena*, notch; *Pecten*, a genus.] In form like *Aviculopecten*, but the hinge is furnished with a series of small cartilage pits throughout its entire length. Type *C. crenulatus*.
amplus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 81, Chemung Gr.

crenulatus, Hall, 1843, (Pecten crenulatus), Geo. Sur. 4th Dist. N. Y., p. 265, and Pal. N. Y., vol. 5, pt. 1, p. 82, Chemung Gr.
glaber, Hall, 1843, (Lima glabra), Geo. Sur. 4th Dist. N. Y., p. 255, and Pal. N. Y., vol. 5, pt. 1, p. 85, Chemung Gr.
hallanus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 231, Subcarboniferous.
impolitus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 83, Chemung Gr.

FIG. 802.—*Crenipecten crenulatus*. FIG. 803.—*Crenipecten crenulatus*. Hinge-line.

leon, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 88, Chemung Gr.
liratus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 87, Chemung Gr.
micropterus, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 86, Chemung Gr.
obsoletus, Hall, 1843, (Lima obsoleta), Geo. Sur. 4th Dist., N. Y., p. 265, and Pal. N. Y., vol. 5, pt. 1, p. 84, Chemung Gr.

FIG. 804.—*Crenipecten retiferus*.

retiferus, Shumard, 1858, (Lima retifera), Trans. St. Louis Acad. Sci., vol. 1, p. 214, and Geo. Sur. Ill., vol. 5, p. 588, Coal Meas.

winchelli, Meek, 1875, (Aviculopecten winchelli), Ohio Pal., vol. 2, p. 296, Waverly Gr.
Ctenodonta, Salter, 1851, syn. for Tellinomya.
abrupta, see Tellinomya abrupta.
angela, see Tellinomya angela.
astartiformis, see Tellinomya astartiformis.
contracta, see Tellinomya contracta.
gibberula, see Tellinomya gibberula.
hartsvillensis, see Tellinomya hartsvillensis.
hubbardi, syn. for Nuculites sulcatus.
iphigenia, see Tellinomya iphigenia.
logani, see Tellinomya logani.
Cucullæa, Lamarck, 1801, Syst. An. [Ety. *Cucullus*, a hood.] Not a Paleozoic genus.
opima, Hall, 1843, syn. for Nucula lirata.
CUNEAMYA, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 90. [Ety. *cuneus*, a wedge; *Mya*, a genus.] Shell large, equivalve, inequilateral ventricose; beaks prominent, incurved; cardinal line straight, ligament external; lunule and escutcheon

eon; pallial line simple. Type *C. miamiensis*.

coriformis, n. sp. Shell large, having a length in some specimens of three inches, and a height of two inches; larger at the anterior end, and coniformly tapering to the posterior point; beaks large, high, pointed and inrolled above the cardinal line; cardinal line straight from the top of the lunule three-fourths of the length of the shell, the posterior part forming a wing-like appendage of the shell; escutcheon dis-

FIG. 805.—*Cuneamya coriformis*. Right valve, below medium size.

tinged and well marked; lunule heart-shaped, very large, wide and deep, margins angular; the anterior end of the shell rapidly slopes backward from the lower extremity of the lunule to the basal line; an obtuse angle is formed at the base of the lunule (this is better shown in the illustrations by the figure of the right valve than by the anterior view); a cincture or furrow, arising at the point of the beaks, and very gradually widening, reaches the basal line anterior to the middle of the shell; anterior umbonal ridge very prominent; posterior umbonal slope prominent, flattened on the outer face so as to form an obtuse angle toward posterior cardinal wing; basal line slightly curved, with a sinus at the cincture surface; concentrically lined. Distinguished from *C. miamiensis* by the remarkably large lunule, better defined cincture, and posterior cardinal wing.

FIG. 806.—*Cuneamya coriformis*. Anterior view, showing lunule.

Found in the Hudson River Group at Cincinnati, Ohio. The specific name is from the heart-shaped lunule. The specimen illustrated is from the collection of Charles Faber.

FIG. 807.—*Cuneamya miamiensis*. Right valve.

curta, Whitfield, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 138, Hud. Riv. Gr.

elliptica, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 317, Hud. Riv. Gr.

miamiensis, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 91, Hud. Riv. Gr.

FIG. 808.—*Cuneamya miamiensis*. Dorsal view.

neglecta, Meek, 1871, (Sedgwickia neglecta,) Proc. Acad. Nat. Sci. Phil., p. 325, and Ohio Pal., vol. 1, p. 142, Hud. Riv. Gr.

parva, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 316, Hud. Riv. Gr. *scapha*, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 92, Hud. Riv. Gr.

CYCLOCONCHA, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 231. [Ety. in allusion to the nearly circular form of the shell.] Equivalve, subequilateral, sub-circular, concentrically lined; cardinal teeth near the middle, with a long lateral tooth on each side. Type *C. mediocardinalis*.

FIG. 809.—*Cycloconcha mediocardinalis*.

mediocardinalis, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 231, Hud. Riv. Gr.

CYPRICARDELLA, Hall, 1858, Trans. Alb. Inst., vol. 4, p. 17. [Ety. diminutive of *Cypricardia*.]

Shell ovate, subelliptical or subquadrate, closed; surface concentrically striated; two cardinal teeth in right valve, one beneath the beak, triangular, the posterior one more slender, and turned obliquely backward, leaving a triangular pit for the tooth from the other valve; long, narrow groove in the anterior cardinal margin apparently for a projection from the left valve; posterior side beveled from above, edge thin, ligament external, occupying a deep cavity; muscular impressions distinct, shallow; pallial impression simple. Type *C. subelliptica*.

bellistriata, Conrad, 1842, (Microdon bellistriatus,) Jour. Acad. Nat. Sci., vol. 8, p. 247, Ham. Gr.

FIG. 810.—*Cypricardella bellistriata*.

81, Jour. Cin.
p. 317, Hud.

eld, 1875, Ohio
Riv. Gr.

s. Dorsal view.
edgwickia neg-
Sci. Phil., p.
1, p. 142, Hud.

Jour. Cin. Soc.
Hud. Riv. Gr.
1875, Ohio Pal.,
Gr.

874, Cin. Quar.
[Ety. in al-
cular form of the
quilateral, sub-
lined; cardinal
side, with a long
side. Type C.

dicardinalis.

iller, 1874, Cin.
p. 231, Hud.

8, Trans. Alb.

Cypricardella
striata.

valve, one be-
ar, the posterior
turned obliquely
angular pit for
er valve; long,
interior cardinal
a projection
terior side bev-
thin, ligament
deep cavity;
distinct, shal-
simple. Type

(Microdon bel-
Nat. Sci., vol. 8,

complanata, Hall, 1870, (Microdon com-
planatus.) Prelim. Notice Lam. Shells,
p. 33, and Pal. N. Y., vol. 5, pl. 42, fig.
22, and pl. 74, figs. 14 to 19, Ham. Gr.
connata, Walcott, 1885, Monogr. U. S. Geo.
Sur., vol. 8, p. 250, Subcarboniferous.
gregaria, Hall, 1870, (Microdon gregarius.)
Prelim. Not. Lam. Shells, p. 32, and Pal.
N. Y., vol. 5, pl. 73, figs. 1-6, and pl. 74,
figs. 1-4, Ham. Gr.

macrostriata, Walcott, 1885, Monogr.
U. S. Geo. Sur., vol. 8, p. 180, De-
vonian.

major, Hall, 1885, Pal. N. Y., vol. 5, p.
307, Up. Held. Gr.

nucleata, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 17, and Geo. Sur. Iowa, p. 663,
Warsaw Gr.

oblonga, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 18, and Bull. Am. Mus. Nat.
Hist., p. 65, Warsaw Gr.

plicata, see Gonioptera plicata.

quadrata, White & Whitfield, 1862, Proc.
Bost. Soc. Nat. Hist., vol. 8, p. 300,
Kinderhook Gr.

reservata, Hall, 1870, (Microdon reserva-
tus.) Prelim. Notice Lam. Shells, p. 33,
and Pal. N. Y., vol. 5, pl. 74, figs. 11-13,
Waverly Gr.

subelliptica, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 17, and Geo. Sur. Iowa, p. 664,
Warsaw Gr.

tenuistriata, Hall, 1870, (Microdon tenu-
istriatus.) Prelim. Notice Lam. Shells,
p. 32, and Pal. N. Y., vol. 5, pl. 73, figs.
23 to 30, and pl. 74, figs. 20, 21, Ham. Gr.

CYPRICARDIA, Lamarck, 1801, Syst. An. sans
Vert. [Ety. from the two genera Cy-
prina and Cardium.] Oblong, oblique
posterior ridge; umbones anterior, de-
pressed; ligament external, in deep,
narrow grooves; cardinal teeth two,
lateral one, in each valve, sometimes
obscure; muscular impressions two,
oval, placed below the extreme anterior
and posterior ends of the cardinal line;
pallial line simple. Typical C. obesa.

angusta, see Cypricardites angustus.

angustata, Vanuxem, syn. for Amnigenia
catskillensis.

choteauensis, Swallow, 1863, Trans. St.
Louis Acad. Sci., vol. 2, p. 96, Wav-
erly or Choteau Gr.

contracta, see Sphenotus contractus.

indianensis, see
Cypricardina
indianensis.

insecta, Dawson,
1868, Acad.
Geol., p. 303,
Carboniferous.

leidy, Lea, see
Leaia, leidy.
obsoleta, see
Cypricardites
obsoletus.

occidentalis, Hall, 1852, Stans. Ex. to
Great Salt Lake, p. 412, Coal Meas.

FIG. 811.—Cypricardina obesa.

occidentalis, Swallow, 1863, Trans. St.
Louis Acad. Sci. This name was pre-
occupied. See U. swallowana.

pikensis, Swallow, 1863, Trans. St. Louis
Acad. Sci., vol. 2, p. 95, Coal Meas.

plicatula, Swallow, 1858, Trans. St. Louis
Acad. Sci., vol. 4, p. 205, Mid. Coal
Meas.

primigenia, see Modiolopsis primigenia.

randolphensis, see Sanguinolites randolph-
ensis.

rhombica, see Cytherodon rhombens.

rigida, see Sphenotus rigidus.

shumardana, Swallow, 1863, Trans. St.
Louis Acad. Sci., vol. 2, p. 95, St. Gen-
evieve limestone.

subplana, see Edmondia subplana.

swallowana, S. A. Miller, 1883, 2d Ed. Am.
Pal. Foss., p. 310, Coal Measures of
Harrison County, Missouri. Proposed
instead of C. occidentalis, Swallow,
1863, in Trans. St. Louis Acad. Sci.,
p. 96.

undulata, Gurley, 1883, New. Carb. Foss.,
p. 3, Coal Meas. Publication in-
valid.

ventricosa, Hall, 1860, 13th Rep. N. Y.
Mus. Nat. Hist., p. 110, Kinderhook Gr.

wheeleri, see Schizodus wheeleri.

CYPRICARDINIA, Hall, 1859, Pal. N. Y., vol.
3, p. 266. [Ety. Cypricardina, from its
resemblance to Cypricardia.] General
form of Cypricardia; inequilateral;
oblique posterior ridge; umbones an-
terior, elevated; concentrically grooved,
sometimes cancellated; postero-cardinal
margin sometimes alate. Type C. lamel-
losa.

arcuata, Hall, 1885, Pal. N. Y., vol. 5,
p. 486, Chemung Gr.

arata, Hall,
1867, 20th
Rep. N. Y.
Mus. Nat.
Hist., p. 385,
Niagara Gr.

carbonaria,

Meek, 1871, FIG. 812.—Cypricardina dis-
tincta. Left valve, long
specimen.

Nat.Sci.Phil.,
p. 163, and Ohio Pal., vol. 2, p. 342,
Coal Meas.

concentrica, Hall, 1859, Pal. N. Y., vol. 3,
p. 268, Low. Held. Gr.

consimilis, Hall,
1885, Pal. N. Y.,
vol. 5, p. 486,
Waverly Gr.

crassa, Hall, 1859,
Pal. N. Y., vol. 3,
p. 268, Low. Held.
Gr.

(?) cylindrica, Hall
& Whitfield, 1872,
24th Rep. N. Y. Mus. Nat. Hist., p. 190,
Carboniferous Gr.

distincta, Billings, 1874, Pal. Foss., vol. 2,
p. 56, Gaspe limestone, No. 8, De-
vonian.

FIG. 813.—Cypricardi-
nia distincta. Left
valve, short spec-
imen.

- dornata*, Hall, 1859, Pal. N. Y., vol. 3, p. 267, Low. Held. Gr.
- indenta*, Conrad, 1842, (Cypricardites indentus,) Jour. Acad. Nat. Sci., vol. 8, p. 244, Up. Held. Gr.
- indianensis*, Hall, 1858, (Cypricardia indianensis,) Trans. Alb. Inst., vol. 4, p. 18, and Bull. Am. Mus. Nat. Hist., p. 58, Warsaw Gr.
- inflata* var. *subaequalis*, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 189, U. Held. Gr.
- lamellosa*, Hall, 1859, Pal. N. Y., vol. 3, p. 266, Up. Held. Gr.
- planulata*, Conrad, 1842, (Pterinea planulata,) Jour. Acad. Nat. Sci., vol. 8, p. 251, and Pal. N. Y., vol. 5, p. 484, Low. Held. Gr.
- sublamellosa*, Hall, 1859, Pal. N. Y., vol. 3, p. 267, Low. Held. Gr.
- subovata*, Miller & Dyer, 1878, Cont. to Pal. No. 2, p. 10, Niagara Gr.
- sulcifera*, Winchell, 1863, (Sanguinolites sulcifera,) Proc. Acad. Nat. Sci. Phil., p. 14, and Pal. N. Y., vol. 5, p. 487, Waverly Gr.
- CYPRICARDITES**, Conrad, 1841, Ann. Geo. Rep. N. Y., p. 51. [Ety. from resemblance to the genus *Cypricardia*.] Equivalve, profoundly inequilateral; external flattened ligamental area; hinge with four or five short oblique cardinal teeth; anterior one largest and most prominent; lateral teeth two, short and remote from the cardinal teeth; two muscular scars; surface concentrically lined with marks of growth. Type *C. curtus*. If the genus can stand, it must be based on this type (all other species are referred to other genera), because this species alone has a hinge-line like the one Conrad made.
- acutumbonus*, Billings, 1866, (Cyrtodonta acutumbona,) Catal. Sil. Foss. Antic., p. 49, Anticosti Gr.
- alta*, see *Modiomorpha alta*.
- alveatus*, Conrad, 1843, Geo. Rep. 3d Dist. N. Y., Ham. Gr.
- amygdalinus*, Hall, 1847, (Ambonychia amygdalina,) Pal. N. Y., vol. 1, p. 165, Black Riv. and Trenton Grs.
- angustus*, Hall, 1843, (Cypricardia angusta,) Geo. Rep. 4th Dist. N. Y., p. 76, Clinton Gr.
- angustatus*, syn. for *Amnigenia catskillensis*.
- angustifrons*, syn. for *Modiolopsis modiolaris*.
- anodontoides*, see *Modiolopsis anodontoides*.
- anticostiensis*, Billings, 1866, (Cyrtodonta (?) anticostiensis,) Catal. Sil. Foss. Antic., p. 14, Hud. Riv. Gr.
- bisulcata*, see *Grammysia bisulcata*.
- breviusculus*, Billings, 1859, (Cyrtodonta breviuscula,) Can. Nat. and Geo., vol. 4, p. 446, Chazy Gr.
- canadensis*, Billings, 1858, (Cyrtodonta canadensis,) Can. Nat. and Geo., vol. 3, p. 434, Black Riv. and Trenton Grs.
- carinatus*, see *Goniophora carinata*.
- carinatus*, Meek, 1872, (Dolabra carinata,) Proc. Acad. Nat. Sci. Phil., p. 326, and Ohio Pal., vol. 1, p. 135, Hud. Riv. Gr. This name was preoccupied.
- cariniferus*, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 246, syn. for *Goniophora chemungensis*.
- catskillensis*, see *Amnigenia catskillensis*.
- chemungensis*, see *Goniophora chemungensis*.
- concentrica*, see *Modiomorpha concentrica*.
- cordiformis*, Billings, 1858, (Cyrtodonta cordiformis,) Can. Nat. and Geo., vol. 3, p. 437, Black Riv. and Trenton Grs.
- corrugatus*, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 244, Ham. Gr.
- curtus*, Conrad, 1841, Ann. Rep. N. Y., p. 53, Hud. Riv. Gr.
- elongatus*, see *Cimitaria elongata*.
- emma*, Billings, 1862, (Cyrtodonta emma,) Pal. Foss., vol. 1, p. 150, Hud. Riv. Gr.
- ferrugineus*, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 116, Clinton Gr.
- ganti*, Safford, 1869, (Cyrtodonta ganti,) Geo. of Tenn., p. 287, Trenton and Hud. Riv. Grs.
- hainesi*, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 147, Hud. Riv. Gr.
- harrietta*, Billings, 1862, (Cyrtodonta harrietta,) Pal. Foss., vol. 1, p. 149, Hud. Riv. Gr.
- haynana*, Safford, 1869, (Cyrtodonta haynana,) Geo. of Tenn., p. 287, Trenton and Hud. Riv. Grs.
- hindi*, Billings, 1862, (Cyrtodonta hindi,) Pal. Foss., vol. 1, p. 151, Hud. Riv. Gr.
- huronensis*, Billings, 1858, (Cyrtodonta huronensis,) Can. Nat. and Geo., vol. 3, p. 432, Black Riv. and Trenton Grs.
- indentus*, see *Cypricardia indenta*.
- inflatus*, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 246, syn. for *Cypricardia indenta*.
- inflatus*, Emmons, 1842, (Nuculites inflatus,) Geo. Rep. N. Y., p. 395, Trenton Gr.
- insularis*, Billings, 1866, (Cyrtodonta insularis,) Catal. Sil. Foss. Antic., p. 14, Hud. Riv. Gr.
- islandicus*, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 189. Proposed instead of *Cypricardites ventricosus*, Hall, 1859, which was preoccupied, Low. Held. Gr.

FIG. 814. — *Cypricardites* hainesi. Left valve.

FIG. 815. — *Cypricardites* hainesi. Interior of left valve, two teeth injured at.

carinata.
Dolabra carinata,
 Phil., p. 326, and
 Hud. Riv. Gr.
 pied.
 Jour. Acad. Nat.
 for Gonolophora
 catskillensis.
 shora chemung-
 pha concentrica.
 8, (Cyrtodonta
 and Geo., vol. 3,
 Trenton Grs.
 Jour. Acad. Nat.
 Gr.
 Rep. N. Y., p.
 ngata.
 cyrtodonta emma,
 Hud. Riv. Gr.
 field, 1875, Ohio
 ton Gr.
 cyrtodonta ganti,
 7, Trenton and
 374. Cin. Quar.
 7, Hud. Riv. Gr.
 2, (Cyrtodonta
 vol. 1, p. 149,

 14. — Cypricardites
 esi. Left valve.
 58, (Cyrtodonta
 and Geo., vol. 3,
 Trenton Grs.
 ia indenta.
 inflatus, Conrad,
 1842, Jour.
 Acad. Nat. Sci.,
 vol. 8, p. 246,
 syn. for Cypricardina in-
 denta.
 inflatus, Em-
 mons, 1842,
 (Nuculites in-
 flatus,) Geo.
 Rep. N. Y., p.
 395, Trenton Gr.
 (Cyrtodonta in-
 flatus, Antic., p. 14,
 st Ed. Am. Pal.
 used instead of
 us, Hall, 1859,
 Low. Held. Gr.

lowensis, Owen, 1840, (Cardium lowense.)
 Rep. on Mineral lands, pl. 17, fig. 8,
 Calciferous Gr.
 latus, Hall, 1847, (Modiolopsis latus,) Pal.
 N. Y., vol. 1, p. 160, Trenton Gr.
 leucothea, Billings, 1862, (Cyrtodonta
 leucothea,) Pal. Foss., vol. 1, p. 46, Black
 Riv. Gr.
 marcellensis, see Lunulicardium marcel-
 lense.
 megambonus, Whitfield, 1878, Ann. Rep.
 Geo. Sur. Wis., p. 73, and Geo. Wis.,
 vol. 4, p. 210, Trenton Gr.
 modiolaris, Emmons, syn. for Modiolopsis
 nasuta.
 mytiloides, see Modiomorpha mytiloides.
 nasuta, see Modiolopsis nasuta.
 niota, Hall, 1861, Geo. Rep. Wis., p. 29,
 and Geo. Wis., vol. 4, p. 208, Tren-
 ton Gr.
 obliquus, Meek & Worthen, 1868, Geo.
 Sur. Ill., vol. 3, p. 311, Galena Gr.
 oblongus, Conrad, syn. for Modiomorpha
 concentrica.
 obsoletus, Hall, 1843, (Cypricardina obso-
 leta,) Geo. Rep. 4th Dist. N. Y., pl. 8,
 fig. 3, Clinton Gr.
 obtusus, Hall, 1847, (Ambonychia ob-
 tusa,) Pal. N. Y., vol. 1, p. 167, Black
 Riv. and Trenton Gr.
 ovata, syn. for Modiolopsis modiolaris.
 plebeius, Billings, 1866, (Cyrtodonta ple-
 beia,) Catal. Sil. Foss., Antic., p. 14,
 Hud. Riv. Gr.
 ponderosus, Billings, 1862, (Cyrtodonta
 ponderosa,) Pal. Foss., vol. 1, p. 150,
 Hud. Riv. Gr.
 quadrangularis, Whitfield, 1878, Jour.
 Cin. Soc. Nat. Hist., vol. 1, p. 138, Hud.
 Riv. Gr.
 quadrilateralis, Hall, 1867, 20th Rep. N. Y.
 Mus. Nat. Hist., p. 388, Niagara Gr.
 radiatus, Conrad, 1841, Ann. Rep. N. Y.,
 p. 53, Ham. Gr. Not recognized.
 rectus, Conrad, 1841, Ann. Rep. N. Y., p.
 52, Up. Held. Gr.
 rectirostris, Hall, 1861, Geo. Rep. Wis., p.
 29, Trenton Gr.
 recurvus, see Cimitaria recurva.
 rotundatus, Hall, 1861, Geo. Rep. Wis.,
 p. 29, and Geo. Wis., vol. 4, p. 208,
 Trenton Gr.
 rugosus, Billings, 1858, (Cyrtodonta ru-
 gosa,) Can. Nat. and Geo., vol. 3, p. 432,
 Black Riv. Gr.
 rugosus, see Gonolophora rugosa.
 saffordi, Hall, 1852, (Palaearca saffordii,) 12th
 Rep. N. Y. Mus. Nat. Hist., p. 11, and
 Geo. of Tenn., p. 287, Low. Held. Gr.
 sectifrons, see Phthonia sectifrons.
 sigmoideus, Billings, 1858, (Cyrtodonta
 sigmoidea,) Can. Nat. and Geo., vol. 3,
 p. 438, Black Riv. Gr.
 sinuatus, see Modiolopsis sinuata.
 spiniferus, Billings, 1858, (Cyrtodonta spi-
 nifera,) Can. Nat. and Geo., vol. 3, p.
 435, Black Riv. Gr.
 sterlingensis, Meek & Worthen, 1866,
 (Dolabra sterlingensis,) Proc. Acad.

Nat. Sci. Phil., p. 260, and Geo. Sur.
 Ill., vol. 3, p. 339, Hud. Riv. Gr.
 subalatus, see Modiomorpha subalata.
 subangulatus, Hall, 1847, (Edmondia sub-
 angulata,) Pal. N. Y., vol. 1, p. 156,
 Black Riv. and Trenton Grs.
 subcarinatus, Billings, 1858, (Cyrtodonta
 subcarinata,) Can. Nat. and Geo., vol.
 3, p. 433, Black Riv. Gr.
 subspatulatus, Hall, 1847, (Modiolopsis
 subspatulata,) Pal. N. Y., vol. 1, p. 159,
 Black Riv. and Trenton Grs.
 truncatus, see Sphenotus truncatus.
 unguatus, Billings, 1866, (Cyrtodonta un-
 gulata,) Catal. Sil. Foss. Antic., p. 15,
 Hud. Riv. Gr.
 ventricosus, Hall, 1847, (Edmondia ven-
 tricosus,) Pal. N. Y., vol. 1, p. 155, Tren-
 ton Gr.
 ventricosus, Hall, 1859, (Palaearca ventri-
 cosa,) Pal. N. Y., vol. 3. This name
 was preoccupied. See Cypricardites is-
 landicus.
 vetustus, Hall, 1847, (Cardiomorpha ve-
 tusta,) Pal. N. Y., vol. 1, p. 154, Tren-
 ton Gr.
 winchelli, Safford, 1869, (Cyrtodonta win-
 chelli,) Geo. Tenn., p. 287, Trenton and
 Hud. Riv. Grs.
 Cyrtodonta, syn. for Cypricardites.
 acutumbona, see Cypricardites acutum-
 bonus.
 anticostiensis, see C. anticostiensis.
 breviscula, see C. brevisculus.
 canadensis, see C. canadensis.
 cordiformis, see C. cordiformis.
 emma, see C. emma.
 ganti, see C. ganti.
 harrietta, see C. harrietta.
 hayniana, see C. haynana.
 hindi, see C. hindi.
 huronensis, see C. huronensis.
 insularis, see C. insularis.
 leucothea, see C. leucothea.
 normanensis, Safford. Not defined.
 plebeia, see Cypricardites plebeius.
 ponderosa, see C. ponderosus.
 rugosa, see C. rugosus.
 saffordi, see C. saffordi.
 sigmoidea, see C. sigmoideus.
 spinifera, see C. spiniferus.
 subcarinata, see C. subcarinatus.
 unguata, see C. unguatus.
 winchelli, see C. winchelli.
 CYTHEROIDON, Hall & Whitfield, 1873, in 23d
 Rep. N. Y., pl. 14, figs. 19-21. [Ety.
 Cythere, a genus; odous, tooth.] Ovate,
 pointed posteriorly; beaks pointed;
 sharp, oblique, umbonal ridge; cardinal
 line short; subcircular anterior and
 posterior muscular scars distinct; hinge
 area strong with angular teeth or crenu-
 lations beneath the beaks, pallial line
 simple, surface concentrically lined.
 Type C. nasutus.
 appressus, Conrad, (Nuculites appressus,)
 1842, Jour. Acad. Nat. Sci., vol. 8, p.
 248, and Pal. N. Y., vol. 5, pl. 75, figs.
 3-9, Ham. Gr.

chemungensis, Conrad, 1842, (*Nuculites chemungensis*.) Jour. Acad. Nat. Sci., vol. 8, p. 247, and Pal. N. Y., vol. 5, pl. 75, figs. 37-40, Chemung Gr.
cuneus, Hall, 1883, Pal. N. Y., vol. 5, pl. 75, figs. 27-30, Waverly Gr.

FIG. 816.—*Cytherodon rhombeus*.

ellipticus, Hall, 1870, (*Schizodus ellipticus*.) Prelim. Notice Lam. Shells, p. 96, and Pal. N. Y., vol. 5, pl. 75, figs. 13-15, Ham. Gr.
gregarius, Hall, 1883, Pal. N. Y., vol. 5, pl. 75, figs. 41-45, Chemung Gr.
nasutus, Hall, 1883, Pal. N. Y., vol. 5, pl. 75, figs. 10-12, Ham. Gr.
oblatus, Hall, Pal. N. Y., vol. 5, pl. 75, figs. 41-45, Chemung Gr.
pauper, Hall, 1883, Pal. N. Y., vol. 5, pl. 75, figs. 24-26, Chemung Gr.
 (?) *placidus*, Billings, 1874, Pal. Foss., vol. 2, p. 137, Up. Sil.
quadrangularis, Hall, 1870, (*Schizodus quadrangularis*.) Prelim. Notice Lam. Shells, p. 96, and Pal. N. Y., vol. 5, pl. 75, figs. 31-36, Chemung Gr.
rhombeus, Hall, 1843, (*Cypriocardia rhombea*.) Geo. Rep. 4th Dist. N. Y., p. 291, and Pal. N. Y., vol. 5, pl. 75, figs. 19-23, Subcarboniferous.
socialis, Billings, 1874, Pal. Foss., vol. 2, p. 138, Up. Sil.
tumidus, Hall, 1870, (*Schizodus tumidus*.) Prelim. Notice Lam. Shells, p. 94, and Pal. N. Y., vol. 5, pl. 75, figs. 1-2, Up. Held. Gr.

FIG. 817.—*Dextlobia ovata*.

beak incurved forward; left valve less inflated; hinge-line having a thickened cartilage plate, bearing a linear posterior groove. Type *D. ovata*.

DEXIOBIA, Winchell, 1863, Proc. Acad. Nat. Sci., p. 10. [Ety. *dexios*, on the right side; *bia*, strength.] Inequivalve, inequilateral, area undefined, right valve very ventricose, umbo prominent;

halli, Winchell, 1863, Proc. Acad. Nat. Sci., p. 11, Marshall Gr.
ovata, Hall, 1858, (*Cardiomorpha ovata*.) Geo. Rep. Iowa, p. 522, Kinderhook Gr.
whitlii, Winchell, 1863, Proc. Acad. Nat. Sci., p. 11, Marshall Gr.

DOLABRA, McCoy, 1844, Syn. Carb. Foss. Ireland, p. 64. [Ety. *dolabra*, a mallet or pickaxe.] Obliquely ovate, gibbous; left valve larger than the right; beaks large, obtuse, nearer the anterior than posterior end; hinge-line straight, shorter than the shell, not crenulated; a flat, narrow ligamental area the length of the hinge-line, widest between the beaks; anterior end narrower than the posterior, rounded; no byssal sinus or furrows; ventral margin slightly convex; posterior end obliquely truncated, slope flattened; surface smooth or finely striated.

alpina, Hall, 1858, Geo. Rep. Iowa, p. 716, Coal Meas.

carinata, see *Cypriocardites carinatus*.
sterlingensis, see *Cypriocardites sterlingensis*.

FIG. 818.—*Ectenodesma birostratum*.

Dystactella, Hall, 1883, Pal. N. Y., vol. 5, p. 4, (Plates and Explanations,) synonym for *Clinopiastha*.

insularis, see *Clinopiastha insularis*.

subnasuta, see *Clinopiastha subnasuta*.

telliniformis, see *Clinopiastha telliniformis*.

ECTENODESMA, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 4. (Plates and Explanations.)

[Ety. *ektēnos*, stretched out; *dema*, a ligament.] Body ovate, oblique; height greater than length; both valves more or less convex; hinge-line longer than the length of the shell; byssal sinus shallow; oblique lateral tooth; ligamental area narrow, striated; surface rayed; distinguished from *Glyptodesma* by having the anterior wing more produced, and both wings more acute at their extremities. Type *E. birostratum*.
birostratum, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 242, Chemung Gr.

EDMONDIA, DeKoninck, 1844, Desc. Anim. Foss., Carb. Belg., p. 66. [Ety. proper

name.] Shell equivalve, inequilateral, tumid, short, oblong or rounded, closed all around; dorsal and ventral margins slightly convex; beaks tumid, with an impressed lunette between them; surface with concentric striae; notec.n, but an internal lamellar cartilage support, much dilated within the cavity of the beaks, the broad end forming the slits in casts coinciding with the edges of the anterior lunette, and the posterior end running nearly parallel to and close within the hinge-line; dorsal margins erect and simple; two simple adductor impressions, often with an accessory impression over each, pallial scar simple, entire. Type *E. unioniformis*.

FIG. 819.—*Edmondia aspenwallensis*. Right valve.

aquimargin-
alis, Win-
chell, 1862,
(*Cardinia*
aquima r-
ginalis,) *P*
ro c.
Acad. Nat.
Sci., p. 413,
Marshall Gr.

anomala,
Dawson,

1867, Acad. Geo., p. 303, Carb.
aspenwallensis, Meek, 1871, Hayden's
Rep. Sur. Wyoming, p. 299, and Pal. E.
Neb., p. 216, Coal Meas.
bicarinata, Winchell, 1863, Proc. Acad.
Nat. Sci., p. 13, Marshall Gr. Prof. Hall
regards this as a syn. for *Sanguinolites*
rigidus.

binumbonata, Winchell,
1862, Proc. Acad. Nat.
Sci., p. 414, Marshall Gr.
burlingtonensis, White &
Whitfield, 1862, Proc.
Bost. Soc. Nat. Hist., vol.
8, p. 301, and Pal. N. Y.,
vol. 5, p. 390, Kinder-
hook Gr.

calhouni, see *Pleurophorus*
calhouni.

circularis, Walcott, 1885,
Monogr. U. S. Geo. Sur.,
vol. 8, p. 246, Carbonif.
erous.

concentrica, see *Astartella*
concentrica.

depressa, Hall, 1870, Prelim. Notice Lam.
Shells, p. 91, and Pal. N. Y., vol. 5, pl.
64, fig. 32, Waverly Gr.

ellipsis, Hall, 1885, Pal. N. Y., vol. 5, p.
392, Waverly Gr.

elliptica, Winchell, 1863, Proc. Acad. Nat.
Sci., p. 13, Marshall Gr.

gibbosa, Swallow, 1858, Trans. St. Louis
Acad. Sci., vol. 1, p. 189, Permian Gr.

glabra, Meek, 1872, Pal. E. Neb. p. 214,
Coal Meas.

harti, Dawson, 1868, Acad. Geol., p. 303,
Carb.

FIG. 820.—*Edmondia aspenwallensis*. Cardinal view.

hawni, Swallow, 1858, Trans. St. Louis
Acad. Sci., vol. 1, p. 209, Coal Meas.

illinoisensis, Worthen, 1884, Bull. No. 2
Ill. St. Mus. Nat. Hist., p. 18, and Geo.

Sur. Ill., vol. 8, p. 122, Keokuk Gr.

ledoides, Winchell, 1860, Rep. Low. Pen-
insula Mich., p. 96, Ham. Gr.

mactroides, Winchell, 1860, Rep. Low.
Peninsula Mich., p. 96, Ham. Gr.

marionensis, Swallow, 1860, Trans. St.
Louis Acad. Sci., vol. 1, p. 654, Cho-
teau Gr.

medon, Walcott, 1884, Monogr. U. S. Geo.
Sur., vol. 8, p. 245, Subcarboniferous.

mortonensis, Geinitz, 1896, (Astarte mor-
tonensis,) Carb. und Dyas in Neb., p. 17,
Coal Meas.

nebraskensis, Geinitz, 1896, (Astarte ne-
braskensis,) Carb. und Dyas in Neb., p. 16,
and Pal. E. Neb., p. 214, Coal Meas.

nilesi, Winchell & Marcy, 1865, Proc.
Bost. Soc. Nat. Hist., p. 97, Niagara Gr.

nitida, Winchell, 1863, Proc. Acad. Nat.
Sci., p. 12, Marshall Gr.

nuptialis, Winchell, 1863, Proc. Acad. Nat.
Sci., p. 12, Marshall Gr.

obliqua, Hall, 1885, Pal. N. Y., vol. 5, p.
38, Chemung Gr.

otoensis, Swallow, 1858, Trans. St. Louis
Acad. Sci., vol. 1, p. 189, Permian Gr.

peroblonga, Meek & Worthen, 1890, Proc.
Acad. Nat. Sci. Phil., p. 249, and Geo.

Sur. Ill., vol. 5, p. 583, Coal Meas.

philipi, Hall, 1870, Prelim. Notice Lam.
Shells, p. 90, and Pal. N. Y., vol. 5, pl.
64, figs. 6-18, Chemung Gr.

pinonensis, Meek, 1877, U. S. Geo. Expl.
40th parallel, vol. 4, p. 46, Devonian.

radiata, see *Clinopistha radiata*.

reflexa, Meek, 1872, Pal. E. Neb., p. 213,
Coal Meas.

rhomboides, Hall, 1883, Pal. N. Y., vol. 5,
pl. 64, figs. 7-8, Chemung Gr.

semiorbiculata, Swallow, 1858, Trans. St.
Louis Acad. Sci., vol. 1, p. 190, Per-
mian Gr.

strigillata, Winchell, 1863, Proc. Acad. Nat.
Sci., p. 12, Marshall Gr.

subangulata, see *Cypriocardites subangu-*
latus.

subcarinata, Hall, 1885, Pal. N. Y., vol. 5,
pl. 64, fig. 31, Chemung Gr.

subnasuta, Hall, 1883, Pal. N. Y., vol. 5,
pl. 64, figs. 5-6, Chemung Gr.

subovata, Hall, 1885, Pal. N. Y., vol. 5, p.
389, Chemung Gr.

subplana, Hall, 1858, Cypriocardia sub-
plana, Trans. Alb. Inst., vol. 4, p. 19,
and Bull. Am. Mus. Nat. Hist., p. 60,
Warsaw Gr.

subtruncata, see *Cuneamya subtruncata*.

subtruncata, Meek, 1872, Pal. E. Neb., p.
215, Coal Meas.

tapetiformis, Meek, 1875, (E. tapesiformis,) *O*
Ohio Pal., vol. 2, p. 304, Waverly Gr.

tenuistriata, Hall, 1885, Pal. N. Y., vol. 5,
p. 393, Chemung Gr.

transversa, Hall, 1885, Pal. N. Y., vol. 5,
p. 389, Chemung Gr.

- undata*, see *Grammysia undata*.
undulata, Hall, 1870, Prelim. Notice Lam. Shells, p. 91, and Pal. N. Y., vol. 5, pl. 64, figs. 1-4, Chemung Gr.
unioniformis, Phillips, 1856, (Isocardia unioniformis,) Geol. Yorkshire, vol. 2, p. 209, and Geo. Sur. Ill., vol. 2, p. 346, Coal Meas.
varsoviensis, Worthen, 1884, Bull. No. 2 Ill. St. Mus. Nat. Hist., p. 18, and Geo. Sur. Ill., vol. 8, p. 121, Keokuk Gr.
ventricosa, see *Cypriocardites ventricosus*.
ELYMELLA, Hall, 1885, Pal. N. Y., vol. 5, p. 50. [Ety. *elymos*, a case.] Equivalve, inequilateral, ovate, elliptical; anterior end short, rounded; posterior end narrower, rounded; beaks closely incurved; umbo prominent; cardinal line short; umbonal slope prominent in the upper part, not defined below; surface concentrically lined. Type *E. nuculoides*.
fabalis, Hall, 1885, Pal. N. Y., vol. 5, p. 502, Ham. Gr.
levata, Hall, 1885, Pal. N. Y., vol. 5, p. 504, Ham. Gr.
nuculoides, Hall, 1885, Pal. N. Y., vol. 5, p. 503, Ham. Gr.
patula, Hall, 1885, Pal. N. Y., vol. 5, p. 505, Waverly Gr.
Entolium, Meek, 1865, Cal. Geo. Sur., vol. 2. [Ety. *entos*, inside; *leion*, smooth.] If synonymous with *Pernopecten*, then the latter has priority; but if distinct, then probably it is not a Paleozoic genus, as the type is from rocks of Jurassic age.
Eodon, Hall, 1877, 1st Ed. Am. Pal. Foss., p. 244. Proposed instead of *Microdon*, Conrad, which was preoccupied.
EOPTERIA, Billings, 1865, Pal. Foss., vol. 1, p. 221. [Ety. *eos*, dawn; *pteron*, a wing.] Prof. Billings said if *Euchasma* is the same as *Eopteria*, then he desired *Eopteria* to be withdrawn from science. Winged as in *Pterinea*, both valves equally convex and gaping; ligament external. Type *E. typica*.
 (?) *ornata*, Billings, 1865, Pal. Foss., vol. 1, p. 307, Quebec Gr.
richardsoni, Billings, 1865, Pal. Foss., vol. 1, p. 306, Quebec Gr.

FIG. 821.—*Eopteria richardsoni*.

- typica*, Billings, 1865, Pal. Foss., vol. 1, p. 221, Quebec Gr.
EUCHASMA, Billings, 1865, Pal. Foss., vol. 1, p. 360. [Ety. *eu*, well; *chasma*, a hollow.] Strongly convex, triangular, in-

equilateral, equivalve subcordiform, gaping, posterior extremity flattened, hinge short, ligament external. Type *E. blumenbachi*.

blumenbachi, Billings, 1859, (Conocardium blumenbachi,) Can. Nat. and Geo., vol. 4, p. 350, Quebec Gr.

EUCHONDRIA, Meek, 1874, Am. Jour. Sci., 3d series, vol. 7, p. 445.

Like *Aviculopecten* in form, but with an unsymmetrical subrostral cartilage pit and unequal crenulations on the two sides. Type *E. neglecta*.

neglecta, Geinitz, 1866,

FIG. 822.—*Euchondria neglecta*. Right valve enlarged two diameters.

(*Pecten neglectus*, Carb. and Dyas in Neb., p. 33, and Geo. Sur. Ill., vol. 5, p. 599, Coal Meas.)

EUMICROTUS, Meek, 1864, syn. for *Pseudomonotis*.

hawni, see *Pseudomonotis hawni*.

hawni var. *ovata*, see *Pseudomonotis hawni* var. *ovata*.

hawni var. *sinuata*, see *Pseudomonotis hawni* var. *sinuata*.

EUTHYDESMIA, Hall, 1885, Pal. N. Y., vol. 5, p. 32.

[Ety. *euthus*, straight; *desmia*, a ligament.] Equivalve, inequilateral, broadly subovate, with a subulate cardinal expansion; cardinal line straight; anterior end short; surface concentrically lined; hinge-line marked by a continuous ligamental groove. Type *E. subtextile*.

FIG. 824.—*Euthydesmia subtextile*.

subtextile, Hall, 1843, (Astarte subtextilis,) Geo. Sur. 4th Dist. N. Y., p. 245, Portage Gr.

Exochorhynchus, Meek, 1864, Pal. Up. Mo. [Ety. *exochos*, prominent; *rhynchus*, beak.] This name was suggested as a probable genus or subgenus to include *Sedgwickia altirostrata*.

FORDILLA, Barrande, 1881, Acephales. Etudes Loc. et Comp., pl. 361, and Bull. U. S. Geo. Sur., No. 30, p. 123.

A minute bilvalve, somewhat resembling a *Modiolopsis* or an *Orthonotella*. Type *F. troyensis*.

troyensis, Barrande, 1886, Bull. U. S. Geo. Sur., vol. 30, p. 125, Up. Taconic.

FIG. 825.—*Fordilla troyensis*.

subcordiform,
emity flattened,
external. Type

1859, (Conocar-
n. Nat. and Geo.,
ir.

FIG. 822.—Eu-
chondria, neg-
lecta. Right
valve enlarged
two diameters.

Pecten neglec-
tus.) Carb. and
Dyas in Neb., p.
33, and Geo.
sur. Ill., vol. 5, p.
339, Coal Meas.
syn. for Pseudo-

s hawni.
domonotis hawni

Pseudomonotis

824.—Euthydesma
subtextile.

straight; anterior
centrically lined;
a continuous lig-
E. subtextile.
tarte subtextilis.)
Y., p. 245, Port-

4, Pal. Up. Mo.
ment: rhynchos,
is suggested as a
genus to include

FORDILLA, Bar-
rande, 1881,
Acephales.
Etudes Loc.
et Comp.,
pl. 361, and
Bull. U. S.
Geo. Sur.,
No. 30, p. 123.
A minute bi-
nubling a Modio-
bellula. Type F.

, Bull. U. S. Geo.
p. Taconic.

Gervillia, DeFrance, 1820, Dict. Sci. Nat.,
xviii. [Ety. proper name.] Type *G.*
anceps. This genus is probably un-
known in the American Palæozoic
rocks.

auricula, see *Monopteria auricula*.

longa, see *Avicula longa*.

longispina, see *Monopteria longispina*.

strigosa, see *Pterinea strigosa*.

sulcata, see *Bakevellia sulcata*.

Glossites, Hall, 1885, Pal. N. Y., vol. 5, p.

49. [Ety. *glosse*, the tongue.] Equi-
valve, inequilateral, elliptical; anterior
end short, margin declining from the
beak and curving below; posterior end
large, broadly rounded; beaks small,
appressed; cardinal line long, gently
arcuate; umbonal slope not defined;
surface marked concentrically; ligament
external; lunule distinct; muscular im-
pressions shallow. Type *G. lingualis*.

amygdalinus, Winchell, 1863, (Sanguino-
lites amygdalinus.) Proc. Acad. Nat.
Sci. Phil., p. 13, and Pal. N. Y., vol. 5,
p. 501, Waverly Gr.

depressus, Hall, 1885, Pal. N. Y., vol. 5, p.

496, Chemung Gr.

ellipticus, Hall, 1885, Pal. N. Y., vol. 5, p.

498, Chemung Gr.

lingualis, Hall, 1885, Pal. N. Y., vol. 5, p.

497, Chemung Gr.

patulus, Hall, 1885, Pal. N. Y., vol. 5, p.

501, Chemung Gr.

procerus, Hall, 1885, Pal. N. Y., vol. 5, p.

499, Chemung Gr.

rudicula, Hall, 1885, Pal. N. Y., vol. 5, p.

498, Chemung Gr.

subnasutus, Hall, 1885, Pal. N. Y., vol. 5,

p. 500, Chemung Gr.

subtenuis, Hall, 1885, Pal. N. Y., vol. 5, p.

495, Ham. Gr.

teretis, Hall, 1885, Pal. N. Y., vol. 5, p.

494, Up. Held. Gr.

GLYPTOCARDIA, Hall, 1885,
Pal. N. Y., vol. 5, p. 35.

[Ety. *glyptos*, sculptured;
cardia, the heart.] Shell

small, equi-
valve, inequi-
lateral, broadly elliptical

or subcircular; beaks incurved; sur-
face plicated and marked with concentric
striae; no area beneath the beaks.

Type *G. speciosa*.

speciosa, Hall, 1843, (*Avicula speciosa*.)
Geo. Sur. 4th Dist. N. Y., p. 243, and
Pal. N. Y., vol. 5, p. 426, Ham. and
Portage Grs.

GLYPTODESMA, Hall, 1883, Pal. N. Y., vol. 5,
pt. 1, p. 4. (Plates and Explanations.)

[Ety. *glyptos*, sculptured; *dema*, a liga-
ment.] Aviculiiform, ligamental area

striated, continuous, hinge with two
strong lateral teeth, and numerous ir-
regular transverse plications along the
cardinal margin; surface concentrically
striated. Type *G. erectum*.

cruciforme, Conrad, 1841, (*Avicula cruci-
formis*.) Ann. Rep. N. Y., p. 54,
Ham. Gr.

erectum, Conrad, 1842, (*Avicula erecta*.)
Jour. Acad. Nat. Sci., vol.

8, p. 238, and Pal. N. Y.,
vol. 5, p. 153, Ham. Gr.

erectum var. *obliquum*, Hall,
1883, Pal. N. Y., vol. 5, pt.

1, p. 155, Ham. Gr.

occidentale, Hall, 1883, Pal.
N. Y., vol. 5, pt. 1, p. 157,
Up. Held Gr.

subrectum, Whitfield, 1882,
(*Actinodesma subrectum*.) Ann. N. Y.
Acad. Sci., vol. 2, p. 215, Ham. Gr.

FIG. 827.
Glyptodesma
erectum.

FIG. 828.—*Glyptodesma erectum*. Mold of left
valve, showing pallial line, muscular scar, tu-
bercles to the interpallial area representing
points of muscular attachment.

GONIOPHORA, Phillips, 1848, Mem. Geo. Sur.
Gt. Brit., vol. 2, p. 264. [Ety. *gonia*, an
angle; *phoros*, bearing.] *Goniophorus*

was used by Agassiz for a genus of
Echinoderms in 1840. Equi-
valve, very inequilateral, rhomboidal or trape-
zoidal, obliquely truncate behind,
rounded in front; cardinal line straight;
beaks small, umbo prominent, and slope
continued as a ridge to the post-inferior
margin; oblique, undefined sinus from
anterior to the beaks to basal margin;
surface concentrically lined; hinge
with an oblique fold or tooth in the
left valve beneath the beak, and a cor-
responding depression in the right
valve; ligament external, attached by
one or more grooves; anterior muscu-
lar impression deep, situated anterior
to the beak; posterior muscular im-
pression shallow, situated on the poste-
rior cardinal slope; pallial line simple.
Type *G. cymbiformis*.

acuta, Hall, 1870, (*Sanguinolites acutus*.)
Prelim. Notice Lam. Shells, p. 37, and
Pal. N. Y., vol. 5, pl. 43, figs. 1-3,
Ham. Gr.

alata, Hall, 1885, Pal. N. Y., vol. 5, p. 294,
Schoharie grit.

bellula, Billings, 1874, Pal. Foss., vol. 2, p.
136, Up. Sil.

carinata, Conrad, 1841, (Cypricardites *carinatus*.) Ann. Rep. Geo. Sur. N. Y., p. 53, and Pal. N. Y., vol. 5, pl. 44, figs. 6-8, Ham. Gr.
chemungensis, Vanuxem, 1842, Cypricardites *chemungensis*.) Geo. Rep. N. Y., p. 181, and Pal. N. Y., vol. 5, pl. 44, figs. 18-22, Chemung Gr.
consimilis, Billings, 1874, Pal. Foss., vol. 2, p. 135, Up. Sil.
crassa, Whiteaves, 1888, Pal. Foss., vol. 3, p. 9, Guelph Gr.
glabra, Hall, 1883, Pal. N. Y., vol. 5, pl. 44, figs. 9-17, syn. for *G. glaucus*.
glaucus, Hall, 1870, (Sanguinolites *glaucus*.) Prelim. Notice Lam. Shells, p. 38, and Pal. N. Y., vol. 5, p. 299, Ham. Gr.
hamiltonensis, Hall, 1870, (Sanguinolites *hamiltonensis*.) Prelim. Notice Lam. Shells, p. 30, and Pal. N. Y., vol. 5, pl. 43, figs. 8-21, Ham. Gr.
mediocria, Billings, 1874, Pal. Foss., vol. 2, p. 137, Up. Sil.
minor, Hall, 1885, Pal. N. Y., vol. 5, p. 305, Chemung Gr.
perangulata, Hall, 1870, (Sanguinolites *perangulatus*.) Prelim. Notice Lam. Shells, and Pal. N. Y., vol. 5, pl. 34, figs. 1-7, Up. Held. Gr.

FIG. 829.—*Gonolophora chemungensis*.

plicata, Hall, 1858, (Cypricardella *plicata*.) Trans. Alb. Inst., vol. 4, p. 18, and Bull. Am. Mus. Nat. Hist., p. 66, Warsaw Gr.
rugosa, Conrad, 1841, (Cypricardites *rugosus*.) Ann. Rep. N. Y., p. 53, and Pal. N. Y., vol. 5, p. 297, Ham. Gr.
speciosa, Hall, 1879, Desc. New. Spec. Foss., p. 17, and 11th Rep. Geol. Indiana, p. 317, Niagara Gr.
subrecta, Hall, 1885, Pal. N. Y., vol. 5, p. 304, Chemung Gr.
transiens, Billings, 1874, Pal. Foss., vol. 2, p. 134, Up. Sil.
trigona, Hall, 1885, Pal. N. Y., vol. 5, p. 302, Chemung Gr.
truncata, Hall, 1883, Pal. N. Y., vol. 5, pl. 44, fig. 15, Ham. Gr.
 GOSSELETTIA, Barrois, 1881, Ann. Soc. Geol. du Nord, vol. 8, p. 176. [Ety. proper name.] Shell subtriangular, truncate on the anterior side, subulate posteriorly; ligamental area wide, longitudinally striate; cardinal teeth below the beak strong; lateral teeth elongate; surface with concentric striae.

retusa, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 266, Ham. Gr.

FIG. 830.—*Gosselettia triquetra*. Right valve.

triquetra, Conrad, 1838, (Pterinea *triquetra*.) Ann. Rep. Geo. N. Y., p. 116, and Pal. N. Y., vol. 5, pt. 1, p. 265, Ham. Gr.

GRAMMYSIA, DeVerneuil, 1847, Bull. Soc. Geo. France, 2d ser., vol. 4, p. 696.

[Ety. *gramme*, a line of writing; *Mys*, a mussel shell, in allusion to the transverse furrows which cross the valve from the umbones to the middle of the ventral margin.] Equivalve, inequilateral; shell thick, oblong; anterior side short, contracted by a deep oval lunette beneath the beak; posterior end elliptically rounded; hinge-line straight; two large adductor impressions in each valve, anterior

rounded, posterior pear-shaped; pallial scar entire; cartilage external, short, in the anterior part of a deep depression formed by the inflexion of the hinge margins; an oblique furrow extends from the beak to about the middle of the ventral margin. Type *G. bisulcata*.
acadica, Billings, 1874, Pal. Foss., vol. 2, p. 140, Up. Sil.
alveata, Conrad, 1841, (Posidonia *alveata*.) Ann. Rep. N. Y., p. 53, and Pal. N. Y., vol. 5, pl. 57, figs. 1-2, and pl. 60, Ham. Gr.
arcuata, Conrad, 1841, (Posidonia *arcuata*.) Ann. Rep. N. Y., p. 53, and Pal. N. Y., vol. 5, p. 373, Ham. Gr.
bisulcata, Conrad, 1838, (Pterinea *bisulcata*.) 1841, (Cypricardites *bisulcata*.) Ann. Rep. N. Y., p. 116, and Pal. N. Y., vol. 5, pl. 51, figs. 1-18, Ham. Gr.
canadensis, Billings, 1874, Pal. Foss., vol. 2, p. 51, Gaspé sandstone, Up. Sil.
caswelli, Foerste, 1885, Bull. Sci. Lab. Denison University, p. 92. Not properly defined.

chemungensis, Pitt, 1874, Bul. Buff. Soc. Nat. Hist., Chemung Gr.
circularis, Hall, 1870, Prelim. Notice Lam. Shells, p. 51, and Pal. N. Y., vol. 5, pl. 57, figs. 3-6, Ham. and Chemung Grs.
communis, Hall, 1885, Pal. N. Y., vol. 5, p. 378, Chemung Gr.
constricta, Hall, 1870, Prelim. Notice Lam. Shells, p. 58, and Pal. N. Y., vol. 5, pl. 59, figs. 13-20, Ham. Gr.
cuneata, Hall, 1883, Pal. N. Y., vol. 5, pl. 62, figs. 1-9, Ham. Gr.
duplicata, Hall, 1885, Pal. N. Y., vol. 5, p. 380, Chemung Gr.
elliptica, Hall, 1870, Prelim. Notice Lam. Shells, p. 53, and Pal. N. Y., vol. 5, pl. 58, figs. 1-12, Chemung Gr.
erecta, Hall, 1870, Prelim. Notice Lam. Shells, p. 52, and Pal. N. Y., vol. 5, p. 363, Ham. Gr.
glabra, Hall, 1885, Pal. N. Y., vol. 5, p. 369, Chemung Gr.
globosa, Hall, 1870, Prelim. Notice Lam. Shells, p. 57, and Pal. N. Y., vol. 5, p. 372, Ham. Gr.
hamiltonensis, syn. for *G. bisulcata*.

FIG. 831.—*Grammysia hannibalensis*. Dorsal view.

lirata, Hall, 1870, Prelim. Notice Lam. Shells, p. 57, and Pal. N. Y., vol. 5, pl. 59, figs. 6-12, Ham. Gr.
magna, Hall, 1870, Prelim. Notice Lam. Shells, p. 50, and Pal. N. Y., vol. 5, p. 362, Ham. Gr.

minor, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 3, p. 174, Fig. 832.—*Grammysia hannibalensis*. Right side view.

nodocostata, Hall, 1870, Prelim. Notice Lam. Shells, p. 50, and Pal. N. Y., vol. 5, pl. 55, figs. 1-11, Ham. Gr.

obsoleta, Hall, 1870, Prelim. Notice Lam. Shells, p. 60, and Pal. N. Y., vol. 5, pl. 59, figs. 21-27, Ham. Gr.
ovata, Hall, 1885, Pal. N. Y., vol. 5, p. 358, Up. Held. Gr.

parallela, Hall, 1870, Prelim. Notice Lam. Shells, p. 59, Ham. Gr.

plena, Hall, 1885, Pal. N. Y., vol. 5, p. 382, Waverly Gr.

præcursor, Hall, 1870, Prelim. Notice Lam. Shells, p. 54, and Pal. N. Y., vol. 5, pl. 59, fig. 1, Schoharie grit.

remota, Billings, 1874, Pal. Foss., vol. 2, p. 139, Up. Sil.
rhomboidalis, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 248, and Geo. Sur. Ill., vol. 3, p. 439, Ham. Gr.

rhomboides, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 72, and Ohio Pal., vol. 2, p. 302, Waverly Gr.
rustica, Billings, 1874, Pal. Foss., vol. 2, p. 139, Up. Sil.
secunda, Hall, 1870, Prelim. Notice Lam. Shells, p. 54, and Pal. N. Y., vol. 5, pl. 59, figs. 2-5, Up. Held. Gr.
subarcuata, Hall, 1870, Prelim. Notice Lam. Shells, p. 61, and Pal. N., vol. 5, pl. 61, figs. 10-22, Chemung Gr.
undata, Hall, 1883, Pal. N. Y., vol. 5, p. 379, Chemung Gr.
ventricosa, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 73, and Ohio Pal., vol. 2, p. 303, Waverly Gr.
Gryphorhynchus, Meek, 1864, Am. Jour. Sci. and Arts. Not. defined.

ILIONIA, Billings, 1875, Can. Nat. and Geol., vol. 8, p. 301. [Ety. proper name.] Irregularly ovate, compressed; one extremity larger than the other, with beaks turned toward the larger end; concave depression from the umbones to the posterior ventral margin; sub-ovate muscular impression in the upper half of the posterior extremity. Type *I. canadensis*.

canadensis, Billings, 1875, Can. Nat. and Geol., vol. 8, p. 301, Corniferous Gr.

costulata, Whiteaves, 1884, Pal. Foss., vol. 3, p. 15, Guelph Gr.

galtensis, Whiteaves, 1884, Pal. Foss., vol. 3, p. 15, Guelph Gr.

FIG. 833.—*Ilionia galtensis*.

sinuata, Hall, 1859, (Anatina sinuata,) Pal. N. Y., vol. 3, p. 265, Low. Held. Gr.

Inoceramus, Sowerby, 1818, Min. Conch., vol. 2. This genus is unknown in American Palæozoic rocks.

chemungensis, see *Mytilarca chemungensis*.
mytilimeris, see *Plethomytilus mytilimeris*.
oviformis, see *Plethomytilus oviformis*.

ISCHYRINIA, Billings, 1866, Catal. Sil. Foss. Antic., p. 16. [Ety. *ischyros*, strong.] Equivalve, inequilateral; two strong ridges radiating from the beak in the interior of each valve. Type *I. winchelli*.

FIG. 834.—*Ischyria winchelli*.

plicata, Billings, 1866, Catal. Sil. Foss. Antic., p. 52, Anticosti Gr.

winchelli, Billings, 1866, Catal. Sil. Foss. Antic., p. 16, Hud. Riv. Gr.

Isocardia, Klein, 1753, Tent. Meth. Ostr. [Ety. *isos*, like; *kardia*, the heart.] This

is an existing littoral genus that burrows in the sand. It is not known in the Palaeozoic rocks.

- (f) *curta*, Shumard, 1858, Trans. St. Louis Acad. Nat. Sci., vol. 1, p. 206, Chocteau Gr.
jennae, Winchell, 1863, Proc. Acad. Nat. Sci., p. 17, Marshall Gr.
unioniformis, see *Edmondia unioniformis*.
Leda, Schumacher, 1817, syn. for *Nuculana*.
barrisi, White & Whitfield, syn. for *Palaeoneilo nuculiformis*.
bellistriata, see *Nuculana bellistriata*.
brevirostris, see *Nuculana brevisrostris*.
curta, see *Nuculana curta*.
dens-mamillata, see *Nuculana dens-mamillata*.
gibbosa, see *Yoldia gibbosa*.
knoxensis, see *Yoldia knoxensis*.
levistriata, see *Yoldia levistriata*.
nuculiformis, see *Palaeoneilo nuculiformis*.
obscura, see *Nuculana obscura*.
ohioensis, Hall, syn. for *Nuculana pandoriformis*.
oweni, see *Yoldia oweni*.
pandoriformis, see *Nuculana pandoriformis*.
polita, see *Nuculana polita*.
rushensis, see *Yoldia rushensis*.
saccata, see *Nuculana saccata*.
subscitula, see *Yoldia subscitula*.
LIOPTERIA, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 4. (Plates and Explanations.) [Ety. *leios*, smooth; *Pleria*, a genus.] Aviculoid, resembling in form *Actinopteria*; anterior extremity auriculate; wing large, extremity produced; test without proper rays; ligament external; ligamental area marked by fine parallel longitudinal striae; hinge with one or two oblique, slender, lateral teeth; the cavity of the beak partially separated from the anterior end by a short partition. Type *L. dekayi*.
bigsbyi, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 165, Ham. Gr.
chemungensis, Vanuxem, 1842, (Avicula *chemungensis*.) Geo. Rep. 3d Dist. N. Y., p. 182, and Pal. N. Y., vol. 5, p. 172, Chemung Gr.
conradi, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 159, Ham. Gr.

FIG. 835.—*Liopteria dekayi*. Cust showing vertical plate anterior to the beaks.

Dist. N. Y., p. 181, and Pal. N. Y., vol. 5, p. 158, Marcellus Shale.

- dekayi*, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 164, Ham. Gr.
gabbii, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 169, Ham. Gr.
greeni, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 160, Ham. Gr.
laevis, Hall, 1843, (Avicula *laevis*.) Geo. Rep. 4th Dist. N. Y., p. 181, and Pal. N. Y., vol. 5, p. 158, Marcellus Shale.
leai, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 168, Ham. Gr.
linguiformis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 173, Chemung Gr.
mittelli, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 166, Ham. Gr.
nitida, Hall, 1883, syn. for *L. chemungensis*.
oweni, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 170, Ham. Gr.
raffinesquii, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 161, Ham. Gr.
savi, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 162, Ham. Gr.
torreyi, Hall, Fig. 836.—*Liopteria raffinesquii*.
troosti, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 167, Ham. Gr.
LEPTODESMA, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 4. (Plates and Explanations.) [Ety. *leptos*, slender; *dema*, a ligament.] Like *Liopteria*, except the anterior end is nasute and acute, instead of auriculate and rounded; hinge-line narrow, with a slender, lateral tooth posterior to the beak; ligament external; test with concentric striae. Type *L. potens*.
acutirostrum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 234, Chemung Gr.
agassizi, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 182, Chemung Gr.
alatum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 218, Chemung Gr.
aliforme, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 220, Chemung Gr.
arciforme, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 229, Chemung Gr.
aviforme, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 224, Chemung Gr.
becki, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 185, Chemung Gr.
billingsi, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 192, Chemung Gr.
biton, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 222, Chemung Gr.
cadmus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 201, Chemung Gr.
clitus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 210, Chemung Gr.
complanatum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 227, Chemung Gr.
corydon, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 212, Chemung Gr.
creon, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 202, Chemung Gr.
curvatum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 196, Up. Chemung Gr.
demus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 203, Chemung Gr.
disparile, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 186, Up. Chemung Gr.
extenuatum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 207, Chemung Gr.

vol. 5, pt. 1, p.
 l. N. Y., vol. 5,
 r. N. Y., vol. 5,
 r. l. chemung-
 vol. 5, pt. 1, p.

 Liopteria rafin-
 squil.
 mung Gr.
 Y., vol. 5, pt. 1,
 N. Y., vol. 5,
 Explanations.)
 na, a ligament.)
 be anterior end
 tead of auricu-
 ge-line narrow,
 tooth posterior
 external; test
 Type L. potens.
 Pal. N. Y., vol.
 g Gr.
 Y., vol. 5, pt.
 Y., vol. 5, pt.
 N. Y., vol. 5, pt.
 N. Y., vol. 5,
 Gr.
 N. Y., vol. 5, pt.
 Y., vol. 5, pt. 1,
 N. Y., vol. 5, pt.
 Y., vol. 5, pt. 1,
 N. Y., vol. 5, pt.
 Y., vol. 5, pt. 1,
 Pal. N. Y., vol. 5,
 Gr.
 N. Y., vol. 5, pt.
 Y., vol. 5, pt. 1,
 N. Y., vol. 5,
 ing Gr.
 Y., vol. 5, pt.
 N. Y., vol. 5,
 ung Gr.
 Pal. N. Y., vol. 5,
 Gr.

flaccidum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 225, Chemung Gr.
 hector, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 209, Chemung Gr.
 jason, Hall, 1884, Hall, Pal. N. Y., vol. 5, pt. 1, p. 213, Chemung Gr.

FIG. 837.—*Leptodesma hector*.

lesleyi, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 223, Up. Chemung Gr.

lichas, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 232, Chemung Gr.

longispinum, Hall, 1843, (Avicula longispina,) Geo. Rep. 4th Dist. N. Y., p. 262, and Pal. N. Y., vol. 5, pt. 1, p. 179, Chemung Gr.

loxias, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 204, Chemung Gr.

lysander, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 216, Chemung Gr.

maclurii, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 228, Chemung Gr.

marcellense, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 175, Marcellus Shale.

matheri, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 193, Chemung Gr.

medon, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 197, Chemung Gr.

mentor, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 205, Chemung Gr.

mortoni, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 190, Chemung Gr.

mytiliforme, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 235, Chemung Gr.

naviforme, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 200, Chemung Gr.

nereus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 217, Chemung Gr.

orcus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 215, Chemung Gr.

orodes, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 206, Up. Chemung Gr.

orus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 219, Chemung Gr.

parallela, Simpson, 1889, Dict. of Pa. Foss., p. 332, Chemung Gr.

patulum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 226, Chemung Gr.

pelops, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 214, Up. Chemung Gr.

phaon, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 230, Chemung Gr.

potens, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 188, Up. Chemung Gr.

Liopteroideis, Simpson, 1889, Dict. of Pa. Foss., p. 331, Chemung Gr.
 lepidum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 195, Chemung Gr.

FIG. 838.—*Leptodesma hector*.

potens var. juvenis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 189, Chemung Gr.

propinquum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 231, Chemung Gr.

protectum, Conrad, 1842, (Avicula protecta,) Jour. Acad. Nat. Sci. Phil., vol. 8, p. 238, and Pal. N. Y., vol. 5, pt. 1, p. 183, Chemung Gr.

quadratum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 233, Chemung Gr.

robustum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 181, Chemung Gr.

rogersi, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 176, Ham. Gr.

rude, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 221, Chemung Gr.

shumardi, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 180, Chemung Gr.

sociale, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 187, Chemung Gr.

spinigerum, Conrad, 1842, (Avicula spinigera,) Jour. Acad. Nat. Sci. Phil., vol. 8, p. 237, and Pal. N. Y., vol. 5, pt. 1, p. 177, Chemung Gr.

stephani, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 194, Up. Chemung Gr.

transversum, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 167, Chemung Gr.

truncatum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 211, Chemung Gr.

umbonatum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 198, Chemung Gr.

umbonatum var. depressum, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 199, Chemung Gr.

LEPTODOMUS, McCoy, 1844, Synopsis Carb. Foss. Ireland, p. 66. [Ety. *leptos*, slender; *domus*, house.] Shell thin, short, oblong, tumid, subequivalve, inequilateral; beaks large, incurved; anterior side short, obtusely rounded, slightly gaping; deep ovate lunette between the beaks; posterior end broad, rounded, gaping, slope compressed, sides sulcated parallel with the ventral margin; dorsal margin inflected so as to form a lunette as long as the hinge-line; no hinge teeth; muscular impressions faint.

arata, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 152, Up. Silurian.

FIG. 839.—*Leptodomus canadensis*.

canadensis, Billings, 1874, Pal. Foss., vol. 2, p. 54, Gaspe limestone No. 8, Devonian.

clavata, Winchell, 1862, Proc. Acad. Nat. Sci., p. 415, Portage Gr.

granosus, see *Allorisma granosum*.

- topekensis*, see *Sedgwickia topekensis*.
undulatus, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 81, and Geo. Wis., vol. 4, p. 293, Niagara Gr.
mainensis, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 118, Low. Held. Gr.
pembrokeensis, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 118, Low. Held. Gr.
percingulatus, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 119, Low. Held. Gr.
Lima, Brugueire, 1791, Encycl. Meth. and Deshayes, 1824, Descrip. de Coquilles fossiles des environs de Paris. [Ety. *lima*, a file.] Not a Palaeozoic genus.
chesterensis, Worthen, not recognized.
glabra, see *Crenipecten glaber*.
macroptera, see *Limoptera macroptera*.
obsoleta, see *Crenipecten obsoletus*.
retifera, see *Crenipecten retiferus*.
rugostriata, see *Aviculopecten rugistriatus*.
LIMOPTERA, Hall, 1870, Prelim. Notice Lam. Shells, p. 15, Up. Held. Gr. [Ety. *Lima*, a genus; *pteron*, a wing.] Large, inequivalve, inequilateral, subquadrate, alate posterior, auriculate anterior; ligamental area large, common, longitudinally striate; hinge with an oblique posterior tooth and cardinal folds beneath the beak; anterior impression deep, posterior large, pallial line simple formed of a series of small pits; inter-pallial area pitted for the attachment of umbonal muscles; test radiated. Type *L. pauperata*.
cancellata, Hall, 1870, Prelim. Notice Lam. Shells, p. 16, and Pal. N. Y., vol. 5, pt. 1, p. 244, Ham. Gr.
curvata, Hall, 1870, Prelim. Notice Lam. Shells, p. 18, and Pal. N. Y., vol. 5, pt. 1, p. 250, Ham. Gr.
- LITHOPHAGA**, Lamarck, 1812, Hist. An. sans Vert. [Ety. *lithos*, stone; *phago*, I eat.] Not American Palaeozoic.
illinoisensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., p. 38, Keokuk Gr. Proposed instead of the form identified as *L. lingualis* of Phillips.
lingualis, Phillips, 1836, (Modiola lingualis.) Geol. Yorkshire, vol. 2, p. 209. Not American.
pertenuis, Meek & Wortlen, 1865, Proc. Acad. Nat. Sci. Phil., p. 245, and Geo. Sur. Ill., vol. 5, p. 539, St. Louis Gr.
Littorina wheeleri, see *Schizodus wheeleri*.
Lucina, Bruguiere, 1792, Encyclop. Meth. [Ety. mythological name.] Type *L. pennsylvanica*. Not an American Palaeozoic genus.
billingsana, see *Paracyclas billingsana*.
elliptica, see *Paracyclas elliptica*.
elliptica var. *occidentalis*, see *Paracyclas elliptica* var. *occidentalis*.
hamiltonensis, see *Paracyclas hamiltonensis*.
lirata, see *Paracyclas lirata*.
occidentalis, Billings, 1859, Assiniboine and Saskatchewan Ex. Exped. This name was preoccupied by Morton for an Eocene species, see *Paracyclas billingsana*.
ohioensis, see *Paracyclas ohioensis*.
retusa, see *Paracyclas retusa*.
varysburgia, see *Paracyclas varysburgensis*.
wyomingensis, see *Paracyclas wyomingensis*.
- LUNULICARDIUM**, Munster, 1840, Beitrage zur Petrefaktenkunde, 3d heft, p. 69. [Ety. *lunula*, a little moon; *Cardium*, a genus.] Equivalve, inequilateral, subelliptical, subcircular, or trigonal; posterior side obliquely truncate, margin often reflexed and produced; beaks pointed; cardinal line marked by a lunate hiatus; surface radiated and concentrically marked; ligament external.
acutirostrum, syn. for *L. ornatum*.

FIG. 840.—*Limoptera macroptera*.

macroptera, Conrad, 1838, (Lima macroptera.) Ann. Rep. N. Y., p. 117, and Pal. N. Y., vol. 5, pt. 1, p. 246, Ham. Gr.
obsoleta, Hall, 1870, Prelim. Notice Lam. Shells, p. 18, and Pal. N. Y., vol. 5, pt. 1, p. 249, Ham. Gr.
pauperata, Hall, 1870, Prelim. Notice Lam. Shells, p. 16, and Pal. N. Y., vol. 5, pt. 1, p. 243, Up. Held. Gr.
sarmenticia, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 167, Devonian.

FIG. 841.—*Lunulicardium curtum*.

curtum, Hall, 1884, Pal. N. Y., vol. 5, pl. 71, figs. 18-23, Up. Held. Gr.

Hist. An. sans
phago, I eat.]

1882, Bull. No. 1,
p. 38, Keokuk
of the form
of Phillips.
(Modiola lin-
e, vol. 2, p. 209.

en, 1865. Proc.
p. 245, and Geo.
St. Louis Gr.
us wheeleri.
Encyclop. Meth.
ame.] Type L.
American Paleo-

s billingsana.
elliptica.
see Paracyclas
lis.
ashamiltonensis.
ta.
Assiniboine and
bed. This name
Morton for an
Paracyclas bill-

oioensis.
usa.
las varysburgen-
cyclas wyoming-

1840, Beitrage zur
heft, p. 69. [Ety.
Pardium, a genus.]
al, subelliptical,
l; posterior side
margin often re-
beaks pointed;
y a lunate hiatus;
d concentrically
ernal.
ornatum.

um curtum.

N. Y., vol. 5, pl. 71,
Gr.

fragile, Hall, 1843, (Avicula fragills,) Geo. Rep. 4th Dist. N. Y., p. 222, and Pal. N. Y., vol. 5, pl. 71, figs. 1-14, Genesee Shale.

fragosum, Meek, 1877, (Posidonomya fragosa,) U. S. Geo. Expl. 40th Parallel, vol. 4, p. 92, Carboniferous.

FIG. 842.—Lunulicardium marcellense.

marcellense, Vanuxem, 1842, (Cypriocardites marcellensis,) Geo. Rep. 3d Dist. N. Y., p. 146, and Pal. N. Y., vol. 5, pl. 71, figs. 15-16, Marcellus Shale.

orbiculare, Hall, 1885, Pal. N. Y., vol. 5, p. 436, Marcellus Shale.

ornatum, Hall, 1843, (Pinnopsis ornata,) Geo. Rep. 4th Dist. N. Y., p. 244, and Pal. N. Y., vol. 5, p. 437, Portage Gr. rude, Hall, 1884, Pal. N. Y., vol. 5, pl. 71, fig. 17, Marcellus Shale.

transversum, Hall, 1885, Pal. N. Y., vol. 5, p. 439, Chemung Gr.

Lyonsia, Turton, 1822. Not found in Palaeozoic rocks.

concaua, see Sedgwickia concava. LYRIOPECTEN, Hall, 1884, Pal. N. Y., vol. 5, p. 3. (Plates and Explanations.) [Ety. lyron, a lyre; Pecten, a genus.] Distinguished from Aviculopecten by the shorter hinge-line and very small anterior wing; surface with strong rays. Type L. magnificus.

alternatus, Simpson, 1889, Dict. Foss. Pa., p. 366, and Trans. Am. Phil. Soc., p. 446, Chemung Gr.

anomiiformis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 53, Up. Held. Gr.

cymbalon, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 47, Ham. Gr.

dardanus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 41, Up. Held. Gr.

fasciatus, Hall, 1884, (Pernopecten fasciculatus,) Pal. N. Y., vol. 5, pt. 1, p. 55, Chemung Gr.

interradiatus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 44, Ham. Gr.

FIG. 843.—Lyriopecten orbiculatus.

macrodontus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 46, Up. Held. Gr.

magnificus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 51, Up. Chemung Gr.

orbiculatus, Hall, 1843, (Avicula orbiculata,) Geo. Rep. 4th Dist., N. Y., p. 202, and Pal. N. Y., vol. 5, pt. 1, p. 42, Ham. Gr.

paralleodontus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 40, Up. Held. Gr.

polydorus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 50, Chemung Gr.

priamus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 54, Chemung Gr.

solox, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 56, Up. Chemung Gr.

tricostatus, Vanuxem, 1842, (Avicula tricostata,) Geo. Sur. 3d Dist. N. Y., p. 179, and Pal. N. Y., vol. 5, pt. 1, p. 48, Chemung Gr.

LYRODESMA, Conrad, 1841, Ann. Geo. Rep. N. Y., p. 51. [Ety. lyra, a harp; desma, a ligament.] Equivalve, inequilateral, semicircular; hinge plate with 6 to 8 angular, crenulated teeth radiating from beneath the beak upon a more or less rounded platform. Type L. planum.

cincinnatiense, Hall, 1871, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 227, Hud. Riv. Gr.

planum, Conrad, 1841, Ann. Geo. Rep. p. 51, Hud. Riv. Gr.

FIG. 844.—Lyrodesma poststriatum.

poststriatum, Emmons, 1842, (Nuculana poststriata,) Geo. Rep. N. Y., p. 399, and Pal. N. Y.,

vol. 1, p. 151, Black Riv. Gr. pulchellum, Hall, 1847, Pal. N. Y., vol. 1, p. 302, Hud. Riv. Gr.

MACRODON, Lycett, 1845, Murch. Geo. Chelt. [Ety. macros, long; odous, a tooth.] Shell oblong, very inequilateral, moderately tumid, a byssal sinus in the anterior third of the ventral margin anterior edges of the adductor impressions prominent; hinge teeth at the anterior end

few, slightly oblique or nearly at right angles to the hinge-line beneath the beak, becoming more oblique toward the anterior end; posterior part of the hinge-line, from beak to anal angle, occupied by one to three long lateral teeth.

carbonarius, Cox, 1857, (Arca carbonarius,) Geo. Sur. Ky., vol. 3, p. 567, Coal Meas.

chemungensis, Hall, 1870, Prelim. Notice Lam. Shells, p. 14, and Pal. N. Y., vol. 5, pl. 51, figs. 11-16, Chemung Gr. cochlearis, Winchell, 1863, Proc. Acad. Nat. Sci., p. 16, Marshall Gr. Prof. Hall suggests that it is a syn. for M. parvus.

FIG. 845.—Macrodon obsoletus.

curtus, Hartt, 1868, Acad. Geol., p. 302, Carb.
delicatus, Meek & Worthen, 1870, Proc. Acad. Nat. Sci., p. 40, and Geo. Sur. Ill., vol. 5, p. 575, Up. Coal Meas.
hamiltoniae, Hall, 1870, Prelim. Notice Lam. Shells, p. 13, and Pal. N. Y., vol. 5, pl. 51, figs. 1-10, Ham. Gr.
hardingi, Hartt, 1868, Acad. Geol., p. 302, Carb.
micronema, Meek & Worthen, 1866, Proc. Acad. Nat. Sci., p. 261, Kaskaskia Gr.
obsoletus, Meek, 1871, Reg. Rep. University W. Va., p. 5, and Pal. Ohio, vol. 2, p. 334, Coal Meas.

FIG. 846.—*Matheria tener*.

ovatus, Hall, 1870, Prelim. Notice Lam. Shells, p. 15, and Pal. N. Y., vol. 5, p. 351, Waverly Gr.
parvus, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., vol. 8, p. 299, Kinderhook Gr.
sangamonensis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 123 Coal Meas.
shubenacadiensis, Hartt, 1868, Acad. Geo., p. 302, Carb.
tenuistriatus, Meek & Worthen, 1867, Proc. Chi. Acad. Sci., vol. 1, p. 17, Up. Coal Meas.
truncatus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 243, Carboniferous.
MATHERIA, Billings, 1858, Can. Nat. and Geo., vol. 3, p. 440. [Ety. proper name.] Equivalve, inequilateral; beaks anterior; two small, obtuse, cardinal teeth in the left valve, and one in the right; no lateral teeth; two muscular impressions; ligament external. Type *M. tener*.
tener, Billings, 1858, Can. Nat. and Geo., vol. 3, p. 440, Trenton Gr.
MEGALOMUS, Hall, 1852, Pal. N. Y., vol. 2, p. 343. [Ety. *megas*, great; *omos*, shoulder.] Large, equivalve, concentrically lined, longitudinal; umbones anterior incurved; shell thick and along the hinge-line thickened on the interior; muscular impression large and deep, with two small circular pits above. Type *M. canadensis*.
canadensis, Hall, 1852, Pal. N. Y., vol. 2, p. 343, Guelph Gr.
compressus, Nicholson & Hinde, 1874, Can. Jour., vol. 14, p. 159, Niagara Gr.
MEGAMBONIA, Hall, 1859, Pal. N. Y., vol. 3, p. 273. [Ety. *megas*, great; *ambon*, the boss of a shield.] Equivalve, inequilateral, subovoid, gibbous in the middle and toward the umbones; anterior side

lobed or alate; muscular impression large; posterior cardinal margin expanded or alate; hinge-line crenulated anteriorly; teeth numerous; surface concentrically lined, and sometimes with radiating striae. Type *M. suborbicularis*.

FIG. 847.—*Megalomus canadensis*.

aviculoidea, Hall, 1850, Pal. N. Y., vol. 3, p. 274, Low. Held. Gr.
bellistriata, Hall, 1859, Pal. N. Y., vol. 3, p. 467, Oriskany sandstone.
cancellata, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 153, Up. Silurian.
cardiiformis, Hall, 1843, (Pterinea cardiiformis,) Geo. Rep. 4th Dist. N. Y., p. 172, and Pal. N. Y., vol. 5, p. 515, Cornif. Gr.

FIG. 848.—*Megambonia cardiiformis*.

cordiformis, see *Mytilarca cordiformis*.
jamesi, see *Ambonychia jamesi*.
lamellosa, Hall, 1859, Pal. N. Y., vol. 3, p. 467, Oriskany sandstone.
lata, Hall, 1859, Pal. N. Y., vol. 3, p. 277, Low. Held. Gr.
lyoni, syn. for *Cardiopsis radiata*.
mytiloidea, Hall, 1859, Pal. N. Y., vol. 3, p. 276, Low. Held. Gr.

muscular impres-
sion cardinal margin
hinge-line crenu-
late numerous; sur-
red, and sometimes
Type M. subor-

canadensis.

Pal. N. Y., vol. 3,
r.
Pal. N. Y., vol. 3,
stone.
Can. Nat. and Geo.,
lurian.
3, (Pterinea cardii-
4th Dist. N. Y., p.
vol. 5, p. 515. Cor-

cardiiformis.

ca cordiformis.

ia jamesi.

Pal. N. Y., vol. 3,
stone.

N. Y., vol. 3, p. 277.

is radiata.

Pal. N. Y., vol. 3,
Gr.

oblonga, Hall, 1859, Pal. N. Y., vol. 3, p. 277, Low. Held. Gr.
obscura, Hall, 1859, Pal. N. Y., vol. 3, p. 277, Low. Held. Gr.
occuldualis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 173, Devonian.
ovata, Hall, syn. for Plethomytilus mytilimeris.
ovoidea, Hall, syn. for Plethomytilus mytilimeris.
rhomboldea, Hall, 1859, Pal. N. Y., vol. 3, p. 275, Low. Held. Gr.
spinneri, Hall, 1859, Pal. N. Y., vol. 3, p. 274, Low. Held. Gr.
striata, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 153, Up. Silurian.
subcardiformis, Hall, syn. for M. cardiiformis.
suborbicularis, Hall, 1859, Pal. N. Y., vol. 3, p. 273, Low. Held. Gr.
Megaptera, Meek & Worthen, 1866. The name was preoccupied.
Microdon, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 3, p. 247. This name was applied by Agassiz to a genus of fish in 1833, and was also preoccupied for a genus of insecta. Hall proposed Eodon, in 1877, but Whitfield has shown that *M. bellistriatus* is a Cypricardella, and hence the latter name has priority.
bellistriatus, see Cypricardella bellistriata.
complanatus, see Cypricardella complanata.
gregarius, see Cypricardella gregaria.
reservatus, see Cypricardella reservata.
tenuistriatus, see Cypricardella tenuistriata.

FIG. 849.—*Modiella pygmaea*.

MODIELLA, Hall, 1884, Pal. N. Y., vol. 5, p. 4. (Plates and Explanations.) [Ety. *modus*, a measure; *ellus*, diminutive.] Subrhomboidal, narrowed and arcuate in front, broadly expanding posteriorly; two well-marked muscular impressions connected by a simple pallial line; surface with radiating striae. Type M. *pygmaea*.
pygmaea, Conrad, 1842, (Pterinea *pygmaea*.) Jour. Acad. Nat. Sci., vol. 8, p. 251, and Pal. N. Y., vol. 5, pl. 76, figs. 9-20, Ham. Gr.
Modiola, Lamarck, 1801, Syst. An. sans Vert. [Ety. *modiolus*, a small measure or drinking vessel.] Oblong, inflated in front, umbones anterior, obtuse, no teeth; pedal impressions three, the central one elongated. Type M. *modiolus*. Not a Palaeozoic genus. Species are only left here for want of material to properly determine the generic relations.
avonia, Dawson, 1868, Acad. Geol., p. 301, Subcarboniferous.
concentrica, see Modiomorpha *concentrica*.
illinoisensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 16, and Geo. Sur. Ill., vol. 3, p. 125, St. Louis Gr.
lingualis, see Lithophaga *lingualis*.
metella, see Mytilops *metella*.

minor, Lea, 1852, Jour. Acad. Nat. Sci., 2d series, vol. 2, Coal Meas. Not determinable.

FIG. 850.—*Modiola modiolus*. aa, Anterior adductors; aa', posterior adductors; uu, pp', pedal muscles; pp, byssal muscles; f, foot; b, byssus; m, pallial line.

nevadensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 239, Subcarboniferous.

obtus, see Modiolopsis *obtus*.

pooli, Dawson, 1868, Acad. Geol., p. 301, Low. Carb. Founded upon a small cast destitute of characters.

præcedens, see Mytilops *præcedens*.

wyomingensis, Lea, 1852, Jour. Acad. Nat. Sci., 2d series, vol. 2, p. 205, Coal Meas. Not recognized, but probably an Anthracomya.

MODIOLOPSIS, Hall, 1847, Pal. N. Y., vol. 1, p. 157. [Ety. *Modiola*, a genus of shells; *opsis*, appearance; from its resemblance to *Modiola*.] Equivalve, inequilateral, elongated, broader posteriorly; umbones anterior; cardinal teeth short, oblique; single, deep, subcircular anterior muscular impression; ligament external; no area; surface concentrically lined. Type M. *modiolaris*.

adrasia, Billings, 1862, Pal. Foss., vol. 1, p. 45, Black Riv. Gr.

anodontoides, Conrad, 1847, (Cypricardites *anodontoides*.) Pal. N. Y., vol. 1, syn. for M. *sinuata*.

arcuata, Hall, 1847, Pal. N. Y., vol. 1, p. 159, Trenton Gr.

aviculoides, Hall, 1847, Pal. N. Y., vol. 1, p. 161, Trenton Gr.

cancellata, Walcott, 1879, Trans. Alb. Inst., vol. 10, p. 22, Utica Slate Gr.

capax, n. sp. Shell very large, oblong; cardinal and basal lines behind the beaks subparallel; basal margin slightly contracted by an undefined cincture arising below the beaks; posterior end broadly rounded; depressed in front of

the beaks; anterior end rounded; beaks large, obtuse, and extending beyond the hinge-line; umbones large; surface marked with concentric lines of growth, and with strong transverse lines over the umbonal region, some of which extend nearly to the basal line, the anterior ones curve a little forward in passing over the umbones. Collected by the author in the Hud. Riv. Gr. at Versailles, Indiana.

carinata, Hall, 1847, Pal. N. Y., vol. 1, p. 160, Trenton Gr.

carrollensis, Worthen, 1882, Bull. No. 1, Ill. St. Mus. Nat. Hist., Galena Gr. Proposed instead of *M. subnasuta* of Meek & Worthen, 1870, Proc. Acad. Nat. Sci., p. 41, which was preoccupied.

cincinnatiensis, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 88, Utica Slate.

FIG. 851.—*Modiolopsis capax*.

concentrica, Hall & Whitfield, 1872, Ohio Pal., vol. 2, p. 86, Hud. Riv. Gr.

curta, Hall, 1847, Pal. N. Y., vol. 1, p. 297, Hud. Riv. Gr.

dicteus, Hall, 1867, 20th Rep. N. Y. Mus. Nat. Hist., p. 385, Niagara Gr.

(?) *dubia*, Hall, 1859, Pal. N. Y., vol. 3, p. 264, Low. Held. Gr.

exilis, Billings, 1874, Pal. Foss., vol. 2, p. 132, Up. Sil.

fabia, Emmons, 1842, Geo. Rep. 2d Dist. N. Y., p. 395, and Pal. N. Y., vol. 1, p. 158, Black Riv., Trenton, and Hud. Riv. Grs.

FIG. 852.—*Modiolopsis faberi*.
Left valve.

behind the middle, and much contracted in front of the beaks; hinge-line nearly straight from the anterior end to the middle; it then becomes arcuate to near the posterior end, which is abruptly rounded; basal line slightly sinuate at the anterior third from the cincture, directed downward and backward from the anterior part of the umbo; beaks

strong, projecting anteriorly beyond the hinge-line; umbones high, subangular, and gradually declining toward the postero-basal line; anterior muscular impression very large, circular, deep, and situate at the anterior end, in front of and below the beaks; dorsal ligament very large. Distinguished from *M. concentrica* in its general outline, more elongate form, more prominent beaks, and higher and more angular umbones. Hud. Riv. Gr., at Cincinnati, O. The specimen figured is from the collection of Charles Faber.

FIG. 853.—*Modiolopsis faberi*.
Cardinal view, showing greatest thickness of shell toward the posterior.

gesneri, Billings, 1862, Pal. Foss., vol. 1, p. 43, Trenton and Black Riv. Grs.

latus, see *Cypricardites latus*.

maia, Billings, 1862, Pal. Foss., vol. 1, p. 44, Trenton Gr.

meyeri, Billings, 1862, Pal. Foss., vol. 1, p. 42, Trenton Gr.

modiolaris, Conrad, 1838, (Pterinea *modiolaris*.)

Ann. Geo. Rep. N. Y., p. 118, and Pal. N. Y., vol. 1, p. 294, Hud. Riv. Gr.

modioliformis, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 294, Trenton Gr.

mytiloides, Hall, 1847, Pal. N. Y., vol. 1, p. 157, Black Riv. and Trenton Grs.

nais, Billings, 1862, Pal. Foss., vol. 1, p. 45, Black Riv. Gr.

FIG. 854.—*Modiolopsis modiolaris*. Hinge and pallial line and muscular impression.

nasuta, Conrad, 1841, (Cypricardites *nasutus*.) Ann. Rep. N. Y., p. 52, and Pal. N. Y., vol. 1, p. 159, Trenton and Hud. Riv. Grs.

nuculiformis, see *Tellinomya nuculiformis*.

obtusa, Hall, 1847, (Modiola *obtusa*.) Pal. N. Y., vol. 1, p. 40, Birdseye Gr.

occidens, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 77, Trenton Gr.

priorly beyond the
high, subangular,

Modiolopsis faberi.
view, showing
thickness of shell
at posterior.

end, in front of
dorsal ligament
shed from M. con-
al outline, more
prominent beaks,
angular umbones,
cinnatli, O. The
on figured is from
ection of Charles

Billings, 1862, Pal.
vol. 1, p. 43, Tren-
Black Riv. Grs.
e Cypricardites

Billings, 1862, Pal.
vol. 1, p. 44, Tren-

Billings, 1862, Pal.
vol. 1, p. 42, Tren-

as, Conrad, 1838,
sea modiolaris,
Y., p. 118, and
294, Hud. Riv. Gr.
t Worthen, 1868,
3, p. 294, Tren-

Pal. N. Y., vol. 1,
Trenton Grs.

l. Foss., vol. 1, p.

liaris. Hinge and
lar impression.

Cypricardites na-
Y., p. 62, and Pal.
Trenton and Hud.

mya nuculiformis.
diola obtusa,) Pal.
rdseye Gr.
5, Monogr. U. S.
7, Trenton Gr.

orthonota, Conrad, 1839, (Unio orthono-
tus,) Ann. Rep. N. Y., p. 66, and Geo.
Rep. 4th Dist. N. Y., pl. 2, figs. 8 and 9,
Medina sandstone.

orthonota, Meek & Worthen, 1868, Geo.
Sur. Ill., vol. 3. This name was preoc-
cupied. See *M. rectiformis*.

ovata, Hall, 1852, Pal. N. Y., vol. 2, p.
101, Clinton Gr.

parallela, see *Orthodesma parallelum*.

parviuscula, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 446, Chazy Gr.

perlata, Hall, 1870, 28th Rep. N. Y. Mus.
Nat. Hist., p. 172, Niagara Gr.

perovata, see *Modiomorpha perovata*.

pholadiformis, Hall, 1851, Lake Sup. Land
Dist., vol. 2, p. 213, Hud. Riv. Gr.

plana, Hall, 1861, Geo. Rep. Wis., p. 30,
Trenton Gr.

pogonipensis, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 78, Trenton Gr.

prisca, Walcott, 1887, Am. Jour. Sci. and
Arts, 3d ser., vol. 34, p. 191, Up. Ta-
conic. Not a *Modiolopsis*.

primigenia, Conrad, 1838, (Unio primi-
genius,) Ann. Rep. N. Y., p. 66, and
Geo. Rep. 4th Dist. N. Y., pl. 2, fig. 3,
Medina sandstone.

recta, Hall, 1867, 20th Rep. N. Y. Mus.
Nat. Hist., p. 386, Niagara Gr.

rectiformis, Worthen, 1882, Bull. No. 1,
Ill. St. Mus. Nat. Hist., p. 38, Tren-
ton Gr. Proposed instead of *M. ortho-*

nota, Meek & Worthen, 1868, Geo. Sur.
Ill., vol. 3, p. 295, which was preoc-
cupied.

rhomboidea, Hall, 1860, Can. Nat. and
Geo., vol. 5, p. 148, Up. Sil.

rudis, Billings, 1874, Pal. Foss., vol. 2, p.
133, Up. Sil.

sinuata, Emmons, 1842, (Cypricardites
sinuatus,) Geo. Rep. 2d Dist. N. Y., p.
399, and Pal. N. Y., vol. 1, p. 298, Hud.
Riv. Gr.

striata, Billings, 1866, Catal. Sil. Foss.
Antic., p. 48, Anticosti Gr.

subalata, Hall, 1852, Pal. N. Y., vol. 2, p.
84, Clinton and Niagara Grs.

subcarinata, Hall, 1852, Pal. N. Y., vol.
2, p. 601, Clinton Gr.

subnasuta, Hall, 1860, Can. Nat. and Geo.,
vol. 5, p. 148, Up. Sil.

subnasuta, Meek & Worthen. See *M. can-*

cellata.

subrhomboidea, Simpson, 1889, Trans.
Am. Phil. Soc., p. 450, and Dict. Foss.,
Pa., p. 411, Clinton Gr.

subspatulata, see *Cypricardites subspatu-*

latus.

superba, Hall, 1861, Geo. Rep. Wis., p.
31, Trenton Gr.

terminalis, Hall, 1847, Pal. N. Y., vol. 1,
p. 318, Hud. Riv. Gr.

trentonensis, Hall, 1847, Pal. N. Y., vol. 1,
p. 161, Trenton Gr.

truncata, Hall, 1847, Pal. N. Y., vol. 1, p.
296, Hud. Riv. Gr.

undulostriata, Hall, 1852, Pal. N. Y., vol.
2, p. 284, Niagara Gr.

unionoides, Meek, 1871, (Anodontopsis
unionoides,) Am. Jour. Sci. and Arts,
3d ser., vol. 2, p. 299, and Ohio Pal.,
vol. 1, p. 141, Hud. Riv. Gr.

varia, Billings, 1874, Pal. Foss., vol. 2, p.
56, Low. Held. Gr.

FIG. 855.—*Modiolopsis versaillesensis*. Left
valve.

versaillesensis, S. A. Miller, 1874, Cin.
Quar. Jour. Sci., p. 150, Hud. Riv. Gr.

FIG. 856.—*Modiolopsis versaillesensis*. Hinge-
line and muscular impression.

MODIOMORPHA, Hall, 1870, Prelim. Notice
Lam. Shells, p. 72. [Ety. contracted from
Modiola, a genus; *morphe*, form.] Equi-
valve, subovate, larger posteriorly, com-
pressed; beaks small; sinus oblique
and constricting the base; surface con-
centrically undulated; single tooth in
the left valve, and corresponding socket
in the other; no lateral teeth; ligament
external. Type *M. concentrica*.

affinis, Hall, 1885, Pal. N. Y., vol. 5, p.
284, Ham. Gr.

alta, Conrad, 1841, (Cypricardites *alta*.)
Ann. Rep. N. Y., p. 52, and Pal. N. Y.,
vol. 5, pl. 37, figs. 1-16, Ham. Gr.

altiformis, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 169, Devonian.

ambigua, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 239, Carboniferous.

arcuata, Hall, 1884, Pal. N. Y., vol. 5, pl.
36, fig. 21, Ham. Gr.

chemos, Clarke, 1885, Bull. U. S. Geo.
Sur., No. 16, p. 30, Genesee shales.

clarens, Hall, 1885, Pal. N. Y., vol. 5, p.
273, Up. Held. Gr.

complanata,
Hall, 1870,
Prelim. Notice
Lam. Shells, p.
73, and Pal.
N. Y., vol. 5,
p. 272, Up.
Held. Gr.

concentrica, Con-
rad, 1838, (Pte-
rinea *concentrica*.) Ann. Rep. Geo. Sur.
N. Y., p. 116, and Pal. N. Y., vol. 5, pl.
34, figs. 9-10, Ham. Gr.

concentrica, Con-
rad, 1838, (Pte-
rinea *concentrica*.) Ann. Rep. Geo. Sur.
N. Y., p. 116, and Pal. N. Y., vol. 5, pl.
34, figs. 9-10, Ham. Gr.

concentrica, Con-
rad, 1838, (Pte-
rinea *concentrica*.) Ann. Rep. Geo. Sur.
N. Y., p. 116, and Pal. N. Y., vol. 5, pl.
34, figs. 9-10, Ham. Gr.

concentrica, Con-
rad, 1838, (Pte-
rinea *concentrica*.) Ann. Rep. Geo. Sur.
N. Y., p. 116, and Pal. N. Y., vol. 5, pl.
34, figs. 9-10, Ham. Gr.

concentrica, Con-
rad, 1838, (Pte-
rinea *concentrica*.) Ann. Rep. Geo. Sur.
N. Y., p. 116, and Pal. N. Y., vol. 5, pl.
34, figs. 9-10, Ham. Gr.

concentrica, Con-
rad, 1838, (Pte-
rinea *concentrica*.) Ann. Rep. Geo. Sur.
N. Y., p. 116, and Pal. N. Y., vol. 5, pl.
34, figs. 9-10, Ham. Gr.

concentrica, Con-
rad, 1838, (Pte-
rinea *concentrica*.) Ann. Rep. Geo. Sur.
N. Y., p. 116, and Pal. N. Y., vol. 5, pl.
34, figs. 9-10, Ham. Gr.

FIG. 857.—*Modiomorpha*
concentrica.

cymbula, Hall, 1870, Prelim. Notice Lam. Shells, p. 75, and Pal. N. Y., vol. 5, pl. 36, figs. 19-20, Ham. Gr.

desiderata, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 240, Carboniferous.

hyale, Hall, 1870, Prelim. Notice Lam. Shells, p. 79, and Pal. N. Y., vol. 5, pl. 41, figs. 28-30, Waverly Gr.

inornata, Billings, 1874, Pal. Foss., vol. 2, p. 52, Devonian.

linguliformis, Hall, 1883, Pal. N. Y., vol. 5, pl. 34, figs. 15-17, Up. Held. Gr.

macilenta, Hall, 1870, Prelim. Notice Lam. Shells, p. 76, and Pal. N. Y., vol. 5, pl. 37, fig. 17, and pl. 39, figs. 17-21, Ham. Gr.

mytiloides, Conrad, 1841, (Cypricardites mytiloides,) Ann. Rep. Geo. N. Y., p. 52, and Pal. N. Y., vol. 5, p. 277, Ham. Gr.

neglecta, Hall, 1883, Pal. N. Y., vol. 5, pl. 41, figs. 12-13, Chemung Gr.

oblonga, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 170, Devonian.

obtusa, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 171, Devonian.

perovata, Meek & Worthen, 1865, (Modiolopsis perovata,) Proc. Acad. Nat. Sci. Phil., p. 246, and Geo. Sur. Ill., vol. 3, p. 438, Ham. Gr.

pintoensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 240, Carboniferous.

planulata, Hall, 1870, Prelim. Notice Lam. Shells, p. 74, syn. for M. mytiloides.

ponderosa, Hall, 1870, (Sanguinolites ponderosus,) Prelim. Not. Lam. Shells, p. 35, and Pal. N. Y., vol. 5, pl. 34, fig. 11, Up. Held. Gr.

putillus, Hall, 1883, Pal. N. Y., vol. 5, pl. 41, figs. 1-2, Schoharie grit.

quadrula, Hall, 1870, Prelim. Notice Lam. Shells, p. 77, and Pal. N. Y., vol. 5, pl. 41, figs. 18-26, Chemung Gr.

recta, Hall, 1885, Pal. N. Y., vol. 5, p. 286, Ham. Gr.

recurva, Hall, 1883, Pal. N. Y., vol. 5, pl. 41, fig. 17, Chemung Gr.

regularis, Hall, 1885, Pal. N. Y., vol. 5, p. 270, Schoharie grit.

rigida, Hall, 1883, Pal. N. Y., vol. 5, pl. 41, figs. 10, 11, 14-16, Chemung Gr.

rigidula, Simpson, 1889, Trans. Am. Phil. Soc., p. 449, and Dict. Foss. Pa., p. 415, Chemung Gr.

schoharie, Hall, 1884, Pal. N. Y., vol. 5, p. 269, pl. 34, fig. 13, Schoharie grit.

subalata, Conrad, 1841, Cypricardites subalatus,) Ann. Rep. N. Y., p. 83, and Pal. N. Y., vol. 5, pl. 39, figs. 1-16, Ham. Gr.

subalata var. chemungensis, Hall, 1885, Pal. N. Y., vol. 5, p. 284, Chemung Gr.

subangulata, Hall, 1885, Pal. N. Y., vol. 5, p. 287, Chemung Gr.

tioga, Hall, 1885, Pal. N. Y., vol. 5, p. 291, Chemung Gr.

MONOPTERIA, Meek & Worthen, 1866, Proc. Chi. Acad. Nat. Sci., vol. 1, p. 20. [Ety. *monos*, single; *pteron*, a wing.] Aviculoid, obliquely produced, angular pos-

teriorly, rounded in front, subequivalve, both valves convex; posterior wing slender, produced, anterior one obsolete or drawn back between the beaks, in a deep lunule; no byssal emargination, but a little gaping in the lunule; muscular impressions faint; cardinal area narrow, with few longitudinal cutlivalve furrows; hinge edentulous. Type M. longispina.

auricula, Stevens, 1858, (Gervillia auricula,) Am. Jour. Sci. and Arts, 2d ser., vol. 25, p. 205, Coal Meas.

gibbosa, Meek & Worthen, 1866, Proc. Chi. Acad. Sci., p. 20, Coal Meas.

longispina, Cox, 1857, (Gervillia longispina,) Geo. Sur. Ky., vol. 3, p. 568, Coal Meas.

FIG. 858.—*Monopteria gibbosa*.

marian, White, 1874, Rep. Invert. Foss., p. 22, and Geo. Sur. W. 100th Mer., vol. 4, p. 151, Carboniferous.

MONOTIS, Bronn, 1824, System Urweltlicher Konchylien. [Ety. *monos*, one; *otos*, ear.] Obliquely oval, compressed, radiated; anterior side short, rounded; posterior slightly eared. Type M. salinaria.

FIG. 859.—*Monotis gregaria*.

elevata, see Panenka elevata.

gregaria, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 38, and Geo. Sur. Ill., vol. 5, p. 573, Coal Meas.

halli, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 185, Permian Gr.

hawni, see Pseudomonotis hawni.

poulsoni, see Panenka poulsoni.

princeps, see Aviculopecten princeps.

radialis, Phillips, 1834, (Pecten radialis,) see Pseudomonotis radialis.

radians, see Panenka radians.

septentrionalis, Haughton, 1857, Jour. Roy. Dub. Soc., vol. 1, (?) Gr.

speluncaria, Schlotheim, 1816, Denkschriften d. k. Ak. d. Wiss. zu Munchen, p. 30, (Gryphites speluncarius,) Permian Gr. Probably not American.

variabilis, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 187, Permian Gr.

MYALINA, DeKoninck, 1844, Desc. Anim. Foss. Carb. Belg., p. 125. [Ety. *Mya*, a genus of shells; *inus*, like.] Subrhomboidal, inequilateral, inequivalve, oblique, slightly sinuous in front for the passage of the byssus; beaks pointed, nearly terminal; surface smooth, or concentrically marked; hinge edentulous; ligamental area broad and furrowed parallel to the hinge-line; muscular and pallial impressions apparently as in *Pteria*; shell structure prismatic. Type M. lamellosa.

ont, subequivalve,
posterior wing
anterior one obsolete
between the beaks, in a
small emargination,
the lunule; mus-
cular; cardinal area
longitudinal cartilage
reticulate. Type M.

(*Gervillia muric-*
and Arts, 2d ser.,
Meas.

858.—*Monopteria*
gibbosa.
ep. Invert. Foss.,
W. 100th Mer.,
iferous.

stem Urweltlicher
ones, one; *ous, olos*,
compressed, ra-
short, rounded;
ed. Type M. sa-

Panenka elevata.
Meek & Worthen,
c. Acad. Nat. Sci.
38, and Geo. Sur.
5, p. 573, Coal

low, 1858, Trans.
Acad. Sci., vol. 1,
Permian Gr.

is hawni.
oulsoni.

ten princeps.
(*Pecten radialis*),
dialis.

dians.
ton, 1857, Jour.
t, (?) Gr.

1816, Denkschrif-
n. zu Munchen, p.
ncarius.) Permian
merican.

3, Trans. St. Louis
87, Permian Gr.

844, Desc. Anim.
125. [Ety. *Mya*, a
like.] Subrhomb-

inequivalve, ob-
s in front for the
s; beaks pointed,
surface smooth; or con-
edentulous;
ad and furrowed
ge-line; muscular
ns apparently as
ucture prismatic.

angulata, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 455, and Geo.
Sur. Ill., vol. 2, p. 300, Kaskaskia Gr.

apachesi, Marcou, 1858, Geol. North
America, p. 44, Subcarboniferous.

aviculoidea, Meek & Hayden, 1860, Proc.
Acad. Nat. Sci. Phil., p. 184, and Pal.
Up. Mo., p. 51, Permian Gr.

aviculoidea, Winchell, see M. rara.

concava, Swallow, 1858, (*Mytilus conc-*
avus.) Trans. St. Louis Acad. Sci., vol.

1, p. 188, Permian Gr.

concentrica, Meek & Worthen, 1860, Proc.
Acad. Nat. Sci. Phil., p. 456, and Geo.

Sur. Ill., vol. 2, p. 281, Warsaw Gr.

congeneris, Walcott, 1885, U. S. Geo. Sur.,
vol. 8, p. 237, Subcarboniferous.

cuneiformis, Gurley, 1883, New Carb. Foss.,
p. 4. Publication invalid.

deltoides, Gabb, 1859, Proc. Acad. Nat.
Sci. Phil., p. 297, Subcarboniferous.

imbricaria, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 412, Marshall Gr.

iowensis, Winchell, 1865, Proc. Acad.
Nat. Sci., p. 127, Chemung Gr.

kansasensis, Shumard, 1858, Trans. St.
Louis Acad. Sci., vol. 1, p. 213, Coal

Meas.

keokuk, Worthen, 1875, Geo. Sur. Ill., vol.
6, p. 524, Keokuk Gr.

meliniformis, Meek & Worthen, 1866,
Proc. Chi. Acad. Sci., vol. 1, p. 19, Coal

Meas.

michiganensis, Winchell, 1862, Proc.
Acad. Nat. Sci., p. 411, Marshall Gr.

monroensis, Worthen, 1884, Bull. No. 2,
Ill. St. Mus. Nat. Hist., p. 15, and

Geo. Sur. Ill., vol. 8, p. 127, St.
Louis Gr.

mytiliformis, Hall, 1852, Pal. N. Y., vol.
2, p. 100, Clinton Gr.

nemesis, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 3, p. 237, Subcarbonif-

erous.

nessus, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 238, Subcarbonif-

erous.

perattenuata, Meek & Hayden, 1858,
Trans. Alb. Inst., vol. 4, p. 77, and

Geo. Sur. Ill., vol. 5, p. 582, Coal
Meas.

permiana, Swallow, 1858, (*Mytilus per-*
manus.) Trans. St. Louis Acad. Sci.,

vol. 1, p. 187, and Pal. Up. Mo., p. 52,
Permian Gr.

perniformis, Cox, 1857, Geo. Sur. Ky.,
vol. 3, p. 569, Coal Meas.

perineiformis, Winchell, 1862, Proc.
Acad. Nat. Sci., p. 412, Marshall Gr.

rara, Winchell, 1870, Proc. Am. Phil.
Soc., p. 390, Marshall Gr. Proposed

instead of M. aviculoidea, Winchell,
1862, which was preoccupied.

recta, Shumard, 1858, Trans. St. Louis
Acad. Sci., vol. 1, p. 212, Permian Gr.

recurvirostris, Meek & Worthen, 1860,
Proc. Acad. Nat. Sci. Phil., p. 456, and

Geo. Sur. Ill., vol. 2, p. 344, Up. Coal
Meas.

squamosa, Sowerby, 1827, Trans. Geo.
Soc. Lond., 2d ser., vol. 3, Permian Gr.
subquadrata, Shumard, 1855, Geo. Rep.
Mo., p. 207, Coal Meas.

FIG. 860.—*Myalina recurvirostris*.

stludovici, Worthen, 1873, Geo. Sur. Ill.,
vol. 5, p. 540, St. Louis Gr.

swalovi, McChesney, 1860, New Pal. Foss.,
p. 57, and Geo. Sur. Ill., vol. 2, p. 341,
Coal Meas.

FIG. 861.—*Myalina subquadrata*.

MYTILARCA, Hall, 1870, Prelim. Notice Lam.
Shells, p. 19. [Ety. from the two gen-
era *Mytilus* and *Arca*.] Equivalve, in-
equilateral, mytiliform; beaks terminal;
hinge short; ligamental area striated;
cardinal teeth beneath the beak small,
oblique; posterior teeth small, oblique,
and at the extremity of the hinge; an-
terior muscular scar umbonal, and pos-
terior one near the postero-basal mar-
gin; pallial line entire, simple; surface
not unfrequently with fine, obscure ra-
diating striae. Type M. chemungensis.

arenacea, see *Plethomytilus arenaceus*.
attenuata, Hall, 1870, Prelim. Notice Lam. Shells, p. 23, and Pal. N. Y., vol. 5, pt. 1, p. 260, Chemung Gr.
canadensis, Billings, 1874, Pal. Foss., vol. 2, p. 52, Gaspe limestone No. 8, Devonian.
carinata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 259, Chemung Gr.
chemungensis, Conrad, 1842, (*Inoceramus chemungensis*,) Jour. Acad. Nat. Sci. Phil., vol. 8, p. 246, and Pal. N. Y., vol. 5, pt. 1, p. 258, Chemung Gr.

FIG. 862.—*Mytilarca chemungensis*.

cordiformis, Hall, 1859, (*Megambonia cordiformis*,) Pal. N. Y., vol. 3, p. 278, Low. Held. Gr.
dubia, Walcott, 1885, Monogr. U. S. Geol. Sur., vol. 8, p. 168, Devonian.
fibristriata, White & Whitfield, 1862, (*Mytilus fibristriatus*,) Proc. Bost. Soc. Nat. Hist., vol. 8, p. 296, and Pal. N. Y., vol. 5, pt. 1, p. 264, Kinderhook Gr.
gibbosa, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 262, Up. Chemung Gr.
knappi, see *Plethomytilus knappi*.
lata, Hall, 1884, (*Mytilops lata*,) Pal. N. Y., vol. 5, pt. 1, p. 262, Chemung Gr.
mytilimeris, see *Plethomytilus mytilimeris*.
nitida, Billings, 1874, Pal. Foss., vol. 2, p. 53, Gaspe limestone No. 8, Devonian.
occidentalis, White & Whitfield, 1862, (*Mytilus occidentalis*,) Proc. Bost. Soc. Nat. Hist., vol. 8, p. 297, and Pal. N. Y., vol. 5, pt. 1, p. 263, Kinderhook Gr.
oviformis, see *Plethomytilus oviformis*.
percarinata, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 202, Up. Held. Gr.
ponderosa, see *Plethomytilus ponderosus*.
pyramidata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 256, Schoharie grit.
radiata, see *Byssopteria radiata*.
regularis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 260, Chemung Gr.
sigillum, Hall, 1870, 28th Rep. N. Y. Mus. Nat. Hist., p. 174, Niagara Gr.
simplex, Hall, 1884, (*Mytilops simplex*,) Pal. N. Y., vol. 5, pt. 1, p. 261, Chemung Gr.
umbonata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 257, Chemung Gr.
MYTILOPS, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 4. [Ety. from resemblance to

Mytilus.] Resembles externally *Modiola* and *Lithodomus*. Hinge-line narrow, oblique, extending about half the length of the shell, beaks terminal. Type *M. præcedens*.

FIG. 863.—*Mytilops precedens*. Left valve.

lata, see *Mytilarca lata*.
metella, Hall, 1870, Prelim. Notice Lam. Shells, p. 1, and Pal. N. Y., vol. 5, pt. 1, p. 268, Chemung Gr.

FIG. 864.—*Mytilops precedens*.

præcedens, Hall, 1870, (*Modiola præcedens*,) Prelim. Not. of Lam. Shells, p. 1, and Pal. N. Y., vol. 5, pt. 1, p. 267, Chemung Gr.

simplex, see *Mytilarca simplex*.

Mytilus, Linnæus, 1758, Syst. Nat., 10th ed. [Ety. *Mytilus*, the fish mussel.] This genus does not, so far as known, exist in Palæozoic rocks. Most of the species referred to it belong to *Myalina* and *Mytilarca*.

concavus, see *Myalina concava*.

fibristriatus, see *Mytilarca fibristriata*.

occidentalis, see *Mytilarca occidentalis*.

ottawensis, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 211, Up. Coal Meas.

permianus, see *Myalina permiana*.

squamosus, Sowerby, 1839, Trans. Geol. Soc. Lond., vol. 4, Permian Gr. Probably not American.

tenuiradiatus, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 211, Up. Coal Meas.

whitfieldanus, Winchell, 1862, Proc. Acad. Nat. Sci., p. 413, syn. for *Mytilarca fibristriata*.

Naïadites, Dawson, 1860, Acad. Geol., but not defined. The name was used for a genus of plants by Buckman in 1843. The fossils were defined by Salter in 1861, under the name of *Anthracomya angulatus*, see *Anthracomya angulata*.

arenaceus, see *A. arenacea*.

carbonarius, see *A. carbonaria*.

elongatus, see *A. elongata*.

levis, see *A. levis*.

obtusius, see *A. obtusa*.

ovalis, see *A. ovalis*.

NUCULA, Lamarck, 1801, Syst. An. sans Vert., p. 87. [Ety. *nucula*, a little nut.] Equivalve, inequilateral oval, or oblong

ternally Modiola
ange-line narrow,
at half the length
minimal. Type M.

ens. Left valve.

dim. Notice Lam.
N. Y., vol 5, pt.

præcedens.

(Modiola præce-
Lam. Shells, p. 1,
5, pt. 1, p. 267,

plex.

yst. Nat., 10th ed.
h mussel.] This
as known, exist
lost of the species
to Myalina and

ncava.

a fibristriata.
a occidentalis.
1858, Trans. St.
1, p. 211, Up. Coal

permiana.

39, Trans. Geol.
rmian Gr. Prob-

, 1858, Trans. St.
p. 211, Up. Coal

1862, Proc. Acad.
Mytilarca fibri-

Acad. Geol., but
ne was used for a
Buckman in 1843.
ined by Salter in
of Anthracomya.
mya angulata.

ca.

onaria.

a.
yst. An. sans Vert.,
a, a little nut.]
ral oval, or oblong

closed all round, without external liga-
mentary facets; beak directed back-
ward; cartilage internal, placed in a
pit under the beak; teeth numerous,
very long. Type *N. nucleus*.

anodontoides, Meek, 1871, Reg. Rep. Uni-
versity W. Va. Coal Meas.

arata, see *Nuculana arata*.

bellatula, Hall, 1843, syn. for *N. belli-*
striata.

bellistriata, Conrad, 1841, (*Nuculites*
bellistriatus.) Ann. Rep. N. Y., p. 40,
and Geo. Rep. 4th Dist. N. Y., p. 197,
Ham. Gr.

beyrichia, Schlotheim, as identified by
Geinitz. See *Nucula parva*.

corballiformis, Hall, 1870, Prelim. Notice
Lam. Shells, p. 2, and Pal. N. Y., vol. 5,
pl. 46, figs. 24-37, Ham. and Chemung
Grs.

FIG. 865.—*Nucula*
cobboldie.

cylindricus, syn. for
Cardiomorpha mis-
souriensis.

diffidens, Hall, 1885,
Pal. N. Y., vol. 5, p.

322, Chemung Gr.

donaciformis, see *Tellinomya donaciformis*.

globularis, Hall, 1885,
Pal. N. Y., vol. 5, p.

322, Chemung Gr.

fabula, see *Clidophorus fabula*.

hians, Hall, 1860, 13th Rep. N. Y. Mus.
Nat. Hist., p. 110, Kinderhook Gr.

houghtoni, see *Tellinomya houghtoni*.

hubbardi, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 417, Marshall Gr. Syn. for
Nuculites sulcatinus.

illinoisensis, Worthen, 1884, Bull. No. 2,
Ill. St. Mus. Nat. Hist., p. 15, and Geo.
Sur. Ill., vol. 8, p. 128, St. Louis Gr.

insularis, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 241, Carbonifer-

ous.

iowensis, White & Whitfield, 1862, Proc.
Acad. Nat. Sci., vol. 8, p. 298. Syn. for
Tellinomya houghtoni.

kazanensis, as identified by Geinitz is *Nu-*
culana bellistriata.

lamellata, Hall, 1883, Pal. N. Y., vol. 5,
pl. 51, figs. 18-20, Ham. Gr.

levata, see *Tellinomya levata*.

levatiformis, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 241, Carbonifer-

ous.

lineata, see *Tellinomya lineata*.

lineolata, Hall, 1843, Geo. Rep. 4th Dist.
N. Y., p. 246, Portage Gr.

lirata, Conrad, 1842, (*Nuculites liratus*.)
Jour. Acad. Nat. Sci., vol. 8, p. 250, and

Pal. N. Y., vol. 5, pl. 45, figs. 17-27,
Ham. Gr.

machariiformis, see *Tellinomya machari-*
formis.

mactriiformis, see *Tellinomya mactriiformis*.

mercensis, syn. for *Cardiomorpha mis-*
souriensis.

microdonta, Winchell, 1863, Proc. Acad.
Nat. Sci., p. 16, Marshall Gr.

minima, Foerste, 1885, Bull. Sci. Lab.
Denison Univ., p. 93. Not properly
defined

minuta, Owen, 1840, Rep. on Min. Lands,
Devonian. The name was preoccupied
by De France in 1825.

nasuta, see *Nuculana nasuta*.

neda, Hall & Whitfield, 1872, 24th Rep.
N. Y. Mus. Nat. Hist., p. 191, Up.
Held. Gr.

niotica, Hall & Whitfield, 1872, 24th
Rep. N. Y. Mus. Nat. Hist., p. 190,
Ham. Gr.

obliqua, see *Palaeoconcha obliqua*.

oblonga, Hall, syn. for *Nuculites oblon-*
gatus.

obsoleta, McChesney, 1860, Pal. Foss., p. 89,
Coal Meas. Not recognized.

parva, McChesney, 1860, New Pal. Foss.,
p. 54, and Geo. Sur. Ill., vol. 5, p. 589,
Coal Meas.

perumbonata, White, 1879, Bull. U. S.
Geo. Sur., vol. 5, No. 2, p. 217, and
Cont. to Pal., No. 6, p. 136, Carbonif-

erous.

poststriata, see *Lvrodosma poststriatum*.

randalli, Hall, 1870, Prelim. Notice Lam.
Shells, p. 3, and Pal. N. Y., vol. 5, pl.

45, figs. 5-16, Ham. and Chemung
Grs.

rectangula, McChesney, 1860, Desc. New
Pal. Foss., p. 74, Ham. Gr.

rescuensis, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 172, Devonian.

sectoralis, Winchell, 1862, Proc. Acad.
Nat. Sci., p. 418, Marshall Gr.

shumardiana, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 16, and Bull. Am. Mus. Nat.

Hist., p. 57, Warsaw Gr.

stella, see *Tellinomya stella*.

subelliptica, Hall, 1883, Pal. N. Y., vol. 5,
pl. 45, fig. 23, Ham. Gr.

umbonata, Hall,
1883, Pal. N. Y.,
vol. 5, pl. 47, figs.

51 and 52, Che-
mung Gr.

varicosa, Hall, 1870,
Prelim. Notice
Lam. Shells, p. 2, and Pal. N. Y., vol. 5,
pl. 46, figs. 12-23, Ham. Gr.

ventricosa, Hall, 1858, Geo. Sur. Iowa, p.
716, Coal Meas.

NUCULANA, Link, 1807, Rost. Samml., vol. 3,
p. 155. [Ety. like a shell of the genus

Nucula.] Equivalve, inequilateral, pro-
duced behind; beaks sometimes di-
rected posteriorly; lunule often present;
rounded in front; post-umbonal slope
defined; surface concentrically lined;
hinge with a line of small teeth inter-
rupted by a triangular cartilage pit be-
neath the beak; muscular impressions
two, small; pallial line, simple, or
slightly sinuous. Type *N. emargin-*
ata.

arata, Hall, 1852, (*Nucula arata*.) Stansb.
Exped. to Gt. Salt Lake, p. 413, Coal
Meas.

FIG. 866.—*Nucula ventri-*
cosa.

bellistriata, Stevens, 1858, (*Leda bellistriata*), Am. Jour. Sci., vol. 25, p. 261, and Geo. Sur. Iowa, p. 717, Coal Meas.

FIG. 867.—*Nuculana bellistriata*. Left valve.

Prelim. Notice Lam. Shells, p. 6, and Pal. N. Y., vol. 5, pls. 38, 39, Ham. Gr.

curta, Meek, 1861, (*Leda curta*), Proc. Acad. Nat. Sci. Phil., p. 144, and Geo.

Sur. Ill., vol. 2, FIG. 868.—*Nuculana bellistriata*. Cardinal view. Gr.

densmamillata, Stevens, 1858, (*Leda densmamillata*), Am. Jour. Sci., vol. 25, p. 261, Marshall Gr.

diversa, Hall, 1883, Pal. N. Y., vol. 5, pl. 47, figs. 31–37, Ham. Gr.

nasuta, Hall, 1858, (*Nucula nasuta*), Trans. Alb. Inst., vol. 4, p. 17, and Bull. Am. Mus. Nat. Hist., p. 57, Warsaw Gr.

nuculiform's, see *Palæoneilo nuculiformis*. obesa, White, 1879, Bull. U. S. Geo. Sur., vol. 5, No. 2, p. 216, and Cont. to Pal., No. 6, p. 136, Carboniferous.

ohioensis, Hall, syn. for *N. pandoriformis*. obscura, Hall, 1885, (*Leda obscura*), Pal. N. Y., vol. 5, p. 331, Ham. Gr.

pandoriformis, Stevens, 1858, (*Leda pandoriformis*), Am. Jour. Sci., vol. 25, p. 261, Waverly Gr.

perstriata, Hall, 1883, Pal. N. Y., vol. 5, pl. 47, figs. 42–44, syn. for *N. rostellata*.

rostellata, Conrad, 1841, (*Nuculites rostellatus*), Ann. Rep. Geo. N. Y., p. 50, Ham. Gr.

saccata, Winchell, 1863, (*Leda saccata*), Proc. Acad. Nat. Sci., p. 16, Marshall Gr.

vaseyana, McChesney, 1860, (*Nuculites vaseyanus*), Desc. New. Pal., Foss., p. 73, Ham. Gr.

NUCULITES, Conrad, 1841, Ann. Geo. Rep. N. Y., p. 49. [Ety. *Nucula*, a genus of shells.] Equivalve, inequilateral, transverse; anterior end rounded; posterior truncate or pointed; beak, anterior; cardinal line arcuate; post-umbonal slope rounded or angular; surface concentrically lined, hinge with a row of transverse narrow teeth from the anterior to the posterior muscular scar; ligament external; anterior scar deep and separated from the cavity of the shell by a clavicle; posterior scar elongate; pallial line simple. Type *N. oblongatus*.

altus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 251, Devonian.

appressus, see *Cytherodon appressus*.

bellistriatus, see *Nucula bellistriata*.

carinatus, Hall, 1860, Can. Nat. and Geol., vol. 5, p. 151, Up. Sil.

chemungensis, see *Cytherodon chemungensis*.

concentricus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 248, Coal Meas.

constrictus, see *Palæoneilo constrictus*.

cuneiformis, Conrad, 1841, Ann. Rep. N. Y., p. 50, and Pal. N. Y., vol. 5, pl. 47, figs. 13–16, Ham. Gr.

emarginatus, see *Palæoneilo emarginatus*.

faba, see *Modiolopsis faba*.

filosus, see *Palæoneilo filosa*.

inflatus, see *Cypricardites inflatus*.

lamellosus, Conrad, 1841, Ann. Geo. Rep. N. Y., p. 50, Up. Sil.

liratus, see *Nucula lirata*.

mactroides, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 249, Marshall Gr.

maximus, see FIG. 869.—*Nuculites oblongatus*. Interior of large left valve.

maxima, see FIG. 870.—*Nuculites oblongatus*. Cast of interior of right valve, showing muscular scars and pallial line.

multilineatus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 251, Ham. Gr.

FIG. 870.—*Nuculites oblongatus*. Cast of interior of right valve, showing muscular scars and pallial line.

nyssa, Hall, 1870, Prelim. Notice Lam. Shells, p. 5, and Pal. N. Y., vol. 5, pl. 47, figs. 25–30, Ham. Gr.

oblongus, see *Clidophorus oblongus*.

oblongus, see *Clidophorus oblongus*.

oblongatus, Conrad, 1841, Ann. Geo. Rep. N. Y., p. 50, and Pal. N. Y., vol. 5, pl. 47, figs. 1–12, Ham. Gr.

planulatus, see *Clidophorus planulatus*.

poststriatus, see *Lyrodessa poststriatum*.

radiatus, see *Pholadella radiata*.

rostellatus, see *Nuculana rostellata*.

scitula, syn. for *Clidophorus planulatus*.

submarginatus, see *Tellinopsis*, submarginata.

sulcatinus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 250, Marshall Gr.

triangularis, Hall & Whitfield, 1877, U. S. Geo. Expl., 40th parallel, vol. 4, p. 248, Devonian.

triqueter, Conrad, 1841, Ann. Rep. N. Y., p. 50, and Pal. N. Y., vol. 5, pl. 47, figs. 17–24, Ham. Gr.

vaseyanus, see *Nuculana vaseyana*.

yoldiiformis, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 24, Had. Riv. Gr.

It is not a *Nuculites*.

NYSSA, Hall, 1870, Prelim. Notice Lam. Shells, but defined by Whitfield, 1882, Ann. N. Y., Acad. Sci., vol. 2, p. 244.

[Ety. mythological name.] Shell bivalve, very oblique and transversely ovate in form; posterior hinge plate

pellistriata.
n. Nat. and Geol.,
rodon chemung-
1842, Jour. Acad.
Coal Meas.
o constricta.
1841, Ann. Rep.
N. Y., vol. 5, pl.
r.
ilo emarginata.
sa.
sa.
s inflatus.
, Ann. Geo. Rep.

—Nuculites oblon-
Interior of large left

1842, Jour. Acad.
1, Ham. Gr.
nyssa, Hall, 1870,
Prelim. Notice
Lam. Shells, p.
5, and Pal. N.
Y., vol. 5, pl.
47, figs. 23-30,
Ham. Gr.
oblongus, see Cli-
dophorus ob-
longus.
oblongatus, Con-
Rep. N. Y., p. 50,
p. 47, figs. 1-12,

rus planulatus.
ma poststriatum.
radiata.
rostellata.
orus planulatus.
nopsis, subemargi-

2, Jour. Acad. Nat.
Marshall Gr.
Whitfield, 1877, U. S.
illel, vol. 4, p. 248,

Ann. Rep. N. Y.,
vol. 5, pl. 47, figs.

vaseyana.
79, Jour. Cin. Soc.
24, Hud. Riv. Gr.

im. Notice Lam.
Whitfield, 1882,
Sci., vol. 2, p. 244.
name.] Shell bi-
and transversely
terior hinge plate

narrow, bearing from one to four long, slender, ridge-like teeth; anterior plate broad, marked by numerous, small, point-like teeth, with intermediate depressions, arranged somewhat radiating from the middle of its inner border; adductor muscles two, one at each extremity; pallial line entire; ligament internal. Type *N. arguta*.

FIG. 871.—*Nyassa arguta*. Left valve.

arguta, Hall, 1870, Prelim. Notice Lam. Shells, p. 28, and Pal. N. Y., vol. 5, pl. 53, figs. 9-20, Ham. Gr.
elliptica, Hall, 1870, Prelim. Notice Lam. Shells, p. 30, and Pal. N. Y., vol. 5, pl. 34, fig. 8, Up. Held. Gr.

FIG. 872.—*Nyassa arguta*. Interior of right valve.

parva, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 173, Devonian.
recta, Hall, 1870, Prelim. Notice Lam. Shells, p. 29, and Pal. N. Y., vol. 5, pl. 53, figs. 1-8, Ham. Gr.
subulatu, Hall, 1870, Prelim. Notice Lam. Shells, p. 29, and Pal. N. Y., vol. 5, pl. 53, figs. 21-26, Ham. Gr.

Opisthoptera, Meek. Not defined.
Orthodesma, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 93. [Ety. *orthos*, straight; *desma*, a ligament.] Elongated, ventricose; cardinal line straight posterior to the beaks, and contracted anterior; ligament external; posterior scar elongate, anterior smaller; pallial line simple; surface concentrically lined. Type *O. rectum*.

byrnesi, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 76, Hud. Riv. Gr.
contractum, Hall, 1847, (Orthonota *contracta*.) Pal. N. Y., vol. 1, p. 300, Hud. Riv. Gr.
cuneiforme, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 314, Hud. Riv. Gr.

curvatum, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 95, Hud. Riv. Gr.

faberi, n. sp. Shell large, posterior part of the cardinal line very slightly elevated, and anterior part contracted in front of the beaks; anterior end rounded, posterior end more abrupt;

basal line contracted in the central part by an undefined cincture arising below the umbones; beaks anterior, obtuse; umbones low and poorly defined; shell unusually high and thin for species in this genus; surface concentrically furrowed. This species bears some resemblance to a *Modiolopsis*, but it is doubtless an *Orthodesma*. Collected by Mr. Charles Faber in the upper part of the Hud. Riv. Gr., at Versailles, Indiana.

FIG. 873.—*Orthodesma faberi*.

mickelboroughi, Whitfield, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 139, Hud. Riv. Gr.

occidentale, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 3, p. 316, Hud. Riv. Gr.

parallelum, Hall, 1847, (Modiolopsis *parallela*.) Pal. N. Y., vol. 1, p. 158, Hud. Riv. Gr.

rectum, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 94, Hud. Riv. Gr.

FIG. 874.—*Orthodesma rectum*.

subovale, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 82, Hud. Riv. Gr.

Orthonota, Conrad, 1841, Ann. Rep. N. Y., p. 50. [Ety. *orthos*, straight; *notos*, the back.] Transversely elongate; margins subparallel; cardinal line straight; two cardinal teeth; no lateral teeth; ligament external; umbonal ridge oblique. Type *O. undulata*.
angulifera, (?) McCoy, 1850, Brit. Pal. Rocks, p. 276, Up. Sil.

carinata, Conrad, 1841, Ann. Rep. N. Y., p. 51, and Pal. N. Y., vol. 5, pl. 78, figs. 34-35, Ham. Gr.

contracta, see *Orthodesma contractum*.
curta, Hall, 1845, Geo. Rep. 4th Dist. N. Y., p. 76, and Pal. N. Y., vol. 2, p. 86, Clinton and Niagara Gr.

ensiformis, Hall, 1870, Prelim. Notice Lam. Shells, p. 89, and Pal. N. Y., vol. 5, pl. 78, fig. 36, Ham. Gr.

incerta, Billings, 1874, Pal. Foss., vol. 2, p. 130, Up. Sil.

parallela, see *Orthodesma parallelum*.

parvula, Hall, 1870, Prelim. Notice Lam. Shells, p. 88, and Pal. N. Y., vol. 5, pl. 78, figs. 29-32, Ham. Gr.
 phaselia, Winchell, 1863, Proc. Acad. Nat. Sci., p. 12, Marshall Gr.
 pholadia, Conrad, 1838, (Pterinea pholadis,) Ann. Geo. Rep. N. Y., p. 118, Hud. Riv. Gr.
 rectidorsalis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 412, Marshall Gr.
 rigida, Hall, 1885, Pal. N. Y., vol. 5, p. 481, Chemung Gr.
siliquoidea, see *Palaeosolen siliquoideus*.
simulans, Billings, 1874, Pal. Foss., vol. 2, p. 131, Up. Sil.
 (?) *speciosa*, Billings, 1874, Pal. Foss., vol. 2, p. 130, Up. Sil.
undulata, Conrad, 1841, Ann. Rep. N. Y., p. 51, and Pal. N. Y., vol. 5, pl. 78, figs. 37-42, Ham. Gr.

FIG. 875.—*Orthonota undulata*.

ventricosa, see *Spathella ventricosa*.
venusta, Billings, 1874, Pal. Foss., vol. 2, p. 129, Up. Sil.

ORTHONOTELLA, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 117. [Ety. *orthos*, straight; *notos*, back; *ellus*, diminutive.] Very small, inequilateral, inequivalve, more or less elliptical; beak anterior; surface smooth or concentrically lined; hinge straight behind the beaks; ligament external. Type *O. faberi*.

FIG. 876.—*Orthonotella faberi*. Greatly enlarged.

faberi, S. A. Miller, 1882, Jour. Cin. Soc. Nat. Hist., vol. 5, p. 117, Hud. Riv. Gr.

Ostrea, Linnaeus, 1758, Syst. Nat. 10th ed., p. 696. [Ety. *ostrea*, an oyster.] Inequivalve, irregular in shape, with a single adductor muscle. Not a Palaeozoic genus, though a species has been founded upon a single valve and called *O. patercula*.

patercula, Winchell, 1865, Proc. Acad. Nat. Sci., p. 124, and 4th Ann. Rep. U. S. Geo. Sur., p. 288, Burlington Gr.
 PALAENATINA, Hall, 1870, Prelim. Notice Lam. Shells, p. 84. [Ety. *palaaios*, ancient; *Anatina*, a genus.] Transversely elongate; gaping; left valve the larger; oblique constriction; hook-like process anterior to the beaks; no lateral

teeth; surface concentrically lined. Type *P. typus*.

angusta, Hall, 1885, Pal. N. Y., vol. 5, p. 490, Chemung Gr.

quadrata, see *Prorhynchus quadratum*.

sinuata, Hall, 1885, Pal. N. Y., vol. 5, p. 491, Chemung Gr.

solenoides, Hall, 1885, Pal. N. Y., vol. 5, p. 489, Chemung Gr.

typus, Hall, 1870, Prelim. Notice

Lam. Shells, p. 85, and Pal.

N. Y., vol. 5, pl.

79, figs. 26-39,

Chemung Gr.

Palaearca, syn. for

Cypricardites.

saffordi, see *Cypricardites saffordi*.

ventricosa, see *Cypricardites ventricosus*.

PALAEOCARDIA, Hall, 1867, 20th Rep. N. Y.

Mus. Nat. Hist., p. 389. [Ety. *palaaios*,

ancient; *kardia*, a heart.] Cordiform,

obliquely subovate, ventricose; um-

bones gibbous; beaks prominent in-

curved; hinge-line very short; surface

marked with radiating striae. Type *P.*

cordiformis.

cordiformis, Hall, 1867, 20th Rep. N. Y.

Mus. Nat. Hist., p. 389, Niagara Gr.

PALAECONCHA, n. gen. [Ety. *palaaios*, an-

cient; *conche*, shell.] Shell small, equi-

valve, inequilateral, oblique, varying

from subcircular or subovoid to mytili-

form; height equal to or greater than

length; closed all around; without ex-

ternal evidence of ligaments; anterior

side more or less truncated and round-

ing into the base below; beaks ele-

vated, projecting beyond cardinal line

without incurving; cardinal line

straight or slightly arching; some evi-

dence points to an anterior and pos-

terior muscular scar near the ends of

the cardinal line; pallial line simple;

no lateral teeth and probably edentu-

lous; surface smooth. Type *P. faberi*.

faberi, n. sp. Shell small, height greater

than length, very slightly oblique,

closed all around; beaks projecting

high above the hinge-line without

incurving; sur-

face smooth,

variable in

size; a large

specimen has a

height of 0.20

inch, and

length 0.14

inch, a small

specimen is

about two-

thirds less. This species is distin-

guished from *P. obliqua* by its greater

proportional height, more prolonged

beak, less oblique form, and generally

larger size. Collected in the upper

part of the Hud. Riv. Gr., at Versailles,

Indiana, and in Butler County, Ohio.

FIG. 877.—*Palaenatina typus*. FIG. 878.—*Palaeoconcha faberi*. Magnified 5 diam.

entrically lined.

N. Y., vol. 5,

is quadratum.

N. Y., vol. 5, p.

l. N. Y., vol. 5,

77.—Palaeonella typus.

saffordi.

es ventricosus.

20th Rep. N. Y.

p. [Ety. palaios,

tr.] Cordiform,

ventricose; um-

y short; surface

striae. Type P.

20th Rep. N. Y.

Niagara Gr.

Ety. palaios, an-

Shell small, equi-

oblique, varying

obovoid to mytili-

or greater than

nd; without ex-

aments; anterior

ed and rounded

below; beaks ele-

nd cardinal line

cardinal line

ching; some evi-

terior and pos-

near the ends of

lial line simple;

probably edentu-

Type P. taberi.

ll, height greater

slightly oblique,

beaks projecting

hinge-line without

incurving; sur-

face smooth,

variable in

size; a large

specimen has a

height of 0.20

inch, and

length 0.14

inch, a small

specimen is

about two-

species is distin-

gua by its greater

more prolonged

m, and generally

d in the upper

Gr., at Versailles,

r County, Ohio.

obliqua, Hall, 1845, (Nucula obliqua,) Am. Jour. Sci., vol. 48, p. 292, and Ohio Pal., vol. 1, p. 139, Hud. Riv. Gr. PALÆONEILLO, Hall, 1870, Prelim. Notice Lam. Shells, p. 6. [Ety. palaios, ancient; Neilo, a genus.] Nuculiform, posterior end extended, sulcus along the umbonal slope; surface concentrically striated or ribbed; hinge-line arcuate, crenulate, not interrupted beneath the beak by a ligamental pit; ligament external; anterior and posterior adductor scars distant; pedal scars within the umbonal cavity. Type P. constricta.

FIG. 878.—Palaeonello bedfordensis.

attenuata, Hall, 1870, Prelim. Notice Lam. Shells, p. 12, and Pal. N. Y., vol. 5, pl. 50, figs. 34-39, Waverly Gr.

barrisi, White & Whitfield, 1862, (Leda barrisi,) Proc. Bost. Soc. Nat. Hist., vol. 8, p. 298, syn. for P. nuculiformis.

bedfordensis, Meek,

1875, Pal. Ohio, vol. 2,

p. 298, Waverly Gr.

bisulcata, Hall, 1870,

Prelim. Notice Lam.

Shells, p. 10, and Pal.

N. Y., vol. 5, pl. 50,

figs. 13-14, Ham. Gr.

brevis, Hall, 1870, Pre-

lim Notice Lam. Shells, p. 10; and Pal.

N. Y., vol. 5, pl. 50,

figs. 24-33, Chemung Gr.

carbonaria, see Yoldia carbonaria.

constricta, Conrad, 1842,

(Nuculites constrictus),

angusta, Hall,

1885, Pal. N. Y.,

vol. 5, p. 344,

Chemung Gr.

arata, Hall, 1883,

Pal. N. Y., vol.

5, pl. 50, fig. 23,

Ham. Gr.

FIG. 880.—Palaeo-

nello bedforden-

sis. Magnified.

FIG. 881.—Palaeonello constricta.

Left valve enlarged, showing

crenulations of hinge-line.

Jour. Acad. Nat. Sci., vol.

8, p. 249, and Pal. N. Y.,

vol. 5, pl. 48, figs. 1-15, Chemung Gr.

constricta var. flexuosa, Hall, 1883, Pal.

N. Y., vol. 5, pl. 48, figs. 16-20, Ham. Gr.

dubia, Hall, 1885, Pal. N. Y., vol. 5, p.

348, Up. Held. Gr.

elongata, Hall, 1883, Pal. N. Y., vol. 5, pl.

48, fig. 39, Chemung Gr.

emarginata, Conrad, 1841, (Nuculites

emarginata.) Ann. Rep. N. Y., p. 50,

and Geo. Wis., vol. 4, p. 337, Ham. Gr.

flosa, Conrad, 1842, (Nuculites flosus,) Jour. Acad. Nat. Sci., vol. 8, p. 250, and Pal. N. Y., vol. 5, pl. 49, figs. 33-38, Chemung Gr.

fecunda, Hall, 1870, Prelim. Notice Lam. Shells, p. 8, and Pal. N. Y., vol. 5, pl. 49, figs. 13-24, Ham. Gr.

maxima, Conrad,

1841, (Nuculites

maximus.) Ann.

Rep. N. Y., p.

50, and Pal.

N. Y., vol. 5, pl.

48, figs. 29-38,

Ham. Gr.

FIG. 882.—Palaeonello max-

ima. Right valve.

Prelim. Notice Lam. Shells, p. 8, and Pal.

N. Y., vol. 5, pl. 49, figs. 25-32, Ham. Gr.

nuculiformis, Stevens, 1858, (Leda nuculiformis,) Am. Jour. Sci. and Arts, 2d

ser., vol. 25, p. 262, Waverly Gr.

parallela, Hall & Whitfield, 1870, 23d Rep.

N. Y. Mus. Nat. Hist., p. 241, Wa-

verly Gr.

perplana, Hall, 1870, Prelim. Notice Lam.

Shells, p. 12, and Pal. N. Y., vol. 5, pl.

50, figs. 15-22, Ham. Gr.

plana, Hall, 1870, Prelim. Notice Lam.

Shells, p. 7, and Pal. N. Y., vol. 5, pl.

48, figs. 21-28, Ham. Gr.

similis, Whitfield, 1882, Ann. N. Y. Acad.

Sci., vol. 2, p. 217, Erie shale, Por-

tage (?) Gr.

tenuistriata, Hall, 1870, Prelim. Notice

Lam. Shells, p. 9, and Pal. N. Y., vol. 5,

pl. 49, figs. 1-12, Ham. Gr.

truncata, Hall, 1883, Pal. N. Y., vol. 5, pl.

50, figs. 40-41, Chemung Gr.

virginica, Hall, 1885, Pal. N. Y., vol. 5, p.

340, Ham. Gr.

PALEOPINNA, Hall, 1883, Pal. N. Y., vol. 5,

pt. 1, p. 4. (Plates and Explanations.)

FIG. 882.—Palaeonello max-

ima. Right valve.

FIG. 880.—Palaeo-

nello bedforden-

sis. Magnified.

FIG. 881.—Palaeonello constricta.

Left valve enlarged, showing

crenulations of hinge-line.

FIG. 883.—Palaeopinna flabellum.

[Ety. palaios, ancient; Pinna, a genus.] Shells large, gaping in front; hinge-line straight, ligamental area narrow, longitudinal groove and slight oblique furrow extending backward from the beak; beak anterior, terminal, directed forward; test more convex, and with finer rays than on the ordinary Pinna, and also finely marked with concentric striae of growth. Type P. flabellum.

- flabellum, Hall, 1884, Pal. N. Y., vol. 5, p. 1, p. 240, Oriskany Gr.
 recurva, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 241, Up. Held. Gr.
- PALÆOSOLEN, Hall, 1885, Pal. N. Y., vol. 5, p. 46. [Ety. *palaio*, ancient; *Solen*, a genus.] Shell in form like *Solen*; dorsal and ventral margins subparallel; anterior end short, rounded; posterior end elongate, truncate; gaping; beaks small, appressed; cardinal line straight; umbonal slope prominent; surface concentrically marked. Type *P. siliquoides*.
- siliquoides, Hall, 1870, (Orthonota siliquoides,) Prelim. Not. Lam. Shells, p. 89, and Pal. N. Y., vol. 5, p. 483, Ham. Gr.
- PANENKA, Barrande, 1881, Syst. Sil. d. l. Boheme, vol. 6, p. 128. Equivalve, inequilateral, elliptical or subcircular, beaks prominent, incurved; cardinal line straight or arcuate; test thin; surface concentrically lined.
- abrupta, Hall, 1885, Pal. N. Y., vol. 5, p. 423, Ham. Gr.
 alternata, Hall, 1885, Pal. N. Y., vol. 5, p. 416, Up. Held. Gr.
 costata, Hall, 1885, Pal. N. Y., vol. 5, p. 419, Marcellus Shale.
 degener, Hall, 1885, Pal. N. Y., vol. 5, p. 424, Ham. Gr.
 dichotoma, Hall, 1885, Pal. N. Y., vol. 5, p. 416, Schoharie grit.
 elevata, Conrad, 1848, (Monotis elevata,) Proc. Acad. Nat. Sci., vol. 3, p. 23, Chemung Gr.
 equilatera, Hall, 1885, Pal. N. Y., vol. 5, p. 419, Marcellus Shale.
 hero, Hall, 1885, Pal. N. Y., vol. 5, p. 418, Marcellus Shale.
 lincklaeni, Hall, 1885, Pal. N. Y., vol. 5, p. 420, Marcellus Shale.
 mollis, Hall, 1885, Pal. N. Y., vol. 5, p. 420, Marcellus Shale.
 multiradiata, Hall, 1885, Pal. N. Y., vol. 5, p. 417, Up. Held. Gr.
 potens, Hall, 1885, Pal. N. Y., vol. 5, p. 422, Ham. Gr.
 poulsoni, Conrad, 1848, (Monotis poulsoni,) Proc. Acad. Nat. Sci., vol. 3, p. 23, Chemung Gr.
 radians, Conrad, 1842, (Pterinea radians,) Jour. Acad. Nat. Sci., p. 252, and Pal. N. Y., vol. 5, p. 422, Ham. Gr.
 retusa, Hall, 1885, Pal. N. Y., vol. 5, p. 421, Ham. Gr.
- robusta, Hall, 1885, Pal. N. Y., vol. 5, p. 424, Portage Gr.
 speciosa, Hall, 1843, (Avicula speciosa,) Geo. Rep. 4th Dist. N. Y., p. 243, Portage Gr.
 ventricosa, Hall, 1885, Pal. N. Y., vol. 5, p. 417, Marcellus Shale.
- Panopea*, Menard de la Groye, 1807, Ann. du Mus. 9. [Ety. mythological name.] *cooperi*, see *Chenomya cooperi*.
- PARACARDIUM, Barrande, 1881, Syst. Sil. de la Boheme, vol. 6, p. 137. [Ety. *para*, allied to; *Cardium*, a genus.] Equivalve, inequilateral, subcircular or subelliptical; posterior side subtruncate; surface marked with fine radii and concentric striae; the margin of a small cardinal area under the beaks is crenulated.
- doris, Hall, 1885, (Cardiola doris,) Pal. N. Y., vol. 5, p. 428, Portage Gr.
- PARARCA, Hall, 1885, Pal. N. Y., vol. 5, p. 36. [Ety. *para*, allied to; *Arca*, a genus.] Equivalve, inequilateral, transversely subelliptical or rhomboidal; anterior end short, rounded; cardinal line about half the length of the valves, arching at the beaks; surface marked by radii and concentric striae; hinge narrow, with a series of minute crenulations. Type *P. venusta*.
- erecta, Hall, 1885, (Cardiola erecta,) Pal. N. Y., vol. 5, p. 432, Waverly Gr.
 neglecta, Hall, 1885, Pal. N. Y., vol. 5, p. 432, Waverly Gr.
 precedens, Hall, 1885, Pal. N. Y., vol. 5, p. 429, Up. Held. Gr.
 sao, Hall, 1885, (Cardiola sao,) Pal. N. Y., vol. 5, p. 430, Chemung Gr.
 transversa, Hall, 1885, (Cardiola transversa,) Pal. N. Y., vol. 5, p. 429, Chemung Gr.
 venusta, Hall, 1885, Pal. N. Y., vol. 5, p. 431, Chemung Gr.
- PARACYCLAS, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 171. [Ety. *para*, allied to; *Cyclas*, a genus.] Equivalve, subequilateral, suborbicular or subelliptical; anterior end regularly rounded; posterior end rounded or subtruncate, more produced than the anterior; beaks small and low; hinge-line short, post-cardinal slope sometimes subalate; surface marked concentrically; ligament supported internally on each side by a narrow plate, which leaves in the east two diverging grooves directed forward from the beak; muscular impression on the post-umbonal slope; pallial line a little within the margin of the shell. Type *P. elliptica*.
- billingsana, S. A. Miller, 1883, 2d Ed. Am. Pal. Foss., p. 311, Devonian. Proposed instead of *Lucina occidentalis*, Billings, 1859, Assiniboine and Sas. Ex. Exped., p. 187, figs. b and c, which name was preoccupied.
- chemungensis, Hall, 1885, Pal. N. Y., vol. 5, p. 443, Chemung Gr.
 elevata, Hall, 1883, Pal. N. Y., vol. 5, pl. 72, figs. 37 to 41, Schoharie grit.
 elliptica, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 171, and Pal. N. Y., vol. 5, pl. 72, figs. 23-30, Cornif. Gr.
 elliptica var. occidentalis, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat.

FIG. 884.—Panenka speciosa.

roye, 1807, Ann.
hological name.]

1881, Syst. Sil.
3, p. 137. [Ety.
dium, a genus.]
al, subcircular or
or side subtrun-
with fine radii
the margin of a
nder the beaks is

diola doris,) Pal.
ortage Gr.

N. Y., vol. 5, p. 36,
; Arca, a genus.]
eral, transversely
nchoidal; anterior
ardinal line about
e valves, arching
marked by radii
; hinge narrow,
ute crenulations.

diola erecta,) Pal.
Waverly Gr.

N. Y., vol. 5, p.

Pal. N. Y., vol. 5,

la sao,) Pal. N. Y.,
ng Gr.

, (Cardiola trans-
ol. 5, p. 429, Che-

N. Y., vol. 5, p.

, Geo. Rep. 4th

Ety. para, allied to;

nivalve, subequilat-

subelliptical; ante-

ended; posterior end

ate, more produced

beaks small and

post-cardinal slope

surface marked

ent supported in-

side by a narrow

in the cast two di-

ected forward from

impression on the

pallial line a little

of the shell. Type

r, 1883, 2d Ed. Am.

vonian. Proposed

identalis, Billings,

d Sas. Ex. Exped.,

, which name was

Hist., p. 189, and Pal. N. Y., vol. 5, pl.
72, figs. 31-33, Up. Held. Gr.

FIG. 885.—*Paracyclas elliptica* var. *occidentalis*.

erecta, Hall, 1885, Pal. N. Y., vol. 5, p.
445, Chemung Gr.

fissa, Hall, 1883, Pal. N. Y., vol. 5, pl. 72,
figs. 35, 36, Schoharie grit.

hamiltonensis, Winchell, 1866, (Lucina
hamiltonensis,) Rep. Low. Pen. Mich.,
p. 95, Ham. Gr.

ignota, Hall, 1883, Pal. N. Y., vol. 5, pl.
72, fig. 34, Chemung Gr.

lirata, Conrad, 1838, (Posidonia lirata,)
Ann. Rep. N. Y., p. 116, and Pal. N. Y.,
vol. 5, pl. 72, figs. 1-19, Corniferous Gr.

ohioensis, Meek, 1871,
Proc. Acad. Nat. Sci.
Phil., p. 62, and
Ohio Pal., vol. 1, p.
248, Cornif. Gr.

FIG. 886.—*Paracyclas peroccidensis*, Hall &
ohioensis. Whitfield, 1877,
U. S. Geo. Expl. 40th Parallel, vol. 4,
p. 248, Devonian.

retusa, Hall, 1843, (Lucina ? retusa,) Geo.
Rep. 4th. Dist. N. Y., p. 246, Por-
tage Gr.

rotunda, Hall, 1885, Pal. N. Y., vol. 5,
p. 444, Chemung Gr.

sabini, White, 1876, Proc. Acad. Nat. Sci.,
p. 31, Chemung Gr.

tenuis, Hall, 1883, Pal. N. Y., vol. 5, pl.
72, figs. 20-22, Ham. Gr.

varysburgensis, Williams, 1887, (Lucina
varysburgia,) Bull. 41, U. S. Geo.
Sur., Portage Gr.

FIG. 887.—*Pernopecten aviculatus*.

wyomingensis, Williams, 1887, Bull. 41,
U. S. Geo. Sur., Portage Gr.

Pecten, Mueller, 1776. This genus is un-
known in the Palaeozoic rocks.

acutialatus, see *Aviculopecten acutialatus*.
armigerus, see *A. armigerus*.

aviculatus, see *Pernopecten aviculatus*.
broadheadi, syn. for *Aviculopecten car-*
boniferus.

cancellatus, see *Aviculopecten cancellatus*.

carboniferus, see *A. carboniferus*.

clevelandicus, see *A. clevelandicus*.

coloradoensis, see *A. coloradoensis*.

convexus, see *A. convexus*.

crenulatus, see *Crenipecten crenulatus*.

dolabriformis, see *Aviculopecten dolabri-*
formis.

duplicatus, see *A. duplicatus*.

hallianus, D'Orbigny, 1847, syn. for *Avi-*
culopecten cancellatus.

halli, see *A. halli*.

hawni, Geinitz, 1866, Carb. und Dyas, p.
36, syn. for *A. carboniferus*.

missouriensis, see *A. missouriensis*.

neglectus, see *Euchondria neglecta*.

occidentalis, see *A. occidentalis*.

providencensis, see *A. providencensis*.

radialis, see *Pseudomonotis radialis*.

ringens, see *Aviculopecten ringens*.

striatus, see *A. striatus*.

tenuilineatus, see *Streblopteria tenuilineata*.

utahensis, see *Aviculopecten utahensis*.

Pernachactas, Castelnau, 1843, Syst. Sil., p.
44. Not recognized.

FIG. 888.—*Pernopecten limiformis*. Hinge-line.

PERNOPECTEN, Winchell, 1865, Proc. Acad.
Nat. Sci. Phil., p. 125. [Ety. from the
shells *Perna* and *Pecten*.] Shell like *Pecten*

FIG. 889.—*Pernopecten limiformis*.
aviculatus, Swallow
1858, (Pecten avicu-
lus,) Trans. St. Louis

Acad. Sci., p. 213, and
Geo. Sur. Ill., vol. 5, p. 588, Coal Meas

cooperensis, Shu-
mard, 1885, (Avicu-
la cooperensis,)
Geo. Rep. Mo., p.
206, Waverly or
Choteau Gr.

fasciculatus, see *Ly-*
riopecten fasciatus.

limiformis, White &
Whitfield, 1862,
(*Aviculopecten li-*
maformis.) Proc.

Bost. Soc. Nat.
Hist., vol. 3, p. 295, Marshall Gr.

limatus, Winchell, 1865, Proc. Acad. Nat.
Sci., p. 126, Marshall Gr.

FIG. 890.—*Pernopecten shumardianus*.

- shumardanus, Winchell, 1865, Proc. Acad. Nat. Sci. Phil., p. 126, and Geo. Sur. Ill., vol. 2, p. 453, Kinderhook Gr.
- PHOLADELLA**, Hall, 1870, Prelim. Notice Lam. Shells, p. 63. [Ety. diminutive of the recent genus *Pholas*.] Equivalve, elongated; valves inflated; beaks anterior, incurved; basal margin constricted; escutcheon and lunule; surface ribbed. Type *P. newberryi*.
- constricta*, Hall, 1883, Pal. N. Y., vol. 5, pl. 78, figs. 26-27, Ham. Gr.
- cuneata*, see *Promacrus cuneatus*.
- decussata*, Hall, 1883, Pal. N. Y., vol. 5, pl. 78, fig. 28, syn. for *Promacrus cuneatus*.
- newberryi*, Hall, 1870, Prelim. Notice Lam. Shells, p. 65, and Pal. N. Y., vol. 5, pl. 78, fig. 25, Waverly Gr.

FIG. 891.—*Pholadella newberryi*.

- ornata*, Hall, 1870, Prelim. Notice Lam. Shells, p. 64, syn. for *P. radiata*.
- parallela*, Hall, 1883, Pal. N. Y., vol. 5, pl. 78, figs. 22-24, Ham. Gr.
- radiata*, Conrad, 1842, (*Nuculites radiatus*), Jour. Acad. Nat. Sci., vol. 8, p. 248, and Pal. N. Y., vol. 5, pl. 78, figs. 15-21, Ham. Gr.
- truncata*, Hall, 1870, Prelim. Notice Lam. Shells, p. 64, syn. for *P. radiata*.
- Pholadomya elongata*, see *Allorisma elongatum*.
- PHTHONIA**, Hall, 1870, Prelim. Notice Lam. Shells, p. 70. Equivalve, elongate-ovate, wider posteriorly; beaks obscure; surface radiated and concentrically marked; no teeth; ligament external. Type *P. sectifrons*.
- cylindrica*, Hall, 1883, Pal. N. Y., vol. 5, pl. 78, figs. 1-4, Ham. Gr.
- lirata*, Hall, 1883, Pal. N. Y., vol. 5, pl. 78, fig. 14, Ham. Gr.
- nitida*, Hall, 1885, Pal. N. Y., vol. 5, p. 477, Chemung Gr.
- nodocostata*, Hall, 1870, Prelim. Notice Lam. Shells, p. 71, and Pal. N. Y., vol. 5, pl. 78, figs. 5-9, Ham. Gr.

FIG. 892.—*Phthonia sectifrons*. Left valve.

- sectifrons*, Conrad, 1842, (*Cypricardites sectifrons*), Jour. Acad. Nat. Sci., vol. 8, p. 245, and Pal. N. Y., vol. 5, pl. 78, figs. 10-13, Ham. Gr.

- truncata*, Hall, 1885, Pal. N. Y., vol. 5, p. 478, Chemung Gr.
- PINNA**, Linnaeus, 1758, Syst. Nat. 10th Ed. [Ety. *pinna*, a wing.] Shell long, triangular, equivalve; beaks terminal, pointed; posterior end broad, truncate, gaping; a subtriangular, posterior muscular impression, and a small reniform one at the beaks; cartilage long, narrow, internal, supported by a slender ridge close within the cardinal edges; no teeth; shell of one internal laminated layer, and an external vertically fibrous layer. Type *P. squamosa*. A living genus that sometimes attains a length of two feet, and ranges from low water to sixty fathoms. It moves vertically, partly buried in sand, with knife-like edges erect. The byssus has been mixed with silk, spun and knit into gloves.
- adamsi*, syn. for *Pinna peracuta*.
- consimilis*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 230, Subcarboniferous.
- hinrichsana*, White & St. John, 1868, Trans. Chi. Acad. Sci., p. 122, St. Louis Gr.
- inexpectans*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 235, Subcarboniferous.
- ludlovi*, Whitfield, 1876, in Ludlow's Carroll to Yellowstone Park, p. 143, Coal Meas.
- marshallensis*, Winchell, 1865, Proc. Acad. Nat. Sci., p. 120, Marshall Gr.

FIG. 893.—*Pinna squamosa*.

- maxvillensis*, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 221, Kaskaskia Gr.
- missouriensis*, Swallow, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 97, Kaskaskia Gr.
- peracuta*, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 214, and Pal. E. Neb., p. 198, Coal Meas.
- stludovici*, Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 326, St. Louis Gr.
- subspatulata*, Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 524, Warsaw Gr.
- Pinnopsis*, syn. for *Lunulicardium*.
- acutirostra*, syn. *Lunulicardium ornatum*.
- ornatus*, see *Lunulicardium ornatum*.
- Placunopsis*, Morris & Lycett, 1853, Monogr. Foss. Great Oolite. [Ety. *Placuna*, a genus; *opsis*, resemblance.] Suborbicular, upper valve convex, radiately striated or taking the form of the surface to which it adheres; lower valve flat; ligamental groove submarginal; muscular impression subcentral. Type *P. jurensis*. Not a Paleozoic genus.

N. Y., vol. 5, p.

Nat. 10th Ed.
Shell long, tri-
saks terminal,
broad, trun-
gonal, posterior
and a small reni-
cartilage long,
rted by a slen-
n the cardinal
of one internal
n external ver-
Type P. squa-
that sometimes
feet, and ranges
y fathoms. It
tly buried in
lges erect. The
with silk, spun

racuta.
Monogr. U. S.
236, Subcarbon-

St. John, 1868,
i., p. 122, St.

5, Monogr. U. S.
235, Subcarbon-

n Ludlow's Cur-
rk, p. 143, Coal

1865, Proc. Acad.
hall Gr.

uanosa.

1882, Ann. N. Y.
21, Kaskaskia Gr.
1863, Trans. St.
2, p. 97, Kaskas-

Trans. St. Louis
214, and Pal. E.

as.
883, Geo. Sur.
Louis Gr.

1875, Geo. Sur.
raaw Gr.
cardium.

ardium ornatum.
um ornatum,

ett, 1853, Monogr.
[Ety. *Placuna*, a
lance.] Suborb-
convex, radiately
form of the sur-
res; lower valve
ve submarginal:
subcentral. Type
Palaeozoic genus.

Species are left here for want of material to determine their generic relations.

FIG. 894.—*Placunopsis* *PLETHOMYTILUS*, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 4. (Plates and Explanations.) (Ety. *pletho*, to be full; *Mytilus*, a genus.) Mytiloid, gibbous; ligamental area finely striated; no cardinal teeth; lateral teeth small, oblique; test, with concentric striae; differs from *Mytilacra* in its true hinge-line and the absence of teeth. Type P. ponderosus.

carbonaria, Meek & Worthen, 1866, Proc. Chi. Acad. Sci., vol. 1, p. 13, Up. Coal Meas. recticardinalis, Meek, 1875, Ohio Pal., vol. 2, p. 331, Coal Meas.

FIG. 895.—*Plethomytilus ponderosus*.

arenaceus, Hall, 1870, (*Mytilarca arena-*
cea.) Prelim. Notice Lam. Shells, p.
20, and Pal. N. Y., vol. 5, pt. 1, p. 253,
Schoharie grit.
knappi, Hall, 1884, Pal. N. Y., vol. 5, pt.
1, p. 256, Ham. Gr.
mytilimeris, Conrad, 1842, (*Inoceramus*
mytilimeris), Jour. Acad. Nat. Sci.,
vol. 8, p. 246, Low. Held. Gr.

oviformis, Conrad, 1842, (*Inoceramus*
oviformis,) Jour. Acad. Nat. Sci., vol.
8, p. 246, and Pal. N. Y., vol. 5, pt. 1,
p. 255, Ham. Gr.

ponderosus, Hall, 1870, (*Mytilarca pon-*
derosa,) Prelim. Notice Lam. Shells, p.
21, and Pal. N. Y., vol. 5, pt. 1, p. 254,
Up. Held. Gr.

PLEUROPHORUS, King, 1844, Ann. Mag. Nat.
Hist., vol. 14, p. 313. [Ety. *pleuron*, a
rib; *phoros*, bearing.] Inequilateral,
longitudinally oblong or subovate; two
cardinal teeth in each valve, alternately
interlocking and divergent; one pos-
terior lateral tooth in each valve, the
receiving tooth in the left valve; an-
terior adductor scar deep, and bounded
posteriorly by a ridge; pallial line sim-
ple. Type P. costatus.

angulatus, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci. Phil., p. 247, and Geo.
Sur. Ill., vol. 6, p. 529, Coal Meas.

calhouni, Meek & Hayden, 1858, (Ed-
monia calhouni,) Trans. Alb. Inst., vol.
4, p. 80, and Pal. Up. Mo., p. 62, Per-
mian Gr.

chesterensis, Worthen, 1884, Bull. No. 2,
Ill. St. Mus. Nat. Hist., p. 16, and Geo.
Sur. Ill., vol. 8, p. 123, Kaskaskia Gr.

costatiformis, Meek & Worthen, 1865,
Proc. Acad. Nat. Sci. Phil., p. 247, and
Geo. Sur. Ill., vol. 3, p. 535, Keokuk Gr.

meeki, Walcott, 1885, Monogr. U. S. Geo.
Sur., vol. 8, p. 246, Carboniferous.

minus, Worthen, 1884, Bull. No. 2, Ill.
St. Mus. Nat. Hist., p. 17, and Geo. Sur.
Ill., vol. 8, p. 124, St. Louis Gr.

monroensis, Worthen, 1884, Bull. No. 2,
Ill. St. Mus. Nat. Hist., p. 17, and Geo.
Sur. Ill., vol. 8, p. 125, St. Louis Gr.

oblongus, Meek, 1872, Pal. E. Neb., p. 212,
Coal Meas.

occidentalis, Meek & Hayden, 1862, Trans.
Alb. Inst., vol. 4, p. 80, and Pal. Up.
Mo., p. 35, Coal Meas.

pallasi, as identified by Geinitz, is P.
oblongus.

permanus, Swallow,
1858, Trans. St.
Louis Acad. Sci.,
vol. 1, p. 192, Per-
mian Gr.

quadrucostatus, Daw-
son, 1868, Acad.
Geo., p. 304, Car-
boniferous.

simplicis, as identified by Geinitz, is P.
subcuneatus.

subcostatus, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci. Phil., p. 246, and Geo.
Sur. Ill., vol. 2, p. 347, Up. Coal Meas.

subcuneatus, Meek & Hayden, 1858,
Trans. Alb. Inst., vol. 4, p. 81, and Pal.
Up. Mo., p. 61, Permian Gr.

(?) subellipticus, Meek, 1867, Am. Jour.
Sci., vol. 44, p. 181, and Pal. E. Neb.,
p. 211, Coal Meas.

tropidophorus, Meek, 1875, Ohio Pal., vol.
2, p. 338, Coal Meas.

FIG. 896.—*Pleurophorus* *costatiformis*.

Pleurorhynchus, Phillips, syn. for *Conocardium*.

antiquum, see *Conocardium antiquum*.

attenuatum, see *Conocardium attenuatum*.

crassifrons, see *Conocardium crassifrons*.

cuneus, see *Conocardium cuneus*.

trigonale, Hall, see *Conocardium subtrigonale*.

vomer, see *Conocardium vomer*.

Posidonia, Bronn, see *Posidonomya*.

alata, see *Posidonomya alata*.

alveata, see *Grammysia alveata*.

arcuata, see *Grammysia arcuata*.

clathrata, see *Posidonomya clathrata*.

distans, see *Posidonomya distans*.

lirata, see *Paracyclus lirata*.

moorei, see *Posidonomya moorei*.

perstriata, see *Posidonomya perstriata*.

Posidonomya, Bronn, 1837, Leth. Geogn.

[Ety. *Poseidon*, a mythological name; *Mya*, a genus.] Shell thin, obliquely oval, subtruncate at one end, equivalve, compressed, concentrically furrowed, hinge-line short and straight, edentulous. Type *P. becheri*.

alata, Hall, 1843, (*Posidonia* (?) *alata*), Geo. 4th Dist. N. Y., p. 72, and Pal. N. Y., vol. 2, p. 87, Clinton Gr.

FIG. 897.—*Posidonomya becheri*.

ambigua, Winchell, 1863, Proc. Acad. Nat. Sci., p. 10, Marshall Gr.

clathrata, Lea, 1853, (*Posidonia clathrata*), Jour. Acad. Nat. Sci., 2d ser., vol. 2, p. 205, Coal Meas.

devonica, Walcott, 1885, Monogr. U. S. Geo. Sur. vol. 8, p. 179, Devonian.

distans, Lea, 1853, (*Posidonia distans*), Jour. Acad. Nat. Sci., 2d ser., vol. 2, p. 205, Coal Meas.

fracta, Meek, 1875, Ohio Pal., vol. 2, p. 333, Coal Meas.

fragosa, see *Lunulicardium fragosum*.

lævis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 178, Devonian.

mesambonata, Winchell, 1862, Proc. Acad. Nat. Sci., p. 420, Marshall Gr.

moorei, Gabb, 1859, (*Posidonia moorei*), Proc. Acad. Nat. Sci., p. 297, Coal Meas.

perstriata, Lea, 1853, (*Posidonia perstriata*), Jour. Acad. Nat. Sci., 2d ser., vol. 2, p. 205, Coal Meas.

rhomboidea, Hall, 1852, Pal. N. Y., vol. 2, p. 284, Niagara Gr.

romingeri, Winchell, 1862, Proc. Acad. Nat. Sci., p. 420, Marshall Gr.

striata, Stevens, 1858, Am. Jour. Sci., vol. 25, p. 265, Coal Meas.

whiteana, Winchell, 1862, Proc. Acad. Nat. Sci., p. 420, Marshall Gr.

PRÆCARDIUM, Barrande, 1881, Syst. Sil. de la Bohême, vol. 6, p. 141. [Ety. *præ*, before; *Cardium*, a genus.] Equivalve, inequilateral, elliptical or trigonal;

beaks prominent, incurved; surface radiated, and concentrically lined; posterior to the beaks a small area carries a series of vertical nearly parallel teeth.

vetustum, Hall, 1843, (*Cardium vetustum*), Geo. Rep. 4th Dist. N. Y., p. 245, and Pal. N. Y., vol. 5, p. 427, Portage Gr.

FIG. 899.—*Præcardium vetustum*.

PRISCONAIA, Conrad, 1867, Am. Jour. Conch., vol. 3. [Ety. proper name.]

Equivalve, inequilateral, and distinguished from *Anthracostraea*, which it much resembles, by having lateral teeth. Type *P. ventricosa*.

ventricosa, Conrad, 1867, Am. Jour. Conch., vol. 3, Coal Meas.

PROMACRUS, Meek, 1871, Am. Jour. Conch., vol. 7, p. 4. [Ety. *pro*, forward; *makros*, long.]

Similar to *Sanguinolites*; anterior end much produced, narrowly rounded; posterior end produced, obliquely truncate; beaks appressed; cardinal margin nearly straight behind the beaks, and declining in front; umbonal slope angular, extending to the basal extremity; surface concentrically lined, and sometimes plicated anteriorly; ligament external. Type *P. andrewsi*.

andrewsi, Meek, 1871, Am. Jour. Conch., vol. 7, p. 4, Waverly Gr.

cuneatus, Hall, 1870, (*Pholadella cuneata*), Prelim. Not. Lam. Shells, p. 66, and Pal. N. Y., vol. 5, p. 510, Waverly Gr.

missouriensis, see *Sanguinolites missouriensis*.

nasutus, see *Sanguinolites nasutus*.

PRORHYNCHUS, Hall, 1885, Pal. N. Y., vol. 5, p. 48. [Ety. *pro*, forward; *rhynchos*, beak.]

Left valve the larger and more gibbous; anterior end truncate, angular or nasute at the antero-dorsal extremity; posterior end broad, margin truncate or broadly rounded; beaks low; cardinal line straight, extending the entire length of the dorsal margin, and alate at both ends; umbonal slope subangular; surface concentrically lined; strong lateral tooth, ligament external. Type *P. quadratum*.

angulatum, Hall, 1885, Pal. N. Y., vol. 5, p. 493, Chemung Gr.

nasutum, Hall, 1885, Pal. N. Y., vol. 5, p. 493, Chemung Gr.

quadratum, Hall, 1883, (*Palæanatina quadrata*), Pal. N. Y., vol. 5, p. 492, Chemung Gr.

FIG. 900.—*Prothyris meeki*.

PROTHYRIS, Meek, 1869, Proc. Acad. Nat. Sci. Phil., p. 172. [Ety. *pro*, forward; *thyris*, an orifice.]

Equivalve, inequilateral, extremely elongate; cardinal and basal margins subparallel; anterior

erved; surface
rically lined;
as a
les
lled

Car-
rep.
and
127,

FIG. 899.
Præcardium
vetustum.

7, Am. Jour.
proper name.]
al, and distin-
al, which it mnch
eral teeth. Type

17, Am. Jour.
as.

n. Jour. Conch.,
forward; mak-
o *Sanguinolites*;
duced, narrowly
d produced, ob-
laks appressed;
straight behind
ining in front-
r, extending to
surface concen-
times plicated
external. Type

m. Jour. Conch.,
r.

Pholadella cune-
n. Shells, p. 68,
510, Waverly Gr.
olites missouri-

s nasutus.

Pal. N. Y., vol.
rward; *rhynchos*,
larger and more
truncate, angular
antero-dorsal ex-
broad, margin
rounded; beaks
aight, extending
e dorsal margin,
; umbonal slope
concentrically
tooth, ligament
dratum.

al. N. Y., vol. 5,

al. N. Y., vol. 5,

3, (Palæanatina
vol. 5, p. 492,

CHYRIS, Meek,
1869, Proc. Acad.
Nat. Sci. Phil., p.
172. [Ety. *pro*, for-
ward; *thyris*, an
orifice.] Equi-
valve, inequilat-
e; cardinal
parallel; anterior

end rounded or subtruncate, with
a deep notch in the antero-ven-
tral margin; posterior end rounded,
lanceolate, or truncate; cardinal line
straight or slightly arcuate; cardinal
slope sometimes subulate; umbonal
slope rounded, undefined or subangular;
surface concentrically lined. Type *P.*
elegans.

aiata, Hall, 1885, Pal. N. Y., vol. 5, p.
461, Chemung Gr.

elegans, Meek, 1871, Am. Jour. Conch.,
vol. 7, p. 5, Coal Meas.

extata, Hall, 1885, Pal. N. Y., vol. 5, p.
402, Chemung Gr.

lanceolata, Hall, 1883, Pal. N. Y., vol. 5,
pl. 76, figs. 2 to 8, Ham. Gr.

meeki, Winchell, 1875, Ohio Pal., vol. 2,
p. 305, Waverly Gr.

planulata, Hall, 1883, Pal. N. Y., vol. 5,
pl. 76, fig. 1, Ham. Gr.

PROTOMYA, Hall, 1885, Pal. N. Y., vol. 5, p.
52. [Ety. *protos*, first; *Mya*, a genus.]

Equivalve, inequilateral, elongate,
ovate-elliptical; anterior end broadly
rounded; posterior end narrower,
rounded; beaks incurved; umbo
prominent; cardinal line long, nearly
straight; umbonal slope gibbous above,
not defined below; surface concentric-
ally lined; ligament external; mus-
cular impressions circular; anterior one
strong and near the margin. Type *P.*
oblonga.

oblonga, Hall, 1885, (Cardiomorpha ob-
longa), Pal. N. Y., vol. 5, p. 509, Ham. Gr.

PSEUDOMONOTIS, Beyrich, 1862, Zeit. der
Deutsch., Geol. Gesellsch., vol. 14.

[Ety. *pseudes*, false; *Monotis*, a genus.]
Suborbicular, plano-convex, left valve
convex, right valve flat or slightly con-
cave; not auriculate; beaks subcentral,
slightly oblique, unequal, left elevated,
gibbous, incurved, right small; hinge
short, narrow, edentulous; cartilage
cavity under the beaks; byssal notch
of right valve narrow, deep, and
separated from the hinge by a small
rudimentary ear, which does not pro-
ject beyond the margin; adductor mus-
cular scar large, subcentral; impres-

FIG. 901.—*Pseudomonotis*
hawni.

hawni var. *ovata*, Meek & Hayden, 1865,
(*Eumicrotis hawni* var. *ovata*), Pal. Up.
Mo., p. 55, Permian Gr.

hawni var. *sinuata*, Meek & Worthen,
1866, (*Eumicrotis hawni* var. *sinuata*),
Geo. Sur. Ill., vol. 2, p. 338, Up. Coal
Meas.

radialis, (?) Phillips, 1834, (Pecten radi-
alis), Encyc. Meth., vol. 4, Coal Meas.

PTERINEA, Goldfuss, 1826, Germ. Petref.

[Ety. *pteron*, a wing.] Transversely
triangular, oblique, inequivalve, very in-
equilateral, left valve most convex,
beaks near the small anterior end;
hinge-line long, straight, forming a
small anterior and large falciform pos-
terior wing, with a linear, flattened,
marginal cartilage facet, longitudinally
striated; shell thick, calcareous; two
long, slightly diverging, posterior, lat-
eral teeth, beneath the hinge in one
valve and one in the other; a few
short, cardinal teeth radiating beneath
and in front of the beaks; anterior im-
pression very strong just in front of the
beak, posterior impression larger, but
faintly marked, superficial; pallial scar
simple; shallow byssal concavity. Type
P. levis.

FIG. 902.—*Pterinea demissa*.

appressa, Conrad, 1838, Ann. Rep. N. Y.
Not defined.

arenacea, Hall, 1877. Proposed, but not
defined.

aviformis, Conrad, 1842, (Avicula avi-
formis), Jour. Acad. Nat. Sci., vol. 8, p.
243, Trenton Gr.

avis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1,
p. 105, Chemung Gr.

bellilineata, Billings, 1866, Catal. Sil. Foss.
Antic., p. 15, Hud. Riv. Gr.

bisulcata, see *Grammysia bisulcata*.

brisa, Hall, 1867, 20th Rep. N. Y. Mus.
Nat. Hist., p. 384, syn. for *P. stricocosta*.

cancellata, Barris, 1879, (Avicula cancel-
lata), Proc. Dav. Acad. Sci., vol. 2, p.
286, Corniferous limestone.

cardiiformis, see *Megambonia cardiiformis*.
cardinata, Winchell, 1862, Proc. Acad. Nat. Sci., p. 412, Marshall Gr.
carinata, Goldfuss, see *Ambonychia carinata*.
chemungensis, Conrad, 1842, (Avicula chemungensis,) Jour. Acad. Nat. Sci., vol. 8, p. 243, and Pal. N. Y., vol. 5, pt. 1, p. 98, Chemung Gr.
concentrica, Conrad, 1838, Ann. Rep. N. Y. Not defined.
consimilis, Hall, 1883, Pal. N. Y., vol. 5, pt. 1, p. 100, Chemung Gr.
corrugata, James, 1874, (Avicula corrugata,) Cln. Quar. Jour. Sci., vol. 1, p. 239, Hud. Riv. Gr.
crenistriata, Winchell, 1862, (Cardiopsis crenistriata,) Proc. Acad. Nat. Sci., p. 417, Marshall Gr.
crenulata, see *Crenipecten crenulatus*.
cuneata, see *Sphenotus cuneatus*.
curiosa, Billings, 1866, Catal. Sil. Foss. Antic., p. 51, Anticosti Gr.
cyrtodontoides, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 95, Niagara Gr.
demissa, Conrad, 1842, (Avicula demissa,) Jour. Acad. Nat. Sci., vol. 8, p. 242, and Pal. N. Y., vol. 1, p. 202, Hud. Riv. Gr.
dispanda, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 97, Chemung Gr.
elliptica, Hall, 1847, (Avicula elliptica,) Pal. N. Y., vol. 1, p. 162, Trenton Gr.

FIG. 903.—*Pterinea flabellum*.

flabellum, Conrad, 1842, (Avicula flabella,) Jour. Acad. Nat. Sci., vol. 8, p. 238, and Pal. N. Y., vol. 5, pt. 1, p. 93, Up. Held. and Ham. Grs.
grandis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 91, Up. Held. Gr.

honeymani, Hall, 1860, (Avicula honeymani,) Can. Nat. and Geol., vol. 5, p. 153, and Acad. Geol., p. 604, Up. Sil.
insueta, Emmons, 1842, (Avicula insueta,) Geo. Rep. 2d Dist. N. Y., p. 390, and Pal. N. Y., vol. 1, p. 291, Utica Slate and Hud. Riv. Grs.
interstriata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 96, Chemung Gr.
modiolaris, see *Modiolopsis modiolaris*.
morganensis, see *Avicula morganensis*.
mucronata, Ulrich, 1870, Jour. Cln. Soc. Nat. Hist., vol. 2, p. 24, Hud. Riv. Gr.
neglecta, McChesney, 1861, New Palaeozoic Fossils, p. 88, Niagara Gr.
newarkensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 165, Devonian.
orbicularis, see *Ambonychia orbicularis*.
pholadis, see *Orthonota pholadis*.
pinguis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 92, Up. Held. Gr.
pintoensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 234, Subcarboniferous.
planulata, see *Cypriocardinia planulata*.
proflifica, Billings, 1866, Catal. Sil. Foss. Antic., p. 10, Hud. Riv. Gr.
prora, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 102, Chemung Gr.
punctulata, Conrad. Not defined. See *Cimitaria recurva*.
pygmæa, see *Modiella pygmæa*.
radicans, see *Panienka radicans*.
reproba, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 106, Chemung Gr.
reversa, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 104, Chemung Gr.
reversa var. avis, see *P. avis*.
revoluta, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 95, Niagara Gr.
rigida, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 101, Chemung Gr.
rugosa, Conrad, 1841, (Avicula rugosa,) Ann. Geo. Rep. N. Y., and Geo. Rep. 4th Dist. N. Y., pt. 26, fig. 2, Waterlime Gr.
smallis, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 214, Marcellus Shale.
spinalata, Winchell, 1865, Proc. Acad. Nat. Sci., p. 124, Burlington (?) Gr.
striacosta, McChesney, 1861, (Ambonychia striacosta,) New Pal. Foss., p. 88, Niagara Gr.
strigosa, White & Whitfield, 1862, Proc. Bost. Soc. Nat. Hist., p. 31, Marshall Gr.
suborbicularis, see *Pterinopecten suborbicularis*.
subpapyracea, Meek & Worthen, 1866, Proc. Chi. Acad. Sci., p. 21, Ham. Gr.
thebesensis, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 354, Niagara Gr.
thisbe, Billings, 1866, Catal. Sil. Foss. Antic., p. 52, Anticosti Gr.
trentonensis, Conrad, 1842, (Avicula trentonensis,) Jour. Acad. Nat. Sci., vol. 8, p. 240, and Pal. N. Y., vol. 1, p. 161, Trenton Gr.
triquetra, see *Gosseletia triquetra*.
undata, see *Ambonychia undata*.

Avicula honey-
Geol., vol. 5, p.
604, Up. Sil.
Avicula insueta,
Y., p. 399, and
291, Utica Slate

al. N. Y., vol. 5,
r.
s modiolaris.
morganensis,
Jour. Cin. Soc.
Hud. Riv. Gr.
61, New Palaeo-
ara Gr.

35, Monogr. U. S.
5, Devonian.
ia orbicularis.
pholadis.

N. Y., vol. 5, pt.
s, Monogr. U. S.
234, Subcarbonif-

ia planulata.
Ontal. Sil. Foss.
r. Gr.

Y., vol. 5, pt. 1,
ot defined. See

gmæa.
ians.

N. Y., vol. 5, pt.
r.
N. Y., vol. 5, pt.

r.
avis.

larcy, 1865, Mem.
p. 95, Niagara Gr.
Y., vol. 5, pt. 1,

(*Avicula rugosa*),
p. and Geo. Rep.
26, fig. 2, Water-

Ann. N. Y. Acad.
rcellus Shale.
805, Proc. Acad.
ington (?) Gr.
1861, (Ambo-
ew Pal. Foss., p.

field, 1862, Proc.
p. 31, Marshall Gr.
inopecten subor-

Worthen, 1868,
p. 21, Ham. Gr.
orthen, 1868, Geo.
64, Niagara Gr.
Catal. Sil. Foss.
ti Gr.

342, (*Avicula tren-*
Nat. Sci., vol. 8,
Y., vol. 1, p. 161,

a triquetra.
a undata.

undulata, Meek & Worthen, 1868, Geo.
Sur. Ill., vol. 3, p. 456, Kinderhook Gr.
varistriata, Billings, 1866, Catal. Sil. Foss.
Antic., p. 50, Anticosti Gr.
volans, Winchell & M. rey, 1865, Mem.
Bost. Soc. Nat. Hist., p. 95, Niagara Gr.
PTERINOPECTEN, Hall, 1883, Pal. N. Y., vol.
5, pt. 1, p. 3. (Plates and Explana-
tions.) [Ety. *Pterinea*, a genus; *Pec-*
ten, a genus.] Valves more or less
convex; radiated and bearing concen-
tric lines of growth; hinge-line long,
straight; wings not well defined, being
simple expansions of the upper lateral
margins to the hinge-line. Type *P.*
undosus.

FIG. 904.—*Pterinopecten undosus*.

conspectus, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 68, Ham. Gr.
crenicostatus, Hall, 1884, Pal. N. Y., vol.
5, pt. 1, p. 78, Chemung Gr.
dignatus, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 62, Marcellus Shale.
dispanus, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 76, Chemung Gr.
erectus, Hall, 1884, Pal. N. Y., vol. 5, pt.
1, p. 77, Chemung Gr.
exfoliatus, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 61, Marcellus Shale.
flitextus, Hall, 1884, Pal. N. Y., vol.
5, pt. 1, p. 67, Ham. Gr.
hermes, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 64, Ham. Gr.
hoosacensis, Walcott, 1885, Monogr.
U. S. Geo. Sur., vol. 8, p. 232,
Subcarboniferous.
imbecilis, Hall, 1884, Pal. N. Y., vol.
5, pt. 1, p. 75, Chemung Gr.
insons, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 59, Up. Held. Gr.
intermedius, Hall, 1884, Pal. N. Y.,
vol. 5, pt. 1, p. 68, Ham. Gr.
invalidus, see *Aviculopecten in-*
validus.
lætus, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 63, Marcellus Shale.
multiradiatus, Hall, 1884, Pal. N. Y.,
vol. 5, pt. 1, p. 57, Up. Held. Gr.
neptunus, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 79, Chemung Gr.
nodosus, Hall, 1884, Pal. N. Y., vol. 5, pt.
1, p. 60, Up. Held. Gr.
reflexus, Hall, 1884, Pal. N. Y., vol. 5, pt.
1, p. 58, Up. Held. Gr.
regularis, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 70, Ham. Gr.

spio, Walcott, 1885, Monogr. U. S. Geo.
Sur., vol. 8, p. 233, Subcarboniferous.
spondylus, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 65, Ham. Gr.

FIG. 905.—*Pterinopecten undosus*.

strictus, Hall, 1884, Pal. N. Y., vol. 5, pt.
1, p. 74, Chemung Gr.
suborbicularis, Hall, 1884, (*Pterinea sub-*
orbicularis.) Geo. Rep. 4th Dist. N. Y.,
p. 204, and Pal. N. Y., vol. 5, p. 80,
Chemung Gr.

terminalis, see *Aviculopecten terminalis*.
undosus, Hall, 1884, Pal. N. Y., vol. 5, pt.
1, p. 72, Ham. Gr.

vertumnus, Hall, 1884, Pal. N. Y., vol. 5,
pt. 1, p. 71, Ham. Gr.

PTERONITELLA, Billings, 1874, Pal. Foss., vol.
2, p. 141. [Ety. diminutive of *Ptero-*
nites.] Resembles *Pterinea*, but pos-
sesses in front of the beaks several
small, anterior, cardinal teeth, and close
beneath the hinge-line several more or
less elongated posterior teeth. Type *P.*
venusta.

curta, Billings, 1874, Pal. Foss., vol. 2, p.
143, Low. Held. Gr.

FIG. 906.—*Pteronitella venusta*. b. Hinge-line.

oblonga, Billings, 1874, Pal. Foss., vol. 2,
p. 143, Low. Held. Gr.
venusta, Billings, 1874, Pal. Foss., vol. 2,
p. 142, Low. Held. Gr.

PTERONITES, McCoy, 1844, Syn. Carb. Foss.
Ireland, p. 81. [Ety. *pteron*, a wing.]
Subtriangular, depressed, hinge-line as
long as the shell; beaks terminal, or
nearly so, forming a very narrow, ob-

tusely pointed anterior end, from which the ventral margin extends to the broad posterior end; left valve most convex; internally a very small tooth under the beak of the right valve, and a very slender, posterior, lateral tooth close to the hinge-line the whole length. Type *P. angustatus*.

gayensis, Dawson, 1868, Acad. Geo., p. 301, Subcarboniferous.

gayensis var. *ornatus*, Dawson, 1883, Rep. on Redpath Mus., No. 2, p. 14, Subcarboniferous.

inoptatus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 239, Chemung Gr.

newarkensis, Meek, 1871, Proc. Acad. Nat. Sci., p. 162, Waverly Gr.

FIG. 907.—*Pteronites profundus*.

profundus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 237, Up. Chemung Gr.

rostratus, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 238, Chemung Gr.

aspergenensis, Whitfield, 1882, Bull. Am. Mus. Nat. Hist., No. 3, p. 56, Warsaw Gr.

FIG. 908.—*Ptychodesma knappanum*. Right side.

Form modioloid; hinge having a wide ligamental area, grooved by the successive growth of the ligament, as in *pectunculus*. Type *P. knappanum*.

FIG. 909.—*Ptychodesma knappanum*. Enlargement of one side of ligamental area.

knappanum, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 192, Up. Held. Gr.

mincr, Hall, 1885, Pal. N. Y., vol. 5, p. 353, Chemung Gr.

nanum, Hall, 1885, Pal. N. Y., vol. 5, p. 353, Chemung Gr.

FIG. 910.—*Ptychopteria beecheri*.

Ptychopteria, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 3. [*Ety. ptyche*, fold; *Pteria*, a genus.] Differs from *Actinopteria* in the nasute anterior extremity, and large, straight wing marked by a strong longitudinal fold. Hinge-line narrow, linear; furnished with one or two linear, oblique, cardinal and lateral teeth; surface with fine rays. It bears about the same relation to *Actinopteria* that *Leptodesma* does to *Liopteria*. Type *P. eugenia*.

alata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 139, Chemung Gr.

beecheri, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 143, Chemung Gr.

FIG. 911.—*Ptychopteria beecheri*.

elongata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 141, Chemung Gr.

eucrate, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, 133, Chemung Gr.

eudora, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 138, Chemung Gr.

eugenia, Hall, 1883, Pal. N. Y., vol. 5, pl. 23, figs. 17-20, Chemung Gr.

expansa, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 152, Chemung Gr.

falcata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 136, Up. Chemung Gr.

galene, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 142, Chemung Gr.

gibbosa, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 149, Up. Chemung Gr.

lata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 145, Up. Chemung Gr.

lobata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 150, Up. Chemung Gr.

perlata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 147, Up. Chemung Gr.

protic, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 129, Chemung Gr.

protoformis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 235, Subcarboniferous.

salamanca, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 131, Chemung Gr.

. Y., vol. 5, p.

becheri.

Pal. N. Y., vol. 5, p. 132, Chemung Gr. sinuosa, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 130, Up. Chemung Gr. spatulata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 144, Up. Chemung Gr. spio, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 137, Chemung Gr. thalia, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 148, Up. Chemung Gr. thietis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 135, Chemung Gr. trigonalis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 140, Chemung Gr. vanuxemi, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 151, Up. Chemung Gr. PYANOMYA, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318. [Ety. *pyanos*, a bean; *Mya*, a genus.] Equivalve, elongate, inequilateral, fragile, edentulous; ligament external. Type *P. gibbosa*.

Ptychopteria becheri.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

N. Y., vol. 5, pt.

sao, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 132, Chemung Gr.

sinuosa, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 130, Up. Chemung Gr.

spatulata, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 144, Up. Chemung Gr.

spio, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 137, Chemung Gr.

thalia, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 148, Up. Chemung Gr.

thietis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 135, Chemung Gr.

trigonalis, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 140, Chemung Gr.

vanuxemi, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. 151, Up. Chemung Gr.

PYANOMYA, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318. [Ety. *pyanos*, a bean; *Mya*, a genus.] Equivalve, elongate, inequilateral, fragile, edentulous; ligament external. Type *P. gibbosa*.

P. gibbosa.

faberi, n. sp. Shell small, equivalve, inequilateral; length twice as great as height; cardinal and basal lines subparallel; anterior end sharply rounded into the subelliptical base; posterior end broadly rounded; valves ventricose in the middle; beaks obtuse; umbonal ridge prominent, subangular, distinctly defined, and directed to the postero-basal margin; ligament external; hinge-line straight behind the beaks and inclined in front; no escutcheon or lunule. Surface marked very faintly by concentric lines of growth. Distinguished from *P. gibbosa* by the angular umbonal ridge, less acute anterior end and straight cardinal line behind the beaks, and other particulars. Hud. Riv. Gr., Cincinnati, Ohio. Collected by Charles Faber.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

SANGUINOLARIA, Lamarek, 1801, Syst. An. sans Vert. [Ety. from the type *Solen sanguinolentus*.] Oval, compressed, rounded in front, attenuated and slightly gaping behind; hinge teeth $\frac{2}{3}$ small; siphonal inflection deep, connected with the pallial line; ligament external, on very prominent fulcrum. Type *S. sanguinolentus*. Typical species *S. diphos*. Not American Paleozoic. Species left under this name for want of material to determine generic relations.

FIG. 914.—*Sanguinolites diphos*.

leptogaster, Winchell, 1863, Proc. Acad. Nat. Sci., p. 18, Marshall Gr.

rostrata, Winchell, 1865, Proc. Acad. Nat. Sci., p. 129, Marshall Gr.

sectoralis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 422, Marshall Gr.

septentrionalis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 421, Marshall Gr.

similis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 421, Marshall Gr.

SANGUINOLITES, McCoy, 1844, Synop. Carb. Foss., Ireland, p. 47. [Ety. *Sanguinolites*, a genus; *lithos*, stone.] Subequivalve, oblong, elongated, margins subparallel or a little arched upward; sides compressed or diagonally gibbous from the beak backward; beaks small, anterior; hinge nearly as long as the shell, margin inflected to form a long posterior lunette; surface wrinkled; large, oval adductor impression in front of the beak surmounted by a small retractor; posterior adductor large, superficial; cartilage external; pallial impression entire; shell thin.

obliquus, see *Goniophora acuta*.

aeolus, see *Sphenotus aeolus*.

amygdalinus, see *Glossites amygdalinus*.

arciformis, see *Sphenotus arciformis*.

borealis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 415, Marshall Gr.

brookfieldensis, Dawson, 1883, Rep. on Redpath Museum, p. 11, Subcarboniferous.

burlingtonensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 14, and Geo. Sur. Ill., vol. 8, p. 129, Burlington Gr.

obliquus, see *Goniophora acuta*.

aeolus, see *Sphenotus aeolus*.

amygdalinus, see *Glossites amygdalinus*.

arciformis, see *Sphenotus arciformis*.

borealis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 415, Marshall Gr.

brookfieldensis, Dawson, 1883, Rep. on Redpath Museum, p. 11, Subcarboniferous.

burlingtonensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 14, and Geo. Sur. Ill., vol. 8, p. 129, Burlington Gr.

obliquus, see *Goniophora acuta*.

aeolus, see *Sphenotus aeolus*.

amygdalinus, see *Glossites amygdalinus*.

arciformis, see *Sphenotus arciformis*.

borealis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 415, Marshall Gr.

brookfieldensis, Dawson, 1883, Rep. on Redpath Museum, p. 11, Subcarboniferous.

burlingtonensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 14, and Geo. Sur. Ill., vol. 8, p. 129, Burlington Gr.

FIG. 912.—*Pyanomya faberi*. Right valve and dorsal view.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

gibbosa, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 318, Hud. Riv. Gr.

FIG. 913.—*Pyanomya gibbosa*. Left valve and dorsal view.

PYRENOMCEUS, Hall, 1852, Pal. N. Y., vol. 2, p. 87. [Ety. *pyrenos*, Nucula; *omoios*, similar; from its resemblance in general form to the shells of the genus *Nucula*.] Equivalve, inequilateral; umbones prominent, beak elevated; muscular impression near the anterior extremity; general form of *Nucula* without the teeth that characterize that genus, or the clavicle of a *Clidophorus*. Type *P. cuneatus*.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

cuneatus, Hall, 1852, Pal. N. Y., vol. 2, p. 87, Clinton Gr.

clavulus, see *Sphenotus clavulus*.
combensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 175, Devonian.
concentricus, Winchell, 1862, (*Cardinia concentrica*), Proc. Acad. Nat. Sci., p. 413, Marshall Gr.
cylindricus, Winchell, 1863, Proc. Acad. Nat. Sci., p. 13, Marshall Gr.
flavius, see *Sphenotus flavius*.
glaucaus, see *Goniophora glaucaus*.
gracilis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 175, Devonian.
hamiltonensis, see *Goniophora hamiltonensis*.
ida, Hall, 1870, Prelim. Notice Lam. Shells, p. 43, and Pal. N. Y., vol. 5, pl. 65, fig. 20, Ham. Gr.
iowensis, Winchell, 1863, Proc. Acad. Nat. Sci., p. 14, Marshall Gr.
jejunus, Winchell, 1863, Proc. Acad. Nat. Sci., p. 15, Marshall Gr.
marshallensis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 415, Marshall Gr.
missouriensis, Swallow, 1860, (Solen (?) *missouriensis*), Trans. St. Louis Acad. Sci., vol. 1, p. 655, Waverly or Choctau Gr.
multistriatus, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 14, and Geo. Sur. Ill., vol. 8, p. 129, Keokuk Gr.
nenia, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 249, Subcarboniferous.
naiadiformis, Winchell, 1870, Proc. Am. Phil. Soc., vol. 12, p. 255, Marshall Gr.
nasutus, Meek, 1871, Am. Jour. Conch., vol. 7, Kinderhook Gr.
obliquus, Meek, 1871, Proc. Acad. Nat. Sci., p. 213, and Ohio Pal. vol. 2, p. 306, Waverly Gr.
perangulatus, see *Goniophora perangulata*.
ponderosus, see *Modiomorpha ponderosa*.
randolphensis, Worthen, 1883, (*Cypri-cardia randolphensis*), Geo. Sur. Ill., vol. 7, p. 326, Kaskaskia Gr.
retusus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 247, Subcarboniferous.
salteri, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 248, Subcarboniferous.
sanduskiensis, Meek, 1871, Proc. Acad. Nat. Sci., p. 68, and Ohio Pal., vol. 1, p. 209, Up. Held. Gr.
securis, Winchell, 1870, Proc. Am. Phil. Soc., vol. 12, p. 255, Marshall Gr.
simplex, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 248, Subcarboniferous.
solenoides, see *Sphenotus solenoides*.
striatus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 249, Subcarboniferous.
strigatus, Winchell, 1865, Proc. Acad. Nat. Sci., p. 127, Marshall Gr.
subtortuosus, see *Sphenotus subtortuosus*.
subtruncatus, Hall, 1885, Pal. N. Y., vol. 5, p. 508, Chemung Gr.
sulciferus, see *Cypriocardia sulcifera*.
tethys, Billings, 1874, Pal. Foss., vol. 2, p. 50, Gaspe limestone No. 8, Devonian.
undatus, Hall, 1870, Prelim. Notice Lam. Shells, p. 41, and Pal. N. Y., vol. 5, pl. 80, figs. 5, 6, Chemung Gr.

unioniformis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 414, Marshall Gr.
valvulus, see *Sphenotus valvulus*.
Schizodus, King, 1844, Ann. Mag. Nat. Hist., vol. 14, p. 313. [Ety. *schizo*, I split; *odus*, a tooth.] Shell oval or subtriangular; anterior side rounded, shorter than the other; posterior side tapering, truncate at the extremity, umbonal ridge extending to the postero-basal region; beaks prominent; surface smooth or with concentric striae; hinge with two smooth cardinal teeth in the right valve and three in the left; middle tooth of the left valve bifid, and fitting between two of the right valve; free margin smooth. Type *S. truncatus*.
aqualia, Hall, 1885, Pal. N. Y., vol. 5, p. 459, Waverly Gr.
amplus, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 41, and Geo. Sur. Ill., vol. 5, p. 579, Coal Meas.
cayuga, Hall, 1870, Prelim. Notice Lam. Shells, p. 95, syn. for *Cytherodon appressus*.

chesterensis, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 457, and Geo. Sur. Ill., vol. 2, p. 301, Kaskaskia Gr.
circulus, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 11, and Geo. Sur. Ill., vol. 8, p. 109, St. Louis Gr.
contractus, Hall, 1885, Pal. N. Y., vol. 5, p. 451, Ham. Gr.
cuneatus, Meek, 1875, Ohio Pal., vol. 2, p. 336, Coal Meas.
curtiformis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 253, Subcarboniferous.
curtus, Meek & Worthen, 1866, Proc. Chi. Acad. Sci., p. 18, Coal Meas.
degener, Hall, 1885, Pal. N. Y., vol. 5, p. 456, Chemung Gr.
deparcus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 252, Subcarboniferous.
depressus, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 11, and Geo. Sur. Ill., vol. 8, p. 109, St. Louis Gr.
ellipticus, see *Cytherodon ellipticus*.
eminens, Hall, 1885, Pal. N. Y., vol. 5, p. 457, Chemung Gr.

FIG. 916.—*Schizodus medmensis*.

FIG. 917.—Hinge of *Schizodus truncatus*.

gregarius, see *Cytherodon gregarius*.
magnus, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 9, and Geo. Sur. Ill., vol. 8, p. 107, Kaskaskia Gr.
medinensis, Meek, 1871, Proc. Acad. Nat. Sci. Phil., vol. 23, p. 165, and Ohio Pal., vol. 2, p. 299, Waverly Gr.

1862, Proc. Acad. Nat. Sci. Phil., p. 42, and Geo. Sur. Ill., vol. 5, p. 581, Coal Meas.

S. truncatus, N. Y., vol. 5, p. 41, and Geo. Sur. Ill., vol. 5, p. 581, Coal Meas. Notice Lam. Cytherodon ap-

then, 1870, Proc. Acad. Nat. Sci. Phil., p. 41, and Geo. Sur. Ill., vol. 5, p. 581, Coal Meas. Notice Lam. Cytherodon ap-

FIG. 916.—*Schizodus medius*.

o. Sur. Ill., vol. 8,

Pal. N. Y., vol. 5,

Ohio Pal., vol. 2, p.

1885, Monogr. U. S. 253, Subcarbonif-

en, 1866, Proc. Chi. Meas.

al. N. Y. vol. 5, p.

1885, Monogr. U. S. 252, Subcarbonif-

1884, Bull. No. 2, Ill. p. 11, and Geo. Sur. Louis Gr.

on ellipticus.

al. N. Y., vol. 5, p.

us, see *Cytherodon*

arius.

is, Worthen, 1884, No. 2, Ill. St.

s. Nat. Hist., p. 9, Geo. Sur. Ill., vol.

107, Kaskaskia Gr. ensis, Meek, 1871, Sci. Phil., vol. 23, p.

vol. 2, p. 299, Wa-

moorei, n. sp. Shell very large, sub-rhomboidal, height and length sub-equal; anterior side straight from the beaks and at right angles to the posterior side, and then rounded into the basal line; basal margin regularly rounded; posterior side sloping at right angles to the anterior side from the beaks and abruptly rounding into the basal margin; beaks prominent, rising above the cardinal line, obtuse, and situate a little anterior to the middle of the shell; umbonal slope broadly rounded and undefined; pallial line strongly marked, pitted, and placed near the margin from one muscular scar to the other; anterior and posterior muscular scars subtrigonal and moderately impressed; a wide vascular impression, somewhat cordate, occupies the central area of the shell, extending from the anterior to the posterior muscular scars; one strong tooth in the right valve directed a little forward, with a socket on each side, the other tooth undefined; surface nearly smooth, showing fine concentric lines of growth. Found by Henry Moores, of Columbus, Ohio, at Carbon Hill, Hocking Valley, in the Coal Measures, and now in the collection of Charles Faber.

FIG. 918.—*Schizodus moorei*. Right valve, posterior part broken off.

nauvoensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 10, and Geo. Sur. Ill., vol. 8, p. 108, Keokuk Gr.

orbicularis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 181, Devonian.

ovatus, Meek & Hayden, 1853, (Axinus ovatus,) Proc. Acad. Nat. Sci. Phil., p. 262, and Pal. Up. Mo., p. 59, Permian Gr.

patulus, Hall, 1885, Pal. N. Y., vol. 5, p. 457, Chemung Gr.

perelegans, Meek & Worthen, 1870, Proc. Acad. Nat. Sci. Phil., p. 42, and Geo. Sur. Ill., vol. 5, p. 581, Coal Meas.

FIG. 919.—*Schizodus moorei*. Interior of right valve, showing pallial line and place of subcordate muscular impression and hinge-teeth.

pintoensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 253, Subcarboniferous.

quadrangularis, see *Cytherodon quadrangularis*.

randolphensis, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 110, Kaskaskia Gr.

rossicus, Verneuil, 1845, Geo. Russ., vol. 2, p. 309, Permian Gr.

subtrigonalis, Meek, 1871, Proc. Acad. Nat. Sci., p. 166, Waverly Gr.

triangularis, Swallow, 1858, Trans. St. Louis Acad. Sci., p. 193, Permian Gr.

tumidus, see *Cytherodon tumidus*.

ulrichi, Worthen, (in press,) Geo. Sur. Ill., vol. 8, p. 110, Up. Coal Meas.

varsoviensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 10, and Geo. Sur. Ill., vol. 8, p. 107, Keokuk Gr.

wheeleri, Swallow, 1862, (Littorina wheeleri,) Trans. St. Louis Acad. Sci., vol. 1, p. 658, and Pal. E. Neb., p. 209, Coal Meas.

SEDGWICKIA, McCoy, 1844, Snyop. Carb. Foss. Ireland, p. 61. [Ety. proper name.]

Nearly equivalve, inequilateral, depressed, oblong, or suboval, very thin; anterior side not quite closed, often gibbous; posterior side longer, more compressed, and gaping; beaks prominent, tumid, incurved; posterior umbonal slopes rounded, or forming an oblique ridge, separated from the postero-dorsal region by a shallow sulcus; lunule distinct; hinge edentulous; cardinal margin inflected so as to form a narrow false area behind the beaks; surface concentrically marked. Type *S. attenuata*.

altirostrata, Meek & Hayden, 1858, (Alorisma (?) altirostratum,) Proc. Acad. Nat. Sci. Phil., p. 263, and Pal. Up. Mo., p. 41, Coal Meas.

(?) compressa, Meek, 1872, Proc. Acad. Nat. Sci. Phil., p. 324, and Ohio Pal., vol. 1, p. 144, Hud. Riv. Gr.

concava, Meek & Hayden, 1858, (Lyonsia concava,) Trans. Alb. Inst., vol. 4, p. 82, and Pal. Up. Mo., p. 41, Coal Meas.

(?) divaricata, Hall & Whitfield, 1875, Ohio Pal., vol. 2, p. 89, Hud. Riv. Gr.

(?) fragilis, Meek, 1872, Proc. Acad. Nat. Sci. Phil., p. 323, and Ohio Pal., vol. 1, p. 143, Hud. Riv. Gr.

(?) lunulata, Whitfield, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 140, Hud. Riv. Gr. neglecta, see Cuneamya neglecta.

subarcuata, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 251, and Geo. Sur. Ill., vol. 3, p. 537, Keokuk Gr.

FIG. 920.—*Sedgwickia topekensis*.

topekensis, Shumard, 1858, (Leptodomus topekensis,) Trans. St. Louis Acad. Sci., vol. 1, p. 208, and Pal. Up. Mo., p. 40, Coal Meas.

Solemya, Lamarck, 1818, Hist. Nat. An. sans Vert., vol. 5. See *Solenomya*—the correct orthography, first used by Menke, 1828, Syn. Meth. Edit.

SOLENYA, Linnaeus, 1758, Syst. Nat., 10th ed. [Ety. *Solen*, a tube or pipe.] Shell very long; subcylindrical; ends gaping; hinge teeth two in each valve; ligament external; anterior scar elongated; posterior oblong; pallial line extending beyond the adductors. Type *S. siliqua*. Not a Palaeozoic genus.

FIG. 921.—*Solen siliqua*. One-third diam.

missouriensis, see *Sanguinolites missouriensis*.

permianus, see *Solenopsis permianus*. priscus, Winchell, 1862, Proc. Acad. Nat. Sci., p. 423, Portage Gr.

quadrangularis, Winchell, 1862, Proc. Acad. Nat. Sci., p. 422, Marshall Gr.

scalpriformis, see *Solenopsis scalpriformis*. *SOLENOMYA*, Lamarck, 1818, (*Solemya*.) Hist. Nat. Anim. sans Vert., vol. 5, p. 488. [Ety. from the resemblance to the two genera *Solen* and *Mya*.] Elongate, ob-

long, equivalve, very inequilateral, posterior end the shorter; dorsal and ventral margins subparallel; ends rounded and gaping; surface covered with a thick, horny periostraca, extending in jagged portions beyond the ventral margin; beaks minute; cartilage forming a thick, triangular mass behind the beaks, supported internally by an oblique ensiform plate; long anterior margin simple, erect, without teeth; posterior adductor small, ovate, within the cartilage pit, anterior impression large, comma-shaped. Type *S. australis*.

anodontoides, Meek, 1875, Ohio Pal., vol. 2, p. 339, Coal Meas.

biarmica, Verneuil, 1845, Geo. Russ. and Ural Mountains, Permian Gr. This species does not occur in this country. curta, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 242, Subcarboniferous.

iowensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 13, and Geo. Sur. Ill., vol. 8, p. 132, St. Louis Gr.

monroensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 13, and Geo. Sur. Ill., vol. 8, p. 131, St. Louis Gr.

radiata, Meek & Worthen, 1860, (*Solemya radiata*.) Proc. Acad. Nat. Sci. Phil., p. 457, and Geo. Sur. Ill., vol. 2, p. 349, Coal Meas.

recurvata, Swallow, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 208, Up. Coal Meas.

soleniformis, Cox, 1857, Geo. Sur. Ky., vol. 3, p. 573, Coal Meas.

varsoviensis, Worthen, 1884, Bull. No. 2, Ill. St. Mus. Nat. Hist., p. 12, and Geo. Sur. Ill., vol. 8, p. 131, Keokuk Gr.

FIG. 922.—*Solenomya vetusta*.

vetusta, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 66, and Ohio Pal., vol. 1, d. 206, Up. Held. Gr.

SOLENOPHIS, McCoy, 1844, Carb. Foss. Ireland, p. 47. [Ety. *Solenopsis*, resembling a shell of the genus *Solen*.] Elongated; beaks compressed, anterior; depression in front of beaks; surface concentrically marked. Type *S. minor*.

permianus, Swallow, 1858, (*Solen permianus*.) Trans. St. Louis Acad. Sci., vol. 1, p. 190, Permian Gr.

scalpriformis, Winchell, 1862, (*Solen scalpriformis*.) Proc. Acad. Nat. Sci., p. 422, Marshall Gr.

solenoides, Gelnitz, 1866, (*Clidophorus solenoides*.) Carb. und Dyas in Neb., p. 25, and Pal. E. Neb., p. 223, Coal Meas.

FIG. 923.—*Solenopsis solenoides*.

[SOL.

inequilateral, dorsal and ventral ends parallel; ends of surface covered by periostraca, extensions beyond the minute; cartilaginous mass distorted internally; plate; long anterior end; without or small, ovate, bit, anterior im-

Ohio Pal., vol.

Geo. Russ. and ...nian Gr. This in this country. ...nogr. U. S. Geo. ...carboniferous. ...Bull. No. 2, Ill. ...13, and Geo. Sur. ...Louis Gr. ...1884, Bull. No. 2, ...p. 13, and Geo. ...St. Louis Gr. ...n, 1860, (Solemya ...Nat. Sci. Phil., p. ...vol. 2, p. 349,

, Trans. St. Louis ...p. 208, Up. Coal

, Geo. Sur. Ky, ...as.

1884, Bull. No. 2, ...ist., p. 12, and ...8, p. 131, Keokuk Gr. ...vetusta, Meek, ...1871, Proc. Acad. Nat. Sci. Phil., p. 66, and Ohio Pal., vol. 1, d. 206, Up. Held. Gr.

Carb. Foss. Ire-

a. 923.—*Solenopsis solenoides*.

; depression in ...ce concentrically ...or.

1858, (Solen per- ...Louis Acad. Sci., ...n Gr.

ll, 1862, (Solen ...Acad. Nat. Sci., p.

66, (Clidophorus ...nd Dyas in Neb., ...Neb., p. 223, Coal

SPATHIELLA, Hall, 1885, Pal. N. Y., vol. 5, p. 33. [Ety. *spathe*, a spathe; *ellus*, diminutive.] Equivalve, very inequilateral, wider behind, transversely sub-cylindrical; anterior end short, narrowly rounded; beaks subanterior, small; umbonal slope rounded or sub-angular; surface concentrically lined. Type S. typica.

typica, Hall, 1885, Pal. N. Y., vol. 5, p. 407, Chemung Gr.

ventricosa, White & Whitfield, 1862, (Orthonota ventricosa.) Proc. Bost. Soc. Nat. Hist., vol. 8, p. 297, and Pal. N. Y., vol. 5, p. 408, Waverly Gr.

SPHENOLIUM, n. gen. [Ety. *sphen*, wedge; *leion*, smooth.] Shell large, equivalve, inequilateral, elongate, cuneiform, ventricose; umbones prominent; beaks incurved at the anterior end; cardinal line at an angle of fifty or sixty degrees from the basal line, and appearing wing-like toward the posterior end; lunule present; no escutcheon; ligament external; muscular scars and hinge-line unknown. Type S. cuneiforme.

cuneiforme, S. A. Miller, 1881, (Orthonota cuneiforme.) Jour. Nat. Soc. Nat. Hist., vol. 3, p. 314, Hud. Riv. Gr. faberi, n. sp. Shell below the medium size for species in this genus; beaks unite over the hinge-line near the anterior end; anterior end, pointed, rounded; hinge-line rising poste-

FIG. 924.—*Sphenolium faberi*. Left valve.

riorly into a wing-like expansion; posterior end prolonged at the postero-basal margin; basal margin subelliptical; umbones high and gradually tapering to the postero-basal margin; surface concentrically lined. Collected by Charles Faber in the Hud. Riv. Gr., at Cincinnati, Ohio.

FIG. 925.—*Sphenolium richmondense*.

richmondense, n. sp. Shell large, cuneiform, ventricose, beaks incurved at the anterior end, pointed; umbones high, defined; cardinal line at a high angle,

having a wing-like posterior end; anterior end rounded below the lunule. Distinguished from S. cuneiforme, which it much resembles, by its shorter form and more angular umbones. Possibly more specimens may show a gradation from one form to the other, and if so, this specific name will fall into synonymy. Collected by Charles Faber in the upper part of the Hud. Riv. Gr., at Richmond, Indiana.

FIG. 926.—*Sphenolium richmondense*. Anterior view.

SPHENOTUS, Hall, 1885, Pal. N. Y., vol. 5, p. 33. [Ety. *sphen*, wedge; *ous*, ear.] Equivalve, very inequilateral, elongate; anterior end short; posterior end obliquely truncate; cardinal line, long, straight; umbonal ridge extending to the post-inferior extremity; surface concentrically lined; two short teeth beneath the beak of the right valve, and one or two slender lateral teeth; ligament external, contained in a groove; anterior muscular scar strongly marked; posterior scar shallow; pallial line simple. Type S. arciformis.

FIG. 927.—*Sphenotus æolus*.

æolus, Hall, 1870, (Sanguinolites æolus.) Prelim. Not. Lam. Shells, p. 46, and Pal. N. Y., vol. 5, p. 404, Waverly Gr. arciformis, Hall, 1870, (Sanguinolites arciformis.) Prelim. Not. Lam. Shells, p. 40, and Pal. N. Y., vol. 5, p. 399, Ham. Gr. arcuatus, Hall, 1885, Pal. N. Y., vol. 5, p. 400, Chemung Gr.

clavulus, Hall, 1870, (Sanguinolites clavulus.) Prelim. Not. Lam. Shells, and Pal. N. Y., vol. 5, p. 401, Chemung Gr. contractus, Hall, 1843, (Cypricardia contracta.) Geo. Sur. 4th Dist. N. Y., p. 292, and Pal. N. Y., vol. 5, p. 399, Chemung Gr.

cuneatus, Conrad, 1838, (Pterinea cuneata.) Ann. Rep. Geo. N. Y., p. 116, and Pal. N. Y., vol. 5, p. 396, Ham. Gr.

flavius, Hall, 1870, (Sanguinolites flavius.) Prelim. Not. Lam. Shells, p. 47, and Pal. N. Y., vol. 5, p. 403, Waverly Gr. rigidus, White & Whitfield, 1862, (Cypricardia rigida.) Proc. Bost. Soc. Nat. Hist., vol. 8, p. 300, and Pal. N. Y., vol. 5, p. 402, Waverly Gr.

signatus, Hall, 1885, Pal. N. Y., vol. 5, p. 405, Waverly Gr.

solenoides, Hall, 1870, (Sanguinolites solenoides.) Prelim. Not. Lam. Shells, p. 38, and Pal. N. Y., vol. 5, p. 398, Ham. Gr.

subtortuosus, Hall, 1870, (Sanguinolites subtortuosus,) Prelim. Not. Lam. Shells, p. 41, and Pal. N. Y., vol. 5, p. 397, Ham. Gr.
 telamon, Hall, 1885, Pal. N. Y., vol. 5, p. 406, Waverly Gr.
 truncatus, Conrad, 1842, (Cypricardites truncatus,) Jour. Acad. Nat. Sci., vol. 8, p. 244, Ham. Gr.
 undatus, Hall, 1885, Pal. N. Y., vol. 5, p. 506, Chemung Gr.
 valvulus, Hall, 1870, (Sanguinolites valvulus,) Prelim. Not. Lam. Shells, p. 46, and Pal. N. Y., vol. 5, p. 403, Waverly Gr.

FIG. 928.—*Spirodomus insignis*.

SPIRODOMUS, Beecher, 1886, 39th Rep. N. Y. Mus. Nat. Hist. [Ety. *speira*, spire; *demos*, house.] Equivalve, elongate-spiral; beaks terminal; muscular impressions at the two extremities; no hinge-line. Type *S. insignis*.

insignis, Beecher, 1886, 30th Rep. N. Y. Mus. Nat. Hist., Waverly Gr.

STREBLOPTERIA, McCoy, 1851, Ann. Mag. Nat. Hist., 2d series, vol. 7, p. 170, and Brit. Pal. Rocks, p. 482. [Ety. *streblos*, turned the wrong way; *pteron*, a wing.] Pectinoid, ovate, or rounded, obliquely extended toward the anterior side; posterior wing rectangular, anterior ear small, deeply defined; surface smooth or radiately ridged; large, faintly marked muscular impression behind the middle; short, narrow tooth posterior to the beaks; ligament confined to a narrow, simple facet on the hinge margin. Type *S. laevigata*.

FIG. 929.—*Streblopteria similis*.
Right valve.

similis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 230, Carboniferous.

tenuilineata, Meek & Worthen, 1860, (Pecten *tenuilineatus*,) Proc. Acad. Nat. Sci., Phil., p. 452, and Geo. Sur. Ill., vol. 2, p. 334, Coal Meas.

TECHNOPHORUS, n. gen. [Ety. *technē*, art; *phoros*, bearing.] Shell small, equivalent, inequilateral; anterior end short, broadly rounded; two or more furrows arising near the beak extend to the postero-basal margin; beak small, upright; surface concentrically lined; umbonal rib in front of the beak represented in the cast by a transverse sulcus; no external ligament, escutcheon, or lunule. Type *T. faberi*.

faberi, n. sp. Shell small, equivalent, inequilateral, a little longer than high; anterior end short, broadly rounded; base more narrowly rounded in the anterior and central part; the postero-basal part slightly produced; cardinal line straight or nearly so; beak extremely small and standing upright, like a little point projecting beyond the cardinal line; valves convex in the umbonal region; two furrows or cinctures arising near the beak in the umbonal region, which gradually widen, are directed to the postero-basal margin, and above these the postero-dorsal part of the shell

is somewhat wing-like; surface marked by very fine concentric lines; the casts show a deep sulcus directly in front of the beak for the reception of an umbonal rib, or support on the interior of the shell. Hud. Riv. Gr., near

Sharonville, Hamilton County, Ohio. Collected by Mr. Charles Faber.

FIG. 930.—*Technophorus faberi*. The right hand figure shows the left valve with a small piece broken from the posterior end; the left hand figure represents a well-preserved cast; the central figure presents a cardinal view.

Tellina, Linnæus, 1758, Syst. Nat., 10th ed. [Ety. *telline*, a sort of mussel.] This genus unknown in the Palæozoic rocks.

(?) *ovata*, Hall, 1843, Geo. Rep. 4th Dist. N. Y. Syn. for *Palæoneilo maxima*.

TELLINOMYA, Hall, 1847, Pal. N. Y., vol. 1, p. 151. [Ety. from the resemblance to the genera *Tellina* and *Mya*.] Nearly equilateral, generally transverse, anterior side largest; beaks approximate, not prominent; hinge-line with a double series of bent teeth connected by smaller ones beneath the beak; ligament posterior, external, on a fulcrum; no striated area or cartilage pit; muscular impressions strong not bounded by elevated lines; pallial line simple. Type *T. nasuta*.

abrupta, Billings, 1862, (Ctenodonta *abrupta*,) Pal. Foss., vol. 1, p. 46, Black Riv. Gr.

æquilatera, Hall, 1852, Pal. N. Y., vol. 2, p. 330, Coralline limestone.

alta, Hall, 1861, Geo. Rep. Wis., p. 27, and Geo. Sur. Ill., vol. 3, p. 309, Trenton Gr.

anatiniformis, see *Pterotheca anatiniformis*.

angela, Billings, 1865, (Ctenodonta *angela*,) Pal. Foss., vol. 1, p. 221, Quebec Gr.

, equivalve, in-
than high; an-
rounded; base
in the anterior
postero-basal
cardinal line
beak extremely
right, like a little
and the cardinal
the umbonal re-
cinctures arising
umbonal region,
are directed to
gin, and above
part of the shell
what wing-like;
marked by very
concentric lines;
s show a deep
directly in front
beak for the re-
of an umbonal
support on the
r of the shell.
Riv. Gr., near
a County, Ohio.
les Faber.

ri. The right hand
with a small piece
end; the left hand
preserved; const; the
cardinal view.

yst. Nat., 10th ed.
mussel.] This ge-
Paleozoic rocks.
eo. Rep. 4th Dist.
oneilo maxima.
al. N. Y., vol. 1, p.
resemblance to the
ya.] Nearly equi-
ansverse, anterior
approximate, not
ne with a double
connected by smaller
lak; ligament pos-
a fulcrum; no
lage pit; muscular
not bounded by
lial line simple.

2, (Ctenodonta ab-
ol. 1, p. 46, Black

Pal. N. Y., vol. 2,
estone.

Rep. Wis., p. 27,
ol. 3, p. 309, Tren-

erotherca anatin-

, (Ctenodonta an-
ol. 1, p. 221, Que-

angustata, Hall, 1860, Can. Nat. and Geo.,
vol. 5, p. 152, Up. Sil.
astartiformis, Salter, 1859, (Ctenodonta
astartiformis,) Can. Org. Rem., Decade
1, p. 39, Black Riv. Gr.
attenuata, Hall, 1860, Can. Nat. and Geo.,
vol. 5, p. 151, Up. Silurian.
cingulata, Ulrich, 1879, Jour. Cin. Soc.
Nat. Hist., vol. 2, p. 23, Hud. Riv. Gr.
contracta, Salter, 1859, (Ctenodonta con-
tracta,) Can. Org. Rem. Decade 1, p. 37,
Black Riv. and Trenton Gr.
curta, Hall, 1852, Pal. N. Y., vol. 2, p. 86,
Clinton Gr.
donaciformis, Hall, 1847, (Nucula ? dona-
ciformis,) Pal. N. Y., vol. 1, p. 316, Tren-
ton Gr.
dubia, Hall, 1847, Pal. N. Y., vol. 1, p.
153, Black Riv. and Trenton Grs.
elliptica, Hall, 1852, Pal. N. Y., vol. 2, p.
102, Clinton Gr.
gibberula, Salter, 1859, (Ctenodonta gib-
berula,) Can. Org. Rem. Decade 1, p.
38, Black Riv. and Trenton Grs.
gibbosa, Hall, 1847, Pal. N. Y., vol. 1, p.
153, Black Riv. and Trenton Grs.
hamburgensis, Walcott, 1885, Monogr.
U. S. Geo. Sur., vol. 8, p. 76, Tren-
ton Gr.
hartsvillensis, Safford, 1859, (Ctenodonta
hartsvillensis,) Geo. of Tenn., p. 287,
Nashville Gr.

FIG. 931.—Tellino-
mya infla.

hilli, S. A. Miller, 1874,
Cin. Quar. Jour. Sci.,
p. 230, Hud. Riv. Gr.
houghtoni, Stevens, 1858,
(Nucula houghtoni),
Am. Jour. Sci., and
Arts, 2d ser., vol. 25,
p. 262, Marshall or Waverly Gr.
inflata, Hall, 1861, Geo. Rep. Wis., p. 26,
Trenton Gr.
iphigenia, Billings, 1862, (Ctenodonta
iphigenia,) Pal. Foss., vol. 1, p. 152,
Hud. Riv. Gr.
lata, Hall, 1852, Pal. N. Y., vol. 2, p. 85,
Clinton Gr.
levata, Hall, 1847, (Nucula levata,) Pal.
N. Y., vol. 1, p. 150, Black Riv., Tren-
ton, and Hud. Riv. Grs.
logani, Salter, 1851, (Ctenodonta logani),
Rep. Brit. Assoc., p. 36, Hud. Riv. Gr.
machæiformis, Hall, 1843, (Nucula machæ-
ariformis,) Geo. Rep. 4th Dist. N. Y., p. 76,
and Pal. N. Y., vol. 2, p. 85, Clinton Gr.
mactrififormis, Hall, 1843, (Nucula mac-
trififormis,) Geo. Rep. 4th Dist., N. Y., p.
76, Clinton Gr.

nasuta, Hall,
1847, Pal.
N. Y., vol. 1,
p. 152, Black
Riv. and Tren-
ton Grs.

nucleiformis,
Hall, 1859,
Pal. N. Y.,
vol. 3, p. 263, Low. Held. Gr.
nuculiformis, Hall, 1847, (Modiolopsis

FIG. 932.—Tellinomya
nasuta.

nuculiformis,) Pal. N. Y., vol. 1, p. 298,
Hud. Riv. Gr.
ovata, Hall, 1861, Geo. Rep. Wis., p. 28,
Trenton Gr.
pectunculoides, Hall, 1871, 24th Rep.
N. Y. Mus. Nat. Hist., p. 228, Hud.
Riv. Gr.
protensa, Hall, 1852, Stans. Ex. to Gt.
Salt Lake, p. 412, Coal Meas.
sanguinolarioidea, Hall, 1847, Pal. N. Y.,
vol. 1, p. 152, Trenton Gr.
stella, Winchell, 1862, (Nucula stella),
Proc. Acad. Nat. Sci., p. 417, Mar-
shall Gr.
subnasuta, see Clinopistha subnasuta.
ventricosa, Hall, 1861, Geo. Rep. Wis., p.
27, and Geo. Sur. Ill., vol. 3, p. 307,
Trenton Gr.

TELLINOPSIS, Hall, 1870, Prelim. Notice
Lam. Shells, p. 80. [Ety. resembling a
shell of the genus Tellina.] General
form like Tellina; beaks small, subcen-
tral, directed backward; ligament ex-
ternal; surface smooth or obscurely
marked; ligament external; muscular
impression shallow. Type T. sub-
emarginata.

submarginata,
Conrad, 1842,
(Nuculites
submarginata,
Journ.
Acad. Nat. Sci.,
vol. 8, p. 249,
and Pal. N. Y.,
vol. 5, p. 464,
Ham. Gr.

Ungulina, Daudin, 1802, Bosc. Hist. Nat.
Coq. 3. [Ety. ungulina, like a hoof.]
suborbicularis, see Cardiomorpha suborbic-
ularis.

Unio orthonotus, see Modiolopsis orthonota.
primigenius, see Modiolopsis primigenia.

VANUXEMIA, Billings, 1858, Rep. of Progr.
Geo. Sur. Can., p. 186. [Ety. proper
name.] Ovate; beaks terminal or sub-
terminal; posterior extremity rounded,
anterior more or less acuminate; two
muscular impressions; anterior teeth
variable in number sometimes curved
and striated; posterior lateral teeth
from two to four. Type V. inconstans.

FIG. 934.—Vanuxemia bayfieldi. Interior of
left valve, showing the striated teeth.

bayfieldi, Billings, 1858, Rep. of Progr.
Geo. Sur. Can., p. 187, Hud. Riv. Gr.
dixonensis, Meek & Worthen, 1866, Proc.
Chi. Acad. Sci., p. 16, Trenton Gr.

- inconstans*, Billings, 1858, Rep. of Progr. Geo. Sur. Can., p. 186, Black Riv. and Trenton Grs.
- montrealensis*, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 444, Chazy Gr.
- tomkinsi*, Billings, 1860, Can. Jour., vol. 6, p. 357, Up. Held. Gr.
- Venus mohegan*, Castelnau, 1843, Syst. Sil., p. 44. Not recognized.
- VERTUMNIA**, Hall, 1884, Pal. N. Y., vol. 5, pt. 1, p. xii. Proposed as a subgenus of *Pterinea*, distinguished by having the right valve convex, and the left flat or concave; hinge area narrow. The species referred to it are *Pterinea avis*, *P. reprobata*, and *P. reversa*.
- YOLDIA**, Muller, 1842, Kroyer's Nat. Tid., vol. 4, p. 91. [Ety. proper name.] Shell ovate or subelliptical, subequilateral, compressed; posterior side narrower than the other; surface smooth, striate, or obliquely sculptured, and covered with a polished epidermis; margins smooth within; inner laminae pearly; hinge plates small, numerous on each side of the beaks; cartilage pit under the beaks; pallial line sinuous. Type *Y. myalis*.
- carbonaria*, Meek, 1871, Rep. Reg. University W. Va., p. 6, and Ohio Pal., vol. 2, p. 336, Coal Meas.
- gibbosa*, McChesney, 1859. The name was preoccupied. See *T. rushensis*.
- knoxensis*, McChesney, 1865, (*Leda knoxensis*,) Expl. Pal. Foss., pl. 2, Coal Meas.
- levistriata*, Meek & Worthen, 1860, (*Leda levistriata*,) Proc. Acad. Nat. Sci. Phil., p. 457, and Geo. Sur. Ill., vol. 2, p. 282, St. Louis Gr.
- oweni*, McChesney, 1860, (*Leda oweni*,) Desc. New. Pal. Foss., p. 52, Coal Meas.
- polita*, McChesney, 1859. The name was preoccupied, see *Y. knoxensis*.
- rushensis*, McChesney, 1865, (*Leda rushensis*,) Expl. Pal. Foss., pl. 2, Coal Meas.
- stevensoni*, Meek, 1871, Rep. Reg. University W. Va., p. 6, and Ohio Pal., vol. 2, p. 335, Coal Meas.
- subscitula*, Meek & Hayden, 1858, (*Leda subscitula*, Trans. Alb. Inst., vol. 4, p. 79, and Pal. E. Neb., p. 205, Permian Gr.
- valvulus*, Hall & Whitfield, 1872, 24th Rep. N. Y. Mus. Nat. Hist., p. 190, Up. Held. Gr.

FIG. 635.—*Yoldia myalis*.

SUBKINGDOM ARTICULATA.

THE Articulata are the most numerous of all living animals, and abound alike on land and sea. They are divided into Classes, Subclasses, Orders, and Suborders. Many of them possess intelligence, arising from ganglionic centers, and in the summer season provide their food for winter. Several living orders are unknown in Palæozoic rocks; this may have resulted, however, from want of preservation. The fossils belong to the Classes Annelida, Crustacea, Arachnida, Myriapoda, Insecta.

CLASS ANNELIDA.

The Annelida have the bodies divided into segments, which are generally furnished with jointed appendages. The living forms are distributed in four Orders, but no such division is practicable with the Palæozoic fossils, where generally only the internal jaws, called Conodonts or worm-burrows, are found preserved. The Conodonts may be the internal jaws of Crustacea, as seems to the author most probable; but there is no ground for referring them to fish, as has been done by some authors. The class may be divided as follows:

CONODONTS.—*Arabellites*, *Distacodus*, *Drepanodus*, *Eunicites*, *Glycerites*, *Lumbriconereites*, *Nereidavus*, *Oenonites*, *Polygnathus*, *Prioniodus*, *Staurcephalites*.

The name was
shensis.

G. 035.—Yoldia
myalia.

eo. Sur. Ill., vol.

, (Leda oweni),
s., p. 52, Coal

The name was
oxensis.

865, (Leda rush-
ss., pl. 2, Coal

Rep. Reg. Uni-
d Ohio Pal., vol.

den, 1858, (Leda
Inst., vol. 4, p.

b., p. 205, Per-

field, 1872, 24th
Hist., p. 190, Up.

TA.

and abound alike

s, and Suborders.

and in the sum-

are unknown in

of preservation.

ida, Myriapoda,

are generally fur-

d in four Orders,

re generally only

preserved. The

the author most

has been done by

Glycerites, Lum-

ioniodus, Stauro-

WORM-BURROWS.—Arenicolites, Gyrichnites, Myrianites, Monocraterion, Ne-
mapodia, Nereites, Palaeochorda, Scolithus, Walcottia.

ORDER TUBICOLA.—Conchicolites, Cornulites, Salterella, Serpula, Serpulites,
Spirorbis.

ORDER UNCERTAIN.—Protoscolex.

ARABELLITES, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 377. [Ety. *Ara-
bella*, an existing genus; *lithos*, stone.]
Jaws with an extremely prominent an-
terior hook, and a row of smaller teeth
on a wide base, sickle-shaped jaws, and
also subquadrate forms, with a straight
upper edge of small teeth. Type *A.*
hamatus.

ascialis, Hinde, 1879, Quar. Jour. Geo. Soc.
Lond., vol. 35, p. 378, Hud. Riv. Gr.

cervicornis, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 379, Hud.
Riv. Gr.

cornutus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 377, Hud. Riv. Gr.
crenulatus, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 379, Hud.
Riv. Gr.

cristatus, see *Eunicites cristatus*.
cuspidatus, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 378, Hud.
Riv. Gr.

elegans, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 382, Clinton Gr.
gibbosus, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 378, Hud.
Riv. Gr.

hamatus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 377, Hud.
Riv. Gr.

lunatus, Hinde, 1879, Quar. Jour. Geo. Soc.
Lond., vol. 35, p. 378, Hud. Riv. Gr.

obliquus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 379, Hud. Riv. Gr.

ovalis, Hinde, 1879, Quar. Jour. Geo. Soc.
Lond., vol. 35, p. 378, Hud. Riv. Gr.

pectinatus, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 379, Hud.
Riv. Gr.

politus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 385, Ham. Gr.

quadratus, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 379, Hud.
Riv. Gr.

rectus, Hinde, 1879, Quar. Jour. Geo. Soc.
Lond., vol. 35, p. 378, Hud. Riv. Gr.

scutellatus, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 379, Hud.
Riv. Gr.

similis, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 382, Niagara Gr.

similis var. *arcuatus*, Hinde, 1879, Quar.
Jour. Geo. Soc. Lond., vol. 35, p. 385,
Ham. Gr.

ARENICOLITES, Salter, 1856, Quar. Jour. Geo.

Soc., vol. 13, p. 199. [Ety. *arena*, sand;
colo, I inhabit; *lithos*, stone.] Circular
holes which appear in twos on the sur-
face of sandstones, and have the ap-

pearance of worm-burrows, like those
of the *Arenicola*. Type *A. sparsus* or
A. didyma.

sparsus, Salter, 1856, Quar. Jour. Geo.
Soc., vol. 13, p. 203, Clinton Gr.

spiralis, Torell, 1868, as identified by Bill-
ings, Pal. Foss., vol. 2, p. 77, Up. Taconic.
woodi, Whitfield, 1882, Geo. Wis., vol. 4,
p. 177, Potsdam Gr.

Aulacodus obliquus, see *Lumbriconereites*
obliquus.

CONCHICOLITES, Nicholson, 1872, Am. Jour.
Sci. and Arts, 3d ser., vol. 3, p. 202.
[Ety. *concha*, shell; *colo*, I dwell; *lithos*,
a stone.] Tubes conical, slightly curved,
walls thin, composed of imbricating
rings. Type *C. gregarius*. Prof. Hall
and others regard this genus as a syn-
onym for *Cornulites*.

corrugatus, Nicholson, 1873, Lond. Geo.
Mag., vol. 10, p. 55, Hud. Riv. Gr.

flexuosus, Hall, 1847, (Tentaculites *flexuo-*
sus.) Pal. N. Y., vol. 1, p. 92, Trenton
and Hud. Riv. Grs.

FIG. 936.—*Conchicolites flexuosus*, on *Stropho-*
mena alternata.

intermedius, Nicholson, 1874, (Ortonia
intermedia.) Geo. Mag., n. s., vol. 1, p.
199, Ham. Gr.

minor, Nicholson, 1873, (Ortonia *minor*.)
Lond. Geo. Mag., vol. 10, p. 56, Hud.
Riv. Gr.

CORNULITES, Schlotheim, 1820, Petrefakten-
kunde, p. 378. [Ety. *cornu*, horn; *lithos*,
stone.] Tube gradually tapering, con-
ical, slightly flexuous, small end usu-
ally curved, and attached to some for-
eign body; walls thick, cellular,
composed of numerous imbricating
rings, their widest edge next the slender
base; external surface annulated,
finely striated longitudinally; inner
surface and casts scalariform, with two
or three longitudinally impressed fur-
rows. Type *C. serpularius*.

arcuatus, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 276, Niagara Gr.

bellistriatus, Hall, 1888, Pal. N. Y., vol. 7, p. 20, Low. Held. Gr.

carbonarius, Gurley, 1883, New. Carb. Foss., p. 8, Kinderhook Gr. The publication is not such as required by the rules of nomenclature.

chrysalis, Hall, 1888, Pal. N. Y., vol. 7, p. 20, Low. Held. Gr.

cingulatus, Hall, 1888, Pal. N. Y., vol. 7, p. 20, Low. Held. Gr.

clintoni, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 184, Clinton Gr. This name was proposed instead of *C. flexuosus*, which is preoccupied, when *Conchicolites* is regarded as synonymous with *Cornulites*.

contractus, Ringueberg, 1884, Proc. Acad. Nat. Sci., p. 148, Niagara Gr. Syn. for *C. proprius*.

distans, Hall, 1852, (Tentaculites *distans*.) Pal. N. Y., vol. 2, p. 184, Clinton Gr.

flexuosus, Hall, 1852, Pal. N. Y., vol. 2, p. 98, Clinton Gr.

flexuosus var. *gracilis*, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 155, Niagara Gr.

nodosus, Ringueberg, 1884, Proc. Acad. Nat. Sci., p. 149, Niagara Gr.

proprius, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 182, Niagara Gr.

tribulus, Hall, 1888, Pal. N. Y., vol. 7, p. 20, Ham. Gr.

DIPTACODUS, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 357. [Ety. *diptaxo*, to doubt; *odous* tooth.] Small, curved tooth, with a sharp edge on both the outer and inner curve; base expanded. Type *D. incurvus*.

incurvus, Pander, 1856, (Machairodus *incurvus*.) Monogr. d. foss. Fische. d. Silur. syst., p. 23, Hud. Riv. Gr.

DREPANODUS, Pander, 1856, Monogr. d. foss. Fische. d. Silur. Syst., p. 20. [Ety. *drepans*, sickle; *odous* tooth.] Small, curved, spine-like tooth, nearly circular in section; base expanded. Type *D. arcuatus*.

arcuatus, Pander, 1856, Monogr. d. Foss. Fische. d. Silur. Syst., p. 20, Hud. Riv. Gr.

Eotrophonia, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 91. Not satisfactorily defined.

setigera, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 91. Not satisfactorily defined, and specimen too poor for definition.

EUNICITES, Ehlers, 1868, Palaeontographica, vol. 17, p. 145. [Ety. *Eunice*, a Nereid; *lithos*, stone.] Minute, variously formed, denticulated jaws of annelids or crustaceans. Type *E. avitus*.

FIG. 937.—*Cornulites arcuatus*.

FIG. 938.—*Cornulites distans*.

alveolatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 384, Ham. Gr.

chiomorphus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 381, Clinton Gr.

clintonensis, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 381, Clinton Gr.

compactus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 384, Ham. Gr.

contortus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 375, Hud. Riv. Gr.

coronatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 381, Clinton Gr.

cristatus, Hinde, 1879, (Arabellites *cristatus*), Quar. Jour. Geo. Soc., vol. 35, p. 378, Hud. Riv. Gr.

digitatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 378, Hud. Riv. Gr.

gracilis, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 376, Hud. Riv. Gr.

major, see *Oenonites major*.

nanus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 384, Ham. Gr.

palmaris, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 384, Ham. Gr.

perdentatus, see *Lumbriconereites perdentatus*.

simplex, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 376, Hud. Riv. Gr.

timidus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 384, Ham. Gr.

GLYCERITES, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 380. [Ety. genus *Glyceria*; *lithos*, stone.] Jaws consisting of a simple curved hook with a wide base, without smaller teeth. Type *G. sulcatus*.

calceolus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 384, Clinton Gr.

sulcatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 360, Hud. Riv. Gr.

sulcatus, var. *excavatus*, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 380, Hud. Riv. Gr.

Gordia marina, see *Palaeochorda marina*.

GYRICHNITES, Whiteaves, 1883, Trans. Roy. Soc. Can., p. 109. [Ety. *gyros*, a circle; *ichnos* a track.] Trails supposed to have been made by an annelid. Type *G. gaspensis*.

gaspensis, Whiteaves, 1883, Trans. Roy. Soc. Can., p. 109, Mid. Devonian.

Helminthoidichnites, Fitch, see *Palaeochorda marina*.

tenuis, see *Palaeochorda tenuis*.

LUMBRICONEREITES, Ehlers, 1868, Palaeontographica, vol. 17, p. 159. [Ety. *Lumbriconereis*, a genus; *lithos*, stone.] Distinguished from *Eunice* by having a well-defined basal extension. Type *L. deperditus*.

Quar. Jour. Geo.
384, Ham. Gr.
179, Quar. Jour.
35, p. 381, Clin-

79, Quar. Jour.
35, p. 381, Clin-

Quar. Jour. Geo.
384, Ham. Gr.
Quar. Jour. Geo.
p. 375, Hud.

Quar. Jour. Geo.
p. 381, Clin-

Arabellites cris-
to. Soc., vol. 35,

Quar. Jour. Geo.
p. 376, Hud.

Quar. Jour. Geo.
p. 376, Hud.

or.
Quar. Jour. Geo.
384, Ham. Gr.

Quar. Jour. Geo.
384, Ham. Gr.

onereites perden-

Quar. Jour. Geo.
35, p. 376, Hud.

Quar. Jour. Geo.
384, Ham. Gr.

Quar. Jour. Geo.
380. [Ety. genus

Jaws consisting
hook with a wide
teeth. Type G.

Quar. Jour. Geo.
384, Clinton Gr.

Quar. Jour. Geo.
p. 380, Hud.

as, Hinde, 1879,
Lond., vol. 35, p.

horda marina.

1883, Trans. Roy.
ty. gyro, a circle;
supposed to have
annelid. Type G.

1883, Trans. Roy.
Devonian.

see Palaeochorda.

tenuis.

1868, Palaeonto-
59. [Ety. Lambri-
e, stone.] Distin-
ites by having a
tension. Type L.

armatus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 383, Clin-
ton Gr.

basalis, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 383, Clin-
ton Gr.

dactyloides, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 380, Hud.
Riv. Gr.

obliquus, Eichwald, 1854, (Sphagodus obli-
quus), Bull. d. l. Soc. Imp. d. Nat. d.
Moscou, p. 110, Hud. Riv. and Clin-
ton Gr.

perdentatus, Hinde, 1879, (Euniceites per-
dentatus), Quar. Jour. Geo. Soc., vol.
35, p. 375, Clinton Gr.

triangularis, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 383, Clin-
ton Gr.

Machairodus, Pander, 1856. This name was
preoccupied. See *Distacodus*.

incurvus, see *Distacodus incurvus*.

MONOCRATERION, Torell, 1860, *Acta univer-*
sitatis lundensis, p. 13. [Ety. *monos*,
one; *kraterion*, small basin.] Borings in
the rock resembling *Scolithus*, except
in having a tunnel-shaped enlargement
at the upper end.

lesleyi, Prime, 1878, Geo. Sur. Pa. DD. p.
79, *Calceiferous* (?) Gr.

MYRIANITES, Murchison, 1839, Sil. Syst., p.
700. [Ety. *Myrias*, a myriad; *lithos*, stone.]
Trails lying together in great numbers,
more or less corrugated upon the edges,
and resembling delicate wave lines upon
the surface of the rock. Type M. ma-
cleali.

murchisoni, Emmons, 1844, *Taconic syst.*,
p. 44, Up. Taconic.

sillimani, Emmons, 1844, *Taconic Syst.*,
p. 44, Up. Taconic.

NEMAPODIA, Emmons, 1844, *Taconic Syst.*, p.
68. [Ety. *nema*, a thread; *pous*, a foot.]
Trail consisting of a series of depres-
sions marked by numerous short paral-
lel fine lines; the trail is flexuous, and
the short, fine lines have the direction
of the trail. Type N. tenuissima.

tenuissima, Emmons, 1844, *Taconic syst.*,
p. 68, Up. Taconic.

NEREIDAVUS, Grinnell, 1877, Am. Jour. Sci.
and Arts, 3d ser., vol. 14, p. 229. [Ety.
Nereis, genus; *avis*, grandfather.] Mi-
nute denticulated teeth or jaws. Type
N. varians.

solitarius, Hinde,
1879, Quar. Jour.
Geo. Soc. Lond.,
vol. 35, p. 385,
Ham. Gr.

variens, Grinnell,
1877, Am. Jour.

Sci. and Arts, 3d ser., vol. 14, p. 229,
Hud. Riv. Gr.

NEREITES, Murchison, 1839, Sil. Syst., p. 700.
[Ety. from a resemblance to the track
of the *Nereis*.] Long, convoluted trails;
each side equally crenulated; crenula-
tions oval or pointed on the margin,

and often traceable to the center of
the trail. Type N. cambrensis.

deweyi, Em-
mons, 1844,
Taconic
Syst., p. 69,
Up. Ta-
conic.

gracilis, Em-
mons, 1844,
Taconic
Syst., p. 69,
Up. Ta-
conic.

jacksoni,
Emmons,
1844, Taconic
Syst., p. 69,
Up. Taconic.

lancoolatus,
Emmons, 1844, *Taconic Syst.*, p. 69, Up.
Taconic. This may belong to *Nereog-*
grapsus, as suggested by Emmons.

loomisi, Emmons, 1844, *Taconic Syst.*, p.
69, Up. Taconic.

pugnis, Emmons, 1844, *Taconic Syst.*, p.
69, Up. Taconic.

robustus, Emmons, 1856, (*Nereograpsus ro-*
bustus), Am. Geol., p. 111, Up. Taconic.

OENONITES, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 376. [Ety. *Oenone*,
a genus; *lithos*, stone.] Jaws with a
more or less curved anterior hook; fol-
lowed by a series of smaller teeth, simi-
lar in character to those of the existing
genus *Oenone*. Type O. curvidens.

amplus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 382, Clinton Gr.

carinatus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 377, Hud. Riv. Gr.

cuneatus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 377, Hud. Riv. Gr.

curvidens, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 376, Hud. Riv. Gr.

fragilis, Hinde, 1879, Quar. Jour. Geo. Soc.
Lond., vol. 35, p. 382, Clinton Gr.

inæqualis, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 376, Hud. Riv. Gr.

infrequens, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 382, Niagara Gr.

major, Hinde, 1879, (*Euniceites major*),
Quar. Jour. Geo. Soc., vol. 35, p. 374,
Hud. Riv. and Clinton Gr.

rostratus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 376, Hud. Riv. Gr.

serratus, Hinde, 1879, Quar. Jour. Geo.
Soc. Lond., vol. 35, p. 376, Hud. Riv. Gr.

ORTONIA, Nicholson, 1872, Lond. Geo. Mag.,
vol. 9. Synonym for *Conchicolites*, if
indeed both are not synonyms for
Cornulites.

conica, syn. for *Conchicolites flexuosus*.

intermedia, see *Conchicolites intermedius*.

minor, see *Conchicolites minor*.

PALAECHORDA, McCoy, 1848, Quar. Jour.
Geol. Soc., vol. 4, p. 224. [Ety. *palaios*,
ancient; *chorda*, intestine.] Trail very
long, cylindrical, chord-like, frequently

FIG. 940.—*Nereites deweyi*.

FIG. 939.—*Nereidavus varians*. Magnified 8 diam.

- crossing itself, without order, surface smooth. Type *P. minor*.
marina, Emmons, 1844, (*Gordia marina*), p. 68, and Am. Geol., p. 103, Up. Taconic.
prima, Whitfield, 1877, Prelim. Rept. Pal. Black Hills, p. 7, and Geol. Black Hills of Dakota, p. 331, Potsdam Gr.
tenuis, Fitch, 1849, (*Helminthoidichnites tenuis*), Trans. Agr. Soc., and Am. Geol., p. 103, Up. Taconic.
- PLANOLITES**, Nicholson, 1873, Proc. Roy. Soc., No. 144. [*Ety. planos*, wanderer; *lithos*, stone.] Irregularly cylindrical, tortuous casts of supposed worm-tubes. Type *P. vulgaris*.
vulgaris, Nicholson, 1873, Proc. Roy. Soc., No. 144, and Pal. Prov. of Ontario, p. 42, Clinton Gr.
- POLYGNATHUS**, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 361. [*Ety. polys*, many; *gnathos*, jaw.] Minute variously formed teeth and minute tuberculated plates. Type *P. dubius*.
coronatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 365, Ham. Gr.
crassus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 365, Ham. Gr.
cristatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 366, Ham. Gr.
curvatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 366, Ham. Gr.
dubius, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 362, Ham. Gr.
duplicatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 364, Ham. Gr.
erianis, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 366, Ham. Gr.
immersus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 364, Ham. Gr.
linguiformis, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 367, Ham. Gr.
nasutus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 364, Ham. Gr.
palmatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 367, Ham. Gr.
pennatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 366, Ham. Gr.
princeps, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 365, Ham. Gr.
punctatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 367, Ham. Gr.
radiatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 364, Ham. Gr.
serratus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 365, Ham. Gr.
simplex, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 367, Ham. Gr.
solidus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 365, Ham. Gr.
truncatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 366, Ham. Gr.
tuberculatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 366, Ham. Gr.
- PRIONIONUS**, Pander, 1856, Monogr. d. Foss. Fische d. Silur. Syst., p. 28. [*Ety. prionion*, small saw; *odous*, tooth.] Basal portion narrow supporting numerous, delicate denticles and an elongated tapering tooth which extends below the basal portion. Type *P. elegans*.
abbreviatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 359, Ham. Gr.
acicularis, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 360, Ham. Gr.
alatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 361, Ham. Gr.
angulatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 360, Waverly Gr.
armatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 360, Ham. Gr.
clavatus, Hinde, 1879, Quar. Jour. Geo. Soc., vol. 35, p. 360, Ham. Gr.
elegans, Pander, 1856, Monogr. d. Foss. Fische d. Silur. Syst., p. 29, Hud. Riv. Gr.
erraticus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 359, Ham. Gr.
fureatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 358, Hud. Riv. Gr.
panderi, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 361, Ham. Gr.
politus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 358, Hud. Riv. Gr.
radicans, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 350, Hud. Riv. Gr.
spicatus, Hinde, 1879, Quar. Jour. Geo. Soc. Lond., vol. 35, p. 361, Ham. Gr.
- PROTOSCOLEX**, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 89. [*Ety. protos*, first; *skolex*, worm.] Long, slender, numerous segments, both ends obtusely pointed. Type *P. covingtonensis*.
covingtonensis, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 89, Utica Slate Gr.
ornatus, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 90, Utica Slate Gr.
simplex, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 91, Utica Slate Gr.
tenuis, Ulrich, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 90, Utica Slate Gr.
- SALTERELLA**, Billings, 1861, Pal. Foss., vol. 1, p. 17. [*Ety. proper name.*] Small, slender, elongate, conical tubes, consisting of several hollow cones placed one within another, the last one forming the chamber of habitation; surface concentrically or longitudinally striated. Type *S. rugosa*.
billingsi, Safford, 1869, Geo. of Tenn., p. 289, Trenton Gr.
obtusa, Billings, 1861, Pal. Foss., vol. 1, p. 18, Up. Taconic.
pulchella, Billings, 1861, Pal. Foss., vol. 1, p. 18, Up. Taconic.
rugosa, Billings, 1861, Pal. Foss., vol. 1, p. 17, Up. Taconic.
- SCOLITHUS**, Haldeman, 1840, Supp. to Monograph of Limnæades. [*Ety. skolex*, worm;

FIG. 942.—*Salterella rugosa*.

extends below
P. elegans.

Quar. Jour. Geo.
359, Ham. Gr.
Quar. Jour. Geo.
360, Ham. Gr.
Quar. Jour. Geo.
361, Ham. Gr.
Quar. Jour. Geo.
360, Waverly Gr.
Quar. Jour. Geo.
360, Ham. Gr.
Quar. Jour. Geo.
361, Ham. Gr.

Monogr. d. Foss.
29, Hud. Riv. Gr.
Quar. Jour. Geo.
359, Ham. Gr.
Quar. Jour. Geo.
358, Hud. Riv. Gr.
Quar. Jour. Geo.
361, Ham. Gr.

Quar. Jour. Geo.
358, Hud. Riv. Gr.
Quar. Jour. Geo.
361, Ham. Gr.
8, Jour. Cin. Soc.

89. [Ety. *serp.*]
Long, slender, nu-
both ends obtuse.

vingtonensis.
1878, Jour. Cin. S.

89, Utica Slate

our.
vol.

Gr.
our.
vol.

Gr.
our.
vol.

Gr.
861,
17.

ate, Fig. 941.—Proto-
scotex ornatus.

mother, the last one
er of habitation;

y or longitudinally
ted. Type S. ru-

si, Safford, 1869,
of Tenn., p. 289,
ton Gr.

, Billings, 1861, Pal.
vol. 1, p. 18, Up-
nic.

lla, Billings, 1861,
Foss., vol. 1, p. 18,
Taconic.

, Billings, 1861, Pal.
vol. 1, p. 17, Up-
nic.

340, Supp. to Mon-
[Ety. *skotex*, worm;

lithos, stone.] Merely worm furrows,
without organic characters.
canadensis, Billings, 1862, Pal. Foss. vol.
1, p. 90, Potsdam Gr.

FIG. 943.—*Scollithus canadensis*.

linearis, Hall, 1847, Pal. N. Y., vol. 1, p.
2, Potsdam Gr.

verticalis, Hall, 1852, Pal. N. Y., vol. 2, p.
6, Medina sandstone.

tuberosus, Miller & Dyer, 1878, Cont. to
Pal. No. 2, p. 5, Hud. Riv. Gr.

woodi, Whitfield, 1880, Ann. Rep. Geo.
Sur. Wis., p. 45, Potsdam Gr.

Serpula, Linnaeus, 1758, Syst. Nat., 10th ed.
p. 786. [Ety. *serpo*, to creep.] Tube cal-
careous, procumbent, variously curved

or spirally coiled, growing singly or in
groups, attached to marine bodies,

capable of receiving the entire animal;
aperture at the larger extremity sim-
ple and rounded. Type S. *vermicularis*.

insita, White, 1878, Proc. Acad. Nat. Sci.,
p. 37, and Cont. to Pal., No. 8, p. 171,
Coal Meas.

omphalodes, see *Spirorbis omphalodes*.
valvata, see *Spirorbis valvatus*.

SERPULITES, Murchison, 1839, Murch. Sil.
Syst., p. 608. [Ety. *Serpula*, a genus of
annelids.] Tubes smooth, arched, slightly

calcareous, glossy; having two small
longitudinal tubes at opposite points of
the circumference, stronger than the
rest of the shell, and prolonged at the
posterior end. Type S. *longissimus*.

annulatus, Dawson, 1868,
Acad. Geol., p. 312, Car-
boniferous.

dissolutus, Billings, 1862, Pal.
Foss., vol. 1, p. 56, Tren-
ton Gr.

hortonensis, Dawson, 1868,
Acad. Geol., p. 312, Car-
boniferous.

inelegans, Dawson, 1868, Acad. Geol., p.
312, Carboniferous.

murchisoni, Hall, 1861, Geo. Rep. Wis.,
p. 48, Potsdam Gr.

murrayi, Dawson, 1883, Rep. on Redpath
Mus., No. 2, p. 13, Carboniferous.

splendens, Billings, 1850, Can. Nat. and
Geo., vol. 4, p. 470, Chazy Gr.

Sphagodus obliquus, see *Lambriconerites*
obliquus.

SPIRORBIS, Lamarck, 1801, Syst. An. sans
Vert., p. 320. [Sig. spiral-whorl.] Tube
calcareous, solitary, coiled; flat, dex-
tral or sinistral, attached by one side
to some foreign object. Type *Serpula*
spirorbis of Linnaeus.

ammon, Winchell, 1866, Rep. Low. Pen-
insula Mich., p. 97, Ham. Gr.

angulatus, Hall, 1862, 15th Rep. N. Y. St.
Mus. Nat. Hist., p. 84, Ham. Gr.

angulatus, Dawson, 1868. The name was
preoccupied.

annulatus, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 34, and Bull. Am. Mus. Nat.
Hist., p. 92, Warsaw Gr.

annulatus var. *nodulosus*, see S. *nodu-
losus*.

anthracosia, Whitfield, 1881, Am. Jour.
Sci. and Arts, 3d ser. vol. 21, p. 128,
Coal Meas.

arietinus, Dawson, 1869, Rep. of Progr.
Geo. Sur. Can., p. 14, Coal Meas.

arkonensis, Nicholson, 1874, Geo. Mag.,
vol. 1, p. 199, Ham. Gr.

carbonarius, Dawson, 1845,
Quar. Jour. Geo. Soc.,
vol. 1, p. 320, Coal Meas.

cincinnatiensis, Miller &
Dyer, 1878, Jour. Cin.
Soc. Nat. Hist., vol. 1,
p. 38, Hud. Riv. Gr.

flexuosus, Hall, 1863, Trans. Alb. Inst.,
vol. 4, p. 224, Niagara Gr.

inornatus, Hall, 1863, Trans. Alb. Inst.,
vol. 4, p. 224, Niagara Gr.

kinderhookensis, Gurley, 1883, New Carb.
Foss., p. 9. Publication not sufficient.

laxus, Hall, 1859, Pal. N. Y., vol. 3, p.
340, Low. Held. Gr.

nodulosus, Hall, 1858, Trans. Alb. Inst.,
vol. 4, p. 34, and Bull. Am. Mus. Nat.
Hist., p. 93, Warsaw Gr.

obesus, Winchell, 1866, Rep. Low. Pen-
insula Mich., p. 97, Ham. Gr.

omphalodes, Goldfuss, 1826, Germ. Pe-
trel., Up. Held. and Ham. Grs.

orbiculostoma, Swallow, 1858, Trans.
St. Louis Acad. Sci., p. 181, Per-
mian Gr.

spinuliferus, Nicholson, 1875, Pal. Prov.
Ont., p. 83, Ham. Gr.

valvatus, Goldfuss, 1826, (*Serpula valvata*).
Not American.

STAUROCEPHALITES, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 383. [Ety.
Staurocephalus, an existing genus; *lithos*,
stone.] Jaws of more or less elongated
compressed denticulate plates resem-
bling those of the genus *Staurocephalus*.
Type S. *niagarensis*.

niagarensis, Hinde, 1879, Quar. Jour.
Geo. Soc. Lond., vol. 35, p. 383, Nia-
gara Gr.

FIG. 945.—*Spiror-*

Trans. Alb. Inst., vol. 4, bicarbonarius.

p. 224, Niagara Gr.

inornatus, Hall, 1863, Trans. Alb. Inst.,

vol. 4, p. 224, Niagara Gr.

kinderhookensis, Gurley, 1883, New Carb.

Foss., p. 9. Publication not sufficient.

laxus, Hall, 1859, Pal. N. Y., vol. 3, p.

340, Low. Held. Gr.

nodulosus, Hall, 1858, Trans. Alb. Inst.,

vol. 4, p. 34, and Bull. Am. Mus. Nat.

Hist., p. 93, Warsaw Gr.

obesus, Winchell, 1866, Rep. Low. Pen-

insula Mich., p. 97, Ham. Gr.

omphalodes, Goldfuss, 1826, Germ. Pe-

trel., Up. Held. and Ham. Grs.

orbiculostoma, Swallow, 1858, Trans.

St. Louis Acad. Sci., p. 181, Per-

mian Gr.

spinuliferus, Nicholson, 1875, Pal. Prov.

Ont., p. 83, Ham. Gr.

valvatus, Goldfuss, 1826, (*Serpula valvata*).

Not American.

STAUROCEPHALITES, Hinde, 1879, Quar. Jour.

Geo. Soc. Lond., vol. 35, p. 383. [Ety.

Staurocephalus, an existing genus; *lithos*,

stone.] Jaws of more or less elongated

compressed denticulate plates resem-

bling those of the genus *Staurocephalus*.

Type S. *niagarensis*.

niagarensis, Hinde, 1879, Quar. Jour.

Geo. Soc. Lond., vol. 35, p. 383, Nia-

gara Gr.

FIG. 946.—*Walcottia rugosa*.

WALCOTTIA, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 39. [Ety. proper name.] A rugose, flexuous, worm-like furrow. Type *W. rugosa*.
 cookana, Miller & Dyer, 1878, Cont. to Pal., No. 2, p. 11, Hud. Riv. Gr.
rugosa, Miller & Dyer, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 39, Hud. Riv. Gr.

CLASS CRUSTACEA.

THERE is such an immense diversity among Crustaceous animals, it has been found necessary to make subclasses, orders, and suborders, to give intelligible definitions to the classification. They are generally covered with a peculiar calcareous secretion or integument, constituting a cutaneous skeleton, inclosing the soft parts of the body. Segments are united by a membrane, giving flexibility to the armor. There being no way to increase the integument by growth, it is cast off at stated periods, and a new one is secreted to cover the enlarged body. The subclasses are Cirripedia, Entomostraca, Xyphosura, Edriophthalmata, and Podophthalmata.

The Subclass Cirripedia includes only a single order which bears the same name. The animals, when mature, are attached to submarine objects, and are inclosed in a shell composed of several calcareous plates, from an opening in which articulated cirri are exerted and retracted when the animal is alive in search of prey. The common barnacle, which frequently covers the bottoms of ships so as to impede their progress across the ocean, is a representative of this order.

The Subclass Entomostraca is divided into several orders, only three of which are Palæozoic, viz.: Ostracoda, Phyllopoda, and Trilobita. The Ostracoda are minute animals inclosed in a little bivalve shell; the feet and antennæ are protruded between the lower edges of the valves. The Cypris, Daphnia, and Polyphemus are living examples of this order. The Phyllopoda are so named on account of the broad and leaf-like feet. Some of them are covered with a bivalve shell, and others are without such protection. The Palæozoic are bivalve shells. The Estheria, which abound in pools and springs, belong to this order. The Trilobita possessed a cephalic shield, a trilobed thorax composed of segments, which were flexible and allowed the animal to double itself up, and a tail-piece called the pygidium. The order became extinct in the Palæozoic era.

The Subclass Xyphosura has an anterior subcrescentiform carapace, inclosing the cephalothoracic organs, and a posterior abdominal piece, from which a tail spine projects. The upper surface is convex, and the lower concave. There are three orders—Amphipeltida, Euripterida, and Xyphosura. Only a fragment of the shell of the Amphipeltida is known. The Euripterida is also an extinct order. A common form of the Xyphosura is the Limulus, or Horseshoe Crab, which is common on the shores of the tropical seas.

The Subclass Edriophthalmata has the head distinct from the thoracic segments, and therefore has no cephalothorax. The head has a pair of simple compound eyes, not pedunculated. The Palæozoic orders are Amphipoda and Anisopoda.

r, 1878, Jour. Cin.
1, p. 39. [Ely.
rugose, flexuous,
type *W. rugosa*.
er, 1878, Cont. to
d. Riv. Gr.
1878, Jour. Cin.
1, p. 39, Hud

The Order Amphipoda consists of animals that live in water, burrow in sand, or become parasitic on fishes. The abdomen is well developed, and bears limbs for leaping or swimming. They always swim on their sides. The common sand-hopper on the shore of the sea belongs to this order. The Order Anisopoda has a long body, convex above and flattened below, and has affinities with the Isopoda, of which the common wood-louse is an example.

The Subclass Podopthalmata has compound eyes at the extremity of a pair of movable stalks; the head and thorax are generally united, covered by a single piece of shell, and called the cephalothorax; this includes the antennæ, eyes, mouth, jaws, feet, etc. The remaining segments form an abdomen, which frequently terminates in a caudal fin. Common examples are the squill and the small edible crab. There are several orders in this subclass, only three of which are Palæozoic, viz.: Phyllocarida, Decapoda, Tetracapoda.

imals, it has been
e intelligible def-
ecular calcareous
nclosing the soft
flexibility to the
is cast off at stated
The subclasses are
phthalmata.

The Phyllocarida has cephalic, thoracic, and abdominal segments. The carapace has no regular hinge. The living representative is *Nebalia*, which inhabits the sea at moderate depths. The Decapoda are stalk-eyed, and the head and thoracic segments are united in a cephalothorax, incased in a common shell, and have the brauchial organs inclosed on the sides of the cephalothorax. The true thoracic legs are almost always ten, whence the name of the order. The Tetracapoda have their relations with the Decapoda.

n bears the same
jects, and are in-
opening in which
alive in search of
s of ships so as to
order.

SUBCLASS AND ORDER CIRRIPIEDIA.

FAMILY BALANIDÆ.—Palæocrusia, Protobalanus.

FAMILY LEPADIDÆ.—Lepidocoleus, Strobilepis, Turrilepas.

ly three of which
he Ostracoda are
antennæ are pro-
Daphnia, and Po-
so named on ac-
th a bivalve shell,
valve shells. The
r. The Trilobita
, which were flex-
led the pygidium.

SUBCLASS ENTOMOSTRACA.

ORDER OSTRACODA.

FAMILY BEYRICHIIDÆ.—Beyrichia, Beyrichona, Hipponicharion, Primitia.

FAMILY CYPRIDÆ.—Candona.

FAMILY CYTHERIDÆ.—Cytherella, Cytheropsis.

FAMILY FABERIIDÆ.—Faberia.

FAMILY LEPERDITHIDÆ.—Aparchites, Isochilina, Leperditia.

ORDER PHYLLOPODA.

FAMILY ESTHERIIDÆ.—Estheria, Leaia, Schizodiscus.

FAMILY UNCERTAIN.—Lepidilla, Lepiditta.

arapace, inclosing
rom which a tail
cave. There are
a fragment of the
extinct order. A
ab, which is com-

ORDER TRILOBITA.

FAMILY ACIDASPIDÆ.—Acidaspis.

FAMILY AGLASPIDÆ.—Aglaspis.

FAMILY AGNOSTIDÆ.—Agnostus, Microdiscus, Shumardia.

FAMILY ASAPHIDÆ.—Asaphus, Barrandia, Megalaspis, Nileus, Ogygia, Symphysurus.

FAMILY BATHYURIDÆ.—Asaphiscus, Bathyurellus, Bathyuriscus, Bathyurus.

FAMILY BRONTEIDÆ.—Bronteus.

FAMILY CALYMENIDÆ.—Calymene, Homalonotus.

thoracic segments,
simple compound
and Anisopoda.

- FAMILY CERAURIDÆ.—Ceraurus, Sphærocoryphe, Sphærexochus.
 FAMILY CONOCORYPHIDÆ.—Bailiella, Chariocephalus, Conocoryphe, Harttia, Menocephalus, Prototypus.
 FAMILY CYPHASPIDÆ.—Cyphaspis.
 FAMILY DICELLOCEPHALIDÆ.—Dicellocephalus, Pterocephalia, Ptychaspis.
 FAMILY ENCRINURIDÆ.—Amphion, Encrinurus.
 FAMILY ELLIPSOCEPHALIDÆ.—Ellipsocephalus.
 FAMILY HARPIDÆ.—Harpes.
 FAMILY ILLÆNIDÆ.—Illænurus, Illænus.
 FAMILY LICHIDÆ.—Lichas, Terataspis.
 FAMILY OLENIDÆ.—Dolichometopus, Oryctocephalus, Telephus, Triarthrella, Triarthrus.
 FAMILY PARADOXIDÆ.—Anopolenus, Atops, Bathynotus, Elliptocephala, Mesonacis, Olenoides, Paradoxides.
 FAMILY PHACOPIDÆ.—Dalmanites, Phacops.
 FAMILY PROETIDÆ.—Harpides, Phæthonides, Phillipsia, Proetus.
 FAMILY PTYCHOPARIDÆ.—Agraulus, Crepicephalus, Liostracus, Longocephalus, Loganellus, Ptychoparia, Solenopleura.
 FAMILY REMOPLEURIDÆ.—Remopleurides.
 FAMILY TRINUCLEIDÆ.—Ampyx, Dionide, Eudymionia, Trinucleus.
 FAMILY AFFINITY UNCERTAIN.—Pemphigaspis.
 TRACKS SUPPOSED TO BE CRUSTACEAN.—Asaphoidichnus, Climachtichnites, Diplichnites, Protichnites, Rusichnites.

SUBCLASS XIPHOSURA.

ORDER AMPHIPELTIDA.

- FAMILY AMPHIPELTIDÆ.—Amphipeltis.

ORDER EURYPTERIDA.

- FAMILY ECHINOGNATHIDÆ.—Echinognathus.
 FAMILY EURYPTERIDÆ.—Anthraconectes, Dolichopterus, Eurypterella, Eurypterus, Pterygotus, Stylonurus.
 FAMILY HEMIASPIDÆ.—Bunodella.

ORDER XIPHOSURA

- FAMILY BELINURIDÆ.—Belinurus, Euproops, Protolinulus.
 FAMILY CYCLIDÆ.—Cyclus, Dipeltis.

SUBCLASS EDRIOPHTHALMATA.

ORDER AMPHIPODA.

- FAMILY DIPLOSTYLIDÆ.—Diplostylus.

ORDER ANISOPODA.

- FAMILY ACANTHOTELSONIDÆ.—Acanthotelson.

SUBCLASS PODOPTHALMATA.

ORDER PHYLLOCARIDA.

FAMILY CERATIOCARIDÆ.—Ceraticaris, Colpocaris, Echinocaris, Elymlocaris, Ribeiria (?), Solenocaris, Tropicocaris.

FAMILY DISCINOCARIDÆ.—Dipterocaris, Spathiocaris.

FAMILY PINACARIDÆ.—Dithyrocaris, Mesothyra.

FAMILY PROTOCARIDÆ.—Protocaris.

FAMILY RACHURIDÆ.—Rachura.

FAMILY RHINOCARIDÆ.—Rhinocaris.

ORDER DECAPODA.

FAMILY ANTHRACARIDÆ.—Anthrapalæmon.

FAMILY CARIDIDÆ.—Palæopalæmon.

FAMILY PALÆOCARIDÆ.—Gampsonyx, Palæocaris.

ORDER TETRADECAPODA.

FAMILY ARCHÆOCARIDÆ.—Archæocaris.

FIG. 947.—*Acanthotelson eveni*. A, dorsal view; st, stylet; t, telson; B, side view; a and z, antennae; f, anterior leg; C, enlarged stylet.

FIG. 948.—*Acanthotelson eveni*. Enlarged; H, anterior legs and antennae; S, punctures left by spines on the lower side.

Acantholoma, syn. for *Acidaspis*.
spinosa, syn. for *Acidaspis tuberculata*.

ACANTHOTELSON, Meek & Worthen, 1860, Proc. Acad. Nat. Sci., p. 47. [Ety. *akantha*, spine; *telson*, end.] Superior antennae as long as the inferior, flagella longer than the peduncles; head about the length of the two anterior

thoracic segments; thoracic and abdominal segments about the same length; anterior thoracic legs longer than the others; telson simple, long, spine-like, laterally compressed; stylets with second segments longer than first, and similar to the telson. Type *A. stimpsoni*. eveni, Meek & Worthen, 1868, Am. Jour. Sci., vol.

FIG. 949.—*Acanthotelson stimpsoni*. Dorsal view.

FIG. 950.—*Acanthotelson stimpsoni*. Enlarged 3 diam.; t, a protuberance that may be an eye.

46, p. 28, and Geo. Sur. Ill., vol. 3, p. 551, Coal Meas.

inequalis, Meek & Worthen, syn. for *Palæocaris typus*.

stimpsoni, Meek & Worthen, 1865, Proc. Acad. Nat. Sci., p. 47, and Geo. Sur. Ill., vol. 2, p. 401, Low. Coal Meas.

ACIDASPIIS, Murchison, 1839, Sil. Syst., p. 658. [Ety. *akis*, a point; *aspis*, shield.] Cephalic shield, semicircular, margin thickened and spinous, lateral angles produced in spines; glabella convex, narrow in front, not reaching the margin, two lobes on each side, and having a large spine projecting backward, eyes prominent; thorax with eight segments; pleurae wide, and terminating in spines; pygidium small, axis short, of two joints, sides depressed, one segmental furrow, long spine extending backward from the margin at each side, and smaller spines from the other parts of the margin. Type A. brighti.

FIG. 951.—*Aeidaspis anchoralis*. Cephalic shield.

anchoralis, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 349, Hud. Riv. Gr.

callicera, Hall, 1888, Pal. N. Y., vol. 7, p. 69, Up. Held. Gr.

ceralepta, Anthony, 1838, (Cera^{ph}cephala *ceralepta*), Am. Jour. Sci., vol. 34, p. 379. Not defined so as to be recognized.

cincinnatiensis, Meek, 1873, Ohio Pal., vol. 1, p. 167, Hud. Riv. Gr.

crocotus, Locke, 1843, Am. Jour. Sci., vol. 44, p. 347, and Ohio Pal., vol. 1, p. 165, Hud. Riv. Gr. The word is misspelled; it should be *crocotus*.

danai, Hall, 1862, Geo. Sur. Wis., p. 423, Niagara Gr.

eripis, see *Terataspis eripis*.

fimbriata, Hall, 1879, Desc. New Spec. Foss., p. 20, and 11th Rep. Geo. and Nat. Hist. Ind., p. 334, Niagara Gr.

grandis, see *Terataspis grandis*.

halli, Shumard, 1855, Geo. Sur. Mo., p. 200, Trenton Gr.

hamata, Conrad, 1841, (Dicanurus *hamatus*), Ann. Rep. N. Y., p. 48, and Pal. N. Y., vol. 3, p. 371, Low. Held. Gr.

horani, Billings, 1859, Rep. of Progr. Geo. Sur. Can., p. 341, Trenton Gr.

ida. avn. for *Acidaspis danai*.

onealli, S. A. Miller, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 86, Hud. Riv. Gr.

parvula, Walcott, 1877, 31st Rep. N. Y. St. Mus. Nat. Hist., p. 69, Trenton Gr.

romingeri, Hall, 1888, Pal. N. Y., vol. 7, p. 71, Ham. Gr.

spiniger, see *Bathyrus spiniger*.

trentonensis, Hall, 1847, Pal. N. Y., vol. 1, p. 240, Trenton Gr.

FIG. 953.—*Aeidaspis onealli*. Pygidium.

tuberculata, Conrad, 1840, Ann. Rep. N. Y., p. 205, and Pal. N. Y., vol. 3, p. 368, Low. Held. Gr.

AGLASPIIS, Hall, 1862, Can. Nat. and Geo., vol. 7, p. 443, and 16th Rep. N. Y. St. Mus. Nat. Hist., p. 181. [Ety. *aglaos*, bright; *aspis*, shield.] Cephalic shield somewhat semielliptical, wider than long, sinus in front; glabella narrow, conical; eyes prominent, and situate anterior to the middle; thorax having eight segments; pygidium small, and terminating in a single spine. Type A. barrandii.

barrandii, Hall, 1862, Can. Nat. and Geo., vol. 7, p. 443, and 16th Rep. N. Y. St. Mus. Nat. Hist., p. 181, Potsdam Gr.

eatoni, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 52, and Geo. of Wis., vol. 4, p. 192, Potsdam Gr.

AGNOSTOS, Brongniart, 1822, Hist. Nat. Crust. Foss., p. 38. [Ety. *agnostos*, obscure.] Body elongate, elliptical; cephalic shield and pygidium subequal, subrotund, or longer than wide, convex rim on the border; glabella convex; no eyes; no facial sutures; two thoracic segments. Type A. pisiformis.

acadicus, Hartt, 1868, Acad. Geol., p. 655, St. John Gr.

acadicus var. *declivis*, Matthew, 1885, Trans. Roy. Soc. Can., p. 70, St. John Gr.

acutitubus, Matthew, 1885, Trans. Roy. Soc. Can., p. 73, St. John Gr.

americanus, Billings, 1860, Can. Nat. and Geol., vol. 5, p. 301, and Pal. Foss., p. 395, Up. Taconic.

bidens, Meek, 1873, 6th Rep. Hayden's Geo. Sur. Terr., p. 463, and Monog. U. S. Geo. Sur., vol. 8, p. 26, Prospect Mountain Gr., Up. Taconic.

canadensis, Billings, 1860, Can. Nat. and Geol., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 397, Up. Taconic.

coloradoensis, Shumard, 1861, Am. Jour. Sci. and Arts, 2d ser., vol. 32, p. 218, Up. Taconic.

communis, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 288, Prospect Mountain Gr., Up. Taconic.

? *di-parilis*, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 179, Potsdam Gr.

fabius, Billings, 1865, Pal. Foss., vol. 1, p. 298, Up. Taconic.

galba, Billings, 1865, Pal. Foss., vol. 1, p. 297, Up. Taconic.

interstrictus, White, 1874, Rep. Invert. Foss., p. 7, and Geo. Sur. W. 100th Mer., vol. 4, p. 38, Up. Taconic.

? *joc-pha*, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 178, Potsdam Gr.

latus, see *Beyrichia lata*.

lobatus, see *Microdiscus lobatus*.

maladenis, Meek, 1873, Hylen's Sur. Terr., p. 464. Not properly defined.

neon, Hall & Whitfield, 1877, U. S. Geo. Expl., 4th parallel, vol. 4, p. 229, Prospect Mountain Gr., Up. Taconic.

840, Ann. Rep. N. Y., vol. 3, p.

. Nat. and Geo., h Rep. N. Y. St. 81. [Ety. *aglus*, Cephalic shield al, wider than glabella narrow, ent, and situate e; thorax having idium small, and e spine. Type A.

n. Nat. and Geo., h Rep. N. Y. St. 1, Potsdam Gr. Ann. Rep. Geo. Geo. of Wis., vol.

1822, Hist. Nat. Ety. *agnostos*, obte, elliptical; ce-gidium subequal, an wide, convex glabella convex; ares; two thoracic isiformis. cad. Geol., p. 655,

Matthew, 1885, p. 70, St. John Gr. 1885, Trans. Roy. John Gr. 1860, Can. Nat. and and Pal. Foss., p.

h Rep. Hayden's and Monog. U. S. p, Prospect Mount-

30, Can. Nat. and and Pal. Foss., vol.

, 1861, Am. Jour., vol. 32, p. 218,

Whitfield, 1877, U. S. al. vol. 4, p. ain Gr., Up. Ta-

16th Rep. N. Y. 179, Potsdam Gr. l. Foss., vol. 1, p.

l. Foss., vol. 1, p.

1874, Rep. Invert. o. Sur. W. 100th e. Taconic. th R-p. N. Y., St. 8, Potsdam Gr.

lobatus. H-yden's Sur. perly d. fined. 1877, U. S. Geo. pl. 4, p. 229, Pros- p. Taconic.

nobilis, Ford, 1872, Am. Jour. Sci., 3d ser., vol. 3, p. 421, Up. Taconic. obtusilobus, Matthew, 1885, Trans. Roy. Soc. Can., p. 72, St. John Gr. orion, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 397, Up. Taconic.

? parilis, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 179, Potsdam Gr. partitus, Matthew, 1885, Trans. Roy. Soc. Can., p. 68, St. John Gr.

prolongus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, p. 230, Prospect Mountain Gr., Up. Taconic.

regulus, Matthew, 1885, Trans. Roy. Soc. Can., p. 67, St. John Gr.

rex, Barrande. Probably not an American species, but illustrative of the genus.

richmondensis, Walcott, 1884, Monogr. U. S. Geo. Sur., vol. 8, p. 24, Prospect Mountain Gr., Up. Taconic.

seclusus, Walcott, 1884, Monogr. U. S. Geo. Sur., vol. 8, p. 25, Prospect Mountain Gr., Up. Taconic.

similis, Hartt, 1868, Acad. Geol., p. 656, St. John Gr.

tessella, Matthew, 1885, Trans. Roy. Soc. Can., p. 71, St. John Gr.

tumidosus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 231, Up. Taconic.

umbo, Matthew, 1885, Trans. Roy. Soc. Can., p. 71, St. John Gr.

vir, Matthew, 1885, Trans. Roy. Soc. Can., p. 69, St. John Gr.

vir var. concinnus, Matthew, 1885, Trans. Roy. Soc. Can., p. 70, St. John Gr.

AGRAULUS, Hawle & Corda, 1847, Prodr. einer Monographie der bomischen Trilobiten, p. 142. [Ety. *agraul*, dwelling in the fields.] Body elongate-ovate; cephalic shield large, semicircular to lunate, with a wide margin, that merges in the cheeks; glabella convex, narrowed and rounded in front, conoidal, three or four lateral furrows on each side, margined in front, neck furrow distinct; eyes small, distant from glabella or submarginal; facial sutures, beginning near the lateral posterior angles, are directed forward, curving over the eyes to the anterior margin, nearly in parallel lines; cheeks small, narrow; sixteen thoracic segments, axial lobe convex; pygidium small, rounded, three segments; hypostoma oval, truncated anteriorly. Type A. reticephalus.

affinis, Billings, 1874, Pal. Foss., vol. 2, p. 72, Up. Taconic.

articephalus, Matthew, 1885, Trans. Roy. Soc. Can., p. 75, St. John Gr.

bipunctatus, Shumard, 1863, (Arionellus bipunctatus,) Trans. St. Louis Acad. Sci., vol. 2, p. 101, Potsdam Gr. Poorly defined; probably belongs to another genus.

convexus, Whitfield, 1877, (Arionellus convexus,) Geo. Sur. Wis., vol. 4, p. 190, Potsdam Gr. Founded upon a fragment, and may belong to another genus.

cylindricus, Billings, 1860, (Arionellus cylindricus, Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 406, Up. Taconic or St. John Gr.

globosus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 61, Up. Taconic.

hallanus, Matthew, 1887, Trans. Roy. Soc. Can., p. 132, St. John Gr.

planus, Shumard, 1861, (Arionellus planus,) Am. Jour. Sci. and Arts, 2d series, vol. 32, p. 219, Potsdam Gr.

pustulatus, Walcott, 1879, (Arionellus pustulatus,) 31st Rep. N. Y. St. Mus. Nat. Hist., p. 68, Chazy Gr.

quadrangularis, Whitfield, 1884, (Arionellus quadrangularis,) Bull. Am. Mus. Nat. Hist., vol. 1, p. 139, Up. Taconic.

socialis, Billings, 1874, Pal. Foss., vol. 2, p. 71, Up. Taconic or St. John Gr.

strenuus, Billings, 1874, Pal. Foss., vol. 2, p. 71, Up. Taconic or St. John Gr.

subclavatus, Billings, 1860, (Arionellus subclavatus,) Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 406, Up. Taconic or Quebec Gr.

texanus, Shumard, 1861, (Arionellus texanus,) Am. Jour. Sci. and Arts, 2d ser., vol. 32, p. 218, Potsdam Gr. or Up. Taconic.

tripunctatus, Whitfield, 1876, R-p. Recon. Up. Mo. to Yel. Nat. Park, p. 141, Potsdam Gr. or Up. Taconic.

whitfieldanus, Matthew, 1887, Trans. Roy. Soc. Can., p. 130, St. John Gr.

whitfieldanus var. compressus, Matthew 1887, Trans. Roy. Soc. Can., p. 130, St. John Gr.

woosteri, Whitfield, 1878, Geo. Sur. Wis., vol. 4, p. 189, Potsdam Gr.

AMPHION, Pander, 1830, Beitrage zur Geognosie des Russischen Reiches, p. 139. [Ety. mythological name.] Cephalic shield short, transverse; glabella convex or subrectangular, three pairs of furrows, front inclosing a small forehead lobe; eyes small; facial suture, behind the eyes, ending on the exterior margin in advance of the rounded angles; thorax 15 to 18 articulations; pleurae without grooves; pygidium with short axis and pleurae with free terminations; labrum pointed, Fig. 956—Amphion canadensis. Pygidium.

convex, Whitfield, 1877, (Arionellus convexus,) Geo. Sur. Wis., vol. 4, p. 190, Potsdam Gr. Founded upon a fragment, and may belong to another genus.

cylindricus, Billings, 1860, (Arionellus cylindricus, Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 406, Up. Taconic or St. John Gr.

globosus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 61, Up. Taconic.

hallanus, Matthew, 1887, Trans. Roy. Soc. Can., p. 132, St. John Gr.

planus, Shumard, 1861, (Arionellus planus,) Am. Jour. Sci. and Arts, 2d series, vol. 32, p. 219, Potsdam Gr.

pustulatus, Walcott, 1879, (Arionellus pustulatus,) 31st Rep. N. Y. St. Mus. Nat. Hist., p. 68, Chazy Gr.

quadrangularis, Whitfield, 1884, (Arionellus quadrangularis,) Bull. Am. Mus. Nat. Hist., vol. 1, p. 139, Up. Taconic.

socialis, Billings, 1874, Pal. Foss., vol. 2, p. 71, Up. Taconic or St. John Gr.

strenuus, Billings, 1874, Pal. Foss., vol. 2, p. 71, Up. Taconic or St. John Gr.

subclavatus, Billings, 1860, (Arionellus subclavatus,) Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 406, Up. Taconic or Quebec Gr.

texanus, Shumard, 1861, (Arionellus texanus,) Am. Jour. Sci. and Arts, 2d ser., vol. 32, p. 218, Potsdam Gr. or Up. Taconic.

tripunctatus, Whitfield, 1876, R-p. Recon. Up. Mo. to Yel. Nat. Park, p. 141, Potsdam Gr. or Up. Taconic.

whitfieldanus, Matthew, 1887, Trans. Roy. Soc. Can., p. 130, St. John Gr.

whitfieldanus var. compressus, Matthew 1887, Trans. Roy. Soc. Can., p. 130, St. John Gr.

woosteri, Whitfield, 1878, Geo. Sur. Wis., vol. 4, p. 189, Potsdam Gr.

AMPHION, Pander, 1830, Beitrage zur Geognosie des Russischen Reiches, p. 139. [Ety. mythological name.] Cephalic shield short, transverse; glabella convex or subrectangular, three pairs of furrows, front inclosing a small forehead lobe; eyes small; facial suture, behind the eyes, ending on the exterior margin in advance of the rounded angles; thorax 15 to 18 articulations; pleurae without grooves; pygidium with short axis and pleurae with free terminations; labrum pointed, Fig. 956—Amphion canadensis. Pygidium.

convex, Whitfield, 1877, (Arionellus convexus,) Geo. Sur. Wis., vol. 4, p. 190, Potsdam Gr. Founded upon a fragment, and may belong to another genus.

cylindricus, Billings, 1860, (Arionellus cylindricus, Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 406, Up. Taconic or St. John Gr.

globosus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 61, Up. Taconic.

hallanus, Matthew, 1887, Trans. Roy. Soc. Can., p. 132, St. John Gr.

planus, Shumard, 1861, (Arionellus planus,) Am. Jour. Sci. and Arts, 2d series, vol. 32, p. 219, Potsdam Gr.

pustulatus, Walcott, 1879, (Arionellus pustulatus,) 31st Rep. N. Y. St. Mus. Nat. Hist., p. 68, Chazy Gr.

quadrangularis, Whitfield, 1884, (Arionellus quadrangularis,) Bull. Am. Mus. Nat. Hist., vol. 1, p. 139, Up. Taconic.

socialis, Billings, 1874, Pal. Foss., vol. 2, p. 71, Up. Taconic or St. John Gr.

strenuus, Billings, 1874, Pal. Foss., vol. 2, p. 71, Up. Taconic or St. John Gr.

subclavatus, Billings, 1860, (Arionellus subclavatus,) Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 406, Up. Taconic or Quebec Gr.

texanus, Shumard, 1861, (Arionellus texanus,) Am. Jour. Sci. and Arts, 2d ser., vol. 32, p. 218, Potsdam Gr. or Up. Taconic.

tripunctatus, Whitfield, 1876, R-p. Recon. Up. Mo. to Yel. Nat. Park, p. 141, Potsdam Gr. or Up. Taconic.

whitfieldanus, Matthew, 1887, Trans. Roy. Soc. Can., p. 130, St. John Gr.

whitfieldanus var. compressus, Matthew 1887, Trans. Roy. Soc. Can., p. 130, St. John Gr.

woosteri, Whitfield, 1878, Geo. Sur. Wis., vol. 4, p. 189, Potsdam Gr.

AMPHION, Pander, 1830, Beitrage zur Geognosie des Russischen Reiches, p. 139. [Ety. mythological name.] Cephalic shield short, transverse; glabella convex or subrectangular, three pairs of furrows, front inclosing a small forehead lobe; eyes small; facial suture, behind the eyes, ending on the exterior margin in advance of the rounded angles; thorax 15 to 18 articulations; pleurae without grooves; pygidium with short axis and pleurae with free terminations; labrum pointed, Fig. 956—Amphion canadensis. Pygidium.

FIG. 955. *Agraulus cylindricus*.

FIG. 954. *Aagnostus rex*.

FIG. 956—*Amphion canadensis*. Pygidium.

- canadensis, Billings, 1859, Can. Nat. and Geo., vol. 4, p. 381, Chazy Gr.
 convexus, Billings, 1865, Pal. Foss., vol. 1, p. 322, Quebec Gr.
 insularis, Billings, 1865, Pal. Foss., vol. 1, p. 290, Quebec Gr.
 julius, Billings, 1865, Pal. Foss., vol. 1, p. 290, Quebec Gr.
 matutinus, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 222, Potsdam Gr.
multisegmentatus, see *Encrinurus multisegmentatus*.
 nevadensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 94, Chazy Gr.
 salteri, Billings, 1861, Can. Nat. and Geo., vol. 6, Calciferous Gr.
 westoni, Billings, 1865, Pal. Foss., vol. 1, p. 321, Quebec Gr.

AMPHIPELTIS, Salter, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 75. [Ety. *amphi*, on both sides; *peltis*, provided with a shield or buckler.] Carapace oblong, oval, rounded in front, more truncate behind; thorax with 9 segments, 5 project beyond the carapace and 4 concealed beneath it; tail-piece semicircular, as wide as the abdomen, and as long as the last three segments taken together. Type A. paradoxus.

FIG. 957.—*Amphipeltis paradoxus*.

paradoxus, Salter, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 76, and Acad. Geo., p. 523, Up. Devonian.
AMPYX, Dalman, 1827, Über die palæozöen oder die sogenannten Trilobiten, p. 53. [Ety. *ampyx*, head-band.] Cephalic shield somewhat trigonal; glabella large, prominent, narrow behind, and projecting upward and forward anteriorly; cheeks flattened, posterior angles produced; no eyes or facial sutures; thoracic segments 5 or 6, flattened, sides straight, divided by a diagonal pleural groove; pygidium subtrigonal, nearly as large as the cephalic shield; one anterior segmental furrow; axis faintly marked with transverse furrows. Type A. nasutus.

halli, Billings, 1861, Pal. Foss., vol. 1, p. 24, Chazy Gr.
lævisculus, Billings, 1865, Pal. Foss., vol. 1, p. 295, Quebec Gr.
normalis, Billings, 1865, Pal. Foss., vol. 1, p. 295, Quebec Gr.
rutilius, Billings, 1865, Pal. Foss., vol. 1, p. 296, Quebec Gr.
semicostatus, Billings, 1865, Pal. Foss., vol. 1, p. 297, Quebec Gr.

Angelina hitchcocki, see *Prototypus hitchcocki*.

Anomocere, Angelin, 1852, Pal. Scand., p. 24. This genus is not yet known in America.

(f) *parvum*, Walcott, 1885, Mon. U. S. Geo. Sur., vol. 8, p. 59, Up. Taconic. This species is founded on a fragment of the cephalic shield and the generic reference is only provisional.

ANOPOLENUS, Salter, 1864, Quar. Jour. Geo. Soc., vol. 20, p. 236, and vol. 21, p. 477. [Ety. *a*, without; *ops*, an eye; *Olenus*, agenus.] Elongated, depressed; cephalic shield semicircular with prolonged spines, and clavate glabella having 4 pairs of furrows; fixed cheeks, large, punctate, strongly margined, each a quarter of a circle in shape, and reaching nearly to the front of the glabella, against which the long eyes abut; thence the facial suture curves outward, and is marginal in front; the long eye-lobe, which forms the margin of the fixed cheeks, reaches quite to the glabella in front, and nearly to the posterior angle below; free cheeks are a narrow band

FIG. 959.—*Anopolenus venustus*.

margined and reaching only three-fourths down the fixed cheek; pygidium wide, expanded, but narrower than the thorax, widely marginate, and serrated by 6 or 8 marginal spines. Type A. henrici.

venustus, Billings, 1874, Pal. Foss., vol. 2, p. 73, Up. Taconic.

ANTHRACONECTES, Meek & Worthen, 1868, Am. Jour. Sci., vol. 46, p. 21, and Geo. Sur. Ill., vol. 3, p. 544. [Ety. *anthrax*, coal; *nectos*, swimming.] Distinguished from *Eurypterus* by the absence of lateral spines at the articulations of the legs, which terminate in single points, and in the great length and simple extremity of the mesial appendage of its operculum, as well as in the possession of two little spatulate supplementary pieces. Type A. mazonensis.

mazonensis, Meek & Worthen, 1868, Am. Jour. Sci. and Arts, vol. 46, p. 21, and Geo. Sur. Ill., vol. 3, p. 544, Coal Meas.

FIG. 960.—*Anthraconectes mazonensis*. Hypostoma enlarged to show the scale-like sculpturing.

[AMP.—ANT.]

prototypus hitch-

Pal. Scand., p.
t yet known ins, Mon. U. S. Geo.
Taconic. This
a fragment of the
he generic refer-
al.

Quar. Jour. Geo.
and vol. 21, p. 477.
an eye; *Olenus*,
epressed; cephalic
with prolonged
labella having 4
ed cheeks, large,
margined, each a
shape, and reach-
t of the glabella,
long eyes abut;
nture curves out-
l, and is marginal
ont; the long eye-
, which forms the
gin of the fixed
eks, reaches quite
the glabella in
t, and nearly to
posterior angle
ow; free cheeks
a narrow band
hing only three-
ed cheek; pygid-
ed, but narrower
widely marginate,
8 marginal spines.

Pal. Foss., vol. 2.

Fig. 960.—*Anthraconecetes mazonensis*. Hypostoma enlarged to show the scale-like sculpturing.

culum, as well as
two little spatu-
pieces. Type A.

Worthen, 1868, Am.
col. 46, p. 21, and
3, p. 544, Coal

Fig. 961.—*Anthraconecetes mazonensis*. *a, b, c*, Legs, crushed and broken; *h*, ends of dorsal half of body segments; *m*, hypostoma; *P*, swimming paddle broken; *t*, natural articulation; *g*, basal joints of same; *x*, enlarged surface markings; *M*, mesial appendage of operculum; 1, 2, 3, articulations; *z*, lateral alae of operculum; *s, s*, accessory pieces; *o*, position of mouth.

ANTHRACOPALEMON, Salter, 1861, Quar. Jour.

Fig. 962.—*Anthracopalemon gracilis*. Dorsal view somewhat enlarged.

front margin serrate; outer antennæ have wide, square basal joints; second and third joints not much oblique; the rest about as broad as long; ab-

domen of six joints, as broad as long, pleuræ except the second pointed; telson broad, appendages to the penul-

Fig. 963.—*Anthracopalemon gracilis*. Caudal parts and one abdominal segment in advance of the telson—all enlarged. *a*, Small, terminal palette; *d, d*, two accessory lamellæ; *c, c*, lateral lamellæ or fins.

timate joint double on each side, subtrigonal, broad, lateral fins divided. Type A. grossarti.

gracilis, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 50, and Geo. Sur. Ill., vol. 2, p. 407, Coal Meas. hillanus, Dawson, 1877, Geo. Mag., vol. 4, p. 56, Coal Meas.

APARCHITES, Jones, 1889, Ann. and Mag. Nat. Hist., 6th ser., vol. 3, p. 384. [Ety. *aparche*, first.] In form like *Lepeditia*, but smaller and without ocular or muscular spot, and having no overlap on the ventral margin. Type A. whitevesi. whitevesi, Jones, 1889, Ann. and Mag. Nat. Hist., 6th ser., vol. 3, p. 384, Trenton Gr. ARCHÆOCARIS, Meek, 1872, Proc. Acad. Nat. Sci. Phil., p. 335. [Ety. *archaios*, ancient; *karis*, shrimp.] Cephalothorax about equaling in length $3\frac{1}{2}$ segments, subtrigonal, pointed in front, truncated and sinuous posteriorly; abdomen with six imbricating segments; telson as long as $3\frac{1}{2}$ abdominal segments with a stylet on each side. Type A. vermiformis.

Fig. 964.—*Archæocaris vermiformis*.

- vermiformis, Meek, 1872, Proc. Acad. Nat. Sci. Phil., p. 335, and Ohio Pal., vol. 2, p. 321, Subcarboniferous.
- Arctinurus*, Castelnau, 1843, Syst. Syl., p. 21, syn. for *Lichas*.
- Arctusina*, Barrande, 1852, Syst. Sil. Boh. Not yet known as an American genus.
- (?) *americana*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 62, Potsdam Gr. Founded upon a fragment of a cephalic shield which does not belong to this genus.
- Arges*, Goldfuss, 1839, Nova Acta Phys. Acad. Caes. Leop. Nat. Cur. Not American.
- Arionellus*, Barrande, 1852, Syst. Sil. Boh., syn. for *Agraulus*.
- bipunctatus*, see *Agraulus bipunctatus*.
- convexus*, see *Agraulus convexus*.
- cylindricus*, see *Agraulus cylindricus*.
- owenii*, see *Crepicephalus owenii*.
- planus*, see *Agraulus planus*.
- pustulatus*, see *Agraulus pustulatus*.
- quadrangularis*, see *Agraulus quadrangularis*.
- subclavatus*, see *Agraulus subclavatus*.
- texanus*, see *Agraulus texanus*.
- tripunctatus*, see *Agraulus tripunctatus*.
- ARISTOZOE*, Barrande, 1872, Syst. Sil. Boh., vol. 1, p. 477. [Ety. *aristos*, best; *zoon*, animal.] Carapace bivalve; test thin; hinge-line straight; ventral margin grooved and reflected; tubercle near anterior margin. Type *A. bisulcata*.
- rotundata*, Walcott, 1887, Am. Jour. Sci. and Arts, 3d ser., vol. 4, p. 193, Up. Taconic.
- troyensis*, Ford, 1873, (Leperditia troyensis,) Am. Jour. Sci. and Arts, 3d ser., vol. 6, p. 138, Up. Taconic.
- ASAPHISCUS*, Meek, 1873, 6th Rep. Hayden's Geo. Sur. Terr., p. 485. [Ety. from the genus *Asaphus*.] Distinguished from *Asaphus* by having nine thoracic segments, a conical and well-defined glabella, without lateral lobes, the furrow at the anterior margin of the head, and less arcuate eyes more remote from the glabella; distinguished from *Bathurellus* by having its conical glabelladepressed, and the margin of the head, in front, first convex, and sloping forward into a transverse mesial furrow, and then rising in a convex margin; the mesial lobe of the pygidium is longer, and the free margins narrower, less flattened and slate.
- Type *A. wheeleri*.
- bradleyi*, Meek, 1873, 6th Rep. Hayden's Geo. Sur. Terr., p. 484, Up. Taconic.

FIG. 965.—*Asaphiscus wheeleri*.

wheeleri, Meek, 1873, 6th Rep. Hayden's Geo. Sur. Terr., p. 485, and Geo. Sur. W. 100th Mer., vol. 4, p. 43, Up. Taconic.

ASAPHOIDICHNUS, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 217. [Ety. *Asaphus*, a genus; *ichnos*, form; *ichnos*, track.] A track supposed to have been made by a crustacean.

Type *A. trifidus*.

dyeri, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 219, Utica Slate Gr.

trifidus, S. A. Miller, 1880, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 218, Utica Slate Gr.

ASAPIUS, Brongniart, 1822, Hist., Nat. Crust. Foss., p. 17. [Ety. *asaphus*, uncertain, obscure.] Body somewhat elliptical, sides straightened; cephalic shield and pygidium nearly equal and somewhat semielliptical; glabella contracted between the eyes; eyes large, smooth; facial sutures extending forward and outward in advance of the eyes, and then curving to the middle of the front margin, posteriorly extending obliquely outward, and cutting the posterior margin of the cephalic shield within the lateral angles; thoracic segments 8, with wide, nearly straight pleural grooves; axis of pygidium, when traceable, elongate conic, segments usually indistinct. Type *A. cornigerus*.

acantholeurus, see *Dalmanites acantholeurus*.

alacer, Billings, 1866, Catal. Sil. Foss. Antic., p. 20, Hud. Riv. Gr.

aspectans, see *Dalmanites aspectans*.

astragaboles, Green, 1834, Am. Jour. Sci., vol. 25, p. 325. Probably founded upon the pygidium of a *Phacops*.

barrandi, Hall, 1851, Lake Sup. Land Dist., p. 210, Birdseye Gr.

canadensis, Chapman, 1856, Can. Jour. vol. 2, p. 47, Trenton Gr.

canalis, Conrad, 1847, Pal. N. Y., vol. 1, p. 25, Chazy Gr.

caribouensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 98, Quebec Gr.

FIG. 966.—*Asaphoidichnus trifidus*.

Rep. Hayden's
, and Geo. Sur.
p. 43, Up. Ta-

caudatus, Green, syn. for *Dalmanites*
limulurus. (?)

cordieri, Castelnau, syn. for *Dalmanites*
limulurus.

coryceus, see *Proetus coryceus*.

crypturus, Green, 1834, Trans. Geo. Soc.
Pa., vol. 1, p. 37. Not an *Asaphus*;
form not determined.

(?) *curiosus*, Billings, 1865, Pal. Foss.,
vol. 1, p. 318, Quebec Gr.

denticulatus, see *Dalmanites denticulatus*.

✕ *ditmarsie*, Honeyman, 1879, Proc. Nova
Scotia Inst., vol. 5, p. 18, Low. Sil.

diurus, Green, 1830, Am. Jour. Sci., vol.
39, p. 40, Niagara Gr. Probably the
fragment of a *Dalmanites*.

edwardsi, Castelnau, syn. for *Dalmanites*
limulurus.

extans, see *Bathyurus extans*.

gigas, Dekay, 1825,
(*Isotelus gigas*,)
Ann. Lyc. Nat. Hist.

N. Y., vol. 1, p. 174,
and Pal. N. Y., vol.

1, p. 231, Trenton
and Hud. Riv.

Gr.

(?) *goniocercus*, Meek,
1873, Hayden's Geo.

Sur. Terr., p. 480,
Quebec Gr. or Up.

Taconic. Probably
a *Megalaspis*.

(?) *goniurus*, Billings,
1860, Can. Nat., vol.

5, p. 301, Up. Ta-
conic. Not defined

so as to be recog-
nized.

grattonensis, Meek &
Worthen, 1870, Proc. Acad. Nat. Sci.,

p. 54, Hud. Riv. Gr.

halli, Conrad, syn. for *Dalmanites boothi*.

halli, Chapman, 1858, Ann. and Mag.
Nat. Hist., 3d ser., vol. 2, p. 14, Tren-
ton Gr.

hausmani, Brongniart, as identified by
D'Archiac and Verneuil. Not Ameri-
can.

hincsi, Salter, 1859, Ann. and Mag. Nat.
Hist., 3d ser., vol. 4, p. 2, Trenton Gr.

homalonotoides, Walcott, 1877, 31st Rep.
N. Y. St. Mus. Nat. Hist., p. 71, Tren-
ton Gr.

(?) *huttoni*, Billings, 1865, Pal. Foss., vol.
1, p. 271, Quebec Gr. or Up. Taconic.

(?) *ilienooides*, Billings, 1860, Can. Nat.
vol. 5, p. 301, Up. Taconic.

iowensis, Owen, 1852, Geo. Wis., Iowa,
and Minn., p. 577, Trenton Gr.

latiostatus, syn. for *Dalmanites anchiops*.

(?) *latimarginatus*, Hull, 1847, Pal. N. Y.,
vol. 1, p. 253, Utica Slate Gr.

limulurus, see *Dalmanites limulurus*.

marginalis, Hall, 1847, Pal. N. Y., vol. 1,
p. 24, Chazy Gr.

megalapthalmus, Troost, 1840, 5th, Geo.
Tenn. Niagara Gr. Not clearly defined,
but probably a *Dalmanites*.

megistus, Locke, 1841, (*Isotelus megis-
tus*,) Trans. Am. Geo. and Nat., p. 221,
Trenton and

Hud. Riv. Grs.

micurus, see *Dal-
manites micu-
rus*.

(?) *morrisi*, Bill-
ings, 1865, Pal.

Foss., vol. 1, p.
272, Quebec Gr.

or Up. Taconic.

murchisoni, Castel-
nau, syn. for *A.*

gigas.

myrmecophorus, see
Dalmanites

*myrmecopho-
rus*.

nasutus, see *Dal-
manites nasu-
tus*.

nodulatus, Hall,
1847, Pal. N. Y.,

vol. 1, p. 248.

Not defined so
as to establish a species.

notus, Billings, 1866, Catal. Sil. Foss.
Antic., p. 24, Hud. Riv. Gr.

obtusum, Hall, 1847, Pal. N. Y., vol. 1, p.
24, Chazy Gr.

pelops, Billings, 1865, Pal. Foss., vol. 1, p.
317, Quebec Gr. or Up. Taconic. Not

an *Asaphus*.

platycephalus, Stokes, 1822, Trans. Geo.
Soc. Lond., 2d ser., vol. 1, p. 258, Tren-
ton Gr.

platypleurus, Green, 1837, Am. Jour. Sci.,
vol. 32, p. 169, Low. Sil. Not very

clearly defined.

pleuropteryx, see *Dalmanites pleuropteryx*.

polepterus, Green, 1838, Am. Jour. Sci.,
vol. 34, Keokuk Gr. Probably a *Phil-
lipia*.

qualiticaudatus, Billings, 1865, Pal. Foss.,
vol. 1, p. 272, Quebec Gr. or Up. Ta-
conic. Not an *Asaphus*.

roningeri, Walcott, 1876, 28th Rep. N. Y.
Mus. Nat. Hist., p. 96, Black Riv. and
Trenton Gr.

selenurus, see *Dalmanites selenurus*.

stokesi, see *Proetus stokesi*.

susae, Calvin, 1882, Geo. Wis., vol. 4, p.
236, Trenton Gr.

tetragonocephalus, Green, 1834, Am. Jour.
Sci., vol. 25, p. 336. Not an *Asaphus*,
and the relations not clear.

trentonensis, see *Lielas trentonensis*.

triangulatus, Whitfield, syn. for *A. ho-
malonotoides*.

trimblii, Green, 1837, Jour. Acad. Nat.
Sci. Phil., vol. 7, Niagara Gr.

vetustus, Hall, 1847, (*Ogygia vetustus*,)
Pal. N. Y., vol. 1, p. 227, Birds-
eye Gr.

vigilans, Meek & Worthen, 1870, (*Isote-
lus vigilans*,) Proc. Acad. Nat. Sci.
Phil., p. 53, and Geo. Sur. Ill., vol. 6,
p. 497, Hud. Riv. Gr.

FIG. 968.—*Asaphus megis-
tus*.

FIG. 967.—*Asaphus*
gigas.

FIG. 966.—*Asaph-*
oldichnus trifid-
us.

Catal. Sil. Foss.

v. Gr.

aspects.

Am. Jour. Sci.,

ly founded upon

cops.

ake Sup. Land

Gr.

858, Can. Jour.

Gr.

al. N. Y., vol. 1,

85, Monogr. U. S.

Quebec Gr.

A. ditmarsie, Honeyman, acc. to Vogdes & (Proc. ~~Acad. Nat. Sci.~~ ^{Nova Scotia} ~~Inst.~~ ^{Inst.}
Nat. Sc., 1888 Vol. 7 Pt. 1) = *A. aculeatus* ~~Green~~

wetherilli, Green, syn. for *Dalmanites limulurus*.

wisconsinensis, Walcott, 1876, 28th Rep. N. Y. Mus. Nat. Hist., p. 97, Trenton Gr. Atrons, Emmons, 1844, Taconic System, p. 64, and Am. Geol. p. 115. [Ety. *a*, absence of; *ops*, an eye.] Cephalic shield semicircular, anterior and lateral edges turned upward, posterior angles rounded, convex; glabella subquadrate, convex, appearing as a continuation of the central lobe, two lateral furrows on each side, neck segment well defined; facial suture beginning at the antero-lateral part of the cephalic shield, runs nearly parallel with the anterior margin to the front of the glabella, when it turns at right angles and runs parallel with the glabella to the posterior margin; no eyes; thoracic segments 17, axial nearly as wide as the lateral lobes, narrowing gradually to the pygidium, armed with a row of short spines, lateral lobes with a row of tubercles on the median line; pygidium small, somewhat semielliptical, flat, axial lobe with a single ring. Type A. *trilineatus*.

fischeri, Billings, 1885, (*Triarthrus fischeri*), Pal. Foss., vol. 1, p. 201, Quebec Gr. or Up. Taconic.

miser, Billings, 1861, (*Conocephalites miser*), Pal. Foss., vol. 1, p. 12, Up. Taconic.

trilineatus, Emmons, 1844, Taconic System, p. 64, and Am. Geol., p. 115, Up. Taconic.

BAILLIÉLLA, Matthew, 1884, Trans. Roy. Soc. Can., vol. 2, pl. 1. [Ety. proper name.] Proposed as a subgenus and founded on *Conocoryphe baileyi*.

BARRANDIA, McCoy, 1849, Ann. Nat. Hist. 2d ser. vol. 4, p. 409. [Ety. proper name.] Ovate, depressed; glabella with incomplete axial furrows and no distinct lobes; eyes large, sub-central; facial

FIG. 909—*Atrons trilineatus*.

suture cutting the posterior margin about the middle, and in front of the eyes arching forward, first outward and then inward; pleurae falcate, with a fulcrum close to the axis, grooved, not faceted; pygidium with short axis and smooth sides. Type B. *cordai*.

(?) *maccoyi*, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 96, Trenton Gr.

Barrandia, Hall, 1860. The name was pre-occupied by McCoy in 1849; beside, it is a syn. for *Elliptocephala*.

thompsoni, see *Elliptocephala thompsoni*, *vermontana*, see *Elliptocephala vermontana*.

BATHYNOTUS, Hall, 1860, 3d Rep. N. Y. St. Mus. Nat. Hist., p. 117. [Ety. *bathus*, ample; *notus*, back.] Cephalic shield somewhat semielliptical, with posterior

FIG. 970—*Bathynotus holopyga*. Long eye-lobes crushed down.

angles produced in very long spines; glabella transversely lobed; eye-lobe narrow, elongate, extending from opposite the antero-lateral angle of the glabella obliquely backward nearly to the posterior margin; facial suture passes nearly around the extended eye-lobe, and cuts the margin before reaching the posterior extension of the eye-lobe; anteriorly it passes in front of the glabella without, as it appears, cutting the front margin; free cheeks united in front; thirteen thoracic segments; middle lobe prominent, twice as wide as the lateral lobes; articulations strong, each bearing a central node; pleurae short, each terminating in a spine, the last pair being prolonged far beyond the pygidium; pygidium short, middle lobe with three annulations, lateral lobes flat and plain; hypostoma having an obtuse angle; at the front margin of the doublure, the latter being cut away to permit the extension to cross it, behind the doublure it is transversely quadrangular. Type B. *holopyga*.

holopyga, Hall, 1859, (*Peltura holopyga*), 12th Rep. N. Y. St. Mus. Nat. Hist., p. 61 and Pal. N. Y., vol. 3, p. 528, Up. Taconic.

the name was proposed in 1840; beside, it is a

phala thompsoni. Cephalo vermont-

and Rep. N. Y. St. 17. [Ety. bathys, Cephalic shield al, with posterior

ga. Long eye-lobes with spines.

very long spines; lobed; eye-lobe extending from the outer angle of the backward nearly to the eye; facial suture the extended eye-margin before reaching the eye; the eye appears, cutting, the cheeks united in the middle segments; mid-ventral node; pleurae long in a spine, the longest far beyond the middle segments; lateral hypostoma having the front margin the latter being the extension to obscure it is trans- Type B. holo-

Peltura holopyga.) Mus. Nat. Hist., vol. 3, p. 528, Up.

BATHYURELLUS, Billings, 1865, Pal. Foss., vol. 1, p. 262. [Ety. diminutive of *Bathyrus*.] Form oblong, ovate; cephalic shield convex, lunate; glabella conical or pointed, without furrows; eyes lunate; facial suture in front of the eye, curving outward, then straight forward or inward on approaching the margin behind the eye, running outward sub-parallel to the neck furrow, and cutting the margin before reaching the outer angle; thorax, nine segments; axis of pygidium short, not strongly grooved, side lobes with short ribs, and a broad, smooth border all around, sometimes concave. Type B. abruptus and B. nitidus.

abruptus, Billings, 1865, Pal. Foss., vol. 1, p. 263, Quebec Gr. or Up. Taconic.

bradleyi, Meek, see *Asaphiscus bradleyi*. expansus, Billings, 1865, Pal. Foss., vol. 1, p. 318, Quebec Gr. or Up. Taconic.

formosus, Billings, 1865, Pal. Foss., vol. 1, p. 266, Quebec Gr. or Up. Taconic.

fraternus, Billings, 1865, Pal. Foss., vol. 1, p. 267, Quebec Gr. or Up. Taconic.

litoreus, Billings, 1865, Pal. Foss., vol. 1, p. 320, Quebec Gr. or Up. Taconic.

marginatus, Billings, 1865, Pal. Foss., vol. 1, p. 264, Quebec Gr. or Up. Taconic.

nitidus, Billings, 1865, Pal. Foss., vol. 1, p. 265, Quebec Gr. or Up. Taconic.

rarus, Billings, 1865, Pal. Foss., vol. 1, p. 320, Quebec Gr. or Up. Taconic.

truncatus, Meek, 1873, Hayden's Geo. Sur. Terr., p. 465. Not satisfactorily defined.

vallidus, Billings, 1865, Pal. Foss., vol. 1, p. 268, Quebec Gr. or Up. Taconic.

wheeleri, Meek, see *Asaphiscus wheeleri*.

BATHYURISCUS, Meek, 1873, 6th Rep. Hayden's U. S. Geo. Sur. Terr., p. 484. [Ety. from *Bathyrus*.] Ovate; head semicircular; glabella straight or slightly expanded in front, three or four pairs of furrows; eyes elongate; facial sutures cut the

anterior margin of the head each side of the greatest expansion of the glabella, and extend obliquely inward to the anterior bases of the eyes; encircling the latter, they extend obliquely outward, cutting the posterior margin so as to leave a narrow, elongate lateral limb; thorax from 7 to 9 segments; axis strong; pleural groove broad; pygidium semicircular, axis strong, and crossed by several furrows which cross the lateral lobes. Type B. haydeni.

FIG. 972.—*Bathyriscus productus*.

FIG. 971.—*Bathyriscus nitidus*.

haydeni, Meek, 1873, (*Bathyriscus haydeni*), 6th Rep. Hayden's U. S. Geo. Sur. Terr., p. 482, Up. Taconic.

howelli, Walcott, 1886, Bull. U. S. Geo. Sur. No. 30, p. 216, Up. Taconic.

productus, Hall & Whitfield, 1877, (*Ogygia producta*), Geol. Expl. 40th Par., vol. 4, p. 244, Up. Taconic.

BATHYURUS, Billings, 1850, Can. Nat. and Geo., vol. 4, p. 365. [Ety. *bathys*, deep; *oura*, tail.] Elliptical, sides straight; cephalic shield lunate, posterior angles produced in spines; glabella subquadrate, rounded anteriorly, convex, furrows obscure, neck segment distinct; eyes large, smooth, semilunar; facial sutures curving forward anteriorly, and posteriorly directed straight backward from the eye, and then, abruptly curving outward, cut the cephalic shield half-way to the genal angle; nine thoracic segments, axial lobe narrower than lateral lobes, and gradually tapering; pleurae furrowed; pygidium smaller than the head, segments closely united, border flattened and smooth; hypostoma not forked. Type B. extans.

amplimarginatus, Billings, 1850, Can. Nat. and Geo., vol. 4, p. 365, Calciferous Gr.

angelini, Billings, 1850, Can. Nat. and Geo., vol. 4, p. 468, Chazy Gr.

arcuatus, Billings, 1865, Pal. Foss., vol. 1, p. 205, Quebec Gr. or Up. Taconic.

armatus, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 321, Quebec Gr. or Up. Taconic.

bituberculatus, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 321, Quebec Gr. or Up. Taconic.

breviceps, Billings, 1865, Pal. Foss., vol. 1, p. 262, Quebec Gr. or Up. Taconic.

capax, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 321, Quebec Gr. or Up. Taconic.

caudatus, Billings, 1865, Pal. Foss., vol. 1, p. 261, Quebec Gr. or Up. Taconic.

coniens, Billings, 1850, Can. Nat. and Geo., vol. 4, p. 366, Calciferous Gr.

(?) congeneris, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 92, Quebec Gr. or Up. Taconic.

cordal, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 321, Calciferous Gr.

cratiformis, Dwight, 1884, Am. Jour. Sci. and Arts, 3d ser., vol. 27, p. 253, Calciferous Gr.

cybele, Billings, 1850, Can. Nat. and Geo., vol. 4, p. 366, Calciferous Gr.

dubius, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 321, Quebec Gr. or Up. Taconic.

extans, Hall, 1847, (*Asaphus* (?) *extans*), Pal. N. Y., vol. 1, p. 228, Lower Trenton Gr.

gregarius, Billings, 1865, Pal. Foss., vol. 1, p. 363, Up. Taconic.

haydeni, see *Bathyriscus haydeni*.

longispinus, Walcott, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 94, Black River and Trenton Grs.

minganensis, Billings, 1865, Pal. Foss., vol. 1, p. 353, Calciferous Gr.
 nero, Billings, 1865, Pal. Foss., vol. 1, p. 260, Quebec Gr. or Up. Taconic.
 oblongus, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 321, Quebec Gr. or Up. Taconic.
 parvulus, Billings, 1861, Pal. Foss., vol. 1, p. 16, Up. Taconic.
 perplexus, Billings, 1865, Pal. Foss., vol. 1, p. 364, Potsdam Gr. or Up. Taconic.
 perspicator, Billings, 1865, Pal. Foss., vol. 1, p. 205, Quebec Gr. or Up. Taconic.
 poronipensis, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 243, Quebec Gr. or Up. Taconic.
 quadratus, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 320, Quebec Gr. or Up. Taconic.
 saffordii, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 321, Quebec Gr. or Up. Taconic.
 seelyi, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 339, Birdseye Gr.
 senectus, Billings, 1861, Pal. Foss., vol. 1, p. 16, Up. Taconic.
 serratus, Meek, 1873, 6th Rep. Hayden's Geo. Sur. Terr., p. 480, Potsdam Gr. or Up. Taconic.
 ?simillimus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 93, Quebec Gr. or Up. Taconic.
 smithi, Billings, 1862, Pal. Foss., vol. 1, p. 56, Black Riv. Gr.
 solitarius, Billings, 1865, Pal. Foss., vol. 1, p. 362, Up. Taconic.
 spiniger, Hall, 1847, (Acidaspis spiniger,) Pal. N. Y., vol. 1, p. 241, Black River and Trenton Gr.
 stonemani, Vogdes, 1884, 12th Rep. Geo. and Nat. Hist. Minn., p. 8, Trenton Gr.
 strenuus, Billings, 1865, Pal. Foss., vol. 1, p. 204, Quebec Gr. or Up. Taconic.
 taurifrons, Dwight, 1884, Am. Jour. Sci. and Arts, 3d ser., vol. 27, p. 252, Calciferous Gr.
 timon, Billings, 1865, Pal. Foss., vol. 1, p. 261, Quebec Gr. or Up. Taconic.
 ?tuberculatus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 91, Quebec Gr. or Up. Taconic.
 vetulus, Billings, 1865, Pal. Foss., vol. 1, p. 365, Potsdam Gr. or Up. Taconic.
BELINURUS, Konig, 1825, Icones Fossilium Sectiles, p. 230. [Ety. *belos*, dart; *oura*, tail.] Cephalo-thoracic shield sub-crescentiform, more than twice as wide as long, lateral angles pointed; ocular ridge surrounds a transversely subelliptical area, within which there is a crown-shaped area, surrounded by a ridge; eyes small and at the lateral extremities of the subelliptical area; mesial lobe narrow, and contracted toward each end; lateral lobes wide, flattened on the margin and serrate on the edge; telson tapering to a point. Type *B. bellulus*.

FIG. 973.—*Bathyrurus smithi*.

danæ, see *Euproops danæ*.
 laccel, Packard, 1885, Am. Naturalist, vol. 19, p. 291, Coal Meas.

FIG. 974.—*Belinurus bellulus*. *e*, Position of eye, at the lateral extremity of a transversely elliptical area.

BEYRICHTIA, McCoy, 1844, Syn. Sil. Foss. Ireland, p. 57. [Ety. proper name.] Carapace equivalve, oblong, extremities rounded, ventral border semicircular, dorsal straight; valves wider at the caudal than the cephalic extremity, more or less convex, impressed with transverse furrows. Type *B. kloedeni*.
aquilatera, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 158, and Acad. Geol., p. 609, Up. Silurian.
americana, Shumard, 1858, (Cythere americana,) Trans. St. Louis Acad. Sci., vol. 1, p. 22, Up. Coal Meas.
areolaris, Rean, 1886, Lond. Geo. Mag., p. 438, Low. Held. Gr.
atlantica, Billings, 1865, Pal. Foss., vol. 1, p. 360, Quebec Gr. or Up. Taconic.
bella, Walcott, 1883, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 213, Trenton Gr.
chambersi, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 234, Hud. River Gr.
ciliata, Emmons, 1855, American Geo., p. 219, Hud. Riv. Gr.
cincinnatiensis, see *Primitia cincinnatiensis*.
clathrata, Jones, 1858, Ann. and Mag. Nat. Hist., 3d series, vol. 1, p. 212, Niagara Gr.
decora, Billings, 1866, Catal. Sil. Foss. Anticosti Gr., p. 67, Anticosti Gr.
dacon, Clarke, 1855, Bull. U. S. Geo. Sur., No. 16, p. 7, Fig. 976.—*Beyrichia duryi*, Genesee yl. Magnified 35 diam. Shale.
duryi, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 232, Hud. Riv. Gr.

FIG. 975.—*Beyrichia chambersi*. Mag. 12 diam.

nae.
m. Naturalist, vol.

FIG. 976.—*Beyrichia duryi*. Interior of left valve, magnified 25 diam.

s. e. Position of eye,
y of a transversely

44, Syn. Sil. Foss.
ty. proper name.]
e, oblong, extremi-
al border semicir-
ant; valves wider at
cephalic extremity,
ex, impressed with
Type B. klødeni.
90, Can. Nat. and
and Acad. Geol., p.

1858, (Cythere amer-
eas Acad. Sci., vol.
Meas.
Lond. Geo. Mag., p.

5, Pal. Foss., vol. 1,
or Up. Taconic.
35th Rep. N. Y. St.
213, Trenton Gr.
nberal, S. A. Miller,
74, Cin. Quar. Jour.
i., vol. 1, p. 234,
ud. River Gr.
ta, Emmons, 1855,
merican Geo., p. 219,
ud. Riv. Gr.
nnatiensis, see Pri-
itia cincinnatensis.
Ann. and Mag. Nat.
ol. 1, p. 212. Niag.

FIG. 978.—*Beyrichia duryi*. Magnified 25 diam.

74, Cin. Quar. Jour.
ud. Riv. Gr.

fœtoidea, White & St. John, 1868, Trans.
Chl. Acad. Sci., p. 126, Up. Coal Meas.
granulosa, Hall, 28th Rep. N. Y. St. Mus.
Nat. Hist., p. 186, Niagara Gr.

FIG. 977.—*Beyrichia duryi*. Interior of left valve, magnified 25 diam.

Hist., 6th ser., vol. 3, p. 379, Low.
Held. Gr.

lata, Vanuxem, 1842, (Agnostus latus.)
Geo. Rep. N. Y., p. 80, and Pal. N. Y.,
vol. 2, p. 301, Clinton Gr.

lithofactor, White & St. John, 1868, Pre-
lim. Notice of New Foss., Coal Meas.
logani, see Primitia logani.

logani var. leperditoides, see Primitia leper-
ditoides.

logani var. reniformis, see Primitia reni-
formis.

maccoyana, Jones, 1855, Ann. and Mag.
Nat. Hist., 2d ser., vol. 16, p. 88, Onon-
daga Gr.

notata, Hall, 1859, Pal. N. Y., vol. 3, p.
379, Low. Held. Gr.

notata var. ventricosa, Hall, 1859, Pal.
N. Y., vol. 3, p. 380, Low. Held. Gr.

novascotia, Jones & Kirby, 1884, Lond.
Geo. Mag., 3d ser., vol. 1, p. 356, Car-
boniferous.

occidentalis, Walcott, 1885, Monogr. U. S.
Sur., vol. 8, p. 204, Devonian.

oculifera, Hall, 1871, 24th Rep. N. Y. St.
Mus. Nat. Hist., p. 232, Hud. Riv. Gr.

oculina, Hall, 1859, Pal. N. Y., vol. 3, p.
378, Low. Held. Gr.

pennsylvanica, Jones, 1858, Ann. and
Mag. Nat. Hist., 3d ser., vol. 1, p. 253,
Onondaga Gr.

persulcata, Ulrich, 1879, Jour. Cin. Soc.
Nat. Hist., vol. 2, p. 12, Hud. Riv. Gr.

petrifactor, White & St. John, 1868,
Trans. Chi. Acad. Sci., p. 125, St.
Louis Gr.

petrifactor var. velata, White & St. John,
1868, Trans. Chi. Acad. Sci., p. 126, St.
Louis Gr.

plagiosa, Jones, 1858, Ann. and Mag. Nat.
Hist., 3d ser., vol. 1, p. 243, Niagara Gr.

punctulifera, Hall, 1862, 15th Rep. N. Y.
St. Mus. Nat. Hist., p. 83, Ham. Gr.

pustulosa, Hall, 1860, Can. Nat. and
Geo., vol. 5, p. 157, and
Acad. Geol., p. 609, Up.

Silurian.

quadrilobata, Hall & Whit-
field, syn. for Beyrichia

regularis.

regularis, Emmons, 1855,
Am. Geo., p. 219, Hud. Riv. Gr.

richardsoni, S. A. Miller, 1874, Cin. Quar.
Jour. Sci., vol. 1, p. 347, Hud. Riv. Gr.

granulata, Hall,
1859, Pal. N. Y.,
vol. 3, p. 377,
Low. Held. Gr.
jonesi, Dawson,
1868, Acad.
Geol., p. 312,
Carboniferous.

klødeni var. aca-
dica, 1880, Ann.
and Mag. Nat.

Hist., 6th ser., vol. 3, p. 379, Low.
Held. Gr.

lata, Vanuxem, 1842, (Agnostus latus.)
Geo. Rep. N. Y., p. 80, and Pal. N. Y.,
vol. 2, p. 301, Clinton Gr.

lithofactor, White & St. John, 1868, Pre-
lim. Notice of New Foss., Coal Meas.

logani, see Primitia logani.

logani var. leperditoides, see Primitia leper-
ditoides.

logani var. reniformis, see Primitia reni-
formis.

maccoyana, Jones, 1855, Ann. and Mag.
Nat. Hist., 2d ser., vol. 16, p. 88, Onon-
daga Gr.

notata, Hall, 1859, Pal. N. Y., vol. 3, p.
379, Low. Held. Gr.

notata var. ventricosa, Hall, 1859, Pal.
N. Y., vol. 3, p. 380, Low. Held. Gr.

novascotia, Jones & Kirby, 1884, Lond.
Geo. Mag., 3d ser., vol. 1, p. 356, Car-
boniferous.

occidentalis, Walcott, 1885, Monogr. U. S.
Sur., vol. 8, p. 204, Devonian.

oculifera, Hall, 1871, 24th Rep. N. Y. St.
Mus. Nat. Hist., p. 232, Hud. Riv. Gr.

oculina, Hall, 1859, Pal. N. Y., vol. 3, p.
378, Low. Held. Gr.

pennsylvanica, Jones, 1858, Ann. and
Mag. Nat. Hist., 3d ser., vol. 1, p. 253,
Onondaga Gr.

persulcata, Ulrich, 1879, Jour. Cin. Soc.
Nat. Hist., vol. 2, p. 12, Hud. Riv. Gr.

petrifactor, White & St. John, 1868,
Trans. Chi. Acad. Sci., p. 125, St.
Louis Gr.

petrifactor var. velata, White & St. John,
1868, Trans. Chi. Acad. Sci., p. 126, St.
Louis Gr.

plagiosa, Jones, 1858, Ann. and Mag. Nat.
Hist., 3d ser., vol. 1, p. 243, Niagara Gr.

punctulifera, Hall, 1862, 15th Rep. N. Y.
St. Mus. Nat. Hist., p. 83, Ham. Gr.

pustulosa, Hall, 1860, Can. Nat. and
Geo., vol. 5, p. 157, and
Acad. Geol., p. 609, Up.

Silurian.

quadrilobata, Hall & Whit-
field, syn. for Beyrichia

regularis.

regularis, Emmons, 1855,
Am. Geo., p. 219, Hud. Riv. Gr.

richardsoni, S. A. Miller, 1874, Cin. Quar.
Jour. Sci., vol. 1, p. 347, Hud. Riv. Gr.

granulata, Hall,
1859, Pal. N. Y.,
vol. 3, p. 377,
Low. Held. Gr.

jonesi, Dawson,
1868, Acad.
Geol., p. 312,
Carboniferous.

klødeni var. aca-
dica, 1880, Ann.
and Mag. Nat.

Hist., 6th ser., vol. 3, p. 379, Low.
Held. Gr.

lata, Vanuxem, 1842, (Agnostus latus.)
Geo. Rep. N. Y., p. 80, and Pal. N. Y.,
vol. 2, p. 301, Clinton Gr.

lithofactor, White & St. John, 1868, Pre-
lim. Notice of New Foss., Coal Meas.

logani, see Primitia logani.

logani var. leperditoides, see Primitia leper-
ditoides.

logani var. reniformis, see Primitia reni-
formis.

maccoyana, Jones, 1855, Ann. and Mag.
Nat. Hist., 2d ser., vol. 16, p. 88, Onon-
daga Gr.

notata, Hall, 1859, Pal. N. Y., vol. 3, p.
379, Low. Held. Gr.

notata var. ventricosa, Hall, 1859, Pal.
N. Y., vol. 3, p. 380, Low. Held. Gr.

novascotia, Jones & Kirby, 1884, Lond.
Geo. Mag., 3d ser., vol. 1, p. 356, Car-
boniferous.

occidentalis, Walcott, 1885, Monogr. U. S.
Sur., vol. 8, p. 204, Devonian.

oculifera, Hall, 1871, 24th Rep. N. Y. St.
Mus. Nat. Hist., p. 232, Hud. Riv. Gr.

oculina, Hall, 1859, Pal. N. Y., vol. 3, p.
378, Low. Held. Gr.

pennsylvanica, Jones, 1858, Ann. and
Mag. Nat. Hist., 3d ser., vol. 1, p. 253,
Onondaga Gr.

persulcata, Ulrich, 1879, Jour. Cin. Soc.
Nat. Hist., vol. 2, p. 12, Hud. Riv. Gr.

petrifactor, White & St. John, 1868,
Trans. Chi. Acad. Sci., p. 125, St.
Louis Gr.

petrifactor var. velata, White & St. John,
1868, Trans. Chi. Acad. Sci., p. 126, St.
Louis Gr.

plagiosa, Jones, 1858, Ann. and Mag. Nat.
Hist., 3d ser., vol. 1, p. 243, Niagara Gr.

punctulifera, Hall, 1862, 15th Rep. N. Y.
St. Mus. Nat. Hist., p. 83, Ham. Gr.

pustulosa, Hall, 1860, Can. Nat. and
Geo., vol. 5, p. 157, and
Acad. Geol., p. 609, Up.

Silurian.

quadrilobata, Hall & Whit-
field, syn. for Beyrichia

regularis.

regularis, Emmons, 1855,
Am. Geo., p. 219, Hud. Riv. Gr.

richardsoni, S. A. Miller, 1874, Cin. Quar.
Jour. Sci., vol. 1, p. 347, Hud. Riv. Gr.

granulata, Hall,
1859, Pal. N. Y.,
vol. 3, p. 377,
Low. Held. Gr.

jonesi, Dawson,
1868, Acad.
Geol., p. 312,
Carboniferous.

klødeni var. aca-
dica, 1880, Ann.
and Mag. Nat.

Hist., 6th ser., vol. 3, p. 379, Low.
Held. Gr.

lata, Vanuxem, 1842, (Agnostus latus.)
Geo. Rep. N. Y., p. 80, and Pal. N. Y.,
vol. 2, p. 301, Clinton Gr.

lithofactor, White & St. John, 1868, Pre-
lim. Notice of New Foss., Coal Meas.

logani, see Primitia logani.

logani var. leperditoides, see Primitia leper-
ditoides.

logani var. reniformis, see Primitia reni-
formis.

maccoyana, Jones, 1855, Ann. and Mag.
Nat. Hist., 2d ser., vol. 16, p. 88, Onon-
daga Gr.

notata, Hall, 1859, Pal. N. Y., vol. 3, p.
379, Low. Held. Gr.

notata var. ventricosa, Hall, 1859, Pal.
N. Y., vol. 3, p. 380, Low. Held. Gr.

novascotia, Jones & Kirby, 1884, Lond.
Geo. Mag., 3d ser., vol. 1, p. 356, Car-
boniferous.

occidentalis, Walcott, 1885, Monogr. U. S.
Sur., vol. 8, p. 204, Devonian.

oculifera, Hall, 1871, 24th Rep. N. Y. St.
Mus. Nat. Hist., p. 232, Hud. Riv. Gr.

oculina, Hall, 1859, Pal. N. Y., vol. 3, p.
378, Low. Held. Gr.

pennsylvanica, Jones, 1858, Ann. and
Mag. Nat. Hist., 3d ser., vol. 1, p. 253,
Onondaga Gr.

persulcata, Ulrich, 1879, Jour. Cin. Soc.
Nat. Hist., vol. 2, p. 12, Hud. Riv. Gr.

petrifactor, White & St. John, 1868,
Trans. Chi. Acad. Sci., p. 125, St.
Louis Gr.

petrifactor var. velata, White & St. John,
1868, Trans. Chi. Acad. Sci., p. 126, St.
Louis Gr.

plagiosa, Jones, 1858, Ann. and Mag. Nat.
Hist., 3d ser., vol. 1, p. 243, Niagara Gr.

punctulifera, Hall, 1862, 15th Rep. N. Y.
St. Mus. Nat. Hist., p. 83, Ham. Gr.

pustulosa, Hall, 1860, Can. Nat. and
Geo., vol. 5, p. 157, and
Acad. Geol., p. 609, Up.

Silurian.

quadrilobata, Hall & Whit-
field, syn. for Beyrichia

regularis.

regularis, Emmons, 1855,
Am. Geo., p. 219, Hud. Riv. Gr.

richardsoni, S. A. Miller, 1874, Cin. Quar.
Jour. Sci., vol. 1, p. 347, Hud. Riv. Gr.

granulata, Hall,
1859, Pal. N. Y.,
vol. 3, p. 377,
Low. Held. Gr.

jonesi, Dawson,
1868, Acad.
Geol., p. 312,
Carboniferous.

klødeni var. aca-
dica, 1880, Ann.
and Mag. Nat.

Hist., 6th ser., vol. 3, p. 379, Low.
Held. Gr.

lata, Vanuxem, 1842, (Agnostus latus.)
Geo. Rep. N. Y., p. 80, and Pal. N. Y.,
vol. 2, p. 301, Clinton Gr.

lithofactor, White & St. John, 1868, Pre-
lim. Notice of New Foss., Coal Meas.

logani, see Primitia logani.

logani var. leperditoides, see Primitia leper-
ditoides.

logani var. reniformis, see Primitia reni-
formis.

maccoyana, Jones, 1855, Ann. and Mag.
Nat. Hist., 2d ser., vol. 16, p. 88, Onon-
daga Gr.

notata, Hall, 1859, Pal. N. Y., vol. 3, p.
379, Low. Held. Gr.

notata var. ventricosa, Hall, 1859, Pal.
N. Y., vol. 3, p. 380, Low. Held. Gr.

novascotia, Jones & Kirby, 1884, Lond.
Geo. Mag., 3d ser., vol. 1, p. 356, Car-
boniferous.

occidentalis, Walcott, 1885, Monogr. U. S.
Sur., vol. 8, p. 204, Devonian.

oculifera, Hall, 1871, 24th Rep. N. Y. St.
Mus. Nat. Hist., p. 232, Hud. Riv. Gr.

oculina, Hall, 1859, Pal. N. Y., vol. 3, p.
378, Low. Held. Gr.

pennsylvanica, Jones, 1858, Ann. and
Mag. Nat. Hist., 3d ser., vol. 1, p. 253,
Onondaga Gr.

persulcata, Ulrich, 1879, Jour. Cin. Soc.
Nat. Hist., vol. 2, p. 12, Hud. Riv. Gr.

petrifactor, White & St. John, 1868,
Trans. Chi. Acad. Sci., p. 125, St.
Louis Gr.

petrifactor var. velata, White & St. John,
1868, Trans. Chi. Acad. Sci., p. 126, St.
Louis Gr.

plagiosa, Jones, 1858, Ann. and Mag. Nat.
Hist., 3d ser., vol. 1, p. 243, Niagara Gr.

punctulifera, Hall, 1862, 15th Rep. N. Y.
St. Mus. Nat. Hist., p. 83, Ham. Gr.

pustulosa, Hall, 1860, Can. Nat. and
Geo., vol. 5, p. 157, and
Acad. Geol., p. 609, Up.

Silurian.

quadrilobata, Hall & Whit-
field, syn. for Beyrichia

regularis.

regularis, Emmons, 1855,
Am. Geo., p. 219, Hud. Riv. Gr.

richardsoni, S. A. Miller, 1874, Cin. Quar.
Jour. Sci., vol. 1, p. 347, Hud. Riv. Gr.

granulata, Hall,
1859, Pal. N. Y.,
vol. 3, p. 377,
Low. Held. Gr.

jonesi, Dawson,
1868, Acad.
Geol., p. 312,
Carboniferous.

klødeni var. aca-
dica, 1880, Ann.
and Mag. Nat.

Hist., 6th ser., vol. 3, p. 379, Low.
Held. Gr.

lata, Vanuxem, 1842, (Agnostus latus.)
Geo. Rep. N. Y., p. 80, and Pal. N. Y.,
vol. 2, p. 301, Clinton Gr.

lithofactor, White & St. John, 1868, Pre-
lim. Notice of New Foss., Coal Meas.

logani, see Primitia logani.

logani var. leperditoides, see Primitia leper-
ditoides.

logani var. reniformis, see Primitia reni-
formis.

maccoyana, Jones, 1855, Ann. and Mag.
Nat. Hist., 2d ser., vol. 16, p. 88, Onon-
daga Gr.

notata, Hall, 1859, Pal. N. Y., vol. 3, p.
379, Low. Held. Gr.

notata var. ventricosa, Hall, 1859, Pal.
N. Y., vol. 3, p. 380, Low. Held. Gr.

novascotia, Jones & Kirby, 1884, Lond.
Geo. Mag., 3d ser., vol. 1, p. 356, Car-
boniferous.

occidentalis, Walcott, 1885, Monogr. U. S.
Sur., vol. 8, p. 204, Devonian.

oculifera, Hall, 1871, 24th Rep. N. Y. St.
Mus. Nat. Hist., p. 232, Hud. Riv. Gr.

oculina, Hall, 1859, Pal. N. Y., vol. 3, p.
378, Low. Held. Gr.

pennsylvanica, Jones, 1858, Ann. and
Mag. Nat. Hist., 3d ser., vol. 1, p. 253,
Onondaga Gr.

persulcata, Ulrich, 1879, Jour. Cin. Soc.
Nat. Hist., vol. 2, p. 1

- flabellifer, Goldfuss, Nova. Acta. Acad. Caes. Leop. Nat. Cur., vol. 16, p. 360, Up. Silurian.
- insularia, Billings, 1866, Catal. Sil. Foss. Antic., p. 66, Auticosti Gr.
- laphani, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 88, and Geo. Wis., vol. 4, p. 310, Niagara Gr.
- lunatus, Billings, 1857, Rep. of Progr. Geo. Sur. Can., p. 338, Trenton Gr.
- niagarensis, Hall, 1852, Pal. N. Y., vol. 2, p. 314, Niagara Gr.
- occusus, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., vol. 1, p. 104, Niagara Gr.
- pompilius, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 123, Low. Held. Gr.
- tullius, Hall, 1888, Pal. N. Y., vol. 7, p. 12, Ham. Gr.
- Bumastus*, Murchison, 1839, Sil. Syst. Not American, though I have illustrated the genus because so many have referred *Illænus* to it.

FIG. 381.—*Bumastus barriensis*.

- barriensis*, see *Illænus ioxus*.
- trentonensis*, see *Illænus trentonensis*.
- BUNODELIA**, Matthew, 1888, Trans. Roy. Soc. Can., p. 56. Body ovate-elongate, trilobed longitudinally; cephalic shield subtriangular, with rounded angles; composed of a glabella, fixed cheeks and movable (?) cheeks; glabella broadly cylindrical and rounded in front; fixed cheeks expanded in front, and having ear-shaped lateral lobes defined by an impressed line which may have been movable; thorax, so far as known, had seven segments, and consisted of an elongate cylindro-conical body, having triangular lappets or pleurae attached at the sides. Type B. *horrida*.
- horrida*, Matthew, 1888, Trans. Roy. Soc. Can., p. 56, Up. Silurian or Low. Devonian.
- CALYMENE**, Brongniart, 1822, Hist. Nat. Crust. Foss., p. 7. [Ety. *kekalymenos*, concealed.] Cephalic shield sublunate, margin thickened, distinctly defined; glabella convex, narrower in front than behind, three lateral furrows on each side, the posterior one deep, neck segment well defined, eyes, small, prominent, hiant, near the glabella furrows, and slightly anterior to the middle; facial sutures cut the margin, in front of the eyes and curving slightly over each eye, defining a semicircular

- eye-lobe, they extend to the lateral angles, each of which is exactly bisected; anteriorly they are connected by a rostral suture, thorax of thirteen segments, axis most convex, lateral lobes wider than axis, bent down with large facets; pygidium semi-oval, axis prominent, seven to eleven segments, margin entire. Type C. *blumenbachi*.
- anchiops*, see *Dalmanites anchiops*.
- becki*, see *Triarthrus becki*.
- blumenbachi*, Brongniart, 1822, Hist. Nat. Crust. Foss., p. 11. American form called *C. niagarensis*.
- bucklandi*, syn. for *Ceraurus pleurexanthemus*.
- bufo*, see *Phacops bufo*.
- callicephala*, Green, 1832, Monograph Trilobites, p. 30, and Pal. N. Y., vol. 1, p. 238, Trenton and Hud. Riv. Grs.
- canerata*, Conrad, 1842, Jour. Acad. Nat. Sci., vol. 8, p. 278, and Pal. N. Y., vol. 2, p. 337, Coralline limestone.
- christyi*, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 119, Hud. Riv. Gr.
- clintoni*, Vanuxem, (Hemicypturus clintoni,) Geo. Rep. 3d Dist. N. Y., p. 479, Clinton Gr.
- conradi*, Emmons, 1856, Am. Geol., p. 236, Hud. Riv. Gr.
- crassimarginata*, see *Proetus crassimarginatus*.
- mammillata*, Hall, 1861, Geo. Rep. Wis., p. 50, Trenton Gr.
- marginalis*, see *Proetus marginalis*.
- multicosta*, Hall, 1847, Pal. N. Y., vol. 1, p. 228, Birdseye and Trenton Gr.
- nasuta*, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 131, Niagara Gr.

FIG. 382.—*Calymene callicephala*. Rolled specimen and the under side of cephalic shield with hyposoma in place.

- niagarensis*, Hall, 1843, Geo. Rep. 4th Dist. N. Y., p. 102, and Pal. N. Y., vol. 2, p. 307, Niagara Gr. This is the American variety of *C. blumenbachi*.
- nupera*, see *Phacops nupera*.
- odontocephala*, syn. for *Dalmanites selenurus*.
- phlyctainodes*, see *Encrinurus phlyctainodes*.
- platys*, Green, 1832, Monograph of Trilobites, p. 32, and Illust. Devon. Foss., pl. 1, Schoharie grit.
- rostrata*, Vogdes, 1880, Proc. Acad. Nat. Sci., p. 176, Clinton Gr.
- rowii*, see *Proetus rowii*.
- rugosa*, Shumard, 1855, Geo. Rep. Mo., p. 200, Low. Held. Gr.

to the lateral an-
s exactly bisected;
connected by a
x of thirteen seg-
nux, lateral lobes
t down with large
semi-oval, axin
eleven segments,
e C. blumenbachi.
s anchiops.

cki.
t, 1822, Hist. Nat.
American form

raurus pleurexan-

1832, Monograph
Pal. N. Y., vol. 1,
Hud. Riv. Grs.

2, Jour. Acad. Nat.
and Pal. N. Y., vol.
limestone.

3th Rep. N. Y. St.
19, Hud. Riv. Gr.
semicrypturus clin-
Dist. N. Y., p. 479,

6, Am. Geol., p. 236,

oetus crassimargin-

31, Geo. Rep. Wis.,

s marginalis.

Pal. N. Y., vol. 1,

Trenton Gr.

Jour. Cin. Soc. Nat.

Niagara Gr.

phala. Rolled spec-
e of cephalic shield

43, Geo. Rep. 4th

and Pal. N. Y., vol.

Gr. This is the

f C. blumenbachi.

pera.

Dalmanites selen-

crinurus phlyctai-

Monograph of Trilo-

lust. Devon. Foss.,

Proc. Acad. Nat.

Gr.

ii.

5, Geo. Rep. Mo., p.

senaria, Conrad, 1841, syn. for C. calli-
cephala.

spinifera, not defined.

trilocata, Hall, 1843, Geo. Rep. 4th Dist.
N. Y., p. 74, Clinton Gr.

CANDONA, Baird, 1845, Trans. Berw. Nat.
Club, vol. 2, p. 152. A living genus,

and probably not Palæozoic. Like
Cypris, except the lower antennæ pos-

sess no tuft of setæ, and the second
pair of jaws are destitute of a branchial

appendage. The shell is also usually
longer and narrower. Type C. lucena.

(?) elongata, Jones & Kirby, 1884, Lond.
Geo. Mag., 3d ser., vol. 1, p. 356, Car-
boniferous.

CERATOCARIS, McCoy, 1849, Ann. and Mag.
Nat. Hist., 2d ser., vol. 4, p. 412. [Ety.

keration, pod; karis, shrimp.] Cara-
pace bivalve, dorsal line angulated

simplex, Clarke, 1885, Bull. U. S. Geo.

Sur., No. 16, p. 43, Ham. Gr.

sinuata, Meek & Worthen, 1868, Am. Jour.

Sci., vol. 46, p. 22, and Geo. Sur. Ill.,

vol. 3, p. 540, Coal Meas.

strigata, see Solenocaris strigata.

Ceratocephala, Warder, not defined so as to

be recognized.

ceralepta, Anthony, a fragment of the
tail of a Ceraurus pleurexanthemus, or

of an Acidaspis.

goniata, Warder, a fragment of a Dal-
manites, or an Acidaspis.

CERAURUS, Green, 1832, Monograph Trilo-
bites, p. 84. [Ety. keras, horn; oura,

tail.] Cephalic shield crescentiform,
trilobed, posterior angles extended into

spines; glabella subquadrate, rounded
and prominent in front, three lateral

furrows on each side; eyes faceted

minutely; facial
sutures, commenc-

ing at the anterior
margin, passing

close to the anterior
corners of the gla-

bella curve around
the eyes, from

which points they
extend outward,

and then deflect a
little backward, and

cut the lateral
margins forward of

the neck furrow, if
it were extended;

cheeks generally
scrobiculate; thorax

of ten or twelve
segments, axial lobe

narrower than the

lateral lobes; pleurae flattened for a

distance, and then curve downward

and backward; pygidium small, seg-
ments terminating in digitations or

spines; labrum oblong truncate, with
a pair of furrows and small lateral

auricles. Type C. pleurexanthemus.

(?) apollo, Billings, 1860, (Cheirurus
apollo,) Can. Nat. and Geol., vol. 5,

p. 67, Quebec Gr. or Up. Taconic.

bimucronatus, see Ceraurus niagarensis.

crosoth, see Acidaspis crosoth.

(?) eryx, Billings, 1860, (Cheirurus eryx,)
Can. Nat. and Geol., vol. 5, p. 67, Que-

bec Gr. or Up. Taconic.

(?) glaucus, Billings, 1865, (Cheirurus
glaucus,) Pal. Foss., vol. 1, p. 323, Que-

bec Gr. or Up. Taconic.

icarus, Billings, 1860, (Cheirurus icarus,)
Can. Nat. and Geol., vol. 5, p. 67, Hud.

Riv. Gr.

insignis, see Ceraurus niagarensis.

meekanus, n. sp., Hud. Riv. Gr. Pro-

posed instead of C. icarus, Meek, in

Ohio Pal., vol. 1, p. 162, and plate 14,

figs. a, b, and c. Meek referred this

form to C. icarus of Billings, but it is

distinguished by the form of the gla-

FIG. 983.—Ceratlocaris sinuata. Outline.

with a slight furrow beneath it on
each side; sides semielliptical, much

elongated from before backward, evenly
convex, ventral margin gently convex,

posterior end truncated obliquely; on
each side near the anterior end, low

down, is an ocular spot; surface
marked with fine, imbricating striae.

Type C. solenoides.

aculeata, Hall, 1859, Pal. N. Y., vol. 3,
p. 422, Waterlime Gr.

acuminata, Hall, 1859, Pal. N. Y., vol. 3,
p. 422, Waterlime Gr.

armata, syn. for Echinocaris punctata.

bradleyi, see Colpocaris bradleyi.

becheri, Clarke, 1885, Bull. U. S. Geo.
Sur., No. 16, p. 44, Ham. Gr.

deweyi, Hall, 1859, (Onchus deweyi,) Pal.
N. Y., vol. 2, p. 320, Niagara Gr.

elytroides, see Colpocaris elytroides.

grandis, Pohlman, 1881, Bull. Buf. Soc.
Nat. Hist., vol. 4, p. 19, Waterlime Gr.

longicauda, see Echinocaris longicauda.

maccoyana, Hall, 1859, Pal. N. Y., vol.
3, p. 421, Waterlime Gr.

punctata, see Echinocaris punctata.

pusillus, Matthew, 1889, Trans. Roy. Soc.
Can., vol. 6, p. 49, Low. Held. Gr.

bella, by the furrows, structure of the thorax and form of central lobe, and by the pygidium, beside occurring in higher rocks and growing to a much larger size.

- (?) *mercurius*, Billings, 1865, (*Cheirurus mercurius*,) Pal. Foss., vol. 1, p. 285, Quebec Gr. or Up. Taconic.
niagarensis, Hall, 1867, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 427, Niagara Gr.
numitor, Billings, 1866, (*Cheirurus numitor*,) Catal. Sil. Foss. Antic., p. 27, Hud. Riv. Gr.
nuperus, Billings, 1866, (*Cheirurus nuperus*,) Catal. Sil. Foss. Antic., p. 60, Anticosti Gr.
 (?) *perforator*, Billings, 1865, (*Cheirurus perforator*,) Pal. Foss., vol. 1, p. 287, Quebec Gr. or Up. Taconic.

FIG. 984. — *Ceraurus pleurexanthemus*.

- pleurexanthemus*, Green, 1832, Monog. Trilobites, p. 84, and Pal. N. Y., vol. 1, p. 242, Trenton and Hud. Riv. Gr.
 (?) *polydorus*, Billings, 1865, (*Cheirurus polydorus*,) Pal. Foss., vol. 1, p. 286, Quebec Gr. or Up. Taconic.
pompilius, Billings, 1865, (*Cheirurus pompilius*,) Pal. Foss., vol. 1, p. 181, Chazy or Black Riv. Gr.
 (?) *prolificus*, Billings, 1865, (*Cheirurus prolificus*,) Pal. Foss., vol. 1, p. 285 and 325, Quebec Gr. or Up. Taconic.
pustulosus, syn. for *Ceraurus pleurexanthemus*.
rarus, Walcott, 1877, 31st Rep. N. Y. St. Mus. Nat. Hist., p. 68, Trenton Gr.
satyrus, Billings, 1865, (*Cheirurus satyrus*,) Pal. Foss., vol. 1, p. 324, Chazy Gr.
 (?) *sol*, Billings, 1865, (*Cheirurus sol*,) Pal. Foss., vol. 1, p. 288, Quebec Gr. or Up. Taconic.
 (?) *solitarius*, Billings, 1865, (*Cheirurus solitarius*,) Pal. Foss., vol. 1, p. 206, Quebec Gr. or Up. Taconic.
tarquinius, Billings, 1863, (*Cheirurus tarquinius*,) Proc. Port. Soc. Nat. Hist., vol. 1, p. 121, Upper Silurian.
vigilans, see *Encriurus vigilans*.
 (?) *vulcanus*, Billings, 1865, (*Cheirurus vulcanus*,) Pal. Foss., vol. 1, p. 284, and 324, Quebec Gr. or Up. Taconic.

CHARIOCEPHALUS, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 175. [Ety. *charis*, charming or graceful; *kephale*, head.] Cephalic shield broad; cheeks moderately convex toward the eyes; glabella regularly convex and marked

by transverse furrows; eyes large, facial sutures cutting the contour of the front at or near the center as in *Agraulus*, but distinguished by the character of the palpebral lobe, large eye, and form of the cheek. Type *C. whitfieldi*.

- tumifrons*, Hall & Whitfield, 1877, U. S. Geo. Expl., 40th parallel, vol. 4, p. 224, Potsdam Gr.
whitfieldi, Hall, 1863, 10th Rep. N. Y. St. Mus. Nat. Hist., p. 175, Potsdam Gr.
Cheirurus, Beyrich, 1845, syn. for *Ceraurus*.
apollo, see *Ceraurus apollo*.
eryx, see *Ceraurus eryx*.
glaucus, see *Ceraurus glaucus*.
icarus, see *Ceraurus icarus*.
mercurius, see *Ceraurus mercurius*.
numitor, see *Ceraurus numitor*.
nuperus, see *Ceraurus nuperus*.
perforator, see *Ceraurus perforator*.
polydorus, see *Ceraurus polydorus*.
pompilius, see *Ceraurus pompilius*.
prolificus, see *Ceraurus prolificus*.
satyrus, see *Ceraurus satyrus*.
sol, see *Ceraurus sol*.
solitarius, see *Ceraurus solitarius*.
tarquinius, see *Ceraurus tarquinius*.
vulcanus, see *Ceraurus vulcanus*.

FIG. 985. — *Climachitichnites wilsoni*.

CLIMACHITICHNITES, Logan, 1860, Can. Nat. and Geol., vol. 5, p. 279. [Ety. *klimax*, ladder; *ichnos*, footstep.] A continuous

geminispinosa, Hartt, 1868, (*Conocephalites geminispinosus*.) Acad. Geol., p. 653, St. John Gr.

FIG. 987.—*Conocoryphe kingi*.

Can., p. 119, St. John Gr.
Coronura, Hall, 1888, Pal. N. Y., vol. 7, p. 32. Founded upon the variation in the spines of the pygidium of *Dalmanites*.
Corycephalus, Hall, 1888, syn. for *Dalmanites*.

CREPICEPHALUS, Owen, 1852, Geo. Sur. Wis. Iowa, and Minn., p. 876. [Ety. *krepsis*, horseshoe; *kephale*, head.] Glabella rather flat, slipper-shaped, tapering and slightly acuminate anteriorly, with a faint ridge in the median line; two small depressions, and a posterior furrow partially divide the glabella; facial sutures run nearly parallel to the margin of the glabella, and join a thickened, cord-like, anterior, narrow border, inclosing a convex area, narrower in front than at the sides; pygidium large; axial lobe has four segments, side-lobes bounded by a slightly concave border, which widens posteriorly, and terminates in long spines, and of which the confines are almost rectangular, with rounded corners. Type *C. iowensis*.

angulatus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 220, Potsdam Gr.

anytus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th parallel, vol. 4, p. 219, Potsdam Gr.

angusta, Walcott, 1886, Bull. U. S. Geo. Sur., No. 30, p. 208, Up. Taconic.

centralis, Whitfield, 1877, Rep. on the Pal. of Black Hills, p. 10, and Geo. Black Hills, p. 341, Potsdam Gr.

diadematus, Hall, 1863, (*Conocephalites diadematus*.) 16th Rep. N. Y. St. Mus. Nat. Hist., p. 167, Potsdam Gr.

eos, Hall, 1863, (*Conocephalites eos*.) 16th Rep. N. Y. St. Mus. Nat. Hist., p. 151, Potsdam Gr.

gibbesi, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 50, Potsdam Gr.

granulosus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 214, Potsdam Gr.

kingi, Meek, 1870, Proc. Acad. Nat. Sci. Phil., vol. 22, p. 63, and Rep. on 40th Parallel, p. 20, Up. Taconic.

quadrans, Hall & Whitfield, 1877, (*Crepicephalus quadrans*.) Geo. 40th Parallel, vol. 4, p. 238, Up. Taconic.

walcotti, Matthew, 1884, Trans. Roy. Soc.

haguei, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 210, Potsdam Gr.

iowensis, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 576, Potsdam Gr.

FIG. 988.—*Crepicephalus lilianus*. Cephalic shield without the movable cheeks.

FIG. 989.—*Crepicephalus lilianus*. Pygidium.

lilianus, Walcott, 1886, Bull. U. S. Geo. Sur., vol. No. 30, p. 207, Up. Taconic.

maculosus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 215, Potsdam Gr.

miniscensis, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., pl. 1, fig. 14, Potsdam Gr.

montanensis, Whitfield, 1876, Rep. Recon. Up. Mo. to Yel. Nat. Park, p. 141, Potsdam Gr.

nitidus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 212, Potsdam Gr.

onustus, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., Potsdam Gr.

oweni, Meek & Hayden, 1861, (*Arionellus* (*Crepicephalus*) *oweni*.) Proc. Acad. Nat. Sci., vol. 13, p. 436, Potsdam Gr.

planus, Whitfield, 1877, Rep. on Pal. of Black Hills, p. 11, and Geol. Black Hills, p. 343, Potsdam Gr.

quadrans, see *Conocoryphe quadrans*.
simulator, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 218, Potsdam Gr.

unisulcatus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 216, Potsdam Gr.

vulcanus, Billings, 1861, (*Conocephalites vulcanus*.) Pal. Foss., vol. 1, p. 14, Potsdam Gr.

wisconsinensis, see *Lonchocephalus wisconsinensis*.

Crypheus, Green, 1837, Jour. Acad. Nat. Sci., vol. 7, syn. for *Dalmanites*. Not well defined, and the name was preoccupied for a genus of Coleoptera in 1835.

boothi, see *Dalmanites boothi*.

calliteles, see *Dalmanites calliteles*.

greeni, syn. for *Dalmanites calliteles*.

Cryptolilius, syn. for *Trinucleus*.

tesselatus, see *Trinucleus concentricus*.

Cybele, Loven, 1845, p. 110, Ofversigt Vetensk. Acad. Handl., p. 110.

punctata, Hall, 1852. This species belongs to the genus *Encrinurus*, and the specific name being preoccupied, the name is changed to *E. ornatus*.

1877, U. S. Geo.
4, p. 210, Pots-

Geo. Sur. Wis.,
6, Potsdam Gr.

989.—*Crepheopha-*
willanus. Pygid-

Bull. U. S. Geo.
7, Up. Taconic.
field, 1877, U. S.
el, vol. 4, p. 215,

Geo. Sur. Wis.,
1, fig. 14, Pots-

1878, Rep. Re-
Nat. Park, p. 141,

field, 1877, U. S.
el, vol. 4, p. 212,

Ann. Rep. Geo.

1861, (Arionellus)
) Proc. Acad. Nat.
Potsdam Gr.

Rep. on Pal. of
and Geol. Black
Gr.

the quadrans.
field, 1877, U. S.
el, vol. 4, p. 218,

field, 1877, U. S.
el, vol. 4, p. 216,

(Conocephalites
vol. 1, p. 14, Pots-

chocephalus wis-

our. Acad. Nat.
Dalmanites. Not
name was preoc-
of Coleoptera in

othi.
calliteles.
es calliteles.
ucleus.

concentricus.
110, Oversigt
l., p. 110.

is species belongs
urus, and the spec-
occupied, the name
tus.

Cyclus americanus, Packard, 1885, Am. Nat.,
vol. 19, p. 293, Coal Meas. Not de-
fined so as to be recognized.

CYPHASIS, Burmeister, 1843, Die Organ der
Trilobiten, p. 103. [Ety. *cyphos*, con-
vex; *asis*, shield.] Cephalic shield
semicircular, posterior angles produced
in spines, margin thickened; glabella
very convex, ovoid, no furrows, but
with two small pyriform basal lobes
bounded by deep furrows; eyes small,
semilunate; cheeks broad; facial su-
tures proceed in a nearly straight line,
from the anterior margin to the eyes, and
are then directed to the posterior an-
gles; thorax 10 to 17 segments, rounded
at their extremities; axis tapering;
pygidium small, axis short, lateral lobes
depressed. Type *C. ceratophthalmus*.

brevimarginatus, Walcott, 1885, Monogr.

FIG. 990.—*Cyphas-*
pis christyi.

girardeauensis, Shumard, 1855, Geo. Rep.
Mo., p. 197, Trenton Gr.

hybrida, Hall, 1888, Pal.
N. Y., vol. 7, p. 144,

Up. Held. Gr.
laevis, Hall, 1876, (Phil-
lipsia *laevis*.) Illust.

Devon. Foss., pl. 21,
Chemung Gr.

minuscula, Hall, 1876,
(*Phillipsia minus-*

cula.) Illust. Devon.
Foss., pl. 20, Up. Held.

Gr.
ornata, Hall, 1876,

(*Phillipsia ornata*), FIG. 991.—*Cyphas-*
Illust. Devon. Foss., *pis girardeau-*
pl. 21, Ham. Gr. *ensis*.

ornata var. *baccata*, Hall, 1888, Pal. N. Y.,
vol. 7, p. 146, Ham. Gr.

stephanophora, Hall, 1888, Pal. N. Y.,
vol. 7, p. 142, Up. Held. Gr.

Cythere, Muller, 1785, Entomotraca sue In-
secta, etc., p. 63. The type is *C. fla-*
vida a living species. The genus is un-
known in Paleozoic rocks.

americana, see *Beyrichia americana*.

carbonaria, see *Leperditia carbonaria*.

cincinnatiensis, see *Cytheropsis cincinnati-*
ensis.

crassimarginata, see *Cytheropsis crassimar-*
ginata.

irregularis, see *Cytheropsis irregularis*.

nebraskensis, see *Cytheropsis nebraskensis*.

okeni, see *Leperditia okeni*.

simplex, see *Cytheropsis simplex*.

sublaevis, see *Leperditia sublaevis*.

subrecta, see *Leperditia subrecta*.

CYTHARELLA, inflata.

I find in the
Acadian Geol-
ogy, p. 206, a
small Entomos-
traca, from the FIG. 992.—*Cytherella in-*
Coal Meas. of *flata*.
Nova Scotia, fig-
ured under this name, but without
any description or reference to any
other author. The figures are repro-
duced.

Cytherella glandella, see *Cytheropsis gland-*
della.

Cytherina, Lamarck, 1818, Anim. sans Vert.
t. v, p. 125. [Ety. diminutive of
Cythere.] A synonym for *Cythere*,
which is not a Paleozoic genus.

alta, see *Leperditia alta*.

crenulata, see *Cytheropsis crenulata*.

cylindrica, see *Isochilina cylindrica*.

fabulites, see *Leperditia fabulites*.

spinosa, see *Beyrichia spinosa*. Not Reuss
in 1844.

subcylindrica, see *Cytheropsis subcylin-*
drica.

subelliptica, see *Cytheropsis subelliptica*.

CYTHEROPSIS, McCoy, 1849, Ann. and Mag.

Nat. Hist., 2d. ser., vol. 4, p. 249. [Ety.
Cytheropsis, resembling *Cythere*.] Dis-

tinguished from *Cythere*, which now
swarm in the sea, by the great thick-
ness of the valves, and in having eye
or muscle spots. Type *C. aldensis*.

cincinnatiensis, Meek, 1872, (*Cythere cin-*
cincinnatiensis.) Proc. Acad. Nat. Sci., p.
331, and Ohio Pal., vol.

1, p. 158, Hud. Riv. Gr.

concinna, see *Primitia*
concinna.

crassimarginata, Win. FIG. 993.—*Cyther-*
chell, 1862, (*Cythere* *opsis cincinnati-*
crassimarginata.) Proc. *ensis*.

Acad. Nat. Sci., p. 429, Marshall Gr.

crenulata, Emmons, 1856, (*Cytherina*
crenulata.) Am. Geol., p. 220, Tren-
ton Gr.

FIG. 994.—*Cytheropsis glandella*.

glandella, Whitfield, 1882, (*Cytherellina*
glandella.) Bull. Am. Mus. Nat. Hist.,
vol. 1, p. 94, Warsaw Gr.

irregularis, S. A. Miller, 1878, (*Cythere*
irregularis.) Jour. Cin. Soc. Nat. Hist.,
vol. 1, p. 106, Hud. Riv. Gr.

nebraskensis, Geinitz, 1866, (*Cythere ne-*
braskensis.) Carb. und Dyas in Neb., p.
2, Coal Meas.

rugosa, Jones, 1858, Ann. Nat. Hist., 3d
ser., vol. 1, p. 240, Black Riv. Gr.

silliqua, Jones, 1858, Ann. Nat. Hist., 3d
ser., vol. 1, p. 249, Elack Riv. Gr.

simplex, White & St. John, 1868, (Cythere simplex,) Trans. Chi. Acad. Sci., p. 127, St. Louis Gr.

subcylindrica, Emmons, 1856, (Cytherina subcylindrica,) Am. Geo., p. 220, Trenton Gr.

subelliptica, Emmons, 1856, (Cytherina subelliptica,) Am. Geo., p. 220, Black Riv. Gr.

Dalmania, Emmrich, 1845. This name having been preoccupied for a genus of insects, Dalmanites has been substituted, though many authors prefer to use Odontochile, a name proposed by Corda.

DALMANITES, (Emmrich, 1845, *Dalmania*.) Barrande, 1852, Syst. Syl. Boh., vol. 1. [Ety. proper name.] Cephalic shield subulate, with lateral angles produced into spines; glabella widest anteriorly, rounded in front, with a highly convex anterior subelliptical lobe, three lateral furrows on each side; eyes prominent, subreniform, lenses numerous, situated posteriorly; facial sutures, curving slightly from the anterior margin, and each, following the curvature of the eye to the posterior part by a sigmoidal flexure, reach the lateral margin very slightly posterior to the eye itself; thorax with eleven segments, axis most convex, lateral lobes wider and more or less flattened; pygidium subtriangular, usually extended posteriorly into a spine, segments numerous. Type *D. caudatus*.

acantholeurus, Conrad, 1841, (Asaphus *acantholeurus*.) Ann. Rep. N. Y., p. 48, and Illust. Devon. Foss., pl. 19, Onondaga limestone.

FIG. 995.—*Dalmanites achates*.

achates, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 63, Trenton Gr.

ageria, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 57, and Illust. Devon. Foss., pl. 12, Up. Held. Gr.

anchiops, Green, 1832, (Calymene *anchiops*.) Monograph of Trilobites, p. 35, and Illust.

Devon. Foss., pl. 9, Schoharie grit.
anchiops var. *armatus*, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 56, Schoharie grit.

anchiops var. *serratus*, syn. for *D. anchiops*.
aspectans, Conrad, 1841, (Asaphus *aspectans*.) Ann. Rep. N. Y., p. 49, and Illust. Devon. Foss., pl. 13, Up. Held. Gr.

barrisi, Hall, 1888, Pal. N. Y., vol. 7, p. 48, Ham. Gr.

bebryx, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 61, Trenton Gr.

bicornis, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 196, Niagara Gr.

bifidus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 63, Up. Held. Gr.

boothi, Green, 1837, (Cryphaeus *boothi*.) Am. Jour. Sci., vol. 32, p. 343, and Pal. N. Y., vol. 7, p. 42, Ham. Gr.

breviceps, Hall, 1866, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 223, Hud. Riv. Gr.

callicephalus, Hall, 1847, (Phacops *callicephalus*.) Pal. N. Y., vol. 1, p. 247, Trenton Gr.

calliteles, Green, 1837, (Cryphaeus *calliteles*.) Am. Jour. Sci. and Arts, vol. 32, p. 346, and Illust. Devon. Foss., pl. 16, Ham. Gr.

calypso, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 61, and Illust. Devon. Foss., pl. 13, Up. Held. Gr.

carleyi, Meek, 1872, Am. Jour. Sci., 3d ser., vol. 3, p. 424, and Ohio Pal. vol. 1, p. 170, Hud. Riv. Gr.

comis, Hall, 1888, Pal. N. Y., vol. 7, p. 41, Up. Held. Gr.

concinnus, Hall, 1876, Illust. Devon. Foss., pl. 10, Schoharie grit.

concinnus var. *serrula*, Hall, 1888, Pal. N. Y., vol. 7, p. 30, Up. Held. Gr.

coronatus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 58, and Illust. Devon. Foss., pl. 12, Ham. Gr.

cuyahogae, Claypole, 1884, Geol. Mag., 3d ser., vol. 1, p. 303, Waverly Gr.

danae, Meek & Worthen, 1866, Proc. Acad. Nat. Sci. Phil., p. 264, and Geo. Sur. Ill., vol. 3, p. 363, Niagara Gr.

dentatus, Barrett, 1876, Am. Jour. Sci. and Arts, vol. 11, p. 200, Low. Held. Gr.

denticulatus, Conrad, 1841, (Asaphus *denticulatus*.) Ann. Rep. N. Y., p. 48, and Illust. Devon. Foss., pl. 10, Up. Held. Gr.

emarginatus, Hall, 1876, Illust. Devon. Foss., pl. 10, Up. Held. Gr.

epicrates, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 119, Low. Held. Gr.

erina, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 62, Up. Held. Gr.

helena, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 61, Up. Held. Gr.

intermedius, Walcott, 1877, 31st Rep. N. Y. St. Mus. Nat. Hist., p. 69, Trenton Gr.

laticaudatus, Hall, 1847. This name is erased from the list.

limulurus, Green, 1832, (Asaphus *limulurus*.) Monograph Trilobites, p. 48, and Pal. N. Y., vol. 2, p. 303, Niagara Gr.

logani, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 156, and Acad. Geol., p. 608, Up. Silurian.

FIG. 997.
Dalmanites calliteles.

h Rep. N. Y. St.
Up. Held. Gr.
ryphaeus boothii,
p. 343, and Pal.
am. Gr.
th Rep. N. Y. St.
847,
us.)
247,

837,
es),
arts,
ust.
16,

15th
Nat.
ust. Fig. 996.—Dal-
manites eul-
liceptus.

m. Jour. Sci., 3d
1 Ohio Pal. vol. 1,

Hall, 1888, Pal.
vol. 7, p. 41, Up.
Gr.

us, Hall, 1876,
Devon. Foss., pl.
chaharie grit.

us var. serrula,
1888, Pal. N. Y.,
p. 30, Up. Held.

us, Hall, 1862,
Rep. N. Y. St.
Nat. Hist., p. 58,
s. pl. 12, Ham. Gr.
34, Geol. Mag., 3d
verly Gr.

en, 1865, Proc.
p. 264, and Geo.
Niagara Gr.
Am. Jour. Sci.
10, Low. Held. Gr.
1841, (Asaphus
Rep. N. Y., p. 48,
Foss., pl. 10, Up.

6, Illust. Devon.
d. Gr.

Proc. Port. Soc.
9, Low. Held. Gr.

Rep. N. Y. St.
2, Up. Held. Gr.

h Rep. N. Y. St.
Up. Held. Gr.
1877, 31st Rep.
Hist., p. 69, Tren-

This name is

(Asaphus limul-
obolites, p. 48, and
303, Niagara Gr.
Nat. and Geo.,
cad. Geol., p. 608,

macrops, Hall, 1862, 15th Rep. N. Y. St.
Mus. Nat. Hist., p. 59, Up. Held. Gr.
meeki, Walcott, 1885, Monogr. U. S. Geo.
Sur., vol. 8, p. 207, Lower Devonian.
micurus, Green, 1832, (Asaphus micru-
rus,) Monograph Trilobites, p. 56, and
Pal. N. Y., vol. 3, p. 359, Low. Held. Gr.

FIG. 998.—*Dalmanites limulus*.

myrmecophorus, Green, 1835, (Asaphus
myrmecophorus,) Supp. to Monograph
of Trilobites, p. 16, and Illust. Devon.
Foss., pl. 13, Up. Held. Gr.
nasutus, Conrad, 1841, (Asaphus nasu-
tus,) Ann. Rep. N. Y., p. 48, and Pal.
N. Y., vol. 3, p. 362, Low. Held. Gr.
ohioensis, Meek & Worthen, 1871, Proc.
Acad. Nat. Sci. Phil., p. 91, and Ohio
Pal., vol. 1, p. 234, Up. Held. Gr.
phacoptyx, Hall, 1888, Pal. N. Y., vol. 7,
p. 31, Up. Held. Gr.
pleione, Hall, 1862, 15th Rep. N. Y. St.
Mus. Nat. Hist., p. 62, and Illust. Devon.
Foss., pl. 16, Up. Held. Gr.
pleuropteryx, Green, 1832, (Asaphus
pleuropteryx,) Monograph Trilobites,
p. 55, and Pal. N. Y., vol. 3, p. 356,
Low. Held. Gr.
pygmaeus, Hall, 1888, Pal. N. Y., vol. 7,
p. 56, Up. Held. Gr.
regalis, Hall, 1876, Illust. Devon. Foss.,
pl. 11, Schoharie grit.
selenurus, Eaton, 1832, (Asaphus selen-
urus,) Geo. Text Book, p. 31, and Illust.
Devon. Foss., pl. 12, Corniferous Gr.
tridens, Hall, 1859, Pal. N. Y., vol. 3, p.
361, Low. Held. Gr.
tridentiferus, Shumard, 1855, Geo. Rep.
Mo., p. 199, Low. Held. Gr.
troosti, Safford, Not defined.
verrucosus, Hall, 1863, Trans. Alb. Inst.,
vol. 4, p. 218, Niagara Gr.

vigilans, Hall, 1861, Rep. Prog. Geo. Sur.
Wis., p. 51, Niagara Gr.
werthneri, Foerste, 1885, Bull. Sci. Lab.
Denison Univ., p. 116. Not well de-
fined.

DICELLOCEPHALUS, Owen, 1852, Geo. Sur.
Wis., Iowa, and Min., p. 573, (written
by Owen Dikelocephalus.) [Ety. *dikella*,
mattock; *kephale*, head.] Cephalic shield
sublunate or semicircular, rather flat;
glabella oblong; sides parallel, rounded
in front, transverse behind; three lat-
eral furrows on each side, the posterior
two crossing the glabella and dividing
it into three lobes; facial sutures arise
in the center of the anterior border,
run parallel with the same to the front
of the eyes, are then directed backward,
make a sigmoid flexure over each eye,
and when near the neck segment ab-
ruptly curve laterally, reaching the pos-
terior border near the posterior spines
of the cephalic shield; thorax with nine
segments, axis narrower than lateral
lobes; pygidium with a flattened bor-
der, produced posteriorly on each side,
and rounded in the middle; axis ex-
tended only about half the length, with
four to six segments, which become ob-
solete on the lateral lobes. Type *D.*
minnesotensis.

affinis, Billings, 1865, Pal. Foss., vol. 1, p.
197, Quebec Gr. or Up. Taconic.
angustifrons, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 42, Potsdam Gr.
barabuensis, Whitfield, 1878, Ann. Rep.
Geo. Sur. Wis., p. 63, and Geo. Wis.,
vol. 4, p. 201, Low. Magnesian Gr.
belli, Billings, 1860, Can. Nat. and Geo.,
vol. 5, p. 301, Quebec Gr. or Up. Taconic.
bilobatus, Hall & Whitfield, 1877, U. S.
Geo. Expl. 40th Parallel, vol. 4, p. 226,
Potsdam Gr.
(?) corax, Billings, 1865, Pal. Foss., vol. 1,
p. 334, Quebec Gr. or Up. Taconic.
crassimarginatus, Whitfield, 1882, Geo.
Wis., vol. 4, p. 344, Potsdam Gr.
cristatus, Billings, 1860, Can. Nat. and
Geo., vol. 5, p. 301, Quebec Gr. or Up.
Taconic.
devinei, Billings, 1865, Pal. Foss., vol. 1,
p. 195, Quebec Gr. or Up. Taconic.
eatonii, Whitfield, 1878, Ann. Rep. Geo.
Sur. Wis., p. 65, and Geo. Wis., vol. 4,
p. 202, Low. Magnesian Gr.
expansus, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 45, Potsdam Gr.
finalis, Walcott, 1885, Monogr. U. S. Geo.
Sur., vol. 8, p. 89, Up. Taconic.
flabellifer, Hall & Whitfield, 1877, U. S.
Geo. Expl. 40th Parallel, vol. 4, p. 227,
Potsdam Gr.
(?) flagricaudus, White, 1874, Rep. Invert.
Foss., p. 12, and Geo. Sur. W. 100th Mer.,
vol. 4, p. 60, Quebec Gr. or Up. Taconic.
gothicus, Hall & Whitfield, 1877, U. S. Geo.
Expl. 40th Parallel, vol. 4, p. 242, Up.
Taconic. Probably a syn. for *Olenoides*
wahsatchensis.

granulosus, see *Ptychaspis granulosus*.
hisingeri, Billings, 1865, Pal. Foss., vol. 1, p. 196, Quebec Gr. or Up. Taconic.
inexpectans, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 90, Quebec Gr. or Up. Taconic.
iole, Walcott, 1885, Monogr. U. S. Geo. Sur. vol. 8, p. 43, Potsdam Gr. or Up. Taconic.
latifrons, Shumard, 1863, Trans. St. Louis Acad. Sci., vol. 2, p. 101, Potsdam Gr.
lodensis, Whitfield, 1880, Ann. Rep. Geo. Sur. Wis., p. 51, and Geo. Wis., vol. 4, p. 189, Potsdam Gr.
magnificus, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 301, Quebec Gr. or Up. Taconic.
marconi, Whitfield, 1884, Bull. Am. Mus. Nat. Hist., vol. 1, p. 139, Up. Taconic.
marica, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 44, Potsdam Gr.
megalops, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 301, Quebec Gr. or Up. Taconic.

minicensis, see *Ptychaspis minicensis*.

FIG. 989.—*Diellocephalus minnesotensis*.

minnesotensis, Owen, 1852, Rep. Wis., Iowa, and Minn., p. 574, Potsdam Gr.
minnesotensis var. *limbatus*, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 141, Potsdam Gr.
misa, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 144, Potsdam Gr.
missisquoi, Billings, 1865, Pal. Foss., vol. 1, p. 199, Quebec Gr. or Up. Taconic.
mulicinctus, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 226, Potsdam Gr.
nasutus, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 44, Potsdam Gr.
osceola, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 146, Potsdam Gr.
oweni, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 301, Quebec Gr. or Up. Taconic.
pauper, Billings, 1865, Pal. Foss., vol. 1, p. 200, Quebec Gr. or Up. Taconic.
pepinensis, Owen, 1852, Geo. Wis., Iowa, and Minn., p. 574, Potsdam Gr.
planifrons, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 301, Quebec Gr. or Up. Taconic.
pogonipensis, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Parallel, vol. 4, p. 243, Potsdam Gr.
quadriceps, see *Olenoides quadriceps*.
richmondensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 41, Potsdam Gr.
roomeri, Shumard, 1861, Am. Jour. Sci., vol. 32, p. 220, Potsdam Gr.
selectus, Billings, 1865, Pal. Foss., vol. 1, p. 199, Quebec Gr. or Up. Taconic.

sesostris, see *Ptychaspis sesostris*.
spiniger, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 143, Potsdam Gr.
wahsatchensis, see *Olenoides wahsatchensis*.

Dicranurus, syn. for *Acidaspis*.

hamatus, see *Acidaspis hamata*.

DIONIDE, Barrande, 1847, in Lith. Proc.

[Ety. from the mythological name *Dione*.] Body oval, tapering behind, trilobed, faintly convex; cephalic shield short, wide, semicircular, or crescentiform, produced at the postero-lateral angles into spines; glabella short, wide, strongly convex, smooth, no lateral furrows; two longitudinal furrows, making it trilobate; cheeks wide, with perforated margin; no eyes; no facial suture; hypostoma elliptical, with two bow-shaped wings in front, and posterior margin entire; six thoracic segments, with nodes on each side of the axial lobe; pygidium subtriangular, rounded behind axis, with numerous segments, and lateral lobes with radial furrows. Type *D. formosa*.

(?) *perplexa*, Billings, 1866, Catal. Sil. Foss. Antic., p. 67, Anticosti Gr.

DIPLETIS, Packard, 1885, Am. Nat., vol. 19, p. 291. [Ety. *dis*, double; *pelle*, small shield.] Cyclur-like in form; suborbicular, flattened, disk-like, sloping from the median area to the edge; divided in two parts, a cephalic shield and abdomen; integument thin, showing no segments. Type *D. diplo-discus*.

diplo-discus, Packard, 1885, Am. Nat., vol. 19, p. 291, Coal Meas. Poorly defined.

Dipleura, Green, syn. for *Homalonotus*.

dekayi, see *Homalonotus dekayi*.

DIPLOCHNITES, Dawson, 1863, Am. Jour. Sci. and Arts, 3d ser., vol. 5, p. 19. [Ety. *diploca*, double; *ichnos*, foot-print.] Consisting of two rows of impressions, each about an inch long and one-fourth of an inch wide, placed close together, while the rows are six inches apart, and the intermediate space smooth, as if a flat body had been drawn over it. Type *D. ænigma*.

ænigma, Dawson, 1863, Am. Jour. Sci. and Arts, 3d ser., vol. 5, p. 19, Coal Meas.

DIPLOSTYLUS,

Salter,

1863, Quar.

Jour. Geo.

Soc., vol.

19, p. 76.

[Ety. *Dip-*

lostylus,

double

tail, in al-

lusion to

the two

pairs of ap-

pendages

to the telson.]

body segments

arched, and

with minute

pleuræ; tail

segment large,

triangular,

FIG. 1000.—*Diplostylus dawsoni*. a, Tail, nat. size; b, terminal joint, enlarged.

to the telson.] Carapace unknown; body segments arched, and with minute pleuræ; tail segment large, triangular,

esostriis.
16th Rep. N. Y. St.
Mus. Nat. Hist.,
Potsdam Gr.
1863, p. 100. *Wahsatchi-*

pis.
amata.
In Lith. Proc.
theological mine
apering behind,
x; cephalic shield
lar, or crescenti-
ne postero-lateral
bellula short, wide,
th, no lateral fur-
row furrows, mak-
a wide, with per-
eyes; no facial
diptical, with two
front, and poste-
six thoracic seg-
each side of the
m subtriangular,
with numerous
lobes with radial
mora.

36, Catal. Sil. Foss.
i Gr.
Am. Nat., vol.
double; *pelle*, small
n form; suborbic-
like, sloping from
the edge; divided
elic shield and ab-
thion, showing no
pygidiscus.

85, Am. Nat., vol.
Poorly defined.
Homalonotus.
s dekayi.
63, Am. Jour. Sci.
l. 5, p. 19. [Ety.
foot-print.] Con-
impressions, each
and one-fourth of
d close together,
six inches apart,
space smooth, as
en drawn over it.

Am. Jour. Sci. and
19, Coal Meas.

Diplostylus dawsoni.
at size; b, terminal
argued.

Carapace unknown;
and with minute
large, triangular,

spinous, with two pairs of simple, ovate
appendages. Type *D. dawsoni*.
dawsoni, Salter, 1863, Quar. Jour. Geo.
Soc., vol. 19, p. 77, and Acad. Geol., p.
207, Coal Meas.

DIPTEROCARIS, Clarke, 1883, Am. Jour. Sci.
and Arts, 3d ser., vol. 25, p. 121. [Ety.
dipteros, two-winged; *karia*, shrimp.]
Carapace elongate, divided along the
major axis into two wings; greatest
width anteriorly; wings united medially
for one-third to one-fifth the length
of the carapace; ankylosed, but separated
toward the ends. Surface marked
concentrically. Type *D. pennidædali*.

pennidædali, Clarke, 1883, Am. Jour. Sci.
and Arts, 3d ser., vol. 25, p. 122, Chem-
mung Gr.

peacervæ, Clarke, 1883, Am. Jour. Sci.
and Arts, 3d ser., vol. 25, p. 123, Chem-
mung Gr.

procne, Clarke, 1883, Am. Jour. Sci. and
Arts, 3d ser., vol. 25, p. 122, Chemung Gr.

FIG. 100.—*Dithyrocaris carbonaria*.
Telson and styliets.

DITHYROCARIIS, Scouler,
1844, Syn. Carb.
Foss., Ireland &
McCoy, 1855, Brit-
ish Pal. Rocks, p.
181. [Ety. *dithyros*,
having two valves;
karia, shrimp.] Car-
apace semioval; an-
terior end rounded,
sometimes notched;
posterior end sub-
truncate, with lat-

eral angles produced into spines; sur-
face with faint imbricating striae, mar-
gins thickened and corrugated, with
three longitudinal ridges, one in the
middle extending the entire length, the
others not reaching the margin; tail
terminating in three tri-
angular spines. Type *D.*

scouleri.
belli, see *Mesothyra belli*.
carbonaria, Meek & Worthen,
1870, Proc. Acad. Nat. Sci.
Phil., p. 55, and Geo. Sur.
Ill., vol. 5, p. 618, Coal Meas.
neptuni, see *Mesothyra nep-*
tuni.

Dolichocephala, Clappole, 1883,
Proc. Am. Phil. Soc., p. 238,
syn. for *Stylonurus*.

lacoana, syn. for *Stylonurus*
excelsior.

DOLICHOMETOPUS, Angelin, 1852, Paleontol-
ogia Scandinavica. [Ety. *dolichos*,
long; *metope*, panel or space between
two hollows.] Cephalic shield with tu-
mid margin; eyes large, narrow, lunate;
glabella wider in front, smooth, no lateral
furrows; neck furrow-marked;
facial sutures, beginning at the poste-
rior margin near the lateral angles, are
directed toward the eyes, passing
which, they diverge to the anterior
margin; pygidium semicircular, strongly

convex, margin entire, axis almost
semicylindrical, with two or more fur-
rows. Type *D. suecicus*. It is doubt-
ful about this being an American
genus, as the identifications have been
made alone on the pygidium.

? *convexus*, Billings, 1865, Pal. Foss., vol.
1, p. 269, Quebec Gr. or Up. Taconic.

? *gibberulus*, Billings, 1865, Pal. Foss.,
vol. 1, p. 269, Quebec Gr. or Up. Taconic.

? *rarus*, Billings, 1865, Pal. Foss., vol. 1, p.
352, Calciferous Gr.

DOLICHOPTERUS, Hall, 1859, Pal. N. Y., vol.
3, p. 414. [Ety. *dolichos*, long; *pteron*,
wing.] Cephalic, thoracic, and caudal
portions similar to *Eurypterus*; post-
oral plate lyrate or cordiform lyrate;
central thoracic appendage from the
first thoracic segment, strong, thick,
and simple, in its anterior part; ante-
rior feet composed of strong, thick
joints, with curved terminal spines;
natatory organs having the joints elon-
gate, the seventh and eighth little
dilated, and the terminal palette ex-
tremely developed. Type *D. macro-*
chirus.

macrochirus, Hall, 1859, Pal. N. Y., vol.
3, p. 414, Waterlime Gr.

mansfieldi, Hall, 1877, Trans. Am. Phil.
Soc., p. 621, Lower Coal Meas.

ECHINOCARIS, Whitfield, 1880, Am. Jour.
Sci. and Arts, 3d ser., vol. 19, p. 34.
[Ety. *echinos*, sea urchin; *karia*, shrimp.]
Carapace bivalve; valves subovate,
united dorsally by a straight hinge,
anterior, posterior, and basal margins
rounded; surface marked by longitu-
dinal ridges or representative nodes or
ridges; abdomen naked, composed of
several segments and a caudal plate,
which is produced into an elongated

FIG. 102.—*Echinocaris punctata*.

spine, with a lateral movable spine on
each side; posterior margin of the ab-
dominal segments bearing spines.
Type *E. sublaevis*.

condylepis, Hall, 1888, Pal. N. Y., vol. 7,
p. 173, Chemung Gr.

longicauda, Hall, 1863, (Ceratiocaris longi-
cauda,) 16th Rep. N. Y. St. Mus. Nat.
Hist., p. 73, Genesee Slate.

multinodosa, Whitfield, 1880, Am. Jour.
Sci. and Arts, 3d ser., vol. 19, p. 38, Erie
shales.

- punctata, Hall, 1863, (Ceratiocaris punctata,) 16th Rep. N. Y. St. Mus. Nat. Hist., p. 74, Ham. Gr.
- pustulosa, Whitfield, 1880, Am. Jour. Sci. and Arts, 3d ser., vol. 19, p. 38, Erie shales.
- socialis, Beecher, 1884, Rep. of Prog. Pa. Geo. Sur. PPP, p. 10, Chemung Gr.
- sublevis, Whitfield, 1880, Am. Jour. Sci. and Arts, 3d ser., vol. 19, p. 36, Erie shales.
- whitfieldi, Clarke, 1885, Bull. U. S. Geo. Sur. No. 16, p. 45, Ham. Gr.
- wrightiana, Dawson, 1881, (Equisetides wrightianus,) Quar. Jour. Geo. Soc., vol. 37, p. 301, Portage Gr.
- ECHINOGNATHUS, Walcott, 1882, Am. Jour. Sci. and Arts, 3d ser., vol. 23, p. 213. [Ety. *echinos*, sea urchin; *gnathos*, jaw.] Founded upon fragments; endognathary limbs (one or more pairs) formed of eight or nine joints, six of which carry long, backward curving spines, articulated to their posterior side; terminal joint slender, elongate, acuminate; surface of body with scale-like markings. Type *E. clevelandi*.
- clevelandi, Walcott, 1882, Am. Jour. Sci. and Arts, 3d ser., vol. 23, p. 213, Utica Slate Gr.
- ELLIPTOCEPHALUS, Zenker, 1833, Beitrage zur Naturgeschichte der Urwelt, p. 51. [Ety. *ellipsis*, ellipse; *kephale*, head.] Broadly ovate; cephalic shield semicircular, depressed, without spines; glabella subquadrangular, rounded in front, without transverse furrows; eyes oblong, lunate, narrow, projecting outward; facial sutures short, commencing at the anterior margin, in front of the eyes, and curving over them toward the posterior angles; thoracic segments twelve, axis nearly as broad as lateral lobes; pygidium small, semicircular, trilobed. Type *E. hoffi*.
- ?curtus, Whitfield, 1877, Ann. Rep. Geo. Sur. Wis., p. 58, and Geo. Wis., vol. 4, p. 191, Potsdam Gr. Founded upon a fragment of the cephalic shield, and the generic reference is very doubtful.
- ELLIPTOCEPHALA, Emmons, 1844, Taconic System, p. 21. [Ety. *ellipsis*, ellipse; *kephale*, head.] Ovate; cephalic shield lunate, more than twice as wide as long, posterior angles produced in spines; groove and border on the anterior and lateral margins; glabella nearly equal in width throughout, and marked with three pairs of furrows; eyes large, elongate, semilunate, extending from near the base of the shield more than half way to the anterior margin; hypostoma broadly ovate; thirteen or fourteen articulations in the thorax, axis convex, lateral lobes flattened, last segments directed backward; pygidium narrow, elongated, axis acutely pointed. Type *E. asaphoides*. This generic name can stand in accordance with rule n of

the British Association of 1842, and the established laws of nomenclature adhered to by reputable scientists since that time, notwithstanding it is recommended to naturalists in selecting names to avoid such as too closely approximate words already adopted. It is true the masculine form of the word was preoccupied by Zenker, but the same can be said of Goniophora of Phillips, for Agassiz had preceded him in using the word Goniophorus; Schizodon was used for a mammal before King used Schizodus for a Lamelli-branch; Gray used Acrophylla before Nicholson used Acrophylum; and we might mention a hundred other instances where generic names, differing only in gender or termination, have been introduced and accepted by the best naturalists, and have come into such general use as to constitute part of the nomenclature of science. Olenellus can not be used to supplant Elliptocephala upon any ground of discovery, definition, or law.

- /asaphoides, Emmons, 1844, Taconic System, p. 21, and Pal. N. Y., vol. 1, p. 256, Up. Taconic.
- gilberti, Meek, 1874, (Olenellus gilberti,) Rep. Invert. Foss., p. 7, and Geo. Sur. 100th Mer., vol. 4, p. 44, Up. Taconic.
- howelli, Meek, 1875, (Olenellus howelli,) Rep. Invert. Foss., p. 8, and Geo. Sur. 100th Mer., vol. 4, p. 47, Up. Taconic.
- iddingsi, Walcott, 1885, (Olenellus iddingsi,) Monogr. U. S. Geo. Sur., vol. 8, p. 28, Up. Taconic.
- thompsoni, Hall, 1859, (Olenellus thompsoni,) 12th Rep. N. Y. St. Mus. Nat. Hist., p. 59, Up. Taconic.

FIG. 1003.—Elliptocephala thompsoni.

- undulostriata, Hall, 1847, (Olenus undulostriatus,) Pal. N. Y., vol. 1, p. 258, Up. Taconic. Poorly defined.
- ELYMOCARIS, Beecher, 1884, Rep. Pa. Geo. Sur. PPP, p. 13. [Ety. *elymos*, pod; *karis*, shrimp.] Carapace bivalve; elongate, longitudinally subquadrangular, dorsal line nearly as long as the valves; margins thickened; optic node near the anterior end, behind which are two elevations; two segments in the abdomen; telson a short, broad spine, with two lateral spines, crenulated on

1. *O. Masonae*, Lister Peach, 2. *J. G. S.*, Vol. 2. (1894) p. 674.
J. Walcott

a of 1842, and the nomenclature adopted by the scientists since, adding it is recommended in selecting a too closely adopted. It form of the word Zenker, but the of Goniophora of had preceded him Goniophorus; Schizomammal before a for a Lamellibrachya before phyllum; and we hundred other in names, differing termination, have accepted by the have come into to constitute part of science. Olenid to supplant Eleny ground of dis-

law.
1844, Taconic Sys-
Y., vol. 1, p. 256,

Olenellus gilberti,
p. 7, and Geo. Sur.
44, Up. Taconic.
Olenellus howelli,
p. 8, and Geo. Sur.
47, Up. Taconic.

FIG. 1004.—Encrinurus egani.

Ellipticocephala
composita.

1847, (Olenus undu-
Y., vol. 1, p. 258,
y defined.

84, Rep. Pa. Geo.
[Ety. elymos, pod;
pace bivalve; elon-
subquadrangular,
long as the valves;
optic node near the
d which are two
ments in the ab-
short, broad spine,
nes, crunulated on

the inner margins for fimbria. Type
E. siliqua.

carisella, Hall, 1888, Pal. N. Y., vol. 7, p.
181, Ham. Gr.

siliqua, Beecher, 1884, Rep. Pa. Geo. Sur.
PPP, p. 13, Chemung Gr.

Embolamus rotundatus, Rominger, syn. for
Bathyriscus howelli.

spinosa, Rominger, syn. for Olenoides
spinosus.

ENCINURUS, Emmrich, 1845, Neues Jahrb. f.

Mineral, p. 42. [Ety. en, prefix; krino,
parted; oura, tail.] Cephalic shield

semielliptical, tuberculated, lateral an-
gles produced into spines; glabella pyrri-

form, three furrows at each side toward
the base; cheeks flattened, triangular;

eyes in the middle of the cheeks, ele-
vated on foot-stalks; facial suture behind

the eye cuts the outer margin in front
of the angles; thorax with eleven seg-

ments; pygidium triangular, lateral
lobes with about eight segments, de-

flected, sometimes pointed; axis nar-
row, convex, with numerous segmental

lines. Type E. punctatus.

deltoides, Shumard, 1855, Geo. Sur. Mo.,
p. 198, Up. Sil.

egani, S. A. Miller,
1880, Jour. Cin. Soc.

Nat. Hist., vol. 2, p.
254, Niagara Gr.

elegantulus, Billings,
1866, Catal. Sil. Foss.

Antic., p. 62, Anti-
costi Gr.

excedrens, Safford,
Not defined.

laevis, Angelin, 1852,
(Cryptonymus lae-
vis,) Palaeontologia

Scandinavica, p. 4,
Up. Sil.

mirus, Billings, 1865,
Pal. Foss., vol. 1, p.

292, Quebec Gr. or Up. Taconic.

multisegmentatus, Portlock, 1843, (Am-
phion multisegmentatus,) Rep. Geo. of

Londonderry, etc., Anticosti Gr.

nereus, Hall, 1867, 20th
Rep. N. Y. St. Mus.

Nat. Hist., p. 425,
Niagara Gr.

ornatus, Hall & Whit-
field, 1875, Ohio Pal.

vol. 2, p. 154, Niag-
ara Gr.

phlyctainodes, Green, 1837, (Calymene
phlyctainodes,) Am. Jour. Sci. and

Arts, vol. 32, p. 167, and Pal. N. Y., vol.
2, p. 314, Niagara Gr.

punctatus, Wallenberg, 1821, Nova Acta
Soc. Upsal., Anticosti Gr.

trentonensis, Walcott, 1877, Rep. N. Y.
St. Mus. Nat. Hist., p. 68, Tren-

ton Gr.

varicostatus, Walcott, 1877, 31st Rep. N. Y.
St. Mus. Nat. Hist., p. 69, Trenton Gr.

vigilans, Hall, 1847, (Ceraurus vigilans,)

Pal. N. Y., vol. 1, p. 245, Black Riv. and
Trenton Grs.

Endymion, Billings, 1862. The name being
preoccupied for a genus of plants, the
author proposed Endymionia.

meeki, see Endymionia meeki.

ENDYMIONIA, Billings, 1865,
Pal. Foss., vol. 1, pp.

93, 281. [Ety. proper
name.] Cephalic

shield semioval, con-
vex; glabella ovate,

convex, an elongate
oval tubercle on each

side; thorax of six or
seven segments, axis

convex, side lobes flat,
groove crossing them

diagonally; pygidium semioval,
trilobed and divided by furrows into

segments; distinguished from Trinucleus
by the absence of a punctured border

on the head shield, and from Ampyx
by the form of the glabella, which has

a tubercle on each side, and is destitute
of a rostrum. Type E. meeki.

meeki, Billings, 1862, (Endymion meeki,)
Pal. Foss., vol. 1, pp. 93, 281, Quebec

Gr. or Up. Taconic.

Enoploura, Wetnerby, 1878, Jour. Cin. Soc.
Nat. Hist., vol. 1, p. 163. Proposed

instead of Anomalocystites, upon the
ground that it is a Crustacean, instead

of a Cystidean.

ESTHERIA, Ruppel, and Straus-Durckheim,
1837, Mus. Senckenberg., vol. 2, p.

119. [Ety. proper name.] Carapace
valves oval, globose, with a definite

hinge-line, well marked umbones and
concentric ridges, valves inequilateral,

subtrigonal or subovate, umbo near an-
terior end. Type E. dahalacensis. A

living genus, and probably not Paleozoic.

pulex, Clarke, 1882, Am. Jour. Sci. and
Arts, 3d ser., vol. 23, p. 466, Ham. Gr.

EUPROPS, Meek, 1867, Am. Jour. Sci., vol.
43, p. 394. [Ety. eu, very; pro, for-

ward; ops, eye.] Cephalo-thoracic
shield crescentic, more than twice as

wide as long, convex, lateral angles
terminating in spines; posterior margin

concave, from the lateral angles
two-thirds of the distance to the middle,

the central part being straight or
slightly concave; the ocular ridge sur-

rounds a crown-shaped or subquadr-
angular area, occupying the central third

of the shield; the sides are slightly
concave, in front there is a central

emargination, and posteriorly the ridge
is continued in a spine, on each side,

directed back over the abdomen; eyes
small, compound, located at the antero-

lateral angles of the crown-shaped cen-
tral area; mesial lobe small, narrowing

forward and reaching the ocular ridge,
in a linear carina; it bears a tubercle on

the posterior part; abdomen trans-
versely subelliptical, mesial lobe nar-

FIG. 1006.—Endymionia meeki.

FIG. 1005.—Encrinurus egani. Cephalic shield.

row; lateral lobes wide, flattened on the margins; segments defined by linear ridges, which are produced beyond the flattened borders in curved mucronate spines; telson subtriangular, gradually tapering. Type *E. danie*.
 collecti, White, 1884, 13th Rep. Ind. Geo. Sur. Nat. Hist., p. 172, Coal Meas.
danie, Meek & Worthen, 1865, (Bellinurus *danie*.) Proc. Acad. Nat. Sci. Phil., p. 43, and Geo. Sur. Ill., vol. 2, p. 395, Coal Meas.

FIG. 1007.—*Euproops danie*. e, Eyes; p, pits; a, node.

longispina, Packard, 1885, Am. Naturalist, vol. 19, p. 291, Coal Meas.
EURYPTERELLA, Matthew, 1888, Trans. Roy. Soc. Can., p. 60. [Ety. diminutive of *Eurypterus*.] Minute; body ovate-elongate, obscurely divided into three regions, and faintly trilobed; head subtriangular, rounded at the outer corners, emarginate behind, seemingly composed of three anchylosed segments; thorax subquadrate, four segments, first one with a median ridge; abdomen elongately triangular, several segments, produced in a long, flexible tail; surface tuberculated. Type *E. ornata*.
ornata, Matthew, 1888, Trans. Roy. Soc. Can., p. 60, Lower Devonian.
EURYPTERUS, DeKay, 1825, Ann. Lyc. Nat. Hist. N. Y., vol. 1, p. 375. [Ety. *euros*, breadth; *pteron*, wing.] Body ovate-lanceolate, gradually attenuate behind, terminating in a spiniform tail; carapace on the upper side entire; eyes two distant, sessile, within the margin of the carapace, two simple oculiform tubercles or cornes situated subcentrally; thoracic and caudal portions composed of thirteen joints, the first narrow and the last prolonged in a triangular spine, with serrated edges; the first two articulations are anchylosed on the lower side, and from the central part a locomotive appendage is directed backward to the 3d or 4th articulation, terminating in two slender processes; mouth

central, beneath the carapace, surrounded by four pairs of jointed feet and a fifth larger pair; the three anterior pairs are similar; several joints bear a small articulating spine at the distal extremities, and the terminal joint consists of a spine; the fourth pair is longer, more slender, without spines, except on the terminal joint; the fifth pair are natatory, longer, more dilated, and placed beneath the posterior part of the carapace, basal joints composed of broad rhomboidal plates covering the posterior part of the carapace, over the inner edges of which there is a longitudinally ovate plate, at the anterior sinuate margin of which is the entrance to the mouth. Type *E. remipes*.

beecheri, Hall, 1884, Geo. Sur. Pa. PPP, p. 30, Chemung Gr.
boylei, Whiteaves, 1884, Pal. Foss., vol. 3, p. 42, Guelph Gr.
dekayi, Hall, 1859, Pal. N. Y., vol. 3, p. 411, Waterlime Gr.
erlensis, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 196, Low. Held. Gr.
giganteus, Pohlman, 1882, Bull. Buff. Soc. Nat. Sci., vol. 4, p. 41, Waterlime Gr.
grandis, Grote & Pitt, 1875, (Eusarcus *grandis*.) Bull. Buff. Soc. Nat. Hist., vol. 3, p. 17, Waterlime Gr.
lacustris, Harlan, 1834, Trans. Geo. Soc. Penn., vol. 1, p. 98, and Pal. N. Y., vol. 3, p. 407, Waterlime Gr.
lacustris var. *robustus*, Hall, 1859, Pal. N. Y., vol. 3, p. 410, Waterlime Gr.
mazonensis, Meek & Worthen, 1868, Am. Jour. Sci., vol. 46, p. 21, and Geo. Sur. Ill., vol. 3, p. 544, Coal Meas.
microphthalmus, Hall, 1859, Pal. N. Y., vol. 3, p. 407, Low. Held. Gr.
pachychirus, Hall, 1859, Pal. N. Y., vol. 3, p. 412, Waterlime Gr.
pennsylvanicus, Hall, 1877, Proc. Am. Phil. Soc., p. 621, Carboniferous.
potens, Hall, 1884, Geo. Sur. Pa. PPP, p. 37, Carboniferous.

prominens, Hall, 1884, Proc. Am. Ass. Sci., vol. 33, p. 420, and Pal. N. Y., vol. 7, p. 157, Clinton Gr.

pulicaris, Salter, 1863, Quar. Jour. Geo. Soc., vol. 19, p. 78, and Acad. Geol., p. 523, Up. Devonian.

pustulosus, Hall, 1859, Pal. N. Y., vol. 3, p. 413, Waterlime Gr.

remipes, DeKay, 1825, Ann. Lyc. Nat. Hist. N. Y., p. 375, and Pal. N. Y., vol. 3, p. 404, Waterlime Gr.

FIG. 1008.—*Eurypterus remipes*. Diagram of small specimen.

carapace, sur-
s of jointed feet
the three anterior
eral joints bear a
ine at the distal
e terminal joint
the fourth pair is
, without spines,
al joint; the fifth
nger, more dilated,
the posterior part
al joints composed
plates covering the
carapace, over the
there is a longitu-
at the anterior sin-
ch is the entrance
E. remipes.

Geo. Sur. Pa. PPP,
4, Pal. Foss., vol. 3,

N. Y., vol. 3, p.

882, Ann. N. Y.

196, Low. Held. Gr.

382, Bull. Buff. Soc.

41, Waterlime Gr.

1875, (Eusarcus

Soc. Nat. Hist.,

me Gr.

Trans. Geo. Soc.

and Pal. N. Y., vol.

Gr.

e, Hall, 1859, Pal.

Waterlime Gr.

Worthen, 1868, Am.

21, and Geo. Sur.

al Meas.

1859, Pal. N. Y.,

Held. Gr.

59, Pal. N. Y., vol.

Gr.

1877, Proc. Am.

rboniferous.

p. Sur. Pa. PPP, p.

4, Proc. Am. Ass.

and Pal. N. Y., vol.

Diagram of small

men.

N. Y., p. 375, and

404, Waterlime Gr.

Diagram of small

men.

Diagram of small

men.

Diagram of small

men.

Diagram of small

men.

Diagram of small

men.

Diagram of small

men.

Diagram of small

men.

Diagram of small

men.

scorpionis, Grote & Pitt, 1875, (Eusarcus
scorpionis.) Bull. Buff. Soc. Nat. Hist.,
vol. 3, p. 1, Waterlime Gr.

stylus, Hall, 1884, Geo. Sur. Pa., PPP, p.
34, Low. Coal Meas.

tetragonophthalmus, Fischer, 1839, Bull.
Soc. Imper. Nat. Moscou., Water-
lime Gr.

Eusarcus, Grote & Pitt, 1875, Bull. Buff. Soc.
Nat. Hist., vol. 3, p. 1, syn. for Eu-
rypterus.

grandis, see Eurypterus grandis.

scorpionis, see Eurypterus scorpionis.

FABERIA, n. gen. [Ety. proper name.]

Minute crustaceans inclosed in a shell
with openings on the edge for the pro-
trusion of the feet and antennae. They
are referred to the Ostracoda, because
the test is like that of Leperditia and
Beyrichia, but they are distinguished
by being closed in a single shell; they
are evidently globose, depressed or vari-
able in form. Type F. anomala.

anomala, n. sp. Minute, subcircular in
outline, and flattened on each side;
thickness about one-fourth the diam-
eter; one edge somewhat sharp-ned; a

FIG. 1000.—*Faberia anomala*.
Three views, mag. 5 diam.

slit or open-
ing, about
eight times
as long as
wide, exists
on the thick-
er edge of
the shell;
and at less
than the thickness of the shell distant
from the slit, there is a circular open-
ing on the edge of the shell, and below
this reaching nearly to the thinner edge
of the shell, there is a very narrow slit
that does not seem to penetrate the
test. Found in the upper part of the
Hud. Riv. Gr., in Butler County, Ohio,
and now in the collection of Charles
Faber.

HARPES, Goldfuss, 1839, Nova Acta Physico
medica Academiae Caesarea Leopoldinae
Caroline Naturae Curiosorum, vol. 19,
p. 358. [Ety. harpe, a hook or sickle.]
Cephalic shield horseshoe-shaped,

very convex cen-
trally, flatly ex-
panded on the ex-
ternal margin, and
posterior angles
produced in long
spines; glabella
very prominent,
short, front sub-
quadrate, posterior
part contracted, a
curved lateral fur-
row on each side
separating two el-
liptical lobes from the posterior half;
eyes small, near the anterior part of
the glabella; facial sutures along the
posterior angles, curving through the

FIG. 1010.—*Harpes*
angula.

eyes and then to the antero-lateral mar-
gins, thoracic segments numerous.
Type H. unguia.

antiquatus, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 408, Chazy Gr.

consuetus, Billings, 1860, Catal. Sil. Foss.
Antic., p. 64, Anticosti Gr.

dentoni, Billings, 1863, Can. Nat. and
Geo., vol. 8, p. 36, Hud. Riv. Gr.

escanabae,

Hall, 1851,

Geo. Lake

Sup. Land

Dist., vol. 2,

p. 211, Tren-
ton Gr.

granti, Bill-
ings, 1865,

Pal. Foss.,

vol. 1, p. 326,

Quebec Gr.

ottawensis,

Billings,

1865, Pal.

Foss., vol. 1, FIG. 1011.—*Harpes ottawensis*.

p. 182, Trenton Gr.

HARPIDES, Beyrich, 1840, Untersuchungen

Trilobiten als Fort. [Ety. from resem-
blance to the genus *Harpes*.] Cephalic

shield semicircular, margin wide and flat,

spines at posterior lateral angles; gla-
bella short, narrow, granular; lobe on

each side at base; cheeks have radi-
ating striae; eyes small, joining the

front end of the glabella by a small
ridge; thorax with 22 segments, pleurae

three times as wide as the axis. Type
H. rugosus. Only fragments have been

referred to this genus in America.

atlanticus, Billings, 1865, Pal. Foss., vol.

1, p. 281, Quebec Gr. or Up. Taconic.

concentricus, Billings, 1865, Pal. Foss.,

vol. 1, p. 292, Quebec Gr. or Up. Ta-
conic.

? deseriens, Billings, 1865, Pal. Foss., vol.

p. 335, Quebec Gr. or Up. Taconic,

HARTT, Walcott 1884, Bull. U. S. Geo. Sur.

vol. 2, p. 263. [Ety. proper name.]

Distinguished from *Conocorypha* by

having a lobe or elevation in the front

of the glabella, small pygidium, and

sloping front to the cheeks and frontal

lobes. Type H. matthewi.

matthewi, Hartt, 1868,

(*Conocoryphites*

matthewi.) Acad.

Geol., p. 646, St.

John Gr.

Hauemannia, Hall, 1888, FIG. 1012.—*Harttia*

synonym for Dal-

manites.

Hemicrypturus, Green, syn. for *Asaphus*.

clintoni, Vanuxem, 1843, Geo. Rep. 3d

Dist. N. Y., p. 79, Clinton Gr. Gen-
eric relation not determined.

rasumowski, syn. for *Asaphus expansus*.

HIPONICHIARION, Matthew, 1885, Trans. Roy.

Soc. Can., p. 64. Breadth nearly equals

the length; broadly semi-elliptical to-

FIG. 1011.—*Harpes ottawensis*.

p. 182, Trenton Gr.

HARPIDES, Beyrich, 1840, Untersuchungen

Trilobiten als Fort. [Ety. from resem-
blance to the genus *Harpes*.] Cephalic

shield semicircular, margin wide and flat,

spines at posterior lateral angles; gla-
bella short, narrow, granular; lobe on

each side at base; cheeks have radi-
ating striae; eyes small, joining the

front end of the glabella by a small
ridge; thorax with 22 segments, pleurae

three times as wide as the axis. Type
H. rugosus. Only fragments have been

referred to this genus in America.

atlanticus, Billings, 1865, Pal. Foss., vol.

1, p. 281, Quebec Gr. or Up. Taconic.

concentricus, Billings, 1865, Pal. Foss.,

vol. 1, p. 292, Quebec Gr. or Up. Ta-
conic.

? deseriens, Billings, 1865, Pal. Foss., vol.

p. 335, Quebec Gr. or Up. Taconic,

HARTT, Walcott 1884, Bull. U. S. Geo. Sur.

vol. 2, p. 263. [Ety. proper name.]

Distinguished from *Conocorypha* by

having a lobe or elevation in the front

of the glabella, small pygidium, and

sloping front to the cheeks and frontal

lobes. Type H. matthewi.

matthewi, Hartt, 1868,

(*Conocoryphites*

matthewi.) Acad.

Geol., p. 646, St.

John Gr.

Hauemannia, Hall, 1888, FIG. 1012.—*Harttia*

synonym for Dal-

manites.

Hemicrypturus, Green, syn. for *Asaphus*.

clintoni, Vanuxem, 1843, Geo. Rep. 3d

Dist. N. Y., p. 79, Clinton Gr. Gen-
eric relation not determined.

rasumowski, syn. for *Asaphus expansus*.

HIPONICHIARION, Matthew, 1885, Trans. Roy.

Soc. Can., p. 64. Breadth nearly equals

the length; broadly semi-elliptical to-

the length; broadly semi-elliptical to-

the length; broadly semi-elliptical to-

the length; broadly semi-elliptical to-

the length; broadly semi-elliptical to-

the length; broadly semi-elliptical to-

the length; broadly semi-elliptical to-

the length; broadly semi-elliptical to-

the length; broadly semi-elliptical to-

ward the base, flattened, crossed by three symmetrical ridges; the middle one is inconspicuous. Type *H. eos*. eos, Matthew, 1885, Trans. Roy. Soc. Can., p. 64, St. John Gr.

FIG. 1013.—*Holometopus angelini*. Head and side view.

Holometopus, Angelin, 1852, Palæontologia Scandinavica. [Ety. *holos*, entire; *metopon*, space between the eyes.] Cephalic shield semicircular; glabella long, narrow, convex, widened in front, separated from cheeks by deep furrows; cheeks tumid; eyes small, situated well to the posterior; neck furrow distinct; facial sutures curved a little outward, in front of the eye. Type *H. limbatus*. angelini, Billings, 1862, Pal. Foss., p. 95, Quebec Gr. or Up. Taconic.

Homalonotus, König, 1825, Icones. Foss. Sectiles, p. 4. [Ety. *homalos*, on the same level; *notos*, back.] Cephalic shield hyperbolic, anterior angle subacute, margins rounded, surface convex; glabella subquadrate, shorter wider posteriorly, no furrows; eyes opposite the

FIG. 1014.—*Homalonotus delphinocephalus*.

central part of the glabella, small; facial suture, from the anterior angle to the margin, following the border of the margin, and curving like the letter S, it reaches the eye, and by a like curve passes to the posterior lateral angle;

thoracic segments 13, axis wider than the lateral lobes, which have subtruncate ends, with large distinct facets; pygidium hyperbolic and terminating in a spine. Type *H. knighti*.

atlas, Castelnau, 1843, Syst. Sil., p. 20. Not recognized.

dawsoni, Hall, 1860, Can. Nat. and Geo., vol. 5, p. 155, and Acad. Geol., p. 607, Up. Silurian.

dekayi, Green, 1832, (Dipleura dekayi.) Monograph Trilobites, p. 79, and Illust. Devon. Foss., pl. 25, Ham. Gr.

delphinocephalus, Green, 1832, (Trimerus delphinocephalus.) Monograph of Trilobites, p. 82, and Pal. N. Y., vol. 2, p. 309, Niagara Gr.

giganteus, Castelnau, 1843, Syst. Sil., p. 20. Not recognized.

herculeanus, Castelnau, 1843, Syst. Sil., p. 20. Not recognized.

jacksoni, Green, 1837, (Trimerus jacksoni.) Ann. Jour. Sci., vol. 32, p. 347, Up. Sil.

knighti, König, 1825, Icones. Foss. Sectiles, pl. 7, fig. 85, Low. Held. Gr.

major, Whitfield, 1885, Bull. Am. Mus. Nat. Hist., vol. 1, p. 193, Oriskany Gr.

vanuxemi, Hall, 1859, Pal. N. Y., vol. 3, p. 352, Low. Held. Gr.

ILLÆNURUS, Hall, 1863, 16th Rep. N. Y. Mus. Nat. Hist., p. 176. [Ety. from the genus *Illanus*; *oura*, tail.] Body broadly elliptical; cephalic shield short, convex, semi-elliptical; glabella subquadrate, convex, smooth, without distinct dorsal furrow; palpebral lobe marginal; cheeks wide; facial suture nearly vertical, slightly diverging, anterior to the eye; movable

cheeks wide and short; thoracic segments convex, central lobe wide, lateral lobes narrow, pygidium short, narrow, subelliptical, convex in front, more curved behind. Type *I. quadratus*.

convexus, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 66, and Geo. Wis., vol. 4, p. 203, Low. Mag. Gr.

eurekensis, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 97, Potsdam Gr.

quadratus, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 176, Potsdam Gr.

ILLÆNUS, Dalman, 1828, ueber die Palæaden oder die sogenannten Trilobiten, p. 51.

[Ety. *illaino*, to look awry, to squint.] Cephalic shield very convex and like one-fourth of a sphere, with the anterior margin slightly produced; glabella defined only as a slight convexity, between subparallel lines, on the posterior part of the shield; eyes semilunate, near the lateral margins smooth; facial suture makes a gentle curve from the antero-lateral margin to the eye, and then to the margin midway of the lateral lobes of the thorax; thoracic segments 9 or ten, broad; pygidium much like the cephalic shield. Type *I. crassicauda*.

ambiguus, Foerste, 1885, Bull. Sci. Lab. Denison Univ., p. 106, Niagara Gr.

3, axis wider than
which have subtrun-
ge distinct facets;
ic and terminating
l. knighti.

Syst. Sil., p. 20. Not

an. Nat. and Geo.,
cad. Geol., p. 607,

(*Dipleura dekayi*),
es, p. 79, and Illust.
Ham. Gr.

en, 1832, (*Trimerus*
Monograph of Trilo-
l. N. Y., vol. 2, p.

443, Syst. Sil., p. 20.

1843, Syst. Sil., p.

Trimerus jacksoni,
32, p. 347, Up. Sil.
Icones. Foss. Sec-
ow. Held. Gr.

5, Bull. Am. Mus.
193, Oriskany Gr.
Pal. N. Y., vol. 3,

r.

6th Rep. N. Y. Mus.
Ety. from the genus

Body broadly ellip-
short, convex, semi-

subquadrate, convex,
inct dorsal furrow;

inal; cheeks wide;
vertical, slightly di-

the eye; movable
ort; thoracic seg-

al lobe wide, lateral

ium short, narrow,
x in front, more

oe *I. quadratus*.
1878, Ann. Rep. Geo.

1 Geo. Wis., vol. 4,
r.

1885, Monogr. U. S.
97, Potsdam Gr.

16th Rep. N. Y. St.
76, Potsdam Gr.

ueber die Palaäden
n Trilobiten, p. 51.

awry, to squint.]
y convex and like

ere, with the an-
le produced; gla-

a slight convexity,
lines, on the pos-

ield; eyes semilu-
l margins smooth;

a gentle curve from
margin to the eye,
gin midway of the

thorax; thoracic
broad; pygidium
dic shield. Type *I.*

americanus, Billings, 1859, Can. Nat. and
Geol., vol. 4, p. 371, Trenton Gr.

angusticollis, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 376, Black Riv. Gr.

arcturus, Hall, 1847, Pal. N. Y., vol. 1, p.
23, Chazy and Black Riv. Grs.

acutus, Billings, 1865, Pal. Foss., vol. 1,
p. 279, Quebec Gr.

armatus, Hall, 1867, 20th Rep. N. Y. St.
Mus. Nat. Hist., p. 418, Niagara Gr.

barriensis, Murch. 1839, Sil. Syst. The
species formerly identified with this is
Illeenus ioxus.

bayfieldi, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 369, Chazy Gr.

clavifrons, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 379, Chazy and Black
Riv. Grs.

conifrons, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 378, Black Riv. Gr.

conradi, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 372, Black Riv. Gr.

consimilis, Billings, 1865, Pal. Foss., vol.
1, p. 277, Quebec Gr.

consobrinus, Billings, 1865, Pal. Foss., vol.
1, p. 280, Quebec Gr.

cornigerus, Hall, 1872, 24th Rep. N. Y.
St. Mus. Nat. Hist., p. 186, Niagara Gr.

crassicauda, Wahlenberg, 1821, (Ento-
mostracites *crassicauda*,) Nov. Act. Soc.
Upsal., vol. 8, p. 27, and Pal. N. Y.,
vol. 1, p. 229, Trenton and Galena Grs.

FIG. 1015.—*Illeenus globosus*. Two views.

cuniculus, Hall, 1867, 20th Rep. N. Y. St.
Mus. Nat. Hist., p. 421, Niagara Gr.

daytonensis, Hall & Whitfield, 1875, Ohio
Pal., vol. 2, p. 119, Niagara Gr.

fraternus, Billings, 1865, Pal. Foss., vol. 1,
p. 276, Quebec Gr.

globosus, Billings, 1859,
Can. Nat. and Geo.,
vol. 4, p. 367, Chazy
Gr.

grastonensis, Meek &
Worthen, 1869, Proc.
Acad. Nat. Sci. Phil.
and Geo. Sur. Ill., vol.

6, p. 508, Niagara Gr.
grandis, Billings, 1859,
Can. Nat. and Geo., vol. 4, p. 380, Hud.

Riv. Gr. and Mid. Sil.
herricki, Foerste, 1887, 15th Rep. Geo. and
Nat. Hist. of Minn., p. 479, Trenton Gr.

imperator, Hall, 1861, Rep. of Progr.
Wis., p. 49, and 20th Rep. N. Y. St.
Mus. Nat. Hist., p. 332, Niagara Gr.

incertus, Billings, 1865, Pal. Foss., vol. 1,
p. 332, Quebec Gr.

indeterminatus, Walcott, 1877, 31st Rep.
N. Y. St. Mus. Nat. Hist., p. 70, Black
Riv. Gr.

insignis, Hall, 1864, 20th Rep. N. Y. St.
Mus. Nat. Hist., p. 331, Niagara Gr.

ioxus, Hall, 1867, 20th Rep. N. Y. St.
Mus. Nat. Hist., p. 420, Niagara Gr.

latidorsatus, Hall, 1847, Pal. N. Y., vol. 1,
p. 230, Trenton Gr.

madisonianus, Whitfield, 1882, Geo. Wis.,
vol. 4, p. 307, Niagara Gr.

milleri, Billings, 1859, Can. Nat. and Geo.,
vol. 4, p. 375, Black Riv. and Tren-
ton Grs.

minnesotensis, Foerste, 1887, 15th Rep.
Geo. and Nat. Hist. of Minn., p. 478,
Trenton Gr.

niagarensis, Whitfield, 1880, Ann. Rep.
Geo. Sur. Wis., p. 68, Niagara Gr.

orbicaudatus, Billings, 1859, Can. Nat. and
Geo., vol. 4, p. 379, Hud. Riv. Gr. and
Mid. Sil.

ovatus, Conrad, 1843, (Thaleops *ovatus*,)
Proc. Acad. Nat. Sci. Phil., vol. 1, p.
332, and Pal. N. Y., vol. 1, p. 259, Black
Riv. Gr.

pterocephalus, Whitfield, 1878, Ann. Rep.
Geo. Sur. Wis., p. 87, and Geo. Wis.,
vol. 4, p. 309, Niagara Gr.

simulator, Billings, 1865, Pal. Foss., vol. 1,
p. 327, Quebec Gr.

taurus, Hall, 1861, Rep. of Progr. Wis.
Sur., p. 49, and Geo. Sur. Ill., vol. 3, p.
320, Trenton and Galena Grs.

trentonensis, Emmons, 1842, (Bumastus
trentonensis,) Geo. Rep. N. Y., p. 390,
and Pal. N. Y., vol. 1, p. 230, Tren-
ton Gr.

tumidifrons, Billings, 1865, Pal. Foss., vol.
1, p. 278, Quebec Gr.

vindex, Billings, 1865, Pal. Foss., vol. 1, p.
179, Chazy Gr.

worthenianus, syn. for *Illeenus insignis*.

ISOCHILINA, Jones, 1858, Can. Org. Rem.,
Decade 3, p. 197. [Ety. *isos*, equal;
cheilos, lip.] Equivalve, the margins of
the valves meeting uniformly, not over-
lapping, as in *Leperditia* greatest con-
vexity central or toward the anterior
end, eye tubercle present; muscular
spot not distinct, externally. Type *I.*
ottawa.

FIG. 1017.—*Illeenus taurus*.

taurus, Hall, 1861, Rep. of Progr. Wis.
Sur., p. 49, and Geo. Sur. Ill., vol. 3, p.
320, Trenton and Galena Grs.

trentonensis, Emmons, 1842, (Bumastus
trentonensis,) Geo. Rep. N. Y., p. 390,
and Pal. N. Y., vol. 1, p. 230, Tren-
ton Gr.

tumidifrons, Billings, 1865, Pal. Foss., vol.
1, p. 278, Quebec Gr.

vindex, Billings, 1865, Pal. Foss., vol. 1, p.
179, Chazy Gr.

worthenianus, syn. for *Illeenus insignis*.

ISOCHILINA, Jones, 1858, Can. Org. Rem.,
Decade 3, p. 197. [Ety. *isos*, equal;
cheilos, lip.] Equivalve, the margins of
the valves meeting uniformly, not over-
lapping, as in *Leperditia* greatest con-
vexity central or toward the anterior
end, eye tubercle present; muscular
spot not distinct, externally. Type *I.*
ottawa.

FIG. 1016.—*Illeenus globosus*. Slide view.

armata, Walcott, 1883, 35th Rep. N. Y. St. Mus. Nat. Hist., p. 213, Trenton Gr.

FIG. 1018.—*Isotelina jonesi*.

cylindrica, Hall, 1852, (Cytherinacylindrica,) Pal. N. Y., vol. 2, p. 14, Medina Gr.

gracilis, Jones, 1858, Can. Org. Rem. Decade 3, p. 98, Black Riv. and Trenton Grs.

jonesi, Wetherby, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 80, Trenton Gr.

labrosa, Jones, 1859, Ann. and Mag. Nat. Hist., 6th ser., vol. 3, p. 383, Low. Held. Gr.

ottawa, Jones, 1858, Can. Org. Rem., Decade 3, p. 97, Black Riv. Gr.

Isotelus, DeKay, 1825, Annals Lyceum Nat. Hist. N. Y., vol. 1, p. 174. [Ety. *isos*, equal; *telos*, end.] A subgenus of *Asaphus*.

canalis, see *Asaphus canalis*.

gigas, see *Asaphus gigas*.

maximus, see *Asaphus megistus*.

megistus, see *Asaphus megistus*.

vigilans, see *Asaphus vigilans*.

LEAIA, Jones, 1862, App. to Mon. Foss. Estheria, p. 116. [Ety. proper name.]

Carapace bivalve, subquadrate, thin, horny, truncated and slightly curved behind, rounded in front, straight on the dorsal edge; surface concentrically ridged and finely reticulated in the furrows; each valve crossed by one, two, or three ridges; the first and most conspicuous crosses from the anterior part of the umbo to the antero-ventral angle; the second, when it exists, reaches the postero-ventral angle, and the third lies along the dorsal margin. Type *L. leidy*.

leidy, Lea, 1856, (Cypricardia leidy,) Proc. Acad. Nat. Sci., vol. 7, p. 341, Coal Meas.

FIG.—1019.—*Leptacarinata*. B1, right valve; B2, enlarged; B3, dorsal view; c, left valve.

tricarinata, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 541, Coal Meas.

✕ *LEPERDITIA*, Rouault, 1851, Bull. Soc. Geo. France, 2d ser., t. 8, p. 377. [Ety. *lepis*, scale; *ditia*, double.] Carapace bivalve,

inequivalve, right valve larger than the left, and overlapping the ventral border, and to some extent the anterior and posterior borders of the left valve; valves smooth, convex, horny, oblong, longer than broad, bean-shaped, inequilateral, posterior half the broader; dorsal border straight; ventral border semi-circular. Type *L. britannica*.

alta, Conrad, 1843, (Cytherina alta,) Geo. Rep. 3d Dist. N. Y., p. 112, and Pal. N. Y., vol. 3, p. 373, Low. Held. Gr.

amygdalina, Jones, 1858, Can. Org. Rem., Decade 3, p. 97, Chazy Gr.

angulifera, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 197, Low. Held. Gr.

anna, Jones, 1858, Can. Org. Rem., Decade 3, p. 96, Hud. Riv. Gr.

anticostiana, Jones, 1858, (L. canadensis var. *anticostiana*.) Can. Org. Rem., Decade 3, p. 95, Hud. Riv. Gr.

arctica, Jones, 1856, Ann. and Mag. Nat. Hist., 2d ser., vol. 17, p. 87, Up. Sil.

argenta, Walcott, 1886, Bull. U. S. Geo. Sur., No. 30, p. 146, Up. Taconic.

billingsi, Jones, 1881, Ann. and Mag. Nat. Hist., 5th ser., vol. 18, Trenton Gr.

bivertex, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 11, Utica Slate Gr.

bivia, White, 1874, Rep. Invert. Foss., p. 11, and Geo. Sur. W. 100th Mer., vol. 4, p. 58, Quebec Gr.

byrnesi, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 123, Utica Slate Gr.

cæcigena, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 202, Hud. Riv. Gr.

canadensis, Jones, 1858, Ann. Nat. Hist., 3d ser., vol. 1, p. 244, Chazy to Trenton Gr.

capax, Safford. Not defined.

carbonaria, Hall, 1858, (Cythere carbonaria,) Trans. Alb. Inst., vol. 4, p. 33, and Bull. Am. Mus. Nat. Hist., p. 94, Warsaw Gr.

cayuga, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 83, Cornif. Gr.

concinnulla, Billings, 1865, Pal. Foss., vol. 1, p. 299, Quebec Gr. or Up. Taconic.

crepiformis, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 10, Hud. Riv. Gr.

cylindrica, Hall, 1871, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 231, Utica Slate and Hud. Riv. Grs.

dermatoides, Walcott, 1887, Am. Jour. Sci. and Arts, 3d ser., vol. 34, p. 192, Up. Taconic.

ebinina, Dwight, 1889, Am. Jour. Sci. and Arts, 3d ser., vol. 38, p. 144, Up. Taconic.

faba, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 186, Niagara Gr.

gigas, see *Asaphus gigas*.

maximus, see *Asaphus megistus*.

megistus, see *Asaphus megistus*.

vigilans, see *Asaphus vigilans*.

LEPERDITIA, Rouault, 1851, Bull. Soc. Geo. France, 2d ser., t. 8, p. 377. [Ety. *lepis*, scale; *ditia*, double.] Carapace bivalve,

inequivalve, right valve larger than the left, and overlapping the ventral border, and to some extent the anterior and posterior borders of the left valve; valves smooth, convex, horny, oblong, longer than broad, bean-shaped, inequilateral, posterior half the broader; dorsal border straight; ventral border semi-circular. Type *L. britannica*.

alta, Conrad, 1843, (Cytherina alta,) Geo. Rep. 3d Dist. N. Y., p. 112, and Pal. N. Y., vol. 3, p. 373, Low. Held. Gr.

amygdalina, Jones, 1858, Can. Org. Rem., Decade 3, p. 97, Chazy Gr.

angulifera, Whitfield, 1882, Ann. N. Y. Acad. Sci., vol. 2, p. 197, Low. Held. Gr.

anna, Jones, 1858, Can. Org. Rem., Decade 3, p. 96, Hud. Riv. Gr.

anticostiana, Jones, 1858, (L. canadensis var. *anticostiana*.) Can. Org. Rem., Decade 3, p. 95, Hud. Riv. Gr.

arctica, Jones, 1856, Ann. and Mag. Nat. Hist., 2d ser., vol. 17, p. 87, Up. Sil.

argenta, Walcott, 1886, Bull. U. S. Geo. Sur., No. 30, p. 146, Up. Taconic.

billingsi, Jones, 1881, Ann. and Mag. Nat. Hist., 5th ser., vol. 18, Trenton Gr.

bivertex, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 11, Utica Slate Gr.

bivia, White, 1874, Rep. Invert. Foss., p. 11, and Geo. Sur. W. 100th Mer., vol. 4, p. 58, Quebec Gr.

byrnesi, S. A. Miller, 1874, Cin. Quar. Jour. Sci., vol. 1, p. 123, Utica Slate Gr.

cæcigena, S. A. Miller, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 202, Hud. Riv. Gr.

canadensis, Jones, 1858, Ann. Nat. Hist., 3d ser., vol. 1, p. 244, Chazy to Trenton Gr.

capax, Safford. Not defined.

carbonaria, Hall, 1858, (Cythere carbonaria,) Trans. Alb. Inst., vol. 4, p. 33, and Bull. Am. Mus. Nat. Hist., p. 94, Warsaw Gr.

cayuga, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 83, Cornif. Gr.

concinnulla, Billings, 1865, Pal. Foss., vol. 1, p. 299, Quebec Gr. or Up. Taconic.

crepiformis, Ulrich, 1879, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 10, Hud. Riv. Gr.

cylindrica, Hall, 1871, 24th Rep. N. Y. St. Mus. Nat. Hist., p. 231, Utica Slate and Hud. Riv. Grs.

dermatoides, Walcott, 1887, Am. Jour. Sci. and Arts, 3d ser., vol. 34, p. 192, Up. Taconic.

ebinina, Dwight, 1889, Am. Jour. Sci. and Arts, 3d ser., vol. 38, p. 144, Up. Taconic.

faba, Hall, 1876, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 186, Niagara Gr.

gigas, see *Asaphus gigas*.

maximus, see *Asaphus megistus*.

megistus, see *Asaphus megistus*.

vigilans, see *Asaphus vigilans*.

Named after M. Leperdit of Rennes.
See Annals Nat. Hist. Soc. G. vol. 5, p. 340,
1851.

ve larger than the
the ventral border,
the anterior and
left valve; valves
y, oblong, longer
ed, inequilateral,
broader; dorsal
border semi-
ttanica.
herina alta, Geo.
p. 112, and Pal.
Low. Held. Gr.
Can. Org. Rem.,
y Gr.
1882, Ann. N. Y.
p. 197, Low.
Org. Rem., Dec-
Gr.
8, (L. canadensis
n. Org. Rem., Dec-
y Gr.
n. and Mag. Nat.
p. 87, Up. Sil.
Bull. U. S. Geo.
p. Taconic.
n. and Mag. Nat.
Trenton Gr.
Journ. Cin. Soc.
1, Utica Slate Gr.
White, 1874, Rep.
t. Foss., p. 11, and
Sur. W. 100th
vol. 4, p. 58, Que-
Gr.
S. A. Miller,
Cin. Quar. Jour.
tica Slate Gr.
1881, Jour. Cin.
4, p. 262, Hud.
1858,
ser.,
y to
med.
Cy-
ans.
33,
Nat. Fro. 1021.—Le-
Gr. perditia ce-
eligena, Nat.
15th ural size and
Nat. magnified.
Gr.
65, Pal. Foss., vol.
or Up. Taconic.
9, Jour. Cin. Soc.
0, Hud. Riv. Gr.
24th Rep. N. Y.
p. 231, Utica Slate
1887, Am. Jour.
vol. 34, p. 192,
Am. Jour. Sci.
ol. 38, p. 144, Up.
ep. N. Y. St. Mus.
gara Gr.

fabulites, Conrad, 1843, (Cytherina fabu-
lites,) Proc. Acad. Nat. Sci. Phil., p. 332,
Trenton Gr.
fonticola, Hall, 1867, 20th Rep. N. Y. St.
Mus. Nat. Hist., p. 428, Niagara Gr.
gibbera, Jones, 1856, Ann. and Mag. Nat.
Hist., 2d ser., vol. 17, p. 90, Niag-
ara Gr.
gibbera var. scalaris, see L. scalaris.
gracilis, see Isochilina gracilis.
hudsonica, Hall, 1859, Pal. N. Y., vol. 3,
p. 375, Low. Held. Gr.
jonesi, Hall, 1859, Pal. N. Y., vol. 3, p.
372, Low. Held. Gr.
josephana, Jones, 1858, (L. canadensis
var. josephana,) Can. Org. Rem., Dec-
ade 3, p. 94, Black Riv. to Trent-
ton Gr.
labrosa, Jones, 1858, (L. canadensis var.
labrosa,) Can. Org. Rem., Decade 3, p.
93, Chazy Gr.
louckana, Jones, 1858, (L. canadensis var.
louckana,) Can. Org. Rem., Decade 3, p.
93, Black Riv. Gr.
marginata, Keyserling, 1846, Wissen-
schaftliche Beobachtungen, etc., Niag-
ara Gr.
minutissima, Hall, 1871, 24th Rep. N. Y.
St. Mus. Nat. Hist., p. 231, Utica Slate
and Hud. Riv. Gr.
morganii, Safford. Not defined.
nana, Jones, 1858, (L. canadensis var.
nana,) Can. Org. Rem., Decade 3, p. 92,
Calceiferous Gr.
okeni, Munster, 1830, (Cythere okeni.)
Jahrbuch für Min., Geo. und Petrif.
Carboniferous.
ottawa, see Isochilina, ottawa.
ovata, Jones, 1858, Ann. and Mag. Nat.
Hist., 3d ser., vol. 1, p. 252, Black
Riv. Gr.
pauquettana, Jones, 1853, (L. canadensis
var. pauquettana,) Can. Org. Rem., Dec-
ade 3, p. 94, Black Riv. Gr.
parasitica, Hall, 1859, Pal. N. Y., vol. 3,
p. 276, Low. Held. Gr.
parvula, Hall, 1859, Pal. N. Y., vol. 3,
p. 376, Low. Held. Gr.
pennsylvanica, Jones, 1858, Ann. and
Mag. Nat. Hist., 3d ser., vol. 1, p. 251,
Clinton Gr.
spinulifera, Hall, 1860, 13th Rep. N. Y.
St. Mus. Nat. Hist., p. 92, Ham. Gr.
radiata, Ulrich, 1879, Jour. Cin. Soc. Nat.
Hist., vol. 2, p. 9, Utica Slate Gr.
rotundata, Walcott, 1885, Monogr. U. S.
Geo. Sur., vol. 8, p. 206, Devonian.
scalaris Jones, 1858, (L. gibbera var. scalaris),
Ann. and Mag. Nat. Hist., 3d
ser., vol. 1, p. 250, Waterlime Gr.
seneca, Hall, 1862, 15th Rep. N. Y. St.
Mus. Nat. Hist., p. 84, Ham. Gr.
sinuata, Hall, 1860, Can. Nat. and Geo.,
vol. 5, p. 158, Up. Silurian.
spinulifera, Hall, 1862, 15th Rep. N. Y.
St. Mus. Nat. Hist., p. 83, Up. Held Gr.
sublevis, Shumard, 1855, (Cythere sub-
levis,) Geo. Rep. Mo., p. 195, Low.
Magnesian Gr.

troyensis, see Aristozoe troyensis.
turgida, Billings, 1865, Pal. Foss., vol. 1,
p. 299, Quebec Gr.
ventralis, Billings, 1865, Pal. Foss., vol. 1,
p. 300, Quebec Gr.
unicornis, Ulrich, 1879, Jour. Cin.
Soc. Nat. Hist., vol. 2, p. 10, Utica
Slate Gr.
LEPIDILLA, Matthew, 1885, Trans. Roy. Soc.
Can., p. 62. [Ety. lepis, a scale.] Bi-
valve; hinge-line straight, projecting
from the general contour of the shell;
umbo and hinge-line separated from the
valve by a sinus, behind which there is
a foramen. Type L. anomala.
anomala, Matthew, 1885, Trans. Roy. Soc.
Can., p. 62, St. John Gr.
LEPIDITTA, Matthew, 1885, Trans. Roy. Soc.
Can., p. 61. [Ety. lepis, scale; ditto,
double.] Minute, obliquely semicir-
cular, wider on the anterior half, hinge
straight; umbones in the middle, low.
Type L. alata.
alata, Matthew, 1885, Trans. Roy. Soc.
Can., p. 61, St. John Gr.
curta, Matthew, 1885, Trans. Roy. Soc.
Can., p. 62, St. John Gr.

LEPIDOCOLEUS,
Faber, 1886,
Jour. Cin.
Soc. Nat.
Hist., vol. 9,
p. 15. [Ety.
lepis, scale;
koleos,
sheath.]
Body elon-
gate, com-
posed of two

FIG. 1022.—Leptidocoleus jamesi.
Mag. 2 diam. Lebanon spec-
imen.

series of thin, imbricating, angular plates,
interlocking and over-
lapping along the basal
edges; plates small,
more or less triangular
in outline; one side al-
ways longer than either
of the others; one side
usually sigmoidal; entire
outer surface marked
with striæ. Type L.
jamesi.

FIG. 1023.—Leptidocoleus jamesi.
St. Faber's
Cincinnati
specimen.

jamesi, Hall & Whitfield,
1875, (Plumulites jamesi),
Ohio Pal., vol. 2, p. 106,
Hud. Riv. Gr.
LICHAS, Dalman, 1826, Über
die Palæaden oder die
Sogenannten Trilo-
biten, p. 71. [Ety. my-
thological name.] Body
subovate, flat, granu-
lated; cephalic shield
somewhat lunate, often
pointed in front; gla-
bella large, convex, a furrow curving
inward and backward from the anterior
third on each side, and cutting off or
partly inclosing two oval spaces; cheeks
small; eyes large, reniform; eye-line

cutting the outer margin in front of the angles; thorax of ten segments; pleuræ flat, falcate, each with a furrow not reaching the margin; pygidium, side lobes flat, two falcate ribs on each side projecting beyond the margin, each with a mesial duplicating groove, middle lobe, semielliptical, pointed. Type *L. laciniatus*.

FIG. 1024.—*Lichas faberi*. Mag. 2 diam.

armatus, Hall, 1862. Preoccupied; changed to *L. eriopis*.
bigsbii, Hall, 1859, Pal. N. Y., vol. 3, p. 364, Low. Held. Gr.
boltoni, Bigsby, 1825, (*Paradoxides boltoni*,) Jour. Acad. Nat. Sci., vol. 4, p. 365, and Pal. N. Y., vol. 2, p. 311, Niagara Gr.
boltoni var. *occidentalis*, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 223, and 11th Rep. Ind. Geo. Sur., p. 344, Niagara Gr.
breviceps, Hall, 1863, Trans. Alb. Inst., vol. 4, p. 222, and 11th Rep. Ind. Geo. Sur., p. 343, Niagara Gr.
canadensis, Billings, 1866, Catal. Sil. Foss. Antic., p. 65, Antic. Gr.
champlainensis, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 342, Birdseye Gr.
cucullus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 266, and Geo. Sur. Ill., vol. 3, p. 299, Trenton Gr.
decipiens, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 104, Niagara Gr.
dracon, Hall, 1888, Pal. N. Y., vol. 7, p. 85, Up. Held. Gr.
emarginatus, Hall, 1879, 28th Rep. N. Y. St. Mus. Nat. Hist., p. 199, Niagara Gr.
eriopis, see *Terataspis eriopis*.

faberi, n. sp. Broadly elliptical, granulated; head somewhat crescentiform, slightly pointed in front, very convex, posterior angles terminating in short, obtuse spines; glabella very convex, divided into three lobes; central lobe contracted in the middle, widely expanded in front, and less expanded behind, and a slight furrow cuts off a small lobe from the postero-lateral angles; lateral lobes reniform; and another small lobe is separated from the posterior part of the cheeks by a stronger furrow; eyes prominent, reniform, and directed backward; occipital ring wide; axial lobe of thorax wider than the lateral lobes; pygidium lacinate, axis with two narrow articulations in front, and a longer posterior one that slopes backward and becomes confluent with the expanded border; lateral lobes composed of three expanded articulations, which terminate in acute points, and are marked in the central part by a groove for three-fourths of their length, which is represented by a rib on the under side; central lobe grooved in like manner, and bifid at the posterior extremity. The pygidium will readily distinguish it from *L. trentonensis*, beside the broader axial lobe of the thorax and somewhat different cephalic shield. Hud. Riv. Gr. at Cincinnati, Ohio. The specimen illustrated is from the collection of Charles Faber.

FIG. 1025.—*Lichas faberi*. Large and small pygidium.

grandis, see *Terataspis grandis*.
gryps, Hall, 1888, Pal. N. Y., vol. 7, p. 84, Up. Held. Gr.
harrisi, S. A. Miller, 1878, Jour. Cin. Soc. Nat. Hist., vol. 1, p. 106, Hud. Riv. Gr.
hispidus, Hall, 1888, Pal. N. Y., vol. 7, p. 77, Up. Held. Gr.
hylaëus, Hall, 1888, Pal. N. Y., vol. 7, p. 81, Up. Held. Gr.
jukesii, Billings, 1865, Pal. Foss., vol. 1, pp. 282 and 335, Quebec Gr.
minganensis, Billings, 1865, Pal. Foss., vol. 1, p. 181, Chazy or Black Riv. Gr.
nereus, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 226, Niagara Gr.
obvius, Hall, 1868, 20th Rep. N. Y. St. Mus. Nat. Hist., p. 424, Niagara Gr.
ptyonurus, Hall, 1888, Pal. N. Y., vol. 7, p. 86, Niagara Gr.
pugnax, Winchell & Marcy, 1865, Mem. Bost. Soc. Nat. Hist., p. 103, Niagara Gr.

elliptical, granu-
at crescentiform,
ont, very convex,
inating in short,
lla very convex,
bes; central lobe
iddle, widely ex-
less expanded be-
urrow cuts off a
ne postero-lateral
eniform; and an-
parated from the
ne cheeks by a
s prominent, reni-
ckward; occipital
e of thorax wider
; pygidium lacini-
arrow articulations
er posterior one
and becomes con-
d border; lat-
of three expanded
terminate in acute
ked in the central
or three-fourths of
s represented by a
side; central lobe
nner, and bifid at
mity. The pygid-
inguish it from L.
the broader axial
d somewhat differ-
Hud. Riv. Gr. at
the specimen illus-
d of Charles

FIG. 1026.—*Lichas*
trentonensis.

pastulosus, Hall, 1859, Pal. N. Y., vol. 3,
p. 366, Low. Held. Gr.

superbus, Billings,
1875, Can. Nat. and
Geol., vol. 7, p. 239,
Up. Held. Gr.

trentonensis, Conrad,
1842, (Asaphus tren-
tonensis,) Jour.
Acad. Nat. Sci., vol.
8, p. 277, and Pal.
N. Y., vol. 1, p. 235,
Black Riv. and Tren-
ton Grs.

LIOTRACUS, Angelin,
1852, Palaeontologia
Scandinavica, p. 23.
[Ety. *leiostrakos*,
smooth-shelled.]

Body elongate; test smooth or with mi-
croscopic punctures; glabella elevated,
furrows faint; dorsal furrow faint in
front; fixed cheek arched downward
at the sides; front
limb concave; occi-
pital ring aculeate; head
at the genal angle
rounded; ends of the
pleurae of the thorax
rounded; pygidium
minute, having few
segments. Type *L. ac-
uleatus*.

aurora, Hartt, 1868, (Cono-
cephalites *aurora*.)
Acad. Geol., p. 653, St.
John Gr.

linnarsoni, Brogger, 1878,
Paradoxides skifrene vid Krekling, p.
47, St. John Gr.

linnarsoni var. *alata*, Matthew, 1887,
Trans. Roy. Soc. Can., p. 147, St.
John Gr.

neglectus, Hartt, 1868, (Conocephalites
neglectus.) Acad. Geol., p. 652, St. John
Gr. Probably a syn. for *L. tener*.

ouangondianus, Hartt, 1868, (Conocephal-
ites *ouangondianus*.) Acad. Geol., p.
648, St. John Gr.

ouangondianus var. *gibbus*, Matthew,
1887, Trans. Roy. Soc. Can., p. 140, St.
John Gr.

ouangondianus var. *immarginata*, Mat-
thew, 1887, Trans. Roy. Soc. Can., p. 139,
St. John Gr.

ouangondianus var. *planus*, Matthew,
1887, Trans. Roy. Soc. Can., p. 140, St.
John Gr.

quadratus, Hartt, 1868, (Conocephalites
quadratus.) Acad. Geol., p. 654, St.
John Gr.

tener, Hartt, 1868, (Conocephalites *tener*.)
Acad. Geol., p. 652, St. John Gr.

Lisogocaris, Clarke, syn. for *Spathiocaris*.

lutheri, see *Spathiocaris lutheri*.

LONGICHOEPHALUS, Owen, 1852, Geo. Wis.,
Iowa, and Minn., p. 575. [Ety. *longus*,
long; *kephale*, head.] Cephalic shield,
having a wide frontal limb; posterior

FIG. 1027.—*Liotra-
chus aculeatus*.

Large and small
m.

grandis.

N. Y., vol. 7, p. 84,

, 1878, Jour. Cin.
ol. 1, p. 106, Hud.

Pal. N. Y., vol. 7, p.

Pal. N. Y., vol. 7, p.

Pal. Foss., vol. 1, pp.
Gr.

, 1865, Pal. Foss.,
or Black Riv. Gr.

6th Rep. N. Y. St.
26, Niagara Gr.

6th Rep. N. Y. St.
24, Niagara Gr.

, 1883, Pal. N. Y., vol. 7,

Marcy, 1865, M. m.
hist., p. 103, Niagara

angle of each cheek terminating in a
spine; glabella short, subquadrate, or
truncato-conical, highly arched; two or
three obscure furrows on each side;
base projected backward, in a spine of
greater or less length, in the median
line, over the thoracic segments; facial
sutures cut the anterior margin in front
of the eyes, and gently curve outward
and then inward to the anterior angles
of the palpebral lobes; thence curving
to the base of the eyes, they are di-
rected backward and slightly outward
to the posterior margin; pygidium sup-
posed to be semilunar, with little or no
border, and having four segments in
the axial lobe. Type *L. chippewensis*.

chippewensis,
Owen, 1852,
Geo. Wis., Iowa,
and Minn., p.
576, Potsdam
Gr.

hamulus, Owen,
1852, Geo. Wis.,
Iowa, and
Minn., p. 576,
Potsdam Gr.

wisconsinensis, Owen, 1852, Geo. Wis.,
Iowa, and Minn., p. 576, and 16th Rep.
N. Y. St. Mus. Nat. Hist., p. 146, Pots-
dam Gr.

LOGANELLUS, Devine, 1863, Can. Nat. and
Geo., vol. 8, p. 95. [Ety. proper name.]
General form ovate; cephalic shield
lunate; glabella convex, conical, two or
three oblique furrows on each side;
facial suture behind the eye curving
outward, and cutting the posterior mar-
gin inside the angle and in front of the
eye, curving outward to the frontal
margin; thorax broad, side lobes flat,

FIG. 1028.—*Lonchoceph-
alus chippewensis*.

FIG. 1029.—*Loganel-
us quebecensis*.

pleurae about
twelve; groove
running along the
middle nearly to
the extremities;
pygidium with a
well-defined axis,
side lobes de-
pressed, and with
four to six ribs;
distinguished from
Olenus by having
the facial suture
curved outward in
front of the eye.

Type *L. quebecensis*. This is one of
the forms often referred to *Conocoryphe*
or to *Ptychoparia*, but the genus may
be worth preserving.

quebecensis, Devine, 1863, Can. Nat. and
Geo., vol. 8, p. 95, Quebec Gr. or Up.
Taconic.

MEGALASPIS, Angelin, 1852, Palaeontologia
Scandinavica. [Ety. *megale*, great; *aspis*,
shield.] Body subelliptical; cephalic
shield obtusely pointed in front,
genal angles spined; glabella convex,

FIG. 1030.—*Megalaspis belemnura*. Pygidium.

expanded anteriorly, no lateral furrows, eyes large and close, posterior; facial sutures commencing at the anterior apex of the shield, curving laterally, and then contracting to the eyes in front and making a sigmoidal flexure, cut the posterior margin midway between the dorsal furrows and the genal spines; pygidium subtriangular, outer margin bordered, and terminating in a spine. Type *M. limbata*.

belemnura, White, 1874, Rep. Invert. Foss., p. 11, and Geo. Sur. W. 100th Mer., vol. 4, p. 59, Quebec Gr. or Up. Taconic.

FIG. 1031.—*Mesonacis vermontana*.

MENOCEPHALUS, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 577. [Ety. *meno*, strength; *kephale*, head.] Cephalic shield semicircular, with a narrow border all around; glabella highly convex, hemispherical or ovate, with a broadly rounded front, sometimes showing two inconspicuous lateral furrows

on each side; cheeks tumid; eyes distant from the middle of the glabella; facial suture cuts the front margin a little inside a line drawn lengthwise of the body and through the eye, and cuts the posterior margin a little outside this line; thoracic segments six or seven, axis convex, tapering a little narrower than the side lobes; pygidium semicircular, axis and side lobes divided by segmental furrows. Type *M. minnesotensis*.

?*globosus*, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 408, Quebec Gr. or Up. Taconic.

minnesotensis, Owen, 1852, Geo. Sur. Wis., Iowa, and Minn., p. 577, Potsdam Gr.

(?) *salteri*, Devine, 1863, Can. Nat. and Geo., vol. 8, p. 210, and Pal. Foss., vol. 1, p. 203, Up. Taconic.

FIG. 1032.
Menocephalus salteri.

?*sedgwicki*, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 301, and Pal. Foss., vol. 1, p. 407, Quebec Gr. or Up. Taconic.

MESONACIS, Walcott, 1885, Am. Jour. Sci. and Arts, 3d ser., vol. 29, p. 328. [Ety. *mesos*, middle; *akis*, point, spear.] Head and first fourteen segments like Elliptocephala, and the pygidium and ten posterior segments like Paradoxides. Type *M. vermontana*.

vermontana, Hall, 1859, (Olenus vermontanus,) 12th Rep. N. Y. St. Mus. Nat. Hist., p. 60, Up. Taconic, Georgia Gr.

MESORHYNCHUS, Hall, 1888, Pal. N. Y., vol. 7, p. lvi. [Ety. *mesos*, middle; *rhynchus*, small door.] Carapace subquadrate; valves in contact at the apices of two broad, subtriangular extensions, situated on the dorsal line opposite the eye nodes, forming a broad and short anterior or rostral cleft, and a long posterior cleft; test broadly infolded on the lower surface, thickened and produced into a conspicuous and acute posterior spine; posterior margin incurved and produced into a short spine at the dorsal line; surface with a single strong carina on each valve; abdomen consisting of two somites, of which the posterior is the longer; post-abdomen with a broad caudal plate, which is produced into a relatively short telson; lateral spines long and setaceous. Type *M. oceani*. *belli*, Woodward, 1870, (Dithyrocaris belli,) Geo. Mag., vol. 8, p. 106, Mid. Devonian.

neptuni, Hall, 1863, (Dithyrocaris neptuni,) 16th Rep. N. Y. St. Mus. Nat. Hist., p. 75, Ham. Gr.

oceani, Hall, 1888, Pal. N. Y., vol. 7, p. 187, Portage Gr.

spumæa, Hall, 1888, Pal. N. Y., vol. 7, p. 193, Ham. Gr.

veneris, Hall, 1888, Pal. N. Y., vol. 7, p. 193, Ham. Gr.

tumid; eyes dis-
e of the glabella;
e front margin a
wn lengthwise of
the eye, and cuts
a little outside
segments six or
tapering a little
lobes; pygidium
side lobes divided
Type M. min-

0. Can. Nat. and
d Pal. Foss., vol.
or Up. Taconic.

FIG. 1032.
Merocephalus
salteri.

60, Can. Nat. and
d Pal. Foss., vol.
or Up. Taconic.
Am. Jour. Sci.
29, p. 328. [Ety.
point, spear.]
een segments like
the pygidium and
ats like Paradox-
ontana.

(Olenus vermont-
Y. St. Mus. Nat.
ic, Georgia Gr.
l. N. Y., vol. 7, p.
lle; *thuris*, small
quadrate; valves
es of two broad,
ons, situated on
te the eye nodes,
short anterior or
ng posterior cleft;
on the lower sur-
produced into a
e posterior spine;
ved and produced
the dorsal line;
strong carina on
consisting of two
e posterior is the
a with a broad
produced into a
; lateral spines
Type M. oceani.
0, (Dithyrocaris
8, p. 106, Mid.

Dithyrocaris nep-
C. St. Mus. Nat.

N. Y., vol. 7, p.

N. Y., vol. 7, p.

N. Y., vol. 7, p.

MICRODISCUS, Emmons, 1856, Am. Geol., p. 116. [Ety. *mikros*, small; *diskos*, quoit.] Subelliptical; cephalic shield semicircular; glabella narrow, convex, rounded in front, more or less pointed behind, without furrows or occipital groove; cheeks more or less convex, no eyes or trace of sutures; thorax with four articulations, axis narrow, convex, lateral lobes wider, depressed; pygidium shorter than the cephalic shield, sub-trigonal or rounded posteriorly, trilobed, axis divided into four or six segments, and having a border. Type M. quadricostatus.

connexus, Walcott, 1887, Am. Jour. Sci. and Arts, 3d ser., vol. 34, p. 194, Up. Taconic.

dawsoni, Hartt, 1888, Acad. Geo., p. 654, St. John Gr.

lobatus, Hall, 1847, (Agnostus lobatus,) Pal. N. Y., vol. 1, p. 258, Up. Taconic.

meeki, Ford, 1876, Am. Jour. Sci. and Arts, 3d ser., vol. 11, p. 371, Up. Taconic.

parkeri, Walcott, 1886, Bull. U. S. Geo. Sur. No. 30, p. 157, Up. Taconic.

FIG. 1033.—Micro-
discus quadri-
costatus. Mag-
nified 5 diam.

NILEUS, Dalman, 1826, Über die Palæozoen oder die Sogenannten Trilobiten, p. 49. [Ety. mythological name.] Cephalic shield twice as wide as long, convex, lateral angles broadly rounded; glabella subquadrate, undefined anteriorly, no lateral furrows, convex, sloping in all directions from the central part; facial sutures in front, nearly parallel with, and almost reaching, the anterior margin, each forming a sigmoid flexure to the anterior part of the eye, then forming a semicircular eye-lobe from the posterior angle of the eye, and directed laterally to the posterior margin within the broadly rounded angle of the cephalic shield; eyes very large, lunate, with many lenses; eight thoracic segments, indistinctly trilobate, axial lobe the broader; pygidium twice as wide as long, not trilobate, no segments, broadly rounded posteriorly. Type N. armadillo.

affinis, Billings, 1865, Pal. Foss., vol. 1, p. 275, Quebec Gr. or Up. Taconic.

FIG. 1034.
Nileus
macrops.

macrops, Billings, 1865, Pal. Foss., vol. 1, p. 273, Quebec Gr. or Up. Taconic. scrutator, Billings, 1865, Pal. Foss., vol. 1, p. 274, Quebec Gr. or Up. Taconic.

Nothozoe, Barrande, Whitfield referred some ovate bodies found in the Potsdam sandstone, without characteristics, to this genus, under the name of Nothozoe vermontana. See Bull. Am. Mus. Nat. Hist., 1884, vol. 1, p. 144.

Nuttainia, syn. for Trinucleus.

concentrica, see Trinucleus concentricus.

sparsa, syn. for Homalonotus dekeyi.

Odontoccephalus, Conrad, 1840, Am. Geo. Rep.

N. Y. Not properly defined.

selenurus, see Dalmanites selenurus.

Odontochile, syn. for Dalmanites.

OGYRIA, Brongniart, 1822, Hist. Nat. Crust. Foss., p. 28. [Ety. mythological name.]

Flat or slightly convex; cephalic shield semicircular; glabella wider in front, with three lateral furrows on each side; eyes large, lunate, affixed centrally near the glabella, facial suture marginal in front, curving like the letter S, and terminating posteriorly midway between the outer angle of the cephalic shield and thoracic axis; thorax with narrow axis, pointed pleure, grooved, obscure, and remote fulcrum; eight segments; pygidium many segments, sides with radiating furrows, the interstices of which are divided by half rays. Type O. guettardi.

klotzi, Rominger, 1887, Proc. Acad. Nat. Sci. Phil., p. 12, Potsdam Gr.

parabola, Hall & Whitfield, syn. for Bathyriscus productus.

problematica, Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 63, Potsdam Gr.

serrata, Rominger, syn. for Olenoides nevadensis.

producta, Hall & Whitfield, see Bathyriscus productus.

spinosa, see Olenoides spinosus.

vetusta see Asaphus vetustus.

Olenellus, Hall, 1862, 15th Rep. N.Y. St. Mus.

Nat. Hist., p. 86, syn. for Elliptocephala.

asaphoides, see Elliptocephala asaphoides.

gilberti, see Elliptocephala gilberti.

howelli, see Elliptocephala howelli.

iddingsi, see Elliptocephala iddingsi.

vermontanus, Hall, 1859, see Mesonacis vermontana.

OLENOIDES, Meek, 1877, Geol. Expl. 40th Par., vol. 4, p. 25. [Ety. *Olenus*, and *oides* form.]

Ovate, head large, semicircular; glabella straight or slightly expanded in front; three pairs of furrows; eyes elongate; facial sutures extend obliquely outward from the anterior base of the eyes and cut the frontal margin; posteriorly they cut the margin at the pleural angle, and run subparallel to the margin to the posterior end of the eye; thorax with eight or more segments; axis strong, pleural groove broad, and lobes well defined; pygidium marked transversely on the

axis, and lateral segments directed backward. Type *O. nevadensis*.

FIG. 1035.—*Olenoides typicalis*.

fordi, Walcott, 1887, Am. Jour. Sci. and Arts, 3d ser., vol. 34, p. 195, Up. Taconic.
laevis, Walcott, 1886, Bull. U. S. Geo. Sur., No. 30, p. 187, Up. Taconic.

nevadensis, Meek, 1870, (Paradoxides nevadensis,) Proc. Acad. Nat. Sci. Phil., p. 62, and Geol. Expl. 40th Par., vol. 4, p. 23, Up. Taconic.

quadriceps, Hall & Whitfield, 1877, (Dicelloccephalus quadriceps,) Geol. Expl. 40th Par., vol. 4, p. 240, Up. Taconic.

spinus, Walcott, 1885, (Ogygia spinosa,) Mon. U. S. Geo. Sur., vol. 8, p. 63, Up. Taconic.

stissingensis, Dwight, 1889, Am. Jour. Sci. and Arts, 3d ser., vol. 38, p. 147, Up. Taconic.

typicalis, Walcott, 1886, Bull. U. S. Geo. Sur. No. 30, p. 183, Up. Taconic.

wahsatchensis, Hall & Whitfield, 1877, (Dicelloccephalus wahsatchensis,) Geol. Expl. 40th Par., vol. 4, p. 241, Up. Taconic.

Olenus, Dalman, 1826, Über die Palæaden oder die sogenannten Trilobiten, p. 54. Not an American genus.

? *logani*, see *Loganellus quebecensis*.

thompsoni, see *Elliptocephala thompsoni*.

undulostriatus, see *Elliptocephala undulostriata*.

vermontana, see *Mesonacis vermontana*.

ORYCTOCEPHALUS, Walcott, 1886, Bull. U. S. Geo. Sur. No. 30, p. 210. [Ety. *orykto*, furrowed; *kephale*, head.] Glabella ob-

long, transversely lobed; eyes central, narrow, ocular ridges connecting them with the axial furrow about the glabella; facial suture marginal in front, and cutting the posterior margin within

FIG. 1036.—*Oryctocephalus primus*. Cephalic shield.

the postero-lateral angles; free cheeks spinous; pygidium with segmented axis and pleural lobes; margin spinous. Type *O. primus*.

primus, Walcott, 1886, Bull. U. S. Geo. Sur. No. 30, p. 210, Up. Taconic.

PALEOCARIS, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 48.

[Ety. *palaio*, ancient; *karis*, shrimp.] Inner and outer pairs of antennae of nearly equal length

FIG. 1037.—*Oryctocephalus primus*. Pygidium.

FIG. 1038.—*Palaeocaris typus*. 3 diam.

the former each bearing a well developed accessory appendage; peduncles of both pairs shorter than the flagella; head about as long as the first two abdominal segments; thoracic legs long and slender, anterior pair not chelate; telson long, tapering, and horizontally flattened; styl-

FIG. 1039.—*Palaeocaris typus*. Caudal part 4 diam.

lets with first joint very small, second double, and also flattened horizontally. Type *P. typus*.

typus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 49, and Geo. Sur. Ill., vol 2, p. 405, Coal Meas.

ed; eyes central, connecting them by about the glau margin in front, for margin within

primus. Cephalo-

les; free cheeks with segmented margin spinous.

1937. — Oryctocephalus primus. Pygidium.

early equal length'

diam.

ing a well developed; peduncles of both pairs shorter than the flagella; head about as long as the first two abdominal segments; thoracic legs long and slender, anterior pair not chelate; telson long, tapering, and horizontally flattened; sty-
ry small, second
ied horizontally.

1865, Proc. Acad. and Geo. Sur. Mens.

PALÆOCREUSIA, Clarke, 1888, Pal. N. Y., vol. 7, p. 210. [Ety. *palaaios*, ancient; *Creusia*, a genus.] Capitulum ovate, patelliform, surface conical; apex truncated by a horizontal plane, forming a large central aperture; surface striated; basis tubuliform, subcylindrical or cup-shaped. Type *P. devonica*.

devonica, Clarke, 1888, Pal. N. Y., vol. 7, p. 210, Up. Held. Gr.

PALÆOPALÆMON, Whitfield, 1880, Am. Jour. Sci. and Arts, 3d ser., vol. 19, p. 40. [Ety. *palaaios*, ancient; *palæmon*, a genus.] Shrimp-like, thoracic carapace narrowed, but not rostrate in front and keeled on the back and sides; abdomen, six segments terminated by an elongated, triangular, and pointed telson; segments arched; pleuræ smooth, not lobed or expanded, extremities rounded; sixth segment bearing caudal flaps, one on each side, composed of five visible elements, the outer four apparently anthylosed to form a triangular plate on each side of the telson; thoracic ambulatory appendages elongated, smooth and filiform, except the upper second joint, which is laterally compressed; abdominal appendages short; antennæ large and strong. Type *P. newberryi*.

newberryi, Whitfield, 1880, Am. Jour. Sci. and Arts, 3d ser., vol. 19, p. 41, Erics shales.

PARADOXIDES, Brongniart, 1822, Hist. Nat. Crust. Foss., p. 31. [Ety. *paradoxos*, marvelous, paradoxical.] Cephalic shield lunate, margin thickened, not reflexed; glabella clavate or oval, moderately convex, enlarged anteriorly, three curved furrows cross it, dividing it into four parts; fixed cheeks tumid; eyes oblong, lunate, distant and opposite the second division of the glabella; facial suture, cutting the margin in front of the eye and curving S-like to the eye, and curving in like manner

FIG. 1040.—*Paradoxides bohemicus*.

to the posterior margin directly behind the eye; movable cheek tumid and prolonged in a spine; thorax 16 to 20; segments, axis convex, narrower than the lateral lobes, lateral lobes flattened and turned backward; pygidium circular or oval; axis segmented, short, lateral lobes flattened and projected backward. Type *P. tessini*.

abeneacus, Matthew, 1885, Trans. Roy.

Soc. Can., p. 78, St. John Gr.

acadicus, Matthew, 1883, Trans. Roy. Soc. Can., p. 103, St. John Gr.

acadicus var. *suricus*, Matthew, 1885, Trans. Roy. Soc. Can., p. 77, St. John Gr.

arcuatus, Harlan, 1835, Trans. Geo. Soc., syn. for *Triarthrus becki*.

barberi, N. H. Winchell, 1885, 13th Ann. R.p. Geo. Sur. Minn., p. 67, Potsdam Gr. Not a *Paradoxides*.

bennetti, Salter, 1850, Quar. Jour. Geo. Soc., vol. 15, p. 552, Up. Taconic.

boltoni, see *Lichna boltoni*.

FIG. 1041.—*Paradoxides harlani*.

decorus, Billings, 1874, Pal. Foss., vol. 2, p. 75, Up. Taconic.

eatonii, syn. for *Triarthrus becki*.

etemnicus, Matthew, 1883, Trans. Roy. Soc. Can. pp. 92, 271, St. John Gr.

etemnicus var. *breviatus*, Matthew, 1883, Trans. Roy. Soc. Can., p. 99, St. John Gr.

etemnicus var. *malicetus*, Matthew, Trans. Roy. Soc. Can., p. 101, St. John Gr.

etemnicus var. *pontificalis*, Matthew, 1883, Trans. Roy. Soc. Can., p. 102, St. John Gr.

etemnicus var. *quacoensis*, Matthew, 1883, Trans. Roy. Soc. Can., p. 102, St. John Gr.

etemnicus var. *suricoides*, Matthew, 1883, Trans. Roy. Soc. Can., p. 106, St. John Gr.

harlani, Green, 1834, Am. Jour. Sci., vol. 25, p. 336, Up. Taconic.

lamellatus, Hartt, 1868, Acad. Geol., p. 656, St. John Gr.

lamellatus var. *loricatus*, Matthew, 1883, Trans. Roy. Soc. Can., p. 106, St. John Gr.

micmac, Hartt, 1868, Acad. Geol., p. 657, and Trans. Roy. Soc. Can., vol. 2, p. 101, St. John Gr.

- nevadensis*, Meek, see *Olenoides nevadensis*.
quadripinnotus, Emmons, syn. for *Bathynotus holopyga*.
regina, Matthew, 1887, Am. Jour. Sci. and Arts, 3d ser., vol. 33, p. 380, and Trans. Roy. Soc. Canada, p. 115, St. John Gr.
tenellus, Billings, 1874, Pal. Foss., vol. 2, Up. Taconic.
thompsoni, see *Elliptocephala thompsoni*.
triarthrus, Harlan, syn. for *Triarthrus becki*.
vermontana, see *Mesonacis vermontana*.
Pellura, M. Edwards, 1840, Hist. Nat. Crust., t. 3, p. 344. Type *P. scarabaeoides*.
holopyga, see *Bathynotus holopyga*.
PEMPHIGASPIS, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 221. [Ety. *pemphix*, pustule; *aspis*, shield.] Founded upon part of a shield somewhat resembling the pygidium of a trilobite; a narrow, straight, annulated axis extends to the margin posteriorly; side lobes wider, ovate and ventricose. Type *P. bullata*.
bullata, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 211, Potsdam Gr.
PHACOPS, Emmrich, 1839, de Trilobites, Dissertatio Inauguralis, p. 19. [Ety. *phakos*, lens; *ops*, eye.] Form compact, glabella inflated and expanded in front; the two front pairs of furrows are obscure; eyes large, numerous lenses; genal rounded; pleurae rounded, pygidium moderate, of few (often coalesced) segments with an even border, never produced. Type *P. latifrons*.

bufo, Green, 1832, (Calymene bufo.) Monograph of Trilobites, p. 41, and Illust. Devon. Foss., pl. 8, Ham. Gr.
bombifrons, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 67, Up. Held. Gr.
cacaponia, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 68, Up. Held. Gr.
callicephalus, see *Dalmanites callicephalus*.
cristata, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 67, Up. Held. Gr.
cristata var. *pipa*, Hall, 1888, Pal. N. Y., vol. 7, p. 18, Up. Held. Gr.
hudsonica, Hall, 1859, Pal. N. Y., vol. 3, p. 355, Low. Held. Gr.
laticaudus, see *Dalmanites laticaudus*.
logani, Hall, 1859, Pal. N. Y., vol. 3, p. 353, Low. Held. Gr.
nupera, Hall, 1843, (Calymene nupera.) Geo. Rep. 4th Dist. N. Y., p. 262, and Illust. Devon. Foss., pl. 8, Chemung Gr.
orestes, Billings, 1860, Can. Nat. and Geo., vol. 4, p. 65, Mid. Sil.
rana, Green, 1832, (Calymene bufo var. rana.) Monograph of Trilobites, p. 42, and Illust. Devon. Foss., pl. 7, Ham. Gr.
trajanus, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 124, Low. Held. Gr.
trisulcata, Hall, 1943, (Calymene (?) trisulcata.) Geo. Rep. 4th Dist. N. Y., p. 74, and Pal. N. Y., vol. 2, p. 300, Clinton Gr.
PHÆTHONIDES, Angelin, 1878, Paleontologia Scandinavica, p. 21. [Ety. *phæthon*, radiant.] Head shield resembling Cyphaspis, the frontal area more concave, and lateral glabellar furrows stronger and generally duplicate; thorax having seven or more narrow segments; axis wide; pygidium resembling Proetus, relatively large, 8 to 12 annulations on the axis, and 8 or 9 on the pleurae; these extend to the margin, and are duplicate the entire length. Type *P. stokesi*.
arenicolus, Hall, 1888, Pal. N. Y., vol. 7, p. 134, Up. Held. Gr.
cyclurus, Hall, 1888, Pal. N. Y., vol. 7, p. 137, Low. Held. Gr.
denticulatus, Meek, 1877, (Proetus denticulatus.) Geol. Expl. 40th Par., p. 49, Devonian.
gemmeus, Hall, 1888, Pal. N. Y., vol. 7, p. 136, Low. Held. Gr.
varicella, Hall, 1888, Pal. N. Y., vol. 7, p. 135, Up. Held. Gr.
PHILLIPSIA, Portlock, 1843, Rep. Geol. Londonderry, p. 305. [Ety. proper name.] Cephalic shield sub-semicircular, angles terminating in spines; glabella subcylindrical, not contracted at base, three furrows on each side; eyes large, reniform, reticulated; thorax of 9 segments having pleural grooves and distinct facets; pygidium semioval, axis and lateral lobes furrowed, margin entire, smooth. Type *P. gemmulifera*.

auriculatus, Hall, 1862, (Proetus auriculatus.) 15th Rep. N. Y. St. Mus. Nat. Hist., p. 79, Waverly Gr.
bufo, Meek & Worthen, 1870, Proc. Acad. Nat. Sci., p. 52, and Geo. Sur. Ill., vol. 5, p. 528, Keokuk Gr.
cliftonensis, Shumard, 1858, Trans. St. Louis Acad. Sci., vol. 1, p. 227, Coal Meas.
coronata, Hall, syn. for *Cyphaspis ornata*.
doris, Hall, 1860, (Proetus doris.) 13th Rep. N. Y. St. Mus. Nat. Hist., p. 112, Waverly Gr.
howi, Billings, 1863, Can. Nat. and Geol., vol. 8, p. 209, Carboniferous.

ymene bufo var.
Trilobites, p. 42,
Foss., pl. 7.

Proc. Port. Soc.
p. 124, Low.

ymene (?) trisul-
cat. N. Y., p. 74,
2, p. 300, Clin-

1878, Paleonto-
21. [Ety. *phae-*
shield resembling
area more con-
labellar furrows
ally duplicate;
or more narrow
; pygidium re-
tively large, 8 to
axis, and 8 or 9
e extend to the
elicate the entire
ssi.

Pal. N. Y., vol. 7,

al. N. Y., vol. 7,

77, (Proetus den-
40th Par., p. 49,

al. N. Y., vol. 7,

al. N. Y., vol. 7,

Rep. Geol. Lon-
y. proper name.]

FIG. 1042. — *Phyllip-
sia gemmulifera*.

15th Rep. N. Y.
79, Waverly Gr.
en, 1870, Proc.
and Geo. Sur.
kuk Gr.

1858, Trans. St.
1, p. 227, Coal

Cyphasps or-

(*stus doris*), 13th
at. Hist., p. 112,

. Nat. and Geol.,
ferous.

insignis, Winchell, 1863, Proc. Acad. Nat.
Sci., p. 24, Burlington Gr.
laevis, see *Cyphasps laevis*.

lodiensis, Meek, 1875, Ohio
Pal., vol. 2, p. 323, Wa-
verly Gr.

major, Shumard, 1858,
Trans. St. Louis Acad.
Sci., vol. 1, p. 226, and
Pal. E. Neb., p. 238, Coal
Meas.

meramecensis, Shumard,
1855, Geo. Rep. Mo., p.
190, Archimedes lime-
stone or Keokuk Gr.

minuscule, see *Cyphasps*
minuscule.

missouriensis, Shumard, 1858, Trans. St.
St. Louis Acad. Sci., vol. 1, p. 225, Coal
Meas.

ornata, Hall, see *Cyphasps ornata*.

perannulata, Shumard, 1858, Trans. St.
Louis Acad. Sci., vol. 1, p. 296, Per-
mian Gr.

portlocki, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci., p. 268, and Geo. Sur.
Ill., vol. 5, p. 525, Keokuk Gr.

rockfordensis, Winchell, 1865, Proc. Acad.
Nat. Sci., p. 133, Kinderhook Gr.

sangamonensis, Meek & Worthen, 1865,
Proc. Acad. Nat. Sci., p. 271, and
Geo. Sur. Ill., vol. 5, p. 615, Coal
Meas.

scutula, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci., p. 270, and Geo. Sur.
Ill., vol. 5, p. 612, Coal Meas.

stevensoni, Meek, 1871, Reg. Rep. Uni-
versity W. Va., Kaskaskia Gr.

swallowi, Shumard, 1855, (Proetus *swal-*
lovi), Geo. Rep. Mo., p. 196, Wa-
verly Gr.

tennesseensis, Winchell, 1869, Geo. of
Tenn., p. 445, Waverly Gr.

tuberculata, Meek & Worthen, 1870,
Proc. Acad. Nat. Sci., p. 52, Burling-
ton Gr.

vindobonensis, Hartt, 1868, Acad. Geol.,
p. 313, Carboniferous.

Phyllites, Cozzens, 1848. Not identified.

chionensis, Cozzens, 1848. Not identified,
but probably the fragment of a *Dal-*
manites.

Platymotus, syn. for *Lichas*.

lolloni, see *Lichas bolloni*.

trentonensis, see *Lichas trentonensis*.

Plumulites, Barrande, syn. for *Turrilepas*.

devonicus, see *Turrilepas devonicus*.

gracillimus, see *Turrilepas gracillimus*.

jamesi, see *Lepidocoleus jamesi*.

newberryi, see *Turrilepas newberryi*.

Prestwichia, Woodward, 1867, Quar. Jour,
Geo. Soc. Lond., vol. 23. Not known
in America.

ericiensis, see *Protolimulus ericiensis*.

PRIMITIA, Jones, 1865, Ann. and Mag. Nat.
Hist., 3d ser., vol. 16, p. 415. [Ety.
primitia, first of the kind.] Carapace
minute; bivalve, equivalve, convex

oblong; hinge straight; surface of each
valve impressed, on the dorsal region
either medially or toward the anterior
extremity, with a vertical sulcus,
variable in size. Type *P. strangu-*
lata.

acadian, Matthew, 1885, Trans. Roy. Soc.
Can., p. 68, St. John Gr.

aqualis, Jones & Hall, 1886, Ann. and
Mag. Nat. Hist., 5th ser., vol. 17, p.
411, Low. Held. Gr.

cincinnatiensis, S. A. Miller,
1875, (Beyrichia *cincinnati-*
ensis), Cin. Quar. Jour.

Riv. Gr., vol. 2, p. 350, Hud. *Primitia*
cincinnati-
ensis.

concinna, Jones, 1858, (Cyther-
opsis concinna), Ann. and
Mag. Nat. Hist., 3d ser., vol. 1, p. 249,
Black Riv. Gr.

cristata, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 59, Calcifer-
ous Gr.

gregaria, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 58, Calcifer-
ous Gr.

logani, Jones, 1858, (Beyrichia *logani*),
Ann. and Mag. Nat. Hist., 3d ser., vol.
1, p. 244, Chazy Gr.

leperditoides, Jones, 1858, (Beyrichia
logani var. *leperditoides*), Can. Org.
Rem., Decade 3, p. 91, Chazy Gr.

mundula, Jones, 1855, Ann. and Mag.
Nat. Hist., 2d ser., vol. 16, p. 90, Low.
Devonian.

muta, Jones, 1865, Ann. and Mag. Nat.
Hist., 3d ser., vol. 16, p. 425, Up.
Sil.

reniformis, Jones, 1858, (Beyrichia *logani*
var. *reniformis*), Can. Org. Rem., De-
cade 3, p. 91, Chazy Gr.

rugulifera, Jones, 1858, (Beyrichia *ru-*
gulifera), Ann. and Mag. Nat. Hist., 3d
ser., vol. 1, p. 242, Niagara Gr.

scaphoides, Jones, 1889, Ann. and Mag.
Nat. Hist., 6th ser., vol. 3, p. 377, Low.
Devonian.

seeleyi, Whitfield, 1889, Bull. Am. Mus.
Nat. Hist., vol. 2, p. 60, Calcifer-
ous Gr.

sigillata, Jones, 1858, (Beyrichia *sigil-*
lata), Ann. and Mag. Nat. Hist., 3d ser.,
vol. 1, p. 242, Niagara Gr.

PROETUS, Steininger, 1830, Bemerkungen
über die Versteinerungen welche im
Uebergangs-Gebirge der Eifel, p. 4.
[Ety. mythological name.] Subellip-
tical; cephalic shield semicircular, mar-
gin thickened; glabella very convex,
parabolic, rounded anteriorly, no lat-
eral furrows; neck furrow well marked;
eyes prominent, smooth, close to gla-
bella; facial suture, on a line with the
eyes in front, curves gently backward
and reaches the posterior margin,
within the genal angle; thoracic seg-
ments 10, convex, lateral lobes, with
an oblique indentation; pygidium tri-

Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

1.8 2.0 2.2 2.5
2.8 3.2 3.6 4.0

10 01

FIG. 1046.—*Proetus alaricus*.

lobed, segmented, semicircular; axis very convex, short. Type *P. cuvieri*.
alaricus, Billings, 1860, Can. Nat. and Geo., vol. 5, p. 68, Hud. Riv. Gr.
angustifrons, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 70, Schoharie grit.
auriculatus, see *Phillipsia auriculatus*.
canaliculatus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 73, Up. Held. Gr.
clarus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 71, Up. Held. Gr.
conradi, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 69, and Illust. Devon. Foss., pl. 20, Schoharie grit.
coryceus, Conrad, 1842, (Asaphus *coryceus*), Jour. Acad. Nat. Sci., vol. 8, p. 277, and Pal. N. Y., vol. 2, p. 315, Niagara Gr.
crassimarginatus, Hall, 1843, (Calymene *crassimarginata*), Geo. Rep. 4th Dist. N. Y., p. 172, and Illust. Devon. Foss., pl. 20, Up. Held. Gr.
curvimarginatus, Hall, 1888, Pal. N. Y., vol. 7, p. 94, Up. Held. Gr.
davenportensis, Barris, 1879, Proc. Dav. Acad. Sci., vol. 2, p. 287, syn. for *P. prouti*.
delphinulus, Hall, 1888, Pal. N. Y., vol. 7, p. 111, Up. Held. Gr.
denticulatus, see *Phæthonides denticulatus*.
doris, see *Phillipsia doris*.
ellipticus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci., p. 267, and Geo. Sur. Ill., vol. 3, p. 460, Kinderhook Gr.
folliceus, Hall, 1888, Pal. N. Y., vol. 7, p. 101, Up. Held. Gr.
granulatus, Wetherby, 1881, Jour. Cin. Soc. Nat. Hist., vol. 4, p. 81, Kansas Gr.
haldemani, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 74, Ham. Gr.
hesione, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 70, and Illust. Devon. Foss., pl. 20, Schoharie grit.
jejunus, Hall, 1888, Pal. N. Y., vol. 7, p. 124, Ham. Gr.
junius, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 122, Low. Held. Gr.
latimarginatus, Hall, 1888, Pal. N. Y., vol. 7, p. 97, Up. Held. Gr.
loganensis, Hall & Whitfield, 1877, U. S. Geo. Exp. 40th Par., vol. 4, p. 264, Waverly Gr.
longicaudus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 108, and Illust. Devon. Foss., pl. 20, Ham. Gr.
macrobius, Billings, 1863, Proc. Port. Soc. Nat. Hist., vol. 1, p. 123, Low. Held. Gr.

macrocephalus, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 77, Ham. Gr.
marginalis, Conrad, 1839, (Calymene *marginalis*), Ann. Geo. Rep. N. Y., p. 66, and Illust. Devon. Foss., pl. 21, Tully limestone.
microgemma, Hall, 1888, Pal. N. Y., vol. 7, p. 109, Up. Held. Gr.
missouriensis, Shumard, 1855, Geo. Rep. Mo., p. 110, Waverly or Choctaw Gr.
nevadae, Hall, 1888, Pal. N. Y., vol. 7, p. 129, Low. Devonian.
occidens, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 80, Ham. Gr.
ovifrons, Hall, 1888, Pal. N. Y., vol. 7, p. 110, Up. Held. Gr.
parviusculus, Hall, 1860, 13th Rep. N. Y. St. Mus. Nat. Hist., p. 120, and 24th Rep., p. 223, Hud. Riv. Gr.
peroccidens, Hall & Whitfield, 1877, U. S. Geo. Exp. 40th Par., vol. 4, p. 262, Waverly Gr.
phocion, Billings, 1874, Pal. Foss., vol. 2, p. 65, Gaspe limestone No. 8, Devonian.
planimarginatus, Meek, 1871, Proc. Acad. Nat. Sci. Phil., p. 89, and Ohio Pal., vol. 1, p. 223, Up. Held. Gr.
protuberans, Hall, 1859, Pal. N. Y., vol. 3, p. 351, Low. Held. Gr.
prouti, Shumard, 1863, Trans. St. Louis Acad. Fig. 1047.—*Proetus phocion*.
rowii, Green, 1838, (Calymene *rowii*), Am. Jour. Sci., vol. 33, p. 406, and Illust. Devon. Foss., pl. 21, Ham. Gr.
spurlocki, Meek, 1872, Am. Jour. Sci., 3d ser., p. 426, and Ohio Pal., vol. 1, p. 161, Hud. Riv. Gr. The young of an Asaphus.
stenopyge, Hall, 1888, Pal. N. Y., vol. 7, p. 110, Up. Held. Gr.
stokesi, Murchison, 1839, (Asaphus *stokesi*), Sil. Syst., p. 625, and Pal. N. Y., vol. 2, p. 316, Niagara Gr.
swallowi, see *Phillipsia swallowi*.
tumidus, Hall, 1888, Pal. N. Y., vol. 7, p. 113, Up. Held. Gr.
verneuli, Hall, 1861, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 73, and Illust. Devon. Foss., pl. 20, Up. Held. Gr.

PROTICHTITES, Owen, 1852, Jour. Geo. Soc., vol. 8, p. 214. [Ety. *protos*, first; *ichnos*, foot-print; *lithos*, stone.] It consists of two rows of tracks or depressions, several inches apart; each row consists of numerous irregular and subcircular depressions, sometimes elongated; between the rows there is frequently a groove, and at other times, the surface

1862, 15th Rep.
Nat. Hist., p. 77,

39, (*Calymene mar-*
Rep. N. Y., p. 63,
Foss., pl. 21, Tully

88, Pal. N. Y., vol.
Gr.

d, 1855, Geo. Rep.
averly or Cho-

al. N. Y., vol. 7, p.

5th-Rep. N. Y. St.
0, Ham. Gr.

al. N. Y., vol. 7, p.

30, 13th Rep. N. Y.
p. 120, and 24th
Riv. Gr.

hitfield, 1877, U. S.
r., vol. 4, p. 262,

Fig. 1047.—*Protetus phocion*.

(*Calymene rowii*),
l. 33, p. 406, and
pl. 21, Ham. Gr.
Am. Jour. Sci., 3d
Pal., vol. 1, p. 161,
young of an *Asa-*

Pal. N. Y., vol. 7,

1839, (*Asaphus*
625, and Pal. N. Y.,
ra Gr.

swallowi.

Pal. N. Y., vol. 7, p.

15th Rep. N. Y. St.
73, and Illust. Dev.
ld. Gr.

52, Jour. Geo. Soc.,
r. *protos*, first; *ichnos*,
stone.] It consists

cks or depressions,
; each row consists
lar and subcircular

lines elongated; be-
rimes is frequently a
r times the surface

has apparently been pressed smooth.
Type *P. septemnotatus*.

FIG. 1049.—*Protobalanus septemnotatus*.

alternans, Owen, 1852,
Jour. Geo. Soc., vol.
8, pl. 14, Potsdam
Gr.

latius, Owen, 1852, Jour.
Geo. Soc., vol. 8, pl.
11, Potsdam Gr.

lineatus, Owen, 1852,
Jour. Geo. Soc., vol.
8, pl. 13, Potsdam
Gr.

logananus, Marsh,
1869, Am. Jour. Sci.
and Arts, 2d ser.,
vol. 48, Potsdam Gr.

multinotatus, Owen,
1852, Jour. Geo. Soc.,
vol. 8, pl. 12, Pots-
dam Gr.

octonotatus, Owen,
1852, Jour. Geo. Soc.,
vol. 8, pl. 10, Pots-
dam Gr.

septemnotatus, Owen,
1852, Jour. Geo. Soc.,
vol. 8, pl. 9, Pots-
dam Gr.

PROTOBALANUS, Whit-
field, 1888, Pal. N. Y.,
vol. 7, p. lxii. [Ety.
protos, first; *Bala-*

nus, genus.] Shell ovate about the
basis; composed of 12 plates of which
the carina is largest and most elevated;
rostrum small; lateral alia five on each
side; radial areas between the lateral alia
broad. Type *P. hamiltonensis*.

hamiltonensis, Whitfield, 1888, Pal. N. Y.,
vol. 7, p. 209, Ham. Gr.

PROTOCARIS, Walcott, 1884, Bull. U. S. Geo.
Sur., vol. 2, p. 283. [Ety. *protos*, first;
karis, shrimp.] Carapace without evi-
dence of a dorsal suture, rounded on
the dorsal line, and bent downward on
the sides; no rostrum; body many
jointed, 31 segments extending out
from beneath the carapace, the last
segment broader than the preceding,
and terminating in two spines. Type
p. marshi.

marshi, Walcott, 1884, Bull. U. S. Geo.
Sur., vol. 2, p. 283, Georgia Gr.

PROTOLIMULUS, Packard, 1886, Mem. Nat.
Acad. Sci., p. 150. [Ety. *protos*, first;
Limulus, a genus.] Cephalothorax
large, subsemicircular; genal angles pro-
duced; cephalic appendages small;
terminal segments of the posterior mem-
bers foliaceous; abdomen composed of
six (?) segments, including the large
caudal spine. Type *P. eriensis*.

eriensis, Williams, 1885, (*Prestwichia*
eriensis), Am. Jour. Sci. and Arts, 3d
ser., vol. 30, p. 46, Chemung Gr.

PROTOTYPUS, Walcott, 1886, Bull. U. S. Geo.
Sur., No. 30, p. 211. [Ety. *protos*, first;
typus, type.] Body ovate; head broad,

semicircular; glabella large, sides par-
allel, rounded in front, no furrows;
frontal limb narrow, in front of the

FIG. 1049.—*Protocaris marshi*.

glabella and bordered; fixed cheeks
crossed in front of the eyes by an ocular
ridge; eyes large, reniform; occipital
ring narrow; movable cheeks curved
on the outer margin and terminating

FIG. 1050.—*Prototypus hitchecocki*.

in spines; facial suture passing very
little outward from the eye to the an-
terior margin, which it cuts at nearly
right angles; behind the eye, it passes
obliquely outward and backward, with

slight curvature, to just within the cheek spine; thorax, twelve segments, strongly trilobed; pleura straight, broadly channelled and pointed; pygidium small, semielliptical, and marked by three furrows on the small axis and lateral areas. Type *P. hitchcocki*.

hitchcocki, Whitfield, 1884, (*Argellina hitchcocki*.) Bull. Am. Mus. Nat. Hist., vol. 1, p. 148, Up. Taconic.

PTEROCEPHALIA, Roemer, 1849, Texas, mit naturwissensch. Anhang. Bonn., and afterward in 1852, Kreid von Texas, p. 92. [Ety. *pteron*, wing; *kephale*, head.] Cephalic shield semicircular, nearly flat; glabella less than half the length of the head shield, with a flat, wing-like projection in front; two or three furrows on each side; neck furrow distinct, facial sutures directed nearly straight back from the anterior margin to the eye, after passing which it is directed at an angle laterally of about forty-five degrees to the posterior margin; eyes situate nearly opposite the posterior lobe of the glabella; pygidium subcircular, margin flattened and produced; axial lobe narrow, about ten segments. Type *P. sanctisabae*.

laticeps, Hall & Whitfield, 1877, (*Conocephalites laticeps*.) Geo. Expl. 40th Par., vol. 4, p. 221, Potsdam Gr.

occidens, Walcott, 1884, Mon. U. S. Geo. Sur., vol. 8, p. 58, Potsdam Gr.

sanctisabae, Roemer, 1849, Texas, mit naturwissensch. Anhang., and in 1852, Kreid von Texas, p. 92, Potsdam Gr.

PTERYGOTUS, Agassiz, 1830, Murch. Sil. Syst., p. 605. [Ety. *pteron*, wing; *ous*, ear.] Distinguished from *Eurypterus* by having eyes marginal instead of within the

FIG. 1051.—*Pterygotus problematicus*.

carapace, twelve segments instead of thirteen in the body, a bilobate caudal extremity and chelate antennae at the anterior part of the carapace. Type *P. problematicus*.

acuticaudatus, Pohlman, 1882, Bull. Buf. Soc. Nat. Sci., vol. 4, p. 42, Waterlime Gr.

buffaloensis, Pohlman, Bull. Buf. Soc. Nat. Hist., vol. 4, p. 17, Waterlime Gr.

cobbi, Hall, 1859, Pal. N. Y., vol. 3, p. 417, Waterlime Gr.

cummingsi, Grote & Pitt, 1875, Bull. Buf. Soc. Nat. Hist., vol. 4, p. 18, Waterlime Gr.

globicaudatus, Pohlman, 1882, Bull. Buf. Soc. Nat. Hist., vol. 4, p. 42, Waterlime Gr.

macrophthalmus, Hall, 1859, Pal. N. Y., vol. 3, p. 418, Waterlime Gr.

osborni, Hall, 1859, Pal. N. Y., vol. 3, p. 419, Waterlime Gr.

quadraticaudatus, Pohlman, 1882, Bull. Buf. Soc. Nat. Hist., vol. 4, p. 43, Waterlime Gr.

PTYCHASPIS, Hall, 1843, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 170. [Ety. *ptyche*, fold; *aspis*, shield.] Cephalic shield broad, with wide depressed convex cheeks; glabella cylindrical, convex, transversely lobed, prominent in front; eyes anterior to the middle; facial suture cutting the anterior border almost in front of the eye, and from below the eye it proceeds obliquely to the base a little without the center of the cheek, leaving the movable cheek near the size of the fixed cheek; movable cheek subtrapezoidal, border thickened, and extended backward in a spine. Type *P. miniacensis*.

barabuenensis, Winchell, 1864, Am. Jour. Sci. and Arts, 2d ser., vol. 37, p. 230, Potsdam Gr.

granulosa, Owen, 1852, (*Dikelocephalus granulosa*.) Geo. Wis., Iowa, and Minn., p. 575, Potsdam Gr.

miniscensis, Owen, 1852, (*Dikelocephalus miniscensis*.) Geo. Wis., Iowa, and Minn., p. 574, Potsdam Gr.

minuta, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 55, and Geo. of Wis., vol. 4, p. 186, Potsdam Gr.

pustulosa, Hall & Whitfield, 1877, U. S. Geo. Expl. 40th Par., vol. 4, p. 223, Potsdam Gr.

sesostris, Billings, 1865, (*Dikelocephalus sesostris*.) Pal. Foss., vol. 1, p. 198, Quebec Gr. or Up. Taconic.

speciosa, Walcott, 1879, 32d Rep. N. Y. St. Fig. 1052.—*Ptychaspis sesostris*.

striata, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 55, Potsdam Gr.

PTYCHOPARIA, Corda, 1847, Prodrom. einer Monographie der böhmischen Trilobiten, p. 141. [Ety. *ptyche*, fold.] Cephalic shield semilunar; genal angle spined or pointed; glabella narrow anteriorly, elevated; furrows four, distinct, directed forward; facial sutures widely separated, extending and converging

n, 1882, Bull. Buf.
p. 42, Waterline Gr.
Bull. Buf. Soc.
17, Waterline Gr.
N. Y., vol. 3, p. 417,

itt, 1875, Bull. Buf.
4, p. 18, Water-

n, 1882, Bull. Buf.
vol. 4, p. 42, Water-

, 1859, Pal. N. Y.,
ime Gr.

al. N. Y., vol. 3, p.

nlman, 1882, Bull.
vol. 4, p. 43, Water-

16th Rep. N. Y. St.
170. [Ety. *ptyche*.

] Cephalic shield
depressed convex
cylindrical, convex,
prominent in front;
middle; facial suture
border almost in
and from below the
quely to the base a
enter of the cheek,
cheek near the size
movable cheek sub-
thickened, and ex-
a spine. Type P.

l, 1864, Am. Jour.
er., vol. 37, p. 230,

52, (Dikelocephalus
is, Iowa, and Minn.,

52, (Dikelocephalus
Wis., Iowa, and
am Gr.

78, Ann. Rep. Geo.
Geo. of Wis., vol.

tr.
Whitfield, 1877, U. S.

ar., vol. 4, p. 223,

FIG. 1062.—Ptychopteria
seosstris.

78, Ann. Rep. Geo.
Potsdam Gr.

847, Prodr. einer
bohmischen Trilo-

ptyche, fold.] Ce-
lunar; genal angle

glabella narrow an-
tunnous four, distinct,

facial sutures widely
ang and converging

forward from the eyes, so as to intersect
the anterior margin within a point
where a line would cut it if drawn
through each eye parallel with the axis
(Corda's figure makes the facial sutures
cut the margin laterally, in a line drawn
at right angles to the anterior end of
the glabella); these lines extend them-
selves from the eyes to the posterior
margin by making a double curve, and
cut the margin within or near the lat-
eral angles; fixed cheek arched down-
ward at the sides; occipital ring spined;
thoracic segments fourteen; ends of
pleurae pointed or rounded; pygidium
medium size, six or seven articulations
in the axis; surface of test with minute
punctures or scattered tubercles. Type
P. striata.

affinis, Walcott, 1884, Mon. U. S. Geo. Sur.,
p. 54, Potsdam Gr.

anatina, Hall, 1863, (Conocephalites ana-
tinus,) 16th Rep. N. Y. St. Mus. Nat.
Hist., p. 158, Potsdam Gr.

(?) annectans, Walcott, 1884, Mon. U. S.
Geo. Sur., vol. 8, p. 91, Pogonip Gr.

antiquatus, Salter, 1859, (Conocephalites
antiquatus,) Jour. Geo. Soc., vol. 15, p.
554, Up. Taconic.

arenosa, Billings, 1861, (Conocephalites
arenosus,) Pal. Foss., vol. 1, p. 15, Pots-
dam Gr.

billingsi, Shumard, 1861, (Conocephalites
billingsi,) Am. Jour. Sci. and Arts, vol.
32, p. 220, Potsdam Gr.

binodosa, Hall, 1863, (Conocephalites bi-
nodosa,) 16th Rep. N. Y. St. Mus. Nat.
Hist., p. 160, Potsdam Gr.

breviceps, Walcott, 1884, Mon. U. S. Geo.
Sur., vol. 8, p. 49, Potsdam Gr.

calceferus, Walcott, 1879, (Conocephalites
calceferus,) 32d Rep. N. Y. St. Mus. Nat.
Hist., p. 129, Calceferous Gr.

calymenoides, Whitfield, 1877, Geo. Sur.
Wis., vol. 4, p. 179, Potsdam Gr.

clavata, Walcott, 1877, Am. Jour. Sci., 3d
ser., vol. 34, p. 198, Up. Taconic.

cordillerae, Rominger, 1887, (Conocephal-
ites cordillerae,) Proc. Acad. Nat. Sci.
Phil., p. 12, Potsdam Gr.

depressa, Shumard, 1861, (Conocephalites
depressus,) Am. Jour. Sci., vol. 32, p.
219, Potsdam Gr.

dissimilis, Walcott, 1884, Mon. U. S. Geo.
Sur., vol. 8, p. 51, Up. Taconic, Pros-
pect Mountain Gr.

eryon, Hall, 1863, (Conocephalites eryon,)
16th Rep. N. Y. St. Mus. Nat. Hist., p.
157, Potsdam Gr.

explanata, Whitfield, 1882, (Conocephal-
ites explanatus,) Geo. Sur. Wis., vol. 4,
p. 181, Potsdam Gr.

fitchi, Walcott, 1887, Am. Jour. Sci., 3d
ser., vol. 34, p. 197, Up. Taconic.

hartii, Walcott, 1879, (Conocephalites
hartii,) 32d Rep. N. Y. St. Mus. Nat.
Hist., p. 130, Calceferous Gr.

housensis, Walcott, 1886, Bull. U. S. Geo.
Sur., No. 30, p. 201, Up. Taconic.

laeviceps, Walcott, 1884, Mon. U. S. Geo.
Sur., p. 54, Potsdam Gr.

(?) linnarsoni, Walcott, 1884, Mon. U. S.
Geo. Sur. Terr., vol. 8, p. 47, Up. Ta-
conic.

minor, Shumard, 1863, (Conocephalites
minor,) Trans. St. Louis Acad. Sci., vol.
2, p. 105, Potsdam Gr.

minuta, Bradley, 1860, (Conocephalites
minutus,) Am. Jour. Sci., 2d ser., vol.
30, p. 242, Potsdam Gr.

nasuta, Hall, 1863, (Conocephalites nasu-
tus,) 16th Rep. N. Y. St. Mus. Nat.
Hist., p. 155, Potsdam Gr.

occidentalis, Walcott, 1884, Mon. U. S.
Geo. Sur., vol. 8, p. 51, Potsdam Gr.

oweni, Hall, 1863, 16th Rep. N. Y. St.
Mus. Nat. Hist., p. 155, Potsdam Gr.

patersoni, Hall, 1863, (Conocephalites pa-
tersoni,) 16th Rep. N. Y. St. Mus. Nat.
Hist., p. 159, Potsdam Gr.

pernasuta, Walcott, 1884, Mon. U. S. Geo.
Sur., vol. 8, p. 49, Potsdam Gr.

perseus, Hall, 1863, (Conocephalites per-
seus,) 16th Rep. N. Y. St. Mus. Nat.
Hist., p. 153, Potsdam Gr.

piochensis, Walcott, 1886, Bull. U. S. Geo.
Sur., No. 30, p. 201, Up. Taconic.

(?) prospectensis, Walcott, 1884, Mon.
U. S. Geo. Sur., vol. 8, p. 46, Up. Ta-
conic—Prospect Mountain Gr.

(?) quadrata, Whitfield, 1880, (Conocephal-
ites quadratus,) Geo. Sur. Wis.,
vol. 4, p. 180, Potsdam Gr.

rogersi, Walcott, 1884, Bull. U. S. Geo.
Sur., vol. 2, p. 283, Up. Taconic.

shumardi, Hall, 1863, (Conocephalites
shumardi,) 16th Rep. N. Y. St. Mus.
Nat. Hist., p. 154, Potsdam Gr.

similis, Walcott, 1884, Monogr. U. S. Geo.
Sur., vol. 8, p. 52, Potsdam Gr.

similis var. robusta, Walcott, 1884, Mon.
U. S. Geo. Sur., vol. 8, p. 53, Pots-
dam Gr.

subcoronata, Hall & Whitfield, 1877,
(Conocephalites subcoronatus,) Geo.
40th Par., vol. 4, p. 237, Up. Taconic.

teucer, Billings, 1861, (Conocephalites
teucer,) Geo. Vt., vol. 2, p. 951, Geor-
gia Gr.

verrucosa, Whit-
field, 1884,
(Conocephalites
verrucosus,) Bull. Am. Mus.
Nat. Hist., vol.
1, p. 139, Pots-

FIG. 1063.—Ptychoparia
zenkeri.

winona, Hall, 1863, (Conocephalites wi-
rona,) 16th Rep. N. Y. St. Mus. Nat.
Hist., p. 161, Potsdam Gr.

zenkeri, Billings, 1860, (Conocephalites
zenkeri,) Can. Nat. and Geo., vol. 5,
and Pal. Foss., vol. 1, p. 398, Up. Ta-
conic.

REMOPLEURIDES, Portlock, 1843, Rep. Geol.
Lond., p. 254. [Ety. *remus*, oar; *pleura*,
rib.] Cephalic shield subcircular or
transversely subelliptical; glabella large,

convex, oval, narrower anteriorly and abruptly bent down over the front; eyes large, semilunar, reaching the neck segment; rostral suture marked; free cheeks, small, narrow, subtriangular, and produced posteriorly in spines; thorax with ten segments, axial lobe very wide, and gradually tapering posteriorly; side lobes narrow, pleura short, falcate, directed backward; pygidium small and terminating in two short spines. Type *R. colbi*.

FIG. 1054. — *Remopleurides striatulus*. a, b, Hypostoma.

affinis, Billings, 1865, Pal. Foss., vol. 1, p. 325, Quebec Gr. *canadensis*, Billings, 1865, Pal. Foss., vol. 1, p. 182, Chazy Gr. *panderi*, Billings, 1865, Pal. Foss., vol. 1, p. 293, Quebec Gr. *schlothelimi*, Billings, 1865, Pal. Foss., vol. 1, p. 294, Quebec Gr. *striatulus*, Walcott, 1875, Cin. Quar. Jour. Sci., vol. 2, p. 347, Trenton Gr. *Rhabdichnites*, Dawson, 1873, Am. Jour. Sci. and Arts, 3d ser., vol. 5, p. 20. A name proposed for certain markings on the rocks which are not the remains of organisms.

RHACHURA, Scudder, 1878, Proc. Bost. Soc. Nat. Hist., vol. 19, p. 296. [Ety. *rachis*, ridge; *oura*, tail.] Type *R. venosa*.

venosa, Scudder, 1878, Proc. Bost. Soc. Nat. Hist., vol. 19, p. 296, Coal Meas.

RHINOCARIS, Clarke, 1888, Pal. N. Y., vol. 7, p. lviii. [Ety. *rhine*, file; *karis*, shrimp.] Cephalothorax univalvular, laterally appressed; outline as in *Ceratiocaris*; anterior extremity produced into a narrow, vertically flattened prora, continuous with substance of the carapace; axial line with a low ridge; abdomen composed of not less than four subcylindrical somites; post-abdomen bearing three spines, of which the telson is elongate and conical, and the cercopods flattened. Type *R. columbina*.

columbina, Clarke, 1888, Pal. N. Y., vol. 7, p. 195, Ham. Gr.

scaphoptera, Hall, 1888, Pal. N. Y., vol. 7, p. 197, Ham. Gr.

RIBERIA, Sharp, 1853, Jour. Geo. Soc., vol. 9, p. 157. [Ety. proper name.] Elongated, laterally compressed in the form of a Pholas; open at both ends and along the pedal margin, with a thick, transverse, internal plate near the anterior extremity, behind which is a cor-

rugated boss for the attachment of a muscle. Type *R. pholadiformis*.

calcifera, Billings, 1865, Pal. Foss., vol. 1, p. 340, Calciferous Gr.

compressa, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 344, Birdseye Gr.

longiuscula, Billings, Pal. Foss., vol. 1, p. 341, Calciferous Gr.

ventricosa, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 344, Birdseye Gr.

FIG. 1056. — *Rusichnites carbonarius*.

RUSICHNITES, Dawson, 1861, Can. Nat. and Geo., vol. 1, p. 363. Ety. *rusos*, wrinkled; *ichnos*, track.] Supposed by the author to be the track of a Crustacean, like the *Limulus*, and consisting of two undulated,

rounded, contiguous furrows; but the type resembles a fucoid quite as much, if not more, than it does a track, and if related to *Rusophycus*, which is clearly a fucoid, then it should be referred to the vegetable kingdom. Type *R. acadicus*.

acadicus, Dawson, 1861, Can. Nat. and Geo., vol. 1, p. 363, and Acad. Geol., p. 410, Coal Meas.

carbonarius, Dawson, 1868, Acad. Geol., p. 257, Carboniferous.

Salteria, Walcott, 1884. The name was preoccupied, and is a synonym for *Balliella*.

Sao, Barrande, 1846, and Syst. Sil. Boh., vol. 2. Type *Sao hirsuta*, a primordial form unknown in America.

? *lamottensis*, Whitfield, 1886, Bull. Am. Mus. Nat. Hist., vol. 1, p. 334, Birdseye Gr.

SCHIZODIACUS, Clarke, 1888, Pal. N. Y., vol. 7, p. 62. [Ety. *schiza*, cleft; *diskos*, quoit.] Carapace valves separable along the hinge; outline circular or ovate, narrow posteriorly; surface convex or depressed, elevated at the beaks, which are prominent, slightly incurved, and situated anteriorly, hinge-line equalling in length the greatest diameter of the carapace; edge parallel, not gaping, surface concentrically wrinkled. Type *S. capsula*.

capsula, Clarke, 1888, Pal. N. Y., vol. 7, p. 207, Ham. Gr.

SHUMARDIA, Billings, 1862, Pal. Foss., vol. 1, p. 92. [Ety. proper name.] Cephalic shield semicircular; glabella convex, subcylindrical, no eyes, pygidium semielliptical, axis cylindro-conical, ribbed, side lobes ribbed, distinguished from *Agnostus* by the ribs on the pygidium. Type *S. granulosa*.

FIG. 1057. — *Shumardia granulosa*.

e attachment of a
califormis.

5, 6.
0.

1, 2.
3.

6. Fig. 1055.—*Ribbia*
1, 2. *caliciformis*

Pal. Foss., vol. 1, p.

1886, Bull. Am. Mus.

344, Birdseye Gr.

NITES, Dawson,

1, Can. Nat. and

o., vol. 1, p. 363.

r. rufos, wrinkled;

nos, track.] Sup-

posed by the author

to be the track of a

crustacean, like the

mulus, and consist-

ing of two undulated

furrows; but the

could quite as much

does a track, and if

thus, which is clearly

could be referred to

kingdom. Type R.

361, Can. Nat. and

and Acad. Geol. p.

1868, Acad. Geol.,

us.

4. The name was

is a synonym for

d Syst. Sil. Boh., vol.

ica, a primordial form

eld, 1886, Bull. Am.

vol. 1, p. 334, Birds-

888, Pal. N. Y., vol.

schiza, cleft; disks,

glacialis, Billings, 1865, Pal. Foss., vol. 1,

p. 283, Up. Taconic.

granulosa, Billings, 1862, Pal. Foss., vol. 1,

p. 92, Quebec Gr. or Up. Taconic.

SOLENOCARIS, Meek, 1872, Proc. Acad. Nat.

Sci. Phil., p. 355. [Ety. *solen*, a genus

of shells; *karia*, shrimp.] Posterior

end subtruncated backward from below,

but not sinuous, as in *Colpocaris*; ocu-

lar tubercle obsolete, and no indica-

tions of having its valves ankylosed

along the dorsal margin. Type *S. strigata*.

This name was preoccupied for

another crustacean by Young, in 1868,

in Proc. Nat. Hist. Glasgow, vol. 1,

p. 171.

III., vol. 8, p. 153, St. Louis Gr.

strigata, Meek, 1872, Proc. Acad. Nat. Sci.

Phil., p. 335, and Ohio Pal., vol. 2, p.

321, Waverly Gr.

SOLENOPLEURA, Angelin, 1852, Palaeontolo-

gia Suecica, p. 26. [Ety. *selene*, moon;

pleuron, side.] Body ovate, test granu-

lated or tuberculous; head wide, semi-

circular; glabella prominent, furrows

distinct, dorsal furrows deep and con-

tinuous; fixed cheeks, elevated in the

middle, front limb convex; occipital

ring bearing a tubercle; genal angle

pointed; thoracic segments fourteen;

ends of pleuræ bluntly rounded; pygidium

medium size, few segments. Type

S. canaliculata.

aeolica, Whiteaves, 1887, Trans. Roy. Soc.

Can., p. 157, St. John Gr.

aeolica var. elongata, Matthew, 1887,

Trans. Roy. Soc. Can., p. 159, St.

John Gr.

communis, Billings, 1874, Pal. Foss., vol.

2, p. 72, Up. Taconic.

formosa, Hartt, 1868, (Conocephalites

formosus,) Acad. Geol., p. 654, St.

John Gr.

halli, Hartt, 1868, (Conocephalites halli,)

Acad. Geol., p. 654, St. John Gr.

nana, Ford, 1878, Am.

Jour. Sci. and Arts,

3d ser., vol. 15, p.

126, Up. Taconic.

orestes, Hartt, 1868,

(Conocephalites

orestes,) Acad.

Geol., p. 649, St.

John Gr.

robbi, Hartt, 1868,

(Conocephalites

robbi,) Acad. Geol.,

ra nana. Magnified.

p. 648, St. John Gr.

thyrsites, Hartt, 1868, (Conocephalites

thyrsites,) Acad. Geol., p. 653, St.

John Gr.

FIG. 1058.—*Solenocaris strigata*.
Cast of interior of left valve.

st ludo vici,
Worthen,
1884, Bull.
No. 2, Ill.
St. Mus.
Nat. Hist.,
p. 3, and
Geo. Sur.

Ill., vol. 8, p. 153, St. Louis Gr.

strigata, Meek, 1872, Proc. Acad. Nat. Sci.

Phil., p. 335, and Ohio Pal., vol. 2, p.

321, Waverly Gr.

SOLENOPLEURA, Angelin, 1852, Palaeontolo-

gia Suecica, p. 26. [Ety. *selene*, moon;

pleuron, side.] Body ovate, test granu-

lated or tuberculous; head wide, semi-

circular; glabella prominent, furrows

distinct, dorsal furrows deep and con-

tinuous; fixed cheeks, elevated in the

middle, front limb convex; occipital

ring bearing a tubercle; genal angle

pointed; thoracic segments fourteen;

ends of pleuræ bluntly rounded; pygidium

medium size, few segments. Type

S. canaliculata.

aeolica, Whiteaves, 1887, Trans. Roy. Soc.

Can., p. 157, St. John Gr.

aeolica var. elongata, Matthew, 1887,

Trans. Roy. Soc. Can., p. 159, St.

John Gr.

communis, Billings, 1874, Pal. Foss., vol.

2, p. 72, Up. Taconic.

formosa, Hartt, 1868, (Conocephalites

formosus,) Acad. Geol., p. 654, St.

John Gr.

halli, Hartt, 1868, (Conocephalites halli,)

Acad. Geol., p. 654, St. John Gr.

nana, Ford, 1878, Am.

Jour. Sci. and Arts,

3d ser., vol. 15, p.

126, Up. Taconic.

orestes, Hartt, 1868,

(Conocephalites

orestes,) Acad.

Geol., p. 649, St.

John Gr.

robbi, Hartt, 1868,

(Conocephalites

robbi,) Acad. Geol.,

ra nana. Magnified.

p. 648, St. John Gr.

thyrsites, Hartt, 1868, (Conocephalites

thyrsites,) Acad. Geol., p. 653, St.

John Gr.

FIG. 1059.—*Solenopleura halli*.
Magnified.

ra nana. Magnified.

p. 648, St. John Gr.

thyrsites, Hartt, 1868, (Conocephalites

thyrsites,) Acad. Geol., p. 653, St.

John Gr.

tumida, Walcott, 1887, Am. Jour. Sci. and

Arts, 3d ser., vol. 34, p. 196, Up. Taconic.

SPATHIOCARIS, Clarke, 1882, Am. Jour. Sci.

and Arts, 3d ser., vol. 23, p. 477. [Ety.

spathe, spathe; *karia*, shrimp.] Cara-

pace in one piece, oblong elliptical,

convex, apical point near the focus of

the ellipse, from which point a cleft

extends backward, widening to the margin.

Type *S. emersoni*.

emersoni, Clarke, 1882, Am. Jour. Sci.

and Arts, 3d ser., vol. 23, p. 478, Port-

age Gr.

lutheri, Clarke, 1882, (Lisgocaris lutheri,)

Am. Jour. Sci. and Arts, 3d ser., vol.

23, p. 478, Ham. Gr.

SPHÆREXOCHUS, Beyrich, 1845, Enber einige

Bohm. Tril., p. 21. [Ety. *sphaira*, ball;

exochos, prominent.] Cephalic shield

very convex, almost globular; cheeks

not scrobiculate; glabella nearly spher-

ical, three furrows on each side, two

upper obscure, lower strong and curved

down to the neck furrow; eyes faceted

minutely; facial suture ending on the

external margin near the angles, in

front continuous and submarginal;

thorax 11 joints; pygidium 3 segments,

free at their ends; labrum subtriangular,

with a marginal furrow; no rostral

shield. Type *S. mirus*.

canadensis, Billings, 1866, Catal. Sil. Foss.

Antic., p. 64, Anticosti Gr.

mirus, not American.

FIG. 1060.—*Sphaerexochus*
parvus. Upper and side
view of glabella.

parvus, Billings,

1865, Pal. Foss.,

vol. 1, p. 180,

Chazy or Black

Riv. Gr.

romingeri, Hall,

1867, 20th Rep.

N. Y. St. Mus.

Nat. Hist., p. 425, Niagara Gr.

SPHÆROCORYPHE, Angelin, 1852, Palaeontolo-

gia Scandinavica. [Ety. *sphaira*, ball;

koryphe, top of the head.] Cephalic

shield convex, genal

glabella spheroidal an-

teriorly, two lateral

furrows; eyes promi-

nent; facial sutures

cut the lateral mar-

gins posteriorly; tho-

rax with ten segments,

axial lobe narrower

than lateral lobes;

pleuræ terminate in

short spines; pygidium

composed of three

segments, anterior one

produced in two

spines. Type *S. gran-*

ulata.

robusta, Walcott, 1875,

Cin. Quar. Jour. Sci.,

vol. 2, p. 273, Trenton

Gr.

salteri, Billings, 1866, Catal. Sil. Foss.

Antic., p. 63, Anticosti Gr.

FIG. 1061.—*Sphaerocoryphe*
robusta.

- STROBILEPS**, Clarke, 1888, Pal. N. Y., vol. 7, p. 63. [Ety. *strobilos*, cone-shaped; *lepis*, scale.] Capitulum composed of four vertical ranges of plates having in general a trihedral form, but varying in size and contour; each plate articulated with or overlapping the next preceding; anterior extremity terminated by a large, circular, conical plate; plates thick and ornamented. Type *S. spinigera*.
- spinigera*, Clarke, 1888, Pal. N. Y., vol. 7, p. 112, Ham. Gr.
- STYLONURUS**, Page, 1850, Geological Text Book, p. 190. [Ety. *stylos*, a mast or spar; *oura*, tail.] General form like *Eurypterus*, but distinguished by the peculiar development of the two posterior foot pairs; these are alike, long, thin, and consist of 9 segments, of which the two last form a small claw; the posterior pair reach to the middle of the long posterior spine. Type *S. powriei*.
- excelsior*, Hall, 1884, 36th Rep. N. Y. St. Mus. Nat. Hist., p. 77, Catskill Gr.
- SYMPHYRUS**, Goldfuss, 1843, Neues Jahrb. f. Mineral. [Ety. *symphysis*, growing together; *oura*, tail.] Elliptical; genal angles rounded; cephalic shield semicircular, convex; glabella convex, subquadrate, smooth, no lateral furrows; eyes lunate; facial sutures arching in front of the glabella, and cutting the posterior part of the cephalic shield near the genal angles; 8 thoracic segments; pygidium somewhat semicircular, no segments, border flattened. Type *S. laeviceps*.
- goldfussi* (?), Walcott, 1885, Monogr. U. S. Geo. Sur., vol. 8, p. 96, Trenton Gr. This species is founded on a glabella with fixed cheeks, and probably it does not belong to this genus.
- TELEPHUS**, Barrande, 1852, Syst. Sil. Boh. [Ety. mythological name.] Glabella strongly convex, margined by deeply impressed bow-shaped furrows; neck segment tumid; cheeks subtriangular, or subrescentiform, small posteriorly, wider in front; pygidium small, strongly convex, hemispherical, margin tumid, axis with three segments. Type *T. fractus*. Only recognized in America by fragments of glabella.
- americanus*, Billings, 1865, Pal. Foss., vol. 1, p. 291, Quebec Gr.
- TERATASPIS**, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 223. [Ety. *teras*, marvel; *aspis*, shield.] Distinguished from *Acidaspis*, which it resembles by the prominent ellipsoidal frontal lobe of the glabella, posterior spines of the lateral lobes and nodes of the occipital ring, and from *Lichas* by the spines of the pygidium being themselves bearers of lateral spines. Type *T. grandis*.
- grandis*, Hall, 1862, 15th Rep. N. Y. St. Mus. Nat. Hist., p. 82, and Illust. Devon.
- Foss., pl. 17, (*Lichas grandis*), Schuchardte grit.
- erliops*, Hall, 1863, (*Lichas erliops*), 16th Rep. N. Y. St. Mus. Nat. Hist., p. 226, and Illust. Devon. Foss., pl. 19, 1 p. Held. Gr.
- Thaleops*, Conrad, syn. for *Iliaenus*.
- ovata*, see *Iliaenus ovatus*.
- THIRATHRELLA**, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 177. [Ety. diminutive of *Triarthrus*.] Glabella elongate, semioval, with the fixed cheeks wide and spreading in the posterior limb, and very narrow in front, an obscure indentation at the margin; general expression like *Triarthrus*. Type *T. auroralis*.
- auroralis*, Hall, 1863, 16th Rep. N. Y. St. Mus. Nat. Hist., p. 177, Potsdam Gr.
- TRIAURUS**, Green, 1832, Monograph of Trilobites, p. 87. [Ety. *triarthrus*, three-jointed.] Subelliptical; cephalic shield somewhat semicircular or sublunate; glabella moderately convex, sides straight, rounded in front, deeply trilobate on each side by the lateral furrows, with a prominent occipital groove near the base, and occipital ring, from the center of which a spine sometimes arises; eyes small and placed on the antero-lateral margin; free cheeks forming a narrow rim; thorax with from 13 to 16 articulations; central axis convex, wider than the lateral lobes; pygidium with 5 to 7 segments in the axis and one or two less in the lateral lobes. Type *T. becki*.
- becki*, Green, 1832, Monograph of Trilobites, p. 87, and Pal. N. Y., vol. 1, p. 237, Utica Slate Gr.
- canadensis*, Smith, 1861, Can. Jour., vol. 6, p. 275, Utica Slate Gr.
- fischeri*, see *Atops fischeri*.
- glaber*, Billings, 1859, Can. Nat. and Geol., vol. 4, p. 382, and Can. Geol., p. 202, Utica Slate Gr.
- spinous*, Billings, 1857, Rep. of Progr. Geo. Sur. Can., Fig. 1062.—Tri. p. 340, and Can. Geol., p. arthrus becki. 202, Utica Slate Gr.
- Trimerus**, syn. for *Homalonotus*.
- delphinocephalus*, see *Homalonotus delphinocephalus*.
- jacksoni*, see *Homalonotus jacksoni*.
- TRINUCLEUS**, Lhwyd, (or, as he spelt it, Lhwydd,) 1698, Phil. Trans., vol. 20, p. 279. [Ety. *trinucleus*, three-kerneled.] Cephalic shield highly convex, a wide border impressed with several rows of deep puncta and posterior angles, terminating in spines; glabella pyriform, pointed behind, no lateral furrows; cheeks convex; no eyes or facial sutures; neck furrow distinct; thorax with six articulations, axis narrow,

as grandis,) Scho-
nichas eriopis,) 16th
Nat. Hist., p. 226,
Foss., pl. 19, Up.

llaenus.
tus.
16th Rep. N. Y. St.
77. [Ety. diminutive.
Glabella elongate,
fixed cheeks wide
the posterior limb,
front, an obscure
margin; general ex-
rus. Type T. auro-

16th Rep. N. Y.
st., p. 177, Pots-

32, Monograph of
Ety. triarthrus, three-
falc; cephalic shield
ular or subulate;
ly convex, sides
front, deeply trilo-
by the lateral fur-
cent occipital groove
occipital ring, from
a spine sometimes
and placed on the
rgin; free cheeks
rim; thorax with
ticulations; central
r than the lateral
ith 5 to 7 segments
e or two less in the
e T. becki.

Mon-
p. 87,
l. 1, p.
1861,
p. 275,

heri.
p. Can.
ol. 4, p.
ol., p.

57, Rep.
p. Can., Fig. 1002.—Tri-
eol., p. arthrus becki.

halonotus.
Homalonotus delphi-

notus jacksoni.
(or, as he spelt it,
hil. Trans., vol. 20, p.
leus, three-kerneled.)
ighly convex, a wide
with several rows of
posterior angles, ter-
es; glabella pyriform,
no lateral furrows;
no eyes or facial su-
ow distinct; thorax
ations, axis narrow,

convex; side lobes wide, flat, straight,
pleural groove not reaching the margin;
pygidium subtriangular, margin de-
flected, axis conical, about six furrows;
side lobes flat, with about the same
number of furrows. Type T. concentri-
cus.

FIG. 1063.—Trinucleus concentricus.

bellulus, Ulrich, 1878,
Jour. Cin. Soc. Nat.
Hist., vol. 1, p. 99. The
young of T. concentri-
cus.
concentricus, Eaton, 1832,
(Nuttainia concentrica.)
Geo. Text-Book, p. 128,
and Pal. N. Y., vol. 1,
p. 249, Trenton to Hud.
Riv. Gr.

TROPIDOCARIS, Beecher, 1884, Geo. Sur. Pa.,
vol. PPP, p. 15. [Ety. tropis, a keel;
kari, a shrimp.] Carapace bivalve,
semioval or semielliptical, obliquely
truncated behind; valves about twice
as long as wide, having one or more
longitudinal ridges; cephalic region
indicated by elevations at the anterior
end; optic node situate on a ridge;
two segments of the abdomen. Type
T. bicarinata.

alternata, Beecher, 1884, Geo. Sur. Pa.,
vol. PPP, p. 19, Waverly Gr.
bicarinata, Beecher, 1884, Geo. Sur. Pa.,
vol. PPP, p. 16, Chemung Gr.
interrupta, Beecher, 1884, Geo. Sur. Pa.,
vol. PPP, p. 18, Chemung Gr.

TURRILEPAS, Woodward, 1865, Quar. Jour.
Geol. Soc., vol. 21, p. 480. [Ety. turris,
tower; lepas, scale.] Elongate, cone-
shaped bodies, composed of from 4 to
6 vertical ranges of scale-like, subtri-
angular plates covered with elevated
concentric lines; plates of middle
range convex and bearing a median
carina. Type T. wrightana.
cancellatus, Hall, 1888, Pal. N. Y., vol. 7,
p. 216, Up. Held. Gr.
devonicus, Clarke, 1882, (Plumulites de-
vonicus,) Am. Jour. Sci., 3d ser., vol.
24, p. 55, Ham. Gr.
flexuosus, Hall, 1888, Pal. N. Y., vol. 7,
p. 215, Up. Held. Gr.
foliatus, Hall, 1888, Pal. N. Y., vol. 7, p.
218, Ham. Gr.
gracillimus, Ringueberg, 1888, (Plumu-
lites gracillimus,) Proc. Acad. Nat. Sci.
Phil., p. 136, Niagara Gr.
newberryi, Whitfield, 1882, (Plumulites
newberryi,) Ann. N. Y. Acad. Sci., vol.
2, p. 217, Portage Gr.
nitidulus, Hall, 1888, Pal. N. Y., vol. 7, p.
218, Ham. Gr.
squama, Hall, 1888, Pal. N. Y., vol. 7, p.
217, Ham. Gr.
tener, Hall, 1888, Pal. N. Y., vol. 7, p.
219, Ham. Gr.

ZACANTHOIDES, Walcott, 1888, Am. Jour.
Sci., 3d ser., vol. 38, p. 165. Proposed
to receive Olenoides levis, O. spinosus,
O. flagricaudatus, and O. typicalis,
but not defined.

CLASS ARACHNIDA.

THE animals, forming the class Arachnida, include the spiders, scorpions, and many offensive parasites and microscopic forms. They are generally possessed of four pairs of legs attached to the anterior division of the body, but have no antennae. The Paleozoic fossils are nearly all referred to an extinct order, Anthracomarti, but a few are referred to the living orders, Pedipalpi and Scorpiones. The Pedipalpi have arm-like prehensile organs, terminating in a movable claw, annulated abdomen, and long flexible limbs. They inhabit tropical countries, and have a forbidding aspect. The Scorpiones have large palpi or arm-like prehensile organs, terminated by a pair of nippers, and an elongated, tail-like abdomen, which ends in a sharp claw; and when the animal is in motion, this is carried over the back in a threatening manner. The poison glands are situated at the base of the claw, and when the animal stings, a portion of the venom is thrown into the wound. The scorpions are inhabitants of tropical countries. The Order Anthracomarti is defined as follows: Body more or less depressed; cephalothorax and abdomen distinctly separable; cephalothorax frequently made up in large part of pedigerous segments,

more or less wedge-shaped, and visible above as well as below, the arrangement of which corresponds to that of the coxæ. The abdomen forms a single mass, and is composed of a variable number of visible segments, ranging from four to nine. Palpi not much longer than the legs, simply terminated.

ORDER ANTHRACOMARTI.

FAMILY ARCHITARIDÆ.—*Anthracomartus*, *Architarbus*, *Geraphrynus*.

FAMILY ARTHROLYCOSIDÆ.—*Arthrolycosa*.

FAMILY POLIOCHERIDÆ.—*Poliochera*.

ORDER PEDIPALPI.

FAMILY GERALINURIDÆ.—*Geralinura*.

ORDER SCORPIONES.

FAMILY EOSCORPIONIDÆ.—*Eoscorpium*, *Mazonia*.

ANTHRACOMARTUS, Karsch, 1882, *Zeitschr. deutsch. geol. Gesellsch.*, p. 556. [Ety. *anthrax*, coal; *Martos*, proper name.] Cephalothorax quadrate, the front square or scarcely convex, about half the size of the abdomen; coxæ radiating from a broad triangular sternal plate, the base of which forms the posterior margin; sides of body constricted so as to show a distinct though slight separation of cephalothorax and abdomen; abdomen orbicular, composed of seven segments of similar length throughout. Type A. *volkelianus*.

pustulatus, Scudder, 1884, *Proc. Am. Acad. Arts and Sci.*, p. 13, *Low. Coal Meas.*

trilobitus, Scudder, 1884, *Proc. Am. Acad. Arts and Sci.*, p. 13, *Coal Meas.*

ARCHITARBUS, Scudder, 1868, *Geo. Sur. Ill.*, vol. 3, p. 568. [Ety. *archaios*, ancient; *tarbos*, object of alarm.] Cephalothorax orbicular, broadly rounded in front, much smaller than the abdomen, but not separated from it by a marked lateral constriction; coxæ radiating from a central pit; abdomen oval, composed of nine segments, of which those on the basal half are very much shorter than the others, and on the dorsal surface are forced still more closely together by the large post-thoracic plate; no abdominal append-

FIG. 1064.—*Architarbus rotundatus*.

ages. Type A. *rotundatus*.
rotundatus, Scudder, 1868, *Geo. Sur. Ill.*, vol. 3, p. 568, *Coal. Meas.*

ARTHROLYCOSA, Harger, 1874, *Am. Jour. Sci. and Arts*, 3d ser., vol. 7, p. 219.

[Ety. *arthron*, a joint; *lykos*, a spider.] Cephalothorax orbicular, twice as large as the abdomen; Coxæ radiating from a central pit; abdomen oval much narrower at the base than the cephalothorax, with no longitudinal sculptur-

FIG. 1065.—*Arthrolycosa antiqua*.

ing, and composed of seven segments; no abdominal appendages. Type A. *antiqua*.
antiqua, Harger, 1874, *Am. Jour. Sci. and Arts*, 3d ser., vol. 7, p. 219, *Coal Meas.*

EOSCORPIUS, Meek & Worthen, 1868, *Am. Jour. Sci. and Arts*, 2d. ser., vol. 46, p. 25, and *Geo. Sur. Ill.*, vol. 3, p. 560. [Ety. *eos*, dawn; *scorpius*, a scorpion.] Cephalothorax quadrangular, somewhat wider behind than long; mesial and lateral furrows between which the surface bears granules; mandibles stout, without teeth or serrations; movable finger curved and sharp at the point; legs stout, divisions long; abdomen twice as long as cephalothorax; segments gradually increase in size to the sixth, while the seventh and last is 2½ times as long as the sixth, but rapidly contracts, and is truncated for the attachment of the tail; the anterior margin of each of the first six segments is rounded; the three tail segments preserved are stout, oblong, and covered with granules; the comb-like organ shows 11 or 12 divisions. Type E. *carbonarius*.

arrangement of
angle mass, and is
om four to nine.

eraphrynus.

, 1874, Am. Jour.
ser., vol. 7, p. 219.

FIG. 1065.—*Arthrolycosa antiqua*.

no abdominal ap-
antiqua.

74, Am. Jour. Sci.
vol. 7, p. 219, Coal

Worthen, 1868, Am.
2d. ser., vol. 46, p.
Ill., vol 3, p. 560.
scorpius, a scorpion.]
quadrangular, some-
than long; mesial
between which the
anules; mandibles
or serrations; un-
nd sharp at the point;
ns long; abdomen
as cephalothorax;
increase in size to
e seventh and last
g as the sixth, but
nd is truncated for
the tail; the anterior
the first six segments
three tail, segments
oblong, and covered
the comb-like organ
divisions. Type E.

carbonarius, Meek & Worthen, 1868, Am.
Jour. Sci. and Arts, 2d ser., vol. 46, p.
24, and Geo. Sur. Ill., vol. 3, p. 560,
Coal Meas.

FIG. 1066.—*Eoscorpius carbonarius*. Natural size;
a, body segment enlarged; c, comb; d, same
enlarged; m, mandibles; p, pincers.

GERALINURA, Scudder, 1884, Proc. Am.
Acad. Arts and Sci., p. 13. [Ety. *geras*,
old; *linon*, linen; *oura*, tail.] Cephalo-
thorax ovate, the front rounded, one-
third as broad as hinder portion; palpi
large and robust, with interior spines;
first two pairs of legs slender, the
hinder stout and broad; abdomen
composed of nine joints, the basal three
rather short, the others subequal and
longer. Type *S. carbonaria*.

carbonaria, Scudder, 1884, Proc. Am.
Acad. Arts and Sci., p. 13, Coal Meas.

GERAPHRYNUS, Scudder, 1884, Proc. Am.
Acad. Arts and Sci., p. 13. [Ety. *geras*,
old; *Phrynus*, a genus.] Cephalothorax
fusiform, angulated in front, nearly as
large as the abdomen; coxae radiating
from a median line; palpi slenderer
than the legs, longer than the cephalo-
thorax, springing from its extreme front,
and of uniform size throughout; ab-
domen subfusiform, composed of nine
segments, rounded behind, with no
constriction at the base; a large tri-
angular post-thoracic plate, crowding
the middle of the first five short seg-
ments out of a straight transverse line;

readily distinguished from *Architarbus*
by its produced and angulate cephalo-
thorax. Type (*E. carbonarius*).

carbonarius, Scudder, 1884, Proc. Am.
Acad. Arts and Sci., p. 13, Coal Meas.

MAZONIA, Meek & Worthen, 1868, Geo. Sur.
Ill., vol. 3, p. 563. [Ety. proper name.]
Cephalothorax moderately convex, sub-
quadrangular; anterior lateral margins
rounded, and anterior margin truncated
on each side of a small mesial trian-
gular projection; mesial furrow extends
forward from the posterior margin,
widening and deepening to the front,
where it occupies one-third of the
breadth, and is partly filled by the oc-
ciferous prominence, which bears on
each side a large eye; eyes circular,
convex, arranged for looking obliquely
forward, outward, and upward; seven
or eight abdominal segments, the last
one truncated for the tail. Type *M.*
woodana.

woodana, Meek & Worthen, 1868, Geo.
Sur. Ill., vol. 3, p. 563, Coal Meas.

POLIOCHERA, Scudder, 1884, Proc. Am. Acad.

Arts and Sci., p. 13. [Ety. *polios*, hoary;
cheras, to be bereft.] Cephalothorax
scarcely longer than broad, slightly
narrowing anteriorly, the front square;
coxae radiating from a median line;
legs stout, moderately long; abdomen
full, at base as broad as the cephalo-
thorax, broadening slightly behind, fully
rounded, composed of four segments,
the first segment about one-third the
length of the others, which are equal;
no abdominal appendages. Type *P.*
punctulata.

punctulata, Scudder, 1884, Proc. Am.
Acad. Arts and Sci., p. 13, Coal Meas.

PROSCORPIUS, Whitfield, 1885, Bull. Am. Mus.

Nat. Hist., vol. 1, p. 183. [Ety. *pro*, be-
fore; *scorpius*, a genus.] Cephalothorax
with large dorsal eye-lobe; eyes small,
one on each side of the median line;
lateral eyes on ridges, as in living scor-
pions; sixth ventral segment of the
preabdomen, counting from behind,
large, equal in length and breadth to
the corresponding dorsal segment; an-
terior walking limb terminating in a
bifid claw; postabdomen not reversed
as in living scorpions. Type *P. osborni*.
Good authorities say this is merely an
Eurypterus, with no affinity or resem-
blance to a scorpion. With this view
the author coincides.

osborni, Whitfield, 1885, Bull. Am. Mus.
Nat. Hist., vol. 1, p. 184, Waterlime Gr.
Synonym, probably, for *Eurypterus*
remipes.

CLASS MYRIAPODA.

THE animals composing the Class Myriapoda are elongated, and composed of numerous segments, all of which are substantially alike except the first and last. The articulations of the body each bear one or two pairs of jointed legs. The common centipede and long-jointed worms, with numerous legs, found in damp places and on trunks of trees, some of which coil up when alarmed, are examples. Only a few Palæozoic fossils are referred to this Class, and these belong to extinct Orders.

ORDER ARCHIPOLYPODA.

FAMILY ARCHIULIDÆ.—Archilus, Trichiulus, Xylobius.

FAMILY EUPHOBERIDÆ.—Acantherpestes, Amynilespes, Anthracerpes, Eileticus, Euphoberia.

FAMILY UNCERTAIN.—Archæoscolex.

ORDER PROTOSYNGNATHA.

FAMILY PALÆOCAMPIDÆ.—Palæocampa.

ACANTHERPESTES, Meek & Worthen, 1868, Geo. Sur. Ill., vol. 3, p. 559. [Ety. *akantha*, a spine; *erpestes*, a creeper.] Spines bifurcate at tip, and arranged in dorsal, pleurodorsal, and lateral rows; segments three, or more than three times as broad as long. Type A. major.

major, Meek & Worthen, 1868, (Euphoberia major,) Am. Jour. Sci. and Arts, 2d. ser., vol. 46, p. 26, and Geo. Sur. Ill., vol. 3, p. 558, Coal Meas.

FIG. 1067.—*Acantherpestes major*. Fragment. s, spine; n, base of spine.

AMYNILESPES, Scudder, 1885, in Zittel's Handbuch der Pal., p. 729. [Ety. *amuno*, to keep off; *iluspaomai*, to crawl.] Spines simple, arranged in dorsolateral

rows; segments four times as broad as long. Type A. wortheni.

wortheni, Scudder, 1885, in Zittel's Handbuch der Pal., vol. 2, p. 729, Coal Meas.

FIG. 1068.—*Amynilespes wortheni*.

ANTHRACERPES, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 51. [Ety. *anthrax*, coal; *erpo*, to creep, in allusion to its carboniferous age and probable habits.] Founded upon an articulated body of nineteen segments and part of another. The last segment terminates in three or four short, slender, hair-like or spine-like appendages. Below the middle of each segment there is a small prominence, marking the spiracles, or breathing apertures, which pertain to the Myriapoda. Type A. typus.

typus, Meek & Worthen, 1865, Proc. Acad. Nat. Sci. Phil., p. 51, and Geo. Sur. Ill., vol. 2, p. 409, Coal Meas.

ARCHÆOSCOLEX, Matthew, 1888, Trans. Roy. Soc. Can., p. 59. [Ety. *archaios*, ancient; *scolex*, worm.] Cylindrical, tapering behind the middle, and also at the three anterior segments; head small, somewhat conical; thorax of three joints, increasing in width back-

d, and composed of
t the first and last,
ted legs. The com-
und in damp places
re examples. Only
belong to extinct

Anthracerpes, Ei-

ur times as broad as
rtheni.

1068.—*Amyntilipes*
wortheni.

roc. Acad. Nat. Sci.
[Ety. *anthrax*, coal;
in allusion to its car-
e and probable hab-
ed upon an articulated
teen segments and
her. The last seg-
ates in three or four
, hair-like or spine-
es. Below the mid-
segment there is a
ence, marking the
breathing apertures,
a to the Myriapoda.
s.
Worthen, 1865, Proc.
Sci. Phil., p. 51, and
, vol. 2, p. 409, Coal

Matthew, 1888, Trans.
, p. 59. [Ety. *archaios*,
, worm.] Cylindrical,
the middle, and also
rior segments; head
conical; thorax of
easing in width back-

ward, but decreasing in length; limbs
tapering, posterior pair as long as the
thorax, and larger and stronger than
those in front; abdomen of eleven vis-
ible seg-
ments, those
at the an-
terior end
wider and
those of the
middle long-
er; basal seg-
ment has an
oblong scar near the posterior edge;
crust chitinous. Type *A. corneus*.

corneus, Matthew, 1888, Trans. Roy. Soc.
Can., p. 59, Devonian.

FIG. 1069.—*Archilulus*
xylobioides. Anter-
ior part enlarged.

ARCHILULUS, Scudder,
1868, Mem. Bost.
Soc. Nat. Hist.,
vol. 2, p. 231, and
Acad. Geol., p.
496. [Ety. *ar-*
chaïos, ancient;
ioulos, wood-louse.]
Segments entire,
varying much in

relative proportions, but generally from
two to three times broader than long,
furnished with only a few papillae, per-
haps supporting spiny hairs. Type *A.*
xylobioides.

xylobioides, Scudder, 1868, Mem. Bost.
Soc. Nat. Hist., vol. 2, p. 236, and Acad.
Geol., p. 496, Coal Meas.

EILETICUS, Scudder, Mem. Bost. Soc. Nat.
Hist. [Ety. *eiletikos*, rolling one's self.]
No spines, but large, low tubercles, ser-
ially arranged; segments few, less
than twice as broad as long. Type *E.*
anthracinus.

anthracinus, Scudder, Mem. Bost. Soc.
Nat. Hist., Coal Meas.

EUPHOBERIA, Meek & Worthen, 1868, Am.
Jour. Sci. and Arts, 2d ser., vol. 46, p.
26. [Ety. *eu*, very; *phoberos*, formid-
able.] Head semicircular; body long,
slender, very slightly tapering, and
terminating abruptly; segments sev-
enty-five or more on the ventral side

FIG. 1070.—*Euphoberia armigera*. Part of a large
specimen.

and half as many on the dorsal; dorsal
half of the segments rounded, and each
supporting three or four spines, curved
slightly backward, and arranged in

rows on the back, spines spinuliferous;
ventral half of the segments each bears
a pair of small slender-jointed legs.
Type *E. armigera*.

FIG. 1071.—*Euphoberia armigera*. Entire specimen.

anguilla, Scudder, 1880, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 177, Coal Meas.
armigera, Meek & Worthen, 1868, Am.
Jour. Sci. and Arts, 2d ser., vol. 46, p.
26, and Geo. Sur. Ill., vol. 3, p. 556,
Coal Meas.

FIG. 1072.—*Euphoberia armigera*. A, part of an
individual; B, enlarged surface pitting.

carri, Scudder, 1880, Mem. Bost. Soc. Nat.
Hist., vol. 3, p. 171, Coal Meas.

fiabellata, Scudder, 1880, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 174, Coal Meas.

granosa, Scudder, 1880, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 168, Coal Meas.

horrida, Scudder, 1880, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 158, Coal Meas.

major, see *Acantherpestes major*.

PALÆOCAMPA, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci. Phil., p. 52. [Ety.
palaïos, ancient; *kampe*, a caterpillar.]
Head small; segments ten, similar, sub-
equal, and each bearing a pair of stout
clumsy legs, and four bunches of cylin-
dric needles or spines; bunches seated
on mammillae, and arranged in dorso-
pleural and lateral rows, needles or
spines, exceedingly slender, scarcely ta-
pering, blunt at tip, and longitudinally
serrated. Type *P. anthrax*.

anthrax, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci. Phil., p. 52, and Geo.
Sur. Ill., vol. 2, p. 410, Coal Meas.

TRICHIULUS, Scudder, 1884, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 290. [Ety. *trichos*,
hair; *ioulos*, wood-louse.] Segments en-
tire, from three to five times broader
than long, closely covered with pa-
pillae, arranged in definite series longi-
tudinally, and transversely supporting
long, sweeping hairs. Type *T. villosus*.
ammonitiformis, Scudder, 1884, Mem.
Bost. Soc. Nat. Hist., vol. 3, p. 292, Coal
Meas.

nodulosus, Scudder, 1884, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 292, Coal Meas.

villosus, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 291, Coal Meas.
Xylobius, Dawson, 1860, Quar. Jour. Geo. Soc., vol. 16, p. 268. [Ety. *xylobius*, living in wood.] Body crustaceous, cylindrical, elongate, rolling spirally; segments thirty or more, anterior ones smooth, posterior ones furrowed; legs small, numerous; labrum quadrilateral, divided by notches or joints into three portions; mandibles two-jointed, last ovate and pointed; eyes ten or more on each side. Type *X. sigillarie*.
dawsoni, Scudder, 1888, Mem. Bost. Soc. Nat. Hist., vol. 2, p. 236, and Acad. Geol., p. 496, Coal Meas.

fractus, Scudder, 1868, Mem. Bost. Soc. Nat. Hist., vol. 2, p. 234, and Acad. Geol., p. 496, Coal Meas.
mazonius, Zittel, 1885, Handbuch der Pal., p. 730, Coal Meas.
sigillarie, Dawson, 1860, Quar. Jour. Geo. Soc., vol. 16, p. 271, Coal Meas.
similis, Scudder, 1868, Mem. Bost. Soc. Nat. Hist., vol. 2, p. 234, and Acad. Geol., p. 496, Coal Meas.

FIG. 1073. — *Xylobius sigillarie*. a. Organ with palpus, pertaining to the mouth, enlarged.

CLASS INSECTA.

INSECTS are possessed of head, thorax, and abdomen. Three pairs of legs and one pair of antennæ belong to them in their perfect state. They are the highest and most complicated class of articulated animals, and abound almost everywhere. No living order, in this class, is known from the Palæozoic rocks. Indeed, the fossils consist almost wholly of fragments of wings showing little else than venation; but they have been studied by Scudder, until he has classified them into an Order and Families, to the general satisfaction of entomologists, and all must concede he has accomplished a very difficult task. The Order Palæodictyoptera, which includes the Orthopteroid and Hemipteroid Palæodictyoptera, has been defined as follows: Body generally elongated; mouth parts variously developed; antennæ filiform; thoracic joints subequally developed; legs moderately large; meso and metathoracic wings closely similar, equally membranous; the six principal veins always developed; the marginal simple, and forming the costal border; the mediastinal generally simple or with superior branches only; the other veins usually dichotomize; stout and well defined cross veins rare; membrane generally reticulate; wings in repose lying on the abdomen; the anal area of hind wings, though usually of great distal extension, never plaited, though sometimes broadly folded; abdomen usually long and slender, the last joint often furnished with simple articulated appendages.

ORDER PALÆODICTYOPTERA.

FAMILY GERARIDÆ.—*Adiphlebia*, *Gerarus*, *Megathentomum*, *Polyernus*.

FAMILY HEMERISTIDÆ.—*Chrestotes*, *Hemeristia*, *Lithentomum*.

FAMILY HOMOTHETIDÆ.—*Anthracothremma*, *Cheliphlebia*, *Didymophleps*, *Eucænus*, *Genentomum*, *Genopteryx*, *Gerapompus*, *Homothetus*.

FAMILY MYLACRIDÆ.—*Lithomylacris*, *Mylacris*, *Necymylacris*, *Paromylacris*, *Promylacris*.

FAMILY PALÆOBLATTINIDÆ.—*Archymylacris*, *Etblattina*, *Gerablattina*, *Oryctoblattina*, *Petrablattina*.

FAMILY PALÆOPTERINIDÆ.—*Aethophlebia*, *Dieconeura*, *Miamia*, *Propteticus*.

FAMILY PALÆEPHEMERIDÆ.—*Ephemerites*, *Geraphemera*, *Platephemera*.

8, Mem. Bost. Soc.
p. 234, and Acad.
Meas.

FIG. 1073.—*Xylobius sigillarius*. a. Organ with palpus, pertaining to the mouth, enlarged.

Acad. Geol., p. 496,

ee pairs of legs and
ey are the highest
almost everywhere.
ocks. Indeed, the
le else than neuro-
sified them into an
gists, and all must
r Palæodictyoptera,
optera, has been de-
eriously developed;
erately large; meso-
; the six principal
costal border; the
he other veins usu-
membrane generally
rea of hind wings,
sometimes broadly
furnished with sim-

um, Polyernus.
um.

ia, Didymophleps,
omothetus.

acris, Paromylacris,

ina, Gerablattina,

Miamia, Propteticus.

Platephemera.

FAMILY PHTHANOCORIDÆ.—Phthanocoris.

FAMILY PROTOPHASMIDÆ.—Haplophlebium, Paolia, Titanophasma.

FAMILY XENONEURIDÆ.—Geroneura, Xenoneura.

FAMILY UNCERTAIN.—Archegogryllus, Dyscritus.

SUPPOSED INSECT TRAILS.—Haplotichnus, Plangtichnus, Treptichnus.

ADIPHLEBIA, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 345. [Ety. a, privative; *dis*, double; *phlebion*, vein.] Body rather stout; wings rather broad; all the nervules simple, arising from their stems near the base of the wings; subparallel and longitudinal. Type A. lacoana.

lacoana, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 345, Coal Meas.

ATHROPHLEBIA, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 338. [Ety. *athro*, strange; *phleps*, a vein.] Interno-median vein terminating before the middle of lower border, emitting a single main branch, beyond its middle which is superior, and which, with median fork of externo-median and larger part of main scapular branch, form a continuous adventitious vein crossing principal nervules of the wing; ultimate offshoots of externo-median vein arise indifferently from the main vein and the principal branch, and are parallel and similar to the offshoots of the veins above. Type A. singularis.

singularis, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 338, Coal Meas.

ANTHRACOTHREMA, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 327. [Ety. *anthrax*, coal; *thremma*, reared.] Body stout; prothorax several times broader than long; wings subequal and elongated; scapular vein arcuate and nearly reaching the tip; externo-median vein with numerous parallel branches, mostly simple. Type A. robusta.

robusta, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 327, Coal Meas.

ARCHEGOGRYLLUS, Scudder, 1868, Proc. Bost. Soc. Nat. Hist., vol. xi, p. 401. [Ety. *archegos*, first in time; *gryllus*, a cricket.] Relations not clearly understood. Type A. priacus.

priacus, Scudder, 1868, Proc. Bost. Soc. Nat. Hist., vol. 11, p. 401, and Mem. Bost. Soc. Nat. Hist., vol. 3, p. 323, Coal Meas.

ARCHIMYLACRIS, Scudder, 1868, Acad. Geol., p. 388. [Ety. *arche*, beginning; *My-lacris*, cockroach.] Mediastinal area com-

paratively short; scapular terminat- ing below the tip, and with the ext- ern o - m e d i a n, which is com- paratively small, occupying less than half the

wing; internomedian vein compara- tively long. Type A. acadicum.

FIG. 1074.—*Archimylacris acadicum*.

acadicum, Scudder, 1868, Acad. Geol., p. 388, Coal Meas.

parallelum, Scudder, 1879, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 85, Coal Meas.

BLATTINA, Burmeister, 1838, Handbuch der Entomologie. [Ety. *Blatta*, a cockroach.] A living genus of cockroaches, raised to the rank of a family, and by some naturalists to the rank of an order, to which the name Dictyoptera has been applied. It is not a Palæozoic genus.

bretonensis, see Mylacris bretonense.

fascigera, see Gerablattina fascigera.

heeri, see Mylacris heeri.

sepulta, see Petrablattina sepulta.

venusta, see Etoablattina venusta.

CHELIPHIL BIA, Scudder, 1884, Mem. Bost. Soc.

Nat. Hist., vol. 3, p. 328. [Ety. *cheli*, forked; *phlebion*, vein.] Body rather slender, but wings large and coarse, without cross veins, interno-median vein extending far toward the tip of the wing with many oblique branches. Type C. elongata.

carbonaria, Scudder,

1884, Mem. Bost.

Soc. Nat. Hist., vol. 3, p. 328, Coal Meas. Fig. 1075.—Chelli-

3, p. 328, Coal Meas. phlebia carbonaria.

elongata, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 328, Coal Meas.

CHRESTOTES, Scudder, 1868, Geo. Sur. Ill., vol. 3, p. 567. [Ety. *chrestotes*, good of its kind.] Wings short, broad, well

FIG. 1076.—*Chrestotes lapidea*.

rounded; vena scapularis throws several branches downward, commencing before the middle of the wing, and with its branches occupies the upper two-

fifths of the upper wing, and perhaps more of the lower; remainder of wing occupied by the longitudinally divaricating branches of the next two veins; anal area in upper wing distinctly set off at the basal portion of the wing. Type *C. lapidea*.

lapidea, Scudder, 1868, *Geo. Sur. Ill.*, vol. 3, p. 567, *Coal Meas.*

DIDYMOPLERS, Scudder, 1878, *Proc. Bost. Soc. Nat. Hist.*, vol. 19, p. 300. [Ety. *didymos*, double; *phleps*, vein.] All the veins and branches above the intermedian longitudinal and nearly parallel; nearly all the lower half of the wing being occupied by the oblique branches of the intermedian vein. Type *D. contusa*. *contusa*, Scudder, 1878, (*Termescontusus*), *Bost. Soc. Nat. Hist.*, vol. 19, p. 300, and *Mem. Bost. Soc. Nat. Hist.*, vol. 3, p. 330, *Coal Meas.*

DIEONEURA, Scudder, 1884, *Mem. Bost. Soc. Nat. Hist.*, vol. 3, p. 336. [Ety. *dieko*, to extend through; *neuron*, a vein.] Externo-median vein simple; intermedian vein important, arcuate, extending far toward the extremity of the lower margin. Type *D. rigida*.

arcuata, Scudder, 1884, *Mem. Bost. Soc. Nat. Hist.*, vol. 3, p. 336, *Coal Meas.*

rigida, Scudder, 1884, *Mem. Bost. Soc. Nat. Hist.*, vol. 3, p. 336, *Coal Meas.*

DYSCRITUS, Scudder, 1868, *Lond. Geo. Mag.*, vol. 5, p. 176. [Ety. *dyscritos*, hard to determine.] Founded on a fragment of the middle part of a wing without proper definition. Type *D. vetustus*.

vetustus, Scudder, 1868, *Lond. Geo. Mag.*, vol. 5, p. 176, *Devonian*.

ENCÆNUS, Scudder, 1884, *Mem. Bost. Soc. Nat. Hist.*, vol. 3, p. 325. [Ety. *en*, very; *kainos*, new, strange.] Body stout, thoracic segments twice as broad as long; abdomen ovate; fore wings with the mediastinal vein straight, terminating before the apical third of the wing with numerous straight branches; scapular with similar branches ending half way between the mediastinal and the tip; externo-median important with distant branches. Type *E. ovalis*.

ovalis, Scudder, 1884, *Mem. Bost. Soc. Nat. Hist.*, vol. 3, p. 325, *Coal Meas.*

FIG. 1077.—*Ephemerites affinis*.

EPHEMERITES, Scudder, 1868, *Geo. Sur. Ill.*, vol. 3, p. 571. [Ety. *Ephemeris*, a

living genus.] The genus was not defined. It is probably a neuropteroid Palaeodictyoptera. Type *E. simplex*, *affinis*, Scudder, 1868, *Geo. Sur. Ill.*, vol. 3, p. 572, *Coal Meas.*

FIG. 1078.—*Ephemerites gigas*.

gigas, Scudder, 1868, *Geo. Sur. Ill.*, vol. 3, p. 571, *Coal Meas.*

primordialis, Scudder, 1878, *Proc. Bost. Soc. Nat. Hist.*, vol. 19, p. 248, *Coal Meas.*

FIG. 1079.—*Ephemerites simplex*.

simplex, Scudder, 1868, *Geo. Sur. Ill.*, vol. 3, p. 571, *Coal Meas.*

ETOBLOTTINA, Scudder, 1882, *Proc. Bost. Soc. Nat. Hist.*, vol. 21, p. 391. [Ety. *etos*, true; *Blattina*, a genus.] Mediastinal area comparatively short; scapular not reaching tip of wing and with the externo-median, which is comparatively large, occupying less than half the wing; intermedian vein comparatively long. Type *E. mazonana*.

FIG. 1080.—*Etoblattina primaeva*. from Saarbrück in Europe for comparison.

lesquereuxi, Scudder, 1870, *Mem. Bost. Soc. Nat. Hist.*, vol. 3, p. 67, *Coal Meas.*

genus was not de-
by a neuropteroid
Type *E. simplex*,
Geo. Sur. Ill., vol.

Geo. Sur. Ill., vol.
1878, Proc. Bost.
Soc. Nat. Hist., vol. 3, p. 248, Coal

merites simplex.

1868, Geo. Sur. Ill.,
Meas.

1882, Proc. Bost.
Soc. Nat. Hist., vol. 3, p. 391. [Ety.
z, a genus.] Media-
stinal vein short; scapu-
lar vein of wing and with
n, which is compara-
tively less than half
the median vein compar-
ed to *E. mazonana*.

E. primaeva. From
type for comparison.

1879, Mem. Bost.
Soc. Nat. Hist., vol. 3, p. 67, Coal

mazonana, Scudder, 1882, Proc. Bost. Soc.
Nat. Hist., vol. 21, p. 391, Coal Meas.

FIG. 1081.—*Etioblattina venusta*.

venusta, Lesque-
reux, 1860, (Blat-
tina *venusta*,)
Geo. Sur. Ark.,
vol. 2, p. 314, Coal
Meas.
Bost. Soc. Nat. Hist., vol. 3, p. 329.
[Ety. *genos*, race, kind; *entomon*, insect.]
Wings large, elongated with coarse ve-
nation and abundant cross veins; medi-
astinal vein very long, with numerous
branches to the costa; other branches
very distant and stout; the externo-m-
edian separated more widely than usual
from the scapular, especially in the
hind wing. Type *G. validum*.

validum, Scudder, 1884, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 329, Coal Meas.

FIG. 1082.—*Genentomum validum*.

GENOPTERYX, Scudder, 1884, Mem. Bost.
Soc. Nat. Hist., vol. 3, p. 327. [Ety.
genos, kind, race; *pteryx*, a wing.]
Interno-median vein with branches very
similar to those of the externo-median
vein, the outermost in close proximity
to the innermost branches of the latter.
Type *G. constricta*.

constricta, Scudder, 1884, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 327, Coal Meas.

GERABLATTINA, Scudder, 1879, Mem. Bost.
Soc. Nat. Hist., vol. 3, p. 110. [Ety.
geras, old; *Blattina*, a genus of insects.]
Mediastinal area comparatively long;
scapular and externo-median area to-
gether occupy less than half the wing,
the branches of both superior; interno-
median vein comparatively long. Type
G. balteata.

balteata, Scudder,

1879, Mem. Bost.

Soc. Nat. Hist., vol.

3, p. 110, and Perm.

or Up. Carb. Flora

of Pa., p. 104, Up.

Coal Meas. or Perm.

ian.

fascigera, Scudder, 1879, (Blattina *fascig-*

era.) Mem. Bost. Soc. Nat. Hist., vol.

3, p. 113, Coal Meas.

GEREPHEMERA, Scudder, 1868, Lond. Geo.
Mag., vol. 5, p. 175. [Ety. *geras*, old;
Ephemera, a genus of insects.] Founded
on the fragment of a tip of the wing.
Definition incomplete. Type *G.*
simplex.

FIG. 1083.—*Gerablattina balteata*.

simplex, Scudder, 1868, Lond. Geo. Mag.,
vol. 5, p. 175, Upper Devonian.

GERAPOMPUS, Scudder, 1884, Mem. Bost.
Soc. Nat. Hist., vol. 3, p. 326. [Ety.
geras, old; *pompus*, an escort.] Body
slender, the prothorax as long as broad;
fore wings well rounded, the media-
stinal arcuate like the costa, with infre-
quent simple branches; scapular end-
ing near the tip. Type *G. blattinoides*.
blattinoides, Scudder, 1884, Mem. Bost.
Soc. Nat. Hist., vol. 3, p. 326, Coal
Meas.

extensus, Scudder, 1884, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 326, Coal Meas.

GERARUS, Scudder, 1868, Mem. Bost. Soc.
Nat. Hist., vol. 3, p. 344. [Ety. *geraros*,
of stately bearing.] Body slender, taper-
ing anteriorly; wings slender; mediastinal
vein variable; branches of scapular
vein numerous, more or less longitudi-
nal, simple or forked, occupying much
more space than the branches of any
other vein. Type *G. danæ*.

danæ, Scudder, 1868, (Miami *danæ*,)
Geo. Sur. Ill., vol. 3, p. 566, Coal Meas.

FIG. 1084.—*Gerarus danæ*.

mazonus, Scudder, 1884, Mem. Bost.
Soc. Nat. Hist., vol. 3, p. 344, Coal Meas.
vetus, Scudder, 1884, Mem. Bost. Soc. Nat.
Hist., vol. 3, p. 344, Coal Meas.

GERONEURA, Matthew, 1888, Trans. Roy.
Soc. Can., p. 57. [Ety. *geros*, old; *neura*,
a vein.] Anterior wing of the body
elliptical elongate, venation strongly
marked, scapular ridge conspicuous;
mediastinal vein close to the scapular,
but curves outward at the extremity;
scapular vein and its branches cover a
triangular area terminating at the
apical end of the wing; main scapular
terminates near the end of the costal
edge; externo-median vein throws off
two branches, the first one stronger
than the main vein, and the second one
goes with a sinuosity toward the base
of the apical margin; nerves regular
and simple. Type *G. wilsoni*.

wilsoni, Matthew, 1888, Trans. Roy. Soc.
Can., p. 57, Lower Devonian.

HAPLOPHLEBIUM, Scudder, 1867, Can. Nat.
and Geo., 2d ser., vol. 3, p. 202, and
Proc. Bost. Soc. Nat. Hist., vol. 11, p.
150. [Ety. *haplos*, simple; *phlebia*,
a vein.] Wing with simple neurulation
and intercostal spaces filled with mi-
nute reticulations without any cross
veins; wing long and slender.
Type *H. barnesi*.

barnesi, Scudder, 1867, Can. Nat. and Geol. 2d ser., vol. 3, p. 202, and Acad. Geol., p. 386, Coal Meas.

FIG. 1085.—*Haptophlebitum barnesi*. A fern covers part of the wing.

longipeunius, Scudder, 1884, Proc. Amer. Acad., vol. 20, p. 172, Coal Meas.

HAPLOTICHNUS, n. gen. [Ety. *haplo*, plainness, simplicity; *ichnos*, track.] Simple, small, half-cylindrical trails running in any direction. Supposed to have been made by the larva or pupa of some paleodictyopterous insect. Type *H. indianensis*.

indianensis, n. sp. A simple half-cylindrical trail, needle-like in size, running in straight or crooked lines, or crossing itself. Found in the upper part of the Kaskaskia Group, at the Whetstone quarries in Orange County, Indiana.

FIG. 1086.—*Haptotichnus indianensis*.

The remains of insects found in the Palaeozoic rocks occur under such circumstances as to induce the belief they were more or less aquatic in their habits, and frequented swamps and shores of bays and inlets. The Whetstone quarries of Orange County, Indiana, are yellowish white, slaty mud-rocks resembling, in appearance, the Solenhofen slates, but coarser in texture. They are limited in extent, and may be fairly presumed to represent the muddy shore of some bay or internal sea of Subcarboniferous age. The slaty layers are covered more or less upon the upper surface with trail-furrows, and on the under surface with elevated lines, showing the trails were made in mud, which afterward hardened, and was then covered with a thin deposit of mud which was tracked and hardened and covered, and so on in one series after another throughout the whole thickness of the slaty deposit. Many of the living Dictyoptera are

aquatic in their habits in the larva and pupa state, and it is not until the perfect insect is about to emerge from the skin of the pupa that it creeps out of the water on the muddy shore or stones, or climbs the stems and leaves of aquatic plants, and from this position the imago springs into an aerial habitat. The trails on the Whetstone slates were evidently made by animals, and

all the evidence seems to indicate they were made by insects, though the evidence may not be either clear or conclusive in the latter respect. Under these circumstances the author has selected three common but distinct trails, and given them generic names; viz., *Haptotichnus*, *Plangtichnus*, and *Treptichnus*.

HEMERISTIA, Dana, 1864, Am. Jour. Sci. and Arts, 2d ser., vol. 37, p. 34. [Ety. *hemera*, day; *istia*, house.] Scapular branch strongly arcuate, at its base distant from the main stem, and at first taking the course of its basal offshoot. Type *H. occidentalis*.

occidentalis, Dana, 1864, Am. Jour. Sci. and Arts, 2d ser., vol. 37, p. 34, Coal Meas.

HOMOTHETUS, Scudder, 1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202. [Ety. *homo*, similar; *thetos*, placed.]

FIG. 1087.—*Homothetus fossilis*.

Mediastinal vein extremely long, scarcely surpassed by the scapular, and with scarcely any branches to the costa; externo-median vein with only a few branches in the outer fourth of the wing; interno-median vein similar to the last. Type *H. fossilis*.

fossilis, Scudder, 1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202, and Acad. Geol., p. 525, Upper Devonian.

Libellula, Linnaeus. Not a Palaeozoic genus. *carbonaria*, see *Cheliphebia carbonaria*.

LITHENTOMUM, Scudder, 1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202. [Ety. *lithos*, stone; *entomon*, an insect.] Main

FIG. 1088.—*Lithentomum hartii*.

scapular branch with a single, or at most two branches, which are almost wholly longitudinal. Type *L. hartii*.

abits in the larva and is not until the pupa that it emerges from the water the muddy shore or, or climbs the stems of aquatic plants, and from this the imago springs to an aerial habitat. Trails on the Whetstones are evidently made by animals, and seem to indicate they are insects, though the trails are either clear or rather respect. Under these the author has common but distinct then generic names; s, Plangichnus, and

1864, Am. Jour. Sci., vol. 37, p. 34, Coal

1867.—Homothetus fossilis.

inal vein extremely passed by the scapularly any branches to no-median vein with in the outer fourth arno-median vein sim-

Type H. fossilis.

1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202, and Acad. Geol., p. 525, Upper Devonian.

1867.—Megathentomum hartti.

with a single, or at times, which are almost al. Type L. hartti.

hartti, Scudder, 1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202, and Acad. Geol., p. 525, Upper Devonian.

LITHOMYLACRIS, Scudder, 1879, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 48. [Ety. lithos, stone; Mylakis, a kind of roach.] Mediastinal and scapular areas together occupying more than half the wing; externo-median area small, compressed, scarcely expanding apically. Type L. angustum.

FIG. 1089.—Lithomyleris angustum. angustum, Scudder, 1879, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 48, Coal Meas.

pittstonianum, Scudder, 1879, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 50, Coal Meas.

simplex, Scudder, 1879, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 51, Coal Meas.

MEGATHENTOMUM, Scudder, 1868, Geo. Sur. Ill., vol. 3, p. 570. [Ety. megathos, largeness; entomon, an insect.] Wings of great size, remarkably broad and rounded; veins distant; simple, infrequent divarications, and cross neuration of delicate, irregular veinlets; the wing is also dotted with larger and smaller spots. Type M. pustulatum.

FIG. 1090.—Megathentomum pustulatum.

pustulatum, Scudder, 1868, Geo. Sur. Ill., vol. 3, p. 570, Coal Meas.

MAMIA, Dana, 1864, Am. Jour. Sci. and Arts, 2d ser., vol. 37, p. 34. [Ety. proper name.] Scapular vein close to the mediastinal; straight, main branch arising near the middle of the wing, and nowhere distant from the main stem.

Type M. bronsoni.

bronsoni, Dana, 1864, Am. Jour. Sci. and Arts, 2d ser., vol. 37, p. 34, Coal Meas. danae, see Gerarus danae.

MYLACRIS, Scudder, 1868, Geo. Sur. Ill., vol. 3, p. 568. [Ety. Mylakis, a kind of cockroach.] Wings broad, mediastinal and scapular areas together occupying

less than half the wing; scapular area larger than the mediastinal. Type M. anthracophilum.

anthracophilum, Scudder, 1868, Geo. Sur. Ill., vol. 3, p. 568, Coal Meas.

antiquum, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 300, Coal Meas.

bretonense, Scudder, 1874, (Blattina bretonensis), Can. Nat., vol. 7, p. 271, Coal Meas.

carbonum, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 304, Coal Meas.

heeri, Scudder, 1874, (Blattina heeri), Can. Nat., vol. 7, p. 272, Coal Meas.

lucifugum, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 301, Coal Meas.

mansfieldi, Scudder, 1880, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 47, Coal Meas.

ovale, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 308, Coal Meas.

pennsylvanicum, Scudder, 1880, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 44, Coal Meas.

priscovolans, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 307, Coal Meas.

NECYMYLACRIS, Scudder, 1880, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 53. [Ety. nekus, dead; Mylakis, a roach.] Some of the apical branches of the mediastinal vein arise beyond the base of the wing, and scarcely partake in the radiate arrangement of the others. Type N. laeoanum.

heros, Scudder, 1880, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 54, Coal Meas.

laeoanum, 1880, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 53, Coal Meas.

ORYCTOBLATTINA, Scudder, 1885, Proc. Acad. Nat. Sci. Phil., p. 37. [Ety. oruktos, quarried; Blattina, a genus.] Principal veins widely separated at base; scapular area surpassing apex, and with externo-median occupying more than half the wing, the branches of latter inferior; interno-median vein comparatively short. Type O. occidua.

occidua, Scudder, 1885, Proc. Acad. Nat. Sci. Phil., p. 37, Coal Meas.

PAOLIA, Smith, 1871, Am. Jour. Sci. and Arts, 3d ser., vol. 1, p. 44. [Ety. proper name.] Wings long, slender, branches of veins dichotomizing strongly, and running in a longitudinal direction, so that the externo-median branches oc-

FIG. 1091.—Mylacriscus anthracophilum.

FIG. 1092.—Mylacriscus lucifugum.

- cupy only a slight portion of the lower margin. Type *P. vetusta*.
 gurleyi, Scudder, 1884, Proc. Amer. Acad., vol. 20, p. 173, Coal Meas.
 lacoana, Scudder, 1884, Proc. Amer. Acad., vol. 20, p. 173, Coal Meas.
 superba, Scudder, 1884, Proc. Amer. Acad., vol. 20, p. 173, Coal Meas.
 vetusta, Smith, 1871, Am. Jour. Sci. and Arts, 3d ser., vol. 1, p. 44, Coal Meas.
 PAROMYLACRIS, Scudder, 1885, Proc. Acad. Nat. Sci. Phil., p. 35. [Ety. *paros*, before, or forefather; *Myllakris*, a kind of roach.] Body much arched; pronotal shield more than twice as broad as long; wings extremely broad; mediastinal area large and extended, and with the scapular occupying half the wing; externo-median area expanding apically. Type *P. rotundum*.
 rotundum, Scudder, 1885, Proc. Acad. Nat. Sci. Phil., p. 35, Coal Meas.
 PETRABlattina, Scudder, 1876, Can. Geol., vol. 3, p. 88. [Ety. *petra*, stone; *Blattina*, a genus.] Scapular and externo-median areas together covering more than half the wing; the externo-median vein directed toward and terminating near the middle of the inner border of the wing, branches superior; interno-median vein very short. Type *P. sepulta*.
 sepulta, Scudder, 1876, (*Blattina sepulta*), Can. Nat. and Geol., vol. 8, p. 88, Coal Meas.
 PHTHANOCORIS, Scudder, 1884, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 58. [Ety. *phthano*, first; *kore*, pupa.] Front wing differentiated from the hind wing; corium distinct from the membrane,

FIG. 1093.—*Plangtichnus erraticus*.

narrow clavus; no embolus or cuneus; mediastinal and scapular veins widely separated at base. Type *P. occidentalis*.
 occidentalis, Scudder, 1884, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 58, and Mem.

Bost. Soc. Nat. Hist., vol. 3, p. 348, Coal Meas.

PLANGTICHNUS, n. gen. [Ety. *Plangtos*, wandering; *ichnos*, track.] A zigzag, half-cylindrical, broken trail, running in any and every direction; sometimes dotted or sunk deeper at the angles than at other places, or most depressed between the angles in some cases. Supposed to have been made by the larva or pupa of some Palaeodictyopterous insect. See remarks under *Haplotichnus*. Type *P. erraticus*.

erraticus, n. sp. A simple, irregularly zigzag, half-cylindrical, broken trail, running in any and every direction, depressed in spots deeper than the general trail. Collected in the upper part of the Kaskaskia Group at the Whetstone quarries, in Orange County, Indiana.

PLATYPHEMERA, Scudder, 1867, Can. Nat.

and Geol., 2d ser., vol. 3, p. 202. [Ety. *platys*, flat; *ephemera*, an insect.] Founded upon the fragment of an upper wing,

FIG. 1094.—*Platyphemera antiqua*.

showing nervation and a heavy cross vein near the base between two middle veins, from which new prominent veins arise; ancient May-flies, in which the lower externo-median stem seems to be formed on the same plan as the upper stem. Type *P. antiqua*.
 antiqua, Scudder, 1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202, and Acad. Geol., p. 524, Devonian.

POLYERNUS, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 343. [Ety. *polys*, many; *ernos*, a scion.] Body moderately stout; wings rather broad; mediastinal vein extending nearly to the tip of wing; branches of scapular vein inequidistant at origin, longitudinal, closely crowded and ramose, yet hardly more important than the externo-median vein. Type *P. complanatus*.
 complanatus, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 343, Coal Meas.
 laminarum, Scudder, 1884, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 343, Coal Meas.
 PROMYLACRIS, Scudder, 1885, Proc. Acad. Nat. Sci. Phil., p. 34. [Ety. *pro*, before; *myllakris*, a kind of roach.] Body much arched; wing broad; mediastinal and scapular areas together not occupying more than a third of the wing; scapular area smaller than the mediastinal, the vein running obliquely to the costal margin. Type *P. ovale*.

ovale, Scudder, 1885, Proc. Acad. Nat. Sci. Phil., p. 34, Coal Meas.

, vol. 3, p. 348, Coal

[Ety. *Platys*, track.] A zigzag, broken trail, running in every direction; sometimes deeper at the angles, or most depressed in some cases, been made by the same Palaeodictyopter-marks under Haploptericus.

simple, irregularly circular, broken trail, and every direction, deeper than the one collected in the upper Kaskaskia Group at the Whetstone quarries, in Orange County,

er, 1867, Can. Nat.

1094.—*Platypheura antiquorum*.

a and a heavy cross between two mid-which new prominent May-flies, in external and scapular veins as in Palaeoptera; external and scapular veins amalgamated at base with the scapular, branching beyond the middle, internal and scapular divided at base in two branches. Type X. *antiquorum*, Scudder, 1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202, and Acad. Geol., p. 323, Upper Devonian.

1884, Mem. Bost. Soc. p. 343. [Ety. *polya*, division.] Body moderately broad; extending nearly to the base of scapular vein origin, longitudinal, and ramose, yet hardly more than the external. P. complanatus. Scudder, 1884, Mem. Bost. Soc. p. 343, Coal Meas. Scudder, 1884, Mem. Bost. Soc. p. 343, Coal Meas. Scudder, 1885, Proc. Acad. Nat. Hist., vol. 3, p. 334. [Ety. *pro*, before; *roch*, reach.] Body much broader; mediastinal and scapular not occupying the whole of the wing; scapular vein beyond the mediastinal, obliquely to the costal margin.

Scudder, 1867, Proc. Acad. Nat. Hist., vol. 3, p. 348, Coal Meas.

PROPTETICUS, Scudder, 1881, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 334. [Ety. *pro*, early; *ptetik*, winged.] Scapular vein widely separated from the mediastinal, arcuate, main branch arising near the base of the wing, parting widely from the main stem. Type P. infernus. Scudder, 1881, Mem. Bost. Soc. Nat. Hist., vol. 3, p. 334, Coal Meas. TENNESSEE, Linnæus, 1748, *Systema Naturæ*, p. 610. Not a Palaeozoic genus. *contusus*, see *Didymophleps contusa*.

FIG. 1095.—*Treptichnus bifurcus*. The trails are larger than shown by the figure.

TITANOPHAMA, Brongniart, as recognized by Scudder. Wing very large, moderately slender; neuration moderately

abundant; scapular vein beginning to branch in the middle of the basal half of the wing.

jucunda, Scudder, 1884, Proc. Amer. Acad., vol. 20, p. 109, Coal Meas.

TREPTEICINUS, n. gen. [Ety. *treptos*, to be turned about; *ichnos*, track.] A zigzag, half-cylindrical, continuous trail, forked at each angle, and running in any direction; each line is prolonged in the direction in which the animal moved, at the angle, so as to form a short fork or projection. Supposed to have been made by the larva or pupa of some Palaeopteron insect. See remarks under *Haplotichnus*. Type T. *bifurcus*.

bifurcus, n. sp. A zigzag, half cylindrical, continuous trail, quite evenly depressed, and forked at each angle; the bifurcation takes place in the direction in which the animal moved, but generally is less sunken than the trail, and sometimes shows simply a dot disconnected with the angle. Collected in the upper part of the Kaskaskia Group at the Whetstone quarries in Orange County, Indiana.

XENONEURA, Scudder, 1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202. [Ety. *xenos*, new, strange; *neura*, a vein.] Mediastinal and scapular veins as in Palaeoptera; external and scapular veins amalgamated at base with the scapular, branching beyond the middle, internal and scapular divided at base in two branches. Type X. *antiquorum*.

antiquorum, Scudder, 1867, Can. Nat. and Geol., 2d ser., vol. 3, p. 202, and Acad. Geol., p. 323, Upper Devonian.

FIG. 1096.—*Xenoneura antiquorum*.

SUBKINGDOM VERTEBRATA.

THIS is the highest division of the Animal Kingdom, and, until within the last twenty years, the essential character upon which the subkingdom was based was the possession of a bony or cartilaginous internal skeleton, having a spine or vertebral column. Since that time the class known as Tunicates, or Ascidians, which have no bony skeleton, has been referred to it; and the *Amphioxus lanceolatus*, a little, slender, transparent creature, having only a gelatinous cord, no brain cavity, and colorless blood, which was regarded as the lowest type of fishes, and had been elevated into an order called *Leptocardia*, is now taken out of the Class Pisces and referred to a separate class called *Acrania*. The Classes *Tunicata* and *Acrania* are not, however, known in Palæozoic rocks. Another class, called *Cyclostomata*, consisting of long, cylindrical, worm-like bodies, with a tough skin, destitute of scales, pectoral and ventral fins, but having a fin at the extremity of the body without any rays, and having a cartilaginous skeleton, and which includes the various species of lampreys that inhabit fresh water, and are also found in the ocean, and which have generally been regarded as an order of fishes, is unknown in Palæozoic rocks. Some authors would, however, place the *Conodonts* in this class; but if they do not belong to the *Annelida*, then there are stronger reasons for believing they belong to *Crustacea* than for thinking they should be referred to the *Cyclostomata*.

No Palæozoic fossil from an animal as highly organized as the lowest mammal or a bird has ever been found. The fossils are confined to the lowest organizations of fishes, batrachians, and reptiles. The lower forms of fishes have only a cartilaginous cord, resembling the embryonic state of fishes having an osseous skeleton. In higher forms the spine consists of bony vertebræ, united in such manner as to allow flexibility and strength by reason of attaching muscles, and also to protect a spinal nerve that passes through it.

CLASS PISCES, OR FISHES.

The Class Pisces has been divided into four subclasses—*Ganoidea*, *Selachia*, *Dipnoa*, and *Teleostia*. The *Teleostia* have been divided into eleven orders, and these into twenty suborders. This subclass embraces a very large majority of the living fishes. All of them have a complete bony vertebral column and skull. Nearly all edible fishes belong to this subclass. Many have a naked skin, but the majority are covered with horny scales of various forms. When the scales are smooth the fish are said to belong to the *Cycloidea*; when the hinder margins of the scales are denticulated they belong to the *Ctenoidea*. Fishes have pectoral fins, which are called the representatives of anterior limbs; and ventral fins, representatives of posterior limbs; and also dorsal, caudal, and anal fins. This subclass is unknown in Palæozoic rocks. Agassiz divided the fish into four groups—*Cycloids*, *Ctenoids*, *Placoids*, and *Ganoids*—based on the character of the scales; but as a single fish has been found bearing two of these types of scales, and as it is now

known that the whole structure of animals must be taken into consideration in classification, his system, like that of every other based on a single character, has given way to more perfect knowledge of animal life and physical structure. And it is quite probable, if the characters of Devonian fishes could be completely ascertained, as we know the living forms, they would all be found to belong to extinct subclasses. They are judged, however, only from meager fragments of the ossified parts, and arranged by homologies with the existing species, and classed in orders where the affinities seem most strongly to arrange them.

SUBCLASS GANOIDEA.

The Ganoidea (from *ganos*, brightness, in allusion to the enameled armor with which some of them are covered) commences in the Devonian strata, where the fossil remains soon become abundant, and continue to occur from that time forward to the present, though very few families now exist. Agassiz included as Ganoids all fish covered, in whole or in part, with bony plates; but some of the living genera were found to belong to the Teleostia, and later classification has been held to include all fossil species falling within the original definition of Agassiz and part of the living forms. The dermal skeleton consists of smooth, bony plates, covered with enamel. In some cases they are rhomboidal, arranged edge to edge in oblique transverse rows; in other cases the scales are rounded; and in a few species the skin is naked. There is much diversity in the skeletons, and all shades of ossification in the vertebral column and skull from cartilaginous to perfect bone. The subclass has been divided into seven orders, viz.: Chondrostea, Halecomorpha, Ginglymoda, Pycnodonta, Crossopterygia, Acanthodea, and Placodermata.

The Order Chondrostea includes the sturgeons of fresh and salt water, and the paddle-fish or spoon-bill cut of the Mississippi River and its tributaries.

The Order Halecomorpha (shad-like) is generally united with the Ginglymoda, under the name of the Holostea; but is distinguished by having large, round scales, no shingle-like fulcra on the fins, and in having the vertebræ concave at both ends, as in the Teleostia. The only living genus is the *Amia*, called bow-fin, mud-fish, dog-fish, etc. It is common to the lakes and sluggish rivers. The order is not certainly known in Palæozoic rocks.

The Order Pycnodonta has a short, vertically-flattened body, covered with rhomboid scales and peculiar dermal ribs. Tail either heterocercal or homocercal.

The Order Ginglymoda has a bony skeleton, rhomboid scales, and shingle-like fulcra on the fins. The vertebræ are convex in front and concave behind, forming ball and socket joints; tail heterocercal, and ventral fins between the pectorals and anals. This order is represented by the gar-pikes, which are common in American rivers.

The Order Crossopterygia is represented by two genera in the African waters, and fossils are referred to it back in geological time as far as the Devonian. The scales may be cycloid or rhomboid; the throat is protected by two or more plates; the caudal fin is diphyccercal; dorsal fin is divided in two or more divisions; pectorals and ventrals have a scaly axis; no fulcra.

The Order Acanthodea had cartilaginous skulls, heterocercal tails, rhomboidal scales, and were armed with a spine before each fin, and are said to occupy a place between the Ganoidea and Selachia. They are all Palæozoic.

The Order Placodermata had the head and thoracic region inclosed in sculptured, bony plates. In some the tail was naked, in others it was covered with ganoid scales; in some the fins were inclosed in plates, but the vertebræ were not ossified. This order includes the oldest fish remains known to the geologist.

SUBCLASS SELACHIA.

The word Selachia is derived from *selachos*, the Greek word for shark. This subclass is also called Elasmobranchia and Chondropterygia, and it includes the living sharks, rays, and skates. The skeleton is cartilaginous, and the plates of the skull are united without sutures. There are pectoral and ventral fins, and the caudal fin is usually heterocercal. The surface of the body is naked or covered with calcified papillæ, comparable with teeth, and even spinous. The placoid scales sometimes form a sculptured armor. The dermal spines found fossil are collectively known as Ichthyodolulites. The teeth are never inserted into the jaws, but are sustained in their position by the strong skin of the gums. They sometimes have obtuse crowns, and form a pavement for both jaws; in other cases the teeth are conical, sharp, arranged in rows, with the apices pointed backward. The subclass is divided into the Holocephala and Plagiostomata. The Holocephala is represented in the existing seas by the Family Chimæridæ, and, it is said, combines some of the characters of the Selachia, Ganoidea, and Batrachia. The Plagiostomata is divided into two orders, the Squalina and Raiina. The vertebræ are well developed, and the skin is covered with plates, shields, or spines. The Order Squalina includes the ocean sharks and dog-fishes. The Order Raiina includes the skates and rays of the present seas; one of them is called the saw-fish, and another produces dangerous electrical discharges.

SUBCLASS DIPNOA.

This subclass is said to furnish a connecting link from the Ganoidea to the Batrachia. In external appearance the fish are ganoid-like. The body is long, eel-like, covered with scales, and terminates in a compressed caudal fin with weak fin-rays. The head is broad and flat. There are two orders, Monopneumonia and Dipneumonia. The Monopneumonia includes the Ceratodidæ, some of which are living in Australia, and they are common in the Mesozoic rocks, but the existence of them in the Palæozoic rocks is very doubtful.

The Order Dipneumonia contains the living Family Sirenidæ, which contains two genera, the *Lepidosiren*, from the rivers of Brazil, and the *Protopterus*, from tropical Africa. There is little reason to believe this order is represented in Palæozoic rocks, though *Ctenodus* and *Dipterus* have been referred to it.

SUBCLASS GANOIDEA.

ORDER ACANTHODEA.

FAMILY ACANTHODIDÆ.—*Acanthodes*.

ORDER CHONDROSTEA.

FAMILY CHONDROSTEIDÆ.—*Asterosteus*, *Macropetalichthys*.

FAMILY PALÆONISCIDÆ.—*Chirolepis*, *Mecolepis*, *Palæoniscus*, *Rhadinichthys*.

ORDER CROSSOPTERYGIA.

FAMILY CELOCANTHIDÆ.—Cēlocanthus.

FAMILY CROSSOPTERYGIDÆ.—Ceratodus, Conchodus, Ctenodus, Heliodus, Onychodus, Peplorhina.

FAMILY DIPTERIDÆ.—Dipterus, Gnathorhiza, Ptyonodus.

FAMILY HOLOPTYCHIDÆ.—Glyptolepis, Holoptychius.

FAMILY PHANEROPLEURONIDÆ.—Phaneropleuron.

FAMILY RHIZODONTIDÆ.—Eusthenopteron, Rhizodus.

ORDER GINGLYMODA.

FAMILY LEPIDOSTEIDÆ.—Acrolepis, Amblypterus, Eurylepis.

ORDER PLACODERMATA.

FAMILY CEPHALASPIDÆ.—Acanthaspis, Acantholepis, Cephalaspis.

FAMILY COCCOSTEIDÆ.—Coccosteus, Dinichthys, Liognathus.

FAMILY PTERASPIDÆ.—Diplaspis, Palæaspis.

FAMILY PTERICHTHYIDÆ.—Aspidichthys, Bothriolepis, Pterichthys.

FAMILY UNCERTAIN.—Mycterops.

ORDER PYCNODONTA.

FAMILY PYCNODONTIDÆ.—Platysomus.

FAMILY UNCERTAIN.—Ectosteomachus.

SUBCLASS SELACHIA.

DIVISION HOLOCEPHALA.

ORDER CHIMEROIDIDEA.

FAMILY CHIMEROIDIDÆ.—Cyrtacanthus, Liognathus, Machæracanthus, Ptyctodus, Rhinodus, Rhynchodus.

DIVISION PLAGIOSTOMATA.

ORDER SQUALINA.

FAMILY COCHLIODONTIDÆ.—Chitonodus, Cochliodus, Cymatodus, Deltodopsis, Deltodus, Deltoptychius, Helodus, Orodus, Orthopleurodus, Petrodus, Platyodus, Pœcilodus, Psephodus, Sandalodus, Stenopterodus, Tæniodus, Tomodus, Trigonodus, Vaticinodus, Xystrodus. The Cochliodontidæ commenced at the base of the Subcarboniferous, reached their greatest development in the same geological system, and only one genus, Orthopleurodus, is found as high as the Coal Measures.

FAMILY HYBODONTIDÆ.—Agassizodus, Apedodus, Bathychilodus, Carcharopsis, Cladodus, Diplodus, Hybochilodus, Janassa, Lambdodus, Liodus, Mesodmodus, Orodus, Peripleurodus, Phœbodus, Pristichilodus, Stenmatodus, Thrinacodus, Polyrhizodus.

FAMILY PETALODONTIDÆ.—Antliodus, Calapodus, Cholodus, Chomatodus, Ctenopetalus, Ctenoptychius, Dactylodus, Desmiodus, Fissodus, Harpacodus, Lisgodus, Peltodus, Peripristis, Petalodus, Petalorhynchus, Polyrhizodus, Pristodus, Tannodus, Venustodus.

FAMILY ICTHYODORULITES.—Acondylacanthus, Amacanthus, Anaclitacanthus, Asteroptychius, Batacanthus, Bythilacanthus, Compacanthus, Ctenacanthus, Cyrtacanthus, Drepanacanthus, Edestus, Erimacanthus, Eunemacanthus, Gampsacanthus, Gisacanthus, Glymmatacanthus, Gyrcacanthus, Homacanthus, Leercacanthus, Listracanthus, Macheracanthus, Marracanthus, Oracanthus, Orthacanthus, Physonemus, Pnigacanthus, Stenacanthus, Xystracanthus.

ORDER RAIINA.

FAMILY PSAMMODONTIDÆ.—Copodus, Psammodus.

ACANTHASPIS, Newberry, 1875, Ohio Pal., vol. 2, p. 36. [Ety. *akantha*, spine; *aspis*, shield.] Cranium plates some-

what quadrangular at one end, then abruptly bending to one side, and prolonged to an acute point; surface carinated and tuberculated. Type A. armata.

armata, Newberry, 1875, Ohio Pal., vol. 2, p. 37, Up. Held. Gr.

ACANTHODES, Agassiz, 1833, Recherches sur les Poiss. Foss., vol. 1, p. 19. [Ety. *akantha*, spine.] Fish lepidoid, mouth wide; lower jaw longer than the upper; teeth brush-like; scales small; dor-

sal fin opposite anal;

pectoral large;

first ray of each fin strong, large, stiff; rays of caudal fin close. Type A. bronni.

affinis, Whiteaves, 1899, Trans. Roy. Soc. Can., vol. 6, p. 77, Low. Devonian.

concinus, Whiteaves, 1889, Trans. Roy. Soc. Can., vol. 6, p. 77, Low. Devonian.

ACANTHOLEPIS, Newberry, 1875, Ohio Pal., vol. 2, p. 38. [Ety. *akantha*, spine; *lepis*, scale.] Tuberculated cranial or dermal plates, having a prevailing

spatulate outline; some are thin and have the appearance of large, elongated, unsymmetrical scales; others are stronger and produced into points that sometimes become spines. Type A. pustulosa.

pustulosa, Newberry, 1875, Ohio Pal., vol. 2, p. 38, Up. Held. Gr.

ACONDYLACANTHUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 432. [Ety. *akondylos*, without bony knobs; *akantha*, spine.] Fin rays long, gradually tapering, laterally compressed, moderately curved posteriorly; lateral faces longitudinally fluted; costae smooth, enameled, increasing by bifurcation and implantation; posterior face excavated longitudinally, without median keel; postero-lateral angles bearing a row of downward hooked denticles; pulp cavity occupying the posterior half of the spine. Type A. gracilis.

aequicostatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 434, Keokuk Gr.

gracilis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 433, Waverly or Kinderhook Gr.

mudgianus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 244, Up. Coal Meas.

nuperus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 242, Up. Coal Meas.

occidentalis, Newberry & Worthen, 1868, (Leptacanthus occidentalis),

FIG. 1098.—Acantholepis pustulosus. Reduced outlines of four plates, probably in their relative positions.

FIG. 1097.—Acanthaspis armata. Plate bearing spine from left side of cranium.

FIG. 1099.—Acondylacanthus gracilis. Side view of spine mag. 1/2 diameter.

thus, Anacriticacanthus, Ctenomacanthus, Eumacanthus, Gyrocantus, anthus, Marracanthus, Stenacanthus, Stenacanthus,

some are thin and lanceolate; others are broad and pointed at the tip. Type A.

1875, Ohio Pal., vol. 38, Up. Held. Gr. Anacanthus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 432. [Ety. *anacanthus*, without bony *akantha*, spine.] Fin long, gradually tapering, laterally compressed, deeply curved posteriorly; lateral faces longitudinally striated; costae smooth, rounded, increasing by bimodal and implantation; anterior face excavated; finally, without meel; postero-lateral

FIG. 1009.—Acondylacanthus gracilis. Side view of spine magn. 1/2 diameter.

Newberry & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 434.

Geo. Sur. Ill., vol. 2, p. 116, St. Louis Gr. rectus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 241, Up. Coal Meas. xiphias, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 244, Keokuk Gr. Acrolepis, Agassiz, 1830, Recherches sur les Poiss. Foss., vol. 2, p. 79. [Ety. *akros*, sharp.]

FIG. 1100.—Acrolepis sedgwicki. Magnified scale.

strongly keeled and sulcated diagonally. Type A. sedgwicki. hortonensis, Dawson, 1868, Acad. Geol., p. 254, Subcarboniferous. Agonichthys, Newberry, syn. for Macropetalichthys. manni, see Macropetalichthys manni. sullivanti, see Macropetalichthys sullivanti. Agassizodus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 311. [Ety. proper name; *odous*, tooth.] Teeth variable, transversely elongated, base usually produced; crown traversed by a crest, raised into several summits, the central one often large. Type A. variabilis. corrugatus, Newberry & Worthen, 1870, (Orodus corrugatus.) Geo. Sur. Ill., vol. 4, p. 358, Coal Meas. scitulus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 322, Coal Meas.

FIG. 1101.—Agassizodus variabilis.

variabilis, Newberry & Worthen, 1870, (Lophodus variabilis.) Geo. Sur. Ill., vol. 4, p. 361, Coal Meas. virginianus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 321, Coal Meas. Amacanthus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 464. [Ety. *ama*, backward; *akantha*, spine.] Dorsal spine firmly implanted, curved forward, laterally compressed, posterior face truncated and longitudinally keeled or denticulate along the median line; rounded and tuberculated in the concave anterior face; lateral surface covered with tuberculate costae. Type A. gibbosus.

gibbosus, Newberry & Worthen, 1866, (Homacanthus gibbosus.) Geo. Sur. Ill., vol. 2, p. 113, St. Louis Gr. Amelyrtus, Agassiz, 1833, Recherches sur les Poissons Fossiles, t. 1, p. 28. [Ety. *ambly*, blunt; *pteron*, fin.] All fins large and composed of numerous rays; pectoral very large; anal broad; dorsal opposite the anal point of the ventral, which is far back; little rays on the superior lobe of the heterocercal tail; head blunt; scales medium, rhomboid. Type A. macropterus. macropterus, Agassiz, 1836, Recherches sur les Poissons Fossiles, vol. 2, p. 28, Coal Meas. Anacriticacanthus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 442. [Ety. *anacritic*, leant upon; *akantha*, spine.] Fin spine recumbent or imbedded along its inferior extent, laterally compressed, subovate in transverse section; exposed part constricted along the line of union with the base. Type A. semicostatus. semicostatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 443, Burlington Gr. Antiodus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 33. [Ety. *antia*, a depression; *odous*, a tooth.] Teeth transversely elliptical, compressed, concavo-convex; crown similar to that of Petalodus; root short or obsolete. Type A. mucronatus. cucullus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 41, Keokuk Gr. gracilis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 393, Warsaw Gr. minutus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 43, Keokuk Gr. mucronatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 38, St. Louis Gr. parvulus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 38, Burlington Gr. perovalia, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 393, Warsaw Gr. politus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 42, Keokuk Gr. robustus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 39, Kaskaskia Gr. sarcinulus, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 356, Burlington Gr. similis, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 41, Keokuk Gr. simplex, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 44, Burlington Gr. sulcatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 45, Keokuk Gr. Apedodus, Leidy, 1856, Jour. Acad. Nat. Sci., 2d ser., vol. 3, p. 162. [Ety. *apedos*, level, smooth; *odous*, tooth.] Flattened lancet-shaped teeth. Type A. priscus. priscus, Leidy, 1856, Jour. Acad. Nat. Sci., 2d ser., vol. 3, p. 162, Chemung Gr. Aspichthys, Newberry, 1873, Ohio Pal., vol. 1, p. 322. [Ety. *aspis*, shield; *ichthys*, fish.] Dorsomedian plate of the carapace similar to that of Pte-

richthys, but many times larger and covered with large, hemispherical, smooth, enameled tubercles. Type *A. clavatus*.

clavatus, Newberry, 1873, Ohio Pal., vol. 1, p. 323, Portage Gr.

FIG. 1102.—*Aspidichthys clavatus*.

Aspidodus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 92, syn. for *Psephodus*.

convolutus, see *Psephodus convolutus*.

crenulatus, see *Psephodus crenulatus*.

Astrocanthus siderius, see *Bythiacanthus siderius*.

ASTEROPTYCHIUS, McCoy, 1854, British Pal. Rocks, p. 615. [Ety. *aster*, star; *ptych*, wrinkle.] Bony fin-ray compressed, long, slender, gradually tapering to a point at the distal end, and abruptly tapering at the striated proximate end or base of insertion; sides moderately convex, converging to the anterior edge, which is strongly, keeled; posterior face with a moderate cavity, each lateral edge having a row of small, pointed teeth, directed upward; sides with smooth, thread-like ridges, separated by broader, flat, longitudinally striated spaces on which are irregularly scattered, smooth, spinous tubercles. Type *A. ornatus*.

bellulus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 439, Coal Meas.

kekuk, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 436, Kekuk Gr.

studenovi, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 437, St. Louis Gr.

tenellus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 248, Up. Coal Meas.

tenuis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 438, Kaskaskia Gr.

triangularis, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 370, Burlington Gr.

vetustus, St. John and Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 435, Waverly or Kinderhook Gr.

ASTEROSTEUS, Newberry, 1875, Ohio Pal., vol. 2, p. 35. [Ety. *aster*, star; *osteon*, bone.] Head long, narrow, broadening in the occipital region; surface covered by a sheet of tuberculated enamel;

nasal pits strongly marked; condyle-like posterior projections. Type *A. stenocephalus*.

stenocephalus, Newberry, 1875, Ohio Pal., vol. 2, p. 36, Corniferous str.

BATACANTHUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 468. [Ety. *batos*, prickly bush; *akantha*, spine.] Spines long, tapering, curved forward; apex obtuse; transverse section subcircular or oval, with anterior angle and posterior face; lateral surfaces rounded, covered with stellate tubercles with intercostal sulci; base moderately inserted; pulp cavity subcentral. Type *B. baculiformis*.

baculiformis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 469, Kekuk Gr.

necis, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 253, Kekuk Gr.

stellatus, Newberry & Worthen, 1866, (*Drepanacanthus* (?) *stellatus*.) Geo. Sur. Ill., vol. 2, p. 125, Kekuk Gr.

BATHYCHILODUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 252. [Ety. *bathys*, deep; *cheilos*, lip; *odus*, tooth.] Teeth minute, laterally elongated, subelliptical, sinuous in front; median cusp with cutting edges flanked by a pair of diverging denticles of similar shape, with a minute denticle between the lateral and median cusps.

Type *B. macissaei*.

macissaei, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 252, Middle Devonian.

FIG. 1103.—*Bathychilodus macissaei*.

BOTHRIOLEPIS, Eichwald, 1840, Bull. Soc. St. Petersburg. [Ety. *bothrion*, a furrow;

FIG. 1104.—*Bothriolepis canadensis*.

lepis, a scale.] Cephalic shield somewhat semielliptical in outline and covered with plates, as in *Pterichthys* and

marked; condyle-
ctions. Type A.

ry, 1875, Ohio Pal.,
s (tr.
& Worthen, 1875,
6, p. 468. [Ety.
akantha, spine.]
g, curved forward;
verse section sub-
lith anterior angle
; lateral surfaces
lith stellate tuber-
sules; base moder-
cavity subcentral.

& Worthen, 1875,
6, p. 469, Ke-

Worthen, 1883, Geo.

53, Keokuk Gr.
& Worthen, 1866,
stellatus.) Geo. Sur.

Keokuk Gr.
& Worthen, 1875,
6, p. 252. [Ety.

lip; *odous*, tooth.]
ally elongated, sub-
in front; median
edges flanked by a
denticles of similar
minute
e lat-
usps.

n &
Geo. Sur. FIG. 1103.—*Ha-*
middle
mucronat.

d, 1840, Bull. Soc. St.
boltrion, a furrow;

lepis canadensis.

cephalic shield some-
l in outline and cov-
as in *Pterichthys* and

Asterolepis, but distinguished by the
course of the furrows and shape of
the plates; it has longer articulating
plates in the limb or arm, and has been
otherwise distinguished, though closely
related to both genera. Type B. or-
natus.

canadensis, Whiteaves, 1880. (*Pterichthys*
canadensis.) Am. Jour. Sci. and Arts,
3d. ser., vol. 20, p. 135, Up. Devo-
nian.

BYTHIACANTHUS, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 444. [Ety.
bythios, deep; *akantha*, spine.] Fin
spines deeply imbedded, laterally com-
pressed, exposed part recumbent,
tuberculated; posterior face low, keeled;
pulp cavity forming a deep channel in
the posterior side of the base. Type B.
vanhornii.

siderinus, Leidy, 1873, (*Asteracanthus* *sider-*
inus.) Ext. Vert. Fauna, p. 313, St.
Louis Gr.

vanhornii, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 445, St.
Louis Gr.

CALODONTUS, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 403. [Ety. *kalos*,
beautiful; *odous*, tooth.]

Teeth in general form
like *Petalodus*, but dis-
tinguished by the tur-
gid, subconical, unsym-
metrical crown. Type C.
apicalis.

apicalis, St. John & Wor-
then, 1875, Geo. Sur. Ill.,
vol. 6, p. 403, Middle
Coal Meas.

FIG. 1105.—*Calo-*
podus apica-
lis. Mag. 2
diam.

CARCHAROPSIS, Agassiz, 1843, *Recherches sur*
les Poissons Fossiles, vol. 3, p. 313.
[Ety. *carcharopsis*, shark-like.] Prin-
cipal cusps very strong, erect, com-
pressed in front, rounded behind,
broadly expanded at base; lateral an-
gles sharp, crenulated; extremities oc-
cupied by isolated, conical, lateral
denticles; coronal faces smooth or
faintly striated vertically; base in out-
line like *Cladodus*, moder-
ately produced
and broadly
rounded poste-
riorly; anterior
face sinuous,
terminating
below in a
ridge; inferior
surface exca-
vated. Type C.
prototypus.

FIG. 1106.—*Carcharopsis*
wortheni.

berry, 1866, Geo. Sur. Ill., vol. 2, p. 69,
Subcarboniferous.

CEPHALASPIS, Agassiz, 1836, *Recherch. Pois.*
Foss., t. 2, p. 135. [Ety. *kephale*, head;
asps, shield.] Entire skeleton external;
head shield very large, subcrescent-

iform when depressed but in better,
condition showing an arching over
the top of the
head, cov-
ered with
discoidal,
sculptured,
bony plates,
with the
crescent
horns di-
rected back-
ward; eyes
large, ellip-
tical, on each
side of the
upper cen-
tral part of
the head;
body rapidly
tapering, an-
gular on top,
and presenting a jointed appearance
somewhat like a trilobite; dorsal, anal,
and caudal fin, the latter like a paddle
or oar. Type C. *lyelli*.

FIG. 1107.—*Cephalaspis lyelli*.
Head shield depressed, and
showing the jointed angular
outline of part of the body.

campbeltoneensis, Whiteaves, 1881, Can.
Nat., vol. 10, Devonian.

dawsoni, Lankaster, 1870, London Geo.
Mag., Devonian.

CERATODUS, Agassiz, 1833, *Recherches sur*
les Poissons Fossiles, t. 1, p. 129. [Ety.
keras, horn; *odous*, tooth.] Teeth large,
thick, longer than wide, very porous;
crown transversely sulcated. Type C.
latissimus.

favosus, Cope, 1884, Pal. Bull., No. 39, p.
28, Permian.

pancristatus, Cope, 1877, Proc. Am. Phil.
Soc., p. 54, Permian.

vinslovi, Cope, 1876, Proc. Am. Phil. Soc.,
p. 410, Permian.

CHIRODUS, McCoy, 1848, Ann. and Mag. Nat.
Hist., vol. 2, p. 130. [Ety. *cheir*, the
hand; *odous*, tooth.] Tooth fan-shaped,
thick, flattened; anterior broad, margin
deeply divided into lobes; inner nearly
straight margin has a small, recurved,
thumb-like lobe projecting nearly at
right angles from the middle of its
length, preventing the mesial junction
of the titlers of each side of the jaw;
inner marginal lobe the longer; surface
minutely punctured. Type C. *pesianæ*.
Not definitely known in America.

acutus, Newberry, 1857, Proc. Acad. Nat.
Sci., vol. 8, p. 99, Coal Meas. Too poorly
defined to warrant recognition.

CHIROLEPIS, Agassiz, 1833, *Recherches sur*
les Poissons Fossiles, t. 1, p. 128. [Ety.
cheir, hand; *lepis*, scale.] Bones of the
head sculptured; shoulder-bone and
fins osseous; pectorals large, reaching
near the ventral fin, and ventral reach-
ing near the anal fin; dorsal fin small
and opposite the posterior part of the
anal fin; tail well-developed, principally
on the lower side; scales small, sculp-

- tured, and ranged diagonally in wavy lines. Type *C. trailli*.
canadensis, Whiteaves, 1881, Am. Jour. Sci. and Arts., 3d ser., vol. 21, p. 496, Up. Devonian.
- Chironodus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 109. [Ety. *chiton*, a smock or coat; *odus*, tooth.] Mandibular posterior teeth trapezoidal, arched in the direction of inrollment; median teeth narrow, inrolled longitudinally; maxillary posterior teeth subquadrilateral, arched, and inrolled along the outer margin. Type *C. spingeri*.
- antiquus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 116, Low. Burlington Gr.
- latus*, Leidy, 1856, (Cochliodus *latus*.) Trans. Am. Phil. Soc., vol. 11, p. 87, pl. 5, fig. 17, Keokuk Gr.
- liratus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 119, St. Louis Gr.
- rugosus*, Newberry & Worthen, 1866, (Pœcilodus *rugosus*, *P. ornatus*, and *P. convolutus*.) Geo. Sur. Ill., vol. 2, pp. 94, 95; vol. 4, p. 368, Keokuk Gr.
- spingeri*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 112, Up. Burlington Gr.
- tribulis*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 117, Keokuk Gr.
- Cholodus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 3, p. 415. [Ety. *cholos*, defective; *odus*, tooth.] Distinguished from *Peltodus* and *Fissodus*, by the eccentrically lobed crest and extreme downward prolongation of the lateral extremities of the coronal fold in the convex face. Type *C. inequalis*.
- inequalis*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 3, p. 416, Coal Meas.
- Chomatodus*, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 107. [Ety. *choma*, a pile or heap; *odus*, tooth.] Teeth transversely much elongated, compressed, and depressed; crown having the homologous parts of *Petalodus*, and the form and structure of *Polyrhizodus*; root short, sometimes obsolete, undivided. Type *C. linearis*.
- affinis*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 54, Keokuk Gr.
- angularis*, see *Tanaodus angularis*.
- arcuatus*, St. John, 1870, Proc. Am. Phil. Soc., vol. 2, p. 435, and Pal. E. Neb., p. 243, Coal Meas.
- chesterensis*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 363, Kaskaskia Gr.
- comptus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 356, Burlington Gr.
- costatus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 85, Keokuk Gr.

FIG. 1108.—*Cholodus inequalis*. Convex face.

- cultellus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 52, Kaskaskia Gr.
- elegans*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 86, Keokuk Gr.
- gracillimus*, see *Tanaodus gracillimus*.
- inconstans*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 360, St. Louis Gr.
- incrassatus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 359, St. Louis Gr.
- insignis*, Leidy, 1856, Trans. Am. Phil. Soc., vol. 11, p. 87, St. Louis Gr.
- linearis*, Agassiz, 1843, (Pœmmodus *linearis*.) Recherches Poiss. Foss., t. 3, p. 108, Subcarb.
- loriformis*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 58, Keokuk Gr.
- molaris*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 56, Keokuk Gr.
- multiplacatus*, see *Tanaodus multiplacatus*.
- obscurus*, see *Tanaodus obscurus*.
- parallelus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 358, Warsaw Gr.
- pusillus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 53, Keokuk Gr.
- varsoviensis*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 393, Warsaw Gr.
- venustus*, Leidy, see *Venustodus leidyi*, where the specific name is made to designate the genus, and the author the specific name, contrary to the rules of nomenclature; also see *Venustodus venustus*.
- Cladodus*, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 196. [Ety. *klados*, twig; *odus*, tooth.] Teeth with broad, horizontal, semicircular, thick, bony, coarsely fibrous base, rounded behind, truncated in front; crown divided into long, sharp, subulate, conical points, arranged along the straight truncated edge of the base; medial cone much larger than the secondary ones, of which latter the external cones are the larger; all the cones striated longitudinally, and either circular in section or with simple cutting edges, slightly compressed. Type *C. mirabilis*.
- acuminatus*, Newberry, 1857, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 99, and Ohio Pal., vol. 2, p. 45, Subcarboniferous.
- alternatus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 265, Waverly or Kinderhook Gr.
- angulatus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 24, Keokuk Gr.

FIG. 1109.—*Chomatodus incrassatus*.

FIG. 1110.—*Cladodus acuminatus*.

& Worthen, 1866,
2, p. 52, Kaskas-

Worthen, 1866, Geo.
6, Keokuk Gr.
lus gracillimus.
& Worthen, 1875,
ol. 6, p. 360, St.

Fig. 1109.—*Chomatodus*
incrassatus.

St. Louis Gr.
3, (*Psammodus* line-
Pois. Foss., t. 3, p.

& Worthen, 1866,
2, p. 58, Keokuk Gr.
& Worthen, 1866, Geo.
56, Keokuk Gr.
aodus multiplicatus.
as obscurus.
& Worthen, 1875,
ol. 6, p. 358, War-

& Worthen, 1866,
2, p. 53, Keokuk Gr.
in & Worthen, 1875,
3, p. 393, Warsaw Gr.
Venustodus leidyi,
name is made to des-
and the author the
strary to the rules of
o see *Venustodus ve-*

3, *Recherches sur les*
t. 3, p. 196. [Ety.
, tooth.] Teeth with
aemicircular, thick,
brous base, rounded
in front; crown dis-
sharp, subulate, con-
ed along the straight
f the base; medial
than the secondary
ter the external cones
ll the cones striated
d either circular in
mple cutting edges,
d. Type *C. mirabilis*.
acuminatus. New-
berry, 1857, Proc.
Acad. Nat. Sci.
Phil., vol. 8, p. 99,
and Ohio Pal., vol.
2, p. 45, Subcarbo-
niferous.

alternatus, St. John
& Worthen, 1875,
Geo. Sur. Ill., vol.
6, p. 265, Waverly
or Kinderhook Gr.
ry & Worthen, 1866,
vol. 2, p. 24, Keo-

bellifer, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 270, Burlington Gr.
carinatus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 279, Coal Meas.
concinus, Newberry, 1875, Ohio Pal.,
vol. 2, p. 48, Portage Gr.
costatus, Newberry & Worthen, 1866, Geo.
Sur. Ill., vol. 2, p. 27, Kaskaskia Gr.
deflexus, Newberry & Worthen, 1870,
Geo. Sur. Ill., vol. 4, p. 355, Burling-
ton Gr.
eccentricus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 272, St.
Louis Gr.
elegans, Newberry & Worthen, 1870, Geo.
Sur. Ill., vol. 6, p. 354, St. Louis Gr.
euglypheus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 274, St.
Louis Gr.
exiguus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 261, Waverly or Kin-
derhook Gr.
exilis, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 258, Waverly or Kin-
derhook Gr.
ferox, Newberry & Worthen, 1866, Geo.
Sur. Ill., vol. 2, p. 26, St. Louis Gr.
fulleri, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 276, Coal Meas.
gomphoides, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 269, Burling-
ton Gr.
gracilis, Newberry & Worthen, 1866, Geo.
Sur. Ill., vol. 2, p. 30, Coal Meas.
grandis, Newberry & Worthen, 1866, Geo.
Sur. Ill., vol. 2, p. 29, Kaskaskia Gr.
hertzeri, Newberry, 1875, Ohio Pal., vol.
2, p. 46, Portage Gr.
intercostatus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 267, Burling-
ton Gr.
isclypus, Newberry & Worthen, 1870, Geo.
Sur. Ill., vol. 4, p. 354, St. Louis Gr.
laminoides, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 30, Keokuk Gr.
magnificus, Tuomey, 1858, 2d Rep. Geo.
Ala., p. 39, Kaskaskia Gr.
micropus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 21, Keokuk Gr.
mortifer, Newberry & Worthen, 1866, Geo.
Sur. Ill., vol. 2, p. 22, Coal Meas.
newmaui, Tuomey, 1858, Geo. Ala., p. 39,
Kaskaskia Gr.
occidentalis, Leidy, 1859, Proc. Acad. Nat.
Sci. Phil., Up. Coal Meas.
pandatus, St. John & Worthen, 1875, Geo.
Sur. Ill. vol. 6, p. 278, Coal Meas.
parvulus, Newberry, 1875, Ohio Pal., vol.
2, p. 43, Portage Gr.
pattersoni, Newberry, 1875, Ohio Pal.,
vol. 2, p. 47, Waverly Gr.
politus, Newberry & Worthen, 1875, Geo.
Sur. Ill., vol. 2, p. 27, Kaskaskia Gr.
prænuntius, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 270, Burling-
ton Gr.
raricostatus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 271, Keo-
kuk Gr.

robustus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 20, Keokuk Gr.
romingeri, Newberry, 1875, Ohio Pal., vol.
2, p. 49, Waverly Gr.
spinosus, Newberry & Worthen, 1866,
Geo. Sur., Ill., vol. 2, p. 22, St. Louis Gr.
springeri, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 259, Waverly
or Kinderhook Gr.
stenopus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 23, St. Louis Gr.
subulatus, Newberry, 1875, Ohio Pal., vol.
2, p. 47, Cuyahoga shale over the Berea
grit.
succinctus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 265, Waverly or
Kinderhook Gr.
turritus, Newberry & Worthen, 1866, Geo.
Sur. Ill., vol. 2, p. 28, Keokuk Gr.
vanhornii, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 273, St.
Louis Gr.
wachsmuthi, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 263, Waverly or
Kinderhook Gr.
zygopus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 25, Kaskas-
kia Gr.

Climaxodus, McCoy, 1848, Ann. and Mag.
Nat. Hist., 2d ser., vol. 2. [Ety. *klimax*,
ladder; *odus*, tooth.] Tooth longer
than wide, gradually narrowing toward
the front, with nearly straight sides;
anterior part of the crown crossed by
broad, imbricating, transverse ridges, at
right angles to its length; surface mi-
nutely punctured. Type *C. imbricatus*.
Not definitely known in America.

brevis, Newberry, 1837, Proc. Acad. Nat.
Sci., vol. 8, p. 100, Coal Meas. Too
poorly defined to warrant recognition.
Coccostræus, Agassiz, 1836, *Recherch. Pois.*
Foss., vol. 2, p. 302. [Ety. *kokkos*, berry;
osteon, bone.] Head rounded; body
triangular, with long vertebrate tail,
like a rudder, the whole compared in
form, by Hugh Miller, to a boy's kite;
head and body covered with tuber-
culated bony plates; central front plate
like the keystone of an arch; the pos-
terior body plate is large, saddle-wise
toward the center, pointed behind; on
the ridge there is a longitudinal groove
ending in a perforation, a little behind
the apex. It is this plate which has
been described as *C. occidentalis*, but it
does not show groove or perforation.
Type *C. decipiens*.

acadiens, Whiteaves, 1881, Can. Nat., vol.
10, Upper Devonian.
occidentalis, Newberry, 1875, Ohio Pal.,
vol. 2, p. 32, Up. Held. Gr.

COCHLORODUS, Agassiz, 1843, *Recherches sur*
les Poissons Fossiles, t. 3, p. 113. [Ety.
kochlias, anything spiral; *odus*, tooth.]
Lower jaw thick, short, bony, V-shaped,
bearing on each ramus two obliquely
twisted and obtusely ridged semi-
cylindrical teeth, strongly inrolled on

the outer margin, convex above, concave below, with porous grinding surfaces, as in *Psammodus*, from the termination of the vertical medullary canals. Type *C. contortus*.

FIG. 1111.—*Cochliodus contortus*.

for *Sandalodus levissimus*.
latus, see *Chitonodus latus*.
leidyi, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 127, Kaskaskia Gr.
nitidus, see *Deltopychius nitidus*.
nobilis, Newberry & Worthen, syn. for *Chitonodus latus*.
obliquus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 126, St. Louis Gr.
occidentalis, see *Deltodus occidentalis*.
vanhornii, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 120, St. Louis Gr.

COELACANTHUS, Agassiz, 1836, *Recherches sur les Poissons Fossiles*, t. 2, p. 170. [Ety. *koilos*, hollow; *akantha*, spine.] Head plates sculptured; scales large, imbricated, sculptured, arranged diagonally; two small dorsal fins supported on interspinous bones, the anterior one a little forward of the ventral fin, and the posterior one nearly opposite the anal fin; caudal fin equi-lobate, and near its extremity a minute supple-

FIG. 1112.—*Coelacanthus elegans*.

mental canal; vertebral column cartilaginous, but neural arches and fin-rays bony; teeth small, numerous, conical. Type *C. granulatus*.

elegans, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 98, and Ohio Pal., vol. 1, p. 339, Coal Meas.

ornatus, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 98, and Ohio Pal., vol. 1, p. 340, Coal Meas.

robustus, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 98, and Ohio Pal., vol. 1, p. 340, Coal Meas.

costatus, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 364, Burlington Gr.
crassus, Newberry & Worthen, 1860, syn.

COMPSACANTHUS, Newberry, 1857, Proc. Acad. Nat. Sci., vol. 8, p. 99, and Ohio Pal., vol. 1, p. 331. [Ety. *compos*, elegant; *akantha*, a spine.] Spines small, gently curved backward; exposed part smooth, polished; section circular; single row of remote, depressed hooks on the posterior median line. Type *C. laevis*.

laevis, Newberry, 1857, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 99, and Ohio Pal., vol. 1, p. 332, Coal Meas.

FIG. 1114.—*Compsacanthus laevis*.

Conchiopsis, syn. for *Coelacanthus*.
anguliferus, syn. for *Coelacanthus elegans*.
exanthematicus, syn. for *Peplorhina anthracina*.

filiferus, syn. for *Coelacanthus elegans*.

FIG. 1115.—*Conchodus plicatus*.

CONCHODUS, McCoy, 1848, Ann. and Mag. Nat. Hist., 2d ser., vol. 2. [Ety. *conchos*, shell; *odus*, tooth.] Teeth large, somewhat semicircular, pointed in front, subtruncate behind, deeply concave on the grinding surface; internal margin straight, thickened, edge abruptly deflected; external border convex, much raised, undulato-plicate, ridge larger in front, smaller posterior; under surface polished, minutely porous. Type *C. ostreiformis*.

plicatus, Dawson, 1868, Acad. Geol., p. 209, Coal Meas.

COPODUS, Agassiz, MSS., 1859, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 227. [Ety. *kopis*, broad, curved knife; *odus*, tooth.] Teeth bilaterally symmetrical, spanning the jaw without mesial suture, arranged in single, longitudinal series from behind backward; lateral borders converging anteriorly; coronal region arched; rim at base; anterior and posterior walls vertical, channelled; inferior surface concave; porous beneath the enamel.

FIG. 1116.—*Copodus cornutus*. Maxillary form. *a*, Triturating surface; *b*, transverse profile; *c*, longitudinal profile.

Type *C. cornutus*.

erry, 1857, Proc.
8, p. 99, and Ohio
[Ety. *compus*, ele-
me.] Spines small,
ward; exposed part
section circular;
e, depressed hooks
dian line. Type C.

Proc. Acad. Nat.
99, and Ohio Pal.,
leas.

canthus laevis.

acanthus.
elacanthus elegans.
ior Peplorhina an-

acanthus elegans.

Coxichodus, Mc-
Coy, 1848,
Ann. and
Mag. Nat.
Hist., 2d.
ser., vol. 2.
[Ety. *con-*
chos, shell;
odus, tooth.]
Teeth large,
ircular, pointed in
behind, deeply con-
ing surface; internal
thickened, edge ab-
external border con-
undulato-plicate,
ont, smaller poste-
e polished, minutely
treiformis.

68, Acad. Geol., p.

Fig. a

Fig. b

Fig. c

Fig. 1116.—*Copodus*
cornutus. Maxil-
lary form. a, Tritu-
rating surface; b,
transverse profile;
c, longitudinal pro-
file.

ace. Type C. var-

pusillus, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 231, Kaskaskia Gr.

Fig. a

Fig. b

Fig. c

Fig. 1117.—*Copodus* *cornutus*. Mandibular
form. a, Triturating
surface; b, transverse
profile; c, longitudinal
profile.

transverse scales or tubercles; con-
triated base rapidly tapering, finely
striated. Type C. *tenuistriatus*.

angulatus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 118, Kaskas-
kia Gr.

burlingtonensis, St. John & Worthen,
1875, Geo. Sur. Ill., vol. 6, p. 426,
Burlington Gr.

butterfi, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 240, Lower Coal
Meas.

cannaliratus, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 239, Kaskas-
kia Gr.

costatus, see *Eunemacanthus costatus*.

coxanus, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 233, Keo-
kuk Gr.

deflexus, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 234, St. Louis Gr.

elegans, Tuomey, 1858, Geo. Ala., p. 38,
Kaskaskia Gr.

excavatus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 428, Keo-
kuk Gr.

formosus, Newberry, 1873, Ohio Pal., vol.
1, p. 328, Waverly Gr.

furcicaratus, Newberry, 1875, Ohio Pal.,
vol. 2, p. 54, Waverly Gr.

gemmatus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 429, St.
Louis Gr.

gracillimus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 126, St.
Louis Gr.

grado-costatus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 425, Burling-
ton Gr.

harrisoni, St. John & Worthen, 1883, Geo.

Sur. Ill., vol. 7, p. 236, St. Louis Gr.

keokuk, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 427, Keokuk Gr.

latispinosus, Whiteavis, 1881, Can. Nat.

and Geol., vol. 10, Upper Devonian.

marshi, Newberry, 1873, Ohio Pal., vol. 1,

p. 326, Coal Meas.

mayi, Newberry & Worthen, 1870, Geo.

Sur. Ill., vol. 4, p. 372, Burlington Gr.

parvulus, Newberry, 1875, Ohio Pal., vol.

2, p. 55, Cleveland shale.

pellensis, St. John & Worthen, 1883, Geo.

Sur. Ill., vol. 7, p. 237, St. Louis Gr.

pugiluculus, St. John & Worthen, 1875,

Geo. Sur. Ill., vol. 6, p. 430, St. Louis Gr.

sculptus, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 421, Waverly or

Kinderhook Gr.

similis, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 431, Kaskaskia Gr.

speciosus, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 424, Waverly or

Kinderhook Gr.

spectabilis, St. John & Worthen, 1875,

Geo. Sur. Ill., vol. 6, p. 420, Waverly or

Kinderhook Gr.

triangularis, Newberry, 1873, Ohio Pal.,

vol. 1, p. 329, Waverly Gr.

varians, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 422, Waverly or

Kinderhook Gr.

Fig. 1118.—*Clennacanthus triangularis*.

vetustus, Newberry, 1873, Ohio Pal., vol.

1, p. 326, Waverly Gr.

wrighti, Newberry, 1884, 35th Rep. N. Y.

Mus. Nat. Hist., p. 206, Ham. Gr.

CTENODUS, Agassiz, 1843, Recherches sur les

Poissons Fossiles, t. 3, p. 137. [Ety.

ktenos, comb; *odus*, tooth.] Tooth

somewhat fan-like, with closely ser-

rated edges, very porous and sulcated;

position in the jaw unknown. Type

C. *cristatus*.

dialophus, Cope, 1878,

Proc. Am. Phil. Soc.,

vol. 17, p. 528, in Pal.

Bull. No. 29, Permian.

fossatus, Cope, 1877,

Proc. Am. Phil. Soc.,

p. 54, Permian.

gurleianus, Cope, 1877,

Proc. Am. Phil. Soc.,

p. 55, Permian.

ohioensis, Cope, 1874,

Proc. Acad. Nat. Sci.

Phil., p. 91, and Ohio

Pal., vol. 1, p. 410,

Coal Meas.

periprion, Cope, 1878,

Pal. Bull. No. 29,

in Proc. Am. Phil. Soc., vol. 17, p. 527,

Permian.

Fig. 1119.—*Cteno-*
odus serratus.

porrectus, Cope, 1878, Pal. Bull. No. 29, in Proc. Am. Phil. Soc., vol. 17, p. 527, Permian.

psyllus, Cope, 1878, Pal. Bull. No. 26, in Proc. Am. Phil. Soc., vol. 17, p. 191, Permian.

reticulatus, Newberry, 1875, Ohio Pal., vol. 2, p. 60, Coal Meas.

serratus, Newberry, 1875, Ohio Pal., vol. 2, p. 59, Coal Meas.

CTENOPETALUS, Agassiz, 1869, Catal. Foss. Fish, Collection of Earl of Eoniskillen, in Geo. Mag., vol. 6. [Ety. *ktenos*, comb; *petalos*, broad, full-grown.] The serrated or denticulated crest distinguishes it from *Petalodus*, which it much resembles, and to which it bears about the same relation as *Petalodus* does to *Anthiodus*. Type C. *serratus*.
bellulus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 398, St. Louis Gr.
limatulus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 399, Kaskaskia Gr.

FIG. 1120.—*Ctenopetalus occidentalis*. Concave face.

medius, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 400, Kaskaskia Gr.

occidentalis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 401, Coal Meas.

vinosus, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 396, Keokuk Gr.

CTENOPTYCHIUS, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 99. [Ety. *ktenos*, comb; *ptyche*, wrinkle.] Teeth small, highly polished, strongly compressed, rounded or obtusely pointed; edge divided into several strong denticulations; base of crown with a few imbricating folds of ganoin; bony root, oblong, flattened in the same direction as the crown. Type C. *apicalis*.

cristatus, Dawson, 1868, Acad. Geol., p. 209, Coal Meas.

FIG. 1121.—*Ctenoptychius cristatus*. Natural size and magnified.

digitatus, Leidy, 1856, Trans. Am. Phil. Soc., vol. 11, St. Louis Gr.

pertenuis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 382, Kaskaskia Gr.

semicircularis, see *Peripristis semicircularis*.

stevensoni, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 383, Coal Meas.
CYMATODUS, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 363. [Ety. *cymatos*, wavy; *odous*, tooth.] Teeth small, oblong, or elliptical, thin, forming a flat or arched plate, of which the crown surface is transversely undulated and uniformly punctate; under surface flat, smooth, at the posterior end bearing a narrow, strap-shaped, oblique root. Type C. *oblongus*.
oblongus, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 364, Up. Coal Meas.

CYRTACANTHUS, Newberry, 1873, Ohio Pal., vol. 1, p. 306. [Ety. *kurtos*, curved; *akantha*, spine.] Spine curved, tuberculated, single row of conical denticles set on the posterior side, limited to the upper portion, and increasing in size from below upward. Type C. *dentatus*.
dentatus, Newberry, 1873, Ohio Pal., vol. 1, p. 307, Up. Held. Gr.

DACTYLODUS, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 33. [Ety. *daktylos*, finger; *odous*, tooth.] Crown as in *Petalodus*; root conspicuous, and divided into a number of radicles. Type D. *princeps*.

concaus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 390, St. Louis Gr.

excavatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 392, Kaskaskia Gr.

inflexus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 48, Kaskaskia Gr.

lobatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 47, St. Louis Gr.

minimus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 391, St. Louis Gr.

princeps, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 45, St. Louis Gr.

FIG. 1123.—*Dactylodus minimus*. Concave face.

& Worthen, 1875, p. 383, Coal Meas. & Worthen, 1870, 4, p. 363. [Ety. *ous*, tooth.] Teeth elliptical, thin, formula, of which the transversely unduly punctate; under h, at the posterior crow, strap-shaped, C. oblongus.

& Worthen, 1870, 4, p. 364, Up. Coal

FIG. 1122.—*Cyracanthus dentatus*.

& Worthen, 1875, 6, p. 392, Kaskas-

FIG. 1123.—*Dactylopus concavus*. Convex face.

391, St. Louis Gr. & Worthen, 1866, 2, p. 45, St. Louis Gr.

DELTOIDOPSIS, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 158. [Ety. from resemblance to Deltodus.] Coronal contour and general aspect near Deltodus, distinguished by the differentiation of the median ridge of the anterior coronal prominence, which approaches Cochliodus or Chitonodus. Type D. angusta. affinis, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 160, Warsaw Gr. angusta, Newberry & Worthen 1870, (D. Iodius angustus,) Geo. Sur. Ill., vol. 4, p. 368, Kaskaskia Gr. bialveatus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 169, Burlington Gr. convexa, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 169, Up. Burlington Gr. convoluta, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 163, Up. Burlington Gr. exornata, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 168, Warsaw Gr. inflexa, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 167, Keokuk Gr. keokuk, St. John & Worthen, Geo. Sur. Ill., vol. 7, p. 169, Keokuk Gr. stimplicatus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 161, St. Louis Gr. DELTODUS, Agassiz, 1859, MSS., and Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 93. [Ety. *della*, triangle; *odus*, tooth.] Teeth large, thick, strong, triangular, more or less arched, sometimes inrolled from the longer and more acute angle to the opposite margin; crown surface arched or marked by 1-3 prominent ridges from the basal margin toward the longer angle. Type D. sublaevis. *latus*, Newberry & Worthen, syn. for Chitonodus *latus*. *angularis*, Newberry & Worthen, syn. for Orthopleurodon carbonarius. *angustus*, see Deltodopsis *angusta*. *cinctus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 146, Warsaw Gr. *angulatus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 99, Kaskaskia Gr. *complanatus*, see Sandalodus *complanatus*. *fasciatus*, see Tæniodus *fasciatus*. *gandis*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 101, Keokuk Gr. Probably syn. for Sandalodus *levisimus*. *intermedius*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 153, St. Louis Gr. *latus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 145, Keokuk Gr. *litoni*, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 367, Subcarboniferous. *occidentalis*, Leidy, 1856, (Cochliodus *occidentalis*.) Trans. Am. Phil. Soc., vol. 11, p. 87, Warsaw and St. Louis Grs. *pavus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 151, St. Louis Gr. *powellii*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 154, Carboniferous.

propinquus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 56, Coal Meas. *rhomboides*, Newberry & Worthen, syn. for Sandalodus *spatulatus*. *spatulatus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 100, Burlington Gr. *stellatus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 97, Keokuk Gr. *trilobus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 148, Warsaw Gr. *undulatus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 98, Keokuk Gr. DELTOPTYCHUS, Agassiz, 1859, MSS., and St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 89. [Ety. *della*, triangle; *ptyx*, a wrinkle.] Posterior teeth of lower jaw trigonal, strongly built, and arched in the direction of inrollment; coronal contour in three divisions, narrowing toward the outer extremity; those of the upper jaw subspatulate, inrolled on the outer margin, acute posteriorly. Type D. *acutus*. *expansus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 98, St. Louis Gr. *nitidus*, Leidy, 1856, (Cochliodus *nitidus*.) Trans. Am. Phil. Soc., vol. 11, p. 87, Kaskaskia Gr. *primus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 93, Up. Burlington Gr. *varsoviensis*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 96, Warsaw Gr. *wachsmuthi*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 93, Keokuk Gr.

FIG. 1124.—*Delloptychus wachsmuthi*.

DESMODUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 337. [Ety. *desmos*, a ligament; *odus*, a tooth.] This name was applied to a genus of bats, in 1826, by Prinz, Neu. Wied. in Beitrage zur Naturg. Brasiliens. Teeth occurring in rows, small, robust; crown laterally elongated, arched vertically, median cusp with lateral crests; base constricted and produced. Type D. *tumidus*. *costelliferus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 341, St. Louis Gr. *flabellum*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 343, Keokuk Gr. *ligniformis*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 342, Keokuk Gr. *minusculus*, Newberry & Worthen, 1866, (Orodus *minusculus*.) Geo. Rep. Ill., vol. 2, p. 67, Keokuk Gr.

FIG. 1125.—*Desmodus costelliferus*. Convex aspect.

tumidus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 339, St. Louis Gr.
DINICHTHYS, Newberry, 1873, Ohio Pal., vol. 1, p. 313, and vol. 2, p. 3. [Ety. *deinos*, terrible; *ichthys*, a fish.] Cranium composed of thick bony plates, strengthened with internal arches ankylosed together, occipital bone in the type species three inches in thickness; relatively small maxillaries bearing a number of acute, conical, ankylosed teeth, which interlocked with a similar series on the mandibles; premaxillaries large, strong, triangular plates or teeth; mandibles of great length, flattened and spatulate behind, turning up anteriorly to form a strong triangular tooth, with its fellow of the opposite mandible, interlocked with the great, divergent, premaxillary teeth; vital parts of the body covered with large, thick plates which formed a carapace. Type *D. terrelli*.
hertzeni, Newberry, 1873, Ohio Pal., vol. 1, p. 316, Portage Gr.

FIG. 1126.—*Dinichthys hertzeni*.

terrelli, Newberry, 1873-75, Ohio Pal., vol. 1, p. 313, and vol. 2, p. 3, Portage Gr.
DIPLOSPIS, Matthew, 1888, Can. Rec. Sci., vol. 2, p. 251. [Ety. *diplos*, double; *aspis*, shield.] Small, having plates on the head, back, and sides, and one ventral plate; plates bearing very fine ridges. Type *D. acadica*.
acadica, Matthew, 1838, Can. Rec. Sci., vol. 2, p. 251, Up. Silurian or Low. Devonian.
DIPLODUS, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 204. [Ety. *diploos*, double; *odus*, a tooth.] This name was used by Rafinesque for a genus of Sparidae in 1810, Indice d'Litologia Siciliana. Teeth having a flattened or rounded base, from which spring two lateral and sometimes a small central denticle; each jaw bore several hundred teeth in radiating rows, the points projecting inward. They belong to sharks possessed of spines, described under the names of *Oracanthus* and *Xenacanthus*. Type *D. gibbosus*.
acinacica, Dawson, 1860, Acad. Geol., p. 211, and Can. Nat. Geol., vol. 5, Coal Meas.
compressus, Newberry, 1857, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 99, and Ohio Pal., vol. 1, p. 335, Coal Meas.
duplicatus, see *Thrinacodus duplicatus*.

gracilis, Newberry, 1857, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 99, and Ohio Pal., vol. 1, p. 335, Coal Meas.

FIG. 1127.—*Diploodus latus*.

incurvus, see *Thrinacodus incurvus*.
latus, Newberry, 1857, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 99, and Ohio Pal., vol. 1, p. 336, Coal Meas.
penetrans, Dawson, 1860, Acad. Geol., p. 211, and Can. Nat. and Geol., vol. 5, Coal Meas.
DIPTERUS, Sedgwick & Murchison, 1835, Geo. Trans., 2d ser., vol. 3. [Ety. *dipteros*, two-winged.] Diptera is an order of insects established by Linnaeus. Small fusiform fishes; heads compressed, tails heterocercal; two dorsal fins opposite two similar anal fins, the second of each the larger; a strongly marked lateral line; scales circular, thickest in the middle, variously curved with concentric lines or longitudinal ridges. Type *D. brachypterygius*.

FIG. 1128.—*Dipterus sherwoodi*.
sherwoodi, Newberry, 1875, Ohio Pal., vol. 2, p. 61, Catskill Gr.

DREPANACANTHUS, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 120. [Ety. *drepas*, a sickle; *akantha*, spine.] Fin spines compressed laterally, gradually tapering to an acute point, curved forward; anterior margin with a row of flattened or conical tubercles; lateral surfaces with tubercles in longitudinal rows; posterior margin without hooks, sometimes with tubercles. Type *D. gemmatulus*.
anceps, see *Xystacanthus anceps*.
gemmatulus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 123, Keokuk Gr.
reversus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 456, St. Louis Gr.
stellatus, see *Batacanthus stellatus*.
ECTOSTE-RACHIS, Cope, 1880, Pal. Bull. No. 32, p. 19. [Ety. *ektos*, without; *osteon*, bone; *rachis*, a ridge, backbone.] Base of the skull consists of ossified parachordals, which embrace the chorda dorsalis posteriorly, and are continued for a short distance posteriorly as a tube; anteriorly the chordal groove is open; trabeculae not ossified; cranial structure embryonic; above and in front of the opening for the chorda the neural canal enters the groove; parachordals subtriangular. Type *E. nitidus*.
ciceronius, Cope, 1883, Pal. Bull. No. 36, in Proc. Am. Phil. Soc., p. 628, Permian.

37, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., 1899, and Ohio Pal., 1899.

Meas.
us, Newberry, 1857, Proc. Acad. Nat. Sci. Phil., vol. 8, p. 350, and Ohio Pal., vol. 1, p. 336, Coal Meas.

trans, Dawson, 1860, Acad. Geol., p. 211, and Can. Nat. and Geol., vol. 5, Coal Meas.

us Murchison, 1835, ser., vol. 3, [Ety. d.] Diptera is an ablished by Linneus. shes; heads com- roceral; two dorsal

simi- the gly line; ckect vari- con- ongi- Type us.

erry, Fig. 1128.—Dipte- ol. 2, rus sherwood.

Worthen, vol. 2, p. 120. [Ety. *akantha*, spine.] Fin laterally, gradually te point, curved for- argin with a row of al tubercles; lateral ercles in longitudinal argin without hooks, tubercles. Type D.

anthus anceps.

ry & Worthen, 1866, 2, p. 123, Keokuk Gr. Worthen, 1875, Geo. 456, St. Louis Gr. thus stellatus.

us, 1880, Pal. Bull. No. 19, Permian. [Ety. *edentes*, a devourer.] Maxillary bone segmented; segments beveled anteriorly and excavated posteriorly for co-adaptation; teeth resembling those of *Carcharodon*, one co-ossified with each maxillary segment. Type E. vorax.

nitidus, Cope, 1880, Pal. Bull. No. 32, p. 19, Permian.

Edestus, Ledy, 1856, Jour. Acad. Nat. Sci., 2d ser., vol. 3, p. 159. [Ety. *edentes*, a devourer.] Maxillary bone segmented; segments beveled anteriorly and excavated posteriorly for co-adaptation; teeth resembling those of *Carcharodon*, one co-ossified with each maxillary segment. Type E. vorax.

giganteus, Newberry, 1888, Ann. N. Y. Acad. Sci., vol. 4, p. 1, Coal Meas.

heinrichsi, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 350, Coal Meas.

minor, Newberry, 1866, Geo. Sur. Ill., vol. 2, p. 84, Coal Meas.

vorax, Ledy, 1856, Jour. Acad. Nat. Sci. Phil., vol. 3, 2d series, p. 159, Coal Meas.

FIG. 1129.—*Edestus vorax*.

Elonichthys peltigerus, see *Paleoniscus peltigerus*.

ERISMACANTHUS, McCoy, 1848, Ann. and Mag. Nat. Hist., 2d series, vol. 2, p. 119.

[Ety. *ereisma*, a prop or stay; *akantha*, spine.] Spine of three parts; one compressed, finely striated, which entered the flesh; the second short, compressed, rapidly tapering, curved backward, sides with longitudinal ridges, and two rows of downward curved teeth on the posterior concave margin; the third, a prop-like part extending forward nearly at right angles with the base, arched, compressed at the basal half, depressed distally, and covered with tubercles and some spines on the under side. Type E. jonesi.

maccoyanus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 461, St. Louis Gr.

EUNEMACANTHUS, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 120. [Ety. *eu*, beautiful; *nema*, a line; *akantha*, spine.] Distinguished from *Ctenacanthus* by the plain dorsal ridge, tuberculated intercostal sulci, and upward direction of the denticles on the angles of the posterior face. Type E. costatus.

costatus, Newberry & Worthen, 1866, (Ctenacanthus costatus), Geo. Sur. Ill., vol. 2, p. 120, St. Louis Gr.

Eurylepis, Newberry, 1856, Proc. Acad. Nat. Sci. Phil. [Ety. *eury*, broad; *lepis*,

scale.] Small; body fusiform; head obtuse; tail elongated, lobes unequal; fins small, with delicate fulcra; dorsal and anal fins opposite, and far back on the body; ventrals near middle of abdomen; cranial surface tubercular; maxillary, mandibular, and jugular plates corrugated; scales smooth, ornamented, or serrated; teeth numerous, conical, short. Type E. tuberculata.

corrugata, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., vol. 1, p. 350, Coal Meas.

granulata, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., vol. 1, p. 352, Coal Meas.

insculpta, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., vol. 1, p. 351, Coal Meas.

lineata, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., vol. 1, p. 353, Coal Meas.

minima, Newberry, 1873, Ohio Pal., vol. 1, p. 353, Coal Meas.

ornatissima, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., vol. 1, p. 352, Coal Meas.

ovoidea, Newberry, 1856, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., vol. 1, p. 351, Coal Meas.

striolata, Newberry, 1873, Ohio Pal., vol. 1, p. 355, Coal Meas.

tuberculata, Newberry, 1856, Proc. Acad. Nat. Sci., and Ohio Pal., vol. 1, p. 350, Coal Meas.

FIG. 1130.—*Eurylepis tuberculata*.

EUSTHENOPTERON, Whiteaves, 1881, Am. Jour. Sci. and Arts, 3d ser., vol. 21, p. 495. [Ety. *eu*, very; *sthenes*, stout; *pteron*, a fin.] Fin rays of anal and second dorsal fins supported by three osselets articulated to a broad inter-spinous apophysis; vertebral centers not ossified; caudal osselets articulated to modified haemal spines. Type E. foordi.

foordi, Whiteaves, 1881, Am. Jour. Sci. and Arts, 3d ser., vol. 21, p. 495, Upper Devonian.

FISSODUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 413. [Ety. *fissus*, split; *odus*, tooth.] Teeth small, in the form of root and general contour

like *Peltodus*, but distinguished by the cleft condition of the crest. Type *F. bifidus*.

bifidus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 414, Kaskaskia Gr.

FIG. 1131.—*Peltodus bifidus*.

Gampsacanthus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 471. [Ety. *gampsos*, curved; *akantha*, spine.] Spines long, laterally compressed, tapering, costate, with larger and smaller tubercles; posterior margin denticulate; base expanded; pulp cavity large. Type *G. typus*.

FIG. 1132.—*Gampsacanthus typus*. Side view of a spine magnified 2 diam., and transverse section.

latus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 474, Keokuk Gr.

squamosus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 473, St. Louis Gr.

typus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 472, St. Louis Gr.

Gibacanthus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 440. [Ety. *geison*, a border; *akantha*, a spine.] Spine curved posteriorly, anterior angle a simple raised keel; lateral faces bearing longitudinal rows of tubercles; posterior face longitudinally keeled. Type *G. stellatus*.

bullatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 441, Kaskaskia Gr.

stellatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 440, St. Louis Gr.

FIG. 1133.—*Glymmatacanthus irishii*. Fragment of spine.

Glymmatacanthus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 446. [Ety. *glymmatos*, engraved; *akantha*, spine.] Fin ray vertically elongated, posteriorly arched, laterally compressed; lateral faces covered with stellate or striated tubercles. Type *G. irishii*.

irishii, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 447, Kinderhook or Waverly Gr.

petrodoides, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 250, Kaskaskia Gr.

rudis, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 249, Keokuk Gr.

Glyptrolepis, Agassiz, 1836, Poiss. Foss., vol. 2, p. 179. [Ety. *glyptos*, sculptured; *lepis*, scale.] Fins long, sometimes pendulous; anterior dorsal opposite ventral, and posterior dorsal opposite anal; tail fin long, spreading below; shoulder bones huge; teeth minute; scales of great size in proportion to the animal, and deeply sculptured. Type *G. elegans*.

microlepidotus, Agassiz, 1836, Poiss. Foss., vol. 2, p. 179, Devonian.

quebecensis, Whiteaves, 1880, Trans. Roy. Soc. Can., vol. 6, p. 77, Low. Devonian.

Gnathorhiza, Cope, 1883, Proc. Am. Phil. Soc., vol. 20, p. 629. [Ety. *gnathos*, jaw; *rhiza*, root.] Founded upon some ganoid teeth. The definition is too meager for identification, and the genus may never again be recognized. Type *G. serrata*.

serrata, Cope, 1883, Proc. Am. Phil. Soc., vol. 20, p. 629, Permian.

Gyracanthus, Agassiz, 1833, Recherches sur les Poissons Fossiles, t. 1, p. 87.

[Ety. *gyros*, a circle; *akantha*, spine.] Fin spines very large, gradually tapering to the apex, and slightly arched backward; inserted base small, rapidly tapering; posterior margin feebly armed with two rows of small denticles; surface of the sides covered with very oblique ridges, which meet at an angle on the anterior face. Type *G. formosus*.

alleni, Newberry, 1873, Ohio Pal., vol. 1, p. 331, Cuyahoga shale.

compressus, Newberry, 1873, Ohio Pal., vol. 1, p. 230, Cuyahoga shale.

cordatus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 251, Keokuk Gr.

duplicatus, Dawson, 1868, Acad. Geol., p. 210, Coal Meas.

magnificus, Dawson, 1868, Acad. Geol., p. 210, Sub-carboniferous.

Harpacodus, Agassiz, 1869, Catal. Foss. Fish. Collection of Earl of Enniskillen, and St. John & Worthen, in Geo. Sur. Ill., vol. 6, p. 354. [Ety. *harpe*, a

FIG. 1134.—*Gyracanthus allenii*. Anterior face.

orthen, 1875, Geo.
447, Kinderhook or

& Worthen, 1883,
7, p. 250, Kaskas-

orthen, 1883, Geo.
49, Keokuk Gr.

1836, Poiss. Foss.,
glyptos, sculptured;

ng, sometimes pen-
soral opposite ven-

orsal opposite anal;
ng below; shoulder

minute; scales of
rtion to the animal,

ured. Type G. el-

gassiz, 1836, Poiss.
170, Devonian.

teaves, 1880, Trans.
vol. 6, p. 77, Low.

e, 1883, Proc. Am.
20, p. 620. [Ety.

iza, root.] Founded
ine teeth. The def-

esiger for identifica-
nus may never again

Type G. serrata.

1883, Proc. Am. Phil.
629, Permian.

z, 1833, Recherches
Fossiles, t. 1, p. 87.

; akan-
spines

ally ta-
x, and

back-
base

bering;
feebly

ows of
surface

d with
which

on the
pe G.

, 1873,
p. 331,

wberry,
ol. 1, p.

ale.
& Wor-

ur. Ill.,
nk Gr.

, 1868,
0, Coal

, 1868,
0, Sub-

, 1860,
 Collec-

nniskil-
karl-

& Worthen, in Geo.
p. 354. [Ety. harpe, a

hook; *odous*, tooth.] Teeth laterally elongated, vertically arched, gently curved outward in the concave face; margins nearly parallel; crown compressed along the crest; serrated, expanded below; convex face low, opposite face concave; coronal borders produced inbeveled; base strong, obliquely produced; lateral angles well defined. Type H. dentatus, or more properly, H. occidentalis.

FIG. 1135.—Harpe-
codius occiden-
talis.

compactus, St. John &
Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 355,
Kaskaskia Gr.
occidentalis, St. John &
Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 355,
St. Louis Gr.

HELIODUS, Newberry, 1875, Ohio Pal., vol. 2, p. 62. [Ety. *helios*, sun; *odous*, tooth.] Distinguished from *Dipterus* by having the upper palato teeth united, forming a rounded, semicircular, triturating plate, bearing radiating tuberculated ridges. Type H. lesleyi. Dr. Traquair, of England, regards *Heliodus* as a synonym for *Palsadaphus*, Van Beneden & DeKoninck, 1864, Bull. Acad. Belg., vol. 17, p. 143.

FIG. 1136.—*Heliodus lesleyi*.

lesleyi, Newberry, 1875, Ohio Pal., vol. 2, p. 64, Chemung Gr.

HELODUS, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 104. [Ety. *helos*, a nail or rudder; *odous*, tooth.] Transversely elongate, crown convex, elevated along the middle into an obtuse, circular ridge, sometimes divided into a line of several compressed cones diminishing from the center; surface porous as in *Psammodus*; margin of the crown raised in the middle on both the inner and outer sides, and it and the root vertically plicated. Type H. simplex.

angulatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 83, Burlington Gr. biformis, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 77, Waverly or Kinderhook Gr.

carbonarius, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 75, Coal Meas.

compressus, Newberry & Worthen, 1870,

Geo. Sur. Ill., vol. 4, p. 360, Burlington Gr. *compressus*, see *Hybocladodus compressus*.

coniculus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 75, Burlington Gr.

consolidatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, syn. for *Chitonodus latus*.

crenulatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 82, Keokuk Gr.

denhumani, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 76, Keokuk Gr.

denticulatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 81, Keokuk Gr.

elytra, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 78, Keokuk Gr.

gibbosus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 79, Keokuk Gr.

gibbus, Leidy, 1856, Trans. Am. Phil. Soc., vol. 11, p. 87, Keokuk Gr.

limax, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 80, Burlington Gr.

nobilis, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, same as *Chitonodus latus*.

placenta, see *Psephodus placenta*.

politus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 79, Keokuk Gr.

rugosus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 4, p. 359, Coal Meas.

sulcatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 82, Keokuk Gr.

undulatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 82, Keokuk Gr.

HOLOTRYCHIUS, Agassiz, 1836, Recherches sur les Poissons Fossiles, t. 2, p. 179.

[Ety. *holos*, entire; *ptyx*, wrinkle.] Body thick, short, rounded, bones of the head granulated; scales large, very thick, subrhomboidal, rounded, imbricating, composed of numerous bony layers, exposed surface marked with large, longitudinal, flexuous wrinkles and tubercles; teeth small, numerous, conical, longitudinally sulcated at base; tail heterocerical, caudal fin triangular, obliquely truncated; dorsal fin opposite a similar anal one close to the base of the caudal; ventral behind the middle of the body. Type H. gigantens.

americanus, Leidy, 1856, Jour. Acad. Nat. Sci., 2d series, vol. 3, p. 169, Catskill Gr.

nobilissimus, Agassiz, as identified by Hall, 1843, Geo. Rep. 4th Dist. N. Y., is described as H. americanus.

taylori, Hall, 1843, (Sauripteria taylori), Geo. Rep. 4th Dist. N. Y., p. 282, Catskill Gr.

FIG. 1138.—Holotrychius americanus. Single tooth.

FIG. 1137.—*Helodus gibbosus*.

FIG. 1134.—Gyracanthus alieni. Anterior face.

Homacanthus, Agassiz, 1845, Poiss. Foss. [Ety. *homos*, similar; *akantha*, spine.]

gibbosus, see *Amacanthus gibbosus*.

gracilis, Whiteaves, 1880, Trans. Roy. Soc. Can., vol. 6, p. 77, Low. Devonian.

rectus, see *Marracanthus rectus*.

Hybocladodus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 284. [Ety. *hybos*, hump; *Cladodus*, a genus.] Teeth small, strongly cuspidate, base resembling that of a *Cladodus*, being elliptical and broadly expanded, with a more or less prominent antero-posteriorly compressed median cone, both surfaces of which are plicated and resemble the crown of a *Hybodus*; anterior face nearly straight, curved laterally, terminating below in a well defined marginal border or ridge, posterior margin broadly rounded, inferior surface excavated immediately behind the marginal border, with a beveled space extending along the posterior margin, superior face more or less convex and beveled to the posterior edge; both coronal surfaces vertically marked with plicæ. Type *H. plicatilis*.

compressus, Newberry & Worthen, 1866, (*Helodus compressus*.) Geo. Sur. Ill., vol. 2, p. 78, Burlington Gr.

intermedius, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 237, Keokuk Gr.

FIG. 1139.—*Hybocladodus plicatilis*.

nitidus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 288, Kaskaskia Gr.

plicatilis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 286, Burlington Gr.

tenuicostatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 286, Keokuk Gr.

JANASSA, Munster, 1839, Beitrage Petrefaktenkunde, vol. 1, and Agassiz in Poiss. Foss., t. 3, p. 375. [Ety. mythological name.] Teeth have a tabulated structure and enameled, wavy crown; small in front and larger toward the posterior part of the jaw; jaw-bone rough and granular. Type *J. angulata*.

gurliana, Cope, 1877, (*Strigillina gurliana*), Proc. Am. Phil. Soc., p. 191, Permian.

linguiformis, Cope, 1877, (*Strigillina linguiformis*), Proc. Am. Phil. Soc., p. 53, Permian.

LAMBODUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 280. [Ety. *Lambda*, a Greek letter; *odous*, tooth.] Teeth small, base posteriorly produced and laterally expanded, broadest behind the cornua; a single strong, slightly sigmoidally curved, recurved, eccentric cornua arises from the anterior angle of the base, terminates in a sharp apex, compressed in front, broadly rounded behind, with more or less distinct cutting edges and vertical costæ. It is distinguished from *Cladodus* by the

single coronal cornua, and the absence of lateral denticles; the basal portion bears some resemblance to *Thrinacodus*.

Type *L. costatus*.

calceolus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 281, Burlington Gr.

calceolus var. *robustus*, St. John & Worthen, 1866, Geo. Sur. Ill., vol. 6, p. 282, Keokuk Gr.

costatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 280, Burlington Gr.

FIG. 1140.—*Lambodus costatus*.

hamulus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 283, Kaskaskia Gr.

reflexus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 284, Kaskaskia Gr.

transversus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 282, St. Louis Gr.

LEACRANTHUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 475. [Ety.

lekroi, the antlers of a stag; *akantha*, spine.] Spines long, tapering, curved, laterally compressed, stellate tubercles irregularly disposed; base thin, expanded; pulp cavity large; apex transversely expanded and armed with strong denticles. Type *L. unguiculus*.

unguiculus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 476, St. Louis Gr.

Leptacanthus, Agassiz, 1837, Poiss. Foss., vol. 3. [Ety. *leptos*, slender; *akantha*, spine.] *occidentalis*, see *Aconylacanthus occidentalis*.

Lionus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 335. [Ety. *leios*, smooth; *odous*, tooth.] Teeth resembling *Orodus*; crown arched, laterally and vertically; basal margins constructed and sharply defined from the base; apex with obscurely defined lateral crests; convex in either face; anterior face produced beneath the median cone, and both faces occupied with faint vertical sulci, producing obscure secondary prominences; surface smooth, punctate, or verrucose; base as in *Orodus*, relatively deep. Type *L. calcaratus*.

calcaratus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 336, Burlington Gr.

calcaratus var. *gros-sipunctatus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 337, Keokuk Gr.

LIognathus, Newberry, 1873, Ohio Pal., vol. 1, p. 306. [Ety. *lis*, smooth; *gnathos*, the jaw.] Jaw the only part yet known; spatulate, dentate only at and near the anterior extremity; resembles *Coccoosteus*. Type *L. spatulatus*.

spatulatus, Newberry, 1873, Ohio Pal., vol. 1, p. 306, Up. Held. Gr.

FIG. 1141.—*Liodus calcaratus*.

Liodus calcaratus.

Liodus calcaratus.

Liodus calcaratus.

Liodus calcaratus.

Liodus calcaratus.

ua, and the absence
; the basal portion
ance to *Thrinacosodus*.

Worthen, 1875, Geo.
281, Burlington Gr.
us, St. John & Wor-
r. Ill., vol. 6, p. 282.

Wor-
r. Ill.,
riling-

Fig. 1140.—*Lamb-
dodus costatus*.
Sur.

Kaskaskia Gr.
Worthen, 1875, Geo.
284, *Kaskaskia* Gr.
hn & Worthen, 1875,
6, p. 282. *St. Louis* Gr.
hn & Worthen, 1875,
6, p. 475. [Ety.
s of a stag; *akantha*,
long, tapering, curved,
sed, stellate tubercles
sed; base thin, ex-
vity large; apex trans-
and armed with strong
L. unguiculus.
in & Worthen, 1875,
vol. 6, p. 476, St.

1837, Poiss. Foss., vol.
ender; *akantha*, spine.]
ndylacanthus occiden-

Worthen, 1875, Geo.
p. 335. [Ety. *leios*,
tooth.] Teeth resem-
rown arched, laterally
basal margins com-
rply defined from the
th obscurely defined
convex in either face;
roduced beneath the
d both faces occupied
al sulci, producing ob-
prominences; surface
e, or verrucose; base as
tively deep. Type L.

calcaratus, St. John
& Worthen, 1875,
Geo. Sur. Ill., vol.
6, p. 336, Bur-
lington Gr.
calcaratus var. *gros-*
sipunctatus, St.
John & Worthen,
Ill., vol. 6, p. 337, Keo-

erry, 1873, Ohio Pal.,
Ety. *lis*, smooth; *gnathos*,
w the only part yet
dentate only at and
or extremity; resembles
Type L. *spatulatus*.
erry, 1873, Ohio Pal., vol.
feld. Gr.

Lisogodus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 363. [Ety. *ligos*, a
spade; *odus*, tooth.] Teeth laterally

FIG. 1142.—*Lisogodus spatulatus*.

abbreviated strong; crown thick, sharp-
created, and sometimes obscurely ser-
rated; basal margins well defined; base

FIG. 1145.—*Machæracanthus peracutus*.

vertical to the crown, rectangular,
prolonged, equal to the elevation of the
crown; inferior surface well defined
from either face above; and generally
slightly beveled from the concave to the
opposite border; coronal surface ena-
meled, worn crest striato-punctate.
Type L. *curtus*.

FIG. 1143.—*Lisogodus curtus*.

curtus, St. John & Wor-
then, 1875, Geo. Sur. Ill.,
vol. 6, p. 364, Burlington
Gr.
selliniformis, St. John &
Worthen, 1875, Geo. Sur.
Ill., vol. 6, p. 366, St.

Louis Gr.
serratus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 365, Burlington Gr.
Listracanthus, Newberry & Worthen, 1870,
Geo. Sur. Ill., vol. 4, p. 371. [Ety. *list-*
ron, shovel; *akantha*, spine.] Spines
small, gently arched, flattened, thin;
sides marked by numerous sharp, lon-
gitudinal carinae, edges set with diver-
gent, slender, acute teeth; most numer-
ous on the convex margin; and largest
base expanded and obliquely truncated.
Type L. *hystrix*.

hildrethi, New-
berry, 1875,
Ohio Pal.,
vol. 2, p. 56,
Coal Meas.

hystrix, New-
berry & Wor-
then, 1870, Geo. Sur. Ill., vol. 4, p. 372,
Coal Meas.

Lophodus, Newberry & Worthen, 1870, Geo.
Sur. Ill., vol. 4. This name was preoc-
cupied by Romanowsky in 1864.

variabilis, see *Agassizodus variabilis*.
Machæracanthus, Newberry, 1857, Bull.
Nat. Inst., p. 6, and Ohio Pal., vol. 1,

p. 302. [Ety. *machaira*, a saber; *akan-*
tha, a spine.] Spines large, flattened,
curved, aneplital, unsymmetrical; edges
and point acute; base narrowed, with a
rough and irregular extremity; central
cavity reaching nearly to the apex; ex-
ternal surface enameled, smooth or
punctate, and striate microscopic struc-
ture dense. Type M. *major*.

major, Newberry, 1857, Bull. Nat. Inst.,
p. 6, and Ohio Pal., vol. 1, p. 304, Up-
held. Gr.

peracutus, Newberry, 1857, Bull. Nat.
Inst., p. 6, and Ohio Pal., vol. 1, p. 305,
Up. Held. Gr.

sulcatus, Newberry, 1857, Bull. Nat. Inst.,
p. 6, and Ohio Pal., vol. 1, p. 305, Up-
held. Gr.

MACROPETALICHTHYS,
Norwood & Owen,
1846, Am. Jour. Sci.,
2d ser., vol. 1, p.
367. [Ety. *makros*,
large; *petalos*, ex-
panded or spread
out; *ichthys*, fish.]
Cranium composed
of large polygonal
plates, united by
double sutures; sur-
face enameled, tu-
bercle, ornament-
ed; eye orbits con-
spicuous; nasal plate wedge-shaped;
occipital plate oblong, emarginate be-
hind, and prolonged anteriorly, where
it meets the nasal plate. Type M.
rapheidolabis.

FIG. 1140.—*Macropetalichthys sullivanti*. One-fifth natural size.

manni, Newberry, (*Agassichthys manni*)
1857, Bull. Nat. Inst., p. 3, Up. Held. Gr.
rapheidolabis, Norwood & Owen, 1846,
Am. Jour. Sci., 2d ser., vol. 1, p. 367,
Up. Held. Gr.

sullivanti, Newberry, 1857, (*Agassichthys*
sullivanti,) Bull. Nat. Inst., p. 3, and
Ohio Pal., vol. 1, p. 294, Up. Held. Gr.
MARRACANTHUS, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 465. [Ety. *mar-*
ron, a spade; *akantha*, spine.] Dorsal
spine nearly straight, or with a forward
curvature, obtusely terminated, rounded
in front, truncated behind, or rounded
into the posterior face, which is longi-
tudinally ridged in apparent continu-
ity with the lateral costæ; lateral face
and anterior margin longitudinally
ridged, the costæ being tuberculated,
those in front more or less strongly

FIG. 1144.—*Listracanthus hystrix*.

developed, with their apices directed upward, and especially in their upper part, where they gradually increase in size, forming strong, more or less deflected hooks, transversely carinated; intercostal spaces minutely ridged and striato-punctate; base moderately inserted, forming a comparatively thin plate, more or less laterally expanded posteriorly from the angular ridge in front, with more or less prominent marginal angles behind; pulp-cavity moderately large, similar in section to the body, and occupying the posterior two-thirds of the spine. In costation and the expanded base it is like *Amacanthus*, but distinguished in all other respects. Type *M. rectus*.

FIG. 1147.—*Marracanthus rectus*. Anterior part of spine.

rectus, Newberry & Worthen, 1866, (*Homacanthus* (?) *rectus*), *Geo. Sur. Ill.*, vol. 2, p. 115, St. Louis Gr.

Mecolepis, Newberry, 1857, *Proc. Acad. Nat. Sci.*, vol. 8, p. 96. [Ety. *mekos*, large; *lepis*, a scale.] Heterocerac lepidoids of small size; body fusiform; head obtuse; tail elongated; lobes unequal; fins small, provided with delicate fulcra; dorsal opposite anal, both far back on the body; crania corrugated or tuberculated; opercular maxillary and hyoid plates ornamented; scales smooth or ornamented; posterior margin serrated; scales of median line crenulated; two rows of scales extending back to near anal fin; teeth conical, short, brush-like. Distinguished from *Palaeoniscus* by small size, posterior position of dorsal fin, and the high lateral scales. Type *M. corrugata*. Probably a syn. for *Palaeoniscus*, but not figured, and species poorly defined.

corrugata, Newberry, 1856, *Proc. Acad. Nat. Sci.*, vol. 8, p. 96, Coal Meas.

granulata, Newberry, 1856, *Proc. Acad. Nat. Sci.*, vol. 8, p. 97, Coal Meas.

insculpta, Newberry, 1856, *Proc. Acad. Nat. Sci.*, vol. 8, p. 97, Coal Meas.

lineata, Newberry, 1856, *Proc. Acad. Nat. Sci.*, vol. 8, p. 97, Coal Meas.

ornatissima, Newberry, 1856, *Proc. Acad. Nat. Sci.*, vol. 8, p. 97, Coal Meas.

ovoidea, Newberry, 1856, *Proc. Acad. Nat. Sci.*, vol. 8, p. 97, Coal Meas.

serrata, Newberry, 1856, *Proc. Acad. Nat. Sci.*, vol. 8, p. 97, Coal Meas.

tuberculata, Newberry, 1856, *Proc. Acad. Nat. Sci.*, vol. 8, p. 96, Coal Meas.

Mesomodius, St. John & Worthen, 1875, *Geo. Sur. Ill.*, vol. 6, p. 290. [Ety. *mesodme*, something between; *odous*, tooth.] Teeth laterally elongated; base

consisting of one inferior flattened; posterior obliquely produced, massive plate, of which the posterior face slopes downward and slightly backward, at an obtuse angle, to the posterior crown face; anterior face slightly produced along the shoulder, which extends parallel with the base of the crown, vertical or beveled, and occupied by a more or less prominent median protuberance, which extends to the edge of the inferior surfaces; both faces are more or less roughened or pitted, lateral angles truncated or rounded, and more or less constricted above, equaling the lateral diameter of the crown; crown rising along the anterior border, sharply constricted in front and laterally, and well defined, sometimes constricted from the posterior basal face, nearly equaling the base in antero-posterior diameter; but more or less compressed along the crest, which rises into a more or less prominent median or submedian cusp, vertical or laterally deflected and recurved, usually compressed, with distinct, sometimes sharp, cutting edges; the lateral portions of the crown denticulated, extremities bearing slightly more prominent cusps than intermediate spaces; both faces ridged vertically; outer face of median cone often strongly buttressed; coronal surface enameled. Type *M. exculptus*.

explanatus, St. John & Worthen, 1875, *Geo. Sur. Ill.*, vol. 6, p. 293, Waverly or Kinderhook Gr.

exculptus, St. John & Worthen, 1875, *Geo. Sur. Ill.*, vol. 6, p. 291, Waverly or Kinderhook Gr.

ornatus, St. John & Worthen, 1875, *Geo. Sur. Ill.*, vol. 6, p. 294, Burlington Gr.

MYCTEROPS, Cope, 1888, *Am. Nat.*, p. 876. [Ety. *mukter*, nose; *ops*, eye.] Founded upon the cast of the cranial and nuchal buckler of a placoderm fish; the eye-holes resemble those of *Cephalaspis*, and they are separated by a nose-hole, which is divided by a narrow bridge. Type *M. ordinata*.

ordinata, Cope, 1888, *Am. Nat.*, p. 876, Coal Meas.

Onchus, Agassiz, 1837, *Recherches sur les Poissons Fossiles*. [Ety. *onchos*, bent, hooked like a talon or arrow-barb.]

deweyi, see *Ceraticaris deweyi*.

ONYCHOLUS, Newberry, 1857, *Bull. Nat. Inst.*, p. 5, and Ohio Pal., vol. 1, p. 296. [Ety. *onyx*, a claw; *odous*, tooth.] Cranium composed of a great number of plates covered with an enameled and tuberculated surface; jaws set with numerous conical, acute, recurved teeth; maxillary forming a low trian-

inferior flattened; produced, massive posterior face slopes slightly backward, at an angle the posterior crown slightly produced; which extends part of the crown, vertex and occupied by a prominent median process extends to the edge of the face; both faces are thickened or pitted, laterally or rounded, and restricted above, equal diameter of the crown; the anterior border, rounded in front and laterally, sometimes converging posterior basal face, the base in anterior; but more or less the crest, which rises as prominent median process, vertical or laterally curved, usually conical, sometimes sharp, the lateral portions of the mandible, extremities more prominent cusps the spaces; both faces the outer face of median buttressed; coronal. Type M. exculptus.

St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 293, Waverly or

FIG. 1148.—Mesododus exculptus.

St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 294, Burlington

1888, Am. Nat., p. 876, [Ety. *onchos*, bent, on or arrow-barb.]

Paris deweyi. Newberry, 1857, Bull. Nat. Inst., p. 5, Up. Held. Gr.

Ohio Pal., vol. 1, p. 1, a claw; *odous*, tooth.]

bed of a great number of teeth with an enameled surface; jaws set with small, acute, recurved teeth forming a low triangular

gle; dentary bones posteriorly acute, where they are overlapped by the articular portions of the mandibles, long and narrow, curving upward to the

FIG. 1149.—Mycterops ordinata.

symphysis, where they support an inter-mandibular arch of bone, to which is attached a series of large, curved, conical teeth; body covered with imbricated circular scales. Type O. sigmoides.

FIG. 1150.—Onychodus sigmoides. One-half nat. size of inter-mandibular crest with 6 teeth.

hopkinsi, Newberry, 1857, Bull. Nat. Inst., p. 5, Chemung Gr. sigmoides, Newberry, 1857, Bull. Nat. Inst., p. 5, and Ohio Pal., vol. 1, p. 299, Up. Held. Gr.

FIG. 1151.—Onychodus sigmoides. Fragment of the right mandible.

ORACANTHUS, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 13. [Ety. *orais*, beautiful; *akantha*, spine.] Dorsal rays large, conical, without solid base, hollow, walls thin, surface tuberculated; no posterior rows of denticles. Type O. milleri.

abbreviatus, Newberry, 1857, Bull. Nat. Inst., p. 5, Up. Held. Gr.

consimilis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, syn. for O. vetustus.

fragilis, Newberry, 1857, Bull. Nat. Inst., p. 5, Up. Held. Gr.

granulatus, Newberry, 1857, Bull. Nat. Inst., p. 5, Up. Held. Gr.

multiseriatus, Newberry, 1857, Bull. Nat. Inst., p. 5, Up. Held. Gr.

(?) obliquus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 477, Keokuk Gr.

pnigeus. This species is made the type of the genus *pnigeacanthus*. See P. *deltoides*.

rectus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 257, Kaskaskia Gr.

vetustus, Leidy, 1856, Jour. Acad. Nat. Sci. Phil., 2d ser., vol. 3, p. 162, St. Louis Gr.

ORODUS, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 97. [Ety. *orais*, beautiful; *odous*, tooth.] Teeth laterally elongated, middle more elevated than extremities, forming an obtuse transverse cone; longitudinal diameter greatest and marked by a medial ridge with oblique secondary ridges. Type O. cinctus.

alleni, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 310, Coal Meas.

carinatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 307, Keokuk Gr.

corrugatus, see Agassizodus corrugatus.

dædaleus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 301, Waverly or Kinderhook Gr.

decussatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 300, Waverly or Kinderhook Gr.

elegantulus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 64, Burlington Gr.

fastigiatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 306, Burlington Gr.

major, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 302, Burlington Gr.

mammillaris, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 66, Keokuk Gr.

minusculus, see Desmiodus minusculus.

minutus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 68, Keokuk Gr.

multicarinatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 62, Waverly or Kinderhook Gr.

neglectus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 308, St. Louis Gr.

ornatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 65, Keokuk Gr.

parallelus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 295, Waverly or Kinderhook Gr.

parvulus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 309, St. Louis Gr.

plicatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 63, St. Louis Gr.

tuberculatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 66, Burlington Gr.

Waverly or Kinderhook Gr.

Waverly or Kinderhook Gr.

Waverly or Kinderhook Gr.

Waverly or Kinderhook Gr.

Waverly or Kinderhook Gr.

Waverly or Kinderhook Gr.

Waverly or Kinderhook Gr.

FIG. 1152.—Orodus mammillaris.

turgidus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 310, Kaskaskia Gr.

FIG. 1153.—*Orodus variabilis*.

variabilis, Newberry, 1875, Ohio Pal., vol. 2, p. 50, Waverly Gr.

varicosatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 304, Burlington Gr.

whitii, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 297, Waverly or Kinderhook Gr.

ORTHACANTHUS, Agassiz, 1843, Poiss. Foss., t. 3, p. 330, [Ety. *orthos*, straight; *akantha*, spine.] Spines straight or gently curved; two or more rows of denticles on the posterior face. Type *O. cylindricus*.

FIG. 1154.—*Orthacanthus gracilis*.

arcuatus, Newberry, 1857, (Pleuracanthus arcuatus,) Proc. Acad. Nat. Sci. Phil., p. —, and Ohio Pal., vol. 1, p. 332, Coal Meas.

gracilis, Newberry, 1875, Ohio Pal., vol. 2, p. 56, Coal Meas.

quadriseptatus, Cope, 1877, Pal. Bull. No. 26, in Proc. Am. Phil. Soc., vol. 17, p. 192, Permian.

ORTHOPLEUROS, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 190. [Ety. *orthos*, straight; *pleuron*, side; *odus*, tooth; in allusion to the straight postero-lateral border of the maxillary posterior tooth.] Posterior teeth of upper jaw subspatulate in outline; postero-lateral border straight, or nearly so, and probably gently curved downward and inward at the outer extremity, and gently arched in the same direction, terminating posteriorly in an acute angle or spur, whence the inner margin, which is greatly thickened or massive, is broadly rounded into and merges with the thin antero-lateral border toward the extremity; coronal surface occupied by a prominent principal fold or ridge rising nearest the straight border, and flanked on the anterior slope by an obscure secondary ridge; the punctate enamel forms a narrow fold along the thickened straight border; teeth supposed to have occupied a similar position on the mandibles, distinguished by their trigonal outline, somewhat strong and spiral inrollment of the extremity, toward which the antero and postero-lateral borders regularly converge, inner margin more or less obliquely rounded, and sigmoidally curved from front toward the posterior angle; coronal surface pre-

sented a more or less well-defined plane; anterior fold, abruptly broken down on that side, where the coronal enamel forms a wide belt sharply defined from the deep basal rim, and limited behind by the more or less deep longitudinal depression from which rises the alate posterior lobe, which is limited exteriorly by a narrow fold of enamel separating the crown from the basal portion of the tooth; mandibular median or second teeth characterized by their triangular outline, rather strong inrollment of the outer extremity; straight postero-lateral border, which is similarly enameled to the antero-lateral border of last above described posterior dental plates; antero-lateral border rapidly and irregularly converging from the subacute angle of the broad, slightly arched inner margin; coronal surface forming a broad, low arch, or nearly plane transversely. Type *O. carbonarius*.

carbonarius, Newberry & Worthen, 1866, (Sandulodus carbonarius,) Geo. Sur. Ill., vol. 2, p. 104, Up. Coal Meas.

convexus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 193, Coal Meas.

novomexicanus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 195, Sub-carboniferous.

PALÆASPI, Clappole, 1885, Quar. Jour. Geo. Soc. Lond. [Ety. *palaos*, ancient; *aspis*, shield.] Plates or scutes ornamented. Only single dorsal plates discovered. Type *P. americana*.

americana, Clappole, 1885, Quar. Jour. Geo. Soc. Lond., Up. Silurian or Low. Devonian.

truncata, Clappole, 1885, Quar. Jour. Geo. Soc. Lond., Up. Silurian or Low. Devonian.

PALÆORATIS, Leidy, 1856, Trans. Am. Phil. Soc., vol. 11, p. 87. [Ety. *palaos*, ancient; *batis*, a prickly kind of roach or ray.] Type *P. insignis*.

insignis, Leidy, 1856, Trans. Am. Phil. Soc., vol. 11, p. 87, Keokuk Gr.

PALÆONISCUS, Agassiz, 1833, Recherches sur les Poissons Fossiles, t. 1, p. 4. [Ety. *palaos*, ancient; *oniscus*, a wood-louse.] Small, fusiform, deep between ventral and pectoral fins; tail heterocercal, forked, upper lobe longer and narrower than lower; fins small; jaws large; teeth minute; scales rhomboidal, smooth or striated. Type *P. fultus*.

alberti, see Rhadinichthys alberti.

brainerdi, Thomas, 1853, Bost. Soc. Nat. Hist., vol. 4, Ohio Pal., vol. 1, p. 346, Berea grit.

browni, Jackson, 1851, Rep. on Albert Coal Mine, Coal Meas.

cairnesi, see Rhadinichthys cairnesi.

gracilis, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 347, Coal Meas.

jacksoni, Dawson, 1877, Can. Nat. Quar. Jour. Sci., vol. 8, Carboniferous.

less well-defined, abruptly broken where the coronal belt sharply deep basal rim, and the more or less depression from late posterior lobe, anteriorly by a narrow separating the crown of the tooth; a or second teeth their triangular out-inrollment of the eight postero-lateral similarly enameled in order of last above dental plates; antero-dially and irregularly the subacute angle of arched inner margin forming a broad, plane transversely.

s.
ry & Worthen, 1866, (narius,) Geo. Sur. Jp. Coal Meas.
Worthen, 1883, Geo. 193, Coal Meas.
John & Worthen, vol. 7, p. 195, Sub-
1885, Quar. Jour. Geo. alaios, ancient; aspis, scutes ornamented. al plates discovered.

1885, Quar. Jour. Silurian or Low.

1885, Quar. Jour. Silurian or Low. De-
56, Trans. Am. Phil. [Ety. palaios, an- kly kind of roach or gnus.

56, Trans. Am. Phil. Keokuk Gr.
1833, Recherches sur les, t. 1, p. 4. [Ety. niscus, a wood-louse.] deep between ventral; tail heterocercal, longer and narrower small; jaws large; scales rhomboidal. Type P. fultus.
chthys alberti.

1853, Bost Soc. Nat. Pal., vol. 1, p. 346,

51, Rep. on Albert Meas.
chthys cairnesi.
& Worthen, 1870, Geo. 347, Coal Meas.
1877, Can. Nat. Quar. Carboniferous.

leidyanus, Lea, 1853, Jour. Acad. Nat. Sci., 2d ser., vol. 2, Coal Meas.
modulus, see Rhadinichthys modulus.
peltigerus, Newberry, 1857, (Elonichthys peltigerus,) Proc. Acad. Nat. Sci., vol. 8, p. 98, and Ohio Pal., vol. 1, p. 345, Coal Meas.

FIG. 1155.—Palaeoniscus peltigerus.

scutigerus, Newberry, 1857, Proc. Acad. Nat. Sci. Phil., Coal Meas.
Peltodus, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 362. [Ety. pelle, a half-moon shield; odous, tooth.] Teeth small and low, round oval or elliptical in outline, arched above in both directions, concave or flattened below; crown surface most strongly arched from front to rear, highest near the anterior margin; more or less evenly punctate throughout; under surface bony and rough; margins thin and irregular where the teeth are separated, thickened and even along the lines of contact when closely set. They are less flat, smooth and pavement-like than Psammodus, and less convoluted than Coeliodus. Type P. unguiformis.
plicomphalus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 411, Kaskaskia Gr.
quadratus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 410, St. Louis Gr.
transversus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 412, Coal Meas.
unguiformis, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 363, Coal Meas.
Pelorhina, Cope, 1873, Proc. Acad. Nat. Sci. Phil., p. 343. [Ety. pelos, a robe; Rhine, a kind of dog-fish.] Type P. anthracina.
anthracina, Cope, 1873, Proc. Acad. Nat. Sci. Phil., p. 343, Coal Meas.
arctica, Cope, 1877, Proc. Am. Phil. Soc., p. 55, Permian Gr.
Periplectrodus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 324. [Ety. peri, near by; Plectrodus, a genus.] Base expanded laterally or compressed; symmetrically inrolled from within outward, inferior surface excavated; crown

consisting of transverse, strong, median cusps, flanked by denticles; one on either side, and then regularly increase in size from the outer to the inner extremity or with age; coronal cusps enameled, smooth or vertically striated. Type P. warreni.

compressus, St. John & Worthen, 1875, Geo. Sur. Ill. vol. 6, p. 326, St. Louis Gr.

expansus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 327, Kaskaskia Gr.

warreni, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 325, Burlington Gr.

PERIPRISTIS, Agassiz, 1870, Proc.

Am. Phil. Soc., vol. 11, p. 334. [Ety. peri, around; prista, saw.] Small, crown compressed, acuminate, serrate, curved laterally; coronal cavity; root as in Petalodus, crown and coronal cavity covered with ganoine. Type P. semicircularis.

semicircularis, Newberry & Worthen, 1866, (Ctenoprists semicircularis,) Geo. Sur. Ill., vol. 2, p. 72, Coal Meas.

Petalodus, Owen 1840, Odontography, p. 60. [Ety. petalos, spread out; odous, tooth.] Teeth transversely elongated, compressed, thin, petal-shaped, cutting edge serrated; base of crown with imbricating folds of enamel, descending lower on the posterior than anterior face; root large, oblong, truncated below; lower edge obtuse, tumid. Type P. hastingsi.

FIG. 1157.—Petalodus alleghaniensis.

alleghaniensis, Leidy, 1856, Jour. Acad. Nat. Sci., 2d series, vol. 3, p. 161, and Geo. Sur. Ill., vol. 2, p. 35, Coal Meas.
curtus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 6, p. 355, Keokuk Gr.
destructor, Newberry & Worthen, 1866, syn. for P. alleghaniensis.

hybridus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 394, St. Louis Gr.

linguifer, Newberry & Worthen, 1856, Geo. Sur. Ill., vol. 2, p. 37, Kaskaskia Gr. proximus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 395, Coal Meas.

PETALORHYNCHUS, Agassiz, 1855, in British Pal. Rocks. [Ety. *petalos*, spread out; *rhynchos*, a beak.] Teeth small, crown compressed, thin, concavo-convex, petal-shaped; higher and narrower than *Petalodus*; imbricating folds on posterior face forming a short transverse band, not extending to the lateral angles of the crown; root long, undivided. Type *P. sagittatum*.

distortum, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 406, St. Louis Gr.

pseudosagittatum, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 405, St. Louis Gr.

spatulatum, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 408, St. Louis Gr.

striatum, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 40, Burlington Gr.

PETRODUS, McCoy, 1848, Ann. and Mag. Nat. Hist., 2d series, vol. 2, p. 132. [Ety. *petros*, a rock; *odous*, a tooth.] Conical; base round or subtrigonal; apex rudely pointed; sides radiatingly ridged; osseous base wider than the crown. Type *P. patelliformis*.

acutus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 72, Coal Meas.

occidentalis, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 70, Coal Meas.

pustulosus, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 389, Burlington Gr.

PHANEROPLEURON, Huxley, 1871, 10th Decade Geo. Sur. of Gt. Britain. [Ety. *phaneros*, open; *pleuron*, side.]

curtum, Whiteaves, 1881, Am. Jour. Sci. and Arts, 3d ser., vol. 21, p. 495, Upper Devonian.

PHÆBODUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 251. [Ety. mythological name; *odous*, a tooth.] Teeth small; base irregularly elliptical, strongly produced in front and faintly excavated at the median line; the antero-inferior angles approximate, and laterally curve to the rounded extremities; broadly though irregularly rounded behind; the angles in front are occupied by a strong, lateral, pad-like prominence, which is more or less distinctly bilobed and beveled to the deeply excavated inferior surface; posterior margin slightly buried; postero-superior surface moderately convex, and surmounted by a laterally elongated, well-defined prominence, which is situated nearly midway between the base of the crown and the posterior border, to which the surface abruptly slopes, and extending

laterally nearly half the diameter of the base; the coronal region consists of three strong cusps, of which the exterior pair are largest, strongly diverging and moderately recurved or nearly vertical, antero-posteriorly compressed or suboval in section, apparently without distinct cutting edges; median cone similar in shape, erect, more or less produced in front and continued to the shallow median depression in the border; a rudimentary denticle between the median and lateral cusps.

FIG. 1158.—*Phæbodus sophie*.

Type *P. sophie*.

sophie, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 251, Devonian.

PHYSONEMUS, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 176. [Ety. *physa*, bladder; *nema*, thread.] Dorsal spine strong, laterally compressed, deeply imbedded, curved; apex directed toward the front; lateral faces bearing costæ and tubercles; pulp cavity large; base notched. Type *P. subteres*.

altonensis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 454, St. Louis Gr.

carinatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 452, Waverly or Kinderhook Gr.

chesterensis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 455, Kaskaskia Gr.

depressus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 452, Waverly or Kinderhook Gr.

falcatus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 252, St. Louis Gr.

gigas, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 373, Burlington Gr.

parvulus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 453, Keokuk Gr.

FIG. 1159.—*Platyodus lineatus*. Crown surface.

proclivis, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 451, Waverly or Kinderhook Gr.

PLATYODUS, Newberry, 1875, Ohio Pal., vol. 2, p. 58. [Ety. *platys*, broad; *odous*, tooth.] Teeth elliptical in outline; crown arched in both directions; sur-

the diameter of the region consists of of which the ex-
st, strongly diver-
recurved or nearly
teriorly compressed
n, apparently with-
edges; median cone
erect, more or less
and
allow
n the
ntary
e me-
usps.

FIG. 1158.—Phreobodus sophie.

Worthen, 1875, Geo.
51, Devonian.

1843, Recherches sur
s, t. 3, p. 176. [Ety.
a, thread.] Dorsal
erally compressed,
rved; apex directed
lateral faces bearing
pulp cavity large;
e P. subteres.

Worthen, 1875,
6, p. 454, St.

Worthen, 1875, Geo.
52, Waverly or Kin-

Worthen, 1875,
6, p. 455, Kaskas-

Worthen, 1875, Geo.
52, Waverly or Kin-

Worthen, 1883, Geo.
52, St. Louis Gr.

Worthen, 1870, Geo.
p. 373, Burling-

Worthen, 1875, Geo.
53, Keokuk Gr.

Crown surface.

Worthen, 1875, Geo.
p. 451, Waverly or

1875, Ohio Pal., vol.
platys, broad; odous,
typical in outline;
oth directions; sur-

face punctate in undulate lines, but
without folds or ridges. Type P. lin-
eatous.

lineatus, Newberry, 1875, Ohio Pal., vol.
2, p. 58, Waverly Gr.

PLATYSONUS, Agassiz, 1833, Recherches sur
les Poissons Fossiles, t. 1, p. 6. [Ety.
platys, broad; soma, body.] Rhom-
boidal, compressed; dorsal and anal
fins nearly equal, opposite; pectorals
small; ventrals small; teeth clavate;
crown dilated, flattened; base slender,
constricted at the base of the ganoine;
scales large, oblong, articular internal
ridge at anterior edge; beveled spine
at the upper corner received in a
notch of the adjoining scale. Type P.
striatus.

circularis, Newberry & Worthen, 1870,
Geo. Sur. Ill., vol. 4, p. 347, Coal
Meas.

Pleuracanthus, Agassiz, 1843, Poiss. Foss.,
vol. 3, p. 66. [Ety. pleura, side; akan-
tha, spine.] The genus was founded
upon a spine supposed to belong to the
Order Raiina. It is serrated on one
edge, curved at the base, and furrowed
on the inferior side. The species named
in this genus from America are too
poorly defined to warrant recognition.
Type P. laevissimus.

arcuatus, see Orthacanthus arcuatus.

biserialis, Newberry, 1857, Proc. Acad.
Nat. Sci., vol. 8, p. 100, Coal Meas.

dilatatus, Newberry, 1857, Proc. Acad. Nat.
Sci., vol. 8, p. 100, Coal Meas.

PNIGEOACANTHUS, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 480. [Ety. from
the specific name in Oracanthus pnigeus;
akantha, spine.] Spine short, conical,
laterally compressed; base broadly ex-
panded before and behind, without in-
sertion, rapidly tapering to the obtuse
apex, which is directed posteriorly;
transverse section elliptical, rounded
into the slightly sigmoidally curved an-
terior border and concave posterior
margin; pulp cavity very large, ex-
tending nearly to the tip; lateral walls
very thin, slightly thickened in the
margins; external surface occupied by
irregularly disposed, radiatingly scul-
ptured tubercles, sometimes arranged in
obscure or interrupted longitudinal and
diagonal order. The type is Oracanthus
pnigeus, of Newberry & Worthen,
which St. John named Pnigeacanthus
deltoides.

deltoides, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 480, Keokuk Gr. But
why should this species not be Pnigea-
canthus pnigeus?

trigonalis, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 259, St. Louis Gr.

POIKILODONTUS, Agassiz, 1843, Recherches Poiss.
Foss., vol. 3, p. 174. [Ety. poikilos, var-
iegated; odous, tooth.] Teeth as in
Cochliodus; terminal tooth obliquely
trigonal, convoluted; median tooth nar-

row, convoluted; all teeth wrinkled at
right angles to the articular edges; sur-
face porous. Type P. jonesi.

carbonarius, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 139, Coal
Meas.

cestriensis, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 135, Kaskas-
kia Gr.

ornatus, Newberry & Worthen, 1866, Geo.
Sur. Ill., vol. 2, p. 95, syn. for Chitonon-
odus rugosus.

rugosus, see Chitonodus rugosus.

springeri, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 138, Subcarbon-
iferous.

stludovici, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 132, St.
Louis Gr.

varsoviensis, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 131, War-
saw Gr.

wortheni, St. John, 1883, Geo. Sur. Ill.,
vol. 7, p. 136, Kaskaskia Gr.

POLYRHIZODUS, McCoy, 1848, Ann. and Mag.
Nat. Hist., 2d series, vol. 2, p. 125.

[Ety. polys, many; rhiza, root; odous,
tooth.] Crown like Petalodus, but
more elongated, transversely lower and
thicker; root divided into numerous
short, robust radicles. Type P.
magnus.

amplus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 387, St. Louis Gr.

carbonarius, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 389, Coal
Meas.

dentatus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 50, Kaskas-
kia Gr.

littoni, Newberry & Worthen, 1870, Geo.
Sur. Ill., vol.
4, p. 357, St.
Louis Gr.

modestus, New-
berry, 1875,
Ohio Pal., vol.
2, p. 50, Cleve-
land shale.

FIG. 1160.—Polyrhizodus littoni. Concave face.

nanus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p.
386, Keokuk Gr.

piasensis, St. John &
Worthen, 1875,
Geo. Sur. Ill., vol.
6, p. 386, Warsaw
Gr.

ponticulus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 51, Kaskas-
kia Gr.

porosus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 49, Burling-
ton Gr.

truncatus, Newberry & Worthen, 1870,
Geo. Sur. Ill., vol. 4, p. 357, Burling-
ton Gr.

williamsi, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 384, Keo-
kuk Gr.

PRISTICLADODUS, McCoy, 1855, British Pal. Rocks, p. 642. [Ety. from the two genera *Pristis* and *Cladodus*.] Teeth resemble

FIG. 1162.—*Pristicladodus springeri*.

Cladodus; median and lateral cusps strong; lateral edges sharp and more or less undulated. Type *P. dentatus*.

springeri, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 255, Waverly or Kinderhook Gr.

PRISTODUS, Agassiz. [Ety. *pristis*, a saw; *odus*, a tooth.] This genus has been only doubtfully identified in America.

(?) *acuminatus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 402, Waverly or Kinderhook Gr.

PSAMMODUS, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 112. [Ety. *psammos*, sand; *odus*, tooth.] Teeth quadrilateral or trapezoidal in general outline, variable, usually thick and massive; the coronal region presents a more or less plane surface, according to the position the form occupied upon the jaws, always arched, generally moderately in the longitudinal direction or from behind forward, transversely concave (maxillary teeth), or more or less convex (mandibular teeth), sometimes raised into a low ridge along the exterior lateral border, also along the articular inner border, or showing a more or less wide convexity in the latter region, and sometimes presenting a more or less well-defined transverse prominence in mature maxillary form; the marginal limits of the crown are well defined, rounded along the exterior of lateral border, and usually in-beveled, and almost always making an angulation at the articular inner border and along the anterior and posterior margins, the enamel extending well down, and more or less distinctly defined from the coarse, vermicularly pitted base which constitutes the greater part of the height of the tooth; in front and behind, the basal wall is nearly exactly vertical to the plane of the coronal surface, and moderately channeled or concave; the inner articular face is also vertical and slightly excavated, presenting generally at one or the other extremity an obliquely truncated articular facet for co-adaptation with the contiguous tooth of the opposite series, the extent and obliquity of the truncation varying greatly according to the species; the exterior lateral border, in typical forms, shows an expansion of the basal portion beyond the coronal limits, increasing in breadth and terminating in a more or

less produced spur at the postero-outer angle of the tooth; the coronal surface exhibits under an ordinary lens a distinct, vertical, prismatic structure, each of the vertical columns inclosing a medullary tube, the appearance of which at the surface produces the exceedingly minute punctation usually observed in these teeth; the exceedingly elegant vermiculose rugosity exhibited in the less worn surfaces of certain species is produced by the wrinkling of the enamel or external layer, and which apparently has no other relation to the medullary tubes than to rudely define them in irregular and transverse or longitudinal rows,

FIG. 1163.—*Psammodus crassidens*. a, Median dental plates; b, transverse profile; c, longitudinal profile.

the punctæ rarely confluent, and the rugose appearance becoming obsolete or more or less obscured over the more exposed parts of the triturating surface; the impression also prevails that the tendency to rugosity of the coronal surface increases with age, since this appearance, so far as observed, seems to be most prevalent and conspicuous in large individuals belonging to the series which have received several accessions, the innermost individuals of which have suffered little from the abrading effects of trituration while in use; but it is not an essential character, as some species evidently always remained quite smooth in their coronal areas; the

at the postero-outer
the coronal surface
ordinary lens a dis-
tinct structure, each
column inclosing a
the appearance of
a produces the ex-
punctuation usually
teeth; the exceed-
culose rugosity ex-
worn surfaces of
produced by the
amel or external
apparently has no
medullary tubes
in them in irregular
longitudinal rows,

crassidens. a, Median
transverse profile; c, longitu-

confluent, and the
becoming obsolete
curved over the more
tritulating surface;
so prevails that the
ity of the coronal
with age, since this
is observed, seems to
and conspicuous in
belonging to the series
and several accessions,
dividuals of which
from the abrading
on while in use; but
al character, as some
ways remained quite
coronal areas; the

inferior surface is plane, in a general
way conforming to that of the crown,
and even possessing distinctive charac-
teristics as applied to species; it shows
in the perfect state a rather dense,
thin layer, perhaps in degree rather
than structurally differing from the more
cellulose middle layer composing the
bulk of the base, and usually marked by
more or less distinct longitudinal
grooves, or smooth and faintly keeled
nearest the inner articular border.
Type *P. porosus*.

angularis, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 107, Kaskas-
kia Gr.

antiquus, Newberry, 1857, Bull. Nat.
Inst., Up. Held. Gr.

bretonensis, Whiteaves, 1881, Can. Nat.,
vol. 10, Carboniferous.

FIG. 1161.—*Psammodus porosus*.

celatus, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 217, St. Louis Gr.

crassidens, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 218, St.
Louis Gr.

glyptus, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 209, Up. Burling-
ton Gr.

FIG. 1165.—Diagram of *Psammodus springeri*.
a, Mandibular series; b, transverse profile; c,
longitudinal profile.

grandis, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 211, Keokuk Gr.

lovianus, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 207, Burlington Gr.

plenus, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 213, St. Louis Gr.

FIG. 1166.—Diagram of *Psammodus springeri*.
a, Maxillary series; b, transverse profile; c,
longitudinal profile.

porosus, Agassiz, 1843, Recherch. Poiss.
Foss., t. 3, p. 112, Kaskaskia Gr.

FIG. 1167.—Hypothetical diagram, showing me-
dian dental plates of *Psammodus springeri*.

reticulatus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 109, Kaskas-
kia Gr.

FIG. 1168.—Hypothetical diagram of *Psammodus*
turgidus.

rhomboides, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 110, syn. for
Sandalodus laevissimus.

- semicylindricus*, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 109, syn. for *Sandalodus laevisimus*.
- springeri*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 202, Upper Burlington Gr.
- tumidus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 205, Up. Burlington Gr.
- turgidus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 206, Keokuk Gr.
- Psephenodus*, Agassiz, 1862, Quar. Jour. Geo. Soc., vol. 18, p. 102. [Ety. *psephos*, a pebble; *odus*, a tooth.] Heavy, more or less spirally inrolled triturating or crushing plates invest the median range of the rami of the jaws; they are trapezoidal in outline, with undulated articular surfaces. Type *P. magnus*.
- convolutus*, Newberry & Worthen, 1866, (*Aspidodus convolutus*), Geo. Sur. Ill., vol. 2, p. 94, Kaskaskia Gr.
- crenulatus*, Newberry & Worthen, 1866, (*Aspidodus crenulatus*), Geo. Sur. Ill., vol. 2, p. 93, Kaskaskia Gr.
- cunulatus*, see *P. lunulatus*.
- latus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 72, St. Louis Gr.
- lunulatus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 74, (misprinted *cunulatus*), Kaskaskia Gr.
- obliquus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 66, Waverly or Kinderhook Gr.
- placenta*, Newberry & Worthen, 1866, (*Helodus placenta*), Geo. Sur. Ill., vol. 2, p. 80, Waverly or Kinderhook Gr.
- reticulatus*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 417, Waverly or Kinderhook Gr.
- symmetricus*, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 71, Waverly or Kinderhook Gr.
- PTERICHTHYS**, Agassiz, 1835, Recherches sur les Poissons Fossiles, t. 1, p. 302. [Ety. *pteron*, fin; *ichthys*, fish.] The outline of this genus reminded Hugh Miller of the figure of a man rudely drawn in black on a gray ground, the head cut off at the shoulders, the arms spread at full as in the attitude of swimming, the body rather long than otherwise, and narrowing from the chest downward, one of the legs cut away at the hip-joint, and the other as if to preserve the balance, placed directly under the center of the figure, which it seems to support. The under part of the body was flat, the upper rose toward the center into a roof-like ridge, and both under and upper were covered with a strong armor of plates; the plates on the under side are divided by a longitudinal suture and a transverse suture, and they would cut at right angles were it not for a lozenge-shaped plate in the center; there are therefore five plates on the under side, all of which are

thickly tuberculated; the upper side is covered with a large, long, hexagonal plate in the central part, that is surrounded by a row of unequal and variously formed plates, all of which are strongly tuberculated; the cephalic shield is rounded in front, and truncated behind where it joins the body carapace, having a transverse median opening; nuchal region occupied by a plate somewhat like the lateral view of a

FIG. 1169.—*Pterichthys milleri*.

- coronet or crown; one post-median plate, another in front, one lateral occipital on each side, two lateral and one postero-lateral on each side, and an angular plate on each postero-lateral side articulating with the limb; the oblong carapace is covered by the large, hexagonal antero-median plate, and a smaller posterior median dorsal, and two dorso-lateral plates on each side; tail thick, conical, covered with rhomboidal scales; surface covered with granules. Type *P. milleri*.
- canadensis*, see *Bothriolepis canadensis*.
- norwoodensis*, Owen, syn. for *Macropetalichthys rapheidolabis*.
- rugosus*, Clapp, 1883, Proc. Am. Phil. Soc., p. 664, Upper Chemung Gr.
- PRYONODUS**, Pander Über die Ctenodipteren des Devonischen Systems, p. 48. [Ety. *ptyktos*, folded; *odus*, tooth.] Elongated; base expanded, subconical; crown flattened or furrowed; enameled; tubes in transverse furrows, with low intervening ridges. Type *P. obliquus*.
- calceolus*, Newberry & Worthen, 1866, (*Rinodus calceolus*), Geo. Sur. Ill., vol. 2, p. 106, Ham. Gr.
- Pryonodus*, Cope, 1877, Proc. Am. Phil. Soc., p. 192. [Ety. *ptyon*, a fan; *odus*, a tooth.]
- paucicristatus*, Cope, 1877, Proc. Am. Phil. Soc., p. 54, Permian.
- vinslovi*, Cope, 1877, Proc. Acad. Nat. Sci. Phil., p. 410, Permian.

; the upper side is
e, long, hexagonal
l part, that is sur-
unequal and vari-
all of which are
ed; the cephalic
ront, and truncated
ns the body charac-
erse median open-
occupied by a plate
lateral view of a

Pygopterus, Agassiz, 1833, Poiss. Foss., t. 1,
p. 10. [Ety. *pyge*, rump; *pteron*, fin.]
Body large, elongate, ovate; fins large,

FIG. 1170.—*Pyctodus calceolus*. Side view.

with fulcral scales; anal fin long, nearly
opposite dorsal; ventrals small; pecto-
rals small, falcate; caudal large, notched;
upper jaw longer than the lower; endo-
skeleton strong; scales small, rhom-
boidal. Type *P. humboldti*. Not defi-
nitely known from America.

scutellatus, Newberry, 1857, Proc. Acad.
Nat. Sci., vol. 8, p. 98, Coal Meas. Too
poorly defined to warrant recognition.

FIG. 1171.—*Pygopterus mandibularis*. Outside and
under surface of scale magnified.

RHADINICHTHYS, Traquair, 1877, Quar. Jour.
Geo. Soc. Lond., vol. 33, p. 548. [Ety.
rhadinos, slender; *ichthys*, fish.] Body
slender; jaws with a row of incurved
lanteries, outside of which there are
smaller teeth; principal rays of pectoral
fin as in *Pygopterus*; dorsal far back,
nearly opposite the anal. Type *R.*
ornatissimus.

FIG. 1172.—*Rhadinichthys alberti*.

alberti, Jackson, 1851, (Palaeoniscus *alberti*.) Rep. on the Albert Coal-mine,
New Brunswick, Coal Meas.

cairnsi, Jackson, 1851, (Palaeoniscus
cairnsi.) Rep. on Albert Coal-mine,
New Brunswick, Coal Meas.

modulus, Dawson, 1877, (Palaeoniscus
modulus.) Can. Nat. and Quar. Jour.
Sci., vol. 8, Carboniferous.

Rhizonus, Owen, 1840, Odontography.
[Ety. *rhiza*, a root; *odus*, tooth.] Jaws
massive, bearing large, compressed,
double-edged teeth, with sulcated bases

in each dental bone, and numerous
smaller ones; scales large, rotundato-
quadrate, thin, inner surface concentric-

ally lined; outer
surface tubercu-
late. Type *R. hibi-*
berti.

angustus, Newberry,
1857, Proc. Acad.
Nat. Sci. Phil., vol.
8, p. 99, Coal Meas.
Poorly defined.

hardingi, Dawson,
1868, Acad. Geol.,
p. 254, Subcarbon-
iferous.

incurvus, Newberry, 1857, Proc. Acad.
Nat. Sci., vol. 8, p. 99, Coal Meas.
Poorly defined.

FIG. 1173.—*Rhizodus occidentalis*. Scale.

lancifer, Newberry, 1857, Proc. Acad. Nat.
Sci. Phil., vol. 8, p. 99, and Ohio Pal.,
vol. 1, p. 342, Coal Meas.

occidentalis, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 19, Coal Meas.
quadratus, Newberry, 1873, Ohio Pal.,
vol. 1, p. 343, Coal Meas.

reticulatus, Newberry &
Worthen, 1870, Geo. Sur.
Ill., vol. 4, p. 349, Coal
Meas.

Rhynchodus, Newberry, 1873,
Ohio Pal., vol. 1, p. 307.

[Ety. *rhynchos*, beak; *odus*,
tooth.] Teeth somewhat
half-circular, compressed,
exterior margins curved;

one cornua produced, the
other obtuse; straight side the tritrat-
ing or cutting edge. Type *R. secans*.

FIG. 1174.—*Rhynchodus frangens*.

crassus, Newberry, 1873, Ohio Pal., vol.
1, p. 312, Up. Held. Gr.

thys milleri.

; one post-median
front, one lateral
side, two lateral
lateral on each side,
ate on each postero-
ulating with the
g carapace is cov-
a hexagonal antero-
a smaller posterior
l two dorso-lateral
; tail thick, conical,
oidal scales; surface
ules. Type *P. mil-*

olepis canadensis.
syn. for *Macropetal-*
pis.

83, Proc. Am. Phil.
Chemung Gr.

er die Ctenodipteri-

ens Systems, p. 48.

ed; *odus*, tooth.]

panded, subconical;

narrowed; enameled;

ree furrows, with

ridges. Type *P.*

& Worthen, 1866,

Geo. Sur. Ill., vol.

7, Proc. Am. Phil.

pyton, a fan; *odus*,

, 1877, Proc. Am.

ermian.

, Proc. Acad. Nat.

ermian.

- excavatus, Newberry, 1877, Geo. of Wis., vol. 2, p. 397, Ham. Gr.
 frangens, Newberry, 1873, Ohio Pal., vol. 1, p. 311, Up. Held. Gr.
 secans, Newberry, 1873, Ohio Pal., vol. 1, p. 310, Up. Held. Gr.
Rinodus, Newberry & Worthen, 1866, syn. for *Ptyctodus*.
calceolus, see *Ptyctodus calceolus*.
SANDALODUS, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 102. [Ety. *sandalon*, a sandal; *odous*, tooth.] Teeth thick, strong, subtriangular or club-shaped, with one or two pointed extremities; twisted and arched; base concave, surface punctate. Type *S. parvulus*.
angustus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 103, Keokuk Gr.
carbonarius, see *Orthopleurodus carbonarius*.
complanatus, Newberry & Worthen, 1866, (Deltodus *complanatus*), Geo. Sur. Ill., vol. 2, p. 98, Upper Burlington Gr.
crassus, Newberry & Worthen, 1870, Geo. Sur. Ill., vol. 4, p. 369, syn. for *S. spatulatus*.
grandis, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 105, syn. for *S. levissimus*.
levissimus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 104, Keokuk Gr.
minor, Newberry & Worthen, 1866, (Trigonodus *minor*), Geo. Sur. Ill., vol. 2, p. 112, Keokuk Gr.
parvulus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 102, St. Louis Gr. In part *Stenopterodus parvulus*.
spatulatus, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 103, St. Louis Gr.

FIG. 1175.—*Sandalodus spatulatus*.

- Sauripteris*, Hall, 1843, Geo. Rep. 4th Dist. N. Y. [Ety. *sauros*, lizard; *pteron*, wing.]
taylori, see *Holoptychius taylori*.
Sicarius extinctus, Leidy, 1855, Proc. Acad. Nat. Sci., vol. 7. Not satisfactorily defined.
STEMMATODUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 328. [Ety. *stemma*, a wreath; *odous*, tooth.] Teeth variable, anomalous, some triangular with three or more rows of denticles, others simple with narrower base and a single row of coronal cusps. Type *S. chiriformis*.
bicristatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 331, Burlington Gr.
bifurcatus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 330, Burlington Gr.

- chiriformis*, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 330, Burlington Gr.
compactus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 334, Kaskaskia Gr.
keokuk, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 334, Keokuk Gr.
simplex, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 332, Burlington Gr.
symmetricus, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 333, Burlington Gr.
STENACANTHUS, Leidy, 1856, Jour. Acad. Nat. Sci., 2d ser., vol. 3, p. 102. [Ety. *stenos*, narrow; *akantha*, a spine.] Narrow denticulated spine. Type *S. nitidus*.
nitidus, Leidy, 1856, Jour. Acad. Nat. Sci., 2d ser., vol. 3, p. 102, Carboniferous.
STENOPTERODUS, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 100. [Ety. *stenos*, narrow; *pteron*, wing; *odous*, tooth.] Teeth distinguished by their long elliptical outline, strongly arched and spiral inrollment of the outer extremity. Crown with a lobe in the direction of inrollment. Type *S. planus*.
elongatus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 106, Warsaw Gr.
parvulus, Newberry & Worthen, 1866, (*Sandalodus parvulus*), Geo. Sur. Ill., vol. 2, p. 102, St. Louis Gr.
planus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 102, Upper Burlington Gr.
Strigillina, Cope, syn. for *Janassa*.
gurleiana, see *Janassa gurleiana*.
linguiformis, see *Janassa linguiformis*.
TÆNIODUS, De Koninck, MSS., and St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 75. [Ety. *tænia*, ribbon; *odous*, tooth.] Trapezoidal, arched from within outward, inrolled obliquely outward and forward; distinguished from *Pseudophodus* by the differentiation of the coronal contour of the maxillary median forms. Type *T. contortus*.
fasciatus, Newberry & Worthen, 1870, (*Deltodus fasciatus*), Geo. Sur. Ill., vol. 4, p. 361, Keokuk Gr.
obliquus, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 78, Kaskaskia Gr.
regularis, St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 77, Warsaw Gr.
TANAODUS, St. John & Worthen, 1875, Geo. Sur. Ill., vol. 6, p. 367. [Ety. *tanaos*, long; *odous*, a tooth.] In the laterally extended linear outline of the crown it resembles *Chomatodus*; in the disproportionate depth of the coronal surfaces and marginal position of the root it resembles *Antliodus* from which it is distinguished by the linear outline of the crown and the inferior surface of the root. Type *T. gracillimus*.
angularis, Newberry & Worthen, 1866, (*Chomatodus angularis*), Geo. Sur. Ill., vol. 2, p. 55, Coal Meas.

& Worthen, 1875,
p. 330, Burling-

& Worthen, 1875,
p. 334, Kaskas-

Worthen, 1875, Geo.
34, Keokuk Gr.
Worthen, 1875, Geo.
32, Burlington Gr.
& Worthen, 1875,
p. 333, Burling-

1856, Jour. Acad.
ol. 3, p. 162. [Ety.
tha, a spine.] Nar-
pine. Type S. ni-

our. Acad. Nat. Sci.,
2, Carboniferous.
in & Worthen, 1883,
7, p. 100. [Ety.
eron, wing; *odous*,
distinguished by their
ine, strongly arched
ent of the outer ex-
t with a lobe in the di-
nt. Type S. planus.
Worthen, 1883, Geo.
106, Warsaw Gr.
& Worthen, 1886,
ulus.) Geo. Sur. Ill.,
ouis Gr.
Worthen, 1883, Geo.
102, Upper Burling-

for Janassa.
a gurleiana.
assa linguiformis.
MSS., and St. John
(Geo. Sur. Ill., vol. 7,
enia, ribbon; *odous*,
l, arched from within
obliquely outward
distinguished from Pse-
differentiation of the
f the maxillary me-
T. contortus.

& Worthen, 1870,
(s.) Geo. Sur. Ill., vol.
Gr.

Worthen, 1883, Geo.
78, Kaskaskia Gr.
& Worthen, 1883, Geo.
77, Warsaw Gr.

Worthen, 1875, Geo.
p. 367. [Ety. *tanos*,
th.] In the laterally
outline of the crown
matodus; in the dis-
th of the coronal sur-
al position of the root
tliodus from which it
by the linear outline
d the inferior surface
e T. gracillimus.

ry & Worthen, 1866,
ularis.) Geo. Sur. Ill.,
Meas.

bellicinctus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 376, Kaskas-
kia Gr.

depressus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 378, Kaskas-
kia Gr.

FIG. 1176.—*Tanaisiodus gra-*
cillimus.

gracillimus,
Newberry &
Worthen,
1866, (Choma-
todus *gracilli-*
mus.) Geo.
Sur. Ill., vol.

2, p. 51, Burlington Gr.

grossipicatus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 375, Kaskas-
kia Gr.

multiplicatus, Newberry & Worthen,
1866, (Chomatodus *multiplicatus*.) Geo.
Sur. Ill., vol. 2, p. 57, Burlington Gr.

obscurus, Leidy, 1856, (Chomatodus ob-
scurus.) Trans. Am. Phil. Soc., vol. 11,
p. 87, Keokuk Gr.

polymorphus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 380, Kaskas-
kia Gr.

prænuntius, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 371, St. Louis Gr.

pumilus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 369, St. Louis Gr.

sculptus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 373, St. Louis Gr.

sublunatus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 368, St.
Louis Gr.

THORACODUS, Cope, 1883, Proc. Acad. Nat.
Sci., p. 108. [Ety. *thoracos*, protected;
odous, tooth.] Jaws plate-like, divided
on middle line, each half with trans-
verse grooves and ridges, and a smooth
border all round. Type T. eurydinus.

eurydinus, Cope, 1883, Proc. Acad. Nat.
Sci. Phil., p. 108, Permian Gr.

THRINACODUS, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 289. [Ety.
thrinakos, three-pronged; *odous*, tooth.]
Teeth small; base produced posteriorly
in a long sometimes twisted vertically
flattened, or laterally compressed, cla-
vate plate, longer than wide, anterior
face narrow, and abruptly beveled from
the basal line of the crown; posterior
extremity more or less obtusely rounded;
inferior surface narrow, plain or faintly
excavated; superior surface gently
convex, concave antero-posteriorly, or
corresponding to the curvature of the
inferior surface; from the antero-super-
ior extremity of the base spring three
more or less relatively stout, nearly
equal, trenchant, acutely pointed, re-
curved cusps the exterior pair diver-
gent, the central one more or less verti-
cal, slightly sigmoidally curved, trans-
verse section sublenticular, compressed
in front, rounded behind, with simple
cutting edges, and more or less strongly
costate in either face. Allied to Diplo-
dus. Type T. nanus.

duplicatus, Newberry & Worthen, 1866,
(Diplodus *duplicatus*.) Geo. Sur. Ill.,
vol. 2, p. 61, Keokuk Gr.

incurvus, Newberry & Worthen, 1866,
(Diplodus *incurvus*.) Geo. Sur. Ill.,
vol. 2, p. 62, Keokuk Gr.

nanus, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 289, Waverly or Kin-
derhook Gr.

TOMODUS, Agassiz, 1859, MSS. and St. John
& Worthen, 1883, Geo. Sur. Ill., vol. 7,
p. 171. [Ety. *tomos*, sharp; *odous*,
tooth.] Distinguished from *Xystrodus*
by the great convexity of the coronal
ridge, abrupt articular border, and ab-
sence of transverse punctae on the trit-
urating surface. Type T. convexus.

limitaris, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 173, Upper Burling-
ton Gr.

Trigonodus, Newberry & Worthen, 1866,
Geo. Sur. Ill., vol. 2, p. 111. [Ety. *trig-*
onos, three-cornered; *odous*, tooth.] Syn.
for *sandalodus*.

major, Newberry & Worthen, 1866, Geo.
Sur. Ill., vol. 2, p. 112, syn. for *Sanda-*
lodus complanatus.

minor, see *Sandalodus minor*.

VATICINODUS, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 80. [Ety. *vati-*
cinus, prophetic; *odous*, tooth.] Pos-
terior teeth distinguished from *Deltop-*
tichius by the absence of the secondary
lobe, the anterior part of the tooth for-
ward of the posterior prominence being
plain, as in *Stenopterodus*. Type V.
vetustus.

carbonarius, St. John & Worthen,
1883, Geo. Sur. Ill., vol. 7, p. 88, Coal
Meas.

discrepanus, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 83, Upper Burling-
ton Gr.

lepis, St. John & Worthen, 1883, Geo. Sur.
Ill., vol. 7, p. 88, Coal Meas.

similis, St. John & Worthen, 1883,
Geo. Sur. Ill., vol. 7, p. 86, Kaskas-
kia Gr.

simplex, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 84, St. Louis Gr.

vetustus, St. John & Worthen, 1883, Geo.
Sur. Ill., vol. 7, p. 82, Waverly or Kin-
derhook Gr.

VENUSTODUS, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 344. [Ety. *ve-*
nostus, beautiful; *odous*, tooth.] Teeth
laterally elongated, vertically arched;
crown constricted at the base, defined
by imbricating folds; crest elevated,
uniform, or with median prominence;
denticulations lateral; base forming a
shallow plate. Type V. *robustus*.

argutus, St. John & Worthen, 1875,
Geo. Sur. Ill., vol. 6, p. 352, Kaskas-
kia Gr.

leidy, St. John & Worthen, 1875, Geo.
Sur. Ill., vol. 6, p. 350, St. Louis Gr.
This name is a syn. for V. *venustus*.

robustus, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 345, Burlington Gr.

tenuicristatus, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 348, Keokuk Gr.

variabilis, St. John & Worthen, 1875, Geo.

Sur. Ill., vol. 6, p. 346, Burlington Gr.

venustus, Leidy, 1856, (Chomatodus venustus,) Trans. Am. Phil. Soc. Phil.,

vol. 11, p. 87, St. Louis Gr.

XYSTRACANTHUS, Leidy, 1859, Proc. Acad.

Nat. Sci. Phil., p. 3. [Ety. *xystra*, a

tool for scraping; *akantha*, spine.] Dis-

tinguished from *Physonemus* by the

slender, straight outline, and less pre-

ponderance of the antero-inferior shoulder.

Type *X. arcuatus*.

acinaciformis, St. John & Worthen, 1875,

Geo. Sur. Ill., vol. 6, p. 459, Coal Meas.

anceps, Newberry & Worthen, 1866, (Dre-

panacanthus anceps,) Geo. Sur. Ill., vol.

2, p. 122, Coal Meas.

arcuatus, Leidy, 1859, Proc. Acad. Nat.

Sci. Phil., p. 3, Up. Coal Meas.

mirabilis, St. John & Worthen, 1875,

Geo. Sur. Ill., vol. 6, p. 458, Coal Meas.

XYSTRodus, Agassiz, MSS., 1859, and St.

John 1870, Proc. Am. Phil. Soc., vol.

11, p. 436. [Ety. *xystra*, an instrument

for scraping; *odus*, tooth.] Mandibular

posterior teeth triangular; great

transverse breadth of the inner margin

as compared with the longitudinal di-

ameter; coronal surface plain, de-

pressed, and alated posteriorly; max-

illary posterior teeth cuneiform, and

narrow transverse diameter at the inner

margin. Type *X. striatus*.

bellulus, St. John & Worthen, 1883, Geo.

Sur. Ill., vol. 7, p. 183, Coal Meas.

imitatus, St. John & Worthen, 1883,

Geo. Sur. Ill., vol. 7, p. 180, St.

Louis Gr.

inconditus, St. John & Worthen, 1883,

Geo. Sur. Ill., vol. 7, p. 179, Keo-

kuk Gr.

occidentalis, St. John, 1870, Paleontology

of Eastern Nebraska, p. 244, syn. for

Orthopleurodus carbonarius.

simplex, St. John & Worthen, 1883, Geo.

Sur. Ill., vol. 7, p. 178, Upper Burling-

ton Gr.

verus, St. John & Worthen, 1883, Geo.

Sur. Ill., vol. 7, p. 181, Kaskaskia Gr.

CLASS BATRACHIA.

[Ety. *batrachos*, frog.]

THE Batrachia live a double life—that is, both on land and in water—and are called Amphibia. [*Amphi*, on both sides, around; *bios*, life.] They approach the fishes in their early stages of growth, and resemble the true Reptilia in their more mature development. All possess lungs, but during their young or larval condition they are always furnished with branchiæ, and in some orders these remain throughout the life of the animal. They form a distinct transition from aquatic fishes to exclusively air-breathing reptiles. They are all strictly oviparous, although in some species the eggs are retained in or upon the body of the parent until the young have attained some degree of growth. After leaving the egg, the animals undergo a series of transformations before arriving at their complete or perfect state. In their early stage they are known as tadpoles, little, fish-like animals, with broad heads, sack-like body, and long, compressed tail. The mouth is at the lower part of the front of the head, and is furnished with a pair of horny jaws, with which they feed upon the animalculæ that furnishes the food. They are vertebrated animals, with cold blood and naked skin, and undergo a metamorphosis or change of condition from an aquatic respiration by gills to an atmospheric respiration by lungs, and a consequent alteration in general structure and mode of life.

1859, and St. Phil. Soc., vol. 1, an instrument [with.] Mandibular; great the inner margin longitudinal dis- face plain, de- anteriorly; max- cuneiform, and eter at the inner tus.

then, 1883, Geo. Coal Meas. Worthen, 1883, 7, p. 180, St.

Worthen, 1883, 7, p. 179, Keo-

70, Paleontology p. 244, syn. for narius.

then, 1883, Geo. 8, Upper Burling-

then, 1883, Geo. 1, Kaskaskia Gr.

In the tadpole and the genera which retain their gills through life, the substance between the vertebræ is soft, and contained in cup-like hollows formed by the concave articular surfaces of contiguous bones, precisely as in fishes. The lower orders are fish-like in possessing permanent branchiæ, the limbs are reduced to a rudimentary condition, and the tail is flattened and surrounded by a fin. In the higher orders the limbs are more and more developed and fitted for terrestrial progression, until they are capable of active motion and the animals can take their habitual residence in trees. The spinal column in some is composed of a continuous *chorda dorsalis*, inclosed in a fibrous sheath, but furnished with bony superior and inferior arches for the protection of the spinal cord and principal blood-vessels. In others the vertebræ are articulated by a sort of ball-and-socket joint. The vertebræ are usually furnished with long, transverse processes which appear to take the place of ribs; ribs are generally deficient. In those having a *chorda dorsalis* the skull is formed of a simple cartilaginous capsule, with which the *chorda* is completely continuous, and the only indications of ossification are in the lateral portions of the occipital bone. In the higher forms the skull is completely ossified; it is always of a broad and flattened form, with enormous, large orbits, and possesses one constant character which distinguishes the skull of a Batrachia from that of a Reptile; namely, the occipital bone is always furnished with two lateral condyles that fit into corresponding sockets in the first vertebra of the neck. The bones of the upper jaw and palate form a broad arch, which is always firmly attached to the skull; the maxillary and intermaxillary bones assist in the formation of the edge of the mouth, and are much developed, transversely expanding the general form of the skull without involving any enlargement of the brain cavity, which is very small.

All Batrachia have teeth on the palate; the salamanders have them also in both the upper and lower jaws, the frogs in the upper only, and the toads in neither. The jaw teeth are always slender, sharp-pointed, and closely set. The frog has about forty on each side of the upper jaw; the salamander has about sixty above and below; the palatine teeth are generally arranged transversely parallel to the jaw teeth. The hind legs of the frog are developed for leaping, and it has no useless tail; the body is contracted into a short space, and the few vertebræ are united into a single immovable piece, unprovided with ribs. The water salamanders, or newts, have a long tail, a slender flexible body, and all their organs are fitted for aquatic life. The structure of the bones is more compact and calcareous, and less transparent and flexible, than in fishes. The bones of the skull have their margins in contact, and occasionally united, but never overlapping. The hyoid bone changes largely in those genera undergoing metamorphosis in accordance with the development of the respiratory organ.

The Batrachia are generally distinguished from the Reptilia by the absence of a scaly covering. The skin of aquatic genera is soft and smooth, and constantly moistened by the cutaneous secretions; in land genera, as frogs and toads, the glands of the skin secrete a thick, whitish fluid. The cuticle is shed frequently. A few species are covered with horny scales.

They begin life with the single heart and gills of fishes; but as their metamorphosis goes on, the heart assumes the compound character necessary for the pulmonary respiration of the reptiles. In the development of the nervous system and the organs of the senses, they exhibit a slight advance upon the fishes. In the first

in water—and are They approach the stilia in their more g or larval condi- ders these remain tion from aquatic oviparous, although the parent until the e egg, the animals te or perfect state. animals, with broad s at the lower part y jaws, with which are vertebrated ani- phosis or change of respiration by lungs, e.

stages the circulation through the branchial apparatus is exactly the same as in the fishes; but later pulmonary arteries make their appearance, lungs are developed, and aerial respiration commences.

The class has been divided into five orders, namely: Amphipneusta, Anura, Urodella, Abranchia, and Apoda. None of the Palæozoic fossil families are referred to any of these orders except the Cocytinidæ, and the correctness of that reference is exceedingly doubtful. All other Palæozoic fossils are referred to orders which have become extinct. The change, either by progression into higher classes of the vertebrate kingdom, or by retrogression to an inferior state, is strongly marked. The Urodella, to which the Cocytinidæ are referred, have long, slender bodies, four limbs, which are sometimes very small, and occasionally the toes are furnished with claws, and a long, persistent tail; no external branchiæ, but in some species there is a branchial aperture on each side of the neck, within which are the branchial arches, with their lamina; lungs well developed, skin smooth, or covered with warty prominences, and furnished with numerous glands, which secrete an acrid, viscid fluid. In general form they resemble the lizards, which belong to the Reptilia. The aquatic and land salamanders belong to this order.

The first Batrachia are found in the Coal Measures. They increase rapidly in numbers, and spread out in progressive evolution through the Permian Group, and reach their highest development and largest size in the Triassic, and since that time they have declined or retrograded, and now constitute a very inferior grade of the Vertebrata. The Animal Kingdom has been divided into classes and orders upon a basis which constitutes, as it is supposed, a natural system, and the more recent study of embryology has demonstrated that this natural system corresponds with the phases of embryonic history in all, or nearly all, its parts, and the study of Palæontology has proven beyond all peradventure that there has been a succession of organic types from the earliest geological time to the present, which is stamped upon the embryonic growth of living animals, and coincides with the grades established by the natural system of classification.

It follows that when the Palæozoic orders are distinct from the living, the class has been more comprehensive than the definition given by zoologists. Indeed, all the fossils can not be strictly embraced within the prescribed limits of the Batrachia. Many of them might be included within the Reptilia, because they combine Batrachian and Reptilian characters, and where the latter prevail probably they should be classed with the Reptilia. Some of them, however, rise a step higher in the animal system, and include Batrachian and Mammalian characteristics, and for this reason it is urged by the evolutionists that the Mammalia descended from the Batrachia, without having passed through the Reptilian stage. In other words, what is here included is a comprehensive type of animal existence not limited by the bounds which define the living Batrachia.

The arrangement of the fossils into orders and families must be regarded as provisional, and only approximating the present learning, for the following reasons, in addition to those which will be apparent to the specialist: 1. There are several synonyms of orders and families. 2. No one has published a complete classification from which the author could compile the learning. 3. The author has never had an opportunity to study the fossils of this class and have an opinion of his own to assert or defend.

the same as in the
s are developed,

usta, Anura, Uro-
es are referred to
that reference is
to orders which
her classes of the
strongly marked.
ender bodies, four
re furnished with
ne species there is
are the branchial
, or covered with
secrete an acrid,
belong to the Rep-

increase rapidly in
rmian Group, and
and since that time
ferior grade of the
es and orders upon
d the more recent
n corresponds with
and the study of
s been a succession
, which is stamped
h the grades estab-

the living, the class
gists. Indeed, all
ts of the Batrachia.
hey combine Batra-
robably they should
step higher in the
eristics, and for this
ended from the Ba-
n other words, what
not limited by the

must be regarded as
following reasons, in
. There are several
complete classification
uthor has never had
inion of his own to

ORDER ARCHEGOSAURIA.

FAMILY ARCHÆGOSAURIDÆ.—Brachydectes, Hylerpeton.

ORDER GANOCEPHALA.

FAMILY COLOSTEIDÆ.—Amphibamus, Colosteus, Sauropleura.

ORDER LABYRINTHODONTIA.

FAMILY BAPHETIDÆ.—Baphetes.

FAMILY TREMATOSAURIDÆ.—Cricotus.

ORDER MICROSAURIA.

FAMILY DIPLOCAULIDÆ.—Diplocaulus.

FAMILY EOSAURIDÆ.—Eosaurus.

FAMILY MOLGOPHIDÆ.—Molgophis, Pleuroptyx.

FAMILY PELIONIDÆ.—Hylonomus, Pelion.

FAMILY PHLEGETHONTIDÆ.—Phlegethontia.

FAMILY PTYONIDÆ.—Ceraterpeton, Hyphasma, Oestoccephalus, Ptyonius, Thyrsideum.

FAMILY TUDITANIDÆ.—Dendrerpeton, Leptophractus, Tuditanus.

ORDER PELYCOSAURIA.

FAMILY BOLOSOURIDÆ.—Bolosaurus, Chilonyx, Lysorophus.

FAMILY DIADECTIDÆ.—Diadectes, Helodectes.

FAMILY CLEPSYDROPSIDÆ.—Archæobolus, Clepsydrops, Dimetrodon, Ectocynodon, Embolophorus, Empedias, Metarmosaurus, Pariotichus, Thero-pleura.

FAMILY EDAPHOSAURIDÆ.—Edaphosaurus, Pantylus.

ORDER RHACHITOMA.

FAMILY ERYOPSIDÆ.—Acheloma, Anisodexis, Eryops, Ichthyacanthus, Trimerorachis, Zatrachys.

ORDER URODELLA.

FAMILY COCYTINIDÆ.—Cocytinus.

ORDER AND FAMILY UNCERTAIN.—Chirotherium, Collettosaurus, Nothodon, Ophiacodon, Sauropus, Sphæropezium, Sphenacodon, Thenaropus.

ACHELOMA, Cope, 1882, Pal. Bull. No. 35, and Proc. Am. Phil. Soc., p. 455. [Ety. α, without; *cheloma*, a notch.] Mandible without angular process; teeth subequal, rather larger anteriorly; pterygoid bone ending in a free, recurved edge anterior to the quadrate bone; palatines and pterygoids nar-

row; palatal foramen wide, posterior border of cranium entire; without notch on the external side of the epiotic bone; vertebræ rhachitomous. Type A. cumminsi.

cumminsi, Cope, 1882, Pal. Bull. No. 35, and Proc. Am. Phil. Soc., p. 456, Permian.

AMPHIBAMUS, Cope, 1865, Proc. Acad. Nat. Sci. Phil., p. 134. [Ety. *amphi*, both; *bama*, a step; from its two modes of

inuous; neural spines of caudal vertebræ well developed; (?) centra. Type A. *grandiceps*.

FIG. 1178.—*Amphibamus grandiceps*. Two diam.

progressing, swimming, and walking.] Teeth small, simple, equal on margins of jaws; sclerotic plates on eye; table of vertex produced; no horns; preopial bones distinct; tarsus cartilag-

Jour. Geo. Soc. London, vol. 10, p. 207, and Acad. Geol., p. 359, Coal Meas. *Bolosaurus*, Cope, 1878, Pal. Bull. No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 506. [Ety. *bolos*, a lump; *saurus*, a liz-

grandiceps, Cope, 1865, Proc. Acad. Nat. Sci. Phil., p. 134, and Geo. Sur. Ill., vol. 2, p. 135, Coal Meas.

ANISODEXIS, Cope, 1882, Pal. Bull. No. 35, and Proc. Am. Phil. Soc., p. 459.

[Ety. *anisos*, unequal; *dexis*, a bite.] Teeth on premaxillary; maxillary, and dentary bones of unequal lengths, some very large, others very small; dental inflections straight, nearly reaching the pulp-cavity; cranial bones sculptured; vertebræ rhachitinous. Type A. *imbricarius*.

imbricarius, Cope, 1882, Pal. Bull. No. 35, and Proc. Am. Phil. Soc., p. 459, Permian.

ARCHÆOBELUS, Cope, 1877, Proc. Am. Phil. Soc., vol. 17, p. 192. [Ety. *archaios*, ancient; *belos*, a weapon.] Maxillary bone with a large, hollow tooth, with two opposite shallow grooves at the base; crown hollow; skeleton unknown. Type A. *vellicatus*.

vellicatus, Cope, 1877, Proc. Am. Phil. Soc., vol. 17, p. 192, Permian.

BAPHETES, Owen, 1853, Jour. Geo. Soc. London, vol. 10, p. 207. [Ety. *bapto*, I dip or dive, a diving animal.] Teeth conical, curved; outer series one or two lines in diameter, inner series three lines or more; implanted and ankylosed in shallow sockets; lower third of teeth longitudinally striated; cranial bones corrugated; head broad. Type B. *planiceps*.

minor, Dawson, 1870, Can. Nat. and Geol., Coal Meas.

planiceps, Owen, 1853, Jour. Geo. Soc. London, vol. 10, p. 207, and Acad. Geol., p. 359, Coal Meas.

of caudal vertebrae. Type

ops, Cope, 1865, Acad. Nat. Sci. p. 134, and Geol. Ill., vol. 2, p. 135, Meas.

is, Cope, 1882, Pal. No. 35, and Proc. Phil. Soc., p. 459. anisos, unequal; a bite.] Teeth on axillary; maxillary and dentary bones unequal lengths, very large, others small; dental fangs straight, reaching the cavity; cranial bones sculptured; vertebral rachitiformous. Type A. imbricarius.

arius, Cope, 1882, Bull. No. 35, and Am. Phil. Soc., 1899, Permian. BELUS, Cope, 1877, Am. Phil. Soc., 17, p. 192. [Ety. belos, ancient; belos, weapon.] Maxillary with a large, hollow tooth, with two shallow grooves on the base; crown low; skeleton unknown. Type A. velli.

atus, Cope, 1877, Am. Phil. Soc., 17, p. 192, Permian. OWEN, 1853, Jour. Soc. London, vol. 10, p. 207. [Ety. bapto, to dip or dive, a diving animal.] Teeth conical, outer series of two lines in meter, inner series of three lines or more; planted and anchored in shallow sockets; over third of teeth longitudinally striated; cranial bones corrugated; head broad. Type B. planiceps.

r, Dawson, 1870, Can. Rept. Geol., Coal Measures. PLANICEPS, Owen, 1853, London, vol. 10, p. 207, p. 359, Coal Meas.

8, Pal. Bull. No. 29, Phil. Soc., vol. 17, p. 507, lump; sauros, a lizard.

ard.] Teeth fixed in shallow alveoli, with crowns expanded transversely to the jaw, swollen at the base, apex low and divided vertically; the postero-internal half in the maxillary series is low and horizontal, the antero-external portion forms a curved cusp; in the lower jaw the relative position of the ledge and cusp is reversed. Type B. striatus.

FIG. 1179.—Baphetes planiceps.

rapidens, see Chilonyx rapidens.

striatus, Cope, 1878, Pal. Bull. No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 507, Permian.

FIG. 1180.—Brachydeutes newberryi.

BRACHYDEUTES, Cope, 1868, Proc. Acad. Nat. Sci. Phil., p. 214. [Ety. brachys, short; deutes, a biter.] Rami short, stout; teeth subequal, elongate, cylindric cones with acute tips turned posteriorly, pulp-cavity large; skeleton unknown. Type B. Newberryi.

newberryi, Cope, 1868, Proc. Acad. Nat. Sci. Phil., p. 214, and Ohio Pal., vol. 2, p. 388, Coal Meas.

FIG. 1181.—Ceraterpeton tenuicorne.

CERATERPETON, Huxley. [Ety. keras, horn; erpeton, reptile.] Teeth simple, equal on outside of jaws; angles of intercalary bones produced into horn-like processes; cranial bones sculptured,

vertebrae undivided; carpus and tarsus osseous.

lineopunctatum, Cope, 1875, Ohio Pal., vol. 2, p. 372, Coal Meas.

tenuicorne, Cope, 1875, Ohio Pal., vol. 2, p. 372, Coal Meas.

CHIROTERIUM, Kaup, 1835, in Leonhard und Bronn Neues Jahrbuch für Mineralogie. [Ety. cheir, the hand; therion, beast.] Represented by foot impressions only. Toes robust, the internal shorter and divergent from the others. Sole (or palum) short, wide. Type C. barthi.

reiteri, Moore, 1873, Am. Jour. Sci. and Arts, 3d ser., vol. 5, p. 292, Coal Meas.

CHILONYX, Cope, 1883, Proc. Am. Phil. Soc., vol. 20, p. 631. [Ety. cheilos, lip; onyx, claw.] Long diameter of the crowns of the teeth transverse to the jaw, and each crown contracting to a slightly incurved apex; maxillary teeth short; temporal fossae roofed; superior surface of cranium divided in areas by grooves. Type C. rapidens.

rapidens, Cope, 1878, (Bolosaurus rapidens), Proc. Am. Phil. Soc., vol. 17, p. 506, and vol. 20, p. 631, Permian.

CLEPSYDROPS, Cope, 1876, Proc. Acad. Nat. Sci. Phil., p. 407. [Ety. klepsydra, an hour-glass; ops, appearance.] Intercentra present; neural spines only elongate posteriorly; premaxillary teeth not especially elongate; one or two long maxillary teeth; no grinding teeth. Type C. colletti.

colletti, Cope, 1876, Proc. Acad. Nat. Sci. Phil., p. 407, Permian.

gigas, see Dimetrodon gigas.

leptocephalus, Cope, 1884, Pal. Bull. No. 39, p. 30, Permian.

limbatus, Cope, 1877, Proc. Am. Phil. Soc., p. 196, Permian or Triassic.

macrospandylus, Cope, 1884, Pal. Bull. No. 39, p. 35, Permian.

natalis, Cope, 1878, Pal. Bull., No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 509, Permian.

pedunculatus, Cope, 1877, Proc. Am. Phil. Soc., p. 63, Permian.

vinslovi, Cope, 1877, Proc. Am. Phil. Soc., p. 62, Permian.

COCYRINUS, Cope, 1871, Proc. Am. Phil. Soc., p. 177, and Ohio Pal., vol. 2, p. 360. [Ety. mythological name.] Vertebrae and ribs osseous; teeth on the premaxillary bone; none on the maxillary; axial hyal with basi-hyal on each side united with corresponding ceratohyal at the end of which is an element in position of stylohyal; haemal or basal branchi-hyals three, the anterior two each supporting one pleural branchi-hyal and the third supporting one; haemal branchi-hyal on the inner side of the ceratohyal, approaching the me-

dian line, and with elongate pleural element. Type *C. gyrinoides*.

FIG. 1182.—*Cocyotinus gyrinoides*.

- gyrinoides, Cope, 1874, Trans. Am. Phil. Soc., and Ohio Pal., vol. 2, p. 364, Coal Meas.
- COLLETOSAURUS, Cox, 1874, Geo. Sur. Ind., 5th Ann. Rep., p. 247. [Ety. proper name; *sauros*, a lizard.] Founded upon tracks having five digits, and supposed to be related to Batrachians and Salamanders. Type *C. indianensis*.
- indianensis, Cox, 1873, Geo. Sur. Ind., 5th Ann. Rep., p. 247, Coal Meas.
- COLOSTEUS, Cope, 1869, Trans. Am. Phil. Soc., p. 22. [Ety. *kolos*, imperfect; *osteon*, a bone.] No vertebral centra, spines, or sclerotic bones; short ribs; two pairs of short limbs; three sculptured pectoral bones; abdominal region protected by scales in chevron; ? anterior teeth longer than posterior, basal half incised sulcate, except two behind the dentary. Type *C. foveatus*.
- crassiscutatus*, Cope, syn. for *C. scutellatus*.
- foveatus, Cope, 1869, Trans. Am. Phil. Soc., p. 24, and Ohio Pal., vol. 2, p. 406, Coal Meas.
- marshi*, see *Ptyonius marshi*.
- pauciradiatus, Cope, 1874, Trans. Am. Phil. Soc., p. 10, and Ohio Pal., vol. 2, p. 408, Coal Meas.
- scutellatus, Newberry, 1856, (Pygopterus scutellatus,) Proc. Acad. Nat. Sci. Phil., p. 98, and Ohio Pal., vol. 2, p. 407, Coal Meas.
- CRICOTUS, Cope, 1876, Proc. Acad. Nat. Sci. Phil., p. 405. [Ety. *krikotos*, ringed.] Centra undivided, equal to the disciform intercentra in the caudal region, intercentra a little smaller in the dorsal region; neural spines and zygapophyses; developed foramen; chordæ dorsalis persistent; teeth equal, except probably the palatines; limbs short, a facial lyra. Type *C. heteroclitus*.
- crassidiscus, Cope, 1884, Pal. Bull. No. 39, p. 25, Permian.
- discochordus, Cope, syn. for *C. heteroclitus*.
- gibsoni, Cope, 1877, Pal. Bull. No. 26, and Proc. Am. Phil. Soc., vol. 17, p. 185, Permian.
- heteroclitus, Cope, 1876, Proc. Acad. Nat. Sci. Phil., p. 405, Permian.
- hypantiricus, Cope, 1884, Pal. Bull. No. 39, p. 30, Permian.

DENDRERPETON, Owen, 1853, Quar. Jour. Geo. Soc., vol. 9, p. 58. [Ety. *dendron*, a tree; *erpeton*, a lizard, from the circumstance under which the reptile was found.] Teeth in double series; outer simple, flattened, conic; inner series conical, with inflected folds of cement; teeth on the vomer; skull-bones corrugated; body protected below with ovate or rhomboidal bony scales, imbricated, horny scales above; fore-limbs the larger; tail natatory; vertebrae biconcave; neural arches and bones ossified. Type *D. acadianum*.

adianum, Owen, 1853, Quar. Jour. Geo. Soc., vol. 9, p. 58, and Acadian Geology, p. 362, Coal Meas.

Fig. 1183.—*Dendrerpeton acadianum*. (a) Cross section of tooth magnified.

- obtusum*, see *Tuditatus obtusum*.
- oweni, Dawson, 1883, Quar. Jour. Geo. Soc., vol. 19, p. 469, and Acad. Geol., p. 368, Coal Meas.
- DIAPLECTES, Cope, 1878, Pal. Bull. No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 505. [Ety. *dia*, crosswise; *dektor*, a biter.] Teeth with much compressed crowns, with bracket-shaped edge, longer axis transverse to the jaws, edges of crowns obtuse, no sculpture on the face. Alveoli not separated. External alveolar border more elevated than the internal, inner alveolar border pierced by a fossa behind the inner extremity of each tooth. Type *D. sidæopellicus*.
- latibuccatus*, see *Empedias latibuccatus*.
- molaris*, see *Empedias molaris*.
- phaseolinus*, see *Empedias phaseolinus*.
- sideropellicus, Cope, 1878, Pal. Bull. No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 505, Permian.
- DIMETRODON, Cope, 1878, Pal. Bull. No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 512. [Ety. *dimetros*, two measures; *odus*, tooth.] Dentition with enormously long incisors and two or three long maxillaries; the pubic bone not distinct from ischium; humerus with trochlear condyles and a defined proximal articular surface; neural spines of dorsal and lumbar vertebrae enormously elongate; intercentra present. Type *D. inclivus*.
- cruciger, Cope, 1878, Am. Naturalist, vol. 12, p. 830, Permian.
- gigas, Cope, 1878, Am. Nat., p. 327, (Clepsydrops gigas,) and Proc. Am. Phil. Soc., vol. 17, p. 515, Permian.

1853, Quar. Jour.
1858. [Ety. *dendron*,
ard, from the cir-
ch the reptile was
uble series; outer
nic; inner series
d folds of cement;
skull-bones corru-
l below with ovate
scales, imbricated,
; fore-limbs the
; vertebrae bicon-
and bones ossified.

3, Quar. Jour. Geo.
Acadian Geology,

(a) Cross section of

s obtusus.
Quar. Jour. Geo.
and Acad. Geol., p.

Pal. Bull. No. 29,
il. Soc., vol. 17, p.
osswise; *decklos*, a
much compressed
-shaped edge, longer
the jaws, edges of
sculpture on the
separated. External
are elevated than the
olar border pierced
the inner e "em-
Type D. sid. copel-

dius latibuccatus.
molaris.

dius phaseolinus.
1878, Pal. Bull. No.
Phil. Soc., vol. 17, p.

8, Pal. Bull. No. 29,
il. Soc., vol. 17, p.
ros, two measures;
ntition with enor-
s and two or three
the pubic bone not
ium; humerus with
and a defined proxi-
face; neural spines
nbar vertebrae enor-
intercentra present.

Am. Naturalist, vol.
n.

n. Nat., p. 327, (Clep-
and Proc. Am. Phil.
5, Permian.

incisivus, Cope, 1878, Pal. Bull. No. 29,
and Proc. Am. Phil. Soc., vol. 17, p.
512, Permian.

rectiformis, Cope, 1878, Pal. Bull. No. 29,
and Proc. Am. Phil. Soc., vol. 17, p.
514, Permian.

semiradicatus, Cope, 1881, Bull. U. S. Geo.
Sur. Terr., vol. 6, No. 1, p. 80, Permian.

DIPLOCAULUS, Cope, 1877, Pal. Bull. No. 28,
and Proc. Am. Phil. Soc., vol. 17, p.
187. [Ety. *diploos*, double; *kaulos*, stem.]
Vertebral centra contracted medially,
perforated by the foramen chordæ dor-
salis, co-ossified with the neural arch
and supporting transverse processes;
zygosphe articulation; two rib articu-
lations, one below the other; axis and
atlas united by a long zygosphen which
is not roofed by the zygantum; no
neural spine, atlas insegmented; arch
extended into the foramen magnum;
squamosal region developed into a
horn. Skull sculptured. Type D. sala-
mandroides.

magnicornis, Cope, 1882, Pal. Bull. No.
35, and Proc. Am. Phil. Soc., p. 453,
Permian.

salamandroides, Cope, 1877, Pal. Bull.
No. 26, and Proc. Am. Phil. Soc., vol.
17, p. 187, Permian.

ECROCYNODON, Cope, 1878, Pal. Bull. No.
29, and Proc. Am. Phil. Soc., p. 508.
[Ety. *ektos*, eternal; *kuon*, dog; *odous*,
tooth.] Cranium short, wide, large
post frontal bones, large orbit; bones
sculptured but no lyra; teeth rhizo-
dont, crowns elongated, compressed,
anterior and posterior cutting edges;
one between the orbit and nostril
larger and longer than the others, and
lying outside of the closed dentary
bone; mandibular symphysis not sutu-
ral but ligamentous. Type E. ordi-
natus.

aguti, Cope, 1882, Pal. Bull. No. 35, and
Proc. Am. Phil. Soc., p. 451, Permian.
ordinatus, Cope, 1878, Pal. Bull. No. 29,
and Proc. Am. Phil. Soc., p. 508, Per-
mian.

EDAPHOSAURUS, Cope, 1882, Pal. Bull. No.
35, and Proc. Am. Phil. Soc., p. 448.
[Ety. *edaphos*, pavement; *sauros*, a liz-
ard.] Temporal fossae not overroofed;
cranial bones not sculptured; mandib-
ular and maxillary teeth subequal;
mandibular ramus expanded inward and
supporting numerous teeth; pterygoid
or malar bones supporting a dense body
of teeth corresponding to those in the
lower jaw; teeth subconical. Neural
spines greatly elongate, hollow. Type
E. pogonias.

microdus, Cope, 1884, Pal. Bull. No. 39, p.
37, Permian.

pogonias, Cope, 1882, Pal. Bull. No. 35, and
Proc. Am. Phil. Soc., p. 449, Permian.

EMBLOPHORUS, Cope, 1878, Pal. Bull. No.
29, and Proc. Am. Phil. Soc., vol. 17, p.
518. [Ety. *ballo*, I throw; *em*, into;

phoros, bearing.] Neural arch co-ossif-
ied, zygapophyses and diapophyses
well developed; centra not ochordal;
intercentra narrowed and transversely
extended; ribs two-headed, the capitulum
is received into a fossa of the pos-
terior border of the intercentrum, in
advance of the vertebra which supports
the diapophysis, to which the tubercu-
lum is attached. Type E. fritillus.

fritillus, Cope, 1878, Pal. Bull. No. 29,
and Proc. Am. Phil. Soc., vol. 17, p. 518,
Permian.

EMPEDIAS, Cope, 1883, Proc. Am. Phil. Soc.,
vol. 20, p. 63. Proposed instead of
Empedocles of Cope in 1878, which was
preoccupied. Teeth with elongate
crowns, with flat grinding surface but
bracket-shaped in transverse vertical
section, arranged transversely to the
longaxis of the jaws; no canines; incisors
wearing chisel-shaped; temporal fossa
covered; vertebrae with hyposphen
on the posterior and hypantum on the
anterior face and short quadrate neural
spines. Type E. alatus.

alatus, Cope, 1878, (Empedocles alatus.)
Proc. Am. Phil. Soc., vol. 17, p. 516,
Permian.

fissus, Cope, 1883, (Empedocles fissus.)
Proc. Am. Phil. Soc., p. 634, Permian.

latibuccatus, Cope, 1878, (Diadectes lati-
buccatus.) Proc. Am. Phil. Soc., vol. 17,
p. 505, Permian.

molaris, Cope, 1878, (Diadectes molaris.)
Am. Nat., vol. 12, p. 565, and Pal. Bull.
No. 32, p. 10, Permian.

phaseolinus, Cope, 1880, (Diadectes pha-
seolinus.) Pal. Bull. No. 32, p. 9, Per-
mian.

Empedocles, Cope, 1878, Proc. Am. Phil. Soc.,
vol. 17, p. 18. The name was preoc-
cupied, see Empedias.

alatus, see Empedias alatus.

fissus, see Empedias fissus.

EOMSAURUS, Marsh, 1862, Can. Nat. and Geo.,
vol. 7, and

Acadian Geol.,
p. 382. [Ety.
eos, the dawn;
sauros, a sea-
fish.] Founded
upon vertebrae,
with biconcave
centra and free
neural arch,
and closed not
ochordal fora-
men. Type E.
acadianus.

acadianus, Marsh,
1862, Can. Nat.
FIG. 1184.—Eosaurus acadi-
and Geol., vol. 7, anus. 1/2 Diam. a, Trans-
verse section.
7, and Acad. Geol., p. 382, Coal Meas.

Epicordylus, Cope, 1878, Pal. Bull. No. 29,
and Proc. Am. Phil. Soc., vol. 17, p. 515.
[Ety. *ept*, upon; *kordylos*, a water-lizard.]
Syn. for Eryops.

- erythroliticus*, see *Eryops erythroliticus*.
Eryops, Cope, 1877, Proc. Am. Phil. Soc., vol. 17, p. 188. [Ety. *eryo*, I protect; *ope*, view.] Vertebrae rhaichitinous throughout; teeth of external series equal; some larger ones on the palatine bones; table of cranium produced, bounded by a notch on each side; no horns nor mucous grooves; pelvic elements co-ossified; no foramen. Type *E. megacephalus*.
erytholeticus, Cope, 1878, (*Epicordylus erytholeticus*.) Proc. Am. Phil. Soc., p. 515, Permian.
ferricolus, Cope, 1878, (*Parioxys ferricolus*.) Proc. Am. Phil. Soc., p. 521, Permian.
megacephalus, Cope, 1877, Proc. Am. Phil. Soc., vol. 17, p. 188, Permian or Triassic.
platypus, Cope, 1877, (*Ichthyacanthus platypus*.) Proc. Am. Phil. Soc., vol. 17, p. 574, Coal Meas.
reticulatus, Cope, 1881, Am. Nat., p. 1020, Permian.
EURYTHORAX, Cope, 1875, Ohio Pal., vol. 2, p. 401. [Ety. *euryx*, broad; *thorax*, the breast.] Established on a thoracic shield, having broad, smooth surfaces on the outer borders for the contact of the overlapping margins of the lateral plates. Subround, with a large excavation from the posterior margin on each side; narrowed portion left in the middle behind has a convex outline; no sculpture. Type *E. sublaevis*.
sublaevis, Cope, 1871, Proc. Am. Phil. Soc., p. 177, and Ohio Pal., vol. 2, p. 402, Coal Meas.

FIG. 1185.—*Hylerpeton dawsoni*. Mandible and portion of cranial bone.

- HELODECTES*, Cope, 1880, Pal. Bull. No. 32, p. 11. [Ety. *helos*, a nail; *dektes*, a biter.] Two rows of subround molariform teeth in each jaw. Type *H. paridens*.
isaaci, Cope, 1880, Pal. Bull. No. 32, p. 12, Permian.
paridens, Cope, 1880, Pal. Bull. No. 32, p. 11, Permian.
HYLERPETON, Owen, 1882, Quar. Jour. Geo. Soc., vol. 18, p. 5. [Ety. *hyle*, wood; *erpeton*, reptile.] Teeth simple, bluntly conical, with large pulp-cavity; about 13 on one side of a jaw; two of the anterior ones of the upper jaw twice as large as the others and deeply sunk in

- the jaw. Length of lower jaw $\frac{1}{2}$ inch; bones of skull puncto-striate. Type *H. dawsoni*.
curtidentatum, Dawson, 1876, Am. Jour. Sci. and Arts, vol. 12, Coal Meas.
dawsoni, Owen, 1882, Quar. Jour. Geo. Soc., vol. 18, p. 5, and Acadian Geology, p. 380, Coal Meas.
longidentatum, Dawson, 1876, Am. Jour. Sci. and Arts, vol. 12, Coal Meas.
HYLONOMUS, Dawson, 1860, Quar. Jour. Geo. Soc., vol. 16, p. 268. [Ety. *hyle*, wood; *nomos*, an abode; forest dweller.] Cranial bones thin, smooth; parietal bones arched; about 28 teeth in each maxillary, elongated, conical, set in a single series, in a furrow, protected externally by an alveolar ridge; teeth longer in intermaxillaries and extremities of mandibles than elsewhere; vertebrae ossified, biconcave, with spinous processes; ribs long and curved; pelvis large; ilium long, expanded below, ischium expanded; pubis expanded, triangular where it joins the ischium, round and arched toward the symphysis; femur thick, nearly straight; tibia short, stout; fibula slender; phalanges broad. Dermal covering of ovate bony scales. Type *H. lyelli*.
acidentatus, Dawson, 1860, Quar. Jour. Geo. Soc., vol. 16, p. 268, and Acad. Geol., p. 378, Coal Meas.

FIG. 1186.—*Hylonomus acidentatus*. Maxillary bone magnified; a, natural size.

- lyelli*, Dawson, 1860, Quar. Jour. Geo. Soc., vol. 16, p. 268, and Acad. Geol., p. 370, Coal Meas.
wymani, Dawson, 1860, Quar. Jour. Geo. Soc., vol. 16, p. 268, and Acad. Geol., p. 378, Coal Meas.
HYPHASMA, Cope, 1875, Proc. Acad. Nat. Sci., p. 18, and Ohio Pal., vol. 2, p. 387. [Ety. *hyphasma*, a web.] Vertebrae osseous; posterior dorsals with fan-like neural spines, ventral armature consisting of rhomboidal scuta, forming packed rows arranged in chevrons, directed backward, on top of which are the usual rod-like scales arranged in packed chevrons with the angle directed forward. Type *H. laevis*.

FIG. 1187.—*Hyphasma laevis*.

- laevis*, Cope, 1875, Proc. Acad. Nat. Sci., p. 18, and Ohio Pal., vol. 2, p. 387, Coal Meas.
ICHTHYACANTHUS, Cope, 1877, Pal. Bull. No. 24, and Proc. Am. Phil. Soc., p. 573

lower jaw $\frac{1}{2}$ inch;
to-atriate. Type H.

on, 1876, Am. Jour.
12, Coal Meas.

, Quar. Jour. Geo.
and Acadian Geology,

on, 1876, Am. Jour.
12, Coal Meas.

360, Quar. Jour. Geo.

[Ety. *hyle*, wood;
forest dweller.] Cra-

ooth; parietal bones
teeth in each maxil-

lical, set in a single
protected externally

teeth longer in
d extremities of man-

ere; vertebrae ossi-

spinous processes;
rved; pelvis large;

aded below, ischium
expanded, triangular

ischium, round and
e symphysis; femur

ht; tibia short, stout;
alanges broad. Der-

ovate bony scales.

Fig. 1188.—*Hylonomus*
iscedentatus. Maxil-

lary bone magnified; a,
natural size.

(Geo. Soc., vol. 16, p.
Geol., p. 370, Coal

860, Quar. Jour. Geo.
8, and Acad. Geol., p.

375, Proc. Acad. Nat.
Ohio Pal., vol. 2, p. 387.

a web.] Vertebrae os-

dorsals with fan-like

ventral armature con-

oid scuta, forming

anged in chevrons, di-

on top of which are

ke scales arranged in

with the angle directed

H. laevis.

ama laevis.

Proc. Acad. Nat. Sci.,
Pal., vol. 2, p. 387, Coal

pe, 1877, Pal. Bull. No.
m. Phil. Soc., p. 573.

[Ety. *ichthys*, a fish; *akantha*, a spine.]
Founded on the posterior dorsal and
caudal vertebrae and adjacent parts.
Posterior limbs well-developed, with
tibia, fibula, osseous tarsus and five
digits; ribs elongate, simple, curved;
abdominal armature in bristle-like rods,
in anteriorly directed chevrons; dorsal
vertebrae short, with simple neural
spines; tall large, vertebrae ossified and
furnished with slender chevron bones,
which terminate in a haemal spine;
neural spines slender, directed back-
ward, caudal series somewhat resem-
bling that of a fish; centra amphicoelous.
Type *I. ohioensis*.

ohioensis, Cope, 1877, Pal. Bull. No. 24,
and Proc. Am. Phil. Soc., p. 573, Coal
Meas.

platypus, see *Eryops platypus*.

LEPTOPHRACUS, Cope, 1873, Proc. Acad.
Nat. Sci., p. 340, and Ohio Pal., vol. 2, p.

399. [Ety. *leptos*, delicate; *phrakto*, ar-

mored.] Founded on various portions
of the cranium; jaws bear large teeth,

round in section at the base, but with
compressed, acute apex, and with cut-

ting edge on anterior face; enamel deli-

cately grooved; there is a large elon-

gate tooth in the upper jaw in the
position of a canine; sculpture of the

cranium little marked; lower jaw
marked with inosculating grooves.

Type *L. obsoletus*.

lineolatus, Cope, 1877, Pal. Bull. No. 24,
and Proc. Am. Phil. Soc., p. 576, Coal
Meas.

obsoletus, Cope, 1873, Proc. Acad. Nat.
Sci., p. 341, and Ohio Pal., vol. 2, p.

400, Coal Meas.

LYSOROPHUS, Cope, 1877, Pal. Bull. No. 26,
and Proc. Am. Phil. Soc., vol. 17, p.

187. [Ety. *lysos*, free; *orophos*, roof.]
Founded upon the centra. Vertebrae

amphicoelous, perforated by the foramen
chordae dorsalis; neural arch freely ar-

ticulated to the centrum; floor of neu-
ral canal deeply excavated; no pro-

cesses or costal articulations on the
centrum, which is excavated by longi-

tudinal fossae; centrum not shortened.
Type *L. tricarinatus*.

tricarinatus, Cope, 1877, Pal. Bull. No.
26, and Proc. Am. Phil. Soc., vol. 17, p.

187, Permian.

METARMOSAURUS, Cope, 1878, Pal. Bull., No.
29, and Proc. Am. Phil. Soc., vol. 17, p.

516. [Ety. *meta*, down; *harmos*, a joint;
sauros, lizard.] Founded upon verte-

brae. Centrum shorter than wide,
deeply biconcave; diapophyses project

below the base of the neural arch, are
short, with small tubercular facet; ca-

pitular facet; facet for intercentrum
small, and is excavated at the anterior

extremity of the base of the centrum;
neural canal large; articular faces of

anterior zygopophyses directed down-
ward and outward. Type *M. fossatus*.

fossatus, Cope, 1878, Pal. Bull., No. 29,
and Proc. Am. Phil. Soc., vol. 17, p. 516,
Permian.

FIG. 1188.—*Molgophis brevicostatus*.

MOLGOPHIS, Cope, 1868, Proc. Acad. Nat.
Sci., p. 220. [Ety. *molges*, a salamander;

ophis, serpent.] Body long, serpentine,
without dermal armature; vertebrae

long, broad, with prominent zygophy-

ses and moderate neural lines; ribs
large, curved, with tubercle and head

on the dilated extremity. Type *M. ma-*
crurus.

brevicostatus, Cope, 1875, Ohio Pal., vol.
2, p. 369, Coal Meas.

macrurus, Cope, 1868, Proc. Acad. Nat.
Sci., p. 220, and Ohio Pal., vol. 2, p. 368,

Coal Meas.

wheatleyi, Cope, 1875, Ohio Pal., vol. 2,
p. 369, Coal Meas.

NOTHODON, Marsh, 1878, Am. Jour. Sci. and
Arts, 3d ser., vol. 15, p. 410. [Ety. *no-*

thos, apurious; *odous*, tooth.] Type *N.*
lentus.

lentus, Marsh, 1878, Am. Jour. Sci. and
Arts, 3d ser., vol. 15, p. 410, Permian.

FIG. 1189.—*Oestocephalus rectidens*.

OESTOCEPHALUS, Cope, 1868, Proc. Acad. Nat.
Sci. Phil., p. 217, and Ohio Pal., vol. 2,

p. 380. [Ety. *oistos*, an arrow; *kephale*,
the head.] Form slender, snake-like;

caudal vertebrae with dilated and sculp-

tured neural and haemal spines; cran-

ium lanceolate; teeth numerous, sub-

equal; no pectoral shields; abdomen
protected by bristle-like rods, which

converge forward; no scales; a pair of
weak posterior limbs; branchial

bones present. Type *O. remex*.

amphimimus, Cope, 1868, Proc. Acad. Nat.
Sci., syn. for *O. remex*.

- pectinatus*, Cope, see *Ptyonius pectinatus*.
rectidens, Cope, 1874, Trans. Am. Phil. Soc., and Ohio Pal., vol. 2, p. 380, Coal Meas.
remex, Cope, 1808, Proc. Acad. Nat. Sci. p. 217, (*Sauropleuria remex*), Ohio Pal., vol. 2, p. 381, Coal Meas.
serrula, see *Ptyonius serrula*.
vinchellianus, see *Ptyonius vinchellianus*.
OPHIACODON, Marsh, 1878, Am. Jour. Sci. and Arts, 3d ser., vol. 15, p. 411. [Ety. *ophiakos*, belonging to serpents; *odous*, tooth.] Type *O. grandis*.
grandis, Marsh, 1878, Am. Jour. Sci. and Arts, 3d ser., vol. 15, p. 411, Permian.
mirus, Marsh, 1878, Am. Jour. Sci. and Arts, 3d ser., vol. 15, p. 411, Permian.
Ornithichnites, Hitchcock, 1836, Am. Jour. Sci. and Arts, vol. 29. [Ety. *ornithos*, a bird; *ichnos*, a footprint.] This is not properly a generic name. No bird-tracks are known in Palaeozoic rocks. The Batrachian tracks referred to it belong to another genus.
culbertsoni, King, 1845, Am. Jour. Sci. and Arts, vol. 48, p. 345, Coal Meas.
gallindoides, King, 1845, Am. Jour. Sci. and Arts, vol. 48, p. 344, Coal Meas.
PANTYLUS, Cope, 1881, Bull. U. S. Geo. Sur. Terr., vol. 6, No. 1, p. 79. [Ety. *pan*, all; *tylos*, a knob.] Founded upon the crania; ossification complete, leaving only orbits, nostrils, and parietal fontanel; surface sculptured; mandible with an angular process; teeth conic, obtuse, larger anteriorly; mandible supporting several rows of teeth, which oppose a pavement of obtuse teeth on the palatine or anterior part of pterygoid bones; quadratojugal and malar bones well developed; no lyra or mucous grooves. Type *P. cordatus*.
cordatus, Cope, 1881, Bull. U. S. Geo. Sur. Terr., vol. 6, p. 79, Permian.
PARIOTICHUS, Cope, 1878, Pal. Bull., No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 508. [Ety. *pareia*, the cheek; *teichos*, a wall.] Founded on the cranium. Temporal fossae were covered by a roof continuous with the postorbital region; zygomatic arch extends low down; orbits lateral; muzzle short, with terminal nares; teeth rooted, crowns obtuse, with outting edge. Type *P. brachyops*.
brachyops, Cope, 1878, Pal. Bull., No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 508, Permian.
megalops, Cope, 1883, Pal. Bull., No. 36, and Proc. Am. Phil. Soc., vol. 20, p. 630, Permian.
Parioxys, Cope, 1878, Pal. Bull., No. 29, and Proc. Am. Phil. Soc., vol. 17, p. 521. [Ety. *parcia*, cheek; *oxys*, sharp.] Syn. for *Eryops*.
ferricolus, see *Eryops ferricolus*.
PELION, Wyman, 1868, Proc. Acad. Nat. Sci. Phil., p. 211. [Ety. proper name.] Founded upon an inferior view of part of the skeleton; head as broad as long,

semielliptical; angles of mandibles project backward; mandibular rami slender, curved; vertebrae have centra as broad as long, and medially contracted; fore limbs stout; humeri long, thickened proximally, flattened and dilated distally; ulna and radius united proximally, narrowing the arm proximally while expanded distally; left hand exhibits four digits, of which the third from the inner is the longer; number of phalanges is 2, 3, 4?; carpus not osseous. Type *P. lyelli*.

FIG. 1190.—*Pelion lyelli*.

lyelli, Wyman, 1858, (*Raniceps lyelli*), Am. Jour. Sci. and Arts, 2d ser., vol. 25, p. 158, and Ohio Pal., vol. 2, p. 389, Coal Meas.

FIG. 1191.—*Philegethonlia linearis*.

PHILEGETHONTIA, Cope, 1871, Proc. Am. Phil. Soc., p. 177. [Ety. *Philegethon*, a burning river of hell.] Head elongate,

of mandibles pro-
nubular rami slender
have centra as
medially contracted;
numeri long, thick-
attened and dilated
radius united proxi-
the arm proximally
tally; left hand ex-
of which the third
the longer; number
47; carpus not os-
ill.

lyelli.

(*Raniceps lyelli*,) Am.
2d ser., vol. 25, p. 158,
2, p. 389, Coal Meas.

linearis.

pe, 1871, Proc. Am.
7. [*Phlegethon*, a
hell.] Head elongate,

triangular; body and tail extremely
elongate; dorsal vertebræ without ribs,
caudals without dilated spines; no
vertebræ armature; no limbs. Type
P. linearis.

linearis, Cope, 1871, Proc. Am. Phil. Soc.,
p. 177, and Ohio Pal., vol. 2, p. 367,
Coal Meas.

serpens, Cope, 1871, Proc. Am. Phil. Soc.,
p. 177, and Ohio Pal., vol. 2, p. 367,
Coal Meas.

PLEUROPTYX, Cope, 1875, Ohio Pal., vol. 2,
p. 370. [*Ety. pleura*, a rib; *ptyx*, a fold,
wing.] Founded upon the vertebral
column and ribs; vertebræ of moderate
length, zygapophyses well developed,
short neural spine in the dorsal region,
not sculptured; ribs short, stout, and
support a wing on the posterior or con-
vex border, which expands downward,
and then abruptly contracts to the
shaft; it is broad and truncate, and in-
cludes a medullary cavity partially filled
with cancellated tissue. Type P. clavatus.

clavatus, Cope, 1875, Ohio Pal., vol. 2, p.
370, Coal Meas.

PYTONIUS, Cope, 1875, Ohio Pal., vol. 2, p.
373. [*Ety. python*, a fan.] Form elon-

FIG. 1192.—*Ptyonius serrula*.

gate, with
long tail and
lanceolate
cranium;
limbs weak,
only poste-
rior known;
three pec-
toral shields;
a b o m e n
protected by
packed os-
seous rods
arranged in
chevron with
angle direct-
ed forward;
neural and
h æ m a l
spines of
caudal ver-
tebræ ex-
panded and
fan-like; ribs
well devel-
oped; teeth
small, num-
erous, sim-
ple or groov-
ed. Type P.
nummifer.

marshi, Cope,
1875, (*Colos-
teus marshi*,)
Trans. Am.
Phil. Soc.,
vol. 14, p. 24,
and Ohio

Pal., vol. 2, p. 375, Coal Meas.

nummifer, Cope, 1875, Ohio Pal., vol. 2,
p. 374, Coal Meas.

pectinatus, Cope, 1868, (*Sauropleuræ pec-
tinata*, Proc. Acad. Nat. Sci., p. 216, and
Ohio Pal., vol. 2, p. 377, Coal Meas.

serrula, Cope, 1871, (*Osteocephalus ser-
rula*,) Proc. Am. Phil. Soc., p. 177, and
Ohio Pal., vol. 2, p. 379, Coal Meas.

vinchellanus, Cope, 1871, (*Osteocephalus
vinchellanus*,) Proc. Am. Phil. Soc., p.
177, and Ohio Pal., vol. 2, p. 376, Coal
Meas.

Pygopterus, Agassiz, 1833, Recherch. Poiss.
Foss.

scutellatus, see *Colosteus scutellatus*.

Raniceps, Wyman, 1858, Am. Jour. Sci. and
Arts. The name was preoccupied by
Cuvier, and Pelion has been substituted.

lyelli, see Pelion lyelli.

Rhachitonus, Cope, 1878, Pal. Bull. No.
29, and Proc. Am. Phil. Soc., vol. 17, p.
526. [*Ety. rachis*, the backbone; *tonos*,
cut.] Syn. for *Eryops*.

valens, syn. for *Eryops megacephalus*.

SAUROPLEURA, Cope, 1868, Proc. Acad. Nat.
Sci. Phil., p. 215, and Ohio Pal., vol. 2,
p. 402. [*Ety. sauros*, a lizard; *pleuron*,
a rib.] Vertebræ and ribs well devel-

oped; limbs four, large; five digits in
the fore foot; carpus cartilaginous; ven-
tral armature of closely arranged rhom-
boidal scuta, arranged in lines closely
placed in chevrons, with the angle an-
terior; teeth of *Labyrinthodont* type,
with deeply inflected enamel and acute
apex. Type S. digitata.

digitata, Cope, 1868, Proc. Acad. Nat. Sci.
Phil., p. 216, and Ohio Pal., vol. 2, p.
403, Coal Meas.

longipes, see *Tuditatus longipes*.

newberryi, Cope, 1875, Ohio Pal., vol. 2,
p. 404, Coal Meas.

pectinata, see *Ptyonius pectinatus*.

remex, see *Osteocephalus remex*.

SAUROPOUS, Lea, 1849, Trans. Am. Phil. Soc.,
vol. 10. [*Ety. sauros*, a lizard; *pous*,
foot.] Founded upon tracks; five robust
toes, the inner having the same direc-
tion as the others, and not divergent as
in *Chirotherium*; palm or sole short,
wide. Type S. primævus.

FIG. 1193.—*Sauropus primævus*.

primævus, Lea, 1849, Trans. Am. Phil.
Soc., vol. 10, Coal Meas.

sydnensis, Dawson, 1868, Acad. Geol., p.
358, Coal Meas.

unguifer, Dawson, 1872, Geo. Mag. Lond.,
vol. 9, Coal Meas.

- SPHENACODON**, Marsh, 1878, *Am. Jour. Sci.*, and *Arts*, 3d ser., vol. 15, p. 410. [Ety. *sphen*, a wedge; *akis*, a barb; *odon*, tooth.] Type *S. ferox*.
- ferox*, Marsh, 1878, *Am. Jour. Sci.* and *Arts*, 3d ser., vol. 15, p. 410, Permian.
- SPHEROPEZIUM**, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, p. 345. [Ety. *sphaira*, sphere; *pezia*, sole of the foot.] Founded upon tracks representing a round depression for the ball of the foot, and five depressions for digits. Type *S. leptodactylum*.
- leptodactylum*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, p. 345, Coal Meas.
- ovidactylum*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, p. 347, Coal Meas.
- pachydactylum*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, p. 346, Coal Meas.
- therodactylum*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, p. 346, Coal Meas.
- THENAROPUS**, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, p. 343. [Ety. *thenaros*, palm of the hand; *pous*, foot.] Founded upon tracks. Type *T. heterodactylus*.
- heterodactylus*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, Coal Meas.
- leptodactylus*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, Coal Meas.
- ovidactylus*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, Coal Meas.
- pachydactylus*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, Coal Meas.
- spherodactylus*, King, 1845, *Am. Jour. Sci.* and *Arts*, vol. 48, Coal Meas.
- THEROPLKURA**, Cope, 1878, *Pal. Bull.* No. 29, and *Proc. Am. Phil. Soc.*, vol. 17, p. 519. [Ety. *thero*, to burn; *pleura*, side.] Pelycosaurian reptiles with free neural arch, and a capitular costal articulation on the centrum, and no known inter-centrum; neural spines not elongate; teeth equal. Type *T. retroversa*.
- obtusidens*, Cope, 1880, *Pal. Bull.* No. 32, p. 4, Permian.
- retroversa*, Cope, 1878, *Pal. Bull.* No. 29, and *Proc. Am. Phil. Soc.*, vol. 17, p. 519, Permian.
- triangulata*, Cope, 1878, *Pal. Bull.* No. 29, and *Proc. Am. Phil. Soc.*, vol. 17, p. 520, Permian.
- uniformis*, Cope, 1878, *Pal. Bull.* No. 29, and *Proc. Am. Phil. Soc.*, vol. 17, p. 519, Permian.
- THYRSIDIUM**, Cope, 1875, *Ohio Pal.*, vol. 2, p. 365. [Ety. *thyrsos*, a rod with leaves.] Founded upon a latero-inferior view of the spinal column; diapophyses enlarged, fan-like; centra contracted; abdomen protected by hair-like rods in chevron, with angle directed forward. Type *T. fasciculare*.
- fasciculare*, Cope, 1875, *Ohio Pal.*, vol. 2, p. 365, Coal Meas.
- TRIMERORACHIS**, Cope, 1878, *Pal. Bull.* No. 29, and *Proc. Am. Phil. Soc.*, vol. 17, p. 524. [Ety. *trimeros*, tripartite; *rachis*,

the backbone.] Centrum rhachitomous, represented by three cortical ossifications of the chorda-sheath, a median inferior and two lateral; the lateral are distinct, and in contact with the neurapophyses above, and the posterior border of the median segment in front; neural arch joins the lateral elements, and is in slight contact with the lateral summits of the inferior element; the halves of the neural arch are co-ossified, and support zygapophyses, but no neural spine; cranial bones sculptured; parasphenoid flat; external nostrils large, superior; angle of mandible little produced; glenoid cavity transverse; deep internal pterygoid fossa; no coronoid process; symphy. short; teeth conic, two series in the upper jaw, the large ones anterior; ribs short, heads expanded. Type *T. insignis*.

bilobatus, Cope, 1883, *Pal. Bull.* No. 36, and *Proc. Am. Phil. Soc.*, vol. 20, p. 629, Permian.

insignis, Cope, 1878, *Pal. Bull.* No. 29, and *Proc. Am. Phil. Soc.*, vol. 17, p. 524, Permian.

TUDITANUS, Cope, 1871, *Proc. Am. Phil. Soc.*, p. 177, and *Ohio Pal.*, vol. 2, p. 391. [Ety. proper name.] Cranium broad, flat, bones sculptured; teeth on premaxillary and maxillary bones of nearly equal sizes; three pectoral shields, sculptured externally; form lizard-like; two pairs of medium liabae; no chevron abdominal rods. Type *T. punctulatus*.

FIG. 1194.—*Tuditanus radiatus*.

brevirostris, Cope, 1874, *Trans. Am. Phil. Soc.*, vol. 14, p. 10, and *Ohio Pal.*, vol. 2, p. 393, Coal Meas.

huxleyi, Cope, 1874, *Trans. Am. Phil. Soc.*, p. 10, and *Ohio Pal.*, vol. 2, p. 397, Coal Meas.

longipes, Cope, 1874, (*Sauropleuria longipes*) *Trans. Am. Phil. Soc.*, vol. 14, p. 10, and *Ohio Pal.*, vol. 2, p. 398, Coal Meas.

entrum rhachito-
three cortical os-
sorda-sheath, a me-
lateral; the lateral
contact with the
, and the posterior
segment in front;
e lateral elements,
et with the lateral
rior element; the
arch are co-ossi-
apophyses, but no
bones sculptured;
external nostrils
of mandible little
cavity transverse;
oid fossa; no coro-
ny's short; teeth
the upper jaw, the
ribs short, heads
insignis.

Pal. Bull. No. 36,
Soc., vol. 20, p. 629,

al. Bull. No. 29, and
c., vol. 17, p. 524,

Proc. Am. Phil. Soc.,
Pal., vol. 2, p. 391.

] Cranium broad,
red; teeth on pre-
lary bones of nearly
e pectoral shields,
ly; form lizard-like;
n limbs; no chevron
type T. punctulatus.

anus radiatus.

74, Trans. Am. Phil.
Soc., and Ohio Pal., vol.

Trans. Am. Phil. Soc.,
Pal., vol. 2, p. 397, Coal

(Sauropleura longi-
Phil. Soc., vol. 14, p.
Soc., vol. 2, p. 398, Coal

mordax, Cope, 1875, Ohio Pal., vol. 2, p. 395,
syn. for Ceraterpeton punctolluatum.

obtusum, Cope, 1868, (Dendererpeton ob-
tusum,) Proc. Acad. Nat. Sci. Phil., p.
213, and Ohio Pal., vol. 2, p. 396, Coal
Meas.

punctulatus, Cope, 1874, Trans. Am. Phil.
Soc., vol. 14, p. 10, and Ohio Pal., vol.
2, p. 392, Coal Meas.

radiatus, Cope, 1874, Trans. Am. Phil.
Soc., vol. 14, p. 10, and Ohio Pal., vol.
2, p. 394, Coal Meas.

tabulatus, Cope, 1877, Proc. Am. Phil.
Soc., p. 577, Coal Meas.

ZATRACHYS, Cope, 1878, Pal. Bull. No. 29,
and Proc. Am. Phil. Soc., vol. 17, p.

523. [Ety. *sa*, an intensive; *trachys*,
rough.] Teeth in single series, and
anchylosed to the bottom of a shallow
groove, the external boundary of which
is most prominent, so the attachment
of the teeth is shortly pleurodont;
teeth have conic crowns and basal
grooves; cranium sculptured, its table
with a notch on each side; two occip-
ital condyles; no intercalary horns.
Type *Z. serratus*.

apicalis, Cope, 1881, Am. Naturalist, vol.
15, p. 1020, Permian.

serratus, Cope, 1878, Pal. Bull. No. 29, and
Proc. Am. Phil. Soc., vol. 17, p. 523,
Permian.

GLOSSARY

— OF —

SPECIFIC NAMES IN USE IN NORTH AMERICAN PALÆONTOLOGY.

Every adjective specific name must agree in gender with the genus to which it belongs; hence, the masculine, feminine, and neuter endings are indicated. Nouns do not change the termination, but remain the same, no matter what the gender of the genus may be. The names of persons and places are not included in this Glossary, because they should be known by the terminations they take when reduced to specific names.

Abacus—A table divided into squares.
Abbreuiatus, a, um—Abbreviated, shortened.
Abnormis, e—Abnormal.
Abruptus, a, um—Abrupt, broken.
Abscissus, a, um—Steep, abrupt.
Acanthophorus, a, um—Thorn-bearing.
Acanthoptera—Spine-wing.
Acinctus, a, um—Girded.
Acervulosus, a, um—Many clustered together.
Aceruus—A heap considered as a body.
Acicula—A small pin or needle.
Acicularis, e—Full of small pins or needles, acicular.
Aciculatus, a, um—Like a small needle.
Acidentatus, a, um—Needle-toothed, sharp-toothed.
Acies—The edge or sharp point.
Acinaeiformis, e—Scimitar-shaped.
Acinus—Any berry, or the kernel in the berry.
Acis—A small javelin.
Acmea—Edge, point.
Acrocarpus, a, um—Pointed fruit.
Actuarius, a, um—Swift, agile.
Aculeatus, a, um—Thorny, pointed, sharp.
Aculeolatus, a, um—Thorny.
Acuminatus, a, um—Sharp-pointed.
Acus—A pin or needle.
Acutangulus—Acute angle.
Acuticosta—Sharp rib.
Acutidactylus—Sharp-fingered.
Acutifolius, a, um—Having acute leaves.
Acutiratus, a, um—Sharp-ridged.
Acutiplicatus, a, um—Sharp-plicated.
Acutiradiatus, a, um—Sharp-rayed.
Acutirostris—Sharp beak.
Acutulus, a, um—Somewhat pointed.
Acutus, a, um—Acute, sharpened.
Adiantites—From resemblance to *Adiantum*.
Adiantoides—Like *Adiantum*.

Adductus, a, um—Stretched, contracted.
Adjunctivus, a, um—Joined, united.
Adjunctus, a, um—Joined, connected.
Adnascens—Growing upon.
Adnatus, a, um—Adnate.
Adorabilis, e—Worthy of adoration.
Adultus, a, um—Adult.
Aduncus, a, um—Bent inward, hooked.
Aegilops—An acorn.
Æmulus, a, um—Emulous, vying with.
Enigma—Obscure, a riddle.
Equalis, e—Equal, like.
Equibrachiatus, a, um—Equal-armed.
Equicostatus, a, um—Equal-ribbed.
Equidistans—Equidistant.
Equilateralis, e—Equilateral.
Equiradiatus, a, um—Equal-rayed.
Equivalvis, e—Equal-valved.
Equus, a, um—Plain, even, level, equal.
Affinis, e—Related, or near to.
Agellus—A small field.
Agglomeratus, a, um—Gathered into a mass.
Aggregatus, a, um—Aggregated.
Agilis, e—Agile, nimble.
Agrarius, a, um—Pertaining to fields or country.
Agrestis, e—Pertaining to the country.
Alatus, a, um—Winged.
Albus, a, um—White.
Alcicornis, e—Elk-horned.
Alectiformis, e—In form like *Alecto*.
Aliger, gera, gerum—Bearing wings.
Alsus, a, um—Cold.
Alternans—Alternating.
Alternatus, a, um—Alternate.
Alterniradiatus, a, um—Alternately rayed.
Alternistriatus, a, um—Alternately striated.
Alternus, a, um—Alternate.
Altilis, e—Flattened.
Altidosatus, a, um—High-backed.
Altiplicatus, a, um—Having high plications.
Altirostris—High beak.

Altus, a, um—High, great, deep.
 Alveatus, a, um—Hollowed out like a trough.
 Alveolaris, e—Small-channeled.
 Alveolatus, a, um—Hollowed out.
 Alveolus—A small cavity.
 Amarus, a, um—Bitter, brackish.
 Ambiguus, a, um—Doubtful, changeable.
 Amoenus, a, um—Pleasant, charming.
 Amphibolus, a, um—Ambiguous.
 Amplexicaulis, e—Embracing the stalk or stem.
 Amplexus—An encircling, surrounding.
 Ampliatus, a, um—Enlarged.
 Amplus, a, um—Ample, spacious, roomy.
 Ampullaceus, a, um—In the form of a flask.
 Amygdaliformis, e—In form like the almond.
 Anabathra—A ladder.
 Analogus, a, um—Analogous.
 Anatiformis, e—Like Anatifa.
 Anatinus, a, um—Of or pertaining to the duck.
 Ancypus—Double, two-headed, doubtful.
 Anchoralis, e—Of or pertaining to an anchor.
 Ancilla—A hand-maid.
 Anellus—A little ring.
 Anguineus, a, um—Serpent-like.
 Angularis, e—Angular, cornered.
 Angulatus, a, um—Having corners.
 Angulosus, a, um—Full of corners.
 Angustatus, a, um—Narrowed.
 Angustifolius, a, um—Narrow-leaved.
 Angustipinna—A narrow feather.
 Angustus, a, um—Narrow, straight, not specious.
 Annectans—Connected together.
 Annularifolius, a, um—Having ring-shaped leaves.
 Annulatus, a, um—Annulated, ringed.
 Annuliferus, a, um—Ring-bearing.
 Anomalus, a, um—Anomalous, not coming under the rule.
 Anonymus, a, um—Nameless.
 Anticeptus, a, um—Anticipated.
 Antennarius, a, um—Pertaining to the antennae.
 Antheloideus, a, um—Like Anthelia.
 Anthracinus, a, um—Coal-black.
 Antiquarius, a, um—Pertaining to antiquity.
 Antiquatus, a, um—Antiquated, ancient.
 Antiquus, a, um—Ancient, old.
 Annularis, e—Relating to the signet ring.
 Annularius, a, um—Of or pertaining to the signet ring.
 Annulatus, a, um—Annulated, ringed.
 Annulus—A ring.
 Apertus, a, um—Opened, uncovered.
 Apicalis, e—Sharp-pointed.
 Apicatus, a, um—Having a pointed termination.
 Aplatus, a, um—Flattened.
 Appendiculatus, a, um—Having lateral appendages.
 Approximatus, a, um—Approximated, approaching near to.
 Aprinus, a, um—Pertaining to a wild boar.
 Aquilinus, a, um—Pertaining to the eagle, aquiline.
 Arachniformis, e—In form like a spider.
 Arachnoideus, a, um—Like a cobweb.

Araneolus—A small spider.
 Aratus, a, um—Plowed.
 Arboreascens—Tree-like, arborescent.
 Arhoreus, a, um—Pertaining to a tree.
 Arbuscula—A shrub.
 Arcanus, a, um—Closed, shut up.
 Archimedeiformis, e—Archimedeiform.
 Arcticus, a, um—Arctic.
 Arctifossa—Close-wrinkle.
 Arctiporus, a, um—Having narrow pores.
 Arctostriatus, a, um—Closely striated.
 Arctirostratus, a, um—Narrow-beaked.
 Arctisegmentus, a, um—Having narrow segments.
 Arctus, a, um—Closed, pressed together, short.
 Arcuatus, a, um—Bent, curved like a bow.
 Arcuosus, a, um—Full of arches, bent over.
 Arenaceus, a, um—Sandy.
 Arenarius, a, um—Pertaining to sand.
 Arenosus, a, um—Sandy.
 Areolatus, a, um—Divided into irregular squares, or angular spaces.
 Argentarius, a, um—Of or pertaining to silver.
 Argentus, a, um—Of or made of silver.
 Argentarius, a, um—Of or belonging to Silver City.
 Argutus, a, um—Sharply defined, distinct.
 Arietinus, a, um—Of or from a ram.
 Armatus, a, um—Armed, equipped.
 Armiger, era, erum—Armed, war-like.
 Armosus, a, um—Many armed.
 Arrectarius, a, um—Erect.
 Arrectus, a, um—Erect, steep.
 Arrosus, a, um—Gnawed.
 Articulus, a, um—Furnished with joints, articulated.
 Articulosus, a, um—Full of knots, or divisions.
 Artemisiifolius, a, um—Like the plant Artemisia.
 Arundinaceus, a, um—Like a reed.
 Aspectans—Expected, looked for.
 Asper, era, erum—Rough, uneven.
 Asperatus, a, um—Roughened, irregular.
 Aspersus, a, um—Scattered, dispersed.
 Asprattilis, e—Rough.
 Assimilis, e—Similar.
 Atavus—Ancestor.
 Attenuatus, a, um—Made thin, attenuated.
 Attritus, a, um—Worn.
 Aucella—A little bird.
 Audaculus, a, um—Bold.
 Augustatus, a, um—Majestic.
 Augustus, a, um—August.
 Auleticus, a, um—That is suitable for a pipe.
 Aureatus, a, um—Adorned.
 Auricula—The ear.
 Auritus, a, um—Eared.
 Australis, e—Southern.
 Auxiliarius, a, um—Helping.
 Avicula—A small bird.
 Avitus, a, um—Ancestral.
 Avus—Grandfather.
 Bacca—A berry, a small, round fruit.
 Bacillum—A small staff.
 Baculiformis, e—Staff-shaped.

- Baccula—A staff or cudgel.
 Balanoides—Like Balaanus.
 Balanus—An acorn.
 Balteatus, a, um—Belted.
 Barbatus, a, um—Bearded.
 Barydactylus, a, um—Heavy-fingered.
 Basalis, e—Pertaining to the base.
 Basalticus, a, um—Basaltic.
 Basilaris, e—Relating to the base.
 Basilicus, a, um—Splendid.
 Belemnura—Having a tail like a dart.
 Bellarugosus, a, um—Beautifully wrinkled.
 Bellatrema—Beautiful opening.
 Bellatulus, a, um—Pretty, neat.
 Bellicinctus, a, um—Beautifully banded.
 Bellicosus, a, um—Warlike.
 Bellifer, era, erum—Warlike.
 Bellilineatus, a, um—Beautifully lined.
 Bellipunctus, a, um—Beautifully dotted.
 Bellistriatus, a, um—Beautifully striated.
 Bellulus, a, um—Very beautiful, pretty.
 Bellus, a, um—Beautiful.
 Bertholletiformis, e—Like Bertholletia.
 Biacutus, a, um—Two-pointed.
 Bialveatus, a, um—Two-channelled.
 Biangulatus, a, um—Two-angled.
 Bicarinatus, a, um—Two-keeled.
 Bicarpus, a, um—Two-fruited.
 Biceps—Two-headed.
 Bieinctus, a, um—Two-banded.
 Bi clavatus, a, um—Two-clubbed.
 Bicorniger, era, erum—Two-horned.
 Bicornis, e—Two-horned.
 Bicornutus, a, um—Two-horned.
 Bicostratus, a, um—Two-ribbed.
 Bicristatus, a, um—Double-peaked or two-crested.
 Bienspidatus, a, um—Two-pointed.
 Bidens—Having two teeth, two-pronged.
 Bidentatus, a, um—Double-toothed.
 Bidorsalis, e—Double-backed.
 Bidorsatus, a, um—Having a double back.
 Bifarius, a, um—In two ways or parts, double.
 Bifidus, a, um—Cleft into two parts.
 Bifidus, a, um—Bifid, cloven in two parts.
 Bifissus, a, um—Cleft into two parts.
 Bifolius, a, um—Two-leaved.
 Biforatus, a, um—Two-holed or double-doored.
 Biformatus, a, um—Two-shaped.
 Biformis, e—Two-formed.
 Bifrons—With two foreheads.
 Bifurca—A two-pronged fork.
 Bifurcatus, a, um—Bifurcated, forked.
 Bifureus, a, um—Two-pronged.
 Bimbricatus, a, um—Double-imbricated.
 Bijugus, a, um—Yoked two together.
 Bilabialis, a, um—Two-lipped.
 Bilamellatus, a, um—Having double lamellae.
 Bilateralis, e—Two-sided.
 Bilineatus, a, um—Two-lined.
 Biliratus, a, um—Two-furrowed.
 Bilix—Woven with a double thread, two-threaded.
 Bilobatus, a, um—Two-lobed.
 Bilobus, a, um—Two-lobed.
 Bimestialis, e—Having two middle parts.
- Bimacronatus, a, um—Two-point'ed.
 Binervis, e—Two-nerved.
 Binodus—Double knot.
 Binumbonatus, a, um—Having double umbones.
 Bipartitus, a, um—Two-parted.
 Bipennis, e—Two-winged.
 Biplicatus, a, um—Two-plicated, or in two folds.
 Bipunctatus, a, um—Bipunctate.
 Bipyramidalis, e—Double-pyramidal.
 Bisectus, a, um—Divided.
 Biserialis, e—In two series.
 Biseriatus, a, um—Having two rows or series.
 Biserrulatus, a, um—Double-serrulated.
 Bisinuatus, a, um—Having two depressions or furrows.
 Bispinulatus, a, um—Two-spined.
 Bispiralis, e—Two-whorled.
 Bistriatus, a, um—Two-striated.
 Bisulcatus, a, um—Two-furrowed.
 Bisulcus, a, um—Cloven.
 Bituberculatus, a, um—Double-tuberculated.
 Biturbinatus, a, um—Double-turbinated.
 Bivertex—Double head.
 Bivus, a, um—Having two ways or passages.
 Bivittatus, a, um—Two-banded.
 Bivolvis, e—Two-rolled.
 Blatta—A cockroach or moth.
 Blattinoides—Like Blattina.
 Bombifrons—Having a hollow front.
 Borassifolius, a, um—Leaved like Borassus.
 Borealis, e—Northern.
 Bovidens—Ox tooth.
 Bovipellalis, e—Ox-footed.
 Brachialis, e—Having arms.
 Brachiatus, a, um—Having arms.
 Brachium—An arm.
 Brachynotus, a, um—Short-ridged.
 Brachyops—Short sight.
 Bracteatus, a, um—Covered with plates, beautiful.
 Breviceps—Short head.
 Brevicornis, e—Short-horned.
 Brevicostatus, a, um—Short-ribbed.
 Breviculus, a, um—Somewhat shortened.
 Brevicurvatus, a, um—Short-curved.
 Brevifolius, a, um—Short-leaved.
 Brevilineatus, a, um—Short-lined.
 Brevilobatus, a, um—Short-lobed.
 Brevilobus—Short lobe.
 Brevimarginatus, a, um—Short-margined.
 Brevinodus—Short node or short knot.
 Breviplicatus, a, um—Short-plicated.
 Breviposticus, a, um—Made short behind.
 Breviradiatus, a, um—Short-rayed.
 Brevis, e—Short.
 Brevisulcatus, a, um—Short-furrowed.
 Brevisculus, a, um—Very short.
 Brisa—Grape-skins.
 Bryonoides—Like moss.
 Buccinum—A trumpet or horn.
 Bucculentus, a, um—Wide-mouthed.
 Bufo—A toad.
 Bulbaceus, a, um—Bulbous.

Bulbosus, a, um—Bulbous.
 Bulbus—A bulb.
 Bulmiformis, e—Like *Bulimus*.
 Bulla—A round object, bubble.
 Bullatus, a, um—Studded with knobs.
 Bulloides—Like a bubble.
 Bullulatus, a, um—Little -sided.
 Bursa—A purse.
 Bursiformis, e—Purse-shaped.
 Cadens—Falling, terminating.
 Caduceus—The herald's staff.
 Cæcigenus, a, um—Born blind.
 Calamen—A bass relief.
 Cælator—A carver.
 Cælatus, a, um—Engraved, carved.
 Cespitosus, a, um—Turf-like.
 Calamitoideus, a, um—Like a Calamite.
 Calamus—A reed.
 Calantica—A covering for the head.
 Calathus—A wicker basket.
 Calcaratus, a, um—Spurred, spur-shaped.
 Calcariformis, e—Like a spur.
 Calceolus—A small shoe.
 Calciferus, a, um—Calciferous.
 Calculus—A small stone.
 Caliculus—A small cup.
 Calix—A cup.
 Callicephalus, a, um—Having a beautiful head.
 Calliteles—A beautiful tail.
 Callosus, a, um—Thick-skinned, callous.
 Calycinus—A little calyx.
 Calycularis, e—Like a little cup or flower-bud.
 Calyenloides—Like a little cup.
 Calyculus—A flower-cup.
 Calymenoides—Like Calymene.
 Calyx—The cup of a flower.
 Cameratus, a, um—Arched.
 Cameriferus, a, um—Chambered.
 Cammarus—A lobster.
 Campaniformis, e—Bell-formed.
 Campanulatus, a, um—Bell-shaped.
 Camurus, a, um—An arch, turned inward.
 Canaliculatus, a, um—Channeled, canalculated.
 Canalis—A channel or groove.
 Cancellatus, a, um—Cross-barred, cancellated.
 Cancellus, a, um—Finely cancellated or latticed.
 Canna—A reed.
 Canneus, a, um—Made of reeds.
 Canniformis—Like Canna.
 Cannaliratus, a, um—Reed-furrowed.
 Cannula—A small reed.
 Capax—Large, spacious.
 Capillaceus, a, um—Similar to hair, stringy.
 Capillaris, e—Of or pertaining to the hair.
 Capillatus, a, um—Having hair.
 Capillosus, a, um—Very hairy.
 Capitalis, e—Relating to the head.
 Capitatus, a, um—Having a head.
 Capitellum—A small head.
 Capitollinus, a, um—Pertaining to the capitol, a tower.
 Caponiformis, e—Capon-form d.
 Capreolus—Prope, stays.

Capularis, e—Pertaining to a coffin.
 Capuloides—Like a capulus.
 Capulus—A coffin or a handle.
 Caput-serpentis—Serpent-head.
 Caput-testudinis—Turtle-head.
 Carabus—A small wicker boat.
 Carbonarius, a, um—Of or relating to coal.
 Carcharidens—Dog-fish tooth.
 Cardiiformis, e—Like a cardium.
 Cardinalis, e—Of or pertaining to a door-hinge, or principal.
 Cardinatus, a, um—Jointed, fitted to, hinged.
 Cardinens, a, um—Of or pertaining to a door-hinge.
 Carica—A kind of fig.
 Carinatus, a, um—Keeled.
 Cariniferus, a, um—Keel-bearing.
 Carnosus, a, um—Fleshy.
 Carus, a, um—Precious, valued.
 Casanea—A chestnut.
 Cactatus, a, um—Frail, easily broken.
 Catastomus, a, um—Gaping at the lower end.
 Catenoides—Chain-like.
 Catennulatus, a, um—A little chain.
 Catilloformis, e—Dish-shaped.
 Catilloides—Dish-like.
 Catillus—A small dish.
 Catinus—A bowl.
 Caudagalli—Tail of a cock.
 Caudatus, a, um—Having a tail.
 Cauliculus—Small stalk or stem of a plant.
 Cavernosus, a, um—Full of hollows.
 Cavifolius, a, um—Hollow-leaved.
 Cavumbilicatus, a, um—Having a hollow umbilicus.
 Cavumbona—Hollow umbo.
 Cavus, a, um—Hollow, concave.
 Celator—A concealer, hider.
 Celebris, a, um—Abundant.
 Celer—Swift, fleet.
 Cellulosus, a, um—Full of cells.
 Celsipora—High pore.
 Celus, a, um—High.
 Centennialis, e—The 100th year.
 Centralis, e—In the middle, central.
 Centratus, a, um—Central.
 Centrilineatus, a, um—Central-lined.
 Centronatus, a, um—Having knots or points.
 Centrosus, a, um—In the central point.
 Cerasiformis, e—Like a dried cherry.
 Cerithioides—Like *Cerithium*.
 Cervicornis, e—Deer-horned.
 Cervinus, a, um—Pertaining to a deer.
 Cessator—An idler, loiterer.
 Cetratus, a, um—Shield-bearing.
 Chærophylloides—Like *Chærophylum*.
 Chiriformis, e—Hand-shaped.
 Chiromorphus, a, um—Hand-formed.
 Chromaticus, a, um—Chromatic, colored.
 Chrysalis—Chrysalis.
 Cicatricosus, a, um—Full of scars.
 Ciceronius—Having warts.
 Ciliatus, a, um—Haired on the margin, fringed.
 Cinctosus, a, um—Full of bands, girded.
 Cinctulus—A small girt.

o a coffin.
 us.
 ndle.
 head.
 head.
 boat.
 relating to coal.
 oth.
 ardium.
 aining to a door.
 painted, fitted to,
 pertaining to a
 l.
 bearing.
 valued.
 asily broken.
 ing at the lower
 ttle chain.
 aped.
 ck.
 ng a tail.
 or stem of a plant.
 l of hollows.
 ow-leaved.
 Having a hollow
 mbo.
 concave.
 ider.
 lant.
 of cells.
 0th year.
 dle, central.
 ral.
 Central-lined.
 Having knots or
 he central point.
 a dried cherry.
 ithium.
 ured.
 aining to a deer.
 erer.
 l-bearing.
 e Chærophyllum.
 shaped.
 Hand-formed.
 Chromatic, colored.
 all of scars.
 arts.
 ed on the margin,
 of bands, girded.
 rt.

Cinctura—A girdle.
 Cinetus, a, um—Banded, girded.
 Cinetulus, a, um—Girded.
 Cingulatus, a, um—Encircled with lines,
 girded.
 Cingulosus, a, um—Covered with lines or
 zones.
 Cingulum—A zone.
 Circinatus, a, um—Compassed, rounded.
 Circinctus, a, um—Encompassed.
 Circularis, e—Circular, round.
 Circulus—A circle.
 Circumliratus, a, um—Circular-lined.
 Cistella—A small box.
 Cistula—A little chest or coffer.
 Citus, a, um—Swift, speedy.
 Clarus, a, um—Clear, brilliant, distinct.
 Clathratus, a, um—Cross-barred, latticed.
 Clausus, a, um—Closed up.
 Clava—A stick.
 Clavacoides, a, um—Club-shaped.
 Clavatus—A little club.
 Clavatus, a, um—Knotted, club-shaped.
 Clavicula—A small twig.
 Clavifrons—Having a club-like front.
 Claviger—A club-bearer.
 Clavigerus, a, um—Club-bearing.
 Clavis—A bar.
 Clavulus—A little club, a small swelling.
 Clavus—A nail, spike.
 Clinatus, a, um—Inclined, bent.
 Clinocameratus, a, um—Curve-chambered.
 Clipeatum—Furnished with a shield.
 Clipeiformis, e—Shield-like.
 Clivus, a, um—Full of hills.
 Clivulus, a, um—Having little hills.
 Clivulus—A little hill.
 Clymenioides—Like Clymenia.
 Clypeatus, a, um—Armed with a shield.
 Clytis—Celebrated.
 Coalescens—Growing together.
 Coalitus, a, um—Grown together.
 Coaptus, a, um—Closely joined.
 Coarctatus, a, um—Compressed, joined.
 Cochlearis, e—In the form of a snail shell.
 Cochleatus, a, um—Spiral.
 Cochleola—A small snail.
 Cognatus, a, um—Near to, cognate.
 Cohærens—Adhering together.
 Collatus, a, um—Joined together, collected.
 Collectus, a, um—Collected.
 Colliculus—A little hill.
 Colligatus, a, um—Bound together, fastened.
 Collinus, a, um—Hilly.
 Colon—The great intestine.
 Colubrellus—A little snake.
 Colubrinus, a, um—Like a snake.
 Colubrosus, a, um—Winding.
 Columnella—A small column.
 Columnellatus, a, um—Pillared.
 Columnaris, e—Columnar.
 Comes—A companion.
 Comis, e—Friendly, nice, delicate.
 Communis, e—Common.
 Commutatus, a, um—Changed, altered.
 Comosus, a, um—Hairy.
 Compactilis, e—Pressed together.
 Compactus, a, um—Compact.
 Compertus, a, um—Discovered, ascertained.

Complanatus, a, um—Levelled, smoothed.
 Complexatus, a, um—Encircled.
 Complexus, a, um—Surrounded, encircled.
 Compressus, a, um—Compressed.
 Computus, a, um—Ornamented, elegant.
 Conatus—An effort.
 Concavus, a, um—Concave.
 Concentricus, a, um—Concentric.
 Concinnulus, a, um—Small and beautiful.
 Concinnus, a, um—Beautiful, neat.
 Conditus, a, um—Joined.
 Confectus, a, um—Completed.
 Conferus, a, um—Pressed close together.
 Confervoides—Like Conferva.
 Confirmatus, a, um—Made firm, established.
 Conflexus, a, um—Bent.
 Confluens—Running together, blended.
 Conformalis, e—Similar.
 Confragosus, a, um—Rough, uneven.
 Confragus, a, um—Rough.
 Confusus, a, um—Mixed together, confused.
 Congener, eris—Congeneric.
 Congestus, a, um—Accumulated, heaped.
 Conglobatus, a, um—Gathered in a round
 mass.
 Conglomeratus, a, um—Gathered together.
 Congregatus, a, um—Assembled together.
 Congregalis, e—Uniting together.
 Congruens—Corresponding, coinciding, run-
 ing together.
 Coniculus—A little cone.
 Conicus, a, um—Conical, cone-shaped.
 Conifollis—An inflated cone.
 Conifer, era, erum—Bearing conical fruit.
 Coniformis, e—Cone-shaped.
 Conifrons—Having a conical front.
 Coniger, era, erum—Bearing fruit of a con-
 ical form.
 Conjugans—Joined, united.
 Conjunctivus, a, um—Connecting.
 Connatus, a, um—Connate, united.
 Connivens—Dissembling, closing.
 Conoideus, a, um—Somewhat conical.
 Consimilis, e—Wholly similar.
 Consobrinus—A cousin, relative, remotely
 allied.
 Consolidatus, a, um—Consolidated.
 Consolidus, a, um—Very firm.
 Consors—Living in common.
 Conspicuus, a, um—Visible, conspicuous.
 Constans—Standing firm.
 Constellatus, a, um—Very starry.
 Constrictostriatus, a, um—Constricted and
 striated.
 Constrictus, a, um—Constricted.
 Consuetus, a, um—Customary, related to.
 Contextus, a, um—Entwined.
 Continens—Holding together.
 Contractus, a, um—Contracted.
 Conritus, a, um—Worn out.
 Conulatus, a, um—Having little cones.
 Conulus—A little cone.
 Conus—A cone.
 Convergens—Converging.
 Convexus, a, um—Convex.
 Convolutus, a, um—Rolled up, spiral-
 whorled.
 Convolvans—Rolled together.
 Coralliferus, a, um—Coral-bearing.

Corallinum—Like red coral.
 Coralloides—Like coral.
 Corbis—A basket.
 Corbula—A little basket.
 Corbulfiformis, e—Like a basket.
 Cordatoovatus, a, um—Cordate-ovate.
 Cordatus, a, um—Cordate, heart-shaped.
 Cordiformis, e—Heart-shaped.
 Coriaceus, a, um—Coriaceous, having the texture of rough skin.
 Coriformis, e—Like Coris.
 Corinthius, a, um—Corinthian.
 Corium—A leather strap, bark.
 Corniculum—A little horn.
 Corniger, era, erum—Horned.
 Cornuformis, e—In the form of a horn.
 Cornu—A horn.
 Cornulum—A little horn.
 Cornutiformis, e—Horn-shaped.
 Cornutus, a, um—Horned.
 Coronarius, a, um—Of or belonging to a wreath.
 Coronatus, a, um—Crowned.
 Corpulentus, a, um—Corpulent.
 Corrugatus, a, um—Corrugated, wrinkled.
 Corticeus, a, um—Covered with bark.
 Corticosus, a, um—Having thick bark.
 Corylus—A hazel.
 Cosciniformis, e—Like Coscinium.
 Costa—A rib.
 Costalis, e—Ribbed.
 Costatiformis, e—Rib shaped.
 Costatulus, a, um—Small ribbed.
 Costatus, a, um—Having ribs, ribbed.
 Costelliferus, a, um—Bearing faint ribs.
 Crassatus, a, um—Thickened.
 Crassibrachiatus, a, um—Thick-armed.
 Crassiscardinalis, e—Having a thick hinge.
 Crassicauda—Thick-tail.
 Crassicaulis, e—Having a thick stem.
 Crassicostatus, a, um—Thick-ribbed.
 Crassidens—Having a thick tooth.
 Crassidiscus—A thick disk.
 Crassifrons—Having a thick front.
 Crassimarginatus, a, um—Thick-margined.
 Crassinervis, e—Having thick or dense nerves.
 Crassiradiatus, a, um—Having thick rays.
 Crassitestus, a, um—Like a thick vessel or pot-lid.
 Crassolaris, e—Thickened.
 Crassus, a, um—Thick.
 Cratera—A bowl.
 Crateriformis, e—Cup shaped.
 Cratellus, a, um—Composed of lattice-work.
 Cratis—Wicker work.
 Crebescens—Frequent, increasing.
 Crebratus, a, um—Made thick, close.
 Crebripora—Having the pores very close.
 Crebrirama—Having dense branches.
 Crebrisepius, a, um—Having many septa.
 Crebristriatus, a, um—Closely striated.
 Crenatocinctus, a, um—Notched around.
 Crenatus, a, um—Crenated, notched.
 Crenistriatus, a, um—Having wrinkled lines.
 Crenulatus, a, um—Crenulated.
 Crepidula—A small sandal.
 Crepiformis, e—Boot-shaped.
 Cretaceous, a, um—Clalk-like.

Crebriformis, e—Full of openings like a sieve.
 Cribrarius, a, um—Pertaining to a sieve.
 Cribrosus, a, um—Full of holes like a sieve.
 Crineus, a, um—Hairy.
 Crispatus, a, um—Curled, crisped.
 Crispus, a, um—Curled, wavy.
 Cristatus, a, um—Tufted, crested.
 Cristula—A small crest.
 Cristulatus, a, um—Small-tufted.
 Crossotus, a, um—Fringed.
 Crotaliformis, e—Shaped like a bell.
 Crotalum—A bell, a rattle.
 Cruciatu, a, um—Cross-shaped, twisted.
 Cruciferous, a, um—Cross-bearer.
 Cruciformis, e—Cruciform.
 Cruciger, era, erum—Cross bearer.
 Crustosus, a, um—Crusted.
 Crustula—A little shell, crust.
 Cryptatus, a, um—Concealed.
 Cryptodens—Hidden tooth.
 Cucullus—A cap, covering.
 Culeus—A leather bag.
 Culmula—A little stalk or stem.
 Culmus—A stem.
 Cultellatus, a, um—Like a little knife.
 Cultellus—A small knife.
 Cultiactylus, a, um—Elegantly fingered.
 Cultratus, a, um—Knife-formed.
 Cultriformis, e—Shaped like a pruning-knife.
 Cumulatus, a, um—Heaped.
 Cumulus—A heap.
 Cuneatus, a, um—Wedge-form d.
 Cuneiformis, e—Wedge-shaped.
 Cuneolus—A little wedge.
 Cuneus—A wedge.
 Cuniculosus, a, um—Full of caves.
 Cuniculus—A cradle, cavity.
 Cunule—A little cradle.
 Curiosus, a, um—Curious.
 Curticardinalis, e—Short-hinged.
 Curtidentatus, a, um—Short-toothed.
 Curtilobus, a, um—Short-lobed.
 Curtirostratus, a, um—Short-beaked.
 Curtus, a, um—Shortened.
 Curvatus, a, um—Curved.
 Curvidens—Having curved teeth.
 Curvijuncturus, a, um—Joining in a curve.
 Curvilineatus, a, um—Having curved lines.
 Curvirostrum—A bent beak.
 Cuspidatus, a, um—Pointed.
 Cyathus—A cup.
 Cyathiformis, e—Cup-shaped.
 Cyclas—Of a round form.
 Cyclonemioides—Like a Cyclonema.
 Cyclopora—Round pore.
 Cycloptera—Circle-wing.
 Cyclopteroides—Like Cyclopterus.
 Cyclostegium—Circular covering.
 Cyclostigma—Having round scars, round-dotted.
 Cyclostomus, a, um—Having a round mouth.
 Cylindraceus, a, um—Like a cylinder.
 Cylindricus, a, m—Cylindrical.
 Cymatoides—Wave-like.
 Cymbalum—A cymbal.

of openings like a
ining to a sieve.
of holes like a

ed, crisped.
wavy.
l, crested.

all-tufted.
red.
d like a bell.
little.
s-shaped, twisted.
oss-bearer.
orm.
ross bearer.
sted.
l, crust.
cealed.
ooth.
ing.
k or stem.

ke a little knife.
ife.
Elegantly fingered.
ife-formed.
ped like a pruning-
teaped.

edge-form d.
ge-shaped.
edge.

Full of caves.
cavity.
dle.
rious.
hort-hinged.
Short-toothed.
hort-lobed.
Short-beaked.
tened.
rved.
urved teeth.
m—Joining in a curve.
—Having curved lines.
nt beak.
Pointed.

p-shaped.
form.
ke a Cyclonema.
pore.
ving.
e Cyclopteris.
lar covering.
g round scars, round-

um—Having a round

—Like a cylinder.
Cylindrical.
-like.
bal.

Cymbiformis, e—Boat-shaped.
Cymbium—A small drinking cup.
Cymbula—A small boat.
Cymosus, a, um—Full of shoots.
Cynodon—Dog-tooth.
Cyrtiniformis, e—Like Cyrtina.
Cyrtodontoides—Like Cyrtodonta.
Cyrtolites—A curved stone.
Cysticus—A little bladder.

Dactyliformis, e—Finger-shaped.
Dactylodus, a, um—Finger-toothed.
Dactyloides—Like thimble punctures.
Dactylus—Growing like a finger.
Debilis, e—Weak, feeble.
Decabrachiatas, a, um—Ten-armed.
Decadactylus, a, um—Ten-fingered.
Decemplicatus, a, um—Ten-plicated.
Decipiens—Deceiving, doubtful.
Declivis, e—Sloping.
Decoratus, a, um—Decorated.
Decornis—Without horns.
Decorosus, a, um—elegant.
Decorticatus, a, um—Barked, decorticated.
Decorus, a, um—Seemly, suitable, beautiful.
Decrescens—Decreasing, growing less.
Decurrens—Decurring, hanging down.
Decursus, a, um—Downward, running down.
Decurtatus, a, um—Curtailed.
Decussatus, a, um—Arranged in pairs that cross each other.
Defiguratus, a, um—Disfigured.
Deflectus, a, um—Deflected.
Deflexus, a, um—Bent, turned aside.
Deformatus, a, um—Deformed.
Deformis, e—Deformed, ugly-shaped.
Degener, eris—Degenerate, unlike the ancestors.
Delicatulus, a, um—Quite delicate.
Delicatus, a, um—Delicate, thin.
Delphinoccephalus—Dolphin-headed.
Deltoideus, a, um—Like the Greek letter Delta.
Deminutivus, a, um—Diminutive.
Demissus, a, um—Hanging down.
Demum—At last, solely.
Denarius, a, um—Containing ten.
Densifolius, a, um—Dense-leaved.
Densmamillatus, a, um—Having mammillated teeth.
Densus, a, um—Dense, thick.
Dentalium—A plow-share.
Dentatus, a, um—Toothed.
Denticulatus, a, um—Denticulated, having small teeth.
Dentilineatus, a, um—Tooth-lined.
Denudatus, a, um—Denuded.
Depareus, a, um—Very scarce.
Depauperatus, a, um—Impoverished.
Deperditus, a, um—Impoverished.
Depressus, a, um—Depressed.
Desertus, a, um—Deserted, forsaken.
Desideratus, a, um—Desired, rare.
Desmopieura—A side band.
Desquamatus, a, um—Scaled off.
Devexus, a, um—Sloping.
Diadematus, a, um—Wearing a diadem.

Dialophus—Through the neck.
Dianthus, a, um—Double-flowered.
Diatretus, a, um—Pierced with holes.
Dichotomus, a, um—Divided.
Dictyopteroides—Like Dictyopteris.
Dictyota—Net-worked.
Dictyum—A net.
Difficilis, e—Difficult, rough.
Diffidens—Diffident, distrusting.
Diffuens—Flowing every way, loose.
Diffusus, a, um—Diffused, extended.
Digitalis, e—Belonging to the finger.
Digitatus, a, um—That has fingers, toes, or claws.
Dignatus, a, um—Excellent.
Digonus, a, um—Two-angled.
Dikrocheilus, a, um—Two-edged.
Dilatatus, a, um—Dilated, widened.
Dilatus, a, um—Spread out.
Diluculum—Day-break, dawning of day.
Diminutivus, a, um—Diminutive.
Dimorphus, a, um—Two-formed.
Diplostegioides—Like Diplostegium.
Diplostesta—Having two tests.
Disciformis, e—Shaped like a quoit.
Discoidalis, e—Discoidal.
Discoideus, a, um—Discoid, disk-like.
Discophorus—Disk-bearer.
Discrepans—Different.
Disculus—A little disk.
Discus—A quoit.
Disjunctus, a, um—Separated, disjoined.
Dispalans—Straggling, stray.
Dispandus, a, um—Spread out, stretched.
Dispanus, a, um—Stretched out.
Dispar—Different.
Disparilis, e—Different, unequal.
Dispassus, a, um—Extended, spread out.
Dispersus, a, um—Dispersed.
Dissectus, a, um—Cut up, dissected.
Dissimularis, e—Dissimilar, unlike.
Dissolutus, a, um—Weak, broken.
Distans—Distant, standing apart.
Distensus, a, um—Distended.
Distinctus, a, um—Distinct.
Distortus, a, um—Distorted, crooked, irregular.
Divaricans—Severed, straddling.
Divaricatus, a, um—Divaricated, wide apart.
Divergens—Diverging.
Diversifolius, a, um—Diverse-leaved.
Diversus, a, um—Diverse, different, unlike.
Divisus, a, um—Dividing.
Docens—A teacher.
Dodecadactylus, a, um—Twelve-fingered.
Dolabriformis, e—Like a mattock or pick-axe.
Dolatus, a, um—Hewed.
Dolorosus, a, um—Wretched.
Donaciformis, e—Like a Donax.
Dorsalis, e—Dorsal.
Dorsatus, a, um—High-backed.
Dotis—An ornament.
Drepanaspis—Having a sickle-shield.
Dubius, a, um—Doubtful.
Dumalis, e—Bushy.
Dumosus, a, um—Bushy.
Duodenarius, a, um—Containing twelve.
Duplicatus, a, um—Duplicated, doubled.
Daplicostatus, a, um—Double-ribbed.

Eboreus, a, um—Made of ivory.
Ebracteatus, a, um—Without scales or bracts.
Eburneolus, a, um—Of ivory.
Eccentricus, a, um—From the center.
Echinatus, a, um—Set with spines.
Ectypus, a, um—Engraved in relief, embossed.
Edax—Voracious.
Edentulus, a, um—Toothless.
Egenus, a, um—Destitute of, very poor.
Elegans—Elegant, handsome.
Elegantissimus, a, um—Very handsome.
Elegantulus, a, um—Quite elegant.
Elevatus, a, um—Elevated.
Ellipticus, a, um—Elliptical.
Elongatus, a, um—Elongated.
Elytra—The wing covering.
Elytroides—Like the elytra of beetles.
Emaceratus, a, um—Thin.
Emaciatulus, a, um—Emaciated, thin.
Emarginatus, a, um—Notched.
Eminens—Prominent, standing out in relief.
Eminulus, a, um—Projecting a little.
Enormis, e—Very large.
Ensiformis, e—Sword-formed.
Eos—The dawn.
Epidermatulus, a, um—Covered with a crust or skin.
Equilaterus, a, um—Equal-sided.
Equisetiformis, e—Like Equisetum.
Erectifolius, a, um—Having leaves erect.
Erectipora—Having erect pores.
Erectus, a, um—Erect, straight.
Erodus, a, um—Eroded, jagged, gnawed.
Erosus, a, um—Eroded, bitten away.
Erraticus, a, um—Wandering, erratic.
Erythroliticus—Red stone.
Escharoides—Like Eschara.
Eucharis, e—Graceful, beautiful.
Euconnus—Perfect cone.
Euginum—Fertile.
Euglypheus, a, um—Well-carved, distinctly marked.
Euomphaloides—Like Euomphalus.
Euphemia—Of good omen.
Euruteines—Extending widely.
Euzona—Beautifully girdled.
Evax—An exclamation of delight.
Exacutus, a, um—Pointed.
Exanthematus, a, um—Covered with eruptions.
Excavatus, a, um—Made hollow, excavated.
Excellens—Excellent, high-rising.
Excelsior—Elevated, lofty.
Excelsus, a, um—Elevated, high.
Excerptus, a, um—Selected, picked out.
Excrescens—Growing out, increasing.
Exculptus, a, um—Adorned, chiseled out.
Exfoliatus, a, um—Exfoliated.
Exiguus, a, um—Small, petty.
Exilis, e—Thin, lean, slender, creeping.
Eximius, a, um—Choice, select, excellent.
Exornatus, a, um—Adorned.
Exortivus, a, um—Eastern.
Expansus, a, um—Expanded, widely spread.
Expatius, a, um—Spread out.
Explanatus, a, um—Made plain, spread out.
Explicatus, a, um—Unfolded, spread out.

Explorator—A scout, an examiner.
Exporrectus, a, um—Smooth, stretched out.
Exsculptus, a, um—Carved.
Exsertus, a, um—Projecting, thrust forth.
Exsul—A wanderer.
Extas—Standing out.
Extensus, a, um—Stretched out, extended.
Extenuatus, a, um—Made thin, slender, drawn out.
Extumidus, a, um—Swelled up.
Exutus, a, um—Divested, stripped off.
Faba—A bean.
Fabalis, e—Bean-stalks.
Fabula—A little bean.
Fabulites—A little stone-bean.
Facetus, a, um—Elegant.
Falcatus, a, um—Hooked.
Falciformis, e—Like a sickle, pruning-knife, or hook.
Fallax—Deceptive.
Falx—A hook, pruning-knife, or sickle.
Famelicus, a, um—Famished.
Fasciatus, a, um—Banded.
Fascicularis, e—Small-bundled.
Fasciculatus, a, um—Bundled.
Fasciculus—A bundle.
Fascigerus, a, um—Bearing fascies.
Fastigatus, a, um—Sloping up to a point.
Faustus, a, um—Fortunate, lucky.
Favositoides, a, um—Like Favosites.
Favosus, a, um—Honeycomb-like.
Fax—A torch, taper.
Fecundus, a, um—Fruitful.
Felix, icis—Fertile.
Fenestella—A little window.
Fenestelliformis, e—Like Fenestella.
Fenestratus, a, um—Reticulated, having open windows.
Fenestrula—A little window.
Ferox—Fierce, hardy, stout.
Ferratus, a, um—Hard as iron, covered with iron.
Ferriculus—Iron distaff.
Ferrugineus, a, um—Of the color of iron, rusty.
Fertilis, e—Fertile, fruitful.
Ferus, a, um—Wild, cruel, fierce.
Festinatus, a, um—Hastened, before the time.
Fibratus, a, um—Having small threads hanging to it.
Fibristriatus, a, um—Fiber-lined.
Fibrosus, a, um—Full of fibers or threads.
Ficoides—Like a fig.
Ficus—A fig.
Fidelis, e—Sure, faithful.
Filiciformis, e—Fern-like.
Filicosta—Having thread-like costa.
Filicula—Fern of trees, wall-fern.
Filiculme—Thread-straw.
Filiformis, e—Filiform.
Filistriatus, a, um—Having thread-like striae.
Filitextilis, e—Woven like thread.
Filitextus, a, um—Woven like thread.
Filosus, a, um—Thready.
Fimbriatus, a, um—Fringed, jagged, scalloped.

examiner.
 outh, stretched out.
 d.
 ing, thrust forth.
 ed out, extended.
 le thin, slender,
 ed up.
 stripped off.
 bean.
 l.
 kle, pruning-knife,
 knife, or sickle.
 shed.
 d.
 ndled.
 ndled.
 ing fascies.
 ng up to a point.
 ate, lucky.
 like Favosites.
 comb-like.
 ful.
 dow.
 e Fene-stella.
 Reticulated, having
 dow.
 tout.
 is iron, covered with
 the color of iron,
 ful.
 el, fierce.
 astened, before the
 wing small threads
 her-lined.
 of fibers or threads.
 l.
 ke.
 ad-like costa.
 wall-fern.
 w.
 Having thread-like
 like thread.
 ven like thread.
 y.
 ringed, jagged, scal-

Fiscellostriatus, a, um—Having divided
 strise.
 Fiscellus—A small basket woven of slender
 twigs.
 Fissicosta—Having divided costæ.
 Fissilis, e—Split.
 Fissiplica—Having divided plications.
 Fissuratus, a, um—Fissured.
 Fissurellus, a, um—Having a little cleft.
 Fissus, a, nm—Divided, cleft, split.
 Fistulosus, a, um—Full of holes, spongy.
 Flabellatus, a, um—Fan-like.
 Flabellifer, era, erum—That bears a fan.
 Flabelliformis, e—Shaped like a fan.
 Flabellites—A stone fan.
 Flabellum—A fan.
 Flaccidus, a, um—Withered, hanging, flag-
 ging, flaccid.
 Flagellaria, e—Like a whip.
 Flagellum—A whip.
 Flagricaudus, a, um—Whip-tailed.
 Flavus, a, um—Golden, yellow.
 Flexicaulis, e—Having a flexible stem.
 Flexifolius, a, um—Having recurved leaves.
 Flexilis, e—Pliant, flexible.
 Flexuosus, a, um—Flexuous, full of turns.
 Florealis, e—Flower-like.
 Floridus, a, um—Flowery, adorned with
 flowers, gay.
 Florifer, era, erum—Flower-bearing.
 Floriformis, e—Flower-shaped.
 Flos—A flower.
 Fluctus—A wave, a billow.
 Fluctuosus, a, um—Full of waves, wavy,
 veiny.
 Fluitans—Flowing, floating.
 Fecundus, a, um—Fruitful, abundant.
 Feculoideus, a, um—Like a tumor.
 Foliaceus, a, um—Foliateous, like leaves,
 leafy.
 Foliatius, a, um—Leaved, having leaves.
 Foliosus, a, um—Leafy, full of leaves.
 Folium—A leaf.
 Folliculus—A small sack.
 Follis—A leather sack.
 Fonticola—Fountain-dwelling.
 Fontinalis—A fountain or spring.
 Formosus, a, um—Beautiful, handsome.
 Fornacula—A little oven.
 Fornax—A furnace.
 Fornicatus, a, um—Arched, vaulted over.
 Forulatus, a, um—Having narrow furrows.
 Fossatus, a, um—Dug out.
 Fossilis, e—That may be dug out of the
 earth, fossil.
 Fossula—A little trench or ditch.
 Foveatus, a, um—Pitted.
 Fractus, a, um—Broken, effeminat.
 Fragarioides—Like a strawberry.
 Fragilis, e—Brittle, frail.
 Fragosus, a, um—Fragile.
 Frangens—A breaker.
 Fraternalis, a, um—Brotherly, fraternal.
 Fraxiniformis, e—Like fraxinus.
 Frequentatus, a, um—Frequent.
 Frinilla—A small bird.
 Fritillus—A dice-box.
 Frondosus, a, um—Full of leaves.
 Frutex—A shrub.

Fruticosus, a, um—Shrubby, full of shoots.
 Fucoides—Like Fucus.
 Fulcratus, a, u. a—Stayed with props.
 Fulgidus, a, um—Shining.
 Fulgur—A thunder-bolt.
 Funatus, a, um—Corded.
 Fungosus, a, um—Spongy.
 Fungulus—A small mushroom.
 Funiculus—A small cord or line.
 Furcatus, a, um—Forked.
 Furcicaratus, a, um—Forked and keeled.
 Furtivus, a, um—Secret, hard to find.
 Fusibrachiatus, a, um—Having fusiform
 arms.
 Fusiformis, e—Fusiform, tapering at both
 ends.
 Fustiformis, e—Club-formed.
 Fustia—A club, staff.
 Futilis, e—Trivial.

Galeatus, a, um—That wears a helmet.
 Galericalatus, a, um—Having a small cov-
 ering.
 Galerum—A cap, hat, or tuft of feathers.
 Gallinuloides—Like a pullet.
 Gemellipara—Twin-bearing.
 Geminispinosus, a, um—Twin-spined.
 Gemma—A young bud, a gem.
 Gemmatus, a, um—Budded, set with gems.
 Gemmicula—A little bud.
 Gemmifer, era, erum—That bears buds or
 gemmules.
 Gemmiformis, e—Shaped like a bud.
 Gemmula—A little bud.
 Geniculatus, a, um—Knotted, jointed.
 Geniculosus, a, um—Knotty.
 Genitivus, a, um—Natural, belonging to the
 same stock.
 Geometricus, a, um—Geometrical.
 Germanus, a, um—Near of kin.
 Gibber, era, erum—Bossed, hunchbacked.
 Gibberosus, a, um—Badly hunchbacked.
 Gibberulus, a, um—Somewhat hunch-
 backed.
 Gibberus, a, um—Humpbacked.
 Gibbosus, a, um—Gibbous, full of hunches,
 or humped.
 Gibbus, a, um—Hunched, gibbous.
 Giganteus, a, um—Giant-like, very large.
 Gigas—A giant.
 Glabellus, a, um—Smooth.
 Glaber, bra, brum—Smooth, bare.
 Glacialis, e—Frozen, icy.
 Gladiolus—A small sword.
 Glandulosus, a, um—Full of kernels, gland-
 ular.
 Glandulus, a, um—Having kernels, gland-
 ular.
 Glans—An acorn, chestnut, or pellet.
 Glanscerasi—Fruit of the cherry-tree.
 Glansfagea—Fruit of the beech-tree.
 Globatus, a, um—Made round.
 Globosus, a, um—Round as a ball, globose.
 Globularis, e—Globular.
 Globuliformis, e—Globe-shaped.
 Globulus—A little ball.
 Glomeratus, a, um—Confused, out of order.
 Gloriosus, a, um—Glorious.
 Glyptus, a, um—Sculptured.

Gomphoidea—Like a stake or club.
 Gomphus—A pile, stake, or club.
 Goniocercus, a, um—Angular, tailed.
 Goniolobus, a, um—Having angular lobes.
 Goniopteroides—Like Goniopterus.
 Goniurus, a, um—Angular-tailed.
 Gonopleura—Angular rib.
 Gothicus, a, um—Gothic.
 Gracilentus, a, um—Slender, thin.
 Gracilis, e—Small, slender, thin, weak.
 Gracilius, a, um—More slender.
 Gracillimus, a, um—Very slender, thin, or weak.
 Gracillistriatus, a, um—slender, striated.
 Gradatus, a, um—Made with steps.
 Gradiocosta—Having steps and ribs.
 Gradiocostatus, a, um—Having steps and ribs.
 Gramineus, a, um—Grassy or belonging to grass.
 Grandævus, a, um—Very old.
 Grandiceps—Big-headed.
 Grandiculus, a, um—Rather large.
 Grandifolius, a, um—Large-leaved.
 Grandis, e—Grand, large.
 Graniferus, era, erum—That bears grains of corn.
 Granilineatus, a, um—Lined with granules.
 Granilineus, a, um—Granule-lined.
 Granosus, a, um—Full of grains or kernels.
 Granulatus, a, um—Granulated, granular.
 Granuliferus, era, erum—Granule-bearing.
 Granulostriatus, a, um—Having granular striae.
 Granulosus, a, um—Covered with small granules.
 Graphicus, a, um—Perfect, excellent, done to the life, written on.
 Gratosus, a, um—Agreeable.
 Gratus, a, um—Acceptable.
 Gravis, e—Weighty, full, old.
 Graviusculus, a, um—Rather deep.
 Gregalis, e—Of the common sort.
 Gregarius, a, um—Of the common sort, common, gregarious.
 Grossiplicatus, a, um—Thick-plaited.
 Grumus—A little heap.
 Gypseus, a, um—Covered or plastered with gypsum.
 Gyraacanthus—Round spine.
 Gyrinoides—Like a tadpole.
 Gyrocera—Circular horn.

Hæsitans—Doubting.
 Hallioides—Like Hallotus.
 Hamatilis, e—Furnished with hooks.
 Hamatus, a, um—Crooked, hooked.
 Hamulosus, a, um—Full of hooks.
 Hamulus—A small hook.
 Harpago—A hook.
 Hastatus, a, um—Bearing spears, halbert-shaped.
 Hastifolius, a, um—Spear or lance-leaved.
 Hastula—A little spear.
 Helicoides—Like a helix.
 Helicteres—A round, smooth spire.
 Heliolitiformis, e—Like Heliolites.
 Helios—The sun.

Hemicyclus—A half-circle.
 Hemicyllindrus—A half-cylinder.
 Hemiplica, us, a, um—Half-plaited.
 Hemisphericus, a, um—Hemispherical.
 Hemiteloides—Like Hemiteles.
 Hemitrypa—Having half-openings.
 Herbaceus, a, um—Grassy.
 Herculeus, a, um—Belonging to Hercules, large of its kind.
 Heterocinctus, a, um—Irregularly girded or banded.
 Heteroclitus, a, um—Extraordinary.
 Heterodactylus, a, um—Irregular-toed or irregularly fingered.
 Heterophyllus, a, um—Irregularly or differently leaved.
 Heteropora—Having irregular pores.
 Heteropteris—Irregular fern.
 Hexadactylus, a, um—Six-fingered.
 Hexagonus, a, um—Having six angles.
 Hexagonalis, e—Hexagonal.
 Hians—Gaping, disjointed.
 Hipparionyx—A colt's hoof.
 Hirsutus, a, um—Rough, hairy, shaggy.
 Hirtus, a, um—Rough, hairy, shaggy.
 Hispidus, e, um—Rough, bristly, rugged.
 Hylucis, a, um—Gaping, cleft.
 Holopyga—Entire rump, whole back.
 Homalonotoidea—Like Homalonotus.
 Horizontalis, e—Horizontal.
 Horridus, a, um—Rough, bristly.
 Hospitalis, e—Of a guest, hospitable.
 Humerosus, a, um—Humped, full of humps.
 Humerulus—A little shoulder.
 Humilis, e—Small, poor.
 Hyalina—Of glass.
 Hybrida—Intermediate between two species, a hybrid.
 Hydraulicus, a, um—Hydraulic.
 Hymenophylloides—Like Hymenophyllites.
 Hyperbolæus, a, um—Extreme.
 Hyperboreus, a, um—Very far north.
 Hypniformis, e—Like Hypnum.
 Hystricosus, a, um—Thorny.
 Hystriculus, a, um—Somewhat covered with spines.
 Hystrix—Covered or beset with spines.

Ichthyoderma—Having a fish-skin.
 Ichthyolepis—Having fish-scales.
 Icosidactylus, a, um—Twenty-fingered.
 Idoneus, a, um—Suitable.
 Ignobilis, e—Ignoble, strange, unknown.
 Ignorabilis, e—Overlooked, unknown.
 Ignotus, a, um—Unknown, strange.
 Illicifolius, a, um—Oak-leaved.
 Illænoides—Like an Illænus.
 Illibatus, a, um—Unimpaired.
 Imago—An image, picture, also a sheath.
 Imbecillus, a, um—Feeble, frail.
 Imbricarius, a, um—Having imbrications.
 Imbricato-articulatus, a, um—Having imbricated articulations.
 Imbricatus, a, um—Laid one on another like tiles, imbricated.
 Imitator—A resembler.
 Immaturus, a, um—Immature, abortive.
 Immersus, a, um—Immersed.

le.
cylinder.
self-plaited.
Hemispherical.
sittels.
openings.
y.
onging to Hercules,
regularly girded or
traordinary.
irregular-toed or ir-
regularly or diff-
regular pores.
fern.
x-fingered.
ing six angls.
nal.
ed.
roof.
y, hairy, shaggy.
y, hairy, shaggy.
y, bristly, rugged.
cleft.
whole back.
Homalonotus.
atal.
y, bristly.
t, hospitable.
aped, full of humps.
oulder.
between two spe-
hydraulic.
ke Hymenophyl-
xtreme.
very far north.
ypnum.
ony.
ewhat covered with
set with spines.
g a fish-skin.
sh-scales.
twenty-fingered.
le.
trange, unknown.
ed, unknown.
wn, strange.
eaved.
gnus.
aired.
are, also a sheath.
ple, frail.
iving imbrications.
um—Having im-
d one on another
ature, abortive.
ersed.

Impar—Odd, unequal, disproportioned.
Imparilis, e—Different.
Imperator—A commander.
Implexus, a, um—Interlaced, interwoven.
Implicatus, a, um—Wrapped together, en-
tangled.
Impolitus, a, um—Rough, unpolished.
Impositus, a, um—Laying over.
Impressus, a, um—Impressed.
Improcerus, a, um—Undersized, not tall.
Impudicus, a, um—Shameless, immodest.
Inaequalis, e—Uneven, unequal.
Inequalis, e—Unequal.
Inequatus, a, um—Unequal.
Inaequicostatus, a, um—Unequally ribbed.
Inaequidactylus, a, um—Unequal-fingered.
Inaequilateralis, e—Inequilateral.
Inaequuplicatus, a, um—Unequally rayed.
Inaequistriatus, a, um—Unequally stri-
ated.
Inceptus, a, um—An undertaking, in-
cipient.
Incertus, a, um—Uncertain, inconstant.
Incills, e—Belonging to, or like a trench,
furrow, or gutter.
Incipiens—The beginning.
Incisivus, a, um—Having the quality of
cutting or biting.
Inciso-lobatus, a, um—Cut into lobes.
Incisus, a, um—Incised.
Inclinatus, a, um—Inclined, bent.
Inclinis, e—Bending.
Incluspora—Having inclosed perforations.
Inclusus, a, um—Closed up.
Incompletus, a, um—Incomplete.
Incomptus, a, um—Untrimmed, rough.
Inconditus, a, um—Irregular, disordered.
Inconspicuus, a, um—Not conspicuous.
Inconstans—Not constant.
Inconsuetus, a, um—Unusual.
Incrassatus, a, um—Thickened.
Increbescens—Abundant.
Incrustans—Incrusting.
Incultus, a, um—Neglected.
Incurvus, a, um—Incurved.
Indagator—A diligent hunter.
Indagatus—Encircling.
Indentatus, a, um—Indented, notched.
Indentus, a, um—Indented, notched.
Indeterminatus, a, um—Not determined.
Indolatus, a, um—Unhewn.
Inelegans—Unadorned.
Inermis, e—Unarmed.
Inexpectans—Not expected.
Infelix—Useless, unhappy, miserable.
Infernus—Underground, the lower.
Infertus, a, um—Filled up.
Inferus—Below, underground.
Inflatus, a, um—Spread, swollen, inflated.
Inflexus, a, um—Bowed, made crooked.
Informis, e—Shapeless, rude.
Infrequens—Rare, infrequent.
Infula—A band, an ornament.
Infundibularius, a, um—Pertaining to a
funnel.
Infundibuliformis, e—Funnel-shaped.
Infundibulum, e—A funnel, hopper.
Ingens—Very large, huge, prodigious.
Ingentior—Larger, enormous.

Inopinatus, a, um—Unexpected, un-
thought of.
Inops—Poor, friendless, unburied.
Inoptatus, a, um—Undesired, not wanted.
Inordinatus, a, um—Disordered.
Inornatus, a, um—Unadorned.
Insculptus, a, um—Engraven, carved.
Insectus, a, um—Uncut.
Insertivus, a, um—Inserted.
Insignis, e—Marked, naturally remark-
able.
Insignitus, a, um—Marked, clear.
Insitna, a, um—Inserted, introduced.
Insculptus, a, um—Engraved.
Insolens—Unusual, rare.
Insolitus, a, um—Rare, hard to find.
Insous—Harmless.
Inspeciosus, a, um—Not handsome.
Insuperatus, a, um—Unexp cted.
Instabilis, e—Not firm, changing.
Insuetus, a, um—Unusual.
Insularis, e—Upon an island.
Intectus, a, um—Uncovered.
Integrifolius, a, um—Whole-leaved.
Intercalaris, e—Intercalated.
Intercedens—Intervening.
Intercellatus, a, um—Being intercellular.
Intercostalis, e—Lined between costae.
Intercostatus, a, um—Ridged between
ribs.
Interlineatus, a, um—Interlined.
Intermedius, a, um—Intermediate, the
middle.
Intermittens—Intermitting, ceasing for a
time.
Internascens—Growing between.
Internodius, a, um—Space between two
knots or joints.
Interplicatus, a, um—Plicated between.
Interruptus, a, um—Broken asunder, inter-
rupted.
Intersculptus, a, um—Engraved in the
middle.
Interscapularis, e—Spaced between the
shoulder pieces.
Int-rstinctus, a, um—Divided.
Interstrialis, e—Having striae between.
Interstrictus, a, um—Drawn together.
Intertextus, a, um—Interwoven, interlaced.
Intervescicula—Having little vesicles be-
tween.
Intextus, a, um—Plaited, woven.
Intortus, a, um—Twirled, entangled, curled.
Intralineatus, a, um—Lined, between lines.
Inutilis, e—Not useful, very poor, insign-
ificant.
Invaginatus, a, um—Invaginated, sheathed,
enwrapped.
Invalidus, a, um—Weak, feeble.
Invenustus, a, um—Unhandsome.
Inversus, a, um—Inverted.
Investis, e—Unclothed.
Involutus, a, um—Involute.
Irrasus, a, um—Unpolished, not smooth.
Irregularis, e—Irregular.
Islandicus, a, um—From an island.
Ischypus, a, um—Strong-footed.
Isosceles—Having equal legs.
Isogramma—Equal weight.

Jaculum—A dart, javelin.
Jejunus, a, um—That has not eaten, hungry.
Jubatus, a, um—Crested.
Jucundus, a, um—Pleasant, agreeable, delightful.
Jugalis, e—Yoked together.
Juglans—A walnut.
Jugosus, a, um—Full of ridges, mountainous.
Junceus, a, um—Made of bulrushes, like a bulrush.
Junciformis, e—Shaped like a bulrush.
Junctus, a, um—Joined, coupled.
Juvenis, e—Young.

Labecula—A little spot.
Labialis, a, um—Lipped.
Labiosus, a, um—Full lipped.
Labrosus, a, um—Having large lips, bordered.
Labyrinthicus, a, um—Labyrinthine.
Laceratus, a, um—Torn, mangled, ragged.
Lachrymosus, a, um—Full of tears.
Laciniatus, a, um—Fringed.
Laciniosus, a, um—Full of plaits, jagged, crumpled.
Lactuca—Lettuce.
Lacunosus, a, um—Full of holes, pitted, uneven.
Lacus—A vat, a basin.
Lacustris, e—Pertaining to a lake or swamp.
Lætus, a, um—Fertile, pleasant, agreeable.
Lævicosta—Having a smooth rib.
Lævicostatus, a, um—Smooth-ribbed.
Læviculus, a, um—Nearly smooth.
Lævigatus, a, um—Planed, made smooth.
Lævis, e—Smooth.
Lævisimus, a, um—Very smooth.
Lævistriatus, a, um—Having smooth striæ.
Læviusculus, a, um—Quite smooth.
Lægena—A flask.
Laguncula—A little flask.
Lamellatus, a, um—Having thin plates.
Lamellosus, a, um—In very thin plates.
Laminatus, a, um—Laminated.
Lamnoides—Like Lamna.
Lanatus, a, um—Woolly.
Lanceolatus, a, um—Spear-shaped.
Lanceifer, era, erum—Lance-bearer.
Lanceifolius, a, um—Lance-leaved.
Lanosus, a, um—Woolly.
Lapicida—A stone-cutter.
Lapideus, a, um—Consisting of stone.
Lapillus—A little stone, a pebble.
Laqueatus, a, um—Arched, vaulted, fluted, paneled.
Largissimus, a, um—Very large, the largest.
Largus, a, um—Plentiful, large.
Laricinus, a, um—Resembling the larch-tree.
Larvatus, a, um—Frightened, masked.
Latealatus, a, um—Broad-winged.
Lateralis, e—Belonging to the side.
Lateralis, a, um—Of or belonging to the sides.
Latericrescens—Side-growing.
Laterniformis, e—Shaped like a lantern.
Latiannulatus, a, um—Having wide annulations.

Latibrachiatus, a, um—Wide-armed.
Latibuccatus, a, um—Wide-cheeked.
Laticeps—Broad head.
Latocosta—Having wide ribs.
Laticostatus, a, um—Wide-ribbed.
Latidactylus—Wide-fingered.
Latidorsatus, a, um—Wide-backed.
Latifasciatus, a, um—Wide-bundled, or wide-banded.
Latifolius, a, um—Broad-leaved.
Latifrons—Having a wide front.
Latijunctus, a, um—Wide-jointed.
Latimarginatus, a, um—Broad-margined.
Latior—Wider.
Latipes—Broad-footed.
Latiradius, a, um—Wide-rayed.
Latispinosus, a, um—Wide-spined.
Latissimus, a, um—Very wide, the widest.
Latitruncatus, a, um—Broadly truncated.
Lativentrus, a, um—Having a wide cavity.
Latus, a, um—Broad, wide, large.
Latusculum—A little side.
Lautus, a, um—Neat, elegant, splendid.
Laxatus, a, um—Made wider, extended, dilated.
Laxus, a, um—Loose, slack, spacious.
Ledoides—Like Leda.
Lens—A lentil.
Lenticularis, e—Lens-shaped, lenticular.
Lentiformis, e—Lens-shaped.
Lentus, a, um—Flexible, pliant, sluggish.
Leperditioides—Like Leperditia.
Lepidodendrifolius, a, um—Having leaves like *Lepidodendron*.
Lepidorachus, e—Having a scaly ridge.
Lepidus, a, um—Pretty.
Lepis—A scale.
Leptænoides—Like *Leptæna*.
Leptocephalus, a, um—Slender-headed.
Leptodactylus, a, um—Slender-toed.
Leptoderma—A thin skin.
Leptogaster—A smooth belly.
Leptonotus, a, um—Slender-backed.
Levatus, a, um—Lifted up.
Læviculus, a, um—Very small.
Levigatus, a, um—Smooth.
Levinodatus, a, um—Having smooth knots.
Levis, e—smooth.
Lichenoides—Like lichen.
Lichenoidens, a, um—Like a lichen.
Ligoniformis, e—Like a mattock.
Liliiformis, e—Shaped like a lily.
Lima—A file.
Limabrachiatus, a, um—File-armed.
Limatulus, a, um—Neat, fine, polished, like a little file.
Limatus, a, um—Polished, neat, elegant.
Limax—A snail, slug.
Limbatus, a, um—Bordered.
Limiformis, e—Lima-shaped.
Limitaris, e—Bounded, limited.
Limulus—Limulus, tail.
Lineanodus, a, um—Having lined knots.
Linearifolius, a, um—Having linear leaves.
Linearis, e—Pertaining to a line, linear.
Linearis, a, um—Belonging to lines.
Lineatoides—Like *lineatus*, a specific name.
Lineatus, a, um—Drawn out, lined.
Lineolatus, a, um—Fine-lined.

de-armed.
 e-checked.
 e.
 e-ribbed.
 e.
 e-backed.
 e-bundled, or wide.
 eaved.
 e-front.
 e-jointed.
 e-road-margined.
 eayed.
 e-spined.
 e-wide, the widest.
 e-ly truncated.
 e-ng a wide cavity.
 e, large.
 e-ant, splendid.
 e-der, extended, di-
 e-k, spacious.
 e-ped, lenticular.
 e-ped.
 e-pliant, sluggish.
 e-reditia.
 e-n—Having leaves
 e a scaly ridge.
 e-na.
 e-ender-headed.
 e-ender-toed.
 e.
 e-elly.
 e-er-backed.
 e.
 e-mall.
 e-n.
 e-ling smooth knots.
 e.
 e-ke a lichen.
 e-natlock.
 e-ke a lily.
 e-File-armed.
 e-fine, polished, like
 e, neat, elegant.
 e.
 e-ped.
 e-ped.
 e-imited.
 e-ling lined knots.
 e-ving linear leaves.
 e-o a line, linear.
 e-ving to lines.
 e-s, a specific name.
 e-out, lined.
 e-lined.

Lineopora—Having line-pores, lined with perforations.
 Lineopunctatus, a, um—Line-punctured or line-dotted.
 Lingualis, e—Tongue-shaped.
 Lingulifer, era, erum—Tongue-bearing.
 Linguiformis, e—Tongue-shaped.
 Lingulatus, a, um—Tongue-shaped, linguulate.
 Lintheum—A napkin, girdle.
 Lioderma—A smooth skin.
 Liosoma—A smooth body.
 Liratus, a, um—Furrowed.
 Lithofactor—Stone-maker.
 Litoreus, a, um—On the shore or sea-side.
 Lobatus, a, um—Lobed.
 Locellus—A little purse or bag.
 Loculosus, a, um—Full of holes or distinct places, partitioned.
 Lonchitis—Spleenwort, the fern "Adders-tongue."
 Longevus, a, um—Ancient, aged.
 Longicameratus, a, um—Long chambered.
 Longicaudatus, a, um—Long-tailed.
 Longicollis—Long-ridged.
 Longicostalis, e—Long-ribbed.
 Longidactylus, a, um—Long-fingered.
 Longidentatus, a, um—Long-toothed.
 Longifolius, a, um—Long-leaved.
 Longipes—Long-footed.
 Longirostris—Having a long proboscis.
 Longispinus, a, um—Long-spined.
 Longispira—Having a long spire.
 Longissimus, a, um—Very long, the longest.
 Longiusculus, a, um—Rather long.
 Longulus, a, um—Rather long.
 Longus, a, um—Long.
 Loriformis, e—Like a thong or whip.
 Lotoblastus—Lotus bud.
 Lucifugus, a, um—Light-shunning.
 Lunatus, a, um—Made like a half-moon, horned.
 Lunulatus, a, um—Crescentiform.
 Luxus, a, um—Dislocated.
 Lycoperdon—Puff-ball shaped.
 Lynx—An animal called a lynx.
 Lyra—A harp.
 Lyratfolius, a, um—Having lyre shaped leaves.

Macer, era, crum—Lean, meager.
 Machæiformis, e—Sword-shaped.
 Macilentus, a, um—Meager, thin, lean.
 Microcephalus, a, um—Long-headed.
 Macrochirus, a, um—Long handed.
 Macroductylus, a, um—Long-fingered.
 Macrodentus, a, um—Long-toothed.
 Macrolepidotus, a, um—Having long scales.
 Macrolineatus, a, um—Long-lined.
 Macromphalus, a, um—Having a large umbilicus.
 Macronotus, a, um—Long known.
 Macropetalus, a, um—Having long flower leaves.
 Macrophorus, a, um—Long-bearing.
 Macrophyllus, a, um—Long-leaved.
 Macropleura—Having long sides.
 Macropora—Having long pores.
 Macrops—Having large eyes.

Macropterus, a, um—Long-winged, or large-finned.
 Macrophthalmus, a, um—Long-eyed.
 Macrospira—Having a long spire.
 Macrospondylus, a, um—Having long vertebrae.
 Macrostomus, a, um—Having a long mouth.
 Macrostriatus, a, um—Having long striae.
 Macrostylus, a, um—Having long spines or columns.
 Macrothyris—Having a long foramen.
 Macrurus, a, um—Long-tailed.
 Mactrifornis, e—Shaped like Mactra.
 Mactroides—Like Mactra.
 Maculatus, a, um—Spotted, speckled.
 Maculosus, a, um—Full of spots, spotted.
 Magister—A chief, master.
 Magnicornis, e—Large-horned.
 Magnicostatus, a, um—Large-ribbed.
 Magnificus, a, um—Magnificent, stately.
 Magnifolius, a, um—Large-leaved.
 Magnisulcatus, a, um—Deep-furrowed.
 Magniventrus, a, um—Large-bellied.
 Magnoliformis, e—Shaped like magnolia.
 Magnus, a, um—Great, large.
 Major—Greater, larger.
 Majus, a, um—Greater, larger.
 Malvaceus, a, um—Like or pertaining to mallows.
 Mamillanus, a, um—Swelling, protuberant.
 Mammatus, a, um—Covered with protuberances.
 Mammiferus, a, um—Teat or nipple bearing.
 Mammillaris, e—Mammillated.
 Mammillatus, a, um—Covered with nipples.
 Maniformis, e—Hand-like.
 Mantleula—A little wallet.
 Manus—A hand.
 Marcidus, a, um—Hanging, flagging, withered.
 Marginalis, e—Marginal.
 Marginatus, a, um—That has a border, broad rim, or margin.
 Marginicinctus, a, um—Having a banded margin.
 Marinus, a, um—Inhabiting the sea.
 Maritimus, a, um—Of or belonging to the sea.
 Masculus, a, um—Stout, hardy, masculine.
 Materiarius, a, um—Of or belonging to timber.
 Maturus, a, um—Ripe, mature.
 Matutinus, a, um—In the morning.
 Maximus, a, um—Greatest, largest.
 Medialis, e—Middle.
 Medianus, a, um—Middle.
 Mediocris, e—Middling, ordinary.
 Medius, a, um—Middle, ordinary.
 Medullaris, e—In the marrow or middle part, like a pith.
 Megacephalus, a, um—Large-headed.
 Megalops—Having large eyes.
 Megambonatus, a, um—Having a great umbo.
 Megambonus, a, um—Having a large umbo.
 Megastomus, a, um—Having a large mouth.
 Megastylus, a, um—Having large spines.
 Megistus, a, um—Very large.
 Melaniformis, e—Shaped like Melania.

- Melanoides—Like Melania.
 Melliformis, e—Purse-shaped.
 Melo—An apple-shaped melon.
 Meloncus, a, um—Like a small melon.
 Meloniformis, e—Melon-shaped.
 Melonoides—Like a melon.
 Membranaceus, a, um—Like a parchment, skinny.
 Meniscus, a, um—A crescent-shaped body.
 Merianopteroides—Like Merianopteria.
 Meristoides—Like Merista.
 Mesacosta—Having middle ribs.
 Mesacostalis, e—Middle-ribbed.
 Mesambonatus, a, um—Having a middle umbo.
 Mesastrialla, e—Middle striated.
 Mesalle, e—Middle parted.
 Mesolobus—Having a middle lobe.
 Meta—Any thing in a conical form.
 Metallicus, a, um—Metallic.
 Metula—A little butt or small pyramid.
 Mica—A crumb or little thing.
 Micans—Stretching out, glittering.
 Microbasalis, e—Having a small base.
 Microcarpus, a, um—Small-fruited.
 Microdentus, a, um—Small-toothed.
 Microdus, a, um—Having small teeth.
 Microlobus, a, um—Small-lobed.
 Micronema—A small thread.
 Microphorus, a, um—Small-bearing.
 Microphyllus, a, um—Small-leaved.
 Micropleura—Having a small rib.
 Micropterus, a, um—Small-winged.
 Microphthalmus, a, um—Small-eyed.
 Micropus—Small foot.
 Microscopicus, a, um—Microscopic.
 Microstigma—Small dot.
 Microstylus—Small spine or pale.
 Micurus, a, um—Small-tailed.
 Micula—A little crumb or grain.
 Millebrachiatus, a, um—Many-armed.
 Milleporaceus, a, um—Having innumerable pores.
 Millepunctatus, a, um—Many-dotted.
 Mimicus, a, um—Mimic.
 Minimus, a, um—The least or smallest.
 Minor—Less, smaller.
 Minuens—Diminishing, making less.
 Minus, a, um—Less.
 Minusculus, a, um—Rather less, rather small.
 Minutisectus, a, um—Finely marked or divided.
 Minutissimus, a, um—Very minute.
 Minutulus, a, um—Very small.
 Minutus, a, um—Diminished, small, minute.
 Mirabilis, e—Extraordinary, wonderful, strange.
 Mirus, a, um—Wonderful, astonishing, extraordinary.
 Miser, era, erum—Wretched, unfortunate.
 Mitella—A head-band, a kind of turban.
 Mithrax—A precious stone.
 Mitigatus, a, um—Tamed, civilized, softened.
 Mitis, e—Ripe, flexible, placid.
 Mitra—A head-band, turban.
 Mixtus, a, um—Mixed.
 Modestus, a, um—Moderate, modest.
 Modiolaris, e—Like Modiola, or a small measure.
 Modioliformis, e—Like a small measure.
 Modulatus, a, um—Symmetrical, well-proportioned.
 Modulus—A small measure.
 Molaris, e—Pertaining to grinding.
 Molestus, a, um—Troublesome, difficult.
 Mollis, e—Flexible, delicate, effeminate.
 Moniliferus, a, um—Bead-bearing.
 Moniliformis, e—Like a necklace.
 Monostigma—Single dot.
 Monstruosus, a, um—Strange, monstrous.
 Monticola—A dweller in the mountains.
 Monticuliferus, a, um—Little mountain-bearing.
 Monticulus—A small mountain.
 Morbillianus, a, um—Measly, spotted.
 Mordax—Biting, given to biting.
 Morsum—That which is bitten off.
 Mortifer, era, erum—Deadly.
 Mucro—A sharp point or edge.
 Mucronatus, a, um—Pointed.
 Mucrospinus, a, um—Sharp-spined.
 Multatenuatus, a, um—Much attenuated.
 Multibrachiatus, a, um—Many-armed.
 Multicallatus, a, um—Much plastered.
 Multicameratus, a, um—Many-chambered.
 Multicarinatus, a, um—Many-keeled.
 Multicaulis, e—Many-stalked.
 Multicinctus, a, um—Many-girded or banded.
 Multicornis, e—Many-horned.
 Multicostatus, a, um—Many-ribbed.
 Multicosta—Having many ribs.
 Multifasciatus, a, um—Many-bundled.
 Multigranulosus, a, um—Many-grained.
 Multigrumus, a, um—Much heaped up.
 Multilamella—Having many thin plates.
 Multilamellosus, a, um—Having many lamellae.
 Multilineatus, a, um—Many-lined.
 Multiliratus, a, um—Many-furrowed.
 Multinodosus, a, um—Many-noded.
 Multinotatus, a, um—Having many marks or tracks.
 Multiplicatus, a, um—Many-folded.
 Multipora—Having many pores.
 Multiporatus, a, um—Having many pores or openings.
 Multipunctatus, a, um—Many dotted or punctured.
 Multiradiatus, a, um—Many-rayed.
 Multiramosus, a, um—Having many branches.
 Multisectus, a, um—Having many divisions or divided folds.
 Multisegmentatus, a, um—Having many segments.
 Multiseptus, a, um—Having many divisions.
 Multiseriatus, a, um—Having many rows or series.
 Multisinuatus, a, um—Many-furrowed.
 Multispinosus, a, um—Many-spined.
 Multistriatus, a, um—Many-striated.
 Multituberculatus, a, um—Having many tubercles.
 Multitubulatus, a, um—Having many pipes.
 Multivolvus, e—Many whorled or rolled.

cola, or a small

small measure.
metrical, well-pro-

re.
grinding.
some, difficult.
te, effeminate.
bearing.
ecklace.

ange, monstrous.
the mountains.
Little mountain-

untain.
sly, spotted.
biting.
bitten off.
dly.
edge.
ted.
arp-spined.
Much attenuated.
Many-armed.
Much plastered.
Many-chambered.
Many-keeled.
lked.
Many-girded or

rned.
any-ribbed.
y ribs.
any-bundled.
Many-grained.
uch heaped up.
uch thin plates.
Having many la-

any-lined.
ny-furrowed.
any-noded.
aving many marks

any-folded.
y pores.
aving many pores

Many dotted or

any-rayed.
— Having many

ing many divisions

um—Having many

ing many divisions.
aving many rows or

Many-furrowed.
Many-spined.
Many-striated.

um—Having many

Having many pipes.
horled or rolled.

Mummiformis, e—Resembling a mummy.
Mundus, a, um—Neat, trim, delicate.
Mundulus, a, um—Neat, trim, delicate.
Muralis, e—Of or belonging to a wall.
Muricatus, a, um—Full of sharp points, pointed.
Musculosus, a, um—Full of muscles.
Mutabilis, e—Inconstant, variable.
Mutatus, a, um—Altered, changed.
Mutus, a, um—Dumb, silent.
Myriophyllus, a, um—Many-leaved.
Myrmecophorus, a, um, wart-bearing.
Mytiliformis, e—Like Mytilus.
Mytilimeris, e—Pertaining to Mytilus.
Mytiloides—Like Mytilus.

Nacrea—Iridescent, like mother-of-pearl.
Nactus, a, um—Obtained, stumbled upon.
Naiadiformis, e—Like a water-nymph.
Nanus—A dwarf.
Nassa—A net, weel, wicker-basket.
Nassula—A little bag-net.
Nasutus, a, um—Large-nosed.
Natalis, e—Native, produced, natural.
Natator—A swimmer.
Naticoides—Like Natica.
Nautiloides—Like Nautilus.
Navalis, e—Of or belonging to ships, naval.
Navicella—A small vessel.
Naviformis, e—Ship-formed.
Navigolum—A little boat.
Nebulosus, a, um—Full of mist, hazy.
Neels—Death.
Neglectus, a, um—Neglected, overlooked.
Nervatus, a, um—Full of nerves or fibers.
Nervosus, a, um—Full of fibers, sinewy.
Neuropteroides, a, um—Like Neuropteris.

Nexilis, e—Knit, tied or wreathed together, twining.
Nexus, a, um—Linked together, interlaced.
Nitela—Brightness, splendor.
Nitens—Shining, neat, beautiful.
Nidulus, a, um—Somewhat spruce, rather trim.

Nitidus, a, um—Neat, shining, polished.
Nobilis, e—Famous, celebrated, noble.
Nobilissimus, a, um—Most celebrated.
Nodobrachiatus, a, um—Knotty-armed.
Nodocarinatus, a, um—Knotty-keeled.
Nodocostatus, a, um—Knotty-ribbed.
Nodocosta—Having knotty ribs.
Nodolonsatus, a, um—Knotty-backed.
Nodomarginatus, a, um—Knotty-margined.
Nodosarius, a, um—Knotty.
Nodostriatus, a, um—Having knotty striae.
Nodosus, a, um—Knotty, full of knots.
Nodulatus, a, um—Knotted.
Noduliferus, a, um—Knot or node bearing.
Nodulostriatus, a, um—Having small knotty striae.
Nodulosus, a, um—Full of little nodes or knots.
Normalis, e—Made by the square or rule.
Notabilis, e—noteworthy, remarkable, extraordinary.
Notans—Noting, marking.
Notatus, a, um—Marked, branded, noted, dotted.

Nothus, a, um—Spurious, not genuine, of mixed breed.

Notus, a, um—Well known, notorious.

Nuciformis, e—Nut-shaped.

Nucleatus, a, um—Deprived of the kernel, stoned.

Nucleiformis, e—Kernel-shaped.

Nucleolatus, a, um—Like a little nut.

Nucleus—A kernel, nut.

Nuculiformis, e—Shaped like Nucula.

Nuculoides—Like Nucula.

Nudus, a, um—Naked, uncovered, empty, alone.

Numerosus, a, um—Numerous, manifold.

Nummifer, era, erum—Coin or disk bearing.

Nummiformis, e—Coin-shaped.

Nummularius, a, um—Of or pertaining to money.

Nummularis, e—Like a little coin.

Nuntius—A messenger.

Nuperus, a, um—Late, newly come or taken, recent.

Nuptialis, e—Nuptial.

Nutans—Nodding, bending backward and forward.

Nutrix—A nurse, the breast or pap.

Nux—A nut.

Nymphalis, e—Of or belonging to a fountain.

Obcordatus, a, um—Inversely heart-shaped.

Obesus, a, um—Fat, plump, swollen.

Oblatus, a, um—Showing, exhibiting.

Obliquatus, a, um—Bent, oblique.

Obluinodus—Oblique-knot.

Oblivus, a, um—Oblique, sidewise.

Oblongifolius, a, um—Oblong-leaved.

Oblongus, a, um—Rather long, oblong.

Obmaximus, a, um—Large in front.

Obovatus, a, um—Inversely ovate.

Obpyramidalis, e—Inversely pyramidal.

Obscurus, a, um—Hidden, not understood, obscure.

Obsolescens—Grown old.

Obsoletus, a, um—Antiquated, obsolete.

Obtectus, a, um—Covered, disguised.

Obtusidens—Blunt-toothed.

Obtusifolius, a, um—Obtuse-leaved.

Obtusilobus, a, um—Obtuse-lobed.

Obtusiplicatus, a, um—Obtuse-plaited.

Obtusaipira—Having a blunt spire.

Obtusus, a, um—Blunted, obtuse.

Obuncus, a, um—Bent in, hooked.

Obvius, a, um—Meeting, laying open, exposed.

Occasus, a, um—Crushed, stricken to the ground.

Occidaneus, a, um—Western.

Occidens—The west, western.

Occidentalis, e—Western.

Oceanus, a, um—Of or belonging to the ocean.

Ocellatus, a, um—Having little eyes.

Octobrachiatus, a, um—Eight-armed.

Octocostatus, a, um—Eight-ribbed.

Octonar'us, a, um—Of the number eight.

Octonotatus, a, um—Having eight marks or tracks.

Ocul'us, a, um—Having eyes.

- Oculiferus, a, um—Eye-bearing.
 Oculinus, a, um—Like an eye.
 Odontopteroides—Like Odontopteris.
 Offula—A small piece.
 Oliviformis, e—Shaped like an olive.
 Oligospiratus, a, um—Having few whorls.
 Olla—A pot.
 Ollicula—A little pot.
 Omphaloides—Like a navel or boss.
 Onustus, a, um—Filled, loaded, burdened.
 Ophioglossoides—Like Ophioglossus.
 Opimus, a, um—Fertile, fruitful, fat, large, plump.
 Oppletus, a, um—Filled.
 Oppositus, a, um—Opposite, placed before.
 Optatus, a, um—Wished, desired, longed for.
 Opusculum—A little fabric.
 Orbicaudatus, a, um—Having a circular tail.
 Orbicella—A little circle.
 Orbicularis, e—Circular, orbicular.
 Orbiculatus, a, um—Of a round or circular form, orbiculate.
 Orbiculostoma—Having a circular mouth.
 Orbicula—Having round pores.
 Ordinatus, a, um—Set in order, regular, ranged in rows.
 Oreopteroides—Like Oreopteris.
 Organum—An instrument, implement, or pipe.
 Oriens—Rising, beginning.
 Orientalis, e—Eastern.
 Originarius, a, um—Original.
 Ornatisimus, a, um—Very ornate, highly adorned.
 Ornatus, a, um—Adorned, embellished.
 Ornigranulus, a, um—Having granules.
 Ornithenoides—Like bird-tracks.
 Orthambonites—Having a straight umbo.
 Orthidoideus, a, um—Like Orthis.
 Orthonotus, a, um—Straight-backed.
 Osculum—A pretty little month.
 Ostiolatus, a, um—Having small openings.
 Ovals, e—Oval, egg-shaped.
 Ovatifolius, a, um—Ovate-leaved.
 Ovatiopora—Having oval pores.
 Ovatus, a, um—Shaped like an egg, ovate.
 Ovibos—The musk ox.
 Ovidactylus—Having ovate toes.
 Oviformis, e—Egg-shaped.
 Ovoidactylus—Having ovoid toes.
 Ovoides—Having an egg shape, ovoid.
 Ovoidens, a, um—Having a form like an egg, ovoid.

Pabulocrinus—Crinoid-food. A word founded on the erroneous opinion that crinoids lived on Gasteropoda.
 Pacator—A peace-maker.
 Pachylirus—Having a thick hand.
 Pachydactylus, a, um—Having thick fingers or thick toes.
 Pachyderma—A thick skin.
 Pachynervis, e—Having thick veins or thick nerves.
 Pachypteroides—Like Pachypteris.
 Pachyteta—Having a thick shell.
 Palaeotrochus—Ancient Trochus.
 Paliformis, e—Shovel-like or stake-like.
 Palmatiformis, a, um—Divided like a hand.
 Palmatus, a, um—Marked with the palm of a hand, palmate.
 Palmpes—Broad-footed.
 Palpebra—The eyelid.
 Paludiformis, e—Shaped like Paludina.
 Palum—A pale, stake.
 Pandatus, a, um—Bent, bowed down in the middle.
 Pandoriformis, e—Shaped like Pandora.
 Pandus, a, um—Bent, crooked, curved.
 Panicum—A grain, panic-grass.
 Pannus, a, um—Ragged, tattered.
 Pannosus, a, um—Full of rags, ragged.
 Papilioniformis, e—Shaped like a butterfly.
 Papillatus, a, um—Bud-shaped, covered with papilli.
 Papillosus, a, um—Full of buds, verrucose.
 Papulatus, a, um—Covered with nipples, warty.
 Papulosus, a, um—Full of pimples.
 Paradoxicus, a, um—Paradoxical.
 Paradoxus, a, um—Strange, contrary to received opinions.
 Paralius, a, um—That grows by the seaside.
 Parallelus, a, um—Parallel.
 Paralleodontus, a, um—Having parallel teeth.
 Parasiticus, a, um—Parasitic.
 Paridens—Having equal teeth.
 Parilis, e—Equal, like, proportionate.
 Partitus, a, um—Proportionably divided.
 Parvibrachiatus, a, um—Small-armed.
 Parvinodus—Having a small knot.
 Parvirostris—Having a little beak.
 Parvispira—Having a small spire.
 Parvituba—Having a small tube.
 Parviusculus, a, um—Quite small.
 Parvulipora—Having small pores.
 Parvulus, a, um—Very small.
 Parvus, a, um—Small, narrow, short, little.
 Patellarius, a, um—Belonging to a plate, plated.
 Patellifer, a, um—Dish-bearer.
 Patelliformis, e—Dish-shaped.
 Patens—Open, wide, extending, spreading.
 Paternus, a, um—Paternal.
 Patulus, a, um—Standing open or opened, wide, large.
 Paucicristatus, a, um—Few-crested.
 Paucidactylus, a, um—Few-fingered.
 Paucinodus, a, um—Having few nodes.
 Pauciradiatus, a, um—Few-rayed.
 Pauciramus, a, um—Having few branches.
 Pauciseptus, a, um—Having few septa.
 Pauper—Poor, small, impoverished.
 Pauperatus, a, um—Poor, impoverished.
 Pauperculus, a, um—Poor.
 Pecteniformis, e—Shaped like a Pecten.
 Pectenoides, a, um—Like a Pecten.
 Pectinaceus, a, um—Of or belonging to a comb, or to the Pecten.
 Pectinatus, a, um—Sloping two ways like a comb.
 Pectinellus, a, um—Like a little comb.
 Pectiniferus, a, um—Comb-bearing.
 Pectunculoides—Like Pectunculus.
 Peculiaris, e—Peculiar, remarkable, singular.

ded like a hand.
with the palm of

d like Paludina.

owed down in the

like Pandora.
ooked, curved.
grass.
attered.
f rags, ragged.
ed like a butterfly.
-shaped, covered

of buds, verrucose,
red with nipples,

of pimples.
adoxical.
ge, contrary to re-

grows by the sea-

nel.

—Having parallel

sitic.
teeth.

proportionate.

tionably divided.

—Small-armed.

small knot.

little beak.

small spire.

small tube.

quite small.

small pores.

small.

narrow, short, little.

longing to a plate,

bearer.

shaped.

tending, spreading.

pal.

ing open or opened,

Few-crested.

Few-fingered.

aving few nodes.

Few-rayed.

aving few branches.

aving few septa.

mpoverished.

or, impoverished.

poor.

ed like a Pecten.

Like a Pecten.

or belonging to a

ten.

ping two ways like

e a little comb.

omb-bearing.

Pectunculus.

remarkable, singular.

Peduncularis, e—Of or belonging to a little foot.

Pedunculatus, a, um—Little-footed.

Pelagicus, a, um—Belonging to the sea.

Pellicula—A small skin or hide.

Pellucidus, a, um—Clear, transparent.

Peloris—A shell fish.

Peltatus, a, um—Armed with shields.

Peltigerus, a, um—Shield-bearing.

Pendens—Hanging, depending.

Pendulus, a, um—Hanging down, pendent,

pendulous.

Penetrans—Piercing, penetrating.

Penicilliformis, e—Brush or pencil-shaped.

Penicillus—A painter's brush or pencil.

Pennatus, a, um—Winged, feathered.

Pennatulus, a, um—Provided with wings.

Penniformis, e—Feather-shaped.

Pentadactylus, a, um—Five-fingered.

Pentagonus, a, um—Pentagonal.

Pentalobus, a, um—Five-lobed.

Pentaspinus, a, um—Five-spined.

Peraeetus, a, um—Very sharp, very acute.

Peramplus, a, um—Very large.

Perangulatus, a, um—Very angular.

Perannulatus, a, um—Many-ringed, very

annular.

Perantiquus, a, um—Very ancient.

Perarctus, a, um—Very close, small, or

slender.

Perasper, a, um—Very rough.

Perattenuatus, a, um—Very attenuated,

drawn out.

Percarinatus, a, um—Very strongly keeled.

Percingulatus, a, um—Encircled with many

lines, many-girded.

Peridentatus, a, um—Many-toothed.

Peregrinus, a, um—Strange, foreign.

Perelegans—Very neat, very elegant.

Perextensus, a, um—Very much extended.

Perforator—A borer through.

Perforatus, a, um—Bored through.

Perfossulatus, a, um—Having many little

ditches.

Pergibbosus, a, um—Very gibbous.

Pergracilis, e—Very slender.

Perhumerosus, a, um—Having angular

shoulders.

Perinflatus, a, um—Much inflated, swol-

len.

Periphris—A round saw.

Perizomatus, a, um—Girdled, banded.

Perlamellosus, a, um—Very lamellose, hav-

ing very thin plates.

Perlatus, a, um—Very wide.

Permargatus, a, um—Large-bordered.

Permultus, a, um—Very many.

Pernasutus, a, um—Very nasute.

Perniformis, e—Shaped like a Perna.

Pernodosus, a, um—Very nodose, knotty.

Perobliquus, a, um—Very oblique.

Peroblongus, a, um—Somewhat oblong.

Peroccidens—From the far West.

Perornatus, a, um—Very ornate.

Perovalis, e—Rather oval.

Perovatus, a, um—Very ovate, or nearly

round.

Perparvus, a, um—Very small.

Perplanus, a, um—Very plain.

Perplexus, a, um—Confused, entangled,

intricate.

Perplicatus, a, um—Interlaced, entangled,

many-folded.

Perpusillus, a, um—Very small.

Perrostellatus, a, um—Having a very little

beak.

Persicaria—A genus of plants.

Persimilis, e—Very similar.

Persinuatus, a, um—Very sinuate or chan-

neled.

Personatus, a, um—Masked, assumed, dis-

guised.

Persiphonatus, a, um—Having a large si-

phuncle.

Perspectivus, a, um—Thoroughly viewed.

Perspicator—Sharp-sighted.

Perspinulatus, a, um—Having many little

thorns or spines.

Perstrialis, e—Having many striae.

Perstriatus, a, um—Very much striated.

Persulcatus, a, um—Very much furrowed.

Pertenuis, e—Very thin, small, or fine.

Pertextus, a, um—Interwoven.

Pertinax—That holds fast, clings to.

Perumbonatus, a, um—Having a very con-

vex umbo.

Perumbrosus, a, um—Very shady.

Perundatus, a, um—Very wavy.

Perundulatus, a, um—Very wavy.

Perverus, a, um—Turned around.

Pervetus, a, um—Very old.

Pervetustus, a, um—Very old.

Pervicax—Immovable, stern.

Pervolutus, a, um—Very much rolled.

Pescapreoli—Having a stock supported by

a small tendril.

Pescervæ—Having deer-feet.

Pesovis—Having sheep-feet.

Petasisformis, e—Cap-shaped.

Petechialis, e—Spotted.

Petilus, a, um—Thin, slender.

Petrifactor—Stone-maker.

Petrodoides—Like Petrodus.

Pexatus, a, um—Clothed in a garment with

a nap on it.

Pharovicinus, a, um—Near the light-house.

Phaseolus—A kidney-bean.

Phaseolinus—Like a bean.

Phlyctainodes—Pimply, pustulous.

Phoca—A seal, sea-dog.

Pholadiformis, e—Like Pholas.

Pholadis—Like a Pholas.

Phragmoceras—Partitioned horn.

Phycoides—Like sea-weed.

Piger, gra, grum—Sluggish.

Pileatus, a, um—Covered with a cap.

Pileiformis, e—Cap-shaped.

Pileolus—A skull-cap, a little cap.

Pileolum—A little cap.

Pileus—A cap or hat.

Pilosus, a, um—Hairy, shaggy.

Pinaster—A wild pine.

Pinguis, e—Fat, plump, fertile.

Pinnatifidus—Having cleft pinnae.

Pinnatus, a, um—Feathered, plumed,

winged.

Pinniformis, e—Like Pinna.

Piscator—A fisher.

- Pisiformis, e—Pea-shaped.
 Pistilliformis, e—In the form of a pestle.
 Pistillus—A pounder, pestle.
 Pisum—A pea.
 Placenta—A cake.
 Placidus, a, um—Placid, smooth.
 Plagiosus, a, um—Full of wounds or stripes.
 Planiceps—Flat-headed.
 Planicosta—Having flat ribs.
 Planidorsalis, e—Flat or smooth-backed.
 Planidorsatus, a, um—Flat or smooth-backed.
 Planifrons—Having a plane front.
 Planimarginatus, a, um—Flat-margined.
 Planiramosus, a, um—Having flat branches.
 Planirostris—Having a smooth beak.
 Planispira—Having a flat spire.
 Planistria—Having flat striae.
 Planistriatus, a, um—Having flat striae.
 Planobasalis, e—Having a flat base.
 Planobrachiatus, a, um—Smooth-armed.
 Planoconvexus, a, um—Flat, convex.
 Planocostatus, a, um—Flat-ribbed.
 Planodiscus—Flat disk.
 Planodorsalis, e—Smooth or flat-backed.
 Planodorsatus, a, um—Having a smooth or flat back.
 Planogyrratus, a, um—Flat-whorled.
 Planorbiformis, e—Like Planorbis.
 Planosulcatus, a, um—Plane-furrowed.
 Planovolvis, e—Flat-whorled.
 Planulatus, a, um—Rather flat.
 Planumbonus, a, um—Having a smooth umbo.
 Planus, a, um—even, level, flat, plane.
 Platymarginatus, a, um—Flat-margined.
 Platybasis—Having a flat base.
 Platycephalus—Having a flat head.
 Platynervis, e—Flat-nerved.
 Platynotus—Having a flat ridge or back.
 Platypleurus—Having flat sides.
 Platyplus—Broad-footed.
 Platyrrachis—Having a flat rachis.
 Platyrs—Broad.
 Platystigma—Having flat scars, dots, or pits.
 Platystomus, a, um—Having a broad mouth.
 Plebeiformis, e—Like a plebeian.
 Plebeius, a, um—Common.
 Pleiopleura—Having wide ribs.
 Pleiissimus, a, um—The largest.
 Plenus, a, um—Full, plump.
 Pleurexanthemus—Having the pleura extending out.
 Pleurites—The side, lateral.
 Pleurodictyoides—Like Pleurodictyum.
 Pleuropletha—Having the side behind.
 Pleuroptera—Having side wings.
 Pleuropteryx—Having side wings.
 Pleurovimeus, a, um—Having side wicker-work.
 Plicatellus, a, um—Having small folds.
 Plicatilis, e—That may be folded, flexible.
 Plicatulus, a, um—Having little plications or folds.
 Plicatus, a, um—Plaited, folded.
 Pliciferus, a, um—Fold-bearing or plaited.
 Plicomphalus—Folded in the middle.
 Pluma—A small feather.
 Plumarius, a, um—Embroidered with feathers.
 Plumosus, a, um—Full of feathers, feathery.
 Plumula—A little feather.
 Plumulosus, a, um—Full of feathers.
 Pluriradialis, e—Many-rayed.
 Pocillatus, a, um—Little-cupped.
 Pocilliformis, e—Cup-shaped.
 Pocillum—A little cup.
 Poculum—A cup, bowl, or goblet.
 Pogonias—A kind of comet.
 Politus, a, um—Polished, smoothed.
 Polydactylus, a, um—Many-fingered.
 Polygonus, a, um—Having many angles, polygonal.
 Polygyrratus, a, um—Many coiled or whorled.
 Polymorphus, a, um—Many-formed.
 Polyphyllus, a, um—Many-leaved.
 Polypleurus, a, um—Having many ribs.
 Polysporus—Having many spores.
 Polystomellus, a, um—Having many little mouths.
 Ponderosus, a, um—Heavy, ponderous.
 Ponticulus—A little bridge.
 Porcatus, a, um—Ridged, furrowed.
 Porosus, a, um—Full of pores.
 Porrectus, a, um—Extended, stretched, or spread out.
 Posticus, a, um—Posterior.
 Postremus, a, um—The last, hindmost, worst.
 Poststriatus, a, um—Having a striated posterior.
 Potens—Powerful.
 Poterium—A drinking vessel, a cup.
 Praecedens—Going before, surpassing.
 Praeceptus, a, um—Anticipated, going before.
 Praecursor—A forerunner.
 Praelongus, a, um—Very long.
 Praematurus, a, um—Very early, untimely, premature.
 Praemorsus, a, um—Bitten off, jagged.
 Praenuntius, a, um—That foretells, or forebodes.
 Praeumbonus, a, um—Very protuberant.
 Prateriformis, e—Prateriform.
 Pravus, a, um—Crooked, deformed, distorted.
 Preciosus, a, um—Precious, splendid.
 Precius, a, um—That brings forth ripe grapes before other vines.
 Pressulus, a, um—Somewhat pressed in, compressed.
 Pressus, a, um—Pressed.
 Pretiosus, a, um—Precious, valuable.
 Primaevus, a, um—Primeval.
 Primarius, a, um—One of the first, remarkable, principal.
 Primigenius, a, um—First of its kind, original, primitive.
 Primitivus, a, um—First of its kind, primitive.
 Primordialis, e—Primordial, original, first of all.
 Primus, a, um—The first.
 Princeps—The first, chief, original, principal.
 Principalis, e—First, original, principal.
 Prius, a, um—Ancient, old.

bordered with
 feathers, feathery.
 feathers.
 ed.
 ipped.
 ed.
 goblet.
 t.
 smoothed.
 y-fingered.
 ng many angles,
 coiled or whorled.
 y-formed.
 y-leaved.
 ng many ribs.
 y spores.
 aving many little
 y, ponderous.
 e.
 furrowed.
 ores.
 led, stretched, or
 r.
 last, hindmost,
 ing a striated pos-
 sessel, a cup.
 y, surpassing.
 icipated, going be-
 long.
 y early, untimely,
 n off, jagged.
 foretells, or fore-
 y protuberant.
 form.
 d, deformed, dis-
 ns, splendid.
 bring forth ripe
 res.
 ewhat pressed in,
 us, valuable.
 eval.
 of the first, remark-
 first of its kind,
 of its kind, primi-
 dial, original, first
 , original, principal.
 ginal, principal.
 , old.

Pristiniformis, e—An ancient form.
Pristinus, a, um—Primitive, early.
Pristis—Any sea monster or saw-fish.
Problematicus, a, um—Problematical, un-
 settled, uncertain.
Proboscidiæ, e—Having a proboscis.
Proboscidiatus, a, um—Having a proboscis.
Procerus, a, um—High, tall.
Proclivis, e—Sloping, steep.
Productus, a, um—Drawn out, produced.
Profundus, a, um—Deep, profound.
Projectus, a, um—Thrown out, projected.
Prolatus, a, um—Brought forth, extended,
 enlarged.
Prolificus, a, um—Prolific, fruitful.
Prolifer, era, erum—Prolific, productive,
 fruitful.
Prolixus, a, um—Stretched far out, long,
 broad.
Prolongatus, a, um—Prolonged.
Prolongus, a, um—Prolonged, stretched
 out.
Prominulus, a, um—Projecting a little,
 rather prominent.
Promissus, a, um—Hanging down, putting
 forth.
Pronis, e—Bent forward, inclined down-
 ward.
Pronus, a, um—Turned forward, bent or
 inclined.
Propinquus, a, um—Near, hard by, re-
 lated to.
Proporoides—Like *Propora*.
Proprius, a, um—Peculiar, proper.
Prora—The prow of a ship.
Proteiformis, e—Having many shapes.
Protensus, a, um—Stretched out.
Protexus, a, um—Closely woven.
Protuberans—Projecting, protuberant.
Proximus, a, um—Nearest.
Pseudogaleatus—False *Galeatus*.
Pseudolineatus, a, um—False-lined.
Pseudomarginalis, e—False-margined.
Pseudomurrayanus, a, um—False *Murray-*
anus.
Pseudosagittatus—False *Sagittatus*.
Psilophleus—Having rough bark.
Pterineiformis, e—Shaped like *Pterinea*.
Pterocephalus—Having a winged head.
Pteroides—Wing-like.
Pterotus, a, um—Winged, feathered.
Pudicus, a, um—Shamefaced, modest.
Pugniunculus—A small dagger.
Pugnax—War-like, combative.
Pugnus—A fist, a handful.
Pulcellus, a, um—Beautiful little.
Pulchellus, a, um—Beautiful little, or some-
 what beautiful.
Pulcher, a, um—Beautiful.
Pulex—A flea.
Pulicaris—Like a flea.
Pulmoneus, a, um—Spongy like the lungs.
Pumilus, a, um—Dwarfish, diminutive,
 little.
Punctatus, a, um—Punctured, dotted.
Punctiferus, a, um—Puncture-bearing.
Punctifrons—Dotted in front.
Punctillatus, a, um—Finely dotted.
Punctipora—Having dotted pores.

Punctolineatus, a, um—Having dotted or
 pitted lines or furrows.
Punctostriatus, a, um—Having pricked or
 dotted striae.
Punctulatus, a, um—Marked with small
 spots.
Punctuliferus, a, um—Bearing punctures
 or dots.
Pusillus, a, um—Very small, petty, in-
 significant.
Pustulatus, a, um—Blistered, covered with
 pustules.
Pustuliferus, a, um—Bearing blisters or pus-
 tules.
Pustulosus, a, um—Full of blisters, pimples,
 or pustules.
Puteatus, a, um—Having little pits or wells.
Puteolatus, a, um—Pitted.
Putillus—A child or dwarf.
Pygmaeus, a, um—Dwarfish.
Pyramidalis, e—Pyramidal, pointed like a
 pyramid.
Pyramidatus, a, um—Pyramidal, made like
 a pyramid.
Pyriformis, e—Pyriform, pear-shaped.
Pyxidatus, a, um—Box-like.
Pyxidicula—A small box.
Pyxidiformis, e—Box-shaped.
Quadrangularis, e—Quadrangular.
Quadrangulatus, a, um—Quadrangular.
Quadrans—A quarter or a fourth part.
Quadraticaudatus, a, um—Square-tailed.
Quadratifolius, a, um—Quadrately-leaved.
Quadratus, a, um—Four-cornered, squared,
 quadrate.
Quadribrachiatus, a, um—Having four arms.
Quadriceps—Square-headed.
Quadriceinctus, a, um—Four banded or
 girdled.
Quadriconatus, a, um—Four-ribbed.
Quadrilateralis, e—Quadrilateral, four-
 sided.
Quadrimumeronatus, a, um—Having four
 sharp points or spines.
Quadrupartitus, a, um—Four-parted.
Quadriseriatus, a, um—Having four series.
Quadrispinus, a, um—Four-spined.
Quadrissulcatus, a, um—Four-furrowed.
Quadrivolvus, e—Four-whorled.
Quadrula—A little square.
Quasillus—A little basket.
Quaternarius, a, um—Containing four, qua-
 ternary.
Quatuordecembrachialis, e—Having four-
 teen arms.
Quercifolius, a, um—Oak-leaved.
Quincuncialis, e—Made in the form of a
 quincunx.
Quinquelobus, a, um, five-lobed.
Quinquenodus, a, um—Having five nodes
 or knots.
Quinquepartitus, a, um—Five-parted.
Quinquesulcatus, a, um—Five-furrowed.
Racematus, a, um—Having clusters.
Racemosus, a, um—Full of clusters, clus-
 tering.
Radians—Radiating, glittering.

- Radiatoplicatus**, a, um—Rayed and plaited.
Radiatus, a, um—Rayed.
Radicans—Rooting.
Radiciformis, e—Root-like.
Radicosus, a, um—Full of roots.
Radicula—A small root.
Ramifer, era, erum—Branch-bearing.
Ramosissimus, a, um—Very branchy.
Ramosus, a, um—Full of branches, ramose.
Ramulosus, a, um—Full of little branches.
Ramulus—A little branch.
Rana—A frog.
Ranunculus—A tadpole.
Rapax—Grasping, rapacious.
Raphanus—A radish-root.
Rapheidolabis—Needle-like forceps.
Rapidens—Having grasping teeth.
Raptor—A robber.
Raricosta—Having few ribs.
Raricostatus, a, um—Having few ribs.
Rarinervis, e—Few-nerved or few-veined.
Raripora—Having few pores.
Rarispinus—Having few spines.
Rarus, a, um—Having wide interstices, thin, scattered, rare.
Recedens—Falling back, receding.
Receptaculum—A receptacle.
Rectangularis, e—Rectangular.
Rectangulus, a, um—Rectangular.
Rectiannulatus, a, um—Having straight annulations.
Recticameratus, a, um—Straight-chambered.
Recticardinalis, e—Having a straight cardinal line.
Rectidorsatus, a, um—Straight-backed.
Rectidens—Having straight teeth.
Rectiformis, e—Straight-formed.
Rectilatera—Having straight sides.
Rectilateralis, e—Straight-sided.
Rectilaterarius, a, um—Straight-sided.
Rectilinea—Having straight lines.
Rectinodus, a, um—Having a straight knot or node.
Rectiplicatus, a, um—Having straight plaits or folds.
Rectirostris—Straight beaked.
Rectirostrus, a, um—Straight-beaked.
Rectiseptatus, a, um—Having straight septae.
Rectistriatus, a, um—Having straight furrows.
Rectistylus, a, um—Having straight stems or styles.
Rectus, a, um—Straight.
Recurvatus, a, um—Curved backward.
Recurvirostris—Having a recurved beak.
Recurvus, a, um—Turned back, bent or curved back.
Reflexus, a, um—Bending backward, reflexed.
Regalis, e—Regal, splendid.
Regius, a, um—Regal, majestic.
Regularis, e—Regular, according to a rule, of or belonging to a bar.
Regulatus, a, um—Regulated.
Reliquus, a, um—Remaining.
Remex—A rower, oarsman.
Remibrachiatus, a, um—Paddle-armed.
Remipes—Oar-footed.
Remotiseptum—Having distant barriers or walls.
Remotus, a, um—Removed, distant, remote.
Remus—An oar.
Reniformis, e—Kidney-shaped.
Repandus, a, um—Bent backward.
Repens—Creeping, crawling.
Repertus, a, um—Discovered, hit upon.
Repositus, a, um—Restored, kept, remote, distant.
Reservatus, a, um—Reserved.
Restrictus, a, um—Drawn back, bound up.
Resupinatus, a, um—Lying on one's back, bent backward.
Resupinoides—Like a resupinate form.
Reticularis, e—Reticulated.
Reticulatus, a, um—Made like a net, net-like, reticulated.
Retiferus, a, um—Net-bearing.
Retiformis, e—Net-formed.
Retorquatus, a, um—Turned back.
Retractilis, e—Drawn back.
Retrorsus, a, um—Turned backward, in reversed order.
Retroversus, a, um—Turned backward, in reversed order.
Retusus, a, um—Beaten back, blunt, dull.
Reversus, a, um—Turned about, reversed.
Revolutus, a, um—Rolled back, revolved.
Rhabdocarpus, a, um—Rod-fruited or long-fruited.
Rhombeus, a, um—Rhomboidal.
Rhombicus, a, um—Rhombic.
Rhombiferus, a, um—Rhomb-bearing.
Rhomboidalis, e—Rhomboidal.
Rhomboides—Rhomb-like.
Rhomboidens, a, um—Lozenge-shaped, rhomboid.
Rhombolinearis, e—Rhomb-lined.
Rhynchonelliformis, e—Like Rhynchonella.
Riciniformis, e—Like a tike or tick.
Ricinula—A little tick.
Rictum—The mouth wide open.
Rigens—Stiffened, standing upright.
Rigidus, a, um—Hard, inflexible, rigid.
Rimosus, a, um—Full of cracks, or fissures.
Ringens—Gaping.
Robustus, a, um—Strong, of hard wood.
Robustus, a, um—Strong, robust.
Rostellatus, a, um—Little-beaked.
Rostellum—A little beak.
Rostratus, a, um—Beaked, curved at the end.
Rota—A wheel.
Rotadentatus, a, um—Wheel-toothed.
Rotallinea—Having a round line.
Rotatorius, a, um—Whorled.
Rotatus, a, um—Wheel-shaped.
Rotulatus, a, um—Rounded.
Rotuliformis, e—Little wheel-shaped.
Rotuloides—Like a little wheel.
Rotulus—A little wheel.
Rotundatus, a, um—Rounded.
Rotundifolius, a, um—Round-leaved.
Rotundilobus, a, um—Round-lobed.
Rotundispira—Having a round spire.
Rotundus, a, um—Wheel-shaped, circular, rotund.
Rubellus, a, um—Reddish.

[RAD.—RUB.]

tant barriers or
distant, remote.

ped.
ckward.
g.
ed, hit upon.
l, kept, remote,

ed.
back, bound up.
g on one's back,

pinate form.
like a net, net-

ing.

ed back.
k.
backward, in re-

med backward, in
back, blunt, dull.
about, reversed.
back, revolved.
od-fruited or long-

oboidal.
mbic.
omb-bearing.
oboidal.
e.
—Lozenge-shaped,

mb-lined.
like Rhynchonella.
like or tick.

e open.
ing upright.
nflexible, rigid.
cracks, or fissures.

g, of hard wood.
g, robust.
e-beaked.

ed, curved at the

heel-toothed.
nd line.

orded.
shaped.
ded.
wheel-shaped.
e wheel.

ounded.
ound-leaved.
ound-lobed.
ound spire.
et-shaped, circular,

sh.

Ruber, bra, brum—Red, ruddy.
Rudicula—A wooden spoon, a spatula.
Rudis, e—Rough, unwrought, unpolished.
Rugatinus, a, um—Having little folds or
plaits.
Rugatus, a, um—Having little wrinkles.
Rugicosta—Having wrinkled ribs.
Rugilineatus, a, um—Having wrinkled lines.
Rugiplicatus, a, um—Having wrinkled
plates.
Rugistriatus, a, um—Having wrinkled striae.
Rugosiusculus, a, um—Covered with small
wrinkles.
Rugosus, a, um—Wrinkled, shriveled.
Rugulatus, a, um—Having wide furrows.
Ruguliferus, a, um—Wrinkle-bearing.
Ruidus, a, um—Rough.
Rusticellus, a, um—Somewhat rustic.
Rusticus, a, um—Rural, rustic, rough.

Saccatus, a, um—That is put in a bag, like
a little bag.

Sacculus—A little bag.

Sagittarius, a, um—Of or belonging to an
arrow.

Sagittatus, a, um—Discharging arrows,
barbed like an arrow.

Salamandroides—Like a salamander.

Salebrosus, a, um—Rough, rugged, uneven.

Saliginoides—Like willow wood.

Salisburioides—Like Salisbury.

Samariformis, e—Like elm-seed.

Sanguinolariodeus, a, um—Like Sanguino-
laria.

Sarcinula—A little bundle.

Sarculus—A little hoe.

Sarmenticius, a, um—Of or belonging to
twigs.

Sarmentosus, a, um—Full of twigs or little
branches.

Saxifragifolius, a, um—Leaved like Saxi-
fraga.

Saxivadus, a, um—Creeping over stone.

Scaber, era, erum—Rough, scurfy.

Scaberrimus, a, um—Very rough, scurfy.

Scabiosus, a, um—Scabby, rough, scurfy.

Scabriculus, a, um—Rough.

Scabrosus, a, um—Rough.

Scalariformis, e—Ladder-like.

Scalaris, e—Of or belonging to a flight of
steps, or a ladder.

Scalatus, a, um—Having stairs.

Scalenus, a, um—Unequal-sided, scalene.

Scalpriformis, e—Lancet-shaped.

Scapha—A skiff or boat.

Scintilla—A spark.

Scissilis, e—Split, cleft, or rent.

Scitulus, a, um—Handsome, pretty, elegant.

Scobiniformis, e—Rasp-like.

Scobina—A rasp.

Scolopendrites—Stone-scolopendrium.

Scoparius—A sweeper.

Scorpionis, e—Of or belonging to a scor-
pion.

Serinium—A case, chest, or box.

Scriptiferus, a, um—Writing-bearing.

Scrutator—A searcher, investigator.

Sculptilis, e—Formed or produced by carv-
ing or graving.

Sculptus, a, um—Engraved, sculptured,
carved.

Scutatus, a, um—Armed with a shield.

Scutellatus, a, um—Armed with a little
shield.

Scutelliformis, e—Waiver-shaped.

Scutigerus, a, um—Shield-bearing.

Scutulus, a, um—Lozenge-shaped, check-
ered.

Scyphulus—A small cup.

Scyphus—A cup, a goblet.

Secalinus, a, um—Like small grain.

Secans—A cutter.

Secretus, a, um—Severed, separated, se-
creted.

Sectifrons—Having a divided front.

Sectoralis, e—Like a sector, or cutter.

Secundus, a, um—Following.

Securiformis, e—Ax or hatchet shaped.

Securis—An ax or hatchet.

Segmentatus, a, um—Ornamented with
strips, trimmed, made of pieces.

Selaginoides—Like Selago.

Selago—A plant.

Selectus, a, um—Culled, selected, chosen.

Selenurus—Having a crescent tail.

Selluliformis, e—Like a little seat or stool.

Semicarinatus, a, um—Half-circular.

Semicircularis, e—Half-circular.

Semicostatus, a, um—Half-ribbed.

Semicylindricus, a, um—Half-cylindrical.

Semiellipticus, a, um—Half-elliptical.

Semifasciatus, a, um—Half-bundled or
banded.

Semina—Seed.

Seminosus, a, um—Full of seeds.

Semiorbiculatus, a, um—Half-orbicular.

Semplicatus, a, um—Half-plaited.

Sempunctatus, a, um—Half-dotted.

Semiradiatus, a, um—Half-rayed.

Semiradicatus, a, um—Half-rooted.

Semireductus, a, um—Half bent back.

Semireticulatus, a, um—Half-reticulated.

Semirotundus, a, um—Half-round, semicir-
cular.

Semistriatus, a, um—Half-striated.

Senarius, a, um—Consisting of six.

Senectus, a, um—Aged, very old.

Senex—Old, aged.

Sentosus, a, um—Full of thorns, thorny.

Separatus, a, um—Separated.

Septatus, a, um—Divided with partitions or
septa.

Septemnotatus, a, um—Seven-marked.

Septentrionalis, e—Northern.

Septoris, e—Having seven mouths.

Septus, a, um—Inclosed, enveloped, sur-
rounded.

Septulus, a, um—Buried in deep sleep,
slumbering.

Seriatas, a, um—In series.

Sericeus, a, um—Silken.

Serotinus, a, um—Backward, late.

Serpens—Creeping, crawling.

Serpillifolius, a, um—Thyme-leaved.

Serpuloides—Like Serpula.

Serpuloides, a, um—Snake-like or Serpula-
like.

Serratus—A small saw.

- Serratus*, a, um—Saw-shaped, serrated.
Serrula—A small saw.
Serrulatus, a, um—Like a little saw.
Servilis, e—Of or belonging to a slave, paltry.
Sesquiplicatus, a, um—Once and a half plaited.
Setaceus, a, um—Hairy.
Setiferus, a, um—Bristle-bearing, having coarse hair.
Setigerus, a, um—Bristle-bearing, having coarse hair.
Sexarmatus, a, um—Six-armed.
Sexlobatus, a, um—Six-lobed.
Sexplicatus, a, um—Six-plaited.
Sexradiatus, a, um—Six-rayed.
Sextans—A sixth part.
Sicula—A dagger, sickle, or scythe.
Sidereus, a, um—Of or belonging to the stars, starry.
Sigaretoides—Like *Sigaretus*.
Sigillarioides—Like *Sigillaria*.
Sigillatus, a, um—Adorned with little images or figures.
Sigillum—A sign, mark.
Sigmoideus—Like the Greek letter Sigma.
Sigmoideus, a, um—Like the Greek letter Sigma.
Signatus, a, um—Marked, designated.
Silicula—A little pod.
Siliqua—A pod.
Siliquoideus, a, um—Like a pod.
Similior—Similar.
Simillimus, a, um—Very similar.
Similis, e—Like, resembling, similar.
Simplex—Simple, plain.
Simplicitas—Simpleness, simplicity.
Simulans—Imitating, copying.
Simulator—A copier, imitator.
Simulatrix—A transformer.
Singularis, e—Alone, solitary, singular.
Singularitas—Singleness, being alone or single.
Sinistrorsus, a, um—Toward the left side.
Sinuatus, a, um—Hollowed out, excavated, having depressions.
Sinuosus, a, um—Full of bendings, curves, or folds, sinuous.
Sirpus—A rush, bulrush.
Smilacifolius, a, um—Smilax-leaved.
Sobrina—A cousin.
Socialis, e—Of or belonging to companionship, social.
Solarioides—Like *Solarium*.
Soleniformis, e—Solen-shaped.
Solenoides—Like Solen.
Solidirostris—A solid beak.
Solidissimus, a, um—Very firm or solid.
Solidulus, a, um—Solid.
Solidus, a, um—Firm, compact, solid.
Solitarus, a, um—Lonely, solitary.
Solus, a, um—Alone, single, sole.
Solutus, a, um—Separated, loosened.
Sordidus, a, um—Small, sordid, paltry.
Sororecula—A little sister.
Sparsipora—Having few pores.
Sparsus, a, um—Scattered, separated, dispersed.
Spartarius, a, um—Of or belonging to a broom.
Spathatus, a, um—Spatula-shaped.
Spacious, a, um—Ample, of great extent, spacious.
Spatulatus, a, um—Blade-shaped, spatulate.
Speciosus, a, um—Handsome, beautiful, splendid.
Spectabilis, e—Visible, admirable, remarkable.
Sphæricus, a, um—Of or belonging to a ball, spherical.
Sphærior—A little ball or pill.
Sphærodactylus—Spherical-toed or fingered.
Sphæroidalis, e—Spheroidal.
Sphærolatus, a, um—A widened sphere.
Sphenophylloides—Like *Sphenophyllum*.
Sphenopteroides—Like *Sphenopteris*.
Spicatus, a, um—Pointed, spiked.
Spiculatus, a, um—Having little points.
Spiculus, a, um—Pointed.
Spinalatus, a, um—Spine-winged.
Spiniferus, a, um—Thorn-bearing, thorny, spiny.
Spinigerus, a, um—Thorn-bearing, thorny, spiny.
Spinobrachiatus, a, um—Having spines on the arms.
Spinoclavatus, a, um—Club-spined.
Spinoporus—Having spines and pores.
Spinosulus, a, um—Somewhat thorny.
Spinosus, a, um—Full of thorns, thorny, prickly.
Spinotenticulatus, a, um—Having spine-feelers.
Spinulicosta—Having spines and ribs.
Spinuliferus, a, um—Spine-bearing.
Spinulosus, a, um—Full of little thorns.
Spinula—A little thorn.
Spiralis, e—Spiral.
Spiratus, a, um—Spiral.
Spiriferoides—Like *Spirifera*.
Spiroema—Having spiral threads or lines.
Spirorbis—Spiral-whorl.
Spissiseptus, a, um—Having crowded or numerous septa.
Spissus, a, um—Thick, crowded, compact, dense.
Splendens—Splendid, bright.
Splendidus, a, um—Bright, shining.
Spondyliiformis, e—Shaped like *Spondylus*.
Spondylus—A vertebra, spondyle.
Spongiaxis—Sponge-axis.
Spongilla—A little sponge.
Sponsus, a, um—Promised, betrothed.
Spurius, a, um—Illegitimate.
Squalodens—A kind of fish-tooth.
Squamifer, era, erum—Scale-bearing.
Squamiformis, e—Scale-like.
Squamosus, a, um—Covered with scales, scaly.
Squamula—A little scale.
Stabilis, e—Firm, stable, durable.
Stachyoides—Like *Stachys*.
Stamineus, a, um—Full of threads, thready.
Stella—A star.
Stellaris, e—Of or belonging to a star, starry.
Stellatimulcatus, a, um—Star-furrowed.
Stellatus, a, um—Covered with stars, starred.
Stellifer, era, erum—Star-bearing, starry.
Stellifolius, a, um—Star-leaved.

shaped.
 of great extent,
 shaped, spatulate.
 some, beautiful,
 admirable, re-
 belonging to a ball,
 pill.
 1-toed or fingered.
 al.
 dened sphere.
 phenophyllum.
 phenopteris.
 spiked.
 g little points.
 winged.
 n-bearing, thorny.
 n-bearing, thorny,
 Having spines on
 sub-spined.
 nes and pores.
 what thorny.
 of thorns, thorny.
 n—Having spine-
 nes and ribs.
 ne-bearing.
 of little thorns.
 fera.
 al threads or lines.
 laying crowded or
 crowded, compact,
 right.
 ht, shining.
 ed like Spondylus.
 spondyle.
 a.
 ge.
 ged, betrothed.
 mate.
 fish-tooth.
 scale-bearing.
 like.
 overed with scales,
 e.
 , durable.
 ys.
 of threads, thready.
 zing to a star, starry.
 a—Star-furrowed.
 d with stars, starred.
 ar-bearing, starry.
 -leaved.

Stelliformis, e—Star-shaped.
 Stenoccephalus—Having a narrow head.
 Stenopus—Having a narrow foot.
 Stigmatus, a, um—Branded.
 Stigmosus, a, um—Full of brand-marks,
 branded.
 Stillativus, a, um—Dropping.
 Stipatus, a, um—Crowded together, sur-
 rounded, compressed.
 Stoloniferus, a, um—Bearing a useless sucker
 or water-shoot.
 Stragulus, a, um—Covered.
 Stramineus, a, um—Made of straw.
 Strenuus, a, um—Vigorous, strenuous.
 Striatellus, a, um—Finely channeled.
 Striatiformis, e—Shaped like *Striatus*, another
 species.
 Striatocostatus, a, um—Striæ-ribbed.
 Striatolineatus, a, um—Striæ-lined.
 Striatomarginatus, a, um—Having a striated
 margin.
 Striatopora—Having striæ and pores.
 Striatulus, a, um—Somewhat striated, or
 having small striæ.
 Striatura—Being channeled or fluted, a
 fluting.
 Striatopora—Having striated pores.
 Striatulus, a, um—Furrowed, striated.
 Strictus, a, um—Drawn tight, bound, pressed
 together.
 Strigatus, a, um—Furrowed, channeled,
 grooved, fluted.
 Strigillatus, a, um—Furrowed, fluted.
 Strigosus, a, um—Lean, thin, meager.
 Striobrachiatus, a, um—Having grooved
 arms.
 Striolatus, a, um—Very minutely striated.
 Strix—A furrow, channel, groove.
 Strophium—A twisted girdle, a band.
 Strophomenoides—Like *Strophomena*.
 Styliola—A truncated column.
 Stylus—A pointed instrument, stake, or pale.
 Subabbreviatus, a, um—Somewhat abbrevi-
 ated.
 Subaculeatus, a, um—Somewhat prickly.
 Subæqualis, e—Subequal.
 Subæquatus, a, um—Somewhat equal.
 Subæquilaterus, a, um—Somewhat equal-
 sided.
 Subalatus, a, um—Somewhat winged.
 Subangularis, e—Somewhat angular.
 Subangulatus, a, um—Somewhat angulated.
 Subarcuatus, a, um—Somewhat curved or
 arcuate.
 Subattenuatus, a, um—Somewhat drawn
 out or attenuated.
 Subcæspitosus, a, um—Somewhat cespitose.
 Subcancellatus, a, um—Subcancellated.
 Subcarbonarius—Below the coal.
 Subcardiiformis, e—Somewhat heart-shaped.
 Subcarinatus, a, um—Somewhat keeled.
 Subcavus, a, um—Somewhat excavated,
 hollowed out.
 Subcentralis, e—Subcentral.
 Subcircularis, e—Subcircular.
 Subclavatus, a, um—Somewhat club-shaped.
 Subcompressus, a, um—Subcompressed.
 Subconcausus, a, um—Subconcaue.
 Subconicus, a, um—Subconical.

Subconoideus, a, um—Somewhat conoidal.
 Subconstrictus, a, um—Subconstricted.
 Subcordiformis, e—Somewhat heart-shaped.
 Subcoronatus, a, um—Somewhat adorned.
 Subcorpulentus, a, um—Somewhat corpu-
 lent.
 Subcrassus, a, um—Somewhat thick.
 Subcrenulatus, a, um—Somewhat crenu-
 lated.
 Subcuneatus, a, um—Somewhat wedge-
 shaped.
 Subcuspidatus, a, um—Somewhat pointed.
 Subcylindricus, a, um—Somewhat cylin-
 drical.
 Subcymbiformis, e—Somewhat boat-shaped.
 Subdecussatus, a, um—Somewhat arranged
 in pairs that cross each other.
 Subdemissus, a, um—Somewhat hanging
 down.
 Subdepressus, a, um—Somewhat depressed.
 Subelegans—Somewhat elegant.
 Subellipticus, a, um—Subelliptical.
 Subemarginatus, a, um—Slightly emargin-
 ated.
 Subfalcatus, a, um—Somewhat scythe-
 shaped, subfalcate.
 Subfurcatus, a, um—Somewhat forked.
 Subfusiformis, e—Somewhat spindle-
 shaped.
 Subglobosus, a, um—Somewhat globose.
 Subgracilis, e—Somewhat slender.
 Subhorridus, a, um—Somewhat rough.
 Subimbricatus, a, um—Somewhat imbrici-
 cated.
 Subimpressus, a, um—Somewhat engraved.
 Sublævis, e—Nearly smooth.
 Sublamellosus, a, um—Somewhat in thin
 plates.
 Sublineatus, a, um—Somewhat striated.
 Subliratus, a, um—Somewhat lined.
 Subluuatus, a, um—Somewhat lunate.
 Submarginatus, a, um—Somewhat mar-
 gined.
 Submicronatus, a, um—Somewhat sharp-
 pointed.
 Submutans—Somewhat changing.
 Subnasutus, a, um—Somewhat nasute.
 Subnervosus, a, m—Somewhat veiny.
 Subnodosus, a, um—Somewhat knotty or
 nodose.
 Suborbicularis, e—Somewhat orbicular or
 orb-shaped.
 Suborbiculatus, a, um—Somewhat orbicular.
 Subovalis, e—Suboval.
 Subovatus, a, um—Subovate.
 Suboviformis, e—Somewhat egg-shaped.
 Subpapillosus, a, um—Somewhat papillose.
 Subpapyraceus, a, um—Somewhat like
 Papyrus, the paper-reed.
 Subplanus, a, um—Somewhat flat.
 Subplicatus, a, um—Somewhat plaited.
 Subpulchellus, a, um—Somewhat handsome.
 Subquadrans—Somewhat squared.
 Subquadratus, a, um—Somewhat squared.
 Subramosus, a, um—Somewhat ramose.
 Subramulosus, a, um—Somewhat branchy.
 Subrectus, a, um—Somewhat straight.
 Subretiformis, e—Somewhat net-shaped, or
 net-like.

- Subrhomboidus**, a, um—Somewhat rhomb-like.
Subrigidus, a, um—Somewhat rigid.
Subrotundatus, a, um—Somewhat rounded.
Subrugosus, a, um—Somewhat wrinkled.
Subscalaris, e—Somewhat ladder-shaped.
Subscutulus, a, um—Somewhat hand-some.
Subsidiuus, a, um—Sinking down, settling.
Subsinuatus, a, um—Somewhat sinuated.
Subsinuosus, a, um—Somewhat sinuous.
Subspatulatus, a, um—Somewhat spatula-shaped.
Subsphericus, a, um—Subspherical.
Subspinulosus, a, um—Somewhat spiny.
Subsplunulosus, a, um—Somewhat covered with small spines.
Substellatus, a, um—Somewhat starred.
Substellatus, a, um—Somewhat finely striated.
Subsulcatus, a, um—Somewhat furrowed.
Subtenuatus, a, um—Somewhat banded.
Subtensus, a, um—Extended underneath, bent.
Subtextilis, e—Somewhat like net-work.
Subtextus, a, um—Woven under, affixed.
Subtilis, e—Fine, thin, slender, delicate.
Subtilitus, a, um—Fine, thin.
Subtilistratus, a, um—Finely striated.
Subtortilis, e—Somewhat twisted.
Subtortuosus, a, um—Somewhat tortuous.
Subtrigona—Somewhat three-angled.
Subtrigonalis, e—Somewhat three-angled, subtrigonal.
Subtruncatus, a, um—Somewhat shortened.
Subtubulatus, a, um—Somewhat pipe or tube formed.
Subtumidus, a, um—Somewhat tumid.
Subturbintus, a, um—Somewhat top-shaped.
Subulatus, a, um—Awl-shaped.
Subumbonatus, a, um—Somewhat protuberant.
Subumbrosus, a, um—Somewhat umbrella-like.
Subundatus, a, um—Somewhat waved.
Subundiferus, a, um—Somewhat wave-bearing.
Subvadius, a, um—Somewhat creeping.
Subvaricosus, a, um—Subvaricose.
Subventricosus, a, um—Subventricose.
Subvesicularis, e—Subvesicular.
Succinctus, a, um—Girded, contracted, succinct.
Succulens—Succulent, sappy.
Sulcatus, a, um—Small-furrowed.
Sulcatus, a, um—Furrowed.
Sulciferus, a, um—Furrow-bearing.
Sulcomarginatus, a, um—Having the margin furrowed.
Sulcopicatus, a, um—Grooved along the middle of the plications.
Superbus, a, um—Superior, excellent, superb.
Superlatus, a, um—Extravagant, excessive, exaggerated.
Supracingulatus, a, um—Encircled or girdled in the upper part.
Supraplanus, a, um—Flat above.
- Surgens**—Rising.
Symmetricus, a, um—Symmetrical.
- Tabulatus**, a, um—Floored, tabulated.
Tæniopteroides—Like *Tæniopteris*.
Tæniopterolideus, a, um—Like *Tæniopteris*.
Tantillus, a, um—So little, such a little thing.
Tapetiformis, e—Formed like tapestry.
Turdus, a, um—Slow, sluggish.
Taxinus, a, um—Like the yew-tree.
Tectorius, a, um—Of or belonging to a cover, rough cast.
Tegulatus, a, um—Tiled, thatched.
Tegulum—A covering, thatch.
Telliniformis, e—Like *Tellina*.
Telum—A dart, spear, or javelin.
Temerarius, a, um—Accidental, casual.
Tenax—Holding fast, gripping, tenacious.
Tenellus, a, um—Somewhat delicate, young.
Tener, era, erum—Delicate, tender, young.
Teneris, e—Delicate.
Tenerrimus, a, um—Very tender, very delicate.
Tenerus, a, um—Tender, delicate.
Tentaculatus, a, um—Having feelers.
Tenuannulatus, a, um—Having slight annulations.
Tenuibrachiatus, a, um—Slender-armed.
Tenuicarinatus, a, um—Finely keeled.
Tenuiceps—Having a slender head.
Tenuicinctus, a, um—Finely girded.
Tenuicornis, e—Slender-horned.
Tenuicostatus, a, um—Fine-ribbed.
Tenuicosta—Having fine ribs or costae.
Tenuicristatus, a, um—Slender-peaked.
Tenuidactylus, a, um—Slender-fingered.
Tenuidens—Having slender teeth.
Tenuidiscus—Having a thin-disk.
Tenuifilum—Fine thread.
Tenuifolius, a, um—Slender-leaved, narrow-leaved.
Tenuilamelatus, a, um—Having thin plates.
Tenuilinentus, a, um—Fine-lined.
Tenuiliratus, a, um—Fine-lined.
Tenuimarginatus, a, um—Thin-margined.
Tenuimuralis, e—Thin-walled.
Tenuinervis, e—Thin-veined, slender-nerved.
Tenuiradiatus, a, um—Slender-rayed.
Tenuiradius—Having slender rays.
Tenuiramosus, a, um—Having slender branches.
Tenuis, e—Thin, fine, slender, narrow.
Tenuisculptus, a, um—Finely engraved.
Tenuisepius, a, um—Having thin septa.
Tenuissimus, a, um—Very thin or slender.
Tenuistriatus, a, um—Fine-lined.
Terebra—A borer, an auger.
Terebralis, e—Like an auger.
Terebriformis, e—Shaped like *Terebra*, or like an auger.
Teres—Rounded, well turned, smooth, polished.
Teretiformis, e—Of a long, round shape.
Terminalis, e—Terminal.
Tersus, a, um—Neat, wiped off, nice.
Tessellatus, a, um—Checkered, tessellated.

metrical.
 , tubulated.
 , lopteris.
 —Like Taniop-
 e, such a little
 ke tapestry.
 ish.
 ew-tree.
 belonging to a
 hatched.
 ch.
 inn.
 uvelin.
 ental, casul.
 ing, tenacious.
 delicate, young.
 e, tender, young.
 tender, very del-
 delicate.
 ing feelers.
 Having slight an-
 Slender-armed.
 finely keeled.
 under head.
 finely girded.
 horned.
 ne-ribbed.
 ribs or costae.
 slender-peaked.
 slender-fingered.
 der teeth.
 thin-disk.
 .
 slender-leaved, nar-
 Having thin plates.
 fine-lined.
 e-lined.
 —Thin-margined.
 alled.
 veined, slender-
 slender-rayed.
 slender rays.
 — Having slender
 ender, narrow.
 finely engraved.
 ying thin septa.
 ry thin or slender.
 fine-lined.
 ager.
 uger.
 ed like Terebra, or
 rned, smooth, pol-
 ng, round shape.
 .
 ped off, nice.
 ckered, tessellated.

Testudinarius, a, um—Arched like a tor-
 toise shell.
 Tetragonocephalus—Having a quadrangular
 head.
 Tetragonophthalmus—Having square eyes.
 Tetragonum—A quadrangle.
 Tetraptix—Having four foli-
 s.
 Tetricus, a, um—Forbidding, stern.
 Textiligerus, a, um—Web-bearing.
 Textilis, e—Woven, plaited, textile.
 Textus, a, um—Woven, fabricated.
 Therodactylus, a, um—Hinge-toed.
 Thallyformis, e—Shaped like Thallus, frond-
 like.
 Tholus—A rotunda or cupola.
 Tiariformis, e—Shaped like a tiara or tur-
 ban.
 Torquis—A necklace, wreath, or ring.
 Tortallinea—Twisted line.
 Tortuosus, a, um—Full of turns or crooks,
 tortuous.
 Tortus, a, um—Twisted, distorted.
 Transiens—Transient.
 Transitionis, e—A passing over.
 Translatus, a, um—Carried over, trans-
 ported.
 Transsectus, a, um—Cut across.
 Transversalis, e—Transverse, crosswise.
 Transversus, a, um—Transverse, crosswise,
 wider than long.
 Triangularis, e—Of or belonging to a triangle,
 triangular.
 Triangulatus, a, um—Triangulated.
 Triarthrus—Having three joints.
 Tribulis—One of the same tribe.
 Tribulosus, a, um—Full of thorns or thistles.
 Tricarinatus, a, um—Three-keeled.
 Tricenarius, a, um—Of or containing thirty.
 Trichoideus, a, um—Hair-like.
 Trichomanoides—Like Trichomanes, the
 maiden-hair fern.
 Tricungulatus, a, um—Three-banded.
 Tricornis, e—Three-horned.
 Tricostatus, a, um—Three-ribbed.
 Tricuspidatus, a, um—Three-pointed.
 Tridactylites—Having three fingers.
 Tridactylus, a, um—Three-fingered.
 Tridens—Having three teeth, times, or
 prongs.
 Tridifterus, a, um—Bearing three teeth
 prongs.
 Tridigitatus, a, um—Three-fingered.
 Trifoliatus, a, um—Three-leaved.
 Trifolius, a, um—Three-leaved.
 Trigonalis, e—Trigonal.
 Trigonolepis—Having triangular scales.
 Trigonostomus, a, um—Having a triangular
 mouth.
 Trigonus, a, um—Trigonal.
 Trilineatus, a, um—Three-lined.
 Triliratus, a, um—Three-lined.
 Trilix—Triple-twilled.
 Trilobatus, a, um—Three-lobed.
 Trilobus, a, um—Three-lobed.
 Trilocularis, e—Three-chambered.
 Trinervis, e—Three-veined.
 Trinodus, a, um—Having three knots.
 Trinucleus—Having three kernels.
 Triplunatus, a, um—Three-winged.

Triplecatellus, a, um—Having three plica-
 tions in one fold.
 Triplicatus, a, um—Three-plaited.
 Triplistriatus, a, um—Three-lined.
 Tripunctatus, a, um—Three-dotted.
 Triquetrus, a, um—Three-cornered, tri-
 angular.
 Triradiatus, a, um—Three-rayed.
 Triserialis, e—In three series.
 Triserratus, a, um—Three-notched.
 Trisinuatus, a, um—Three-furrowed.
 Trisulcatus, a, um—Three-furrowed.
 Trisutura—Having three sutures.
 Trituberculatus, a, um—Having three tu-
 bercles.
 Trivolvus, e—Three-whorled.
 Trochiformis, e—Shaped like Trochus.
 Trochiscus—A small, round ball, a pill.
 Tropidophorus, a, um—Keel-bearing.
 Tradiferus, a, um—Pike-bearing.
 Truncatulus, a, um—Somewhat truncated.
 Truncatus, a, um—Truncated, cut short.
 Tuber—A hump, bump, or protuberance.
 Tuberculatus, a, um—Tuberculated, cov-
 ered with tubercles.
 Tuberculosus, a, um—Full of tubercles.
 Tuberosus, a, um—Full of humps or pro-
 tuberances.
 Tubiformis, e—Pipe, tube, or trumpet-
 formed.
 Tubiporoides—Like Tubipora.
 Tubularis, e—Hollow like a pipe.
 Tubulatus, a, um—Formed like a pipe or
 tube.
 Tubulostriatus, a, um—Having tube-like
 striae.
 Tubulosus, a, um—Abounding in tubes.
 Tubulus—A small pipe or tube.
 Tumidifrons—Swelling front.
 Tumidosus, a, um—High-swelling.
 Tumidulus, a, um—Swollen, tumid.
 Tumidus, a, um—Swollen, tumid.
 Tumulosus, a, um—Full of hills, hilly.
 Tumulus—A mound.
 Tunicatus, a, um—Coated, covered with
 skin or peel.
 Turbidus, a, um—Confused, disordered
 turbid.
 Turbinatus, a, um—Turbinated, cone-shaped.
 Turbiniformis, e—Top-shaped.
 Turgidus, a, um—Swollen, inflated, turgid.
 Turricula—A little tower, a turret.
 Turritella—A little tower.
 Turritiformis, e—Tower-like.
 Turritus, a, um—Fortified with towers.
 Tutus, a, um—Safe, secure, examined.
 Typicalis, e—Typical.
 Typus—The type.
 Tyrans—A tyrant.

Uber—A teat, pap, or udder.
 Umbella—A parasol, umbrella.
 Umbelliferus, a, um—Umbrella-bearing.
 Umbilicatus, a, um—Made like an umbil-
 icus.
 Umbonatus, a, um—Having a shield, em-
 bossed.
 Umbraculum—A shade, umbrella.
 Umbrosus, a, um—Shady, umbrageous.

Uncinatus, a, um—Barbed, furnished with hooks or tenters.

Uncus, a, um—Hook-curved, barbed.

Undans—Waving.

Undatus, a, um—Wavy.

Undosus, a, um—Full of waves, billowy.

Undulatus, a, um—Diversified as with waves, undulated.

Undulostriatus, a, um—Having wavy striae.

Undulosus, a, um—Full of undulations, wavy.

Ungiculus, a, um—Having claw-like processes.

Ungulifer, era, erum—Claw-bearing.

Unguliformis, e—Claw-shaped.

Ungula—A claw, talon, hoof.

Ungulatus, a, um—Having claws or hoofs.

Unguloideus, a, um—Hoof-like or claw-like.

Uniangulatus, a, um—One-angled.

Unicarinatus, a, um—One-keeled.

Unicornis, e—One-horned.

Unicostatus, a, um—One-ribbed.

Unicus, a, um—One and no more, single, sole.

Uniformis, e—Having only one shape, uniform.

Unilargus, a, um—One large, of one size.

Unilobatus, a, um—One-lobed.

Unioniformis, e—Like Unio.

Unionoides—Like Unio.

Uniserialis, e—Having a single row or series.

Unispinus—Having one spine.

Unisulcatus, a, um—Having one furrow.

Unitus, a, um—United.

Ureus, a, um—Of or belonging to the tail.

Urniformis, e—Urn-shaped.

Urophyllus, a, um—Sharp-leaved.

Utriculus—A little matrix, a bud or hull.

Vadosus, a, um—Full of shadows.

Vagens—Wandering, vagrant.

Valens—Vigorous.

Validus, a, um—Strong, powerful.

Vallorus, a, um—Intrenched.

Valvatiformis, e—Like folding doors, or like *Valvata*.

Valvulus—A pod, like the shell of a bean.

Varians—Varying, varied.

Variabilis, e—Changeable, variable.

Varicus, a, um—Straddling.

Varicosus, a, um—Having threads or lines enlarged, varicose.

Varicostatus, a, um—Variably ribbed.

Variolatus, a, um—Variable width or distances apart.

Variolosus, a, um—Full of changes.

Variopora—Having different pores.

Varistriatus, a, um—Having variable striae.

Varius, a, um—Diverse, manifold, different, various.

Varus, a, um—Bent, stretched or grown apart.

Vascularius, a, um—Vascular, consisting of small vessels.

Vasiformis, e—Vase-shaped.

Vastator—A desolator, ravager.

Vaticinus, a, um—Prophetic.

Vellicatus, a, um—Vellicated, pinched.

Velix—Swift, fleet, fitted for motion.

Velutinus, a, um—Velvety.

Velum—A sail, awning, curtain, veil.

Venatus, a, um—Veined.

Venosus, a, um—Full of veins, veiny.

Ventralis, e—Ventral.

Ventricosus, a, um—Bulging out, ventricose.

Venulosus, a, um—Full of small veins.

Venustus, a, um—Lovely, charming.

Venusus, a, um—Lovely, beautiful, graceful.

Verbenifolius, a, um—Leaved like *Verbena*.

Vermicularis, e—Worm-shaped.

Vermiculus—A little worm, grub.

Verrucosus, a, um—Full of warts, rough, rugged.

Versiformis, e—Changing its form, changeable.

Vertebralis, e—Somewhat like vertebrae.

Vertebratus, a, um—Articulated, jointed,

vertebrated, like a backbone.

Verticalis, e—Vertical.

Verticillatus, a, um—Whorled.

Verticillus—The whorl of a spindle.

Verus, a, um—True, real, genuine.

Vesicularis, e—Vesicular.

Vesiculatus, a, um—Vesicled.

Vesiculosus, a, um—Full of blisters or vesicles, vesiculose.

Vesperalis, e—Belonging to the evening.

Vestitus, a, um—Covered, clothed, adorned.

Veterator—One who has grown old.

Vetulus, a, um—Old.

Vetustus, a, um—Old, ancient.

Vexabilis, e—Disturbed, vexed, troublesome.

Viatricus, a, um—Of or belonging to a journey.

Viator—A wayfarer, traveler.

Vicinus, a, um—Near, neighboring, kindred.

Victus, a, um—Conquered, vanquished.

Vigilans—Watchful, vigilant.

Villosus, a, um—Hairy, shaggy, rough.

Viminalis, e—Bearing twigs for plaiting.

Vinctus, a, um—Bounded, fettered, girded.

Vinculatus, a, um—Bound.

Vindex—A defender.

Vinosus, a, um—Full of wine.

Viola—The violet.

Virgatus, a, um—Made of twigs, twig-like.

Virgo—A maid.

Virgosus, a, um—Full of twigs.

Virgulatus, a, um—Striped, like a small rod.

Virguncula—A little maid.

Vittatus, a, um—Bound with a fillet, banded.

Volans—Flying.

Volutus, a, um—Rolled, turned around, whorled.

Vomer—A plowshare.

Vomerium—A plowshare.

Vorax—Ravenous, voracious.

Vorticellatus, a, um—Whorled.

Vulgatus, a, um—General, usual, common.

Xiphias—A sword-fish.

Xylobioides—Like *Xylobius*.

Yoldiiformis, e—Like *Yoldia*.

Zaphrentiformis, e—Shaped like *Zaphrentis*.

Zizac—Slanting in straight lines from side to side, having sharp turns.

Zonatus, a, um—Zoned, belted.

Zonulatus, a, um—Small-girdled.

Zygopus—With joined feet.

veins, veiny.
 ng out, ventricose.
 f small veins.
 ly, charming.
 beautiful, graceful.
 ved like Verbena.
 shaped.
 m, grub.
 of warts, rough,
 its form, change-
 t like vertebrae.
 ticulated, jointed,
 lbone.
 orled.
 f a spindle.
 , genuine.
 ,
 cled.
 of blisters or vesic-
 y to the evening.
 i, clothed, adorned.
 grown old.
 acient.
 vexed, troublesome.
 belonging to a
 veler.
 ighboring, kindred.
 ed, vanquished.
 ilant.
 shaggy, rough.
 wigs for plaiting.
 ed, fettered, girded.
 nd.
 wine.
 of twigs, twig-like.
 f twigs.
 ed, like a small rod.
 ld.
 with a fillet, banded.
 d, turned around,
 re.
 cious.
 orled.
 ral, usual, common.
 obius.
 Yoldia.
 aped like Zaphrentis.
 ight lines from side
 p tins.
 l belted.
 ll-girdled.
 feet.

INDEX OF GENERA.

In addition to alphabetically indexing all the Palaeozoic genera in this work, and placing in italics those which have been used but do not belong to North America, the gender of each genus is designated as follows: *m*, for masculine; *f*, for feminine; *n*, for neuter.

	PAGE.		PAGE.		PAGE.
ACAMBONA, <i>f</i>	332	Ambocelia, <i>f</i>	332	Anthracosia, <i>f</i>	463
Acanthaspis, <i>f</i>	586	Ambonychia, <i>f</i>	460	Antlidus, <i>m</i>	587
Acantherpestes, <i>m</i>	572	Ammonites	432	Apachites, <i>m</i>	529
Acanthocladia, <i>f</i>	291	Amnigenia, <i>f</i>	461	Apedodus, <i>m</i>	587
Acanthoclema, <i>n</i>	291	Ampheristocrinus, <i>m</i>	223	Aphlebia, <i>f</i>	106
Acanthodes, <i>m</i>	586	Amnylespes, <i>m</i>	572	Aplocystites, <i>m</i>	224
Acanthograptus, <i>m</i>	170	Amphibamus, <i>m</i>	618	Arabellites, <i>m</i>	517
Acantholepis, <i>f</i>	586	Amphicelia, <i>f</i>	461	Arachnocrinus, <i>m</i>	224
Acantholoma	525	Amphidema	461	Arachnophyllum, <i>n</i>	172
Acanthophyton, <i>n</i>	104	Amphigenia, <i>f</i>	333	Araucarites, <i>m</i>	106
Acanthotelson, <i>n</i>	525	Amphion, <i>m</i>	527	Arca	463
Acervularia, <i>f</i>	170	Amphipeltis, <i>f</i>	528	Archaeobelus, <i>m</i>	618
Acheloma, <i>n</i>	617	Amphoracrinus, <i>m</i>	223	Archaeocaris, <i>f</i>	529
Acidaspis, <i>f</i>	528	Amplexopora, <i>f</i>	291	Archaeocidaris, <i>f</i>	224
Actis	395	Amplexus, <i>m</i>	171	Archaeocrinus, <i>m</i>	225
Aclisina, <i>f</i>	395	Ampullaria	396	Archaeopteris, <i>f</i>	106
Acondylacanthus, <i>m</i>	586	Ampyx, <i>m</i>	528	Archegogyllus, <i>m</i>	575
Acrocrinus, <i>m</i>	216	Amnydalocystites, <i>m</i>	223	Archaeocyathellus, <i>m</i>	154
Acroculia	395	Anaclitacanthus, <i>m</i>	587	Archaeocyathus, <i>m</i>	154
Acrogenia, <i>f</i>	291	Anarthrocanna, <i>f</i>	105	Archaeoscolex, <i>m</i>	572
Acrolepis, <i>f</i>	587	Anastrophia, <i>f</i>	333	Archimedes, <i>m</i>	292
Acrophylum, <i>n</i>	170	Anatina, <i>f</i>	461	Archimediopora	292
Acrothele, <i>f</i>	332	Anazyga, <i>f</i>	334	Archimylacris, <i>f</i>	575
Acrotreta, <i>f</i>	332	Ancyrocrinus, <i>m</i>	223	Architarbus, <i>n</i>	570
Actinoceras, <i>n</i>	431	Aneimites, <i>m</i>	105	Architulus, <i>m</i>	573
Actinocrinus, <i>m</i>	216	Angelina	528	Arctinurus	530
Actinodema, <i>n</i>	459	Angellum, <i>n</i>	462	Arenicolites, <i>m</i>	517
Actinopteria, <i>f</i>	459	Anisodexis, <i>m</i>	618	Arethusina	530
Actinotrypa, <i>f</i>	291	Anisophyllum, <i>n</i>	172	Arges, <i>m</i>	530
Adiphebia, <i>f</i>	575	Anisotrypa, <i>f</i>	292	Arionellus, <i>m</i>	530
Agilops	332	Annularia, <i>f</i>	105	Aristophycus	106
Aethophlebia, <i>f</i>	575	Anodontopsis, <i>f</i>	462	Aristozoe, <i>f</i>	530
Agaricia	170	Anomocare	528	Arthracantha, <i>f</i>	225
Agaricocrinus, <i>m</i>	220	Anomalocrinus, <i>m</i>	223	Arthracia, <i>f</i>	107
Agassichthys	587	Anomalocystites, <i>m</i>	224	Arthroclema, <i>n</i>	293
Agassizocrinus, <i>m</i>	221	Anomalodontia, <i>f</i>	462	Arthrocyclops, <i>f</i>	570
Agassizodus, <i>m</i>	587	Anomaloides	224	Arthronema, <i>n</i>	293
Agelacrinus, <i>m</i>	221	Anomia	334	Arthropycus, <i>n</i>	107
Agiaspis, <i>f</i>	526	Anomites	334	Arthropora, <i>f</i>	293
Agnostus, <i>m</i>	526	Anomphalus, <i>m</i>	396	Arthrostigma, <i>n</i>	107
Agraulus, <i>m</i>	527	Anopolenus, <i>m</i>	528	Arthrostylus, <i>m</i>	293
Aleclo	291	Anthaspidella, <i>f</i>	153	Artisia, <i>f</i>	107
Alethopteris, <i>f</i>	104	Antholithes, <i>m</i>	106	Asaphiscus, <i>m</i>	530
Allagecrinus, <i>m</i>	222	Anthophylum, <i>n</i>	172	Asaphoidichnus, <i>m</i>	530
Allocrius, <i>m</i>	222	Anthracerpes, <i>m</i>	572	Asaphus, <i>m</i>	530
Allocystites, <i>m</i>	222	Anthracomartus, <i>m</i>	570	Ascoceras, <i>n</i>	432
Alloprosallocrius, <i>m</i>	222	Anthracomya, <i>f</i>	462	Ascodictyon, <i>n</i>	293
Allorisma, <i>n</i>	459	Anthraconectes, <i>m</i>	528	Asolanus	107
Alveolites, <i>m</i>	170	Anthracopalaemon, <i>m</i>	529	Aspidella, <i>f</i>	389
Amacanthus, <i>m</i>	587	Anthracoptera, <i>f</i>	463	Aspidichthys, <i>m</i>	587
Amblypterus, <i>m</i>	587	Anthracopupa, <i>f</i>	396	Aspidocrinus, <i>m</i>	225

IMAGE EVALUATION TEST TARGET (MT-3)

**Photographic
Sciences
Corporation**

23 WEST MAIN STREET
WEBSTER, N.Y. 14560
(716) 872-4503

18 20 22 25
36 32 28

10 01

	PAGE.		PAGE.		PAGE.
<i>Aspidodus</i> , m	588	<i>Beinertia</i> , f	109	<i>Callotrypa</i> , f	296
<i>Aspidopora</i> , f	293	<i>Belemnocrinus</i> , m	229	<i>Calophyllum</i> , n	175
<i>Asplenites</i> , m	107	<i>Belemnosporgia</i> , f	155	<i>Calopodus</i> , m	589
<i>Astarte</i> , f	463	<i>Belinurus</i> , m	534	<i>Calymene</i> , f	536
<i>Astartella</i> , f	463	<i>Bellerophon</i> , m	396	<i>Calypptograptus</i> , m	175
<i>Asteracanthus</i> , m	588	<i>Berenices</i> , f	294	<i>Camarella</i> , f	338
<i>Asterias</i>	226	<i>Bergeria</i>	109	<i>Camarium</i> , n	338
<i>Asterocarpus</i> , m	107	<i>Beyrichia</i> , f	534	<i>Camaroceras</i> , n	432
<i>Asterocrinus</i>	226	<i>Beyrichona</i> , f	535	<i>Camarocladia</i> , f	156
<i>Asterophycus</i> , n	107	<i>Billingsia</i>	338	<i>Camarocrinus</i> , m	230
<i>Asterophyllites</i> , m	107	<i>Billingsia</i>	398	<i>Camaphoria</i> , f	338
<i>Astropteris</i> , f	108	<i>Blastoidocrinus</i> , m	229	<i>Cameratheca</i> , f	389
<i>Asteroptychus</i> , m	588	<i>Blastophycus</i> , n	109	<i>Campanulites</i>	230
<i>Asterosteus</i> , m	588	<i>Blattina</i> , f	575	<i>Campophyllum</i> , n	176
<i>Astraea</i>	172	<i>Blothrophyllum</i> , n	174	<i>Candons</i> , f	537
<i>Astræophyllum</i> , n	172	<i>Blumenbachium</i>	155	<i>Caninia</i>	176
<i>Astræospongia</i> , f	154	<i>Bolboporites</i> , m	174	<i>Canistocrinus</i>	230
<i>Astrios</i>	226	<i>Bolosaurus</i> , m	618	<i>Cannapora</i> , f	176
<i>Astrocerium</i> , n	172	<i>Bornia</i> , f	109	<i>Capulus</i> , m	399
<i>Astroconia</i> , n	154	<i>Rothriolepis</i> , f	588	<i>Carabocrinus</i> , m	230
<i>Astrocrinites</i>	226	<i>Bothrodendron</i> , m	109	<i>Carbonarca</i> , f	468
<i>Astropolithon</i> , m	108	<i>Botryllopora</i> , f	294	<i>Carcharopsis</i> , f	589
<i>Astylocrinus</i>	226	<i>Brachiocrinus</i> , m	229	<i>Cardinia</i> , f	469
<i>Astylospongia</i> , f	154	<i>Brachiospongia</i> , f	155	<i>Cardiocarpon</i> , n	111
<i>Atactopora</i> , f	293	<i>Brachydictes</i> , m	619	<i>Cardiola</i> , f	469
<i>Atactoporella</i> , f	293	<i>Brachymerus</i>	338	<i>Cardiomorpha</i> , f	469
<i>Atazocrinus</i>	226	<i>Brachyphyllum</i> , n	109	<i>Cardiopsis</i> , f	470
<i>Ateleocystites</i> , m	226	<i>Brachyprion</i>	338	<i>Cardiopsis</i> , f	112
<i>Atelestocrinus</i> , m	226	<i>Brongniartia</i>	535	<i>Cardium</i> , n	470
<i>Athyris</i> , f	334	<i>Bronteus</i> , m	535	<i>Carinaropsis</i> , f	399
<i>Atopa</i> , f	532	<i>Bruckmannia</i>	109	<i>Carinopora</i> , f	296
<i>Atrypa</i> , f	335	<i>Bucanella</i> , f	398	<i>Carpolithes</i> , m	112
<i>Atulacodus</i>	517	<i>Bucania</i> , f	398	<i>Caryocrinus</i> , m	231
<i>Aulacophyllum</i> , n	172	<i>Bucanophyllum</i> , n	174	<i>Caryocystites</i>	231
<i>Aulocopina</i> , f	154	<i>Bulimella</i>	399	<i>Caryophyllia</i>	176
<i>Aulophyllum</i> , n	173	<i>Bulimorpha</i> , f	399	<i>Casuarinites</i>	113
<i>Aulopora</i> , f	173	<i>Bumastus</i>	536	<i>Catenipora</i>	176
<i>Aulosteges</i> , m	337	<i>Bunodella</i> , f	536	<i>Catilocrinus</i> , m	231
<i>Avicula</i> , f	463	<i>Bursacrinus</i> , m	229	<i>Caulerites</i> , m	113
<i>Aviculopecten</i> , m	465	<i>Buscopora</i> , f	295	<i>Caulopteris</i> , f	113
<i>Aviculopinna</i> , f	468	<i>Byssopteria</i> , f	468	<i>Caulopora</i> , f	156
<i>Azinura</i>	173	<i>Bythograptus</i> , m	174	<i>Celluloxylon</i> , n	113
<i>Azinus</i>	468	<i>Bythotrephes</i> , f	109	<i>Centrocrinus</i>	231
<i>Azophyllum</i> , n	173	<i>Bvthiacanthus</i> , m	589	<i>Centronella</i> , f	339
		<i>Bythopora</i> , f	295	<i>Cephalaspis</i> , f	589
<i>BACTRITES</i> , m	432			<i>Ceramella</i> , f	296
<i>Bactropora</i> , f	294	<i>Cacabocrinus</i>	229	<i>Ceramopora</i> , f	296
<i>Balera</i> , f	108	<i>Calamites</i> , m	110	<i>Ceramoporella</i> , f	297
<i>Balliella</i> , f	532	<i>Calamocladus</i> , m	110	<i>Ceraterpton</i> , n	619
<i>Bakevella</i> , f	468	<i>Calamodendron</i> , n	110	<i>Ceratiocaris</i> , f	537
<i>Balanocrinus</i>	226	<i>Calamophycus</i> , n	110	<i>Ceratocephala</i>	537
<i>Baphetes</i> , m	618	<i>Calamopora</i>	174	<i>Ceratodus</i> , m	589
<i>Barandia</i> , f	532	<i>Calamostachys</i> , m	110	<i>Ceraurus</i> , m	537
<i>Barandia</i>	532	<i>Calapocia</i> , f	174	<i>Cerlocrinus</i> , m	231
<i>Barycrinus</i> , m	226	<i>Calathium</i> , n	155	<i>Cerionites</i> , m	156
<i>Baryphyllum</i> , n	174	<i>Calathocrinus</i>	230	<i>Ceripora</i> , f	297
<i>Batacanthus</i> , m	588	<i>Calaurops</i> , f	400	<i>Chænocardia</i> , f	470
<i>Bathychilodus</i> , m	588	<i>Calcarina</i> , f	155	<i>Chænomya</i> , f	470
<i>Bathynotus</i> , m	532	<i>Calceocrinus</i> , m	230	<i>Chainodictyon</i> , n	297
<i>Bathyrellus</i> , m	533	<i>Calceola</i> , f	175	<i>Chariocephalus</i> , m	538
<i>Bathyuriscus</i> , m	533	<i>Calcisphæra</i> , f	155	<i>Charionella</i>	339
<i>Bathyrus</i> , m	533	<i>Callipteridium</i> , n	111	<i>Cheliphlebia</i> , f	575
<i>Batocrinus</i> , m	227	<i>Callipteria</i> , f	111	<i>Chemnitzia</i> , f	400
<i>Batospongia</i> , f	154	<i>Calocystites</i> , m	230	<i>Chetetes</i> , m	176
<i>Batostoma</i> , n	294	<i>Callograptus</i> , m	175	<i>Chilonyx</i> , m	619
<i>Batostomella</i> , f	294	<i>Callonema</i> , n	399	<i>Chiloporella</i> , f	297
<i>Batrachia</i> , f	155	<i>Callopora</i> , f	295	<i>Chilotrypa</i> , f	297
<i>Bechera</i> , f	108	<i>Calloporella</i> , f	296	<i>Chirocrinus</i>	231

	PAGE.
<i>f.</i>	296
<i>n.</i>	175
<i>m.</i>	589
<i>f.</i>	536
<i>ptus, m.</i>	175
<i>f.</i>	338
<i>n.</i>	338
<i>ss, n.</i>	432
<i>dia, f.</i>	156
<i>mus, m.</i>	230
<i>oria, f.</i>	338
<i>a, f.</i>	389
<i>les</i>	230
<i>illum, n.</i>	176
<i>f.</i>	537
<i>f.</i>	176
<i>us</i>	230
<i>a, f.</i>	176
<i>sis, f.</i>	399
<i>us, m.</i>	230
<i>a, f.</i>	468
<i>sis, f.</i>	589
<i>f.</i>	469
<i>pon, n.</i>	111
<i>f.</i>	469
<i>rppha, f.</i>	469
<i>s, f.</i>	470
<i>ris, f.</i>	112
<i>n.</i>	470
<i>sis, f.</i>	399
<i>a, f.</i>	296
<i>es, m.</i>	112
<i>us, m.</i>	231
<i>ites</i>	231
<i>llia</i>	176
<i>ites</i>	113
<i>a</i>	176
<i>inus, m.</i>	231
<i>es, m.</i>	113
<i>is, f.</i>	113
<i>ra, f.</i>	156
<i>ylon, n.</i>	113
<i>us</i>	231
<i>ella, f.</i>	339
<i>pis, f.</i>	589
<i>a, f.</i>	296
<i>ora, f.</i>	296
<i>orella, f.</i>	297
<i>ton, n.</i>	619
<i>aris, f.</i>	537
<i>phala</i>	537
<i>as, m.</i>	589
<i>s, m.</i>	537
<i>us, m.</i>	231
<i>es, m.</i>	156
<i>a, f.</i>	297
<i>ardia, f.</i>	470
<i>hya, f.</i>	470
<i>ictyon, n.</i>	297
<i>ephalus, m.</i>	538
<i>lla</i>	339
<i>lebia, f.</i>	575
<i>tsia, f.</i>	400
<i>s, m.</i>	176
<i>x, m.</i>	619
<i>rella, f.</i>	297
<i>pa, f.</i>	297
<i>us</i>	231

	PAGE.
<i>Chirodus, m.</i>	589
<i>Chirolepis, f.</i>	589
<i>Chirurus</i>	538
<i>Chirosporgia, f.</i>	156
<i>Chirotherium, n.</i>	619
<i>Chiton, m.</i>	400
<i>Chitonodus, m.</i>	590
<i>Chloephycus</i>	113
<i>Cholaster, f.</i>	231
<i>Cholodus, m.</i>	590
<i>Chomatodus, m.</i>	590
<i>Chondrites, m.</i>	114
<i>Chonetes, m.</i>	339
<i>Chonophyllum, n.</i>	177
<i>Chonostegites, m.</i>	177
<i>Chrestotes, f.</i>	575
<i>Cimataria, f.</i>	472
<i>Cladodus, m.</i>	590
<i>Cladograptus, m.</i>	178
<i>Cladopora, f.</i>	178
<i>Clathrocellia, f.</i>	389
<i>Clathropora, f.</i>	297
<i>Cleodictya, f.</i>	157
<i>Clepsydrops, f.</i>	619
<i>Clidophorus, m.</i>	470
<i>Clidocrinus, m.</i>	231
<i>Climactichnites, m.</i>	538
<i>Climacograptus, m.</i>	178
<i>Climaxodus, m.</i>	591
<i>Clinopiatha, f.</i>	472
<i>Clodermia</i>	389
<i>Clisiophyllum, n.</i>	179
<i>Cliospora, f.</i>	400
<i>Clonograptus, m.</i>	179
<i>Clonopora, f.</i>	298
<i>Clotrocrinus, m.</i>	232
<i>Clymenia</i>	432
<i>Cnemidium, n.</i>	157
<i>Cococrinus, m.</i>	232
<i>Cocosteus, m.</i>	591
<i>Cochliodus, m.</i>	591
<i>Cocytinus, m.</i>	619
<i>Codaster, m.</i>	232
<i>Codonites, m.</i>	233
<i>Codonochilus, n.</i>	400
<i>Coelacanthus, m.</i>	592
<i>Coelocrinus, m.</i>	233
<i>Coelocaulis, f.</i>	298
<i>Coeloconus, m.</i>	298
<i>Crochirus, m.</i>	233
<i>Coelospora, f.</i>	340
<i>Conites, m.</i>	179
<i>Conostroma, n.</i>	157
<i>Coleolus, m.</i>	389
<i>Coleophyllum, n.</i>	179
<i>Coleoprius, m.</i>	390
<i>Colletosaurus, m.</i>	620
<i>Colosteus, m.</i>	620
<i>Colpocaris, f.</i>	539
<i>Colpoceras, n.</i>	432
<i>Columnaria, f.</i>	179
<i>Columnospira, f.</i>	180
<i>Comarocystites, m.</i>	233
<i>Combophyllum, n.</i>	180
<i>Compasacanthus, n.</i>	592
<i>Compasaster, m.</i>	234
<i>Compocrinus, m.</i>	234
<i>Conchicolites, m.</i>	517

	PAGE.
<i>Conchiopsis</i>	592
<i>Conchodus, m.</i>	592
<i>Conchopeltis, f.</i>	400
<i>Conchites</i>	432
<i>Conocardium, n.</i>	472
<i>Conocephalites, m.</i>	539
<i>Conocerac, n.</i>	432
<i>Conocoryphe, f.</i>	539
<i>Conocrinus</i>	234
<i>Conophyllum</i>	180
<i>Conopterium, n.</i>	157
<i>Conostichus, m.</i>	114
<i>Conotubularia</i>	432
<i>Constellaria</i>	180
<i>Conularia, f.</i>	390
<i>Conulites</i>	432
<i>Copodus, m.</i>	592
<i>Cordaianthus, m.</i>	114
<i>Cordaicarpus, m.</i>	114
<i>Cordaistrobilus, m.</i>	114
<i>Cordaites, m.</i>	114
<i>Cordylocrinus, m.</i>	234
<i>Cornulites, m.</i>	517
<i>Coronocrinus, m.</i>	234
<i>Coronura</i>	540
<i>Corycephalus</i>	540
<i>Coscinnella, f.</i>	298
<i>Cosciniun, n.</i>	298
<i>Coscinopora</i>	157
<i>Coscinotrypa, f.</i>	298
<i>Cotyledonocrinus, m.</i>	234
<i>Crania, f.</i>	341
<i>Craspedophyllum, n.</i>	180
<i>Crateripora</i>	298
<i>Cremacrinus</i>	234
<i>Crenatopteris, f.</i>	115
<i>Crenipecten, m.</i>	473
<i>Crepicephalus, m.</i>	540
<i>Crepidophyllum, n.</i>	180
<i>Crepidopora, f.</i>	299
<i>Cricotus, m.</i>	620
<i>Crinocystites, m.</i>	234
<i>Crisosoma</i>	234
<i>Crisina, f.</i>	299
<i>Crisinella, f.</i>	299
<i>Cromyocrinus, m.</i>	234
<i>Crumenacrinites</i>	234
<i>Cruziana, f.</i>	115
<i>Cryphaeus</i>	540
<i>Cryptoblastus</i>	234
<i>Cryptoceras</i>	432
<i>Cryptolithus</i>	540
<i>Cryptonella, f.</i>	342
<i>Cryptopora, f.</i>	299
<i>Cryptozoon, n.</i>	157
<i>Ctenacanthus, m.</i>	592
<i>Ctenocrinus, m.</i>	234
<i>Ctenodonta</i>	473
<i>Ctenodus, m.</i>	593
<i>Ctenopetalus, m.</i>	594
<i>Ctenoptychius, m.</i>	594
<i>Cucullia</i>	473
<i>Cuneamya, f.</i>	473
<i>Cupellacrinus</i>	235
<i>Cupulocrinus, m.</i>	235
<i>Cyathaxonia, f.</i>	180
<i>Cyathocrinus, m.</i>	235

	PAGE.
<i>Cyathophycus, n.</i>	158
<i>Cyathophyllum, n.</i>	181
<i>Cyathopora</i>	182
<i>Cyathospongia, f.</i>	158
<i>Cybele</i>	540
<i>Cyclaster</i>	267 237.
<i>Cycloconcha, f.</i>	474
<i>Cyclocystoides, m.</i>	237
<i>Cyclograptus, m.</i>	182
<i>Cyclolites, m.</i>	183
<i>Cyclonema, n.</i>	400
<i>Cyclopore, f.</i>	299
<i>Cycloporella, f.</i>	299
<i>Cyclopteris, f.</i>	115
<i>Cyclora, f.</i>	401
<i>Cyclostigma, n.</i>	116
<i>Cyclostoma</i>	401
<i>Cylindrocellia, f.</i>	158
<i>Cyclus</i>	541
<i>Cymatodus, m.</i>	594
<i>Cymoglossa, f.</i>	116
<i>Cyphaspis, f.</i>	541
<i>Cypriacardia, f.</i>	474
<i>Cypriocardia, f.</i>	475
<i>Cypriocardinia, f.</i>	475
<i>Cypriocardites, m.</i>	476
<i>Cyrtacanthus, m.</i>	594
<i>Cyrtia, f.</i>	342
<i>Cyrtina, f.</i>	342
<i>Cyrtoceras, n.</i>	432
<i>Cyrtocerina, f.</i>	436
<i>Cyrtodonta</i>	477
<i>Cyrtolites, m.</i>	401
<i>Cyrtoneura, f.</i>	402
<i>Cystiphorolites, m.</i>	183
<i>Cystiphyllum, n.</i>	183
<i>Cystocrinus, m.</i>	237
<i>Cystodictya, f.</i>	299
<i>Cystopora, f.</i>	300
<i>Cystostylus, m.</i>	184
<i>Cythere, f.</i>	541
<i>Cythereella, f.</i>	541
<i>Cytherellina, f.</i>	541
<i>Cytherina, f.</i>	541
<i>Cytherodon, m.</i>	477
<i>Cytheropsis, f.</i>	541
<i>Cytocrinus, m.</i>	237
<i>Dactylocrinus, m.</i>	594
<i>Dactylophycus, n.</i>	116
<i>Dadoxylon, n.</i>	116
<i>Daemocrinus</i>	237
<i>Dalmanites, m.</i>	542
<i>Dalmanites, m.</i>	542
<i>Daneites, m.</i>	116
<i>Dania, f.</i>	184
<i>Dawsonella, f.</i>	402
<i>Dawsonia, f.</i>	184
<i>Decadactylocrinites</i>	237
<i>Decadocrinus</i>	237
<i>Decenia, f.</i>	116
<i>Dekayella, f.</i>	184
<i>Dekayia, f.</i>	184
<i>Deltacrinus, m.</i>	237
<i>Elatodopsis, f.</i>	595
<i>Elatodus, m.</i>	595
<i>Delthyris</i>	343

	PAGE.		PAGE.		PAGE.
<i>Deltotychius</i> , m . . .	595	<i>Dolichocephala</i> . . .	545	<i>Eozoon</i> , n . . .	159
<i>Denderpeton</i> , n . . .	620	<i>Dolichocephalus</i> , m . . .	545	<i>Ephemerites</i> , m . . .	576
<i>Dendrocrinus</i> , m . . .	238	<i>Dolichopterus</i> , m . . .	545	<i>Epicordylus</i> , m . . .	621
<i>Dendrograptus</i> , m . . .	184	<i>Donacrinites</i> . . .	240	<i>Equisetites</i> , m . . .	118
<i>Dendrophycus</i> , n . . .	117	<i>Dorycrinus</i> , m . . .	240	<i>Equisetum</i> . . .	118
<i>Dendropora</i> , f . . .	185	<i>Drepanacanthus</i> , m . . .	596	<i>Eremopteris</i> , f . . .	118
<i>Dentalina</i> , f . . .	158	<i>Drepanodus</i> , m . . .	518	<i>Eretmocrinus</i> , m . . .	242
<i>Dentalium</i> , n . . .	402	<i>Duncanella</i> , f . . .	187	<i>Eridophyllum</i> , n . . .	187
<i>Desmodus</i> , m . . .	595	<i>Dyscritus</i> , m . . .	576	<i>Eridopora</i> , f . . .	301
<i>Desmiophyllum</i> , n . . .	117	<i>Dystactella</i> , f . . .	478	<i>Erismacanthus</i> , m . . .	597
<i>Dexiobia</i> , f . . .	478	<i>Dystactophycus</i> , n . . .	117	<i>Erisocrinus</i> , m . . .	243
<i>Diadectes</i> , m . . .	620	<i>Dystactospongia</i> , f . . .	158	<i>Eryops</i> , f . . .	622
<i>Diamesopora</i> , f . . .	300	<i>EATONIA</i> , f . . .	345	<i>Eschara</i> , f . . .	301
<i>Dicelloccephalus</i> , m . . .	543	<i>Eccyliomphalus</i> , m . . .	402	<i>Escharopora</i> , f . . .	301
<i>Dicellomus</i> . . .	343	<i>Echinocaris</i> , f . . .	545	<i>Esteria</i> , f . . .	547
<i>Dichocrinus</i> , m . . .	239	<i>Echinocystites</i> , m . . .	241	<i>Etmophyllum</i> , n . . .	159
<i>Dichograptus</i> . . .	185	<i>Echinodiscus</i> , m . . .	241	<i>Etblattina</i> , f . . .	576
<i>Dichotrypa</i> , f . . .	300	<i>Echinocrinites</i> , m . . .	241	<i>Eucalyptocrinus</i> , m . . .	243
<i>Dicraniscus</i> . . .	343	<i>Echinognathus</i> , m . . .	546	<i>Euchasma</i> , n . . .	480
<i>Dicranograptus</i> , m . . .	185	<i>Echinus</i> . . .	241	<i>Euchondria</i> , f . . .	480
<i>Dicranophyllum</i> , n . . .	117	<i>Ectenocrinus</i> , m . . .	242	<i>Eucladocrinus</i> , m . . .	244
<i>Dicranopora</i> , f . . .	300	<i>Ectenodesma</i> , n . . .	478	<i>Eugaster</i> , m . . .	244
<i>Dicranurus</i> . . .	544	<i>Ectenodictyon</i> , f . . .	159	<i>Eulima</i> . . .	403
<i>Dictyocrinus</i> , m . . .	240	<i>Ectocynodon</i> , m . . .	621	<i>Eumetria</i> , f . . .	346
<i>Dictyolites</i> . . .	117	<i>Ectosteorachis</i> , m . . .	596	<i>Eumicrotia</i> . . .	480
<i>Dictyonema</i> , n . . .	185	<i>Edaphosaurus</i> , m . . .	621	<i>Eunema</i> , n . . .	403
<i>Dictyophyton</i> , n . . .	158	<i>Edestus</i> , m . . .	597	<i>Eunemacanthus</i> , m . . .	597
<i>Dictyopteria</i> , f . . .	117	<i>Edmondia</i> , f . . .	478	<i>Eunicites</i> , m . . .	518
<i>Dictyostroma</i> , n . . .	158	<i>Edrioaster</i> , m . . .	242	<i>Eumomphalus</i> , m . . .	403
<i>Didymograptus</i> , m . . .	185	<i>Edriocrinus</i> , m . . .	242	<i>Eupachycrinus</i> , m . . .	245
<i>Didymopora</i> , f . . .	576	<i>Edriospongia</i> , f . . .	159	<i>Eupherbia</i> , f . . .	573
<i>Didymophyllum</i> , n . . .	117	<i>Eichwaldia</i> , f . . .	345	<i>Euproops</i> , f . . .	547
<i>Dieconeura</i> , f . . .	576	<i>Eliticus</i> , m . . .	573	<i>Eurydictya</i> , f . . .	301
<i>Dignomia</i> , f . . .	343	<i>Elasacrinus</i> . . .	242	<i>Eurylepis</i> , f . . .	597
<i>Dimetrodon</i> , m . . .	620	<i>Elasmophyllum</i> , n . . .	187	<i>Eurypterella</i> , f . . .	548
<i>Dinichthys</i> , m . . .	596	<i>Eleutherocrinus</i> , m . . .	242	<i>Eurypterus</i> , m . . .	548
<i>Dinobolus</i> , m . . .	343	<i>Elkania</i> , f . . .	346	<i>Eurythorax</i> , m . . .	622
<i>Dionide</i> , f . . .	544	<i>Ellipsocephalus</i> , m . . .	546	<i>Eusarcus</i> , m . . .	549
<i>Dipeltis</i> , m . . .	544	<i>Elliptocephala</i> , f . . .	546	<i>Euspiropora</i> , f . . .	301
<i>Diphyphyllum</i> , n . . .	186	<i>Elonichthys</i> . . .	597	<i>Euspirocrinus</i> , m . . .	246
<i>Diplazites</i> . . .	117	<i>Elymella</i> , f . . .	480	<i>Eusthenopteron</i> , m . . .	597
<i>Diplocaura</i> . . .	544	<i>Elymocarid</i> , f . . .	546	<i>Euthydesma</i> , n . . .	480
<i>Diplichnites</i> , m . . .	544	<i>Embolamus</i> , m . . .	547	<i>Evactinopora</i> , f . . .	301
<i>Diplocaulus</i> , m . . .	621	<i>Embolophorus</i> , m . . .	621	<i>Exochorhynchus</i> . . .	480
<i>Diploceras</i> , n . . .	436	<i>Emmonsia</i> , f . . .	187	<i>FABRIA</i> , f . . .	549
<i>Diploclema</i> , n . . .	300	<i>Empedias</i> , m . . .	621	<i>Favastrea</i> . . .	187
<i>Diplodus</i> , m . . .	596	<i>Empedocles</i> , m . . .	621	<i>Favicella</i> , f . . .	301
<i>Diplograptus</i> , m . . .	186	<i>Enallopora</i> , f . . .	300	<i>Faviphyllum</i> , n . . .	187
<i>Diplophyllum</i> . . .	187	<i>Encrinurus</i> , m . . .	576	<i>Favistella</i> , f . . .	188
<i>Diplopore</i> , f . . .	300	<i>Encrinurus</i> , m . . .	547	<i>Favosites</i> , m . . .	188
<i>Diplostegium</i> , n . . .	117	<i>Endoceras</i> , n . . .	436	<i>Favositopora</i> . . .	189
<i>Diplostylus</i> , m . . .	544	<i>Endolobus</i> . . .	437	<i>Fenestella</i> , f . . .	302
<i>Diplothea</i> , f . . .	391	<i>Endothyra</i> , n . . .	159	<i>Fenestralia</i> , f . . .	305
<i>Diplotrypa</i> , f . . .	187	<i>Endymion</i> . . .	547	<i>Fenestrapora</i> , f . . .	305
<i>Dipterocaris</i> , f . . .	545	<i>Endymionia</i> , f . . .	547	<i>Ficoidites</i> . . .	118
<i>Dipterus</i> , m . . .	596	<i>Enoploura</i> , f . . .	547	<i>Filicites</i> . . .	118, 189
<i>Discina</i> , f . . .	344	<i>Entolium</i> , n . . .	480	<i>Fissodus</i> , m . . .	597
<i>Disclites</i> . . .	436	<i>Eocidarid</i> , f . . .	242	<i>Fistulipora</i> , f . . .	305
<i>Discolites</i> . . .	402	<i>Eocystites</i> , m . . .	242	<i>Flabellaria</i> . . .	118
<i>Discophycus</i> , n . . .	117	<i>Eodon</i> , m . . .	480	<i>Flustra</i> , f . . .	306
<i>Discophyllum</i> , n . . .	187	<i>Eophyton</i> , n . . .	118	<i>Forbesiocrinus</i> , m . . .	246
<i>Discosorus</i> , m . . .	438	<i>Eopteria</i> , f . . .	480	<i>Fordilla</i> , f . . .	480
<i>Discotrypa</i> , f . . .	300	<i>Eosaurus</i> , m . . .	621	<i>Fucoides</i> . . .	118
<i>Distacodus</i> , m . . .	518	<i>Eoscorpius</i> , m . . .	570	<i>Fungispongia</i> , f . . .	160
<i>Dithyrocaris</i> , f . . .	545	<i>Eospongia</i> , f . . .	159	<i>Fusispira</i> , f . . .	404
<i>Dolabra</i> , f . . .	478	<i>Eotrochus</i> , m . . .	403	<i>Fusillina</i> , f . . .	160
<i>Dolatocrinus</i> , m . . .	240	<i>Eotrophonia</i> . . .	518	<i>Fusus</i> . . .	404

	PAGE.
.....	159
.....	576
.....	821
.....	118
.....	118
.....	118
.....	242
.....	187
.....	301
.....	597
.....	243
.....	622
.....	301
.....	301
.....	547
.....	159
.....	576
.....	243
.....	480
.....	480
.....	244
.....	244
.....	403
.....	348
.....	480
.....	403
.....	597
.....	518
.....	403
.....	245
.....	573
.....	547
.....	301
.....	597
.....	548
.....	548
.....	622
.....	549
.....	301
.....	246
.....	597
.....	480
.....	301
.....	480
.....	549
.....	187
.....	301
.....	137
.....	188
.....	188
.....	189
.....	302
.....	305
.....	305
.....	118
.....	118, 189
.....	597
.....	305
.....	118
.....	306
.....	246
.....	480
.....	118
.....	180
.....	404
.....	180
.....	404

	PAGE.
<i>Galium</i>	119
<i>Gampsacanthus</i> , <i>m</i>	598
<i>Gaurocrinus</i> , <i>m</i>	248
<i>Genetomnum</i> , <i>n</i>	577
<i>Gennæocrinus</i> , <i>m</i>	247
<i>Genopteryx</i> , <i>f</i>	577
<i>Geoporia</i>	189
<i>Gerablattina</i> , <i>f</i>	577
<i>Geralinura</i> , <i>f</i>	571
<i>Gerapompis</i> , <i>m</i>	577
<i>Gerarus</i> , <i>m</i>	577
<i>Geraplyurus</i> , <i>m</i>	571
<i>Gerephemera</i> , <i>f</i>	577
<i>Geroneura</i> , <i>f</i>	577
<i>Gervillia</i> , <i>f</i>	481
<i>Gissacanthus</i> , <i>m</i>	598
<i>Glauconome</i> , <i>f</i>	307
<i>Glossites</i> , <i>m</i>	481
<i>Glossoceras</i> , <i>n</i>	437
<i>Glossograptus</i> , <i>m</i>	189
<i>Glossotrypa</i> , <i>f</i>	307
<i>Glycerites</i> , <i>m</i>	518
<i>Glymmatacanthus</i> , <i>m</i>	598
<i>Glyptaster</i> , <i>m</i>	247
<i>Glyptocardia</i> , <i>f</i>	481
<i>Glyptocrinus</i> , <i>m</i>	247
<i>Glyptocystites</i> , <i>m</i>	248
<i>Glyptodendron</i> , <i>n</i>	119
<i>Glyptodesma</i> , <i>n</i>	481
<i>Glyptolepis</i> , <i>f</i>	598
<i>Glyptotrypa</i>	307
<i>Gnathorhiza</i> , <i>f</i>	598
<i>Gomphoceras</i> , <i>n</i>	437
<i>Gomphocystites</i> , <i>m</i>	249
<i>Goniasteroidocrinus</i> , <i>m</i>	249
<i>Goniasterites</i> , <i>m</i>	438
<i>Gonioceras</i> , <i>n</i>	441
<i>Goniocella</i>	348
<i>Goniophora</i> , <i>f</i>	481
<i>Gonioteria</i>	119
<i>Goniotrypa</i> , <i>f</i>	307
<i>Gordia</i>	119-518
<i>Gorgonia</i> , <i>f</i>	307
<i>Gosseletia</i> , <i>f</i>	482
<i>Grammysia</i> , <i>f</i>	482
<i>Granatocrinus</i> , <i>m</i>	250
<i>Graphocrinus</i> , <i>m</i>	251
<i>Graptodictya</i> , <i>f</i>	307
<i>Graptolithus</i> , <i>m</i>	19
<i>Gryphorhynchus</i>	483
<i>Gullemites</i> , <i>m</i>	119
<i>Gypidia</i> , <i>f</i>	343
<i>Gyracanthus</i> , <i>m</i>	598
<i>Gyrichnites</i> , <i>m</i>	518
<i>Gyroceras</i> , <i>n</i>	441
<i>HADROCRINUS</i> , <i>m</i>	252
<i>Hadrophylum</i> , <i>n</i>	191
<i>Haimenophylum</i> , <i>n</i>	191
<i>Hallia</i> , <i>f</i>	191
<i>Halionia</i> , <i>f</i>	119
<i>Halysites</i> , <i>m</i>	191
<i>Halysocrinus</i>	252
<i>Haplocrinus</i> , <i>m</i>	252
<i>Haplophlebum</i> , <i>n</i>	577
<i>Haplotichnus</i> , <i>m</i>	578
<i>Harlania</i> , <i>f</i>	119
<i>Harmodites</i>	191

	PAGE.
<i>Harpacodus</i> , <i>m</i>	598
<i>Harpes</i> , <i>m</i>	549
<i>Harpides</i> , <i>m</i>	549
<i>Hartia</i> , <i>f</i>	549
<i>Harumannia</i>	548
<i>Helicopora</i> , <i>f</i>	308
<i>Helicotoma</i> , <i>f</i>	405
<i>Heliodus</i> , <i>m</i>	599
<i>Heliolites</i> , <i>m</i>	191
<i>Heliophycus</i> , <i>n</i>	119
<i>Heliophyllum</i> , <i>n</i>	192
<i>Helminthoidichnites</i> , <i>m</i>	518
<i>Helodectes</i> , <i>m</i>	622
<i>Helodus</i> , <i>m</i>	599
<i>Helopora</i> , <i>f</i>	308
<i>Hemeristia</i> , <i>f</i>	578
<i>Hemicosmites</i> , <i>m</i>	252
<i>Hemicrypturus</i>	549
<i>Hemicystites</i> , <i>m</i>	252
<i>Hemipronites</i>	346
<i>Hemitrypa</i> , <i>f</i>	309
<i>Hernodia</i> , <i>f</i>	309
<i>Heterocrinus</i> , <i>m</i>	252
<i>Heterocystites</i> , <i>m</i>	253
<i>Heterodictya</i> , <i>f</i>	309
<i>Heterophrenis</i> , <i>f</i>	193
<i>Heterospongia</i> , <i>f</i>	180
<i>Heteroschisma</i> , <i>n</i>	253
<i>Heterotrypa</i>	193
<i>Hindella</i> , <i>f</i>	346
<i>Hindia</i> , <i>f</i>	180
<i>Hippocrinus</i>	346
<i>Hippodophycus</i> , <i>n</i>	120
<i>Hipponicharion</i> , <i>m</i>	549
<i>Hippothoa</i>	309
<i>Holocystites</i> , <i>m</i>	253
<i>Holometopus</i> , <i>m</i>	550
<i>Holopea</i> , <i>f</i>	405
<i>Holopella</i> , <i>f</i>	406
<i>Holoptychius</i> , <i>m</i>	599
<i>Homacanthus</i>	600
<i>Homalonotus</i> , <i>m</i>	550
<i>Homocrinus</i> , <i>m</i>	255
<i>Homothetus</i> , <i>m</i>	578
<i>Homotrypa</i> , <i>f</i>	309
<i>Homotrypella</i> , <i>f</i>	310
<i>Hornera</i> , <i>f</i>	310
<i>Hortholus</i>	442
<i>Houghtonia</i>	193
<i>Huronia</i> , <i>f</i>	442
<i>Hybochinus</i> , <i>m</i>	255
<i>Hybocladodus</i> , <i>m</i>	606
<i>Hybocrinus</i> , <i>m</i>	255
<i>Hybocystites</i> , <i>m</i>	255
<i>Hydnoceras</i>	442
<i>Hydreionocrinus</i> , <i>m</i>	256
<i>Hylerpeton</i> , <i>n</i>	622
<i>Hydonomus</i> , <i>m</i>	622
<i>Hydræonophyllites</i> , <i>m</i>	120
<i>Hyphasma</i> , <i>n</i>	622
<i>Hyolithellus</i> , <i>m</i>	391
<i>Hyolithes</i> , <i>m</i>	391
<i>Hypanthocrinites</i>	256
<i>Hystericrinus</i>	256
<i>Hystrispongia</i> , <i>f</i>	160
<i>ICHNOPHYCUS</i> , <i>n</i>	120
<i>Ichthyacanthus</i> , <i>m</i>	622

	PAGE.
<i>Ichthyocrinus</i> , <i>m</i>	256
<i>Ichthyorachis</i> , <i>f</i>	310
<i>Icosidactylocrinites</i>	258
<i>Idiophyllum</i> , <i>n</i>	120
<i>Idiotrypa</i> , <i>f</i>	310
<i>Illionia</i> , <i>f</i>	483
<i>Illænurus</i> , <i>m</i>	550
<i>Illænus</i> , <i>m</i>	550
<i>Inachus</i>	406
<i>Inocaulis</i> , <i>m</i>	193
<i>Inoceramus</i>	483
<i>Intrapora</i> , <i>f</i>	310
<i>Intricaria</i> , <i>f</i>	310
<i>Iocrinus</i> , <i>m</i>	258
<i>Iphidea</i> , <i>f</i>	346
<i>Ischadites</i>	160
<i>Ischyrinia</i> , <i>f</i>	483
<i>Isocardia</i>	483
<i>Isocillina</i> , <i>f</i>	551
<i>Isonema</i> , <i>n</i>	406
<i>Isotelus</i> , <i>n</i>	552
<i>Isotrypa</i> , <i>f</i>	310
<i>JANASSA</i> , <i>f</i>	600
<i>KNORRIA</i> , <i>f</i>	120
<i>Koninekia</i> , <i>f</i>	348
<i>Kutorgina</i> , <i>f</i>	347
<i>LABECHIA</i> , <i>f</i>	310
<i>Lambdodus</i> , <i>m</i>	600
<i>Lamellipora</i> , <i>f</i>	194
<i>Lampterocrinus</i> , <i>m</i>	257
<i>Lasiocladia</i> , <i>f</i>	160
<i>Leaia</i> , <i>f</i>	552
<i>Lecanocrinus</i> , <i>m</i>	552
<i>Lecracanthus</i> , <i>m</i>	600
<i>Lecythocrinus</i>	257
<i>Leda</i>	484
<i>Lepadocrinus</i> , <i>m</i>	257
<i>Lepeditia</i> , <i>f</i>	552
<i>Lepetopsis</i> , <i>f</i>	406
<i>Lepidechinus</i> , <i>m</i>	258
<i>Lepidesthes</i> , <i>f</i>	258
<i>Lepidilla</i> , <i>f</i>	553
<i>Lepiditta</i> , <i>f</i>	553
<i>Lepidocidaris</i> , <i>f</i>	258
<i>Lepidocoleus</i> , <i>m</i>	553
<i>Lepidocystis</i> , <i>m</i>	121
<i>Lepidodendron</i> , <i>n</i>	121
<i>Lepidodiscus</i>	258
<i>Lepidolites</i> , <i>m</i>	160
<i>Lepidophloios</i> , <i>m</i>	122
<i>Lepidophyllum</i> , <i>n</i>	123
<i>Lepidostrobos</i> , <i>m</i>	123
<i>Lepidoxylon</i> , <i>n</i>	124
<i>Lepocrinites</i>	258
<i>Leptacanthus</i>	600
<i>Leptæna</i> , <i>f</i>	347
<i>Leptobolus</i> , <i>m</i>	348
<i>Lepoecelia</i> , <i>f</i>	348
<i>Leptodesma</i> , <i>n</i>	484
<i>Leptodomus</i> , <i>f</i>	485
<i>Leptomitus</i> , <i>m</i>	160
<i>Leptophloeum</i> , <i>n</i>	124
<i>Leptophractus</i> , <i>m</i>	623
<i>Leptopora</i> , <i>f</i>	194
<i>Leptopterion</i> , <i>n</i>	161

	PAGE.		PAGE.		PAGE.
Leptotrypa, f.	311	Macrodon, m.	487	Myrianites, m.	519
Lescuropteris, f.	124	Macropetalichthys, m.	601	Myrtillocrinus, m.	261
Lesleya, f.	125	Macrostachya, f.	126	Mytilarca, f.	493
Libellula, f.	573	Macrostylocrinus, m.	259	Mytilops, f.	494
Lichas, m.	553	Madrepora	195	Mytilus, m.	494
Lichenalia, f.	311	Malocystites, m.	259		
Lichenocrinus, m.	258	Mariacrinus, m.	259	Naiadites	494
Lichenotrypa, f.	312	Marracanthus, m.	601	Natica	413
Licroryphus, n.	125	Marsupiocrinus, m.	260	Naticopsis, f.	413
Lima, f.	486	Martinia, f.	352	Nautilus, m.	443
Limaria, f.	194	Matheria, f.	488	Nebulipora, f.	198
Limoptera, f.	486	Mathevia, f.	392	Necymylacris, n.	579
Lindstromia, f.	194	Mazonia, f.	571	Nelimenia	445
Lingula, f.	349	Mecolepis, f.	602	Nemagraptus, m.	198
Lingulelasma, n.	351	Meekella, f.	353	Nemapodia, f.	519
Lingulella, f.	352	Meekopora, f.	312	Nemataxis, m.	313
Lingulepis, f.	352	Megalaspis, f.	555	Nematophycus, n.	127
Lingulops, f.	352	Megalagraptus, m.	195	Nematophyllum, n.	127
Limpora	194	Megalomus, m.	488	Nematopora, f.	313
Annarsonia, f.	352	Megalopteris, f.	126	Nematoxylon, n.	127
Liolena, n.	310	Megambonia, f.	488	Nephropteris	127
Liodus, m.	600	Meganteris	353	Nereidavus, m.	519
Liognathus, m.	600	Megaphyton, n.	126	Nereites, m.	519
Lioptra, f.	484	Megaptera	489	Nerioptris, f.	127
Liorhynchus, n.	347	Megastroma, n.	161	Neuropteris, f.	127
Liostracus, m.	555	Megathentomum, n.	579	Nicholsonella, f.	313
Lisogocaris	555	Megistocrinus, m.	260	Nileus, m.	557
Lisgodus, m.	601	Melia	443	Nipterocrinus, m.	262
Listracanthus, m.	601	Melocrinus, m.	261	Nodosinella, f.	161
Lithentomum, n.	578	Melonites, m.	261	Næggerathia, f.	129
Lithodictyon	125	Menocephalus, m.	556	Nothodon, m.	623
Lithomylacris, n.	579	Merista, f.	353	Nothozoe	557
Lithophaga, f.	486	Meristella, f.	353	Nucleocrinus, m.	262
Lithostrotion, n.	194	Meristina, f.	354	Nucleospira, f.	355
Litorina	406-486	Mesodmodus, m.	602	Nucula, f.	494
Lituites, m.	442	Meonacis, f.	556	Nuculana, f.	495
Loculipora, f.	312	Mesothyrta, f.	556	Nuculites, m.	496
Loftusia, f.	161	Mespilocrinus, m.	261	Nulipora	161
Loganellus, m.	555	Mefarmosaurus, m.	623	Nuttainia	557
Lonchocephalus, m.	555	Metoptoma, f.	409	Nyassa, f.	496
Lonchopteris, f.	125	Miamia, f.	579	Nyctopora, f.	198
Lonsdalia, f.	194	Michelinia, f.	196		
Lophodus	601	Microceras, n.	410	OBOLELLA, f.	355
Lophophyllum, n.	194	Microcylus, m.	196	Obolellina	355
Lophospira, f.	406	Microdiscus, m.	557	Obolus, m.	355
Loxonema, n.	406	Microdoma, f.	410	Odontocephalus	557
Lucina, f.	486	Microdon	489	Odontochile	557
Lumbriconereites, m.	518	Microspongia, f.	161	Odontopteris, f.	129
Lunaipora, f.	195	Millepora	196	Odontotrypa, f.	313
Lunulicardium, n.	486	Mitoclema, n.	313	Oenites, m.	519
Lunulites	161	Modiola, f.	489	Oestoecephalus, m.	623
Lycopodiolites	125	Modiolopsis, f.	489	Ogygia, f.	557
Lycopodites, m.	125	Modiomorpha, f.	491	Oldhamia, f.	198
Lyellia, f.	195	Møllerina, f.	161	Olenellus, m.	557
Lyonsia	487	Molgophis, m.	623	Olenoides, m.	557
Lyriocrinus, m.	258	Monocraterion, n.	519	Olenus, m.	558
Lyriodictya, f.	161	Monograptus, m.	196	Oligocarpia, f.	130
Lyriopecten, m.	487	Monomerella, f.	354	Oligoporus, m.	263
Lyrodesma, n.	487	Monopteria, f.	492	Olivianites	263
Lyropora, f.	312	Monotis, f.	492	Ollacrinus	263
Lysocestites, m.	259	Monotrypa	196	Omphalotrochus, m.	413
Lysocephalus, m.	623	Monotrypella, f.	196	Omphyma, f.	198
		Monticulipora, f.	197	Onchus	602
MACHÆRACANTHUS, m.	601	Murchisonia, f.	410	Oncoceras, n.	445
Machairoodus	519	Myalina, f.	492	Onychaster, m.	264
Maclurea, f.	408	Myelodactylus, m.	261	Onychocrinus, m.	264
Macrochilus, n.	408	Myclerope, f.	602	Onychodus, m.	602
Macrochilina, f.	409	Mylacris, n.	579	Ophiacodon, m.	624

PAGE.		PAGE.		PAGE.		PAGE.	
519	<i>Ophieta, f.</i>	413	<i>Paleschara, f.</i>	313	<i>Phyllodictya, f.</i>	315	
261	<i>Opisthoptera</i>	497	<i>Palmacites</i>	131	<i>Phyllograptus, m.</i>	200	
493	<i>Oracanthus, m.</i>	603	<i>Panenka, f.</i>	500	<i>Phyllopora, f.</i>	315	
494	<i>Orbicula</i>	356	<i>Panopæa</i>	500	<i>Phylloporina</i>	315	
494	<i>Orbiculoidea, f.</i>	356	<i>Pantylus, m.</i>	624	<i>Phyllopteris, f.</i>	133	
	<i>Orbitulites</i>	161	<i>Paolia, f.</i>	579	<i>Physotocrinus, m.</i>	269	
494	<i>Ormathichnus, m.</i>	414	<i>Paracardium, n.</i>	500	<i>Physonemus, m.</i>	606	
413	<i>Ormoceras, n.</i>	445	<i>Paracyclas, f.</i>	500	<i>Physophycus, n.</i>	133	
413	<i>Ormoxyton, n.</i>	130	<i>Paradoxidæ, m.</i>	559	<i>Physospongia, f.</i>	163	
443	<i>Ormithichnites, m.</i>	624	<i>Pararca, f.</i>	500	<i>Phytolithus</i>	133	
198	<i>Orodus, m.</i>	603	<i>Pariotichus, m.</i>	624	<i>Phytopsis, f.</i>	134	
579	<i>Orthacanthus, m.</i>	604	<i>Parioxya, m.</i>	624	<i>Pileopsis</i>	415	
445	<i>Orthis, f.</i>	356	<i>Pariocrinus</i>	267	<i>Pileotrypa, f.</i>	315	
198	<i>Orthisina, f.</i>	360	<i>Paromyiacris, n.</i>	580	<i>Piliolites</i>	561	
519	<i>Orthoceras, n.</i>	445	<i>Pasceolus, m.</i>	162	<i>Piloceras, n.</i>	453	
313	<i>Orthodesma, n.</i>	497	<i>Patella, f.</i>	415	<i>Pinacotrypa, f.</i>	315	
127	<i>Orthogoniopteris, f.</i>	130	<i>Pattersonia, f.</i>	163	<i>Pinna, f.</i>	502	
127	<i>Orthonema, n.</i>	414	<i>Pecten</i>	131	<i>Pinnatopora</i>	315	
313	<i>Orthonota, f.</i>	497	<i>Pecton</i>	501	<i>Pinnopsis</i>	502	
127	<i>Orthonotella, f.</i>	498	<i>Pelion, m.</i>	624	<i>Pinnularia, f.</i>	134	
127	<i>Orthonychia</i>	414	<i>Peltodus, m.</i>	605	<i>Pisocrinus, m.</i>	269	
519	<i>Orthopleurodus, m.</i>	604	<i>Pellura</i>	560	<i>Piscinopsis, f.</i>	502	
519	<i>Orthopora, f.</i>	313	<i>Pemphigaspis, f.</i>	560	<i>Plangtichnus, m.</i>	580	
127	<i>Orthostoma, n.</i>	414	<i>Pentacrinites</i>	267	<i>Planolites</i>	520	
127	<i>Ortonia</i>	519	<i>Pentagonia, f.</i>	360	<i>Planorbis</i>	415	
313	<i>Oryctoblattina, f.</i>	579	<i>Pentagonites</i>	267	<i>Plasmopora, f.</i>	200	
557	<i>Oryctocephalus, m.</i>	558	<i>Pentamerella, f.</i>	360	<i>Plataphemera, f.</i>	580	
262	<i>Ostrea</i>	498	<i>Pentamerus, m.</i>	361	<i>Playceras, n.</i>	415	
161			<i>Pentremites, m.</i>	267	<i>Platycrinus, m.</i>	270	
129			<i>Peplorhina, f.</i>	605	<i>Platycystites, m.</i>	272	
623	<i>PACHYCRINUS, m.</i>	265	<i>Perichocrinus</i>	269	<i>Platyodus, m.</i>	606	
557	<i>Pachydictya, f.</i>	313	<i>Periplectrodus, m.</i>	605	<i>Platymotus</i>	561	
262	<i>Pachylocrinus</i>	265	<i>Peripristis, m.</i>	605	<i>Platyschisma, n.</i>	418	
355	<i>Pachyphyllum</i>	130	<i>Pernachactas</i>	501	<i>Platysomus, m.</i>	607	
494	<i>Pachyphyllum, n.</i>	198	<i>Pernopecten, m.</i>	501	<i>Platystoma, n.</i>	418	
495	<i>Pachypora, f.</i>	199	<i>Peronopora, f.</i>	199	<i>Platystrophia</i>	332	
496	<i>Pachypteris, f.</i>	130	<i>Petalichnus, m.</i>	452	<i>Plasmopora, f.</i>	200	
161	<i>Palaecis, f.</i>	161	<i>Petalodus, m.</i>	605	<i>Plectambonites</i>	362	
557	<i>Palaecacmea, f.</i>	414	<i>Petalorhynchus, n.</i>	606	<i>Plectrotylus</i>	418	
496	<i>Palaeanatina, f.</i>	498	<i>Petalotrypa, f.</i>	314	<i>Plethomytilus, m.</i>	503	
198	<i>Palaearca</i>	498	<i>Petigopora, f.</i>	314	<i>Pleuracanthus, m.</i>	607	
	<i>Palaesaspis, f.</i>	604	<i>Petrablattina, f.</i>	580	<i>Pleurocystites, m.</i>	272	
355	<i>Palaester, m.</i>	266	<i>Petraia, f.</i>	199	<i>Pleurodictyum, n.</i>	200	
355	<i>Palaesterina, f.</i>	266	<i>Petraster, m.</i>	269	<i>Pleuronotus, m.</i>	419	
355	<i>Palaechinus, m.</i>	604	<i>Petrodus, m.</i>	606	<i>Pleurophorus, m.</i>	503	
557	<i>Palaebatis, m.</i>	573	<i>Phacelopora, f.</i>	314	<i>Pleuroptyx, f.</i>	625	
557	<i>Palaecocampa, f.</i>	498	<i>Phacops, f.</i>	560	<i>Pleurorhynchus</i>	504	
129	<i>Palaecocardia, f.</i>	558	<i>Phænopora, f.</i>	314	<i>Pleurotomaria, f.</i>	419	
313	<i>Palaecocaris, f.</i>	498	<i>Phæthonides, m.</i>	560	<i>Plicatula</i>	362	
519	<i>Palaecoconcha, f.</i>	519	<i>Phaneropleuron, n.</i>	606	<i>Plumulina, f.</i>	134	
623	<i>Palaechorda, f.</i>	266	<i>Phanerotinus, m.</i>	415	<i>Plumulites, m.</i>	561	
557	<i>Palaecocoma, f.</i>	267	<i>Pharetrella, f.</i>	392	<i>Plungeacanthus, m.</i>	607	
198	<i>Palaecocrinus, m.</i>	559	<i>Phæbodius, m.</i>	606	<i>Poecilodus, m.</i>	607	
557	<i>Palaecocrisia, f.</i>	199	<i>Phillipsia, f.</i>	560	<i>Poliocera, f.</i>	571	
557	<i>Palaecocylus, m.</i>	267	<i>Phillipsastrea, f.</i>	199	<i>Polygonites</i>	453	
558	<i>Palaecocystites, m.</i>	161	<i>Philocrinus</i>	269	<i>Polydylasma</i>	200	
130	<i>Palaecomano, f.</i>	499	<i>Philegethontia, f.</i>	624	<i>Polyernus, m.</i>	580	
263	<i>Palaeonello, f.</i>	604	<i>Pholadella, f.</i>	502	<i>Polygnathus, m.</i>	520	
263	<i>Palaeoniscus, m.</i>	555	<i>Pholadomya</i>	502	<i>Polyphemopsis, f.</i>	423	
263	<i>Palaecopalemon, m.</i>	130	<i>Pholidocidaris, m.</i>	269	<i>Polypora, f.</i>	315	
413	<i>Palaephyeus, n.</i>	199	<i>Pholidops, f.</i>	362	<i>Polyporites, m.</i>	134	
198	<i>Palaephyllum, n.</i>	499	<i>Phractopora, f.</i>	315	<i>Polyrhizodus, m.</i>	607	
602	<i>Palaepinna, f.</i>	131	<i>Phragmoceras, n.</i>	452	<i>Polyspora</i>	134	
445	<i>Palaopteris</i>	500	<i>Phragmodictya, f.</i>	163	<i>Porambonites, m.</i>	362	
264	<i>Palaesolen, m.</i>	162	<i>Phragmolites</i>	415	<i>Porcellia, f.</i>	424	
264	<i>Palaesopongia, f.</i>	199	<i>Phragmostoma, n.</i>	415	<i>Portes</i>	200	
602	<i>Palaotrochis</i>	414	<i>Phthanocoris, f.</i>	580	<i>Porocrinus, m.</i>	273	
624	<i>Palaotrochus, m.</i>	131	<i>Phthonia, f.</i>	502	<i>Posidonia, f.</i>	504	
	<i>Palaoxyria, f.</i>						

	PAGE.		PAGE.		PAGE.
<i>Posidonomya</i> , <i>f</i>	504	<i>Ptychocrinus</i> ,	276	<i>Saccammina</i> , <i>f</i>	184
<i>Poterocrinus</i> , <i>m</i>	273	<i>Ptychodesma</i> , <i>n</i>	508	<i>Saccocrinus</i> , <i>m</i>	278
<i>Præcardium</i> , <i>n</i>	504	<i>Ptychonema</i> , <i>n</i>	201	<i>Saccospongia</i> , <i>f</i>	104
<i>Præopora</i> , <i>f</i>	201	<i>Ptychoparia</i> , <i>f</i>	564	<i>Særichnites</i> , <i>m</i>	453
<i>Prestwichia</i> ,	561	<i>Ptychophyllum</i> , <i>n</i>	201	<i>Sagenaria</i> ,	138
<i>Primitia</i> , <i>f</i>	561	<i>Ptychopteria</i> , <i>f</i>	508	<i>Sagenella</i> , <i>f</i>	321
<i>Prioniodus</i> , <i>m</i>	520	<i>Ptyctodus</i> , <i>m</i>	610	<i>Salterella</i> , <i>f</i>	520
<i>Prionotus</i> ,	201	<i>Ptyonius</i> , <i>m</i>	625	<i>Salteria</i> ,	566
<i>Prisonalia</i> , <i>f</i>	504	<i>Ptyonodus</i> , <i>m</i>	610	<i>Sandalodus</i> , <i>m</i>	612
<i>Prismopora</i> , <i>f</i>	317	<i>Pugiunculus</i> ,	392	<i>Sanguinolaria</i> , <i>f</i>	509
<i>Pristicladodus</i> , <i>m</i>	608	<i>Pupa</i> , <i>f</i>	424	<i>Sanguinolites</i> , <i>m</i>	509
<i>Pristodus</i> , <i>m</i>	608	<i>Pyanomya</i> , <i>f</i>	509	<i>Sao</i> ,	566
<i>Productella</i> , <i>f</i>	362	<i>Pycnocrinus</i> , <i>m</i>	276	<i>Saportæa</i> , <i>f</i>	138
<i>Productus</i> , <i>m</i>	363	<i>Pycnostylus</i> , <i>m</i>	202	<i>Sarcinula</i> , <i>f</i>	203
<i>Proetus</i> , <i>m</i>	561	<i>Pygopterus</i> , <i>m</i>	611-625	<i>Sauripteria</i> ,	612
<i>Promacrus</i> , <i>m</i>	504	<i>Pygorhynchus</i> , <i>n</i>	277	<i>Sauropleura</i> , <i>f</i>	625
<i>Promylacris</i> , <i>n</i>	580	<i>Pyrenomæus</i> , <i>m</i>	509	<i>Sauropus</i> , <i>m</i>	625
<i>Propeticus</i> , <i>m</i>	581			<i>Sceavogyra</i> , <i>f</i>	425
<i>Prorynchus</i> , <i>n</i>	504	<i>Quenstedtia</i> ,	202	<i>Scalaripora</i> , <i>f</i>	321
<i>Proscorpius</i> ,	571			<i>Scalites</i> , <i>m</i>	425
<i>Protæra</i> , <i>f</i>	201	<i>RAMIPORA</i> , <i>f</i>	320	<i>Scaphiocrinus</i> , <i>m</i>	279
<i>Protaster</i> , <i>m</i>	276	<i>Raniceps</i> ,	625	<i>Scenella</i> , <i>f</i>	392
<i>Protasterina</i> ,	276	<i>Raphistoma</i> , <i>n</i>	424	<i>Scenellopora</i> , <i>f</i>	322
<i>Prothyris</i> , <i>f</i>	504	<i>Rastrites</i> , <i>m</i>	202	<i>Sceptropora</i> , <i>f</i>	322
<i>Protichnites</i> , <i>m</i>	562	<i>Rauffella</i> , <i>f</i>	163	<i>Schizambon</i> ,	370
<i>Protoblechnum</i> , <i>n</i>	134	<i>Receptaculites</i> , <i>m</i>	163	<i>Schizoblastus</i> , <i>m</i>	281
<i>Protocaris</i> , <i>f</i>	563	<i>Remopleurides</i> , <i>m</i>	565	<i>Schizobolus</i> , <i>m</i>	370
<i>Protoerisina</i> , <i>f</i>	317	<i>Renssæleria</i> , <i>f</i>	366	<i>Schizocrania</i> , <i>f</i>	371
<i>Protocyathus</i> , <i>m</i>	163	<i>Reptaria</i> , <i>f</i>	320	<i>Schizocrinus</i> , <i>m</i>	281
<i>Protograptus</i> , <i>m</i>	201	<i>Retepora</i> , <i>f</i>	320	<i>Schizodus</i> , <i>m</i>	510
<i>Protollimulus</i> , <i>m</i>	563	<i>Reteporina</i> , <i>f</i>	320	<i>Schizodiscus</i> , <i>m</i>	566
<i>Protomya</i> , <i>f</i>	505	<i>Retiocrinus</i> , <i>m</i>	277	<i>Schizopteria</i> ,	138
<i>Protoscolex</i> , <i>m</i>	520	<i>Retiograptus</i> , <i>m</i>	202	<i>Schenaster</i> , <i>m</i>	281
<i>Protopongia</i> , <i>f</i>	163	<i>Retiolites</i> , <i>m</i>	202	<i>Schutzia</i> , <i>f</i>	139
<i>Protostigma</i> , <i>f</i>	135	<i>Retzia</i> , <i>f</i>	366	<i>Scoliostruma</i> , <i>n</i>	425
<i>Prototaxites</i> , <i>m</i>	135	<i>Rhabdaria</i> , <i>f</i>	164	<i>Scolithus</i> , <i>m</i>	520
<i>Prototypus</i> , <i>m</i>	563	<i>Rhabdichnites</i> ,	566	<i>Scolopendrites</i> , <i>m</i>	139
<i>Prontella</i> , <i>f</i>	318	<i>Rhabdocarpus</i> , <i>m</i>	136	<i>Scyphia</i> ,	164
<i>Psammodus</i> , <i>m</i>	608	<i>Rhachiopteria</i> , <i>f</i>	137	<i>Scyphocrinus</i> , <i>m</i>	281
<i>Psaronius</i> , <i>m</i>	135	<i>Rhachitonus</i> , <i>m</i>	625	<i>Scythalocrinus</i> ,	281
<i>Psæphodus</i> , <i>m</i>	610	<i>Rhachura</i> , <i>f</i>	566	<i>Sedgwickia</i> , <i>f</i>	511
<i>Pseudocrania</i> , <i>f</i>	366	<i>Rhacophyllum</i> , <i>n</i>	137	<i>Selaginites</i> , <i>m</i>	139
<i>Pseudomonotis</i> , <i>f</i>	505	<i>Rhadinichthys</i> , <i>m</i>	611	<i>Selenoides</i> ,	164
<i>Pseudopocopter a</i> , <i>f</i>	135	<i>Rhynidietya</i> , <i>f</i>	320	<i>Selenopora</i> , <i>f</i>	322
<i>Pseudophorus</i> , <i>m</i>	424	<i>Rhniopora</i> , <i>f</i>	321	<i>Semicoscinium</i> , <i>n</i>	322
<i>Pallophyton</i> , <i>n</i>	136	<i>Rhinocaris</i> , <i>f</i>	566	<i>Septopora</i> , <i>f</i>	322
<i>Pterichthys</i> , <i>m</i>	610	<i>Rhizodus</i> , <i>m</i>	611	<i>Serpula</i> , <i>f</i>	521
<i>Pterinea</i> , <i>f</i>	505	<i>Rhizograptus</i> , <i>m</i>	202	<i>Serpulites</i> , <i>m</i>	521
<i>Pterinopecten</i> , <i>m</i>	507	<i>Rhizolites</i> ,	138	<i>Shumardia</i> , <i>f</i>	566
<i>Pterocephalus</i> , <i>n</i> , <i>f</i>	564	<i>Rhizomopteria</i> , <i>f</i>	138	<i>Sicarius</i> ,	612
<i>Pteronaulinus</i> , <i>m</i>	453	<i>Rhizomorpha</i> , <i>f</i>	138	<i>Sidamina</i> ,	453
<i>Pteronitella</i> , <i>f</i>	507	<i>Rhodocrinus</i> , <i>m</i>	277	<i>Sigillaria</i> , <i>f</i>	139
<i>Pteronites</i> , <i>m</i>	507	<i>Rhombodictyon</i> , <i>n</i>	164	<i>Sigillarioides</i> , <i>m</i>	141
<i>Pteropora</i> ,	318	<i>Rhombopora</i> , <i>f</i>	321	<i>Sigillariostrobus</i> , <i>m</i>	141
<i>Pterotheca</i> , <i>f</i>	392	<i>Rhopalonaria</i> , <i>f</i>	321	<i>Siphonia</i> ,	164
<i>Pterocrinus</i> , <i>m</i>	276	<i>Rhynchodus</i> , <i>m</i>	611	<i>Siphonocrinus</i> , <i>m</i>	281
<i>Pterygotus</i> , <i>f</i>	564	<i>Rhynchonella</i> , <i>f</i>	367	<i>Siphonotreta</i> , <i>f</i>	371
<i>Ptilocarpus</i> , <i>m</i>	136	<i>Rhynchospira</i> ,	370	<i>Skenidium</i> , <i>n</i>	371
<i>Ptilonella</i> ,	318	<i>Rhynchotreta</i> , <i>f</i>	370	<i>Smithia</i> , <i>f</i>	203
<i>Ptilodictya</i> , <i>f</i>	318	<i>Rhynobolus</i> ,	370	<i>Solenocaris</i> , <i>f</i>	567
<i>Ptilograptus</i> , <i>m</i>	201	<i>Ribeiria</i> , <i>f</i>	566	<i>Solarium</i> ,	425
<i>Ptilonaster</i> ,	276	<i>Rinodus</i> ,	612	<i>Solemya</i> ,	512
<i>Ptilophyton</i> ,	136	<i>Romingeria</i> , <i>f</i>	203	<i>Solen</i> , <i>m</i>	512
<i>Ptilopora</i> , <i>f</i>	319	<i>Rotaria</i> , <i>f</i>	164	<i>Soleniscus</i> , <i>m</i>	425
<i>Ptiloporella</i> , <i>f</i>	319	<i>Rotella</i> , <i>f</i>	425	<i>Solenocaris</i> ,	566
<i>Ptiloporina</i> , <i>f</i>	319	<i>Rotularia</i> ,	138	<i>Solenochilus</i> , <i>n</i>	453
<i>Ptilotrypa</i> , <i>f</i>	320	<i>Rusichnites</i> , <i>m</i>	566	<i>Solenomya</i> , <i>f</i>	512
<i>Ptychaspis</i> , <i>f</i>	564	<i>Rusophycus</i> , <i>n</i>	138	<i>Solenopleura</i> , <i>f</i>	567

PAGE.		PAGE.		PAGE.		PAGE.	
164	Solenopsis, f.	512	Streptosolen, m.	165	Thallistigma, n.	327	
278	Solenoula, f.	141	Streptospongia, f.	165	Thamniscus, m.	327	
164	Sorocladus, m.	141	Striatopora, f.	205	Thamnolictya, f.	166	
453	Spathella, f.	513	Stricklandia, f.	379	Thamnograptus, m.	207	
138	Spathiocaris, f.	507	Stricklandina, f.	379	Thamnopora, f.	328	
321	Spatiopora, f.	323	Strigillina, f.	612	Thamnotrypa, f.	328	
520	Sphaerexochus, m.	567	Strobilocystites, m.	283	Theca, f.	393	
566	Sphaerocoryphe, f.	567	Strobilospongia, f.	165	Thecia, f.	207	
612	Sphaerocrinus, m.	282	Strobilus, m.	146	Thecostegites, m.	207	
509	Sphaerocystites, m.	282	Stromatocerium, n.	165	Thenaropus, m.	620	
509	Sphaerolites, m.	203	Stromatopora, f.	165	Theropleura, f.	626	
566	Sphagodus, m.	521	Strombodes, m.	205	Thoracodus, m.	613	
138	Sphenacodon, m.	626	Strophalosia, f.	379	Thrinacodus, m.	613	
203	Sphenolum, n.	513	Strophites, m.	427	Thyrasidium, n.	626	
612	Sphenophyllum, n.	141	Strophodonta, f.	379	Thysanocrinus, m.	286	
625	Sphenopteris, f.	142	Strophomena, f.	381	Titanophasma, f.	581	
625	Sphenopterium, m.	164	Strophonella, f.	383	Tomodus, m.	613	
425	Sphenothallus, m.	143	Strophostylus, m.	427	Trachomaticnus, m.	454	
321	Sphenotus, m.	513	Strotocrinus, m.	283	Trachydoria, n.	428	
425	Spheropezium, n.	626	Strotopora, f.	326	Trachypora, f.	207	
279	Sphragiopora, f.	323	Stylaster, f.	205	Trachyum, n.	166	
392	Spirangium, n.	144	Stylifer, m.	427	Trematodus, m.	428	
322	Spirifer, f.	371	Styliola, f.	392	Tremataster, m.	286	
322	Spiriferina, f.	377	Stylcnurus, m.	568	Trematella, f.	328	
370	Spirigera, m.	377	Subretepora, f.	326	Trematis, f.	385	
281	Spirodorus, m.	514	Subulites, m.	427	Trematoceras, n.	454	
370	Spirophyton, n.	144	Sulcopora, f.	326	Trematocrinus, m.	287	
371	Spirorbis, m.	521	Symphylurus, m.	568	Trematodiscus, m.	455	
281	Spirula, m.	454	Synbathocrinus, m.	284	Trematopora, f.	328	
510	Spongia, m.	165	Syncladia, f.	326	Trematospira, f.	385	
566	Sporangites, m.	144	Syntrielsma, n.	383	Treptichnus, m.	581	
138	Sporocystis, m.	144	Syngocorinus, m.	285	Triacrinus, m.	287	
281	Squamaster, m.	282	Syringodendron, n.	146	Triarthrella, f.	568	
139	Staphylopteris, f.	144	Syringolites, m.	205	Triarthrus, m.	568	
425	Stauraccephalites, m.	521	Syringophyllum, m.	166	Trichiulus, m.	573	
520	Stauroraptus, m.	203	Syringopora, f.	206	Trichomanites, m.	147	
139	Steganoecrinus, m.	282	Syringostroma, n.	166	Trichophycus, n.	147	
164	Stellipora, f.	203	Syringothyris, f.	384	Trichospongia, f.	166	
281	Stemmatocrinus, m.	282	Syringoxylon, n.	146	Tricelocrinus, m.	287	
281	Stemmatodus, m.	612	Tæniaster, m.	285	Trigonocarpus, n.	147	
511	Stemmatopteris, f.	144	Tæniodictya, f.	327	Trigonodus, m.	613	
139	Stenacanthus, m.	612	Tæniodus, m.	612	Trigonotreta, f.	386	
164	Stenaster, m.	282	Tæniophyllum, n.	146	Trimerella, f.	386	
322	Stenocrinus, m.	283	Tæniopora, f.	327	Trimerorhachis, f.	626	
322	Stenopora, f.	203	Tæniopteris, f.	146	Trimerus, m.	568	
322	Stenopterodus, m.	612	Tæniopora, f.	327	Trinucleus, m.	568	
521	Stenoschisma, n.	377	Talarocrinus, m.	285	Triphyllopteris, f.	148	
521	Stenotheca, f.	392	Tanaodus, m.	612	Triplesia, f.	387	
566	Stephanocrinus, m.	283	Tanurus, m.	146	Trochila, f.	428	
612	Stereocrinus, m.	283	Taxocrinus, m.	285	Trochoceras, n.	455	
453	Sternbergia, f.	145	Technocrinus, m.	285	Trocholites, m.	456	
139	Stictopora, f.	323	Technophorus, m.	514	Trochonema, n.	428	
141	Stictoporella, f.	325	Teleocrinus, m.	285	Trochophyllum, n.	207	
141	Stictotrypa, f.	325	Telephus, m.	568	Trochus, m.	428	
164	Stigmara, f.	145	Tellina, m.	514	Troostocrinus, m.	287	
281	Stigmarioides, m.	145	Tellinomya, f.	514	Tropidoleptus, m.	387	
371	Stigmatopora, f.	325	Tellinopsis, f.	515	Tropidopora, f.	329	
371	Straparollina, f.	426	Temnochilus, n.	454	Tryblidium, n.	429	
203	Straparollus, m.	426	Tentaculites, m.	393	Tuberculopora, f.	329	
567	Streblopteris, f.	514	Terataspis, f.	568	Tubipora, f.	207	
425	Streblotrypa, f.	325	Teratichnus, m.	454	Tuditatus, m.	626	
512	Strephochetus, m.	165	Terebratula, f.	384	Turbo, m.	429	
512	Strephodes, m.	204	Terebratulites, m.	385	Turbonilla, m.	429	
425	Streptaxis, m.	427	Termes, m.	581	Turripelas, m.	569	
566	Streptelasma, n.	204	Tetradium, n.	206	Turritella, f.	429	
453	Streptoceras, n.	454	Tetragraptus, m.	206			
512	Strobilepis, f.	468	Textularia, f.	166			
567	Streptorhynchus, n.	378	Thalops, f.	568	ULODENDRON, n.	148	
					Ungulina,	515	

	PAGE.		PAGE.		PAGE.
<i>Unio</i> ,	515	<i>Vitulina</i> , <i>f</i>	387	<i>Xystracanthus</i> , <i>m</i>	614
<i>Unitrypa</i> ,	329	<i>Volkmannia</i> , <i>m</i>	148	<i>Xystrodus</i> , <i>m</i>	614
<i>Uphantænia</i> , <i>f</i>	168				
<i>VALVULINA</i> , <i>f</i>	167	<i>WALCHIA</i> , <i>f</i>	148	<i>YOLDIA</i> , <i>f</i>	516
<i>Vanuxemia</i> , <i>f</i>	515	<i>Walcottia</i> , <i>f</i>	522		
<i>Vasocrinus</i> , <i>m</i>	287	<i>Waldheimia</i> , <i>f</i>	387	<i>ZAPHRENTIS</i> , <i>f</i>	208
<i>Vaticinodus</i> , <i>m</i>	613	<i>Whittleseyia</i> , <i>f</i>	148	<i>Zaptychius</i> , <i>m</i>	430
<i>Venus</i> ,	518	<i>Worthenopora</i> , <i>f</i>	330	<i>Zatrachys</i> , <i>n</i>	627
<i>Venustodus</i> , <i>m</i>	613	<i>XENOCRINUS</i> , <i>m</i>	287	<i>Zeacrinus</i> , <i>m</i>	288
<i>Vermipora</i> , <i>f</i>	207	<i>Xenoneura</i> , <i>f</i>	581	<i>Zittellelia</i> , <i>f</i>	167
<i>Vertumnia</i> , <i>f</i>	516	<i>Xenophora</i> , <i>f</i>	430	<i>Zonites</i> , <i>m</i>	430
<i>Vesicularia</i> , <i>f</i>	208	<i>Xylobius</i> , <i>m</i>	574	<i>Zygospira</i> , <i>f</i>	388

s, m	614
	614
	516
	208
	430
	627
	288
	167
	430
	388

FIRST APPENDIX, 1892.

As THIS work was intended for the use of beginners in the pursuit of geological and palaeontological information, as well as for the most advanced students, it has been urged that I should have accented the technical words, with a view of bringing about correct and uniform pronunciation.

This criticism may be well taken, but there are only a few words commonly mispronounced, and a few examples will suffice to correct the pronunciation of most of these.

Words ending in *ceras*, *crinus*, *pora*, and *lepis* are accented on the antepenult, as *Orthoc'eras*, *Cyrtoc'eras*, *Gomphoc'eras*, *Actinoc'rinus*, *Xenoc'rinus*, *Platyc'rinus*, *Monticulip'ora*, *Leptop'ora*, *Chirol'epis*, *Bothriol'epis*, etc. Words ending in *ites* have the *i* long and the accent on the penult, as *Cypricardi'tes*, *Dalmani'tes*, *Favosi'tes*, *Litui'tes*, *Trocholi'tes*, etc. Words ending in *nema*, *mena*, etc., have the accent on the penult, as *Cyclone'ma*, *Loxone'ma*, *Strophome'na*, *Calyme'ne*, etc. We say *Ath'yris*, *Cari'na*, *Cerat'odus*, *Cœlentera'ta*, *Epithe'ca*, *Onych'odus*, *Palæaspis*, and *Syringoth'yris*.

I have seen no reason to change the established nomenclature of the Groups of rocks as set forth in the geological part of this work, notwithstanding there may be some who apply the word Cambrian to rocks indiscriminately from the Taconic to the Devonian. It seems to be a word that is easily pasted over ignorance, and some use it for that reason.

VEGETABLE KINGDOM.

THERE have been very few fossil plants described since the publication of this work.

BYTHOTREPHIS pergracilis, Dawson, 1889, Trans. Roy. Soc. Can., vol. 7, p. 54, Up. Taconic.

Cruziana carleyi, James, 1885, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 155. Not recognized.

DACTYLOIDES bulbosus, Hall, 1886, 39th Rep. N. Y. St. Mus. Nat. Hist., p. 160. Obscure markings supposed to represent marine algae, graptolites, or traces of a spongoid substance.

DICRANOPHYLLUM. Type *D. gallicum*.

LEPIDODENDRON cliftonense, Dr. 1891, Bull. G. S. A., vol. 2, p. 52. Sur. murrayanum, Dawson, 1891, circle G. S. A., vol. 2, p. 532, Coal Meas.

Nematophyllum was preoccupied by McCoy for a genus of corals in 1851, and hence could not be used again by Fontaine & White as a generic name. I now propose for *Nematophyllum* of Fontaine & White in Permian or Upper Carboniferous Flora of West Virginia and S. W. Pennsylvania, p. 35, the generic name *Nematophyllites* with *N. angustus*, Fontaine & White as the type.

RUSOPHYCUS, from its etymology, should be spelled *Rhysophycus*.

acadicum, Dawson, 1861 (*Rusichnites acadicus*), Can. Nat. and Geol., vol. 1, p. 363, and Acad. Geol., p. 410, Coal Meas.

- carbonarium, Dawson, 1868 (Rusichnites carbonarius), Acad. Geol., p. 257, Carboniferous.
- clintonense, Dawson, 1890 (Rusichnites clintonensis), Quar. Jour. Geol. Soc., vol. 46, p. 598, Clinton Gr.
- grenvillense, Dawson, 1890 (Rusichnites grenvillensis), Quar. Jour. Geol. Soc., vol. 46, p. 598, Up. Taconic.
- SPHENOPHYLLUM vetustum, Newberry, 1889, Jour. Ctn. Soc. Nat. Hist., vol. 12, p. 55, Up. Held. Gr.
- SPHENOPTERIS salisburyi, Lesquereux, 1887, Franklin Soc. Rep. on R. I., p. 69, Coal Meas.
- SPIRAXIS, Newberry, 1884, Ann. N. Y. Acad. Sci., vol. 3, p. 217. An elongated spiral cast, supposed to represent a fucoiid. Type S. major. Another species, S. randalli, is also described. Both from the Chemung Gr.
- TRIGONOCARPUM ambiguum, Lesquereux, 1890, Dict. of Foss. Pa., vol. 3, p. 1213, Coal Meas.
- VOLKMANIA brevistachys, Lesquereux, 1890, Dict. of Foss. Pa., vol. 3, p. 1253, Coal Meas.

ANIMAL KINGDOM.

SUBKINGDOM PROTOZOA.

THE animals of this Subkingdom are supposed to have consisted of protoplasm, in a cell or cells, capable of secreting an outer wall, but without hardened tissues or alimentary organs. I arrange the palaeozoic sponges in this Subkingdom and in the Class Porifera, because I see no reason to suppose such forms as Pattersonia, Dystactospongia, Archæocyathus, Strephochetus, etc., were any more highly developed than Rhizopoda. It may be part of the Class has shown such variation and development as to point to a higher organization, which is in accordance with all intelligent views of evolution, but I see no evidence to warrant raising the Class to the rank of a Subkingdom in the animal scale, as some have done. In any view presented by those who claim that the Porifera occupy a place between Protozoa and Cœlenterata, I discover no ground for raising it to the rank of a Subkingdom, if any regular grade is to be maintained, in the classification of animals.

ERRATA.—For family "Leptonitidæ," on page 153, read "Family Leptomitidæ.—Leptomitus," and strike "Leptomitus" from those enumerated under "Family Affinity Uncertain." On page 161 change the specific names under Palæacis to the feminine gender.

ACANTHODICTYA, Hinde, 1889, Trans. Roy. Soc. Can., vol. 7, p. 47. [Ety. *akantha*, a spine; *dictyon*, net.] Subcylindrical, skeletal mesh-work of longitudinal and transverse spicular strands or fibres; longitudinal strands composed of somewhat loosely arranged fascicles of elongated overlapping spicules, and the spicules of the slender transverse fibres are, as a rule, disposed in a single series. From the outer surface of the sponge, numerous spicular rays project outward at right angles. Anchored, probably, by a basal prolongation of the longitudinal strands. Some of the elongated longitudinal spicules are cruciform, and their transverse rays form the cross fibres. The general structure resembles Cyathospongia, but is char-

acterized by the presence of the projecting surface rays. Type A. hispida, described in the same place, from Upper Taconic rocks.

Actinodictya placenta, Hall, 1892, 9th Ann. Rep. State Geologist, p. 59, Chemung Gr. Not entitled to recognition for want of illustration.

ACTINOSTROMA fenestratum, Nicholson, 1889, Monog. Brit. Stromatoporoids, p. 146, and A. tyrrelli, 1891, Ann. and Mag. Nat. Hist., vol. 7, p. 317, Devonian.

Biopella, Wallace, 1878, Am. Jour. Sci., vol. 115, p. 369. [Ety. *bios*, life; *palla*, a ball.] Founded upon very imperfect and certain material from the geodes of the Kuk Group, with the statement "there is uncertainty as to the action of species," and yet B.

R. I., p. 69, Coal

ann. N. Y. Acad.
elongated spiral
present a fucoid.
other species, S.
med. Both from

n, Lesquereux,
vol. 3, p. 1213,

s, Lesquereux,
vol. 3, p. 1253,

of protoplasm,
ordered tissues or
gdom and in the
as Pattersonia,
more highly de-
much variation and
ordance with all
ising the Class to
e. In any view
een Protozoa and
bkingdom, if any

Family Leptomi-
umerated under
ames under Palæ-

presence of the pro-
Type A. hispida,
place, from Upper

ll, 1892, 9th Ann.
p. 59, Chemung Gr.
nition for want of

n, Nicholson, 1889,
atoporoids, p. 146,
l, Ann. and Mag.
317, Devonian.
Am. Jour. Sci., vol.
bios, life; *palla*, a
on very imperfect
al from the geodes
p, with the state-
uncertainty as to
ecies." and yet B.

keokuk, *B. grandis*, *B. wortheni*, *B. wood-*
mani, *B. hecklei*, *B. hyntli*, *B. aliei*, and
B. palmata are named, with almost char-
acterless descriptions, without illustra-
tion.

CIDAROSPONGIA, Gurley, 1884, New Carb.
Foss. Bull., No. 2, p. 4. Round hemi-
spherical bodies, with flattened irregular
base, full hemispherical top or dome,
which, along the margin, is distinctly
divided into twelve lobes, being quite
regular in size and extending fully one-
half the distance from the margin to
the center of the dome, where they be-
come obsolete, leaving a smooth, slightly
depressed, central, circular area, whose
diameter is about one-half that of the
body. The top of the dome is punc-
tured by two well-defined circular open-
ings, which are situated close together,
and at the margin of the smooth cir-
cular area. During growth the dome
becomes elevated, and a lower or basal
portion becomes developed, extending
from the flattened base to the margin
of the dome. This lower portion is
somewhat irregularly marked by con-
centric lines of growth. The sides, be-
low the dome, are moderately straight.
Type C. ella. Described at the same
place, from the Coal Meas.

CRYPTOZOON *steeli*, Brainard & Seely, 1890,
Bull. Am. Mus. Nat. Hist., vol. 3, p. 6,
Calcliferous Gr.

Cryptodictya alleni, Hall, 1892, 9th Ann. Rep.
State Geologist, p. 60, Chemung Gr. Not
entitled to recognition for want of illus-
tration.

CUCUMULITES, Gurley, 1884, New Carb. Foss.
Bull., No. 2, p. 2. Body consisting of a
thin, punctate, elongate, tubular shell,
which is slightly arcuate and expand-
ing; being closed and rounded at the
larger extremity and somewhat pointed
with a well-defined terminal or oral
opening at the smaller extremity. Sur-
face ornamented by more or less nu-
merous pointed elevations or tubercles,
which along the central portion of the
shell become arranged in longitudinal
rows, breaking up and becoming irregu-
lar in crossing the larger rounded end
and toward the smaller extremity. Type
C. tuberculatus. Described at the same
place with C. tricar-
inatus from the War-
saw Group.

Cyathophycus siluriana,
James. Too poorly
defined to be recog-
nized.

CYATHOSPONGIA *quebec-*
ensis, Dawson &
Hinde, 1889, Trans.
Roy. Soc. Can., vol.
7, p. 44, Up. Taconic.

CYCLOSPONGIA, S. A. Miller, 1891, Advance
Sheets 17th Rep. Geo. Sur. Indiana,
p. 5. [Ety. *kuklos*, a circle; *spongia*,

sponge.] Sponge, circular, button-shaped
or discoid, and consisting of numerous
thin, calcareous laminae, having a con-
centric structure and filled with minute
canals or interstices. The structure has
some resemblance to that of *Strephochetus*
richmondense, but the laminae are much
thinner, and the interlaminae spaces are
much less marked,
and no vertical tubes
have been found with-
in them. That was a
free sponge; this one
is supposed to have
been attached to
some other object.
Type C. discus. De-
scribed in the same
place, from the Up-
per Helderberg Gr.

FIG. 1190. — *Cyclo-*
spongia discus.
Upper side.

Dictyophyton amalthæa, *D. halli*, *D. randalli*,
D. sceptrum, *D. scitum*, *D. tomaculum*, *D.*
vascellum, Hall, 1892, 9th Rep. State
Geologist, N. Y., pp. 56 to 58, Chemung
Gr. Such descriptions, without illus-
trations, are not entitled to recognition.

HALICHONDITES, Dawson, 1889, Trans. Roy.
Soc. Can., vol. 7, p. 52. Oval or irregu-
lar masses of small simple spicules,
imbedded in patches of pyrite, and,
without any definite arrangement or
root, spicules may indicate the pres-
ence of a halichondroid sponge. In the
best preserved specimens the spicules
appear to be biacerate and more slender
and pointed than in *Lasiothrix*, and they
seem to be in two series, inclined at a
very oblique angle to each other. In
some specimens elongated spaces, with
well-defined margins, are covered with
thin films of pyrites, which may have
resulted from the replacement or in-
crustation of a mass of minute spicules,
of which traces remain in some places.
He observed that sponges having origi-
nally much keratose or other dense
animal matter, would naturally aggre-
gate in and around themselves a greater
quantity of pyrite than those of a more
purely siliceous character. Type H.
confusus. Named at the same place,
from the Upper Taconic.

Hyalostelia metissica, Hinde, 1889, Trans.
Roy. Soc. Can., vol. 7, p. 49, Up. Ta-
conic. Founded upon a confused mass
of supposed spicules.

LASIOTHRIX, Hinde, 1889, Trans. Roy. Soc.
Can., vol. 7, p. 50. [Ety. *lasios*, shaggy;
thrix, hair.] Sponge small, depressed
oval in outline, the outer surface cov-
ered by a layer of longitudinally ar-
ranged, apparently simple, acerate spic-
ules; beneath this is another layer of
spicules disposed transversely. From
the base of the sponge several simple
elongated spicules extend. Type L.
curvicostata. Described in the same
place, from the Upper Taconic. L.
flabellata is also described.

FIG. 1195. — *Cyclo-*
spongia discus.
Lower side.

- PALEACIS cavernosa*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 4, Waverly Gr.
- PROTOSPONGIA coronata*, P. cyathiformis, P. delicatula, P. mononema, P. polynema, and P. tetranema, Dawson & Hinde, 1889, Trans. Roy. Soc. Can., vol. 7, pp. 37 to 43, Up. Taconic.
- Ptychosstylus*, Gurley, 1884. The name was preoccupied by Gabb in 1865.
- RECEPTACULITES elrodi*, S. A. Miller, 1892, Advance Sheets of 18th Rep. Geo. Sur. Ind., p. 3, Up. Held. Gr.
- Rhombedictyon globosus*, James. Too poorly defined to be recognized.
- Spiræ gathus*, Hinde. Synonym for *Archæocyathus*.
- Stephanella sancta*, Hinde, 1891, Lond. Geo. Mag., vol. 8, p. 22. Radiating lines of pyrites supposed to represent spicules. No generic or specific characters.
- Stromatopora ludlowensis* and *S. tubularis*, James. Too poorly defined to warrant any recognition.
- SYRINGOSTROMA. Type *S. densum*.

SUBKINGDOM CÆLENTERATA.

THE Order Rugosa has been called Tetracoralla, because the septa are said to be some multiple of four; and the Tabulata, Hexacoralla, because the septa are said to be a multiple of six. The internal cavity of a Graptolite, so far as I have been able to discover, is divided longitudinally into three or four departments, which is altogether different from the structure of the living Hydrozoa, where the interior part of the body consists of a single undivided cavity. It may be the Order Graptolida should be raised to the rank of a Class, because it is so different in composition and structure from the living Hydrozoa.

ERRATA.—On page 168, insert "Family Calceolidæ.—Calceola," and remove "Baryphyllum" to the "Cyclolitidæ." On page 169, for "Family Helioporidæ" read "Family Heliolitidæ." On page 177, strike out "Chonophyllum validum." In the eleventh line from the top of page 189, read "F. alpenensis" for "F. dumosus." In the sixth line from the bottom of page 194, read "1669" for "1869." In the first line on top of page 196, read "Description" for "Describes." In the third and seventh lines from the top of page 199, read "Hamilton" for "Chemung." In the fifteenth and sixteenth lines from the bottom of page 201, read "Edwards & Haime, 1850," for "Lonsdale, 1839, Sil. Syst., p. 691, and E. & H." For the word "lamellæ," in the definition of Zaphrentis, on page 208, read "septa," and add Clifford as an author of the genus with Rafinesque. On page 209, under Z. elliptica, read "Pal., No. 8," instead of Pal., No. 6."

FIG. 1197.—Amplexus blairi.

- ACTINOCYSTIS variabilis*, Whiteaves, 1892, Cont. to Can. Pal., p. 271, Devonian.
- Alveolites roemeri* refer to Cladopora.
- AMPLEXUS blairi*, A. bicostatus, A. corniculum, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 8. The blairi and corniculum from the Chouteau limestone and bicostatus from the Burlington Group.
- cinctus, S. A. Miller, 1892, 18th Rep. Geo. Sur. Ind., p. 5, Niagara Group.

- coralloides*, probably, not an American species.
- CHETTES ponderosus*, Rominger, 1892, Am. Geol., vol. 10, p. 61, Up. Held. Gr.
- CENOGRAPTUS*, Hall, 1868, 20th Rep. N. Y. Mus. Nat. Hist., pp. 210, 211, 251. [Ety. *koinos*, living together; *grapho*, I write.] Polypary compound, developed bilaterally from the initial point; cellules on one side of slender branches, which are developed on one or two sides of a long slender axis or rachis, the free extremities of which are likewise celluliferous. Not branching dichotomously. Type C. divergens.
- divergens, Hall, 1859 (Graptolithus divergens), Pal. N. Y., vol. 3, p. 509, Hud. Riv. Gr.
- gracilis, Hall, 1847 (Graptolithus gracilis), Pal. N. Y., vol. 1, p. 274, Utica Slate.

es. Too poorly
ym. for Archæ-
191, Lond. Geo.
diating lines of
esent spicules.
characters.
and *S. tubularis*,
ined to warrant
ensum.

ATA.

a are said to be
e septa are said
as I have been
ments, which is
ere the interior
the Order Grap-
erent in composi-

la," and remove
ly *Helioporidæ* "
phyllum validum."
sis" for "F. du-
ad "1669" for
tion" for "Des-
99, read "Ham-
e bottom of page
yst., p. 691, and
tis, on page 208,
esque. On page
"

not an American

minger, 1892, Am.
Up. Held. Gr.
20th Rep. N. Y.
10, 211, 251. [Ety.
; *grapho*, I write.]
developed bilater-
point; cellules on
ranches, which are
two sides of a long
a, the free extremi-
wise celluliferous.
omously. Type *C.*

Graptolithus diver-
bl. 3, p. 509, Hud.
Graptolithus gracilis),
274, Utica Slate.

surcularis, Hall, 1868, 20th Rep. N. Y.
Mus. Nat. Hist., p. 210, Utica Slate.
COLUMNARIA disjuncta, Whiteaves, 1892, Cont.
to Can. Pal., p. 269, Devonian.
Crepidophyllum is a synonym for *Crasped-*
ophyllum, and the species under it
should be referred to *Craspedophyllum*.
CYATHOPHYLLUM athabascense, Whiteaves,
1891, Cont. to Can. Pal., vol. 1, p. 202,
Devonian. *C. petraoides* and *C. waska-*
sense, 1892, pp. 264 and 265, Devonian.
C. juvenc, refer to *Heliophyllum juvenc*.
C. panicum refer to *Diphyphyllum panicum*.
CYSTELASMA, S. A. Miller, 1891, Advance
Sheets 17th Rep. Geo. Sur. Ind., p. 12,

FIG. 1168.—*Cystelasma* wrinkled or con-
lanesvillense. Summit
and side views.

[Ety. *kustia*, a cav-
ity; *elasma*, lamellæ.]
Corallum simple, ir-
regularly turbinate
or cylindrical, con-
sisting of an outer
wall, transversely
stricted, which is
connected by obli-
que plates, irregularly disposed, that
give to the interior cystose chambers of
unequal size and irregular shape. No
septa or regular tabulle. Structure ves-
icular. Type *C. lanesvillense*, described
in the same place from the Warsaw Gr.
CYSTIPHYLLUM greenii, S. A. Miller, 1892,
Advance Sheets 18th Rep. Geo. Sur.
Ind., p. 4, Up. Held. Gr.

DAWSONIA are small, conical or bell-shaped,
and each has a minute spine at the
summit. They have not been found in
connection with any Graptolite.

DICTYONEMA pertenne and *D. scalariforme*,
Foerste, 1887, Bull. Denison Univ.,
vol. 2, p. 107, Niagara Gr.

DIPHYPHYLLUM panicum, Winchell, 1866,
(*Cyathophyllum panicum*.) Rep. Low.
Penin. Mich., p. 90, Ham. Gr.

DIPLOTRYPA westoni, Ulrich, 1889, Micro-
palæontology of Can., pt. 2, p. 30, Hud.
Riv. Gr.

Graptolithus alatus, *G. crucifer*, *G. headi*,
G. logani, *G. octobrachiatus* refer to
Loganograptus. *G. divergens* and *G.*
gracilis refer to *Cœnograptus*.

Hadrophyllum apatum, Cummins, 1891,
2d Ann. Rep. Geo. Sur. Texas, p. 552,
Coal Meas.

FIG. 1199.—*Leptopora gorbyi*. Summit views.

HELIOPHYLLUM juvenc, Rominger, 1876,
(*Cyathophyllum juvenc*.) Foss. Corals,
p. 101, Ham. Gr.

parvulum, Whiteaves, 1891, Cont. to Can.
Pal., p. 203, Devonian.

LEPTOPORA gorbyi, S. A. Miller, 1891, Ad-
vance Sheets 17th Rep. Geo. Sur. Ind.,
p. 6, Chouteau limestone.

LOGANOGRAPTUS, Hall, 1868, 20th Rep. N. Y.
St. Mus. Nat. Hist., pp. 207 and 251.

FIG. 1200.—*Leptopora*
gorbyi. Basal
view.

[Ety. proper name; *grapho*, I
write.] Polypary compound,
growing bilaterally from the
initial point, and consisting of
four, eight or more simple
stipes numerously avided
near the base and below the
commencement of the cells;
furnished with a corneous cen-
tral disc. Type *L. headi*. To
this genus is also referred the species
described as *Graptolithus alatus*, *G. cru-*
cifer, *G. logani* and *G. octobrachi-*
atus.

MICROCYNCLUS blairi, S. A. Miller, 1892, Ad-
vance Sheets 18th Rep. Geo. Sur. Ind.,
p. 7, Chouteau
limestone.

MONOTRYPELLA con-
fluens, Foerste,
1887, Bull. Deni-
son Univ., vol.
2, p. 172, Niag.
ara Gr.

FIG. 1201.—*Microcynclus*
blairi.

unيجا, Whiteaves, 1891, Cont. to Can. Pal.,
vol. 1, p. 214, Devonian.

Monticulipora clintonensis, *crustulata*, *cleave-*
landi, *milfordensis* and *nicholsoni*, James,
1888, Jour. Cin. Soc. Nat. Hist., vol. 11,
pp. 15 to 36, Hud. Riv. Gr. Some of
these are synonyms for species de-
scribed under other genera, by Ulrich.
Others are not defined so as to be re-
cognized. Whether or not any of them
will stand is a question. *M. faleri* and
ohioensis by the same party, 1884, same
Journal, vol. 3, p. 137, Hud. Riv. Gr.
Not recognized.

M. molesta, Nicholson, seems to be a syn-
onym for *M. mammulata*.
parasitica, var. *plana*, Ulrich, 1889, Micro-
palæontology of Canada, pt. 2, p. 29,
Hud. Riv. Gr.

OLDHAMIA antiqua was described in 1844, in
Jour. Geo. Soc. Dublin, vol. 3, p. 60.

PACHYPHYLLUM devoniense, Edwards &
Haime, 1851, Polyp. Foss. Terr. Pal.,
p. 397, Devonian.

PRASOPORA parvula, Foerste, 1887, Bull.
Denison Univ., vol. 2, p. 170, Niagara
Group.

Stenopora ohioensis, Foerste, 1887, Bull. Deni-
son Univ., vol. 2, p. 85, Coal Meas.

STRIATOPORA gorbyi, S. A. Miller, 1892, Ad-
vance Sheets 18th Rep. Geo. Sur. Ind.
p. 7, Niagara Gr.

ZAPHRENTIS calyculus, *Z. chouteauensis*, *Z.*
declinis, *Z. exigua*, *Z. tantilla*, and
Z. tenella, S. A. Miller, 1891, Advance
Sheets 17th Rep. Geo. Sur. Ind., pp. 10,
11, and 12. All from the Chouteau lime-
stone except *Z. declinis*, which is from
the Keokuk Gr.

SUBKINGDOM ECHINODERMATA.

CLASS CRINOIDEA.

THE only systematic classification of crinoids into families, so far as learning has extended, is based upon the number and arrangement of the plates in the calyces. See "The Structure, Classification, and Arrangement of American Palæozoic Crinoids into Families," 16th Rep. Geo. Sur., Indiana, p. 302. The number of basal plates is first in importance. There are no rudimentary basals. Second in importance is the presence or absence of subradials. Third, the presence or absence of regular interradials. Fourth, the structure of the azygous side. Families having only two basal plates are confined to the Subcarboniferous and Carboniferous. Those having three basals extend from the Lower Silurian to the Subcarboniferous. Those having four basals extend from the Lower Silurian to the Devonian. And those having five basals extend from the earliest crinoids throughout Palæozoic time.

GENERA HAVING TWO BASALS.

FAMILY ACROCRINIDÆ.—Acrocrinus.

FAMILY DICHOCRINIDÆ.—Dichocrinus, Cotyledonocrinus, Talarocrinus.

FAMILY PTEROTOCRINIDÆ.—Pterotocrinus.

GENERA HAVING THREE BASALS, NO SUBRADIALS, REGULAR INTERRADIALS.

FAMILY ACTINOCRINIDÆ.—Actinocrinus, Agaricocrinus, Alloprosallocrinus, Amphorocrinus, Batocrinus, Blaiocrinus, Cylicocrinus, Dorycrinus, Eretmocrinus, Gennæocrinus, Megistocrinus, Phisetocrinus, Saccocrinus, Steganocrinus, Strotocrinus, Teleocrinus.

FAMILY ARTHRACANTHIDÆ.—Arthracantha.

FAMILY DOLATOCRINIDÆ.—Alloocrinus, Dolatocrinus, Hadrocrinus, Stereocrinus.

FAMILY PLATYCRINIDÆ.—Coccocrinus, Cordyloocrinus, Eucladoocrinus, Macrostylocrinus, Marsupioocrinus, Platycrinus.

GENERA HAVING THREE BASALS, SUBRADIALS, NO REGULAR INTERRADIALS.

FAMILY AMPHERISTOCRINIDÆ.—Ampheristocrinus, Closterocrinus.

FAMILY ICHTHYOCRINIDÆ.—Ichthyocrinus, Lecanocrinus, Mespilocrinus.

GENERA HAVING THREE BASALS, SUBRADIALS, AND REGULAR INTERRADIALS.

FAMILY TAXOCRINIDÆ.—Forbesocrinus, Onychocrinus, Taxocrinus.

GENERA HAVING THREE BASALS, NO SUBRADIALS, NO REGULAR INTERRADIALS. THE FAMILIES ARE ANOMALOUS AND HAVE NO NEAR AFFINITY WITH EACH OTHER.

FAMILY CALCEOCRINIDÆ.—Calceocrinus, Deltacrinus, Halysiocrinus.

FAMILY SYNATHOCRINIDÆ.—Synathocrinus.

FAMILY ZOPHOCRINIDÆ.—Zophocrinus.

GENERA HAVING FOUR BASALS, NO SUBRADIALS, REGULAR INTERRADIALS.

FAMILY EUCALYPTOCRINIDÆ.—Callierinus, Eucalyptocrinus, Hypanthocrinus.

FAMILY MELOCRINIDÆ.—Compsocrinus, Mariaocrinus, Melocrinus, Technocrinus.

FAMILY XENOCRINIDÆ.—Xenocrinus.

GENERA HAVING FIVE BASALS, FIVE SUBRADIALS, NO REGULAR INTERRADIALS.

FAMILY AGASSIZOCRINIDÆ.—Agassizocrinus.

FAMILY CYATHOCRINIDÆ.—Abrotoocrinus, Arachnocrinus, Bursacrinus, Carabocrinus, Cyathocrinus, Graphiocrinus, Palæocrinus.

FAMILY DENDROCRINIDÆ.—Dendrocrinus, Ottawacrinus.

FAMILY ERISOCRINIDÆ.—Erisocrinus, Menocrinus, Stemmatocrinus.

FAMILY EUPACHYCRINIDÆ.—Æsiocrinus, Delocrinus, Eupachyocrinus, Ulocrinus.

FAMILY MEROOCRINIDÆ.—Merocrinus.

FAMILY POTERIOCRINIDÆ.—Atelestocrinus, Baryocrinus, Cœliocrinus, Euspirocrinus, Goniocrinus, Homocrinus, Hydreionocrinus, Poteriocrinus, Scaphiocrinus, Vasocrinus, Zeacrinus.

GENERA HAVING FIVE BASALS, FIVE SUBRADIALS, REGULAR INTERRADIALS.

FAMILY GLYPTASTERIDÆ.—Cyphocrinus, Glyptaster, Lampteroocrinus, Thyasocrinus.

FAMILY GAUROCRINIDÆ.—Gaurocrinus, Retiocrinus.

FAMILY RHODOCRINIDÆ.—Archæocrinus, Goniasteroidocrinus, Lyriocrinus, Rhaphanoocrinus, Rhodocrinus.

GENERA HAVING FIVE BASALS, NO SUBRADIALS, REGULAR INTERRADIALS.

FAMILY CLEIOCRINIDÆ.—Cleioocrinus.

FAMILY GLYPTOCRINIDÆ.—Cupuloocrinus, Glyptocrinus, Pycnoocrinus, Schizocrinus, Siphonocrinus.

GENERA HAVING FIVE BASALS, NO SUBRADIALS, NO REGULAR INTERRADIALS.

FAMILY ANOMALOCRINIDÆ.—Anomalocrinus.

FAMILY BELEMNOCRINIDÆ.—Belemnocrinus.

FAMILY CATILLOCRINIDÆ.—Catilloocrinus.

FAMILY GAZACRINIDÆ.—Gazacrinus.

FAMILY HAPLOCRINIDÆ.—Allageocrinus, Haploocrinus.

FAMILY HETEROCRINIDÆ.—Ectenocrinus, Heterocrinus, Iocrinus, Obiocrinus.

FAMILY HYBOCRINIDÆ.—Hybocrinus.

FAMILY MISSOURICRINIDÆ.—Missouricrinus.

GENERA BELONGING TO ANOMALOUS FAMILIES.

FAMILY PISOCRINIDÆ.—Pisocrinus. This family has five basals, followed by three plates, that are both radial and subradial in position.

FAMILY EDRIOCRINIDÆ.—Edriocrinus. The base is solid in this family, and is followed by five radials.

FAMILY CAMAROCRINIDÆ.—Camarocrinus. Distinct from all other families.

FAMILY ANCYROCRINIDÆ.—Ancyrocrinus. Distinct from all other families.

GENERA UNCERTAIN AS TO FAMILY AFFINITY.

Aspidocrinus, Brachioocrinus, Coronocrinus, Cystocrinus, Nipterocrinus, Pachycrinus, and the fossil described by Hall as Myrtillocrinus americanus.

ORDER CYSTOIDEA.

FAMILY STRIBALOCYSTIDÆ.—Stribalocystites.

ABROTCRINUS, Miller & Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, p. 30, and 16th Rep. Geo. Sur. Ind., p. 350. [Ety. *abrotos*, immortal; *krinon*, lily.] Calyx bowl-shaped, depressed below; basals 5, occupying a shallow concavity; subradials 5, as high as wide; first radials pentagonal, wider than high, truncated horizontally the entire width of the plates, sutures gaping; brachial or second radials constricted in the middle and bearing upon the upper sloping sides the free arms; arms bifurcate frequently and bear pinules; no regular inter-radials. First azygous plate in line with the first radials, horizontally truncated above and having a gaping suture; second plate constricted in the middle, and followed by a single series of plates. Type *A. cymosus*, which is described at the same place, from the Keokuk Gr.

FIG. 1202.—*Abrotocrinus cymosus*.

ACTINOCRINUS blairi and *A. brittsi*, S. A. Miller, 1892, Adv. Sheets 18th Rep. Geo. Sur. Ind., pp. 35,

36, Burlington Gr., and *A. chouteauensis*, p. 18, from the Chouteau limestone, and *A. fossatus*, p. 40, from the Burlington Gr.

FIG. 1203.—*Actinocrinus chouteauensis*. Azygous view.

FIG. 1204.—*Actinocrinus chouteauensis*. Summit view.

FIG. 1205.—*Actinocrinus chouteauensis*. Basal view.

grandis, Miller & Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, p. 25, and 16th Rep. Geo. Sur. Ind., p. 346, Keokuk Gr. *nodosus*, S. A. Miller, 1891, Bull. No. 4, Geo. Sur. Mo., p. 33, Burlington Gr.

puteatus, Rowley & Hare, 1891, Kansas City Scientist, vol. 5, p. 101, Burlington Gr.

sedaliensis, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 16, Burlington Gr.

senarius, Hall, 1860, Supp. Geo. Rep. Iowa, p. 25, syn. for *Physetocrinus ornatus*.

ESIOCRINUS, Miller & Gurley, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 14. [Ety. *aisios*, auspicious, coming at good time;

us, Ohioerinus.

ES.
als, followed by
is family, and is
ther families.
ther families.

V.
rocrinus, Pachyc-
us.

and A. chouteau-
ne Chouteau lime-
us, p. 40, from the

g. 1204.—Actinoerinus chouteauensis. Summit view.

grandis, Miller & Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, p. 25, and 16th Rep. Geo. Sur. Ind., p. 346, Keokuk Gr.
nodosus, S. A. Miller, 1891, Bull. No. 4, Geo. Sur. Mo., p. 33, Burlington Gr.
hare, 1891, Kansas

er, 1892, Advance
eo. Sur. Ind., p. 16,

pp. Geo. Rep. Iowa,
etocrinus ornatus.
Gurley, 1890, Jour.
vol. 13, p. 14. [Ety.
oming at good time;

krinon, lily.] Column pentagonal, calyx bowl-shaped, plates smooth or granular. Basals five, forming a pentagonal, flattened or concave disc. Subradials large, four hexagonal, one heptagonal, and curving upward half the height of the calyx. First radials 5, truncated above; one or more brachials in each ray supporting strong arms, composed of a single series of plates; arms 10, bearing pinnules. No regular interradials. An azygous interradial rests upon a subradial, between two first radials, and is followed by two plates that connect with the proboscis. Proboscis long, composed of four series of plates bearing numerous transverse fissures on the sides of the plates. Type *A. magnificus*, described at the same place, from the Up. Coal Meas. *A. harii* was described at the same time, and *A. basilicus* was described in Desc. New Gen. and Spec. Echinodermata, p. 53, from the Up. Coal Meas. All of which were republished in 16th Rep. Geo. Sur. Indiana, pp. 337, 338, and 369.

FIG. 1206.—Actinoerinus magnificus.

lykinsi, Butts, 1891, Kansas City Scientist, vol. 5, p. 144, Coal Meas.

ÆTHOCYSTITES, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 9.

FIG. 1207.—Aethocystites sculptus.

form an obconical cup, commencing from a small column. In the second range there are five elongated plates.

The third range consists of five much shorter plates. The plates are ornamented with wrinkles, and bear tubular ridges radiating from a central point, in the middle range, which follow the longitudinal sutures or center of the first and third ranges, and have porous connection with the interior of the body. The plates do not possess pores after the manner of *Holocystites* or *Caryocrinus*, and no pores have been determined, except as above stated. Type *A. sculptus*. Described at the same place, from the Niagara Gr.

AGANASTER, Miller & Gurley, 1890, Desc. New Gen. and

FIG. 1208.—Aganaster gregarius. Dorsal side.

Spec. Echinodermata, p. 57, republished in 16th Rep. Geo. Sur. Ind., p. 372. [Ety. *agan*, very much; *aster*, star.] Circular disc with five long, narrow rays. Dorsal side covered with small polygonal plates, not inter-

rupted by the presence of the rays, thus showing the disc has a greater depth than the rays have. Rays narrow, convex, spine-bearing, and composed of plates arranged opposite each other. Ventral side has a deep central disc and ten oral plates. Type *A. gregarius*. Described by Meek and Worthen as *Protaster gregarius*.

FIG. 1209.—*Aganaster gregarius*. Dorsal side of arm, magnified.

AGARICOCRINUS blairi and *A. sampsoni*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., pp. 20 and 21, Chouteau limestone, and *A. chouteauensis* and *A. germanus*, pp. 42 and 43, Chouteau limestone.

dissimilis, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 55, Keokuk Gr. And also at the same time and place, from the same Group, *A. gorbyi* and *A. indianensis*.

decoratus, Rowley & Hare, 1891, Kansas City Scientist, vol. 5, p. 117. Burlington Gr.

splendens, Miller & Gurley, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 18, republished in 17th Rep. Geo. Sur. Ind., p. 55, Keokuk Gr.

AGELACRINUS blairi, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 12, Keokuk Gr.

FIG. 1210.—*Allocrinus benedicti*. Basal view.

ALLOCRINUS benedicti, S. A. Miller, 1891, 17th Rep. Geo. Sur. Ind., p. 37, Niagara Gr.

ALLOPROSALLOCRINUS gurleyi, S. A. Miller, 1891, 17th Rep. Geo. Sur. Ind., p. 58, Keokuk Gr.

FIG. 1211.—*Alloprosallocrinus gurleyi*. Basal and side views.

ARACHNOCRINUS canadensis, Whiteaves, 1891, Cont. to Can. Pal., p. 208, Devonian.

ARCHÆOCIDARIS legrandensis, Miller & Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, p. 59, Kinderhook Gr.

ATELEOCYSTITES, Billings, is probably a good genus, and not a synonym for *Anomalocystites*. In that case, *Ateleocystites huxleyi* would be the type, and the

genus may include *A. balanoides*, which does not seem to belong to Hall's genus *Anomalocystites*.

BARYCRINUS blairi and *B. boonvillensis*, S. A. Miller, Bull. No. 4, Geo. Sur. Mo., p. 25, Keokuk Gr.

FIG. 1212.—*Barycrinus princeps*.

latus, Hall, 1861, (*Cyathocrinus latus*), Proc. Bost. Soc. Nat. Hist., p. 292, Burlington Gr.

princeps, Miller and Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, p. 52, Keokuk Gr.

FIG. 1213.—*Barycrinus princeps*. Azygous view.

stellifer, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 46, Keokuk Gr.

balanoides, which
ing to Hall's genus

B. boonvillensis,
4, Geo. Sur. Mo.,

us princeps.

athocrinus latus),
Itist., p. 292, Bur-

Gurley, 1890, Desc.
ec. Echinodermata,

uceps. Azygous view.

ller, 1892, Adva
Geo. Sur. Ind., p. 46,

BATOCRINUS agnatus, B. boonvillensis, B. crawfordsvillensis, B. decoris, B. gorbui, B. gurleyi, B. mediocris, B. pulchellus, B. spergenensis, B. venustus, S. A. Miller, 17th Rep. Geo. Sur. Ind., pp. 53, 60 to 68. All from the Keokuk Gr. except B. decoris and B. spergenensis, which are from the Warsaw Gr.

FIG. 1214.—Batocrinus
icosidactylus.

gurleyi, and sweeti are from the Keokuk Gr., B. davisii from the Kaskaskia Gr., and the others from the Burlington Gr.

cantonensis, B. facetus, B. jucundus, B. marinus, B. poculum, Miller and Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, pp. 19, 20, 34 to 36, B. jucundus and B. marinus were published in the Jour. Cin. Soc. Nat. Hist., vol. 13, pp. 19 and 20, and all were republished in the 16th Rep. Geo. Sur. Ind., pp. 340, 341, 352, 353, and 354. B. poculum is from the Kinderhook Gr., the others from the Keokuk Gr.

calvini, Rowley, 1890, Am. Geol. vol. 5, p. 146, Burlington Gr.

britti and B. comparilis, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., pp. 32 and 33, Burlington Gr.

FIG. 1216.—Blairocrinus arrosus. Basal and summit views.

and B. decrepitus, p. 34, Keokuk Gr., and B. blairi, p. 39, Burlington Gr. divalis, S. A. Miller, 1892, Advance

bulbosus, B. davisii, B. gurleyi, B. inflatus, B. rotadentatus, B. sweeti, and B. abscissus, Rowley and Hare, 1891, Kansas City Scientist, vol. 5, pp. 102 and 114 to 117. B. abscissus,

FIG. 1215.—Batocrinus irreg-
ularis.

Sheets 18th Rep. Geo. Sur. Ind., p. 22, Keokuk Gr., and at same place republished B. iccoidactylus and B. irregularis.

BELEMNOCHINUS sampsoni, S. A. Miller, 1890, Bull. No. 4, Geo. Sur. Mo., p. 26, Burlington Gr.

BLAIROCRINUS, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 69. [Ety. proper name; *krinon*, lily.] Calyx low, saucer-shaped, but not depressed at the base. Surface deeply sculptured and bears radial ridges, though the inter-radial ridges are not sunken; vault elevated above the arm openings, more or less convex above, with a short subcentral proboscis, having an opening on top, surrounded by numerous small plates. Basals, 3, forming a flat hexagonal disc; primary radials, 3 x 5;

FIG. 1217.—Blairocrinus trijugis. Basal, summit, and side views.

secondary radials, 1 x 10, axillary; tertiary radials, 1 x 20; regular inter-radials, one large plate resting upon the first radials, followed by one or two ranges of two plates each, and these by two elongated plates that connect with the plates of the vault. First azygous plate in line with the first radials, followed by ranges of two plates until they connect with the plates of the vault. Type B. trijugis, described at the same time and place from the Chouteau limestone.

arrosus and B. bullatus, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 41, Chouteau limestone.

CALCEOCRINUS should be corrected, p. 230, ninth line from the top, so as to read "three" instead of "four" anchylosed plates.

indianensis, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 35, Niagara Gr.

CALLICRINUS, D'Orbigny, 1850, Prodrome Pal. Stratigraphique, 1, p. 45. [Ety. *kallos*, beautiful; *krinon*, lily.] Body oblong, cylindrical; calyx cup-shaped, base excavated for the insertion of the column. Basals 4, unequal, cuneate; nosubradials; radials 3 x 5; first one hexagonal, transverse, arcuate below; second one short, quadrangular; third, pentagonal, axillary; secondary radials, 2 x 10, the second axillary and supporting the arms, which are composed of a double series of plates bearing pinnules; first inter-radial large, decagonal, bearing two elongated plates in the second series, like Eucalyptocrinus; vault and proboscis as in Eucalyptocrinus; surface

deeply sculptured or bearing a more or less developed spine on each radial and interradial plate. Type *C. costatus*. *acanthinus*, Ringueberg, 1890, Ann. N. Y. Acad. Sci., vol. 5, p. 302, Niagara Gr. *hechleri*, Wachsmuth & Springer, 1892, Am. Geol., vol. 10, p. 140. Not defined so as to be recognized.

FIG. 1218.—*Caryocrinus indianensis*. Side and summit views.

CARYOCRINUS indianensis, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 19, Niagara Gr.

CLIOCRINUS, p. 231, is, probably, incorrectly defined. See my remarks, p. 323, 16th Rep. Geo. Sur. Ind.

CODASTER gracillimus and *C. grandis*, Rowley and Hare, 1891, Kansas City Scientist, p. 99, Burlington Gr.

CONONITES inopinatus, Rowley and Hare, 1891, Kansas City Scientist, pp. 100 and 118, Burlington Gr.

CYATHOCRINUS benedicti, C. gurleyi, C. labyrinthicus, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., pp. 47 to 49, *C. benedicti*, from the Niagara Gr. and the other two species from the Keokuk Gr.

FIG. 1219.—*Cyathocrinus gurleyi*.

multibrachiatus var. *squamosus*, Hall, 1872, in a note to photographic plate No. 5, sometimes distributed with Desc. New Spec. Crin. from the Carb. rocks of the Misc. Valley, Keokuk Gr. *opimus*, Miller and Gurley, 1890, Desc.

New Gen. and Spec. Echinodermata, p. 28, and 16th Rep. Geo. Sur. Ind., p. 348, Keokuk Gr.

FIG. 1220.—*Cylicocrinus canaliculatus*.

CYLIPOCRINUS, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 31. [Ety. *kulir*, *ikos*, a cup; *krinon*, a lily.] Calyx urn-shaped, truncated below. Basals 3, expanded; primary radials, 3 x 5,—the first one very large; second one, small, quadrangular; third, small, pentagonal; secondary radials 2 or more; arms 10. Regular interradials consist of one large plate followed by very small ones between the arm-bases that connect with the vault-plates; first azygous plate in line with the first radials, followed by three plates in the second and in the third ranges, which are connected with smaller plates that connect with the vault. The vault is convex and covered with minute plates, except the ambulacral grooves, which are open and have serrated edges as if protected by some kind of cilia. The primary radials resemble those in some species of *Batocrinus*; the azygous area resembles *Saccocrinus*; the vault is different from that in all other known genera of crinoids. Type *Cylicocrinus canaliculatus*, described at the same place from the Niagara Gr.

FIG. 1221.—*Cyphocrinus gorbyi*. Basal and summit views.

CYPHOOCRINUS, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 50. [Ety. *kuphos*, bowed down; *krinon*, lily.] Calyx obconoidal or obpyramidal as high as the first interradials, then rapidly expands and curves downward until the periphery and ambulacral orifices are directed below a horizontal line. Basals 5, forming a pentagonal disc; sub-radial, 5, hexagonal, except one which is truncated by the first azygous plate; it is heptagonal; primary radials 3 x 5; 3 of the first ones heptagonal,

c. Echinodermata.
p. Geo. Sur. Ind.,

as canaliculatus.

ler, 1892, Advance
Geo. Sur. Ind., p. 31.
up; *krinon*, a lily.]
truncated below.
; primary radials,
very large; second
ular; third, small,
ary radials 2 or
regular interradials
plate followed by
ween the arm-bases
e vault-plates; first
ne with the first ra-
three plates in the
third ranges, which
smaller plates that
ault. The vault is
with minute plates,
cral grooves, which
errated edges as if
kind of cilia. The
emble those in some
s; the azygous area
nus; the vault is
in all other known
Type *Cylicocrinus*
ibed at the same
ara Gr.

rybi. Basal and sum-
ws.

ler, 1892, Advance
Geo. Sur. Ind., p. 50.
ved down; *krinon*,
bidal or obpyramidal
t interradials, then
d curves downward
and ambulacral ori-
below a horizontal
ming a pentagonal
exagonal, except one
by the first azygous
nal; primary radials
st ones heptagonal,

and the other two hexagonal; second
radials quadrangular; third radials pent-
agonal, and bear on the upper slop-
ing sides secondary
radials; regular inter-
radials numerous; the
first one large and
resting between the
short upper sloping
sides of the first radials;

FIG. 122.—*Cyphocrinus gorbui*. Side
view of calyx.

It is followed, in the
second range, by two
plates, and by three
plates in succeeding ranges until they
unite with the plates of the vault; inter-
secondary radials present; first azygous
plate large, truncates a subradial, and
is followed by three or four plates in
each succeeding range, until they unite
with the plates of the vault; vault con-
vex and covered with more or less num-
erous plane, convex, or spinous
plates; a large spinous plate occupies
the center of the vault, and the anal
orifice, without any prominence, is on
the azygous side of it. Type *Cypho-*
crinus gorbui, described at the same
place from the Niagara Gr.

DELOCINUS, Miller and Gurley, 1890, Jour.
Cin. Soc. Nat. Hist., vol. 13, p. 9, and
16th Rep. Geo. Sur. Ind., p. 333. [Ety.
delos, manifest; *krinon*, lily.] Calyx
basin-shaped; arms broad, composed of
a double series of interlocking plates;
column round; surface smooth or granu-
lous; plates thick; basals 5, occupying
a concavity and forming a cone in the
interior; subradials 5, large, inflexed
below the middle, regularly arched, and
the upper part terminating in an acute
angle; first radials wider than high,
truncated above and separated from the
second radial on the outer face by a
gaping suture, but immediately within,
a straight crenated ridge extends from
one outer angle of the plates to the
other, having a furrow on each side so

FIG. 123.—*Delocrinus hemisphericus*.

as to form a toothed hinge on which the
second plate articulates; behind this
hinge, in the middle part of each plate,
there is a socket for the reception of a

tooth-like projection; second radials or
brachials produced externally in a strong
spine; no regular interradials; one
azygous interradial resting on a sub-
radial and followed by another piece
above the top of the calyx. Type *D.*
hemisphericus.

craig, Worthen, 1875, (*Eupachyrinus*
craig), Geo. Sur. Ill., vol. 6, p. 527, Coal
Meas.

fayettensis, Worthen, 1873, (*Eupachyri-*
nus fayettensis), Worthen, Geo. Sur.
Ill., vol. 5, p. 565, Up. Coal Meas.

hemisphericus, Shumard, 1858, (*Poteri-*
ocrinus hemisphericus), Trans. St. Louis
Acad. Sci., vol. 1, p. 221, and 16th Rep.
Geo. Sur. Ind., p. 335, Up. Coal
Meas.

inflexus, Geinitz, 1866, (*Cyathocrinus*
inflexus), Carb. und Dyas, in Neb.,
p. 62, and White's Cont. to Pal., No. 6,
p. 123, Up. Coal Meas.

missouriensis, Miller and Gurley, 1890,
Jour. Cin. Soc. Nat. Hist., vol. 13, p. 14.
Up. Coal Meas.

DENDROCRINUS nodobrachiatus, Ringueberg,
1890, Ann. N. Y. Acad. Sci., vol. 5,
p. 303, Niagara Gr.

DICHOCRINUS blairi, S. A. Miller, 1891, Ad-
vance Sheets 17th Rep. Geo. Sur. Ind.,
p. 36, Keokuk Gr.

cinctus, Miller & Gurley,
1890, Jour. Cin. Soc. Nat.
Hist., vol. 13, p. 21, and
16th Rep. Geo. Sur. Ind.,

FIG. 124.—*Dichocrinus cinctus*. Two of the views
are magnified.

p. 342, Kinderhook or Waverly Gr.

humberti, S. A. Miller, 1891, Bull. No. 4,
Geo. Sur. Mo., p. 26, and Advance Sheets
17th Rep. Geo. Sur. Ind., p. 36, Ke-
okuk Gr.

parvulus, S. A. Miller, 1891, Bull. No. 4,
Geo. Sur. Mo., p. 27, Keokuk Gr.

ulrichi, Miller & Gurley, 1890, Desc.
New Gen. and Spec. Echinodermata,
p. 48, and 16th Rep. Geo. Sur. Ind.,
p. 366, Keokuk Gr.

DOLATOCRINUS has only three basals, as
shown by a specimen belonging to Mr.
Gurley.

DORYCRINUS amonius, and *D. confragosus*,
S. A. Miller, 1890, Bull. No. 4, Geo. Sur.
Mo., p. 34, Burlington Gr.

elegans, S. A. Miller, 1892, Advance Sheets
18th Rep. Geo. Sur. Ind., p. 17, Bur-
lington Gr.

inflatus, Rowley & Hare, 1891, Kansas City Scientist, p. 114, Burlington Gr.
ECHINODISCUS sampsoni, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 76, Keokuk Gr.

FIG. 1225.—*Echinodiscus sampsoni*.

EOCIDARIS blairi, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 73, Keokuk Gr.

FIG. 1226.—*Eocidaris blairi*.

FIG. 1227.—*Eocidaris blairi*. Showing spines.

ERETMOCRINUS lyonanus, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 59, Keokuk Gr.
prægravis, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 37, Keokuk Gr.
Erisocrinus inflexus. See *Delocrinus inflexus*.

FIG. 1228.—*Eretmocrinus lyonanus*. Side and basal views.

EUCALYPTOCRINUS ellipticus, E. elrodi, E. gorbyi, E. subglobosus, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., pp. 37 to 40, Niagara Gr.
muralis, Ringueberg, 1890, Ann. N. Y. Acad. Sci., vol. 5, p. 305, Niagara Gr.
lindahli, Wachsmuth & Springer, 1892, Am. Geol., vol. 10, p. 139. Not defined so as to be recognized.

EUPACHYCRINUS craigi. See *Delocrinus craigi*.
fayettensis. See *Delocrinus fayettensis*.
hemisphericus. See *Delocrinus hemisphericus*.

FIG. 1229.—*Eupachyrcrinus harii*.

harii, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 71, Up. Coal Meas.
magister, Miller & Gurley, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 4, and 16th Rep. Geo. Sur. Ind., pp. 328, 371, Up. Coal Meas.
sphaeralis, Miller & Gurley, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 5, and 16th Rep. Geo. Sur. Ind., p. 329, Up. Coal Meas.

FIG. 1230.—*Eupachyrcrinus tumulosus*. Azygous and basal views.

tumulosus, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 70, Kaskaskia Gr.
FORBESOCRINUS elegantulus, S. A. Miller, 1891, Bull. No. 4, Geo. Sur. Mo., p. 40, Keokuk Gr.
speciosus, Miller & Gurley, 1890, Desc. New Gen. and Spec. Echinodermata,

FIG. 1231.—*Forbesocrinus speciosus*.

p. 27, and 16th Rep. Geo. Sur. Ind., p. 347, Keokuk Gr.
spinifer instead of "spiniger."

See *Delocrinus*
crinus fayettensis.
Delocrinus hemi-

crinus harli.

1891, Advance Sheets
ur. Ind., p. 71, Up.

Gurley, 1890, Jour.
at., vol. 13, p. 4, and
r. Ind., pp. 328, 371,

Gurley, 1890, Jour.
at., vol. 13, p. 5, and
ur. Ind., p. 329, Up.

is tumulosus. Azygous
al views.

Miller, 1891, Advance
Geo. Sur. Ind., p. 70,

atulus, S. A. Miller,
Geo. Sur. Mo., p. 40,

Gurley, 1890, Desc.
Spec. Echinodermata,

crinus speciosus.

Rep. Geo. Sur. Ind.,
r.
"spiniger."

GAZACRINUS S. A. Miller, 1892, Adv. Sheets
18th Rep. Geo. Sur. Ind., p. 49. [Ety.

FIG. 1232.—*Gazacrinus inornatus*.

radials 2 x 10; arms composed of a single series of flattened plates, one plate in each radial area; vault sustained by a specialized frame-work, with ambulacral canals connecting the arms with a central orifice. Type *Gazacrinus inornatus*. Described at the same place, from the Niagara Gr.

FIG. 1233.—*Gazacrinus inornatus*.

GLYPTASTER lockportensis, Ringneberg, 1890, Ann. N. Y. Acad. Sci., vol. 5, p. 304, Niagara Gr.

GONIASTEROIDOCRINUS tuberosus is from the Keokuk Gr. It is re-defined in the Advance Sheets 17th Rep. Geo. Sur. Ind., p. 51, and several errors respecting it are corrected.

GONIOCRINUS, Miller & Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, p. 32, and 16th Rep. Geo. Sur. Ind., p. 351. [Ety. *gonia*, an angle; *krinon*, a lily.] Calyx small, basin-shaped; plates convex or angular. Basals 5, small, extending beyond the column. Sub-

FIG. 1234.—*Gonioasteroidocrinus tuberosus*. Showing the pendulous arms.

radials five, about the same size as the basals; first radials larger, wider than long, and supporting on the slightly concave upper faces, a little shorter

than the width of the plates, the brachials; brachials 3 in each ray, flanged at the sides; arms resembling *Scaphiocrinus*; no regular interradials; azygous interradials, consisting of a series of plates, the first one like a first radial, and resting upon the upper truncated face of a subradial, which is followed by plates very much like the brachials, which form a convex, arm-like appendage that curves in toward the proboscis at or above the base of the free arms. A small azygous plate also exists on the right side of the area resting between the upper sloping sides of two subradials and the under sloping side of a first radial, and the azygous plate which truncates a subradial. Column pentagonal, bearing

FIG. 1235.—*Gonioocrinus sculptilis*. Natural size and magnified.

FIG. 1236.—*Gonioocrinus sculptilis*. Left side view and azygous side. Both magnified a little.

thinner plates; canal pentagonal. Type *G. sculptilis*, which is described at the same place, from the Waverly or Kinderhook Gr.

GRANATOCRINUS aplatus, *G. concinnulus*, *G. excavatus*, *G. exiguus*, *G. pyriformis*, Rowley & Hare, 1891, Kansas City Scientist, vol. 5, pp. 99, 100, 117, 118, Burlington Gr.

GRAPHIOCRINUS must be restored, as described by De Koninck & Lehon, and the redefinition of Wachsmuth & Springer wholly set aside. The generic formula is as follows: Basals 5; radials 2 x 5; anal 1; arms 10; none bifurcate. There is not a shadow of reason or evidence for supposing De Koninck and Lehon were mistaken in their diagnosis.

Holocystites *adipatus*, H. benedicti, H. colletti, H. commodus, H. gorbvi, H. indianensis, H. madisonensis, H. ornatisimus, H. papulosus, H. parvulus, H. parvus, H. scitulus, H. spungleri, H. subovatus, H. wykoiffi, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., pp. 13

FIG. 1237.—*Holocystites subovatus*. Summit.

to 18, Niagara Gr. *amplus*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 8, Niagara Gr. **Hydrionocrinus** *pentagonus*, Miller & Gurley, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 17, and 16th Rep. Geo. Sur. Ind., p. 330, Up. Coal Mens.

FIG. 1238.—*Hydrionocrinus pentagonus*.

Hyptiocrinus *typus*, Wachsmuth & Springer, 1892, Am. Geo., vol. 10, p. 138. Not defined so as to be recognized.

Ichthyocrinus *conoideus*, Ringueberg, 1890, Ann. N. Y. Acad. Sci., vol. 5, p. 305, Niagara Gr.

greenii, S. A. Miller, 1892, 18th Rep. Geo. Sur. Ind., p. 52, Keokuk Gr.

Idiocrinus *elongatus* and *I. ventricosus*, Wachsmuth & Springer, 1892, Am. Geo., vol. 10, p. 135. Not defined so as to be recognized.

Lecanocrinus *tennesseensis*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 41, Niagara Gr.

Mariacrinus *aureatus*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 34, Niagara Gr.

granulosus, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 35, Niagara Gr.

Melocrinus *æqualis*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 48, Niagara Gr.

oblongus, *M. parvus*, Wachsmuth & Springer, 1892, Am. Geo., vol. 10, p. 143. Not defined so as to be recognized.

Melonites was preoccupied when Owen & Norwood used it, and Meek & Worthen proposed instead of it *Melonechinus*.

Menocrinus has five basals, and does not have any near affinity with *Platycrinus*, nor belong to the same family.

Missouricrinus, S. A. Miller, 1890, Bull. No. 4, Geo. Sur. Mo., p. 31. [Ety.

proper name; *krinon*, lily.] Calyx obconoidal or basin-shaped; plates smooth or granulous; basals 5, forming a small cup; no subradials; no regular interradials; primary radials 1x5, wider than high, and separated from the brachials by an external gaping suture; brachials axillary, except in the ray

FIG. 1239.—*Missouricrinus admonitus*.

opposite the azygous side; arms resembling those in *Scaphiocrinus*; first azygous interradial rests between two primary radials and truncates a basal plate; column pentagonal. Type *M. admonitus*, which was described at the same time, from the Burlington Gr.

Myelodactylus *gorbyi*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 72, Niagara Gr.

Nipterocrinus has five basals.

FIG. 1240.—*Onychaster asper*.

Onychaster *asper*, *O. confragosus*, *O. demissus*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 74, Keokuk Gr.

Onychocrinus *cantonensis*, Miller & Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, p. 41, and 16th Rep. Geo. Sur. Ind., p. 358, Keokuk Gr.

ulrichi, Miller & Gurley, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 17, and 16th Rep. Geo. Sur. Ind., p. 330, Keokuk Gr.

FIG. 1241.—*Onychaster demissus*.

norwoodi, Meek & Worthen should be restored, as it is not a synonym for *O. exculptus*.

Ottawacrinus, p. 265, first line at top of page read subradial instead of "basal."

Paleocystites, p. 267, is described in Decade 3.

Pentremites *basilaris* and *P. broadheadi*

ly.) Calyx ob-
; plates smooth
forming a small
o regular inter-
s 1 x 5, wider
ated from the
gaping suture;
pt in the ray

s admonitus.

side; arms re-
phioerinus; first
ata between two
uncates a basal
onal. Type M.
described at the
Burlington Gr.

A. Miller, 1891,
Rep. Geo. Sur.

als.

ter asper.

fragosus, O. de-
1891, Advance
Sur. Ind., p. 74,

s, Miller & Gur-
Gen. and Spec.
and 16th Rep.
Keokuk Gr.

ley, 1890, Jour.
ol. 13, p. 17, and
nd., p. 339, Keo-

demissus.

then should be
synonym for O.

t line at top of
stead of "basal."
described in Dec-

d P. broadheadi

are described on p. 150, and *P. clavatus*
and *P. hemisphericus*, on p. 157; *P.*
gemmiformis, p. 553; *P. nodosus*, p. 155;
P. sampsoni, p. 551; and *P. spinosus*,
p. 154, of vol. 4, Trans. St. Louis Acad.
Sci. *P. cherokeus* was not defined by
Troost, and as Roemer described *P. sul-*
catus in 1852, it was too late for Hall to
call the same species *P. cherokeus* in
the Geo. Sur. Iowa, in 1858.

laterniformis, see *Troostocrinus laterni-*
formis.

Pisocrinus benedicti, *P. campana*, *P. gorbyi*,
S. A. Miller, 1891, Advance Sheets 17th
Rep. Geo. Sur. Ind., pp. 28 to 32, Ni-
agara Gr. *P. gem-*
miformis is redefi-
ned at the same
place and the ge-
neric characters dis-

FIG. 1242.—*Pisocrinus benedicti*. Side and basal views.

pyriformis was described in 1884.

PLATYCRINUS *absentivus*, *P. equiternus*,
P. allophyllus,
P. annosus, *P.*
brittsi, *P. ollic-*
ula, from the
Chouteau lime-
stone; *P. ac-*
clivus, *P. batli-*
ola, *P. blairi*,
P. broadheadi,
P. carchesium,
P. concinnus, *P. gorbyi*, *P. lautus*, *P. oc-*
cidental, *P. pulcellus*, *P. rotundus*,
P. sampsoni, *P. sulcatus*, from the Bur-

FIG. 1243.—*Pisocrinus gorbyi*. Summit and side view showing arm-blades.

lington Gr.; and *P. æternalis*, *P. amabi-*
lis, *P. boonvillensis*, *P. pentagonus*,
from the Keokuk Gr., Bull. No. 4, Geo.
Sur. Mo., pp. 8 to 23, S. A. Miller,
1890.

alabamensis, S. A. Miller, 1891, Advance
Sheets 17th Rep. Geo. Sur. Ind., p. 50,
Kaskaskia Gr.

altidorsatus, *P. corbuliformis*, *P. mar-*
ginatus, *P. pisum*, *P. planobasalis*,
from the Burlington Gr.; *P. curry-*
villensis, *P. insolens*, from the Chou-
teau limestone; Rowley & Hare,
1891, Kansas City Scientist, pp. 97,
98, 113.

caducus, S. A. Miller, 1892, Advance
Sheets 18th Rep. Geo. Sur. Ind., p. 13,
Keokuk Gr., and at same place *P. chou-*
teauensis and *P. colletti* from the Chou-
teau limestone.

POTERIOCRINUS agnatus, *P. amœnus*, *P.*
boonvillensis, *P. coryphæus*, S. A.
Miller, 1891, Advance Sheets 17th Rep.
Geo. Sur. Ind., pp. 42 to 45, Keokuk
Group.

arcanus, *P. cantonensis*, *P. crawfordsvill-*
ensis, *P. granilineus*, *P. subramosus*,
P. verus, from the Keokuk Gr.; *P. ge-*
nista, *P. legrandensis*, *P. scopæ*, *P.*
spartarius, from the Kinderhook Gr.;
Miller and Gurley, 1890, Jour. Cin. Soc.
Nat. Hist., vol. 13, pp. 23, 24, and Desc.

New Gen. and Spec. Echinodermata,
pp. 29, 37 to 40, 49, republished, 18th
Rep. Geo. Sur. Ind., pp. 343, 344, 348, 355
to 358, 365.

brittsi, S. A. Miller, 1890, Bull. No. 4, Geo.
Sur. Mo., p. 30, Keokuk Gr.

meekianus refer to *Cyathocrinus meeka-*
nus, Chouteau limestone.

waltersi, Rowley and Hare, 1891, Kan-
sas City Scientist, p. 101, Burlington
Gr.

Protaster gregarius, see *Aganaster gregarius*.

Ruopocrinus benedicti, S. A. Miller, 1892,
Advance Sheets 18th Rep. Geo. Sur.
Ind., p. 15, Keokuk Gr.

celatus, *R. sculptus*, Miller and Gurley,
1890, Desc. New Gen. and Spec. Echi-
nodermata, pp. 42, 43, and 16th Rep.

FIG. 1244.—*Sacocrinus howardi*. Side, azygous, and summit views.

Geo. Sur. Ind., pp. 359, 360, Kinder-
hook Gr.

nanus is from the Kinderhook Gr.
parvus, S. A. Miller, 1891, Bull. No. 4,
Geo. Sur. Mo., p. 39, Keokuk Gr.

SACCOCRINUS benedicti and *S. howardi*,
S. A. Miller, 1892, Advance Sheets
18th Rep. Geo. Sur. Ind., pp. 29 and 30,
Niagara Gr.

gorbyi, S. A. Miller, 1891, Advance Sheets
17th Rep. Geo. Sur. Ind., p. 57, Ni-
agara Gr.

SCAPHOCRINUS bellus, *S. bonoensis*, *S. dis-*
parilis, *S. granuliferus*, *S. graphicus*, *S.*
lacunosus, *S. manns*,
S. priemorsus, *S. re-*
pertus, Miller &
Gurley, 1890, Jour.
Cin. Soc. Nat. Hist.,
vol. 13, p. 24, and
Desc. New Gen. and
Spec. Echinoder-

mata, pp. 29, 45 to
52, republished in
16th Rep. Geo. Sur.
Ind., pp. 345, 349, 362
to 367, Keokuk Gr.

boonvillensis, *P. con-*
strictus, S. A. Miller,
1891, Bull. No. 4, Geo.
Sur. Mo., pp. 37, 38,
Keokuk Gr.

gorbyi, *S. porrectus*, *S.*
sampsoni, S. A. Mil-
ler, 1891, Advance

Sheets 17th Rep.
Geo. Sur. Ind., pp. 42, 46; *S. sampsoni*
from the Chouteau limestone, the others
from the Keokuk Gr.

FIG. 1245.—*Scaphocrinus gorbyi*.

Geo. Sur. Ind., pp. 42, 46; *S. sampsoni*
from the Chouteau limestone, the others
from the Keokuk Gr.

lyoni, and *S. maniformis*, S. A. Miller, 1892, Advance Sheets Geo. Sur. Ind., p. 45, Keokuk Gr.

FIG. 1244.—*Schenaster legrandensis*.

Sur. Ind., p. 371, Kinderhook Gr.
STEGANOCRINUS benedicti, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 27, Keokuk Gr.

FIG. 1247.—*Schenaster legrandensis*. Part of ventral side magnified $6\frac{1}{2}$ diameters.

STEPHANOCRINUS elongatus, S. hammelli, *S. obpyramidalis*, S. A. Miller, 1891, 17th Rep. Geo. Sur. Ind., pp. 22 to 26, Niagara Gr. *S. osgoodensis* is redescribed and the genus is discussed at the same place.
cornetti, S. A. Miller, 1892, Advance

FIG. 1248.—*Steganocrinus benedicti*.

Sheets 18th Rep. Geo. Sur. Ind., p. 12, Niagara Gr.

STRIBALOCYSTITES, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 20. [Ety. *stribalos*, close pressed, thick, in allusion to the thick tumid plates; *kustis*, bladder.] Body rudely

subovate or subelliptical, and covered by about five series of tumid plates.

FIG. 1249.—*Stribalocystites gorbyi*. Side, summit, and basal views.

Basals 4, unequal; second series of plates 6, unequal; fourth and fifth series irregular and covering the summit; no arms; orifice near the summit on the azygous side, and another on the left near the summit, both being above the third range of plates; sometimes there is a central orifice. Type *S. tumidus*, which is described at the same place, from the Niagara Gr.

gorbyi, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 11, Niagara Gr.

SYNBATHOCRINUS blairi, S. A. Miller, 1891, Bull. No. 4, Geo. Sur. Mo., p. 32, Keokuk Gr.

wachsmuthi, Meek & Worthen, 1869, Proc. Acad. Nat. Sci., p. 67, and Geo. Sur. Ill., vol. 5, p. 437, Burlington Gr.
TAXOCRINUS spinifer, Hall, 1861, Proc. Bost. Soc. Nat. Hist., p. 318, Burlington Gr.

FIG. 1251.—*axocrinus subovatus*.

TECHNOCRINUS spinulosus is the type of the genus.

TROOSTOCRINUS laterniformis, Owen & Shumard, 1850 (*Pentremites laterniformis*), Jour. Acad. Nat. Sci. 2d ser., vol. 2, p. 66, Kaskaskia Gr.

nitidulus, Miller & Gurley, 1890, Desc. New Gen. and Spec. Echinodermata, p. 58, and 16th Rep. Geo. Sur. Ind., p. 373, St. Louis Gr.

wachsmuthi, Gurley, 1884, New Carb. Foss. Bull. No. 2, p. 1, Warsaw Gr.

ULOCRINUS, Miller & Gurley, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 6, and 16th Rep. Geo. Sur. Ind., p. 330. [Ety. *oulos*, solid, substantial; *crinus*, lily.] Calyx globular or pyramidal, large, plates more or less convex, smooth or granular. Basals 5, forming a pentagonal disc or low cup; columnar opening pentagonal; subradials 5, very large; first radials large, pentagonal, upper face projects over the interior of the calyx so as to make a broad articu-

cal, and covered
of tumid plates.

rbyl. Side, summit,
ws.

second series of
fourth and fifth

1250. — *Stribalocys-
es tumidus*. Side
d basal views.

being above the
; sometimes there
Type *S. tumidus*,
at the same place,

2, Advance Sheets
Ind., p. 11, Niagara

S. A. Miller, 1891,
r. Mo., p. 32, Keo-

& Worthen, 1869,
ci., p. 67, and Geo.
437, Burlington Gr.
all, 1861, Proc. Bost.
c. Nat. Hist., p. 318,
Burlington Gr.

ovatus, Miller &
urley, 1890, Desc.
ew Gen. and Spec.
chinodermata, p. 26,
d 16th Rep. Geo.
r. Ind., p. 347,
eokuk Gr.

ocrinus spinulosus
the type of the
enus.

ormis, Owen & Shu-
mites lateriformis),
ci. 2d ser., vol. 2, p.

Gurley, 1890, Desc.
e. Echinodermata, p.
Geo. Sur. Ind., p. 373,

, 1884, New Carb.
p. 1, Warsaw Gr.
Gurley, 1890, Jour.
t., vol. 13, p. 6, and
r. Ind., p. 330. [Ety.
antial; *krinon*, lily.]
r pyramidal, large,
s convex, smooth or
p, forming a pentag-
cup; columnar open-
snbradials 5, very
s large, pentagonal,
s over the interior of
make a broad articu-

lating face for the first brachial; no
regular interradials. A quadrangular
azygous plate placed obliquely forms
part of the calyx, and a small plate

FIG. 1252.—*Uloerinus buttsi*.

rests upon its upper angle at the top of
the calyx, and projects slightly above
the top of the first radials; column
round. Type *U. buttsi*, described at
the same place, from the Upper Coal
Meas. *U. kansansensis* is also described
at the same place, from the same rocks.

ZEACRINUS commaticus, *Z. pocillum*, S. A.

Miller, 1891, Bull. No. 4, Geo. Sur. Mo.,
pp. 28, 36, Keokuk Gr.

dubius, Miller & Gurley, 1890, Desc. New
Gen. and Spec. Echinodermata, p. 44,
and 16th Rep. Geo. Sur. Ind.,
p. 361, Keokuk Gr.

iaggi, Kowley & Hare, 1891,
Kansas City Scientist, p. 103,
Burlington Gr.

ZOPHOCRINUS, S. A. Miller, 1891,
Advance Sheets 17th Rep. Geo.
Sur. Ind., p. 32. [Ety. *zophos*,
dark, obscure; *krinon*, lily.]
Body ovate or pear shaped,
and covered by two circles of
plates and the vault. Basals
or first circle of plates 3, form-
ing an obconoidal cup, higher
than wide; two of the plates
are of equal size, and quadran-

gular; the other is larger and pen-

FIG. 1254.—*Zophocrinus howardi*. Side and
summit views.

FIG. 1253.—*Uloerinus kansansensis*.

tagonal; second circle of plates 4,
3 pentagonal and 1 quadrangular;
they are horizontally truncated at
on top, and bear a circle of num-
erous pinnules surrounding a
convex vault. Seven plates con-
stitute the test of the calyx. It
possessed a column with a small
columnar canal. Type *Z. how-
ardi*, described at the same place,
from the Niagara Gr.

The paper entitled "Descriptions of New Species of Crinoidea, from investigations of the Iowa Geological Survey, Preliminary Notice, by James Hall," dated February 25, 1861, has never been published, as required by the laws of nomenclature, and is not, therefore, entitled to recognition. It is a private pamphlet, that was never kept for sale, and was not generally distributed among those conversant with the subject, and contains no figures of any of the organisms, which, with such meagre and imperfect descriptions as the text contains, would be absolutely necessary to enable an expert paleontologist to determine the organism intended to be named. Hall says, on page 10, that he published the abstract of the descriptions to get ahead of the publications of the Geological Survey of Illinois. A very selfish excuse for such poor work, and it does not add any vitality to the publication, which was never entitled to recognition under the laws of nomenclature, as shown on page 95 of this work. One genus and two species have been redefined in valid publications, and they will stand as of the date of the redefinition. The following names, never having appeared with valid definitions, should be stricken out, as of no more value than the names in a private catalogue: *Actinocrinus carica*, sometimes referred to *Eretmocrinus*; *A. ovatus*, *A. multibrachiatus* var. *echinatus*, *A. lucina*, *A. thetis*, *A. thos*, *A. quaternarius* var. *spiniferus*, *A. themis*; *A. remibrachiatus*, sometimes referred to *Eretmocrinus*; *A. tenuiradiatus*, sometimes referred to *Strotocrinus* or *Teleiocrinus*; *A. eryx*, *A. (Calathocrinus) erodus*, sometimes referred to *Strotocrinus* or *Teleiocrinus*; *A. (Calathocrinus) inculptus*, sometimes referred to *Strotocrinus* or *Teleiocrinus*; *A. (Calathocrinus) althea*, sometimes referred to *Teleiocrinus*; *A. lagena*, *A. thalia*; *A. maluta*, sometimes referred to *Eretmocrinus*; *A. maluta* var. *attenuata*, sometimes referred to *Eretmocrinus*; *A. (?) tenuidictus*, *A. securis*, *A. infrequens*, *A. locellus*, *A. doris*, *Platycrinus olla* (name preoccupied), *P. regalis*, *P. glyptus*, *P. calyculus*, *P. nodobrachiatus* (name preoccupied), *P. parvinodus*, *P. eminus*, *P. aqualis* (*P. aqualis* in Geo. Sur. Ill., vol. 5, p. 456), *Synbathocrinus papillatus*, *Rhodocrinus wachsmuthi*, *Heterocidaria keokuk*, and *H. levispinus*. All of the above, including *Lepidocrinus*, *L. imbricatus*, and *Protaster (?) barrisi*, which have been since defined, in proper publications, are

printed in the above named private pamphlet, on about nine pages. In other words, thirty-eight new species and two new genera are defined on about nine pages, without a single figure of any kind, and without mentioning a locality from which any of them came, or informing any one of the age of the rocks further than to write "Burlington limestone."

Wachsmuth (who has probably seen some of the so-called types) has condemned, as synonyms of other species, the above named *Actinocrinus multibrachia'us* var. *echinatus*, *A. thetis*, *A. thoas*, *A. quadrarius* var. *spiniferus*, *A. themis*, *A. eryx*, *A. lagina*, *A. securis*, *A. locellus*, *A. doria*, *Platycrinus olla*, *P. glyptus*, *P. calyculus*, *P. nodobrachiatus*, and *Synbathocrinus papillatus*.

Had I been able to see a copy of the pamphlet, none of the names would have appeared as valid in the first edition of this work; but none of my correspondents had ever seen one, and I borrowed most of the names from references made to them by Meek & Wachsmuth. More recently Wm. F. E. Gurley, of Danville, Illinois, has been able to obtain a copy of all except the last page of the pamphlet, and I have been allowed to examine it, with the above result.

SUBKINGDOM MOLLUSCOIDA.

CLASS BRYOZOA.

SOME genera were by accident placed in two families in this Class. Correct by striking *Peronopora* out of the *Batostomellidæ*; *Eridopora*, *Lichenotrypa*, *Sagenella*, and *Selenopora* from the *Ceramoporidæ*; *Coscinella*, *Reptaria*, and *Semiopora* from the *Fenestellidæ*; *Anisotrypa* from the *Rhabdomesontidæ*; *Heliotrypa* from the *Stictoporidae*; *Criscinella* from the *Thamniscidæ*; *Acanthoclema*, *Amplexopora*, *Atactopora*, *Bactropora*, *Chilotrypa*, *Nemataxis*, and *Tropidopora* from the *Trematoporidae*. Strike out the family *Labechiidæ* because it belongs, probably, to the Protozoa. Place *Semiopora* with the *Ptylodietyonidæ*; *Sagenella* and *Reptaria* with the *Tubuliporidae*.

The bryozoium is sometimes called *cœnœcium* (*koinos*, common; *oikos*, house) or *polyzoarium*, especially when *Polyzoa* is used for the Class instead of *Bryozoa*. *Gymnolæmata*, in the last line on page 289, is from, *gymnos*, naked; *laimos*, the throat.

ARTHROCLEMA *arimatum*, *A. cornutum* and *Arthrostylus conjunctus* and *A. obliquus*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 12, pp. 193 and 194 and pp. 189 and 190, Trenton Gr.

Ceramopora concentrica and *C. whitei*, James, 1888, Jour. Cin. Soc. Nat. Hist., vol. 11, p. 38. Not recognized.

CHAINODICTYON, at the top of p. 297, is from *chaino*, gaping; and *dictyon*, net.

DIASPORINA, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 12, p. 177. [Ety. diminutive of *Diastopora*.] Zoarium bifoliate, flabellate; zoecia tubular, prostrate, not entirely immersed; apertures constricted, oblique, the anterior side not elevated; interspaces, striated. Type *D. flabellata*, described, at the same time, from the Trenton Gr. He also described at the same time from the Trenton Gr. *Enallopora mundula* under the name of *Mitoclema mundulum*. He gives his reasons for not using the ge-

neric name *Enallopora* in vol. 8, Geo. Sur. Ill. p. 683.

DRYMOTRYPA, Ulrich, 1890, Geo. Sur. Ill., vol. 8, p. 399. [Ety. *drymos*, a thicket; *trypa*, perforation.] Zoarium dichotomous, reverse striated; zoecia in ranges, tubular, thick walled in tangential sections, springing from a thin plate; superficial apertures angular, oval within. Type *D. diffusa*, Hall's *Retepora diffusa*, and includes Hall's *Thamniscus cisseis* and *T. niagarensis*.

Fenestella hemitrypa, see *Hemitrypa proutiana*.

FISTULIPORA *laxata*, Ulrich, 1889, Micro-palæontology of Canada, p. 37, Hud. Riv. Gr. The genus *Fistulipora* is recorded by some authors to the Alcyonaria, because the larger pores increase by cœnenchymal gemmation, a method of increase said to be unknown among Bryozoa.

Glyptotrypa or p. 307, read *Glyptopora*.

In other words, the pages, without a which any of them write "Burlington

has condemned, as *ia'us* var. *echinatus*, *lag'na*, *A. recuris*, *rach'atus*, and *Syn-*

es would have ap-
pendents had ever
to them by Meek &
s, has been able to
been allowed to ex-

DIDA.

Class. Correct by
enotrypa, Sagenella,
and Semiopora from
teliotrypa from the
ema, Amplexopora,
ra from the Tremat-
gs, probably, to the
a and Reptaria with

on; *oikos*, house) or
instead of Bryozoa.
l; *laimos*, the throat.

opora in vol. 8, Geo.

890, Geo. Sur. Ill., vol.
umos, a thickset; *trupa*,
rium dichotomous, re-
cia in ranges, tubular,
tangential sections,
thin plate; superficial
oval within. Type D.
epora diffusa, and in-
mniscus cisseis and T.

see Hemitrypa prout-

Ulrich, 1889, Micro-
Canada, p. 37, Hud.
nus *Fistulipora* is re-
authors to the Alcy-
e larger pores increase
gemmation, a method
to be unknown among

07, read *Glyptopora*.

HELOPORA alternata, H. mucronata, Ulrich,
1890, Jour. Cin. Soc. Nat. Hist., vol. 12,
p. 192, Trenton Gr.

NEMATOPORA conferta, N. granosa, N. ovalis,
Ulrich, 1890, Jour. Cin. Soc. Nat. Hist.,
vol. 12, pp. 196, 197, Trenton Gr.

PACHYDICTYA emaciata, P. obesa, P. turgida,
Foerste, 1887, Bull. Denison Univ., vol.
2, pp. 162 to 165, Niagara Gr.

pumula, P. triserialis, Ulrich, 1890, Jour.
Cin. Soc. Nat. Hist., vol. 12, pp. 186, 187,
and in Micropaleontology, pp. 42, 43,
P. hexagonalis and P. magnipora, Tren-
ton Gr.

PALESCHARA quadrangularis, Nicholson, 1874,
(Chetetes quadrangularis.) Rep. Pal.
Prov. Ont., p. 61, Devonian.

PINACOTRYPA marginata, Whiteaves, 1892,
Cont. to Can. Pal., p. 278, Devonian.

POLYPORA manitobensis, Whiteaves, 1892,
Cont. to Can. Pal., p. 280, Devonian.

RHINIDICTYA exigua, R. humilis and R.
minima, Ulrich, 1890, Jour. Cin. Soc.
Nat. Hist., vol. 12, pp. 183 to 185, Tren-
ton Gr.

RHOMBOPORA multipora, Foerste, 1887, Bull.
Denison Univ., vol. 2, p. 72, Coal Meas.

STICTOPORELLA rigida, Ulrich, 1890, Jour.
Cin. Soc. Nat. Hist., vol. 12, p. 188,
Trenton Gr.

STOMATOPORA monilliformis, Whiteaves, 1891,
Cont. to Can. Pal., vol. 1, p. 212, De-
vonian; S. tenuissima, Utica Gr. and
S. turgida, Hud. Riv. Gr., Ulrich, 1890,
Jour. Cin. Soc. Nat. Hist., vol. 12,
pp. 175 and 176.

SUBRETEPORA halli and S. sublaxa, Ulrich,
1890, (Philloporina halli and P. sublaxa,) Jour.
Cin. Soc. Nat. Hist., vol. 12, pp.
179 and 181, Trenton Gr. Ulrich gives
his reasons for not using Subretepora
in vol. 8, p. 686, Geo. Sur. Ill.

VINELLA, Ulrich, 1890, Jour. Cin. Soc. Nat.
Hist., vol. 12, p. 173. [Ety. proper
name.] Zoarium attached to foreign
bodies, consisting of exceedingly slen-
der, ramifying, thread-like tubes, occa-
sionally radiately arranged; surface
faintly lined longitudinally; a row of
widely separated pores along the sur-
face of the tubes. Type V. repens, de-
scribed at the same time from the
Trenton Gr.

CLASS BRACHIOPODA.

ONE valve of the shells in this Class is always larger than the other. A line drawn vertically from the beak to the base will divide the shell into two equal parts. The flattened space between the beaks is called the hinge area or the cardinal area; the aperture in one of the beaks is called the foramen; and the triangular plate in front of the foramen and sometimes forming part of its circumference is called the deltidium, but in some genera it does not exist. When there are no teeth, as in Crania, the valves are held together by the adductor muscles. The shells are found in all Groups of rocks, from the Taconic to the most recent; but the Class seems to have reached its maximum development in the Devonian, and to have slowly declined since the Carboniferous age.

Beecher says the main characters common to the Class are, the bivalve shell, the pedicled or fixed condition, the animal composed of two pallial membranes intimately related to the shell, a visceral sac, and two arms or appendages near the mouth. The extreme range of variation does not eliminate any of these features, and consequently no univalve or multivalve forms are found, nor any strictly free swimming species, nor growths or modifications adapting the organism to a Pelagic life. All Brachiopods have a common form of embryonic shell called the protegulum. (Ety. *pro*, early; *tegos*, a covering.) The protegulum is semicircular or semielliptical in outline, with a straight or arcuate hinge-line, and no hinge area. A slight posterior gaping is produced by the pedicel valve being usually more convex than the brachial. The modifications noted are apparently due to accelerated growth, by which characters primarily nealagic become so advanced in the development of the individual as to be impressed finally upon the embryonic shell.

- Acrotlele dichotoma* refer to *Acrotreta* dichotoma.
- Acrotreta gultelmi* refer to *Discinopsis gultelmi*.
- Anoplia*, Hall, 1892, Pal. N. Y., vol. 8, p. 309. Proposed as a subgeneric name with *Leptaena* (?) nucleolata, Hall, as the type. It is only necessary to say it has no place in nomenclature as a subgeneric name. If it is a name which deserves retention, it should be in a generic sense, and probably it should be placed in that rank.
- ATHYRIS angelica* var. *occidentalis*, Whiteaves, 1891, Cont. to Can. Pal., vol. 1, p. 227, Devonian.
- ashlandensis*, Herrick, 1888, Bull. Denison Univ. vol. 4, p. 24, Waverly Gr.
- brittsi*, and *A. ottervillensis*, S. A. Miller, 1892, 18th Rep. Geo. Sur. Ind., p. 60, Ham. Gr.
- ATRYPA calvini*, and *A. reticularis*, var. *niagarensis*, Nettleroth, 1889, Kentucky Foss. Shells, pp. 89, 92, Niagara Gr.
- deflecta*, Hall, see *Zygospira deflecta*.
- ellipsoidea*, Nettleroth, 1889, Kentucky Foss. Shells, p. 90, Up. Held Gr.
- missouriensis*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 61, Ham. Gr.
- Aulosteges spondyliiformis*, see *Strophalosia spondyliiformis*.
- BARROISELLA*, Hall, 1892, Pal. N. Y., vol. 8, p. 62. [Ety. proper name.] Lingula-like shell; pedicel-valve bears a high cardinal area, appearing as a thickened triangular plate, divided by a broad pedicel-groove, and having at each basal angle a boss or condyle which served as a muscular fulcrum or as a point of articulation; the interior has a subquadrate depressed area in continuation of the pedicel-groove, and from its antelateral angles diverge two linear depressions, which extend about one-fourth the length of the shell; from outside and behind the extremities of these depressions begins a pair of long, curved furrows, composed of two shorter curves, the posterior rounding over the extremities of the linear depressions; the anterior and longer curves gradually approximate, and nearly meet at about one-third the length of the shell from the anterior margin; these furrows are accompanied by low ridges along their inner margin. A low median ridge, with elevated edges, begins at the posterior umbonal impression, and continues to the center of the valve, widening near its anterior extremity; behind its termination there is a pair of indistinct muscular impressions. In the brachial valve the beak is scarcely prominent, and the muscular markings are essentially as in the opposite valve, but more sharply developed; beneath the beak there is a faint umbonal scar. The long compound lateral curves have a greater degree of curvature than in the pedicel-valve, and their posterior portion incloses a thickened area, which is continued into a low median septum that bifurcates in the middle of the valve, and has extending from the angle an intercalated ridge. Type *Lingula subspatulata*, Meek & Worthen.
- BILLINOSELLA*, Hall, 1892, Pal. N. Y., vol. 8, p. 230. [Ety. proper name.] Shell Orthis-like, transverse; subquadrate or semicircular in outline; contour concavo or plano convex; surface striate; pedicel-valve the more convex; cardinal area moderately high, vertical or slightly incurved; delthyrium covered by a convex plate, which may be minutely perforated at the apex; teeth well developed, but dental plates are continued along the bottom of the umbonal cavity, inclosing a small subelliptical muscular area near the apex. In the brachial valve the cardinal area is greatly inclined, making an obtuse angle with that of the opposite valve; delthyrium partly covered by a convex deltidium, which never attains the development seen in the opposite valve, and is sometimes absent. Type *Orthis pepina*, Hall. It probably includes *Streptorhynchus primordiale*, Whitfield, *Orthis grandeva* and *Orthis laurentina*, Billings.
- CAMARELLA bernensis*, and *C. owatonnaensis*, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 328, Trenton Gr.
- minor, Walcott, 1890, 10th Ann. Rep. U. S. Geo. Sur. p. 614, Up. Taconic.
- CHONETES mantobensis*, Whiteaves, 1892, Cont. to Can. Pal., p. 281, Devonian.
- subquadrate, Nettleroth, 1889, Kentucky Foss. Shells, p. 67, Up. Held Gr.
- tumida, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 36, Waverly Gr.
- complanata*, *dawsoni*, *reversa*, refer to *Chonostrophia*.
- CHONOSTROPHUS*, Hall, 1892, Pal. N. Y., vol. 8, p. 312. [Ety. *chonos*, a cup; *pektos*, combed.] Shell like *Chonetes*, the cardinal margin of the pedicel-valve bearing a row of erect spines; beak compressed, leaving a flattened area or cicatrix from attachment in early growth; surface ornamented with a double series of concentric lines or wrinkles, having the appearance of the engraving on a machine-turned watch-case, and strongest on the umbonal and central part; these wrinkles are crossed by concentric growth-lines, and sometimes by finer radiating lines more or less flexuous. Casts of the pedicel-valve show the impression of a short median septum dividing two broad obcordate flabelliform muscular scars, from the outer margin of which there are radiating vascular scars; impressions of a narrow cardinal area and exceedingly small teeth also occur on

curvature than in and their posterior thickened area, which low median septum the middle of the extending from the d ridge. Type *Lin-*

2, Pal. N. Y., vol. 8, per name.] Shell rae; subquadrate or line; contour convex; surface striate; ore convex; cardinal high, vertical or delthyrium covered which may be mit at the apex; teeth at Dental plates are e bottom of the uming a small subellipnear the apex. In the cardinal area is making an obtuse the opposite valve; covered by a convex never attains the n the opposite valve, absent. Type *Orthis* probably includes imordiale, Whitfield, a and *Orthis lau*

and *C. owatonsensis*, ill. Minn. Acad. Nat. Trenton Gr.

90, 10th Ann. Rep. 614. Up. Taconic. is, Whiteaves, 1892, p. 281, Devonian.

oth, 1889, Kentucky Up. Held. Gr.

888, Bull. Denison, Waverly Gr.

reversa, refer to Cho-

92, Pal. N. Y., vol. 8, *onos*, a cup; *pektos*, like *Chonetes*, the car-e pedicle-valve bear-et spines; beak com-a flattened area or attachment in early ornamented with a concentric lines or the appearance of the achine-turned watch- on the umbonal and e wrinkles are crossed with-lines, and some-diating lines more or asts of the pedicel-mpression of a short dividing two broad orm muscular scars, argin of which there ular scars; impres- cardinal area and teeth also occur on

the internal cast. Type *Chonetes fischeri*, Norwood & Pratt.

CHONOSTROPHIA, Hall, 1892, Pal. N. Y., vol. 8, p. 310. [Ety. *chonos*, a cup; *strophe*, turning around.] Shell like a reversed *Chonetes*, concavo-convex, the pedicel-valve being slightly convex about the umbo, but becoming broadly concave over the pallial region; outline and contour like *Leptena*; valves extremely tenuous and compressed; surface covered with fine, alternating or fasciculate striae. In the pedicel-valve the upper margin of the cardinal area bears a row of hollow spines of the same structure and arrangement as in *Chonetes*. The delthyrium is narrow, and appears to be more or less completely closed. The teeth are quite strong, and rest upon the bottom of the valve; between them arises a low median septum, extending one-third or one half the length of the valve, dividing a subcordate muscular area, the outer margins of which are distinctly elevated. In the brachial valve the crural plates are united to form a bilobed cardinal process. On the inner surface it ends abruptly; internal pallial region finely papillose; shell-substance fibrous, punctate. Type *Chonetes reversa*, Whitfield. To the same genus is referred *Chonetes complanata*, *C. dawsoni*, and *C. holderbergia*, Hall, 1892, Pal. N. Y., vol. 8, p. 353, Low. Held. Gr.

CHRISTIANA, Hall, 1892, Pal. N. Y., vol. 8, p. 298. [Ety. proper name.] Shell resembling *Leptena*, usually longitudinally elongated, sometimes semielliptical in outline; normally concavo-convex; surface smooth or covered with fine radiating lines, crossed by stronger concentric plications; cardinal area of the pedicel-valve high; delthyrium probably closed by a convex plate; teeth divergent, and from their bases extend the elevated margins of two linguiform muscular scars, traversing the shell almost the entire length; these diductor scars inclose two elongate adductors. In the brachial valve the cardinal process is bipartite, each of the lobes being grooved behind; the crural plates are long and divergent, terminating in elevated extremities or crura. The lower moiety of these plates is produced on each side of an elevated muscular ridge, curving slightly inward on the sides, then outward on approaching the anterior margin of the valve, each branch recurving and passing backward, parallel to the median axis, as far as the base of the cardinal process. The interspaces are divided transversely at about one-third their length from the hinge-line, by a lower ridge. The four areas thus inclosed represent the posterior and anterior scars of the

adductor muscles. Type *Leptena subquadrata*, Hall.

CLITAMBONITES, Pander, 1830, Beitrage zur Geognosie des russ. Reiches, p. 70. [Ety. *klitos*, a sloping place; *ambon*, any rising; *tilhos*, stone.] Shells with a sub-semicircular marginal outline; convex or subpyramidal; hinge-line straight, and forming the greatest diameter of the shell; pedicel-valve elevated; cardinal area high, vertical, or sometimes incurved and crossed by a broad delthyrium, with a perforate deltidium; dental lamelle strongly developed, converging and uniting in the median line; median septum about half the length of the valve; muscular impressions obscure; cardinal area developed in the brachial valve, delthyrium filled with a callosity; dental sockets large; crural plates low; thickened transverse area in the umbonal region; surface striate; shell-substance impunctate. Type *Pronites adscendens*, Pander. It includes *Orthisina vernaulti*, Eichwald, and *Hemipronites americanus*, Whitfield.

CONOTRETA, Walcott, 1890, Advance Sheets Biological Soc. Washington. [Ety. *konos*, cone; *tretos*, perforated.] Structure calcareo corneous. Five narrow ridges radiate from the apex toward the front, or the internal side of the dorsal or conical valve, the central one of which joins the thickened apex, which is supposed to have been perforated by a siphonal tube. In *Acrotreta* there is an elongated muscular scar extending from each side of the siphonal tube obliquely forward. Type *C. rusti*, which is described at the same place from the Trenton Gr.

CRANIA agaricina, *C. pulchella*, Hall, 1892, Pal. N. Y., vol. 8, p. 180, Low. Held. Gr., and *C. granosa* and *C. favincola* at same place, Ham. Gr.

blairi, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 56, Chouteau limestone, and *C. greenii* from Up. Held. Gr.

columbiana, Walcott, 1888, Proc. U. S. Nat. Mus. p. 441, Up. Taconic. Not defined so as to be recognized.

halli, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 328, Trenton Gr.

radicans, see *Strophalosia radicans*.

CRANIELLA (Ehler, 1888, Bull. de la Soc. d'Etudes Scientif. d'Angers, p. 37. [Ety. diminutive of *Crania*.]) Shell somewhat irregular; outline subcircular or subquadrangular; ventral valve thin, adhering by its entire surface; dorsal valve conoidal, more or less elevated; apex subcentral, posterior; interior of the dorsal valve without a well-defined border; impressions of the adductors large, distinct, four in number, of which the posterior two are quite distant, the two subcentrals smaller, closely approximate or even confluent; from near each

- of the posterior impressions starts a broad vascular sinus, strongly sinuous near its point of departure, narrowing gradually in following the contour of the valve, emitting from its marginal side dichotomizing secondary branches. Type *C. medianensis*, Ehler. It probably includes *Crania hamiltonia*, Hall, and *C. greenii*, S. A. Miller.
- ulrichi, Hall, 1892, Pal. N. Y., vol. 8, p. 181, Trenton Gr.
- CRYPTONELLA ovalis, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 76, Ham. Gr.
- traversensis, Winchell, 1866, (Terebratula traversensis,) Rep. Low. Penin. Mich., p. 95, Ham. Gr.
- Derbya, Waagen, 1884, Paleontologica Indica, Ser. 13, vol. 1, pp. 576, 591 to 607. [Ety. proper name.] This genus, according to Hall, is distinguished from *Streptorhynchus* by the presence of a median septum in the pedicel valve. He notes no other generic difference. Type *D. regularis*, Waagen. Hall refers to this genus *Orthis keokuk*, Hall, *Orthis robusta*, Hall, *Hemipronites* (*Streptorhynchus*) *crassus*, Meek & Hayden, and regards *Hemipronites lasallensis* and *H. richmondi*, McChesney, as synonyms for *H. crassus*. I consider *Orthis robusta*, Hall, more closely related to *Hemipronites crassus*, Meek, than either *H. lasallensis* or *H. richmondi*, McChesney, and if one is to be retained as a species all three should be, and in no event is either one of them congeneric with *Orthis keokuk*. Hence, I see no propriety in the use of the word *Derbya* as applied to American fossils, and as I have seen no typical specimens belonging to the genus, I express no opinion as to its value. Hall, however, describes *Derbya broadheadi*, *D. bennetti*, *D. cymbula*, *D. affinis*, and *D. biloba*, from the Up. Coal Meas., Pal. N. Y., vol. 8, pp. 347 to 350, and on p. 346 *D. ruginosa*, from the Keokuk Gr., and *D. costatula* from the Kaskaskin Gr. All of these may be referred to *Streptorhynchus*, and it is not too much to say that Meek would have classed all of them as synonyms for *Streptorhynchus crassus*.
- DISCINA concordensis, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 328, Trenton Gr.
- keokuk, Gurley, 1884, New Carb. Foss., p. 6, Keokuk Gr.
- sampsoni, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 80, Chouteau limestone.
- DISCINOPSIS, Matthew, 1892, Pal. N. Y., vol. 8, p. 105. [Ety. from resemblance to Discina.] Shell subcircular, depressed conical; apex excentric; apex of the pedicel-valve truncated, with a circular aperture; interior having a pair of diverging furrows from the beak, that converge toward the anterior margin, and inclose a thickened area, which, in the subumbonal region, is apparently free and projects like a shelf, from beneath which the foramen probably opened; shell substance tenuous, apparently corneous; surface marked concentrically, and also with radiating striae. Type *Acrotreta gullelmi*, Matthew.
- EATONIA variabilis, Whiteaves, 1891, Cont. to Can. Pal., vol. 1, p. 233, Devonian.
- Hallina, Winchell & Schuchert, 1892, Am. Geo., vol. 9, p. 291. [Ety. proper name.] Shells small, articulate, rostrate, biconvex, semiplicate; pedicel opening bounded laterally by incomplete deltidial plates; calcified brachial supports long and much like Waldheimia. Type *H. saffordi*, described at the same place with *nicolleti*, from Trenton Gr. Neither one defined so as to be recognized.
- Koninekina americana, Swallow, may be stricken from the list as it is not a Koninekina, and is too poorly defined to be recognized.
- KUORGINA labradorica var. swantonensis, Walcott, 1889, Proc. Nat. Mus., vol. 12, p. 36, Up. Taconic.
- LEPTENA charlotte, Winchell & Schuchert, 1892, Am. Geo., vol. 9, p. 288, Trenton Gr., and *Plectambonites gibbosus*, from Galena Gr. Not defined so as to be recognized.
- minnesotensis, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 329, Trenton Gr., and *præcosis*, *recedens* and *saxea*, Hud. Riv. Gr.
- LEPTENISCA, Beecher, 1890, Am. Jour. Sci. and Arts, Ser. 3, vol. 40, p. 239. [Ety. diminutive of Leptæna.] Shell concavo-convex, attached to foreign objects by calcareous cementation of the ventral beak; valves articulated by teeth and sockets; dorsal valve concave; interior with broad spiral impression on each side of the median line, making a single volution; adductor impressions small; cardinal line narrow, bearing in the center two prominent, bilobed, cardinal processes, separated to admit the vertical septum in the opposite beak; ventral valve convex, area elongate, triangular; fissure covered with a pedicel sheath; cardinal muscular scar supported on or limited by two elevated lamellæ; cavity of beak divided by a vertical septum, on each side of which, in the anterior half, is a small, adductor scar; shell structure punctate. Type *L. concava*.
- adnascens and *L. tangens*, Hall, 1892, Pal. N. Y., vol. 8, p. 352, Low. Held Gr.
- concava, Hall, 1857, (Leptæna concava,) 10th Rep. N. Y. St. Nat. Hist., p. 47, and Pal. N. Y., vol. 3, p. 197, Low. Held Gr.
- LEPTELLA, Hall, 1892, Pal. N. Y., vol. 8, p. 293. [Ety. leptos, thin.] Shell small, concavo-convex, semicircular or semi-

area, which, in the is apparently free shelf, from beneath probably opened; nuous, apparently marked concentric-diating striae. Type Matthew.

teaves, 1891, Cont. p. 233, Devonian. Schuchert, 1892, Am. Ety. proper name.] ate, rostrate, bicon-pedicle opening by incomplete deltidial brachial sup-like Waldheimia. described at the same from Trenton Gr. so as to be recog-

Swallow, may be ist as it is not a too poorly defined

var. swantonensis, Nat. Mus., vol. 12,

chell & Schuchert, 9, p. 288, Trenton mites gibbosus, from defined so as to be

on, 1892, Bull. Minn. 3, p. 329, Trenton ecedens and saxeae,

890, Am. Jour. Sci. 40, p. 239. [Ety. na.] Shell concavo-foreign objects by tion of the ventral lated by teeth and ve concave; internal impression on dian line, making a ductor impressions narrow, bearing in ninent, bilobed, car- arated to admit the the opposite beak; ex, area elongate, covered with a ped- muscular scar sup- ed by two elevated beak divided by a each side of which, is a small, adductor re punctate. Type

ens, Hall, 1892, Pal. Low. Held Gr.

Leptaena concava,) Nat. Hist., p. 47, 3, p. 197, Low.

Pal. N. Y., vol. 8, thin.] Shell small, micircular or semi-

elliptical; hinge-line straight; pedicel-valve evenly convex; cardinal area moderately high, delthyrium nearly covered by a convex plate; teeth inconspicuous; brachial valve slightly concave; cardinal area high, delthyrium filled with the cardinal process which is divided behind by a median groove; on the cardinal margin this process is double but less conspicuous than the crural plates, which are arched and highly elevated above the hinge-line; they are short, terminate abruptly, and inclose deep sockets; visceral area flattened or concave; its anterior margin forming a double visceral area is divided by a broad median ridge, and its surface covered with five sharp radiating lines, which end at the line of deflection. Type *Leptaena sordida*, Billings.

LINDSTROMELLA, Hall, 1892, Pal. N. Y., vol. 8, p. 134. [Ety. proper name.] Shells with outline contour and pedicel characters as in Orbiculoidea; brachial valve with a faint median septum and two strong approximating ridges, beginning behind a transverse line passing through the apex, and rapidly converging to meet the median septum; anterior adductor scars lying between these ridges and the median septum; a circular scar at the posterior extremity of each ridge. Type *L. aspidium*, p. 178, Ham. Gr.

LINGULA atra, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 16, Waverly Gr.

deflecta, Winchell & Schuchert, 1892, Am. Geo., vol. 9, p. 284, Galena Gr. Not defined so as to be recognized.

exilis, see Lingulodiscina exilis. gannensis, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 17, Waverly Gr.

gorbyi, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 55, Chouteau limestone.

macconnelli, Walcott, 1888, Proc. U. S. Nat. Mus., p. 441, Up. Taconic. Not properly defined.

meeki, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 18, Waverly Gr.

norwoodi, see Lingulops norwoodi.

parrishi, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 53, Up. Coal Meas.

riciniformis var. galenensis, Winchell & Schuchert, 1892, Am. Geo., vol. 9, p. 284, Galena Gr. Not defined so as to be recognized.

sedaliensis, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 54, Chouteau limestone.

subspatulata, refer to Barroisella subspatulata.

tighti, Herrick, 1887, Bull. Denison Univ., vol. 2, p. 43, Coal Meas.

triangulata, Nettleroth, 1889, Ky. Foss. Shells, p. 34, Up. Held. Gr.

waverlyensis, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 18, Waverly Gr.

LINGULELASMA galenensis, Winchell & Schuchert, 1892, Am. Geo., vol. 9, p. 285, Galena Gr. Not defined so as to be recognized. Ulrich, Schuchert and others persist in using the word Lingulasma, which has no more meaning than Lasma or Linguma would have. I spelled the word Lingulelasma, as it should have been coined, supposing the misspelling was accidental; but, probably, Mr. Ulrich's spelling should be recognized, and if so, then the word should be stricken from science, as a meaningless compound, under rules on page 98.

LINGULODISCINA, Whitfield, 1890, Bull. Am. Mus. Nat. Hist., p. 121. [Ety. *Lingula* and *Discina*.] Upper valve linguloid in character, having a terminal beak, the accretions of growth being along the lateral and basal margins; lower valve having its growth-lines nearly equal on all sides of the initial point and perforated on the cardinal side by a byssal slit as in Discina; shell structure as in Lingula and Discina. Type *L. exilis*. *exilis*, Hall, 1860, (*Lingula exilis*), 13th Rep. N. Y. St. Mus. Nat. Hist., p. 77, and Pal. N. Y., vol. 4, p. 7, Ham. Gr.

LINGULOPS granti, Hall, 1892, Pal. N. Y., vol. 8, p. 173, Niagara Gr.

norwoodi, instead of Lingula norwoodi.

MIMULUS, Barande, 1879, Systeme Silurien du Centre de la Boheme, vol. 5, p. 109. [Ety. *mimulus*, a mime, an imitator.] *Spirifera* (?) *waldronensis*, Miller & Dyer, was referred to *Triplesia* by Hall, without a knowledge of the interior, and he now refers it to *Mimulus*, though the internal characters of that genus are unknown.

MONOMERELLA egani, M. greenii, M. kingi, M. ortonii, Hall, 1892, Pal. N. Y., vol. 8, pp. 174, 175, Niagara Gr.

NEWBERRIA, Whiteaves, 1891, Cont. to Can. Pal., vol. 1, p. 236. [Ety. proper name.] Shells elongate-ovoid, having the general contour and external aspect of Rensseleria and Amphigenia, but without the strongly radiate striate surface of the former genus. The convexity of the valves is greatest in the umbonal region, and the surface is distinctly flattened over the lateral slopes, leaving the median portion of the valves very prominent; the pedicel valve has the rostrum produced and incurved; the apex truncated by a circular foramen; deltidial plates not determined; the teeth are comparatively small, projecting forward and gently upward, free at their extremities, and supported by narrow dental plates which join the interior of the valve above the middle of its depth, and are continued forward as slender ridges upon the inner surface, which gradually merge into the shell. In the bottom of the rostral and umbonal cavity there is a broad, scarcely defined,

- muscular area, from which radiate a series of vascular ridges and depressions; the strongly marked pair of adductors are situated posteriorly, just within this muscular area; lying in front of these is a single (rarely divided) elongate adductor impression which often extends forward to the center of the shell; on each side of the muscular impressions is a thickened triangular area, very narrow at its origin in the umbonal region, widening anteriorly and produced into two divergent furrows (four in all), which extend over the pallial region, in some instances almost to the anterior margin. In the brachial valve there are two short, divergent, crural plates, which are not united at their bases to form a hinge-plate, as in *Rensseleria*; a low median ridge arises between them, passing for a short distance along the bottom of the valve, separating the obovate, narrowly flabelliform muscular scars of the adductor muscles. These scars are characterized by the strong striation of their surfaces; surface smooth, or with obscure radiating striae; distinguished from *Rensseleria*, which has strong radiating striae on the surface, and preserves two broad, strong, dental plates on the interior of the pedicel-valve, which reach nearly to the bottom of the rostral and post umbonal cavity, leaving a narrow space for the muscular area, quite unlike that of the corresponding valve of *Newberria*. It is from this narrow cavity, produced by the encroachment of these strong dental plates, that we have the narrow rostral casts of *Rensseleria*. The thickened strong hinge-plate, which supports the crura in the brachial valve of *Rensseleria*, does not exist in *Newberria*; the spoon-shaped process found in *Amphigenia* does not exist in *Newberria*. Type *N. johanni*.
- condoni*, McChesney, 1867, (*Rensseleria condoni*), Trans. Chi. Acad. Sci., vol. 1, p. 36, Oriskany Gr.
- claypolei*, Hall, 1891, Advance Sheets 10th Ann. Rep. N. Y. St. Geo., Ham. Gr.
- johanni*, Hall, 1867, (*Rensseleria johanni*), Pal. N. Y., vol. 4, p. 385, Up. Held. Gr.
- levis*, Meek, 1868, (*Rensseleria levis*), Trans. Chi. Acad. Sci., p. 108, Devonian.
- missouriensis*, Swallow, 1891, Advance Sheets 10th Ann. Rep. N. Y. St. Geo., Ham. Gr.
- NUCLEOSPIRA indianensis*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 79, Ham. Gr.
- OBOLELLA misera* and *O. pretiosa*, refer to the genus *Linnarssonella*.
- OEHLERTELLA*, Hall, 1892, Pal. N. Y., vol. 8, p. 132. A proposed subgeneric name to include *Discina pleurites*.
- ORBICULOIDEA herzeri*, Cuyahoga Shales, *O. numulus*, Low. Held. Gr., and *O. ovals*, Trenton Gr., Hall, 1892, Pal. N. Y., vol. 8, pp. 177, 178.
- ORTHIS arcuaria*, Hud. Riv. Gr.; *O. holstoni*, *O. loricula*, *O. saffordi*, Trenton Gr.; *O. cweni*, Keokuk Gr.; *O. senecta*, Clinton Gr., and *O. superstes*, Chemung Gr., Hall, 1892, Pal. N. Y., vol. 8, pp. 340 to 342.
- benedicti*, S. A. Miller, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 78, Niagara Gr.
- germana*, Winchell & Schuchert, 1892, Am. Geo., vol. 9, p. 290, Galena Gr.
- neesai*, Trenton Gr.; *proavita*, Hud. Riv. Gr. Not defined so as to be recognized.
- goodwini*, Nettleroth, 1889, Kentucky Foss. Shells. p. 39, Up. Held. or Ham. Gr.
- inequalis*, see *Streptorhynchus inequalis*.
- linneyi*, Nettleroth, 1889, Kentucky Foss. Shells, p. 41, Hud. Riv. Gr.
- manitobensis*, Whiteaves, 1892, Cont. to Can. Pal., p. 283, Devonian.
- missouriensis*, Swallow, is the same described afterward as *O. theimii*.
- corpulenta*, *macrior*, *rogata*, *tersus*, *minnesotensis*, *petric*, *Sardeson*, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, pp. 330, 332, Trenton & Hud. Riv. Gr. The first four seem to be synonyms for *O. testudinaria*, but possibly they are not.
- Orthisina alberta*, Walcott, 1883, Proc. U. S. Nat. Mus., p. 442, Up. Taccnic. Not properly defined.
- Paterina* is a generic name proposed for such shells as *Kutorgina labradorica*, without any distinct generic characters being pointed out. See Am. Jour. Sci. and Arts, Ser. 3, vol. 41, p. 343.
- PENTAMERELLA thusnelda*, Nettleroth, 1889, Kentucky Foss. Shells, p. 51, Up. Held. Gr.
- PENTAMERUS colletti*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 77, Waterlime Gr.
- complanatus*, Nettleroth, 1889, Kentucky Foss. Shells, p. 53, Niagara Gr.
- decussatus*, Whiteaves, 1891, Can. Record Sci., p. 295, Up. Sil.
- globulosus*, Nettleroth, 1889, Kentucky Foss. Shells, p. 54, Niagara Gr.
- knotti*, Nettleroth, 1889, Kentucky Foss. Shells, p. 56, Niagara Gr.
- uniplicatus*, Nettleroth, 1889, Kentucky Foss. Shells, p. 63, Niagara Gr.
- PHOLIDOPS calceola*, P. patina, Hall, 1892, Pal. N. Y., vol. 8, p. 182, Corniferous limestone.
- POLYTUCHIA*, Hall, 1892, Pal. N. Y., vol. 8, p. 230. [Ety. *polus*, many; *toichos*, the wall of a house.] Shell small, subtriangular in contour; hinge-line straight; pedicel-valve with a high, nearly vertical cardinal area, obliquely striated; delthyrium covered with a convex plate; dental lamellae widely separated, descend for a distance vertically, and then bend inward to the median line,

Gr., Hall, 1892, Pal. 77, 178.
 Riv. Gr.; O. holstoni, Fordi, Trenton Gr.; k Gr.; O. senecta, superstes, Chemung. N. Y., vol. 8, pp. 177, 178.
 ler, Advance Sheets. Ind., p. 78, Ni.
 & Schuchert, 1892, p. 290, Galena Gr.; Gr.; *proavita*, Hud. ed so as to be recog-
 h, 1889, Kentucky 30, Up. Held. or
 orhynchus inaequale. 1889, Kentucky Foss. Riv. Gr.
 aves, 1892, Cont. to Devonian.
 w., is the same des- O. thelmii.
 rogata, tersus, min- Sardeson, 1892, Bull. Sci., vol. 3, pp. 330, d. Riv. Gr. The first mononyms for O. testu- ly they are not.
 ott, 1883, Proc. U. S. Up. Taccnic. Not
 name proposed for: utorgina labradorica, et generic characters. See Am. Jour. Sci. ol. 41, p. 343.
 lida, Nettleroth, 1889, Shells, p. 51, Up.
 A. Miller, 1891, Ad- Rep. Geo. Sur. Ind., r.
 roth, 1889, Kentucky Niagara Gr.
 es, 1891, Can. Record
 th, 1889, Kentucky Niagara Gr.
 889, Kentucky Foss. ara Gr.
 th, 1889, Kentucky Niagara Gr.
 patina, Hall, 1892, p. 182, Corniferous
 2, Pal. N. Y., vol. 8, s, many; *toichos*, the Shell small, subtri- hinge-line straight; a high, nearly verti- obliquely striated; ed with a convex llae widely separated, tance vertically, and to the median line,

forming with the deltidium a subrostral vault; the inner spoon-shaped plate, spondylium, is supported by a stout median septum and two smaller lateral septa, which meet it at the lines of angulation; the umbonal cavity is divided into five chambers; brachial valve shallow, depressed-convex; cardinal area narrow; delthyrium broad; dental sockets widely separated; crural plates narrow and nearly parallel to the hinge-line; cardinal process simple, linear, prominent, and at its union with the crural plates there is a subtriangular thickening supported by a median septum; surface striated; no fold or sinus. Type P. apicalis, Whitfield, described at the same place from the Calcliferous Gr.
 PRODUCTELLA minneapolis, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 332, Trenton Gr. It does not belong to this genus.
 pyxidata is from the Kinderhook Gr. semiglobosa, Nettleroth, 1889, Kentucky Foss. Shells, p. 70, Up. Held. Gr.
 PRODUCTUS blairi, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 79, Chouteau limestone.
 nodocostatus, P. raricostatus and P. rush- villensis, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 19 to 23, Waverly Gr. pileiformis, McChesney, 1859, New Pal. Foss., p. 40, Kaskaskia Gr.
 PSEUDOCRANIA anomala, Hall says, is a mis- nomer, that the shell described is a streptorhynchoid, whatever that means.
 RENNELERIA condoni, R. johanni and R. levis, refer to Newberria.
 RETZIA ashlandensis, Herrick, 1888, (Rhynchospira ashlandensis,) Bull. Denison Univ., vol. 3, p. 25, Waverly Gr. circularis, R. plicata and R. triangularis, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 61, Chouteau limestone.
 sobrina, Beecher & Clark, 1889, Mem. N. Y. St. Mus., p. 61, Niagara Gr.
 RHINOBOLOS, (misspelled Rhynobolus,) Hall has shown, in Pal. N. Y., vol. 8, p. 44, why this genus is distinct from Trimerella, and that it should be restored with R. galtensis as the type, and he defined R. davidsoni from the Niagara Gr.
 RHYNCHONELLA alleghania, Williams 1887, Bull. No. 41, U. S. Geo. Sur., p. 87, Waverly Gr.
 belliformis, R. louisvillensis, R. rugicosta, R. saffordi var. depressa, R. tenuistri- ata, Nettleroth, 1889, Kentucky Foss. Shells, pp. 73 to 82; louisvillensis and tenuistriata from the Up. Held. Gr., the others from the Niagara Gr.
 colletti, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 57, Niagara Gr., and R. kokomoensis from the Waterlime Gr.
 levis, R. medialis, R. striata, Simpson, 1889, Trans. Am. Phil. Soc., pp. 443,

444; levis from the Clinton Gr., the others from the Waverly Gr.
 minnesotensis and sancta, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 333, Trenton Gr.
 Rhynchospira ashlandensis, see Retzia ash- landensis.
 ROEMERELLA, Hall, 1892, Pal. N. Y., vol. 8, p. 137. Proposed as a subgenus to in- clude Discina grandis.
 SKENIDIUM, (should be spelled Scenidium,) anthonense, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 333, Trenton Gr.
 SCHIZOCRANIA, helderbergia, Low. Held. Gr. S. schucherti, Hud. Riv. Gr., Hall, 1892, Pal. N. Y. vol. 8, p. 179.
 SIPHONOTRETA minnesotensis, Hall, 1892, Pal. N. Y., vol. 8, p. 177, Trenton Gr.
 SPIRIFERA byrnesi, S. davisii, S. dubia, S. foggi, S. hobbsi, S. knappana, S. macconathi, Nettleroth, 1889, Kentucky Foss. Shells, pp. 109 to 122; foggi from the Niagara Gr., the others from the Up. Held. Gr. capax is from the Kinderhook Gr., and is not a synonym for S. parryana.
 carteri was so poorly defined that it could not be recognized, and it is now claimed by some that it is the same as Syringothyris typus. The name should be dropped from the list.
 deltoideus, S. tenuispinus, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 27, Waverly Gr.
 disjuncta var. occidentalis, Whiteaves, 1891, Cont. to Can. Pal., vol. 1, p. 222, Devonian.
 subventricosa, McChesney, 1860, New Pal. Foss., p. 44, 1865, pl. 1, fig. 4, Coal Meas.
 texta, see Syringothyris texta.
 winchelli, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 46, Waverly Gr.
 SPIRAIFERA depressa, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 47, Wa- verly Gr.
 STREPTORHYNCHUS inaequale, Hall, 1858, (Orthis inaequalis,) Geo. Iowa, p. 490, Kinderhook Gr.
 subaulcatum, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 335, Trenton Gr.; desideratum, (Orthothetes desider- atus,) Hall, 1892, Pal. N. Y., vol. 8, p. 345, Waverly Gr., and S. ulrichi, p. 351, Kaskaskia Gr.
 STRICKLANDINIA louisvillensis, Nettleroth, 1889, Kentucky Foss. Shells, p. 63, Ni- agara Gr.
 subquadrata, Herrick. Not properly de- fined.
 STRINGOCEPHALUS burtoni, DeFrance, 1827, Dict. des Sci. Naturelles, vol. 51, p. 102, and 1891, Cont. to Can. Pal., vol. 1, p. 235, Devonian.
 STROPHALOSIA keokuk, Beecher, 1890, Am. Jour. Sci. and Arts, Ser. 3, vol. 40, p. 244, Keokuk Gr.
 radicans, Winchell, 1866, (Crania radi- cans,) Rep. Low. Penin. Mich., p. 92, Ham. Gr.

- rockfordensis, Hull, 1892, Pal. N. Y., vol. 8, p. 353, Up. Devonian.
- scintilla, Beecher, 1890, Am. Jour. Sci. and Arts, Ser. 3, vol. 40, p. 243, Chouteau limestone.
- spondyliiformis, White & St. John, 1898, (Aulosteges spondyliiformis), Trans. Chl. Acad. Sci., p. 118, Coal Meas.
- STROPHODONTA, instead of Strophodonta. [Ety. *stropheus*, the socket in which the door moves; *odon*, tooth.]
- STROPHOMENA acanthoptera, Whiteaves, 1891, Can. Record Sci., p. 294, Up. Sil.
- conradi, S. winchelli, Hall, 1892, Pal. N. Y., vol. 8, p. 344, Trenton Gr.
- emaciata, scotfieldi, septata, Winchell & Schuchert, 1892, Am. Geo., vol. 9, pp. 285, 287, Trenton Gr., and planodorsata, Hud. Riv. Gr. Not properly defined.
- halli, iniquassa, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 334, Trenton Gr.
- SYRINGOTHYRIS randalli, Simpson, 1889, Trans. Am. Phil. Soc., p. 441, Chemung Gr.
- texta, Hall, 1857, (Spirifer textus), 10th Rep. N. Y. St. Mus. Nat. Hist., p. 169, Waverly Gr.
- TEREBRATULA gorbyi, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 77, Keokuk Gr.
- inconstans, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 24, Waverly Gr.
- inornata is from the Coal Meas., and described on p. 48.
- occidentalis, S. A. Miller, 1892, 18th Rep. Geo. Sur. Ind., p. 59, Chouteau limestone.
- traversensis, see Cryptonella traversensis.
- TREMATOSPIRA helena, Nettleroth, 1889, Kentucky Foss. Shells, p. 137, Niagara Gr.
- TRIPLESIA, or Triplasia as it should be spelled, can not be changed into Triplicia as suggested by Hall in 1892, Pal. N. Y., vol. 8, p. 209. In that event Diceraniscus would have priority. But Triplasia among the insects does not interfere with Triplasia among the fossil brachiopods.
- ZYGOSPIRA aquila, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 335, Trenton Gr.
- deflecta, Hall, 1847, (Atrypa deflecta), Pal. N. Y., vol. 1, p. 140, Trenton Gr.
- kentuckiensis, Nettleroth, 1889, Kentucky Foss. Shells, p. 137, Hud. Riv. Gr.
- uphami, Winchell & Schuchert, 1892, Am. Geo., vol. 9, p. 291, Galena Gr.

SUBKINGDOM MOLLUSCA.

CLASS PTEROPODA.

- COLEOLOIDES, Walcott, 1889, Proc. U. S. Nat. Mus., vol. 12, p. 37. [Ety. *Coleolus*, a genus; *oides*, like.] Shell slender, elongate, cylindrical, straight or slightly curved, apparently thin; surface marked by very fine, slightly oblique, longitudinal striae. Type *C. typicalis*, described at the same place from the Up. Taconic. A poor definition without illustration.
- CONULARIA, above the septum, is often broken off, and it has been suggested that it may have fallen off during the life of the animal; hence the shell is sometimes called deciduous, from *decido*, I fall off.
- gracilis, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 48, pl. 8, fig. 2; vol. 3, pl. 6, fig. 13, Waverly Gr. The name was preoccupied by Hall in 1847. As the species appears to be distinct from all others, I now propose the specific name *herricki*. The species will then be called *C. herricki*, after Prof. C. L. Herrick.
- intertexta, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 63, Keokuk Gr.
- salinensis, Whiteaves, 1891, Cont. to Can. Pal., vol. 1, p. 244, Devonian.
- sampsoni, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 80, Chouteau limestone.
- HYOLITHES lanceolatus, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 63, Chouteau limestone.
- similis, Walcott, 1889, Proc. U. S. Nat. Mus., vol. 12, p. 38, Up. Taconic. Not illustrated.
- terravivus, Walcott, 1889, Proc. U. S. Nat. Mus. vol. 12, p. 37, Up. Taconic. Not illustrated.

Rep. Geo. Sur. Ind.,

1888, Bull. Denison
Waverly Gr.

Coal Meas., and de-

ller, 1892, 18th Rep.
59, Chouteau lime-tonella traversensis.
Nettleroth, 1889,
Shells, p. 137, Ni-as it should be
changed into *Trip-*
oy Hall in 1892, Pal.
189. In that event
have priority. But
he insects does not
tasia among the fos-Sardeson, 1892, Bull.
Sci., vol. 3, p. 335,Atrypa defl-cta), Pal.
Trenton Gr.roth, 1889, Kentucky
Hud. Riv. Gr.& Schuchert, 1892,
291, Galena Gr.

CA.

propose the specific
the species will then
cki, after Prof. C. L.iller, 1892, Advance
Geo. Sur. Ind., p. 63,s, 1891, Cont. to Can.
Devonian.iller, 1891, Advance
Geo. Sur. Ind., p. 80,
e.S. A. Miller, 1892,
18th Rep. Geo. Sur.

au limestone.

9, Proc. U. S. Nat.
B, Up. Taconic. Not1889, Proc. U. S. Nat.
7, Up. Taconic. Not

CLASS GASTROPODA.

THE better authors use Gastropoda, and I prefer it to Gasteropoda. The masticatory apparatus or chitinous band, bearing minute teeth and forming a rasp-like ribbon, is called the odontophore, (*odous*, tooth; *phero*, I carry.) When the aperture is round or entire, the shell is called holostomatous, (*holos*, whole; *stoma*, mouth;) but when the aperture possesses a notch, more or less prolonged, for the respiratory siphon, or forming a trough-like extension for the skirt of the mantle, as in *Fusispira*, the shell is called siphonostomatous, (*siphon*, a tube; *stoma*, mouth.) The margin of the aperture is sometimes called the peristome, (*peri*, around; *stoma*, mouth), or peritreme, (*peri*, around; *trema*, an opening.) The notch separating the outer lip from the first whorl is called the posterior canal, and the notch separating it from the end of the columella is the anterior canal. When the columella is solid, it is called imperforate; and when hollow, perforated. The shells are sometimes composed of aragonite; at other times there is an inner layer of aragonite and an outer layer of calcite; and, especially in the younger shells, there is a horny epidermis which may disappear with age. It is said that aragonite is much more readily displaced in fossilization than calcite, which may account for the casts of *Murchisonia* associated with the shells of *Cyclonema*, in the same Lower Silurian strata, and other like phenomena in other genera and species of shells.

The new genus *Helenia* is supposed to belong to the Dentalidæ; *Fusispira* should be referred to the Subulitidæ; *Metoptoma* and *Palæacmæa* to the Patellidæ.

ACLISINA bellilineata, S. A. Miller, 1891,
17th Rep. Geo. Sur. Ind., p. 85, Chouteau limestone.

swallowana, on p. 395, is magnified three and a half diameters.

BELLEROPHON gorbyi, S. A. Miller, 1891, 17th Rep. Geo. Sur. Ind., p. 84, Hud. Riv. Gr.

incomptus, B. nodocostatus, B. ourayensis, B. rugocleurus, B. tenuilineatus,

Gurley, 1884, New. Carb. Foss. Bull. No. 2, pp. 8 to 11, Coal Meas.

subcordiformis, Herrick, 1887, Bull. Denison Univ., vol. 2, p. 18, Subcarboniferous.

CALLONEMA clarki, Nettleroth, 1889, Kentucky Foss. Shells, p. 175, Up. Held. Gr.

CARINAROPSIS deleta and *phalera*, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 335, Trenton Gr.

CONCHOPELTIS obtusa, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 336, Trenton Gr.

DENTALIUM granvillense, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 92, Waverly Gr.

ECCYLIOMPHALUS perkinsi, Whitfield, 1890, Bull. Am. Mus. Nat. Hist., vol. 3, p. 30, Calciferous Gr.

triangulus, Whitfield, 1890, Bull. Am. Mus. Nat. Hist., vol. 3, p. 29, Calciferous Gr.

ECCYLIOMPHALUS comes, Hall, syn. for *Eccyliomphalus laxus*.

flexistriatus and *E. maskusi*, Whiteaves, 1891, Cont. to Can. Pal., vol. 1, pp. 242, 243, Devonian.

manitobensis, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 100, Devonian.

sampsoni, Nettleroth, 1889, Kentucky Foss. Shells, p. 182, Up. Held. Gr.

FLEMINGIA, DeKoninck, 1881, Annales du Musee Royal D'Histoire Naturelle de Belgique, tome 6, p. 93. [Ety. proper name.]

Shell conical, spire acute, spiral whorls nearly flat externally; circumference more or less angular; aperture often depressed, and angular at the outer margin; peristome not continuous, external border oblique, slender, and sharp; columella slender, slightly twisted over, and giving place to the formation of an umbilical fossette more or less large and imperforate; shell thin; surface smooth or covered with irregular striae, obliquely crossed with more distant lines. Type *F. prisca*. This genus is nearly related to *Eotrochus*.

carbonaria, Meek & Worthen, 1866, (Trochita (?) carbonaria), Proc. Acad. Nat. Sci., p. 270, Kaaskaskia Gr. This species is figured and described by DeKoninck as

- a *Flemingia*, and yet it much resembles an *Eotrochus*.
- (?) *stultus*, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 45, Waverly Gr. This species may belong to *Eotrochus*.
- FUSIPIRA* (?) *spicula*, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 336, Trenton Gr.
- HELENIA*, Walcott, 1889, Proc. U. S. Nat. Mus., vol. 12, p. 30. [Ety. proper name.] Shell an elongate, narrow, flattened, curved tube; transverse section and aperture elliptical; surface marked by transverse, concentric, imbricating lines of growth. Type *H. bella*, which is described at the same place from the Up. Taconic, without illustration.
- HELICOTOMA similis*, Whitfield, 1890, Bull. Am. Mus. Nat. Hist., vol. 3, p. 31, Calciferous Gr.
- HOLOPEA cassina* is from the Calciferous Gr. *hubbardi*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 64, Hud. Riv. Gr.
- perundosa*, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 336, Trenton Gr.
- LEPETOPSIS parrishi*, Gurley, 1884, New Carb. Foss. p. 7, Up. Coal Meas.
- MACLUREA Manitobensis*, Whiteaves, 1880, Trans. Roy. Soc. Can., p. 75, Trenton Gr.
- Macrochilus carinatum*, see *Macrochilina carinata*.
- MACROCHILINA blairi*, S. A. Miller, 1891, 17th Rep. Geo. Sur. Ind., p. 84, Chouteau limestone.
- carinatum*, Nettleroth. The name was preoccupied. See *M. Nettlerothana*.
- nettlerothana*, n. sp. Proposed instead of *M. carinatum* of Nettleroth, in Kentucky Foss. Shells, p. 180, Up. Held. Gr., which name was preoccupied.
- MEGLOPTOMA*. Type *M. pileus*.
- explanata*, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 336, Trenton Gr.
- MURCHISONIA archiacana* and *M. dowlingi*, Whiteaves, 1892, Cont. to Can. Pal., p. 315, Devonian.
- hammelli*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 65, Hud. Riv. Gr.
- obelisca* is from the Calciferous Gr.
- NATICOPSIS remex* is from the Coal Meas.
- PALEACMEA cingulata*, Whiteaves, 1892, Cont. to Can. Pal., p. 311, Devonian.
- PLATYCERAS boonvillense*, S. A. Miller, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 82, Keokuk Gr.
- compressum*, Nettleroth, 1889, Kentucky Foss. Shells, p. 162, Up. Held. Gr.

FIG. 1255.—*Holopea hubbardi*.

- cyrtolites* is from the Burlington Group.
- milleri*, Nettleroth, 1889, Kentucky Foss. Shells, p. 165, Up. Held. Gr.
- missouriense*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 82, Burlington Gr.
- nasutum*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 82, Chouteau limestone.
- pettiscense*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 81, Burlington Gr.
- romingeri*, Walcott, 1888, Proc. U. S. Nat. Mus., p. 42, Up. Taconic. Not illustrated.
- PLATYSTOMA broadheadi*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 86, Chouteau limestone.
- PLEUROTOMARIA arabella*, Nettleroth. The name was preoccupied. See *P. nettlerothana*.
- clivosa*, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 337, Trenton Gr.
- difficilis*, Whitfield, 1890, Bull. Am. Mus. Nat. Hist., vol. 3, p. 33, Calciferous Gr.
- goniostoma*, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 98, Devonian.
- hurii*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 83, Up. Coal Meas.
- infranodosa*, Whiteaves, 1892, Cont. to Can. Pal., p. 313, Devonian.
- nettlerothana*, n. sp. Proposed instead of *P. arabella*, of Nettleroth, in Kentucky Foss. Shells, p. 171, which name was preoccupied, Up. Held. Gr.
- proctori*, Nettleroth, 1889, Kentucky Foss. Shells, p. 173, Up. Held. Gr.
- sedallensis*, S. A. Miller, 1891, 17th Rep. Geo. Sur. Ind., p. 83, Chouteau limestone.
- strigillata*, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 86, Waverly Gr.
- PORCELLIA Manitobensis*, Whiteaves, 1892, Cont. to Can. Pal., p. 318, Devonian.
- RAPHISTOMA tyrrelli*, Whiteaves, 1892, Cont. to Can. Pal., p. 314, Devonian.
- STRAPAROLLUS blairi*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 86, Chouteau limestone.
- SUBULITES* was described by Emmons, 1842, Geo. Sur. 2d Dist. N. Y., p. 392.
- benedicti*, S. A. Miller, 1891, 17th Rep. Geo. Sur. Ind., p. 84, Niagara Gr.
- TRYBLIDIUM conicum* is from the Chazy Gr.
- exsertum*, validum, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 337, Trenton Gr.
- indianense*, S. A. Miller, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 85, Hud. Riv. Gr.
- madisonense*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 64, Hud. Riv. Gr.

CLASS CEPHALOPODA.

THE living animals of this Class are carnivorous, but it is too strong a presumption to conclude that all fossil Cephalopoda were carnivorous. The evidence does not warrant the presumption. The mouth of the shell or aperture may be round, elliptical, lunate, T-shaped, or of almost any other form, and the shell may gradually expand to the aperture, be contracted behind the aperture, or more or less contracted to the aperture. The ventral side may be indicated by the shape of the aperture, either by an emargination or a sinus, but the position of the siphuncle does not always indicate it; for in *Orthoceras* the siphuncle may be central, or in the growth of the shell it may cross the central line. The external shell is generally destroyed and for this reason it has been supposed to have been composed of aragonite rather than calcite; but the fact that the shell often appears as if run together in fossilization does not indicate such a distinction.

ACTINOCERAS hindii, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 101, Devonian.

ASCO CERAS indianense, Newell, 1888, Proc. Bost. Soc. Nat. Hist., vol. 23, p. 484, Niagara Gr.

APSIDOCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 280. [Ety. *apsis*, the felloe of a wheel; *keras*, horn.] Loosely coiled, smooth, costated, or tuberculated gyroceran shells, with flattened abdomens. The whorls in section are triangular, the dorsum forming the internal apex of the outline; siphons near the venter and nummuloidal. The sutures have broad ventral lobes, saddles at the lateral angles, broad lobes on the sides and dorsal saddles; there is frequently a line of heavy tubercles on each of the lateral angles of the whorls; they are all large shells, and the abdomen is frequently hollow or fluted along the center. Type *A. magnificum*.

insigne, Whiteaves, 1889, Trans. Roy. Soc. Can., p. 82, Trenton Gr.

magnificum, Billings, 1857, (*Gyroceras magnificum*,) Rep. of Progr. Geo. Sur. Can., p. 307, Hud. Riv. Gr.

Asymptoceras newloni, see *Solenochilus newloni*.

CYRTO CERAS boycei is from the Chazy Gr. *dardanum* and *fosteri* are illustrated on pl. 16 in 20th Rep. N. Y. St. Mus. Nat. Hist.

hertzeri, refer to *Hexamoceras hertzeri*, howardi, Niagara Gr., and *C. thompsoni*, Hud. Riv. Gr., S. A. Miller, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 60.

indianense, S. A. Miller, 1891, Advance

Sheets 17th Rep. Geo. Sur. Ind., p. 88, Niagara Gr.

manitobense, Whiteaves, 1880, Trans. Roy. Soc. Can., p. 80, Trenton Gr.

nashvillense, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 87, Niagara Gr.

occidentale, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 103, Devonian.

saffordii, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 88, Hud. Riv. Gr.

Discites was preoccupied in 1708 by Walch, and again used by Schlotheim in 1820, before DeHaan used it in 1825 or McCoy in 1844. Hyatt in 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 292, proposed *Discitoceras* for the shells included in McCoy's genus. All the species therefore under the name *Discites* in this work should be referred to *Discitoceras*.

DOMATOCERAS, Hyatt, 1891, 2d Ann. Rep. Geo. Sur. Texas, p. 342. The generic characters can not be determined from the definition. *D. umbilicatum* is defined at the same place from the Coal Meas. as the type.

ENDOCERAS crassisiphonatum, Whiteaves, 1891, Trans. Roy. Soc. Can., vol. 9, p. 79, Trenton Gr.

ENDOLOBUS gibbosus, Hyatt, 1891, 2d Ann. Rep. Geo. Sur. Texas, p. 353, Coal Meas.

EPHIPIOCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 290. [Ety. *ephippion*, a saddle; *keras*, horn.] Generic definition very obscure. Type *E. ferratum*, described by Cox as *Nautilus ferratus*; and to the same genus *Nautilus divinus* should be referred if the

- description is such as to establish it, for the species do not properly belong to *Nautilus*.
- Gastrioceras compressum*, see *Goniates compressus*.
- GOMPHOCERAS angustum*, Newell, 1888, Proc. Bost. Soc. Nat. Hist., vol. 23, p. 475, Niagara Gr.
- clarki*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 88, St. Louis Gr.
- hertzeri*, see *Hexamoceras hertzeri*.
- lineare*, Newell, 1888, Proc. Bost. Soc. Nat. Hist., vol. 23, p. 473, Niagara Gr.
- manitobense*, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 102, Devonian.
- parvulum*, Whiteaves, 1891, Can. Rec. of Sci., p. 298, Up. Sil.
- projectum*, Newell, 1888, Proc. Bost. Soc. Nat. Hist., vol. 23, p. 476, Niagara Gr.
- wabashense*, Newell, 1888, Proc. Bost. Soc. Nat. Hist., vol. 23, p. 470, Niagara Gr.
- GONIATITES baylorensis*, White, 1891, Bull. U. S. Geo. Sur. No. 77, p. 19, Permian Gr.
- brownensis*, 1891, S. A. Miller, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 90, Waverly Gr.
- compressus*, Hyatt, 1891, (Gastrioceras compressum,) 2d An. Rep. Geo. Sur. Texas, p. 355, Coal Meas.
- gorbyi*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 90, Chouteau limestone.
- greenii*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 76, Knobstone Gr.
- indianensis*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 90, Waverly Gr.
- leviculus*, Miller & Faber, 1892, Jour. Cin. Soc. Nat. Hist., vol. 14, p. 167, St. Louis Gr.
- limatus*, Miller & Faber, 1892, Jour. Cin. Soc. Nat. Hist., vol. 14, p. 166, St. Louis Gr.
- missouriensis*, Miller & Faber, 1892, Jour. Cin. Soc. Nat. Hist., vol. 14, p. 164, Up. Coal Meas.
- occidentalis*, Miller & Faber, 1892, Jour. Cin. Soc. Nat. Hist., vol. 14, p. 166, Coal Meas.
- sciotoensis*, Miller & Faber, 1892, Jour. Cin. Soc. Nat. Hist., vol. 14, p. 165, Waverly Gr.
- GONIOCERAS lambi*, Whiteaves, 1891, Trans. Roy. Soc. Can., vol. 9, p. 86, Trenton Gr.
- GYROCERAS canadense*, G. *glicinctum*, G. *submammillatum*, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, pp. 106, 107, Devonian.
- magnificum*, refer to *Apsidoceras magnificum*.
- HEXAMOCERAS*, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 278. [Ety. *hex*, six; *meros*, part; *keras*, horn.] This genus is distinguished from *Gomphoceras* only by the aperture. In *Gomphoceras* there is only one lateral branch on each side of a longer median aperture, which gives the mouth a T-shape; in *Hexamoceras* there are three branches on each side of the median aperture, increasing in length toward the summit. Type *H. panderi*.
- cacabiforme*, Newell, 1888, Proc. Bost. Soc. Nat. Hist., vol. 23, p. 481, Niagara Gr.
- delphicum*, Newell, 1888, Proc. Bost. Soc. Nat. Hist., vol. 23, p. 479, Niagara Gr.
- hertzeri*, Hall & Whitfield, 1875, (Cyrtoceras hertzeri,) Ohio Pal., vol. 2, p. 150, Niagara Gr.
- HOMALOCERAS*, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 104. [Ety. *homalos*, level; *keras*, horn.] Shell consisting of a slender tube, which is broadly and strongly arcuate, curved in the same plane, and much flattened laterally, its venter or outer border being very narrow, truncated, and depressed in the center; sutural line consisting of two very narrow saddles, with an equally narrow sinus between them on the venter, a broadly concave sinus or lobe on each of the sides, and a rather narrow saddle on the dorsum; siphuncle cylindrical, exogastric, and placed near the venter or outer and convex margin; body-chamber long, occupying about one-third of the entire length. Type *H. planatum*, which is described at the same place from the Devonian.
- LITUITES eatoni*, L. *eatonii* var. *cassinensis*, L. *internistriatus*, L. *seelyi*, are from the Calceferous.
- magnificus*, refer to *Apsidoceras magnificum*.
- MEDLICOTTIA*, Waagen, 1879, Pal. Indica, Ser. 13, pp. 39, 83. [Ety. proper name.] Shell discoid and volutions deeply embracing; lobes divided by single linguiform marginal saddles, or trifoliate or divided saddles; ventral lobes deep and undivided; the first pair of saddles are narrow, long, and the margins cut by several lobes and saddles growing progressively longer internally; numerous auxiliary lobes are generated from the marginal divisions in the outlines of the first pair of saddles, and from the division of large magnosellarian saddles near the umbilicus. Type *M. primis*.
- copei*, White, 1889, Am. Nat., vol. 23, p. 117, and Bull. U. S. Geo. Sur., No. 77, Permian Gr.
- METACOCERAS*, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 268. Nautiloid shells, with a wide umbilicus perforated in the middle; volutions subquadangular, bearing nodes on the dorsolateral sides, slightly embracing; siphuncle eccentric on the dorsal side. Type *M. sangamonense*.
- cavatiforme*, M. *dubium*, M. *hayi*, M. *inconspicuum*, M. *walcotti*, Hyatt, 1891,

branch on each side aperture, which gives e; in *Hexamoceras* anches on each side aperture, increasing in summit. Type II.

1888, Proc. Bost. Soc. p. 481, Niagara Gr. 1888, Proc. Bost. Soc. vol. 23, p. 479, Ni-

tfeld, 1875, (Cyrtoc- o Pal., vol. 2, p. 150,

es, 1890, Trans. Roy. 104. [Ety. *homalos*, Shell consisting of which is broadly and curved in the same attenuated laterally, its order being very narrow and depressed in the me consisting of two les, with an equally seen them on the ven- cave sinus or lobe on and a rather narrow um; siphuncle cylin- and placed near the and convex margin; ng, occupying about entire length. Type ch is described at the the Devonian.

toni var. *cassinensis*, L. seelyi, are from the

Apsidoceras mag-

, 1879, Pal. Indica, [Ety. proper name.] volutions deeply em- ided by single lingu- iddles, or trifoliate or ventral lobes deep the first pair of saddles and the margins cut and saddles growing er internally; num- ers are generated from sions in the outlines of saddles, and from large magnosellarian umbilicus. Type M.

Am. Nat., vol. 23, p. S. Geo. Sur., No. 77,

1883, Proc. Bost. Soc. 2, p. 268. Nautiloid de umbilicus perfo- e; volutions subquad- nodes on the dorso- ightly embracing; si- on the dorsal side. nense.

um, M. hayi, M. in- walcotti, Hyatt, 1891,

2d Ann. Rep. Geo. Sur. Texas, pp. 334 to 340, Coal Meas planorbiforme, Meek & Worthen, 1860, Proc. Acad. Nat. Sci. Phil., p. 469, and Geo. Sur. Ill., vol. 2, p. 386, (Nautilus planorbiformis,) Upper Coal Meas.

FIG. 1236.—*Metacoceras cavatiforme*.

sangamonense, Meek & Worthen, 1860, (Nautilus sangamonensis,) Proc. Acad. Nat. Sci. Phil., p. 470, and Geo. Sur. Ill., vol. 2, p. 386, Up. Coal Meas.

NAUTILUS bisulcatus, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 20, Wa-

verly Gr. *forbesanus*, refer to *Temnochilus forbes-*

anum. *kelloggi* is from the Calciferous. *parallelus* is Beecher's species.

planorbiformis, see *Metacoceras planorbi-*

forme. *quadrangularis*, McChesney, should be *quadrangulus*. It is now referred to the genus *Tainoceras*, and is therefore *T. quadrangulum*.

sangamonensis, see *Metacoceras sanga-*

monense. *toddi*, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 72, Up. Coal Meas.

ONCOCERAS *gibbosum*, Whiteaves, see O. *whiteavesi*.

magnum, Whiteaves, 1889, Trans. Roy. Soc. Can., p. 79, Trenton (?) Gr.

whiteavesi, n. sp. Trenton (?) Gr. Proposed instead of O. *gibbosum*, Whiteaves, in Trans. Roy. Soc. Can., 1889, p. 80, which name was preoccupied.

ORTHOCERAS *brainerdi* is from the Calciferous.

canadense, Whiteaves, 1891, (Sactoceras *canadense*,) Trans. Roy. Soc. Can., vol. 9, p. 85, Trenton Gr.

colletti, Up. Coal. Meas., O. *franklinense*, Niagara Gr., and O. *gorbyi*, Hud. Riv. Gr., S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., pp. 65 to 68.

harri, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 87, Up. Coal Meas.

obstructum, Newell, 1888, Proc. Bost. Soc. Nat. Hist., vol. 23, p. 467, Ni-

agara Gr. *selkirkense*, Whiteaves, 1891, Trans. Roy. Soc. Can., vol. 9, p. 82, Trenton Gr.

semiplanatum, Whiteaves, 1891, Trans. Roy. Soc. Can., vol. 9, p. 81, Tren-

ton Gr. *tyrrelli*, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 100, Devonian.

winnepegense, Whiteaves, 1891, Trans. Roy. Soc. Can., vol. 9, p. 82, Tren-

ton Gr. *Phacoceras*, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 292. [Ety. *phakos*,

a lentil; *keras*, horn.] The characters ascribed to it do not distinguish it from *Discitoceras*. Type *P. oxystomum*. In 1891 the same author described *P. dumbli* from the Coal Meas. in 2d Ann. Rep. Geo. Sur. Texas, p. 347.

PHRAGMOCERAS *missouriense*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 89, Chouteau lime-

stone. *POPANOCERAS*, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 337. [Ety. *popanus*,

a round, flat cake; *keras*, horn.] Volutions involute, compressed and costate, or marked by furrows; the lobes and saddles are numerous and club-shaped; the ventral lobes are divided by prominent narrow, siphonal saddles, carrying small funnel lobes. Three or more pairs of lobes are divided by marginal saddles, either single or double, the terminations of the lobes being either bifid or trifid. Type *P. kinganum*.

walcotti, White, 1889, Am. Nat., vol. 23, p. 117, and Bull. U. S. Geo. Sur., No. 77, p. 21, Permian Gr.

POTERICERAS, McCoy, 1854, British Pal. Foss. p. 321. [Ety. *poterion*, a cup; *keras*, horn.] Shell short, fusiform; section circular; mouth contracted; septa simple; siphon subcentral, moniliform; distinguished from *Gomphoceras* and allied genera by having an entire and simple aperture. Type *P. ellipticum*.

apertum and *P. nobile*, Whiteaves, 1889, Trans. Roy. Soc. Can., vol. 8, pp. 77, 78, Trenton (?) Gr.

gracile, Whiteaves, 1891, Trans. Roy. Soc. Can., vol. 9, p. 87, Trenton Gr.

missouriense, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 70, Chouteau limestone.

Ptychites cumminsi, see *Waagenoceras cum-*

minsi. *Sactoceras canadense*, see *Orthoceras cana-*

dense. *SOLENOCHILUS blairi*, S. A. Miller, 1892, Ad-

advance Sheets 18th Rep. Geo. Sur. Ind., p. 75, Chouteau limestone.

newloni, Hyatt, 1891, (Asymptoceras *newloni*,) 2d Ann. Rep. Geo. Sur. Texas, p. 346, Coal Meas.

rockfordense, S. A. Miller, 1891, Advance

- Sheets 17th Rep. Geo. Sur. Ind., p. 89, Waverly Gr.
- STREPTODISCUS, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 71. [Ety. *streptos*, twisted; *diskos*, quoit.] Proposed instead of Trematodiscus of Meek & Worthen, which was preoccupied. Type *S. stygialis*.
- indianensis, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 71, Keokuk Gr.
- ohioensis, Miller & Faber, (Trematodiscus ohioensis,) 1892, Jour. Cin. Soc. Nat. Hist., vol. 14, p. 168, Waverly Gr.
- TEMNOCHILUS conchiferum T. crassum, and T. depressum, Hyatt, 1891, 2d Ann. Rep. Geo. Sur. Texas, pp. 329 to 333, Coal Meas.
- forbesanum instead of Nautilus forbesanus, on p. 444.
- TETRAGONOCERAS, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 105. [Ety. *tetra*, four; *gonia*, angle; *keras*, horn.] Shell subspiral, recurved, coiled loosely on the same plane, making a lax volution toward the apex, but nearly straight anteriorly; transverse section quadrangular; siphuncle cylindrical, exogastric, marginal, and placed in the middle of the venter. Type *T. gracile*, which is described at the same place from the Devonian.
- Trematodiscus, see Streptodiscus.
- TROCHOCERAS maccharlesi, Whiteaves, 1889, Trans. Roy. Soc. Can., p. 81, Trenton (?) Gr.
- WAAGENOCERAS, Gemmellaro, 1888, Gior. Sci. Nat. Ed. Econ., vol. 19, p. 11. [Ety. proper name; *keras*, horn.] Shell discoid; volutions embracing and bearing transverse constrictions; septa numerous and complex. Type *W. stachei cumminsi*, White, 1889, Am. Nat., vol. 23, p. 117, and Bull. U. S. Geo. Sur., No. 77, p. 20 Permian Gr.
- TAINOCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 269. [Ety. *tainia*, a head-band; *keras*, horn.] Whorls discoidal; section quadrate; two lateral rows of tubercles, and two rows on the outer side in the later stages of growth; siphon above the center; sutures have ventral, lateral, and dorsal lobes, but no annular lobes. Type *T. quadrangulum*. It is very much like Metacoceras, if not a synonym.
- cavatum, Hyatt, 1891, 2d Ann. Rep. Geo. Sur. Texas, p. 341, Coal Meas.

LAMELLIBRANCHIATA.

THIS Class has gradually improved in the Geological ages, and is now in the maximum of its development. The lobes of the animal are right and left, and unite along the dorsal or hinge-line of the shell, and extend laterally in the form of mantles to the pallial line. The depth of the pallial sinus indicates the size of the siphonal muscles. The shell opens along the ventral side. The external hinge ligament consists of horny fibres that are found preserved, sometimes, almost as well as the shell, in Lower Silurian rocks. Some shells have an internal ligament or cartilage between the hinges of the two valves, located in pits or cartilage furrows on the faces of the hinge. The anterior adductor is in front of the mouth, and the posterior adductor behind the umbo, near the termination of the intestine. Sometimes scars are left by the pedal muscles used for projecting and retracting the foot. When there are no teeth or crenulations on the hinge, the shell is edentulous. The teeth beneath the umbo are the cardinal teeth, and those anterior or posterior to the umbo are lateral teeth. Some shells are composed of aragonite, others of calcite, and others have an outer layer of calcite and an inner one of aragonite, and generally there is an epidermis or outer horny coating. The outer calcite layer is secreted at the margin of the mantle or circumference of the shell and is prismatic, while the inner layers of aragonite are secreted in the form of laminæ by the sides of the mantle.

The families Unionidæ and Ostreidæ, I think, are not palæozoic. Anthracosia, Priscaonia, and Cardinia might be referred to the Cardiniidæ; Clinopistha and Solenomya to the Solenomyidæ.

ter. Type *T. gracile*, found at the same place as *n. p.*

Leptodiscus.
lesi, Whiteaves, 1889, Can., p. 81, Trenton Gr.

mellaro, 1888, Gior., vol. 19, p. 11. [Ety. *as*, horn.] Shell embracing and bearing plications; septa numerous. Type *W. stachei*, 1889, Am. Nat., vol. 23, U. S. Geo. Sur., No. 1, Gr.

1883. Proc. Bost. Soc., p. 269. [Ety. *tainia*, *eras*, horn.] Whorls quadrate; two lateral ribs, and two rows on the later stages of above the center; lateral, lateral, and dorsal lobes. Type *n.* It is very much if not a synonym. 191, 2d Ann. Rep. Geo. Sur., Coal Meas.

s, and is now in the right and left, and laterally in the form of indicates the size of the The external hinge sometimes, almost as an internal ligament s or cartilage furrows of the mouth, and the the intestine. Some- d retracting the foot. edentulous. The teeth posterior to the umbo of calcite, and others and generally there is e secreted at the mar- atic, while the inner sides of the mantle.

ozoic. Anthracosia, e; Clinopistha and

ALLORISMA *convexum* and *A. cooperi*, Herrick, 1888, Bull. Denison Univ., vol. 3, pp. 72, 74, and *A. consanguinatum*, *A. cnyahoga*, vol. 4, pp. 28, 29, Waverly Gr.

ANODONTOPSIS *affinis*, Whiteaves, 1892, Cont. to Can. Pal., p. 303, Devonian.

Arca ornata, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 83, Waverly Gr.

AVICULA *circulus*, refer to *Pernopecten circulus*.

ohioensis, Herrick, 1887, (Gervillia *ohioensis*), Bull. Denison Univ., vol. 2, p. 36, Coal Meas.

recta, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 115, Waverly Gr.

subspatulata, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 30, Waverly Gr.

AVICULOPECTEN *cooperi*, *A. granvillensis*, and *A. perelongatus*, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 50, Waverly Gr.

scalaris and *sorer*, Herrick, 1887, Bull. Denison Univ., vol. 2, pp. 26, 27, Coal Meas.

sculptilis, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 92, Coal Meas.

Carbonarca occidentalis, refer to *Edmondia occidentalis*.

CARDIOPSIS *teuicostata*, Whiteaves, 1892, Cont. to Can. Pal., p. 307, Devonian.

CLIDOPHORUS *consuetus*, Ulrich, 1892, 19th Rep. Geo. Sur. Minn., p. 223, Galena Gr.

CLINOPISTHA *striata*, Nettleroth, 1889, Kentucky Foss. Shells, p. 200, Up. Held. Gr.

FIG. 1257.—*Clonychia rhomboidea*. Lateral and anterior views of a cast of a right valve.

CLONYCHIA, Ulrich, 1892, Am. Geo., vol. 10, p. 97. [Ety. *kleio*, I close; *onyx*, a claw.] Distinguished from *Ambonychia* by the absence of radiating plications or striae, the absence of a byssal opening in the anterior end, the less central position of the muscular scars, and by the absence of distinct hinge-teeth. Type *Ambonychia lamellosa*, Hall, and including *A. erecta*, *A. attenuata*, *A. mytiloides*, *A. undata*, and *A. amygdalina*, and

also *C. rhomboidea*, described at the same place from the Trenton Gr.

CONOCARDIUM *alternistriatum*, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 42, Waverly Gr.

elrudi, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 95, Niagara Gr.

exiguum and *C. parvulum* S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 94, Ham. Gr.

indianense S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 94, Keokuk Gr.

CRENIPecten *foerstii*, Herrick, 1887, Bull. Denison Univ., vol. 2, p. 28, Coal Meas.

senilis and *C. subcardiformis*, Herrick, 1888, Bull. Denison Univ., vol. 3, pp. 53, 54, Waverly Gr.

CYPRICARDELLA *gorbyi*, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 92, Keokuk Gr.

producta, Whiteaves 1892, Cont. to Can. Pal., p. 309, Devonian.

CYPRICARDINIA *scitula*, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 38, Waverly Gr.

CYPRICARDITES *halli*, Nettleroth, 1889, Kentucky Foss. Shells, p. 206, Hud. Riv. Gr.

cingulata, *C. germanus*, *C. glabellus*, *C. nanus*, *C. obtusifomis*, *C. sardesoni*, *C. tenellus*, Ulrich, 1892, 19th Rep. Geo. Sur. Minn., pp. 231 to 239, Trenton Gr.

hindi and *C. sterlingensis* refer to *Whitella hindi* and *Whitella sterlingensis*.

luculentus, *minnestensis*, *triangularis*, *vicinus*, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 338, Trenton and Hud. Riv. Gr.

modestus, *C. oviformis*, *C. terminalis*, Ulrich, 1892, Am. Geo., vol. 10, pp. 98 to 100, Trenton Gr.

Cyrtodonta hindi, see *Whitella hindi*.

huronensis, see *Cypricardites huronensis*.

Dolabra sterlingensis, see *Whitella sterlingensis*.

EDMONDIA *occidentalis*, Swallow, 1860, (*Cardinia occidentalis*), Trans. St. Louis Acad. Sci., vol. 1, p. 655, Chouteau limestone.

sulcifera, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 30, Waverly Gr.

Entolium attenuatum, Herrick, see *Pernopecten attenuatus*.

Gervillia ohioensis, see *Avicula ohioensis*.

GLOSSITES *manitobensis*, Whiteaves, 1892, Cont. to Can. Pal., p. 310, Devonian.

GLYPTODESMA *cancellatum*, Nettleroth, 1889, Kentucky Foss. Shells, p. 227, Up. Held. Gr.

GONIODON, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 84. [Ety. *gonia*, an angle; *odon*, tooth.] Shell equivaive, very inequilateral, gibbous, not gaping; resembling *Paleoneilo*, but the hinge is continuous, slightly flexed be-

neath the beaks, without true teeth (?); but the hinge margin of both valves zigzagged by sharp incisions, into which corresponding projections of the opposite valve fit closely; the series of denticulations thus formed is continuous, but the size of the excisions diminishes before and behind the beaks; posterior adductor scar nearly terminal. Type *G. ohioensis*, which is described at the same place from the Waverly Gr.

GRAMMYSIA blairi, S. A. Miller, 1891, Advance Sheets 17th Rep. Geo. Sur. Ind., p. 93, Chouteau limestone.
famelica, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 35, Waverly Gr.
herricki, n. sp. Proposed instead of *G. ovata*, Herrick, in Bull. Denison Univ., vol. 4, p. 35, pl. 3, fig. 12, Up. Subcarboniferous.
ovata, Herrick, was preoccupied, see *G. herricki*.

compressed, with a sulcus extending toward the basal margin. Type *I. truncata*, which with *I. elongata* is described at the same place from the Hud. Riv. Gr.

ovalis, Ulrich, 1892, 19th Rep. Geo. Sur. Minn., p. 242, Hud. Riv. Gr.
LIOPTERIA halli, L. nasuta, L. newberryi, and L. ortonii, Herrick, 1888, Bull. Denison Univ., vol. 3, pp. 60, 61, and vol. 4, pp. 29, 114, Waverly Gr., except the last, which is from the Keokuk Gr.

LEPTODESMA scutella, Herrick, 1888, Bull. Denison Univ., vol. 3, p. 59, Waverly Gr.

LIMOPTERA, the type is *L. macroptera*.

LYRIOPECTEN nodocostatus, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 32, Waverly Gr.

MACRODON newarkensis, M. striatocostatus, Herrick, 1888, Bull. Denison Univ., vol. 4, pp. 36, 37, and M. triangularis, vol. 3, p. 74, Waverly Gr.
pygmaeus, Whiteaves, 1892, Cont. to Can. Pal., p. 299, Devonian.

MATHERIA rugosa, Ulrich, 1892, 19th Rep. Geo. Sur. Minn., p. 241, Trenton Gr.

MEGALODON, Sowerby, 1827, Genera of Recent and Foss. Shells. [Ety. *mega*, large; *odus*, tooth.] Shell oblong, smooth or keeled; ligament external; hinge teeth 1 x 2, thick; one posterior lateral tooth; anterior adductor impression deep, with a raised margin and a small pedal scar behind it; beaks subspiral. Type *M. cucullatus*.
subovatus, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 97, Devonian.

Modiola waverlyensis, see *Mytilopsis waverlyensis*.

MODIOLOPSIS alata, M. angustata, M. milleri, M. parva,

M. simulatrix, M. subparallela, Ulrich, 1890, Am. Geo., vol. 5, pp. 274 to 283, Hud. Riv. Gr.; and M. oviformis, from the Trenton Gr.; and M. oblonga, M. pulchella, and M. subtruncata, from the Utica Slate.

charlestownensis, Nettleroth, 1889, Kentucky Foss. Shells, p. 218, Up. Held. Gr.

concava, M. similis, Ulrich, 1892, 19th Rep. Geo. Sur. Minn., pp. 225, 227, Trenton Gr., and M. subelliptica, p. 226, Galena Gr.

dychei, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 77, Hud. Riv. Gr.

MODIOMORPHA attenuata, Whiteaves, 1890, Trans. Roy. Soc. Can., vol. 8, p. 96, Devonian, M. compressa, M. tumida, and

FIG. 1258.—*Ischyrodonta truncata*. a, Outline view of a left valve; b, internal view of same; c, d, and e, three views of a cast.

ISCHYRODONTA, Ulrich, 1890, Am. Geo., vol. 6, p. 173. [Ety. *ischyros*, strong; *odus*, tooth.] Short or elongated, thick bivalve shells, having small subterminal beaks, with the hinge straight or slightly arcuate and extended posteriorly; hinge-plate wide and strong, without posterior lateral teeth, but with a strong cardinal tooth in the left valve, and two nearly as strong in the right; just in front of them a pair of subcircular, large and deep anterior muscular impressions, and between these and the teeth another very small pair; posterior scar large, but faintly marked, ovate, situated in the postero-cardinal region; pallial-line simple; ligament probably internal; in casts the beaks are prominent,

a sulcus extend-
basal margin. Type
with *I. elongata* is
same place from the

19th Rep. Geo. Sur.
d. Riv. Gr.

usuta, L. newberryi,
Herrick, 1888, Bull.
vol. 3, pp. 60, 61, and
4, Waverly Gr., ex-
ch is from the Keo-

Herrick, 1888, Bull.
vol. 3, p. 59, Wa-

s. L. macroptera.

tatus, Herrick, 1888,
iv., vol. 4, p. 32, Wa-

is, M. striatocostatus,
Bull. Denison Univ.,

and M. triangularis,
p. 74, Waverly Gr.

is, Whiteaves, 1892,
to Can. Pal., p. 299,
nian.

a rugosa, Ulrich, 1892,
Rep. Geo. Sur. Minn.,
Trenton Gr.

on, Sowerby, 1827,
ra of Recent and Foss.

s. [Ety. *megas*, large;
tooth.] Shell ob-

smooth or keeled;
vent external; hinge

1x2, thick; one pos-
terolateral tooth; an-

adductor impression
with a raised margin

a small pedal scar
and it; beaks subspir-

M. cucullatus.

us, Whiteaves, 1890,
s. Roy. Soc. Can., vol.

97, Devonian.

waverlyensis, see Myti-

is alata, M. angus-

M. milleri, M. parva,

subparallela, Ulrich,

vol. 5, pp. 274 to 283,

and M. oviformis, from

and M. oblonga, M.

M. subtruncata, from

Nettleroth, 1889, Ken-

ells, p. 218, Up. Held.

is, Ulrich, 1892, 19th
Minn., pp. 225, 227,

M. subelliptica, p. 226,

Miller, 1892, Advance
Geo. Sur. Ind., p. 77,

ata, Whiteaves, 1890,
Can., vol. 8, p. 96, De-

M. parvula, 1892, Cont. to Can. Pal.,
p. 296, Devonian.

Monotis is not an American paleozoic
genus.

MYALINA trigonalis, Whiteaves, 1892, Cont.
to Can. Pal., p. 294, Devonian.

MYTILARCA inflata, Whiteaves, 1892, Cont.
to Can. Pal., p. 293, Devonian.

MYTILOPS waverlyensis, Herrick, 1888,
(Modiola waverlyensis,) Bull. Denison
Univ., vol. 3, p. 63, Waverly Gr.

NUCULA herzeri, Nettleroth, 1889, Ken-
tucky Foss. Shells, p. 221, Up. Held.
Gr.

manitobensis, Whiteaves,
1892, Cont. to Can. Pal.,
p. 301, Devonian.

NUCULANA similis, and N.
spatulata, Herrick, 1888,
Bull. Denison Univ., vol.

3, p. 79, Waverly Gr.

ORECARDIA, Herrick, 1888,
Bull. Denison Univ., vol.

4, p. 41. [Ety. *oraios*,
produced at a fit season;

kardia, heart.] Shell in-

equilateral, inequivalve,

ventricose, strongly curved, acute, elevated
beak, which inclines forward at the

apex; hinge-line extended, produced
posteriorly, furnished with a thickened

ridge or cartilage plate; the beaks are
separated from the hinge by a pseudo-

area which is elevated, and more or
less arched under the beak; surface

marked with radiating lines. Type O.
ornata, which, with O. cornuta, is de-

scribed at the same place from the Wa-
verly Gr.

ORTHODESMA minnesotense, O. saffordi,
Ulrich, 1892, 19th Rep. Geo. Sur. Minn.,
pp. 228, 229, Trenton Gr.

ORTHONOTA corrugata, Whiteaves, 1890,
Trans. Roy. Soc. Can., vol. 8, p. 98, De-
vonian.

Ostrea patricula, Winchell, in the opinion
of Whitfield, is from the Cretaceous of

New Jersey, which by some accident
was mixed with Waverly or Burling-

ton fossils.

PALAEONEILO consimilis, P. curta, P. ignota,
Herrick, 1888, Bull. Denison Univ.,
vol. 4, pp. 43, 44, Waverly Gr.

Panænka is a Polish or Bohemian word, sig-
nifying the same as the Latin *puella*.

a little girl. It is not formed accord-
ing to the rules of nomenclature, and

should be discarded.

grandis, Whiteaves, 1891, Can. Rec. Sci.,
p. 402, Corniferous Gr.

PARACYCLAS elongata, and P. ooterlonii,
Nettleroth, 1889, Kentucky Foss. Shells,
p. 210, 212, Up. Held. Gr.

PERNOPECTEN attenuatus, Herrick, 1887,
(Entolium attenuatum,) Bull. Denison
Univ., vol. 2, p. 24, Waverly Gr.

circulus, Shumard, 1885, (Avicula cir-
culus,) Geo. Rep. Mo., p. 206, Chouteau
limestone.

PLETHOCARDIA, Ulrich, 1892, 19th Rep. Geo.
Sur. Minn., p. 243. [Ety. *pletho*, to be

full; *kardia*, heart.] Shell thin, oblique,
tumid, with the margins closed; beaks

large, prominent, spirally enroled, and
curving forward; narrow, deep escutch-

eon posterior to the beaks; bifid cardin-

al tooth projects forward and down-

ward; single lateral tooth at the

posterior extremity of the hinge-line;

anterior muscular scar deep at the

antero-dorsal angle, margined by a

curved ridge extending from the under

side of the cardinal tooth. Type P.

FIG. 1250.—Plethocardia umbonata.

umbonata, from the Trenton Gr., de-
scribed at the same place. P. sub-
erecta is also described from the Ga-
lena Gr.

POSIDONOMYA fragilis, Herrick, 1888, Bull.
Denison Univ., vol. 3, p. 39, Wa-
verly Gr.

PROMACRUS missouriensis, Swallow, 1860,
(Solen (?) missouriensis,) Trans. St. Louis
Acad. Sci., vol. 1, p. 655, Chouteau

limestone. It is not a Sanguinolites.

nasutus, Meek, 1871, (Sanguinolites
nasutus,) Am. Jour. Conch., vol. 7,
Chouteau limestone.

truncatus, Herrick, 1888, Bull. Denison
Univ., vol. 3, p. 60, Waverly Gr.

PTERINEA lobata, Whiteaves, 1892, Cont. to
Can. Pal., p. 292, Devonian.

PTERINOPECTEN ashlandensis, and P. cari-
niferus, Herrick, 1888, Bull. Denison
Univ. vol. 3, p. 58, and vol. 4, p. 33,

Waverly Gr.

sedaliensis, S. A. Miller, 1891, Advance
Sheets 17th Rep. Geo. Sur. Ind., p. 93,
Chouteau limestone.

PTERONITES obliquus, Herrick, 1888, Bull.
Denison Univ., vol. 3, p. 58, Wa-
verly Gr.

PTYCHOPTERIA aequalvalvis, Whiteaves, 1891,
Cont. to Can. Pal., vol. 1, p. 239, De-
vonian.

mesocostalis, Williams, 1887, Bull. U. S.
Geo. Sur., No. 41, p. 35, Portage Gr.

oboleta, Simpson, 1889, Trans. Am.
Phil. Soc., p. 448, Chemung Gr.

SANGUINOLITES missouriensis and S. nasutus
refer to Promacrus.

senilis, Herrick, 1888, Bull. Denison
Univ., vol. 3, p. 66, Waverly Gr.

SCHIZODUS affinis, Herrick, 1887, Bull. Deni-
son Univ., vol. 2, p. 41, Coal Meas.

harii, S. A. Miller, 1891, Advance Sheets

17th Rep. Geo. Sur. Ind., p. 91, Up. Coal Meas.

harlanensis, Herrick, 1888, Bull. Denison

FIG. 1260.—*Technophorus divaricatus*. Left valve natural size and magnified 3 diameters.

and *S. subcircularis*, 1887, vol. 2, pp. 36, 41, 42, Coal Meas.

wheeleri, Swallow, 1862, (*Cypricardia* (?) *wheeleri*), is in Trans. St. Louis Acad. Sci., vol. 2, p. 96.

SOLENOMYA cuyahogensis, Herrick, 1888, Bull. Denison Univ., vol. 4, p. 115, Waverly Gr., and *S. meekana*, and *S. subradiata*, 1887, vol. 2, p. 30, Coal Meas.

SPATHELLA subelliptica, Whitelaw, 1892, Cont. to Can. Pal., p. 298, Devonian.

STREBLOPTERIA gracilis, *S. media*, *S. squamosa*, Herrick, 1888, Bull. Denison Univ., vol. 3, pp. 56, 57, Waverly Gr.

TECHNOPHORUS extenuatus, Ulrich, 1892, 19th Rep. Geo. Sur. Minn., p. 222, Trenton Gr., and *T. divaricatus*, *T. flistriatus*, *T. subacutus*, Am. Geo., vol. 10, pp. 101 and 102, Trenton Gr.

TELLINOMYA diminuens, and *T. cuneata*, Simpson, 1889, Trans. Am. Phil. Soc., p. 453, Clinton Gr.

compressa *T. nitida*, *T. planodorsata*, *T. subrotunda*, Ulrich, 1892, 19th Rep. Geo. Sur. Minn., pp. 215 to 219, Trenton Gr.; *T. intermedia*, p. 218, Galena Gr.; and *T. recurva* and *T. similis*, pp. 220, 221, Hud. Riv. Gr.; and *T. longa*, Am. Geo., vol. 10, p. 103, Black Riv. Gr.

candens, *lepida*, Sardeson, 1892, Bull. Minn. Acad. Nat. Sci., vol. 3, p. 339, Trenton Gr.

WHITELLA, Ulrich, 1891, Am. Geo., vol. 6, p. 176. [Ety. proper name.] Shell large, thin, obliquely quadrangular or suboval, equivalve, inequilateral, more or less ventricose, closed all around; beaks prominent, incurved; cardinal line straight or slightly convex, the edges inflected to form a sharply defined es-

cutcheon extending beyond the beaks nearly to the anterior extremity of the shell; area finely striated longitudinally; hinge-line straight, one-half or two-thirds the length of the shell, with four or five oblique teeth in front of the beaks; ligament probably external and internal; two simple adductor impressions, posterior one faint, pallial-line entire; surface concentrically lined. Type *W. obliquata*, described at the same place from the Hud. Riv. Gr.; and *W. compressa* and *W. scotlandi*, from the Trenton Gr.

concentrica, Ulrich, 1892, 19th Rep. Geo. Sur. Minn., p. 247, Trenton Gr.; *W. praecepta*, p. 246, Galena Gr.; and *W. sulcodorsata*, p. 248, Hud. Riv. Gr. *hindi*, Billings, 1862, (*Cyrtodonta hindi*), Pal. Foss., vol. 1, p. 151, Hud. Riv. Gr. *sterlingensis*, Meek & Worthen, 1866, (*Dolabra sterlingensis*), Proc. Acad. Nat. Sci., p. 260, and Geo. Sur. Ill., vol. 2, p. 339, Hud. Riv. Gr.

FIG. 1261.—*Whitella obliquata*. *a*, Left side of a cast; *b*, and *c* anterior and cardinal views of a cast; *d*, left side of another cast; *e*, hinge and part of the muscular impressions.

g beyond the beaks
rior extremity of the
striated longitudin-
straight, one-half or
gth of the shell, with
e teeth in front of the
robably external and
ple adductor impres-
ne faint, pallial-line
concentrically lined.
ta, described at the
the Hud. Riv. Gr.;
sa and W. sciofieldi,
Gr.
1892, 19th Rep. Geo.
17, Trenton Gr.; W.
Galena Gr.; and W.
8, Hud. Riv. Gr.
(Cyrtodonta hindi,
p. 151, Hud. Riv. Gr.
& Worthen, 1866,
ensis.) Proc. Acad.
, and Geo. Sur. Ill.,
Riv. Gr.

Figure 2. a, Left side of a
and cardinal views of
another cast; c, hinge
lar impressions.

SUBKINGDOM ARTICULATA.

CLASS ANNELIDA.

THE Tubicola are invested in tubes to which they are not muscularly attached, and the paleozoic forms show no muscular scars. The tubes seem to have been composed of calcite, and are generally found attached to some other object at the apex or on one side. The jaws of the Conodonts are minute, glossy black, and chitinous or horny, instead of being composed of calcite. Most of them, probably, belong to the masticatory apparatus of Crustaceans.

- ARABELLITES aciculatus, and A. hindei, James, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 148, Hud. Riv. Gr.
CORNULITES sublevis, Whiteaves, 1891, Cont. to Can. Pal., vol. 1, p. 210, Devonian.
POLYGNATHUS wilsoni, James, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 148, Hud. Riv. Gr.
PRIONIODUS dychei, James, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 147, Hud. Riv. Gr.
SABELLARITES, Dawson, 1890, Quar. Jour.

Geo. Soc., vol. 46, p. 605. [Ety. *Sabellar*, a genus; *ites*, from *lithos*, stone.] Elongated tubes composed of grains of sand and calcareous organic fragments associated with carbonaceous, flocculent matter, indicating a horny or membranous sheath. Type *S. trentonensis*, which is described at the same place from the Trenton Gr., with *S. phosphaticus* from the Up. Taconic. *SPIRORBIS* is generally more or less elevated in the last whorl, and the aperture is always turned up.

CLASS CRUSTACEA.

THE integument of the Crustacea (*crusta*, a crust,) is called chitinous; the segments or somites (*soma*, body,) are arranged longitudinally, and united transversely by a membrane. The Cirripedia (*cirrus*, a curl; *pes*, foot,) are attached to submarine objects by the anterior end or metamorphosed head. The articulated cirri are exerted and retracted from an opening at the posterior extremity. The Eutomostrea (*entomos*, cut into; *ostrakon*, a shell,) have been defined as follows: Animals aquatic, covered with a shell or carapace of a horny consistency, formed of one or more pieces, in some genera resembling a cuirass or buckler, and in others a bivalve shell, which completely or in great part envelops the body and limbs of the animal. In other genera the animal is inverted with a multivalve carapace, like jointed-plate armor; the branchiae are attached either to the feet or to the organs of mastication; the limbs are jointed and more or less setiferous. The animals, for the most part, undergo a regular moulting or change of shell as they grow; in some cases this amounts to a species of transformation. The Ostracoda (*ostrakon*, a shell,) have the valves united on the back by a membrane or ligament, and the valves are closed by an adductor muscle, the place of attachment being indicated by a pit, group of spots, or tubercle. Many genera have been recently described, some of them, appar-

ently, not ranking higher than species. The central part of the cephalic shield of the Trilobita (*treis*, three; *lobos*, a lobe,) is called the glabella; the grooves at the sides of it are called axal furrows, the one at the rear the neck furrow; the fixed cheeks are on each side of the glabella, and separated from the free cheeks by the facial suture, though the facial suture is absent in Trinucleus and some other genera; the central part of the thorax is the axis, and the side lobes are the pleurae; the dividing line is the axal furrow; the segments of the pygidium are anchylosed. The Xiphosura (*xiphos*, a sword; *oura*, a tail,) have a broad, convex buckler, compound, subcentral eyes, and ocelli in front; the mouth has a small labrum and six pairs of appendages; the telson, or terminal segment, is ensiform.

- ACANTHOTELSON magister, Packard, 1886, Mem. Nat. Acad. Sci., vol. 3, p. 127, Carboniferous.
- ACIDASPIDES ortonii, Foerste, 1887, Bull. Denison Univ., vol. 2, p. 90, Niagara Gr. perarmata, Whiteaves, 1891, Can. Rec. Sci., p. 300, Up. Sil.
- ÆCHIMINA, Jones & Holl, 1869, Ann. and Mag. Nat. Hist., Ser. 4, vol. 3, p. 217. [Ety. *aichme*, a sharp point.] Valves thick, straight at the hinge, rounded at the ends, convex at the ventral border, and outdrawn at the surface into a broad-based and sharp-pointed hollow cone, which either involves all the surface or rises from the postero-dorsal or centro-dorsal region. Type *Æ. cuspidata*.
- abnormis, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 183, Niagara Gr., and *Æ. marginata*, p. 184, from the Ham. Gr.
- AGNOSTUS desideratus, Walcott, 1889, Proc. U. S. Nat. Mus., vol. 12, p. 39, Up. Taconic.
- latus*, see *Bollia lata*.
- AGRAULOS redpathi, Walcott, 1890, 10th Ann. Rep. U. S. Geo. Sur., p. 654, Up. Taconic.
- AMPYX americanus, Safford & Vogdes, Proc. Acad. Nat. Sci., Trenton Gr.
- ANOMALOCARIS, Whiteaves, 1892, Can. Rec. Sci., vol. 5, p. 205. [Ety. *anomalos*, unlike; *karis*, shrimp.] A phyllocarid crustacean; body from 9 to 13 segments, exclusive of the caudal segment, each bearing a pair of slender, narrowly elongated and acutely pointed, simple and probably branchial appendages of the nature of uropods or foot-gills; posterior terminal segment margined with three pairs of caudal spines, one terminal, the other two lateral. Type *A. canadensis*, described at the same place from rocks of uncertain age, probably Up. Taconic.
- APARCHITES concinnus, Jones, 1858, (Cytheropsis concinna,) Ann. & Mag. Nat. Hist., Ser. 3, vol. 1, p. 249, Black Riv. Gr.
- inornatus, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 182, Up. Held. Gr., and *A. oblongus*, p. 137, Hud. Riv. Gr. mitis, Jones, 1891, Cont. to Can. Micro-Pal., p. 91, Devonian; and *A. mundulus* and *A. tyrrelli*, p. 62, from the Chazy Gr.
- ARISTOZOE canadensis, Whitfield, 1890, Ann. N. Y. Acad. Sci., p. 505, Trenton Gr.
- ASAPHUS has a lip-plate, labrum, or hypostoma, (*hypo*, under; *stoma*, mouth;) that is wide and deeply forked behind. The jointed limbs and branchial filaments shown in the illustration of the Oxford specimen in the Jour. of the Cin. Soc. of Nat. Hist., as well as the description of them, are largely imaginary, as the specimen does not show the characters.
- ATOPS reticulata, Walcott, 1890, (Conocoryphe reticulata,) 10th Ann. Rep. U. S. Geo. Sur., p. 649, Up. Taconic.
- AVALONIA, Walcott, 1890, 10th Ann. Rep. U. S. Geo. Sur., p. 640. Type *A.manuelensis*, described at the same place from the Up. Taconic.
- BAIRDIA, McCoy, 1846, Synop. Foss. Ireland, p. 164. [Ety. proper name.] Carapace varying from a broadly triangular to a narrow elongate subtriangular form, with extremities more or less acute; surface smooth and setiferous or finely punctate; no central tubercle; lucid spots well marked; margins thin and trenchant; when closed the edges of the right valve lie within those of the left; interior of the marginal borders, except on the dorsal edge, cased with a narrow lamelliform plate, as in Cypris, except that a slight fold or notch is frequently apparent at the angles of the hinge-line; the dorsal edge of the right valve is quite simple, and, in the closed carapace, underlies the dorsal edge of the left valve, which is larger and overlapping; ventral margin incurved. Type *B. curta*.
- anticostiensis, Jones, 1890, Quar. Jour. Geo. Soc., vol. 46, p. 548, Hud. Riv. Gr.
- cestriensis, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 210, Kaskaskia Gr.; and *B. leguminoides*, p. 197, Ham. Gr.
- BARYCHILINA, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 198. [Ety. *barys*, heavy, thick; *cheilos*, a lip.] Carapace small, subrhomboidal or ovate; valves thick, unequal, the right the larger, overlapping the left except in the pos-

cephalic shield of the grooves at the furrow; the fixed free cheeks by the some other genera; the pleuræ; the diaphragm ankylosed. The convex buckler, com- small labrum and six

Whitfield, 1890, Ann. 505, Trenton Gr. te, labrum, or hy- der; *stoma*, mouth; deeply forked behind. and branchial fila- e illustration of the n the Jour. of the list., as well as the , are largely imagin- n does not show the

Mcott, 1890, (Conoc- 10th Ann. Rep. 640, Up. Taconic. 90, 10th Ann. Rep. 640. Type A. man- at the same place

ic. Synop. Foss. Ire- ty. proper name.] from a broadly tri- v elongate subtrian- extremities more or smooth and setifer- atate; no central oots well marked; trenchant; when the right valve lie left; interior of the cept on the dorsal narrow lamelliform except that a slight quently apparent at nge-line; the dorsal ve is quite simple, carapace, underlies he left valve, which erlapping; ventral Type B. curta.

1890, Quar. Jour. 548, Hud. Riv. Gr. 90, Jour. Cin. Soc. p. 210, Kaskaskia aminoides, p. 197,

90, Jour. Cin. Soc. p. 198. [Ety. *barus*, s, a lip.] Carapace al or ovate; valves e right the larger, except in the pos-

terior half of the more or less convex dorsal side; the edges of the valves in this portion of the back are smooth, and resemble a pair of thick lips; edges of both valves thick and smooth all around, that of the right valve much the heavier; a sharply defined, narrow or rounded umbilical pit; surface striate. Type B. punctostriata, which, with B. punctostriata var. curta and B. pulchella, are described at the same place from the Up. Held. or Ham. Gr.

BATHYURELLUS, Whitfield, 1890, Bull. Am. Mus. Nat. Hist., vol. 3, p. 38, Up. Taconic or Calciferous.

BATHYURISCUS dawsoni, Walcott, 1888, Proc. U. S. Nat. Mus., p. 446, Up. Taconic.

Bathyrurus seelyi, refer to *Bolbocephalus seelyi*.

BEECHERELLA, Ulrich, 1891, Am. Geo., vol. 8, p. 198. [Ety. proper name.] Carapace small, elongate, boat-shaped to ovate, moderately convex, more or less inequivalve; dorsal margin varying from nearly straight to strongly convex; back sometimes flattened, with a sharply defined carina on one or both valves, giving them a triangular shape in cross sections; in other cases the dorsal slope is convex; antero-dorsal extremity acuminate, often drawn out into a long spine; spine strongest on the right valve, sometimes absent on the left; posterior extremity acuminate or rounded; ventral edges convex or straightened in the middle; hingement simple, dorsal edge of right valve thickened, and in the central part overlapping the left valve. Type B. carinata, described at the same place, from the Low. Held. Gr., with B. cristata, B. angulata, B. navicula, B. ovata, B. subtumida, and B. subtumida var. intermedia, from the same rocks.

BEYRICHTIA ciliata, refer to *Ctenobolbina ciliata*.

clavigera and B. clavigera var. clavifracta, Jones, 1891, Cont. to Micro-Pal., p. 65, Chazy Gr.; and B. quadrifida, p. 66, Trenton Gr.; and B. tuberculata var. strictispiralis, p. 77, Up. Sil. Also B. clarkii, 1890, Quar. Jour. Geo. Soc., vol. 46, p. 17, Low. Held. Gr.; and B. diffusa, p. 546, Anticosti Gr.; and B. halli, p. 15, Waterlime Gr.; and B. hamiltonensis, p. 19, Ham. Gr.; and B. kalmadini, p. 538, Ham. Gr., and B. subquadrata, p. 537, Corniferous Gr. *duryi*, refer to *Ctenobolbina duryi*.

lyoni, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 190, Up. Held. or

Ham. Gr.; also B. tricollina, Ham. Gr.; and B. radiata var. cestrionensis and B. simulatrix, pp. 204, 205, Kaskaskia Gr. *richardsoni*, refer to *Depranella richardsoni*.

symmetrica, refer to *Bollia symmetrica*. tuberculata, Kløden, 1834, (Battus tuberculatus,) Verst. d. Mark Brandenburg, p. 115, Up. Sil.

BOLBOCEPHALUS, Whitfield, 1890, Bull. Am. Mus. Nat. Hist., vol. 3, p. 36. [Ety. *bolbos*, bulb; *kephale*, head.] Head large, semicircular, including the movable cheeks; glabella proportionally large, bulbous or subspherical, expanded in front of the eyes, and marked by a single, very indistinct furrow near the posterior; fixed cheeks narrow; eyes large, semicircular, elevated; lateral limbs narrow from front posteriorly; frontal limb linear; facial

FIG. 1262.—*Beecherella carinata*. Right valve, interior and dorsal views of same. Magnified 20 diameters.

suture cutting the posterior margin within the genal angle behind, and passing closely around the glabellar lobe in front; movable cheeks triangular, marginal, spined; pygidium semicircular, lobed, and transversely furrowed. Type B. seelyi.

seelyi, Whitfield, 1886, (Bathyrurus seelyi,) Bull. Am. Mus. Nat. Hist., vol. 1, p. 339, Calciferous Gr.

truncatus, Whitfield, 1890, Bull. Am. Mus. Nat. Hist., vol. 3, p. 37, Calciferous Gr.

BOLLIA, Jones & Holl, 1886, Ann. and Mag. Nat. Hist., ser. 5, vol. 17, p. 360. [Ety. proper name.] Valves oblong, with rounded and nearly equal ends; straight on the back, more or less outcurved on the ventral edge; surface punctate and bearing a lobular elevation on each side

- of a median bay-like sulcus, constituting two irregular, obliquely transverse lobes, which converge downward and meet near the middle of the ventral region by a low, narrow, bent isthmus, sinuous in the adult, but more or less simply curved in the young state. The dorsal portions of this horseshoe lobe project outward; there is a strong semilunar ridge at each end of the valve, parallel with the marginal border, which has a slight outer rim. Type B. *uniflexa*.
- bilobata*, Jones, 1890, Quar. Jour. Geo. Soc., vol. 46, p. 540, Corniferous Gr.; and B. *hindei*, p. 540, Ham. Gr.; and B. *semilunata*, p. 548, Hud. Riv. Gr.
- granifera*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 205, St. Louis Gr.; and B. *obesa*, p. 189, Up. Held. or Ham. Gr.; B. *persulcata* and B. *pumila*, pp. 116, 117, Hud. Riv. Gr.
- lata*, Vanuxem, 1842, (*Agnostus latus*.) (Geo. Rep. N. Y., p. 80, and Pal. N. Y., vol. 2, p. 301, Clinton Gr.)
- symmetrica*, Hall, 1852, (*Beyrichia symmetrica*.) Pal. N. Y., vol. 2, p. 317, Niagara Gr.
- ungula*, Claypole, 1889, Am. Geo., vol. 4, p. 338, Marcellus limestone.
- BRONTEUS senescens*, Hall, 1892, 8th Ann. Rep. St. Geo. N. Y., Chemung Gr.
- BYRNOCYPRIS*, Brady, 1880, Rep. Ostracoda of the Challenger, p. 45. [Ety. *buthos*, the depth of the sea; *Cypris*, a genus.] Carapace smooth, more or less reniform; left valve much larger than the right, which it overlaps both on the dorsal and ventral margins.
- devonica*, B. *indianensis*, B. *punctulata*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 196, Up. Held. or Ham. Gr.
- hindstromi* and B. *cotusa*, Jones, 1890, Quar. Jour. Geo. Soc., vol. 46, p. 548, Anticosti Gr.
- CALYMENE* has a subquadrate hypostoma, with two short spines posteriorly; the surface, when well preserved, is granular or tubercular.
- multicosta* is from the Chazy Gr.
- vogdesi*, Foerste, 1887, Bull. Denison Univ., vol. 2, p. 95, Niagara Gr.
- CARCINOSOMA*, Claypole, 1890, Am. Geo., vol. 6, p. 400. [Ety. *karkinos*, a crab; *soma*, the body.] Proposed instead of *Eurysoma*, on p. 259. Body ovate, narrower in front, abruptly tapering behind into a cylindrical abdomen ending in a spiniform tail; head-shield entire, roundly triangular, bluntly pointed in front; thoracic segments 6 or more, ending on each side in a backwardly directed point; abdominal segments 4, subquadrate; beyond these is a sharply triangular spine; appendages consist of 5 pairs of organs, the first 4 of which taper rapidly to a point and are furnished with spinous processes; the swimming feet are thicker and longer than the others, and consist of three segments. Type C. *newlini*, described at the same time on p. 260, from the Waterlime Gr.
- CERATOCARIS* had numerous body-rings, posterior to the bivalve carapace, carrying lunellar appendages, behind which there was a pointed telson and two lateral spines.
- Conocoryphe* is a synonym for *Atops*.
- CTENORHOLMINA*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 108. [Ety. *ktenos*, comb; *bolbos*, bulb.] Carapace small, elongate, suboval, strongly convex; the posterior two-fifths more or less bulbous or subglobular, and separated from the remainder by a deep, narrow sulcus, extending in a gentle curve from the dorsal margin more than half the distance across the valves toward the postero-ventral border; the anterior three-fifths often with another oblique but less impressed sulcus; valves equal, dorsal margin straight, hinge simple, ventral edge thick, and the true contact margins generally with a row of small spines on each side; in a lateral view both are concealed by a frill or flattened border, usually mistaken for the true contact edges; surface generally granulous. Type C. *ciliata*.
- alata*, C. *bispinosa*, C. *ciliata* var. *curta*, C. *ciliata* var. *emaciata*, *tumida*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, pp. 108 to 111, Hud. Riv. Gr.; and C. *antespinosa*, C. *informis*, C. *minimu*, C. *papillosa*, pp. 186 to 188, Up. Held. or Ham. Gr.; and C. *punctata*, p. 186, from the Niagara Gr.
- ciliata*, Emmons, 1855, (*Beyrichia ciliata*.) Am. Geo., p. 219, Hud. Riv. Gr.
- duryi*, S. A. Miller, 1874, (*Beyrichia duryi*.) Cin. Quar. Jour. Sci., vol. 1, p. 232, Hud. Riv. Gr.
- CYCLUS*, DeKoninck, 1841, Mem. Acad. Sci. Bruxelles, vol. 14, p. 18. [Ety. *kuklos*, circle.] Carapace longer than wide, somewhat hemispherical, narrow, smooth border, indented behind shield; divided down its center by a dorsal ridge, from which radiate ribs or transverse wrinkles. Type C. *radialis*.
- americana*, Packard, 1886, Mem. Nat. Acad. Sci., vol. 3, p. 143, Coal. Meas.
- Cypridina*, Edwards-Milne, 1838, Lamarck's Anim. Sans Vert., vol. 5, p. 178. [Ety. from the genus *Cypris*.] They have two eyes situated toward the middle of their bivalve test, and a caudal appendage at the posterior border, armed with spines disposed as the teeth of a comb. Edwards mentioned no type when founding the genus upon the living *Ostracoda*, but it is very evident the genus is not known in Palaeozoic rocks.
- herzert*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 209, Keokuk Gr.

ing feet are thicker
e others, and con-
ents. Type C. new-
the same time on
sterlime Gr.

umerous body-rings,
valve carapace, car-
ppendages, behind
pointed telson and

m for Atops.

1890, Jour. Cin. Soc.
p. 108. [Ety. *klenos*.

] Carapace small,
strongly convex; the

more or less bulky-
and separated from

deep, narrow sulcus,
ntle curve from the

than half the dis-
valves toward the

der; the anterior
with another oblique

ed sulcus; valves
gin straight, hinge

valve thick, and the
ns generally with a

es on each side; in
are concealed by a

border, usually mis-
contact edges; surface

s. Type C. ciliata.

ciliata var. curta, C.
ta, tumida, Ulrich,

. Nat. Hist., vol. 13,
d. Riv. Gr.; and C.

formis, C. minima,
6 to 188, Up. Held.

C. punctata, p. 186,
r.

(Beyrichia ciliata),
ud. Riv. Gr.

r, 1874, (Beyrichia
r. Jour. Sci., vol. 1,

r.

41, Mem. Acad. Sci.
p. 18. [Ety. *kuklos*,

longer than wide,
spherical, narrow,

ented behind shield;
center by a dorsal

radiate ribs or trans-
pe C. radialis.

1886, Mem. Nat.
p. 143, Coal. Meas.

ne, 1838, Lamarck's
t., vol. 5, p. 178.

enus Cypris.] They
situated toward the

bivalve test, and a
at the posterior

spines disposed as
mb. Edwards men-

found the genus
Ostracoda, but it is

enus is not known in
Jour. Cin. Soc. Nat.

9, Keokuk Gr.

Cythere ohioensis, see *Cytheropsis ohio-*
ensis.

CYTHERELLA, Jones, 1849, Monog. Entom.
Cret., p. 28. [Ety. diminutive of *Cythere*.]

Carapace oblong, compressed, smooth or
pitted; no terminal denticulations;

contact margins of the right (larger)
valve grooved or lobed on its inner

edge for the reception of a flange pre-
sented by the contact margin of the

left (smaller) valve; both groove and
flange stronger at the posterior than at

the anterior portion of the valves. The
lucid spots resemble those in Cyprid-

ina.

ovatifomis, Ulrich, 1890, Jour. Cin.
Soc. Nat. Hist., vol. 13, p. 209, Kas-

kaskia Gr.

Cytherina phaseolus, see *Leperditia phaseolus*.

CYTHERELLINA, Jones & Holl, 1869, Ann. and
Mag. Nat. Hist., vol. 3, p. 215. [Ety.

from *Cytherella*.] Carapace valves
elongate, convex, smooth, thick, ex-

cavated internally, with undulating con-
tours. Type C. siliqua.

glandella, instead of *Cytheropsis glandella*.

CYTHEROPSIS *concinna*, see *Aparchites con-*
cinnus.

ohioensis, Herrick, 1888, (Cythere ohio-
ensis), Bull. Denison Univ., vol. 4, p.

60, Waverly Gr.

DALMANITES troosti, Safford, 1889, (Chasmo-
pops troosti), Proc. Acad. Nat. Sci., Tren-

ton Gr.

DEPRANELLA, Ulrich, 1890, Jour. Cin. Soc.
Nat. Hist., vol. 13, p. 117. Carapace

small, high, subelliptical in outline,
dorsal border straight, terminating ab-

solutely at each end; ventral border
nearly straight or gently convex, round-

ing almost evenly at the ends; a
sickle-shaped ridge runs nearly parallel

with the posterior and ventral edges,
and is sometimes produced beyond the

postero-dorsal border; dorsal slope
with two or more strong tubercles or

ridges; the two valves meet equally at
the ventral edge. Type D. crassinoda,

which is described at the same place,
from the Birdseye Gr., with D. ampla,

D. elongata, D. macer, D. nitida, pp. 119
to 121, Chazy Gr.

richardsoni, S. A. Miller, 1874, (Beyrichia
richardsoni.) Cin. Quar. Jour. Sci., vol.

1, p. 347, Hud. Riv. Gr.

DIONIDE is from N. Jahrb. fur Miner., Hft. 4,
p. 391, and Syst. Sil. Boh., p. 640.

ELLIPTOCEPHALA broggeri, Walcott, 1889,
(Olenellus broggeri.) Proc. U. S. Nat.

Mus., vol. 12, p. 41, Up. Taconic.

ELPE, Barrande, 1872, Syst. Sil. Boh., vol. 1,
Supp. p. 510. Type E. pinguis.

tyrelli, Jones, 1891, Cont. to Micro-Pal.,
p. 93, Devonian.

ENCRINURUS exceedensis, Safford, 1889, Proc.
Acad. Nat. Sci., Trenton Gr.

thresheri, Foerste, 1887, Bull. Denison
Univ., vol. 2, p. 101, Niagara Gr.

ENTOMIS, Jones, 1861, Mem. Geol. Sur. Gt.

Brit., Geol. Edinb., p. 137. Carapace
ovate-oblong, bean-like; valves indented

by a transverse furrow, which begins on
the dorsal margin, at about one-third of

its length from the anterior extremity,
and reaches half-way or more across

the valve; surface bearing in front of
the sulcus a tubercle or spine which is

sometimes wanting; anterior border not
indented. Type E. tuberosa.

madisonensis, Ulrich, 1890, Jour. Cin. Soc.
Nat. Hist., vol. 13, p. 107, Hud. Riv. Gr.;

and E. waldronensis, p. 183, Niagara
Gr.

rhomboidea, Jones, 1890, Quar. Jour. Geo.
Soc., vol. 46, p. 20, Ham. Gr.

EURVCHILINA, Ulrich, 1889, Micropaleontol-
ogy of Can., p. 52. [Ety. *eury*, from

cheilos, lip.] Valves somewhat semicir-
cular or semi-elliptical; dorsal line

straight; subcentral sulcus and a node
behind it; broad border, often striated;

hinge simple; surface reticulate, gran-
ulose, or smooth. Type E. reticulata,

which is described at the same place
from the Trenton Gr., and also E. man-

itobensis.

æqualis, E. granosa, E. longula, E. obesa,
E. subradiata, Ulrich, 1890, Jour. Cin.

Soc. Nat. Hist., vol. 13, p. 126 to 129,
Chazy and Birdseye Gr.

striatmarginata, S. A. Miller, (Beyrichia
striatmarginata,) 1874, Cin. Quar. Jour.

Sci., vol. 1, p. 233, Hud. Riv. Gr.

Eurysona, Clappole, 1890, Am. Geol., vol. 6,
p. 259. The name was preoccupied.

See Carcinosoma.

newlini, see Carcinosoma newlini.

GRIFFITHIDES, Portlock, 1843, Rep. Geol.
Londonderry, p. 310. [Ety. proper

name.] Distinguished from Phillipsia,
which it closely resembles by the pyr-

iform or tumid glabella and small,
smooth, lunate eyes. Type G. longiceps.

bufo, Meek & Worthen, 1870, Proc. Acad.
Nat. Sci., p. 52, and Geo. Sur. Ill., vol. 5,

p. 528, Keokuk Gr.

portlocki, Meek & Worthen, 1865, Proc.
Acad. Nat. Sci., p. 268, and Geo. Sur.

Ill., vol. 5, p. 525, Keokuk Gr.

sedaliensis, Vogdes, 1888, Trans. N. Y.
Acad. Sci., vol. 7, p. 276, Waverly Gr.

HALLIELLA, Ulrich, 1890, Jour. Cin. Soc. Nat.
Hist., vol. 13, p. 184. [Ety. proper name.]

Valves similar to Primitia, but with a
larger sulcus, narrow at the dorsal edge,

and widening as it extends downward;
posterior lobe smaller than the an-

terior; the latter generally divided at
or near the straight dorsal edge; sur-

face ornamented or smooth; ventral
edge thick. Type H. retifera, described

at the same place from the Up. Held.
or Ham. Gr.

HARPIDES. Type H. hospes.

ISOCHILINA amii, I. labelliosa, I. grandis var.
latimarginata, I. ottawa var. intermedia,

and I. whiteavesi, Jones, 1891, Cont. to
Can. Micropaleontology, pp. 68 to 78,

- Chazy and Trenton Gr.; and *I. bellula* and *I. dawsoni*, p. 92, Devonian.
- ambiana*, *I. ampla*, *I. kentuckiensis*, *I. safordi*, *I. subnodosa*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, pp. 177 to 180, Birdseye and Trenton Gr.; and *I. rectangularis*, p. 182, Up. Held. or Ham. Gr.
- fabacea* and *I. lineata*, Jones, 1890, Quar. Jour. Geo. Soc., vol. 46, p. 21, Ham. Gr.
- JONESELLA*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 121. [Ety. proper name.] Carapace small, ovate, moderately convex; valves equal, their outline and general aspect much as in *Primitia*, but differing in having a simple or more or less divided prominent ridge on the posterior two-thirds, more or less curved. Type *J. crepiformis*.
- crassa*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 123, Trenton Gr.; *J. digitata* and *J. pedigera*, p. 122, Hud. Riv. Gr.
- crepiformis*, Ulrich, 1870, Jour. Cin. Soc. Nat. Hist., vol. 2, p. 10, Hud. Riv. Gr.
- Karlia*, Walcott, 1888, Proc. U. S. Nat. Mus., p. 444. [Ety. from the Christian name Karl, which is a violation of all rules, and should be disregarded.] Form elongate-oval, convex; head longitudinally semicircular, deeply marked by the dorsal furrows; glabella clavate, broadly expanded in front, with or without faint glabellar furrows; occipital furrow well defined; fixed cheeks subtriangular; posterior furrow broad; eye lobe small; free cheeks narrow; hypostoma with a thick, rounded anterior margin that is extended into the large lateral wings, the sides of which extend one-half way back on the oval, convex body; posterior marginal rim strong, and separated from the body by a well-defined sulcus; thorax with seven segments; axis with a central spine on each segment; pleural lobes with a broad groove; anterior lateral ends of pleurae faceted; pygidium short, transverse, four to five segments in the axis, lateral lobes slightly grooved; surface granulose. Type *K. minor*, described at the same place from the Up. Taconic, and also *K. stephenensis*.
- ISOXYS*, Walcott, 1890, 10th Ann. Rep. U. S. Geo. Sur., p. 925. Carapace large; dorsal margin slightly curved; dorsal angles produced into sharp points; anterior and posterior ends alate, subequal in outline, and merging into the rounded ventral margin, without forming an angle; marginal rim narrow; valves equal; surface smooth. Type *I. chilhoweana*, Up. Taconic.
- KIRBYA*, Jones, 1850, Trans. Tyneside Nat. Field Club, vol. 4, p. 129. [Ety. proper name.] Carapace valves compressed, thick, oblong, impressed with a subcentral pit and raised into ridges, some concentric with the margin, associated sometimes with longitudinal wrinkles and by a reticulate ornament; valves suboblong, higher behind than before; extremities rounded, one more obliquely than the other; dorsal border straight, and its ends subacute; ventral border nearly straight in its middle third, and broadly curved at the ends; hinge simple; ventral edge of the dextral valve overlaps slightly that of the other. Subcentral pit variable. Type *K. permiana*.
- lindahli*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 207, St. Louis Gr., and *K. venosa*, p. 208, Kaskaskia Gr.; *K. parallela*, *K. semimuralis*, *K. subquadrata*, p. 192, Up. Held. or Ham. Gr. Mr. Ulrich also identifies *K. oblonga* and *K. tricolliata*, Jones & Kirby, pp. 206 and 207, Kaskaskia Gr., and *K. costata*, McCoy, p. 208, St. Louis Gr.
- walcotti*, Jones, 1890, (*Primitia*?) (*walcotti*), Quar. Jour. Geo. Soc., vol. 46, p. 543, Ham. Gr.
- KLOEDONTIA*, Jones & Holl, 1886, Ann. and Mag. Nat. Hist., ser. 5, vol. 17, p. 347. [Ety. proper name.] Carapace valves smooth, convex, impressed with two short, vertical furrows on the dorsal region, and a third smaller furrow defines a narrow semi-lune at the front end of the valve. Type *K. wilkensana*. *Beyrichia notata* and *B. notata* var. *ventricosa*, Hebert, 1859, Pal. N. Y., vol. 3, pp. 379, 380, Held. Gr., are referred by Jones to this genus.
- LEPERDITIA* was derived from a proper name. There is generally a tubercular eye-spot near the hinge, below and behind which there is a slight inflation and a vertical groove, extending from the dorsal margin part way across the valves.
- aequilatera*, *fimbriata*, *dorsicornis*, *granilirata*, *germana*, *millepunctata*, *inflata*, *mundula*, *sulcata*, and *tumida*, Ulrich, 1892, Am. Geol., vol. 10, pp. 264 to 269, Birdseye, Trenton, and Hud. Riv. Grs.
- appressa*, *L. cecigena* var. *frankfortensis*, *L. linneyi*, *L. tumidula*, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, pp. 174 to 177, Trenton Gr.; *L. nicklesi*, p. 200, Warsaw Gr.; *L. subrotunda*, p. 181, Up. Held. or Ham. Gr.
- balthica* var. *primæva*, *L. obscura*, Jones, 1891, Cont. to Micropal., p. 70, Trenton Gr.; *L. balthica* var. *guelphica*, *L. hisingeri* var. *egena*, var. *fabulina*, var. *gibbera*, *L. caeca*, *L. phaseolus* var. *guelphica*, *L. selwyni*, *L. whiteavesi*, pp. 80 to 89, Niagara and Guelph Gr.; *L. exigua*, p. 94, Devonian.
- canadensis* var. *nana*, syn. for *L. canadensis*.
- claypolei*, Jones, 1890, Quar. Jour. Geo. Soc., vol. 46, p. 25, Hud. Riv. Gr.
- crepiformis*, see *Jonesella crepiformis*.
- frontalis*, Jones, 1890, Quar. Jour. Geo. Soc., vol. 46, p. 547, Anticosti Gr.

margin, associated longitudinal wrinkles ornament; valves behind than before; one more obliquely subborder straight, late; ventral border middle third, and the ends; hinge of the dextral slightly that of the pit variable. Type

Jour. Cin. Soc. Nat. St. Louis Gr., and Kaskaskia Gr.; K. curialis, K. subquadri- feld. or Ham. Gr. entifies K. oblonga ones & Kirby, pp. askin Gr., and K. 08, St. Louis Gr. rmitia(?) walcotti), oc., vol. 46, p. 543,

oll, 1886, Ann. and r. 5, vol. 17, p. 347. Carapace valves pressed with two ews on the dorsal smaller furrow de- lune at the front type K. wilkensana. nd B. notata var. 0, Pal. N. Y., vol. 3, Held. Gr., are re- his genus.

ed from a proper herally a tubercular hinge, below and is a slight inflation ve, extending from art way across the

dorsicornis, grani- lepnata, inflata, and tumida, Ulrich, 10, pp. 264 to 269, and Hud. Riv. Grs. var. frankfortensis, lula, Ulrich, 1890, Hist., vol. 13, pp. Gr.; L. nicklesi, p. subrotunda, p. 181, Gr.

L. obscura, Jones, pal., p. 70, Trenton guelphica, L. his- var. fabulina, var. L. phaseolus var. ni, L. whitevesi, a and Guelph Gr.; onian.

n. for L. canadensis. Quar. Jour. Geo. Hud. Riv. Gr. la crepiformis. Quar. Jour. Geo. Anticosti Gr.

hisingeri, Schmidt, 1873, Mem. Acad. Imp. Sci. St. Petersburg, ser. 7, vol. 31, p. 16, Niagara Gr.

marginata, Schmidt, 1873, Mem. Acad. Imp. Sci. St. Petersburg, ser. 7, vol. 31, p. 19, Niagara Gr.

uana, syn. for L. canadensis.

parasitica, refer to Beyrichia parasitica.

phaseolus, Hisinger, 1831, Anteckn. Phys. Geogr., vol. 5, pp. 110 to 135, Niagara Gr.

punctulifera, see Primitiopsis punctulifera. subcylindrica, Ulrich, 1889, Micropal. of Can., p. 49, Hud. Riv. Gr.

LEPIDOCOLEUS scales or plates are supposed to form the capitulum, (caput, head; applied to a barnacle, from its being supported on a peduncle,) but it has been suggested that they are the scales of the peduncle, and the capitulum is unknown.

LACHAS bicornis and L. robbinsi, Ulrich, 1890, Ann. Geol., vol. 10, pp. 271 and 272, the first Hud. Riv. Gr., and second Galena Gr.

champlainensis is from the Chazy Gr.

faberi is a synonym for L. halli. At the time of the publication of this book I was not aware of the publication of Foerste's species.

halli, Foerste, 1888, Bull. Denison Univ., vol. 3, p. 118, Hud. Riv. Gr.

MACROCAUS, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 78.

[Ety. makros, long; karia, shrimp.] Carapace bivalved, united dorsally with a strong ligament; valves long, narrow, and ornamented with anastomosing striae; they are pointed on the dorsal side in front, and on the ventral side at the posterior end, while in the middle part the dorsal and ventral sides are subparallel; abdomen consisting of twelve or more segments, which

FIG. 1263.—*Macrocaris gorbyi*. Carapace valves, abdominal segments and postabdomen.

very slowly taper from the fourth or fifth to the postabdomen; postabdomen consisting of a short, expanding plate, with a central ridge or line of division.

The genus *Strigocaris* is known only from the carapace valves, and the posterior ends are subtruncated from the ventral side to the dorsal side, while in this genus the posterior ends of the valves are subtruncated from the dorsal to the ventral side. Type *M. gorbyi*, described at the same place from the Keokuk Gr.

MACROCYPRI, Brady, 1867, Intellectual Observer, vol. 12, p. 119. [Ety. makros, long; Cypria, a genus.] Carapace subcylindrical or long triangular, and often Bairdia-like, generally elongate, attenuated at the extremities; valves thin, smooth, unequal, with beveled plates within the ends, more or less sinuate on the ventral margin, the right larger than the left, and overlapping dorsal hinge-line flexuous.

subcylindric, Jones, 1890, Quar. Jour. Geo. Soc., vol. 46, p. 549, Anticosti, Gr.

FIG. 1264.—*Mesothyra gurleyi*. Postabdomen.

MESOTHYRA gurleyi, S. A. Miller, 1892, Advance Sheets 18th Rep. Geo. Sur. Ind., p. 77, Kinderhook Gr.

MICRODISCUS bellinmarginatus, Shaler & Foerste, 1888, Bull. Mus. Comp., p. 35, Up. Taconic.

helena, Walcott, 1889, Proc. U. S. Nat. Mus., vol. 12, p. 40, Up. Taconic.

MOOREA, Jones & Kirby, 1869, Ann. and Mag. Nat. Hist., ser. 4, vol. 3, p. 225. [Ety. proper name.] Carapace valves simple,

thick, flattened, longer on the dorsal than the ventral margin, without any subcentral pit, and ornamented with narrow, rounded ridges, following more or less closely and completely the marginal contour. Type *M. silurica*.

bicornuta, Ulrich, 1890, Jour. Cin. Soc. Nat. Hist., vol. 13, p. 191, Ham. Gr.;

and *M. granosa*, p. 206, Kaskaskia Gr. kirbyi, Jones, 1890, Quar. Jour. Geo.

Soc., vol. 46, p. 542, Corniferous Gr.

NORMOZOE was described in 1872, Syst. Sil. Boh., vol. 1, Supp., p. 536. Type *N. pollens*.

OCTONARIA, Jones, 1887, Ann. and Mag. Nat. Hist., ser. 5, vol. 19, p. 404. [Ety.

octonarius, a, um, consisting of eight; an adjective.] If any regard to the

rules of nomenclature is to be observed, such generic names must be disregarded. The characters ascribed to the genus are, probably, like other characters, ascribed by the same author to other genera, of no more than specific value. It would not be advisable, therefore, to propose a substantive name for the adjective. Ulrich defined four species, *curta*, *ovata*, *clavigera*, and *stigmata*, in *Jour. Cin. Soc. Nat. Hist.*, vol. 13, pp. 193 to 195; but it is only necessary to translate such names to show the absurdity. *Octonaria curta*, of eight short; *Octonaria ovata*, of eight ovate, etc. But the rules of nomenclature absolutely require a substantive for a generic name. If authors, who do not know how to distinguish an adjective from a noun, would consult some one who does before publishing their new generic names, we might be spared the trouble of indexing such work.

Ogygoris, Walcott, 1888, *Proc. U. S. Nat. Mus.*, p. 446. [Ety. from resemblance to *Ogygia*.] Distinguished from *Ogygia* by having a well-defined ocular ridge and a narrow palpebral lobe. Type *O. klotzi*.

klotzi, instead of *Ogygia klotzi*.

Olenellus broggeri, see *Elliptocephala broggeri*.

Olenoides curticei, Walcott, 1888, *Proc. U. S. Nat. Mus.*, p. 443, *Up. Taconic. desideratus* and *O. elli*, Walcott, 1890, 10th Ann. Rep. U. S. Geo. Sur., pp. 642, 644, *Up. Taconic.*

Pachydomella, Ulrich, 1890, *Jour. Cin. Soc. Nat. Hist.*, vol. 13, p. 197. [Ety. *pachys*, thick; *domus*, house; *ellus*, diminutive.] Carapace ventricose; valves thick and strong, the left much the largest, its thick edges overlapping the right valve on all sides; dorsal side strongly arched, ventral edge more nearly straight, ends subequal; a faintly impressed, subcentral umbilical pit. Type *P. tumida*, described at the same place from the *Up. Held* or *Ham. Gr.*

Paradoxides walcotti, Shaler & Foerste, 1888, *Bull. Mus. Comp. Zool.*, vol. 16, p. 36, *Up. Taconic.*

Phacops has the facial sutures uniting in front and abruptly curving from the eyes to the lateral margins. The hypostoma is convex and subtriangular. *pulchella*, Foerste, 1887, *Bull. Denison Univ.*, vol. 2, p. 99, *Niagara Gr.*

Piletonoides immaturus, *P. occidentalis*, *P. spinosus*, Herrick, 1888, *Bull. Denison Univ.*, vol. 4, pp. 57 to 59, *Waverly Gr.*

Phillipsia has an hypostoma longitudinally convex, winged antero-laterally, and terminating in an obtuse point behind. The type of the genus is *P. kelli* instead of *P. gemmulifera*.

bufo, refer to *Griffithides bufo*.

consors, Herrick, 1888, *Bull. Denison Univ.*, vol. 4, p. 53, *Keokuk Gr. nodocostata*, Hare, 1891, *Kansas City Sci.*, vol. 5, p. 33, *Up. Coal. Meas.*

portlocki, refer to *Griffithides portlocki*. *sampsoni*, Vogdes, 1888, *Trans. N. Y. Acad. Sci.*, vol. 7, p. 246, *Keokuk Gr.*

præcursor, Herrick, 1888, *Bull. Denison Univ.*, vol. 3, p. 29, *Waverly Gr.*

serraticandata, Herrick, 1888, *Bull. Denison Univ.*, vol. 4, p. 52, *Coal Meas.*

shumardi, Herrick, 1887, *Bull. Denison Univ.*, vol. 2, p. 69. Proposed instead of *Proetus missouriensis* of Shumard.

truncleata, Herrick, 1887, *Bull. Denison Univ.*, vol. 2, p. 64, *Coal. Meas.*

PLACENTULA, Jones & Holl, 1886, *Ann. and Mag. Nat. Hist.*, ser. 5, vol. 17, p. 407.

[Ety. *placentula*, a little cake.] Valves suborbicular, nearly semicircular on the ventral border, straight on the dorsal margin inside, but projecting with unequal and variable angles at the outer dorsal region; surface flat or slightly convex, surrounded by a raised rim, which slopes down suddenly outside the edge of the valve; this rim incloses a depressed and reticulated area, and in or near the antero-dorsal region there is a small depression defined by a raised, loop-like border. Type *P. excavata*.

inornata, and *P. marginata*, Ulrich, 1890, *Jour. Cin. Soc. Nat. Hist.*, vol. 13, p. 124, *Hud. Riv. Gr.*

POLYCOPE, Sars, 1865, *Översigt af Norges Marine Ostracoder.*

sublenticularis, Jones, 1890, *Quar. Jour. Geo. Soc.*, vol. 46, p. 550, *Anticosti Gr.*

PONTOCYPRIS illinoisensis, Ulrich, 1890, *Jour. Cin. Soc. Nat. Hist.*, vol. 13, p. 107, *Hud. Riv. Gr.*; and *P. acuminata*, p. 210, *Waverly Gr.*

PRIMITIA centralis, *P. perminima*, Ulrich, 1890, *Jour. Cin. Soc. Nat. Hist.*, vol. 13, p. 130, *Utica Slate*; and *P. glabra*, *P. impressa*, *P. medialis*, *P. milleri*, *P. nodosa*, *P. rudis*, pp. 131 to 136, *Hud. Riv. Gr.*; and *P. nitida*, *P. sculptilis*, pp. 135, 136, *Trenton Gr.*; and *P. cestrinensis*, *P. cestrinensis* var. *caldwellensis*, *P. granimarginata*, *P. simulans*, and *P. subequata*, pp. 201, 202, *Kaskaskia Gr. clarkei*, Jones, 1890, *Quar. Jour. Geo. Soc.*, vol. 46, p. 535, *Corniferous Gr.*; *P. walcotti*, p. 543, *Corniferous Gr.*; *P. billingsi*, p. 547, *Clinton Gr.*; *P. seminulum*, p. 5, *Ham. Gr.*; *P. ulrichi*, and *P. whitfieldi*, pp. 6, 9, *Utica Slate*.

lativia and *P. parallela*, ulrich, 1889, *Micro-Pal. of Can.*, pp. 51, 52, *Hud. Riv. Gr.*

minuta, Eichwald, 1854, (*Cypridina minuta*), *Bull. Imp. Soc. Nat. Moscow*, vol. 27, p. 99, *Hud. Riv. Gr.*

mundula var. *effusa*, Jones, 1891, *Cont. to Can. Micro Pal.*, p. 64, *Chazy Gr.*; and *P. incisa*, p. 64, *Trenton Gr.*; and *P. scitula*, p. 91, *Devonian*.

PRIMITIOPSIS, Jones, 1887, *Sil. Ostrac. Goth-*

1888, Bull. Denison
Keokuk Gr.
191, Kansas City Sci.,
al. Meas.
ithides portlocki.
1888, Trans. N. Y.
246, Keokuk Gr.
1888, Bull. Denison
Waverly Gr.
k, 1888, Bull. Deni-
52, Coal Meas.
1887, Bull. Denison
Proposed instead of
sis of Shumard.
1887, Bull. Denison
Coal. Meas.
1886, Ann. and
r. 5, vol. 17, p. 407.
little cake.] Valves
semicircular on the
sight on the dorsal
projecting with un-
angles at the outer
edge flat or slightly
d by a raised rim,
suddenly outside the
this rim incloses a
ulated area, and in
dorsal region there is
defined by a raised,
type P. excavata.
1890, Ulrich, 1890,
Hist., vol. 13, p. 124,
Översigt af Norges
1890, Quar. Jour.
p. 550, Anticosti Gr.
sis, Ulrich, 1890,
Hist. vol. 13, p. 107,
d P. acuminata, p.
perminima, Ulrich,
Nat. Hist., vol. 13,
and P. glabra, P.
malis, P. milleri, P.
p. 131 to 136, Hud.
titida, P. sculptilis,
n Gr.; and P. cestrin-
var. caldwelensis,
P. simulans, and P.
202, Kaskaskia Gr.
ar. Jour. Geo. Soc.,
ferous Gr.; P. wal-
ferous Gr.; P. bil-
n Gr.; P. seminun-
P. ulrichi, and P.
Utica Slate.
ulrichi, Ulrich, 1889,
pp. 51, 52, Hud.
54, (Cypridina mi-
Nt. Moscow, vol.
Gr.
Jones, 1891, Cont.
p. 64, Chazy Gr.;
Trenton Gr.; and
onian.
7, Sil. Ostrac. Goth-

land, p. 5. [Ety. from the resemblance
to Primitia.] Like Primitia externally,
except that the anterior end has a
specially smooth area, corresponding
with an internal portion which is par-
titioned off from the rest of the cavity
by a cross wall. Type P. planifrons.
punctulifera, Hall, 1860, (Leperditia punc-
tulifera,) 13th Rep. N. Y. St. Mus. Nat.
Hist., p. 92, Ham. Gr.
PROETUS determinatus, Foerste, 1887, Bull.
Denison Univ., vol. 2, p. 91, Niagara Gr.
longicaudus, Hall, synonym for Phillipsia
major.
minutus, Herrick, 1888, Bull. Denison
Univ., vol. 4, p. 56, Waverly Gr.
PTYCHOPARIA attleboroensis, Shaler & Foerste,
1888, Bull. Mus. Comp. Zool., vol. 16,
p. 39, Up. Taconic.
metisensis, Walcott, 1890, 10th Ann. Rep.
U. S. Geo. Sur., p. 651, Up. Taconic.
mucronata, Shaler & Foerste, synonym for
Atops trilineata.
Rusichnites, synonym for Rusophycus and
R. acadicus; R. carbonarius, R. clinton-
ensis, and R. grenvillensis should be
referred to Rusophycus.
Sao, Barrande, 1846, Notice preliminaire,
p. 13, and Syst. Sil. Boh., vol. 1, p. 382.
[Ety. mythological name.] Body ovate,
trilobation marked; head subsemicir-
cular; glabella prominent, well defined;
dorsal furrows deep; three lateral fur-
rows, between which are lobes in relief,
separated on the summit by a longitudi-
nal furrow; facial sutures cut the
frontal border and arch outwardly to
the anterior projections of the eyes, and
the posterior branches arch in like man-
ner to a point at the interior of the genal
angle; the eye arch is prolonged in re-
lief toward the front of the glabella;
thorax seventeen segments; pygidium
small, two articulations. Type S. hir-
suta.

lamontensis is from the Chazy Gr.

FIG. 125.—*Schmidtdella crassimarginata*. Interior and exterior views of right valve and anterior and ventral views, magnified 10 diameters.

SCHMIDTDELLA, Ulrich, 1892, Am. Geo., vol.
10, p. 269. [Ety. proper name.] Carapace
small, rounded, moderately convex and
near *Aparchites* among the *Leperditidae*;
valves inflated in the dorsal region,
which projects shoulder-like over
and out from a nearly straight hinge-
line; right valve slightly the larger, its

ventral edge overlapping that of the
left; no sulcus or tubercles. Type
S. crassimarginata, described at the
same place from the Birdseye limestone.
SOLENOCARIS having been preoccupied before
Meek used it, Vogdes proposed *Strig-
ocaris* in 1889, Ann. N. Y. Acad. Sci.,
vol. 5, p. 34.
SOLENOPLEURA bombifrons, Matthew, 1887,
Trans. Roy. Soc. Can., vol. 4, p. 156, Up.
Taconic.
harveyi and *S. howleyi*, Walcott, 1889,
Proc. U. S. Nat. Mus., vol. 12, p. 45,
Up. Taconic.
STREPULA, Jones & Holl, 1886, Ann. and Mag.
Nat. Hist., ser. 5, vol. 17, p. 403. [Ety.
diminutive of *strepus*, a stirrup, from the
loop-like pattern of the ridges.] Carap-
pace valves slightly convex, suboblong,
with rounded ends, or semielliptical,
and bear narrow ridges that run into
the slightly thickened dorsal margin;
the intervening furrows form broad
valleys, and a subcentral tubercle or
lobular swelling is sometimes present;
the chief ridge is a free supramarginal
lamina, standing outward and down-
ward, and hiding the real marginal
edge in the side view; the edge of the
bivalved carapace is narrow, ovate,
cross-barred at the sides with ridges,
some straight and parallel, some ob-
liquely divergent. Type *S. concentrica*.
lunatifera, Ulrich, 1889, Micro-Pal. of Can.,
p. 56, Hud. Riv. Gr.
sigmoidalis and *S. plantaris*, Jones, 1890,
Quar. Jour. Geo. Soc., vol. 46, pp. 11,
540, Ham. Gr.
Tetradella, Ulrich, 1890, Jour. Cin. Soc. Nat.
Hist., vol. 13, p. 112. Carapace oblong
or subquadrate, never tumid, hinge-line
straight; surface depressed, a ridge fol-
lows the margin from the posterior to
the antero-dorsal angle; in the inclosed
space two narrow ridges traverse the
valves in a vertical or oblique direction
from the dorsal edge, to the posterior
half of the ventral portion of the sub-
marginal ridge, uniting with it; the
union of these ridges is supposed to be
the most significant character of the
genus. Type *T. quadrilobata*, which is
a straight synonym for *Beyrichia* regu-
laris of Emmons. Mr. Ulrich refers to
this genus *Beyrichia oculifera* and *B.*
chambersi. I do not agree with him in
respect to the genus, or in what he
esteems the important characters.
subquadrans, Ulrich, 1890, Jour. Cin. Soc.
Nat. Hist., vol. 13, p. 115, Trenton Gr.
TURRIERAS canadensis, Woodward, 1889,
Lond. Geo. Mag., 3d ser., vol. 6, p. 274,
Utica slate.
ULRICHIA, Jones, 1890, Quar. Jour. Geo. Soc.,
vol. 46, p. 543. [Ety. proper name.] In
form like *Primitia*, except there is no
sulcus across the valves, but instead a
tubercle on each side of the position in
which the sulcus occurs in *Primitia*; a

very weak generic distinction. Type
U. conradi, described at the same place
from Thedford, Canada.
confluens and U. emarginata, Ulrich, 1890,

Jour. Cin. Soc. Nat. Hist., vol. 13, p.
203, Kaskaskia Gr.
Zacanthoides eutoni, Walcott. Not defined.

CLASS ARACHNIDA.

In this Class (*arachne*, a spider,) the body is divided into segments, some of which are provided with articulated appendages, and a pair of ganglia is developed in each somite. The integument is hardened with chitine. The segments of the head and thorax are united, generally, into a cephalothorax. Instead of antennæ there are chelicere (*chela*, a claw; *keras*, a horn,) or mandibles (*mandibulum*, a jaw.) There are maxillæ (*maxillæ*, jaws,) carrying long jointed maxillary palpi, (*palpo*, I feel.)

ARCHITARBUS elongatum, Scudder, 1890,
Mem. Bost. Soc. Nat. Hist., vol. 4, p.
449, Coal Meas.
rotundatus, read A. rotundatum.
GERATARBUS lacoel and G. scabrum, Scudder,
1890, Mem. Bost. Soc. Nat. Hist., vol. 4,
p. 448, Coal Meas.

GREOPHONUS carboarius, Scudder, 1890,
Mem. Bost. Soc. Nat. Hist., vol. 4, p.
454, Coal Meas. This is Fig. 1074, on
page 575, over the name Archimylacris
acadicum.
KUSTARACHNE tenuipes, Scudder, 1890, Mem.
Bost. Soc. Nat. Hist., vol. 4, p. 450, Coal
Meas.

CLASS MYRIOPODA.

THIS Class (*myrios*, countless; *podes*, feet,) has the mouth on the under side, provided with mandibles and maxillæ. The head bears a pair of antennæ,

ACANTHERPESTES inequalis, Scudder, 1890,
Mem. Bost. Soc. Nat. Hist., vol. 4, p.
424, Coal Meas.
AMYNILYPES, instead of Arynilesipes.
ARCHIULUS glomeratus, Scudder, 1890, Mem.
Bost. Soc. Nat. Hist., vol. 4, p. 436, Coal
Meas.
EILETICUS inequalis, Scudder, 1890, Mem. Bost.
Soc. Nat. Hist., vol. 4, p. 421, Coal Meas.
EUPHOBERIA cuspidata, Scudder, 1890, Mem.
Bost. Soc. Nat. Hist., vol. 4, p. 426, Coal
Meas.
granosa is represented by Fig. 1072.
hystricosus, Scudder, 1890, Mem. Bost. Soc.
Nat. Hist., vol. 4, p. 426, Coal Meas.
simplex, Scudder, 1890, Mem. Bost. Soc.
Nat. Hist., vol. 4, p. 426, Coal Meas.

spinulosus, Scudder, 1890, Mem. Bost.
Soc. Nat. Hist., vol. 4, p. 430, Coal
Meas.
tracta, Scudder, 1890, Mem. Bost. Soc.
Nat. Hist., vol. 4, p. 433, Coal Meas.
ILYODES divisa and I. elongata, Scudder,
1890, Mem. Bost. Soc. Nat. Hist., vol. 4,
p. 422, Coal Meas.
LATZELIA primordialis, Scudder, 1890, Mem.
Bost. Soc. Nat. Hist., vol. 4, p. 419, Coal
Meas.
PALENARTHUS impressus, Scudder, 1890,
Mem. Bost. Soc. Nat. Hist., vol. 4, p.
422, Coal Meas.
XYLOBIUS frustulentus, Scudder, 1890, Mem.
Bost. Soc. Nat. Hist., vol. 4, p. 438, Coal
Meas.

CLASS INSECTA.

THERE are six principal veins in typical wings arising from the anterior and posterior root, named as follows: The anterior vein at the margin of the wing is the marginal vein; this is followed by the mediastinal and scapular veins, that cut the

c. Hist., vol. 13, p.
ott. Not defined.

anterior margin toward the extremity of the wing; the externomedian vein is directed toward the tip of the wing, and the internomedian and aual veins terminate in the posterior margin of the wing.

segments, some of
anglia is developed
e segments of the
instead of antennæ
andibulum, a jaw.)
ary palpi, (*palpo*, I

us, Scudder, 1890,
at. Hist., vol. 4, p.
his is Fig. 1074, on
name Archimylacris
Scudder, 1890, Mem.
, vol. 4, p. 450, Coal

ARCHILEOSCOLEX corneus, Matthew, 1889,
Trans. Roy. Soc. Can., vol. 4, p. 59, Low.
Devonian.

ARCHIMYLACRIS acadica is represented by
Fig. 1075. Read A. parallela for A. par-
allelum.

paucinervis, Scudder, 1890, Mem. Bost.
Soc. Nat. Hist., vol. 4, p. 411, Coal Meas.
ENCÆNUS, read Eucænus on p. 576, and in
the Index on p. 658.

EPHEMERITES affinis, E. gigas, E. primordialis,
and E. simplex, are probably not insects.
Some of them are plants.

ETOBILATTINA mazona, instead of E. mazon-
ana.

occidentalis, Scudder, 1890, Mem. Bost. Soc.
Nat. Hist., vol. 4, p. 410, Coal Meas.

LITHOMYLACRIS is feminine, and read L.
angusta and L. pittstonana for L. an-
gustum and L. pittstonanum.

pauperata, Scudder, 1890, Mem. Bost. Soc.
Nat. Hist., vol. 4, p. 409, Coal Meas.

MYLACRIS is feminine, and read M. anthra-
cophila, M. antiqua, M. bretonensis, M.
lucifuga, M. ovalis, M. pennsylvanica,
instead of anthracophilum, antiquum,
bretonense, lucifugum, ovale, and penn-
sylvanicum.

NECUMYLACRIS is feminine; read N. lacoana
for N. lacoanum.

PAROMYLACRIS is feminine; read P. rotunda
for P. rotundum.

ampla, Scudder, 1890, Mem. Bost.
Soc. Nat. Hist., vol. 4, p. 408, Coal
Meas.

PROMYLACRIS is feminine; read P. ovalis
for P. ovale.

rigida and P. testudo, Scudder, 1890, Mem.
Bost. Soc. Nat. Hist., vol. 4, p. 403, Coal
Meas.

TITANOPHASMA, Brongniart, 1882, Comptes
rend. Acad. 95, 1228, instead of Titan-
ophama on page 581. The Type is T.
fayoli.

SUBKINGDOM VERTEBRATA.

CLASS PISCES.

he under side, pro-
tentæ,

1890, Mem. Bost.
ol. 4, p. 430, Coal

, Mem. Bost. Soc.
433, Coal Meas.
elongata, Scudder,
e. Nat. Hist., vol. 4,

Scudder, 1890, Mem.
, vol. 4, p. 419, Coal

us, Scudder, 1890,
at. Hist., vol. 4, p.

Scudder, 1890, Mem.
, vol. 4, p. 438, Coal

the anterior and pos-
of the wing is the
veins, that cut the

THE work by Agassiz, "Recherches sur les Poissons fossiles"—(Text,) Tome 1-5;
(Atlas,) Tome 1-5—was published in parts from 1833 to 1844, and very irregularly.
The copy in the Public Library at Cincinnati does not show when published. Wood-
ward & Sherborn, in "A Catalogue of British Fossil Vertebrata, 1890," have pub-
lished the dates at which each plate and part of the text was published. The list
covers about five pages. From this list of dates I make the following corrections:
Acanthodes, 1833; Acrolepis, 1834; Ceratodus, 1838; Chirolepis, 1835; Chomatod-
us, 1838; Coccosteus, 1842; Rep. Brit. Assoc., p. 85, Cochliodus, 1838; Ctenodus,
1838; Ctenacanthus, 1837; Ctenoptychius, 1838; Glyptolepis, 1843; Gyracanthus,
1837; Helodus, 1838; Holoptychius, 1838, Murch. Sil. Syst., p. 599, type H. no-
bilissimus; Oracanthus, 1837; Orodus, 1838; Palæoniscus, Blainville, 1818, Dict.
Hist. Nat., vol. 27, p. 320; Phyonemus, named but not described until 1855, Brit.
Pal. Foss., p. 638; Platysomus, 1835; Pleuracanthus, 1837; Pœcilodus, named but
not described until 1855, Brit. Pal. Foss., p. 638; Psephodus was described by
Morris & Roberts; Pterichthys, 1843; Pygopteris, 1834, type P. mandibularis.

ACANTHODES semistriatus, Woodward, 1892,
Geo. Mag., vol. 9, p. 3, Low. Devon-
ian.

Acantholepis, Newberry, 1875, was preoccu-
pied by Mayer in 1861 in Hymenoptera.
See Eczematolepis.

Actinophorus, Newberry, 1880, Pal. Fish N.

Am., p. 174. The name was preoccu-
pied by Creutz in 1799. See Tegeolepis.
clarkii, see Tegeolepis clarkii.

ANTLIODUS arcuatus, Newberry, 1880, Pal.
Fish N. Am., p. 208, St. Louis Gr.

ASTEROPTYCHIUS elegans, Newberry, 1880,
Pal. Fish N. Am., p. 176, Waverly Gr.

- BOTHRIOLEPIS leidyi*, Newberry, 1889, Pal. Fish N. Am., p. 111, Catskill Gr.; and *B. minor*, p. 112, Chemung Gr.
- CALLOGNATHUS*, Newberry, 1889, Pal. Fish N. Am., p. 69. [Ety. *kallós*, beautiful; *gnathos*, jaw.] Small fishes, of which only the mandibles are known; these are from one to three inches in length; the posterior end of the dentary bone flat, thin, spatulate, smooth; the anterior half narrower, thicker, and ornamented; the upper edge closely set with numerous subequal, conical, obtuse, blunt-pointed teeth. Type *C. regularis*, described at the same place from the Huron Shale, as well as *C. serratus* from the Cleveland Shale.
- Carcharopsis* was not defined by Agassiz, and the forms subsequently defined under that name belong to *Dicrenodus*.
- Cephalaspis laticeps* and *C. campbelltonensis*, Traquair, 1890, Lond. Geo. Mag., 3d ser., vol. 7, p. 16. Not defined so as to be recognized.
- CLADODUS carinatus*, Newberry, 1889, Pal. Fish N. Am., p. 103. Too poorly defined to warrant recognition; beside the name was preoccupied.
- kepleri*, Newberry, 1889, Pal. Fish N. Am., p. 103, and *C. terrelli*, p. 170, and *C. tunidus*, p. 172, Cleveland Shale.
- CELOSOMUS*, Newberry, 1889, Pal. Fish N. Am., p. 188. [Ety. *koilos*, hollow; *osteon*, bone.] Fishes of large size allied to *Dendrodus* and *Rhizodus*; only a coracoid, mandible, and tooth are known. The bones consist of a thin shell of dense osseous tissue inclosing large cavities, once doubtless filled with cartilage; the coracoid is about a foot in length and an inch and a half in diameter at the middle, and the central cavity is as large, relatively, as in the long bones of birds, the shell which surrounded it being but from one-eighth to one-quarter of an inch in thickness; the dentary bone is about one foot in length, two and a half inches wide in the middle, where it is one and a quarter inches in thickness, and four inches wide at the posterior end, where it was doubtless joined to the angular and articular elements; on the outside the posterior half is excavated to form a deep sulcus for the reception of the motor muscle; on the inside the jaw is flattened and gently arched downward to the rounded lower edge; the upper side bears on the outside a subacute toothless ridge; within and below this is a wide shoulder with seven broad, shallow pits, in which were planted the rounded bases of the teeth; teeth strong, conical, straight, acute and smooth above, plicated below; complicated interior structure as in *Dendrodus*. Type *C. ferox*, described at the same place from Carboniferous rocks.
- CTENACANTHUS angustus*, Newberry, 1889, Pal. Fish N. Am., p. 181, Esrea Grit; and *C. clarkii*, and *C. compressus*, p. 168, Cleveland Shale; and *C. cylindricus*, p. 202, Keokuk Gr.; and *C. littoni*, p. 201, St. Louis Gr.; and *C. randalli*, p. 105, Olean conglomerate. And he refers *C. parvulus* to *Hoplonchus*, a genus established by Davis in 1875, in Quar. Jour. Geo. Soc. Lond., with *H. elegans* as the type. See p. 169.
- CTENODUS wagneri*, Newberry, 1889, Pal. Fish N. Am., p. 172, Cleveland Shale.
- DICRENODUS*, Romanovsky, 1853, Bull. Soc. Imp. Moscou, p. 408. This name has priority over *Carcharopsis*, which was not defined by Agassiz, nor by any other one until 1883, and it has priority over *Pristicladodus*, both of which names Woodward & Sherborn say are synonyms.
- Dinichthys*, Newberry, 1873, Ohio Pal. vol. 1, p. 313. This name was preoccupied by Hitchcock, among the fishes, in 1868. See *Ponerichthys*.
- corrugatus*, *D. curtus*, *D. gouldi*, *D. hertzeri*, *D. intermedius*, *D. minor*, *D. precursor*, *D. terrelli*, *D. tuberculatus*. See *Ponerichthys*.
- Diplodus* was preoccupied in 1810, by Rafinesque, for a genus of Sparidae, before Agassiz used the word. See *Dissodus*.
- acinaces*, *D. compressus*, *D. gracilis*, *D. latus*, *D. penetrans*, *D. problematicus*. See *Dissodus*.
- DIPLOGNATHUS*, Newberry, 1878, Ann. N. Y. Acad. Sci., vol. 1, p. 188, and Pal. Fish N. Am., p. 159. [Ety. *diploos*, double; *gnathos*, the jaw.] Dentary bones long and slender, flattened, straight, spatulate behind, were originally covered with cartilage; anterior and exposed portions rising into points which diverge from the symphysis, giving a forked extremity to the lower jaw; conical acute teeth formed from the jaw tissue are set along the outer margin of the mandibles and on the inside of the divergent extremities beyond the symphysis; the teeth are recurved, and formed a kind of forked rake; a deep pit in each dentary bone marks the point of insertion of a powerful ligament, which bound the rami together and prevented splitting. Type *D. mirabilis*, which was described at the same place from the Cleveland Shale.
- DIPTEUS fiabelliformis*, *D. levis*, *D. minutus*, *D. nelsoni*, *D. radiatus*, Newberry, 1889, Pal. Fish N. Am., pp. 89 to 119, Chemung Gr.
- Dissodus* n. gen. [Ety. *dissos*, double; *odous*, tooth.] Proposed for *Diplodus*, Agassiz, 1843, Recherches sur les Poissons Fossiles, t. 3, p. 204, which name was preoccupied. The type is *D. gibbosus*, and the American species are *D. acinaces*, *D. compressus*, *D. gracilis*, *D. latus*, *D. penetrans*.
- problematicus*, Woodward, (*Diplodus problematicus*,) 1892, Geo. Mag., vol. 9, p. 2, Low. Devonian.

C. compressus, p. 169; and *C. cylindricus*, Gr.; and *C. littori*, Gr.; and *C. randalli*, p. 169. And he also to Hoplonchus, a fish by Davis in 1875, in Loc. Lond., with H. e. See p. 169.

Newberry, 1889, Pal. Fish. N. Am., p. 116, Cleveland Shale. See p. 169. This name has priority, which was given by Agassiz, nor by any other, and it has priority both of which names are born say are syn-

1873, Ohio Pal. vol. 1, p. 188, and Pal. Fish. N. Am., p. 116, Cleveland Shale. This name was preoccupied by Blyth, among the fishes, in 1868.

D. gouldi, *D. hertzeri*, *D. minor*, *D. precursor*, *D. us*. See Ponerichthys. Named in 1810, by Rafinesque, of Sparidae, before the word. See Dissodus. *D. gracilis*, *D. latus*, *D. problematicus*. See Dissodus.

Newberry, 1878, Ann. N. Y. Acad. Sci., vol. 1, p. 189, and Pal. Fish. N. Am., p. 116, Catskill Gr. [Ety. *diploos*, double; *dentary* bones long, straight, spatulate, originally covered with enamel, and marked by fine denticulate, longitudinal sutures, which divide the surface into broad, nearly equal bands or flattened ridges; the sutures are most numerous below, but terminate in succession above, so that few reach the conical pointed summit. Type *G. politus*. *G. politus*, Newberry, 1889, (Heteracanthus *politus*.) Pal. Fish. N. Am., p. 66, Ham. Gr.]

GANOBYNCHUS, Traquair, 1873, Lond. Geo. Mag., vol. 10, p. 552. [Ety. *ganos*, brightness; *rhynchus*, beak.] This genus was founded on the fragment of a snout 1½ inches long and 3 inches wide, supposed to belong to a fish 4 or 5 feet long. The country from which it came and its geological age are wholly unknown. Type *G. woodwardi*.

woodwardi, Newberry, 1889, Pal. Fish. N. Am., p. 95, Chemung Gr. GLYPTASPIS, Newberry, 1889, Pal. Fish. N. Am., p. 157. [Ety. *glyptos*, sculptured; *aspis*, shield.] Placoderm fishes of large size, protected by thick bony plates, of which those of the plastron were probably 5 in number; the middle one is lance-shaped or subrhomboidal, its central portion tuberculated, its margins sloped off and smoothed or striated by the overlap of the lateral ventral plates. The upper part of the body carried a number of lanceolate or ellipsoidal plates, of which the central parts are ornamented, the margins smooth and sloped down to thin edges; other parts unknown. Type *G. verrucosa*.

verrucosa, Newberry, 1889, Pal. Fish. N. Am., p. 158, Cleveland Shale. GLYPTOPOMUS, Agassiz, 1844, Monographie des

ECZEMATOLEPIS, n. gen. [Ety. *eczema*, a cutaneous eruption; *lepis*, scale.] Proposed for Acantholepis, Newberry, 1875, Ohio Pal., vol. 2, p. 38, which name was preoccupied. The type is *E. pustulosa*, Newberry, 1875, (Acantholepis *pustulosa*.) Ohio Pal., vol. 2, p. 38, Up. Held. Gr.

Eurylepis, Newberry, 1880, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., vol. 1, p. 350. This name was preoccupied by Blyth, among the reptiles, in 1854. See Haplolepis.

corrugata, *E. granulata*, *E. insculpta*, *E. lineata*, *E. minima*, *E. ornatissima*, *E. ovoidea*, *E. striolata*, *E. tuberculata*. See Haplolepis.

GAMPHACANTHUS, n. gen. [Ety. *gamphos*, bent; *akantha*, spine.] Proposed for Heteracanthus, Newberry, 1889, Pal. Fish. N. Am., p. 65, which name was preoccupied. Pectoral (?) spines eight inches or more in length, robust, with a posterior opening reaching to or near the summit; base compressed, one and a half inches wide, obliquely rounded below; shaft curved forward, regularly arched transversely, covered with highly polished enamel, and marked by fine denticulate, longitudinal sutures, which divide the surface into broad, nearly equal bands or flattened ridges; the sutures are most numerous below, but terminate in succession above, so that few reach the conical pointed summit. Type *G. politus*.

politus, Newberry, 1889, (Heteracanthus *politus*.) Pal. Fish. N. Am., p. 66, Ham. Gr.

GANOBYNCHUS, Traquair, 1873, Lond. Geo. Mag., vol. 10, p. 552. [Ety. *ganos*, brightness; *rhynchus*, beak.] This genus was founded on the fragment of a snout 1½ inches long and 3 inches wide, supposed to belong to a fish 4 or 5 feet long. The country from which it came and its geological age are wholly unknown. Type *G. woodwardi*.

woodwardi, Newberry, 1889, Pal. Fish. N. Am., p. 95, Chemung Gr.

GLYPTASPIS, Newberry, 1889, Pal. Fish. N. Am., p. 157. [Ety. *glyptos*, sculptured; *aspis*, shield.] Placoderm fishes of large size, protected by thick bony plates, of which those of the plastron were probably 5 in number; the middle one is lance-shaped or subrhomboidal, its central portion tuberculated, its margins sloped off and smoothed or striated by the overlap of the lateral ventral plates. The upper part of the body carried a number of lanceolate or ellipsoidal plates, of which the central parts are ornamented, the margins smooth and sloped down to thin edges; other parts unknown. Type *G. verrucosa*.

verrucosa, Newberry, 1889, Pal. Fish. N. Am., p. 158, Cleveland Shale.

GLYPTOPOMUS, Agassiz, 1844, Monographie des

Poissons fossiles du Vieux Gres Rouge on Systeme Devonien des Iles Britanniques et de Russie, p. 77. Type *G. minor*, of which Huxley figured and described nearly an entire specimen in 1866, in Brit. Organic Remains, dec. 12, (Mem. Geo. Sur.,) p. 4.

sayrei, Newberry, 1878, Ann. N. Y. Acad. Sci., vol. 1, p. 189, and Pal. Fish. N. Am., p. 116, Catskill Gr.

Goniodus, Newberry, 1880, Pal. Fish. N. Am., p. 67. This name was twice preoccupied, first by Agassiz in 1836, and by Dunker, in 1848, in the mollusca. See Xenodus *hertzeri*, Newberry, 1889, see Xenodus *hertzeri*.

GORGONICHTHYS, Claypole, 1892, Am. Geo., vol. 10, p. 1. [Ety. *Gorgon*, mythological name; *ichthys*, fish.] Founded upon the lower left mandible, which is similar to that in Ponerichthys, and the point of a premaxillary tooth, behind which there is a tooth terminating downward in two blunt processes, the larger one in front, which distinguishes it from the upper cutting blade in Ponerichthys. Type *G. clarki*, described at the same place from the Cleveland Shale.

GYRACANTHUS incurvus, Traquair, 1890, Geo. Mag., vol. 7, p. 21, Low. Devonian.

sherwoodi, Newberry, 1889, Pal. Fish. N. Am., p. 119, Catskill Gr.; and *G. inornatus*, p. 177, Waverly Gr.

HAPLOLEPIS, n. gen. [Ety. *haplos*, simple; *lepis*, scale.] Proposed for Eurylepis, Newberry, 1880, Proc. Acad. Nat. Sci. Phil., and Ohio Pal., vol. 1, p. 350. Type *H. tuberculata*, Newberry, 1886, (Eurylepis *tuberculata*.) Proc. Acad. Nat. Sci., and Ohio Pal., vol. 1, p. 350, Coal Meas. The species are *H. corrugata*, *H. granulata*, *H. insculpta*, *H. lineata*, *H. minima*, *H. ornatissima*, *H. ovoidea*, *H. striolata*, *H. tuberculata*.

HARPACANTHUS, Traquair, 1886, Ann. and Mag. Nat. Hist. A sickle-shaped spine, with *H. fimbriatus*, from Scotland, as the type. Cited by Newberry in Pal. Fish. N. Am., p. 203.

Heteracanthus, Newberry, 1889, Pal. Fish. N. Am., p. 65. The name was preoccupied in 1836 by Diesing, among the Vermes. See Gamphacanthus.

politus, Newberry, see Gamphacanthus *politus*.

HOLONEMA, Newberry, 1889, Pal. Fish. N. Am., p. 92. [Ety. *holos*, entire; *nema*, thread.] A placoderm fish of medium size, having the body inclosed in armor made up of polygonal plates, of which the external surface is entirely covered by radiating lines of enamel; the central plate of the plastron is coffin shaped, pointed before, broadest near the anterior end, where the sides are produced into prominent lateral angles; from this point backward it narrows to a truncated end, which is half as wide as the greatest

- breadth. Other parts unknown. Type *H. rugosum*.
- rugosum*, Claypole, 1883, (*Pterichthys rugosa*.) Proc. Am. Phil. Soc., vol. 20, p. 664, Chemung Gr.
- HOLOPTYCHUS** *granulatus*, H. *pustulosus*, H. *tuberculatus*, Newberry, 1889, Pal. Fish N. Am., pp. 100, 101, Chemung Gr.; and H. *halli* and H. *radiatus*, pp. 114, 115, Catskill Gr.
- ICANODUS**, n. gen. [Ety. *ikanos*, befitting; *odous*, tooth.] Proposed for *Tomodus*, Agassiz, MSS. and St. John & Worthen, 1883, Geo. Sur. Ill., vol. 7, p. 171, which name was preoccupied among the fishes by Trautschold, in 1879. Type I. *limitaris*.
- limitaris*, St. John & Worthen, 1883, (*Tomodus limitaris*.) Geo. Sur. Ill., vol. 7, p. 173, Up. Burlington Gr.
- JANASSA** has for its type *J. bituminosa*.
- Labodus marginatus*, Newberry, 1889, Pal. Fish N. Am., p. 198, St. Louis Gr. The genus *Labodus* was not defined by Agassiz. Davis used it in 1883, but in 1889, Woodward & Sherborn classed it as a synonym for *Copodus*. The species illustrated by Newberry will belong to some other genus, probably not yet defined.
- Liognathus*, Newberry, 1873, Ohio Pal., vol. 1, p. 306. The name was preoccupied among the fishes by Lacepede, in 1802. See *Lispognathus*.
- spatulatus*, see *Lispognathus spatulatus*.
- LISPOGNATHUS**, n. gen. [Ety. *lispos*, smooth; *gnathos*, jaw.] Proposed for *Liognathus*, Newberry, 1873, Ohio Pal., vol. 1, p. 306, which name was preoccupied. Type *L. spatulatus*.
- spatulatus*, Newberry, 1873, (*Liognathus spatulatus*.) Ohio Pal., vol. 1, p. 306, Up. Held. Gr.
- MAZODUS**, Newberry, 1889, Pal. Fish N. Am., p. 178. [Ety. *maza*, a barley-cake; *odous*, tooth.] Teeth of Elasmobranch or Selachian fishes, often of large size, thick and massive, with an ovoid, elliptical, or angular outline; upper surface arched in both directions, smooth or finely granulated; under surface concave, coarsely pitted, and variously furrowed and lobed; sides marked by irregular, often pustulous ridges; interior similar throughout, showing irregular, vertical, calcigerous tubes or columns closely compacted into a dense, hard, and enamel-like tissue; mandibles 7 to 8 inches long, and 1½ inches to 1½ inches wide. Type *M. kepleri*.
- kepleri*, Newberry, 1889, Pal. Fish N. Am., p. 180, Cuyahoga Shale.
- MILLERICHTHYS**, n. gen. Proposed for *Pterichthys*, Agassiz, 1843, Poiss. Foss., vol. 2, p. 302, which name was preoccupied among the fishes by Swainson, in 1839. [Ety. the name is proposed in honor of Hugh Miller, who was really the first to fully characterize and illustrate the genus in his work on "The Old Red Sandstone."] Type *M. milleri*. It is doubtful whether or not this genus is represented in North America.
- MYLOSTOMA**, Newberry, 1883, Trans. N. Y. Acad. Sci., vol. 2, p. 146, and Pal. Fish N. Am., p. 161. [Ety. *mylos*, a grinder; *stoma*, mouth.] Teeth consist of strong and massive tables of bony tissue, becoming more dense and enamel-like toward the triturating surface; they apparently formed several pairs on both the upper and lower jaws; the principal plates of the lower jaw had long oval or spatulate crowns, 3 to 6 inches in length by 1 to 2 inches wide, and half an inch or more in thickness, supported by strong, vertical, spatulate bones, which projected downward and backward, terminating posteriorly in thin, rounded margins. The upper surface of the crown is raised into a more or less prominent tubercle, which is situated near the exterior margin, and slightly anterior to the middle; another pair are triangular in outline; the dental plates of the upper jaw are tabular and consist of very dense tissue. Type *M. variabile*.
- terrelli*, *M. variabile*, Newberry, 1883, Trans. N. Y. Acad. Sci., vol. 2, p. 146, and Pal. Fish N. Am., p. 164, Cleveland Shale.
- OSTOPHORUS**, n. gen. Proposed for *Sphenophorus*, Newberry, 1889, Pal. Fish N. Am., p. 92, which name was preoccupied by Schoenherr, in 1838, among the Coleoptera. [Ety. *ostos*, an arrow; *phoros*, bearing.] The clavicle is a flattened bone, 6 inches or more in length by 1½ inches in width at the middle, narrowing to either end; the anterior margin strongly reflexed; the exterior surface is marked by many rows of relatively large arrowhead-like tubercles, closely set one behind the other, the points directed forward; other parts unknown. Type *O. lilleyi*.
- lilleyi*, Newberry, 1889, (*Sphenophorus lilleyi*.) Pal. Fish N. Am., p. 92, Chemung Gr.
- ONYCHODUS ortonii**, Newberry, 1889, Pal. Fish N. Am., p. 71, Ham. Gr.
- PETALORHYNCHUS**, Agassiz, MSS. only, Newberry & Worthen, 1866, Geo. Sur. Ill., vol. 2, p. 32.
- PHANEROPLEURON**, Huxley, 1859, Anderson's Dura. Den., p. 67. Type *P. andersoni*.
- PHGODUS politus**, Newberry, 1889, Pal. Foss. N. Am., p. 173, Cleveland Shale.
- PHYLOLEPIS**, Agassiz, 1844, Poiss. Foss. Vieux Gres Rouge, etc., p. 67. [Ety. *phyllon*, a leaf; *lepis*, scale.] Thin, more or less elliptical scales. Type *P. concentricus*.
- delicula*, Newberry, 1889, Pal. Fish N. Am., p. 97, Chemung Gr.

in his work on
stone." Type M.
ful whether or not
resented in North

1883, Trans. N. Y.
146, and Pal. Fish
y. *mulos*, a grinder;
th consist of strong
s of bony tissue,
se and enamel-like
ing surface; they
several pairs on
d lower jaws; the
the lower jaw had
ate crowns, 3 to 6
1 to 2 inches wide,
more in thickness,
e, vertical, spatulate
ted downward and
ting posteriorly in
ins. The upper sur-
s raised into a more
tubercle, which is
terior margin, and
the middle; another
in outline; the den-
per jaw are tabular
dense tissue. Type

e, Newberry, 1883,
Sci., vol. 2, p. 146, and
p. 164, Cleveland

Proposed for Sphe-
erry, 1889, Pal. Fish
ch name was pre-
herr, in 1838, among
Ety. *oistos*, an arrow;
The clavicle is a flat-
s or more in length
dth at the middle,
r end; the anterior
flexed; the exterior
by many rows of
rowhead-like tubere-
e behind the other
ed forward; other
type O. lilleyi.

1889, (Sphenophorus
N. Am., p. 92, Che-

ewberry, 1889, Pal.
Ham. Gr.

assiz, MSS. only,
hen, 1866, Geo. Sur.

ey, 1859, Anderson's
7. Type P. ander-

ewberry, 1889, Pal.
73, Cleveland Shale.
1844, Poiss. Foss.
e, etc., p. 67. [Ety.
lepis, scale.] Thin,
cal scales. Type P.

, 1889, Pal. Fish N.
ng Gr.

PHYSONEMUS altonensis, see *Stethacanthus*
altonensis.

stellatus, Newberry, 1889, Pal. Fish N.
Am., p. 200, St. Louis Gr.

PONERICHTHYS, n. gen. [Ety. *poneros*,
wicked; *ichthys*, a fish.] Proposed for
Dinichthys, Newberry, 1873, Ohio Pal.,
vol. 1, p. 313, and vol. 2, p. 3, which
was preoccupied among the fishes by
Hitchcock, in 1888. Type P. *terrelli*.

corrugatus, P. *curtus*, P. *gouldi*, P. *inter-*
medius, P. *minor*, Newberry, 1889, Pal.
Fish N. Am., pp. 149 to 156, Cleveland
Shale.

hertzeri, Newberry, 1873, (Dinichthys
hertzeri.) Ohio Pal., vol. 1, p. 316, Por-
tage Gr.

præcursor, Newberry, 1889, Pal. Fish N.
Am., p. 51, Corniferous Gr.

terrelli, Newberry, 1873, (Dinichthys *ter-*
relli.) Ohio Pal., vol. 1, p. 313, and vol.
2, p. 3, Portage Gr.

tuberculatus, Newberry, 1889, Pal. Foss.
N. Am., p. 98, Chemung Gr.

Pristicladodus is said to be a synonym for
Dicrenodus, by Woodward & Sherborn.
PAISTONUS, Agassiz, MSS. only, Davis, 1883,
Trans. Roy. Soc. Dub., vol. 1, p. 519.

PROTODUS, Woodward, 1892, Lond. Geo.
Mag., vol. 9., p. 1. [Ety. *protos*, first;
odus, tooth.] A Selachian tooth; dental
crown consists of a single robust, solid,
conical cusp, invested with gano-den-
tine; root large, undivided, laterally
expanded and antero-posteriorly com-
pressed. Type P. *jexi*, described at the
same place from the Low Devonian.

Pterichthys was preoccupied by Swainson
among the fishes in 1839. See Miller-
ichthys.

rugosus, see *Holonema rugosum*.

RINZODUS anceps, Newberry, 1889, Pal.
Fish N. Am., p. 191, St. Louis Gr.

RHYNCHODUS greenei, Newberry, 1889, Pal.
Fish N. Am., p. 51, Ham. Gr.

Sphenophorus, Newberry, 1889, Pal. Fish N.
Am., p. 92. The name was preoccupied
among the Coleoptera, by Schoenherr,
in 1838. See *Oestophorus*.

STETHACANTHUS, Newberry, 1889, Pal. Fish
N. Am., p. 198. [Ety. *stethos*, the breast;
akantha, a spine.] Pectoral spines of
medium or large size, unsymmetrical,
(rights and lefts,) broadly falcate in
outline, the conical summit compressed,
with anterior and posterior margins
rounded; below the solid summit the
posterior margin is opened by a deep
sulcus, of which the walls, of unequal
thickness, terminate posteriorly in thin
and fragile edges; anterior border gen-
tly concave, about one third its length
from the base rising into a strong, often
tumid, shoulder; basal portion narrow
and compressed, terminating in a car-
tilaginous condyle for articulation. In
life the posterior sulcus was occupied
by the base of the pectoral fin. Type
S. *altonensis*.

altonensis, St. John & Worthen, 1875,
(*Physonemus altonensis*.) Geo. Sur.
Ill., vol. 6, p. 454, St. Louis Gr.

tumidus, Newberry, 1889, Pal. Fish N.
Am., p. 198, Berea Grit.

TEGEOLEPIS, n. gen. [Ety. *tegeos*, a roof;
lepis, scale.] Proposed for *Actinophorus*,
Newberry, 1889, Pal. Fish N. Am., p.
174, which was preoccupied among the
Coleoptera, by Crentz, in 1790. Tile-
scaled ganoids, of medium or large
size, long and slender; body cylindrical;
head pointed, bony; teeth numerous,
conical, acute; fins without fulcrum, deli-
cate, many-rayed; scales narrow, quad-
rangular, thin. Type T. *clarkii*.

clarkii, Newberry, 1889, (Actinophorus
clarkii.) Pal. Fish N. Am., p. 175, Cleve-
land Shale.

TITANICHTHYS, Newberry, 1889, Pal. Fish
N. Am., p. 130. [Ety. mythological
name; *ichthys*, a fish.] Cranium of
more gigantic size than that of *Poner-*
ichthys, being about 4 feet broad at the
occiput, 3 feet or more in length, trian-
gular in outline, and marked by incised
lines; supra-scapulas or supra-clavicles
rhomboidal or trapezoidal in outline,
from 15 to 20 inches in diameter;
clavicles 2 feet in length by eight inches
in breadth, turned forward and nar-
rowed at the lower end; mandibles 2 or
3 feet in length, subcylindrical or sub-
triangular rods curved upward ante-
riorly and furrowed like a gouge. Type
T. *agassizi*, which, with T. *clarkii*, is de-
scribed at the same place from the
Cleveland Shale.

Tomodus, Agassiz, MSS., and St. John &
Worthen, 1883, Geo. Sur. Ill., vol. 7, p.
171, was preoccupied among the fishes
by Trantschold, in 1879. See *Icanodus*.

limitaris, see *Icanodus limitaris*.

TRACHOSTEUS, Newberry, 1889, Pal. Fish N.
Am., p. 166. [Ety. *trachys*, rough; *osteon*,
a bone.] Placoderm fishes inclosed in
defensive armor, consisting of a number
of large, but relatively thin, bony
plates, of which the outer enameled
surface is thickly set with high conical
tubercles, that are acute, rounded, or
cupped at the summit; the spaces be-
tween these tubercles are radiately
lined; under jaws consisted of cartilagi-
nous, angular, and articular parts with
dense bony dentary portions; dental
bones straight, posterior end spatulate,
anterior third or exposed part carries a
row of slender, conical, acute teeth
along its upper margin; premaxillaries
subtriangular in outline, anterior face
arched, and terminating below in an
acute point; posterior edge horizontal,
and carrying slender, pointed teeth,
which matched with a portion of those
of the mandible; eye orbits relatively
large and round, encircled by a ring
composed of four sclerotic plates, those
on one side narrower than the others;

surface in part tuberculated. Type *T. clarkii*, which is described at the same place from the Cleveland Shale.

XENODUS, n. gen. [Ety. *xenos*, strange; *odus*, tooth.] Proposed for *Goniodus*, Newberry, 1880, Pal. Fish. N. Am., p. 67, which was preoccupied among the fishes, by Agassiz, in 1836, and among the Mollusca by Dunker, in 1848. Teeth numerous, composing a roughened pavement, small, variable in size and

form; generally subtriangular in outline, depressed, with the central portion elevated into an obtuse angular ridge of denser tissue, and having a polished surface; other portions of the crown and the lateral margins roughened by a vermicular pitted or corrugated marking; the lower surface rough and bone-like. Type *X. hertzeri*, which was described at the same place from the Ham. Gr.

CLASS BATRACHIA.

CERATERPETON, Huxley, 1867, Trans. Roy. Irish Acad., vol. 24, p. 354. Type *C. galvani*.

GLOSSARY.

ERRATA.

Altilis, e—Fattened, instead of flattened.
Coriformis, e—Heart-shaped, instead of like *Coris*.
Craticulus, instead of *Cratiulus*.
Eugenium, noble; instead of *Euginum*, fertile.

Macronotus, a, um—Having a long back, instead of long known.
Ornognanulus, instead of *Ornigranulus*.
Vadosus, a, um—Full of shallows, instead of shadows.

INDEX OF GENERA.

ERRATA.

Eucænus, instead of *Encænus*.
Lithomylacris, f.
Mylacris, f.

Necomylacris, f.
Paromylacris, f.
Promylacris, f.

subtriangular in outline, the central portion of the angular ridge of the crown having a polished surface, the margins of the crown roughened by corrugated markings, the base rough and bone-colored, which was derived from the

354. Type *C. galvani*.

having a long back, in-
volved.
of *Ornigranulus*.
of shallows, instead of

SECOND APPENDIX

TO

NORTH AMERICAN GEOLOGY AND PALÆONTOLOGY.

OCTOBER, 1897.

[THIS Appendix will be mailed to any one on receipt of \$1.00; and hereafter it will be mailed with the bound volume, including the First Appendix, on receipt of \$5.00. Address, S. A. Miller, Court and Walnut, Cincinnati, Ohio.]

THE Report of the Geological Survey of Ohio, Vol. VII, was published in February, 1895. The title page is dated 1893. The map which faces the title-page bears the date of 1894, made by the engraver. The article on "Recent Changes in Nomenclature," by W. A. Kellerman, on page 80a, is dated "Ohio State University, January, 1895." The volume was not published in parts, but made its first appearance in the latter part of February, 1895.

The plates and manuscript on Palæontology, for the Eighteenth Report of the Geological Survey of Indiana, were completed and delivered to the State printer in the spring of 1892. Early in June of that year, Mr. Beachler, who was in the employment of Mr. Wachsmuth, asked one of the assistant State geologists for the privilege of examining the plates and manuscript, which was refused; he then tried to see them in the hands of the State printer, but again failed. This led the State geologist, Professor S. S. Gorby, to order the work printed as advance sheets of the Report. Some light is thrown upon the subject by the letter published in Bulletin No. 4, p. 37, of the Illinois State Museum of Natural History. The printing was commenced in July, but was delayed, for some reason, until August. I went to Indianapolis and read the last of the proof on the 28th of August, and took home, to Cincinnati, on that night, a copy of the work without a cover. It was to be issued on the 1st day of September, and hence bears the date "September, 1892." Mr. E. T. J. Jordan, the assistant State geologist, informed me by letter that he filled the mailing-list on the first day of September, and I was also assured by Professor S. S. Gorby that it was done on that day, and I received two copies by mail, one of which I gave to Charles L. Faber, in the first week of September. Later a box of the books was sent to me by freight, which I commenced to distribute in September. From these I sent twenty copies to K. A. Blair, at Sedalia, Missouri, who was at McClellan Springs for his health; but he has the written evidence that he received them on the 14th day of October, 1892. The statements, therefore, of Mr. Wachsmuth, in *Crinoidea Camerata*, pages 200 to 203, that the advance sheets of the 18th Indiana Report were not printed until October 26, 1892, are untrue.

In 1894, Vols. IV and V of the Missouri Geological Survey on Paleontology appeared. They would not, probably, be worth mentioning were it not for the fact that every intelligent definition is taken literally from the writings of the late Professor Meek, in Paleontology of Ohio, Paleontology of Eastern Nebraska, Proc. Acad. Nat. Sci. Phil., or from some other deceased paleontologist, without using quotation marks, or otherwise giving the credit to the real author.

In 1879, Mr. Ulrich figured and described, in the Jour. Cin. Soc. Nat. Hist., what he called *Tellinomya cingulata*. In 1895, Ohio Geology, Vol. VII, page 680, he figured and described another and different fossil, for the species, and said that the hinge teeth "were misrepresented in the original description and figures." In the latter instance we have a specimen of *Tellinomya peetunculoides*, and the species would have been known from the first publication, had it not been for the misrepresentations in the original description and figures. In the same Journal he figured and described what he called *Crytolites nitidulus*. In 1897, Geo. of Minn., Vol. III, page 866, he figured and described for it another fossil, and called it *Crytolitina nitidula*, and said: "The original description and figures are incorrect, where they differ from the present work on the species." In the latter case, we have the fragment of a cast of *Crytolites carinatus*, and it would have been known from the first publication, if the original description and figures had not been erroneous or fictitious. *Stomatopora proutana* was described and illustrated, in 1882, in the Jour. Cin. Soc. Nat. Hist., Vol. V, page 39, plate 1. In 1886, in the 14th Ann. Rep. Minn. Sur., page 59, he described, without illustration, *Ropalonaria pertenuis*. In 1893, Geo. Sur. Minn., Vol. III, page 117, he classed *Ropalonaria pertenuis* as a synonym for *Stomatopora proutana*; and, on page 116, plate 1, and in Jour. Cin. Soc. Nat. Hist., Vol. XII, page 175, he described and illustrated *Stomatopora proutana* under the name of *Stomatopora tenuissima* n. sp. These instances may indicate what reliance, if any, is to be placed on his "original description and figures." The synonymy in Ohio Geology, Vol. VII, and in Geo. Sur. Minn., Vol. III, if not appalling, is without a parallel in natural history.

Definitions of invertebrate palæozoic fossils, without illustration, are worse than worthless. They are worthless, because no one, other than the author, can determine whether fossils collected elsewhere than at the typical locality belong to the proposed species, and no comparison can be made with them. They are obstacles because they demand attention to which they are not entitled, and therefore they are much worse than catalogue names, which are simply worthless. I have condemned them in this Appendix, and feel that even that much recognition is going too far. It would probably be better not to notice them at all. No well-conducted journal, at this day, will admit such matter to its columns.

When three or more species have been described in any genus, the genus may be divided into groups; but giving names to such groups, for the purpose of having them applied to the species, is not binomial, or consistent with any judicious system of nomenclature. There is no science in subgeneric or group names, and they should be wholly disregarded.

VEGETABLE KINGDOM.

PLANTÆ.

- Archæophyton newberryanum*, Britton. Applied to films of graphite.
- BYTHOGRAPTUS latus*, Hall, has been shown (?) to belong to the Algae by Whitfield, Bull. Am. Mus. Nat. Hist., vol. 6, p. 331. He further suggests that *Bythocladius* would be more appropriate. I can not understand how an Algae can be mistaken for a Graptolite, or a Graptolite mistaken for an Algae.
- CALLITHAMNOPSIS*, Whitfield, 1894, Bull. Am. Mus. Nat. Hist., vol. 6, p. 354. [Ety. *Callithamnion*, a genus of Algae; *opsis*, resemblance.] Frond articulate, branched, branches opposite in pairs, or in whorls near the upper end of the joints, and composed of single joints between bifurcations. Type *C. fruticosus*.
- fruticosus*, Hall, 1865, (*Oldhamia fruticosus*), Can. Org. Rem. Decade 2, p. 50, Trenton Gr.
- CHÆTOCLADUS*, Whitfield, 1894, Bull. Am. Mus. Nat. Hist., vol. 6, p. 356. [Ety. *chaite*, hair; *klados*, stem or branch.] Marine plants with jointed cylindrical stems giving off whorls of hair-like filaments at given distances. Type *C. plumula*, described from the same place from the Trenton Gr.
- CHÆTOMORPHA*, A living genus among the Algae.
- (?) *prima*, Whitfield, 1894, Bull. Am. Mus. Nat. Hist., vol. 6, p. 355, Trenton Gr. It is hardly possible that this species should belong to a living genus.
- CHONDrites*, cuneatus, Whiteaves, 1897, Pal. Foss., vol. 3, p. 140, Low. Sil.
- cupressinus*, Whiteaves, 1896, Can. Rec. Sci., vol. 6, p. 388, Low. Sil.
- gracillimus*, Whiteaves, 1896, Can. Rec. Sci., vol. 6, p. 389, Low. Sil.
- patulus*, Whiteaves, 1896, Can. Rec. Sci., vol. 6, p. 387, Low. Sil.
- DACTYLOPORUS*, Herzer, 1893, Am. Geol., vol. 12, p. 289. [Ety. *dactylos*, finger; *poros*, pore.] A fungus with trunk, pileus and sporiferous arrangement. Type *D. archæus*, described at the same place from the Coal Meas.
- DICTYOLITES*, Penhallow, 1893, Proc. Nat. Mus., vol. 16, p. 113. [Ety. *dictyon*, a net; *lithos*, stone.] Fronds plane, membranaceous, and regularly dichotomous, the ultimate ramuli generally bifid. Midrib none, margins regular. Type *D. fasciolus*, described at the same place with *D. maximus*, Devonian.
- Glyptodendron catonense* seems to have been a mistake, and the name may as well be stricken out; for even if it is a Cephalopod, as has been asserted, it is too poor to retain a specific name.
- HALISERITES*, Sternberg, Fronds plane, membranaceous, costate, and dichotomous throughout; ramuli more or less linear with simple terminations. Sporangia in groups lateral to the midrib. *chondriformis* and *lineatus*, Penhallow, 1893, Proc. Nat. Mus., vol. 16, p. 113, Devonian.
- INCOLARIA*, Herzer, 1893, Am. Geol., vol. 11, p. 365. [Ety. *incola*, an inhabitant of a place.] A fungus in fissures of bark sending forth rounded, overlapping mycelia. Type *I. securiformis*, described at the same place from the Coal Meas.
- LEPIDODENDRON corrugatum* is from the Low. Coal Meas.
- NEUROPTERIS caudata*, White, 1893, Bull. No. 98, U. S. Geo. Sur., p. 87, Coal Meas.
- Jenneyi*, White, 1893, Bull. No. 98, U. S. Geo. Sur., p. 82, Coal Meas.
- PECOPTERIS lesquereuxi*, White, 1893, Bull. No. 98, U. S. Geo. Sur., p. 65, Coal Meas.
- PRIMICORALLINA*, Whitfield, 1894, Bull. Am. Mus. Nat. Hist., vol. 6, p. 357. Articulated marine plants, consisting of elongated cylindrical fronds, composed of a central longitudinal axis, which is jointed and hollow in the fossil condition, and supports whorls of jointed pinnules from each joint; pinnules decomposed. Type *P. trentonensis*, described at the same place from the Trenton Gr.
- PROTOSALVINIA*, Dawson, 1888, Geo. Hist. of Plants, p. 84. [Ety. *protos*, first; *Salvinia*, a genus.] Plants with rhizocarpean affinities of which the detached disks are macrospores and the cellular envelopes sporocarps. Type *P. (Sporangites) huronensis*, and includes *Sporangites bilobatus*.
- clarkei*, Dawson, 1888, Geo. Hist. of plants, p. 84, Devonian.
- PSILOPHYTON grandis*, Penhallow, 1893, Proc. Nat. Mus., vol. 16, p. 113, Devonian.
- RUSOPHYCUS chesterense*, Miller and Gureley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 54, Kaskaskia Gr. See *R. montanense*.

- montanense, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 54, Burlington Gr. A singular mixing up of definitions occurred in printing. The definition of *chesterense* belongs to *montanense*, as may be seen by the illustrations, and that of *montanense* to *chesterense*.
- SPHENOPTERIS *lucoci*, White, 1893, Bull. No. 98, U. S. Geo. Sur., p. 65, Coal Meas.
- Spiraxis*, Newberry, 1884. The name was preoccupied by Adams, in 1850.
- WALCHIA *imbricula*, Dawson, 1888, Geo. Hist. of Plants, p. 138, Permian.
- WINCHELLINA, Herzer, 1893, Am. Geol., vol. 11, p. 285. [Ety. proper name.] Cell bundles encased by a thick periderm with an inner tissue of oblong subquadrate cells with thick walls simulating a transverse section of Carboniferous fossil pine. Type *W. fascina*, described at the same place from the Coal Meas.

ANIMAL KINGDOM.

SUBKINGDOM PROTOZOA.

- Anomalospongia*, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 68. This name was proposed to supersede *Anomaloidea*. The difficulty is that it is founded upon fragments and no generic characters are determined. There is not a shadow of reason to suppose the fragments were ever siliceous. They are crystalline calcite. The material is precisely like that of erinoids and other echinoderms, found in the same layers of rocks. The fossil has no relation to sponges.
- AFROCCELLEA, Rauff, 1895, Palaeontographica, vol. 43, p. 39.
- winnepegensis, Rauff, 1895, Palaeontographica, vol. 43, p. 269, and Pal. Foss. Can., vol. 3, p. 145, Low. Sil.
- CLATHROMICTYON *jewetti*, Girty, 1896, 14th Rep. N. Y. St. Geol., p. 208, Low. Held. Gr.
- CRYPTOZOON *boreale*, Dawson, 1896, Can. Rec. Sci., p. 207, Trenton Gr.
- lachutense*, Dawson. Not defined so as to be recognized.
- occidentale*, Dawson, 1896, Can. Rec. Sci., p. 207, Taconic.
- Cyathophycus siluriana*, James, Syn. for *Trichophycus lanosum*.
- Cylindrocella*, Ulrich. There are no generic characters ascribed to this name, and the proposed species are without specific characters.
- GIRVANELLA *antiqua*, Dawson. Not defined so as to be recognized, but probably a *Strephochetus*.
- GLOBIGERINA, D'Orbigny, 1825, Foraminifères de Vienne. There is not a shadow of probability that the species below referred to this genus belong to it.
- aurita*, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 111, Up. Taconic.
- cambrica*, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 111, Up. Taconic.
- didyma, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 111, Up. Taconic.
- grandis*, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 111, Up. Taconic.
- Heterospongia*, Ulrich. This genus and the species belong to specimens of *Monticulipora* altered by silicification.
- Hyalostelia marcellia*, Clarke. Not defined so as to be recognized.
- LOFTSIA is regarded by Waagen as an Eocene Hydrozoan in the family Sphaeractinidae.
- LYSACTINELLA, Girty, 1896, 14th Rep. N. Y. St. Geol., p. 207. [Ety. *lys*, loose; *aktin*, spicule; *ellus*, diminutive.] Spherical to sub-spherical, sessile, without anchoring spicules. Spicules hexacts, pentacts, tetracts, simple or ornate. Type *L. gebhardi*, described at the same place, from the Low. Held. Gr., and also *L. perelegans*.
- MICROSPONGIA *subrotunda*, James, 1891, Jour. Cin. Soc. Nat. Hist., vol. 14, p. 55, Hud. Riv. Gr.
- Mallerina*, Ulrich, Syn. for *Calcisphaera*.
- greenii*, Ulrich, Syn. for *Calcisphaera robusta*.
- ORBULINA, D'Orbigny, 1845, Foraminifères de Vienne, p. 22. There is not a shadow of probability that the species below belong to this genus.
- ingens*, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 110, Up. Taconic.
- intermedia*, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 110, Up. Taconic.
- ovalis*, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 110, Up. Taconic.
- PALÆOSACOUS, Hinde, 1893, Lond. Geo. Mag., 3d ser., vol. 10, p. 57. [Ety. *palaios*, ancient; *sakkos*, bag.] Cylindrical or

Dawson, 1888, Geo.
38, Permian.
1893, Am. Geol.,
ty. proper name.]
ed by a thick peri-
r tissue of oblong
th thick walls sim-
section of Carbo-
Type *W. fascina*,
me place from the

sack-like sponges with thin walls of
rhombic meshes. Type *P. dawsoni*,
described at the same place from the
Quebec Gr.

Fig. 1296.—*Palenels cavernosa*, sum-
mit and convex side.

RAUFFELLA fucoida, Sanderson, 1896, Bull.
Minn. Acad.
Nat. Sci., vol. 4,
p. 78, St. Peter
Sandstone.

Fig. 1297.—*Receptaculites dixonensis*,
basal view.

Nat. Hist., p. 47, Galena Gr.

RECEPTACULITES dixonensis is de-
scribed and fig-
ured, in Ohio
Geol., vol. 7, p.
419.

dixonensis, Mil-
ler and Gurley,
1896, Bull. No.
11, Ill. St. Mus.

eatoni should be stricken out as it is only
a catalogue name.

monticulatus,
Hall, seems to
be a synonym
for *R. infundi-
bulliformis*.

Streptospongia,
Ulrich, Syn. for
Strephochetus.

SYRINGOTROMA,
barretti, Girty,
1896, 14th
Rep. N. Y. St.
Geol., p. 296,
Low. Held. Gr.

centrotum, Girty,
1896, 14th
Rep. N. Y. St.
Geol., p. 296,
Low. Held. Gr.

consmile, Girty, 1896, 14th Rep. N. Y.
St. Geol., p. 297, Low. Held. Gr.

foveolatum, Girty, 1896, 14th Rep. N. Y.
St. Geol., p. 295, Low. Held. Gr.

microporum, Girty, 1896, 14th Rep. N.
Y. St. Geol., p. 296, Low. Held. Gr.

Trichospongia hystrix, Whiteaves, 1897,
Pal. Foss. Can., vol 3, p. 147, Low. Sil.

Fig. 1298.—*Receptaculites dixonensis*, lateral view.

1894, Trans. N. Y.
p. 111, Up. Taconic.
1894, Trans. N. Y.
p. 111, Up. Taconic.
1894, Trans. N. Y.
p. 111, Up. Taconic.
This genus and
ing to specimens of
ered by silicification.
Clarke. Not defined
ized.

by Waagen as an
ban in the family

, 1896, 14th Rep.
p. 267. [Ety. *luc*, I
le; *ellus*, diminutive.]
pherical, sessile, with-
sacles. Spicules hex-
tracts, simple or or-
bhardi, described at
from the Low. Held.
erelegans.

tunda, James, 1891,
Nat. Hist., vol. 14, p.

yn for *Calcsipherna*.
n. for *Calcsipherna* ro-

y, 1845, Foraminiferes
There is not a shadow
at the species below
nus.

1894, Trans. N. Y.
4, p. 110, Up. Taconic.
ew, 1894, Trans. N. Y.
4, p. 110, Up. Taconic.
894, Trans. N. Y. Acad.
10, Up. Taconic.

1893, Lond. Geo. Mag.,
p. 57. [Ety. *palaios*,
bag.] Cylindrical or

SUBKINGDOM CŒLEENTERATA.

CLASS ANTHOZOA.

THAT the *Monticuliporidae* are true corals, and have their nearest relationship with the *Favositidae*, there has been no doubt among palaeontologists since the great work on the subject by Dr. Waagen, in *Palaeontologia Indica*. The classification of the family with the Bryozoa and the vast synonymy of generic and specific names by Ulrich have made it necessary to draw upon Dr. Waagen for some of the structural features that distinguish bryozoa from corals, that the student may have an opportunity to apply the present state of learning to the investigation of these fossil forms.

The Bryozoon is an animal of rather high organization and very short-lived. The only mode of propagation is by gemmation, and this only in one way, by protruding one of the walls of the mother cell while it is yet in a young state, and afterward partitioning off the protruding part. Each one produces only one or two gems, and these only in the peripheral part of the colony or apex of the branches, and the abode is completed as soon as the gem has attained full size. The fully-developed animals do not produce gems, and the mother animal can

always be distinguished. The gemmation is restricted to the side opposite the aperture of the cell. There is free communication between the young feeding animals and the old ones for the passage of the nutritive fluid, so that more or less pores must remain open, in the walls of the single cells, by which the animals communicate with each other. The animal is only for a short time a feeder, after which it assumes a latent vitality, in which its chief function is restricted to the thickening of the walls of the lodge, and thus strengthening the stony parts of the whole colony. The size of the lodge is fixed, and can not be extended beyond the length of the animal. If tabulæ occur, they are not a sign of progressive growth of the zooid, but represent the retrograde metamorphosis to which the zooids are subject,—a shrinking of the animal occurs, and in this shrinking process the animal deposits from time to time diaphragms behind its ever-contracting body. The circumstance that the greater part of the animals of a colony is in a state of latent life, the functions of taking and digesting food being performed only by a few animals, at the top of the branches, in arborescent colonies, brings with it the other peculiarity, that all the animals of a colony are in intimate connection and communication together. This communication is brought about by numerous capillary tubes and by large openings in the walls of the lodges, by which a free communication of all the animals of a colony is established. The calcareous substance of which the walls of the cells are built is composed of very thin fibers placed vertically to the surfaces of the wall, so that in sections cutting the single cells transversely, a concentric arrangement of the fibers can never be observed. These fibers leave interstices between them at intervals, producing numerous capillary tubes, by which the walls are always pierced in great numbers. In a longitudinal section of an arborescent specimen all the cells may be seen to take their origin at an imaginary axis, and slowly ascend and bend sideways to the surface of the branch. There is no splitting of the cells, and no such distinction between the central and peripheral part of a branch as always occurs, as Nicholson says, in the *Monticuliporidae*. (Struct. and Affin. Montic., page 32.)

The animal of a colony of corals is of low organization and long-lived. Propagation is by fissiparity and by different methods of gemmation. The animals never stop producing gems, but develop them at all times of life and at different levels all over the colony, which produces the small calices intermingled with the larger ones. The gemmation takes place indiscriminately on all sides. The animals are all self-feeding, and perform all their vital functions during their whole lifetime. They constantly undergo a certain process of renovation and each one deposits large masses of sclerenchyma behind itself, and thus slowly ascends within its tube, sometimes chambering off the dead and useless parts of the corallum by diaphragms or tabulæ.

In the *Monticuliporidae*, the gemmation is generally intermural, and the young animal has no part in common with the mother animal, and appears as if it were only filling a void space between several old animals. The gemmation commences with a thickening of the primary plate or primordial wall. After this thickening has been completed, the primary mural plate of the new individual to be developed begins to be formed. In the middle of the originally dark thickening, light-colored wall-substance appears, surrounded by dark lines, indicating the primary wall of the new individual. In this state no hollow for the reception of the animal exists.

opposite the aper-
g feeding animals
more or less pores
animals communi-
er, after which it
the thickening of
the whole colony.
the length of the
wth of the zooid,
s are subject,—a
e animal deposits
The circumstance
of latent life, the
a few animals, at
other peculiarity,
communication to-
lary tubes and by
nication of all the
f which the walls
lly to the surfaces
esely, a concentric
ave interstices be-
y which the walls
of an arborescent
maginary axis, and
There is no split-
nd peripheral part
iporidae. (Struct.

long-lived. Propo-
on. The animals
e and at different
ermingled with the
ides. The animals
eir whole lifetime.
one deposits large
s within its tube,
um by diaphragms

ral, and the young
pears as if it were
mation commences
ter this thickening
ual to be developed
ening, light-colored
the primary wall of
the animal exists.

The third state of development is reached when the light-colored wall-substance, in the middle of the gem, becomes perforated, thus forming the hollow for the reception of the animal. The gemmation is identical with that occurring in the *Favositidae*, except that the thickenings of the primary mural plate inaugurating the formation of gems is more frequent in the *Monticuliporidae* than in the *Favositidae*, which causes the corallites at the surface to be of more unequal size. The primary wall appears, in most cases, as a dark line hemmed in on both sides by light-colored masses of sclerenchyma composed of oblique, symmetrically-arranged fibers without capillary tubes or any other openings through the walls to connect the animals with each other. The secondary thickenings are always composed of successive reversed conical layers of sclerenchyma, which appear, in tangential sections, as concentric rings surrounding the visceral cavities of the single corallites. When the primary plate is not preserved, these rings may surround the visceral cavity, and without understanding the state of preservation they have been called "marginal rings." The smaller corallites are young ones or gems, which have not attained their full development, and have been produced by intermural gemmation. As the animals inhabiting the corals never stop producing gems, it is only natural that the gems produced just before the whole colony ceased augmenting, in thickness or length, appear in very different states of development on the surface of the corallum, some as small pores, but otherwise similar to the mother animals, and some even only as thickenings of the primordial wall, and projecting as little spines above the upper termination of that wall. Thus, in all stages of growth, large pores, small pores, and little spines, called "spiniform corallites" by Nicholson, and as a synonym "acanthopores," are observable on the surface of the *Monticuliporidae*. The tabulae are usually closer in the early growth of the gems than in the more adult state, and the better fed and more vigorous the animal, the closer will be the tabulae. The difference, therefore, in the number of the tabulae in a given corallite is not of specific importance. As the so-called "spiniform corallites" are nothing but newly-developed gems, they are never of specific importance. Sometimes the primary wall is not well preserved, and is represented by black spots, which have also been called "spiniform corallites," but a paleontologist ought to be able to understand the state of the preservation of his specimen. Rudimentary septa have been observed in some species of *Monticulipora* similar to those existing in the *Chetidae*. The appearance as of an operculum, which is not uncommon in the corallites, is caused by a secondary infiltration, into the cells, of carbonate of lime, which assumes a spheroidal shape, and possesses no organic structure of any kind. No such organ as an operculum has ever been found in the corallites.

Fissiparity is a mode of growth that does not occur in the Bryozoa, but is not of rare occurrence in the *Monticuliporidae*. It can be very easily distinguished from gemmation in longitudinal sections of the corallites, by the circumstance, that the internal space of the new animal forms part of the internal space of the mother animal. The longitudinal wall begins at one of the tabulae, and the animal is literally split in two by a wall which stretches from one side of its body cavity to the other.

The family *Monticuliporidae* must be removed from the sub-class *Alcyonaria* to the sub-class *Zoantharia*, and will include *Dekayia* (if it is a genus), *Dianulites*, *Monotrypella*, *Monticulipora*, *Peronopora*, possibly *Stenopora*, and some of the genera, if

they are valid genera, proposed by Ulrich, among the Bryozoa; viz., *Atactoporella*, *Batostoma*, *Batostomella*, *Homotrypa*, and *Homotrypella*. The *Favositidae* will include *Alveolites*, *Calapoecia*, *Chonostegites*, *Cladopora*, *Cenites*, *Dendropora*, *Favosites*, *Leptopora*, *Lunatipora*, *Michelinia*, *Nyctopora*, *Pachypora*, *Pleurodietyum*, *Romingeria*, *Sphaerolites* (?), *Striatopora*, *Syringolites*, *Trachypora*, and *Vernipora*.

The family *Fistuliporidae* must be removed from the Bryozoa to the sub-class *Aleyonaria*, and will include *Callopora*, *Callotrypa*, *Cœlocaulis*, *Evactinopora*, *Favicella*, *Fistulipora*, *Prasopora*, and *Strotopora*. The important distinction between the two families, *Monticuliporidae* and *Fistuliporidae*, is in the mode of gemmation, being intermural in the former, and coenenchymal in the latter. There is no coenenchyma in the *Monticuliporidae*. The coenenchyma consists of parallel tubes, parallel to the polypites, and each coenenchymal tube possesses numerous tabulae. It has been supposed that the *Fistuliporidae* are dimorphic; that is, occupied by two different sets of animals, of which one, the *siphonozooids*, dwelt in the coenenchymal tubes, whilst the other, the *autozooids*, occupied the larger calices. In the coenenchymal gemmation a number of these coenenchymal tubes apparently unite to form together a new autozoid, so that several reduced individuals become blended together into a single perfect one. This coenenchymal gemmation occurs in the *Helioporidae*, and shows the coral nature of the fossil.

ACROPHYLLUM. The central boss is formed solely by the elevation of the successive tabulae; and no vertical plates take part in its formation as is the case in *Clisiophyllum*.

pluriradiale, Nicholson, instead of *Clisiophyllum pluriradiale*.

AMPLEXUS. The corallum is always simple, subcylindrical or cylindro-conical and more or less twisted. It is covered with an epitheca having encircling lines of growth, and accretion ridges or constrictions are more or less developed. Septa short, never reach the center. Tabulae extend completely across the visceral chamber and occupy the central area. The fossula consists of a lateral depression of the tabulae. Calice circular, moderately deep, with a thin margin.

annulatus, Whitfield. This name was preoccupied by Verneuil and Haime in Bull. Soc. Geol. de France, in 1850. See A. whitfieldi.

geniculatus, Worthen, 1890, Geo. Sur. Ill., vol. 8, p. 82, Kaskaskia Gr.

rockfordensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 53, Kinderhook Gr.

whitfieldi, n. sp. This name is proposed for *A. annulatus*, Whitfield, 1878, Ann. Rep. Geo. Sur. Wis., p. 80, and Geo. Wis., vol. 4, p. 314, Niagara Gr., *annulatus* was preoccupied in 1850, by Verneuil and Haime.

AULOPORA *trentonensis*, Winchell and Schuchert, 1893, Geo. Sur. Minn., vol. 3, p. 95, Trenton Gr.

AXOPHYLLUM. Simple, turbinute, epitheca complete. Columella large, composed of numerous vertical, spirally twisted lamellae; in longitudinal section, it appears as a cylindrical cellular mass of large size, and it pierces a central area occupied by strong, distant tabulae, surrounded by an accessory wall. The space between this inner wall and the outer wall is occupied by dissepiments, making an exterior zone of large vesicles. Septa extend to the center of the visceral chamber.

Azygograptus walcotti, Lapworth, 1896, Jour. Geol., vol. 4, p. 69. Definition worthless.

BRYOGRAPTUS, Lapworth, 1880, Ann. and Mag. Nat. Hist., vol. 5, p. 164. [Ety. *bryon*, moss; *grapho*, I write.] Type B. kjerulfi.

lentus, Matthew, 1894, Trans. N. Y. Acad. Sci. vol. 14, p. 270, St. John Gr. *multiramosus*, Gurley, 1896, Jour. Geol., vol. 4, p. 64. Definition worthless.

Bythograptus, Hall. This is an alga (?). See Bull. Am. Mus. Nat. Hist., vol. 6, p. 357.

CALYPTOGRAPTUS *miconemmatodes* and *radiatus*, Spencer, 1884, Bull. Mus. Univ. Mo., p. 29, Niagara Gr.

CAMPOPHYLLUM. The septa are not as well developed as they are in *Cynthophyllum*, and the tabulae are exposed over a larger central area. The calice is deep.

kansasense, Miller and Gurley, 1893, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 67, Up. Coal Meas.

iz., *Atactoporella*,
tidæ will include

Favosites, *Lepto-*
Romingeria, *Sphæ-*

a to the sub-class

Stenopora, *Favicella*,

between the two

remmation, being

no coenenchyma

tubes, parallel to

ulæ. It has been

by two different

enchymal tubes,

the coenenchymal

to form together

ded together into

e *Helioporidæ*, and

turbinate, epitheca
lla large, composed
al, spirally twisted
dinal section, it ap-
pal cellular mass of
perces a central area
g, distant tabulæ,
cessory wall. The
inner wall and the
ed by dissepiments,
zone of large vesid-
to the center of
er.

Lapworth, 1896,
p. 69. Definition

th, 1880, Ann. and
ol. 5, p. 164. [Ety.
p, I write.] Type B.

894, Trans. N. Y.
p. 270, St. John Gr.
1896, Jour. Geol.,
ition worthless.

This is an alga (?).
Nat. Hist., vol. 6,

onematodes and ru-
34, Bull. Mus. Univ.
Gr.

septa are not as well
are in Cyathophyl-
are exposed over
rea. The calice is

d Gurley, 1893, Bull.
s. Nat. Hist., p. 67,

texanum, Shumard. Not defined so as
to be recognized.

Caryocaris oblongus and *curvilinearis*, Gurley,
1896, Jour. Geol., vol. 4, p. 87, Calcif-
erous Gr. Named as shrimps, but de-
scribed as Graptolites, and such no-
menclature should be thrown out. The
Phyllopod genus *Caryocaris* is fully de-
scribed in Monogr. Brit. Palæoz., Phyl-
locarida Pal. Soc., 1892, p. 89.

CERATOSTIGMA, Clarke, 1894, 13th Rep. St.
Geol. N. Y., p. 178. Supposed to be-
long to the Aleyonaria, though of
doubtful nature. Type *C. papillata*,
described, at the same place, from the
Hain. Gr.

CHETES carbonarius, Worthen, refer to
Stenopora carbonaria.

clavacoides. See *Leptotrypa clava-*
coides.

ortoni, refer to *Monticulipora ortoni*.

perantiquus, Whiteaves, 1897, Pal. Foss.,
vol. 3, p. 238, Low. Sil.

CHONOSTEGITES clappi, Edwards and Haine,
1857, Pol. Foss. d. Terr. Pal., p. 299,
Up. Held. Gr.

CLADONCHUS, McCoy, 1847, Ann. Nat.
Hist., vol. 20, p. 14-227. [Ety. *klados*,
a twig; *chonos*, a funnel.] Corallum
compound, cylindrical, walls of vari-
able stoutness, always developed in the
form of upright branching and reticu-
late colonies rising from one corallite,
and bifurcating, at various angles, in
the different species; and there is free
communication between the parent
and the young corallites, by a prolon-
gation of the visceral chamber of the
former, and extending throughout the
latter; and all are kept in union by
the epitheca. Epitheca smooth, or
with delicate annulations of growth.
The space between it and the en-
dotheca is occupied by more or less
dense sclerenchyma. It is distin-
guished from *Aulopora* by the erect
habit of growth, regular, angular mode
of branching, slender, equal stem-like
tubes, and abruptly dilated terminal
cups bent in nearly opposite directions.
It frequently encircles crinoid col-
umns, and sends off the erect corallites
at right angles to the column. Type
C. tenuicollis.

CLADOPORA clarkei, Girty, 1896, 14th Rep.
N. Y. St. Geol., p. 306, Low. Held. Gr.
helli, Girty, 1896, 14th Rep. N. Y. St.
Geol., p. 306, Low. Held. Gr.

CLIMACOGAPTUS, *antiquus*, *kamptotheca*, *lat-*
icaulis, and *oligotheca*, Gurley, 1896,
Jour. Geol., vol. 4, p. 74. Definitions
worthless.

phylliporus, Gurley, 1896, Jour. Geol.,
vol. 4, p. 77. Group not given. Poor
definition.

CLISTOPHYLLUM. Epitheca complete, and
marked with constrictions and accre-
tion ridges. Surface of the central
boss marked with spirally bent or

straight lamellæ which are attached to
the primary septa by delicate dissep-
iments and pass upward to a median
columnar test on the crown of the
boss. Central area formed by vertical,
spirally-twisted or straight lamellæ
and by vesicular tabulæ directed up-
ward and inward. Intermediate area
formed by an outward extension of
the tabulæ in large, nearly horizontal
vesicles. External area formed by
minute vesicular tissue, vesicles ar-
ranged in oblique rows directed out-
ward and upward.

pluriradiale refer to *Aerophyllum pluri-*
radiale.

CLONOGAPTUS proximatus, Matthew, 1895,
Trans. N. Y. Acad. Sci., vol. 14, p. 265,
St. John Gr.

COENITES. The type is *C. juniperinus*.

Constellaria varia, Ulrich, refer to *Stelli-*
pora varia.

CYATHAXONIA has primary and secondary
septa, no interocular tabulæ in the in-
termediate area, and the spaces are open
from the superior to the inferior part
of the corallum. There are a few intersep-
tal dissepiments near the periphery.

CYATHOPHYLLUM. The tabulæ never ex-
tend completely across the visceral
chamber as they do in *Zaphrentis*, and
the septa are symmetrically developed,
regularly arranged, and are not inter-
rupted by any well-defined fossula.

juvenile, refer to *Heliophyllum* juvenile.

DAWSONIA monodon and tridens, Gurley,
1896, Jour. Geol., vol. 4, p. 88, Calcif-
erous Gr.

Dekayia, I have long regarded as founded
upon weathered and altered fragments
of *Monticulipora*, but out of deference
to the opinions of others have cata-
logued the species; but I have been
unable to distinguish any specific dif-
ference between *D. aspera* and *D. mul-*
tispinosa, or between *D. pelliculata*, *D.*
appressa, and *D. paupera*, and no two
specimens are exactly alike. Many
more names might be applied to these
weathered and altered corals with
equally as much propriety. It is not a
valid genus. Any species of *Monticu-*
lipora may present all the characters
attributed to *Dekayia*. I have two or
three species of *Dekayia* and *Monticu-*
lipora on the same specimen.

Dekayella, Ulrich, Syn. for *Monticulipora*.
obscura, Ulrich, Syn. for *Monticulipora*
ulrichi.

peruviana, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 270, Trenton Gr., refer to
Monticulipora.

robusta, Foord, refer to *Monticulipora*
robusta.

DENDROGAPTUS arundinaceus, Hall, 1847,
(*Graptolithus arundinaceus*), Pal. N. Y.,
vol. 1, pl. 74, Utica Slate.
unilateralis, Gurley, 1896, Jour. Geol.,
vol. 4, p. 84. Definition worthless.

- Desmograptus devonicus* and *macrodictyum*, Gurley, 1896, Jour. Geol., vol. 4, p. 83. Not properly defined.
- DIANULITES*, Eichwald, 1829, Zool., special vol. 1, p. 180, and fully defined by Dybowski, 1877, Chætid. d. ostbalt. Silurform., p. 14. Type *D. petropoli-tana*, which was subsequently made the type of *Diplotrypa*.
- discoides*, Ulrich, 1893 (Mesotrypa discoides), Geo. Sur. Minn., vol. 3, p. 260, Galena Gr.
- dubia*, Ulrich, 1890 (Diplotrypa (?) dubia), Geo. Sur. Ill., vol. 8, p. 459, Hud. Riv. Gr.
- limitaris*, Ulrich, 1893 (Diplotrypa limitaris), Geo. Sur. Minn., vol. 3, p. 286, Galena Gr.
- neglecta*, Ulrich, 1893 (Diplotrypa neglecta), Geo. Sur. Minn., vol. 3, p. 287, Galena Gr.
- patella*, Ulrich, 1890 (Diplotrypa patella), Geo. Sur. Ill., vol. 8, p. 458, Hud. Riv. Gr.
- quebecensis*, Ami., 1892 (Diplotrypa quebecensis), Can. Record Sci., p. 101, Quebec Gr.
- rotunda*, Ulrich, 1893 (Mesotrypa rotunda), Geo. Sur. Minn., vol. 3, p. 262, Galena Gr.
- selkirkensis*, Whiteaves, 1897 (Mesotrypa selkirkensis), Pal. Foss., vol. 3, p. 162, Low. Sil.
- Dicellograptus gurleyi* and *polytheatus*, Gurley, 1896, Jour. Geol. vol. 4, p. 70. Definitions worthless.
- DICRANOGRAPTUS* arkansensis and parvungulus, Gurley, 1892, Geo. Rep. Ark., p. 416. Group not given.
- diapason*, Gurley, 1896, Jour. Geol., vol. 4, p. 73. Definition worthless.
- DICTYONEMA actinotum*, Gurley, 1896, Jour. Geol., vol. 4, p. 82. Definition worthless.
- blairi*, Gurley, 1896, Jour. Geol., vol. 4, p. 82, Chouteau limestone. Although the definition is worthless, I know the form, and it has not the slightest resemblance to Dictyonema or any other Graptolite. It consists of a granular carbonaceous mineral that has replaced more or less a Bryozoon.
- crassum*, Girty, 1896, 14th Rep. N. Y. St. Geol., p. 288, Low. Held. Gr.
- Dichograptus remotus*, Gurley, 1896, Jour. Geol., vol. 4, p. 64. Definition worthless.
- DIDYMOGRAPTUS* bipunctatus, Gurley, 1896, Jour. Geol., vol. 4, p. 65, Calciferous Gr.
- convexus*, Gurley, 1896, Jour. Geol., vol. 4, p. 67. Group not given.
- geminus*, Hisinger. Probably not an American species.
- perflexus*, Gurley, 1896, Jour. Geol., vol. 4, p. 66. Definition worthless.
- sagitticaulis*, Gurley. Not defined.
- DIPHYPHYLLUM* stokesi, instead of Lithostrotion stokesi, and refer to Lower Silurian.
- DIPLOGRAPTUS*, *folium*, Hisinger. Probably not an American species.
- ruedemanni*, Gurley, 1896, Jour. Geol., vol. 4, p. 307, Utica Slate.
- stenosus*, Gurley, 1896, Jour. Geol., vol. 4, p. 78. Definition worthless.
- trifidus*, Gurley, 1890, Geo. Sur. Ark., vol. 3, p. 417, Low. Sil.
- Diplotrypa* is a synonym for Dianulites, to which the species must be referred.
- DUNCANELLA* rudis, Girty, 1896, 14th Rep. N. Y. St. Geol., p. 299, Low. Held. Gr.
- Emmonsia*, seems to be a Syn. for Favosites.
- ERIDOPHYLLUM* is retained by some as distinct from Diphyphyllum on the ground that the corallites are united at intervals by numerous, lateral outgrowths of the epitheca.
- FAVOSITES*, *cervicornis*, *cristatus*, and *reticulatus*, are not American.
- conradi*, Girty, 1896, 14th Rep. N. Y. St. Geol., p. 304, Low. Held. Gr.
- gothlandicus* and *troosti* are Upper Silurian, not Devonian species.
- GRAPTOLITHUS* acanthonotus, Gurley, 1896 (Tetragraptus acanthonotus), Jour. Geol., vol. 4, p. 65, Calciferous Gr.
- arundinaceus*. See Dendrograptus arundinaceus.
- HADROPHYLLUM* tenneseense, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 87, Keokuk Gr.
- Fig. 1299. — Hadrophyllum tenneseense, lateral and summit views.
- HELIOPHYLLUM* juvene, Rominger, instead of Cythophyllum juvene.
- INOCAULIS* canadensis, Whiteaves, 1897, Pal. Foss. Can., vol. 3, p. 149, Low. Sil.
- Leptograptus macrotheca*, Gurley, 1896, Jour. Geol., vol. 4, p. 69. Definition worthless.
- Lichenaria*, Winchell and Schuchert, 1893, Geo. Sur. Minn., vol. 3, p. 83. [Ety. as given by the authors, "Lichen, tree-moss; and aria, the latter portion of Columnaria, its most likely relative," which makes an absurd compound.] It may be a synonym for Columnaria, but it is too poorly defined to be recognized.
- typa*, Winchell and Schuchert. Too poorly defined to be recognized.
- LITHOSTROTION*, published 1869, instead of 1869. Corallites surrounded with an epitheca. Primary septa extend from the outer wall, nearly or quite to the columella. Columella compact, styliform. Central area formed by irregular, somewhat elevated tabulae. Interseptal loculi filled with dissepiments, producing in longitudinal sections a series of small lenticular cells, arranged in layers which are directed upward and outward. Growth by calicular gemination or lateral budding.

Linsinger. Probable species.

1896, Jour. Geol., late.

Jour. Geol., vol. 4, ethless.

Geo. Sur. Ark., 11.

for Dianulites, to be referred.

, 1896, 14th Rep.

0. Low. Held. Gr. Syn. for Favosites.

d by some as disphyllum on the

allites are united erous, lateral out-

een.

status, and reticu-

can.

14th Rep. N. Y. St. field. Gr.

ti are Upper Silu-

species.

atus, Gurley, 1896

thoniferous), Jour.

califerous Gr.

ndrograptus arnu-

ense, Miller and

Gurley, 1895,

Bull. No. 7,

Ill. St. Mus.

Nat. Hist., p.

87, Keokuk

Gr.

HELIOPHYLLUM

Juvene, Ro-

tyathophyllum ju-

Whiteaves, 1897, Pal.

149, Low. Sil.

t, Gurley, 1896,

p. 69. Definition

d Schuchert, 1893,

pl. 3, p. 83. [Ety.

ors, "Leichen, tree-

e latter portion of

it likely relative,"

bsurd compound.]

n for *Columaria*,

defined to be rec-

Schuchert. Too

recognized.

d 1869, instead of

rounded with an

septa extend from

ly or quite to the

lla compact, styli-

formed by irregu-

lated tabulae. Inter-

with dissepiments.

tudinal sections a

ular cells, arranged

e directed upward

owth by calicular

al budding.

junceum, may be taken from the list, as it is not American.

stokesi, refer to *Diphyphyllum stokesi* and to Lower Silurian.

LOPHOPHYLLUM profundum, Worthen, 1890, Geo. Sur. Ill., vol. 8, p. 79, Coal Meas.

MICHELINIA branneri, Miller and Gurley, 1893, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 68, Coal Meas.

MONOGRAPTUS beecheri, Girty, 1896, 14th Rep. N. Y. St. Geol., p. 288, Low. Held. Gr.

MONOTRYPA cumulata, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 307, Trenton Gr.

intabulata, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 305, Galena Gr.

magna, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 304, Trenton Gr.

nodosa, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 306, Hud. Riv. Gr.

rectinuralis, Ulrich, Syn. for *Monticulipora undulata*.

MONOTRYPELLA appressa, Ulrich, 1890, Geo. Sur. Ill., vol. 8, p. 453, Ham. Gr.

crassimuralis, Ulrich, 1890, Geo. Sur. Ill., vol. 8, p. 452, Hud. Riv. Gr.

MONTICULIPORA arborea, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 220, Galena Gr.

cannonensis, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 221, Galena Gr.

consimilis, Ulrich, Syn. for *M. cincinnatensis*.

eccentrica, James, 1894, Jour. Cin. Soc. Nat. Hist., vol. 16, p. 185, Hud. Riv. Gr.

falsi, James, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 138, Hud. Riv. Gr.

incompta, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 219, Trenton Gr.

inflecta, Ulrich, 1890, (*Heterotrypa inflecta*.) Geo. Sur. Ill., vol. 8, p. 414, Hud. Riv. Gr.

lamellosa, Ulrich, 1890, Geo. Sur. Ill., vol. 8, p. 408, Hud. Riv. Gr.

ohioensis, James, 1884, Jour. Cin. Soc. Nat. Hist., vol. 7, p. 137, Hud. Riv. Gr.

prolifera, Ulrich, 1890, (*Heterotrypa prolifera*.) Geo. Sur. Ill., vol. 8, p. 413, Hud. Riv. Gr.

singularis, Ulrich, 1890, (*Heterotrypa singularis*.) Geo. Sur. Ill., vol. 8, p. 415, Hud. Riv. Gr.

winchelli, Ulrich, 1890, Geo. Sur. Ill., vol. 8, p. 408, Ham. Gr.

Nebulipora, McCoy, Syn. for *Monticulipora* and *N. papallata*, is not American.

OLDHAMIA fruticosa, refer to *Callithamnopsis fruticosa*.

PERONOPORA, Nicholson, 1881, Struct. and Affin. Montic., p. 215. [Ety. *perone*, a buckle or clasp; *poros*, pore.] Corallum laminar, forming undulating fronds; corallites large and small, the latter interspersed among the former, and sometimes aggregated into clusters. Walls of the corallites thickened, and seemingly fused together. Type *Monticulipora frondosa*, and including *M. decipiens*, if it is a good species. I

think this subgenus of Nicholson should be raised to the rank of a genus, but I am not sure that any other species than the two mentioned belong to it.

PHYCNOGRAPTUS, Gurley, 1890, Jour. Geol., vol. 4, p. 89. [Ety. *phukos*, seaweed; *grapho*, I write.] Long, slender, flexuous, segmented stems, with a single central pit in each segment. Type *P. brachymera*, described at the same place. Group not given.

PRASOPORA insularis, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 251, Galena Gr.

lenticularis, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 253, Trenton Gr.

PROTAREA magna, Whiteaves, 1897, Pal. Foss., vol. 3, p. 155, Low. Sil.

verneuili was defined in less than four lines, and has never been illustrated. It was found in the range of *Protarea retusta*, about forty-five miles from Cincinnati, and, no doubt, is a synonym for it. It may be stricken from the list of fossils.

STELLIPORA parva, Ulrich, 1890, (*Constellaria parva*.) Geo. Sur. Ill., vol. 8, p. 424, Hud. Riv. Gr.

varia, Ulrich, 1893, (*Constellaria varia*.) Geo. Sur. Minn., vol. 3, p. 311, Galena Gr.

STENOPORA. Wnagen says there are no mural pores, and that Nicholson and Etheridge mistook accidental grooves for them.

carbonaria, Worthen, instead of *Chetetes carbonaria*.

Stephanograptus crassicaulis and *exilis*, Gurley, Jour. Geol., vol. 4, p. 68. Definitions worthless.

STREPTELASMA breve, Ulrich, 1893, Geo. Sur. Minn., vol. 2, p. 92, Trenton Gr.

parasiticum, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 89, Trenton Gr.

robustum, Whiteaves, 1896, Can. Rec. Sci., vol. 6, p. 351, and Pal. Foss., vol. 3, p. 153, Low. Sil.

STYLASTREA anna is described and figured in Ohio Geol., vol. 7, p. 420.

SYRINGOPORA occidentalis, Meek, 1877, Expl. 40th Parallel, vol. 4, p. 50, Carboniferous.

pacallata, Etheridge, is not American, beside the name was preoccupied for a Carboniferous species.

TETRADUM pacchi var. canadense, Foord. Syn. for *Stromatopora completa*.

THAMNOGRAPTUS allisii, Whiteaves, 1897, Pal. Foss. Can., vol. 3, p. 148, Low. Sil.

ZAPHRENTIS. The word "septa" is preferred to "lamellae" in the definition. The corallum is usually more or less curved, and coated with a thin epithec.

The fossula varies from the concave or ventral side to the convex or dorsal

Fig. 1270.—*Zaphrentis chelycutus*. Front view, showing calyx, and side view.

Fig. 1271.—*Zaphrentis chontenchensis*. Front view, showing calyx, and side view.

side; it is large and deep, and is formed by the coalescence of a greater or less number of the septa which unite near the center of the visceral chamber and form the walls of the fossette. The septa are usually very thick at the margin of the corallum, and there is

generally one secondary septum between each pair of primary septa.

Fig. 1272.—*Zaphrentis exilum*. Front view, showing calyx, and side view, magnified two diameters.

Fig. 1273.—*Zaphrentis tantilla*. Front view, showing calyx, and side view, magnified two diameters.

calcariformis, Hall, is from the Warsaw Gr. instead of Up. Held. Gr. *denticulatus*, Goldfuss, 1826, (*Anthophyllum denticulatum*,) Petref. Germ. t. 1, p. 46, Ningura Gr.

Fig. 1274.—*Zaphrentis tenella*. Front view, showing radiating septa, and side view.

knappi, Hall. Syn. for *Z. deformis*. Exactly the same description.

SUBKINGDOM ECHINODERMATA.

ACACROCINUS, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 515. [Ety. *akakos*, simple; *krinon*, lily.] Basals three; primary radials three by five; arms ten; azygous and regular interradsial areas connect with the vault. Type *A. elrodi*.

elrodi, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 515, Niagara Gr.

ACROCINUS *amphora*, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 808, St. Louis Gr.

ACTINOCRINIDÆ. As to this family see Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 73.

ACTINOCRINUS, *adolescens*, Wachsmuth and Springer, 1897, (*Teleocrinus adolescens*,) N. Am. Crin. Cam., vol. 2, p. 635, Burlington Gr.

albersi, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 46, Chouteau Gr.

amplus, instead of *Saccocrinus amplus*.

arrosus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 15, Burlington Gr.

asterius, McChesney, 1860, Desc. New

Pal. Foss., p. 13, Burlington Gr., is not a syn. for *B. verrucosus*.

Fig. 1275.—*Actinocrinus bischoffi*, azygous view.

Fig. 1273.
Actinoecrinus *zaphrentis* n. sp.
 Front view, showing
 only x, and side
 view, magnified
 two diameters.

s from the War-
 saw Gr. instead
 of Up. Held. Gr.
denticulatus, Gold-
 fus, 1826, (*Antho-
 phyllum denticu-
 latum*.) Petref.
 Germ. t. 1, p. 46,
 Niagara Gr.
knappi, Hall. Syn.
 for *Z. deformis*.
 Exactly the
 same descrip-
 tion.

MATA.

Burlington Gr., is not
 a new species.

Actinoecrinus *bischoffi*,
 new.

augustatus, Miller and Gurley, 1894,
 Bull. No. 3, Ill. St. Mus. Nat. Hist.,
 p. 11, Keokuk
 Gr.

Fig. 1276.—*Actinoecrinus*
botruosus, azygous
 side.

bischoffi, Miller
 and Gurley,
 1896, Bull. No.
 10, Ill. St. Mus.
 Nat. Hist., p.
 8, Burlington
 Gr.

botruosus, Miller
 and Gurley,
 1895, Bull. No.
 6, Ill. St. Mus.
 Nat. Hist., p.
 22, Keokuk Gr.

calyculoides, see
Batoecrinus *ca-*

lyculoides.
concinus, refer to *Shumardocrinus* *con-*

cinnus.
corniculum, refer to *Doryecrinus* *cornicu-*

lum.
daphne is from the Keokuk Gr.

denticulatus, Wachsmuth and Springer,
 1897, (*Cactoc-*

erinus denticu-
latus.) N. Am.
 Crin. Cam., vol. 2,
 p. 606, Bur-

lington Gr.
erraticus, Mil-

ler and Gur-
ley, 1894, Bull.
 No. 3, Ill. St.
 Mus. Nat. Hist.,
 p. 14, Burlin-

gton Gr.
excerptus, Hall,
 1861, is also in

Bost. Jour. Nat. Hist., p. 276.

extensus, Wachsmuth and Springer, 1897,
 (*Cactocrinus* *extensus*.) N. Am. Crin.

Cam., vol. 2, p. 616, Burlington Gr.

foveatus, Miller and Gurley, 1895, Bull.
 No. 7, Ill. St. Mus. Nat. Hist., p. 49,

Burlington Gr.

Fig. 1278.—*Actinoecrinus* *fossatus*, azygous
 and side views.

gibsoni, Miller and Gurley, 1894, Bull.
 No. 3, Ill. St. Mus. Nat. Hist., p. 10,

Keokuk Gr.

glans. See Mem. Am. Mus. Nat. Hist.,
 vol. 1, p. 10, and illustration.

gracilis, Wachsmuth and Springer, 1897,
 N. Am. Crin. Cam., vol. 2, p. 572, Bur-

lington Gr.

griffithi, Wachsmuth and Springer. Syn.
 for *Steganoecrinus* *albersi*.

inflatus, refer to *Amphoecrinus* *inflatus*.

jessleae, Miller and Gurley, 1896, Bull.
 No. 8, Ill. St. Mus. Nat. Hist., p. 32,

Burlington Gr.

limabrachiatus is also in Mem. Am.
 Mus. Nat. Hist., vol. 1, p. 5, and illus-

trated.

lobatus, in Geo. Sur. Ill., vol. 8, p. 97, is
A. augustatus.

mutata var. *attenuatus*. See *Batoecrinus*
attenuatus.

monticuliferus, Miller and Gurley, 1894,
 Bull. No. 5, Ill. St. Mus. Nat. Hist., p.

46, Keokuk Gr.

multiramosus, Wachsmuth and Springer.
 Syn. for *Actinoecrinus* *gilsoni*.

obesus, Keyes, 1895, Mo. Geo. Sur., vol.
 4, p. 187, Burlington Gr.

opuseculum is also in Mem. Am. Mus.
 Nat. Hist., vol. 1, p. 9, and illustrated.

pettisensis, Miller and Gurley, 1896, Bull.

Fig. 1279.—Actinoecrinus *pettisensis*, azygous
 and opposite views.

No. 10, Ill. St. Mus. Nat. Hist., p. 6,
Burlington Gr.

plagiosus, Miller and Gurley, 1894, Bull.
 No. 3, Ill. St. Mus. Nat. Hist., p. 18,

Burlington Gr.

pollubrum, Miller and Gurley, 1896, Bull.
 No. 9, Ill. St. Mus. Nat. Hist., p. 26,

Burlington Gr.

proboscoidialis, Hall, refer to *Batoecrinus*
proboscoidialis.

quaternarius is also in Mem. Am.
 Mus. Nat. Hist., vol. 1, p. 7, and illus-

trated.

ramulosus, refer to *Batoecrinus* *ramu-*
losus.

sampsoni, Miller and Gurley, 1896, Bull.
 No. 10, Ill. St. Mus. Nat. Hist., p. 5,

Burlington Gr.

securis is figured in Geo. Sur. Ill., vol 5,
 pl. 9, see also p. 328.

senectus, Miller and Gurley, 1897, Bull.
 No. 12, Ill. St. Mus. Nat. Hist., p. 36,

Chouteau Gr.

sobrinus, Miller and Gurley, 1896, Bull.
 No. 10, Ill. St. Mus. Nat. Hist., p. 10,

Burlington Gr.

spectabilis, Miller and Gurley, 1896, Bull.
 No. 10, Ill. St. Mus. Nat. Hist., p. 9,

Burlington Gr.

subpulchellus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 13, Burlington Gr.

Fig. 1280.—*Actinoecrinus subpulchellus*, azygous and opposite views.

subscitulus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 12, Burlington Gr.

thetis, is figured in Mem. Am. Mus. Nat. Hist., vol. 1, p. 6.

tuberculosis, Wachsmuth and Springer, N. Am. Crin. Cam., vol. 2, p. 573, Burlington Gr.

validus. See *Steganoecrinus validus*.

AESIOCRINUS angulatus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 59, Up. Coal Mens.

AESIOCYSTITES, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 13. [Ety. *aistos*, auspicious; *kustis*, a bladder.] Body highly convex or hemispherical; free, not parasitic. Plates and arms curve over the margin. Interbranchial plates, nonimbricating. Arms five, convex, radiate from the center, and composed externally of a double series of alternating and interlocking plates, that cover a deep, angular furrow. Ovarian pyramid in the larger interbranchial area near the margin. Type, *A. priscus*.

Fig. 1281.—*Aesloecystites priscus*. Two summit and one lateral view.

priscus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 14, Trenton Gr.

AGARICOERINUS. The type of this genus is *A. tuberosus*, and for remarks see Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 9.

adamsensis, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 24, Burlington Gr.

arcula, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 30, Keokuk Gr.

Fig. 1282.—*Agaricoerinus arcula*, summit view.

conicus, Wachsmuth and Springer, 1897, N. Am. Crinoiden Camerata, vol. 2, p. 501, Keokuk Gr.

eris is from the Keokuk Gr.

excavatus is also in Mem. Am. Mus. Nat. Hist., vol. 1, p. 26, and illustrated.

Fig. 1283.—*Agaricoerinus bluhli*, basal and summit views.

helice is from the Keokuk Gr.

hodgsoni, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 23, Burlington Gr.

illinoisensis, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 25, Burlington Gr.

iowensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 5, Keokuk Gr.

keokukensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 7, Keokuk Gr.

Fig. 1284.—*Agaricoerinus bluhli*, lateral view.

ornotrema is also in Mem. Am. Mus. Nat. Hist., vol. 1, p. 24, and illustrated; but as the word was not properly made, it was changed to *bellatrema*.

pentagonus is also in Mem. Am. Mus. Nat. Hist., vol. 1, p. 25, and illustrated. *profundus*, Miller and Gurley, 1895, Bull.

Fig. 1285.—*Agaricoerinus profundus*, basal view.

No. 6, Ill. St. Mus. Nat. Hist., p. 26, Keokuk Gr.

tugurium, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 28, Keokuk Gr.

ACASSIZOERINUS ovalis, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 36, Kaskaskia Gr.

nd Springer, 1897.
amerata, vol. 2, p.

ck Gr.
m. Am. Mus. Nat.
nd illustrated.

s blairi, basal and
ews.

uk Gr.
Gurley, 1896, Bull.
Nat. Hist., p. 23,

and Gurley, 1896,
Mus. Nat. Hist., p.

(Gurley, 1897, Bull.
Nat. Hist., p. 5,

. 1284.—*Agaricoerinus*
blairi, internal view.

24, and illustrated;
not properly made,
ellatrema.

Mem. Am. Mus.
25, and illustrated.
Gurley, 1895, Bull.

us profundus,
w.

Nat. Hist., p. 26,

Gurley, 1895, Bull.
Nat. Hist., p. 28,

Miller and Gurley,
Ill. St. Mus. Nat.
ia Gr.

Fig. 1286.—*Agaricoerinus profundus*,
summit view.

Fig. 1287.—*Agaricoerinus profundus*,
azygous side of a small specimen.

Fig. 1288.—*Agaricoerinus sampsoni*, basal
view, enlyx and arms.

Fig. 1289.—*Agaricoerinus tugurium*,
basal view.

Fig. 1290.—*Agaricoerinus tugurium*, summit
view.

Fig. 1291.—*Agaricoerinus tugurium*, azygous
side view.

Fig. 1292.—*Agassizocrinus* ovals,
enlyx and arms, azygous view.

AGELACRINUS fuberi, S. A. Miller, 1894, Jour.
Cin. Soc. Nat. Hist.,
vol. 17, p. 156, Hud.
Riv. Gr.

legrandensis, Miller
and Gurley, 1894,
Bull. No. 5, Ill. St.
Mus. Nat. Hist., p. 15
Kinderhook Gr.

Fig. 1293.
Agelacrinus blairi,
magnified twodi-
ameters.

pulaskiensis, Miller and
Gurley, 1894, Bull.
No. 5, Ill. St. Mus.
Nat. Hist., p. 16 Kaskaskia Gr.

AGELADISCUS n. gen. [Ety. *agyle*, herd; *diskos*, quoit.] This name is proposed instead of *Echinodiscus*, Worthen and Miller, which was preoccupied by Breynius. It will include *A. optatus*, *A. kaskascensis*, and *A. sampsoni*. *Ageladiscus* will be a much better word, because the fossils belong to the *Agelacriniidae*, and have no near affinity to the *Echinida*.

ALLAGECRINUS americanus, Rowley, 1895, Am. Geol., vol. 16, p. 219, Chouteau Gr.

ALLOPHOSALLOCRINUS celsus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 47, Warsaw Gr.

AMPHORACRINUS blairi, Miller and Gurley, 1890, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 26, Burlington Gr.

inflatus, Hall, 1860, (Actinocrinus inflatus,) Supp. Geo. Iowa, p. 20, and Mem. Am. Mus. Nat. Hist., vol. 1, p. 22, Burlington Gr.

jessieae, Miller and Gurley, 1896, Bull.

Fig. 1291.—*Amphoracrinus jessieae*, azygous and basal views.

No. 10, Ill. St. Mus. Nat. Hist. p. 21, Chouteau Gr.

planobasalis was described in 1860 in the Supp. to Geo. of Iowa.

sampsoni, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 27, Chouteau Gr.

sedaliensis, Miller, and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 54, Chouteau Gr.

Fig. 1295.—*Amphoracrinus sedaliensis*, azygous and basal views.

viminalis refer to the Keokuk Gr.

Aorocrinus, Wachsmuth and Springer, Syn. for *Dorycerinus*. Type *Dorycerinus immaturus*.

Aristocrinus, Rowley, 1895, Am. Geol., vol. 16, p. 217. [Ety. *aristos*, best; *krinon*, lily.] Type *Taocrinus concavus*. Imperfectly defined, probably a synonym.

ARCHAEOCRINUS asperatus, Miller and Gur-

Fig. 1296.—*Archaeocrinus asperatus*, basal and summit views.

ley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 19, Trenton Gr.

Fig. 1297.—*Archaeocrinus asperatus*, azygous side.

knoxensis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 34, Trenton Gr.

parvus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 21, Trenton Gr.

Fig. 1298.—*Archaeocrinus parvus*, basal, azygous, and summit views.

peculiaris, Miller and Gurley, 1894, Bull.

Fig. 1299.—*Archaeocrinus peculiaris*, lateral and azygous views.

No. 5, Ill. St. Mus. Nat. Hist., p. 17, Trenton Gr.

Fig. 1300.—*Archaeocrinus peculiaris*, basal view.

ARTHRAACANTHIA depressa, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 751, Chemung Gr.

ASTROCYSTITES, Whiteaves, 1897, Can. Rec. Sci., p. 287. [Ety. *aster*, star; *kystis*, bladder.] General form obovate, terminating in a column below. Summit, in structure, much like that in *Esio-cystites*, with five radiating grooves

cusula, Miller and Gurley, 1895, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 8, Keokuk Gr.

elstula, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 18, St. Louis Gr.

clavigerus is from the Keokuk Gr. ello, Hall, 1861, (Actinoerinus ello,) Bost. Jour. Nat. Hist., vol. 7, p. 262, Burlington Gr.

cloella, Hall, 1861, (Actinoerinus cloella,) Bost. Jour. Nat. Hist., vol. 7, p. 266, Burlington Gr.

cognatus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 60, Burlington Gr.

commendabilis, Miller and Gurley, 1895, (Eretmocrinus commendabilis,) Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 25, Keokuk Gr.

complanatus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 27, Burlington Gr.

consanguineus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 63, Burlington Gr.

copiosus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 51, Warsaw Gr.

corbulis, Hall, 1861, (Actinoerinus corbulis,) Bost. Jour. Nat. Hist., vol. 7, p. 265, Burlington Gr.

Fig. 1305.—*Batoerinus curiosus*, azygous side.

curiosus, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 6, Keokuk Gr.

delicatulus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 12, Keokuk Gr.

depressus, Keyes, 1895, (Eretmocrinus depressus,) Geol. Sur. Mo., vol. 4, p. 176, Burlington Gr.

douglasi, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 29, Keokuk Gr.

enodatus, Miller and Gurley, 1896, Bull.

Fig. 1300.—*Batoerinus curiosus*, basal view.

No. 10, Ill. St. Mus. Nat. Hist., p. 46, Burlington Gr.

Fig. 1307.—*Batoerinus enodis*, basal and azygous views.

enodis, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 25, Burlington Gr.

faberi, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 18, Burlington Gr.

folliculus, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 15, Burlington Gr.

formaceus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 24, Burlington Gr.

gemmiformis, Hall, 1860, (Actinoerinus gemmiformis,) Supp. Geo. Sur. Iowa, p. 23, Burlington Gr.

germanus, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 5, Burlington Gr.

Fig. 1308.—*Batoerinus glaber*, opposite azygous area.

glaber, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 32, Burlington Gr.

heteroclitus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 31, Keokuk Gr.

hodgsoni, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 8, Burlington Gr.

honorabilis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 10, Keokuk Gr.

ignotus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 28, Keokuk Gr.

imparilis, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 20, Burlington Gr.

Fig. 1309.—*Batoerinus honorabilis*, opposite azygous side.

Nat. Hist., p. 46,

is, basal and azygous

Gurley, 1896, Bull. Nat. Hist., p. 25,

ley, 1896, Bull. No. Hist., p. 18, Bur-

Gurley, 1896, Bull. Nat. Hist., p. 15,

Gurley, 1895, Bull. Nat. Hist., p. 24.

1890, (Actinoerinus p. Geo. Sur. Iowa, p.

Gurley, 1896, Bull. Nat. Hist., p. 5.

glaber, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 32, Burlington Gr.

heteroglitus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 31, Keokuk Gr.

hodgei, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 31, Keokuk Gr.

130.—Batoerinus habellum, opposite azygous side.

inconspicuous, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 25, Keokuk Gr.

incultus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 21, Burlington Gr.

incipitatus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 30, Keokuk Gr.

insolens, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 35, Burlington Gr.

insperatus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 22, Burlington Gr.

insuetus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 14, Keokuk Gr.

intermedius, Wachsmuth and Springer, 1881, (Eretmoerinus intermedius,) Proc. Acad. Nat. Sci. Phil., p. 348, Keokuk Gr.

jesseni, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 20, Burlington Gr.

konincki, Shumard, 1855, (Actinoerinus konincki,) Geo. Sur. Mo., p. 194, Burlington Gr.

Fig. 130.—Batoerinus habellum, azygous side.

labellum, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 21, Keokuk Gr.

laciniosus, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 14, Keokuk Gr.

letus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 24, Burlington Gr.

lerna, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 12, Keokuk Gr.

laura is figured in Mem. Am. Mus. Nat. Hist., vol. 1, p. 17.

lepidus is figured in Mem. Am. Mus. Nat. Hist., vol. 1, p. 16.

leucosia, Hall, 1861, (Actinoerinus leucosia,) Bost. Jour. Nat. Hist., vol. 7, p. 261, Burlington Gr.

levigatus, Miller and Gurley, 1896, Bull.

Fig. 132.—Batoerinus levis, azygous and summit views.

No. 10, Ill. St. Mus. Nat. Hist., p. 20, Burlington Gr.

levis, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 23, Burlington Gr.

lyonatus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 18, St. Louis Gr.

magnilleus, Lyon and Casseday, 1859, (Eretmoerinus magnilleus,) Am. Jour. Sci., vol. 28, p. 241, Keokuk Gr.

minor, Wachsmuth and Springer, 1897, (Eretmoerinus minor,) N. Am. Crin. Cam., vol. 2, p. 301, Burlington Gr.

modestus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 30, Keokuk Gr.

modulus, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 16, Burlington Gr.

nanus, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 17, Burlington Gr.

neglectus, Meek and Worthen, 1869, Proc. Acad. Nat. Sci., p. 355, and Geo. Sur. Ill., vol. 5, p. 377, Burlington Gr.

nitens, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 15, Burlington Gr.

nitidulus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 17, Keokuk Gr.

nodosarius, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 22, Burlington Gr.

nodosus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 5, Burlington Gr.

nodulosus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

oblatus is figured in Mem. Am. Mus. Nat. Hist., vol. 1, p. 12.

obtusus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 22, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 5, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

obtusus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 6, Burlington Gr.

originarius, Wachsmuth and Springer, 1881, (*Eretmocrinus originarius*.) Proc. Acad. Nat. Sci., p. 348, Keokuk Gr.

Fig. 1315.—*Batoerinus parilis*, calyx and arms, azygous view of calyx.

parilis, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 17, Burlington Gr.

peculiaris, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 18, Keokuk Gr.

Fig. 1316.—*Batoerinus pettisensis*, azygous and opposite views.

pettisensis, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 19, Burlington Gr.

Fig. 1317.—*Batoerinus pileus*, lateral view.

pileus, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 18, St. Louis Gr.

planus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus.

Nat. Hist., p. 37, Burlington Gr.
politus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 31, Burlington Gr.

polydactylus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 5, Keokuk Gr.

proboesidialis, Hall, instead of *Actino-*
crinus proboscidialis.

procerus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 33, Keokuk Gr.

prodiginis, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 30, Keokuk Gr.

proximus, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 7, Burlington Gr.

ramulosus, Hall, 1858, (*Actinoerinus ramulosus*.) Geo. Sur. Iowa, p. 615, Keokuk Gr.

regalis, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 14, Burlington Gr.

Fig. 1318.—*Batoerinus prodiginis*, azygous side.

reliquus, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 22, Burlington Gr.

semibrauchiatus var. *expansus*, Wachsmuth and Springer, Syn. for *E. cassidynus*.

remotus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 43, Burlington Gr.

repertus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 70, Burlington Gr.

repositus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 45, Burlington Gr.

reservatus, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 25, Burlington Gr.

robustus, Wachsmuth and Springer, 1897, (*Loboerinus robustus*.) N. Am. Crin. Cam., vol. 2, p. 436, Keokuk Gr.

rotuliformis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 26, Burlington Gr.

rudis, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 52, Keokuk Gr.

rugosus, Wachsmuth and Springer, 1897, (*Eretmocrinus*.) N. Am. Crin. Cam., vol. 2, p. 402, Burlington Gr.

rusticellus, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 23, Burlington Gr.

rusticus, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 28, Burlington Gr.

saccellus, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 40, Burlington Gr.

sacculus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 52, Warsaw Gr.

sagetownensis, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 54, Burlington Gr.

salemensis, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 8, Warsaw Gr.

Gurley, 1896, Bull.
Nat. Hist., p. 14,

Wachsmuth and Springer, 1897, Bull.
Nat. Hist., p. 22,

Wachsmuth and Springer, 1897, Bull.
Nat. Hist., p. 43,

Gurley, 1896, Bull.
Nat. Hist., p. 70,

Gurley, 1896, Bull.
Nat. Hist., p. 45,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

Gurley, 1897, Bull.
Nat. Hist., p. 25,

sampsoni, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 7,
Keokuk Gr.

scitulus, Miller and Gurley, 1897, Bull.
No. 12, Ill. St. Mus. Nat. Hist., p. 19,
Burlington Gr.

scyphus, Miller and Gurley, 1894, Bull.
No. 3, Ill. St. Mus. Nat. Hist., p. 23,
Burlington Gr.

sedaliensis, Miller and Gurley, 1896,
Bull. No. 10, Ill. St. Mus. Nat. Hist.,
p. 71, Burlington Gr.

selectus, Miller and Gurley, 1896, Bull.
No. 10, Ill. St. Mus. Nat. Hist., p. 37,
Burlington Gr.

senex, Miller and Gurley, 1897, Bull. No.
12, Ill. St. Mus. Nat. Hist., p. 21, Bur-
lington Gr.

serratus, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 27,
Keokuk Gr.

sharonensis, Miller and Gurley, 1897,
Bull. No. 12, Ill. St. Mus. Nat. Hist.,
p. 18, Burlington Gr.

shepardi, Rowley, 1893, Am. Geol., vol.
12, p. 305. Not defined so as to be
recognized.

signatus, Miller and Gurley, 1896, Bull.
No. 9, Ill. St. Mus. Nat. Hist., p. 10,
Keokuk Gr.

simplex, Wachsmuth and Springer,
1897, (Dizygocirrus indianensis var.
simplex,) N. Am. Crin. Can. vol. 2, p.
416, Keokuk Gr.

solitarius, Miller and Gurley, 1896, Bull.
No. 8, Ill. St. Mus. Nat. Hist., p. 22,
Burlington Gr.

speciosus, Miller and Gurley, 1896, Bull.
No. 10, Ill. St. Mus. Nat. Hist., p. 47,
Burlington Gr.

spinosis, Miller and Gurley, 1895, Bull.
No. 6, Ill. St. Mus. Nat. Hist., p. 5,
Keokuk Gr.

spurius, Miller and Gurley, 1896, Bull.
No. 9, Ill. St. Mus. Nat. Hist., p. 20,
Burlington Gr.

stelliformis, Miller and Gurley, 1896,
Bull. No. 9, Ill. St. Mus. Nat. Hist.,
p. 9, Keokuk Gr.

strenuus, Miller and Gurley, 1896, Bull.
No. 9, Ill. St. Mus. Nat. Hist., p. 11,
Keokuk Gr.

subaequatus, Miller and Gurley, 1896,
Bull. No. 10, Ill. St. Mus. Nat. Hist.,
p. 72, Burlington Gr.

sublevis, Miller and Gurley, 1896, Bull.
No. 10, Ill. St. Mus. Nat. Hist., p. 41,
Burlington Gr.

subovatus, Miller and Gurley, 1896,
Bull. No. 10, Ill. St. Mus. Nat. Hist.,
p. 50, Burlington Gr.

subrotundus, Miller and Gurley, 1896,
Bull. No. 10, Ill. St. Mus. Nat. Hist.,
p. 48, Burlington Gr.

subscitulus, Miller and Gurley, 1896,
Bull. No. 10, Ill. St. Mus. Nat. Hist.,
p. 51, Burlington Gr.

superbus, n. sp. Proposed instead of
Eretmocirrus cassidagynus, Miller and
Gurley in Bull. No. 3, Ill. St. Mus.
Nat. Hist., p. 17, which name is pre-
occupied in this genus. (By typo-
graphical error, it was also printed
Eretmocirrus lymanus at same place.)
Burlington Gr.

tuberculatus, Wachsmuth and Springer,
1897, N. Am. Crin. Can., vol. 2, p. 379,
Burlington Gr.

variabilis, Miller and Gurley, 1896, Bull.
No. 10, Ill. St. Mus. Nat. Hist., p. 58,
Burlington Gr.

varsouviensis, Worthen, 1882, (Eretmo-
cirrus varsouviensis,) Geo. Sur. Ill.,
vol. 7, p. 306, Warsaw Gr.

venustus, Miller and Gurley, 1895,
Bull. No. 7, Ill. St. Mus. Nat. Hist.,
p. 12, Keokuk Gr.

verneuilianus, Shumard, 1855, (Actino-
cirrus verneuilianus,) Geo. Sur. Mo., p.
193, Burlington Gr.

veterator, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 8,
Keokuk Gr.

vetustus, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 10,
Keokuk Gr.

vicinus, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 34,
Keokuk Gr.

wetherbyi, Miller and Gurley, 1895, Bull.
No. 6, Ill. St. Mus. Nat. Hist., p. 11,
Keokuk Gr.

Fig. 1319.—*Butoerchus*
spherosus, azygous
side.

Fig. 1320.—*Belemnocirrus*
sampsoni.

Fig. 1321.—*Belemnocirrus*
sampsoni.

Fig. 1322.—*Belemnocirrus*
sampsoni.

Fig. 1323.—*Belemnocirrus*
sampsoni.

Fig. 1324.—*Belemnocirrus*
sampsoni.

Fig. 1325.—*Belemnocirrus*
sampsoni.

Fig. 1326.—*Belemnocirrus*
sampsoni.

subrotundus, Miller and Gurley, 1896,
Bull. No. 10, Ill. St. Mus. Nat. Hist.,
p. 48, Burlington Gr.

subscitulus, Miller and Gurley, 1896,
Bull. No. 10, Ill. St. Mus. Nat. Hist.,
p. 51, Burlington Gr.

superbus, n. sp. Proposed instead of
Eretmocirrus cassidagynus, Miller and
Gurley in Bull. No. 3, Ill. St. Mus.
Nat. Hist., p. 17, which name is pre-
occupied in this genus. (By typo-
graphical error, it was also printed
Eretmocirrus lymanus at same place.)
Burlington Gr.

tuberculatus, Wachsmuth and Springer,
1897, N. Am. Crin. Can., vol. 2, p. 379,
Burlington Gr.

variabilis, Miller and Gurley, 1896, Bull.
No. 10, Ill. St. Mus. Nat. Hist., p. 58,
Burlington Gr.

varsouviensis, Worthen, 1882, (Eretmo-
cirrus varsouviensis,) Geo. Sur. Ill.,
vol. 7, p. 306, Warsaw Gr.

venustus, Miller and Gurley, 1895,
Bull. No. 7, Ill. St. Mus. Nat. Hist.,
p. 12, Keokuk Gr.

verneuilianus, Shumard, 1855, (Actino-
cirrus verneuilianus,) Geo. Sur. Mo., p.
193, Burlington Gr.

veterator, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 8,
Keokuk Gr.

vetustus, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 10,
Keokuk Gr.

vicinus, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 34,
Keokuk Gr.

wetherbyi, Miller and Gurley, 1895, Bull.
No. 6, Ill. St. Mus. Nat. Hist., p. 11,
Keokuk Gr.

Fig. 1320.—*Belemnocirrus*
sampsoni.

Fig. 1321.—*Belemnocirrus*
sampsoni.

Fig. 1322.—*Belemnocirrus*
sampsoni.

Fig. 1323.—*Belemnocirrus*
 

Fig. 1324.—*Belemnocirrus*
 

Fig. 1325.—*Belemnocirrus*
 

Fig. 1326.—*Belemnocirrus*
 

Fig. 1327.—*Belemnocirrus*
 

Fig. 1328.—*Belemnocirrus*
 

Fig. 1329.—*Belemnocirrus*
 

Fig. 1330.—*Belemnocirrus*
 

Fig. 1331.—*Belemnocirrus*
 

Fig. 1332.—*Belemnocirrus*
 

Fig. 1333.—*Belemnocirrus*
 

Fig. 1334.—*Belemnocirrus*
 

Fig. 1335.—*Belemnocirrus*
 

Fig. 1336.—*Belemnocirrus*
 

Fig. 1337.—*Belemnocirrus*
 

Fig. 1338.—*Belemnocirrus*
 

Fig. 1339.—*Belemnocirrus*
 

Fig. 1340.—*Belemnocirrus*
 

Fig. 1341.—*Belemnocirrus*
 

Fig. 1342.—*Belemnocirrus*
 

Fig. 1343.—*Belemnocirrus*
 

Fig. 1344.—*Belemnocirrus*
 

Fig. 1345.—*Belemnocirrus*
 

Fig. 1346.—*Belemnocirrus*
 

Fig. 1347.—*Belemnocirrus*
 

Fig. 1348.—*Belemnocirrus*
 

Fig. 1349.—*Belemnocirrus*
 

Fig. 1350.—*Belemnocirrus*
 

Fig. 1351.—*Belemnocirrus*
 

Fig. 1352.—*Belemnocirrus*
 

Fig. 1353.—*Belemnocirrus*
 

Fig. 1354.—*Belemnocirrus*
 

Fig. 1355.—*Belemnocirrus*
 

Fig. 1356.—*Belemnocirrus*
 

Fig. 1357.—*Belemnocirrus*
 

Fig. 1358.—*Belemnocirrus*
 

Fig. 1359.—*Belemnocirrus*
 

Fig. 1360.—*Belemnocirrus*
 

Fig. 1361.—*Belemnocirrus*
 

Fig. 1362.—*Belemnocirrus*
 

Fig. 1363.—*Belemnocirrus*
 

Fig. 1364.—*Belemnocirrus*
 

Fig. 1365.—*Belemnocirrus*
 

Fig. 1366.—*Belemnocirrus*
 

Fig. 1367.—*Belemnocirrus*
 

Fig. 1368.—*Belemnocirrus*
 

Fig. 1369.—*Belemnocirrus*
 

Fig. 1370.—*Belemnocirrus*
 

Fig. 1371.—*Belemnocirrus*
 

Fig. 1372.—*Belemnocirrus*
 

Fig. 1373.—*Belemnocirrus*
 

Fig. 1374.—*Belemnocirrus*
 

Fig. 1375.—*Belemnocirrus*
 

Fig. 1376.—*Belemnocirrus*
 

Fig. 1377.—*Belemnocirrus*
 

Fig. 1378.—*Belemnocirrus*
 

Fig. 1379.—*Belemnocirrus*
 

Fig. 1380.—*Belemnocirrus*
 

Fig. 1381.—*Belemnocirrus*
 

Fig. 1382.—*Belemnocirrus*
 

Fig. 1383.—*Belemnocirrus*
 

Fig. 1384.—*Belemnocirrus*
 

Fig. 1385.—*Belemnocirrus*
 

Fig. 1386.—*Belemnocirrus*
 

Fig. 1387.—*Belemnocirrus*
 

Fig. 1388.—*Belemnocirrus*
 

Fig. 1389.—*Belemnocirrus*
 

Fig. 1390.—*Belemnocirrus*
 

Fig. 1391.—*Belemnocirrus*
 

Fig. 1392.—*Belemnocirrus*
 

Fig. 1393.—*Belemnocirrus*
 

Fig. 1394.—*Belemnocirrus*
 

Fig. 1395.—*Belemnocirrus*
 

Fig. 1396.—*Belemnocirrus*
 

Fig. 1397.—*Belemnocirrus*
 

Fig. 1398.—*Belemnocirrus*
 

Fig. 1399.—*Belemnocirrus*
 

Fig. 1400.—*Belemnocirrus*
 

Fig. 1401.—*Belemnocirrus*
 

Fig. 1402.—*Belemnocirrus*
 

Fig. 1403.—*Belemnocirrus*
 

Fig. 1404.—*Belemnocirrus*
 

Fig. 1405.—*Belemnocirrus*
 

Fig. 1406.—*Belemnocirrus*
 

Fig. 1407.—*Belemnocirrus*
 

Fig. 1408.—*Belemnocirrus*
 

Fig. 1409.—*Belemnocirrus*
 

Fig. 1410.—*Belemnocirrus*
 

Fig. 1411.—*Belemnocirrus*
 

Fig. 1412.—*Belemnocirrus*
 

Fig. 1413.—*Belemnocirrus*
 

Fig. 1414.—*Belemnocirrus*
 

Fig. 1415.—*Belemnocirrus*
 

Fig. 1416.—*Belemnocirrus*
 

Fig. 1417.—*Belemnocirrus*
 

Fig. 1418.—*Belemnocirrus*
 

Fig. 1419.—*Belemnocirrus*
 

Fig. 1420.—*Belemnocirrus*
 

Fig. 1421.—*Belemnocirrus*
sampson

ventral side, with an arm in the anterior part. Type *B. wetherbyi*. *wetherbyi*, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 9, Trenton Gr.

Fig. 1321.—*Botryocrinus wetherbyi*, dorsal views of two specimens, and ventral view.

BLATROCRINUS spinosulus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 28, Chouteau Gr.

Cactocrinus, Wachsmuth and Springer, Syn. for *Batrocrinus*. Type *Batrocrinus proboscideialis*.

denticulatus, Wachsmuth and Springer. See *Actinocrinus denticulatus*.

obesus, Keyes. See *Actinocrinus obesus*.

extensus. See *Actinocrinus extensus*.

CALCEOCRINUS. This name was applied by Hall, in 1852—*Publ. N. Y.*, vol. 2, p. 352—to the basal plates of a crinoid from the Niagara Gr., without proposing a specific name. In 1860—13th Rep. N. Y. St. Mus. Nat. Hist., p. 122—he described and illustrated *Cheirocrinus chrysalis* from the Niagara Gr. The name *Cheirocrinus* was preoccupied. Shumard, in 1866, discovered that *Cheirocrinus* was identical with *Calceocrinus* and classified the species under that name. This being the fact, the type became *Calceocrinus chrysalis* from 1860, not from 1852. There is no excuse for the recent synonyms proposed for this genus.

bidentatus, *contractus*, *balli*, typus. Ringueberg, 1889, Ann. N. Y. Acad. Sci., vol. 4, p. 388, Niagara Gr.

Fig. 1322.—*Calceocrinus kentuckiensis*, anterior and azygous views.

kentuckiensis, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 29, Trenton Gr. See *Castocrinus*.

CALLICRINUS beaulei, Wachsmuth and Springer, 1897, N. Am. Crin. Cont., vol. 1, p. 355, Niagara Gr.

Camptocrinus, Wachsmuth and Springer, Syn. for *Dichocrinus*.

cirriifer. See *Dichocrinus cirriifer*.

myelodaetylus. See *Dichocrinus myelodaetylus*.

CARABOOCRINUS ovalis, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 25, Trenton Gr.

Fig. 1323.—*Carabocrinus ovalis*, azygous and opposite views.

CARYOOCRINUS bulbulus, Miller and Gurley, 1894, Bull. No. 5, Ill.

Fig. 1324.—*Caryocrinus bulbulus*, anterior and posterior views.

St. Mus. Nat. Hist., p. 11, Niagara Gr. *ellipticus*, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 10, Niagara Gr.

Fig. 1325.—*Caryocrinus ellipticus*, anterior and summit views.

hammelli, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 65, Niagara Gr.

kentuckiensis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 59, Niagara Gr.

Fig. 1326.—*Caryocrinus kentuckiensis*, anterior, posterior, and summit views.

milliganae, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 63, Niagara Gr.

CASTOCRINUS, Ringueberg, 1889, Ann. N. Y. Acad. Sci., vol. 4, p. 388. This name was proposed for one of the divisions, to be made of the genus *Calceocrinus*, with *Calceocrinus furcillatus* as the type, and to include, *C. rugosus*, *C. articulatus*, *C. inaequalis*, and a new species, *C. hillingtonensis*, described, in the same place, from the Trenton Gr. It would seem that the name should stand, (it was accidentally overlooked in the first appendix.) and, if so, it will include *Calceocrinus kentuckiensis*.

ichocrinus myelo-

Fig. 1324.—*Cyathocrinus ovalis*, azygous and opposite views.

St. Mus. Nat. Hist., p. 11, Niagara Gr.

ellipticus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Hist., p. 10, Niagara Gr.

ellipticus, anterior and summit views.

Gurley, 1896, Bull. Nat. Hist., p. 65,

and Gurley, 1895, Mus. Nat. Hist., p.

blairi, azygous and opposite views.

d Gurley, 1896, Mus. Nat. Hist.

erg, 1889, Ann. N. Y. Mus., p. 388. This name is one of the divisions of the genus *Calceocrinus*, *furcillatus* as the name, *C. rugosus*, *C. equalis*, and a new species, described, in from the Trenton formation that the name was accidentally first appendix.) and, of *Calceocrinus ken-*

CODASTER blairi, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 86, Chouteau Gr.

Fig. 1327.—*Codaster blairi*, basal, lateral, and summit views.

jessiae, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 89, Chouteau Gr.

whitei, refer to *Codonites whitei*.

CONOXITES conicus, in Geo. Sur. Ill., vol. 8, p. 201, Whitfield says, in Mem. Mus., is a syn. for *C. whitei*.

whitei, Hall, 1861, (*Codaster whitei*.) is figured, by Whitfield, in Mem. Am. Mus. Nat. Hist., vol. 1, p. 36, and was collected in the Kinderhook Gr.

COELOCRINUS dilatatus, Hall, is figured in Mem. Am. Mus. Nat. Hist., vol. 1, p. 33.

CATHOCRINUS andersoni, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 30, Keokuk Gr.

blairi, Miller and Gurley, 1895, Bull. No. 7, p. 67, and Bull. No. 8, p. 50, Ill. St. Mus. Nat. Hist., Chouteau Gr.

brittsi, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 70, Burlington Gr.

chouteauensis, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 68, and Bull. No. 9, p. 38, Chouteau Gr.

faberi, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 85, Burlington Gr.

inacquadactylus, Whitfield. See *C. maxvillensis*.

labyrinthicus. See *Poteriocrinus labyrinthicus*.

macadamisi, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 69, St. Louis Gr.

maxvillensis, Whitfield, 1895, Ohio Geo. vol. 7, p. 465. Proposed instead of *C. inacquadactylus* of Whitfield, which was preoccupied.

meekanus, Shumard, 1855, (*Poteriocrinus meekanus*.) Geo. Rep. of Mo., p. 188, Chouteau Gr.

signatus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 32, Keokuk Gr.

tumidulus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 31, Keokuk Gr.

waldronensis, redescribed and refigured in Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 49.

CYCLOCYSTOIDES, illinoisensis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 61, Hudson Riv. Gr.

CYCLOCRINUS indianensis, Miller and Gur-

Fig. 1331.—*Cythoecrinus indianensis*, basal, azygous, and lateral views.

ley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 31, Niagara Gr.

CYPHOCRINUS. The advance sheets of the 18th Rep. Geo. Sur. Indiana were printed in August, 1892, and were mailed and distributed on the 1st day of September, 1892, as can be proven by the State geologist, at that time, and by his assistants and by many who received the work. The statements, therefore, which appear in North American Crinoida Camerata, by Wachsmuth and Springer, vol. 1, pp. 200 to 203, that it was published October 26, 1892, are untrue.

Diaboloecrinus, Wachsmuth and Springer, Syn. for *Archaeocrinus*.

hieroglyphicus, Wachsmuth and Springer, Syn. for *Archaeocrinus asperatus*.

perplexus, Wachsmuth and Springer, Syn. for *Archaeocrinus knoxensis*.

DICRINOCRINUS bozemanensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 44, Keokuk Gr.

cirifer, Wachsmuth and Springer, 1897, (*Camptocrinus cirifer*.) N. Amer. Crin. Cam., vol. 2, p. 780, Kaskaskia Gr.

delicatulus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 766, Kinderhook Gr.

huntsville, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 773, St. Louis Gr.

Fig. 1332.—*Dichocrinus blairi*.

- myelodactylus, Wachsmuth and Springer, 1897, (Camptocrinus myelodactylus.) N. Am. Crin. Cam., vol. 2, p. 779, Keokuk Gr.
- oblongus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 759, Warsaw Gr.
- pendens, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 774, Burlington Gr.
- superstes, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 766, Kaskaskia Gr.
- Dizygocrinus*, Wachsmuth and Springer, syn. for *Batoerinus*. Type *Batoerinus indianensis*.
- indianensis* var. *simplex*. See *Batoerinus simplex*.
- cantonensis*, Wachsmuth and Springer, Syn. for *Batoerinus labellum*.
- mutabilis*, Wachsmuth and Springer, Syn. for *Batoerinus vicinus*.
- DOLATOERINUS* unplus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 45, Ham. Gr.
- apertus, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 48, and Bull. No. 9, p. 49, Ham. Gr.
- approximatus, Miller and Gurley, 1894, Bull. No. 4, Ill. St. Mus. Nat. Hist., p. 25, Ham. Gr.
- argutus, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 41, Ham. Gr.
- arrosus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 52, Ham. Gr.

Fig. 1333.—*Dolatoerinus asper*, basal and summit views.

- asper*, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 47, Ham. Gr.

Fig. 1334.—*Dolatoerinus aspratilis*, basal and summit views.

- aspratilis*, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 49, Ham. Gr.
- aureatus*, Miller and Gurley, 1894, Bull.

- No. 4, Ill. St. Mus. Nat. Hist., p. 24, Ham. Gr.
- basiliensis*, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 43, Ham. Gr.

Fig. 1335.—*Dolatoerinus basiliensis*, basal and summit views.

- bellarugosus*, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 43, Ham. Gr.
- bellulus*, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 57, Ham. Gr.
- bullaceus*, Miller and Gurley, 1894, Bull. No. 4, Ill. St. Mus. Nat. Hist., p. 22, Ham. Gr.
- celatus*, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 46, Ham. Gr.

Fig. 1336.—*Dolatoerinus charlestownensis*, basal and summit views.

- charlestownensis*, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 44, Ham. Gr.
- cistula*, Miller and Gurley, 1896, Bull.

Fig. 1337.—*Dolatoerinus corporosus*, zygous side.

- No. 9, Ill. St. Mus. Nat. Hist., p. 46, Ham. Gr.
- corporosus*, Miller and Gurley, 1895,

Nat. Hist., p. 24,
Gurley, 1896, Bull.
Nat. Hist., p. 43,

dissimularis, basal and
summit views.

and Gurley, 1896,
Mus. Nat. Hist.,

Gurley, 1895, Bull.
Nat. Hist., p. 57,

Gurley, 1894, Bull.
Nat. Hist., p. 22,

Gurley, 1896, Bull.
Nat. Hist., p. 46,

arlestownensis, basal
views.

Miller and Gurley,
Ill. St. Mus. Nat.
Hist., p. 54.
Gurley, 1896, Bull.

corporosus, azygous

Nat. Hist., p. 46,

and Gurley, 1895,

Bull. No. 6, Ill. St. Mus. Nat. Hist., p.
50, Ham. Gr.

Fig. 138.—*Dolotoerinus corporosus*, basal
view.

dispar, Miller and Gurley, 1896, Bull.
No. 9, Ill. St. Mus. Nat. Hist., p. 40,
Ham. Gr.

Fig. 139.—*Dolotoerinus dissimularis*, basal and
summit views.

dissimularis, Miller and Gurley, 1896,
Bull. No. 9, Ill. St. Mus. Nat. Hist., p.
54 Ham. Gr.

excavatus, Wachsmuth and Springer,
Syn. for *D. grandis*.

Fig. 140.—*Dolotoerinus exornatus*, basal and lateral
views.

grandis, Miller and Gurley, 1894, Bull.
No. 4, Ill. St. Mus. Nat. Hist., p. 14,
Ham. Gr.

greeni, Miller and Gurley, 1894, Bull.
No. 4, Ill. St. Mus. Nat. Hist., p. 28,
Ham. Gr.

harmelli, Miller and Gurley, 1895, Bull.
No. 6, Ill. St. Mus. Nat. Hist., p. 52,
Ham. Gr.

icosidactylus, Wachsmuth and Springer,
syn. for *D. greeni*.

indianensis, Miller and Gurley, 1896,
Bull. No. 8, Ill. St. Mus. Nat. Hist., p.
40, Ham. Gr.

ingueculi, Miller and Gurley, 1896, Bull.
No. 9, Ill. St. Mus. Nat. Hist., p. 51,
Ham. Gr.

lineolatus, Miller and Gurley, 1894, Bull.
No. 4, Ill. St. Mus. Nat. Hist., p. 27,
Ham. Gr.

lyoni, Miller and Gurley, 1896, Bull. No.
9, Ill. St. Mus. Nat. Hist., p. 44, Ham.
Gr.

lyoni, Wachsmuth and Springer, 1897.
The name was preoccupied.

major, Wachsmuth and Springer, Syn.
for *D. spinosus*.

magnificus, Miller and Gurley, 1894,
Bull. No. 4, Ill. St. Mus. Nat. Hist., p.
5, Ham. Gr.

marshi var. *hamiltonensis*, Wachsmuth
and Springer, Syn. for *D. ornatus* var.
asperatus.

neglectus, Miller and Gurley, 1894, Bull.
No. 12, Ill. St. Mus. Nat. Hist., p. 37,
Ham. Gr.

nodosus, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 56,
Ham. Gr.

ornatus var. *asperatus*, Miller and Gur-
ley, 1894, Bull. No. 4, Ill. St. Mus.
Nat. Hist., p. 16, Ham. Gr.

peculiaris, Miller and Gurley, 1896, Bull.
No. 9, Ill. St. Mus. Nat. Hist., p. 55,
Ham. Gr.

preciosus, Miller and Gurley, 1896, Bull.
No. 9, Ill. St. Mus. Nat. Hist., p. 41,
Ham. Gr.

pulehellus, Miller and Gurley, 1895, Bull.
No. 6, Ill. St. Mus. Nat. Hist., p. 55,
Ham. Gr.

succulus, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 58,
Ham. Gr.

sulcibrosus, Miller and Gurley, 1895, Bull.
No. 7, Ill. St. Mus. Nat. Hist., p. 59,
Ham. Gr.

spinosus, Miller and Gurley, 1894, Bull.
No. 4, Ill. St. Mus. Nat. Hist., p. 8,
Ham. Gr.

stellifer, Miller and Gurley, 1894, Bull.
No. 4, Ill. St. Mus. Nat. Hist., p. 20,
Ham. Gr.

Fig. 141.—*Dolotoerinus vasculum*, basal view.

tuberculatus, Wachsmuth and Springer,
Syn. for *D. bellulus*.

vasculum, Miller and Gurley, 1895, Bull.

No. 6, Ill. St. Mus. Nat. Hist., p. 53, Ham. Gr.

Fig. 1312.—*Dohioerinus vasculum*, azygous side.

venustus, Miller and Gurley, 1894, Bull. No. 4, Ill. St. Mus. Nat. Hist., p. 23, Ham. Gr.

DORYCERINUS *alabamensis*, Miller and Gur-

Fig. 1313.—*Dorycerinus amoenus*, azygous and side views.

ley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 15, Keokuk Gr.

Fig. 1314.—*Dorycerinus elegans*, azygous and summit views.

Fig. 1315.—*Dorycerinus faberi*, azygous and opposite views.

Ill. St. Mus. Nat. Hist., p. 48, Keokuk Gr.

parvus, refer to the St. Louis Gr. *pendens* is illustrated in Mem. Am. Mus. Nat. Hist., vol. 1, p. 18.

precursor is better referred to *Batocrinus*.

Fig. 1316.—*Dorycerinus simpsoni*, azygous and opposite views.

simpsoni, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 17, Burlington Gr.

subaeiformis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 30, Burlington Gr.

tricornis, Hall, 1858, (*Actinoerinus tricornis*.) Geo. Sur. Iowa, vol. 1, p. 569, and Am. Mus. Nat. Hist., vol. 1, p. 19, Burlington Gr.

ECHINOXYSTES was used in 1861, in the Edinburgh Phil. Jour., vol. 13, p. 106.

Echinodiscus, Worthen and Miller, 1883. This name was preoccupied by Breynius, in 1732, as the 7th genus in his *Dissertation Physien de Polythalamis*, p. 63. See *Ageladiscus*.

EMPEROCERINUS, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 42.

[Ety. *emperor*, deformed; *kri-non*, lily.] Basals three, unequal; subradials five, three heptagonal, two hexagonal. Primary radials, two in each ray. Regular interradials, one or more. Azygous plates, two or more. Ten ambulacral openings to the vault. Type *E. indianensis*.

indianensis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 43, Niagara Gr.

Emperocrinus. It has been shown, in Bulletins 7 to 10 of the Ill. St. Mus. Nat. Hist., that this genus is founded upon characters of specific value only, and that it is a synonym of *Batocrinus*. This has unfortunately resulted in a few preoccupied names, requiring others to be substituted in their places. I illustrate two of the most strongly-marked species that have been referred to this genus, and a comparison of them, with those illustrated under *Batocrinus*, will show they are congeneric.

adultus, *attenuatus*, *calyculoides*, *calyculoides* var. *nodosus*, *clio*, *claelia*, *commendabilis*, *corbulis*, *coronatus*, *depressus*, *gemmiformis*, *intermedius*, *konincki*, *leucosia*, *magnificus*, *minor*, *neglectus*, *originarius*, *ramulosus*, *remibrachiatus*, *rugosus*, *varsoviensis*, and *vernemiliannus*, all belong to *Batocrinus*.

Fig. 1318.—*Emperocrinus indianensis*, azygous side view.

Gurley, 1896, Bull.
Nat. Hist., p. 17.

and Gurley, 1897,
Mus. Nat. Hist., p.

(*Actinoerinus tri-*
pora, vol. 1, p. 569.
Hist., vol. 1, p. 19.

used in 1861, in
Ill. Jour., vol. 13.

and Miller, 1883.
occupied by Brey-

7th genus in his
de *Polythalamitis*.

and Gurley, 1895.

Fig. 1347.—*Emperoerinus*
indianensis, basal
view.

gous plates, two or
more. Ten am-
bulacral open-
ings to the vault.
Type *E. indian-*
ensis.

indianensis, Miller
and Gurley, 1895,
Bull. No. 6, Ill.
St. Mus. Nat.

Gr.

been shown, in
of the Ill. St. Mus.

s genus is founded
of specific value

a synonym of *Bo-*
unfortunately re-

reoccupied names,
to be substituted in

strate two of the
ked species that

to this genus, and

em, with those il-

toerinus, will show

congeneric.

calycu-
calyculoides var. *nor-*

io, *cloelia*, *commen-*
corbulis, *coronatus*,

gemmiformis, *in-*
s, *konincki*, *leucostia*,

us, *minor*, *neglectus*,

us, *ramulosus*, *reni-*
us, *rugosus*, *varso-*
and *verneuilianus*,
ing to *Batoerinus*.

Fig. 1349.—*Eretmoerinus commendabilis*, col-
umn, calyx and arms.

Fig. 1350.—*Eretmoerinus praevalis*.

cassedayanus, Miller and Gurley. See

Batoerinus superbus.

depressus, Wachsmuth and Springer,

Fig. 1351.—*Eucalyptocrinus elrodii*, basal view.

1897, was preoccupied by
Keyes in 1895.
expansus, Keyes. Not intel-
ligently defined.

Fig. 1352.—*Eucalyptocrinus el-*
rodii, side view.

Fig. 1353.—*Eucalyptocrinus gorbyi*,
side view.

granuliferus, Wachsmuth and
Springer. Syn. for *E. com-*
mendabilis.

EUCALYPTOCRINUS lindahli, Wachs-
muth & Springer,
Syn. for *E. wor-*
theni.

milliganae, Miller
and Gurley, 1896,
Bull. No. 10, Ill.

St. Mus. Nat. Hist., p. 88, Niag-
ara Gr.

ventricosus, Wachs-
muth and Springer, 1897, N.

Am. Crin. Cam., vol. 1, p. 341, Niag-
ara Gr.

wortheni, Miller and Gurley, 1894, Bull.
No. 3, Ill. St. Mus. Nat. Hist., p. 53,
Niagara Gr.

Eucheirocrinus, Meek and Worthen,
1873, Geo. Sur. Ill. vol. 5, p. 443. This

name was suggested as a possibility,
rather than defined. If it is to stand,

it has priority over *Deltacrinus*, and,
properly spelled, the type would be

Eucheirocrinus wachsmuthi. Clearly it
can not stand, if the law of nomen-
clature is applied.

EUCALDOCRINUS millebrachiatus var. *im-*
maturus, Wachsmuth and Springer,
N. Am. Crin. Cam., vol. 2, p. 722, Bur-

lington and Keokuk Gr.

EUPACHYCRINUS orbicularis is figured in

Fig. 1354.—*Eucalypto-*
crinus sub-
globosus.

Mem. Am. Mus. Nat. Hist., vol. 1, p. 30.

purvus, Miller and Gurley, 1893, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 60, Up. Coal Mens.

sanctiludovici, refer to *Hydreimnerius sanctiludovici*.

Eutrochocrinus, Wachsmuth and Springer, Syn for *Batoerinus*. Type *Batoerinus christyi*.

FORBESOCRINUS communis, refer to the Keokuk Gr.

Fig. 1355.—*Forbesocrinus elegantulus*, azygous and opposite views.

greeniei, Miller and Gurley, 1896, Bull.

No. 9, Ill. St. Mus. Nat. Hist., p. 57, Keokuk Gr.

jerseyensis Miller and Gurley, 1896, Bull. No. 8, Ill.

St. Mus. Nat. Hist., p. 58, Warsaw Gr.

kelloggi, refer to the Keokuk Gr.

macdamsi, Miller and Gurley, 1896, Bull. No. 9, Ill.

St. Mus. Nat. Hist., p. 58, Keokuk Gr.

multibrachiatus, Illustrated Bull. No. 8, Ill. St.

Mus. Nat. Hist., p. 55.

pyriformis, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p.

47, Keokuk Gr.

tardus, refer to the Keokuk Gr.

washingtonensis, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 54, Keokuk Gr.

GAZACRINUS. The advance sheets of the 18th Rep. Geo. Sur. Ind. were printed in August, 1892, and were mailed and distributed, on the 1st day of September, 1892, as can be proved by the State Geologist and his assistants, and by many who received the work. The statements therefore, which appear in North American Crinoidea Cameraata, by Wachsmuth and Springer, vol. 1, pp. 202 and 203, that the work was

published October 28, 1892, are untrue.

Fig. 1357.—*Forbesocrinus washingtonensis*, opposite side.

Gilbertsoerinus disparvus, Wachsmuth and Springer, Syn. for *Gonistasteroidocrinus lyonnus*.

indianensis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 38, Ham. Gr.

Fig. 1358.—*Gilbertsoerinus greeniei*, basal and lateral views.

greeniei, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus.

Nat. Hist., p. 35, Ham. Gr.

spinigerus, instead of *Gonistasteroidocri-*

nus spinigerus. See Ohio Geol., vol. 7, p. 447.

GLYPTASTER milliganae, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 87, Niagara Gr.

GLYPTOCRINUS billingsi, Wachsmuth and Springer, 1897, (Periglyptocrinus billingsi.) N. Am. Crin. Cam., vol. 1, p. 277, Trenton Gr.

mercensis, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 28, Trenton Gr.

typus, Wachsmuth and Springer, 1897, (Tanacrinus typus.) N. Am. Crin. Cam., vol. 1, p. 186, Hud. Riv. Gr.

GONISTASTEROIDOCRINUS faberi, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 84, Keokuk Gr.

lyonnus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 55, Keokuk Gr.

papillatus is figured in Mem. Am. Mus. Nat. Hist., vol. 1, p. 36.

spinigerus, refer to *Gilbertsoerinus spinigerus*.

GRANATOCRINUS leda, refer to *Pentremites leda*.

magnibasis, Rowley, 1895, Am. Geol., vol. 16, p. 220, Burlington Gr.

mutabilis, Rowley, 1893, Am. Geol., vol. 12, p. 306, Chouteau Gr.

sphaeroidalis, Miller and Gurley, 1894,

26, 1892, are un-

washingtonensis, op-
te.

s, Wachsmuth and
onasteroidocrinus

and Gurley, 1895,
Mus. Nat. Hist., p.

greeni, Miller
and Gurley,
1895, Bull. No.
6, Ill. St. Mus.
Nat. Hist., p.
35, Ham. Gr.

spinigerus, instead of Goniatroidocri-
Ohio Geol., vol. 7,

Miller and Gurley,
Ill. St. Mus. Nat.
Gr.

Wachsmuth and
Triglyptocrinus bil-
l. Cam., vol. 1, p.

and Gurley, 1894,
Mus. Nat. Hist., p.

and Springer, 1897,
s.) N. Am. Crin.
Hud. Riv. Gr.

Laberi, Miller and
No. 10, Ill. St. Mus.
Tokuk Gr.

Gurley, 1894, Bull.
Nat. Hist., p. 55

in Mem. Am. Mus.
36.

Albertocrinus spinifer to Pentremites

1895, Am. Geol.
ington Gr.

1893, Am. Geol., vol.
Gr.

and Gurley, 1894

Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 05, Kaskaskia Cr.
winslowi, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 06, Burlington Cr.

GRAPHIOCRINUS may not be an American genus. The species referred by the authors to *Scaphiocrinus* should be retained in that genus.

Heterocidaris Hall, 1861, Desc. New Crin.
Not defined so as to be recognized.

keokuk and *terispina*, Hall. Not defined so as to be recognized.

HEXACRINUS, Austin, 1843, Monogr. Recent and Foss. Crin., p. 48. [Ety., *hex*, six; *crinus*, lily.] Strongly resembles *Platycrinus*. Basals 3, subequal. First radials subquadrangular, and form the sides of the calyx, with a narrow facet for a second radial, which in some cases is axillary, and from which the free arms arise. In other cases there is a third primary radial. Arms ten, and bearing armlets. Vault more or less convex. Type *H. melo*.

leai, instead of *Platyerinus leai*.
occidentalis, Wachsmuth and Springer,
1897, N. Am. Crin. Cam. vol. 2, p. 745,
Ham. Gr.

Holocystites asper, Miller and Gurley,
1895, Bull. No. 7, Ill. St. Mus. Nat.
Hist., p. 84, Niagara Cr.

gyrinus, Miller and Gurley, 1894, Bull.
No. 5, Ill. St. Mus. Nat. Hist., p. 5,
Niagara Cr.

Fig. 1359.—*Holocystites setulus*, side and summit views.

Fig. 1390.—*Holocystites splendens*, left anterior view.

sphaeroidalis, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 85, Niagara Gr.
splendens, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 7, Niagara Gr.

HYDREINOCRINUS
crussidiscus, Mil-
ler and Gurley,
1894, Bull. No. 3,
Ill. St. Mus. Nat.
Hist., p. 43, Up.
Coal Mens.

granuliferus, Miller and Gurley, 1894.

Bull. No. 3, Ill. St. Mus. Nat. Hist., p.
44, Up. Coal Mens.

Fig. 1361 — *Holocystites gyrinus*, ventral view.

noduliferus, Miller and Gurley, 1894,
Bull. No. 3, Ill. St. Mus. Nat. Hist., p.
41, Up. Coal Mens.
sanctiludovici, Worthen, figured in Bull.
No. 3, Ill. St. Mus. Nat. Hist., p. 40.
subsinuatus, Miller and Gurley, 1894,
Bull. No. 3, Ill. St. Mus. Nat. Hist., p.
40, Up. Coal Mens.

Hyptiocrinus, Wachsmuth and Springer, Syn. for *Cyphocrinus*, as I learn from N. Am. Crin. Cam., vol. 1, p. 200. *Cyphocrinus* was published and distributed Sept. 1, 1892, and the American Geologist, which contained the unintelligible definition of *Hyptiocrinus*, was received by me on the 11th day of September, and I suppose, therefore, it was distributed on the 9th at Minneapolis.

ICHTHYOCRINUS clarkensis, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 43, Warsaw Gr.

spinulosus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 44, Keokuk Gr.

Idiocrinus, Wachsmuth and Springer, Syn. for *Gazacrinus*, as I learn from N. Am. Crin. Cam., vol. 1, p. 202. *Gazacrinus* was published and distributed Sept. 1, 1892, and before the Sept. No. of the Am. Geologist, which contained the unintelligible definition of *Idiocrinus* was printed.

elongatus and *I. ventricosus*, Wachsmuth and Springer, Syn. for *Gazacrinus inornatus*.

INDIANOCRINUS, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 83. [Ety. proper name; *krinon*, lily.] Basals 5. No subradials. Primary radials one by four. Arms four. No regular interradials. Azygous interradial rests between the upper sloping sides of two basals, and is followed by two plates at the top of the calyx. Type *I. punctatus*.

punctatus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 83, Niagara Gr.

FIG. 1302.—*Indanoecrinus punctatus*, basal, summit, and lateral views, magnified two diameters.

FIG. 1303.—*Leannocrinus greenel*, azygous side.

LECANOCRINUS greenel, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 52, Niagara Gr.

oswegoensis, Miller and Gurley, 1894, Bull. No. 4, Ill. St. Mus. Nat. Hist., p. 33, Niagara Gr.

LEPIDERTHES wortheni, Jackson, 1896, Bull. Geo. Soc. Am., vol. 7, p. 207, Keokuk Gr.

Lobocrinus, Wachsmuth and Springer, Syn. for *Batoecrinus*. Type *Batoecrinus nashvillei*.

robustus. See *Batoecrinus robustus*.

spiniferus, Wachsmuth and Springer, Syn. for *Batoecrinus marinus*.

MACROSTYLOCYRINUS indianensis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 33, Niagara Gr.

Macrocrinus, Wachsmuth and Springer, Syn. for *Batoecrinus*. Type *Batoecrinus konineki*.

MARSUPIOCRINUS striatus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 732, Niagara Gr.

MEGIOTOCRINUS expansus, Miller and Gur-

ley, 1894, Bull. No. 4, Ill. St. Mus. Nat. Hist., p. 35, Ham. Gr.

hemisphaericus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 44, Ham. Gr.

indianensis, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 27, Ham. Gr.

ornatus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 42, Ham. Gr.

spinulosus, Lyon, instead of *spinulosus*. *whitell* is illustrated in Mem. Am. Mus. Nat. Hist., vol. 1, p. 27.

MELOCRINUS calvinii, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 300, Ham. Gr.

gracilis, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 298, Ham. Gr.

gregeri, Rowley, 1893, Am. Geol., vol. 12, p. 303, Ham. Gr.

lyllyi, Rowley, 1894, Am. Geol., vol. 13, p. 151, Ham. Gr.

oblongus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 300, Niagara Gr.

parvus, Wachsmuth and Springer, Syn. for *M. equalis*.

roemerii, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 301, Niagara Gr.

sampsoni, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 55, Chouteau Gr.

tersus, Rowley, 1894, Am. Geol., vol. 13, p. 152, Ham. Gr.

tiffanyi, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 299, Ham. Gr.

MELONITES giganteus, Jackson, 1896, Bull. Geo. Soc. Am., vol. 7, p. 172, Keokuk Gr.

indianensis, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 5, St. Louis Gr.

septenarius, Whitfield, 1896, Bull. Geo. Soc. Am., vol. 7, p. 182, Warsaw Gr.

MESPILOCRINUS scitulus is described in Bost. Jour. Nat. Hist., vol. 7, p. 321.

MITROCRINUS, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 22. [Ety. *mitra*, a turban; *krinon*, lily.]

Calyx depressed; vault elevated. Basals three, equal. Primary radials three by six. Secondary radials. No

FIG. 1304.—*Mitrocrinus wetherbyi*, basal and summit views.

subradials. Regular interradials three. Azygous interradials three or more,

Ill. St. Mus. Nat. Hist., p. 22.

and Gurley, 1895, Bull. Nat. Hist., p. 42.

and Gurley, 1895, Bull. Nat. Hist., p. 42.

end of spinulosus. Mem. Am. Mus. Nat. Hist., p. 27.

Wachsmuth and Springer, 1897, Bull. Nat. Hist., p. 298.

and Springer, 1897, Bull. Nat. Hist., p. 298.

Am. Geol., vol. 12, p. 301.

Am. Geol., vol. 13, p. 301.

and Springer, 1897, Bull. Nat. Hist., p. 301.

and Springer, 1897, Bull. Nat. Hist., p. 301.

Gurley, 1895, Bull. Nat. Hist., p. 55.

Am. Geol., vol. 13, p. 299.

and Springer, 1897, Bull. Nat. Hist., p. 299.

and Gurley, 1894, Bull. Nat. Hist., p. 321.

and Gurley, 1894, Bull. Nat. Hist., p. 321.

1896, Bull. Geo. Soc. Am., vol. 7, p. 321.

is described in Bull. Geo. Soc. Am., vol. 7, p. 321.

and Gurley, 1894, Bull. Nat. Hist., p. 321.

urban; *krinon*, lily.] Bull. Nat. Hist., p. 321.

Primary radials. No secondary radials. No

terherbyl, basal and

ews.

interradials three, s three or more,

the first one resting on a single basal. Type, *M. wetherbyi*.

wetherbyi, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 22. Trenton Gr.

MYELODACTYLUS possessed long side plates extending from the outer rim and gradually tapering to the center, where they formed a funnel-shaped cavity resembling the closed umbilicus of a Goniatite. Mr. J. S. Fig. 1395.—*Myelodactylus gorbbyi*, side and anterior views. Side plates removed.

Indiana, has a specimen of *M. gorbbyi*, with the side plates in position. The genus has no near relation to a erinoid. Probably, it is nearer *Cycloegypoides* than to any other genus of the *Echinodermata*.

XYLOCHORIS greenii, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 62, Up. Held. Gr.

venustus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 63, Up. Held. Gr.

OLIGOPORUS bellulus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 7, Keokuk Gr.

blairi, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 6, Keokuk Gr.

missouriensis, Jackson, 1896, Bull. Geo. Soc. Am., vol. 7, p. 184, Keokuk Gr.

sulcatius, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 8, St. Louis Gr.

Fig. 1396.—*Onychaster asper*, large specimen showing outer integument.

OXYCHORIS asteriiformis was described in Bost. Jour. Nat. Hist., vol. 7, p. 320.

norwoodi is not a Syn. for *O. exculptus* and is from the Keokuk Gr.

puryus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 52, Kaskaskia Gr.

pulaskiensis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 40, Kaskaskia Gr.

PALEASTER wykoffi, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 46, Hud. Riv. Gr.

PALECHINUS elegans is the type of the genus.

Pentatrematiles, Roemer, is a Syn. for *Pentremites*. Roemer simply attempted to correct the word, which was derived from *penta*, five; and *trema*, foramen. *PENTREMITES* ledn, instead of *Grannatoerinus ledn*.

lycorinus was described in the 15th Rep. N. Y. St. Mus. Nat. Hist.

sampsoni, Hambach, Syn. for *Grannatoerinus roemeri*.

Periglyptocrinus, Wachsmuth and Springer, Syn. for *Glyptocrinus*.

billingsi, Wachsmuth and Spring. See *Glyptocrinus billingsi*.

PETALOCRINUS, Weller, 1890, Journal of Geology, vol. 4, p. 166. The genus is valid, but the definition is so doubtful that, without seeing specimens, I am unwilling to undertake to give the generic characters. Type *P. mirabilis*, described, at the same place, from the Niagara Gr., and *P. major* proposed for a fragment.

Phidocrinus barydactylus, Keyes, 1895, Geo. Sur. Mo., vol. 4, p. 220. Not intelligently defined.

PHOLIOCRINUS meeki, Jackson, 1896, Bull. Geo. Soc. Am., vol. 7, p. 210, Keokuk Gr.

PHYSETOCRINUS lobatus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 590, Burlington Gr.

sampsoni, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 14, Burlington Gr.

Fig. 1397.—*Physetoerinus sampsoni*, azygous and opposite views.

PISOERINUS baccula, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 79, Niagara Gr.

Fig. 1398.—*PISOERINUS baccula*, internal, basal, and summit views.

milligame, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 80, Niagara Gr.

PLATYCRINUS agassizi, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 669, Kinderhook Gr.

bozemanensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 42, Keokuk Gr.

Fig. 1399.—*PISOERINUS milligame*, lateral view of a perfect specimen without sutures, basal view of same magnified two diameters.

Fig. 1399.—*PISOERINUS milligame*, basal, side, and summit views, showing sutures of the calyx.

IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N. Y. 14580
(716) 872-4503

1.8
2.0
2.2
2.5
2.8
3.2
3.6
4.0
4.5
5.0
5.6
6.3
7.1
8.0
9.0
10.0
11.2
12.5
14.0
16.0
18.0
20.0
22.5
25.0
28.0
31.5
36.0
40.0
45.0
50.0
56.0
63.0
71.0
80.0
90.0
100.0

1.8
2.0
2.2
2.5
2.8
3.2
3.6
4.0
4.5
5.0
5.6
6.3
7.1
8.0
9.0
10.0
11.2
12.5
14.0
16.0
18.0
20.0
22.5
25.0
28.0
31.5
36.0
40.0
45.0
50.0
56.0
63.0
71.0
80.0
90.0
100.0

bridgerensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 43, Keokuk Gr.

Fig. 1371.—*Platyterinus aequifurcus*.

Fig. 1372.—*Platyterinus allophyllus*.

ensula, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 76, Burlington Gr.

Fig. 1373.—*Platyterinus annosus*.

Fig. 1374.—*Platyterinus blairi*.

elimatus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 74, Chouteau Gr.

Fig. 1375.—*Platyterinus ensula*, basal, azygous and opposite views.

concinclus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 63, Burlington Gr.

constritus, refer to the Keokuk Gr.

cortina, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 40, Chouteau Gr.

davisi, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 684, Burlington Gr.

doughlassi, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 86, Burlington (?) Gr.

elegans is illustrated in Mem. Am. Mus. Nat. Hist., vol. 1, p. 3.

excavatus is illustrated in Mem. Am. Mus. Nat. Hist., vol. 1, p. 3.

formosus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 72, Burlington Gr.

formosus var. *approximatus*, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 60, Burlington Gr.

geometricus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 697, Burlington Gr.

germanus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 78, Chouteau Gr.

graphicus, refer to the Keokuk Gr.

hodgsoni, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 63, Burlington Gr.

huntville, Wachsmuth and Springer, Syn. for *P. alabamensis*.

illinoisensis, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 62, Burlington Gr.

inornatus, McChesney, 1859, Desc. New Spec. Foss., p. 6, Burlington Gr., is not a Syn. for *P. burlingtonensis*.

missouriensis, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 73, Chouteau Gr.

modestus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 77, Burlington Gr.

nodostratus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 698, Burlington Gr.

perasper, Meek and Worthen. Not recognized.

peculiaris, Wachsmuth and Springer, N. Am. Crin. Cam., vol. 2, p. 700, Burlington Gr.

pettisensis, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 73, Chouteau Gr.

prattensis is on page 569, Trans. St. Louis Acad. Sci., vol. 1,

richfieldensis, refer to the Keokuk Gr.

semifusus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 77, Burlington Gr.

sharonensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 42, Burlington Gr.

spinifer, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 708, Burlington Gr.

spinifer, var. *elongatus*, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 709, Burlington Gr.

striobrachiatus is illustrated in Mem. Am. Mus. Nat. Hist., vol. 1, p. 4.

subscitulus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 62, Burlington Gr.

suleiferus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 75, Burlington Gr.

tugurium, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 70, Burlington Gr.

Fig. 1377.—*Platyterinus tugurium*, basal and azygous views.

vaseellum, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 32, Keokuk Gr.

Gurley, 1896, Bull.
Nat. Hist., p. 63.

h and Springer.
osis.
and Gurley, 1896.
Mus. Nat. Hist.,

, 1859, Desc. New
Burlington Gr., is not
tonensis.

and Gurley, 1895.
Mus. Nat. Hist.,

Gurley, 1895, Bull.
Nat. Hist., p. 77.

uth and Springer.
Cam., vol. 1, p. 698.

Worthen. Not rec-

h and Springer, N.
2, p. 700, Burling-

and Gurley, 1895,
Mus. Nat. Hist.,

39, Trans. St. Louis

o the Keokuk Gr.
Gurley, 1895, Bull.

s Nat. Hist., p. 77.

and Gurley, 1897,
St. Mus. Nat. Hist.,

and Springer, 1897,
vol. 2, p. 708, Bur-

gatus, Wachsmuth
N. Am. Crin. Cam.,

ington Gr.
illustrated in Mem.

st., vol. 1, p. 4.
and Gurley, 1896,

Mus. Nat. Hist., p.

l Gurley, 1895, Bull.
s. Nat. Hist., p. 75,

l Gurley, 1895, Bull.
s. Nat. Hist., p. 70,

Fig. 1379.—*Platyocrinus vaseletum*, basal and internal views.

and Gurley, 1895,
Mus. Nat. Hist., p.

verrucosus, illustrated. See N. Am.
Crin. Cam., vol. 2, p. 705.

Fig. 1378.—*Platyocrinus vaseletum*, basal, internal, and summit views.

Fig. 1379.—*Pleurocystites mercerensis*, dorsal and basal views.

5, Ill. St. Mus. Nat. Hist.,
p. 24, Trenton Gr.

POTERIOCRINUS alberst, Miller
and Gurley, 1896, Bull.

No. 9, Ill. St. Mus. Nat.
Hist., p. 29, Kaskaskia Gr.

altonensis, Miller and Gurley,
1895, Bull. No. 7, Ill.

St. Mus. Nat. Hist., p. 62,
St. Louis Gr.

arrectarius, Miller and Gurley,
1896, Bull. No. 9, Ill.

St. Mus. Nat. Hist., p. 33, St. Louis Gr.

Fig. 1380.—*Poterocrinus kentuckiensis*, lateral view.

Fig. 1381.—*Poterocrinus alberst*, azygous and opposite views.

Fig. 1382.—*Poterocrinus blairi*, azygous and opposite views.

blairi, Miller and Gurley, 1895, Bull. No.
7, Ill. St. Mus. Nat. Hist., p. 61, Bur-
lington Gr.

Fig. 1384.—*Poterocrinus altonensis*, azygous and opposite views.

Fig. 1385.—*Poterocrinus altonensis*, view opposite azygous side.

bozemmen-

sis, Miller

and Gur-

ley, 1896,

Bull. No.

10, Ill. St.

Mus. Nat.

Hist. p. 82,

Burling-

ton (?) Gr.

broadheadi,

Miller and

Gurley,

1895, Bull.

No. 7, Ill.

St. Mus.

Fig. 1386.—*Poterocrinus circumtextus*, azygous and opposite views.

Nat. Hist., p. 63, Chou-
teau Gr.

circumtextus, Miller and
Gurley, 1894, Bull. No.

5, Ill. St. Mus. Nat.
Hist., p. 31, Keokuk Gr.

Fig. 1387.—*Poterocrinus labyrinthicus*, view of two specimens.

corycia, refer to Keokuk
Gr.

crineus, refer to Keokuk
Gr.

doughlassi, Miller and Gurley, 1896, Bull.

Fig. 1388.—*Poterocrinus corycia*, refer to Keokuk Gr.

Fig. 1389.—*Poterocrinus crineus*, refer to Keokuk Gr.

Fig. 1390.—*Poterocrinus doughlassi*, Miller and Gurley, 1896, Bull.

Fig. 1391.—*Poterocrinus corycia*, refer to Keokuk Gr.

Fig. 1392.—*Poterocrinus crineus*, refer to Keokuk Gr.

Fig. 1393.—*Poterocrinus doughlassi*, Miller and Gurley, 1896, Bull.

No. 10, Ill. St. Mus. Nat. Hist., p. 83, Burlington (?) Gr.
hammondi, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 35, Kinderhook Gr.
labyrinthicus, S. A. Miller, 1891, (Cynthocrinus labyrinthicus,) Adv. sheets 17th Rep. Geo. Sur. Ind., p. 48 and final Report, p. 659, Keokuk Gr.

Fig. 1388.—*Poteriocrinus lautus*, azygous and opposite views.

lautus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 30, Keokuk Gr.
maccabei, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 34, Kinderhook Gr.
maccabei var. *decrepitus*, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 36, Kinderhook Gr.
neglectus, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 31, Keokuk Gr.
pleias, refer to the Keokuk Gr.
pulaskiensis, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 39, Kaskaskia Gr.
sampsoni, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 65, Chouteau Gr.
vagus, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 46, Kaskaskia Gr.
Prochivocrinus, Ringueberg, Syn. for *Calceocrinus*.

Fig. 1389.—*Pterotocrinus wetherbyi*, azygous and basal views.

PTEROTOCRINUS wetherbyi, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 44, Kaskaskia Gr.

Fig. 1390.—*Reticocrinus alveolatus*, azygous side.
RETICOCRINUS alveolatus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 26, Trenton Gr.

RHODOCRINUS barrisi var. *striatus*, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 231, Burlington Gr.

blairi, Miller and Gurley, 1896, Bull. No. 9, Ill. St. Mus. Nat. Hist., p. 37, Chouteau Gr.

bozemanensis, Miller and Gurley, 1897,

Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 40, Keokuk Gr.

Fig. 1391.—*Rhodocrinus blairi*, basal, azygous, and opposite views.

bridgerensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 41, Keokuk Gr.

douglassi, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 39, Keokuk Gr.

truncatus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 231, Burlington Gr.

tuberculatus, Wachsmuth & Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 232, Burlington Gr.

vesperalis, White. This species was described from the Upper Coal Measures west of Humboldt, Kansas. It is now asserted that it came from the Trenton Group of Tennessee, and is an *Archæocrinus*. If so, the definition was wholly erroneous, and the specific name must be disregarded.

wortheni var. *urceolatus*, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 221, Burlington Gr.

Saccocrinus amplus, refer to *Actinocrinus amplus*.

umbrosus, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist. p. 24, Niagara Gr.

SAMPSONOCRINUS, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 51. [Ety. proper name; *krinon*, lily.] Body globose. Basals three, one pentagonal, two hexagonal. Primary radials, two or three in each radial series, and differing, in this respect, in the same specimen. Secondary and tertiary radials. Interradials connect with the vault, and in three of the areas the first plate abuts upon the basals, by which arrangement eight plates unite with the basal disk. Proboscis subcentral. Type *S. hemisphericus*.

Fig. 1393.—*Sampsonocrinus hemisphericus*, basal and summit views.

Mus. Nat. Hist., p.

Miller and Gurley, 1897,
Mus. Nat. Hist.,
Gurley, 1897, Bull.
Nat. Hist., p. 39,

1892.—*Rhodocrinus*
crinus, azygous and op-
posite views.
Burlington Gr.

This species was de-
scribed in the Trenton
Coal Measures
of Kansas. It is now
known from the Trenton
formation, and is an Archæo-
zoic definition was
given, and the specific
name regarded.

actinocrinus, Wachsmuth
& Springer, 1897, N. Am. Crin. Cam.,
vol. 1, p. 39, Burlington Gr.

Miller and Gurley, 1895, Bull.
Nat. Hist., p. 24,

Miller and Gurley, 1895,
Mus. Nat. Hist., p.
name; *krinon*, lily.]
Basals three, one pen-
gonal. Primary ra-
dials in each radial series,
this respect, in the
Secondary and ter-
tiary radials connect
in three of the areas
upon the basals, by
eight plates unite
the Proboscis subcen-
tralis.

Miller and Gurley, 1897,
Mus. Nat. Hist.,
Gurley, 1897, Bull.
Nat. Hist., p. 39,

hemisphericus, Miller and Gurley, 1895,
Bull. No. 7, Ill. St. Mus. Nat. Hist.,
p. 51, Chouteau Gr.

Fig. 1894.—*Sainpsonocrinus hemisphericus*, azygous and lateral views.

SCAPHIOCRINUS. This genus must be re-
stored, as originally defined by Hall,
with *S. simplex* as the type. The spe-
cies described by Worthen as *Potero-
crinus*, and referred to *Scaphiocrinus* in
the body of this work, may be restored
to *Poteroocrinus*. The work of Wach-
smuth on this genus may be wholly set
aside.

regina, refer to the Keokuk Gr.

arrosus, Miller and Gurley, 1894, Bull.
No. 3, Ill. St. Mus. Nat. Hist., p. 39,
Keokuk Gr.

carinatus, is illustrated in Mem. Am.
Mus. Nat. Hist., vol. 1, p. 31.

lyrioepes, refer to the Keokuk Gr.

martinensis, Miller and Gurley, 1894,
Bull. No. 3, Ill. St. Mus. Nat. Hist., p.
38, Kaskaskia Gr.

notatus, Miller and Gurley, 1896, Bull.
No. 9, Ill. St. Mus. Nat. Hist., p. 34,
Kinderhook Gr.

subcarinatus, refer to the Keokuk Gr.

subtortuosus, refer to the Keokuk Gr.

tortuosus is illustrated in Mem. Am.
Mus. Nat. Hist., vol. 1, p. 32.

SHUMARDOCRINUS. Miller and Gurley, 1895,
Bull. No. 7, Ill. St. Mus. Nat. Hist., p.
40. [Ety. proper name; *krinon*, lily.]
Basals three. Radials two by five.
Regular interradials one. Azygous
interradials three. Interradial areas
connect with the vault. No proboscis.
Orifice subcentral. Type *S. concinnus*.
concinnus, Shumard, 1855, (Actinocrinus
concinnus), Geo. Sur. Mo., p. 189, Bur-
lington Gr.

SIPHONOCRINUS pentagonus, Wachsmuth
and Springer, 1897, N. Am. Crin. Cam.,
vol. 1, p. 213, Niagara Gr.

STEGANOCRINUS albersi, Miller and Gurley,
1897, Bull. No. 12, Ill. St. Mus. Nat.
Hist., p. 33, Burlington Gr.

blairi, Miller and Gurley, 1897, Bull. No.
12, Ill. St. Mus. Nat. Hist., p. 35,
Burlington Gr.

globosus, Wachsmuth and
Springer, 1897, N. Am. Crin. Cam.,
vol. 2, p. 585, Kinderhook Gr.

griffithi, Miller and Gurley, 1897, Bull.

Fig. 1895.—*Stephanocrinus hammetti*, summit and side views.

No. 12, Ill. St. Mus. Nat. Hist., p. 34,
Burlington Gr.

Fig. 1896.—*Stephanocrinus osgoodensis*, side and summit views.

sharonensis, Miller and Gurley, 1897,
Bull. No. 12, Ill. St. Mus. Nat. Hist.,
p. 32, Burlington Gr.

spargenensis, Miller and Gurley, 1896,
Bull. No. 8, Ill. St. Mus. Nat. Hist., p.
29, St. Louis Gr.

validus, Meek and Worthen, 1860, (Ac-
tinocrinus *validus*), Proc. Acad. Nat.
Sci., p. 384, and Geo. Sur. Ill., vol. 2, p.
200, Burlington Gr.

STEREOCRINUS barrisi, Wachsmuth and
Springer, 1897, N. Am. Crin. Cam., vol.
1, p. 326, Ham. Gr.

indianensis, Miller and Gurley, 1897,
Bull. No. 12, Ill. St. Mus. Nat. Hist., p.
38, Ham. Gr.

Fig. 1897.—*Stribalocystites sphaeroidalis*, anterior, basal, and summit views.

STRIBALOCYSTITES sphaeroidalis, Miller and
Gurley, 1895, Bull. No. 6, Ill. St. Mus.
Nat. Hist., p. 58, Niagara Gr.

STROTOCRINUS agilops is illustrated in
the Mem. Am. Mus. Nat. Hist., vol. 1,
p. 20.

blairi, Miller and Gurley, 1895, Bull. No.
7, Ill. St. Mus. Nat. Hist., p. 48, Bur-
lington Gr.

ornatus, Miller and Gurley, 1896, Bull.
No. 8, Ill. St. Mus.

Nat. Hist., p. 30, Bur-
lington Gr.

regalis is illustrated in
Geo. Sur. Ill., vol. 5,
p. 357.

venustus, Miller and
Gurley, 1894, Bull.
No. 3, Ill. St. Mus.

Nat. Hist., p. 26, Bur-
lington Gr.

SYNBATHOCRINUS angu-
laris, Miller and Gur-
ley, 1894, Bull. No. 5,
Ill. St. Mus. Nat. Hist.,
p. 42, Keokuk Gr.

illinoisensis, Miller &
Gurley, 1896, Bull.
No. 8, Ill. St. Mus.

Nat. Hist., p. 53, Bur-
lington Gr.

wachsmuthi, Meek and Worthen, 1860,

Fig. 1898.—*Synbathocrinus blairi*, opposite and azygous views.

- Proc. Acad. Nat. Sci., p. 67, and Geo. Sur. Ill., vol. 5, p. 437, Burlington Gr.
- TALAROOCRINUS decornis**, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 788, St. Louis Gr.
- patel, Miller and Gurley 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 44, Kaskaskia Gr.
- subglobosus, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 2, p. 789, St. Louis Gr.
- trijugis, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 45, St. Louis Gr.
- Tanaocrinus**, Wachsmuth and Springer, Syn. for Glyptocrinus.
- typus. See Glyptocrinus typus.
- TAXOOCRINUS colletti**, White, 1881, (T. multibrachiatus var. colletti,) is a good species.
- concavus, Rowley, 1893, Am. Geol., vol. 12, p. 304, and vol. 13, p. 153, Ham. Gr.
- crawfordsvillensis, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 49, Keokuk Gr.
- juvenis is illustrated in Mem. Am. Mus. Nat. Hist., vol. 1, p. 35.
- splendens, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 61, Keokuk Gr.
- ungula, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 59, Keokuk Gr.

Fig. 1399.—*Taxocrinus wetherbyi*, basal view, azygous side down.

wetherbyi, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 41, Kaskaskia Gr.

THALAMOOCRINUS, Miller and Gurley, 1895,

Fig. 1400.—*Taxocrinus wetherbyi*, lateral view.

Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 81. [Ety. *thalamos*, a small house; *krinon*, lily.] Body pear-

shaped or fusiform, and covered by three ranges of plates and a small vault. First circle, five plates, equal. Second circle, five plates. Third circle, six plates. Type *T. ovatus*.

cylindricus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 82, Niagara Gr.

ovatus, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 81, Niagara Gr.

Fig. 1402.—*Thalamocrinus ovatus*, lateral, azygous, and summit views, magnified two diameters.

Teleocrinus, Syn. for Strotocrinus.

adolescens. See Actinocrinus adolescens.

THYLACOCRINUS, Oehlert, 1878, Bull. Soc. Geol. de France, Tome VIII, p. 6. [Ety. *thylakos*, a bag; *krinon*, lily.] The French Devonian species are very large, and the calyx, as high as the secondary radials, seems to be constructed as in Rhodocrinus. Above this there are intersecondary and intertertiary radials, and twenty or more large arms. Type *T. ramuloti*.

clarkei, Wachsmuth and Springer, 1897, N. Am. Crin. Cam., vol. 1, p. 248, Ham. Gr.

Fig. 1403.—*Thysanocrinus milligane*, basal, azygous, and opposite side views.

THYSANOOCRINUS milligane, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 51, Niagara Gr.

ULOOCRINUS blairi, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 57, Up. Coal Meas.

occidentalis, Miller & Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 58, Up. Coal Meas.

ZEACRINUS bellulus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 34, Kaskaskia Gr.

Fig. 1404.—*Zeacrinus grandiculus*, azygous side.

blairi, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 66, Keokuk Gr.

cylindricus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 38, Kaskaskia Gr.

doverensis, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 35, Kaskaskia Gr.

, and covered by plates and a small five plates, equal plates. Third circle *T. oratus*. and Gurley, 1895, Mus. Nat. Hist., p.

2.—*Thalamocrinus* us, lateral, azygous, summit views, magnified diameters.

protoerinus. *erinus* *adolescens*. rt, 1878, Bull. Soc. me VIII, p. 6. [Ety. *krinon*, lily.] The species are very large, as the secondary be constructed as in ove this there are d intertertiary rator more large arms.

and Springer, 1897, vol. 1, p. 248, Ham.

milligane, basal, azygous side views.

que, Miller and Gur-

at.

ra

er

4,

St.

p.

&

ll.

s.

p.

l-

r-

ll,

St.

Fig. 1404.—*Zeacrinus*

grandiculus, azygous side.

and Gurley, 1895, Bull.

Mus. Nat. Hist., p.

er.

Miller and Gurley,

p. 5, Ill. St. Mus. Nat.

Kaskaskia Gr.

ler and Gurley, 1896,

Ill. St. Mus. Nat. Hist.,

Kaskaskia Gr.

durabilis, Miller and Gurley, 1895, Bull. No. 6, Ill. St. Mus. Nat. Hist., p. 48, Kaskaskia Gr.

grandiculus, Miller and Gurley, 1894, Bull. No. 5, Ill. St. Mus. Nat. Hist., p. 32, Kaskaskia Gr.

kentuckiensis, Miller and Gurley, 1896, Bull. No. 8, Ill. St. Mus. Nat. Hist., p. 57, Kaskaskia Gr.

maniformis is figured in Geo. Sur. Iowa, p. 682.

merope, refer to the Keokuk Gr.

moorei is reproduced in Ohio Geol., vol. 7, p. 483.

nitidus, Miller and Gurley, 1894, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 36, Kaskaskia Gr.

obesus, Miller and Gurley, 1894, Bull.

No. 3, Ill. St. Mus. Nat. Hist., p. 35, Kaskaskia Gr.

paternus, refer to the Keokuk Gr.

peculiaris, Miller and Gurley, 1896, Bull.

No. 8, Ill. St. Mus. Nat.

Hist., p. 34, Kaskaskia

Gr.

pulaskiensis, Miller and

Gurley, 1895, Bull. No.

5, Ill. St. Mus. Nat.

Hist., p. 47, Kaskaskia

Gr.

salemensis, Miller and

Gurley, 1894, Bull. No.

5, Ill. St. Mus. Nat.

Hist., p. 37, Keokuk Gr.

scoparius is figured in

Mem. Am. Mus. Nat. Hist., vol. 1, p. 34.

Fig. 1405.—*Zeacrinus pulaskiensis*, azygous and opposite views

SUBKINGDOM MOLLUSCOIDA.

CLASS BRYOZOA.

SEVERAL genera included among the Bryozoa, in the original text of this work, belong to the polyp corals. See remarks under Coelenterata. In 1889, when the work was in part printed, and most of the article on the Bryozoa was in the galleys, but not yet made up into pages, part of the printed pages of Vol. VIII of the Illinois Geo. Sur. was presented to me, and, on the representation that the volume would be published at about the same time that my work would be, I was induced to open up the galleys and insert the Bryozoa from the Illinois Report. This caused several genera, by oversight, to appear in different families, and practically destroyed the family arrangement, and it, probably, also led to the accidental placing of the words "Subkingdom Mollusca" over the word Brachiopoda. While, therefore, I published the species from the Illinois Report as "in press," the fact is, that the volume was not published for a year or more after my work was published. Such family names as *Amplexoporidae*, *Batostomellidae*, *Rhabdomesontidae*, etc., must be stricken out; but as so many genera of polyp corals are printed with the Bryozoa, I will not, at this time, undertake a family arrangement of the Class, which, at best, could only be approximately correct.

Microscopic sections, in the hands of a scholar, are no doubt of great service in exposing the life history of the Bryozoa; but there are some who can make two or three genera and five or six species out of a single specimen, and it will not do to place much confidence in their microscopic work.

Amplexopora is not a valid genus and *affinis*, *pustulosa*, *septosa*, *superba*, and *winchelli* may be referred to *Monticulipora*; and *canadensis*, *cingulata*, and its synonym *robusta*, may be referred to *Batostomella*, if they are valid species. *Anototichia*, Ulrich, 1890, Geo. Sur. Ill., vol.

8, p. 381. Too poorly defined to be recognized. The species *impolita* and *ponderosa* are also too poorly defined to be recognized.

Archimedes grandis, Ulrich, Syn. for *A. wortheni*.

negligens, Ulrich, Syn. for *A. owenianus*.

- ARTHROCLEMA striatum** Ulrich, 1883, Geo. Sur. Minn. vol. 3, p. 198, Trenton Gr.
- ARTHIOPORA bifurcata**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 178, Trenton and Galena Gr.
- reversa**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 178, Trenton Gr.
- Aspidopora**, Ulrich, Syn for Prasopora.
- elegantula**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 258, Galena Gr., refer to Prasopora elegantula.
- ASTROPORITES**, Lambe, 1896, Can. Jour. Sci., *A. ottawensis* is described as the type from the Trenton Gr.
- Atactopora septosa** is a Monticulipora.
- ATACTOPORELLA crassa**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 225, Galena Gr.
- insueta**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 224, Trenton Gr.
- ortoni**, refer to Monticulipora ortoni.
- ramosa**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 226, Trenton Gr.
- BAROSTOMA decipiens**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 298, Trenton Gr.
- humile**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 294, Galena Gr.
- magnopora**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 291, Trenton Gr.
- minnesotense**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 297, Trenton Gr.
- montuosum**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 293, Trenton Gr.
- varium**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 292, Trenton Gr.
- Buscopora**, Ulrich, Syn. for Fistulipora.
- BYTHOPORA aleicornis**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 264, Trenton Gr.
- Bythotrypa**, Ulrich. Too poorly defined to be recognized.
- CALLOPORA ampla**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 281, Trenton Gr.
- angularis**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 277, Trenton Gr.
- cincinnatiensis**, Ulrich, Syn. for Fistulipora occidentalis.
- crenulata**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 284, Trenton Gr.
- dumalis**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 282, Trenton Gr.
- goodluensis**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 282, Galena Gr.
- minutissima**, instead of Leioclema minutissimum.
- onealli**, James, Syn. for Callopora sigillarioides.
- pulchella**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 283, Trenton Gr.
- punctata**, instead of Leioclema punctatum.
- Calloporella**, Ulrich, Syn. for Prasopora.
- nodulosa**, Ulrich. Too poorly defined to be recognized.
- CERAMOPHYLLA**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 331. No generic characters are given by which it can be distinguished. Type *C. frondosa*.
- frondosa**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 331, Trenton Gr.
- CERAMOPORA inclusa**, Ulrich, 1893, (Ceramoporella,) Geo. Sur. Minn., vol. 3, p. 329, Trenton Gr.
- interporosa**, Ulrich, 1893, (Ceramoporella,) Geo. Sur. Minn., vol. 3, p. 330, Hud. Riv. Gr.
- Ceramoporella**, Ulrich, Syn. for Ceramopora.
- CREPIOPORA perampla**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 323, Trenton Gr.
- spatiosa**, Ulrich. Too poorly defined to be recognized.
- subequata**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 322, Trenton Gr.
- Cyatodictya**, Ulrich, seems to be a synonym for Stictopora.
- DIAMESOPORA trentonensis**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 330, Trenton Gr.
- Discotrypa**, Ulrich, Syn. for Prasopora.
- Eridopora minima**, Ulrich, Syn. for Pileotrypa denticulata.
- ERIDOTRYPA**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 264. [Ety. *eridos*, in dispute., *trypa* a perforation.] Zoonia ramose, branches slender. Zoonia more or less oblique, with thick walls, the tubes intersected by diaphragms only. The latter may be wanting in the axial region, are in most cases absent for a short distance within the apertural edge, but always present and closest together in the turn from the axial into the narrow peripheral region. Mesopores with close-set diaphragms, varying in number; sometimes abundant, at other times very few. Acanthopores small, never numerous, sometimes wanting. Type *E. mutabilis*.
- exigua**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 266, Galena Gr.
- mutabilis**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 265, Galena Gr.
- ESCHAROPORA angularis**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 168, Trenton Gr.
- confluens**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 171, Trenton Gr.
- limitaris**, Ulrich, 1893, Geo. Sur. Minn., vol. 3, p. 172, Trenton Gr.
- EVACTINOPORA radiata** is from the Keokuk Gr.
- FENESTELLA**. The nonporiferous side of the branches is a basal plate on which the cells take their origin, and is usually longitudinally striated. There is no axis. There are two rows of cells on the branches, on each side of a high keel. Sometimes there are accessory pores on the keel. There are no pores on the dissepiments. Species are distinguished by the size of the branches and of the fenestrules, and from the arrangement of the cell openings. Age thickens the basal plate, and sometimes closes part of the pores, and there is great difference in the appearance in different states of preservation.
- bigeneris**, Ulrich, Syn. for *F. perplexa*.

ich, 1893, (Ceram-
Minn., vol. 3, p.

1893, (Ceramopo-
Minn., vol. 3, p. 330,

Syn. for Ceramo-

Ulrich, 1893, Geo.
323, Trenton Gr.
poorly defined to

1893, Geo. Sur.
Trenton Gr.

s to be a synonym

sis, Ulrich, 1893,
3, p. 330, Trenton

for Prasopora.

ch, Syn. for Pileo-

3, Geo. Sur. Minn;
eridas, in dispute.,

] Zonaria ramose,
occeia more or less
walls, the tubes in-
fragments only. The
ing in the axial re-
cesses absent for a
thin the apertural
present and closest
en from the axial
peripheral region.
se-set diaphragms,
sometimes abun-
s very few. Acan-
er numerous, some-
e *E. mutabilis*.

(Geo. Sur. Minn.,
Gr.

3, Geo. Sur. Minn.,
Gr.

Ulrich, 1893, Geo.
p. 168, Trenton Gr.

3, Geo. Sur. Minn.,
on Gr.

3, Geo. Sur. Minn.,
on Gr.

from the Keokuk

poriferous side of
sal plate on which
origin, and is usu-
striated. There
are two rows of
es, on each side of
ometimes there are
the keel. There
the dissepiments.
nished by the size
of the fenestrules,
gement of the cell
thickens the basal
s closes part of the
great difference in
different states of

for *F. perplexa*.

patellifera, Ulrich, Syn. for *F. variopora*.
sculptilis, Ulrich, Syn. for *F. stellata*.

Fistulipora normalis, Ulrich, Syn. for
Lichenalia substellata.

oweni, James, is probably a *Dianoropora*,
and may be so referred.

Glyptopora megastoma, Ulrich, Syn. for *Cos-
cinium keyserlingi*.

HELOPORA elegans, Ulrich, 1893, Geo. Sur.
Minn., vol. 3, p. 194, Hud. Riv. Gr.

harrisi, James, 1893, Geo. Sur. Minn.,
vol. 3, p. 195, Hud. Riv. Gr.

quadrata, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 193, Galena Gr.

Hemiphragma, Ulrich. Syn. for *Bato-
stoma*.

tenuimurale, Ulrich, Syn. for *Batostoma
irrasum*.

HOMOTRYPA callosa, Ulrich, 1893, Geo. Sur.
Minn., vol. 3, p. 243, Galena Gr.

insignis, Ulrich, Syn. for *H. subramosa*.
intercalaris, Ulrich, 1893, Geo. Sur. Minn.,

vol. 3, p. 238, Trenton Gr.

separata, Ulrich, Syn. for *H. minnesoten-
sis*.

similis, Foord, 1883, Cont. Micro Pal.
Cambro. Sil. Rocks, p. 10, Trenton Gr.

tuberculata, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 240, Trenton Gr.

HOMOTRYPELLA multiporata, Ulrich, 1893,
Geo. Sur. Minn., vol. 3, p. 230, Trenton

Gr.
mundula, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 232, Galena Gr.

ovata, Ulrich, 1893, Geo. Sur. Minn., vol.
3, p. 231, Galena Gr.

rustica, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 234, Hud. Riv. Gr.

subgracilis, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 230, Trenton Gr.

LEPTOTRYPA acervulosa, Ulrich, 1893, Geo.
Sur. Minn., vol. 3, p. 318, Galena Gr.

claviformis, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 319, Trenton Gr.

informis, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 317, Trenton Gr.

Leioclema, Ulrich, Syn. for *Callopora*. The

principal character is the presence of

"spineform corallites" or "acantho-
pores," which, as Waagen has shown,

are merely the newly-developed gems
in the intermural development or frag-

ments of the secondary walls.

Leioclema, Foerste. Not defined so as
to be recognized as a genus and *L. Ohio-*

ensis is too poorly defined to be recog-
nized as a species.

Lichenotrypa cavernosa, Ulrich, Syn. for *L.
longispin*.

Lyropora oralis, Ulrich, Syn. for *L. quin-
cuncialis*.

ranosculum, Ulrich, Syn. for *L. lyra*.
Mesotrypa, Ulrich, Syn. for *Dianulites*.

spinosa, Ulrich, Syn. for *Prasopora para-
sitica*.

Nematopora quadrata, Ulrich, Syn. for *N.
ovalis*.

NICHOLSONELLA laminata, Ulrich, 1893, Geo.
Sur. Minn., vol. 3, p. 315, Trenton Gr.

pulchra, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 314, Trenton Gr.

PACHYDICTYA elegans, Ulrich, 1893, Geo.
Sur. Minn., vol. 3, p. 154, Galena Gr.

PHLEOPORA incipiens, Ulrich, 1893, Geo.
Sur. Minn., vol. 3, p. 174, Trenton Gr.

wilmingtonensis, Ulrich, 1893, Geo. Sur.
Minn., vol. 3, p. 175, Hud. Riv. Gr.

PHYLLODICTYA varia, Ulrich, 1893, Geo. Sur.
Minn., vol. 3, p. 144, Trenton Gr.

PHYLLOPORA. The basal plate on the non-
poriferous side is composed of fine

capillary tubes and is longitudinally
striated or granulated. The cells as-

cend from their origin a short distance
along the basal plate and then bend

directly outward. The branches are
anastomosing.

Pinacotrypa, Ulrich, Syn. for *Fistulipora*.

POLYPORA. The nonporiferous side is oc-
cupied by the basal plate, composed of

one or more layers of capillary fibers,
and is longitudinally striated. On the

obverse side, mostly the inner one,
there are from three to ten longitu-

dinal rows of pores; they are round,
and usually the margin is elevated. No

keel. The cells originate at the axis
of the branches, and rise in an oblique

direction upward and outward. Dis-
sepiments without cells.

simulatrix, Ulrich, Syn. for *P. hallana*.

PROBOSCINA tumulosa, Ulrich, Syn. for *Sto-
matopora frondosa*.

PTILODICTYA whitfieldi, Foerste, Syn. for
Phenopora expansa.

RHINDICTYA grandis, Ulrich, 1893, Geo.
Sur. Minn., vol. 3, p. 136, Trenton Gr.

humilis, Ulrich, Syn. for *Pachydietya
pumula*.

neglecta, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 130, Galena Gr.

pediculata, Ulrich, 1893, Geo. Sur. Minn.,
vol. 3, p. 137, Trenton Gr.

Rhabdomesontida, Vine, Syn. for *Petalopo-
ridae*, Waagen, which includes *Rhomi-
bopora*.

RHOMBOPORA. Waagen says the little
granules or spinules on the surface

are due to the method of preserva-
tion. The cause of this character is

the accumulation of sparry matrix
at the mouths of such very narrow

canals, which often protrudes in hem-
ispherical masses and is of very gen-
eral occurrence in widely different

fossils.

RHOPALONARIA pertennis, Ulrich, Syn. for
Stomatopora proutana.

SEMICOSONTUM obliquatum, Ulrich, Syn. for
Fenestella obliquatum.

SEPTOPORA biserialis, instead of *Synocladia
biserialis*.

rectistyla, instead of *Synocladia recti-
styla*. It is described and figured in

Ohio Geol., vol. 7, p. 467.

Spatiopora, Ulrich, Syn. for *Actatopora*.

iovensis, Ulrich, 1893, not defined so as
to be recognized.

labeculosa, Ulrich, 1893, *Geo. Sur. Minn.*, vol. 3, p. 320, Trenton Gr.
STICTOPORELLA dumosa, Ulrich. Not defined so as to be recognized.
STOMATOPORA caudensis, Whiteaves, 1897, *Pal. Foss.*, vol. 3, p. 161, Low. Sil.
pertensis, Ulrich, Syn. for *S. proutana*.
tenuissima, Ulrich, Syn. for *S. proutana*.
STROMATOTRYPA, Ulrich, 1893, *Geo. Sur. Minn.*, vol. 3, p. 301. [Ety. *stroma*, layer; *trypa*, opening.] Zonarium consisting of one or more thin layers on foreign bodies. Zoecial tubes short, few diaphragms, proximal end scarcely prostrate, oval in cross section; walls thin, containing one or more constricted, bead-like tubuli to each zoecium. Apertures oval, separated by depressed interspaces, the peristomes minutely papillose. Mesopores abundant, beginning on the basal lamina,

decreasing in size with age, closely tabulated, the diaphragms finely punctured; mouths rarely visible, closed by a common dermal sheet. True ananthopores wanting. Type *S. ovata*.
ovata, Ulrich, 1893, *Geo. Sur. Minn.*, vol. 3, p. 302, Trenton Gr.
SYNOCLADIA. This is a Permian genus in the Eastern hemisphere, and is not known in America.
biseriatis, refer to *Septopora biserialis rectistyla*, refer to *Septopora rectistyla*.
THAMNISCUS is a Permian genus and many species referred to it probably belong elsewhere.
Trigonolictya, Ulrich. Not defined so as to be recognized.
UNITRYPA conferta, Ulrich, Syn. for *U. acanthis*.
retrota, Ulrich, Syn. for *U. tegulata*.

CLASS BRACHIOPODA.

DR. WAAGEN proposed, in *Palaeontologia Indica*, to subdivide the *Arthropomata* into three Suborders, *Kampylopegmata*, *sive Terebratulacea*, comprising the families *Terebratulidae*, *Thecideidae*, *Rhynchonellidae*, and *Stringocephalidae*; *Helicopegmata*, *sive Spiriferaceae*, comprising the families *Atrypidae*, *Nucleospiridae*, *Athyridae*, and *Spiriferidae*; *Aphaneropegmata*, *sive Productacea*, comprising the families *Strophomenidae*; and *Productidae*. As a matter of course, the words *Terebratulacea*, *Spiriferaceae*, and *Productacea* are much to be preferred as Subordinal names over *Kampylopegmata*, *Helicopegmata*, and *Aphaneropegmata*. He proposed to divide the Order *Lyopomata* into the Suborders, *Gastropegmata*, *sive Craniacea*, comprising the family *Craniidae*; *Daikaulia*, *sive Discinacea*, comprising the families *Discinidae*, and *Siphonotretidae*; *Mesokaulia*, *sive Lingulacea*, comprising the families *Obolidae*, *Trimerellidae*, and *Lingulidae*. And it is equally clear that *Craniacea*, *Discinacea*, and *Lingulacea* are to be preferred as Subordinal names to *Gastropegmata*, *Daikaulia*, and *Mesokaulia*. But how different are the views of Professor James Hall, who uses *Articulata* instead of *Arthropomata*, and *Inarticulata* instead of *Lyopomata*, and uses no Subordinal names or family names! He says the classification even into families, if attempted, must be arbitrary, procrustean, and embarrassing to the student, without any corresponding benefit, and that, in the present state of our knowledge, it is better to avoid them altogether. I agree with Professor Hall, that the Subordinal names may be dispensed with; but as the grouping into families is probably correct, in some respects, I prefer to retain the family names for the purpose of approximating the truth in classification.

Professor Hall, however, uses subgeneric names, which Thomson and Nicholson in their work on the "Generic Types of the Palaeozoic Corals," say "are nearly useless, if not absolutely obstructive, in actual practice." And Professor Hall has proposed to divide the genus *Orthis* into fourteen groups, viz.: *Orthis*, *Plectorthis*,

with age, closely
apophyses finely
rarely visible,
on dermal sheet.
wanting. Type 8.

o. Sur. Minn., vol.

Permian genus in
here, and is not

otopora biserialis
opora rectistyla.

i genus and many
probably belong

Not defined so as

elch. Syn. for U.

for U. tegulata.

side the *Arthropo-*
comprising the fam-
alidae; *Helicopeg-*
spiridae, *Athyridae*,
the families *Stroph-*
rebratulacea, *Spiri-*
names over *Kam-*
p divide the Order
prising the family
cinidae, and *Sipho-*
Obolidae, *Trimer-*
scinacea, and *Lina-*
ta, *Daikaulia*, and
es Hall, who uses
popomata, and uses
even into families,
the student, with-
our knowledge, it is
at the Subordinal
s probably correct,
urpose of approxi-

son and Nicholson
" say "are nearly
Professor Hall has
Orthia, *Plectorthia*,

Dinorthia, *Plesionmys*, *Hebertella*, *Orthostrophia*, *Platyostrophia*, *Heterorthia*, *Bilobites*, *Dalmanella*, *Rhipidomella*, *Schizophoria*, *Orthotichia*, and *Euteletra*. Any genus found in the Paleozoic rocks, in which there are more than two defined species may be divided into groups, in like manner and for as good reasons. The coining of technical names for groups may be made to exhaust the Greek Lexicon, but to me they do not appear consistent with binomial nomenclature, or of such value to the science as to overcome the burthen they place upon it.

I purchased a copy of Pal. N. Y., vol. 8, pt. 2, in August, 1897, immediately after hearing that it had been published. Mr. Jacob Van Deloo, at the same time, kindly sent me the last reports of the State Geologist of N. Y., from which it is evident, that some part at least of vol. 8, pt. 2, was published as early as, or earlier than, 1896, and I have arbitrarily assumed that, as the date of the genera and species, though the dedication of the book is dated November 29, 1894. The old genera are subdivided into new genera and subgenera, which I have catalogued without time to give them much examination, as this Appendix was prepared before I received the books, and the matter is inserted in the manuscript.

ACROTHELE matthewi var. costata, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 128, Up. Taconic.

AMBOCHELIA gemmula, McChesney, 1850, Desc. New Spec. Foss., p. 41, Conl Mens.

spinosa, Clarke, 1894, 13th Rep. St. Geol. N. Y., p. 177, Ham. Gr.

ANASTROPHIA (?) scottii, Winchell and Schuchert, 1895, Geo. Sur., Minn., vol. 3, p. 383, Galena Gr. *Camarella* (?)

ATHYRIS densa, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 364, St. Louis Gr.

differentius, instead of *A. differens*. *prinstana*, Billings, may be referred to *Hindella*, instead of *Meristella*.

ATRYPA laticorrugata, Foerste, Syn. for *A. reticularis*.

ATRYPINA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 161. Type *Leptocella imbricata* and including *Colospira disparilis*.

BARRANDELLA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, pp. 241-368. Type *B. areyi*, described at the same place from the Clinton Gr.

CAMARELLA. In the ventral valve the dental plates form a small chamber below the beak; in the dorsal valve the crura are fixed to two strong septal plates, which unite to form a very long and strong median septum.

CAMAROSPIRA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 82. Genus founded on *Camorphoria eucharis*, Pal. N. Y., vol. 4, p. 368, as the type.

CAMAROTECTIA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 189. Type *Rhynchonella congregata*, and including *R. altalis*, *R. plena*, *R. fringilla*, *R. glacialis*, *R. equiradiata*, *R. obtusiplicata*, and all the species included under the name *Stenoschisma* in Pal. N. Y., vol. 4.

CAPPELLINTA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 248. This genus is like *Pentamerus*, but has the valves reversed. Type *C.*

mica, described at the same place, from the Niagara Gr.

CATAZYGIA, Hall, 1896, Pal. N. Y., vol. 8, p. 157. This name is proposed as a subgenus of *Zygospira*, with *Zygospira heudi* as the type.

CHONETES pulchellus is described and figured in Bull. Denison Univ., vol. 3, p. 37. *reversus* is figured and described, in 1895, in Ohio Geol., vol. 7, p. 443.

CLINTONELLA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 159. [Ety. proper name.] Shells small suboval. Valves sub-

equally biconvex, greatest convexity oblique. Pedicle valve with a small umbo, compressed laterally, apex incurved. Cardinal area replaced by a wide triangular delthyrium, without a trace of deltidial plates. Umbo merges anteriorly into a sinus, which makes a flexure at the margin and bears two plications which reach the beak, and from four to eight radial plications on the lateral slopes. Beak in brachial valve inconspicuous. Surface with concentric striae. Shell substance, fibrous, impunctate. Type *C. vagabunda*, described at the same place from the Clinton Gr.

CLITHYRIS, King, 1850, Perm. Foss. Eng., p. 137. [Ety. *klein*, I close; *thuris*, a small door.] Lenticular, minutely punctured, laminar. Spirals pectinated. Dental plates large and separated. Oranal base perforated. Foramen situated at the point of the umbone, and open inferiorly by the fissure. Type, *C. pectinifera*. It includes *Athyris americana*, *A. hirsuta* and *A. sublemellosa*.

CONCHIDIUM, Linne, 1753, Museum Tensionianum, p. 90. Type *C. biloculare*. Hall includes in this genus *Pentamerus nyctius*, *P. tenuicosta*, *P. knappi*, *P. colletti*, *P. decussatus*. *C. nettlerothi*,

proposed by Hall in Pal. N. Y., vol. 8, pt. 2, p. 234, for *P. knighti* of Nettleroth, in Kentucky Foss. Shells, p. 57, and *Gypidula wagniformis*.
crassiplica, greenel, and georgie, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 369, the first two Niagara Gr.; the last, Clinton Gr.

Fig. 1466.—*Crania albersti*, magnified from $\frac{1}{24}$ to $\frac{3}{2}$ diameters.

Crania albersti, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, p. 154, Hud. Riv. Gr.

carbonaria is described and figured, in 1896, in Ohio Geol., vol. 7, p. 484.

chesterensis, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 47, Kaskaskia Gr.

dubia, Foerste. Too poorly defined to be recognized.

granulosa is described and figured in Geo. Sur. Minn., vol. 3, p. 373.

minutula, Winchell and Schuchert, 1895, (Schizotreta minutula.) Geo. Sur. Minn., vol. 3, p. 366, Hud. Riv. Gr.

reversa, Sardeson, 1896, Bull. Minn. Acad. Sci., vol. 4, p. 77, St. Peter Sandstone.

Cranicella clintonensis, Foerste. Too poorly defined to be recognized.

ulrichti, Hall, Pal. N. Y., vol. 8, pp. 153, 181, Syn. for *Cranicella hulli*.

Fig. 1467.—*Cranicella ovalis*, dorsal and side views.

CYCLOPHITNA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 206. Founded on *Rhynchospira nobilis*.

CYCLOSPIRA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 146. This genus is founded on *Orthis bicutata*, Emmons, Geol. N. Y. 2d District, p. 395. The name was used by permission without definition in Geo. Sur. Minn., vol. 3, p. 469.

spursiplica, Foerste. Too poorly defined to be recognized.

CYRTIA radians, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 362, Clinton Gr.

CYRTIA burlingtonensis, Rowley, 1893, Am. Geol., vol. 12, p. 308, Burlington Gr. *lachrymosa*, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 362, Waverly Gr.

Fig. 1468.—*Discina sumpsoni*, large and small dorsal valve, and east of ventral valve.

umbonata var. *alpenensis*, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 362, Ham. Gr.

Discina. Dall and Hall have shown that

this genus is not known from the paleozoic rocks, and that nearly all the

species referred to it belong to Orbiculoiden.

illinoisensis, Miller and Gurley, refer to Orbiculoiden Illinoisensis. *meekana* is described and illustrated, in 1896, in Ohio Geol., vol. 7, p. 483.

munda, Miller and Gurley, refer to Orbiculoiden munda.

EATONIA. Waagen distinguishes this genus by the absence of dental plates in the ventral valve, strong median septum in the dorsal valve, and the four crural processes.

coulteri, Miller and Gurley, 1893, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 72, Oriskany Gr.

Entelates, Fischer de Waldheim, 1830, Oryctographi du Gouvern. de Moscou, p. 144. Not defined so as to be recognized by any one, and, without any proper excuse, Waagen has proposed to supplant Syntrielasma with it. He might have used any other catalogue name, and he would have had his followers.

HALLINA, Winchell and Schuchert, 1895, Geo. Sur. Minn., vol. 3, p. 471. (It first appeared by name without proper definition in 1892, in Am. Geol., vol. 9, p. 291.) [Ety. proper name.] Shells small, articulate, rostrate, biconvex, and semiplicate. Pedicle opening usually, bounded laterally by incomplete deltidial plates. Calcified brachial supports longer than half the length of the dorsal valve. Crural plates of the dorsal valve probably coalesce. Shell structure fibrous, impunctate. Type *H. saffordi*.

nicolleti, Winchell and Schuchert, Syn. for *Atrypa exigua*.

saffordi, Winchell and Schuchert, 1895, Geo. Sur. Minn., vol. 3, p. 473, Birds-eye limestone.

HIPPARIOSYX, Vanuxem, 1842, Geo. of N. Y. Rep. 3d Dist., p. 29. [Ety. *hippos*, horse; *onyx*, nail, hoof.] Shell large, subhemispherical. Pedicle valve slightly convex or concave; hinge-line short, straight; cardinal area low; beak retrorse; delthyrium broad, and covered by an imperforate convex deltidium; teeth large, and supported by lamellae which extend to the bottom of the umbonal cavity, and are produced into strong ridges that surround a large muscular area composed of broad, flabellate diductors, enclosing an elongate or cordate adductor impression. Slight median septum in both valves. No cardinal area in brachial valve. General appearance externally like *Streptorhynchus*, to which genus it has sometimes been referred. Type *H. proximus*.

proximus, Vanuxem, 1842, Geo. N. Y. Rep. 3d Dist., p. 124, Oriskany sandstone.

HUSTEDIA, Hall, 1896, Pal. N. Y., vol. 8,

belong to Orbicula-

Gurley, refer to

ensis.

and illustrated, in

ol. 7, p. 483.

and the four crural

guishes this genus

ntal plates in the

g median septum

and the four crural

Gurley, 1893, Bull.

Nat. Hist., p. 72,

Waldheim, 1890,

uv. de Moscou, p.

ns to be recog-

and, without any

gen has proposed

ism with it. He

y other catalogue

have had his fol-

Schuchert, 1895,

ol. 3, p. 471. (It

me without proper

Am. Geol., vol. 9,

er name.) Shells

ostrate, biconvex,

dicle opening usu-

ally by incomplete

nicified brachial

n half the length

Crural plates of

probably coalesce.

rous, impunctate.

Schuchert, Syn.

Schuchert, 1895,

ol. 3, p. 473, Birds-

1842, Geo. of N.

20. [Ety. hippos,

of.] Shell large,

Pedicle valve

concave; hinge-

t; cardinal area

lethyrium broad,

perforate convex

ge, and supported

extend to the bot-

l cavity, and are

g ridges that sur-

cular area com-

bellate diductors,

e or cordate ad-

Slight median

ves. No cardinal

ve. General ap-

like Streptorhyn-

it has sometimes

H. proximus.

1842, Geo. N. Y.

4, Oriskany sand-

al. N. Y., vol. 8,

pt. 2, p. 120. [Ety. proper name.]

This genus is externally like *Eumetria*,

but differs in the internal structure.

Type *Terebratula mormoni*.

HYATTELLA, Hall, 1896, Pal. N. Y., vol. 8,

pt. 2, p. 61. Genus founded on *Atrypa*

congesta, Pal. N. Y., vol. 2, p. 67, as

the type.

KUTORGINA ambigua, instead of *Obolella*

ambigua.

LEIORHYNCHUS newberryi is from the

Chemung Gr.

LEPTENA charlotte, Winchell and Schu-

chert, Syn. for *Strophomena halli*.

gibbosa, Winchell and Schuchert, 1895,

(*Plectambonites gibbosa*.) Geo. Sur.

Minn., vol. 3, p. 416, Galena Gr.

producta, Hall, 1896, (*Plectambonites*

producta.) Pal. N. Y., vol. 8, pt. 2, p.

360, Niagara Gr.

sordida, refer to *Leptella sordida*.

LINGULA antiqua, refer to *Lingulella* an-

tiqua.

beltrami, Winchell and Schuchert, 1895,

Geo. Sur. Minn., vol. 3, p. 351, Hud.

Riv. Gr.

clathrata, Winchell and Schuchert, 1895,

Geo. Sur. Minn., vol. 3, p. 345, Trenton

Gr.

compta, Hall, 1892, Pal. N. Y., vol. 8,

p. 171, Ham. Gr.

deflecta, Winchell and Schuchert, 1895,

Geo. Sur. Minn., vol. 3, p. 348, Hud.

Riv. Gr.

dolata, Sarsen, 1890, Bull. Minn. Acad.

Sci., vol. 4, p. 95, Magnesian Gr.

flabellula, Hall, 1892, Pal. N. Y., vol. 8,

p. 172, Waverly Gr.

hurbuti, described and illustrated in

Geo. Sur. Minn., vol. 3, p. 347.

indianensis, Miller and Gurley, 1893,

Bull. No. 3, Ill. St. Mus. Nat. Hist., p.

69, Keokuk Gr.

lamellata, Hall, is from the Niagara Gr.

lingulata, Hall, 1892, Pal. N. Y., vol. 8,

p. 173, Clinton Gr.

modesta, Ulrich, Syn. for *L. norwoodi*.

norwoodi, refer to *Lingulops norwoodi*.

paliformis, refer to *Lingulella paliformis*.

paracetus, Hall, 1892, Pal. N. Y., vol. 8,

p. 172, Waverly Gr.

scutella, Hall, 1892, Pal. N. Y., vol. 8, p.

171, Chemung Gr.

teniola, Hall, 1892, Pal. N. Y., vol. 8, p.

18, Clinton Gr. Proposed instead of

L. lamellata in Pal. N. Y., vol. 2, p. 55.

waverlyensis, Herrick, is the same that

others have identified with *L. scotica*

of Davidson.

LINGULELLA inflata var. ovalis, Matthew,

1894, Trans. N. Y. Acad. Sci., vol. 14,

p. 127, Up. Taconic.

LINGULOPS granti, Hall, 1892, Pal. N. Y.,

vol. 8, p. 173, Niagara Gr.

LIORHYNCHUS lesleyi, Hall, 1896, Pal. N.

Y., vol. 8, pt. 2, p. 368, Up. Devonian.

LISSEPLEURA, Whitfield, 1896, Bull. Am.

Mus. Nat. Hist., vol. 8, p. 232. [Ety.

lisos, smooth; *pleura*, rib.] A brachio-

podous shell, more or less inequivalve,

with a small imperforate beak; sur-

face radiately ribbed; ribs smooth,

without interspaces. Shell substance

fibrous. Ventral valve with a spoon-

shaped cavity in the beak, formed by

the dental plates, and a deep, bilobed,

muscular imprint in front of it. Dor-

sal valve with a strong median sep-

tum. Type *Rhynchonella aquivalvis*,

Hall. Low. Held. Gr.

MARGINIFERA, Wagon, 1887, Paleontologia

Indica, p. 17. [Sig. bearing a margin.]

Externally somewhat like *Productus*,

but the shell margin readily breaks off

and exposes a thick, prominent, shelly

ridge, placed vertically on the internal

surface of the dorsal valve, and by

which the visceral part of that valve

is girt. In the ventral valve there is

a similar ridge, developed within the

wings only. In this way the visceral

part of the shell is perfectly chambered

off from the remainder of the shell.

These ridges are smooth, striated, or

eremulated. Type *M. splendens*. Pro-

fessor Hall doubted the generic value

of these distinguishing characters,

while I regard them as of more impor-

tance than the characters ascribed to

Orthothetes or *Derbya*, or many other

proposed genera among the Brachio-

pods.

lasallensis, instead of *Productus lasal-*

ensis.

splendens, instead of *Productus splen-*

dens.

wabashensis, instead of *Productus wa-*

bashensis.

MARTINIA was described on page 139, in-

stead of 128. It is also distinguished

from *Spirifera* by having no dental

plates in the ventral valve.

MEKKELLA occidentalis, Newberry, 1861,

(*Streptorhynchus occidentalis*.) Ives

Rep. Col., p. 126, Up. Coal Meas.

pyramidalis, Newberry, 1861, (*Strepto-*

rhynchus pyramidalis.) Ives Rep. Col.,

p. 126, Up. Coal Meas.

MERISTA bella, refer to *Meristella bella*.

tennesseensis, Hall, 1896, Pal. N. Y.,

vol. 8, pt. 2, p. 365, Up. Sil.

MERISTELLA walcottii, Hall, 1896, Pal. N. Y.,

vol. 8, pt. 2, p. 365, Oriskany Gr.

METAPLASTIA, Hall, 1896, Pal. N. Y., vol. 8,

pt. 2, p. 56. A genus based on *Spirif-*

era pyxidata, Pal. N. Y., vol. 3, p. 428,

as the type.

NECLEOSPIRA rotundata is described and

illustrated in Ohio Geol. vol. 7, p. 413.

OBOLELLA ambigua, refer to *Kutorgina* am-

bigua.

cingulata, refer to *Kutorgina cingulata*.

OBOLUS pristinus, Matthew, 1894, Trans.

N. Y., Acad. Sci., vol. 14, p. 112, Up.

Taconic.

pulcher, Matthew, 1889, Can. Rec. Sci.,

p. 306, St. John Gr.

ORBICULOIDEA, D'Orbigny, 1850, not 1847.

- Hall has shown in Pal. N. Y., vol. 8, p. 128, that *O. morristi*, Davidson, is the type of the genus, and that it includes nearly all the forms described as *Discina* in the palaeozoic rocks. Diagnosis. Shells subcircular or subelliptical, inequivalve. Pedicle valve depressed convex, or flattened, with the apex slightly elevated and inclined posteriorly. On the exterior a narrow pedicle furrow begins just below and behind the apex, extends over a greater or less portion of the radius of the valve, and at its distal end is produced into a short tubular siphon, which traverses the substance of the shell obliquely backward, emerging on the interior surface, where it produces a narrow groove, which terminates before reaching the margin. On the interior there is a thickened ridge corresponding with the external groove. The apex of the larger or brachial valve is directed backward, and on the interior there is a longitudinal ridge or septum extending backward. Shell substance corneous, lamellae appearing phosphatic. Surface marked by fine lamellose concentric striae and radiating lines.
- conica*, refer to *Schizotreta conica*.
illinoisensis, Miller and Gurley, 1893, (*Discina illinoisensis*.) Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 70, Coal Meas.
munda, Miller and Gurley, 1893, (*Discina munda*.) Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 71, Coal Meas.
- ORISKANIA**, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 269. Type *O. navicella*, figured at the same place from the Oriskany Gr.
- ORTHIDIUM**, Hall, 1892, Pal. N. Y., vol. 8, p. 244. Shell small, externally resembling *Scenidium*. Pedicle valve the more convex, broad open delthyrium, strong teeth, inconspicuous dental plates. Brachial valve less convex, cardinal area narrow, dental sockets developed. Crural plates short, erect, coalesced with the cardinal process, which is a vertical, transverse, subrescentic plate, at the base of which the shell is excavated. Muscular scar quadruplicate. Surface bearing radiating striae fold and sinus. Type *O. gemmicula*.
gemmicula, instead of *Orthis gemmicula*.
ORTHIS acutiloba, Ringueberg, Syn. for *O. biloba*.
billingsi is referred to *Protorthis billingsi*.
charlottae, Winchell, Syn. for *O. pectinella*.
circularis, Winchell, Syn. for *O. subaequata*.
fausta and *var. squamosus*, synonyms for *O. rugiplicata*.
flabellites, Hall, 1892, Pal. N. Y., vol. 8, p. 227, Niagara Gr. Instead of *O. flabellum*, which is not an American species.
- futilis*, Sardeson, 1897, Am. Geol., vol. 19, Hud. Riv. Gr.
glypta, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 359, Niagara Gr.
ignota, Sardeson, 1897, Am. Geol., vol. 19, p. 99, Hud. Riv. Gr.
inaequalis, refer to *Streptorhynchus inaequale*.
kaskaskiensis, refer to *Streptorhynchus kaskaskiense*.
maerior, Sardeson, Syn. for *O. emacerata*.
media, Winchell, Syn. for *O. subaequata*.
meedsi, Winchell and Schuchert, Syn. for *O. minnesotensis*.
nichelini var. *nevadensis*, Meek, 1877, Expl. 40th Parallel, vol. 4, p. 63, Carboniferous.
minneapolis, Winchell, Syn. for *O. subaequata*.
porrecta, Sardeson, 1897, Am. Geol., vol. 19, p. 104, Hud. Riv. Gr.
proarita, Winchell and Schuchert, Syn. for *O. petre*.
proximus is referred to *Hipparionyx proximus*.
quacoensis is referred to *Protorthis quacoensis*.
richmondi is *Streptorhynchus richmondi*.
sweeneyi, Winchell, 1895, Geo. Sur. Minn., vol. 3, p. 426, Trenton Gr.
whitfieldi, Winchell, 1895, Geo. Sur. Minn., vol. 3, p. 437, Hud. Riv. Gr.
- ORTHO RHYNCHULA**, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 181. This genus is founded on *Orthis linneyi*, Nettleroth.
Orthothetes, Fischer de Waldheim, 1830, Oryctographie. This genus was not defined as required by the laws of nomenclature, but is revived by Waagen, in *Palaeontologia Indica*, p. 807, for such forms as *Spirifera crenistria*, Phillips, and *Strophomena arctostriata*, Hall, and *Streptorhynchus pandora*, Billings. The name can not stand as used by Waldheim, or as used by Evans in 1829, nor by reason of being a catalogue name. And though Waagen defined it in 1887, he had no right to use the word. He says it is not to be distinguished by any external characters, but the cardinal process is small and not supported by septa, but instead in general there is a septum in the dorsal valve and none in the ventral.
bellulus, Clarke, 1894, 13th Rep. St. Geol. N. Y., p. 176, Marcellus Shale, refer to *Streptorhynchus bellulum*.
desideratus, Hall, refer to *Streptorhynchus desideratum*.
- PARASTROPHIA**, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 221. Type *Atrypa (Camarella) hemiplicata*, and including *Pentamerus reversus*.
divergens, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 366, Hud. Riv. Gr.
greeniei, latiplicata, and multiplicata, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 367, Niagara Gr.
PARAZYGIA, Hall, 1896, Pal. N. Y., vol. 8,

Am. Geol., vol.
N. Y., vol. 8, pt.
Am. Geol., vol.
Streptorhynchus in-
for *O. emacerata*.
for *O. subequata*.
Schuchert, Syn. for
sis, Meek, 1877,
vol. 4, p. 63, Car-
Syn. for *O. sub*-
97, Am. Geol., vol.
Gr.
Schuchert, Syn.
Hipparionyx prox-
o Protorthis qua-
ynchus richmondi.
1895, Geo. Sur.
Trenton Gr.
1895, Geo. Sur.
Hud. Riv. Gr.
1896, Pal. N. Y.,
1. This genus is
megi, Nettleroth.
Waldheim, 1830,
is genus was not
by the laws of
revived by Waa-
ia Indica, p. 607.
pirifera crenistria,
mena arclostriata,
ynchus pandora,
can not stand as
as used by Evans
n of being a cat-
ough Waagen de-
ad no right to use
it is not to be dis-
ternal characters,
cess is small and
ta, but instead in
ptum in the dorsal
e ventral.
3th Rep. St. Geol.
ellus Shale, refer
pellulum.
r to Streptorhyn-
8, Pal. N. Y., vol.
e *Atrypa* (*Cama-*
d including *Penta-*
Pal. N. Y., vol. 8,
iv. Gr.
and multiplicata,
., vol. 8, pt. 2, p.
al. N. Y., vol. 8,

pt. 2, p. 127. This genus is founded on
Trematospira lirsuta, Pal. N. Y., vol.
4, p. 274.
PENTAGONIA is recognized by Hall, in his
latest work, as a good genus.
PENTAMERUS oblongus var. corrugatus,
Weller, 1896, Jour. Geo., vol. 4, p. 171,
Niagara Gr.
pesovis is described and illustrated in
Ohio Geol., vol. 7, p. 414.
PUOLIDOPS cincinnatiensis is inequivalve,
and the pedicle valve has a subcircular
foramen.
greenel, Miller and Gurley, 1897, Bull.
No. 12, Ill. St. Mus. Nat. Hist., p. 48,
Ham. Gr.
Plectambonites, Syn. for *Leptena*.
PRODUCTELLA marquesti, Rowley, 1894, Am.
Geol., vol. 14, p. 153, Ham. Gr.
PRODUCTUS inflatus, McChesney, 1859, Desc.
New Spec. Foss., p. 40, Coal Meas.
lasallensis, refer to *Marginifera lasallen-*
sis.
longus, Meek, 1877, Expl. 40th Parallel,
vol. 4, p. 67, Col Meas.
pileiformis, McChesney, 1859, Desc. New
Spec. Foss., p. 40, Kaskaskia Gr.
splendens, refer to *Marginifera splendens*.
tubulospinus, McChesney, 1859, Desc.
New Spec. Foss., p. 37, Up. Coal Meas.
wabashensis, refer to *Marginifera wabash-*
ensis.
wilberinus, McChesney, 1859, Desc. New
Spec. Foss., p. 36, Coal Meas.
PROTORHYNCHA, Hall, 1896, Pal. N. Y., vol.
8, pt. 2, p. 180. This genus is founded
on *Atrypa dubia*, Hall.
PROTORTHIS, Hall, 1892, Pal. N. Y., vol. 8,
p. 231. [Ety. *protos*, first; *Orthis*, a
genus.] Shells small, transversely sub-
quadrate or semicircular. Hinge-line
straight, equal to the greatest width of
the valves. Valves unequally biconvex
or subplano-convex, the pedicle-valve
being the larger. Cardinal area nar-
row on both valves, but wider on the
pedicle valve; delthyrium broad, closed
below by a concave plate; teeth pres-
ent. Brachial valve with a delthyrium,
dental sockets obscure, crural plates
small. Surface plicated, and having
interstitial radii and concentric striae,
sinus and fold. Shell substance fibrous.
Type *P. billingsi*.
billingsi, instead of *Orthis billingsi*.
quacoensis, instead of *Orthis quacoensis*.
PROTOZYGA, Hall, 1896, Pal. N. Y., vol. 8,
pt. 2, p. 149. This genus is founded on
Atrypa exigua, Pal. N. Y., vol. 1, p. 141.
PTYCHOSPIRA, Hall, 1896, Pal. N. Y., vol. 8,
pt. 2, p. 112. Like *Retzia*, but having
a coarsely and sparsely plicated sur-
face. Type *P. ferita*, and includes *Ret-*
zia sexplicata, which is illustrated on
pl. 50, Figs. 13, 14.
PUGNAX, Hall, 1896, Pal. N. Y., vol. 8, pt. 2,
p. 202. A subgenus of *Rhynchonella*,
with *R. acuminata* as the type, and in-
cluding *R. pugnax*, *R. reniformis*, *R.*

alta, *R. missouriensis*, *R. striatocostata*,
R. explanata, *R. mutata*, *R. ottumwa*,
R. uta, *R. eatoniiformis*, and *Camaro-*
phoria scallariana.
Rafinesquina, Hall, 1892, Pal. N. Y., vol. 8,
p. 280, Syn. for *Strophomena*.
lata. See *Strophomena lata*.
RENSELERIA formosa is described and il-
lustrated in Ohio Geol., vol. 7, p. 413.
RETICULARIA, McCoy, 1844, Synop. Carb.
Foss., p. 143. [Ety. *reticulum*, a little
net.] Shells rounded, orbicular, or elon-
gately or transversely oval. Hinge-
line shorter than the greatest width of
the shell. Surface covered by fine hair-
like spines, arranged in concentric rows,
and representing double tubes, that
pass below the surface of the shell, but
do not pass through it. Shell fibrous.
The muscular impressions of the ven-
tral valve are in an elongately oval
groove. There are no partitions, den-
tal plates, or median septum in the
valve. In the dorsal valve there are no
partitions, septum, or shelly support
of the dental sockets. No hinge-plate.
The crura are fixed with a broad base
to the inner side of the dental sockets,
and extend down to the frontal region,
where they abruptly bend up to form
the first volution of the spiral. No
lateral branch. Apex of the spiral di-
rected laterally or toward the hinge-
line. Type *R. lineata*, instead of *Spiri-*
feria lineata.
RHYNCHONELLA *aequalvalvis*, refer to *Lisso-*
pleura aequalvalvis.
hydraulica and varicosta are described
and illustrated in Ohio Geol., vol. 7,
pp. 414 and 421.
RHYNCHOTREMA, Hall, 1860, 13th Rep. N. Y.
St. Mus. Nat. Hist., p. 68. [Ety. *rhyn-*
chos, beak; *trema*, an opening.] Hall
proposed to distinguish this from *Rhyn-*
chonella on the ground that the car-
dinal area of the ventral valve results
from the bending inward and coales-
cing of the deltidial plates, but Waa-
gen has suggested that the absence of
dental plates in the ventral valve is
of far greater importance, and estab-
lishes the genus. Type *R. capax*, and
including *R. increbescens*, *R. dentata*, *R.*
speciosa, and *Trematospira quadriplicata*,
S. A. Miller, heretofore referred to
Rhynchotrema.
SCENIDIUM *halli*, as figured in Pal. N. Y.,
vol. 8, pl. VII A., is a syn. for *Sceni-*
dium anthonense.
SCHIZAMBON (?) *dodgei*, Winchell and
Schuchert, 1895, Geo. Sur. Minn., vol.
3, p. 361, Trenton Gr.
lockei, Winchell and Schuchert, Syn.
for *Trematis terminalis*.
? *canadensis*, Ami, 1892, Pal. N. Y., vol.
8, p. 116, Utica Slate.
SCHIZOTREMA, Kutorga, 1848, Ueber die
Siphonotretæ; Verhandl. der russ.
kais. mineral. Gesellsch. zu St. Pe-

- tersburg, p. 272. [Ety. *schiza*, a cleft; *tretum*, perforated.] The external, slit-shaped pedicle aperture is the reverse of that in *Siphonotreta*. The apex of the beak is high, eccentric, and posterior to it there is a narrow elliptical slit, which extends for one-half the face of the cone, and then merges into the inner siphon. The brachial valve is depressed convex, or even flat; its beak is sharply defined, depressed, and directed toward the cardinal margin, but not marginal. Type *S. elliptica*. conica, instead of Orbiculoidea conica. minutula, Winchell and Schuchert. See *Crania minutula*.
- SELENELLA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 270. Type *S. gracilis*, figured at the same place from the Niagara Gr.
- SEMINULA, McCoy, 1844, Synop. Carb. Foss., p. 150. He spelled the word, in 1855, *Semiluna*. Shell smooth, subpentagonal, valvexsinuate. Muscular impressions consisting of two pairs of very narrow elongate scars. Type *S. ambigua*. It includes *Athyris trinuclea* and *A. subtilita*.
- dawsoni, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 364, Carboniferous.
- rogersi, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 364, Up. Held. Gr.
- Siphonotreta scotica* is referred by Hall to *Schizambon* (?) *canadensis*.
- SPIRIFERA aciculifera, Rowley, 1893, Am. Geol., vol. 12, p. 307, Chouteau Gr.
- canandaigue, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 360, Ham. Gr.
- clavatula, McChesney, is a good species.
- crispata, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 360, Niagara Gr.
- lineata, Martin, refer to *Reticularia lineata*.
- macbridei, Calvin, 1892, Bull. State Univ. Iowa, vol. 2, p. 165, Ham. Gr.
- mundula, Rowley, 1893, Am. Geol., vol. 12, p. 307, Burlington Gr.
- norwoodi is described in Expl. 40th Parallel, vol. 4, p. 39.
- perplexa, McChesney, is a good species from the Coal Meas.
- pulchra, Meek, refer to *Spiriferina pulchra*. It is described in Expl. 40th Parallel, vol. 4, p. 85.
- solidirostris is described in Bull. Denison Univ., vol. 3, p. 47.
- subventricosa, McChesney, 1859, Desc. New Spec. Foss., p. 44, Coal Meas.
- transversa, McChesney, is a good species, and is on p. 42.
- urbana, Calvin, 1892, Bull. State Univ. Iowa, vol. 2, p. 165, Ham. Gr.
- williamsi, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 361., Chemung Gr.
- SPIRIFERINA gonionota, Meek, 1877, Expl. 40th Parallel, vol. 4, p. 84, Carboniferous.
- STREPTORHYNCHUS crenistria, Phillips, (misspelled on p. 378, *crenistriatum*), is not an American species.
- desideratum, Hall, 1892, (Orthothetes desideratus,) Pal. N. Y., vol. 8, p. 345, Waverly Gr.
- flabellum and hydraulicum are described and illustrated in Ohio Geol., vol. 7, pp. 410 and 421.
- inequale, Hall, 1858, (Orthis inequalis,) Geo. Rep. Iowa, p. 490, Chouteau or Kinderhook Gr.
- kaskaskiense, McChesney, 1859, (Orthis kaskaskiensis,) Desc. New Spec. Foss., p. 31, Kaskaskia Gr.
- occidentalis. See *Meekella occidentalis*.
- pyramidalis. See *Meekella pyramidalis*.
- richmonda, McChesney, 1859, (Orthis richmonda,) Desc. New Spec. Foss., p. 32, Coal Meas.
- winchelli, Hall, 1892, (Strophomena winchelli,) Pal. N. Y., vol. 8, p. 344, Galena Gr.
- STRICKLANDINIA triplexiana, Foerste. Not defined so as to be recognized.
- STROPHALOSIA beecheri, Rowley, 1893, Am. Geol., vol. 12, p. 308, Chouteau Gr.
- STROPHODONTA, Hall, instead of *Strophodonta*. (Ety. *stropheos*, heart; *odous*, tooth:)
- STROPHOMENA billingsi, Winchell and Schuchert, Syn. for *Streptorhynchus rectum*.
- emaciata, Winchell and Schuchert, 1895, Geo. Sur. Minn., vol. 3, p. 399, Galena Gr. It looks like an *Orthis*, and is not a *Strophomena*.
- hanoverensis, Foerste. Not defined so as to be recognized.
- lata, Whiteaves, 1896, (Rafinesquina lata,) Can. Rec. Sci., vol. 6, p. 172, Low Sil.
- planodorsata, Winchell and Schuchert, Syn. for *Streptorhynchus planumbonum*.
- schofieldi, Winchell and Schuchert, Syn. for *Streptorhynchus subsulcatum*.
- septata, Winchell and Schuchert, Syn. for *Streptorhynchus subtentum*.
- trentonensis, Winchell and Schuchert, Syn. for *Streptorhynchus subtentum*.
- STROPHONELLA costatula, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 359, Niagara Gr.
- crassa, Rowley, 1894, Am. Geol. vol. 13, p. 153, Ham. Gr.
- SYNTROPHIA, Hall, 1896, Pal. N. Y. vol. 8, pt. 2, p. 216. The type is *Triplexia lateralis*, Whitfield, and intended to include *Orthis barabensis*, *Stricklandinia arachne*, and *S. arethusa*.
- SYRINGOTHYRIS missouri, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 363, Chouteau Gr.
- TRIMERELLA gallensis refer to *Rhinobolus gallensis*.
- TRIPLASIA ulrichi, Winchell and Schuchert, 1895, Geo. Sur. Minn., vol. 3, p. 409, Hud. Riv. Gr.
- WHITFIELDIA, Hall, 1896, Pal. N. Y., vol. 8, pt. 2, p. 58. A genus founded on *Atrypa nitida* or *Meristina nitida*, Pal. N. Y., vol. 2, p. 266, as the type.

1892, (Orthothetes
Y., vol. 8, p. 345,

um are described
Ohio Geol., vol. 7,

(Orthis inæqualis),
490, Chouteau or

ney, 1859, (Orthis
New Spec. Foss.,

kella occidentalis.
kella pyramidalis.
ey, 1859, (Orthis
New Spec. Foss., p.

Strophomena win-
ol. 8, p. 344, Galena

ana, Foerste. Not
recognized.

Rowley, 1893, Am.
Chouteau Gr.

instead of Stropho-
leas, heart; odous,

Winchell and Schu-
eptorhynchus rec-

nd Schuchert, 1895,
ol. 3, p. 399, Galena
n *Orthis*, and is not

Not defined so as

1896, (Rafinesquina
, vol. 6, p. 172, Low

ll and Schuchert,
ynchus planumbo-

nd Schuchert, Syn.
s-subsulcatum.

Schuchert, Syn. for
btentum.

ll and Schuchert,
chus subtentum.

a, Hall, 1896, Pal.
p. 359, Niagara Gr.

Am. Geol. vol. 13,

, Pal. N. Y. vol. 8,
type is *Triplexia lat-*
nd intended to in-
nsis, *Stricklandinia*
husa.

i, Hall, 1896, Pal.
2, p. 363, Chou-

refer to Rhinobo-

hell and Schuchert,
inn., vol. 3, p. 409,

1896, Pal. N. Y., vol.
genus founded on

eristina nitida, Pal.
as the type.

ZYGOSPIRA uphami, Winchell and Schu-
chert, 1895, Geo. Sur. Minn., vol. 3, p.
468, Galena Gr.

aquila, Sardeson, Syn. for *Atrypa exigua*.
putilla, Hall, 1896, Pal. N. Y., vol. 8, pt.
2, pp. 157-365, Hud. Riv. Gr.

SUBKINGDOM MOLLUSCA.

CLASS PTEROPODA.

COLEOLUS clintonensis, Foerste. Not de-
fined so as to be recognized.

CONULARIA is an Order established by
Miller and Gurley, 1893, Bull. No. 11,
Ill. St. Mus. Nat. Hist., p. 22, where
it is claimed that the Order may not
belong to the Pteropoda or to the Gas-
tropoda, and as the Order was anni-
hilated, in Palæozoic times, it may
even belong to an extinct class in the
Subkingdom Mollusca. The family
Conulariidae is discussed at the same
place.

CONULARIA *bilineata*, Foerste. Too poorly
defined to be recognized.

blairi, Miller and Gurley, 1893, Bull. No.
3, Ill. St. Mus. Nat. Hist., p. 73, Chou-
teau Gr.

chesterensis is illustrated in Geo. Sur.
Ill., vol. 8, pl. 11.

gratiosa, Miller and Gurley, 1893, Bull.
No. 3, Ill. St. Mus. Nat. Hist., p. 74,
St. Louis Gr.

greenei, Miller and Gurley, 1896, Bull.
No. 11, Ill. St. Mus. Nat. Hist., p. 27,
Keokuk Gr.

missouriensis is from the Keokuk Gr.
newberryi is illustrated in Ohio Pal., vol.
2, pl. 18.

roeperi, Miller and Gurley, 1896, Bull.
No. 11, Ill. St. Mus. Nat. Hist., p. 26,
Coal Meas.

sedaliensis, Miller and Gurley, 1896,
Bull. No. 11, Ill. St. Mus. Nat. Hist.,
p. 28, Burlington Gr.

spergenensis, Miller and Gurley, 1893,
Bull. No. 3, Ill. St. Mus. Nat. Hist., p.
74, St. Louis Gr.

ENCHOSTOMA, Miller and Gurley, 1896, Bull.
No. 11, Ill. St. Mus. Nat. Hist., p. 29.
[Ety. *enchos*, sword; *stoma*, blade.]
Shell smooth, elongate, lanceolate;
transverse section more or less rounded
or narrowly subovate. Shell substance
thin, solid, flexible, horny, lime-phos-
phate. Type *E. lanceolatum*.

lanceolatum, S. A. Miller, 1892, Advance
Sheets 18th Rep. Geo. Sur. Ind., p. 63,
Chouteau Gr.

Hyolithellus is generally regarded as a
Brachiopod, and the same that was
named by Hall, *Discinella*.

HYOLITHES *alatus*, Whiteaves, 1892, Cont.
to Can. Pal., p. 342, Devonian.

ceratophilus, Clarke, 1894, 13th Rep. St.
Geol. N. Y., p. 172, Up. Held Gr.

dubius, Miller and
Faber, 1894, Jour.

Cin. Soc. Nat.
Hist., vol. 17, p.

155, Hud. Riv. Gr.

gracillor, Matthew,
1894, Trans. N. Y.

Acad. Sci., vol.
14, p. 130, Up. Ta-

conic.

versaillesensis, Mil-
ler and Faber,

1894, Jour. Cin. Soc. Nat. Hist., vol.
17, p. 155, Hud. Riv. Gr.

Fig. 1409.—*Hyolithes*
versaillesensis, mag-
nified 4 diameters.

CLASS GASTROPODA.

ANTHRACOPUPA *ohioensis* is described and
illustrated in Ohio Geol., vol. 7, p. 491.

Fig. 1410.—*Aclisina*
bellilineata.

Archinacella, Ulrich, 1897,
Geo. Sur. Minn., vol.

3, p. 821. No generic
character is men-

tioned that will distin-
guish it from *Trybli-*

dium, and, if not a full synonym, in
all aspects it can not have the force of

more than a catalogue name. Part of
the species referred to it belong to
Tryblidium, and others may belong to
Carinaropsis.

cingulata. See *Tryblidium cingulata*.
depressa. See *Tryblidium depressum*.

powersi. See *Tryblidium powersi*.
richmondensis, Syn. for *Tryblidium in-*

dianense.
rotunda. See *Tryblidium rotundum*.

rugatina, Syn. for *Tryblidium indianense*.
semicarinata, Syn. for *Carinaropsis patelliformis*.
simulatrix, Syn. for *Carinaropsis patelliformis*.
subrotunda, Syn. for *Tryblidium validum*.
ASTRALITES, Whiteaves, 1892, Cont. to Can. Pal., p. 323. [Ety. from the living genus *Astraliu*.] Shell conical, imperforate, flattened at the base; periphery subangular, fringed with a thin, regularly lobate or sinuate lateral expansion; columella encircled with a single narrow but prominent spiral fold, which is represented by a deep spiral groove in casts of the interior. Type *A. fimbriatus*, described at the same place, from the Devonian.
BELLEROPHON alternodosus is illustrated and described in Ohio Geol., vol. 7, p. 479.
bilineatus, Ulrich, Syn. for *B. troosti*.

Fig. 1411.—*Bellerophon gorbii*, front and dorsal views.

blairi, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 21, Chouteau Gr.
capax, Ulrich, Syn. for *B. mohri*.
cincinnatiensis, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, p. 29, Hud. Riv. Gr.
clausus, Ulrich, Syn. for *B. troosti*.
exiguus, Foerste, 1896, Ohio Geol., vol. 7, p. 548, Niagara Gr.
globularis, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., p. 28, Hud. Riv. Gr.
lindsleyi is described and figured in Geo. Sur. Minn., vol. 3, p. 889.
opertus, Foerste, Syn. for *B. exiguus*.
recurvus, Ulrich, Syn. for *B. cincinnatiensis*.
sedaliensis, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 21, Chouteau Gr.
similis, Ulrich, Syn. for *B. platystoma*.
subangularis, Ulrich, Syn. for *Bellerophon cincinnatiensis*.
subglobulus, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 917, Trenton Gr.
BUCANIA crassa, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 893, Hud. Riv. Gr.
elliptica, Ulrich, Syn. for *B. intexta*.
emmonsii, Ulrich, Syn. for *B. halli*.
frankfortensis, Ulrich, Syn. for *Bellerophon lindsleyi*.
halli, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 886, Trenton Gr.
imbricata, Ulrich, 1897, (Salpingostoma imbricata,) Geo. Sur. Minn., vol. 3, p. 902, Hud. Riv. Gr.

miconema, Ulrich, Syn. for *Bellerophon lindsleyi*.
minnesotensis, Ulrich, Syn. for *B. halli*.
nana, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 895, Trenton Gr.
nashvillensis, Ulrich, Syn. for *Bellerophon lindsleyi*.
obsoleta, Ulrich, 1897, (Tetraota obsoleta,) Geo. Sur. Minn., vol. 3, p. 880, Trenton Gr.
peracuta, Ulrich. Not defined so as to be recognized.
richmondensis, Ulrich, 1897, (Salpingostoma richmondensis,) Geo. Sur. Minn., vol. 3, p. 903, Hud. Riv. Gr.
rugatina, Ulrich, Syn. for *Bellerophon lindsleyi*.
sculptilis, Ulrich, 1897, (Salpingostoma sculptilis,) Geo. Sur. Minn., vol. 3, p. 902, Trenton Gr.
simulatrix, Ulrich, Syn. for *Bucania expansa*.
singularis, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 894, Trenton Gr.
subangulata, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 891, Trenton Gr.
sublata, Ulrich, Syn. for *B. halli*.
Bucanopsis, Ulrich, Syn. for *Bucania*.
curvifera, Ulrich, Syn. for *Bucania costata*.
Bucanospira, Ulrich. Proposed without intelligent definition.
CARINAROPSIS acuta, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 928, Black Riv. Gr.
explanata, Ulrich. Not defined so as to be recognized.
minima, Ulrich, Syn. for *C. phalera*.
Clathrospira, Ulrich, Syn. for *Pleurotomaria*.
conica, Ulrich, Syn. for *Pleurotomaria subconica*.
convexa, Ulrich, Syn. for *Pleurotomaria subconica*.
Cælocaulus. A subgenus of *Murchisonia*, proposed by Oehlert in 1888, but not recognized by any American paleontologist.
negeletus, Ulrich, founded on a fragment of *Murchisonia gracilis*.
œhlerti. See *Murchisonia œhlerti*.
CONCHOPELTIS compressa, Ulrich, 1897, (Scenella compressa,) Geo. Sur. Minn., vol. 3, p. 840, Trenton Gr.
Conradella, Ulrich, Syn. for *Cyrtolites*.
bellula, Ulrich, Syn. for *Cyrtolites elegans*.
fimbriata. See *Cyrtolites fimbriatus*.
grandis, *obliqua*, *similis*, and *triangularis*, Ulrich, Synonyms for *Cyrtolites compressus*.
CYCLONEMA gracile, *humerosum*, *inflatum*, *limatum*, *mediale*, *pulchellum*, (preoccupied,) *simulans*, *sublere*, and *transversum*, Ulrich, Synonyms for *Cyclonema bilix*.
pulchellum, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 19, Keokuk Gr.
textile, Ulrich, 1897, (Strophostylus tex-

for Bellerophon

yn. for B. halli.
Sur. Minn., vol.

Syn. for Bellerophon

(Tetranotha obso-
n., vol. 3, p. 880,

defined so as to

, 1897, (Salpingo-
) Geo. Sur. Minn.,
iv. Gr.

for Bellerophon

7, (Salpingostoma
Minn., vol. 3, p.

for Bucania ex-

, Geo. Sur. Minn.,
on Gr.

1897, Geo. Sur.
Trenton Gr.

or B. halli.
for Bucania.

for Bucania cos-

Proposed without
1. 1897, Geo. Sur.

, Black Riv. Gr.
not defined so as to

for C. phalera.

Syn. for Pleuroto-

for Pleurotomaria

for Pleurotomaria

s of Murchisonia,
t in 1888, but not
American paleon-

ded on a fragment
ilis.

onia oehlerti.

sa, Ulrich, 1897,
) Geo. Sur. Minn.,
on Gr.

for Cyrtolites.
for Cyrtolites ele-

ites fimbriatus.
e, and triangularis,
for Cyrtolites com-

erosum, inflatum,
ulchellum, (preoc-

ubleve, and trans-

onyms for Cyclo-

and Gurley, 1896,
Mus. Nat. Hist.,

Strophostylus tex-

tilis,) Geo. Sur. Minn., vol. 3, p. 1064,
Trenton Gr.

CYRTOLITES dilatatus, Ulrich, 1897, Geo.
Sur. Minn., vol. 3, p. 865, Black River
Gr.

disjunctus, Ulrich, Syn. for C. ornatus.
fimbriatus, Ulrich, 1897, (Conradella
fimbriata,) Geo. Sur. Minn., vol. 3, p.
907, Trenton Gr.

nitidulus, Ulrich, Syn. for C. carinatus.
parvus, Ulrich, Syn. for C. carinatus.

retroreus, Ulrich, Syn. for C. ornatus.
subacutus, Ulrich, 1897, (Oxydiscus sub-

acutus,) Geo. Sur. Minn., vol. 3, p.
913, Trenton Gr.

youngi, Foerste. Not defined so as to
be recognized.

CYRTONELLA horrida, Clarke, 1893, 12th
Ann. Rep. N. Y. St. Geol., p. 48.
Probably a Platyceras, but too poorly
defined for recognition.

CYRTOSPIRA, Ulrich, Syn. for Subulites.
bicaurata, tortilis, and wykoffensis. See
Subulites.

DENTALIUM martini is described and illus-
trated in Ohio Geol., vol. 7, p. 423.

DYERIA, Ulrich, Syn. for Bucania.

ECCYLIOMPHALUS contiguus, Ulrich, 1897,
Geo. Sur. Minn., vol. 3, p. 1037, Tren-
ton Gr.

subrotundus, Ulrich, Syn. for Eccyli-
omphalus undulatus.

ECCYLIPTERUS beloitensis, Ulrich, Syn. for
Ophileta owenana.

EOTOMARIA, Ulrich, Syn. for Pleurotomaria.
canalifera, see Pleurotomaria canalifera.
elevata, see Scallites elevatus.

labiosa, Ulrich, Syn. for Pleurotomaria
canalifera.

vicina, Ulrich, Syn. for Pleurotomaria
dryope.

EUCONOSPIRA, Ulrich, Syn. for Pleuroto-
maria.

planibasalis, Ulrich, Syn. for Pleuroto-
maria missouriensis.

EUNEMA brevispira, Whiteaves, 1892, Cont.
to Can. Pal., p. 320, Devonian.

clathratulum, Whiteaves, 1892, Cont. to
Can. Pal., p. 322, Devonian.

speciosum, Whiteaves, 1892, Cont. to
Can. Pal., p. 321, Devonian.

subspinosum, Whiteaves, 1892, Cont. to
Can. Pal., p. 321, Devonian.

EUOMPHALUS subtrigonalis, Whiteaves,
1892, Cont. to Can. Pal., p. 328, De-
vonian.

winonensis, Sardeson, 1896, Bull. Minn.,
Acad. Sci., vol. 4, p. 96, Magnesian Gr.

FUSISPIRA angusta, Ulrich, 1897, Geo. Sur.
Minn., vol. 3, p. 1079, Trenton Gr.

convexa, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1078, Trenton Gr.

intermedia, Ulrich, Syn. for F. inflata.

nobilis, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1078, Trenton Gr.

planulata, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1078, Trenton Gr.

schucherti, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1078, Black Riv. Gr.

subbrevis, Ulrich, Syn. for F. inflata.

sulcata, Ulrich, Syn. for F. schucherti.

GYRONEMA, Ulrich, Syn. for Cyclonema.

duplicatum, Ulrich, Syn. for Pleuroto-
maria percarinata.

HELEIONOPSIS, Ulrich, Syn. for Tryblidium.

subcarinata, Ulrich, may be a Carinini-
opsis.

HELICOTOMA declivis is figured in Geo. Sur.
Minn. vol. 3, p. 1036, from the Trenton
Gr.

marginata, Ulrich. Not defined so as to
be recognized.

peccatonica, Sardeson, 1896, Bull. Minn.
Acad. Sci., vol. 4, p. 97, Magnesian Gr.

HOLOPEA amphi, Ulrich, 1897, Geo. Sur.
Minn., vol. 3, p. 1065, Trenton Gr.

appressa, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1065, Trenton Gr.

concinnulla, Ulrich, 1897, Geo. Sur.
Minn., vol. 3, p. 1066, Trenton Gr.

excelsa, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1067, Trenton Gr.

grandis, Miller and Gurley, 1896, Bull.

Fig. 1412.—*Holopea grandis*, lateral view.

No. 11, Ill. St. Mus. Nat. Hist., p. 19,
Keokuk Gr.

insignis, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1065, Trenton Gr.

newtonensis is described and illustrated
in Ohio Geol., vol. 7, p. 477.

parvula, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1067, Trenton Gr.

rotunda, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1066, Trenton Gr.

similis, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1066, Trenton Gr.

supraplana, Ulrich, 1897, Geo. Sur.
Minn., vol. 3, p. 1068, Trenton Gr.

HORMOTOMA was suggested by Salter as a
subgenus of Murchisonia, with Mur-
chisonia gracilis as the type, in 1859,
Can. Org. Rem. Dec. 1, p. 18. It has
not come into use among paleontolo-
gists.

salteri, Ulrich, Syn. for Murchisonia gra-
cilis.

subangulata, Ulrich, Syn. for Murchi-
sonia gracilis.

trentonensis, Ulrich, Syn. for Murchi-
sonia bellicincta.

winnipegeensis. See Murchisonia winni-
pegensis.

KOKENIA, Ulrich, Syn. for Bucania.

LIOSPIRA, Ulrich, Syn. for Raphistoma.

- abrupta*, Ulrich, Syn. for *Raphistoma lapicida*.
angustata and *obtusa*, Ulrich, Synonyms for *Pleurotomaria progne*.
angulata, *mundula*, *subconca*, and *rugata*, Ulrich, Synonyms for *Pleurotomaria eugenia*.
decipiens, refer to *Raphistoma decipiens*.
persimilis, Ulrich, Syn. for *Pleurotomaria helena*.
Lophospira abnormis, Ulrich, Syn. for *Murchisonia medialis*.
acuminata. See *Murchisonia acuminata*.
ampla. See *Murchisonia ampla*.
centralis. See *Murchisonia centralis*.
concinna, *fillmorensis*, *obliqua*, *perforata*, *procera*, and *pulchella*, Ulrich, Synonyms for *Murchisonia milleri*.
conoidea. See *Murchisonia conoidea*.
conradana, Ulrich, Syn. for *Murchisonia ventricosa*.
decursa, *producta*, and *tenuistriata*, Ulrich, Synonyms for *Murchisonia perangulata*.
humilis. See *Murchisonia humilis*.
knoxvillensis. See *Trochonema knoxvillense*.
lirata. See *Murchisonia lirata*.
medialis and *medialis* var. *burginensis*. See *Murchisonia medialis*.
notabilis. See *Trochonema notabile*.
oweni. See *Murchisonia oweni*.
peracuta. See *Murchisonia peracuta*.
perlamellosa. See *Murchisonia perlamellosa*.
quadrilucata. See *Murchisonia quadrilucata*.
saffordi. See *Murchisonia saffordi*.
spironema. See *Murchisonia spironema*.
trochonemoides. See *Trochonema trochonemoides*.
LOXONEMA altivolve, Whiteaves, 1892, Cont. to Can. Pal., p. 334, Devonian.
cingulatum, Whiteaves, 1892, Cont. to Can. Pal., p. 336, Devonian.
gracillimum, Whiteaves, 1892, Cont. to Can. Pal., p. 337, Devonian.
parvulum and *plicatum* are described and illustrated in Ohio Geol., vol. 7, pp. 424 and 486.
regularis, refer to *Macrochilina regularis*.
winnipegense, Whiteaves, 1893, Can. Rec. Sci., p. 326, Trenton Gr.
MACLUREA crassa, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1040, Trenton Gr.
depressa, Ulrich, Syn. for *Maclurea bigsbyi*.
knoxvillensis, Ulrich. Not defined so as to be recognized.
nitida, Ulrich, Syn. for *Maclurea bigsbyi*.
Maclurina, Ulrich, Syn. for *Maclurea*.
Fig. 1413.—*MACROCHILINA* *prisca* and *blairi*. *MACROCHILINA* *prisca* and *blairi* are described and illustrated in Ohio Geol., vol. 7, pp. 424 and 478.
epulehlla, Whiteaves, 1892, Cont. to Can. Pal., p. 340, Devonian.
regularis, instead of *Loxonema regularis*. It is described and illustrated in Ohio Geol., vol. 7, p. 485.
Meekospira, Ulrich, Syn. for *Polyphemopsis subconica*. See *Subulites subconica*.
MURCHISONIA *ampla*, Ulrich, 1897, (*Lophospira ampla*.) Geo. Sur. Minn., vol. 3, p. 981, Hud. Riv. Gr.
acuminata, Ulrich, 1897, (*Lophospira acuminata*.) Geo. Sur. Minn., vol. 3, p. 973, Hud. Riv. Gr.
argylensis, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 97, Shakopee Dolomite.
centralis, Ulrich, 1897, (*Lophospira centralis*.) Geo. Sur. Minn., vol. 3, p. 979, Trenton Gr.
conoidea, Ulrich, 1897, (*Lophospira conoidea*.) Geo. Sur. Minn., vol. 3, p. 976, Trenton Gr.
elevata, Ulrich, 1897, (*Lophospira elevata*.) Geo. Sur. Minn., vol. 3, p. 977, Trenton Gr.
humilis, Ulrich, 1897, (*Lophospira humilis*.) Geo. Sur. Minn., vol. 3, p. 968, Trenton Gr.
indianensis, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 18, Keokuk Gr.
lirata, Ulrich, 1897, (*Lophospira lirata*.) Geo. Sur. Minn., vol. 3, p. 968, Utica Gr.
medialis, Ulrich, 1897, (*Lophospira medialis*.) Geo. Sur. Minn., vol. 3, p. 973, Trenton Gr.
ohlerti, Ulrich, 1897, (*Cœlocaulus ohlerti*.) Geo. Sur. Minn., vol. 3, p. 1020, Galena Gr.
oweni, Ulrich, 1897, (*Lophospira oweni*.) Geo. Sur. Minn., vol. 3, p. 980, Trenton Gr.
peracuta, Ulrich, 1897, (*Lophospira peracuta*.) Geo. Sur. Minn., vol. 3, p. 976, Trenton Gr.
perlamellosa, Ulrich, 1897, (*Lophospira perlamellosa*.) Geo. Sur. Minn., vol. 3, p. 985, Hud. Riv. Gr.
putilla, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 98, Oneota Dolomite.
pygmæa, Rowley, 1895, Am. Geol., vol. 16, p. 222, Chouteau Gr. Very poorly defined.
quadrilucata, Ulrich, 1897, (*Lophospira quadrilucata*.) Geo. Sur. Minn., vol. 3, p. 967, Hud. Riv. Gr.
saffordi, Ulrich, 1897, (*Lophospira saffordi*.) Geo. Sur. Minn., vol. 3, p. 982, Trenton Gr.
spironema, Ulrich, 1897, (*Lophospira spironema*.) Geo. Sur. Minn., vol. 3, p. 983, Black Riv. Gr.
textilis, Ulrich, 1897, (*Schizolophia textilis*.) Geo. Sur. Minn., vol. 3, p. 992, Hud. Riv. Gr.
winnipegensis, Whiteaves, 1897, (*Hormotoma winnipegensis*.) Pal. Foss., vol. 3, p. 192, Low. Sil.

Fig. 1413.—*MACROCHILINA* *prisca* and *blairi*. *MACROCHILINA* *prisca* and *blairi* are described and illustrated in Ohio Geol., vol. 7, pp. 424 and 478.

1892, Cont. to Can.
Monema regulare.
 illustrated in Ohio
 Polyphemosis.
 subconica.
 Ulrich, 1897, (Lopho-
 spir. Minn., vol. 3
 1897, (Lophospira
 Minn., vol. 3, p.
 1896, Bull. Minn.
 4, p. 97, Shakopee
 (Lophospira cen-
 Minn., vol. 3, p. 979,
 (Lophospira co-
 Minn., vol. 3, p. 976,
 (Lophospira el-
 Minn., vol. 3, p. 977,
 (Lophospira hu-
 Minn., vol. 3, p. 968,
 and Gurley, 1896,
 Mus. Nat. Hist.,
 Lophospira lirata,) 3, p. 968, Utica Gr.
 (Lophospira med-
 Minn., vol. 3, p. 973,
 1897, (Cœlocaulus
 Minn., vol. 3, p.
 Lophospira oweni,) 3, p. 980, Tren-
 7, (Lophospira per-
 Minn., vol. 3, p. 976,
 1897, (Lophospira
 Sur. Minn., vol. 3,
 1896, Bull. Minn.
 4, p. 98, Oneota
 95, Am. Geol., vol.
 u Gr. Very poorly
 1897, (Lophospira
 o. Sur. Minn., vol.
 Gr.
 7, (Lophospira saf-
 Minn., vol. 3, p. 982,
 1897, (Lophospira
 Sur. Minn., vol. 3, p.
 (Schizolopha tex-
 Minn., vol. 2, p. 992,
 teaves, 1897, (Horg-
 ensis,) Pal. Foss,
 Sil.

Naticopsis inornata, Whiteaves, 1892, Cont.
 to Can. Pal., p. 333, Devonian.
manitobensis, Whiteaves, 1892, Cont. to
 Can. Pal., p. 332, Devonian.
 orton and ziczac are described and illus-
 trated in Ohio Geol., vol. 7, pp. 489 and 477.
 ventrica, Norwood and Pratt, instead
 of *ventricosa*.
Omospira, Ulrich, Syn. for *Scalites*.
laticincta. See *Scalites laticinctus*.
 ✓ *Oxyochilus nitidulus*, Clarke, 1894, 13th
 Rep. St. Geol. N. Y., p. 172, Marcellus
 Shale.
OPHILETA alturensis, Sardeson, 1896, Bull.
 Minn. Acad. Nat. Sci., vol. 4, p. 98,
 Oneota Dolomite.
 angularis, Ulrich, 1897, (Ophileta an-
 gularis,) Geo. Sur. Minn., vol. 3, p.
 1031, Black Riv. Gr.
 complanata var. *nana* is in Expl. 40th
 Parallel, instead of 4th.
 fausta, Sardeson, 1896, Bull. Minn. Acad.
 Nat. Sci., vol. 4, p. 75, St. Peter Sand-
 stone.
 subluxa, Ulrich, 1897, (Ophileta sub-
 laxa,) Geo. Sur. Minn., vol. 3, p. 1030,
 Trenton Gr.
Ophileta, Ulrich, Syn. for *Ophileta*.
Oxydiscus subacutus. See *Cyrtolites suba-*
cutus.
Owenella, Ulrich, Syn. for *Bellerophon*.
PALEAOMELA humilis, Ulrich, 1897, Geo.
 Sur. Minn., vol. 3, p. 837, Trenton Gr.
Pulcopupa, Foerste, and also the type *P.*
abrupta. Too poorly defined to be rec-
 ognized.
PALAGIELLA, Matthew,
 1894, Trans. N. Y.
 Acad. Sci., vol. 14,
 p. 131, Up. Taconic.
 Shell few, whorled,
 discoid, flattened,
 outer edge angular,
 slightly tumid be-
 low; upper and
 lower lips arched
 forward in the middle. Type *P. at-*
lantoides, described
 at the same place
 from the Up. Taconic.
PLATYCERAS cyrtolites
 is from the Burling-
 ton Gr.
depressum, Ulrich.
 Not defined so as to
 be recognized.
indianense, Miller and
 Gurley, 1897, Bull.
 No. 12, Ill. St. Mus.
 Nat. Hist., p. 48,
 Ham. Gr.
parvulum, Whiteaves,
 1892, Cont. to Can.
 Pal., p. 331, Devon-
 nian.
squalodens is de-
 scribed and illus-
 trated in Ohio Geol., vol. 7, p. 423.

Fig. 145.—*Platyceras indianense*, lateral view of a large specimen.

Fig. 144.—*Platyceras boonvillense*.

vetulum, Sardeson, 1896, Bull. Minn.
 Acad. Nat. Sci., vol. 4, p. 76, St. Peter
 Sandstone.
wisconsinensis, Ulrich. Not defined so
 as to be recog-
 nized.

PLATYSCHISMA. The
 type is *P. cir-*
roides.

PLATYSTOMA tum-
idum, White-
 eaves, 1892. The
 name was preoc-
 cupied by Meek
 and Worthen in
 1860. See *P. whiteavesi*.

Fig. 143.—*Platyceras missouriense*.

Fig. 147.—*Platystoma broadheadi*, front and summit views.

whiteavesi, n.
 sp. Proposed
 instead of *P.*
tumidum,
 Whiteaves,
 in Cont. to
 Can. Pal., p.
 331, pl. 43,
 fig. 12, De-

voman, Pentamerus Point, Lake Man-
 itoba.
Plethospira, Ulrich, Syn. for *Pleuroto-*
maria.
striata. See *Pleurotomaria striata*.
PLEUROTOMARIA niens, Sardeson, 1896, Bull.
 Minn. Acad. Nat. Sci., vol. 4, p. 76, St.
 Peter Sandstone.

Fig. 148.—*Pleurotomaria haril*, front view, showing aperture and height of shell.

canalifera, Ulrich, 1897, (Eotomaria ca-
 nalifera,) Geo. Sur. Minn., vol. 3, p.
 1002, Trenton Gr.
gilitexta, Foerste. The name was preoc-
 cupied.
margaritoides, Whiteaves, 1897, Pal.
 Foss., vol. 3, p. 190, Low. Sil.
minima, Rowley, 1895, Am. Geol., vol. 16,
 p. 222, Chouteau Gr. Poorly defined.
providencis, Broadhead, 1896, Am. Jour.
 Sci., vol. 152, p. 237, Ham. Gr.
spenceri, Whiteaves, 1892, Cont. to Can.
 Pal., p. 341, Devonian.
stokesiana, Whiteaves, 1897, Pal. Foss.,
 vol. 3, p. 190, Low. Sil.
striata, Ulrich, 1897, (Plethospira stri-
 ata,) Geo. Sur. Minn., vol. 3, p. 1011,
 Hud. Riv. Gr.
ventricosa, Ulrich, 1897, (Seelya ventri-
 cosa,) Geo. Sur. Minn., vol. 3, p. 1009,
 Calciferous Gr.
POLYPHEMOPSIS melanoides is described
 and illustrated in Ohio Geol., vol. 7,
 p. 478.

Protowartha, Ulrich, Syn. for *Bellerophon concinna* and *subcompressa*, Synonyms for *Bellerophon morrowensis*.

granistriata and *planodorsata*, Ulrich, Synonyms for *Bellerophon globularis oboea*, *pervoluta*, and *rectangularis*, Ulrich, Synonyms for *Bellerophon bilobatus*.

✓ *PNEUDOPHORUS* tectiformis, Whiteaves, 1892, Cont. to Can. Pal., p. 330, Devonian.

RAPHISTOMA affine, Foerste, 1895, Ohio Geol., vol. 7, p. 550, Niagara Gr. decipiens, Ulrich, 1897, (*Liospira decipiens*.) Geo. Sur. Minn., vol. 3, p. 998, Trenton Gr.

leiosomellum, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 99, Oneota Dolomite.

lewistonense, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 99, Oneota Dolomite.

owenii, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 100, Oneota Dolomite.

peracutum, Ulrich, Syn. for *R. lenticulare richmondensis*, Ulrich, Syn. for *R. lenticulare*.

ruddum, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 101, Shakopee Dolomite.

Raphistomina, Ulrich, Syn. for *Raphistoma R. denticulata*, *modesta*, and *rugata*, Synonyms for *R. lapidea*.

Salpingostoma imbricata. See *Bucania imbricata*.

richmondensis. See *Bucania richmondensis*.

sculptilis. See *Bucania sculptilis*.

SCALITUS elevatus, Ulrich, 1897, (*Eotomaria elevata*.) Geo. Sur. Minn., vol. 3, p. 1005, Trenton Gr.

laticinctus, Ulrich, 1897, (*Omospira laticincta*.) Geo. Sur. Minn., vol. 3, p. 945, Trenton Gr.

SCENELLA affinis, Syn. for *Conchopeltis compressa*.

beloitensis, Syn. for *Conchopeltis minnesotensis*.

compressa. See *Conchopeltis compressa*.

magnifica. Not defined so as to be recognized.

radialis, Syn. for *Conchopeltis obtusa*.

Schizolopha, Ulrich, Syn. for *Murchisonia moorei*, Syn. for *Murchisonia multigruma*.

textilis. See *Murchisonia textilis*.

Seelya, Ulrich, Syn. for *Pleurotomaria mundula*, Ulrich. Not recognized; probably distorted specimens of *Cyclonema ventricosa*. See *Pleurotomaria ventricosa*.

Solenospira, Ulrich, Syn. for *Murchisonia*.

STENOITHECA unguiformis. See *Tryblidium unguiforme*.

✓ *STRAPAROLLINA* obtusa, Whiteaves, 1892, Cont. to Can. Pal., p. 328, Devonian.

STRAPAROLLUS incarinatum, Foerste, 1895, Ohio Geol., vol. 7, p. 552, Niagara Gr. intralobatus, Sardeson, 1896, Bull. Minn.

Acad. Nat. Sci., vol. 4, p. 101, Calciferous—Oneota Dolomite.

missouriensis, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 20, Chouteau Gr.

STROPHOSTYLUS textilis. See *Cyclonema textile*.

SUBULITES beloitensis, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1072, Trenton Gr.

bicurvatus, Ulrich, 1897, (*Cyrtospira bicurvata*.) Geo. Sur. Minn., vol. 3, p. 1074, Trenton Gr.

canadensis, Ulrich, Syn. for *Subulites elongatus*.

conradi, Ulrich, Syn. for *S. elongatus*.

directus, Foerste, Syn. for *S. gracilis*.

dlxonensis, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1071, Trenton Gr.

exactus, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 101, Calciferous Gr.

nanus, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1072, Trenton Gr.

parvus, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1072, Trenton Gr.

pergracilis, Ulrich. Not defined so as to be recognized.

planilateralis, Foerste. Too poorly defined to be recognized.

regularis, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1072, Trenton Gr.

subconicus, Ulrich, 1897, (*Meekospira subconica*.) Geo. Sur. Minn., vol. 3, p. 1080, Hud. Riv. Gr.

tortilis, Ulrich, 1897, (*Cyrtospira tortilis*.) Geo. Sur. Minn., vol. 3, p. 1074, Trenton Gr.

wykoffensis, Ulrich, 1897, (*Cyrtospira wykoffensis*.) Geo. Sur. Minn., vol. 3, p. 1074, Trenton Gr.

Tetranota, Ulrich, Syn. for *Bucania*.

macra and *securinata*, Synonyms for *Bucania bidorsata*.

obsoleta. See *Bucania obsoleta*.

Trepospira, Ulrich, Syn. for *Pleurotomaria*.

TROCHONEMA altum, Ulrich, Syn. for *Pleurotomaria niota*.

arctatum, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1054, Trenton Gr.

bellulum, Ulrich, Syn. for *B. eccentricum*.

eccentricum, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1049, Trenton Gr.

fragile, Ulrich, Syn. for *T. umbilicatum*.

knoxvillense, Ulrich, 1897, (*Lophospira knoxvillensis*.) Geo. Sur. Minn., vol. 3, p. 989, Trenton Gr.

maddisonense and *rugosum*, Ulrich, Synonyms for *T. umbilicatum*.

nitidum, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1053, Utica Gr.

notabile, Ulrich, 1897, (*Lophospira notabilis*.) Geo. Sur. Minn., vol. 3, p. 990, Black Riv. Gr.

obsoletum, Ulrich, 1897, Geo. Sur. Minn., vol. 3, p. 1054, Trenton Gr.

retrosum, Ulrich, Syn. for *T. subcrassum*.

4, p. 101, Calcifer-
te.
and Gurley, 1896,
Mus. Nat. Hist.,

See Cyclonema

Ulrich, 1897, Geo.
1072, Trenton Gr.
97, (Cyrtospira bi-
Minn., vol. 3, p.

yn. for Subulites

or S. elongatus.

for S. gracilis.

7, Geo. Sur. Minn.,
on Gr.

1896, Bull. Minn.
4, p. 101, Calcifer-

Geo. Sur. Minn., vol.
Gr.

Geo. Sur. Minn.,
ton Gr.

ot defined so as to

e. Too poorly de-
ed.

7, Geo. Sur. Minn.,
ton Gr.

1897, (Meekospira
ur. Minn., vol. 3, p.

, (Cyrtospira tor-
in., vol. 3, p. 1074.

1897, (Cyrtospira
Sur. Minn., vol. 3,

for Bucania.

ta, Synonyms for

a obsoleta.

n. for Pleurotomma

Ulrich, Syn. for Pleu-

7, Geo. Sur. Minn.,
ton Gr.

n. for B. eccentrici-

, 1897, Geo. Sur.
9, Trenton Gr.

or T. umbilicatum.

1897, (Lophospira
p. Sur. Minn., vol.

Gr.
psum, Ulrich, Syn-

licatum.

7, Geo. Sur. Minn.,
a Gr.

7, (Lophospira no-
Minn., vol. 3, p.

97, Geo. Sur. Minn.,
ton Gr.

yn. for T. subcras-

robbinsi, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1053, Trenton Gr.

salteri, Ulrich, 1897, Geo. Sur. Minn.,
vol. 3, p. 1053, Trenton Gr.

simile, Ulrich. Not defined.

subcrassum, Ulrich, 1897, Geo. Sur.
Minn., vol. 3, p. 1051, Trenton Gr.

trochomenoides, Ulrich, 1897, Geo. Sur.
Minn., vol. 3, p. 990, Trenton Gr.

vagrans, Ulrich, Syn. for T. beloitense.

TRYBLIDIUM cingulatum, Ulrich, 1897, (Ar-
chinneella cingulata,) Geo. Sur. Minn.,

vol. 3, p. 829, Trenton Gr.

depressum, Ulrich, 1897, (Archinnucella
depressa,) Geo. Sur. Minn., vol. 3, p.

830, Trenton Gr.

madisonense, refer to Cyrtoceras madi-
sonensis.

modestum. Too poorly defined to be rec-
ognized.

powersi, Ulrich, 1897, (Archinnucella pow-
ersi,) Geo. Sur. Minn., vol. 3, p. 829,

Trenton Gr.

repertum, Sardeson, 1896, Bull. Minn.
Acad. Nat. Sci., vol. 4, p. 102, Shakopee

Dolomite.

rotundum, Ulrich, 1897, (Archinnucella
rotunda,) Geo. Sur. Minn., vol. 3, p.

835, Hud. Riv. Gr.

striatum, Ulrich, 1897, (Helelonopsis
striatum,) Geo. Sur. Minn., vol. 3, p. 827,

Hud. Riv. Gr.

unguiforme, Ulrich, 1897, (Stenotheca
unguiformis,) Geo. Sur. Minn., vol. 3,

p. 843, Trenton Gr.

CLASS CEPHALOPODA.

It is very clear that the genus *Nautilus* does not exist in Paleozoic rocks, and that the species referred to that genus belong elsewhere. *Lituites* is unknown from North America. The early generic references of authors are frequently erroneous. Professor Hyatt, who has given the subject profound study, has divided the Class into a great many genera, and has cited foreign authors, whose works are not accessible to me, and I am frequently unable to understand what are relied upon as generic characters. I have reproduced such definitions as I have had an opportunity to examine, though characters, which he regards as of generic value, I sometimes think are of no more than specific importance.

APHELECERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 293. [Ety. *apheles*, smooth; *keras*, horn.] No type specified. Much like *Discitoceras*, and said to include *D. disciforme*.

APHETOCERAS, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 447. [Ety. *aphetos*, free; *keras*, horn.] Shell smooth, coiled in the same plane, but not in contact. Whorls in section, compressed, elliptical, or oviform, the venter narrower than the dorsum. Siphuncle subventral. Type *A. americanum*.

americanum, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 447, Calciferous (?) Gr.

attenuatum, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 449, Calciferous Gr.

boreale, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 448, Quebec Gr.

complanatum, instead of *Lituites* complanatus.

farnsworthi, Billings, 1861, (*Lituites farnsworthi*.) Pal. Foss., vol. 1, p. 21, Calciferous Gr.

ASOCERAS costulatum, Whitelaves, 1896, Can. Rec. Sci., vol. 6, p. 394, Low. Sil. gibberosum, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 102, Onkota Dolomite.

BARRANDEOCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 299. [Ety. proper name; *keras*, horn.] Large umbilicus, and compressed, slightly costated, or smooth whorls. Ventral side narrower than dorsal; whorls barely in contact; siphon above the center; septa deeply concave, and sutures with ventral saddles, lateral lobes, and dorsal saddles, without annular lobes. Type *B. natator*.

convolvans, Hall, 1847, (*Lituites convolvans*.) Phil. N. Y., vol. 1, p. 53, Black Riv. Gr.

elrodi, White, 1882, (*Gyroceras elrodi*.) 11th Ann. Rep. Geo. Ind., p. 356, Niagara Gr.

minganense, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 451, Chazy Gr.

natator, Billings, 1859, (*Nautilus natator*.)

- Can. Nat. and Geol., vol. 4, p. 466, Chazy Gr.
- Cameroceras hennepini*. See *Endoceras hennepini*.
- CENTROCERAS**, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 283. [Ety. *ken-tro-n*, a spur; *keras*, horn.] Whorls compressed; abdomen hollow, sometimes narrow, with a row of tubercles on each side. Sutures have deep lateral dorsal and ventral lobes, the latter V-shaped. Type *C. marcellense*.
- ammonis*, Hall, 18'9, (*Discites ammonis*,) Pal. N. Y., vol. 5, pt. 2, p. 425, Up. Held. Gr.
- marcellense*, Vanuxem, 1842, (*Goniatites marcellensis*,) Geo. Sur. 3d Dist. N. Y., p. 146, Marcellus Shale.
- CYCLOGASTEROCERAS**, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 498. He says the genus is defined in 4th Ann. Rep. Geo. Sur. Texas, but it is not in my copy. He describes *Cyclogasteroceras canaliculatum* in Proc. Am. Phil. Soc., vol. 32, p. 498, from the Carboniferous.
- COLOCERAS**, Hyatt, 1893, 4th Ann. Rep. Geo. Sur. Texas, p. 449. Type *C. globatum*.
- globulare*, Hyatt, 1893, 4th Ann. Rep. Geo. Sur. Texas, p. 452, Coal Meas.
- CYCLOLITUITES**, Remele, but I have no reference to his work.
- americanus*, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 505. Group of rocks not referred to. No illustration.
- CYRTOCERAS** *clintonense*, Foerste, 1895, Ohio Geol., vol. 7, p. 534, Niagara Gr. cretaceum is described and illustrated in Ohio Geol., vol. 7, p. 429.
- dresbachense*, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 102, Oneota Dolomite.
- dunleithense*, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 30, Trenton Gr.
- eatonense*, Claypole, 1878, (*Glyptodendron eatonense*,) Am. Jour. Sci., vol. 115, p. 302, Niagara Gr. The specimen is so poor that the specific name is of no value.
- featherstonhaughii*, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 807, Trenton Gr.
- houghtoni*, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 807, Trenton Gr.
- kansasense*, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 88, Up. Coal Meas.
- laticurvatum*, Whiteaves, 1896, Can. Rec. Sci., vol. 6, p. 365, Low. Sil.
- maximum*. See *Nephriticeras maximum*, instead of *Nautilus maximus*.
- metellus*. See *Melonoceras metellus*.
- minneapolis*, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 808, Trenton Gr.
- norwoodi*, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 809, Trenton Gr.
- scofieldi*, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 810, Trenton Gr.
- shumardi*, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 810, Trenton Gr.
- undulatum*, refer to *Holoceras undulatum*.
- CYRTOCERINA madisonensis**, instead of *Tryblidium madisonense*.
- schoolcrafti*, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 774, Trenton Gr.
- DELTOCERAS**, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 449. [Ety. *delto-s*, a scroll; *keras*, horn.] Shell similar to *Aphetoceras*, but more complicated. Whorls similar in section, but grow more rapidly in the ventro-dorsal diameters. Siphuncles large and ventral. Whorls in contact in the earlier epembryonic stages or throughout the ephebic stage. A departure from the spiral regularly takes place in the gerontic stage or earlier; sometimes the entire ephebic stage is free. No impressed zone. Type *D. planum*.
- planum*, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 450, Calceferous Gr.
- Discites*. This name was twice preoccupied before DeHaan used it in 1825, in his monograph, page 41. His definition was without illustration, and too brief to describe a genus. McCoy, in 1844, used the word in his Synop. Carb. Foss., p. 17, and for the genus defined by McCoy, Hyatt has proposed *Discitoceras*.
- ammonis*. See *Centroceras ammonis*.
- marcellensis*. See *Centroceras marcellense*.
- DISCITOCERAS**, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 292. It includes species with quadrangular whorls, having the abdomen slightly convex, sides flattened, the dorsum very gibbous, and having a slight impression. The young are ridged longitudinally, with prominent transverse striae, not subspinous. The sutures have ventral and lateral lobes and broad dorsal saddles, with small annular lobes. Siphon above the center. Living chamber from one-fourth to three-fourths of a volution. Aperture with a deep ventral sinus. Type *D. costellatum*. The *Discites* must be referred to this genus.
- DISCOCERAS**, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 500. [Ety. *diskos*, quoit; *keras*, horn.] Discoid, planorbicular, concave on both sides. Volutions three or more, slightly embracing, with transverse section nearly circular. Surface marked by distant costae, which cross the sides obliquely backward, curving more strongly on the periphery, indicating a profound sinus in the lip. Imbricating striae of growth between the costae. Type *D. antiquissimum*.
- canadense*, Whiteaves, 1897, Pal. Foss., vol. 3, p. 227, Low. Sil.
- grastonense*, Meek and Worthen, instead of *Lituites grastonensis*.
- marshi*, Hall, instead of *Lituites marshi*.

aliocheras undul-
is, instead of Try-

1897, Geo. Sur.
Trenton Gr.

Proc. Am. Phil.
[Ety. *dellos*, a
Shell similar to
more complicated.
section, but grow
ventro-dorsal diam-
large and ventral.
the earlier epim-
throughout the epi-
ture from the spiral
e in the gerontic
netlines the entire
e. No impressed
um.

Proc. Am. Phil.
Calciferous Gr.
us twice preoccu-
used it in 1825, in
e 41. His defini-
stration, and two
genus. McCoy, in
n his Synop. Carb.
the genus defined
s proposed *Dicito-*

cheras ammonis.
ntrocheras marcel-

3, Proc. Bost. Soc.
s. 292. It includes
gonal whorls, hav-
ightly convex, sides
um very gibbous,
impression. The
ngitudinally, with
se striae, not sub-
s have ventral and
ad dorsal saddles,
r lobes. Siphon
Living chamber
three-fourths of
ure with a deep
be *D. costellatum*.
e referred to this

Proc. Am. Phil.
00. [Ety. *diakos*,
Discoid, planor-
both sides. Volu-
s, slightly embrac-
section nearly cir-
ked by distant coe-
sides obliquely
more strongly on
eating a profound
abrificating striae of
e coste. Type *D.*

1897, Pal. Foss.,
il.
Worthen, instead
nsis.

of *Lituites marshi*.

multicostatum, Whitfield, instead of *Lit-*
uites multicostatus.

ortoni, instead of *Lituites ortoni*.
Domatoceras, Hyatt. The whorl in sec-
tion is distinctly hexagonal in a ma-
ture state.

militarium, Hyatt, 1893, 4th Ann. Rep.
Geo. Sur. Texas, p. 445, Coal Meas.
simplex, Hyatt, 1893, 4th Ann. Rep. Geo.
Sur. Texas, p. 441, Coal Meas.

Edaphoceras, Hyatt, 1883, Proc. Bost.
Soc. Nat. Hist., vol. 22, p. 288. [Ety.
edaphos, a seat; *keras*, horn.] Young
shells arcuate until a late stage of
growth, with whorls fusiform in sec-
tion, and sutures with dorsal and ven-
tral lobes and angular lateral saddles;
but the siphon shifts from the venter,
where it is in the larva, to near the
center. The adult is close coiled, with
flattened sides and broad lateral sad-
dles. Type *E. niotense*.

niotense, Meek and Worthen, 1865,
(*Temnochilus niotense*.) Proc. Acad.
Nat. Sci. Phil., p. 260, and Geo. Sur.
Ill., vol. 5, p. 523, Keokuk Gr.

Endoceras. The type is *E. subcentrale*.
aulema, Clarke, 1897, Geo. Sur. Minn.,
vol. 3, p. 770, (Nanno *aulema*.) Trenton
Gr.

consuetum, Sardeson, 1896, Bull. Minn.
Acad. Nat. Sci., vol. 4, p. 103, Shak-
opee Dolomite.

henneplini, Clarke, 1897, (*Cameroceras*
henneplini.) Geo. Sur. Minn., vol. 3, p.
779, Galena Gr.

Endolobus gibbosus, Hyatt, refer to *Stea-*
roceras gibbosum.

Ephippioceras, Hyatt, 1883, Proc. Bost.
Soc. Nat. Hist., vol. 22, p. 290. [Ety.
ephippion, a saddle; *keras*, horn.] Shell
coiled and having subacute, prominent
ventral saddles; broad lateral lobes;
subacute lateral saddles near the
shoulders, and broad, shallow dorsal
lobes. Septa creased or raised into a
median ridge between the two saddles.
Type *E. ferratum*.

montgomeryensis, instead of *Nautilus*
montgomeryensis.

Eury stomites, Schröder, 1891, Pal. Abh.
Dames et Kayser, vol. 5, p. 26. [Ety.
eurya, broad; *stoma*, mouth; *lithos*,
stone.] Discoid, much like *Tarphy-*
ceras. Siphuncle subventral in the
nepionic and aneanian substages, be-
coming extracentroventral in all the
later stages of development, or it may
remain nearer the venter. Growth
more rapid than in *Tarphyceras*, fewer
whorls in the same diameter, and
ventro-dorsal diameter longer. Whorl
may be rounded in early life, but ac-
quires a more or less flattened ven-
ter and primitive lateral and ill-defined
umbilical zones. Umbilical perfora-
tion large, and the impressed zone is
a contact furrow not generated until
the whorls come in contact. The con-

tact furrow is deeper than in *Tarphy-*
ceras, body chamber free and variable
in length. The aperture has lateral
crests, most prominent opposite the
centers of the lateral zones. Sutures
straight or sinuous. Type *E. kelloggi*.
apollo, instead of *Lituites apollo*.
gibbosum, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 443, Calciferous Gr.
Imperator, instead of *Lituites imper-*
ator.

kelloggi, Whitfield, 1886, (*Nautilus kel-*
loggi.) Bull. Am. Mus. Nat. Hist., vol.
1, p. 328, Calciferous Gr.

plicatus, Whiteaves, 1896, Can. Rec. Sci.,
vol. 6, p. 395, Low. Sil.

robertsoni, instead of *Lituites robert-*
soni.

rotundus, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 443, Calciferous Gr.

undatus, Emmons, 1842, (*Inachus unda-*
tus.) Geo. Rep. N. Y. 394, and Pal.
N. Y., vol. 1, p. 52, Black Riv. Gr.

undatus, var. *occidentalis*, Hall, 1861,
(*Lituites undatus* var. *occidentalis*.)
Rep. of Progr. Wis., p. 38, Black Riv.
Gr.

virginianus, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 444, Calciferous Gr.

Gastrioceras branneri. See *Goniattites bran-*
neri.

Glyptioceras, Hyatt, 1883, Proc. Bost. Soc.
Nat. Hist., vol. 22, p. 328. [Ety. *glyphus*,
the notch in an arrow; *keras*, horn.]
Whorls in section semilunar, trape-
zoidal, or compressed; abdomen broad
and convex; sides divergent outwardly,
and frequently costated. Sutures with
acute angular lobes and saddles, and
siphonal saddle frequently bottle-
shaped. Siphonal saddles small, and
occupy only the apex of the straight-
sided, deep ventral lobes. First pair
of saddles spatulate. Type *G. cren-*
istria.

summinsi, Hyatt, 1893, 4th Ann. Rep.
Geo. Sur. Texas, p. 467, Coal Meas.

ineisum, Hyatt, 1893, 4th Ann. Rep. Geo.
Sur. Texas, p. 471, Coal Meas.

Gomphoceras amphora, Hyatt, and scioto-
ense are described and illustrated in
Ohio Geol., vol. 7, pp. 428 and 533.

indianense, Miller and Faber, 1894, Jour.
Cin. Soc. Nat. Hist., vol. 17, p. 137,
Hud. Riv. Gr.

mitriforme, Clarke, 1894, 13th Rep. St.
Geol. N. Y., p. 171, Marcellus Shale.

ortoni, Foerste, 1895, Ohio Geol., vol. 7,
p. 533, Niagara Gr.

Goniattites blairi, Miller and Gurley, 1896,
Bull. No. 11, Ill. St. Mus. Nat. Hist.,
p. 35, Chouteau Gr.

branneri, Smith, 1897, (*Gastrioceras*
branneri.) Marine Foss. from Coal
Meas., p. 47, Lower Coal Meas.

olkhornensis, Miller and Gurley, 1896,
Bull. No. 11, Ill. St. Mus. Nat. Hist.,
p. 37, Coal Meas.

fultonensis, Miller and Gurley, 1896, Bull.

No. 11, Ill. St. Mus. Nat. Hist., p. 39, Coal Meas.

Fig. 1419.—*Goniatites greenenstlensis*, internal and ventral views.

greenenstlensis, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 44, St. Louis Gr.

Fig. 1420.—*Goniatites illinoisensis*, internal and ventral views.

illinoisensis, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 42, Coal Meas.

jessleae, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 46, Chouteau Gr.

kansasensis, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 43, Up. Coal Meas.

Fig. 1421.—*Goniatites kentuckiensis*, showing outer shell.

kentuckiensis is redescribed and illustrated in Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 40.

louisianensis Rowley, 1895, Am. Geol., vol. 16, p. 221, Chouteau Gr.

lunatus, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 41, Coal Meas.

montgomeryensis, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 38, Coal Meas.

parrishii, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 36, Up. Coal Meas.

Fig. 1422.—*Goniatites subcavus*, internal and ventral views.

subcavus, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 45, Coal Meas.

GYROCERAS baeri, refer to *Trochoceras baeri*, where it was first referred.

columbiense and *seminodosum* are described and illustrated in Ohio Geol., vol. 7, pp. 430 and 431.

elrodi, refer to *Barrandoceras elrodi*.

inelegans, *liratum*, and *subliratum*, refer to *Nephriticeras*.

undulatum refer to *Halloceras undulatum*.

HALLOCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 284. [Ety. proper name; *keras*, horn.] Whorls triangular in section, abdomen broad, sides divergent, dorsum narrow. Thick costae or large nodes on the angles of the sides. Sutures have ventral and lateral lobes. Siphon small and near the venter. Type *H. undulatum*. *undulatum*, Vanuxem, 1842, (*Cyrtoceras undulatum*,) Geol. Rep. N. Y., p. 139, and Pal. N. Y., vol. 5, pt. 2, p. 378, Up. Held. Gr.

LITOCERAS, Hyatt, 1894, Proc. Am. Phil. Soc. vol. 32, p. 474. [Ety. *litos*, plane; *keras*, horn.] Shells discoid. Siphuncle dorsal or below the center in adults, but ventral in the neanic and earlier stages. Umbilicus of good size. Whorls larger and broader than in *Schroederoceras*, and have in the ephibic stage similar abdomens and convex, divergent sides, without umbilical shoulders. The aperture is less compressed than in *Schroederoceras*, but not flaring as in *Trocholites*. Hyponomie sinus smaller and shallower than in *Schroederoceras*, and the contact furrow broader and deeper. Sutures have deep dorsal lobes, saddles on the lines of involution, and broad lateral lobes. Type *L. whiteavesi*. Not *L. versutum*, as in the first definition of the genus, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 268.

Gurley, 1896, Bull.
Nat. Hist., p. 41,

and Gurley,
Ill. St. Mus. Nat.
Hist., p. 36.

Gurley, 1896, Bull.
Nat. Hist., p. 36.

cervus, lateral and
ventral.

Gurley, 1896, Bull.
Nat. Hist., p. 45,

to Trochoceras
first referred.

modosum are de-
scribed in Ohio Geol.,
vol. 1.

Discoceras elrodi.
subliratum, refer

Discoceras undula-

Proc. Bost. Soc.
Nat. Hist., p. 284. [Ety.

, horn.] Whorls
abdomen broad,
umbilical narrow. Thick

on the angles of
have ventral and
small and near

undulatum.

1842, (Cytroceras
Rep. N. Y., p. 139,
vol. 5, pt. 2, p. 378,

Proc. Am. Phil.
[Ety. *litos*, plane;

discoid. Siphon
center in adults,
umbilical and earlier

of good size,
broader than in
the ephebic

and convex, di-
st umbilical shoul-
ders less compressed

as, but not flaring
Hyponomic sinus
than in *Schroeder-*

furrow broader
have deep dorsal
lines of involu-

crural lobes. Type *L.*
versutum, as in the

genus, 1883,
Nat. Hist., vol. 22, p.

biangulatum, Hyatt, 1894, Proc. Am.
Phil. Soc., vol. 32, p. 479, Calciferous Gr.

hercules, Billings, 1856, (Nautilus her-
cules,) Rep. Geol. Sur. Can., p. 300,
Hud. Riv. Gr.

insolens, Billings, 1865, (Nautilus ins-
olens,) Pal. Foss., vol. 1, p. 258, Calcif-
erous Gr.

versutum, Billings, 1865, (Nautilus versu-
tum,) Pal. Foss., vol. 1, p. 259, Quebec Gr.

whiteavesi, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 475, Calciferous Gr.

Lituites, Breynius, 1732, Dissertatio Phys-
ica de Polythalamis, p. 27, Montfort,
1806. This genus is unknown in Amer-
ica, and therefore all the references of
American authors to it have been er-
roneous. Professor Hyatt has referred
most of the species to other genera,
and all of them must be so distributed.

americanus, D'Orbigny, Syn. for Bar-
randoceras convolvans.

apollo, refer to Eurytomites apollo.

bickmoreanus. See Plectoceras bickmore-
anum.

complanatus. See Aphetoceras complan-
atum.

convolvans. See Barrandoceras con-
volvans.

entoni. See Schroedero-ceras entoni.

entoni var. cassinensis. See Schroedero-
ceras cassinense.

farnsworthi. See Aphetoceras farns-
worthi.

farnsworthi, in part. See Tarphyceras
farnsworthi.

grastonensis. See Discoceras grastonense.

imperator. See Eurytomites imperator.

internistriatus. See Trocholithes interni-
striatus.

magnificum. See Aspidoceras magnificum.

marshi. See Discoceras marshi.

multicostatus. See Discoceras multicos-
tatum.

niagarensis, Spencer. Too poorly de-
fined to be recognized.

ortoni. See Discoceras ortoni.

palinurus. See Schroedero-ceras pali-
nurus.

pluto, Billings. Not defined so as to be
recognized.

robertsoni. See Eurytomites robertsoni.

seeleyi. See Tarphyceras seeleyi.

undatus. See Eurytomites undatus.

undatus var. occidentalis. See Eurytom-
ites undatus var. occidentalis.

MITROCERAS, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 503. [Ety. *mitra*, tur-
ban; *keras*, horn.] This name is pro-
posed instead of *Trochoceras*, with
Trochoceras gebhardi as the type, on
the ground that Barrande had pre-
occupied *Trochoceras* for a different
genus, though Barrande and Hall
thought their species belonged to the
same genus.

MELONOCERAS, Hyatt, 1883, Proc. Bost.
Soc. Nat. Hist., vol. 22, p. 280. [Ety.
melon, goat; *keras*, horn.] Arcuate

cones, section compressed, ovate; dor-
sum wider than venter. Siphon near
the venter. Sutures have ventral and
dorsal saddles, and slight lateral lobes.
Living chamber short. Aperture en-
tirely open, and partially subtriangular
or contracted and pear-shaped. Type
M. prematurum.

metellus, Billings, 1865, (Cytroceras
metellus,) Pal. Foss., vol. 1, p. 191,
Quebec Gr.

prematurum, Billings, 1866, (Phragmo-
ceras prematurum,) Can. Nat. and
Geol., vol. 5, p. 173, Black Riv. Gr.

Naumo, Clarke, 1894, Am. Geologist, vol.
14, p. 205. A generic name, signifying
a player upon the flute, applied to
what has been generally called the
siphuncle of an *Endoceras*, or what I
regard as the body chamber, and years
ago described under the names of *E.*
egani, *E. bristolense*, and *E. inequi-*
abile. Type *N. aulema*.

aulema. See *Endoceras aulema*.

NAUTILUS *acervus*. See *Nephriticeras*
acervum.

buchinum. See *Nephriticeras buchinum*.

cancellatus, McChesney, not a *Nautilus*.

cervus. See *Nephriticeras cervum*.

champlainense. See *Tarphyceras cham-*
plainense.

cornutum. See *Rhadinoceras cornutum*.

divisus. See *Ephippioceras divisum*.

ferratus. See *Ephippioceras ferratum*.

hercules. See *Litoceras hercules*.

hyatti. See *Rhadinoceras hyatti*.

insolens. See *Litoceras insolens*.

jason. See *Plectoceras jason*.

kelloggi. See *Eurostomites kelloggi*.

liratus. See *Nephriticeras liratum*.

liratus var. juvene. See *Nephriticeras*
juvene.

magister. See *Nephriticeras magister*.

maximum. See *Nephriticeras maximum*.

montgomeryensis. See *Ephippioceras*
montgomeryense.

natator. See *Barrandoceras natator*.

nodocarinatus, McChesney, 1859, Desc.
New Spec. Foss., p. 66, Coal Meas., is
a good species, but not a *Nautilus*.

oriens. See *Nephriticeras oriens*.

ortoni, pauper, and subquadrangularis
are described and illustrated, in Ohio
Geol., vol. 7, pp. 481, 486, and 487.

ponderosus. See *Titanoceras ponderosum*.

quadrangulus, McChesney, instead of
quadrangularis. See *Tainoceras quad-*
rangulum.

subliratus. See *Nephriticeras subliratum*.

versutum. See *Litoceras versutum*.

NEPHRITICERAS, Hyatt, 1883, Proc. Bost.
Soc. Nat. Hist., vol. 22, p. 300. [Ety.
nephrites, kidney-shaped; *keras*, horn.]
Whorls kidney-shaped or elliptical.
Umbilicus large, impressed zone on
the dorsum. Siphon nannuloid and
above the center. Sutures have broad
ventral saddles and slight ventral
lobes. Dorsal V-shaped annular lobes

in the impressed zone. Living chamber from one fourth to half a volution.

Type *N. bucinum*.

- acream, Hall, 1879, (*Nautilus acreams*), Pal. N. Y., vol. 5, pt. 2, p. 417, Ham. Gr. bucinum, Hall, 1876, (*Nautilus bucinum*), Illust. Dev. Foss., pl. 60, and Pal. N. Y., vol. 5, pt. 2, p. 412, Ham. Gr. cavum, Hall, 1879, (*Nautilus cavum*), Pal. N. Y. vol. 5, pt. 2, p. 416, Ham. Gr. inelegans, Meek, 1871, (*Gyroceras inelegans*), Proc. Acad. Nat. Sci., p. 89, and Ohio Pal., vol. 1, p. 232, Up. Held. Gr. juvene, Hall, 1879, (*Nautilus liratus* var. juvenis), Pal. N. Y., vol. 5, pt. 2, p. 411, Ham. Gr. liratum, Hall, 1880, (*Gyroceras liratum*), 13th Rep. N. Y. Mus. Nat. Hist., p. 104, Marcellus Shale. magister, Hall, 1879, (*Nautilus magister*), Pal. N. Y., vol. 5, pt. 2, p. 422, Ham. Gr. maximum, Conrad, 1838, (*Cyrtoceras maximum*), Ann. Rep. N. Y., p. 117, and Pal. N. Y., vol. 5, pt. 2, p. 418, Ham. Gr. oriens, Hall, 1876, (*Nautilus oriens*), Illust. Dev. Foss., pl. 61, and Pal. N. Y., vol. 5, pt. 2, p. 420, Ham. Gr. subliratum, Hall, 1876, (*Gyroceras subliratum*), Illust. Dev. Foss., pl. 58, and Pal. N. Y., vol. 5, pt. 2, p. 409, Ham. Gr. ONCOCERAS carveri, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 801, Trenton Gr. douglassi, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 801, Galena Gr. minnesotense, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 798, Galena Gr.

ORTHOCERAS albersi, Miller and Faber, 1894,

Fig. 1423.—*Orthoceras albersi*.

Jour. Cin. Soc. Nat. Hist., vol. 17, p. 140, Hud. Riv. Gr.

- beltrami, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 789, Galena Gr. caldwelli, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 31, Up. Held. Gr. crawfordi, Foerste. Too poorly defined to be recognized. dantonense, Foerste. Too poorly defined to be recognized. erraticum, Foerste. Too poorly defined to be recognized. fenestrulatum, Clarke, 1894, 13th Rep. St. Geol. N. Y., p. 168, Marcellus Shale. geneva, Clarke, 1894, 13th Rep. St. Geol. N. Y., p. 168, Corniferous Gr. hanoverense, Foerste. Too poorly defined to be recognized. ignotum, Foerste. Too poorly defined to be recognized. incarcerationum, Clarke, 1894, 13th Rep. St. Geol. N. Y., p. 170, Marcellus Shale. inceptum, Foerste. Too poorly defined to be recognized. lata-nummulatum, Foerste. Too poorly defined to be recognized. leseuri, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 785, Trenton Gr. ludlowense, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, p. 139, Hud. Riv. Gr. minnesotense, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 77, St. Peter Sandstone. nicolleti, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 784, Trenton Gr. nova-carlisle, Foerste. Too poorly defined to be recognized. nuntioides, Clarke, 1894, 13th Rep. St. Geol. N. Y., p. 170, Marcellus Shale. perroti, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 785, Hud. Riv. Gr. spyroceroides, Foerste. Too poorly defined to be recognized. staffordense, Clarke, 1894, 13th Rep. N. Y. St. Geol., p. 169, Marcellus Shale. turgido-nummulatum, Foerste. Too poorly defined to be recognized. youngi, Foerste, 1895, Ohio Geol., vol. 7, p. 537, Niagara Gr. PHRAGMOCERAS prematurum, refer to Melonoceras prematurum. PLEOCERAS corniculum, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 103, Oneota Dolomite. winchelli, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 787, Calciferous Gr. [It is spelled newton-winchelli, but as that is not binomial I have dropped the middle name newton, and probably the whole name should be stricken out for not conforming to the binomial system.] PLEOTOCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 268. [Ety. *plektos*, twisted or plaited; *keras*, horn.] Whorls quadrate, abdomen narrower than dorsum, and sides convergent outward. Surface having costae curved

, Geo. Sur. Minn.,
Gr.
Gurley, 1896, Bull.
Nat. Hist., p. 31,
Too poorly defined
Too poorly de-
ced.
Too poorly defined
e, 1894, 13th Rep.
8, Marcellus Shale.
13th Rep. St. Geol.
ferous Gr.
Too poorly de-
ced.
Too poorly defined to
e, 1894, 13th Rep.
p. 170, Marcellus
Too poorly defined
erste. Too poorly
ized.
Geo. Sur. Minn.,
on Gr.
and Faber, 1894,
t. Hist., vol. 17, p.
on, 1896, Bull. Minn.
l. 4, p. 77, St. Peter
7, Geo. Sur. Minn.,
on Gr.
erste. Too poorly
ized.
1894, 13th Rep. St.
Marcellus Shale.
Geo. Sur. Minn.,
Riv. Gr.
e. Too poorly de-
ced.
1894, 13th Rep. N. Y.
arcellus Shale.
Foerste. Too poorly
ized.
Ohio Geol., vol. 7,
rum, refer to Mælo-
m.
Sardeson, 1896,
Nat. Sci., vol. 4, p.
te.
7, Geo. Sur. Minn.,
ferous Gr. [It is
schelli, but as that
have dropped the
ton, and probably
should be stricken
forming to the bi-
33, Proc. Bost. Soc.
p. 268. [Ety. *plek-*
ted; *keras*, horn.]
abdomen narrower
sides convergent
aving coste curved

posteriorly and crossing the abdomen.
Siphon ventral and holochonoidal.
Type *P. jason*.
bickmoreanum, Whitfield, 1885, (Litu-
ites bickmoreanus.) Bull. Am. Mus.
Nat. Hist., vol. 1, p. 191, Niagara
Gr.
jason, Billings, 1859, (Nautilus jason,)
Can. Nat. and Geol., vol. 4, p. 164,
Chazy Gr.
obscurum, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 499, Black Riv. Gr.
POTEROCCERAS jerseyense, Miller and Gur-
ley, 1896, Bull. No. 11, Ill. St. Mus.
Nat. Hist., p. 32, Kinderhook Gr.
PYCNOCCERAS, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 454. [Ety. *puknos*,
close; *keras*, horn.] This genus is very
much like *Spherocceras*. Type *P. aper-*
tum.
apertum, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 455, Calciferous Gr.
caliciforme, Hyatt, 1894, Proc. Am.
Phil. Soc., vol. 32, p. 456, Calciferous
Gr.
REMELECCERAS, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 525. [Ety. proper
name; *keras*, horn.] Wide umbilicus,
perforated, shallow contact furrow,
having V-shaped sutures. Transverse
section elliptical. Type *R. impres-*
sium.
clarkense, Miller and Gurley, Bull. No.

Fig. 1421.—Remelecceras clarkense, internal view.

12, Ill. St. Mus. Nat. Hist., p. 49, Keo-
kuk Gr.
impressum, Hyatt, 1894, Proc. Am. Phil.
Soc., vol. 32, p. 525, Keokuk or Wa-
verly Gr.

RHADIACCERAS, Hyatt, 1894, Proc. Am.
Phil. Soc., vol. 32, p. 530. [Ety. *radi-*
nos, slender; *keras*, horn.] More
slender than *Nephritoceras*; discoid
whorls compressed el-
liptical, or rounded,
and have an impressed
zone in mature shells.
Type *R. cornu-*
lulum.
cornulum, Hall, 1876,
(Nautilus cornulum,)
Illust. Dev. Foss., pl. 60,
and Pal. N. Y., vol. 5,
pt. 2, p. 414, Ham. Gr.
hyatti, Beecher,
1888, (Nautilus hyatti.) Pal.
N. Y., vol. 7, p. 37, Ham.
Gr.
SCHROEDEROCERAS, Hyatt, 1894, Proc. Am.
Phil. Soc., vol. 32, p. 458. [Ety. proper
name; *keras*, horn.] Type *S. angula-*
tum. It includes forms that have been
referred to *Lituites*, and is related to
Litoceras, and *Trocholites*. Umbilical
perforation large; whorls few; sutures
sinuous; living chamber short.
casinense, Whitfield, 1886, (Lituites
eatoni var. casinensis.) Bull. Am. Mus.

Fig. 1425.—Remelecceras clarkense, dorsal view.

Fig. 1426.—Solenochilus henryvillense, lateral view.

Nat. Hist., vol. 1, p. 332, Calciferous
Gr.
eatoni, Whitfield, 1886, (Lituites eatoni.)

Bull. Am. Mus. Nat. Hist., vol. 1, p. 331, Calciferous Gr.
 pallinurus. Billings, instead of *Lituites pallinurus*.

SOLENOCHILUS kentuckiense, Hyatt, 1893, 4th Am. Rep. Geo. Sur. Texas, p. 461, Coal Meas.

henryvillense, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 51, Keokuk Gr.

FIG. 1427.—*Solenochilus henryvillense*, dorsal view.

STEAROCERAS, Hyatt, 1893, Geo. Sur. Texas, 4th Rep., p. 422. Distinguished from *Endolobus* by the deep, narrow umbilici; slight, shallow lobes on the venter, and small dorsal and annular lobes. Type *S. gibbosum*.

gibbosum, Hyatt, 1890, (*Endolobus gibbosus*,) Geo. Sur. Texas, 2d Rep., p. 353, Coal Meas.

TAINOCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 269. [Ety. *tainia*, a head-band; *keras*, horn.] Discoid, whorls quadrate, and having on each side, and also on the abdomen, two rows of tubercles. Siphon above the center. Type *T. quadrangulum*.

duttoni, Hyatt, 1893, Geo. Sur. Texas, 4th Rep., p. 401, Coal Meas.

quadrangulum, McChesney, 1865, (*Nautilus quadrangulus*,) Desc. New Pal. Foss., p. 65, Coal Meas.

TARPHYCERAS, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 433. [Ety. *tarphus*, close; *keras*, horn.] Resembles *Eury-stomites*, but more discoidal, more numerous, and more slowly growing whorls, longer living chamber; the whorls sometimes flattened on the abdomen and approximating a quadrangular form, and the aperture is like that of *Trocholites*, with a deep broad hypomic sinus. Type *T. prematurum*.

meoini, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 435, Calciferous Gr.

champlainense, Whitfield, 1886, (*Nautilus champlainensis*,) Bull. Am. Mus. Nat. Hist., vol. 1, p. 329, Calciferous Gr.

extensum, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 438, Calciferous Gr.

farnsworthi, Billings, 1861, (*Lituites farnsworthi*, in part,) Pal. Foss., vol. 1, p. 21, Calciferous Gr.

macdonaldi, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 435, Calciferous Gr.

prematurum, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 437, Calciferous Gr.

seeleyi, Whitfield, 1886, (*Lituites seeleyi*,) Bull. Am. Mus. Nat. Hist., vol. 1, p. 330, Calciferous Gr.

TEMNOCHILUS greenense, Miller and Gurley, 1897, Bull. No. 12, Ill. St. Mus. Nat. Hist., p. 52, Kaskaskia Gr.

THINOCERAS, Hyatt, 1893, 4th Ann. Rep. Texas, p. 430. Shells large and retaining the longitudinal ridges. Sutures similar to those of *Discitoceras*, except that they correlate with the broader venter and lateral zones of the whorls, the lobes being broader and shorter, especially on the zones, than in *Discitoceras*. Type *T. depressum*.

kentuckiense, Hyatt, 1893, 4th Ann. Rep. Geo. Sur. Texas, p. 432, Subcarboniferous.

depressum, Hyatt, 1893, 4th Ann. Rep. Geo. Sur. Texas, p. 430, Subcarboniferous.

TITANOCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 289. [Ety. *Titan*, mythological name; *keras*, horn.] The whorl has a narrower abdomen than in *Apsidoceras*, and longer abdoindorsal diameter, and is more compressed or shield-shaped. There is a narrow impressed zone on the dorsum, and an undivided, narrow dorsal lobe. Type *T. ponderosum*.

ponderosum, White, 1872, (*Nautilus ponderosus*,) Pal. E. Neb., p. 236, Coal Meas.

TREMATOCERAS ohioense is described and illustrated in Ohio Geol., vol. 7, p. 426.

TRIPTEROCERAS, Hyatt, 1883, Proc. Bost. Soc. Nat. Hist., vol. 22, p. 287. [Ety. *tripter*, a rubbing tool; *keras*, horn.] Lateral saddles acute; venter flattened and broader than the dorsum which forms the apex of the subtriangular section. Siphon ventral, nummuloidal; whorl arcuate in the young, but straight in the full-grown. Type *T. hastatum*.

hastatum, instead of *Orthoceras hastatum*, and to this genus may also be referred *O. planoconvexum*.

oweni, Clarke, 1897, Geo. Sur. Minn., vol. 3, p. 792, Trenton Gr.

TROCHOCERAS bneri, instead of *Gyroceras bneri*.

TROCHOLITES canadensis, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 486, Trenton Gr.

dyceri, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 489, Hud. Riv. Gr.

internastriatus, Whitfield, 1886, (*Lituites internastriatus*,) Bull. Am. Mus. Nat. Hist., vol. 1, p. 332, Calciferous Gr.

TROCHOLITOCERAS, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 480. [Ety. from *Trocholites*, a genus; *keras*, horn.] This is distinguished from *Trocholites* only by having the siphuncle ventral of the center in the earlier substages of development. Type *T. walcotti*.

walcotti, Hyatt, 1894, Proc. Am. Phil. Soc., vol. 32, p. 480, Calciferous Gr.

CLASS LAMILLIBRANCHIATA.

386. (Lituites see-
s. Nat. Hist., vol.
s Gr.

, Miller and Gur-
12, Ill. St. Mus.
iskaskia Gr.

893, 4th Ann. Rep.
is large and retain-
al ridges. Sutures
f *Discitoceras*, ex-
rrelate with the
lateral zones of the
being broader and
on the zones, than
pe *T. depressum*.

1893, 4th Ann. Rep.
432, Subcarbonif-

893, 4th Ann. Rep.
430, Subcarbonif-

83, Proc. Bost. Soc.
289. [Ety. *Titan*,
; *keras*, horn.] The
wer abdomen than
longer abdomino-
nd is more com-
haped. There is a
one on the dorsum,
narrow dorsal lobe.

1872, (Nautilus
Neb., p. 236, Coal

a is described and
Geol., vol. 7, p. 426.
1883, Proc. Bost.
122, p. 287. [Ety.
tool; *keras*, horn.]
cute; venter flat-
than the dorsum
pex of the subtri-
phon ventral, num-
euate in the young,
full-grown. Type

Orthoceras hastat-
genus may also be
vexum.

(Geo. Sur. Minn.,
on Gr.

tend of *Gyroceras*

, Hyatt, 1894, Proc.
2, p. 486, Trenton Gr.
oc. Am. Phil. Soc.,
Riv. Gr.

field, 1886, (Lit-
s.) Bull. Am. Mus.
332, Calciferous Gr.
t, 1894, Proc. Am.
p. 480. [Ety. from
is; *keras*, horn.]
d from *Trocholites*
siphuncle ventral
e earlier substages
ype *T. walcotti*.
Proc. Am. Phil.
Calciferous Gr.

Actinomya, Ulrich, Syn. for *Modiolopsis*.
kentonensis, Syn. for *Modiolopsis cin-*
cinnatiensis.

subcarinata, Syn. for *Modiolopsis mod-*
ioliformis.

Allodesma, Ulrich, 1894, Geo. Sur. Minn.,
p. 617. Probably founded on a poor
cast of an *Orthodesma*. There are no
characters ascribed to it that can be
called generic. The author of the ge-
nus refers it to the *Cycloconchidae*, and
attempts to establish it, upon the forms
he had shortly before described, with
confidence, under the name of *Modio-*
opsis subelliptica.

Allonychia, Ulrich, Syn. for *Ambonychia*.
Founded on *Ambonychia jamesi*.

ovata, Syn. for *Ambonychia jamesi*.

subrotunda, Syn. for *Ambonychia jamesi*.

ALLORISMA andrewsi and *A. maxvillensis* are
illustrated in Ohio Geol., vol. 7, p. 475.
pleuropistha, refer to *Pholadella pleuro-*
pistha.

AMBONYCHIA affinis, Ulrich, Syn. for *A.*
planistriata.

casei, refer to *Opisthoptera casei*.

cincinnatiensis, Miller and Faber, 1894,
Jour. Cin. Soc. Nat. Hist., vol. 17, p.
24, Hud. Riv. Gr.

excavata, Ulrich, 1895, (*Clionychia ex-*
cavata.) Ohio Geol., vol. 7, p. 651, Hud.
Riv. Gr.

obesa, Ulrich, 1895, (*Byssonychia obesa*.)
Ohio Geol., vol. 7, p. 630, Hud. Riv. Gr.
This species was founded upon poor
casts that have generally been regarded
as *A. radiata*, but it may be a distinct
species.

perangulata, Ulrich, 1895, (*Psilonychia*
perangulata.) Ohio Geol., vol. 7, p. 649,
Hud. Riv. Gr. This species is of very
doubtful value.

subundata, Ulrich, 1895, (*Clionychia sub-*
undata.) Ohio Geol., vol. 7, p. 651, Utica
Slate Gr.

tenuistriata, Ulrich, 1894, (*Byssonychia*
tenuistriata.) Geol. of Minn., p. 500,
Hud. Riv. Gr.

ANOMALODONTA plicata, Ulrich, Syn. for
Ambonychia costata.

Auoptera, Ulrich, Founded upon poor
casts of *Angellum cuneatum*.

miseneri, Ulrich, Syn. for *Angellum cun-*
eatum.

Anthracoptera longa, Dawson, 1894, (*Nai-*
adites longus.) Can. Rec. Sci., Coal
Mens.

mytiloides, Dawson, 1894, (*Naiadites my-*
tiloides.) Can. Rec. Sci., Coal Mens.

ARISTERELLA, Ulrich, 1894, Geo. Sur. Minn.,
p. 524. [Ety. *aristera*, the left; *ella*,
diminutive.] The left valve is smaller
than the right. Founded on small
shells of uncertain relations. Type *A.*
nitidula, described at the same place
from the Trenton Gr.

ASTHENODONTA Whiteaves, 1893, Trans.
Roy. Soc. Can., p. 23. [Ety. *asthene's*,
weak; *odontos*, a tooth.] A large shell
having an outline similar to *Margari-*
tana margaritifera. Type *A. westoni*,
described at the same place from the
Coal Mens.

AVICULA whitfieldi, Foerste, Syn. for *Cyp-*
ricardites ferru-
gineus, Hall and
Whitfield, which
is not an *Avicula*
or a *Cypriocardites*.

AVICULOPECTEN utah-
ensis is figured in
Expl. 40th Par-
allel, vol. 4, p. 95.

BLATRELLA, Miller and
Gurley, 1896, Bull.
No. 11, Ill. St.
Mus. Nat. Hist.,
p. 6. [Ety. proper

name.] Equi-
valve, inequali-
teral, elliptical, subovate, or subcir-
cular. Beaks anterior incurved. Um-

Fig. 142.—*Blatrella sedallensis*, right valves of
two casts.

Fig. 143.—*Blatrella sedallensis*, right valves of
two casts.

bones high. Cardinal line straight,
posterior to the beaks. Margins closed.
Ligament exter-
nal. A pit be-
neath the beak
of the right
valve, a single
tooth anterior,
and a bifid tooth
posterior. Type
B. sedallensis.

sedallensis, Miller and Gurley, 1896, Bull.

Fig. 144.—*Blatrella sedallensis*, cardinal
view.

No. 11, Ill. St. Mus. Nat. Hist., p. 7, Chouteau Gr.

Fig. 1431.—*Blutella sedallensis*, hinge-line and outside of same valve.

BODMANNIA, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, p. 23. [Ety. proper name.] Equivalve, profoundly inequilateral; ventricose; general outline like *Cypriocardites*; beaks anterior, incurved; hinge-line short, at a high angle to the base of the shell; ligament external; shell thin, concentrically lined. Type *B. inequetum*.

insuetum, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, page 23, Hud. Riv. Gr.

ventricosa, instead of *Edmondia ventricosa*, in Pal. N. Y., vol. 1, p. 155, which has been frequently referred to *Cypriocardites*.

BYSSONYCHIA, Ulrich, Syn. for *Ambonychia acutirostris*, *bynesi*, and *imbricata*, Synonyms for *Ambonychia costata*. *alveolata*, *præcurva*, and *suberecta*, Synonyms for *Ambonychia radiata*. *cultrata*, *grandis*, and *richmondensis*, Synonyms for *Ambonychia robusta*. *obesa*, refer to *Ambonychia obesa*. *tenuistriata*, refer to *Ambonychia tenuistriata*. *vera*, Syn. for *Ambonychia cincinnatensis*.

Fig. 1432.—*Chænomyna longa*, left side view.

Fig. 1433.—*Chænomyna longa*, cardinal view.

CHÆNOMYNA longa, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 8, Chouteau Gr.

CLEIDOPHORUS major, Ulrich, refer to *Lyrodesma major*.

CLINOPISTNA antiqua, Whiteaves, 1897, Pal. Foss., vol. 3, p. 185, Low. Sil.

Clionychia, Ulrich, is a synonym for *Ambonychia*.

excavata, refer to *Ambonychia excavata nitida*, Syn. for *Ambonychia lamellosa subundata*, refer to *Ambonychia subundata*.

COLPOMYA, Ulrich, 1894, Geo. Sur. Minn., p. 522. [Ety. *kolpos*, sinus; *Mya*, a genus.] This genus agrees in form, muscular scars, and pallial line with *Modiolopsis*, but is said to differ somewhat, in the cardinal

Fig. 1434.—*Conocardium illianense*.

tooth. Type *C. constricta*, described at the same place from the Trenton Gr.

demissa, Ulrich, 1894, Geo. Sur. Minn., p. 524, Trenton Gr.

Fig. 1435.—*Conocardium parvulum* and *Conocardium exiguum*, magnified two diameters.

CORALLIDOMUS, Whitfield, 1895, Ohio Geol., vol. 7, p. 493. [Ety. *korallion*, a coral; *domos*, a house.] Shell oblong, somewhat modioliform; ligament external; anterior and posterior muscular scars; integral pallial line; burrowing habit of life. Type *C. concentricus*, described at the same place from the Hud. Riv. Gr.

Ctenodonta albertina, *ciagulata*, and *modiosonensis*, Ulrich, Synonyms for *Tellinomya pectunculoides*. *caltrini*, refer to *Tellinomya caltrini*.

carinata, refer to *Tellinomya carinata*.

cuneiformis, refer to *Tellinomya cuneiformis*.

filistriata, Syn. for *Tellinomya levata*.

intermedia, *recurva*, *compressa*, and *similis*, Ulrich, Synonyms for *Tellinomya alta*.

medialis, Ulrich, Syn. for *Tellinomya nitida*.

oviformis, Ulrich. It may be a *Tellinomya*, but the characters given do not describe a species.

perminuta, Ulrich, Syn. for *Palæoconcha obliqua*.

scofieldi, refer to *Tellinomya Scofieldi*.

simulatrix, Ulrich. Not defined so as to be recognized.

socialis, refer to *Tellinomya socialis*.

subnata, Syn. for *Tellinomya ovata*.

CUNÆMYA oblonga, Ulrich, 1894, Geo. Sur. Minn., p. 623, Galena Gr.

Whiteaves, 1897, Pal.
Low. Sil.

Synonym for *Ambonychia*

ambonychia excavata.
ambonychia lamellosa.
ambonychia subun-

Geo. Sur. Minn.,
sinus; *Mya*, a ge-

134.—*Conocardium*
h. flaccidum.

bed at the same
con Gr.

Geo. Sur. Minn.,

CORALLIDOMUS,
Whitfield.
1895, Ohio
Geol., vol. 7,
p. 493. [Ety.
corallion, a
coral; *domos*,
a house.]
Shell ob-

long, some-
times external;
or muscular scars;
; burrowing habit
ventricosus, described
from the Hud. Riv.

gulgata, and *madri-*
rich, Synonyms for
pectunculoides.

to *Tellinomya cul-*

er to *Tellinomya*

fer to *Tellinomya*

i. for *Tellinomya*

curva, *compressa*,
Ulrich, Synonyms
mya alta.

ch, Syn. for *Telli-*
nomya.

ch. It may be a
but the characters
describe a species.
rich, Syn. for *Pa-*
boliqua.

er to *Tellinomya*

rich. Not defined
d.

mya socialis.

linomya ovata.

ch, 1894, Geo. Sur.

Gr.

uncatula, Ulrich, 1894, Geo. Sur. Minn.,
p. 622, Galena Gr.

Cycloconcha ovata, Ulrich, Syn. for *Cyclo-*
concha mediocardinalis.

Cymatonomata, Ulrich, Syn. for *Orthodesma*.
attenuata, *contracta*, and *typicalis*, Syn-
onyms for *Orthodesma cylindricum*.

productifrons, Ulrich, refer to *Ortho-*
desma productifrons.

recta, Ulrich, probably synonymous with
Orthodesma cylindricum.

semistriata, Ulrich, refer to *Orthodesma*
semistriatum.

CYPRICARDELLA

eximia.

Miller and

Gurley,

1896, Bull.

No. 11, Ill. Fig. 1436.—*Cypricardella gor-*
St. Mus. Nat. Hist.,

Nat. Hist.,
p. 15, Chouteau Gr.

CYPRICARDITES affinis, Ulrich, 1894, (*Cyr-*
todonta affinis.) Geo. Sur. Minn., p.
540, Trenton Gr.

amplus, Ulrich, 1894, (*Cyrtodonta amplus*.)
Geo. Sur. Minn., p. 538, Trenton Gr.

billingsi, Ulrich, 1894, (*Cyrtodonta bill-*
ingsi.) Geo. Sur. Minn., p. 538, Tren-
ton Gr.

caswelli, Foerste, Ohio Geol., vol. 7, p.
561, Niagara Gr. It has no resem-
blance to a *Cypricardites*.

descriptus, Sardeson, 1896, Bull. Minn.
Acad. Nat. Sci., vol. 4, p. 70, St. Peter
Sandstone.

dignus, Sardeson, 1896, Bull. Minn. Acad.
Nat. Sci., vol. 4, p. 71, St. Peter
Sandstone.

finitimus, Sardeson, 1896, Bull. Minn.
Acad. Nat. Sci., vol. 4, p. 70, St. Peter
Sandstone.

fragosus, Sardeson, 1896, Bull. Minn.
Acad. Nat. Sci., vol. 4, p. 70, St. Peter
Sandstone.

glabellus, Ulrich, Syn. for *Cypricardites*
niota.

islandicus. This specific name will be
dropped, if Hall's *C. ventricosus* of
1847 is, as we believe, a *Bodmannia*.

minnesotensis, Sardeson, Syn. for *Cypric-*
ardites niota.

miseneri, Ulrich, 1895, (*Ischyrodonta*
miseneri.) Ohio Geol., vol. 7, p. 675,
Hud. Riv. Gr.

ovalis, Ulrich, 1895, (*Ischyrodonta ovalis*.)
Ohio Geol., vol. 7, p. 674, Hud. Riv.
Gr.

persimilis, Ulrich, 1894, (*Cyrtodonta*
persimilis.) Geo. Sur. Minn., p. 544,
Trenton Gr.

sterlingensis, Meek and Worthen, refer to
Whitella sterlingensis.

subovatus, Ulrich, 1894, (*Cyrtodonta*
subovata.) Geo. Sur. Minn., p. 536,
Trenton Gr.

sulcodorsatus, Ulrich, 1894, Geo. Sur.
Minn., p. 626, Hud. Riv. Gr.

ventralis, Ulrich, 1894, (*Saffordia ven-*

tralis.) Geo. Sur. Minn., p. 626, Hud.
Riv. Gr.

Cyrtodonta affinis, Ulrich, refer to *Cypric-*
ardites affinis.

amplus, Ulrich refer to *Cypricardites*
amplus.

billingsi, Ulrich, refer to *Cypricardites*
billingsi.

gibbera, and *obesa*, Ulrich, Synonyms for
Cypricardites rotundatus.

jaesvillensis, Ulrich, Syn. for *Cypricar-*
ardites hronensis.

parra, Ulrich, Syn. for *Modiolopsis*
plana.

persimilis, Ulrich, refer to *Cypricardites*
persimilis.

rotulata, Ulrich, Syn. for *Cypricardites*
niota.

suborata, refer to *Cypricardites subo-*
vatus.

DOLABRA. The type of the genus is *D.*
angusta.

sterlingensis, refer to *Whitella sterling-*
ensis.

EDMONDIA albersi, Miller and Gurley, 1896,
Bull. No. 11, Ill. St. Mus. Nat. Hist.,
p. 8, Chouteau Gr.

Fig. 1437.—*Edmondia albersi*, right valve.

retusta, Whiteaves, 1897, Pal. Foss., vol.
3, p. 187, Low. Sil. Probably the cast
of a *Lyrodonta*.

ELYMELLA missouriensis, Miller and Gur-
ley, 1896, Bull. No. 11, Ill. St. Mus.
Nat. Hist., p. 15, Chouteau Gr.

Fig. 1438.—*Elymella missouriensis*, left valve
and cardinal view.

Eudodesma, Ulrich, Syn. for *Orthodesma*.
compressum, Ulrich, refer to *Orthodesma*
compressum.

cuneatum, Ulrich, refer to *Orthodesma*
cuneatum.

postulatum, Ulrich, refer to *Orthodesma*
postulatum.

undatum, Ulrich, refer to *Orthodesma*
undatum.

Eridonychia, Ulrich, Syn. for *Ambonychia*;
and *E. apicalis*, *crenata*, and *paucicos-*
tata, Synonyms for *Ambonychia cos-*
tata.

Eurymya, Ulrich, Syn. for *Modiolopsis*.
Goniophora dubia, Hall, instead of *Modiolopsis dubia*.

Fig. 1439.—*Grammysia blairi*, view of the two valves.

Ischyrodonta, Ulrich, Syn. for *Cypricardites*.
decepiens and *truncata*, Ulrich, Synonyms for *Cypricardites hainesi*.
elongata, Ulrich. Probably a *Cypricardites*.
miseneri, Ulrich, refer to *Cypricardites miseneri*.
modioliformis, Ulrich. Probably a *Modiolopsis*, but the name is preoccupied.
oralis, Ulrich, refer to *Cypricardites ovalis*.

LEPTODOMUS. The type is *L. fragilis*.

Liopteria speciosa, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 13, Chouteau Gr.
subovata, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 13, Chouteau Gr.

Fig. 1440.—*Liopteria speciosa*, left valve.

Lucina livonensis, Clarke. Not defined so as to be recognized.
Lunulicardium grande, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 5, Chouteau Gr.
lincolatum, Clarke, 1894, 13th Rep. St. Geol. N. Y., p. 175, Ham. Gr.
livoniae, Clarke, 1894, 13th Rep. St. Geol., N. Y., p. 175, Ham. Gr.
retrosum, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 6, Chouteau Gr.

LYRODESMA *acuminatum*, Ulrich, 1894, Geo. Sur. Minn., p. 609, Trenton Gr.
cannonense, Ulrich, 1894, Geo. Sur. Minn., p. 610, Trenton Gr.
conradi, Ulrich, Syn. for *Lyrodesma cincinnatiense*.
grande, Ulrich, too obscure for determination.
inornatum, Ulrich, Syn. for *Lyrodesma planum*.

Fig. 1441.—*Lunulicardium retrosum*, right valve.

major, instead of *Cleidophorus major*.
subplanum, Ulrich, Syn. for *Lyrodesma cincinnatiense*.

Fig. 1442.—*Macrodon blairi*, right and left valves.

MACRODON *blairi*, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 11, Chouteau Gr.

Fig. 1443.—*Macrodon facetus*, cardinal view and right valve.

facetus, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 10, Chouteau Gr.
pettensis, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 11, Chouteau Gr.

MEGAMBONIA *aviculoidea*, Hall, refer to *Pterinea aviculoidea*.

Modiolodon, Ulrich, Syn. for *Modiolopsis*.
declivus, Ulrich, Syn. for *Modiolopsis subrecta*.

gibbus, Ulrich, refer to *Modiolopsis gibba*.
obtusius, Ulrich, Syn. for *Modiolopsis modiolaris*.
oviformis, Ulrich, refer to *Modiolopsis oviformis*.

oviformis var. *ampla*, Ulrich, Syn. for *Modiolopsis oviformis*.

patulus, Ulrich, refer to *Modiolopsis patula*.

subovalis, Ulrich, Syn. for *Modiolopsis unionoides*.

subrectus, Ulrich, refer to *Modiolopsis subrecta*.

MONOTORSIS *affinis*, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 72, St. Peter Sandstone.

angustifrons, Whiteaves, 1897, Pal. Foss., vol. 3, p. 183, Low. Sil.

arguta, Ulrich, 1894, Geo. Sur. Minn., p. 506, Trenton Gr.

chattfieldensis, Ulrich, 1894, Geo. Sur. Minn., p. 508, Trenton Gr.

consimilis, Ulrich, 1894, Geo. Sur. Minn., p. 505, Syn. for *M. similis*.

contigua, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 71, St. Peter Sandstone.

dubia, Hall, refer to *Goniophora dubia*, Trenton Gr.

excellens, Ulrich, 1894, Geo. Sur. Minn., p. 511, Hud. Riv. Gr.

fountainensis, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 71, St. Peter Sandstone.

gibba, Ulrich, 1894, (*Modiolodon gibbus*), Geo. Sur. Minn., p. 522, Trenton Gr.

gregalis, Sardeson, 1896, Bull. Minn.

Modiophorus major.
n. for *Lyrodesma*

right and left valves.

and Gurley, 1896,
Mus. Nat. Hist.,

s, cardinal view and
ve.

Gurley, 1896, Bull.
Nat. Hist., p. 10,

Gurley, 1896, Bull.
Nat. Hist., p. 11,

a, Hall, refer to

n. for *Modioloopsis*.

n. for *Modioloopsis*

Modioloopsis gibba.

. for *Modioloopsis*

er to *Modioloopsis*

Ulrich, Syn. for

ais. .

er to *Modioloopsis*

n. for *Modioloopsis*

er to *Modioloopsis*

rdeson, 1896, Bull.

ci., vol. 4, p. 72, St.

es, 1897, Pal. Foss.,

Sil.

Geo. Sur. Minn., p.

4, 1894, Geo. Sur.

on Gr.

4, Geo. Sur. Minn.,

similis.

1896, Bull. Minn.

4, p. 71, St. Peter

Goniophora dubia,

4, Geo. Sur. Minn.,

on, 1896, Bull. Minn.

4, p. 71, St. Peter

Modiolodon gibbus.)

522, Trenton Gr.

1896, Bull. Minn.

Acad. Nat. Sci., vol. 4, p. 71, St. Peter Sandstone.

litoralis, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 71, St. Peter Sandstone.

nana, Ulrich, 1894, Geo. Sur. Minn., p. 507, Galena Shales.

obsoleta, Ulrich, 1894, Geo. Sur. Minn., p. 506, Trenton Gr.

oweni, Ulrich. Not defined so as to be recognized.

putula, Ulrich, 1894, (*Modiolodon patulus*.) Geo. Sur. Minn., p. 521, Galena Gr.

postica, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 71, St. Peter Sandstone.

senecta, Sardeson, 1896, Bull. Minn. Acad. Nat. Sci., vol. 4, p. 72, St. Peter Sandstone.

subrecta, Ulrich, 1895, (*Modiolodon subrectus*.) Ohio Geol., vol. 7, p. 653, Hud. Riv. Gr.

Fig. 144.—*Mytilarca jessiae*, two left valves.

MYTILARCA jessiae, Miller and Gurley, 1896, Bull. No. 11, Ill. St. Mus. Nat. Hist., p. 14, Chouteau Gr.

mytiliformis, Foerste, 1895, Ohio Geol., vol. 7, p. 559, Niagara Gr.

perearinata is illustrated in Ohio Geol., vol. 7, p. 422.

Nauidites longus. See *Anthracopectera longa*.

mytiloides. See *Anthracopectera mytiloides*.

NUCULITES ferrugineum, Foerste. Not defined so as to be recognized.

subcuneatus, Clarke, 1894, 13th Rep. St. Geol. N. Y., p. 173, Ham. Gr.

triangulus, Hall and Whitfield, instead of *triangularis*.

ORISTHOPTERA, Meek, is defined by Ulrich, 1895, Ohio Geol., vol. 7, p. 642, with *Ambonychia casei* as the type.

alternata, Ulrich, Syn. for *O. fissicosta*.

ampla, Ulrich, Syn. for *Ambonychia costata*.

casei, instead of *Ambonychia casei*.

extenuata, Ulrich, 1895, Ohio Geol., vol. 7, p. 645, Hud. Riv. Gr.

fissicosta, Meek, as defined by Ulrich, 1895, Ohio Geol., vol. 7, p. 643, Hud. Riv. Gr.

laticostata, Ulrich, 1895, Ohio Geol., vol. 7, p. 646, Hud. Riv. Gr.

notabilis, Ulrich, 1895, Ohio Geol., vol. 7, p. 648, Hud. Riv. Gr. Doubtful species.

obliqua, Ulrich, 1895, Ohio Geol., vol. 7, p. 646, Hud. Riv. Gr.

ORTHODESMA affine, Whiteaves, 1897, Pal. Foss., vol. 3, p. 184, Low Sil.

Fig. 145.—*Orthodesma cymbula*, right and left valves and cardinal view.

ashmani, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, p. 146, Hud. Riv. Gr.

canaliculatum, Ulrich, 1894, Geo. Sur. Minn., p. 520, Hud. Riv. Gr.

cylindricum, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, p. 22, Hud. Riv. Gr.

cymbula, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, p. 22, Hud. Riv. Gr.

ellipticum, Ulrich, 1895, Ohio Geol., vol. 7, p. 667, Hud. Riv. Gr.

grande, Ulrich, 1895, (*Psiloco-*

concha grandis.) Ohio

Geol., vol. 7, p. 665, Hud. Riv. Gr.

minimum, Ulrich, 1895, (*Psiloco-*

concha minima.) Ohio Geol., vol. 7, p. 669, Hud. Riv. Gr.

minnesotense, Ulrich, 1894, (*Psiloco-*

concha minnesotensis.)

Geo. Sur. Minn., p. 531, Galena Gr.

parvum, Ulrich, 1895, Ohio Geol., vol. 7, p. 660, Hud. Riv. Gr.

productifrons, Ulrich, 1895, (*Cymatonta*

productifrons.) Ohio Geol., vol. 7, p. 665, Hud. Riv. Gr.

scaphula, Miller and

Faber, 1894, Jour.

Cin. Soc. Nat. Hist., vol. 17, p. 145, Hud. Riv. Gr.

Fig. 147.—*Orthodesma scaphula*, right valve and cardinal view.

Cin. Soc. Nat. Hist., vol. 17, p. 145, Hud. Riv. Gr.

schucherti, Ulrich, 1894, *Geo. Sur. Minn.*, p. 518, *Galena Gr.*
semistriatum, Ulrich, 1895, (*Cymatona semistriata*.) *Ohio Geol.*, vol. 7, p. 663, *Hud. Riv. Gr.*
sinuatum, Ulrich, 1894, (*Rhytmya sinuata*.) *Geo. Sur. Minn.*, p. 619, *Galena Gr.*
subangulatum, Ulrich, *Syn. for O. recurvum*.
tenuistriatum, Ulrich, 1895, (*Psiloconcha tenuistriata*.) *Ohio Geol.*, vol. 7, p. 668, *Hud. Riv. Gr.*

Ortonella, Ulrich, *Syn. for Cypricardites*.
PALEONEILO similis is illustrated in *Ohio Geol.*, vol. 7, p. 453.

Palaopteria, Whiteaves, *Syn. for Pterinea*.
PALEOSOLEN occidentalis, Miller and Gurley, 1896, *Bull. No. 11, Ill. St. Mus. Nat. Hist.*, p. 16, *Chouteau Gr.*

Fig. 1449.—*Palaosolen occidentalis*, cardinal view and left valve.

PHYSETOMYA, Ulrich, 1895, *Ohio Geol.*, vol. 7, p. 693. [Ety. *physetos*, inflated; *Mya* a genus.] Shell thin, elongate, inflated anteriorly, tapering posteriorly; base arcuate. Beaks in front of the middle, incurved; umbones rounded. Escutcheon and lunule. Surface lined concentrically. Type *P. acuminata*.

acuminata, Ulrich, 1895, *Ohio Geol.*, vol. 7, p. 693, *Hud. Riv. Gr.*

Pinna maxvillensis is illustrated in *Ohio Geol.*, vol. 7, p. 474.

POSTIDONOMYA lasallensis, Miller and Gurley, 1896, *Bull. No. 11, Ill. St. Mus. Nat. Hist.*, p. 12, *Coal Meas.*

PROLOBELLA, Ulrich, Fig. 1449.—*Postidonomya lasallensis*, left valve; same magnified.
Syn. for Pterinea.

striatula, refer to *Pterinea striatula*.
Psiloconcha, Ulrich, *Syn. for Orthodesma elliptica*, refer to *Orthodesma ellipticum*.
grandis, Ulrich, refer to *Orthodesma grande*.
inornata, *sinuata*, and *subrecta*, Ulrich, *Synonyms for Orthodesma subovale*.
minima, Ulrich, refer to *Orthodesma minimum*.
minnesotensis, Ulrich, refer to *Orthodesma minnesotense*.
tenuistriata, Ulrich, refer to *Orthodesma tenuistriatum*.

PSILONYCHIA, Ulrich, *Syn. for Ambonychia*.
perangulata, Ulrich, refer to *Ambonychia perangulata*.

PTERINEA aviculoidea, Hall, instead of *Megambonia aviculoides*.

cincinnatiensis, Miller and Faber, 1874, *Jour. Cin. Soc. Nat. Hist.*, vol. 17, p. 25, *Hud. Riv. Gr.*

parvula, Whiteaves, 1897, (*Palaopteria parvula*.) *Pal. Foss. Can.*, vol. 3, p. 181, *Low. Sil.*

rugatula, Miller and Faber, 1874, *Jour. Cin. Soc. Nat. Hist.*, vol. 17, p. 26, *Hud. Riv. Gr.*

similis, Whitfield, is illustrated in *Ohio Geol.*, vol. 7, p. 445.

striatula, Ulrich, 1894, (*Prolobella striatula*.) *Geo. Sur. Minn.*, p. 532, *Galena Gr.*

Rhytmya, Ulrich, *Syn. for Orthodesma compressa*, *convexa*, and *radiata*, Ulrich, *Synonyms for Orthodesma ashmani*.

ahana, Ulrich, *Syn. for Orthodesma mickelboroughi*.

producta, Ulrich, *Syn. for Orthodesma scaphula*.

recta, Whiteaves, near *Orthodesma rectum*, but probably a distinct species of *Orthodesma*.

sinuata, Ulrich, refer to *Orthodesma sinuatum*.

Saffordia, Ulrich, *Syn. for Cypricardites sulcodorsata*, Ulrich, refer to *Cypricardites sulcodorsata*.

Fig. 1451.—*Schizodus harti*, interior of left valve.

Fig. 1452.—*Schizodus harti*, left valve
ventralis, Ulrich, refer to *Cypricardites ventralis*.

SANGUINOLITES. The type is *S. angulatus*.

l, instead of Me-
and Faber, 1874,
ist., vol. 17, p. 25,

07, (*Palaopteria*
n., vol. 3, p. 181,

ber, 1874, Jour.
vol. 17, p. 26,

1450. — *Pterino-*
ecten sedallensis,
right valves, one
ing broken off.

Orthodesma.
radiata, Ulrich,
ssima ashmani.
for *Orthodesma*

for *Orthodesma*

Orthodesma rec-
tistinct species of

to *Orthodesma*

Cypricardites.
fer to *Cypricar-*

terior of left valve.

ril, left valve
to *Cypricardites*
e is *S. angulatus*.

Schizodus sedallensis, Miller and Gurley,

Fig. 1453.—*Schizodus harli*, cardinal view.

1896, Bull. No. 11,
Ill. St. Mus. Nat.
Hist., p. 17, Chou-
teau Gr.

SPHENOLITUM parallelum,
Ulrich, 1894, Geo.
Sur. Minn., p. 624,
Trenton Gr.

striatum, Ulrich, 1894, Fig. 1451.—*Schizodus*
sedallensis, right
valve.

Fig. 1455.—*Sphenolium cuneiforme*, part of shell broken off and
hinge and ligamental furrows.

SPHENOTUS sinuatus, Miller and Gurley,
1896, Bull. No. 11, Ill. St. Mus. Nat.
Hist., p. 9, Chouteau Gr.

Fig. 1456.—*Sphenotus sinuatus*, right and left
valves.

TECHNOPHORUS cincinnatiensis, Miller and
Faber, 1894, Jour. Cin. Soc. Nat. Hist.,
vol. 17, p. 147, Hud. Riv. Gr.
punctostriatus, Ulrich, Syn. for *T. cin-*
cinnatiensis.

TELLINOMYA absimilis, Sardeson, 1896,
Bull. Minn. Acad. Nat. Sci., vol. 4, p.
74, St. Peter Sandstone.

calvini, Ulrich, 1894, (*Ctenodonta cal-*
vini.) Geo. Sur. Minn., p. 596, Hud.
Riv. Gr.

Fig. 1457.—*Technophorus cincinnatiensis*, nat-
ural size and magnified four diameters.

carinata, Ulrich, 1894, (*Ctenodonta cari-*
nata.) Geo. Sur. Minn., p. 589, Galena
Gr.

clintonensis, Foerste, 1895, Ohio Geol.,
vol. 7, p. 563, Clinton Gr.

compressa, *intermedia*, *recurra*, and *similis*,
Ulrich, Synonyms for *T. alta*.

cuneiformis, Ulrich, 1894, (*Ctenodonta*
cuneiformis.) Geo. Sur. Minn., p. 587,
Trenton Gr.

minima, Foerste, 1895, Ohio
Geol., vol. 7, p. 563, Niaga-
ra Gr.

novicia, Sardeson, 1896,
Bull. Minn. Acad. Nat.
Sci., vol. 4, p. 74, St. Pe-
ter Sandstone.

scofieldi, Ulrich, 1894,
(*Ctenodonta scofieldi*.)
Geo. Sur. Minn., p. 593,
Trenton Gr.

socialis, Ulrich, 1894,
(*Ctenodonta socialis*.)
Geo. Sur. Minn., p. 594,
Trenton Gr.

socialis, Foerste, 1895. The name was
preoccupied.

VANUXEMIA abrupta, Ulrich, 1894, Geo.
Sur. Minn., p. 560, Galena Gr.

crassa, *media*, and *suberecta*, Ulrich, Syn-
onyms for *Cypricardites rotundatus*.
decipiens, Ulrich, Syn. for *Modiolopsis*
plana.

subrotunda, Ulrich, 1894, Geo. Sur.
Minn., p. 559, Trenton Gr.

umbonata, Ulrich, 1894, Geo. Sur. Minn.,
p. 558, Trenton Gr.

wortheni, Ulrich, Syn. for *Cypricardites*
rectirostris.

WHITELLA rugatina, Ulrich, 1894, Geo.
Sur. Minn., p. 569, Trenton Gr.

subcarinata, Ulrich, 1894, Geo. Sur.
Minn., p. 572, Galena Gr.

Whiteavesia, Ulrich, Syn. for *Modiolopsis*.
kentonensis, Ulrich, Syn. for *Modiolopsis*
cincinnatiensis.

SUBKINGDOM ARTICULATA.

CLASS ANNELIDA.

I AM most fully convinced that the Conodonts are not the teeth of Annelids, but belong to the masticatory apparatus of Crustaceans. Several different forms belonged to a single animal. It is idle work to give specific names to these fragments, and a load upon the science. I reprint the names, but with no idea that they represent species.

<p>ARABELLITES procursus, Foerste, 1888, Am. Geol., vol. 2, p. 417, Hud. Riv. Gr.
 EUNICITES confinis, Foerste, 1888, Am. Geol., vol. 2, p. 418, Hud. Riv. Gr.
 fulcatus, Foerste, 1888, Am. Geol., vol. 2, p. 418, Hud. Riv. Gr.
 pululus, Foerste, 1888, Am. Geol., vol. 2, p. 418, Hud. Riv. Gr.
 LUMBRICONEREITES austini, Foerste, 1888,</p> | <p>Am. Geol., vol. 2, p. 417, Hud. Riv. Gr.
 OENONITES deripiens, Foerste, 1888, Am. Geol., vol. 2, p. 417, Hud. Riv. Gr.
 SPIRORBIS anthracosia is described and illustrated in Ohio Geol., vol. 7, p. 492. blairi, Miller and Gurley, 1895, Bull. No. 7, Ill. St. Mus. Nat. Hist., p. 89, Chouteau Gr.</p> |
|---|--|

CLASS CRUSTACEA.

<p>ACIDASPIS brevispinosa, Foerste. Too poorly defined to be recognized.
 AGNOSTUS fallax var. trilobatus, Matthew, 1895, Trans. N. Y. Acad. Sci., vol. 15, p. 216, St. John Gr.
 fissus var. trifissus, Matthew, 1895, Trans. N. Y. Acad. Sci., vol. 15, p. 231, St. John Gr.
 levigatus var. ciceroides, Matthew, 1895, Trans. N. Y. Acad. Sci., vol. 15, p. 234, St. John Gr.
 levigatus var. mamilla, Matthew, 1895, Trans. N. Y. Acad. Sci., vol. 15, p. 234, St. John Gr.
 levigatus var. terranovicus, Matthew, 1895, Trans. N. Y. Acad. Sci., vol. 15, p. 233, St. John Gr.
 nathorsti var. confluens, Matthew, 1895, Trans. N. Y. Acad. Sci., vol. 15, p. 233, St. John Gr.
 ALUTA, Matthew, 1895, Trans. N. Y. Acad. Sci., vol. 15, p. 198. [Ety. aluta, leather.] Small oval or ovate bivalves, like <i>Aparchites</i>, but having a soft, flexible test and finely punctate. Type <i>A. flexilis</i>, described at the same place from the St. John Gr.
 APARCHITES arrectus, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 646, Trenton Gr.</p> | <p>chatfieldensis, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 646, Trenton Gr.
 ellipticus, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 644, Trenton Gr.
 fimbriatus, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 645, Hud. Riv. Gr.
 granilabiat, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 644, Trenton Gr.
 millepunctatus, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 645, Trenton Gr.
 parvulus, Jones, 1897, Pal. Foss. Can., vol. 3, p. 230, Low. Sil.
 secunda, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 136, Up. Taconic.
 ARGES consanguineus. See <i>Lichas consanguineus</i>.
 wesenbergensis var. paulianus. See <i>Lichas paulianus</i>.
 ARISTOZOE canadensis, Whitfield, 1895, Ohio Geol., vol. 7, p. 462, Trenton Gr.
 ASAPHUS. If <i>Isotelus</i> is to rank as a genus, it will include nearly all American species referred to <i>Asaphus</i>.
 canalis is described and figured in Bull. Am. Mus. Nat. Hist., vol. 1, p. 336.
 ulrichi, Clarke, 1894, (Ptychopyge ulrichi), Geo. Sur. Minn., vol. 3, p. 709, Trenton Gr. Founded on fragments of the pygidium.</p> |
|--|--|

- vigilans*, Meek and Worthen, refer to *Nileus vigilans*.
AVALONTIA acuticler, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 140, Up. Taconic.
BATHYURUS schucherti, Clarke, 1894, Geo. Sur. Minn., vol. 3, p. 724, Trenton Gr. *stonemani*. See *Proetus stonemani*.
Bergonia, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 145. Proposed as a subgenus under *Protolenus*.
BEYRICHOXA ovata, planata, rotundata, and triangula, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 134, Up. Taconic.
BEYRICHIA bicornis, Ulrich, 1894, (*Dicranella bicornis*.) Geo. Sur. Minn., vol. 3, p. 665, Trenton Gr.

Fig. 1458.—*Beyrichia ham-melli*, magnified 12 diameters.

- hammelli*, Miller and Faber, 1894, Jour. Cin. Soc. Nat. Hist., vol. 17, p. 157, Hud. Riv. Gr. initials, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 658, Trenton Gr.
marginata, Ulrich, 1894, (*Dicranella marginata*.) Geo. Sur. Minn., vol. 3, p. 666, Trenton Gr.
simplex, Ulrich, 1894, (*Dicranella simplex*.) Geo. Sur. Minn., vol. 3, p. 666, Trenton Gr.
spinosa, Ulrich, 1894, (*Dicranella spinosa*.) Geo. Sur. Minn., vol. 3, p. 665, Trenton Gr.
BOLLIA subaequata, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 669, Galena Gr. *typa*, Ulrich, 1894, (*Dilobella typa*.) Geo. Sur. Minn., vol. 3, p. 673, Trenton Gr. *unguloides*, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 669, Galena Gr.
BRONTES manitobensis, Whiteaves, 1892, Cont. to Can. Pal., p. 347, Devonian.
BUMASTUS trentonensis, Emmons, 1842, Geo. Rep. N. Y., p. 390, and Geo. Sur. Minn., vol. 3, p. 718, Trenton Gr.
BYTHOCEPRIS curta, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 680, Trenton Gr.
granti, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 680, Trenton Gr.
robusta, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 690, Trenton Gr.
CALYMENE callicephal may be confined to the Hud. Riv. Gr. and *senaria* Conrad, may be applied to the related Trenton form, according to some authors.
neapolitana, Clarke, 1892, Am. Jour. Sci., 3d ser., vol. 43, p. 57, Upper Devonian.
vogdesi, Foerste, Syn. for *C. niagarensis*.
CARINOSOMA ingens, Clappole, 1894, Am. Geol., vol. 13, p. 77, Waterlime Gr.
CERATOCARIS monroei, Whitfield, 1896, Bull. Am. Mus. Nat. Hist., vol. 8, p. 301, Low. Held. Gr.
poduriformis, Whitfield, 1896, Bull. Am. Mus. Nat. Hist., vol. 8, p. 302, Low. Held. Gr.

- Ceratopsis*, Ulrich, Syn. for *Peyrichia*.
chamberi var. *robusta*, Ulrich, Syn. for *Beyrichia chamberi*.
CERAURUS elintoni, Foerste, 1895, Ohio Geol., vol. 7, p. 527, Niagara Gr.
millerranus, Miller and Gurley, 1893, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 80, Hud. Riv. Gr.
scofieldi, Clarke, 1894, (*Cyrtometopus scofieldi*.) Geo. Sur. Minn., vol. 3, p. 735, Trenton Gr.
trentonensis, Clarke, 1894, (*Pseudo-spherexochus trentonensis*.) Geo. Sur. Minn., vol. 3, p. 734, Trenton Gr.
CYRIPODITES, Matthew, 1895, Trans. N. Y. Acad. Sci., vol. 15, p. 205. [From a supposed resemblance to *Cirripedes*.] Small calcareous plates of irregular contour and relief associated with *Eocypites*, and supposed to be covering plates of *Cirripedes*. Type *C. cambrensis*, described at the same place from the St. John Gr.
Conolichas, Dames, a subgenus. See *Lichas*.
CTEXONOLINA is probably a synonym for *Beyrichia*.
fulerata, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 674, Trenton Gr.
CYBELE winchelli, Clarke, 1894, Geo. Sur. Minn., vol. 3, p. 742, Galena Gr.
CYPHASPIS bellula, Whiteaves, 1892, Cont. to Can. Pal., p. 340, Devonian.
clintonense, Foerste. Too poorly defined to be recognized.
(?) galenensis, Clarke, 1894, Geo. Sur. Minn., vol. 3, p. 759, Galena Gr.
Cyrtometopus, Angelin, 1854. Subgenus. See *Ceraurus*.
scofieldi. See *Ceraurus scofieldi*.
CYTHERELLA, Jones, 1848, Monog. Entom. Cret. Form., p. 28. [Ety. diminutive of the genus *Cythere*.] Type *C. ovata*.
subrotunda, Ulrich. Not defined so as to be recognized.
DALMANITES dolphi, Clarke. Not defined so as to be recognized.
eboraceus, Clarke, 1894, (*Pterygometus eboraceus*.) Geo. Sur. Minn., vol. 3, p. 728, Trenton Gr.
schmidti, Clarke, 1894, (*Pterygometus schmidti*.) Geo. Sur. Minn., vol. 3, p. 729, Trenton and Galena Gr.
Drepanella, Ulrich, should be spelled *Drepanella*.
Dicranella, Ulrich, Syn. for *Beyrichia*.
bicornis, Ulrich. See *Beyrichia bicornis*.
marginata, Ulrich. See *Beyrichia marginata*.
simplex, Ulrich. See *Beyrichia simplex*.
spinosa, Ulrich. See *Beyrichia spinosa*.
Dilobella, Ulrich, Syn. for *Bollia*.
typa, Ulrich. See *Bollia typa*.
DREPANELLA bigeneris, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 672, Trenton Gr.
bilateralis, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 671, Trenton Gr.
ECHINOCARIS multinodosa, pustulosa, and sublevis are described and figured in Ohio Geol., vol. 7, pp. 458 and 459.

- ELPE* ulrichi, Foerste, 1895, Ohio Geol., vol. 7, p. 532, Niagara Gr. Probably a *Cytheropsis*.
- ELYMOCARIS* hindei, Jones and Woodward, 1894, Lond. Geo. Mag., 4th ser., vol. 1, p. 292, Ham. Gr.
- ENCERIMURUS* cristatus, Clarke, 1894, Geo. Sur. Minn., vol. 3, p. 741, Hud. Riv. Gr.
- vanulus*, Clarke, 1894, Geo. Sur. Minn., vol. 3, p. 739, Trenton Gr.
- ENTOMOCARIS*, Whitfield, 1896, Bull. Am. Mus. Nat. Hist., vol. 8, p. 299. [Ety. *entomus*, cut up; *karis*, a shrimp.] Carapace ovate in outline, bivalvular, with a strong hiatus in front and rounded behind; hinge-line straight for about half its length. Rostrum not known. Abdomen composed of fourteen or more segments, three or four of which may be naked. The post-abdomen bears three spines, the central one or telson, elongate and slender, and the lateral ones (*cercopods*) flattened and articulated to the caudal plate. Type *E. telleri*.
- telleri*, Whitfield, 1896, Bull. Am. Mus. Nat. Hist., vol. 3, p. 300, Low. Held. Gr.
- EURVCHILINA* subequata, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 663, Trenton Gr.
- symmetrica*, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 663, Trenton Gr.
- ventrosa*, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 662, Galena Gr.
- EURYPTERUS* erlensis is figured and described in Ohio Geol., vol. 7, p. 416.
- kokomoensis*, Miller and Gurley, 1896, Bull. No. 10, Ill. St. Mus. Nat. Hist., p. 90, Waterlime Gr.
- Gerasaphes* ulrichiana, Clarke, 1894, Geo. Sur. Minn., vol. 3, p. 710. This subgenus and species are founded upon some fragments of *Proetus spurlocki*, described by Meek, in 1872, in Am. Jour. of Sci., and redescribed and illustrated in Ohio Pal., vol. 1, p. 161. The specimens are not very rare, and have been considered by the best authorities as the young of *Asaphus megistos*, and the species is called the young of *Asaphus* in N. Am. Geol. and Pal., p. 562. If it is a distinct species, it will wear the name of *spurlocki* and not *ulrichiana*.
- GRIFFITHIDES* ornata, Vogdes, 1895, Proc. Cal. Acad. Sci., vol. 4, p. 589, Low. Coal Mens.
- HALLIELLA* lubiosa, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 656, Galena Gr.
- HARPES* (subgenus *Harpina*) minnesotensis, Clarke, 1894, Geo. Sur. Minn., vol. 3, p. 755, Galena Gr.
- (subgenus *Harpina*) rutellum, Clarke, 1894, Geo. Sur. Minn., vol. 3, p. 757, Galena Gr.
- ILLAENUS* danielsi, Miller and Gurley, 1893, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 76, Niagara Gr.
- ovatus*, Foerste, Syn. for *I. ovatus*.
- minnesotensis*, Foerste, Syn. for *Nileus vigilans*.
- orbicundatus* of Billings may be a *Bumastus*.
- ISOCHELINA* scoffieldi, Ulrich, 1894, (Macrotonella scoffieldi,) Geo. Sur. Minn., vol. 3, p. 684, Trenton Gr.
- ISOTELUS* should probably rank as a genus instead of a subgenus of *Asaphus*, and made to include most of the American *Asaphus*.
- vigilans*. See *Nileus vigilans*.
- JOSSELLA* obscura, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 668, Galena Gr.
- KRAUSELLA*, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 691. [Ety. proper name; *ellus*, diminutive.] Carapace small, somewhat elongate, subelliptical, obscurely triangular or semi-ovate in outline, the dorsal margin more convex than the ventral, the latter straight or but gently convex; with moderately thick and very unequal valves; right valve the smaller, drawn out posteriorly into a strong, spine-like process; left valve overlapping the right all around. Type *K. inequalis*.
- arcuata*, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 692, Trenton Gr.
- inequalis*, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 691, Trenton Gr.
- LEPERDITIA* angulifera, is described and illustrated in Ohio Geol., vol. 7, p. 418.
- anticostana* is a Chazy and Trenton species, not Hud. Riv.
- canalis*, Ulrich, (*Leperditella canalis*,) Geo. Sur. Minn., vol. 3, p. 637, Trenton Gr.
- maera*, Ulrich, (*Leperditella maera*,) Geo. Sur. Minn., vol. 3, p. 638, Trenton Gr.
- minor*, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 138, Up. Taconic.
- persimilis*, Ulrich, (*Leperditella persimilis*,) Geo. Sur. Minn., vol. 3, p. 637, Trenton Gr.
- primeva*, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 138, Up. Taconic.
- Leperditella*, Ulrich, Syn. for *Leperditia*.
- canalis*, *maera*, and *persimilis*, Ulrich. See *Leperditia*.
- LEPIDITTA* auriculata and sigillata, Matthew, 1894, Trans. Roy. Soc. Can., vol. 11, p. 99, Up. Taconic.
- LICHAS* byrnesianus, Miller and Gurley, 1893, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 78, Niagara Gr.
- consanguineus*, Clarke, 1894, (Arges consanguineus,) Geo. Sur. Minn., vol. 3, p. 746, Low. Held. Gr.
- cornutus*, Clarke, (*Conolichas cornutus*,) 1894, Geo. Sur. Minn., vol. 3, p. 749, Trenton Gr.
- hanoverensis*, Miller and Gurley, 1893, Bull. No. 3, Ill. St. Mus. Nat. Hist., p. 78, Niagara Gr.
- paulianus*, Clarke, 1894, (Arges wesenbergensis var. paulianus,) Geo. Sur. Minn., vol. 3, p. 744, Trenton Gr.

Syn. for *Nileus*

may be a Bu-

ch, 1894, (Macro-
eo. Sur. Minn.,
Gr.

rank as a genus
of *Asaphus*, and
of the American

gilians.

h, 1894, Geo. Sur.
Galena Gr.

Geo. Sur. Minn.,
y, proper name;

Carapace small,
subelliptical, ob-

semi-ovate in out-
rigin more convex

latter straight or
with moderately

und valves; right
drawn out poste-

spine-like process;
ing the right all

aequalis,
Geo. Sur. Minn.,

on Gr.

4, Geo. Sur. Minn.,
on Gr.

is described and
Geol., vol. 7, p. 418.

ry and Trenton spe-

perditella canalis,
vol. 3, p. 637, Tren-

perditella macra,
vol. 3, p. 638, Tren-

1894, Trans. N. Y.
p. 138, Up. Taconic.

Lepeditella persi-
Minn., vol. 3, p. 637,

1894, Trans. N. Y.
p. 138, Up. Taconic.

n. for Lepeditella.
persimilis, Ulrich.

and sigillata, Mat-
Roy. Soc. Can., vol.

nic.

Miller and Gurley,
Ill. St. Mus. Nat.

on Gr.

ke, 1894, (Arges con-
Sor. Minn., vol. 3, p.

r.

onolichas cornutus,
Minn., vol. 3, p. 749,

r and Gurley, 1893,
t. Mus. Nat. Hist., p.

1894, (Arges wesen-
aulianus.) Geo. Sur.

44, Trenton Gr.

Macrotonella, Ulrich, Syn. for *Isochilina*, to
which refer *M. scollfieldi*.

MEMACOA, Matthews, 1894, Trans. N. Y.

Acad. Sci., vol. 14, p. 141. [Ety. proper
name.] Glabella prominent, cylindrical,

extending to the front of the shield;
eye-lobes continuous, and a short pos-

terior extension of the dorsal suture.
The front area is like *Zacanthoides*, but

different in the posterior extension.
The long eye lobes and short posterior

extension of the suture resemble *El-*
lipsoccephalus. Type *M. matthewi*, de-

scribed at the same place with *M.*
matthewi var. *ingens*, *M. recuera*, and

M. plana, from the Up. Taconic.

Microdiscus schucherti, Matthew, 1895,
Trans. N. Y. Acad. Sci., vol. 15, p. 238,

St. John Gr.

MOOREA angularis, Ulrich, 1894, Geo. Sur.
Minn., vol. 3, p. 682, Trenton Gr.

perplexa, Ulrich, 1894, Geo. Sur. Minn.,
vol. 3, p. 683, Trenton Gr.

punctata, Ulrich, 1894, Geo. Sur. Minn.,
vol. 3, p. 682, Trenton Gr.

NILEUS vigilans, instead of *Asaphus* (*Iso-*
telus) vigilans.

PALEOPALEMON newberryi, is described and
illustrated in Ohio Geol., vol. 7, p. 461.

Platynotopus is a subgenus of *Lichas*.

PLUMULITES manuelensis, Matthew, 1895,
Trans. N. Y. Acad. Sci., vol. 15, p.

200, St. John Gr.

POLYCOPE has a bivalve circular or ovate
carapace, without beak or sinus, and

is described on page 122. A living
genus. Type *P. orbicularis*.

PRIMITIA celata, Ulrich, 1894, Geo. Sur.
Minn., vol. 3, p. 653, Trenton Gr.

constricta, Ulrich, 1894, (Primitiella con-

stricta,) Geo. Sur. Minn., vol. 3, p. 647,
Trenton Gr.

duplicata, Ulrich, 1894, Geo. Sur. Minn.,
vol. 3, p. 654, Trenton Gr.

fillmorensis, Ulrich, (Primitiella fill-

morensis,) Geo. Sur. Minn., vol. 3, p.

649, Trenton Gr.

fusiformis, Matthew, 1894, Trans. N. Y.
Acad. Sci., vol. 14, p. 137, Up. Taconic.

gibbera, Ulrich, 1894, Geo. Sur. Minn.,
vol. 3, p. 655, Hud. Riv. Gr.

limbata, Ulrich, (Primitiella limbata,) Geo. Sur. Minn., vol. 3, p. 648, Trenton Gr.

mammata, Ulrich, 1894, Geo. Sur. Minn.,
vol. 3, p. 652, Trenton Gr.

micula, Ulrich 1894, Geo. Sur. Minn.,
vol. 3, p. 653, Galena Gr.

minutissima, Ulrich, 1894, Geo. Sur.
Minn., vol. 3, p. 651, Trenton Gr.

oculata, Matthew, 1894, Trans. N. Y.
Acad. Sci., vol. 14, p. 136, Up. Taconic.

sanctipauli, Ulrich, 1894, Geo. Sur.
Minn., vol. 3, p. 652, Trenton Gr.

simulans, Ulrich, (Primitiella simulans,) Geo. Sur. Minn., vol. 3, p. 648, Trenton Gr.

tumidula, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 655, Hud. Riv. Gr.

uphami, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 651, Galena Gr.

PRIMITIELLA, Ulrich, Syn. for *Primitia*, to which refer *constricta*, *fillmorensis*, *limbata*, *simulans*.

PROETUS determinatus, Foerste. Too poorly defined to be recognized.

mundulus, Whiteaves, 1892, Cont. to Can. Pal., p. 350, Devonian.

placidus, Vogdes, 1896, Cal. Acad. Sci., p. 197, Chouteau Gr.

stonemani, Vogdes, 1881, (Bathyrus stonemani,) 12th Ann. Rep. Minn., p. 8, Devonian.

PROTAGRAULUS, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 138. [Ety. *protos*, first; *Agraulius*, a genus.] Glabella depressed to the level of the cheeks, scarcely traceable except at the posterior end. Cheeks slightly arched down at the front and sides. Eye-lobes long. Anterior ends of the facial sutures approximate. Type *P. priscus*, described at the same place from the Up. Taconic.

PROTOLEUS, Matthew, 1892, Bull. Nat. Hist. New Brunswick, p. 34. [Ety. *protos*, first; *Olenus* a genus.] Type *P. elegans*. St. John Gr.

PSEUDOSPIEREXOCCHUS trentonensis, Clarke. See *Ceraurus trentonensis*.

Pterygomotopus, a subgenus of *Dalmanites*, to which refer *eboracensis* and *schmidti*.

Psychoparia, a synonym for *Atops*, to which all the species should be referred.

Psychopage, a subgenus of *Asaphus*, to which refer *ulrichi*, if it is to be regarded as a species.

SCHMIDTELLA affinis, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 641, Galena Gr.

brevis, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 642, Trenton Gr.

cambrica, Matthew, 1894, Trans. N. Y. Acad. Sci., vol. 14, p. 137, Up. Taconic.

incompta, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 642, Trenton Gr.

subrotunda, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 643, Trenton Gr.

umbonata, Ulrich, 1894, Geo. Sur. Minn., vol. 3, p. 641, Trenton Gr.

SPHEREXOCCHUS pisum, Foerste. Too poorly defined to be recognized.

Thaleops, Conrad, 1843, Proc. Acad. Nat. Sci. Phil., vol. 1, p. 332, is regarded as a subgenus of *Iliaenus*. But it would be better if all subgeneric names were dropped.

TERRILEPAS newberryi is described and illustrated in Ohio Geol., vol. 7, p. 463.

CLASS ARACHNIDA.

- ANTHRACOMARTUS woodruffi, Scudder, 1893, Insect Fauna, R. I. Coal Field, p. 9, Coal Meas. | MAZONTIA mendica, Scudder, 1895, Cont. to Can. Pal., vol. 2, p. 83, Coal Meas.

CLASS MYRIAPODA.

- ARCHILUS euphoerioides, Scudder, 1895, Cont. to Can. Pal., vol. 2, p. 59, Coal Meas. | IYELLI, Scudder, 1895, Cont. to Can. Pal., vol. 2, p. 60, Coal Meas.

CLASS INSECTA.

- ANTHRACOBLETTINA americana, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 129, Coal Meas.
virginiensis, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 130, Permian.
ETOBLETTINA accubita, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 88, Low. Permian.
angusta, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 100, Low. Permian.
aperta, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 80, Low. Permian.
arcta, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 97, Low. Permian.
balteata, Scudder, 1879, Mem. Bost. Soc. Nat. Hist., p. 110, Low. Permian.
benedicti, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 84, Low. Coal Meas.
clarki, Scudder, 1893, Insect Fauna, R. I. Coal Field, p. 14, Coal Meas.
clintoniana, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 86, Coal Meas.
communis, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 93, Low. Permian.
debilis, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 71, Low. Permian.
defossa, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 108, Low. Permian.
detecta, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 75, Low. Permian.
eakiniana, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 88, Low. Permian.
exigua, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 76, Low. Permian.
exilis, Scudder, 1893, Insect Fauna, R. I. Coal Field, p. 17, Coal Meas.
expugnata, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 102, Low. Permian.
expulsata, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 89, Low. Permian.
expuncta, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 79, Low. Permian.
executa, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 96, Low. Permian.
exsensa, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 86, Low. Coal Meas.
fasciata, Scudder, 1889, Proc. Bost. Soc. Nat. Hist., vol. 24, p. 47, Low. Coal Meas.
fossa, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 70, Low. Coal Meas.
funeraria, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 78, Low. Permian.
funesta, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 85, Low. Coal Meas.
gorhami, Scudder, 1893, Insect Fauna, R. I. Coal Field, p. 16, Coal Meas.
gracilentia, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 95, Low. Coal Meas.
gratiosa, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 90, Low. Permian.
hastata, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 94, Low. Coal Meas.
hilliana, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 99, Low. Coal Meas.
hustoni, Scudder, 1889, Proc. Bost. Soc. Nat. Hist., vol. 24, p. 53, Low. Coal Meas.

- packardii, Scudder, 1893, Insect Fauna, R. I. Coal Field, p. 11, Coal Meas.
- ORYCTOBLATTINA laquata**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 133, Up. Coal Meas.
- PARALOGUS**, Scudder, 1893, Insect Fauna, R. I. Coal Field, p. 20. [Ety. *paralogos*, unlooked for.] Fore wings three or more times as long as wide, costal margin nearly straight, hinder margin arcuate. Mediastinal vein simple, close to the outer margin. Scapular vein near to and parallel with the mediastinal vein; emits an offshoot below the middle of the basal half, which immediately divides, the upper branch bearing a few longitudinal offshoots, the other many arcuate offshoots. Externomedian vein sinuate. Internomedian and anal veins originate from a single stem, and each bears numerous arcuate branches. The reticulation is quadrangular. Type *P. aschnoides*, described at the same place from the Coal Meas.
- PAROMYLACRIS clintoniana**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 53, Low. Coal Meas.
- pluteus**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 54, Coal Meas.
- triangularis**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 52, Coal Meas.
- PETRABlattina hastata**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 141, Low. Permian.
- POROBLATTINA complexinervis**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 139, Low. Permian.
- fossa**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 137, Low. Permian.
- gratiosa**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 136, Low. Permian.
- longinqua**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 135, Low. Coal Meas.
- ohioensis**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 138, Low. Coal Meas.
- PROGONOBLATTINA columbiana**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 131, Low. Coal Meas.
- PROMYLACRIS harei**, Scudder, 1895, Rev. Am. Foss. Cockroaches, p. 48, Up. Coal Meas.
- RHAPHIDIOPSIS**, Scudder, 1893, Insect Fauna, R. I. Coal Field, p. 10. [Ety. from the resemblance to *Rhaphidia*.] Distinguished from *Corydaloides* by the nonfalcate form of the relatively shorter wings, the excessive breadth of the hind wings, and the less numerous and much more distant scapular branches. Type *R. diversipenna*, described at the same place from the Coal Meas.

SUBKINGDOM VERTEBRATA.

CLASS PISCES.

- ASPIDICHTHYS notabilis**, Whiteaves, 1892, Cont. to Can. Pal., p. 354, Devonian.
- BRONTICHTHYS**, Clappole, 1894, Am. Geol., vol. 14, p. 379. [Ety. *Brontes*, mythological name; *ichthys* fish.] A fish-spine. Type *B. clarkei*, described at the same place from the Cleveland Shale.
- CLADODUS clarki, rivipetrosi, and sinuatus**, Clappole, 1893, Am. Geol., vol. 11, p. 327, Cleveland Shale.
- magnificus**, Clappole, 1894, Am. Jour., vol. 14, p. 137, Cleveland Shale.
- CLADOSELACHE newberryi**, Dean, 1893, Trans. N. Y. Acad. Sci., vol. 13, p. 115, Waverly Gr.
- COCOOSTEUS cuyahogae**, Clappole, 1895, Ohio Geol., vol. 7, p. 615, Cleveland Shale.
- halmodeus**, Clarke, 1894, 13th Rep. St. Geol., N. Y., p. 161, Marcellus Shale.
- CTENACANTHUS acutus**, Eastman, 1897, Am. Jour. Sci., vol. 154, p. 13, Keokuk Gr.
- DINICHTHYS canadensis, clarkei, gracilis, and lincolni**. See Ponerichthys.
- prentis clarkei**, Clappole. Not binomial.
- MONOCLADODUS**, Clappole, 1893, Am. Geol., vol. 11, p. 329. [Ety. *monos*, single; *Cladodus*, a genus.] Distinguished from *Cladodus* by the teeth, which consist of a single cusp each, without lateral denticles. Type *M. clarki*, which, with *M. pinnatus*, is described at the same place from the Cleveland Shale.
- PETALODUS securiger**, Hay, 1895, Jour. Geol., vol. 3, p. 561, Coal Meas., Syn. for *P. alleghaniensis*.
- TITANICHTHYS attenuatus**, Wright, 1895, Ohio Geol., vol. 7, p. 612, Cleveland Shale.
- brevis**, Clappole, 1896, Am. Geol., vol. 16, p. 167, Cleveland Shale.

Scudder, 1895, Rev.
Sci., p. 141, Low.

Scudder,
Cockroaches, p.

Rev. Am. Foss.
Low. Permian.

Rev. Am. Foss.
Low. Permian.

1895, Rev. Am.
p. 135, Low. Coal

, Rev. Am. Foss.
Low. Coal Meas.

biana, Scudder,
Cockroaches, p.

lder, 1895, Rev.
ss, p. 48, Up. Coal

, 1893, Insect
eld, p. 10. [Ety.

e to *Rhaphidia*.]
rydaloides by the

the relatively
essive breadth of

ne less numerous
istant scapular

liversipenna, de-
place from the

TAMIOBATIS, Eastman, 1897, Am. Jour.
Sci., vol. 154, p. 85. [Ety. *Tamias* a
genus; *batis*, a ray.] Founded upon
what is supposed to be the cranium
of a skate. Type *T. retustus*, described
at the same place from rocks of De-
vonian or Subcarboniferous Age.

PONERICHTHYS canadensis, Whiteaves, 1892.
(*Dinichthys canadensis*.) Cont. to Can.
Pal., p. 353, Devonian.
clarkei, gracilis, and lincolni, Claypole,
1893. (*Ponerichthys clarkei*, etc.) Am.
Geol., vol. 12, p. 275, Cleveland and
Marcellus Shales.

CLASS BATRACHIA.

ALLOPUS littoralis, Marsh, 1894, Am. Jour.
Sci., vol. 148, p. 83, Coal Meas. Tracks
made by an animal having five toes
on its fore feet and four on its hind
feet.

BAROPUS lentus, Marsh, 1894, Am. Jour.
Sci., vol. 148, p. 83, Coal Meas. Tracks
made by an animal having four toes on
both its fore and hind feet.

DROMOPUS agilis, Marsh, 1894, Am. Jour.
Sci., vol. 148, p. 82, Coal Meas. Tracks

made by an animal having long, slender
digits, terminated by sharp claws.

LIMNOPUS vagus, Marsh, 1894, Am. Jour.
Sci., vol. 148, p. 82, Coal Meas. Tracks
made by an animal having four toes on
its fore feet and five on the hind feet.

NANOPUS caudatus, Marsh, 1894, Am. Jour.
Sci., vol. 148, p. 82, Coal Meas. Tracks
made by an animal having three toes
on its fore feet and four on those be-
hind.

TA.

154, p. 13, Keo-

-kei, gracilis, and
athys.

ole. Not bino-

1893, Am. Geol.,

. monos, single;
Distinguished

te teeth, which
sp each, without

type *M. clarki*,
tus, is described

m the Cleveland

ay, 1895, Jour.
il Meas., Syn. for

, Wright, 1895,
612, Cleveland

Am. Geol., vol.
hale.