IMAGE EVALUATION TEST TARGET (MT-3)

Photographic
Sciences
Corporation

23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1983

**Th
to**

**Th
po
of
filr**

Or
be
the
sio
oth
firs
sio
or

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☒ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

**Th
sha
TIM
wh**

**Ma
dif
ent
beg
rig
req
me**

The copy filmed here has been reproduced thanks to the generosity of:

Entomology Research Library
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque de recherche entomologique
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

rata
)

elure,
à

NORTH AMERICAN TACHINIDÆ.

GONIA.

BY PROF. S. W. WILLISTON, NEW HAVEN, CT.

The genus *Gonia* is one easily recognized by reason of its peculiar structure of the head and antennæ; unfortunately the separation of species is a much less simple matter. As is so frequently the case, isolated speci-

with numerous, not very strong bristles; face in profile nearly perpendicular, the epistoma but slightly projecting; the oral margin with a row of bristles, not ascending on the edges of the median facial depression. Cheeks broad; bare, or with short bristles. Antennæ elongate; third joint two or three times as long as the second in the female, from four to eight times in the male; arista stout, composed of three joints, the second and third elongate, and joining each other at an angle, as though broken. Eyes bare. Palpi cylindrical. Scutellum with six strong bristles on its border. Abdomen broad oval or ovate; first and second segments each with a pair of median posterior bristles and one extreme lateral posterior one; third segment with a row on the posterior border. No spinule at tip of auxiliary vein; fourth vein obtusely angulated, without stump of vein, terminating in the costa a considerable distance before the tip; posterior cross-vein much nearer the angle than to the anterior cross-vein.

Head yellow, except the occiput, palpi yellow; thorax black, margins of dorsum and scutellum often yellowish; legs black. Larvæ, so far as known, parasitic on hymenoptera and lepidoptera.

Table of Species.

- 1.—Abdomen wholly black, or at most only somewhat reddish on the sides of base; claws and pulvilli of male small. . FRONTOSA Say.
Abdomen largely or wholly red or yellow. 2
- 2.—Face and antennæ golden yellow; claws of male small. . SENILIS, n. sp.
Face and antennæ not golden yellow. 3
- 3.—Sides of face, pleuræ and femora with yellow hair; claws and pulvilli of male small. PORCA, n. sp.

BY G. H. FRENCH, CARBONDALE, ILL.

Homchadena Elda, n. sp.

Expanse 1.30 inches. Apex very slightly produced, the posterior angle rounded; eyes naked; palpi oblique, projecting beyond the head in their natural position about the width of the eyes; tibiae unarmed; abdomen equalling the hind wings; covering of thorax mostly scales. Color, warm gray, slightly wine tinted; basal half line black, not very distinct, strongly angulated outward on the subcostal vein, a black basal dash crossing the lower end of this line, the space enclosed between the costa

OVT

?Box

ced, the posterior
beyond the head in
e unarmed; abdo-
ly scales. Color,
not very distinct,
black basal dash
between the costa

mens frequently furnish grounds for much more decided opinions than a richer material will sustain. I describe in the present paper five species that present characters sufficiently decisive to render their recognition not doubtful; other species I forbear to name till my material is richer. It will be seen that I make little use of the structure of the arista, a character that has been relied upon much in the European species; but I am satisfied that it is a variable one in individuals of the same species, and is, moreover, distinctly sexual. There is a distinct variation in different species in the position of the anterior cross-vein, as regards the termination of the auxiliary; of how much value the character is I am not yet decided.

Generic characters.—Moderately large species, not very bristly. Head very large, inflated; front very broad, only a little narrower in the male, with numerous, not very strong bristles; face in profile nearly perpendicular, the epistoma but slightly projecting; the oral margin with a row of bristles, not ascending on the edges of the median facial depression. Cheeks broad; bare, or with short bristles. Antennæ elongate; third joint two or three times as long as the second in the female, from four to eight times in the male; arista stout, composed of three joints, the second and third elongate, and joining each other at an angle, as though broken. Eyes bare. Palpi cylindrical. Scutellum with six strong bristles on its border. Abdomen broad oval or ovate; first and second segments each with a pair of median posterior bristles and one extreme lateral posterior one; third segment with a row on the posterior border. No spinule at tip of auxiliary vein; fourth vein obtusely angulated, without stump of vein, terminating in the costa a considerable distance before the tip; posterior cross-vein much nearer the angle than to the anterior cross-vein.

Head yellow, except the occiput, palpi yellow; thorax black, margins of dorsum and scutellum often yellowish; legs black. Larvæ, so far as known, parasitic on hymenoptera and lepidoptera.

Table of Species.

- 1.—Abdomen wholly black, or at most only somewhat reddish on the sides of base; claws and pulvilli of male small. . FRONTOSA Say.
Abdomen largely or wholly red or yellow.....2
- 2.—Face and antennæ golden yellow; claws of male small. . SENILIS, n. sp.
Face and antennæ not golden yellow.....3
- 3.—Sides of face, pleuræ and femora with yellow hair; claws and pulvilli of male small.....PORCA, n sp.

- Sides of face with black bristles ; pleuræ not with yellow hair.
- 4.—Abdomen reddish yellow, with or without small black spots ; claws there
and pulvilli of male small.....SEQUAX, n. sp. not m
- Abdomen largely black ; claws and pulvilli of male large. . EXUL, n. sp. All

GONIA FRONTOSA Say, J. Acad. Phil. vi., 175 ; Compl. Wr. ii., 365. United

Gonia philadelphica Macquart, Dipt. Exot. ii., 3, 51, 6. and no

? *Gonia albifrons* Walker, List, etc., iv., 798. pollino

♀. Length 9.5-10 m.m. Face and front light yellow, with a silvery. third se
or slightly golden sheen ; front a little translucent on the sides when seen in mos
from above ; face on the sides with short black bristles, and a number of third to
longer ones near the sides of the median depression ; median depression third to
in its greatest width about as wide as the least width of the sides of the as long
face ; cheeks with short black bristly hairs only. Antennæ blackish ones in
brown, the second joint and base of third yellowish ; second joint about Englan
one-third the length of the third ; second joint of the arista not more than Carolin
three-fifths the length of the third. Palpi yellow. Dorsum of thorax red, wit
black ; the humeri, post-alar callosities, and scutellum in large part, luteous A speci
yellowish ; dorsum, when seen from behind, distinctly gray pollinose, the ban
leaving four slender stripes. Bristles of scutellum long, reaching to base have th
of third abdominal segment. Abdomen deep shining black ; the third the thir
segment with a very narrow, the third with a narrow, the fourth with a other sp
moderately broad, basal white-pollinose cross band. Wings nearly hyaline other sp
behind ; the costal, subcostal and first basal cells very distinctly tinged respect
with brown. in havi
than th

♂. Length 9-10 m.m. Front distinctly narrower, and more project- not a
ing below. Median facial depression deeper, larger, and longer, broader space
below than the sides of the face. Antennæ larger and longer, the second larger
joint shorter, not more than one-fifth or one-sixth as long as the third ; Th
arista shorter than the joint, its second joint curved, three-fifths, or three- decide
fourths as long as the third. Dorsum of the thorax a little more hairy. upon
Abdomen more elongate or egg-shaped, the sides of the second and third is alw
segments sometimes reddish. Tarsal joints of front feet a little less disci- I t
form, the claws a little longer. " larg
not us
phica
sibly
have
the f

This description is drawn from twelve females, mostly collected to-
gether in Fairmount Park, Philadelphia, by Mr. E. Keen, and eight males
from Minnesota, collected together. They are all closely related, and
their absolute identity seems certain. Among these specimens, however,

ow hair.
 spots; claws
 SEQUAX, n. sp.
 . . . EXUL, n. sp.

W. ii., 365.
 i., 6.

with a silvery;
 es when seen
 a number of
 n depression
 e sides of the
 nœ blackish
 joint about
 ot more than
 m of thorax
 part, luteous
 y pollinose,
 hing to base
 the second
 urch with a
 arly hyaline
 ictly tinged
 ore project-
 er, broader
 the second
 the third;
 s, or three-
 ore hairy.
 and third
 less disci-

ected to-
 ight males
 ated, and
 however,

there is no absolute uniformity in the length of the arisal joints, though not much discrepancy.

All the rest of my specimens, from various localities throughout the United States, are larger, reaching twelve and a half millimetres in length and none of them as small as the largest of the ones described. The pollinose bands of the abdomen are usually broader on the second and third segments, the wings in none so strongly infuscated proximally, and in most the infuscation is more or less obsolete; the antennæ vary from red to black, the second joint of the arista in the female varies from one-third to three-fourths as long as the third joint; in the male from one-half as long to longer. The dorsal thoracic stripes vary from very slender ones in Colorado specimens, to broader and more distinct ones in New England, North Carolina, and Southern Californian ones. Specimens from Carolina and New England have the basal part of the venter yellowish red, with moderately large spots on the sides of the basal segments above. A specimen from Wyoming has the thoracic dorsum strongly pollinose, and the bands of the abdomen unusually broad. Two others from California have the second joint of the arista very short, not a third of the length of the third, the wings scarcely at all infuscated in front, and, unlike all the other specimens, the cheeks do not have black hairs, but yellow pile; other specimens from California, however, are like the eastern ones in this respect. Yet another, rubbed (female) specimen, from Connecticut, differs in having the sides of the face distinctly narrower, considerably narrower than the median portion. The bristly hairs on the sides of the face vary not a little in size; those of the described specimens leave a more bare space below on the inner side, that is less, or not at all, apparent in the larger specimens.

The value of these various characters I confess myself unable to decide. Only this much seems certain, no, or little reliance can be placed upon the comparative lengths of the arisal joints; and the second joint is always longer and more bent in the male than in the female.

I think there can be little doubt but that this is Say's species; his "large, obsolete, reddish brown spot near the base" of the abdomen is not usually, though frequently, present. The synonymy of *G. philadelphia* seems equally certain. The species must be closely allied to, possibly identical with *G. fasciata* Meigen, of Europe, the larvæ of which have been found in nests of *Bombus terrestris* and *Megilla reclusa*, where the flies deposit their eggs. A specimen from the Boston Society of

Natural History, somewhat teneral with red antennæ, bears the label "From *Pyrrharctia isabella*, Apr. 24."

Altogether, I have examined about seventy-five specimens, from New England, New York, Pennsylvania, North Carolina, Indiana, Minnesota, Montana, Colorado, Wyoming, Northern and Southern California.

GONIA SENILIS, n. sp.

♂. Length 10 m.m. Front and face golden yellow, not very changeable, the vertex on either side sub-translucent when seen from above; sides of the face considerably narrower than the median depression, clothed with short black bristles. Antennæ wholly pure orange-yellow, the tip of arista brownish, second joint very short, third very long, six or eight times longer than the second; second joint of the arista shorter than the third. Thorax not densely pollinose, when seen from behind, with four slender stripes, scutellum luteous at tip only. Abdomen black, sides broadly red, forming a moderately broad lateral posterior margin, the first and reaching to about the middle of the third segment on each side; pollinose bands broad. Wings nearly hyaline, anterior cross-veins nearly opposite the middle of the subcostal section of the costa. Claws and pulvilli small.

One specimen, Western Kansas. Although I have but the single specimen, I venture to describe this species. The color of the antennæ seems evidently normal; should it really be darker in other specimens, however, the other characters given will, I believe, render the species recognizable.

GONIA PORCA, n. sp.

♂. Length 13 m.m. Face and front light yellow with a silvery sheen, the front, except in a very oblique light, oily translucent on the sides; sides of the face without the usual short black bristles, except above; the largest part, as also the cheeks, is clothed with long fine hairs or pile; the usual row along the sides of the median depression. Median depression wider than the sides of the face below. Antennæ black, the third joint five or six times longer than the second; second joint of the arista shorter than the third. Dorsum of thorax thinly pollinose, stripes not very distinct; the sides more or less and the scutellum luteous yellow. Pleuræ with long fine yellow hairs or pile. Abdomen yellowish red, shining; pollinose bands narrow, that on the second ve

bears the lab
mens, from N
iana, Minneso
alifornia.
arrow ; first segment, under the scutellum, the second with a median
ripe, the third with a stripe more or less dilated behind, and the fourth
chiefly, black. The anterior femora behind with long fine light yellow
air. Wings nearly hyaline, the costal cells scarcely infuscated.
Two specimens, Mt. Hood, Or. The moderately abundant, long, light
ellow fine hair on face, pleuræ and four anterior femora is characteristic.

GONIA EXUL, n. sp.

not very chan
en from abov
lian depressio
e orange-yell
very long, six
rista shorter th
om behind, w
omen black,
terior margin
segment on
terior cross-v
e costa. Cla
but the sin
of the anten
other specime
nder the spec
♀. Length 12, 13 m. m. Front and face light yellow, with a semi-
ranslucent appearance as though oiled ; when seen obliquely, with a sil-
ery sheen ; sides of the face with numerous, short, black bristles, on the
inner part, near the sides of the median depression, with three or four
moderately stout bristles ; median depression small and narrow, not as
wide as the sides ; cheeks without black bristles. Antennæ blackish
brown, second joint and base of third yellow ; second joint about one-
third the length of the third ; arista variable, third joint not twice the
length of the second. Dorsum of thorax gray pollinose, when seen from
behind leaving four moderately broad stripes, the margins more or less
and the scutellum yellow. Abdomen yellowish red ; first segment under
the scutellum, the second with a broad median stripe, the third except the
anterior angles, or, rather, with a large triangle reaching the whole width
of the segment behind and touching in front more or less broadly, the
dorsum of the fourth wholly black ; pollen of the whole abdomen variable
in different reflections, but most conspicuous on the base of the third and
the whole of the fourth segments. Wings nearly hyaline, the veins yel-
lowish, except posteriorly.

Ten specimens, Connecticut, Massachusetts, New York.

y with a silv
anslucent on t
bristles, exc
with long f
lian depressio
low. Antenn
second ; seco
thorax thinly
nd the scutell
pile. Abdom
the second ve
a. Males from California, that I cannot distinguish, have the antennæ
wholly black, the third joint five or six times the length of the second, the
face strongly silvery pollinose, and the claws and pulvilli very large, con-
siderably longer than the terminal joint of the tarsi. In some, the black
triangle on the third abdominal segment is much smaller. A female
with them seems to agree in all respects with the Eastern specimens.

b. A female specimen from California, with the abdomen marked
similarly, has the third joint of the antennæ shorter, not more than twice
the length of the second, the second joint of the arista very short, and
the bristles on the sides of the median depression not parallel with the
edge, but divergent, whisker-like.

c. Two female specimens from Arizona differ distinctly in having a broader and more projecting face, smaller eyes, whiter and more dense pollen on face, thorax and abdomen. Another specimen with these two, however, though agreeing in the latter respects, has the face of the usual width.

d. A male specimen from North Park is of a distinct species, though closest allied to the present. The sides of the face are unusually narrow, the front short, convex and almost horizontal, the abdomen marked more like that of *G. senilis*, the size smaller (12 m.m.), the claws and pulvilli large.

GONIA SEQUAX, n. sp.

♂. Length 12-13 m.m. Like the males of *exul*, except that the abdomen is either wholly yellowish red, with a black spot under the scutellum, and a brownish hind margin to the third segment, or with a slender black stripe on the second, and a small triangle on the third and fourth segments. The claws and pulvilli are very much smaller, much shorter than the terminal joints of the tarsi.

Three specimens, California.

NOTES ON THE GENUS EXOPROSOPA.

BY D. W. COQUILLET, LOS ANGELES, CAL.

In my paper on "The North American Genera of Anthracina," which appeared in the last volume of the CAN. ENTOMOLOGIST (pp. 157-159), I proposed the name *Velocia* for those species of *Exoprosopa* which have four submarginal cells in each wing. Through the kindness of Dr. Williston, I have been put in possession of a paper by Prof. Camillo Rondani, entitled "Dipterorum Species et Genera Aliqua Exotica," published in 1863 in the Archivio per la Zoologia, vol. iii., and on pages 56 and 57 of this paper Prof. Rondani proposes the name *Hyperalonia* for the above group; this name being the earlier, must supersede the one proposed by me.

In the same paper Prof. Rondani proposes the name *Argyrosipila* for those species of *Exoprosopa* in which the third and fourth veins are joined together before the margin of the wing, instead of reaching the margin