IMAGE EVALUATION TEST TARGET (MT-3)

Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1983

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured covers/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- ☐ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☒ Bound with other material/
Relié avec d'autres documents
- ☒ Tight binding may cause shadows or distortion
along interior margin/
Liaison serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- ☐ Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- ☐ Additional comments:/
Commentaires supplémentaires:

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☒ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
☐	☐	☒	☐	☐	☐
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:

Library
Agriculture Canada

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol $\rightarrow$ (meaning "CONTINUED"), or the symbol ∇ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Bibliothèque
Agriculture Canada

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole $\rightarrow$ signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

ONTARIO AGRICULTURAL COLLEGE
EXPERIMENT STATION.

BULLETIN LXXXIV.

EXPERIMENTS WITH SPRING GRAINS

BY THOMAS SHAW, PROFESSOR OF AGRICULTURE, AND
C. A. ZAVITZ, B.S.A., EXPERIMENTALIST.

PUBLISHED BY THE DEPARTMENT OF AGRICULTURE

December 15, 1899.

TORONTO
PRINTED BY WARWICK & SONS

MINISTER OF AGRICULTURE

HON. JOHN DRYDEN, TORONTO.

Ontario Agricultural College and Experimental Farm, Guelph,
under control of the Minister of Agriculture.

JAMES MILLS, M.A.		President.
THOMAS SHAW..		Professor of Agriculture and Farm Superintendent.
A. E. SHUTTLEWORTH, B.A. Sc.		Professor of Chemistry.
J. HOYES PANTON, M.A., F.G.S.		Professor of Natural History and Geology.
F. C. GRENSIDE, V.S.		Professor of Veterinary Science.
H. H. DEAN, B.S.A.		Professor of Dairy Husbandry.
E. LAWRENCE HUNT, B.A.		Assistant Resident and Mathematical Master.
CAPTAIN WALTER CLARKE..		Instructor in Drill and Gymnastics.
O. A. ZAVITZ, B.S.A..		Experimentalist.
H. B. SHARMAN, B.S.A.		Assistant Chemist.
A. McOALLUM.		Bursar.

ADVISORY BOARD.

C. C. JAMES, M.A., Secretary		Deputy Minister of Agriculture, Toronto
JOHN I. HOBSON, Chairman		Mosborough, County of Wellington

BULLETIN LXXXIV.

EXPERIMENTS WITH SPRING GRAINS.

(140 Varieties grown side by side for four years.)

The great difference in point of yield of different varieties of grain of the same species has been observed by farmers everywhere. The wisdom of growing only the best varieties at all times will also be generally conceded, and more especially in times when prices rule low and margins are close, will such a line of action be in a sense imperative. In former years the husbandman was restricted in his choice to but few varieties, but now through the work of the Agricultural Experiment Stations, the best varieties that can be obtained are being made easily accessible to every one.

This Bulletin relates to spring grains, foreign and native, grown under similar conditions for the past four years, and it gives important comparisons with reference to them. These comparisons relate, (1) to the yield of grain per acre; (2) weight of grain per measured bushel; (3) weight of straw per acre; (4) date of maturity; (5) strength of straw; (6) length of plant, and (7) color of grain and other peculiarities of growth. Important particulars are also given in reference to results obtained from sowing various kinds of grain at different dates. As rust prevailed to a considerable extent, we have taken it into account when speaking of spring wheat and oats. During 1892, 67 varieties of barley, 63 of spring wheat, 116 of oats and 61 of peas, were grown upon the station grounds and under the same conditions. Full particulars relating to these will be found in the Annual Report. The information contained in the bulletin has been obtained from tests made with 37 varieties of barley, 22 of spring wheat, and 81 of oats. These have been grown side by side for the past four years and under similar conditions. It also includes the results obtained from 6 leading varieties of peas, out of 20 varieties grown for two years. The plots varied from one fiftieth to one-hundredth of an acre in the different years.

In the tables given below, the yields of grain per acre are estimated from the yields per plot. The standard weight per measured bushel was used with the different classes of grains, and when this bulletin is compared with the corresponding one of last year, viz., LXXXI, it will be observed that some varieties in the list for 1891 have dropped out and others have taken their places. It will also be noticed that the remarks on the different varieties vary slightly, owing to a difference in behavior, caused doubtless by variations in soil and climate.

Oats. Table I relates to the ten varieties of oats which have given the highest yields for the four years ending with 1892.

Variety.	Color of grain.	Amount of rust, 1892.	Height of plant, average, 4 years.	Date of maturity, 1892.	Weight per mea- sured bushel, 3 years.	Straw per acre, 4 years.	Grain per acre, bushel 34 lb., 4 years.
Joanette Black.....	Black..	None	in. 41.0	Aug. 7	lb. 35.8	tons 2.7	bush. 83.0
Chenailles Black.....	" ..	"	44.0	6	35.8	2.8	79.1
Black Etampes.....	" ..	"	40.8	7	36.4	2.6	79.0
Siberian	White..	Slight.....	54.0	6	36.7	2.5	77.2
Houdan Black.....	Black..	None	41.8	6	36.4	2.1	77.1
Danebrog.....	White..	Very slight.	54.8	8	33.7	2.4	75.4
Oderbrucker.....	" ..	Slight.....	49.0	7	30.2	2.5	75.1
Improved Besthorne...	" ..	Very slight.	50.3	8	33.5	2.0	74.7
Probsteler.....	" ..	Average...	53.8	8	33.0	2.5	74.5
Pringle's Progress.....	" ..	Slight.....	47.3	4	30.1	2.3	74.2

The varieties grown in 1889 and 1890 were sown April 22nd, those grown in 1891, April 29th, and those grown in 1892, April 18th. It should be borne in mind that all the leading varieties, except the last mentioned, that is Pringle's Progress, are foreign. They were imported for our station from Europe in 1889. Of the 81 varieties grown during the four years mentioned, 15 only were from a Canadian source, and these include nearly all the old varieties. The Joanette Black, Chenailles Black, Black Etampes and Houdan Black, are very much the same in all essential characteristics. So much so that the three first mentioned appear to be one and the same variety, though probably grown for some time past in different localities in France. These varieties are therefore more suitable for sowing on good rich land. They stand up fairly well, are not coarse, and are almost entirely free from rust tendencies. They are also inclined to shell easily when ripe. The hull is unusually thin, and is easily removed in threshing. The 81 varieties grown for four years gave an average yield in 1892 of 53 bushels per acre. Of these the 10 which gave the highest yields for four years made an average of 76.9 bushels per acre, and the 10 which gave the lowest yields 44.2 bushels. The fourteen best yielding varieties for the four years have all a spreading head. Of the 27 varieties grown for two years only, the Golden Giant and Magnet stand first and second respectively. The former gave an average yield of 89 bushels per acre, and the latter 86.4 bushels. Vick's American Banner came third, with an average yield of 86.3 bushels per acre. Of the 8 newly obtained varieties which have been tested for one year only, the American Beauty stands first. It gave 76.7 bushels per acre.

which have
1892.

Spring Wheat. Table II relates to the varieties of spring wheat which have given the highest yields for the four years ending 1892.

Straw per acre, 4 years.	Grain per acre, bushel 3/4 lb., 4 years.
tons.	bush.
2.7	83.0
2.8	79.1
2.6	79.0
2.5	77.2
2.1	77.1
2.4	75.4
2.5	75.1
2.0	74.7
2.5	74.5
2.3	74.2

Variety.	Seed for 1889. obtained from.	Bearded or bald.	Height of plant, average, 4 years.	Strength of straw, average, 3 years.	Date of maturity.	Weight per measured bushel, 3 years.	Straw per acre, 3 years.	Grain per acre (bush. 60 lb. 4 years.
			in.		Aug.	lb.	tons.	bush.
Herison Bearded...	France...	Bearded	40.5	Medium	11	63.5	1.7	26.0
Pringle's Champion	Germany	"	42.2	Medium	11	60.8	1.8	23.6
Saxonka.....	Russia...	"	40.6	Medium	12	60.2	1.5	22.6
Holben's Improved	Germany	Bald...	41.6	Strong.	16	59.0	1.7	22.0
Bart Trimenia.....	Greece...	Bearded	38.5	Strong.	13	62.8	1.5	21.8
Summer.....	Germany	"	36.8	Medium	11	57.6	1.6	20.1
Ordinary Bearded								
March.....	France...	"	38.3	Strong.	13	58.1	1.7	20.6
Konigsburg.....	Russia...	"	40.5	Strong.	13	62.1	1.5	20.4
Odessa Ghirka.....	Russia...	Bald...	40.3	Medium	14	59.9	1.6	20.2
Nenhert.....	Germany	"	43.4	Strong.	16	55.0	1.6	18.6

22nd, those
April 18th.
except the
They were
81 varieties
from a Cana-
dian varieties.
The
and Houdan
varieties. So
one and the
in different
suitable for
not coarse,
they are also
ly thin, and
own for four
per acre. Of
years made an
the lowest
eties for the
es grown for
t and second
bushels per
Banner came
p. Of the 8
e year only,
per acre.

The varieties grown in 1889 were sown April 18th, those in 1890 and 1891, April 25th, and those in 1892, April 23rd. While the Herison Bearded comes first in point of yield, stands fairly stiff in the straw, and has been almost entirely free from rust, the club-shaped heads are somewhat uneven. Pringle's Champion produces fairly strong straw, has a well shaped head, is not much liable to rust, and produces a fine sample of grain. The 22 varieties grown for four years gave an average of 19.2 bushels per acre in 1892. Of these the 5 which gave the highest yields for four years made an average of 23.2 bushels per acre, and the 5 which gave the lowest yields, 13.1 bushels. Of the 21 varieties grown for three years, the Red Fern comes first. It has given a yield of 31.4 bushels per acre. The White Russian, with a yield of 28.7 bushels, comes second. Of the 10 varieties grown for two years the McCarlin stands first. It has given a yield of 29.7 bushels per acre. The Rio Grande, with a yield of 28.4 bushels, comes second.

Hulless Barley.—Six varieties of Hulless barley have been grown for three years. The average yield per acre for that time has been 48 bushels, of 60 pounds per measured bushel, or 44.78 bushels of 48 pounds per bushel. If we compare the three leading varieties with the three best sorts possessed of hull, we find there is an advance in yield of 1.3 bushels per acre in favor of the former. The average yields respectively, are 54.8 bushels per acre, and 53.5 bushels. The Hungarian, from Hungary, as the name would indicate, stands first on the list, and in several respects it is a very promising variety.

Barley.—Table III relates to the ten varieties of barley which gave the highest yields for the four years ending 1892, and it gives the results obtained from the common six-rowed variety of Ontario.

Variety.	Seed for 1889, obtained from.	No. of rows per head.	Height of plant, average, 4 years.	Strength of straw, average 3 years.	Date of maturity, 1892.	Weight per mea- sured bushel, 3 years.	Straw per acre, 4 years.	Grain per acre bush. (48 lb.) 4 years.
Mandscheuri.....	Russia ..	6	41.0	Strong..	July 30	50.8	1.8	57.6
French Chevalier..	France ..	2	36.3	Medium	Aug. 3	52.0	1.8	54.0
Empress	England.	2	39.0	Medium	" 7	52.0	2.0	62.8
Scotch Improved...	Ontario..	6	39.0	Medium	July 29	52.3	1.3	51.3
Selected Chevalier.	England.	2	39.5	Weak..	Aug. 3	51.8	2.0	50.8
Thanet.....	"	2	38.5	Weak..	" 3	52.0	1.9	50.5
Improved Cheyne..	"	2	39.0	Medium	" 3	52.4	2.0	50.3
Oderbrucker.....	Germany	6	38.8	Strong..	July 29	52.9	1.5	49.9
Kalina.....	Sweden..	2	37.5	Medium	Aug. 3	52.6	1.8	49.9
Hallett's Pedigree	Germany	2	36.3	Strong..	" 7	52.8	1.7	49.7
Common Ontario, 6- rowed.....	Ontario..	6	36.3	Medium	July 29	53.1	1.3	46.7

The varieties grown in 1889 were sown April 15; those in 1890 and 1892, April 19, and those in 1891, April 30. It will be observed that the common Ontario six-rowed barley, so universally grown, has the seventeenth place in the order of yield. Of the eleven varieties grown for two years, the California Brewing stands first, and it also has first place amid the 63 varieties grown in 1892 when it yielded 61.5 bushels per acre. The Kinna-Kulla, a two-rowed variety, imported from Sweden four years ago, and which possesses the property of standing up well in a remarkable degree, is constantly improving in yield, insomuch that in 1892, it gave 58.6 bushels per acre, and stands second in point of yield. The Dutch bill, which bears a close resemblance to the Kinna-Kulla, is also still in the straw. The yield in 1892 was 47.8 bushels.

Peas.—Table IV relates to the ten varieties of peas which have given the highest yields during the two years 1890 and 1892.

Owing to imperfect germination of the seed in 1891, the results for that year were considered to be incomplete to be of any special value. The varieties grown in 1890 were sown April 24th, and those sown in 1892, April 23. Only two of the imported varieties, viz., the Princess Royal and Early Racehorse, have a prominent place in the list. The Mummy pea, popular in many parts of Ontario, gave an average of 33.7 bushels for the two years mentioned, and an average weight of 63.8 pounds. Of the sixty-one varieties grown

barley which
and it gives
of Ontario.

1892, the Princess Royal gave the largest yield, viz., 46.2 bushels, and the Early Racehorse the heaviest weight per measured bushel, viz., 62.8 pounds.

Variety.	Seed of 1890 obtained from.	Date of maturity.	Weight per measured bushel.	Straw per acre.	Grain per acre.
			lb.	tons.	bush.
Prussian Blue	Ontario	Aug. 11	62.7	1.94	43.39
Black Eyed Marrowfat	"	" 17	61.2	1.66	39.90
Princess Royal	England	" 15	60.7	1.45	39.15
White Eyed Marrowfat	Ontario	" 15	62.1	1.32	38.73
Multipliers	"	" 21	61.8	1.69	37.90
Early Racehorse	England	" 12	63.5	1.45	35.89

Dates of Seeding.—Table v relates to the average results from seeding oats, spring wheat, barley and peas at different dates in 1891 and 1892.

Dates of Seeding.	Oats.		Spring wheats.		Barleys.		Peas.	
	Weight per measured bushel.	Grain per acre (bu 34 lb).	Weight per measured bushel.	Grain per acre (bu 60 lb).	Weight per measured bushel.	Grain per acre (bu 48 lb).	Weight per measured bushel.	Grain per acre (bu 60 lb).
	lb.	bush.	lb.	bush.	lb.	bush.	lb.	bush.
April 22nd	33.0	52.7	60.7	14.4	48.7	26.1	57.9	19.5
May 1st	31.9	56.9	58.8	12.2	47.0	27.6	59.2	21.1
May 9th	29.2	53.7	56.7	8.6	46.0	27.2	60.7	21.8
May 18th	27.5	41.8	54.2	5.4	42.7	18.3	61.0	18.5
May 26th	24.4	25.3	53.0	3.9	41.9	11.8	61.2	9.7

In every instance, except that of the peas, which were grown only in 1892, and of which but two varieties were used, the figures represent the average of two varieties sown in 1891, and also in 1892, since they represent the average results of four experiments with the same kinds of grain. It will be observed (1) that oats, spring wheat and barley gave the heaviest weights per measured bushel from the first seeding, and there was a gradual decrease in weight with each successive seeding. With peas the results were exactly the reverse, as the weight of the grain continually increased with each seeding. (2) That while in the case of spring wheat, the highest yields were obtained from the first seeding, in that of oats and barley these came from the second seeding, and in that of peas from the third seeding. But when the tests for 1892 only are compared, we find that the best yields obtained in every instance came from the first seeding, except with peas.

Remarks.—The soils where these plots were grown may be termed a mild clay loam, with a somewhat open sub-soil. The field containing the plots grown in 1889 and 1890 was low lying, and that in which the plots were grown in 1891 and 1892 was elevated.

Conclusions.

1. That considerably higher yields may be expected from several of the foreign varieties of oats, than from the old varieties.

2. That of all the varieties of oats grown for four years, the Joannette Black stands first in point of yield among the black varieties and the Siberian among the white varieties.

3. That the Herison Bearded and Pringle's Champion still retain first and second places in point of yield, among all the varieties grown for four years.

4. That fifteen imported varieties of barley have given a larger yield for four years than the common six-rowed variety.

5. That of all the varieties of barley grown for four years, the Mandscheuri and French Chevalier still hold first and second places respectively, in point of yield.

6. That of twenty varieties of peas grown for two years the Prussian Blue stands at the head of the list in point of yield, and of sixty-one varieties grown in 1892, the Princess Royal gave the largest yield per acre, and the Early Racehorse the heaviest weight per measured bushel.

7. That while it is important that cereal spring grains should as a rule be sown as soon as possible after the ground has become sufficiently dry in the spring, it is more important relatively in the case of spring wheat than in that of oats and barley, while no special advantage seems to follow very early the sowing of peas.

Distribution of Seed.—We have a limited quantity of seed for distribution, and will supply any of the following varieties at moderate prices, and in the order in which the applications are received, viz *Oats*—Joannette Black, Black Etampes, Siberian, Probsteler, Poland White, Improved Waterloo, Bavarian, White Abundance and Golden Giant. *Peas*—Prussian Blue, Tall White Marrowfat, Canada Cluster, Prince Albert and D'Auvergne. *Barley*—Mandscheuri and Odenbrucker. Apply to the Professor of Agriculture, Guelph, Ont.

Some of the varieties will be distributed in smaller lots, through the medium of the Ontario Agricultural and Experimental Union. For circulars apply to the Secretary, O. A. Zavitz, Experimental Station, Guelph.

a
ng
ch

ral

the
ack

tain
own

rgen

, the
laces

the
nd of
argest
t per

uld as
suffi
e case
pecial

for dis
oderat
l, viz
Poland
Golden
a Clus
d Ode
t.

through
Union
ntal St