**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1981

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured covers/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☐ Covers restored and/or laminated/
Couverture restaurée et/ou pelliculée
- ☒ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☐ Bound with other material/
Relié avec d'autres documents
- ☐ Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- ☐ Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- ☐ Additional comments:/
Commentaires supplémentaires:

- ☐ Coloured pages/
Pages de couleur
- ☐ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☐ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☐ Pages detached/
Pages détachées
- ☐ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	14X	18X	22X	26X	30X
☐	☐	☐	☒	☐	☐
12X	16X	20X	24X	28X	32X

The copy filmed here has been reproduced thanks to the generosity of:

Library of Congress
Photoduplication Service

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol → (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Library of Congress
Photoduplication Service

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole → signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

37
1

from the *Canadian Naturalist*, Vol. VIII. No. 6.)

LOWER CARBONIFEROUS FISHES OF NEW BRUNSWICK.

By PRINCIPAL DAWSON, LL.D., F.R.S.

The recent sinking of a shaft on the property of the Beliveau Albertite and Oil Company on the Petitecodiac River, has exposed a new and interesting deposit of fossil fishes in the rich bituminous shales of that district, which contain the remarkable deposits of Albertite, described in my *Acadian Geology*, second edition, p. 231 *et seq.* The bed affording these fossils is a dark brown bituminous shale; and I am informed by Mr. E. B. Chandler, to whom I am indebted for an interesting collection of the fish remains, was from four to five feet thick. The specimens thus presented, with those previously in my collection, and one kindly given to me by Mr. F. Adams, of this University, and the valuable memoirs recently published by Dr. Newberry in the *Ohio Reports*, and by Dr. Traquair in the *Journal of the Geological Society*, enable me now to give a revision of the fishes of this locality, as described by Dr. Jackson in his Report of 1851 on the Albert mine, which I was unable to do in the second edition of *Acadian Geology*, owing to the small number of specimens to which at that time I had access.

In the collections in my possession, I recognize, in all, five species, three of them very small, and two of larger size. Of these, one, which is unusually well preserved and is the smallest of the whole, appears to be new, and I shall begin by describing it.

Palæoniscus (Rhadinichthys) Modulus, n. s.—Length, five to six centimetres; greatest breadth, 15 to 17 millimetres—the proportion of length to breadth being about five to one and a half. Head, oval and obtuse; details not preserved, except that the bones are sculptured with fine waving lines. Body gracefully curved, and upper lobe of tail long and slender. Pectoral fins small, with stout, unjointed rays. Ventral not distinctly preserved, but apparently small and nearer to pectorals than to anal. Dorsal and anal of moderate size and opposite each other. Caudal very heterocercal, with the lower lobe sharply pointed. Fins with well developed fuleral spines, especially large at the base of the caudal. Scales of the sides rhombic, coarsely toothed on the posterior edges and elaborately sculptured with flat, scaly ridges,

corresponding to the teeth of the edge. The ridges are arranged in an upper and lower series, the latter oblique to the former, so that each scale has the appearance of being composed of two distinct portions. Lower surface of scales smooth, with a few furrows corresponding to the ridges above, and the posterior edges similarly serrate. Caudal scales narrowly rhombic, pointed, and with a few central lines. The back is protected with about ten large oval scales between the head and the dorsal. They are sculptured with waving lines, curving with the edges, and are apparently truncate and serrate behind. The fish figured by Jackson, Pl. II, Fig. 5, but not named, probably belongs to the above species.

Fig. *Palaeoniscus Modulus*, N. S.

- (a) Outline, natural size.
- (b) Series of Scales enlarged, seen from inside. The lower row are those on mesial line.
- (c) Surface of exposed part of scale from side and upper lobe of tail, showing sculpture, enlarged.
- (d) One of the dorsal scales, enlarged.

This beautiful and elaborately ornamented little fish is a perfect model in miniature of that type of lower carboniferous Palaeoniscids to which it belongs, and which has recently been separated by Dr. Traquair in the genus or subgenus *Rhadinichthys*. For this reason, I have given it the specific name *modulus*. To the same genus belong the two next species, described by Jackson, of which I shall give merely distinctive marks.

P. Alberti, Jackson, is larger than the preceding. The scales have more numerous striae. The dorsal scales are rounded pos-

teriorly. The posterior edge of the anal fin approaches nearly to the caudal, and extends considerably behind the posterior edge of the dorsal.

P. Guirnsii, Jackson.—About the same size with the last, but more slender, and the head less obtuse in front. Scales thick and with few striae, and less numerous serrations. Dorsal scales pointed posteriorly. Anal fin somewhat remote from caudal and opposite dorsal.

A specimen collected by Mr. Ells, of the Geological Survey, indicates a fish of the same general form with *P. Alberti*, but about six inches long. The outline of this fish is well seen, but the details are not sufficiently clear to show if it differs in these from the smaller species.

The next species and perhaps the following one, belong to the genus *Elonichthys* of Giebel. They are much larger than the preceding.

P. Brownii, Jackson, is deep in form, with large dorsal and anal, the latter reaching almost to base of caudal. Scales of body broad and with numerous fine horizontal striato-punctate furrows, which turn abruptly upward at the anterior side of each scale. A nearly perfect specimen, collected by Mr. Ells, shows that the head was of moderate size, and the body about ten inches long and three and a quarter inches wide, the breadth at the dorsal fin being as great as at the shoulders, giving a sort of rectangular form to the fish, whose breadth suddenly diminishes toward the tail.

The crystalline lens of the eye of Mr. Ells's specimen is preserved in calcite. Under the microscope it shows concentric laminae and coarse bands or rods with indistinct denticulations; the structure being similar to that in the crystalline lens of the modern ganoid *Amia ocellicauda*. This is the first instance known to me of the preservation of the structure of the crystalline lens in a palaeozoic fish.

P. Jacksoni, n. s.—A species figured, but not described, by Jackson, is represented by many fragments in my collection. It is the largest of these fishes, reaching a length of 15 inches. It may be distinguished from the last by its more slender form, its small anal fin, more remote from the caudal, and by the character of the scales, which have many horizontal striae, and have in the broader ones a few deep and strong serrations posteriorly.

t.c. H. O. Apr. 27, 12

The whole of these fishes have been preserved entire, the body being perfectly flattened and thrown into attitudes which imply that they were imbedded when living or immediately after death. The material in which they are contained is shown, by its microscopical and chemical characters, to have been a vegetable muck or mud, and the fish were either overwhelmed by it in the manner of a bursting bog, or were stifled by the non-oxygenated water mixed with this mud, and suddenly killed and imbedded in the accumulating sediment. That they occur in this perfect state and in a limited thickness of the deposit, may imply that at certain times they were overwhelmed by the irruption of this fetid organic mud into the water in which they lived. The bed is low down in the Lower Carboniferous series, being the equivalent of the Horton series of Nova Scotia; so that these fishes are among the oldest that we know in the Carboniferous period; but we know, from the Horton beds, that many far larger and predaceous ganoids were their contemporaries. No remains of these have however as yet been found in the Albert or Beliveau beds, which were probably deposited in limited fresh-water basins, perhaps not ordinarily accessible to the larger fishes.

Sir Philip Egerton* and Dr. Traquair† have both remarked on the similarity of these fishes to those found in the Lower Carboniferous of Scotland, and Dr. Newberry has described very similar species from the Carboniferous of Illinois and Ohio.‡

* Journal of Geological Society, 1853.

† *ib.* 1877.

‡ Report on Illinois, Vol. II; Palaeontology of Ohio, Vol. I.

-1899.

ire, the body
which imply
y after death.
n, by its mi-
a vegetable
by it in the
oxygenated
ud imbedded
this perfect
y imply that
ption of this
l. The bed
e equivalent
e fishes are
period ; but
ger and pre-
nins of these
liveau beds,
basins, per-

h remarked
the Lower
s described
and Ohio.†

ol. 1.