+QK110
.E18

Chis
250

A
MANUAL OF BOTANY
FOR
The Northern States.

COMPRISING

GENERIC DESCRIPTIONS OF ALL PHENOGAMOUS AND CRYPTO-
GAMOUS PLANTS TO THE NORTH OF VIRGINIA, HITH-
ERTO DESCRIBED; WITH REFERENCES TO
THE NATURAL ORDERS OF LIN-
NEUS AND JUSSIEU.

Each Genus is further illustrated by short Descriptions of
its most common Species.

BY THE MEMBERS OF THE BOTANICAL CLASS IN
WILLIAMS' COLLEGE, (Mass.)

From a Manuscript System,

COMPILED BY THE AUTHOR OF

RICHARD'S BOTANICAL DICTIONARY.

ALBANY:

PRINTED BY WEBSTERS AND SKINNERS.

1817.

+2K10
E12

TO THE AUTHOR

OF
RICHARD'S BOTANICAL DICTIONARY.

DEAR SIR,

WE consider ourselves happy in having an opportunity to express our gratitude for the assiduity and care, which you have manifested for our improvement in Natural Science. As the course of Lectures on Mineralogy, which you have conducted so entirely to the satisfaction of the Faculty of this College, and so much to the benefit of your class, is nearly completed, we are gratified with the prospect of attending your course of Lectures on Botany—Knowing that our improvement in that branch of Natural Science would be greatly facilitated by a systematic description of vegetables, and being destitute of such a system, we render you our thanks for the one, which you have been pleased, gratuitously, to present to us for publication.

Accept, dear Sir,

The assurances of our respect and esteem.

Williams' College, 8th April, 1817.

H. Walker Bishop,

Dorus L. Clark,

William Eastman,

Luther Hamilton,

Lyman James,

Medad Pomeroy,

Emory C. Washburn,

Selah R. Arms,

Daniel D. Barnard,

Homer Bartlett,

Chandler Bates,

Charles Dillingham,

William A. Hallock,

Gerard H. Hallock,

Anson T. Hooker,

Cyrus M. Lazell,

Lucas Morgan,

Joseph P. Mosher,

William Richards,

Hervey Smith,

Thomas Spring,

B. Horace Starkweather,

Lemuel P. Bates,
George W. Benedict,
Daniel I. Betts,
David L. Coe,
Joshua N. Danforth,
Ebenezer Emmons,
Joseph H. Estabrook,
Charles Fitch,
Parker L. Hall,
Gardner Hayden,
Leicester Lloyd,
Thomas Peck,
William A. Porter,
John B. Skinner,
Royal W. Smith,
William Wells,
John Whilton,
John C. Brigham,
Nelson Brown,
Elijah H. Burrit,
Samuel Dickinson,

Alvan Alvord,
Charles Baker,
Dwight Baldwin,
Johnson Baldwin,
Andrew Burnham,
Benjamin F. Clark,
Gardner Dorrance,
Jonathan Ely,
Simon C. Ewers,
Abner Forbes,
Mason Fussell,
William Gildersleeve,
Edward Hooker,
William W. Hunt,
Royal Joy,
William C. Kittrige,
John C. Morgan,
Isaac Oakes,
Elijah Thayer,
Alvan Wheeler.

PREFACE.

EVERY one, who has passed through the first struggles of a student in practical botany, recollects the embarrassments in which he was perpetually involved, by the multitude of foreign genera contained in the European systems.— Doctor Bigelow's Boston Flora proves the very great relief which is afforded to the student by a system of plants limited to his own particular district. But there is a middle way in this as in other things. The northern section of the United States is sufficiently contracted for this object; and the common cultivated exotics ought certainly to be included.

Another obstacle to the progress of the student is the very concise generic characters given in most of the works of this nature. If he sets out with the right class, is led to the right order, and fixes at last upon the right genus, his work is reduced to a manageable state. To guide him through those three steps is the object of this Manual. To effect this with ease to the learner, the common English names, the most striking specific characters, together with a few popular references, and the natural orders of Linneus and Jussieu, have been annexed to each generic description.

The generic characters of the phenogamous plants are translated from Persoon, and arranged principally upon his method. Additions and corrections were made, however, on comparing the translation with the systems of Linneus, Pursh, Muhlenberg, Willdenow, Bigelow, the Hortus Kewensis, the Encyclopædia, and the Synopsis of American Genera.

The descriptions of cryptogamous plants are chiefly from Sprengel, and Doctor Smith's translation of Acharius. The orders, Algae and Fungi, are mostly from Turton and Linneus.

This work should be accompanied by the Botanical Dictionary, published at New-Haven, which was translated from the French of Professor Richard; and it is so constructed that it may be bound in the same volume.

LINNEAN SYSTEM OF VEGETABLES.

All Vegetables are divided into twenty-two* CLASSES. These CLASSES are divided into ORDERS. ORDERS are divided into GENERA. GENERA are divided into SPECIES. SPECIES are frequently changed into *varieties*. Varieties, however, are more properly within the province of the Gardener, than of the Botanist; at least the method of procuring varieties.

When a Botanist sees a plant, which he never saw before, and wishes to know its name and uses; he proceeds as follows.

1. He takes the unknown flower in his hand (no unknown plant can be ascertained without the flower) and compares its parts with the description of each class, until he finds the class to which it belongs.

2. He then goes to the orders of that class and finds its order in the same way.

3. Next he goes to the genera of that order, and reads their descriptions, until he finds the genus to which it belongs.

4. At last looks over the species of that genus, until he finds the exact description of his plant.

5. Thus he finds the Apple to be CLASS 12, ORDER 5, GENUS *Pyrus*, SPECIES *Malus*.

* Linneus divided them into 24 classes. But farther discoveries, since his death, have proved the classes Polyadelphia and Polygamia to be too uncertain and variable to be any longer retained. Persoon, therefore, and other eminent Botanists, have rejected them. See these classes in the Dictionary.

LINNEAN CLASSES.

1. MONANDRIA, 1 stamen, or one sessile anther in the flower.
2. DIANDRIA, 2 stamens, or 2 sessile anthers.
3. TRIANDRIA, 3 stamens, or 3 sessile anthers.
4. TETRANDRIA, 4 stamens, or 4 sessile anthers.
5. PENTANDRIA, 5 stamens, or 5 sessile anthers.
6. HEXANDRIA, 6 stamens, or 6 sessile anthers.
7. HEPTANDRIA, 7 stamens, or 7 sessile anthers.
8. OCTANDRIA, 8 stamens, or 8 sessile anthers.
9. ENNEANDRIA, 9 stamens, or 9 sessile anthers.
10. DECANDRIA, 10 stamens, or 10 sessile anthers.
11. DODECANDRIA, 12 to 19 stamens, or sessile anthers.
12. ICOSANDRIA, about 20, or more, standing on the calyx.
13. POLYANDRIA, always 20 or more, on the receptacle.
14. DIDYNAMIA, 4 stamens, 2 of them uniformly the longest.
15. TETRADYNAMIA, 6 stamens, 4 of them uniformly the longest.
16. MONADELPHIA, stamens united by their filaments in one set, anthers remaining separate.
17. DIADELPHIA, stamens united by their filaments in two sets (sometimes in one set) flowers papilionaceous.
18. SYNGENESIA, stamens 5, united by their anthers in one set, flowers compound.
19. GYNANDRIA, stamens stand on the germ, style, or stigma, separate from the base of the calyx and corol.
20. MONOECIA, stamens and pistils in separate flowers, on the same plant.
21. DIOECIA, stamens and pistils on separate plants.
22. CRYPTOGAMIA, stamens and pistil so obscure that the plants can only be classed by natural families.

ORDERS OF EACH CLASS.

Class	1st.	2d.	3d.	4th.	5th.	6th.	7th.	8th.	10th.	13th	16th.
1	Mon.	Dig.	—	—	—	—	—	—	—	—	—
2	Mon.	Dig.	Tri.	—	—	—	—	—	—	—	—
3	Mon.	Dig.	Tri.	—	—	—	—	—	—	—	—
4	Mon.	Dig.	—	Tet.	—	—	—	—	—	—	—
5	Mon.	Dig.	Tri.	Tet.	Pen.	—	—	—	—	—	—
6	Mon.	Dig.	Tri.	Tet.	—	Hex.	—	—	—	—	—
7	Mon.	Dig.	—	Tet.	—	—	Hep.	—	—	—	—
8	Mon.	Dig.	Tri.	Tet.	—	—	—	—	—	—	—
9	Mon.	—	Tri.	—	—	Hex.	—	—	—	—	—
10	Mon.	Dig.	Tri.	—	Pen.	—	—	—	—	—	—
11	Mon.	Dig.	Tri.	Tet.	Pen.	—	—	—	—	—	—
12	Mon.	—	—	—	Pen.	—	—	—	Pol.	—	—
13	Mon.	Dig.	—	—	Pen.	—	—	—	—	Pol.	—
14	Gym.	Ang.	—	—	—	—	—	—	—	—	—
15	Silic.	Siliq.	—	—	—	—	—	—	—	—	—
16	—	—	Tri.	—	Pen.	—	—	—	Dec. Pol.	—	—
17	—	—	—	—	—	Hex.	—	Oct.	Dec.	—	—
18	Pol. Acq.	Pol. Sup.	Pol. Frus.	Pol. Nec.	Pol. Seg.	—	—	—	—	—	—
19	Mon.	Dia.	—	—	Pen.	Hex.	—	—	—	—	—
20	Mon.	Dia.	Tri.	Tet.	Pen.	Hex.	—	—	Pol. Mon.	—	—
21	—	Dia.	Tri.	Tet.	Pen.	Hex.	—	Oct.	—	Pol. Mon.	—
22	Filices.	Musci.	Hepaticae.	Algae.	Lichenes Fungi.	—	—	—	—	—	—

Note. The numbers at the head of this page are used to express the orders directly under them.

EXPLANATIONS FOR THE PRECEDING PAGE.

- Mon. Monogynia, 1 style, or 1 sessile stigma.
 Dig. Digynia, 2 styles, &c.
 Tri. Trigynia, 3 styles, &c.
 Tet. Tetragynia, 4 styles, &c.
 Pen. Pentagynia, 5 styles, &c.
 Hex. Hexagynia, 6 styles, &c.
 Hep. Heptagynia, 7 styles, &c.
 Dec. Decagynia, 10 styles, &c.
 Pol. Polygynia, more than 10 styles, &c.
 Gym. Gymnospermia, seeds naked.
 Ang. Angiospermia, seeds in capsules.
 Silic. Siliculosa, having pods whose length and breadth are nearly equal.
 Siliq. Siliquosa, having pods whose lengths are more than double their breadths.

In the 16th, 17th, 19th, 20th, 21st classes, the names and characters of preceding classes are taken for orders. In which, Mon. Monandria. Dia. Diandria. Tri. Triandria. Tet. Tetrandria. Pen. Pentandria. Hex. Hexandria. Oct. Octandria. Dec. Decandria. Pol. Polyandria. Mon. Monadelphia.

In the 18th class. 1. Pol. *Æq.* Polygamia *Æqualis*. 2. Pol. Sup. Polygamia *Superflua*. 3. Pol. Frus. Polygamia *Frustranea*. 4. Pol. Nec. Polygamia *Necessaria*. 5. Pol. Seg. Polygamia *Segregata*.

The 1st order in the 18th class is distinguished by having *all the florets* perfect. The 2d, by having those of the *disk* perfect, while those of the *ray* are pistillate. The 3d, by having those of the *disk* perfect, while those of the *ray* are neutral. The 4th, by having those of the *disk* staminate, while those of the *ray* are pistillate. The 5th, by having the *florets* all perfect, while each *floret* has a *perianth* of its own.

In the 22d class, the orders are distinguished by natural family characters. 1. *Filices*, (ferns) which bear fruit on the back of the leaves, or in which some part of the leaves seem as it were metamorphosed into a kind of fruit-bearing-spike. The appendix to this order includes the *Pteriodes*, which bear fruit on a peculiar appendage. 2. *Musci*, (mosses) which bear, on leafy stems and branches, one-celled capsules, opening at the top, where they are covered by a peculiar lid. 3. *Hepaticae*, (liverworts) which bear, on herbaceous fronds, four-celled capsules opening with four valves. 4.

Algae, (seaweeds, &c.) which bear, in an aquatic or gelatinous frond, vesiculous or filamentous fruit. 5. *Lichenes*, (lichens) which bear, on fibrous, compact or gelatinous fronds, scutellae, patellulae, peltae, thalamia, disks or pilidia, containing the fructification. 6. *Fungi*, (mushroom, &c.) which are destitute of herbage, consisting of a spungy, pulpy, leathery or woody substance, and bear fruit in a naked dilated membrane, or within the substance of the plant.

EXPLANATIONS OF THE CHARACTERS ANNEXED TO THE GENERIC DESCRIPTIONS.

The *first number* following the generic description is the number of the natural order of Linneus, to which the genus belongs; the *second number* is that of Jussieu. By referring to the natural orders of Linneus in the Botanical Dictionary, the qualities of plants may be so far ascertained, as they depend on these natural affinities.

☉ annual. ♂ biennial. ♀ perennial. ♀ woody.

p, when the corol is purple. r, red. w, white. y, yellow. g, greenish. e, exotic.

* Placed before one or more generic nouns, at the end of an order, indicates that these genera sometimes vary from the classes and orders to which they belong, and are found to possess the characters of the class and order under which their names are thus set down.

A.

SYSTEM OF GENERA

FOR THE

NORTHERN STATES.

CLASS I. MONANDRIA.

ORDER I. MONOGYNIA.

SALICORNIA. Calyx inflated, entire: corol o: seed 1, inclosed in the calyx. 12. 29—*virginica*, (samphire) branches undivided—*herbacea*, divided.

☺.

HIPPURIS. Calyx superior, obsolete, with a 2-lobed margin: corol o: seed 1: stigma simple. 15. 6.—*Vulgaris*, (maretail) leaves linear, whorled. 2.

* *Scirpus*, *Cyperus*.

ORDER II. DIGYNIA.

CALLITRICHE. Calyx inferior, 2-leaved: capsule membranaceous and margined: (flowers sometimes monoecious, and by some the calyx is called the corol.) 12. 6—*autumnalis*, (water-starwort) leaves all linear—*intermedia*, stem-leaves linear only—*verna*, flowers androgynous. ☺.

BLITUM. Calyx 3-cleft, berry-like: corol o: seed 1. 12. 29.—*capitatum*, (blite) heads alternate—*virgatum*, heads scattered. ☺. r.

CINNA. Calyx a 1-flowered glume, 2-valved: corol a 2-valved glume: seed 1. 4. 10—*arundinacea*, panicle large, many-flowered. 2.

* *Uniola*.

B

CLASS II. DIANDRIA:

ORDER I. MONOGYNIA.

A. Corol inferior, 1-petalled regular: border, except in *Jasminum*, 4-parted.

JASMINUM. Corol salver-form, 5 to 8—cleft: berry 2-seeded, each seed solitary, arilled. 44. 37.—*officinale*, (jasmine) leaves pinnate. h. w. e.

LIGUSTRUM. Calyx 4-toothed: corol with ovate divisions: berry 4-seeded. 44. 37—*vulgare* (prim) panicled. h. w.

SYRINGA. Corol salver-form: capsule 2-celled. 44. 37—*vulgaris*, (lilac) leaves cordate. h. p. and w. e.

B. Corol inferior, 1-petalled, irregular: fruit capsular.

UTRICULARIA. Calyx 2-leaved, equal: corol ringent, spurred: capsule 1-celled, globular. 24. 34—*vulgaris* (bladder-wort) floats with air-blisters. 24. y.

CATALPA. Corol 5-cleft: calyx 2-leaved: capsule 2-celled. 40. 45—*syringacifolia*, (catalpa tree) leaves cordate. h. w. p. y.

JUSTICIA. Calyx simple or double, 5-parted: corol ringent or nearly equal: capsule 2-celled, busting with an elastic claw: partitions transverse. 40. 36—*adhatoda*, (malabar nut) leaves lance-ovate. h. p.

GRATIOLA. Calyx 7-leaved, the 2 outer ones spreading: corol 4-cleft, 2-lipped, reversed: stamens 4; (2 of them barren) stigma 2-lipped: capsule 2-celled. 40. 40—*officinalis*, (hedge-hyssop) leaves slightly toothed. [This is the *aurea* of Muhlenberg.] ☉. y.

VERONICA. Calyx 4-parted: corol 4-cleft, lower division smaller: capsule 2-celled. 40. 35—*serpyll-*

CLASS II. ORDER II.

49

lifolia, (speedwell) racemes terminal. $\frac{1}{2}$. w. b.
Scutellata, racemes lateral. $\frac{1}{4}$. w. b.

C. Corol inferior, 1-petalled, irregular: seeds
 naked.

LYCOPUS. Calyx tubular, half 5-cleft: corol
 tubular, 4-cleft, nearly equal, 1 division emargin-
 ate: stamens distant: seeds 4, retuse. 42. 39—*eu-*
ropeus, (water-horehound) leaves below gash-
 toothed, above serrate. $\frac{1}{4}$ —*virginicus*, leaves
 equally serrate, base narrow. $\frac{1}{4}$ -w.

MONARDA. Calyx cylindric striated, 5-toothed:
 corol ringent, upper lip linear, involving the fila-
 ments. 42. 39—*didyma*, (Oswego tea) heads in
 whorls. $\frac{1}{4}$. r.—*allophylla*, (mountain mint) leaves
 acute-serrati. $\frac{1}{4}$. p.

ROSMARINUS. Corol ringent, upper lip 2-part-
 ed: filaments long, curved, simple, with a tooth.
 42. 39—*officinalis* (rosemary) leaves hairy beneath.

SALVIA. Calyx tubular, 2-lipped, underlip 2-
 toothed: corol ringent: filaments transversely af-
 fixed to a pedicel. 42. 39—*officinalis*, (sage) leaves
 crenate. $\frac{1}{4}$. b. p. e.—*urticaefolia*, leaves doubly-
 serrate. $\frac{1}{4}$. b.

COLLINSONIA. Calyx tubular, 2-lipped: corol
 unequal, underlip many-cleft, capillary: one per-
 fect seed. 40. 39—*canadensis*, (horse balm) leaves
 heart-ovate. $\frac{1}{4}$. y.

D. Corol superior.

CIRCAEA. Calyx 2-leaved: corol 2-petalled:
 capsule hispid, 2-celled, not gaping; cells 1-seed-
 ed. 48. 88—*lutetiana*, (enchanter's nightshade)
 leaves ovate, a little toothed. $\frac{1}{4}$. w.

* Cunila, Schoenus, Scirpus, Verbena.

ORDER II. DIGYNIA.

ANTHOXANTHUM. Calyx glume 2-valved, 1-

flowered: corol glume 2-valved, acuminate, awned: seed 1. 4. 10—*odoratum*, (sweet vernal grass) spiked; floret larger than awn. 24—An American variety, *altissimum*, is larger and of a darker green. Ives.

* *Holcus*, *Festuca*, *Saccharum*.

CLASS III. TRIANDRIA.

ORDER I. MONOGYNIA.

A. *Flowers superior.*

VALERIANA. Calyx 0, or with an extremely small margin: corol 1-petalled, 5-cleft, base gibbous: seed 1: stamens, 1, 2, 3 and 4, exsert. 48. 56—*pauciflora*, (valerian) root-leaves pinnate, stem-leaves ternate. 24. r.

FEDIA. Calyx 3 to 6-toothed: corol 5-parted: nut 2 or 3-celled: seed naked, or crowned with a tooth. 48. 56—*radiata*, (lamb lettuce) leaves spatulate. ☉. b.

IRIS. Calyx, spathe 2-valved: corol 6-parted, divisions alternately reflected: stigmas petal-like, 6. 18.—*pumila*, (dwarf flower-de-luce) scape shorter than the leaves. 24. b.—*virginica*, (wild iris or flag) stem 2-cornered, with alternate leaves. 24. y. b.—*gracilis* (Boston iris) petals beardless, inner ones erect: leaves linear, flat: germ triangular, twice grooved on each side. Bigelow. This plant is very abundant near New-Haven as well as Boston.

B. *Flowers inferior.*

COMMELINA. Calyx, spathe cordate: corol 6-petalled: nectaries 3, cross-form, inserted on peculiar filaments: capsule sub-globose, 3-celled. 6. 13—*communis* (day-flower) stem creeping. ☉. b. e.

LEPTANTHUS. Calyx, spathe 1-flowered: corol long-tubular; border 6-parted with the stamens on

the divisions : capsule 3-celled, many-seeded, gaping at the angles. 3. 1—*reniformis*, leaves round-cordate ; spathe. 2. w.

XYRIS. Calyx, glume 2-valved, in a head : corol 3-petalled, equal, crenate : capsule 3-valved, many-seeded, 6. 13—*anceps*, (yellow-eyed grass) culm 2-edged. 2. y.

C. *Flowers grassy : valves of the calyx glume-like : corol none.*

SCHOENUS. Spikelets sub-convolute, acute ; scales heaped in fascicles, outer ones dry, shining : corol 0 : seed 1, roundish, naked or surrounded with bristles. 3. 9—*albus*, (bog-rush) leaves bristle-form. 2. —*glomeratus*, leaves flat. 2.

CYPERUS. Glumes chaffy, scales imbricated 2-ways : seed single, beardless : spikelets compressed. 3. 9—*spathaceus*, (galingale) axillary racemes. 2. —*rotundus*, umbel doubly-compound. 2.

SCIRPUS. Glumes chaffy, scales imbricated every way : seed single, naked, surrounded with hairs or bristles. 3. 9—*macrostachius*, (club-rush) panicle clustered, leafy, terminal. 2. —*triqueter*, spikelets lateral, sessile, bunched. 2. —*retrofractus*, umbels simple. 2. —*acutus*, culm round, very tall. 2.

ERIOPHORUM. Glume chaffy, imbricated every way : seed beset round with very long dense woolly hairs. 3. 9—*cyperinum*, (cotton grass) panicles thrice-compound, one growing from the top of another. 2. —*angustifolium*, leaves channelled, triangular at top—*polystachium*, leaves flat, spikes peduncled. 2.

TRICHOPHORUM. Calyx, scales imbricated every way : seed beset with capillary bristles, which at length project out, always 6 in number : spikelets ovatis. 3. 9—*eriphorum*, (light-hair) culm leafy : spike crowded.

D. *Flowers grassy: valves of the calyx glume-like: corol 2-valved.*

SPARTINA. Calyx 2-valved, compressed, one of them keeled and longer than the other: corol 2-valved. 3. 9—*cynosuroides*, (salt-grass) peduncles rough. 2—*juncea*, peduncles smooth. 2—*glabra*, valves of calyx mostly glabrous. 2.

CENCHRUS. Involucre divided, echinate, 3 or 4-flowered: glumes 2-valved, 2-flowered: corol 2-valved, awnless: style 2-cleft, 4, 10—*echinatus* (hedgehog grass) spike conglomerate.

* *Juncus*, *Galium*.

ORDER II. DIGYNIA.

The proper Grasses.

A. *Flowers perfect.*

1st. *Glume 1-flowered.*

ORYZOPSIS. Calyx 2-valved, lax, obovate: corol teretish-ovate, leathery: valves 2, outer one awned at the apex: appendages 2, linear, chaffy, 4. 10—*asperifolia*, (mountain rice) leaves rough.

PANICUM. Calyx 3-valved, the third valve dorsal and very minute: corol 2-valved: inflorescence various. 4. 10—*crus-galli*, (cockfoot grass) spike doubly-compound. ☺—*glaucum*, (panic grass) spike terete. ☺—*verticillatum*, spike whorled. ☺—*latifolium*, leaves broad and hairy at the neck of the sheath. 2—*nitidum*, panicle very branching, flowers minute.

DIGITARIA. Calyx 2 or 3-valved, concave; the outer one very small or none, the inmost one of the length of the corol: corol 2-valved, oblong-ovate, awnless: style very long: spikes digitate, linear: flowers in pairs. 4. 10—*sanguinalis*, (finger grass) culm creeping, flowers imbricate. ☺—*dactylon*, scions creeping, spikes finger-form. 2.

ARISTIDA. Calyx 2-valved: corol 1-valved, with 3 awns at the tip. 4. 10—*dichotoma*, (beard grass) culm forked.

ALOPECURUS. Calyx 2-valved: corol 1-valved, simple at the tip; sometimes awned at the base. 4. 10—*pratensis*, (foxtail grass) culm erect. 2/ —*geniculatus*, culm geniculate.

PHALARIS. Calyx 2-valved, valves keeled, nerved, equal in length, including the 2-valved pilose corol. 4. 10—*arundinacea*, (ribbon grass) panicle oblong, compact, leaves particoloured. 2/ e.

HORDEUM. Calyx lateral, 2-valved, 1 or 2-flowered; florets in threes, the middle one sessile, lateral ones often barren: corol 2-valved, acute, outer valve awned. 4. 10—*vulgare* (barley) florets in 2 erect rows. e.—*jubatum*, involucre bristle-form.

MILIUM. Calyx 2 valved, 1-flowered, ventricose: corol 2-valved, very short: stigmas pencil-form. 4. 10—*effusum*, (millet) panicked, awnless. e.

AGROSTIS. Calyx 2-valved, 1-flowered, valves acute: corol 2-valved: stigmas longitudinally hispid or plumose, (florets spreading) 4. 10—*vulgaris* (red top) panicle spreading, glume calyx twice as long as corol. 2/ —*alba*, panicle lax, culm creeping—*stricta*, panicle long, stiff and strait. 2/.

SACCHARUM. Calyx involucre with long wool, 2-valved: corol 1 or 2-valved: stigmas 1 to 3. 4. 10—*officinarum* (sugar cane) flowers panicked, leaves flat. 2/.

MUHLENBERGIA. Calyx 1-valved, very minute: corol 2-valved, base hairy, outer valve awned at the top: seed 1, oblong acuminate (Flowers panicked) 4. 10—*diffusa*, glabrous: culm weak: panicle slender, compact. 2/ —*erecta*, pubescent; culm erect and firm: panicle lax.

LEERSIA. Calyx 0: corol 2-valved, closed; valves compressed, boatform, awnless. 4. 10—*ory-*

zoides (cut grass) spikelets spreading. 2. — *virginica*, spikelets close-pressed—*lenticularis*, spikelets imbricate, glumes ciliate.

TRICHODIUM. Calyx 2-valved, valves nearly equal, acute; keel with small spines: corol 1-valved, awnless, shorter than calyx: stigmas nearly sessile, sub-hispid. Panicked. 4. 10—*laxiflorum*, (thin grass) culm erect, branched, slender. 2.

PHLEUM. Calyx indurated, 2-valved, sessile; linear, truncate, bicuspidate: corol inclosed. 4. 10. — *pratense* (timothy grass) spike long, cylindric. 2. 2d. Glumes 2 or 3-flowered; panicked.

AIRA. Calyx glossy, 2-valved, 2-flowered: corol 2-valved, awnless, or awned at the base. 4. 10. — *cespitosa*, (hair grass) corol with a strait awn—*flexuosa*, awned, geniculate—*obtusata*, awnless.

3d. Glumes 2-flowered or more; close-spiked.

ELYMUS. General calyx involucre-like, 4-leaved, 2-flowered; partial calyx lateral, 2-valved, aggregate-ternate, many-flowered. (In a spike.) 4. 10—*virginicus* (lime-grass) involucre striate, long-awned. 2.

ELEUSINE. Calyx awnless, many-flowered: inflorescence an unilateral digitate spike: corol 2-valved, awnless. 4. 10—*indica*, (dog-tail grass) spikelets about 6-flowered.

LIMNETIS, or DACTYLIS. Calyx 2-valved, one valve smaller: corol 2-valved, awnless, compressed, carinate: style long. Panicles strait; spikes lateral-flowered, florets imbricate. 4. 10—*glomerata*, (orchard grass) a panicle condensed into a spike-form head. 2.

SECALE. Calyx 2-valved, 2 or many-flowered, opposite, solitary: glumes linear-lanceolate, smooth or channelled both sides. 4. 10—*cereale*, (rye) glumes rough-ciliate.

TRITICUM. Calyx 2-valved, about 3-flowered,

alternate ; florets obtusish and pointed : glumes beardless or interruptedly bearded : spikelets shortish. 4. 10—*hybernum*, (winter wheat) nearly awnless. ♂ —*aestivum*, (summer wheat) long-awned. ☹.

LOLIUM. Calyx 1-leaved, permanent, many-flowered : florets in a 2-rowed simple spike. 4. 10—*perenne*, (darnel-grass) spike awnless, ♀ —*temulentum*, spike awned. ☹.

ATHEROPOGON. Calyx 1-valved, 2-flowered : corol 2-valved, awned. 4. 10—*apludoides* (hairy beard-grass) leaves very narrow. ♀.

4th. *Glumes many-flowered ; in panicles.*

UNIOLA. Spikelets flat, 2-edged, ovate : calyx many-glumed : corol 2-valved, awnless, inner one smallest : stamens 1 to 3 : appendages 2, somewhat 2-horned. 4. 10—*spicata* (sea rush-grass) panicle spike-form, pointing one way. ♀ —*latifolia*, panicle lax. ♀. Monandrous.

BRIZA. Calyx 2-valved, many-flowered : spikelets of the panicle 2-ranked : valvelets inflated heart-form, obtuse ; inner one minute. 4. 10—*canadensis* (quake-grass) culm erect, panicle lax.

SORGHUM. Polygamous. Florets in pairs ; one perfect and sessile, the other staminate or neutral and pedicelled. 4. 10—*saccharatum*, (broom-corn) panicle somewhat whorled, spreading : seed oval. ☹.

POA. Calyx 2-valved, many-flowered : corol ovate ; valves 2-coloured, acutish, scarious at their margins : spikelets of the panicle ovate, awnless. 4. 10—*pratensis*, (meadow-grass) panicle diffused, spikelets 4-flowered. ♀ —*trivialis*, panicle diffused, spikelets 3-flowered. ♀ —*compressa*, (blue-grass) culm compressed, panicle pointing one way. ♀ —*repens*, culm creeping, spikelets 6 to 10-flowered. ☹.

FESTUCA. Calyx 2-valved, many-flowered; spikelets oblong, teretish, or diverging; awnless or furnished with a terminal awn. 4. 10—*pratensis* (fescue-grass) panicle pointing one-way, spikelets linear. 2—*lucida*, panicle nodding, spikelets ovate-lanceolate. 2—*fluitans*, panicle erect. 2.

BROMUS Calyx 2-valved; spikelets oblong, terete-2 ranked; valves awned below the tip. 4. 10.—*secalinus*, (ches-) awnless, panicle spreading—*mollis*, (broom-grass) florets imbricate, panicle compact. ♂.

AVENA. Calyx 2-valved, many-flowered; valves with a twisted awn on the back: glumes membranaceous and somewhat follicle-like. 4. 10—*sativa*, (oats) panicle, calyx 2-seeded. ☉.

ARUNDO. Calyx 2-valved, one or many-flowered, glabrous: florets heaped together: corol surrounded with wool. 4. 10—*phragmites*, (reed-grass) calyx 5-flowered. 2.

B. Flowers polygamous.

1st. Glume 1-flowered.

ANDROPOGON. Calyx glume 2-valved: corol 3-valved: flowers in pairs; 1 sessile perfect, the other peduncled, staminate, awnless—rarely neutral. 4. 10—*nutans*, (beard-grass) panicle compact, nodding, calyx bristly.

2d. Glume 2-flowered, or more. Panicle.

HOLEUS. Calyx glume 2-valved, opaque, nervous: corol smaller than calyx, 2-valved, the outer one-awned under the tip: nectary linear 2-parted: stigma subsessile. 4. 10—*lanatus*, (soft-grass) glumes hairy. 2—*odoratus*, glumes 3-flowered, pointed, awnless. 2.

* Alsine.

CLASS III. ORDER III.

17

ORDER III. TRIGYNIA.

A. Corol 5-petalled.

HOLOSTEUM. Calyx 5-leaved: capsule 1-celled, nearly cylindric, bursting at the top. 22. 82—*succulentum*, (succulent chickweed) leaves oval, fleshy:

B. Corol 3-petalled.

LECHEA. Calyx 3-leaved: capsule 3-celled, 3-valved, with 3 inner valvalets: seed 1. 22. 81—*major*, (pinweed) flowers dull, scattered—*minor*, flowers dull, panicked.

C. Corol none.

MOLLUGO. Calyx 5-leaved, inferior: capsule 3-celled, 3-valved, many-seeded. 22. 82—*verticillata*, (carpet-weed) leaves whorled, stem prostrate.

☉. w.

QUERIA. Calyx 5 leaved, inferior: capsule 1-celled: seed 1. 22. 82—*canadensis*, stem forked, leaves lance-ovate. 24.

PROSERPINACA. Calyx 3-parted superior: nut 3-sided, 3-celled, crowned by the calyx. 15. 22—*palustris*, (mermaid weed) leaves lanceolate, serrate above, and divided below, water. 24. g.

* *Zanthoxylon*, Alsine.

CLASS IV. TETRANDRIA.

ORDER I. MONOGYNIA.

A. Flowers 1 petalled, inferior, calyx 4-cleft.

CALLICARPA. Corol 4-cleft, tubular: berry 4-seeded. 43. 38—*Americana*, (Bermuda mulberry) cymes sessile, leaves oval serrate. 42. r.

PLANTAGO. Corol 4-cleft reflexed: capsule 2-celled, opening transversely: stamens very long. 54. 31—*major*, (plantain) leaves broad ovate 24. w. —*lanceolata*, (ribwort) leaves lanceolate. 24.

LYCIUM. Corol tubular, having the throat closed by the beards of the filaments: berry 2-celled,

many-seeded. 28. 41—*carolinianum*, (box thorn) leaves oblong-spatulate, sharp. $\frac{1}{2}$. p.

FRASERA. Corol 4 cleft, spreading; segments with bearded glands in the middle: capsule compressed, sub-margined, 2-valved: seeds few, imbricated—*walteri*, (pyramid flower) leaves oblong-oval, opposite and whorled. y.

BARTONIA, or CENTAURELLA. Corol 4-cleft, bell-form, permanent: capsule 1-celled, 2-valved, many-seeded.—*Paniculata*, (screw-stem) stem square. w.

HOUSTONIA. Corol salver-form: capsule 2-celled, 2-valved, 2-seeded. 47. 57—*coerulea*, (Venus' pride) root-leaves ovate. $\frac{1}{2}$. b.

B. *Flowers 1-petalled, superior.*

MITCHELLA. Calyx 4-toothed: corols 2 on each germ, tubular: berry double, 4-seeded: stigmas 4. 48. 57—*repens*, (partridge-berry) stem decumbent, leaves roundish. $\frac{1}{2}$. w.

SANGUISORBA. Calyx inferior, 2-leaved: corol superior, 4 parted: capsule between the calyx and corol. 54. 92—*medica*, (burnet saxifrage) stamens longer than corol, spike cylindric. y.

CEPHALANTHUS. Inflorescence in a head: general calyx none: proper calyx superior: corol funnel-form: receptacle globular, hairy: capsules 2 to 4-partible: seed solitary, oblong. 48. 57—*occidentalis*, (button bush) flowers in heads. $\frac{1}{2}$. w.

LINNAEA. Calyx double; that of the fruit 2-leaved, that of the flower 5-parted: corol bell-form: stamens somewhat didynamous: berry 3-celled, dry. 48. 58—*borealis*, stem creeping, flowers in twins. $\frac{1}{2}$. w. r.

RUBIA. Calyx 4-toothed: corol 4 or 5-cleft, bell-form: berries 2, one-seeded, (stamens 4-5; leaves stellate.) 47. 57—*tinctoria*, (madder) leaves whorled, stem square, prickly. $\frac{1}{2}$. y.

GALIUM. Calyx 4-toothed: corol flat: fruit dry: seeds 2, roundish: (leaves stellate.) 47. 57—*tinctorium*, (bedstraw) stem whorls in sixes, branch whorls in fours. 2. w.—*apharine*, fruit prickly, whorls in eights. ☉. w.—*verum*, fruit smooth, whorls in eights. 2. y.—*brachiatum*, whorls in fours. 2. w.—*asprellum*, whorls all in sixes, stem decumbent. 2. w.

DIPSACUS. Common calyx many-leaved; proper calyx 1-leaved superior: receptacle chaffy: seed 1, crowned: flowers aggregate. 48. 56—*fullo-num*, (teasel) chaff hooked. ♂. e.—*sylvesiris*, chaff strait. ♂. b.

C. *Flowers 4-petalled, inferior:*

AMMANIA. Calyx tubular, plaited, 8-toothed: petals 4, or none, on the calyx: capsule 4-celled—(sometimes 2-celled and 8-stamened.) 17. 91—*humilis*, (tooth cup) stem procumbent. ☉. r.

CORNUS. Calyx 4-toothed: drupe with a 2-celled nut. Some species have a 4-leaved involucre. 45. 58—*florida*, (dogwood) involucre large, petal-like. 2. y. w.—*paniculata*, cymes paniced. 2. w.—*alba*, cymes flat. 2. w.—*circinata*, leaves orbicular, acuminate—branches warty.

LUDWIGIA. Calyx 4-parted, the divisions long, permanent: capsule 4-cornered, 4-celled, perforated at the top, many-seeded—(petals sometimes wanting.) 17. 88—*alternifolia*, (seed-box) glubrous. ☉. y.—*filosa*, rough-haired. 2. y.

E. *Flowers not complete.*

POTHOS. General calyx a spathe: perianth 0: spadix simple, covered with flowers: petals 4, wedge-form: berries globose, 2-seeded, inclosed in a spongy receptacle. 2. 7—*foetida*, (skunk-cabbage) stemless, leaves oval 2. p.

ISNARDIA. Calyx bell-form, 4-parted, superior:

corol o : capsule 4-celled, 4-cornered, many-seeded, surrounded by the calyx. 17. 91—*palustris*, (water-purslane) creeping, leaves ovate. ☺. g.

RIVINA. Corol 4-petalled, inferior, permanent : calyx o : berry 1-seeded : seed lenticular, rough, (stamens 4, 8 or 12) 12. 29—*laevis*, racemes simple, leaves ovate. r.

* *Convallaria*, Cardamine, Prinos.

ORDER II. DIGYNIA.

HAMAMELIS. Involucre 3-leaved : perianth 4-leaved : petals 4, very long, linear : nut 2-celled, 2-horned. 54. 78—*virginica*, (witch hazel) leaves undulate-toothed, ob-ovate. Flowers in autumn, and perfects its seed the following spring. h. y.

* *Cuscuta*, *Swertia*.

ORDER IV. TETRAGYNIA.

ILEX. Calyx 5-toothed ; corol 4-parted, wheel-form ; style o ; berry 4-celled, cells 1-seeded. 43. 95—*opaca*, (holly) leaves evergreen, repand, spinous. h. g.—*canadensis*, leaves deciduous, ovate, mostly entire. h. w. y.

SAGINA. Calyx 4-leaved ; petals 4 ; capsule 4-celled, 4-valved, many-seeded. 22. 82—*procumbens* (pearl-wort) stem procumbent. h. w.

POTAMOGETON. Calyx o : petals 4 ; style o : seeds 4. 15. 6.—*natans*, (pond-weed) upper leaves with rounded bases—*fluitans*, upper leaves lance-oval.

CLASS V. PENTANDRIA.

ORDER I. MONOGYNIA.

A. *Flowers 1-petalled, inferior ; having 4 naked seeds.* Rough-leaved plants.

PULMONARIA. Calyx prismatic 5-angled : corol funnel-form, with an open throat : seeds roundish, obtuse. 41. 42—*virginica*, (lungwort) leaves ovate, panicle terminal.

LITHOSPERMUM. Calyx 5-parted : corol funnel-form, with an open throat : seeds ovate, pointed, stoney : stamens and pistils inclosed. 41. 42—*officinale*, (stone-seed, or gromwell) seeds smooth. 24. y.—*arvense*, seeds wrinkled.

CYNOGLOSSUM. Calyx 5-parted : corol funnel-form, vaulted, throat closed : seeds depressed, affixed laterally to the style within. 41. 42—*officinale*, (hound-tongue) leaves sessile. 24. p.—*amplexicaule*, upper leaves clasping. 24. w. b.

ANCHUSA. Calyx 5-parted : corol funnel-form, vaulted, throat closed : seeds marked at the base, and their surface generally veined. 41. 42—*officinalis*, (bugloss) spikes one-sided, leaves lanceolate. 24. e.

MYOSOTIS. Calyx half-5-cleft : corol salver-form, curved, 5-cleft, vaulted, throat closed, the lobes slightly emarginate : seeds smooth or echinate. 41. 42—*scorpioides* (scorpion grass) racem-
ed, leaves lance-oval. 24. b.

HELIOTROPIMUM. Calyx tubular, 5-toothed : corol salver-form, 5-cleft, with teeth or folds between the divisions ; throat open. (Spikes recurved, involute.) 41. 42.—*indicum* (turnsole) leaves heart-ovate, acute. 24. b. e.

ONOSMODIUM. Calyx deeply 5-parted : corol oblong, somewhat bell-form, with the acute divisions converging, the upper part of the corol being ventricose and half-5-cleft, throat open : anthers sessile enclosed : styles exsert. (Spikes revolute.) 41. 42—*hispidum*, (false gromwell) leaves oval-lanceolate, papillose-punctate.

BORAGO. Corol wheel-form, the throat closed with rays. 41. 42—*officinalis*, (borage) leaves alternate. e.

SYMPHITUM. Limb or upper part of corol, tubular-swelling, the throat closed with subulate rays.

41. 42—*officinalis*, (comfrey) leaves decurrent. 24. w. e.

ECHIUUM. Corol mostly irregular, throat naked, large: stigma cleft. 41. 42—*vulgaris*, (viper's bugloss) stem bristly, spikes lateral. b.

B. *Flowers 1 petalled, inferior; seeds covered with a 1-celled capsule: calyx 5-parted or 5-toothed.*

HYDROPHYLLUM. Corol bell-form, 5-cleft, with 5 longitudinal honey-bearing grooves inside: capsule globose, 2-valved: stigma 2-cleft. 41. 42—*virginicum*, (water-leaf) leaves pinnate-pinnatifid. 24. w.—*canadense*, leaves lobed. 24. w.

ANAGALLIS. Corol wheel-form, deeply 5-parted: capsule opening transversely: stamens hairy. 20. 34—*arvensis*, (scarlet pimpernel) stem square, procumbent; leaves dotted. ☉. r.

LYSIMACHIA. Corol wheel-form: capsule globular, 10-valved, mucronate: stigma obtuse. (In some species the filaments are united at the base.) 20. 34—*ciliata*, (loosestrife) petioles ciliated. 24. y. *thyrsifolia*, racemes lateral. 24.—*hybrida*, leaves opposite, lanceolate, acute at the base—*stricta*, stem squarer, racemes terminal 24. y.—*quadrifolia*, leaves whorled or in pairs nearly opposite, with axillary 1-flowered peduncles. 24. y. Near the outlet of the Beaver Pond (New-Haven) I found this species in abundance, with 2, 3, 4, 5 and 6 leaves together; though Linneus allows but 4.

DODECANTHEON. Corol wheel-form, reflexed capsule oblong: stamens on the inside of the tube: stigma obtuse. 20. 34.—*meadia*, (false-cowslip) leaves repand-toothed, umbels many-flowered. 24. p.

MENYANTHES. Corol hairy: capsule ovate: stigma 2-cleft. 21. 34—*trifoliata*, (buck bean) leaves 3 on a petiole. 24. w.

PRIMULA. Umbellets involucred: tube of corol

cylindric, throat open, divisions of corol emarginate : capsule 1-celled with a 10-cleft mouth : stigma globular. 21. 34—*vulgaris*, (primrose) scape 1-flowered. 24.—*veris*, (cowslip) scape many-flowered. 24.

HOTTONIA. Corol salver-form : stamens on the tube of the corol : stigma globular. 21. 34—*palustris*, (feather-leaf) flowers whorled. 24.

—— Differ from the above in having 2-celled capsules.

SPIGELIA. Corol funnel-form : twin capsules 4-valved, 2-celled : stigma simple : seeds many. 47. 46—*marilandica*, (pink-root) leaves lance-ovate, sessile, opposite. 24. p.

CONVOLVULUS. Corol funnel-form, plaited : stigma 2-cleft or double : cells of the capsule 2 or 3 ; each 2-seeded. 29. 43—*sepium*, (morning glory) hind lobes of the arrow-form leaf truncate. 24. p. w.—*stans*, stem erect. 24. w.—*batatas*, (sweet potatoe) leaves heart-halbert-form. 24. p. w.—*tricolor*, leaves lance-ovate. ☉.

DATURA. Calyx tubular, angled, caducous : corol funnel-form, plaited : capsule 4-valved, 4-celled, smooth or thorny. 28. 41—*stramonium*, (thorn-apple) pericarp spinous. ☉. w. b.

NICOTIANA. Corol funnel-form, limb plaited : stigma notched : stamens inclined : capsules 2-celled, 2-valved. 28: 41—*tabacum*, (tobacco) leaves decurrent. ☉. p.

VERBASCUM. Corol wheel-form, somewhat irregular : stamens declined, hairy : capsules 2-celled, 2-valved ; valves inflexed : many-seeded. 28. 41.—*thapsus*, (mullein) leaves decurrent. ☉. y. —*blattaria*, (cockroach mullein) leaves clasping. ☉. y.

CHIRONIA. Calyx somewhat bell-form : corol

salver-form or wheel-form : anthers becoming spiral ; filaments on the tube of the corol : pistil declined : pericarp 4-celled. 20. 40—*angularis*, (centaury) leaves clasping, stem acute-angled. ☉. p. —*venosa*, stem somewhat decumbent, leaves lanceolate. 24. w.

ATROPA. Corol bell-form : stamens distant : berry globular, 2-celled, sitting on the calyx. 28. 41—*belladonna*, (deadly nightshade) leaves ovate. 24. e.

PHACELIA. Corol somewhat bell-form, 5-cleft : stamens exsert : capsule 2-celled, 2-valved, 4-seeded. 41. 42—*fimbriata*, corol fringed. ☉. b. —*parviflora*, corol entire. ☉. b.

HYOSCYAMUS. Corol funnel-form, obtuse, irregular : stamens inclined : capsule 2-celled, covered with a lid. 28. 41—*niger*, (henbane) leaves clasping, pinnate. ☿. y. p. e.

——— *Differ from the above in having 3-celled capsules.*

DIAPENSIA. Calyx 5-leaved, imbricated with leaflets : corol salver-form ; limb flat, 5-cleft : stamens crowning the tube of the corol : capsule 3-valved, many-seeded. 21. 43—*lapphonica*, leaves linear. 24. w.

PHLOX. Calyx prismatic : corol salver-form, with a tube somewhat curved : filaments unequal in length : stigma 3-cleft : capsule 3-celled, 1-seeded. 20. 44—*paniculata*, (lichnedia) segments of corol rounded. 24. p. —*subulata*, lobes of corol notched. 24. w. p.

POLEMONIUM. Calyx-half 5-cleft : corol wheel-form, 5-parted : stamens on 5 valves closing the bottom of the tube : stigma 3-cleft. 29. 44—*repens*, (Greek-valerian) leaves pinnate, root creeping. 24. b.

IPOMOEA. Corol funnel or bell-form, with 5 plaits : stigma globe-headed, papillose : capsule 2 or 3-celled, many-seeded. 29. 43—*quamoclit*, (cypress vine) leaves pinnatifid. ☺. p. r.

PYXIDANTHERA. Divisions of calyx lying over each other, chaffy-membranaceous : corol bell-form, 5-parted : filaments lamellar : cells of anthers opening transversely : stigmas 3 : style thick. 21. 43—*barbulata*, (box-flower) leaves lance-wedge-form. h. w.

——— *Differ from the above in having 5-celled capsules.*

AZALEA. Corol tubular, half-5-cleft, somewhat oblique : stamens on the receptacle : stigma obtuse, usually ending with 5 short papillae. 18. 50—*viscosa*, (wild honey-suckle) tube of corol twice as long as the divisions. h. w. —*nitida*, tube of corol not twice as long as the divisions. h. w. y.

VINCA. Corol salver-form, twisted, border 5-cleft, with oblique divisions ; throat 5-angled : seed naked, oblong ; follicles 2, erect, terete, narrow. 30. 47—*minor*, (periwinkle) stem procumbent, lance-oval.

——— *Differ from the above in having 2-celled berries.*

SOLANUM. Calyx permanent : corol bell or wheel-form, 5-lobed, plaited : anthers thickened, with 2 pores at the top : berry containing many seeds. 28. 41—*tuberosum*, (potatoe) leaves interruptedly pinnate. y. w. p.—*nigrum*, (nightshade) leaves angular. ☺. w.—*dulcamara*, (bittersweet) climbing, leaves halbert-form. h. b.

PHYSALIS. Corol bell or wheel-form : anthers converging : berries contained in an inflated calyx : seeds numerous. 28. 41—*alkekengi*, (winter-cher-ry) leaves in pairs, entire, acute.

CAPSICUM. Corol wheel form : berry juiceless : anthers converging : calyx angular. 28. 41—*annuum*, (red pepper) flowers solitary. ☺. w. e.

C. *Flowers 1-petalled, half-inferior : calyx 5-cleft.*

SAMOLUS. Corol salver-form, 5-lobed with intermediate scales fencing the stamens between them : capsule 1-celled, 5-toothed ; many-seeded. 21. 34—*valerandi*, (brookweed) leaves obtuse, racemes many-flowered. ♀. w.

D. *Flowers 1-petalled superior : calyx 5-cleft.*—
(Stems woody.)

LONICERA. Corol tubular, 5-cleft, unequal : berry 2 or 4-celled : seeds many. 48. 58—*semper-virens*, (trumpet honey-suckle) tube of corol dilating upwards. ♀. r. e.—*dioica*, corols ringent, gibbous at the base. ♀. y. p.

XYLOSTEUM. Corol tubular, border 5-parted, nearly equal : berries in pairs, whose bases are connate, or single coadunate ; 2-celled. 48. 58—*ciliatum*, (fly honey-suckle) peduncle 2-flowered, berries distinct ; leaves entire, hairy. ♀. y. w.—*solonis*, two flowers seated on one germ ; berry double, not distinct : leaves oblong-ovate, villose. Found on the White Mountains, (N. H.) by Dr. D. Solon. C. H. Smith.

DIERVILLA. Calyx oblong : corol twice as long as calyx, funnel-form, border 5-cleft, spreading : capsule oblong 4-celled, many-seeded. 48. 58—*canadensis*, (yellow honey-suckle) leaves serrate. ♀. y.

(Stems not woody.)

CAMPANULA. Corol bell-form, closed at the bottom by valves bearing the stamens : stigma 3 to 5-cleft : capsule 3 to 5-celled, opening by lateral pores. 29. 52—*perfoliata*, (bell-flower) leaves clasping. ☺. b.—*rotundifolia*, root leaves-round-

ish-cordate, stem-leaves linear. $\mathcal{L}$. b.—*media*, (Canterbury bells) capsules 5-celled, flowers erect. δ .

LOBELIA. Corol irregular, often irregularly slit-
ted: anthers cohering and somewhat curved: stig-
ma simple: capsule 2 or 3-celled: 29. 52—*cardi-
nalis*, (cardinal flower) spike pointing one-way. $\mathcal{L}$.
r.—*inflata*, capsules inflated. δ . w. b.—*pallida*,
leaves toothed. $\mathcal{L}$. w. b.—*syphilitica*, leaves with
sinuses reflexed. $\mathcal{L}$. b.

MIRABILIS. Corol funnel-form, coarctate be-
low: calyx inferior: germ between the calyx and
corol: stigma globular. 54. 32—*jalapa*, (four o'-
clock) flowers peduncled, leaves glabrous. e.

TRIOSTEUM. Calyx of the length of the corol:
corol tubular, 5-lobed: berry 3-celled, 3 seeded.—
48. 58—*perfoliatum*, (feverwort) leaves connate,
flowers whorled. $\mathcal{L}$. p.

E. *Flowers 5-petalled, inferior. (Stems woody.)*

RHAMNUS. Calyx cup-form, 4 or 5-cleft; corol
consists of scales fencing in the stamens and in-
serted on the calyx (sometimes wanting); stigma
either simple, 2 or 4-cleft; berry 3 or 4-seeded;
(sometimes polygamous or dioecious.) 43. 95.—
catharticus (buck-thorn) spines terminal. $\mathfrak{h}$ —*alni-
folius*, no spines. $\mathfrak{h}$.

CEANOTHUS. Petals bay-like, vaulted, standing
in the cup-form calyx; berry, or capsule, dry, 3-
grained. 43. 95.—*americanus* (New-Jersey tea)
leaves oval, serrate, downy beneath. $\mathfrak{h}$. w.

CELASTRUS. Calyx flat: corol spreading: cap-
sule 3-angled, 3-celled, berry-like: stigma 3-cleft:
seeds calyptred. 43. 95.—*scandens* (staff tree)
climbing: fruit berry-like, red, permanent, open-
ing with 3-valves. $\mathfrak{h}$. w-y.

EUONYMUS. Calyx 5-parted, flat: corol flat:

capsule 5-angled, 5-celled, 5-valved, coloured; seeds calyptred. 43. 95—*atropurpureus* (spindle tree) peduncles many-flowered. $\frac{1}{2}$ p.—*americanus*, peduncles about 3-flowered. $\frac{1}{2}$ y.

VITIS. Calyx 5-toothed: petals cohering at the tip hood-like, withering: berry 5-seeded, globular. (Often dioecious.) 46. 72—*vulpina* (grape vine) leaves heart-form, tooth-serrate. $\frac{1}{2}$ —*riparia*, leaves short, 3-cleft, petioles hairy. $\frac{1}{2}$ —*rotundifolia*, leaves heart kidney-form. $\frac{1}{2}$.

AMPELOPSIS. Calyx 5-toothed: petals reflexed, spreading: berry 5-seeded, globular. 46. 72—*quinquefolia* (false grape) leaves in fives. $\frac{1}{2}$ g.

ITEA. Calyx 5-cleft, bearing the 4 long linear reflexed petals: stigma capitate, 2-lobed: capsule 2-valved; the valves bearing the seeds and having inflexed margins. 18. 50—*virginica*, leaves ovate, serrate; racemes terminal. $\frac{1}{2}$ w.

(Stems not woody.)

IMPATIENS. Calyx 2-leaved: corol irregular spurred: anthers cohering: capsule elastic, 5-valved. 24. 73—*nolli-me-tangere* (jewel weed) peduncles many-flowered. $\odot$. y.—*balsamina* (touch-me-not, or balsam weed) peduncles 1-flowered.

VIOLA. Calyx 5-leaved: corol irregular, with a horn behind (sometimes the horn is wanting:) anthers attached by a membranous tip: capsule 1-celled, 3-valved. 29. 80.—STEMLESS. *Cucullata*, (violet) leaves cordate, flowers inverted—*lanceolata*, leaves lanceolate—*palmata*, leaves palmate—*pedata*, leaves pedate—*primulifolia*, leaves heart-oblong, petioles membrane-winged. STEMMED — *Pubescens*, leaves cordate, hairy—*canadensis*, leaves cordate, glabrous—*tricolor*, leaves tooth-crenate. e.

CLAYTONIA. Calyx 2-valved: stigma 3-cleft: capsule 1-celled, 3-valved, 3-seeded. 13. 86—*virginica*, leaves lance-linear. $\frac{1}{2}$ w-r.

F. *Flowers 5-petalled, superior.*

RIBES. Corol and stamens inserted on the calyx: style 2-cleft: berry many-seeded. 36. 85—*rubrum* (currant) flowers flattish. h. g. e.—*nigrum*, flowers oblong. h. g. y—*floridum*, flowers cylindric. h. w. y.

HEDERA. Petals oblong: berry 5-seeded, surrounded by the calyx: style simple. 46. 58—*helix*, (European ivy) leaves 5-angled. flowers umbelled. h. e.

G. *Corols wanting.*

GLAUX. Calyx inferior, 5-lobed, coloured: capsule 1-celled, 5-valved, 5-seeded, surrounded by the calyx. 17. 91—*maritima* (sea-milkwort) grows in salt water. u.

THESIMUM. Calyx superior, half-5-cleft, bearing the stamens: nut 1-seeded, covered with the permanent calyx. 31. 24—*umbellatum* (false toad-flax) leaves oblong. u. w. g.

ORDER II. DIGYNIA.

A. *Follicles 2: corol 1-petalled, with a 5-parted border.*

PERIPLOCA. Nectary pitcher-form, putting out 5 threads, surrounding the stamens: corol wheel-form: stigma capitate, 5-cornered. 30. 47—*graeca* (follicle vine) flowers terminal. u. p.

APOCYNUM. Corol bell-form: stamens alternating with 5 filamentous pointed nectaries: stigma broad almost sessile: follicles long linear. 30. 47—*androsemifolium* (dog-bane) cymed, corol spreading. u. r.—*cannabinum*, (Indian hemp) paniced, corol erect. u. y. g.

B. *Capsules 1 or 2-celled, 2-valved: corol 1-petalled: calyx 5-cleft.*

SWERTIA. Corol wheel-form with nectariferous

pores at the base of the divisions: capsule 1-celled. 20. 46—*pusilla*. (false gentian) stem 1-flowered.

☉. b.—*corniculata*, corol 4-horned. ☉. g-y.

GENTIANA. Corol with a tubular base, without pores: capsule 1-celled, oblong: columellas 2, longitudinal. 20. 46—*saponaria*. (gentian) corols 5-cleft. 2. b.—*crinita*, corols 4-cleft, tetrandrous. b.

CUSCUTA. Corol 5-cleft: capsule 2-celled, dividing transversely at the base: seeds binate. 29. 43—*americana* (dodder) stem twining, leafless, yellow, parasitic. w.

C. Corol 5-petalled.

HEUCHERA. Calyx inferior: corol on the calyx: capsule 2-beaked, 2 celled. 13. 84—*americana* (allum-root) leaves radical, cordate. 2. r.

PANAX. Polygamous. Umbelled. Involucre many leaved: calyx 5-toothed, superior: berry heart-form, 2-seeded: calyx, in the staminate flowers, entire. 46. 59—*quinquefolia*, (ginseng) leaflets quinate 2. g.—*trifolia*, leaves ternate. 2. w.

D. Corol o. (Stem herbaceous, except *Atriplex halimus*.)

SALSOLA. Calyx 5-leaved with a capsular base: capsule 1-celled, with a spiral seed. 12. 29—*kali*, (salt-wort) thorny, flowers axillary.

CHENOPODIUM. Calyx 5-leaved, 5-angled, inferior: seed 1, lens-like, invested by the calyx. 12. 29—*album*, (pigweed) leaves deltoid, erose. ☉. g.—*rubrum*, leaves heart-deltoid, sinuate-toothed—*hybridum*, leaves cordate, tooth-angled—*botrys* (oak of Jerusalem) leaves sinuate-pinnatifid. ☉. g.—*anthelminticum* (wormseed) leaves ovate-oblong, toothed. g.

BETA. Calyx 5-leaved: seed kidney-form within the fleshy substance of the base of the calyx. 12.

29—*vulgaris* (red beet) lower leaves ovate. ♂. e.
—*cicla*, (white beet) leaves ternate. ♂. e.

ATRIPLEX. Calyx 5-leaved, inferior: style 2-parted: seed 1, depressed. Pistillate flowers are intermixed, in which the calyx is 2-leaved. 12. 29
—*patula*, (orach) stem procumbent. ☉.—*halimus*, stem erect, very branching. ♀. (Stem woody.)

ULMUS. Calyx bell-form withering, border 5-cleft: seed 1, enclosed in a flat membranaceous samara. (Stamens vary from 4 to 8.) 53. 99—*americana*, (elm) flowers pedicelled; a variety, *pendula*, has drooping branches. ♀.—*fulva*, (slippery elm) flowers sessile.

CELTIS. Polygamous. Calyx 5-parted: styles thickish, spreading: drupe 1-seeded: staminate flowers beneath the perfect ones, with 6-parted calyxes: stamens 6. 53. 99—*occidentalis* (nettle tree) leaves ovate rough.

E. Plants UMBELLIFEROUS. Flowers 5-petalled,
2-seeded.

—— With general and partial involucre.

ERYNGIUM. Involucre many-leaved: flowers in heads, all fertile: receptacle chaffy: fruit crowned, 2-partible. 45. 60—*maritimum* (sea-holly) leaves radical spinous. b.

CORIANDRUM. Corols radiate, with petals inflected, emarginate: general involucre 1-leaved; partial one halved: fruit globular. 45. 60—*sativum*, (coriander) fruit globose. w. e.

HYDROCOTYLE. Umbels simple and very small: flowers all fertile: involucre 4-leaved: petals entire: fruit half-orbicular, compressed. 45. 60—*americana*, (water-navelwort) leaves crenate, reniform. w.

SANICULA. Flowers of the disk abortive: umbels crowded into head-like heaps: seeds prickly.

45. 60—*marilandica*, (sanicle, blackroot) forked, leaves about 7-lobed. ♂. w.

HERACLEUM. General involucre caducous : flowers unequal : petals inflexed, emarginate : fruit oval, compressed, striate, margined, emarginate. 45. 60—*lanatum*, (cow parsley) leaves ternate, woolly beneath. ♀. w.

DAUCUS. Involucres pinnatifid : flowers somewhat radiate, those of the disk abortive : fruit hispid with hairs. (Umbels funnel-form.) 45. 60—*carota*, (carrot) seed bristly. ♂. w. e.

ANGELICA. Petals incurved : corols equal : calyx 5-toothed : style reflexed : fruit roundish, solid, 3-winged both sides. 45. 60—*triquinata*, (angelica) terminal leaflet sessile. ♀.

SISON. Petals lanceolate, inflected : umbels with few peduncles : fruit ovate, striate : involucres about 4-leaved. 45. 60—*canadense*, (hone-wort) umbel irregular, leaves ternate. ♀. w.

FERULA. Petals heart-form : fruit oval, flat-compressed, with 3 lines on both sides : flowers uniform, all fertile : general involucre caducous, partial ones many-leaved. (Generally tall.) 45. 60—*villosa*, (giant fennel) leaves triternate, leaflets serrate. ♀. w.

LIGUSTICUM. Involucre membranaceous : calyx 5-toothed : petals equal, involute, entire : fruit 3 or 5-ribbed each side, oblong. 45. 60—*levisticum*, (lovage) leaves manifold, upper one irregularly cut. ♀. e.

AMMI. Involucres pinnatifid : corols radiate, all perfect : fruit smooth or striate : umbels lax. 45. 60—*capillaceum*, (bishop weed) leaves capillary, many-cleft.

Sium. Involucres many-leaved : petals heart-form : fruit somewhat ovate, striate, compressed. 45. 60—*latifolium*, (water parsnip) leaves pinnate, umbels terminal. ♀. w.

SELINUM. Involucre reflexed : perianth entire : petals heart-form, equal : fruit flat-compressed, striated in the middle, oval-oblong. 45. 60—*canadense*, (milk-parsley) glabrous, leaves doubly-pinnate.

CONIUM. Partial involucre halved, somewhat 3-leaved : fruit somewhat globular, marked with 5 lines, crenate at both ends. 45. 60—*maculatum*, (poison hemlock) stem spotted, leaves tripinnate. ♂. w.

——— *With a partial, and without a general, involucre.*

CHAEROPHYLLUM. Involucre concave, reflexed, 5 to 8-leaved : petals inflexed, heart-form : fruit oblong, smooth or striated, very glabrous. 45. 60—*claytoni*, (sweet cicely) petioles 3-parted, leaflets pinnatifid-lobed. ♀.

CICUTA. Involucre 3 to 5-leaved : fruit subovate, grooved. 45. 60—*maculata*, (cowbane) stipules 2-lobed.

IMPERATORIA. Fruit roundish, compressed, gibbous in the middle, bound at the margin with a wing : petals inflex-emarginate. 45. 60—*ostruthium*, (master-wort) leaves biternate. e.

SCANDIX. Umbel simple : flowers rayed : involucre many-leaved, emarginate, or gash-pinnate : fruit somewhat hairy, terminated with a very long beak. 45. 60—*cerefolium*, (chervil) leaves tripinnate. ☉. w.

——— *Without a partial involucre, and rarely having a general one.*

SMYRNIUM. Fruit somewhat compressed, gibbous, striate : petals acuminate, keeled.—(Fruit becoming black.) 45. 60—*aurum*, (alexanders) leaves pinnate. ♀. y.—*integerrimum*, leaves ternate or biternate. ♀. y.

PASTINACA. Fruit oval, flat-compressed : petals involute, entire. 45. 60—*sativa*, (parsnip) leaves pinnate. ☉. y. e.

ANETHUM. Fruit ovatish, compressed, striate : petals involute, entire. 45. 60—*foeniculum*, (fennel) fruit ovate. ♀. e.—*graveolens*, (dill) fruit flattened. ☉. e.

APIUM. Fruit ovate, striate : involucre 1-leaved : petals equal. 45. 60—*graveolens*, (celery) stem-leaves wedge-form. ♂. w. e.

CARUM. Fruit ovate-oblong, striate : involucre about 1-leaved : petals keel-form, inflex-emarginate. 45. 60—*carui*, (caraway) stem branching. ♂. e.

* Polygonum, Scleranthus, Asclepias and Cynanchum, are usually placed here ; but they are carried to the 19th class on the authority of Doct. Smith.

ORDER III. TRIGYNIA.

A. Corols superior, 5-cleft.

VIBURNUM. Calyx 5-parted : berry or drupe 1-seeded. 43. 58—*opulus*, (snowball) leaves 3-lobed, petioles glabrous. ♀.—*acerifolium*, (maple-leaf) leaves 3-lobed, petioles hairy. ♀.—*lentago*, (sheep-berry,) leaves ovate, acuminate. ♀.

SAMBUCUS. Calyx 5-parted : berry 3-seeded. 43. 68—*canadensis*, (black-berried elder) cymes 5-parted. ♀. w.—*pubescens*, (red-berried elder) somewhat racemed. ♀. w.

B. Corols inferior, 5-petalled.

STAPHYLEA. Calyx 5-parted : capsules inflated, connate : nuts globular, having a cicatrice. 23. 95.—*trifoliata*, (bladder nut) leaves ternate. ♀. w.

RHUS. Calyx 5-parted : berry 1-seeded. 43. 94—*copallinum*, (sumach) petioles winged. ♀.—*vernix*, (poison sumach) glabrous panicle few-flowered : ♀.—*glabrum*, glabrous, panicle dense. ♀

—*typhhinum*, hairy. $\frac{1}{2}$. —*radicans*, (poison ivy) parasitic. $\frac{1}{2}$.

SAROTHTA. Calyx 5-parted : capsule 1-celled, 3-valved, coloured. 20. 82—*gentianoides*, (pine-weed) leaves minute, close-pressed. γ .

ALSINE. Calyx 5-leaved : petals equal : capsule 1-celled, 3-valved, many-seeded ; having a free central columella. 22. 82—*media*, (chick-weed) stem procumbent, leaves long-ovate. ☉. w.

* Euphorbia.

ORDER IV. TETRAGYNIA.

PARNASSIA. Calyx 5-parted : corol 5-petalled : nectaries 5 : heart-form, ciliate, with globular tips : capsule 4-valved, 2-celled : seed membranaceous-margined. 14. 64. $\frac{1}{4}$. —*caroliniana*, (parnassus-grass) leaves radical, roundish.—*asarifolia*, radical leaves kidney-form. $\frac{1}{4}$.

ORDER V. PENTAGYNIA.

ARALIA. Umbellets involucred : perianth 5-toothed, superior : petals 5 : berry crowned, 5-celled ; cells 1-seeded. 46. 59.—LEAVES TERNATE —*racemosa*, (spikenard) stem branching. $\frac{1}{4}$. w.—*nudicaulis*, (wild sarsaparella) scape with umbels. $\frac{1}{4}$. w.—LEAVES PINNATE—*hispida*, umbelled. w.—*spinosa*, paniced. $\frac{1}{2}$. w.

LINUM. Calyx 5-leaved : corol 5-petalled : capsule 5-valved, 10 celled : seeds solitary : filaments thickening at the base. 14. 73—*usitatissimum*, (flax) calyx-leaves 3-nerved. ☉. b. e.—*virginianum*, radical leaves ovate. γ .

DROSERA. Calyx 5-cleft : corol 5-petalled : capsule 1-celled, 3 to 5-valved at the top, seeds many. (Sometimes 6-styled. The leaves are always beset with glandular hairs resembling dew.) 14. 64 —*rotundifolia*, (sundew) leaves orbicular. w.—*longifolia*, leaves obovate.

STATICE. Calyx 1-based, entire, plaited, scarious, inferior : corol 5-petalled : seed 1. 48. 35—*limonium*, (marsh-rosemary) leaves obovate, mucronate. $\mathcal{U}$. b.

* *Spergula*, *Cerastium*.

ORDER XIII. POLYGYNIA.

XANTHORHIZA. Calyx 0 : petals 5 : nectaries 5, pedicelled : capsules 1-seeded, about 5 in number. 26. 61—*aprifolia*, (yellow root) leaves pinnate. $\mathcal{h}$.

CLASS VI. HEXANDRIA.

ORDER I. MONOGYNIA.

A. *Flower with a perianth and corol, without a spathe.*

TRADESCANTIA. Calyx inferior, 3-leaved : corol 3-petalled : filaments with jointed heads : capsules 3-celled. 6. 13—*virginica*, (spiderwort) leaves lanceolate, umbels terminal. $\mathcal{U}$. b. e.

BERBERRIS. Calyx inferior, 6-leaved : petals 6, with 2 glands at the claw of each : style 0 : berry 1-celled, 2-seeded. (Stigma navelled : stamens spring up on being irritated.)—*canadensis*, (barberry) branches dotted, thorns in threes, flowers racemed. $\mathcal{h}$. y.

CAULOPHYLLUM. Calyx inferior, 6-leaved : petals 6, opposite to the leaves of the calyx : drupe 1-seeded : anther 2-celled, gaping at the margin. 24. 78—*thruictroides*, (poppoose-root) leaves more than decompound. $\mathcal{U}$. y-g.

PRINOS. Calyx inferior, 6-cleft : corol wheel-form, 6-cleft ; berry 6-seeded. 43. 95—*verticillatus*, (winter-berry) leaves obovate-lanceolate, acuminate. (Berries red, permanent.) $\mathcal{h}$. w.—*glaber* ? leaves wedge-lanceolate, obtusish, glabrous. $\mathcal{h}$.

CLEOME. Calyx 4-leaved : petals 4, ascending : glands 3, one at each sinuate division of the calyx

except the lowest: stamens from 6 to 20, or more: capsule silique-like, 1-celled, 2-valved. Tetradynamous. 25. 64—*dodecandra*, (false-mustard) leaves ternate, stamens 10 to 14. ☼. w.

FLOERKEA. Calyx 3-leaved, corol 3-petalled: style 2-cleft: capsule bladder-like, 2-grained. 15. 22—*proserpinacoides*, (false-mermaid) peduncles long, solitary. ☼. g-y.

B. *Flowers with a spathe or glume, without a perianth.*

AMARYLLIS. Corol superior, 6-petalled, unequal: filaments unequal in proportion or direction, declined, inserted in the throat of the tube. 9. 17—*atamasco*, (atamask lily) corol liliaceous, stamens declined. 2. r.

HYPOXIS. Glume-like spathe 2-valved: corol superior, 6-parted, permanent: capsule elongated, narrow at the base: seed roundish. 10. 17—*erecta*, (star grass) scape 2 to 4-flowered. 2. y.

ALLIUM. Spathe many-flowered: petals inferior, ovate, spreading, sessile. (Flowers in close umbels or heads.) 9. 16—*porrum*, (leek) umbelliferous, leaves flat. 2. —*ascalonicum*, (eschallots) umbelliferous, leaves awlform. 2. e.—*triccum*, (garlic) umbelliferous, scape half-terete. 2. —*cepa*, (onion) scape bulging, longer than the terete leaves. 2. e.—*schoenoprasium*, (chives) scape and leaves terete, nearly equal. 2. e.—*sativum*, (common garlic) bulb-bearing, leaves flat. 2. e.—*canadensis*, (wild onion) bulb-bearing, leaves linear. 2.

GALANTHUS. Petals 3, concave, superior: nectaries 3, small, emarginate: stigma simple. 6. 17—*nivalis*, (snow-drop) leaves linear, keeled radical; flowers single. 2. w.

NARCISSUS. Petals 6, equal, superior: nectary bell-form, 1-leafed, including the stamens. 9. 17—

jonquilla, (jonquil) petals thrice as long as nectary. 24. e.—*pseudo-narcissus*, (daffodil) petals equalling nectaries. 24. e.

PONTEDERIA. Corol inferior, 6-cleft, 2-lipped : capsule fleshy, 3-celled, many-seeded : three stamens commonly inserted on the tip, and three on the tube, of the corol. 6. 17—*cordata*, (pickerel weed) leaves heart-oblong. 24. b.

C. *Flowers without spathe, perianth or glume.*

AGAVE. Corol erect, superior, tubular, or funnel-form : filaments erect, longer than the corol : capsule triangular, many-seeded. 10. 15—*virginica*, (agave) stemless, leaves serrate, scape simple 24. g-y.

ALETRIS. Corol superior, funnel-form, wrinkled, somewhat 6-cleft ; with the stamens inserted on the base of its divisions : capsule 3-celled, many-seeded, half-inferior when mature. 10. 16—*aurea*, (false aloe) flowers spiked, nearly sessile, corols rough, leaves lance-sword-form. 24. y-w.

HEMEROCALLIS. Corol 6-parted, tubular-funnel-form : stamens declined : stigma small, simple. 10. 17—*flava*, (day lily) segments of corol acute. 24. y-e.—*fulva*, 3 inner segments of corol obtuse. 24. y-e.

DRACAENA. Corol inferior, 6-parted ; or 6-petalled cohering at the base : filaments somewhat thickened in the middle : berry with 3 one-seeded cells. 11. 12—*borealis*, (dragoness plant) nearly stemless, leaves oval. 24. w.

CONOSTYLUS. Corol more than half-inferior, 6-cleft, permanent, with branching hairs : anthers erect : style conic, 3-parted : stigma simple : capsule opening at top, 3-celled, many-seeded.—*americana*, corol woolly within, scape corymb-panicled, leaves grass-sword-form. Discovered in New-Jersey by Pursh.

ERYTHRONIUM. Corol liliaceous, inferior, 6-petalled; petals reflexed, having 2 tubercle-form nectaries at the base of the three inner alternate petals. 11. 14—*dens-canis*, (dog-tooth violet, or adder-tongue) scape naked, leaves 2, glabrous spotted. 2. y.

ASPARAGUS. Corol superior, 6-parted, erect; the three inner petals reflexed at the apex: berry 3-celled; many-seeded. 11. 12—*officinalis*, (asparagus) leaves fascicled, bristle-form. 2. y. e.

HYACINTHUS. Corol roundish or bell-form, equal 6-cleft: 3 nectariferous pores at the top of the germ: stamens inserted in the middle of the corol: cells somewhat 2-seeded. 10. 16—*botryoides*, (grape hyacinth) corol globular, leaves channelled. 2. b.—*orientalis*, (garden hyacinth) corol funnel-form, leaves keeled. 2.

LILIUM. Corol inferior liliaceous 6-petalled; petals with a longitudinal line from the middle to the base: capsule with the valves connected by hairs crossing as in a sieve. 10. 14—*philadelphicum*, (red lily) flowers about 2 or 3, erect; leaves mostly whorled. r-y.—*canadense*, (yellow lily) flowers pendant, leaves whorled. y.—*superbum*, (superb lily) leaves whorled below; scattered above; flowers pendant revolute. y.

TULIPA. Corol 6-petalled, liliaceous: style none. 10. 14—*suaveolens*, (tulip) petals obtuse, stem hairy. e.—*gesneriana*, petals obtuse, stem glabrous. e.

CONVALLARIA. Corol inferior 6-cleft: berry 3-celled, spotted before ripening. 11. 12—*racemosa*, (clustered solomon-seal) corol wheel-form, raceme terminal, leaves alternate sub-sessile. 2. w.—*multiflora*, peduncles axillary, leaves alternate somewhat clasping. 2. w.—*bifolia*, tetrandrous; corol wheel-form, 4-cleft, leaves 2 heart-ovate. 2. w.

UVULARIA. Corol inferior, 6-petalled, with a

nectariferous hollow at the base of each petal : filaments very short. 11. 14—*perfoliata*, (bellwort) leaves perfoliate. 2. y.—*sessilifolia*, leaves sessile, lance-ovate. 2. y.

ORONTIUM. Receptacle spadix-like, cylindric, covered with florets : spathe 0 : corol 6 petalled, naked : style 0 : follicle 1-seeded. 2. 7—*aquaticum*, (floating arum) leaves lance-ovate, glabrous, tapering base. 2. y.

NARTHECIUM. Corol 6-petalled, inferior, spreading, permanent : filaments thread-form, hairy : capsule prismatic : seed appendaged at both ends. 10. 16—*americanum*, (false asphodel) having a clasping bract. 2. y.

ORNITHOGALUM. Corol 6-petalled, erect, permanent, spreading above the middle : filaments dilated or subulate at the base : capsule roundish, angled, 3-celled : seed roundish, naked. 10 16—*umbellatum*, (star of Bethlehem) peduncles longer than bracts, corymbs few-flowered. 2. w.

ACORUS. Receptacle spadix-like, cylindric, covered with florets : corol 6-petalled) naked : corol 0 : capsule 3-celled. 2. 7—*calamus*, (sweet flag) spike projecting from the side of a sword-form leaf. 2.

D. *Flowers without corol.*

JUNCUS. Glume 2-valved : calyx 6-leaved, permanent : stigmas 1 or 3-celled, 3-valved, many-seeded. 5. 13—*tenuis*, (rush grass) leaves channeled, panicle cymed. 2.—*sylvaticus*, leaves flat, hairy, panicle cymed. 2.—*campestris*, leaves flat, hairy, spikelets terminal. 2. (Very small.)—*effusus*, (bull-rush) culm round, smooth, panicle lateral. 2.

* *Trientalis*, *Polygonum*, *Laurus*, *Corydalis*.

ORDER II. DIGYNIA.

ORYZA. Calyx, glume 2-valved, 1-flowered :

corol 2-valved, adhering to the seed. 4. 10—*sativa*, (rice) culm jointed, leaves clasping, panicle terminal ☺. e.

* *Ulmus*.

ORDER III. TRIGYNIA.

MEDEOLA. Calyx 0 : corol deeply 6-parted, revolute : berry 3-seeded. 11. 12—*virginica*, (Indian cucumber) leaves in distant whorls. 24. w.

MELANTHIUM. Polygamous. Calyx 0 : corol wheel-form, 6-parted or 6-petalled, each part with 2 glands at the base : stamens from the lengthened claws of the petals : capsule ovate, somewhat 3-cleft at the apex, 3-celled : seeds membrane-winged. numerous. 10. 13—*virginicum*, (black-flower) flowers paniced, petals oval. 24. y.

VERATRUM. Polygamous. Calyx 0 : corol 6-petalled, without glands : capsules 3, many-seeded. (The staminate flowers have the rudiment of a pistil.) 10. 13—*viride*, (itchweed, or swamp hellebore) panicle decompound, leaves nerved. 24. g.—*luteum*, (white hellebore) raceme simple. 24. w.

HELONIAS. Calyx 0 : corol 6-petalled : styles distinct : capsules 3-celled, 3-horned, few-seeded. 10. 13—*latifolia*, (helonias) leaves lance-awlform, scape nearly leafless, spike short. 24. p.

SCHEUCHZERIA. Calyx 6-parted inferior : corol 0 : anthers linear : stigmas sessile, lateral : capsules 3, inflated, 1 or 2-seeded. (Sometimes the calyx is taken for the corol.) 5. 13—*palustris*, (less flowering-rush) capsules 3, 5, or 6, stigmas lateral.

ZIGADENUS. Calyx 6-leaved, spreading. with 2 glands above the narrow base of each leaf : stamens inserted in contact with the germ : capsules membranaceous, 3-celled, many-seeded. (Sometimes the calyx is taken for the corol.)—*glaberrimus*.

mus. (zigadene) leaves lance-linear, narrow, long.
 24. b

TRILLIUM Calyx 3-leaved, inferior, spreading: corol 3-petalled: berry 3-celled, many-seeded. 11. 12—*cernuum*, (wake-robin) flowers peduncled, pendant. p-w.—*rhomboideum*, flowers peduncled, erect or erectish; Var. *erectum*, *album*, and *grandiflorum* p. or w.

TRIGLOCHIN. Calyx 3-leaved, inferior; corol 3-petalled, resembling the calyx: stigmas sometimes 6: capsules dehiscent at the base. 5. 13—*maritimum*, (arrow-grass) capsule 6-celled, grooved.

RUMEX. Calyx 3-leaved: petals 3, converging; stigma many-cleft: seed 1, naked, three-sided. 12. 28—*crispa*, (dock) valves ovate entire, all bearing grains. 24.—*obtusifolia*, valves toothed, mostly bearing grains. 24.—*aquatica*, valves ovate entire, grains obsolete. 2.—*britannica*, valves entire, all bearing grains, leaves obsoletely sheathed. 24.—*acetosella*, (sorrel) leaves hastate. 24.

ORDER XIII. POLYANDRIA.

ALISMA. Calyx 3-leaved: petals 3: capsules numerous, 1-seeded. 5. 13—*plantago*, (water-plantain) leaves heart-ovate. 24. w.

CLASS VII. HEPTANDRIA.

ORDER I. MONOGYNIA.

TRIENTALIS. Calyx 7-leaved: corol 7-parted, equal, flat: berry juiceless: number of stamens variable. 20. 34—*europaica*, (chick-wintergreen) leaves lanceolate, clustered at the top. 24. w.

ÆSCULUS. Calyx inflated, 4 or 5-toothed: corol 4 or 5-petalled, inserted on the calyx, unequal, pubescent: capsule 3-celled: seeds large, chestnut-form. 23. 66—*hippocastanum*, (horse-chestnut) leaves 7-fringed, corol 5-petalled. 2. c.

CLASS VIII. ORDER I.

43

ORDER IV. TETRAGYNIA.

SAURURUS. Calyx an ament with 1-flowered scales: corol 0: germs 4: berries 4, 1-seeded. 2. 6—*cernuus*, (lizard-tail) scales of the ament petiolated, many-spiked. 24.

* *Polygonum orientale*.

CLASS VIII. OCTANDRIA.

ORDER I. MONOGYNIA.

A. Flowers superior.

OXYCOCCUS. Calyx superior 4-cleft: corol 4-parted, the divisions nearly linear, revolute: filaments converging, anthers tubular, 2-parted: berry many-seeded. 18. 51—*vulgaris*, (cranberry) stem creeping, leaves acutish, revolute. 2. r-w. —*macrocarpus*, stems rising, leaves obtuse, flat. 2.

GAURA. Calyx 4-cleft, tubular: corol 4-petalled, ascending towards the upper side: nut 1-seeded. 17. 88—*biennis*, (virginian loosestrife) leaves lanceolate, toothed. 8. r.

EPILOBIUM. Calyx 4-cleft, tubular: corol 4-petalled: capsule oblong and of great length: seeds feathered. 17. 88—*palustre*, (willow-herb) stem teretes, leaves lanceolate toothletted. 2. r. —*tetragonum*, stem grooved—4-sided, leaves serrulate. 2. r.—*angustifolium*, leaves lance-linear, entire, scattered. 2. p-b.

OENOTHERA. Calyx 4-cleft, tubular, caducous, divisions deflected: petals 4, inserted on the calyx: stigma 4-cleft: capsule 4-celled, 4-valved: seeds not feathered. 17. 88—*biennis*, (scabish, or tree-primrose) leaves lance-ovate, stamens shorter than corol. Large. 8. y.—*humila*, leaves entire, obtuse, capsule obovate. Small. 2. y.

B. Flowers inferior.

RHEXIA. Calyx pitcher-form, 4 or 5-cleft: pe-
E

tals 4, oblique, inserted on the calyx: anthers declined: capsule 4-celled, within the calyx, setose: seeds numerous, cochleate. 17. 90—*virginica*, (deer-grass) leaves 3-nerved, ciliate-serrate. 24. p.

ACER. Polygamous. Calyx 5-cleft: corol 4 or 5-petalled: samaras 2, united at the base, 1-seeded. 23. 66—*saccharinum*, (sugar-maple) leaves acuminate-toothed, 5-lobed-palmate, sinuate, base sub-cordate. 2. y.—*rubrum*, (soft-maple) leaves sub-5-lobed, gash-toothed, unequally. 2. r.

JEFFERSONIA. Calyx 5-leaved, coloured, caducous: corol 8-petalled: capsule obovate, sub-peduncled, 1-celled, dehiscent below: seeds many, oblong, arilled at the base—*diphylla* (twin-leaf) stemless, peduncle naked, 1-flowered. 24. w.

DIRCA. Calyx 0: corol tubular, limb obsolete, unequal: stamens longer than the tube of the corol: berry, or drupe, 1-seeded. (The corol may be taken for a coloured calyx.) 31. 25—*palustris*, (leatherwood) leaves oval, flowers axillary, 2 or 3 in a hairy bud-like involucre. 2. w.

TROPOEOLUM. Calyx 4 or 5-cleft, coloured, spurred: petals 4 or 5, unequal: nuts leathery, sulcate. 23. 73—*majus*, (nasturtion) leaves peltate. 24. y. e.

* *Chrysosplenium* varies into the 2d order of this class.

ORDER III. TRIGYNIA.

CARDIOSPERMUM. Calyx 4-leaved: petals 4: nectary 4-leaved, unequal: capsules 3, connate, inflated. 23. 65—*halicacabum*, (heart-seed) climbing, leaflets gash-lobed. 2. w.

POLYGONUM. Calyx inferior, 5-parted, coloured: corol 0: seed 1, angular, covered with the calyx. (Stamens and pistils vary in number—The calyx in some species might be taken for a corol.) 12. 28—*aviculare*, (knotgrass) flowers axillary,

stem procumbent. ☉. w.—*hydrophiper*, (water-pepper) stamens 6, styles 2; spike filiform, nodding, leaves spotless. ☉. w-p.—*persicaria*, (persicaria) stamens 6, styles 2, peduncles smooth, stipules ciliate. ☉. w-p.—*pensylvanicum*, styles 2, peduncles bristly, stipules not ciliate. ☉. p.—*scandens*, climbing, leaves cordate. 2. w.—*convolvulus*, twining, leaves heart-arrowform. ☉. w.—*arifolium*, stem prickly, leaves hastate. 2. r.—*virginicum*, stamens about 5, styles 2, coloured calyx 4-cleft, leaves broad-ovate ciliate. 2. w.—*maritimum*, flowers axillary, stem erect. w.—*sagittatum*, stem decumbent, prickly, leaves arrow-form. ☉. w-p.—*orientale*, stamens 7, styles 2, leaves broad-ovate, stipules hairy. Very tall. ☉. r. e.—*fagopyrum*, (buckwheat) stem smooth, branching. e.

CLASS IX. ENNEANDRIA.

ORDER I. MONOGYNIA.

LAURUS. Calyx 4 to 6-parted: corol o: nectar-ies 3, two-bristled glands, surrounding the germ: drupe 1-seeded. (Stamens vary from 3 to 14—often diœcious—The calyx may be taken for a corol.) 12. 27—*sassafras*, (sassafras) leaves 3-lobed and simple. 2. y.—*benzoin*, (spice-bush) leaves simple, acute both ends. 2. y.

ORDER III. TRIGYNIA.

RHEUM. Calyx none: corol 6-cleft, permanent; seed 1, 3-sided. 12. 28—*palmatum*, (rhubarb) leaves palmate roughish. e.

CLASS X. DECANDRIA.

ORDER I. MONOGYNIA.

A. *Flowers many-petalled, irregular.*

PODALYRIA. Calyx 5-cleft, somewhat 2-lipped; corol papilionaceous, wings of the length of the

banner: legume inflated, smooth, many-seeded. 32. 93—*tinctoria*, (wild indigo) stipules obsolete, leaves wedge-form. 24. y.—*australis*, stipules twice as long as petiole. 24. b.

CERCIS. Calyx 5-toothed, gibbous below: corol papilionaceous, banner under the wing, short: legume oblong. 33. 93—*canadensis*, (judas-tree) leaves cordate. 2. r.

CASSIA. Calyx 5-leaved: corol 5-petalled: anthers, 3 lower ones beaked: legume membranaceous. 33. 93—*chamaecrista*, (cassia) pinnate leaflets many-paired. 20. y.—*marilandica*, pinnate leaflets 8-paired. 24. y.

RHODORA. Calyx 5-toothed: corol 3-petalled: stamens declined: capsule 5-celled. 18. 50—*canadensis*, (rhodora) umbelled, leaves oval entire. 2. r.

B. Flowers 5-petalled, regular.

CLETHRA. Calyx 5-parted, permanent: style permanent: stigma 3-cleft: capsule 3-celled, 3-valved, surrounded by the calyx. (Spiked) 18. 51—*alnifolia*, (sweet pepper-bush) leaves obovate, serrate. 2. w.

PYROLA. Calyx 5 parted: anthers with 2 pores: capsule 5-celled, dehiscent at the angles. 18. 51—*umbellata*, (bitter wintergreen) peduncles umbel-corymb, leaves lance-obovate. 24. w.—*maculata*, peduncles 2-flowered, leaves lance-ovate, whitish near the midrib. 24. w.—*secunda*, raceme many-flowered turned to one side, leaves ovate serrulate. 24. w.—*rotundifolia*, (shin-leaf) raceme many-flowered, leaves round. 24. w.

Ruta. Calyx 5-parted: petals concave: receptacle surrounded by 10 nectariferous dots: capsule lobed. (Petals sometimes 4, and stamens 8.) 26. 81—*graveolens*, (rue) leaves decompose or more, petals entire.

DIONAEA. Calyx 5-leaved : stigma fringed : capsule 1-celled, gibbous : seeds numerous. (Petals sometimes 6.) 14. 73—*muscipula*, (Venus' fly-trap) radical leaves with a terminal appendage resembling a rat-trap. 24. w.

C. *Flowers 1-petalled.*

ANDROMEDA. Calyx 5-parted, inferior : corol ovate or roundish, with a 5-cleft reflexed mouth : capsule 5-celled, with partitions contrary. (Stamens sometimes 8.) 18. 51.—*paniculata*, (white-bush) corols roundish, panicled, leaves lance-obovate, stem whitish crack-barked. 2. w.—*calyculata*, (leather-leaf) racemes one-sided, leafy, leaves lance-oblong, dotted. 2. w.

RHODODENDRON. Calyx 5-parted : corol somewhat funnel-form and oblique : stamens declining : capsule 5-celled, 8. 50—*maximum* (rosebay) corol wheel-form, umbel dense terminal : leaves paler beneath. 2. r.

KALMIA. Calyx 5-parted : corol wheel-salverform, with 10 horns beneath and 10 cavities within : capsule 5-celled. 18. 50—*latifolia*, (broad-leaf laurel) corymbs terminal, leaves leathery ovate-oval. 2. w-r.—*angustifolia*, (narrow-leaf laurel) corymbs lateral, plant small. 2. r.

VACCINIUM. Calyx 5-toothed or 5-parted : corol bell or pitcher-form, 5-cleft, the divisions reflected : filaments inserted on the germ with the corol : berry 4 or 5-celled, many-seeded. (The foreign species are sometimes octandrous.) 18. 51—*corymbosum*, (whortleberry) corymbs bracted, corol sub-cylindric, leaves pubescent beneath. 2. w-p.—*resinosum*, racemes bracted, corol ovate, leaves sprinkled with resinous dots. 2. p.—*tenellum*, raceme sessile, leaves smoothish, serrulate, oboval. 2. w.

EPIGAEA. Calyx double, outer 3-leaved, inner 5-parted: corol salver-form; capsule 5-celled, many-seeded. 18. 51—*repens*, (trailing arbutus) leaves heart-ovate, stem creeping. $\frac{1}{2}$. w-r.

GUALTHERIA. Calyx inferior, double, outer 2-leaved, inner 5-cleft: corol ovate; capsule 5-celled, invested with the inner, berried calyx: nectary 10-pointed. 18. 51—*procumbens* (spicy winter-green) leaves obovate. $\frac{1}{2}$. w.

ARBUTUS. Calyx inferior, 5-parted: corol ovate, pellucid at the base: berry 5-celled. 18. 51—*uva-ursi*, (bear-berry) leaves wedge-obovate, stem procumbent. $\frac{1}{2}$. r.

LEDUM. Calyx 5-cleft: corol flat, 5-parted: capsule 5-celled, dehiscent at the base. (Stamens sometimes 5.) 18. 50—*latifolium*, (labrador-tea) leaves oblong, downy beneath, stamens about 5. $\frac{1}{2}$. w.—*buxifolium*, leaves ovate, glabrous, corymbs terminal. $\frac{1}{2}$. w.

D. Calyx none.

MONOTROPA. Corol 10-petalled: 5 outer with nectariferous hollows at the base: capsules 5-valved. (A fifth part of the carpogenation sometimes wanting.)—*lanuginosa*, (beech-drops) flowers spik-ed, woolly. w. Whole plant yellowish white.—*uniflora*, (bird's nest) stem 1-flowered. w. Whole plant ivory white.

* Geranium, Rhexia, Pertulacca, Lythrum.

ORDER II. DIGYNIA.

HYDRANGEA. Calyx 5-toothed, superior: corol 5-petalled: capsule 2-celled, 2-beaked, dehiscent between the beaks. 13. 85—*arborescens*, (hydrangea) stamens 8 to 10, leaves oblong-ovate, toothed. $\frac{1}{2}$. w. Vid. Hortensia.

SAXIFRAGA. Calyx 5-parted: corol 5-petalled: capsule 1-celled, 2-beaked; many-seeded. 13. 84—*vernalis*, (early saxifrage) leaves oval crenate.

24. w.—*pensylvanica*, leaves lance-oblong, toothed.
24. w.

TIARELLA. Calyx 5-parted: corol inserted on the calyx; 5-petalled, petals entire: capsule 1-celled: 2-valved, 1 valve larger. 13. 84—*cordifolia*, (mitre-wort) leaves cordate, petals longer than the calyx. 24. w.

MITELLA. Calyx 5-cleft: petals 5 on the calyx pinnatifid: capsule 1-celled, 2-valved, valves equal. 13. 84—*diphylla*, (false sanicle) leaves cordate, scape-like stem 2-leaved. 24. w.

CHRYSOSPLENIUM. Calyx 4 or 5-cleft, half-superior, coloured: corol o: capsule 2-beaked; 1-celled, many-seeded. (Lateral flowers 8-stamened.) 13. 86—*oppositifolium*, (golden saxifrage) leaves opposite, heart-form, crenate. 24.

DIANTHUS. Calyx inferior, cylindrical, 1-leafed, with 4 to 8 scales at the base: petals 5 with claws: capsule cylindrical, 1-celled, dehiscent at the top. 22. 82—*caryophyllus*, (pink) petals crenate-lobed, scales of calyx ovate acute. 24. r.—*plumarius*, petals many-cleft, scales of calyx ovate. 24. r.—*barbatus*, (sweet-william) flowers aggregate, scales of the calyx as long as the tube. 24.

SAPONARIA. Calyx inferior; 1-leafed, tubular, without scales: petals 5, with claws: capsule oblong, 1-celled. 22. 82—*officinalis*, (soap-wort) calyx cylindric, leaves lance-oval. ☺. r.—*vaccaria*, calyx conic, 5-angled, leaves ovate sessile. ☺. r.

SCLERANTHUS. Calyx inferior, 1-leafed: corol o: seeds 1 or 2 inclosed in the calyx: stamens inserted on the calyx. 22. 86—*annuus*, (knavell) stem spreading, calyx becoming sharp-toothed. ☺. g.

ORDER III. TRIGYNIA.

ARENARIA. Calyx inferior, spreading, 5-leaved: petals 5, entire: capsule 1-celled, many-seed-

ed. 22. 82—*lateriflora*, (sandwort) stem simple, corol longer than calyx. 24. w.—*serpyllifolia*, stem branched, corol shorter than calyx. 3. w.—*stricta*, leaves linear-subulate, calyx striate. 24. y.—*canadensis*, stem prostrate glabrous, stamens 5, capsule globose. ☉. r.

STELLARIA. Calyx 5-leaved, inferior, spreading: petals 5, deeply cleft, mostly to the claw, spreading: capsule ovate, 1-celled, many-seeded, 6-toothed at the top. 22. 82—*pubera*, (starwort) petals longer than calyx, leaves sessile ovate ciliate. 24. w.—*alsine*, petals shorter than calyx, stem decumbent. w.

SILENE. Calyx 1-leaved, inferior, conic: petals 5, with claws appendaged at the mouth: capsule imperfectly 3-celled. 22. 82—*pensylvanica*, (catch-fly) viscid-pubescent, root-leaves wedge-form, stem-leaves lanceolate. w-r.—*virginica*, panicle forked, calyx downy and cylindric when in flower. 24. w-p.—*antirrhina*, peduncles 3-cleft, petals emarginate, leaves lanceolate. ☉. w.

CUCUBALUS. Calyx inferior, 1-leaved, bell-form, inflated: petals 5 with claws, not appendaged at the mouth: capsule 3-celled. 22. 82—*stellatus*, (bladder-campion) leaves in fours, petals many-cleft. 24. w.—*behen*, stem forked, calyx bladder-like. 24. w.

HORTENSIA. Flowers deformed: florets solitary: calyx 5-toothed, minute: corol 5-petalled. The gay flowers composing the cyme, have a large coloured, permanent petal-like 5-leaved calyx, and a minute caducous 4 or 5-petalled corol: stamens 8, 10 or 11. 13. 85—*speciosa*, (changeable hydrangea) leaves broad-ovate, tapering to both ends, serrate, cymed. g-r. This is the plant so common in flower pots, improperly called hyder-india.

ORDER V. PENTAGYNIA.

SEDUM. Calyx inferior 5-cleft: petals 5: 5 nectariferous scales at the base of the germ: capsules 5. 13. 83—*telephium*, (liveforever, or orpine) leaves tooth-serrate, stem erect, corymb leafy. 24. e.—*portulacoides*, creeping, leaves round-spatulate. 2. w-e.—*anacampteros*, (stonecrop) decumbent, leaves wedge-form. e.

PENTHORUM. Calyx 5-10-cleft: petals 5 or 0: capsule 5-cuspidate, 5-celled. 13. 83—*sedoides*, (Virginia orpine) stem angular, leaves serrate, alternate. 24. y.

OXALIS. Calyx 5-leaved inferior: petals 5, cohering by the claws: capsule 5-celled, 5-cornered, dehiscent at the corners: stamens, with 5 shorter: outer ones adhering at their bases. 14. 73—*violacea*, (wood sorrel) stemless, scape 2-cleft, umbelled. 24. p.—*acetosella*, stemless, scape 1-flowered, root granular. 24. w-p.—*dillenii*, stem erect, peduncles as long as leaves. ☺. y.—*stricta*, stem erect, peduncles shorter than the leaves. y.—*corniculata*, stem decumbent, rooting. ☺. y.

SPERGULA. Calyx 5-leaved inferior: petals 5, entire: capsule ovate 1-celled, 5-valved. (Stamens often 5.) 22. 82—*arvensis*, (corn-spurry) leaves in whorls, peduncles reflexed in maturity, seed kidney-form. ☺. w.—*saginoides*, leaves opposite, linear, smooth, awnless, stem decumbent, flowers pentandrous. Persoon asks whether this is not a variety of *sagina procumbens*. Professor Ives considers all the American species of *spargula* and *sagina*, as a natural assemblage of plants, which ought to be united in one genus.

AGROSTEMMA. Calyx 1-leaved coriaceous: petals 5 with claws, border obtuse, entire: capsule 1-celled, many-seeded. 22. 82—*githago*, (cockle) bristly, petals entire. ☺. r.—*coronaria*, (rose-cockle) downy, petals emarginate. ♂. r.

CERASTIUM. Calyx 5-leaved : petals 5, 2-cleft : capsule 1-celled, dehiscent at top, tooth-like. 22. 82—*vulgatum*, (mouse-ear chickweed) petals equaling the calyx, plant hirsute. w.—*viscosum*, viscid, petals hardly longer than calyx. w.—*semidecandrum*, pentandrous, petals shorter than calyx. w.

LYCHNIS. Calyx 1-leaved, oblong, smooth : petals 5, with claws : the limb somewhat 2-cleft : capsule 1 or 5-celled. 22. 82—*chalcedonica*, (campion) flowers in fastigate heads. $\frac{1}{2}$. e.

* *Phytolacca*.

ORDER X. DECAGYNIA.

PHYTOLACCA. Calyx 0 : corol 5-petalled, calyx-like, inferior : berry 10-celled, 10-seeded. 54. 89—*decandra*, (pokeweed) leaves ovate acute, racemes simple. $\frac{1}{2}$. w. berries red.

CLASS XI. DODECANDRIA.

ORDER I. MONOGYNIA.

ASARUM. Calyx 3 or 4-cleft, superior : corol 0 : stigma 6-cleft : capsule coriaceous, crowned with the calyx. 11. 23—*canadense*, (wild ginger, or white snake-root) calyx reflexed, leaves in pairs, heart-renaliform. $\frac{1}{2}$. p. root aromatic.—*virginicum*, leaves solitary, glabrous, cordate. $\frac{1}{2}$.

HUDSONIA. Calyx tubular, 3-leaved, inferior : corol 5-petalled : stamens 15 : capsule 1-celled, 3-valved, 3-seeded. 18. 51—*ericoides*, (false-heath) leaves subulate-needle-form, hirsute, peduncles solitary, filiform. $\frac{1}{2}$. y.

PORTULACCA. Calyx 2-cleft, inferior : corol 5-petalled : capsule 1-celled, opening transversely : columella 5, filiform. 13. 86—*oleracea*, (purslane) leaves wedge-form, succulent, flowers sessile. $\odot$. y.

LYTHRUM. Calyx 6, 8, 10 or 12-toothed, inferior : petals 5 or 6 on the calyx : capsule 2-celled,

many-seeded, covered. 17. 91—*verticillatum*, (milk-willowherb) flowers whorled, leaves opposite, generally in threes, lanceolate. 24. p.

DECUMARIA. Calyx 10-leaved, superior : corol 10-petalled : capsule about 8 or 10-celled, seeds numerous. 19. 89—*sarmentosa*, (decumary) stem sarmentose, lower leaves round, upper ones lance-ovate. 1/2. w.

TALINUM. Calyx inferior, 2 or 5-leaved : corol 5-petalled : capsule 1-celled, 3-valved, many seeded, seeds arillate : columella globose—*teretifolium*, (taliny) leaves cylindric, flesh ycorymbs terminal. 24. p.

ORDER II. DIGYNIA.

AGRIMONIA. Calyx 5-toothed, invested with an outer one : petals 5 : seeds 2, in the bottom of the calyx. 35. 92—*eupatoria* (agrimony) leaves interruptedly pinnate. 24. y.

* Delphinium.

ORDER III. TRIGYNIA.

EUPHORBIA. Calyx 1-leaved, inflated : petals 4 or 5, standing on the calyx : capsule 3-grained. 38. 96—*maculata*, (spotted spurge) stem prostrate, forked, leaves hairy spotted. ☹. w.—*helioscopia*, (sun spurge) umbel 5-rayed, then 3-rayed, forked, leaves wedge form. ☹.—*polygonifolia*, (knot spurge) leaves entire, stem decumbent. 24.

RESEDA. Calyx 1-leafed, 4 to 6-parted : petals in many divisions : capsule 1-celled, dehiscent at the top : seed reniform (stamens 11 to 15 : styles 3, 5, or none) 54. 64—*odorata*, (mignonette) leaves entire and 3-lobed, calyx long as petals, styles often. 24. y. e.—*luteola*, (dyers' weed) leaves lanceolate, entire, flat, calyx 4-cleft. Found growing wild near New-Haven, by professor Ives.

ORDER XIII. POLYGYNIA.

SEMPERVIVUM. Calyx 9 to 12-parted : petals 8 to 12 : capsules 12, many-seeded. (Stamens 16 or 20.) 13. 83—*tectorum*, (house leek) leaves ciliate, offsets spreading. 24.—*arboreum*, branched woody. h.

CLASS XII. ICOSANDRIA.

ORDER I. MONOGYNIA.

PHILADELPHUS. Calyx 4-5-parted, superior : corol 5-petalled : style 4-cleft : capsule 4-5-celled, many-seeded ; seed arilled. 19. 89—*coronarius* (false syringa, or mock orange) leaves slightly toothed. h. w.

PRUNUS. Calyx 5-cleft, inferior : corol 5-petalled : nut of the drupe smooth with prominent seams at the sutures. 36. 92—*borealis*, fruit ovate, umbels sub-corymbed, sessile, few-flowered, leaves acuminate, petioles with 2 glands. h. w.—*virginiana*, (wild cherry) flowers in erect compact racemes, leaves doubly-toothed, petioles with about 4 glands. h. w.—*serotina* (choak-cherry) racemes lax, serratures simple, lower ones sub-glandular. h. w.—*cerasus* (common cherry) umbels sub-sessile, leaves lance-ovate. h. w. e.—*domestica*, (common plum) peduncles sub-solitary, leaves lance-ovate. e.

CACTUS. Calyx superior, 1-leafed, imbricate : petals numerous, in many series, the inner ones larger ; stigma many-cleft : berry 1-celled, many-seeded, navelled. 13. 85—*opuntia*, (prickly pear) nearly stemless, leaves very thick, succulent and proliferous. y.

MYRTUS. Calyx superior, 5-cleft : petals 5 : berry 2 or 3-celled, many-seeded. 19. 89—*communis*, (myrtle) flowers solitary. h. w.

AMYGDALUS. Calyx 5-cleft, inferior : petals 5 : drupe with a nut perforated with pores : flowers

sessile. 36. 92—*persica*, (peach) leaves serrate, flowers solitary, sessile. $\frac{1}{2}$. r. e. A variety, *nectarina*, has glabrous fruit.

* *Lythrum*.

FROM ORDER II. DYGINIA, TO ORDER V. PENTAGYNIA.

CRATAEGUS. Calyx superior, 5-cleft: petals 5: styles generally 3 or 5: berry mealy: seeds 2 to 5, bony. 36. 92.—*coccinea*, (thorn bush) leaves heart-ovate, petioles and calyx glandular. $\frac{1}{2}$. w.—*cordata*, leaves heart-ovate, petioles and calyx without glands. $\frac{1}{2}$. w.—*crus-galli*, leaves wedge-obovate, styles 2. $\frac{1}{2}$. w.—*oxyacantha*, (quick-set) leaves obtuse, 3-cleft, styles 3. $\frac{1}{2}$.

SORBUS. Calyx 5-cleft, superior: petals 5: styles 2 to 3: berry pomaceous: seeds 2 or 3 cartilaginous. 36. 92.—*americana*, (mountain ash, or service tree) corymbed, leaves pinnate, leaflets serrate. $\frac{1}{2}$.

ARONIA. Calyx 5-toothed: petals 5: fruit pomaceous: berry 5 or 10-celled; cells 1 or 2-seeded; seeds cartilaginous. 36. 92.—*botryaphium*, (shad-flower) leaves oblong-oval cuspidate, flowers racemed, petals linear. $\frac{1}{2}$. w.—*arbutifolia*, (choak-berry) corymbed; leaves obovate crenate-serrate, midrib glandular above. $\frac{1}{2}$. w.—*ovalis*, leaves round-oval, flowers racemed, petals obovate, ciliate. $\frac{1}{2}$. These species were formerly arranged under the genus *pyrus*.

MESPILUS. Calyx 5-cleft, superior, divisions generally foliaceous: corol 5-petalled: styles 2-5: drupe or berry with from 2 to 5 bony seeds.—36. 92.—*pyracantha*, (medlar, or evergreen thorn) thorny, leaves lance-ovate, crenate, calyx obtuse. $\frac{1}{2}$. —*germanica*, (common medlar) without thorns. $\frac{1}{2}$.

ORDER V. PENTAGYNIA.

PYRUS. Calyx 5-cleft, superior : corol 5-petalled : pome 5-celled, many-seeded. 36. 92—*communis*, (pear) leaves ovate, flowers corymbed. $\frac{1}{2}$. w. e. —*malus*, (apple) leaves ovate, flowers in sessile umbels. $\frac{1}{2}$. r-w. e.—*cydonia*, (quince) leaves entire, flowers solitary. $\frac{1}{2}$. w. e.—*coronaria*, (crab-tree) leaves cordate, angular, flowers corymbed. $\frac{1}{2}$. r-w.

SPIRÆA. Calyx 5-cleft, inferior, spreading : corol 5-petalled : capsule 2-valved within, many-seeded. 36. 92—*alba*, (steeple bush, or hand-hack) leaves wedge-lanceolate, serrate, glabrous, panicled. $\frac{1}{2}$. w.—*tomentosa*, leaves lanceolate, downy, racemes compound. $\frac{1}{2}$. p.—*trifoliata*, (Bowman's root, or Indian physic) leaves ternate, flowers subpanicled. $\frac{1}{4}$. w.

MESEMBRYANTHEMUM. Calyx superior, 5-cleft : petals numerous, linear, cohering at the base : capsule fleshy, many-seeded, turbinate. 13. 87—*cristallinum*, (ice-plant) very branching, leaves ovate with ice-like papillæ. w. e.

ORDER XIII. POLYGYNIA.

ROSA. Calyx urn-form, 5-cleft, fleshy, contracted towards the top : petals 5 : seeds numerous, bristly, fixed to the sides of the calyx within. 35. 92. —*caroliniana*, (swamp-rose) germ globular hispid, peduncles somewhat corymbed and hispid, stem with stipular prickles. $\frac{1}{2}$. r.—*rubiginosa*, (sweet-brier) germs ovate, peduncle and germ hispid, petioles and stems with recurved prickles. $\frac{1}{2}$. r.—*lucida*, (wild rose) germ globular-depressed, somewhat bristly, leaves oblong-oval, glabrous, flowers mostly in pairs, $\frac{1}{2}$. r.—*parviflora*, germ globular-depressed, bristly, prickles on the stipule strait, leaves oval, flowers in pairs. $\frac{1}{2}$. r.

RUBUS. Calyx 5-cleft, corol 5-petalled : pistils

numerous: berry composed of many juices, 1-seeded, acines. 35. 92—*occidentalis*, (black-raspberry) leaflets 3, downy beneath, stem prickly, petioles cylindric. $\frac{1}{2}$. w.—*odoratus*, (flowering raspberry) leaves simple, palmate. $\frac{1}{2}$. r.—*procumbens*, (creeping blackberry) procumbent, peduncles and petioles hook-prickly. $\frac{1}{2}$. w.—*strigosus*, (red-raspberry) without prickles, strongly bristled, leaflets in threes or fives, pinnate, white down beneath. $\frac{1}{2}$. w.—*villosus*, (high blackberry) leaflets 5 or 3, oval-pointed, sharp-serrate, downy both sides, stem and petioles prickly. $\frac{1}{2}$. w.—*saxatilis*, (brier herb) stem herbaceous, creeping, leaflets 3, nearly glabrous. $\frac{1}{4}$. w.—*ideus*, (garden raspberry) leaflets 5 and 3, stem prickly, petioles channelled. $\frac{1}{2}$. w-e.

DALIBARDA. Calyx 5-cleft: corol 5-petalled: styles long, caduceous, 5 to 8; berry composed of dry granulations. 35. 92—*repens*, (dalibard) leaves cordate, stoloniferous. $\frac{1}{4}$. w.—*fragarioides*, leaves ternate, crenate-lobed, ciliate. $\frac{1}{4}$. y.

FRAGARIA. Calyx 10-cleft, 5 alternate divisions smaller: corol 5-petalled: receptacle ovate, berry-like, caduceous. 35. 92—*virginiana*, (American strawberry) calyx spread, down on petioles erect, on peduncles close. $\frac{1}{4}$. w.—*vesca*, (garden strawberry) calyx reflexed, down on petioles much spread. $\frac{1}{4}$. w.

POTENTILLA. Calyx 10-cleft, 5 alternate divisions smaller: corol 5-petalled: seeds awnless, roundish, rugose, fixed to a dry small receptacle. 35. 92—*humila*, (five-finger) nearly stemless, hairy, erect, peduncles 1-flowered shorter than petioles. $\frac{1}{4}$. y.—*reptans*, stem creeping, leaflets obovate serrate, peduncles 1-flowered. $\frac{1}{4}$. y.—*simplex*, stem erect, simple, hairy, leaves oblong-oval, peduncles lateral, 1-flowered, long. $\frac{1}{4}$. y.—*canadensis*, stem ascending, rough-haired. $\frac{1}{4}$. y.—*norwegica*, leaves

ternate, stem forked. 24. y.—*argentea*, (silver cinquefoil) leaflets in 5 wedge-form divisions, silvery whitish, prostrate. 24. y.

GEUM. Calyx 10-cleft, 5 alternate divisions smaller: corol 5-petalled: seeds with a bent awn: receptacle columnar, villous. 35. 92—*rivale*, (avens, or herb bennet) flowers drooping, awns bent in the middle, root-leaves lyrate. 24. p.—*virginianum*, flowers erect, awns twisted near the end, stem-leaves ternate. 24. y-w.

COMARUM. Calyx 10-cleft, 5-alternate divisions smaller: petals 5, smaller than the calyx: receptacle ovate, spongy, permanent, villous. 35. 92—*palustre*, (marsh fivefinger) leaflets in threes, fives, or sevens, serrate. 24. p.

CALYCANTHUS. Calyx 1-leafed, pitcher-shaped, scurfy, with coloured divisions: corol none: styles many, with glandular stigmas: seeds many, tailed, in the succulent calyx. 35. 92—*floridus*, (Carolina allspice) branches spreading, leaves broad-oval. 2. p. e.

CLASS XIII. POLYANDRIA.

ORDER I. MONOGYNIA.

CHELIDONIUM. Calyx 2-leaved: corol 4-petalled: silique-like capsule 1-celled, linear: seeds crested, many. 27. 62—*majus*, (celandine) leaves pinnate-lobed, umbels axillary. 24. y.

ACTAEA. Calyx 4-leaved, caduceous: corol 4-petalled: berry 1-celled: seeds half orbicular: 26. 61—*spicata*, (neclace weed, or bane-berry) racemes terminal. 24. w.

ASCYRUM. Calyx 4-leaved: 2 interior leaflets, cordate, large: corol 4-petalled: filaments disposed in 4 parcels: capsule oblong, 1-celled, 2 valved, included in the calyx leaves. 20. 68—*cruce-andree*, (St. Andrew's cross, or St. Peter's wort)

stem terete, leaves ovate, panicle dichotomous. $\frac{1}{2}$. y.—*stans*, stem ancipital, leaves oblong, flowers vary into the 3d order. y.

CISTUS. Calyx 5-leaved, 2 of them smaller, corol 5-petalled : capsule 3-valved, opening at the top. 20. 80—*canadensis*, (rock-rose) stamens prostrate on the petals, leaves alternate, lanceolate. $\frac{1}{4}$. y. Late in autumn this plant sends off curved ice crystals from near the root, of a very singular structure.

SARRACENIA. Calyx double, 3 and 5-leaved : corol 5-petalled : stigma peltate, covering the stamens : capsule 5-celled. 13. 22—*purpurea*, (side-saddle flower) leaves cup-form with bordered mouths, radical. $\frac{1}{4}$. p. Found in abundance in the lakes on the Catskill mountains near Benton's turn-pike.

CITRUS. Calyx 5-cleft : petals 5, oblong : filaments dilated at the base, and several parcels : berry 9 to 18 celled—Polyadelphous. 18. 70—*aurantium*, (orange) leaves oval with winged petioles. $\frac{1}{2}$. w. e.—*medica*, (lemon) leaves ovate with linear petioles. $\frac{1}{2}$. w. e. A variety *limon* (lemon) has the fruit roundish and small.

PAPAVER. Calyx 2-leaved, caducous : corol 4-petalled : capsule 1-celled, dehiscent by pores under the permanent stigma. 27. 62—*somniferum*, (poppy) leaves clasping. ☺. p. e.—*rheas*, leaves pinnatifid. ☺. r.

TILIA. Calyx 5-parted : corol 5-petalled : capsule 5-celled, globular, coriaceous, dehiscent at the base, one-seeded. 37. 79—*pubescens*, (bass-wood) leaves truncate at the base and oblique, somewhat downy. $\frac{1}{2}$. w.—*americana*, leaves cordate, glabrous. $\frac{1}{2}$. w.

SANGUINARIA. Calyx caducous, 2-leaved : corol about 8-petalled : capsule pod-like, ovate, 1-

celled. 27. 62—*canadensis*, (blood-root) scape 1-flowered, leaves lobed. 2. w.

PODOPHYLLUM. Calyx 3-leaved: corol about 9-petalled: berry 1-celled, crowned with the stigma. 27. 61—*peltatum*, (wild mandrake) leaves peltate, pedate. 2. w.

CAPPARIS. Calyx 4-leaved, coriaceous, caducous: petals 4: stamens long: berry coriaceous, peduncled, 1-celled, globular or pod-like—*spinosa*, (caper-bush) peduncles solitary, 1-flowered, leaves oval, alternate. w-r. e.

NYMPHAEAE. Calyx 4 to 7-leaved, larger than the petals: corol many-petalled: stigma marked with radiated lines: berry many-celled, many-seeded. 13. 22—*odorata*, (pond-lily) leaves round-cordate, leaves of calyx about as long as the petals. 2. w.—*advena*, leaves cordate, petals wedge-form, much shorter than the calyx. 2. y.

ARGEMONE. Calyx 3-leaved: corol 6-petalled: capsule semi-valved. 27. 62—*mexicana*, (horned poppy) leaves cut-pinnatifid, thorny. ☹. y.—variety, *alba*, has white flowers.

FROM ORDER II. DIGYNIA—TO ORDER V. PENTAGYNIA.

FOTHERGILLIA. Calyx truncate, entire: corol 0: filaments very long, clavate: germ bifid: capsule 2-celled or 2-horned: seed bony, solitary. 50. 99—*alnifolia*, (Fothergill's bush) leaves wedge-form, spikes terminal. h.

PAEONIA. Calyx 5-leaved: petals 5: style 0: stigmas 2 or 3: capsules pod-like, many-seeded. 26. 61—*officinalis*, (peony, erroneously called pina) leaves decompound. 2. r. In gardens the stamens usually change to petals.

HYPERICUM. Calyx 5-parted, divisions sub-ovate: corol 5-petalled: filaments often united at the

base in 3 or 5 sets : styles 2 to 5 : capsules roundish with a number of cells equal to the number of styles. 20. 68—*perforatum*, (St. John's wort) stem 2-edged, leaves obtuse having pellucid dots. 24. y.—*parviflorum*, corymbs terminal, forked, leaves sessile. 24. y.—*virginicum*, flowers 9-stamened, terminal, leaves clasping. 24. p.—*canadense*, flowers peduncled axillary solitary, leaves sessile, stem square. y.

DELPHINIUM. Calyx 0 : corol 5-petalled, unequal : nectary 2-cleft, horned behind : capsules 1 or 3, pod-like. 26. 61—*staphisagria*, (larkspur) leaves palmate obtuse-lobed, nectary 4-leaved. b. e.—*azureum*, leaves linear-many-cleft, spiked, spur curved. 24. p. e.

ACONITUM. Calyx 0 : petals 5, upper one vaulted : nectaries 2, hooded, peduncled, recurved : capsules 3 or 5, pod-like. 26. 61—*napellus*, (monk's hood) leaves 5-parted with 3-parted divisions. 24. p. e.

AQUILEGIA. Calyx 0 : petals 5 : nectaries 5 alternating with the petals and ending in horns beneath : capsules 5, distinct. 26. 61—*canadensis*, (columbine) nectary horns nearly strait. 24. r-y.—*vulgaris*, nectary horns incurved. 24. b-p.

CIMICIFUGA. Calyx 4-leaved : petals (or nectaries) 4, pitcher-form, cartilaginous : capsules 4 : seeds scaly. (Petals, pistils and calyx-leaves, variable.) 26. 62—*americana*, (bug-bane) flowers spiked, leaves decom-pound, pistils 5 or 6. 24. w.

NIGELLA. Calyx none : petals 5 : nectaries 5, three-cleft, within the corol : capsules 5, convex, 26. 61—*damascena*, (fennel flower) involucre 5-leaved, longer than the flower. ☺. b.

* Reseda, Ascyrum, Caltha, Helleborus, Anemone, Hepatica.

ORDER XIII. POLYGYNIA.

HYDRASTIS. Calyx 0 : corol 3-petalled : berry composed of 1-seeded acines. 26. 61—*canadensis*, (orange-root) leaves binate, palmate, petioled. 2. w.

CLEMATIS. Calyx 0 : petals, 3, 4, 5, or 6 : seeds compressed : styles permanent, becoming long tails. (Some species are diœcious.) 26. 61—*virginiana*, (virgin's bower) diœcious, climbing, leaves ternate with climbing petioles. 2. w.

THALICTRUM. Calyx 0 : petals 4 or 5 : filaments very long : seeds without tails, striate, terete. (Some species are diœcious.) 26. 61—*dioicum*, (meadow rue) diœcious, leaves heart-roundish with obtuse lobes, peduncles axillary, shorter than leaves. 2. r. w.—*cornuti*, diœcious, leaflets 3-cleft, panicles terminal. 2. w.—*polygamum*, hairy and slender, leaves more than decompound, panicle terminal. 2. w.

HELLEBORUS. Calyx 0 : petals 5 or many : nectaries tubular, 2-lipped : capsules 5 or 6, erectish, compressed, many-seeded. 26. 61—*trifoliatu*s, (gold thread) scape 1-flowered, leaves ternate with roundish crenate leaflets. 2. w. Root creeping and golden yellow.

CALTHA. Calyx 0 : petals 5 to 9 : capsules numerous, many seeded : nectaries 0 : (styles variable in number.) 26. 61—*palustris*, (American cowslip) leaves heart-reniform, crenate. 2. y.

ANEMONE. Calyx 0 : petals 5 to 9 : seeds numerous, naked. 26. 61—*nemorosa*, (wind-flower) stem simple, 1-flowered, 3-leaved with gashed leaflets. 2. r-w.—*thalictroides*, (rue-anemone) stem simple, stem-leaves 6 whorled petioled, root-leaves long-peduncled. 2. w.—*virginiana*, stem forked, many-flowered, petals acute. 2. g-y.

HYDROPELTIS. Calyx 6-leaved somewhat converging : corol 0 : capsules oblong, somewhat fleshy, numerous, 2-celled, 1 or 2-seeded. 26. 61—*purpurea*, (water-shield) leaves oval, peltate, floating. 24. p.

HEPATICIA. Calyx 3-leaved : petals 6 to 9 : seeds naked. 26. 61—*triloba*, (liverleaf) calyx-leaves broad-ovate obtuse, leaves 3-lobed obtuse. 24. w. r. Var.—*acuta*, calyx-leaves acute, leaves 3 to 5-lobed acute. Perhaps this variety ought to constitute a new species. The specimens found by professor Dewey, near Williams' College, are much firmer in their texture than the common kind, and differ materially in their general habit.

ANNONA. Calyx 3-leaved : corol 6-petalled : stigma obtuse : berries 1-celled, many seeded, roundish with a scaly bark. 52. 76—*glabra*, (custard apple) leaves lance-ovate, fruit conic glabrous. e.

LIRIODENDRON. Calyx 3-leaved : corol 6 or 9-petalled, liliaceous : seeds in a samara, imbricate on a strobile-like spike. 52. 75—*tulipifera*, (tulip tree, or white wood) leaves 3-lobed, with the middle lobe abrupt. 2. w.

MAGNOLIA. Calyx 3-leaved : corol 6 to 9-petalled : capsules numerous, imbricate on a strobile-like spike, 2 valved : seeds berried, pendulous. 52. 75—*glauc*, (magnolia, or beaver tree) leaves oval obtuse glaucous beneath, petals obovate. 24. w.

NELUMBium. Calyx 4 or 5-leaved : petals 5 or more : nuts half immersed in a truncate receptacle, and crowned with the permanent style. 26. 61—*luteum*, (nelumba) leaves peltate, peduncles and petioles glabrous. 24. y.

RANUNCULUS. Calyx 5-leaved : petals 5, with claws and a nectariferous pore or scale on the inside of each : seeds numerous. 26. 61—*abortivus*, (crow-foot) radical leaves cordate crenate, stem-leaves

subternate. 24. y.—*acris*, (butter-cup, or crowfoot) lower leaves 3-parted many-cleft, upper ones linear. 24. y.—*sceleratus*, (water crowfoot) lower leaves palmate, upper ones digitate. 24. y.—*pensylvanicus*, calyx reflexed, leaves ternate 3-cleft hairy beneath. ☺. y.—*fluviatilis*, leaves all under water, capillary. y-w.—*fascicularis*, leaves ternate-pinnate, root fascicled. 24. y.—*bulbosus*, calyx reflexed, peduncles grooved, root bulbous. 24. y.—*repens*, scions creeping, leaves linear entire. 24. y.

TROLLIUS. Calyx 0 : petals from 5 to 14 : capsules many, ovate, many-seeded. 26. 61.—*americanus*, (globeflower) petals about 5 obtuse, nectary shorter than the stamens. 24. y.

ATRAGENE. Calyx 2 or 4-leaved, sometimes involucred : petals numerous, outer ones largest : seeds tailed. 26. 61.—*americana*, (atragene) leaves in fours, leaflets cordate. Flowers large. 24. p.

ADONIS. Calyx 4-5-leaved : petals 5 or more, without nectariferous pores : seeds awnless. 26. 61.—*autumnalis*, (pheasant's eye) fruit sub-cylindric, petals 5 to 8. e.

CLASS XIV. DIDYNAMIA.

ORDER I. GYMNOSPERMIA.

A. *Calyxes 5-parted with the divisions or teeth nearly equal.*

TEUCRIUM. Corol deep-cleft on the upper side and without an upper lip ; lower lip 3-cleft, the middle division rounded : stamens and pistils incurved, stamens exsert. 42. 39.—*canadense*, (woodsage, or wild germander) whorls 6-leaved, leaves lance-ovate, serrate, spike crowded. 24. r.—*virginicum*, bracts shorter than peduncle, leaves ovate. 24. r.

ISANTHUS. Calyx bell-form : corol 5-parted,

tube strait, narrow; divisions ovate equal: stamens nearly equal: stigma linear, recurved. 42. 39—*coeruleus*, (blue gentian) viscid-hairy: leaves lance-ovate, tapering to both ends. ☺. b.

VERBENA. Corol funnel-form with a curved tube, border 5-cleft nearly equal: calyx 5-cleft: seeds 2 or 4: sometimes but 2 stamens or 2 barren ones.—*urticifolia*, (vervain) spikes filiform, leaves undivided ovate serrate. 2. w-p.—*hastata*, leaves hastate below and lanceolate above. 2. w-b. *angustifolia*, spikes filiform, leaves narrow lance-linear. Small. 2. r.—*aubletia*, spikes lax, leaves 3-cleft, erect. ☺. r.—*spuria*, stems many, leaves many-cleft jagged. 2. w-r.—*pinnatifida*, leaves pinnatifid.

LAVANDULA. Calyx ovate subdentate, bracts under-studded: corol resupinate: stamens in the tube. 42. 39—*spicata*, (lavander) leaves sessile lance-linear, spike leafy.

LAMIUM. Corol with the upper lip entire, vaulted; under lip 2-lobed; throat with a tooth at each margin. 42. 39—*amplexicaule*, (dead nettle) floral leaves sessile clasping gashed. ☺. p.

BRACHYSTEMUM. Corol with the upper lip sub-emarginate, lower lip obtusely 3-lobed: filaments very short nearly equal: seed oblong-cylindric. 42. 39—*virginicum*, (Virginian thyme) leaves lanceolate, flowers in small heads fastigate, stem square. 2. w.

PYCNANTHEMUM. Involucre bract-like, many-leaved, under small heads of flowers: calyx tubular, striate: corol with the upper lip sub-entire, lower lip 3-cleft: stamens nearly equal. 42. 39—*aristatum*, (wild basil, or mountain mint) bracts linear acuminate, leaves lance-oval with short petioles acuminate, plant downy. 2. w-p.—*incanum*, bracts bristle-form, leaves oval petioled acuminate,

downy. 2/. w-p. Both species have a pleasant pungent taste.

NEPETA. Calyx dry : striated : corol with a longish tube, under lip with the middle division crenate, throat with a reflected margin : stamens approximate. 42. 39—*cataria* (catmint) flowers whorled, leaves cordate tooth-serrate. 2/. w-b.

HYSSOPUS. Corol with the under lip 3-parted, its intermediate division sub-crenate : stamens strait, distant. 42. 39—*officinalis* (hyssop) leaves lanceolate obtuse entire. 2/. b. e.—*nepetoides*, leaves sub-cordate acute-toothed. 2/. y.

MENTHA. Corol nearly equal, 4-cleft : broadest division emarginate : stamens erect, distant. 42. 39—*viridis*, (spear mint) spikes interrupted, leaves sessile, bracts bristle-form. 2/. p.—*piperita*, (peppermint) spikes obtuse, leaves petioled, ovatis. 2/. b.—*borealis*, (Canada mint) leaves petioled projecting. 2/. b-w.

STACHYS. Corol with the upper lip vaulted, the lower lip reflexed at the sides, the middle division largest, emarginate : stamens reflexed towards the sides after discharging the pollen. 42. 39—*aspera* (false horehound) stem erect, with hairs pointing backward, leaves lance-cordate, spike whorled. 2/. r.

SATUREJA. Calyx tubular, striate : corol with divisions nearly equal : stamens distant. 42. 39.—*hortensis*, (summer savory) peduncles axillary 1 or 3-flowered, leaves lance-ovate. ☺. b. e.

LEONURUS. Calyx 5-angled, 5-toothed, corol with the upper lip villose, flat, entire ; lower lip 3-parted, middle division undivided. 42. 39—*cardiaca*, (motherwort) upper leaves 3-lobed, entire. 2/. w.

MARRUBRIUM. Calyx salver-form, rigid, marked with 10 lines : corol with the upper lip 2-cleft,

linear, strait. 42. 39—*vulgare*, (horehound) calyx with 10-hooked teeth, leaves toothed. 4. w.—*su-
phinum*, (nodding horehound) leaves round-cordate,
crenate, rugose-veined, teeth of the calyx 5, strait.
4. b-p.

CUNILA. Calyx cylindric, 5-toothed, throat vil-
lose, corol ringent; upper lip erect, flat emargi-
nate: 2 of the stamens usually barren. 42. 39.—*pu-
leginoides*, (penny-royal) leaves oblong 2-toothed,
flowers whorled. ☉. b.

GLECHOMA. Calyx 5-cleft: each pair of an-
thers approaching so as to exhibit the form of a
cross. 42. 39.—*hederacea*, (ground ivy) leaves kid-
ney-form crenate, stem square procumbent. 4. b.

B. Calyces 2-lipped.

MELISSA. Calyx dry, flattish above, with the
upper lip sub-fastigiate: corol with the upper lip
somewhat vaulted, 2-cleft; lower lip with the mid-
dle lobe cordate. 42. 39.—*officinalis*, (balm) whorls
half-round, bracts pedicelled. 4. w.—*nepeta*, (low
calamint) branching hairy, leaves ovate obtuse. 4.
w-r.

TRICHOSTEMA. Corol with the upper lip fal-
cate: the under lip 3-parted with the middle divi-
sion small, oblong: filaments very long-exsert. 42.
39.—*dichotoma*, (blue-curly) stem forked. ☉. b.

DRACOCEPHALUM. Corol with the throat inflat-
ed; the upper lip concave, the middle division of
the lower lip roundish, notched. 42. 40.—*denticula-
tum*, (dragon head) flowers spiked, remote, leaves
lance-obovate. 4. p.—*virginianum*, flowers spiked,
crowded, leaves lance-linear. 4. p.—*variegatum*,
spikes 4-cornered short, leaves oblong. p.

SCUTELLARIA. Calyx with an entire mouth,
which is closed with a lid after the corol falls out:
tube of corol bent. 42. 39.—*lateriflora*, (skull-cap)

branching, leaves smooth with a rough keel, racemes lateral and leafy. 24. b. Said to be highly efficacious in the hydrophobia.—*galericulata*, leaves heart-lanceolate, crenate, wrinkled, flowers axillary. 24. b.

ORIGANUM. Calyxes collected into a 4-sided strobile-like cone, with broad intervening bracts: corol with the upper lip erect, flat: under lip 3-parted, divisions nearly equal. 42. 39—*marjorana*, (marjoram) spike ternate, leaves oval obtuse. ☺. p. w. e.—*vulgare*, spikes panicled, bracts longer than calyx. p.

THYMUS. Calyx with the throat closed with hairs: corol with the upper lip flat, emarginate; lower lip longer. 42. 39—*vulgaris*, (thyme) flowers in whorled spikes, leaves ovate revolute. p. e.—*serpyllus*, (mother of thyme) flowers in heads, stem decumbent, leaves flat. p. e.

CLINPODIUM. Involucre many-bristled, bract-like, under the whorl: perianth 2-lipped: corol with the upper lip flat, obcordate, strait. 42. 39—*vulgare*, (field thyme) stem simple, leaves above hairy remotely toothed. 24. p.

PRUNELLA. Calyx with the upper lip dilated: filaments 2-forked with an anther on one of the points: stigma 2-cleft. 42. 39—*pensylvanica*, (self-heal, or heal-all) leaves lance-ovate toothed, upper lip of calyx 3-awned. 24. p. A variety, *ovata*, has ovate leaves, and *lanceolata*, lanceolate leaves and blue flowers.

PHRYMA. Calyx 5-toothed: seed single. (The under lip of calyx shorter, 2-toothed.) 40. 39—*leptostachya*, (phryma) leaves ovate serrate, spike slender terminal. 24. p.

ORDER II. ANGIOSPERMIA.

A. *Calyxes 2-cleft, or 2-leaved.*

OBOLARIA. Corol 4-cleft, bell-form : capsule 1-celled, 2-valved, many-seeded : stamens from the divisions of the corol. (Persoon says the calyx is 5-cleft—Muhlenberg says it is 2-leaved.) 40. 35—*virginica*, (penny-wort) spikelets terminal, leaves opposite coloured. ☹. r-w.

B. *Calyxes 4-cleft.*

OROBANCHE. Calyx of 2-lobed lateral leaflets : corol ringent : capsule 1-celled, 2-valved, many-seeded : glands under the base of the germ. 40. 35 *uniflora*, (cancer-root, broom-rape) stem 1-flowered. b-w.—*virginiana*, stem branched. p.

BARTSIA. Calyx lobed, emarginate, coloured : corol less than calyx ; upper lip longest : capsule 2-celled : seed angled. 40. 35—*coccinea*, (painted cup) leaves cut-pinnatifid, bract 3-cleft. ☼. r. & y.

RHINANTHUS. Calyx inflated : corol ringent ; the upper lip mostly compressed, the under lip spread 3-parted : capsule 2-celled, obtuse, compressed. 40. 35—*crista-galli*, (yellow-cockscomb) calyx glabrous, upper lip of corol 2-toothed at the apex, stem spotted. ☺. y-p.—*virginicus*, corol with spreading throat, leaves sinuate-toothed. y.

EUPHRASIA. Calyx cylindric : corol 2-lipped ; upper lip 2-cleft, lower lip 3-lobed, with the divisions 2-cleft : lower anthers lobed spinose. 40. 35—*officinalis*, (eye-bright) lower lip emarginate, leaves ovate toothed. ☺.

MELAMPYRUM. Corol with the upper lip compressed, the margin folded back : capsule 2-celled, oblique, dehiscent on one side : seeds 2, gibbous. 40. 35—*lineare*, (cow-wheat) leaves linear-lanceo-

late, the upper ones toothed, bristled at the base, flowers axillary. ☼. y-w.

SCHWALBEA. Calyx tubular-inflated; upper division small, lower division large emarginate: corol ringent: capsule ovate, compressed, 2-celled, 2-valved with a double partition: seeds many, chaffy. 40. 40—*americana*, (chaff-seed) stem square, hairy, leaves lanceolate. 2. p.

C. *Leaves 5-cleft or 5-leaved.*

ZAPANIA. Flowers capitate: calyx 5-toothed: corol 5-cleft: stigma head-peltate, oblique: pericarp bladder-like, vanishing, connecting the two seeds. 40. 40—*lanceolata*, (fog-fruit) leaves lance-linear sharp-serrate, stem creeping.—*nodiflora*, leaves wedge-form toothed, stem creeping.

SCROPHULARIA. Calyx 5-cleft: corol sub-globose resupinate, middle division of lower lip reflexed: capsule 2-celled. 40. 40—*marylandica*, (fig-wort) leaves cordate serrate, stem blunt angled. 2. p.—*glabrata*, leaves doubly-serrate. 2. p.

ERINUS. Calyx 5-leaved: corol with the limb equally 5-cleft, lobes emarginate; upper lip very short, reflexed: capsule 2-celled. 40. 35—*africanus*, (erinus) flowers lateral sessile, leaves lanceolate, slightly toothed. p.

BIGNONIA. Calyx 5-cleft cup-form: corol with the throat bell-form, 5-cleft, ventricose beneath: capsule silique-like, 2-celled: seed membrane-winged. 40. 45—*radicans*, (trumpet-flower) stem rooting, leaves pinnate. h. r.

RUELLIA. Calyx 5-parted, often 2-bracted: corol somewhat bell-form, limb 5-lobed: stamens conjugate: capsule tapering to both ends, toothed, elastically dehiscent: seeds few. 40. 36—*strepens*, (ruel) leaves ovate entire, peduncles 2-flowered short. 2. p.

BUCHNERA. Calyx obsoletely 5-toothed : corol with the limb in 5 equal divisions, the lobes cordate : capsule 2-celled. 40. 34—*americana*, (buchnera) spikes remotely flowered, leaves lanceolate 3-nerved. 2. b.

ANTIRRHINUM. Calyx 5-leaved or deeply 5-parted, the two lower divisions remote : corol ringent, spurred or with a prominent base, the gape closed with a prominent palate : capsule ovate, 2-valved, dehiscent at the apex. 40. 40—*linaria*, (snap-dragon) flowers in terminal spikes, leaves crowded together lance-linear. 2. y.—*canadense*, (toad-flax) flowers racemed, leaves alternate remote. 3. p.

GERARDIA. Calyx 5-cleft : corol 2-lipped ; lower lip 3-parted with emarginate lobes, middle one 2-parted : capsule 3-celled, dehiscent at the base. 40. 40—*flava*, (false foxglove) hairy, leaves lanceolate gash-pinnatifid below, flowers axillary opposite. 2. y.—*quercifolia*, (oak-leaf foxglove) glabrous, stem panicle-like, pinnatifid leaves with lanceolate divisions, upper leaves entire. 2. y.—*purpurea*, (dwarf foxglove) leaves linear, very branching. 3. p.—*pedicularia*, (lousewort foxglove) hairy, leaves pinnatifid doubly-serrate. 2. y.

DIGITALIS. Calyx 5-parted : corol bell-form, ventricose, 5-cleft : capsule ovate, 2-celled. (Flowers racemed.) 40. 40—*purpurea*, (foxglove) corol obtuse with an entire upper lip, leaves hairy. 3. p.

PEDICULARIS. Calyx 5-cleft or obliquely truncate : corol ringent : capsule 2-celled, mucronate, oblique : seeds numerous, coated. (Leaves many-cleft.) 40. 35—*canadensis*, (lousewort) calyx obliquely truncate, spike leafy short, upper lip of corol with 2 bristles. 2. y. & p.—*gladiata*, (high heal-all) spikes leafy rough-haired, flowers alternate. 2. y-p.

MIMULUS. Calyx prismatic, 5-toothed : corol ringent, upper lip folded back upon its sides : stigma thick : capsule 2-celled, many-seeded. 40. 45—*ringens*, (monkey-flower) leaves sessile lanceolate glabrous. 24. b.

CHELONE. Calyx 5-cleft or 5-leaved : corol ringent, inflated ; the upper lip emarginate obtuse, under lip slightly 3-cleft : the rudiment of a smooth filament between the two tallest stamens : capsule 2-celled. 40. 45—*glabra*, (snake-head) leaves lanceolate serrate, upper ones opposite. 24. w.—*obliqua*, leaves all opposite. 24. p.

PENTSTEMON. Calyx 5-cleft or 5-leaved : corol ringent inflated : the rudiment of a bearded filament between the two tallest stamens : capsule 2-celled. (Taken from the last genus.) 40. 35—*pubescens*, (beard-tongue) stem hairy, upper lip short. 24. p.

MARTYNIA. Calyx 5-cleft : corol ringent almost bell-form : capsule woody and barked, with a hooked beak ; 4-celled, 2-valved. 40. 42—*proboscidea*, (unicorn-plant) stem branched, leaves cordate, fruit hook-beaked. 24. w.

CLASS XV. TETRADYNAMIA.

ORDER I. SILICULOSA.

All plants of this class have flowers with 4-leaved calyxes and cruciform corols.

A. Silicle without a notch at the end.

DRABA. Silicle oval-oblong, valves flattish, parallel to the partition : style none. 39. 63—*verna*, (whitlow-grass) leaves lance-oblong acutish subserrate, with short stiff hairs, scape naked, petals 2-cleft. 24. w.—*caroliniana*, leaves ovate hirsute, scape naked simple smooth, silicle linear. 24. w.

BUNIAS. Silicle somewhat 4-sided, not opening

by valves, caducous ; 2 or 4-celled, wrinkled. 39. 63—*edentula*, (sea rocket) leaves obovate sinuate, silicle with 2 smooth one-seeded toothless joints. ☉. Bigelow.

B. *Silicle with a notch at the end.*

ALYSSUM. Filaments of the two shorter stamens marked with a tooth : silicle emarginate, often hairy. 39. 63—*sativum*, (madwort) leaves clasping, flowers corymbed. ☉. y.

COCHLEARIA. Silicle thick, rugose, many-seeded, 2-valved ; the valves gibbous obtuse. 39. 63—*officinalis*, (scurvy grass) root-leaves round-cordate petioled, stem-leaves sessile oblong. ♂. w.—*armoracia*, (horse-radish) root-leaves lanceolate, crenate. ♀. w. e.

LEPIDIUM. Calyx spreading : corol regular : silicle emarginate, cordate, many-seeded : valves carinate, partition contrary. 39. 63—*sativum*, (pepper-grass) leaves oblong, many-cleft, glabrous. ☉. w. e.—*virginicum*, leaves below pinnatifid, above serrate. ♀. w.

THLASPI. Calyx spreading : silicle emarginate, obcordate, many-seeded ; valves resemble two boats with the keels outward. 39. 63—*bursa-pastoris*, (shepherd's-purse) silicle wedge-obcordate, root-leaves pinnatifid. ☉. w.—*campestre*, (mithridate mustard) silicle roundish-cordate, leaves arrow-form. ☉. w.

ORDER II. SILIQUOSA:†

A. *Calyx leaves converging or closed upon the corols, when the flowers are mature.*

ARABIS. Glands 4, one within each leaflet of the calyx, of the size of the reflected scale : silique

† Plants of this order resemble each other so nearly, that long descriptions are necessary.

compressed, torulose, subdivaricate. 39. 63—*thaliana*, (wall-cress) radical leaves oblong petioled, cauline leaves lanceolate sessile, stem erect rough-haired at the base, petals twice as long as the calyx.

☉. w.—*tuberosa*, stem simple, leaves obtuse lanceolate toothed, lower leaves sub-petioled, root tuberous-fibrous. 24. w.—*lyrata*, leaves glabrous, radical leaves lyrate, stem leaves linear-lanceolate sub-falcate. ☼. w.—*hispidula*, radical leaves liontooth-lyrate, stem-leaves lanceolate with forked hairs, stem branchy glabrous. 24. w.—*canadensis*, leaves lanceolate toothed glabrous, or with very soft hairs. w.

TURRITIS. Calyx converging, erect: corol erect: silique very long, angled, striate. 39. 63—*laevigata*, (tower-mustard) leaves glabrous, radical ones obovate serrate, cauline ones lance-linear entire clasping. ♂. w.—*hirsuta*, leaves all hispid, cauline ones clasping, silique 4-angled. 24. w.

DENTARIA. Calyx longitudinally converging; silique bursting elastically with revolute valves: the partition thick and fungus-like: stigma emarginate. (Roots always fleshy with tooth-like processes.)—*diphylla*, (tooth-root) stem with two ternate leaves nearly opposite and unequally gashed. 24. w-y.—*laciniata*, leaves triternate, leaflets 3-parted linear toothed, racemes terminal many-flowered. 24. w-p.

HESPERIS. Calyx closed, shorter than the claws of the petals: petals when bent obliquely, are linear or obovate: silique sub-terete: stigmas forked, with converging apices. 39. 63—*pinnatifida*, (rocket) leaves sharply serrate, upper ones lanceolate, lower one pinnatifid, siliques shortish sub-peduncled, 24. w.—*tristis*, stem branched bristly spreading. y. e.—*matronalis*, (dame's violet) stem simple erect. leaves lance-ovate denticulate. 24. w. e.

ERYSIMUM. Calyx closed; silique columnar,

exactly 4-sided. 39. 63—*officinale*, (hedge-mustard) leaves runcinate, siliques pressed close to the stem. ☉. y.—*barbarea*, (wintercress) lower leaves lyrate, with a roundish terminal lobe, upper leaves obovate toothed. ♄. y.

BRASSICA. Calyx erect, converging : partition extending beyond the valves of the silique : seed globose : glands between the short stamens and pistil, and between the long stamens and calyx. 39. 63.—*oleracea*, (cabbage) leaves smooth glaucous repand-lobed. ♂. e.—*napus*, (rape) leaves smooth, upper ones lance-cordate clasping, lower ones lyrate toothed, silique torulose. ♂. e.—*rapa*, (turnip) root orbicular depressed, radical leaves lyrate scabrous, cauline ones entire smooth. ♂. e.

RAPHANUS. Calyx closed, setose : silique torose, somewhat jointed, terete, not opening by valves, 1 or 2-celled : glands between the short stamens and pistil, and between the long stamens and calyx. 39. 63.—*raphanistrum*, (white radish) leaves lyrate, silique 2-celled when immature, and not jointed, in maturity 1-celled and jointed. ☉. w. or y. e.—*sativus*, (garden radish) leaves lyrate, silique terete torose 2-celled. ☉. e. Var. *fusiformis*, root spindle form ; *globosa*, root globose ; *niger*, root blackish and sharp-tasted.

CHEIRANTHUS. Calyx closed, two of the leaflets gibbous at the base : petals dilated : silique when young with a glandular tooth each side : stigma 2-lobed : seed flat. 39. 63—*chiri*, (stock-july-flower) leaves lanceolate acute glabrous. e.

B. *Calyx-leaves spread, not lying closed upon the corol when the flowers are mature.*

CARDAMINE. Calyx leaves spreading but little : stigma entire : a single gland between each of the short stamens and the calyx : silique long bursting

elastically with revolute valves. 39. 63—*pensylvanica*, (American watercress) branching, glabrous, leaves pinnate, leaflets angle toothed roundish-oblong obtuse, siliques narrow erect. w.—*virginica*, leaves pinnate, leaflets lanceolate 1-toothed or eared at the base. (In habit much resembles the shepherd's purse.) 24. w.—*hirsuta*, leaves without stipules, leaflets roundish mostly repand-toothed petioled, flowers tetrandrous. w.

SISYMBRIUM. Calyx and corol spreading : silique bursting, with a short terete beak, valves straitish. 39. 63—*nasturtium*, (English watercress) silique declined, leaves pinnate with sub-cordate leaflets. 24. r. Southern states.—*amphibium* (water-radish) silique declined oblong ovate, leaves pinnatifid serrate, petals longer than calyx. 24. w.

SINAPIS. Calyx spreading : corol with strait claws : glands between the short stamens and pistil, and between the long stamens and calyx : partition extending beyond the valves of the silique, ensiform. 39. 63. *nigra*, (mustard) silique glabrous, leaves linear-lanceolate at top. ☺. y. e.—*alba*, silique hispid, leaves pinnatifid. e.

CLASS XVI. MONADELPHIA.

ORDER III. TRIANDRIA.

SISYRINCHIUM. Spathe 2-leaved : perianth 0 : corol superior, 6-cleft, tubular : style 1 : capsule 3-celled. 6. 18—*anceps*, (blue-eyed grass) culm 2-edged, leaves ensiform, fruit globular glabrous. 24. b.

ORDER V. PENTANDRIA.

PASSIFLORA. Calyx 5-parted, coloured : corol 5-petalled, on the calyx : nectary, a triple filamentous crown within the petals : berry pedicelled. 34. 97—*coerulea*, (passion flower) leaves palmate 5-

lobed. b. South America.—*lutea*, leaves cordate 3-lobed. y. Southern States.

ERODIUM. Calyx 5-leaved: corol 5-petalled: nectariferous scales 5-alternating with the filaments: arils 5, 1-seeded, awned, beaked at the base of the receptacle; awn spiral, bearded within. (Taken from the geranium.) 14. 73.—*cicutarium*, (stork's bill) peduncles many-flowered, leaves pinnate with sessile gash-pinnatifid leaflets, stem prostrate rough-haired. ☹. p.

* *Lysimachia*, *Linum*, *Anagallis*.

ORDER X. DECANDRIA.

GERANIUM. Calyx 5-leaved: corol 5-petalled, regular: nectariferous glands 5, adhering to the base of the long filaments: arils 5, 1-seeded, awned, beaked at the head of the receptacle: awn naked, strait. 14. 73.—*maculatum*, (spotted cranebill, or crowfoot) stem dichotomous erect, leaves gash-5-parted, peduncles 2-flowered. ♀. p.—*robertianum*, (herb-robert) calyx 10-angled pilose, peduncle 2-flowered, leaves in 3 or 5 three-cleft pinnatifid divisions. ☹. p.—*columbinum*, peduncles thrice as long as the 5-parted leaves, capsules glabrous. ☹. b.

MIMOSA: Calyx 5-toothed: corol 5-cleft, 5-petalled, or 0: stamens capillary, very long, 4 to 10 or more, sometimes not united: legume sometimes jointed. 33. 93.—*brachyloba*, (sensitive plant) unarmed, leaves doubly pinnate 8-paired. ♀. r. w. e.—*glandulosa*, unarmed, leaves doubly-pinnate 12-paired. ♀. w. e.—*sensitiva*, prickly. e.

ORDER XIII. POLYANDRIA.

GORDONIA. Calyx simple, 5-leaved or 5-parted: corol 5-petalled adhering at the base: style 5-cornered, stigma 5-cleft: capsule 5-celled: co-

lumella columnar: seeds 2 with leaf-like wings. 37. 74—*lasianthus*, (leaf-seed) leaves leathery glabrous, flowers long-peduncled. $\frac{1}{2}$. w. Southern states.—*franklinia*, flowers sessile. $\frac{1}{2}$. w.

SIDA. Calyx simple, angular, 5-cleft: style many-parted: capsules many, 1 or 3-seeded. (Pedicel articulated under the apex.) 37. 74—*spinosa*, (Indian-mallows) leaves lance ovate, toothed, stipules bristle-form. ☉. y.—*abutilon*, leaves heart-roundish. ☉. y.—*rhombifolia*, leaves lance-wedge-form. $\frac{1}{2}$. y.

HIBISCUS. Calyx double, outer one many-leaved, inner one about 5-cleft: stigmas 5: capsule 5 or 10-celled, many-seeded. 37. 74—*moscheutos*, (hibiscus, or marsh-mallow) leaves ovate acuminate serrate and downy beneath, petiole bearing the flower. $\frac{1}{4}$. w-p.—*palustris*, (marsh hibiscus) leaves ovate sub-trilobed, flowers axillary. $\frac{1}{4}$. p.—*syriacus*, (althaea frutex, or syrian mallows) leaves wedge-ovate 3-lobed toothed. $\frac{1}{2}$. p. Southern states.—*esculentus*, (okra) leaves cordate 5-lobed, outer calyx 10 to 12-parted, caducous. ☉. w.e.

MALVA. Calyx double, outer one 3-leaved, inner one 5-cleft: capsules many, 1-seeded. 37. 74.—*rotundifolia*, (low mallows) stem prostrate, leaves heart-reniform doubly-crenate or 5 to 7 crenate-lobed. $\frac{1}{4}$. b-p.—*caroliniana*, stem prostrate, leaves 5-lobed cut-toothed. ☉.—*sylvestris*, (common mallows) stem erect, leaves with 7 sharpish lobes. $\frac{1}{4}$. b.—*americana*, leaves ovate rough. ☉. y.

ALTHAEA. Calyx double, outer one 6 or 9-cleft: capsules many, 1-seeded. 37. 74.—*officinalis*, (holly-hock or marsh mallows) leaves oblong-ovate, toothed, 3-lobed, downy. $\frac{1}{4}$. p-w. e.—*rosea*, (common holly-hock) leaves cordate crenate 5 to 7 angled rugose. ♂. p. y. w. e.

Gossypium. Calyx double, outer one 3-cleft:

CLASS XVII. ORDER VI to VIII. 73

capsule 4-celled, seeds involved in wool. 37. 74.—
herbaceum, (cotton) leaves 5-lobed mucronate. ☉.
 w. e.

CLASS XVII. DIADELPHIA.

FROM ORDER VI, HEXANDRIA, TO ORDER VIII,
 OCTANDRIA.

FUMARIA. Calyx 2-leaved, caducous : corol irregular, spurred at the base : filaments 2, each with 3 anthers : capsule drupe-like, 1-celled, 1-seeded, not opening by valves ; seed affixed to the side of the cell. 24. 62.—*officinalis*, (fumitory) leaves more than decomposed with leaflets wedge-lanceolate gashed, stem with spreading branches. ☉. r.

CORYDALIS. Calyx 2-leaved : corol ringent 1 or 2-spurred : filaments 2, membranaceous, each with 3 anthers : capsule silique-like, many-seeded. 24. 62.—*cucullaria*, (colic weed) stemless, corol 2-spurred, scape naked, leaves more than decomposed. 24. y-w. Those found near Williams' College are mostly hexandrous—they have also a two-leaved bract, so near the calyx in the immature state, as to give the appearance of a 4-leaved calyx.—*rosea*, 1-spurred, leaves pinnate-pinnatifid, silique one and a half longer than peduncle. 24. r.y.

POLYGALA. Calyx 5-leaved, unequal, 2 of the leaflets wing-like, larger, coloured : corol irregular (or rather, calyx 3-leaved, corol imperfectly papilionaceous) capsule obcordate, 2-celled, 2-valved. Keel of corol sometimes appendaged. 33. 35.—KEEL APPENDAGED.—*paucifolia*, (evergreen snake-root) keel crested, flowers terminal about in threes, leaves ovate near the flowers. 24. r.—*rubella*, (milkwort) keel crested, raceme nearly bractless, wings of corol 3-nerved obtuse longer than keel, stem-leaves linear obtuse, radical ones ovate. 24. r.

—*incarnata*, keel crested, spiked, stem branched erect, leaves alternate subulate. r.—KEEL NOT APPENDAGED.—*sanguinea*, spikes in heads, peduncles scurfy, stem angled, erect, branched. r.—*verticillata*, leaves narrow-linear whorled, flowers with spaces between them. w.r.—*senega*, (seneca snake-root) spike filiform terminal, stem simple, leaves lance-oblong. $\mathcal{U}$. w.

ORDER X. DECANDRIA.

A. Stamens united in one set.

AMORPHA. Calyx bell-form 4 or 5-cleft: banner ovate, concave, wings and keel 0: legume 2-seeded, falcate. 32. 93.—*fruticosa*, (false indigo) somewhat shrubby, leaves glandular petioled, calyx 4-toothed. $\frac{1}{2}$. b.p.

LUPINUS. Corol 2-lipped: anthers, 5 oblong and 5 roundish: legume coriaceous, torulose. 32. 93.—*perennis* (lupine) leaves divided, calyxes not appendaged alternate. $\mathcal{U}$. b.—*villosus*, leaves simple, calyxes appendaged half-whorled. $\mathcal{U}$. w.p.—*filosus*, leaves divided, calyxes appendaged whorled.

CRYPTOLARIA. Corol with the banner cordate, large: keel acuminate: filaments connate, with a dorsal fissure: style curved; legume pedicelled, turgid. 32. 93.—*sagittalis*, (rattle-box) leaves lance-oblong, stipules acuminate decurrent, opposite to the racemes. ☺. y.—*parviflora*, leaves lanceolate, upper stipules decurrent very short opposite to the racemes.

GENISTA. Calyx 2-lipped, 2 upper teeth very short: banner oblong, reflexed back from the pistil and stamens. 32. 93.—*tinctoria*, (dyer's kneed-grass) leaves lanceolate smooth, branches striate. e.

ARACHIS. Calyx 2-lipped: corol upside down: legume gibbous, torulose, veiny, coriaceous. 32. 93

hypogaea, (pea-nut) stem procumbent, leaves pinnate abrupt. ☹. y.

B. *Stamens in 2 sets ; generally 9 in one set, and 1 alone in the other.*

1. *Legume many-seeded, stigma pubescent.*

PISUM. Calyx with the divisions leaf-like, about equal : banner protruding 2 folds : style compressed, carinate, villose above : legume without down at the suture. 32. 93—*sativum*, (pea) petioles cylindric, stipules rounded at the base and notched. ☹. w. e. Var. *umbellatum*, bouquet pea, peduncles many-flowered terminal : *quadratum*, garden pea, fruit cinereous : *humile*, dwarf pea, leaflets bijugous.—*maritimum*, (sea pea) petioles flattish above, stem angular, stipules somewhat arrow-form. 24. w.

LATHYRUS. Calyx with the two upper divisions shorter : style flat, villose above, broader towards the top. (Stem mostly winged, leaflets 2 or more terminated by a divided tendril.) 32. 93—*odoratus*, (sweet pea) peduncles 2-flowered, leaflets ovate-oblong. ☹. p. e.—*venosus*, stem not winged. ☹.

VICIA. Calyx emarginate above, 2-toothed ; 3 strait long teeth below : banner emarginate : stigma bearded transversely on the lower side. 32. 93.—*americana*, (wild vetch) peduncles many-flowered shorter than the leaves, stipules half-arrow-form. 24. p.—*pusilla*, peduncles axillary 1-flowered, leaves about in sixes, stipules half-arrow-form. y. w.—*mitchilla*, peduncles many-flowered, leaflets about 7-paired retuse-mucronate, legume 2-seeded. w. p. This species was found last summer by professor Mitchill of New-York, and sent to the Italian naturalist, C. S. Rafinesque ; who, discovering it to be a new species, gave it this specific name.—*sativa*, (tare) legumes sessile axillary mostly in pairs, stipules toothed. b-w.—*faba*, (gar-

den bean) legumes nearly sessile axillary 3 together swelling at the seeds, stipules sagittate. ☺. e.

PHASEOLUS. Keel, stamens and style spirally twisted together. 32. 93—STEMS TWINING.—*vulgaris*, (common poled-bean) racemes solitary shorter than the leaves, peduncles in pairs. ☺. p. w. e. *helvolus*, (wild bean) flowers in heads, leaflets 3-cornered oblong. ☺. p.—STEM ERECT.—*nanus*, (bush-bean) bracts larger than calyx, stem smooth. ☺. w. p. e. All of this genus are usually called kidney-beans.

DOLICHOS. Banner with two oblong parallel processes at the base, compressing the wings beneath them. 32. 93—*purpureus*, (cowhage) stem twining glabrous, petioles hairy. ☺. p.—*phaseolus*, stem erect, seeds globular, with coloured hilums. w.

ROBINIA. Calyx 4-cleft, upper division 2-parted: banner large, reflexed, roundish: legume gibbous, elongated. 32. 93—*viscosa*, (locust tree) racemes erect, branches and legumes glandular-viscid. h. r.—*hispida*, stem bristly without thorns. h. p.—*pseudo-acacia*, (false-acacia) racemes pendant, legumes smooth, stipules spinous. h. w.

INDIGOFERA. Calyx spreading: keel with a subulate spur both sides: legume linear, small, terete or quadrangular. 32. 93—*tinctoria*, (indigo) leaves pinnate, racemes shorter than the leaves, stem somewhat woody. e.

2. *Legume many-seeded, stigma without pubescence.*

VEXILLARIA.* Banner very large, spreading, overshadowing the wings: calyx bell-form or tu-

* This name is given as a substitute for that of the 1712th genus of Persoon, which is so severely censured by Doctor Smith in Rees' Cyclopædia.

bular : legume linear, very long, acuminate. 32. 93.—*virginiana*, (butterfly-weed) leaves ternate oblong pubescent, calyxes twinned. b.—*mariana*, leaves ternate, calyx cylindric, bracts lance-linear. w.p.

GALACTIA. Calyx 4-toothed, 2-bracted : petals all oblong, banner broader incumbent : stigma obtuse : legume terete : seed roundish. 32. 93.—*glabella* (milk-way plant) glabrous, leaves oblong-ovate somewhat emarginate. 2. p.

GLYCINE. Calyx 2-lipped, the tooth of the lower lip longest : the end of the keel curling upwards, and, in appearance, pushing back the banner : style incurved with the keel : legume oblong compressed. 32. 93.—*angulata*, (wild bush-bean) leaves ternate, side ones 2-lobed, end one ovate. w.p.—*monica* (pea-vine or wild liquorice) stem twining, hairy, leaves ternate ovate. 3. b-w.—*aphos* (ground-nut) pinnatifid leaves with 7 lance-ovate leaflets, stem twining, root tuberous. 2. p.

GALEGA. Calyx with subulate teeth nearly equal : legume torulose, teretish. (Very glabrous pinnate leaves.) 32. 93.—*virginica* (goat's rue) calyx woolly, leaflets oval-oblong. 2. p.

3. Legumes few-seeded.

MEDICAGO. The keel deflected from the direction of the banner : legume compressed, cochleate. 32. 93.—*sativa* (lucerne clover) leaflets oblong toothed, peduncles racemed. 2. b. c.—*lupulina*, procumbent, leaflets obovate, legume reniform 1-seeded. 8. y.—*maculata*, leaflets obcordate toothed, stipules toothed. 3. y.

ASTRAGALUS. Calyx 5-toothed : keel obtuse : legume somewhat 2-celled, incurved at the suture below. 32. 93.—*canadensis* (milk vetch) spreading, leaflets (about 21) glabrous, legume sub-cylindric.

2. y.—*carolinianus*, erect, leaflets (about 41) hairy beneath, legumes swelled. 2. y.—*secundus*, procumbent, legumes pointing one-way. 2. p.

4. *Legumes about 1-seeded.*

TRIFOLIUM. Flowers sub-capitate : legume included in the calyx, not opening by valves, 2 to 4-seeded. (Leaves always ternate.) 32. 93—*pratense* (red clover) heads dense-ovate, 4 of the calyx-teeth equal, stipules awned. 2. r.—*rapens* (white clover) creeping, heads umbelled, leaflets emarginate serrulate. 2. w.—*arvense* (field clover, or rabbit-foot) heads cylindric, calyx-teeth bristle-form longer than corol, leaflets narrow-obovate. ☺. r-w.

MELILOTUS. Flowers racemed : calyx tubular, 5-toothed : keel simple, shorter than the wings and banner : legume rugose, longer than the calyx. (Taken from the trifolium.) 32. 93—*officinalis* (melilot clover) stem erect, stipules lance-subulate. w.

LESPEDEZA. Calyx 5-parted, divisions nearly equal : keel of the corol transversely obtuse : legume lens-form, unarmed, 1-seeded. (Leaves always ternate. Taken from the hedysarum.) 32. 93—*capitata* (bush-clover) erect, leaves sessile with oblong leaflets, little heads in terminal heaps, legume much smaller than the calyx. p. A variety, *angustifolia*, leaves lance-linear, peduncles long. —*polystachia*, erect and very downy, leaflets round ovate, legume and calyx equal. w.—*procumbens*, procumbent, hairy, leaflets ovate. 2. p.

5. *Legumes jointed, or in lomentis.*

HEDYSARUM. Calyx 5-cleft : keel of corol transversely obtuse : loment many-jointed, joints 1-seeded, compressed, generally hispid. (Most, or all, the species in the Northern States have ternate leaves.) 32. 93—*canadense* (bush-trefoil) leaves

lance-oblong, stipules filiform, flowers racemed. 2l. p.—*violaceum*, leaves oval obtuse, racemes umbelled, flowers in pairs. 2l. p.—*divergens*, leaves oblong obtuse, racemes longer than petioles, flowers in pairs. 4l. p.—*nudiflorum*, leaves roundish-ovate pointed, scape paniced glabrous. 4l. p.—*rotundifolium*, stem prostrate hairy, leaves orbicular. 4l. p.—*canescens*, leaves roundish, stem angular ciliate bristly. 2l. p.

STYLOSANTHES. Calyx tubular, very long, bearing the corol which originates higher than the top of the germ : loment two-jointed, hooked. (Sometimes the pod is in a single joint.) 32. 93—*elator* (pencil-flower) leaflets lanceolate, bracts ciliate, flowers spiked. y.

AESCHYNOMENE. Calyx 2-lipped : loment compressed, one suture strait, another lobed ; joints truncate, 1-seeded. 32. 93—*hispida* (false sensitive plant) petioles with bristly tubercles, leaflets many-paired linear obtuse. 32. y-r.

POLYADELPHOUS PLANTS.

The genera *Ascyrum* and *Hypericum* were placed in the class *polyadelphia* by Linneus, and are still retained in that class by many authors. We, following Persoon's method, have removed them to the class *polyandria*, and left the class *polyadelphia* out of the system.

CLASS XVIII. SYNGENESIA.

ORDER I. POLYGAMIA ÆQUALIS.

A. Florets ligulate.

TRAGOPOGON. Calyx simple, many-leaved : receptacle naked : egret plumose and stiped. † 45. 53

† Pull off the corol, stamens and styles ; whatever is left on the germ is the egret. In some flowers the stipe to sti-

—*dandelion* (goat's beard) stemless, leaves erisiform, flower with a scape. $\mathcal{U}$. y.—*horrifolium*, having a stem. δ . p.

APARGIA. Calyx imbricate: receptacle naked: egret plumose, sessile. (Taken from the leontodon.) 49. 53—*autumnalis* (false hawkweed) leaves lanceolate smoothish tooth-pinnatifid, scape branching. y. Dr. Bigelow supposes it came to Boston from Europe.

LEONTODON. Calyx double: receptacle naked: egret plumose, stiped. 49. 53—*taraxacum*, (dandelion) leaves radical runcinate, outer calyx-leaves reflexed. $\mathcal{U}$. y.

PRENANTHES. Florets in a simple series (or in one circular row:) calyx calyced: receptacle naked: egret simple, sub-sessile. 49. 53—*alba* (white lettuce) florets numerous, leaves angular and halbert-form toothed, flowers nodding. $\mathcal{U}$. w.—*altissima*, florets about 5, leaves 3-lobed. $\mathcal{U}$. y-r.—*cordata*, leaves cordate toothed ciliate. $\mathcal{U}$. y.

LACTUCA. Calyx imbricate, cylindric, with the margin membranaceous: receptacle naked: egret simple, stiped: seed smooth. 49. 63—*sativa* (garden lettuce) leaves rounded, those of the stem cordate, corymbd. y.—*elongata* (tall lettuce) leaves smooth beneath, lower ones runcinate clasping, upper ones lanceolate. $\mathcal{U}$. w. Dr. Bigelow calls this the fire-weed; but the *fire-weed* is a species of *senecio*.

HIERACIUM. Calyx imbricate, ovate; egret simple, sessile: receptacle naked or sub-pillose.—(From white becoming yellowish.) 49. 53—*venosum*

ped egret will not appear before full maturity. Pull off all the germs; if nothing is left, the receptacle is naked. But whatever is left belongs to the receptacle, and gives it the character of chaffy, hairy or bristly.

(veiny hawkweed) scape naked branched, leaves radical prostrate obovate ciliate entire with coloured veins. $\mathcal{U}$. y.—*kalmii*, stem many-flowered, leaves small lanceolate toothed. $\mathcal{U}$. y.—*scabrum*, panicle glandular hairy. $\mathcal{U}$. y.

SONCHUS. Calyx imbricate inflated : egret pilose, sessile : receptacle naked. 49. 53—*oleraceus* (sow thistle) peduncles woolly umbelled, calyx glabrous, leaves runcinate denticulate. ☺. y.—*palidus*, raceme terminal compound, leaves lance-ensiform clasping toothed. y.—*palustris*, calyx and peduncles bristly, somewhat umbelled, leaves runcinate with sagittate base. $\mathcal{U}$.

CYNARA. Receptacle bristly : calyx dilated, imbricate ; scales with fleshy base, emarginate, pointed : egret plumose, sessile. 49. 54—*scolymus*, (garden artichoke) leaves prickly or unarmed pinnate and undivided, calyx-scales ovate. b. e.

HYOSERIS. Calyx sub-globular, calyced, the exterior very small : receptacle naked : egret unequal, outer simple, pilose, inner membranaceous : seed near the margin often involved in the scales of the calyx. 49. 53—*prenanthoides*, (swine lettuce) stem-leaves clasping, radical ones sub-lyrate. y.

KRIGIA. Calyx simple, many-leaved : egret 5 membranous leaves with 5 alternating bristles : receptacle naked. 49. 53—*virginica*, (dwarf-dandelion) scape 1-flowered, leaves lance-lyrate, glabrous. ☺. y.

CICHORIUM. Calyx calyced : egret many chaffy leaves : receptacle somewhat chaffy. 49. 53—*intibus*, (succory or endive) flowers sessile in pairs, leaves runcinate. $\mathcal{U}$. b.

B. *Florets tubulous ; flowers capitate.*

LIATRIS. Calyx imbricate, oblong : egret plumose, coloured : receptacle naked. (Flowers pur-

ple.) 49. 54—*spicata* (gay feather) leaves linear entire glabrous ciliate at base dotted, flowers spiked, calyx-scales linear. 2. p.—*scariosa*, raceme terminal, calyx-scales obovate. 2. p.—*squarrosa*, peduncles 1-flowered. 2. p.

VERNONIA. Calyx imbricate, ovate; egret double, exterior chaffy and short, interior capillary: receptacle naked: stigma 2-cleft, shortish. 49. 55—*noveboracensis* (flat-top) corymb fastigiate, stem glabrous, leaves lanceolate. 2. p.—*fracata*, stem pubescent. 2. r.

CNICUS. Calyx imbricate, with prickly scales: receptacle villose: egret plumose. (Taken from the carduus.) 49. 54—*lanceolatus* (thistle) leaves decurrent hispid pinnatifid with 2-lobed spreading divisions, stem hairy, calyx with lance-thorny scales. ♂. p.—*arvensis* (Canada thistle) leaves sessile glabrous tooth-pinnatifid edged with spines, panicked, scales of calyx minute lanceolate with short soft bristly spines. ♂. p.—*altissimus*, radical leaves pinnatifid, stem leaves lance-oblong toothed ciliate. 2. p.

CARTHAMUS. Calyx ovate, imbricate with scales, ovatish-leafy at apex: egret chaff-hairy or none: receptacle chaff-bristly. 49. 54—*coeruleus* (false-saffron) egret hairy, leaves thorny-toothed. 2. b. e. *tinctorius*, egret none, leaves serrate prickly. ☹. y. e.

OXOPORDON. Calyx imbricate with sharp-mucronate scales: egret pilose, caducous: receptacle alveolate: seed transversely sulcate. 49. 54—*acanthium* (cotton thistle) leaves sinuate downy thorny oblong, decurrent.

CARDUUS. Calyx ovate, imbricate with prickly scales: receptacle villose: egret pilose. 49. 54—*pectinatus* (comb-tooth thistle) leaves decurrent lanceolate pectinate. ♂. p.—*marianus*, leaves clasping, spines channelled doubly-thorned. ☹.

ARCTIUM. Calyx globose, with scales hooked at apex: egret chaff-bristly: receptacle chaffy.

49. 54—*lappa* (burdock) stem-leaves cordate. ♂. p.

SPARGANOPHORUS. Calyx sub-globose, imbricate with unequal scales recurve-spreading at the lips: receptacle naked: seed with a cup-like crown, sub-cartilagenous. 49. 55—*verticillatus* (crown-cup) leaves linear-setaceous whorled, about 1-flowered.

C. Florets tubulous; flowers discoid.

BIDENS. Calyx calyced, nearly equal; sometimes the flower is furnished with one or two radiate florets: receptacle chaffy, flat: egret 2, 3 or 4 awns with reflexed and erect lateral prickles: seed 4-cornered. 49. 55—WITH 1 OR 2 RADIATE FLORETS.—*bipinnata*, (burr-marigold or cuckold) leaves doubly-pinnate. ☺.—*chrysanthemoides*, leaves oblong tapering to both ends, calyx shorter than ray. ☺. y.—*cernua*, leaves lanceolate, outer calyx longer than the florets. ☺. y.—WITHOUT RADIATE FLORETS.—*frondosa*, outer calyx 6 times longer than the florets, lower leaves pinnate, upper ones serrate. ☺. y.—*connata*, stem-leaves in threes, floral ones lance-oblong. ☺. y.

KUHNTIA. Calyx imbricate, cylindrical: egret plumose, sessile: receptacle naked. 49. 55—*eupatoroides*, (false boneset) leaves serrate, corymb compact. ♀. w.—*critonia*, leaves entire, corymb spreading. ♀. w.

EUPATORIUM. Calyx imbricated (rarely simple) oblong: style long, cloven half way down: egret pilose or rough papillose: receptacle naked. 49. 55—FLORETS 6 OR MORE.—*perfoliatum*, (boneset, thoroughwort) leaves connate-perfoliate downy. ♀. w.—*purpureum*, leaves petioled 4 or 5 in a whorl lance-ovate, stem hollow. ♀. p.—*verticillatum*, leaves 3 or 4 in a whorl with wedge-form base

stem solid glabrous. 2l. p.—FLORETS 3 TO 5.—*scs-silifolium*, leaves clasping dotted beneath. 2l. w.—*verbenifolium*, leaves sessile, florets 5, calyx hairy. 2l. w.

MIKANIA. Calyx 4 or 6-leaved, equal ; 4 or 6-flowered ; receptacle naked : egret pilose. 49. 55—*scandens* (climbing boneset) stem glabrous, leaves cordate spread-toothed. w.

CACALIA. Calyx cylindric, oblong ; somewhat calyced at the base : receptacle naked : egret pilose. (Leaves mostly succulent.) 49. 55—*suaveolens* (wild caraway) leaves hastate-sagittate. 2l. w.—*atriplicifolia*, radical leaves cordate, stem-leaves deltoid. 2l. w.—*reniformis*, radical leaves repand-toothed. 2l. w.

ORDER II. POLYGAMIA SUPERFLUA.

A. Flowers discoid.

CONYZA. Calyx imbricate with scales sub-linear ovate ; or hemispheric and sub-scarious : egret simple, capillary : receptacle naked. Florets of the ray with a 3-cleft border, not composing a ray in appearance. 49. 55—*camphorata*, (plowman's wort) leaves lance-ovate sessile, corymb crowded. ☹. p.—*bifrons*, leaves spatulate clasping. 2l.

GNAPHALIUM. Calyx imbricate, with the marginal scales rounded, scarious, shortish, glossy, coloured : receptacle naked : egret pilose or plumose. (Florets often all perfect.) 49. 55—*plantagineum*, (mouse-ear, life everlasting) small, corymb compact, root-leaves spatulate, shoots procumbent, dioecious. 2l. w.—*margaritaceum*, (common life everlasting) leaves lance-linear acuminate, corymb fastigiate, odour pleasant. 2l. w.—*uliginosum*, (cudweed) stem much-branched, leaves lance-linear tapering to both ends, racemes terminal crowded. ☹. w.

ARTEMISIA. Calyx imbricate, with scales rounded, converging : egret 0 : receptacle somewhat villose or nakedish. (Flowers mostly rounded.) 49. 55—**LEAVES PINNATIFID**—*absinthium* (wormwood) divisions of radical leaves obtuse, of stem-leaves acutish. 2. e.—*vulgaris*, divisions of leaves all acute and toothed. 2.—**LEAVES PINNATE**—*abrotanum* (southern wood) lower leaves doubly-pinnate, upper ones pinnate filiform. 2. e.—*pontica*, stem-leaves doubly-pinnate with linear leaflets, flowers nodding. 2.

TANACETUM. Calyx imbricate, hemispheric : scales acuminate : rays obsolete, 3-cleft : egret somewhat marginal : receptacle naked. (Flowers corymbed.) 49. 55—*vulgare*, (tansy) leaves doubly-pinnate. 2. y. e.

B. Flowers radiate.

CHRYSANTHEMUM. Calyx hemispherical, imbricate, with the scales membranous at the margin : receptacle naked : egret none, or a narrow margin. 49. 55—*leucanthemum*, (ox-eyed daisy) leaves clasping obtuse cut-pinnatifid, radical ones obovate. 2. w.—*parthenium*, (feverfew) corymbed, leaves compound, leaflets ovate. 2. e.

TAGETES. Calyx simple, 1-leafed, 5-toothed, tubular : florets of the ray 5, permanent : egret 5 erect awns : receptacle naked. 49. 55—*erecta*, (African marigold) stem erect, calyx angular, leaves pinnate. 2. y. e.—*patula*, (French marigold) stem spreading, calyx smooth. 2. y. e.

BELLIS. Calyx hemispherical, scales equal : egret 0 : receptacle naked conical : seed obovate. 49. 55—*perennis*, (garden daisy) scape 1-flowered, leaves obovate. 2. b-w. e.

PYRETHRUM. Calyx hemispherical, imbricate with scales acutish and scarious margined : egret

a marginal ring : receptacle naked. (Taken from *Chrysanthemum*.) 49. 55—*serotinum*, (wild feverfew) leaves lanceolate, branches corymbed. 2.

ARNICA. Calyx with leaflets equal : egret simple : receptacle naked. The florets of the ray usually have 5 filaments without anthers. 49. 55—*montana*, (arnic) leaves ovate entire having 2 opposite cauline ones. 2. y. e.

DORONICUM. Calyx-scales equal, in two rows, longer than the disk : egret simple : receptacle naked : seeds of the ray naked and destitute of egret. 49. 55—*nudicaule*, (leopard's bane) stem almost leafless. 2. y.

INULA. Calyx imbricate or squarrose : egret simple : anthers ending in 2 bristles at the base : receptacle naked : ray florets numerous. 49. 55—*helenium*, (elecampane) leaves clasping ovate downy beneath, calyx-leaves ovate. 2. y.—*mariana*, leaves sessile lance-oblong glandular-toothed tapering base. 2. y.

ERIGERON. Calyx imbricate : egret pilose : receptacle naked : florets of the ray linear, very narrow, numerous. 49. 55—*bellidifolium*, (early flea-bane) root-leaves obovate serrate, stem-leaves lanceolate entire. 2. p. w.—*philadelphicum*, (blue flea-bane) stem-leaves half-clasping lanceolate subserrate, ray-florets as long as those of the disk, stem many-flowered. 2. p. b.—*canadense*, (Canada flea-bane) leaves lanceolate ciliate, lower ones tooth-serrate, paniced, stem and flowers hairy, flowers small numerous. 2. w.

SOLIDAGO. Calyx-scales imbricate, closed : egret simple : receptacle naked, furrowed with dots or punctures : ray-florets about 5. (Yellow.) 49. 55—RACEMES POINTING ONE WAY.—*altissima*, (golden rod) stem erect hairy tall, leaves lanceolate rough wrinkled, lower ones deep-serrate, raceme

panicled. 2. y.—*rugosa*, very rough, lower leaves with close serratures, racemes spread-panicled. y.—*canadensis*, stem downy, leaves rough serrate, racemes panicled recurved, ligulate, florets abbreviated. 2. y.—RACEMES ERECT.—*bicolor*, leaves oval, lower ones serrate, stem and leaves hairy, calyx-leaves obtuse. 2. w-y.—*lanceolata*, stem and leaves glabrous, branching, leaves lance-linear entire, corymbs terminal. 2. y.—*laevigata*, stem and leaves smooth, leaves fleshy, peduncles scaly downy. 2. y.—*caesia*, (a variety of *flexicaulis*) glabrous, leaves acuminate, racemes axillary. y.

SENECIO. Calyx cylindric; leaflets with withering, blackish tips, and a few small calycle leaflets at the base: egret simple: receptacle naked. 49. 55 —*aureus*, (rag-wort) root-leaves heart-ovate serrate petioled, stem-leaves pinnatifid-toothed with lanceolate terminal divisions. 2. y.—*hieracifolius*, ray obscure, leaves clasping oblong unequally toothed. ☼. y.—*vulgaris*, ray obscure, flowers scattered, leaves sinuate-pinnatifid clasping toothed. ☼. y.—*elongatus*, glabrous, root-leaves spatulate, stem-leaves pinnatifid. 2.

TUSSILAGO. Calyx simple, scales equal, and equalling the disk, sub-membranous: receptacle naked: pistillate florets ligulate or without teeth: egret simple, sessile. (Sometimes polygamous.) 49. 55—*farfara*, (colt's foot) scape 1-flowered scaly, at length come out radical heart-angular leaves. 2. y.

ASTER. Calyx imbricate, the inferior scales spreading: egret simple: receptacle naked, scrobiculate. (Florets of the ray more than 10, except in *solidaginoides*, and a few other species.) 49. 55. LEAVES LANCE-LINEAR ENTIRE.—*solidaginoides*, (star-flower) leaves obtuse with scabrous margins, branches fastigate-corymbed, flowers sessile with

5-flowered rays, calyx-scales whitish green-tipped. $\mathcal{U}$. w.—*linarifolius*, leaves keeled dotted rough cuspidate, fastigiate, stem sub-decumbent. p.—*subulatus*, glabrous, leaves subulate, stem paniced, flowers small many. w.b.—*salicifolius*, glabrous, stem paniced at top, calyx-scales acute with spreading tips. b.—*novae-angliae*, leaves clasping eared at the base hairy, calyx-scales lax coloured lanceolate. p.—*cyaneus*, leaves clasping smooth, stem paniced glabrous, branches racemed, inner calyx-scales with coloured tips. b.—LEAVES OVATISH ENTIRE—*cornifolius*, glabrous, leaves acuminate short-petioled with rough margin. w.—*amygdalinus*, leaves tapering to the base, margin rough, fastigiate. w. LEAVES SERRATE CORDATE—*undulatus*, leaves clasping, lower ones with winged petioles. b.—*cordifolius*, leaves all with winged petioles. p.w. LEAVES SERRATE OR TOOTHED OVATE.—*corymbosus*, fastigiate. w.—*chinensis*, (china aster) calyx leafy, branches 1-flowered. e. LEAVES LANCEOLATE, LOWER ONES SERRATE—*funiceus*, leaves clasping, paniced—*diffusus*, leaves glabrous, stem hairy, much-branched. w.

HELENIUM. Calyx 1-leafed, many parted : egret s chaffy leaves : receptacle naked in the disk, chaffy in the ray : florets of the ray half-3-cleft : seed villose. (Leaves decurrent.) 49. 55—*autumnale*, (false-sunflower) leaves serrate glabrous. $\mathcal{U}$. y.

BOLTONIA. Calyx imbricate : egret tooth-awned, somewhat 2-horned : receptacle hemispherical, alveolate : florets of the ray numerous : seed flat-compressed, margined. 49. 55—*asteroides*, (false-chamomile) leaves entire, seeds oval glabrous. $\mathcal{U}$. r-y.—*glastifolia*, lower leaves serrate, seeds obcordate. $\mathcal{U}$. r-y.

MATRICARIA. Calyx flat, imbricate, scales scarious at the margin : receptacle naked, cone-cylin-

dric : egret none. 49. 55—*chamomilla*, (wild chamomile) leaves pinnate and doubly-pinnate.

ANTHEMIS. Calyx hemispherical, nearly equal : egret 0, or a membranous margin : florets of the ray more than 5 : receptacle chaffy, the chaffs flat, with a rigid acuminate apex. 49. 55—*cotula*, (mayweed) receptacle bristly, leaves doubly-pinnate, leaflets 3-parted. ☉. w.—*arvensis*, divisions of leaves lance-linear. ☉. w.—*nobilis*, (chamomile) receptacle chaffy, leaflets 3-parted linear-subulate. 24. e.

ACHILLEA. Calyx imbricate, ovate : egret 0 : receptacle chaffy : florets of the ray 4 to 10, dilated. (Flowers corymbed.) 49. 55—*millefolia*, (yarrow) leaves doubly-pinnatifid, stem grooved. 24. w.

VERBESINA. Calyx in double order : receptacle chaffy : florets of the ray about 5 : egret awned. 49. 55—*sigesbeckia*, (crown-beard) leaves decurrent opposite lance-ovate serrate. 24. y.

ORDER III. POLYGAMIA FRUSTRANEA.

RUDBECKIA. Calyx with a double order of scales : receptacle chaffy, conic : egret a 4-toothed margin or 0. 49. 55—*laciniata*, (cone flower) leaves pinnate, stem-leaflets oblong, with the egret. 24. y.—*purpurea*, leaves lance-ovate alternate undivided, ray-petals 2-cleft. 24. p.—*triloba*, leaves spatulate, lower ones 3-lobed, upper ones undivided. 8. y.

HELIANTHUS. Calyx imbricate, sub-squarrose, leafy : receptacle flat, chaffy : egret 2-leaved, caducous. 49. 55—*annuus*, (sunflower) leaves all cordate 3-nerved, flowers drooping. ☉. y.—*decapetalus*, leaves ovate, stem smooth below, peduncles scabrous, ray about 10-petalled. 24. y.—*divaricatus*, leaves opposite sessile ovate-oblong, panicle forked. 24. y.—*altissimus*, leaves alternate lanceolate, petioles ciliate. 24. y.—*tuberosus*, (Jerusalem arti-

choke) lower leaves heart-ovate, root tuberous. 24. y.

COREOPSIS. Calyx double, also many-leaved : seeds compressed, emarginate : receptacle chaffy : egret 2-horned. 49. 55.—*verticillata*, (tickseed sunflower) leaves pinnate three or five in a whorl. 24. w.—*trichosperma*, pinnate leaves with 5 or 7 leaflets, seed wedge-form. y.—*auriculata*, leaves ovate entire, lower ones ternate. 24. y.

CENTAUREA. Calyx various, mostly imbricate, roundish : egret simple, various : receptacle bristly : corols of the ray funnel-shaped, longer, irregular. 49. 54.—*cyanus*, (blue-bottle) leaves linear entire lower ones toothed. ☉. b. e.—*jacea*, leaves lanceolate entire, radical ones a little toothed. 24. b.—*benedicta*, (blessed thistle) leaves sub-decurrent, calyx with branchy thorns.—*centaureum*, (tall blue-bottle) leaves pinnate, leaflets doubly-serrate. e.

ORDER IV. POLYGAMIA NECESSARIA.

SILPHIUM. Calyx squarrose, scales leafy, broad : egret 2-horned-margin : receptacle chaffy : seeds compressed, obcordate, margined. 49. 55.—*perfoliatum*, (ragged cup) leaves opposite connate, stem 3-sided. 24. y.—*ternatum*, leaves in threes near the middle of the stem, paniced. 24. y.

POLYMNIA. Calyx double, exterior 4-5-leaved, interior 10-leaved, leaflets concave : egret 0 : receptacle chaffy. 49. 55.—*canadensis*, (leaf-cup) leaves denticulate, lower ones pinnatifid. 24. w.—*vedalia*, leaves 3-lobed. 24. y.

IVA. Calyx 3-leaved : styles 2, long : seed naked, obtuse : receptacle hairy : ray 5-flowered. 49. 55.—*frutescens*, (highwater shrub) leaves lanceolate rough-dotted deep-serrate. 2. g.

ORDER V. POLYGAMIA SEGREGATA.

ELEPHANTOPUS. Common calyx 3-leaved, partial calyx 4-flowered : florets ligulate : receptacle naked : egret setaceous. 49. 55—*carolinianus*, (elephant foot) stem simple pilose, leaves narrowed at the base. 4. p. Southern states.

CLASS XIX. GYNANDRIA.

ORDER I. MONANDRIA—OR ORDER II. DIANDRIA.

Calyx 5-leaved coloured, (or corol 5-petalled.)

A. *With a spur under the lower lip.*

ORCHIS. Calyx ringent-like, the upper leaflets vaulted : lip dilated, spur long : anther terminal adnate. 7. 21—*spectabilis*, (scarlet orchis) lip of nectary emarginate, spur as long as germ, stem leafless. r.—*orbiculata*, (round-leaved orchis) lip entire, side-leaflets of calyx spreading, spur longer than germ. g.—*fimbriata*, lip 3-parted, divisions wedge-form fringed, spur longer than germ. 4. p. *psychodes*, lip 3-parted fringe-like, spur as long as germ. 4. y-g.—*ciliaris*, lip 3-parted lanceolate fringed, root-bulb undivided, spur long, 4. y.

LIMODORUM. Calyx somewhat spreading : lip spreading, elongated at the base into a spur : anther a terminal lid, caducous. 7. 21—*unifolium*, (limodore) has but one leaf. w.

SATYRIUM. Calyx ringent-like, upper leaflet vaulted : lip linear, parted or sub-entire : spur short, sub-inflated and somewhat 2-lobed : anther terminal, adnate. 7. 21—*bracteatum*, (satyrion) lip bifid at the apex, bract twice as long as the flower. g.

B. *Lower lip without a spur.*

NEOTTIA. Calyx ringent-like, the outer side leaflets united before about the inflated base of the lip: anther parallel to the acuminate style, affixed behind. 7. 21—*pubescens*, (ladies' traces) radical leaves ovate netted with white veins, scape and flowers hairy, spike oblong. 24. w.—*cernua*, root-bulbs branched, stem leafy sheathed, lip of nectary entire, spike spiral. 24. w.—*tortilis*, lip 3-lobed, middle division crenate, spike spiral, leaves radical-linear. w.—*aestivalis*, stem leafy.

EPIPACTIS. Calyx erect, spreading: anther a permanent lid: lip flat, pendulous, 2-cleft at the apex. 7. 21—*convallaroides*, (lily orchis) stem with 2 roundish cordate leaves, lip obtuse 2-lobed.

CYMBIDIUM. Calyx 4 or 5-leaved, erect or spreading: lip concave at the base, with a spreading border: anther a deciduous lid: pollen globose. 7. 21—*corallorhizon*, (coral root) lip oblong acute undivided, stem leafless, sheathed. y-w. Plant white—*odontorhizon*, lip ovate obtuse, stem leafless sheathed. w-p. Plant white—*pulchellum*, (grass-pink) radical leaf ensiform, lip erect with expanded border. 24. p.—*hyemale*, leaves radical 5-nerved, lip obovate with crenate margin. y-g.

MALAXIS. Calyx spreading, turned upside down by the twisting of the peduncle when the petals open: lip ascending, concave, spreading: anther a lid. 7. 21—*liliafolia* (marsh-twayblade) leaves 2-ovate-lanceolate, lip wedge-obovate pointed. 24. p-y.—*ophioglossoides*, leaf 1 heart-ovate, lip 2-cleft. 24. w.

ARETHUSA. Calyx somewhat ringent-like: leaflets somewhat converging: anther a permanent lid: pollen powdery-granulated. 7. 21—*bulbosa* (arethusa) leafless, 1-flowered, root globose. 24. p.—*ophioglossoides*, scape 1 or 2-flowered with 2 leaves, root fibrous. 24. p.

CLASS XIX. ORDER II, V AND VI. 99

ORDER II. DIANDRIA.

CYPRIPEDIUM. Calyx coloured, 4-leaved, spreading : lip inflated, hollow : style with a terminal lobe ; and a petal-like appendage on the upper side. 7. 21—*pubescens* (yellow ladies' slipper) stem leafy, style-lobe obtuse, shoe-form nectary shorter than calyx. 2. y.—*spectabile* (ladies' slipper) stem leafy, style-lobe obtuse, nectary longer than calyx. 2. p.—*humile*, stemless, leaves radical. 2. p.

ORDER V. PENTANDRIA.

ASCLEPIAS. Follicles 2 : nectaries 5, concave, containing a little horn : petals reflexed. 30. 47—STEMS SIMPLE, LEAVES OPPOSITE—*syriaca*, (milk-weed, or silk-weed) leaves lance-oblong sharp downy beneath, umbels downy. 2. w-p.—*quadrifolia*, four largest leaves near the middle of the stem. 2. w-r.—*purpurascens*, leaves ovate downy beneath, nectaries upside down, umbels erect. 2. p.—*obtusifolia*, leaves clasping round-obtuse oval-oblong. 2. p.—STEM SIMPLE, LEAVES WHORLED OR SCATTERED—*verticillata*, leaves mostly whorled narrow-linear. 2. y-w.—*tuberosa*, leaves scattered lance-oblong rough-haired. 2.—*incarnata*, stem branching, leaves lanceolate, hairy. 2. r.

CYNANCHUM. Calyx 5-toothed : corol wheel form : nectary cylindric, 5-toothed : follicles 2. 30. 47—*obliquum* (choak-dog) twining, leaves cordate, corymbs axillary. p.

ORDER VI. HEXANDRIA.

ARISTOLOCHIA. Calyx 0 : corol superior, 1-petalled, ligulate, inflated at the base : capsule 6-celled, many-seeded. 11. 23—*serpentaria* (birth-wort) stem zigzag, scaped, lips of corol lanceolate. 2.—

sipho, stem twining, peduncles bracted, limb of corol 3-cleft. h. y.

* *Arum* was formerly placed here, but is now removed to class *monoecia*.

CLASS XX. MONOECIA.

ORDER I. MONANDRIA.

FLUVIALIS. Staminate flowers—calyx 0 : corol 0 : anther sessile. Pistillate flowers—calyx 0 : corol 0 : style filiform : stigma 2-cleft : capsule 1-seeded. 15. 6. (Flowers axillary.)—*flexilis*, (river-nymph) leaves in sixes linear 2-toothed at the apex. ☹.—*fragilis*, leaves in threes or opposite prickly-toothed. ☹.

CHARA. Staminate flowers—calyx 0 : corol 0 : anther globose, sessile. Pistillate flowers—calyx 0 : corol 0 : style 0 : stigmas 5 : berry 1-celled, many-seeded. (Aquatics.) 15. 6—*vulgaris* (chara) branchlets terete, joints leafy, leaflets oblong-subulate—*foliosa*, the leaflets linear.

NAJAS.—Staminate flowers—calyx 2-lobed : corol 4-cleft : anthers sessile, cohering. Pistillate flowers—calyx 0 : corol 0 : style subulate : stigma 2 or 3-cleft : capsule 1 to 4-seeded. 15. 6—*canadensis* (water nymph) small filiform smooth, leaves narrow-linear.

ORDER II. DIANDRIA.

LEMNA. Staminate flowers—calyx 1-leafed : corol 0. Pistillate flowers—calyx 1-leafed : corol 0 : capsule 1-celled, 2-seeded. 54. 6—*minor* (duck-meat) leaves oval flat both sides cohering at their bases, root solitary—*polyrhiza*, leaves oval flat cohering 2 or 3 together, roots fascicled. Leaves greenish-purple ; appear like flaxseed, scattered upon the surface of stagnant waters, with the roots

growing immediately from the under surface of the leaves, and the flowers from their upper surfaces. Found in great abundance near Williams' College.

PODOSTEMUM. Staminate flowers—calyx 0: corol 0: stamens affixed to a pedicel. Pistillate flowers—calyx 0: corol 0: germ ovate: stigma 1, sessile: capsule 2-celled, 2-valved, many-seeded. 54.6—*ceratophyllum* (thread-foot) stem filiform floating, leaves pinnate, flowers axillary.

ORDER III. TRIANDRIA.

TYPHA. Ament cylindric. Staminate flowers—calyx obsolete, 3-leaved: corol 0. Pistillate flowers—below the staminate: calyx 0: corol 0: seed 1, pedicelled; the pedicels surrounded at the base with long hairs resembling egret. 3. 8—*latifolia* (cat-tail, or reed mace) staminate and pistillate spikes near or adjoining, leaves sub-ensiform. 24.—*angustifolia*, staminate and pistillate spikes remote, leaves half-cylindric. 24.

SPARGANIUM. Staminate flowers—ament roundish: calyx 3-leaved: corol 0. Pistillate flowers—calyx 3-leaved: corol 0: stigma 2-cleft: drupe juiceless, 1-seeded. 3. 8—*ramosum* (burr-reed) leaves 3-sided at the base, common peduncle branched. 24. g-w.—*simplex*, common peduncle simple. 24.

ZEA. Staminate flowers—calyx-glume 2-flowered, awnless: corol-glume awnless. Pistillate flowers—calyx-glume 2-valved: style 1, filiform, pendulous: seed solitary, immersed in an oblong receptacle. 4. 10—*mays* (Indian corn) leaves entire linear-lanceolate keeled. ☉.g. Var. *precox* (summer corn) small and ripens in the summer.

TRIPSACUM. Staminate flowers—calyx-glume 2-flowered, outer one staminate, inner one neutral: corol-glume membranaceous. Pistillate flowers—

calyx-glume 1-flowered, covered with a 1-leafed involucre perforated with hollows : corol-glume 2-valved : styles 2 : seed 1. 4. 10—*dactyloides* (sesame grass) spikes ternate. 4.

COIX. Staminate flowers—in remote spikes : calyx-glumes 2-flowered, awnless : corol-glume awnless. Pistillate flowers—calyx-glume 2-flowered : corol-glume awnless : style 2-parted : seed covered with the bone-like calyx. 4. 10—*lacryma* (Job's tear) culm branched, leaves wrinkled, seeds ovate.

CAREX. Aments imbricate (usually in cylindric spikes.) Staminate flowers—calyx-scales solitary : corol 0. Pistillate flowers—calyx-scales solitary : corol inflated, monopetalous, 2-toothed at the apex : stigmas 2 or 3 : nut 3-sided, inclosed in the permanent corol, which becomes an utriculus-like permanent aril. 3. 9—STIGMAS 2, STAMENS AND PISTILLS IN DISTINCT SPIKES—*coespitosa*† (bog sedge) pistillate spikes about in threes, lower one very short, aril longer than the obtuse scale—*crinita*, staminate spikes in pairs, pistillate spikes in fours, aril shorter than the awned scale—STIGMAS 2, SPIKES SEVERAL, ANDROGYNOUS, WITH STAMINATE FLORETS ABOVE—*stipata*, aril bi-cuspidate plano-convex nerved, culm very rough, spikelets oblong about in fives—*paniculata*, aril with ciliate-serrate margin, spikes paniced.—LIKE THE LAST, EXCEPT WITH PISTILLATE FLORETS ABOVE—*scirpoides*, spikelets about in fours, aril ciliate-serrate—*festucacea* (fescue sedge) spikelets about in eights, aril ciliate-serrate.—STIGMAS 3, SPIKES ANDROGYNOUS WITH STAMINATE FLORETS ABOVE—*pedunculata*,

† Fifty-five species of *Carex* have been examined and described in the Northern States. It is therefore very difficult to select the most common, in conformity with our plan.—We have inserted none but those found most common near Yale College by the author, and near Williams' College by Professor Dewey.

spikes about in fours, aril obovate 3-sided obtuse.—
 STIGMAS 3, STAMENS AND PISTILS ON DISTINCT
 SPIKES, STAMINATE ONES SOLITARY, PISTILLATE
 ONES SESSILE OR WITH INCLOSED PEDUNCLES—
varia, pistillate spikes short about in threes, aril
 shorter than the scale—*tentaculata*, pistillate spikes
 in threes, aril a little longer than the scale—*lupu-*
lina, pistillate spikes in threes, aril mucronate sev-
eral times longer than the scale—oligocarpa, pis-
tillate spikes in pairs, aril obovate beaked longer
than the scale—folliculata, pistillate spike sub-soli-
lary about 6-flowered.—LIKE THE LAST, EXCEPT
 WITH LONG PEDUNCLES AND LONG SHEATHS—
plantaginea (plantain sedge) pistillate spikes in
 fours, culm with leaf-pointed bracts, radical leaves
 broad 3-nerved—*granularis*, pistillate spikes in
 threes, aril nerved globular inflated sub-emargin-
 ate, scale lance-ovate—*conoidea*, pistillate spikes in
 pairs, aril oblong-conic obtuse equalling the awned
 scale.—STIGMAS 3, STAMINATE SPIKES MANY—
pellita, staminate spikes in pairs, aril bifid, shorter
 than the oblong awned scale—*lacustris*, staminate
 spikes in fours, aril 2-forked. A variety, *gigantea*,
 is very large—*vecicaria*, staminate spikes in threes,
 aril oblong inflated bi-cuspidate.

SCLERIA. Staminate flowers—calyx-glume 2 or
 6-valved, many flowered : corol-glumes awnless.
 Pistillate flowers—calyx like the staminate : corol
 0 : stigmas 1 to 3 : nut coloured sub-globose. 3. 9.
 —*verticillata* (whip-grass) flowers spiked, culm 3-
 sided glabrous, fruit globose.

COMPTONIA. Staminate flowers—ament cylin-
 dric with calyx-scales : corol 2-petalled : filaments
 2-forked. Pistillate flowers—spike or ament ovate,
 corol 6-petalled : styles 2 : nut ovate. 50, 99—*as-*
plenifolia (sweet-fern) leaves alternately pinnatifid
 hairy beneath. $\frac{1}{2}$. g.

SERPICULA. Staminate flowers—calyx 4-toothed : corol 4-petalled. Pistillate flowers—calyx 4-parted : nut tomentose. 15. 88—*verticillata* (little snake-weed) leaves whorled prickly serrate. w.

ERIOCAULON. Common calyx many-leaved : florets many in an imbricate head : corol 3 to 6-petalled : stamens higher than the germ : capsule 3-celled : seed 1, crowned with the corol. Central flowers staminate. 6. 13—*pellucidum* (pipe-wort) stem 6 or 7-angled, stem and root pellucid, leaves awl-form 5-nerved. Grows mostly under water—*decangulare*, stem 10-angled—*gnaphaloides*, culm sub-compressed.

ORDER IV. TETRANDRIA.

BOEHMERIA. Staminate flowers—calyx 4-parted : corol 0 : nectary 0. Pistillate flowers—calyx 0 : corol 0 : styles 2 : seed 1, compressed. (Flowers capitate.) 53. 98—*cylindrica* (round-head) leaves opposite glabrous. 2.—*lateriflora*, leaves alternate rough. 2.

URTICA. Staminate flowers—calyx 4-leaved : corol 0 : nectary central cyathiform. Pistillate flowers—calyx 2-leaved (2-valved :) corol 0 : seed 1, glossy. 53. 98—*dioica*, (nettle) leaves opposite cordate coarse-serrate, spike-panicked, dioecious. 2. g.—*urens*, leaves opposite oval 3 to 5-nerved, racemes simple. ☉. g.—*humila*, leaves opposite ovate, racemes 2-parted shortish sub-corymbed. ☉. —*procera*, leaves opposite cordate, petioles ciliate. 2.—*canadensis*, leaves alternate heart-ovate, aments branched 2-ranked erect. 2.—*whitlowi*, (Albany flax) leaves alternate heart-ovate 3-nerved, upper ones opposite, panicle forked, root tuberous. 2. About 6 feet high. Discovered near Albany by Mr. Charles Whitlow.

PARIETARIA. Polygamous. Perfect flowers—

calyx 4-cleft inferior: corol 0: stamens elastic: style 1: seed 1. Pistillate flowers—calyx 2-leaved: seed covered with the dried elongated calyx. 53. 98—*pensylvanica*, (pellitory) involucre longer than the flower, leaves lance-oblong with opaque dots. 2.

Morus. Staminate flowers—calyx 4-parted: corol 0. Pistillate flowers—calyx 4-leaved: corol 0: styles 2: calyx berried: seed 1. 53. 98—*alba*, (mulberry) leaves heart-oblique smooth. 2. w.—*rubra*, leaves cordate downy beneath. 2. w.

ALNUS. Staminate flowers—ament composed of wedge-form truncate 3-flowered receptacles: calyx a scale: corol 4-parted. Pistillate flowers—calyx 2-flowered scales: corol 0: seed compressed ovate wingless. 50. 99—*serrulata*, (alder) leaves obovate acuminate, stipules oval obtuse. 2. g.

Buxus. Staminate flowers—calyx 3-leaved: petals 2: germ a mere rudiment. Pistillate flowers—calyx 4-leaved: petals 3: styles 3: capsules 3-beaked, 3-celled: seeds 2. 38. 96—*sempervirens*, (box) leaves ovate. 2. Var. *angustifolia*, leaves lanceolate. Var. *suffruticosa*, small, leaves obovate.

ORDER V. PENTANDRIA.

XANTHIUM. Staminate flowers—common calyx imbricate: corol 5-cleft, funnel-form: receptacle chaffy. Pistillate flowers—involucre 2-leaved, 1-flowered: corol 0: drupe dry, muricate, 2-cleft: nut 2-celled—*strumarium*, (sea burdock) stem unarmed, leaves cordate 3-nerved, fruit an oval burr. —*orientale*, leaves 3 to 5-lobed.

AMBROSIA. Staminate flowers—common calyx 1-leaved: corol 1-petalled, 5-cleft, funnel-form: receptacle naked. Pistillate flowers—calyx 1-leaved, entire, the swelling part 5-toothed, 1-flowered: corol 0: nut from the indurated calyx, 1-seeded. 54.

32—*elator*, (common hogweed) leaves doubly-pinnate smoothish with ciliate petioles, racemes terminal paniced. ☹.—*artemisifolia*, branches fastigiate. ☹.—*trifida*, leaves 3-lobed, fruit 6-spined. ☹.

AMARANTHUS. Staminate flowers—calyx 3 or 5-leaved: corol 0: stamens 3 or 5. Pistillate flowers.—calyx and corol as the staminate: styles 3: capsule 1-celled, opening transversely: seed 1. 54. 30—*oleraceus*, (pot amaranth) glomerules axillary ramose, leaves rugose oblong very obtuse emarginate. ☹. g.—*hybridus*, leaves lance-ovate, racemes decompound crowded erect. ☹. g-p.—*hypochondriacus*, leaves lance-oblong mucronate. ☹. p.—*albus*, glomerules axillary, leaves obovate retuse, stem 4-sided. ☹.—*melancholicus*, (love-lies bleeding) glomerules axillary peduncled, leaves lance-ovate coloured. ☹. p. e.

ORDER VI. HEXANDRIA?

ZIZANIA. Staminate flowers—calyx 0: corol-glume 2-valved, awnless, with pistillate flowers intermixed. Pistillate flowers—calyx 0: corol-glume 2-valved, cuculate, awned: style 2-parted: seed 1 invested in the plaited corol. 4. 10—*clavulosa*, (water oats) panicles branching, pistillate flowers above, bract clavate.

ORDER XIII. POLYANDRIA.

MYRIOPHYLLUM. Staminate flowers—calyx 4-cleft: petals 4, caducous: stamens 4, 6 or 8. Pistillate flowers—calyx and corol like the staminate: pistils 4: style 0: seeds 4, having a bark. 15. 6—*spicatum*, (water millfoil) leaves all pinnate capillary, lower ones whorled, all under water. 24.

SAGITTARIA. Staminate flowers—calyx 3-leaved: corol 3-petalled: filaments mostly 24. Pistillate flowers—calyx and corol as in the staminate:

pistils many : capsules aggregate, 1-seeded. 5. 13
 —*sagittifolia*, (arrow-head) leaves lanceolate acuminate sagittate with strait lanceolate hind lobes. 2. w.

CALLA. Spathe ovate becoming expanded : spadix covered with the fructification : stamens intermixed. Staminate flowers—calyx and corol 0 : anthers sessile. Pistillate flowers—calyx and corol 0 : berries 1-celled crowned with the short style. 2. 7—*palustris*, (water arum) leaves cordate, spathe ovate. 2. w.

ARUM. Spathe cuculate : spadix not entirely covered with the fructification ; being more or less naked above, with pistillate flowers beneath and staminate in the middle ; (sometimes a few are staminate beneath :) berry mostly 1-seeded : generally cirrose-glandular beneath. 2. 7—*triphyllum*, (Indian turnip) leaves ternate, with ovate entire leaflets, naked part of spadix very long clavate. 2. p. y. g. Near Williams' college mostly dioecious. —*virginicum*, leaves heart-sagittate acute with obtusish hind lobes, naked part of spadix very short. In wet places.

POTERIUM. Staminate flowers—calyx 4-leaved : corol 4-parted : stamens 30 to 50. Pistillate flowers—calyx and corol like the staminate : pistils 2 : berry from the indurated tube of the corol. 54. 92 —*sanguisorba*, (burnet) stem sub-angled, filaments exsert. 2. e.

QUERCUS. Staminate flowers—calyx sub-5-cleft : stamens 5 to 10. Pistillate flowers—calyx 1-leaved, entire, scabrous, (a woody cup :) styles 2 to 5 : nut coriaceous, surrounded at the base by the permanent calyx. 2. 50. 99—LEAVES OBOVATE, TOOTHED —*montana*, (rock-oak, or chesnut oak) leaves nearly equal-toothed.—*bicolor*, (swamp oak) leaves unequally toothed with entire base.—

LEAVES LANCEOLATE TOOTHED.—*castanea*, (yellow oak) leaves acuminate.—LEAVES 3-LOBED AT THE END—*nigra*, (black jack) leaves wedge-form.—LEAVES SINUATE WITH MUCRONATE LOBES, OBLONG—*discolor*, (two-coloured oak) downy leaves.—*rubra*, (red-oak) leaves glabrous obtuse, with acutish lobes, calyx flat.—*coccinea*, (scarlet oak) leaves glabrous deep-sinuate, calyx with tapering base.—*palustris*, (pin oak) leaves glabrous deep-sinuate downy at the axils of the veins, calyx with flat-turbinate base.—DIFFER FROM THE LAST IN BEING OBOVATE.—*tinctoria*, (black oak) leaves pubescent beneath with obsoletely toothed lobes—*ilicifolia*, (scrub oak) leaves 3 to 5-lobed downy beneath.—LEAVES SINUATE WITH AWNLESS LOBES—*alba*, (white oak) leaves downy with entire obtuse lobes.

JUGLANS. Staminate flowers—ament imbricate : calyx a scale : corol 6-parted : filaments 4 to 18. Pistillate flowers—calyx 4-cleft superior : corol 4-cleft : styles 2 : drupe coriaceous or woody : nut sulcate. $\frac{1}{2}$. 50. 94—LEAFETS 7 OR UNDER, ACUMINATE.—*alba*, (hickory) leaflets scabrous pubescent beneath lanceolate, fruit sub-quadrangular.—*squamosa*, (shag-bark walnut) leaflets soft-pubescent lance-oblong, fruit ovate—*glabra*, (pignut) leaflets ovate glabrous with resinous dots beneath, fruit obcordate.—LEAFETS 9 OR OVER—*nigra*, (black walnut) leaflets lanceolate sub-pubescent, fruit globular.—*amara*, (bitter nut) leaflets lanceolate acuminate about 9, odd one sub-sessile, putamen thin.—*cinerea* (butter-nut) leaves lance oblong with soft pubescence and viscous petioles, fruit oblong.

FAGUS. Staminate flowers—ament roundish : calyx 5-cleft, bell-form : stamens about 12. Pistillate flowers—calyx 4-toothed, setose : germs 2 : nuts 2 inclosed in the calyx becoming coriaceous echinate. 50. 99—*ferruginea* (beach) leaves ovate

serrate acuminate pubescent beneath. $\frac{1}{2}$. g.w.—*sylvatica*, leaves glabrous obsoletely serrate. $\frac{1}{2}$.

CASTANEA. Polygamous. Staminate flowers—ament naked, linear: corol 5-petalled: stamens 10 to 20. Pistillate flowers—calyx 5 or 6-leaved, mucricate: germs 3: stigmas pencil-form: nuts 3, with coriaceous putamen, inclosed in the calyx becoming echinate. 50. 9—*vesca-americana*, (chestnut) leaves lance-oblong glabrous, sinuate-serrate. $\frac{1}{2}$.

BETULA. Staminate flowers—ament imbricate, scales peltate 3-flowered: calyx a scale: stamens 10 to 12. Pistillate flowers—calyx a 2-flowered scale: seed 1, winged. 50. 99—*papyracea*, (canoe birch, or paper birch) leaves ovate acuminate doubly-serrate, veins hirsute beneath. $\frac{1}{2}$.—*populifolia*, (white birch, broom birch) leaves deltoid long acuminate very glabrous. $\frac{1}{2}$.—*lenta*, (black birch) leaves heart-ovate sharp-serrate acuminate, petioles and nerves hairy beneath. $\frac{1}{2}$. A variety, *lutea*, (yellow birch) bark yellowish.

CARPINUS. Ament imbricate. Staminate flowers—calyx-scales ciliate: stamens 10. Pistillate flowers—calyx-scales 2-flowered: corol 3-cleft: nut ovate sulcate. 50. 99—*americana*, (hornbeam) scales of the pistillate ament 3-parted with the intermediate division lance-ovate, leaves unequally serrate. $\frac{1}{2}$.

OSTRYA. Ament imbricate. Staminate flowers—calyx a scale: filaments ramose. Pistillate flowers—ament naked: capsules inflated, imbricate, with one seed at its base. 50. 99—*virginica*, (hop hornbeam) leaves oblong-ovate acuminate. $\frac{1}{2}$. g.

CORYLUS. Staminate flowers—ament imbricate: calyx a scale: stamens about 8. Pistillate flowers—calyx 2-parted, lacerated: styles 2: nut ovate, surrounded by and included in the permanent leaf-

like calyx. 50. 99—*americana*, (common hazlenut) calyx of the fruit bell-form with a dilated border, leaves cordate acuminate. $\frac{1}{2}$. g.—*rostrata*, (beaked hazlenut) calyx of the fruit elongated into a tubular beak, leaves oblong-ovate. $\frac{1}{2}$. g.

PLATANUS. Ament globose. Staminate flowers—corol scarcely apparent, anthers growing around the filaments. Pistillate flowers—calyx many-leaved : style with a recurved stigma : seed roundish, crowned with the mucronate style, with egret-like hairs at the base. 50. 99—*occidentalis*, (buttonwood, or plane tree) leaves quinquangular pubescent. $\frac{1}{2}$. g.

LIQUIDAMBAR. Staminate flowers—ament conic, surrounded with a 4-leaved involucre : filaments numerous. Pistillate flowers—ament globose, surrounded with a 4-leaved involucre : calyx 1-leaved, pitcher-form, 2-flowered : styles 2 : capsules 2, surrounded at the base by the calyx, 1-celled, many-seeded. 51. 99—*styraciflua*, (sweetgum) leaves palmate-lobed with the axils of the veins hairy. $\frac{1}{2}$.

ORDER XVI. MONADELPHIA.

PINUS. Staminate flowers—calyx 4-leaved : stamens many : anthers naked. Pistillate flowers—calyxes in strobiles or cones, scales 2-flowered : pistil 1 : nut with a membranaceous wing. (Perhaps more properly a samara.) $\frac{1}{2}$. 51. 100—LEAVES EVERGREEN, 2 IN A FASCICLE.—*resinosa*, (yellow pine) cones not half as long as the leaves—SAME, WITH 3 LEAVES IN A FASCICLE.—*rigida*, (pitch-pine) spines on the scales reflexed—SAME, WITH 5 LEAVES IN A FASCICLE—*strobis*, (white pine) strobiles cylindric longer than leaves.—LEAVES DECIDUOUS FASCICLED—*pendula*, (black larch) strobile oblong—*microcarpa*, (hacmatack)

strobile roundish, few-flowered.—LEAVES SOLITARY, SCALES OF THE CONE SMOOTH AND TAPERING.—*balsamea*, (balsam fir) leaves flat emarginate.—*canadensis*, (hemlock tree) leaves flat denticulate two ranked.—*alba*, (single spruce) leaves 4-sided incurved, strobile sub-cylindric with obovate scales.—*nigra*, (double spruce) leaves 4-sided strait erect, strobile ovate with oval scales.

CUPRESSUS: Staminate flowers—ament imbricate: calyx a scale: corol 0: anthers 4, sessile. Pistillate flowers—ament strobilaceous: calyx a 1-flowered scale: corol 0: stigmas 2, with concave points: nut angled. 51. 100—*thyoides* (white cedar) branchlets compressed and covered with 4 imbricated rows of short leaflets, cones minute and angular. $\frac{1}{2}$.

THUJA. Staminate flowers—ament imbricate: calyx a scale: corol 0: anthers 4. Pistillate flowers—ament a strobile: calyx a 2-flowered scale: corol 0: nut 1, surrounded with a marginal wing. 51. 100—*occidentalis* (arbor vitae†) branches spreading, cones smooth with obtuse scales. $\frac{1}{2}$.

ACALYPHA. Staminate flowers—calyx 3 or 4-leaved: stamens 8 to 16. Pistillate flowers—calyx and corol like the staminate: styles 3, 2-cleft: capsule tricoccous, 3-celled. (Having large bracts.) 38. 95—*virginica*, (three seed mercury) leaves lance-oblong. ☺.

RICINUS. Staminate flowers—calyx 5-parted: stamens numerous. Pistillate flowers—calyx 3-parted: styles 3, 2-cleft: capsules echinate, 3-celled, 3-seeded. 38. 96—*communis* (palma-christi, or castor oil plant) leaves palmate peltate. ☺. e.

PHYLLANTHUS. Staminate flowers—calyx 5 or

† This tree is also called white cedar in Dr. Hosack's catalogue; to which and to Phelps' Calendar, we generally refer, as the standards for English names.

6-parted : filaments united in a column ; anthers 3 and generally several barren filaments. Pistillate flowers—calyx and corol like the staminate : nectary with a 12-angled margin : styles 3 : capsule tricocous. 38. 96—*obovata*, (leaf-flower) leaves obovate obtusish. ☹.

MOMORDICA. Staminate flowers—calyx 5-cleft : corol 5-parted : filaments 3. Pistillate flowers—style 3-cleft : berry gourd-like and bursting elastically : seeds compressed. 34. 97—*balsamina*, (balsam apple) leaves cut-palmate, fruit angular. ☹. e. —*echinata*, fruit setose-echinate, leaves heart-5-lobed. ☹.

SICYOS. Staminate flowers—calyx 5-toothed : corol 5-parted : filaments 3. Pistillate flowers—styles 3-cleft : berry gourd-like, 1-seeded. 34. 97—*angulata*, (single-seed cucumber) leaves angular. ☹. w.

CUCURBITA. Staminate flowers—calyx 5-toothed : corol 5-cleft : filaments 3. Pistillate flowers—calyx and corol like the staminate : pistil 3-cleft : pomaceous berry large, 3 to 5-celled : seeds thickened at the margin. 34. 97—*lagenaria*, (gourd) leaves cordate round-obtuse hairy, fruit clavate with a woody rind. e.—*verrucosa*, (warted squash) leaves cordate deeply 5-lobed with the middle lobe narrowed at its base, fruit clavate warty. e.—*melo-pheo*, (flat squash) leaves cordate obtuse sub-5-lobed, fruit depress-bossed. e.—*pheo*, (pumpkin) leaves cordate obtuse sub-5-lobed, fruit roundish or oblong smooth. e. Var. *potiro*, fruit depressed or flattened—*citrullus*, (water-melon) leaves 5-lobed, lobes sinuate-pinnatifid obtuse, fruit oval smooth. e.

CUCUMIS. Staminate flowers—calyx 5-toothed : corol 5-parted : filaments 3. Pistillate flowers—calyx and corol like the staminate : stigmas 3

thick, 2-parted : berry with sharpish seeds. 34. 97—*sativus*, (cucumber) angles of leaves strait, fruit oblong rough. ☼. y. e.—*melo*, (musk-melon) angles of leaves round, fruit torulose. e.

MELOTHRIA. Staminate flowers—calyx 5-toothed; corol bell-form : filaments 3. Pistillate flowers—style 1 : stigmas 3 : berry 3-celled, many-seeded. 34. 97—*pendula*, (creeping cucumber) leaves 5-lobed, peduncles 1-flowered. ☼. y.

CLASS XXI. DIOECIA.

ORDER II. DIANDRIA.

VALLISNERIA. Staminate flowers—spathe 2-parted : spadix covered with florets : perianth 3-parted. Pistillate flowers—spathe 2-cleft, 1-flowered : perianth 3-parted, superior : corol 3-petaled : capsule 1-celled, many-seeded. (Stamens 2 and 6) 1. 22—*americana*, (tape-grass) leaves linear, peduncles strait. 24.—*spiralis*, peduncle of the fruit spiral, leaves linear with tapering base.

SALIX. Staminate flowers—ament cylindric : calyx a scale, with a nectariferous gland at the base. Pistillate flowers—ament and calyx like the staminate : style 2-cleft : capsule 1-celled, 2 valved : seed with egret-like down. 2. 50. 99.—LEAVES SERRATE—*nigra*, (brittle-joint willow) leaves lanceolate acuminate glabrous with hairy petioles, staminate flowers 4-stamened and in perfection after the leaves appear.—*conifera*, (cone willow) leaves lance oblong remotely serrulate glabrous above and downy beneath, stipules lunate.—*eriocephala*, (swamp willow) twigs downy, leaves oblong-oval sub-retuse at base, aments downy.—*vitellina*, (yellow willow) leaves lance-ovate acute glabrous above—*lucida*, leaves glabrous oblong acuminate with glandular serratures, staminate flowers in perfec-

tion after leaves appear.—*babylonica*, (weeping willow) leaves glabrous lance-linear, branches pendant. e.—LEAVES ENTIRE—*tristis*, leaves lanceolate tapering to both ends, downy beneath—*repens*, (creeping willow) stem depressed, leaves lance-oval.

FRAXINUS. Perfect flowers—calyx 0, or 4-parted : corol 0, or 4-petalled : pistil 1 : samara 1-seeded with a lanceolate wing. Pistillate flowers on a distinct plant : calyx, corol, pistils same as staminate. h. 44. 37—*americana* (white ash) leaflets subovate, acuminate entire—*juglandifolia*, (walnut-leaved ash) leaflets ovate serrate.

ORDER III. TRIANDRIA.

FIGUS. Common receptacle fleshy enclosing the apetalous florets, either in the same or in a distinct individual. Staminate flowers—calyx 3-parted.—Pistillate flowers—calyx 5-parted : pistil 1, lateral : seed 1, covered with the closed permanent somewhat fleshy calyx. 53. 98—*carica*, (fig tree) leaves palmate about 3-lobed, fruit glabrous pear-form.

* Carex.

ORDER IV. TETRANDRIA.

HIPPOPHAE. Staminate flowers—calyx 2-parted : corol 0. Pistillate flowers—calyx and corol like the staminate : style 1 : berry 1-seeded. 16. 24—*canadensis*, (sea-buckthorn) leaves ovate.

MYRICA. Staminate flowers—ament oblong : calyx an ovate scale : corol 0. Pistillate flowers—calyx and corol like the staminate : styles 2 : drupe or berry 1-seeded. 50. 99—*cerifera*, (bay-berry) leaves lanceolate slightly toothed. h.—*gale*, (sweet gale) leaves lanceolate slightly serrate. h.

BROUSSONETIA. Staminate flowers—ament cylindric : calyx 4-parted. Pistillate flowers—ament

globose: calyx tubulous, 3 or 4-toothed: germ becoming fleshy, clavate, prominent: styles lateral: seed 1, covered by the calyx. 50. 99.—*papyrifera*, (paper mulberry) leaves lobate or undivided. $\frac{1}{2}$. e.

ORDER V. PENTANDRIA.

HAMILTONIA. Perfect flowers—calyx superior 5-cleft: corol 0: nectary with a 5-toothed disk: stamens 5: pistil 1: fruit a drupe. Staminate flowers like the perfect, except in wanting the pistil—*oleifera*, (American oil-nut) leaves oblong, racemes spike-like hairy. $\frac{1}{2}$.

ZANTHOXYLUM. Staminate flowers—calyx 5-parted: corol 0: stamens 3 to 5. Pistillate flowers—pistils 3 to 5: capsules equal to the number of pistils, 1-seeded. 46. 94.—*fraxineum*, (prickly ash, or tooth-ache tree) leaves pinnate, leaflets ovate obscurely serrate, umbels axillary. $\frac{1}{2}$. Among a great number of these plants which I examined near Yale and Williams' Colleges, I have never been able to find a staminate flower.

SPINACIA. Staminate flowers—calyx 5-parted: corol 0. Pistillate flowers—styles 4: seed 1, within the indurated calyx. 12. 29.—*oleracea*, (spinach) fruit sessile, unarmed or sessile. ☺. w. e.

ACNIDA. Staminate flowers—calyx 5-parted: corol 0. Pistillate flowers—calyx 3-parted: style 0: stigmas 3, sessile: capsule 1-seeded. 53. 29.—*cannabina*, (water-hemp) leaves lanceolate, capsules slender acute-angled. ☼.—*ruscocarpa*, capsules obtuse-angled. ☺.

HUMULUS. Staminate flowers—calyx 5-leaved: corol 0. Pistillate flowers—calyx 1-leaved, entire, oblique, spreading: styles 2: seed 1, within the leaf-like calyx: inflorescence strobile-form. 53. 98.—*lupulus*, (hop) stem twining dextrorsum

CANNABIS. Staminate flowers—calyx 5-parted.

116 CLASS XXI. ORDERS VI, VIII TO XIII.

Pistillate flowers—calyx 5-leaved, entire, gaping laterally : styles 2 : nut 2-valved, within the closed calyx. 53. 98—*sativa*, (hemp) stem hairy, leaves digitate, staminate flowers solitary. ☺.

NYSSA. Perfect flowers—calyx superior, 5-parted : corol 0 : pistil 1 : fruit a drupe. Staminate flowers—stamens 10. 12. 24—*biflora*, (tupelo tree, or pepperidge) leaves ovate-oblong entire, peduncle of perfect flowers 2-flowered. $\frac{1}{2}$. —*triflora*, (sour gum) leaves oval, peduncle about 3-flowered. $\frac{1}{2}$.

ORDER VI. HEXANDRIA.

SMILAX. Staminate flowers—calyx 5-leaved : corol 0. Pistillate flowers—calyx and corol like the staminate : styles 3 : berry 3-celled : seeds 2. 11. 12—*rotundifolia*, (green briar) stem prickly, leaves cordate acuminate 5 to 7-nerved, climbing. $\frac{1}{2}$. w-y —*peduncularis*, (Jacob's ladder) stem terete unarmed : leaves ovate acuminate 9-nerved, peduncles of fertile umbels longer than the leaves. $\frac{1}{2}$. g.—*herbacea*, stem and leaves unarmed, leaves 7-nerved, climbing with tendrils. $\frac{1}{2}$. w-y.—*sylvestrica*, (true sarsaparilla) stem prickly-terete, leaves lance-oblong 3-nerved mucronate.

DIOSCOREA. Staminate flowers—calyx 6-parted : corol 0. Pistillate flowers—styles 3 : capsule 3-celled, compressed : seed 2, membranaceous. (Leaves generally alternate.) 11. 12—*villosa*, (yam root) leaves downy beneath, cordate. $\frac{1}{2}$.

FROM ORDER VIII. OCTANDRIA, to ORDER XIII.

POLYANDRIA.

POPULUS. Staminate flowers—ament cylindric : calyx a torn scale : corol turbinate, oblique, entire. Pistillate flowers—ament, calyx and corol like the staminate : stigma 4-cleft : capsule 2-celled : seed

with egret-like hairs. (Leaves having a tremulous motion.) $\frac{1}{2}$. 50. 99—*tremuloides*, (white poplar) leaves heart-roundish abruptly acuminate serrate-toothed hairy edges—*trepida*, (high poplar) leaves round-oval smooth both sides, unequally sinuate-toothed. (Young leaves downy.)—*angulata*, (balm of gilead) leaves cordate obtusely hook-toothed glabrous, branches with winged angles—*balsamifera*, (balsam poplar) leaves ovate close serrate whitish or rusty beneath, stipules gummy, buds resinous—*dilatata*, (lombardy poplar) leaves deltoid. e.

DROSYPYROS. Staminate flowers—calyx 4 to 6-cleft dilated : corol pitcher-form, 4 to 6 cleft : stamens 8, 16 or 29. Pistillate flowers—calyx and corol like the staminate : styles 4-cleft : berry 8 to 12-seeded. 18 41—*virginiana*, (date plumb) leaves ovate obtusish glabrous. $\frac{1}{2}$. w.

GLEDITSCHIA. Perfect flowers—calyx 4-cleft : corol 3-petalled : pistil 1 : capsule legume-like. Staminate flowers—calyx 3-leaved ; corol 3-petalled. Pistillate flowers—calyx 5-leaved : corol 5 petalled. 33. 93—*triacanthos*, (honey-locust) leaflets 10-paired, leaves doubly-pinnate. $\frac{1}{2}$. w.

DATISCA. Staminate flowers—calyx 5 leaved ; corol 0 : anthers sessile, about 15. Pistillate flowers—calyx superior 2-toothed : styles 3 : capsules 3-angled, 3-horned 1-celled, pervious. 54.—*hirta*, (false hemp) leaves pinnate, stem hirsute. $\frac{1}{2}$.

MENISPERMUM. Staminate flowers—calyx 2-leaved : petals 4 or 6 outer and 8 inner : stamens 16. Pistillate flowers—corol like the staminate : 8 barren stamens : germs 2 or 3 : berries paired, 1-seeded. 11. 77—*canadense*, (moonseed) stem twining, leaves peltate cordate round angular. $\frac{1}{2}$. w. —*virginicum*, leaves sub-peltate cordate lobed pubescent beneath $\frac{1}{2}$ w.

ORDER XVI. MONADELPHIA.

JUNIPERUS. Staminate flowers--ament ovate : calyx a scale : stamens 3. Pistillate flowers--calyx 3-parted : petals 3 : styles 3 : berry 3-seeded, consisting of the three unequal tubercles of the calyx. (Nut bony, 1-celled with balsamy glands at the base.) 51. 100--*virginiana*, (red cedar,) leaves united by their bases in threes. $\frac{1}{2}$.--*communis*, (juniper) leaves ternate (or three ranked) spreading mucronate longer than the berry--*sabina*, (savin) leaves opposite obtuse imbricate 4-ways. $\frac{1}{2}$.

TAXUS. Staminate flowers--calyx about 4-bud-leaves : corol 0 : stamens many, anthers peltate, 8-cleft. Pistillate flowers--style 0 : receptacle acorn-cupform ; nut ovate. 51. 100--*canadensis*, (yew) leaves linear pointing 2-ways, margin revolute. $\frac{1}{4}$.

POLYGAMOUS PLANTS.

Some authors still retain the class of Polygamia, but we, following Persoon, have distributed the genera belonging to this class as follows :--*Andropogon* to class 3, order 2. *Holcus* to 3, 2. *Parax* to 5, 2. *Atriplex* to 5, 2. *Celtis* to 5, 2. *Veratrum* to 6, 3. *Melanthium* to 6, 3. *Acer* to 8, 1. *Mimosa* to 16, 10. *Parietaria* to 20, 4. *Fraxinus* to 21, 2. *Nyssa* to 21, 5. *Gleditschia* to 21, 13.

CLASS XXII. CRYPTOGRAMIA.

ORDER I. FILICES.

Ferns are annulated, when each capsule has a kind of chain passing around it at about right angles with the suture. When the fruit is ripe the chain in straitening opens the capsule ; it being attached by its ends to two contiguous edges of it.

A. *With annulated capsules.*1. *Without involucre.*

ACROSTICHUM. Capsules numerous, covering the whole lower surface of the frond. (The fertile leaves differ in shape from the barren. This genus should be examined in the young state; for the scattered fruit of other genera, by becoming confluent when old, often cover the frond.) 55. 5—*aureum* (fork-fern) frond pinnate with alternate leaflets, fruit on the upper leaflets 2.

POLYPODIUM. Capsules disposed in round scattered dots, on various parts of the lower surface of the frond. 55. 5—*vulgare*, (polypod) frond pinnatifid with oblong obtuse sub-serrate lobes, root with chaffy scales—*incanum*, same, with lobes lanced entire—*virginianum*, root smooth—*connectile*, frond doubly-pinnate, ciliate, fruit-dots minute—*calcareum*, frond ternate and doubly-pinnate.

2. *With involucre.*

ONOCLEA. Fruit-dots indeterminate, capsules covering the whole lower surface of the frond: involucre formed by turning in or rolling back the margin of the leaf, which opens inwards, in maturity, towards the midrib. (The fertile leaves are contracted and narrower than the barren ones.) 55. 5—*sensibilis*, (sensitive polypod) barren frond pinnate, fertile one doubly-pinnate—*struthiopteris*, fronds doubly-pinnatifid, divisions roundish.

BLECHNUM. Capsules in uninterrupted lines running parallel to the midrib of the frond on both sides: involucre opens inwards.† (The fruit of

† The involucre opens *inwards*, when it opens or separates, so as to leave the capsules naked on the side next to the midrib.

this genus, when ripe, often covers the whole lower surface of the frond like the *acrostichum*; it must therefore be examined in the young state.) 55. 5—*borealis*, (Roman fern) barren frond pinnatifid, fertile one pinnate with acuminate leaflets—*serrulatum*, frond pinnate, leaflets serrate tapering to both ends.

PTERIS. Capsules arranged in a continued line along the very margin of the frond: involucre opening inwards. (When the leaves are extremely small, the rows of capsules on opposite sides meet and cover the lower surface like the *onoclea*.) 55. 5—*aquilina*, (common brake) leaflets pinnate with lanceolate repand-toothed divisions, upper ones smallest—*gracilis*, leaflets lanceolate obtuse alternate, lower ones pinnatifid—*caudata*, leaflets doubly-pinnate, subdivisions sub-linear—*atropurpurea*, frond doubly-pinnate, subdivisions sub-linear, lines of the capsules somewhat interrupted.

VITTARIA. Capsules in an uninterrupted line along the disk or margin of the frond: involucre double, one of which opens inwards, the other outwards. (Leaves always simple.) 55. 5—*lineata*, (ribbon fern) frond linear very long pendant.

ASPLENIUM. Capsules in lines parallel to each other, situated exactly *upon* the secondary veins of the frond: involucre opening inwards. (By the secondary veins is meant those of the middle part of the disk, not the larger ones at the margin or midrib. The parallel lines of capsules may stand obliquely with respect to the midrib, but their direction must be parallel to each other.) 55. 5—**FROND DOUBLY-PINNATE BELOW.**—*montanum*, (spleen wort) leaflets pinnatifid—*ruta-muraria*, frond only pinnate at top with obtuse leaflets.—**FROND PINNATE ONLY**—*thelypteroides*, leaflets deeply-pinnatifid with tooth-serrate divisions—*angustifolium*,

base of upper leaflets truncate, of lower ones rounded—*ebeneum*, leaflets serrulate, upper ones eared, lower ones cordate—*melanocaulon*, leaflets crenate with wedge-form base. FROND SIMPLE—*rhyzophyllum*, (walking-leaf) frond lanceolate with a rooting apex.

SCOLOPENDRIUM. Capsules in strait or zigzag parallel lines *between* the secondary veins (or ribs) of the frond: involucre double, opening on the top of the lines of capsules and folding down towards each side. (In the genus *asplenium*, the lines of capsules lie across the veins of the frond; but in this they are in line with the veins and attached along their sides or between them.)—*vulgare*, (caterpillar fern) capsules between 2 veins.

WOODWARDIA. Capsules in oblong or oval spots arranged in regular rows on both sides of the midrib: involucre vaulted and opening inwards. (When the leaves have large segments, the capsules are arranged along the midribs of the segments also.) 55. 5—*angustifolia*, (kidney-fruit fern) frond pinnate, segments linear acute entire, barren ones sub-serrate—*virginica*, segments pinnatifid.

ADIANTUM. Capsules disposed in oblong spots arranged along the margin of the frond: involucre is formed by turning back the margin of the frond over the capsules, and it opens inwards. (The lines of oblong spots are generally along that margin, which may be considered the end of the leaf or of the segments of the leaf.) 55. 5—*pedatum*, (maiden-hair) frond pedate, branches pinnate, leaflets halved.

ASPIDIUM. Capsules in scattered roundish spots on various parts of the whole lower surface of the frond: involucre a kidney-form or round membrane, fastened to the frond in or near the center of the fruit-dot, and opening on all sides. (The

involucre, when a little opened, is strictly peltate.)

55. 5. FROND TERNATE—*crucitarium*, (shield-fern) leaflets pinnatifid.—FROND PINNATE WITH UNDIVIDED LEAFLETS—*acrostichoides*, leaflets ciliate-serrate. FROND PINNATE WITH PINNATIFID LEAFLETS—*noveboracensis*, divisions of leaflets subentire ciliate obtusish—*cristatum*, divisions of leaflets tooth-serrate obtuse. FROND DOUBLY-PINNATE.—*felixmas*, leaflets serrate. Var. *erosum*, leaflets erose-serrate—*marginale*, leaflets crenate decurrent.

ATHYRIUM. (Taken from *aspidium*.) Capsules in scattered roundish or oblong spots on various parts of the whole lower surface of the frond: involucre a roundish membrane fastened to the frond at one side of the fruit-dot and opening on the other. (The involucre is generally fastened inwards and opens outwards towards the margin of the frond.) 55. 5. FROND PINNATE—*thelypteris*, (snuff-box fern) divisions of leaflets acute. FROND DOUBLY-PINNATE, LEAFLETS DOUBLY-SERRATE—*felix-femina*, fruit-dots strait-oblong—*asplenoides*, fruit-dots lunulate. SAME, WITH LEAFLETS SERRATE—*bulbiferum*, leaflets opposite, fruit-dots round. SAME, WITH LEAFLETS NOTCHED OR TOOTHED—*tenue*, leaflets decurrent, toothed—*rufidulum*, leaflets crenate with united bases, chaff-bristly beneath.

DICKSONIA. Capsules in small round scattered fruit-dots at the margin of the frond: involucre double; one part is formed of the thin margin of the frond turned over upon the fruit-dots, the other is from the frond inward side of the fruit-dots, each part opening opposite to where it is fastened to the frond. 55. 5—*pilosiuscula*, (fine-haired fern) frond doubly-pinnate, leaflets pinnatifid.

WOODSIA. Involucre perianth-form, with a hairy margin: capsules pedicelled included within the

involucre : fruit-dots roundish, scattered. 55. 5—*hyperbolea*, (calyxed fern) frond pinnate, leaflets 3-parted—*ilvensis*, frond doubly-pinnatifid, divisions oblong.

CHEILANTHES. Fruit-dots roundish, distinct, situated at the margin of the frond : involucre a roundish membrane formed by turning over the margin of the frond upon the fruit dots, and opening inwards. (The margin is always crenate, and each fruit-dot has its own involucre perfectly separate from the others.) 55. 5—*vestita*, (lip-fern) frond doubly-pinnate with pinnatifid leaflets, hairy both sides.

B. *With capsules not annulated.*

SCHIZAEA. Capsules top-form, striate longitudinally and radiately, situated upon the back part of a narrow spike-form appendage of the frond, surrounded with or interspersed among, bundles of hairs. (The appendages, bearing the capsules are generally digitate and occupy the summit of the frond.) 55. 5—*tortuosa*, (one-sided fern) frond simple, spikes one-way.

OSMUNDA. Capsules globose, pedicelled, radiate-striate or wrinkled, having a hinge at the joining of the two valves, which resembles part of the jointed ring of annulated ferns : the capsules either occupy the whole lower surface of the frond, or a panicle-raceme. (The parts of the frond occupied by the fruit are always more contracted than the barren parts.) 55. 5—*cinnamomea*, (flowering fern) frond doubly-pinnatifid, fertile ones distinct and panicle-racemed—*interrupta*, barren leaflets on the frond with the fertile ones, above and below and sometimes among them—*spectabilis*, frond doubly-pinnate, leaflets sharp-serrate—*regalis*, frond bipinnate, terminating in several racemes, very branching and without hairs.

LYGODIUM. Capsules radiate-striated or wrinkled, opening laterally, longitudinally or transversely ; they are situated upon spikelets which proceed from the margin of the frond, each capsule being hid under a scale which falls off frequently in ripening. 55. 5—*palmatum*, (fringed fern) leaves palmate.

APPENDIX TO THE ORDER FILICES.

PTEROIDES.

Plants related to the ferns in their general habit ; but in some degree approaching the habit of pines, umbelliferous and culminiferous plants. The young leaves are never spiral ; in which particular they differ from most of the proper ferns.

BOTRYCHIUM. Capsules coriaceous, globose, one-celled, smooth ; they are disposed in spikes or racemes, generally open lengthwise, sometimes irregularly. 55. 5—**FROND 3-PARTED, BIPINNATIFID**—*gracile*, (grape-fern) divisions of the leaflets sub-pinnatifid and sub-dentate—*dissectum*, divisions 2-parted, 2-toothed at the apex—*virginicum*, divisions obtuse, sub-3-toothed.—**SAME, BIPINNATE**—*fumaroides*, leaflets lunate crenate.—**SOMEWHAT DOUBLY-TERNATE**—*obliquum*, leaflets serrulate, unequally cordate.

OPHIOGLOSSUM. Capsules round, 1-celled, opening transversely ; they are placed on a somewhat jointed spike in two close rows. 55. 5—*vulgatum*, (adder-tongue fern) frond ovate simple, spikes about an inch long.

PSILOTUM. Capsules globose, 3-celled, 3-seeded, opening upwards with three pores ; they are sessile on the branches and axillary. 55. 5—*dichotoma*, (naked fern) stem naked 3-sided forked.

LYCOPodium. Capsules mostly kidney-form or

roundish, 2 or 4-valved opening elastically; they are placed under separate scales in a spike, or sometimes in the axils of leaves. (Very leafy, their stems being generally covered with 2, 3 or 4 rows of narrow simple entire leaves.) 55. 5—*complanatum*, (club-moss, or ground pine) leaves acute 2-rowed united, spikes in pairs or in fours peduncled erect, stem creeping—*obscurum*, leaves scattered 6-rowed, shoots erect, branches radiately forked, spikes nearly sessile cylindric solitary terminal—*rupestre*, leaves scattered ending in a filament, spikes 4-sided sessile.

EQUISETUM. Fruit placed under peltate bodies, which are arranged in whorls forming a spike-form raceme; four spiral filaments surround the seed (probably) which resemble green globules. (Fertile plants mostly leafless, the stems of all are jointed with toothed sheaths at every joint, and usually longitudinally striated and hollow.) 55. 5—*hyemale*, (scouring rush) stem naked very rough, sometimes branched at the base, spike terminal—*arvense*, (field horsetail) fertile stem simple naked, sheaths broad sharp black distinct—*sylvaticum*, (woods horsetail) fertile stem with a few whorls of branches, teeth at the joints broad-pointed.

ORDER II. MUSCI.†

A. Capsules without a lid.

PHASCUM. Capsule furnished with calyptra, but

† So far the most common species of each genus have been given. Perceiving that the work is swelling beyond what was intended; nothing more will be given under the five remaining orders, than is absolutely necessary for studying the genera. Perhaps another edition may appear, which will include all the known species of plants in the Northern States. The manuscript was prepared last summer, under the immediate inspection of Professor Ives of Yale College; and it is now ascertained, that it would not extend this book beyond another hundred pages.

having no lid it falls off closed. (The smallest of mosses, never half an inch high, and more commonly not the eighth of an inch, and almost sessile.) 56. 4—*crispum*, (earth moss) having a stem, leaves lance-acuminate.

B. *Capsules with a lid and without a peristome.*

SPHAGNUM. Calyptré breaks off transversely, leaving the lower half on the capsule: lid caducous. (Found only in swamps. Becomes reddish when old.)—*latifolium*, (peat-moss) leaves ovate obtuse close-imbricate.

GYMNOSTOMUM. Calyptré wholly falling off: lid caducous—*pyriforme*, (toothless moss) leaves ovate acute spread, capsule pear-form, lid obtuse pointed.

C. *Capsule with a lid and furnished with a single peristome, proceeding from the outer surface of the capsule, or from both inner and outer.*

1. *Teeth fewer than 16.*

TETRAPHIS. Teeth 4, taking their origin from both the inner and outer surface of the capsule. 56. 4—*pellucida*, (four-toothed moss) nearly transparent in various parts.

SPLACHNUM. Teeth 8: capsule with an umbrella-form apophysis below its base. 56. 4—*amullaceum*, (umbrella moss) apophysis blubber-like.

2. *Teeth always 16, and neither split, connected, nor in pairs.*

GRIMMIA. Calyptré smallish: peduncle terminal. 56. 4—*controversa*, (awl-cap moss) leaves lance-linear, serrate at top—*alpicola*, leaves obtusish, capsule obconic nearly sessile.

MASCHALOCARPUS. Calyptræ small : peduncle arillary : stem sarmentose. 56. 4—*julaceum*, (creeping moss) ascending, leaves ovate acute nearly as long as the peduncle.

3. *Teeth always 16, and split.*

DICRANUM. Each tooth split part of the way from the tip towards the base, incurved and solid. (Muhlenberg found 23 species in North America.) 56. 4—*bryoides*, (fork moss) leaves 2-rowed compressed, capsule terminal erect.

TRICHOSTOMUM. Each tooth split quite down to the base, erect and filiform. 56. 4—*pallidum*, (hair-mouth moss) stem simple short, leaves capillary and sheathing.

4. *Teeth 16, 32 or 64, connected.*

POLYTRICHUM. Tips of the teeth all meet in a flat membrane, which extends over most of the mouth of the capsule : calyptræ mostly composed of hair. 56. 4—*juniperinum*, (hair-cap moss) capsule square, leaves lance-linear with rough keels.

5. *Teeth 16, 32, or 64, in pairs.*

DIDYMODON. Capsule without any apophysis below its base. (The *splachnum*, having teeth sometimes in pairs, is best distinguished from this genus by its apophysis.) 56. 4—*lineare*, (double-tooth moss) leaves linear.

D. *Capsule with a lid, and furnished with a single peristome proceeding only from the inner surface of the capsule.*

BARBULA. Peristome consists of twisted filaments. 56. 4—*apiculata*, (spiral-tooth moss) stem short, leaves remotish lance-oblong.

DIPHASCUM. Teeth vein-like, lying on a conic plaited membrane ; capsule inflated almost sessile

upon the root. 56. 4—*foliosum*, (bladder mouth) leaves linear acuminate.

E. *Capsules with a lid, and furnished with a double peristome, the outer one consisting of 8 or 16 distinct teeth.*

1. *Inner peristome consisting of teeth or filaments not connected at their bases.*

ORTHOTRICHUM. Inner peristome consists of 8 or 16 fine hair-like filaments, vaulted inwards. 56. 4—*anomalum*, (curve-haired moss) branched, leaves lanceolate keeled obtusish, calyptrae hairy toothed base—*striatum*, leaves acute keeled spread, calyptrae entire.

NECKERA. Outer peristome consists of 16 teeth, the inner one of 16 filaments, erect; peduncles axillary. (Sarmentose.) 56. 4—*pinnata*, (strait-haired moss) decumbent branched, leaves 2-rowed, capsule immersed in the sheath—*macrospora*, peduncles long.

TIMMIA. Outer teeth very short; inner ones 16, long and membranaceous, in distinct pairs: peduncles terminal. 56. 4—*polystachoides*, (many-spiked moss) leaves lance-linear serrate spread, lid depressed in the centre.

CLIMACIUM. Inner peristome consists of 16 filaments, disposed in pairs; each pair is united at top and connected below by transverse fibres: peduncles axillary. 56. 4—*dendroides*, (tree moss) a projecting collar below the outer teeth.

2. *Inner peristome consisting of teeth or filaments connected towards the base by a membrane.*

MEesia. Inner peristome consisting of 16 reticulate flat teeth: peduncle terminal. 56. 4—*longisetia*, (net-tooth moss) having a very long awn.

LESKEA. Outer teeth long-acuminate; inner

peristome consists of a delicate membrane at the base, and upwards becoming uniform acuminate teeth: peduncle axillary. (Greatly resembles Hypnum, but the ciliae between the inner teeth of the hymenium will distinguish it, if a very high magnifier be used.)—*rostrata*, (goose-neck moss) having a long beak. (Muhlenberg found 16 species.)

BARTRAMIA. Inner peristome consists of a delicate membrane terminating in 16 acuminately divided teeth disposed in pairs: peduncles terminal, except *halleriana*. 56. 4—*arcuata*, (bow-moss) branched, leaves serrate striate. (Muhlenberg found 5 species.)

MNIUM. Inner peristome consists of a membrane terminating in pointed teeth which are much perforated with large holes compared with the size of the teeth, having between them fine hairs: peduncles terminal. 56. 4—*cuspidatum*, (pricked-tooth moss) holes through the teeth smallish. (Muhlenberg found 11 species.)

BRYUM. Inner peristome consists of a membrane terminating in pointed teeth not perforated, having between them very fine delicate hairs: peduncles terminal. 56. 4—*conciideum*, (thread moss) leaves entire spread, capsules erect pearform—*nutans*, leaves serrate at top, capsule pendant. (About 12 species have been found in North America.)

HYPNUM. Inner peristome consists of a membrane terminating in pointed teeth not perforated, having between them very fine hairs, to be seen with a good magnifier: peduncles axillary. 56. 4—*serpens*, (sleeping moss) creeping, leaves 1-nerved at bottom, entire. (Muhlenberg examined 38 species in North America.)

3. *Inner peristome consisting of a membrane, or of united filaments.*

FONTINALIS. Inner peristome consisting of a reticulate membrane with longitudinal fibres. 56. 4—*antephyretica*, (water moss) leaves of the sheath obtuse, stem branched—*capillacea*, leaves acute, peduncle long.

BUXBAUMIA. Outer teeth extremely short and truncate; inner peristome consists of 2 or more cylinders of filaments slightly connected together, the inner cylinder being the longest: capsule oblique: peduncle radical. 56. 4—*aphylla*, (leafless moss) capsule peduncled.

F. *Capsule with a lid, and furnished with a double peristome, outer one consisting of 16 connected teeth.*

FUNARIA. Outer teeth oblique and connected at the tips: inner are 16 filaments lying flat: peduncles terminal. 56. 4—*hygrometrica*, (rope moss) leaves inflected, peduncle becomes twisted by drying.

ORDER III. HEPATICAE.†

JUNGERMANNIA. Capsules 4-valved with chain-like seed-dispersers, peduncled, arising from a bell-form calyx. (An apophysis adjoins the capsule, and it has a silvery filamentous peduncle, which often grows several inches in one night. The peduncle rises from a bell-form calyx, which is more permanent than the fruit. In this calyx the fruit lies concealed a while, at length it opens and discloses a small shining black globule, which rises on the silvery peduncle and bursts into 4

† The plants of this order have the fructification much like that of Mosses; but the frond or leafy part sometimes resembles that of Lichens.

valves.) 57. 3.—*forella*, (false moss) branches flowering in the middle, flowers sub-sessile, leaves obovate. (Muhlenberg found 27 species.)

MARCHANTIA. Capsules peduncled, bursting at their summits: seeds attached to elastic fibres: the capsules are placed beneath umbrella-like stars or cones. (It agrees with some species of *jungermannia* in external appearance, but it is of a much firmer texture. The frond is always leafy, lobed, furnished with a strong midrib and beset beneath with villous roots. The disk of the leaf consists of a beautiful green network.) 57. 3.—*polymorpha*, (brook liverwort) frond obtusely lobed. (In the spring some of this species send up ovate anthers or buds on pellucid filaments from the disk of the frond. Near the middle of the summer the umbrellas appear, bearing the fruit under the rays.)

ANTHOCEROS. Capsules horn-like, issuing from a calyx with toothed reflexed margin, covered at first with a brown villous caducous calyptra; afterwards splitting into valves and presenting the columella with the seeds attached: around the calyx, within the substance of the leaf, and in a proper involucre, are inclosed heaps of reddish minute anthers or buds. (The substance of the frond is made up of beautiful 6-sided cells, each of which is marked on the surface with a small protuberance.) 57. 3.—*levis*, (wax liverwort) frond flat, crenate—*punctatus*, frond sinuate-lobed obtuse.

RICCIA. Capsules sessile, globular, 1-celled, crowned with an appearance of withered anthers; they are lodged in the substance of the leaf, and to be discovered only by the darker colour of the place where situated. (Usually found with *Lemna* in ponds and stagnant waters.) 57. 3.—*fluitans*, (gaping-liverwort) fronds repeatedly forked, linear,

smooth, reticulate obtuse, cleft at tips—*natans*, frond obcordate flat with long compressed radicles.

ORDER IV. ALGAE.

FUCUS. Germs imbedded in the gelatinous substance of the frond, aggregate : seeds produced in clustered tubercles, which burst at their summits. 57. 2—*edulis*, (dying seaweed) frond a roundish short stem expanding into a wedge obovate obtuse succulent leaf. An excellent vermifuge. (See Doct. A. R. Griffin's inaugural dissertation, before the New-York college of physicians.)

ULVA. Frond membranaceous or gelatinous : fruit in semi-transparent vesicular membranes within the substance or under the general cuticle of the frond, either solitary or clustered. 57. 2—*purpurea*, (seaweed) lance-oblong, flat, margin wavy entire.

CONFERVA. Consisting of herbaceous tubes or fibres, generally separated within by transverse partitions : seeds scattered through the joints, or contained in solitary closed tubercles attached to the tubes or fibres. 57. 2—*nitida*, (green-hair, frog's spittle) filaments simple lubricous shining, joints cylindric—*fluvialis*, (brook green-hair) filaments branched alternately.

ORDER V. LICHÈNES.†

A. *Receptacle none ; propag. naked, scattered or aggregate.*

LEPRARIA. Efflorescence sub-globular, scattered loosely over a crustaceous base, which it often

† This order is wholly copied from the system of Acharius, translated by Doct. Smith. Acharius has since subdivided *Parmelia*, *Baemyces*, &c. by giving generic names to the sections of species. Sprengel says, these three first genera are merely the first rudiments of other genera, and should be rejected.

covers entirely : frond crustaceous, indeterminate. 57. 2—*chlorina*, (leprous-like lichen*) greenish yellow or dark deep yellow, thick, effuse.

SPILOMA. Efflorescence in shapeless superficial masses, woolly and powdery, rigid ; frond crustaceous mostly indeterminate, but sometimes bordered, thin and uniform. 57. 2—*melanopia*, (efflorescent lichen) blackish flat sub-confluent.

B. *Receptacles† composed of a peculiar compact hardish substance, not in any manner formed out of the frond, from which it differs in colour, being often black.*

1. *Receptacle open.*

VARIOLARIA. Receptacles seated on a thick tartarous crust, which supplies them with an accessory border ; their disk is occupied with an assemblage of powdery substance : the frond is generally determinate. This genus was supposed to have no receptacle, until Dr. Smith detected the error. 57. 2—*faginea*, (small-pox lichen) frond leprous thin glabrous grey, receptacles white.

OPEGRAPHIA. Clefts black, sessile, oblong or linear, simple, confluent or branched, strait or zig-

* Young botanists who have not been informed to what department of nature the word *lichen* applies, may be told, that the following substances are lichens. The spots on rocks and stones which are white, yellow, black, &c. sprinkled over with dots resembling fly-dirt upon window-glass. Also the whitish green patches on old fences, trees, &c. and the substance on trees of a fibrous structure and firm texture, usually called moss. It differs from moss in being more tough and hard, and generally less green.

† As the student will find nothing on lichens like those organs denominated the fructification of phenogamous plants, he may be told that by receptacles are here meant those specks or dots mentioned in the last note, or those saucer-form substances on the large greenish patches on fences.

zag, bordered : their disk flat or slightly concave, naked or powdery, narrow, linear, rarely dilated : their border is of their own substance, often very narrow and scarcely perceptible, sometimes tumid, elevated, and sometimes inflexed : frond crustaceous, uniform, thin, generally indeterminate, occasionally leprous. 57. 2—*scripta*, (lettered lichen) clefts zigzag.

LECIDEA. Spangles sessile, with a flattish or convex disk : frond various, crustaceous ; uniform and limited, scattered and indeterminate, leafy, membranous, or stellate. 57. 2—*immersa*, (spangled lichen) greenish white, spangles black.

CALICIUM. Puffs turbinate, lenticular, or nearly globose : generally elevated on a rigid simple stalk of a different substance from the crust : frond crustaceous, leprous or tartarous, uniform or scattered, powdery and granulated or solid and chequered, or somewhat leafy and imbricated. This genus was confounded with the *fungi*, till distinguished by Persoon. 57. 2—*strigonellum*, (fungus lichen) crust none.

2. Receptacles closed, at length opening.

GYROPHORA. Buttons superficial flattish : frond leafy, peltate, between membranous and leafy. 57. 2—*vellea*, (button lichen) brownish grey, peltate.

BATHELIUM. Receptacles sessile, sub-globose, opening by a deciduous lid, hollow, from 1 to 3 cells lined with seeds : frond crustaceous, indeterminate, uniform.

VERRUCARIA. Tubercles elevated, discharging their chained seeds by a terminal pore, or else they fall out entire : frond thin crustaceous, uniform. Very minute. 57. 2—*nitida*, (mite lichen) pale green, tubercles crowded.

ENDOCARPON. Hollows hidden in the substance

of the frond, each opening by a little mouth : seeds enclosed in cells : frond membranous, somewhat crustaceous, of a determinate figure ; so united to the substance of the receptacles as to appear of a piece, excepting a little difference of colour. 57. 2—*miniatum*, (hidden lichen) peltate, pale, with spread-lobes.

C. *Receptacles formed of the frond itself, with which it agrees in substance and colour : seeds (excepting of the sphaerophoron) enclosed in proper cells.*

1. *Receptacles compound, either pierced or gaping.*

THELOTREMA. Receptacles compound ; the outer ones wart-like, elevated, of the substance of the crust ; inner ones somewhat globular, with a concave disk : frond a firm, continued, indeterminate crust. 57. 2—*pertusus*, (crust lichen) warts hemispheric with 1 or 2 pores.

2. *Receptacles nearly globular, closed ; at length bursting.*

SPHAEROPHORON. Cellules nearly globose, each enclosing a roundish ball of naked compact seeds, which turns to black powder, and is discharged by the torn orifice, leaving the cellule hollow and empty : frond shrubby, branched, solid, rigid, tartarous or minutely fibrous within ; with a smooth cartilaginous coat. A very elegant genus resembling corol. 57. 2—*globiferus*, (seed-ball lichen) shrubby branched cylindric brownish smooth.

ISIDIUM. Receptacle of the seeds terminal, sub-discoid, solid, various in thickness, covered with, and shining through, the crust of the papillae of the frond, which at length bursts, and each receptacle appears coloured and partly exposed : globules also either terminate each papilla or are nearly ses-

sile on the crust itself ; which are coloured and finally fall out, leaving pits at the summits of the branches : frond crustaceous, leprous and tartarous, cracked, rather indeterminate ; either papillary only, or throwing up solid branches. Resembles sphaerophoron ; but differs in having a base, terminal globules, &c. 57. 2—*corallinus*, (coral lichen) warty-branched, tartarous, grey.

3. *Receptacles orbicular, open ; their border of the colour of the frond, surrounding a variously-coloured disk.*

URCEOLARIA. Shields or spangles cup-form, variously coloured, sunk into the substance of the crust, or of its fragments or warts : disk concave, rarely becoming flattish by age, but never elevated above the crust : proper border often scarcely manifest, of the colour of the disk ; the accessory border sometimes formed of the crust, elevated like a ring surrounding the disk of each shield : frond crustaceous, tartarous, uniform, determinate, chequered or cracked, rarely figured or lobate. Often found on stones, &c. and some species are inseparable from the stones on which they grow. 57. 2—*bryophilla*, (pitcher-shield lichen) greyish with glaucous-black shields.

PARMELIA. Shields superficial or elevated, thick or sub-membranous, flattish, convex, or concave, crowned with a free accessory border : frond various, crustaceous, leafy, branched or laciniated, cartilaginous, membranous or gelatinous. A vast genus. One of its best characters is, that the shield has no border of its own substance ; but always has an accessory one of the substance of the crust. 57. 2—*herbacea*, (shield lichen) bright green with fulvous shields.

4. *Receptacles flattened, nearly destitute of a border; their upper side entirely covered with a thin seed-bearing disk, of considerable solidity at the surface.*

STICTA. Shields on the upper side of the frond, sessile, close-pressed, orbicular, membranaceous; their disk flat, rarely somewhat convex; their border accessory, thin, entire, free, slightly-tumid, rarely fringed, of the colour of the frond; pits among the down of the under side, scattered, sessile, minute, coloured: their disk roundish powdery; border elevated, a little inflated, entire or jagged, sometimes obliterated: frond leafy, membranous, or in some measure leathery, lobed or jagged, downy or shaggy beneath. The pits on the back of the frond form the best of its characters. 57. 2—*sylvatica*, (pit-back lichen) glaucous-brown, shields marginal brown.

PELTIDEA. Targets closely attached to the frond about the margin on the upper or under side, rarely lateral or towards the middle, each consisting entirely of a flattish (rarely concave) coloured disk: border accessory, very thin and scarcely discernible, united with the frond and of the same colour: frond rather leathery, with downy veins and fibres beneath, rarely naked; the circumference lobed, its barren lobes depressed, fertile ones elongated, ascending, naked at the base. Best distinguished from the sticta by having prominent veins or ribs underneath, and no pits. 57. 2—*venosa*, (target lichen) grey with divergent lobes, targets round brown.

CETRARIA, targets shield-like, thickish, flat, close-pressed, sessile, near the margin; their circumference loose, rounded, deformed or wavy, disk coloured, slightly convex: border accessory,

very narrow, entire or crenate, of the colour of the frond : frond leafy, membranous or sub-cartilaginous, rigid, smooth and naked on both sides ; sometimes cellular or pitted ; lobed and jagged, often crisped at the edges. The receptacles are rather between shields and targets ; well distinguished from *Peltidea* by its smooth, polished, membranous, sub-rigid frond. 57. 2—*islandica*, (iceland lichen) chesnut-greenish, ciliate-prickly.

CORNICULARIA. Orbs shield-like, terminal, peltate, thickish, cartilaginous, orbicular, flat ; at length rather convex, uneven, irregular ; disk generally of the colour of the frond ; border scarcely any, except in a young state a straight accessory one, entire, naked, or radiated, at length reflexed : frond cartilaginous, rigid, smooth, rather solid, spongy within, of a shrubby tufted branched habit, the branches acute. Resembles the *Parmelia* in the frond and the *Usnea* in the fruit. 57. 2—*tristis*, (horned lichen) brown, branches cylindric—*muricata*, horns lateral and terminal.

USNEA. Orbs sub-coriaceous, flattened, peltate, naked and smooth on both sides, mostly much dilated ; disk first concave, then flat, even, subsequently rather convex and cracked, or warty, scarcely coloured ; border either none or accessory, which is entire or toothed in the circumference, very often radiated : knobs on the same, or on a different plant, which are sessile, lateral, scattered ; at first shield-like and sometimes in a manner bordered, finally convex and warty, coloured. This is the filamentous lichen, tapering, more or less jointed, having a central elastic medullary thread. It is sometimes called tree-moss. 57. 2—*plicata*, (filamentous lichen, or tree-moss) pendant, palish white, orbs greenish white.

5. *Receptacles convex, more or less globose, clothed with a seed-bearing layer, or disk, sessile on the branches, papillary processes, or peculiar stalks of the frond ; terminal and permanent.*

STEREOCAULON. Knobs sub-turbinate, at first furnished with a proper border, then globose, without stalks, simple or clustered, coloured, permanent; disk at first flat and small, but gradually occupying the whole; the border of the same uninterrupted substance and colour, entire, at length obliterated: frond solid, almost woody, caulescent, erect, branched, mostly clothed with scales, rough and fibrous. 57. 2—*ramulosus*, (knobbed lichen) fibrous rough shrubby scale, knobs dark brown.

BAEMYCES. Knobs capitate, nearly globose, with reflexed and scarcely bordered edges, terminating their own proper fruit-stalks, simple or clustered, coloured, permanent; disk properly none, the whole globose surface being covered with the thin, solid, seed-bearing coat; border none: frond either subcrustaceous, softish, granulated, indeterminate, rarely figured; or consisting of a cartilaginous leafy and lobed crust: fruit-stalks elongated out of the very substance of the frond, simple or branched, tubular or nearly solid, dilated or tapering upwards, sometimes barren. 57. 2—*cocciferus*, (scarlet-mouth lichen) stem and cup greenish grey, knobs scarlet, sessile on the edges of the cup.

ORDER VI. FUNGI:

AGARICUS. Fungus with gills underneath. 58
1—*campestris*, (eatable mushroom) gills pink-colour, stem white, with volva.

MERULIUS. Fungus with veins underneath. 58.
1—*cornucopoides*, (veined fungus) stem grooved, pileus dull-yellow.

BOLETUS. Fungus with pores underneath. 58. 1—*igniarius*, (touch-wood) form of a horse's hoof.

THAELAEOPHORA. Fungus warty underneath, the substance corky. 58. 1—*caryophyllea*, (pink fungus) brown with radiate lines and white edge.

HYDNUM. Fungus prickly underneath. 58. 1—*chrysorhizum*, (paper punk) membranaceous flat spreading stemless, root yellow filiform, extending along the grains of decaying timber. Torrey.

MORELLUS. Fungus reticulate or cellular above, smooth underneath. 58. 1—*esculentus*, (morel) without volva, cellular outside, margin attached to stem.

HELVELLA. Fungus turbinate, plaited, wrinkled, inflated, deformed or concave, smooth, elastically ejecting seeds from the upper surface. 58. 1—*calyciformis*, (smooth-cup fungus) ochre, glabrous, scurfy, stem short.

CLAVARIA. Fungus cup-form, erect, elongated, simple or branched: seeds dispersed over the whole surface, or collected in tubercles opening at top. 58. 1—*cornea*, (club fungus) red-orange, simple or cloven.

CLATHRUS. Fungus roundish, cancellated with fleshy branches interwoven with each other. 58. 1—*cancellatus*, (latticed fungus) pilens open lenticled.

PEZIZA. Fungus concave above; hemispherical or bell-form; seeds contained in the cup. 58. 1—*coccinea*, (dish fungus) scarlet within, buff without.

CYATHUS. Fungus bell-form or cylindric, bearing lentiform capsules within. 58. 1—*striatus*, (bird's nest fungus) dark brown, villous.

LYCOPERDON. Fungus becoming powdery and fibrous within: seeds attached internally to the fibres. 58. 1—*bovistum*, (puff-ball) white becoming blackish, spherical.

FULIGO. Fungus with a cellular-fibrous bark; the fibres penetrating in a reticulate manner through a fibrous mass. 58. 1—*flava*, (soot fungus) yellow.

STEMONITIS. Fungus covered with a fugaceous or membranaceous bark, filled with seminiferous fibres elastically expanding themselves. 58. 1—*fascicularis*, (brittle-bark puff-ball) steel-blue, head sub-turbinate, stem very short.

AECIDIUM. Fungus furnished with membraneous cases, glabrous on both sides, filled with naked seeds not cohering 58. 1—*cornutum*, (cased fungus) yellow, cases long curved olive grey.

SPHAERIA. Fungus furnished with cases filled with roundish naked gelatinous seeds. 58. 1—*mori*, (globule fungus) clustered scarlet small.

SCLEROTIUM. Fungus quite simple, globular-oblong, tough, hardish, gaping a little at length in the center, covered with an inseparable bark never opening above. 58. 1—*semen*, (barked fungus) sub-globular, blackish, rugged.

UREDIO. Fungus parasitical, consisting of a mealy powder, destitute of a case, growing from under the cuticles of leaves and stems, bursting at last, with an even margin. 58. 1—*frumenti*, (blight, or grain-rust) linear oblong, black-brown, growing from under the cuticle of culms—*flava*, (yellow rust) sub-globular yellow, growing from under the cuticle of leaves.

TUBERCULARIA. Fungus gelatinous: cup tubercled: stalk thick, filled, closely appressed, bearing the seeds on the upper surface. 58. 1—*vulgaris*, (tubercled fungus) convex rugged with a sub-repand-tooth margin.

MUCOR. Fungus fugacious, heads at first semi-transparent, then becoming opaque, fixed to simple or branched stalks. 58. 1—*mucedo*, (common mould) globular, stem simple grey.

MONILA. Consisting of moniliform filaments clustered into a head. 58. 1—*coespitosa*, (bead-fungus) branched with ternate spikes.

XYLOSTROMA. Fungus leathery, expanded, deformed, penetrating, smooth, even : seeds among the fibres, globular minute. 58. 1—*giganteum*, (leather punk, or oak leather) fibres parallel, the whole fungus filling the interstices between the cleavages of decaying wood.

FINIS.

INDEX OF THE GENERA.

A.	Page.	Alyssum	73
Acalypha	111	Amaranthus	106
Acer	44	Amaryllis	37
Achillæa	95	Ambrosia	105
Acnida	115	Ammania	19
Aconitum	61	Ammi	32
Acorus	40	Amorpha	80
Acrostichum	119	Ampelopsis	28
Actæa	58	Amygdalus	54
Adiantum	121	Anagallis	22
Adonis	64	Anchusa	21
Æcidium	141	Andromeda	47
Æschynomene	85	Andropogon	16
Æsculus	42	Anemone	62
Agaricus	139	Anethum	34
Agave	38	Angelica	32
Agrimonia	53	Annona	63
Agrostemma	51	Anthemis	95
Agrostis	13	Anthoceros	131
Aira	14	Anthoxanthum	9
Aletris	38	Antirrhinum	71
Alisma	42	Apargia	86
Allium	37	Apium	34
Alnus	105	Apocynum	29
Alopecurus	13	Aquilegia	61
Alsine	35	Arabis	73
Althæa	78	Arachis	80

INDEX.

143

Aralia	35	Boehmeria	104
Arbutus	48	Boletus	140
Arctium	89	Boltonia	94
Arenaria	49	Borago	21
Arethusa	98	Botrychium	124
Argemone	60	Brachystemum	65
Aristida	13	Brassica	75
Aristolochia	99	Briza	15
Arnica	92	Bromus	16
Aronia	55	Broussonetia.	114
Artemisia	91	Bryum	129
Arum	107	Buchnera	76
Arundo	16	Bunias	72
Asarum	52	Buxbaumia	130
Asclepias	99	Buxus	105
Ascyrum	58		
Asparagus	39	C.	
Aspidium	121	Cacalia	90
Asplenium	120	Cactus	54
Aster	93	Calla	107
Astragalus	83	Calicium	134
Athyrium	122	Callicarpa	17
Atragene	64	Callitriche	7
Atriplex	31	Caltha	62
Atropa	24	Calycanthus	58
Atheropogon	15	Campanula	26
Avena	16	Cannabis	115
Azalea	25	Capparis	60
		Capsicum	26
B.		Cardamine	75
Baemyces	139	Cardiospermum	44
Barbula	127	Carduus	88
Bartonia	18	Carex	102
Bartramia	129	Carpinus	109
Bartsia	69	Carthamus	88
Bathelium	134	Carum	34
Bellis	91	Cassia	46
Berberis	36	Castanea	109
Beta	50	Catalpa	8
Betula	109	Caulophyllum	36
Bidens	89	Ceanothus	27
Bignonia	70	Celastrus	27
Blechnum	119	Celtis	31
Blitum	7	Cenchrus	12

Centaurea	96	Conyza	90
Centaurella	18	Coreopsis	96
Cephalanthus	18	Coriandrum	31
Cerastium	52	Cornicularia	138
Cercis	46	Cornus	19
Cetraria	137	Corydalis	79
Chærophyllum	33	Corylus	109
Chara	100	Cratægus	55
Cheilanthes	123	Crotolaria	80
Cheiranthus	75	Cucubalus	50
Chelidonium	58	Cucumis	112
Chelone	72	Cucurbita	112
Chenopodium	30	Cunila	67
Chironia	23	Cupressus	111
Chrysanthemum	91	Cuscuta	30
Chrysosplenium	49	Cyathus	140
Cichorium	87	Cymbidium	98
Cicuta	33	Cynanchum	99
Cimicifuga	61	Cynara	87
Cinna	7	Cynoglossum	21
Circaea	9	Cyperus	11
Cistus	59	Cypripedium	99
Citrus	59	D.	
Clathrus	140	Dactylis	14
Clavaria	140	Dalibarda	57
Claytonia	28	Datisca	117
Clematis	62	Datura	23
Cleome	36	Daucus	32
Clethra	46	Decumaria	53
Climacium	128	Delphinium	61
Clinopodium	68	Dentaria	74
Cnicus	88	Dianthus	49
Cochlearia	73	Diapensia	24
Coix	102	Dicksonia	122
Collinsonia	9	Dicranum	127
Comarum	58	Didymodon	127
Commelina	10	Diervilla	26
Comptonia	193	Digitalis	71
Conferva	132	Digitaria	12
Conium	33	Dionaea	47
Conostylus	58	Dioscorea	116
Convallaria	39	Diospyros	117
Convolvulus	23	Diphascum	127

INDEX.

145

Dipsacus	19	Fraxinus	114
Dirca	44	Fucus	132
Dodecatheon	22	Fuligo	141
Dolichos	82	Fumaria	79
Doronicum	92	Funaria	130
Draba	72		
Dracaena	38	G.	
Dracocephalum	67	Galactia	83
Drosera	35	Galanthus	37
		Galega	83
E.		Galium	19
Echium	22	Gaultheria	48
Elephantopus	97	Gaura	43
Eleusine	14	Genista	80
Elymus	14	Gentiana	30
Endocarpon	134	Geranium	77
Epigaea	48	Gerardia	71
Epilobium	43	Geum	58
Epipactis	98	Glaux	29
Equisetum	125	Glechoma	67
Erigeron	92	Gleditschia	117
Erinus	70	Glycine	83
Eriocaulon	104	Gnaphalium	90
Eriophorum	11	Gordonia	77
Erodium	77	Gossypium	78
Eryngium	31	Gratiola	8
Erysimum	74	Grimmia	126
Erythronium	39	Gymnostomum	126
Eupatorium	89	Gyrophora	134
Euphorbia	53		
Euphrasia	69	H.	
Euonymus	27	Hamamelis	20
		Hamiltonia	115
F.		Hedera	29
Fagus	108	Hedysarum	84
Fedia	10	Helenium	94
Ferula	32	Helianthus	95
Festuca	16	Heliotropium	21
Ficus	114	Helleborus	62
Floerkea	37	Helonias	41
Fluvialis	100	Helvella	140
Fontinalis	130	Hemerocallis	38
Fothergillia	60	Hepatica	63
Fragaria	57	Heracleum	32
Frasera	18	Hesperis	74

Heuchera	30	Juncus	40
Hibiscus	78	Jungermannia	130
Hieracium	86	Juniperus	118
Hippophae	114	Justicia	8
Hippuris	7		K.
Holcus	16	Kalmia	47
Holosteum	17	Krigia	87
Hordeum	13	Kuhnia	89
Hortensia	50		L.
Hottonia	23	Lactuca	86
Houstonia	18	Lamium	65
Hudsonia	52	Lathyrus	81
Humulus	115	Laurus	45
Hyacinthus	39	Lavandula	65
Hydnum	140	Lechea	17
Hydrangea	48	Lecidea	134
Hydrastis	62	Ledum	48
Hydrocotyle	31	Leersia	13
Hydropeltis	64	Lemna	100
Hydrophyllum	22	Leontodon	86
Hyoscyamus	24	Leonurus	66
Hyoseris	87	Lepidium	73
Hypericum	60	Lepraria	132
Hypnum	129	Leptanthus	10
Hypoxis	37	Leskea	128
Hyssopus	66	Lespedeza	84
	I.	Liatris	87
Ilex	20	Ligusticum	32
Impatiens	28	Ligustrum	8
Imperatoria	33	Lilium	39
Indigofera	82	Limnetis	14
Inula	92	Limodorum	97
Ipomaea	25	Linnaea	18
Iris	10	Linum	35
Isanthus	64	Liquidambar	110
Isidium	135	Liriodendron	63
Isnardia	19	Lithospermum	21
Itea	28	Lobelia	27
Iva	96	Lolium	15
	J.	Lonicera	26
Jasminum	8	Ludwigia	19
Jeffersonia	44	Lupinus	80
Juglans	108	Lychnis	52

INDEX.

147

Lycium	17	Morellus	140
Lycoperdon	140	Morus	105
Lycopodium	124	Mucor	141
Lycopus	9	Muhlenbergia	13
Lygodium	124	Myosotis	21
Lysimachia	22	Myrica	114
Lythrum	52	Myriophyllum	106
	M.	Myrtus	54
Magnolia	63		N.
Malaxis	98	Najas	100
Malva	78	Narcissus	37
Marchantia	131	Narthecium	40
Marrubium	66	Neckera	128
Martynia	72	Nelumbium	63
Maschalocarpus	127	Neottia	98
Matricaria	94	Nepeta	66
Medeola	41	Nicotiana	23
Medicago	83	Nigella	61
Meesia	128	Nymphaea	60
Melampyrum	69	Nyssa	116
Melanthium	41		O.
Melilotus	84	Obolaria	69
Melissa	67	Oenothera	43
Melothria	113	Onoclea	119
Menispermum	117	Onopordon	88
Mentha	66	Onosmodium	21
Menyanthes	22	Opegrapha	133
Merulius	139	Ophioglossum	124
Mesembryanthemum	56	Orchis	97
Mespilus	55	Origanum	68
Mikania	90	Ornithogallum	40
Milium	13	Orobanche	69
Mimosa	77	Orontium	40
Mimulus	72	Orthotrichum	128
Mitabilis	27	Oryza	40
Mitchella	18	Oryzopsis	12
Mitella	49	Osmunda	123
Mnium	129	Ostrya	109
Mollugo	17	Oxalis	51
Momordica	112	Oxycoccus	43
Monarda	9		P.
Monila	142	Panax	30
Monotropa	48	Paeonia	60

Panicum	12	Poterium	107
Papaver	59	Pothos	19
Parietaria	104	Prenanthes	86
Parmelia	136	Primula	22
Parnassia	35	Prinos	36
Passiflora	76	Proserpinaca	17
Pastinaca	34	Prunella	68
Pedicularis	71	Prunus	54
Peltidea	137	Psilotum	124
Penthorum	51	Pteris	120
Pentstemon	72	Pulmonaria	20
Periploca	29	Pycnanthemum	65
Peziza	140	Pyrethrum	91
Phacelia	24	Pyrola	46
Phalaris	13	Pyrus	56
Phascum	125	Pyxidanthera	25
Phaseolus	82	Q.	
Philadelphus	54	Quercus	107
Phleum	14	Queria	17
Phlox	24	R.	
Phryma	68	Ranunculus	63
Phyllanthus	111	Raphanus	75
Physalis	25	Reseda	53
Phytolacca	52	Rhamnus	27
Pinus	110	Rheum	45
Pisum	81	Rhexia	43
Plantago	17	Rhinanthus	69
Platanus	110	Rhododendron	47
Poa	15	Rhodora	46
Podalyria	45	Rhus	34
Podophyllum	60	Ribes	29
Podostemum	101	Riccia	131
Polemonium	24	Ricinus	111
Polygala	79	Rivina	20
Polygonum	44	Robinia	82
Polymnia	96	Rosa	56
Polypodium	119	Rosmarinus	9
Polytrichum	127	Rubia	18
Pontederia	38	Rubus	56
Populus	116	Rudbeckia	95
Portulacca	52	Ruellia	70
Potamogeton	20	Rumex	42
Potentilla	57	Ruta	46

INDEX.

149

S.

Saccharum	13
Sagina	20
Sagittaria	106
Salicornia	7
Salix	113
Salsola	30
Salvia	9
Sambucus	34
Samolus	26
Sanguinaria	59
Sanguisorba	18
Sanicula	31
Saponaria	49
Sarothra	35
Sarracenia	59
Satureja	66
Satyrion	97
Saururus	43
Saxifraga	48
Scandix	33
Scheuchzeria	41
Schizaea	123
Schoenus	11
Schwalbea	70
Scirpus	11
Scleranthus	49
Scleria	103
Sclerotium	141
Scolopendrium	121
Scrophularia	70
Scutellaria	67
Secale	14
Sedum	51
Selinum	53
Sempervivum	54
Senecio	93
Serpicula	104
Sicyos	112
Sida	78
Silene	50
Silphium	96
Sinapis	76

Sison	32
Sisymbrium	76
Sisyrinchium	76
Sium	32
Smilax	116
Smyrnium	33
Solanum	25
Solidago	92
Sonchus	87
Sorbus	55
Sorghum	15
Sparganium	101
Sparganophorus	89
Spartina	12
Spergula	51
Sphaeria	141
Sphaerophoron	135
Sphagnum	126
Spigelia	23
Spiloma	133
Spinacia	115
Spiraea	56
Splachnum	126
Stachys	66
Staphylea	34
Statice	36
Stellaria	50
Stemonitis	141
Stereocaulon	139
Sticta	137
Stylosanthes	85
Swertia	29
Symphitum	21
Syringa	8
T.	
Tagetes	91
Talinum	53
Tanacetum	91
Taxus	118
Tetraphis	126
Teucrium	64
Thaëlaephora	140
Thalictrum	62

Thelotrema	135	Utricularia	8
Thesium	29	Uvularia	39
Thlapsi	73		V.
Thuja	111	Vaccinium	47
Thymus	68	Valeriana	10
Tiarella	49	Vallisneria	113
Tilia	59	Variolaria	133
Timmia	128	Veratrum	41
Tradescantia	36	Verbascum	23
Tragopogon	85	Verbena	65
Trichodium	14	Verbesina	95
Trichostema	67	Vernonia	88
Trichostomum	127	Veronica	8
Tricophorum	11	Verrucaria	134
Trientalis	42	Vexillaria	82
Trifolium	84	Viburnum	34
Triglochin	42	Vicia	81
Trillium	42	Vinca	25
Triosteum	27	Viola	28
Tripsacum	101	Vitis	28
Triticum	14	Vittaria	120
Trollius	64		W.
Tropaeolum	44	Woodsia	122
Tubercularia	141	Woodwardia	121
Tulipa	39		X.
Turritis	74	Xanthium	105
Tussilago	93	Xanthorhiza	36
Typha	101	Xylosteum	26
		Xylostroma	142
		Xyris	11
Ulmus	31		Z.
Ulva	132	Zanthoxylum	115
Uniola	15	Zapania	70
Urceolaria	136	Zea	101
Uredo	141	Zigadenus	41
Urtica	104	Zizania	106
Usnea	138		

ENGLISH INDEX.

	<i>Page.</i>	Bear-berry	48
Adder tongue	39	Beard-grass	13, 16
African marigold	91	Beard-tongue	72
Agave	38	Beaver tree	63
Agrimony	53	Bed straw	19
Albany flax	104	Beech	108
Alder	105	Beech drops	48
Alexanders	33	Beet	31
Allum root	30	Bell-flower	26
Amaranth	106	Bell-wort	40
Althaea frutex	78	Bermuda-mulberry	17
American cowslip	62	Birch	109
American oil nut	115	black	109
American water-cress	76	canoe or paper	109
Angelica	32	white or broom	109
Apple	56	yellow	109
Arbor vitæ	111	Birds nest	48
Arethusa	98	Birth-wort	99
Arnic	92	Bishop-weed	32
Arrow grass	42	Bitter nut	108
Arrow head	107	Bitter sweet	25
Artichoke	87	Bitter winter-green	46
Ash	114	Black-berry	57
Asparagus	39	Black flower	41
Atamask lily	37	Black-jack	108
Atragene	64	Black root	32
Avens	58	Bladder-campion	50
Balm	67	Bladder mouth	128
Balm of Gilead	117	Bladder-nut	34
Balsam apple	112	Bladder wort	8
Balsam fir	111	Blessed thistle	96
Balsam weed	28	Blight	141
Bane-berry	58	Blite	7
Barberry	36	Blood-root	60
Barley	13	Blue bottle	96
Bass-wood	59	Blue curls	67
Bay-berry	114	Blue-eyed grass	76
Bean	82	Blue gentian	65

Blue grass	15	Caper bush	60
Bog rush	11	Caraway	34
Bog sedge	102	Cardinal flower	27
Boneset	89	Carolina allspice	58
Borage	21	Carpet weed	17
Boston iris	10	Carrot	32
Bowman's root	56	Cassia	46
Box	105	Castor-oil plant	111
Box-flower	25	Catalpa tree	8
Box thorn	18	Catchfly	50
Brake	120	Catmint	66
Brier-herb	57	Cat-tail	101
Brittle bark puff-ball	141	Cedar	111, 118
Brook-weed	26	Celandine	58
Broom corn	15	Celery	34
Broom grass	16	Centaury	24
Broom-rape	69	Chaff seed	70
Buchnera	71	Chamomile	95
Buck-bean	22	Changeable hydrangea	50
Buck-thorn	27	Chara	100
Buck-wheat	45	Cherry	54
Bugbane	61	choak	54
Bugloss	21	common	54
Bullrush	40	white	54
Burdock	89	Chervil	33
Burnet	107	Chestnut	109
Burnet-saxifrage	18	Chestnut oak	107
Burr-marigold	89	Chess	16
Burr-reed	101	Chick-weed	35
Bush clover	84	Chick-winter-green	42
Bush trefoil	84	China aster	94
Butter-cup	64	Chives	37
Butterfly weed	83	Choak berry	55
Butternut	108	Choak-dog	99
Button bush	18	Climbing-boneset	90
Button wood	110	Clover	84
Cabbage	75	Club-moss	125
Camphion	52	Club rush	11
Canada mint	66	Cockfoot grass	12
Canada thistle	88	Cock-roach mullein	23
Cancer root	69	Cockle	51
Canterbury bells	27	Colic weed	79

INDEX.

153

Colt's foot	93	Decumary	53
Columbine	61	Deer grass	44
Comb-tooth thistle	88	Dill	34
Comfrey	22	Dock	42
Common mould	141	Dodder	30
Cone flower	95	Dog bane	29
Coriander	31	Dog tail grass	14
Corn spurry	51	Dog tooth violet	39
Corol root	98	Dog wood	19
Cotton	79	Dragoness plant	38
Cotton grass	11	Dragon head	67
Cotton thistle	88	Duck meat	100
Cowbane	33	Dwarf dandelion	87
Cowhage	82	Dwarf foxglove	71
Cow-parsley	32	Dyer's kneed grass	80
Cowslip	23	Dyer's weed	53
Cow-wheat	69	Dying sea weed	132
Crab-tree	56	Eatable mushroom	139
Cranberry	43	Elder black berried	34
Cranbill	77	red berried	34
Creeping cucumber	113	Elecampane	92
Crow-foot	63 & 77	Elephant foot	97
Crown-beard	95	Elm	31
Crown-cup	89	Enchanter's nightshade	9
Cuckold	89	Endive	87
Cucumber	113	English water-cress	76
Cudweed	90	Erinus	70
Currant	29	Eschallots	37
Custard apple	63	Evergreen snakeroot	79
Cut grass	14	Evergreen thorn	55
Cypress vine	25	Eye bright	69
Daffodil	38	False acacia	82
Daisy	91	False aloe	38
Dalibard	57	False asphodel	40
Dame's violet	74	False boneset	89
Dandelion	86	False chamomile	94
Darnel grass	15	False cowslip	22
Date plum	117	False foxglove	71
Day flower	10	False gentian	30
Day lily	38	False grape	28
Deadly night-shade	24	False gromwell	21
Dead nettle	65	False hawk weed	86

False heath	52	Flax	35
False hemp	117	Fleabane	92
False horehound	66	Floating arum	40
False indigo	80	Flower de luce	10
False mermaid	37	Fog fruit	70
False mustard	37	Follicle vine	29
False saffron	83	Fothergill's bush	60
False sanicle	49	Four o'clock	27
False sensitive plant	85	Fox glove	71
False sun-flower	94	Fox tail grass	13
False syringa	54	French marigold	91
False toad-flax	29	Frog's spittle	132
Feather leaf	23	Fumitory	79
Fennel	34	Fungus	
Fennel flower	61	barked	141
Fern		bead	142
adder tongue	124	birdnest	140
calyxed	123	cased	141
caterpillar	121	club	140
fine haired	122	dish	140
flowering	123	globule	141
fork	119	latticed	140
fringed	124	pink	140
grape	124	smooth cup	140
kidney fruit	121	soot	141
lip	123	tuberled	141
naked	124	veined	139
one sided	123	Galingale	11
ribbon	120	Garlic	37
Roman	120	Gayfeather	88
shield	122	Gentian	30
snuff-box	122	Giant fennel	32
Fescue grass	12	Ginseng	30
Fescue sedge	102	Globe flower	64
Feverfew	91	Goat's beard	86
Feverwort	27	Goat's rue	83
Field thyme	68	Golden rod	92
Fig tree	114	Golden saxifrage	49
Fig wort	70	Gold thread	62
Finger grass	2	Gourd	112
Five finger	57	Grain rust	141
Flat top	88	Grape vine	28

INDEX.

155

Grass

arrow	42
beard	13 & 16
blue	15
blue eyed	76
broom	16
cockfoot	12
cotton	11
cut	14
darnel	15
deer	44
dog tail	14
dyer's kneed	80
fescue	16
finger	12
foxtail	13
hair	14
hairy beard	15
hedgehog	12
knot	44
lime	14
meadow	15
orchard	14
panic	12
parnassus	35
pepper	73
quake	15
reed	16
ribbon	13
rush	40
salt	12
scorpion	21
scurvy	73
sea rush	15
sesame	102
soft	16
star	37
sweet vernal	10
tape	113
thin	14
timothy	14
whip	103

N 2

Grass

whitlow	72
yellow eyed	11
Grass pink	98
Greek valerian	24
Green brier	116
Green hair	132
Gromwell	21
Ground ivy	67
Ground nut	83
Ground pine	125
Hacmatack	110
Hair grass	14
Hairy beard grass	15
Hard-hack	56
Hazle nut	110
Heal all	68
Heart seed	44
Hedgehog grass	12
Hedge hyssop	8
Hedge mustard	75
Hellebore swamp	41
white	41
Helonias	41
Hemlock	33
Hemlock tree	111
Hemp	116
Henbane	24
Herb bennet	58
Herb robert	77
Hibiscus	78
Hickory	103
High heal all	71
High water shrub	96
Hog weed	106
Holly	20
Holly hock	78
Hone wort	32
Honey locust	117
Honey suckle	26
fly	26
trumpet	26

Honey suckle		Lamb lettuce	16
yellow	26	Larch	110
Hop	115	Larkspur	61
Hop horn beam	109	Laurel	47
Hore hound	67	broad leaved	47
Horn beam	109	narrow leaved	47
Horned poppy	60	Lavander	65
Horse balm	9	Leaf cup	96
Horse chesnut	42	Leaf flower	112
Horse radish	73	Leaf seed	78
Horse tail	125	Leather leaf	47
Hound tongue	21	Leather punk	142
House leek	54	Leather wood	44
Hyacinth	39	Leek	37
garden	39	Lemon	59
grape	39	Leopard's bane	92
Hydrangea	48	Less flowering rush	41
Hyssop	66	Lettuce garden	86
Ice plant	56	tall	86
Indian corn	101	white	86
Indian cucumber	41	Lichen button	134
Indian hemp	29	coral	136
Indian mallows	78	crust	135
Indian physic	56	efflorescent	133
Indian turnip	107	filamentous	138
Indigo	82	fungus	134
Itch weed	41	horned	138
Ivy	29	Iceland	138
Jacob's ladder	116	knobbed	139
Jasmine	8	leprous like	133
Jerusalem artichoke	95	lettered	134
Jewel weed	28	mitre	134
Job's tear	102	pit back	136
Jonquil	38	pitcher shield	136
Judas' tree	46	scarlet mouth	139
Juniper	118	seed ball	135
Knawell	49	shield	136
Knot grass	44	small pox	133
Knot spurge	53	spangled	134
Labrador tea	48	target	136
Ladies' slipper	99	Lichnedia	24
Ladies' traces	98	Life everlasting	90

INDEX.

157

Light hair	11	May weed	95
Lilac	8	Meadow grass	15
Lily	39	Meadow rue	62
Lily orchis	98	Medlar	55
Lime grass	14	Melilot clover	84
Limodore	97	Mermaid weed	17
Little snake weed	104	Mignonette	53
Live forever	51	Milk parsley	33
Liver leaf	63	Milk vetch	83
Liver wort	131	Milk way plant	83
brook	131	Milk weed	99
gaping	131	Milk willow herb	53
wax	131	Milk wort	79
Lizard tail	43	Millet	13
Locust tree	82	Mithridate mustard	73
Lombardy poplar	117	Mitre wort	49
Loose strife	22	Mock orange	54
Louse wort	71	Monkey flower	72
Louse wort fox glove	71	Monk's hood	61
Lovage	32	Moon seed	117
Love lies bleeding	106	Morel	140
Low calamint	67	Morning glory	23
Lucerne clover	83	Moss	
Lung wort	20	awl cup	126
Lupine	80	bow	129
Madder	18	creeping	127
Mad wort	73	curve haired	128
Magnolia	63	double tooth	127
Maiden hair	121	earth	126
Malabar nut	8	false	131
Mallows	78	fork	127
Maple	44	four toothed	126
Maple leaf	34	goose neck	129
Mare's tail	7	hair cap	127
Marjoram	68	hair mouth	127
Marsh five finger	58	leafless	130
Marsh hibiscus	78	many speckled	128
Marsh mallow	78	net tooth	128
Marsh mallows	78	peat	126
Marsh rosemary	36	pricked tooth	129
Marsh tway blade	98	rope	130
Master wort	33	sleeping	120

Moss		Oak of Jerusalem	36
spiral tooth	127	Oats	16
trait haired	128	Okra	78
thread	129	Onion	37
toothless	126	Orach	31
tree	128	Orange	59
umbrella	126	Orange root	62
water	130	Orchard grass	14
Mother of thyme	68	Orchis	97
Mother wort	66	round leaved	97
Mountain ash	55	scarlet	97
Mountain mint	9 & 65	Orpine	51
Mountain rice	12	Oswego tea	9
Mouse ear	90	Ox-eyed daisy	91
Mouse ear chickweed	55	Painted cup	69
Mulberry	105	Palma christi	111
Mullein	23	Panic grass	12
Mushroom	139	Paper punk	140
Musk melon	113	Parnassus grass	35
Mustard	76	Parsnip	34
Myrtle	54	Partridge berry	18
Nasturtion	44	Passion flower	76
Neclace weed	58	Pea	81
Nelumba	63	Peach	55
Nettle	104	Pea nut	81
Nettle tree	31	Pear	56
New-Jersey tea	27	Pearl wort	20
Nightshade	25	Peat moss	126
Nodding horehound	67	Pea vine	83
Oak	107	Pellitory	105
black	108	Pencil flower	85
pin	108	Penny royal	67
red	108	Penny wort	69
rock or chesnut	107	Peony	60
scarlet	108	Pepperidge	116
scrub	108	Pepper grass	73
swamp	107	Pepper mint	66
two coloured	108	Periwinkle	25
white	108	Persicaria	45
yellow	108	Pheasant's eye	64
Oak leaf foxglove	71	Phryma	68
Oak leather	142		

INDEX.

159

Pickereel weed	38	Rabbit foot	84
Pig weed	30	Radish	75
Pine	110	Ragged cup	96
pitch	110	Rag wort	93
white	110	Rape	73
yellow	110	Raspberry	57
Pine weed	35	Rattle box	80
Pink	49	Red clover	84
Pink root	23	Red pepper	26
Pin weed	17	Red top	13
Pipe wort	104	Reed grass	16
Plane tree	110	Reed mace	101
Plantain	17	Rhodora	46
Plantain sedge	103	Rhubarb	45
Plowman's wort	90	Ribbon grass	13
Plum	54	Rib wort	17
Poison hemlock	33	Rice	41
Poison ivy	35	River nymph	100
Poison sumach	34	Rocket	74
Poke weed	52	Rock rose	59
Polypod	119	Rose	56
Pond lily	60	Rose bay	47
Pond weed	20	Rose cockle	51
Poplar	117	Rosemary	9
balsam	117	Round head	104
high	117	Rue	46
lombardy	117	Rue anemone	62
white	117	Ruel	70
Poppoose root	36	Rush grass	40
Poppy	59	Rye	14
Potatoe	25	Sage	9
Prickly ash	115	Salt grass	12
Prickly pear	54	Salt wort	30
Prim	8	Samphire	7
Primrose	23	Sand wort	50
Puff ball	140	Sanicle	32
Pumpkin	112	Sassafras	45
Purslane	52	Satyrion	97
Pyramid flower	18	Savin	118
Quake grass	15	Saxifrage	48
Quick set	55	Scabish	43
Quince	56	Scarlet pimpernel	22

Scorpion grass	21	Spear mint	66
Scouring rush	125	Speedwell	9
Screw stem	18	Spice bush	45
Scurvy grass	73	Spicy winter green	48
Sea buckthorn	114	Spider wort	36
Sea burdock	105	Spikenard	35
Sea holly	31	Spinach	115
Sea milk-wort	29	Spindle tree	28
Sea pea	81	Spleen wort	120
Sea rocket	73	Spotted cranebill	77
Sea rush-grass	15	Spruce	111
Sea weed	132	single	111
Sedge	102	double	111
Seed box	19	Spurge	53
Self heal	68	knot	53
Seneka snake root	79	spotted	53
Sensitive plant	77	sun	53
Sensitive polypod	119	Squash	112
Service tree	55	flat	112
Sesame grass	102	warted	112
Shad flower	55	Staff tree	27
Sheep berry	34	St. Andrew's cross	58
Shepherd's purse	73	Star flower	93
Shin leaf	46	Star grass	37
Side saddle flower	59	Star of Bethlehem	40
Silk weed	99	Star wort	50
Silver cinquefoil	58	Steeple bush	56
Single seed cucumber	112	St. John's wort	61
Skull cap	67	Stock july flower	76
Skunk cabbage	19	Stone crop	51
Slippery elm	31	Stone seed	21
Snake head	72	Stork's bill	77
Snap dragon	70	St. Peter's wort	58
Snow ball	34	Strawberry	57
Snow drop	37	Succory	87
Soap wort	49	Succulent chick weed	17
Soft grass	16	Sugar cane	13
Solomon's seal	39	Sugar maple	44
Sorrel	42	Sumach	34
Sour gum	116	Summer corn	101
Southern wood	91	Summer savory	66
Sow thistle	87	Sundew	35

INDEX

161

Sunflower	95	True sarsaparilla	116
Swamp rose	56	Trumpet flower	70
Sweet brier	56	Tulip	39
Sweet cicely	33	Tulip tree	63
Sweet fern	103	Tupelo tree	116
Sweet flag	40	Turnip	75
Sweet gale	114	Turnsole	21
Sweet gum	110	Twin leaf	44
Sweet pea	81	Unicorn plant	72
Sweet pepper bush	46	Valerian	10
Sweet potatoe	23	Veiny hawk weed	87
Sweet vernal grass	10	Venus' fly trap	47
Sweet william	49	Venus' pride	18
Swine lettuce	37	Vervain	65
Syrian mallows	78	Vetch	31
Taliny	53	Violet	28
Tall blue bottle	96	Virginian loosestrife	43
Tansy	91	Virginian orpine	51
Tape grass	113	Virginian thyme	65
Tare	81	Virgin's bower	62
Teasel	19	Wake robin	42
Thin grass	14	Walking leaf	121
Thistle	88	Wall cress	74
Thorn apple	23	Walnut black	108
Thornbush	55	shagbark	108
Thorough wort	89	Walnut leaved ash	114
Threadfoot	101	Water arum	107
Three seed mercury	111	Water crowfoot	64
Thyme	68	Water hemp	115
Tick seed sunflower	96	Water horehound	9
Timothy grass	15	Water leaf	22
Toad flax	71	Water melon	112
Tobacco	23	Water milfoil	106
Tooth ache tree	115	Water navel wort	31
Tooth cup	19	Water nymph	100
Tooth root	74	Water oats	106
Touch me not	28	Water parsnip	32
Touch wood	140	Water pepper	45
Tower mustard	74	Water plantain	42
Trailing arbutus	48	Water purslane	20
Tree moss	138	Water radish	76
Tree primrose	43	Water shield	63

Water star wort	7	Wild onion	37
Weeping willow	113	Wild sarsaparilla	35
Wheat		Wild vetch	81
summer	15	Willow	113
winter	15	brittle joint	113
Whip grass	103	cone	113
White ash	114	creeping	114
White bush	47	weeping	114
White cedar	111	yellow	113
White clover	84	Willow herb	43
White lettuce	86	Wind flower	62
White radish	75	Winter berry	36
White snake root	52	Winter cherry	25
White wood	63	Winter cress	75
Whitlow grass	72	Winter green	46
Whortleberry	47	Witch hazel	20
Wild basil	65	Wood sage	64
Wild bush bean	83	Wood sorrel	51
Wild caraway	90	Wormseed	30
Wild chamomile	95	Wormwood	91
Wild feverfew	92	Yam root	116
Wild flag	10	Yarrow	95
Wild germander	64	Yellow cockscomb	69
Wild ginger	52	Yellow eyed grass	11
Wild honeysuckle	25	Yellow root	36
Wild indigo	46	Yellow rust	141
Wild iris	10	Yew	18
Wild liquorice	83	Zigadene	42
Wild mandrake	60		

ERRATA.

THE Address. For "Mason Fussell," read *Fris-sell*.

The Preface. For "Turton and Linneus," read *Turton's Linneus*.

On the 5th page, for "generic nouns," read *generic names*.

Correct the following by the help of the Index :

AGAVE. For "*virginca*," read *virginica*.

ALISMA. Over this genus erase "*POLYANDRIA*," and insert *POLYGYNIA*.

ANCHUSA. For $\frac{1}{2}$ read $\frac{1}{4}$.

ANDROPOGON. For "stainate," read *staminate*.

ATRIPLEX. Erase "(stem woody)" and insert (*stems woody*) over the genus *Ulmus*, as in page 26.

BROMUS *secalinus*. Read *calyx awnless*, &c.

CARUM. In the asterisked genera following this genus, set a period after *Scleranthus*.

CELASTRUS. For "climbing," read *twining*.

CITRUS, Var. *limon*. For "(lemon)" read (*lime*.)

CLAVARIA. For "cup-form," read *club-form*.

CORNUS *alba*. For $\frac{1}{4}$ read $\frac{1}{2}$. Over this genus insert, D. *Flowers 4-petalled, superior*.

DACTYLIS. This genus must be considered as following *Sorghum* on the next page.

DODECANTHEON. Read *DODECATHEON*.

ELYMUS. Strike out the words "4-leaved, 2-flowered" and "ternate."

GRATIOLA. For "Calyx 7-leaved, the 2 outer ones spreading:" read, *Calyx 5-leaved, sometimes with a 2-leaved calycle*.

INULA. For "*halenium*," read *helenium*.

LEPTANTHUS. Erase "3. 1." at the end of the generic description.

LONICERA *dioica*. For $\mathcal{L}$ read h .

MAGNOLIA. For $\mathcal{L}$ read h .

MARRUBIUM. Read MARRUBIUM.

PYXIDANTHERA. For h read $\mathcal{L}$.

RUBUS. For "juices," read *juicy*.

SMYRNIUM. In both species for h read $\mathcal{L}$.

SPIRAEA. For "(hand-hack)" read (*hard-hack*.)

THLASPI. For "roundish-cordate," read *roundish-obcordate*.

TRIFOLIUM. Read 1 to 4-seeded. And for "repens" read *repens*.

VERONICA. For h read $\mathcal{L}$.

ZAPANIA. In the section placed over this genus, for "Leaves," read *Calyxes*.

The reader will not be surprised at this list of errors, when he is told, that the author resides 34 miles from the printer, and never saw any of the proof sheets.

The alterations in the characters of the genera, *Elymus* and *Gratiola*, were caused by a perusal of the first numbers of that incomparable work, Elliot's Botany of South Carolina and Georgia.

New York Botanical Garden Library

QK110 .E18

Eaton, Amos/A manual of botany for the n gen

3 5185 00087 3511

Made in Italy

06-09 MIN

8 032919 990075

www.colibrisystem.com