4+x
1.7
10.3

MEMOIRS

OF THE

CARNEGIE MUSEUM.

VOL. VII.

NO. 3.

A CATALOG OF THE OPHIDIA FROM SOUTH AMERICA AT PRESENT
(JUNE, 1916) CONTAINED IN THE CARNEGIE MUSEUM
WITH DESCRIPTIONS OF SOME NEW SPECIES.

BY LAWRENCE EDMONDS GRIFFIN.

The recent reptilia in the Carnegie Museum represent collections made at various times and places by collectors, who incidentally in connection with other undertakings preserved such specimens belonging to this class as came to hand. No systematic collecting of these animals was undertaken by any of the expeditions which have gone out from the Carnegie Museum to the American tropics, except by Mr. G. A. Link, Sr., in the Isle of Pines. The collection made by Mr. Link has already been reported upon in the Annals of this Museum.¹ Although the collections have been acquired for the most part as the result of somewhat desultory collecting, they nevertheless include a considerable number of interesting species, some of which appear to be new to science.

Many specimens were preserved by Mr. John D. Haseman during his extensive travels in South America from 1907 to 1910. For a brief outline of the journeyings of Mr. Haseman the reader is referred to the articles published in the seventh volume of the Annals of the Carnegie Museum.² Mr. Haseman went to

¹ "The Reptiles and Amphibians of the Isle of Pines." By Thomas Barbour, Annals Carnegie Museum, Vol. X, 1916, pp. 297-308, Pl. XXVIII.

² W. J. Holland, "The Carnegie Museum Expedition to Central South America, 1907-1910," Annals of the Carnegie Museum, VII, 1910-1911, pp. 283-286.

John D. Haseman, "A Brief Report upon the Expedition of the Carnegie Museum to Central South America," *l. c.*, pp. 287-299.

C. H. Eigenmann, "A List of Localities at which Mr. Haseman Collected," *l. c.*, pp. 299-314.

South America primarily to collect fishes. Incidentally he preserved a considerable number of such reptilia as he encountered. The localities at which he obtained reptilia are given herewith in alphabetical order:

Arequa, Paraguay.

Asumpcion, Bolivia.

Bom Jesus de Lapa, Bahia, Brazil.

Cacequy (Rio Ibicuy into Uruguay) Rio Grande do Sul, Brazil.

Cidade de Matto Grosso, Brazil.

Entre Rios, Minas Geraes, Brazil.

Jacarehy, Rio Parahyba, São Paulo, Brazil.

Lagoa de João Pereira, near Barra, Rio São Francisco, Minas Geraes, Brazil.

Mogy das Cruces, Rio Tieté, São Paulo, Brazil.

Muniz Freire, Espiritu Santo, Brazil.

Penedo, Alagoas, Mouth of Rio São Francisco, Brazil.

Piracicaba, Rio Paraíba, São Paulo, Brazil.

Puerto Suarez, Eastern Bolivia (near Corumbá).

Rio Doce, Minas Geraes, Brazil.

Rio de Janeiro, Brazil.

Rio Mamoré, Bolivia, below the mouth of the Rio Guaporé.

Santarem, Brazil.

São Antonio de Guaporé, Matto Grosso, Brazil.

São Cruz, Campos de Matto Grosso, Brazil.

São João del Rey, Minas Geraes, Brazil.

São Luis de Cáceres, Rio Paraguay, Matto Grosso, Brazil.

São Matias, Bolivia.

Sete Lagoas (into Rio das Velhas, into Rio São Francisco), Minas Geraes, Brazil.

Urucum Mountains, Matto Grosso, Brazil.

Villa Bella, Bolivia.

Xiririca, Rio Ribeira de Iguape, São Paulo, Brazil.

Mr. and Mrs. H. H. Smith collected in the Province of Santa Marta, Colombia, from 1898 to 1901. The snakes contained in their collections were taken at Bonda (150 ft. alt.); Cacagualito (1,500 ft.); Minca (2,000 ft.); Masinga (2,000 ft.); Valparaiso (4,500 ft.); Las Nubes (4,500 ft.); El Libano (6,000 ft.).³ The labels in their collections usually give the month in which the specimen was captured, but very rarely the year. It is probable, however, that most of the reptiles of the Smith Collection were taken in the year 1901.

³ The localities at which Mr. and Mrs. H. H. Smith collected in Colombia are described in the *Annals of the Carnegie Museum*, Vol. VI, 1909, pp. 74-76.

The snakes cataloged by me under the Department Numbers 1841 to 1873 are without original data or field-labels, but they are positively known to have been received from South America, and are undoubtedly a part of the collections of H. H. Smith from northern Colombia.

The collections sent in by Señor Don José Steinbach were all taken in the Department of Santa Cruz de la Sierra, Provincia del Sara, Bolivia, except a small part of his sendings, which are labelled as from Puerto Suarez, Bolivia. Only two definite localities are given on his labels, Santa Cruz de la Sierra, and Las Juntas. The altitudes given range from 250 to 450 meters above sea-level.

A small collection of snakes made by Mr. Thomas LeBoutelier was given to the Museum in 1909 by Mrs. A. Marshall Bell of Pittsburgh. It is unfortunate that the maker of this collection was not careful in labelling the specimens. Most of them have no other locality than "South America" assigned to them, while in a few cases the locality given is manifestly incorrect.

Miss Lola Vance collected a few reptiles at Tarma, Peru. The altitude of this place is 6,000 feet.

A few serpents were also collected by Mr. J. O. Kerby at Massasão and Pranguina on the Amazon, and donated by him to the Carnegie Museum.

For the greater convenience of those who may refer to this paper I have arranged the species of each genus in alphabetical order, when more than one species is listed, and the genera of each family and subfamily are also arranged in like manner. In the statistical tables I have introduced some modifications of customary usage. Under "upper labials" the figure outside of the parenthesis indicates the total number of shields, while the figures within the parenthesis tell which supralabials are in contact with the eye. The formula for the temporals is written with a comma, instead of with the plus-sign. When there is a difference in the number or arrangement of the shields of the left and right sides, both are given, the formula for the left side being always written above that for the right side.

Class **REPTILIA.**

Order **SQUAMATA.**

SUBORDER **SERPENTES.**

Family **TYPHLOPIDÆ.**

Genus **Helminthophis** Peters.

1. **Helminthophis bondensis** sp. nov.

Rostral half the width of the head, rounded posteriorly, forming a broad suture with the frontal, extending to a line connecting the posterior margins of

the eyes; frontal broader than the rostral, but little longer than the diameter of the eye, in contact with the prefrontal and the ocular; prefrontal in contact with frontal, ocular, and subocular; no preoculars; a subocular; eye under the posterior and lower part of the ocular; four upper labials, the first the largest, second and third in contact with the subocular, the second also in contact with the upper nasal, which reaches the subocular. Twenty-two scale-rows around the body; diameter of body contained forty-five times in the length. Tail shorter than broad, ending in a point, but without spine.

Brown, each scale darker in the center; a little lighter beneath. Tip of head lighter with fine brown markings.

This species resembles *H. albirostris* Peters more closely than any other. It is distinguished by having one, instead of two suboculars, by the position of the eye under the ocular shield, and by the second upper labial being separated from the prefrontal by the nasal.

Type, ♂, No. 216, Carnegie Museum Catalog of Ophidia; Bonda, Colombia: H. H. Smith (coll.), May.

COUNTS AND MEASUREMENTS.

Scale-rows.....	22
Upper labials.....	4
Preoculars.....	0
Suboculars.....	1
Diameter of body.....	4 mm.
Total length.....	180 mm.
Length of tail.....	2 mm.

Genus *Typhlops* Oppel.

2. *Typhlops reticulata* (Linnæus).

Anguis reticulata LINNÆUS, Syst. Nat., Ed. XII, I, 1766, p. 391.

Typhlops reticulatus BOULENGER, Cat. Snakes, I, 1893, p. 27.

The collection contains two specimens of this species:

No. 17, ♂, Province del Sara, Bolivia, Elev. 350 M., Steinbach coll., Feb., 1911.

No. 334, ♀, Villa Bella, Bolivia, Haseman coll., Oct. 8, 1909.

COUNTS AND MEASUREMENTS.

	(No. 17.)	(No. 334.)
Scale-rows.....	20	20
Upper labials.....	4	4
Preoculars.....	1	1
Total length in mm.....	257	334
Length of tail in mm.....	8.5	4
Diameter in mm.....	8	8

Family BOIDÆ Gray.

Genus *Boa*⁴ Linnæus.3. *Boa cooki* (Gray).*Corallus cookii* GRAY, Zool. Misc., 1842, p. 42.*Corallus cookii* BOULENGER, Cat. Snakes, I, 1893, p. 99.

The five specimens in the collection are listed as follows:

No. 137, ♂, Bonda, Colombia, Elev. 150 ft., Smith coll., June.

No. 139, ♂, Bonda, Colombia, Elev. 150 ft., Smith coll., Sept. 6 (in house).

No. 142, ♂, Bonda, Colombia, Elev. 150 ft., Smith coll., June.

No. 204, ♀, Bonda, Colombia, Elev. 150 ft., Smith coll., July.

No. 1860, ♂, South America (Colombia, H. H. Smith).

COUNTS AND MEASUREMENTS.

	(No. 137.)	(No. 139.)	(No. 142.)	(No. 204.)	(No. 1860.)
Anal.	1	1	1	1	1
Scale-rows.	43	39	43	42	41
Gastrosteges.	270	269	270	272	271
Urosteges.	116	114	114	115	114
Total length in mm.	904	1,596	1,395	567	593
Length of tail in mm.	170	328	290	112	110

4. *Boa hortulana* (Linnæus).*Coluber hortulanus* LINNÆUS, Mus. Ad. Frid., 1754, p. 37.*Boa hortulana* LINNÆUS, Syst. Nat., Ed. XII, 1766, I, p. 374.*Corallus hortulanus* BOULENGER, Cat. Snakes, I, 1893, p. 101.

The two specimens in our possession are the following:

No. 370, ♂, Entre Rios, Brazil, Haseman coll., June 4, 1908.

No. 371, Entre Rios, Brazil, Haseman coll., June 4, 1908 (head only).

COUNTS AND MEASUREMENTS.

	(No. 370)
Anal.	1
Scale-rows.	53
Gastrosteges.	266
Urosteges.	109
Total length in mm.	1,616
Length of tail in mm.	300

⁴ See Stejneger, Proc. U. S. Nat. Mus., XXIV, 1902, p. 184.

Genus *Constrictor* Laurenti.5. *Constrictor constrictor* (Linnaeus).*Boa constrictor* LINNÆUS, Syst. Nat., Ed. XII, 1766, I, p. 373.*Boa constrictor* BOULENGER, Cat. Snakes, I, 1893, p. 117.

The specimens in the collection of the Carnegie Museum are listed as follows:

No. 238, ♂, South America, LeBoutelier Collection.

No. 239, South America, LeBoutelier Collection (head only).

No. 240, South America, LeBoutelier Collection (head only).

No. 241, ♂, South America, LeBoutelier Collection.

No. 242, South America, LeBoutelier Collection (head only).

No. 243, ♂, South America, LeBoutelier Collection.

No. 1782, ♀, Bonda, Columbia, H. H. Smith coll., (Skin).

No. 1859, ♂, South America, J. H. Smith coll.

No. 2022, ♂, Cacagualito, Colombia, Mrs. H. H. Smith.

COUNTS AND MEASUREMENTS.

	(No. 238.)	(No. 241.)	(No. 243.)	(No. 1782.)	(No. 1859.)	(No. 2022.)
Anal.	1	1	1	1	1	1
Scale-rows.	95	85	85	86	85	88
Gastroteges.	237	233	243	235	239	246
Urostege.	53	60	54	51	62	57
Total length in mm.	542	514	576	1,000	578	727
Length of tail in mm.	55	56	60	175	64	90

Genus *Epicrates* Wagler.6. *Epicrates cenchria* (Linnaeus).*Boa cenchria* LINNÆUS, Mus. Ad. Frid., II, 1764, p. 41; Syst. Nat. Ed. XII, 1766, I, p. 374.*Epicrates cenchris* BOULENGER, Cat. Snakes, I, 1893, p. 94.

No. 26, ♂, Las Juntas, Bolivia, Elev. 50 M., José Steinbach coll., Dec. 1913.

No. 115, ♂, Santa Cruz de la Sierra, José Steinbach coll.

No. 136, ♂, Minca, Colombia, Elev. 2,000 ft., H. H. Smith coll., June.

No. 320, ♂, Urucum Mts., Matto Grosso, Brazil, Haseman coll., May 2, 1909.

COUNTS AND MEASUREMENTS.

	(No. 26.)	(No. 115.)	(No. 136.)	(No. 320.)
Anal.	1	1	1	1
Scale-rows.	46	43	46	44
Gastroteges.	257	233	243	234
Urostege.	56	46	53	51
Total length in mm.	593	1,154	390	557
Length of tail in mm.	65	103	43	58

6a. *Epicrates cenchria* var. *fusca* (Gray).

Cliftia fusca GRAY, Catalogue, 1849, p. 99.

Epicrates cenchris, var. B, BOULENGER, Cat. Snakes, I, 1893, p. 96.

No.	Sex.	Locality.	Elevation.	Collector.	Anal.	Scale-rows.	Gastro-steges.	Uro-steges.	Total Length in Min.	Length Tail in Min.
1861		S. America.....		H. H. Smith	1	50	247	50		
2025	♂	Cacagualito, Colombia...	1,500 ft.	Mrs. H. H. Smith	1	49	247	49	1,790	160

Genus *Eunectes* Wagler.

7. *Eunectes murinus* (Linnaeus).

Boa murina LINNÆUS, Syst. Nat., Ed. XII, 1766, I, p. 374.

Eunectes murinus BOULENGER, Cat. Snakes, I, 1893, p. 115.

No.	Sex.	Locality.	Collector.	Anal.	Scale-rows.	Gastro-steges.	Uro-steges.	Upper Labials.	Preoculars.	Postoculars.	Total Length in Min.	Length Tail in Min.
217		S. America	LeBoutelier	1	54	239	69	17	1	2	725	98
218		"	"	1	60		71	17	1	3	685	98
219		"	"	1	58	248	63	16	1	2	700	89
220		"	"	1	63	247	68	16	1	4	687	95
221		"	"	1	62		58	15	1	3	590	74
222		"	"	1	62	248	66	17	1	3	570	76
223		"	"	1	54	241	67	16	1	2	600	82
224		"	"	1	58		65	15	1	3	707	101
225		"	"	1	54	241	67	16	1	3	723	102
226		"	"	1	58	240	68	17	1	4	617	83
228		"	"	1	65	247	69	16	1	3	728	101
229		"	"	1	58	244	70	16	1	3	728	100
230		"	"	1	62	242	70	17	1	3	700	97
231		"	"	1	56	254	64	16	1	2	725	97
232		"	"	1	57	246	67	17	1	3	620	77
233		"	"	1	66	255	67	16	1	2	716	100
234		"	"	1	60	244	67	16	1	4	698	99
235		"	"	1	62	246	68	18	1	2	702	101
236		"	"	1	60	242	69	15	1	2	720	105
237	♀	Chile (?)	"	1	60	245	69	17	1	3	706	91

8. *Eunectes notæus* Cope.

Eunectes notæus COPE, Proc. Acad. Nat. Sci. Philada., 1862, p. 70.

Eunectes notæus BOULENGER, Cat. Snakes, III, 1896, p. 594.

The only specimen of this species at present in the collection (No. 379) was taken by Mr. John D. Haseman near Santarem, in a swamp between the Rio Amazonas and Tapajos, on Dec. 7, 1909. When captured the snake was found to have a bird in its stomach. It is a female.

COUNTS AND MEASUREMENTS.

Anal.....	1
Scale-rows.....	48
Gastrosteges.....	227
Urosteges.....	49+
Upper labials.....	14
Preoculars.....	1
Postoculars.....	3
Total length in mm.....	1,280
Length of tail in mm.....	143

Family COLUBRIDÆ Boulenger.

SERIES A. AGLYPHÆ.

Subfamily COLUBRINÆ Boulenger.

Genus *Aporophis* Cope.9. *Aporophis flavifrenatus* (Cope).

Lygophis flavifrenatus COPE, Proc. Acad. Nat. Sci. Philada., 1862, p. 80.

Aporophis flavifrenatus BOULENGER, Cat. Snakes, II, 1894, p. 158.

This species is represented in the collection by a single individual (No. 363) taken by J. D. Haseman on February 1, 1909, at Cacequy, Rio Grande do Sul. It is a female.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	17
Gastrosteges.....	156
Urosteges.....	77/77
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	638
Length of tail in mm.....	162

10. *Aporophis lineatus* (Linnæus).

Coluber lineatus LINNÆUS, Mus. Ad. Frid., 1754, p. 30, Pl. XII, fig. 1, and Pl. XX, fig. 1; Syst. Nat., Ed. XII, 1766, I, p. 582.

Aporophis lineatus BOULENGER, Cat. Snakes, II, 1894, p. 158.

There are at hand two specimens of this species, one (No. 5) taken in the Province del Sara, Bolivia, by José Steinbach in January, 1912, the other (No. 346) taken by Haseman at Bom Jesus de Lapa, Bahia, Brazil. Both specimens are females.

COUNTS AND MEASUREMENTS.

	(No. 5.)	(No. 346.)
Anal.	1/1	1/1
Scale-rows.	19	19
Gastrosteges.	165	171
Urosteges.	77/77	71/71
Upper labials.	8(4.5)	8(4.5)
Preoculars.	1	1
Postoculars.	2	2
Temporals.	1, 2	1, 2
Total length in mm.	675	300
Length of tail in mm.	178	68

11. *Aporophis melanocephalus* sp. nov.

Maxillary teeth about twenty, followed after a considerable interspace by two enlarged, compressed teeth below the posterior border of the eye; mandibular teeth subequal. Head narrow, not much wider than neck; snout narrow, high; eye large, pupil round. Body cylindrical, ventrals rounded; scales smooth, without pits, in fifteen rows. Rostral broader than deep, visible from above; internasals as long as wide, almost as long as the prefrontals; frontal once and a half as long as broad, sides straight and nearly parallel, longer than its distance from the end of the snout, considerably shorter than the parietals; nostril between two nasals; loreal considerably deeper than long; one preocular; two postoculars; temporals 1, 2; eight upper labials, the fourth and fifth bordering the eye; five lower labials in contact with the anterior chin-shields, which are as long as the posterior.

Back and sides of body and tail uniform dark brown; each scale has a dark brown center and light brown edge. The upper surface of the head is nearly black, this color shading into the brown of the body on the neck. A black vertebral stripe five scale-rows wide joins the black of the occiput. A black lateral stripe partly covering the second and third rows of scales extends from the temples along the neck. These three stripes quickly merge into the brown body color. Close to the head they are separated by white lines one scale-row wide, which become darker as they are followed caudad, and also soon merge into the color of the body. There is a small white spot back of the eye on the outer part of the parietal, and a similar spot in front of the eye on the lateral part of the prefrontal. The upper lip and ventral surfaces of the head and body are white, without dark markings.

The type, No. 18, of the Catalog of the Reptilia in the Carnegie Museum, is a female, taken at Las Juntas, Bolivia, 250 M. above sea-level, by José Steinbach in December, 1913. It is unique.

COUNTS AND MEASUREMENTS.

Anal.	1/1
Scale-rows.	15
Gastrosteges.	144
Urosteges.	72/72
Upper labials.	8(4.5)
Preoculars.	1
Postoculars.	2
Temporals.	1, 2
Total length in mm.	291
Length of tail in mm.	47

Genus *Atractus* Wagler.12. *Atractus badius* (Boie).

Brachyorrhos badius BOIE, Isis, 1827, p. 540.

Atractus badius BOULENGER, Cat. Snakes, II, 1894, p. 308.

The two specimens at hand were taken by H. H. Smith, one, No. 201, a male, at Minca at an elevation of 2,000 ft. above sea-level, the other, No. 215, a female, at El Libano at an elevation of 6,000 ft. Both localities are in the Province of Santa Marta, Colombia. The female specimen was taken in May, the male in July. The latter specimen has lost the tip of the tail.

COUNTS AND MEASUREMENTS.

	(No. 201.)	(No. 215.)
Anal.	1	1
Scale-rows.	17	17
Gastrosteges.	152	146
Urosteges.	30/30+	28/28
Upper labials.	7(3.4)	7(3.4)
Preoculars.	0	0
Postoculars.	2	2
Temporals.	1, 2	1, 2
Total length in mm.	270+	442
Length of tail in mm.	32	49

No. 201 is uniform dark brown above, with about forty pairs of small light spots making a row on either side of the back. The belly is nearly covered with dark brown, square, or oblong, spots. The frontal shield is longer than wide.

No. 215 is reddish brown above, with transverse black spots which show a tendency to produce bars, but are too irregular to actually do so. Each scale of the ventral surface carries two lateral and one or two median large, black spots, the whole forming two definite lateral stripes and a less definite median one.

13. *Atractus tæniatus*⁵ sp. nov. (Plate XXVIII, figs. 1-3.)

Eight maxillary teeth. Symphysials separated from the chin-shields by the first lower labials. Snout broad, rounded; rostral considerably broader than deep, the part visible from above as long as the suture between the internasals; internasals very small, pentagonal, the combined width of the internasals and posterior nasals (with which the internasals are in contact) being just equal to the width of the prefrontals; prefrontals as broad as long, entering the orbit; frontal once and a quarter as broad as long, longer than the prefrontals, as long as its distance from the end of the snout, much shorter than the parietals; loreal twice as long as deep, entering the orbit; two postoculars, the lower very small, both in contact with the anterior end of the parietal; temporals narrow, 1, 2; six upper labials, the third and fourth entering the orbit, the third to the last once and one-half to twice as long as deep; four lower labials in contact with each of the single pair of chin-shields, which are twice as long as broad.

Dark brown above, with a darker vertebral stripe; a light bar across the temporals and parietals; edge of upper lip light (white?); white below, chin spotted with brown, and a large brown spot occupying the center of each gastrostege, forming a midventral brown stripe; urosteges irregularly spotted with brown.

This species resembles *A. roulei* Despax; the two are compared below.

A. tæniatus.

Rostral wider than deep, portion visible from above considerable.
Internasals pentagonal.
Prefrontals as long as broad, shorter than the frontal.
Frontal once and a quarter as broad as long.
Loreal twice as long as deep.
Two postoculars.
Pupil round.
Four lower labials in contact with chin-shields.
Six upper labials.
Scales in fifteen rows.
Light band across head.
Urosteges lightly spotted with brown.
Scale formula, 1-15-152-24/24

A. roulei.

Rostral as wide as deep, little visible from above.
Internasals subtriangular.
Prefrontals almost as wide as long, as long as the frontal.
Frontal once and a half as broad as long.
Loreal twice as long as deep.
One postocular.
Pupil subelliptic.
Three lower labials in contact with chin-shields.
Six upper labials.
Scales in fifteen rows.
No light band across head.
Urosteges uniformly brown.
Scale formula, 1-15-154-22/22

The type, which is unique, is No. 117 of the Catalog of Reptiles in the Carnegie

⁵ *ravio* = 3 fillet, or head-band.

Museum. It was taken by José Steinbach near Santa Cruz de la Sierra, Bolivia. The collector failed on his label to indicate the date of capture. It is a male.

COUNTS AND MEASUREMENTS.

Anal.....	1
Scale-rows.....	15
Gastrosteges.....	152
Urosteges.....	24/24
Upper labials.....	6
Preoculars.....	0
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	218
Length of tail in mm.....	22

Genus *Elaphe* Fitzinger.14. *Elaphe corais* (Boie).

Coluber corais BOIE, Isis, 1827, p. 537.

Coluber corais BOULENGER, Cat. Snakes, II, 1894, p. 31.

The Carnegie Museum possesses three examples of this species (Nos. 143, 144, and 149) all of which were captured by H. H. Smith, in the month of June at Bonda, Province of Santa Marta, Colombia. No. 144 is a female, No. 149 is a male; the sex of No. 143 I have been unable to determine.

COUNTS AND MEASUREMENTS.

	(No. 143.)	(No. 144.)	(No. 149.)
Anal.....	1	1	1
Scale-rows.....	17	17	17
Gastrosteges.....	212	214	203
Urosteges.....	81/81	82/82	88/88
Upper labials.....	8(4.5)	9(5.6)	8(4.5)
Preoculars.....	1	1	1
Postoculars.....	2	2	2
Temporals.....	2, 2	2, 2	2, 2
Total length in mm.....	1,762	1,582	1,832
Length of tail in mm.....	375	353	422

15. *Elaphe dichroa* (Peters).

Herpetodryas dichroa PETERS, Mon. Berl. Akad., 1863, p. 284.

Coluber dichrous BOULENGER, Cat. Snakes, II, 1894, p. 30.

The only example of this species at present in the Museum is No. 329, which was taken by J. D. Haseman at Entre Rios, Brazil, on June 4, 1908. It is a female.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	15
Gastrosteges.....	181
Urosteges.....	96/96
Upper labials.....	8(3.4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 2
Total length in mm.....	402
Length of tail in mm.....	100

Genus *Dimades* Gray.16. *Dimades plicatilis* (Linnæus).

Coluber plicatilis LINNÆUS, Mus. Ad. Frid., 1754, p. 23, Pl. VI, fig. 1; Syst. Nat., Ed. XII, 1766, I, p. 376.

Dimades plicatilis BOULENGER, Cat. Snakes, II, 1894, p. 186.

The single specimen we have is No. 378 of the Catalog of Reptiles in the Carnegie Museum. It was taken by J. D. Haseman at Santarem, Brazil, on December 7, 1909. It is a female, and was captured under a log in the forested lowlands between the Amazon and Tapajos rivers in the act of brooding forty-eight eggs, five of which together with one of the unhatched young were collected by Mr. Haseman and are preserved as C. M. No. 354. Mr. Haseman states that "Sucury" and "Sucurujaba" are the local names of this serpent.

The shells of the eggs are membranous, thin, and white. The eggs measure 35 mm. in length and 25 mm. in width. Considering the advanced stage of development of the young, there is still a large amount of yolk in the eggs. These eggs seem to have passed through about half of the period of incubation. The unhatched young is 160 mm. long; the length of its tail is 35 mm. The central portion of the back, three scales wide, is light brown; on each side is a dark brown stripe covering the third scale row and the contiguous half of the scale row on either side. A row of small dark dots lies on each side of the vertebral stripe. The intervals between the dorsal and lateral bands, and the ventral surface, are white. The adult female has well-defined lateral dark stripes on the same scale rows as the young, but the dorsal surface between these is uniformly colored except for the two rows of small dark brown spots. The ventral surface of the adult bears four rows of large brown dots.

COUNTS AND MEASUREMENTS.

Anal	1/1
Scale-rows	15
Gastrosteges	139
Urosteges	37/37
Upper labials	7(3)
	8(3.4)
Preoculars	1
Postoculars	2
Temporals	1, 1
Total length in mm.	1,373
Length of tail in mm.	194

Genus *Coronella* Laurenti.17. *Coronella micropholis* (Cope).

Lampropeltis micropholis COPE, Proc. Acad. Nat. Sci. Philada., 1860, p. 257.

Coronella micropholis BOULENGER, Cat. Snakes, II, 1894, p. 203.

The specimen, which is a female (No. 2036), was taken by Mrs. H. H. Smith at Cacagualito, Colombia.

COUNTS AND MEASUREMENTS.

Anal	1
Scale-rows	23
Gastrosteges	228
Urosteges	43/43
Upper labials	7(3.4)
Preoculars	1
Postoculars	2
Temporals	2, 2
Total length in mm.	732
Length of tail in mm.	128

Genus *Drymobius* Cope.18. *Drymobius bifossatus* (Raddi).

Coluber bifossatus RADDI, Mem. Soc. Ital. Modena, XVIII, (Fis.) 1820, p. 333.

Drymobius bifossatus BOULENGER, Cat. Snakes, II, 1894, p. 10.

The collection possesses two specimens:

No. 127, ♀. From Santa Cruz de la Sierra, Bolivia. (Steinbach coll.)

No. 381, ♂. From Rio de Janeiro, Brazil. (Haseman coll., July 1, 1908.)

The color of the skin of No. 127 is almost uniform.

COUNTS AND MEASUREMENTS.

	(No. 127.)	(No. 381.)
Anal.	1/1	1/1
Scale-rows.	15	15
Gastrosteges.	179	167
Urosteges.		97/97
Upper labials.	8(4.5)	8(4.5)
Preoculars.	1	1
Postoculars.	2	2
Temporals.	2, 2	2, 2
Total length in mm.		1,638
Length of tail in mm.		455

19. *Drymobius boddaerti* (Santzen).

Coluber boddaerti SANTZEN, Meyer's Zoöl. Archiv, II, 1796, p. 59.

Drymobius boddaerti BOULENGER, Cat. Snakes, II, 1894, p. 11.

There are twenty specimens in the collection, all of which are presumed to have come from Colombia, and to have been collected by Mr. and Mrs. H. H. Smith, except one (No. 29) which was taken by José Steinbach at Las Juntas, Bolivia. Nos. 176-183 inc., and No. 186 are from Bonda, where they were captured by the Smiths in 1901 at dates ranging from May to September. Nos. 202-203 were taken by the same parties at Valparaiso, Colombia (elev. 4,500 ft.); Nos. 2006, 2009, and 2020 at Cacagualito, Colombia (elev., 1,500 ft.) by Mrs. H. H. Smith. Nos. 1866-1868 inclusive, and Nos. 1872-1873 are a part of a collection, which, as stated in the introductory pages of this paper, had no field data attached to it, but are known to have been sent in by Mr. H. H. Smith while laboring in Colombia.

Both anterior and posterior temporals are frequently divided into two or three scales in longitudinal series.

Nos. 202 and 203 are young specimens of the variety *rappii*. The prevailing color of the upper surface is dark brown. Numerous white bars cross the back, alternating with a series of shorter white bars on each side which reach the ventral surface. The labials are black-margined; the chin and throat are checkered with black and white. No. 2006 is uniform olive-green above, shading to brown on the tail; the lower surface is white, with faint cloudy markings on the chin and throat. No. 2020 is brown above, with faintly defined darker stripes along the middle of the back and each side. The vertebral band is outlined by a black line in the posterior part of the body. The lateral stripes become darker and more distinct on the tail.

COUNTS AND MEASUREMENTS.

Numbr.	Anal.	Scale-rows.	Gastrosteges.	Urosteges.	Upper Labials.	Preoculars.	Postoculars.	Temporals.	Total Length in Mm.	Length Tail in Mm.	Sex.
No. 176.....	1/1	17	198	120/120	9(4.5.6)	1	2	2, 2	1,364	402	♀
" 177.....	1/1	17	172	97/97	9(4.5.6)	1	2	2, 2	916	268	
" 178.....	1/1	17	189	96/96	9(4.5.6)	1	2	2, 2	1,140	305	♀
" 179.....	1/1	17	191	86/86	9(4.5.6)	1	2	2, 2	1,180	288	♀
" 180.....	1/1	17	188	94/94	9(4.5.6)	1	2	2, 2	1,163	307*	♀
" 181.....	1/1	17	174	100/100	9(5.6)						
" 182.....	1/1	17	175	95/95	9(4.5.6)	1	2	2, 2	925	265	
" 183.....	1/1	17	188	118/118	9(4.5.6)	1	2	2, 2	1,023	287	♂
" 186.....	1/1	17	186	88/88	9(4.5.6)	1	2	2, 2	1,222	367	♂
" 29.....	1/1	17	193	53/53	9(4.5.6)	1	2	2, 2	975	242	♀
" 202.....	1/1	17	179	55/55	9(4.5.6)	1	2	2, 2	802	137*	♀
" 203.....	1/1	17	194	118/118	9(4.5.6)	1	2	2, 2	384	67*	♂
" 1866.....	1/1	17	180	82/82	9(4.5.6)	1	2	2, 2	505	152	♀
" 1867.....	1/1	17	187	94/94	9(4.5.6)	1	2	2, 2	985	240	♀
" 1868.....	1/1	17	188	94/94	9(4.5.6)	1	2	2, 2	1,005	266	♀
" 1872.....	1/1	17	198	121/121	9(4.5.6)	1	2	2, 2	1,030	307	♀
" 1873.....	1/1	17	196	119/119	9(4.5.6)	1	2	2, 2	1,133	343	♀
" 2006.....	1/1	17	189	118/118	9(4.5.6)	1	2	2, 2	1,270	372	♂
" 2009.....	1/1	17	189		9(4.5.6)	1	2	2, 2	826	250	♂
" 2020.....	1/1	17	185	93/93	9(4.5.6)	1	2	2, 2	1,020	93*	♀
									1,110	287	♀

* Tip of tail lost.

20. *Drymobius rhombifer* (Günther).*Coryphodon rhombifer* GUENTHER, P. Z. S., 1860, p. 236.*Drymobius rhombifer* BOULENGER, Cat. Snakes, II, 1894, p. 14.

The collection contains four specimens as follows:

No. 188, ♀, Minca, Colombia, 2,000 ft. (H. H. Smith coll.), July. Skin.

No. 213, ♀, Bonda, Colombia, 150 ft. (H. H. Smith coll.), June.

No. 1865, ♀, Colombia, S. America (H. H. Smith coll.).

No. 1871, ♂, Colombia, S. America (H. H. Smith coll.). Tip of tail lost.

No. 213 has, in addition to the ordinary markings, a brown spot in the center of each gastrostegae, all together making a midventral stripe. Each of these specimens possesses a greater number of urosteges than is given for the species in Boulenger's Catalog, *i. e.* 84 to 96.

COUNTS AND MEASUREMENTS.

Number.	Anal.	Scale-rows.	Gastrosteges.	Urosteges.	Upper Labials.	Preoculars.	Postoculars.	Temporals.	Total Length in Mm.	Length Tail in Mm.
188	1/1	17	151	100/100	9(4.5.6)	1	2	2, 2		
213	1/1	17	160	102/102	9(4.5.6)	1	2	2, 2	474	130
1865	1/1	17	157	100/100	9(4.5.6)	1	2	2, 2	375	103
1871	1/1	17	152	97/97*	9(4.5.6)	1	2	2, 2	1,272	385

* Tip of tail lost.

Genus *Helicops* Wagler.21. *Helicops angulata* (Linnæus).

Coluber angulatus LINNÆUS, Mus. Ad. Frid., 1754, p. 23, Pl. XV, fig. 1; Syst. Nat., Ed. XII, 1766, I, p. 377.

Helicops angulatus BOULENGER, Cat. Snakes, I, 1893, p. 278.

The only specimen in the Museum (No. 279) is a male from the LeBoutelier collection, which simply bears the label "South America."

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	19
Gastrosteges.....	117
Urosteges.....	74/74
Upper labials.....	9(4)
	8(4)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 4
Total length in mm.....	300
Length of tail in mm.....	88

22. *Helicops carinicauda* (Wied) var. *infratæniata* Jan.

Coluber carinicaudus WIED, Beitr. Naturgesch. Bras., I, 1825, p. 300, Plate.

Helicops infratæniatus JAN, Arch. Zool. Anat. Phys., III, 1865, p. 245 and Icon.

Gén., 28, Pl. III, fig. 3 (1868).

Helicops carinicauda, var. B, BOULENGER, Cat. Snakes, I, 1893, p. 277.

No. 355, ♂, Cacequy, Rio Grande do Sul (Haseman coll.), Jan. 31, 1909.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	19
Gastrosteges.....	128
Urosteges.....	61/61
Upper labials.....	8(3.4)
	8(4)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	543
Length of tail in mm.....	128

23. *Helicops leopardina* (Schlegel).*Homalopsis leopardina* SCHLEGEL, Phys. Serp., II, 1837, p. 358.*Helicops leopardinus* BOULENGER, Cat. Snakes, I, 1893, p. 278.

The Museum possesses eleven examples of this species, as follows:

- No. 25, ♂. Las Juntas, Bolivia, coll. J. Steinbach, December, 1913.
 No. 321, ♂. Puerto Suarez, Bolivia, coll. J. D. Haseman, May 7, 1909.
 No. 322, ♀. São Antonio de Guaporé, coll. J. D. Haseman, July 4, 1909.
 No. 323, ♂. Puerto Suarez, Bolivia, coll. J. Steinbach, Jan., 1909.
 No. 324, ♂. Aregua, Paraguay, coll. J. D. Haseman, April 7, 1909.
 No. 325, ♂. Puerto Suarez, Bolivia, coll. J. D. Haseman, May 6, 1909.
 No. 326, ♂. Puerto Suarez, Bolivia, coll. J. D. Haseman, May 6, 1909.
 No. 327, ♀. São Luiz de Cáceres, coll. J. D. Haseman, May 25, 1909.
 No. 338, ♂. Santarem, Brazil, coll. J. D. Haseman, Dec. 7, 1909.
 No. 357, ♂. Santarem, Brazil, coll. J. D. Haseman, December 15, 1909.
 No. 358, ♂. Santarem, Brazil, coll. J. D. Haseman, December 15, 1909.

The hypapophyses of the posterior trunk vertebræ of *H. leopardina* are variably developed. Some specimens show scarcely a keel, some a well-developed plate, some an intermediate condition. No. 338 has no loreal shields; the loreal is absent on the left side of No. 357, and the right nasal of this specimen is in contact with the preocular.

COUNTS AND MEASUREMENTS.

Number.	Anal.	Scale-rows.	Gastro-stages.	Uro-stages.	Upper Labials.	Pre-oculars.	Post-oculars.	Temporals.	Total Length in Mm.	Length Tail in Mm.
No. 321.....	1/1	19	114	74/74	8(4)	1	2	{2, 2 (1, 2)	430	136
" 322.....	1/1	19	115	62/62	8(3.4)	1	2	1, 2	245	65
" 323.....	1/1	19	112	73/73	8(4)	2	2	1, 2	420	129
" 324.....	1/1	19	119	64/64	8(4)	{1 (2)	2	{2, 2 (1, 2)	196	48
" 325.....	1/1	19	116	68/68	8(4)	1	2	2, 2	295	82
" 326.....	1/1	21	112	64/64	8(4)	1	2	2, 2	261	72
" 327.....	1/1	19	119	72/72	8(4)	1	2	{2, 2 (1, 2)	365	97
" 338.....	1/1	19	120	87/87	8(3.4)	1	2	{2, 3 (1, 2)	294	95
" 357.....	1/1	19	117	89/89	8(4)	{1 (2)	2	1, 2	568	201
" 358.....	1/1	19	115	72/72	8(4)	{1 (2)	2	{2, 2 (1, 2)	354	105
" 25.....	1/1	19	122	83/83	8(4)	1	2	1, 2	300	96

24. *Helicops modesta* Günther.

Helicops modestus GUENTHER, Ann. & Mag. Nat. Hist. (3), VII, 1861, p. 425.

Helicops modestus BOULENGER, Cat. Snakes, I, 1893, p. 277.

The specimens at hand are listed as follows:

No. 335, ♂, collected by J. D. Haseman at Jacarehy, São Paulo, Brazil, July 15, 1908.

No. 336, ♂, collected by J. D. Haseman at Jacarehy, São Paulo, Brazil, July 15, 1908.

No. 348, ♀, collected by J. D. Haseman at Mogy das Cruzes, São Paulo, July 20, 1908.

No. 349, ♂, collected by J. D. Haseman at Mogy das Cruzes, São Paulo, July 20, 1908.

Nos. 348 and 349 are uniform black above, with no traces of stripes. The scales of the anterior half of the body scarcely show traces of keels, which are poorly developed in the posterior half. These two specimens were captured with a seine in the Rio Tieté. Nos. 335 and 336 are reddish brown above, with three darker stripes. They were taken in a pond near the Rio Parahyba.

No. 348 has no traces of hypapophyses on the posterior trunk vertebræ; the hypapophyses of these vertebræ of No. 349 are only low keels.

COUNTS AND MEASUREMENTS.

	(No. 335.)	(No. 336.)	(No. 348.)	(No. 349.)
Anal.	1/1	1/1	1/1	1/1
Scale-rows.	19	19	19	19
Gastrosteges.	121	125	120	125
Urosteges.	70/70	54/54	59/59	55/55
Upper labials.	8(3.4)	8(4)	8(4)	8(4)
Preoculars.	1	1	1	{ 1
				{ 2
Postoculars.	2	2	2	2
Temporals.	{ 2, 2	2, 2	2, 2	{ 2, 3
	{ 2, 3			{ 2, 2
Total length in mm.	381	285	316	587
Length of tail in mm.	111	65	73	121

25. *Helicops polylepis* Günther.

Helicops polylepis GUENTHER, Ann. Mag. N. H. (3), VII, 1861, p. 426.

Helicops polylepis BOULENGER, Cat. Snakes, I, 1893, p. 280.

No. 319, ♂, Santarem, Brazil, J. D. Haseman coll., December 12, 1909.

No. 328, ♂, São Luiz de Cáceres, J. D. Haseman coll., May 25, 1909.

No. 360, ♂, Santarem, Brazil, J. D. Haseman coll., December 9, 1909.

No. 2038, ♂, São Antonio de Guaporé, J. D. Haseman coll., July 30, 1909.

Specimen No. 2038 was taken in the Rio Guaporé, and according to Haseman's note is called locally "Cobra de Agua." It presents some variations from the description given by Boulenger. The rostral is separated from the internasal by the nasal. The frontal is once and two-thirds as long as broad, as wide in front as behind, with perfectly straight and parallel sides. It is shorter than its distance from the end of the snout, and shorter than the parietals. The anterior chin-shields are longer and considerably larger than the posterior. There are no keels on the scales of the outer row except in the posterior fourth of the trunk, and on the tail. The color of the lower surface is dark brown, with only a few widely separated small yellow spots. This appears to be the largest specimen of the species which has been recorded.

Specimens 319, 328, and 360 agree more nearly, but not entirely, with Boulenger's description. The eye of all is longer than one-half the frontal shield; this character is either more variable than usual or depends to some extent upon the treatment of the specimen. The lower surfaces of Nos. 319 and 360 are almost black, with round white (originally yellow?) spots on one or both extremities of most of the gastrosteges, forming conspicuous rows. The lower surface of No. 328 presents only a few small and indistinct light spots. The dorsal surface of this specimen is darker than that of any of the others, with more distinct black markings. The three middle rows of dorsal spots tend in all the specimens to run into each other and produce a pattern of diagonal crossing lines.

Hypapophyses are not present at all on the posterior trunk vertebræ of two of these specimens, and are scarcely indicated in the others.

No. 319 was found under a barrel on the bank of the Amazon. No. 360 was caught in a fish net in the Amazon. A small fish was found in the stomach of No. 328.

COUNTS AND MEASUREMENTS.

	(No. 319.)	(No. 328.)	(No. 360.)	(No. 2038.)
Anal.....	1/1	1/1	1/1	1/1
Scale-rows.....	23	25	23	25
Gastrosteges.....	131	119	130	122
Urosteges.....	71/71	70/70	79/79	28/28*
Upper labials.....	8(4)	8(4)	8(3.4) (8(4)	8(4)
Preoculars.....	1	1	1	1
Postoculars.....	2	2	2	2
Temporals.....	1, 2	1, 2	1, 2 2, 4	2, 3
Total length in mm.....	309	344	407	940
Length of tail in mm.....	83	96	116	127*

* Tip of tail lost.

Genus *Herpetodryas* Boie.26. *Herpetodryas carinatus* (Linnæus).

Coluber carinatus LINNÆUS, Mus. Ad. Frid., 1754, p. 31; Syst. Nat., Ed. XII, 1766, I, p. 384.

Herpetodryas carinatus BOULENGER, Cat. Snakes, II, 1894, p. 73.

No. 2, ♂, Province del Sara, Bolivia, Steinbach coll. (elev. 400 M.), Sept., 1911.

No. 3, ♀, Province del Sara, Bolivia, Steinbach coll. (elev. 350 M.), Jan., 1912.

No. 150, ♂, Bonda, Colombia, H. H. Smith coll. (elev. 150 ft.), June.

No. 367, ♂, Villa Bella, Bolivia, J. D. Haseman coll., Oct. 7, 1909.

Nos. 2 and 3 belong to the variety *flavolineatus* Jan. Nos. 150 and 367 are of the variety *bicarinatus* (Wied).

COUNTS AND MEASUREMENTS.

	(No. 2.)	(No. 3.)	(No. 150.)	(No. 367.)
Anal.....	1/1	1/1	1/1	1/1
Scale-rows.....	12	12	12	12
Gastrosteges.....	153	155	157	152
Urosteges.....	130/130	120/120	126/126	136/136
Upper labials.....	9(4.5.6)	9(4.5.6)	{ 8(4.5) 9(5.6)	9(4.5.6)
Preoculars.....	1	1	1	1
Postoculars.....	2	2	2	2
Temporals.....	1, 2	1, 1	1, 2	{ 1, 2 1, 1
Total length in mm.....	1067	937	1605	1330
Length of tail in mm.....	405	366	521	455

27. *Herpetodryas fuscus* (Linnæus).

Coluber fuscus LINNÆUS, Mus. Ad. Frid., 1754, p. 32, Pl. XVII, fig. 1; Syst. Nat., Ed. XII, 1766, I, p. 383.

Herpetodryas fuscus BOULENGER, Cat. Snakes, II, 1894, p. 75.

No. 365, ♂, Villa Bella, Bolivia, J. D. Haseman coll., October 8, 1909.

No. 1428, ♂, "Massasao, River Amazon" *vide* J. O. Kerby, who presented it to the Museum.

No. 1428 belongs to the variety *saturninus* (Linnæus).

COUNTS AND MEASUREMENTS.

	(No. 365.)	(No. 1428.)
Anal.....	1	1
Scale-rows.....	10	10
Gastrosteges.....	150	156
Urosteges.....	129/129	120/120
Upper labials.....	9(4.5.6)	9(4.5.6)
Preoculars.....	1	1
Postoculars.....	2	2
Temporals.....	1, 1	1, 1
Total length in mm.....	702	385
Length of tail in mm.....	350	128

Genus *Leptophis* Bell.28. *Leptophis ahætulla* (Linnaeus).

Coluber ahætulla LINNÆUS, Syst. Nat., Ed. XII, 1766, I, p. 387.

Leptophis liocercus BOULENGER, Cat. Snakes, II, 1894, p. 113.

The Carnegie Museum possesses four specimens.

No. 23, ♀, Las Juntas, Bolivia, 250 M., J. Steinbach coll., December, 1913.

No. 268, ♀, South America, LeBoutelier coll.

No. 314, ♂, Puerto Suarez, J. Steinbach coll., December, 1908.

No. 366, ♂, Villa Bella, Bolivia, J. D. Haseman coll., October 10, 1909.

Nos. 23, 314, and 366 have lost the tips of their tails. A right loreal is present in the case of No. 268. No. 366 was found to have the intestines stuffed with winged termites, and the stomach of No. 314 contained four small frogs.

COUNTS AND MEASUREMENTS.

	(No. 23.)	(No. 268.)	(No. 314.)	(No. 366.)
Anal.....	1/1	1/1	1/1	1/1
Scale-rows.....	15	15	15	15
Gastrosteges.....	166	160	166	158
Urosteges.....	129/129*	145/145		145/145*
Upper labials.....	8(4.5)	9(5.6)	8(4.5)	9(5.6)
Preoculars.....	1	1	1	1
Postoculars.....	2	2	2	2
Temporals.....	1, 2	1, 2	1, 2	1, 2
Total length in mm.....	846	1170	1100	1338
Length of tail in mm.....	430*	437	150*	514*

* End of tail lost.

29. *Leptophis bocourti* Boulenger.

Leptophis bocourti BOULENGER, P. Z. S., 1898, p. 116.

This species is represented by but a single specimen, No. 2011, which was collected by Mrs. H. H. Smith at Cacagualito, Colombia (elev. 1,500 ft.). It is a female.

The specimen agrees well with Boulenger's description, except in the greater number of gastrosteges and the smaller number of urosteges. The terminal shield, or spine, of the tail is formed normally, yet the tip of the tail is thicker than is usual in the genus. It is therefore possible that the tip has been amputated and healed, and that the reduced number of urosteges of this specimen is thus accounted for. The color is a uniform, very dark green on the upper surface, without the dots and lines which Boulenger describes; light green below, except the lower jaw, which is yellow, with a green border covering about half of the lower labials.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	15
Gastrosteges.....	179
Urosteges.....	103/103
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	1212
Length of tail in mm.....	355

30. *Leptophis nigromarginatus* (Günther).

Ahatulla nigromarginata GUENTHER, Ann. & Mag. Nat. Hist. (3), XVIII, 1866, p. 28.

Leptophis nigromarginatus BOULENGER, Cat. Snakes, II, 1894, p. 112, Pl. III, fig. 3.

There are two females of this species in the collection, which were received from Mr. J. O. Kerby, and have attached to them labels which indicate that they were taken at Pranguina on the Amazon River. They bear the Catalog Numbers 2007 and 2008. The tip of the tail of No. 2007 has been lost.

COUNTS AND MEASUREMENTS.

	(No. 2007.)	(No. 2008.)
Anal.....	1/1	1/1
Scale-rows.....	15	15
Gastrosteges.....	158	162
Urosteges.....	111/111	157/157
Upper labials.....	8(4.5)	9(5.6)
		8(4.5)
Preoculars.....	1	1
Postoculars.....	2	2
Temporals.....	1, 1	1, 1
	1, 2	1, 2
Total length in mm.....	940	1077
Length of tail in mm.....	317	392

31. *Leptophis occidentalis* (Günther).

Ahatulla occidentalis GUENTHER, P. Z. S., 1859, p. 412.

Leptophis occidentalis BOULENGER, Cat. Snakes, II, 1894, p. 111, Pl. III, fig. 2.

The Carnegie Museum has five specimens of this species listed as follows:

No. 140, ♀, Bonda, Colombia, H. H. Smith coll., June.

No. 141, ♀, Bonda, Colombia, H. H. Smith coll., August.

No. 1095, ♀, Bonda, Colombia, H. H. Smith coll., August 25.

No. 1869, ♂, Colombia, H. H. Smith coll.

No. 1870, ♀, Colombia, H. H. Smith coll.

COUNTS AND MEASUREMENTS.

	(No. 140.)	(No. 141.)	(No. 1065.)	(No. 1869.)	(No. 1870.)
Anal.....	1/1	1/1	1/1	1/1	1/1
Scale-rows.....	15	15	15	15	15
Gastrosteges.....	184	177	176	168	183
Urosteges.....	158/158	164/164	146/146	152/152	158/158
Upper labials.....	8(4.5)	8(4.5)	8(4.5)	9(5.6)	8(4.5)
			9(5.6)		
Preoculars.....	1	1	1	1	1
Postoculars.....	2	2	2	2	2
Temporals.....	1, 2	1, 2	1, 2	1, 2	1, 2
Total length in mm.....	1323	1486	1463	1800	1475
Length of tail in mm.....	460	575	505	621	504

32. *Leptophis rostralis* Lönnberg.

Leptophis rostralis LÖNNBERG, Ann. & Mag. Nat. Hist. (7), X, 1902, p. 458.

No. 1862 is the only specimen as yet representing this species in the Carnegie Museum. It is slightly defective, having lost the end of the tail. It is a female, and was undoubtedly taken by H. H. Smith in Colombia.

The specimen agrees with Lönnberg's description in all the principal features, though showing some variation. The principal divergence is that in our specimen the preocular is in contact with the frontal. The portion of the rostral visible from above equals one-half the length of the internasals. The internasals are almost (five-sixths) as long as the prefrontals. The frontal is considerably longer than its distance from the end of the snout, but shorter than the parietals. The anterior chin-shields are shorter than the posterior. The color of the upper surface is a uniform dark brown without any visible markings. The upper lip and the lower surface of the head are whitish. The scales of the ventral surface appear to have been dark green; the posterior margin of each is light. The specimen has not been well preserved to show color, and the loss of the data regarding it is unfortunate. The number of gastrosteges exceeds that of the type.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	15
Gastrosteges.....	180
Urosteges.....	86/86*
Upper labials.....	9(5.6)
	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	373
Length of tail in mm.....	94*

* Tip of tail lost.

Genus *Liophis* Wagler.33. *Liophis albiventris* (Jan).

Liophis reginae, vars. *albiventris* and *quadrilineata* JAN, Arch. Zool. Anat. Phys., II, 1863, p. 294; Icon. Gén., 16, 1866, Pl. VI, figs. 2 and 3.

Liophis albiventris BOULENGER, Cat. Snakes, II, 1894, p. 130.

This species is represented by a single male specimen from the LeBoutelier collection, which has been assigned the Catalog Number 278. The label attached to it by the collector states that it came from Sipan, South America.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	17
Gastrosteges.....	153
Urosteges.....	74/74
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	1
Temporals.....	1, 2
Total length in mm.....	495
Length of tail in mm.....	127

34. *Liophis almadensis* (Wagler).

Natrix almadensis WAGLER, in Spix, Serp. Bras., 1824, p. 30, Pl. X, fig. 3.

Liophis almadensis BOULENGER, Cat. Snakes, II, 1894, p. 134.

No. 112, ♂, Santa Cruz de la Sierra, Bolivia, J. Steinbach collector.

No. 352, ♂, São João del Rey, Minas Geraes, Brazil, J. D. Haseman coll., May 17, 1908.

COUNTS AND MEASUREMENTS.

	(No. 112.)	(No. 352.)
Anal.....	1/1	1/1
Scale-rows.....	19	19
Gastrosteges.....	158	161
Urosteges.....	60/60	70/70
Upper labials.....	8(4.5)	8(4.5)
Preoculars.....	1	1
Postoculars.....	2	2
Temporals.....	1, 2	1, 2
Total length in mm.....	352	485
Length of tail in mm.....	71	103

35. *Liophis elæoides* sp. nov.

Tail $5\frac{1}{2}$ to $6\frac{1}{3}$ times in total length. Eye large, its length being equal to the distance from the front of the eye to the middle of the nostril; head broad and flat; snout bluntly pointed. Rostral broader than deep, visible from above; internasals

as long as broad or longer, as long as, or longer than, the prefrontals; frontal once and a quarter to once and a half as long as broad, shorter than its distance from the rostral (this is true of the mature specimens, the frontal of small specimens is longer than its distance from the end of the snout); frontal as long as, or shorter than, the parietals; loreal deeper than long; one pre- and two postoculars; temporals 1, 2; eight upper labials, the fourth and fifth bordering the orbit; five lower labials in contact with the anterior chin-shields which are as long as, or longer than, the posterior.

Scales in 19 rows: gastrosteges 160-167, obtusely angulate laterally; anal divided; urosteges in 49-54 pairs.

Uniform dark green above, this color extending to the outer ends of the gastrosteges. During life the color is undoubtedly of an olive tone, due to the color of the horny scales, most of which are rubbed off in our specimens. There are no spots, nor light or dark edges to scales; neither have the young a dark nuchal band. The lower surface is a uniform yellowish white. The upper lip is light green, lighter than the upper part of the head, but not white. (Type, C. M. Cat. Rept., No. 32.)

The Museum has fourteen specimens of this species, all of which were collected by Mr. José Steinbach in the Provincia del Sara, Bolivia, and most of them bear the label Santa Cruz de la Sierra. They bear in the Catalog of Reptiles of the Museum the numbers 32 (type), 44, 59, and 91-102 inclusive. All after No. 32 are regarded as paratypes. The type has a label giving the date of capture as February, 1913. No. 59 is labelled as having been taken in October, 1911. No dates are given on any of the other labels.

COUNTS AND MEASUREMENTS.

No.	Num- ber.	Sex	Anal.	Scales rows.	Gastro- steges.	Uro- steges.	Upper Labials.	Pre- oculars.	Post- oculars.	Tem- porals.	Total Length in Mm.	Length Tail in Mm.
No.	32	♂	1/1	19	163	49/49	8(4.5)	1	2	1, 2	612	102
"	44	♀	1/1	19	164	50/50	8(4.5)	1	2	1, 2	400	69
"	59	♂	1/1	19	160	49/49	8(4.5)	1	2	1, 2	615	109
"	91	♂	1/1	19	162	50/50	8(4.5)	1	2	1, 2	255	41
"	92	♂	1/1	19	162	54/54	8(4.5)	1	2	1, 2	541	95
"	93	♂	1/1	19	165	54/54	7(4.5)	1	2	1, 2	522	94
"	94	♂	1/1	19	167	49/49	8(4.5)	1	2	1, 6	680	106
"	95	♂	1/1	19	163	53/53	8(4.5)	1	2	1, 2	315	50
"	96	♂	1/1	19	164	49/49	8(4.5)	1	2	1, 2	475	83
"	97	♂	1/1	19	163	50/50	8(4.5)	1	2	1, 2	425	73
"	98	♂	1/1	19	161	49/49	8(4.5)	1	2	1, 2	393	65
"	99	♂	1/1	19	162	50/50	8(4.5)	1	2	1, 2	612	97
"	100	♂	1/1	19	162	52/52	8(4.5)	1	2	1, 2	297	48
"	102	♂	1/1	19	161	50/50	8(4.5)	1	2	1, 2	242	40

36. *Liophis melanostigma* (Wagler).

Natrix melanostigma WAGLER, in Spix, Serp. Bras., 1824, p. 17, Pl. IV, fig. 2.

Liophis melanostigma BOULENGER, Cat. Snakes, II, 1894, p. 142.

No. 340, ♀, Xiririca, Rio Ribeira, J. D. Haseman coll., December 10, 1908.

This is the only representative of this species in the Museum at the present time.

A black zigzag band occupies the vertebral region. The lower parts of the sides are black, this lateral band being broken into large spots on the neck. A light brown band separates the dorsal and lateral dark stripes. The third and fourth rows of scales are marked by an indistinct series of small white spots. A pair of conspicuous white spots lies on the nape just behind the occiput.

Internasals slightly longer than broad; frontal twice as long as broad, considerably longer than its distance from the end of the snout, shorter than the parietals.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	17
Gastrosteges.....	149
Urosteges.....	86/86
Upper labials.....	9(4.5.6)
	8(3.4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	770
Length of tail in mm.....	241

37. *Liophis melanotus* (Shaw).

Coluber melanotus SHAW, Zoölogy, III, 1802, p. 534.

Liophis melanotus BOULENGER, Cat. Snakes, II, 1894, p. 134.

The Carnegie Museum possesses three examples of this species, as follows:

No. 214, ♀, Bonda, Colombia, H. H. Smith coll., June, 1901.

No. 1863, ♂, South America. (No doubt collected by H. H. Smith in Colombia.)

No. 1864, ♂, South America. (No doubt collected by H. H. Smith in Colombia.)

Specimen 214 has a row of black spots covering the tips of the second row of scales in the anterior half of the trunk. The others show traces of the same markings.

COUNTS AND MEASUREMENTS.

	(No. 214.)	(No. 1863.)	(No. 1864.)
Anal.	1/1	1/1	1/1
Scale-rows.	17	17	17
Gastrosteges.	158	155	152
Urosteges.	58/58	36/36*	59/59
Upper labials.	8(4.5)	8(4.5)	8(4.5)
Preoculars.	1	1	1
Postoculars.	2	2	2
Temporals.	1, 2	1, 2	1, 2
Total length in mm.	434	470	356
Length of tail in mm.	85	65*	80

* End of tail lost.

38. *Liophis pæcilogyrus* (Wied).*Coluber pæcilogyrus* WIED, Beitr. Naturgesch. Bras., I, 1825, p. 371, Plate.*Liophis pæcilogyrus* BOULENGER, Cat. Snakes, II, 1894, p. 131.

There are fourteen specimens at hand, which are referable to this species. Seven of them come from the Provincia del Sara, Bolivia, and were collected by José Steinbach, who neglected except in two cases to indicate the date of capture. They are listed as follows:

- No. 7, ♂, Provincia del Sara, Bolivia, J. Steinbach, Jan., 1912.
 No. 53, ♀, Provincia del Sara, Bolivia, J. Steinbach, Jan., 1913.
 No. 105, ♀, Santa Cruz de la Sierra, J. Steinbach.
 No. 106, ♀, Santa Cruz de la Sierra, J. Steinbach.
 No. 111, ♂, Santa Cruz de la Sierra, J. Steinbach.
 No. 113, ♂, Santa Cruz de la Sierra, J. Steinbach.
 No. 274, ♂, South America, LeBoutelier Collection.
 No. 275, ♀, South America, LeBoutelier Collection.
 No. 277, ♂, South America, LeBoutelier Collection.
 No. 312, ♂, Santa Cruz de la Sierra, J. Steinbach.
 No. 332, ♂, Entre Rios, Brazil, J. D. Haseman, June 4, 1908.
 No. 333, ♂, Cidade de Matto Grosso, J. D. Haseman, July 1, 1909.
 No. 351, ♂, Entre Rios, Brazil, J. D. Haseman, June 4, 1908.
 No. 359, ♀, Penedo, Alagoas, Brazil, J. D. Haseman, March 22, 1908.

Specimens No. 7, 53, 105, 106, 113, and 312, from the Province del Sara, Bolivia, are all very dark on the upper surface, and uniformly colored. The edges of the dorsal and lateral scales are black, the centers olive. When the surface of a

specimen is dry, the scales appear brownish with an iridescent sheen. The scales of the upper surface of the head are similarly light brown or olive with dark margins. The upper labials are white. The lower surface is either entirely white, or some of the gastrosteges have more or less complete black anterior margins. Nos. 7 and 113, young specimens, are as uniformly marked as the older ones. There is no nuchal collar, nor are any transverse bars visible. The length of the tail of these specimens is contained $6\frac{3}{8}$ to 7 times in the total length. The number of gastrosteges and urosteges varies within very narrow limits and about the minimum for *L. pacilogyrus*. It is possible that the specimens represent a well defined subspecies.

I feel doubtful about the identification of No. 111. No pits are visible on the scales, and the length of the tail is contained six and one-half times in the total length. The dorsal scales are mostly black, but some are white, or white-edged, so arranged as to form numerous narrow light cross-bars or reticulations. Most of the gastrosteges are entirely or partly brown.

COUNTS AND MEASUREMENTS.

No.	Sex.	Anal.	Scale-rows.	Gastrosteges.	Urosteges.	Upper Labials.	Pre-oculars.	Post-oculars.	Temporals.	Total Length in Min.	Length of Tail in Min.
No. 7....	♂	1/1	19	149	41/41	8(4.5)	1	2	1, 2	277	42
" 53....	♀	1/1	19	148	38/38	8(4.5) { 7(3.4)	1	2	1, 2	510	71
" 105....	♀	1/1	19	142	42/42	8(4.5)	1	2	1, 2	535	77
" 106....	♀	1/1	19	146	46/46	8(4.5)	1	2	1, 2	512	80
" 111....	♂	1/1	19	148	44/44	8(4.5)	1	2	1, 2	251	39
" 113....	♂	1/1	19	146	41/41	8(4.5)	1	2	1, 1, 2	225	33
" 274....	♂	1/1	19	160	61/61	8(4.5)	1	2	1, 2	697	142
" 275....	♀	1/1	19	174	57/57	8(4.5)	1	2	1, 2	882	145
" 312....	♂	1/1	21	157	54/54	8(4.5)	1	2	1, 2	480	83
" 332....	♂	1/1	19	162	62/62	8(4.5)	1	2	1, 2	494	69
" 333....	♂	1/1	19	153	43/43	8(4.5)	1	2	1, 2	275	52
" 351....	♂	1/1	19	167	56/56	8(4.5)	1	2	1, 2	285	40
" 359....	♀	1/1	19	153	49/49	8(4.5)	1	2	1, 2	513	90
										800	132

39. *Liophis reginae* (Linnæus).

Coluber reginae LINNÆUS, Mus. Ad. Frid., 1754, p. 24, Pl. XIII, fig. 3; Syst. Nat., Ed. XII, 1766, p. 378.

Liophis reginae BOULENGER, Cat. Snakes, II, 1894, p. 137.

The Carnegie Museum has three specimens of this species, No. 19, ♂, No. 24, ♀, and No. 27, ♂, which were taken by José Steinbach at Las Juntas, Bolivia, in December, 1913.

COUNTS AND MEASUREMENTS.

	(No. 19.)	(No. 24.)	(No. 27.)
Anal.	1/1	1/1	1/1
Scale-rows.	17	17	17
Gastrosteges.	140	149	146
Urosteges.	70/70	71/71	76/76
Upper labials.	8(4.5)	8(4.5)	8(4.5)
Preoculars.	1	1	1
Postoculars.	2	2	2
Temporals.	1, 2	1, 2	1, 2
Total length in mm.	201	729	612
Length of tail in mm.	46	182	162

40. *Liophis viridis* Günther.

Liophis viridis GÜNTHER, Ann. & Mag. Nat. Hist. (3), IX, 1862, p. 58, Pl. IX, fig. 2.

Liophis viridis BOULENGER, Cat. Snakes, II, 1894, p. 135.

The specimens in the Carnegie Museum are listed as follows:

No. 281, ♂, South America, LeBoutelier collection.

No. 316, ♀, Lagoa de João Pereira, Brazil, J. D. Haseman coll., December 23, 1907.

No. 344, ♀, Bom Jesus de Lapa, Bahia, Brazil, J. D. Haseman coll.

No. 344 is defective, having in part lost its tail.

COUNTS AND MEASUREMENTS.

	(No. 281.)	(No. 316.)	(No. 344.)
Anal.	1/1	1/1	1/1
Scale-rows.	19	19	19
Gastrosteges.	182	198	192
Urosteges.	72/72	76/76	41/41*
Upper labials.	8(4.5)	8(4.5)	8(4.5)
Preoculars.	1	1	1
Postoculars.	2	2	2
Temporals.	1, 2	1, 2	1, 2
Total length in mm.	402	597	528
Length of tail in mm.	85	125	70*

* Tip of tail lost.

Genus *Lystrophis* Cope.41. *Lystrophis histricus* (Jan).

Heterodon histricus JAN, Arch. Zoöl. Anat. Phys., II, 1863, p. 224.

Lystrophis histricus BOULENGER, Cat. Snakes, II, 1894, p. 152.

The only specimen of this species in the possession of the Carnegie Museum (No. 331) was taken at São Matias, Bolivia, by J. D. Haseman, May 7, 1909. It is a male.

Snout, to level of eyes, brown; a darker brown band across the eyes, followed by another, which is interrupted in the middle, and ends on each side back of the

mouth. The bands across the head are separated by very narrow light lines. The chevron-shaped band of the nape is broad and almost joins the parietal bands. The remainder of the upper surface of the body is crossed by similar brown, black-edged, chevron-shaped bands, which almost reach the gastrosteges and are separated by slightly narrower light bars. A small dark spot lies in the center of each light bar between the ventral ends of the dark bands. There are a few faint dark spots on the light ventral surface. The total number of dark bands, including the nuchal and tail bands, is twenty-six.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	19
Gastrosteges.....	142
Urosteges.....	31/31
Upper labials.....	7(3.4)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	176
Length of tail in mm.....	21

42. *Lystrophis semicinctus* (Duméril & Bibron).

Heterodon semicinctus DUMÉRIL & BIBRON, *Erp. Gén.*, VII, 1854, p. 774.

Lystrophis semicinctus BOULENGER, *Cat. Snakes*, II, 1894, p. 153.

There are fifteen specimens of this species in the collection, all taken by José Steinbach in the Province del Sara, Bolivia. It is worthy of note that in the case of the specimen bearing the catalog number 85 both loreals enter the orbit below the preocular. The list of species is as follows:

- No. 12, ♀, Provincia del Sara, J. Steinbach coll., Feb., 1911.
- No. 14, ♂, Provincia del Sara, J. Steinbach coll., Feb., 1911.
- No. 15, ♂, Provincia del Sara, J. Steinbach coll., Feb., 1911.
- No. 31, ♀, Provincia del Sara, J. Steinbach coll., Feb., 1913.
- No. 33, ♂, Provincia del Sara, J. Steinbach coll., Sept., 1913.
- No. 36, ♀, Provincia del Sara, J. Steinbach coll., Sept., 1913.
- No. 39, ♂, Provincia del Sara, J. Steinbach coll., Sept., 1913.
- No. 51, ♂, Provincia del Sara, J. Steinbach coll., Jan., 1913.
- No. 84, ♂, Santa Cruz de la Sierra, José Steinbach coll.
- No. 85, ♂, Santa Cruz de la Sierra, José Steinbach coll.
- No. 86, ♂, Santa Cruz de la Sierra, José Steinbach coll.
- No. 87, ♀, Santa Cruz de la Sierra, José Steinbach coll.

No. 88, ♀, Santa Cruz de la Sierra, José Steinbach coll.

No. 89, ♀, Santa Cruz de la Sierra, José Steinbach coll.

No. 90, ♂, Santa Cruz de la Sierra, José Steinbach coll.

COUNTS AND MEASUREMENTS.

Num- ber	Sex	Anal.	Scale- rows	Gastro- steges	Uro- steges	Upper labials	Pre- oculars	Post- oculars	Tempo- rals	Total Length in Mm.	Length of tail in Mm.
No. 12.....	♀	1/1	21	151	27/27	8(4.5)	1	2	1, 2	416	45
" 14.....	♀	1/1	21	154	35/35	8(4.5)	$\begin{Bmatrix} 2 \\ 1 \end{Bmatrix}$	$\begin{Bmatrix} 2 \\ 1 \end{Bmatrix}$	1, 2	426	58
" 15.....	♂	1/1	21	151	35/35	$\begin{Bmatrix} 8(5) \\ 7(4.5) \end{Bmatrix}$	2	2	1, 2	488	65
" 31.....	♀	1/1	21	155	25/25	8(4.5)	2	2	1, 2	516	57
" 33.....	♀	1/1	21	151	27/27	$\begin{Bmatrix} 9(4.5) \\ 8(4.5) \end{Bmatrix}$	1	2	1, 2	178	17
" 36.....	♀	1/1	21	154	31/31	8(4.5)	1	2	1, 2	488	53
" 39.....	♀	1/1	21	155	30/30	8(4.5)	1	2	1, 2	175	19
" 51.....	♀	1/1	21	154	32/32	8(4.5)	2	2	1, 2	270	32
" 84.....	♀	1/1	21	149	28/28	8(4.5)	1	2	1, 2	237	25
" 85.....	♀	1/1	21	153	37/37	8(4.5)	1	2	1, 2	148	18
" 86.....	♀	1/1	21	156	36/36	8(4.5)	$\begin{Bmatrix} 2 \\ 1 \end{Bmatrix}$	$\begin{Bmatrix} 2 \\ 1 \end{Bmatrix}$	1, 2	235	27
" 87.....	♀	1/1	21	150	26/26	8(4.5)	1	2	1, 2	178	19
" 88.....	♀	1/1	21	140	25/25	8(4.5)	2	2	1, 2	217	22
" 89.....	♀	1/1	21	158	31/31	$\begin{Bmatrix} 8(5) \\ 8(4.5) \end{Bmatrix}$	1	2	1, 2	526	59
" 90.....	♂	1/1	21	150	36/36	8(4.5)	2	2	1, 2	500	70

Genus *Phrynonax* Cope.

43. *Phrynonax fasciatus* (Peters).

Spilotes fasciatus PETERS, Mon. Berl. Akad., 1869, p. 443.

Phrynonax fasciatus BOULENGER, Cat. Snakes, II, 1894, p. 21.

This species is represented by a single specimen taken by Mrs. H. H. Smith at Cacagualito, Colombia, in May, 1901. It is a female and bears the number 2026 in the Catalog of Reptiles in the Carnegie Museum.

COUNTS AND MEASUREMENTS.

Anal.....	1
Scale-rows.....	23
Gastrosteges.....	209
Urosteges.....	116/116
Upper labials.....	8(4.5.6)
	7(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 2
Total length in mm.....	1240
Length of tail in mm.....	299

Genus *Rhadinæa* Cope.44. *Rhadinæa merremi* (Wied).*Coluber merremii* WIED, Reise Bras., II, 1821, p. 121.*Rhadinæa merremii* BOULENGER, Cat. Snakes, II, 1894, p. 168.

The Carnegie Museum has two specimens, bearing the Catalog Nos. 350 and 353, both of which were captured by Mr. J. D. Haseman at Entre Rios, Brazil, on June 4, 1908. They are males.

COUNTS AND MEASUREMENTS.

	(No. 350.)	(No. 353.)
Anal.....	1/1	1/1
Scale-rows.....	17	17
Gastrosteges.....	145	145
Urosteges.....	47/47	44/44
Upper labials.....	8(4.5)	8(4.5)
	9(4.5.6)	
Preoculars.....	1	1
Postoculars.....	2	2
Temporals.....	1, 2	1, 2
Total length in mm.....	842	175
Length of tail in mm.....	147	31

45. *Rhadinæa occipitalis* (Jan).*Enicognathus occipitalis* JAN, Arch. Zool. Anat. Phys., II, 1863, p. 267.*Rhadinæa occipitalis* BOULENGER, Cat. Snakes, II, 1894, p. 175.

The only specimen we possess is a female (C. M. No. 6) which was caught by Mr. José Steinbach in the Province del Sara, Bolivia, at an elevation of 350 M. in January, 1912.

COUNTS AND MEASUREMENTS.

Anal.....	1/1
Scale-rows.....	15
Gastrosteges.....	181
Urosteges.....	70/70
Upper labials.....	8(3.4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	507
Length of tail in mm.....	119

46. *Rhadinæa orina* sp. nov.

Eighteen small maxillary teeth followed after a short interspace by two much larger. Eye rather large. Length of parietals equal to their distance from the

internasals (in one case slightly more); rostral a little broader than deep, visible from above; internasals broader than long, or as long as broad, much shorter than the prefrontals; frontal once and two-thirds to twice as long as wide, longer than its distance from the end of the snout, shorter than the parietals; loreal once and a half to twice as deep as long; one preocular and two postoculars; temporals 1, 2; eight upper labials, the fourth and fifth bordering the orbit; five lower labials in contact with the anterior chin-shields, which are as long as, or (usually) considerably longer than, the posterior.

Scales in seventeen rows; anal divided; gastrosteges 156 to 161; urosteges in 52-60 pairs. Length of tail five and one-third to six times in the total length.

Olive above, the scales tipped or margined with black. The younger specimens have a chevron-shaped black band crossing the back of the head, the point lying on the parietal shields, and the ends passing just behind the angle of the mouth. This is followed by a rather distinct narrow light nuchal collar, behind which are a number of distinct narrow dark bands separated by much narrower light spaces. On the posterior part of the trunk and on the tail are four indistinct, longitudinal, dark stripes, two being dorsal and one along the middle of each side. The upper surface of the head is dark brown. The upper lip and the lower surface are yellowish, the anterior margins of the gastrosteges are brown, the brown line being widest at the outer ends of the scales. The markings are most distinct in the younger specimens. The older ones are nearly uniform olive above, and yellowish beneath, with only faint traces of the dark bars on the anterior part of the body and of the lines on the posterior extremity.

This species is very closely related to *R. merremi*.

Type, C. M. No. 264; paratypes, C. M. Nos. 263, 265, 266, 267.

The specimens belonging to this species form a part of the LeBoutelier Collection presented to the Museum by Mrs. A. Marshall Bell, and are labelled as from "the Sierras of Bolivia."

COUNTS AND MEASUREMENTS.

	(No. 263 ♀.)	(No. 264 ♂.)	(No. 265 ♀.)	(No. 266 ♂.)	(No. 267 ♂.)
Anal.....	1/1	1/1	1/1	1/1	1/1
Scale-rows.....	17	17	17	17	17
Gastrosteges.....	161	157	156	157	159
Urosteges.....	60/60	55/55	55/55	55/55	52/52
Upper labials.....	8(4.5)	8(4.5)	8(4.5)	8(4.5)	8(4.5)
Preoculars.....	1	1	1	1	1
Postoculars.....	2	2	2	2	2
Temporals.....	1, 2	1, 2	1, 2	1, 2	1, 2
Total length in mm.....	336	337	274	235	243
Length of tail in mm.....	63	60	50	43	41

Genus *Spilotes* Wagler.47. *Spilotes pullatus* (Linnæus).

Coluber pullatus LINNÆUS, Mus. Ad. Frid., 1754, p. 35, Pl. XX, fig. 3; Syst. Nat., Ed. XII, 1766, p. 388.

Spilotes pullatus BOULENGER, Cat. Snakes, II, 1894, p. 23.

The eight specimens we possess were all taken in Colombia by Mr. H. H. Smith. They are listed as follows:

No. 145, Bonda, Colombia, elev. 150 ft., June.

No. 146, ♀, Bonda, Colombia, elev. 150 ft., June.

No. 147, ♂, Bonda, Colombia, elev. 150 ft., August.

No. 148, ♂, Bonda, Colombia, elev. 150 ft., June.

No. 187, ♀, Bonda, Colombia, elev. 150 ft., August.

No. 1781, ♂, Masinga, Colombia, elev. 2,000 ft., August 16.

No. 2039, ♂, Bonda, Colombia, elev. 150 ft., June.

No. 2040, ♀, Minca, Colombia, elev. 2,000 ft., July 27, 1899.

COUNTS AND MEASUREMENTS.

Num- ber.	Sex.	Anal.	Scale- rows.	Gastro- steges.	Uro- steges.	Upper Labials.	Pre- oculars.	Post- oculars.	Tem- porals.	Total Length in Mm.	Length of Tail in Mm.
No. 145 ...			16		114/114	8(4.5) 9(5.6)	1	2	2, 1 1, 1	1,930	527
" 146 ...	♀	1	18	226	122/122	8(4.5)	1	2	1, 1	1,880	450
" 147 ...	♂	1	16	215	129/129	8(4.5)	1	2	2, 1	2,135	571
" 148 ...	♂	1	16	210	122/122	7(3.4)	1	2	1, 1	1,875	528
" 187 ...	♀	1	16	223	121/121	7(3.4)	1	2	1, 1	760	184
" 1781 ...	♂	1	16	213	117/117	8(4.5)	1	2	1, 1	2,010	562
" 2039 ...	♂	1	16	231	117/117	7(3.4)	1	2	1, 1	532	125
" 2040 ...	♀	1	16	220	113/113	8(4.5)	1	2	1, 1	520	116

Genus *Tropidodipsas* Günther.48. *Tropidodipsas spilogaster* sp. nov. (Plate XVIII, figs. 4-6.)

Maxillary short, the thirteen or fourteen maxillary teeth decreasing a little in front and more behind; the posterior mandibular teeth decrease gradually. Head very distinct from neck; eye large, its length equal to the distance from the front of the orbit to the center of the nostril, protuberant, with a small vertically elliptic pupil. Body cylindrical, slender; scales smooth, without pits.

Rostral much broader than deep, just visible from above; nasal divided; internasals a trifle more than half as long as the prefrontals; prefrontals entering the orbit; frontal as broad as long, shorter than its distance from the end of the

snout, much shorter than the parietals, twice as wide as the supra-ocular; supra-ocular much broader behind than in front; loreal oblong, once and a fifth to once and a half as long as deep, entering the orbit; no preocular; one crescentic postocular; temporals 1, 2; seven upper labials, the third and fourth entering the orbit; two pairs of short chin-shields, the anterior a little the longer and as broad as long, the posterior broader than long.

Scales smooth, in 15 rows; anal entire; gastrosteges 147-157; urosteges in 44 to 47 pairs.

Upper surface reddish brown, with a dorsal series of black spots narrowly margined with white; the anterior three or four spots are several times as large as the others and extend upon the edges of the gastrosteges. Between the first and second, and the second and third spots, the interspaces are almost white (red in life?). The sides of the body and tail bear smaller and more numerous black spots which may reach to the edges of the gastrosteges. The scales of the lighter areas are finely stippled with brown and black.

A black fleur-de-lys-like marking on the parietals and frontals (on specimen 47 a nearly uniform dark rounded spot); lips, chin, and lower surface white (pink in life?) with black dots; most of these on specimen 42 are very small, but on 47 they are nearly all large square checks. The urosteges are nearly covered by black checks.

The type (C. M. No. 42) and the paratype (C. M. No. 47) were both collected in the Province del Sara, Bolivia, at an elevation of 350 meters, by Mr. José Steinbach, the former in November, the latter in December, of the year 1912. Both specimens are males.

COUNTS AND MEASUREMENTS.

	(No. 42 ♂.)	(No. 47 ♂.)
Anal.....	1	1
Scale-rows.....	15	15
Gastrosteges.....	147	157
Urosteges.....	44/44	47/47
Upper labials.....	7(3.4)	6(3.4) 7(3.4)
Preoculars.....	0	0
Postoculars.....	1	1
Temporals.....	1, 2	1, 2
Total length in mm.....	297	288
Length of tail in mm.....	56	51

Genus *Xenodon* Boie.49. *Xenodon colubrinus* Günther.

Xenodon colubrinus GUENTHER, Catalog of Colubrine Snakes, 1858, p. 55.

Xenodon colubrinus BOULENGER, Cat. Snakes, II, 1894, p. 146.

The only representative of this species is a specimen which is derived from the LeBoutelier collection (C. M. No. 280) and which has no other indication of the locality than the label "South America." It is a male.

COUNTS AND MEASUREMENTS.

Anal.....	1
Scale-rows.....	19
Gastrosteges.....	147
Urosteges.....	43/43
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	436
Length of tail in mm.....	58

50. *Xenodon merremi* (Wagler).

Ophis merremii WAGLER, in Spix, Serp. Bras., 1824, p. 47, Pl. XVII.

Xenodon merremii BOULENGER, Cat. Snakes, II, 1894, p. 150.

There are three specimens at hand, all of which are females. They are cataloged as follows:

C. M. No. 52, ♀, Prov. del Sara, Bolivia, J. Steinbach coll., January, 1913.

C. M. No. 56, ♀, Prov. del Sara, Bolivia, J. Steinbach coll., October, 1911.

C. M. No. 315, ♀, Puerto Suarez, Bolivia, J. Steinbach coll., January, 1909.

COUNTS AND MEASUREMENTS.

	(No. 52.)	(No. 56.)	(No. 315.)
Anal.....	1/1	1	1/1
Scale-rows.....	19	19	19
Gastrosteges.....	151	146	152
Urosteges.....	32/32	35/35	36/36
Upper labials.....	7(3.4)	7(3.4)	7(4)
Preoculars.....	1	1	1
Postoculars.....	3	3	3
Temporals.....	1, 2	1, 2	1, 2
Total length in mm.....	740	835	514
Length of tail in mm.....	78	102	59

The anal of No. 56 is single, but shows indications of a fusion of two scales. This specimen has only four lower labials in contact with the anterior chin-shields.

51. *Xenodon neuwiedi* Günther.

Xenodon neuwiedi GUENTHER, Ann. & Mag. Nat. Hist. (3), XII, 1863, p. 354, Pl. V, fig. C.

Xenodon neuwiedi BOULENGER, Cat. Snakes, II, 1894, p. 148.

No. 1234, ♂, no data.

No. 1235, ♂, no data.

COUNTS AND MEASUREMENTS.

	(No. 1234.)	(No. 1235.)
Anal.....	1/1	1/1
Scale-rows.....	21	21
Gastrosteges.....	169	168
Urosteges.....	59/59	64/64
Upper labials.....	8(4.5)	8(4.5)
Preoculars.....	1	1
Postoculars.....	2	2
Temporals.....	1, 2	1, 2
Total length in mm.....	330	478
Length of tail in mm.....	54	83

52. *Xenodon severus* (Linnæus).

Coluber severus LINNÆUS, Mus. Ad. Frid., 1754, p. 25, Pl. VIII, fig. 1; Syst. Nat., Ed. XII, 1766, I, p. 379.

Xenodon severus BOULENGER, Cat. Snakes, II, 1894, p. 149.

The only specimen in the collection is a male (C. M. No. 128) taken by José Steinbach at Santa Cruz de la Sierra, Bolivia. Mr. Steinbach furnished no date upon his label.

	(No. 128.)
Anal.....	1
Scale-rows.....	21
Gastrosteges.....	133
Urosteges.....	37/37
Upper labials.....	8(4.5)
Preoculars.....	2 1
Postoculars.....	2 3
Temporals.....	1, 2
Total length in mm.....	385
Length of tail in mm.....	48

SERIES B. OPISTHOGLYPHA.

Subfamily DIPSADOMORPHINÆ Boulenger.

Genus *Clelia* (Oxyrhopus) Fitzinger.53. *Clelia bitorquata* (Günther).

Tachymenis bitorquatus GUENTHER, Ann. & Mag. Nat. Hist. (4), IX, 1872, p. 19.

Oxyrhopus bitorquatus BOULENGER, Cat. Snakes, III, 1896, p. 104, Pl. VI, fig. 1.

The four specimens of this species in the Museum are listed as follows:

C. M. No. 272, ♀, "South America," LeBoutelier Collection.

C. M. No. 343, ♂, Bom Jesus de Lapa, Bahia, Brazil, J. D. Haseman coll., no date.

C. M. No. 369, ♂, Piracicaba, J. D. Haseman coll., Sept. 7, 1908.

C. M. No. 376, ♀, Tarma, Peru, Miss Lola Vance coll., no date.

Nos. 272, 343, and 369 have black cross-bands arranged in threes the entire length of the body and tail. No. 376 has two black bands on the neck, two sets of threes on the anterior part of the body, traces of a third set, and the remainder of the upper surface (originally) red with black tipped scales.

COUNTS AND MEASUREMENTS.

	(No. 272.)	(No. 343.)	(No. 369.)	(No. 376.)
Anal.....	1	1	1	1
Scale-rows.....	19	19	19	19
Gastrosteges.....	208	206	205	207
Urosteges.....	76/76	63/63	70/70	75/75
Upper labials.....	8(4.5)	8(4.5)	8(4.5)	8(4.5)
Precoculars.....	1	1	1	1
Postoculars.....	2	2	2	2
Temporals.....	2, 3	2, 3	2, 3	2, 3
Total length in mm.....	808	656	1150	574
Length of tail in mm.....	152	111	200	110

54. *Clelia clelia* (Daudin).

Coluber clelia DAUDIN, Rept., VI, 1803, p. 330, Pl. LXXVIII.

Oxyrhopus clelia BOULENGER, Cat. Snakes, III, 1896, p. 108.

C. M. No. 330, ♂, Entre Rios, Brazil, J. D. Haseman coll., June 4, 1908.

C. M. No. 380, ♂, Rio Mamoré, Bolivia, J. D. Haseman coll., Sept. 19, 1909.

In specimen No. 330 there is no loreal.

COUNTS AND MEASUREMENTS.

	(No. 330.)	(No. 380.)
Anal.....	1	1
Scale-rows.....	19	19
Gastrosteges.....	239	230
Urosteges.....	76/76	77/77
Upper labials.....	7(3.4)	7(3.4)
Preoculars.....	1	1
Postoculars.....	2	2
Temporals.....	2, 3	2, 3
Total length in mm.....	623	1,857
Length of tail in mm.....	99	320

55. *Clelia doliata* (Duméril & Bibron).

Oxyrhopus doliatus DUMÉRIL & BIBRON, Erp. Gén., VII, 1854, p. 1020.

Oxyrhopus doliatus BOULENGER, Cat. Snakes, III, 1896, p. 106.

The collection contains one male of this species, which formed a part of the LeBoutelier Collection, and, as was the case with all of the specimens in that collection, has no definite locality-label, being simply marked as from "South America." The tip of the tail of the specimen has been lost.

Eye nearly one-half its distance from the end of the snout; the portion of the rostral visible from above one-half as long as its distance from the frontal; frontal once and a quarter as long as broad, a little shorter than its distance from the end of the snout, and a very little shorter than the parietals.

Broad black cross-bands extending to the gastrosteges, posteriorly making complete annuli. These are much broader than the light interspaces, almost touching along the middle of the back, narrower on the sides, though still wider than the light bands. The markings are intermediate between varieties A and B of Boulenger (*loc. cit.*).

COUNTS AND MEASUREMENTS.

	(No. 273.)
Anal.....	1
Scale-rows.....	19
Gastrosteges.....	201
Urosteges.....	*
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 3
Total length in mm.....	835
Length of tail in mm.....	60*

* Tip of tail lost.

56. *Clelia euprepa* sp. nov. (Plate XXVIII, figs. 7-9.)

Thirteen solid maxillary teeth followed after a short space by two slightly enlarged grooved fangs. Eye large, its length three-fifths to two-thirds of the distance from the eye to the tip of the snout. Snout broad, rounded, scarcely projecting beyond the lower lip. Rostral one-half as deep as broad, the portion visible from above one-half or less the length of the internasal; internasals two-thirds as long as the prefrontals; frontal once and a third as long as wide, pentagonal, longer than its distance from the end of the snout, shorter than the parietals; loreal once and a half to nearly twice as long as deep; one preocular, just reaching the top of the head, separated from the frontal; two postoculars; temporals 2, 3 (2, 2 on the left side of No. 109); eight upper labials, the third, fourth, and fifth bordering the orbit. The third upper labial touches the edge of the orbit by its corner only. Nine lower labials, four in contact with the anterior chin-shields, which are longer than the posterior.

Nineteen rows of scales; anal undivided; gastrosteges 225-230; urosteges in 94-105 pairs.

The body and tail bear narrow, irregular black rings, separated by narrower white spaces which are probably bright red in life. The outlines of the black markings are extremely irregular, many being united by diagonal black bars, while others are broken along the back. Very few of the ventral shields of the dark rings are completely black, the color extending half or two-thirds across a shield and then stopping abruptly. Most of the light dorsal and lateral scales are tipped with brown. The scales of the upper surface of the head and the lower labials are black in the center with light (red ?) edges. A large white (red ?) spot crosses the nape just behind the parietals.

Type, C. M. No. 109; paratype, C. M. No. 108.

The Museum possesses two males of this species, collected by José Steinbach at Santa Cruz de la Sierra, Bolivia, and numbered as above.

COUNTS AND MEASUREMENTS.

	(No. 108.)	(No. 109.)
Anal.....	1	1
Scale-rows.....	19	19
Gastrosteges.....	230	225
Urosteges.....	105/105	94/94
Upper labials.....	8(3.4.5)	8(3.4.5)
Preoculars.....	1	1
Postoculars.....	2	2
Temporals.....	2, 3	{ 2, 2 2, 3
Total length in mm.....	577	797
Length of tail in mm.....	135	170

57. *Clelia peruviana* sp. nov.

Eleven subequal solid maxillary teeth followed after a short space by two slightly enlarged grooved fangs. Eye small, two-fifths as long as its distance from the end of the snout. Snout rounded, scarcely projecting beyond the lower lip. Rostral two-thirds as deep as broad, the portion visible from above equal to the suture between the internasals, not quite one-third the distance between it and the frontal; internasals one-half as long as the prefrontals; frontal once and one-fifth as long as broad, as long as its distance from the end of the snout, a little shorter than the parietals; loreal once and two-thirds as long as deep; one preocular, separated, but not widely, from the frontal, and reaching the upper surface of the head; two postoculars; temporals 2, 3; eight upper labials, the fourth and fifth entering the orbit; four lower labials in contact with the anterior chin-shields, which are longer than the posterior.

Scales in 19 rows; anal entire; 179 gastrosteges; 70 pairs of urosteges.

Top of the head orange, with a dark patch on the frontal and parietals. The back and sides of the anterior third of the body are uniform black. Back of this the body is marked by broad black rings separated by narrow light (red?) spaces in which each scale is tipped or margined with black. The black annuli are a little narrower below than above. In the posterior part of the body and tail the lower surface is almost entirely black. Passing forward, the black color becomes limited more and more to the posterior margins of the ventral shields. The gastrosteges of the anterior quarter have only their outer edges black-margined. The under surface of the head, the lips, and the throat are uniformly yellowish.

The type (C. M. No. 377) is unique. It is a male, and was collected at Tarma, Peru, by Miss Lola Vance, who neglected to give the date of capture.

COUNTS AND MEASUREMENTS.

(No. 377.)

Anal.....	1
Scale-rows.....	19
Gastrosteges.....	179
Urosteges.....	70/70
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 3
Total length in mm.....	691
Length of tail in mm.....	148

58. *Clelia petolaria* (Linnæus).

Coluber petolaris LINNÆUS, Mus. Ad. Frid., 1754, p. 35, Pl. IX, fig. 2; Syst. Nat., Ed. XII, 1766, I, p. 387.

Oxyrhopus petolaris BOULENGER, Cat. Snakes, III, 1896, p. 101.

The specimens in the Carnegie Museum are listed as follows:

C. M. No. 22, ♂, Las Juntas, Bolivia, elev. 250 M., Steinbach coll., Dec., 1913.

C. M. No. 75, ♀, Santa Cruz de la Sierra, Bolivia, Steinbach coll.

C. M. No. 107, ♂, Santa Cruz de la Sierra, Bolivia, Steinbach coll.

C. M. No. 110, ♂, Santa Cruz de la Sierra, Bolivia, Steinbach coll.

The specimen No. 107 is slightly defective, having lost the tip of its tail. Nos. 22, 107, and 110 have broad black bars across the back; the abdomens are uniformly light.

No. 22 has nineteen bars, three to four times as wide as the light interspaces. No. 107 has nineteen bars, two to three times as wide as the interspaces. No. 110 has 28 bars, three to four times as wide as the very narrow interspaces.

COUNTS AND MEASUREMENTS.

	(No. 22.)	(No. 75.)	(No. 107.)	(No. 110.)
Anal.....	1	1	1	1
Scale-rows.....	19	19	19	19
Gastrosteges.....	197	200	205	200
Urosteges.....	82/82	80/80	54/54*	85/85
Upper labials.....	8(4.5)	8(4.5)	8(4.5)	8(4.5)
Preoculars.....	1	1	1	1
Postoculars.....	2	2	2	2
Temporals.....	2, 3	2, 3	2, 3	2, 3
Total length in mm.....	313	164	633	255
Length of tail in mm.....	63	38	54*	54

* Tip of tail lost.

59. *Clelia rhombifer* (Duméril & Bibron).

Oxyrhopus rhombifer DUMÉRIL & BIBRON, Erp. Gén., VII, 1854, p. 1018.

Oxyrhopus rhombifer BOULENGER, Cat. Snakes, III, 1896, p. 103.

C. M. No. 11, ♂, Provincia del Sara, Bolivia, Steinbach coll., March, 1912.

C. M. No. 45, ♂, Provincia del Sara, Bolivia, Steinbach coll., December, 1912.

C. M. No. 57, ♂, Provincia del Sara, Bolivia, Steinbach coll., December, 1911.

C. M. No. 64, ♂, Provincia del Sara, Bolivia, Steinbach coll., December, 1911.

C. M. No. 65, ♂, Provincia del Sara, Bolivia, Steinbach coll., December, 1911.

C. M. No. 66, ♂, Provincia del Sara, Bolivia, Steinbach coll., December, 1911.

COUNTS AND MEASUREMENTS.

Num- ber.	Sex.	Anal.	Scale rows	Gastro- steges.	Uro- steges.	Upper labials.	Pre- oculars.	Post- oculars.	Tem- porals.	Total Length in mm.	Length Tail in mm.
No. 11.....	♂	1	19	196	75/75	8(4.5)	1	2	2, 3	580	125
" 45.....	♂	1	19	201	87/87	8(4.5)	2	2	2, 2	375	80
" 57.....	♂	1	19	203	69/69	8(3.4, 5)	1	2	2, 3	661	118
" 64.....	♂	1	19	206	66/66	8(4.5)	1	2	2, 3	633	117
" 65.....	♂	1	19	195	77/77	8(4.5)	1	2	2, 3	336	68
" 66.....	♂	1	19	198	77/77	8(3.4, 5)	1	2	2, 3	615	130

60. *Clelia trigemina* (Duméril & Bibron).

Oxyrhopus trigeminus DUMÉRIL & BIBRON, *Erp. Gén.*, VII, 1854, p. 1013.

Oxyrhopus trigeminus BOULENGER, *Cat. Snakes*, III, 1896, p. 104.

The only specimen we have at present is a female from the LeBoutelier Collection, which is simply labelled as from "South America." C. M. No. 271.

Frontal a trifle shorter than its distance from the tip of the snout, much shorter than the parietals. The triads of dark brown bars are widely separated, but the individual bars in each are close and not very distinctly marked, as the intervening red scales are heavily colored with dark brown.

COUNTS AND MEASUREMENTS.

(No. 271.)

Anal.....	1
Scale-rows.....	19
Gastrosteges.....	199
Urosteges.....	64/64
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 3
Total length in mm.....	688
Length of tail in mm.....	120

Genus *Pseudoboa* Schneider.61. *Pseudoboa coronata* Schneider.

Pseudoboa coronata SCHNEIDER, *Hist. Amph.*, II, 1801, p. 286.

Oxyrhopus coronatus BOULENGER, *Cat. Snakes*, III, 1896, p. 111.

The species is represented in the collection by a male (C. M. No. 116) taken at Santa Cruz de la Sierra by José Steinbach, who failed to give any further information.

Eye two-fifths as long as the snout; frontal slightly broader than long, shorter than its distance from the end of the snout; anterior chin-shields a little shorter

than the posterior; pits of scales extremely small and difficult to see. The color (in alcohol) is very light, almost white above, except for the dark brown on the front of the head and the nape.

COUNTS AND MEASUREMENTS.

	(No. 116.)
Anal.....	1
Scale-rows.....	17
Gastrosteges.....	191
Urosteges.....	76
Upper labials.....	7(3.4)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 2
Total length in mm.....	297
Length of tail in mm.....	64

Genus *Erythrolamprus* Wagler.62. *Erythrolamprus æsculapii* (Linnæus).

Coluber æsculapii LINNÆUS, Mus. Ad. Frid., 1754, p. 29, Pl. XI, fig. 2; Syst. Nat., Ed. XII, 1766, I, p. 380.

Erythrolamprus æsculapii BOULENGER, Cat. Snakes, III, 1896, p. 200.

There are fourteen specimens of this species in the collection. All belong to the variety having double annuli. The upper surface of the head is largely black, without distinct bands. The specimens are cataloged as follows:

- C. M. No. 129, ♂, Valparaiso, Colombia, H. H. Smith coll., April.
- C. M. No. 130, ♀, Bonda, Colombia, H. H. Smith coll., May.
- C. M. No. 131, ♀, Bonda, Colombia, H. H. Smith coll., June.
- C. M. No. 132, ♀, Bonda, Colombia, H. H. Smith coll., May.
- C. M. No. 133, ♀, Bonda, Colombia, H. H. Smith coll., June.
- C. M. No. 134, ♀, Bonda, Colombia, H. H. Smith coll., May.
- C. M. No. 135, ♂, Bonda, Colombia, H. H. Smith coll., May.
- C. M. No. 270, ♀, South America, LeBoutelier Collection.
- C. M. No. 342, ♀, Entre Rios, Brazil, J. D. Haseman coll., June 4, 1909.
- C. M. No. 1096, ♂, Bonda, Colombia, H. H. Smith coll., August.
- C. M. No. 1841, ♀, South America, LeBoutelier Collection.
- C. M. No. 1842, ♀, South America, LeBoutelier Collection.
- C. M. No. 1843, ♀, South America, LeBoutelier Collection.
- C. M. No. 2035, ♂, Cacagualito, Colombia, Mrs. H. H. Smith coll., no date given.

COUNTS AND MEASUREMENTS.

No.	Sex.	Anal.	Scale-rows.	Gastro-stages.	Uro-stages.	Upper Labials.	Pre-oculars.	Post-oculars.	Temporals.	Total Length in Mm.	Length of Tail in Mm.
No. 129...	♂	1/1	15	192	60/60	7(3.4)	{ 2 1	2	1, 2	507	83
" 130...	♀	1/1	15	188	47/47	7(3.4)	{ 1 1	2	1, 2	516	70
" 131...	♀	1/1	15	193	52/52	7(3.4)	{ 1 2	2	1, 2	867	130
" 132...	♀	1/1	15	185	47/47	7(3.4)	1	2	1, 2	804	120
" 133...	♀	1/1	15	193	37/37	7(3.4)	1	2	1, 2	798	87
" 134...	♀	1/1	15	190	48/48	7(3.4)	1	2	1, 2	859	113
" 135...	♂	1/1	15	201	61/61	7(3.4)	2	2	1, 2	780	146
" 270...	♀	1/1	15	193	36/36	7(3.4)	1	2	1, 2	558	58
" 342...	♀	1/1	15	203	48/48	7(3.4)	1	2	1, 2	756	96
" 1096...	♂	1/1	15	195	59/59	7(3.4)	2	2	1, 2	784	131
" 1841...	♀	1/1	16	191	3/3*	7(3.4)	1	2	1, 2	709	5*
" 1842...	♀	1/1	15	179	51/51	7(3.4)	1	2	1, 2	842	125
" 1843...	♀	1/1	15	192	51/51	7(3.4)	{ 1 2	2	1, 2	810	115
" 2035...	♂	1/1	15	192	60/60	7(3.4)	1	2	1, 2	732	128

* The tail is lost.

Genus *Himantodes* Duméril & Bibron.63. *Himantodes cenchoa* (Linnæus).*Coluber cenchoa* LINNÆUS, Syst. Nat., Ed. XII, 1766, I, p. 389.*Himantodes cenchoa* BOULENGER, Cat. Snakes, III, 1896, p. 84.

The specimens in the Carnegie Museum are cataloged as follows:

C. M. No. 189, ♀, Bonda, Colombia, H. H. Smith coll., September 1.

C. M. No. 190, ♂, Bonda, Colombia, H. H. Smith coll., June.

C. M. No. 362, ♂, Muniz Freire, Espiritu Santo, Brazil, Haseman coll., June 18, 1908.

C. M. No. 375, ♂, Tarma, Peru, Miss Lola Vance coll., no date.

COUNTS AND MEASUREMENTS.

	(No. 189.)	(No. 190.)	(No. 362.)	(No. 375.)
Anal.	1/1	1/1	1/1	1/1
Scale-rows.	17	17	17	17
Gastrosteges.	243	236	240	237
Urosteges.	145/145	142/142	167/167	157/157
Upper labials.	8(4.5)	8(4.5)	8(4.5)	8(3.4.5)
	8(3.4.5)	8(3.4.5)		9(4.5.6)
Preoculars.	1	1	2	2
		2	2	
Postoculars.	2	2	2	2
			3	
Temporals.	2, 3	2, 3	2, 2, 3	2, 3
	3, 3		2, 3	3, 3
Total length in mm.	902	820	1285	1057
Length of tail in mm.	251	240	381	334

Genus *Tantilla* Baird & Girard.64. *Tantilla melanocephala* (Linnaeus).

Coluber melanocephalus LINNÆUS, Mus. Ad. Frid., 1754, p. 24, Pl. XV, fig. 2;
Syst. Nat., Ed. XII, 1766, I, p. 378.

Homalocranium melanocephalum BOULENGER, Cat. Snakes, III, 1896, p. 215.

The only example of this serpent which we possess is a male, taken by Mr. J. D. Haseman at Santa Cruz, in the province of Matto Grosso, Brazil, on June 22, 1909. It bears the catalog number 337.

COUNTS AND MEASUREMENTS.

(No. 337.)

Anal.....	1/1
Scale-rows.....	15
Gastrosteges.....	152
Urosteges.....	63/63
Upper labials.....	7(3.4)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 1
Total length in mm.....	180
Length of tail in mm.....	43

65. *Tantilla semicincta* (Duméril & Bibron).

Homalocranium semicinctum DUMÉRIL & BIBRON, Erp. Gén., VII, 1854, p. 862.

Homalocranium semicinctum BOULENGER, Cat. Snakes, III, 1896, p. 219.

The specimens in the Carnegie Museum are cataloged as follows:

No. 200, ♀, Bonda, Colombia, H. H. Smith coll., August.

No. 210, ♂, Bonda, Colombia, H. H. Smith coll., June.

No. 1094, ♂, Bonda, Colombia, H. H. Smith coll., August.

No. 1844, ♀, South America, Mr. and Mrs. H. H. Smith coll., no date.

No. 2024, ♀, Cacagualito, Colombia, Mrs. H. H. Smith, no date.

No. 2037, ♂, Cacagualito, Colombia, Mrs. H. H. Smith, no date.

Nos. 200, 210, 2024, 2037, and 1844 are all dark brown above with narrow white bars or half-bars. The lower surfaces are white. The tip of the snout is yellow touched with brown; there is a yellow spot on the lip behind the eye. The stomach of no. 1094 contained a centipede 153 mm. long.

COUNTS AND MEASUREMENTS.

No.	Num-ber.	Sex.	Anal.	Scale-rows.	Gastro-steges.	Uro-steges.	Upper Labials.	Pre-oculars.	Post-oculars.	Temporals.	Total Length in Mm.	Length of Tail in Mm.
No. 200...		♀	1/1	15	178	59/59	7(3.4)	1	2	1, 1	482	98
" 210...		♀	1/1	15	170	68/68	7(3.4)	1	2	1, 1	465	110
" 1094...		♂	1/1	15	170	64/64	7(3.4)	1	2	1, 1	425	95
" 1844...		♂	1/1	15	180	13/13*	7(3.4)	1	2	1, 1	437	22*
" 2024...		♂	1/1	15	175	10/10*	7(3.4)	1	2	1, 1	448	17*
" 2037...		♀	1/1	15	172	57/57	7(3.4)	1	2	1, 1	200	41

* Tip of tail lost.

Genus *Leptodeira* Fitzinger.66. *Leptodeira annulata* (Linnæus).*Coluber annulatus* LINNÆUS, Mus. Ad. Frid., 1754, p. 34, Pl. VIII, fig. 2.*Leptodira annulata* BOULENGER, Cat. Snakes, III, 1896, p. 97.

We have twenty-one specimens of this snake, fifteen from Bolivia, and six from Colombia. The Bolivian specimens were all taken in the Province del Sara by J. Steinbach, and the Colombian specimens by Mr. and Mrs. H. H. Smith. In the following table the first fifteen numbers are those of the Bolivian examples,

COUNTS AND MEASUREMENTS.

No	Num-ber.	Sex.	Anal.	Scale-rows.	Gastro-steges.	Uro-steges.	Upper Labials.	Sub-oculars.	Pre-oculars.	Post-oculars.	Temporals.	Total Length in Mm.	Length of Tail in Mm.
	9	♀	1/1	19	188	61/61*	8(3.4.5)	0	{2 1	{2 1	1, 2	675	138*
"	37 70	♀ ♂	1/1 1/1	19 19	189 191	73/73 87/87	8(3.4.5) 9(4.5.6) 8(4.5)	0	1	2	1, 2	413	85
"	71	♀	1/1	19	187	73/73	7(3.4)	0	1	2	1, 2	502	122
"	72	♂	1/1	21	189	80/80	8(4.5)	1	1	2	1, 2	645	138
"	73	♂	1/1	19	170	74/74	8(4.5)	1	1	2	1, 2	680	150
"	74	♂	1/1	19	186	92/92	8(4.5) 8(3.4.5)	0	1	2	1, 2	680	148
"	76	♂	1/1	19	196	55/55*	8(4.5)	{0 1	1	2	1, 2	245 603	49 105*
"	77	♂	1/1	19	196	85/85	8(4.5)	1	1	2	1, 2	485	111
"	78	♂	1/1	19	190	87/87	7(3.4)	0	1	2	1, 1	610	150
"	79	♀	1/1	19	184	24/24*	8(4.5)	1	1	2	1, 2	555	52*
"	80	♂	1/1	19	189	90/90	(head injured)					705	178
"	81	♀	1/1	19	191	73/73*	8(4.5)	1	{2 1	{2 1	1, 1	700	140*
"	82	♂	1/1	19	191	85/85	8(4.5)	1	{1 1	{2 3	1, 2	480	118
"	83	♂	1/1	19	191	84/84	8(3.4.5)	0	1	2	1, 2	616	155
"	191	♂	1/1	21	180	88/88	8(4.5)	1	1	2	1, 2	561	154
"	192	♂	1/1	21	178		8(4.5)		2	2	1, 1	502	71*
"	193	♂	1/1	21	178	93/93	8(4.5)		1	2	1, 2	491	132
"	194	♂	1/1	19	173		8(4.5)		1	2	1, 2	520	75*
"	205	♂	1/1	21	177	88/88	8(4.5)		1	2	1, 2	404	103
"	2023	♂	1/1	21	176	79/79	8(4.5)		1	2	1, 2	805	181

* Tip of tail lost.

and the last six represent the material from Colombia. The Colombian material was collected from June to September at Bonda, except No. 2023, which was taken at Cacagualito. No dates accompany the specimens received from Mr. Steinbach, except in the case of No. 9, which was taken in January, and No. 37, which bears the label "August, 1913."

Genus *Oxybelis* Wagler.

67. *Oxybelis acuminatus* (Wied).

Coluber acuminatus WIED, Abbild. Nat. Bras., 1822, and Beitr. Nat. Bras., I, 1825, p. 322.

Oxybelis acuminatus BOULENGER, Cat. Snakes, III, 1896, p. 192.

The specimens in the Carnegie Museum are cataloged as follows:

No. 41, ♂, Provincia del Sara, Bolivia, Steinbach coll., November, 1912.

No. 171, ♀, Bonda, Colombia, H. H. Smith coll., 1901.

No. 172, ♀, Bonda, Colombia, H. H. Smith coll., June, 1901.

No. 173, ♀, Bonda, Colombia, H. H. Smith coll., July, 1901.

No. 174, ♂, Bonda, Colombia, H. H. Smith coll., July, 1901.

No. 175, ♀, Bonda, Colombia, H. H. Smith coll., August, 1901.

No. 184, ♀, Bonda, Colombia, H. H. Smith coll., August, 1901.

No. 1845, ♂, Colombia, H. H. Smith.

No. 1846, ♀, Colombia, H. H. Smith.

No. 1847, ♀, Colombia, H. H. Smith.

No. 1848, ♂, Colombia, H. H. Smith.

No. 1849, ♂, Colombia, H. H. Smith.

No. 1850, ♂, Colombia, H. H. Smith.

No. 2010, ♀, Cacagualito, Colombia, Mrs. H. H. Smith.

COUNTS AND MEASUREMENTS.

No.	Sex.	Anal.	Scale-rows.	Gastro-stages.	Uro-stages.	Upper labials.	Pre-oculars.	Post-oculars.	Temporals.	Total Length in Mm.	Length Tail in Mm.
41	♂	1/1	17	186	158/158	9(4.5.6)	1	2	1, 2	1,083	434
171	♀	1/1	17	185	161/161	9(4.5.6)	1	2	1, 2	1,165	458
172	♀	1/1	17	179	159/159	(94.5.6)	1	2	1, 2	1,390	530
173	♀	1/1	17	182	153/153	9(4.5.6)	1	2	1, 2	1,020	388
174	♂	1/1	17	182	170/170	9(4.5.6)	1	2	1, 2	1,172	472
175	♀	1/1	17	183	153/153	9(4.5.6)	1	2	1, 2	1,136	435
184	♀	1/1	17	189	159/159	9(4.5.6)	1	2	1, 2	1,260	481
1845	♂	1/1	17	..	166/166	9(4.5.6)	1	2	1, 2	1,240	483
1846	♀	1/1	17	185	160/160	9(4.5.6)	1	2	1, 2	1,090	318
1847	♀	1/1	17	179	146/146	9(4.5.6)	1	2	1, 2	1,180	445
1848	♂	1/1	17	180	173/173	9(4.5.6)	1	2	1, 2	1,105	454
1849	♂	1/1	17	184	159/159	9(4.5.6)	1	2	1, 2	1,128	441
1850	♂	1/1	17	183	159/159	9(4.5.6)	1	2	1, 2	455	162
2010	♀	1/1	17	176	159/159	8(4.5)	1	2	1, 2	1,194	471

68. *Oxybelis fulgidus* (Daudin).

Coluber fulgidus DAUDIN, Rept., VI, 1803, p. 352, Pl. LXXX.

Oxybelis fulgidus BOULENGER, Cat. Snakes, III, 1896, p. 191.

The only specimen we have is a female taken at Cacagualito, Colombia, by Mrs. H. H. Smith. It is C. M. No. 2027.

COUNTS AND MEASUREMENTS.

(No. 2027.)

Anal.....	1/1
Scale-rows.....	17
Gastrosteges.....	208
Urosteges.....	158/158
Upper labials.....	10(5.6.7)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	1708
Length of tail in mm.....	592

Genus *Philodryas* Wagler.69. *Philodryas nattereri* Steindachner.

Philodryas nattereri STEINDACHNER, Sitzber. Ak. Wien, LXII, 1870, p. 345, Pl. VII, figs. 1-3.

Philodryas nattereri BOULENGER, Cat. Snakes, III, 1896, p. 134.

A single male, captured by J. D. Haseman at Bom Jesus de Lapa, Bahia, Brazil. No other information accompanies the specimen. It bears the number 345 in the Catalog of the Museum.

Brown above, darkest on the head and neck. Faint brown lines run along the centers of the first and fourth rows of scales on each side of the body.

COUNTS AND MEASUREMENTS.

(No. 345.)

Anal.....	1
Scale-rows.....	21
Gastrosteges.....	209
Urosteges.....	132/132
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 2
Total length in mm.....	540
Length of tail in mm.....	166

70. *Philodryas olfersi* (Lichtenstein).*Coluber olfersii* LICHTENSTEIN, Verz. Doubl., 1823, p. 104.*Philodryas olfersii* BOULENGER, Cat. Snakes, III, 1896, p. 129.

The specimens in the Carnegie Museum are listed as follows:

No. 10, ♂, Provincia del Sara, Bolivia, Steinbach coll., March, 1912.

No. 13, ♂, Provincia del Sara, Bolivia, Steinbach coll., February, 1911.

No. 16, ♂, Provincia del Sara, Bolivia, Steinbach coll., February, 1911.

No. 62, ♂, Provincia del Sara, Bolivia, Steinbach coll., October, 1911.

No. 101, ♂, Santa Cruz de la Sierra, Bolivia, Steinbach, no date.

No. 103, ♂, Santa Cruz de la Sierra, Bolivia, Steinbach, no date.

No. 269, ♂, Brazil, LeBoutelier Collection.

No. 374, ♂, Tarma, Peru, Miss Lola Vance coll., no date.

COUNTS AND MEASUREMENTS.

No.	Num- ber.	Sex.	Anal.	Scale- rows.	Gastro- steges.	Uro- steges.	Upper Labials.	Pre- oculars.	Post- oculars.	Tem- porals.	Total Length in Mm.	Length Tail in Mm.
No. 10	10	♂	1/1	19	189	108/108	8(4.5)	1	2	1, 2	491	138
" 13	13	♂	1/1	19	193	103/103	8(4.5)	1	2	1, 2	1,275	299
" 16	16	♂	1/1	19	188	112/112	8(4.5)	1	2	1, 2	570	166
" 62	62	♂	1/1	19	185	80/80*	8(4.5)	1	2	1, 2	496	120*
" 101	101	♂	1/1	19	192	96/96	8(4.5)	1	2	1, 2	407	99
" 103	103	♂	1/1	19	188	112/112	8(4.5)	1	2	1, 2	960	282
" 269	269	♂	1/1	19	176	104/104	8(4.5)	1	2	1, 2	900	255
" 374	374	♂	1/1	19	189	72/72*	8(4.5)	1 1	2 3	1, 2	480	108*

* Tip of tail lost.

71. *Philodryas schottii* (Schlegel).*Xenodon schottii* SCHLEGEL, Phys. Serp., II, 1837, p. 91, Pl. III, figs. 8 and 9.*Philodryas schottii* BOULENGER, Cat. Snakes, III, 1896, p. 130.

The only example of this species at hand is a female (No. 104) taken by Mr. José Steinbach at Santa Cruz de la Sierra, Bolivia.

COUNTS AND MEASUREMENTS.

	(No. 104.)
Anal.	1/1
Scale-rows.	19
Gastrosteges.	193
Urosteges.	99/99
Upper labials.	7(3.4)
Preoculars.	1
Postoculars.	2
Temporals.	1, 2
Total length in mm.	1515
Length of tail in mm.	379

Genus *Rhinostoma* Duméril & Bibron.72. *Rhinostoma guianense* (Troschel).

Heterodon guianensis TROSCHER, in Schomburgk, Reise Brit. Guian., III, 1848, p. 653.

Rhinostoma guianense BOULENGER, Cat. Snakes, III, 1896, p. 114.

There are thirteen examples of this species in the Museum of which it may be confidently asserted that they were all secured in Colombia by Mr. and Mrs. H. H. Smith, when engaged in collecting in that part of the world. All the specimens which have locality labels attached came from Bonda, and the dates of those upon which there are records of the time of capture show that they were taken from April to the first week in September.

Nos. 211, 212, 1857, and 1858 are of the light-colored variety with a dark brown collar on the nape.

COUNTS AND MEASUREMENTS.

Num- ber.	Sex.	Anal.	Scale- rows.	Gastro- steges.	Uro- steges.	Upper Labials.	Pre- oculars.	Post- oculars.	Tem- porals.	Total Length in Mm.	Length Tail in Mm.
No. 163...	♀	1	19	200	61/61	8(4.5)	1 2	2 2	2, 3	916	151
" 164...	♂	1	19	196	57/57	8(4.5)	1 1	1 2	2, 3	835	134
" 165...	♂	1	19	201	63/63	8(4.5)	1	2	2, 3	825	140
" 166...	♂	1	19	199	60/60	8(4.5)	1	2	2, 3	836	140
" 167...	♂	1	19	191	69/69	8(4.5)	1 1	4 3	2, 3	611	124
" 168...	♂	1	19	206	62/62	8(4.5)	1	2	2, 3	956	157
" 169	♂	1	19	185	70/70	9(5.6) (8(4.5))	1	3	3, 4	553	114
" 170...	♂	1	19	197	57/57	8(4.5)	1	2	2, 3	554	86
" 211...	♂	1	19	199	55/55+5	8(4.5)	1	2	2, 3	375	58
" 212...	♂	1	21	192	73/73	9(4.5, 6) (8(4.5))	1	2	3, 3	344	62
" 1856...	♂	1	19	201	52/52+6	8(4.5)	1	2	2, 3	540	83
" 1857...	♂	1	19	193	70/70	8(4.5)	1	2	2, 3	442	80
" 1858*	..	..	..	..			..	..	..	335*	65*

* Mutilated.

Genus *Tachymenis* Wiegmann.73. *Tachymenis peruviana* Wiegmann.

Tachymenis peruviana WIEGMANN, Nov. Acta Ac. Leop.-Carol., XVII, I, 1835, p. 252, Pl. XX, fig. 1.

Tachymenis peruviana BOULENGER, Cat. Snakes, III, 1896, p. 118.

The specimen (C. M. No. 347) is a female, and was collected by Mr. J. D. Haseman at Bom Jesus de Lapa, Bahia, Brazil.

COUNTS AND MEASUREMENTS.

	(No. 347.)
Anal.....	1/1
Scale-rows.....	19
Gastrosteges.....	146
Urosteges.....	48/48
Upper labials.....	8(4.5)
Preoculars.....	1
Postoculars.....	2
Temporals.....	2, 3
Total length in mm.....	193
Length of tail in mm.....	33

Genus *Thamnodynastes* Wagler.74. *Thamnodynastes nattereri* (Mikan).

Coluber nattereri MIKAN, Delect. Faun. Flor. Bras., 1820, Plate.

Thamnodynastes nattereri BOULENGER, Cat. Snakes, III, 1896, p. 116.

C. M. No. 48, ♂, Provincia del Sara, Bolivia, José Steinbach coll., Dec., 1912.

C. M. No. 339, ♂, Santarem, Brazil, J. D. Haseman coll., December 7, 1909.

C. M. No. 361, ♀, Rio Doce, J. D. Haseman coll., May 20, 1908.

No. 339 was captured in the water of a swamp between the Rio Tapajos and the Amazon. All except the two outer rows of scales are strongly keeled. It conforms to the characters of *T. nattereri* except that there are seventeen instead of nineteen rows of scales. No. 361 was captured in a mill-race.

COUNTS AND MEASUREMENTS.

	(No. 48.)	(No. 339.)	(No. 361.)
Anal.....	1/1	1/1	1/1
Scale-rows.....	19	17	19
Gastrosteges.....	156	148	155
Urosteges.....	74/74	76/76	64/64
Upper labials.....	8(4.5)	8(4.5)	8(3.4.5)
Preoculars.....	1	2	1
Postoculars.....	2	2	2
Temporals.....	2, 3	2, 3	2, 3
Total length in mm.....	568	588	581
Length of tail in mm.....	142	156	112

SERIES C. PROTEROGLYPHA.

Subfamily ELAPINÆ Boulenger.

Genus *Elaps* Schneider.75. *Elaps colombianus* sp. nov.

Seven upper labials, third and fourth bordering the orbit; snout broad and rounded, not projecting; sixth and seventh labials the largest of the series; rostral just visible from above, much broader than deep; internasals much shorter and narrower than prefrontals; first lower labials in contact behind the symphysial; posterior nasal in contact with the preocular. Eye two-thirds to three-quarters as long as its distance from the mouth; frontal once and a quarter to once and two-fifths as long as broad, much broader than the supraoculars, slightly shorter than its distance from the end of the snout (longer than this distance in the young specimen No. 2031), much shorter than the parietals; parietals longer than their distance from the internasals; one pre- and two postoculars; temporals 1, 1; third upper labial larger than the fourth; four lower labials in contact with the anterior chin-shields, which are a little shorter than the posterior.

Scales in 15 rows; anal divided; gastrosteges 187-207; urosteges in 33-44 pairs.

Body with 12 to 14 black rings, three to four scales wide, edged with narrow yellow rings one scale wide; the latter margined with narrow irregularly outlined black rings. The last are variably developed, from a row of scales tipped with black, to two or three rows of all black scales. The red interspaces are more than twice as long as the length of the black triads, with usually black tipped scales. Top of the head black from snout to parietals; posterior edges of parietals, the temporals, fifth, sixth, and part of seventh upper labials crossed by a yellow band, which passes partly or completely around the lower jaw and is widest at the sides of the head. Anterior half of lower jaw black. The first black annulus is separated from the parietals by not more than one scale. A narrow black ring is only faintly indicated behind this first or nuchal annulus. The last annulus of the body crosses the anal scales. On the tail are four to six black annuli, separated by narrow red spaces, in which all the scales may be black-margined, or the central scales are black, forming a narrow black bar which may extend only to the sides or across the lower surface of the tail. The tail pattern is, thus, a broad black annulus, a narrow red ring, a narrow black cross-bar or ring, a narrow red ring, a broad black annulus, etc.

This species seems not distantly related to *Elaps fulvius* (Linnæus).

Type, C. M. No. 197; paratypes, C. M. Nos. 198, 2031, 2033.

There are four specimens representing this species in our collection, which are cataloged as follows:

No. 197, ♀, Minca, Colombia, H. H. Smith coll., June.

No. 198, ♀, Bonda, Colombia, H. H. Smith coll., June.

No. 2031, ♂, Cacagualito, Colombia, Mrs. H. H. Smith, no date.

No. 2033, ♀, Cacagualito, Colombia, Mrs. H. H. Smith, no date.

COUNTS AND MEASUREMENTS.

	(No. 197.)	(No. 198.)	(No. 2031.)	(No. 2033.)
Anal.....	1/1	1/1	1/1	1/1
Scale-rows.....	15	15	15	15
Gastrosteges.....	207	206	187	203
Urosteges.....	33/33	36/36	44/44	33/33
Upper labials.....	7(3.4)	7(3.4)	7(3.4)	6(2.3)
Preoculars.....	1	1	1	7(3.4)
Postoculars.....	2	2	2	1
Temporals.....	1, 1	1, 1	1, 1	2
Total length in mm.....	623	508	343	1, 1
Length of tail in mm.....	56	55	44	597
Annuli.....	13+5	12+4	12+6	64

76. *Elaps corallinus* Wied.

Elaps corallinus WIED, N. Acta Ac. Leop.-Carol., XI, 1820, p. 198, pl. IV.

Elaps corallinus BOULENGER, Cat. Snakes, III, 1896, p. 420.

There are four specimens in the collection, as follows:

C. M. No. 199, ♂, Cacagualito, Colombia, H. H. Smith coll., Oct. 3, 1901.

C. M. No. 261, ♀, LeBoutelier Collection, South America.

C. M. No. 341, ♀, São Antonio de Guaporé, J. D. Haseman coll., August 5, 1909.

C. M. No. 1236, ♂, no data.

COUNTS AND MEASUREMENTS.

	(No. 199.)	(No. 261.)	(No. 341.)	(No. 1236.)
Anal.....	1/1	1/1	1/1	1/1
Scale-rows.....	15	15	15	15
Gastrosteges.....	191	200	213	195
Urosteges.....	47/47	43/43	29/29	44/44
Upper labials.....	7(3.4)	7(3.4)	7(3.4)	7(3.4)
Preoculars.....	1	1	1	1
Postoculars.....	2	2	2	2
Temporals.....	1, 1	1, 1	1, 1	1, 1
Total length in mm.....	373	597	637	605
Length of tail in mm.....	55	83	52	82
Annuli.....	14+7	16+6	24+6	15+6

77. *Elaps frontalis* Duméril & Bibron.*Elaps frontalis* DUMÉRIL & BIBRON, Erp. Gén., VII, 1854, p. 1223.*Elaps frontalis* BOULENGER, Cat. Snakes, III, 1896, p. 427.

This species is represented by a female (No. 356) taken at Sete Lagoas by Mr. J. D. Haseman, May 4, 1908.

COUNTS AND MEASUREMENTS.

	(No. 356.)
Anal.....	1/1
Scale-rows.....	15
Gastrosteges.....	226
Urosteges.....	21/21
Upper labials.....	7(3.4)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 1
Total length in mm.....	726
Length of tail in mm.....	38

78. *Elaps hollandi** sp. nov. (Plate XXVIII, figs. 10-12.)

Seven upper labials, third and fourth bordering the orbit, the sixth largest, seventh next in size; rostral just visible from above; internasals about half as long as the prefrontals; first lower labials in contact behind the symphysial; posterior nasal in contact with the preocular. Eye slightly longer than its distance from the mouth; frontal considerably shorter than the parietals, shorter than its distance from the end of the snout, once and a half as long as broad, hexagonal, considerably wider than the supraocular; parietals longer than their distance from the internasals; rostral nearly twice as broad as deep; one pre- and two postoculars; temporals 1, 1, both large and broad; four lower labials in contact with the anterior chin-shields, which are shorter than the posterior. Tail short, rather bluntly pointed.

Scales in 15 rows; anal divided; gastrosteges 185-204; urosteges in 17-22 pairs.

Body and tail with black annuli arranged in threes, a few of the scales of the interspaces tipped with black. The central black annulus is little wider than the other two, covering four to five and a half gastrosteges while the narrower annuli cover three to four. On No. 206 the three annuli of the tail are similar to those of the body; on No. 207 there are but two caudal annuli. The upper and lower surfaces of the head are black as far as the fourth series of scales back of the parietals; the black of the upper surface is broken by a narrow yellow band passing

* Named in honor of Dr. W. J. Holland, Director of the Carnegie Museum, whose kind interest has stimulated so many scientific workers.

from lip to lip just behind the eyes. This band covers most of the fifth and sixth upper labials, the upper postocular, the anterior temporal, the back of the supraoculars and frontal, and the anterior third of the parietals. The center of the chin is light.

There are two specimens in the collection, cataloged as follows:

C. M. No. 206, type, ♂, Bonda, Colombia, H. H. Smith coll., June, 1901.

C. M. No. 207, paratype, ♀, Bonda, Colombia, H. H. Smith coll., June, 1901.

COUNTS AND MEASUREMENTS.

	(No. 206.)	(No. 207.)
Anal.	1/1	1/1
Scale-rows.	15	15
Gastrosteges.	185	204
Urosteges.	22/22	17/17
Upper labials.	7(3.4)	7(3.4)
Preoculars.	1	1
Postoculars.	2	2
Temporals.	1, 1	1, 1
Total length in mm.	366	318
Length of tail in mm.	30	17
Sets of annuli.	7+1	8+1

79. *Elaps mipartitus* Duméril & Bibron.

Elaps mipartitus DUMÉRIL & BIBRON, *Erp. Gén.*, VII, 1854, p. 1220.

Elaps mipartitus BOULENGER, *Cat. Snakes*, III, 1896, p. 431.

C. M. No. 208, ♀, Valparaiso, Colombia, H. H. Smith, April, 1901.

C. M. No. 209, ♀, Las Nubes, Colombia, H. H. Smith, December, 1901.

Our specimens differ from Boulenger's diagnosis in not having the anterior temporal scale noticeably narrowed. In No. 209 the anterior and posterior temporal shields are fused into one. In both specimens the frontal shield lacks a little of being as long as its distance from the tip of the snout.

The head is black to back of the eyes, the black patch including all or nearly all of the fourth upper labial, and half of the supraoculars and frontal. The yellow band extends to the posterior ends of the parietals. The chin is yellow with scattered, irregularly shaped, dark flecks. The following is copied from the field label:

"This snake when alive has the head and tail almost precisely alike—bright red. It coils up with its head underneath its tail and threatens with its tail."

COUNTS AND MEASUREMENTS.		
	(No. 208.)	(No. 209.)
Anal.	1/1	1/1
Scale-rows.	15	15
Gastrosteges.	252	249
Urosteges.	27/27	24/24
Upper labials.	7(3.4)	7(3.4)
Preoculars.	1	1
Postoculars.	2	1
		2
Temporals.	1, 1	1
Total length in mm.	361	477
Length of tail in mm.	26	32

80. *Elaps narduccii* Jan.

Elaps narduccii JAN, Arch. Zoöl. Anat. Phys., II, 1863, p. 222.

Elaps narduccii BOULENGER, Cat. Snakes, III, 1896, p. 433.

C. M. No. 8, ♀, Provincia del Sera, Bolivia, 350 M., Steinbach coll., Jan., 1912.

C. M. No. 114, ♂, Santa Cruz de la Sierra, Bolivia, Steinbach coll.

Some variability in the proportions of the head shields of this species may be deduced from the published descriptions. In our specimen No. 8 the width of the frontal is equal to that of the supraocular, and the posterior chin-shields are longer than the anterior. In No. 114 the frontal is much wider than the supraocular, and the anterior chin-shields are the longer.

COUNTS AND MEASUREMENTS.		
	(No. 8.)	(No. 114.)
Anal.	1/1	1/1
Scale-rows.	15	15
Gastrosteges.	316	279
Urosteges.	18/18	26/26
Upper labials.	7(3.4)	7(3.4)
Preoculars.	1	1
Postoculars.	2	2
Temporals.	1, 1	2, 1
		1, 1
Total length in mm.	541	467
Length of tail in mm.	21	30

81. *Elaps princeps* Boulenger.

Elaps princeps BOULENGER, Ann. Mag. Nat. Hist. (7), Vol. XV, 1905, p. 456.

C. M. No. 126, ♂, Santa Cruz de la Sierra, Bolivia, Steinbach coll.

This snake was collected at the same place and by the same person as the

original specimens described by Boulenger. Our specimen is considerably larger than any of those, but otherwise is very like them. The Carnegie Museum specimen differs structurally only in that the eye is slightly larger (three-quarters as long as its distance from the mouth); the frontal is once and a quarter as long as broad, and the parietals are a little longer than their distance from the internasals.

COUNTS AND MEASUREMENTS.

(No. 126.)

Anal.....	1/1
Scale-rows.....	15
Gastrosteges.....	221
Urosteges.....	22/22
Upper labials.....	7(3.4)
Preoculars.....	1
Postoculars.....	2
Temporals.....	1, 2
Total length in mm.....	1602
Length of tail in mm.....	65

Family AMBLYCEPHALIDÆ Günther.

Genus *Cochliophagus* Duméril & Bibron.82. *Cochliophagus catesbyi* (Sentzen).

Coluber catesbyi SENTZEN, Meyer's Zoöl. Archiv, II, 1796, p. 66.

Leptognathus catesbyi BOULENGER, Cat. Snakes, III, 1896, p. 449.

C. M. No. 20, ♂, Las Juntas, Bolivia, Steinbach coll., Dec., 1913.

C. M. No. 21, ♂, Las Juntas, Bolivia, Steinbach coll., Dec., 1913.

C. M. No. 28, ♂, Las Juntas, Bolivia, Steinbach coll., Dec., 1913.

C. M. No. 2034, ♂, Brazil, no data.

COUNTS AND MEASUREMENTS.

	(No. 20.)	(No. 21.)	(No. 28.)	(No. 2034.)
Anal.....	1	1	1	1
Scale-rows.....	13	13	13	13
Gastrosteges.....	176	179	175	193
Urosteges.....	94/94	78/78	87/87	93/93
Upper labials.....	{9(4.5.6) 8(3.4.5)	8(4.5)	{8(4.5.6) 8(4.5)	8(4.5)
Preoculars.....	2	2	2	2
Postoculars.....	1	2	2	2
Temporals.....	1, 2	1, 2	1, 2	1, 2
Total length in mm.....	530	208	186	747
Length of tail in mm.....	141	50	45	197*

* Tip of tail lost.

Family VIPERIDÆ Bonaparte.

Subfamily CROTALINÆ Oppel.

Genus *Lachesis* Daudin.83. *Lachesis lanceolatus* (Lacépède).*Coluber lanceolatus* LACÉPÈDE, Serp., II, 1789, pp. 80, 121, Pl. V, fig. 1.*Lachesis lanceolatus* BOULENGER, Cat. Snakes, III, 1896, p. 535.

COUNTS AND MEASUREMENTS.

No.	Sex	Ann.	Scales rows.	Gastro- steges.	Uro- steges.	Upper Labials.	Scales between Supra- oculars.	Scales between Subocu- lar and Labials.	Total Length in Mm.	Length Tail in Mm.
No. 43.....	♀	1	23	181	60/60	8	7	1	550	72
" 121.....	♂	1	23	185	66/66	8	6	1	690	101
" 159.....	♂	1	25	214	70/70	7	8	1	395	57
" 244.....	♂	1	27	196	53/53	7	6	1	1,045	116
" 245.....	♂	1	25	190	63/63	7	7	1	805	110
" 246.....	♂	1	23	202	65/65	8	7	1	820	115
" 247.....	♀	1	25	191	60/60	7	8	1	960	137
" 248.....	♂	1	25	187	58/58	7	6	1	840	130
" 249.....	♂	1	27	180	51/51	7	6	1	885	108
" 250.....	♂	1	25	188		7	7	1	840	..*
" 252.....	♂	1	23	208	54/54	8	7	1	832	91
" 253.....	♂	1	25	181	61/61	7	10	2	923	130
" 254.....	♂	1	25	184	62/62	7	7	1	705	105
" 255.....	♂	1	25	185	59/59	7	7	1	975	143
" 257.....	♂	1	25	184	63/63	7	6	1	857	123
" 258.....	♂	1	27	...		..	..	..	880	118
" 313.....	♀	1	25	182	62/62	8	7	1	535	74
" 372.....	♂	1	27	190	62/62	7	6	1	595	80
" 373.....	..	1	23	184	56/56	..	..	..	369	47
" 2019.....	♂	1	25	218	69/69	7	6	1	565	76

* Tip of tail lost.

The twenty specimens of this species in the collection are cataloged as follows:

C. M. No. 43, ♀, Provincia del Sara, Bolivia, 350 M., Steinbach coll., December, 1912.

C. M. No. 121, ♂, Santa Cruz de la Sierra, Bolivia, Steinbach coll.

C. M. No. 159, ♂, Bonda, Colombia, 150 ft., H. H. Smith coll., August.

C. M. No. 244, ♂, South America, LeBoutelier Collection.

C. M. No. 245, ♂, South America, LeBoutelier Collection.

C. M. No. 246, ♂, South America, LeBoutelier Collection.

C. M. No. 247, ♀, Chili, LeBoutelier Collection.

C. M. No. 248, ♂, Colombia, LeBoutelier Collection.

C. M. No. 249, ♂, Brazil, LeBoutelier Collection.

C. M. No. 250, ♂, Venezuela, LeBoutelier Collection.

- C. M. No. 252, ♀, Peru, LeBoutelier Collection.
C. M. No. 253, ♂, Paraguay, LeBoutelier Collection.
C. M. No. 254, ♂, Paraguay, LeBoutelier Collection.
C. M. No. 255, ♂, Paraguay, LeBoutelier Collection.
C. M. No. 257, ♂, Montevideo, Uruguay, LeBoutelier Collection.
C. M. No. 258, ♂, Argentina (?), LeBoutelier Collection.
C. M. No. 313, ♀, Santa Cruz de la Sierra, Bolivia, Steinbach coll.
C. M. No. 372, ♂, Tarma, Peru, Miss Lola Vance coll.
C. M. No. 373, Tarma, Peru, Miss Lola Vance coll.
C. M. No. 2019, ♂, Cacagualito, Colombia, 1500 ft., Mrs. H. H. Smith coll.

84. *Lachesis lansbergii* (Schlegel).

Trigonocephalus lansbergii SCHLEGEL, Mag. de Zoöl., 1841, Rept. pl. I.

Lachesis lansbergii BOULENGER, Cat. Snakes, III, 1896, p. 546.

There are eighteen specimens of this species in the collection all of which came from the Province of Santa Marta in Colombia, although the last four mentioned in the following list have no data accompanying them. These belong to the lot, the numbers of which run from C. M. No. 1841 to 1873, mentioned in the introductory notes.

- C. M. No. 151, ♂, Bonda, Colombia, H. H. Smith coll., July.
C. M. No. 152, ♀, Bonda, Colombia, H. H. Smith coll., May.
C. M. No. 153, ♂, Bonda, Colombia, H. H. Smith coll., May.
C. M. No. 154, ♀, Bonda, Colombia, H. H. Smith coll., August 31.
C. M. No. 155, ♂, Bonda, Colombia, H. H. Smith coll., August 29.
C. M. No. 156, ♀, Bonda, Colombia, H. H. Smith coll., June.
C. M. No. 157, ♀, Bonda, Colombia, H. H. Smith coll., May.
C. M. No. 158, ♀, Bonda, Colombia, H. H. Smith coll., June.
C. M. No. 160, ♀, Bonda, Colombia, H. H. Smith coll., May.
C. M. No. 185, ♂, Bonda, Colombia, H. H. Smith coll., August.
C. M. No. 1093, ♀, Bonda, Colombia, H. H. Smith coll., August.
C. M. No. 1851, ♂, South America (Colombia) (Mr. & Mrs. H. H. Smith coll.).
C. M. No. 1852, ♀, South America (Colombia) (Mr. & Mrs. H. H. Smith coll.).
C. M. No. 1853, ♀, South America (Colombia) (Mr. & Mrs. H. H. Smith coll.).
C. M. No. 1855, ♂, South America (Colombia) (Mr. & Mrs. H. H. Smith coll.).
C. M. No. 2016, ♀, Cacagualito, Mrs. H. H. Smith coll., no date.
C. M. No. 2017, ♂, Cacagualito, Mrs. H. H. Smith coll., no date.
C. M. No. 2018, ♀, Cacagualito, Mrs. H. H. Smith coll., no date.

Our Colombian series of *L. lansbergi* compels a modification of the formula of the species to include a wider variability, as follows:

Scales in 23-27 rows; anal single; gastrosteges 147-159; urosteges 28-36, as shown by the accompanying table.

COUNTS AND MEASUREMENTS.

Num- ber.	Sex.	Anal.	Scale- rows.	Gastro- steges.	Uro- steges.	Upper Labials.	Scales between Supra- oculars.	Scales between Suboc- ulars and Labials.	Total Length in Mm.	Length of Tail in Mm.
No. 151....	♂	1	25	158	34	9	6	1	376	45
" 152....	♂	1	25	152	32	9	5	1	412	49
" 153....	♂	1	25	155	35	9	5	1	317	38
" 154....	♂	1	25	153	35	9	5	1	440	56
" 155....	♂	1	23	147	33	9	4	1	374	44
" 156....	♂	1	25	147	28	10	3	1	288	27
" 157....	♂	1	25	150	31	9	6	1	301	32
" 158....	♂	1	23	153	33	9	6	1	399	47
" 160....	♂	1	23	153	29	9	4	1	254	24
" 185....	♂	1	25	154	31	9	5	1	452	53
" 1093....	♂	1	25	152	31	9	5	1	570	62
" 1851....	♂	..	..	..	..	9	6	1	..*	..
" 1852....	♂	1	25	158	28	{ 9	5	1	290	26
" 1853....	♀	1	25	151	29	{ 10	5	1	409	41
" 1855....	♂	1	25	151	31	{ 9	5	1	196	19
" 2016....	♀	1	23	154	34	{ 9	4	1	436	55
" 2017....	♂	1	23-24	154	36	{ 10	5	1	225	23
" 2018....	♀	1	25	151	32	{ 10	5	1	420	50

* Mutilated.

85. *Lachesis mutus* (Linnaeus).

Crotalus mutus LINNÆUS, Syst. Nat., Ed. XII, 1766, I, p. 373.

Lachesis mutus BOULENGER, Cat. Snakes, III, 1896, p. 533.

We have but one specimen representing this species in the collection (C. M. No. 125). It consists only of the head and tail of a very large individual, collected by José Steinbach at Santa Cruz de la Sierra, Bolivia.

86. *Lachesis newwiedi* (Wagler).

Bothrops newwiedi WAGLER, in Spix, Serp. Bras., 1824, p. 56, Pl. XXII, fig. 1.

Lachesis newwiedi BOULENGER, Cat. Snakes, III, 1896, p. 542.

The specimens in the collection are listed as follows:

No. 1, ♂, Province del Sara, Bolivia, J. Steinbach coll., Sept., 1911.

No. 4, ♀, Province del Sara, Bolivia, J. Steinbach coll., Jan., 1912.

No. 34, ♂, Province del Sara, Bolivia, J. Steinbach coll., Sept., 1913.

No. 35, ♂, Province del Sara, Bolivia, J. Steinbach coll., Sept., 1913.

- No. 38, ♂, Province del Sara, Bolivia, J. Steinbach coll., Sept., 1913.
 No. 40, ♀, Province del Sara, Bolivia, J. Steinbach coll., Sept., 1913.
 No. 46, ♀, Province del Sara, Bolivia, J. Steinbach coll., Dec., 1912.
 No. 49, ♂, Province del Sara, Bolivia, J. Steinbach coll., Dec., 1912.
 No. 50, ♂, Province del Sara, Bolivia, J. Steinbach coll., Jan., 1913.
 No. 54, ♀, Province del Sara, Bolivia, J. Steinbach coll., Jan., 1913.
 No. 55, ♀, Province del Sara, Bolivia, J. Steinbach coll., Oct., 1911.
 No. 58, ♀, Province del Sara, Bolivia, J. Steinbach coll., Oct., 1911.

COUNTS AND MEASUREMENTS.

No.	Num-ber.	Sex.	Anal.	Scale-rows.	Gastro-steges.	Uro-steges.	Upper Labials.	Scales between Subrostrals and Labials.	Scales between Subrostrals and Labials.	Total Length in Mm.	Length Tail in Mm.
No. 1.....	♂	1	23	169	48/48	8	6	2	442	61	
" 4.....	♀	1	25	170	45/45	9	8	2	827	93	
" 34.....	♀	1	24	172	49/49	8	7	1	288	40	
" 35.....	♀	1	25	171	41/41	8	6	1	467	55	
" 38.....	♀	1	25	178	45/45	8	6	2	775	83	
" 40.....	♀	1	24	180	46/46	8	7	1	438	52	
" 46.....	♀	1	25	166	41/41	7	5	2	438	48	
" 49.....	♂	1	25	177	53/53	8	8	2	410	53	
" 50.....	♀	1	25	175	50/50	8	8	1	558	84	
" 54.....	♀	1	23	174	51/51	8	8	2	548	70	
" 55.....	♀	1	25	179	49/49	8	7	2	264	34	
" 58.....	♀	1	27	183	42/42	8	5	2	502	53	
" 60.....	♀	1	25	180	53/53	8	6	1	580	83	
" 61.....	♀	1	25	181	45/45	8	8	2	630	82	
" 67.....	♀	1	23	170	49/49	8	6	1	365	50	
" 68.....	♀	1	23	171	50/50	8	8	2	354	45	
" 69.....	♀	1	25	179	46/46	8	7	2	277	33	
" 119.....	♀	1	25	184	49/49	8	8	2	455	57	
" 120.....	♀	1	25	176	45/45	10	7	2	334	29	
" 122.....	♀	1	23	167	51/51	8	7	1	897	109	
" 123.....	♀	1	23	175	45/45	7	7	1	314	40	
" 317.....	♀	1	23	174	52/52	8	7	2	380	52	
" 318.....	♀	1	23	184	47/47	8	5	2	425	49	
" 364.....	♀	1	25	171	43/43	8	9	1	815	80	

- No. 60, ♂, Province del Sara, Bolivia, J. Steinbach coll., Oct., 1911.
 No. 61, ♂, Province del Sara, Bolivia, J. Steinbach coll., Oct., 1911.
 No. 67, ♀, Province del Sara, Bolivia, J. Steinbach coll., Oct., 1911.
 No. 68, ♂, Province del Sara, Bolivia, J. Steinbach coll., Oct., 1911.
 No. 69, ♂, Province del Sara, Bolivia, J. Steinbach coll., Oct., 1911.
 No. 119, ♂, Province del Sara, Bolivia, J. Steinbach coll., no date.
 No. 120, ♀, Province del Sara, Bolivia, J. Steinbach coll., no date.

- No. 122, ♀, Province del Sara, Bolivia, J. Steinbach coll., no date.
 No. 123, ♀, Province del Sara, Bolivia, J. Steinbach coll., no date.
 No. 317, ♂, Asumpcion, Bolivia, J. D. Haseman coll., June 14, 1909.
 No. 318, ♂, Campos de Matto Grosso, Braz., Haseman coll., June 20, 1909.
 No. 364, ♀, São João del Rey, Minas Geraes, Brazil, Haseman, no date.

87. *Lachesis peruvianus* Boulenger.

Lachesis peruvianus BOULENGER, Ann. & Mag. Nat. Hist. (7), XII, 1903, p. 334.

The Museum has only one example of this species, a female (C. M. No. 373), which was collected by Miss Lola Vance at Tarma, Peru (elev. 6,000 ft.).

COUNTS AND MEASUREMENTS.

	(No. 373.)
Anal.....	1
Scale-rows.....	23
Gastrosteges.....	184
Urosteges.....	5 + 51/51
Upper labials.....	7
Total length in mm.....	369
Length of tail in mm.....	47

Genus *Crotalus* Linnaeus.

88. *Crotalus terrificus* (Laurenti).

Caudisona terrifica Laurenti, Syn. Rept., 1758, p. 93.

Crotalus terrificus Boulenger, Cat. Snakes, III, 1896, p. 573.

There are thirteen specimens of this species in the collection, as follows:

- C. M. No. 30, ♀, Provincia del Sara, Bolivia, José Steinbach, February, 1913.
 C. M. No. 63, ♀, Provincia del Sara, Bolivia, José Steinbach, October, 1911.
 C. M. No. 118, ♀, Santa Cruz de la Sierra, Bolivia, Steinbach, no date.
 C. M. No. 124, Santa Cruz de la Sierra, Bolivia, Steinbach, head only, mutilated.
 C. M. No. 161, ♀, Bonda, Colombia, H. H. Smith, May.
 C. M. No. 162, ♀, Bonda, Colombia, H. H. Smith, May.
 C. M. No. 259, ♂, Argentina, LeBoutelier Collection.
 C. M. No. 276, ♀ juv., South America, LeBoutelier Collection.
 C. M. No. 1854, ♂, South America, H. H. Smith Coll. (?).
 C. M. No. 2012, ♂, Cacagualito, Colombia, Mrs. H. H. Smith.
 C. M. No. 2013, ♀, Cacagualito, Colombia, Mrs. H. H. Smith.
 C. M. No. 2014, ♀, Cacagualito, Colombia, Mrs. H. H. Smith.
 C. M. No. 2015, ♀, Cacagualito, Colombia, Mrs. H. H. Smith.

COUNTS AND MEASUREMENTS.

Number.	Anal.	Scale-rows.	Gastrosteges.	Urosteges.	Upper Labials.	Scale-rows Between Supra-oculars.	Scale-rows Between Eye and Supra-labials.	Total Length in Mm.	Length of Tail in Mm.	Remarks.
30	1	27	161	27+2/2	13	2	3	620	52	Four rattles.
63	1	29	176	19+2/2	11					
118	1	29	175	21+1/1	14	3	3	598	46	Three rattles.
					14	5	4	376	30	One rattle.
124					13					
					14	3	4			Head only.
					15					
161	1	29		25+3/3	13	3	3	1180+	178	Ten rattles.
162	1	27		23+1/1	13	3	3	1100	100	Skin.
259	1	27	173	20+1/1	14	3	3	1149	115	Nine rattles.
276	1	27	176	22+2/2	16	2	3	295	23	Young.
					17					
1854	1	29	170	29+1/1	13	3	3	364	36	Young.
2012	1	25	161	30	14	3	3	390	41	One rattle.
2013	1	29	174	27+1/1	13	3	3	420	37	One rattle.
2014	1	25	168	30+1/1	15	3	3	410	43	One rattle.
					14					
2015	1	29	179	30+1/1	14	3	3	390	40	One rattle.

In addition to the specimens cataloged above there are contained in the Carnegie Museum three specimens which were received in the LeBoutelier Collection, which are very probably wrongly attributed to the localities given on the original labels. One is a specimen of *Natrix fasciata fasciata* (Linnaeus), C. M. No. 262, labeled as having been obtained in the "sierras of Bolivia." This species has been found as far south as Costa Rica, but has never been reported hitherto from any point south of that country, and in view of the careless labeling of the LeBoutelier Collection I hesitate to record the species as having been actually obtained at the locality indicated.

There are also two specimens of *Agkistrodon piscivorus* (Lacépède), C. M. Nos. 251 and 260, the first labeled "Peru," the second labeled "Brazil." The occurrence of this ophidian in the localities cited is more than doubtful.

EXPLANATION OF PLATE XXVIII.

- Fig. 1. *Atractus taniatus* Griffin. Type, C. M. No. 117. $\times 4$.
FIG. 2. Do. Dorsal View of Head. $\times 4$.
FIG. 3. Do. Ventral View of Head. $\times 4$.
FIG. 4. *Tropidodipsas spilogaster* Griffin. Type, C. M. No. 5085. $\times 2$.
FIG. 5. Do. Dorsal View of Head. $\times 2$.
FIG. 6. Do. Ventral View of Head. $\times 2$.
FIG. 7. *Clelia euprepa* Griffin. Type, C. M. No. 109. $\times 2$.
FIG. 8. Do. Dorsal View of Head. $\times 2$.
FIG. 9. Do. Ventral View of Head. $\times 2$.
FIG. 10. *Elaps hollandi* Griffin. Type, C. M. No. 206. $\times 2$.
FIG. 11. Do. Dorsal View of Head. $\times 2$.
FIG. 12. Do. Ventral View of Head. $\times 2$.

SOUTH AMERICAN OPHIDIA.