Author: Thomas, Joseph James

**Title: The absorption of nitrogen, phosphorus, and
potassium by Pennsylvania cigar-leaf tobacco as
modified by environmental conditions**

Place of Publication:

Copyright Date: 1935

Master Negative Storage Number: MNS# PSt SNPaAg031.8

<104141> * *OCLC* Form:manuscript item 2 Input:CHF Edit:FMD
 008 ENT: 980112 TYP: s DT1: 1935 DT2: LAN: eng
 035 (OCoLC)38092080
 037 PSt SNP aAg031.8 \$bPreservation Office, The Pennsylvania State
 University, Pattee Library, University Park, PA 16802-1805
 090 20 Thesis 1935d \$bThoma,JJ \$cst*7341038 \$cax+(Archival)
 090 20 Microfilm D344 reel 31.8 \$cmc+(service copy, print master, archival
 master)
 100 1 Thomas, Joseph James.
 245 14 The absorption of nitrogen, phosphorus, and potassium by Pennsylvania
 cigar-leaf tobacco as modified by environmental conditions \$ca thesis
 by Joseph James Thomas.
 260 \$c1935.
 300 81 leaves \$bill. \$c29 cm.
 502 Thesis (Ph. D.)--Pennsylvania State College.
 504 Bibliography: leaves 76-81.
 533 Microfilm \$bUniversity Park, Pa. : \$cPennsylvania State University
 \$d1997. \$e1 microfilm reel ; 35 mm. \$f(USAIN state and local literature
 preservation project. Pennsylvania) \$f(Pennsylvania agricultural
 literature on microfilm).
 590 Archival master stored at National Agricultural Library, Beltsville, MD
 : print master stored at remote facility.
 590 This item is temporarily out of the library during the filming process.
 If you wish to be notified when it returns, please fill out a Personal
 Reserve slip. The slips are available in the Rare Books Room, in the
 Microforms Room, and at the Circulation desk.
 650 0 Tobacco \$xAnalysis.
 650 0 Tobacco \$xComposition.
 830 0 USAIN state and local literature preservation project. \$pPennsylvania.
 830 0 Pennsylvania agricultural literature on microfilm.

**BEST COPY
AVAILABLE**

The Pennsylvania State College

The Graduate School

Department of Agricultural and Biological Chemistry

The Absorption of Nitrogen, Phosphorus, and Potassium by
Pennsylvania Cigar-Leaf Tobacco as Modified
by Environmental Conditions

A Thesis

by

Joseph James Thomas

Submitted in partial fulfillment

for the degree of

Doctor of Philosophy

August 1935

Approved: August 1, 1935

W. B. Healy
Professor of Soil and Phytochemistry

1935

R. A. Blustein
Head of the Department

Table of Contents

	Page
List of Tables	iv
List of Figures	vi
I. Introduction	1
II. Historical	2
Description of the Soils	3
III. Object of These Investigations	6
IV. Review of Literature	7
A. General	7
B. Quality of Tobacco in Relation to Fertilizer Practices	9
C. Tobacco Researches in Pennsylvania	10
D. The Interrelationship of the Absorption of Mineral Nutrients as Influenced by Different Fertilizer Treatments	11
V. Methods	14
A. Field Practices	14
B. Chemical Practices	15
VI. Presentation of Data	19
A. General	19
B. The Plan of the Lancaster Experimental Tobacco Plots	19
C. Schedule of Cultural and Chemical Practices	24
D. Precipitation Data	24

Table of Contents (continued)

	Page
E. The Yield, Dry Weight, and Field Observations	28
F. The Chemical Data	40
1. The Soil	40
2. The Plant	40
VII. Discussion of Results	58
A. General	58
B. The Soil	59
C. Rainfall	60
D. Sampling for Chemical Analysis	61
E. The Effects of Different Field and Fertilizer Treatments	61
1. On Tobacco	61
2. On Wheat	69
3. On Clover	70
4. Recapitulation	70
F. The Absorption of Nitrogen, Phosphorus, and Potassium considered from Liebig's "Law of the Minimum"	72
VIII. Summary	73
IX. Acknowledgments	75
X. Bibliography	76

List of Tables	Page
A. Economic Statistics of Pennsylvania Tobacco Production.	20
I. The Field Treatments of the Experimental Plots of the Three-Year Rotation System.	22, 23
II. The Monthly Precipitation for the Lancaster Area from 1932 to 1934 Inclusive, in Comparison with Normal Precipitation.	26
III. The Tobacco, Wheat, and Clover Yield for the Years 1932, 1933, and 1934.	29, 30
III A. Field Notes of the 1932 and 1934 Tobacco Crops.	38, 39
III B. Quality Data of the Cured Tobacco Crops for the Years 1932, 1933, and 1934.	41, 42
IV. The Time of Seeding, Transplanting, Topping, Harvesting, and Sampling the Tobacco Crops for 1932, 1933, and 1934.	25
V. The Chemical, Physical, and Colloidal Analyses of the Soil from the Lancaster Experimental Station, Compared with a Typical Analysis of a Hagerstown Soil.	43, 44

List of Tables (continued)	Page
VI. Precipitation Records from the Time of Transplanting to Maturity as Related to the Intervals of Sampling and Topping.	27
VII. The Weight of Average Middle Leaves of the 1932 and 1933 Tobacco Crops.	35
VIII. The Nitrogen Content of the 1932 and 1933 Tobacco Crops.	46
IX. The Phosphorus Content of the 1932 and 1933 Tobacco Crops.	47
X. The Potassium Content of the 1932 and 1933 Tobacco Crops.	48
XI. The Weight of Average Leaves of the 1934 Tobacco Crop.	36
XII. The Nitrogen Content of the 1934 Tobacco Crop.	49
XIII. The Phosphorus Content of the 1934 Tobacco Crop.	50
XIV. The Potassium Content of the 1934 Tobacco Crop.	51

List of Figures	Page
I. The Lancaster Experimental Tobacco Plots Showing the Various Soil Divisions.	21
II. The Tobacco Yields of the 31 Plots of the Three-Year Rotation for the Years 1932, 1933, and 1934.	31
III. The Wheat Yields of the 31 Plots of the Three-Year Rotation for the Years 1933 and 1934.	32
IV. The Clover Yields of the 31 Plots of the Three-Year Rotation for the Years 1933 and 1934.	33
V. The Dry Weights of the Average Middle Leaves.	37
VI. The Percentages of N, P, and K in 1932 1933, and 1934.	52
VII. The Milligrams of N, P, and K in the Average Middle Leaf in 1932, 1933, and 1934.	53
VIII. The Percentages of N, P, and K in 1934. IX.	54, 55
X. The Milligrams of N, P, and K in the XI. Average Middle Leaf in 1934.	56, 57

THE ABSORPTION OF NITROGEN, PHOSPHORUS, AND POTASSIUM BY
PENNSYLVANIA CIGAR-LEAF TOBACCO AS MODIFIED
BY ENVIRONMENTAL CONDITIONS

I. - Introduction

Pennsylvania annually has a tobacco crop valued at several million dollars. In recent years federal economic policies have tended to reduce acreage and at the same time increase the price, improve the quality, and maintain the value of the crop. The result of these tendencies can be seen by an inspection of Table A which shows an estimated reduction of twenty-six thousand acres of tobacco harvested over a period of four years at a loss of approximately one million dollars.

Because of the present limited acreage and the demand for a better grade of cigar-leaf tobacco at a higher price, it has been deemed advisable to investigate the effect of the various fertilizers on the yield and quality of this important crop. This investigation involves specifically a study of the time-absorption of the three major nutrients-nitrogen, potassium, and phosphorus-by the growing tobacco plant under the influence of various fertilizer combinations.

II. - Historical

Tobacco investigations in Pennsylvania were initiated in 1893 by Dr. William Frear of the Pennsylvania Agricultural Experiment Station. This work was done throughout the various tobacco growing sections of the state. It dealt with fertilizers, curing, variety and strain tests, some shelter-tent experiments with Sumatra tobacco, and economic studies of the cost of tobacco production. In 1916, in cooperation with the United States Department of Agriculture, experiment stations were established at Ephrata and at Lock Haven. The previous work was continued and additional investigations were made on steam sterilization of seed beds, the use of fertilizers on seed beds, seed cleaning, rate of seeding, fumigation, fertilizer treatments, various cultural practices, variety and strain tests, and high and low nicotine breeding. (56,58)^a

A new location for the tobacco experimental station with advantages of accessibility, publicity, and centralization was founded in 1930 nearer the city of Lancaster. The experimental field, consisting of ten acres, is located on the Lefever farm near Roseville, just off the Oregon Pike, on the so-called back road to Eden, one mile from Lancaster city. The field is nearly level with a slight southern exposure, and the soil is of the Hagesstown silt loam type. There has been no fertilizer or lime used on this field for more than twenty years although tobacco was grown there in

1929. The plots were laid out and soil samples taken from each of the plots of the three and four year rotations. A detailed soil survey was made at the same time.

Description of the Soils

Figure I is a map of the experimental field showing the various soil divisions. This map points out two pre--dominating types of soil- Hagerstown silt loam and Hagers--town clay loam. Of the former there are four distinct phases:¹

A. Hagerstown Silt Loam

The Hagerstown silt loam as mapped on the tobacco experimental field is typical of this type as mapped in other parts of the state. The brown silt loam surface soil extends to a depth of 8 to 10 inches. There is practically no change in texture until a depth of 13 to 15 inches is reached but the last five inches is a pale yellow silt loam uncolored with organic matter. This pale yellow layer is a part of the A₂ horizon the upper part has been mixed with the A, and A₀ layers by plowing.

The B horizon is a reddish-yellow friable clay loam which becomes heavier and redder as depth increases, grading into a brick red granular heavy clay. The depth to the bed rock was not determined. It was not encountered at any place on the plots with a 40 inch soil auger.

Quartz fragments were noticed here and there on the plots. This mineral is found in seams or veins within the beds of limestone.

B. Hagerstown Silt Loam, Deep Phase.

This phase differs from the typical in that the dark layer which constitutes the surface soil ranges from 15 inches to 30 inches deep. It occupies one good sized area at the foot of a slope and is partly colluvial in origin. The center of this area is where the 30 inch borings were obtained which thin out to 15 inches on the outer edges.

A pale yellow silt loam was encountered between the brown and the reddish yellow clay loam horizons. The latter was found at from 25 to 36 inches from the top.

C. Hagerstown Silt Loam, Heavy Subsoil Phase.

This separation is very similar to the normal type except that the brown surface soil rests directly upon the reddish yellow clay loam B horizon. The dull brick red friable clay is encountered at about 24 inches below the surface.

The soil in most cases seems to separate the clay loam member of the series from the typical silt loam member.

D. Hagerstown Silt Loam, Shallow Phase.

The shallow phase as the name implies is characterized by a shallow brown surface soil which does not exceed 6 inches in depth and averages only about 4 inches. This grades through a reddish-yellow clay loam to the usual brick red clay.

The shallow phases only occur in two very

small patches in Tier III of the four year rotation. The long narrow area probably marks the location of an old back furrow.

E. Hagerstown Clay Loam.

The Hagerstown clay loam occupies the higher land in the central part of the tract. It is characterized by a reddish-brown clay loam surface soil which grades at about 8 to 10 inches into a granular heavy red clay subsoil. Fragments of quartz are present in a few places but nowhere in any appreciable amounts.

The clay loam member of the Hagerstown series is a B-C soil rather than an A-B-C one. It represents what is left after the surface soil has been removed by erosion.

Information relative to climate, agricultural practices, and relation of soils to geology are well discussed in Soil Survey of Lancaster County. (32)

Table V² gives the results of the chemical, physical, and colloidal analyses of composited soil samples from the three year rotation. One composite consists of samples from each of the thirty-one plots on Tiers I and II of the three year rotation; the other, from each of the thirty-one plots of Tier III of the three year rotation. There is also given a comparative analysis by Thomas (62) of a typical Hagerstown soil.

-
- a. Numbers in parentheses indicate bibliography.
 1. Descriptions of the soils taken from Soil Survey Report by Dr. A. L. Patrick and C. C. Lewis.
 2. cf. bibliography, Claffey (8).

III. - Object of These Investigations

The object of these investigations is: first, to study the rate and quantity of absorption of nitrogen, potassium, and phosphorus by the tobacco plant when the soil is fertilized with various combinations of these nutrients with and without manure; second, to show the relation of this absorption to the yield and quality of tobacco.

IV. - Review of Literature

A. General

Soil, climate, seed, and cultural practices constitute an important quartet of factors which determines the type, kind, and quality of tobacco that is produced in any part of the world. Since soil and climate are fixed for local areas, each tobacco growing section produces its own characteristic type of tobacco. Any change in these two factors produces a profound effect on the finished product. For this reason, tobacco researches are more or less localized in each region where a particular type of tobacco is grown. For example, in the Connecticut valley where fine shade-grown wrappers are produced, the results of researches with respect to the kinds and amounts of fertilizers to be employed would not apply to the heavy type of cigar-filler leaf that is produced in Pennsylvania, or to the burly cigarette types produced in the Southern tobacco areas of the United States. Garner (27) has fully discussed these relationships and their chemical aspects recently.

The literature on the several phases of tobacco research- growth, mineral metabolism, fermentation, and manufacture- is voluminous so that only a portion of it can be covered here. The roles of the several mineral nutrients in plant metabolism have been ably discussed in several texts on plant biochemistry, plant physiology, and agronomy (50, 60, 69, 75). For the latest theories on the mineral nutrition

of plants see (9).

The tobacco plant is characterized by its relatively high ash content which usually places tobacco in the category of soil exhausting crops (72). The composition of this ash is readily affected by a change in the composition of the soil solution chiefly brought about by the addition of fertilizers to the soil (1). Since burning quality is a function of the mineral constituents of the leaf, the quality of the tobacco varies directly with the fertilizer treatments (34, 35, 36, 37). It is conclusive that the types and amounts of fertilizers used affect the yield (1, 46, 53). Therefore the ideal fertilizer for tobacco is one which gives the optimum ratio of mineral nutrients in the soil solution, and in such a concentration that both the yield and the quality of the tobacco are at a maximum. In other words, a physiologically balanced nutrient solution should be provided.

Because the tobacco plant is a heavy feeder, and because the growing plant shows the effect of any deficiency of certain essential elements, it constitutes a good indicator for the availability of fertilizer elements in the soil (30, 31, 44, 45, 51). Davidson (19) gives analyses of parts of the tobacco plant at different stages of growth. Kosutany (40) gives the same data with dry weight. The reports of the Connecticut station (2, 3, 4, 5, 6, 7, 8, 38, 39, 52) are most comprehensive in their discussions of experimentation regarding fertilizer problems,

cultural practices, mineral metabolism, and disease control. These apply especially to tobacco production in the Connecticut valley. Anderson (6) discusses at length and describes fully experiments dealing with potash problems in growing quality tobacco. The nitrogen nutrition of tobacco has been taken up by Garner (28) and by Beaumont (13). McMurtrey (46) recently described fertilizer tests using varying forms and amounts of nitrogen and potassium. Ames and Boltz (1), discussing the influence of fertilizers on the composition and quality of tobacco, have described conditions which are especially applicable to Pennsylvania cigar-leaf production. Garner (29) and Brown (16) have described the effects of other crops in the rotation on tobacco. Beaumont (14) has described field experiments; Moss (53) dealt with the fertilization of flue-cured tobacco.

B. Quality of Tobacco in Relation to Fertilizer Practices.

As the result of the work of Schlösing (61), Nessler (54), Van Bemmelen (70), Mayer (47, 48, 49), Barth (11), Behrens (15), Garner (26), and Haley (34, 35, 36, 37), the importance of the presence of potassium in both organic and inorganic combinations in the tobacco leaf has been established in relation to burning quality. That the presence of excessive amounts of chlorine exerts a deleterious effect on burning quality has been proved by Mayer (47, 48, 49), Fesca (20), Garner (26), Ames and Boltz (1), Olson (58),

Jenkins (39). Garner (26) showed a relationship between water extracts and burning quality, while Graham and Carr (33) demonstrated that ether, alcohol, and hot water extractions of leaves could be correlated with burning quality.

C. Tobacco Researches in Pennsylvania.

For a general description of Pennsylvania tobacco production--soils, climate, types of tobacco grown, and cultural practices--see Frear (25). Some of the facts (18, 21, 23, 24, 25, 34, 35, 36, 37, 55, 56, 57, 58, 59, 71) which have been established regarding the Pennsylvania cigar-leaf tobacco from 1893 to the present time are: development of strain^s and varieties which are adapted to Pennsylvania conditions, and which have the desirable features of size, shape, and number of leaves, disease resistance, and high and low nicotine content; proper sterilization of seed beds to control diseases; proper means of seed cleaning, rate of seeding, seed bed fertilization and fumigation; proper planting distances and topping heights; and proper curing and sweating methods. It has been shown also that the chemical composition of the leaf as influenced by fertilizer treatment is efficacious in determining the burning quality; that high yield and good quality of tobacco can be obtained with the application of commercial fertilizers which may or may not be supplemented with manure; that manure alone does not give as satisfactory results as a combination of manure and commercial fertilizers.

D. The Interrelationship of the Absorption of Mineral Nutrients as Influenced by Different Fertilizer Treatments.

Liebig (42, 43) in 1843 stated in what is now known as "The Law of the Minimum",

"By the deficiency or absence of one necessary constituent, all others being present, the soil is rendered barren for all those crops to the life of which that one constituent is indispensable."

It therefore followed that nutrient elements are absorbed in certain definite proportions in such a manner that a limit having been reached by one, the absorption of the others is retarded. It also follows that if the rate of absorption of the nutrients be plotted against time, the curves for a completely balanced fertilized crop will always be on a higher plane than those for a crop which has been fertilized by an unbalanced supplement--provided, of course, all other growth factors are constant.

Thomas, in a series of papers, (63, 64, 65, 66, 67, 68) has discussed Liebig's "Law of the Minimum" and has applied it to the interpretation of his work with apple trees. Lagatu and Maume, (41) however, maintain that the corrolary of Liebig's law is not capable of generalization as a result of their work with the grape. They concluded that the depression in yields produced by the application of an incomplete fertilizer is not due to a depression of the absorption of the remaining elements but, on the contrary, to a nutritional lack of balance owing to an increased absorption of these elements. Thomas (61), on the

other hand, accounts for this apparent anomaly by stating that the element omitted from the fertilizer is already present in the soil solution in such a concentration that its critical concentration in the plant is never reached--that the whole problem is one of relativity. Bartholomew (12), working with tomato plants in nutrient sand cultures, maintains that though theoretically possible to set up a physiologically balanced nutrient solution, it would be difficult to adjust the varying concentrations which would be necessary to keep pace with growth changes and demands; that nutrient solutions, varying greatly in composition, produce similar results; that the amounts (Mgms.) of nitrogen, phosphorus, and potassium absorbed by plants, receiving all elements essential for growth, were controlled to a great extent by the concentrations of those elements in the nutrient media; that a deficiency of potassium in the nutrient media may result in an increased absorption of nitrogen and phosphorus by the leaves of the tomato plant; that the use of percentage composition of a plant may lead to erroneous conclusions; that Liebig's "Law of the Minimum" is not applicable to fertilization with potassium salts. Carolus (17) also working with tomatoes on field plots receiving various fertilizers, concluded that the lack of N, P, or K in field fertilizer applications did not result in as marked changes in composition with respect to those elements as ^{occurs} ~~the curves~~ under nutrient solutions. The effect of the scarcity or abundance of one ingredient in the

fertilizer on the absorption of the other ingredients was as significant in many cases as the effect of the ingredient on its own absorption. The inclusion of N or P in the fertilizer resulted in an increased absorption of all of the constituents determined. The complete lack of potassium in the 4-10-0 mixture was not as detrimental to growth and absorption as was the presence of the maximum addition in a 4-10-12 mixture. The author states his work neither refutes nor substantiates the application of Liebig's "Law of the Minimum" to potassium fertilization.

V. - Methods

A. Field Practices.

In 1932 the new location for the tobacco experimental station at Lancaster was used for the first time. The work dealt with the three year rotation of tobacco, wheat, and clover with the exception of plots 29, 30, 31 on which alfalfa replaced clover. At this time tobacco was grown on tier I, clover on tier II, and tier III was bare.

After the proper sterilization of the seed beds (57), the Swarr-Hibshman strain of the Pennsylvania seed-leaf was sowed, transplanted, topped, harvested, and cured. The usual cultural practices of growing cigar-filler tobacco in Lancaster County were followed (25).

Plots, one twenty-first of an acre in area, were laid out; the plants were placed at twenty-six inch intervals in rows forty-two inches apart. Proper methods in the control of insects and diseases were followed as they were needed. Throughout the growing season field observations were made. Weights for the yield and criteria of the quality were determined after curing and stripping.

The fertilizer treatments of the three year rotation are as follows:

1. Varying amounts and kinds for the tobacco.
2. A uniform treatment of 300 pounds of 16 per cent superphosphate per acre for the wheat.
3. Clover, no treatment.

Manure is applied approximately one month before the tobacco is transplanted; the other fertilizers, with the exception of nitrate of soda, which is applied as a side dressing approximately one month after transplanting, are broadcast about one week before transplanting. However, plots 16, 17, and 18 are fertilized in the row at the time of transplanting.

The following is a list of fertilizers used:

1. Nitrogen

- a. Two-thirds from cotton-seed meal.
- b. One-third, nitrate of soda, except on plot 14 where two-thirds of the nitrogen comes from nitrate of soda.

2. Phosphoric acid

Sixteen per cent superphosphate

3. Potash

Sulphate of potash

4. Manure

Stock yard with very little straw

5. Lime

Hydrated

B. Chemical Practices.

Samples of tobacco leaves for chemical analyses were taken every ten days throughout the growing season, starting when the plants were of such a size that eleven leaves constituted a sample large enough for the several analyses. The first sampling was usually forty to

fifty days after transplanting, and continued until harvest, with the exception of 1932 when eleven plants from each plot were allowed to remain in the field several days after harvesting. Each of these samples consisted of eleven leaves, the sixth leaf from the bottom or the eighth from the top, taken from eleven different representative plants from each plot. Care was exercised that no plant was sampled more than once during any growing season. The sample was immediately weighed, packed in dry ice, and shipped to State College where it was placed in an oven at 70 to 80 ° C. for twelve to twenty-four hours. It was again weighed, the whole leaf (lamina and midrib), ground in a Wiley mill so that more than ninety percent passed a sixty-mesh screen, about ninety-eight percent passed a forty-mesh screen, and a hundred percent passed a twenty-mesh screen. The sample was then sealed and the chemical analyses for moisture, nitrogen, phosphorus, and potassium were made.

(1) Moisture analysis

A one gram sample was dried in vacuo over concentrated sulphuric acid to constant weight (10).

(2) Nitrogen analysis

The Kjeldahl-Gunning-Arnold modification Method to include nitrates (10).

(3) Preparation of solution for phosphorus and potassium.

A five gram sample in a porcelain crucible was placed in a muffle furnace regulated so as not to exceed

a temperature of 480°C . for twenty-four hours. The ash was then taken up in dilute nitric acid and dilute hydrochloric acid, filtered, and washed with hot dilute nitric acid (1 + 99). The filter paper and the residue were replaced in the crucible and burned in the muffle furnace at a full red heat. This ash was then taken up with a mixture of concentrated hydrochloric and nitric acids, boiled, diluted, filtered, and washed with hot dilute nitric acid (1 + 99). The filtrate and washings were made up to a volume of 500 milliliters. A 50 ml. aliquot was then taken for the potassium determination, and a 100 ml. aliquot for the phosphorus determination.

(4) Phosphorus

The volumetric method, with precipitation at 45° to 50°C . was followed (10).

(5) Potassium

To the 50 ml. aliquot in a porcelain dish was added 1 ml. of dilute sulphuric acid (1 + 1). The solution was evaporated to dryness on the steam bath. The dish was then placed in the muffle furnace and cautiously heated until the residue was white. The residue was taken up with 1 ml. of concentrated hydrochloric acid, and hot water added, stirring until solution was complete. Then 5 mls. of platinic chloride ($.02 \text{ gm. per ml.}$) were added and the solution evaporated on the steam bath just to dryness. The residue was taken up with 25 mls. of acid--ulated alcohol (2.25 normal hydrochloric acid in eighty

percent alcohol). The mixture was thoroughly stirred. After standing about one-half hour, the contents of the dish were filtered through a weighed gooch crucible, washed with eighty percent alcohol, with Lindo-Gladding solution, again with eighty percent alcohol, dried at 100°C for one-half hour and then weighed.

All results are given on the moisture-free basis. All calculations are made to the elements N, P, and K. The weight of the "Average Middle Leaf" is the mean weight in grams of the eleven leaves constituting each sample. The milligrams of nutrient per average middle leaf was obtained by multiplying the weight in milligrams of the average middle leaf by the percentage of that nutrient.

VI. - Presentation of Data

All Chemical Results are expressed on the moisture-free basis.

A. General

Table A gives figures on the acreage harvested, average yield per acre, total production, average price per pound, and the valuation of Pennsylvania tobacco for the period 1930 to 1934. These data were taken from the Yearbooks of Agriculture, United States Department of Agriculture, 1931 to 1935. Over 90 per cent of the tobacco produced in Pennsylvania is grown in the Lancaster area.

B. The Plan of the Lancaster Experimental Tobacco Plots.

Figure I is a map of the experimental field with the various soil divisions indicated. The three-year rotation area is divided into three tiers of thirty-one plots each, so that every year the three crops--tobacco, wheat, and clover--are represented, as follows:

	<u>Tier I</u>	<u>Tier II</u>	<u>Tier III</u>
<u>1932</u>	Tobacco	Clover	Bare
<u>1933</u>	Wheat	Tobacco	Clover
<u>1934</u>	Clover	Wheat	Tobacco

Table I shows the field and fertilizer treatments that are given to the thirty-plots of the three-year rotation. These various fertilizer combinations and practices were designed to show the effects of varying amounts and kinds of nitrogen, of varying amounts of phosphorus and

Table A
Economic Statistics of Pennsylvania Tobacco Production.

Year	Acreage Harvested	Yield per acre	Production	Ave. price per pound	Value
	Acres	Pounds	1000 pounds	cents	Dollars
1930	41,300	965	39,854	6.4	2,550,656
1931	40,900	1410	57,669	7.4	4,265,060
1932	42,100	1100	46,316	5.2	2,408,432
1933	25,300	1050	26,563	5.5	1,460,965
1934	15,200	1152	17,504	7.5	1,312,800

Data taken from Yearbooks of Agriculture, U. S. Dept. of Agriculture, 1931-1935.

Fig. 1. The Lancaster Experimental Tobacco Plots Showing the Various Soil Divisions.

Table I

The field treatments of the experimental plots, Tiers I, II, and III of the three-year rotation system. All figures are given on the basis of one acre.

Plot Numbers for Tiers I, II, III	Manure	Fertilizer	Formula	Lime	Fertilizer Applied	Reference
	tons	lbs.		lbs.		
1	----	1,000	6-8-4	----	Broadcast	N P 1/3 K ^b
2	----	1,000	6-8-8	----	"	N P 2/3 K
3	----	1,000	6-8-12	----	"	N P K
4	----	1,000	6-8-16	----	"	N P 4/3 K
5 ^c	----	1,000	0-8-12	----	"	P K
6	----	1,000	3-8-12	----	"	1/2 N P K
7	----	1,000	9-8-12	----	"	3/2 N P K
8	----	1,000	6-0-12	----	"	N N K
9	----	1,000	6-4-12	----	"	N 1/2 P K
10 ^c	----	1,000	0-8-12	----	"	P K
11	----	500	6-8-12	----	"	1/2 N 1/2 P 1/2 K
12	----	1,500	6-8-12	----	"	3/2 N 3/2 P 3/2 K
13	----	1,000	6-8-12	1,000	"	N P K L
14 ^a	----	1,000	6-8-12	----	"	N P K
15 ^c	----	1,000	0-8-12	----	"	P K
16	----	500	6-8-12	----	In Row	1/2 N 1/2 P 1/2 K-R
17	----	1,000	6-8-12	----	" "	N P K-R
18	----	1,500	6-8-12	----	" "	3/2 N 3/2 P 3/2 K-R
19 ^c	----	-----	-----	----	-----	Check - 0
20	----	1,000	0-8-12	----	Broadcast	P K
21	10	500	0-8-12	----	"	1/2 P 1/2 K M

Table I (continued)

Plot Numbers for Tiers I, II, III	Manure	Fertilizer	Formula	Lime	Fertilizer Applied	Reference
	tons	lbs.		lbs.		
22	10	1,000	3-8-12	----	Broadcast	1/2 N P K M
23	10	1,000	3-8-8	----	"	1/2 N P 2/3 K M
24	10	500	3-8-12	----	"	1/4 N 1/2 P 1/2 K M
25 ^c	----	1,000	0-8-12	----	"	P K
26	10	1,000	3-8-16	----	"	1/2 N P 4/3 K M
27	10	-----	-----	-----	-----	M
28	20	-----	-----	-----	-----	2 M
29 ^d	-----	-----	-----	-----	-----	Check - 0
30 ^{c,d}	-----	1,000	0-8-12	-----	Broadcast	P K
31	-----	1,000	6-8-12	-----	"	N P K

Reference Key: N = 60 lb. nitrogen (N₂) /acre
P = 80 lb. phosphoric acid (P₂O₅) /acre
K = 120 lb. potash (K₂O) /acre
L = 1,000 lb. lime (CaCO₃) /acre
M = 10 tons manure/acre

- a. - Two-thirds of nitrogen as nitrate of soda.
- b. - Fertilizer broadcast unless indicated by symbol "R" which designates row application.
- c. - Replicates.
- d. - Alfalfa in rotation instead of clover.

potassium, of varying the amounts of the same fertilizer, of row versus broadcast application, of using manure with and without varying fertilizers, of demonstrating the effect of lime, and of replacing clover with alfalfa. Spaced at definite intervals are six replicate plots which receive a basic fertilizer treatment. The purpose of these is to show the variation in the natural fertility of the soil. There are two plots which receive no fertilizer treatment, designated as checks.

C. Schedule of Cultural and Chemical Practices.

Table IV gives the dates and the age in days from transplanting, of seeding, transplanting, topping, harvesting, and sampling the tobacco crops for the years 1932, 1933, 1934. It can be observed from the table that the growing period in 1932 was longer than that of either 1933 or 1934. This can be partially explained by the influence of climatic factors, and partially by the necessity of harvesting earlier in 1933 and 1934 because of the rapid spread of diseases. In 1932, eleven plants on each plot were allowed to remain in the field for seven days, after harvesting. The leaves taken from these plants constituted the sixth sampling, and were designated as being post-mature.

D. Precipitation Data

Table II gives the amounts of rainfall at monthly intervals for the years 1932, 1933, and 1934. The departure from normal is also indicated. Table VI gives the incidences and amounts of rainfall relative to the cultural

Table IV

The time of seeding, transplanting, topping, harvesting and sampling the tobacco crops for 1932, 1933 and 1934.

	1932		1933		1934	
	Date	Age in Days	Date	Age in Days	Date	Age in Days
Seed Bed Sowed	April 15	-52	April 10	-61	April 10	-59
Transplanted	June 6	0	June 10	0	June 8	0
Topped	Aug. 22	77	Aug. 20	71	Aug. 7	60
Harvested	Sept. 12	98	Aug. 28	79	Aug. 24	77
First Sampling	July 25	49	July 29	49	July 24	46
Second Sampling	Aug. 6	61	Aug. 8	59	Aug. 3	56
Third Sampling	Aug. 17	72	Aug. 18	69	Aug. 13	66
Fourth Sampling	Aug. 29	84	Aug. 28	79	Aug. 24	77
Fifth Sampling	Sept. 7	93	-----	--	-----	--
Sixth Sampling	Sept. 19	105	-----	--	-----	--

Age of plants refers to days from transplanting date.

Table II

Precipitation records compiled by the U. S. Weather Bureau for the Lancaster area from 1932 to 1934 inclusive, in comparison with normal precipitation.

Month	1932	Departure from normal	1933	Departure from normal	1934	Departure from normal
	inches	inches	inches	inches	inches	inches
January	4.29	+1.05	1.89	-1.39	2.41	-0.83
February	1.89	-0.82	2.53	-0.18	2.10	-0.61
March	6.19	+2.86	6.13	+2.80	4.55	+1.22
April	2.41	-2.08	5.07	+1.58	3.98	+0.49
May	3.99	-0.76	4.86	+1.63	4.35	+1.12
June	4.28	+0.31	2.66	-1.31	3.51	-0.46
July	3.61	-0.27	9.13	+5.25	5.41	+1.53
August	5.23	-1.43	13.94	+10.11	4.39	+0.59
September	1.67	-1.69	4.52	+1.16	9.83	+6.47
October	8.39	+5.28	2.16	-0.95	1.36	-1.75
November	4.13	+1.60	0.63	-1.90	3.40	+0.87
December	3.14	+0.02	2.74	-0.38	2.58	-0.54
Total	48.22	+8.45	56.22	+16.45	47.87	+8.10

Table VI

Precipitation records from the time of transplanting to maturity as related to the intervals of sampling and topping. These figures are based on observations made by the U. S. Weather Bureau for the Lancaster area for 1932, 1933 and 1934.

Time Interval	Year					
	1932		1933		1934	
	Interval in days	Rainfall in inches	Interval in days	Rainfall in inches	Interval in days	Rainfall in inches
Transplanting to 1st sampling	0-43	5.78	0-43	11.40	0-45	6.61
1st sampling to 2nd sampling	49-60	2.12	49-58	4.11	46-55	1.37
2nd sampling to 3rd sampling	61-71	3.56	59-68	2.31	56-65	2.38
3rd sampling to 4th sampling	72-83	1.08	69-79	7.52	66-77	0.99
4th sampling to 5th sampling	84-92	1.32				
5th sampling to 6th sampling	93-105	0.00				
Total	0-105	13.86	0-79	25.34	0-77	11.35
Transplanting to Topping	0-76	12.34	0-70	17.89	0-59	6.34
Topping to Harvest	77-98	1.52	71-79	7.45	60-77	3.01
Total	0-98	13.86	0-79	25.34	0-77	11.35

and chemical practices of transplanting, topping, harvesting, and sampling for the years 1932, 1933, and 1934.

E. Yield, Dry Weight, and Field Observations.

1. The Yields

a. Tobacco.

Table III gives the tobacco yields of the thirty-one plots of the three-year rotation for the years 1932, 1933, and 1934. Also shown is the average tobacco yield for each plot the three years, the average yield for all plots for each year and for the entire period. Figure II graphically presents these data.

b. Wheat.

Table III also gives the wheat yields of the thirty-one plots of the three-year rotation for the years 1933 and 1934. Also shown is the average yield for each plot for the two-year period, the average for each year, and for the two-year period. Figure III graphically presents these data. No data are given for wheat in 1932 because the experiment was not begun in sufficient time to allow for its being planted.

c. Clover.

Table III also gives the clover yields of the thirty-one plots of the three-year rotation for the years 1933 and 1934. Also shown is the average yield for each plot for the two-year period, the average for each year, and the average for the two-year period. Figure IV graphically presents these data. No data are given for clover in

Table III

The Tobacco, Wheat and Clover Yields for the Years 1932, 1933 and 1934. All Figures are given on the basis of one acre.

Plot No.	Tobacco				Wheat			Clover		
	Tier I	Tier II	Tier III	Ave.	Tier I	Tier II	Ave.	Tier I	Tier III	Ave.
	1932	1933	1934		1933	1934		1934	1933	
	Lb.	Lb.	Lb.		Bu.	Bu.		Lb.	Lb.	
1	1783	1218	1743	1581	27.30	30.00	28.7	6680	6720	6700
2	1911	1365	1890	1722	30.40	26.40	28.4	7230	7860	7545
3	1925	1407	1964	1765	28.00	30.50	29.3	7390	7860	7625
4	1827	1099	1617	1514	30.10	32.50	31.3	6980	6720	6850
5	1281	1070	1502	1284	23.80	26.60	25.2	6880	6720	6800
6	1596	1218	1386	1400	26.60	31.30	29.0	6950	6720	6835
7	1596	1218	1533	1449	27.60	30.80	29.2	7490	7860	7675
8	1554	966	1313	1278	29.80	28.00	28.9	6780	6720	6750
9	1638	1155	1481	1425	28.70	26.80	27.8	6490	5040	5765
10	1512	1008	1449	1323	29.20	25.30	27.3	6870	7860	7365
11	1743	1050	1470	1418	27.40	27.60	27.5	6450	5040	5745
12	1806	1281	1670	1586	30.40	33.10	31.8	6190	5880	6035
13	1638	1134	1817	1530	28.40	26.40	27.4	6230	5880	6055
14	1743	1134	1743	1540	30.80	26.30	28.6	6190	5880	6035
15	1554	1134	1418	1369	27.30	31.20	29.3	5670	5040	5355
16	1869	1239	1691	1600	37.50	28.20	32.9	6640	6720	6680
17	2352	1386	1974	1904	33.60	34.30	34.0	6120	5880	6000
18	2583	1512	2048	2048	27.60	30.90	29.3	6090	5880	5985
19	1050	987	1355	1131	20.60	30.70	25.7	5040	5040	5040
20	1680	1155	1701	1512	24.90	27.30	26.1	5680	5040	5360
21	1764	1260	1869	1631	33.20	31.10	32.2	6310	5880	6095
22	2205	1428	1917	1843	30.40	27.30	28.9	6420	5880	6150
23	2056	1260	1869	1728	26.20	33.50	29.9	6450	5880	6165

Table (continued)
III

Plot No.	Tobacco				Wheat			Clover		
	Tier I	Tier II	Tier III	Ave.	Tier I	Tier II	Ave.	Tier I	Tier III	Ave.
	1932	1933	1934		1933	1934		1934	1933	
	Lb.	Lb.	Lb.		Bu.	Bu.		Lb.	Lb.	
24	1869	1260	1764	1631	35.40	27.10	31.3	6090	5040	5565
25	1680	1260	1901	1614	31.10	26.00	28.6	5990	6720	6355
26	2163	1134	1869	1722	32.90	37.00	35.0	5300	5040	5170
27	1869	1260	1691	1607	33.90	35.70	34.8	5990	5880	5935
28	1932	1407	2037	1792	34.70	35.40	35.1	6010	5880	5945
29	1155	966	1439	1187	24.10	36.90	30.5	5690	5880	5785
30	1659	1134	1712	1502	25.90	28.70	27.3	5990	5040	5515
31	1785	1260	2091	1725	32.90	29.60	31.3	6240	5880	6060
Ave.	1767	1205	1708	1560	29.40	30.08	29.74	6339	6108	6224

Fig. II. The Tobacco Yields of the 31 Plots of the Three-year Rotation for the Years 1932, 1933 and 1934.

The average yield is shown for each plot and for each year, as well as the mean yield for the entire period.

Fig. III. The Wheat Yields of the 31 Plots of the Three-year Rotation for the Years 1933 and 1934.

The average yield is shown for each plot and for each year, as well as the mean yield for the two years.

Fig. IV. The Clover Yields of the 31 Plots of the Three-year Rotation for the Years 1933 and 1934.

The average yield is shown for each plot and for each year as well as the mean yield for the two years.

1932 because weeds constituted the bulk of the yield, and hence the figures are not accurate.

2. The Dry Weights (Tobacco).

Table VII contains the dry weights of the average middle leaf from representative plots in 1932 and 1933, for samples taken at the intervals indicated in Table IV. Table XI gives the same data for the thirty-one plots in 1934 and also the average dry weights at the several samplings. Figure V graphically presents the contents of Tables VII and XI.

3. Field Data

a. During Growth.

Table III A presents the cost of the fertilizer and manure (based on 1932 prices, not including the cost of hauling, mixing, or applying), observation on the growth, the color, the prevalence of diseases, and the state of maturity, at harvest, for the tobacco on the several plots of the three-year rotation for the years 1932 and 1934. The 1933 crop is not included because the disease, wildfire, infected the plants about the seventieth day after transplanting. Two severe storms, one July 26, and the other, August 23, rapidly spread the causative organism and necessitated harvesting before the plants had properly matured. Commercially the tobacco crop was a complete failure. In 1934 almost the same situation arose, except that the crop had practically matured so that the early harvesting prevented the further spread of and damage

Table VII

The weight of average middle leaves of the 1932 and 1933 tobacco crops as based on samples taken from representative experimental plots at different intervals during the growing season. The results are expressed on the moisture-free basis.

Plot No.	Treatment	Year	Sampling					
			First	Second	Third	Fourth	Fifth	Sixth
			Dry Matter gms	Dry Matter gms	Dry Matter gms	Dry Matter gms	Dry Matter gms	Dry Matter gms
1	N P 1/3 K	1932	2.68	4.87	6.99	7.88	9.35	9.09
		1933	1.55	5.28	6.25	7.62	----	----
3	N P K	1932	2.71	4.77	8.09	----	10.00	10.14
		1933	2.32	5.96	7.60	8.78	-----	-----
5	P K	1932	1.52	2.92	5.43	7.76	8.08	9.58
		1933	3.32	3.23	5.41	6.99	----	----
8	N K	1932	1.52	2.85	4.89	7.29	7.46	9.57
		1933	1.92	4.09	5.42	7.63	----	----
19	Check	1932	----	----	----	----	----	----
		1933	2.34	3.13	5.49	7.24	----	----

Table XI

The weight of average leaves of the 1934 tobacco crop as based on samples taken from the experimental plots at different intervals during the growing season. The results are expressed on moisture-free basis.

Plot No.	Sampling			
	First	Second	Third	Fourth
	Gm. per Leaf	Gm. per Leaf	Gm. per Leaf	Gm. per Leaf
1	5.56	7.20	9.81	12.27
2	5.27	6.02	9.91	14.06
3	4.65	6.69	11.82	11.91
4	4.82	6.20	9.59	12.92
5	3.60	5.00	8.51	11.60
6	4.28	4.97	8.34	12.29
7	4.58	6.18	8.53	11.42
8	4.27	5.76	6.80	10.11
9	5.43	5.87	8.02	9.66
10	4.37	5.83	8.86	10.82
11	3.60	5.26	8.37	12.07
12	5.45	6.94	8.41	11.79
13	4.40	6.75	9.77	13.29
14	4.05	6.92	8.90	12.17
15	4.29	6.83	7.58	10.64
16	4.17	6.62	7.18	12.05
17	3.67	7.02	8.89	12.73
18	5.13	5.93	11.33	14.01
19	3.38	5.31	9.01	11.75
20	4.21	6.98	9.40	11.90
21	3.68	5.50	8.60	12.64
22	4.84	7.38	10.84	13.19
23	3.33	6.17	9.25	13.46
24	4.29	7.14	10.19	12.56
25	5.10	6.86	10.01	10.75
26	4.76	6.70	9.24	13.86
27	3.77	6.60	8.85	14.77
28	4.52	7.12	10.86	13.79
29	3.46	6.67	9.41	13.96
30	4.76	7.35	11.32	14.76
31	4.97	7.60	10.23	14.57
Average	4.41	6.43	9.28	12.51

Fig. V. The Dry Weights of the Average Middle Leaves.

The figures are based on samples taken throughout the growing season from representative plots in 1932 and 1933, and from the 31 plots in 1934. The average for 1934 is also given. The first sample was taken in 1932 on the 49th. day after transplanting; in 1933, on the 49th. day, and in 1934, on the 46th. day.

Table III A

Field Notes of the 1932 and 1934 Tobacco Crops^a.

Plot No.	Cost of Fertilizer ^b	Growth		Color		Diseases		Maturity at Harvest	
		1932	1934	1932	1934	1932 ^c	1934 ^d	1932	1934
1	\$14.60	5	5	D	D	0	3 ^e	2	
2	15.64	7	5	D	D	0	3	2	
3	17.70	5	6	D	D	0	2	2	
4	22.64	5	4	D	M	0	2	2	
5	7.29	1	4	M	M	2	1		
6	12.49	4	3	D	D	0	2		
7	23.05	4	5	V D	V D	0	2		
8	14.05	2	2	M	M	0	1		
9	15.83	4	4	D	D	0	1	0	
10	7.29	2	4	M	M	0	1	1	
11	8.85	5	4	D	D	0	1	1	
12	26.55	5	5	V D	V D	0	2	0	
13	17.70	5	5	D	V D	0	2	1	
14	16.32	5	5	D	D	0	2	2	
15	7.29	2	4	M	M	0	2	0	
16	8.85	6	5	D	D	0	2	2	
17	17.70	7	6	D	D	0	1	2	
18	26.55	8	6	V D	D	0	1	2	
19	-----	0	2	L(K)	L	3	2	0	
20	7.29	4	4	D	M	0	2	1	
21	27.29	5	5	D	D	0	2	2	
22	32.40	7	5	V D	D	0	1	2	
23	28.83	7	5	D	D	0	2	2	
24	26.25	5	5	D	D	0	2	2	
25	7.29	5	5	D	M	0	3	2	

Table III A (cont.)

Plot No.	Cost of Fertilizer ^b	Growth		Color		Diseases		Maturity at Harvest	
		1932	1934	1932	1934	1932 ^c	1934 ^d	1932	1934
26	\$28.68	7	5	D	V D	0	2	2	
27	20.00	5	5	D	D	2	3	1	
28	40.00	5	6	D	I D	2	2	1	
29	-----	1	4	L(K)	L	2	3	0	
30	7.29	4	5	D	D	0	2	2	
31	17.70	5	7	D	D	0	1	2	

- a. Commercially, the 1933 crop was a total failure because of damage by wildfire, spread by severe storms.
- b. Does not include cost of hauling, mixing and applying. 1932 prices. Cost per acre.
- c. Mosaic in 1932.
- d. Wildfire leaf spots in 1934.
- e. Plots 1, 2 and 3 severely damaged by wildfire in 1934 because of proximity to infected Rustica plots.
- f. Plots 14 and 19 in 1934—considerable ring spots.

Key

Growth	Color (Green)	Diseases	Maturity at Harvest
0-Very Sorry	L-Light	0-None	0-Unripe
1-Very Poor	M-Medium	1-A Very Few	1-Almost Ripe
2-Poor	D-Dark	2-Few	2-Ripe
3-Rather Poor	V D-Very Dark	3-Numerous Infections	
4-Fairly Good	I D-Intensely Dark		
5-Good	(K)-Potash Hunger		
6-Very Good			
7-Fine			
8-Excellent			

by wildfire, and at the same time lowered only slightly the commercial value of the crop. However, the tobacco on plots 1, 2, and 3 were rather severely damaged because of their proximity to the initial source of infection--the Rustica plots.

b. After Curing.

Table III B gives data on several criteria of quality--viz, burn (interval of holding fire, spread of fire, and character of ash), elasticity, body, color, and the percentages of wrappers, fillers, and seconds for the years 1932 and 1934. No data ^{are} ~~is~~ given for 1933 for the reasons outlined in "a" of this section. The data for 1934 should be interpreted in the light of the observations given in the same section.

F. The Chemical Data.

1. The Soil.

Table V presents the results of the chemical, physical, and colloidal analyses of two soil samples taken from each of the thirty-one plots of Tiers I and II of the three-year rotation; the other, from each of the thirty-one plots of Tier III of the three-year rotation. Also given is a comparative analysis of a typical Hagerstown soil. These data are from Claffey (18) and from Thomas (62).

2. The Plant.

a. Nitrogen

b. Phosphorus

c. Potassium

Table III B

Quality Data of the Cured Tobacco Crops for the Years 1932, 1933^a and 1934.

Plot No.	Burn		Elasticity		Body		Color		Per cent Wrappers			Per cent Fillers			Per cent Seconds ^b		
	1932	1934	1932	1934	1932	1934	1932	1934	1932	1933	1934	1932	1933	1934	1932	1933	1934
1	5		3	2	M	M	L	D	80.1	6.8	36.7	19.9	46.6	38.6	46.6	24.7	
2	5		4	2	M	M	L	D	82.4	6.1	50.0	17.6	44.6	35.6	49.2	14.4	
3	5		4	5	M	M	L	D	83.6	5.9	69.6	16.4	44.8	21.3	49.3	9.1	
4	5		4	5	M	M	D	D	86.2	7.6	72.7	13.8	46.2	24.7	46.2	2.6	
5	6		3	2	H	T	D	D	75.4	7.8	71.3	24.6	51.0	28.0	41.2	0.7	
6	7		4	2	H	H	D	D	79.0	6.9	65.1	21.0	48.3	28.8	44.8	6.1	
7	6		4	5	H	M	D	D	79.0	6.9	73.9	21.0	50.0	24.7	43.1	1.4	
8	6.5		2	2	H	H	D	D	74.3	8.7	71.2	25.7	50.0	27.2	41.3	1.6	
9	7		4	4	M	M	L	L	80.8	9.0	75.9	19.2	45.5	22.7	45.5	1.4	
10	5.5		2	3	H	M	D	L	75.0	8.3	68.9	25.0	50.0	29.7	41.7	1.4	
11	5		4	4	M	H	D	D	81.9	8.0	75.0	18.1	44.0	24.3	48.0	0.7	
12	5		4	3	H	M	D	L	81.4	8.2	75.4	18.6	37.7	23.9	54.1	0.7	
13	5		4	5	H	H	D	D	76.9	9.3	79.7	23.1	38.9	19.7	51.8	0.6	
14	4		3	2	H	H	D	L	79.5	9.2	77.1	20.5	35.2	21.7	55.6	1.2	
15	6		2	2	H	M	D	L	79.7	9.2	75.5	20.3	40.8	23.0	50.0	1.5	
16	5		3	4	H	M	D	L	83.1	6.8	74.6	16.9	35.6	24.2	57.6	1.2	
17	6		6	5	M	M	L	L	85.7	10.6	74.4	14.3	28.8	24.5	60.6	1.1	
18	7		6	5	M	M	L	L	84.5	11.1	80.0	15.5	34.7	19.5	54.2	0.5	
19	3		0	1	T	H	D	D	40.0	8.4	63.5	60.0	45.8	33.4	45.8	3.1	
20	4		3	2	H	M	D	L	80.0	9.1	71.6	20.0	47.3	27.8	43.6	0.6	
21	5		3	2	H	M	D	L	79.8	10.0	69.7	20.2	36.7	29.2	53.3	1.1	
22	6		4	4	M	H	D	D	84.4	13.2	67.4	15.2	42.7	31.8	44.1	0.8	
23	4		4	2	M	H	L	D	87.7	13.4	71.4	12.3	33.3	26.4	53.3	2.2	
24	4.5		4	4	M	H	L	D	80.9	11.7	76.1	19.1	36.7	21.2	51.6	2.7	

Table III B (continued)

Plot No.	Burn		Elasticity		Body		Color		Per cent Wrappers			Per cent Fillers			Per cent Seconds ^b		
	1932	1934	1932	1934	1932	1934	1932	1934	1932	1933	1934	1932	1933	1934	1932	1933	1934
25	6		3	4	M	M	L	L	78.7	8.3	73.4	21.3	33.3	22.7	58.4	3.9	
26	5		4	4	M	M	L	L	84.5	7.5	82.0	15.5	44.4	16.9	48.1	1.1	
27	5		3	3	H	H	D	D	80.0	13.3	63.3	20.0	36.7	23.0	50.0	13.7	
28	4		4	4	H	M	D	L	77.2	11.9	74.2	22.8	43.3	23.2	44.8	2.6	
29	4		2	3	T	T	D	L	63.6	8.7	59.1	36.4	39.1	32.1	52.2	8.8	
30	6.5		4	2	H	T	L	L	83.5	11.1	73.6	16.5	37.1	24.5	51.8	1.9	
31	6		4	4	T	M	L	L	82.3	13.3	74.9	17.7	36.7	24.6	50.0	0.5	

a. Commercially, the 1933 crop was a total failure because of wildfire damage, spread by severe storms.

b. In 1933, damaged wrappers are considered seconds.

In 1934, Plots 1, 2 and 3 were badly damaged by wildfire because of proximity to infected Rustica plots.

Key

Burn	Elasticity	Body	Color
	6--Excellent		
	5--Very Good		
10--Excellent	4--Good	H--Heavy	L--Light Brown
0--Very Poor	3--Fair	T--Thin	D--Dark Brown
	2--Poor	M--Medium	
	1--Very Poor		
	0--None		

Table V

The Physical, Chemical and Colloidal Analysis of the Soil from the Lancaster Experimental Plots, Compared with a Typical Analysis of a Hagerstown Soil.

Physical and Chemical Analysis			
Constituent	Claffey Tiers I and II	Claffey Tier III	Thomas
Classification	Silty Clay Loam	Silty Clay Loam	Silt Loam Clay Loam
Surface Soil to	8"	8"	9.50"
Fine Gravel	0.37%	0.12%	-----
Coarse Sand	0.19%	0.29%	7.41%
Medium Sand	0.88%	2.41%	5.01%
Fine Sand	1.34%	2.14%	3.40%
Very Fine Sand	8.01%	9.30%	12.66%
Silt	61.83%	52.87%	50.41%
Clay	23.01%	24.18%	15.75%
Organic Matter	4.35%	3.79%	5.30% ^a
Hygroscopic Moisture	1.62%	2.25%	-----
Hygroscopic Coefficient	4.75	6.03	-----
Wilting Point	6.50%	9.00%	-----
SiO ₂	69.65%	68.82%	72.33%
R ₂ O ₃	17.49%	21.64%	-----
Al ₂ O ₃	12.88%	15.14%	10.26%
Fe ₂ O ₃	4.61%	6.50%	3.69%
MgO	0.70%	0.75%	0.67%
CaO	2.11%	2.14%	0.68%
Na ₂ O	0.47%	0.38%	2.26%
K ₂ O	0.72%	0.66%	3.93%
P ₂ O ₅	0.17%	0.18%	0.10%
Mn ₂ O ₃	0.50%	0.46%	0.26%
N ₂	0.04%	0.05%	-----
H ₂ O at 100° C	1.62%	2.25%	-----
Organic Matter	3.01%	3.01%	5.30% ^a
pH	8.12	8.12	-----

Table V (cont.)
Colloidal Phase Analysis

Constituent	Claffey Tiers I and II	Claffey Tier III	Thomas
Colloidal Content			
A. Mechanical Analysis	8.64%	11.74%	
B. Hydrometer	39.10%	39.10%	
C. Hygroscopic Coefficient	15.09%	20.34%	
SiO ₂	45.52%	34.45%	
Al ₂ O ₃	13.69%	19.97%	
Fe ₂ O ₃	37.69%	35.95%	
R ₂ O ₃	51.62%	55.94%	
Combined H ₂ O	5.86	9.61	
Mol. SiO ₂			
Ratio Al ₂ O ₃	5.27	2.92	
Mol. SiO ₂			
Ratio Fe ₂ O ₃	2.99	2.55	
Mol. SiO ₂			
Ratio R ₂ O ₃	1.91	1.36	
Ca as CaCO ₃	3.63 M.E. ^b	3.45 M.E. ^b	
Ca	7.60 M.E.	7.65 M.E.	
Mg	2.54 M.E.	2.60 M.E.	
K	0.77 M.E.	0.77 M.E.	
Na	0.16 M.E.	0.17 M.E.	
NH ₄	0.14 M.E.	0.15 M.E.	
H	0.00 M.E.	0.00 M.E.	
Total M.E.	11.22	11.34	
% Saturation	100	100	
Lime Req.	---	---	
Ca extd. as Bicarbonate	1.95 M.E.	1.89 M.E.	

a. Ignition Loss.

b. (M.E.) Millequivalents of Replaceable Bases per 100 gm. of Soil.

Tables VIII, IX, and X give both percentages and milligrams per leaf of N, P, and K of samples taken from representative plots throughout the growing seasons of 1932 and 1933, and of samples taken on the dates indicated in Table IV. Tables XII, XIII, and XIV present the same data for the year 1934.

Figure VI presents graphically the percentage absorption of N, P, and K by the plants during the time intervals indicated in Table IV, on plots 1, 3, 5, and 8 for the years 1932, 1933, and 1934; and plot 19, for the years 1933 and 1934. Figure VII presents a similar picture except that the unit for absorption is the number of milligrams of N, P, and K absorbed per average middle leaf on plants taken from the above mentioned plots during the same intervals.

Figures VIII and IX present graphically the percentage absorption of N, P, and K by the plants, during the time intervals indicated in Table IV, on plots other than 1, 3, 5, 8, and 19 for the year 1934. Figures X and XI portray the same absorption except that the unit for absorption is the number of milligrams of N, P, and K absorbed per average middle leaf on plants taken from the above mentioned plots during the same intervals.

Table VIII

The nitrogen content of the 1932 and 1933 tobacco crops as based on samples taken from representative experimental plots at different intervals during the growing season. The results are expressed on the moisture-free weight of the average middle leaf.

Plot No.	Treatment	Year	Sampling											
			First		Second		Third		Fourth		Fifth		Sixth	
			Nitrogen per cent	mgm.	Nitrogen per cent	mgm.	Nitrogen per cent	mgm.	Nitrogen per cent	mgm.	Nitrogen per cent	mgm.	Nitrogen per cent	mgm.
1	N P 1/3 K	1932	4.21	113	4.47	217	4.20	294	3.86	304	3.49	326	2.86	260
		1933	5.34	83	4.02	212	3.92	245	4.19	319	-----	---	-----	---
3	N P K	1932	4.17	113	4.40	210	4.17	337	-----	---	4.08	408	4.19	425
		1933	5.18	120	4.31	257	3.40	258	3.01	264	-----	---	-----	---
5	P K	1932	4.10	62	4.26	124	3.93	213	3.66	284	3.73	301	4.04	387
		1933	3.89	129	3.86	125	3.28	177	2.96	207	-----	---	-----	---
8	N K	1932	4.48	68	4.29	122	4.02	197	3.51	256	4.15	310	4.20	402
		1933	4.33	83	4.36	178	4.54	246	3.74	285	-----	---	-----	---
19	Check	1932	-----	---	-----	---	-----	---	-----	---	-----	---	-----	---
		1933	4.49	105	4.83	151	4.40	242	3.61	261	-----	---	-----	---

Table IX

The phosphorus content of the 1932 and 1933 tobacco crops as based on samples taken from representative experimental plots at different intervals during the growing season. The results are expressed on the moisture-free weight of the average middle leaf.

Plot No.	Treatment	Year	Sampling											
			First		Second		Third		Fourth		Fifth		Sixth	
			Phosphorus per cent	mgm.	Phosphorus per cent	mgm.	Phosphorus per cent	mgm.	Phosphorus per cent	mgm.	Phosphorus per cent	mgm.	Phosphorus per cent	mgm.
1	N P 1/3 K	1932	.208	5.6	.218	10.6	.187	13.1	.178	14.0	.208	19.4	.188	17.1
		1933	.444	6.9	.240	12.7	.253	15.8	.285	21.7	----	----	----	----
3	N P K	1932	.217	5.9	.229	10.9	.192	15.5	----	----	.185	18.5	.170	17.2
		1933	.420	9.7	.262	15.6	.245	18.6	.224	19.6	----	----	----	----
5	P K	1932	.180	2.7	.199	5.8	.177	9.6	.151	11.7	.169	13.7	.174	16.7
		1933	.260	8.6	.280	9.0	.217	11.7	.230	16.1	----	----	----	----
8	N K	1932	.199	3.0	.203	5.8	.177	8.7	.146	10.6	.181	13.5	.162	15.5
		1933	.350	6.7	.235	9.6	.272	14.7	.240	18.3	----	----	----	----
19	Check	1932	----	----	----	----	----	----	----	----	----	----	----	----
		1933	.314	7.3	.295	9.2	.264	14.5	.228	16.5	----	----	----	----

Table X

The potassium content of the 1932 and 1933 tobacco crops as based on samples taken from representative experimental plots at different intervals during the growing season. The results are expressed on the moisture-free weight of the average middle leaf.

Plot No.	Treatment	Year	Sampling											
			First		Second		Third		Fourth		Fifth		Sixth	
			Potassium	Potassium	Potassium	Potassium	Potassium	Potassium	Potassium	Potassium	Potassium	Potassium	Potassium	Potassium
			per cent	mgm.	per cent	mgm.	per cent	mgm.	per cent	mgm.	per cent	mgm.	per cent	mgm.
1	N P 1/3 K	1932	3.27	88	3.11	151	1.88	131	1.59	125	2.85	266	1.88	171
		1933	3.02	47	2.20	116	2.80	175	2.23	170	-----	----	-----	----
3	N P K	1932	3.04	82	3.34	159	3.18	257	-----	----	2.53	253	2.07	210
		1933	3.65	85	3.23	193	3.75	285	3.43	301	-----	----	-----	----
5	P K	1932	2.99	45	3.12	91	2.77	150	2.38	185	3.15	255	2.87	275
		1933	3.26	108	3.41	110	3.36	182	3.18	222	-----	----	-----	----
8	N K	1932	2.93	45	3.09	88	2.89	141	2.19	160	2.88	215	2.23	213
		1933	3.63	70	3.89	159	4.19	227	3.35	256	-----	----	-----	----
19	Check	1932	-----	----	-----	----	-----	----	-----	----	-----	----	-----	----
		1933	3.13	73	2.75	86	3.08	169	3.00	217	-----	----	-----	----

Table X II

The nitrogen content of the 1934 tobacco crop as based on samples taken from the experimental plots at different intervals during the growing season. The results are expressed on the moisture-free weight of the average middle leaf.

Plot No.	Sampling							
	First		Second		Third		Fourth	
	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf
1	5.13	286	4.82	347	4.69	460	4.54	557
2	4.71	248	5.21	314	4.74	470	4.88	686
3	5.19	241	4.95	331	5.00	591	4.27	509
4	4.25	205	4.56	283	5.20	499	4.43	573
5	4.56	164	4.87	244	4.76	405	4.01	466
6	4.36	186	4.97	247	5.16	430	4.31	530
7	5.08	233	4.91	303	5.31	453	4.41	503
8	4.49	192	4.80	277	5.45	370	4.55	460
9	4.79	260	4.94	290	5.30	425	4.47	432
10	4.26	186	4.72	275	4.78	424	3.77	407
11	5.08	183	4.78	251	5.05	422	4.43	535
12	4.28	233	4.77	331	4.84	407	4.33	510
13	5.04	222	4.97	336	5.10	498	4.60	587
14	4.21	171	4.94	342	5.00	445	5.25	638
15	4.85	208	4.61	315	4.85	367	4.79	510
16	4.50	188	4.85	321	5.29	380	4.81	580
17	5.11	188	4.86	341	5.21	464	4.81	612
18	4.46	229	5.01	297	5.10	577	4.55	638
19	5.23	177	4.64	246	5.04	454	4.93	580
20	4.29	181	4.75	332	4.80	451	4.54	540
21	4.83	178	4.60	253	4.74	407	4.71	596
22	4.53	220	4.69	346	4.93	535	4.56	602
23	5.10	170	4.83	298	5.03	465	4.81	648
24	4.40	189	4.75	339	5.09	518	4.40	553
25	4.87	248	4.83	338	4.94	494	4.73	508
26	4.51	214	5.15	345	5.13	474	4.38	607
27	4.88	184	4.54	299	5.10	452	4.60	679
28	4.45	201	4.75	339	5.16	560	4.33	597
29	5.30	183	5.21	348	5.11	481	4.83	675
30	4.51	215	5.06	372	5.27	597	4.98	735
31	5.01	249	4.91	373	5.00	511	4.66	679
Mean	4.72	207.2	4.85	312.0	5.04	467.3	4.57	575.2

Table XIII

The phosphorus content of the 1934 tobacco crop as based on samples taken from the experimental plots at different intervals during the growing season. The results are expressed on the moisture-free weight of the average middle leaf.

Plot No.	Sampling							
	First		Second		Third		Fourth	
	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf
1	.319	17.7	.269	19.4	.250	24.5	.241	29.6
2	.269	14.2	.295	17.8	.251	24.8	.260	36.6
3	.327	15.2	.265	17.7	.268	31.7	.224	26.6
4	.220	10.6	.233	14.4	.257	24.6	.230	29.8
5	.271	9.7	.240	12.0	.226	19.2	.214	24.8
6	.216	9.3	.266	13.2	.271	22.6	.209	25.7
7	.295	13.5	.239	14.8	.257	21.9	.212	24.2
8	.218	9.3	.245	14.1	.280	19.0	.209	21.1
9	.250	14.7	.260	15.2	.259	20.8	.213	20.6
10	.231	10.1	.265	15.4	.241	21.4	.207	22.4
11	.304	10.9	.242	12.7	.243	20.3	.234	28.3
12	.220	11.9	.253	17.6	.242	20.4	.219	25.8
13	.295	13.0	.246	16.6	.243	23.7	.215	27.5
14	.227	9.2	.263	18.2	.252	22.5	.237	28.9
15	.307	13.2	.266	18.2	.276	20.9	.237	25.2
16	.258	10.8	.263	17.4	.259	18.6	.247	29.7
17	.329	12.1	.270	18.9	.272	24.2	.232	29.5
18	.244	12.5	.275	16.3	.265	30.0	.229	32.1
19	.346	11.7	.277	14.7	.261	23.5	.226	26.5
20	.240	10.1	.262	18.3	.267	25.1	.238	28.4
21	.301	11.1	.264	14.6	.279	24.0	.238	30.1
22	.265	12.9	.269	19.9	.279	30.3	.227	30.0
23	.341	11.4	.271	16.7	.283	26.2	.233	31.4
24	.255	10.9	.284	20.3	.294	30.0	.227	28.5
25	.312	15.9	.291	19.9	.279	27.9	.238	25.6
26	.255	12.1	.291	19.5	.289	26.7	.237	32.9
27	.358	13.5	.272	17.9	.290	25.6	.249	36.8
28	.267	12.1	.289	20.6	.324	35.2	.234	32.3
29	.341	11.8	.280	18.7	.286	26.9	.249	34.8
30	.286	13.6	.293	21.5	.287	32.5	.246	36.4
31	.281	13.9	.283	21.5	.276	28.3	.230	33.5
Mean	.279	12.2	.267	17.2	.268	24.9	.230	28.9

Table XIV

The potassium content of the 1934 tobacco crop as based on samples taken from the experimental plots at different intervals during the growing season. The results are expressed on the moisture-free weight of the average middle leaf.

Plot No.	Sampling							
	First		Second		Third		Fourth	
	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf	Per cent	Mgm. per Leaf
1	3.11	173	2.54	183	1.82	179	1.74	213
2	3.23	170	2.88	174	2.32	230	2.53	355
3	3.76	175	3.52	236	3.45	407	2.49	296
4	4.68	226	4.43	275	3.54	340	3.37	403
5	3.63	131	3.23	161	3.49	297	3.66	425
6	4.09	175	2.65	132	3.72	310	3.49	428
7	4.16	191	3.36	208	2.85	243	3.33	380
8	3.54	151	3.57	206	3.48	237	3.46	350
9	4.27	232	3.40	200	3.41	274	3.25	314
10	3.35	147	2.97	173	2.92	259	3.04	329
11	3.33	120	2.72	143	2.99	251	3.19	385
12	4.53	247	3.60	250	3.09	259	3.01	356
13	4.46	196	3.74	252	3.98	389	3.88	515
14	3.76	152	3.22	223	2.97	265	2.92	343
15	3.81	164	2.72	186	4.04	306	3.01	320
16	2.38	99	3.06	203	2.79	200	2.77	334
17	3.48	128	3.28	230	2.85	253	2.98	380
18	3.71	190	3.56	211	3.32	376	3.46	484
19	3.17	107	2.58	137	2.58	233	2.69	316
20	3.59	151	3.10	216	3.66	344	3.24	386
21	3.74	138	3.26	179	3.66	315	2.69	340
22	5.05	245	3.67	271	4.30	467	3.65	482
23	4.21	140	3.55	219	4.06	375	3.65	491
24	4.80	206	4.35	311	3.68	375	3.48	438
25	3.97	202	3.68	252	3.88	388	3.55	382
26	4.92	234	4.21	282	4.30	397	4.24	587
27	4.12	155	4.34	286	3.93	348	3.51	519
28	4.60	208	4.35	310	4.64	504	4.41	609
29	4.06	140	3.70	247	3.10	292	2.96	414
30	4.36	208	4.47	329	3.86	437	3.51	518
31	3.78	188	3.85	293	3.75	383	3.62	528
Mean	3.92	173.8	3.47	225.1	3.43	320.4	3.25	407.1

Fig. VI. The Percentages of N, P and K in 1932, 1933, and 1934.

The figures are based on analyses of samples taken throughout the growing season from representative plots. Topping and harvesting dates are indicated. To obtain absolute values for P, divide by 10.

Fig. VII. The Milligrams of N, P and K in the Average Middle Leaf, in 1932, 1933 and 1934.

The figures are based on analyses of samples taken throughout the growing season from representative plots. Topping and harvesting dates are indicated. To obtain absolute values for P, divide by 10.

Fig. VIII. The Percentages of N, P and K in 1934.

The figures are given for those plots not shown in Fig. VI, and are based on analyses of samples taken throughout the growing season. Topping and harvesting dates are indicated. To obtain absolute values for P, divide by 10.

Fig. IX. The Percentages of N, P and K in 1934.

The figures are given for those plots not shown in Figs. VI and VIII, and are based on analyses of samples taken throughout the growing season. Topping and harvesting dates are indicated. To obtain absolute values for P, divide by 10.

Fig. X. The Milligrams of N, P and K in the Average Middle Leaf in 1934.

The figures are given for those plots not shown in Fig. VII, and are based on analyses of samples taken throughout the growing season. Topping and harvesting dates are indicated. To obtain absolute values for P, divide by 10.

Fig. XI. The Milligrams of N, P and K in the Average Middle Leaf in 1934.

The figures are given for those plots not shown in Figs. VII and X, and are based on analyses of samples taken throughout the growing season. Topping and harvesting dates are indicated. To obtain absolute values for P, divide by 10.

n.b.- Plot 23 should be $1/2N$ P $2/3K$ M.

VII. - Discussion of Results

A. General.

Observations for three years in an experiment dealing with soil and plant relationships in actual field practices, where all variables except the one being studied are not subject to absolute control by the investigator, should be very carefully used in the interpretation of results.

Caution should be exercised because:

1. Climatic factors, notably rainfall, may be more influential in effect than is the variable being studied.
2. The variation in the natural fertility of the soil from field plot to field plot may account for observed differences.
3. Where the plant itself is being studied, the overshadowing of such things as translocation, age of plant and degree of maturity, diseases, cultural practices, and inaccuracy of sampling, may mask the effect of the imposed conditions.

In such experiments, therefore, the usual procedure is to repeat the experiment many times, over a long period of time, in an attempt to overcome the inaccuracies mentioned above. Statistical handling of the data then points to the significant factors that are causing differences.

In view of the above facts this study must be considered the first portion of a long-time experiment. Final

verification of results must await the completion of the several cycles through which this study is scheduled to go.

B. The Soil.

A high percentage of the tobacco grown in Pennsylvania is produced on the Hagerstown series of soils which, as a whole, is considered to be very productive. Tiers I, II, and III of the three-year rotation are located on several types of this Hagerstown series. It is difficult to determine the role that the unequal distribution of these several types of the Hagerstown soil over the experimental field plays in the interpretation of the data from these plots. The chemical, physical, and colloidal analyses, given in Table V, of the soil from Tiers I and II compared with that from Tier III, would indicate that the difference in inherent fertility of the different plots is not marked. However, the real criterion of variation in inherent fertility is the data from the replicate plots 5, 10, 15, 20, 25, and 30, all of which receive the same basic treatment of 1000 lbs. of 0-8-12 per acre. It must be noted that plot 30 differs from the other replicates in that alfalfa replaces clover in the rotation.

A study of the data from these replicate plots reveals that on the average the inherent fertility ranges from plot 5, the lowest in natural fertility, to plot 25, the highest in natural fertility, and decreases slightly to plot 30. If a line be drawn through the peaks of these plots, as they are shown in Figure II, this variation in fertility can be

ascertained. The only method used in this work to discount the variation in fertility was to refer the specific data under observation to the nearest replicate, considering that the natural fertility proceeds from replicate to replicate, as the slope of the line joining them. The no-treatment plots, 19 and 29, invariably give tobacco yields which are the lowest of all 31 plots on each tier, showing that the inherent fertility is of a comparatively low order.

The data in Table V show several pertinent facts about the soil which deserve mention: the relatively high pH--8.12, the low potassium and organic matter contents, and the high calcium content.

C. Rainfall.

The role of water in the mineral nutrition of plants and in plant metabolism cannot be too strongly emphasized. The overshadowing effects of rainfall, both amount and distribution, are very difficult to isolate from the imposed conditions in field experiments. The rainfall for each of the three years--1932, 1933, and 1934, during which this work was conducted, was above normal. Of these periods, the rainfall of 1932 and 1934 deviated slightly from normal while that of 1933 deviated greatly. With regard to the time-distribution of rain, the 1932 growing season was comparatively long because of dry intervals especially at the beginning of the season just after transplanting; the 1933 distribution was fairly good, but the quantities were great; the 1934 distribution, on the other hand, was conducive to

good growth.

D. Sampling for Chemical Analysis.

The expediency of the method of sampling--eleven leaves taken at the same relative position on the plant from eleven different plants on each plot--might be questioned because the effects of maturity and age of the plants, the effects of cultural practices, such as topping and suckering, the effects of diseases, the effects of translocation of mineral nutrients within the plant, and the effect of previous crops, are reflected in the samples, together with the effects of the various fertilizer treatment. This method of sampling was adopted because of lack of facilities for handling either larger numbers of leaves or whole plants.

E. The Effects of Different Field and Fertilizer Treatments.

1. On Tobacco.

These effects will first be discussed under the several sub-groupings into which the various treatments fall. Then they will be considered as a whole. The criteria which will be used for ascertaining the differences caused by treatment are the yield, the quality, the growth characteristics, the dry weight, and the absorption of nitrogen, phosphorus, and potassium considered from the viewpoints of both percentage and milligrams absorbed per leaf.

a. When the amount of nitrogen in the fertilizer is varied, phosphorus and potassium being constant: plot 5, P K; 6, $1/2N$ P K; 3, N P K; 7, $3/2N$ P K.

From the viewpoints of yield, dry weight,

quality, and growth characteristics, the treatment of plot 3 undoubtedly is the best of this group. According to the percentage absorption, no apparent correlation exists between the amount of nitrogen applied, and the amount of nitrogen, phosphorus, and potassium absorbed by the plant; according to milligrams per leaf, plot 3 is conspicuous as having absorbed the most nitrogen, phosphorus, and potassium. This somewhat indicates the fact that the amount of nitrogen, phosphorus, and potassium absorbed is not necessarily proportional to the amount applied, but that the ratio or balance of the nutrients in the fertilizer is a more important consideration. It also indicates that for purposes of comparison of treatments, the total amount of nutrients absorbed is probably a better criterion than is a percentage comparison(12).

b. When the amount of phosphorus in the fertilizer is varied, nitrogen and potassium being constant: plot 8, N K; 9, N 1/2P K; 3, N P K.

With this kind of treatment plot 3 is again the best when judged from yield, dry weight, quality and growth characteristics data. On the percentage basis, no apparent correlation exists between the amount of nitrogen absorbed and phosphorus applied, but there is some correlation when judged by the milligram standard. The absorption of phosphorus, from both a percentage and a milligram per leaf standard, increases in the plant as it is applied to the soil, indicating that a lack of phosphorus is a very

important limiting factor in the growth of tobacco on this particular soil. As far as potassium absorption is concerned, percentage viewpoint, it will be noticed that potassium for the most part exists in greater percentages on the lower phosphorus plots; by the milligram standard, the data seems to indicate that the potassium absorption increases as the phosphorus applications increase. This can be accounted for by the increase in dry weight that accompanies increasing applications of phosphorus. Undoubtedly, phosphorus is one of the most important of the limiting factors in growth.

c. When the amount of potassium in the fertilizer is varied, nitrogen and phosphorus being constant: plot 1, N P 1/3K; 2, N P 2/3K; 3, N P K; 4, N P 4/3K.

Under these conditions plot 3 gives the highest yield; plot 4, the highest quality. There is no correlation for the percentage of nitrogen absorbed and potassium applied. However, there seems to be an inverse relationship between potassium applied and nitrogen absorbed when considering total milligrams. There seems to be no correlation between phosphorus absorbed and potassium applied, though plot 3 did absorb a fair amount of phosphorus. The potassium is absorbed in a direct relationship to the amount applied.

A balance of nutrients in the fertilizer, such as occurs in the treatment of plot 3 (6-8-12), is more favorable to high yield, quality, and absorption of nutrients

than is any other consideration in the fertilization of tobacco on this soil.

d. When the amount of the same fertilizer is varied and when applied in the row and broadcast:
plot 11, 1/2N 1/2P 1/2K-B; 3, N P K-B; 12, 3/2N 3/2P 3/2K-B;
plot 16, " " " -R; 17, " " " -R; 18, " " " -R.

The yields occurred in the following order: 18, 17, 3, 16, 12, 11. In other words, the row treatments of 1500 and 1000 pounds are superior. But 1000 pounds broadcast is also a good treatment as far as yield is concerned. In fact 18 was the highest yielding plot of all plots except in 1934 when it was surpassed by plot 31. This same order with few exceptions holds for quality. In most instances, the amounts of nitrogen, phosphorus, and potassium absorbed vary directly with the amounts applied. Row treatment gave slightly higher amounts of nitrogen absorbed than did broadcast treatment. Phosphorus and potassium, on the other hand, were absorbed about equally well by either treatment.

e. When lime is added, nitrogen, phosphorus, and potassium being constant: plot 3, N P K; 13, N P K L.

Yield and quality data indicate that liming is not beneficial. This is to be expected when the high pH of the soil is considered. The lime plot showed the higher percentage of nitrogen absorbed; higher potassium absorption by both percentage and milligram standards; and less phosphorus taken in by the plant from both angles. Perhaps this

condition can be accounted for by the stimulation by the lime of the micro-organic population of the soil, liberating nitrogen and potassium; but the excess of calcium makes soil phosphorus more unavailable.

f. When $2/3$ of the nitrogen is applied as nitrate of soda, nitrogen, phosphorus, and potassium being constant: plot 3, N P K-- $2/3$ of the nitrogen from cotton seed meal; 14, N P K, $2/3$ of the nitrogen from nitrate of soda.

Yield and quality data favor cotton seed meal. This may probably be accounted for if the organic matter of the cotton seed meal, and the effect of nitrate of soda on calcium availability, are considered. The percentage of nitrogen from the nitrate of soda surpassed that absorbed from the cotton seed meal as the plant matured. The percentage of phosphorus absorbed was increased by cotton seed meal, as were also the percentage and milligrams of potassium.

g. When alfalfa replaces clover in the rotation: 29 versus 19; 30 versus 25; 31 versus 3.

The data for 1932 cannot be considered in this respect because no crop preceded the tobacco. The yield and quality data of 1933 and 1934 would seem to indicate that alfalfa is superior to clover in the rotation. On the whole, more nitrogen, phosphorus, and potassium were absorbed on the alfalfa plots. Perhaps this can be explained by the fact that alfalfa would put more organic matter into this

relatively poor soil, and also because the growth of alfalfa is favored by the high pH of the soil.

h. When manure is applied.

It should be noted that manure adds not only organic matter but also some nitrogen, phosphorus, and potassium to the soil. If the manure be considered a .5-.25-.5 fertilizer, a ten ton application of manure would add 100 lbs. of nitrogen, 50 lbs. of phosphoric acid, and 100 lbs. of potash; a 20 ton application would double these figures. In interpreting the data on both the manure plots, and the fertilizer plus manure plots, these figures should be kept in mind. For example, plot 28 was treated with 20 tons of manure, thus adding 200 lbs. of nitrogen, 100 lbs. of phosphoric acid and 200 lbs. of potash; plot 22 was treated with 10 tons of manure and 1000 lbs. of 3-8-12; thus 130 lbs. of nitrogen, 130 lbs. of phosphoric acid and 220 lbs. of potash were added. It can be seen that both of these plots have more nitrogen, phosphorus, and potassium added in their treatments than do any of the plots treated with fertilizer alone.

1'. ... and varying amounts of manure used: plot 27, M; 28, 2M.

In all cases plot 28 gave a higher yield than plot 27 while the quality was about the same. More nitrogen and more potassium were absorbed by 28 than by 27, while there seems to be but a slight correlation between the amount of phosphorus absorbed and the amount of manure

applied. The phosphorus absorption was relatively high in both cases.

2'. When a constant amount of manure is used,

a'. ...and varying amounts of commercial fertilizer applied: plot 24, $1/4N$ $1/2P$ $1/2K$ M; 22, $1/2N$ P K M.

A comparison of these two plots indicates that on the average the more commercial fertilizer added with manure, the greater the yield, and the better the quality. As far as the percentage of nitrogen absorbed is concerned, there seems to be no correlation with the treatment. But from the milligram viewpoint more nitrogen seems to have been taken in with the increase in fertilizer. However, so much nitrogen is present in the manure that the additional 15 lbs. on plot 22 do not seem to make much difference. There seems to be no apparent correlation with phosphorus absorption, but more potassium is absorbed where the greater amount of fertilizer is applied.

b'. ...and varying amounts of nitrogen in the commercial fertilizer applied: plot 21, $1/2P$ K M; 24, $1/4N$ $1/2P$ $1/2K$ M.

The difference between these two plots with respect to yield and quality is very slight. This is apparent when one considers that the difference in the treatment of these two plots is only 15 lbs. of nitrogen. The correlation between nitrogen and phosphorus absorption

and treatment is not apparent, but there does seem to be more potassium absorbed ^{where} the greater amount of nitrogen was used.

c'. ...and varying amounts of potassium in the commercial fertilizer applied: plot 23, $1/2N$ P $2/3K$ M; 22, $1/2N$ P K M; 26, $1/2N$ P $4/3K$ M.

Plot 22 gave the highest yield indicating that the balance of nutrients in the fertilizer is important. The quality differences among these three plots is not marked. On plot 22 the milligrams of nitrogen absorbed were slightly higher than on the other two plots, as were also the milligrams of absorbed phosphorus. A correlation seems to exist again between the amount of potassium applied and the amount absorbed.

In all of these cases where manure was supplemented with commercial fertilizer the amounts of the fertilizing elements were all comparatively high so that a slight variation in the amounts of any one of them did not seem to have any marked effect on the criteria which were studied. Perhaps phosphorus, here again, is the only limiting growth factor although the nitrogen is relatively high for the production of good cigar-leaf tobacco. There seems to be no limit to the amount of potassium which can be applied when one considers that potassium probably governs quality more than does any one mineral element.

1. When nothing is applied, when manure alone is applied, when fertilizer alone is applied, and when

fertilizer and manure are applied.

A study of the data from the untreated plots shows the advantage of using some kind of fertilizer treatment. This can be seen when plots 29 and 19 are compared with all the other plots with respect to yield, growth, quality, and susceptibility to disease. When manure alone is used, a 20 ton application gives fair results, while a 10 ton application gives only mediocre results. Commercial fertilizer alone, on the other hand, gave the best results of this study especially when the ratio of fertilizing element was 6-8-12 and applied in amounts of 1000 or 1500 lbs. both in the row and broadcast. However, when commercial fertilizer is used with manure the increases are marked and approach very closely the high yield and quality that result from the use of commercial fertilizer alone. From a practical standpoint, it is advisable to use manure, if it is available and inexpensive, supplemented with a commercial fertilizer high in phosphorus and potassium.

2. The effects of the various fertilizer treatments on wheat.

The only factor studied with regard to wheat was the yield. All plots received a treatment of 300 lbs. of 16% superphosphate. The carry-over effects of the tobacco fertilization was most marked on the manure plots. Surprisingly, the carry-over effects of the row treatment, which one would expect to be detrimental, were not marked. However, the data are rather meager for any definite

conclusions to be drawn with respect to wheat.

3. The effects of the various fertilizer treatments on clover.

There seems to be but little correlation between the clover yields and the treatments which were given to the tobacco two years prior in the rotation. The best yields of clover were obtained from the first 7 or 8 plots which had received commercial fertilizer alone.

4. Recapitulation.

It seems advisable to give here a resume of the lengthy discussion above.

The highest yields were obtained where 1500 and 1000 lbs. of 6-8-12 were used, followed by a combination of fertilizer and manure, 20 tons of manure alone, and 1000 lbs. of 6-8-12 broadcast. Where alfalfa was in the rotation and 1000 lbs. of 6-8-12 applied the yield was also high. Cotton seed meal appears to be a superior carrier of nitrogen than is nitrate of soda. Liming is not gainful on the soil studied.

The highest percentage of wrappers, which is indicative of quality, ^{was} were obtained from those plots treated with fertilizer plus manure, where large amounts of potassium were applied, and where the fertilizer was applied in the row. The other criteria of quality--burn, elasticity, body, and color of the cured leaves--also indicate that row treatments, large amounts of potassium, and fertilizer plus manure are best. With regard to the field observations--

growth, color of the green plant, and degree of maturity--row application, commercial fertilizer plus manure, and alfafa included in the rotation were the most favorable forms of treatment. Susceptibility to disease seems to be related to the amount of potassium applied. This is in accord with observations made by McMurtry (57) and Rohde (73, 74).

The data seem to indicate that the greatest amount of nitrogen is absorbed by the plant under the following conditions: where one element such as potassium or phosphorus is omitted from the fertilizer; where alfafa occurs in the rotation, and where manure is applied. Phosphorus is perhaps the most limiting factor in the growth of the tobacco plant on the soils which were used in this study. Phosphorus was absorbed to the greatest extent by the plant when nitrogen and potassium were omitted from the fertilizer treatment; when manure was used; when alfafa occurred in the rotation; and when a combination of commercial fertilizer and manure were applied. Potassium was absorbed by the plant almost in direct ratio to the amount applied in the fertilizer. In the order mentioned, twenty tons of manure, commercial fertilizer plus manure, 1000 lbs. 6-8-16, and 1000 lbs. 6-8-12 with alfafa in the rotation gave the greatest amounts of potassium in the leaf.

It seems that the absolute amounts of nitrogen, phosphorus, and potassium absorbed per leaf was a better criterion to follow than was the percentage composition of

the leaf when the interrelationships of absorption were being studied.

The wheat and clover data show no significant differences.

F. The Absorption of Nitrogen, Phosphorus, and Potassium
Considered from Liebig's "Law of the Minimum".

In this study, the dry weight, and the percentages and milligrams per leaf of nitrogen, phosphorus, and potassium of samples taken during the growing season from plots 1, 3, 5, and 8, for three years, and from plot 19 for two years were used. Plot 1 is considered to have been a N P plot, although 40 lbs. of potash had been added; plot 3 is considered the complete nutrient plot---N P K; plot 5, a P K plot; plot, 8, a N K plot; and plot 19, as a check plot.

A study of the data seems to reveal the following points: that on the average, the dry weight production for three years follows Liebig's Law; in 1932 as far as absorption of the nutrients is concerned, Liebig's law holds in the majority of cases; in 1933 and 1934 the variations are so great that the law is neither substantiated nor is it invalidated. In field experiments, Liebig's "Law of the Minimum" cannot be followed ideally because many uncontrollable variables enter into the picture of the absorption processes of the several nutrients.

VIII. - Summary

Under the limitations of the experimental conditions-- observations taken over the relatively short period of three years, variations of soil fertility, and the effects of diseases--the following facts appear to have been established about the fertilization of tobacco:

1. Phosphorus is the greatest limiting factor in the growth of the plant on the soil studied.
2. 6-8-12 seems to be the best ratio of fertilizer elements.
3. Fifteen hundred and 1000 lbs. of 6-8-12, applied in the row, give the best yields with good quality. A combination of 10 tons of manure and 1000 lbs. of 3-8-12 constitute a good treatment. Likewise, 1000 lbs. of 6-8-12, broadcast, gave good results, especially when alfalfa occurs in the rotation.
4. The potassium absorption by the plant varies directly with the amounts applied in the fertilizer.
5. The amount of nitrogen absorbed is the highest where either phosphorus or potassium is omitted from the fertilizer, where alfalfa is used in the rotation, and where manure alone is used.

6. The phosphorus absorption is greatest where either nitrogen or potassium is omitted from the fertilizer, where alfalfa is used, where manure is used alone, and where fertilizer plus manure is used.
7. Liebig's "Law of the Minimum" does not hold in all cases studied.

IX. - Acknowledgments

The author takes this opportunity to express his sincere appreciation to Dr. D. E. Haley for his many helpful suggestions during the course of this work; to Mr. Otto Olson for his kind cooperation in the conduct of these experiments; and to several other friends who aided in the preparation of this manuscript.

X. - Bibliography

1. Ames, J. W., and Bolts, G. E.
1915. Tobacco, Influence of Fertilizers on Composition and Quality.
Ohio Agr. Expt. Sta. Bull. 285.
2. Anderson, P. J.
1929. Soil Reaction Studies on the Connecticut Tobacco Crop.
Jour. Amer. Soc. Agro. 21: 156.
3. Anderson, P. J., and Swanback, T. R.
1928. Report of Tobacco Substation.
Conn. Agr. Expt. Sta. Bull. 299.
4. Anderson, P. J., Swanback, T. R., Street, O. E., et al.
1930. Tobacco Substation at Windsor Report for 1929.
Conn. Agr. Expt. Sta. Bull. 311.
5. Anderson, P. J., Swanback, T. R., Street, O. E., et al.
1931. Tobacco Substation at Windsor Report for 1930.
Conn. Agr. Expt. Sta. Bull. 326.
6. Anderson, P. J., Swanback, T. R., and Street, O. E.
1932. Potash Requirements of the Tobacco Crop.
Conn. Agr. Expt. Sta. Bull. 334.
7. Anderson, P. J., Swanback, T. R., and Street, O. E.
1932. Tobacco Substation at Windsor Report for 1931.
Conn. Agr. Expt. Sta. Bull. 335.
8. Anderson, P. J., Swanback, T. R., and Street, O. E.
1934. Tobacco Substation at Windsor Report for 1933.
Conn. Agr. Expt. Sta. Bull. 359.
9. Annual Review of Biochemistry.
1932-1935. Vol. 1-4.
Stanford Univ. Press, Stanford Univ. Cal.
10. Association of Official Agricultural Chemists.
1931. Official and Tentative Methods of Analysis.
A. O. A. C. Washington, D. C.
11. Barth, M.
1891. Untersuchungen von in Klassegezogenen Tabaken und Einige Beziehungen Zwischen der Qualitat des Tabaks und Seiner Zusammensetzung.
Landw. Ver. Sta. 39: 81-104.
12. Bartholomew, R. P., Watts, V. M., Janesen, George.
1933. The Effect of Variations in the Nutrient Media upon the Nitrogen, Phosphorus, and Potassium Content on Plants with Special Reference to the Tomato.
Univ. of Ark. Agr. Expt. Sta. Bull. 288.
13. Beaumont, A. B., Larsinos, G. J., Pickenbrock, P., and Nelson, P. R.
1931. The Assimilation of Nitrogen by Tobacco.
Jour. Agr. Res. 43: No. 6.
14. Beaumont, A. B., and Snell, M. E.
1931. Field Experiments with Tobacco.
Mass. Agr. Expt. Sta. Bull. 271.

Bibliography (cont.)

15. **Bahrens, J.**
1894. Weitere Beitrage zur Kenntnis der Tabakpflanze.
V. Der Anatomische Bau und die Bestandteile des
Tabakblatts in ihrer Beziehung zur Brennbarkeit.
Landw. Ver. Sta. 43: 271-301.
16. **Brown, D. E., and Lunn, W. M.**
1925. Cropping Tests with Tobacco.
Univ. Of Maryland. Agr. Expt. Sta. Bull. 275.
17. **Carolus, R. L.**
1933. Tomato Fertilization. II. The Effect of
Different Fertilizer Ratios on the Chemical Com-
position of Tomatoes.
Va. Truck Expt. Sta. Bull. 81.
18. **Claffey, L. W.**
1933. A Chemical Study of the Soils of the New
Tobacco Experiment Station at Lancaster, Pennsylvania.
Thesis: College Library, State College, Pennsylvania.
19. **Davidson, R. J.**
1895. Analyses of Parts of Tobacco Plant at Different
Stages of Growth.
Va. Agr. Expt. Sta. Bull. 50.
20. **Fesca, M.**
1888. Ueber Kultur, Behandlung und Zusammensetzung
Japan ischer Tabake.
Landw. Jahrb. 17: 329-372.
21. **Frear, W.**
1888. Pennsylvania Tobacco Soils.
Pa. Agr. Expt. Sta. Bull. 7.
22. **Frear, W.**
1903. Experiments in Growing Sumatra Tobacco under
Shelter-Tents.
Pa. Agr. Expt. Sta. Bull. 72.
23. **Frear, W.**
1909. Shelter-Tent Experiments with Sumatra Tobacco.
Pa. Agr. Expt. Sta. Bull. 89.
24. **Frear, W., et al.**
1915-16. Tobacco Experiments.
Pa. Agr. Expt. Sta. Annual Report.
25. **Frear, W., and Hibshman, E. K.**
1922. The Production of Cigar-Leaf Tobacco in
Pennsylvania.
U. S. Dept. of Agr. Farmers' Bull. 416.
26. **Garner, W. W.**
1907. The Relation of the Composition of the Leaf
to the Burning Qualities of Tobacco.
U. S. Dept. Agr. Bur. Plant Ind. Bull. 105.
27. **Garner, W. W., Bacon, C. W., and Bowling, Jr.**
1934. Cigarette and Cigar Tobaccos. Relationship
of Production Conditions to Chemical and Physical
Characteristics.
I. E. C. 26: 970.

Bibliography (cont.)

28. Garner, W. W., Bacon, C. W., Bowling, J. D., and Brown, D. E.
1934. The Nitrogen Nutrition of Tobacco.
U. S. Dept. Agr. Bull. 414.
29. Garner, W. W., Luan, W. M., Brown, D. E.
1925. Effects of Crops on the Yields of Succeeding Crops in the Rotation with Special Reference to Tobacco.
Gen. Agr. Res. 30: 1095.
30. Garner, W. W., McMurtry, J. E. Jr., Bowling, J. D. Jr., and Moss, E. G.
1930. Magnesium and Calcium Requirements of the Tobacco Crop.
Gen. Agr. Res. 40: 145.
31. Garner, W. W., McMurtry, J. E. Jr., and Moss, E. G.
1923. Sand-drown, a Chlorosis of Tobacco Due to Magnesium Deficiency, and the Relation of Sulphates and Chlorides to the Disease.
Gen. Agr. Res. 23: 27.
32. Gilbert, B. D., Cobb, W. B., Moffitt, E. L., and Cox, J. F.
1916. Soil Survey of Lancaster County, Pennsylvania.
U. S. Dept. Agr. Bur. of Soils. July, 1916.
33. Graham, Vere, and Carr, R. H.
1924. Chemical Factors Determining the Quality of Tobacco.
Jour. Amer. Chem. Soc. 46: No. 3.
34. Haley, D. E.
1929. The Chemical Approach to the Study of Problems of Tobacco Fertilization.
Jour. Amer. Soc. Agro. 21: No. 2.
35. Haley, D. E., Longenecker, J. B., and Olson, O.
1931. Composition and Quality of Pennsylvania Cigar-Leaf Tobacco as Related to Fertilizer Treatment.
Plant Phys. 6: 177.
36. Haley, D. E., Hasset, E. S., and Olson, O.
1928. A Study of Certain Constituents of the Leaf and Their Relation to the Burning Qualities of Tobacco.
Plant Phys. 3: 155.
37. Haley, D. E., and Olson, Otto.
1929. On the Relation of Potassium to Iron in the Combustion of Cigar-Leaf Tobacco.
Science 70: 17.
38. Jenkins, E. H.
1892. Experiments in Growing Tobacco with Different Fertilizers.
Conn. Agr. Expt. Sta. Ann. Rept. 1-24.

Bibliography (cont.)

39. Jenkins, E. H.
1914. Studies on the Tobacco Crop of Connecticut.
Conn. Agr. Expt. Sta. Bull. 150.
40. Kostutany.
1882. Chem Physiologische Untersuchung der Tobak--
sorten Ungarns.
Budapest B. 41.
41. Lagatu, H., et Maure, L.
1928. Antagonisme du Calcaire a l'Egard de l'absorp--
tion de la Potasse par la Vigne. Communication faite
le 22 Juin, 1928, a Bordeaux, au Congres International
du Vin et du Pire Maritime.
42. Liebig, Justus.
1843. Chemistry in Its Application to Agriculture and
Physiology.
Ed. III. (Out of Print)
43. Liebig, Justus.
1863. Les Loix Naturelles de l'Agriculture.
Paris. Ed. 2. Chapt. II.
44. McMurtrey, J. E. Jr.
1929. The Effect of Boron Deficiency on the Tobacco
Plants in Aerated and Un-aerated Solutions.
Jour. Agr. Res. 38: 371.
45. McMurtrey, J. E. Jr.
1933. Distinctive Effects of the Deficiency of Certain
Essential Elements on the Growth of Tobacco Plants in
Solution Cultures.
U. S. Dept. of Agr. Bull. 340.
46. McMurtrey, J. E. Jr., Lunn, W. M., and Brown, D. E.
1934. Fertilizer Tests with Tobacco.
Univ. of Md. Agr. Expt. Sta. Bull. 358.
47. Mayer, A.
1879. Uber die Verbrennlichkeit und den Chlorgehalt--
gedungten Tabaks.
Central-Blatt fur Agrikulturchemie 8: 814
48. Mayer, A.
1890. Tabakdungungsversuche mit Beurteilung der
Qualitat des Erzeugnisses.
Landw. Ver. Sta. 38: 126.
49. Mayer, A.
1890. Uber die Verbrennlichkeit des Tabaks.
Landw. Ver. Sta. 38: 127.
50. Miller, E. C.
1931. Plant Physiology.
McGraw-Hill Book Co., Inc. N. Y. and London.
51. Morgan, M. F.
1929. Tobacco as an Indicator Plant in Soil Studies.
Jour. Amer. Soc. Agro. 21: No. 2.
52. Morgan, M. F., Anderson, P. J., Dorsey, Henry.
1929. Soil Reactions and Liming as Factors in
Tobacco Production in Connecticut.
Conn. Agr. Expt. Sta. Bull. 306.

Bibliography (cont.)

53. Moss, E. G., McMurtrey, J. E. Jr., Lunn, W. M., and Carr, J. M.
1927. Fertilizer Tests with Flue-cured Tobacco.
U. S. Dept. Agr. No. 12.
54. Kessler, J.
1867. Der Tabaks Seine Bestand-Teile Undeene Behandlung.
(Mannheim)
55. Olson, Otto.
1922. Preparing Cigar-Leaf Tobacco for Marketing.
Pa. Agr. Expt. Sta. Leaflet 16.
56. Olson, O.
1923. Results of Tobacco Experiments in Pennsylv--
vania, 1912-1922.
Pa. Agr. Expt. Sta. Bull. 179.
57. Olson, Otto.
1923. Tobacco Seed Beds.
Pa. Agr. Expt. Sta. Leaflet 17.
58. Olson, O., and Haley, D. E.
1920. Results of Tobacco Experiments in Pennsylv--
vania 1922-1927.
Pa. Agr. Expt. Sta. Bull. 240.
59. Orton, C. R., and Olson, O.
1925. I. Tobacco Resistant to Black Root-Rot in
Pennsylvania. II. A Method of Eliminating Ex--
perimental Error in Comparative Field Tests.
Pa. Agr. Expt. Sta. Bull. 193.
60. Russell, Sir Edward John.
1932. Soil Conditions and Plant Growth.
Longmans-Green & Co. London and N. Y.
61. Schlosing, Th.
1860. Uber die Verbrennlichkeit des Tabaks.
Landw. Ver. Sta. 3: 98.
62. Thomas, Walter.
1923. Ultimate Analysis of the Mineral Constituents
of a Hagerstown Silty Clay Loam Soil and Occurrence
in Plants of Some of the Elements Found.
Soil Science, Vol. XV. No. 1.
63. Thomas, Walter.
1929. Balanced Fertilizers and Liebig's Law of the
Minimum.
Science. Vol. 70. No. 1816.
64. Thomas, Walter.
1930. The Conception of Balance with Respect to the
Absorption of Nitrogen, Phosphorus, and Potassium
by Plants and the Influence of the Level of Nutri--
tion.
Science. Vol. 72. No. 1869.
65. Thomas, Walter.
1930. The Feeding Power of Plants.
Plant Phys. 5: No. 4.

Bibliography(cont.)

66. Thomas, Walter.
1932. Composition of Current and Previous Season's Branch Growth in Relation to Vegetative and Reproductive Responses in *Pyrus Malus*.
Plant Phys. 7: 391
67. Thomas, Walter.
1932. The Reciprocal Effects of Nitrogen, Phosphorus, and Potassium as Related to the Absorption of These Elements by Plants.
Soil Science. Vol. 33. No. 1.
68. Thomas, Walter.
1933. Absorption, Utilization, and Recovery of Nitrogen, Phosphorus, and Potassium by Apple Trees Grown in Cylinders and Subjected to Differential Treatment with Nutrient Salts.
Jour. Agr. Res. Vol. 47. No. 8
69. Totttingham, W. E.
1934. Plant Biochemistry.
Burgess Publ. Co., Minneapolis, Minn.
70. Van Bemmelen, J. M.
1899. Über die Zusammensetzung der Asche der Tabakblätter in Beziehung zu ihrer Guten oder Schlechten Qualität Insbesondere zu ihrer Brennbarkeit.
Landw. Ver. Sta. 37: 409.
71. Whitney, W.
1894. Tobacco Soils of Pennsylvania.
U. S. Dept. Agr. Report. 1894.
72. Wolff, Emil.
1880. Aschen-Analysen von landund forstwirtschaftlichen Producten, Fabrik-Abfällen und wildwachsenden Pflanzen.
Wiegandt, Hempel, und Paray. Berlin.
73. Rohde, Gustav.
1935. Disturbances of the absorption of Light Rays by Plants Due to Potash Deficiency.
Unpublished Data. Berlin-Zehlendorf.
74. Rohde, Gustav.
1935. Symptoms of Potash Deficiency on Potatoes.
Unpublished Data. Berlin.
75. Dutcher, R. A., and Haley, D. E.
1932. Agricultural Biochemistry.
John Wiley and Sons, Inc. New York.

**End of
Title**