COLUMBIA LIBRARIES OFFSITE
HEALTH SCIENCES STANDARD

HX00018279

RECAP

Columbia University
in the City of New York

College of Physicians and Surgeons

Library

Digitized by the Internet Archive
in 2010 with funding from
Columbia University Libraries

<http://www.archive.org/details/atlasofhumananat12told>

AN ATLAS
OF
HUMAN ANATOMY

FOR STUDENTS AND PHYSICIANS

BY
CARL TOLDT, M.D.

ASSISTED BY
PROFESSOR ALOIS DALLA ROSA, M.D.

Adapted to English and American and International Terminology

BY
M. EDEN PAUL, M.D. BRUX., M.R.C.S., L.R.C.P.

FIRST SECTION

A. THE REGIONS OF THE HUMAN BODY

B. OSTEOLOGY

(FIGURES 1 TO 377 AND INDEX)

REVISED EDITION

NEW YORK
REBMAN COMPANY
141, 143 AND 145 WEST 36TH STREET

Copyright, 1919, by
REEMAN COMPANY

Q M 25
T 572

v. 1-2

cop. 3

TRANSLATOR'S PREFACE

THE science of human anatomy is purely descriptive in its methods, the field it covers is not very extensive, and its boundaries are sharply limited; it is, therefore, one of the few sciences in which something closely verging on finality and completeness has been attained. Even, however, if no new anatomical data are likely to be forthcoming, there is yet scope for originality in the method of presentation of those data of which the science now consists; and originality of this kind Professor Toldt's "Atlas of Human Anatomy" exhibits in a high degree. In the many admirable manuals of human anatomy now extant in English, the illustrations, even when numerous, as they are often, and when good, as they are occasionally, form a mere supplement—usually a very imperfect supplement—to the text. Atlases of anatomy, and useful atlases, also exist in English, but all are quite fragmentary. Some, like the well-known and valuable, but somewhat antiquated, "Illustrations of Dissections," consist of a series of pictures of selected regions carefully prepared on the cadaver: these are models for the imitation of the student in his own dissecting work, but are not of much value for private study. Others, like Bellamy's English edition of Braune's atlas of frozen sections of the human body, present a small number of anatomical facts from a striking and unfamiliar point of view. But among English works, an accurate pictorial representation of *all* the data of human anatomy, carefully drawn to scale from actual specimens, and arranged suitably for systematic study, has hitherto been lacking.

Whilst a true knowledge of anatomy, a knowledge that will through life supply the needs of the physician and the surgeon in their practical work, can be obtained only in the dissecting-room, the student's labours with scalpel and forceps must be preceded and supplemented by systematic private study. Now, for this purpose, the textual descriptive treatise is not alone sufficient; or, if sufficient, it is so at an excessive expenditure of time and labour. Both in his work preparatory to dissection and in his revision of his anatomical knowledge subsequent to dissection, the energy of the student will be enormously economized if he has at hand a graphic representation of every structure named and described in his systematic treatise. An increased use of the visual or graphic method, both in the acquirement and in the revivification of knowledge, is a feature of the age in all educational departments; but this English translation of Professor Toldt's work is, as far as the English-speaking races are concerned, the first adequate application of the method to the study of human anatomy.

In speaking of the finality and completeness of anatomical science, one exception must be made, and this exception relates to anatomical terminology, which, though nearly completed, has not yet attained finality. Had there been a universal anatomical nomenclature—a nomenclature, that is, adopted by, or even fully intelligible to, anatomists of all nationalities—an English edition of this work would have been superfluous. Anatomy, however, like all other sciences, has suffered from the dispersion of tongues that ensued on the Renaissance, when the good and the evil of mediævalism became inextricably confounded, and were cast away together, and the inestimable gift of a language common to the learned of all lands was lost for ever. The German-speaking peoples have a fairly complete and fairly pure Latin anatomical nomenclature, needing, however, to be eked out here and there by the vernacular; whilst in England, as in France, a strange and bastard dialect, half Latin and half vernacular, has come into use. Uncouth jargon as it is, being current and familiar, it is not likely in England and America ever to be replaced by the more consistent terminology in use in the anatomical schools of Germany and Austria; I have, however, in this English edition of the "Atlas of Anatomy" retained the terminology of the original side by side with the English translation, distinguishing between the two by a difference of type.

In some cases, in the nomenclature used by the author, terms are met with which have no counterpart in English anatomical terminology: either because the author regards as normal a structure which English anatomists regard as a variety; or, and far more commonly, because the structure in question, though normal, is unimportant, and English anatomists have therefore neglected to name it. Sometimes, in such cases, I have given a literal English translation of the Latin name used by the author; sometimes, however, a periphrasis has been required to explain what the structure is, or to account for the absence of an English name, and this periphrasis, when lengthy, has been printed as a foot-note. In all such cases, an *asterisk* is prefixed both to the Latin name and to its English equivalent, to indicate to the reader that there is something unusual in the terms employed.

I must further point out that in a few instances the author's nomenclature actually conflicts with that commonly used in England, so that the literal translation of the author's name for a certain structure is applied in England to a structure totally different. For instance, what the author calls *canalis pterygopalatinus* is in England called the *posterior palatine* or *palatomaxillary canal*, while the *pterygopalatine canal* of English anatomists is called by the author *canalis pharyngeus*. But for this warning, beginners might imagine such divergencies to be due to carelessness on the part of the translator or to errors of the press.

A further difficulty has arisen from the fact that English anatomical nomenclature is itself not yet finally settled, nor even wholly consistent. Not merely is the same structure often known by several names; but, which is worse, the same

TRANSLATOR'S PREFACE

name is sometimes applied to two different structures. Reform is therefore needed, but it is not the part of a translator to undertake it, and I have perforce been content to follow the authorities. My leading authority has been the tenth edition of Quain's "Elements of Anatomy," but I have also had Macalister's "Text-book of Human Anatomy" in constant requisition. From these works I have, when more than one name is used to denote any structure, taken all those in common use, the order in which the alternative names have been printed showing most often the relative frequency of employment; in a few cases, however, where a name less commonly used has appeared to me distinctly preferable for any reason to an alternative name more commonly used, I have given the less usual but preferable name the precedence. To this small extent only have I been influenced by my own views in the matter of anatomical terminology; and, with the exception of those names which for the reason already furnished are preceded by an asterisk, all the terms in the English nomenclature are in use by one or more of the leading English authorities.

As regards the terminology employed in the United States of America, the contributions of the scientific investigators of that country to anatomy have, owing to the early perfection of this branch of study, been far less extensive than in the case of the other sciences ancillary to medicine; and the science of anatomy was for the most part taken bodily over, text-books, terminology, and all complete. A few differences, however, exist, and I have therefore collated my manuscript with that useful little work, Young's "Synopsis of Human Anatomy," and any divergent terms in use in America only have been inserted in my translation, and distinguished by the addition of the letters "U.S."

A considerable number of the references to the figures will be found to be in the English nomenclature only. These are either cases in which the English and the International descriptive terms were identical, and the printing of both was therefore superfluous; or else cases in which in the original the reference was wholly in German.

Measurements given in the original in centimetres have in all cases been reduced to inches. In illustrations of foetal parts the age of the foetus is given in months from the date of fertilization of the ovum. On the Continent, however, the period of utero-gestation is usually reckoned as ten "months" of four weeks each; not, as with us, as nine calendar months. To avoid mistake, I have in all such cases after the word "month" or "months" added in parentheses the words "months of four weeks each."

I cannot dismiss mention of the works of reference I have employed without alluding to the German-English "Dictionary of Medical Terms," by Treves and Lang—a book invaluable to all those engaged in the translation of German medical works.

Since this Atlas is intended for the use of beginners, as well as for that of advanced students of human anatomy and of practitioners of medicine, I may fitly conclude this preface with a few words on the general principles of anatomical nomenclature. For descriptive purposes the body is regarded as being in the upright posture, with the arms extended by the sides, and the hands fully supinated, so that the palms look forward. With this attitude kept in mind, the meaning of the terms *superior* and *inferior*, *anterior* and *posterior*, *external* and *internal*, is obvious. Sometimes, however, descriptive terms of another kind are used, to remove the confusion liable to arise from the adoption by man of an attitude different from that of all the other vertebrata, and to homologize the nomenclature of human with that of comparative anatomy. Thus, *cephalic* and *caudal* in comparative anatomy correspond respectively with *superior* and *inferior* in human anatomy; *ventral* and *dorsal*, with *anterior* and *posterior*. Dividing the body into right and left halves by a vertical *median plane*, which cuts the surface of the body at the *median line*, *medial* or *mesial* and *lateral* correspond respectively with *internal* and *external* in denoting position respectively nearer to, or more remote from, the median plane. Other terms in frequent use are *superficial* and *deep*, *central* and *peripheral*, *proximal* and *distal*; these are self-explanatory.

In some cases descriptive terms applied to portions of certain structures denote the relation of these portions to other structures, as when we speak of the *vertebral* and the *sternal* extremities of the ribs, or the *acromial* and the *sternal* extremities of the clavicle. Terms of similar import are *radial* and *ulnar* applied to structures of the forearm; *tibial* and *fibular* (or *peroneal*) of the leg; *palmar* and *dorsal* of the hand; *plantar* and *dorsal* of the foot; *flexor* and *extensor* of any of the extremities. It is to be noted that *internal* and *external* are sometimes used in a sense different from that previously explained, being employed to denote the interior and exterior positions respectively, either in relation to the general axis of the body or to the axis of one of its cavities. In this sense, for instance, we may speak of the *internal* and the *external* tables of the cranial vault, or of the *internal* and the *external* oblique muscles of the abdomen; but it is, as a rule, better to use the words *inner* and *outer* to denote this relation, and to reserve *internal* and *external* for position in respect to the median plane.

Finally we have to explain the terms used to denote certain directions, more especially the direction of certain sections: these are *horizontal* and *vertical*, requiring no definition; *sagittal*, denoting a dorso-ventral direction either in or parallel to the median plane; and *frontal* or *coronal*, which are synonymous terms, denoting direction in a transverse vertical plane.

The definition of many of the terms used in descriptive anatomy, such as *condyle* and *tuberosity*, *process* and *tubercle*, *sinus* and *cavity*, *ligament*, *tendon*, and *aponeurosis*, would be superfluous, since the student will best gain an accurate notion of their meaning by an examination of the structures to which they are respectively applied.

M. EDEN PAUL.

ALDERNEY, August, 1903.

REGIONES
CORPORIS HUMANI

THE REGIONS
OF THE HUMAN BODY

FIG. 1.—ANTERIOR SURFACE OF THE BODY.

Regions of the Human Body.

FIG. 1a.—ANTERIOR SURFACE OF THE BODY.

Regiones Corporis Humani.

FIG. 2.—POSTERIOR SURFACE OF THE BODY.

Regions of the Human Body.

FIG. 2a.—POSTERIOR SURFACE OF THE BODY.

Regiones Corporis Humani.

¹ Known also as the *infraclavicular triangle* or *infraclavicular fossa*, and sometimes called *Morvenheimer's space*.

FIG. 3.—HEAD AND NECK.

Regions of the Head and Neck.

FIG. 3a.—HEAD AND NECK.

Regiones Capitis et Colli.

FIG. 4.—MALE PERINEAL REGION.

FIG. 5.—FEMALE PERINEAL REGION.

Perineal Region.

FIG. 4a.—MALE PERINEAL REGION.

FIG. 5a.—FEMALE PERINEAL REGION.

Regio Perinealis.

OSTEOLOGIA

OSTEOLOGY

THE STRUCTURE OF THE BONES

FIG. 6.—PORTION OF A CROSS-SECTION THROUGH THE COMPACT TISSUE OF A LONG BONE.

FIG. 7.—PORTION OF A LONGITUDINAL SECTION THROUGH THE COMPACT TISSUE OF A LONG BONE.

FIG. 8.—DIAGRAM OF THE STRUCTURE OF BONE.

FIG. 9.—PART OF THE MIDDLE SEGMENT OF THE FEMUR FROM WHICH THE PERIOSTEUM HAS BEEN PARTIALLY REMOVED.

The medullary canal (cavum medullare) has been opened, and the bone-marrow (medulla ossis) is seen.

Periosteum and Bone-Marrow (Medulla Ossium).

FIG. 10.—PROXIMAL PORTION IN
FRONTAL SECTION.

FIG. 11.—DISTAL PORTION IN
SAGITTAL SECTION.

Humerus—Arm-bone: Substantia compacta et substantia spongiosa ossium—Compact and cancellous tissue of the bones.

Ossa longa—Long bones.

FIG. 12.—PROXIMAL PORTION IN
SAGITTAL SECTION.

Ulna.

FIG. 13.—DISTAL PORTION IN
FRONTAL SECTION.

FIG. 14.—PROXIMAL PORTION.

Radius.

FIG. 15.—DISTAL PORTION.

Substantia compacta et substantia spongiosa ossium—Compact and cancellous tissue of the bones.
Ossa longa—Long bones.

FIG. 16.—PROXIMAL PORTION IN
FRONTAL SECTION.

FIG. 17.—DISTAL PORTION IN FRONTAL SECTION.

Femur—Thigh-bone: Substantia compacta et substantia spongiosa ossium—Compact and cancellous tissue of the bones.

Ossa longa—Long bones.

FIG. 18.—PROXIMAL PORTION IN
FRONTAL SECTION.

FIG. 19.—DISTAL PORTION IN
FRONTAL SECTION.

Tibia—Shin-bone: Substantia compacta et substantia spongiosa ossium—Compact and cancellous tissue of the bones.

Ossa longa—Long bones.

FIG. 20.—THIRD LUMBAR VERTEBRA IN HORIZONTAL SECTION.

FIG. 21.—SIXTH DORSAL VERTEBRA IN SAGITTAL SECTION.

FIG. 22.—BODY OF THE SECOND LUMBAR VERTEBRA IN FRONTAL SECTION.

Vertebræ : Substantia compacta et substantia spongiosa ossium—Compact and cancellous tissue of the bones.

Ossa brevia—Short bones.

FIG. 23.—OS CUNEIFORME III.. THE EXTERNAL CUNEIFORM BONE.

FIG. 24.—TALUS, THE ASTRAGALUS, IN SAGITTAL SECTION.

FIG. 25.—CALCANEUM, THE OS CALCIS, IN SAGITTAL SECTION.

Ossa tarsi—Tarsal bones: Substantia compacta et substantia spongiosa ossium—Compact and cancellous tissue of the bones.

Ossa brevia—Short bones.

FIG. 26.—NUTRIENT FORAMEN AND NUTRIENT CANAL OF THE RIGHT TIBIA.
SEEN FROM BEHIND.

By sawing away a portion of the shaft the whole length of the nutrient canal has been opened up.

Nutrient Foramen and Nutrient Canal of a Long Bone.

FIG. 27.—OUTER SURFACE OF LEFT ILIUM.

By the removal of the outer compact layer and the cancellous tissue of a portion of the bone, the nutrient canals have been displayed. The bristles projecting towards the right above and below show that the nutrient canals into which they have been inserted open on the inner surface of the bone—that turned away from the observer.

FIG. 28.—OS PARIETALE, PARIETAL BONE, PREPARED TO SHOW THE DIPLOË AND THE COMPACT INNER TABLE. SEEN FROM WITHOUT.

Substantia compacta, substantia spongiosa, et canales nutricii—Compact tissue, cancellous tissue, and nutrient canals.

Ossa plana—Flat bones.

FIGS. 29 AND 30.—TWO STAGES IN THE INTRACARTILAGINOUS OSSIFICATION OF LONG BONES, AS SHOWN BY LONGITUDINAL SECTIONS OF THE PHALANGES OF A HUMAN FŒTUS.

FIGS. 31 AND 32.—TWO STAGES IN THE INTRACARTILAGINOUS OSSIFICATION OF THE EPIPHYSIS OF A LONG BONE, AS SHOWN BY LONGITUDINAL SECTIONS OF THE DISTAL PORTIONS OF METACARPAL BONES.

FIG. 33.—INTRACARTILAGINOUS OSSIFICATION OF A SHORT BONE.

A section of the cuboid bone of a new-born child.

FIG. 34.—INTRAMEMBRANOUS OSSIFICATION OF THE BONES OF THE CRANIAL VAULT.

Upper half of the parietal bone of a human fœtus in the eleventh week of intra-uterine life.

FIG. 35.—ANTERIOR ASPECT.

FIG. 36.—VIEWED FROM THE LEFT SIDE,
THE ARM HAVING BEEN REMOVED.

Skeleton humanum—The human skeleton.

SKELETON TRUNCI
THE AXIAL SKELETON

FIG. 37.—THE VERTEBRAL COLUMN AS A WHOLE. CLASSIFICATION AND NOMENCLATURE OF THE VERTEBRÆ.

Columna vertebralis—The spinal column.

FIG. 38.—SEEN FROM THE LEFT SIDE.

FIG. 39.—SEEN FROM ABOVE.

FIG. 40.—SEEN FROM BEFORE.

FIG. 41.—SEEN FROM BEHIND.

Vertebræ: Vertebra thoracalis VI.—Sixth dorsal vertebra.

FIG. 42.—SEEN FROM THE LEFT SIDE.

FIG. 43.—SEEN FROM BEFORE.

FIG. 44.—SEEN FROM ABOVE.

FIG. 45.—SEEN FROM BELOW.

Vertebræ: Vertebra cervicalis V.—Fifth cervical vertebra.

Vertebrae: Vertebra lumbalis II.—Second lumbar vertebra.

FIG. 49.—VERTEBRA THORACALIS I.—FIRST DORSAL VERTEBRA. SEEN FROM BEFORE.

FIG. 50.—VERTEBRA THORACALIS XI.—ELEVENTH DORSAL VERTEBRA. SEEN FROM THE LEFT SIDE.

FIG. 51.—VERTEBRA THORACALIS XII.—TWELFTH DORSAL VERTEBRA. SEEN FROM THE LEFT SIDE.

FIG. 52.—VERTEBRA LUMBALIS V.—FIFTH LUMBAR VERTEBRA. SEEN FROM THE LEFT SIDE.

Vertebræ: Transitional forms of the dorsal and lumbar vertebræ.

FIG. 53.—THE ATLAS, OR FIRST CERVICAL VERTEBRA. SEEN FROM ABOVE.

FIG. 54.—EPISTROPHEUS, THE AXIS, OR SECOND CERVICAL VERTEBRA. SEEN FROM BEFORE.

FIG. 55.—VERTEBRA CERVICALIS VII.—SEVENTH CERVICAL VERTEBRA. SEEN FROM ABOVE.

Vertebræ: The atypical cervical vertebræ.

FIG. 56.—MALE SACRUM (FACIES PELVINA—PELVIC SURFACE). SEEN FROM BEFORE.

FIG. 57.—FEMALE SACRUM (FACIES PELVINA—PELVIC SURFACE). SEEN FROM BEFORE.

Oss sacrum—The sacrum.

FIG. 58.—MALE SACRUM (FACIES DORSALIS—DORSAL SURFACE). SEEN FROM BEHIND.

FIG. 59.—TRANSVERSE SECTION THROUGH THE SACRUM AT THE LEVEL OF THE FIRST SET OF SACRAL FORAMINA.

FIG. 60.—SACRUM AND COCCYX. SEEN FROM THE LEFT SIDE.

FIG. 61.—SACRUM AND COCCYX IN SAGITTAL SECTION THROUGH THE MEDIAN LINE.

FIG. 62.—COCCYX SEEN FROM BEFORE.

FIG. 63.—COCCYX SEEN FROM BEHIND.

Os sacrum et os coccygis—Sacrum and coccyx.

FIG. 64.—UNILATERAL ASSIMILATION AND COMMENCING ANKYLOSIS OF THE FIFTH LUMBAR VERTEBRA WITH THE SACRUM. SEEN FROM BEFORE.

From a boy seventeen years of age. Epiphyses on the bodies of the sacral vertebra, and along the lateral margins of the sacrum.

FIG. 65.—ANKYLOSIS OF THE SACRUM WITH THE COCCYX. SEEN FROM BEHIND.

From a man forty years of age.

Anomalous Ankyloses of the Sacrum.

FIG. 66.—DIAGRAM SHOWING THE RELATIONS OF THE PROTO-VERTEBRÆ TO THE VERTEBRÆ. (FROM VON EBNER.)

FIG. 69.—THE THREE CENTRES OF OSSIFICATION OF A VERTEBRÆ.

Horizontal section through the second and a portion of the first lumbar vertebrae of a human fetus in the beginning of the fifth month (months of four weeks each). (Length of fetus, $5\frac{1}{2}$ inches.)

FIG. 67.—PRIMITIVE CARTILAGES OF THE VERTEBRAL BODIES.

From a sagittal section through the vertebral column of a human fetus in the tenth week. (Length of fetus, $2\frac{1}{4}$ inches.)

FIG. 68.—CENTRES OF OSSIFICATION OF THE BODIES OF THE VERTEBRÆ.

From a sagittal section through the vertebral column of a human fetus in the fourth month (months of four weeks each). (Length of fetus, $4\frac{1}{2}$ inches.)

Development of the Vertebrae.

FIG. 70.—FIRST LUMBAR VERTEBRA OF A NEW-BORN CHILD.

FIG. 71.—FIRST LUMBAR VERTEBRA IN THE EIGHTEENTH YEAR.

FIG. 72.—THE SEVENTH CERVICAL VERTEBRA OF A NEW-BORN CHILD.

FIG. 74.—THE AXIS AT THE END OF THE FIRST YEAR.

FIG. 73.—THE ATLAS AT THE END OF THE FIRST YEAR.

FIG. 76.—THE AXIS IN THE FOURTH YEAR.

FIG. 75.—THE ATLAS IN THE FOURTH YEAR.

FIG. 77.—SEEN FROM BEFORE.

FIG. 78.—SEEN FROM BEHIND.

Development of the Vertebrae: The centres of ossification in the sacrum and coccyx of a child at the age of two months.

(1—7, Costæ veræ, sternal or true ribs ; 8—12, Costæ spuræ, asternal or false ribs ; 11 and 12, Costæ fluctuantes, floating ribs.)

FIG. 79.—THE THORAX SEEN FROM BEFORE.

The Thorax.

FIG. 80.—THE THORAX SEEN FROM BEHIND.

The Thorax.

FIG. 81.—THE TWELVE PAIRS OF RIBS.

Ossa costalia—The ribs.

FIG. 82.—THE TWELVE RIBS OF THE RIGHT SIDE IN THEIR NATURAL POSITION,
SEEN FROM THE RIGHT.

Costæ—The ribs.

FIG. 83.—FIRST (RIGHT) RIB. SEEN FROM ABOVE.

FIG. 85.—FOURTH (RIGHT) RIB. SEEN FROM BEHIND.

FIG. 84.—SECOND (RIGHT) RIB. SEEN FROM ABOVE.

FIG. 86.—SEVENTH (RIGHT) RIB. SEEN FROM WITHIN.

FIG. 87.—TWELFTH (RIGHT) RIB. SEEN FROM WITHIN.

Costæ—The ribs.

¹ In certain pathological conditions the angle between the manubrium and the body of the sternum becomes less obtuse, and therefore more prominent. It is then known as *angulus Ludovici*, or Ludwig's angle.—TR.

FIG. 88.—THE STERNUM SEEN FROM BEFORE.

FIG. 89.—THE STERNUM SEEN FROM THE LEFT SIDE.

FIG. 90.—THE UPPER PORTION OF THE STERNUM WITH THE SUPRASTERNAL BONES (A RARE VARIETY). SEEN FROM BEFORE.

Sternum—The breast-bone.

Epiphysis of the head
Epiphysis capituli

FIG. 91.—POSTERIOR PORTION OF THE SIXTH RIB, IN THE FIFTEENTH YEAR.

Epiphysis of the head
Epiphysis capituli

Epiphysis of the articular facet of the tubercle
Tubercle of the rib
Tuberculum costae

FIG. 92.—POSTERIOR PORTION OF THE SIXTH RIB, IN THE EIGHTEENTH YEAR.

FIG. 93.—DIVIDED PRIMITIVE CARTILAGE OF THE STERNUM.

From a human fetus of two months (months of four weeks each).

FIG. 94.—PRIMITIVE CARTILAGE OF THE STERNUM.

From a human fetus of four months (months of four weeks each).

FIG. 95.—PRIMITIVE CARTILAGE OF THE STERNUM WITH THE FIRST CENTRE OF OSSIFICATION IN THE MANUBRIUM.

From a human fetus in the second half of the sixth month (months of four weeks each).

FIG. 96.—CENTRES OF OSSIFICATION IN THE STERNUM OF A NEW-BORN CHILD.

FIG. 97.—STERNUM OF A BOY AT THE AGE OF ELEVEN YEARS.

Development of the Ribs and the Sternum.

FIG. 98.—SKELETON OF A THORACIC SEGMENT.

FIG. 99.—SKELETON OF A CERVICAL SEGMENT.

FIG. 100.—SKELETON OF A LUMBAR SEGMENT.

FIG. 101.—SKELETON OF A SACRAL SEGMENT.

The Homologous Skeletal Parts of the Segments of the Body.

CRANIUM ET OSSA CRANII

THE SKULL

AND THE BONES OF THE SKULL

FIG. 102.—THE SKULL SEEN FROM BEFORE: NORMA FRONTALIS.

Cranium—The skull.

FIG. 103.—THE SKULL SEEN FROM THE LEFT SIDE; NORMA LATERALIS.

¹ *Canalis basipharyngeus,* basipharyngeal canal: This term is not often used by English anatomists, nor even is the canal itself mentioned by Quain. Macalister, however, in his description of the vomer, writes (p. 233): "In the region of its sphenoidal articulation there are three canals transmitting small vessels in the young skull, which usually become obliterated with advancing age; these are—one median vomerine canal between the vomer and the root of the rostrum, and a lateral on each side between the extremity of the ala vomeris and the vaginal process. These run parallel to, and may communicate with, the pterygopharyngeal [pterygopalatine] canal in the vaginal process." The two lateral canals here mentioned are those called by Toldt *canales basipharyngei.*—Tr.

FIG. 104.—EXTERNAL ASPECT OF THE BASE OF THE SKULL: BASIS CRANII EXTERNA.

FIG. 105.—INTERNAL ASPECT OF THE BASE OF THE SKULL—BASIS CRANII INTERNA: FOSSÆ CRANII ANTERIOR, MEDIA, ET POSTERIOR—THE ANTERIOR, MIDDLE, AND POSTERIOR CRANIAL FOSSÆ. SEEN FROM ABOVE.

FIG. 106.—CALVARIA—THE SKULLCAP, OR ROOF OF THE SKULL. INNER ASPECT.

FIG. 107.—VASCULAR CANALS OF THE DIPLOË OF THE ROOF OF THE SKULL, SHOWN BY REMOVAL OF THE OUTER TABLE OF COMPACT BONE FROM THE FRONTAL BONE AND THE PARIETAL BONE: CANALES DIPLOICI (BRESCHETI). SEEN FROM THE LEFT SIDE.

FIG. 108.—THE SEPARATE BONES OF WHICH THE CRANIUM CEREBRALE OR CRANIUM PROPER CONSISTS.

Ossa cranii cerebrials—Bones of the cranium proper.

FIG. 109.—THE SEPARATE BONES OF WHICH THE CRANIUM VISCERALE (FACIES OSSEA), OR FACIAL PORTION OF THE SKULL, CONSISTS.

Ossa faciei—Bones of the face.

FIG. 110.—ANTERIOR (INTERNAL) ASPECT OF THE OCCIPITAL BONE.

Os occipitale—The occipital bone.

¹ See foot-note to p. 57.

FIG. 111.—POSTERIOR (EXTERNAL) ASPECT OF THE OCCIPITAL BONE.

Os occipitale—The occipital bone.

FIG. 112.—THE OCCIPITAL BONE SEEN FROM BELOW (EXTERNAL BASAL SURFACE).

FIG. 113.—THE OCCIPITAL BONE SEEN FROM ABOVE (INTERNAL BASAL SURFACE).

Os occipitale—The occipital bone.

FIG. 114.—THE PORTIONS OF THE OCCIPITAL BONE FROM A HUMAN FÆTUS AT THE END OF THE SIXTH MONTH (MONTHS OF FOUR WEEKS EACH). SEEN FROM WITHIN.
Body-length, 12 inches.

FIG. 115.—THE OCCIPITAL BONE OF A CHILD AGED FIFTEEN MONTHS. SEEN FROM WITHIN.

¹ The human occipital bone consists of four elements, which are still separate at birth, being united by intervening cartilage; these are, the *basilar* portion (basilar process), the two *condylar* portions, and the *tabular* portion. In comparative anatomy these are known respectively as *basi-occipital*, *exoccipitalis*, and *supra-occipital*. The basi-occipital and the exoccipitals ossify each from a single nucleus; the supra-occipital ossifies from four nuclei, an upper pair and a lower. These soon unite, but leave for some time fissures running in along the superior curved line. Not uncommonly this fissure persists on one or both sides through life, and in rare cases there is a persistent suture running right across and dividing the tabular portion of the occipital bone into two parts (*Sutura mendosa; see Figs. 111 and 114). Of these two parts, the lower, which belongs to the base of the skull and ossifies in cartilage, is the proper *supra-occipital* element, homologous with the *supra-occipital* bone of other vertebrata; whilst the upper, which belongs to the vertex of the skull and ossifies in membrane, represents the *interparietal* bone of many animals. This part alone is entitled to the name *squama occipitalis*, a term, however, little used by English anatomists. The occasional persistence of the suture between the interparietal and supra-occipital elements of the occipital bone is of surgical importance, since, in cases of injury to the back of the head, it is, if present, liable to be mistaken for fracture.—Tr.

¹ Spine for the attachment of the lower head of the external rectus muscle of the eyeball.

² See note to p. 48.

³ See note to p. 59.

⁴ As mentioned in the Preface, the canal called by English anatomists *palatomaxillary* or *posterior palatine canal* is by Toldt called *pterygopalatine canal*. The inner grooved portion of the sphenomaxillary surface, which he here calls the *pterygopalatine groove*, leads down into that canal, but does not form a part of it, since it lies between the palate bone and the superior maxillary bone.—Tr.

FIG. 116.—THE SPHENOID BONE SEEN FROM BEFORE.

FIG. 117.—THE SPHENOID BONE SEEN FROM BEHIND.

Os sphenoidale—The sphenoid bone.

FIG. 118.—THE SPHENOID BONE SEEN FROM ABOVE (CEREBRAL ASPECT).

¹ The Continental nomenclature differs here from that of English anatomists. What is called by the author *facies temporalis alæ magnæ*, the temporal surface of the great wing, is in England known as the *temporozygomatic surface*, being divided by the *infratemporal crest* into an upper *temporal surface* and a lower *sphenomaxillary surface*. As regards the term *crista orbitalis*, the crest, this is not used by Quain at all, while Macalister applies it to the lower margin of the sphenoidal fissure, the free border separating the orbital from the cerebral surface of the great wing. Toldt, on the other hand, as an examination of Fig. 110 shows, means by *crista orbitalis* the posterior margin of the sphenomaxillary fissure, free border separating the orbital from the zygomatic surface of the great wing.—Tr.

² See note to p. 43.

FIG. 119.—THE SPHENOID BONE SEEN FROM BELOW (EXTERNAL ASPECT).

Os sphenoidale—The sphenoid bone.

¹ See note to p. 59.² English anatomists use the terms *pituitary fossa* and *sella turcica* as synonyms; Todd, more accurately, distinguishes between them, meaning by *pituitary fossa* (Fossa hypophyseos) the deep pit on the upper surface of the body of the sphenoid bone which lodges the pituitary body or hypophysis cerebri, and by *sella turcica* the saddle-shaped surface which forms the floor of that fossa.—Tr.FIG. 120.—THE SPHENOID BONE SEEN FROM THE LEFT SIDE (TEMPOROZYGOMATIC SURFACE).¹

FIG. 121.—THE SPHENOID SINUSES IN MEDIAN SAGITTAL SECTION, THE GREATER PART OF THE SPHENOIDAL SEPTUM HAVING BEEN REMOVED. SEEN FROM THE LEFT SIDE.

FIG. 122.—THE SPHENOID SINUSES, EXPOSED FROM ABOVE BY THE REMOVAL OF THE INNER LAMELLA OF COMPACT BONE.

The right sinus is opened from above; the left is unopened.

Os sphenoidale—The sphenoid bone.

FIG. 123.—THE SPHENOID BONE OF A BOY BORN AT FULL TERM, SEEN FROM ABOVE.
Body-length, 19 inches.

FIG. 124.—THE SPHENOID BONE OF A BOY BORN AT FULL TERM, SEEN FROM BELOW.
Body-length, 19 inches.

FIG. 125.—CONCHÆ SPHENOIDALES—THE SPHENOIDAL TURBINATE BONES. SEEN FROM ABOVE.

FIG. 126.—THE RELATION OF THE SPHENOIDAL TURBINATE BONES TO THE INFERIOR SURFACE OF THE SPHENOID BONE IN THE SIXTH YEAR OF LIFE.

¹ What is called the mandibular fossa by Toldt is the posterior non-articular portion of the glenoid fossa (separated from the articular portion by the Glaserian fissure). Its floor is formed by the tympanic plate, and it lodges a portion of the parotid gland.—Tr.

FIG. 127.—THE LEFT TEMPORAL BONE SEEN FROM THE OUTER SIDE (TEMPORAL SURFACE).

FIG. 128.—THE LEFT TEMPORAL BONE SEEN FROM BELOW (EXTERNAL BASAL SURFACE).

Os temporale—The temporal bone.

FIG. 129.—THE LEFT TEMPORAL BONE SEEN FROM WITHIN (CEREBRAL SURFACE).

FIG. 130.—THE LEFT TEMPORAL BONE SEEN FROM ABOVE (INTERNAL BASAL SURFACE).

FIG. 131.—THE LEFT TEMPORAL BONE SEEN FROM BEFORE.

FIG. 132.—THE EXTERNAL WALL OF THE TYMPANUM AND THE MASTOID CELLS DISPLAYED BY A SECTION THROUGH THE LEFT TEMPORAL BONE IN A PLANE PARALLEL WITH THE SQUAMOUS PORTION OF THAT BONE.

The petrosquamous sulcus (along which a sound has been passed) is in this specimen partly bridged over by bone; anteriorly it communicates with the outer surface of the bone by means of a spurious jugular foramen—*foramen jugulare spurium* (Variety).

Os temporale—The temporal bone.

FIG. 133.—VERTICAL SECTION THROUGH THE PETROUS PORTION OF THE LEFT TEMPORAL BONE AND THROUGH THE ANTERIOR PART OF THE SQUAMOUS PORTION. (NOMENCLATURE OF THE SURFACE AND BORDERS OF THE PETROUS PORTION.)

FIG. 134.—AQUEDUCTUS FALLOPII, OR CANAL FOR THE FACIAL NERVE, SHOWN FROM BENEATH BY THE REMOVAL OF A WEDGE-SHAPED PIECE FROM THE PETROUS PORTION OF THE LEFT TEMPORAL BONE.

The canaliculus tympanicus, for the tympanic branch of the glossopharyngeal nerve (Jacobson's nerve), is also opened up throughout its whole length. A sound has been passed through the canaliculus for the auricular branch of the pneumogastric nerve (nerve of Arnold). In the Continental nomenclature this canaliculus is known as the *canaliculus mastoideus*.

Os temporale—Temporal bone.

FIG. 135.—VERTICAL SECTION THROUGH THE LEFT TEMPORAL BONE IN A PLANE PARALLEL WITH THE SUPERIOR BORDER OF THE PETROUS PORTION, AND PASSING THROUGH THE MIDDLE OF THE EXTERNAL AUDITORY MEATUS, TO DEMONSTRATE THE TYMPANIC CAVITY, CAVUM TYMPANI, AND THE ADJOINING PARTS.

FIG. 136. VERTICAL SECTION THROUGH THE LEFT TEMPORAL BONE IN A PLANE PARALLEL WITH THE SUPERIOR BORDER OF THE PETROUS PORTION, THE SECTION PASSING ALONG THE POSTERIOR WALL OF THE EXTERNAL AUDITORY MEATUS, TO DEMONSTRATE THE TYMPANIC CAVITY, CAVUM TYMPANI, AND THE ADJOINING PARTS.

Os temporale—Temporal bone.

FIG. 137.—VERTICAL SECTION THROUGH THE LEFT TEMPORAL BONE, CROSSING OBLIQUELY THE SUPERIOR BORDER OF THE PETROUS PORTION, AND PASSING THROUGH THE ANTERIOR PORTION OF THE MASTOID PROCESS, TO DEMONSTRATE THE TYMPANIC CAVITY, CAVUM TYMPANI, AND THE ADJOINING PARTS (ESPECIALLY THE MASTOID ANTRUM AND THE MASTOID CELLS).

FIG. 138.—VERTICAL SECTION THROUGH THE LEFT TEMPORAL BONE, CROSSING OBLIQUELY THE SUPERIOR BORDER OF THE PETROUS PORTION, AND PASSING THROUGH THE POSTERIOR PORTION OF THE MASTOID PROCESS, TO DEMONSTRATE THE TYMPANIC CAVITY, CAVUM TYMPANI, AND THE ADJOINING PARTS.

A bristle has been passed through the canaliculus tympanicus (the canal for Jacobson's nerve—the tympanic branch of the glossopharyngeal nerve) into the tympanum, and, after traversing this cavity, leaves it by the canaliculus that opens into the groove for the small superficial petrosal nerve.

Os temporale—Temporal bone.

FIG. 139.—THE BONY LABYRINTH, LABYRINTHUS OSSEUS, SHOWN IN THE LEFT PETROUS PORTION. SEEN OBLIQUELY FROM IN FRONT AND BELOW. THE OSSEOUS SEMICIRCULAR CANALS AND ALSO THE CANAL OF THE COCHLEA HAVE BEEN PARTLY OPENED. THE RELATIONS BETWEEN THE AQUEDUCT OF FALLOPIUS AND THE OSSEOUS LABYRINTH ARE CLEARLY SHOWN.

¹ The helicoid depression of the tractus spiralis foraminulentus corresponds with the base of the cochlea, and at the centre of the helix is the foramen centrale cochleæ, the orifice of the central canal of the modiolus. On the significance of this term *Area of the Cochlea*, see also note ³³⁴, p. 956y, in the Appendix to Part VI.; and on the significance of the term *Area of the Facial Nerve*, see note ³³⁸ on the same page.

FIG. 140.—THE INTERNAL AUDITORY MEATUS, MEATUS ACUSTICUS INTERNUS, EXPOSED FROM ABOVE BY THE REMOVAL OF A RIGHT-ANGLED WEDGE FROM THE PETROUS PORTION OF THE LEFT TEMPORAL BONE, DISPLAYING THE FUNDUS OF THE INTERNAL AUDITORY MEATUS, OR RENIFORM FOSSA, DIVIDED BY THE TRANSVERSE CREST, OR CRISTA FALCIFORMIS, INTO SUPERIOR AND INFERIOR FOSSÆ. SEEN FROM BEHIND AND ABOVE.

Os temporale—Temporal bone.

¹ Regarding the signification of the terms *crus simplex* and *crus commune*, see Appendix to Part VI, p. 956, note 229.

FIG. 141.—PORTIONS OF THE OSSEOUS LABYRINTH AND THE TYMPANUM, SHOWN IN THE LEFT TEMPORAL BONE BY A VERTICAL SECTION THROUGH THE PETROUS PORTION IN THE PLANE OF THE SUPERIOR SEMICIRCULAR CANAL. SEEN OBLIQUELY FROM IN FRONT AND WITHIN.

The fenestra ovalis is divided vertically.

FIG. 142.—PORTIONS OF THE OSSEOUS LABYRINTH AND THE TYMPANUM, SHOWN IN THE LEFT TEMPORAL BONE BY A HORIZONTAL SECTION THROUGH THE PETROUS PORTION ALONG THE INTERNAL AND THE EXTERNAL AUDITORY MEATUS. SEEN FROM ABOVE.

Os temporale—Temporal bone

FIG. 143.—SEEN FROM WITHOUT.

FIG. 144.—SEEN FROM WITHIN.

THE THREE PARTS OF THE LEFT TEMPORAL BONE FROM AN EIGHT-MONTHS FÆTUS (MONTHS OF FOUR WEEKS EACH).

Body-length, 1½ inches.

FIG. 145.—THE SQUAMOUS PORTION OF THE TEMPORAL BONE AND THE TYMPANIC RING UNITED. SEEN FROM WITHIN.
From a fetus at term (body-length, 19 inches).

FIG. 146.—THE THREE PORTIONS OF THE LEFT TEMPORAL BONE UNITED. SEEN FROM WITHOUT AND BELOW.
From a new-born male infant (body-length, 21 inches).

Development of the Temporal Bones.

FIG. 147.—THE LEFT TEMPORAL BONE OF A BOY AT THE AGE OF EIGHT MONTHS: FORMATION OF THE TYMPANIC PLATE AND OF THE EXTERNAL AUDITORY MEATUS. SEEN OBLIQUELY FROM WITHOUT AND BELOW.

FIG. 148.—THE LEFT TEMPORAL BONE OF A GIRL AT THE AGE OF THREE YEARS: FORMATION OF THE TYMPANIC PLATE AND OF THE EXTERNAL AUDITORY MEATUS. SEEN OBLIQUELY FROM WITHOUT AND BELOW.

Development of the Temporal Bones.

FIG. 149.—THE LEFT PARIETAL BONE SEEN FROM WITHOUT. EXTERNAL SURFACE:
FACIES PARIETALIS.

Os parietale—Parietal bone.

FIG. 150.—THE RIGHT PARIETAL BONE SEEN FROM WITHOUT. INTERNAL SURFACE: FACIES CEREBRALIS.

Cs parietale—Parietal bone.

FIG. 151.—THE FRONTAL BONE SEEN FROM BEFORE. EXTERNAL SURFACE : FACIES FRONTALIS.

Os frontale—Frontal bone.

¹ Ridges (*juga cerebralia*) and sulci (*impressiones digitatæ*) corresponding respectively to the sulci and convolutions of the superjacent portion of the frontal lobes of the cerebrum.—TR.

FIG. 152.—THE FRONTAL BONE SEEN FROM BEHIND. INTERNAL SURFACE: *FACIES CEREBRALIS*.

Os frontale—Frontal bone.

FIG. 153.—THE FRONTAL BONE SEEN FROM BELOW.

FIG. 154.—THE FRONTAL SINUSES, SINUS FRONTALES, SHOWN BY THE REMOVAL OF THE EXTERNAL TABLE AND THE DIPLOË, AND PARTLY OPENED UP. SEEN FROM BEFORE AND BELOW.

Os frontale—Frontal bone.

FIG. 155.—THE TWO HALVES OF THE FRONTAL BONE FROM A HUMAN FŒTUS IN THE EIGHTH MONTH (MONTHS OF FOUR WEEKS EACH). SEEN FROM BEFORE.

Body-length of fœtus 15 inches.

FIG. 156.—THE FRONTAL BONE OF A GIRL AGED SEVEN YEARS IN WHICH THE FRONTAL SINUSES HAVE BEEN EXPOSED. SEEN OBLIQUELY FROM BEFORE AND FROM THE RIGHT SIDE.

Development of the Frontal Bones.

FIG. 157.—THE ETHMOID BONE SEEN FROM THE LEFT SIDE

FIG. 158.—THE ETHMOID BONE SEEN FROM ABOVE.

FIG. 159.—THE LEFT HALF OF THE ETHMOID BONE SEEN FROM WITHIN. SUPERIOR AND MIDDLE TURBINATE BONES.

FIG. 160.—THE LEFT HALF OF THE ETHMOID BONE SEEN FROM WITHIN.

The anterior portion of the middle turbinate bone of the nose has been removed. The free border of the removed portion is indicated by a dotted line.

FIG. 161.—THE ETHMOID BONE SEEN FROM BEHIND AND BELOW. VERTICAL PLATE OF THE ETHMOID AS UPPER PART OF THE BONY SEPTUM OF THE NOSE.

FIG. 162.—AN ETHMOID BONE THE LEFT LATERAL MASS OF WHICH HAS BEEN REMOVED.

The entire surface of the vertical plate is exposed.

Os ethmoidale—Ethmoid bone.

Lachrymal process
Processus lacrimalis

Ethmoidal process
Processus ethmoidalis

FIG. 163.—INNER SURFACE.

Ethmoidal process
Processus ethmoidalis

Maxillary process
Processus maxillaris

FIG. 164.—OUTER SURFACE.

CONCHA NASALIS INFERIOR—THE INFERIOR TURBinate BONE OF THE LEFT SIDE.

Lachrymal crest
Crista lacrimalis
posterior
Lachrymal groove
Sulcus lacrimalis
Hamular process
Hamulus lacrimalis

FIG. 165.—EXTERNAL SURFACE.

Nasal foramen
Foramen nasale

FIG. 166.—INTERNAL SURFACE.

FIG. 167.—ANTERIOR SURFACE.

FIG. 168.—POSTERIOR SURFACE.

Groove of nasal nerve
Sulcus ethmoidalis

OS LACRIMALE: LACHRYMAL BONE
OF THE LEFT SIDE.

OS NASALE: NASAL BONE
OF THE LEFT SIDE.

Alæ of the vomer
Alæ vomeris

FIG. 169.—SEEN FROM THE LEFT SIDE.

FIG. 170.—SEEN FROM ABOVE.

VOMER—THE VOMER.

Bones of the Nasal Region.

FIG. 171.—LEFT SUPERIOR MAXILLARY BONE: EXTERNAL SURFACE.

FIG. 172.—LEFT SUPERIOR MAXILLARY BONE: INTERNAL SURFACE. VIEW INTO THE MAXILLARY SINUS, OR ANTRUM OF HIGHMORE.

Maxilla—Superior maxilla.

FIG. 173.—THE LEFT SUPERIOR MAXILLARY BONE. EXTERNAL SURFACE.

The dental canals are exposed by partial removal of the superficial plate of bone, and their course is shown by means of bristles passed through them.

FIG. 174.—THE LEFT SUPERIOR MAXILLARY BONE. INTERNAL SURFACE.

The foremost and the hindmost of the dental canals have been exposed by the removal of the superficial plate of bone. By means of bristles passed through the canals the situation of the respective dental foramina is indicated. Most of the inner wall of the antrum of Highmore has been cut away.

Maxilla—Superior maxillary bone.

FIG. 175.—THE LEFT SUPERIOR MAXILLARY BONE. SEEN FROM BELOW.

FIG. 176.—SEEN FROM THE INNER SIDE.

FIG. 177.—SEEN FROM BELOW.

THE LEFT SUPERIOR MAXILLARY BONE OF A FŒTUS AT THE END OF THE SIXTH MONTH (MONTHS OF FOUR WEEKS EACH).

Body-length, 12 inches.

FIG. 178.—SEEN FROM THE INNER SIDE.

FIG. 179. SEEN FROM BELOW.

THE LEFT SUPERIOR MAXILLARY BONE OF A BOY BORN AT FULL TERM.

Body-length, 21 inches.

Maxilla—Superior maxillary bone.

FIG. 180.—INNER SURFACE. FIG. 181.—SEEN FROM BEHIND. FIG. 182.—OUTER SURFACE.
OS PALATINUM.—THE PALATE-BONE (OF THE LEFT SIDE).

FIG. 183.—INNER SURFACE. FIG. 184.—SEEN FROM BEHIND. FIG. 185.—OUTER SURFACE.
THE LEFT PALATE-BONE OF A BOY BORN AT FULL TERM.

Body-length, 21 inches.

FIG. 186.—SEEN FROM BEFORE. FIG. 187.—SEEN FROM BEHIND.
OS ZYGOMATICUM—MALAR BONE (OF THE LEFT SIDE).

Supplementary Bones of the Upper Jaw.

FIG. 188.—THE INFERIOR MAXILLARY BONE SEEN FROM THE LEFT SIDE.

FIG. 189.—THE INFERIOR MAXILLARY BONE SEEN FROM ABOVE.

Mandibula—The inferior maxillary bone, lower jaw, or mandible.

FIG. 190.—THE INFERIOR MAXILLARY BONE SEEN FROM BEFORE.

¹ "The angle of the jaw," writes Quain, "... is more or less rounded off." The specimen from which Fig. 190 was drawn has, however, an abnormal projection at each angle, to which in the German nomenclature the name *processus angularis (var.)* is given. This variety is not described by Quain or by Macalister.—Tr.

FIG. 191.—THE INFERIOR MAXILLARY BONE SEEN FROM BEHIND.

Mandibula—The inferior maxillary bone, lower jaw, or mandible.

FIG. 192.—THE INFERIOR MAXILLARY BONE SEEN FROM THE LEFT SIDE.
The mandibular or inferior dental canal has been exposed by the removal of a portion of the superficial plate of bone.

FIG. 195.—MENTAL REGION OF A NEW-BORN BOY. THE LATERAL HALVES OF THE INFERIOR MAXILLARY BONE HAVE NOT YET UNITED; BETWEEN THEM ARE THE MENTAL OSSICLES. SEEN FROM BEFORE.

FIG. 196.—THE FULLY DEVELOPED MENTAL PROTUBERANCE, IN A CHILD AGED SIX AND A HALF MONTHS. THE MENTAL OSSICLES HAVE UNITED WITH EACH OTHER AND WITH THE LATERAL HALVES OF THE INFERIOR MAXILLARY BONE.

Mandibula—The inferior maxillary bone, lower jaw, or mandible.

FIG. 197.—FACIAL PORTION OF THE SKULL OF A WOMAN AGED EIGHTY-FOUR YEARS, SHOWING ATROPHY OF THE ALVEOLAR PROCESSES OF THE SUPERIOR AND INFERIOR MAXILLARY BONES (NUTCRACKER FACE).

FIG. 198.—SEEN FROM ABOVE.

FIG. 199.—SEEN FROM BELOW.

OS HYOIDEUM—THE HYOID BONE.

FIG. 200.—THE HYOID BONE OF A BOY, STILL-BORN AT FULL TERM.
Body-length, 21 inches.

Senile atrophy of the jaws: Os hyoideum—the hyoid bone.

FIG. 201.—SKULL DIVIDED INTO AN ANTERIOR AND A POSTERIOR PORTION BY A FRONTAL SECTION PASSING THROUGH THE MASTOID PROCESSES. THE FIRST CERVICAL VERTEBRA HAS ALSO BEEN DIVIDED BY THE SECTION, AND LEFT ATTACHED TO THE SKULL.

VIEW OF THE ANTERIOR PORTION OF THE CRANIAL CAVITY. THE ANTERIOR AND MIDDLE CRANIAL FOSSÆ, AS WELL AS THE ANTERIOR PORTION OF THE POSTERIOR CRANIAL FOSSA, SEEN FROM BEHIND. ON THE RIGHT SIDE IS A PARAMASTOID PROCESS ARTICULATING WITH THE ATLAS.

Cavum cranii cerebialis—Cranial cavity.

FIG. 202.—SKULL DIVIDED INTO AN ANTERIOR AND A POSTERIOR PORTION BY A FRONTAL SECTION PASSING THROUGH THE MASTOID PROCESSES AND THE BASILAR PORTION OF THE OCCIPITAL BONE. VIEW OF THE POSTERIOR PORTION OF THE CRANIAL CAVITY. THE DIVISION OF THE LONGITUDINAL SULCUS INTO RIGHT LATERAL AND LEFT LATERAL SULCUS IS SHOWN. THE LATERAL SULCUS PASSES ON EITHER SIDE INTO THE SIGMOID SULCUS, WHICH TERMINATES IN THE JUGULAR FORAMEN.

¹ U.S.: Anterior ethmoidal canal.

² See note to p. 48.

FIG. 203.—THE EXTERNAL WALL OF THE NASAL CAVITY WITH THE TURBINATE BONES AND THE NASAL MEATUS: MEDIAN SAGITTAL SECTION. LEFT SIDE.

¹ See note to p. 48.

FIG. 204.—THE EXTERNAL WALL OF THE NASAL CAVITY: MEDIAN SAGITTAL SECTION. LEFT SIDE.

The middle turbinate bone of the nose has been cut away.

Cavum nasi—Nasal cavity.

FIG. 205.—THE EXTERNAL WALL OF THE NASAL CAVITY AFTER REMOVAL OF THE THREE TURBinate BONES AND THE LATERAL MASS OF THE ETHMOID: MEDIAN SAGITTAL SECTION, LEFT SIDE.

FIG. 206.—THE BONY SEPTUM OF THE NOSE SEEN FROM THE LEFT SIDE: SAGITTAL SECTION OF THE FACIAL PART OF THE SKULL, A LITTLE TO THE LEFT OF THE MEDIAN PLANE.

Cavum nasi—Nasal cavity.

¹ I am indebted to Professor Toldt for the following account of the *Sphenoidal canaliculus, which is accurately described neither by Quain nor by Macalister: "It begins in the scaphoid fossa and divides as it ascends into two branches, the inner of which opens into the Vidian canal, while the outer opens on the cerebral surface of the great wing of the sphenoid bone, between the lingula of the sphenoid bone and the foramen ovale." The outer branch is termed by English anatomists the *foramen of Vesalius*, and transmits an emissary vein: the inner branch gives passage to the *sphenoidal branch* of the otic ganglion, by means of which this ganglion communicates with the Vidian nerve.—Tr.

FIG. 207.—BY MEANS OF A NEARLY HORIZONTAL SECTION PASSING THROUGH THE CENTRE OF THE ENTRANCE TO THE ORBIT, THE UPPER PARTS OF THE NASAL FOSSÆ AND OF THE ORBITS ARE DISPLAYED. SEEN FROM BELOW.

Posteriorly the section passes through the body of the sphenoid bone and the root of the pterygoid process, opening up the Vidian canal through its whole length. A bristle has been passed into the *sphenoidal canaliculus¹ from the Vidian canal. The lateral masses of the ethmoid bone (ethmoidal labyrinths) are divided in the horizontal plane.

Cavum nasi et orbita—The nasal fossæ and the orbits.

¹ See note to p. 92.

FIG. 208.—BY MEANS OF A NEARLY HORIZONTAL SECTION IN A PLANE A LITTLE ABOVE THAT OF THE INFERIOR WALLS OF THE ORBITS, THE LOWER PORTIONS OF THE NASAL FOSSÆ AND OF THE ORBITS HAVE BEEN EXPOSED. SEEN FROM ABOVE.

Posteriorly the section passes through the sphenoidal sinuses, in this specimen exceptionally capacious. The upper orifices of the *sphenoidal canaliculus (see note to p. 92) are indicated by bristles.

Cavum nasi et orbita—Nasal cavity and orbits.

FIG. 209.—BY MEANS OF A SECTION PASSING THROUGH THE MOST ANTERIOR PORTIONS OF BOTH ZYGOMATIC ARCHES, THE NASAL CAVITY AND THE ORBITS ARE DIVIDED TOWARDS THEIR POSTERIOR EXTREMITIES IN THE FRONTAL PLANE.

The anterior portion of the skull thus divided is figured from behind. The three turbinate bones of the nose and the three nasal meatus, as well as the bony septum of the nose, are seen in frontal section. The maxillary sinuses are also opened up, and the communication of these sinuses with the general cavity of the nose is to be seen above the uncinate process.

Cavum nasi et orbita—Nasal cavity and orbits.

FIG. 210.—BY MEANS OF A SECTION PASSING VERTICALLY THROUGH THE POSTERIOR PORTIONS OF BOTH LACHRYMAL BONES, THE ANTERIOR PORTION OF THE NASAL CAVITY AND OF THE ORBITS IS DISPLAYED IN FRONTAL SECTION. SEEN FROM BEHIND.

From the right side of the anterior portion of the skull thus divided, a layer of bone a quarter of an inch in thickness has been removed by a section in a plane parallel with the first section, so as to open up the canal of the nasal duct, canalis nasolacimalis, in its entire length. The sounds passed into the two frontal sinuses indicate the orifices of these sinuses in the nasal fossae. A third sound has been passed through the left canal of the nasal duct from the orbit into the nasal cavity.

Cavum nasi et orbita—Nasal cavity and orbits.

¹ See note to p. 92.

FIG. 211.—ANTERIOR PORTION OF THE SKULL, SEPARATED FROM THE POSTERIOR PORTION BY A FRONTAL SECTION PASSING THROUGH THE TWO EXTERNAL AUDITORY MEATUS.

The view from behind shows the posterior nares with the posterior border of the bony septum of the nose, the bony framework of the oral cavity, and the zygomatic fossæ; further, in frontal section, the tympanic cavities, with portions of the bony labyrinths and the internal auditory meatus.

¹ In the English nomenclature *palatine spine* is an alternative name for the *posterior nasal spine*.—Tr.

² The English nomenclature recognises one *palatine groove* only, that running forwards on the under surface of the hard palate from the inferior orifice of the posterior palatine canal, lodging the large palatine nerve and vessels.—Tr.

FIG. 212.—THE HARD PALATE, PALATUM DURUM, WITH THE ALVEOLAR PROCESS OF THE SUPERIOR MAXILLARY BONE, REMOVED BY A SECTION PASSING HORIZONTALLY THROUGH BOTH SUPERIOR MAXILLARY BONES ABOVE THE FLOOR OF THE NASAL FOSSÆ. SEEN FROM BELOW.

FIG. 213.—THE FLOOR (INFERIOR WALL) OF THE NASAL FOSSÆ WITH THE MAXILLARY SINUSES LYING ON EITHER SIDE, SHOWN BY MEANS OF A HORIZONTAL SECTION THROUGH THE SUPERIOR MAXILLARY BONES. SEEN FROM ABOVE.

FIG. 214.—INTERNAL PORTION OF THE LEFT ORBIT, SHOWN BY THE REMOVAL OF THE EXTERNAL WALL. SEEN FROM THE LEFT SIDE.

FIG. 215.—EXTERNAL PORTION OF THE LEFT ORBIT, SHOWN BY THE REMOVAL OF THE INTERNAL WALL OF THE ORBIT AND OF A PORTION OF THE SUPERIOR MAXILLARY BONE IN A LONGITUDINALLY HEMI-SECTED SKULL.

Orbita—The orbits.

FIG. 216.—THE LEFT MAXILLARY SINUS, OR ANTRUM OF HIGHMORE, SHOWN BY THE REMOVAL OF THE EXTERNAL WALL OF THE BODY OF THE SUPERIOR MAXILLARY BONE AND THE MALAR BONE. SEEN FROM THE LEFT SIDE.

¹ See note to p. 92.

FIG. 217.—THE LEFT SPHENOMAXILLARY FOSSA, SHOWN BY REMOVAL OF THE MALAR BONE, THE TEMPORAL BONE, AND THE GREAT WING OF THE SPHENOID BONE. SEEN FROM THE LEFT SIDE.

A sound has been passed through the Vidian canal, and another through the *sphenoidal canaliculus¹ (exceptionally large in this specimen).

Sinus maxillaris—Maxillary sinus, or antrum of Highmore.—Fossa pterygopalatina—Sphenomaxillary fossa.

FIG. 218.—A LARGE WORMIAN BONE IN THE UPPERMOST PART OF THE LAMBDOID SUTURE.

FIG. 219.—TRANSVERSE SUBDIVISION OF THE SQUAMA OCCIPITALIS. SEVERAL WORMIAN BONES IN THE LAMBDOID SUTURE.

¹ See note to p. 57.

Ossa suturarum—Wormian bones.

FIG. 220.—SIDE-VIEW OF THE SKULL: NORMA LATERALIS.

FIG. 221.—SKULL SEEN FROM ABOVE
NORMA VERTICALIS.FIG. 222.—FRONT VIEW OF THE SKULL:
NORMA FRONTALIS.

TRANSLATOR'S NOTE.—The above measurements do not fully correspond with those used by English craniologists. For this reason a purely literal translation of the German terms has been given.

The Principal Measurements of the Skull.

FIG. 223.—HUMAN EMBRYO, FOUR WEEKS OLD.

The umbilical vesicle has been opened.

¹ See note to p. 57.

FIG. 224.—THE LEFT HALF OF THE SKULL OF A HUMAN FÆTUS AT THE END OF THE FOURTH MONTH (MONTHS OF FOUR WEEKS EACH). SEEN FROM WITHIN. THE CARTILAGINOUS PORTION OF THE PRIMORDIAL CRANIUM (CHONDROCRANIUM) IS COLOURED BLUE.

Body-length, $4\frac{1}{2}$ inches.

Development of the Skull.

Squamous portion of the temporal bone
Squama temporalis

Cartilaginous bar of the mandibular or first
visceral arch (Meckel's cartilage)

The incus
Incus

Mandible
Mandibula

Cartilaginous portion of the primordial cranium
Pars cartilaginea cranii primordialis

Cartilaginous bar of the hyoid or second
visceral arch (primitive cartilage of
the styloid process)

The malleus
Malleus

Tympanic ring
Annulus tympanicus

FIG. 225.—THE SQUAMOUS PORTION OF THE TEMPORAL BONE, THE TYMPANIC MEMBRANE WITH THE MALLEUS, THE INCUS, AND MECKEL'S CARTILAGE, FROM A HUMAN FÆTUS IN THE FIRST HALF OF THE FIFTH MONTH (MONTHS OF FOUR WEEKS EACH). SEEN FROM WITHIN.

Body-length, $5\frac{1}{2}$ inches.

Squamous portion of the temporal bone
Squama temporalis

Parietal bone
Os parietale

Membranous portion of the primordial cranium
Pars membranacea cranii primordialis

Tympanic ring
Annulus tympanicus

Greater wing
Ala magna

Squamous portion of the frontal bone
Squama frontalis

Cartilaginous bar of the mandibular or first
visceral arch (Meckel's cartilage)

Intermembranous part of the supra-occipital
portion of the occipital bone¹
Pars intermembranacea squamæ occipitalis

Cartilaginous portion of the primordial cranium
Pars cartilaginea cranii primordialis

Intracartilaginous part of the supra-occipital
portion of the occipital bone¹
Pars intracartilaginea squamæ occipitalis

Foramen magnum
Foramen occipitale magnum

$\frac{1}{1}$

Lateral portion of the occipital bone (exoccipital)¹
Pars lateralis ossis occipitalis

Basilar portion of the occipital bone (basi-occipital)¹
Pars basilaris ossis occipitalis

Primitive cartilage of the petrous portion
of the temporal bone
Knorpelanlage der Pars petrosa

Primitive cartilage of the hyoid bone
Knorpelanlage des Os hyoideum

Cartilaginous portion of the primordial cranium
Pars cartilaginea cranii primordialis

Mandible
Mandibula

Maxilla
Maxilla

¹ See note to p. 57.

FIG. 226.—THE SKULL OF A HUMAN FÆTUS IN THE MIDDLE OF THE FIFTH MONTH (MONTHS OF FOUR WEEKS EACH). SEEN FROM THE LEFT SIDE AND BELOW.

Body length, $6\frac{1}{2}$ inches. The cartilaginous bars of the visceral arches are displayed, and these, together with the cartilaginous portion of the primordial cranium, are coloured blue. Part of the cartilage of the left side of the inferior maxilla has been removed in order to lay bare a portion of Meckel's cartilage which lies beneath it.

$\frac{1}{1}$

FIG. 227.—THE DRIED SKULL OF A HUMAN FÆTUS IN THE MIDDLE OF THE FOURTH MONTH (MONTHS OF FOUR WEEKS EACH).

Body-length, $3\frac{1}{2}$ inches.

$\frac{2}{1}$

FIG. 228.—THE DRIED SKULL OF A HUMAN FÆTUS AT THE END OF THE SIXTH MONTH (MONTHS OF FOUR WEEKS EACH).

Body-length, $11\frac{1}{2}$ inches.

Development of the Skull.

Anterolateral fontanelle
Fonticulus sphenoidalis

FIG. 229.—THE DRIED SKULL OF AN INFANT BORN AT FULL TERM. SEEN FROM THE LEFT SIDE.

Body-length, $20\frac{1}{2}$ inches. In the anterolateral fontanelle there remains part of the membranous portion of the primordial cranium (membrane of the fontanelle); in the posterolateral fontanelle there remains part of the cartilaginous portion of the primordial cranium.

FIG. 230.—THE DRIED SKULL OF AN INFANT BORN AT FULL TERM. SEEN FROM ABOVE.

Body-length, $20\frac{1}{2}$ inches. In the anterior and in the posterior fontanelle there remain parts of the membranous portion of the primordial cranium.

¹ See note to p. 57.

Development of the Skull.

SKELETON EXTREMITATUM,
SUPERIORIS ET INFERIORIS

THE SKELETON OF THE UPPER
AND LOWER EXTREMITIES
(THE APPENDICULAR SKELETON)

FIG 231.—THE SKELETON OF THE UPPER EXTREMITY.

Skeleton extremitatis superioris—The skeleton of the upper extremity.

FIG. 232.—THE SKELETON OF THE LOWER EXTREMITY.

Skeleton extremitatis inferioris—The skeleton of the lower extremity.

FIG. 233.—THE SHOULDER-GIRDLE AND ITS RELATION TO THE THORAX. SEEN FROM ABOVE.

FIG. 234.—THE SHOULDER-GIRDLE AND ITS RELATION TO THE THORAX. SEEN FROM BEHIND.

Cingulum extremitatis superioris—The shoulder-girdle.

FIG. 235.—THE RIGHT CLAVICLE SEEN FROM ABOVE.

FIG. 23.—THE RIGHT CLAVICLE SEEN FROM BELOW.

FIG. 237.—STERNAL EXTREMITY OF THE RIGHT CLAVICLE OF A FEMALE AGED TWENTY YEARS, WITH A STERNAL EPIPHYSIS. SEEN FROM BEFORE.

Clavicula—The clavicle.

FIG. 238.—THE RIGHT SCAPULA SEEN FROM BEFORE.

FIG. 240.—THE RIGHT SCAPULA SEEN FROM THE OUTER OR AXILLARY ASPECT.

FIG. 239.—THE RIGHT SCAPULA SEEN FROM BEHIND.

Scapula—The shoulder-blade.

FIG. 241.—IN THE FIFTH MONTH OF INTRA-UTERINE LIFE (MONTHS OF FOUR WEEKS EACH).

FIG. 242.—IN THE SECOND YEAR OF LIFE.

FIG. 243.—IN THE FIFTEENTH YEAR OF LIFE.

FIG. 244.—IN THE SEVENTEENTH YEAR OF LIFE.

FIG. 245.—IN THE NINETEENTH YEAR OF LIFE.

Development of the Scapula.

FIG. 246.—THE RIGHT HUMERUS SEEN FROM BEFORE.

FIG. 247.—THE RIGHT HUMERUS SEEN FROM BEHIND.

Humerus—The humerus.

FIG. 250.—FROM A GIRL AGED FOUR MONTHS.

FIG. 252.—FROM A BOY AGED THREE YEARS (HORIZONTAL SECTION.)

FIG. 251.—FROM A BOY AGED TWO YEARS.

THE CENTRES OF OSSIFICATION OF THE PROXIMAL EXTREMITY OF THE HUMERUS.

FIG. 253.—FROM A BOY AGED THIRTEEN YEARS.

FIG. 254.—FROM A GIRL AGED NINETEEN YEARS.

THE COMMON EPIPHYSIS OF THE PROXIMAL EXTREMITY OF THE HUMERUS.

FIG. 255.—FROM A BOY AGED TWO AND A HALF YEARS.

FIG. 256.—FROM A BOY AGED THIRTEEN YEARS.

FIG. 257.—FROM A BOY AGED SEVENTEEN YEARS.

THE EPIPHYSES OF THE DISTAL EXTREMITY OF THE HUMERUS.

Development of the Humerus.

FIG. 258.—ANTERIOR ASPECT.

FIG. 259.—POSTERIOR ASPECT.
THE ULNA OF THE RIGHT SIDE.

FIG. 260.—EXTERNAL ASPECT.

FIG. 261.—TRANSVERSE (HORIZONTAL) SECTION THROUGH THE MIDDLE OF THE ULNA AND THE RADIUS, WITH THE INTEROSSEOUS MEMBRANE, IN SUPINATION.

Ossa antibrachii—The bones of the forearm.

FIG. 262.—ANTERIOR ASPECT.

FIG. 263.—POSTERIOR ASPECT.
THE RADIUS OF THE RIGHT SIDE.

FIG. 264.—INTERNAL ASPECT.

FIG. 265.—THE PROXIMAL EXTREMITIES OF THE BONES OF THE RIGHT FORE-ARM SEEN FROM ABOVE.

FIG. 266.—THE DISTAL EXTREMITIES OF THE BONES OF THE RIGHT FOREARM SEEN FROM BELOW.

FIG. 267.—THE BONES OF THE RIGHT FOREARM, IN SUPINATION.

FIG. 268.—THE BONES OF THE RIGHT FOREARM, IN PRONATION.

The Bones of the Forearm in Supination and Pronation.

Centres of ossification
in the olecranon

Tuberosity of the ulna
Tuberositas ulnae

Double epiphysis of
the olecranon in
the act of uniting
with the shaft

FIG. 269.—FROM A BOY AGED
THIRTEEN YEARS.

FIG. 270.—FROM A BOY AGED
SEVENTEEN YEARS.

THE DEVELOPMENT OF THE EPIPHYSIS OF THE PROXIMAL EXTREMITY OF THE ULNA.

FIG. 271.—FROM A GIRL AGED
SIX YEARS.

FIG. 272.—FROM A YOUNG MAN AGED
NINETEEN YEARS.

THE DEVELOPMENT OF THE EPIPHYSIS OF THE DISTAL EXTREMITY OF THE ULNA.

Centre of ossification in
the head of the radius

FIG. 273.—FROM A BOY AGED FIVE YEARS.

THE DEVELOPMENT OF THE EPIPHYSIS OF THE PROXIMAL EXTREMITY OF THE RADIUS.

Epiphysis of the head of
the radius in the act
of uniting with the
shaft

Tuberosity of the
radius
Tuberositas radii

Centre of ossification in the
distal extremity of the
radius

FIG. 275.—FROM A GIRL AGED
TWO YEARS.

FIG. 276.—FROM A YOUNG MAN AGED
NINETEEN YEARS.

THE DEVELOPMENT OF THE EPIPHYSIS OF THE DISTAL EXTREMITY OF THE RADIUS.

FIG. 277.—PALMAR ASPECT OF THE SKELETON OF THE RIGHT HAND (FACIES VOLARIS).

Skeletum manus—The skeleton of the hand.

FIG. 278.—DORSAL ASPECT OF THE BONES OF THE RIGHT HAND SEEN FROM THE DORSAL SIDE (FACIES DORSALIS).

Skeleton manus—The skeleton of the hand.

FIG. 279.—DISTAL ASPECT.

FIG. 280.—POSTERIOR ASPECT.

OS NAVICULARE MANUS—THE RIGHT SCAPHOID BONE.

FIG. 281.—RADIAL ASPECT.

FIG. 282.—ULNAR ASPECT.

OS LUNATUM—THE RIGHT LUNAR (OR SEMILUNAR) BONE.

FIG. 283.—RADIAL ASPECT.

FIG. 284.—PALMAR ASPECT.

OS TRIQUETRUM—THE RIGHT PYRAMIDAL OR CUNEIFORM BONE.

FIG. 285.—PALMAR ASPECT.

FIG. 286.—POSTERIOR ASPECT.

OS PISIFORME—THE RIGHT PISIFORM BONE.

FIG. 287. THE POSTERIOR ASPECT OF THE RIGHT CARPUS, CONTAINING AN OS CENTRALE.

Ossa carpi—The bones of the carpus: first, superior, or proximal row.

FIG. 288.—DISTAL ASPECT.

FIG. 289.—ULNAR ASPECT.

OS MULTANGULUM MAJUS—THE RIGHT TRAPEZIUM.

FIG. 290.—ULNAR ASPECT.

FIG. 291.—RADIAL ASPECT.

OS MULTANGULUM MINUS—THE RIGHT TRAPEZOID BONE.

FIG. 292.—RADIAL ASPECT.

FIG. 293.—ULNAR ASPECT.

OS CAPITATUM—THE RIGHT OS MAGNUM OR CAPITATE BONE.

FIG. 294.—DISTAL ASPECT.

FIG. 295.—PROXIMAL ASPECT.

OS HAMATUM—THE RIGHT UNCIFORM BONE.

Ossa carpi—The bones of the carpus: second, inferior, or distal row.

FIG. 296.—RADIAL ASPECT OF THE FIVE METACARPAL BONES OF THE RIGHT HAND.

FIG. 297.—ULNAR ASPECT OF THE FIVE METACARPAL BONES OF THE RIGHT HAND.

FIG. 298.—RADIAL ASPECT OF THE BONES OF THE RIGHT INDEX FINGER (MARGO RADIALIS DIGITI SECUNDI).

The Metacarpal Bones of the Hand and the Phalanges of the Fingers.

FIG. 299.—IN THE SECOND YEAR OF LIFE.

CENTRES OF OSSIFICATION OF THE CARPUS (SECTIONS PARALLEL TO THE POSTERIOR SURFACE).

FIG. 300.—IN THE SEVENTH YEAR OF LIFE.

FIG. 301.—MIDDLE FINGER IN THE SECOND YEAR OF LIFE.

FIG. 302.—MIDDLE FINGER IN THE SEVENTH YEAR OF LIFE.

FIG. 304.—BONES OF THE THUMB IN THE FOURTH YEAR OF LIFE.

FIG. 305.—METACARPAL BONE OF THE THUMB IN THE EIGHTEENTH YEAR OF LIFE.

FIG. 303.—EPIPHYSES OF THE BONES OF THE MIDDLE FINGER IN THE ACT OF UNITING WITH THE SHAFTS: SEVENTEENTH YEAR OF LIFE.

Development of the Bones of the Hand.

FIG. 306.—PELVIS MULIEBRIS—THE FEMALE PELVIS. ANTERIOR ASPECT.

The formation of the pelvis out of the sacrum and coccyx and the two innominate bones. The iliac portions of the innominate bones with the base of the sacrum constitute the upper or false pelvis; the pubic and ischiatic portions of the innominate bones with the sacrum and the coccyx constitute the lower or true pelvis; the boundary between the false and the true pelvis corresponds with the upper aperture or entrance of the true pelvis, the line separating the two being known as the brim or inlet of the true pelvis. Regarded as the means of attachment of the lower limb to the trunk, the pelvis is the *cingulum extremitatis inferioris*, or pelvic girdle.

Cingulum extremitatis inferioris—Pelvic girdle.

FIG. 307.—PELVIS VIRILIS—THE MALE PELVIS. ANTERIOR ASPECT.

The upper or false and the lower or true pelvis, pelvis major and pelvis minor. The brim of the true pelvis is divided into a sacral, an iliac, and a pubic portion.

Cingulum extremitatis inferioris—Pelvic girdle.

FIG. 308.—PELVIS VIRILIS—THE MALE PELVIS. POSTERIOR ASPECT.

The pelvic outlet, apertura pelvis (minoris) inferior, in the bony pelvis appears to be bounded on either side by the lower borders of the pubis and the ischium and the greater and lesser sciatic notches, incisurae ischiadicæ major et minor, and behind by the projecting part of the sacrum and by the coccyx. But inasmuch as on either side there are two strong ligaments arising from the sacrum and coccyx, the great and the small sacrosiatic ligaments, ligamenta sacrotuberosum et sacrospinosum, which stretch across the two sciatic notches, and thus enlarge the posterior and lateral walls of the true pelvis, by this means the sciatic notches are filled in, and the outlet of the pelvis is notably diminished in size.

Cingulum extremitatis inferioris—Pelvic girdle.

FIG. 309.—THE MEDIAN-SAGITTAL OR ANTERO-POSTERIOR DIAMETERS OF THE TRUE PELVIS.

FIG. 310.—THE DIAMETERS OF THE PELVIC INLET (*APERTURA PELVIS SUPERIOR*).

The Principal Diameters of the True Pelvis.

FIG. 312.—THE RIGHT HIP-BONE, OUTER SIDE. POSTERIOR ASPECT.

The outer surface of the ilium is seen, divided by the three gluteal lines into areas from which the three gluteal muscles respectively arise. The spine of the ischium projects backwards from the posterior border of the hip-bone; above this spine is the iliosciatic or great sciatic notch, below it is the small sciatic notch. The lowest portion of the bone is formed by the backwardly projecting tubercity of the ischium. The acetabulum is viewed obliquely from behind.

Body of the pubis: The use of this term by English anatomists is a variable one. Macalister, whose terminology here, as usual, is in conformity with that of Continental anatomists, writes: "The pubis consists of a body which forms a little less than one-fifth of the acetabulum . . .", Quain, on the other hand, writes: "The flat portion between the rami [of the pubis] is the body", and Young, in his "Synopsis of Human Anatomy" (U.S.), follows Quain's usage. The Continental application of the term has, however, the advantage in the point of consistency, the body being then, in the case of each of the three elements of the hip-bone, the thickened portion taking part in the formation of the acetabulum, of which the body of the pubis constitutes about one-fifth, the body of the ilium nearly two-fifths, and the body of the ischium the remainder.—Tr.

FIG. 313.—THE RIGHT HIP-BONE, OUTER SIDE. SEEN FROM THE RIGHT.

For this illustration the hip-bone has been so placed that a direct view of the interior of the acetabulum is obtained; this consists of two portions, the rough, nearly circular fossa of the acetabulum, and the horse-shoe shaped articular surface (facies lunata—covered in the recent state with cartilage); opposite the fossa of the acetabulum, the rim bounding the depression is interrupted by the cotyloid notch (incisura acetabuli). The three bones which, separate at first, subsequently unite to form the hip-bone, all take part in the formation of the acetabulum (see note ¹ above).

FIG. 314.—IN THE FIFTH MONTH OF FŒTAL LIFE (MONTHS OF FOUR WEEKS EACH).

FIG. 315.—IN THE SIXTH MONTH OF FŒTAL LIFE (MONTHS OF FOUR WEEKS EACH).

FIG. 316.—IN THE FOURTH YEAR OF LIFE.

FIG. 317.—IN THE FOURTEENTH YEAR OF LIFE.

FIG. 318.—IN THE SEVENTEENTH YEAR OF LIFE.

Development of the Hip-Bone.

FIG. 319.—THE RIGHT FEMUR,
ANTERIOR ASPECT.

FIG. 320.—THE RIGHT FEMUR,
POSTERIOR ASPECT.

Femur—The femur, or thigh-bone.

FIG. 321.—THE RIGHT FEMUR,
INTERNAL ASPECT.

FIG. 322.—THE UPPER PORTION OF THE RIGHT FEMUR: THIRD TROCHANTER (VAR.). SEEN FROM BEHIND.

FIG. 323.—TRANSVERSE SECTION THROUGH THE MIDDLE OF THE SHAFT OF THE RIGHT FEMUR.

FIG. 324.—HORIZONTAL SECTION THROUGH THE PROXIMAL EXTREMITY OF THE RIGHT FEMUR, ABOVE THE SMALL TROCHANTER, TO SHOW THE CALCAR FEMORALE.

FIG. 325.—DISTAL EXTREMITY OF THE RIGHT FEMUR. SEEN FROM THE OUTER SIDE.

FIG. 326.—DISTAL ARTICULAR EXTREMITY OF THE RIGHT FEMUR. SEEN FROM BELOW.

Femur—The femur.

FIG. 327.—FROM A BOY STILL-BORN AT FULL TERM.
Body-length, 21 inches.

FIG. 328.—FROM A BOY AGED NINE AND A HALF MONTHS.

THE CENTRES OF OSSIFICATION OF THE PROXIMAL AND DISTAL EPIPHYSES.

FIG. 329.—FROM A GIRL AGED SEVEN YEARS.

FIG. 330.—FROM A GIRL AGED FIFTEEN YEARS.

THE EPIPHYSES OF THE PROXIMAL EXTREMITY.

FIG. 331.—IN FRONTAL SECTION.

FIG. 332.—IN SAGITTAL SECTION.
THE EPIPHYSES OF THE DISTAL EXTREMITY FROM A GIRL AGED FIFTEEN YEARS.

Development of the Femur.

FIG. 333.—ANTERIOR ASPECT.

FIG. 334.—POSTERIOR ASPECT.

THE RIGHT TIBIA.

Base of the patella—
Basis patellæ

FIG. 335.—ANTERIOR SURFACE.

Base of the patella
Basis patellæ

FIG. 336.—POSTERIOR SURFACE.

THE RIGHT PATELLA.

Ossa cruris—The bones of the leg: the patella, rotula, or knee-pan.

FIG. 338.—FACIES ARTICULARIS SUPERIOR TIBIÆ—THE PROXIMAL ARTICULAR SURFACE OF THE RIGHT TIBIA.

Note to Fig. 338.—*Anterior and Posterior Intercondylar Fossa*.—These terms are used neither by Quain nor by Macalister. The latter speaks of the hollows in question as *depressions for the crucial ligaments*.—Tr.

FIG. 339.—THE DISTAL ARTICULAR SURFACES OF THE BONES OF THE RIGHT LEG.

FIG. 337.—RIGHT TIBIA SEEN FROM THE OUTER SIDE.

FIG. 340.—EXTERNAL ASPECT.

FIG. 341.—INTERNAL ASPECT.

THE RIGHT FIBULA OR PERONEAL BONE.

FIG. 342.—TRANSVERSE SECTION THROUGH THE MIDDLE OF THE BONES OF THE RIGHT LEG, WITH THE INTEROSSEOUS MEMBRANE.

Ossa cruris—Bones of the leg.

FIG. 343.—FROM A BOY STILL-BORN AT FULL TERM.
Body length, 21 inches.

FIG. 344.—FROM A BOY AGED NINE AND A HALF MONTHS.

FIG. 345.—FROM A BOY AGED ONE AND A HALF YEARS.

FIG. 346.—PROXIMAL AND DISTAL PORTIONS OF THE FIBULA OF A GIRL AGED FOUR AND A HALF YEARS.

FIG. 347.—PROXIMAL PORTION OF THE TIBIA OF A GIRL AGED FIFTEEN YEARS. SAGITTAL SECTION.

FIG. 348.—THE DISTAL PORTIONS OF THE BONES OF THE LEG OF A GIRL AGED FIFTEEN YEARS. FRONTAL SECTION.

FIG. 349.—BONES OF THE RIGHT FOOT SEEN FROM THE OUTER SIDE: *FACIES DORSALIS PEDIS* ET *MARGO LATERALIS PEDIS*.

FIG. 350.—THE BONES OF THE RIGHT FOOT SEEN FROM THE INNER SIDE: *MARGO MEDIALIS PEDIS*.

Skeleton pedis—Bones of the foot.

FIG. 351.—THE BONES OF THE RIGHT FOOT SEEN FROM THE PLANTAR SIDE: FACIES PLANTARIS PEDIS.

Skeleton pedis—Bones of the foot.

FIG. 352.—THE BONES OF THE RIGHT FOOT SEEN FROM THE DORSAL SIDE (ARRANGED IN TWO LONGITUDINAL ROWS).

Skeleton pedis. Bones of the foot.

FIG. 353.—SEEN OBLIQUELY FROM WITHIN AND BEFORE.

FIG. 354.—SEEN OBLIQUELY FROM WITHOUT AND BEHIND.

THE RIGHT CALCANEUM OR OS CALCIS.

FIG. 355.—SEEN FROM BELOW.

FIG. 356.—SEEN FROM BEHIND.

THE RIGHT ASTRAGALUS.

FIG. 357.—THE RIGHT ASTRAGALUS WITH AN OS TRIGONUM (VAR.). SEEN FROM BELOW.

Ossa tarsi—Bones of the tarsus.

FIG. 358.—ANTERIOR ASPECT.

FIG. 359.—POSTERIOR ASPECT.

OS NAVICULARE PEDIS—THE RIGHT NAVICULAR OR SCAPHOID BONE OF THE FOOT.

FIG. 360.—EXTERNAL ASPECT.

FIG. 361.—ANTERIOR ASPECT.

OS CUNEIFORME I.—THE RIGHT INTERNAL CUNEIFORM BONE.

INNER ASPECT.

OUTER ASPECT.

INNER ASPECT.

OUTER ASPECT.

FIG. 362.—OS CUNEIFORME II.—THE RIGHT MIDDLE CUNEIFORM BONE.

FIG. 363.—OS CUNEIFORME III.—THE RIGHT EXTERNAL CUNEIFORM BONE.

FIG. 364.—INNER ASPECT.

FIG. 365.—SEEN OBLIQUELY FROM WITHOUT AND BEHIND.

FIG. 366.—OUTER ASPECT.

OS CUBOIDEUM—THE RIGHT CUBOID BONE.

Ossa tarsi—Bones of the tarsus.

FIG. 367.—THE METATARSAL BONES OF THE RIGHT FOOT SEEN FROM THE INNER SIDE.

FIG. 368.—THE METATARSAL BONES OF THE RIGHT FOOT SEEN FROM THE OUTER SIDE.

FIG. 369.—PLANTAR ASPECT OF THE HEAD OF THE METATARSAL BONE OF THE GREAT TOE (CAPITULUM OSSIS METATARSALIS HALLUCIS).

FIG. 370.—THE PHALANGES OF THE SECOND TOE SEEN FROM THE INNER SIDE (MARGO MEDIALIS DIGITI SECUNDI PEDIS).

FIG. 371.—FROM A HUMAN FÆTUS IN THE MIDDLE OF THE NINTH MONTH (MONTHS OF FOUR WEEKS EACH).
Body-length, $17\frac{1}{2}$ inches.

FIG. 372.—FROM A BOY STILL-BORN AT FULL TERM.
Body-length, 22 inches.

FIG. 373.—FROM A BOY AGED TWELVE WEEKS.

FIG. 374.—FROM A BOY AGED THREE YEARS.

Development of the Bones of the Foot.

FIG. 375.—OSSIFICATION OF THE BONES OF THE FOOT IN A GIRL AGED SIX YEARS.

FIG. 376.—FROM A GIRL AGED EIGHT YEARS.

FIG. 377.—FROM A YOUNG MAN AGED EIGHTEEN YEARS (EPIPHYSIS BEGINNING TO UNITE).

EPIPHYSIS OF THE TUBEROSITY OF THE CALCANEUM.

Development of the Bones of the Foot.

INDEX

TO THE

REGIONS OF THE HUMAN BODY

AND TO THE

OSTEOLOGY

INDEX

TO THE REGIONS OF THE HUMAN BODY.

A.
 ABDOMEN, external region of the, 2, 3
 regions of, 2, 3
 Abdominal regions, 2, 3
 Acromial region, 3, 4
 Anal region, 5
 Auricular region, 3, 4
 Axillary region, 2

B.
 Back, medial region of the, 3
 regions of the, 3
 Brachial region, anterior, 2, 3
 external, 2, 3
 internal, 2, 3
 posterior, 3
 Buccal region, 4

C.
 Calcaneal region, 2, 3
 Calf, region of the, 3
 Clavicular region, 4
 Crural region, anterior, 2
 external, 2, 3
 internal, 2
 posterior, 3

D.
 Deltoid region, 2, 3
 Digital regions of the foot, dorsal, 2

E.
 Elbow, anterior region of the, 2
 external region of the, 2, 3
 internal region of the, 2
 posterior region of the, 2, 3
 Epigastric region, 2

F.
 Face, regions of the, 2, 4
 Femoral region, anterior, 2
 external, 2, 3
 internal, 2, 3
 posterior, 2
 Fingers, palmar regions of the, 2, 3
 dorsal regions of the, 3
 Foot, dorsal region of the, 2
 plantar region of the, 3
 Forearm, dorsal region of the, 2
 palmar region of the, 2, 3
 radial region of the, 2
 ulnar region of the, 2, 3
 Fossa axillaris, 2*a*
 carotica, 4*a*
 jugularis, 3*a*
 poplitea, 3*a*
 retromandibularis, 4*a*
 supraclavicularis major, 4*a*
 minor, 4*a*

Fossa, axillary, 2
 infraclavicular, 2, note, 4, note
 retromandibular, 4
 supraclavicular, greater, 4
 lesser, 4

Fovea nuchae, 3*a*
 Frontal region, 2, 4
 Furrow, carotid, 4
 nuchal, 3

G.
 Gluteal region, 3, 5

H.
 Ham, the, 3
 Hand, dorsal region of the, 2, 3
 palmar region of the, 2, 3
 Head, regions of the, 4
 Hip, region of the, 2, 3
 Hyoid region, 4
 Hypochondriac region, 2, 3
 Hypogastric region, 2

I.
 Infraclavicular region, 2, 4
 Inframammary region, 2
 Infra-orbital region, 4
 Infrascapular region, 3
 Inguinal region, 2
 Interscapular region, 3

K.
 Knee, anterior region of the, 2
 posterior region of the, 3

L.
 Labial region, lower, 4
 upper, 4
 Laryngeal region, 4
 Lisfranc, tubercle of, 4*a*
 Lower extremity, regions of the, 2, 3
 Lumbar region, 3

M.
 Malleolar region, external, 3
 internal, 2
 Mammary region, 2
 Mastoid region, 3, 4
 Mental region, 4
 Mesogastric region, 2
 Mohrenheimer's space, 2, note, 4, note

N.
 Nasal region, 2, 4
 Neck, anterior region of the, 2
 external region of the, 2
 regions of the, 4
 Nuchal region, 3, 4

O.
 Occipital region, 3, 4
 Olecranon, region of the, 2, 3
 Oral region, 2, 4
 Orbital region, 2, 4

P.
 Palpebral region, lower, 4
 upper, 4
 Parietal region, 2-4
 Parotidomasseteric region, 4
 Patellar region, 2
 Pectoral regions, 2, 3
 region, external, 2, 3
 Perineal region, 3, 5
 Pubic region, 2
 Pudendal region, 2, 5

R.

Regions of the human body, 1-5

Regio *vel* regiones:

abdominis, 2*a*, 3*a*
 lateralis, 2*a*, 3*a*
 acromialis, 3*a*, 4*a*
 aualis, 5*a*
 antibrachii dorsalis, 2*a*
 radialis, 2*a*
 ulnaris, 2*a*, 3*a*
 volaris, 2*a*, 3*a*
 auricularis, 3*a*, 4*a*
 axillaris, 2*a*
 brachii anterior, 2*a*, 3*a*
 lateralis, 2*a*, 3*a*
 medialis, 2*a*, 3*a*
 posterior, 3*a*
 buccalis, 4*a*
 calcanea, 2*a*, 3*a*
 capitis, 4*a*
 clavicularis, 4*a*
 colli, 4*a*
 anterior, 2*a*
 lateralis, 2*a*, 4*a*
 corporis humani, 1*a*-5*a*
 coxæ, 2*a*, 3*a*
 cruris anterior, 2*a*
 lateralis, 2*a*, 3*a*
 medialis, 2*a*
 posterior, 3*a*
 cubiti anterior, 2*a*
 lateralis, 2*a*, 3*a*
 medialis, 2*a*
 posterior, 2*a*, 3*a*
 deltoidea, 2*a*, 3*a*
 digitorum (manus), 2*a*, 3*a*
 pedis, 2*a*
 dorsales digitorum (manus), 3*a*
 pedis, 2*a*
 dorsalis manus, 2*a*, 3*a*
 pedis, 2*a*
 dorsi, 3*a*
 epigastrica, 2*a*
 extremitatis inferioris, 2*a*, 3*a*
 superioris, 2*a*, 3*a*
 faciei, 2*a*, 4*a*
 femoris anterior, 2*a*
 lateralis, 2*a*, 3*a*
 medialis, 2*a*, 3*a*
 posterior, 3*a*
 frontalis, 2*a*, 4*a*
 genu anterior, 2*a*
 posterior, 3*a*
 glutea, 3*a*, 5*a*
 hyoidea, 4*a*
 hypochondriaca, 2*a*, 3*a*
 hypogastrica, 2*a*
 infraclavicularis, 2*a*, 4*a*
 infraomammalis, 2*a*
 infraorbitalis, 4*a*
 infrascapularis, 3*a*
 inguinalis, 2*a*
 interscapularis, 3*a*
 labialis inferior, 4*a*
 superior, 4*a*
 laryngea, 4*a*
 lumbalis, 3*a*
 malleolaris lateralis, 3*a*
 medialis, 2*a*
 mammarialis, 2*a*
 mastoidea, 3*a*, 4*a*
 mediana dorsi, 3*a*
 mentalis, 4*a*
 mesogastrica, 2*a*

Regio *vel* regiones:

nasalis, 2*a*, 4*a*
 nuchæ, 3*a*, 4*a*
 occipitalis, 3*a*, 4*a*
 olecrani, 2*a*, 3*a*
 oralis, 2*a*, 4*a*
 orbitalis, 2*a*, 4*a*
 palpebralis inferior, 4*a*
 superior, 4*a*
 parietalis, 2*a*, 3*a*, 4*a*
 parotideomasseterica, 4*a*
 patellaris, 2*a*
 pectoris, 2*a*, 3*a*
 lateralis, 2*a*, 3
 perinealis, 3*a*, 5*a*
 plattaris pedis, 3*a*
 pubica, 2*a*
 pudendalis, 2*a*, 5*a*
 retromalleolaris lateralis, 3*a*
 medialis, 2*a*
 sacralis, 3*a*, 5*a*
 scapularis, 3*a*
 sternalis, 2*a*
 sternocleidomastoidea, 4*a*
 subhyoidea, 4*a*
 subinguinalis, 2*a*
 submaxillaris, 4*a*
 submental, 4*a*
 supra-orbitalis, 4*a*
 suprascapularis, 3*a*
 suprasternalis, 4*a*
 suralis, 3*a*
 temporalis, 2*a*, 3*a*, 4*a*
 thyroidea, 4*a*
 trochanterica, 2*a*, 3*a*
 umbilicalis, 2*a*
 ungulares, 2*a*, 3*a*
 urogenitalis, 5*a*
 volares digitorum, 2*a*, 3*a*
 volaris manus, 2*a*, 3*a*
 zygomatica, 4*a*

Retromalleolar region, external, 3
 internal, 2

S.
 Sacral region, 3, 5
 Scapular region, 3
 Space, Mohrenheimer's, 2, note, 4, note
 popliteal, 3
 Sternal region, 2
 Sternocleidomastoid region, 4
 Subhyoid region, 4
 Subinguinal region, 2
 Submaxillary region, 4
 Submental region, 4
 Supra-orbital region, 4
 Suprascapular region, 3
 Suprasternal region, 4

T.
 Temporal region, 2-4
 Thyroid region, 4
 Triangle, deltoideopectoral, 2, 4
 infraclavicular, 2, note, 4, note
 omoclavicular, 4
 Trigonum deltoideopectoral, 2*a*, 4*a*
 omoclavicular, 4*a*
 Trochanteric region, 2, 3
 Tubercle of Lisfranc, 4*a*

U.
 Umbilical region, 2
 Ungual regions, 2, 3
 Upper extremity, regions of the, 2, 3
 Urogenital region, 5

Z.
 Zygomatic region, 4

INDEX

TO THE OSTEOLOGY

Certain names in this Index have an asterisk (*) prefixed; these, as more fully explained in the Translator's Preface, being terms that form part of the English nomenclature used in this work, but which are not commonly employed by English anatomists. To other names a dagger (†) is prefixed; these are Latin names used by the author in the original work, but not included in the official nomenclature of the "Anatomische Gesellschaft."

A.

- ACETABULUM, 129
 Acromion, 108, 110, 111
 †Aditus ad antrum tympanicum, 66, 67
 orbitæ, 48, 98
 Agger nasi, 78, 90
 Air-cells of the Eustachian tube, 66
 Ala magna, 58-61
 ossis ilii, 130
 parvæ, 58-61
 Alæ vomeris, 79, 91, 92
 Alisphenoid, 61
 Alveolæ dentales mandibulæ (lower jaw-bone), 84
 maxillæ (upper jaw-bone), 82
 Ampulla ossea lateralis, 69
 posterior, 69
 superior, 69
 Ampulla, osseous, of the external semicircular canal, 69
 of the posterior semicircular canal, 69
 of the superior semicircular canal, 69
 Angle, acromial, 110
 of the jaw, 84, 85
 of Ludwig, 41
 of the parietal bone, frontal, 72, 73
 mastoid, 72, 73
 occipital, 72, 73
 sphenoidal, 72, 73
 of the pubis, 125
 of the ribs, 37, 40
 of the scapula, inferior, 110
 internal, 110
 superior, 110
 of the sphenoid bone, parietal, 58-60
 of the sternum, 41
 subcostal, 36
 Angulus anterior pyramidis, 65
 costæ, 37, 40
 frontalis (ossis parietalis), 72, 73
 inferior (scapulae), 110
 infrasternalis, 36
 lateralis (scapulae), 110
 Ludovici, 41
 mandibulæ, 84, 85
 mastoidæ (ossis parietalis), 72, 73
 medialis (scapulae), 110
 occipitalis (ossis parietalis), 72, 73
 parietalis (ossis sphenoidalis), 58-60
 posterior pyramidis, 65
 pubis, 125
 sphenoidalis (ossis parietalis), 72, 73
 sterni, 41
 superior pyramidis, 49, 65
 Annulus tympanicus, 70, 103, 104
 Antrum of Highmore, 80-82, 93-95, 97, 99
 orifice of the, 90
 mastoid, 64, 66, 67, 70
 entrance to, 66, 67
 tympanicum, 64, 66, 67, 70
 Apertura *vel* aperture:
 externa aqueductus vestibuli, 63, 68, 70
 canaliculi cochleæ, 62, 63
 mastoidæ, 60

- Apertura *vel* aperture:
 † inferior canaliculi tympanici, 62
 † interna canaliculi cochleæ, 69
 pelvis [minoris] inferior, 126, 127
 superior, 124, 125, 127
 piriformis, 46, 48, 90, 91, 93
 † sinus maxillaris, 90
 sphenoidalis, 58-61, 90-93
 † sinuum frontaliuum, 75, 76, 90, 95
 superior canaliculi tympanici, 66, 67
 thoracis inferior, 36
 superior, 36
 tympanica canaliculi chordæ, 64
 Apex capituli fibulæ, 138
 † ossis coccygis, 32
 sacri, 30, 31
 patellæ, 136
 pyramidis, 63, 64, 67, 70
 Apophysis articularis (articular apophysis), 43
 costalis (costal apophysis), 43
 muscularis (muscular apophysis), 43
 Aquæeductus vestibuli, 69
 Aqueduct of the cochlea, external orifice, 62, 63
 internal orifice, 69
 of Fallopius, 64-69
 deficiency in its tympanic wall, 61
 of the vestibule, 69
 external orifice 63, 68, 70
 Arch of the atlas, anterior, 29
 posterior, 29
 neural, or vertebral, 25, 26
 orbital, 48, 74, 76, 77, 92, 98
 pubic, 124
 of the ribs, 36
 zygomatic, 46-48, 96
 Arcus anterior atlantis, 29
 costarum, 36
 posterior atlantis, 29
 pubis, 124
 superciliaris, 46, 48, 74
 vertebræ, 25, 26
 zygomaticus, 46-48, 96
 *Area cochleæ, 68
 cribrosa media, 68
 superior, 68
 * of the facial nerve, 68
 nervi facialis, 68
 vestibularis inferior, 68
 superior, 68
 Arnold's nerve, causal for, 62, 64, 65
 Arteria nutritia, 20
 protovertebralis, 34
 Artery, intercostal, primitive, 34
 nutrient, 20
 protovertebral, 34
 Articulation, manubrio-gliadiolæ, 41
 Astragalus, 17, 142, 143
 Atlas, 29
 development of, 35
 Atrium meatus medi (atrium of the middle meatus), 90
 Auditory aperture, external, 47, 48, 62
 internal, 63, 65, 68, 70

- Axis of the pelvis, 127
- Axis, the, 29
 - development of, 35
- B.**
- Base of the mandible, 85, 87
 - of the patella, 136
 - of the sacrum, 32
 - of the skull, external aspect, 48
 - internal aspect, 49
- Bases of the metacarpal bones, 122
 - of the metatarsal bones, 145
 - of the phalanges of the fingers, 122
 - of the toes, 145
- Basilar portion of the occipital bone, 54-57, 60
- Basio-occipital portion of the occipital bone, 54-57, 60, and note, p. 57
- Basis cranii externa, 48
 - interna, 49
 - mandibulæ, 85, 87
 - ossis sacri, 32
 - ossium metacarpalium, 122
 - metatarsalium, 145
 - patellæ, 136
 - phalangis manus, 122
 - pedis, 143
- Basisphenoid, 58-61
- Bertin, bones of, 58, 59, 61
- Bodies of the metacarpal bones, 122
 - of the metatarsal bones, 145
 - of the phalanges of the fingers, 122
 - of the toes, 145
 - of the ribs, 40
 - of the vertebrae, 25-27
- Body
 - of the astragalus, 143
 - of the calcaneum, 141, 143
 - of the femur, 132, 133
 - of the fibula, 138
 - of the humerus, 112
 - of the hyoid bone, 87
 - of the ilium, 130
 - of the inferior maxillary bone, 84-87
 - of the ischium, 128, 130
 - of the malar bone, 83
 - of the mandible, 84-87
 - of the maxilla, 80, 81, 87, 89
 - of the os calcis, 141, 143
 - of the pubis, 130
 - of the radius, 115
 - of the sphenoid bone, 58-61
 - of the sternum, 41
 - of the superior maxillary bone, 80, 81, 87, 89
 - of the tibia, 136, 137
 - of the ulna, 114
- Bone, capitate, 119-121
 - central, of the carpus, 120
 - cuboid, 142, 144
 - cuneiform, of foot, external, 17, 142, 144
 - first, 141, 142, 144
 - internal, 141, 142, 144
 - middle, 142, 144
 - second, 142, 144
 - third, 17, 142, 144
 - of hand, 119, 120
- ethmoid, 52, 73
- frontal, 52, 74-76
 - development of, 77
- hip-, 107, 128-130
 - development of, 131
- hyoid, 87
- innominate, 107, 128-130
 - development of, 131
- interparietal, 100, note to p. 57
- lachrymal, 53, 79, 90-92, 95
- lunar, 119, 120
- malar, 53, 83
- marrow, 11
- maxillary, inferior, 46, 47, 53, 84-86
 - development of, 86
- Bone, maxillary, superior, 46, 47, 53, 80, 81
 - development of, 82
- metacarpal, first, 106, 119
- nasal, 53, 79, 90, 91
- navicular, 142, 144
- occipital, 52, 54-56
 - development of, 57, and note
- palate, 53, 83
- parietal, 19, 52, 72, 73
 - development of, 21
- pisiform, 118-120
- premaxillary, 82, 97
- pyramidal, 119, 120
- scaphoid, 119, 120
- sphenoid, 52, 58-60
 - development of, 61
- sphenoidal spongy, 58, 59, 61
 - turbinate, 58, 59, 61
- subcoracoid, 111
- temporal, 52, 62-69
 - development of, 70, 71
- trapezoid, 119-121
- turbinate of the nose, highest, 78
 - inferior, 53, 70, 90, 91, 93-95
 - middle, 78, 90, 94, 95
 - superior, 78, 90, 92, 94
- unciform, 119-121
- Bones, general considerations, 9-21
 - development of, 20, 21
 - flat, 19
 - long, 12-15
 - minute structure, 10
 - short, 16, 17, 21
- of Bertin, 58, 59, 61
- of the carpus, 120, 121
 - development of, 123
- of the cranium proper, 52
- of the face, 53
- of the forearm, 106, 114-116
 - development of, 117
- of the leg, 107, 136-138
 - development of, 139
- metacarpal, 119, 122
 - development of, 123
- metatarsal, 142, 145
 - development of, 146, 147
- sesamoid, of foot, 140, 141
 - of hand, 118
- of the skull, 52-104
- suprasternal, 41
- of the tarsus, 17, 143, 144
 - development of, 146, 147
- Wormian, 100
- Border of the fibula, anterior, 138
 - external, 138
 - internal, 138
 - interosseous, 138
- of the frontal bone, nasal, 76, 77
 - parietal, 74-76
 - sphenoidal, 75, 76
- of the humerus, inner, 112
 - outer, 112
- of the occipital bone, lambdoid, 54-26
 - mastoid, 54-56
- of the parietal bone, anterior, 72, 73
 - frontal, 72, 73
 - inferior, 72, 73
 - occipital, 72, 73
 - posterior, 72, 73
 - sagittal, 72, 73
 - squamous, 72, 73
 - superior, 72, 73
- of the petrous portion of temporal bone, anterior, 65
- of the petrous portion of the temporal bone, posterior, 65
- of the petrous portion of the temporal bone, superior, 49, 65
- of the radius, anterior, 114, 115
 - internal, 114, 115
 - posterior, 114, 115

- Border of the scapula, axillary, 110
 superior, 110
 vertebral, 110
 of the sphenoid, external, 58-60
 frontal, 58, 59
 malar, 58-60
 orbital, 59
 petrous, 59
 posterior, 59
 squamous, 58-60
 of the superior maxillary bone, infra-orbital, 80, 83,
 98
 lacrimal, 80
 of the temporal bone, occipital, 62, 63
 parietal, 62, 63
 sphenoidal, 62-64
 of the tibia, anterior, 136, 138
 external, 136-138
 internal, 136, 138
 of the ulna, anterior, 114
 external, 114
 posterior, 114
 Breschet's canals, 19, 51, 88
 Brim of the pelvis, 124, 125
 iliac portion, 125
 pubic portion, 125
 sacral portion, 30, 125
 Bulla, ethmoidal, 78, 90

C.

- Calcaneum, or os calcis (called by Toldt "Calcaneus"), 17,
 142, 143
 development of, 146, 147
 Calcar femorale, 134, 135
 Calvaria, 50
 Canal or canals (see also "Canaliculus"):
 for Arnold's nerve, external orifice of, 66
 for the auricular branch of the pneumogastric
 nerve, 62, 64, 65
 basipharyngeal (see note to p. 48), 48, 58-60
 of Breschet, 19, 51, 88
 carotid, 62-65, 67, 70, 71, 76
 for the chorda tympani nerve, 64, 65, 67
 dental, 81
 anterior and middle, 81
 inferior, 86
 posterior, 80, 81, 99
 diploic, 19, 51, 88
 ethmoidal, anterior, 76, 90, 91, 98
 posterior, 76, 91, 98
 of the Eustachian tube, 63, 64, 66, 67, 69
 of the facial nerve, 64, 69
 Haversian, 10
 infra-orbital, 80, 81, 95
 internal orbital, anterior, 76, 90, 91, 98
 posterior, 76, 91, 98
 for Jacobson's nerve, 65-67
 inferior orifice, 62
 superior orifice, 66, 67
 malar, 83
 mandibular, 86
 medullary, 11, 18-20
 nutrient, 13, 18, 19
 palatine, accessory, 83, 90, 91, 97
 posterior, 97, 99
 inferior orifice of, 90, 96, 97
 palato-maxillary, 97, 99
 inferior orifice of, 90, 96, 97
 pterygoid (or Vidian), 58, 59, 61, 92, 93, 96, 99
 pterygopalatine, 48, 58-60, 90-92
 sacral, 31, 32
 semicircular, external, 67-69
 posterior, 68, 69
 superior, 68, 69
 spinal, 43
 temporal, 83
 of the tensor tympani muscle, 63, 64, 66, 67, 69
 Vidian (or pterygoid), 58, 59, 61, 92, 93, 96, 99
 Volkmann's, 10, 11

- Canaliculus *vel* canaliculi (see also "Canal"):
 caroticotympanici, 62, 67
 chordæ tympani, 64, 65, 67
 cochleæ (apertura externa), 62, 63
 (apertura interna), 69
 † innominatus, 93
 † mastoideus, 62, 64, 65
 † sphenoidalis, 92, 93, 96, 99
 tympanicus, 65
 Canalis *vel* canales:
 alveolares, 81
 basipharyngeus, 48, 58, 59, 90
 caroticus, 62-65, 67, 70, 71, 96
 condyloideus, 54-57, 88, 89
 diploici [Breschet], 19, 51, 88
 facialis [Fallopil], 64, 65, 67-69
 hypoglossi, 54-57, 88
 incisivus, 80, 91
 infra-orbitalis, 80, 81, 95
 mandibulæ, 86
 musculotubarius, 63, 64
 nasolacrimalis, 93, 95
 nutricius, 13, 18, 19
 palatini, 90, 97
 pharyngeus, 48, 58, 59, 90-92
 pterygoideus [Vidian], 58, 59, 61, 92, 93, 96, 99
 pterygopalatinus, 97, 99
 sacralis, 31, 32
 semicircularis lateralis, 67-69
 posterior, 68, 69
 superior, 68, 69
 vertebralis, 43
 Capitellum of the humerus, 112
 Capitulum costæ, 40
 fibulæ, 138
 humeri, 112
 mandibulæ, 84-86, 96
 ossium metacarpalium, 122
 metatarsalium, 141, 145
 radii, 115
 ulnæ, 114
 Caput femoris, 132-134
 humeri, 112
 † ossis capitati, 121
 tali, 143
 Carpus, 106, 118
 Cartilage, Meckel's, 103
 Cartilages, costal, 39, 40
 Cartilago costalis, 39, 40
 Cavitas glenoidalis, 110, 111
 Cavity, cranial, 88, 89
 medullary, 11, 18-20
 nasal, 90-95
 sigmoid (of the radius), 115
 (of the ulna), great, 114, 115
 small, 114
 thoracic, 43
 tympanic, 67
 † Cavum cranii cerebialis, 88, 89
 medullare, 11, 18-20
 nasi, 90-95
 thoracis, 43
 tympani, 65-70, 96
 Cells, ethmoidal, 76, 78, 90
 of the Eustachian tube, 66
 mastoid, 64, 67
 tympanic, 67
 Cellulæ ethmoidales, 78, 90
 mastoideæ, 64, 67
 pneumaticæ tubariæ, 66
 tympanicæ, 67
 Centra of the vertebrae, 25-27
 Choanæ, 48, 93, 96
 Chorda dorsalis, 34
 tympani nerve, canal for, 64, 65, 67
 orifice of the canal for, 64
 Ciugulum extremitatis inferioris, 107, 124-126
 superioris, 106, 108
 Circumference, articular, of the radius, 115
 of the ulna, 114

- Circumferentia articularis radii, 115
 ulnæ, 114
 Clavicula (the clavicle), 106, 108, 109
 Clivus, 49, 56, 88
 Coccyx, the, 24, 32, 33, 125, 126
 development of, 35
 extremity of the, 32
 Cochlea, 65, 68, 69
 Collum anatomicum, 112
 chirurgicum, 112
 costæ, 40
 femoris, 132-134
 mandibulæ, 84, 85
 radii, 115
 scapule, 110
 tali, 143
 Column, spinal, 24
 development of, 34, 35
 vertebral, 24
 development of, 34, 35
 Concavity of the carpus, 118
 Concha nasalis inferior, 55, 79, 90, 91, 93-95
 media, 78, 90, 94, 95
 superior, 78, 90, 92, 94
 suprema [Santorini], 78
 Conchæ sphenoidales, 58, 59, 61
 Condylar portion of the occipital bone, 54, 56, 57, and note,
 p. 57
 Coudyle of the femur, external, 132, 134
 internal, 132-134
 of the humerus, external, 112
 internal, 112
 of the inferior maxillary bone, 84-86, 96
 of the mandible, 84-86, 96
 Condyles of the occipital bone, 48, 54-56
 Condylus lateralis femoris, 132, 134
 tibias, 136, 137
 medialis femoris, 132-134
 tibias, 136, 137
 occipitalis, 48, 54-56
 +Conjugata diagonalis, 127
 vera, 127
 Conjugate diameter, diagonal, 127
 true, 127
 Cornu, coccygeal, 32
 of the hyoid bone, great, 87
 small, 87
 sacral, 31, 32
 Cornua sphenoidalia, 58, 59, 61
 Corpus calcanei, 141, 143
 costæ, 40
 femoris, 132, 133
 fibulæ, 138
 humeri, 112
 mandibulæ, 84-87
 maxillæ, 80, 81, 87, 99
 ossis hyoidei, 87
 ilium, 130
 ischii, 128, 130
 pubis, 130
 sphenoidalis, 58-61
 ossium metacarpalium, 122
 metatarsalium, 145
 phalangis manus, 122
 pedis, 145
 radii, 115
 sterni, 41
 tali, 143
 tibias, 136, 137
 ulnæ, 114
 vertebræ, 25-27
 Costæ, 38, 40
 development of, 42
 spuriæ, 36
 veræ, 36
 Craniometry, 101
 Cranium, 45-104
 cavity of, 88, 89
 development of, 102-104
 facial portion (cranium viscerale), 53, 96, 97
 Cranium, primordial (cranium primordiale), 102, 103
 proper (cranium cerebrale), 52, 88, 89
 * Crest, buccinator, 84, 85
 ethmoidal, of the palate-bone, 83
 of the superior maxillary bone, 80, 81
 frontal, 49, 50, 75, 88
 of the fundus of the internal auditory meatus, trans-
 verse, 65, 68
 of the ilium, 128-130
 inferior turbinate, of the superior maxillary bone,
 80, 81
 infratemporal, 48, 58-60, and note, p. 59
 lachrymal, 79, 98
 * anterior, see "Ridge, orbital"
 * posterior, see "Crest, lachrymal"
 nasal, of the superior maxillary bone, 91, 95, 97
 obturators, 130
 occipital, external, 48, 55, 56
 internal, 49, 54, 56
 of the occipital bone for the rectus capitis posticus
 major muscle, 56
 of the occipital bone for the rectus capitis posticus
 minor muscle, 56
 * orbital, 59
 * of the septum of the nose, lateral, 93
 * sacral, articular, 31
 * external, 31
 * median, 31
 sphenoidal, 58, 59
 supra-nastoid, 62
 temporal, 46, 74, 77
 transverse, see "Crista falciformis"
 turbinate, inferior, of the palate-bone, 83
 of the superior maxillary bone,
 80, 81
 superior, of the palate-bone, 83
 of the vestibule, 69
 Crista anterior (fibulæ), 138
 (tibias), 136, 138
 buccinatoria, 84, 85
 capituli costæ, 40
 colli costæ, 40
 couchalis (maxillæ), 80, 81
 (ossis palatini), 83
 ethmoidalis (maxillæ), 80, 81
 (ossis palatini), 83
 falciformis, 65, 68
 frontalis, 49, 50, 75, 88
 galli, 49, 78, 88, 94, 95
 iliaca, 128-130
 infratemporalis, 48, 58-60
 interossea (fibulæ), 138
 (radii), 114, 115
 (tibias), 136-138
 (ulnæ), 114
 intertrochanterica, 132-134
 lacrimalis anterior, 80, 98
 posterior, 79, 98
 lateralis (fibulæ), 138
 septi (nasi), 93
 medialis (fibulæ), 138
 musculi recti capitis majoris, 56
 minoris, 56
 supinatoris, 114
 nasalis, 91, 95, 97
 obturatoria, 130
 occipitalis externa, 48, 55, 56
 interna, 49, 54, 56
 orbitalis (alæ magnæ), 59
 sacralis articularis, 31
 lateralis, 31
 medialis, 31
 sphenoidalis, 58, 59
 transversa, 65, 68
 tuberculi majoris, 112
 minoris, 112
 vestibuli, 69
 Crus commune and crus simplex of the semicircular canals,
 69, and note, p. 69
 Cupular portion of the epitympanic recess, 64

D.

- Dens epistrophei, 29
 Depression, digital, 85
 Depressions for the lateral metatarso-phalangeal ligaments, 145
 Pacchionian, 50, 73
 Diameter, conjugate, of the pelvis, diagonal, 127
 true, 127
 Diameters, oblique, of the pelvis, 127
 transverse, of the pelvis, 127
 Diaphysis, 20
 Digi (manus), 118
 pedis, 140, 141
 Diploe, 19, 50, 51
 of the frontal bone, 76, 77
 Disc, epiphysial, 20
 Dorsum of the foot, 140
 of the scapula, 110
 sellæ, 58-61, 96

E.

- Emmence, articular, 62
 deltoid, 112
 frontal, 46, 74, 77, 104
 iliopectineal, 128, 130
 jugular, 54, 56
 olivary, 59-61
 parietal, 72, 104
 papillary, 66, 67, 69
 of the second rib for the serratus magnus, 40
 of the superior semicircular canal, 63, 70
 Eminentia arcuata, 63, 70
 carpi radialis, 118
 ulnaris, 118
 cruciata, 54, 89
 iliopectinea, 128, 130
 intercondyloidea, 136, 137
 papillaris, 66, 67, 69
 pyramidalis, 66, 67, 69
 Entrance to the mastoid antrum, 66, 67
 to the orbit, 48, 98
 Epicondyle of the femur, external, 132, 134
 internal, 132, 133
 Epicondylus lateralis (femoris), 132, 134
 (humeri), 112
 medialis (femoris), 132, 135
 (humeri), 112
 Epiphysis, 20
 Epistropheus, 29
 development of, 35
 Epitympanic recess, 63, 69, 70
 Eustachian canal, 63, 64, 66, 67, 69
 Extremitas acromialis (claviculæ), 109
 sternalis (claviculæ), 109
 Extremity of the clavicle, acromial, 109
 sternal, 109
 Exoccipital, 54, 56, 57, and note, p. 57

F.

- Facet or facets, articular:
 of the acromion, clavicular, 110
 of the atlas, superior, 29
 of the astragalus—
 for the calcaneum, anterior, 143
 middle, 143
 posterior, 143
 for the external malleolus, 143
 for the internal malleolus, 143
 for the navicular bone, 143
 of the clavicle, acromial, 109
 sternal, 109
 of the heads of the ribs, 40
 of the odontoid process, anterior, 29
 of the os calcis, anterior, 143
 cuboid, 143
 internal, 143
 posterior, 143
 for the ribs (sternal), 41
 of the tubercles of the ribs, 40

Facet or facets, articular:

- of the vertebrae, inferior, 25-27
 superior, 25-27

† Facial portion of the skull, 53

- † Facies, anterior femoris, 132, 133
 lateralis humeri, 112
 maxillæ, 80
 medialis humeri, 112
 pyramidis, 65
 articularis acromialis (claviculæ), 109
 acromii, 110
 anterior calcanei, 143
 epistrophei, 29
 calcanea (tali) anterior, 143
 media, 143
 posterior, 143
 capituli costæ, 40
 fibulæ, 138
 carpea, 115
 cuboidea (calcanei), 143
 fibularis, 136, 137
 inferior tibiæ, 137
 vertebrarum, 25-27
 malleolaris (tibiæ), 137
 malleoli (lateralis), 137, 138
 media (calcanei), 143
 navicularis (tali), 143
 ossis temporalis, 62, 70, 71
 patellæ, 136
 posterior (calcanei), 143
 sternalis (claviculæ), 109
 superior tibiæ, 137
 vertebrarum, 25-27
 tuberculi costæ, 40
 auricularis ossis ilium, 128
 sacri, 31, 32
 cerebralis alæ magnæ, 58, 61
 ossis frontalis, 75
 parietalis, 73
 squammæ temporalis, 63, 64
 costalis scapulae, 110
 dorsalis manus, 119
 + ossis sacri, 31, 32, 126
 + pedis, 140
 + radii, 114-116
 scapulae, 110
 ulnæ, 114
 frontalis (ossis frontalis), 74
 inferior pyramidis, 65
 infratemporalis (maxillæ), 80, 99
 + lateralis femoris, 132, 133
 fibulæ, 138
 radii, 114-116
 tibiæ, 136-138
 lunata (acetabuli), 130
 malaris (ossis zygomatici), 83
 malleolaris lateralis (tali), 143
 medialis (tali), 143
 maxillaris ossis palatini, 83
 + medialis femoris, 132, 133
 fibulæ, 138
 tibiæ, 136, 138
 ulnæ, 114
 nasalis maxillæ, 80
 ossis palatini, 83
 orbitalis alæ magnæ, 58-61, 98
 maxillæ, 80
 ossis frontalis, 76
 zygomatici, 83
 osæa, 53
 parietalis (ossis parietalis), 72
 patellaris, 132-134
 pelvina (ossis sacri), 30, 32, 125
 + plantaris pedis, 141
 posterior fibulæ, 138
 humeri, 112
 pyramidis, 65
 tibiæ, 136-138
 sphenomaxillaris (alæ magnæ), 58, 99
 superior (tali), 143

- Facies symphyseos, 128
temporalis alae magnae, 58-61
 ossis frontalis, 74, 77
 zygomati, 83
 squamae temporalis, 62, 63
- † volaris manus, 118
 radii, 114-116
 ulnae, 114, 116
- Femur, the, 14, 107, 132-134
development of, 135
- Fenestra cochleae, 67, 69
ovalis, 67-69
rotunda, 67, 69
vestibuli, 67-69
- Fibres, Sharpey's, 10
- Fibula, the, 107, 138
development of, 139
- Fingers, skeleton of, 118
- Fissura orbitalis inferior, 93, 96, 98
superior, 58, 59, 88, 92, 95
 petro-occipitalis, 48, 49, 88
 petrosquamosa, 63, 64, 69
 petrotympanica [Glaseri], 62, 65, 70, 71
 pterygoidea, 58, 59
 pterygomaxillaris, 99
 sphenio-occipitalis, 60
 sphenopetrosa, 48, 49
 tympanomastoidea, 62, 66, 71
- Fissure, Glaserian, 62, 65, 70, 71
incisor, 48, 82, 97
occipitosphenoidal, 60
orbital, 58, 59, 88, 92, 98
petrobassilar, 48, 49, 98
petrosquamosa, 63, 64, 69
pterygomaxillary, 99
sphenomaxillary, 93, 96, 98
tympanomastoid, 62, 66, 71
- Fontanelle, anterior, 104
anterolateral, 104
posterior, 104
posterolateral, 104
- Fonticulus frontalis [major], 104
mastoides, 104
occipitalis [minor], 104
sphenoidalis, 104
- Foramen or foramina:
alveolar, anterior and middle, 84
 posterior, 80, 81, 99
caecum, 49, 75, 88, 91, 94
condylar, anterior, 54-57, 88
 posterior, 54-57, 88, 89
costotransverse, 26, 29
emissarii occipitalis, 89
emissary, mastoid, 62, 63, 89
 occipital, 89
 of Vesalius, see "Canaliculus sphenoidalis," and note, p. 92
ethmoidale anterius, 76, 90, 91, 98
 posterius, 76, 91, 98
frontale, 76
incisivum, 82, 93, 96, 97
inferior dental, 85
infra-orbital, 80, 81, 98, 99
intervertebral, 24
 of the sacrum, 31
jugular, 48, 49, 88
 bipartite, 88
 divided by an intrajugular process, 88
lacerate, anterior, 58, 59, 88, 92, 98
 middle, 48, 49
 posterior, 48, 49, 88
mandibular, 85
magnum, 48, 54-56, 89
mastoid, 62, 63, 89
mental, 85, 86
nasal, 79
nutricium, 18, 19
 claviculae, 109
 femoris, 132
 fibulae, 138
- Foramen or foramina:
nutricium humeri, 112
 ossis ilii, 19
 radii, 115, 116
 tibiae, 18, 136, 137
 ulnae, 114, 116
- nutrient, 18, 19
of the clavicle, 109
of the femur, 132
of the fibula, 138
of the humerus, 112
of the ilium, 19
of the radius, 115, 116
of the tibia, 18, 136, 137
of the ulna, 114, 116
- obturator, 128-130
optic, 59-61, 88, 98, 99
ovale (of the hip-bone), 128-130
 (of the skull), 59, 61, 96
- palatina minora, 83, 90, 97
palatinum majus, 90, 96, 97
- * palatine, great, 90, 96, 97
parietal, 72, 73, 100
pterygospinosus, 91
rotundum, 38-61, 88, 92, 93, 99
- sacral, anterior, 30
posterior, 31
- singulare, 68
sphenoidal, 58-61, 90-93
sphenopalatine, 90, 91, 98, 99
spinal, 25-27
spinous, 59
of Stenson, 82
stylomastoid, 62, 64, 67, 70, 71
supra-orbital, 74, 76
thyroid, 128-130
transversarium, 26, 29
vertebrale, 25-27
for vertebral artery, 26, 29
of Vesalius, note to p. 92 (see also "Canaliculus sphenoidal")
zygomatofaciale, 83
zygomatoco-orbitale, 83
zygomatocotemporale, 130
- Fossa acetabuli, 130
canine, 80
condylar, posterior, 55
coronoid, 112
cranial, anterior, 49, 88, 95
 middle, 49, 88
 posterior, 49, 88
digastric (of the inferior maxillary bone), 85
 (of the temporal bone), 62, 71
digital, 132, 133
of the external lateral ligament of the ankle-joint, 137, 138
floccular, 63, 70
glandulae lachrymalis, 76, 77, 92, 98
of the Gasserian ganglion, 63
glenoid, of the scapula, 110, 111
 of the temporal bone, 62, 70, 71
hypophyseos, 59-61, and note to p. 60
iliac, 128
of the incus, 61
infraspinous, 110
infratemporal, 48, 96
intercondylar, of the femur, 132, 134
 of the tibia, anterior, 137
intercondyloidea (femoris), 132, 134
 anterior (tibiae), 137
 posterior (tibiae), 137
jugular, of the temporal bone, 62, 65
lachrymal, 76, 77, 92, 98
mandibularis, 62
occipital, inferior, 54, 56
 superior, 54
olecranon, 112
palatine, anterior, 82, 96, 97
pituitary, 59-51
prenasal, 97

- Fossa, pterygoid, 58, 95
 pterygopalatine, 92, 98, 99
 radial, 112
 reniform, 65
 sacci lacrimalis, 98
 scaphoid, 58
 sphenomaxillary, 92, 98, 99
 subarcuata, 63, 70
 subscapular, 110
 supraspinous, 110
 temporal, 47
 trochanterica, 132, 133
 trochlear, 76
 vermic, 89
 zygomatic, 48, 96
- * Fossula fenestra cochleæ, 66, 68
 vestibuli, 66
 of the fenestra ovalis, 66
 rotunda, 66, 68
 ovalis, 66
 of the petrous ganglion, 62, 65
 petrosa, 62, 65
 rotunda, 66, 68
- Fovea articularis ossis temporalis, 62
 superior atlantis, 29
 capitis femoris, 132, 133
 capituli radii, 115
 costalis, inferior, 25
 superior, 25
 transversalis, 25
 dentis, 29
 pterygoidea (processus condyloidei), 84, 85
 sublingualis, 85
 submaxillaris, 85
 trochlearis, 76
- † Foveolæ ethmoidales, 76
 granulares [Pacchioni], 59, 73
- Frons, 46, 47
- Fundus meatus acustici interni, 65
 of the internal auditory meatus, 65
- G.**
- Geniculum of the aqueduct of Fallopius, 65, 67, 68
 canalis facialis, 65, 67, 68
 "German horizontal," 101
- Girdle, pelvic, 107, 124-126
 shoulder, 106, 108
- Glabella, 46, 74
- Gladiolus, 41
- Great wing of the sphenoid, 58-61
- Groove or grooves:
 for Arnold's nerve, 62, 65
 for the auricular branch of the pneumogastric nerve, 62, 65
 basilar, 49, 56, 58
 bicipital, 112
 carotid, 58, 59
 fibular, 136, 137
 of the hamular process, 58
 infra-orbital, 80, 93, 94, 98, 99
 interosseous, of the astragalus, 143
 of the calcaneum, 143
 of the tarsus, 143
- lachrymal, 98
 of the lachrymal bone, 79, 92
 of the superior maxillary bone, 80, 82, 91, 98
- meningeal, 50, 73
 of the middle temporal artery, 62
 mylohyoid, 85
 of the nasal nerve, 79, 91
 obturator, 128
 occipital, 62, 71
 optic, 59
 palatine, 82, 97
 anterior, 80, 91
 posterior, 80
 of the palate-bone, 83
 of the sphenoid bone, 58, and
 note to same

- Groove or grooves:
 palatomaxillary, 80
 of the palate-bone, 83
 of the sphenoid bone, 58, and
 note to same
- preauricular, 128
 plexus, 67
 of the promontory, for the nerves of the tympanic
- * pterygopalatine, note to p. 58
 spiral, 112
 for spinal nerve, 26
 of the subclavian artery, 40
 subcostal, 40
 of the superficial petrosal nerve, great, 63, 64
 small, 63, 64
- of the tendon of—
 the flexor carpi radialis muscle, 121
 the flexor longus hallucis muscle, 141, 143
 the peroneus longus muscle, 140, 141, 143, 144
 the popliteus muscle, 134
 the tibialis posticus muscle, 136
 of the tendons of the peroneal muscles, 138
 of the ulnar nerve, 112
 vertebral (for vertebral artery), 29

H.

- Hamulus lacrimalis, 79, 98
 ossis hamati, 118, 121
 pterygoideus, 58, 60, 95
- Haversian canals, 10
 system of lamellæ, 10, 11
- Head of the astragalus, 143
 of the capitate bone, 121
 of the femur, 132-134
 of the fibula, 138
 of the humerus, 112
 of the inferior maxillary bone, 84-86, 96
 of the mandible, 84-86, 96
 of the os magnum, 121
 of the radius, 115
 of the scapula, 110
 of the ulna, 114
- Heads of the metacarpal bones, 112
 of the metatarsal bones, 141, 145
 of the ribs, 40
- Heart, rudimentary, 102
- Hiatus canalis facialis, 63-67
 Fallopii, 63-67
 maxillaris, 80, 91, 99
 of the sacrum, 31
 semilunaris, 78, 90
 subarcuatus, 63, 70
- Hip-bone, the, 107, 128-130
 development of, 131
- Hook of the unciform bone, 118, 121
- Horseshoe-shaped articular surface of the acetabulum, 130
- Humerus, the, 12, 106, 112
 development of, 113

I.

- Iliac portion of the iliopectineal line, 128
- Ilium, the, 19, 124-126
- * Impressiones digitate, 49, 75, and notes, pp. 49 and 75
- Impression, rhomboid, 109
- Impressio trigemini, 63
- Incisura vel incisura:
 acetabuli, 130
 clavicularis, 41
 costales, 41
 ethmoidalis, 76, 77
 fibularis, 136, 137
 frontalis, 74, 76
 ischiadica major, 128, 129
 minor, 129
 jugularis ossis occipitalis, 56, 89
 temporalis, 63
 sterni, 41
- lacrimalis, 80

Incisura *vel* *incisurae* :

- mandibulae, 83
 - mastoidea, 62, 71
 - nasalis, 80
 - parietalis, 62, 63
 - radialis (ulnae), 114
 - scapulae, 110
 - semilunaris, 114, 115
 - sphenopalatina, 83
 - supra-orbitalis, 74, 76
 - tympanica [Rivini], 62, 64, 70, 71
 - ulnaris (radii), 115
 - vertebralis inferior, 25, 27
 - superior, 27
- Inclination of the pelvis, 127
- Infundibulum, 78
- Interparietal bone, note, p. 57
- Ischium, the, 124-126
- Iter chordae posteriorius, 64, 65, 67

J.

- Jacobson's nerve, caual for, 65-67
- Jaw-bone, lower, 46, 47, 53, 84-86
 - development of, 86
 - upper, 46, 47, 53, 80, 81
- Jaws, senile atrophy of, 87
- Juga alveolaria, 80, 85
 - cerebralia, 49, 75

L.

- Labium externum (cristae iliaceae), 129, 130
 - internum (cristae iliaceae), 128
 - laterale (lineae asperae), 132, 133
 - mediale (lineae asperae), 132, 133
- Labyrinth, bony, 68, 69, 96
 - of the ethmoid bone, 78, 92, 94
 - osseous, 68, 69, 96
- Labyrinthus ethmoidalis, 78, 92, 94
 - osseous, 68, 69, 96
- Lachrymal canal, 93, 95
- Lacunae of bone, 10
- Lamina cribrosa, 78, 90, 91, 94
 - externa of the cranial bones, 19, 50
 - interna of the cranial bones, 19, 50
 - lateralis processus pterygoidei, 58, 60, 61, 99
 - malaris (ossis zygomatici), 83
 - medialis processus pterygoidei, 58, 61
 - orbitalis (ossis zygomatici), 83, 98
 - papyracea, 78, 91, 92, 94
 - perpendicularis, 78, 91, 94, 95
 - of the vertebrae, 25, 26
- Limbus alveolaris mandibulae, 84, 86
 - maxillae, 80-82, 97
- dentalis, 48
- Linca *vel* *lineae* :
 - arcuata (ossis ilii), 128
 - aspera (femoris), 132, 133
 - glutaea anterior, 129, 130
 - inferior, 129
 - posterior, 129
 - intercondyloidea (femoris), 132
 - intermedia (cristae iliaceae), 128, 130
 - intertrochanterica, 132, 133
 - musculares (scapulae), 110
 - mylohyoidea, 85
 - nuchae inferior, 48, 55, 56
 - superior, 55, 56
 - suprema, 55
 - obliqua (mandibulae), 84, 85
 - pectinea (femoris), 132, 133
 - poplitea, 136, 137
 - temporalis inferior, 47, 62, 72
 - (ossis frontalis), 46, 74, 77
 - superior, 47, 72
 - terminalis (pelvis), 124, 125
 - pubica, 125
 - sacralis, 30, 125
 - transverse (fossis sacri), 30

Line or lines :

- curved, of the ilium, inferior, 129
 - middle, 129, 130
 - superior, 129
- of the occipital bone, highest, 55
 - inferior, 48, 55, 56
 - superior, 55, 56
- gluteal, inferior, 129
 - middle, 129, 130
 - posterior, 129
- iliopectineal, iliac portion, 128
 - pubic portion, 128
- intercondylar, 132
- intertrochanteric, anterior, 132, 133
 - posterior, 132, 133
- oblique, external (of the inferior maxillary bone), 84, 85
 - internal (of the inferior maxillary bone), 85
- of the scapula, 110
- of the tibia, 136, 137
- pectineal, 132, 133
- temporal, inferior, 47, 72
 - superior, 47, 72
- trapezoid, 109
- Lingula mandibulae, 84-86
 - sphenoidalis, 59, 99
- Lip of the crest of the ilium, inner, 128
 - outer, 129, 130
- of the linea aspera, inner, 132, 133
 - outer, 132, 133

M.

- Malleolus, external, 137, 138
 - internal, 136, 137
 - lateralis, 137, 138
 - medialis, 136, 137
- Mandibula, 46, 47, 53, 84-86
 - development, 86
- Mauubrium sterni, 47
- Margin, alveolar, of the inferior maxillary bone, 84, 86
 - of the superior maxillary bone, 80-82, 97
 - of bicipital groove, inner, 112
 - outer, 112
 - supra-orbital, 48, 74, 76, 77, 92, 98
- Margo axillaris (scapulae), 110
 - dorsalis radii, 114, 115
 - ulnae, 114
- frontalis ossis parietalis, 72, 73
 - sphenoidalis, 58, 59
- infraglenoidalis (tibiae), 136, 137
- infra-orbitalis, 80, 83, 98
- lacrimalis (maxillae), 80
- lambdoidens, 54-56
- lateralis humeri, 112
 - pedis, 140
- mastoideus, 54-56
- medialis humeri, 112
 - pedis, 140
 - tibiae, 136, 138
- nasalis (ossis frontalis), 76, 77
- occipitalis ossis parietalis, 72, 73
 - temporalis, 62, 63
- orbitalis (ossis sphenoidalis), 59
- parietalis ossis frontalis, 74-76
 - temporalis, 62, 63
- petrosus (ossis sphenoidalis), 59
- sagittalis (ossis parietalis), 72, 73
- sphenoidalis ossis frontalis, 75, 76
 - temporalis, 62-64
- squamosus ossis parietalis, 72, 73
 - sphenoidalis, 58-60
- superior (scapulae), 110
- supra-orbitalis, 48, 74, 76, 77, 92, 98
- vertebralis (scapulae), 110
- volaris radii, 114, 115
 - ulnae, 114
- zygomaticus (ossis sphenoidalis), 58-60
- Massa lateralis (atlantis), 29
- Masces, lateral (of the atlas), 29

Mass, lateral, of the ethmoid bone, 78, 92, 94
 of the sacrum, 30, 31
 Mastoid antrum, 64, 66, 67, 70
 entrance to, 66, 67
 portion of the temporal bone, 62-64, 70, 71
 Maxilla, the, 46, 47, 53, 80, 81
 development of, 82
 Meatus acusticus externus, 47, 66, 69, 71, 96
 internus, 65, 68, 69, 96
 auditory, external, 47, 66, 69, 71, 96
 internal, 65, 68, 69, 96
 nasal, common, 78, 96
 inferior, 90, 94
 middle, 90, 94
 superior, 90, 94
 nasi communis, 78, 96
 inferior, 90, 94
 medius, 90, 94
 superior, 90, 94
 nasopharyngeal, 90
 nasopharyngeus, 90
 Medulla ossium, 11
 Membrane of the anterior fontanelle, 102
 Mesosternum, 41
 Metacarpus, 106, 118
 Metatarsus, 107, 140
 Metasternum, 41

N.

Nares, posterior, 48, 93, 96
 Nasal aperture, anterior, 46, 48, 90, 91, 93
 posterior, 48, 93, 96
 Neck, anatomical, of the humerus, 112
 of the astragalus, 143
 of the femur, 132-134
 of the inferior maxillary bone, 84, 85
 of the mandible, 84, 85
 of the radius, 115
 of the scapula, 110
 surgical, of the humerus, 112
 Necks of the ribs, 40
 † Norma frontalis, 46, 101
 † lateralis, 47, 101
 † verticalis, 101
 Notch, cotyloid, 130
 clavicular, 41
 ethmoidal, 76, 77
 great sciatic, 128, 129
 iliosciatic, 128, 129
 interclavicular, 40
 jugular, of the occipital bone, 56, 89
 of the temporal bone, 63
 lachrymal, 80
 nasal, 80
 parietal, 62, 63
 popliteal, 137
 pterygoid, 58, 89
 of Rivinus, 62, 64, 70, 71
 sciatic, 129
 sigmoid, 84
 small sciatic, 129
 sphenopalatine, 83
 supra-orbital, 74, 76
 suprascapular, 110
 vertebral, inferior, 25, 27
 superior, 27

Notochord, 34
 "Nutcracker face," 87
 Nutrient artery, 20

O.

Occiput, 47
 Odontoid process, 29
 Olecranon, 114, 115
 Optic vesicle, primary, 102
 Orbit, entrance to the, 48, 98
 Orbital orifice, height, 101
 width, 101
 Orbitosphenoid, 61
 Orbits, the, 92-95, 98
 Orifice of the Eustachian tube, tympanic, 66

Os *vel* ossa:
 † acetabuli, 131
 † antibrachii, 106, 114-116
 development of, 127
 brevia, 16, 17
 development of, 21
 capitatum, 119-121
 calcis, 17, 142, 143
 development of, 146, 147
 carpi, 120, 121
 development of, 123
 centrale carpi, 120
 coccygis, 24, 32, 33, 125, 126
 development of, 35
 costale, 38, 39
 coxæ, 107, 128-130
 development of, 13
 cranii, 52-104
 cerebri, 52
 † cruris, 107, 136-138
 development of, 139
 cuboideum, 142, 144
 cuneiforme primum, 141, 142, 146
 secundum, 142, 144
 tertium, 17, 142, 144
 ethmoidale, 52, 78
 faciei, 53
 frontale, 52, 74-76
 development of, 77
 hamatum, 119-121
 hyoideum, 87
 ilium, 19, 124-126
 † ince, 100
 † incisivum, 82, 97
 † infracoracoideum, 111
 innominatum, 107, 128-130
 development of, 131
 interparietale, 100
 ischii, 124-126
 lacrimale, 53, 79, 90-92, 95
 longa, 12-15
 lunatum, 119, 120
 magnum, 119-121
 metacarpale I., 106, 119
 metacarpalia, 119, 122
 development of, 123
 metatarsale I., 107, 142
 metatarsalia, 142, 145
 development of, 146, 147
 multangulum majus, 119-121
 minus, 119-121
 nasale, 53, 79, 90, 91
 naviculare manus, 119, 120
 pedis, 142, 144
 occipitale, 52, 54-56
 development of, 57
 palatinum, 53, 83
 parietale, 19, 52, 72, 73
 development of, 21
 pisiforme, 118-120
 planum, 78, 91, 92, 94
 plana, 19
 pubis, 124-126
 sacrum, 24, 30-33, 125, 126
 development of, 35
 sesamoides manus, 118
 pedis, 140, 141
 sphenoidale, 52, 58-60
 development of, 51
 suprasternalia, 41
 suturum, 100
 tarsi, 17, 143, 144
 temporale, 52, 62-69
 development of, 70, 71
 trigonum, 143
 triquetrum, 119, 120
 zygomaticum, 53, 83
 "Osseous corpuscles," 10
 Ossification, intracartilaginous, 20
 intramembranous, 21

Ossification of the Y-shaped cartilage of the acetabulum, 131
 Osteology, 7 *et seq.*
 Ostium tympanicum tubæ auditivæ, 66

P.

Palate, hard, 48, 94, 96, 97
 Palatum durum, 48, 94, 96, 97
 Paries carotica cavi tympani, 65, 66
 inferior orbitæ, 93, 98, 99
 jugularis cavi tympani, 66
 labyrinthica cavi tympani, 65, 66
 lateralis orbitæ, 98
 mastoidea cavi tympani, 66
 medialis orbitæ, 98
 superior orbitæ, 92, 98
 tegimentalis cavi tympani, 65
 Pars alveolaris mandibulæ, 84-86
 basilaris ossis occipitalis, 54-57
 cupularis recessus epitympanici, 64
 horizontalis ossis palatini, 83, 90, 91, 97
 lateralis ossis occipitalis, 53, 56, 57
 sacri, 30, 31
 mastoidea ossis temporalis, 62-64, 70, 71
 nasalis ossis frontalis, 74, 77
 orbitalis ossis frontalis, 75-77, 94, 98
 perpendicularis ossis palatini, 83, 90, 91, 97, 99
 petrosa ossis temporalis, 62-64, 70, 71
 tympanica ossis temporalis, 62, 64, 65, 71
 Patella, 107, 136
 Pecten ossis pubis, 128
 Pedicle of the neural arch, 25, 26
 Pelvic brim or inlet, 124, 125, 127
 girdle, 107, 124-126
 outlet, 126, 127
 Pelvis, 124, 127
 axis of, 127
 diameters of, 127
 false, 124, 125
 female, 124
 major, 124, 125
 male, 125, 126
 measurements of, 127
 minor, 124, 125, 127
 muliebris, 124
 true, 124, 125, 127
 virilis, 125, 126
 Perichondrium, 20
 Periosteum, 11, 20
 Petrosal ganglion, depression for, 62, 65
 Petrous portion of the temporal bone, 62-64, 67, 70, 71
 Phalanges digitorum manus, 106, 118, 119, 122
 development of, 123
 pedis, 107, 140-142, 145
 development of, 146, 147
 hallucis, 107
 pollicis, 106
 Phalanges of the fingers, 106, 118, 119, 122
 development of, 123
 of the great toe, 107
 of the thumb, 106
 of the toes, 107, 140-142, 145
 development of, 146, 147
 Pit, olfactory, 102
 Plane, nuchal, 48, 55, 100
 occipital, 55, 100
 Planum nuchale, 48, 55, 100
 occipitale, 55, 100
 popliteum, 132
 sternale, 36
 temporale, 47, 72
 Plate, cribriform, 78, 90, 91, 94
 of the ethmoid bone, orbital, 78, 91, 92, 94
 vertical, 78, 91, 94, 95
 of the frontal bone, orbital, 75-77, 94, 98
 of the palate bone, horizontal, 83, 90, 91, 97
 palate, 83, 90, 91, 97
 vertical, 83, 90, 91, 97, 99
 pterygoid, external, 58, 60, 61, 99
 internal, 58, 61
 tympanic, 62, 64, 65, 71
 Ponticulus promontorii, 67

Portion, nasal, of the frontal bone, 74, 77
 Forus acusticus externus, 47, 48, 62
 internus, 63, 65, 68, 70

Postphenoid, 61

Premaxilla, 82, 97

Presphenoid, 61

Presternum, 41

Process or processes:

 accessory, 27

 alar, 78, 94

 articular, of the sacrum, superior, 30-32
 of the vertebrae, inferior, 25-27, 29
 superior, 25-27, 29

 of the astragalus, external, 143
 posterior, 141, 143
 trochlear, 143

 basilar, of the occipital bone, 54-57, 60

 clinoid, anterior, 58-60

 middle, 59, 60

 posterior, 58, 59

 cochleariform, 64-68

 coracoid, 110, 111

 coronoid (of the mandible), 84, 86
 (of the ulna), 114

 costal, 27

 ensiform, 41

 ethmoidal, 79, 90

 of frontal bone, external angular, 75-77

 humeral, of the lachrymal bone, 79, 98

 of the sphenoid bone, 58, 60, 95

 intrajugular, of the occipital bone, 63

 of the temporal bone, 56

 jugular, 54, 56, 89

 lachrymal, 79, 90

 of the malar bone, frontal, 83

 marginal, 83

 orbital, 83, 98

 temporal, 83

 mamillary, 27

 mastoid, 48, 62, 64, 67, 71, 88

 maxillary, of the inferior turbinate bone, 79, 99

 nasal, of the palate bone, 83, 99

 of the maxillary bone (inferior), alveolar, 81-86
 coronoid, 84-86
 (superior), alveolar, 80, 82, 99
 malar, 80, 82
 nasal, 80, 81, 91
 palatine, 80-82, 91,
 91, 97

 odontoid, 29

 of the palate bone, orbital, 83, 90-92, 98, 99

 pyramidal, 83, 97, 99

 sphenoidal, 83, 90, 91

 paramastoid, 88

 paroccipital, 88

 pterygoid, 48, 58-61, 91, 92

 pterygospinous, 60, 91

 spinous, of the sphenoid bone, 58-61, 93

 (of vertebrae), 25-27

 styloid, of the fibula, 138

 of the radius, 115

 of the temporal bone, 48, 62, 64

 of the third metacarpal bone, 122

 of the ulna, 114, 115

 transverse, 25, 26, 28

 tympanic, anterior, 70, 71

 posterior, 70

 unciform, of the unciform bone, 118, 121

 uncinate, 78, 90, 94, 95, 99

 ungual, of the phalanges of the fingers, 122

 of the toes, 141, 145

 vaginal, of the sphenoid bone, 58, 59, 61, 90, 91

 of the temporal bone, 64, 70, 71

 xiphoid, 41

Processus accessorius, 27

 alaris, 78, 94

 alveolaris (maxillæ), 80, 82, 99

 articularis inferior vertebrarum, 25-27, 29

 superior ossis sacri, 30-32

 vertebrarum, 25-27, 29

 clinoideus, anterior, 58-60

Processus clinoidens, medius, 59, 60
 posterior, 58, 59
 cochleariformis, 65, 66, 68
 condyloideus, 84, 85
 coracoidens, 110, 111
 coronoideus mandibulae, 84-86
 ulnae, 114
 costarius, 27
 ethmoidalis, 79, 90
 frontalis, 80, 81, 91
 frontosphenoidalis, 83
 intrajugularis ossis occipitalis, 56
 temporalis, 63
 jugularis, 54, 56, 89
 lacrimalis, 79, 90
 lateralis tali, 143
 tuberis calcanei, 141, 143
 mamillaris, 27
 marginalis, 83
 mastoidens, 48, 62, 64, 67, 71, 88
 maxillaris, 79, 99
 medialis tuberis calcanei, 141, 143
 nasalis ossis palati, 83, 99
 orbitalis, 83, 90-92, 98, 99
 palatinus, 80-82, 90, 91, 97
 paramastoidens, 88
 posterior tali, 141, 143
 pterygoideus, 48, 58-61, 91, 92
 pterygospinosus (Civinini), 60, 91
 pyramidalis, 83, 97, 99
 sphenoidalis, 83, 90, 91
 spinosus, 25-27
 styloideus ossis metacarpalis III., 122
 temporalis, 48, 62, 64
 radii, 115
 ulnae, 114, 115
 temporalis, 83
 transversus, 25, 26, 28
 trochlearis, 143
 tympanicus, anterior, 70, 71
 posterior, 70
 † uncinatus, 78, 90, 94, 95, 99
 † vaginalis, 58, 59, 61, 90, 91
 xiphoideus, 41
 zygomatikus maxillae, 80, 82
 ossis frontalis, 75-77
 temporalis, 62-64, 70, 71
 Prominence of the aqueduct of Fallopius, 66
 of the external semicircular canal, 66, 67
 styloid, 64
 Prominentia canalis facialis, 66
 semicircularis lateralis, 66, 67
 styloidea, 64
 Promontorium (cavi tympani), 66-68
 (columnae vertebralis), 24, 124
 Promontory of the tympanum, 66-68
 sacral, 24, 124
 Protovertebrae, 34
 Protuberance, mental, 84-86
 occipital, external, 48, 55, 56
 internal, 49, 54, 57, 89
 Protuberantia mentalis, 84-86
 occipitalis externa, 48, 55, 56
 interna, 49, 54, 57, 89
 Pterygoid plate, external, 58, 60, 61, 99
 internal, 58, 61
 Pubis, the, 124-126
 Pyramid of the tympanum, 66, 67, 69
 Pyramis [pars petrosa] ossis temporalis, 62-64, 70, 71

R.

Radius, 13, 106, 115, 116
 development of, 117
 Radix arcus vertebrae, 25, 26
 Ramus, inferior ossis ischii, 128, 130
 pubis, 128, 130
 of the ischium, inferior, 128, 130
 superior, 128
 mandibulae, 84-87

Ramus of the pubis, ascending, 128
 descending, 128, 130
 inferior, 128, 130
 superior, 128
 superior ossis ischii, 128
 pubis, 128
 Recess, elliptical, 69
 epitympanic, 64, 69, 70
 spheno-ethmoidal, 90, 92
 Recessus ellipticus, 69
 epitympanicus, 64, 69, 70
 spheno-ethmoidalis, 90, 92
 Ribs, the, 36-40
 asternal, 36, development of, 42
 false, 36
 floating, 36
 sternal, 36
 true, 36
 Ridge or ridges:
 of the carpus, radial, 118
 ulnar, 118
 cruciform, internal, 54
 gluteal, 132
 for the interarticular costocentral ligament, 40
 intermediate, of the crest of the ilium, 128, 130
 interosseous, of the fibula, 138
 interosseous, of the tibia, 136-138
 intertrochanteric, 132, 133
 mylohyoid, 85
 of the neck of the rib, 40
 orbital, 80, 98
 pectoral, 112
 postauricular, 62
 superciliary, 46, 48, 74
 supinator (of the ulna), 114
 transverse, of the sacrum, 30
 of the trapezium, 118, 121
 trapezoid, 109
 Ring, tympanic, 70, 103, 104
 Roof of the skull, 50
 of the tympanum, 65
 Rostrum, sphenoidal, 58-61, 91
 primitive, 61
 Row of teeth, 48

S.

Sacrum, the, 24, 30-33, 125, 126
 development of, 35
 apex of the, 30, 31
 Scapula, the, 106, 108, 110
 development of, 111
 Sella turcica, 60, and footnote
 Semicanal musculi tensoris tympani, 64, 66, 67, 70
 tubae auditivae, 64, 66, 67, 69
 Septa, interalveolar, 82, 84
 Septum between the canal for the tensor tympani muscle
 and the Eustachian canal, 64-68
 canalis musculotubarii, 64, 66, 67
 of the frontal sinuses, 76, 94
 nasi osseus, 46, 91-93, 96
 of the nose, bony, 46, 91-93, 96
 sinuum frontalem, 76, 94
 sphenoidalium, 60, 93
 sphenoidal, 60, 93
 Shaft of the femur, 132, 133
 of the fibula, 138
 of the humerus, 112
 of the radius, 115
 of the tibia, 136, 137
 of the ulna, 114
 Shafts of the metacarpal bones, 145
 of the metatarsal bones, 145
 of the phalanges of the fingers, 122
 of the toes, 145
 of the ribs, 40
 Shoulder-girdle, 106, 108
 Sinciput, 47
 † Sinus cervicalis, 102
 frontalis, 76, 77, 90, 91, 94, 95
 maxillaris, 80-82, 93-95, 97, 99
 posterior (cavi tympani), 64, 66, 67

- Sinus sphenoidalis, 60, 61, 90-93
 tarsi, 140
 tympani, 66, 67
- Skeleton, appendicular, 105-147
 axial, 23-104
 extremities inferioris, 105, 107, 124-147
 superioris, 105, 106, 108-123
 of the foot, 140-142
 development of, 146, 147
 of the hand, 118, 119
 development of, 123
 human, 22
 of the lower limb, 105, 107, 124-127
 manus, 118, 119
 development of, 123
 pedis, 140-142
 development of, 146, 147
 trunci, 23-43
 of the trunk, 23-43
 of the upper limb, 105, 106, 108-123
- Skull (see also "Cranium"):
 cap, 50
 measurements of, 101
 views of:
 base, external, 48
 internal, 49
 front, 46, 101
 roof, inner, 50
 side, 47, 101
 top, 101
- Small wing of the sphenoid, 58-61
- Space, interosseous, of the forearm, 106, 116
 of the leg, 107
- Spaces, intercostal, 36
- Spatia intercostalia, 36
- Spatium interosseum antibrachii, 106, 116
 cruris, 107
- Sphenoid bone, 58-60
 development of, 61
- Spina *vel* spine:
 angularis, 58-61, 93
 ethmoidalis, 59
 frontalis, 74-77, 90, 91
 iliaca anterior inferior, 128, 129
 superior, 128-130
 posterior inferior, 128, 129
 superior, 128, 129
 ischiadica, 128, 129
 mentalis, 84, 85, 96
 musculi recti lateralis, 58
 nasalis anterior, 46, 47, 80, 90, 91, 93, 97
 posterior, 90, 91, 97
 palatina, 82, 97
 scapula, 110
 suprameatum, 62
 trochlearis, 76, 98
 tympanica major, 64, 71
 minor, 64, 70, 71
- Spine, ethmoidal, of the sphenoid bone, 59
 iliac, anterior inferior, 128, 129
 superior, 128-130
 posterior inferior, 128, 129
 superior, 128, 129
 of the ischium, 128, 129
 for the lower head of the external rectus muscle of the eyeball, 58
 mental, 84, 85, 96
 nasal, anterior, of the superior maxillary bone, 46, 47, 90, 91, 93, 97
 of the frontal bone, 74-77, 90, 91
 posterior, 90, 91, 97
 palatine, 90, 91, 97
 peroneal, of the astragalus, 143
 pharyngeal, 56
 pubic, 130
 of the scapula, 110
 of the sphenoid bone, 58-61, 93
 suprameatal, 62
 of the tibia, 136, 137
 trochlear, 76, 98
- Spine, tympanic, anterior, 64, 71
 posterior, 64, 70, 71
- *Spines, palatine, 82, 97
- Spongy bone, sphenoidal, 58, 59, 61
- Squama frontalis, 74-77
 occipitalis, 54-57
 temporalis, 62-64, 70, 71
- Squamous portion of the frontal bone, 74-77
 of the occipital bone, 54-57
 of the temporal bone, 62-64, 70, 71
- Sternum, 41
 development of, 42
- Subiculum promontorii, 67
- Substance of bone, cortical, 19, 20
- Substantia compacta, 11-17, 19
 corticalis, 19, 20
 spongiosa, 11-17, 19
- Sulcus *vel* sulci:
 arteriæ occipitalis, 62, 71
 temporalis medice, 62
 vertebralis, 29
 arteriosi, 50, 73
 calcanei, 143
 canaliculi mastoidei, 62, 65
 carotici, 58, 59
 carpi, 118
 chiasmatis, 59
 costæ, 40
 ethmoidalis, 79, 91
 hamuli pterygoidei, 58
 infra-orbitalis, 80, 93, 94, 98, 99
 intertubercularis, 112
 lacrimalis maxillæ, 80, 82, 91, 98
 ossis lacrimalis, 79, 92
 lateral, 49, 54, 56, 63, 64, 73, 88, 89
 longitudinal, 54, 73, 75, 88, 89
 mallei, 70
 malleolaris (tibiar), 136
 musculi flexoris hallucis longi (calcanei), 141, 143
 (tali), 141, 143
 peronæi longi (calcanei), 140, 141, 144
 (ossis cuboidei), 140, 141, 144
 mylohyoideus, 85
 nervi petrosi superficialis majoris, 63, 64
 minoris, 63, 64
 radialis, 112
 spinalis, 26
 ulnaris, 112
 obturatorius, 128
 olfactorius, 90
 palatini, 82, 97
 paraglenoidalis, 128
 petrosal, inferior, 56, 62, 63, 88
 superior, 63
 petrosquamosus, 64
 petrosus inferior, 88
 ossis occipitalis, 56
 temporalis, 62, 63
 superior, 63
- † præauricularis, 128
 promontorii, 67
- † pterygopalatinus maxillæ, 80
 ossis palatini, 83
 sphenoidalis, 58
- sagittalis (ossis frontalis), 75, 88
 (ossis occipitalis), 54, 89
 (ossis parietalis), 73
- sigmoideus, 49, 56, 63, 64, 88, 89
 subclaviæ, 40
 tali, 143
 transversus ossis occipitalis, 49, 54, 56, 89
 parietalis, 73
- tubæ auditivæ, 48
 tympanicus, 64, 66, 69, 70
 venosus, 50
- Supra-occipital bone, note to p. 57
 portion of the occipital bone, 54-57, and
 note, p. 57
- Surface, articular, of the astragalus, superior, 143
 of the external malleolus, 137, 138

- Surface, articular, of the head of the fibula, 138
 of the internal malleolus, 137
 of the patella, 136
 of the radius, for the ulna, 115
 of the tibia, for the head of the fibula, 136, 137
 inferior or distal, 137
 superior or proximal, 137
- auricular, of the ilium, 128
 of the sacrum, 31, 32
 of the femur, anterior, 132, 133
 patellar, 132-134
 popliteal, 132
 postero-external, 132, 133
 postero-internal, 132, 133
- of the fibula, anterior,
 external, 138
 internal, 138
 posterior, 138
 of the foot, dorsal, 140
 plantar, 141
- of the frontal bone, anterior, 74
 cerebral, 75
 frontal, 74
 orbital, 76
 posterior, 75
 temporal, 74, 77
- of the great wing of the sphenoid bone, cerebral, 58, 61
 of the great wing of the sphenoid bone, orbital, 58-61, 98
 of the great wing of the sphenoid bone, sphenomaxillary, 58, 99
 of the great wing of the sphenoid bone, temporal, 58-61, and note, p. 59
 of the hand, dorsal, 119
 palmar, 118
- of the humerus, antero-external, 112
 antero-internal, 112
 posterior, 112
- of the malar bone, malar, 83
 orbital, 83
 outer, 83
 temporozygomatic, 83
- patellar (of the femur), 132-134
- of the parietal bone, cerebral, 73
 external, 72
 internal, 73
 parietal, 72
 temporal, 47, 72
- of the petrous portion of the temporal bone, anterior, 65
 of the petrous portion of the temporal bone, inferior, 65
 of the petrous portion of the temporal bone, posterior, 65
- popliteal, 132
- of the pubic symphysis, 128
- of the radius, anterior, 114-116
 external, 114-116
 posterior, 114-116
- of the sacrum, dorsal, 31, 32
 pelvic, 30, 32, 125
 posterior, 126
- of the scapular, anterior, 110
 posterior, 110
- of the squamous portion of the temporal bone, cerebral, 63, 64
 of the squamous portion of the temporal bone, temporal, 62, 63
- of the sternum, anterior, 36
- of the superior maxillary bone, anterior, 80
 facial, 80
 nasal, 80
 orbital, 80
 zygomatic, 80, 99
- temporozygomatic, 58-61, and note, p. 59
- of the tibia, external, 136-138
 internal, 136, 138
 posterior, 136-138
- Surface, of the ulna, anterior, 114, 116
 internal, 114
 posterior, 114
- of the vertical plate of the palate-bone, external, 83
 internal, 83
 nasal, 83
- zygomatic, 58-61, and note, p. 59
- Sustentaculum tali, 140, 141, 143
- Sutura coronalis, 46, 47, 49, 50, 88, 100
 ethmoidomaxillary, 98
 frontalis, 77
 fronto-ethmoidalis, 49, 98
 frontolacrimalis, 91, 98
 frontomaxillary, 46, 98
 incisiva, 48, 82, 97
 infra-orbitalis, 80
 intermaxillary, 46
 internasalis, 46, 92
 lacrimoconchal, 90, 95
 lacrimo-ethmoidalis, 91, 98
 lacrimomaxillary, 91, 98
 lamboidea, 47, 50, 89, 100
 mendoza, 55, 57, 104
 nasofrontalis, 46, 90-92, 98
 nasomaxillary, 46, 47
 occipitalis transversa, 100
 occipitomastoid, 47-49, 88, 89, 100
 palatina mediana, 96, 97
 transversa, 90, 91, 96, 97
 palato-ethmoidalis, 99
 palatomaxillary, 98
 parietomastoid, 47, 49, 89, 100
 sagittalis, 50, 88, 100
 sphenomaxillary, 49, 98, 99
 sphenofrontalis, 47
 sphenoorbitalis, 49, 98
 sphenoparietalis, 46, 47, 88
 sphenosquamosa, 46-49, 88
 splinozygomatica, 47, 98
 squamosa, 46, 47, 88
 squamosomastoid, 62, 70, 71
 zygomaticofrontalis, 46, 47, 92
 zygomaticomaxillary, 46, 47
 zygomaticotemporalis, 46, 47
- Suture, coronal, 46, 47, 49, 50, 88, 100
 ethmoidomaxillary, 98
 frontal, 77
 fronto-ethmoidal, 49, 98
 frontolachrymal, 91, 98
 frontomalar, 46, 47, 92
 frontomaxillary, 46, 98
 frontoparietal, 46, 47, 49, 50, 88, 100
 infra-orbital, 80
 intermaxillary, 46
 interparietal, 50, 88, 100
 internasal, 46, 92
 lachrymo-ethmoidal, 91, 98
 lachrymomaxillary, 91, 98
 lachrymotubinal, 90, 95
 lamboidei, 47, 50, 89, 100
 malomaxillary, 46, 47
 mendoza, 55, 57, 104, and note to p. 57
 metopic, 77
 nasofrontal, 46, 90-92, 98
 nasomaxillary, 46, 67
 occipital, transverse, 55, 57, 100, 104, and note to p. 57
 occipitomastoid, 47-49, 88, 89, 100
 occipitoparietal, 47, 50, 89, 100
 palatine, anterior, 48, 82, 97
 median, 96, 97
 middle, 96, 97
 posterior, 90, 91, 96, 97
 transverse, 90, 91, 96, 97
 palato-ethmoidal, 99
 palatomaxillary, 98
 parietomastoid, 47, 49, 89, 100
 petrosphenoidal, 48, 49
 premaxillary, 48, 82, 97
 sagittalis, 50, 88, 100

Suture, sphenothmoidal, 49, 98, 99
 sphenofrontal, 47
 sphenomalar, 47, 98
 sphenoorbital, 49, 98
 sphenoparietal, 49, 47, 88
 sphenosquamous, 46-49, 88
 squamosomastoid, 62, 70, 71
 squamous, 46-48
 temporomalar, 46, 47

Symphysis ossium pubis, 125
 pubic, 125

Synchondrosis epiphyseos, 20
 intersphenoidalis, 61
 intraoccipitalis, anterior, 57
 posterior, 57
 occipitosphenoidal, 49
 sphenoccipitalis, 49
 sternalis, 41

T.

Table, inner, of the bones of the skull, 19, 50
 of the frontal bone, 75-77
 outer, of the bones of the skull, 19, 50
 of the frontal bone, 75, 76

Tabular portion of the occipital bone, 54-57, and note.
 p. 57

Talus, 17, 142, 143

Tarsus, 107, 140

Tegmen tympani, 64, 65, 69-71, 96

Thorax, 36, 37

Tibia, 15, 18, 107, 136, 137

development of, 139

Tissue of bone, cancellous, 11-17, 19
 compact, 11-17, 19

Toes, skeleton of, 140, 141

Tooth-sockets of the lower jaw, 84
 of the upper jaw, 82

*Torus occipitalis, 100

palatinus, 95

Tactus spiralis foraminulentus, 68

Trapezium, the, 119-121

Trochanter, great, 132-134

major, 132-134

minor, 132, 133

small, 132, 133

tertius, 133

third, 133

Trochlea humeri, 112

phalangis manus, 127

pedis, 145

tali, 140, 143

Tuber calcanei, 140, 141, 143, 147

cochleæ, 66-68

frontale, 46, 74, 77, 104

ischadicum, 128-130

maxillare, 80-82, 99

parietale, 72, 104

Tubercle of the atlas, anterior, 29

posterior, 29

carotid, 24

of the cervical vertebræ, anterior, 26

posterior, 26, 29

conoid, 109

infraglenoid, 110

of the ischium, 128-130

of Lisfranc, 40

mental, 84, 85, 96

obturator, anterior, 128

posterior, 128, 130

pharyngeal, 56

of the posterior process of the astragalus,

external, 143

of the posterior process of the astragalus,

internal, 143

of the ribs, 37, 40

scalenæ, 40

of the spine of the tibia, inner, 137

outer, 137

supraglenoid, 110

Tubercle of the tibia, 136, 137

of the tuberosity of the calcaneum, inner, 141, 143
 outer, 141, 143

Tuberculum anterius atlantis, 29

vertebrarum cervicalium, 26

articulare, 62

caroticum, 24

costæ, 37, 40

intercondyloideum laterale, 137

mediale, 137

jugulare, 54, 56

laterale tali, 143

majus, 112

mediale tali, 143

mentale, 84, 85, 96

minus, 112

obturatorium anterius, 128

posterior, 128, 130

ossis multanguli majoris, 118, 121

navicularis manus, 118, 120

pharyngeum, 56

posterior atlantis, 29

vertebrarum cervicalium, 26, 29

pubicum, 130

scalenæ [Lisfranci], 40

sellæ, 59-61

Tuberositas coracoidea, 109

costæ II., 40

costalis, 109

deltoidæ, 112

glutea, 132

iliaca, 128, 129

infraglenoidalis, 110

masseterica, 84, 86

ossis cuboidei, 141, 144

metacarpalis V., 122

metatarsalis I., 141, 145

V., 140, 141, 145

navicularis pedis, 140, 141, 144

pterygoidea, 85

radii, 115, 116

sacralis, 31

supraglenoidalis, 110

tibiae, 136, 137

ulnæ, 114

unguicularis phalangum digitorum (manus),

122

unguicularis phalangum digitorum pedis, 141,

145

Tuberosity, costal, 109

of the cuboid bone, 141, 144

of the femur, external, 132, 134

internal, 132, 133

of the fifth metacarpal bone, 122

of the first metatarsal bone, 141, 145

of the fifth metatarsal bone, 140, 141, 145

of the humerus, great, 112

small, 112

of the ilium, 128, 129

masseteric, 84, 86

of the navicular bone, 140, 141, 144

of the os calcis, 140, 141, 143, 147

of the palate-bone, 83, 97, 99

pterygoid, internal, 85

of the radius, 115, 116

of the sacrum, 31

of the scaphoid bone, 118, 120

of the superior maxillary bone, 80-84, 99

of the tibia, external, 136, 137

internal, 136, 137

of the trapezium, 118, 121

of the ulna, 114

Turbinatè bone, sphenoidal, 58, 59, 61

orbital border of the, 59

Tympanic plate, 62, 64, 65, 71

ring, 70, 103, 104

sinus, 66, 67

spine, anterior, 64, 71

posterior, 64, 70, 71

sulcus, 64, 66, 69, 70

U.

Ulna, the, 13, 106, 114, 116
development of, 117

V.

Vaginal process of the temporal bone, 64, 70, 71
of the sphenoid bone, 58, 59, 61, 90, 91
Vagina processus styloidei, 64, 70, 71
Venter of the scapula, 110
Vertebra or vertebrae:
cervical, 24, 26, 29
seventh, 29
development of, 34, 55
dorsal, 24, 25, 28
first, 28
eleventh, 28
twelfth, 28
lumbar, 24, 27, 28
fifth, 28, 33
prominens, 24
sacral, 24, 30-32
structure, 16
thoracales, 24, 25, 28
thoracalis I., 28
XI., 28
XII., 28
Vertex, 46, 47

Vesicle, optic, primary, 102
umbilical, 102
Vestibulum labyri thi, 65, 68, 69
Vomer, 53, 79, 91, 94, 95

W.

Wall of the orbit, external, 98
inferior, 93, 98
internal, 98
superior, 92, 98
of the tympanum, carotid, 65, 66
inner, 65, 66
jugular, 66
labyrinthine, 65, 66
mastoid, 66
superior (roof), 65
Wing of the ilium, 130
of the sphenoid, great, 58-61
small, 58-61

X.

Xiphisternum, 41

Z.

Zygoma, 46-48, 62-64, 70, 71, 96
Zygomatic width, 101

AN ATLAS
OF
HUMAN ANATOMY
FOR STUDENTS AND PHYSICIANS

BY
CARL TOLDT, M.D.

ASSISTED BY
PROFESSOR ALOIS DALLA ROSA, M.D.

Adapted to English and American and International Terminology

BY
M. EDEN PAUL, M.D. BRUX., M.R.C.S., L.R.C.P.

SECOND SECTION

C. ARTHROLOGY

(FIGURES 378 TO 489 AND INDEX)

REVISED EDITION

NEW YORK
REBMAN COMPANY
141, 143 AND 145 WEST 36TH STREET

Copyright, 1919, by
REBMAN COMPANY

SYNDESMOLOGIA
ARTHROLOGY

THE ARTICULATIONS IN GENERAL

FIG. 378.—SUTURA SERRATA—SERRATED OR DENTATED SUTURE.

FIG. 379.—SUTURA SQUAMOSA—SQUAMOUS OR SCALY SUTURE.

FIG. 380.—HARMONIA—HARMONIC SUTURE.

Synarthrosis, or Continuous Articulation.

FIG. 381.—SYNCHONDROSIS. (THE SPHENO-OCCIPITAL SYNCHONDROSIS OF A GIRL AT THE AGE OF TWO YEARS; MEDIAN SAGITTAL SECTION.)

FIG. 382.—SYMPHYSIS. (THE PUBIC SYMPHYSIS; FRONTAL SECTION, POSTERIOR PORTION.)

FIG. 383.—GOMPHOSIS.

Synarthrosis, or Continuous Articulation.

FIG. 384.—EXTENSION. POSTERIOR ASPECT.

FIG. 385.—FLEXION. LATERAL ASPECT.

DIARTHROSIS (METACARPO-PHALANGEAL ARTICULATION).—GINGLYMUS, OR HINGE JOINT (ARTICULATIONS OF THE FINGERS).

FIG. 386.—ARTICULATIO COCHLEARIS, COCHLEOID OR SCREW GINGLYMUS. (SUPERIOR ARTICULAR SURFACE OF THE LEFT ASTRAGALUS, SEEN FROM BEHIND, SHOWING THE SCREW FORM OF THE ANKLE-JOINT).

Diarthrosis, or Discontinuous Articulation—Articulatio simplex, simple joint.

FIG. 387.—CONDYLARTHROSIS, OR CONDYLOID JOINT. (THE METACARPO-PHALANGEAL ARTICULATION OF THE INDEX-FINGER; LATERAL VIEW.)

FIG. 388.—CONDYLARTHROSIS, OR CONDYLOID JOINT. (RIGHT OCCIPITO-ATLANTAL ARTICULATION; SEEN FROM THE OUTER SIDE.)

The outer half of the lateral mass of the atlas has been removed by a sagittal section passing through the joint.

FIG. 389.—ARTICULATIO SELLARIS—SADDLE JOINT. (ARTICULATIO CARPOMETACARPEA POLLICIS—CARPOMETACARPAL JOINT OF THE THUMB.)

Diarthrosis, or Discontinuous Articulation—Articulatio simplex, simple joint.

FIG. 390.—ENARTHROSIS,¹ OR ARTHRODIA; BALL-AND-SOCKET JOINT. (THE RIGHT SHOULDER-JOINT SEEN FROM BEFORE, THE ANTERIOR HALF OF THE GLENOID FOSSA AND OF THE CAPSULE OF THE JOINT HAVING BEEN REMOVED.)

FIG. 391.—ENARTHROSIS,¹ OR ARTHRODIA; BALL-AND-SOCKET JOINT. (THE RIGHT HIP-JOINT SEEN FROM BEFORE, THE ANTERIOR HALF OF THE ACETABULUM AND OF THE CAPSULE OF THE JOINT HAVING BEEN REMOVED.)

¹ I have departed a little on this page from the author's terminology, which differs slightly from that in use in England. Both the hip and the shoulder joint are classed by him as examples of ARTICULATIO SPHÆROIDEA. *Kugelgelenk*—lit. "ball joint"; one of these, the shoulder-joint, in which the concave articular surface is considerably less than a hemisphere in extent, is called by him ARTHRODIA, *freies Gelenk*—lit. "free joint"; while the other, the hip-joint, in which the concave articular surface is considerably more than a hemisphere in extent, is called ENARTHROSIS, *Nutgelenk*—lit. "nut joint." English anatomists do not, as a rule, draw this distinction. The movements of the hip and the shoulder joint are identical in character, and the greater extent of the rigid portion of the enveloping surface in the one case than in the other has not been considered a difference sufficiently important to warrant a separation into two classes. Hence, in England the terms ENARTHROSIS and ARTHRODIA are applied indifferently to all ball-and-socket joints.—TR.

Diarthrosis, or Discontinuous Articulation—*Articulatio simplex*, simple joint; *articulatio sphæroidea*,¹ ball-and-socket joint.

FIG. 392.—ARTICULATIO COMPOSITA, COMPOUND JOINTS (ARTICULATIONS OF THE HAND).—AMPHIARTHROSIS=ARTHRODIA OR GLIDING JOINT, AS SEEN IN THE INTERCARPAL, CARPO-METACARPAL, AND INTERMETACARPAL JOINT.

The arrows show the dorsal emergence of the axes of the wrist joint and of the transverse carpal articulation respectively.

FIG. 393.—ARTICULATIO TROCHOIDEA, TROCHOID OR PIVOT JOINT.—GINGLYMUS, HINGE JOINT. (PALMAR ASPECT OF THE ELBOW-JOINT.)

The radial half of the capitulum of the humerus has been removed by a sagittal section passing through its centre of curvature.

Diarthrosis, or Discontinuous Articulation.
Amphiarthrosis, arthrodia or gliding joint—Articulatio composita, compound joint.

FIG. 394.—SYNDESMOSIS.—INTERARTICULAR FIBROCARILAGE. (THE STERNOCLAVICULAR ARTICULATION, ANTERIOR ASPECT.)

The right articulation is divided through the middle by a frontal section; in the left, the front portion of the capsule has been removed, and the clavicle has been drawn backwards.

FIG. 395.—SYNDESMOSSES SCAPULÆ, PROPER LIGAMENTS OF THE SCAPULA; SEEN FROM ABOVE.

Syndesmosis, fibrous or gamentous union—Discus articularis, interarticular fibrocartilage.

JUNCTURÆ OSSIIUM TRUNCII
THE ARTICULATIONS OF THE
TRUNK

FIG. 396.—FIBROCARTILAGINES INTERVERTEBRALES, INTERVERTEBRAL DISCS. (SECOND, THIRD, AND FOURTH LUMBAR VERTEBRÆ; SEEN FROM BEFORE.)

Ligamenta columnæ vertebralis—The ligaments of the vertebral column.

FIG. 397.—INTERVERTEBRAL DISC BETWEEN THE SECOND LUMBAR VERTEBRA AND THE THIRD.
(LOWER HALF OF A HORIZONTAL SECTION.)

columnæ vertebralis—The ligaments of the vertebral column.

FIG. 398.—FIBROCARILAGINES INTERVERTEBRALES, INTERVERTEBRAL DISCS. LIGAMENTA FLAVA, OR SUBFLAVA. LIGAMENTA INTERSPINALIA, INTERSPINOUS LIGAMENTS. LIGAMENTUM SUPRASPINALIS, SUPRASPINOUS LIGAMENT. (MEDIAN SECTION THROUGH THE TWELFTH DORSAL AND THE FIRST LUMBAR VERTEBRÆ, LEFT HALF.)

Ligamenta columnæ vertebralis—The ligaments of the vertebral column.

Capsule of the joint of the articular processes
Capsula articulationis intervertebralis

Arch of the first dorsal vertebra
Arcus vertebræ thoracalis I.

FIG. 399.—LIGAMENTA FLAVA, LIGAMENTA SUBFLAVA. ARTICULATIONES INTERVERTEBRALES, INTERVERTEBRAL ARTICULATIONS. OF THESE LATTER THE RIGHT ARE UNOPENED, THE LEFT OPENED. (THE ARCHES OF THE FIRST FOUR DORSAL VERTEBRÆ, SEPARATED FROM THE BODIES BY A FRONTAL SECTION. SEEN FROM BEFORE.)

Ligamenta columnæ vertebralis—The ligaments of the vertebral column.

FIG. 400.—LIGAMENTUM LONGITUDINALE ANTERIUS, THE ANTERIOR COMMON LIGAMENT. LIGAMENTUM LUMBOCOSTALE, LUMBOCOSTAL LIGAMENT. (THE VENTRAL ASPECT OF THE VERTEBRAL COLUMN, THE ANTERIOR HALF OF THE BASE OF THE SKULL AND THE ANTERIOR HALF OF THE PELVIS HAVING BEEN REMOVED.)

Ligamenta columnæ vertebralis—The ligaments of the vertebral column.

FIG. 401.—INTERVERTEBRAL DISCS, INTERSPINOUS LIGAMENTS, SUPRASPINOUS LIGAMENT, LIGAMENTUM NUCHÆ, SPINAL CANAL; INTERVERTEBRAL FORAMINA. VERTICAL AXIS OF THE VERTEBRAL COLUMN. (MEDIAN SECTION THROUGH THE VERTEBRAL COLUMN.)

FIG. 402.—POSTERIOR COMMON LIGAMENT AND ITS RELATION WITH THE POSTERIOR OCCIPITO-AXIAL AND THE DEEP POSTERIOR SACRO-COCYGEAL LIGAMENTS. (THE VERTEBRAL COLUMN WITH THE SPINAL CANAL OPENED BY THE REMOVAL OF THE NEURAL ARCHES; SEEN FROM BEHIND.)

¹ Macalister calls this ligament *ligamentum latum axiale*, the broad axial ligament.—Tr.

FIG. 403.—LIGAMENTUM LONGITUDINALE ANTERIUS, THE ANTERIOR COMMON LIGAMENT OF THE VERTEBRAL COLUMN. (THE FOURTH TO THE TENTH DORSAL VERTEBRÆ; SEEN FROM BEFORE.)

Ligamenta columnæ vertebralis—The ligaments of the vertebral column.

FIG. 404.—LUMBAR AND SACRAL PORTIONS OF THE POSTERIOR COMMON LIGAMENT, AND THE CONNEXIONS OF THIS LIGAMENT WITH THE SPINAL DURA MATER AND WITH THE DEEP POSTERIOR SACROCOCYGEAL LIGAMENT. (THE SACRUM AND THE LUMBAR PORTION OF THE VERTEBRAL COLUMN WITH THE SPINAL CANAL LAID OPEN FROM BEHIND.)

The sacral portion of the dura mater has been drawn to the left side.

Ligamenta columnæ vertebralis—The ligaments of the vertebral column.

FIG. 405.—THE LIGAMENTS CONNECTING THE ANTERIOR SURFACES OF THE SACRUM AND COCCYX: THE ANTERIOR AND THE LATERAL SACROCOCYGEAL LIGAMENTS.

FIG. 406.—THE LIGAMENTS CONNECTING THE POSTERIOR SURFACES OF THE SACRUM AND COCCYX: THE SUPERFICIAL AND DEEP POSTERIOR SACROCOCYGEAL LIGAMENTS, AND THE INTERARTICULAR AND LATERAL SACROCOCYGEAL LIGAMENTS.

THE COCCYX WITH THE TWO LAST SACRAL VERTEBRÆ.

Ligamenta columnæ vertebralis—The ligaments of the vertebral column.

FIG. 407.—LIGAMENTUM NUCHÆ. (THE CERVICAL PORTION OF THE VERTEBRAL COLUMN AND THE POSTERIOR PORTION OF THE CRANIUM; SEEN FROM THE RIGHT SIDE.)

Ligamenta columnæ vertebralis—The ligaments of the vertebral column.

FIG. 408.—ARTICULATIONES CAPITULORUM ET COSTOTRANSVERSARIÆ, THE COSTOCENTRAL AND COSTOTRANSVERSE ARTICULATIONS: LIGAMENTUM CAPITULI COSTÆ INTERARTICULARE, THE INTERARTICULAR COSTOCENTRAL LIGAMENT; LIGAMENTUM COLLI COSTÆ ET LIGAMENTUM TUBERCULI COSTÆ, THE MIDDLE AND POSTERIOR COSTOTRANSVERSE LIGAMENTS. (THE SEVENTH DORSAL VERTEBRA WITH THE VERTEBRAL EXTREMITIES OF THE SEVENTH PAIR OF RIBS; SEEN FROM ABOVE.)

On the left side the costocentral and costotransverse articulations have been opened by a horizontal section through the rib and the vertebral body; on the right side the section passes through the intervertebral disc on a plane just above the attachment of the interarticular costocentral ligament to the ridge between the two articular facets on the vertebral extremity of the rib.

Articulationes costovertebrales—Costovertebral articulations.

¹ This term, *costotransverse foramen*, is also used by English anatomists to denote the foramina in the transverse processes of the cervical vertebrae for the transmission of the vertebral artery.—Tr.

² These are upper and lower costal facets respectively in relation to the articular surface of the head of the rib; but, strictly speaking, what is here called "upper costal facet" is the lower costal facet of the sixth dorsal vertebra; while what is here called "lower costal facet" is the upper costal facet of the seventh dorsal vertebra.—Tr.

FIG. 409.—COSTOCENTRAL AND COSTOTRANSVERSE ARTICULATIONS: ANTERIOR COSTOCENTRAL OR STELLATE LIGAMENT; INTERARTICULAR COSTOCENTRAL LIGAMENT; ANTERIOR SUPERIOR COSTOTRANSVERSE LIGAMENT; AND COSTOTRANSVERSE FORAMINA. POSTERIOR OR INTERNAL INTERCOSTAL APONEUROSES. (THE THIRD TO THE SEVENTH DORSAL VERTEBRÆ WITH THE VERTEBRAL EXTREMITIES OF THE THIRD TO THE SEVENTH RIGHT RIBS; SEEN FROM THE RIGHT AND FROM BEFORE.)

The third, fourth, and fifth costocentral articulations are unopened; the sixth and the seventh have been opened from before. In the sixth articulation, by the partial removal of the bodies of the fifth and sixth dorsal vertebrae, the upper and lower surfaces of the intervertebral disc have been exposed, and the attachment of the disc to the ridge between the two articular facets on the head of the rib has been demonstrated.

Articulationes costovertebrales—Costovertebral articulations.

FIG. 410.—THE LIGAMENTS CONNECTING THE DORSAL VERTEBRÆ WITH THE RIBS POSTERIORLY: POSTERIOR AND POSTERIOR SUPERIOR COSTOTRANSVERSE LIGAMENTS; INTERTRANSVERSE LIGAMENTS; POSTERIOR OR INTERNAL INTERCOSTAL APONEUROSSES; SUPRASPINOUS LIGAMENT. (SIXTH TO TENTH DORSAL VERTEBRÆ WITH THE VERTEBRAL EXTREMITIES OF THE SIXTH TO TENTH RIBS.)

Articulationes costovertebrales—Costovertebral articulations.

Third cervical vertebra
Vertebra cervicalis III.

Costotransverse facet
Fovea costalis transversalis

Capsule of the joint of the articular
processes
Capsula articulationis intervertebralis

Posterior costotransverse
ligament
Lig. tuberculi costæ

Third dorsal vertebra
Vertebra thoracalis III.

Interarticular costocentral ligament
Lig. capituli costæ interarticulare

Capsule of the costocentral articulation
Capsula articulationis capituli

Capsule of the costotransverse articulation
Capsula articulationis costotransversariæ

Anterior superior costotransverse ligament
Lig. costotransversarium anterius

FIG. 411.—THIRD, FOURTH, AND
FIFTH CERVICAL VERTEBRÆ
SEEN FROM THE RIGHT SIDE.

FIG. 412.—THIRD, FOURTH, AND FIFTH DORSAL VERTEBRÆ
SEEN FROM THE RIGHT SIDE.

Superior articular facet
Facies articularis superior

Mamillary process
Processus mamillaris

Accessory process
Processus accessorius

Second lumbar vertebra
Vertebra lumbalis II.

Intervertebral disc
Fibrocartilago intervertebralis

Capsule of the joint of the
articular processes
Capsula articulationis intervertebralis

FIG. 413.—SECOND, THIRD, AND FOURTH LUMBAR VERTEBRÆ SEEN FROM THE RIGHT AND
FROM BEHIND.

THE DIRECTION OF THE ARTICULAR SURFACES AND THE CONNEXIONS OF THE CAPSULES
OF THE JOINTS OF THE ARTICULAR PROCESSES IN THE CERVICAL, DORSAL, AND LUMBAR
VERTEBRÆ RESPECTIVELY.

The Synovial Articulations of the Vertebral Column.

FIG. 414.—*MEMBRANE OF THE STERNUM. ANTERIOR CHONDROSTERNAL LIGAMENTS, COSTO-XIPHOID LIGAMENTS, AND ANTERIOR OR EXTERNAL INTERCOSTAL APONEUROSES. THE RELATION OF THESE LATTER TO THE EXTERNAL INTERCOSTAL MUSCLES AND TO THE PECTORALIS MINOR MUSCLE. (ANTERIOR WALL OF THE THORAX SEEN FROM BEFORE.)

Articulaciones sternocostales—Chondrosternal articulations.

FIG. 415.—POSTERIOR OR INTERNAL INTERCOSTAL APONEUROSES, AND THEIR RELATION TO THE TRIANGULARIS STERNI AND TRANSVERSALIS ABDOMINIS MUSCLES. STERNOCLAVICULAR ARTICULATION. (ANTERIOR WALL OF THE THORAX SEEN FROM BEHIND.)

Articulationes sternocostales—Chondrosternal articulations.

FIG. 416.—CHONDROSTERNAL ARTICULATIONS. INTERCHONDRAL ARTICULATIONS. INTERSTERNAL SYNCHONDROSES. STERNOCLAVICULAR ARTICULATION. (POSTERIOR HALF OF A FRONTAL SECTION THROUGH THE STERNUM, THE CARTILAGES OF THE RIBS, AND THE STERNAL EXTREMITIES OF THE CLAVICLES.)

Articulationes sternocostales—Chondrosternal articulations.

ARTICULATIONES
ET LIGAMENTA CAPITIS

THE ARTICULATIONS
AND LIGAMENTS OF THE HEAD

¹ See note to p. 177.

FIG. 417.—POSTERIOR OCCIPITO-AXIAL LIGAMENT. OCCIPITO-ATLANTAL AND ATLANTO-AXIAL SYNOVIAL JOINTS. (THE THREE UPPERMOST CERVICAL VERTEBRÆ AND THE OCCIPITAL BONE SEEN FROM BEHIND.)

By a frontal section behind the occipital condyles, the squamous portion of the occipital bone and the neural arches have been removed. The dura mater has been cut transversely in the basilar groove, and turned upwards

FIG. 418.—TRANSVERSE LIGAMENT OF THE ATLAS. (THE ATLAS SEEN FROM ABOVE.)

Articulationes atlanto-occipitalis et atlanto-epistropheæ—Occipito-atlantal and atlanto-axial articulations.

FIG. 419.—CRUCIFORM LIGAMENT OF THE ATLAS; LATERAL OR ALAR ODONTOID OR CHECK LIGAMENTS OCCIPITO-ATLANTAL AND ATLANTO-AXIAL SYNOVIAL JOINTS, THE RIGHT CLOSED, THE LEFT OPEN.

The cranial dura mater and the posterior or long occipito-axial ligament¹ have been cut transversely in the basilar groove and turned upwards.

¹ See note to p. 177.

² Accessory occipito-axial ligament. This ligament is not mentioned by the author, though it is well shown in Fig. 419. The accessory ligament is a bundle of fibres strengthening the capsule of the occipito-atlantal joint at its postero-internal angle. It runs downwards and inwards from the back of the occipital condyle to the body of the axis near the base of the odontoid process.—Tik.

FIG. 420.—LATERAL OR ALAR ODONTOID OR CHECK LIGAMENTS. MIDDLE ODONTOID LIGAMENT OR SUSPENSORY LIGAMENT OF THE AXIS.

The transverse ligament of the atlas has been divided in the middle and the parts have been turned outwards; the dura mater and the posterior or long occipito-axial ligament have been entirely removed.

THE ATLAS AND THE AXIS WITH THE ANTERIOR PORTION OF THE OCCIPITAL BONE SEEN FROM BEHIND, A SECTION HAVING BEEN MADE SIMILAR TO THAT IN THE PREPARATION SHOWN IN FIG. 414.

Articulationes atlanto-occipitalis et atlanto-epistrophica—Occipito-atlantal and atlanto-axial articulations.

Odontoid process—Dens epistrophei

FIG. 421.—OCCIPITO-ATLANTAL AND ATLANTO-AXIAL ARTICULATIONS IN FRONTAL SECTION. LATERAL OR ALAR ODONTOID OR CHECK LIGAMENTS AND MIDDLE ODONTOID LIGAMENT OR SUSPENSORY LIGAMENT OF THE AXIS.

The section passes through the middle of the posterior condylar foramina, and divides the summit of the antero-posterior curve of the occipital condyles.

* See note to p. 177.

FIG. 422.—ARTICULATIONS AND LIGAMENTS OF THE ODONTOID PROCESS. STRATIFORM ARRANGEMENT OF THE CRUCIFORM LIGAMENT OF THE ATLAS, THE POSTERIOR OR LONG OCCIPITO-AXIAL LIGAMENT, AND THE DURA MATER. SPHEENO-OCCIPITAL SYNCHONDROSIS. (MEDIAN SECTION THROUGH THE POSTERIOR PORTION OF THE BASE OF THE CRANIUM AND THE THREE UPPERMOST CERVICAL VERTEBRÆ.)

Articulationes atlanto-occipitalis et atlanto-epistrophica—Occipito-atlantal and atlanto-axial articulations.

FIG. 423.—ANTEP OR OCCIPITO-ATLANTAL LIGAMENT AND PETROBASILAR SYNCHONDROSIS. (THE ATLAS WITH THE POSTERIOR PORTION OF THE BASE OF THE SKULL, SEEN FROM BEFORE.)

FIG. 424.—POSTERIOR OCCIPITO-ATLANTAL LIGAMENT. (THE ATLAS WITH THE OCCIPITAL BONE, SEEN FROM BEHIND.)

Articulatio atlanto-occipitalis—Occipito-atlantal articulation.

FIG. 425.—ARTICULATIO MANDIBULARIS, TEMPOROMANDIBULAR OR TEMPOROMAXILLARY ARTICULATION: LIGAMENTA TEMPOROMANDIBULARE ET STYLOMANDIBULARE. EXTERNAL LATERAL AND STYLOMANDIBULAR OR STYLOMAXILLARY LIGAMENTS. LIGAMENTUM STYLOHYOIDEUM, STYLOHYOID LIGAMENT. (RIGHT TEMPOROMANDIBULAR OR TEMPOROMAXILLARY ARTICULATION, SEEN FROM THE OUTER SIDE.)

Articulatio mandibularis—Temporomandibular or temporomaxillary articulation.

FIG. 426.—ARTICULATIO MANDIBULARIS, TEMPOROMANDIBULAR OR TEMPOROMAXILLARY ARTICULATION: LIGAMENTA SPHENOMANDIBULARE ET STYLOMANDIBULARE, SPHENOMANDIBULAR OR INTERNAL LATERAL AND STYLOMANDIBULAR OR STYLOMAXILLARY LIGAMENT. RELATIONS OF THE STYLOMANDIBULAR OR STYLOMAXILLARY LIGAMENT TO STYLOGLOSSUS AND INTERNAL PTERYGOID MUSCLES. LIGAMENTUM PTERYGOSPINOSUM, PTERYGOSPINOUS LIGAMENT. (THE POSTERIOR PART OF THE FACIAL PORTION OF THE SKULL WITH THE ADJOINING PORTION OF THE BASE OF THE SKULL, DIVIDED SAGITTALLY SOMEWHAT TO THE LEFT OF THE MEDIAN PLANE.)

The basilar and condylar portions of the occipital bone have been removed.

Articulatio mandibularis—Temporomandibular or temporomaxillary articulation.

FIG. 427.—ARTICULATIO MANDIBULARIS, TEMPOROMANDIBULAR OR TEMPOROMAXILLARY ARTICULATION: DISCUS ARTICULARIS, INTERARTICULAR DISC OR MENISCUS, AND LIGAMENTUM STYLOMANDIBULARE, STYLOMANDIBULAR OR STYLOMAXILLARY LIGAMENT. RELATIONS OF THE SUPERIOR HEAD OF THE EXTERNAL PTERYGOID MUSCLE TO THE ANTERIOR WALL OF THE CAPSULAR LIGAMENT AND TO THE INTERARTICULAR DISC OR MENISCUS. (RIGHT TEMPOROMANDIBULAR OR TEMPOROMAXILLARY ARTICULATION, DIVIDED IN A PLANE NEARLY APPROACHING THE SAGITTAL; THE INTERNAL PORTION BEING FIGURED.)

The section runs somewhat obliquely forwards and inwards.

FIG. 428.—FIBROCARILAGO BASALIS; FIBROCARILAGE OF THE FORAMEN LACERUM MEDIUM: SYNCHONDROSES SPHENOPETROSA, PETRO-OCIPITALIS, ET SPHERO-OCIPITALIS; PETROSPHENOIDAL, PETROBASILAR, AND OCCIPITOSPHEROIDAL SYNCHONDROSES: LIGAMENTUM PTERYGOSPINOSUM; PTERYGOSPINOUS LIGAMENT. (THE MIDDLE PORTION OF THE BASE OF THE SKULL, SEEN FROM BELOW.)

Articulatio mandibularis, temporomandibular or temporomaxillary articulation—Synchondroses et ligamenta baseos cranii, synchondroses and ligaments of the base of the skull.

JUNCTURÆ OSSIUM
EXTREMITATUM,
SUPERIORIS ET INFERIORIS

THE ARTICULATIONS
OF THE SUPERIOR AND INFERIOR
EXTREMITIES

FIG. 429.—ARTICULATIO STERNOCLAVICULARIS, STERNOCLAVICULAR ARTICULATION: CAPSULA ARTICULARIS, CAPSULE OF THE JOINT; DISCUS ARTICULARIS, INTERARTICULAR FIBRO-CARTILAGE; LIGAMENTA INTERCLAVICULARE ET COSTOCLAVICULARE, INTERCLAVICULAR AND COSTOCLAVICULAR OR RHOMBOID LIGAMENTS.

The left sternoclavicular articulation has been opened by the removal of the anterior wall of the capsular ligament.

¹ The interarticular fibrocartilage is occasionally defective in the centre, and in rare cases even entirely wanting; there is then only one synovial cavity.—Tr.

FIG. 430.—ARTICULATIO STERNOCLAVICULARIS, STERNOCLAVICULAR ARTICULATION: DISCUS ET CAPSULA ARTICULARIS, INTERARTICULAR FIBRO-CARTILAGE AND CAPSULAR LIGAMENT; CAVUM ARTICULARE, SYNOVIAL CAVITY OR CAVITIES; LIGAMENTA INTERCLAVICULARE ET COSTOCLAVICULARE, INTERCLAVICULAR AND COSTOCLAVICULAR OR RHOMBOID LIGAMENTS.

The right sternoclavicular articulation has been divided by a frontal section; in the left, the capsule has been removed and the clavicle has been drawn backwards.

STERNOCLAVICULAR ARTICULATION, SEEN FROM BEFORE.

Articulationes et ligamenta cinguli extremitatis superioris—Articulations and ligaments of the shoulder-girdle.

FIG. 431.—LIGAMENTA ACROMIOCLAVICULARE ET CORACOCLOVICULARE, SUPERIOR ACROMIOCLAVICULAR
LIGAMENT AND CORACOCLOVICULAR LIGAMENT (CONOID PORTION); LIGAMENTA CORACO-ACROMIALE
ET TRANSVERSUM SCAPULÆ SUPERIUS, CORACO-ACROMIAL AND SUPRASCAPULAR LIGAMENTS.
(THE RIGHT SCAPULA WITH THE ACROMIAL HALF OF THE CLAVICLE AND THE SHOULDER-JOINT,
SEEN FROM ABOVE.)

Articulationes et ligamenta cinguli extremitatis superioris—Articulations and ligaments of the shoulder-girdle.

FIG. 432.—ARTICULATIO HUMERI, SHOULDER-JOINT: CAPSULA ARTICULARIS, CAPSULAR LIGAMENT; LIGAMENTUM CORACOHUMERALE, CORACOHUMERAL LIGAMENT; LIGAMENTUM TRANSVERSUM SCAPULAE SUPERIOR ET LIGAMENTUM TRANSVERSUM SCAPULAE INFERIOR, CORACOSCAPULAR OR SUPRASCAPULAR LIGAMENT AND SPINOGLENOID LIGAMENT. (THE RIGHT SHOULDER-JOINT, SEEN FROM BEHIND.)

The acromion has been sawn off, and the coracoid extremity of the coraco-acromial ligament has been turned upwards.

Articulatio humeri—Shoulder-joint.

FIG. 433.—ARTICULATIO HUMERI, SHOULDER-JOINT: LABRUM GLENOIDALE, GLENOID LIGAMENT; RELATIONS OF THE TENDON OF THE LONG HEAD OF THE BICEPS MUSCLE AND OF THE EPIPHYSIAL DISC TO THE SYNOVIAL CAVITY OF THE ARTICULATION. (THE RIGHT SHOULDER-JOINT IN FRONTAL SECTION; POSTERIOR HALF.)

FIG. 434.—ARTICULATIO ACROMIOCLAVICULARIS, ACROMIOCLAVICULAR JOINT: DISCUS ARTICULARIS, INTERARTICULAR FIBROCARILAGE; LIGAMENTUM ACROMIOCLAVICULARE, SUPERIOR ACROMIOCLAVICULAR LIGAMENT. (THE RIGHT ACROMIOCLAVICULAR ARTICULATION IN FRONTAL SECTION; POSTERIOR PORTION.)

Articulatio humeri—Shoulder-joint. Articulatio acromioclavicularis—Acromioclavicular articulation.

FIG. 435.—ARTICULATIONES HUMERI ET ACROMIOCLAVICULARIS, THE SHOULDER-JOINT AND THE STERNOCLAVICULAR ARTICULATION: BURSA MUSCULI SUBSCAPULARIS ET BURSA SUBDELTOIDEA, BURSA OF THE SUBSCAPULARIS⁴ MUSCLE AND SUBDELTOID OR SUBACROMIAL BURSA; VAGINA MUSCOSA INTERTUBERCULARIS, SYNOVIAL SHEATH OF THE BICIPITAL GROOVE; LIGAMENTA CORACOCALVICULARE, CORACO-ACROMIALE, ET TRANSVERSUM SCAPULÆ SUPERIUS; THE CORACOCALVICULAR (CONOID AND TRAPEZOID), CORACO-ACROMIAL, AND CORACOSCAPHAL OR SUPRASCAPHAL LIGAMENT. (RIGHT SHOULDER-JOINT, INJECTED WITH TALLOW; THE ACROMIAL EXTREMITY OF THE CLAVICLE HAS BEEN DRAWN UPWARDS. SEEN FROM BEFORE.)

Articulatio humeri—Shoulder-joint.

FIG. 436.—ARTICULATIO HUMERI, SHOULDER-JOINT: INTRACAPSULAR PORTION OF THE TENDON OF THE LONG HEAD OF THE BICEPS MUSCLE; LABRUM GLENOIDALE, GLENOID LIGAMENT; APERTURE OF COMMUNICATION OF THE BURSA OF THE SUBSCAPULARIS MUSCLE WITH THE SHOULDER-JOINT; RELATIONS OF THE TENDONS OF THE SCAPULAR MUSCLES WITH THE CAPSULE OF THE SHOULDER-JOINT. (RIGHT SHOULDER-JOINT FROM THE OUTER SIDE)

After tallow had been injected into the joint and allowed to harden, the capsular ligament and the surrounding scapular muscles were divided by a circular incision midway between their attachments to the scapula and the humerus, a strip of the capsule, however, being left undivided, where the tendon of the long head of the biceps muscle passes through the joint. The humerus with the distal half of the capsule has been turned upwards.

¹ Projection of the synovial membrane of the elbow-joint, which membrane, after passing downwards between the vertical articular surface of the head of the radius and the inner surface of the orbicular ligament, forms a circular pouch or sac below this ligament around the neck of the radius.—TR.

FIG. 437.—ARTICULATIO CUBITI, ELBOW-JOINT: CAPSULA ARTICULARIS, CAPSULE OF THE JOINT; LIGAMENTA COLLATERALIA, LATERAL LIGAMENTS; LIGAMENTUM ANNULARE RADII, ANNULAR OR ORBICULAR LIGAMENT OF THE RADIUS; *RECESSUS SACCIFORMIS (*see note above*). (RIGHT ELBOW-JOINT, UNOPENED; ANTERIOR OR PALMAR ASPECT.)

The *recessus sacciformis has been injected with tallow.

Articulatio cubiti—Elbow-joint.

FIG. 438.—*ARTICULATIO CUBITI, ELBOW-JOINT: ATTACHMENT OF THE CAPSULE TO THE ANTERIOR SURFACES OF THE HUMERUS AND THE ULNA (ANTERIOR LIGAMENT); LIGAMENTA COLLATERALIA, LATERAL LIGAMENTS; LIGAMENTUM ANNULARE RADII, ANNULAR OR ORBICULAR LIGAMENT OF THE RADIUS. (RIGHT ELBOW-JOINT; ANTERIOR OR PALMAR ASPECT.)*

The capsule has been divided above and below, close to its attachment to the bones, and between the lateral ligaments (*i.e.*, the greater portion of the anterior ligament has been removed); the cut ends of the anterior ligament have been folded back against the bones. The *recessus sacciformis—see note on previous page—has been removed.

FIG. 439.—ARTICULATIO CUBITI, ELBOW-JOINT: CAPSULA ARTICULARIS, CAPSULE OF THE JOINT; LIGAMENTUM COLLATERALE ULNARE, INTERNAL LATERAL LIGAMENT. (RIGHT ELBOW-JOINT; POSTERO-INTERNAL ASPECT.)

¹ See note to p. 204.

FIG. 440.—ARTICULATIO CUBITI, ELBOW-JOINT: CAPSULA ARTICULARIS, CAPSULE OF THE JOINT; LIGAMENTUM COLLATERALE RADIALE, EXTERNAL LATERAL LIGAMENT; LIGAMENTUM ANNULARE RADII, ANNULAR OR ORBICULAR LIGAMENT OF THE RADIUS; *RECESSUS SACCIFORMIS. (RIGHT ELBOW-JOINT; POSTERO-EXTERNAL ASPECT.)

The *recessus sacciformis has been injected with tallow.

Articulatio cubiti—Elbow-joint.

FIG. 441.—ARTICULATIO HUMERO-ULNARIS, HUMERO-ULNAR ARTICULATION. (SAGITTAL SECTION OF THE RIGHT ELBOW-JOINT; THE RADIAL HALF IS FIGURED.)

The section passes through the trochlea and the great sigmoid cavity of the ulna, in a plane vertical to the axis of the trochlea.

Articulatio cubiti—Elbow-joint.

FIG. 442.—ARTICULATIONES RADIO-ULNARES, PROXIMALIS ET DISTALIS; RADIO-ULNAR ARTICULATIONS, SUPERIOR AND INFERIOR: LIGAMENTUM ANNULARE RADII, ANNULAR OR ORBICULAR LIGAMENT OF THE RADIUS; MEMBRANA INTEROSSEA ANTIBRACHII, INTEROSSEOUS MEMBRANE OR LIGAMENT OF THE FOREARM. (THE BONES OF THE RIGHT FOREARM WITH THE RADIO-ULNAR LIGAMENTS; ANTERIOR OR PALMAR ASPECT.)

Articulations of the Bones of the Forearm.

FIG. 443.—THE POSTERIOR OR DORSAL LIGAMENTS OF THE WRIST AND HAND: LIGAMENTA COLLATERALIA CARPI, LATERAL LIGAMENTS OF THE WRIST AND THE CARPUS; SUPERFICIAL POSTERIOR OR DORSAL LIGAMENTS OF CARPUS AND METACARPUS. THE CAPSULES AND THE LATERAL LIGAMENTS OF THE METACARPO-PHALANGEAL AND INTERPHALANGEAL ARTICULATIONS.

In the articulations of the index finger the posterior wall of the capsule has been divided transversely across the middle of the joint, and the ends have been turned upwards and downwards; in the articulations of the thumb and the middle finger the capsule has been divided only between the lateral ligaments, leaving these intact; in the articulations of the ring and little fingers the capsule has not been opened.

Articulationes manus et digitorum—Articulations of the hand and fingers.

¹ These are fibrous plates rather than ligaments properly so called, and, being thickened into fibrocartilage at each side along their attachments to the lateral metacarpo-phalangeal ligaments, they are grooved on the palmar surfaces for the flexor tendon. Macalister calls them *glenoid* ligaments. It is in the lateral fibrocartilaginous portions of these plates that the sesamoid bones of the metacarpo-phalangeal joint of the thumb, and occasionally of some of the other fingers, are developed. Tr.

FIG. 444.—THE ANTERIOR OR PALMAR LIGAMENTS OF THE WRIST AND HAND: THE SUPERFICIAL LIGAMENTS OF THE CARPUS AND THE METACARPUS; THE CAPSULES AND LIGAMENTS OF THE METACARPO-PHALANGEAL AND THE INTERPHALANGEAL ARTICULATIONS. LIGAMENTUM CARPI TRANSVERSUM, ANTERIOR ANNULAR LIGAMENT OF THE WRIST; CANALIS CARPI, CANAL OF THE CARPUS BENEATH THE ANTERIOR ANNULAR LIGAMENT (FOR THE TRANSMISSION OF THE FLEXOR TENDONS). RELATIONS OF THE TENDONS OF THE FLEXOR CARPI ULNARIS AND FLEXOR CARPI RADIALIS MUSCLE TO THE ANTERIOR OR PALMAR CARPAL AND METACARPAL LIGAMENTS. LIGAMENTA ACCESSORIA VOLARIA, ANTERIOR OR PALMAR METACARPO-PHALANGEAL LIGAMENTS; LIGAMENTA CAPITULORUM TRANSVERSA, TRANSVERSE METACARPAL LIGAMENT (ANTERIOR OR PALMAR DISTAL INTERMETACARPAL LIGAMENTS). OSSA SESAMOIDEA, SESAMOID BONES.

In the interphalangeal articulations of the index finger the anterior portions of the capsules between the lateral ligaments have been entirely cut away; in most of the middle finger the capsules have been divided transversely across the middle of the joint and the divided halves of the anterior ligament turned upwards and downwards; in the interphalangeal articulation of the thumb the anterior portion of the capsule has been divided along its attachment to the distal phalanx and the lateral ligaments, and, together with the sesamoid bones embedded in it on each side, has been turned upwards; in the remaining joints the capsule has been left intact.

FIG. 445.—THE ANTERIOR OR PALMAR LIGAMENTS OF THE RIGHT CARPUS, SHOWN BY THE REMOVAL OF THE ANTERIOR ANNULAR LIGAMENT OF THE WRIST: LIGAMENTUM RADIOCARPEUM VOLARE, ANTERIOR OR PALMAR LIGAMENT OF THE WRIST-JOINT; LIGAMENTUM CARPI RADIATUM, ANTERIOR OR RADIATE LIGAMENT OF THE CARPUS.

FIG. 446.—THE ANTERIOR OR PALMAR LIGAMENTS OF THE INTERCARPAL (TRANSVERSE CARPAL) AND CARPO-METACARPAL ARTICULATIONS, AFTER THE ANTERIOR ANNULAR LIGAMENT OF THE WRIST AND THE TENDONS OF THE FLEXOR CARPI RADIALIS AND FLEXOR CARPI ULNARIS HAVE BEEN ENTIRELY REMOVED.

The radiocarpal and distal radio-ulnar articulations have been opened, the triangular fibrocartilage and the internal lateral ligament of the wrist-joint being left intact; the bones of the forearm have been separate: from one another and from the carpus; the pisiprimal articulation, the carpometacarpal articulations of the thumb and the ring finger, and the second intermetacarpal articulation, have been partially opened.

Articulatio manus—Articulations of the hand.

FIG. 447.—THE SHORT POSTERIOR OR DORSAL LIGAMENTS OF THE TRANSVERSE CARPAL ARTICULATION, AND OF THE CARPOMETACARPAL AND INTERMETACARPAL ARTICULATIONS. (THE RIGHT CARPUS WITH THE DISTAL EXTREMITIES OF THE BONES OF THE FOREARM AND THE PROXIMAL EXTREMITIES OF THE METACARPAL BONES.)

The distal radio-ulnar articulation and the radiocarpal and transverse carpal articulations have been opened by the removal of the posterior ligaments, and the bones of the forearm have been drawn a little upwards and away from the carpus.

FIG. 448.—THE ARTICULAR SURFACES AND THE ATTACHMENTS OF THE CAPSULE OF THE RADIOCARPAL ARTICULATION OR WRIST-JOINT; CANALIS CARPI, THE CANAL OF THE CARPUS. (THE RIGHT CARPUS WITH THE METACARPAL BONES; ANTERO-SUPERIOR ASPECT.)

The distal extremities of the bones of the forearm have been turned to the ulnar side, after division of the capsule of the wrist-joint with the exception of the internal lateral ligament.

FIG. 449.—ARTICULATIO RADIO-ULNARIS DISTALIS, DISTAL RADIO-ULNAR ARTICULATION. ARTICULATIO MANUS, ARTICULATIONS OF THE HAND: ARTICULATIONES RADIOCARPEAE ET INTERCARPEAE, RADIOCARPAL ARTICULATION OR WRIST-JOINT AND TRANSVERSE CARPAL ARTICULATION; ARTICULATIONES CARPOMETACARPEAE, CARPOMETACARPAL ARTICULATIONS; ARTICULATIONES INTERMETACARPEAE, INTERMETACARPAL ARTICULATIONS. ARTICULATIONES METACARPO-PHALANGEAE, METACARPO-PHALANGEAL ARTICULATIONS; ARTICULATIONES DIGITORUM MANUS, INTERPHALANGEAL ARTICULATIONS OF THE FINGERS. (THE SKELETON OF THE RIGHT HAND WITH THE DISTAL EXTREMITIES OF THE BONES OF THE FOREARM; POSTERIOR OR DORSAL ASPECT.)

The articulations are all opened by a section in the frontal plane: and whereas in the fingers this plane passes through the joints from side to side, dividing the lateral ligaments; in the thumb, owing to the opposition of this member, the plane of section passes through the joints in a dorsopalmar direction, and divides the dorsal and palmar ligaments.

Articulationes manus et digitorum—Articulations of the hand and fingers.

¹ See note on p. 211.

FIG. 450.—THE ARTICULATIONS OF THE HAND, SEEN IN SAGITTAL OR DORSOPALMAR SECTION, SHOWING THEIR RELATIONS TO THE EPIPHYSIAL DISCS. (THE RADIAL PORTION OF THE DIVIDED RIGHT HAND OF A YOUTH AGED SEVENTEEN YEARS.)

The section traverses the distal extremity of the radius, the carpus, and the metacarpal bone and phalanges of the middle finger.

Articulationes manus et digitorum—Articulations of the hand and fingers.

¹ Intervertebral disc of the sacrococcygeal articulation.—Tr.

FIG. 451.—ARTICULATIO SACRO-ILIACA, SACRO-ILIAC ARTICULATION: LIGAMENTA SACRO-ILIACA ANTERIORA, ANTERIOR SACRO-ILIAC LIGAMENT; LIGAMENTUM ILIOLUMBALE, ILIOLUMBAR LIGAMENT. LIGAMENTA SACROSPINOSUM ET SACROTUBEROSUM, SMALL OR ANTERIOR AND GREAT OR POSTERIOR SACROSCIATIC LIGAMENTS. FORAMINA ISCHIADICA MAJUS ET MINUS, GREAT AND SMALL SACROSCIATIC FORAMINA. (THE PELVIS WITH THE FOURTH AND FIFTH LUMBAR VERTEBRÆ, IN FRONTAL SECTION; POSTERIOR HALF, SEEN FROM BEFORE.)

The ligaments of the right sacro-iliac articulation are intact; those of the left articulation have been removed.

Articulationes et ligamenta cinguli extremitatis interioris—Articulations and ligaments of the pelvic girdle.

¹ See note on p. 216.

FIG. 452.—LIGAMENTA SACROTUBEROSUM ET SACROSPINOSUM, POSTERIOR OR GREAT AND ANTERIOR OR SMALL SACROSCIATIC LIGAMENTS; FORAMINA ISCHIADICA, SACROSCIATIC FORAMINA. MEMBRANA OBTURATORIA, OBTURATOR MEMBRANE OR LIGAMENT; CANALIS OBTURATORIUS, OBTURATOR CANAL. ARTICULATIO SACRO-ILIACA, SACRO-ILIAC ARTICULATION. (THE RIGHT HALF OF A PELVIS DIVIDED IN THE MEDIAN PLANE; SEEN FROM THE INNER SIDE.)

Articulationes et ligamenta cinguli extremitatis inferioris—Articulations and ligaments of the pelvic girdle.

¹ Macalister and some other English anatomists agree with Toldt in naming this ligament the *interosseous* sacro-iliac ligament. That name is, however, misapplied, since the fibres of the ligament do not connect two closely adjacent parallel articular surfaces. True interosseous fine fibres are occasionally found connecting corresponding parts of the auricular surfaces of the sacrum and the ilium. 16.

FIG. 453.—POSTERIOR LIGAMENTS OF THE PELVIC GIRDL: LIGAMENTUM SACROTUBEROSUM, POSTERIOR OR GREAT SACROSPINOUS LIGAMENT, AND ITS RELATION TO THE POSTERIOR LAYER OF THE LUMBAR APONEUROSIS. FORAMINA ISCHIADICA, SACROSPINOUS FORAMINA. LIGAMENTUM SACRO-ILIACUM POSTERIUS LONGUM, LONG OR OBLIQUE (POSTERIOR) SACRO-ILIAC LIGAMENT; LIGAMENTA SACRO-ILIAC INTEROSSEA, POSTERIOR SACRO-ILIAC LIGAMENT (see note above); LIGAMENTUM ILIOLUMBALE, ILIOLUMBAR LIGAMENT. (THE PELVIS WITH THE FOURTH AND FIFTH LUMBAR VERTEBRE; SEEN FROM BEHIND.)

On the right side, the posterior layer of the lumbar aponeurosis has been divided close to its continuation into the posterior or great sacrospinous ligament, and turned outwards; on the left side, this superficial portion of the posterior or great sacrospinous ligament has been cut across, and the divided ends have been turned upwards and downwards. The lowest fibres of the middle layer of the lumbar aponeurosis have on the right side been left intact, but on the left side entirely removed.

Articulationes et ligamenta cinguli extremitatis inferioris—Articulations and ligaments of the pelvic girdle.

FIG. 454.—DEEP POSTERIOR LIGAMENTS OF THE SACRO-ILIAC ARTICULATION: LIGAMENTA SACRO-ILIACA INTEROSSEA, POSTERIOR SACRO-ILIAC LIGAMENTS; LIGAMENTUM SACRO-ILIACUM POSTERIUS BREVE, POSTERIOR (SHORT) SACRO-ILIAC LIGAMENT. (THE RIGHT HALF OF A PELVIS DIVIDED IN THE MEDIAN PLANE; POSTERO-INTERNAL ASPECT.)

The upper portion of the posterior or great sacrospinous ligament has been removed; the long or oblique (posterior) sacro-iliac ligament has been divided transversely in the middle, and the ends have been turned upwards and downwards.

FIG. 455.—HORIZONTAL SECTION THROUGH THE SACRO-ILIAC ARTICULATION; UPPER SURFACE OF LOWER SEGMENT: SUPERFICIAL AND DEEP LAYERS OF THE POSTERIOR SACRO-ILIAC LIGAMENT.

The section was made in a plane at right angles to the long axis of the pelvis, and passes through the middle of the body of the first sacral vertebra.

¹ This forms part of the posterior sacro-iliac ligament of English anatomists. See note on p. 218.—Tr. ² See note on p. 218.

Articulationes et ligamenta cinguli extremitatis inferioris—Articulations and ligaments of the pelvic girdle.

FIG. 456.—SYMPHYSIS OSSIUM PUBIS, PUBIC SYMPHYSIS: LIGAMENTUM PUBICUS SUPERIUS, SUPERIOR PUBIC LIGAMENT; LIGAMENTUM ARCUATUM PUBIS, INFERIOR PUBIC OR SUBPUBIC LIGAMENT; LIGAMENTUM TRANSVERSUM PELVIS, TRANSVERSE LIGAMENT OF THE PELVIS. THE ORIGIN OF THE TENDONS OF THE RECTUS ABDOMINIS MUSCLES FROM THE PUBIS, AND THE RELATION OF THESE TENDONS TO THE PUBIC SYMPHYSIS. (THE ANTERIOR ASPECT OF THE PUBIC SYMPHYSIS.)

The lower extremities of the rectus abdominis muscles have been pulled a little apart.

Fissure of the interpubic disc

Interpubic disc
Lamina fibrocartilaginea interpubica

Pubis
Os pubis

Interlacing tendons¹

¹ The anterior pubic ligament is not mentioned by the author. It consists of two parts, a superficial and a deep. The deep part, which is not shown in any of the figures, is made up of fibres passing transversely from bone to bone in front of the interpubic disc; the fibres of the superficial part are oblique, interlace freely, and are mainly derived from the tendons of the external oblique and rectus muscles of the abdomen, as well as from those of the superficial adductors of the thigh. These interlacing tendinous fibres of the superficial part of the anterior pubic ligament are shown in both the figures on this page. —Tr.

² This ligament is a portion of the deep perineal fascia or triangular ligament of the urethra. The name of transverse ligament of the pelvis, which is rarely used in England, was given to it by Henle. —Tr.

FIG. 457.—HORIZONTAL SECTION THROUGH THE PUBIC SYMPHYSIS OF A NULLIPAROUS WOMAN AGED TWENTY-ONE YEARS; UPPER SURFACE OF LOWER SEGMENT: LAMINA FIBROCARILAGINEA INTERPUBICA, INTERPUBIC DISC; FISSURE IN THE INTERPUBIC DISC. RE-INFORCEMENT OF THE INTERPUBIC ARTICULATION BY THE INTERLACING ON ITS ANTERIOR SURFACE OF THE FIBRES OF THE TENDONS OF ORIGIN OF THE RECTUS ABDOMINIS MUSCLES AND THE TENDONS OF INSERTION OF THE EXTERNAL OBLIQUE MUSCLES.

The plane of section lies in the upper half of the symphysis.

Symphysis ossium pubis—Pubic symphysis.

¹ The slight posterior prominence of the interpubic disc has not received any name from English anatomists. A few transverse fibres connect the pubic bones in this region, forming the *posterior pubic ligament*, which is not mentioned by Todd.—Tr.

² See note on p. 220.

FIG. 458.—SYMPHYSIS OSSIUM PUBIS, PUBIC SYMPHYSIS: TORUS PUBICUS, POSTERIOR PROMINENCE OF THE INTERPUBIC DISC; LIGAMENTUM TRANSVERSUM PELVIS, TRANSVERSE LIGAMENT OF THE PELVIS (see note ² above), WITH THE VENOUS FORAMINA; CONNEXIONS OF THE TRANSVERSE LIGAMENT OF THE PELVIS WITH THE OBTURATOR FASCIA. (THE PUBIC SYMPHYSIS SEEN FROM BEHIND.)

FIG. 459.—SYMPHYSIS OSSIUM PUBIS, PUBIC SYMPHYSIS: LAMINA FIBROCARILAGINEA INTERPUBICA, INTERPUBIC DISC; LIGAMENTUM PUBICUM SUPERIUS, SUPERIOR PUBIC LIGAMENT; LIGAMENTUM ARCUATUM PUBIS, INFERIOR PUBIC OR SUBPUBIC LIGAMENT. (THE PUBIC SYMPHYSIS IN FRONTAL SECTION; ANTERIOR SURFACE OF POSTERIOR SEGMENT.)

Symphysis ossium pubis—Pubic symphysis.

FIG. 460.—ARTICULATIO COXÆ, THE HIP-JOINT: LIGAMENTUM ILIOFEMORALE, ILIOFEMORAL LIGAMENT, OR Y LIGAMENT OF BIGELOW; LIGAMENTUM PUBOCAPSULARE, PUBOFEMORAL LIGAMENT, AND ITS RELATIONS TO THE OBTURATOR MEMBRANE. (THE RIGHT HIP-JOINT, SEEN FROM BEFORE.)

Articulatio coxæ—The hip-joint.

FIG. 461.—ARTICULATIO COXÆ, THE HIP-JOINT: CAPSULA ARTICULARIS, CAPSULAR LIGAMENT; ZONA ORBICULARIS, THE CIRCULARLY DISPOSED FIBRES OF THE CAPSULAR LIGAMENT, FORMING A BAND ROUND THE NECK OF THE FEMUR, WHICH IS MOST DISTINCT BEHIND AND BELOW. RELATION OF THE EPIPHYSIAL DISC OF THE HEAD OF THE FEMUR TO THE HIP-JOINT. (THE RIGHT HIP-JOINT IN FRONTAL SECTION; ANTERIOR SURFACE OF POSTERIOR SEGMENT.)

The section passes through the middle of the cotyloid notch and of the fossa of the interarticular ligament.

Articulatio coxæ—The hip-joint.

FIG. 463.—ARTICULATIO COXÆ, THE HIP-JOINT: THE ACETABULUM; LABRUM GLENOIDALE ET LIGAMENTUM TRANSVERSUM ACETABULI, THE COTYLOID LIGAMENT AND THE TRANSVERSE LIGAMENT OF THE ACETABULUM. LIGAMENTUM TERES FEMORIS, THE INTERARTICULAR OR ROUND LIGAMENT. (VIEW OF THE INTERIOR OF THE SOCKET OF THE RIGHT HIP-JOINT.)

The interarticular or round ligament has been divided close to its attachment to the head of the femur.

FIG. 464.—ARTICULATIO COXÆ, THE HIP-JOINT: CAPSULA ARTICULARIS, THE CAPSULAR LIGAMENT; ZONA ORBICULARIS (see p. 223), AND ITS RELATIONS TO THE PUBOFEMORAL AND ISCHIOCAPSULAR LIGAMENTS. MEMBRANA OBTURATORIA ET CANALIS OBTURATORIIUS, OBTURATOR MEMBRANE OR LIGAMENT AND OBTURATOR CANAL. (THE POSTERO-INTERNAL SIDE OF THE RIGHT HIP-JOINT SEEN FROM BELOW.)

The articular cavity has been injected with tallow.

Articulatio coxæ—The hip-joint.

FIG. 465.—ARTICULATIO COXÆ, THE HIP-JOINT: ZONA ORBICULARIS (see p. 223), AND ITS RELATIONS TO THE ILIOFEMORAL, PUBOFEMORAL, AND ISCHIOCAPSULAR LIGAMENTS; THE THIN PORTIONS OF THE CAPSULE, AND THE COMMUNICATION BETWEEN THE SYNOVIAL CAVITY AND THE BURSA BENEATH THE TENDON OF THE ILIOPSOAS MUSCLE; THE ACETABULUM, WITH THE TRANSVERSE LIGAMENT OF THE ACETABULUM, AND THE INTERARTICULAR OR ROUND LIGAMENT.

After the capsule of the right hip-joint had been prepared from without, the joint was fully flexed; the capsule was then divided by a section in the direction of a line passing from the middle of the upper border of the great trochanter to the inner border of the iliopectineal eminence; the ilium was cut away except for that portion of the bone which contributes to the formation of the acetabulum, and the neck of the femur was sawn across just internal to the distal attachment of the capsule; the interarticular ligament was divided close to the head of the femur, and this latter, together with the intracapsular portion of the neck, was removed; the acetabulum and the remaining proximal portion of the femur were divided in two by a continuation of the section already made through the upper wall of the capsule; the two halves of the acetabulum and the head and neck of the femur were then opened out till the cut surfaces met at a very obtuse angle; so that the capsule and the cotyloid ligament were fully exposed from within. The synovial membrane was dissected off, and the inner surface of the capsule cleaned from fat and cellular tissue.

FIG. 466.—ARTICULATIO GENU, THE KNEE-JOINT: LIGAMENTUM PATELLÆ, PATELLAR LIGAMENT, OR INFRAPATELLAR TENDON; RETINACULA PATELLÆ, LATERAL PATELLAR LIGAMENTS; LIGAMENTA COLLATERALIA, LATERAL LIGAMENTS OF THE KNEE-JOINT. BURSA SUPRAPATELLARIS, THE BURSA BENEATH THE SUPRAPATELLAR TENDON; THE RELATION OF THE UNUSUALLY LARGE SUBCRUREUS MUSCLE TO THE CAPSULE OF THE KNEE-JOINT. ARTICULATIO TIBIOFIBULARIS, SUPERIOR TIBIOFIBULAR ARTICULATION: LIGAMENTA CAPITULI FIBULÆ, ANTERIOR SUPERIOR TIBIOFIBULAR LIGAMENT. (THE RIGHT KNEE-JOINT FROM BEFORE.)

The synovial cavity has been injected with tallow.

Articulatio genu—The knee-joint.

FIG. 467.—ARTICULATIO GENU, THE KNEE-JOINT: LIGAMENTA COLLATERALIA, LATERAL LIGAMENTS OF THE KNEE-JOINT; LIGAMENTA CRUCIATA, CRUCIAL LIGAMENTS; LIGAMENTUM PATELLÆ, PATELLAR LIGAMENT, OR INFRAPATELLAR TENDON. ARTICULATIO TIBIOFIBULARIS, SUPERIOR TIBIOFIBULAR ARTICULATION: LIGAMENTA CAPITULI FIBULÆ, ANTERIOR SUPERIOR TIBIOFIBULAR LIGAMENT. (THE RIGHT KNEE-JOINT SEEN FROM BEFORE AND WITHOUT.)

The capsular ligament has been removed from the front of the joint between the two lateral ligaments, and the patellar ligament has been turned downwards.

Articulatio genu—The knee-joint.

FIG. 468.—ARTICULATIO GENU, THE KNEE-JOINT: CAPSULA ARTICULARIS, CAPSULAR LIGAMENT, AND THE RELATIONS OF THE SUBCRUREUS MUSCLE TO THIS LIGAMENT; BURSA SUPRAPATELLARIS, THE BURSA BENEATH THE SUPRAPATELLAR TENDON. LIGAMENTUM COLLATERALE TIBIÆ, INTERNAL LATERAL LIGAMENT OF THE KNEE-JOINT; LIGAMENTUM PATELLÆ ET RETINACULUM PATELLÆ MEDIALE, PATELLAR LIGAMENT, OR INFRA-PATELLAR TENDON, AND INTERNAL LATERAL PATELLAR LIGAMENT. RELATIONS OF THE EPIPHYSIAL DISCS TO THE ARTICULATION.

The same preparation as that of Fig. 466, seen from within.

Articulatio genu—The knee-joint.

FIG. 469.—ARTICULATIO GENU, THE KNEE-JOINT: CAPSULA ARTICULARIS, CAPSULAR LIGAMENT; BURSA SUPRAPATELLARIS, AND THE RELATIONS OF THE QUADRICEPS EXTENSOR CRURIS MUSCLE TO THE SAME. LIGAMENTUM COLLATERALE FIBULARE, EXTERNAL LATERAL LIGAMENT OF THE KNEE-JOINT. LIGAMENTUM PATELLAE ET RETINACULUM PATELLAE LATERALE, PATELLAR LIGAMENT, OR INFRAPATELLAR TENDON, AND EXTERNAL LATERAL PATELLAR LIGAMENT; THE RELATIONS OF THE LATTER TO THE PLANTARIS MUSCLE AND TO THE OUTER HEAD OF THE GASTROCNEMIUS MUSCLE. BURSA INFRAPATELLARIS PROFUNDA, BURSA BENEATH THE PATELLAR LIGAMENT. ARTICULATIO TIBIOFIBULARIS, SUPERIOR TIBIOFIBULAR ARTICULATION. LIGAMENTA CAPITULI FIBULAE, ANTERIOR SUPERIOR TIBIOFIBULAR LIGAMENT. RELATIONS OF THE EPIPHYSAL DISCS TO BOTH THE JOINTS.

The same preparation as that of Figs. 466 and 468, seen from the outer side. The synovial cavity of the knee-joint has been opened behind the external lateral ligament of the knee-joint, and the bursa beneath the patellar ligament has also been opened.

FIG. 470.—ARTICULATIO GENU, THE KNEE-JOINT: CAPSULA ARTICULARIS, CAPSULAR LIGAMENT; LIGAMENTUM POPLITEUM OBLIQUUM, POSTERIOR LIGAMENT OF THE KNEE-JOINT; RELATIONS OF THIS LIGAMENT TO THE TENDON OF THE SEMIMEMBRANOSUS MUSCLE AND TO THE OUTER HEAD OF THE GASTROCNEMIUS MUSCLE; LIGAMENTUM POPLITEUM ARCUATUM ET RETINACULUM LIGAMENTI ARCUATI, ARCULATE PORTION OF THE POSTERIOR EXTERNAL LATERAL LIGAMENT (MACALISTER), AND SHORT EXTERNAL LATERAL LIGAMENT (QUAIN), OR DIRECT PORTION OF THE POSTERIOR EXTERNAL LATERAL LIGAMENT (MACALISTER). BURSA MUSCULI GASTROCNEMII MEDIALIS, BURSA BETWEEN THE TENDON OF THE SEMIMEMBRANOSUS MUSCLE AND THE TENDON OF THE INNER HEAD OF THE GASTROCNEMIUS MUSCLE, COMMUNICATING WITH THE KNEE-JOINT AND HAVING A COMMON CAVITY WITH THE BURSA BETWEEN THE TENDON OF THE SEMIMEMBRANOSUS MUSCLE AND THE KNEE-JOINT. BURSA MUSCULI POPLITEI, SYNOVIAL SHEATH OF THE TENDON OF THE POPLITEUS MUSCLE. (THE RIGHT KNEE-JOINT, SEEN FROM BEHIND.)

Articulatio genu—The knee-joint.

FIG. 471.—ARTICULATIO GENU, THE KNEE-JOINT: LIGAMENTA CRUCIATA, CRUCIAL LIGAMENTS, AND LIGAMENTUM MENISCI LATERALIS, POSTERIOR OR ASCENDING BAND OF THE EXTERNAL SEMILUNAR FIBROCARILAGE (ELLIS) OR LIGAMENT OF WRISBERG (MACALISTER). (VARIETY¹). COMMUNICATION OF THE SYNOVIAL SHEATH OF THE TENDON OF THE POPLITEUS MUSCLE WITH THE SYNOVIAL CAVITIES OF THE KNEE-JOINT AND OF THE SUPERIOR TIBIOFIBULAR ARTICULATION. (THE RIGHT KNEE-JOINT, SEEN FROM BEHIND.)

The posterior portion of the capsular ligament, between the external and internal lateral ligaments, has been removed, except for a strip left above close to the femoral attachment, which has been turned upwards; the synovial sheath of the tendon of the popliteus muscle has been opened at its inner side, and its posterior wall has been turned outwards with the tendon and the proximal portion of the popliteus muscle.

¹ Though this band is called a variety by Toldt, it is, in fact, usually present, but its situation varies; inasmuch as it passes, sometimes behind (as here), and sometimes before, and sometimes as a double band behind and before, the posterior or internal crucial ligament. Its strength and thickness varies much, but it is seldom entirely wanting.—TK.

FIG. 472.—ARTICULATIO GENU, THE KNEE-JOINT: PARS SYNOVIALIS CAPSULAE ARTICULARIS, SYNOVIAL MEMBRANE OF THE KNEE-JOINT; PLICA SYNOVIALIS PATELLARIS, MUCOUS LIGAMENT; PLICÆ ALARES, ALAR LIGAMENTS; BURSA SUPRAPATELLARIS, BURSA BENEATH THE SUPRAPATELLAR TENDON. (THE RIGHT KNEE-JOINT, SEEN FROM BEFORE AND WITHOUT.)

The joint was first injected, and the injected material allowed to solidify; the capsule of the joint was then prepared from without, the quadriceps extensor cruris muscle being left intact; subsequently the joint was opened by a section passing through the quadriceps muscle and the upper part of the front of the capsule from one lateral ligament to the other close to the attachment of the capsule to the femur, and the anterior wall of the joint together with the distal portion of the quadriceps extensor muscle was turned down. The quadriceps was divided at a somewhat higher level than the capsule, and the bursa beneath the suprapatellar tendon divided in the frontal plane, to show the communication of this bursa with the joint.

Articulatio genu—The knee-joint.

FIG. 473.—ARTICULATIO GENU, THE KNEE-JOINT: THE SEMILUNAR FIBROCARILAGES, THE ALAR LIGAMENTS, AND THE MUCOUS LIGAMENT; THE INSERTION INTO THE SYNOVIAL MEMBRANE OF THE SUBCRUREUS MUSCLE; THE EXTENSION OF THE SYNOVIAL MEMBRANE OF THE KNEE-JOINT TO FORM THE SYNOVIAL SHEATH OF THE TENDON OF THE POPLITEUS MUSCLE AND THE BURSA BETWEEN THAT MUSCLE AND THE POSTERIOR SURFACE OF THE OUTER TUBEROSITY OF THE TIBIA, AND THE RELATION OF THE SYNOVIAL SHEATH OF THE TENDON TO THE EXTERNAL LATERAL LIGAMENT OF THE KNEE-JOINT. (THE PROXIMAL EXTREMITIES OF THE BONES OF THE LEG WITH THE ANTERIOR WALL OF THE CAPSULE OF THE KNEE-JOINT SEEN FROM BEHIND.)

After the joint had been injected, and the injected material allowed to solidify, the capsule of the joint was prepared from without, the quadriceps extensor cruris muscle being left intact: the lateral ligaments and the tendon of the popliteus muscle were then divided, and the capsule was opened behind and on either side at a higher level than the semilunar cartilages, and was divided in front along the line of its reflection on to the anterior surface of the femur; after division of the cruciate ligaments, the femur was removed.

Articulatio genu—The knee-joint.

FIG. 474.—ARTICULATIO GENU, THE KNEE-JOINT: CRUCIAL LIGAMENTS AND LIGAMENT OF WRISBERG; BURSA BENEATH THE INTERNAL LATERAL LIGAMENT OF THE KNEE-JOINT. (THE RIGHT KNEE-JOINT SEEN FROM THE INNER SIDE.)

The capsule was removed, the patellar ligament and the internal lateral ligament were turned downwards: the femur was divided sagittally through the middle of the intercondylar fossa, and the external condyle placed in the position it occupies during extension of the knee-joint, while the internal condyle was turned backwards and rotated on its median axis to the extent of 180° .

Articulatio genu—The knee-joint.

Transverse ligament of the semilunar
fibrocartilages
Lig. transversum genu

Anterior attachment of the internal
semilunar fibrocartilage

External tubercle of the
spine of the tibia
Tuberculum intercondyloideum
laterale
Capsular ligament
Capsula articularis
External
semilunar fibrocartilage
Meniscus lateralis
Synovial sheath of the tendon
of the popliteus muscle
Bursa m. poplitei
Anterior superior tibiofibular
ligament
Ligg. capituli fibulae
Styloid process of the fibula
Apex capituli fibulae

Anterior or external
crucial ligament
Lig. cruciatum anterius

Internal semilunar
fibrocartilage
Meniscus medialis

Ligament of Wrisberg¹
Lig. menisci lateralis (Roberti)

¹ See note to p. 233.

FIG. 475.—THE DISTAL ARTICULAR SURFACES OF THE KNEE-JOINT: THE INTERARTICULAR SEMILUNAR FIBROCARILAGES, AND THE EXTENSION OF THE ANTERIOR EXTREMITIES OF THESE IN THE TRANSVERSE LIGAMENT OF THE SEMILUNAR FIBROCARILAGES. THE COMMUNICATION BETWEEN THE SUPERIOR TIBIOFIBULAR ARTICULATION AND THE SYNOVIAL SHEATH OF THE TENDON OF THE POPLITEUS MUSCLE. (THE PROXIMAL EXTREMITY OF THE LEFT TIBIA WITH THE SEMILUNAR FIBROCARILAGES AND THE HEAD OF THE FIBULA, SEEN FROM ABOVE.)

The capsule was divided above (proximal to) the semilunar fibrocartilages, the crucial ligaments were cut across, and the femur was removed.

FIG. 476.—THE PROXIMAL ARTICULAR SURFACES OF THE KNEE-JOINT, AND THE FEMORAL ATTACHMENTS OF THE CRUCIAL LIGAMENTS.

The distal extremity of the femur removed from the preparation shown in Fig. 475, seen from below.

Articulatio genu—The knee-joint.

FIG. 477.—ARTICULATIO TIBIOFIBULARIS ET SYNDESMOSIS TIBIOFIBULARIS, THE TIBIOFIBULAR ARTICULATIONS. THE INTEROSSEOUS MEMBRANE, OR LIGAMENT, OF THE RIGHT LEG; THE ANTERIOR SUPERIOR AND ANTERIOR INFERIOR TIBIOFIBULAR LIGAMENTS. (SEEN FROM BEFORE.)

The Tibiofibular Articulations.

FIG. 478.—ARTICULATIONES TALOCRURALIS ET TALOCALCANEAE, THE ANKLE-JOINT AND THE ASTRAGALOCALCANEAL ARTICULATION; SYNDESMOSIS TIBIOFIBULARIS, INFERIOR TIBIOFIBULAR ARTICULATION, SHOWING THE INTEROSSEOUS LIGAMENT. THE RELATIONS OF THE DISTAL EPIPHYSIAL DISCS OF THE TIBIA AND THE FIBULA TO THE ANKLE-JOINT. (THE ANKLE-JOINT AND THE ASTRAGALOCALCANEAL ARTICULATION OF THE RIGHT LEG, DIVIDED IN A VERTICAL PLANE CLOSELY APPROXIMATING THE FRONTAL PLANE; POSTERIOR SURFACE OF THE ANTERIOR SEGMENT.)

The plane of section passes through the lowermost parts of the lateral portions of the superior articular surface of the astragalus.

FIG. 479.—THE CAPSULAR LIGAMENTS AND THE SUPERFICIAL POSTERIOR LIGAMENTS OF THE ANKLE-JOINT AND OF THE ASTRAGALOCALCANEAL ARTICULATION. THE RELATIONS OF THE DISTAL EPIPHYSAL DISCS OF THE TIBIA AND FIBULA TO THE ANKLE-JOINT. THE POSTERIOR PORTIONS OF THE LATERAL LIGAMENTS OF THE ANKLE-JOINT: LIGAMENTUM TALOTIBIALE POSTERIUS, THE POSTERIOR ASTRAGALOTIBIAL PORTION OF THE INTERNAL LATERAL, OR DELTOID, LIGAMENT OF THE ANKLE-JOINT; LIGAMENTUM CALCANEOTIBIALE, THE CALCANEOTIBIAL PORTION OF THE INTERNAL LATERAL, OR DELTOID, LIGAMENT OF THE ANKLE-JOINT; LIGAMENTUM CALCANEOFIBULARE, THE MIDDLE BAND, OR CALCANEOFIBULAR PORTION, OF THE EXTERNAL LATERAL LIGAMENT OF THE ANKLE-JOINT. LIGAMENTUM TALOCALCANEUM POSTERIUS, THE POSTERIOR ASTRAGALOCALCANEAL LIGAMENT.—SYNDESMOSIS TIBIOFIBULARIS: LIGAMENTUM MALLEOLI LATERALIS POSTERIUS, THE POSTERIOR INFERIOR TIBIOFIBULAR LIGAMENT. (THE ANKLE-JOINT AND THE ASTRAGALOCALCANEAL ARTICULATION OF THE RIGHT LEG, SEEN FROM BEHIND.)

FIG. 480.—ARTICULATIONES TALOCRURALIS ET TALOCALCANEAE, THE ANKLE-JOINT AND THE ASTRAGALOCALCANEAL ARTICULATION; SYNDESMOSIS TIBIOFIBULARIS, THE INFERIOR TIBIOFIBULAR ARTICULATION; LIGAMENTUM MALLEOLI LATERALIS POSTERIUS, THE POSTERIOR INFERIOR TIBIOFIBULAR LIGAMENT. THE POSTERIOR PORTIONS OF THE LATERAL LIGAMENTS: LIGAMENTUM TALOTIBIALE POSTERIUS, THE POSTERIOR ASTRAGALOTIBIAL PORTION OF THE INTERNAL LATERAL, OR DELTOID, LIGAMENT OF THE ANKLE-JOINT; LIGAMENTUM CALCANEOTIBIALE, THE CALCANEOTIBIAL PORTION OF THE INTERNAL LATERAL, OR DELTOID, LIGAMENT OF THE ANKLE-JOINT; LIGAMENTUM TALOFIBULARE POSTERIUS, THE POSTERIOR BAND, OR ASTRAGALOFIBULAR PORTION, OF THE EXTERNAL LATERAL LIGAMENT OF THE ANKLE-JOINT; LIGAMENTUM CALCANEOFIBULARE, THE MIDDLE BAND, OR CALCANEOFIBULAR PORTION, OF THE EXTERNAL LATERAL LIGAMENT OF THE ANKLE-JOINT. (THE ANKLE-JOINT AND THE ASTRAGALOCALCANEAL ARTICULATION OF THE RIGHT LEG, SEEN FROM BEHIND.)

The thin posterior portion of the capsule of the ankle-joint has been removed. The posterior portion of the capsule of the astragalocalcaneal articulation has been separated from its attachment to the astragalus external to its attachment to the posterior process of that bone, and turned backwards on to the os calcis.

FIG. 481.—THE SUPERFICIAL INTERNAL LIGAMENTS OF THE ANKLE-JOINT AND OF THE ASTRAGALOCALCANEAL ARTICULATION, AND THE RELATION OF THE DISTAL EPIPHYSAL DISC OF THE TIBIA TO THE ANKLE-JOINT. (THE RIGHT TARSA, WITH THE ADJOINING PORTIONS OF THE TIBIA AND FIBULA AND OF THE FIRST METATARSAL BONE; SEEN FROM THE INNER SIDE.)

FIG. 482.—THE DEEP INTERNAL LIGAMENTS OF THE ANKLE-JOINT AND OF THE ASTRAGALOCALCANEAL ARTICULATION. (THE RIGHT TARSUS, WITH THE ADJOINING PORTIONS OF THE TIBIA AND FIBULA AND OF THE FIRST METATARSAL BONE; SEEN FROM THE INNER SIDE.)

The deltoid ligament (internal lateral ligament of the ankle-joint) has been cut across the middle, and the divided ends have been turned up and down. The capsular ligaments of the ankle-joint and of the astragalocalcaneal articulation have been removed, except for the deeper special bands.

FIG. 483.—ARTICULATIO TALOCRURALIS, THE ANKLE-JOINT; SYNDESMOSIS TIBIOFIBULARIS, THE INFERIOR TIBIOFIBULAR ARTICULATION. THE RELATIONS OF THE DISTAL EPIPHYSAL DISCS OF THE TIBIA AND FIBULA TO THE ANKLE-JOINT. ARTICULATIONES INTERTARSEÆ ET TARSO-METATARSEÆ, THE INTERTARSAL AND TARSO-METATARSAL ARTICULATIONS; ARTICULATIONES INTERMETATARSEÆ, THE INTERMETATARSAL ARTICULATIONS. LIGAMENTA TARSII DORSALIA ET TARSO-METATARSEÆ DORSALIA ET LIGAMENTA BASII DORSALIA; THE DORSAL LIGAMENTS OF THE TARSUS, THE DORSAL TARSO-METATARSAL LIGAMENTS, AND THE DORSAL PROXIMAL INTERMETATARSAL LIGAMENTS. (THE RIGHT TARSUS WITH THE ADJOINING PORTIONS OF THE TIBIA AND FIBULA AND OF THE METATARSUS; DORSO-EXTERNAL ASPECT.)

The joints are unopened except for the astragalocalcaneal, astragalonavicular, and naviculocuneiform articulations, which have been partly opened.

Articulationes pedis—The articulations of the foot.

¹ The strongest part of this ligament is situate external, and not superior, to the calcaneocuboid articulation; it would therefore be more appropriately named the *external calcaneocuboid* ligament.—Tr.

² In the English nomenclature these are regarded as two distinct ligaments, not, as by Toldt, as two parts of a single ligament.—Tr.

FIG. 484.—ARTICULATIONES TALOCRURALIS ET TALOCALCANEÆ, THE ANKLE-JOINT AND THE ASTRAGALOCALCANEAL ARTICULATION; LIGAMENTUM MALLEOLI LATERALIS ANTERIUS, ANTERIOR INFERIOR TIBIOFIBULAR LIGAMENT. THE ANTERIOR AND MIDDLE BANDS OF THE EXTERNAL LATERAL LIGAMENT OF THE ANKLE-JOINT (LIGAMENTUM TALOFIBULARE ANTERIUS ET LIGAMENTUM CALCANEOFIBULARE); LIGAMENTA TALOCALCANEÆ, LATERALE. *ANTERIUS, ET INTEROSSEUM: THE EXTERNAL, *ANTERIUS, AND INTEROSSEOUS ASTRAGALOCALCANEAL LIGAMENTS.—ARTICULATIONES INTERTARSEÆ ET TARSOMETARSEÆ, THE INTERTARSAL AND TARSOMETATARSAL ARTICULATIONS; ARTICULATIONES INTERMETARSEÆ, THE INTERMETATARSAL ARTICULATIONS: LIGAMENTA TARSII DORSALIA, LIGAMENTA TARSOMETATARSEÆ DORSALIA, ET LIGAMENTA BASIUM DORSALIA, THE DEEPER DORSAL TARSAL AND TARSOMETATARSAL LIGAMENTS, AND THE DORSAL PROXIMAL INTERMETATARSAL LIGAMENTS. (THE RIGHT TARSUS, WITH THE ADJOINING PORTIONS OF THE TIBIA AND FIBULA AND OF THE METATARSAL BONES; DORSO-EXTERNAL ASPECT.)

The anterior ligament of the ankle-joint and the lateral portion of the capsule of the astragalocalcaneal articulation have been removed.

Articulationes pedis The articulations of the foot.

FIG. 485.—THE SUPERFICIAL LIGAMENTS OF THE PLANTAR SURFACE OF THE RIGHT FOOT AND THE RELATIONS OF THE TENDONS OF THE TIBIALIS ANTICUS AND POSTICUS AND OF THE PERONEUS LONGUS MUSCLES TO THESE LIGAMENTS. LIGAMENTA TARSII ET TAROMETATARSEÆ, THE TARSAL AND THE TAROMETATARSAL LIGAMENTS; LIGAMENTA BASIUM PLANTARIA, LIGAMENTA CAPITULORUM TRANSVERSA, ET LIGAMENTA ACCESSORIA PLANTARIA; THE PLANTAR PROXIMAL INTERMETATARSAL LIGAMENTS, THE TRANSVERSE METATARSAL LIGAMENT, AND THE INFERIOR METATARSOPHALANGEAL LIGAMENTS (see note above); THE RELATIONS OF THESE LIGAMENTS TO THE DIGITAL PROCESSES OF THE PLANTAR FASCIA.

Articulationes pedis—The articulations of the foot.

FIG. 485.—THE DEEP LIGAMENTS OF THE SOLE OF THE FOOT AND THE SESAMOID BONES OF THE METATARSOPHALANGEAL ARTICULATION OF THE GREAT TOE.

In the preparation shown in Fig. 485 the tendons of the muscles were removed, also the long plantar ligament except for its posterior extremity, and the transverse metatarsal ligament and the digital processes of the plantar fascia were cut away. Some of the metatarsophalangeal and interphalangeal articulations have been opened; others have been left intact.

Articulationes pedis—The articulations of the foot.

FIG. 487.—THE ARTICULAR SURFACES OF THE ASTRAGALOCALCANEONAVICULAR ARTICULATION, THE DEEP LIGAMENTS OF THE DORSUM OF THE FOOT, THE METATARSOPHALANGEAL AND INTERPHALANGEAL ARTICULATIONS OF THE TOES. (THE RIGHT FOOT SEEN FROM THE DORSAL SIDE.)

The capsule of the astragalocalcaneal articulation was removed, with the exception of the internal astragalocalcaneal ligament; the capsule of the astragalonavicular articulation was also removed with the exception of its internal and plantar walls; and, after removing the interosseous astragalocalcaneal ligament, the astragalus was rotated inwards (on the internal astragalocalcaneal ligament as a hinge), until its inferior surface looked directly upwards.

¹ See note 1 to p. 245.

² Known also as the *midtarsal* or *mid-tarsal* joint. It is through this joint (the two parts of which are, however, entirely separate articulations) that the foot is divided in Chopart's amputation.

³ It is through the tarsometatarsal articulations that the foot is divided in Lisfranc's amputation.

FIG. 483.—ARTICULATIONES INTERTARSEÆ ET TAROMETATARSEÆ, THE INTERTARSAL AND TAROMETATARSAL ARTICULATIONS, SHOWING CHOPART'S (MIDTARSAL) LINE, AND LISFRANC'S OR HEY'S TAROMETATARSAL LINE: LIGAMENTA TARSII INTEROSSEA ET LIGAMENTA CUNEO-METATARSEÆ INTEROSSEA, THE INTEROSSEOUS LIGAMENTS OF THE TARSUS, AND THE INTEROSSEOUS METATARSOCUNEIFORM LIGAMENTS. ARTICULATIONES INTERTARSEÆ, THE INTERTARSAL ARTICULATIONS. ARTICULATIONES METATARSOPHALANGEÆ, THE METATARSOPHALANGEAL ARTICULATIONS. ARTICULATIONES DIGITORUM PEDIS, THE INTERPHALANGEAL ARTICULATIONS OF THE TOES. THE RELATIONS OF THE EPIPHYSIAL DISCS OF THE METATARSAL BONES AND OF THE PHALANGES OF THE TOES TO THE RESPECTIVE ARTICULATIONS. (HORIZONTAL SECTION THROUGH THE ARTICULATIONS OF THE RIGHT FOOT OF A YOUTH AGED SEVENTEEN YEARS. SUPERIOR SURFACE OF THE LOWER SEGMENT.)

FIG. 489.—THE ARTICULATIONS OF THE RIGHT FOOT OF A YOUTH AGED SEVENTEEN YEARS, SEEN IN SAGITTAL SECTION, AND SHOWING THE RELATIONS OF THESE ARTICULATIONS TO THE EPIPHYSIAL DISCS.

The section passes through the distal extremity of the tibia, the astragalus, the os calcis, the middle cuneiform bone, the second metatarsal bone, and the phalanges of the second toe.

Articulationes pedis—The articulations of the foot.

INDEX

TO THE ARTHROLOGY

Certain names in this Index have an asterisk (*) prefixed; these, as more fully explained in the Translator's Preface, being terms that form part of the English nomenclature used in this work, but which are not commonly employed by English anatomists. To other names a dagger (†) is prefixed; these are Latin names used by the author in the original work, but not included in the official nomenclature of the "Anatomische Gesellschaft."

A.

- ACETABULUM, the, 225, 227
transverse ligament of the, 168, 223, 225, 227
- †Amphiarthroses: intercarpeæ, 169
- Amphiarthrosis, 169
- Ankle-joint, the, 239, 241, 243, 245
- Annulus fibrosus (fibrocartilagineus intervertebralis), 173, 174
- Aponeurosis (see also "Fascia"):
intercostal, anterior or external, 186
posterior or internal, 176, 183, 184, 187
lumbar, see "Fascia, lumbar"
plantar, 246
plantaris, 246
- Arthrodia, 168
- Articular facet of the clavicle, sternal, 198
surface of the carpus, radial, 213
- Articulatio *vel* articulationes:
acromioclavicularis, 201, 202
atlanto-epitrochlearis, 190-192
atlanto-occipitalis, 167, 190-193
calcaneocuboides, 249
capitulum (costarum), 182-184, 185
carpometacarpea, 167, 212-215
pollicis, 167, 212-214
cochlearis, 166
composita, 169
costotransversariae, 182
costovertebrales, 182-184
coxae, 168, 222-227
cubiti, 169, 204-208
cuneonavicularis, 244, 249
digitorum (manus), 166, 210, 211, 214, 215
pedis, 249, 250
ellipsoidea, 167
et ligamenta capitis, 190-196
cinguli extremitatis inferioris, 216-219
cinguli extremitatis superioris, 198, 199
genu, 228-237
hallucis, 246-248
humeri, 168, 200-203
humero-radialis, 169
humero-ulnaris, 208
intercarpea, 169, 212-215
interchondrales, 186, 188
intermetacarpeæ, 169, 212-214
intermetatarsæ, 244, 245, 249
intertarsæ, 244, 245, 249
intervertebrales, 173-175, 185
mandibularis, 194-196
manus, 169, 210-214
metacarpophalangeæ, 166, 167, 210, 211, 214, 215
metacarpophalangea pollicis, 210, 214
metatarsophalangeæ, 249, 250
ossis pisiformis, 212
pedis, 239, 250
pollicis, 210, 211, 214
radiocarpea, 169, 212-215
radio-ulnaris distalis, 209-214
proximalis, 169, 209

Articulatio *vel* articulationes:

- sacro-iliaca, 216, 217, 219
sellaris, 167, 169
simplex, 166-168
sphæroidea, 168
sternoclavicularis, 170, 185-187, 198
sternocostales, 186-188
talocalcanea, 239, 241, 244, 245, 249, 250
talocalcaneonavicularis, 250; articular surfaces of, 248
talocruralis, 239, 241, 243-245
talonavicularis, 241, 249
tarsi transversa (Chopart's), 249
tarso-metatarsæ, 244, 245, 249, 250
tibiofibularis, 228, 229, 231, 233, 238
trochoidea, 169
- Articulation or articulations (see also "Joint"):
acromio-clavicular, 201, 202
and ligaments of the head, 190-196
of the pelvic girdle, 216-219
of the shoulder-girdle, 195-199
astragalocalcaneal, 239, 241, 244, 245, 249, 250
astragalocalcaneonavicular, 250; articular surfaces of, 248
astragalonavicular, 244, 249
atlanto-axial, 190-192
calcaneocuboid, 249
carpal, transverse, 169, 212-215
carpometacarpal, 167, 212-215
of the thumb, 116, 212-214
chondrosternal, 186-188
continuous, 164, 165
costo-central, 182-185
costotransverse, 182
discontinuous, 166-169
of the fingers, 166, 210, 211, 214, 215
of the foot, 239, 250
of the great toe, 246-248
of the hand, 169, 210-214
of the hip, 168, 222-227
humero-radial, 169
humero-ulnar, 208
intercarpal, 169, 212-215
interchondral, 186, 188
intermetacarpal, 169, 212-214
intermetatarsal, 244, 245, 249
interphalangeal, of the fingers, 166, 210, 211, 214, 215
of the thumb, 210, 211, 214
of the toes, 249, 250
intertarsal, 244, 245, 249
intervertebral, 173-175, 185
of the knee, 228-237
of the lower extremity, 216-250
mediotarsal (Chopart's), 249
metacarpophalangeal, 166, 167, 210, 211, 214, 215
of the thumb, 210, 214
metatarsophalangeal, 249, 250
naviculocuneiform, 244, 249
occipito-atlantal, 167, 190-193
pisopyramidal, 212

Articulation or articulations:

- radiocarpal, 169, 212-215
- radio-ulnar, distal, 209-214
 - proximal, 169, 209
- sacrococcygeal, 180, 216, 217
- sacro-iliac, 216, 217, 219
- of the shoulder, 168, 200-203
- sternoclavicular, 170, 185-187, 198
- tarsometatarsal, 244, 245, 249, 250
- temporomandibular, 194-196
- temporomaxillary, 194-198
- of the thumb, 210, 211, 214
- tibiofibular, inferior, 238-241, 244, 245
 - superior, 228, 229, 231, 233, 238
- of the trunk, 171-187
- of the upper extremity, 198-215

B.

- Bicipital groove, synovial sheath of the, 201, 202
- Bigelow, Y ligament of, 222, 224, 226, 227
- Bones, sesamoid, of the foot, 246, 247, 250
 - of the hand, 210, 211, 214
 - of the knee, 232, and note, p. 211
- Bursa beneath the internal lateral ligament of the knee-joint, 236
 - beneath the patellar ligament, 231
 - beneath the suprapatellar tendon, 228, 230, 231, 234, 235
 - between the tendon of the semimembranosus muscle and the knee-joint, 232
 - between the tendon of the semimembranosus muscle and the tendon of the inner head of the gastrocnemius muscle, 232
 - iliopectineal, 227
 - of the odontoid process, 192
 - of the subscapularis muscle, 202, 203
 - subacromial, 202, 203
 - subdeltoid, 202, 203
 - of the tendon of the popliteus muscle, 232, 233, 235, 237
- † Bursa apicis dentis (epistrophei), 192
- ‡ iliopectinea, 227
- † infrapatellaris profunda, 231
- † ligamenti collateralis tibialis, 236
- † muscoli gastrocnemii medialis, 232
- † poplitei, 232, 233, 235, 237
- † semimembranosi, 232
- † subscapularis, 202, 203
- † subdeltoides, 202, 203
- † suprapatellaris, 228, 230, 231, 234, 235

C.

- Canal of the carpus, 211, 213
 - neural, 177
- obturator, 217, 222, 224, 226
- Canalis carpi, 211, 213
 - obturatorius, 177, 222, 224, 226
 - vertebralis, 177
- Capsula *vel* capsulae:
 - articularis, 166-168, 170
 - pars fibrosa, 168
 - synovialis, 168
 - articulationis acromioclavicularis, 201
 - atlanto-epistropheae, 181, 190
 - atlanto-occipitalis, 167, 181, 190, 193
 - capituli, 185
 - carpometa-carpeae pollicis, 210-213
 - costotransversariae, 185
 - coxae, 168, 222, 223, 226, 227
 - pars fibrosa, 224
 - synovialis, 224
 - cubiti, 204-207
 - pars fibrosa, 208
 - synovialis, 208
 - genu, 228, 230-233, 236, 237
 - pars synovialis, 234, 235
 - hallucis, 246-248

Capsula *vel* capsulae:

- articulationis humeri, 199, 200, 203
 - pars fibrosa, 201
 - synovialis, 201, 203
 - mandibularis, 194, 196
 - manus, 214
 - ossis pisiformis, 212
 - radiocarpeae, 213
 - radio-ulnaris distalis, 209, 212
 - sternoclavicularis, 198
 - talocalcaneae, 239-241
 - talocruralis, 240-242
 - pars fibrosa, 250
 - talonavicularis, 242
 - articulationum digitorum (manus), 166, 210, 211
 - pedis, 246-248
 - intervertebralia, 173, 175, 181, 184, 185
 - metacarpophalangearum, 166, 167, 210, 211
 - metatarsophalangearum, 247, 248
 - tarsometatarsarum, 247
- Capsule or capsules (see also "Ligament, capsular"):
- of the acromioclavicular joint, 201
 - of the articulations of the great toe, 246-248
 - of the astragalonavicular articulation, 242
 - atlanto-axial, 181, 190
 - of the carpometacarpal joint of the thumb, 210-213
 - of the costocentral articulations, 185
 - of the costotransverse articulations, 185
 - of the hip-joint, fibrous portion, 168
 - synovial portion, 168
 - of the interphalangeal articulations of the fingers, 166, 210, 211
 - of the interphalangeal articulations of the toes, 246-248
 - of the inferior radio-ulnar articulation, 209-212
 - of the joints of the articular processes of the vertebrae, 173, 175, 181, 184, 185
 - of the metacarpophalangeal articulations, 166, 167, 210, 211
 - of the metatarsophalangeal articulations, 247, 248
 - occipito-atlantal, 167, 181, 190, 193
 - of the pisipyramid articulation, 212
 - of the radiocarpal articulation, 213
 - of the sternoclavicular joint, 198
 - of the tarsometatarsal articulations, 247
 - of the transverse carpal articulation, 214
 - of the wrist-joint, 213

Cartilage, articular, 168

hyaline, of the vertebral bodies, 174

Cartilago articularis, 168

Cavitas glenoidalis scapulae, 201, 203

Cavity, sigmoid, of the radius, 212

great, of the ulna, 208, 209

synovial, 170

Cavum articulare, 170

Chorda obliqua, 209

Communication between the radiocarpal and the pisipyramid articulation, 212, 213

Communication between the wrist-joint and the pisipyramid articulation, 212, 213

Condylarthrosis, 167

Condyles of the femur, 237

Condylus femoris, 237

Continuous articulation, 164, 165

Cooper's ligament, 221, 224

Cotyloid notch, 223-225

D.

Diarthrosis, 166-169

Discontinuous articulation, 166-169

Disc or discs, epiphyseal:

of the femur, distal, 230

proximal, 223

of the fibula, distal, 239, 240, 244

proximal, 231

of the humerus, proximal, 201

of the metatarsal bones, 249, 250

of the os calcis, 240

- Disc or discs, epiphysial :
 of the phalanges of the fingers, 215
 of the middle finger, 215
 of the toes, 249
 of the radius, distal, 215
 of the third metacarpal bone (distal extremity), 215
 of the tibia, distal, 239, 240, 242, 244, 250
 proximal, 230, 231
- Disc, interpubic, 217, 220, 221
- Discs, intervertebral, 172-174, 177, 178, 182, 183, 185
 of the sacrococcygeal articulation,
 216, 217, 180
- Discus articularis, 170
 articulationis acromioclavicularis, 201
 mandibularis, 196
 radio-ulnaris distalis, 212-214
 sternoclavicularis, 160, 188, 198
- Dura mater encephali, 165, 190, 192
 spinalis, 179
- E.**
- Elbow-joint, 169, 204-208
- Enarthrosis, 168
- F.**
- Facies articularis acromii, (tal)
 calcanea, 201
 media, 248
 posterior, 248
 (calcanei) anterior, 248
 media, 248
 posterior, 248
 capituli costæ, 182
 carpæ (radii), 213
 (dentis epistrophei) anterior, 192
 posterior, 192
 fossæ mandibularis, 196
 patellæ, 229, 234, 235
 posterior (ossis navicularis pedis), 248
 sternalis (claviculæ), 198
 auricularis (ossis ilium), 219
 lunata (acetabuli), 168, 224, 225, 227
 patellaris (femoris), 229, 234, 237
- Falciform process of the great sacrosciatic ligament, 217
- Falx inguinalis, 220
- † Fascia lumbalis, 176
 lumbar, 176, 216, 218
 lumbodorsalis, 176, 218
 obturator, 221
 perineal, deep, 220, 221, and note, p. 221
 plantar, 246
 triangular, 220
- Fibrocartilage, interarticular, 170
 of the acromioclavicular
 joint, 201
 of the inferior radio-ulnar
 articulation, 212-214
 of the sternoclavicular joint,
 170, 188, 198
 of the temporomandibular
 articulation, 196
- triangular, 212-214
- Fibrocartilages, semilunar, 229, 233, 235-237
 posterior or ascending band of
 the external, 233, 235-237
 transverse ligament of the, 229,
 237
- Fibrocartilaginous middle portion of the internal calcaneo-
 navicular ligament, 242, 243, 247, 248, and note, p. 248
- Fibrocartilago basalis, 196
- Fibrocartilagine intervertebrales, 172-174, 177, 178, 182,
 183, 185
- Fibrocartilago navicularis, 242, 243, 247, 248, and note,
 p. 248
- Filum terminale, 179
- Fissure of the interpubic disc, 220
- Fold of the synovial membrane of the wrist-joint, 213
- Foramen ischiadicum majus, 176, 216-218
 minus, 176, 216-218

- Foramen, sacrosciatic, great, 176, 216-218
 small, 176, 216-218
- Fossa, glenoid, of the scapula, 201, 203
 of the interarticular ligament, 223
- Fovea articularis superior (atlantis), 167, 190
 capitis femoris, 223
 dentis, 190
- Foveæ costales (corporum vertebrarum), 183
 transversales, 175, 178, 185

G.

- Ginglymus, cochleoid, 166
 screw, 166
- Gliding joints, 169
 intercarpal, 169
- Gomphosis, 165
- *Groove, preauricular, 216

H.

- Harmonia, 164
- Hip-joint, 168, 222-227

I.

- Incisura clavicularis (sterni), 198
 scapulae, 199, 200
 semilunaris (ulnae), 208, 209
 ulnaris (radii), 212
- Interlacing tendons in front of the pubic symphysis, see
 "Ligament, pubic, anterior"
- Interpubic disc, 217, 220, 221
- Intervertebral discs, see "Discs, intervertebral"

J.

- Joint or joints, see also "Articulation" :
 ball-and-socket, 168
 compound, 169
 condyloid, 167
 elbow, 169, 204-208
 hinge, 166, 169
 knee, 228-237
 mediotarsal, 249
 pivot, 169
 saddle, 167, 169
 simple, 166-168
 trochoid, 169
- Juncturæ ossium extremitatis inferioris, 216-250
 superioris, 198-215
- trunci, 171-187

K.

- Knee-joint, 228-237

L.

- Labrum glenoidale, 168
 articulationis coxæ, 223-225, 227
 humeri, 201, 203
- Lamina fibrocartilaginea interpubica, 217, 220, 221
- Ligament or ligaments:
 acromioclavicular, superior, 199, 201
 alar, 234, 235
 of the ankle-joint :
 anterior, 242, 244, 250
 astragalofibular, anterior, 244, 245
 posterior, 239, 241
 astragalonavicular, 242-245
 astragalotibial, anterior, 243
 posterior, 240, 241, 243
 calcaneofibular, 239-241, 244, 245
 calcaneotibial, 239-242
 capsular, 240, 242, 250
 deltoid, 239-245, 248, 249
 lateral, external, 239-241, 244, 245
 internal, 239-245, 248, 249
 posterior, 240
 tibionavicular, 242, 244, 245, 248
 annular, of radius, 169, 204, 205, 207-209

Ligament or ligaments:

- astragalocalcaneal, *anterior, 244, 245, 250
 - capsular, 239-241
 - external, 245
 - internal, 243, 248
 - interosseous, 239, 244, 245, 249, 250
 - posterior, 240, 242
- of the atlas, cruciform, 191, 192
 - transverse, 190-192
- of the base of the skull, 196
- of Bigelow, 222, 224, 226, 227
- calcaneocuboid, dorsal, 244, 245, 248
 - inferior, 242, 243, 245-247, 250
 - plantar, 242, 243, 245-247, 250
 - superior, 244, 245, 248
- calcaneonavicular, internal:
 - inferior portion, 242, 243, 246, 248
 - middle portion, 242, 243, 247, 248, and note p. 248
 - superior portion, 242
- capsular, 166-168, 179 (see also "Capsule or capsules")
 - of the elbow-joint, 204-208
 - of the hip-joint, fibrous portion, 224
 - synovial portion, 224
 - of the shoulder-joint, 199, 200, 203
 - of the temporomandibular articulation, 194, 196
 - of the temporomaxillary articulation, 194, 196
- of the carpus, annular, anterior, 211-213
 - palmar, 212
 - radiate, 212
- carpometacarpal, 210-214
 - anterior, 211, 212
 - dorsal, 210, 213
 - interosseous, 214
 - palmar, 211, 212
 - posterior, 210, 213
- central, of the spinal cord, 179
- coccygeal, lateral, 179, 180
- conoid, 199, 202
- Cooper's, 221, 224
- coraco-acromial, 170, 199, 200, 202
- coracoclavicular, 199, 202
- coracohumeral, 199, 200
- coracoscapular, 170, 199, 200, 202
- costoclavicular, 186, 188, 198
- costocentral, anterior, 176, 178, 182, 183
 - interarticular, 182, 183, 185
 - stellate, 176, 178, 182, 183
- costotransverse, anterior superior, 176, 183
 - middle, 182
 - posterior, 182, 184, 185
 - posterior superior, 184
- cotyloid, 223-225, 227
- crucial, 229, 233, 235-237
 - of the knee, 229, 233, 235-237
- cruciform, of the atlas, 191, 192
- cubocuneiform, dorsal, 241, 245, 248
 - interosseous, 249, 250
 - plantar, 246, 247
- of the elbow-joint, anterior, 204, 205
 - capsular, 204-208
 - external lateral, 204, 205, 207
 - internal lateral, 204-206
 - posterior, 206, 207
- glenoid, 168
 - of the shoulder-joint, 201, 203
- of the hip-joint:
 - capsular, 168, 222, 223, 226, 227
 - cotyloid, 223-225, 227
 - iliofemoral, 222, 224, 226, 227
 - interarticular, 169, 223-225, 227
 - ischiofemoral, 222, 226, 227
 - pubofemoral, 222, 224, 226, 227
 - round, 169, 223-225, 227
 - teres, 169, 223-225, 227
 - transverse, of acetabulum, 168, 223, 225, 227

Ligament or ligaments:

of the hip-joint:

- Y, of Bigelow, 222, 224, 226, 227
- zona orbicularis, 223, 224, 226, 227
- iliofemoral, 222, 224, 226, 227
- iliolumbar, 176, 216, 218, 224
- interarticular, of the hip-joint, 169, 223-225, 227
- interclavicular, 157, 158, 198
- intercuneiform, dorsal, 244, 248
 - interosseous, 249
- intermetacarpal, distal, anterior, 211
 - palmar, 211
 - proximal, anterior, 211, 212
 - dorsal, 210, 213
 - interosseous, 214
 - palmar, 211, 212
 - posterior, 210, 213
- intermetatarsal, distal, dorsal, 248
 - plantar, 246
 - proximal, dorsal, 244, 245, 248, 249
 - plantar, 246, 247
- interosseous, of the forearm, 209
 - of the leg, 228, 229, 231-235, 238-241
- interspinous, 163, 174, 177, 181
- intertransverse, 184
- ischiofemoral, 222, 226, 227
- of the knee-joint, 228-237
 - alar, 234, 235
 - capsular, 228, 230-233, 236, 237
 - crucial, anterior or external, 229, 233, 235-237
 - internal or posterior, 229, 233, 235-237
 - lateral, external, 228, 229, 231-235
 - posterior, 231, 232
 - arcuate portion, of the, 232
 - direct portion of the, 231, 232
 - short, 231, 232
 - internal, 228-230, 232, 233, 235, 236
- mucoous, 234, 235
- patellar, 228-231, 236
 - lateral, external, 228, 231
 - internal, 228, 230
- posterior, 232, 235
- semilunar fibrocartilages:
 - external, 229, 233, 235-237
 - posterior or ascending band of the, 233, 235-237
 - internal, 229, 233, 235-237
 - transverse ligament of the, 229, 237
- Wrisberg's, 233, 235-237
- lateral, of the finger-joints, 166, 210, 211, 214
 - of the metacarpophalangeal articulation of the thumb, 210
 - of the metacarpophalangeal articulations, 166, 167, 210, 214
 - of the metatarsophalangeal articulations, 266-268
 - of the toe-joints, 247, 248
- metacarpal, transverse, 211
- metacarpophalangeal, anterior, 211, 215
 - glenoid, 211, 215
 - palmar, 211, 215
- metatarsal, transverse, 246
- metatarsocuneiform, interosseous, 249
- metatarsophalangeal, 246, 247
 - inferior, 250, and note, p. 246
- mucoous, 234, 235
- naviculocuboid, dorsal, 241, 245, 248
 - plantar, 246, 247
- naviculocuneiform, dorsal, 242-245, 248
 - plantar, 247
- oblique radio-ulnar, 209
- oburator, 117, 218, 222, 224, 226

Ligament or ligaments:

- occipito-atlantal, anterior, 176, 181, 192, 193
 - posterior, 181, 192, 193
- occipito-axial, posterior or long, 177, 190-192
- odontoid, alar, 191, 192
 - check, 191, 192
 - lateral, 191, 192
 - middle, 191, 192
 - suspensory, 191, 192
- orbicular, of the radius, 169, 204, 205, 207-209
- patellar, 228-231, 236
 - lateral, external, 228, 231
 - internal, 228, 231
- of the pelvis, transverse, 220, 221, and note, p. 221
- pisimetacarpal, 211, 212
- pisi-uncinate, 211, 212
- plantar, long, 242, 243, 245, 246, 247
 - short, 247, 250
- pterygospinous, 195, 196
- pubic, anterior, note, p. 220
 - of Astley Cooper, 221, 224
 - inferior, 217, 218, 220, 221
 - posterior, note, p. 221
 - subpubic, 217, 218, 220, 221
 - superior, 220, 221
- pubofoemoral, 222, 221, 226, 227
- rhomboid, 186, 188, 198
- round, of the hip-joint, 169, 223-225, 227
- sacrocccygeal, anterior, 180, 216
 - interarticular, 177, 179, 180
 - lateral, 177, 179, 180
 - posterior, deep, 177, 179, 180
 - superficial, 180, 218
- sacro-iliac, anterior, 216, 217, 219, 244
 - interosseous, note, p. 218
 - long, 218
 - oblique, 218
 - posterior, 218, 219
- sacrosciatic, anterior, 176, 180, 216-219
 - great, 176, 216-219
 - posterior, 176, 216, 219
 - small, 176, 180, 216-219
- sphenomandibular, 195
- sphenomaxillary, 195
- of the spine, 172-181
- sternoclavicular, 186, 187, 198
- stylohyoid, 194, 195
- stylomandibular, 195
- stylomaxillary, 195
- subflava, 173-175, 192
- suprascapular, 170, 199, 200, 202
- supraspinous, 173, 174, 177, 181, 184, 218
- tarsometatarsal, dorsal, 242, 244, 245, 248
 - interosseous, 249
 - plantar, 246, 247
- of the tarsus, dorsal, 244, 245
 - interosseous, 249
 - plantar, 246
- temporomandibular, 194
- temporomaxillary, 194
- tubiofibular, anterior inferior, 238, 244, 245
 - superior, 228, 229, 231, 234, 237, 238
 - posterior inferior, 240, 241
- transverse, of the acetabulum, 168, 223, 225, 227
 - of the atlas, 190-192
- of the transverse carpal articulation, 210-214
 - anterior, 212
 - dorsal, 210, 213
 - interosseous, 213, 214
 - palmar, 212
 - posterior, 210, 213
- trapezoid, 202
- triangular, of the urethra, 220, 221, and note, p. 221
- vaginal, of the toes, 246
- of the vertebral column, 172-181

Ligament or ligaments:

- of the vertebral column, anterior common, 173, 174, 176, 178, 181-183, 216
 - posterior common, 173, 174, 177, 179, 190, 192
 - of the wrist joint, anterior, 211, 212
 - annular, 211-213
 - dorsal, 210
 - lateral, external, 210, 213, 214
 - internal, 210-212, 214
 - palmar, 211, 212
 - posterior, 210
 - Ligamentum *vel* ligamenta:
 - accessoria plantaria, 246, 247, 250
 - volaria, 211, 215
 - acromioclaviculare, 199, 201
 - alaria, 191, 192
 - annulare radii, 169, 204, 205, 207-209
 - apices dentis, 191, 192
 - arcuatum pubis, 217, 218, 220, 221
 - basium (ossium metacarpalium) dorsalia, 210, 213
 - (ossium metacarpalium) interossea, 214
 - volaria, 121, 212
 - (ossium metatarsalium) dorsalia, 244, 245, 248, 249
 - (ossium metatarsalium) plantaria, 246, 247
 - bifurcatum, 245, 248, 249
 - +
 - calcaneocuboideum dorsale, 244, 245, 248
 - plantare, 242, 243, 246, 247, 250
 - calcaneofibulare, 239-241, 244, 245
 - calcaneonavicular dorsale, 242
 - plantare, 242, 243, 246-248
 - calcaneotibiale, 239-242
 - capituli costae interarticular, 182, 183, 185
 - radiatum, 176, 178, 182, 183
 - fibulae, 228, 229, 231, 234, 237, 238
 - capitulorum (ossium metacarpalium) trans-versa, 211
 - (ossium metatarsalium) dorsale, 248
 - trans-versa, 246
 - carpi radiatum, 212
 - transversum, 211-213
 - carpometacarpea dorsalia, 210, 213
 - interossea, 214
 - volaria, 211, 212
 - coccygeum laterale, 179, 180
 - collaterale carpi radiale, 210, 213, 214
 - ulnare, 210-212, 214
 - fibulare, 228, 229, 231-235
 - radiale, 204, 205, 207
 - tibiale, 228-230, 232, 233, 235, 236
 - ulnare, 204-206
 - collateralia articulationum digitorum (manus), 166, 210, 211, 214
 - articulationum digitorum pedis, 247, 248
 - articulationis metacarpo-phalangeae pollicis, 210
 - articulationum metacarpophalan-gearum, 166, 167, 210, 214
 - articulationum metatarsophalan-gearum, 246-248
- colli costae, 182
- columnae vertebrales, 172-181
- conoideum, 199, 202
- coraco-acromiale, 170, 199, 200, 202
- coracoclaviculare, 202
- coracohumeral, 199, 200
- costoclaviculare, 186, 188, 198
- costotransversarium anterior, 176, 183
 - posterior, 184
- costoxiphoidea, 186
- cruciatum anterior, 229, 233, 235-237
 - atlantis, 191, 192

Ligamentum *vel* ligamenta:

- cruciatum posterius, 229, 233, 235-237
- cruciata (genu), 229, 233
- cuboideonaviculare dorsale, 244, 245, 248
 - plantare, 246, 247
- cuneocuboideum dorsale, 244, 245, 248
 - interosseum, 249, 250
 - plantare, 246, 247
- cuneometatarsa interossea, 249
- deltoidium, 242, 243, 249
- flava, 173-175, 192
 - iliofemorale, 222, 224, 226, 227
- iliofemorale, 176, 216, 218, 224
- iliofemorale dorsalis, 210, 213
- intercarpa dorsalis, 213, 214
 - volaria, 212
- interclaviculare, 187, 188, 198
- intercostalia externa, 186
 - interna, 176, 183, 184, 187
- intercuneiformia dorsalis, 244, 248
 - interossea, 249
- interspinalia, 163, 174, 177, 181
- intertransversaria, 184
- ischiocapsulare, 222, 226, 227
- longitudinale anterius, 173, 174, 176, 178, 181-183, 216
 - posterius, 173, 174, 177, 179, 190, 192
- lumbocostale, 176, 216, 218
- malleoli lateralis anterius, 238, 244, 245
 - posterius, 245, 241
- menisci lateralis (Roberti), 233, 235-237
- navicularicuneiformia dorsalis, 242-245, 248
 - plantaria, 247
- nuchae, 177, 181
- patellae, 228-231, 236
 - pisohamatum, 211, 212
 - pisometacarpeum, 211, 212
 - plantare longum, 245-247
 - popliteum arcuatum, 232
 - obliquum, 232, 235
 - pterygospinosum, 195, 196
 - pubicum superius, 220, 221
 - pubocapsulare, 222, 224, 226, 227
 - radiocarpeum dorsale, 210
 - volare, 211, 212
- sacrocoxygeum anterius, 180, 216
 - articulare, 177, 179, 180
 - laterale, 177, 179, 180
 - posterius profundum, 177, 179, 180
 - superficiale, 180, 181
- sacro-iliaca anteriora, 216, 217, 219, 224
 - interossea, 218, 219
- sacro-iliacum posterius breve, 219
 - longum, 218, 219
- sacrospinosum, 176, 180, 216-219
- sacrothoracum, 176, 216-219
- sphenomandibulare, 195
- sternoclaviculare, 186, 187, 198
- sternocostale interarticular, 188
- sternocostalia radiata, 186
- stylohyoideum, 194, 195
- stylomandibulare, 194-196
- supraspinale, 173, 174, 177, 181, 184, 218
- talocalcaneum anterius, 244, 245, 250
 - interosseum, 239, 244, 245, 249, 250
 - laterale, 245
 - mediale, 243, 248
 - posterius, 240, 242
- talofibulare anterius, 244, 245
 - posterius, 239, 241
- talonavicular (dorsale), 242-245
- tarsi dorsalis, 244, 245
 - interossea, 249
 - plantaria, 246
- tarsometatarsa dorsalis, 242, 244, 245, 248
 - interossea, 249

Ligamentum *vel* ligamenta:

- tarsometatarsa plantaria, 246, 247
- temporomandibulare, 194
- teres femoris, 169, 223-225, 227
- tibionaviculare, 242, 244, 245, 248
- transversum acetabuli, 188, 223, 225, 227
 - atlantis, 190-192
 - genu, 229, 237
 - pelvis, 220, 221
 - scapulae inferius, 200
 - superius, 170, 199, 200, 202
 - trapezoideum, 202
 - tuberculi costae, 182, 184, 185
 - vaginalia (digitorum pedis), 246
- Line, Chopart's, 249
 - Hey's, 249
 - Lisfranc's, 249
 - mediotarsal, 249

M.

- †Membrana atlanto-epistrophica, 192
 - atlanto-occipitalis anterior, 176, 181, 192, 193
 - posterius, 181, 192, 193
 - interossea antibrachii, 209
 - cruris, 238
 - obturatoria, 117, 218, 222, 224, 226
 - sterni, 186, 187
 - tectoria, 177, 190-192
- Membrane, intercostal, see "Aponurosis"
- interosseous, of the forearm, 209
 - of the leg, 228, 229, 231-235, 238-241
- oburator, 117, 218, 222, 224, 226
- synovial, of the hip-joint, 223-225, 227; its reflection on the neck of the femur, 224
 - of the knee-joint, 234, 235
- Meniscus lateralis, 229, 233, 235-237
 - medialis, 229, 233, 235-237
- Musculus articularis genu, 228, 230, 231, 235

N.

- Notch, clavicular, of the sternum, 198
 - cotyloid, 223-225
 - suprascapular, 199, 200
- Nucleus pulposus of the intervertebral discs, 173, 174

O.

- Obturator canal, 217, 222, 224, 226
 - fascia, 221
 - ligament, 117, 218, 222, 224, 226
 - membrane, 117, 218, 222, 224, 226
- Ossa sesamoidea manus, 210, 211, 214
 - pedis, 246, 247, 250
- Ossa sesamoidea articulationis genu (var.), 232

P.

- Pad of fat of the fossa acetabuli, 223
- Periosteum, dental, 165
- Plate, fibrous, 246, 247, 250, and notes, pp. 211 and 246
 - glenoid, 246, 247, 250, and notes, pp. 211 and 246
 - sesamoid, 246, 247, 250, and notes, pp. 211 and 246
- Plicae alares, 234, 235
- Plica synovialis (articulationis radiocarpeae), 213
 - patellaris, 234, 235
- Pouch of synovial membrane, circular, of the proximal radio-ulnar articulation, 204, 207, and note, p. 204
 - vertical, of the distal radio-ulnar articulation, 214
- Process, falciform, of the great sacrosacral ligament, 217
- Processus calciformis, 217
- Prominence of the pubic symphysis, posterior, 221
- Promontorium, 177, 216, 217, 219
- Promontory of the sacrum, 177, 216, 217, 219
- Pubic symphysis, 218, 220, 221

R.

- *Recess, sacciform, of the distal radio-ulnar articulation, 214
 of the proximal radio-ulnar articulation, 204, 207, and note, p. 204
 Recessus sacciformis (articulationis radio-ulnaris [distalis]), 214
 (articulationis radio-ulnaris proximalis), 204, 207
 Retinaculum ligamenti arcuati, 231, 232
 patellae lateralis, 228, 231
 mediale, 228, 231

S.

- Sacral promontory, 117, 216, 217, 219
 Sesamoid bones, see "Bones, sesamoid"
 of the foot, 246, 247, 250
 of the hand, 210, 211, 214, and note, p. 211
 Shoulder-joint, 168, 200-203
 Substance, sutural, 164
 Sulcus paraglenoidalis (præauricularis), 216
 Surface, articular, of the acetabulum, 168, 224, 225, 227
 of the acromion, 201
 of the astragalus for the os calcis, anterior, 248
 of the astragalus for the os calcis, middle, 248
 of the astragalus for the os calcis, posterior, 248
 of the atlas, superior, 167, 169
 of the centra for the ribs, 183
 of the heads of the ribs, 182
 of the navicular bone, posterior, 248
 of the odontoid process, anterior, 192
 posterior, 192
 of the os calcis, anterior, 248
 middle, 248
 posterior, 247
 of the patella, 229, 234, 235
 of the transverse processes for the ribs, 175, 178, 185
 of the astragalus, superior articular:
 external malleolar portion, 166
 internal malleolar portion, 166, 241
 tibial portion, 166, 241, 243
 auricular, of the ilium, 219
 patellar, of the femur, 229, 234, 237
 Sutura, 164
 serrata, 164
 squamosa, 164
 Sutural substance, 164
 Suture, varieties of, 164
 Symphysis, 165
 ossium pubis, 218, 220, 221
 sacrococcygea, 180, 216, 217
 Synarthrosis, 164, 165
 Synchrondrosis, 165
 of the base of the skull, 196

Synchrondrosis *vel* synchrondroses

- epiphysial, see "Dise, epiphysial"
 epiphyscos capitis femoris, 223
 humeri, 201
 capituli ossis metacarpalis III., 215
 distalis femoris, 230
 fibulae, 239, 240, 244
 radii, 215
 tibiae, 239, 240, 242, 244, 250
 proximalis fibulae, 231
 tibiae, 230, 231
 tuberi calcanei, 240
 epiphysium ossium metatarsalium, 249, 250
 phalangium digiti II. (manus), 215
 digitorum pedis, 249
 et ligamenta bascos cranii, 196
 interphalangea pedis (var.), 249
 petro-occipitalis, 193, 196
 spheno-occipitalis, 165, 192, 196
 sphenopetrosa, 196
 sternalis, inferior, 188
 superior, 188
 Syndesmosis, 170
 tibiofibularis, 238-241, 244, 245
 Synovial membrane of the shoulder-joint, 202, 203
 sheath of the bicipital groove, 201, 202
 of the tendon of the popliteus muscle, 232, 233, 235, 237: its communication with the knee-joint and the superior tibio-fibular articulation, 233

T.

- Tendon of the long head of the biceps muscle, 201, 202, 203
 †Torus pubicus, 221
 Triangular fibrocartilage, 212-214
 ligament of the urethra, 220, 221, and note, p. 221
 Trochlea tali, 166, 239, 241
 facies malleolaris lateralis, 166
 medialis, 166, 243
 superior, 166, 241, 243
 Trochlear surface of the astragalus, 166, 239, 241

V.

- Vagina mucosa intertubercularis, 201, 202

W.

- Wrist-joint, 169, 212-215

Z.

- Zona orbicularis, 223, 224, 226, 227

COLUMBIA UNIVERSITY LIBRARIES

This book is due on the date indicated below, or at the expiration of a definite period after the date of borrowing, as provided by the library rules or by special arrangement with the Librarian in charge.

[illegible]

COLUMBIA UNIVERSITY LIBRARIES

1010231200

Q. 1. 2. 3.

10104

10104

1-1-2
cop 3

~~Series of the same series~~

AUG 16 1938

RESERVES

NOV 17

DEC 16

Handwritten signature

QM25

T572

v.1-2

cop.3