62K
507
W85

A

0
0
0
8
6
9
8
5
4
0

UC SOUTHERN REGIONAL LIBRARY FACILITY

PRELIMINARY NOTES ON
THE EMBRYOLOGY OF REBOULIA
HEMISPHAERICA
BY
WILLIAM L. WOODBURN

QK
507
W85 Woodburn

Southern Branch
of the
University of California
Los Angeles

Form L 1

QK
507
W85

PRELIMINARY NOTES ON
THE EMBRYOLOGY OF REBOULIA
HEMISPHERICA

WILLIAM L. WOODBURN

NEW YORK

1919

Q K
507
W 85

SOUTHERN BRANCH,
UNIVERSITY OF CALIFORNIA,
LIBRARY,
LOS ANGELES, CALIF.

Preliminary notes on the embryology of *Reboulia hemisphaerica*

WILLIAM L. WOODBURN

(WITH PLATE 19)

The object of this paper is to present a part of a more extended study on the embryology and cytology of *Reboulia hemisphaerica* (L.) Raddi. Only certain stages in the process of fertilization and in the early development of the embryo sporophyte will be considered at this time. The description of various other stages in the life history together with a summary of the previous literature on *Reboulia* will be left for a subsequent report.

FERTILIZATION

The male gamete in the bryophytes may undergo two series of marked morphological changes. The first series includes the steps leading from the non-motile condition of the androcyte (the cell which directly becomes transformed into the mature sperm) through the formation of the actively motile free-swimming sperm. The second series includes the reverse steps and occurs in those cases where a sperm reaches an egg. In this second case the motile sperm becomes again non-motile with a resting nucleus similar to that of the androcyte or androcyte mother-cell. Details of the main stages in the first series have been carefully described for numerous liverworts and mosses. The processes accomplished in the second series, after the sperm reaches the egg, are not so well known. However, soon after

3, 4, 5, 6 and 11) indicate that from the hypobasal cell is developed the foot and from the epibasal cell the stalk and sporangium of the mature sporophyte.

A third division follows in either the middle or apical cell of the tier of three (FIG. 4) which results from the first and second divisions. Compare FIG. 5, which shows a tier or series of four cells, with FIG. 4. Then follows (FIG. 6) a division in the basal or foot cell at right angles to the first three division walls. The foot at this stage has become quite dense in protoplasmic contents.

The order of divisions just described seems to represent the usual conditions. Compare, however, FIGS. 2-6 with FIGS. 7-10. In each section represented by FIGS. 7 and 9 there is a triangular-shaped apical cell, while in FIGS. 8 and 10 both apical and basal cells of triangular shape are present. FIG. 11 represents a slightly different condition, in which the foot has become divided into an irregular group of cells.

In no case do we find the same sequence of early divisions as described by Cavers (2). In speaking of the early divisions of the sporophyte of *Reboulia*, he says, "The transverse basal wall is followed by two sets of nearly equal vertical walls which intersect each other at right angles, so that the embryo shows a regular octant stage."

SUMMARY.

The egg and sperm nuclei are both in a resting condition in the earliest stages of fusion.

Among the Bryophytes little is known concerning the details of nuclear behavior during the stages of fertilization.

The earliest divisions of the zygote are transverse. A longitudinal series or tier of four cells may be formed.

Occasionally, however, both apical and foot cells of triangular outline may be formed, or the basal cell may divide into an irregular group which constitutes the foot.

Of the first two cells formed, the hypobasal cell evidently produces the foot, and the epibasal cell the stalk and sporangium of the mature sporophyte.

LITERATURE CITED

1. Black, Caroline A. The morphology of *Riccia Frostii* Aust. Ann. Bot. 27: 511-532. pl. 37, 38. 1913.
2. Cavers, F. Contributions to the biology of the Hepaticae. Part I.—*Targionia*, *Reboulia*, *Preissia*, *Monoclea*. Leeds and London. March, 1904.
3. Garber, J. F. The life history of *Ricciocarpus natans*. Bot. Gaz. 37: 161-177. pl. 9, 10 + f. 1-4. 1904.
4. Meyer, K. Untersuchungen über den Sporophyt der Lebermoose. Entwicklungsgeschichte des Sporogon der *Corsinia marchantioides*. Bull. Soc. Imp. Nat. Moscou 1911: 263-286. pl. 11 + f. 1-22. 1912.

Explanation of plate 19

The figures were drawn by Miss Mary C. Blair.

FIG. 1. Early stage of fusion of the egg and sperm nuclei, $\times 1775$.

FIG. 2. Telophase of first division of zygote, $\times 565$.

FIG. 3. First division of zygote completed; hypobasal cell "a," and epibasal cell "b" in prophase of subsequent division, $\times 735$.

FIG. 4. Division of epibasal cell completed, showing tier of three cells, $\times 565$.

FIG. 5. Young embryo consisting of tier of four cells, $\times 565$.

FIG. 6. Hypobasal cell divided longitudinally; cell next above in process of longitudinal division, $\times 565$.

FIG. 7. Young embryo with wedge-shaped apical cell, $\times 735$.

FIG. 8. Young embryo with both basal and apical cells triangular in shape, $\times 735$.

FIG. 9. Embryo, older than in any of the preceding figures, with wedge-shaped apical cell $\times 565$.

FIG. 10. Slightly more advanced embryo than in FIG. 9, showing both apical and basal cells triangular in outline, $\times 565$.

FIG. 11. Embryo showing early differentiation of foot "a," seta "b," and capsule "c," $\times 565$.

65587

WOODBURN: EMBRYOLOGY OF REBOULIA

Lithomount
Pamphlet
Binder
Gaylord Bros.
Makers
Syracuse, N. Y.
PAT. JAN 21, 1908

UC SOUTHERN REGIONAL LIBRARY FACILITY

A 000 869 854 0