Digitized by the Internet Archive
in 2010 with funding from
University of Toronto

BRITISH POTAMOGETONS

(POND WEEDS)

THE
POTAMOGETONS
(POND WEEDS)
OF THE
BRITISH ISLES

WITH DESCRIPTIONS OF ALL THE SPECIES, VARIETIES AND HYBRIDS

BY
ALFRED FRYER, A.L.S.
AND
ARTHUR BENNETT, A.L.S.

(The Lucens Group edited from MR. FRYER'S Manuscript Notes by A. H. EVANS, F.Z.S.)

ILLUSTRATED BY
ROBERT MORGAN, F.L.S.
AND OTHERS

199790
12 / 1 / 26

LONDON
L. REEVE & CO. LIMITED
6, HENRIETTA STREET, COVENT GARDEN

1915

LONDON: PRINTED BY WILLIAM CLOWES AND SONS, LIMITED, DUKE STREET, STAMFORD STREET S.E.,
AND GREAT WINDMILL STREET, W.

CONTENTS.

	PLATES	PAGE
INTRODUCTION		vii
KEY TO THE SPECIES		xi
GENERAL DESCRIPTION OF THE GENUS POTAMOGETON		1
BRITISH SPECIES AND HYBRIDS—		
1. natans	1-3	4
2. crassifolius	4-5	8
3. fluitans	6-7	11
4. Kirkii	8-9	16
5. polygonifolius	10-12	18
6. coloratus	13-15	22
7. Billupsii	16-17	25
8. alpinus	18-20, 26 (figs. 6, 7, 8)	27
9. Drucei	21	31
10. Griffithii	22-23	34
11. praelongus	24-26	36
12. perfoliatus	27-28	38
13. crispus	29-30	43
14. Cooperi	31-32	48
15. Bennettii	33	53
17. densus	34	56
The lucens group, edited by A. H. Evans	—	59
18. lanceolatus	39	63
19. graminifolius	36	64
20. heterophyllus	35	65
21. varians	41	67
22. falcatus	40	68
23. involutus	44	69
24. nitens	37-38	69
25. lucens	45-47	71
26. Zizii (= angustifolius)	42	72
27. coriaceus	43	74
28. decipiens	48	75
29. salignus	49	76
Grass-leaved group, by Arthur Bennett	—	77
30. compressus (zosteraefolius)	50	77
31. acutifolius	51	79
32. obtusifolius	52	80
33. Friesii	53	81
34. rutilus	54	82
35. pusillus	55	83
36. Sturrockii	(Fig. 1)	85
37. trichoides	56	85
38. pectinatus	57	86
39. vaginatus	58	87
40. interruptus	59	88
41. marinus	60	89
42. pennsylvanicus	(Fig. 2)	91
INDEX TO GROUPS, SPECIES AND SYNONYMS		93
PLATES.

INTRODUCTION

BY

ARTHUR BENNETT, A.L.S.

THE present work was projected some time ago by my friend the late Mr. Alfred Fryer, who had devoted many years to a most careful study of the British water plants included in the genus *Potamogeton* and their various states and forms, many of which he had kept under continuous observation by means of artificial cultivation. In the first few pages of the work he describes the general characters of the genus, and refers to the manner in which he proposed to deal with some of the problems connected with the systematic arrangement of the British species and natural hybrids. At the time of his death he had completed and passed for press the descriptions of the first seventeen species and hybrids, and had made considerable progress in the preparation of the matter for the next group, the *lucens* group (pp. 59-76). Mr. A. H. Evans, who edited this section of the work after Mr. Fryer's death, wrote as follows in regard to it:—"With the exception of the distribution of the various forms, contributed, as in the earlier parts, by Mr. Arthur Bennett, the whole of this instalment may be taken as the result of the labours of Mr. Fryer on the section of Pondweeds to which he chiefly devoted himself. He left in MS. the introductory to this '*lucens*-group,' and he also sketched out the specific portion as a whole. A few of the articles were quite finished, and I am glad to say that I have found it possible to complete the remainder either from the author's own '*Notes on Pondweeds*' in the '*Journal of Botany*' for various years or from the information which he constantly gave me during the expeditions which we made together in the later part of his life to the localities near Chatteris, where the rarer plants grow or grew. I revised with Mr. Fryer the introductory paragraphs, and feel certain that the whole of the letterpress represents his final ideas on the group. If he had lived a few months longer he might have added further details, especially with regard to the land-forms, or what he considered mere states of the species he recognised; and therefore his former '*Notes on Pondweeds*' should be read in conjunction with the present instalment, though it should be remembered that his mature views are not likely always to be in accord with those of his earlier years, when he was still engaged in struggling with the difficulties of his subject and first observing the growth of the plants in his tanks or in their native waters."

For the remainder of the work, dealing with the grass-leaved group, the present writer alone is responsible. I have had no help from my old friend's MS. All that he left for this section were the names and synonyms of the species, and these I have adopted almost as he left them. Although Mr. Fryer had not taken quite the same keen interest in the grass-leaved *Potamogetons* as he had done in the other groups, it will always be a matter of regret that he did not live to work them out in the same exhaustive manner, and himself complete this monograph which will ever remain a monument to his memory.

Mr. Fryer at one time suggested that the fertilisation of the genus was mainly brought about by small aquatic beetles, but though he was a keen entomologist, he does not seem to have followed this up, and I am not aware of any writer on our aquatic beetles who has suggested any such connection between them. Mr. F. Balfour Brown, in an article on the "Aquatic Coleoptera of the Norfolk Broads,"* does not mention anything with reference to such a connection.

In our own country the first to put the genus in anything like a systematic form was Sir J. E. Smith in his "English Flora," 1824-28, where he described thirteen species, eight of the broad-leaved and five of the grass-leaved species. From that date to the year 1904, when the 9th edition of Babington's Manual of British Botany appeared, the genus has gradually been added to and revised, until our *Potamogetons* have become fairly well known, though so far as their life-histories are concerned there is yet much to be learnt.

The genus was instituted by Tournefort in his "Institutiones rei Herbariae," p. 232, 1700; it was taken up by Linnæus in the first edition of his "Species Plantarum," p. 126, 1753, where he gives twelve species. Lourerio, in his "Flora Cochinchinensis," p. 301, 1790, uses the name "*Hydrogeton heterophyllus*" for a species, perhaps one of the *natans* group. Rafinesque, in the "Medical Reports," New York, vol. v. p. 354 (1808), varies the name to "*Potamogiton*"; and Clairville, in his "Manuel d'Herborisation en Suisse et en Valais," pp. 34, 44 (1811) has "*Potamogetum*." Rafinesque, in Journ. Phys. T. 89, p. 101 (1819), suggests "*Peltopsis*" for *P. perfoliatus*, L. Gay, in Compt. Rend. Acad. vol. xxxviii. p. 702 (1854), has "*Spirillus*" for the American species *P. hybrides* and *spirillus*, and on p. 703 suggests the name *Groenlandia* for *P. densus*, L.; but none of these names have been taken up, as the genus is a very natural one, and does not need to be broken up.

It is spread all over the five divisions of the world, but it is far more common in temperate than in tropical countries. In the Monograph of the Order in Engler's "Das Pflanzenreich," Heft 31, 1907, Dr. Graebner recognises eighty-seven species, with many sub-species, varieties and other forms, and about forty-seven hybrids; but

* Trans. Norf. and Norwich Nat. Hist. Soc. vol. viii. pp. 58-82 (1905).

these numbers are at present merely tentative, as the genus as a whole still requires much careful investigation and research before anything approaching finality can be affirmed. Since 1907 about eleven species and eighteen hybrids and sub-species have been described, with one omitted by Graebner in 1901, namely, *P. Columbianus Sukesdorf*.

Forty-two of these species are limited to one country each, and using A. R. Wallace's Areas of Zoological Distribution,* twenty-two species are confined to the Palaearctic Region, eighteen to the Nearectic, fifteen to the Neotropical, five to the Ethiopian, one to the Oriental and nine to the Australian Region. The remaining twenty-eight species are more or less distributed over all the regions, but one only, *P. pectinatus*, L., is found in all six regions.

According to the Kew arrangement; two species are confined to Europe, six to North Africa and the Orient, three to Northern Asia, nine to China and Japan, two to Malaya, four to Australia, two to Polynesia, two to Tropical Africa, four to the Mascarene Islands, fifteen to North America, one to Central America, one to the West Indies, four to East Tropical South America, five to Temperate South America, while there are no species confined to India, New Zealand, South Africa, or West Tropical America; here again *P. pectinatus*, L., is recorded from all these divisions except Malaya.

Of these the rarest are *P. sibericus*, Ar. Benn (once gathered), *P. Drummondii*, Bentham, gathered by Drummond, c. 1844, and not found since, *P. Griffithii*, Ar. Benn., *P. Curtisii*, Morang, in Florida, and three or four later named species which at present must be held tentative as to distribution, especially the Japanese. These islands are very rich in the genus, and have not yet been thoroughly worked, though there is a probability that two at present endemic North American species occur there. But this is quite in keeping with the character of the Flora, which the late Dr. Asa Gray has commented on.†

The limitation of the species is undoubtedly somewhat difficult, as some of them are exceedingly variable, being split up into petty varieties which are merely the description of individuals, whilst happily others will not lend themselves to be so divided. Considered from the local point of view this splitting up into varieties is of value so long as they are held to be local variations; studied in connection with their environment as to colour, depth, density, chemical composition of the water, etc., they would in time evolve many interesting facts on their origin and evolution.

To any one with a critical faculty and spare time to grow them, the genus would prove very attractive, as although Mr. Fryer grew many of the broad-leaved species,

* A. R. Wallace, "Island Life," ed. p. 2, 31 (1892).

† In Am. Jour. of Science, vol. 11, series 2, p. 135 (1846). Mem. Am. Acad. vol. vi. p. 377 (1859).

he did not grow the grass-leaved species, and these would well repay the labour entailed.

They are not difficult to grow, but if grown in pots submerged, it is well to cover the earth in which they are grown with a thin stratum of clay. If in tubs this is not required, and while they will succeed in almost any soil, a mixture of one-third of sharp sand is beneficial. In the last section, such species as *pectinatus* require more room for rooting than the grass-leaved species, as they form many tuberous roots which extend in every direction. If grown in a pond the margin is better than deeper parts; except for such species as *prælongus*, *lucens*, and *perfoliatus* which are strictly submerged species of deep water, and never form floating leaves.

KEY TO THE SPECIES OF POTAMOGETON.

1. Stipules free from the leaf.
 - a. *With floating leaves.*

Floating leaves various, coriaceous.
Fruit large.
Submerged leaves bladeless *natans*, p. 4, pl. 1-3.
" " with blades *fluitans*, p. 11, pl. 6-7.

Fruit small.
Leaves all pellucid *coloratus*, p. 22, pl. 13-15.
" not " *polygonifolius*, p. 13, pl. 10-12.
" elongated, 15 inches long *Kirkii*, p. 16, pl. 8-9.
Floating leaves ovate *pennsylvanicus*, p. 91, fig. 2.

Floating leaves lanceolate.
Leaves with distinct chain-areolation *lanceolatus*, p. 63, pl. 39.
Submerged leaves all linear, others various.
Leaves petiolate, submerged lanceolate, freely fruiting *heterophyllus*, p. 65, pl. 35.
" " " strap-shaped *graminifolius*, p. 66, pl. 36.
" clasping, " " *falcatus*, p. 68, pl. 40.
" " sub-sterile or wholly so *nitens*, p. 69, pl. 37-38.
Submerged leaves not linear.
Fruit acuminate *alpinus*, p. 27, pl. 18-20, and 26 (figs. 6-8).
" rounded *Zizii* (= *angustifolius*), p. 72, pl. 12.
 - b. *Without floating leaves.*

Stems not compressed.
Leaves cucullate at the apex *prælongus*, p. 36, pl. 24-26.
" perfoliate, not cucullate *perfoliatus*, p. 38, pl. 27-28.
" petiolate, acute or rounded *lucens*, p. 71, pl. 45-47.
" sessile, rounded *decipiens*, p. 75, pl. 43.

Stems compressed.
Leaves usually crisped and serrulate *crispus*, p. 43, pl. 29-30.
Leaves all linear, submerged, fruit not toothed.
5 principal nerves *zosterifolius*, p. 77, pl. 50.
3 " " *acutifolius*, p. 79, pl. 51.

Stems compressed, branched, no secondary nerves.
5-7-nerved, leaves fascicled *Friesii*, p. 81, pl. 53.
3-nerved, not fascicled, glands large translucent *obtusifolius*, p. 80, pl. 52.
1-3-nerved, not fascicled, glands small, dull *pusillus*, p. 83, pl. 55.

Stems slightly compressed, usually simple.
Leaves 3-5-nerved *rutulus*, p. 82, pl. 54.

Stems filiform, branched.
Leaves 3-5 nerved *Sturrockii*, p. 85, fig. 1.
" setaceous, fruit toothed *trichoides*, p. 85, pl. 56.
2. Stipules sub-adnate to the leaf.
With floating and submerged leaves *Griffithii*, p. 34, pl. 22-23.
3. Stipules adnate to the leaf.
Fruit large, style lateral.
Leaves 1-nerved, nut with faint lateral ridges *pectinatus*, p. 86, pl. 57.
" 3-5- " " strong " " *interruptus*, p. 88, pl. 59.
- Fruit small, style faint, nearly central.
Leaf-sheaths much inflated *vaginatus*, p. 87, pl. 58.
" not " *marinus*, p. 89, pl. 60.
4. Stipules absent.
Leaves all opposite, submerged *densus*, p. 56, pl. 34.

Other forms, mostly intermediate between their supposed parents and variable, described in this work, but not included in the above Key:—

crassifolius, p. 8, pl. 4-5.
Billupsii, p. 25, pl. 16-17.
Drucei, p. 31, pl. 21.
Cooperi, p. 48, pl. 31-32.
Bennettii, p. 53, pl. 33.

varians, p. 67, pl. 41.
involutus, p. 69, pl. 44.
coriaceus, p. 74, pl. 43.
salignus, p. 76, pl. 49.

THE POTAMOGETONS OF THE BRITISH ISLES.

POTAMOGETON, *Linnaeus*.

Flowers perfect, sessile, on a terminal or axillary pedunculate spike; perianth segments 4, incurved, unguiculate; anthers 4, sessile, 2-celled, inserted at the base of the claws; carpels 4, with shortly styled or subsessile stigmas, 1-ovuled; fruit a compressed, ovoid, or rounded drupelet, with the embryo incurved.

Perennial aquatic herbs, inhabiting fresh, or rarely brackish or salt water; some species are amphibious when wholly or partially forsaken by the water in summer, then producing stunted growths fully exposed to the air; these are always infertile, but continue the life of the individual plant by thickened subterraneous stolons.

ROOTSTOCK, in its simplest form, a stolon creeping in the soil and producing a stem at every second joint; stolons either few or numerous, nearly simple, or with numerous secondary stolons; perennial or sometimes only annual in species which are reproduced by winter-buds formed on the fruiting branches of the plant.

STEM usually terete, sometimes compressed, or flattened and leaf-like; mostly annual, rarely with the lower part surviving, and throwing out fresh branches a second year.

LEAVES sheathing at the base as in *P. pectinatus*, or narrowly strap-shaped and sessile, as in *P. pusillus*, or membranous, broad, and amplexicaul, as in *P. perfoliatus*, or short stalked, as in *P. lucens*, and submerged in all these species; or the lower membranous and submerged, and the upper stalked, coriaceous and floating, as in *P. heterophyllus*, or the lower reduced to a petiole and midrib, as in *P. natans*; extremely variable in shape (even on the same plant) from linear to orbicular; alternate, or opposite below the peduncles and branches; and all opposite in one species *P. densus*.

STIPULES adnate in the *Pectinati*; or free, and clasping the stem in the rest of the groups, in which, however, the lowest stipule on each stem or branch often carries an adnate leaf, as in *P. coloratus*, or several of the lower stipules, as in *P. crispus*; herbaceous and persistent, or scarious, and soon decaying after their office of protecting the young shoots and leaves has been fulfilled.

growth of the plant; the others are probably due to temporary local conditions. Certainly all such forms as were observed by me for a few years, either growing naturally or cultivated, ultimately reverted to the ordinary typical form figured in our first plate. I have been able to watch plants resembling all those figured except var. *linearis*, and in no instance have I failed to trace a complete reversion to the type.

P. POLYGONIFOLIUS c. *LINEARIS*, Syme, Ex. Club Report, p. 37 (1875); London Catalogue, ed. 8 (1881); Botanical Exchange Club Report (1888), p. 191. "*P. natans*, *L. form.*," Ex. Club Rep., p. 312 (1891).

This early state of *P. natans* was mistaken by the late Dr. Boswell Syme for a form of *P. polygonifolius*; it differs in no respect from typical forms of *P. natans* which I have had under cultivation, except that in its Irish locality, "Long Range, Killarney," the force of the stream keeps it constantly submerged so as to prevent the formation of flowers, and to a great extent of the upper broad floating leaves. In consequence of this arrested development of the flower-spikes the phyllodes are more fully produced, and become more persistent, and the plant more branched. Mr. R. W. Sculley has liberally given me a very complete set of specimens of this form, by means of which it can be traced through all its stages until the typical jointed floating leaves are produced. Figure 5 in plate 2 shows the usual state of "*linearis*."

In the Ex. Club Rept. p. 136 (1886), under the name of "*P. polygonifolius* var. *fluitans*" (corrected by Mr. Bennett to "*pseudo-fluitans*"), a form of *P. natans* is noticed. The Rev. W. R. Linton gathered this in Connemara. This plant, as Mr. Bennett suggests, is *P. natans*, a form intermediate between the state named "var. *lanceolatus*," Fieber, and "*linearis*." This, probably growing in less rapid water, is enabled to produce small but abortive flower-spikes. These slender spikes occasioned some doubt as to its being true *natans*, but I have seen almost as delicate ones produced under unfavourable conditions in stagnant water.

"*P. NATANS* v. *PROLIXUS*, Koch," London Catalog. ed. 8, No. 1497, b. (1886). ed. 9, No. 1580, b. (1895). Koch, Fl. Germ., vol. ii. p. 775 (1844).

PLATE 3. The original specimen is figured on our plate, it was gathered by Mr. T. F. Mott near Leicester, in 1881. The central and right-hand figures represent specimens from the same locality gathered by Mr. E. F. Cooper in 1892; both gatherings were in May, and serve to illustrate the persistency of the local conditions which produced a form unvaried for many years. Here at last we seem to have reached a true, permanent variety; the leaves of the early and later gatherings are alike semipellucid and almost membranous in texture, and agree fairly well with Koch's description, "*Folia submersa saepe numerosa basi apiceque attenuata pellucida, tamen non ita conspicue ut in P. fluitante.*" From this description (assuming that Koch had not a form of *P. fluitans* before him) it is

evident that Koch mistook the not yet emerged upper leaves for strong submerged lower leaves. This state of *P. natans* is rarely to be met with later in the season. Mr. Cooper found a specimen in July with typical *natans*, and I once found it in a densely crowded pond under the shade of trees, as late as July, but then all the growths that had reached the surface were of the usual form. In the Fenland drains it is common in spring, at which season the vegetation is frequently kept submerged by the artificial stream caused by drainage mills; as soon as the water is left at rest the leaves reach the surface and become coriaceous. Mechanical submersion, in one case by crowding, in the other by motion of the water, causes the attenuated shape and pellucid texture of the leaves. In a glass vase I once noticed a membranous floating leaf of *P. nitens* which was accidentally lifted on to the edge of the vase, and so partially exposed to the air; it became opaque and semi-coriaceous in a single day, resuming its transparent membranous texture after being replaced for about the same time in the water. This I repeated experimentally several times, always with the like result.

Seeing, then, that the shape and texture of the leaves of this form are due to temporary conditions, and revert to the ordinary type of the species during the life of the individual plant, I think we may safely erase the "*var. proluxus*" from our lists.

On referring to Koch's description, the first part of it will be seen to apply to a somewhat different state of *P. natans*. "*β. proluxus*, caules et petioli in rivulis valde elongantur et folia in rivulis strenua fluentibus saepe angustiora oblongo-lanceolata fiunt." Specimens well agreeing with this part of Koch's description were found by Mr. C. R. Billups in St. Helen's Canal, Hulme, Lancashire, October 15th, 1886. These have phyllodes 2 ft. long, and petioles 23 in., with narrowly elliptical or spear-shaped lamina 2-2½ in. long. This is the form described as *prolaxus* of Koch in Dr. Morong's Naid. N. America, p. 14. Visiting the Lancashire locality two or three years later, Mr. Billups found the water of less depth, and all the plants of *natans* had reverted to the usual form.

When growing in shallow water, which becomes dried up in summer, *land-forms* are produced, these are reduced to a short-stemmed or nearly stemless tuft of broad coriaceous leaves, but these are preceded in all instances by a pair of the narrow phyllodes, always longer than the broad leaves; if watched from the first these phyllodes will always be seen (plate 2, fig. 3), but as they soon decay they are seldom noticed in the specimens one generally sees in collections. When the *land-form* is produced entirely in the open air it does not flower; all the so-called "*flowering land-forms*" of *P. natans* that I have seen evidently began their growth in water, and this rapidly wasting away they had subsequently assumed the *land-form*.

P. natans has been recorded from all vice-counties except Dorset, South Essex,

study, chiefly of the living plants; in the few instances in which I have not been able to make direct observations I have acknowledged my indebtedness to others.

My friend, Mr. Arthur Bennett, has most generously given me his latest information as to the vice-comital distribution of the species through the British Isles, and such help as to priority of nomenclature as I demanded; at the same time he is in no way responsible for the use, or mis-use, of the valuable aid he has so freely given. I may in some instances use the more generally accepted specific names instead of those which, though prior in date, have neither intelligible descriptions nor traditional usage to support them; thinking it perhaps better not too hastily to set aside well-known names in favour of those which their authors failed to make known by either clear descriptions, good illustrations, or by the distribution of specimens among their contemporaries. I will here merely add that when my friend has published his complete nomenclature of the genus I will gladly be the first to accept his final decision, having perfect confidence in his unequalled knowledge of the subject, and his unrivalled power of specific discrimination in this difficult genus.

1. **POTAMOGETON NATANS**, *Linnaeus*, *Species Plantarum* (1753).

P. NATANS, Smith, Eng. Flor. vol. i. p. 229. Bab. Man. ed. 7, p. 371. Hooker, Stud. Flor. ed. 3, p. 431. Syme, Eng. Bot. vol. ix. p. 26, pl. 1399. Cham. Linn. vol. ii. p. 217, No. 20, Tab. 6, fig. 20. Fieber, Pot. Böehm, p. 22, Tab. 2, fig. 9. Fries Nov. Suec. p. 28. Reich. Icon. Germ. vol. vii. p. 26, pl. 50. Irmisch, Pot. p. 5, Tab. 1, f. 1-16. Boreau, Fl. Centr. Fr. ed. 3, vol. ii. p. 598. Gren. et Godr. Fl. Fr. Morong Naiad. p. 13, pl. 25. E. F. Cooper, Leicestershire Potamogetons, p. 3. Fryer, Journ. Bot. vol. xxiv. p. 237. Lond. Cat. ed. 9, No. 1580. P. Raunkiaer, Danske Blom. p. 68, fig. 31. Ascheron et Graebner Synop. Mittel Europ. Flor. p. 303. (1.) Tiselius, Potamog. Suec. Exsic. Nos. 1-4.

SPECIFIC CHARACTER.—Leaves all coriaceous, floating; upper jointed to the petiole by a narrow, flexible, discoloured part of the midrib extending a little below the lamina; lower reduced to a long narrow petiole and midrib (phyllode), exceeding the upper and ultimately floating.

DESCRIPTION: Rootstock stout, with terete stolons not contracted at the nodes, white with numerous slightly-raised reddish glandular spots subspirally arranged.

STEM stout, terete, ascending from alternate internodes, sparingly branched, often simple, the basal internode faintly glandular, the glands becoming immersed in the second and subsequent internodes.

LEAVES all stalked coriaceous entire flat; upper with a flexible discoloured joint at the base of the lamina, many-ribbed, with the principal ribs prominent on the underside, often with a strong fold at the base, cordate, or abruptly or gradually

narrowed into the petiole; lower reduced to a narrowly linear phyllode convex on the upper side for the lower half but slightly channelled towards the tip, much exceeding the upper leaves in length, ultimately floating, and then sometimes spatulate or clavate towards the apex; these linear leaves are succeeded by others of an intermediate character which are expanded into a short lanceolate or spear-shaped lamina (usually jointless) and so passing into the ordinary coriaceous elliptical to orbicular floating leaves, but soon decaying like the typical linear leaves.

STIPULES very long, often exceeding the internodes, acuminate, scarious or herbaceous, with strong longitudinal fibres, back with two prominent ribs forming a slightly-grooved keel, soon decaying, but the fibrous part persistent.

PEDUNCLES slightly thicker than the stem, terete, slightly flattened towards the base and channelled, narrowed towards the spike, shorter than the subtending leaves, 2-3 times longer than the spike.

SPIKE thick, cylindrical, densely flowered, $1\frac{1}{2}$ -2 inches long.

FRUIT obovate, slightly compressed, usually rounded on the back, rarely with an acute keel when fresh; acutely keeled when dry, with faint lateral ridges; dorsal margin much rounded, upper slightly convex, beak short, facial.

COLOUR shining dark green, or sometimes bright green, the young leaves streaked and tinted with reddish purple.

HABIT very robust, stem 1-6 feet long, growing in dense masses, which often completely hide the surface of the water. The clusters of phyllodes are often produced late in the autumn, and grow slowly throughout the winter; in spring, as the stems lengthen, the intermediate spear-like leaves begin to appear, together with the broad upper floating leaves, the latter being involute until they reach the surface, while the linear and spear leaves are flat throughout their growth. When the mass of vegetation reaches the surface the lower leaves soon decay. The permanent broad upper-leaves emerge from the water while still involute, and about half their length is exposed to the air for a few days, when, their growth being nearly completed, they expand, and, sinking back, float on the surface of the water, every part of which they are enabled to occupy by means of their flexible joint, which allows each leaf to twist in any direction, even reversely parallel to its own petiole. The involute emerged leaves are conspicuous from a considerable distance, and serve to distinguish *P. natans* when growing from its associated species.

Several named varieties of *P. natans* have been described by authors; of these it will be necessary to consider only those which have at some time found a place in the writings and records of British botanists. After a careful study of these forms, and the cultivation of most of them, I have come to the conclusion that, so far as this country is concerned, all these so-called varieties are mere *states*, some of which are merely the natural changes of form, which take place according to the progressive

FLOWER-SPIKE usually emerged until fertilization is accomplished, then it falls back, and the fruit is ripened at or beneath the surface of the water; very rarely fertilization has been observed to occur in wholly submerged spikes.

FLOWERS protogynous, as far as I have been able to observe, and probably fertilized by the pollen being discharged with some force from a contiguous spike, or by the action of insects, or by the wind, or even by being carried by the water itself. In the latter case the occasional fertility of constantly submerged flowers is easily explained.

The FRUIT affords good characters to aid in the discrimination of species, but it is difficult to observe, especially when fresh. When dried the external sculpture becomes more pronounced; care, however, must be taken to found comparisons on perfectly mature examples. Much importance is attached by some botanists to the exact degree of curvature of the embryo, and the relative thickness of the radicular and cotyledonary ends, but these features are variable, as well as the external form; and I believe this slight degree of variation is not solely due to age or degree of maturity, but is often inherent, like the variability of the plants themselves. Hence the distinctions afforded by the fruit in closely allied or "critical" species are of little practical use to the average observer. If a comparison is made of the figures and descriptions of the fruits of Potamogetons given by Chamisso, Fieber, Reichenbach, and Morong, the difficulty I have pointed out will become very evident.

The drupelets are ovoid or subglobose, somewhat compressed, keeled, or rounded on the lower margin, with two fainter lateral keels; the nutlet is 3-keeled; the embryo incurved, uncinate or cochleate, and always with the cotyledonary end somewhat ascending, so as to form part of a spiral; radicular end thickened.

Although the species differ so widely in external appearance, the genus does not readily afford distinct groups, nor subgenera; the arrangement usually followed is that of Koch, who has divided the species into the following groups:—

1. HETEROPHYLLI; comprising the broad-leaved species with coriaceous floating leaves, from *P. natans* to *P. nitens*.

2. HOMOPHYLLI; species with all the leaves membranous and submerged; from *P. lucens* to *P. crispus*.

3. CHLOEPHYLLI; species with strap-shaped, grass-like leaves, from *P. Friesii* to *P. trichoides*.

4. COLEOPHYLLI; species with adnate stipules, the bases of the leaves sheathing the stems.

5. ENANTIOPHYLLI; leaves all opposite, *P. densus*.

Looking at the Potamogetons of Europe these are fairly distinct groups, with the exception of one aberrant species, *P. crispus*, which is in many respects inter-

mediate between the *Homophylli* and *Chloephylli*; but if we try to group the Potamogetons of North America in the same manner we find in *P. spirillus* a species which unites in itself the characters of Groups 1, 3 and 4. Even in Europe the Groups 1 and 2 have some intermediate forms, as *P. alpinus* for instance, which is sometimes without coriaceous floating leaves. In other instances, as *P. decipiens* with coriaceous floating leaves, the intermediate forms are hybrids between species belonging to different groups, and so naturally possess characters derived from each. These points will be considered more fully when the species are described in detail.

In the following pages those characters which either separately or in conjunction seem peculiar to each species are grouped together to form a *specific character*, which is placed before the more detailed description. The student, it is hoped, may be able, by means of these short descriptions and a reference to the plates, to name the ordinary states of any of our British species. The detailed descriptions are drawn up in their present form for the purpose of affording a means of readily making a critical comparison of any one part of the plant throughout the whole series of species without the necessity of reading the whole of the descriptions.

For the more or less permanently differing variations of those species which are capable of producing fertile seed the term *variety* is used; for temporary differences (often more striking than permanent variations) depending on age of the individual or unusual local conditions, and which naturally during growth, or on the plant being restored to more natural surroundings, revert at once to the typical form, the term *state* is applied.

Among the older and generally accepted species we find some, such as *P. nitens* Weber (1787), and *P. fluitans*, Roth (1788), which have recently been proved to be hybrids. To reject these names would lead to great confusion, and would tend greatly to obscure our researches into the origin of many specific or quasi-specific forms; in retaining them I also think it well to name newly-discovered hybrids, prefixing a $\times$ to the generic name ($\times P. fluitans$) to distinguish these forms from true species.

The first described form of such hybrid species is taken as the type; the departures from which are of two kinds:—when an individual plant of one of these hybrids has been observed to sport into a permanently distinct form, the term *variety* may still be applied; but when a distinct result is obtained by the interbreeding of the same parents, or in cases where the parents are reasonably supposed to be the same, the difference is expressed in the following pages by the Latin word *forma*.

No attempt is made at unravelling the intricacies of synonymy; the references to other writers are given solely to direct the student to the best sources of information. The specific descriptions and life-history of each species are made from my own

Denbigh, Flint, West Lancaster, Durham, Isle of Man, Kirkcudbright, and Banff; probably it occurs in all or most of them. It is frequent in Ireland, common throughout Europe and North America, and has been found in several localities in Asia and Africa; although reported from Australia and New Zealand, it is probably absent from Australasia.

Explanation of the Plates. POTAMOGETON NATANS. Plates 1—3.

PLATE 1, fig. 1, fruiting spike; 2, drupelet mag.; 3, longitudinal section; 4, transverse section; 5, section of the stem.

PLATE 2, fig. 1, barren summer shoot, to show phyllodes and underside of floating leaf; fig. 2, land-form, early state with phyllodes. (All the above are drawn from Cambridgeshire specimens); fig. 3, "var. *linearis*," from specimens gathered by Mr. R. W. Sculley at Long Range, Killarney, "from slow-flowing water," selected to show the branched stem with persistent phyllodes and the rarely-formed upper leaves. Specimens from "swift-flowing water" do not show these leaves.

PLATE 3. The lower figure on the right is from Mr. T. F. Mott's original specimen gathered by him, "May, 1881, pond near Scraftoft Long Spinny"; the central and left-hand figures are from the same Leicestershire locality, gathered by Mr. E. F. Cooper, "May 5th, 1892."

2. × **POTAMOGETON CRASSIFOLIUS**, *Fryer*, Journal of Botany, vol. xxviii. p. 321, Tab. 299 (November, 1890).

P. NATANS × *ZIZII*, Aschers. et Graeb. Synop. p. 332 (1897).

SPECIFIC CHARACTER: a barren hybrid between *P. natans* ♀ and forms of *P. coriaceus* ♂ (*P. Zizii*, Auct.). Leaves all stalked, coriaceous when mature; lower leaves fleshy, soon decaying, distant, few in number, upper oblong-oval or obovate, often strongly auricled at the base of the lamina, not completely floating, but often partially submerged like those of *P. coriaceus*, phyllodes spatulate, shorter than or little exceeding the internodes; stipules acute, not winged nor channelled on the back, whitish, with two conspicuous green ribs, horny with scarious edges, or green and herbaceous, the lowest rarely with a short adnate phyllode, or a short ordinary lanceolate lower leaf.

DESCRIPTION: ROOTSTOCK stout, white with reddish glands, stolons with short internodes, scarcely contracted at the joints, deeply rooting.

STEMS stout, terete, 1–5 feet long, simple below, shortly branched above, or simple throughout, irregularly ascending at intervals of several internodes, frequently terminal.

LEAVES all stalked, coriaceous, opaque when mature; lower fleshy and translucent when young, lanceolate, gradually narrowed into the petiole, lowest reduced to linear or spatulate phyllodes shorter than or little exceeding the internodes, all

soon decaying, upper leaves oval, or oblong-oval, rarely elliptical, usually strongly auricled towards the base, with the lamina decurrent below the auricles; lateral ribs translucent, not sunk in the substance of the leaf, nor usually prominent beneath. The lowest leaf of each branchlet is often lanceolate or oblanceolate. All the leaves are thick and fleshy when fresh, but extremely thin when dried, thus resembling those of *P. coriaceus*.

PETIOLE shorter than the lamina, flat above, expanded towards the leaf, herbaceous and veined with transverse veins throughout, thin when dried, like the leaves.

STIPULES acute, horny with scarious edges, white with two conspicuous green ribs, but not winged nor channelled on the back, or green and herbaceous, $1\frac{1}{2}$ –2 in. long; lowest sometimes bearing a short phyllode, or more rarely a somewhat longer ordinary lower leaf.

PEDUNCLE equal, terminal or becoming lateral by continued growth of the branch; equalling or slightly exceeding the stem in thickness, 2–3 times longer than the flower-spike, much shorter than the subtending foliage.

SPIKE cylindrical, dense, $1-1\frac{1}{2}$ in. long, barren, sepals remaining closed, but stigmas exerted, soon decaying.

FRUIT abortive, very rarely formed, but never perfected, whole spike soon decaying.

COLOUR bright green, the young upper leaves often streaked with reddish-purple, like those of *P. natans*.

HABIT: lower leaves permanently submerged, usually decayed by the time of flowering; upper leaves sometimes raised above the water till fully expanded, like those of *P. natans*.

The plant often grows throughout the winter, like *P. natans*, which it resembles in its general habit.

In the Journal of Botany I have described two varieties of this hybrid species (l.c. p. 323), one of which I have had under cultivation for eight years. It never wholly dies down like *P. fluitans*, but the submerged leaves and stems grow slowly all through the winter. Although it produces flower-spikes in abundance, these rarely produce even abortive drupelets, but decay soon after the stigmas are exerted. In this form the upper leaves are regularly spread out and flat, orbicular-ovate, rarely lanceolate, with prominent ribs and with the purple colouring beneath of *P. natans*, and the whole plant is a little stouter in substance when dried than in the typical form.

The "Westmoor variety" I have now decided to name as a distinct form, possibly of somewhat distinct parentage:—

× *P. CRASSIFOLIUS*, FORMA VERRUTA mihi. Plate 5.

Hybrid, *P. natans* ♀ × *P. Zizii* ♂. Stem branched from the base or simple, submerged leaves oblanceolate or lanceolate, very long, sometimes exceeding the floating coriaceous leaves, phyllodes sometimes produced along the stem nearly up to the floating leaves, and then exceeding them in length; floating leaves lanceolate, oval, or oblong-oval, with the petiole often much longer than the lamina, which occasionally bends back as in *P. natans*, but without a perceptible joint, rarely auricled at the base, extremely variable in shape, and length of petiole; peduncle stouter than the stem, greatly exceeding the very short spike. Habit of *P. natans*, or of the flowering branches of *P. Kirkii*. A comparison of the figure of this form on plate 5 and that of *P. Kirkii* on plate 8 will show the necessity of studying the early states of Potamogetons as well as the more perfect or flowering states. Much of the lower part of *f. verruta* might well pass for *P. Kirkii*. The young state, however, is totally different, being nearer that of typical *crassifolius*; while the occasionally elongate phyllodes and narrowly lanceolate lower leaves exceeding the short-petioled upper leaves, in some of the plants, approach the habit of *P. natans*.

The plant figured is from a cultivated root, which was gathered in a shallow ditch at Westmoor, Doddington, Cambridgeshire; it was one of many seedlings which I was able to watch for several seasons. A full account of the circumstances under which they grow will be found in the Journal of Botany, November, 1890, p. 322. Since that was written a single plant was found in a flowering state; it is so totally different in facies that I regret the limits of our work forbid a separate plate. If I had not been able to connect the flowerless and flowering form by numerous intermediate forms, I could not have even guessed that they were of the same origin, and many of them probably from the same cross-fertilized spike. In naming new hybrid-species in this work I am guided solely by their parentage, not by likeness to one type in themselves, nor by difference from already named forms. Else the forms of *verruta* might have had half a dozen names, as they are so dissimilar in appearance, resembling *Kirkii*, *natans*, *Zizii*, and with additional special forms of their own.

The hybrids of Potamogetons are not like the hybrids of less variable plants; in some instances they may be constant enough to one type, but in others the variation seems to have no bounds. We find suggestions of not two, but many species, and it is impossible to suppose that these resemblances are accidental or uncontrolled by law. When by observation and actual experiment the law that governs the variation of cross-bred forms of this genus is discovered, wholly or in part, we shall probably get a clearer view of one of the modes by which "species" may have originated. The plants in this group of forms allied to *P. fluitans* are infertile hybrids, later on we shall have to consider fertile hybrids. In what respect do they differ from "true species"?

DISTRIBUTION: *P. crassifolius* is at present only known from three separate localities near Chatteris, Cambridgeshire. Through the kindness of Mr. J. Baagöe, I have received Danish specimens which suggest a similar parentage, but owing to the extreme variability of some of the hybrids of the *fluitans*-group I dare not name them. All such forms can be determined only by exhaustive observation in the field; comparison of specimens is here useless, or even misleading. No doubt other different-looking hybrids between *Natans* and *Zizii* will be found, but from what has been observed in other extremely variable species there is little doubt that some common character will be found by which they may be separated from *P. fluitans* as clearly as they now are from *P. Kirkii*.

When *Natans* grows intermixed with both *lucens* and *Zizii* it will be necessary to study hybrid seedlings that may be found, from their earliest stages.

Explanation of the Plates. × POTAMOGETON CRASSIFOLIUS, Plates 4 and 5.

PLATE 4. Rootstock, phyllodes, lower and upper leaves; fig. 1, lower stipule with adnate leaf; 2, section of stem, mag. All from Mepal Common, Cambridgeshire, July, 1897.

PLATE 5. *forma verruta, mihi*; a complete barren branch from root upwards; cultivated plant from Westmoor, Doddington, Cambridgeshire, August 8th, 1892. Peduncle and flower-spike, wild, from same locality, September, 1890.

3. × **POTAMOGETON FLUITANS**, Roth. Tentamen Floræ Germanicæ, vol. i. p. 72 (1782), vol. ii. p. 202.

P. FLUITANS, Cham. et Schlecht. Linnæa, vol. ii. p. 219, No. 21 (1827). Fieber, Pot. Böehm, p. 21 (1838). Fries Novit. Suec. ed. 2, p. 29. Koch, Synop. Fl. Germ. ed. 2, p. 776. Reich. Icon. Germ., vol. vii. p. 26, Tab. 49 (excl. *β stagnatilis*). Fryer, Journal Bot., vol. xxvi. p. 273 (1888). Beeby, Journal Bot., July, 1890. Raunkjær Dansk. Blomster, p. 97 (1896). Tiselius, Potamog. Suec. Exsic. No. 6.

SPECIFIC CHARACTER.—Stem branched from the base, lower leaves membranous, lanceolate, persistent; upper oval-lanceolate coriaceous, all long-stalked; stipules horny, herbaceous, persistent; flower-spike barren; a hybrid between *P. natans* ♀ and *P. lucens* ♂.

DESCRIPTION: ROOTSTOCK with crooked stolons, internodes somewhat thickened and sub-tuberous, striking deeply into the soil.

STEM moderately stout, terete, ascending at irregular intervals, much branched from the base, or simple in late growths; lower branches submerged, not flowering, upper branches floating and spreading on the surface of the water.

LEAVES all stalked, lowest reduced to narrowly linear, short phyllodes permanently submerged, followed by linear-lanceolate, or narrowly lanceolate, or elliptical-lanceolate membranous translucent submerged leaves, somewhat undulated,

flat or longitudinally folded, acuminate, with the lamina insensibly narrowed into the petiole; upper leaves coriaceous, floating, oblong-lanceolate, or elliptical-lanceolate, with the numerous lateral ribs usually translucent and immersed in the substance of the leaf on the underside, or with 2-3 on each side prominent towards the base of the lamina; sometimes slightly folded towards the base, but these with a part of the lamina decurrent below the fold.

PETIOLE slightly convex above, always shorter than the lamina.

STIPULES very long, 1-4 in., deeply channelled on the back, or sometimes slightly winged, acute, persistent, at first clasping, afterwards patent; horny in texture, often herbaceous, green or whitish with green veins; rarely with a leaf reduced to a phyllode on the back.

PEDUNCLE 3-4 in. long, a little thicker than the stem, equal, or slightly swollen in the middle.

SPIKE cylindrical, 1-1½ in. long, densely flowered; sepals rarely expanding, and then bright green, or closed and dull green, but the stigmas always exerted.

FRUIT abortive, 3 keeled, the central keel acute.

HABIT: The whole plant rising by the time of flowering, and spreading in fan-like masses on the surface of the water, the leaves never reflexed nor irregularly disposed as in *P. natans*. By this time most of the translucent membranous lower leaves become coriaceous like the true floating leaves which support the flower-spike, only a few of the lower leaves remaining submerged and membranous. In autumn long narrowly linear folded grass-like leaves are often thrown out from the lower part of the stem, especially when the plant has been cut over repeatedly in the summer; but these are quite distinct from the phyllodes which are constantly produced on the young stems in spring. These submerged phyllodes gradually pass into the ordinary membranous lower leaves by intermediate bodkin-pointed leaves with the petiole and midrib thick and fleshy, but expanded in the middle into a narrow wing-like lamina as in *P. lucens*.

COLOUR bright green to olive-green, drying darker.

This plant was first separated from *P. natans* by Roth in 1783, and described in his *Beyträge zur Botanik*, but not named until 1788 in his *Tentamen Floræ Germanicæ*. In reading his excellent description of the species it is necessary to remember that *P. polygonifolius* was at that time still confused with *P. natans*; bearing this in mind Roth's description is so exact and clear that it is not a little remarkable how much error and confusion has gathered around such a well-defined form.

Mr. W. H. Beeby in an excellent paper in the *Journal of Botany* (July, 1890) made the first decisive step towards clearing up the difficulty by showing that the barren hybrid form figured by Reichenbach (*Icon. Flor. Germ. Tab. 49*) exactly

agrees with Roth's description ; and he also points out the difference between the specimens seen by him of the fruit-bearing European plant (often named "*fluitans*") and the plant figured by Reichenbach :—" My contention, therefore, is : First, that the demand for fruit, wherewith to prove that any plant is *P. fluitans* Roth, must be withdrawn, or the evidence whereon the demand is based be divulged ; and secondly, that the name *P. fluitans* Roth must be withheld from the freely-fruited Continental plant, until it is shown to produce the characteristic submerged leaves described by Roth. . . ."

Subsequently Dr. Tiselius generously lent me his fine series of the fruiting plant collected by himself in the River Neckar, and also gave me a very perfect set for my own herbarium. By these means I was enabled to compare the two forms, in both early and mature stages of growth, and I at once saw that the two plants setting the question of fruit altogether aside, were distinct in foliage and habit.

The fruiting plant from the River Neckar* has, indeed, a few translucent submerged leaves, but they are more distinctly stalked than in the plant of Roth, and are all decayed by the time of flowering. No linear-lanceolate leaves nor phyllodes are present. The upper leaves are oval or oblong-oval, with the petiole often greatly exceeding the lamina, which has the lateral ribs raised above the surface on the underside, not immersed in the substance of the leaf as in *P. fluitans*. The lateral ribs are opaque even in young examples, instead of translucent, and the areolation is totally different.

The rootstock is more like that of *P. polygonifolius*, as also are the stipules, and the stem is simple, or rarely with lateral branchlets ; finally, the habit of the fruiting plant is more diffuse, the foliage is more scanty, and the floating leaves do not spread like a fan.

As far as I am capable of judging, the European fruiting plant is not specifically identical with *P. Lonchites* of Tuckerman, neither in foliage nor in fruit ; in the latter both external sculpture and curvature of the embryo seem different to my eyes.

Before leaving the subject I would call attention to Mr. Bennett's Notes on Potamogetons, Journal of Botany, October, 1893 ; he there says :—" We have no certain knowledge of any of Roth's species being preserved in any herbarium ; but there are at Munich specimens in Schreber's herbarium, named as such, and gathered in ' In Seebach, 1775,' and others ' In Seebach, 1782.' It seems to me a reasonable inference that these specimens are from (or seen by) Roth ; the more so

* Afterwards considered by Dr. Morong to be equal to *P. Lonchites*, Tuckerman, var. *Novæboracensis* Morong, Naiadaceæ N. America, p. 21.

because there are other species in the same collection actually received from Roth, and signed by him. They are the plant we call *fluitans* in England (hybrid?), and not the Neckar plant of Schimper and Dr. Tiselius."

I will now try to solve Mr. Bennett's query as to the hybridity of Roth's *P. fluitans*, especially as it will show the nature of the evidence on which I have assumed certain plants described in the following pages to be natural hybrids. This one case clearly stated will save much needless repetition.

P. fluitans is found, according to Mr. Bennett, in "Europe generally"; although having so wide a range, it has never been found in a fruiting state, although it has been repeatedly gathered and observed for over a hundred years; this alone affords strong presumptive, but not positive, evidence of hybrid origin; but if we add to this strong presumptive the more positive evidence afforded by its local distribution and its sudden appearance in new localities in the fens of Cambridgeshire and Huntingdonshire, I think no one can doubt its origin from *P. natans* crossing with *P. lucens*.

When I first found it in 1884 it was growing in an isolated pit only a few yards square with the two supposed parents and *P. crispus*, but as the station is a low, swampy excavation by the side of Vermuyden's Drain, and is liable to be submerged by floods from the upper or "high-lands," I naturally supposed that at some time or other the species had been carried thither from, or by means of, the brook which brings down the upland floods. A repeated search year after year clearly showed that both brook and river above the station of *fluitans* were not tenanted by the species. In the course of two or three years I found other stations where *fluitans* occurred, evidently from seeds, in single plants in ditches and drains which were far removed from, and had no connection with, one another. The next step was to try if this form, which had not been found seeding in a wild state, could be induced to do so under cultivation; it grew freely, but the flower-spikes remained barren, and I began to feel sure that *fluitans* was a hybrid. All doubt was removed when I found a plant growing in a drain which had been recently deepened, and most of the vegetation destroyed, in a spot which had been constantly examined by me for years. This plant when first found was evidently of recent origin, and then confined to a single rootstock, and it was produced under circumstances which I have repeatedly found favourable to the intercrossing of distinct species of Potamogeton; that is, by the drain having the greater part of its vegetation destroyed, so that single flower-spikes grew apart from others of their own species. As a rule, the stigmas are exerted and receptive before the pollen on the same spike is ripe; if no other spike of its own kind is near to supply the necessary pollen, then the stigmas may become fertilized by a contiguous polliniferous spike of some other species. At one time I thought this could only take place by mechanical, and so accidental, means,

but more recent observations show that the ripe pollen is discharged in a shower some little distance around the spike; thrown around it, in fact, with sufficient force to reach many or all of the stigmas in a contiguous spike.

Usually only one or two druplets are thus made fertile, but the occurrence of hybrid seedlings along one ditch in the before-mentioned case of *P. crassifolius* induced me to believe that occasionally the whole or a greater part of a spike was fertilized by the pollen of another species.

P. fluitans never occurs in the fens apart from *P. natans* and *P. lucens*; in like manner *P. crassifolius* never occurs apart from *natans* and *Zizii*; and generally when a drain or ditch has been cleaned out and deepened, so that the vegetation has been for the greater part destroyed, I have found after a year or two seedling, or apparently seedling, plants of hybrid Potamogetons. Short of producing these forms by artificial crossing, it seems impossible to obtain stronger evidence of their hybrid origin. When we further consider how exactly *P. fluitans* is intermediate in many of its characters between *P. natans* and *P. lucens*, as was long ago suggested by Chamisso and Schlechtendal and by Fries, it seems impossible to doubt its being a true hybrid, constantly reproduced from the crossing of these two species.

At present *P. fluitans* is only certainly known as a British plant from the counties of Sussex and Surrey, where it has been found by Mr. Beeby, and from Huntingdon and Cambridge. Mr. Druce sent me specimens from Berkshire, which probably will be found to belong to the fluitans-group, but they are not sufficient to name with certainty.

Out of England this hybrid species is at present known only from "Europe generally." See Mr. Bennett's valuable note in the Journal of Botany for October, 1893, and vol. xxxiii. p. 372 (1895).

Description of the Plates, 6-7. P. FLUITANS Roth.

PLATE 6. Upper part of flowering branch of *P. fluitans*, from a specimen gathered in Huntingdonshire, September, 1893. To get so large a portion in the plate it was necessary to disarrange the fan-like disposition of the upper leaves; all the leaves were strongly undulated on the margin, some of the lower ones so much so as to look serrulate; in this Mr. Morgan has exactly reproduced their natural appearance in the fresh specimen! This undulation disappears to a great extent when the plant is dried under pressure; all these leaves had entire margins, although some of them were almost *crisped* when fresh.

PLATE 7, fig. 1. Part of the rootstock showing young growths with phyllodes and lower membranous leaves; fig. 2, the Land-form, with linear lower leaves; fig. 3, axillary branchlets by means of which the plant is propagated in autumn.

4. × **POTAMOGETON KIRKII**, Syme, Eng. Bot. ed. 3, vol. ix. p. 31 (1869).

Hooker, Stud. Flor. ed. 3, appendix p. 535. A. Bennett, Ex. Club Report, 1895, p. 496 (1897). Fryer, l.c. p. 497.

P. sparganifolius, "Bab. Man. ed. 6, p. 363 (*ex parte*) non Läst! nec Fries!" Syme, E. B. vol. ix. p. 31. Bab. Man. ed. 7, p. 372. Lond. Cat. ed. 8, No. 1484, ed. 9, No. 1585. *P. polygonifolius*, c, Kirkii, Lond. Cat. ed. 7, No. 1219.

SPECIFIC CHARACTER.—Lower leaves submerged, "sessile," very long linear membranous "with several rows of greatly elongate-cancellate areolations along the midrib"; upper leaves like the lower but shorter, lanceolate or oblanceolate, or very long stalked and floating, coriaceous, oblong, or elliptical, or lanceolate. Flower-spike barren, short, on slender long equal peduncles, borne alike on branches with submerged or floating coriaceous leaves. Probably a hybrid; between *P. natans* ♀ and *P. polygonifolius* ♂?

DESCRIPTION: ROOTSTOCK not known.

STEM terete, much branched, "the lower branches barren."

LEAVES: lowest submerged sessile (or stalked?) 6–15 in. long by $1\frac{1}{4}$ in. wide, membranous, pellucid, bright green, attenuated gradually into the base, more abruptly so at the acute apex, entire, with 8–14 lateral ribs with distant irregular transverse veins; midrib bordered by several rows of elongate cancellate areolations forming a more or less conspicuous band along the middle of the leaf (the leaves decayed by the time of flowering?). On the upper fertile branches the leaves are of two kinds, neither of which is essential to the production of flower-spikes, usually submerged and of the same structure and substance as the lower leaves, but more distinctly stalked, the lamina passing insensibly into the petiole, clustering, flabellate; floating leaves coriaceous or subcoriaceous, very long stalked, oblong or elliptical, lamina rounded and abrupt at the base, sometimes slightly auricled, or lanceolate and gradually narrowed into the petiole, lateral ribs opaque, midrib slightly discoloured and prominent at the base as in *P. polygonifolius*, and all the leaves with areolations as in that species.

PETIOLE 3–5 inches long, greatly exceeding the lamina, somewhat herbaceous and indistinguishable from the lamina in the lower, and lanceolate upper leaves.

STIPULES "very long, blunt, not winged on the back," but with two distinct green ribs forming a decided keel, "scarious with numerous fibres," or herbaceous.

SPIKE cylindrical, densely flowered, sepals not expanding, $\frac{3}{4}$ in. long, flowers barren.

PEDUNCLE 3–4 in. long, slender, equal.

COLOUR light green, upper leaves darker green, drying darker, "upper leaves tinged with reddish brown."

HABIT: apparently with linear lower leaves first springing in tufts like the phyllodes of *P. natans*, succeeded by greatly branched stems with or without coriaceous floating leaves on the upper branches; the flower-spikes are produced as in *P. heterophyllus*, independently of the coriaceous leaves.

I have copied much of Dr. Syme's original description, because it would be difficult to improve it in the absence of living specimens, which unfortunately neither of us could obtain. A root grown by anyone accustomed to study living plants of *Potamogeton* would solve much that is now obscure in the life history and relations of this remarkable form.

Probably it is a hybrid; it seems to fall naturally under the *fluitans* group, from some of which some of its states are hardly distinguishable; the lower part of our plate 5, *P. crassifolius forma verruta*, might almost pass for the form of *P. Kirkii* figured in plate 8. I regret we could not figure a flowering example of this state, which is well drawn in Syme's figure, E. Bot. ed. 3, plate 1403.

For a long time English botanists erroneously considered *P. Kirkii* to be the *P. sparganifolius* of Laestadius; this mistake arose from attaching too great importance to the long narrow lower leaves; these, when dry, are often very narrow and perhaps more strongly suggestive of *P. sparganifolius* than those figured in our plate. When we compare the mature plants we find the flowers and spikes are very distinct, those of the latter plant being broad and less densely flowered, with expanded sepals, and fertile, the lower leaves are brighter green, less numerous, not spreading like a fan; and the floating leaves are long-lanceolate and less coriaceous (see Tiselius, Pot. Suec. Exsic. No. 5).

Dr. Tiselius' *P. dubius* (l.c. No. 19) is much more like our plant, but, as he suggests, is rather *P. graminens* $\times$ *natans* than *P. natans* $\times$ *polygonifolius*. Other specimens from Dr. Tiselius, however, of *P. dubius* approach our *P. Kirkii* much more closely, but all seem to lack the peculiar lower submerged leaves.

Mr. J. Baagöe sent me a beautiful form collected by himself in Denmark, with the name of "*P. fluitans forma rivularis Lange*," which seems to connect our species with *P. fluitans*. But it is always difficult, and often unsafe to judge from dried specimens, especially when not prepared in a similar manner. In Mr. Baagöe's specimen the leaves are nearly all reduced to short submerged phyllodes along the whole length of the stem, even up to the base of the terminal flower-spike. This remarkable form may result from *P. fluitans* growing in a rapid stream, or not improbably it may have a somewhat different parentage from typical *P. fluitans* of Roth.

Dr. Syme (l.c. p. 32) says of *P. Kirkii*, "I have some suspicion that it may be a very luxuriant abnormal form of *P. polygonifolius*." In addition to the reasons he then gives for doubting this, we may add the production of the flower-spike in the absence of floating leaves.

LOCALITIES : At present this hybrid-species is only known from Ireland, where it was "found by Mr. Thomas Kirk in the Ballinabrack River at Maam, co. Galway" (Syme), "Loch Neagh by Dr. D. Moore," Lough Corrib, Galway, T. Kirk, Sept. 7th, 1854.

Description of the Plates.

PLATE 8, upper branch with submerged leaves, this state sometimes produces flower-spikes, Maam, co. Galway, Coll. W. R. Linton, August 13th, 1885; Herb. C. Bailey.

PLATE 9, flower-spike on branch with floating leaves, Maam, co. Galway, August 13th, 1885; Herb. E. F. Linton. Maam, co. Galway, Coll. W. R. Linton, August 13th, 1885; Herb. C. Bailey. Lower leaves, Maam, co. Galway, Coll. W. A. Shoolbred, July 5th, 1895; Herb. C. Bailey.

5. **POTAMOGETON POLYGONIFOLIUS**, *Pourret*. Act. Toulouse, iii. p. 325 (1734).

P. oblongus, Viviani, Anal. Bot. 2 (1802); Cham. et Schlecht. Linn. p. 214, No. 19. Fries Novit. ed. 2, p. 29. Fieber, Pot. Böhmens, p. 19. Koch, Fl. Germ. vol. ii. p. 775. Crepin, Fl. Belge Fragm. Fasc. i. p. 25.

P. polygonifolius, Reich. Germ. vol. vii. p. 24, Tab. 44. Bab. Man. ed. 2, p. 371. Hooker, Stud. Flor. ed. 3, p. 431. Syme, E. B. ed. 3, vol. ix. p. 27, pl. 1+00. Bab. Man. ed. 7, p. 371. Hooker, Stud. Flor. ed. 3, p. 431. Druce, Fl. Oxford, p. 282. Tiselius, Pot. Scand. Exsic. fol. 9, 10, 11.

SPECIFIC CHARACTER.—Leaves all stalked, lower submerged, membranous, upper floating, coriaceous or subcoriaceous; no linear phyllodes. Drupelets very small, yellowish-green when fresh, reddish when dry. Petioles of the coriaceous leaves often persistent after the decay of the lamina, and then falsely resembling phyllodes, but may be known by the lacerated end lacking the acute apex of a perfect leaf.

DESCRIPTION : ROOTSTOCK slender to stout, contorted, with many short internodes, with numerous fibrous rootlets, decaying in autumn except three or four terminal internodes, the terminal one bearing a small reddish winter bud; nodes annular, prominent.

STEM terete, ascending at irregular distances from the rootstock, sometimes from every node, or at intervals of 3–4 nodes, rooting at the lower joints and then throwing out stolons. In shallow-water forms the stem is only one or two inches long, or obsolete.

LEAVES all stalked; lower submerged, pellucid, membranous, gradually narrowed into the petiole, often absent in shallow-water forms; lowest rarely narrowly linear-lanceolate, but then being a linear lamina, not a phyllode; upper very long stalked

in deep water, short or long stalked in shallow-water forms, coriaceous, lamina suborbicular to lanceolate, cordate, or abruptly or gradually narrowed into the petiole; midrib and principal lateral ribs somewhat prominent on the underside; midrib conspicuous, prominent and somewhat discoloured at base of the lamina on the upperside, but never narrowed into a joint below the leaf as in *P. natans*.

PETIOLE coriaceous, or herbaceous, slightly convex above, in deep-water forms usually greatly exceeding, but in shallow-water forms often shorter than the lamina.

STIPULES long, subfibro-scarious, not keeled nor channelled on the back, subpersistent.

SPIKE short cylindrical slender, densely flowered and fruited.

PEDUNCLE slender, equal, many times longer than the spike.

FRUIT small, beak short, facial, dorsal margin rounded when fresh, keeled when dry with faint lateral ridges.

COLOUR "dull olive-green, often tinged with reddish brown" (Syme).

HABIT in deep-water forms like that of *P. fluitans*; in shallow-water forms with creeping rhizomes producing short stems at regular intervals, with the leaves all similar and coriaceous; very variable, many intermediate forms occurring between these two extremes, which insensibly pass into one another.

This species was confused with *P. natans* by Linnæus and most of the earlier authors, and by some of the later ones, as Bentham in his *Handbook of the British Flora* (1865). Perhaps this error arose in the first instance from the resemblance of the coriaceous floating leaves of some deep-water forms to those of small states of *P. natans*. The resemblance is merely superficial, and possibly the latter species approaches *P. lucens* almost as nearly as it does *P. polygonifolius*. When the Potamogetons of the whole world are better known I have little doubt that many species will be found falling in a natural arrangement between these two. The whole series of Australasian forms which have been assigned by authors to one or other of these two species will probably be found to belong to neither of them, but to constitute intermediate species. To evolutionists it will seem probable that the European flora has possessed and lost many such intermediates; so really distinct and apart are the two species originally mixed in *P. natans*. *P. polygonifolius* is so variable according to the depth of the water in which it grows, and is so susceptible to the influences of local conditions of every kind, that each district, almost, seems to produce a peculiar form. Hence, as may be supposed, many striking or extreme forms have had varietal names attached to them; or, at least, some attempt has been made to arrange the forms in natural groups. The best of these arrangements is that of the late Dr. Boswell Syme, in his excellent edition of *English Botany*, and is as follows:—

“Var. *genuinus* (Eng. Bot. pl. 1400).

“Lowest leaves narrower and thinner in texture than the upper ones, which are rounded or cordate at the base.

“Var. *pseudo-fluitans* (Journal of Botany, Tab. 342).

“Lower leaves membranous, elliptical-strapshaped, attenuated at each end; floating leaves subcoriaceous, gradually attenuated into the petiole.

“Var. *ericetorum* (our plate 11, lower figures).

“Leaves all similar, subcoriaceous, floating or rising out of the water, longly-stalked, with an oblong-oval or roundish lamina.”—English Botany, ed. 3, p. 28.

Taken in their extreme forms the above “varieties” seem so natural and so distinct that we might suppose they could be advantageously retained in the present work. On looking at a good series of specimens, however, it will at once be evident that the intermediate forms are so numerous and so gradually link “var. *ericetorum*” to “var. *pseudo-fluitans*,” that more than half our series will remain unnamed.

Under cultivation, by gradually increasing the depth of the water, “var. *ericetorum*” speedily becomes “var. *genuinus*,” and the same change has been noticed in plants growing naturally. Mr. Druce, in his “Flora of Oxfordshire,” writes (p. 283):—“In 1885 I noticed the floating form in a small pond . . . and a month later, this being dried up, the plant assumed the heath form”; the reverse result here being obtained.

The plant from the River Leven, Loch Lomond, selected by Dr. Boswell Syme as the type of “var. *pseudo-fluitans*,” is probably the ordinary deep-water “*genuinus*,” with leaves elongated and membranous (even in one upper leaf) by being kept almost continuously submerged by running water. I have not been able to obtain specimens to cultivate from this locality, but from observations made by Mr. Alexander Somerville, of Glasgow, on a very similar form found by him in the Isle of Arran, a form with “membranous lower leaves,” and otherwise exactly agreeing with Dr. Boswell Syme’s description, has been traced through “var. *genuinus*” almost or completely to “var. *ericetorum*.”

One more so-called “variety” has found a place in our British Lists, and so it must be briefly noticed here. It was named “var. *angustifolius*” by Fries (Novit. Flor. Suec. ed. 2, p. 30). I do not understand his description, and therefore do not copy it. The specimens figured in our plate 11 were collected by Mr. Charles Bailey, and named by Mr. Arthur Bennett from type specimens of Fries’ plant. This form also passes insensibly into the other above-named “varieties.” If we accept such transient states as “varieties,” we might as well name a dozen others equally indefinite.

The form figured on plate 12 is not only distinct in appearance from the usual

states of *polygonifolius*, but it remains unaltered when cultivated in stagnant water, under conditions wholly differing from those of its natural station. It was found by Mr. W. H. Beeby in 1890; the following account of it is extracted from his paper on *The Flora of Shetland* (Scottish Naturalist, January, 1891).

“Potamogeton (*polygonifolius* × *heterophyllus*?) Walls (district of). A remarkable plant, occurred plentifully in the Burn from Brousta Loch. It bears considerable resemblance to deep-water states of *P. polygonifolius*, but differs from that species in the shape, and texture when fresh, of the submerged leaves; and also in producing hardly any floating leaves when growing in still, shallow water. The few floating leaves produced are much thinner and less coriaceous. It has been referred with considerable confidence to *P. plantagineus*; while *P. natans* × *heterophyllus* has also been suggested. My own opinion is expressed above as to the origin of the plant, but what it really is must at present be held to be uncertain.”

In the summer of 1888 Mr. R. W. Sculley found a very similar plant in the Long Range, Killarney. On this he writes (Journal of Botany, April, 1890, “Plants found in Kerry”):—“Among these was a large gathering of the *Potamogeton* I had sent him (Mr. Arthur Bennett) in 1888, labelled ‘*P. polygonifolius*, submerged form,’ and about which there was then some doubt; however, after examining last year’s series, Mr. Bennett seems satisfied the plant is rightly named, . . . it is a common plant in most of the Kerry mountain streams.”

Both the Irish and Scottish plants have a remarkable likeness to some forms of *P. coloratus* (*plantagineus*), and at first I strongly inclined to regard them as specifically identical. After further examination, and considering the fact that *P. coloratus* has not been found in the Shetland Isles, I now incline to agree with Mr. Bennett in ranking this form under *P. polygonifolius* for the present. In deference to Mr. Beeby’s opinion as to its being possibly a hybrid, I do not use the term *var.*, but in conformity with my usage in this work prefer to employ the term *forma*, as indicating a possible hybrid origin.

POTAMOGETON POLYGONIFOLIUS, FORMA CANCELLATA mihi.

Rootstock resembling that of *P. polygonifolius*. Stem erect, simple, stipules acute, brownish, scarious, lower soon decaying, exceeding the rather short internodes; leaves all stalked thin, membranous, translucent, lower attenuated gradually into the petiole, lanceolate-strapshaped, or lanceolate, apex blunt, or acute, midrib with several rows of elongate chain-like areolations on each side, with 4–8 lateral ribs connected by distinct coarse ascending or transverse veins, giving a cancellated appearance to the whole leaf. Upper leaves similar in texture and venation to the lower, petiole often exceeding the lamina, which is contracted and auricled a little above the base, alternate like the lower leaves, or opposite below the peduncles and then sometimes subcoriaceous and indistinctly auricled at the base; peduncles

equal, slender, exceeding the subtending foliage, spike short, apparently barren. Colour bright green, drying olive green, tinged with brown or reddish crimson.

The above description is made solely from the Shetland plant; the similar-looking Irish one yet requiring to be tested by cultivation.

DISTRIBUTION in the British Islands:—*P. polygonifolius* has been recorded from all Watson's vice-counties except North Wilts, Bucks, West Suffolk, East Gloucester and Nottingham. In Ireland it is common. The record for Huntingdonshire requires confirmation; possibly it may occur near the Cambridgeshire locality at Gamlingay.

DISTRIBUTION in other countries:—Throughout Europe generally, but more abundant in the west. Mr. Bennett records Morocco, Madeira, Canaries, Angola, Madagascar, Siberia to China, Tibet, India, and Japan. The Australian records are somewhat doubtful. It seems to be a Palearctic species with some few southern outliers.

Explanation of the Plates.

PLATE 10. *P. polygonifolius*, Killearn, Stirlingshire, collected August 4th, 1897, by Mr. R. Kidston and Colonel Stirling; fig. 1, fruiting branch; fig. 2, flower-spike; fig. 3, part of lower leaf, mag.; fig. 4, section of stem; fig. 5, transverse section of fruit; fig. 6, vertical section, mag. All from fresh specimens.

PLATE 11. *P. polygonifolius*, "var. *ericetorum*," lower figures, collected by Mr. R. Kidston, August 26th, 1897, Stirlingshire, from fresh specimens.

P. polygonifolius, "var. *angustifolius* Fries," three upper figures from specimens collected by Mr. Charles Bailey, July 22nd, 1885, in a "peaty pool, Glen Etive, North Argyll.

PLATE 12. *P. polygonifolius*, *forma cancellata*, *mihi*; fig. 1, from specimens collected by Mr. W. H. Beeby, Wales, Shetland; fig. 2, cultivated specimen, grown at Chatteris.

6. **POTAMOGETON COLORATUS**, *Hornemann*, *Fl. Dan.*, Fasc. 25, t. 1449 (1813).

P. coloratus, Cham. et Schlecht, *Linnaea*, vol. ii. p. 194, Tab. 5, fig. 15 (1827). Fieber, *Pot. Böhm.*, p. 18, Tab. 1, fig. 5 (1838), Ascherson et Graeb. *Synop. Mitteleurop.* Fl. p. 309 (1897).

P. plantagineus, Du Croz, ap. Roem. et Schult. *Syst.* iii. p. 504 (1818), Hooker and Arnott, *Brit. Fl.* ed. 8, p. 484 (1860), Reichenbach, *Ich. Fl. Germ.*, vol. vii. p. 25, Tab. 45, et var. b? *pachystachyus*, Tab. 46 (1845). Hooker, *Stud. Flor.* ed. 3, p. 431. *Bab. Man.* ed. 7, p. 371 (1874). Syme, *Eng. Bot.* ed. 3, p. 29, pl. 1401.

P. Hornemannii, Meyer, *Chlor. Hanov.* p. 521 (1836), Koch *Syn. Fl. Germ.* ed. 2, p. 777 (1844). *P. plantagineus*, Tiselius, *Pot. Suec. Exsic.* fol. 7, *forma typica*, fol. 8, *forma terrestris*.

SPECIFIC CHARACTER.—Leaves all stalked, thin, membranous, pellucid, with opaque ribs and numerous transverse veins forming a delicate but conspicuous netting over the whole leaf; upper floating, opposite or nearly so, petiole flat above, concave and widened towards the lamina; lower submerged, persistent, broadly lanceolate, distinctly stalked or with the lamina decurrent to the base of the petiole, or often exactly like the floating leaves. Fruit minute, greenish, blunt when fresh but faintly keeled when dry. Plant growing throughout the winter.

DESCRIPTION: ROOTSTOCK with many irregularly springing slender white or reddish stolons, rooting freely at each node, fibrous, confused, with distinct terminal winter or resting buds, but the growth and extension continues throughout the year.

STEM terete, slender, with short internodes, simple or slightly branched in deep-water forms.

LEAVES all stalked, upper somewhat coriaceous when fresh, but thin, membranous, pellucid with the numerous ribs and transverse veins distinctly showing; very variable in shape and size with no clear distinction between the submerged and floating leaves; upper floating, short or long stalked, elliptical, oval, or lanceolate, rarely suborbicular, sometimes cordate and auricled at the base, many pairs opposite or subopposite, in hot summers sometimes subcoriaceous but then always translucent, with minute areolation over the whole leaf. Lower leaves like the upper, or lanceolate, or oblanceolate, or falcate, stalked, but often with the lamina decurrent at the base of the petiole and so appearing sessile; never reduced to phyllodes, persistent.

PETIOLE flat above, concave and expanding towards the lamina, somewhat herbaceous, always shorter than the lamina, especially in the upper leaves.

STIPULES blunt with a slight mucro, not channelled on the back, herbaceous, pellucid, subpersistent and becoming scarious when old, shorter than or exceeding the internodes, the lowest usually with a short spatulate or lanceolate adnate leaf.

PEDUNCLES slender, gracile, a little thicker than the stem, equal, exceeding the subtending foliage, 2–8 in. long.

SPIKE thin, cylindrical, densely flowered and fruited, 1–1½ in. long, slender in fruit.

FRUIT minute, greenish, dorsal margin blunt when fresh, obscurely keeled when dry, with faint lateral ridges (“dry drupelets very small rounded not keeled green, beak obsolete, $\frac{1}{20}$ in. long,” Students’ Flora, ed. 3), beak very short, subcentral.

HABIT very variable according to soil and depth of water, generally resembling that of *P. polygonifolius* in similar stations, or towards *P. alpinus*, or even *P. lucens*. Growing freely throughout the winter, the “resting buds” seem rather to enable the plant to contend against long continued drought than to serve as true winter buds

like those of *P. alpinus*. The *Land-form* grows freely on dried mud, producing a rosette of leaves at the end of each internode and forming long continuous runners. These tufts of leaves form the bases of new plants on the return of the water, producing growths like fig. 3 on plate 15.

Although an inhabitant of moorish fen districts, it also grows freely on Oxford and Boulder Clays; plants on the latter being extremely vigorous with stems 1-3 feet, and leaves .5 in. long.

Bentham in his *Handbook of the British Flora* ranks this as a variety of *P. natans*, and all the earlier authors confounded it with that species, or later on with *P. polygonifolius*. With the former it has no close relationship, and is always clearly separable from the latter by its persistent rootstock and its abundant lower leaves, and by its minute greenish fruit. If we look only at the general habit it is very like *P. alpinus*, and must come next to that species in any natural arrangement, dissimilar as the fruits of the two species are. When gathered it is impossible to mistake it for any other British Potamogeton, but in the water it simulates *P. lucens* and *P. polygonifolius*, and probably has been recorded, in the absence of fruit, for *P. alpinus*; the reddish colour of the leaves in some localities resembling that of some states of the latter species.

The distribution is remarkable, and probably the stations in which *coloratus* is now found are survivals of a once widely spread bog or fen land which diminished as our climate grew dryer. It is not a little remarkable that where *coloratus* abounds *polygonifolius* is seldom or never present. In the great *Fenland* in the counties of Huntingdon, Cambridge, Norfolk and Lincoln, *polygonifolius* is entirely absent, as far as I have been able to ascertain, and in a large part of this extensive tract, *coloratus* is the most abundant species. Notwithstanding the plentifully produced land-form, this species evidently requires a moist atmosphere, having greatly diminished in numbers around Chatteris during the past succession of dry seasons; it will bear exposure to the air, and to hot sunshine if these conditions are accompanied by occasional rains. This being the case, we can understand its being at present chiefly confined to low-lying lands, and its interrupted distribution from "North Somerset" to "Mid-Ebudes."

In Watson's "Topographical Botany," ed. 2, *P. coloratus* is recorded from twenty-seven vice-counties, from these we must deduct two:—Cardigan, the specimen on which the record rests being, according to Mr. Bennett, *P. polygonifolius*; and Leicester, from which county Mr. E. F. Cooper says it is absent. In addition to the twenty-five remaining vice-counties we may add Hereford! North-East York! Mid-West York. Westmoreland! Dumfries! Selkirk! Roxburgh, and Mid-Ebudes. I have received it from many localities in Ireland, and would be thankful for any additional records of this species, which is more abundant in the British Isles than it

is in the rest of Europe; in the *Cybele Hibernica*, ed. 2 (1898), it is said to be "rather rare," but is recorded from all the districts except X. and XI.

Mr. Arthur Bennett in his "Notes on Pondweeds" in the *Journal of Botany* for March, 1891, gives "Arabia, Socotra, and the West Indies" as extra-European localities. To these Dr. Ascherson now adds "Algeria." The only specimens I have been able to examine from any of these localities are from Socotra, collected by Professor I. Bayley Balfour; they are without flower-spikes, and in foliage and habit resemble *P. polygonifolius*. On the imperfect evidence of these specimens I think they cannot be safely ranged under any European species.

Explanation of Plates 13—15. POTAMOGETON COLORATUS.

PLATE 13. *P. coloratus*, upper branches and fruit-spikes of the typical form, from Parsonware Drove, Benwick, Cambridgeshire, on peat, August, 1897. Fig. 1, section of stem, mag.; fig. 2, section of fruit, mag.

PLATE 14, fig. 1. Flowering branch of form with cordate leaves; fig. 2, submerged leaves of the same, from Pidley Fen, Huntingdonshire, on clay, September, 1888; fig. 3, transverse section of fruit.

PLATE 15, fig. 1. Barren autumnal stem, from Grunty Fen, Cambridgeshire, on peat, October, 1892; fig. 2, flowering branch of a deep-water form, from Ramsey Hollow, Huntingdonshire, on peat, July, 1884; fig. 3, early state, from Welches Dam, Cambridgeshire, on peat, January, 1888; fig. 4, 5, land-forms from Welches Dam, Cambridgeshire, on peat-mud, September, 1890.

7. × **POTAMOGETON BILLUPSII**, *Fryer*, in *Journal of Botany*, vol. xxxi. p. 353 (1893).

× *P. BILLUPSII*, l.c. tt. 337 (excl. fig. 2), 338.

P. COLORATUS × *ZIZII*, Ascherson u. Graebner, *Synop. Mitteleurop. Flora*, vol. i. p. 335 (1897).

SPECIFIC CHARACTER.—Rootstock fibrous, but thickening in autumn into sub-tuberous stolons; floating leaves with long petioles greatly exceeding the somewhat spathulate subcoriaceous or membranous lamina; peduncles slender, equal, longer than the petioles; flower-spike $\frac{1}{2}$ in. long, barren, with the sepals closed but the stigmas exerted, soon decaying; lower submerged leaves on short branches at the base of the stem, mostly stalked, rarely subsessile, translucent, membranous, persistent. Plant growing throughout the year like *P. coloratus*, which it resembles in peduncle and spike. Differing from that species in the subcoriaceous floating leaves with very long petioles, and from *P. Zizii* in the latter character and in growing throughout the winter; it has also much the facies of some of the forms of *P. varians*. A barren hybrid = *P. coloratus* ♀ × *P. "Zizii"* ♂.

DESCRIPTION: ROOTSTOCK irregular, confused, fibrous, reddish, or white, with

numerous slender stolons, thickening into moniliform resting-buds, and frequently with long stolons terminated by reddish winter-buds like those of *P. coloratus*.

STEM slender, terete, ascending, simple above, with a few short submerged branches below, internodes long below, but greatly shortened or obsolete between the floating leaves.

LEAVES: lowest reduced to a linear phyllode shorter than the internode; lower stalked, or subsessile from the lamina being decurrent to the base, submerged, lanceolate, or oblanceolate, or falcate, pellucid, passing gradually by intermediates into the floating leaves; upper with the lamina oblong, elliptical, or lanceolate, very long stalked.

PETIOLE slender, herbaceous, flat, greatly exceeding the lamina, sometimes expanded above.

STIPULES herbaceous, persistent, shorter than the internodes, except at the upper part of the stem, blunt, or truncate, closely clasping when young, afterwards patent and involute; lowest rarely with a narrow strap-shaped adnate leaf shorter than the stipule.

PEDUNCLES slender, equal, greatly exceeding the spike; not axillary, but opposite a single floating leaf, 2-2½ in. long.

SPIKE barren, sepals closed, but stigmas exserted, soon decaying, a little thicker than the peduncle, ½ in. long.

FRUIT none; though flower-spikes are produced freely they are absolutely barren.

COLOUR varying from bright green in the land-form, to yellowish-green, tinged with brown in the young shoots. The floating coriaceous leaves are tinted with reddish-brown in their early stage. The colour of the whole plant varying according to the season and conditions of growth; but usually approaching that of *P. coloratus*.

HABIT: growing throughout the year like *P. coloratus*; the young shoots resemble those of *P. varians*, but these are succeeded by a dense growth of flowering shoots. An inhabitant of shallow, stagnant water, subject to be dried up in summer. Under cultivation this species thrives best when the above conditions are imitated; it then produces its flower-spikes abundantly, but if the water is kept at a uniform depth of 18 inches the flower-spikes are rarely formed.

In the *Journal of Botany* for December, 1893, a full account of the discovery of this hybrid is given, illustrated by two excellent plates by Mr. Morgan. It is necessary to refer the student to that description because the specific character is now limited to the form with coriaceous floating leaves, excluding the supposed state there figured on tab. 337, fig. 2. Notwithstanding continued cultivation for six years, and almost daily watching the plant under different conditions of growth,

I am still unable to determine the range of variation, or to clear up the relationship *P. Billupsii* bears to certain of the Fenland *Zizii* forms. Therefore it seems better to restrict the name of *P. Billupsii* to the form now figured in our plates 16 and 17, which has remained constant in its character, and which is the only form I can with certainty regard as "*Zizii*" $\times$ *coloratus*. By "*Zizii*" I here mean a plant of very doubtful rank, which can only be so named in an aggregate sense; this plant is much nearer the form named *P. lucens* var. *coriaceus*, by Nolte, figured later on in this work as *P. coriaceus*, than it is to typical *P. Zizii*.

The present hybrid-species was found by Parsonware Drove near Benwick, in Cambridgeshire, in August, 1892, growing amongst a mass of "*Zizii*" and *coloratus*. The rootstock was probably two or three years old, and the shoots first observed sprang from a land-form which had been produced when the ditch was dry in the earlier part of the summer. I have succeeded in reproducing the land-form under cultivation in the present summer of 1898, but owing to the rapid drying up of the pond in which it grew I did not succeed in further producing the remarkable states noticed in the original locality, where the plant has now been destroyed by the deepening of the ditch in which it grew.

A plant gathered in September, 1892, near Mepal, probably is a hybrid of similar origin, and no doubt it occurs in other localities in the Fens, but in the absence of flower-spikes it may readily be passed over as small *P. coloratus*, or *P. varians*. Our plates give a good representation of the typical state of the plant when living, but in this case the student will do well to rely more on the descriptions than on the figures. Even dried specimens fail to give the peculiar look of the living plant, which when in flower, or with flower-spikes, cannot be confounded with any other British *Potamogeton*.

Mr. C. R. Billups, my nephew, gave me much valuable assistance in the field-work which resulted in the discovery of this species, which bears his name.

Explanation of Plates 16, 17. $\times$ *P. BILLUPSII*, Fryer.

PLATE 16, fig. 1. Flowering branch, July, 1897, cultivated; fig. 2, part of rootstock with subtuberous stolons, and young stems, June 1898, cultivated.

PLATE 17, fig. 1. Flowerless branches, July 23rd, 1897, grown in water 12-18 inches deep; fig. 2, section of the stem, mag.; fig. 3, submerged leaf to show petiole and venation, nat. size. All from fresh cultivated specimens.

8. *POTAMOGETON ALPINUS*, Balbis, Miscell. Bot. p. 13 (1804).

P. ALPINUS, Ar. Bennett, in Journal of Botany (Aug. 1889). Reichenbach Icon. Flor. Germ. vol. vii. Tab. 32, fig. 57 (1845). Morong Naiad. N. America, p. 19, pl. 30 (1893). Ascherson u. Graeb. Synop. Mitteleurop. Flora, p. 311 (1897). Tiselius,

Pot. Succ. Exsic. fol. 12, "Forma *palustris*," fol. 13, "Forma *palustris*, Fries," fol. 14, "Forma *viridis*, Tis.," fol. 15, "Forma *longifolia*, Tis.," fol. 16, "Forma *borealis*, Tis.," fol. 17, "Forma *gracilior*, Tis.," fol. 18, "Forma *iterata*, Tis." (1894).

P. rufescens, Schrad. in Cham. Adnot. ad Kunth. Fl. Berol. p. 5 (1815). Cham. et Schlecht. in Linnæa, vol. ii. p. 210, Tab. 5, fig. 18 (1827). Fries Novit. Fl. Succ. p. 30 (1828). Reichenbach, Ic. Fl. Germ. vol. vii. p. 20, Tab. 32 (1845). Koch, Synop. Fl. Germ. et Helvet. vol. ii. p. 777 (1844). Bab. Man. ed. 7, p. 372 (1874). Hooker, Stud. Flor. ed. 3, p. 431 (1884).

P. fluitans, Smith, Eng. Flor. ed. 2, p. 231 (1828).

"*P. purpurascens*, Seidl!" Fieber, Pot. Böhmens, p. 16, Tab. 1, fig. 4 (1838).

SPECIFIC CHARACTER.—Rootstock annual, springing from a resting bud, which is deciduous, or attached to a short persistent stolon; leaves strap-shaped, lanceolate, or elliptical, subsessile, or very shortly stalked, all similar, thin, membranous, pellucid, or with the upper or some of them coriaceous and floating, obovate, blunt, short or long-stalked; these are produced alike on barren and flowering shoots; and, unlike those of the preceding species, belong to the vegetative rather than to the reproductive system of the plant, the flower-spikes ripening fruit equally well when they are absent. Fruit reddish-fawn colour, oblique, obovate, lenticular, acutely keeled, with obscure lateral ridges on the semicircular dorsal margin, with a less prominent ridge or keel on the convex ventral margin, beak subcentral, prominent, acute, reflexed. Colour of the whole plant pale, or dark olive-green, tinged with red or purple, especially on the stems and young leaves, and on the richly-coloured resting-buds. The stipules are large, long, patent, or clasping, hyaline, blunt, the lower usually bearing hooded adnate leaves, and the ordinary lower leaves are also frequently hooded, as in *P. praelongus*.

DESCRIPTION: Rootstock annual, growing in late autumn and early in the ensuing year from resting-buds, usually deciduous, which are produced on the ends of summer stolons, or on short stolons springing from the axils of the leaves along the stem, or sessile on the nodes of the roots and stem. In summer the rootstock resembles that of *P. coloratus*, with many irregular stolons, fibrous at the nodes, perishing with the stems in autumn.

STEM terete, simple, slender, ascending, internodes shorter than the leaves, nodes conspicuous, annular in the dried plant.

LEAVES either all similar, strap-shaped-lanceolate, thin, membranous, pellucid, attenuated towards each end, subsessile, or shortly stalked, flat and undulate, or rarely channelled and recurved, alternate, opposite beneath the peduncles, or the upper floating and coriaceous, long or short-stalked, blunt, lanceolate, or obovate, or elliptical, alternate, or opposite below the peduncles. The lower submerged leaves

have a row of elongate cancellate areolations on each side of the midrib, and two principal lateral ribs with many intermediate finer ones connected by ascending transverse veins, forming a somewhat regular quadrangular reticulation over the whole leaf; although narrowed towards the end, the apex is always blunt, often hooded, though rarely so conspicuously as in *P. praelongus*. The lateral ribs of the upper leaves often spring from the midrib for one-third of its length, in other respects the venation is much the same as in the lower leaves.

PETIOLE herbaceous, always shorter than the lamina.

STIPULES hyaline, very large, blunt, herbaceous, usually exceeding the internodes except in the middle of the stem; the lower are often patent and cymbiform, bearing adnate hooded leaves; slightly veined, with two more conspicuous ribs on the back.

PEDUNCLE equal, slender, erect, slightly curved when the fruit is mature.

SPIKE cylindrical, abrupt, densely flowered and fruited, about one-third the length of the peduncle.

FRUIT obliquely obovate, lenticular, dorsal margin semicircular, acutely keeled to the base, with faint lateral ridges, ventral margin convex, with an acute ridge, beak acute, prominent, recurved, colour reddish-fawn, sometimes reddish-chestnut.

COLOUR light yellowish-green, or dark olive-green, tinged with red or purple, especially on the stolons, stems, and young leaves, rarely green untinged with red even when dried. The conspicuous resting-buds are rich deep red or purple.

HABIT: varying in length according to depth of water from two inches to two feet, or more; somewhat like that of *P. coloratus*, and so resembling *P. lucens* in certain stages of growth; the fruit is produced and ripens equally well when only submerged leaves are present. In this respect the habit is much that of the forms of *P. coriaceus*, and differs from that of all the preceding forms, which only produce flower-spikes when coriaceous leaves are formed.

Although this species has been described under many names, and almost all writers have also named so-called varieties, it does not seem to possess any permanently distinct forms in this country. Dissimilar as many of the states are in appearance, all may be obtained from one rootstock in different seasons, and some of the forms which have been distinguished by varietal names, may often be found simultaneously on one rootstock, thus the "var. β , *homophyllus*," of Syme, Eng. Bot. vol. vii. p. 31, described as having "leaves all pellucid, none of them floating" (see our plate 18) is merely the submerged state of "var. α , *genuinus*, lower leaves pellucid; upper leaves subcoriaceous, floating."

The specimen selected for illustration is one of a very beautiful and instructive series collected from a small, rather shallow pit of stagnant water in Delamere Forest, Cheshire, by Mr. C. R. Billups, in 1898 and preceding years; no other *Potamogeton* grows with it, and flowers and fruit are produced alike on shoots with

entirely submerged or with coriaceous floating leaves, the latter also occurring on barren stems. The elongation of the stems and leaves is evidently caused by the crowded vegetation which consists entirely of this one species. On the sides of the pit and in the shallower parts of it quite dwarf forms are found, indeed several states occur as distinct as Syme's two "varieties." Mr. Billups considers that all the plants in this station originally sprang from a single seedling, as no other stations for *P. alpinus* are to be found within several miles of this isolated marl-pit. He also states that he has never met with a permanent rootstock in this species, it apparently being renewed annually from the conspicuous resting-buds (plate 20, fig. 4). This confirms what I have repeatedly observed in plants under cultivation; roots of *alpinus* planted too late to form resting-buds invariably having died in the autumn.

Although this statement is the result of observations carried on for some years, I would caution students not to accept this, or any other statement in this work, without careful examination of the behaviour of living plants of this and other species they may have opportunities of watching throughout the year. The life-history of the best-known Potamogeton is by no means completely understood, and I can readily conceive that local conditions might in some instances produce a more or less perennial rootstock in *P. alpinus*.

This species is one of the most distinct British Potamogetons, and only bears a superficial resemblance when growing to *P. lucens* and *P. praelongus*. The plant it most nearly resembles when in the water is the rufous state of *P. coloratus*, but I have seen no examples which present any difficulty when gathered, although *alpinus* varies in stature from two inches to twice as many feet. I have received it mixed up with specimens of $\times$ *P. nitens* and with the subsequently-described $\times$ *P. Drucei*, which only goes to prove that Potamogetons cannot be named at sight by any but the most accustomed eyes. Even so great a botanist as Bentham placed *alpinus* as a form of *lucens*, an allocation that makes it difficult to comprehend his views of what constitutes a species of the genus Potamogeton, even if we regard his "species" as corresponding to the groups into which other botanists have more or less successfully attempted to arrange the species.

DISTRIBUTION in the British Islands.

P. alpinus has been recorded in seventy-eight of Mr. H. C. Watson's 112 vice-counties, extending from Cornwall to Caithness, and from Carnarvon to East Norfolk. Of these, Cambridgeshire needs confirmation. Professor Babington knew of no specimen, nor could he find one in any other older collections in the University Herbarium. It is not recorded from East Cornwall, North Devon, North Wilts, Isle of Wight, West Kent, South Essex, Hunts, West Gloucester, Glamorgan, Brecon, Radnor, Carmarthen, Cardigan, Montgomery, Merioneth, Flint, Nottingham, West-

moreland, Isle of Man, Ayr, Selkirk, Roxburgh, Linlithgow, Kincardine, Elgin, Easternness, Argyll, North Ebuies, West Ross, East Sutherland, West Sutherland, Outer Hebrides, Orkneys, Shetlands. The *Cybele Hibernica* records it as "rare and local" from all districts except V. and IX.

DISTRIBUTION in other countries.—Throughout Europe; Asia: East Siberia, Thibet, Afghanistan; Africa: Cape of Good Hope; North America: Greenland, Vancouvers Island to Nova Scotia, and widely distributed throughout the Northern United States, but "rare" (Morong).

Explanation of Plates 18—20. POTAMOGETON ALPINUS.

PLATE 18, fig. 1. Flowering stem, with leaves wholly submerged; fig. 2, fruiting spike from same plant, without floating coriaceous leaves; fig. 3, section of stem, mag. Marl-pit in Delamere Forest, Cheshire, gathered by Mr. C. R. Billups, September, 1897.

PLATE 19. Submerged stem, with folded recurved leaves; a state with the look, when growing, of *P. lucens*. Mugdock Reservoir, Strathblane, Stirlingshire, July 18th, 1893, gathered by Col. J. S. Stirling and Mr. R. Kidston.

PLATE 20, fig. 1, 2. "Cultivated plants from winter-buds collected February 29th, 1892. Dried March 31st, 1892," C. R. Billups. Showing early state with leaves all distinctly hooded (split in drying) as in *P. praelongus*, and leaf bearing stipules. The lower figure represents the resting or winter-buds. All from a pond near Knutsford, Cheshire; fig. 3, flowering shoot, with floating coriaceous leaves, Mugdock Reservoir, July 22nd, 1891. Gathered by Col. Stirling and Mr. R. Kidston. (The figures of fruit were accidentally omitted, see Plate 26, fig. 6, 7, 8.)

9. × POTAMOGETON DRUCEI, *mihi*.

P. FLUITANS, Fryer in litt. 1893; Druce, Flora of Berkshire, p. 516 (1897). *P. fluitans*? Druce l.c. "additions, etc." *P. polygonifolius*? Fryer in litt. 1897. *P. natans* × *alpinus*? Druce, Flora, p. 516. *P. fluitans*? *P. alpinus* × *natans* or *P. alpinus* × *polygonifolius*? Druce in Exchange Club Report for 1897, p. 568 (1898).

SPECIFIC CHARACTER.—Rootstock persistent, stem terete, short, stout, simple, internodes much shorter than the petioles. Lower leaves submerged, transparent when young, with conspicuous veins and cross reticulations, lanceolate-elliptical, narrow towards both ends, with a blunt tip. Petiole of all the leaves thick, fleshy, terete, very long, 4–7 inches, usually exceeding the lamina. Upper leaves at first submerged like the lower, but ultimately floating and wholly coriaceous or with a narrow band of beautifully translucent reticulations like the lower, obovate, oblong, oval, or elliptical, with a short blunt mucro, 19–21 ribbed, with three principal pairs of lateral ribs, the innermost springing from the midrib about one-third from its base. Petioles 4–7 in. long, lamina 4–7 in. long by 1½–2 in. broad, with minute

regular elongate areolations over the whole leaf. Stipules very long (3–4 in.), exceeding the internodes, acute, hyaline and subherbaceous, ultimately when decayed, with persistent fibres, not channelled nor ribbed on the back. Peduncles much thicker than the stem, swollen towards the middle, gradually narrowed at each end, 3–4 in. long, spike cylindrical, 2–2½ in. long, numerous but not densely flowered. Flowers expanding, but apparently abortive. Colour of the living plant bright green, the ribs streaked with ivory-white, drying reddish-brown and dark green. The habit of the plant is very peculiar, the leaves not reaching the surface by the elongation of the stems so much as by the remarkable lengthening of the petioles, and the short stems are more or less persistent, sinking down in the autumn and rooting at the joints, from each of which a new shoot is produced, which probably grows throughout the winter.

DESCRIPTION: ROOTSTOCK stout, persistent, rooting with numerous strong fibres at the nodes; growing throughout the winter like *P. natans* (?).

STEM simple, terete, with short internodes, 1–4 in., which are always shorter than and usually greatly exceeded by the petioles; the length of the whole stem 12–18 inches.

LEAVES: lower submerged in their early growth, but ultimately ascending to the surface and becoming more or less coriaceous like the upper, elliptical-lanceolate, narrowed towards both ends, midrib fleshy, thick, prominent on both surfaces with four pairs of conspicuous ribs springing from its base to half its length, the two lower pairs reaching the apex, the others converging into the midrib, a fainter pair sometimes showing near the margin; cross venation bold, conspicuous, with branched ascending anastomosing veins somewhat regularly distributed over the whole leaf; upper like the lower, but ultimately thickly coriaceous, opaque, or with a narrow border along each side beautifully translucent and veined like the submerged leaves; lateral ribs translucent, faintly showing on both surfaces, 19–21 ribbed, cross-venation and minute elongate areolation distinctly showing over the whole surface of the leaf when held up to the light; oblong-elliptical, or obovate, or oval, lamina 5–7 in. long by 1½–2 in. broad, terminating in an abrupt blunt mucro. All the leaves are usually alternate, rarely opposite even beneath the peduncles.

PETIOLE very long, terete, or slightly flattened above when immature, thick, fleshy, usually exceeding the lamina, 2½–8 in. long.

STIPULES 3–4 in. long, usually exceeding the internodes, acute, herbaceous with scarious edges, not channelled on the back, hyaline, green, or with whitish edges, lower usually decayed by the time of flowering, but with persistent fibres.

PEDUNCLE terete, very stout, thicker than the stem, swollen towards the middle, narrowing towards the spike, 4 in. long, usually opposite a single leaf, the petiole of which it about equals.

FLOWER-SPIKE cylindrical, 4 in. long, flowers numerous but not crowded, freely expanding but apparently abortive.

FRUIT not observed.

COLOUR bright green, drying reddish-green like *P. alpinus*; the young leaves often with the ribs and cross nerves ivory-white tinged with green.

HABIT like that of deep-water forms of *P. polygonifolius* or of $\times$ *P. fluitans*, but making a dense growth from the rootstock and branches, which root at each node, late in the autumn.

P. Drucei has the most beautiful foliage of any of our British forms; when the submerged leaves first reach the surface they have the appearance of delicate lace laid on almost invisible glass. The epidermis and parenchyma are perfectly transparent and can only be seen in certain lights, the ordinary appearance of the leaves resembling that of the leaves of *Ouvirandra fenestralis*. As the upper leaves grow older they become opaque, and then resemble those of *P. natans*; some, however, retaining a narrow strip of lacework along the outer margins until they decay.

The difficulty of correctly assigning this species its proper rank in the genus is best shown by the short synonymy given above. Probably Mr. Druce is correct in supposing it to be *alpinus* $\times$ *natans*. It does not agree with any figure or description of Schrader's *P. spathulatus* ("*alpino-natans*") known to me, nor with a specimen I possess so named by good botanists, therefore I have not ventured to place our present plant under Schrader's species, the more especially as it has not yet been satisfactorily decided whether *P. spathulatus* is a form of *alpinus*, or a hybrid between *alpinus* and *natans*, or *alpinus* $\times$ *polygonifolius*.

This plant was first found by Mr. G. C. Druce in the River Loddon, Berkshire, August, 1893, and I have great pleasure in dedicating it to the earnest worker who has done so much to advance British topographical Botany, and who at once recognized the distinctness of the plant from others of the genus.

Unfortunately the first specimens sent me were badly dried and wanted roots and lower leaves. These by themselves I should still name *P. fluitans*, but not typical, as they have a look of *P. polygonifolius* about them not often seen in Roth's plant.

On the evidence of better specimens I with some hesitation referred the plant to *P. polygonifolius*, but required living roots to grow. Even when by the kindness of my friend I succeeded in obtaining these, I strongly inclined to refer the plant to Syme's *P. polygonifolius* var. *pseudo-fluitans*, although unlike his type specimens in the Edinburgh University Herbarium, yet well agreeing with his short description. After growing the roots for some weeks, to my surprise the plant began to make autumnal growths which removed it from *polygonifolius* and to a considerable extent supported Mr. Druce's view of its hybrid origin—*alpinus* $\times$ *natans*.

Perhaps it would have been better to have delayed the publication of this species until its life-history had been more fully worked out; but both artist and author were unwilling to omit the insertion of so beautiful a plant in its proper place in this work.

Explanation of plate.

PLATE 21. $\times$ POTAMOGETON DRUCEI, fig. 1, a barren stem drawn from a fresh specimen gathered by Mr. Druce from the River Loddon, Berkshire, September, 1898, with submerged, and floating coriaceous leaves; the petioles are shortened as numbered in inches on the plate; fig. 2, a young leaf from a cultivated plant, end of September.

10. **POTAMOGETON GRIFFITHII**, *Ar. Bennett*, *Journal of Botany*, vol. xxi. p. 65, Tab. 235 (March, 1883).

P. GRIFFITHII, *Ar. Bennett*, *Report of Bot. Exchange Club*, 1884 (1885). *Fryer*, *Journal of Botany*, vol. xxvi. p. 58 (1888). *P. alpinus* $\times$ *praelongus*, *Ascherson u. Graebner Synop. Mitteleurop. Flora*, p. 317 (1897).

SPECIFIC CHARACTER.—Rootstock stout, deeply rooting, persistent; stem branched, not wholly decaying at the top in winter; lower leaves undulated, entire, strap-shaped, hooded, sessile and slightly amplexicaul, or narrowed gradually to the base and subpetiolate; upper leaves long-stalked, oblanceolate; stipules long, narrow, blunt, lower with an adnate leaf, often exceeding the internodes; peduncles slender, narrowed towards the base of the spike, swollen towards the centre; spike cylindrical, densely flowered, barren; colour green tinged with yellowish-brown, floating leaves green.

DESCRIPTION: Rootstock persistent with stout deeply-rooting stolons.

STEM 1–2 feet long, stout, flexuous, internodes short, nodes conspicuously annular when dry.

LEAVES lower long strap-shaped, undulated, blunt, suddenly narrowed to the somewhat hooded apex, semiamplexicaul, sessile, or narrowed at the base into a short petiole “7–12 in. long, $\frac{1}{2}$ – $\frac{3}{4}$ in. broad,” *Ar. B.*; “11-veined, with 4–5 fainter veins close to the midrib, semitranslucent,” *Ar. B.*; upper leaves “floating, opposite, long-stalked, oblanceolate, gradually attenuated into the long petiole, 13–17-veined, with distinct areolation over the whole surface, more conspicuous towards the midrib,” *Ar. B.*; more opaque than the lower, coarsely cancellate with ascending veins, apex sometimes slightly hooded.

PETIOLE 2–4 in. long, but shorter than the lamina, flat, herbaceous.

STIPULES scarious, translucent bright green, “2 in. long,” *Ar. B.*, nearly as long as, or often exceeding, the internodes, the lower always with a lanceolate, sometimes hooded, adnate leaf; rather persistent.

PEDUNCLE 2-5 in. long, slender, equal, or a little swollen in the middle and narrowed towards the spike.

SPIKE cylindrical, short, $\frac{1}{2}$ -1 in. long, barren.

FRUIT: "young fruit oval, with the beak forming the continuation of the inner edge," *Ar. B.*

COLOUR bright green to dark olive-green, tinged with yellowish-brown.

HABIT "of *P. praelongus*, Wulfen," *Ar. B.*, which it resembles in its winter growth, and in its persistent strongly-rooting rhizome.

It will be noticed that I have used Mr. Bennett's original description as much as possible, appending his initials where I have used his exact words. For years I have tried to procure living specimens to grow and describe, but without success. Mr. Bennett unfortunately lost his original stock, and the roots he sent me some years ago were killed by a severe frost which destroyed many roots of *Potamogetons* growing under artificial conditions. Hence we have been compelled to draw the plates and write the descriptions for this work entirely from dried specimens, and the life-history of this obscure form still remains imperfectly known.

Because the plant seems unable to produce ripe fruit it has been, somewhat hastily, assumed to be a hybrid. On such evidence as we have at present I do not subscribe to that opinion, but incline to think it may be, if not a distinct species, an extreme state of *P. alpinus*.

There are many points in the life-history of the plant, however, which seem to make my opinion almost untenable. The strong deeply-rooting rhizome, with its partial growth throughout winter, and the absence of the characteristic winter-buds, all point to specific difference. In habit or facies it seems much nearer *P. alpinus* than *P. praelongus*, as the "hooded" leaves which have been often insisted upon as showing a relationship to the latter species are found equally in *P. coloratus* and *P. alpinus* in their young, completely submerged state, and also occur to some extent in *P. perfoliatus* and other species.

Again if we take leaf-structure and venation, we get a very near approach to *alpinus*, but all these characters do not seem to me sufficient to out-weigh the difference in habit.

Except in its persistent rootstock, and its growth throughout the winter, I do not see any resemblance of a specific character to *P. praelongus*, while the leaves are far less like those of that species than those of *P. alpinus*.

Against the theory of hybridity we have the testimony of more than one observer that *P. Griffithii* is the only *Potamogeton* growing in Llyn-an-Afon, Carnarvonshire, which is its only known locality.

Although this species has been known for some years, we must leave the

determination of its true character to some enterprising botanist who may gather fresh roots to be tested by cultivation.

Explanation of Plates. POTAMOGETON GRIFFITHII. Plates 22, 23.

PLATE 22, fig. 1. Cultivated plant from Mr. A. Bennett's Herbarium, flowering stem; fig. 2, single lower leaf, with stipule.

PLATE 23, fig. 1. Submerged young branch of cultivated plant, with stipule bearing adnate leaf, Fryer Herb.; fig. 2, young submerged branch, Bennett Herb.

11. POTAMOGETON PRAELONGUS, Wulfen in Roemer Archiv., vol. iii. p. 331 (1805).

P. PRAELONGUS, Cham. in Linnæa, vol. ii. p. 191. Fieber, Pot. Böhmens, p. 15, Tab. 1, fig. 3 (1838). Koch, Synopsis, Fl. Germ. ed. 2, p. 779 (1844). Reich. Icon. Fl. Germ. vol. vii. p. 21, Tab. 33. Bab. Man. ed. vii. p. 374 (1874). Hooker, Stud. Flor. ed. iii. p. 433 (1884). Syme, Eng. Bot. ed. iii. vol. ix. p. 41, pl. 1411 (1869). Morong, Naiadaceæ N. America, p. 32, pl. 39 (1893). Ascherson et Graeb. Synop. Mitteleurop. Flor. p. 315 (1897). Raunkjær, Danske Blomsterpl. p. 78, fig. 41 (1896). Tiselius, Pot. Suecici Exsic. Nos. 85-88. Notulæ, 85, 86, forma *elegans*, Tis. (1895).

SPECIFIC CHARACTER.—Rootstock persistent, with stout white deep-rooting stolons; barren stems wholly submerged, the fertile ones ascending to the surface at the time of flowering, and afterwards sinking to the bottom to ripen the fruit. Leaves all similar, membranous, pellucid, sessile, rounded at the base, oblong or strap-shaped oblong, blunt and hooded at the apex. Fruit very large, dorsal margin semicircular, acutely keeled with a prominent wing often exceeding the beak; lateral ridges obscure, inner margin almost straight, terminating in the conspicuous beak. The peduncles are usually terminal and very long (6-15 in.), always greatly exceeding the densely-flowered and fruited spike ($1\frac{3}{4}$ - $2\frac{3}{4}$ in. long). Soon after flowering the fertile stems become completely submerged, and sinking to the bottom, ripen their fruit late in the autumn.

DESCRIPTION: ROOTSTOCK persistent, striking deep into the ground with stout white far-creeping stolons, rootlets at the internodes very numerous and long.

STEM ascending, flexuous, stout, simple below, branched above, or simple in the barren stems, with short internodes, rarely exceeding the leaves in the flowering shoots.

LEAVES all similar; semi-amplexicaul, oblong-strap-shaped or oblong, from a rounded base, sometimes gradually narrowed from base to apex; blunt, distinctly and constantly hooded at the tip. The midrib is broadly bordered by conspicuous elongate chain-like areolations, lateral ribs numerous, with two principal ones near

the margin arising from the base and terminating at the hooded apex, the intermediate ribs are more slender, and rise from the base of the central and two principal lateral ribs, which themselves are slightly bordered by areolations like the midrib. Cross-venation rectangular, forming a somewhat regular network over the whole leaf. The first-formed leaf at the base of the stem is oblanceolate and the lamina is so narrowed towards the sessile base as to make the midrib falsely resemble a true petiole. All leaves are slightly undulate.

PETIOLE none.

STIPULES hyaline, obtuse, often exceeding the internodes, subpersistent, brown or purple-green, or sometimes white, thin and scarious, then fibrous from decay of the tissue. The lower sometimes with adnate small ordinary leaves.

PEDUNCLE very long, 6-15 in., equal, a little thicker than the stem, greatly exceeding the subtending leaves.

FRUIT (dried) very large, acutely keeled, with the upper part of the keel often expanded into a wing exceeding the beak, ventral margin almost straight, and terminating in the prominent beak.

COLOUR bright green, with darker green stems, in autumn the young half-unfolded leaves are delicately tinted with pale green and violet.

HABIT: *P. praelongus* is very much like *P. lucens*, and is with difficulty distinguished from that species when in the water, except at the time of flowering. The young shoots are produced in autumn, and grow slowly through the winter, rising to the surface in the following spring and flowering in May and June. As soon as fertilization of the spike is complete it becomes submerged, and the whole flowering branch sinks gradually to the bottom of the water, where the fruit slowly matures. The remarkably long peduncles enable the plant to keep its flower-spikes emerged in the event of a sudden rise in level of the stream or river, a provision perhaps necessary from the early season of flowering and its being rather an inhabitant of tidal rivers, streams, and canals, than of stagnant waters.

P. praelongus is more nearly related to *P. perfoliatus* than to any other of our British species, but it is one of the most distinct Potamogetons known, and when gathered is not easily confused with any other plant. In the water, as already stated, the mature plant has so much the facies of *P. lucens* that even the most experienced collector may pass it by for that species. Again, in states with very short leaves, *praelongus* has much the look of *P. perfoliatus*.

Although differing greatly in the length and breadth of the leaves, *praelongus* does not afford any forms, in these islands at least, which can be distinguished as varieties, nor can I find any distinguishing characteristics between southern and northern specimens. Greater divergences from the ordinary form of the plant

occur in gatherings from plants growing side by side in Cambridgeshire, than between the usual type of these and specimens from Shetland.

DISTRIBUTION in the British Islands.

This species is chiefly found in the eastern side of England, and more generally towards the north, ranging from Surrey to Shetland. In the westward side it is found in only one county in Wales, Merioneth; and in Stafford, Salop, Mid-west York, Westmoreland, and Cumberland. In Scotland it ranges from Wigton to the Shetland Isles, and from Mid-Ebudes to South Aberdeen, and is recorded with more or less certainty from twenty-one Scottish vice-counties. In England it ranges from Surrey and Berks to Cheviot-land, being most abundant in Province IV., Ouse, in every county of which it occurs. In all forty-seven counties have been recorded for *P. praelongus*, and probably it will be found growing in many others. In Ireland it seems rare, the few specimens I have seen from there offer no peculiarities. The "Cybele Hibernica" records it as "very rare" from Districts I.—VI. and IX.—XI., XII.

DISTRIBUTION in other countries.—In North America *P. praelongus* ranges from Vancouver's Island to Nova Scotia, extending southwards to Sierra Co., Cal., in the west and New York in the east; in Europe from the Faroes to North Russia, extending southwards to France. It is recorded from West Siberia and Japan.

Explanation of the Plates. POTAMOGETON PRAELONGUS. Plates 23—25.

PLATE 24, fig. 1. Flowering plant from Mepal, Cambridgeshire, June, 1897; 2, a detached flower, mag.; 3, section of the stem, mag.

PLATE 25. Part of submerged barren stem to show autumnal growth, from Jackson's Fen, near Woodwalton, Huntingdonshire, August, 1897.

PLATE 26, fig. 1. Rootstock with submerged autumnal shoots, Woodwalton Fen, Huntingdonshire, September, 1898; 2, fruiting branch, Mepal, Cambridgeshire, June, 1885; 3, fruit, mag.; 4, vertical section, mag.; 5, horizontal section, mag., both these figures show the cavity in the "process," or projection from the interior, around which the embryo is coiled. (Figures 6, 7, 8, mag., are the fruit of *P. alpinus* accidentally omitted on Plate 20.)

12. **POTAMOGETON PERFOLIATUS**, Linn., Species Plantarum, vol. i. p. 126 (1753).

P. PERFOLIATUS, Ray, Synop. Stirp. Brit. ed. 3, p. 149 (1724). Smith, Eng. Flor. ed. 2, p. 233 (1828). Chamisso in Linnæa, vol. ii. p. 188, Tab. 5, fig. 13 (1827). Fries Novit. Flor. Suec. p. 42 (1828). Fieber, Pot. Böhm. p. 14, Tab. 1, fig. 2 (1838). Reichenbach, Icon. Flor. Germ. vol. vii. p. 19, Tab. 29, fig. 53, 54 (1845). Koch, Synop. Flor. Germ. ed. 2, p. 779 (1844). Syme, Eng. Bot. ed. 3, vol. ix. p. 374 (1874). Hooker, Stud. Flor. ed. 3, p. 434 (1884). Morong, Naiad. N. America,

p. 33, pl. 40 (1893). Ascherson u. Graebner, Synop. Flor. Mitteleurop. p. 313 (1897). Raunkiær, Dansk. Blomst. p. 82, fig. 45 a-c. Tiselius, Pot. Suec. Exsic. Fasc. 2, fol. 89, Forma *typica*, Tis., fol. 90, Forma *rotundifolia*, M. K. fol. 91, Forma *densifolia*, Tis., fol. 92, Forma *imbecilla*, Tis. (Notulam 92) (1895).

SPECIFIC CHARACTER.—Rootstock with stout white far-creeping stolons, decaying in winter, except the terminal shoot of two or three joints, which bear winter-buds somewhat resembling those of *P. alpinus*, usually colourless, but of a reddish colour like those of *P. coloratus*, when springing from the axils of the upper leaves, or sometimes becoming so when dried. Stems terete, slender or stout, much branched when in flower, with long or short internodes, but usually shorter than the leaves. Leaves all submersed, similar, membranous, slightly concave, edges with minute spines forming an obscure denticulation along the upper margins which are somewhat undulated and scabrous; all sessile, amplexicaul, oval, ovate, sub-orbicular, or oblong-lanceolate from a cordate base, usually blunt, or with the apex contracted into a blunt mucro; with 5-9 principal ribs often narrowly bordered by elongate chain-like areolations, with numerous fainter intermediate ribs; cross-venation distant, irregular and indistinct. Stipules short, blunt, scarious, soon decaying, the lowest with a small adnate leaf, commonly reduced to a herbaceous auriculate or orbicular expansion towards the apex. Peduncles equal, exceeding the subtending foliage. Fruit oblique-obovate, blunt on the semicircular outer margin, with the central keel acute in immature, but blunt and obscure in ripe drupelets; lateral ridges obscure; inner margin convex, terminating in the subcentral rather prominent slightly recurved beak. Colour light or dark olive-green, often tinged with yellow or purplish-brown; stem white or green.

DESCRIPTION: ROOTSTOCK with stout far-creeping white, or sometimes reddish stolons, these spring irregularly from the same node as the ascending stem, sometimes in pairs; all decaying in autumn except the two or three last-formed joints, the terminal one bearing a winter-bud, which, when growth commences somewhat late in spring, is like the growing bud of *P. alpinus*. Deciduous winter-buds, however, are not produced in the present species. The stolons are furnished at each node with numerous long deep-rooting fibres.

STEM terete, stout or rather slender, simple below, much and repeatedly branched above, with but few barren branches until late in the summer; in deep, running, or tidal waters often 8-10 feet long or more, and then with fewer flowering branches; these are usually produced in pairs springing from the axils of the leaves which subtend the first formed peduncles. The internodes in these deep-water states are greatly elongated, much exceeding the leaves, which on the contrary are shorter and stouter than on plants growing in more tranquil waters; in which the internodes are usually much shorter than the leaves.

LEAVES amplexicaul, sessile, or the lowest rarely petiolate, cordate at the base which is more or less expanded so as to wholly or partially surround the stem; ovate, orbicular, oblong, or lanceolate towards the apex; with 5-9 principal ribs rising from the base of the leaf, each often bordered by coarse elongate chain-like areolations, with many intermediate slender ribs, all connected by rather irregular transverse veins; apex blunt and entire, or narrowed into a short blunt mucro in the more lanceolate forms; rarely hooded, but then distinctly so; margins with minute spine-like denticulations towards the apex, which are more or less deciduous, scabrous, a little undulated; the whole leaf is slightly concave, shining, more or less membranous and translucent, a little stiff and horny in texture. The lowest leaf is sometimes subpetiolate, with an orbicular or lanceolate lamina which is so narrowly decurrent to the base of the midrib as to give the appearance of a distinct petiole.

PETIOLE usually none, or present on the lowest leaf only as above described, broad, herbaceous, being merely a part of the midrib.

STIPULES short, blunt, scarious, usually decaying before the full expansion of the leaf; the lowest often with an adnate leaf, which is usually reduced to a herbaceous auricle, or rarely resembling a small ordinary ovate leaf.

PEDUNCLES equal, about as thick as the stem, axillary, or terminal, exceeding the opposite subtending leaves, often a little curved; about twice the length of the spike, $2-2\frac{1}{2}$ in. long.

SPIKE thick, cylindrical, densely flowered, flowers pedicellate as in *P. praelongus*; $1-1\frac{3}{4}$ in. long.

FRUIT obliquely obovate, beak slightly recurved, subcentral, slightly winged towards the ventral margin, which is gibbous in the upper part and obliquely concave to the base; dorsal margin semicircular, obscurely keeled, with very faint lateral ridges.

HABIT resembling that of *P. praelongus* during the season of growth, but the whole of the plant dying early in autumn and renewed in the following spring from the terminal winter-buds above described, which are generally produced on short stolons which grow freely from barren stems springing up when the fertile branches are matured; these resting stolons of two or three joints with a terminal growing bud are usually subterranean, but are sometimes produced in the axils of the upper leaves.

COLOUR varying from white to reddish green, or to olive-green on the stems; and from light to dark green, or yellowish-green, clear or tinged with purple-brown, on the leaves. The plant sometimes varies in colour on successive stems from the same stolon.

Potamogeton perfoliatus has so considerable an amount of variation both in the

shape and colour of the leaves that at first sight it seems easily separable into distinct varieties, but the examination of a good series, even of dried specimens, shows that the most extreme forms pass into one another so gradually that they are probably nothing more than *states* produced by local and often temporary conditions. An isolated plant growing in a newly dug clay pit afforded unusually good opportunities for observation during several years, and this plant changed so much in the shape, colour, and even texture of the leaves, that I am further induced to think that all our British forms are mere states, not true varieties. When first noticed this single plant was very small, apparently a seedling of the year, and the leaves were elongated, thin, and diaphanous, bearing much resemblance to those drawn on our plate 28, fig. 2. After three or four years the whole plant grew coarser in texture and the leaves were more horny, of a less vivid green, and approximate in shape to those figured in plate 27, fig. 1. Ultimately the plant spread and occupied a considerable part of the little pit in which it grew, and became so like the ordinary state of the species that I ceased to observe it, thinking that it had afforded sufficient proof that the texture and shape of the leaves are too inconstant to afford satisfactory grounds for varietal distinction.

In the ninth edition of *The London Catalogue of British Plants* (1895) Mr. Arthur Bennett has admitted "var. *b. lanceolatus*, Blytt" (Plate 28, fig. 1), on the evidence of specimens collected by Mr. A. Brotherston in a "mill-dam, Selkirk, June, 1876." The greatly elongated leaves of this form seem to be due to artificial submergence of the whole plant, which has evidently been unable to reach the surface of the water. I have little doubt, from the examination of a series of specimens, that this is an extreme state of the plant represented in figure 2 on the same plate, a form not infrequent in cold and clear northern streams. In this form I at first thought that we had a true climatic race or variety, and with the object of testing this supposition I examined Mr. Bennett's series of *P. perfoliatus* from all parts of the world. The evidence thus obtained is far from conclusive, but seems to indicate that the so-called varieties of *P. perfoliatus* are merely states induced in individual plants by local or temporary conditions which are not sufficient to prevent reversion to the type, which itself always seems to accompany the variety.

It is important that we should arrive at just conclusions on the value of the variations presented by each species, and that we should not confound temporary states with permanent varieties which are, in fact, incipient species. In looking over an extensive series of dried specimens of any *Potamogeton*, it is easy to select very distinct-looking forms, and as easy to give varietal names to them; this is the method that has been followed by most of the writers on the genus, and I regret to say is still being followed, with the result of confusing rather than elucidating an already too difficult subject.

Our plates show a sufficiently wide range of leaf variation to enable the student to name any state of *P. perfoliatus* he is likely to meet with, therefore I think it is unnecessary to add a long series of descriptions which I have made of the various states of this species occurring in British waters, and merely to make one or two remarks on points of interest.

On plate 28, fig. 1, we have figured two of the most extreme British forms, which have been referred to the "var. *lanceolatus*" of Blytt. The leaves of fig. 1 are thin pellucid and greatly elongated, probably by long continued artificial submergence. I have gathered equally diaphanous leaves from a plant which afterwards produced those of the usual stiff horny texture. Fig 2 is a form more common in northern than in southern waters, but as the plant already referred to produced elongated leaves, similar to those figured, when it grew under altered conditions, and as these forms with lanceolate leaves are usually, or always, found growing with the ordinary states of the species, I am unwilling to accept *lanceolatus* as a true variety in this country. The fruit as seen in herbariums, and in a lesser degree when fresh, varies slightly in size, in being bluntly or acutely keeled, in the depth of the furrows between the keel and the lateral ridges, and in the beak being almost facial, or central. These characters probably depend upon the degree of maturity of each drupelet, as young fruits are usually sharply keeled, and the beak is usually subcentral rather than facial.

Although so dissimilar in appearance to the greater part of the broad-leaved Potamogetons, *P. perfoliatus* crosses readily with several of them, but I think is always the male parent; hence the *perfoliatus*-hybrids are ranged in this work, after the supposed female parents. As none of the hybrid forms have been produced artificially it is not possible to say with absolute certainty whether this classification is correct, but it has the advantage of ranging together those species and hybrid-species which most closely resemble each other in habit and general facies if observed in all stages of their growth; thus if we take the foliage of the mature plants alone, $\times$ *P. Cooperi* has the look of *P. perfoliatus* rather than of *P. crispus*, but the compressed stem and winter-growth induce me to range it under the latter species as its probable seed parent.

DISTRIBUTION in the British Islands.

P. perfoliatus has been recorded in ninety-seven vice-counties from Cornwall to Shetland; records are wanting for East Cornwall, South Somerset, Isle of Wight, Radnor, Pembroke, Cardigan, Montgomery, Merioneth, Denbigh, Flint, South-East York, Peebles, Haddington, Dunbarton, Cantire. Probably it will be found in most of them; except, perhaps, the Isle of Wight. The "*Cybele Hibernica*" gives it as frequent throughout Ireland.

DISTRIBUTION in other countries. Mr. Arthur Bennett has sent me the follow-

ing account:—"Central America: Guatemala! St. Lucia. Australia: Victoria! Queensland! N.S. Wales! Tasmania. Sandwich Isles. Iceland! Faroes. Scandinavia! to Italy! and Dalmatia! Holland! and Belgium! to Central Russia! and Caucasus! Asia: Siberia! Songoria! Tartary. Turkestan! Tibet to 14,000 feet, and E. Tibet to 15,000 feet. India: Ceylon to Kasmir! Afghanistan! Persia! China. Japan! Arabia! Africa: Algeria! S. Morocco! N. America: Canada to Vancouver's Island! Ontario! to Lewes River 62°! Massachusetts to California!"

Explanation of Plates 27, 28. POTAMOGETON PERFOLIATUS.

PLATE 27, fig. 1. Submerged stem; fig. 2, young terminal short with flower-spike and subtending fertile branchlets; fig. 3, flower, mag.; fig. 4, section of stem, mag.; fig. 5, section of fresh drupelet, mag.; fig. 6, section of dried drupelet, mag. From Mepal, Cambridgeshire.

PLATE 28, fig. 1. "Var. *lanceolatus*, Blytt," mill-dam, Selkirk, June, 1876. A. Brotherston, Herb. Charles Bailey; fig. 2, "var. *lanceolatus*, Blytt," West Forfar, 1892. John Knox, Herb. A. Fryer.

13. POTAMOGETON CRISPUS, Linn., Sp. Pl. ed. 1, p. 126 (1753).

P. CRISPUS, Chamisso et Schlecht. in Linnæa, vol. ii. p. 186, Tab. 5, fig. 12 (1827). Fries Novit. Flor. Suec. ed. 2, p. 43 (1828). Reichenbach, Icon. Flor. Germ. vol. vii. p. 18, Tab. 29, fig. 50 type, Tab. 30, fig. 51, "var. *gemmifer*," fig. 52, "var. *serrulatus*," Schrad. (1845). Koch, Synop. Flor. Germ. ed. 2, p. 779 (1844). Fieber, Pot. Böhmens, p. 31, Tab. 3, fig. 15 (1838). Bab. Man. ed. 7, p. 374 (1874). Hooker, Stud. Flor. ed. 3, p. 434 (1884). Syme, Eng. Bot. ed. 3, vol. ix. p. 43, pl. 1413 (1869). Morong, Naiad. N. America, p. 36, pl. 44 (1893). Ascherson und Graebner, Synop. Flor. Mitteleurop. p. 335 (1897). Raunkiaer, Danske Blomsterpl. p. 83, fig. 46 (1896). Irmisch Potameen, p. 17, Tab. 2, fig. 22-37 (1858). Tiselius, Pot. Suec. Exsiccati, fol. 93, 94, Forma *typica* (1895).

SPECIFIC CHARACTER.—Rootstock usually annual, stem springing laterally in autumn from a deciduous axillary or terminal winter-bud, or from a surviving portion of a stem of the preceding summer's growth, compressed, obscurely quadrangular when mature, channelled on the broad flattened sides. Leaves acutely serrulate, flattened when young, crisped or strongly undulate when mature, 3-5 ribbed, the midrib with many parallel fibres, two principal lateral ribs very near the margin, the outer rib just at the base of the serrulations; linear, strap-shaped, or oblong, blunt or rounded at the apex, or slightly acute, without a distinct mucro. Stipules short, scarious, blunt, or lacerated at the apex, fugitive, several of the lower on both stem and branches often with adnate ordinary leaves and then more

persistent. Peduncles equal, a little curved, exceeding the subtending leaves. Spike short, few-flowered. Fruit obliquely obovate, with a long acute beak equalling the nutlet, 3-keeled with the central keel slightly winged and crested, with a long fleshy tooth at the base, sometimes reduced to a tubercle, but always present in the fresh fruit, lateral ridges or keels obscure. The plant grows throughout the winter, and is usually or always only of annual duration. Colour brownish-green or bright green, sometimes tinged with red or lilac-purple on the young leaves.

DESCRIPTION: ROOTSTOCK probably of annual duration only, from seed, or more frequently springing from an axil towards the centre of a deciduous thyriform winter-bud, with slender terete stolons, somewhat contorted, with strong fibrous roots from the nodes.

STEM compressed, obscurely quadrangular, with the flattened broader sides, from whence the leaves spring, deeply guttered when mature, rounded on the narrow sides; simple below, much and repeatedly branched above in the mature plant, also with short branchlets springing from the axils of the leaves on the main stem; on which the internodes are at first shorter than, afterwards usually exceeding the leaves, very short on the side branches.

LEAVES all similar, sessile, but not amplexicaul, strap-shaped, margins serrate, in the mature leaves often strongly undulate and crisped at the margins, the submerged and young leaves flat and like those of *P. obtusifolius*, but with finely serrulate margins, rounded or acute, obscurely mucronate; 3-5 ribbed, the lateral ribs near the margin, and joining the midrib unequally a little below the apex; midrib bordered by several rows of elongate cancellate areolations. When full-grown a faint rib is perceptible close to the margin, joining the lateral rib some distance below the apex, cross-venation distant, oblique, alternate from the sides of the midrib, not continuous, but terminating at the principal lateral rib, from which a vein rises a little higher up and joins it to the faint marginal rib.

STIPULES soon decaying, but always present on young growths, and more persistent when bearing leaves, scarious, apex blunt at first, afterwards lacerated, short, seldom more than $\frac{3}{4}$ in. long, often many of the lower with adnate leaves.

PETIOLE none, or the lowest leaf with the lamina so narrowly contracted towards the base as to appear petiolate, but then it is merely a naked portion of the midrib.

PEDUNCLE a little compressed and slightly channelled towards the slender curved apex, equalling or exceeding the subtending leaves, $1\frac{1}{2}$ - $2\frac{1}{2}$ in. long, terminal, but ultimately placed in the forks of the branches by the continued growth of the plant, rarely lateral when on the last formed branchlets, and then only by abortion of one of the two branchlets which constantly spring on either side of its base.

SPIKE short, loose, few-flowered, 7-10, few, or densely fruited, and then subglobular as in the *Chloephylli*.

FRUIT oblong, oblique, compressed, with a pointed slightly recurved beak about equalling the nutlet in length, central keel slightly winged and prominent, sometimes a little crested towards the base, which is furnished with a long projecting fleshy tooth, rarely reduced to a tubercle, lateral ridges obscure, blunt; pericarp "thick and corky" (Morong).

COLOUR olive-green, or dark green, the young leaves sometimes clear bright green; often tinged with reddish-brown and with purple or lilac midribs and veins; sometimes the young leaves of the upper branches are tinged with crimson; when in fruit the whole plant often becomes brown or yellow green.

HABIT: growing from late autumn throughout the winter, and flowering from May onwards till September, the fruit-spikes, unlike the rest of the British species, are borne for a considerable time above the surface of the water:—"Probably the fruit of *P. crispus* is matured under water, but it seems to be less deeply submerged than that of most species, and liable to frequent exposure to the air, a condition which the thick corky drupelets can sustain without injury" (Journal of Botany, p. 226).

This species is so marked in character and so uniform in habit under varying conditions of growth that few attempts have been made to give specific or varietal names to any of its states. It will be necessary in this place to notice only two of them: "*P. serratus*," of Hudson, and "var. *cornutus*," of E. F. Linton.

The *P. serratus* (Hudson, Flora Anglica, p. 62 (1862) has been assumed by subsequent authors to be the young state of *crispus*, mistaken by Hudson for a distinct species. From Hudson's description and from his reference to Ray, Synopsis, ed. 3, p. 149, and to Gerard, ed. 2, p. 823, *P. densus* possibly might have been intended, especially as that common species is not otherwise included in his list of British species of Potamogeton. Smith, in his English Flora (ed. 2, p. 234 (1828), relying perhaps rather upon Gerard's figure than his description, and by the sweeping method of excluding all Hudson's synonymy, places *P. serratus* as var. β under *P. crispus*, and has been followed in this course by many subsequent botanists. In 1890, writing in the Journal of Botany, I was inclined to think that we had a flat-leaved form in the Fens which possibly might be a true variety as distinguished from a temporary state; I then wrote (l.c. p. 225), "Authors generally consider that Hudson's *P. serratus* (*P. crispus* var. *serratus*, Lond. Cat. ed. 8) was founded on the young state of *P. crispus*; perhaps, however, it would be better to restrict the name of *P. serratus* to a small flat-leaved form of the species which is distinguished by its finely serrulate flat, narrow, strap-shaped leaves even when the plant is in fruit. Possibly this form may be a true variety of *crispus*, but, as far as I have been able to observe in the Fens, where in some districts it is common, it is more probably a *state* induced by local

conditions. It is usually found in ditches crowded with vegetation, and when shoots from the same rootstock are able to struggle into an unoccupied space on the surface of the water the leaves become crisped and undulated to some extent."

After nearly ten years, during which *P. crispus* has been frequently observed and its changes watched, I am now convinced that the form to which I proposed to restrict the name of *serratus* is merely a *state* which reverts on the same individual to the typical form of the species. Hence, although the supposed variety is figured on our plates, I think it unnecessary to retain the varietal name of "*serratus*."

The second proposed variety, "*cornutus*," was founded on a very imperfect knowledge of the species. The fruit is *always* more or less furnished with a tooth at the base of the keel. This is a character I have never found wanting in any specimen I have examined, either British or foreign! It is true that in individual drupelets the tooth is sometimes reduced to a mere tubercle, but in every spike I have examined some of the fruits are long-horned at the base. Probably these are the more mature fruits. Anyone who has watched the growth and development of these spines or tubercles on the fruit of *Ceratophyllum*, or on the fruits of the grass-leaved section of *Potamogeton*, will not hastily attempt to found varieties on the greater or less development of these variable organs.

Perhaps it will be well, as my description of *P. crispus* in the Journal of Botany (l.c. pp. 225-27) seems to have been overlooked by the writer in the same Journal for 1894 (p. 187), to quote some substantial authority for the existence of the tooth on which Mr. Linton has founded his proposed variety. In 1838, Fieber, in his valuable *Potamogeta Böhmens*, p. 33, in describing the *type* (his *varieties* are founded entirely on the leaves), writes: "Früchte zusammengedrückt, langgehörnt, rundlich-viereckig. Rückenkiel geflügelt, am Grunde höckerig, vorn am Grunde ein spitziger Höcker." Morong, in his *Naiadaceæ of North America* (March, 1893), wrote, p. 36, "Fruit ovate . . . having three rounded keels on the back, the *middle one with a small projecting tooth near the base*."

Here we have the horn, tooth, or tubercle, clearly defined by two of the most eminent writers on the genus, and so we must presume that as no reference is made in Mr. Linton's description of his variety to either of these writers, or to the horn or tubercle which they have described, he was altogether unaware of the existence of any such feature in the *type*; and that his variety is named from the actual presence of a horn rather than from its relative length.

As far as the length goes, from a specimen supplied by Mr. Linton himself, I can safely aver that the horn of his "*cornutus*" is about the average length of the same feature in the majority of specimens; I have seen it much more strikingly developed.

Enough has been said to show why I reject both these "*varieties*," "*serratus*" and "*cornutus*"; but it cannot be repeated nor insisted upon too often, that students

of the genus should examine for themselves and form their own conclusions from what they see in the living plants; no writer on this difficult genus has escaped errors from insufficient means of observation and comparison, and everyone who goes to Nature for his guide will find much to supplant and correct in the observations already made even by writers of the highest repute.

P. crispus, like *P. perfoliatus*, seems to cross readily with very diverse species, but unlike the latter plant it probably is usually the seed-bearing parent. Perhaps in this respect the first hybrid placed under it, $\times P. Cooperi$, = *crispus* $\times$ *perfoliatus* is most open to doubt; but the closely allied $\times P. undulatus$, = *crispus* $\times$ *praelongus*, and $\times P. Bennettii$, = *crispus* $\times$ *obtusifolius* (or *Friesii*?) are so like *crispus* in habit and facies as to make it highly improbable that they are the offspring of any other seed-bearing parent.

The winter-buds by means of which *P. crispus* is generally propagated are very remarkable, and in some degree resemble those of the grass-leaved Potamogetons. They differ, however, in being formed, for the greater part, in the axils of the leaves, rather than at the ends of the branches, and in the growing bud which produces the future plant being axillary instead of terminal, as in the *Chloephylli*. These winter-buds are well described and figured by Sauvageau, in his admirable notes on this genus in the *Journal de Botanique* for 1894, to which I refer the student for a complete account.

DISTRIBUTION in the British Islands.

P. crispus has been recorded from ninety-eight of Watson's vice-counties, and probably will be found in the remaining fourteen; those from which records are wanting are Cardigan, Montgomery, Merioneth, Isle of Man, West Perth, Easternness, Westernness, Argyle, Mid-Ebudes, North Ebudes, West Ross, East Sutherland, West Sutherland, Shetland. The "Cybele Hibernica" records it as occurring throughout Ireland in all the twelve districts, but adds, "not common in some parts of Ireland, as in Kerry and Donegal."

DISTRIBUTION in other countries. To the kindness of Mr. Arthur Bennett I am indebted for the following:—"Europe generally. Asia. In Asia Minor! to China! Siberia! Japan! Sumatra! Tibet! India, thirteen provinces. Africa. Egypt! Abyssinia! to Lake Nyassa—southwards to Natal! Transvaal! Cape Colony! Algeria. Mauritius (Baker's Flora), America, from Canada south to Virginia, west to Arizona. Australia, Victoria—Queensland. South Australia."

Explanation of the Plates 29, 30. POTAMOGETON CRISPUS.

PLATE 29, fig. 1. Mature branch with fruit-spikes and axillary and terminal winter-buds; Mepal, Cambridgeshire, August 25th, 1897; fig. 2, winter-bud; fig. 3, a more compact example towards Reichenbach's "var. *gemmifer*"; fig. 4, section of fresh fruit, mag.; fig. 5, section of stem, mag.

PLATE 30, fig. 1. Young state of the species, usually taken to be the *P. serratus* of Hudson, Sutton, Cambridgeshire, February, 1888; fig. 2, detached stipule with adnate leaves; fig. 3, base of lateral branch with similar stipules; fig. 4, fruiting branch of the flat-leaved state, to which I formerly proposed to restrict the name of "var. *serratus*," Sutton, Cambridgeshire, July, 1888; fig. 5, outline of part of leaf to show venation; fig. 6, winter-bud with young plant springing from an axil towards the base; fig. 7, dried fruit and section, mag.

14. × **POTAMOGETON COOPERI**, Fryer in Bot. Exch. Club Report for 1895 (issued June, 1897), p. 497.

P. UNDULATUS v. *COOPERI*, Fryer, Journal of Botany, vol. xxix. p. 289, pl. 313 (1891). *Potamogeton* × *Cooperi* = *crispus* × *perfoliatus*, Fryer, Exchange Club Report for 1895, p. 497 (June 5th, 1897). *P. perfoliatus* var. vel *P. heterophyllus*, Auct., var. *nitens*, Weber, J. Jackson in Report Botanical Record Club, pp. 150-1 (1880). *P. perfoliatus* var. *Jacksoni*, F. A. Lees, l.c. p. 150. *P. crispus* × *perfoliatus*, J. Baagöe in Botanisk Tidsskrift, 21. Bind, 2 Hefte, 1897, pp. 226-29 (reprint from, 1887). *P. perfoliatus* × *crispus* (*cymatodes*), Ascherson und Graebner, Synopsis der Mitteleuropäischen Flora, p. 337 (August 27th, 1897). C. Raunkjær, Danske Blomsterplanter, p. 104-106 (1896). *P. undulatus* var. *Cooperi*, Cooper in Transactions Leicester Lit. and Phil. Soc. p. 397, October, 1894. *P. undulatus*, A. Bennett in Irish Naturalist (June, 1894). *P. undulatus*, Fryer in Annals Scot. Nat. Hist. April, 1892.

SPECIFIC CHARACTER.—Rootstock perennial, growing throughout the winter; stem compressed like that of *P. crispus*; leaves flat, or slightly folded, undulated, entire, or with a few deciduous spines like those of *P. perfoliatus*; the young leaves are flat at first, narrow, strap-shaped, resembling those of *P. crispus*, but apparently with entire margins, which, however, under a lens, show a fine denticulation; as the season advances, the later-produced leaves are somewhat cymbiform, slightly hooded, scabrous at the margins and often with denticulations visible to the naked eye, all sessile from a rounded base, semiamplexicaul, 5-7 ribbed, 3 principal with 2-4 fainter and intermediate; blunt or slightly acute at the apex. The growth of the plant is often like that of a water-mint, being pyramidical in outline, but when the flower-spikes are produced, the upper branches floating just beneath the surface of the water assume the habit and look of those of *P. decipiens*; or, when the leaves are more closely folded and recurved, of *P. nitens*. Peduncles slender, equal or rarely a little thickened above, shorter than the subtending opposite leaves, but many times exceeding the very short ($\frac{1}{4}$ in.) oblong flower-spike; stigmas exserted, but flowers

always abortive, colour light or dark green, stem and young leaves sometimes beautifully tinged with bright crimson.

DESCRIPTION: ROOTSTOCK persistent, with stout strongly rooting terete stolons, the primary or barren internodes usually much shorter than the secondary or stem-bearing ones, one or more stolons springing from the stem-bearing node.

STEM compressed, channelled on the flattened broad sides, rounded on the narrow sides, which are furnished with a more or less prominent ridge, terete at the base like the stolons, but after the second node compressed and obscurely quadrangular like that of *P. crispus*, but somewhat irregular in shape according to the age or vigour of growth of the plant, a little twisted between the nodes; simple below at first, afterwards with branchlets from the axils of each leaf, after the first flower-spikes are formed repeatedly branched with alternate or opposite branchlets.

LEAVES semiamplexicaul, sessile, strap-shaped, oblong, to ovate or lanceolate from a rounded or cordate base; 5-7 or rarely 9-ribbed, the midrib and the principal conspicuous lateral rib on each side of it bordered by somewhat coarsely cancellate elongate areolations, intermediate ribs fainter, or some of them absent, cross-venation nearly at right angles to the ribs forming a coarse, regular network over the whole leaf; apex blunt, often slightly hooded and rounded, never mucronate, though sometimes narrowly contracted in lanceolate forms.

STIPULES scarious, truncate, soon decaying on the main stem, more persistent on the branches, on which they are a little shorter than, or exceed the internodes, two or three of the lower on both main stem and branchlets bearing adnate, ordinary leaves as in *P. crispus*, or rarely reduced to auricles as in *P. perfoliatus*.

PEDUNCLES slender, equal, straight, a little compressed, sometimes a little curved and thickened towards the middle, both forms occurring on the same plant; $1\frac{1}{2}$ -2 in. long, sometimes a little exceeding the subtending opposite leaves when full grown.

FLOWER-SPIKES very short, cylindrical or ovoid; flowers closed, but stigmas exerted, abortive.

COLOUR light to dark green, or yellowish-green tinged with crimson on the base of the young leaves and midribs, and on the stems and young stolons and winter-buds; generally drying darker.

HABIT in the very young state resembles that of *P. crispus*, the submerged barren shoots towards *P. perfoliatus* or *P. nitens*, the flowering branches like those of *P. decipiens*. By the synonymy it will be seen that at first I described this hybrid species as a variety of *P. undulatus* of Wolfgang, a plant of which I had been able to examine only one specimen, and that by no means a characteristic one. In 1895 Mr. J. Bangöe found a Potamogeton in Denmark which he and Mr. C. Raunkiær, by comparison with typical specimens collected by Wolfgang himself and preserved in the St. Petersburg Museum, identified as the true *P. undulatus* = *P. crispus* $\times$ *praelongus*;

and further, were able to distinguish it from plants erroneously named "*undulatus*" from Leicestershire and Cheshire. These two eminent botanists kindly sent me specimens, living and dried, of the Danish plant, together with a translation into English from Mr. Raunkjær's valuable natural history of the Danish flowering plants (part of which I append). I accepted their decision, and in the Exchange Club Report for 1895 proposed the name of $\times$ *Potamogeton Cooperi* for our British plant, and distributed many specimens amongst English and Continental botanists under that name.

At page 108 of Mr. C. Raunkjær's book, under the heading of *Potamogeton* hybrids, he writes:—" *Potamogeton crispus* $\times$ *praelongus* (= *P. undulatus*, Wolfgang). On an expedition, led by Mr. Baagøe, to the lakes and rivulets of Jylland in the summer of 1895, a *Potamogeton* in Gudenaå, which has proved to be *P. undulatus* of Wolfgang, as it agrees with his description as well as with the specimens of it in the Museum of St. Petersburg. . . . A plant has been found in several places in England which is thought by English botanists to be identical with *P. undulatus*, Wolfg., and which is considered by Mr. Fryer to be *crispus* $\times$ *perfoliatus*. The specimens I have had occasion to examine are rather different from our plant, as well as from the specimens in the St. Petersburg Museum, and several characters seem to prove that the English plant is really *P. crispus* $\times$ *perfoliatus*, but in that case it is not identical with *P. undulatus*, Wolfg., for the latter agrees perfectly with the plant found here in Denmark, and is, as we are about to see, without doubt *P. crispus* $\times$ *praelongus*.

"That *P. crispus* is one parent is evident enough from the form of the stem; in *P. undulatus* the stem is compressed, with a more or less deep furrow on each of the two broad sides; this characteristic form of the stem is also found in *P. crispus*, but not in any other Danish form of *Potamogeton*, nor, as far as I know, in any other species of *Potamogeton*.

"A comparative examination of the structure of the stem and leaves further proves that the other parent must be *P. praelongus*, and not *P. perfoliatus*. In *P. undulatus* numerous vascular bundles are found in the [false] bark, and these bundles are not to be found in *P. crispus*; in *P. perfoliatus* a few feeble bundles only are found, or they are quite lacking; on the contrary, *P. praelongus* has numerous bundles in the bark. As to the structure of the axial cylinder, . . . it appears that in *P. undulatus* it differs from *P. crispus* and approaches *P. praelongus*, and sometimes perfectly resembles the latter on this point, while there is no resemblance at all with *P. perfoliatus*.

"As to the nervation of the leaves, *P. undulatus* is just between *P. crispus* and *P. praelongus*. And at last, as to the margin of the leaves; *P. perfoliatus* has at least most frequently the leaves with dentate margins, and in *P. crispus* the leaves

are always serrulate, and in a hybrid between these two forms a serrulate margin of the leaves might be expected to be found; but both in the Danish Potamogeton and in the specimens of the St. Petersburg Museum of *P. undulatus*, the leaves are quite entire as in *P. praelongus*. The morphological as well as the anatomical proportions seem to prove that *P. undulatus* is *P. crispus* $\times$ *praelongus*.

"The Potamogeton found in England, and by English botanists referred to *P. undulatus*, differs from Wolfgang's and our *P. undulatus* by its serrulate leaves. Mr. Fryer writes that *P. undulatus*, Wolfgang, has also serrulate leaves, but Wolfgang himself says that the leaves have entire margins, and the leaves of the specimens of the St. Petersburg Museum have entire margins, as well as those growing in Denmark. Moreover, the vascular bundles of the stem's bark are also lacking in the English plant; on the contrary, strings are found just beneath the epidermis, as in *P. perfoliatus*. Finally, the English plant has more and more distant strings in the axial cylinder than occur in both *P. crispus* and *P. praelongus*, therefore it can hardly be supposed to be a hybrid between these two species; but it approaches very much to *P. perfoliatus* both in the structure of the axial cylinder as well as in several other characters; for instance, in the structure of the bark of the stem, habit, &c., and therefore I agree with Mr. Fryer in considering it *P. crispus* $\times$ *perfoliatus*, but it cannot in that case, as before mentioned, be referred to *P. undulatus*, Wölg., which is no doubt *crispus* $\times$ *praelongus*."

The above translation was kindly furnished by Mr. Baagöe, and I have only made one or two merely verbal alterations where necessary, to omit references to the figures of structure given in the original work, and to slightly alter the arrangement of the words in accordance with English usage. It is valuable in this place because it clearly shows the reasons for rejecting *P. Cooperi* as a variety of the true *P. undulatus*, as understood by Continental botanists, who have the command of better type specimens than we have in England.

I need remark on one point only, and in spite of all the efforts I have been able to make (very insufficient ones, it is true), I have not been able to clear it up: Did Wolfgang collect and distribute two forms under the name of *P. undulatus*? If so, I think one of them will prove to be *P. Cooperi*.

In my original description of *P. Cooperi* I say, l.c. p. 290, in reference to Wolfgang's description of *P. undulatus*:—"The minutely-toothed edges of the leaves escaped the notice of the describer, but they are present in Wolfgang's specimen in Nolte's Herbarium, and exactly resemble those of the British plant." Now this was also the opinion of an excellent botanist on the botanical staff at the British Museum, who examined the specimen with me. Possibly, however, from the condition of the specimen and owing to its being glued to the paper, a false appearance of "serration" may have been caused.

Except in the case of very carefully selected and well-dried specimens, it is often extremely difficult to name living plants from them, especially when the distinctions between closely allied hybrids such as the two in question have to be ascertained.

No such difficulty exists in distinguishing the living Danish and English plants, they are unlike in both appearance and habit! *P. Cooperi*, when growing, looks like *perfoliatus*, or *nitens*, or *decipiens*. *P. undulatus* might easily be passed over as a robust form of *P. crispus*. Mr. C. Raunkiær sent me living roots, and a drawing was made for this work of a plant grown from them, but unfortunately the number of plates allowed us is insufficient to permit us to publish it. This is the more to be regretted as *P. undulatus* is almost certain to occur in some part of our country. Mr. Raunkiær's lucid description will, however, serve for its identification, should anyone be fortunate enough to meet with it.

In my note-book for 1892 I find written: "Caspary's specimen, Herb. Brit. Mus., '*crispus* × *praelongus*' from Lake Delugi, Podgras, Prussia, is very near if not quite an *undulatus* form." This was written long before the discovery by Mr. Baagøe of the Danish plant and Mr. Raunkiær's identification of that with the plants of Caspary and Wolfgang, and has a certain degree of value as showing the close relationship between the two hybrids *undulatus* and *Cooperi*. In this particular I need only further add that my esteemed friend and correspondent Mr. Baagøe says, in discussing the difference between the British and Continental plants, that our Scottish form of *P. Cooperi* from Stirling approaches the true *undulatus* more than the English forms.

We now come to the question of varieties:—In some of the states of *P. Cooperi* as I first received them, or as I saw them in other Herbariums, the hybrid species seemed to divide itself into two very distinct varieties; but after long-continued cultivation I found the two forms were merely states, and occurred simultaneously on the same rootstock; that is, shoots of a *perfoliatus* aspect, and shoots which more closely approached the *undulatus* of Wolfgang, or rather, in our British examples, large-leaved *nitens* or *decipiens*. And further than this, under cultivation I found it impossible to distinguish the very dissimilar-looking forms from Scotland, Leicester, Chester, and Cambridgeshire; all assumed a uniform aspect. Hence, as in the case of *perfoliatus*, I am not able to divide the forms of this plant into varieties, but am induced to regard the varying forms as merely temporary *states*. Therefore the two varieties, *Cooperi* and *Jacksoni*, proposed in the Journal of Botany for October, 1891, p. 291, are here both included under × *P. Cooperi*; since, as I before said, both forms, that is the *perfoliatus* and *crispus* forms, occur simultaneously on one rootstock.

The species was first noticed, in the Report of the Botanical Record Club

for 1880, p. 150, under the name of *P. perfoliatus* var. *Jacksoni*, Lees, and was found by Mr. J. Jackson in "mid-west Yorkshire."

If I had at first identified this plant with the Leicestershire form named *Cooperi*, I certainly would have retained the name given by Mr. Lees, using it to represent the hybrid-species. But the true character of the species was not seen until its discovery by Mr. E. F. Cooper in the canal at Loughborough, and a little later by Mr. C. R. Billups in the Dee at Chester. Indeed it was chiefly the acuteness of the latter observer that led to the true relationship of the plant being determined.

DISTRIBUTION in the British Islands.

Mid-West Yorks, Leicester, Chester, Cambridgeshire, Gloucester, Salop, Anglesea, Stirling, and in Ireland, Antrim.

Description of the Plates 31, 32, × POTAMOGETON COOPERI.

PLATE 31, fig. 1. Upper part of flowering stem; fig. 2, secondary branch with axillary branchlets; fig. 3, section of stem, mag.; figs. 4 and 5, detached stipules with adnate leaves. From specimens collected by Mr. E. F. Cooper from the Canal, Loughborough, Leicestershire, August, 1897.

PLATE 32, fig. 1. Complete young stem, before flowering; fig. 2, part of flowering stem. From Stirlingshire, collected by Colonel J. S. Stirling and Mr. R. Kidston, September, 1897.

15. × **POTAMOGETON BENNETTII**, Fryer in Journal of Botany, vol. xxxiii. p. 1 (Jan. 1st, 1895).

× *P. BENNETTII*, Fryer, l.c. p. 1, Tab. 348. *P. crispus* × *obtusifolius*, Ascherson u. Graebner, Synopsis Mitteleurop. Flor. p. 349 (1897).

SPECIFIC CHARACTER.—Rootstock growing throughout the year, with slender terete stolons creeping on or near the surface of the soil and rooting at the nodes. Stem compressed, slightly channelled on the broad sides, rounded at the corners, smooth on the surface and less angular than that of *P. crispus*; slender, simple below, repeatedly branched when in flower, and then with a tendency to produce short axillary fascicles of leaves, like *P. Friesii*; always springing in the first instance from the apex of a terminal deciduous winter-bud, as in the *Chloephylli*, afterwards irregularly from the nodes of the stolons thrown out at the base of the primary stem; intermediate in compression and general facies between the stems of *P. crispus* and *P. obtusifolius*. Leaves all similar, alternate, opposite beneath the peduncles only, never beneath the barren branches; linear, acute, rarely a little blunt, narrowed towards the base, which is sometimes furnished with glands on the stem like those of the *Chloephylli*; all sessile, 3 nerved, or rarely with two additional incomplete nerves just within the margin as in *P. crispus*, cross-venation distant,

ascending, simple, or a little branched towards the apex of the leaf; midrib bordered with elongate areolations, and so, wide and conspicuous; margin inconspicuously serrulate in the early stem-leaves, but entire, or with very minute spine-like serrulations towards the apex not visible to the naked eye, nor when dried. Stipules scarious or subherbaceous, with two conspicuous green ribs, truncate, clasping, persistent on the branches, usually decayed on the main stem by the time of flowering; shorter than the internodes on the main stem, equalling or exceeding them on the branches. The one first formed at the base of the stem rarely with an adnate ordinary leaf, and then closely sheathing, green and herbaceous with the upper part scarious. Peduncles slender, equal, falling short of the subtending opposite leaves, but many times longer than the very short ovoid spike. At first terminal, afterwards placed in the fork of the two branches which rise from the axils of the subtending leaves, or lateral through the abortion of one of these. Flower-spike not exceeding $\frac{1}{4}$ in. in length, very few-flowered, sepals remaining closed, but stigmas exerted, always abortive, soon decaying. Colour of the whole plant dark green, tinged with reddish-brown as growth advances towards maturity. Probably a hybrid between *P. crispus* and *P. obtusifolius*, or *P. Friesii*.

DESCRIPTION: Rootstock persistent, growing throughout the year, with slender terete stolons creeping on or near the surface of the soil, strongly rooting at the nodes.

STEM very slender, compressed, the broad sides less deeply channelled than in *P. crispus*, more rounded at the corners, and less angular in outline.

LEAVES all similar, sessile, linear, narrowed towards the base, acute or rarely a little blunt at the apex, 3-5 ribbed, but the outer pair rarely developed, and close to the margin as in *P. crispus*. The two principal lateral ribs forming a complete arch some little distance below the apex; the midrib bordered by rows of fine elongate areolations, and so, conspicuously broad, cross-venation distant, ascending, simple, or a little branched towards the apex of the leaf. Sometimes glands are developed at the base as in the *Chloephylli*. In the autumn the terminal shoots become horny and the tips of the leaves decay, thus forming a deciduous winter-bud intermediate in character between those of *P. crispus* and *P. Friesii*, and resembling the *Chloephylli* in the young stem growing from the apex of the bud, instead of being axillary from near the base as in *P. crispus*. Consequently this species is never propagated by any part of the stem; nor do these horny winter-buds ever form in the axils of the leaves; but, as far as I have been able to observe, are strictly terminal.

PETIOLE none.

STIPULES scarious to herbaceous, closely clasping and the lower part almost surrounding the stem, hyaline, whitish, or pale green, with two prominent green ribs on the back, apex truncate; persistent, but much decayed on the stems by the time

of flowering and falling short of the internodes, on the branches equalling or exceeding them. The first-formed one at the base of the main stem very rarely with an adnate leaf of the ordinary shape.

PEDUNCLE slender, equal, $1-1\frac{1}{2}$ in. long, usually falling short of the opposite subtending leaves, rarely a little exceeding them.

SPIKE very few-flowered, ovoid or roundish, sepals not expanded, but stigmas exserted, flowers always abortive, soon decaying.

FRUIT none.

HABIT intermediate between *P. crispus* and *P. obtusifolius*, resembling the latter in the full-grown plant, but in the young state rather like that of *P. crispus*; growing throughout the winter, the rootstock continually increasing by stolons growing from the stem-bearing nodes; new plants produced by the deciduous winter-buds.

COLOUR dark green, tinged with reddish-brown in the mature plant.

REMARKS: This hybrid species is founded on a plant gathered from Wood Pond, Grangemouth, Stirlingshire, by Messrs. R. Kidston and Col. J. S. Stirling in 1890. This was without flowers and was mistaken by the discoverers for a variety of *P. obtusifolius*; and also was named *P. crispus* by an eminent authority. In 1894, however, the plant was found well in flower, and a plentiful supply of fresh specimens confirmed the opinion the two Scotch naturalists had formed, by repeated examination of the growing plant, that it is in all probability a hybrid between *P. crispus* and *P. obtusifolius*. I am indebted to the generosity of Messrs. Kidston and Stirling for an abundant supply of living plants and for permission to describe and name the plant in the Journal of Botany. On referring to that description it will be seen that the form was supposed to be identical with the imperfectly known and described *P. serrulatus* of Regel and Maack (Tent. Flor. Ussuriensis, p. 139). But subsequently Mr. Bennett has succeeded in identifying that form with *P. Robbinsii*, Oakes, so that our Scottish plant probably was not previously described. In 1897 a further supply of living specimens was sent from Scotland; one of them has been under almost daily observation ever since, and, having flowered freely for two seasons, has confirmed the supposition of its hybrid origin, and also supplied ample materials for the natural history given above.

I have seen no specimens of this hybrid from other localities, but it probably may be found in collections of Potamogetons under the name of *P. obtusifolius*, or of *P. crispus*.

Description of the Plate 33. × POTAMOGETON BENNETTII. Fig. 1, young barren stem from Stirlingshire, collected by Messrs. Kidston and Stirling, September, 1897; fig. 2, flowering stem; fig. 3, flower, on spike, mag.; fig. 4, section of stem, mag.; fig. 5, winter-bud, as seen in autumn; fig. 6, winter-bud, with young plant springing from the apex. All from plants cultivated at Chatteris, 1899.

17. **POTAMOGETON DENSUS**, *Linn.*, Sp. Pl. ed. 1, p. 126 (1853).

P. DENSUS, Cham. et Schlecht. *Linnaea*, vol. ii. p. 160, Tab. 4, fig. 1 (1827). Fieber, *Pot. Böhmens*, p. 13, Tab. 1, fig. 1 (1838). Reichenbach, *Icon. Flor. Germ.* vol. vii. p. 18, Tab. 28, figs. 46-49 (1845). Koch, *Synop. Flor. Germ.* ed. 2, p. 781 (1844). Bab. *Man.* ed. 7, p. 376 (1874). Hooker, *Stud. Flor.* ed. 3, p. 434 (1884). Syme in *Eng. Bot.* ed. 3, p. 44, pl. 1414 (1869). Ascherson u. Graeb. *Synop. Flor. Mitteleurop.* p. 354 (1897). Tiselius, *Pot. Suec. Exsic.* fol. 147 (1897).

? *P. serratum*, Huds. *Flora Anglica*, p. 61 (1762).

SPECIFIC CHARACTER.—Rootstock persistent with stolons creeping on or near the surface of the soil, with many strong deeply-rooting fibres at the nodes, stem terete, simple at first, branched after flowering; leaves all similar, opposite, sessile, the base embracing the stem, in pairs, or rarely in threes, lanceolate or linear lanceolate from a cordate base, flat and ascending, or channelled and reflexed, entire or very minutely serrulate, with somewhat distant spines towards the obtuse apex, 3-5 ribbed: peduncles short, slender, equal, erect and emerged at the time of flowering, afterwards reflexed and submerged, spike usually 4-flowered. Fruit orbicular cochleate, with an acute winged herbaceous keel, lateral ridges obsolete, embryo spiral of $1\frac{1}{2}$ whorls. Colour light to dark green, sometimes tinged with yellow or brown.

DESCRIPTION: Rootstock persistent, growing throughout the winter, with stout stem-like stolons creeping on the surface of the soil, or near it, rooting strongly at the nodes.

STEM terete, slender, simple until flowering, afterwards branched, usually in pairs; one or more stems rising irregularly from successive or alternate nodes; the internodes on the lower part of the stem equal or exceed the leaves in plants growing in deep water, on the upper part falling much short of them, and are sometimes almost obsolete, so that the leaves form densely imbricated clusters; or in shallow water the whole of the internodes are much shorter than the leaves from base to summit.

LEAVES opposite or subopposite, in pairs, rarely in threes, sessile, amplexicaul, the base of one enfolding the other; the base adhering to the stem and so a little sheathing, thence suddenly expanded and subcordate, flat, or more frequently folded and recurved; 3-7 ribbed, midrib broad, with many rows of parallel elongate veins, the two principal lateral ribs joining it at some distance from the apex, the others, when present, seldom extending beyond the middle of the leaf and then joining the lateral ribs, cross-veins few, irregular, and distant; margins scabrous, with minute spine-like denticulations, but apparently entire to the naked eye; flaccid, or rigid, flat, or folded, patent, ascending or recurved, distant or densely imbricated, according to depth of water, or temporary conditions of growth.

STIPULES absent except in a pair of leaves which subtend the peduncle, and in the pair at the base of the branches. "Adnate and adhering to the leaf at the base, and with an oblong free portion" (Syme). The stipules are obscure and difficult to observe, no author I have been able to consult sufficiently describes them, and I have not been able to arrive at any definite conclusion as to their relationship to the stipules in other species of *Potamogeton*; or as to whether they are stipules at all in the sense of the word as used in the preceding descriptions.

PEDUNCLES slender, equal, shorter than the subtending leaves, erect and emerged in flower, strongly recurved and submerged afterwards.

FLOWER-SPIKE very short, 4-flowered.

FRUIT lenticular, ventral margin oblique towards the base, beak slender, central, a little recurved, winged on each side by a decurrent herbaceous growth extending about half-way down the drupelet on the ventral, and to the base on the dorsal margin, on the latter forming a prominent acute winged keel. The nutlet is obtuse on the dorsal margin, showing no trace of a keel; embryo spiral, of $1\frac{1}{2}$ whorls.

COLOUR light green, or dark green, tinged with yellow or brown in boggy stations.

HABIT: growing throughout the year, and flowering and fruiting from May to September. The upper branches frequently throw out long roots, thus forming new plants when the lower stems decay.

REMARKS: *P. densus* is not very nearly allied to any other species in the genus, and probably will be separated under a subgeneric name, at least, when the whole of the *Potamogetons* shall be sufficiently known to make a monograph of the genus possible. Every writer on these plants seems to have recognized the difficulty of placing this species in its proper position. Koch, in his arrangement of the genus, made it the sole occupant of his last division "*Enantiophylli*"; which, following Chamisso and Schlechtendal, he placed after the "*Coleophylli*" or *pectinatus* group. This is apparently about as unnatural a location as could be made, looking at the external characters of the two groups alone. Fieber, whose knowledge of the European species has seldom, if ever, been surpassed, places it in his first division, and follows it by *P. perfoliatus*—the first species in his second division. This certainly seems a little better, but still with the unfortunate result that his arrangement of the genus begins neither with the highest—or most specialized forms—nor with the lowest—or least specialized. Reichenbach has placed it after the "*Chloephylli*," or grass-leaved section, and before *P. crispus*, which is practically what is done in the present work, not because this is at all a natural arrangement, but because it is difficult to know where else to place it. And so the position it occupies in the "London Catalogue of British Plants" is followed, although *P. densus* has no closer relationship to *P. crispus* than it has to *P. pectinatus*. Messrs. Ascherson and Graebner, in their "Flora of Middle Europe," quote the name "*Groenlandia*, Gay Compt. rend. Ac. Sc.

Paris, xxxviii. (1854) 702," in a generic sense, but as in their very comprehensive work they have not adopted it, probably they think it proposed on insufficient grounds.

I regret that I have been able to add little or nothing to the life-history of this species. It is difficult to cultivate, and difficult to observe in the natural stations crowded with vegetation which it usually affects.

Distribution in the British Islands.

P. densus is decidedly English and Eastern in distribution rather than Northern and Western. In England and Wales it has been recorded from all the seventy vice-counties except West and East Cornwall, North (and South?) Devon, Monmouth, Salop, Brecon, Radnor, Pembroke, Cardigan, Montgomery, Merioneth, Carnarvon, Denbigh, and Flint. In Scotland it is recorded only from Lanark, Roxburgh, Haddington and Edinburgh. The "Cybele Hibernica" records it from Districts I. to VII., "Lowland . . . rare and local."

Distribution in other countries.

Mr. Arthur Bennett, to whose unwearied kindness I am constantly indebted, gives me the following list:—Europe: Sweden! Denmark! Norway! France! Belgium! Schleswig! Austria! Spain! Portugal! Italy! Greece! Switzerland! Russia. Lithuania. Urals! Poland! Asia: Himalaya! Afghanistan! Turkey in Asia! Kurdistan! Africa: Tunis! Algeria!

Explanation of Plate 34, POTAMOGETON DENSUS.

Fig. 1, part of stolon with young unbranched stem; fig. 2, upper branched part of stem, with flowers and fruiting spikes, and with a rootlet (the lower branch to the left, with rootlets at the base, is detached from the stem); fig. 3, a flower-spike, mag.; fig. 4, detached leaf, mag. (the lateral ribs are erroneously represented as extending much beyond the middle of the leaf); fig. 5, section of stem, mag.; fig. 6, section of dried fruit, mag.; fig. 7, fresh fruit, mag.; fig. 8, section of same, mag. All drawn from specimens gathered at Mepal, Cambridgeshire, August, 1898.

THE LUCENS-GROUP.

Before proceeding with the specific descriptions of the closely related forms which have been variously described as sub-species, or varieties, or hybrids of *P. lucens*, and which together constitute the *lucens*-group, perhaps the most interesting in the genus, it is necessary to give some account of its treatment by former naturalists; and an explanation of the method adopted in the present work.

Linnaeus (*Species Plantarum*, 1753) described the two extreme forms under the names *P. lucens* and *P. gramineum*; the latter name undoubtedly in part included the *P. heterophyllus* of this work, but as it was confounded by him with *P. obtusifolius*, and as it is somewhat doubtful as to which of the more recent segregates the plant originally described belonged, it is impossible to retain the name in the present work. No doubt *P. lucens* of Linnaeus is the plant described by subsequent authors, and as it has no real varieties in Britain, nor, as far as I have seen, in other countries, but only variable *states* of one species, the original name can be used without danger of including any other "species" or "sub-species."

The next form described as a species was the *P. nitens* of Weber (*Supp. Fl. Holsatiae*, 1787); this is a hybrid and in fact includes several hybrids of varying origin, the result of crosses of *P. heterophyllus*, or some of its segregates, with *P. perfoliatus*. Recent authors usually include hybrids between any species of the *lucens*-group in the aggregate "*P. nitens*." Chamisso and Schlechtendal doubtfully described a *nitens* form as a sub-species of their *P. Proteus* (which included the whole group), under the name of *P. curvifolius*, *Hartm.*

Another species has been generally recognized as the *P. Zizii* of Koch, in *Fl. German*, 1827, and Nolte named a form allied to this *P. lucens* var. *coriaceus*.

One more has been generally accepted as a species belonging to the *lucens*-group:—*P. decipiens* (Nolte in litt.), Koch *Fl. Germ.*, p. 779, sp. 10, 1844. This "species" includes many forms, all hybrids, with *P. perfoliatus* as one parent, and so approaches *P. nitens*.

The following species are usually placed immediately after *P. natans* and the allied forms, a method not adopted in the present work because an endeavour was made to propound a linear arrangement as far as possible—the only natural sequence for restricted groups for the following of Darwin's theory of evolution, not only because there seemed no place for them in the series already described, but also because this group affords the rare instance of a set of plants which at its two extremities presents forms which have not the slightest apparent resemblance to one another, and

yet are connected by a series of graduations so evidently intermediate that we are continually arrested by plants which can with equal propriety be placed under two of the "species" or "sub-species" into which botanists have divided the forms of this remarkable group. Thus the term "species," as used in the following chapters, must not be accepted as bearing the full significance that it has when applied to forms so distinct as *P. natans* and *P. pectinatus*, but merely denotes names by which we may distinguish "convenient bundles,"* and so give as full and as necessary an account of them as of the first, which are at present connected by no linking forms. Freely admitting the difficulty of clear definition, I must in the present instance contend for the advantage that results from the naming lower groups of the large groups with quasi-specific names, since by so doing we not only assist memory by writing *P. falcatus* instead of *P. gramineus*, var. b, but also by such divisions aid our mental conception of the origin of what we may call "distinct, true species."

REMARKS: The species included in the present section of the genus pass by almost imperceptible gradations from *P. lucens* to *P. heterophyllus*, the two extremes of the group; the series of intermediate forms is so complete that it seems to satisfy the demand of the Darwinian postulate that if all the forms that have existed in time and space could be laid before us "all would blend together by steps as fine as those between the finest varieties." The fruit, too, on which many botanists found the species in this genus, passes by similar insensible characters of sculpture and size from the small to the large forms. In fact from the examination of a full series of dried specimens it seems at first sight difficult to avoid the conclusion that they represent one extremely variable species.

As above, Chamisso and Schlechtendal in their admirable monograph of the genus (Linnea, vol. ii. 1827), unable to find distinct specific characters, united all the forms known to them under the name of *Potamogeton Proteus*, ranking under this super-species the previously named European forms, *P. lucens*, Linn., *P. Zizii*, M. et K., and *P. heterophyllus*, Schreb. as sub-species; adding "*P. (Proteus?) curvifolius*, Hartm." doubtfully because the fruit was unknown to them. This latter is a barren hybrid and the fruit remains unknown to this day, although almost every subsequent writer on the genus has described supposed specimens of it.

* Mr. A. H. Evans, in his admirable "Birds" in the "Cambridge Natural History" (p. 15), has put the case of "genus," "species" and "race" so well that I cannot forbear quoting him in full: "It cannot be denied that genera and species are merely 'convenient bundles,' and that divisions of either, if carried too far, defeat the object for which classification is intended. Genera are only more distinct from species, and species from races, because the intervening links have disappeared; and, if we could have before us the complete series which, according to the doctrine of evolution, has at some time existed, neither genus nor species would be capable of definition, any more than are races in many cases; while the same remark will apply to the larger groups."

This treatment of the group seems to have been founded on the examination of herbarium specimens, and so far looks a perfectly natural arrangement, but on studying the living plants and their local distribution and life history it appears to me to completely break down.

With many students of the genus in the present day, I long held the view that the whole group had directly descended from *P. lucens*, Linn., or even that the members were merely extreme varieties constantly produced under favourable conditions, and freely interbreeding with the parent plant and with one another. But after thirty years' observation of countless thousands of the living plants, I can come to no other conclusion, erroneous though it may be, than that the three above named plants have descended from some extinct or unknown ancestor.

Whatever may be the origin of the mass of closely related forms, it is necessary to arrange them as naturally as possible. The first step is to separate the hybrids which have been described and named as true species and, for reasons which will appear later on, to rank them under the parent species they most resemble; the next to assign its proper place to each of the modern species which have been taken from the older aggregates described under the names of *lucens*, *Zizii*, and *heterophyllus*. Then the real difficulty begins: it is to reduce to some sort of order the innumerable forms which may be either crosses between varieties of the aggregate species, or merely extreme varieties of any one of them.

When botanists began to admit that natural hybrids did occur in the genus, many collectors of Pond-weeds regarded all considerable deviations from the type as due to hybridity, and ranked them even as crosses between three or more forms! Such mongrels do occur, but they are very rare; I have not met with above one or two cases where the variation apparently depended on the mixture of more than two forms. The most remarkable thing about the supposed origin of certain forms is that the parentage was assigned on one side to hybrids like *P. nitens*, which no one can find producing a fertile drupelet.* The last difficulty and probable source of error is this, every species throughout the group, and one may almost say every individual, seems at some time or other to undergo a parallel series of variations from the earliest to the latest stages of growth, and it may happen that two segregates may vary away from

* We must bear in mind, however, that many eminent botanists hold an opposite view, and believe that *P. nitens* is occasionally fertilized by one or other of its parent species, and so produces a form which is capable of being fertilized by a third species in which other species are mingled together. This may be so, but it is not borne out by a single example of a perfect fruit of *nitens* in any Herbarium known to me. At one time I agreed with those who believe that such mature fruits do occur, but all the specimens sent to me prove, on careful examination, to be true *P. heterophyllus*, by characters apart from the fruit which presents no distinctive characters that I can discover.

their types but towards the like stage of a third species, and in such a case it is not easy to decide the true rank of such approximating forms.

In spite of this complex state of closely allied forms and interlocking variations some attempt must be made to define the forms that would be good and distinct species if the intermediates were taken away, because it is in such groups as this that the course of nature in creating species can be best studied. Evidently the only plan is to segregate the most distinct types and treat them as if no linking form existed. No one who has had the opportunity of watching the growth and distribution of these plants can possibly suppose them to belong to one species; that is to say, that they are all forms of *P. lucens* or of *P. heterophyllus*. The practice of the earlier botanists of considering forms united by intermediates as necessarily belonging to one species is discredited by the whole life-history of the members of this interesting *lucens*-group. When I first studied them in the living state I thought all the forms had descended from *P. lucens*, of which *P. Zizii* seemed to be a variety still being produced, but a succession of very dry summer evenings led to a knowledge of *Land-forms* and the fact that *P. lucens* never had a land-form even under the most favourable circumstances, but invariably died down as the places in which it grew became dry, while *Zizii* and its segregates flourished and made vigorous growth and tuberous stolons for the next year's growth. Even when exposed on dry land to the full rays of the sun I saw that no recent production of such a variety was possible, and no previous one probable. Here, then, was the first doubt on the descent theory. Two or three years' further investigation showed that *Zizii* and *heterophyllus* were closely allied, and probably crossed with each other freely. The questions then arose, Are these two latter forms one or two "species," and, if distinct at present, has one directly descended from the other? The answer to the first question seemed to be that they were two species; the second question is answered under the head of *P. Zizii*. For reasons there given I came to the conclusion that all these species had a common unknown ancestor. To make one species of these three would entail sinking several generally adopted segregates to the rank of sub-species; as *P. polygonifolius* and *P. coloratus* under *P. natans*, to which they are not even nearly allied, few of our British forms being more distinct than the two forms, and *coloratus* probably belonging to a different section of the genus.

18. × **POTAMOGETON LANCEOLATUS**, Smith, English Botany, pl. 1985 (1808).

P. LANCEOLATUS, Bab. Man. ed. 7, p. 372 (1874). Hooker, Stud. Flor. ed. 3, p. 432 (1884).

DESCRIPTION : ROOTSTOCK very slender, almost filiform ; stolons springing from a tuberous rhizome.

STEM filiform, very slender, branched from the base, with axillary branchlets.

LEAVES all with elongate cancellate venation, glandular at the junction with the stem ; lower elongate, somewhat strap-shaped, gradually narrowed towards the sessile base, rather abruptly towards the rounded or blunt, obscurely mucronate apex. Upper subcoriaceous, stalked, opposite, oval or lanceolate-ovate, somewhat blunt at the apex.

PETIOLE terete, shorter than the subtending leaves.

STIPULES short, narrow, acute, scarious, subpersistent, soon decaying on the main stem.

PEDUNCLE terete, equal, somewhat curved, $\frac{3}{4}$ –1½ inch long, shorter than subtending leaves, stouter than stem.

SPIKE oval, very short, 2–3 lin.

FRUIT imperfect, abortive. Up to the date of writing I have never been able to examine a perfectly developed fruit.

COLOUR dark green.

HABIT : a slender plant resembling *P. Friesii* in its early states, and an extremely small *P. Billupsii* or *P. varians* in its mature state.

REMARKS : *P. lanceolatus*, Smith, is certainly a hybrid, probably *P. heterophyllus*, × *P. pusillus* (not × *graminifolius*, Fries) : *P. pusillus* from Exning, near Newmarket, sometimes has floating subcoriaceous leaves subtending the peduncles, which are netted with chain-like areolations, like those of *P. lanceolatus*. The French plant differs slightly from ours ; it is much larger, and may possibly be *P. heterophyllus* × *Friesii* (that is, assuming that ours is × *pusillus*). *P. Friesii* also has very rarely “floating” leaves with chain-like areolations.

Distribution in the British Islands.

This species has only been found in Great Britain in Cambridge and Anglesea. Ireland : Clare, Galway (var. *hibernicus*, A. Benn).

Distribution in other countries :—

France : Saon et Loire ! Switzerland : Berne ! Lanenberg : Nolte ! Island of Corfu : Letourneux !

Plate 39, POTAMOGETON LANCEOLATUS, was drawn from Cambridgeshire specimens, obtained between Burwell and Wicken in August, 1880. The smaller figure is to show the elongate, nearly linear leaves produced in late autumn (from specimens grown at Croydon by A. Bennett).

19. **POTAMOGETON GRAMINIFOLIUS** (Fr.), Novit. Flor. Suec. ed. 2, p. 36 (1828).

P. GRAMINEUS, Linn., Flor. Lapp. ed. 2, p. 46, Fries, Novit. Flor. Suec. ed. 2, p. 36, var. α , *graminifolius* [p. 37, var. β , *heterophyllus* (1828)]. Ascherson, Synop. Flor. Mitteleurop. p. 231 (1897). Fryer, Journal of Botany, vol. xxx. p. 33, Tab. 317, 318. Koch, Synop. Fl. Germ. ed. 2, p. 777 (1844). *P. heterophyllus*, Schriebl, Fries, Pot. Böhmens, p. 28, Tab. 1, fig. 12 (1838).

DESCRIPTION: ROOTSTOCK persistent, with numerous long, rather stout stolons, often with the terminal internodes thickening to moniliform tubers.

STEM slender, rising irregularly from the nodes, with numerous submerged barren branches at the base, the upper part simple, sparingly branched after flowering.

LEAVES: lower submerged, undulate, entire, not serrate nor denticulate, flat and ascending, or folded and recurved, narrow, attenuated towards both ends, sessile but not amplexicaul, lanceolate acuminate, lowest sometimes reduced to a midrib. Upper long-stalked, petiole usually, often greatly, exceeding the lamina, coriaceous, or rarely membranous, elliptical, oblong, oval, or suborbicular; lamina suddenly contracted into the petiole, or sometimes gradually narrowed at the base and a little decurrent.

PETIOLE of upper leaf usually longer than the lamina.

STIPULES persistent, herbaceous, small, narrow, becoming involute and then appearing acute, larger and broader on autumnal barren stems, cymbiform beneath the peduncles.

PEDUNCLE much thicker than the stem, slightly swollen upwards, usually curved from the base, terminal, longer than the flower spike, often very much so, exceeding or falling short after subtending foliage.

SPIKES dense, cylindrical, $1-1\frac{1}{4}$ in. long, few, usually produced singly, terminal.

FRUIT "roundish or obliquely obovate, $\frac{3}{4}-1\frac{1}{2}$ lin. long by $\frac{3}{4}$ wide to 1 lin. broad, slightly curved or angled on the ventral side, rounded or indistinctly 3-keeled on the back, the obovate forms with an inward basal curve on the face and an indentation running into it from the centre of the sides; style short, obtuse, apical, rarely facial, apex of the embryo nearly touching the base, pointing slightly inside of it." (Morong.) Central keel usually acute when immature, lateral keels always obscure, and when ripe the central one also.

COLOUR light to dark green, sometimes tinged with red or reddish-buff, especially in the young states of the plant.

HABIT much as *P. heterophyllus*.

REMARKS: *P. graminifolius* differs from *P. varians* by the absence of the spatulate, coriaceous, submerged autumnal leaves, and of the membranous *Zizii*-like summer foliage; the early state, too, is very different. I consider it a valid species. From *P. falcatus*, the species to which *P. graminifolius* is most nearly allied, the distinction is equally well marked. In *P. graminifolius* the lower leaves are sessile, instead of semiamplexicaul; the fruit is much larger and less freely produced, the flower-spikes are single, instead of in clusters, and with nearly equal, instead of swollen, peduncles. Again, the early spring states widely remove the two forms, and under no state can *P. graminifolius* be mistaken for *P. nitens*.

[On mature consideration Mr. Fryer decided to assign full specific rank to this form, and spoke of this as his last word on the subject.]

Distribution in the British Islands.

England: Cambridgeshire, Anglesea, Hunts. Scotland: Perth! Shetland! Caithness. Ireland: Frequent in the rivers (Fryer).

The specimens drawn on Plate 36, POTAMOGETON GRAMINIFOLIUS, were from Pidley Fen, Huntingdonshire.

20. POTAMOGETON HETEROPHYLLUS, Schreber, Flora Lipsica, p. 21 (1771).

P. GRAMINEUS, L. Sp. Pl. ed. 1, 127 (1753). E. B. ed. 3 (1869), vol. ix. p. 35, Pl. 1406. Fries, Novit. Flor. Suec. ed. 2, p. 37 (*P. gramineus* var. β , *heterophyllus*, 1828). *P. Proteus heterophyllus*, Cham. et Schlecht. Linnæa, vol. ii. p. 197 (1827). *P. heterophyllus*, Fryer, Journal of Botany, vol. xxv. p. 163 (1887). *P. gramineus*, Raunkiaer, Danske Blomsterplanter, p. 75, Fig. 38 (1896).

DESCRIPTION: ROOTSTOCK persistent, fibrous, with many far-creeping stolons, those produced late in the summer with moniliform tubers, especially those from the barren autumnal growths and from the land-forms.

STEM slender, flexuous, ascending, much and repeatedly branched at the base with submerged branches forming a dense mass of submerged vegetation, upper part often simple with longer internodes until it reaches the surface, then with one or many short flowering branches with or without floating leaves.

LEAVES very variable; lower linear-lanceolate, narrowed towards both ends, acute, often apiculate, undulate, flat, or more frequently folded and a little recurved, 5-7-ribbed, with rather distant ascending more or less regular cross-venation. Upper elliptical to orbicular, lamina usually falling short of the petiole, floating, coriaceous or rarely membranous, often absent in the flowering plant, belonging to the vegetative system, lowest reduced to a midrib, or bodkin-pointed.

PETIOLE narrow, slightly expanding below the lamina, coriaceous, shorter than, equalling, or often greatly exceeding the lamina.

STIPULES acute, with a fine mucro like the leaves, only blunt when this decays, herbaceous or scarious, small and clasping, afterwards patent, usually about half the length of the internodes, but in the upper branchlets sometimes exceeding or falling short of them, often wide and cymbiform below the peduncles, two-ribbed on the back, one or both of the ribs produced beyond the tip into an acute scarious mucro, which soon decays.

PEDUNCLES stout, swollen upwards, especially when the fruit is ripe, 2-4 inches long, rarely produced by local conditions to 6-9 or more inches long (*forma longipedunculata*), but this does not warrant the varietal name given by many authors, as it may occur on any form of the species, and is not constant, being probably due to constantly rising or rapidly running water. I have never seen fully matured fruit on such peduncles in this section of the genus. By the abortion of the upper internodes of the stem 6 or more peduncles start from the same tuft of leaves and stipules.

SPIKE 1-2 in. long, shorter than the peduncles, densely flowered and fruited.

FRUIT: Drupelets small, flattened at the sides; dorsal margin blunt, with prominent keel and ridges, or acutely keeled; upper margin straight, terminated by the short beak; variable.

COLOUR green or olive-green.

HABIT: Variable (*See* Fryer, *Journal of Botany*, 1887, p. 164), somewhat that of *P. Zizii*.

Distribution in the British Islands.

P. heterophyllus has a wide range both in Great Britain and Ireland, occurring from Shetland to the English Channel, with the exception of:—E. Sutherland, Westernness, Linlithgow, Edinburgh, Haddington, Roxburgh, Cheviotland, Isle of Man, Derby, Flint, Merioneth, Montgomery, Cardigan, Pembroke, Radnor, Brecon, Warwick, Worcester, Hereford, Monmouth, Gloucester, Bedford, W. Norfolk, Bucks, Middlesex, W. Kent, W. Sussex, S. Hants, Isle of Wight, Dorset?, S. Wilts, S. Somerset, Devon and Cornwall. Of the 76 Watsonian vice-counties where it is found, comparatively few are on the Eastern borders or in Wales. In Ireland it is found in thirty-one of Praeger's districts.

Distribution in other countries.

Europe: Norway from 71° 2' to Italy. France to Lithuania (Russia). Asia: Siberia from 70° to Southern Manchuria and Japan. N. America: Greenland. Canada from lat. 51° to Quebec and from Ottawa to Vancouver's Island. United States (Maine to N. Carolina and New England to California). S. America: Venezuela.

Plate 35, POTAMOGETON HETEROPHYLLUS, represents the usual plant of the Fens near Chatteris.

21. × **POTAMOGETON VARIANS**, Morong, in Fryer, Journal of Botany, 1889 (February), p. 33. [Plate 287 published subsequently, in March, 1889.]

P. SPATHULÆFORMIS (Robbins), Morong, in Naiad. N. America, p. 26, Plate 35 (1893). *P. varians*, Fryer, Journ. of Bot. vol. xxvii. p. 33.

DESCRIPTION : ROOTSTOCK tuberous.

STEM slender, usually simple below, with a few branches above the middle, not divided into secondary branchlets, or in shallow water with a few branches from the base, and then rarely with very short secondary branchlets.

LEAVES : Lowest submerged leaves reduced to phyllodes, or narrow linear-lanceolate bodkin-pointed ; ordinary submerged leaves sessile or stalked, narrowly lanceolate, attenuated towards each end, or spathulate-oblong, or obtuse, mucronate, flat and ascending, or rarely folded and recurved. Floating leaves alternate, obovate, oblong, spathulate or orbicular, coriaceous, rarely membranous, long stalked, belonging to the barren state of the plant, and never directly sustaining the flower-spike.

PETIOLE very long, coriaceous.

STIPULES narrow, slender, herbaceous, persistent, blunt or contracted into a short mucro when dry.

PEDUNCLES lateral, not necessarily subtended by opposite coriaceous leaves, but usually springing from the stem opposite a membranous leaf, or rarely opposite a stipule only ; slightly thicker than the stem, not swollen upwards, as long as the opposite leaf.

SPIKE $\frac{2}{3}$ –1 in. long, rather slender, not dense.

FRUIT : Drupelets small, flattened and impressed at the sides, nearly circular in outline, with a short subcentral beak ; central keel acute, almost winged ; lateral ridges prominent.

COLOUR reddish-green or light green, drying darker ; the lower part of the stem and young shoots sometimes bright red.

HABIT : In shallow water barren states look like miniature forms of *P. coriaceus* ; the early spring growth is like that of *P. heterophyllus*, but is more diffuse and less submerged. For a full account, see Fryer, as above.

Distribution in the British Islands.

P. varians has been found only in Cambridgeshire, Hunts and Pembroke, as far as Great Britain is concerned. In Ireland in Waterford and Donegal.

Distribution in other countries.

N. America. Mystic Pond, Mass., United States (Tuckerman).

The specimens figured on Plate 41, POTAMOGETON VARIANS, were doubtless obtained from Witcham Meadlands, Cambridgeshire.

22. **POTAMOGETON FALCATUS**, Fryer, Journal of Botany, March, 1889 (vol. xxvii. p. 66).

P. FALCATUS, Fryer, Journal of Botany, vol. xxvii. p. 66, Tab. 286 (March, 1889), vol. xxviii. p. 210 (1890). *P. gramineus* × *nitens*, Ascherson et Graebner, Synop. Flor. Mitteleurop. p. 326 (1897).

DESCRIPTION: ROOTSTOCK with strong, far-creeping stolons, each terminating in autumn with rather elongated moniliform tubers, rooting with strong fibres at the nodes.

STEM terete, ascending, sparingly branched from the base, with distant, alternate ascending branches, the lower of which are permanently submerged, but not forming a dense mass as in *P. gramineus*.

LEAVES: Lower semi-amplexicaul, flat or a little folded when young, oblong-elliptical, gradually contracted from a little below the middle to the somewhat rounded base, rather suddenly attenuated towards each end, acuminate, entire. Upper similar, amplexicaul or rarely stalked, floating, coriaceous, elliptical.

PETIOLE short, when present.

STIPULES persistent, herbaceous, acute, on the stem long and clasping, afterwards patent, falling short of the internodes on the main stems, often exceeding them on the branchlets. Those at the base of the peduncles very large, stout, cymbiform, sustaining the flower-spikes.

PEDUNCLES slender, equal, or nearly so, falling short of the subtending foliage.

SPIKE: Fruiting spike $\frac{2}{3}$ –1 in. long, cylindrical, dense.

FRUIT: Drupelets small, dark green; inner margin rounded, terminated by the prominent subcentral beak; outer margin almost semicircular, acutely keeled; lateral ridges distinct, distant from the central keel.

COLOUR dark green, or reddish-green when young, drying darker, or blackish-green.

HABIT: In its early state *P. falcatus* has the habit and appearance of the *serratus* state of *P. crispus*; later on, it resembles long-leaved forms of *P. heterophyllus*, or luxuriant plants may be readily mistaken for *P. Zizii*. When gathered, young plants are exactly like certain forms of *P. nitens*.

REMARKS: I cannot place *P. falcatus* under *P. nitens*, which is a barren hybrid, and usually has the leaves longitudinally folded and recurved, with 12 ribs, instead of flat and ascending, with 6 ribs (*see* also Fryer, Journal of Botany, vol. xxvii. p. 66). *P. falcatus* is sufficiently distinguished from *P. heterophyllus* by its amplexicaul leaves; from *P. Zizii* by the same character, and by the opaque ribs of the coriaceous floating leaves.

[To the end of his life Fryer was somewhat uncertain as to the specific validity of this form.]

Distribution in the British Islands.

P. falcatus is recorded from Huntingdonshire, Nottingham and E. Perth.

Plate 40, POTAMOGETON FALCATUS, was drawn from specimens obtained at Stocking Fen, near Ramsay, Huntingdonshire.

23. × **POTAMOGETON INVOLUTUS**, Fryer, Journal of Botany, vol. xxxiv. (1896), p. 1, pls. 353, 354.

DESCRIPTION: ROOTSTOCK with stout far-creeping stolons.

STEM terete, simple below, much branched above, 1–5 ft. long.

LEAVES: Lower semi-amplexicaul, longitudinally involute, broadest at the base, tapering gradually to the apex, many-ribbed, with coarse, often prominent ascending reticulations. Upper coriaceous, lanceolate, elliptical, oval or spatulate, produced singly opposite, or in pairs subtending, the peduncles, abundant on both flowering and barren branches.

PETIOLE flattened, equalling or exceeding the lamina.

STIPULES persistent, involute, herbaceous or horny, blunt.

PEDUNCLES equal, often curved, shorter than the subtending leaves.

SPIKE short, abortive, usually with closed, rarely with open flowers.

FRUIT imperfect, compressed, keeled.

COLOUR: Dark green, the stems often reddish.

HABIT that of *P. Zizii*, or in shallow water of certain forms of *P. nitens*.

REMARKS: This plant is a hybrid of *P. coriaceus* and *P. perfoliatus*, so it cannot be a form or variety of *P. nitens*, but must rank as a hybrid species, like *P. decipiens*, *P. lanceolatus*, &c. In the Journal of Botany (loc. cit.) I named it *Potamogeton nitens forma involuta*.

Distribution in the British Islands.

P. involutus has a very limited distribution, occurring only in Cambridgeshire, Hunts and Northamptonshire.

The specimens figured on Plate 44, POTAMOGETON INVOLUTUS, were from Black-bush Drain, Whittlesey, Cambridgeshire.

24. × **POTAMOGETON NITENS**, Weber, Fl. Hols. Suppl., p. 11 (1787).

P. (PROTEUS?) CURVIFOLIUS. Cham. et Schlecht. in Linnæa, vol. ii. p. 205 (1827). *P. nitens*, Fries, Novit. Flor. Suec. p. 34 (1828). Fieber, Pot. Böhmens, p. 30, Tab. 2, fig. 14 (1838). Koch, Synop. Flor. Germ. ed. 2, p. 778 (1844). Bab. Man. ed. 7,

p. 373 (1874). *P. heterophyllus*, sub-sp. *nitens*, Web., Hooker, Stud. Flor. ed. 3, p. 432 (1884). Ascherson et Graebner, Synop. Flor. Mitteleurop. p. 325 (1897).

[DESCRIPTION (from Castle Gregory specimens): Rootstock creeping, and giving rise to several stems, 12–16 in. long.

STEM round, slender, much branched.

LEAVES: Lower membranous, transparent, crowded on the stem; firm, shining and recurved: the lower lanceolate, semi-amplexicaul, pointed at both ends, about $1\frac{1}{2}$ in. long by $\frac{1}{2}$ in. in width, half folded, so as to appear channelled, and finely undulated at the margins. Upper leaves commonly submerged, broader and blunter at the points, narrowed at the base into a short petiole, often floating and coriaceous.

PETIOLE short or absent.

STIPULES persistent, membranous, about one-third the length of the leaf.

PEDUNCLE short and stout, about the length of the upper leaves, slightly thickened upwards.

SPIKE dense, short.

FRUIT: In the Castle Gregory (Irish) specimens (which were hardly mature) small and semicircular, with a greenish rib-like keel, not prominent when fresh, but distinctly visible when half dried: the younger fruit ovate, with a slight prominence on the back. [We now know (p. 61) that the drupelets are never fertile.]

COLOUR: Dark green.

HABIT that of *P. heterophyllus*, but usually differs in the absence of floating leaves.

Distribution in the British Islands.

P. nitens is recorded from Surrey, Cambs, Hunts, Anglesea, S. Lincoln, S. Lancaster? S. Northumberland, Cheviotland, Roxburgh, Fife, Stirling, Mid and E. Perth, Forfar, N. and S. Aberdeen, Elgin, Easternness, Argyle, Clyde Isles, Cantire, S. and Mid Ebudes, E. Ross, West Sutherland, Caithness, Orkney and Shetland. Also from nineteen divisions of Ireland, but so indefinite a form may have been overlooked or erroneously included in the list.

Distribution in other countries.

Europe: From Iceland and Norway to Italy, and from France to Lithuania (Russia). Asia: Japan! N. America: Canada (Ontario!) United States (Michigan! Connecticut! Florida! Massachusetts!)

Plate 37 represents *P. NITENS*, Plate 38, *P. NITENS LATIFOLIUS*. Mr. Fryer has not noted whence the specimens came, or what status he assigns to “*latifolius*,” but see remarks following.]

REMARKS: *P. curvifolius* is a barren hybrid, and is the *P. nitens* of Weber and subsequent writers. It is no one definite form, but is an aggregate of several distinct hybrids, all of which agree in having *P. perfoliatus* as one parent.

Weber's original description in the Supplement to Flor. Holsat. (1787), pp. 5, 6, is as follows :—

“*Potamogeton nitens*. P. foliis lanceolatis oppositis Web. in fossis prope Nostorf. Folia remota, opposita, longe petiolata, lanceolata, petiolo nonnihil longiora, integerrima, superne nitentia. Pedunculus longus, crassiusculus. Spica multiflora.

“Habitus *P. natantis*, sed planta tenera, multo minor, folia angustiora. Idem tamen videtur *Potamogeton* foliis lanceolatis oblongis, petiolis longis, Gronovii, Flor. Virg. p. 139.

“*P. natans* β , Linn. Spec. Plant. i. p. 182.

“Misit simile specimen Upsaliae lectum Cl. Ehrhard sub homine varietatis *Pot. lucentis*. A *lucente* autem quam maxime differt foliis oppositis, multo brevioribus.”

[It will be noticed that Weber's type evidently had floating leaves, and probably resembled *P. involutus*, which was originally described as *P. nitens*, Weber, *forma involuta*, by Fryer, in the Journal of Botany, vol. xxxiv. p. 1. There is, however, no specimen extant, even at Kiel, named by Weber himself. The nearest is one there named *Potamoget. lucens* var. Ehrh., *P. nitens*, Weber, *P. curvifolius*, Hartman (teste Nolte) Bei Kiel, legit Nolte.]

25. POTAMOGETON LUCENS, L., Sp. pl. ed. 1, p. 126 (1753).

P. PROTEUS, α LUCENS, Chamisso et Schlechtendal in Linnæa, vol. ii. p. 196, Tab. 5, f. 16 a.a. (1827). *P. lucens*, Fieber, Pot. Böhmens, p. 24, Tab. 2, f. 10 (1838). Fries, Novit. Flor. Suec. ed. 2, p. 33 (1828). Koch, Synop. Flor. Germ. ed. 2, p. 778 (1844). Bab. Man. ed. 7, p. 373 (1874). Hooker, Stud. Flor. ed. 3, p. 432 (1884). Morong, Naiad, N. America, p. 30, plate 38 (1893). Ascherson et Graebner, Synop. Flor. Mitteleurop, p. 318 (1897). Fryer, Journal of Botany, vol. xxv. (1887), pp. 50, 309 (aquatic and land forms).

DESCRIPTION: ROOTSTOCK with thick, fleshy, subterranean stolons.

STEM stout, branched from a little below the middle; fruiting branches with many lateral branchlets starting at nearly right angles.

LEAVES all submerged, translucent, undulate and finely serrulate at the margins, elliptical to lanceolate or oblong-ovate to rotundate, subsessile, petiolate; lamina narrowed into the petiole, but decurrent to the stem, not amplexicaul, acuminate, mucronate, or cuspidate; lowest often reduced to a thickened midrib.

PETIOLE truly existent, but obscured by the decurrent lamina.

STIPULES two-keeled, very large, long, obtuse, subscarious to herbaceous.

PEDUNCLE stout, thickened upwards.

SPIKE dense, cylindrical.

FRUIT: Drupelets large, "turgid," obtuse on the back, faintly keeled; lateral ridges obscure; beak short, subcentral.

COLOUR yellowish olive-green, or green, rarely blue-green.

HABIT: The largest British pond-weed. All the branches ultimately float near the surface, but remain submerged; the leaves are very variable in size and shape, and rarely have a hooded tip. Coriaceous floating leaves are rarely, if ever, produced.

Distribution in the British Islands.

P. lucens has a very wide range, from Orkney to the South of England, with the exception of:—Shetland, Hebrides? E. Sutherland, E. and W.? Ross, Mid Ebrudes, S. Ebrudes, Cantire, Clyde Isles, Dumbarton, Elgin, Linlithgow, Haddington, Peebles, Isle of Man, Westmoreland, N. and W. Lancaster, Flint, Merioneth, Montgomery, Cardigan, Pembroke, Radnor, Brecon, Monmouth, Isle of Wight, N. Devon, and Cornwall. It may be expected to occur elsewhere. In Ireland it occurs in twenty-seven districts.

Distribution in other countries.

Europe: Norway to Italy and France to Lithuania (Russia), Faroes. Asia: Siberia to India and Afghanistan to China. Africa: Abyssinia to Cape Colony and Mauritius (sub-sp.) to Senegal. N. America: Canada from lat. 51° to Quebec and Nova Scotia. United States (Michigan and Connecticut to Florida, and New York to California and Mexico). S. America: Argentine Republic. Australasia: Murray River, mutual border of N. S. Wales and Victoria! Tambo River, Victoria! with var. *longifolius* (Gay).

Plate 45 represents *P. LUCENS*; Plate 46 the same in its young autumnal state; Plate 47, *P. LUCENS CORNUTUS*.

26. **POTAMOGETON ZIZII**, Koch (= **ANGUSTIFOLIUS**), Bertch & Presl. Röstlin (1821).

P. PROTEUS, β *ZIZII*, Chamisso et Schlechtendal in Linnæa, vol. ii. p. 201, Tab. 5, f. 16, b. c. (1827). *P. lucens*, β *heterophyllus*, Fries, Novit. Flor. Suec. ed. 2, p. 34 (1828). *P. Zizii*, Mert. et Koch, Fieber Pot. Böhmens, p. 26, Tab. 2, fig. 11 (1838). *P. lucens*, sub-sp. *Zizii*, Roth. Hooker, Stud. Flor. ed. 3, p. 433 (1884). Morong, Naid. N. America, p. 29, plate 37 (1893). *P. lucens* var. *heterophyllus*, Fries, Novit. Flor. Suec. ed. 2, p. 34 (1828). *P. Zizii*, Ascherson et Graebner, Synop. Flor. Mitteleurop. p. 319 (1897). *P. Zizii*, Charles Bailey, Manchester Lit. and Phil. Soc. Proceedings, Dec. 12, 1882. Fryer, Journal of Botany, vol. xxv. p. 113.

DESCRIPTION: ROOTSTOCK producing numerous permanently submerged barren branches.

STEM stout, much branched at the base; fruiting branches ultimately ascending to the surface, with many lateral branches starting at acute angles, and often with terminal coriaceous floating leaves on the flowerless as well as on the flowering shoots.

LEAVES: lower, submerged, alternate, translucent, undulate, and finely denticulate towards the tip, sessile or shortly stalked, often longitudinally folded and recurved, or flat and ascending, narrowed towards the base, and narrow in proportion to their length; elliptical to lanceolate, or oblong-ovate, acuminate or mucronate, lowest sometimes reduced to a midrib. Upper alternate or opposite, more distinctly stalked; uppermost floating, coriaceous or subcoriaceous, flat, entire, or membranous, very long stalked, with translucent ribs.

PETIOLE $\frac{1}{2}$ –3 in., with slightly decurrent lamina, which narrows into it.

STIPULES blunt; lower small, clasping; upper large, free above, herbaceous, rarely subscarios, two-keeled or two-winged on the back, upper often cymbiform.

PEDUNCLES longer than the leaves at their base, often produced in clusters, stout, thickened upwards, 3–5 in. long.

SPIKE dense, cylindrical, $1\frac{1}{2}$ – $2\frac{1}{2}$ in. long.

FRUIT: Drupelets small in proportion to the size of the plant, laterally compressed, with a prominent sometimes winged keel, and prominent lateral ridges; beak short, forming a continuation of the straight upper margin.

COLOUR dark green or green, in deep water sometimes yellowish olive-green.

HABIT: The characteristic long-stalked floating leaves, when present, distinguish the present species from *P. lucens*; but the habit of growth is also different, many barren submerged branches being produced, and many of the lower branches on the fruiting shoots remaining deeply submerged even at the time of flowering. The side branchlets too of the flowering branches usually start at acute angles, instead of at right angles as in *P. lucens*. The general habit of growth is like that of *P. heterophyllus*, but the plant is larger, with thinner, coriaceous leaves more gradually attenuated into the petiole.

REMARKS: I use the name *P. Zizii*, as that of *angustifolius* has, according to the Americans, been previously used in the genus.

Distribution in the British Islands.

Absent from Watson's Peninsula and Channel Provinces, and only occurring in Wales in Anglesea, as far as our present information goes. It is found in E. Kent, Surrey, E. Suffolk, W. Norfolk, Cambridge, Hunts, W. Gloucester, Hereford, Stafford, Salop, Anglesea, S. Lincoln, Leicester, Chester, N.E. and S.E. Yorks, Westmoreland with N. Lancaster, Cumberland, Kirkeudbright, Wigton, Lanark, Selkirk, Roxburgh, Berwick, Linlithgow, Fife, Stirling, E. and W. Perth, Forfar, Dumbarton, North Eubudes, E. Ross. Ireland in twenty-two divisions, commonest in the west.

Distribution in other countries.

Europe: Norway to Montenegro, Italy and France to West Russia. Asia: Turkestan! Tibet! China! Himalayas! Kashmir to Upper Burma! Africa: Madagascar! America: Canada from 57° N. lat. to Quebec, and Ontario to British Columbia. Trinidad and Cuba. United States (New England to Florida, and Vermont to California and New Mexico).

Plate 42, POTAMOGETON ZIZII, no doubt represents what Fryer considered the typical form of the Cambridgeshire and Huntingdonshire Fens.

27. **POTAMOGETON CORIACEUS**, Fryer, Journal of Botany, 1889, p. 8.

P. CORIACEUS, Nolte, Bennett in Journal of Botany, vol. xxiv. p. 223. *P. Zizii* *Il. coriaceus*, Ascherson et Graebner, Synop. Flor. Mitteleurop., p. 320 (1897). Fryer, Journal of Botany, vol. xxvii. p. 8 (1889) [full description].

DESCRIPTION: ROOTSTOCK tuberous.

STEM 1-4 ft. long, stout, round, much branched from the base, or simple below in strong shoots produced late in the season. Lower branches permanently submerged; upper branches ultimately ascending to the surface and spreading into numerous branchlets, each of which is terminated by coriaceous floating leaves, often forming a rosette.

LEAVES: Those at the base of the stem often reduced to phyllodes, ordinary submerged leaves stalked or sessile, membranous, with numerous longitudinal ribs connected by transverse veins, entire, without spinous denticulations, usually flat or very slightly undulated, oblong or obovate, mucronate. Upper leaves coriaceous, all stalked, lamina always greatly exceeding the petiole, with 8-10 lateral translucent ribs on either side; obovate or roundish or clavate, rarely elliptical.

PETIOLE shorter than the lamina.

STIPULES large, herbaceous, blunt, slightly winged on the back.

PEDUNCLE very stout, swollen upwards, slightly curved at the time of flowering, usually straight and erect in fruit.

SPIKE stout, cylindrical, dense, 1½ in. long, usually terminal.

FRUIT: Drupelets rather small in proportion to the size of the plant; inner margin nearly straight, terminated by the short beak; outer margin semicircular, with an acute almost winged keel, projecting at right angles from the base of the fruit.

HABIT: That of *P. Zizii*; but the plant has densely coriaceous floating leaves, like those of *P. heterophyllus*, very rarely elliptical, but inclining to orbicular. The lamina greatly exceeds the petiole, thereby differing from that of *P. heterophyllus*.

COLOUR bright green or brownish-green, usually drying darker.

REMARKS : The typical or central form of the *lucens* group aggregated by Chamisso and Schlechtendal into one species, *P. proteus*, will probably be found in *P. coriaceus* (Journal of Botany, vol. xxv. p. 52).

Distribution in the British Islands.

Cambridgeshire.

Distribution in other countries.

Sweden, France, Switzerland, Schleswig-Holstein.

Plate 43, POTAMOGETON CORIACEUS, is drawn from specimens obtained at or near Witeham Meadlands, Mepal, Cambridgeshire.

28. × POTAMOGETON DECIPIENS, Nolte in Koch, Syn. p. 779 (1843-1844).

P. DECIPIENS, Koch, Synop. Flor. Germ. ed. 2, p. 779 (1844) (Nolte in litt. ad amicos). Bab. Man. ed. 7, p. 373 (1874). *P. lucens*, sub-sp. *decipiens* Nolte, Hooker, Stud. Flor. ed. 3, p. 433 (1884). *P. decipiens*, Fryer, Journal of Botany, 1890, p. 137 [full description and notes]. *P. perfoliatus* × *lucens*, Ascherson et Graebner, Synop. Flor. Mitteleurop. p. 329 (1897).

DESCRIPTION : ROOTSTOCK with far-spreading stolons, striking deeply into the soil.

STEM stout, round, simple below, sparingly branched above the middle ; all the branches spreading ultimately at the surface of the water, with widely diverging branchlets.

LEAVES all submerged, semiamplexicaul, or upper sessile ; alternate, opposite at base of the peduncles only ; lowest leaf very rarely reduced to a short strap-shaped phyllode which is rounded at the tip and mucronate, or with a narrow lamina attenuated towards each end. Ordinary leaves all similar, oblong, strap-shaped, rounded and mucronate at the apex, or narrowed and apiculate, upper rarely orbicular, flat or involute at the base, with three principal ribs on each side of the midrib, and fainter intermediate ribs.

PETIOLE very short or absent.

STIPULES blunt, stout, long ; those on the upper branchlets longer than the internodes ; herbaceous, persistent ; those at the base of the stem sometimes dilated towards the tip into a leaf-like lobe, those at the base of the branches often bearing a conspicuous sessile leaf, 1-2 in. long.

PEDUNCLE not swollen upwards, often scarcely thicker than the stem, 2 to 3 times longer than the flower-spike, subtended by two opposite leaves, terminal becoming lateral by the growth of the branch.

SPIKE 1 in. long, barren.

FRUIT : Drupelets resembling those of *P. perfoliatus*, but abortive.

COLOUR bright green or brownish-green, drying darker.

HABIT that of *P. lucens*, but more apt to form large thick beds, by the extension of the strong-growing rootstock.

Distribution in the British Islands.

P. decipiens has a somewhat curious distribution, though it must be remembered that it is a form often confounded with its nearest allies. It occurs over a broad district in the South of England (Somerset, Dorset, S. Hants, W. Sussex), thence it ranges northwards to Surrey, Oxfordshire and Warwickshire, and circles round by Northants, Leicester, Hunts and Cambs to S. Lincoln. On the west it extends to Hereford. In Scotland it has been met with in Roxburgh, Berwick, Stirling, Mid Perth, Forfar and S. Eubodes. In Ireland it occurs in four divisions.

Distribution in other countries.

Europe: Sweden to Tyrol, and France to Lithuania (Russia). Asia: N. India.

Plate 48, POTAMOGETON DECIPIENS, was no doubt drawn from Cambridgeshire specimens obtained at Welches Dam or Mepal.

29. × POTAMOGETON SALIGNUS, Fryer MS.

POTAMOGETON SALICIFOLIUS, Wolfgang, in Roem. et Schultes, Mant. iii. p. 356 (1827). *P. decipiens* var. *salicifolius*, Ar. Bennett.

Many different opinions are held with regard to this form by such eminent authorities on the group as Fryer, Arthur Bennett, and Tiselius—not to mention Hägstrom, Baagoe, Almquist, Babington and others. We can, however, only give here that of Fryer, which is taken from his MS. notes:—"I now make it a hybrid species; it is so very distinct from any other *decipiens* form that I think it deserves segregation (although I first suggested its *decipiens* affinity to my friend Bennett). It will be found in the Potamogeton book under the name of *P. salignus*. The same varietal or specific name ought never to be used twice in the same genus."

[DESCRIPTION: The original description of *P. salicifolius*, Wolfg., is as follows:—"Caule tereti ramoso, foliis omnibus submersis, membranaceis, alternis, oblongo-lanceolatis, obtusicaulis, sessilibus, undulatis trinerviis, sub-denticulatis; stipulis angustis, patentibus, brevibus; pedunculis alaribus, elongatis aequalibus, crassicaulis, folia sub-aequantibus. Wolf. No. 13, Besser in litt. Folia, 3-7 poll., 5-10 lin. lata. In fluvis Wilna non procul a Wilna. Lithuania."

Distribution in the British Islands.

Herefordshire.

From the above note it is evident that Mr. Fryer eventually considered our plant the same as Wolfgang's; but this he formerly doubted. Ascherson and Graebner, Syn. Fl. Mitt. Europ. p. 328 (1897) refer it to *P. alpinus* × *lucens*.—A. Bennett.]

DISTRIBUTION in other countries.

Brandenburg, Silesia, Lauenberg, Lithuania. It is given for Norway by Hartman, and for Sweden and Lapland by Nyman.

[Addendum, Oct., 1914.—I take the opportunity of observing that the justification quoted by Mr. Evans from Mr. Fryer's MS. notes on the previous page for assigning the name *salignis* to this plant, namely, that "the same varietal or specific name ought never to be used twice in the same genus," is in my view insufficient. It would entail so many other alterations in the nomenclature of the genus.

On the question of the position of this *Potamogeton* the Journal of Botany, 1908, p. 251 should be consulted.—A. BENNETT.]

EXPLANATION OF PLATE 49. × *POTAMOGETON SALIGNUS*. Drawn from Herefordshire specimens. With flower spikes and section of stem.

GRASS-LEAVED *POTAMOGETONS*.

By ARTHUR BENNETT, A.L.S.

The following eight species belong to the grass-leaved section of the genus, the *Chleophylli*, Koch, "Whole plant submersed, leaves grass-like, exactly linear, sessile and alternate, stipules free."

The growth in this section is somewhat different to that of those previously described; many of the species produce gemmae, or winter-buds in profusion, others more sparingly; many produce glands at the base of the leaves, which in some are dull, in others shining.

30. **POTAMOGETON ZOSTERAEFOLIUS**, Schum. Enum. Pl. Saell. ed. 1, p. 50 (1801).

P. COMPRESSUS (*ZOSTERAEFOLIUS*), Fries, Novit. Flor. Suec. ed. 2, p. 44 (1828).

Koch, Synop. Flor. Germ. ed. 2, p. 779 (1844). Reichenbach, Icon. Flor. Germ. et Helv. vol. vii. tab. 27 (1845). Fieber, Pot. Böhm. p. 33, tab. 3, fig. 16 (1838). Hooker, Stud. Flor. ed. 3, p. 434 (1884). Syme, Eng. Bot. ed. 3, vol. ix. p. 45, tab. 1415 (1869). Morong, Naid. N. America, p. 37, tab. 45 (1893). Babington,

* Page 76 ended the section edited after the author's death by Mr. A. H. Evans and issued in 1913. Page 77 commences the section for which Mr. Arthur Bennett is solely responsible. See Introduction, page vii.

Man. Brit. Bot. ed. 9, p. 441 (1904). (*P. compressus*) Linn. Sp. Pl. ed. 1, p. 127 (1753). Ascherson et Graebner, Synop. Flor. Mitteleurop. ed. 2, p. 518 (1913). Rannkiær Dansk. Blomst. p. 84, fig. 48, p. 86, fig. 48, A. B. (1896).

DESCRIPTION. Rootstock sometimes persistent, but often the plants are produced by the growth of the winter-buds. The plant, as it grows, gradually produces longer leaves, and branches of these may remain without flowering, while in the next species this is rare.

STEM much flattened, broad, $\frac{1}{8}$ – $\frac{1}{4}$ inch wide. When starting from a winter-bud the first leaves are very short, very obtuse, and with hardly any definite veining.

LEAVES all similar, sessile, linear, usually rounded and cuspidate at the apex, but sometimes acuminate, with 5 principal ribs (rarely 3), and many secondary ones; very variable in length.

STIPULES long, in the lower leaves rounded and cuspidate, in the upper acute with many semi-persistent fibres.

PEDUNCULES usually terminal between the forks of the stem, stout, twice to four times as long as the spikes.

SPIKES oblong, dense-flowered, $\frac{3}{4}$ – $1\frac{1}{2}$ inches long.

SEPALs orbicular-rotund, 1-veined.

FRUIT. Drupelets large, slightly compressed, subreniform, ventrally rather convex, dorsally rounded, bluntly 3-keeled, sometimes with slight bosses, terminated by a central beak.

DISTRIBUTION in the British Islands.

It is recorded in 24 of the Watsonian vice-counties; ranging from Surrey north to Forfar; and from Norfolk to Cheshire and Salop; but it is not on record for Wales or Ireland.

DISTRIBUTION in other countries.

Europe: From Norway and Finland south to Croatia; and from France eastwards to Transylvania; generally distributed in 32 governments of Russia from Archangel to the Caucasus. Asia: Siberia and Japan. N. America: Canada from 57° N. Lat. to Ontario, Brit. Columbia; and in the United States from Maine to Pennsylvania, across to Oregon and California.

The Scotch specimens are generally longer-leaved than the English, and more acute.

Our own specimens are fairly alike in the fruits, but in N. America they vary greatly, and are not represented by the Icones figures. They probably come under the var. *selenocarpus*. Fischer in Ber. Bay. Bot. Ges. vol. xi. 1907.

Whether or not this is the *compressus* of the Species Plantarum, ed. 1, is not a question to be discussed here; at present it seems better to use the nomenclature usually adopted in our books.

EXPLANATION OF PLATE 50. *POTAMOGETON ZOSTERAEOFOLIUS*. Drawn from specimens from Rockland Broad, E. Norfolk. With parts of leaves and section of stem, mag.

31. **POTAMOGETON ACUTIFOLIUS**, Link, in Roem. et Schultz. Sys. vol. iii. p. 513 (1818).

P. ACUTIFOLIUS, Cham. et Schlecht. in Linnæa, vol. ii. p. 180, tab. 4, fig. 9 (1827). Fries, Novit. Flor. Succ. ed. 2, p. 46 (1828). Koch, Synop. Flor. Germ. et Helv. ed. 2, vol. ii. p. 780 (1844). Reichenbach, Icon. Flor. Germ. vol. vii. p. 17, tab. 26 (1845). Fieber, Pot. Böhm. p. 34, tab. 3, fig. 17 (1838). Hooker, Stud. Flor. ed. 3, p. 435 (1884). Syme, Eng. Bot. ed. 3, vol. ix. p. 46, tab. 1416 (1869). Babington, Man. Brit. Bot. ed. 9, p. 519 (1904). Ascherson et Graebner, Synop. Flor. Mitteleurop. ed. 2, p. 441 (1913).

P. cuspidatus, Schrad. ap. Smith, Eng. Flor. ed. 2, p. 235 (1828) et Herbarium.

DESCRIPTION. ROOTSTOCK sometimes persistent, and at others from a winter-bud, but less frequently than in the last species, and the leaves less obtuse from these buds.

STEM flattened to $\frac{1}{8}$ inch, but often less, sometimes tapering in the internodes.

LEAVES all similar, linear 2–5 inches long by 2 lines wide, acute-acuminate with 3 principal ribs, and many (18) secondary ones, or at times the midrib is composed of many (6) compacted ribs and translucent.

STIPULES 1 inch long, brownish-white, acute, persistent with many fine ribs.

PEDUNCULES generally short, $\frac{1}{2}$ inch long, rarely $\frac{3}{4}$ –1 inch, equal.

SEPALs rhomboidal, 1-nerved.

FRUIT. Drupelets large, 2 lines by 1 line, greenish-olive, compressed, slightly compressed on the ventral margin, with a recurved beak forming a continuation and a tooth near the base, half-obovate on the dorsal margin, bluntly 3-keeled and often undulated.

β major, Fieber, l.c.

Leaves up to 6 inches, wider, often $\frac{3}{16}$ inch wide, stem more flattened.

Buckenham Ferry, E. Norfolk.

DISTRIBUTION in the British Islands.

Dorset, Sussex, Kent, Surrey, Herts, Middlesex, Norfolk, Northampton, Warwick, Lincoln, S.E. York.

DISTRIBUTION in other countries.

Europe : Sweden to Italy, and France to Lithuania, but rarer than the last species. Asia : Manchuria (a sub-sp.). Australia : Victoria River, N. Australia.

The last habitat, the only one known south of the Equator, cannot be held to be an introduction; at the same time it is a very remarkable one. Others will perhaps

eventually be found. The specimens in the Kew Herbarium from Australia were named "*P. zosterifolius*, Schum." by the late Baron von Mueller. *P. cuspidatus*, Schrader ex Smith, is generally referred to *zosteracfolius*, but the specimens so named in Smith's herbarium are *acutifolius*, a smaller, more slender plant than *zosteracfolius*, and with very different fruit; it is often in this condition from June to August, while *zosteracfolius* is usually in fruit the end of July, with numerous barren branches.

EXPLANATION OF PLATE 51. POTAMOGETON ACUTIFOLIUS. Drawn from specimens from Buckenham Ferry, E. Norfolk. With fruit spike and part of leaf, mag.

32. **POTAMOGETON OBTUSIFOLIUS**, Mert. et Koch, Dent. Fl. ed. 3, vol. i. p. 855 (1823).

P. OBTUSIFOLIUS, Chamisso et Schlechtendal in Linnæa, vol. ii. p. 178, tab. 4, fig. 7 (1827). Fries, Novit. Flor. Suec. ed. 2, p. 47 (1828). Koch, Synop. Flor. Germ. et Helv. ed. 2, vol. ii. p. 780 (1844). Reichenbach, Icon. Flor. Germ. vol. vii. p. 16, tab. 25 (1845). Fieber, Pot. Böhm. p. 37, tab. 3, fig. 19 (1838). Syme, Eng. Bot. ed. 3, vol. ix. p. 47, tab. 1417 (1869). Hooker, Stud. Flor. ed. 3, p. 435 (1884). Babington, Man. Brit. Bot. ed. 9, p. 442 (1904). Morong, Naid. N. America, p. 40, tab. 48 (1893). Ascherson et Graebner, Synop. Flor. Mitteleurop. ed. 2, p. 521 (1913). *P. gramineus* in Smith, Eng. Fl. ed. 2, vol. i. p. 235 (1828). *P. compressus*, Willd. Berol. Mag. (1809).

DESCRIPTION. ROOTSTOCK growing from winter-buds, or from creeping rhizomes, the young plants smaller, and more delicate than the last two species.

STEM. Four-sided, slightly compressed, repeatedly branched in the ordinary form, less so in the deep-water forms.

LEAVES all similar, linear, sessile narrowed to the base, obtuse, subapiculate, translucent with 3 (rarely 5) ribs and no secondary ribs as in the last two species. The young leaves, especially when produced from a winter-bud, are often without any definite ribs, the whole leaf being made up of translucent chain-like areolation. The leaves often have glands at their base, which are large and translucent.

STIPULES small and subacute, 6-9 lines long.

PEDUNCULES rather slender, equal, $\frac{1}{2}$ to $1\frac{1}{2}$ inches in the deep-water forms.

SPIKES 3-4 lines long, oblong-ovoid, dense-flowered.

SEPALs rhomboidal, or rhomboidal elongate, 1-nerved.

FRUIT. Drupelets obliquely obovate, slightly compressed, convex on the ventral margin, semicircular on the dorsal, 3-keeled, terminated by a short beak, the lower part of the margin sometimes slightly tuberculated.

β fluvialis, Lange et Mortensen, Bot. Tidssk. vol. x. p. 200 (1878).

Stems elongated to 2-4 feet, sparingly branched, internodes elongated, peduncules 1-1½ inches long, spikes ½ inch.

In deep ponds and lakes.

The species is generally with 3-ribbed leaves, but sometimes it occurs with 5, and Fieber figures a form with 7; this I have only seen in Canadian specimens from the far North.

DISTRIBUTION in the British Islands.

From Hants and Kent north to Inverness; in Watsonian counties. Absent from the south-west of England. In Ireland in twenty of Mr. Praeger's divisions.

DISTRIBUTION in other countries.

Europe: From Norway to Italy (Piedmont) and France to the Banat and Lithuania. Asia: Siberia, Mongolia and Persia. N. America: Greenland to 68° 38' N. Lat., Canada. United States: Maine to N. York, and Pennsylvania to Minnesota. S. America: "Ecuador," Graebner in Das Pflanzenreich.

EXPLANATION OF PLATE 52. POTAMOGETON OBTUSIFOLIUS. Drawn from specimens from Estwaite Water, Lake Lancashire. Figs. 2, flower spike; 3, end of leaf; 4 and 5, flowers; 6, ovary; 7, fruit, all mag.

33. **POTAMOGETON FRIESII**, Ruprecht, Beitz. Pfl. d. Russ. Reiches, vol. iv. p. 43 (1845).

P. compressus, Smith, Eng. Fl. vol. i. p. 234 (1824). L. Reich. Icon. Fl. Germ. vol. vii. p. 15, tab. 24 (1845). Fieber, Die Pot. Böhm, p. 36, tab. 3, fig. 18 (1838). *P. pusillus*, var. *major*, Fries, Novit. Flor. Succ. ed. 2, p. 48 (1820), var. *latifolius*, Meyer, Chl. Hann. p. 525 (1836), sub-sp. *Friesii*, Hooker, Stud. Flor. ed. 3, p. 438 (1884). *P. major* (Fr.) Morong, Naiad. N. America, p. 41, tab. 49 (1893). *P. Friesii*, Rupr. Bab. Man. Brit. Bot. ed. 9, p. 442 (1904). *P. mucronatus*, Schrad. Ascherson et Graebner Synop. Flor. Mitteleurop. ed. 2, p. 523 (1893). Syme, Eng. Bot. ed. 3, vol. ix. p. 48, tab. 1418 (1869).

DESCRIPTION. As with the last three species sometimes proceeding from winter-buds, or creeping rhizomes.

STEM semi-compressed, slender, nearly simple, but generally heavy in the axils of the upper leaves, fascicles of small leaves.

LEAVES all similar, linear, sessile, 1½-2½ inches long and 1 line wide, subacute, or obtuse and cuspidate, usually 5, but rarely 3-7-ribbed, biglandular at the base, glands small and generally dull.

PEDUNCULES slightly (or sometimes more so) thickened upwards, 1-1½ inches long.

SPIKES interrupted, few-flowered.

SEPAIS orbicular, 1-veined.

FRUIT. Drupelets small, olive, slightly compressed, convex on the ventral margin, rounded-oval on the dorsal, bluntly 3-keeled with the beak short, central and recurved.

COLOR. Plant dull green, generally keeping its colour when dried.

The least understood of the grass-leaved section, most of the others being so named in different European Herbaria, even as *P. pectinatus*. With us, even as late as in the third edition of the English Botany (1869) Dr. Syme only recorded two stations for this plant; now we have it on record for 48 of the Watsonian counties, and no doubt it is still passed over in others. That it is not always easy to separate it from *P. obtusifolius* is shown by the fact that some of our own botanists and many others in Europe and America still cross name them.

DISTRIBUTION in British Islands.

From Somerset and Dorset, north to Caithness and Orkney; and Norfolk to Cheshire. In Ireland: Limerick, Galway, Westmeath, Mayo, Monaghan.

DISTRIBUTION in other countries.

Europe: Finland, 66° 59' N. Lat. The Faroes. Sweden to Montenegro, and France to Lithuania. Russia: Archangel and Ingria to S.W. Russia. Asia: Siberia and Manchuria. N. America: Canada, Ontario to Alberta, and Assiniboia. U.S.A.: Maine to N. England States, Iowa, Louisiana. S. America: Argentina teste Dr. Stuckert.

The African plant is now referred to *P. pusillus*, L., var. *australis*, Hägstrom.

EXPLANATION OF PLATE 53. POTAMOGETON FRIESII. Drawn from specimens from Buckenham Ferry, E. Norfolk. With fruit spike and part of leaf, mag.

34. **POTAMOGETON RUTILUS**, Wolfgang in Roem. et Schut. Sys. Veg. Mant. vol. iii. p. 362 (1827).

P. CAEPITOSUS, Nolte in Reich. Leon. vol. vii. tab. 23, fig. 40, p. 21 (1845).

P. pusillus, L., var. β , Chamisso et Schlechtendal in Linnæa, vol. ii. p. 172 (1827). Tiselius, Pot. Suec. Exsic. vol. iii. No. 105 (1897).

DESCRIPTION. STEM subcompressed, lower part branched, upper simple, slender.

LEAVES all similar, narrow-linear, lower crowded, 1-3 inches, 3-ribbed without secondary ribs, semirigid, gradually acuminate at the apex, often tinged with red at the base.

STIPULES long 1 inch, subpersistent, acuminate.

PEDUNCLES $1\frac{1}{2}$ -2 inches long, slightly thickened upwards.

SPIKES 3 lines long, few-flowered.

SEPAIS lance-oblong.

FRUIT. Drupelets $\frac{1}{2}$ –1 line long, nearly straight on the ventral margin, the beak forming a continuation, semi-convex on the dorsal margin. Plant elongated with long internodes in the upper part, and sometimes with fascicles of leaves on the axils.

DISTRIBUTION in the British Islands.

England : Anglesea. Scotland : Orkney ?

DISTRIBUTION in other countries.

Europe : Sweden to Hungary, and France to Lithuania, Finland, 62° N. Lat. ; but rare everywhere. Russia : Onegaland to Poland. North America : Canada, Anticosti, Lake Superior. U.S. : N. York, Michigan, Minnesota.

EXPLANATION OF PLATE 54. POTAMOGETON RUTILUS. Drawn from Swedish specimens. Figs. 2, part of stem leaf and stipules ; 3, end of leaf ; 4, flower ; 5, fruit, all mag.

35. **POTAMOGETON PUSILLUS**, Linn. Sp. Pl. ed. 1, p. 127 (1753).

P. PUSILLUS, Chamisso et Schlechtendal in Linnæa, vol. ii. p. 170, tab. 4, fig. 5 (1827). Fries, Novit. Flor. Suec. ed. 2, p. 48 (1828). Fieber, Pot. Böhm. p. 39, tab. 4, fig. 20 (1838). Koch, Synop. Flor. Germ. et Helv. ed. 2, vol. ii. p. 780 (1844). Syme, Eng. Bot. ed. 3, vol. ix. p. 49, tab. 1419 (1869). Hooker, Stud. Flor. ed. 3, p. 435 (1884). Morong, Naid. N. America, p. 45, tab. 53 (1893). Babington, Man. Brit. Bot. ed. 9, p. 442 (1904). Ascherson u. Graebner Synop. Fl. Mitteleurop. ed. 2, p. 525 (1913).

DESCRIPTION. Stems rising from a creeping rhizome as well as (rarely) from winter-buds ; many of these are either carried away, or die naturally.

STEMS subterete, not compressed, sometimes much and densely branched, at times nearly simple, but variable, from 6 to 24 inches.

LEAVES all similar, linear, obtuse or mucronate or acute, 1–3-ribbed, generally biglandular at the base, 1–3 inches long by $\frac{1}{2}$ – $\frac{3}{4}$ line broad, often reticulate between the midrib and other ribs, at times the ribs are nearly obsolete, and the leaf is made up of chain-like areolation only.

STIPULES small, hyaline, obtuse and cymbiform when enclosing the young spikes.

PEDUNCULES slender, equal, 3–9 lines long, but very variable, and often 1–4 inches, rarely thickened in fruit.

SPIKE not (or rarely) interrupted, 3–13-flowered.

SEPALS orbicular, with more than one vein.

FRUIT. Drupelets, small obliquely elliptical, scarcely compressed, acuminate, convex on the ventral margin, semi-obovate on the dorsal, bluntly 3-keeled, with a short central beak, very rarely the base of the keels are slightly tuberculated.

a vulgaris, Fr. Nov. Fl. Succ. ed. 2, p. 48 (1828). Stem sparingly branched, leaves acute or subacute, 3-ribbed, drupelets smooth.

β Bertcholdii, Ascherson, Fl. Brand. p. 664 (1864). Stem branched, leaves broad, 3-ribbed, the secondary ones near the leaf margins, central with reticulate-areolation on either side. Drupelets broad based, the dorsal margin slightly tuberculated. Lake Lancashire; Kent, Norfolk.

γ rigidus, Ar. Benn. Scot. Nat. p. 24 (1883). Stem and leaves rigid, leaves setaceous, subacute, 1-ribbed, involute $1\frac{1}{2}$ –2 inches long. Habit of *P. rutilus*, Wolfg., but leaves shorter, less acute. Orkney, Shetland.

δ tenuissimus, Mert. et Koch, Deut. Fl. ed. 3, p. 1, 857 (1823). Stem filiform, much-branched, leaves 1 ribbed, linear-subulate, acute.

ε similis, Ar. Benn. *P. pusillus*, var. *acuminatus*, Ar. Benn. Jour. Bot. 173 (1907) non Fieber. Stem semi-compressed, leaves linear, long, $2-3\frac{1}{2}$ inches by 2 lines wide. Like *P. acutifolius*, Link. Stirling. Dr. Kidston.

DISTRIBUTION in the British Islands.

Generally distributed, wanting only in three counties.

DISTRIBUTION in other countries.

Europe: Iceland and Norway to Italy and Corsica; France to the Banat and E. Russia; Madeira. Asia: Siberia, 70° N. Lat. to India; Arabia and China. Africa: Algeria to Cape Colony, Egypt, Azores. N. America: Greenland, 69° N. Lat. to Quebec, N. Scotia to B. Columbia. U.S.A.: Maine to Louisiana; N. Jersey to Arizona and California. S. America: Chili (Kunth), Brazils (Schuman) to Argentina (Stuckhart).

EXPLANATION OF PLATE 55. POTAMOGETON PUSILLUS. Drawn from specimens from Loch Marlee, Perth. Figs. 2, part of stem with leaf and stipule; 3, flower; 4, fruit; all mag.

36. POTAMOGETON STURROCKII, Ar. Benn. Scot. Nat. 28 (1883).

P. pusillus, L. subsp. *STURROCKII*, Hooker, Stud. Fl. ed. 3, p. 435 (1884). Ascherson u. Graebner, Synop. Flor. Mitteleurop. ed. 2, p. 530 (1913). Morong, Naiad. N. America, p. 46 (1893).

DESCRIPTION. STEM filiform, branching from the base.

LEAVES all similar, mostly patent in the living plant, narrow-linear, 1–3 inches long by 1 line wide, bright green, translucent, with 3–5 ribs, without any secondary ones.

PEDUNCULES very slender, $1\frac{1}{2}$ –2 inches long, equal.

SPIKE 1 line long, few but dense-flowered. Drupelets smaller than in *pusillus*, convex on both margins, with a short beak.

Whole plant delicate, pellucid, and not conforming to any named *pusillus*, but standing apart.

DISTRIBUTION in Great Britain.

Anglesea, Perth, Stirling, Caithness, Inner Hebrides.

DISTRIBUTION in other countries.

N. America: Sable Island, Canada, U. States. "Novae Angliae," Robbins.

Among the very large number of *pusillus* forms from all parts of the world that have passed through my hands in various European herbaria (as well as our own) there are only two gatherings in America that agree with this plant, and neither Dr. Graebner nor I have seen any European specimens; in the second edition of the Synopsis Florae Mitteleuropeas (1913) he can only give "Schottland" as a locality for it.

P. Sturrockii.

Fig. 1.—1, flower spike; 2, flower; 3, segment of perianth; 4, fruit.

37. **POTAMOGETON TRICHOIDES**, Chamisso et Schlechtendal in Linnæa, vol. ii. p. 175, tab. 4, fig. 6 (1827).

P. TUBERCULATUS, Tenore, ap. Gussone fl. Neap. syll. app. v. p. 4 (1842). *P. monogynus*, Gay, ap. Coss. et Germ. Fl. Paris, supp. p. 89 (1843). *P. Phialae*, Post, Bull. Herb. Boiss. vol. i. p. 409 (1893). Reichenbach, Icon. Flor. Germ. et Helv. vol. vii. tab. 21, fig. 34 (1845). Babington, Bot. Gaz. vol. ii. p. 285 (1850). Caspary, Jour. Linn. Soc. vol. viii. p. 273 (1865). Syme, Eng. Bot. ed. 3, vol. ix. p. 51, tab. 1420 (1869). Ascherson u. Graebner, Synop. Flor. Mitteleurop. ed. 2, p. 532 (1913).

DESCRIPTION. STEMS semiterete, filiform, repeatedly branched.

LEAVES all similar, sessile, setaceous, acute and often curled at the apex; 1, more rarely 3-ribbed.

STIPULES small; subacute, or acute, subscarious.

PEDUNCULES long 2-4½ inches long, slender, equal.

SPIKE oblong-ovoid, or subglobose, few-flowered.

SEFALS suborbicular.

FRUIT. Drupelets small, varying from the ventral margin with a small projection towards the base, with a short straight beak, and the dorsal margin semicircular,

bluntly heeled ; or the ventral margin with a strong tooth, and with the dorsal margin with the heels more or less tuberculated.

COLOUR. Plant deep dull green, drying much darker.

β Trimmeri, Caspary, l.c.

LEAVES 3-ribbed, front more or less tuberculated on the dorsal heels.

DISTRIBUTION in the British Islands.

England : Devon, Sussex, Surrey, Suffolk, Norfolk, Cambridge. Scotland : Perth.

DISTRIBUTION in other countries.

Europe : Sweden to Sicily, and France to Lithuania ; Russia, Ingria, Livonia to Kursk. Asia : Anti-Libanus. Africa : Algeria, Port Natal, Teneriffe (?).

The original specimens in the Berlin Herbarium (and the original description and figure agree) show none of the tuberculation on the back of the fruit, only a projecting tooth on the outer edge, and the leaves are 1-ribbed. For this reason Caspary named the British (Norfolk) specimens var. "*Trimmeri*," after the finder, the Rev. K. Trimmer, author of a Flora of Norfolk. But in the French, Bohemian and probably other plants the ribs vary from 1 to 3, and an extreme sub-species, *P. condylocarpus*, is described by Tausch in the Bot. Zeit. vol. ii. p. 413 (1836), where the tuberculation becomes extreme, and the shape differs. Schumann, in Flora Brasiliensis, iii. 2, 710 (1894), points out that the variation is so gradual that it seems hardly needful to divide the species into varieties. Cosson et Germain, Fl. Paris, 572 (1845), give "3-5 nervures;" this I have not seen, and there are only three in Gay's own specimens at Kew.

EXPLANATION OF PLATE 56. POTAMOGETON TRICHOIDES. Drawn from Cambridge-shire specimens. With parts of leaves, flower spike, and section of stem, all mag.

COLEOPHYLLI, KOCH.

The four following species belong to this section, whole plant submersed, leaves grass-like exactly linear, base vaginate, stipules adnate.

38. POTAMOGETON PECTINATUS, Linn. Sp. Pl. ed. i. p. 127 (1753).

P. PECTINATUS, L. Chamisso et Schlechtendal in Linnæa, vol. ii. p. 164, tab. 4, fig. 2 (1827). Fries, Novit. Flor. Suec. ed. 2, p. 52 (1828). Fieber, Pot. Böhm, p. 43, tab. 4, 24 (1838). Koch, Synop. Flor. Germ. et Helv. ed. 2, vol. ii. p. 781 (1844). Hooker, Stud. Flor. ed. 3, p. 436 (1884). Morong, Naid. N. America, p. 51, tab. 58 (1893). Ascherson u. Graebner, Synop. Flor. Mitteleurop. ed. 2, p. 538 (1913). *P. pectinatus*, L., sub-sp. *eupectinatus*, Syme, Eng. Bot. ed. 3, vol. x. p. 53, tab. 1422 (1869).

DESCRIPTION. Plant from a long creeping rhizome, often with many tuberous expansions.

STEM cylindrical or subcompressed, slender, naked below, and branching above (or branching from a running rhizome), often in a semilateral manner.

LEAVES. Early leaves flat, linear usually 3-ribbed, upper leaves setaceous, 1-ribbed, of two interrupted tubes, many with one or more branches in the axils.

STIPULES rather long and narrow, united to the sheathing petiole of the leaf, with only the apex free.

PEDUNCULES equal, varying in length, 3-6 inches.

SPIKES 1-2 inches long, few-flowered, interrupted, the lowest distant, the upper subapproximate.

SEPALs suborbicular-reniform.

FRUIT. Drupelets large, 2 lines by 1 line, olive-green, tinged with olive-brown, straight or slightly convex on the ventral margin, with a very short beak forming a continuation of the ventral margin. Dorsal margin without ridges, and rounded when fresh; nut with strong lateral ridges.

DISTRIBUTION in the British Islands.

From Cornwall and Kent north to Sutherland and Shetland; in about eighty-eight counties. Ireland: Kerry north to Down and Antrim in about twenty-five counties.

DISTRIBUTION in other countries.

Europe: Iceland, Norway at 70° 25' N. Lat. to Sicily; and France to Bulgaria. Asia: Siberia to China, India, Turkey-in-Asia to Corea. Africa: Algeria to Natal and Transvaal, Madagascar. North America: Greenland, Canada to B. Columbia, U.S. Maine to Florida, and N. York to California. South America: Brazil, Peru, Chili, Patagonia. Australia: Victoria, S. Australia, New Zealand, New Caledonia.

EXPLANATION OF PLATE 57. *POTAMOGETON PECTINATUS*. Drawn from specimens from Lake Lancashire. Figs. 1, plant; 2, sheath; 3, flower; 4, fruit; 5, longitudinal section of fruit, all mag.

39. **POTAMOGETON VAGINATUS**, Turcz. Moscow. Soc. Nat. Bull. vol. xxvii. p. 66 (1856).

P. JUNCTIFOLIUS, Kerner 2, *vaginatus* Ascherson u. Graebner, ed. 2, p. 544 (1913). *P. pectinatus*, L., var. *fluvialilis*, Schüb. et Mart. Flora Wurtemberg, vol. v. p. 112 (1834), teste Hochreutiner. Le Jolis, Notes Pl. Europ. in Mem. Soc. Nat. Cherbourg, vol. vi. p. 294 (1891). Tiselius, Pot. Succ. exsic. pp. 107-110 (1896). Ar. Bennett, Journal of Bot. p. 172 (1907).

DESCRIPTION. Forming thick rhizomes with black inflated sheaths, the lowest without a leaf, the second and third with a short leaf 1 inch long.

STEM stout at the base, with very loose inflated sheaths, 1 inch by 3 lines, persistent to time of flowering. Upper sheaths embracing the stem with obtuse leaves 1-1½ inches long.

LEAVES. Upper leaves linear-setaceous, or setaceous, 1-3-ribbed, "without transverse nerves," Turcz. l.c.

PEDUNCULES equal, 2-6 inches long.

SPIKES interrupted, 2-3 inches long.

FRUIT. Drupelets crowded in whorls, obliquely-ovate, semicircular compressed, faintly 3-keeled on the dorsal margin, beak very short, central.

DISTRIBUTION in the British Islands.

Shetland, Beeby.

DISTRIBUTION in other countries.

Europe: Sweden, 64° N. Lat. to Sodermanland, Finland, Archangel (Russia), Switzerland, Austria (?). Asia: Siberia, Turkestan, Songaria, Japan. N. America: Labrador, Canada, 56° N. Lat., Alberta, Saskatchewan. Africa: Transvaal.

P. vaginatus belongs to the group of *P. marinus*, L. (*filiformis*, Nolte), the first differing considerably from the *pectinatus* group. See Baagoe in Videnh. Midd. Nat. Foren. (1903).

EXPLANATION OF PLATE 58. POTAMOGETON VAGINATUS. Drawn from Swedish specimens as fresh specimens could not be obtained from Shetland. Figs. 1, 2 and 5, plant natural size; 3, stipule; 4, flower; 6, fruit; figs. 3, 4 and 6 mag.

40. POTAMOGETON INTERRUPTUS, Kitab, ap Schultz, Oest Flora, ed. 2, p. 328 (1814).

P. FLABELLATUS, Babington, Man. Brit. Bot. ed. 3, p. 343 (1851). *P. pectinatus*, L., sub-sp. *flabellatus*, Syme, Eng. Bot. ed. 3, vol. ix. p. 53, tab. 1421 (1869). Hooker, Stud. Flor. ed. 3, 436 (1884). Morong, Naid. N. America, p. 53, tab. 60 (1893).

DESCRIPTION. Plant growing from tubers, and a perennial rhizome, creeping in the mud, with short black, rather loose sheaths.

STEM long, stout, simple below, much-branched above, spreading in a more or less fan-shaped manner.

LEAVES. Lower leaves 3-8 inches long by ½-1½ lines wide, 3-5-ribbed, usually decaying by the time the plant is in flower, upper leaves ¾ line broad, 1-3-ribbed.

PEDUNCULES slender, equal, 1-4 inches long.

SPIKE 1-2 inches, often interrupted.

SEPALS reniform-orbicular.

FRUIT. Drupelets 2 lines long, nearly straight on the ventral margin, but gibbous at the apex, with the beak forming a continuation, semicircular on the dorsal margin, central keel pronounced, lateral ridges.

β scoparius, Fryer, Jour. Bot. vol. xxvi. p. 297 (1888). *P. pectinatus*, var. *scoparius*, Wallr., Sch. Crit. p. 68 (1822). *P. marinus*, Hudson, Fl. Angl. ed. 2, p. 77 (1778). Stem naked below when fruiting, much-branched and leafy at the apex, early leaves flat, 3-ribbed, upper setaceous, 1-ribbed. Dispersed over England as far north as Yorkshire.

DISTRIBUTION in British Islands.

Cornwall to Orkney; in fifty-eight counties and vice-counties. Ireland: Kerry to Donegal, in twenty counties.

DISTRIBUTION in other countries.

Europe: Sweden to Italy, and France to the Caucasus. Asia: Siberia, Mongolia, and India. Africa: Egypt, Libyan Desert, Madagascar. N. America: Canada, 49° N. Lat., Alberta, U.S., Pennsylvania, N. California and bordering States. S. America: Argentina, Uruguay, Chili. Australasia: New Zealand, S. Australia (?).

The Rev. K. Trimmer, in his Flora of Norfolk, p. 155, 1866, mentions that "In the rapid current of the Trent by Walton-on-Trent Bridge, where *P. flabellatus* abounds. I have always met with plenty of the broad lower leaves in undecayed condition through the months of July and August."

Ascherson and Graebner, in Synop. Fl. Mitteleurop. ed. 2, p. 541 (1913), separate *flabellatus* and *interruptus* under *pectinatus*, L.; referring the synonymy under *interruptus* to *P. Vaillantii*, Roem. et Sch. Sys. vol. iii. p. 514 (1818), and *pectinatus* var. *dichotamus*, Wallr., Sch. Crit. p. 68 (1822). To me they are simply forms coming (with *flabellatus*) under *interruptus*, so far as the specimens can be relied on at Berlin. Of course much of this will simply be difference of opinion as to the value of the characters, and until the whole genus is more completely investigated it ill becomes any one to be dogmatic in these matters. It is possible, and perhaps probable, that the North American form will have to be separated from the European, but until more specimens are obtained, or the plants studied *in situ*, it is best to leave them under one head.

EXPLANATION OF PLATE 59. POTAMOGETON INTERRUPTUS. Drawn from Cambridge-shire specimens. Fig. 1, early form; 2, later form, both natural size; 3, section of stem; 4 and 5, parts of leaves and stipules; 6, flower spike; 7, flower; 8, fruit spike; 9, longitudinal section of fruit; figs. 3 to 9 mag.

41. POTAMOGETON MARINUS, Linn. Sp. Pl. ed. 1, p. 126 (1753).

P. FILIFORMIS, Pers. Syn. 1, 16, 152 (1805). Nolte, Nov. Fl. Holst. p. 20 (1826). *P. marinus*, Koch, Synop. Flor. Germ. et Helv. ed. 2, vol. ii. p. 781 (1844). Reichenbach, Icon. Flor. Germ. vol. vii. p. 11, tab. 18, figs. 28, 29 (1845). Fries, Novit. Flor. Suec.

ed. p. 54 (1828). *P. filiformis*, Nolte, Syme, Eng. Bot. ed. 3, ix. 55, t. 1424 (1869). Hooker, Stud. Flor. ed. 3, 436 (1884). Morong, Naid. N. America, 50, t. 57 (1893). Babington, Man. Brit. Bot. ed. 9, 443 (1904). Ascherson u. Graebner, Synop. Flor. Mitteleurop. ed. 2, 542 (1913).

DESCRIPTION. Creeping in the mud with slender rhizomes, and throwing up tufts of leaves at intervals.

STEMS slender, terete or subcompressed, filiform, sparingly branched at the base, and simple above.

LEAVES all similar, setaceous, subacute, 1-ribbed.

STIPULES rather long and narrow.

PEDUNCULES equal 4-9 inches long.

SPIKES 2 inches, interrupted, few-flowered in whorls.

SEPALs obovate.

FRUIT. Drupelets small, olive, subcompressed, semioblanceolate on the ventral margin, semicircular on the dorsal, bluntly 3-keeled, with a short, nearly central beak. Plant bright green, generally turning darker in drying. When not in fruit and robust *marinus* is difficult to separate from *pectinatus*, which at times assumes a *filiformis*-like aspect, but when in fruit specimens can be easily separated.

DISTRIBUTION in the British Islands.

Anglesea (Davies herbarium), Berwick north to Shetland, Outer Hebrides; in about 17 counties. Ireland: Galway to Armagh; in 10 of Mr. Praeger's divisions.

DISTRIBUTION in other countries.

Europe: Iceland, Faroes, Norway, 70° 51' N. Lat., Sweden to Piedmont, France (H. Alps), Russia. Asia: Siberia, 70° N. Lat., Pamirs and Mongolia to Turkestan, Persia, India, Japan. Africa: Egypt, Reuwendzani Expedition, 1893-4. N. America: Greenland to 71° N. Lat., Canada, Labrador to Hudson's Bay and B. Columbia, U. States, Maine to Dakota and California. S. America: Chili (Gay); Argentina (Fries). Australia: Victoria, S. Australia, New Zealand.

EXPLANATION OF PLATE 60. POTAMOGETON MARINUS. Drawn from Forfarshire specimens; with longitudinal section of fruit and section of stem, mag.

42. **POTAMOGETON PENNSYLVANICUS**, Cham. et Schlecht., Linn. ii. 227 (1827).

P. CLAYTONII, Tuckerman, Am. Jour. Science, Ser. 1, vol. xiv. p. 38 (1843). *P. Nuttallii*, Chamisso et Schlechtendal, l.c., teste Morong, in Naid. N. America, tab. 29, 2 (1893). Ar. Bennett, Naturalist, x. 10, 373 (1908). Ar. Bennett, Trans. Edinb. Bot. Soc. vol. xxiii. p. 211 (1908).

DESCRIPTION. "Stems slender, compressed, mostly simple, 0-3-2 m. long. Floating leaves opposite, elliptic to obovate, obtuse, short-petioled, 3-8 cm. long, 8-24 mm. wide, many-nerved; submerged leaves linear, 2-ranked, 5-17 cm. long, 2-6 mm. wide, 5-nerved with reticulations on either side of midrib; stipules obtuse, hyaline, not keeled; peduncles 2-13 cm. long; spikes 1-2-5 cm. long; fruit round-obovoid, 2.5-4 mm. long, 2-3 mm. thick, 3-keeled, the sides flat and indistinctly impressed; style short, apical; embryo coiled one and one-third times." Morong, ex Britton, Man. Fl. N. States and Canada, p. 43 (1901).

DISTRIBUTION. From N. Scotia to B. Columbia, and Maine to Georgia, and New Jersey to Washington Territory.

Found by Miss A. E. Vigurs in the canal at Salterhebble Bridge, near Halifax, Yorkshire, in 1907; and, the next year, between there and Elland, three miles up the canal. Most likely introduced with cotton, as it is one of the common species in the United

States, being abundant in the States where cotton is grown. I believe this to be the only instance of the introduction of a *Potamogeton*. The supposed case of *P. crispus*, L., to North America proved a mistake; although said to be "not known to the earlier American botanists," Pursh recorded it in 1814 for several habitats, and says he has seen living specimens, "Canada to Virginia."

I give the earliest name of which there are specimens extant (i.e. at Berlin), but it is probable that Rafinesques' name of *P. epiphydum* in Medical Reports 2nd Hex. vol. v. p. 354 (1808) will eventually have to be accepted.

Dr. Morong, i.e., adopts the name *P. Nuttallii* or the page before the one I quote, but while well knowing the keenness and ability of my late friend, I cannot agree with him in this, though of course I may be mistaken.

P. Pennsylvanicus.

Fig. 2.—1, plant; 2, flower; 3, segment of perianth with stamen; 4, segment of perianth; 5, fruit; 6, ovary.

CORRIGENDA.

Page 47, headline, *for* "Cooperi" *read* "crispus."

Page 56, line 1, "densus" should have been numbered 16, instead of 17, the number of subjects separately described in the work is 41, and not 42 as numbered.

The subjects on Plates 39 to 47 do not follow the same order as in the text. But the references to the Plates in the text are correct.

Plates 20 and 26. Figures 6, 7, 8 on Plate 26 are the fruit of *P. alpinus* omitted on Plate 20.

INDEX.

Names of groups are printed in small capitals, and of synonyms in italics.

The page on which the main description of the plant is given is indicated by black numerals.

In this Index a × is used to indicate the species crossed and not the name of the hybrid.

- acuminatus*, v. (*pusillus*), 84.
acutifolius, 79.
 ——— Plate 51.
alpinus, 27.
 ——— Plates 18–20, 26.
 ——— × *lucens*, 76.
 ——— × *natans*, 31.
 ——— × *polygonifolius*, 31.
 ——— × *praelongus*, 34.
angustifolius, 72.
angustifolius, v. (*polygonifolius*), 20, 22.
australis, v. (*pusillus*), 82.

Bennettii, 47, 53.
 ——— Plate 33.
Bertcholdii, v. (*pusillus*), 84.
Billupsii, 25.
 ——— Plates 16, 17.
borealis, f. (*alpinus*), 28.

caepitosus, 82.
cancellata, f. (*polygonifolius*), 21.
 Chloephylli group, 2, 57, 77.
 Claytonii, 90.
 COLEOPHYLLI group, 2, 57, 86.
coloratus, 21, 22.
 ——— Plates 13–15.
 ——— × *Zizii*, 25.
compressus, 77, 80, 81.
Cooperi, 47, 48.
 ——— Plates 31, 32.
coriaceus, 27, 59, 74.
 ——— Plate 43.
 ——— × *natans*, 8.
 ——— × *perfoliatus*, 69.
cornutus, 72.
 ——— v. (*crispus*), 45.
crassifolius, 8, 15.
 ——— Plates 4, 5.
 ——— f. (*verruta*), 9, 17.

crispus, 43.
 ——— Plates 29, 30.
 ——— × *Friesii*, 54.
 ——— × *obtusifolius*, 53.
 ——— × *perfoliatus*, 48.
 ——— × *praelongus*, 49.
curvifolius, 59, 69, 70.
cuspidatus, 79, 80.
eymatodes, 48.

decipiens, 59, 75.
 ——— Plate 48.
 ——— v. *salicifolius*, 76.
densifolia, f. (*perfoliatus*), 39.
densus, 56.
 ——— Plate 34.
dichotomus, v. (*pectinatus*), 89.
Drucei, 31.
 ——— Plate 21.
 ——— *dubius*, 17.

elegans, f. (*praelongus*), 36.
 ENANTIOPHYLLI group, 2, 57.
epihydrium, 91.
ericetorum, v. (*polygonifolius*), 20, 22.
eupectinatus, sub-sp. (*pectinatus*), 86.

falcatus, 68.
 ——— Plate 40.
filiformis, 90.
flabellatus, sub-sp. (*interruptus*), 88.
fluitans, 11.
 ——— Plates 6, 7.
 ——— f. *rivularis*, 17.
fluitans, 28, 31.
fluvialis, v. (*obtusifolius*), 81.
fluvialilis, v. (*pectinatus*), 87.
Friesii, 81.

Friesii, Plate 53.
 ——— × *crispus*, 47, 54.
 ——— × *heterophyllus*, 63.

gemmifer, v. (*crispus*), 43, 47.
genuinus, v. (*alpinus*), 29.
 ——— v. (*polygonifolius*), 20.
gracilior, f. (*alpinus*), 28.
gramineum, 59.
gramineus, 64, 65, 80.
 ——— × *nitens*, 68.
graminifolius, 64.
 ——— Plate 36.
 GRASS-LEAVED section, 77.
Griffithii, 34.
 ——— Plates 22, 23.

 HETEROPHYLLI group, 2.
heterophyllus, 59, 65.
 ——— Plate 35.
 ——— × *Friesii*, 63.
 ——— × *polygonifolius*, 21.
 ——— × *pusillus*, 63.
 ——— 64.
 ——— f. (*lucens*), 72.
 ——— sub-sp. (*nitens*), 70.
 ——— v. (*perfoliatus*), 48.
hibernicus, v. (*lanceolatus*), 63.
 HOMOPHYLLI group, 2.
homophyllus, v. (*alpinus*), 29.
Horneumanni, 22.

imbecilla, f. (*perfoliatus*), 39.
interruptus, 88.
 ——— Plate 59.
involuta, f. (*nitens*), 69.
involutus, 69.
 ——— Plate 44.
iterata, f. (*alpinus*), 28.

Jacksoni, v. (*perfoliatus*), 48.
junctifolius, 87.

- Kirkii, 10, 16.
 ——— Plates 8, 9.
 lanceolatus, 63.
 ——— Plate 39.
 ——— v. (perfoliatus), 41.
 linearis, v. (natans), 6.
 lonchites, 13.
 longifolia, f. (alpinus), 28.
 Lucens group, 59.
 lucens, 71.
 ——— Plates 45–47.
 ——— f. *alpinus*, 30.
 ——— v. *coriaceus*, 27.
 ——— sub-sp. *decipiens*, 75.
 ——— *heterophyllus*, 72.
 ——— sub-sp. *Zizii*, 72.
 ——— × *alpinus*, 76.
 ——— × *natans*, 11, 14.
 ——— × *perfoliatus*, 75.
 major, 81.
 marinus, 89.
 ——— Plate 60.
 marinus, 89.
 monogyneus, 85.
 mucronatus, 81.
 natans, 4, 19, 24.
 ——— Plates 1–3.
 ——— × *alpinus*, 31.
 ——— × *coriaceus*, 8.
 ——— × *lucens*, 11, 14.
 ——— × *polygonifolius*, 16.
 ——— × *Zizii*, 8.
 natans, v. *prolixus*, 6.
 nitens, 7, 59, 69.
 ——— Plates 37, 38.
 ——— f. *involuta*, 69.
 ——— × *gramineus*, 68.
 Nuttali, 90.
 oblongus, 18.
 obtusifolius, 80.
 ——— Plate 52.
 ——— × *crispus*, 47, 53.
 pachystachyus, v. (*plantagineus*), 22.
 palustris, f. (*alpinus*), 28.
 pectinatus, 86.
 ——— Plate 57.
 ——— sub-sp. *flabellatus*, 88.
 ——— v. *fluciatilis*, 87.
 Pennsylvanicus, 90.
 perfoliatus, 38, 59, 70.
 ——— Plates 27, 28.
 ——— v. *heterophyllus*, 48.
 ——— v. *Jacksoni*, 48.
 ——— × *coriaceus*, 69.
 ——— × *crispus*, 47, 48.
 ——— × *lucens*, 75.
 Phialæ, 85.
 plantagineus, 21, 22.
 polygonifolius, 17, 18, 24.
 ——— Plates 10–12.
 ——— f. *cancelata*, 21.
 ——— v. *pseudo-fluitans*, 33.
 ——— × *alpinus*, 31.
 ——— × *natans*, 16.
 polygonifolius, 16.
 ——— c. *linearis*, 6.
 praelongus, 36.
 ——— Plates 24–26.
 ——— × *alpinus*, 34.
 ——— × *crispus*, 47, 49.
 prolixus, 6.
 Proteus, 59.
 ——— *curvifolius*, 69.
 ——— *heterophyllus*, 65.
 ——— *lucens*, 71.
 ——— *Zizii*, 72.
 pseudo-fluitans, v. (*polygonifolius*), 20, 33.
 pusillus, 83, 84.
 ——— Plate 55.
 ——— v. *australis*, 82.
 ——— × *heterophyllus*, 63.
 purpurascens, 28.
 rigidus, v. (*pusillus*), 84.
 Robbinsii, 55.
 rotundifolia f. (*perfoliatus*), 39.
 rufescens, 28.
 rutilus, 82.
 ——— Plate 54.
 salicifolius, 76.
 salignus, 76.
 ——— Plate 49.
 scoparius, v. (*interruptus*), 89.
 serratum, 56.
 serratus, 45.
 serrulatus, 55.
 ——— v. (*crispus*), 43.
 similis, v. (*pusillus*), 84.
 sparganifolius, 16.
 spathulæformis, 67.
 spathulatus, 33.
 stagnatilis (*fluitans*), 11.
 Sturrockii, 84.
 tenuissimus, v. (*pusillus*), 84.
 trichoides, 85.
 ——— Plate 56.
 Trimmeri, v. (*trichoides*), 86.
 tuberculatus, 85.
 undulatus, 47.
 undulatus, 48.
 ——— v. *Cooperi*, 48.
 vaginatus, 87.
 ——— Plate 58.
 Vaillantii, 89.
 varians, 67.
 ——— Plate 41.
 verruta, f. (*crassifolius*), 9, 17.
 virides, f. (*alpinus*), 28.
 vulgaris, v. (*pusillus*), 84.
 Zizii (= *Angustifolius*), 59, 72.
 ——— Plate 42.
 Zizii II. *coriaceus*, 74.
 Zizii × *coloratus*, 25.
 ——— × *natans*, 8.
 zosteraefolius, 77, 80.
 ——— Plate 50.

R. Morgan del et lith

L. Reeve & C^o London

Viney & Brooks, Day & Son Imp

J. Gardner del.

J. Reeve sc. London

V. & A. Brown & Sons

R Morgan del. et lith.

Grass, 2nd ed. 1841, vol. 1, p. 101, fig. 1.

R. Morgan del. et lit.

J. Reeve & Co. London

Morgan del et lith

Vincent Brooks, Day & Son Imp

L. Reeve & Co London

Fig 2.

Fig 1.

Fig 1.

Fig 2.

R. Morgan del et lit

Windsor, New York, N. Y. 1874

Reeve & Co. London

4. x5.

x8.

5.

3

2

Margar del sulci

J. Reeve & J. L. L. L.

Thymus, Portulaca, & L. L. L.

R. Morgan del. et lith.

Vincent Brooks, Day & Son Lith.

L. Reeve & Co. London.

R. Morgan del. lith.

L. Reeve & Co. London

Vincent Brooks Day & Son Ltd.

F. Mangar delatellii

Vandier Br. Mus. Nat. Hist. Brit. Mus.

L. Reuter & Co.

H. Virgin. del. et lith.

Paspalum

J. E. Smith & Co. Lith.

Paspalum

Botanical Illustration

Eleocharis acicularis (L.) Rostk.

Eleocharis acicularis (L.) Rostk.

Eleocharis acicularis (L.) Rostk.

Arisaema aculeatum

Botany 10. 11. 1850

Small Arisaema, Spadix and Bract

Calamagrostis phaeocephala

2

L. Rees & C. Smith

Journal of Botanical Society of London 14. 1894

x9

Eleocharis acicularis

Eleocharis acicularis

Eleocharis acicularis

F. C. Knight del et lit

Chenopodium *Chenopodium* *Chenopodium*

M. Smith del. et sculp.

E. Knight del. et lith.

J. Reeve & Sons London

Valeriana (L.) Benth.

M. Smith del et lith

L. Reeve & Co. London

William Brown, Day & Son, Edinburgh

M. Smith del et lith.

Vincent Brown & Son Lith.

Stylidium lineare (L.) DC.

George Engelmann

Illustration of *Stylidium lineare* (L.) DC.

University of Toronto
Library

DO NOT
REMOVE
THE
CARD
FROM
THIS
POCKET

Acme Library Card Pocket
Under Pat. "Ref. Index File"
Made by LIBRARY BUREAU

UTL AT DOWNSVIEW

D RANGE BAY SHLF POS ITEM C
39 09 04 17 08 007 0