Photographic Sciences Corporation

**23 WEST MAIN STREET
WEBSTER, N.Y. 14580
(716) 872-4503**

**CIHM/ICMH
Microfiche
Series.**

**CIHM/ICMH
Collection de
microfiches.**

Canadian Institute for Historical Microreproductions / Institut canadien de microreproductions historiques

© 1982

Technical and Bibliographic Notes/Notes techniques et bibliographiques

The Institute has attempted to obtain the best original copy available for filming. Features of this copy which may be bibliographically unique, which may alter any of the images in the reproduction, or which may significantly change the usual method of filming, are checked below.

L'Institut a microfilmé le meilleur exemplaire qu'il lui a été possible de se procurer. Les détails de cet exemplaire qui sont peut-être uniques du point de vue bibliographique, qui peuvent modifier une image reproduite, ou qui peuvent exiger une modification dans la méthode normale de filmage sont indiqués ci-dessous.

- ☐ Coloured covers/
Couverture de couleur
- ☐ Covers damaged/
Couverture endommagée
- ☐ Covers restored and/or laminated/
Couverture restaurée et/cu pelliculée
- ☒ Cover title missing/
Le titre de couverture manque
- ☐ Coloured maps/
Cartes géographiques en couleur
- ☐ Coloured ink (i.e. other than blue or black)/
Encre de couleur (i.e. autre que bleue ou noire)
- ☐ Coloured plates and/or illustrations/
Planches et/ou illustrations en couleur
- ☐ Bound with other material/
Relié avec d'autres documents
- ☐ Tight binding may cause shadows or distortion
along interior margin/
La reliure serrée peut causer de l'ombre ou de la
distortion le long de la marge intérieure
- ☐ Blank leaves added during restoration may
appear within the text. Whenever possible, these
have been omitted from filming/
Il se peut que certaines pages blanches ajoutées
lors d'une restauration apparaissent dans le texte,
mais, lorsque cela était possible, ces pages n'ont
pas été filmées.
- ☐ Additional comments:/
Commentaires supplémentaires:

- ☐ Coloured pages/
Pages de couleur
- ☒ Pages damaged/
Pages endommagées
- ☐ Pages restored and/or laminated/
Pages restaurées et/ou pelliculées
- ☒ Pages discoloured, stained or foxed/
Pages décolorées, tachetées ou piquées
- ☒ Pages detached/
Pages détachées
- ☐ Showthrough/
Transparence
- ☐ Quality of print varies/
Qualité inégale de l'impression
- ☐ Includes supplementary material/
Comprend du matériel supplémentaire
- ☐ Only edition available/
Seule édition disponible
- ☐ Pages wholly or partially obscured by errata
slips, tissues, etc., have been refilmed to
ensure the best possible image/
Les pages totalement ou partiellement
obscurcies par un feuillet d'errata, une pelure,
etc., ont été filmées à nouveau de façon à
obtenir la meilleure image possible.

This item is filmed at the reduction ratio checked below/
Ce document est filmé au taux de réduction indiqué ci-dessous.

10X	12X	14X	16X	18X	20X	22X	24X	26X	28X	30X	32X
☐	☐	☐	☐	☐	☐	☐	☒	☐	☐	☐	☐

The copy filmed here has been reproduced thanks to the generosity of:

Library of Congress
Photoduplication Service

The images appearing here are the best quality possible considering the condition and legibility of the original copy and in keeping with the filming contract specifications.

Original copies in printed paper covers are filmed beginning with the front cover and ending on the last page with a printed or illustrated impression, or the back cover when appropriate. All other original copies are filmed beginning on the first page with a printed or illustrated impression, and ending on the last page with a printed or illustrated impression.

The last recorded frame on each microfiche shall contain the symbol ➡ (meaning "CONTINUED"), or the symbol ▼ (meaning "END"), whichever applies.

Maps, plates, charts, etc., may be filmed at different reduction ratios. Those too large to be entirely included in one exposure are filmed beginning in the upper left hand corner, left to right and top to bottom, as many frames as required. The following diagrams illustrate the method:

L'exemplaire filmé fut reproduit grâce à la générosité de:

Library of Congress
Photoduplication Service

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole ➡ signifie "A SUIVRE", le symbole ▼ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.

QL 469
.G 8

*Prof Joseph Henry
from H. C. Cotton*

[From the Proceedings of the American Association for the Advancement of Science,
Detroit Meeting, August, 1875.]

12
**THE EFFECT OF THE GLACIAL EPOCH UPON THE DISTRIBUTION OF
INSECTS IN NORTH AMERICA. By AUG. R. GROTE, of Buffalo,
N. Y.**

FROM the condition of an hypothesis, the Glacial epoch has been elevated into that of a theory, by the explanations it has afforded to a certain class of geological phenomena. The present paper endeavors to show that certain zoological facts are consistent with the presence in past times of a vast progressive field of ice gradually extending over large portions of the North American Continent and moving from the north to the south. These facts are in the present instance afforded by a study of the Lepidoptera, certain kinds of butterflies and moths now inhabiting the United States and adjacent territories. Before proceeding with the subject, a brief statement of some of the phenomena assumed to have attended the advent of the Glacial Epoch is necessary.

At the close of the Tertiary, the temperature of the earth's surface underwent a gradual change by a continuous loss of heat. The winters gradually became longer, the summers shorter. The tops of granitic mountains in the east and west of the North American continent, now in summer time bare of snow and harboring a scanty flora and fauna, became, summer and winter, covered with congealed deposits. In time the mountain snows consolidated into glacial ice which flowed down the ravines into

the valleys. Meanwhile the northern regions of the continent, which may have inaugurated, submitted extendedly to the same phenomena. Glacial ice, first made on elevations, finally formed at, and poured over lower levels. Glacial streams finally united to form an icy sea whose frozen waters slowly plowed the surface of the rocks; and whose waves, in their movement from north to south, absorbed the local glaciers in their course, and extended over all physical barriers into the Southern States and down the Valley of the Mississippi. To the main Ice-sheet, the Appalachians and Rocky Mountains are supposed to have contributed their local glaciers. Before this frozen deluge the animals must have always retreated. The existing insects of the pliocene must, in submitting to the change of climate which accompanied the advance of the glacier, have quitted their haunts with reluctance, and undergone a severe struggle for existence, no matter how gradually they had been prepared for the encounter. We must expect that multitudes of specific forms ultimately perished of whose remains no traces have been preserved.

Such being a brief statement of the outlines of the opening of the Glacial Epoch, we turn to some facts offered by a study of our existing species of butterflies and moths. The tops of the White Mountains and the ranges of mountain elevations in Colorado, offer us particular kinds of these insects living in an isolated manner at the present day and confined to their respective localities. In order to find insects like them we have to explore the plains of Labrador and the northern portion of the North American Continent, in regions offering analogous conditions to those obtaining on the summits of these mountains. The genera *Oeneis* and *Brenthis* among the butterflies, and *Anarta* and *Agrotis* among the moths, are represented by the same or similar species in all of the above mentioned localities. In the case of the White Mountain Butterfly, *Oeneis semidea*, we have a form sustaining itself on a very limited alpine area on the top of Mount Washington.¹ Although there is some doubt that precisely

¹ See Mr. Seudder's article in the "Geology of New Hampshire," 1,342. Mr. Seudder first pointed out the existence of alpine and subalpine faunal belts on Mount Washington, and interestingly remarks "that if the summit of Mt. Washington were somewhat less than two thousand feet higher it would reach the limit of perpetual snow." Consult also, an earlier paper of great value by Dr. A. S. Packard, Jr., on "The Insect Fauna of the Summit of Mount Washington as compared with that of Labrador" (these Proceedings, Vol. XVI, 151). Dr. Packard, in comparing the climate of the two localities, says: "The seasons correspond very exactly, as the snow melts in the early summer, and ice is formed early in the autumn at about the same dates."

Q-469

.G8

the same form has been discovered in Colorado, the fact remains that butterflies exceedingly like it, though registered by us under different specific names, live in Labrador and Colorado.

Whether the White Mountain Butterfly be, as suspected by Lederer, a local modification of some one of the Labradorian forms or not, the geographical distribution which its genus enjoys cannot be meaningless. The question comes up, with regard to the White Mountain Butterfly, as to the manner in which this species of *Oeneis* attained its present restricted geographical area. How did the White Mountain Butterfly get up the White Mountains? And it is this question that I am disposed to answer by the action attendant on the decline of the Glacial Epoch.

I have before briefly outlined the phenomena attendant on the advance of the Ice-sheet, and I now dwell for a moment on those which must equally be presumed to have accompanied its retirement. Many of the features of its advance were repeated in reverse order on the subsidence of the main Ice-sheet or Glacial sea. The local glaciers appeared again separate from the main body and filled the valleys and mountains and ravines, running thus at variance with the main body of the Glacier, being determined by local topography. A reversal of the temperature shortened the winters and lengthened the summers. Ice-loving kinds of insects, such as our White Mountain Butterfly, hung on the outskirts of the main Ice-sheet, where they found their fitting conditions of temperature and food. The main Ice-sheet had pushed them insensibly before it, and, during the continuance of the Glacial Epoch, the geographical distribution of the genus *Oeneis* had been changed from a high northern region to one which may well have included portions of the Southern States. And, on its decline, the Ice-sheet drew them back again after itself by easy stages; yet not all of them. Some of these butterflies strayed by the way, delayed by the physical nature of the country and destined to plant colonies forever separate from their companions. When the main Ice-sheet left the foot of the White Mountains, on its long march back to the pole where it now seems to rest, some of these wayward, flitting, *Oeneis* butterflies were left behind. These had strayed up behind the local glaciers on Mount Washington, and so became separate from the main body of their companions which journeyed northward, following the retirement of the main Ice-sheet. They found in elevation their

congenial food and climate, and they have followed these gradually to the top of the mountain, which they have now attained and from which they cannot now retreat. Far off in Labrador, the descendants of their ancestral companions fly over wide stretches of country, while they appear to be in prison on the top of a mountain.

I conceive that in this way the mountains generally may have secured their Alpine animals. The Glacial Epoch cannot be said to have expired. It exists even now for high levels above the sea while the Laplander and Esquimaux find it yet enduring in the far North. Our yearly winters are fractions of the Glacial year. Had other conditions been favorable, we might now find Arctic man living on snow-capped mountains in the Temperate zones.

At a height of between 5,600 and 6,200 feet above the sea and at a mean temperature of about forty-eight degrees during a short summer, the White Mountain Butterflies (*Oeneis semidea*), yet enjoy a climate like that of Labrador within the geographical limits of New Hampshire. And in the cases of the moths² an analogous state of things exists. The species *Anarta melanopa* is found on Mount Washington, the Rocky Mountains and Labrador. *Agrotis islandica* is found in Iceland, Labrador, the White Mountains, and, perhaps, Colorado.³ As on islands in the air, these insects have been left by the retiring of the ice-flood during the opening of the Quaternary.

On inferior elevations, as on Mount Katahdin in Maine, where we now find no *Oeneis* butterflies, these may have formerly existed, succumbing at last to a climate gradually increasing in warmth from which they had no escape; while the original colonization in the several instances must have always greatly depended upon local topography.

In conclusion, I have briefly endeavored to show that the present distribution of certain North American insects may have been brought about by the phenomena attendant on the Glacial Epoch. The discussion of matters connected with this theoretic period of the earth's history, still, as it now appears, brings out more and more a clearer conception of its actuality. I hope that

² I have since (*Psyche*, I, 131) recorded the first indication of the occurrence of the Arctic *Larva Rossii* on Mount Washington, from a single specimen taken by Mr. B. Pickman Mann, above the tree line.

³ I believe Dr. Packard's identification of this species, in Prof. Hayden's Reports, is incorrect. The Coloradan species is *Agrotis auxiliaris*, Grote.

my present statements may draw the attention of our zoölogists more fully to the matter, seeing that we have in our own country fields for its full exploration. And I permit myself to believe, that testimony as to the former existence of a long and widely spread winter of the years, is offered in evidence through the frail, brown, *Oeneis* butterflies, that live on the tops of the mountains.

[Printed at the SALEM PRESS, June, 1870.]