

A NEW KALLSTROEMIA FROM TEXAS

LOUIS WILLIAMS

Washington University Fellow in Botany

Kallstroemia hirsuta n. sp.¹ Ascending or erect annual, 1–2 dm. high; stems 1–several from the base, densely hirsute and strigose-pubescent, the hirsute hairs with pustulate bases; leaves 2–4.5 cm. long, oblong to ovate-oblong, petiolate, densely hirsute and strigose-pubescent on both surfaces, the petiole about as long as the lower pair of leaflets; leaflets 3–6 pairs, sessile, obliquely oblong or ovate, 7–14 mm. long, 3–8 mm. wide, obtuse or acute, about equal in size, the terminal more oblique; stipules lance-oblong, 3–4 mm. long, 1–1.5 mm. wide; calyx-lobes linear to linear-lanceolate, 10–13 mm. long, 1.5–2.5 mm. wide, hirsute and strigose-pubescent on the outer surface, the chartaceous margins inrolled in fruit; petals about 5, deltoid-ovate, 18–23 mm. long, probably white or yellowish, not marcescent; fruit compressed-oval in outline, 8–10 mm. wide, 5–6 mm. high, outer surface of carpels not tuberculate but the median portion and the two edges prominent and connected by transverse ridges, densely hirsute, particularly on the raised portions; persistent style or beak 8–10 mm. long, hirsute at the inconspicuously swollen base, stigmatic portion 1.5–2 mm. long, furcate.

TEXAS: Langtry, Valverde County, May, 1913, *C. R. Orcutt*, 6126 (Mo. Bot. Gard. Herb.) TYPE.

This species is most closely allied to *K. grandiflora* Torr., from which it may be separated by the following characters: the entire plant is densely pubescent with hirsute and strigose

¹ *Kallstroemia hirsuta* sp. nov. Annuā 1–2 dm. alta, dense hirsuta et strigoso-pubescente; foliis oblongis aut oblongo-ovatis utrinque pubescentibus; foliolis 3–6 paribus, sessilibus, 7–14 mm. longis, 3–8 mm. latis; stipulis lanceolato-oblongis, 3–4 mm. longis, 1–1.5 mm. latis; lobis calycis linearibus vel lineari-lanceolatis, 10–13 mm. longis, 1.5–2.5 mm. latis; petalis ca. 5, deltoideo-ovatis, 18–23 mm. longis; carpellis non tuberculatis parte medio et marginibus prominentibus; stylo 8–10 mm. longo, ad basem hirsuto.

Issued March 25, 1935.



Williams, Louis O. 1935. "A New *Kallstroemia* from Texas." *Annals of the Missouri Botanical Garden* 22, 49–50. <https://doi.org/10.2307/2394138>.

View This Item Online: <https://www.biodiversitylibrary.org/item/54266>

DOI: <https://doi.org/10.2307/2394138>

Permalink: <https://www.biodiversitylibrary.org/partpdf/11513>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.