

Bonn. zool. Beitr.	Jg. 36	H. 3/4	S. 585—606	Bonn, Oktober 1985
--------------------	--------	--------	------------	--------------------

Conservation status of reptiles in the Canary Islands

by

A. Machado, L.F. López-Jurado & A. Martín

Introduction

On the occasion of the symposium „Herpetologia Canariensis”, organized by the Museum Alexander Koenig (Bonn), November 1—3, 1984, we considered it appropriate to present an actual vision of the conservation status of the Canarian herpetofauna. We also feel compelled to do it because of the recent proliferation of endangered species lists and red data books, completed by international organizations (IUCN, Council of Europe, etc). In various cases, these works only compile information previously extant in publications, which are usually not specific on conservation. This gives base to a distorted view of the true status of the species in nature.

In this context, we present an analysis of the conservation status of the Canarian reptiles (marine species excluded) based on our direct field knowledge and gathered throughout several years of work in all islands. Moreover, we do not have any reference of publications that treat this matter in detail. That of Baez & al. (1984), despite the title, only refers to distribution of species and their natural enemies, and does not discuss specific conservational aspects.

Studies on Canarian reptiles

The literature on Canarian reptiles is rich relative to the variety of the fauna and, in particular, to its high endemic level. Thus, many publications attend to descriptions of new taxa, biogeography and the very interesting island evolutionary aspects. There are also general synthetical works like those of Salvador (1984), Klemmer (1976) or Baez (1984). Nevertheless, and as could be stated by the papers presented in the Herpetologia Canariensis Symposium, in recent years a revival of Canarian reptile studies has taken place. This may be due to the development of new fields of study (ethology, captive breeding, reproduction, physiology, etc) or the emergence of modern and more sophisticated techniques (cytogenetic, biochemical) applied to classical theses. Field work has become more specialized and even limited populations on small islets and offshore rocks are being studied (Joger & Bischoff 1983, Martín 1985, Bischoff 1985, etc). Another important branch of study, that will probably grow quickly in the immediate future, is that of fossil species systematics and past distribution.

In this paper we have included a bibliography of Canarian herpetology, convinced that it gives a better idea of what has been achieved and that it will be of help to anyone interested in Canarian reptiles.

Threatened categories

The geographic area examined in this work is the Canary Islands. This signifies that the assigned conservation categories apply only to Canarian populations. Since the majority of cases refer to endemics, these categories are valid at a taxa status. The only exceptions are: *Hemidactylus turcicus*, an alien recently introduced, *Chalcides viridanus* and *Chalcides sexlineatus* which also inhabit Madeira, but were probably introduced by man (Pasteur, 1981 p. 7).

The universally accepted threatened categories of IUCN (vide Honegger 1978; Blas Aritio 1976; Groombridge 1982) are used, but slightly adapted to an archipelago scale. The time gap for trend evaluation is approximately 15 years.

Endangered (E): Taxa in danger of extinction and whose survival is unlikely if the causal factors continue operating. Included are taxa whose numbers have been reduced to a critical level or whose habitats have been so drastically reduced that they are deemed to be in immediate danger of extinction. (We will differentiate, as Ex taxa or populations that are possibly already extinct).

Vulnerable (V): Taxa believed likely to move into the endangered category in the near future if the causal factors continue operating. Included are taxa of which most, or all, the populations are decreasing due to over-exploitation, extensive destruction of habitat or other environmental disturbance; taxa with populations that have been seriously depleted and whose ultimate security is not yet assured; and taxa with populations that are still abundant but are under threat from serious adverse factors throughout their range.

Rare (R): Taxa with small world (Canarian) populations that are not at present endangered or vulnerable but are at risk. These taxa are usually localised within restricted geographical areas or habitats or are thinly scattered over a more extensive range.

Indeterminate (I): Taxa known to be "endangered", "vulnerable" or "rare" but where there is not enough information to say which of the three categories is appropriate.

Insufficiently known (K): Taxa that are suspected but not definitely known to belong to any of the above categories, because of lack of information.

Not threatened (nt).

Status of reptiles in the Canary Islands

On table I a general list of Canarian reptiles is given and their threatened category symbols are shown for each island, islet or rock, if present.

Gallotia aff. *simonyi*¹⁾ is the only taxon considered endangered. The population on Hierro is restricted to a stock of one hundred specimens living in extreme conditions on a ridge of a cliff face. For a detailed account see Machado (1985).

1) Blas Aritio (1976 p. 50) considers Gran Canarian (!) and Hierro Giant lizards in need of special protection. IUCN Red Data Book (1978) and Council of Europe's threatened amphibians and reptiles list (Honegger 1978: 79) includes only the latter under "endangered" status.

Table I. Status of reptiles in the Canary Islands.

Fam. GEKKONIDAE	H	s1	s2	G	P	T	r1	r2	C	F	I	L	g	m	a	e
<i>Tarentola d. delalandii</i> (Dum. & Bibron, 1836)		R	R		nt	nt	R	R								
<i>Tarentola boettgeri boettgeri</i> Steind., 1891									nt							
<i>Tarentola b. hierrensis</i> Joger & Bischoff 1983	nt															
<i>Tarentola gomerensis</i> Joger & Bischoff, 1983				G												
<i>Tarentola angustimentalis</i> Steind., 1891									nt	R	nt	nt	R	R	R	V
<i>Hemidactylus turcicus</i> (Linné, 1758)						V			I							
Fam. LACERTIDAE	H	s1	s2	G	P	T	r1	r2	C	F	I	L	g	m	a	e
<i>Gallotia simonyi</i> (Steind., 1889)			Ex													
<i>Gallotia</i> aff. <i>simonyi</i>	E															
<i>Gallotia stehlini</i> (Schenkel, 1901)									nt							
<i>Gallotia galloti galloti</i> (Oudart, 1839)						nt										
<i>Gallotia galloti eisentrauti</i> Bischoff, 1982						nt	R									
<i>Gallotia galloti insulanagae</i> Martín, 1985								V								
<i>Gallotia galloti palmae</i> (Boett. & Müller, 1914)					nt											
<i>Gallotia galloti gomerae</i> (Boett. & Müller, 1914)				nt												
<i>Gallotia galloti caesaris</i> (Lehrs, 1914)	nt	R														
<i>Gallotia atlantica</i> (Peters & Doria, 1882)									V	nt	nt	nt	V	V	V	? ²⁾
Fam. SCINCIDAE	H	s1	s2	G	P	T	r1	r2	C	F	I	L	g	m	a	e
<i>Chalcides viridanus viridanus</i> (Gravenh., 1851)	nt						nt	R								
<i>Chalcides v. coeruleopunctatus</i> Salvador, 1975				nt												
<i>Chalcides sexlineatus</i> Steind., 1891									nt							
<i>Chalcides (polylepis) occidentalis</i> (Steind., 1900) ³⁾										R		I				
SYMBOLS H = Hierro s1 = Greater Roque Salmor s2 = Smaller Roque Salmor P = Palma G = Gomera T = Tenerife r1 = Roque de Tierra r2 = Roque de Fuera C = Gran Canaria F = Fuerteventura I = Isla de Lobos L = Lanzarote g = Graciosa m = Mtña. Clara a = Alegranza e = Roque Este																

2) In the British Museum (Natural History) there are specimens of *Tarentola* and *Gallotia* collected by Hugh B. Cott in the Roque del Este, off Lanzarote (on Sept. 20th 1931, after Vølsoe, 1951). We have been on the rock (March 1976, Sept. 1983) and have not seen lizards.

3) Pasteur (1981) mentions that *occidentalis* may be a valid species.



Fig. 1. The "Roque Chico" of Salmor (El Hierro), type locality of the now extinct *Gallotia simonyi*. Photo A. Machado.

The Lizard of Salmor, *G. simonyi* (Steind.), is extinct as confirmed during August 1984. A further record of *G. simonyi* on the Roque de Fuera (Anaga, Tenerife) given by Báez & Bravo (1983) is probably an error (Martín 1985).

All populations inhabiting small islets or rocks offshore have to be classified as 'rare', because of their small size, but deserve extra consideration when those populations have a special taxonomic status, such as *Gallotia galloti insulana-gae* Martín, 1985 (endemic to the Roque de Fuera). In this particular case, the present day threat is collecting and therefore the taxon was placed under category 'vulnerable'.

Threat analysis

Habitat reduction

Since man arrived in the Canary Islands (not previous to 2000 B.C., Castro Alfin 1983: 83) many changes have taken place in the insular environment, especially with the development of modern civilization. In general terms, one may conclude that a group of large size reptiles have suffered a strong decline if not extinction (Kämmer 1982, Machado 1985).

However, most of the remaining reptilian fauna has gained an important advantage as the available habitat has increased due to deforestation and increased aridity in vast sectors of the islands (vide Machado 1976, Castro Alfin 1983). Other favourable factors include the establishment of fruit monocultures (vineyards, *Opuntia*, tomatoes, bananas, etc) and the extensive building of stone fences and walls (fig. 2).

In general, habitat reduction is not a real problem for Canarian reptiles today. As suggested above, many lizards and geckos adapt successfully to agricultural




Fig. 2. The cultivated areas in the intermediate zones of some islands may represent a habitat expansion for thermophilic reptiles. Photo I. Sánchez.

systems, and they may even increase, as sometimes occurs with other groups of vertebrates (v. habitat expansion in birds, Vølsoe 1955). Nevertheless, some special cases are worth mentioning.

Hemidactylus established itself in the Barranco de Santos in the surroundings of the municipal market of Santa Cruz (Baez 1979) but it could be threatened if the works of re-development and canalization of the ravine (= barranco) continue.

The Gran Canarian population of *Gallotia atlantica*⁴⁾ is restricted to a small area of 2.5 x 1 km, and it is possible that this area may become urbanized.

The small Roque del Este, 11 km off Lanzarote, was used as a target by Spanish Navy and Airforce (see footnote 2) till the sixties. The Defense Department was made aware that the largest Spanish colony of *Falco eleonora* breeds there. The bombing was subsequently stopped, but there is no guarantee that it will not be resumed.

Killing, trapping and biocides

Country youths amuse themselves by 'fishing' or killing lizards. Catching is done with the help of a long 'balango'⁵⁾ (= *Avena barbata* Pott. ex Link) whose stem-tip is transformed into a lasso noose. In addition, catapults, gummy-guns and modern air-rifles, — illegally — are still used, with lizards and birds being their

4) Barquín & Martín (1982) consider it as possibly introduced. On the other hand, *Gallotia stehlini* has been recently observed in Fuerteventura (Tiscamanita, F. Domínguez, 1984 pers. comm.) associated with cane-bundles brought for use in tomato-fields from Gran Canaria.

5) The same practice is done in Madeira, and called 'balanco' (Sarmiento, A.A., 1984: Vertebrados da Madeira, vol. 1, Funchal., p. 244)

favorite targets. The impact on lizard populations is not very important, but it represents a mischievous behaviour and sets a bad example for young generations.

Lizards may achieve very high densities and cause considerable economic damage to vineyards, tomatoes and, on a lesser scale, to strawberries and market garden plant seedlings. In the past farmers used tobacco to poison lizards (Viera y Clavijo 1886); more recently, after the use of strychnine sulphate was prohibited, farmers turned to killing them with Fosferno (abandoned), Lannate (extensive), and Furadan and Nemacur (starting), which are products commonly used as fungicides or nematocides in the banana plantations.

The biocide is mixed with tomato or ripe bananas and its effect on the reptiles is almost instantaneous because of its very high toxicity⁶. Its effect may easily extend to birds and other animals (including dogs). One of us (López-Jurado) recorded more than 2000 lizards killed during one tomato crop in a plantation of 5–6 hectares in southern Gran Canaria. In the same area he has placed bait (tomato + Lannate) at 15 spots and collected 92 lizards (87 dead) and 1 skink.

Big plastic bags (fertilizer packets) and deep tin-cans (fig. 3) with bait, water, or oil, are hung or fixed on the boundary walls of fields. Lizards fall inside, sometimes in huge quantities. Tello Marquina (1979) collected 50 specimens in a bag, in six hours. Afterwards they are buried or burned (!). When heavy damage occurs⁷ children may be paid for bringing in dead lizards.

These practices, even though not sufficiently controlled, do not threaten any reptilian taxa and may be regarded as a counterbalance of population explosions due to agriculture activity. However, the use of the mentioned biocides has, in many cases, severe adverse consequences on other species, mainly birds (blackbirds, chaff-chaffs, warblers, etc) and their use should be avoided.

The Regional Agriculture Centre (CRIDA XI) did research to find mechanical, biological and nature-compatible methods to control lizard plagues (Tello Marquina, op.c.) but the project was cancelled before any satisfactory results were obtained.

Introduced species

Natural predators of reptiles in the Canary Islands are kestrels (very efficient), buzzard, shrikes, stonecurlews, ravens, choughs, hoopoes (Pérez Padrón, pers. comm.), Barn and Long-eared owls (these latter two mainly on geckos), and shrews. Dogs, cats, rats and, to a lesser scale, hedgehogs also catch reptiles. The

6) Active agents are *Ethylparation* for Fosferno, *Methomyl* for Lannate, *Carbofurane* for Furadan and *Fenamifos* for Nemacur. Lethal oral doses of commercial product for male rats are: 26 mg/Kg, 113–160 mg/Kg, 23–40 mg/Kg and 32–47 mg/Kg respectively (based on data from Worthing, 1979).

7) Economic loss of 30 % of production in tomato, vegetable and ornamental seedlings; 40 % in vineyards in Tenerife (DIARIO DE LAS PALMAS, January 21, 1979: "Lucha contra los lagartos").





Fig. 3. Tin cans used as lizard traps to control populations in cultivated fields. Photo C. Hernández-Rubio.

Fig. 4. Conservation poster (40 x 30 cm), produced by the Union of Insular Councils. Its message means "El Hierro, where there still is Nature". In smaller text information on the status of *Gallotia simonyi* and its importance to science is given.



importance of predation by the latter mentioned species is low on small reptiles but, as postulated by Machado (1985), could have been the reason for the decline of the large lizards. Fortunately, snakes have not been introduced into the archipelago. If this was the case, considerable impact on reptiles by competition and/or predation would be expected. Competition between goats and rabbits with herbivorous reptiles is of importance only at specific sites.

Collecting and trade

The extinction of *Gallotia simonyi* on its only confirmed locality (smaller Roque de Salmor) was due to overcollection, mainly for scientific purposes but also for commercial interest, and occurred in a period of only 42 years. This threat still exists for all species populations living in reduced zones, like islets or rocks offshore, and it is magnified by their taxonomical status. At present many collectors concentrate on endemics or any type of 'singularity' (v. Compte Sart 1977), as keeping reptiles as pets is an increasing hobby in Europe.

Collecting for science or for other reasons is regulated by law, as discussed below, but not everybody applies for a license, and control on remote sites remains difficult.

Trade in Canary reptiles is nowadays strictly regulated and practically prohibited. The Department of Commerce (and in many cases, foreign authorities) will ask for an ICONA⁸ license, before giving the export/import permit. Some authorized consignments of lizards and frogs were sent from Gran Cana-

ria in 1979, but a strong controversy arose in the public media, and therefore the activity was halted.

Superstition

Townspeople in the Canaries are now much more aware of lizards, in particular, due to the Giant lizard of Hierro, which has become a symbol of conservation (fig. 4). In the country, however, reptiles are just noxious animals ('bichos'), and many women do become very afraid of them, placing poison around their houses.

Geckos still suffer from superstition. Some people believe they can spit or urinate in one's eye causing loss of vision. In northern Gomera (Alonso Herrera, pers. comm.) it is seen as a very dangerous animal. They are considered to like milk and therefore they enter nursing babies' mouths as they sleep, producing asphyxia. Babies are watched and parents may place chewed tobacco at their side to prevent the geckos. When they catch one, they put some tobacco in its mouth, causing a rapid death.

On the contrary, skinks are treated with respect. An elderly person told us (Fuerteventura) that people used to put a living skink inside a piece of cane and keep it hanging around their neck, thinking it was good remedy for coughs and asthma. Sarmiento (1948, p. 239) describes a similar practice from Madeira, using *Lacerta dugesi* M. Edw. On Gomera, the skink was placed alive under the hat and carried along to cure headaches, dandruff and to prevent lice. Mr Florencio Fumero, who was a forest ranger at Vilaflor, Tenerife (died May 1982), did live with a skink permanently in his hat; it should be good for the health.

Legal protection

Spain's first protected animal species list was established by the Decree 2573/1973 of October, 5th. It covered 7 mammals, 44 birds and 3 reptiles (chamaleon and two landturtles). Later, by the Royal Decree 3181/1980 of December, 30th, a much stronger regulation was approved, and the list of strictly protected fauna species greatly increased, following the philosophy of the Convention for the Protection of Wildlife and Natural Habitats (Bern Convention).

In all Spanish territory, the hunting, capture, tenancy, traffic, commerce and exportation of the fauna species listed is prohibited, as is the collecting of eggs or young. The preparation and commerce of stuffed specimens is also prohibited.

In addition, regulations governing collection for scientific, conservation, cultural or educational purposes are given. Under special circumstances, temporary capture or hunting of protected species can be authorized to prevent damage to other species, crops, property, health or public safety. The export of living protected animals or their eggs can be allowed only when reintroduction projects in other countries are involved.

The following reptiles⁹⁾ inhabiting the Canary Islands are protected:

'Salamanquesa costera'	<i>Hemidactylus turcicus</i>
'Salamanquesa común'	<i>Tarentola mauritanica</i>
'Salamanquesa canaria'	<i>Tarentola delalandii</i>
'Lagarto de Haría'	<i>Lacerta atlantica</i>
'Lagarto gigante de Hierro'	<i>Lacerta simonyi</i>
'Lagarto canarión'	<i>Lacerta stehlini</i>
'Lagarto tizón'	<i>Lacerta galloti</i>
'Eslizón canario'	<i>Chalcides viridanus</i>

These are the names that appear in the Royal decree. Spain's authorities probably followed nomenclature established by Salvador (1974) in his field guide (except for *T. delalandii*, which is correctly named as "perinquén"). The list is obviously not consistent with recent nomenclatorial changes and has to be updated. It establishes at least some Spanish official common names for the Canarian reptiles. We do not discuss if they are the most appropriate. On Table II

Table II. Local names of reptiles in the Canary Islands.

barbazul	adult male lizard, Palma
bocinegro	adult male lizard, southern Tenerife
bocín	adult male lizard, southern Tenerife
barbolete	adult male lizard, Palma
cacharón	adult male lizard, NW Palma
cacheros	adult male lizard, La Gomera
lagartija	skink, Gran Canaria
lagartija de la Virgen	skink, southern Gomera
lagartija	small or striped lizards, Palma, Tenerife
lagartijo	common lizard, Hierro
lagarto	giant lizard, Hierro
lebrancho	big lizard (applied to other animals too)
lisa	skinks, western islands
lisneja	skink, Fuerteventura
mero (merito)	black male Hierro common lizard
paperos	adult male lizard, Palma
papazules	adult male lizard, Palma
pen(r)inqué	geckos, Gran Canaria
perenquén*	geckos, Tenerife, Hierro and eastern islands
pijina	juvenile lizard, southern Tenerife
pracan	geckos, Gomera
rayada	female lizard, central Gomera
rolisa	skink, northern Gomera
salamandra	geckos, Palma
tijita	juvenile lizard, Tenerife and Palma
tizo (tizón)	adult male lizard, northern and SE Tenerife
verdino (verdín)	adult male lizard, Tenerife

* Following Wölfel (1965, p. 661) this term should be considered as an aboriginal word.

8) ICONA is the Instituto Nacional para la Conservación de la Naturaleza, an autonomous body inside the Department for Agriculture, Fisheries and Food, governed by a Director General.

9) All marine turtles found in Canarian waters are listed, but we are not considering the sea environment in this paper. For further information on turtles, see Brito Hernández (1982) or López Jurado et al. (1983).

a list of local names compiled by us is given. It shows a great variety and lack of uniformity in denominations.

The Bern convention was signed by the Spanish State (19 Sep. 1979) but has not yet been ratified (Aug. 1985). Regarding land reptiles, *Lacerta simonyi* is included in Appendix II of strictly protected species (no exploitation). Under Appendix III (protected fauna species) all reptiles not included in Appendix II are considered. This last category implies that legislative and administrative measures shall be taken to ensure the protection of these wild species, by regulating any type of exploitation.

The Royal decree which stays in force, is related to the Hunting law of 1970, whose orientation is more towards the hunting activity than to protection of wild species. A General law for conservation of nature is being prepared at present. In this context, an international instrument like the Bern convention, becoming domestic law when ratified, would be of great value.

Apart from Spanish law, there is again a special case with *Gallotia simonyi*, which has been listed as an endangered species by the United States Fish and Wildlife Service (Federal register on February 29, 1984) under the provisions of the Endangered species act of 1973. This declares it illegal for any person subject to the jurisdiction of the United States to take, import or export, possess, carry or conduct any commerce with that lizard.

None of the Canarian reptiles are included in the Convention on International Trade in Endangered Species of Wild Fauna and Flora (C.I.T.E.S.).

Habitat protection overview

The Canary Islands have a high number of protected areas, in accordance with the richness and singularity of its biota. We consider it of interest to give a broad view of these areas (see fig. 6) and the protection of habitat for reptiles.

TEIDE NATIONAL PARK: Established by decree in 1954 and reestablished by law in 1981; volcanic region (fig. 5) with high mountain shrub vegetation in the middle and upper parts of the island of Tenerife (between 2,000 and 3,717 m altitude). Extension 13,500 hectares; four permanent rangers in the field. *Gallotia galloti eisentrauti* is abundant; few geckos and skinks have been seen.

CALDERA DE TABURIENTE NATIONAL PARK in La Palma, established by decree in 1954 and reestablished by law in 1981. Deep depression ranging from 300 to 2,423 m altitude. Covers 4690 hectares mainly of Canary pine forest. Two permanent rangers in the field. Lizards in general uncommon; geckos scarce.

GARAJONAY NATIONAL PARK: Established by law in 1981. Central forested ("laurisilva") region of Gomera (between 700 and 1,487 m altitude). Extension 3,984 hectares; nine permanent rangers in the field. Some lizards, skinks and geckos in the southern part of the park, locally common.

TIMANFAYA NATIONAL PARK: Established by law in 1974 and reestablished in 1981. Covers 5,107 hectares of historic lava flows on Lanzarote. Two permanent rangers in the field. Lizards restricted to kipukas.

EL CANAL Y LOS TILES MAN AND BIOSPHERE RESERVE: Declared by decree in 1984 in the island of La Palma. Sector of deep ravine between 250 and 1,400 m



Fig. 5. Habitat of *Gallotia galloti eisentrauti*, a common species in Teide National Park. Photo A. Machado.

altitude with 511 hectares of laurisilva and pine forest; property of ICONA; one permanent ranger in the field. Lizards scarce, only in upper parts.

DUNAS DE CORRALEJO E ISLA DE LOBOS NATURAL PARK: Established by decree in 1983. Embraces 2,482 hectares of a dune system and a small volcanic island with xerophytic vegetation. Highest point at 127 m; one permanent ranger in the field. Lizards and geckos common in Lobos, but less common in the dunes at Corralejo.

MASPALOMAS NATURE SITE: Has a provisional protection status due to Royal decree 1741 of June 25, 1982. No permanent ranger in the field. 328 hectares of coastal dunes with palms and *Tamarix* vegetation. Very interesting lizard population; skinks and geckos occasional.

The site where *Gallotia* aff. *simonyi* survives in Hierro falls inside a public forest — under ICONA's tutelage — and access has been prohibited since 1975. Permanent vigilance from outside is maintained. However, a project for the declaration of the whole zone including the Roques de Salmor is under administrative process and ready for public hearing. This future Strict Reserve ("Reserva Integral") will probably join the Biogenetic Reserves Network of the Council of Europe.

Action for the future

The situation of the Canarian reptile fauna is not as dangerous as that of other vertebrate groups, or even plants. Actions will be taken in the immediate future to try to recover Hierro's Giant lizard from its extremely endangered status, and its biotope is in the process of being declared a Strict Nature Reserve. Legislation will be actualized soon, and natural habitat protection is entering the general process of land use planning¹⁰).

10) The Land Use Act, presently under consideration, incorporates "Planes Especiales de Protección y Catalogación de Espacios Naturales" (= Special Plans for the Protection and Cataloguing of Naturales Sites) for all islands except Gran Canaria.

Nevertheless, even being optimistic, there are still some other points that deserve attention.

— Educational material on reptiles' role in nature should be prepared not only for school children, but in particular for people in the country.

— New methods for controlling lizard plagues must be applied, stopping the actual and illegal use of strong biocides.

— Any localized reptile population that is somehow singular, should be studied, delimited and, if necessary, protection of its habitat should be sought.

— A Strict Nature Reserve embracing the non-inhabited rocks and islets north of Lanzarote (Roque del Este, Roque del Infierno, Montaña Clara and Alegranza) should be promoted. The same is valid for both Roques de Anaga, off Tenerife.

— Collection of reptiles in small rocks and islets should be strictly controlled to prevent decimation of vulnerable and rare populations.

— A solution for pet-keepers' interests could be found by regulating some form of farming their product, or taking advantage of special cases (plagues, etc).

Zusammenfassung

In dieser Arbeit wird der Schutzstatus der landlebenden Reptilien der Kanarischen Inseln analysiert, basierend auf den Freilandbefahrungen der Verfasser. Zur Beurteilung der Arten werden IUCN Kategorien verwendet. Kritische Arten sowie die Reptilienpopulationen beeinflussenden Faktoren werden erörtert. Letztere beinhalten Habitatveränderungen, das Töten, Abfangen oder Vergiften aus wirtschaftlichen Gründen oder aus Aberglauben, das Sammeln und den Einfluß eingeführter Arten. *Gallotia* aff. *simonyi* ist das einzige als „bedroht“ eingestufte Taxon, allerdings werden die meisten Reptilienpopulationen auf Felsen oder Inselchen als „gefährdet“ angesehen. Bisher vorliegende Studien über kanarische Echsen werden kommentiert, und eine Bibliographie der bisher veröffentlichten Arbeiten wird als Appendix beigelegt. Außerdem werden gesetzliche Bestimmungen, national und international, diskutiert. Soweit es den Schutz der Reptilien betrifft, werden die kanarischen Naturparke und Reservate erwähnt. Aktuelle Schutzprogramme werden kurz erläutert und Vorschläge für zukünftige Aktionen gemacht.

Resumen

En este trabajo se presenta un análisis del estado de conservación de la fauna de reptiles terrestres de las Islas Canarias (España) basado en la experiencia de campo de los autores. Se utilizan las categorías de conservación de la UICN y se discuten las especies críticas y los principales factores de amenaza que afectan a las poblaciones. Entre éstos se tratan las alteraciones del habitat, caza, trampeo y biocidas, especies introducidas, coleccionismo, comercio y supersticiones. El Lagarto gigante de El Hierro (*Gallotia* aff. *simonyi*) es el único taxon que se considera amenazado de extinción, y la mayoría de las poblaciones de reptiles de roques e islotes figuran bajo la categoría de vulnerables o raros. Se comentan en general los estudios existentes sobre reptiles de Canarias y se aporta una bibliografía prácticamente completa sobre Herpetología Canaria. Asimismo se incluye

una lista de nombres locales de los reptiles terrestres y se discuten las disposiciones legales, tanto nacionales como internacionales, que afectan a su conservación. También se expone una panorámica de la protección de hábitat para reptiles obtenida en la red canaria de parques y reservas naturales. Por último se comentan sucintamente los programas de conservación actualmente en marcha y se aportan algunas sugerencias para acciones en el futuro.




Fig. 6. National Parks and Reserves in the Canary Islands.

References

(only non-herpetological, others included in the Appendix)

- Castro Alfin, D. (1983): Historia de las Islas Canarias. De la prehistoria al descubrimiento. — Madrid: Editora Nacional, Cultura y Sociedad, 255 pp.
- Honegger, R.E. (1987): Amphibiens et reptiles menacés en Europe. — Strasbourg: Conseil de l'Europe, Collection sauvegarde de la Nature 15, 127 pp.
- Machado, A. (1976): Introduction to a faunal study of the Canary Islands' laurisilva, with special reference to the groundbeetles. XII, pp. 347—411 in G. Kunkel: Biogeography and ecology in the Canary Islands. Monographia Biologicae 30, The Hague.
- Sarmento, A.A. (1984): Vertebrados da Madeira. 1. Volume (Mamíferos — Aves — Répteis — Batráquios). — Funchal: (2 ed.), Junta Geral do Distrito Autónomo do Funchal, 317 pp.
- Vølsøe, H. (1951): The breeding birds of the Canary Islands. I. Introduction and synopsis of the species. — Vidensk. Medd. fra Dansk. naturh. Foren. 113: 1—153.
- (1955): The breeding birds of the Canary Islands. II. Origin and history of the Canarian avifauna. — Vidensk. Medd. fra Dansk. naturh. Foren. 111: 117—178.
- Wölfel, D.J. (1965): Monumenta Linguae Canariae. Die Kanarischen Sprachdenkmäler. Eine Studie zur Vor- und Frühgeschichte Weißafrikas. — Graz: Akademische Druck u. Verlagsanstalt, 928 pp., 8 Karten.
- Worthing, Ch., ed. (1979): The pesticide manual. — London: British Crop Protection Council, 655 pp.

Antonio Machado, Instituto Nacional para la Conservación de Naturaleza, Aptdo 619, Santa Cruz de Tenerife. — Luis Felipe López-Jurado, Departamento de Biología, Colegio Universitario de Las Palmas. — Aurelio Martín, Departamento de Zoología, Universidad de La Laguna, Islas Canarias.

Appendix

A bibliography of the reptile and amphibian fauna of the Canary Islands

(compiled with the assistance of J. Díaz de Castro, Las Palmas, and supplemented by the editor).

- Acosta Martínez, P. & M. Pellicer Catalán (1976): Excavaciones arqueológicas en la Cueva de la Arena (Barranco Hondo, Tenerife). — Ann. Est. Atlant. 22: 122—184.
- Ahl, E. (1925): Über eine ausgestorbene Riesenschildkröte der Insel Teneriffa. — Z. dt. geol. Ges. 77 A (4): 575—580.
- (1928): Über einen weiteren Fund von *Testudo burchardi* E. Ahl aus Teneriffa. — 2, pp. 445—447 in Burchard, O. & Ahl, E. „Neue Funde von Riesen-Landschildkröten aus Teneriffa. — Z. dt. geol. Ges. 79 A (4), 1927: 439—447.
- Altman, W. (1985): Über die Haltung und Fortpflanzung des Kanarenskinkes, *Chalcides viridianus* (Gravenhorst, 1851). — Sauria 7 (1): 25—26.
- Arnold, E.N. (1973): Relationships of the Palaearctic lizards assigned to the genera *Lacerta*, *Algyroides* and *Psammodromus* (Reptilia: Lacertidae). — Bull. Br. Mus. nat. Hist. (Zool). 25 (8): 289—366.
- Báez, M. (1977): Los reptiles de las Islas Canarias. — Aguayro 93: 3pp., 1 fig., 2 mapas.
- (1979): Sobre la presencia de *Hemidactylus turcicus* en Tenerife (Islas Canarias) (Rept., Gekkonidae). — Bol. Est. Centr. Ecología 8 (15): 77—78.
- (1983): Consideraciones sobre las características zoogeográficas de la fauna de Canarias. — pp. 21—70 in: Instituto de Estudios Canarios: 50 Aniversario (1932—1982). — I. Ciencias... — Santa Cruz de Tenerife: Instituto de Estudios Canarios, 1982.

- (1983): Poblamiento animal de las Islas Canarias. — pp. 25–84 in Báez, M. et al.: Canarias, origen y poblamiento. — S/C de Tenerife: Circulo de Estudios Sociales de Canarias.
- (1984): Anfibios y reptiles. — pp. 259–273 in: Fauna (marina y terrestre) del Archipiélago Canario. — Edirca, S.L. (Las Palmas de Gran Canaria).
- (1985): Datos sobre la termorregulación de *Gallotia galloti* (Sauria, Lacertidae). — Bonn. zool. Beitr. 36: 557–562.
- & T. Bravo (1983): Sobre la presencia de *Gallotia simonyi* (Reptilia, Lacertidae) en el Roque de Fuera (Tenerife). — Vieraia 12, (1–2): 339–348.
- , J.J. Bacallado & A. Martín (1984): Los reptiles de Canarias: importancia científica y problemática conservacionista. — II Reunión Iberoamer. Cons. Zool. Vert., pp. 84–90.
- & R.S. Thorpe (1985): Microevolution of the lizard *Gallotia galloti* within the island of Tenerife. — Bonn. zool. Beitr. 36: 513–515.
- Barquín Diez, E. & W. Wildpret de la Torre (1975): Diseminación de plantas canarias. Datos iniciales. — Vieraia 5 (1–2): 38–60.
- Barquín, J. & A. Martín (1982): Sobre la presencia de *Gallotia* (= *Lacerta*) *atlantica* (Peters y Doria, 1882) en Gran Canaria (Rept., Lacertidae). — Doñana, Acta Vertebrata 9: 377–380.
- Blas Aritio, L. (1976): El libro rojo de la fauna española. — Madrid: Incafo, 191 pp.
- Bedriaga, J.V. (1886): Beiträge zur Kenntnis der Lacertidenfamilie. — Abh. senckenberg. naturforsch. Ges., 14, 1888 [1886]: 427 pp., 1 Tafel.
- Bertin, L. (1945): La faune herpétologique des Iles Atlantiques dans ses rapports avec la paléogéographie de ces archipels. — C. R. somm. Séanc. Soc. Biogéogr. 20 (167/168), 1943: 1–3.
- (1946): Le Peuplement des îles Atlantides en vertébrés hétérothermes. — pp. 87–107 in: Contribution a l'étude du peuplement des îles Atlantides. — Société de Biogéographie, Mem. 8, 1 cart., 3 tabl.
- Bings, W. (1980): Herpetologische Studien auf Teneriffa (Kanarische Inseln). — Salamandra 16 (4): 203–214.
- (1985): Zur früheren Verbreitung von *Gallotia simonyi* auf Hierro, mit Vorschlägen zur Wiederansiedlung. — Bonn. zool. Beitr. 36: 417–427.
- Bischoff, W. (1969): Lacertenbastarde. — Aquar. Terrar. 8: 262–265.
- (1971): *Lacerta g. galloti* Duméril & Bibron 1839, die Eidechse von Teneriffa. — Aquar. Terrar. 18 (9): 308–311.
- (1973): Einiges zur Winterruhe bei Lacerten. — Aquar. Terrar. 20: 350–351.
- (1973): Über eine bemerkenswerte Verhaltensweise von *Lacerta simonyi stehlini*. — Aquarien und Terrarien 20: 278–279.
- (1974): Beobachtungen bei der Pflege von *Lacerta simonyi stehlini* (Sauria, Lacertidae). — Salamandra 10 (3/4): 93–103.
- (1981): Freiland- und Terrarienbeobachtungen an der Omaneidechse, *Lacerta jayakari* BOULENGER, 1887 (Reptilia: Sauria: Lacertidae). — Zeitsch. Kölner Zoo 24 (4): 135–143.
- (1982): Die innerartliche Gliederung von *Gallotia galloti* (Duméril & Bibron 1839) (Reptilia: Sauria, Lacertidae) auf Teneriffa, Kanarische Inseln. — Bonn. zool. Beitr. 33 (2–4): 363–382.
- (1984): Bemerkungen über die endemische Echsenfauna der Kanarischen Inseln. — Sauria 6 (2): 5–11.
- (1985): Bemerkungen zur innerartlichen Variabilität von *Gallotia atlantica* (Peters & Doria, 1882) (Lacertidae). — Bonn. zool. Beitr. 36: 489–506.
- (1985): Die Herpetofauna der Kanarischen Inseln I–VI. — herpetofauna (Weinstadt) 7 (34): 11–22; (35): 27–34; (36): 13–21; (37): 15–24; (38): — ; (39): —.
- (1985): Vorläufige Beschreibung der Eidechse der Kanareninsel Fuerteventura als *Gallotia atlantica mahoratae* ssp. n. (Sauria: Lacertidae). — Salamandra 21 (1): 46–54.
- , H.K. Nettmann, & S. Rykena (1979): Ergebnisse einer herpetologischen Exkursion nach Hierro, Kanarische Inseln. — Salamandra 15 (3): 158–175.
- & P. v. d. Elzen (1982): *Lacerta jayakari* Boulenger. — Suppl. Rev. franç. Aquariol. Herpetol. 2 (82), Fiche 234.
- Boettger, C.R. & L. Müller (1914): Preliminary notes on the local races of some canarian lizards. — Ann. Mag. nat. Hist. (8) 14: 67–78.
- Böhme, W. (1971): Über das Stachelepithel am Hemipenis lacertider Eidechsen und seine systematische Bedeutung. — Z. zool. Syst. Evolforsch. 9: 187–223.
- & W. Bings (1975): Zur Frage des Überlebens von *Lacerta s. simonyi* Steindachner (Sauria: Lacertidae). — Salamandra 11 (1): 39–46.

- & — (1977): Nachträge zur Kenntnis der kanarischen Rieseneidechsen (*Lacerta simonyi*-Gruppe) (Reptilia, Sauria, Lacertidae). — *Salamandra* 13 (2): 105–111.
- & W. Bischoff (1976): Das Paarungsverhalten der kanarischen Eidechsen (Sauria, Lacertidae) als systematisches Merkmal. — *Salamandra* 12 (3): 109–119.
- , H.-K. Nettmann, S. Rykena, & J. Freundlich (1981): Nachweis von *Gallotia simonyi* (Steindachner, 1889) (Reptilia: Lacertidae) aus einer frühmittelalterlichen Fundschicht auf Hierro, Kanarische Inseln. — *Bonn. zool. Beitr.* 32 (1–2): 157–166.
- , R. Hutterer & W. Bings (1985): Die Stimme der Lacertidae, speziell der Kanareneidechsen (Reptilia: Sauria). — *Bonn. zool. Beitr.* 36: 337–354.
- Bory de St-Vincent, J.B.G.M. (1803): *Essais sur les Isles Fortunées et l'antique Atlantide, ou précis de l'histoire générale de l'Archipel des Canaries*. — Paris: Baudouin, Germinal an XI [i-viii], + 522 pp., 7 pls.
- Böttger, O. (1874): Reptilien von Marocco und von den Canarischen Inseln. II. Übersicht der von dem Herrn Dr. C. von Fritsch i. J. 1863, von den Herren Dr. Fr. Noll und Dr. Grenacher im Jahre 1871 und von den Herren Dr. C. von Fritsch und Dr. J.J. Rein im Jahre 1872 auf den Canaren gesammelten Reptilien. — *Abh. senckenb. naturforsch. Ges.* 9, 1873: 121–191.
- (1883): Katalog der Reptilien-Sammlung im Museum der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main. — Frankfurt am Main: Knauer, X, 140 S.
- Boulenger, G.A. (1885): *Catalogue of the lizards in the British Museum (Natural History)*. — Second edition. — Volume 1: Gekkonidae, Eublepharidae, Uroplatidae, Pygopodidae, Agamidae. — London, British Museum (Natural History), xii, 436 pp., 32 pl.
- (1887): *Catalogue of the lizards in the British Museum (Natural History)*, Ed. 2, Vol. 3. — London, 575 pp., 39 pl.
- (1890): On the varieties of *Chalcides ocellatus*, Forsk. — *Ann. Mag. nat. Hist.* (6) 5: 444–445.
- (1891): On Simony's Lizard, *Lacerta simonyi*. — *Proc. zool. Soc. London*, 1891 (13): 201–202.
- (1892): On newly discovered East-African chameleons, with remarks on some other reptiles described by Dr. Steindachner. — *Ann. Mag. nat. Hist.* 9 (6): 72–74.
- (1920): *Monograph of the Lacertidae. Volume I*. — London: Trust. Brit. Museum, x, 352 pp.
- Boulenger, E.G. (1920): On some lizards of the genus *Chalcides*. — *Proc. zool. Soc. Lond.*, 1920 (1): 77–83.
- Bravo, T. (1953): *Lacerta maxima* n. sp., de la fauna continental extinguida en el Pleistoceno de las Islas Canarias. — *Estudios geol. Inst. Invest. geol. Lucas Mallada* 9 (17): 7–34.
- (1954): *Geografía General de las Islas Canarias. Tomo I*. — Santa Cruz de Tenerife: Goya ediciones, 410 pp.
- (1964): *Geografía General de las Islas Canarias. Tomo II. Gran Canaria, Fuerteventura, Lanzarote*. — Santa Cruz de Tenerife: Goya Ediciones, 592 pp.
- (1978): Yacimientos de Vertebrados fósiles en Canarias. — *An. Inst. Est. Can.* 16–20: 42–44.
- Brito Hernández, A. & T. Cruz Simó (1982): Tortugas marinas en Canarias. — *Viera y Clavaja* 11 (1/2), 1981: 319.
- Brongersma, L.D. (1968): Notes upon some turtles from the Canary Islands and from Madeira. — *Proc. K. ned. Akad. Wet.* 71-C (2): 128–136.
- Burchard, O. (1928): Über die Auffindung fossiler Knochenreste einer Riesen-Landschildkröte auf Teneriffa. — 1, pp. 439–444 in Burchard, O. & Ahl, E.: *Neue Funde von Riesen-Landschildkröten aus Teneriffa*. — *Z. dt. geol. Ges.* 79 A (4), 1927: 439–447.
- (1934): *Testudo Burchardii*, E. Ahl. El primer gran fósil descubierto en Canarias. — *Inst. Estud. canarios (Monogr.)* 1, 15 pp.
- Busto y Blanco, F. del (1864): *Topografía médica de las Islas Canarias*. — Sevilla: La Andalucía, xi, 532 pp.
- Calle, A. de la (1975): *Estudio histológico a niveles óptico y electrónico de las fibras nerviosas y células corticales del cerebro de L. galloti*. — Doctoral Thesis, Univ. La Laguna, Tenerife, Spain.
- Camp, C.L. (1923): Classification of the lizards. — *Bull. Amer. Mus. nat. Hist.* 48 (11): 289–481.
- Cano, J., M. Báez, L.F. López-Jurado, & G. Ortega (1984): Caryotype structure in the lizard *Gallotia galloti* in the Canary Islands. — *J. Herpet.* 18 (3): 344–346.
- Carnero, A. & F. Pérez Padrón (1977): Los lagartos de las Islas Canarias. — *Vida silv.* 24: 248–253.
- Compte Sart, A. (1977): La captura de especies raras y la conservación de biotopos restringidos. — *Asturnatura* 3 (3/4), 1975–1976: 87.
- Cott, H.B. (1934): On the ecology of *Hyla arborea* var. *meridionalis* in Gran Canaria, with special reference to predatory habits considered in relation to the protective adaptations of insects. — *Proc. zool. Soc. Lond.* 1934 (2): 311–331.

- Cyren, O. (1933): Biologiskt från Madeira och Kanarieöarna. — Fauna och Flora 1933 (1), 16 pp.
- (1934): Zur Kenntnis der Lacertiden der Iberischen Halbinsel und Makaronesiens. — Göteborgs K. Vetensk. Vitter. Samh. Handl. (B) 4 (1): 1–64.
- De Vera, L.M. (1984): Respuestas fisiológicas a la temperatura en reptiles (*Gallotia galloti*). — Tesis doctoral. Facultad de Medicina. Universidad de La Laguna, Tenerife.
- Díaz Reyes, G. & N. González Henríquez (1982): Origen y evolución de la flora y fauna de las Islas Canarias. — Aguayro 142: 16–22.
- Dubois, A. (1984): Les specimens-Types de *Gallotia galloti* (Oudart, 1839) (Reptiles, Sauriens). — Bull. Soc. Linn. Lyon 53 (1): 27–30.
- Duméril, A.M.C. & G. Bibron (1836): Erpétologie générale ou Histoire naturelle complète des Reptiles. — Tome troisième. — Paris: Roret iv, 517 pp., errata et emendanda, planches 25–36.
- & — (1839): Erpétologie générale ou Histoire naturelle complète des Reptiles. 5. — Paris.
- Evers, A., K. Klemmer, I. Müller-Liebenau, P. Ohm, R. Remane, P. Rothe & R. Zur Strassen (1970): Erforschung der mittelatlantischen Inseln. — Umschau 70 (6): 170–176.
- Fejervary, G.J. von (1914): Über kanarische Lacerten. — Verh. zool.-bot. Ges. Wien 64 (9/10): 320–333.
- (1914): Über die Entwicklung des Farbenkleides bei den Lacerten. (Gedanken zu einer phylogontog. Studie). — Zool. Anz. 43: 533–537.
- Felipe Hernández, J. (1976): Estudio de la corteza cerebral en periodos prenatales del desarrollo embrionario de *L. galloti*. — Doctoral Thesis, Univ. La Laguna, Tenerife, Spain.
- Fernández, J.M. (1966): La fauna entomológica canaria y sus orígenes. — Act. V. Cong. Panafri. Prehist. et l'étude du Quatern., Publ. S. I. A. Excmo. Cabildo Insular 6: 61–68.
- Fernández Ruiz, B., J.M. Pérez Batista & C. Dobao (1978): Ultraestructura de la nefrona del *Chalcides viridanus* (Scincidae). I. El corpúsculo renal. — Morf. Norm. y Patol. secc A. 2: 413–423.
- , —, — & J. García Miranda (1978): Ultraestructura de la nefrona del *Chalcides viridanus*. II. El componente tubular. — Morf. Norm. y Patol. secc A. 2: 425–441.
- Fritsch, K. von (1867): Reisebilder von den Canarischen Inseln. — Petermanns Geograph. Mitt., Erg. Bd. 5 (22): 1–44.
- (1870): Über die ostatlantischen Inselgruppen. — Ber. senckenb. naturf. Ges. 1870: 72–113.
- García Cruz, C.M. (1978): Características anatómicas y biométricas de *Lacerta maxima* Bravo, 1953, (Reptilia: Lacertidae). — Memoria de Licenciatura, Universidad de La Laguna, Tenerife, 192 pp. [not published]
- & A. Marrero Rodríguez (1979): Sobre la distribución geográfica de los yacimientos de vertebrados fósiles de las Islas Canarias. — Vieraea 8 (1) 1978: 95–106.
- García Cruz, G., C.M. Marrero & A. Hernández (1980): Breves consideraciones sobre la filogenia y evolución del grupo *Simonyi* (Reptilia: Lacertidae). — Cuad. Cienc. biol. 6–7 (1977–1978): 55–57.
- García Verdugo, J.M., P. Berbel Navarro, J. Regidor García & C. Lopez Garcia (1983): Ultrastructure of neuronal cell bodies in the dorso-medial cortex of *Lacerta galloti*. — J. Hirnforsch. 24: 307–314.
- , —, — & — (1983): Ultrastructure of neuronal cell bodies in the dorsal cortex of *Lacerta galloti*. — J. Hirnforsch. 24: 485–494.
- , —, — & — (1984): Ultrastructure of neuronal cell bodies in the medial cortex of *Lacerta galloti*. — J. Hirnforsch. 25: 187–196.
- Gasc, P.-P. (1971): Les variations cumulatives dans la région présacrée des sauriens. Application a la reconstitution de *Lacerta goliath* Mertens. — Annals Paléont. (Vertébr.) 57 (1): 131–156.
- Gervais, P. (1839): Zoologie: Animaux d'Afrique. — Extraits P. v. Soc. philom. Paris, 1839: 4.
- (1844): Reptiles des Iles Canaries, recueillis par MM. P. Barker Webb et S. Berthelot, et décrits par M. Paul Gervais. — pp. 1–5 in Webb, P.B. & S. Berthelot: Histoire naturelle des Iles Canaries. — Paris: Bethune.
- Godoy Arcaya, R. (1982): La Isla de El Hierro. Guías raras y completas de territorios y habitantes de España. — Madrid: (E.C.C.E.) Secretaría de Estado de Turismo, Mint. Transp., Tur. y Comunicaciones, 206 pp.
- Gonzalez-Lamas, Z., R.H. Lopez Orge & A.M. Lamas (1980): Study of hemoglobin and A soluble heart peroxidase in *Lacerta galloti* and *L. stehlini* by isoelectric focusing. — In Radola, B. J. (Ed.): Electrophoresis'79. — Berlin; New York: Walter de Gruyter, pp. 841–845.
- , A. de Armas, P. Betancor & E.M. Hevia (1981) Isozymes of superoxide dismutase in liver from *Lacerta stehlini*. — In Allen, A. (Ed.): Electrophoresis'81. — Berlin; New York: Walter de Gruyter, pp. 703–707.

- Gravenhorst, J.L.C. (1851): Über die im Zoologischen Museum der Universität Breslau befindlichen Wirtelschleichen (Pseudosaura), Krüppelfüssler (Brachypoda) und einige andere denselben verwandte Reptilien aus den Zünften der Schleichen und Dickzüngler. — *Nova Acta Acad. Caesar. Leop. Carol.* 23 (1): 291–394.
- Groh, K. (1985): Landschnecken aus quartären Wirbeltierfundstellen der Kanarischen Inseln (Gastropoda). — *Bonn. zool. Beitr.* 36:395–415.
- Groombridge, B. (1982): The IUCN Amphibia-Reptilia Red Data Book. Part 1. Testudines, Crocodylia, Rhynchocephalia. — Gland: IUCN, WWF & UNEP.
- Hartung, G. (1857): Die geologischen Verhältnisse der Inseln Lanzarote und Fuerteventura. — *Neue Denkschr. Allg. schweiz. Ges. ges. Naturw.* 15 (4): iv, 1–163.
- Hecke, A. van (1975): Genezing van een voedselweigerende, ontkalkte *Lacerta galloti*. — *Lacerta* 34 (1): 12–13.
- Heselhaus, R. (1981): Der Lebensraum der Kanarischen Rieseneidechse, *Lacerta simonyi stehlini*. — *Aquar.- u. Terrar.-Z.* 34 (2): 66–67.
- Honegger, R.E. (1975–79): Red Data Book 3, Amphibians and Reptiles. — Gland, IUCN.
- (1978): Amphibiens et reptiles menacés en Europe. — Strasbourg: Conseil de l'Europe, Coll. Sauvegarde de la Nature 15, 127 pp.
- Hutterer, R. (1979): Occurrence of the European free-tailed bat *Tadarida teniotis* on Hierro, Canary Islands. — *Afr. Small Mammal Newsl.* 3: 6–7.
- (1985): Neue Funde von Rieseneidechsen (Lacertidae) auf der Insel Gomera. — *Bonn. zool. Beitr.* 36: 365–394.
- Joger, U. (1984): Morphologische und biochemisch-immunologische Untersuchungen zur Systematik und Evolution der Gattung *Tarentola* (Reptilia: Gekkonidae). — *Zool. Jb. Anat.* 112: 137–256.
- (1984): Taxonomische Revision der Gattung *Tarentola* (Reptilia: Gekkonidae). — *Bonn. zool. Beitr.* 35 (1–3): 129–174.
- (1984): Die Radiation der Gattung *Tarentola* in Makaronesien (Reptilia: Sauria: Gekkonidae). — *Cour. Forsch.-Inst. Senckenberg* 71: 91–111.
- (1985): Biochemical and immunological data on the systematic position of the endemic gecko of the Selvages Islands. — *Bonn. zool. Beitr.* 36: 307–314.
- & W. Bischoff (1983): Zwei neue Taxa der Gattung *Tarentola* (Reptilia: Sauria: Gekkonidae) von den Kanarischen Inseln. — *Bonn. zool. Beitr.* 34 (4): 459–468.
- Kämmer, F. (1982): Beiträge zu einer kritischen Interpretation der rezenten und fossilen Gefäßpflanzenflora und Wirbeltierfauna der Azoren, des Madeira-Archipels, der Ilhas Selvagens, der Kanarischen Inseln und der Kapverdischen Inseln, mit einem Ausblick auf Probleme des Artenschwundes in Makaronesien. — Freiburg im Breisgau: 179 pp.
- Klemmer, K. (1971): Die Echten Eidechsen. — pp. 285–307 in Grzimek, G. (Herausg.): Enzyklopädie des Tierreich. Sechster Band: Kriechtiere; XII. Kapitel. — Zürich: Kindler.
- (1976): The Amphibia and Reptilia of the Canary Islands. — XV, pp. 433–456 in Kunkel, G. (ed.): Biogeography and Ecology in the Canary Islands. — Monographia biologica 30; The Hague: W. Junk Publishers.
- Koenig, A. (1890): Ornithologische Forschungsergebnisse einer Reise nach Madeira und den kanarischen Inseln. — *Journ. Ornith.* 38 (191–192): 257–488.
- (1890): Ueber Teneriffa in zoologischer Beziehung. — *Sb. niederrhein. Ges. Nat. Heilkde. Bonn* A: 3–13, 20–28.
- Kraepelin, K. (1895): Zoologische Ergebnisse einer Frühjahrs-Excursion nach Madeira und den Canarischen Inseln. — *Verh. naturw. Ver. Hamb.* 2 (2), 1894, 1895: 6–17.
- Kreffft, G. (1949): Beobachtungen an kanarischen Inseleidechsen. — *Wschr. Aquar. Terrar. Kde.* 43: 17–21, 41–42, 66–68, 93–94, 114–116.
- (1950): Beiträge zur Kenntnis der kanarischen Echten Eidechsenfauna. — *Zool. Anz.* 145 (Suppl.): 426–444.
- Ledru, A.-P. (1810): Voyage aux îles de Ténériffe, La Trinité, Saint-Thomas, Sainte-Croix et Porto Rico, exécuté par ordre du gouvernement français, depuis le 30 septembre jusqu'au 7 juin 1798, ... — Tome premier. — Paris: Bertrand, 4 p., i–xlvii, 315 pp, 1 carte.
- Lehrs, P. (1909): Studien über Abstammung und Ausbreitung in den Formenkreisen der Gattung *Lacerta* und ihrer Verwandten. — *Zool. Jb. (Syst.)* 28 (1): 81–120.
- (1914): [*Lacerta caesaris*, n. sp.] — *Abst. Proc. zool. Soc. London* 134: 41.
- (1914): Description of a new lizard from the Canary Islands. — *Proc. zool. Soc. London* 1914 (3): 681–684.
- López García, C., A. Molowny & J. Pérez Clausell (1983): Volumetric and densitometric

- study in the cerebral cortex and septum of a lizard (*Iacerta galloti*) using the Timm method. — *Neurosc. Lett.* 40: 13–18.
- , E. Soriano, A. Molowny, J.M. Verdugo, P.J. Berbel & J. Regidor (1983): The Timm positive system of axonic terminals of the cerebral cortex of *Lacerta*. — pp. 137–148 in Grisolina, S. et al. (eds.): *Ramón y Cajal's Contribution to the Neurosciences*. — Amsterdam: Elsevier Science Publishers.
- López-Jurado, L.F. (1985): Los reptiles fósiles de la Isla de Gran Canaria (Islas Canarias). — *Bonn. zool. Beitr.* 36: 355–364.
- & M. Báez (1985): La variación de *Chalcides sexlineatus* en la Isla de Gran Canaria (Islas Canarias). — *Bonn. zool. Beitr.* 36: 315–336.
- & S. Martínez González (1983): Las tortugas en Canarias. — *Aguayro* 147: 29–31.
- , M. Ruiz & L. Dos Santos (1980): Sobre la reproducción del eslizón canario (*Chalcides viridanus*) en la isla de Gran Canaria. — *Doñana Acta vert.* 6 (2) 1979: 225–227.
- Lorenzo Hernández, A. (1975): Estudio histológico a nivel de la microscopía óptica y con el M.E. de los plexos coroideos de *L. galloti*. — Doctoral Thesis, Univ. La Laguna, Tenerife, Spain.
- Loveridge, A. (1947): Revision of the African lizards of the family Gekkonidae. — *Bull. Mus. comp. Zool. Harv.* 98 (1): 1–461.
- Lutz, D. & W. Mayer (1984): Albumin-immunologische und proteinelektrophoretische Untersuchungen zur systematischen Stellung von *Lacerta lepida* Daudin und *Lacerta princeps* Blandford (Sauria, Lacertidae). — *Zool. Anz.* 212 (1–2): 95–104.
- Macau Vilar, F. (1958): Contribución al estudio del Mioceno Canario. — *Bol. R. Soc. Esp. Hist. Nat.* 56: 477–486.
- (1960): Contribución al estudio del Cuaternario de Gran Canaria. — *Ann. estud. atlant.* 6: 117–132.
- Mac Gillivray, J. (1852): Visit to Teneriffe, and ascent to the Peak of Teyde. — *Zoologist* 10: 3441–3449.
- Machado, A. (1985): Sinópsis del plan de recuperación del Lagarto gigante del Hierro. — *Bonn. zool. Beitr.* 36: 471–480.
- (1985): Hypothesis on the reasons for the decline of the large lizards in the Canary Islands. — *Bonn. zool. Beitr.* 36: 563–575.
- (1985): New data concerning the Hierro Giant lizard and the Lizard of Salmor (Canary Islands). — *Bonn. zool. Beitr.* 36: 429–470.
- , L.F. López-Jurado & A. Martín (1985): Conservation status of reptiles in the Canary Islands. — *Bonn. zool. Beitr.* 36: 586–606.
- Maffiote y la Roche, P. (1916): *Historia de las Islas Canarias*. — Edición Ilustrada. — A. J. Benítez Editor. Santa Cruz de Tenerife, 528 pp.
- Manrique y Saavedra, A.M. (1873): *Elementos de Geografía é Historia natural de las Islas Canarias...* — Las Palmas: Ramírez, 98 pp.
- Marrero Rodríguez, A. & C.M. García Cruz (1978): Nuevo yacimiento de restos subfósiles de dos vertebrados extintos de la Isla de Tenerife (Canarias), *Lacerta maxima* Bravo, 1953 y *Canaryomis bravoii* Crus. et Pet., 1964. — *Vieraea* 7 (2) 1977: 165–174.
- Martín, A. (1985): Los lagartos de los roques del norte de Tenerife. — *Bonn. zool. Beitr.* 36: 517–528.
- Martín Esquivel, J.L. (1982): El ecosistema cavernícola en los tubos volcánicos del Valle de Güimar. — Santa Cruz de Tenerife: F.T.C.E., Mon. 1, 68 pp.
- Martín Trujillo, J.M. (1976): Análisis histológico del pallium de Reptiles Squamatas (*L. galloti*, *Ch. viridanus* y *T. delalandii*). — Doctoral Thesis, Univ. La Laguna, Tenerife, Spain.
- Martínez Guijarro, F.J., P.J. Berbel, A. Molowny & C. López García: Apical dendritic spines and axonic terminals in the bipyramidal neurons of the dorsomedial cortex of lizards (*Lacerta*). — *Anat. Embryol.*, in print.
- Martínez Méndez, F. (1964): Resultados de una campaña de exploración paleontológica a Tenerife. — *Publ. Cated. Paleont. Univ. Barcelona* 1: 1–11.
- Martínez Rica, J.P. (1975): Contribución al estudio de la biología de los geconos ibéricos (Rep., Sauria). — *Publ. Cent. pirenaico Biol. exp.* 5, 1974: 1–294.
- (1982): Primeros datos sobre la población de lagarto negro (*Gallotia simonyi simonyi* STEIND.) de la Isla de Hierro. — *Amphibia-Reptilia* 2 (4): 369–380.
- May, W. (1912): Gomera die Waldinsel der Kanaren. Reisetagebuch eines Zoologen. — *Verh. naturw. Ver. Karlsruhe* 24: ix, 214 pp.
- Mehely, L. von (1904): Materialien zur Systematik und Phylogenie der *Muralis*-ähnlichen Lacer-ten. — *Annl. hist.-nat. Mus. natn. hung.* 7: 409–621.

- Menne, H.A.L. (1956): Hagedissen van de Canarische eilanden. — *Lacerta*, 14 (11/12): 81–83.
- Mertens, R. (1921): Zur Kenntnis der geographischen Formen von *Chalcides ocellatus* Forskal (Rep., Lac.). — *Senckenbergiana* 3 (3/4): 116–120.
- (1922): Verzeichnis der Typen in der herpetologischen Sammlung des Senckenbergischen Museums. — *Senckenbergiana* 4 (6): 162–183.
- (1928): Über den Rassen- und Artenwandel auf Grund des Migrationsprinzips, dargestellt an einigen Amphibien und Reptilien. — *Senckenbergiana* 10 (3/4): 81–91.
- (1934): Die Insel-Reptilien, ihre Ausbreitung, Variation und Artbildung. — *Zoologica*, Stuttg. 32 (6): 209 pp.
- (1935): Zoologische Eindrücke von einer atlantischen Inselfahrt. — *Blätt. f. Aquar- u. Terrarien*. 46: 82–89.
- (1942): *Lacerta goliath* n. sp., eine ausgestorbene Rieseneidechse von den Kanaren. — *Senckenbergiana* 25 (4/6): 330–339.
- Mobbs, A.J. (1981): Observations on the herpetofauna of Tenerife. — *Herptile* 6 (3): 6–12.
- Molina Borja, M. (1981): Etograma del lagarto de Tenerife, *Gallotia galloti galloti* (Sauria-Lacertidae). — *Doñana, Acta vert.* 8: 43–78.
- (1985): Spatial and temporal behaviour of *Gallotia galloti* in a natural population of Tenerife. — *Bonn. zool. Beitr.* 36: 541–552.
- , A. Marrero, A. Carnero & F. Pérez (1980): Reptiles y anfibios de Canarias. I. — *Aguayro* 122: 13–14.
- , —, — & — (1980): Reptiles y anfibios de Canarias. 2. Biología y etología de los lacértidos canarios. — *Aguayro* 123: 12–14.
- , —, — & — (1980): Reptiles de las Islas Canarias 3. Daños en la agricultura y enemigos naturales. — *Aguayro* 124: 14 pp.
- , A. Carnero Hernández, F. Pérez Padrón, A. Marrero, J.J. Bacallado, F. Domínguez, I. Sosa Sosa & M. Báez Fumero (1980): Fauna canaria. Reptiles. — *Aguayro* 122 (15–16); 123 (15–16); 124 (15–16); 125 (15–16); 126 p. 15; 126 p. 16; 127 p. 15; 127 p. 16; 128 p. 16. — Fauna canaria. Anfibios. — *ibid.* 129 p. 21, 129 p. 22 [collection of illustr. species sheets; diverse authors].
- Molowny, A. (1980): Estudio de la corteza cerebral de *Lacerta* y otros reptiles con la técnica de Timm. — Doctoral Thesis, Universidad de La Laguna, Tenerife, Spain.
- & C. López García (1978): Estudio citoarquitectónico de la corteza cerebral de reptiles. III. Localización histoquímica de metales pesados y definición de subregiones Timm positivas en la corteza de *Lacerta*, *Chalcides*, *Tarentola* y *Malpolon*. — *Trab. Inst. Cajal Invest. Biol.* 70: 55–74.
- & F. Marín Girón (1972): Estudio citoarquitectónico de la corteza cerebral de reptiles. I. Areas y núcleos corticales en *Lacerta galloti*, *Chalcides* sp. y *Tarentola mauritanica*. — *Trab. Inst. Cajal Invest. Biol.* 64: 125–158.
- Monzón Mayor, M., J.M. Martín Trujillo & C. López García (1977): Estudio citoarquitectónico de los núcleos mesencefálicos de *Lacerta galloti galloti*. — *Trab. Inst. Cajal Invest. Biol.* 69: 205–222.
- Monzón Mayor, M. (1980): Estudio citoarquitectónico de los núcleos mesencefálicos de *Lacerta galloti* (Dum. et Bibron) y otros Reptiles Squamata (*Lacerta stehlinii*, *Chalcides viridanus* y *Tarentola delalandii*). — Doctoral Thesis, Colegio Univer. Las Palmas (Div. Medicina), Las Palmas, Spain.
- Nettmann, H.K. & S. Rykena (1979): Mauergeckos (*Tarentola mauritanica* L.), die ihre Eier im Sand vergraben (Reptilia: Sauria; Gekkonidae). — *Salamandra* 15: 53–57.
- & — (1985): Verhaltens- und fortpflanzungsbiologische Notizen über kanarische und nordafrikanische *Tarentola*-Arten. — *Bonn. zool. Beitr.* 36: 287–305.
- Oudart, P. (1839): Zoologie, Reptiles. Planche unique. — In: Webb, P. & S. Berthelot (1835–1850) *Histoire Naturelle des Iles Canaries*, Tome deuxième, deuxième partie, Zoologie: 1 pl.
- Pasteur, G. (1981): A survey of the species groups of the Old World scincid genus *Chalcides*. — *J. Herpet.* 15 (1): 1–16.
- & B. Giro (1960): Les tarentes de l'ouest africain. II. *Tarentola mauritanica*. — *Bull. Soc. Sci. nat. phys. Maroc* 40 (4): 309–322.
- & S. Salvidio (1985): Notes on ecological genetics of *Gallotia galloti* populations from Tenerife. — *Bonn. zool. Beitr.* 36: 553–556.
- Pellicer Catalán, M. & P. Acosta, (1971): Estratigrafías arqueológicas canarias: La cueva del Barranco de la Arena (Tenerife). — *Anuar. Estud. Atlant.* 17: 265–279 (Madrid — Las Palmas).

- Peters, G. (1961): Die Perleidechse (*Lacerta lepida* Daudin) gehört zum Subgenus *Gallotia* Boulenger. — Mitt. Zool. Mus. Berl. 37 (2): 271–285.
- (1967): Kriechtiere. — In: Urania Tierreich 4: 355–507. Leipzig, Jena, Berlin.
- Peters, W. & G. Doria (1882): Le crociere dell'Yacht "Corsaro" del capitano armatore Enrico d'Albertis. I. Note erpetologiche e descrizioni di una nuova specie di *Lacerta* delle Isole Canarie. — Annali Mus. civ. Stor. nat. Genova 18: 431–434.
- Pérez Batista, M.A. (1980): Estudio a nivel óptico y ultraestructural del hipotálamo de *L. galloti* y *Ch. viridanus*. — Doctoral Thesis, Univ. La Laguna, Tenerife, Spain.
- , B. Fernández Ruiz & I. Suarez (1981): Estudio de los componentes astrocitarios del hipotálamo de *Chalcides viridanus* (Reptil., Scincidae). — Morf. Norm. y Patol. secc A. 5: 47–53.
- , — & — (1982): A striated intercellular material in the hypothalamus of *Lacerta galloti galloti*. — Arch. Biol. (Bruxelles) 93: 243–247.
- Popp, B. (1961): 3 schöne südliche Eidechsen. — Aquar.- u. Terrar.-Z. 14 (5): 150–151.
- Radovanovic, M. (1959): Zum Problem der Speziation bei Inseleidechsen. — Zool. Jb. (Syst.) 86: 395–436.
- Ramos, A. (1980): Tabla de desarrollo embrionario de *Lacerta galloti galloti* (Periodo de organogénesis) y aspectos de su reproducción. — Memoria de Licenciatura. Facultad de Biología. Universidad de La Laguna, Tenerife [not published].
- Regidor, J. (1978): Tipología neuronal y organización de la corteza cerebral de *Lacerta galloti*. Estudio con los métodos de Golgi. — Las Palmas de Gran Canaria: Excma. Mancomunidad de Cabildos, Plan Cultural, 152 pp.
- , J.M. Martín Trujillo, C. López García & F. Marín Giron (1974): Estudio citoarquitectónico de la corteza cerebral de reptiles. II: Tipología dendrítica y distribución neuronal en áreas corticales de *Lacerta galloti*. — Trab. Inst. Cajal Inves. Biol. 66: 1–32.
- Rodríguez de la Fuente, F. (1981): Las Islas Canarias. Los Reptiles y Anfibios. — pp. 259–271 in Enciclopedia Salvat de la Fauna, Tomo IX, Madrid, 300 pp.
- Rogner, M. (1979): E(chs)kursionen zu den Kanarischen Inseln. — Aquarien Mag. 13 (9): 456–461.
- (1981): Einfach zu pflegende Kanarische Rieseneidechsen. — Das Aquarium 15 (139): 38–40.
- (1982): Durch Nachzucht erhalten: Der Gestreifte Kanarenskink. — Aquar. Mag. 16 (5): 284–285.
- (1983): Zur Biologie, Pflege und Zucht einiger Walzenskink-Arten (*Chalcides*). — Aquarium 174: 657–662.
- Salvador, A. (1972): Nota sobre el lagarto negro gigante de Canarias, *Lacerta simonyi*. — Bol. R. Soc. Esp. Hist. Nat. (Biol.) 69: 317–320.
- (1974): Guía de los Anfibios y Reptiles españoles. — Madrid: Instituto Nacional para la Conservación de la Naturaleza, 282 pp.
- , (1975): Los eslizones de la isla de La Gomera. — Bol. Estac. cent. Ecol. 4 (8): 83–85.
- Schenkel, E. (1901): Achter Nachtrag zum Katalog der herpetologischen Sammlung des Basler Museums. — Verh. naturf. Ges. Basel 13 (1): 142–199.
- Schifter, H. (1965): Langjährige Haltung einer *Lacerta stehlini*. — Aquar.- u. Terrar.-Z. 18 (1): 22–24.
- Schneider, H. (1978): Der Paarungsruf des Teneriffa-Laubfrosches: Struktur, Variabilität und Beziehung zum Paarungsruf des Laubfrosches der Camargue (*Hyla meridionalis* Böttger, 1874, Anura, Amphibia). — Zool. Anz. 201 (3/4): 273–288.
- (1981): Fortpflanzungsverhalten des Mittelmeerlaubfrosches (*Hyla meridionalis*) der Kanarischen Inseln (Amphibia: Salientia: Hylidae). — Salamandra 17 (3/4): 119–129.
- (1982): Phonotaxis bei Weibchen des Kanarischen Laubfrosches, *Hyla meridionalis*. — Zool. Anz., Jena 208 (3/4): 161–174.
- (1985): Bioakustische und verhaltensphysiologische Untersuchungen am Laubfrosch der Kanaren (*Hyla meridionalis*). — Bonn zool. Beitr. 36: 277–286.
- Siebenrock, F. (1894): Das Skelet der *Lacerta Simonyi* Steind. und der Lacertidenfamilie überhaupt. — Sber. Akad. Wiss. Wien (Abth. I) 103 (4/5): 205–292.
- Simony, O. (1892): Die Canarischen Inseln, insbesondere Lanzarote und die Isletas. — Schrit. Vereins Verbr. naturwiss. Kenntnisse in Wien 32: 353–398.
- Spencer, J. (1982): The lizard of Salmor. — Island Gazette (The Canary Islands English Monthly), Febr. 1982: 21–22.
- Steindachner, F. (1889): [Bericht über eine von Prof. O. Simony auf den Roques del Zalmor bei Hierro (Canarische Inseln) entdeckte neue Eidechsenart von auffallender Größe, *Lacerta Simonyi* Steind.]. — Anz. Akad. Wiss. Wien 26 (27): 260–262.

- (1891): Über einige neue und seltene Reptilien und Amphibien. — Anz. Akad. Wiss. Wien, 28 (14): 141–144.
- (1891): Über einige neue und seltene Reptilien und Amphibien-Arten. — Sitzber. kais. Akad. Wiss. Wien, math.-nath. Kl. C, I: 299–306.
- (1891): Über die Reptilien und Batrachier der westlichen und östlichen Gruppe der canarischen Inseln. — Ann. naturh. Mus. Wien 6 (3): 287–306.
- (1900): Expedition S.M. Schiff „Pola“ in das Rothe Meer nördliche und südliche Hälfte, 1895/96 und 1897/98. Zoologische Ergebnisse XVII. Bericht über die herpetologischen Aufsammlungen. — Denkschr. kais. Akad. Wiss. Wien, math.-nat. Cl. 69: 325–335.
- Stumpel, A.H.P. (1973): Waarnemingen aan de herpetofauna van Tenerife (Canarische eilanden). — *Lacerta* 31 (9/10): 153–158.
- Tello Marquina, J.C. (1979): Los lagartos (*Lacerta* sp.) una plaga de algunos cultivos de las Islas Canarias. — An. INIA, Ser. Prot. veg. 9: 183–196.
- Thorpe, R.S. (1985): Relative similarity between subspecies of the western Canary Island lizard, *Gallotia galloti*. — Bonn. zool. Beitr. 36: 529–532.
- (1985): Alternative hypotheses for the causation of geographic variation in the western Canary Island lizard *Gallotia galloti*. — Bonn. zool. Beitr. 36: 533–539.
- (1985): Body size, island size and variability in the Canary Island lizards of the genus *Gallotia*. — Bonn. zool. Beitr. 36: 481–487.
- (1985): Extent of racial divergence in the eastern Canary island lizard, *Gallotia atlantica*. — Bonn. zool. Beitr. 36: 507–512.
- , K. Watt & M. Báez (1985): Some interrelationships of the Canary Island lizards of the genus *Gallotia*. — Bonn. zool. Beitr. 36: 577–584.
- Trujillo Trujillo, C.M. (1982): Ontogénesis de los núcleos talámicos de *Gallotia galloti galloti*. Estudio estructural y ultraestructural. — Doctoral Thesis, Univ. La Laguna, Tenerife, Spain.
- & C. López García (1977): Citoarquitectura de los núcleos talámicos en los Reptiles *Lacerta galloti* y *Chalcides viridanus*. — Trab. Inst. Cajal Inv. Biol. 69: 223–238.
- , M.A. Pérez Batista, B. Fernández Ruiz & J. Martín Trujillo (1983): Formación de la pared talámica en *Gallotia galloti galloti* (Reptil., Lacertidae). I. Zona ventricular. — Morf. Norm. y Patol. secc. A. 7 (3): 855–864.
- , —, — & — (1983): Formación de la pared talámica en *Gallotia galloti galloti* (Reptil., Lacertidae). II. Zona intermedia. — Morf. Norm. y patol. secc. A. 7 (3): 1186–1196.
- Valverde, J.A. (1959): La protection de la faune en Espagne: ses problèmes. — pp. 31–43 in Proc. 7th Tech. Meeting IUCN, Athens.
- Viera y Clavijo, J. de (1866): Diccionario de Historia Natural de las Islas Canarias o índice alfabético descriptivo de sus tres reinos animal, mineral y vegetal. — Las Palmas de Gran Canaria: La Verdad, i–lxxxii, 344 pp.
- Vogel, P. (1974): Der Stimmapparat der Reptilien, mit besonderer Berücksichtigung von *Lacerta galloti*. — Verh. naturf. Ges. Bas. 85 (1/2): 135–153.
- Voggenreiter, V. (1985): Ausgewählte Arealarten von Pflanzen und Tieren der Insel Tenerife und ihre ökologisch-chorologische Interpretation. — Bonn. zool. Beitr. 36: 261–276.
- Wermuth, H. (1965): Liste der rezenten Amphibien und Reptilien: Gekkonidae, Pygopodidae, Xantusiidae. — Tierreich, 80: xxii, 246 pp.
- Werner, F. (1902): Über Kanarische Echsen. — Natur und Haus 10: 11–24 (Wien).
- Wevers, A. jr. (1909): *Lacerta Galloti* und *Lacerta Simonyi*. — *Lacerta* 1902 (2): 8.
- Wijffels, L. (1956): *Lacerta simonyi stehlini* in het terrarium. — *Lacerta* 14 (11/12): 83–84.
- Yanes Mendéz, C. (1985): Ontogenia de las áreas basales telencefálicas de *Gallotia galloti eisen-trauti* (Reptilia, Lacertidae). Estudio estructural y ultraestructural. — Doctoral Thesis, Univ. La Laguna, Tenerife, Spain.
- , J.M. Martín Trujillo & C. López García (1977) Citoarquitectura de los núcleos del estriado de un saurio Lacertidae: *Lacerta galloti galloti*. — Trab. Inst. Cajal Inv. Biol. 69: 273–296.
- Zimmermann, H. (1984): Die Echsen von Teneriffa. Lebensraum, Verhalten, Fortpflanzung. — Aquar. Mag., 18 (8): 390–396.



Machado, A and López-Jurado, A Martín L F. 1985. "Conservation status of reptiles in the Canary Islands." *Bonner zoologische Beiträge : Herausgeber: Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn* 36, 585–606.

View This Item Online: <https://www.biodiversitylibrary.org/item/156275>

Permalink: <https://www.biodiversitylibrary.org/partpdf/120013>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: Zoologisches Forschungsmuseum Alexander Koenig

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.