Aristolochia wuana, a New Name in Chinese *Aristolochia* (Aristolochiaceae)

Liu Zhenwen

Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, 650204, People's Republic of China; The Graduate School of the Chinese Academy of Sciences, Beijing, 100049, People's Republic of China.
liuzw@mail.kib.ac.cn

Deng Yunfei

Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, 650204, People's Republic of China; Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, 510650, People's Republic of China.
Corresponding author: yfdeng@scbg.ac.cn

ABSTRACT. A new name, *Aristolochia wuana* Zhen W. Liu & Y. F. Deng, is proposed to replace the later homonym *Aristolochia macrocarpa* C. Y. Wu & S. K. Wu ex D. D. Tao, not *A. macrocarpa* Duchartre. The species may be closely related to *A. griffithii* Hooker f. & Thomson ex Duchartre but differs in its retuse or rounded leaf apex (vs. acute or shortly acuminate), oblong capsule (vs. narrowly cylindrical), and the smooth upper surface of the seeds (vs. rugose).

Key words: *Aristolochia*, Aristolochiaceae, China.

In 1983, a new species of *Aristolochia* L. (Aristolochiaceae) was described from Xizang, China, as *A. macrocarpa* C. Y. Wu & S. K. Wu ex D. D. Tao (Tao, 1983). The species did not appear in the recently published Volume 5 of *Flora of China* (Huang et al., 2003). Furthermore, this name is a later homonym of *A. macrocarpa* Duchartre from West Africa (Duchartre, 1864; Ma, 1992). After studying the specimens of both *A. macrocarpa* Duchartre and *A. macrocarpa* C. Y. Wu & S. K. Wu ex D. D. Tao, it was apparent that the specimens represent two distinct species. According to the classification of Schmidt (1935) and Gonzalez (1999), *A. macrocarpa* Duchartre belongs to subgenus *Paristolochia* (Hutchinson & Dalziel) O. C. Schmidt and *A. macrocarpa* C. Y. Wu & S. K. Wu ex D. D. Tao belongs to subgenus *Siphisia* (Rafinesque) Duchartre. Subgenus *Siphisia* differs from subgenus *Paristolochia* in having six anthers (vs. eight to 10), a 3-lobed gynostemium (vs. 8- to 10-lobed), and fruit flesh that is dehiscent (vs. woody indehiscent). Subgenera *Siphisia* and *Paristolochia* were treated at generic rank as *Isotrema* Rafinesque and *Paristolochia* Hutchinson & Dalziel, respec-

tively, by some authors (Klotzsh, 1859; Huber, 1960, 1985, 1993; Poncey, 1978; Mabberley, 1987). However, the broad circumscription of *Aristolochia* was widely accepted by others (Hou, 1984; Phuphathana-phong, 1985; Hwang, 1988; Ma, 1989, 1992; Gonzalez, 1999; Murata et al., 2001; Huang et al., 2003) and supported by recent molecular studies (Wanke et al., 2006). According to Article 53.1 of the *International Code of Botanical Nomenclature* (McNeil et al., 2006), *A. macrocarpa* C. Y. Wu & S. K. Wu ex D. D. Tao must be replaced, and the following new name is proposed here.

Aristolochia wuana Zhen W. Liu & Y. F. Deng, nom. nov. Replaced name: *Aristolochia macrocarpa* C. Y. Wu & S. K. Wu ex D. D. Tao, Fl. Xizang. 1: 585. 1983, non *Aristolochia macrocarpa* Duchartre, 1864. TYPE: China. Xizang Autonomous Region: Zayü Xian, in woods, 2100 m, 1 Aug. 1973, Qinghai-Xizang (Tibet) Complex Expedition 73-948 (holotype, KUN).

The species has not yet been described in English. The emended full description is provided as follows.

Shrub climbing; stems terete, striate, sparsely gray villous; buds densely grayish brown villous. Petioles 8–18 cm, sparsely gray villous; leaf blade cordate, 18–24 × 16–20 cm, thick papery, adaxially sparsely pubescent along the veins, abaxially reddish brown or white villous, basally palmate, with primary veins 3 to 5, lateral veins in 3 to 4 pairs, tertiary veins prominent abaxially, base cordate to auriculate, margin entire, apex retuse or rounded. Flowers not seen; pedicels 6–7 cm in fruit. Capsule solitary, oblong, ca. 11 × 6 cm,

6-ribbed, dehiscing at tip; seeds ovoid, upper surface plano-convex, smooth, lower surface deeply concave.

Distribution and habitat. *Aristolochia wuana* is known from northern Gaoligong Shan, occurring in southeastern Xizang and northeastern Yunnan in China. The plants were noted from mixed conifer and broad-leaved forest at altitudes from 2000 to 2300 m.

Phenology. The species has been collected in fruit from August to September. No flowering specimens have been seen.

Etymology. The species epithet honors Wu Zhengyi (Wu Chengyih) (13 June 1916–) of the Kunming Institute of Botany, Chinese Academy of Sciences, who first found *Aristolochia wuana* during the expedition to Qinghai-Xizang Plateau led by him in 1970–80s. Prof. Wu is a great expert on the flora of China and has devoted 70 years to the study of the flora and vegetation of China. He has organized several large-scale surveys of plant resources in southwestern China. He is currently the co-chair of the editorial committee of the *Flora of China*, a joint project between the Chinese Academy of Sciences and Missouri Botanical Garden that aims to revise the Chinese vascular plants. In 2007, he won the State Preeminent Science and Technology Award of China for his outstanding achievements in the fields of systematic botany and plant geography, as well as plant diversity, conservation, and sustainable use of plant resources.

Relationships. In the protologue, *Aristolochia wuana* was regarded as similar to *A. moupinensis* Franchet, differing in its leaves $18\text{--}24 \times 16\text{--}20$ cm (vs. $6\text{--}16 \times 5\text{--}12$ cm), retuse apex (vs. acute or acuminate), and larger fruits ca. 11×6 cm (vs. $6\text{--}8 \times 2\text{--}3.5$ cm) (Tao, 1983; Huang et al., 2003). Ma (1989) placed *A. macrocarpa* C. Y. Wu & S. K. Wu ex D. D. Tao in synonymy with *A. griffithii* Hooker f. & Thomson ex Duchartre. In fact, it differs from the latter in its retuse or rounded leaf apex (vs. acute or shortly acuminate), oblong capsule (vs. narrowly cylindrical), and the smooth upper surface of the seeds (vs. rugose).

Additional specimens examined. CHINA. **Xizang:** Mêdog Xian, near Beibeng, 9 Aug. 1974, *Qinghai-Xizang (Tibet) Complex Expedition 74-4304* (KUN); Zayü Xian, Xia Zayü, 3 Sep. 1976, C. Y. Wu et al. 5450 (KUN). **Yunnan:** Gongshan Xian, the trail from the Dulong Township to Zayü, 6 Sep. 1982, *Qinghai-Xizang (Tibet) Complex Expedition 9974* (KUN).

Acknowledgments. We thank Peng Hua (KUN) and Nicholas Turland (MO) for their comments on the manuscript, and K and KUN for allowing us to

examine the specimens. The project was supported by the grant of Wu Zheng-yi's Outstanding Prize from Yunnan Province, China (KIB-WU-2001-04).

Literature Cited

- Duchartre, P. 1864. Aristolochiaceae. Pp. 421–498 in A. de Candolle (editor), *Prodromus Systematis Naturalis Regni Vegetabilis*, Vol. 15, Pt. 1. Masson, Paris.
- Gonzalez, F. 1999. Inflorescence morphology and the systematics of Aristolochiaceae. *Syst. Geogr. Pl.* 68: 159–172.
- Hou, D. 1984. Aristolochiaceae. Pp. 531–553 in *Flora Malesiana*, Ser. I. Spermatophyta, Vol. 10. Nationaal Herbarium Nederland, Leiden.
- Huang, S. M., L. M. Kelly & M. Gilbert. 2003. Aristolochiaceae. Pp. 246–269 in C. Y. Wu, P. H. Raven & D. Y. Hong (editors), *Flora of China*, Vol. 5. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
- Huber, H. 1960. Zur Abgrenzung der Gattung *Aristolochia* L. *Mitt. Bot. Staatssamml. München* 3: 531–553.
- . 1985. Samenmerkmale und Gliederung der Aristolochiaceen. *Bot. Jahrb. Syst.* 107: 277–320.
- . 1993. Aristolochiaceae. Pp. 129–137 in K. Kubitzki, J. G. Rohrer & V. Bittrich (editors), *The Families and Genera of Vascular Plants*, Vol. 2: Flowering Plants, Dicotyledons—Magnoliid, Hamamelid and Caryophyllid Families. Springer, Berlin.
- Hwang, S. M. 1988. *Aristolochia* L. Pp. 199–245 in H. S. Kiu & Y. R. Ling (editors), *Flora Reipublicae Popularis Sinicae*, Tomus 24. Science Press, Beijing.
- Klotzsh, F. 1859. Die Aristolochiaceae des Berliner Herbariums. *Monatsber. Königl. Preuss. Akad. Wiss. Berlin* 1859: 571–625.
- Ma, J. S. 1989. A revision of *Aristolochia* Linn. from E. & S. Asia. *Acta Phytotax. Sin.* 27: 321–364.
- . 1992. A taxonomic revision of genus *Aristolochia* subgenus *Pararistolochia*. *Acta Phytotax. Sin.* 30: 508–514.
- Mabberley, D. J. 1987. *The Plant-Book: A Portable Dictionary of the Vascular Plants*, 2nd ed. Cambridge University Press, Cambridge.
- McNeill, J., F. R. Barrie, H. M. Burdet, V. Demoulin, D. L. Hawksworth, K. Marhold, D. H. Nicolson, J. Prado, P. C. Silva, J. E. Skog, J. H. Wiersma & N. J. Turland (editors). 2006. *International Code of Botanical Nomenclature (Vienna Code)*. *Regnum Veg.* 146.
- Murata, J., D. Ohi-Thoma, S. G. Wu, D. Darnaedi, T. Sugawara, T. Nakanishi & H. Murata. 2001. Molecular phylogeny of *Aristolochia* (Aristolochiaceae) inferred from *matK* sequence. *Acta Phytotax. Geobot.* 52: 75–83.
- Phuphathanaphong, L. 1985. Studies in Thai Flora 4. Aristolochiaceae. *Thai Forest Bull., Bot.* 15: 29–57.
- Poncey, O. 1978. Le genre *Pararistolochia*, Aristolochiaceae d'Afrique tropicale. *Adansonia* 17: 465–494.
- Schmidt, O. C. 1935. Aristolochiaceae. Pp. 204–242 in A. Engler & K. Prantl (editors), *Die natürlichen Pflanzenfamilien*, Vol. 16b, 2nd ed. W. Engelmann, Leipzig.
- Tao, D. D. 1983. Aristolochiaceae. Pp. 584–587 in C. Y. Wu (editor), *Flora Xizangica*, Vol. 1. Science Press, Beijing.
- Wanke, S., F. Gonzalez & C. Neinhuis. 2006. Systematics of pipevines: Combining morphological and fast-evolving molecular characters to investigate the relationships within the subfamily Aristolochiaceae. *Int. J. Pl. Sci.* 167(6): 1215–1227.

Liu, Zhenwen. and Deng, Yunfei. 2009. "Aristolochia wuana, a New Name in Chinese Aristolochia (Aristolochiaceae)." *Novon a journal of botanical nomenclature from the Missouri Botanical Garden* 19, 370–371.

View This Item Online: <https://www.biodiversitylibrary.org/item/124658>

Permalink: <https://www.biodiversitylibrary.org/partpdf/121936>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Permission to digitize granted by rights holder

Rights: <https://www.biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.