
A New Combination in *Tabernaemontana* (Apocynaceae)

A. J. M. Leeuwenberg

Department of Plant Taxonomy, Agricultural University, 37, Generaal Foulkesweg,
P.O. Box 8010, 6700 ED Wageningen, The Netherlands

In preparation of the *Stemmadenia* and *Tabernaemontana* (Apocynaceae) treatments for the upcoming *Flora of the Venezuelan Guayana*, the following new combination is required.

Tabernaemontana cerea (Woodson) Leeuwenb.,
comb. nov. Basionym: *Stemmadenia cerea*
Woodson, Bull. Torrey Bot. Club 75: 557.
1948. *Anartia cerea* (Woodson) Allorge, Mém.
Mus. Natl. Hist. Nat. sér. 2B, 30: 58, pl. 22.
1985. TYPE: Guyana. Georgetown, NW portion

of Kanuku Mts., Mt. Iramaikpang, A. C. Smith
3606 (holotype, MO; isotypes, A, B, G, MG,
NY, P, S, U, W).

Tabernaemontana cerea does not have the 5
wings below the anthers inside the corolla tube, the
most important character to distinguish *Stemmadenia*
from *Tabernaemontana*. *Anartia* is a syn-
onym of *Tabernaemontana* (see Leeuwenberg, J.
Ethnopharm. 10: 5. 1984).



Leeuwenberg, A J M. 1991. "A New Combination in *Tabernaemontana* (Apocynaceae)." *Novon a journal of botanical nomenclature from the Missouri Botanical Garden* 1, 105–105. <https://doi.org/10.2307/3391635>.

View This Item Online: <https://www.biodiversitylibrary.org/item/14661>

DOI: <https://doi.org/10.2307/3391635>

Permalink: <https://www.biodiversitylibrary.org/partpdf/122213>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.