

## AN ABNORMAL SPECIMEN OF *LAMINARIA DIGITATA* (L.) LAMOUROUX

ARTHUR C. MATHIESON, EMERY F. SWAN  
AND RICHARD A. FRALICK<sup>1</sup>

The present report describes the occurrence and morphology of an abnormal specimen of *Laminaria digitata* (L.) Lamouroux. The plant was found (by Emery Swan) detached in a deep tide pool near mean low water at Odiornes Point, Rye, New Hampshire, U.S.A. (70°42'42" W longitude, 43°03'33" N latitude) on May 16, 1971. Figures 1 and 2 illustrate the general morphology of the plant; a single stipe and two attached blades are evident. The overall length of the plant was 208 cm. The maximum dimensions of the fronds were 120 × 98 and 76 × 74 cm respectively, exclusive of the stipe. The larger blade was darker in color, healthier in appearance, and it seemed to be more actively growing, as evidenced by its segmentation (Fig. 2), than the smaller frond.




*Laminaria digitata* is a common plant in the vicinity of Odiornes Point. It grows abundantly on solid rocks from 0 to 14 m below mean low water, with maximum biomass occurring from -2 to 10 m (Mathieson, Hehre and Reynolds, in press). It is speculated that the plant broke off its holdfast and then regenerated a second frond.

Recent studies by Markham (1968) demonstrated extensive regenerative capacities in *Laminaria sinclairii* (Harvey) Farlow, Anderson *et* Eaton. Detached portions of its haptera, as short as 2.5 mm in length, were capable of developing into entire plants. *Laminaria digitata* also exhibits extensive regenerative capacities (unpublished data, A. Mathieson), for new blades can be produced from intact stipes if they are injured or destroyed. Similar regenerative capacities are probably evident in healthy, pelagic specimens. Figure 3 shows a conspicuous swelling of the

---

<sup>1</sup>Published with the approval of the Director of the New Hampshire Agricultural Experiment Station as Scientific Contribution Number







stipe in the abnormal specimen; it probably represents the site of regeneration of the second frond. The swelling is several cm below the transition zone — the area of “active” cell division. Thus, regeneration occurred in an area presumably restricted to elongation and differentiation of cells. The older appearance of the smaller frond suggests that it is the original one. However, in order to preserve the intact specimen a critical examination (section) of the stipe was not made.

Burrows (1956, 1958) reported extensive populations of loose-lying algae (particularly *Chorda filum*, *Desmarestia aculeata*, *Laminaria saccharina* and *Saccorhiza polyschides*) from the subtidal zones at Port Erin Bay, Isle of Man — particularly after periods of calm weather and high insolation. Most of the plants were initially attached. Extensive SCUBA observations near Odiorne Point have never shown quantities of loose-lying *L. digitata* or any other algae — perhaps because of the exposed nature of the site. The accidental deposition of the specimen in a deep tide pool may have provided a relatively stable habitat and allowed regeneration to occur. According to Burrows (1958) members of the loose-lying populations may serve as a means of restocking habitats devoid of reproductive plants. No reproductive sori were found on the specimen of *L. digitata*. The vegetative state of the plant may have been due to the time of its collection, for *L. digitata* exhibits limited reproduction during the spring in New England (Mathieson, Hehre and Reynolds, in press). It should also be noted that several pelagic algae (e.g. *Sargassum*, *Focus* and *Gracilaria*) are reported (Fritsch, 1959) to have limited reproductive capacities.

---

Figure 1. Intact specimen of *Laminaria digitata*, with a single stipe and two attached blades.

Figure 2. Close up of stipe and fronds, showing darker and healthier appearance of larger (left) frond. Active segmentation of the larger frond is also visible.

Figure 3. Conspicuous swelling of the stipe, that may represent the site of regeneration of the second frond.



According to van Overbeek (1940 a, b) auxins may be found in *Macrocystis pyrifera* and other known brown algae. If this were true for *L. digitata*, then the loss of polarity, due to a pelagic state, may explain the regeneration of a second frond rather than a holdfast. Extensive regeneration and the loss of polarity must also occur in other unattached algae that produce spherical morphologies, such as *Cladophora holsatica* Kützinger described by Smith (1950) and the "beach form" of *Ascophyllum nodosum* (L.) Le Jolis ecad *Mackaii* (Turner) Cotton reported by Gibb (1957).

In conclusion the occurrence of an abnormal specimen of *L. digitata* poses a variety of questions concerning the growth and ecology of the species. Additional studies should be carried out to determine the growth, reproduction and ecological significance of pelagic specimens of *L. digitata*.

#### LITERATURE CITED

- BURROWS, E. M. 1956. A preliminary survey of the sublittoral algae of Port Erin Bay, Isle of Man. *Phycol. Bull.* 4: 14-15.
- , 1958. Sublittoral algae populations in Port Erin Bay, Isle of Man. *J. Mar. Biol. Assoc. U.K.* 37: 687-703.
- FRITSCH, F. E. 1959. The structure and reproduction of the algae. Vol. II. Cambridge University Press, 939 pp.
- GIBB, D. C. 1957. The free-living forms of *Ascophyllum nodosum* (L.) Le Jol. *J. Ecol.* 45: 49-83.
- MARKHAM, J. W. 1968. Studies on the haptera of *Laminaria sinclairii* (Harvey) Farlow, Anderson et Eaton. *Syesis* 1: 125-131.
- MATHIESON, A. C., HEHRE, E. J. and REYNOLDS, N. B. Investigations of New England marine algae I: A floristic and descriptive ecological study of the marine algae at Jaffrey Point, New Hampshire. *Nova Hedwigia* (in press).
- OVERBEEK VAN, F. 1940a. Auxin in marine algae. *Plant Physiol.* 15: 291-299.
- SMITH, G. M. 1950. The fresh-water algae of the United States, McGraw-Hill Book Co., N. Y., 719 pp.

JACKSON ESTUARINE LABORATORY  
OF THE UNIVERSITY OF NEW HAMPSHIRE  
DURHAM, N.H., U. S. A. 03824



Mathieson, Arthur C., Swan, E F, and Fralick, Richard A . 1972. "AN ABNORMAL SPECIMEN OF LAMINARIA-DIGITATA." *Rhodora* 74, 127-130.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/14694>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/123695>

**Holding Institution**

Missouri Botanical Garden, Peter H. Raven Library

**Sponsored by**

Missouri Botanical Garden

**Copyright & Reuse**

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.