Validation of Some Suprageneric Names in Podocarpopsida

Alexander B. Doweld

National Institute of Carpology (Gaertnerian Institution), P.O. Box 72, RUS-119517, Moscow, Russian Federation. nicar@aha.ru

James L. Reveal

Professor Emeritus, University of Maryland, College Park, Maryland 20742-5815, U.S.A. Mailing address: 18625 Spring Canyon Road, Montrose, Colorado 81401-7906, U.S.A. jr19@umail.umd.edu

Abstract. Upon learning that the genus Pherosphaera W. Archer bis (1850) was a synonym of the monospecific Tasmanian genus Microcachrys Hooker f. (1845), the family name Microcachrydaceae Doweld & Reveal (1999) was proposed for conservation giving it priority over Pherosphaeraceae Nakai (1938). The long-used name Pherosphaeraceae was consistently misapplied in the sense of Microstrobos J. Garden & L. A. S. Johnson (1951). Accordingly, a new family, Microstrobaceae, is validated and segregated into its own order, Microstrobales. In addition, for those who might wish to recognize the taxon at a lower rank within Podocarpaceae Endlicher, the names Microstroboideae and Microstrobeae are established to replace the already existing but consistently misapplied Pherosphaeroideae Pilger (1903, 1916) and Pherosphaereae Pilger (1903).

Key words: Microcachryaceae, Microcachrys, Microstrobaceae, Microstrobales, Microstrobeae, Microstroboideae, Microstrobos, Pherosphaera, Pherosphaeraceae, Pherosphaereae, Pherosphaeroideae.

In 1999, while validating and proposing several new suprageneric names in Pinophyta Cronquist, Takhtajan & Zimmermann, we proposed Microcachrydaceae Doweld & Reveal for the rare, monospecific Tasmanian genus *Microcachrys* J. D. Hooker (1845). At that time we were not aware that *Pherosphaera* W. Archer bis (1850) was a synonym of *Microcachrys* and that Pherosphaeraceae Nakai (1938) was the correct name for our new family.

The nomenclatural confusion began in 1857 when Joseph Dalton Hooker misapplied Archer's name to plants of a different genus. In this sense, *Pherosphaera* was used until 1951. Aware that the type of *P. hookeriana* W. Archer bis was based on the female reproductive structures of *Microcachrys tetragona* J. D. Hooker, Garden and Johnson proposed *Microstrobos* J. Garden & L. A. S. Johnson. They also published *Microstrobos niphophilus* J. Garden & L. A. S. Johnson as a new species for what had mistakenly been called *P. hookeriana* for nearly a century. When Nakai proposed his family name in 1938, he circumscribed the taxon to include only what is now known as *Microstrobos*.

In 1953, two proposals (Elliott, 1953; Barber, 1953) were published to conserve *Pherosphaera* J. D. Hooker (1857) over *Pherosphaera* W. Archer bis (1850). Later, Florin (1956), aware of the typification problem, placed *Microstrobos* in synonymy under *Pherosphaera* J. D. Hooker and proposed *Pherosphaera niphophila* (Garden & L. A. S. Johnson) Florin (Shinners, 1957). In 1958, the Committee for Spermatophyta rejected conservation of the later homonym and proclaimed *Microstrobos* to be the correct name for what had long been known as *Pherosphaera* (Rickett, 1958). Even so, *Pherosphaera* sensu J. D. Hooker continued to be used (e.g., Gaussen, 1974; Willis, 1973; Wielgorskaya, 1995; Melikyan & Bobrov, 2000).

Reveal and Doweld (2001) have proposed conservation of Microcachrydaceae so that the correct family name for the only genus in the taxon, Microcachrys, would be Microcachrydaceae rather than Pherosphaeraceae. Nonetheless, we recognize the necessity of establishing a family for Microstrobos, and thus propose Microstrobaceae. It is noteworthy that recently Melikyan and Bobrov (2000) used Pherosphaeraceae for this taxon, being unaware, like us, that Pherosphaera was a synonym of Microcachrys (which they placed in Microcachrydaceae). In addition, these authors proposed a new order "Microstrobales" Bobrov & Melikyan; however, Article 16.1 of the Code (Greuter et al., 2000) requires that all typified ordinal names be based on a validly published family name.

By establishing Microstrobaceae and validating Microstrobales following the classification scheme

Novon 11: 395–397. 2001.

Novon

proposed by Melikyan and Bobrov (2000), we hope to remove all references to the long-confused use of the generic name Pherosphaera for Microstrobos. However, we are fully aware that others consider Microstrobos to be a taxon within Podocarpaceae Endlicher (Conran et al., 2000; Hart, 1987; Kelch, 1997, 1998), and therefore validate both Microstroboideae and Microstrobeae. We knowingly-as permitted by the last sentence in Article 34.1 of the Code (Greuter et al., 2000)—take this action firstly to correct the misapplication of Pherosphaera for Microstrobos at the ranks of subfamily and tribe as suggested by Pilger (1903, 1916), and secondly to provide the necessary nomenclature in Podocarpaceae or whatever other family one might wish to use for these taxa.

It is essential to emphasize that judging by the original Pilger (1903) and Nakai (1938) descriptions, the names Pherosphaeraceae, Pherosphaeroideae, and Pherosphaeraee were established for plants recognized now as belonging to the genus *Microstrobos* (= *Pherosphaera* J. D. Hooker, 1857, non W. Archer bis, 1850). We hope that Microcachrydaceae will be conserved over Pherosphaeraceae but acknowledge that these changes will cause some confusion for researchers consulting the older literature. By correcting the names according to the *Code* (conservation of *Pherosphaera* Hooker f. over W. Archer bis was rejected in 1958!), we hope our modifications will promote nomenclatural stability in the future.

Microstrobaceae Doweld & Reveal, fam. nov. TYPE: *Microstrobos* J. Garden & L. A. S. Johnson, Contr. New S. Wales Nat. Herb. 1: 316. 1951.

Arbores sempervirentes, dioicae; folia acicularia, imbricata, in 4–5 lineas (nunquam decussata vel opposita) disposita; strobili masculi sessiles solitarii, grana pollinis 3-saccata; strobili feminei solitarii sessiles, ovoidei, in aggregationes 2–8, bracteae imbricatae, acuminatae; semen unicum, epimatium nullum.

Evergreen dioecious shrubs; leaves scale-like, imbricate, arranged in 4–5 rows but never decussate or opposite; male cones sessile, solitary, pollen with 3 air sacs; female cones solitary, sessile, ovoid, in aggregations of 2–8, bracts imbricate, acuminate; seed 1, epimatium lacking.

Microstrobales Melikyan & Bobrov ex Doweld & Reveal, ord. nov. TYPE: *Microstrobos* J. Garden & L. A. S. Johnson, Contr. New S. Wales Nat. Herb. 1: 316. 1951. Microstrobaceae Doweld & Reveal, Novon 11: 396. 2001. Ab ordinibus Podocarpalibus Saxegothaealibusque epimatiis nullis differt.

An order differing from the Podocarpales and the Saxegothaeales in lacking an epimatium.

Microstroboideae Doweld & Reveal, subfam. nov. TYPE: *Microstrobos* J. Garden & L. A. S. Johnson, Contr. New S. Wales Nat. Herb. 1: 316. 1951. Microstrobaceae Doweld & Reveal, Novon 11: 396. 2001.

Arbores sempervirentes, dioicae; strobili masculi sessiles solitarii, terminales; strobili feminei terminales, solitarii sessiles, bracteae imbricatae, acuminatae; semen erectum, unicum, epimatium nullum.

Evergreen dioecious shrubs; male cones sessile, solitary, terminal; female cones terminal, solitary, sessile, bracts imbricate, acuminate; seed erect, one, epimatium lacking.

Microstrobeae Doweld & Reveal, trib. nov. TYPE: Microstrobos J. Garden & L. A. S. Johnson, Contr. New S. Wales Nat. Herb. 1: 316. 1951. Microstrobaceae Doweld & Reveal, Novon 11: 396. 2001.

Arbores sempervirentes, dioicae; folia acicularia, imbricata, in 4–5 lineas (nunquam decussata vel opposita) disposita; strobili terminales, sessiles, solitarii; semen unicum, erectum, epimatium nullum; numerus chromosomatum haploideus n = 13.

Evergreen dioecious shrubs; leaves scale-like, imbricate, arranged in 4–5 rows but never decussate or opposite; cones terminal, sessile, solitary; seed one, erect, epimatium lacking; n = 13.

Acknowledgments. We are grateful to K. N. Gandhi at Harvard University Herbaria who provided us with some of the literature for this paper, and to a reviewer who provided additional clarifications and suggestions. This research is a contribution to the Systema Spermatophytorum Project of the National Institute of Carpology (Gaertnerian Institution), Moscow. This study is part of the Index Nominum Supragenericorum Plantarum Project from the Norton-Brown Herbarium (MARY) and presented with the cooperation of the International Association for Plant Taxonomy and the National Agricultural Library, Beltsville, Maryland.

Literature Cited

- Archer, W. 1850. Note on *Microcachrys*, Hook. fil., and on a new allied genus of Coniferae of Van Diemen's Land. Hooker's J. Bot. Kew Gard. Misc. 2: 51–52.
- Barber, H. N. 1953. Propositions pour le Congrès de Paris, nos. 28–32. Taxon 2: 214.
- Conran, J. G., G. G. Wood, P. G. Martin, J. M. Dowd, C.

- J. Quinn, P. A. Gadek & R. A. Price. 2000. Generic relationships within and between the gymnosperm families Podocarpaceae and Phyllocladaceae based on an analysis of the chloroplast gene *rbc*L. Austral. J. Bot. 45: 715–724.
- Doweld, A. B. & J. L. Reveal. 1999. Validation of new suprageneric names in Pinophyta. Phytologia 84: 363– 367.
- Elliott, C. G. 1953. Proposal no. 32. Proposal to conserve the genus *Pherosphaera* sensu Hooker. Taxon 2: 141.
- Florin, R. 1956. Nomenclatural notes on genera of living gymnosperms. Taxon 5: 188–192.
- Garden, J. & L. A. S. Johnson. 1951. *Microstrobos*, a new name for a podocarpaceous genus Contr. New S. Wales Nat. Herb. 1: 315–316.
- Gaussen, H. 1974. Les Gymnosperms actuelles et fossiles. Fascicule XIII. Les Podocarpines sauf les *Podocarpus*. Trav. Lab. Forest. Toulouse 2(1) fasc. XIII: 1–174.
- Greuter, W., J. McNeill, F. R. Barrie, H. M. Burdet, V. Demoulin, T. S. Filgueiras, D. H. Nicolson, P. C. Silva, J. E. Skog, P. Trehane, N. J. Turland & D. L. Hawksworth. 2000. International Code of Botanical Nomenclature (Saint Louis Code). Regnum Veg. 138.
- Hart, J. A. 1987. A cladistic analysis of conifers: Preliminary results. J. Arnold Arbor. 68: 269–307.
- Hooker, J. D. 1845. On the Huon Pine, and on *Micro-cachrys*, a new genus of Coniferae from Tasmania; to-gether with remarks upon the geographical distribution of that order in the Southern Hemisphere. London J. Bot. 4: 137–157.

—____. 1857. Coniferae. Bot. Antarct. Voy. 3 [Fl. Tasman.]: 349–359.

- Kelch, D. G. 1997. The phylogeny of the Podocarpaceae based on morphological evidence. Syst. Bot. 22: 113– 131.
- Melikyan, A. P. & A. V. Bobrov. 2000. Morphology of female reproductive structures and an attempt of the construction of phylogenetic system of orders Podocarpales, Cephalotaxales, and Taxales. Bot. Zhurn. (Moscow & Leningrad) 85(7): 50–67.
- Nakai, T. 1938. Indigenous species of conifers and taxads of Korea and Manchuria, and their distribution. Tyosen-Sanrin 158: 1–15.
- Pilger, R. 1903. Taxaceae. Pflanzenr. IV, 5(Heft 18): 1– 124.
- . 1916. Kritische Übersicht über die neuere Literatur betreffend die Familie der Taxaceae. Bot. Jahrb. Syst. 54: 1–43.
- Reveal, J. L. & A. B. Doweld. 2001. A proposal to conserve the family name Microcachrydaceae (Pinophyta). Taxon 50.
- Rickett, H. W. 1958. Report of the Committee for Spermatophyta. Conservation of generic names I. Taxon 7: 184–193 [Pherosphaera: 186].
- Shinners, L. H. 1957. Against conservation of Xanthophyllum and Pherosphaera. Taxon 6: 135–137.
- Wielgorskaya, T. 1995. Dictionary of Generic Names of Seed Plants. Columbia Univ. Press, New York.
- Willis, J. C. 1973. A Dictionary of the Flowering Plants and Ferns, revised by H. K. Airy Shaw, 8th ed. Cambridge Univ. Press, Cambridge.



Doweld, A B and Reveal, James L. 2001. "Validation of some suprageneric names in Podocarpopsida." *Novon a journal of botanical nomenclature from the Missouri Botanical Garden* 11, 395–397. <u>https://doi.org/10.2307/3393150</u>.

View This Item Online: https://doi.org/10.2307/3393150 Permalink: https://www.biodiversitylibrary.org/partpdf/12808

Holding Institution Missouri Botanical Garden, Peter H. Raven Library

Sponsored by Missouri Botanical Garden

Copyright & Reuse Copyright Status: In copyright. Digitized with the permission of the rights holder. License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.