

SCIENTIFIC NOTE:

***CLOEODES EXCOGITATUS* (EPHEMEROPTERA: BAETIDAE) IN NORTHERN CALIFORNIA¹**R.D. Waltz², Peter Ode³, Jon Lee⁴

The small minnow mayfly genus *Cloeodes* Traver is found in many different freshwater habitats throughout the Neotropics, Afrotropics, Asia and southern temperate areas of North America. Various revisionary works (Waltz and McCafferty 1987) and systematic works (Kluge 1991; Waltz 1993; Waltz and McCafferty 1994; McCafferty and Lugo-Ortiz 1995) have provided taxonomic data for the genus and added greatly to our knowledge of the world distribution of the genus.

The genus was recognized as having tropical affinities by Waltz and McCafferty (1987), McCafferty and Waltz (1990), and McCafferty et al. (1992). McCafferty and Lugo-Ortiz (1995) and Lugo-Ortiz and McCafferty (1993, 1994, 1995) further added to the known distribution of the genus in the Western Hemisphere, viz., southwestern North America, Mesoamerica, and South America.

Herein, we report a remarkable, although possibly not unexpected, distribution record for this primarily subtropical and tropical genus from northern California. Larvae of *C. excogitatus* Waltz and McCafferty were taken in the following site in northern California: CA: Mendocino Co., Williams Crk near Covelo, 29-IX-1996. The above site is characterized by fine silt bottoms mixed with cobble. Stream temperatures are warm in the summer months. The above specimens were taken in low flow conditions near the end of the summer dry season.

Lugo-Ortiz and McCafferty (1994, 1995) extended the known range of *C. excogitatus* from its type locality in Oak Creek Canyon, Arizona, southward to Yavapai County (Arizona) and Mexico. The type locale of *C. excogitatus* was previously the most northern report of the genus in North America. This report extends the known range of this species northward into North America by at least 600 miles reaching the Coastal Biotic Province of California (Usinger 1953) at approximately 40 degrees north latitude. Other northward extensions of primarily tropical genera into North America are also known in other mayfly genera, e.g., the baetid genera *Camelobaetidium* Demoulin and *Paracloeodes* Day (see McCafferty et al. 1992).

Voucher specimens are deposited at the Purdue Entomological Research Collection, Purdue University, West Lafayette, IN and the California Academy of Science, San Francisco.

LITERATURE CITED

- Kluge, N. 1991. Cuban mayflies of the family Baetidae (Ephemeroptera). 1. Genera *Callibaetis*, *Cloeodes*, and *Paracloeodes*. Zool. Zh. 12: 128-136. [in Russian].
Lugo-Ortiz, C.R and W.P. McCafferty. 1993. Genera of Baetidae (Ephemeroptera) from Central America. Entomol. News 104: 191-195.

¹ Received October 31, 1997. Accepted December 17, 1997.

² IDNR, Division of Entomology and Plant Pathology, 402 West Washington, Room W-290, Indianapolis, IN 46204.

³ Aquatic Biological Assessment Laboratory, California Department of Fish and Game, 2005 Nimbus Road, Rancho Cordova, CA 95670.

⁴ 2250 Wilson Street, Arcata, CA 95521.

- Lugo-Ortiz, C. R. and W.P. McCafferty.** 1994. New records of Ephemeroptera from Mexico. Entomol. News 105: 17-26.
- Lugo-Ortiz, C.R. and W.P. McCafferty.** 1995. Annotated inventory of the mayflies (Ephemeroptera) of Arizona. Entomol. News 106: 131-140.
- McCafferty, W.P. and C.R. Lugo-Ortiz.** 1995. *Cloeodes hydatum*, n.sp. (Ephemeroptera: Baetidae): An extraordinary, drought tolerant mayfly from Brazil. Entomol News 106: 29-35.
- McCafferty, W.P., R.W. Flowers, and R.D. Waltz.** 1992. The biogeography of Mesoamerican mayflies. pp. 173-193. In: S.P. Darwin and A.L. Weldon (eds.). Biogeography of Mesoamerica: proceedings of a symposium. Tulane Univ. Stud. Zool. Bot., Suppl. Publ. 1.
- McCafferty, W.P. and R. D. Waltz.** 1990. Revisionary synopsis of the Baetidae (Ephemeroptera) of North and Middle America. Trans. Am. Entomol. Soc. 116: 769-799.
- Usinger, R.L.** 1953. Introduction to Aquatic Entomology. Stream and Lake Classifications. pp 7-8 in Aquatic Insects of California, with keys to North American genera and California species. Univ. California Press, Berkeley. 508 pp.
- Waltz, R.D.** 1993. *Cloeodes binocularis* (Ephemeroptera: Baetidae) a new combination for a Neotropical species of *Pseudocloeon* s. auctt. Entomol. News 104: 233-234.
- Waltz, R.D. and W.P. McCafferty.** 1987. Revision of the genus *Cloeodes* Traver (Ephemeroptera: Baetidae). Ann. Entomol. Soc. Am. 80: 191-207.
- Waltz, R.D. and W.P. McCafferty.** 1994. *Cloeodes* (Ephemeroptera: Baetidae) in Africa. Aquat. Insects 16: 165-169.



Waltz, R. D., Ode, P, and Lee, J. 1998. "Cloeodes Excogitatus (Ephemeroptera : Baetidae) In Northern California." *Entomological news* 109, 215–216.

View This Item Online: <https://www.biodiversitylibrary.org/item/20693>

Permalink: <https://www.biodiversitylibrary.org/partpdf/13388>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: American Entomological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.