NOTES

longing to Capparales and, though generally accorded recognition, Thorne (1976) includes *Tovaria* in Capparaceae. Raven (1975) has suggested x = 10 or 11 as basic in Capparaceae, which has a range of numbers from n = 17, 16, 14, 12, 11, 10 and 8, while x = 12 (or possibly 14) seems basic in Cruciferae, in which a wide range of numbers occur. In the remaining families of the order, Resedaceae has n = 10, 6, 12, and 11 and Moringaceae n = 14 in the three species studied. *Tovaria* appears well placed in this alliance, and its base number of x = 14 accords well with Moringaceae and falls within the range recorded for Capparaceae, although n = 14 is unusual in this family, being reported only in two African genera, *Euadenia* and *Buchholzia*. The occurrence of x = 14 in another family of Capparales tends to support Raven's hypothesis of a tetraploid origin with x = 14 as basic for the order.

LITERATURE CITED

- BEHNKE, H. D. & T. J. MABRY. 1977. S-type sieve-element plastids and anthocyanins in Vivianiaceae: evidence against its inclusion into Centrospermae. Pl. Syst. Evol. 126: 371-375.
- BORTENSCHLAGER, S. 1967. Vorläufige Mitteilungen zur Pollenmorphologie in der Familie der Geraniaceen und ihre systematische Bedeutung. Grana Palynol. 7: 400–468.

DAHLGREN, R. 1975. A system of classification of the Angiosperms to be used to demonstrate the distribution of characters. Bot. Not. 128: 119–147.

GOLDBLATT, P. & P. K. ENDRESS. 1977. Cytology and evolution in Hamamelidaceae. J. Arnold Arbor. 58: 67-71.

HUTCHINSON, J. 1959. The Families of Flowering Plants. Ed. 2. Vol. 1. Dicotyledons. Clarendon Press, Oxford.

LEFOR, W. M. 1971. A taxonomic revision of the Vivianiaceae. Ph.D. dissertation, Univ. of Connecticut.

Biol. Sci. Ser. 2(15): 225–255.

RAVEN, P H. 1975. The bases of Angiosperm phylogeny: cytology. Ann. Missouri Bot. Gard. 62: 724-764.

TAKHTAJAN, A. 1969. Flowering Plants: Origin and Dispersal. Transl. by C. Jeffrey. Oliver & Boyd, Edinburgh.

THORNE, R. F. 1976. A phylogenetic classification of the Angiospermae. Evol. Biol. 9: 35-106.

I would like to thank Otto Zöllner for seed of Viviania crenata and Alwyn H. Gentry for the fixed buds of *Tovaria pendula*.

-Peter Goldblatt, B. A. Krukoff Curator of African Plants, Missouri Botanical Garden, 2345 Tower Grove Avenue, St. Louis, Missouri 63110.

ADDITIONS TO THE PANAMANIAN CONVOLVULACEAE

Since the publication of the Convolvulaceae in the *Flora of Panama* (Austin, 1975) several collections have been made of taxa new to the country. There has also been a name change.

1978]

1. Ipomoea ramosissima (Poir.) Choisy in DC., Prodr. 9: 377. 1845. Based on Convolvulus ramosissimus Poir.

Convolvulus ramosissimus Poir. in Lam., Encycl. Méth. Suppl. 3: 468. 1813. New name for C. cymosus Ruiz & Pavón.

C. cymosus Ruiz & Pavón, Fl. Peru 2: 9. tab. 417. f.b. 1799, non Desr., 1796. TYPE: Peru, Huanuco, Ruiz & Pavón (holotype presumably MA, not seen; F, isotype).

These plants are similar to Ipomoea trifida (H.B.K.) G. Don with which they have been confused previously. The flowers are smaller in *I. ramosissima* (13-24 mm long) than in *I. trifida* (30-40 mm long) and the sepals are shaped differently; obovate to elliptic-obovate in the former, ovate in the latter. While *I. ramosissima* is comparatively common in southern South America, it appears to be rare in Central America. It has been reported in Costa Rica under the name *I. quesadana* Standley (1940: 99) (Austin-Smith 1609, F, holotype), and in Guatemala as *I. perplexa* L. O. Williams (1970: 193) (Belize, Bartlett 12868, MICH, holotype, not seen; F, isotype).

DARIÉN: Duke 10217 (MO).

2. Ipomoea indica (Burm. f.) Merrill, Int. Rumph. Herb. Amb. 445. 1917, should replace the binomial *I. acuminata* (Vahl) Roem. & Schult. While Merrill transferred the epithet correctly, he did not typify the name. Because of doubt, various authors (e.g. van Ooststroom, 1953: 465; Verdcourt, 1963: 113) have considered the name suspect. F. R. Fosberg finally unraveled the cryptic comments in the Burman original and typified the epithet. *Ipomoea indica* is based on *Convolvulus indicus flore violaceo* Besler, Hort. Eyst. Aest. Ord. 13, fig. 2. 1613 (chosen lectotype by Fosberg, 1976).

3. Maripa lewissi D. Austin, sp. nov.

Frutex scandens, ramis lignosis teretibus glabris. Folia rigide coriacea, glabra, breviter obtuse acuminataque, 6–18 cm longa, 2–8 cm lata, petiolo 5–15 mm longo, nervo supra sulciformi subtus prominente. Paniculae terminales, basi foliatae, pyramidales, rachi glabra, cymis contractis. Sepala coriacea, interiora 4–5 mm, exteriora 5–6 mm longa, glabrata. Corolla purpurea, 10–12 mm longa. Fructus ellipsoideus, breviter apiculatus, 16–19 mm longus, abortu 1-locularis, 1-spermus, pericarpio coriaceo-lignoso tenui.

Lianas; stems terete, light to dark grey brown. Leaves with petioles 5–15 mm long; blades glabrous, ovate to oblong-elliptic or ovate-lanceolate, 6–18 cm long, 2–8 cm wide, the base rounded, the apex acuminate to obtuse-acuminate, coriaceous, the secondary veins 8–11 pairs. Inflorescences terminal, paniculate-thyrsiform; bracts triangular to triangular-subulate, 1–1.5 mm long, acute, persistent at anthesis and in fruit, glabrous. Flowers with the pedicels 3–4 mm long; sepals broadly ovate, the outer 5–6 mm long, the inner 4–5 mm long, coriaceous, glabrous or with scattered glandular indument; corolla cylindric-funnel form, lavender, 10–12 mm long, the lobes rounded; stamens included, the filaments 6 mm long, inserted 3 mm above the corolla base, the anthers narrowly ovate, basally sagittate, 2 mm long; ovary conic-oblong, 2 mm long, glabrous, the style 9 mm long. Fruits ellipsoid, 16–19 mm in diameter, rounded in cross-section, faintly striate, brown when dry, yellow when living, the calyx loosely appressed

NOTES

against the fruit base; seeds 1, slightly flattened, oblong to ellipsoid, 12–14 mm long, 8–10 mm across, tan when dry.

TYPE: PANAMA. PANAMÁ, El Llano-Cartí road, 8 km N of Pan-American Highway at El Llano, 5 Mar. 1974, Nee & Warmbrodt 10379 (MO, holotype; FAU, isotype).

The species is named for Walter H. Lewis who turned my attention to the tropical members of the Convolvulaceae in 1966. These lianas grow between 300-450 m in the Premontane Wet Forest and bloom in March. Fruiting specimens have been collected between May and December.

This is the only small-flowered (10-12 mm) member of the genus known in Central America. *Maripa panamensis* and *M. nicaraguensis*, which are the two species previously known from the area, have flowers 25–45 mm long. *Maripa lewissii* belongs to sect. *Maripa*, while the other two species in Central America are members of sect. *Mouroucoa* (Austin, 1973). All of the other members of sect. *Maripa* are known from east of the Andes Mountains.

Additional specimens seen: COLÓN: Santa Rita Ridge, 30 Dec. 1971, Dwyer & Gentry 9341 (MO), 19 Sep. 1971, Gentry 1858 (MO). PANAMÁ: El Llano-Cartí road, 8–11 km from Inter-American Highway, 14 Aug. 1975, Mori 7740 (FAU, MO). 9.6 km from Inter-American Highway, 25 May 1975, Mori & Kallunki 6375 (MO). 10–12 km from Inter-American Highway, 31 Oct. 1974, Mori & Kallunki 2929 (FAU, MO). 7–12 km from Inter-American Highway, 18 July 1974, Croat 25176 (MO). SAN BLAS: Inland from airport at Mandinga, 7 Oct. 1966, Duke 8914 (MO).

KEY TO PANAMANIAN Maripa

- - b. Leaves chartaceous to subcoriaceous, reticulate beneath, the veins prominent below; flowers densely sericeous outside; fruits smooth to slightly striate when dry _________. M. nicaraguensis Hemsley
 b. Leaves thick-coriaceous, rugose beneath, the veins not prominent below; flowers

Кеч то Maripa sect. Maripa

a. External sepals appressed pubescent outside.

- aa. External sepals glabrous or with peltate scales outside.
 c. Internal sepals appressed dibrachiate-pubescent; all sepals appressed to base of fruits.

d. Leaves with sunken oil glands above, glabrous below; ovary glabrous ...

- dd. Leaves without oil glands but with peltate scales or glabrous; ovary glabrous or pubescent.

 - ee. Sepals subequal; corolla indument balanced or unbalanced dibrachiate; fruits 25–35 mm long.
 - f. Corolla indument balanced dibrachiate; fruits 30-35 mm long
 - ff. Corolla indument unbalanced dibrachiate; fruits 25 mm long
 - M. janusiana D. Austin

1978]

- cc. Internal sepals glabrous outside or glandular; sepals loosely appressed or spreading in fruit.

 - gg. Flowers 10-12 mm long; fruits 16-19 mm long, ellipsoid, shortly apiculate *M. lewisii* D. Austin

Thanks are due Ron Liesner for pointing out the specimens on which *Maripa lewisii* is based and to Marshall Crosby for loaning the material.

LITERATURE CITED

- AUSTIN, D. F. 1973. The American Erycibeae (Convolvulaceae): Maripa, Dicranostyles, and Lysiostylis I. Systematics. Ann. Missouri Bot. Gard. 60: 306-412.
- -----. 1975. Convolvulaceae. In R. E. Woodson, Jr. & R. W. Schery, Flora of Panama. Ann. Missouri Bot. Gard. 62: 157-224.
- FOSBERG, F. R. 1976. Ipomoea indica taxonomy: a tangle of Morning Glories. Bot. Not. 129: 35-38.

OOSTSTROOM, S. J. VAN. 1953. Convolvulaceae. In Flora Malesiana, 4: 388-512.

STANDLEY, P. C. 1940. Studies of American plants—X. Publ. Field Mus. Nat. Hist., Bot. Ser. 22: 65–129.

VERDCOURT, B. 1963. Convolvulaceae. In Flora of Tropical East Africa. 161 pp.

WILLIAMS, L. O. 1970. Tropical American plant, XI. Fieldiana, Bot. 32: 179-206.

—Daniel F. Austin, Department of Biological Sciences, Florida Atlantic University, Boca Raton, Florida 33431.

TRATTINICKIA ASPERA (BURSERACEAE) REPORTED NEW FOR SOUTH AMERICA

The Flora of Panama treatment for the Burseraceae by D. M. Porter (Ann. Missouri Bot. Gard. 57: 5–27. 1970) reported *Trattinickia aspera* (Standley) Swart to be endemic to Panama. The species has now been discovered in Colombia based on *Garcia-Barriga 3600* from the Commisaria del Putomayo, along the Mocoa Trail between Mocoa and Puerto Limón at an elevation of 500–600 m. The report of this species for Colombia is not at all surprising, but it is unusual that its first report for South America would be based on a collection made near the Ecuadorian border and on the edge of the Amazonian basin.

—Thomas B. Croat, P. A. Schulze Curator of Botany, Missouri Botanical Garden, 2345 Tower Grove Avenue, St. Louis, Missouri 63110.

A NAME CHANGE IN THE ANTILLEAN SOLANACEAE

Jaltomata antillana (Krug & Urban) D'Arcy, comb. nov.

Saracha antillana Krug & Urban, Notizbl. Königl. Bot. Gart. Berlin 1: 80. 1895. SYNTYPES: Jamaica, Harris 5109, 5522 (B, if extant, neither seen).



Austin, Daniel F. 1978. "Additions to the Panamanian Convolvulaceae." *Annals of the Missouri Botanical Garden* 65, 777–780. <u>https://doi.org/10.2307/2398879</u>.

View This Item Online: https://doi.org/10.2307/2398879 Permalink: https://www.biodiversitylibrary.org/partpdf/13545

Holding Institution Missouri Botanical Garden, Peter H. Raven Library

Sponsored by Missouri Botanical Garden

Copyright & Reuse Copyright Status: In copyright. Digitized with the permission of the rights holder. License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.