XXXVI. *On a New Genus of Balanophoreæ from New Zealand, and Two New Species of Balanophora.* By J. D. HOOKER, M.D., F.R.S. & L.S. &c.

Read June 16th, 1859.

SINCE laying before the Linnean Society my observations on the structure and affinities of *Balanophoreæ**, I have received two additional species of *Balanophora*, and a remarkable new genus of the Order, which I shall now proceed to describe under the name of *Dactylanthus*.

DACTYLANTHUS, n. g.

Dioicum. *Rhizoma* tuberosum, lobatum. *Pedunculi* plurimi, squamati, apice spadices plurimos gerentes. Flores in spadices dense conferti, inferiores dissiti. FL. MASC. *Stamen* solitarium sessile, filamentum brevissimo, anthera 2-loculari. FL. FÆM. *Perianthium* ovario globoso subsessili adnatum, lobis 2-3 subulatis, stylo filiformi, stigmatibus simplicibus.

1. DACTYLANTHUS TAYLORI, H. f. (TAB. LXXV. A.)

Hab. Nova Zelandia ad radices *Fagi* et *Pittospori* montibus prope Wanganui, alt. 4000 ped. (*Rev. R. Taylor, M.A.*). *Nom. vern.* *Pua reinga* (Flos Hades).

Rhizoma magnitudine pugilli et ultra, tuberculatum, superficie celluloso, pustulis papillisve nullis, intus sicco rubrum. *Pedunculi* numerosi, 4-6 unc. alti, sicco fragiles, rugulosi, opaci, ferruginei, intus rufescentes, squamis imbricatis laxè tecti, apice in receptaculum floriferum dilatati. *Squamæ* oblongæ, obtusæ, lata basi sessiles, inferiores dissitæ, $\frac{1}{4}$ unc. longæ, brunneæ, superiores sensim majores, supremæ inflorescentiam involucentes, 1-2 unc. longæ, lineari-oblongæ, membranaceæ, albæ medio sordide brunneæ v. rubræ. *Spadices* (v. pedicelli floriferi) 10-30, conferti, erecti, 1-2 unc. longi, cylindracei v. fusiformes, crassitie pennæ corvinæ, stipitati, basi nudi v. floribus deformati bracteolati. *Flores* dense conferti. MASC. *Stamen* solitarium nudum, sessile, filamentum brevissimo crasso; anthera adnata, 2-locularis, transverse oblonga, rimis subverticalibus dehiscens, loculis crassiusculis, inferiores in spica imperfectæ dissitæ stipitatæ. *Pollen* pallidum, globosum. FL. FÆM. sicco nigri, sub $\frac{1}{10}$ unc. longi. *Ovarium* stipitatum, subglobosum. *Perianthium* supra ovarium paulo angustatum, dein bi-, rarius 3-partitum, a dorso compressum, cum stylo connatum; segmentis respectu axeos lateralibus, subulatis, inæqualibus. *Stylus* cylindricus, teres, superne paulo incrassatus, apice obtusus.—“*Planta viva odorem subgratum Melonis, sed demum ingratum humi spirat.*”—*Taylor.*

For a specimen of this singular plant I am indebted to my friend the Rev. R. Taylor, of New Zealand, who brought a fragment of it to England in 1856, and on my pointing out its probable interest, promised to procure more on his return to New Zealand. This he did; and early in the present year I had the pleasure of receiving from him a dried specimen of the female plant, a perfect male inflorescence in a letter, and a pen and ink sketch of the peduncle and flowers, with notes on the same. Owing to the great contraction of the cellular tissue, and blackening and fragility of the only entire specimen, which was unfortunately loosely packed in a box and much damaged, I have had considerable difficulty in so restoring the parts as to enable Mr. Fitch to complete the accompanying Plate, and myself to make the necessary analyses.

* Page 1 of this volume.

The most remarkable character of *Dactylanthus* is its inflorescence, which, instead of presenting the solitary capitulum or spadix of most of its allies, or the branched character of *Sarcophyte*, consists of numerous erect spadices, densely covered with flowers. Of these the males consist of a solitary stamen, not articulate with the spadix, and without any trace of perianth, and hence present by far the most reduced form of male flower in the Order. The female spadices present a crinite appearance from the long flexuose styles of the flowers. The ovary appears to be compressed, furnished with a short thick stipes, and its upper part is contracted into a short thick neck, apparently consisting of the tube of the perianth connate with the base of the style: this contracted part varies much in width and breadth, and sometimes appears as if reduced to the adnate bases of the subulate lobes of the perianth.

I have not been able to ascertain the nature either of the contents of the ovary, which is extremely minute, and crumbles into black powder on being touched, or of the tissues of the rhizome and peduncle, which do not seem to differ materially from those of other *Balanophoreæ*.

The position of *Dactylanthus* is amongst my first group of *Balanophoreæ*, the *Monostyli*; but in many respects it partakes of the structure of various genera of *Distyli*, and is not very nearly related to any described genus. In habit it presents the rhizome of *Balanophora*. In the male flowers reduced to a solitary stamen, it agrees with *Lophophytum* and *Thonningia*, and in the subulate perianth-lobes of the female flower with *Cynomorium*. Upon the whole it may perhaps rank nearest to *Thonningia*, with which it agrees in habit, in some points of structure of the female flower, especially in the tube of the perianth being connate with the style above the ovary, in the absence of imperfect flowers amongst the females, of bracts, and of a perianth to the male flower.

The natives of New Zealand call this plant "Flower of Hades," because it grows almost buried in the soil; they further consider it allied to *Freycinetia*, which it strikingly resembles in the inflorescence.

Mr. Taylor informs me that he has heard vaguely of the existence of another species with blue flowers in the forests of Mount Egmont.

2. *BALANOPHORA HARLANDI*, H. f. Dioica, rhizomate parvo lobato epustulato, pedunculo ima basi bracteis subfoliaceis involucrato, capitulo globoso, floribus fœmineis capitulo sessilibus. (TAB. LXXV. B.)

Hab. Sylvis insulæ Hongkong (*Harland*); in convalle Mont. Gough (*C. Wilford*, Dec. 1858).

Species pusilla, 1-3 pollicaris, ab omnibus differt pedunculo basi tantum squamato; et præterea a *B. dioica*, capitulo globoso, et rhizomate epustulato; a *B. fungosa* capitulo unisexuali globoso, floribus fœmineis capitulo (non pedicello bracteolæ) sessilibus.

3. *BALANOPHORA LOWII*, H. f. Dioica, rhizomate parvo papilloso et pustulato, pedunculo brevi, squamis inferioribus parvis, superioribus magnis numerosis imbricatis concavis capitulum velantibus, capitulo fl. ♂ oblongo, receptaculo columnari, perianthio 4-fido, lobis lateralibus angustioribus. (TAB. LXXV. C.)

Hab. Sylvis umbrosis montis Kina Balou, insulæ Borneo (*H. Low*).

Hooker, Joseph Dalton. 1859. "On a New Genus of Balanophoreæ from New Zealand, and Two New Species of Balanophora." *Transactions of the Linnean Society of London* 22, 425–427.

<https://doi.org/10.1111/j.1096-3642.1856.tb00114.x>.

View This Item Online: <https://www.biodiversitylibrary.org/item/46591>

DOI: <https://doi.org/10.1111/j.1096-3642.1856.tb00114.x>

Permalink: <https://www.biodiversitylibrary.org/partpdf/14297>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.