

A new species of *Conocardium* Bronn, 1834 (Mollusca: Rostroconchia) from the Mississippian of Missouri, U. S. A.

Richard D. Hoare

Department of Geology, Bowling Green State University, Bowling Green, Ohio 43403-0218, U. S. A.

Abstract: The rostroconch *Conocardium formosum* n. sp. is described from the Mississippian (Chesterian) of Missouri. The narrow, elongate shape of the shell is different from all other known North American species of the genus.

Key Words: Mollusca, Rostroconchia, Mississippian, Missouri

Numerous reports on North American Mississippian rostroconchs have appeared, among which are: Hall (1856, 1883); Winchell (1862, 1870); White and Whitfield (1862); Whitfield (1882); Herrick (1888a, b); Miller (1892); Rowley (1900); Beede (1906); Girty (1910); Weller (1916, 1921); Branson (1942, 1958); Pojeta and Runnegar (1976); and Hoare (1990). Species of *Conocardium* s.s. were included in many of these reports but only that of Pojeta and Runnegar (1976) included narrowly elongate forms of the genus. These authors illustrated but did not describe two specimens, one Pennsylvanian and one Mississippian in age, designating both as *Conocardium* aff. *C. elongatum* (Sowerby, 1815). *Conocardium elongatum* is a Lower Carboniferous (Mississippian) species known from England and is quite different from the American Mississippian species described herein.

The holotype is in the collections of the National Museum of Natural History (USNM).

SYSTEMATICS

Class Rostroconchia Pojeta, Runnegar, Morris,
and Newell, 1972

Order Conocardioida Neumayr, 1891

Superfamily Conocardiacea Miller, 1889

Family Conocardiidae Miller, 1889

Genus *Conocardium* Bronn, 1834

Conocardium formosum, new species

Fig. 1

Conocardium aff. *C. elongatum* (Sowerby). Pojeta and Runnegar, 1976, p. 70, pl. 38, figs. 4-7.

Diagnosis. Narrowly elongate shell with distinct separa-




Fig. 1. *Conocardium formosum* n. sp. 1-5, holotype, right lateral, posterior, anterior, dorsal, and ventral views. Carterville Formation, Missouri, USNM 209298 (scale bar = 1 cm).

tion of body and snout; strongly developed folded wing-like structure dorsally and laterally on snout.

Description. Elongate, narrow, body distinct from snout; sides of snout subparallel; umbonal areas extend slightly above hinge line; dorsal margin slopes ventrally from beak area both posteriorly and anteriorly; ventral margin of body convex below beaks becoming straighter towards rostrum and snout, curves sharply ventrally meeting snout area anteriorly; anterior gape subcircular, restricted to anterior extremity; posterior orifice lacking; rostrum suboval in cross section, length unknown; rostral face and body with 29 even-sized, rounded, closely-spaced, radial costae with narrow interspaces followed anteriorly by 10 fine more

widely spaced costae; snout ornamentation begins at beak area as a V-shaped, folded, wing-like structure expanding anteriorly on dorsal and lateral surfaces, marked ventrally by four coarse costae followed dorsally by five or more fine, widely-spaced costae; ventrally, snout ornamentation forms an inverted V-shaped margin with radial costae near anterior gape; inner shelves not visible.

Measurements. Length, not including rostrum, 13.1 mm; width, 5.1 mm; height, 6.8 mm.

Etymology. Latin, *formosum*, beautifully formed.

Type. Holotype, USNM 209298.

Occurrence. Upper Carterville Formation (Mississippian, Chesterian) in mine dump near Duenweg, Missouri.

Discussion. *Conocardium formosum* differs from *C. elongatum* (Sowerby, 1815) by the presence of a more sharply set-off wing-like structure on the snout, a more abruptly curved rostral face, a narrower, more distinctly separated snout from the body, and sides of snout subparallel not converging anteriorly. Other Chesterian species that have been described (e. g. *C. peculiare* Girty, 1910; *C. chesterensis* Weller, 1921) are not as elongate and have much smaller length/height ratios as is the case for all other Mississippian species of the genus known in North America.

The folded wing-like structure on the snout of *Conocardium formosum* is similar to the structure present on the snout of species of the genus *Arceodomus* Pojeta and Runnegar, 1976. *Arceodomus* has a thin, relatively smooth outer shell layer covering the radial costae on the body. No trace of such an outer layer is present on the holotype of *C. formosum*. The ornamentation on the snout of the Pennsylvanian *Arceodomus prolata* Hoare and Mapes, 1990, differs significantly from that of *C. formosum* in having coarser, evenly-sized and spaced radial costae with intervening fine radial and transverse lirae.

LITERATURE CITED

- Beede, J. W. 1906. Pelecypoda. In: *Fauna of the Salem Limestone of Indiana*, pp. 1323-1334. E. R. Cummings and J. W. Beede, Indiana Department of Geology 30th Annual Report.
- Branson, C. C. 1942. Correction of homonyms in the lamellibranch genus *Conocardium*. *Journal of Paleontology* 16:387-392.
- Branson, C. C. 1958. Two Mississippian species of *Conocardium*. *Oklahoma Geology Notes* 18:137-142.
- Bronn, H. G. 1834. *Lethaea Geognostica*. Vol. 1, E. Schweizerbart, Stuttgart. 672 pp.
- Girty, G. H. 1910. New genera and species of Carboniferous fossils from the Fayetteville shale of Arkansas. *New York Academy of Sciences, Annals* 20:189-238.
- Hall, J. 1856. Descriptions of new species of fossils from the Carboniferous limestones of Indiana and Illinois. *Transactions of the Albany Institute* 4:1-36.
- Hall, J. 1883. No title given. *Indiana Department of Geology and Natural History, 12th Annual Report for 1883* pp. 319-375.
- Herrick, C. L. 1888a. The geology of Licking County, Part 4. The Subcarboniferous and Waverly group. *Denison University Scientific Laboratories Bulletin* 3:13-110.
- Herrick, C. L. 1888b. Geology of Licking County, Part 4. List of Waverly fossils continued. *Denison University Scientific Laboratories Bulletin* 4:11-60, 97-123.
- Hoare, R. D. 1990. Mississippian rostroconch mollusks from Ohio. *Journal of Paleontology* 64:725-732.
- Hoare, R. D. and R. H. Mapes. 1990. *Arceodomus prolata* n. sp. (Mollusca, Rostroconchia) from the Pennsylvanian of Texas. *Journal of Paleontology* 64:491-492.
- Miller, S. A. 1889. North American Geology and Paleontology. *Western Methodist Book Concern* 664 pp. (First appendix, 1892, pp. 665-718; second appendix, 1897, pp. 719-793.)
- Miller, S. A. 1892. Paleontology. In: S. S. Girty, pp. 699-705. *Indiana Department of Geology and Natural Resources 16th Annual Report for 1891*.
- Neumayr, M. 1891. Beiträge zu einer morphologischen Eintheilung der Bivalven. *K. Akademie der Wissenschaften zu Wien Denkschriften* 58:781-801.
- Pojeta, J. R., Jr. and B. Runnegar. 1976. The paleontology of rostroconch mollusks and the early history of the phylum mollusca. *U. S. Geological Survey Professional Paper* 968:1-88.
- Pojeta, J. R., Jr., and B. Runnegar, N. J. Morris, and N. D. Newell. 1972. Rostroconchia, a new class of bivalved mollusks. *Science* 177:264-267.
- Rowley, R. R. 1900. Descriptions of new species of fossils from the Devonian and Subcarboniferous rocks of Missouri. *The American Geologist* 25:264-273.
- Sowerby, J. 1812-1815. The mineral conchology of Great Britain; or coloured figures and descriptions of those remains of testaceous animals for shells which have been preserved at various times and depths in the earth, vol. 1, London. 234 pp.
- Weller, S. 1916. Description of a Ste. Genevieve limestone fauna from Monroe County, Illinois. *Contributions of the Walker Museum, University of Chicago* 1:243-265.
- Weller, S. 1921. Geology of Hardin County, Illinois. *Illinois State Geological Survey Bulletin* 41:1-416.
- White, C. A. and R. P. Whitfield. 1862. Observations upon the rocks of the Mississippi Valley which have been referred to the Chemung Group of New York, together with descriptions of new species of fossils from the same horizon at Burlington, Iowa. *Proceedings of the Boston Society of Natural History* 8:289-306.
- Whitfield, R. P. 1882. On the fauna of the Lower Carboniferous limestones at Spergen Hill, Indiana, with a revision of the descriptions of its fossils hitherto published, and illustrations of the species from the original type series. *Bulletin of the American Museum of Natural History* 1:39-97.
- Winchell, A. 1862. Descriptions of fossils from the Marshall and Huron groups of Michigan. *Proceedings of the Academy of Natural Sciences of Philadelphia* pp. 405-430.
- Winchell, A. 1870. Notices and descriptions of fossils, from the Marshall Group of the western states, with notes on fossils from other formations. *Proceedings of the American Philosophical Society* 9:245-260.

Date of manuscript acceptance: 29 November 2001



Hoare, Richard D. 2002. "A new species of *Conocardium* Bronn, 1834 (Mollusca: Rostroconchia) from the Mississippian of Missouri, U.S.A." *American malacological bulletin* 17, 123–124.

View This Item Online: <https://www.biodiversitylibrary.org/item/173085>

Permalink: <https://www.biodiversitylibrary.org/partpdf/144471>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: American Malacological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.