

Mukherjee who was provided morphometric data and colour photographs. The species has been determined as *Creobroter apicalis* as the eye-mark is placed in the middle of the forewing.

There are, in all, 6 species presently under the genus *Creobroter* in India. *C. apicalis* has been reported earlier from Ambenali in Maharashtra (Mukherjee and Hazra 1983). The species is also known from Karnataka (Mukherjee *et al.* 1995) and our collection from Santegully, Kumta, (14.ix.1998 N. Rane) becomes an additional report, but from a definite locality in Karnataka State. All the specimens in our collection are females. Mukherjee *et al.* (1995) also record the examination of 23 females and of one male specimen.

## ACKNOWLEDGEMENTS

We are grateful to Dr. T.K. Mukherjee for

help in specific determination of this mantis, and the authorities of Modern College for facilities. One specimen of this species was collected during DBT-Funded 'Biodiversity Monitoring Project Workshop' organized by the Centre for Ecological Sciences, Indian Institute of Science, Bangalore. We are grateful for financial assistance provided for the Project through Prof. Madhav Gadgil and IISc, Bangalore.

February 29, 2000

H.V. GHATE  
NILESH RANE  
SACHIN RANADE  
Post-Graduate Research Centre,  
Department of Zoology,  
Modern College,  
Shivaji Nagar,  
Pune 411 005,  
Maharashtra, India.

## REFERENCES

- MUKHERJEE, T.K. & A.K. HAZRA (1983): On a small collection of Mantidae (Dictyoptera) from Maharashtra, India, with the description of a new species. *Rec. Zool. Surv. India* 80: 459-465.  
MUKHERJEE, T.K., A.K. HAZRA & A.K. GHOSH (1995): The mantid fauna of India (Insecta: Mantodea). *Oriental Ins.* 29: 185-358.

21. *SISYPHUS LONGIPES* (OLIVER) (COLEOPTERA : SCARABAEIDAE : SCARABAEINAE) — A NEW RECORD FOR ANDAMAN ISLANDS

The Andamans and the Nicobars, situated 1,200 km off the Indian mainland in the Bay of Bengal between 6° and 14° N and 91° and 94° E, though rich in insect fauna with several endemic species, dung beetles are very poorly represented on these islands. Only six species having been reported, namely *Catharsius molossus* L., *Copris spinator* Har., *Onthophagus cervus* F., *O. orientalis* Har., *O. unifasciatus* (Schall.), and *Paraphytus andamanus* Arrow (Arrow 1931, Veenakumari and Prashanth Mohanraj 1994). None of these species, however, belong to the dung roller group. We report the occurrence of *Sisyphus longipes* (Oliver), a dung roller of the

Family Sisyphini from the Andaman Islands. A single specimen was caught in Garacharma, S. Andaman on January 25, 1998.

*S. longipes* has a wide distribution from Sri Lanka through central and eastern India to Burma (= Myanmar) (Arrow 1931). Many elements of the Andaman fauna (eg. a large percentage of the avifauna) are presumed to have arrived on these islands across the much narrower stretches of water that existed between Burma and these islands, as compared to any of the other neighbouring continental areas, during the Pleistocene sea level lowering (Ripley and Beehler 1989). If *S. longipes* had arrived on these




Fig. 1: *Creobroter apicalis* ♀, actual size 30 mm (length)



islands during that period and established itself here, then it must be a very rare species or one with cryptic habits, as we have not found any more specimens during our dung beetle surveys on these islands. This is likely as *S. longipes* is known to inhabit obscure places like the nests of ants (Arrow, 1931). In case the species has not yet established itself on these islands, the specimen collected by us may be part of the waif biota arriving on these islands or a vagrant, just like the dozen or so species of butterflies that Ferrar (1948) identified as vagrants on these islands. Further studies can establish the status

of this species on the Andaman Islands.

## ACKNOWLEDGEMENT

We thank Dr. A. K. Bandyopadhyay, Director, Central Agricultural Research Institute for encouragement.

June 26, 1999

K. VEENAKUMARI  
PRASHANTH MOHANRAJ  
Central Agricultural Research Institute,  
Port Blair 744 101,  
Andaman and Nicobar Islands, India.

## REFERENCES

- ARROW, G. J. (1931): The Fauna of British India including Ceylon and Burma. Coleoptera: Lamellicornia, III (Coprinae). Taylor and Francis, London, pp. 428.
- FERRAR, M. L., (1948): The butterflies of the Andamans and Nicobars. *J. Bombay nat. Hist. Soc.* 47: 470-491.
- RIPLEY, S. D. & B.M. BEEHLER (1989): Ornithogeographic affinities of the Andaman and Nicobar Islands. *J. Biogeogr.* 16: 323-332.
- VEENAKUMARI, K. & PRASHANT MOHANRAJ (1994): *Onthophagus unifasciatus* F. (Coleoptera: Scarabaeidae: Scarabaeinae) - A new record for Andaman Islands. *J. Bombay nat. Hist. Soc.* 91(1): 153-154.

## 22. LARGE SCALE EMERGENCE AND MIGRATION OF THE COMMON EMIGRANT BUTTERFLIES *CATOPSILIA POMONA* (FAMILY : PIERIDAE)

During my journey on June 17, 1999, through the forest tracts between Mahasamund (Dist. H.Q.) to Tumgaon and Jhalap (NH 6), Madhya Pradesh, no less than five to six thousand Common Emigrant butterflies (*Catopsilia pomona*) were observed flying south to north at a moderate height of 0.60 m to 4 m above ground. At that time (1230 hrs to 1330 hrs) the sun was shining. This forest tract surrounds a big man-made reservoir named Kodar and has teak plantation patches in between the forest, on NH 6.

Interestingly, in the teak (*Tectona grandis*) patches, the butterflies were almost absent, whereas in mixed deciduous forest patches they were present in large numbers.

A few Mottled Emigrants (*Catopsilia pyranthe*) and Lime Butterflies (*Papilio demoleus*) were also flying with the Common

Emigrants. It was noted that the swarm of butterflies seemed to be on a northward migration. During my return journey (1600 hrs to 1700 hrs) the sky was heavily clouded and it was drizzling; hardly 200 to 300 butterflies were seen on the same route.

Butterflies usually migrate northward to avoid the southwest monsoon. In this case, the migration may be due to premonsoon rain in the month of June. The locality had moderate rains in the past 15 days, but the monsoon was yet to set in. The large scale emergence and migration appeared to have started three months in advance. Also, the marked absence of the species in teak patches was interesting.

November 18, 1999

A.M.K. BHAROS  
B-101, Gayatri Nagar,  
PO Shanker Nagar,  
Raipur 492 001, Madhya Pradesh, India.



Veenakumari, K. and Mohanraj, Prashanth. 2000. "Sisyphus Longipes (Oliver) (Coleoptera : Scarabaeidae : Scarabaeinae): a new Record for Andaman Islands." *The journal of the Bombay Natural History Society* 97, 298–301.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/189503>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/155324>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Biodiversity Heritage Library

**Copyright & Reuse**

Copyright Status: In Copyright. Digitized with the permission of the rights holder

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.