

- KEARNEY, T. H. and R. H. PEEBLES. 1951. Arizona flora. Univ. California Press, Berkeley.
- KRAUSE, D. L. and K. I. BEAMISH. 1972. Taxonomy of *Saxifraga occidentalis* and *S. marshallii*. Canad. J. Bot. 52:2131-2141.
- . 1973. Notes on *Saxifraga occidentalis* and closely related species. Syesis 6:105-114.
- ORLOCI, L. 1969. Information analysis of structure in biological collections. Nature 223:483-484.
- RAVEN, P. H. and H. J. THOMPSON. 1964. Haploidy and angiosperm evolution. Amer. Naturalist 98:251-252.

A NEW SUBSPECIES OF RHUS CHONDROLOMA (ANACARDIACEAE) FROM MEXICO

DAVID A. YOUNG

Department of Biological Sciences
Union College, Schenectady, New York 12308

During my biosystematic investigation of *Rhus* subg. *Lobadium* (Young, 1975), a new subspecies of *Rhus chondroloma* Standley was discovered. Because the new subspecies and *R. chondroloma* ssp. *chondroloma* are involved in hybrid complexes with *Rhus standleyi* Barkley it is desirable to publish the description of this new taxon before the details of hybridization are presented.

***Rhus chondroloma* Standley ssp. *huajuapanensis* Young, ssp. nov.** (fig. 1). A ssp. *chondroloma* differt: rami et petioli et segmenta rachidis et foliola pubescentes; foliola parviora et numerosiora.

TYPE: Mexico, Oaxaca, distrito Huajuapán, ca 2 mi N of Huajuapán de León, on limestone palm lands along Mex Hwy 190, ca 1620 m, 22 Jan 1974, S. L. Buchmann and D. A. Young 1-96. Holotype: RSA; isotypes: MEXU, TEX, UC, US.

Aromatic evergreen shrub or small tree 1-6 m high, with relatively stout, brownish, densely pubescent twigs, dotted with reddish lenticels. Leaves pinnately compound, 3-9 leaflets (most frequently 7), 7.0-10.0 (8.5; numbers in parentheses are the mean for a particular character) cm long; leaflets coriaceous, entire, slightly revolute, margin white-corneous, conspicuously pallid-veined, bluish-green above, pale green below; terminal leaflet 2.0-4.0 (3.0) cm long, 1.5-4.0 (2.5) cm wide, distinctly obovate, apex obtuse to rounded (rarely retuse to emarginate), base obtuse; lateral leaflets 2.0-4.0 (3.0) cm long, 1.5-2.5 (1.7) cm wide, elliptic to oval, obtuse at both ends, sessile to subsessile; upper surface of leaflets pilose to puberulous, veins densely pubescent, margins ciliate, lower surface pilose, also densely covered with sessile orange-glandular trichomes. Petioles 1.0-3.0 (1.9) cm long, wingless; rachis segments 1.0-2.5 (1.8) cm long, distinctly winged; petioles and rachis




FIG. 1. *Rhus chondroloma* ssp. *huajuapensis*. a, male flower; b, fruits; c, pistil of female flower; d, sepal; e, leaf; f, inflorescence; g, pistil of male flower; h, lower surface of leaflet (dark dots represent glandular trichomes).

segments covered with a dense soft pubescence. Inflorescences small, dense, terminal spikes, ca 1–4 cm long, slightly narrower. Bracts ovate-deltoid, ca 1–2 mm long, 1.5–2.0 mm wide, densely pubescent on outer surface with simple and orange-glandular trichomes, glabrous on inner surfaces. Sepals greenish-pink, rotund, ca 1–2 mm in diameter, sparingly ciliate with simple and orange-glandular trichomes. Petals white, oval, sexually dimorphic (smaller in male-sterile flowers), ca 2.5–4.0 (3.4) mm long, ca 1.5–2.5 (2.0) mm wide, glabrous, not ciliate. Stamens smaller than sepals in male-sterile flowers, slightly longer than sepals in hermaphrodites. Styles 3, more or less distinct. Fruit a drupe, pubescent with reddish-orange glandular trichomes and long simple trichomes, ca 8–10 mm in diameter.

HABITAT AND DISTRIBUTION (fig. 2): Endemic to the limestone soils in the area surrounding Huajuapán de León, Oaxaca, and adjacent Puebla,

Mexico. Some common associates are: *Amelanchier denticulata*, *Tecoma stans*, *Lippia* spp., *Mimosa* spp., and *Ptelea trifoliolata*.

SPECIMENS EXAMINED: Mexico, Oaxaca, Entre Los Cerros Jicota y Amarillo, 2300 m, *Cabrera* 32 (ENCB); between Tlaxiaco and Teposcolula, *Camp* 2320 (ENCB, NY); Cerro Solo, 7 km al NE de Tepeleme, 2350 m, *Cruz Cisneros* 2108 (ENCB, DS, MICH, MSC); in limy soil of xeric hillside in palm land ca 8 mi N of Huajuapán de León, *Rowell, Webster, and Barkley* 17M579 (male) and 17M580 (female) (ARIZ, ENCB, MICH, MSC, TEX, UC); Magdalena Jicotlán, distrito de Coixtlahuaca, 2100 m, *Rzedowski* 26688 (DS, ENCB, MICH, MSC); on limestone 6 mi NW of Huajuapán de León, 6200 ft, *Webster, Miller, and Miller* 11436 (MEXU, GH). Puebla, limestone hills with shrubby vegetation, Mex Hwy 125, 0.9 mi NE of Acatepec, *Denton* 1485 (MICH); Chila, hillslope, *Saunders* s.n. (ENCB).

Key to subspecies of *Rhus chondroloma*

- a. Leaflets mostly trifoliolate (1–7), often unifoliolate, glabrous; terminal leaflet 4.5–6.5 cm long, 2.5–5.0 cm wide, lateral leaflets 2.5–4.5 cm long, 1.5–3.0 cm wide; petioles and rachis segments glabrous; Guerrero and Puebla. *R. chondroloma* ssp. *chondroloma*
- aa. Leaflets mostly 7 (3–9), pubescent; terminal leaflet 3.0–4.0 cm long, 1.5–4.0 cm wide; lateral leaflets 2.0–4.0 cm long, 1.5–2.5 cm wide; branches, petioles and rachis segments pubescent; Oaxaca.
 *R. chondroloma* ssp. *huajuapánensis*




FIG. 2. Geographical distribution of the two subspecies of *Rhus chondroloma*. (Base map used by permission of the editor of Goode Base Maps, copyright by The University of Chicago Department of Geography).

Rhus chondroloma ssp. *huajuapaneensis* differs most noticeably from *R. c.* ssp. *chondroloma* (Standley, 1936) in its markedly pubescent branches, petioles, rachis segments, and leaflets and in its smaller, more numerous leaflets. *Rhus c.* ssp. *huajuapaneensis* appears to flower mainly from May to July, although a few flowers were present on one herbarium specimen examined collected in January. My collections in January were all in bud; mature fruits were present in September. The species is subdioecious; morphologically bisexual flowers were present, but either the anthers or ovules appeared to be nonfunctional. No pollinators were observed.

In the herbarium material examined, this subspecies was consistently identified as *Rhus duckerae* Barkley, and it keys to *R. duckerae* in Barkley's (1937) key. *Rhus c.* ssp. *huajuapaneensis* is quite distinct from the type of *R. duckerae* (US!), which is the only collection of the latter taxon. The status of *R. duckerae* as a distinct species is questionable. Preliminary studies suggest that *R. duckerae* is actually a hybrid between *R. oaxacana* Loesen. and *R. standleyi* (Young, 1975).

ACKNOWLEDGMENTS

I thank Drs. S. Carlquist, C. E. Jones, R. Scogin, and R. F. Thorne for critically reading the manuscript, which represents a portion of my Ph.D. Dissertation submitted to the Claremont Graduate School, Claremont, California. Field studies were supported by National Science Foundation grant (Improving Doctoral Dissertations in the Field Sciences) GB 38327. Thanks are also due to Consejo Nacional de Ciencia y Tecnologia, Mexico City, Mexico, for a collecting permit (No. 1/609-5466), the Editors of the Goode Base Maps for permission to use their copyrighted base map, Susan Payne for Figure 1, and the curators of the following herbaria for loans of specimens: ARIZ, DS, ENCB, GH, MEXU, MICH, MSC, NY, TEX, UC, US.

LITERATURE CITED

- BARKLEY, F. A. 1937. A monographic study of *Rhus* and its immediate allies in North and Central America, including the West Indies. Ann. Missouri Bot. Gard. 24:265-500.
- STANDLEY, P. C. 1936. Studies of American plants. VI. Field Mus. Bot. Ser. 11:145-276.
- YOUNG, D. A. 1975. Systematics of *Rhus* subgenus *Lobadium* section *Styphonia* (Anacardiaceae). Unpublished Ph.D. Dissertation, Claremont Graduate School, Claremont, California.



Young, David A. 1976. "A NEW SUBSPECIES OF RHUS CHONDROLOMA (ANACARDIACEAE) FROM MEXICO." *Madroño; a West American journal of botany* 23, 354–357.

View This Item Online: <https://www.biodiversitylibrary.org/item/185095>

Permalink: <https://www.biodiversitylibrary.org/partpdf/169339>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: California Botanical Society

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.