

A NEW VARIETY OF *PENSTEMON FREMONTII*
(SCROPHULARIACEAE) FROM COLORADO

ROBERT D. DORN
Box 1471, Cheyenne, WY 82003

ROBERT W. LICHVAR
RD #2 Keller Road, West Valley, NY 14171

ABSTRACT

Penstemon fremontii var. *glabrescens*, a new variety from western Colorado, is described and illustrated. The var. *glabrescens* has glabrous or glabrate leaves and narrow deciduous basal leaves unlike the usually pubescent and mostly persistent basal leaves of var. *fremontii*. Habitats are likewise different and var. *glabrescens* is found only along the southern edge of the range of var. *fremontii*.

While conducting a threatened and endangered species inventory in Garfield County, Colorado, we encountered a *Penstemon* that appeared to be a minor variant of *P. fremontii* Torrey & A. Gray. We later found similar material in the Rocky Mountain Herbarium (RM) from an adjacent county. The habitats were identical and the difference from typical *P. fremontii* was the same.

***Penstemon fremontii* T. & G. var. *glabrescens* Dorn & Lichvar, var. nov.** (Fig. 1)—TYPE: USA, Colorado, Garfield Co.: T7S R100W sect. 24 W $\frac{1}{2}$ NW $\frac{1}{4}$ NW $\frac{1}{4}$, ca. 30 km SE of Douglas Pass, 2440 m, shale slope in mountain brush, 7 Jul 1987, *Dorn 4656* (holotype, RM; isotypes, to be distributed).

Differt a var. *fremontii* foliis, praeter interdum margines et costam, glabris vel glabratis.

Perennial from creeping rootstocks. Stems 1.5–3 dm long, short-pubescent to base or nearly so. Leaves opposite, the basal oblanceolate or obovate and sometimes deciduous, transitional to the lanceolate or linear-lanceolate upper leaves, (1–)2–7 cm long, 2–12 mm wide, glabrous, or short-pubescent on midrib and margins, or sometimes the lower blades sparsely pubescent. Inflorescence short-pubescent to glabrate, not glandular. Sepals glabrous, ovate and acuminate, with narrow erose-scarious margins, 3–5 mm long. Corolla blue, 15–22 mm long, the lobes (2–)3–5 mm long, glabrous. Anthers pubescent, 1–1.5 mm long, indehiscent at base, mostly bluish with pale suture margins. Staminode bearded with yellow hairs near tip. Capsules 7–9 mm long.




FIG. 1. *Penstemon fremontii* var. *glabrescens*. Circle shows close-up of stem pubescence.




FIG. 2. Colorado, Utah, Wyoming, and parts of adjacent states showing collection locations of *Penstemon fremontii* var. *fremontii* (circles) and var. *glabrescens* (triangles) based on specimens at COLO, CS, and RM. One symbol may represent more than one collection.

PARATYPES: USA, Colorado, Garfield Co.: T8S R99W sect. 6 NE¼ SW¼, 2500 m, shale slope, 8 Jul 1987, *Lichvar* 7339 (RM). Rio Blanco Co.: Fawn Creek Road, Piceance Basin, 1950 m, 6 Jul 1978, *Peterson et al.* 1178b (CS); T2N R98W sect. 17 SE¼ NW¼ NW¼, 1860 m, 10 Jun 1982, *Baker and Nauman* 82-184 (CS); T3S R95W sect. 11 NE¼, 2135 m, 2 Jul 1982, *Kelley and Riefler* 82-37 (CS); along Piceance Ck., 2.1 km N of Rock School, 1885 m, 20 Jun 1983, *Peterson and Rollins* 83-215 (CS, RM); T2S R95W sect. 26 NE¼ SW¼, 2195 m, 3 Jun 1986, *O'Kane* 2408 (COLO, RM); Piceance Creek Valley, 16 km upstream from Rock School, 18 Jun 1988, *Weber* 17922 (COLO).

In var. *fremontii* at least the lower leaves are almost always distinctly short-hairy, but occasional individuals have leaves with a tendency toward glabrescence (e.g., *Neese* 14102 [RM]), the basal leaves tend to be broad and persistent, and the corolla lobes are often short and narrow. In var. *glabrescens* most of the leaves are

TABLE 1. SELECTED CHARACTERISTICS FOR *PENSTEMON FREMONTII* AND RELATED TAXA.

Species or variety	Lower stems hairy	Leaf width (mm)	Leaves hairy	Inflorescence	Calyx length (mm)	Corolla length (mm)	Anther length (mm)
<i>P. fremontii</i>							
var. <i>fremontii</i>	yes	2-18(-27)	yes	pubescent	2.5-6.5	(12-)14-23(-28)	(1-)1.2-1.5(-1.8)
var. <i>glabrescens</i>	yes	2-12	midrib and margin	pubescent	2-5	14-22	1-1.5
<i>P. gibbensii</i>	yes (no)	2-7(-8)	yes (no)	glandular	3.5-7(-8)	(15-)16-18(-20)	1-1.3
<i>P. saxosorum</i>	no	3-19	no	glabrous	3.5-8	17-25(-30)	1-1.5
<i>P. scariosus</i>	no	2-15(-23)	no	glandular (glabrous)	(3-)4-9(-13)	(15-)16-30(-33)	(1.3-)1.5-2.2(-2.6)

glabrous or glabrate except sometimes for the margins and midvein, the basal leaves tend to be narrow and deciduous, and the corolla lobes are often long and wide.

Variety *fremontii* is typically found associated with *Artemisia tridentata* on relatively well developed soils. Variety *glabrescens* is found on partly barren, south facing slopes of Green River shale with *Amelanchier*, *Chrysothamnus*, and *Holodiscus* on poorly developed soils. The main elevational range of the two varieties is nearly the same, about 1770 to 2500 m, but var. *fremontii* has been found down to 1525 m. The habitats are strikingly different, however, and var. *glabrescens* is found only at the southernmost edge of the range of var. *fremontii* (Fig. 2).

Variety *glabrescens* also resembles *P. gibbensii* Dorn but lacks the glands in the inflorescence and tends to have broader leaves. *Penstemon scariosus* Pennell is also similar but has glabrous stems below the inflorescence and the hairs on the anthers are generally longer and more copious and the anther sacs are longer. *Penstemon saxosorum* Pennell likewise has glabrous stems below the inflorescence. A comparison of selected features for these taxa is presented in Table 1.

ACKNOWLEDGMENTS

We thank Ronald Hartman, curator of RM, for use of those facilities, the curators of COLO and CS for allowing study of their specimens, and the reviewers of the original manuscript, Noel Holmgren and William Weber, for their helpful comments. Field work was supported by Bureau of Land Management Contract #CO-910-CT7-016.

(Received 12 Apr 1989; revision accepted 9 Apr 1990.)



Dorn, Robert D. and Lichvar, Robert W . 1990. "A NEW VARIETY OF
PENSTEMON FREMONTII (SCROPHULARIACEAE) FROM COLORADO."
Madroño; a West American journal of botany 37, 195–199.

View This Item Online: <https://www.biodiversitylibrary.org/item/185620>

Permalink: <https://www.biodiversitylibrary.org/partpdf/171067>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: California Botanical Society

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.