
A New Species of *Pecluma* (Pteridophyta–Polypodiaceae)

Olga G. Martínez

Herbario MCNS, Facultad de Ciencias Naturales, Universidad Nacional de Salta, Av. Bolivia 5150, A4400AWE, Salta, Argentina. martinog@unsa.edu.ar

Elías R. de la Sota

Herbario LP, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s.n. B1900FWA, La Plata, Buenos Aires, Argentina. sota@museo.fcnym.unlp.edu.ar

ABSTRACT. A new species of *Pecluma* M. G. Price, *P. barituensis* O. Martínez & de la Sota, is described from material collected from the northwestern forests and mountain woods in Argentina. This taxon is related to *Pecluma oranensis* (de la Sota) de la Sota, from which it differs with its pinnae narrower at mid-rachis, thereby appearing farther apart, and its auriculiform basal pinnae.

RESUMEN. Se describe y se ilustra una nueva especie de *Pecluma*, *P. barituensis* O. Martínez & de la Sota, sobre material proveniente de las selvas y bosques montanos del noroeste argentino. Este taxón es afín a *Pecluma oranensis* (de la Sota) de la Sota, del que difiere principalmente por sus pinnas medias más delgadas y apareciendo así más distanciadas y las basales auriculiformes.

Key words: Argentina, *Pecluma*, Polypodiaceae.

The genus *Pecluma* M. G. Price (Polypodiaceae) includes more than 30 species that grow in American tropical and subtropical regions. In Argentina, Ponce (1996) quoted eight species, three of which grow in the Argentinean northwest region.

By reviewing Evans's (1969) and Price's (1983) works on the *Polypodium pectinatum-plumula* complex and on *Pecluma* and by floristic studies carried out on the American continents (de la Sota, 1973, 1977, 1995; Tryon & Tryon, 1982; Mickel & Beitel, 1988; Tryon & Stolze, 1993; Moran, 1995; de la Sota & Martínez, 1998; Salino, 1998; Mickel & Smith, 2004; Kessler & Smith, 2005), it is concluded that the specimen described here is a new taxon.

Pecluma barituensis O. Martínez & de la Sota, sp. nov. TYPE: Argentina. Prov. Salta: Dpto. Santa Victoria, Parque Nac. Baritú, 1500 m, 14 July 1999, O. G. Martínez & E. R. de la Sota 693 (holotype, MCNS; isotype, LP). Figure 1.

Plantae epiphyticae raro terrestres, affines *Peclumae oranensis* (de la Sota) de la Sota sed rhizomate breviter repente, frondibus gracilibus usque ad 70 cm longis, cum segmentis medianis plus angustis et plus distantibus et segmentis basalibus manifeste reductis ad auriculas parvulas, sporis 64, clare verrucosis differunt.

Epiphytic plants, rarely terrestrial; rhizome short-creeping, 3–5 mm diam., with clathrate scales that are deltoid to lanceolate, basifixed, brown, ovate to cordate, with entire margin, with and without trichomes at the back, to 1 cm. Fronds 50–70 cm, petioles 1/5–1/4 of the frond's length, terete, dark brown, articulate to the rhizome, with short phyllopo-dia, 1–2 mm; laminae elliptic to lanceolate, papyrus-like, slender, pectinate, 40–50 × 8–10 cm, attenuate at the base, gradually attenuate toward the tip; rachis dark, pilose with abundant appressed trichomes, ctenoid, multicellular, 0.3–0.5 mm, denser adaxially; pinnae to 85 pairs; basal pinnae reduced to auricles; pinnae on mid-rachis distant, linear to linear-lanceolate, perpendicular to the rachis, 4–5 × 0.2–0.5 cm, margin entire with hyaline trichomes, lamina with scattered clavate trichomes and setose trichomes over the costae. Lateral veins 1- or 2-forked, with circular superficial sori, 1 mm diam., with receptacles bearing simple or ramified trichomes, as paraphyses; sporangia glabrous; spores 64, spores monoletе reniform, clearly verrucose.

Geographic distribution and habitat. The new species is an epiphyte and occasionally terrestrial. It was observed as frequent in forests or humid montane woods, between 1200 and 1500 m, in Salta and Jujuy provinces in northwestern Argentina.

This species had been previously identified in Argentina as *Pecluma oranensis* because both species are found in montane forests or yungas. However, *P. barituensis* can be easily distinguished from *P. oranensis*, as can be seen in Table 1.

In addition, although it is not a diagnostic character, it should be noted that the pinnae are consistently outspread. Usually, in other poikilohydric taxa of this




Figure 1. *Pecluma barituensis* O. Martínez & de la Sota. —A. Habit. —B. Close-up of median pinna showing the venation pattern and the sori position. —C. Setose trichome from the rachis. —D. Ctenoid trichomes from the rachis. —E. Pair of pinnae. —F. Simple and ramified trichomes on the sori receptacles. —G. Rhizome scale, drawn without trichomes. —H. Rhizome scales, with trichomes. Drawn from the holotype O. G. Martínez & E. R. de la Sota 693 (MCNS).

Table 1. Morphological comparison of *Peculuma barituensis* and *P. oranensis*

	<i>P. barituensis</i>	<i>P. oranensis</i>
Pinnae on mid-rachis	4–5 × 0.2–0.5 cm	6–7 × 0.6–0.7 cm
Distance between pinnae	usually greater than their width	equal to their width
Basal pinnae	auriculiform	subdilated
Apical pinnae	gradually reduced, 8 to 10 pairs, perpendicular to the rachis	attenuated, 5 to 6 pairs, oblique to the rachis

genus, the pinnae are contracted to reduce the evapotranspirational surfaces, as an adaptation to decreased atmospheric relative humidity.

In this contribution, paraphyses are defined by the criteria given by de la Sota (1986), interpreting the indument of the sori receptacles.

Paratypes. ARGENTINA. **Salta:** Dpto. Anta, Parque Nac. El Rey, Arroyo Quina, *A. Brown 1355* (MCNS); Dpto. Santa Victoria, Parque Nac. Baritú, *O. G. Martínez & V. Aquino 1025* (MCNS), *1028* (MCNS), *1031* (MCNS), *O. G. Martínez, E. R. de la Sota, L. Novara & A. Ganem 685* (MCNS). **Jujuy:** Dpto. Ledesma, Parque Nac. Calilegua, camino a La Mesada y Abra de Cañas, *A. Brown 1714* (MCNS).

Acknowledgments. The authors acknowledge the curators of the consulted herbaria for kindly providing material and María del Carmen Otero for the illustration. This work was supported by grants from the National Council of Scientific and Technological Research (CONICET) and from the National University of Salta, for Project 1434.

Literature Cited

Evans, M. A. 1969. Interspecific relationship in the *Polypodium pectinatum-plumula* complex. *Ann. Missouri Bot. Gard.* 55: 193–293.

Kessler, M. & A. R. Smith. 2005. Seven new species, 13 new combinations, and one new name of Polypodiaceae from Bolivia. *Candollea* 60(1): 271–288.

Mickel, J. T. & J. M. Beitel. 1988. Pteridophyte Flora of Oaxaca, México. *Mem. New York Bot. Gard.* 46: 1–568.

——— & A. R. Smith. 2004. The Pteridophytes of Mexico. *Mem. New York Bot. Gard.* 88.

Moran, R. C. 1995. Polypodiaceae. Pp. 333–366 in G. Davidse, M. Sousa S. & S. Knapp (editors), *Flora Mesoamericana*, Vol. 1. Universidad Nacional Autónoma de México, México, D.F.; Missouri Botanical Garden, St. Louis; The Natural History Museum, London.

Ponce, M. M. 1996. Pteridophyta. Pp. 1–79 in F. O. Zuloaga & O. Morrone (editors), *Catálogo de las Plantas Vasculares de la República Argentina*. I. Monogr. Syst. Bot. Missouri Bot. Gard. 60.

Price, M. G. 1983. *Peculuma*, a new tropical American fern genus. *Amer. Fern J.* 73: 109–116.

Salino, A. 1998. New combinations in *Peculuma*. *Novon* 8: 296–297.

Sota, E. R. de la. 1973. Sinopsis de las Pteridófitas del Noroeste de Argentina, II. *Darwiniana* 18: 173–188.

———. 1977. Pteridophyta. In A. L. Cabrera (editor), *Flora de Provincia de Jujuy*. Colecc. Ci. Inst. Nac. Tecnol. Agropecu. 13(2).

———. 1986. Sobre la posición sistemática de *Polypodium fuscopunctatum* Hook. y *Polypodium percussum* Cav. (Polypodiaceae s. str., Pteridophyta). *Physis*, C 44(106): 19–28.

———. 1995. Novedades para la Flora Argentina. Nueva combinación en Polypodiaceae. *Hickenia* 2: 137.

——— & O. G. Martínez. 1998. Polypodiaceae. *Aport. Bot. Salta, Ser. Fl.* 5(8): 1–27.

Tryon, M. R. & R. G. Stolze. 1993. Polypodiaceae, Pteridophyta of Peru V. *Fieldiana, Bot.*, 32: 70–183.

——— & A. F. Tryon. 1982. *Fern and Allied Plants with Special Reference to Tropical America*. Springer-Verlag, Berlin.



Sota,
Eli

as R. de la and Martínez, Olga Gladys. 2007. "A new species of *Pecluma* (Pteridophyta--Polypodiaceae)." *Novon a journal of botanical nomenclature from the Missouri Botanical Garden* 17, 43–45.

[https://doi.org/10.3417/1055-3177\(2007\)17\[43:ANSOPP\]2.0.CO;2](https://doi.org/10.3417/1055-3177(2007)17[43:ANSOPP]2.0.CO;2).

View This Item Online: <https://www.biodiversitylibrary.org/item/41805>

DOI: [https://doi.org/10.3417/1055-3177\(2007\)17\[43:ANSOPP\]2.0.CO;2](https://doi.org/10.3417/1055-3177(2007)17[43:ANSOPP]2.0.CO;2)

Permalink: <https://www.biodiversitylibrary.org/partpdf/17522>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.