

NOTES ON NEW AND NOTEWORTHY PLANTS. CX

Harold N. Moldenke

SYNGONANTHUS UMBELLATUS f. *PROLIFERENS* Moldenke, f. nov.

Haec forma a forma typica speciei capitulis fructiferis saepe proliferentibus recedit.

This form differs from the typical form of the species in having at least some of its fruiting-heads more or less proliferous with clusters of green, linear, leaf-like structures to about 1 cm. long issuing from the center or from one side.

The type of the form was collected by Bassett Maguire, Julian A. Steyermark, and Celia K. Maguire (no. 53542) on a wet savanna 2 km. south of Río Chibau, on the righthand bank of the Río Chicanán, Bolívar, Venezuela, at 100 meters altitude, the "flowering heads gray-white", on September 1, 1962, and is deposited in the Britton Herbarium at the New York Botanical Garden.

SYNGONANTHUS UMBELLATUS f. *STELLARIS* Moldenke, f. nov.

Haec forma a forma typica speciei bracteis involucrantibus saepe elongatus linearibus foliaceis usque 1 cm. longis stellato-radiatis recedit.

This form differs from the typical form of the species in having some of its fruiting-heads subtended by an involucre or semi-involucre composed of 2—10 stellately radiating, green, foliaceous, linear-elongate bracts to about 1 cm. long.

The type of the form was collected by James L. Zarucchi (no. 2048) in wet open areas in a low-tree forest adjacent to open savannas at Mitú and vicinity along the Río Vaupés at Circassia, Vaupés, Colombia, on September 14, 1976, and is deposited in my personal herbarium.

VERBENA WRIGHTII var. *INTERMEDIA* Moldenke, var. nov.

Haec varietas a forma typica speciei laminis foliorum ovatis irregulariter inciso-lobulatis lobulis antrorsia acutis recedit.

This variety differs from the typical form of the species in having its leaf-blades ovate in outline, marginally merely incised-lobulate, the lobules antrorsely ascending, broad, short, and apically acute.

The type of the variety was collected by Noel H. and Patricia K. Holmgren (no. 6784) in a pinyon-juniper-oak woodland on an open southwest-facing slope, at 5000 feet altitude, on the Baboquivari Trail, Baboquivari Mountains, 19 airline miles east-southeast of Sells, T. 19 S, R. 7 E, m Sec. 15, Pima County, Arizona, on April 13, 1973, and is deposited in the Britton Herbarium at the New York Botanical Garden. The collectors assert that the plant was "locally common to frequent", the corolla pink, withering blue. The calyx characters are those of *V. wrightii* A. Gray, but the foliage is reminiscent of that of *V. gooddingii* var. *nepetifolia* Tidestr.



Moldenke, Harold N. 1978. "NOTES ON NEW AND NOTEWORTHY PLANTS PART 110." *Phytologia* 39, 161–161.

View This Item Online: <https://www.biodiversitylibrary.org/item/47392>

Permalink: <https://www.biodiversitylibrary.org/partpdf/176354>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.