Lecanium pallidior, Ckll. & King, n. sp.

♀ Scale. 3 mm. long, 2 broad, 1½ to 2 high, very convex, very shiny, wrinkled; light yellowish-brown or ochreous, paler at the sides than dorsally. This is a very much paler scale than *fletcheri*, and when boiled in caustic potash it becomes very pale and transparent, while *fletcheri* remains dark brown.

Antennae. Six-segmented, the segments measuring as follows in μ (I.) 34-5I. (2.) 28-3I. (3.) 42-45. (4.) 34. (5.) 23-3I. (6.) 34-42. Mr. King reports a 7-segmented antenna, with the last three segments measuring respectively 12, 16 and 38 μ . This agrees well enough with *fletcheri*, which I find always to have 7 segments, the last three 14-20, 14-17 and 39-42 μ . There is also this in common with *fletcheri*, that the second segment is shorter than the first or third.

Legs. The different legs are similar in type, but the tibiae and tarsi are variable. Four legs measured gave as follows — $(\mu$.)

					(1.)	(2.)	(3.)	(4.)
Coxa.					87.	85.	87.	85.
Femur+	- tro	char	nter.		116.	116.	116.	118.
Tibia.					110.	76.	82.	68.
Tarsus +	- clay	w.			65.	76.	- 90.	79.

TWO NEW COCCIDS FROM BERMUDA.

IN January of this year, the writer in his search for coccids infesting green-house plants, observed a potted plant in one of the conservatories visited, trimmed quite close to its roots, and the remaining stumps well covered with a small clear white scale. The plant in question was recognized at once to be *Cycas revoluta*. Upon enquiry it was learned that this plant was imported from Bermuda last year. Some of the old stems with the coccids were taken for study. One of the species proved to be *Aspidiotus hederae* Vallot, a very common species through the United States, and the other coccid was The first is an anterior leg. These dimensions are very different from those of *L*. *fletcheri*, as will be seen by comparing the statistics given below.

Lecanium fletcheri, Ckll.

Antennae. (µ .) Segme Mass. Canadian	spn.		5. 42. 2	. 6. 7. 0. 17. 42. 4. 14. 39.
Legs. (µ .)	Coxa.	Femur+ troch.	Tibia.	Tarsus + claw.
Mass. spn.	113.	127.	87.	85.
Canadian; front leg.	56.	107.	90.	85.
" middle leg.	121.	144.	96.	93 -
" hind leg.	130.	141.	93.	90.
" hind leg.	121.	130.	93.	96.

The front legs of *fletcheri* appear thus to differ appreciably from the other four as regards the coxa and femur. Both coxa and femur of the middle and hind legs are noticeably longer than in *pallidior*.

Lecanium pallidior was found by Mr. King at Methuen, Mass., Nov. 15, 1898, on small twigs of *Chamaecyparis thyoides* (L). The leaves and small limbs were thickly covered at that date with young larvae. The species belongs, of course, to Eulecanium.

new to the writer, who sent a mount and some of the scales to Prof. Cockerell for determination. He identifies it as Aulacaspis elegans Leon. Described as Howardia elegans, and only known hitherto from Portici Italy, on Cycas revoluta. The & and & scale are unknown. The species differs from typical Aulacaspis by lacking the circumgenital glands. The occurrence of this species is of much interest and the coccids of Bermuda are almost unknown. There are only two other species known to the writer recorded from those islands, Mytilaspi citricola Pack. on orange and lime trees, and Chionaspis citri Comst. on orange. Geo. B.King. Lawrence, Mass., Feb. 27, 1899.



King, George B. 1899. "Two New Coccids From Bermuda." *Psyche* 8, 350–350. <u>https://doi.org/10.1155/1899/45343</u>.

View This Item Online: https://doi.org/10.1155/1899/45343 DOI: https://doi.org/10.1155/1899/45343 Permalink: https://www.biodiversitylibrary.org/partpdf/183253

Holding Institution Smithsonian Libraries and Archives

Sponsored by Smithsonian

Copyright & Reuse Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.