

specially interested in the Greenfield field excursions. Will those who plan to attend kindly *notify the chairman before April 15?*

A detailed announcement of trains, hotel arrangements, etc., will be issued later.

If the Greenfield trip proves satisfactory other similar outings will be arranged.

C. H. KNOWLTON,	}	<i>Committee on Field Excursions.</i>
Hingham, Mass.		
M. L. FERNALD		
F. G. FLOYD		

THE NEW ENGLAND FEDERATION OF NATURAL HISTORY SOCIETIES will hold its annual meeting Friday and Saturday, April 26 and 27, at the building of the Boston Society of Natural History. The exhibition will be open both days and there will be sessions Friday evening and Saturday morning.—J. H. EMERTON.

REPORTS ON THE FLORA OF THE BOSTON DISTRICT,—XIV.

CYPERACEAE.

CAREX, L. TO N.

C. lanuginosa Michx. Swamps and low meadows, common.

C. laxiculmis Schwein. Rich damp woods; occasional near Boston, also found in West Newbury and Andover.

C. laxiculmis Schwein., var. **copulata** (Bailey) Fernald. Haycock Meadow, Medford (*Wm. Boott*, July 5, 1853).

C. laxiflora Lam. Rich woods and meadows; well distributed throughout, but not very common.

C. laxiflora Lam., var. **blanda** (Dewey) Boott. Abundant near Boston, also collected at Framingham; West Newbury, according to Robinson, Fl. Essex Co., 120, 180.



Emerton, J. H. 1912. "The New England Federation of Natural History Societies." *Rhodora* 14, 76–76.

View This Item Online: <https://www.biodiversitylibrary.org/item/14485>

Permalink: <https://www.biodiversitylibrary.org/partpdf/187771>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.