

REPORTS ON THE FLORA OF THE BOSTON  
DISTRICT,— XI.

## CYPERACEAE.

## ERIOPHORUM.

**E. callitrix** Cham. Cold bogs; reported from several scattered stations.

**E. gracile** Roth. Cold bogs and swamps. Nine scattered stations in northeastern and central portions. An early bloomer which has escaped many collectors.

**E. tenellum** Nutt. Boggy meadows, well distributed.

**E. virginicum** L. Bogs and wet meadows; common throughout.

**E. virginicum** L., var. **album** Gray. Danvers (*J. H. Sears*, Sept. 1, 1885); Harland St. meadows, Milton (*J. R. Churchill*, Sept. 1, 1895).

**E. viridi-carinatum** (Engelm.) Fernald. Boggy meadows and swamps. Not reported from the extreme southeastern portion, but abundant elsewhere.

## FUIRENA.

**F. squarrosa** Michx. Shores of Round Pond, Tewksbury, 1854 (*Wm. Boott*) to the present.

## HEMICARPHA.

**H. micrantha** (Vahl) Pax. Sandy shores of ponds, rare; North Andover, Andover, Winchester, Concord, Wayland, Framingham, Sherborn.

## RYNCHOSPORA.

**R. alba** (L.) Vahl. Meadows and bogs; not reported from the extreme southeastern portion, but abundant elsewhere.

**R. alba** (L.) Vahl, var. **macra** Clarke. Fresh Pond, Cambridge (*Wm. Boott*, Aug. 20, 1853); Round Pond, Tewksbury (*M. L. Fernald*, Sept. 18, 1909).



**R. fusca** (L.) Ait. f. Low ground, frequent in northern three quarters of district, doubtless throughout.

**R. glomerata** (L.) Vahl. Moist soil, abundant throughout.

**R. macrostachya** Torr. Great Pond, South Weymouth (*Wm. Boott*, Sept. 2, 1870; *E. & C. E. Faxon*, Sept. 19, 1878; *G. G. Kennedy*, Sept. 9, 1908).

**R. macrostachya** Torr., var. **inundata** (Oakes) Fernald. Great Pond, South Weymouth (*E. Faxon*, Sept. 19, 1878; *G. G. Kennedy*, Sept. 9, 1908).

#### CLADIUM.

**C. mariscoides** (Muhl.) Torr. Margins of ponds and wet places, common throughout. Collected by E. F. Williams in a *salt-marsh* at Scituate.

#### SCLERIA.

**S. reticularis** Michx. Wet sandy and gravelly shore, Winter Pond, Winchester, from 1858 (*Wm. Boott*) to the present.

**S. triglomerata** Michx. Open place in dry woods near a dry watercourse, South Sudbury (*W. P. Rich*, Aug. 8, 1899); Concord (*E. S. Hoar*, according to Dame & Collins, *Fl. Middlesex Co.* 114, 1888).

C. H. KNOWLTON	} Committee on Local Flora.
J. A. CUSHMAN	
WALTER DEANE	
A. K. HARRISON	

THREE ADDITIONAL PLANTS FROM NANTUCKET.—The following plants may be of interest as not being previously recorded from Nantucket. Specimens of *Polypodium vulgare* L. were found by Miss Grace Gardner of Nantucket last summer. *Corallorhiza maculata* Raf. was found by Mr. Nelson, a Cornell student, who spent the summer on the island. In 1909 I found some patches of *Glaux maritima* L. on Tuckernuck Island and in 1910 found an abundance of it in the marshy area about the south inlet of the Island. It spreads over a large area along the shore and is of the small-leaved typical form. Specimens of these three species are in the herbarium of the Nantucket Maria Mitchell Association at Nantucket.—JOSEPH A. CUSHMAN, Boston Society of Natural History.



Knowlton, Clarence Hinckley et al. 1911. "REPORTS ON THE FLORA OF THE BOSTON DISTRICT,— XI." *Rhodora* 13, 104–105.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/14484>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/187870>

**Holding Institution**

Missouri Botanical Garden, Peter H. Raven Library

**Sponsored by**

Missouri Botanical Garden

**Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.