

BROMUS CILIATUS L., var. **denudatus** (Wiegand), n. comb.,—*B. ciliatus*, forma *denudatus* Wiegand, RHODORA, xxiv. 91 (1922).

When Professor Wiegand published var. *denudatus* merely as a form, he spoke of it as occurring "Throughout the range of the typical form" of the species, the plant with the median and upper sheaths villous or pilose. In my own experience, however, I have long considered the plant with glabrous upper sheaths not only as of more boreal general range but as having a strong tendency to shorter and broader leaf-blades; and a study of the collections of the Gray Herbarium and of the New England Botanical Club shows these tendencies to be even stronger than I had supposed. For instance, selecting certain northeastern areas the floras of which have been much collected, I find that these two herbaria have from the selected colder areas: 8 sheets of typical *B. ciliatus* from Newfoundland but 23 of var. *denudatus*; 0 sheets of typical *B. ciliatus* from the Gaspé Peninsula but 19 of var. *denudatus*; from the east coast of Maine 4 of typical *B. ciliatus*, 12 of var. *denudatus*. from Willoughby, Vermont 0 typical *B. ciliatus*, 7 of var. *denudatus*. For selected warmer areas the figures are: central and southern Maine, typical *B. ciliatus* 36, var. *denudatus* 11; central and southern New Hampshire, typical *B. ciliatus* 16, var. *denudatus* 0; Connecticut, typical *B. ciliatus* 10, var. *denudatus* 0. In western Newfoundland var. *denudatus* ascends to the high mountain-meadows, typical *B. ciliatus* being confined to low altitudes; on the Gaspé Peninsula, where no typical *B. ciliatus* has been found, var. *denudatus* ascends to the highest alpine meadows of the Shick-shock Mts. (1000–1125 m.) and in the White Mountain region, where the plant with pilose sheaths is confined to the lower altitudes, var. *denudatus* reaches the Fan of Huntington Ravine.—M. L. FERNALD, Gray Herbarium.

The date of the December issue (unpublished as this goes to press) will be announced later.



Fernald, Merritt Lyndon. 1926. "Bromus ciliatus L., var. denudatus (Wiegand), n. comb." *Rhodora* 28, 20–20.

View This Item Online: <https://www.biodiversitylibrary.org/item/14498>

Permalink: <https://www.biodiversitylibrary.org/partpdf/187885>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.