The rootlets are not formed till afterwards upon the outside of it. Another proof that the *Lycopodia* cannot be said to have a main-root.

4. *The terminal bud becomes inclosed by an envelope (two bud-envelopes, Knospenhüllblättern).* These remind us of the cotyledons of the Dicotyledons. At all events those bud-envelopes are identical with it, which so often occur in the Phanerogamia, *e. g.* in many *Leguminosæ*, as in *Amicia*, *Lathyrus*, *Pisum*, in *Agri-monia*, in *Cunonia capensis*, especially, exactly as in *Selaginella denticulata*, in the *Begonias*, &c. Here these enveloping leaves have usually been looked upon as stipules. It is impossible that they can be such; for they are distinguished from true stipules, *e. g.* in *Roses* by the fact that they always inclose axial buds, and always surrounding the stem at the origin of the internode, they persist or drop off (*Cunonia*), while the stipules, mere appendicular organs, rest upon the leaves. This is alluded to in another place.

5. *The Selaginellæ possess a main-axis.* This however remains very short and divides at once in the terminal bud into two branches, which in like manner always divide again dichotomously. Hence the "caulis dichotomo-procumbens."

6. *The leaf is merely an "outshoot" (Aufreibung) of the stem, which expands into a flattened body.*

7. *In Selaginella denticulata an accessory organ also is found between the axis and the leaf; the import of this is as yet doubtful.* In more recent researches in living *Selaginellæ* I found it also in *Selaginella stellata*, *opus* and *viticulosa*, and of exactly the same structure. It thus becomes probable that it extends throughout the whole of the genus *Selaginella*.

8. Among the significant peculiarities in *Selaginella* is also to be included the extremely regular position of the branch-roots. I mentioned above that they are formed upon the upper side of the stem in *Selag. denticulata*, and that Schleiden, without referring to the species, says that he has observed them on the under side. Since this was printed I have become acquainted with several living *Selaginellæ* in the Berlin Botanic Garden, and I found in one of them, in *Selag. stellata*, Spr., a confirmation of Schleiden's assertion. I thus had an opportunity to seek for the law which prevails here. It proves to be as follows: the roots of the branches make their appearance both on the upper and under side of the stem, but always just inferior to the forked division of a branch, and indeed within its last leaf. If this be on the upper side—and then it is a folium intermedium—there, also, the branch-root appears. But if the leaf be on the under side—and it is then a larger leaf—the branch-root also is formed on the under side always between the stem and leaf, consequently as a true branch.

9. *The oophoridium in Selaginella denticulata, and all those Selaginellæ where it stands at the base of the spike, is always a metamorphosed main-branch.* This is the case in *Selag. ferruminata*, Sprg. (= *elongata*, Klotzsch), *lucidinervia*, Sprg. (= *pedata*, Kl.), *conduplicata*, ej., *distorta*, ej., *Poppigiana*, ej., *sulcata*, ej., *stolonifera*, ej., *deliquescens*, ej., *puberula*, Kl., *laevigata*, Sprg., *articulata*, &c. Whether this law may be modified in those species where antheridia and oophoridia occur on one spike, whether it is here a transformed secondary branch, e. g. in *S. stellata*, Sprg., *pygmæa*, ej., *helvetica*, ej., &c., yet remains to be investigated. In other *Selaginellæ*, the fruit of which is dioecious, where therefore oophoridia and antheridia occur on distinct spikes, the oophoridium appears to be a metamorphosed main-branch, e. g. in *S. selaginoides*—certain it is however that the oophoridium is never a leaf-product.

The oophoridium-spores, more properly the ovules, are formed at the extremity of the vascular bundle which projects free into the interior of the oophoridium as a hollow mother-cell. The course of its development still requires to be more minutely investigated.

10. *The antheridium is a metamorphosed twig-bud* (Zweigknospe) developed simultaneously with the leaves from the axis, a twig here being understood to be a secondary branch.* The antheridium is never a leaf-product. Here refer also the sporangium of *Psilotum* and *Tmesipteris*.

The antheridium-spores originate by fours in a mother-cell, and certainly by cytoblast-formation as understood by Schleiden (formation of a membrane around a nucleus), never by division of the cytoblastema.

EXPLANATION OF PLATES II. TO VI.

PLATE II.

Fig. 1. The germinating spore: *a*, the rootlet; *b*, the terminal bud; *c*, the germ. 250 †.

Fig. 2. The terminal bud so seen that the other branch-bud lies behind the one *a*. 250.

Fig. 3. The same, unfolded artificially. 250.

Fig. 4. The whole germinating plant, removed from the spore: *a*, the point of vegetation for the stem and root; *b*, the germ (*Keim-körper*). 50.

Fig. 5. The germinating plant with the bud-leaves. 10.

Fig. 6. A bud-enveloping leaf, with a branch-bud *a*; *b*, an accessory organ. 250.

Fig. 7. The base of the bud-envelope. 400.

* The word *Zweigknospe*, literally twig-bud, is here used in distinction to *Astknospe* or branch-bud, which latter may be regarded as determining the bifurcation of the main axis, while the former is a secondary ramification.—TR.

† The figures give the magnifying power.

Pouchet, G. and Guerne, Jules de. 1887. "On the food of the sardine." *The Annals and magazine of natural history; zoology, botany, and geology* 19, 323–324. <https://doi.org/10.1080/00222938709460251>.

View This Item Online: <https://www.biodiversitylibrary.org/item/19437>

DOI: <https://doi.org/10.1080/00222938709460251>

Permalink: <https://www.biodiversitylibrary.org/partpdf/18957>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.