lem alii deducant, ego non. Lente et modeste perpendant Medusæ ne virtutem hanc obtineant ex aqua, vel hæc a Medusis immensis scilicet Oceanus minimæ sui parti tanta miracula debbit !”

XLIV.—*Characters of undescribed Chalcidites collected in North America by E. Doubleday, Esq., and now in the British Museum.* By FRANCIS WALKER, F.L.S.

1. *Pteromalus Lausus*, fem. *Niger, abdomen viridi cupreo et purpureo varium, antennæ piceæ, pedes flavo-fulvi, femora nigra, alæ sublimpidæ.*

Corpus breve, crassum, convexum, nigrum, obscurum, scitissime squameum, pilis albis hirtum: caput transversum, breve, thoracis latitudine; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: os piceum: antennæ piceæ, graciles, extrorsum crassiores, thorace paullo longiores; articulus 1^{us} longus, linearis, fulvus; 2^{us} fuscus, cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes sat longi, usque ad 10^{um} curtantes; clava fusiformis, apice ferruginea, articulo 10^o paullo latior et multo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum brevi-obconicum: metathorax brevis, declivis: petiolus brevissimus: abdomen ovatum, nitens, læve, nigro-viride, basi cupreo et purpureo varium, supra planum, subtus profunde carinatum, apice acuminatum, thorace paullo longius vix angustius; segmentum 1^{um} magnum; 2^{um} et sequentia breviora, subæqualia: pedes nigri, simplices, subæquales; trochanteres picei; genua fulva; protarsi fulvi; mesotarsi et metatarsi flavi, apice fusci: alæ sublimpidæ; squamulæ piceæ; nervi fusci; nervus humeralis fulvus ulnari paullo longior, ulnaris valida radiali fere duplo longior, cubitalis radiali paullo brevior; stigma sat magnum, obscurius. (Corp. long. lin. $1\frac{3}{4}$; alar. lin. 3.)

It is allied to *Pteromalus intermedius*, *Eutelus intermedius*, Ent. Mag. ii. 366.

2. *Pteromalus Cratylus*, fem. *Ater, abdomen nigro-æneum, antennæ piceæ, pedes flavi, femora nigra, alæ limpidæ.*

Corpus breve, latum, crassum, convexum, atrum, obscurum, scitissime squameum, parce hirtum: caput transversum, breve, thorace paullo latius; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: antennæ piceæ, subclavatæ, graciles, thorace paullo longiores; articulus 1^{us} fulvus, longus, linearis; 2^{us} cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes mediocres, usque ad 10^{um} curtantes et latescentes; clava conica, acuminata, articulo 10^o duplo longior: thorax brevi-ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum obconicum: metathorax brevis, declivis: petiolus brevissimus: abdomen nigro-æneum, brevi-ovatum, nitens,

læve, supra planum, subtus carinatum, apice acuminatum, thoracis longitudine et latitudine: pedes flavi, simplices, subæquales; coxæ nigrae; femora nigra, apice flava; tarsi apice fusci: alæ limpidæ; squamulæ fulvæ; nervi flavi; nervus humeralis ulnari multo longior, radialis ulnari brevior cubitali vix longior; stigma minutum. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. 2.)

This is also allied to *Pt. intermedius*.

3. *Pteromalus Cassotis*, fem. *Ater, abdomen cupreum, antennæ piceæ, pedes fulvi, tarsi flavi, alæ limpidæ.*

Corpus breve, latum, convexum, atrum, obscurum, scitissime squameum, parce hirtum: caput transversum, breve, thorace latius; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ piceæ, clavatæ, thorace non longiores; articulus 1^{us} fulvus, longus, gracilis; 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes breves, usque ad 10^{um} paullo curtantes et latescentes; clava conica, acuminata, articulo 10^o paullo latior et duplo longior: thorax ovatus: prothorax transversus, brevis: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum subconicum: metathorax mediocris, declivis, obconicus: petiolus brevissimus: abdomen brevi-ovatum, nitens, læve, cupreum, basi viridi cupreo et purpureo micans, supra planum, subtus carinatum, apice acuminatum, thorace brevius non angustius: pedes fulvi, simplices, subæquales; coxæ nigrae; mesotarsi et metatarsi flavi, apice fusci: alæ limpidæ; squamulæ fulvæ; nervi fulvi; nervus humeralis ulnari multo longior, radialis cubitali longior; stigma parvum. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. 2.)

Allied to *Pt. deplanatus*, Nees d'Essenbeck Hym. Ich. aff. ii. 110.

4. *Pteromalus Euthymus*, mas. *Ater, abdomen nigro-cupreum, antennæ piceæ, pedes fulvi, femora picea, tarsi flavi, alæ limpidæ.*

Corpus breve, latum, convexum, nigrum, obscurum, scitissime squameum, parce hirtum: caput transversum, breve, magnum, thorace latius; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ piceæ, clavatæ, thorace non longiores; articulus 1^{us} fulvus, longus, gracilis; 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes usque ad 10^{um} breves, subæquales, discreti; clava conica, acuminata, articulo 10^o paullo latior et duplo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ non bene determinatæ; scutellum subconicum: metathorax mediocris, obconicus, declivis: petiolus brevissimus: abdomen subconicum, depressum, nitens, nigro-cupreum, læve, basi angustum, thorace brevius et multo angustius et humilior; segmentum 1^{um} magnum, 2^{um} et sequentia brevia: pedes fulvi, simplices, subæquales; coxæ nigrae; femora picea, basi et apice fulva; mesotarsi et metatarsi flavi, apice fusci: alæ limpidæ; squamulæ piceæ; nervi fulvi; nervus humeralis ulnari multo longior, radialis ulnari paullo brevior cubitali longior; stigma fuscum, parvum. (Corp. long. lin. $\frac{3}{4}$; alar. lin. $1\frac{1}{2}$.)

Allied to *Psilocera obscura*, Ent. Mag. i. 374.

5. *Pteromalus Dipsas*, fem. *Nigro-æneus, pedes fulvo-picei, femora nigro-picea, tarsi flavi, alæ limpidae.*

Corpus nigro-æneum, convexum, obscurum, scitissime squameum, parce hirtum: caput transversum, breve, thoracis latitudine; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: antennis articulus 1^{us} fulvus, longus, gracilis; 2^{us} fuscus: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum subrotundum: metathorax brevis, declivis: petiolus brevissimus: abdomen ovatum, nitens, læve, supra planum, subtus carinatum, apice acuminatum, thorace paullo brevius et angustius: pedes fulvi, simplices, subæquales; coxæ nigræ; trochanteres picei; femora nigro-picea; mesopedum et metapedum tibiæ fuscae, tarsi flavi apice fusci: alæ limpidae; squamulæ piceæ; nervus humeralis ulnari multo longior, radialis ulnari multo brevior cubitali paullo longior; stigma parvum. (Corp. long. lin. 1; alar. lin. $1\frac{3}{4}$.)

Allied to *Pt. discolor*, Ent. Mag. iii. 473.

6. *Pteromalus Hermeas*, fem. *Niger, antennæ nigræ, pedes fulvi, femora nigra, tarsi flavi, alæ limpidae.*

Corpus angustum, convexum, nigrum, parum nitens, scitissime squameum, parce hirtum: caput transversum, breve, thorace paullo latius; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: antennæ nigræ, graciles, subclavatae, thorace non longiores; articulus 1^{us} longus, linearis; 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes mediocres, usque ad 10^{um} curtantes et latescentes; clava conica, acuminata, articulo 10^o multo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum brevi-obconicum: metathorax brevis, declivis: petiolus brevissimus: abdomen fusiforme, nitens, læve, supra depressum, subtus profunde carinatum, apice acuminatum, thorace longius et angustius: pedes fulvi; coxæ nigræ; trochanteres picei; femora nigra; mesotarsi et metatarsi flavi, apice fusci: alæ limpidae; squamulæ fulvæ; nervi flavi; nervus humeralis ulnari multo longior, radialis ulnari brevior cubitali longior; stigma minimum. (Corp. long. lin. $\frac{3}{4}$; alar. lin. $1\frac{1}{4}$.)

Allied to *Pt. hemipterus*, Ent. Mag. iii. 196.

7. *Pteromalus Epicles*, fem. *Cyaneo-viridis, abdomen cupreo et purpureo varium, antennæ nigræ, pedes fulvi, femora nigro-picea, tarsi albidi, alæ albo limpidae.*

Corpus breve, crassum, latum, convexum, cyaneo-viride, parum nitens, scitissime squameum, parce hirtum: caput transversum, breve, thorace vix latius; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennis articulus 1^{us} longus, gracilis, niger, basi fulvus: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ non bene determinatæ, postice approximatae; scutellum obconicum: metathorax brevis, declivis: petiolus brevissimus:

abdomen ovatum, nitens, læve, cyaneo cupreo et purpureo varium, supra planum, subtus profunde carinatum, apice acuminatum, thorace angustius non longius : oviductus fulvus abdominis basi subtus emissus : pedes fulvi ; coxæ nigro-virides ; trochanteres picei ; femora nigro-picea ; mesotarsi et metatarsi albidii, apice fusci : alæ albido limpidæ ; squamulæ fulvæ ; nervi pallide flavi ; nervus humeralis ulnari multo longior, radialis ulnari brevior cubitali longior ; stigma minutum. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. 2.)

Allied to *Pteromalus altus*, *Eutelus altus*, Ent. Mag. ii. 367.

8. *Pteromalus Damo*, fem. *Cupreus*, abdomen basi viridi varium, pedes fulvi, tarsi flavi, alæ subfulvæ.

Corpus angustum, sublineare, convexum, cupreum, parum nitens, scitissime squameum, parce hirtum : caput ademptum : thorax ovatus : prothorax transversus, brevissimus : mesothoracis scutum longitudine latius ; parapsidum suturæ vix conspicuæ ; scutellum obconicum : metathorax mediocris, obconicus, declivis : petiolus brevissimus : abdomen fusiforme, nitens, læve, basi viridi varium, supra depressum, subtus profunde carinatum, apice acuminatum, thorace paullo angustius et multo longius : pedes fulvi ; coxæ cupreæ ; mesotarsi et metatarsi flavi, apice fusci : alæ subfulvæ ; squamulæ fulvæ ; nervi flavi ; nervus humeralis ulnari multo longior, radialis ulnari brevior cubitali longior ; stigma minutum. (Corp. long. lin. $\frac{3}{4}$; alar. lin. $1\frac{1}{4}$.)

Allied to *Pt. tenuis*, Ent. Mag. ii. 498.

9. *Pteromalus Cæx*, fem. *Nigro-æneus*, abdomen æneum, antennæ fusæ, pedes fulvi, tarsi flavi, alæ limpidæ.

Corpus angustum, nigro-æneum, convexum, nitens, scitissime squameum, parce hirtum : caput transversum, breve, thorace latius ; vertex latus ; frons impressa, abrupte declivis : oculi picei, mediocres, non extantes : antennæ fusæ, clavatæ, thorace non longiores ; articulus 1^{us} fulvus, longus, gracilis ; 2^{us} cyathiformis ; 3^{us} et 4^{us} minuti ; 5^{us} et sequentes transversii, usque ad 10^{um} curtantes ; clava ovata, acuminata, articulo 10^o latior et duplo longior : thorax ovatus : prothorax transversus, brevissimus : mesothoracis scutum longitudine latius ; parapsidum suturæ vix conspicuæ ; scutellum obconicum : metathorax mediocris, declivis, obconicus : petiolus brevissimus : abdomen fusiforme, æneum, læve, supra depressum, subtus profunde carinatum, apice attenuatum et acuminatum, thorace angustius et multo longius : pedes fulvi, simplices, subæquales ; coxæ æneæ ; mesotarsi et metatarsi flavi, apice obscuriores : alæ limpidæ ; squamulæ piceæ ; nervi fulvi ; nervus humeralis ulnari multo longior, radialis ulnari brevior, cubitalis sat longus ; stigma minutum. (Corp. long. lin. $\frac{3}{4}$; alar. lin. $1\frac{1}{4}$.)

Perhaps a variety of *Pt. Damo*.

10. *Pteromalus Doryssus*, mas. *Niger*, antennæ piceæ, pedes fulvi, femora nigro-picea, tarsi flavi, alæ limpidæ.

Corpus angustum, convexum, nigrum, parum nitens, scitissime squameum, parce hirtum : caput transversum, breve, thorace paullo

latius; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: antennæ piceæ, validæ, subfiliformes, thorace non longiores; articulus 1^{us} fulvus, longus, gracilis; 2^{us} cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes usque ad 10^{um} mediocres, subæquales; clava conica, compressa, acuminata, articulo 10^o multo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum obconicum: metathorax sat magnus, obconicus, declivis: petiolus brevissimus: abdomen sublineare, nitens, læve, depressum, thorace multo angustius non brevius: pedes fulvi, simplices, subæquales; coxæ nigræ; femora nigro-picea; meso- et metatarsi flavi, apice fusci: alæ limpidæ; squamulæ piceæ; nervi fulvi; nervus humeralis ulnari multo longior, radialis ulnari brevior, cubitali longior; stigma minutum. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. $2\frac{1}{4}$.)

Allied to *Pteromalus Zonaras*, Mon. Chal. i. 227.

11. *Pteromalus Timæa*, fem. *Ater, abdomen nigro-purpureum, antennæ piceæ, pedes nigri, tarsi flavi, alæ limpidæ.*

Corpus crassum, atrum, convexum, parum nitens, scitissime squameum, parce hirtum: caput transversum, breve, thorace paullo latius; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: antennæ piceæ, clavatæ, thorace non longiores; articulus 1^{us} niger, longus, gracilis; 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes mediocres, usque ad 10^{um} curtantes et latescentes; clava ovata, articulo 10^o latior et duplo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ non bene determinatæ; scutellum brevi-obconicum: metathorax brevis, declivis: petiolus brevissimus: abdomen fusiforme, nigro-purpureum, nitens, læve, basi cyaneo-micans, supra depressum, subtus carinatum, apice acuminatum, thorace paullo longius et multo angustius: pedes nigri; trochanteres picei; genua fulva; protarsi fulvi; mesotarsi et metatarsi flavi, apice fusci: alæ limpidæ; squamulæ piceæ; nervi flavi; nervus humeralis ulnari multo longior, radialis ulnari brevior cubitali longior; stigma minutum. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. 2.)

12. *Pteromalus Orontas*, fem. *Ater, abdomen viride, discus purpureus, antennæ nigræ, pedes nigro-picei, tarsi flavi, alæ limpidæ.*

Corpus angustum, convexum, nigrum, parum nitens, scitissime squameum, parce hirtum: caput transversum, breve, thorace paullo latius; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: antennæ clavatæ, nigræ, thorace non longiores; articulus 1^{us} longus, gracilis, 2^{us} longicyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes mediocres, usque ad 10^{um} latescentes; clava ovata, acuminata, articulo 10^o latior et duplo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum subconicum: metathorax brevis, declivis: petiolus brevissimus: abdomen longiovatum, viride, nitens, læve, supra depressum, subtus carinatum,

apice acuminatum, thorace angustius et longius; discus purpureus: pedes nigri, simplices, subæquales; trochanteres picei; genua fulva; tibiæ piceæ; protarsi fulvi; mesotarsi et metatarsi flavi, apice fusci: alæ limpidæ; squamulæ piceæ; nervi flavi; nervus humeralis ulnari longior, radialis ulnari brevior cubitali longior; stigma minutum. (Corp. long. lin. 1; alar. lin. $1\frac{1}{2}$.)

13. *Pteromalus Dymnus*, fem. *Ater*, abdomen viridi-æneum, antennæ piceæ, pedes nigri, tarsi flavi, alæ limpidæ.

Corpus validum, convexum, nigrum, parum nitens, scitissime squameum, parce hirtum: caput transversum, breve, thoracis latitudine; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: antennæ piceæ, clavatæ, thorace non longiores; articulus 1^{us} longus, gracilis; 2^{us} cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes mediocres, usque ad 10^{um} curtantes; clava conica, articulo 10^o multo longior: thorax ovatus, pilis albis hirtus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ: metathorax brevis, declivis: petiolus brevissimus: abdomen longiovatum, viridi-æneum, nitens, læve, supra depressum, subtus carinatum, apice acuminatum, thorace paullo longius et angustius: pedes nigri, simplices, subæquales; trochanteres picei; genua fulva; protarsi fulvi; mesotarsi et metatarsi flavi, apice fusci: alæ limpidæ, sat latæ; squamulæ piceæ; nervi pallide flavi; nervus humeralis ulnari multo longior, radialis ulnari brevior cubitali longior; stigma fuscum, parvum. (Corp. long. lin. 1; alar. lin. $1\frac{3}{4}$.)

14. *Pteromalus Eryx*, mas. *Ater*, abdomen cyaneum, discus cupreus, pedes flavi, metatibiæ fusco cinctæ, alæ limpidæ.

Corpus sublineare, convexum, nigrum, obscurum, scitissime squameum, parce hirtum, subtus nigro-cyaneum: caput ademptum: thorax longiovatus, pilis albis hirtus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ non bene determinatæ; scutellum brevi-conicum: metathorax mediocris, obconicus, declivis: petiolus brevissimus: abdomen sublineare, cyaneum, nitens, læve, depressum, thorace angustius non longius; discus cupreus: sexualia fulva: pedes pallide flavi, simplices, subæquales; coxæ cyaneæ; trochanteres picei; metatibiæ fusco-cinctæ; protarsi fulvi: alæ limpidæ; squamulæ piceæ; nervi pallide flavi; nervus humeralis ulnari multo longior, radialis ulnari brevior cubitali longior; stigma minutum. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. $2\frac{1}{4}$.)

15. *Pteromalus Hybreas*, fem. *Æneo-ater*, abdomen cupreum, antennæ nigræ, pedes fulvi, tarsi flavi, alæ limpidæ.

Corpus sublineare, æneo-atrum, convexum, parum nitens, scitissime squameum, pilis albis hirtum: caput transversum, breve, magnum, thorace multo latius; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ nigræ, graciles, extrorsum crassiores, thorace non breviores; articulus 1^{us} longus, linearis; 2^{us} cyathiformis, piceus; 3^{us} et 4^{us} minimi; 5^{us} et sequentes

mediocres, usque ad 10^{um} paullatim curtantes; clava conica, acuminata, articulo 10° multo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum brevi-obconicum: metathorax mediocris, obconicus, declivis; petiolus brevissimus: abdomen longiovatum, cupreum, nitens, læve, supra planum, subtus vix carinatum, apice acuminatum, thorace paullo longius vix latius; segmenta subæqualia: pedes fulvi, simplices, subæquales; coxæ nigræ; mesotarsi et metatarsi flavi, apice fusci: alæ limpidæ; squamulæ fulvæ; nervi fulvi; nervus humeralis ulnari multo longior, radialis ulnari vix brevior cubitali multo longior; stigma minutum. (Corp. long. lin. $1\frac{3}{4}$; alar. lin. 3.)

16. *Pteromalus Eurypon*, fem. *Ater*, abdomen cupreum, antennæ piceæ, pedes fulvi, tarsi flavi, alæ limpidæ.

Corpus convexum, atrum, parum nitens, scitissime squameum, pilis albis hirtum: caput transversum, breve, magnum, thorace multo latius; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ piceæ, subclavatæ, thorace longiores; articulus 1^{us} fulvus, longus, gracilis; 2^{us} cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes mediocres, usque ad 10^{um} paullatim curtantes; clava longiconica, acuminata, articulo 10° duplo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum brevi-obconicum: metathorax mediocris, obconicus, declivis: petiolus brevis: abdomen longiovatum, nitens, læve, cupreum, viridi et æneo-varium, supra depressum, subtus carinatum, apice acuminatum, thorace paullo longius et angustius: pedes fulvi, simplices, subæquales: coxæ nigræ; mesotarsi et metatarsi flavi, apice obscuriores: alæ limpidæ; squamulæ piceæ; nervi fulvi; nervus humeralis ulnari multo longior, radialis ulnari paullo brevior cubitali longior; stigma minutum. (Corp. long. lin. $1\frac{1}{4}$; alar. lin. 2.)

[To be continued.]

BIBLIOGRAPHICAL NOTICES.

Flore de l'Algérie, ou Catalogue des Plantes indigènes du Royaume d'Alger. Par G. MUNBY: Paris, 1847. Pp. 120. Six plates.

THIS account of the plants of French Africa is the work of a well-known English botanist who has settled in Algeria. It is a valuable contribution towards our knowledge of the flowering plants of the great Mediterranean province. Our acquaintance with the Algerine flora has hitherto been derived mainly from the admirable 'Flora Atlantica' of Desfontaines, published in 1798. During 1842 and 1843 the botany of Algeria was explored by the Members of the Scientific Commission, under the late M. Bory St. Vincent, sent by the French Government. The botanical researches were conducted chiefly by M. Durieu, and have not as yet been published. Mr. Munby has

Walker, Francis. 1847. "Characters of undescribed Chalcidites collected in North America by E. Doubleday Esq., and now in the British Museum." *The Annals and magazine of natural history; zoology, botany, and geology* 19, 392–398. <https://doi.org/10.1080/037454809494566>.

View This Item Online: <https://www.biodiversitylibrary.org/item/19437>

DOI: <https://doi.org/10.1080/037454809494566>

Permalink: <https://www.biodiversitylibrary.org/partpdf/18968>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.