



First record of *Cranchia scabra* Leach, 1817 (Cephalopoda: Cranchiidae) in the Mediterranean Sea

Antoni Quetglas¹, Francesc Alemany¹, Aina Carbonell¹ & Pilar Sánchez²

KEY WORDS: Cephalopoda, *Cranchia scabra*, first record, Balearic Islands, Mediterranean Sea.

ABSTRACT: The capture of a female specimen of *Cranchia scabra* Leach, 1817, a species not previously recorded in the Mediterranean Sea, is reported from Majorca (Balearic Islands, north-western Mediterranean).

RIASSUNTO: La pubblicazione si riferisce della cattura di una femmina di *Cranchia scabra* Leach, 1817, nei pressi di Maiorca (Isole Baleari, Mediterraneo nordoccidentale). La specie non era stata segnalata in precedenza nel Mediterraneo.

¹ IEO, Centre Oceanogràfic de Balears, Moll de Ponent s/n, Apt. 291, 07080 Palma de Mallorca, Spain.

² CSIC, Institut de Ciències del Mar, Passeig Joan de Borbó s/n, 08039 Barcelona, Spain.

INTRODUCTION

The cranchiids are one of the most important cephalopod families in number of species (NESIS, 1987; VOSS, 1980; VOSS *et al.*, 1992). Such a richness implies a high degree of morphological diversity, that is increased by the presence of sexual dimorphism and great ontogenetic changes in the majority of species. Moreover, many members of this family have been described by larval specimens only. All these factors explain the high confusion existing with the cranchiids and that the majority of the species are still undescribed (VOSS *et al.*, 1992). It is obvious that this family, presently constituted by 13 genera and more than 60 species (VOSS *et al.*, 1992), would undergo important contributions by oncoming studies.

The cranchiids are characterized by the fusion of the mantle to the head at nuchal region and to the funnel at posterolateral corners and by the presence of photophores in the eyes. They are worldwide distributed from epipelagic to upper bathypelagic waters from the Subarctic to the Antarctic (VOSS *et al.*, 1992). In the present note the occurrence of a member of this family, *Cranchia scabra* Leach 1817, is reported for the first time in the Mediterranean Sea. The species is the only Mediterranean member of the subfamily Cranchiinae.

MATERIAL, METHODS AND RESULTS

On March 25th 1998, a female specimen of *C. scabra* (Fig. 1) was caught in a haul carried out by a commercial bottom trawler fishing between 506 m and 769 m of depth in Majorcan waters (Balearic Islands, north-western Mediterranean) between the co-ordinates 39° 15' 02" N – 02° 25' 40" E and 39° 04' 52" N – 02° 25' 29" E. The specimen was identified by external morphological characters (VOSS, 1980; VOSS *et al.*, 1992): mantle with scattered cross-shape tubercles; protruding, small oval eyes; small, paddle-shape fins; funnel large, broad, extending nearly to base of arms, surface papillated; short, nontubercular, inverted V-shape cartilaginous strips extending posteriorly from

anterior apex of funnel-mantle fusions. It was not possible to determine the sexual condition of the specimen because of deteriorated internal organs.

After identification, morphometric characters (Table 1) were measured to the nearest mm and the total weight was taken to the nearest 0.1 g. The beaks were extracted and upper rostral length (URL) and lower rostral length (LRL) were measured to the nearest 0.1 mm using a dissecting microscope. The specimen was afterwards preserved in 70% ethyl alcohol and deposited in the Centre Oceanogràfic de Balears collection with the number COB1/1998.

DISCUSSION

Cranchia scabra has a circumglobal distribution, occurring in tropical and subtropical waters. The species is one of the commonest cranchiids (CLARKE, 1966). Juveniles live in the upper 400 m but larger individuals inhabit deeper waters, being cited up to 2000 m depth (VOSS *et al.*, 1992). In the Atlantic, *C. scabra* is carried in the current system into north temperate waters in the central and eastern ocean to at least 37°N (VOSS, unpublished data). Considering the fact that juveniles occur in the upper ocean layers, they may be expected to be occasionally transported into the Mediterranean Sea (VOSS, pers. comm.).

The cranchiids are poorly represented in the Mediterranean, since only three other species have been reported up to date (BELLO, 1986; MANGOLD and BOLETZKY, 1988). One of these species, *Galiteuthis armata* Joubin 1898, although it is not very frequent, has been caught throughout this sea. Another one, *Teuthowenia megalops* (Prosche 1849), was firstly recorded by SÁNCHEZ (1985) in the north-western Mediterranean and it was subsequently found in the stomach contents of a *Globicephala melaena* Traill 1809, stranded on the French coast of this sea (BELLO, 1992). More recently, BELLO and BIAGI (1999) reported the occurrence of a very large Taoniine cranchiid squid, belonging to either *Galiteuthis* or, more probably, *Megalocranchia*.



ACKNOWLEDGEMENTS

We are very grateful to the crew of the trawler Portocolom who provided the specimen. Special thanks to Nancy Voss (Rosenstiel School of Marine and Atmospheric Science, University of Miami, Florida) for the information facilitated. Thanks are also given to Sandra M. Starbuck for the English version.

REFERENCES

BELLO G., 1986. Catalogo dei Molluschi Cefalopodi viventi nel Mediterraneo. *Boll. Malacol.*, 22: 197–214.

BELLO G., 1992. Addenda al catalogo dei Molluschi Cefalopodi viventi nel Mediterraneo. *Boll. Malacol.*, 28 (1–4): 93–95.

BELLO G. and V. BIAGI, 1999. A large cranchiid squid (Cephalopoda: Teuthoidea) caught in the Mediterranean Sea. *Boll. Malacol.*, in press.

CLARKE M.R., 1966. A Review of the Systematics and Ecology of Oceanic Squids. *Adv. Mar. Biol.*, London; 4: 91–300.

MANGOLD K. and S. V. BOLETZKY, 1988. Mediterranean cephalopod fauna. In: E. R. Trueman and M. R. Clarke (eds.): *The Mollusca*, vol 12. *Paleontology and Neontology of Cephalopods*, pp. 315–330. Academic Press, San Diego.

NESIS K.N., 1987. *Cephalopods of the world*. (Translated by B.S. Levitov). T.F.H. Publications, Neptune City, N.J., 351 pp.

SÀNCHEZ P., 1985. Sobre la presencia de *Teuthowenia megalops* (Prosche, 1849) (Cephalopoda, Cranchiidae) en el Mediterráneo. *Inv. Pesq.*, 49 (2): 315–318.

VOSS N.A., 1980. A generic revision of the Cranchiidae (Cephalopoda; Oegopsida). *Bull. Mar. Sci.*, 30 (2): 365–412.

VOSS N.A. S.J. STEPHEN and ZH. DONG, 1992. Family Cranchiidae Prosche, 1849. In: Sweeney, M.J., C.F.E. Roper, K.M. Mangold, M.R. Clarke and S. v. Boletzky (eds.): “Larval” and Juvenile Cephalopods: A Manual for Their Identification. *Smithson. Contrib. Zool.*, 513: 187–210.




Fig. 1 – Female specimen of *Cranchia scabra* caught in Majorcan waters (Balearic Islands, north-western Mediterranean Sea).

	mm
Mantle length	102
Maximum mantle width	59
Arm length I	35
Arm length II	47
Arm length III	69
Arm length IV	46
Tentacle length	183
Tentacle club length	20
Base of the fin	13
Maximum fin width	17
Head width	32
Eye diameter	17
Total weight (g)	26.0
Nº photophores in the eye	14
Upper rostral length (URL)	2.0
Lower rostral length (LRL)	2.5

Table 1 – Morphometric and meristic data of the female specimen of *Cranchia scabra* caught in Majorcan waters (Balearic Islands, north-western Mediterranean Sea).



Quetglas, Antoni et al. 1999. "First record of *Cranchia scabra* Leach, 1817 (Cephalopoda: Cranchiidae) in the Mediterranean Sea." *Bollettino malacologico* 35, 1–2.

View This Item Online: <https://www.biodiversitylibrary.org/item/208895>

Permalink: <https://www.biodiversitylibrary.org/partpdf/190490>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: Società Italiana di Malacologia

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.