both during cool windy weather with sunny periods, when the temperature hardly crept above 13°C. even in full sunshine, on each occasion vibrating commenced shortly after the sun had emerged from behind a cloud and lasted for 10-15 seconds.

The first time I witnessed this behaviour (15th April, 1977) the butterfly had chosen the underside of a Corylus avellana (Hazel) leaf about 9in. from the ground and due to considerable fluttering caused by the wind I was not too certain which vibrations were caused by the wind and which by the insect itself. On the second occasion (5th May, 1980) the brimstone, a male, settled on the underside of a Verbascum thapus (Great Mullien) leaf, a much firmer anchorage, and I was able to observe the leading edge of both forewings vibrate quite rapidly.

This does seem a logical method of raising body temperature when heat absorption is not available; has anyone similar observations? — C. J. Tubb, 19 Penny's Lane, Fordingbridge,

Hants. SP6 1HQ.

An Early Record of Vanessa atalanta (L). In Cumbria. — In the afternoon of 19th April, 1980 my wife drew my attention to a Red Admiral flying in the sunshine here at Kendal Wood. The specimen was flying in the drive to my house, occasionally settling when it was seen to be in a very worn condition. There are only a few April records for this species in Cumbria. The late Dr. R. C. Lowther of Grange-over-Sands kept detailed phenological records for many years and I note from these that V. atalanta occurred in the early part of the year on only four occasions between 1920 and 1949. The earliest of these concerned a specimen at Arnside on 25th March, 1949.

These early records occasion the thought that V. atalanta may at times hibernate in this country even so far north as Cumbria. Last winter was generally a very mild one which may be significant in the present occurrence. — Dr. Neville L. Birkett, Kendal Wood, New Hutton, Cumbria LA8 0AQ.

A SECOND BRITISH SPECIMEN OF LAMPRONIA FLAVIMITRELLA HUEBNER (LEP.: INCURVARIIDAE). — On the 24th May, 1980, at about 7.30 p.m., I netted an incurvariid of unfamiliar appearance in one of the clearings in Hoads Wood near Ashford, Kent. On returning home, further examination showed it was probably referable to *L. flavimitrella*, a species only once before taken in Britain. I have since submitted the specimen to Dr. J. D. Bradley, who has kindly confirmed the determination. My example, a female of wing expanse 16 mm, has the two transverse bands on forewing uninterrupted. The first British *flavimitrella*, a male, was taken by the late Denzil Ffennell in his garden at Martyr Worthy, Hants., on the 18th May, 1974, and recorded by him in *Ent. Gaz.*, 28:180-182. — J. M. Chalmers-Hunt.

ABNORMAL ABUNDANCE OF THE COMMA: POLYGONIA C-ALBUM L. IN KENT. — At Brookland, Romney Marsh, in a lane coming from the direction of Lydd, on the 5th August, 1979, I saw more than 70 of this butterfly at bramble flowers,



Pfeiffer, Richard. 1980. "An early record of Vanessa atalanta (L.) in Cumbria." *The entomologist's record and journal of variation* 92, 195–195.

View This Item Online: https://www.biodiversitylibrary.org/item/94533

Permalink: https://www.biodiversitylibrary.org/partpdf/194477

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Amateur Entomologists' Society

License: http://creativecommons.org/licenses/by-nc-sa/3.0/

Rights: https://biodiversitylibrary.org/permissions

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.