

Lulworth Skipper: *Thymelicus acteon* Rottemburg. Very common.

POSTSCRIPT

The identifications of *Maniola jurtina*, *Hipparchia fatua* and *Cyaniris semiargus* need further checking. A correction note will be published later if this is found to be necessary.

HYLES LINEATA F. SSP. LIVORNICA ESP.: STRIPED HAWK-MOTH IN THE NEW FOREST. — A fresh specimen of this moth was taken in my M.V. trap at Brockenhurst on April 3rd. Immediately prior to this date a severe south westerly gale had been in force for two days. It seems possible therefore that in view of the early date of its capture the moth may have originated from Spain or North Africa. — A. D. A. RUSSWURM, Coridon, Ober Road, Brockenhurst, Hants. SO4 7ST.

THE PAINTED LADY: CYNTHIA CARDUI L. IN MARCH 1985. — On 14th March 1985, a worn *C. cardui* was feeding on flowers of the winter heather (*Erica carnea*) in my garden. — H. G. PHELPS, 'Green Oak', Potters Hill, Crockerton, Warminster, Wilts, BA12 8AD.

THERA BRITANNICA TURNER: SPUCE CARPET IN KENT. — In 1984, I saw this species for the first time at East Malling when a single specimen came to m.v.l. on 17th September, to be followed by two more on 15th October. Chalmers-Hunt (*The Butterflies and Moths of Kent*, Vol. 3) gives only two other records for the Weald Medway division of the county. There are several rows of *Cupressus* trees near the site of capture, planted as windbreaks or for screening, and these may have induced the moths to colonise the area. — D. A. CHAMBERS, 15 Briar Close, Larkfield, Maidstone, Kent.

EUPITHECIA — DELAYED EMERGENCE. — A number of species of Lepidoptera sometimes delay emergence from pupae until the second (or more) year after pupation (eg. genus *Cucullia*), but I have never before observed this in the genus *Eupithecia*, although I have bred many Pug species. However, on 5th June 1983 I took a worn *Eupithecia insigniata* Hb. (Pinion-spotted Pug) which had obviously laid most of its ova, but produced about half a dozen more before it died. From these, four pupae resulted, two emerging in 1984 on 8th. and 10th. March respectively. Later examination of the two remaining pupae showed one dried up but the other apparently healthy. This was kept outside for the winter 1984/5, brought in to room temperature at the beginning of March 1985 and emerged on 13th March. — Col. D. H. STERLING, "Tangmere", 2 Hampton Lane, Winchester, Hants. SO22 5LF.



Chambers, D. A. 1985. "Thera britannica Turner: spruce carpet in Kent." *The entomologist's record and journal of variation* 97, 93-93.

View This Item Online: <https://www.biodiversitylibrary.org/item/94531>

Permalink: <https://www.biodiversitylibrary.org/partpdf/195792>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Amateur Entomologists' Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.