NOTE ON THE BRITISH SPECIES OF ANOMIA.

By R. WINCKWORTH, M.A.

Read 9th November, 1921.

(PLATE I.)

THE object of the present note is to emphasize the distinctions between the four British species of Anomiidæ. There is nothing new in this; Forbes and Hanley recognize four species, though *squamula* is placed with *ephippium* instead of with *aculeata*; abroad, Dautzenburg and Jensen among others distinguish them conchologically. Ridewood in Phil. Trans., vol. 195, 1903, describes the gill of *A. aculeata* and places this species with *Dimya* in a separate suborder *Dimyacea* (l.c., p. 185): he also figures *A. ephippium* and *A. laqueata*, the latter similar to *patelliformis*.

I have recently examined numerous examples of this group from Plymouth Marine Biological Association and elsewhere. All the species are very variable in shape, thickness, and sculpture: and the young of all four are discouragingly similar in appearance, while the muscular scars are often very faint in small examples; so that the soft parts and above all the gill are a valuable aid to identification. There is no need to section the gill or even to use a microscope; a good lens will at once show sufficient to determine the species, as summarized below.

ANOMIACEA. ANOMIIDÆ.

Genus ANOMIA, L., 1758. Type, *A. ephippium*, L.

1. EPHIPPIUM, L., 1758. Upper or left valve with three distinct muscular scars (Pl. I, f. 8). Gill W-shaped in section, and at once recognizable by a dependent membranous flap of the outer ascending lamella, the fifth lamella of Lacaze Duthiers (Pl. I, f. 1).

Genus MONIA, Gray, 1849. Type, *A. zelandica*, Gray, in Dieffenbach, 1843.

I do not like referring British species to a New Zealand genus, but as I have only seen the shell of *zelandica*, and can see no conchological reason for not including our species under the same genus, I must at present leave the two following species here.

2. PATELLIFORMIS, L., 1761. The original description in Fauna Svecica is not quite satisfactory, and suggests the next species, but the figure given in N. Act. Upsala 1773 confirms the usually accepted identification, and in each case the species is mentioned as received from A. R. Martin. Hanley states that the types in the Linnean cabinet were introduced by the younger Linné.

Upper valve with two muscular scars, which are separate and distinct (Pl. I, f. 9). The typical sculpture may be almost obsolete, and, indeed, is quite absent in very young examples, or may be

exaggerated into prickles. Gill, W-shaped, without the fifth lamella of flap of *ephippium* (Pl. I, f. 2). Further, in addition to the row of ciliated discs at the lower angles of the lamellæ, which are found also in *Anomia*, there are in *Monia* other intermediate rows of ciliated discs along the faces of the filaments. In this species there is one of these in the outer and two in the inner lamella.

3. *SQUAMA*, Gmelin, 1791 = *A. striata*, Lovén, non Bolten = *A. glauca*, Monterosato. Gmelin's species is based on Chemnitz, viii, 697, where the figure is recognizable and the description unmistakable, locality Drontheim. In this species the shell adductor immediately adjoins the byssal muscle on the left side of the animal, so that the two form one continuous scar on the upper valve (Pl. I, f. 10); this part of the shell is a deep green. The gill is of the same kind as that of the last species, but the filaments are typically very much finer and the number of intermediate ciliary rows is two and three, usually, instead of one and two, but may be more numerous in part or throughout (Pl. I, f. 3). Typical examples always show on the upper valve the peculiar striated sculpture of numerous crowded rows of minute radiating scales.

One form of this species to which I give the name of var. *crassa*, as I am unable to identify it with any of the numerous *Anomiidæ* which have been described, is so different that I at first thought it a distinct species both from shell and animal characters. It is very much thicker, particularly in the coarse upper valve: the surface occasionally shows traces of the typical *squama* sculpture, but is usually much encrusted and devoid of sculpture (Pl. I, f. 13-18). It is markedly convex from its habitat on the convex side of *Pecten*. Muscular scar much as in type, but the shell adductor is comparatively more dorsal, forming an anterior prolongation of the byssal scar (Pl. I, f. 11). Gill similar, but the separate filaments larger and coarser, and the intermediate ciliary rows are highly irregular, usually three and four or more (Pl. I, f. 4), while in typical *squama* these rows are regular. The hinge process on the lower valve is enormously developed, and the border of this valve has a wide green margin within. But these and other differences do not seem to me essential, and I have no doubt as to its being a variety of *squama*, modified by its more exposed habitat.

Genus *HETERANOMIA*, nov. Type, *A. squamula*, L. = *A. aculeata*, Müller.

Anomiiform mollusca, in which the gill lamellæ have descending filaments only, each gill being therefore $\cap$ shaped in section. The alimentary and circulatory systems show it to be allied to *Anomia* and *Monia*. The lateral asymmetry is strongly marked, so that the right gill is not much more than half the length of the other.

4. *SQUAMULA*, L. 1758 = *A. aculeata*, Müller. There is no doubt that this is the Linnean species from the habitat on seaweed and

crabs, and the locality, Swedish sea, where *ephippium* is not found. There is no distinction between the smooth and aculeate forms: I have taken wholly smooth, wholly aculeate and intermediate forms on the same *Chlamys opercularis*; nor can I find any essential difference in other varieties, such as *cylindrica*, Gmelin, from the arm of a crab *Inachus*. It is at once distinguished from all other Anomiidæ by the gills, which have no ascending lamellæ (Pl. I, f. 5). Two muscular scars in the upper valve, small adjacent but distinct (Pl. I, f. 12), not showing the radial furrows of *Monia*. The byssal plug is thin brown striate, as in *Monia* and quite different from that of *Anomia*. Jensen further points out (Danish Ingolf Exp., ii, 5, p. 1) that the notch of this species is small and oval and the umbo almost or entirely marginal, while the notch in *patelliformis* is large and triangular and the umbo always a little way off the margin.

EXPLANATION OF PLATE I.

SECTIONS OF GILL. (Diagrammatic. Dots indicate ciliary junctions.)

- FIG. 1. *Anomia ephippium*. $\times 5$.
 „ 2. *Monia patelliformis*. $\times 5$.
 „ 3. *M. squama*. $\times 5$.
 „ 4. *M. squama* var. *crassa*. $\times 5$.
 „ 5. *Heteranomia squamula*. Diagram of gill. $\times 5$.
 „ 6. *H. squamula*. Right side mantle removed, showing reduced gill.
 $\times 3/2$.
 „ 7. *H. squamula*. Left side, mantle removed. $\times 3/2$.

OUTLINES OF MUSCULAR SCARS IN LEFT VALVE.

- FIG. 8. *Anomia ephippium*. Poole.
 „ 9. *Monia patelliformis*. Firth of Forth.
 „ 10. *M. squama*. Oban.
 „ 11. *M. squama* var. *crassa*. Eddystone.
 „ 12. *Heteranomia squamula*. Brighton.

SHELLS.

FIGS. 13-18.—*Monia squama* var. *crassa*. Eddystone.

Winckworth, Ronald. 1922. "NOTE ON THE BRITISH SPECIES OF ANOMIA."
Proceedings of the Malacological Society of London 15, 32–34.
<https://doi.org/10.1093/oxfordjournals.mollus.a063764>.

View This Item Online: <https://www.biodiversitylibrary.org/item/95719>

DOI: <https://doi.org/10.1093/oxfordjournals.mollus.a063764>

Permalink: <https://www.biodiversitylibrary.org/partpdf/202511>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.