# Protective Display and Sound Production in Some New World Arctiid and Ctenuchid Moths<sup>1</sup>

# A. D. BLEST

Department of Zoology and Comparative Anatomy, University College, London<sup>2</sup>

(Text-figures 1-29)

#### 1. INTRODUCTION

ESPITE a large literature on the protective devices of arthropods, there have been few attempts to study their evolution within restricted taxa in relation to particular natural communities. Such studies are to be desired because the growth of knowledge about the behavior of predators in prey/predator situations (see reviews by de Ruiter, 1956; Sheppard, 1959, 1961) has led to a belief that the selection pressures which determine the evolution of protective behavior may be formulable in fairly simple terms, and that the end-products of evolution under given circumstances may be to some extent predictable. For instance, a discussion of Müllerian and Batesian mimicry based on assumptions of this type has been given by Huheey (1961), and of the constraints which may modify the adult lifespans of some hemileucine saturniid moths by Blest (1963b).

Nevertheless, it may be doubted whether many natural situations present a sufficient degree of simplicity for such analyses, and the present more superficial survey of a large fauna may serve to illustrate some of the difficulties of such an approach.

The protective display behavior of arctiid and ctenuchid moths will be described in three main neotropical locations: at Barro Colorado Island, in the Panamá Canal Zone (lowland seasonal

search, New York Zoological Society. <sup>2</sup>Acknowledgments. The many people who have given help in Trinidad and the Panama Canal Zone have already been thanked elsewhere (Blest, 1960), and I can only repeat my indebtedness to them. For hospitality at Santa Clara I am grateful to Sr. C. Hartman. I thank the authorities of the British Museum (Natural History) for access to the arctiid and ctenuchid collections. This study was supported by the U.S. National Institutes of Health (Project RG-7109). Figures 15-29 were drawn by Mr. Roger Earle.

tropical evergreen forest: Bennett, 1963); at Volcán and Santa Clara near the Panamá-Costa Rica border (seasonal lower montane forest at 4,000-5,000 ft. above sea level); and in the Arima valley of Trinidad (transitional between lower montane rain forest and lowland seasonal forest, at an effective altitude of 500-1,800 ft. above sea-level (Beard, 1946; Beebe, 1952).

These two families were chosen because they form a group of allied, numerous and predominantly distasteful species, a large proportion of which have been accessibly monographed for the areas concerned (Forbes, 1939; Kay & Lamont, 1937; Fleming, 1957, 1959).

The nomenclature follows that used in the current arrangement of material in the British Museum (Natural History) collections, and it may be noted that Forbes (1939) lists illustrations in Seitz (1915-17) for most of the species which he records for Barro Colorado and comments on their accuracy.

Although the ctenuchid fauna is a rich one, this group has not been studied in as much detail as the arctiids; a sufficient range of species has been examined to establish the general behavioral continuity between the two families, which also show a wide range of morphologically transitional forms (Forbes, 1939). The use of light traps to secure material has caused the predominantly diurnal Ctenuchinae to be poorly represented. No attempt has been made to study the behavior of the minute and fragile lithosiine arctiids. The behavior of virtually all the dominant species on Barro Colorado has been examined between May 5-September 20, 1961, July 2-September 20, 1962, and the month of July, 1963; only those species which are rare or whose identity is uncertain are omitted. In Trinidad, only the most common species were seen

<sup>&</sup>lt;sup>1</sup>Contribution No. 1058, Department of Tropical Re-

in April, 1961, and work at Volcán and Santa Clara was confined to three weeks in July, 1961.

# 2. MATERIALS AND METHODS

The moths were captured as adults at light traps which were run throughout the night; the insects were removed from the vicinity of the light and placed in separate containers as soon as possible after arrival at the traps. The simple methods for examining protective display behavior have been described elsewhere (Blest, 1957).

The palatability tests employed White-faced Monkeys, (*Cebus albifrons*), and Geoffroy's Pinché (*Oedipomidas spix*) (Canal Zone); and domestic fowl (Santa Clara). The monkeys and marmosets were confined in groups of 4-6 in large outdoor cages.

# 3. MODES OF PROTECTIVE BEHAVIOR

The protective devices can be divided into a small number of behavior patterns found widely throughout the two groups, while occasionally showing marked taxonomic segregation, and a larger number of specialized devices with a much more scattered distribution, usually combined in some way with the basic display types. The behavioral repertoire may be summarized as follows:

I. The Emission of Ultrasonic Signals.-One species, Melese laodamia, has been shown to emit ultrasonic sounds at high intensity when it is roughly handled; the presence of the soundproducing organs on other species, and the emission by them of sounds with a faint audible component, allowed the assumption that they too are primarily generating noises in the ultrasonic range (Blest, Collett & Pye, 1963) and recent observations have confirmed that they do so. The sounds are produced by modified episternites of the metathorax. Those of Melese are modulated between 30 and 90 Kc/sec. Each burst of sound is composed of a train of cycles of modulation which sweep over the main frequency range some forty times/second; the modulation cycles themselves are formed from a series of discrete pulses whose repetition rate may rise to over 1,000/sec. The mechanics and acoustics of the process of sound production are discussed in detail elsewhere, together with their possible function. The latter is, however, quite obscure; the thresholds of sound production to tactile stimulation are very variable, and it is not yet clear what proportion of species will emit them while on the wing, or in what stimulating circumstances (Blest, Collett & Pye, 1963).

Table 1 includes a summary of the thresholds

at which sounds are produced to handling the insects. Species with a low threshold (L) will start to make sounds as soon as they are picked up and held lightly; those with medium thresholds (M) need to be squeezed, or rolled lightly between the fingers; species with high thresholds (H) must be manipulated to the point of mutilation. Those species marked "VH" perform erratically and only a minority of individuals will respond to mutilation. This classification reveals that (a) any mode of protective coloration can be linked to sound production; (b) the advanced mimetic ctenuchids have lost the ability to make sounds; the anatomical condition of the tymbal organs (i.e., their reduction to leathery and densely scaled structures) suggests that they are unlikely to use them in any circumstances; (c) there is within the arctiids, at least, a correlation between threshold of sound production, "toughness," and palatability. This relationship reveals that, predominantly, palatable arctiids have a low threshold for sound emission, while distasteful species are more likely to possess the higher thresholds.

In addition there is another unexplained correlation. While distasteful species with high thresholds for sound production are likely to have their flight periods at any time in the night, palatable species with low thresholds concentrate their flights between 3 and 5 o'clock in the morning. Table 2 summarizes the data for 35 species of arctiid and ctenuchid. The individual moths arriving at an ultra-violet light-trap were recorded hourly and removed from the light and destroyed after they had been scored. The table gives the total number of male individuals for each group of species which were recorded nightly over a three-week period on Barro Colorado Island from June 30-July 25, 1962. It is unlikely that these correlations can be understood until more is known about the emission of ultrasound in free flight, and about the normal flight times of insectivorous bats in the habitats concerned. Several species of arctiid and ctenuchid, including the genera Melese and Halysidota, can be made to reply with bursts of ultrasound to pulses from a bat simulator (Roeder, unpublished data; Pye, Pye, Flynn & Blest, in prep.), but the significance of this response is not yet understood.

II. Classical "Reflex Immobilization."-The use of the term will be restricted, for reasons which have already been discussed (Blest, 1957), to immobilization, or "freezing" which does not involve the assumption of a specialized posture. Characteristically, in both families, it does not outlast actual handling by more than a few seconds, and is given only to a constraining tactile

# TABLE 1. LIST OF SPECIES STUDIED, WITH NOTES ON THEIR DISPLAY TYPES, PALATABILITY AND SOUND PRODUCTION

Conventions: Localities:	B — Barro Colorado S — Santa Clara and Volcan T — Arima Valley, Trinidad	All of the species listed below have
Display Type:	R — Reflex immobilization II — Type II display III — Type III display W — Wing-waving display	specimens in the collections of the British Museum (Natural History). Where insufficient specimens were available for valid comparison, or
Sound Production:	L — Low threshold to tactile stimuli M — Moderate " " " H — High " " " VH — Very High " " "	there were no authenticated speci- mens available, the species is marked 4.

ARCTIIDAE PHAEGOPTERINAE (77 species)	Locality	Display type	Additional components, etc.	Sound production	Palatability and notes
Automolis lineosa Walker	В	R		L-M	Rejected by Cebus
A. lineatus Druce	в	R		L-M	22 22 22 22
A. sp. (unidentified)-I	S	R			Near to A. lineatus
A. reducta Walker	в	R		H	Rejected by Cebus
A. dilutus Felder	В	R	Falls to visual stimuli or touch	Н	37 27 79
A. pauperis Schaus	S	III			Rejected by fowls
A. rutila Stoll	BT	III	Text fig. 9	н	Rejected by Cebus
A. priscilla Rothschild	в	III		н	One female only
A. chionoplaga Dognin	В	III	Text fig. 10	н	Rejected by Cebus
A. sicilia Druce	BS	III		н	<b>33 33 39</b>
A. metallica Joicey	BT	III		н	<b>37 33 33</b>
A. taeniata Guerin	S	III	Text fig. 11	VH	Rejected by fowl
A. vitrea occidentalis Rothschild	в	III		н	Rejected by Cebus
A. vinasia Schaus	S	R		(-)	Two specimens only
A. elissa Schaus	В	III		н	Rejected by Cebus
A. elissoides Rothschild	в	III	Text figs. 7 & 8	н	»» »» »»
A. salma Druce	В	III		?	One male, Aug. 1963
A. critheis Druce	В	R(III)			Rejected by Cebus
A. tegyra Druce	BS	III	Text fig. 12	VH	33 33 33
A. obscurata Schaus	В	R(III)	Cryptic color pattern	L	Palatable
Cissura plumbea Hampson	В	R		н	Rejected by Cebus
Prumala spI	S	R			Accepted by fowl. Near <i>P.</i> suanus Druce, and matches unidentified specimens from Volcan in British Museum collection
Idalus hippia Stoll	в	R		M	
I. daga dares Druce	в	R(III)			Rejected by Cebus
Eupseudosoma aberrans Schaus	в	R(III)		L)	Rejected by Colum
E. aletis Schaus	в	R(III)		LŚ	Rejected by Ceous
Robinsonia similis Rothschild	т	III		M-H	
R. deiopea Druce	в	III	Text fig. 18	М-Н (	
R. sanea Druce	в	III	Text fig. 19	M-H (	Rejected by Cebus
R. variegata Rothschild	в	111	Rest with stripe abdomen expos	d M-H ) sed.	
R. flavomarginata Reich.	В		Convergent to a Pericopidae	small	
Baritius hampsoni Dognin	В	11 '		)	
B. haemorrhoides Schaus	В	II		{	Rejected by Cebus

BS

s

т

в

B. superba Schaus

P. affinis Rothschild

P. punctata Rothschild

Pachydota saduca Druce

II

II)

II

11)

Text fig. 2

fig. 6

No abdominal

component. Text

Rejected by Cebus

# TABLE 1. LIST OF SPECIES STUDIED, WITH NOTES ON THEIR DISPLAY TYPES, PALATABILITY AND SOUND PRODUCTION

ARCTIIDAE PHAEGOPTERINAE (77 species)	Locality	Display type	Additional components, etc.	Sound production	Palatability and notes
Melese laodamia Druce	BT	R	Text fig. 16	L \	M. laodamia rejected by
M. russata H. Edwards	S	R		L	Silver-beaked Tanagers. M.
M. asana Druce	В	R)	Forelegs some-	. L (	laodamia, M. asana and M incerta accepted by both
M. incerta Walker	BT	R	times quivered,	, as L (	primates. M. russata and
			in Bertholdia.		chiriquensis (1 8) accepted
M. chiriquensis Schaus	s	R /	Text lig. 15	_ /	by fowl on majority of occa-
Bertholdia grisescens Rothschild -	S	III		L	Accepted by fowl
B. myosticta Hampson	S	11)	Abdomen not	L	Rejected by fowl
B. fumida Schaus	S	11	raised and qui	v- L	Single specimen
B. albipunctata Schaus	В	11)	ered. Text fig.	5 L	Single specimen
Neritos cotes Druce	В	R		M)	Accorded by primeter
N. suffusa Rothschild	BT	R		M	Accepted by primates
Ochrodota pronapides Druce	BS	R	Text fig. 17	L	Accepted by Cebus & fowl
Phaemolis lepida Schaus	В	R		L)	
Neaxia bella Schaus	В	R		L	
Eriostepta beata Dognin	В	R		L }	Accepted by Cebus
Araeomolis persimilis Rothschild	В	R		L	
Amaxia pyga Schaus	В	R		н/	
Amaxia chaon Druce	В	R		L	et.addde.Dadae
Ammalo helops Cramer	В	R		)	Delected has Calue
Elysius conspersa Walker	В	R		}	Rejected by Ceous
Amastus rumina Druce	S	R			Territoria de la construcción de la
Opharus astur Cramer	BS	R			
Psychophasma erosa Herrich-Schaffer	B	R			
Castrica phalaenoides Drury	В	R		M	
Tricypha imperialis Heylaerts	B	R			Rejected by Cebus
Pelochyta misera Schaus	В	R			
Halysidota interlineata Walker	В	R	Forelegs raised and quivered a in <i>Bertholdia</i> . Abdomen not	1 IS	Rejected by <i>Cebus</i>
H. underwoodi Rothschild	S	R-II	curled. The dis	play	
H. atra Druce	S	R-11 /	of <i>H. interlinea</i> infrequent and transient	uta	
H. cyclozonata Hampshire	В	R		L	
H. iridescens Schaus	В	R		L)	
H. marionensis Schaus	В	R		{	Accepted by Cebus
H. angulata Rothschild	В	R		)	
Glaucostola guttipalpis Walker	В	R		M	
G. reimona Schaus -I	В	R		M	
G. flavida Schaus	В	R		M	Rejected by Cebus
G. metaxantha Schaus -I	В	R		?	One male, July, 1963
Agoraea semivitrea Rothschild	В	II			The statement of the statement of the
Metaxanthia aureiventris Rothschild	В	R			
M. atribasis Rothschild 4	В	R			Agrees with Forbes's des- cription of specimens from Barro Colorado, but not with authenticated speci- mens. Possibly a new species.
BELEMINIINAE (3 species)					men a coolery when opened
Belemnia pavonia Forbes	B	RW		{	Rejected by Cebus
Belemniastis troetschi Druce	в	RW	thorax	,	an an ann an Annaichtean an Annaichtean An annaistean an Annaichtean Annaichtean
Diospage chrysobasis Hampson	В	RW			and the second

# TABLE 1. LIST OF SPECIES STUDIED, WITH NOTES ON THEIR DISPLAY TYPES, PALATABILITY AND SOUND PRODUCTION (CONTINUED)

			and a second		
ARCTIIDAE ARCTIINAE (5 species)	Locality	Display type	Additional components,	Sound production	Palatability and notes
			eic.		L
Ecpantheria laeta Walker	в	II		{	Rejected by Cebus
E. muzina Oberthür	Т	II	Text fig. 3	)	
E. icasia trinitatis Rothschild	В	II		}	Rejected by Cebus
E. orsa Cramer	в	II		)	
Virbia rosenbergi Rothschild	в	II,W	Some slow elev- tion and depres	a }-	Accepted by Cebus
CTENUCHIDAE			sion of the wing	<u>s</u>	
EUCHROMIINAE (35 species)					
Histaea cepheus Cramer	т	RW	Copious reflex bleeding	VH	Sounds made by single male only
Homoeqcera stictosoma Druce	В	RW	Male with ventr valve secretion	ral	Rejected by Cebus
Autochloris jansonis Butler	В	RW	Frothing of per copid type	i	Rejected by Cebus
Sarosa innotata Draudt	В	RW	"Bobbing" of abdomen		Rejected by Cebus
Gymnelia salvini Butler	В	RW			23 33 35
Phoenicoprocta paucipuncta Dyar	в	RW			<b>55 53 37</b>
P. vacillans Walker	Т	RW			
P. insperata Walker	В	RW			
Mesothen pyrrha Schaus	В	RW			
Cosmosoma teuthras Walker	B	RW			Rejected by Cebus
C. batesi Butler	B	RW			
C. hercyna hercynacula Dyar	B	RW			
C. metallescens Menetries	B	RW			
C remotum Walker	B	RW			
C. advena Druce	s	RW			
C. klagesi Bothschild	т	PW/			
C. metathoracica Kava	T	DIA			
C. hogotensis Feld	R	DW7			One male culu
Saurita clusia Druce	שת	DW		· \	One male only
S afflicta Walker	DI	DUI			Rejected by Cebus
S. applicate Walker	DI	RW		{	
Isanthrana archroniformia	B	RW		)	
Staudinger	5	11	flight. Text fig. 4	n	One female only
Psoloptera thoracica Walker	В	RW			Rejected by Cebus
Rhynchopyga cryptoleuca Walker Dycladia vitrina Rothschild	B	RW R			39 99 99
Phoenarete diana Druce	S	II-W	Display posture assymetrical		Rejected by chickens
Macrocneme adonis Druce	В	RW	Male genitalia		All <i>Macrocneme</i> tested with <i>Cebus</i> were rejected
M. thyra Moschler	BT	RW	fig. 1a		
M. cyllarus Druce	В	RW			
M. hampsoni Schrottky	В	RW			
M. auripes Walker	В	RW			
M. chrysotarsia Hampson	В	RW			
Calonotos tiburtus Cramer	В	RW )	Male genitalia	)	Rejected by Cebus
C. metallicus Druce	В	RW	extruded. Text		27 23 27
C. craneae Fleming	т	TW)	fig. 1a		
	and the second second		-		

I	49	):	1	0

# TABLE 1. LIST OF SPECIES STUDIED, WITH NOTES ON THEIR DISPLAY TYPES, PALATABILITY AND SOUND PRODUCTION (CONTINUED)

	1	1			
CTENUCHIDAE CTENUCHINAE (63 species)	Locality	Display type	Additional components,	Sound production	Palatability -and notes
Marecidia achrysa Forbes J	B	RW			Rejected by <i>Cebus</i>
Fumenogaster moza Druce	B	RW			»» »» »»
Chrysostola fulvispher Druce	B	RW			
Ptervgopterus leucomela Walker	B	RW			Mimics general facies of
	5				sphecid wasp. Antennae pal- pate ground when walking, and movement jerky. Reject- ed by <i>Cebus</i>
Androcharta meones Stollinger	В	RW			-
Amycles anthracina Walker	В	RW			
Atyphopsis modesta Butler	В	R			Certain of these very similar
Episcepsis venata Butler	В	R		)	species producesounds when
E. lamia Butler	В	R			they are handled roughly
E. lenaeus Cramer	В	R		}	(M), but the distribution of
E. hypoleuca Hampson	В	R		\-	been studied.
E. frances Dyar	В	R	8	]	All are superficially much
E. gnomoides Schaus	В	R		{	alike, and all those tested
E. capysca Schaus	В	R		)	were rejected by Cebus. Sin-
E. pseudothetis Fleming	Т	R			gle specimens from Barro
E. redunda Schaus	т	R		- }	signed to <i>E. vinasia</i> Schaus, <i>E. klagesi</i> Rothschild and <i>E. demonis</i> Druce have also been taken.
Eucereon aoris Moschler	В	R		L	All species of Eucereontested
E. aeolum Hampson	В	R		L	with Cebus or with domestic
E. obscurum Moschler	В	R	Specialized res posture. Text fig. 14	ting L	fowl were rejected. They in- cluded E. obscurum, E. late- fascia, E. pseudarchias, E.
E. leria Druce	S	R)	Crimson cervic	al M	cora, E. aeolum, E. tarona.
E. tarona Hampson	В	R	hairtufts. Text figs. 20, 21	Μ	E. leria, E. atrigutta, and E. flavicaput
E. formosa Dognin	S	R	Text fig. 27	M	
E. atrigutta Druce	B,S	R		м	
E. venosa Schaus	В	R		M	
E. rosa Walker	В	R		н	
E. maia Druce	B,T	R	Text fig. 29	н	
E. zamorae Dognin	В	R			
E. flavicaput Hampson	B,S	R	Text fig. 24		
E. scyton Cramer	В	R			
E. decora Schaus	S	R	Dorsal abdom hair tufts	inal	
E. xanthura Schaus	В	R			
E. intranotata	В	R			
E. myrtusa Druce	S	R			
E. latefascia Walker	В	R	Text fig. 22		
E. pseudarchias Hampson	В	R			
E. striata Druce	S	R			
E. rogersi Druce	B,S	R	Text fig. 25		
E. tigrata Herrich-Schaffer	S	R	Text fig. 26		
E. costulata Herrich-Schaffer	S	R	Text fig. 28		
E. zephyrus Schaus	В	R			
Heliura balia Hampson	В	R	Text fig. 23	-	
H. rhodophila Walker -	В	R			

CTENUCHIDAE CTENUCHINAE (63 species)	Locality	Display type	Additional components, etc.	Sound production	Palatability and notes
H. banoca Forbes -I	в	R		)	
Delphyre rubricincta Hampson	В	R	Foul odor relea when handled	used	All Heliura and Delphyre
D. atava Druce	В	R	Resembles Episcepsis	ML	rejected by Cebus
D. aurorina Forbes -I	В	R		)	
Correbia lycoides Walker	ВТ	R	Very heavily sclerotized	— j	All are lycid beetle mimics.
C. affinis Druce	В	R		}	and all are rejected by Cebus
Correbidia terminalis Walker	ВТ	R	Weakly sclerotized	— )	
Hyaleucera luctuosa Moschler	в	R-II			
H. fusiformis Walker	в	R-II			
H. gigantea Schaus	BS	R	Resting posture and behavior li <i>Eucereon obscu</i>	e ike urum	
Ptychotrichos zeus Schaus	В	II	Abdomen curle wings not eleva	d, ated	
Napata walkeri Druce	вт	R		L	
N. leucotelus Butler	в	R	All are tough a	nd L	All are rejected by Cebus
Aclytia gynamorpha Hampson	В	R	bear bold patte	rns L	
A. punctata Butler	В	R	when they are a	at L	
A. albistriga Schaus -I	В	R/	rest	L)	
Leucopleura ciarana Schaus -I	В	RW	Genitalia extruc by both sexes	ded	Rejected by Cebus
Xanthopleura troetschi Druce	В	RW		·	

 TABLE 1. LIST OF SPECIES STUDIED, WITH NOTES ON THEIR DISPLAY TYPES, PALATABILITY

 AND SOUND PRODUCTION (CONTINUED)

stimulus. It may be associated with other, more specialized, postures, for example, the extrusion of colored genital processes, of colored tubes from the cervical region, or of abdominal hair tufts (see below). This is the typical protective response of the ctenuchids, and two examples are given in Text-fig. 1.

III. Wing-waving Displays.—The wings, which are protracted from rest, are slowly and gently elevated and depressed. These displays are similar to the saturniid rhythmic displays (Types III & IV) (Blest, 1957). They are found in the North American Diacrisia virgo, in virtually all the mimetic Ctenuchidae as a concomitant of walking, and in Virbia spp, following Type II display. Transitional genera between the Arctiidae and Ctenuchidae (e.g., Belemnia, Belemniastis and Diospage, but not Metaxanthia) also wave their wings as they walk.

IV. Type II Sustained Static Displays.—These have been described for the hemileucine Saturniidae (Blest, 1957). The moth falls from its substrate, curls the abdomen so that the dorsal surface is convex, and places its legs in a characteristic posture; the wings are simultaneously elevated (Text-figs. 2-6). Static displays of this kind may be maintained for periods of a few seconds up to several minutes by saturniids such as *Dirphia* (*Periphoba*) *hircia* (Blest, 1960). Among the present families these displays show more variety: the abdomen is not always curled, and the angle of wing elevation and leg-extension varies widely between species.

V. Type III Sustained Static Displays.—Displays of this type are the converse of the Type II displays; the abdomen is elevated instead of depressed, so that it projects above the folded wings at right angles to the substrate, and the wings are slightly depressed. The abdomen itself is usually bright colored and banded (Text-figs. 7-12).

#### VI. Special Devices:

(i) Extrusion of the Male Genitalia.—The male genital processes may be ornamented with brightly-colored tufts of hair (e.g., Baritius haemorrhoides), or they may be so feebly sclerotized that they assume the color of the blood (e.g., Macrocneme thyra intacta, Calonotos craneae). In species so decorated the genitalia are extruded during display (Text-fig. 1).

(ii) The Display of Abdominal or Cervical

[49:1	0
-------	---

TABLE 2. FREQUENCY WITH WHICH MALE MOTHS WERE CAPTURED AT HOURLY INTERVALS THROUGHOUTTHE NIGHT OVER A THREE-WEEKS PERIOD (JUNE 30-JULY 25, 1962), ON BARRO COLORADO ISLAND

I. ARCTIIDAE Flight Times Species	19:00- 20:00	20:00- 21:00	21:00- 22:00	22:00- 23:00	23:00- 24:00	24:00- 01:00	01:00- 02:00	02:00- 03:00	03:00- 04:00	04:00- 05:00	Totals	Sound Threshold
Melese laodamia	3	1	1	0	1	3	0	1	118	19	147	L
M. incerta	1	2	2	3	1	4	8	33	60	20	134	L
M. asana	1	0	1	0	3	0	0	0	2	2	9	L
All Melese	5	3	4	3	5	7	8	34	180	41		
Automolis obscurata	1	3	0	0	1	0	0	1	4	13	22	L
Halysidota iridescens	0	1	0	0	0	0	1	5	15	7	29	L
Automolis dilutus	1	1	17	20	4	2	0	0	1	1	47	Н
A. critheis	0	3	7	5	0	1	0	0	0	0	16	Silent
A. rutila/A. chionoplaga	2	3	0	0	0	0	0	1	0	0	6	H
A. elissa/A. elissoides	0	0	0	0	0	0	3	0	0	0	3	Н
A. sicilia/A. metallica	1	1	1	0	3	2	2	0	0	0	10	H
A. lineosa	0	0	1	4	2	0	1	0	1	0	9	L-M
All Automolis	4	8	26	29	9	5	6	1	2	1		
Robinsonia variegata/ R. sanea/R. deiopea	1	0	1	2	2	0	2	0	0	1	9	M-H
Amaxia pyga	1	0	0	2	2	10	5	3	2	0	24	н
Elysius conspersa	0	0	0	3	10	5	2	1	0	0	21	Silent
II. CTENUCHIDAE												
Eucereon pseudarchas	0	3	3	6	4	3	3	0	4	0	26	Silent
Aclytia gynamorpha	3	9	6	2	0	0	2	0	0	2	24	L
Episcepsis spp.	3	13	6	6	3	2	3	0	2	2	40	M-silen
Correbidia terminalis	8	24	15	8	5	1	0	1	0	0	62	Silent
Correbia lycoides	7	11	5	0	1	2	0	0	0	0	26	Silent
Delphyre rubricincta	11	8	2	5	0	0	0	0	1	0	27	Silent
Calonotos tiburtus	6	31	3	0	1	0	0	0	0	0	41	Silent
C. metallicus	2	8	1	2	0	0	0	0	0	0	13	Silent
Macrocneme spp.	23	22	2	2	0	1	1	0	0	0	51	Silent
Homoeocera stictosoma	0	0	1	7	2	4	3	1	0	0	18	Silent

Hair Tufts.—Some ctenuchids (e.g., Eucereon decora) have a series of lateral hair tufts on the anterior part of the abdomen, which is concealed

at rest, and only erected during display. Display is of Type II, and the erected hair tufts are therefore readily visible to a predator. Some



TEXT-FIG. 1. a. Calonotos craneae, reflex immobilization, female. b. Macrocneme thyra intacta, male. Reflex immobilization, with slight extrusion of the genitalia.



*Eucereon* spp. and related genera have red or orange cervical hair tufts which are only exposed when the moths are handled.

(iii) Release of Materials from the Ventral Valve.—The males of certain ctenuchids possess a specialized ventral abdominal structure, the ventral valve, which in some species encloses

a yellow or whitish secreted product. In only one of the species observed in this study has any directly protective function been noted for this material, though it may assist in rendering the moths unpalatable. In *Homeocera stictosoma*, light restraint of the insect elicits reflex immobilization, and the instant release of the accumu-



TEXT-FIGS. 7-12. Type III displays. 7, Automolis elissoides at rest; 8, A. elissoides displaying; 9, A. rutila; 10, A. chionoplaga; 11, A. taeniata; 12, A. tegyra.

lated secretion, in the form of a light white flocculent material composed of fine friable fibres which rapidly envelop the animal, and which have a faint but definite phenolic odor. They are dispersed when the insect flies away. The material is not replenished after it has been used, and the ventral valves of animals which have released it remain empty until they die.

(iv) Reflex Bleeding.—In a great many species reflex immobilization is accompanied by reflex bleeding from the joints and prothorax. It is particularly copious in Histaea cepheus and Amastus rumina.

(v) Specialized Secretions. - Several European arctiids are known to possess specialized defensive secretions which are emitted from glands in the prothorax (Bisset, Frazer, Rothschild & Schachter, 1960). In one case (Arctia caja (L.)), the secretions have been shown to contain an unusual higher alkyl choline ester, probably  $\beta$ - $\beta$ -dimethylacrylylcholine, or a near derivative (Bisset *et al.*, *loc. cit.*). It is likely that some of the species under consideration may possess such defensive mechanisms, but they were not specifically searched for in this study.

(vi) Odors.-Most arctiids and ctenuchids which are definitely distasteful have a noticeably unpleasant, musty odor to the human observer, and they are often rejected by Cebus in test situations after being smelled but not tasted. These odors are absent or less noticeable in Blest: Display and Sound Production in Arctiid and Ctenuchid Moths



TEXT-FIG. 13. a. & b. Cervical region of unidentified ctenuchine near to *Eucereon*, to show extrusible lemon-yellow processes.

species which have proved palatable to both *Cebus* and *Oedipomidas* (e.g., *Melese laodamia* and *M. asana*). Odor is not, however, a necessary concomitant of unpalatibility, for *Bertholdia myosticta*, which were consistently rejected by chickens, do not possess a strong smell.

Delphyre rubricincta, which performs a Type II display exhibiting a black and crimson banded abdomen, simultaneously releases a foul odor of short persistence. No visible droplets of fluid have been seen to accompany it, and its origin is consequently for the present unknown. This odor somewhat resembles that of the alarm substance of the large neotropical ants of the genus *Paraponera*. Specimens of *Gymnelia salvini* have a faint but similar odor.

(vii) Frothing. – Belemniastis troetschi and Autochloris jansonis froth vigorously from the thorax when they are handled. Many pericopids produce froth and so do African members of the genus Rhodogastria (Arctiidae) (Carpenter, 1938).

(viii) Other Specialized Components.—Although the general forms of the displays are remarkably stereotyped, the minor components are often modified. For example, species of *Baritius* tend to retain their hold on the substrate while performing a Type II display, and *B.* haemorrhoides and *B. superba* back away from the disturbing stimulus, following a semicircular path. *Bertholdia* spp. place the first pair of legs in a specialized posture and quiver them rapidly (Fig. 5); similar specialized leg postures are also found in Melese asana and M. incerta, and in Halysidota underwoodi.

The ctenuchid Sarosa innotata accompanies an otherwise normal reflex immobilization with vertical bobbing movements of the abdomen, which is yellow with a black tip; the significance of this behavior pattern is obscure, and similarly marked species of Cosmosoma (e.g. C. hercyna hercynacula) do not possess it.

A single specimen of an unidentified ctenuchine near *Eucereon* from Santa Clara extruded a pair of flabby lemon-yellow processes from the cervical region when touched (Text-fig. 13). No perceptible odor was associated with them. *E. leria* and *E. tarona* bent their heads forward to expose crimson hair tufts in the same region.

Perhaps the majority of ctenuchids show some degree of mimicry, hymenopterous and dipterous insects being the most usual models. A general account is given in the various sections of Seitz's "Macrolepidoptera of the World" dealing with the family. Although the flight patterns of these mimetic species often closely resemble those of the various models, their resting behavior is relatively little specialized; the more extreme wasp-mimics, however, (e.g., Pterygopterus leucomela) walk with the jerky movements of their models, and may even palpate the surface of the ground with their antennae in a wasp-



TEXT-FIG. 14. The resting posture of *Eucereon ob*scurum.

1964]



like manner. An account of some ctenuchid flight adaptations is given by Beebe & Kenedy (1957).

VII. Cryptic Coloration.—Although several genera have an inconspicuous appearance, perhaps equivalent to "generalized cryptic coloration" in the saturniids (Blest, 1957) (e.g., Melese, Bertholdia, Elysius, Ammalo) (Text.figs. 15 and 16), very few appear to have specialized procryptic coloration. The exceptions, which have been shown to be palatable (Ochrodota pronapides (Text-fig. 17), Amaxia pyga, Melese chiriquensis), are also striking for their fragility; this correlation between procrypsis and fragility can probably be explained in terms of natural selection, and requires more extensive study (Blest, 1963a).

#### 4. NOTES ON INDIVIDUAL GENERA

Although a list of the species and their behavior is given in Table I, certain genera deserve special discussion.

# I. Automolis.

Forbes (1939) considers that this genus is not homogeneous, and it has been split into a number of species-groups. Its protective behavior is of two kinds: reflex immobilization, and Type III display. Either may be accompanied by sound production, the threshold for which is lowest in those species showing reflex immobilization only, and highest in *A. taeniata*, only one of some two dozen specimens producing an auditory signal. Since *A. taeniata* maintains its display particularly vigorously, it would appear that this relation may be merely a matter of the occlusion of sound production by the competing display reflex.

**TEXT-FIG. 17** 

Despite the uniformity of the Type III displays, the visual patterns which they exhibit vary widely (Text-figs. 7-12). *A. dilutus* apparently mimics an implausibly large lampyrid beetle and does not display. Tactile or visual stimuli cause it to fall to the ground, and the related *A. reducta* behaves in the same way.

All members of the genus tested showed some degree of unpalatability; some species (e.g., A. sicilia, A. vitrea) were occasionally accepted by Cebus, although the majority of individuals were rejected. Most species were consistently rejected (A. metallica, A. diluta, A. critheis, A. elissa, A. elissoides), and survived investigation and rejection by Cebus on the majority of occasions. A. taeniata proved highly distasteful to domestic fowl, and individual birds rapidly learned to ignore them.

#### II. Cissura.

This genus was represented by one species only, *C. plumbeus* on Barro Colorado. It is notable for the coupling of reflex immobilization to sound production at very high threshold, the signals consisting of a brief train of short squeaks. The coloration is aposematic, and comprises longitudinal carmine stripes on a dark slategray ground. This was the only species which all the *Cebus* in the colony learned to ignore on sight, and of which every individual survived investigation during the early encounters.

#### III. Robinsonia.

All members of this genus have Type III displays, and all produce sounds. Two of the species (R. deiopea and R. sanea), (Text-figs. 18 & 19), have uniformly pale yellow abdomens and rest in the normal arctiid position, with the abdomen concealed by the folded wings. R. flavomarginata can be found resting on the undersides of the leaves of low second-growth vegetation in the forest; this and the similar R. variegata show convergence towards the smaller pericopids which may be found in the same habitat, particularly in the concentration of dark pigmentation along the veins of the forewings; they share the pericopid habit of resting with the brightly colored abdomen exposed.

# IV. Eucereon, Delphyre, Heliura, Hyaleucera gigantea.

All the species of this group which have been tested have proved unpalatable. The 27 species from Barro Colorado and Santa Clara assort themselves into the following groups on the basis of their wing patterns; 24 belong to *Eucer*eon, to which three other genera are convergent.

This grouping omits Delphyre atava, which resembles an Episcepsis, and D. aurorina, Heliura banoca and H. rhodophila, which neither resemble each other nor any of the main groups of Eucereon. Group II is more diversified than the others, and no account is taken of the numerous species of Eucereon found in Trinidad.

Each of these types of pattern is distinct, and the members of each group are closely similar to each other. The species in most cases are adequately illustrated by Seitz. The distribution of these species between Santa Clara and Barro Colorado suggests the following conclusions: (i) In most cases each group can be treated as two small subgroups of similar sympatric species, with little overlap. (ii) Each such subgroup functions as a potential Müllerian assemblage. (iii) There is a rather striking pattern of replacement within each group. Thus, in Group I,



TEXT-FIGS. 18 & 19. Robinsonia. 18, R. deiopea; 19, R. sanea.

Group I		Group II		Group	III	Group IV		
leria tarona (Text-figs. 20,	(S) (B) 21).	aeolum aoris pseudarchias latefascia myrtusa rosa Heliura balia Delphyre rubricin (Text-figs. 22, 23	(B) (B) (B) (B) (S) (B) (B) neta (B) ).	zamorae flavicaput scyton decora xanthura rogersi (Text-figs. 2	(B) (BS) (B) (S) (B) (S) (24, 25).	atrigutta formosa tigrata (Text-figs.	(BS) (S) (S) 26, 27).	
Group V		Group V	[	Group	VII	Group	VIII	
striata costulata phaeoprocta intranotata (Text-fig. 28).	(S) (S) (B) (B)	obscurum Hyaleucera gigan (Text-fig. 14).	(BS) stea (BS)	zephyrum	(B)	maia (Text-fig.	(B) 29)	

B = Barro Colorado; S = Santa Clara.

*tarona* on Barro Colorado is replaced in Santa Clara by the very closely similar *leria*; in Group V, *phaeoprocta* and *intranotata* by *costulata* and *striata*. The close similarity between the patterns and coloring found in the two habitats in these distasteful species is in striking contrast to those of the predominantly procryptic families such as the Lasiocampidae and Noctuidae, where the wealth of lichens, mosses and epiphytes at the higher altitude has provoked the evolution of patterns and coloration distinct from the forms prevalent on Barro Colorado (Blest, 1963a).



TEXT-FIG. 21

[49:10



TEXT-FIGS. 22 & 23. Eucereon, Group II. 22, E. latefascia; 23, Heliura balia.





It is possible that the existence of similar groups of "equivalent" species in the two areas is merely a function of the niches available to the larvae, and of the stability of the color patterns within the groups. However, there still remains the possibility that the distribution of aposematic patterns within populations of sympatric species is determined by some feature of predator behavior which has not yet been appreciated. For example, Van der Vecht (1961) has provided some support for regarding the action of local predator populations as selectively critical, in a study of the Indo-Australian wasps of the genera Eumenes, Pareumenes and Pseudeumenes. In this case, Eumenes flavopictus, extending from southern India to New Guinea, is mimicked by species of Pseudeumenes and Pareumenes. All three genera show parallel variation throughout their range, and form in each locality where they simultaneously occur a distinctively patterned Müllerian assemblage. The size of the departure from the pattern of the mainland Eumenes population relates directly to the degree of isolation of the divergent populations. It seems necessary to assume that in each isolate specific local patterns of predation have stabilized the dominant species (Eumenes flavopictus) and have caused convergence with the other two

genera. This interpretation holds whether or not genetic drift is invoked to explain the initial divergence between the island and the mainland populations.

# V. Macrocneme, Calonotos.

In both these ctenuchid genera, the male genital apparatus has been elaborated in certain species to provide a warning signal. Macrocneme thyra intacta and M. adonis belong to a division of the genus in which the male ventral valve is large and bifid. The whole genital apparatus, which is bright yellow in life, is extruded during reflex immobilization. In the remaining species it remains retracted, and the valves are small and usually asymmetrical (Forbes, 1939). A similar trend is found in Calonotos. Males of C. craneae possess supernumerary genital processes and instantly extrude the genital apparatus when restrained. It is feebly sclerotized, bright yellow, and has a fleshy appearance in life. C. tiburtus and C. metallica lack the extra process; the apparatus is normally sclerotized, brown, and is not spontaneously extruded during handling.

5. SYSTEMATIC CONSIDERATIONS Little can be concluded from the distribution

[49:10

Blest: Display and Sound Production in Arctiid and Ctenuchid Moths



TEXT-FIG. 28. Eucereon, Group V. E. costulata.

of display behavior within the two families. Within the Arctiidae, it can be said that no arctiine possesses a Type III display, which is only found in certain Phaegopterinae (Automolis, Robinsonia spp.). Type II displays are found in both subfamilies, and in certain ctenuchids. In general, the imprecise morphological division between the ctenuchids and arctiids, most evident in such transitional genera as Belemnia, Belemniastis and Diospyge, is paralleled by their behavior. Such ctenuchines as Delphyre and Eucereon spp. differ little in their behavior (reflex immobilization or Type II static displays) from arctiids such as Agoraea and Ecpantheria spp. The abandonment of the characteristic arctiid displays and resting and reflex immobilization postures follows the adoption by the ctenuchids of varying degrees of mimicry of nonlepidopterous insects, and reduced wing-scaling.

# 6. RELATIONS BETWEEN ENVIRONMENTAL TEMPERATURE AND DISPLAY BEHAVIOR

It is already clear that the persistence of static displays shows some relationship to environmental temperature. The ambient diurnal temperature on Barro Colorado fluctuates around 27° C at a height of 55 ft. in the forest shade (Allee, 1926), and may reach 34° C during the day. This latter temperature is nearing that of working lepidopterous muscle as measured by Krogh & Zeuthen (1941) for temperate nymphalids and sphingids and by Dorsett (1962) for tropical sphingids. It follows, and observation confirms, that on Barro Colorado only small periods of shivering need precede fully competent flight. Indeed, large and heavy sphingids and saturniids often make successful escape flights without shivering at all. It might be argued, then, that for a species only moderately unpalatable and not physically resilient, selection might well favor escape flights over a protracted display. In



TEXT-FIG. 29. Eucereon, Group VIII. E. maia.

fact, this appears to be the case. The static displays of the Barro Colorado arctiids and ctenuchids are transient, and are only held for periods of seconds; so much so that they have proved difficult to photograph.

At Santa Clara, however, at an altitude of 4,000 ft., the temperatures are lower. Apparently in correspondence, displays and reflex immobilization are long-persisting. Automolis taeniata and A. tegyra maintained Type III displays invariably for periods of seconds, and sometimes for minutes. On Barro Colorado, A. tegyra, and A. metallica which belongs to the same species-group as A. taeniata, both gave only transient displays. At Volcán itself, Halysidota underwoodi gave a lengthy display which added specialized leg postures to reflex immobilization. At Barro Colorado, H. interlineata, closely related to it, assumed reflex immobilization without leg components for no more than brief periods.

The evidence from arctiid and ctenuchid displays alone does not allow a certain distinction between the effects of natural selection for a display length adjusted to the diurnal ambient temperature, and the mere effect of immediate temperature upon the physiological and behavioral responses of the individual moths. Controlled experiments were not possible, but chilling Automolis on Barro Colorado did not seem to lengthen their displays by a substantial amount. The best evidence favoring the first alternative comes from the hemileucine saturniids; there, the palatable forms in general bear evespot patterns, and either display them statically, or perform rhythmic displays. With the exception of Automeris janus, which occurs at a wide range of altitudes, and rare individuals of A. tridens and A. junonia, none of the Barro Colorado Automeris with simple static displays (8 species) perform rhythmic components,

1964]

177

[49:10

while of the Santa Clara Automeris which were seen as adults, two (A. nyctimane and A. subpicta) had low-threshold rhythmic displays, while one (A. montezuma) had rhythmic leg components. A. nyctimane, received in England as pupae from Bogotá, Colombia (8,000 ft.), also had low-threshold rhythmic displays. All were well sustained, in contrast to the transient displays of the Barro Colorado Automeris.

In addition, two species, A. junonia and A. tridens, perform brief static displays on Barro Colorado, and chilling fails to elicit a rhythmic performance. In the Arima valley of Trinidad (diurnal temperature fluctuating around 25° C) a proportion of individuals of both species perform rhythmic displays, as does the related A. liberia. The species whose displays are characteristically of low threshold (those of the A. nyctimane group) are typically either montane, or found at the extremes of the range of the genus, in southern Brazil and Argentina, and in Mexico. A detailed study of their distribution in terms of their environmental temperatures and display performance would be of great interest.

# 7. THE GENERAL ADAPTIVENESS OF DISPLAY AND ESCAPE BEHAVIOR

Probably the most important predators of these small moths are insectivorous birds, notably the numerous species of tropical flycatcher (Tyrannidae) and bats, though nothing is known about the patterns of predation prevailing in the forest canopy. It is likely that birds discriminate more carefully in their choice of prey than the captive monkeys used in these observations, if only for the reason that monkeys caged in groups are in a competitive situation in which prey is eaten to assert dominance, or even to please the experimenter, as well as to satisfy hunger. For example, Silver-beaked Tanagers rejected Melese laodamia<sup>3</sup> while the two primates accepted it, and domestic fowl took the equivalent species (M. russata) at Santa Clara. For these reasons it is remarkable that so many of the species achieved almost complete protection against the caged Cebus. The small number of experimental predators and the erratic appearance of any one species of moth precluded formal palatability tests of the kind so elegantly used by Brower (1958, 1960); tests with Cebus were confined to some dozen presentations to the colony in each day, in which no more than eight arctiids were presented, the remaining species being palatable sphingids, noctuids and geometrids. The moths were pushed through the wire, and in most cases the monkeys took them directly in their hands.

From the observations which resulted, it appears that there are two rather distinct ways in which an unpalatable species may survive investigation by an inexperienced monkey. Cebus pick up small prey in their hands, and smell it before biting it. Arctiids and ctenuchids are usually rejected at this stage. An actively struggling prey is usually picked to pieces, however, even if it is not subsequently eaten; the rejection tends to be followed by manual investigation which is sufficiently vigorous to elicit struggling from those saturniids which have static displays. Arctiids, however, remain passive throughout this phase, and are ultimately dropped. In nature they would fall some 30-90 ft. from the forest canopy to the ground. Even in cage conditions they succeed in creeping away without exciting further attention. Correlated with this behavior is an outstandingly tough and leathery consistency. Such moths (e.g., Automolis metallica and related species) will suffer quite violent investigation without lethal injury.

The alternative method of escape exploits the fact that a monkey when holding a passive prey eventually opens its hands to look at it; species such as *Eucereon* and the euchromiine mimics of Hymenoptera and Diptera terminate their reflex immobilization with instant flight as soon as tactile restraint is removed. *Eucereon aoris* and small *Macrocneme* repeatedly eluded *Cebus* in this way.

It is likely that these two escape mechanisms may succeed against most small primates, and perhaps avian predators as well.

#### 8. DISCUSSION

The main problem raised by these observations may be simply stated: the arctiids and ctenuchids consist of predominantly distasteful insects. So unpalatable are they that most species are rejected even by caged colonies of Cebus. If it is assumed that predators in nature must learn to avoid distasteful species, and that those which search visually learn best bold aposematic patterns of a conventional kind (Cott, 1949), then it follows that selection should tend to promote Müllerian convergence in any one taxonomic group of distasteful animals, given certain simplifying conditions in the ecological situation (Blest, 1963a). These conditions consist, in the main, of the mere absence of various sources of constraint. Convergence, for example, might be prevented were the animals to use visual signals for sexual recognition, in which case selection for character divergence as an isolating

<sup>&</sup>lt;sup>8</sup>Tests with Silver-beaked Tanagers were conducted by Mr. T. S. Collett and Mr. C. T. Collins in Trinidad.

mechanism (Brown & Wilson, 1956) should tend to sustain the opposite change in any one ecological assembly. Interactions between sexual and mimetic demands of just this character have been described by Stride (1956, 1957). Associations between specialized procryptic coloration and highly evolved habitat-selection could also bring about the same result. In the present case neither of these restraints would appear to operate. The majority of species in both families are highly and efficiently distasteful; they are not, in general, cryptic at rest, and since they are fully nocturnal (with the exception of the Ctenuchinae and some Euchromiinae), there is little likelihood that their courtship involves visual signals. Yet the course of evolution within these families seems often to have involved the maximum divergence of color patterns, even within genera. This is particularly striking in the case of the Automolis spp. illustrated in Text-figs. 7-12; all have Type III displays, and some, at least, are unquestionably closely related, (A. chionoplaga, A. priscilla and A. rutila, for example), yet the overall patterns are very distinct. Of the 16 Automolis found on Barro Colorado, only three (A. elissa, A. elissoides and A. salma) are closely similar to each other. On the other hand, there are apparently convergences across genera (some Episcepsis, Delphyre atava and Aclytia ventralis are much alike) or even families (e.g., the arctiid Automolis rutila and part of the ctenuchine genus Aclytia), and even participation in outside mimetic complexes (for example, wasp-mimicry, carried to such extremes as in the ctenuchid genera Pseudosphex (Beebe & Kenedy, 1957), and Pterygopterus, and the association of Correbia and Correbioides spp. (Ctenuchidae) with lycid beetles and their mimics; a North American lycid mimetic assembly is discussed in detail by Linsley, Eisner & Klots (1961). Finally, there are those genera which have maintained a close internal similarity; many of the numerous species of Macrocneme can only be distinguished reliably by genital characters, and the large distasteful genera Eucereon and Episcepsis each contains numerous closely similar species.

These discrepancies between simple prediction and fact need explanation; at the least they suggest the hazards of facile evolutionary speculation. Although the confusing and contradictory situation which is presented could be explained on the assumption that much of the adult coloration results from pleiotropic genes whose adaptive significance lies in other spheres, it is preferable to assume that the adult patterning is adaptive. Given this premise, two types of problem are presented:

(i) The case represented by Automolis, where

intrageneric divergence has been achieved without, in general, any close Müllerian resemblance to co-existent species outside the divergent group.

(ii) The case presented by many ctenuchine and euchromiine species where divergence has implemented Müllerian resemblances to nonlepidopterous insects.

These trends can only be understood in relation to the behavior of predator populations, and the type of information that is needed for their analysis is almost wholly lacking. The selection pressures are perhaps most likely to relate to the distribution of feeding habits within the assemblies of predator species. Research to date has been largely concerned to demonstrate that the learning and discriminatory capacities of single predator species are a sufficient explanation of the mimetic detail which is actually found in nature. It is well known that those highly aposematic and distasteful species which have been studied prove to be the victims of one or more natural predators which are undeterred by devices which are effective against more fastidious enemies (Cott, 1949). For example, whereas highly distasteful hemileucine saturniids are rejected consistently by marmosets, the less fastidious, and, in caged colonies, more highly motivated Cebus will take them, and quite soon learn to rub off the distasteful vestiture of hairs and scales (Blest, 1963b, and unpublished data). It has been correctly recognized that the existence of such predators, immune to the protective mechanisms of particular species, does not invalidate the general evolutionary argument. But the merely partial efficiency of these adaptations has not been seen as an active selective force in its own right.

Avian predators, so far as is known, learn to search for their prey within ecological limits doubtless determined by inborn releasing mechanisms, and they must also learn to avoid distasteful species. In an inexperienced predator population, the discovery and consumption of a palatable prey increases the probability that more prey of the same kind will be taken (de Ruiter, 1956: Blest, 1963b), and, if Tinbergen's (1960) hypothesis of "specific search images" retained over relatively long periods by the members of predator populations proves valid (Gibb, 1962), the behavior of the predator populations themselves is altered by experience. It would follow, then, that the existence of specific search images could provide a source of selection pressures promoting divergence within related aposematic forms, if there were even one species of predator for whom the group as a whole proved palatable. For the greater the population density of visually similar forms, the more rapidly will a specific

search image directed to them be set up; and this adaptively undesirable consequence of Müllerian resemblance will be attenuated if the similarities are broken down and replaced by divergences in the course of evolution. In fact, the situation would necessarily impose simple direct selection for divergent forms, which might or might not be balanced by the factors promoting Müllerian convergence. The plausibility of this hypothesis can only be assessed when much more is known about the pattern of avian predation in tropical forest. In particular, the relatively narrow ecological niches which are said to be characteristic of tropical birds (MacArthur & MacArthur, 1961; Klopfer & MacArthur, 1961; Klopfer, 1962) may prove to have special selective consequences for insects with spatial distributions which are complex in relation to those of their potential predators.

In contrast to what has been claimed for the saturniids (Blest, 1957), there is no clear derivation for the display behavior of these two families. Although there are rhythmic displays of the type which have been assumed to be transitional between flight behavior and static displays (Blest, 1957), there is no evidence of intermediate types; all the displays are static save for the leg-quivering movements found in some Bertholdia, etc., and a few other specialized components, and wing-waving appears only as an accompaniment of walking. The evolutionary origins of the static displays are consequently obscure. Rhythmic displays in which the wings are slowly raised and lowered without locomotion are performed by the North American Diacrisia virgo, but nothing can be said about the relationship which they may bear to the present tropical behavior patterns. Nor, as was hoped when the study was started, is the origin and precise function of sound production suggested by comparative study. All that contemporary species seem able to demonstrate is that when unpalatability has reached a certain degree, the capacity to produce ultrasonic sounds is lost, presumably because it confers no further advantage. At present, too little is known about the treatment of unpalatable prey by bats for their significance to be assessed.

#### 9. SUMMARY

 The protective display behavior of 183 New World species of arctiid and ctenuchid moth from Trinidad and Panama is summarized, and compared with that of the Saturniidae. Their displays are characterized by the independent evolution of a large variety of "specialized" devices, sometimes developed uniquely, but often showing intergeneric convergence. All but a few displays are static and there are virtually no rhythmic components. The most advanced forms seem to rely upon "reflex immobilization" as a defence.

- 2. Most members of the two groups possess tymbal organs on the metathorax, but only a minority of species use them to produce ultrasonic signals when handled, and the ability to do so is lost in the advanced ctenuchids, although some ctenuchines transitional between the two families (e.g., Eucereon, Episcepsis) readily emit sounds. There is no correlation between the mode of protective coloration and the tactile threshold for sound-production, save that in the more palatable phaegopterine arctiids (e.g., Melese, Bertholdia and Ochrodota spp.) the threshold for sound-production is lower than in those which are more distasteful (e.g., Automolis, Robinsonia and Cissura spp.).
- 3. The possible role of environmental diurnal temperature in determining the course of the evolution of protective displays is discussed.
- 4. It is suggested that the families show less simple Müllerian convergence within genera than might be predicted from their almost uniform unpalatability. Reasons for this are discussed, with particular reference to L. Tinbergen's hypothesis of the "specific search images" of birds.

#### 10. LITERATURE

## Allee, W. C.

- 1926. Measurement of environmental factors in the tropical rain forest of Panama. Ecology, **7**, 445-468.
- BEARD, J. S.
  - 1964. The natural vegetation of Trinidad. Oxford.
- BEEBE, W.
  - 1952. Introduction to the ecology of the Arima Valley, Trinidad, B.W.I. Zoologica, **37**, 157-183.
- BEEBE, W., & ROSEMARY KENEDY
  - 1957. Habits, palatability and mimicry in thirteen ctenuchid moth species from Trinidad, B.W.I. Zoologica, 42, 147-158.
- BENNETT, C. F.
  - 1963. A phytophysiognomic reconnaissance of Barro Colorado Island, Canal Zone. Smith. Misc. Coll., 145 (7), 1-8.

# BISSET, G. W., J. F. D. FRAZER, MIRIAM ROTHSCHILD & M. SCHACHTER

1960. A pharmacologically active choline ester and other substances in the garden tiger moth *Arctia caja* (L). Proc. Roy. Soc. Lond, Ser. B, **152**, 255-262. BLEST, A. D.

- 1957. The evolution of protective displays in the Saturnioidea and Sphingidae. Behaviour, 11, 257-310.
- 1960a. The resting position of *Cerodirphia speciosa* (Cramer), (Lepidoptera, Saturniidae): the ritualization of a conflict posture. Zoologica, N.Y., **45**, 81-90.
- 1960b. The evolution, ontogeny and quantitative control of the settling movements of some New World Saturniid moths, with some comments on distance communication by honey-bees. Behaviour, **16**, 188-253.
- 1963a. Relations between moths and predators. Nature, 197, 1046-1047.
- 1963b. Longevity, palatability and natural selection in five species of New World saturniid moth. Nature, **197**, 1183-1186.

BLEST, A. D., T. S. COLLETT & J. D. PYE

- 1963. The generation of ultrasonic signals by a New World arctiid moth. Proc. Roy. Soc. Lond., Ser. B., **158**, 196-207.
- BROWER, JANE V. Z.
  - 1958. Experimental studies of mimicry in some N. American butterflies. I. Danaus plexippus and Limenitis archippus archippus. Evolution, 12, 32-47.
  - Experimental studies of mimicry. 4. The reactions of starlings to different proportions of models and mimics. Amer. Nat., 94, 271-282.

BROWN, W. L., & E. O. WILSON

1956. Character displacement. Systematic Zoology. 5, 49-64.

CARPENTER, G. D. H.

- 1938. Audible emission of defensive froth by insects. Proc. Zool. Soc. Lond., A, 108, 243-252.
- Сотт, Н. В.

1949. Adaptive Coloration in Animals. London.

DE RUITER, L.

1956. Countershading in caterpillars. An analysis of its adaptive significance. Arch. Néerl. Zool., 11, 285-341.

#### FLEMING, H.

- 1957. The Ctenuchidae (moths) of Trinidad, B.W.I. Part I. Euchromiinae. Zoologica, 42, 105-130.
- 1959. The Ctenuchidae of Trinidad, B.W.I. Part II. Ctenuchinae. Zoologica, **44**, 85-104.

Forbes, W. T. M.

1939. The Lepidoptera of Barro Colorado Island, Panamá. Bull. Mus. Comp. Zool. Harvard, 85, No. 4, 99-322. GIBBS, J.

1962. L. Tinbergen's hypothesis of the role of specific search images. Ibis, **104**, 106-111.

HUHEEY, J.

1961. Studies in warning coloration and mimicry. author. III. Evolution of Müllerian mimicry. Evolution, 15, 567-568.

KAYE, W. J., & N. LAMONT

1927. A catalogue of the Trinidad Lepidoptera Heterocera, (moths). Mem. Dept. Agric. Trinidad and Tobago, No. 3, iv-144.

KLOPFER, P. H.

- 1962. Behavioural Aspects of Ecology. Prentice-Hall, N.J., U.S.A.
- KROGH, A., & E. ZEUTHEN
- 1941. The mechanism of flight preparation in some insects. J. Exp. Biol., 18, 1-10.

LINSLEY, E. G., T. EISNER & A. B. KLOTS

1961. Mimetic assemblages of sibling species of Lycid beetles. Evolution, 15, 15-29.

MACARTHUR, R. & J. MACARTHUR

1961. On bird species diversity. Ecology, 42, 594-598.

MACARTHUR R. & P. H. KLOPFER

1961. On the causes of tropical species diversity: niche overlap. Amer. Nat., **95**, 223-226.

SEITZ, A. (ed.)

1915-1917. The Macrolepidoptera of the World. Vol. VI. Stuttgart.

SHEPPARD, P. M.

- 1959. Natural Selection and Heredity. Hutchinsons, London.
- 1961. Some contributions to population genetics resulting from the study of the Lepidoptera. Adv. Genet, **10**, 165-216.
- STRIDE, G. O.
  - 1956.. On the courtship behaviour of Hypolimnas misippus L. (Lepidoptera, Nymphalidae) with notes on the mimetic association with Danaus chrysippus L. (Lepidoptera, Danaidae). Brit. J. Anim. Behav., 4, 52-68.
  - 1957. Investigations into the courtship behaviour of the male of *Hypolimnas misippus* L. (Lepidoptera, Nymphalidae), with special reference to the role of visual stimuli. Brit. J. Anim. Behav. **5**, 153-167.

TINBERGEN, L.

1960. The dynamics of insect and bird populations in pine woods. Arch. Néerl. Zool., 13, 159-273.

VECHT, J. VAN DER

1961. Evolution in a group of Indo-Australian Eumenes (Hymenoptera, Eumenidae). Evolution, 15, 468-477.



Blest, A. D. 1964. "Protective display and sound production in some New World arctiid and ctenuchid moths." *Zoologica : scientific contributions of the New York Zoological Society* 49(10), 161–181. <u>https://doi.org/10.5962/p.203299</u>.

View This Item Online: <a href="https://www.biodiversitylibrary.org/item/208153">https://doi.org/10.5962/p.203299</a> Permalink: <a href="https://www.biodiversitylibrary.org/partpdf/203299">https://www.biodiversitylibrary.org/partpdf/203299</a>

Holding Institution Smithsonian Libraries and Archives

**Sponsored by** Biodiversity Heritage Library

**Copyright & Reuse** Copyright Status: In Copyright. Digitized with the permission of the rights holder Rights Holder: Wildlife Conservation Society License: <u>http://creativecommons.org/licenses/by-nc/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.